

Climate change in the longer term

The foregoing calculations have focussed on the period up to the year 2100; it is clearly more difficult to make calculations for years beyond 2100. However, while the timing of a predicted increase in global temperatures has substantial uncertainties, the prediction that an increase will eventually occur is more certain. Furthermore, some model calculations that have been extended beyond 100 years suggest that, with continued increases in greenhouse climate forcing, there could be significant changes in the ocean circulation, including a decrease in North Atlantic deep water formation.

Other factors which could influence future climate

Variations in the output of solar energy may also affect climate. On a decadal time-scale solar variability and changes in greenhouse gas concentration could give changes of similar magnitudes. However the variation in solar intensity changes sign so that over longer timescales the increases in greenhouse gases are likely to be more important. Aerosols as a result of volcanic eruptions can lead to a cooling at the surface which may oppose the greenhouse warming for a few years following an eruption. Again, over longer periods the greenhouse warming is likely to dominate.

Human activity is leading to an increase in aerosols in the lower atmosphere, mainly from sulphur emissions. These have two effects, both of which are difficult to quantify but which may be significant particularly at the regional level. The first is the direct effect of the aerosols on the radiation scattered and absorbed by the atmosphere. The second is an indirect effect whereby the aerosols affect the microphysics of clouds leading to an increased cloud reflectivity. Both these effects might lead to a significant regional cooling; a decrease in emissions of sulphur might be expected to increase global temperatures.

Because of long-period couplings between different components of the climate system, for example between ocean and atmosphere, the earth's climate would still vary without being perturbed by any external influences. This natural variability could act to add to, or subtract from, any human-made warming; on a century timescale this would be less than changes expected from greenhouse gas increases.

How much confidence do we have in our predictions?

Uncertainties in the above climate predictions arise from our imperfect knowledge of:

- future rates of human-made emissions
- how these will change the atmospheric concentrations of greenhouse gases
- the response of climate to these changed concentrations

Firstly, it is obvious that the extent to which climate will change depends on the rate at which greenhouse gases (and other gases which affect their concentrations) are emitted. This in turn will be determined by various complex economic and sociological factors. Scenarios of future emissions were generated within IPCC WGIII and are described in the annex.

Secondly, because we do not fully understand the sources and sinks of the greenhouse gases, there are uncertainties in our calculations of future concentrations arising from a given emissions scenario. We have used a number of models to calculate concentrations and chosen a best estimate for each gas. In the case of carbon dioxide, for example, the concentration increase between 1990 and 2070 due to the Business-as-Usual emissions scenario spanned almost a factor of two between the highest and lowest model result (corresponding to a range in radiative forcing change of about 50%).

Furthermore, because natural sources and sinks of greenhouse gases are sensitive to a change in climate, they may substantially modify future concentrations (see earlier section: "Greenhouse gas feedbacks"). It appears that, as climate warms, these feedbacks will lead to an overall increase, rather than decrease, in natural greenhouse gas abundances. For this reason, climate change is likely to be greater than the estimates we have given.

Thirdly, climate models are only as good as our understanding of the processes which they describe, and this is far from perfect. The ranges in the climate predictions given above reflect the uncertainties due to model imperfections; the largest of these is cloud feedback (those factors affecting the cloud amount and distribution and the interaction of clouds with solar and terrestrial radiation), which leads to a factor of two uncertainty in the size of the warming. Others arise from the transfer of energy between the atmosphere and ocean, the atmosphere and land