

angle of the occiput; prothorax with a longitudinal intermediate ridge, transversely streaked with yellow and brownish-black; hemelytra sprinkled, and towards the base almost streaked, with black and yellow. In other respects this species resembles *C. striata*.

398. *CORIXA PLANIFRONS* Kirby.—Length of body 4 lines. Two specimens taken with the preceding.

This species differs from the preceding in having the under side of the body black, with two pale spots on each side of the breast, and the anal half of the abdomen pale-yellow. The head is yellow, the vertex is ridged longitudinally and separated from the front by a transverse curvilinear ridge; and from which the anterior part of the face is inflexed, plane or slightly concave; in other respects this species exhibits exactly the same characters with *C. carinata*. They may perhaps be sexual varieties.

[285.] FAMILY NOTONECTIDÆ.

399. *NOTONECTA INSULATA* Kirby.—Length of body $6\frac{2}{3}$ lines. A single specimen taken.

Body underneath black, above yellowish. Head yellowish, with a brownish longitudinal stripe between the eyes, which are reddish; scutellum very black, velvety; hemelytra with the lateral margin brown spotted with yellow; the posterior half of the hemelytra is brown anteriorly, black-brown in the middle, fuliginous at the apex; in this darkened portion anteriorly is a yellowish triangle connected with the last marginal spot, and posteriorly is a white kidney-shaped spot followed by the sooty apex; breast very hairy with longish fuliginous hairs; legs yellowish.

VII. HOMOPTERA.

FAMILY CERCOPIDÆ.

400. *CERCOPIS MARGINELLA* Fabr.—Length of body $3\frac{1}{4}$ lines. Taken with the preceding, and at Carlton-house in April.

Body black, dotted and inscribed with white. Hemelytra embrowned; nervures black; lateral margin sanguine; margin of the abdomen edged with white.