

*Lycaena pseudargiolus*, Bd.-Lec.

The form *Pseudargiolus* is very rare; found mostly in July and August.

Var. *Violacea*, Edw.

This form is common and seems to be found at all times through the spring and summer.

Var. *Neglecta*, Edw.

This also is common and most so during the summer months.

*Lycaena comyntas*, Gdt.

Common, and is found as late as October.

*Lycaena isola*, Reak. or *Alce.*, Edw.

I have taken a few specimens but only in October; have not seen anything of them at any other time.

*Pieris protodice*, Bd.-Lec.

Common through the summer.

*Pieris rapae*, Linn.

Too common from April to October.

*Nathalis iole*, Bdv.

Rare; found only in September.

*Colias eurhytheme*, Bdv.

Very common, from April to November; have taken a few Albinos of this species in October.

Var. *Keewaydin*, Edw.

Not uncommon; a few taken in August, but most in latter part of September and October.

*Colias philodice*, Gdt.

This is our most common butterfly, and is found from early in April to middle of November. It is more common than *C. eurhytheme*.

*Colias caesonina*, Stoll.

Common through summer months.

*Terias nicippe*, Cram.

Rare; have taken but one specimen, but that was perfectly fresh; October.

*Terias lisa*, Bd.-Lec.

Common, but most so in September and October. Have taken white females of this species.

*Papilio ajax*, Linn.

Not uncommon in variety *marcellus*, Bdv., but *telamonides* is rare. This species I found very hard to capture on account of the colour and the skill they have in dodging the stroke of the net.