145. Leptura kerniana Fall.

Taken by Daggett on the Kern River.

146. Leptura plagifera LeConte.

Taken rarely in northern Sierras. Found especially on the eastern sides of the Cascades.

147. Leptura rubida LeConte.

Rare. Taken in Northern California, Coast belt to San Francisco, and in Sierras—a different colour phase in each section.

148. Leptura subargentata Kirby.

Common at Camp Meeker, Sonoma County.

var. ruficeps LeConte.

Found throughout the Sierras.

var rhodopus LeConte.

Breeds in tan bark oak.

var. similis Kirby.

149. Leptura molybdica LeConte.

Fairly common throughout the State on various flowers. Taken by Fall at Pasadena on Ceanothus; by Van Dyke in Sierra County, and the middle and northern Sierras.

150. Leptura læta LeConte.

Found throughout the State; not abundant anywhere, most common in the foothills of the Sierras. Found by Van Dyke on flowers, especially those of California wild holly. Bred by Rivers from live oak. Taken by author at Donner Lake, July 12.

151. Leptura tribalteata LeConte.

Found in Northern California and Sierras as far south as Walker Basin. Taken by author at Donner Lake July 6 on meadow flowers.

152. Leptura coquilletti Fall.

Taken on flowers of *Eriogonum fasciculatum* in June by Ricksecker. Taken by Van Dyke in Sierra Madre Mts., San Bernardino Mts., and also Los Angeles County.