

camera lucida. The whole process of mounting may be accomplished in a minute or little more, or it may be mounted and sketched in five minutes or less. In this way I have mounted and sketched hundreds of wings of Tineidæ, which I hope some day to publish in the pages of the CAN. ENT.

V. T. CHAMBERS.

Colorado Springs, Colorado, Feb., 1876.

---

ENTOMOLOGICAL NOTES.

DEAR SIR,—

*Ceratonia quadricornis*.— I have during the past season, and for the first time in my life, taken a number of the larvæ of *Ceratonia quadricornis* Harr., of a brown color. Out of five taken in one locality, four were of a deep brown. The fifth was of the usual green hue. These larvæ were about half grown when captured. This was early in September, but since then I have taken several other specimens, the majority of which were brown. I should like to learn whether any of your readers have, at any time, made similar captures.

*Doryphora 10-lineata*.— A safe and effective method of destroying this beetle is to drag an ordinary butterfly net over the haulm when the larvæ and beetles are feeding. By this method, and without making any particular effort, I captured on Long Island, this year, over five hundred specimens in about as many minutes. By making a sort of drag net, large enough to be worked by two boys, a much larger number could of course be captured in a given time. I have no doubt that Long Island will be plentifully supplied with *10-lineata* next year; for, although thousands were drowned and washed up on the beach at Coney Island and elsewhere, the number that escaped must have been great. But, although admitting the desirability of ridding ourselves of this pest, I by no means admit that it is the scourge talked about by some Entomologists. As I intimated before, we had plenty of them on this Island, but no one complains of a small crop of potatoes in consequence.

W. V. ANDREWS.

36 Bøerum Place, Brooklyn.

---

ERRATA.—On page 16, line 7 from top, for *Heliophila renipuncta* read *H. unipuncta*.