






Good Seed from a Pit.




Two-man Planter.




Cutting by Hand.

Potatoes Badly Affected with Scab.
They should never be planted without being treated for disease.

Using a Machine for Cutting Seed.




One-man Planter.




An Undesirable Type—Rough and Deep-eyed.



The Type of Tubers to Select for Seed.



Plowing the Potato Field and Dropping the Sets in Every Third Furrow.

Type of Potato which Markets Prefer.
Smooth, shallow-eyed tubers weighing eight ounces each.

place to
as they us
take from
leave the
which can
which hav
have been
as strong
atoes are
and left t
planting.
out where
them with
strong, gre
the slender
These spr
injured in
of any pla
previous to
er growth

This dis
siderable lo
noticed at
before cutt
propagation
treatment i
susceptible
years. It n
produced o
potatoes. A
of the fung
green crop
amount of
manure fro
affected wit
potato scab
two hours i
gallons of w
sacks or bin
This treatm
as it is claim
will weaken
may also be
are dissolv
stand for a t
gallons. Th
half hours.
remembered
left after tre
be fatal to l
solution is s
potatoes. T
Rhizoctonia
in some sect
Powdery s
scab, but as y
in this count
prevent it g
Only seed tha
be planted an
seed should be
are other dise
may be seen a
dark rings or
some fungous
that would in
Potato grower
seed they plan

Some grower
potatoes for s
and is never e
seed end was
it was thrown
cut at a certai
custom is fast
is planted a m
the sets a cer
number of eye
work with the
in size and sha
a machine can
which is suffici
The stand of c
about equal to
hand.

While the s
thin portion, n
eye on it, to wh
the set as thick
one eye must be
Sufficient plant
the young pla
se has a t
tubers that f
fertile, and we
may consist of
to be little a
and the cost o
amount require
high as twenty
sets and distan
feet apart and
thirteen bushels
the sets weigh
nineteen bushels
is about the ave
Freshly-cut t
as much as possi
quickly, in order