found July 15th was placed in the side of a small mound, and was well concealed by mosses and overhanging ferns. The nest was composed of grasses and moss and lined with fruit stems of hair moss. The female flushed from the nest and disappeared among the numerous, small evergreen bushes that grew near the nest, nor did she put in an appearance during the half hour I stayed near. The nest measured, outside diameter 3 inches, inside 1.75, outside depth 2 inches, inside 1 inch, and contained three eggs, white, marked with rufous and cinnamon-brown, chiefly about the larger end.

Tennessee Warbler, Helminthophila paregrina.

Have observed this rather rare species but have never become acquainted with its habits. It arrives late in May or early June.

PARULA WARBLER, Comsopthlypis americana.

A fairly common migrant and summer resident, arriving here the first half May. Its shivering skew-ee-oo song is most often emitted from a high perch in tall trees. The writer has found two nests of this species, one of which was situated forty feet up in a yellow birch tree, the other was twenty feet up in a beech. Both nests were made in *Usnea* lichen where it hung below the limb on which it grew. The lichen had been shaped at a distance of four to six inches below the limbs, and sewn with horse hairs to keep in shape for a nest. The cavity of the nest in the beech measured an inch and one half in both depth and diameter, and contained four eggs, white with rufous markings about the larger end.

The last bird observed during 1902 was on September 17th.

CAPE MAY WARBLER, Dendroica tigrina.

As a rule this species is a rare summer resident. But during the summer of 1902 it was tolerably common. The first spring migrant recorded that season was on May 17th. It frequents young evergreen thickets, and in such places one may hear their seep. seep, seep, seep, of a song and the bird be entirely hidden from view.

YELLOW WARBLER, Dendroica æstiva.

After May 20 this species is a common summer resident along the river valleys, where it breeds in the shrubbery and is especially