Measurements.—Length of body, 3, 17.5 mm.; 9, 19 mm.; of pronotum, 3, 4.5 mm.; 9, 5 mm.; of tegmina, 3, 20.5 mm; 9, 24.5 mm.; of antennæ, 3, 46 mm.; of posterior femora, 3, 17 mm.; 9, 17.5 mm.; of ovipositor, 7 mm.

This dull coloured grasshopper has been found in small numbers in both Vigo and Fulton counties, in upland prairie meadows, where it frequents the tall grasses, usually in company with *Xiphidium strictum*, Scudder.

It is a smaller and more slender bodied insect than the common O. vulgare, Harris, and has a shorter and narrower pronotum and a much smaller ovipositor than that species.

Orchelimum Bruneri, nov. sp.

A species of about the same length, but less robust than O. vulgare, Harris, having the posterior femora armed beneath, and the ovipositor very broad, nearly straight and of more than average length.

Cone of the vertex narrow, moderately elevated, rounded at apex. Tegmina long and narrow, a little shorter than the wings. Posterior femora rather stout, the apex, when appressed, not quite reaching the tip of ovipositor; armed beneath on the apical half with three or four small spines. Cerci of male stout, acuminate, with the internal tooth prominent.

Ovipositor very similar to that of O. gladiator, Bruner, being very long and stout, nearly straight above, and with the under side of apical third sloping rapidly to the acute apex.

Colour of dried specimens.—With the exception of the ovipositor, which is a light reddish-brown, and the usual stripe on occiput and disk of pronotum, the whole body is a pale, transparent brownish-green, the green showing plainly only on the lower half of the side of pronotum and on the meso and metapleura. The reddish-brown dorsal stripe of occiput and pronotum is bordered laterally throughout its *entire* length with a very narrow one of much darker brown. When immersed in alcohol the reddish-brown stripe fades to a yellowish white, leaving the two lateral ones as prominent dark streaks, widest on the central portion of the frontal disk.

Measurements.—Length of body, \mathcal{J} , 18 mm.; \mathcal{Q} , 20.5 mm.; of tegmina, \mathcal{J} , 21 mm.; \mathcal{Q} , 25 mm.; of pronotum, \mathcal{J} and \mathcal{Q} , 4.75 mm.; of hind femora, \mathcal{J} , 16.5 mm.; \mathcal{Q} , 18 mm.; of ovipositor, 10 mm. Described from 2 \mathcal{J} s and 4 \mathcal{Q} s.