

**CIHM
Microfiche
Series
(Monographs)**

**ICMH
Collection de
microfiches
(monographies)**



Canadian Institute for Historical Microreproductions / Institut canadien de microreproductions historiques

© 1998

The copy filmed here has been reproduced thanks to the generosity of:



Special Collections,
D.B. Weldon Library,
University of Western Ontario

The images appearing here are the best quality possible considering the condition and legibility of the original copy and in keeping with the filming contract specifications.

Original copies in printed paper covers are filmed beginning with the front cover and ending on the last page with a printed or illustrated impression, or the back cover when appropriate. All other original copies are filmed beginning on the first page with a printed or illustrated impression, and ending on the last page with a printed or illustrated impression.

The last recorded frame on each microfiche shall contain the symbol \rightarrow (meaning "CONTINUED"), or the symbol ∇ (meaning "END"), whichever applies.

Maps, plates, charts, etc., may be filmed at different reduction ratios. Those too large to be entirely included in one exposure are filmed beginning in the upper left hand corner, left to right and top to bottom, as many frames as required. The following diagrams illustrate the method:



L'exemplaire filmé fut reproduit grâce à la générosité de:


Special Collections,
D.B. Weldon Library,
University of Western Ontario

Les images suivantes ont été reproduites avec le plus grand soin, compte tenu de la condition et de la netteté de l'exemplaire filmé, et en conformité avec les conditions du contrat de filmage.

Les exemplaires originaux dont la couverture en papier est imprimée sont filmés en commençant par le premier plat et en terminant soit par la dernière page qui comporte une empreinte d'impression ou d'illustration, soit par le second plat, selon le cas. Tous les autres exemplaires originaux sont filmés en commençant par la première page qui comporte une empreinte d'impression ou d'illustration et en terminant par la dernière page qui comporte une telle empreinte.

Un des symboles suivants apparaît sur la dernière image de chaque microfiche, selon le cas: le symbole \rightarrow signifie "A SUIVRE", le symbole ∇ signifie "FIN".

Les cartes, planches, tableaux, etc., peuvent être filmés à des taux de réduction différents. Lorsque le document est trop grand pour être reproduit en un seul cliché, il est filmé à partir de l'angle supérieur gauche, de gauche à droite, et de haut en bas, en prenant le nombre d'images nécessaire. Les diagrammes suivants illustrent la méthode.



MICROCOPY RESOLUTION TEST CHART

(ANSI and ISO TEST CHART No. 2)



2.8



5.0

5.6

3.2



6.3

3.6

7.1

8.0

4.0



9.0

10

11.2

12.5

14

16

18

20



APPLIED IMAGE Inc

1653 East Main Street
Rochester, New York 14609 USA
(716) 482 - 0300 - Phone
(716) 288 - 5929 - Fax

A RARE FORM OF EXTRAUTERINE PREGNANCY.

BY

BRICE W. GOLDSBOROUGH, M.D.,

Chief of Staff of the Cambridge Hospital, Cambridge, Md.,

AND

THOMAS S. CULLEN, M.B. (Tor.,)

Associate Professor of Gynecology in the Johns Hopkins University.

On February 28, 1901, Dr. Goldsborough was called in by Dr. I. N. Tannar of Vienna, Maryland, to see what the doctor supposed to be a case of obscure pregnancy. The patient had 1 child 9 years ago. In April, 1900, she missed her period and since then presented the usual signs of pregnancy; nausea, enlarged breasts, increase in size of the abdominal girth. In August, while lifting some boxes, something suddenly gave way in her left side. This occasioned severe pain and she had to remain in bed until November 1. About the middle of September there was a bloody uterine discharge, and accompanying it was considerable pain and nausea. Subsequently, she had several similar discharges which may have been menstrual periods. During the month of November she was able to be out of bed, but had to return in December. Throughout the entire illness she has had a good appetite, has been fairly well nourished. When seen her temperature was 101.5° ; her pulse 140. Immediate removal to the Cambridge Hospital was advised, and on the following day she was driven 23 miles.

On examining the patient under anesthesia the abdomen is seen to be very prominent. There is, however, no bulging in the flanks. The umbilicus is converted into a tumor fully 5 cm. long by 3 cm. broad (Fig. 1). The skin over this appears to be much thinned out and at one point has given way. From this abraded area a chocolate colored fluid is escaping. This is exceedingly offensive. Around the umbilicus the tissue is markedly indurated and pits on pressure. On vaginal examination the cervix is found to be intact, but it is impossible to outline the uterus. Nothing can be detected laterally. After cleansing the abdomen as far as possible, an incision was made just below the sternum and continued down almost to the pubes. The abdominal cavity proper was not exposed; that is to say, none of the abdominal contents came into view. Filling the cavity was a large quantity of chocolate-colored fluid; a



fetus between 6 and 7 months and a large placenta. The placenta was attached low down in the pelvis, was exceedingly friable, but came away without producing any hemorrhage. The walls of the sac were about 4 mm. in thickness and excessively friable. They reminded one very much of granulation-tissue. It is impossible to determine where the pregnancy took place, as the pelvic organs were entirely walled out. It is probable, however, that the uterus ruptured and that the fetus with its membranes



Fig. 1.—The drawing of course is somewhat diagrammatic. It represents a longitudinal section of the body. The fetus with the fetal membranes are lying immediately beneath the abdominal wall and are attached anteriorly to the peritoneum almost from the sternum to the pubes. (B.) At the umbilicus the fetal sac hinges into the hernial opening and at the most prominent point this hernial sac has given way, allowing the fluid to escape externally. At A numerous granulations have formed on the inner surface of the sac. The fetus is well preserved, appears to be about 6 months old and shows slight maceration on the face, on the arms and legs. The site of the placenta is roughly outlined by the dotted lines. The cervix is seen to be normal but on account of the marked distortion, the presence of the abdominal tumor and the oedema it was impossible to outline the uterus or appendages, hence their relation is left hazy. The bladder and rectum are in their normal positions. As will be seen from the drawing a median incision in the abdominal wall would open directly into the sac and in no way involve the general peritoneal cavity.

intact escaped. It has then become surrounded by granulation tissue, this latter extending to the pubes and the abdomen was loosely packed. An incision was closed and drainage. At the operation of Dr. John M. Goldsborough a

Following normal to 101. has shown no regular for 6 days tone. The pain the escape of a gauze drain abdomen was disappeared. On discharge. On to outline the about the size directly behind

Pathological Number 4,744) its accompanying itself is 17 cm. to the heel is 2 no external abdominal moderate quantity peeled off. The in length. It is approximate some places it especially in the geneous, is hem

Histological parts of the pla

escaped into the abdominal cavity. The fetal membranes then become attached to the abdominal wall and to the surrounding structures. After removing the fetus and the membranes, this large sac which extended almost from the sternum to the pubes and laterally filled the entire anterior portion of the abdomen was thoroughly washed out with salt solution and then packed with iodoform gauze. The upper half of the abdomen was closed, the lower half left open to insure thorough drainage. At the time of operation the patient's pulse was 140. The operation occasioned no shock. The anesthetic was given by Dr. John Mace, and assisting the operator were Drs. Briceborough and Guy Steele.

Following the operation the temperature ranged from 101.5° for the first 4 days, since which time it has shown no elevation. The pulse was weak and irregular for 6 days, but since then has regained its normal rate.

The pack was removed on the seventh day with the escape of a moderate amount of discharge. A light gauze drain was then inserted. On March 13 the abdomen was perfectly flat and all evidence of edema had disappeared. On removing the drain there was a slight discharge. On bimanual examination it was now possible to outline the uterus to some extent. The organ was the size of a 2 months' pregnancy and situated posteriorly behind the pubes. It was slightly movable.

Pathological Report (Gynecological Pathological Report No. 4,744). The specimen consists of a foetus with accompanying placenta. The foetus when folded upon measures 17 cm. in length. The distance from the occiput to the heel is 29 cm. The child is well formed, shows no external abnormality and is a female. There is a moderate quantity of hair but the skin has to a great extent macerated and the pigmented layer is readily rubbed off. The umbilical cord appears to be about 8 cm. in length. It shows nothing of interest. The placenta measures approximately 16 x 10 x 5 cm. It is very friable. In places it presents the usual appearance, in others, especially in the depth, the tissue is somewhat homogeneous, is hemorrhagic and suggests breaking down.

Microscopical examination of sections from various parts of the placenta show that it consists almost entirely

of necrotic tissue and canalized fibrin. The contours of the villi are everywhere visible but the nuclei of the epithelial cells as well as those of the stroma of the villi have entirely disappeared. The central portions of numerous villi are partially filled with calcareous plaques. At one point are a moderate number of disintegrated polymorphonuclear leukocytes. Otherwise the entire tissue is devoid of nuclei.

This complete necrosis of the placenta accounts for the ease with which it was peeled off and also for the absence of hemorrhage during its removal.

