






# **IMAGE EVALUATION TEST TARGET (MT-3)**



150mm

6"



**APPLIED IMAGE . Inc**  
1653 East Main Street  
Rochester, NY 14609 USA  
Phone: 716/482-0300  
Fax: 716/288-5989

© 1993, Applied Image, Inc., All Rights Reserved

**CIHM  
Microfiche  
Series  
(Monographs)**

**ICMH  
Collection de  
microfiches  
(monographies)**



**Canadian Institute for Historical Microreproductions / Institut canadien de microreproductions historiques**

**© 1993**

**Technical and Bibliographic Notes / Notes techniques et bibliographiques**

The Institute has attempted to obtain the best original copy available for filming. Features of this copy which may be bibliographically unique, which may alter any of the images in the reproduction, or which may significantly change the usual method of filming, are checked below.

- Coloured covers/  
Couverture de couleur
- Covers damaged/  
Couverture endommagée
- Covers restored and/or laminated/  
Couverture restaurée et/ou pelliculée
- Cover title missing/  
Le titre de couverture manque
- Coloured maps/  
Cartes géographiques en couleur
- Coloured ink (i.e. other than blue or black)/  
Encre de couleur (i.e. autre que bleue ou noire)
- Coloured plates and/or illustrations/  
Planches et/ou illustrations en couleur
- Bound with other material/  
Relié avec d'autres documents
- Tight binding may cause shadows or distortion  
along interior margin/  
La reliure serrée peut causer de l'ombre ou de la  
distortion le long de la marge intérieure
- Blank leaves added during restoration may appear  
within the text. Whenever possible, these have  
been omitted from filming/  
Il se peut que certaines pages blanches ajoutées  
lors d'une restauration apparaissent dans le texte,  
mais, lorsque cela était possible, ces pages n'ont  
pas été filmées.
- Additional comments:/  
Commentaires supplémentaires:

This item is filmed at the reduction ratio checked below/  
Ce document est filmé au taux de réduction indiqué ci-dessous.

10X	14X	16X	18X	22X	26X	30X
12X	14X	16X	20X	J	24X	28X

The copy filmed here has been reproduced thanks to the generosity of:



Library  
Agriculture Canada

The images appearing here are the best quality possible considering the condition and legibility of the original copy and in keeping with the filming contract specifications.

Original copies in printed paper covers are filmed beginning with the front cover and ending on the last page with a printed or illustrated impression, or the back cover when appropriate. All other original copies are filmed beginning on the first page with a printed or illustrated impression, and ending on the last page with a printed or illustrated impression.

The last recorded frame on each microfiche shall contain the symbol ➡ (meaning "CONTINUED"), or the symbol ▽ (meaning "END"), whichever applies.

Maps, plates, charts, etc., may be filmed at different reduction ratios. Those too large to be entirely included in one exposure are filmed beginning in the upper left hand corner, left to right and top to bottom, as many frames as required. The following diagrams illustrate the method:



L'exemplaire filmé fut reproduit grâce à la générosité de:

Bibliothèque  
Agriculture Canada

Les images suivantes ont été reproduites avec le plus grand soin, compte tenu de la condition et de la netteté de l'exemplaire filmé, et en conformité avec les conditions du contrat de filmage.

Les exemplaires originaux dont la couverture en papier est imprimée sont filmés en commençant par le premier plat et en terminant soit par la dernière page qui comporte une empreinte d'impression ou d'illustration, soit par le second plat, selon le cas. Tous les autres exemplaires originaux sont filmés en commençant par la première page qui comporte une empreinte d'impression ou d'illustration et en terminant par la dernière page qui comporte une telle empreinte.

Un des symboles suivants apparaîtra sur la dernière image de chaque microfiche, selon le cas: le symbole ➡ signifie "A SUIVRE", le symbole ▽ signifie "FIN".

Les cartes, planches, tableaux, etc., peuvent être filmés à des taux de réduction différents. Lorsque le document est trop grand pour être reproduit en un seul cliché, il est filmé à partir de l'angle supérieur gauche, de gauche à droite, et de haut en bas, en prenant le nombre d'images nécessaire. Les diagrammes suivants illustrent la méthode.

ONTARIO AGRICULTURAL COLLEGE  
EXPERIMENT STATION

---

BULLETIN LVIII

SPRING GRAINS IN 1890

BY THOMAS SHAW PROFESSOR OF AGRICULTURE

PUBLISHED BY THE DEPARTMENT OF AGRICULTURE

*February 2 1891*

---

TORONTO  
PRINTED BY WARWICK & SONS

O

JAN  
THO  
C. O.  
J. H.

F. C.  
H. H.  
E. L.

THOM  
C. A.  
J. E.  
JAMES

The  
lished b  
charge o  
Ministe

# ONTARIO AGRICULTURAL COLLEGE

## EXPERIMENT STATION.

GUELPH, ONTARIO.

---

### COLLEGE STAFF.

JAMES MILLS, M.A., President, English Literature and Political Economy.  
THOMAS SHAW, Agriculture, Live Stock and Arboriculture.  
C. C. JAMES, M.A., Inorganic, Organic, Agricultural and Analytical Chemistry.  
J. H. PANTON, M.A., F.G.S., Geology, Botany, Zoology, Meteorology and  
Horticulture.  
F. C. GRENSIDE, V.S., Veterinary Anatomy, Pathology, Materia Medica  
and Obstetrics.  
H. H. DEAN, B.S.A., Dairy Husbandry.  
E. L. HUNT, R.A., Assistant Resident and Mathematical Master.

---

### FARM OFFICERS.

THOMAS SHAW, Farm Superintendent and Director of Experiments.  
C. A. ZAVITZ, B.S.A., Assistant Director of Experiments.  
J. E. STORY, Farm Foreman.  
JAMES FORSYTH, Foreman of Horticultural Department.

---

64203

The Bulletins of this Agricultural College and Experiment Station are published by the Department of Agriculture and will be mailed regularly free of charge to all members of Farmers' Institutes. Address, Hon. JOHN DRYDEN, Minister of Agriculture, Toronto.



## BULLETIN LVIII

---

### EXPERIMENTS WITH SPRING GRAINS.

The tendency with every variety of grain that has ever been produced is to deteriorate when grown long upon the same kind of soil and in the same locality. The necessity therefore for the introduction of varieties from other countries, or from distant parts of the same country, and of producing varieties which are entirely new, will always exist. In this fact we find a perpetual argument for the existence of experimental farms.

The object of this Bulletin is to make known to our farmers (1) the varieties of spring grains which grew most successfully at this farm during the summer of 1890, and (2) important comparisons relating to all the varieties grown. These comparisons relate (1) to yield, weight of grain, weight of straw and date of maturing in all the varieties mentioned; (2) to color and other peculiarities of growth, to the structure and weight of the grains, to the relative percentage of the hull and to proneness to injury from rust in some of the varieties named, (3) to the number of varieties tested from different countries and the average yields obtained from them, and (4) to six-rowed and two-rowed barleys respectively, with and without hulls.

The information thus given has been obtained from tests made with 54 varieties of barley, 54 varieties of spring wheat, 20 varieties of pease and 92 varieties of oats. The different varieties of each kind of grain were grown under similar conditions as to soil and treatment. They were sown side by side upon plots containing one-fiftieth of an acre each in the case of the barleys, one-sixtieth of an acre in that of pease, and one-hundredth of an acre in that of the spring wheat and oats.

In the tables given below the yields are based upon the standard weights obtained from the measured bushel, and from samples that were thoroughly cleaned.

**BARLEY.**—The information given in the first table relates to the ten varieties of barley which gave the highest yields in 1889 and 1890. The average yield is given for the two years. The varieties

grown in 1889 were sown on April 15th and those of 1890 on April 19th.

Variety.	Kind.	Country whence obtained.	Bush. per acre.	Lb. of grain per bush.	Tons of straw per acre.	No. of grains per ounce.	Date of maturing.
Cheyne .....	2-rowed.	Germany ..	48.5	53½	2.18	670	Aug. 6
Chevalier .....	2-rowed.	France ..	47.5	52½	1.18	726	" 7
Mandschuri .....	6-rowed.	Russia ..	47.3	51½	1.70	755	July 31
Oderbrucher.....	6-rowed.	Germany ..	45.8	54½	1.42	743	Aug. 1
Empress .....	2-rowed.	England ..	45.4	51½	1.92	720	" 6
Early Black .....	6-rowed.	France ..	44.8	51	1.61	591	July 31
Scotch Improved .....	6-rowed.	Ontario ..	44.4	52	1.27	805	" 31
Selected Chevalier .....	2-rowed.	England ..	43.0	50½	1.75	758	Aug. 7
Common .....	6-rowed.	Ontario ..	41.7	54½	1.29	819	July 31
Golden Drop .....	2-rowed.	Germany ..	41.4	53½	1.84	646	Aug. 5

In addition to these, the following varieties went over 40 bushels per acre, viz.: Hallett's Pedigree (Germany), Phoenix (Germany), Italian Rice (Germany), and Imperial (France). Carter's Prize Prolific is not included in the above list as with us the average yield for the two years was 32.3 bushels per acre on the small plots and the return this year for the acre plot was 31½ bushels, and the weight for the crop of 1890 was 52 lb. per bushel. It appears however to have done much better in many other localities.

**SPRING WHEAT.**—The information given in the second table relates to seven varieties of spring wheat which gave the highest yields in 1890. They were sown April 25th.

Variety.	Country whence obtained.	Bush. per acre.	Lb. of grain per bush.	Tons of straw per acre.	Rust tendency.	Date of maturing.
Herison Bearded .....	France ..	27.2	64	1.90	Very slight...	Aug. 8
White Fyfe .....	Ontario ..	26.9	60	1.89	Considerable.	" 14
Bart Trimenia .....	Greece ..	26.3	64	1.06	Very slight..	" 10
White Russian .....	Ontario ..	22.1	57½	1.91	Considerable.	" 11
Red Fern .....	Ontario ..	21.0	60	1.76	Slight .....	" 11
Pringle's Champion .....	Germany ..	21.2	59½	1.90	Very slight..	" 12
Holben's Improved .....	Germany ..	20.4	58½	1.70	Medium....	" 14

The Herison Bearded has been tested for two years, and was last year ahead of all the foreign varieties and also of all the Canadian

varieties except the Wild Goose. It is a fairly strong grower, stands up well and is almost entirely free from rust. The head is short but compact, and is bearded; the chaff is red and the berry is rather small but plump, and is also red. The Bart Trimenia, though a good yielder and weighs well, is rather coarse in the berry.

**P E A S E.**—The information given in the third table relates to five varieties of peas which gave the highest yields in 1890. They were sown April 24th.

Variety.	Country whence obtained.	Bush. per acre.	Lb. per bush.	Tons of straw per acre.	Date of maturing.
Sweet Jessie .....	England ....	31.16	62 <i>1</i> <sup>2</sup>	.82	Aug. 11
Early Racehorse.....	England ...	29.29	64 <i>1</i> <sup>2</sup>	1.06	" 10
Mummy .....	Ontario.....	28.60	65 <i>1</i> <sup>2</sup>	1.26	" 11
Early Britain.....	England ....	28.37	60	.57	" 11
White Wonder .....	New Zealand.	27.67	64 <i>1</i> <sup>2</sup>	.30	" 5

**O A T S.**—The information given in the fourth table relates to the ten varieties of oats which gave the highest yields for 1889 and 1890. The average yields are given for the two years. They were sown on April 22nd of each year.

Variety.	Country whence obtained.	Bush. per acre.	Lb. per bush.	Tons of straw per acre.	Date of maturing.
Goanette Black .....	France...	71.7	34 <i>1</i> <sup>2</sup>	2.7	July 9
Houdan Black .....	France...	69.1	34 <i>1</i> <sup>2</sup>	2.0	" 9
Chenailles .....	France...	69.1	34 <i>1</i> <sup>2</sup>	2.9	" 8
Oderbrucher .....	Germany.	66.6	31 <i>1</i> <sup>2</sup>	2.3	" 6
Danebrog .....	Germany.	64.0	32	2.7	" 9
Siberian.....	France...	63.6	34 <i>1</i> <sup>2</sup>	2.1	" 2
Black Etampes.....	France...	62.3	36 <i>1</i> <sup>2</sup>	2.5	" 9
Probsteier.....	Germany.	61.8	33	2.6	" 6
Pringle's Progress... .	Ontario..	61.8	28 <i>1</i> <sup>2</sup>	2.7	" 4
Improved Besthorne...	Germany.	61.8	32 <i>1</i> <sup>2</sup>	1.9	" 9

It will be observed that all but one of these best varieties are foreign, although 21 varieties of Canadian oats were grown, including nearly all the old standard varieties. The average of yield of the ten varieties given in the table was 64.1 bushels per acre, while the

average yield of the ten poorest was 27.8 bushels, that is, the ten good varieties yielded 230 per cent. more than the ten poor ones. It is also worthy of note that these good varieties all possessed the spreading head. The tendency to rust is slight in the first, third, fourth, fifth, seventh and ninth, considerable in the tenth and slight to medium in the rest. The color of grain in the first and second is pale black, the third and seventh black, the tenth yellow, and in the others white.

The Goanette Black, Houdan Black, Chenailles and Black Etampes are very much alike in all essential characteristics. The straw is strong and of medium height, or a little less. It stands exceptionally well, is not coarse and is almost entirely free from rust. They are somewhat inclined to shell easily when ripe, as is often the case with valuable grains. They also yield well and weigh well, and have a small percentage of hull as compared with the kernel.

**AVERAGE YIELDS OF BARLEYS.**—The average yields of barleys grown from seed obtained in the five countries named are as follows:

Country.	No. of varieties.	Bush. per acre.
Germany .....	17	35.6
England .....	18	42.2
France .....	7	37.6
Ontario .....	4	38.2
Sweden .....	4	33.5

**AVERAGE YIELDS OF OATS.**—The average yields of oats grown from seed obtained in the seven countries named in the table are as follows:

Country.	No. of varieties.	Bush. per acre.
Ontario .....	21	44.8
France .....	19	47.8
Germany .....	15	38.0
Scotland .....	13	41.8
England .....	9	41.8
Russia .....	5	42.7
Australia .....	4	32.6

**TWO-ROWED AND SIX-ROWED BARLEYS.**—The next table gives information regarding the 54 varieties of barley grown with and

without hulls as to yield, weight of grain per bushel and the size of the grains.

Character of head.	With or without hulls.	No. of varieties.	Bush. per acre.	Lb. per bush.	No. of grains per oz.
Six-rowed {	With hulls ...	9	37.45	51.22	772
	Without hulls	5	31.00	61.80	848
Two-rowed {	With hulls ...	39	37.84	51.87	685
	Without hulls	1	21.90	63.00	478

It should be borne in mind (1) that the average yields obtained in growing these grains is more than the average is likely to prove in the country generally, for they were all grown under very favorable conditions; (2) that similar weights are not likely to be realised unless the grains are thoroughly cleaned; and (3) that the facts in this bulletin will be valuable mainly in localities with soil and climatic conditions similar to those here. The soil where these plots were grown may be termed a mild clay loam, containing a considerable quantity of humus, and it is somewhat low in aspect. The average mean temperature for the past four years for the five months commencing with May 1st has been 61.43° F.

#### CONCLUSIONS.

The following conclusions may safely be drawn from the experiments which have been summarised above:

1. It is highly probable that some of the foreign varieties of barley named in the table will be found to give higher yields than the varieties now grown in Ontario when they have become more generally introduced.
2. It is probable that in some localities the Herison Bearded spring wheat will be found to be an improvement on many of the varieties that are now grown.
3. From present indications, based upon the trial given in the table and upon a trial on a larger scale on the farm, the Mummy pea is likely to become a generally useful variety, although the straw is not highly valued for feeding purposes owing to its coarse habit of growth.
4. That four varieties of French oats, viz., the Goanette Black, the Houdan Black, the Chenailles and the Black Etampes, are likely to prove of much value to the farmers of this province, as they

possess in common and in a marked degree nearly all the most valuable characteristics of oats. Their color will however discount them in the estimation of the oatmeal millers. The behavior of the Oderbrucher, also a white variety from Germany, has been such as to entitle it to the favorable consideration of the farmers. It has been most favorably recommended by the oatmeal millers.

5. Judging from the experience of the past two years the English barleys give on the whole the best results, but some fine growing and yielding varieties come from Germany and France. In reference to oats, the French varieties should be placed first, all things considered, although some kinds from Germany do nearly as well.

6. The average yields obtained from the two-rowed and six-rowed varieties of barley are not far different, nor is there much difference in the average weights of the two classes.

7. The average returns from the foreign varieties are in a majority of instances superior to those from the old standard varieties.

---

valu-  
ount  
r of  
such  
It

glish  
wing  
ence  
con-

owed  
ence

ritiy

