




**IMAGE EVALUATION  
TEST TARGET (MT-3)**



**Photographic  
Sciences  
Corporation**

23 WEST MAIN STREET  
WEBSTER, N.Y. 14580  
(716) 872-4503

**CIHM/ICMH  
Microfiche  
Series.**

**CIHM/ICMH  
Collection de  
microfiches.**



Canadian Institute for Historical Microreproductions / Institut canadien de microreproductions historiques

**© 1985**

Technical and Bibliographic Notes/Notes techniques et bibliographiques

The Institute has attempted to obtain the best original copy available for filming. Features of this copy which may be bibliographically unique, which may alter any of the images in the reproduction, or which may significantly change the usual method of filming, are checked below.

L'Institut a microfilmé le meilleur exemplaire qu'il lui a été possible de se procurer. Les détails de cet exemplaire qui sont peut-être uniques du point de vue bibliographique, qui peuvent modifier une image reproduite, ou qui peuvent exiger une modification dans la méthode normale de filmage sont indiqués ci-dessous.

- Coloured covers/  
Couverture de couleur
- Covers damaged/  
Couverture endommagée
- Covers restored and/or laminated/  
Couverture restaurée et/ou pelliculée
- Cover title missing/  
Le titre de couverture manque
- Coloured maps/  
Cartes géographiques en couleur
- Coloured ink (i.e. other than blue or black)/  
Encre de couleur (i.e. autre que bleue ou noire)
- Coloured plates and/or illustrations/  
Planches et/ou illustrations en couleur
- Bound with other material/  
Relié avec d'autres documents
- Tight binding may cause shadows or distortion along interior margin/  
La reliure serrée peut causer de l'ombre ou de la distorsion le long de la marge intérieure
- Blank leaves added during restoration may appear within the text. Whenever possible, these have been omitted from filming/  
Il se peut que certaines pages blanches ajoutées lors d'une restauration apparaissent dans le texte, mais, lorsque cela était possible, ces pages n'ont pas été filmées.
- Additional comments: / Various pagings.  
Commentaires supplémentaires:

- Coloured pages/  
Pages de couleur
- Pages damaged/  
Pages endommagées
- Pages restored and/or laminated/  
Pages restaurées et/ou pelliculées
- Pages discoloured, stained or foxed/  
Pages décolorées, tachetées ou piquées
- Pages detached/  
Pages détachées
- Showthrough/  
Transparence
- Quality of print varies/  
Qualité inégale de l'impression
- Includes supplementary material/  
Comprend du matériel supplémentaire
- Only edition available/  
Seule édition disponible
- Pages wholly or partially obscured by errata slips, tissues, etc., have been refilmed to ensure the best possible image/  
Les pages totalement ou partiellement obscurcies par un feuillet d'errata, une pelure, etc., ont été filmées à nouveau de façon à obtenir la meilleure image possible.

This item is filmed at the reduction ratio checked below/  
Ce document est filmé au taux de réduction indiqué ci-dessous.

10X	14X	18X	22X	26X	30X
<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
12X	16X	20X	24X	28X	32X

The copy filmed here has been reproduced thanks to the generosity of:




D. B. Weldon Library  
University of Western Ontario  
(Regional History Room)

The images appearing here are the best quality possible considering the condition and legibility of the original copy and in keeping with the filming contract specifications.

Original copies in printed paper covers are filmed beginning with the front cover and ending on the last page with a printed or illustrated impression, or the back cover when appropriate. All other original copies are filmed beginning on the first page with a printed or illustrated impression, and ending on the last page with a printed or illustrated impression.

The last recorded frame on each microfiche shall contain the symbol  $\rightarrow$  (meaning "CONTINUED"), or the symbol  $\nabla$  (meaning "END"), whichever applies.

Maps, plates, charts, etc., may be filmed at different reduction ratios. Those too large to be entirely included in one exposure are filmed beginning in the upper left hand corner, left to right and top to bottom, as many frames as required. The following diagrams illustrate the method:



L'exemplaire filmé fut reproduit grâce à la générosité de:


D. B. Weldon Library  
University of Western Ontario  
(Regional History Room)

Les images suivantes ont été reproduites avec le plus grand soin, compte tenu de la condition et de la netteté de l'exemplaire filmé, et en conformité avec les conditions du contrat de filmage.

Les exemplaires originaux dont la couverture en papier est imprimée sont filmés en commençant par le premier plat et en terminant soit par la dernière page qui comporte une empreinte d'impression ou d'illustration, soit par le second plat, selon le cas. Tous les autres exemplaires originaux sont filmés en commençant par la première page qui comporte une empreinte d'impression ou d'illustration et en terminant par la dernière page qui comporte une telle empreinte.

Un des symboles suivants apparaîtra sur la dernière image de chaque microfiche, selon le cas: le symbole  $\rightarrow$  signifie "A SUIVRE", le symbole  $\nabla$  signifie "FIN".

Les cartes, planches, tableaux, etc., peuvent être filmés à des taux de réduction différents. Lorsque le document est trop grand pour être reproduit en un seul cliché, il est filmé à partir de l'angle supérieur gauche, de gauche à droite, et de haut en bas, en prenant le nombre d'images nécessaire. Les diagrammes suivants illustrent la méthode.



W. J. Gage's Mathematical Series.

---

THE TEACHER'S  
HAND-BOOK OF ALGEBRA ;

CONTAINING

METHODS, SOLUTIONS AND EXERCISES

ILLUSTRATING

THE LATEST AND BEST TREATMENT OF THE ELEMENTS  
OF ALGEBRA.

BY

J. A. McLELLAN, M.A., LL.D.,

HIGH SCHOOL INSPECTOR FOR ONTARIO.

*The object of pure Mathematics, which is another name for Algebra, is the unfolding  
of the laws of the human intelligence.*"—SYLVESTER.

---

FOURTH EDITION—REVISED AND ENLARGED.

---

TORONTO:

W. J. GAGE & COMPANY.

1881.

---

Entered according to Act of Parliament of Canada in the year  
1880 by W. J. GAGE & COMPANY, in the office of the Minister  
of Agriculture.

---

T 713

A  
of  
to  
fo  
  
m  
no  
  
so  
m  
ele  
  
the  
eff  
ad  
  
qu  
he  
  
top  
ord  
Div  
  
A  
pos  
eve  
ent  
exp

## PREFACE.

---

This book—embodying the substance of Lectures at Teachers' Associations—has been prepared at the almost unanimous request of the teachers of Ontario, who have long felt the need of a work to supplement the elementary text-books in common use. The following are some of its special features :

It gives a large number of solutions in illustration of the best methods of algebraic resolution and reduction, some of which are not found in any text-book.

It gives, classified under proper heads and preceded by type-solutions, a great number of exercises, many of them illustrating methods and principles which are unaccountably ignored in elementary Algebras.

It presents these solutions and exercises in such a way that the student not only sees how Algebraic transformations are effected, but also perceives how to form for himself as many additional examples as he may desire.

It shows the student how simple principles with which he is quite familiar, may be applied to the solution of questions which he has thought beyond their reach.

It gives complete explanations and illustrations of important topics which are strangely omitted or barely touched upon in the ordinary books, such as the Principle of Symmetry, Theory of Divisors, Factoring, Applications of Horner's Division, &c.

A few of the exercises are chiefly supplementary to those proposed in the text-books, but the intelligent student will find that even these examples have not been selected in the usual apparently aimless fashion ; he will recognise that they are really expressions of certain laws ; they are in fact proposed with a view



to lead him to investigate these laws for himself as soon as he has sufficiently advanced in his course. Nos. 8, 9, 10 and 11 afford instances of such exercises.

Others of the questions proposed are preparatory or interpretation exercises. These might well have been omitted, were it not that they are generally omitted from the text-books and too often neglected by teachers. Practice in the interpretation of a new notation and in expression by means of it, should always precede its use as a symbolism itself subject to operations. Nos. 23 to 36 of Ex. iii., and nearly the whole of Ex. xv. may serve for instances.

By far the greater number of the exercises are intended for practice in the methods exhibited in the solved examples. As many as possible of these have been selected for their intrinsic value. They have been gathered from the works of the great masters of analysis, and the student who proceeds to the higher branches of mathematics will meet again with these examples and exercises, and he will find his progress aided by his familiarity with them, and will not have to interrupt his advanced studies to learn processes properly belonging to elementary Algebra. In making this selection, it has been found that the most widely useful transformations are, at the same time, those that best exhibit the methods of reduction here explained, so that they have thus a double advantage. A great part of the exercises have, of necessity, been prepared specially for this work.

Articles and exercises have been prepared on the theory of substitutions, on Elimination, &c., but it has finally been decided to hold these over for Pt. ii., which will probably appear if the present work be favorably received.

SECT  
SECT  
SECT

SECT  
SECT

SECT  
SECT  
SECT  
SECT  
SECT

SECT  
SECT  
SECT  
SECT

SIMP  
P

SIMP  
E

EXAM

# CONTENTS.

	PAGE.
CHAPTER I.—SUBSTITUTION, HORNER'S DIVISION, &c.	
SECT. 1.—Numerical and Literal Substitution.....	1
SECT. 2.—Fundamental Formulas and their Applications.....	10
SECT. 3.—Horner's Methods of Multiplication and Division, and their Applications .....	21
CHAPTER II.—PRINCIPLE OF SYMMETRY, &c.	
SECT. 1.—The Principle of Symmetry and its Applications .....	33
SECT. 2.—The Theory of Divisors and its Applications.....	39
CHAPTER III.—FACTORING.	
SECT. 1.—Direct Application of the Fundamental Formulas .....	62
SECT. 2.—Extended Application of the Formulas .....	71
SECT. 3.—Factoring by Parts .....	79
SECT. 4.—Application of the Theory of Divisors.....	83
SECT. 5.—Factoring a Polynome by Trial Divisors.....	90
CHAPTER IV.—MEASURES AND MULTIPLES, &c.	
SECT. 1.—Division, Measures and Multiples .....	101
SECT. 2.—Fractions .....	109
SECT. 3.—Ratios .....	122
SECT. 4.—Complete Squares, &c.....	130
CHAPTER V.	
SIMPLE EQUATIONS OF ONE UNKNOWN QUANTITY.....	139
Preliminary Equations. Resolution by Factors. Fractional Equations. Application of Ratios. Equations involving Surds, Higher Equations, &c.	
CHAPTER VI.	
SIMULTANEOUS EQUATIONS.....	170
Equations of Two Unknown Quantities. Systems of Equations. Application of Symmetry. Equations of Three Unknowns. Systems of Equations.	
CHAPTER VII.	
EXAMINATION PAPERS.....	207

1  
val

2.  
valu

2(s)  
3.  
follow

a  
a  
a  
(  
4  
a

## CHAPTER I.

### SECTION I.—SUBSTITUTION.

#### Exercise i.

1. If  $a = 1$ ,  $b = 2$ ,  $c = 3$ ,  $d = 4$ ,  $x = 9$ ,  $y = 8$ , find the value of the following expressions :—

$$\begin{aligned}
 & 1 - \{1 - (1 - \overline{1-x})\}. \\
 & a - (x-y) - (b-c)(d-a) - (y-b)(x+c). \\
 & x-y[y - (y-a)\{d+c(b-c)\}]. \\
 & (x+d)(y+b+c) + (x-d)(a-b-d) + (y+d)(a-x-d). \\
 & (d-x)^3 + (c+y)^3. \\
 & (a-b)(c^3 - b^2x) - (c-d)(b^3 - a^2x) + (d-b-c)(d^3 - c^3). \\
 & \frac{d-a}{d+a} + \frac{d+c}{d-c} - \frac{2d+b}{d-b}.
 \end{aligned}$$

2. If  $a = 3$ ,  $b = -4$ ,  $c = -9$ , and  $2s = a + b + c$ , find the value of the following expressions :—

$$\begin{aligned}
 & s(s-a)(s-b)(s-c). \\
 & s^2 + (s-a)^2 + (s-b)^2 + (s-c)^2. \\
 & s^2 - (s-a)(s-b) - (s-b)(s-c) - (s-c)(s-a). \\
 & 2(s-a)(s-b)(s-c) + a(s-b)(s-c) + b(s-c)(s-a) + c(s-a)(s-b).
 \end{aligned}$$

3. If  $a = 2$ ,  $b = -3$ ,  $c = 1$ ,  $x = 4\frac{1}{2}$ , find the value of the following expressions :—

$$\begin{aligned}
 & \frac{a^2 - b^2}{a^3 + b^3}, \quad \frac{a^2 + b^2}{a^3 - b^3}, \quad \frac{(a-b)^2}{(a+b)^3}, \quad \frac{(a-b)^3}{(a+b)^2}, \\
 & \frac{a^2 + ab + b^2}{a^2 - ab + b^2}, \quad \frac{a^2 - b^3}{a^3 - b^2}, \quad \frac{x}{2} \left\{ \frac{2x-3}{3} - \frac{3x-1}{4} \right\} \frac{x-1}{2}, \\
 & \frac{(a+b)\{(a+b)^2 - c^2\}}{4b^2c^2 - (a^2 - b^2 - c^2)^2}, \\
 & \frac{a^2(b-c) + b^2(c-a) + c^2(a-b)}{(a-b)(b-c)(c-a)}.
 \end{aligned}$$

4. If  $a = 6$ ,  $b = 5$ ,  $c = -4$ ,  $d = -3$ , find the value of the following expressions:—

$$\sqrt{(b^2 + ac)} + \sqrt{(c^2 - 2ac)}. \quad \sqrt{\{b^2 + ac + \sqrt{(c^2 - 2ac)}\}}.$$

$$\frac{a^2 - \sqrt{(b^2 + ac)}}{2a - \sqrt{(b^2 - ac)}}, \quad \frac{c + \sqrt{(d^2 + c^2)}}{c^3 + 2d(d^2 - c^2)}.$$

5. If  $x = 3$ ,  $y = 4$ ,  $z = 0$ , find the value of:—

$$\{3x - \sqrt{(x^2 + y^2)}\}^2 \{2x + \sqrt{(x^2 + y^2 + z)}\}.$$

$$x^y + y^z + z^x, \quad (x^y)^{x-y} + (y-z)^{y-z} + (z-x)^{z-x}.$$

$$(x^3 - y^3) \div \sqrt[3]{\{3x^3 + 3(3x^2 + 3xy + y^2)y\}}.$$

6. Calculate the values of  $\frac{(x+y+z)^3 - 3(x^3 + y^3 + z^3)}{xyz}$  when

- (a)  $x = 1$ ,  $y = 2$ ,  $z = 3$ .  
 (b)  $x = 2$ ,  $y = 3$ ,  $z = 4$ .  
 (c)  $x = 3$ ,  $y = 4$ ,  $z = 5$ .  
 (d)  $x = 10$ ,  $y = 11$ ,  $z = 12$ .

7. Given  $x = 3$ ,  $y = 4$ ,  $z = -5$ , calculate the values of

$$(x+y+x)^3 - 3(x+y+z)(xy+yz+zx).$$

$$x^2(y+z) + y^2(z+x) + z^2(x+y) + 2xyz.$$

$$x^2(y-z) + y^2(z-x) + z^2(x-y).$$

$$(5x-4z)^2 + 9(4x-z)^2 - (13x-5z)^2.$$

$$(3x+4y+5z)^2 + (4x+3y+12z)^2 - (5x+5y+13z)^2.$$

8. If  $s = a + b + c$ , find the value of

$$(2s-a)^2 + (2s-b)^2 - (2s+c)^2, \text{ given}$$

- (1)  $a = 3$ ,  $b = 4$ ,  $c = 5$ ,                      (2)  $a = 21$ ,  $b = 20$ ,  $c = 29$ ,  
 (3)  $a = 119$ ,  $b = 120$ ,  $c = 169$ ,            (4)  $a = 3$ ,  $b = -4$ ,  $c = 5$ ,  
 (5)  $a = 5$ ,  $b = 12$ ,  $c = -13$ .

9. If  $a = 1$ ,  $b = 3$ ,  $c = 5$ ,  $d = 7$ ,  $e = 9$ ,  $f = 11$ , prove that

$$a+b+c+d+e+f = \left(\frac{a+f}{2}\right)^2.$$

$$\frac{1}{ab} + \frac{1}{bc} + \frac{1}{cd} + \frac{1}{de} + \frac{1}{ef} = \frac{1}{2} \left(\frac{1}{a} - \frac{1}{f}\right).$$

$$\frac{1}{abc} + \frac{1}{bcd} + \frac{1}{cde} + \frac{1}{def} = \frac{1}{4} \left(\frac{1}{ab} - \frac{1}{ef}\right).$$

$$\frac{1}{abcd} + \frac{1}{bcde} + \frac{1}{cdef} = \frac{1}{6} \left(\frac{1}{abc} - \frac{1}{def}\right).$$

$$a^2 + b^2 + c^2 - ab - bc - ca = b^2 + c^2 + d^2 - bc - cd - db = c^2 + d^2 + e^2 - cd - de - ec = d^2 + e^2 + f^2 - de - ef - fd.$$

10. If  $a = 1, b = 2, c = 3, d = 4, e = 5, f = 6, g = 7$ , prove that

$$a + b + c = \frac{1}{2}cd, \quad a + b + c + d = \frac{1}{2}de,$$

$$a + b + c + d + e = \frac{1}{2}ef, \quad a + b + c + d + e + f = \frac{1}{2}fg,$$

$$a^2 + b^2 + c^2 = \frac{ed(e+d)}{ab(a+b)}, \quad a^2 + b^2 + c^2 + d^2 =$$

$$\frac{de(d+e)}{ab(a+b)}, \quad a^2 + b^2 + c^2 + d^2 + e^2 = \frac{ef(e+f)}{ab(a+b)},$$

$$a^2 + b^2 + c^2 + d^2 + e^2 + f^2 = \frac{fg(f+g)}{ab(a+b)}.$$

$$a^3 + b^3 + c^3 = (a + b + c)^2,$$

$$a^3 + b^3 + c^3 + d^3 = (a + b + c + d)^2,$$

$$a^3 + b^3 + c^3 + d^3 + e^3 = (a + b + c + d + e)^2,$$

$$a^3 + b^3 + c^3 + d^3 + e^3 + f^3 = (a + b + c + d + e + f)^2$$

$$a^4 + b^4 + c^4 = \frac{cd(c+d)(c^2d-1)}{bc(b+c)},$$

$$a^4 + b^4 + c^4 + d^4 = \frac{de(d+e)(cde-1)}{bc(b+c)},$$

$$a^4 + b^4 + c^4 + d^4 + e^4 = \frac{ef(e+f)(cef-1)}{bc(b+c)},$$

$$a^4 + b^4 + c^4 + d^4 + e^4 + f^4 = \frac{fg(f+g)(efg-1)}{bc(b+c)},$$

$$c^2 + d^2 = e^2, \quad c^3 + d^3 + e^3 = f^3$$

11. Assume any numerical values for  $x, y$ , and  $z$ , and calculate the values of the following expressions:—

$$(x^5 - 10x^3 + 5x)^2 + (5x^4 - 10x^2 + 1)^2 - (x^2 + 1)^5.$$

$$(x+1)^3 - 2(x+5)^3 - (x+9)^3 + 2(x+11)^3 + (x+12)^3 - (x+16)^3.$$

$$(x^2 - y^2)^2 + (2xy)^2 - (x^2 + y^2)^2$$

$$(x^3 - 3xy^2)^2 + (3x^2y - y^3)^2 - (x^2 + y^2)^3.$$

$$(3x^2 + 4xy + y^2)^2 + (4x^2 + 2xy)^2 - (5x^2 + 4xy + y^2)^2.$$

$$(x-y)^3 + (y-z)^3 + (z-x)^3 - 3(x-y)(y-z)(z-x).$$

Art. I. If  $x =$  any number, as, for example, 3, then  $x^2$  (which =  $x.x$ ) =  $3x$ ,  $x^3$  (which =  $x.x^2$ ) =  $3x^2$ ,  $x^4$  (which =  $x.x^3$ ) =  $3x^3$ , &c. Or  $3 = x, 3x = x^2, 3x^2 = x^3, 3x^3 = x^4$ , &c. Hence prob.

blems like the following may be solved like ordinary arithmetical problems in "Reduction Descending."

## EXAMPLES.

1. Find the value of  $x^2 - 2x - 9$  when  $x = 5$ .

$$\begin{array}{r}
 x^2 - 2x - 9 \\
 5 \\
 \hline
 5x \\
 - 2x \\
 \hline
 3x \\
 5 \\
 \hline
 15 \\
 - 9 \\
 \hline
 6
 \end{array}$$

*Explanation.*

$$x^2 = 5x,$$

$$\therefore x^2 - 2x = 3x = 15, \text{ and}$$

$$\therefore x^2 - 2x - 9 = 15 - 9 = 6.$$

2. Find the value of  $x^4 - x^3 - 4x^2 - 3x - 5$  when  $x = 3$ .

$$\begin{array}{r}
 x^4 - x^3 - 4x^2 - 3x - 5 \\
 3 \\
 \hline
 p_1 \dots\dots\dots 3x^3 \\
 \quad \quad \quad - x^3 \\
 \hline
 r_1 \dots\dots\dots 2x^3 \\
 \quad \quad \quad 3 \\
 \hline
 p_2 \dots\dots\dots 6x^2 \\
 \quad \quad \quad - 4x^2 \\
 \hline
 r_2 \dots\dots\dots 2x^2 \\
 \quad \quad \quad 3 \\
 \hline
 r_3 \dots\dots\dots 6x \\
 \quad \quad \quad - 3x \\
 \hline
 r_3 \dots\dots\dots 3x \\
 \quad \quad \quad 3 \\
 \hline
 p_4 \dots\dots\dots 9 \\
 \quad \quad \quad - 5 \\
 \hline
 r_4 \dots\dots\dots 4.
 \end{array}$$

$$\therefore x^4 - x^3 - 4x^2 - 3x - 5 = 4 \text{ if } x = 3.$$

ne  
w  
ti  
an  
re

*Explanation.*

$$\begin{aligned} x^4 &= 3x^3, \\ \therefore x^4 - x^3 &= 2x^3 = 6x^2, \\ \therefore x^4 - x^3 - 4x^2 &= 2x^2 = 6x, \\ \therefore x^4 - x^3 - 4x^2 - 3x &= 3x = 9, \\ \therefore x^4 - x^3 - 4x^2 - 3x - 5 &= 4. \end{aligned}$$

8. Find the value of  $2x^4 + 12x^3 + 6x^2 - 12x + 10$ .  
Using coefficients only, we have

$$\begin{array}{r} 2+12+6-12+10 \\ \hline -5 \\ \hline p_1 \dots\dots -10 \\ \quad \quad \quad +12 \\ \hline r_1 \dots\dots +2 \\ \quad \quad \quad -5 \\ \hline p_2 \dots\dots -10 \\ \quad \quad \quad +6 \\ \hline r_2 \dots\dots -4 \\ \quad \quad \quad -5 \\ \hline p_3 \dots\dots 20 \\ \quad \quad \quad -12 \\ \hline r_3 \dots\dots 8 \\ \quad \quad \quad -5 \\ \hline p_4 \dots\dots -40 \\ \quad \quad \quad +10 \\ \hline r_4 \dots\dots -30 \end{array}$$

$\therefore$  the quantity =  $-30$  if  $x = -5$ .

**Art. II.** If the coefficients, and also the values of  $x$  are small numbers, much of the above may be done mentally, and the work will then be very compact. Thus, performing mentally the multiplications and additions (or subtractions) of the coefficients, and merely recording the partial reductions  $r_1, r_2, r_3$ , and the result  $r_4$ , the last example would appear as follows:—



$$\begin{array}{r}
 -5 \ ) \ 2 \ +12 \ +6 \ -12 \ +10 \\
 \underline{2} \\
 -4 \\
 \underline{8} \\
 -30
 \end{array}$$

**Art. III.** In the above examples, the coefficients are "brought down" and written below the products  $p_1, p_2, p_3, p_4$ , and are added or subtracted, as the case may require, to get the partial reductions  $r_1, r_2, r_3$ , and the result  $r_4$ . Instead of thus "bringing down" the coefficients, we may "carry up" the products  $p_1, p_2, p_3, p_4$ , writing them beneath their corresponding coefficients, and thus get  $r_1, r_2, r_3, r_4$  in a third (horizontal) line. Arranged in this way Ex. 2 will appear

$$\begin{array}{r|rrrrr}
 3 & 1 & -1 & -4 & -3 & -5 \\
 & & +3 & +6 & +6 & +9 \\
 \hline
 & 1 & +2 & +2 & +3 & 4;
 \end{array}$$

and Ex. 3 will appear

$$\begin{array}{r|rrrrr}
 -5 & 2 & +12 & +6 & -12 & +10 \\
 & & -10 & -10 & +20 & -40 \\
 \hline
 & 2 & +2 & -4 & +8 & -30
 \end{array}$$

Comparing these arrangements with those first given (Ex. 2 and 3), it will be seen that they are figure for figure the same, except that the multiplier is not repeated.

**Art. IV.** When there are several figures in the value of  $x$ , they may be arranged in a column, and each figure used separately, as in common multiplication. Where only approximate values are required, "contracted multiplication" may be used.

4. Find the value of  $3x^5 - 160x^4 + 344x^3 + 700x^2 - 1910x + 1200$ , given  $x = 51$ .

$$\begin{array}{r|rrrrrr}
 1 & 3 & -160 & +344 & +700 & -1910 & +1200 \\
 50 & & 3 & -7 & -13 & 37 & -23 \\
 & & 150 & -350 & -650 & 1850 & -1150 \\
 \hline
 & 3 & -7 & -13 & +37 & -23; & +27
 \end{array}$$

$\therefore$  result is 27.

5. Given  $x = 1.183$ , find the value of  $64x^4 - 144x + 45$  correct to three decimal places.

	64	0	0	-144	+45
1	64	75.712	89.5673	-38.0419	
1	6.4	7.5712	8.9567	-3.8042	
8	5.12	6.0570	7.1654	-3.0434	
8	.192	.2271	.2687	-.1141	
<hr/>					
	64,	75.712,	89.5673,	-38.0419,	-.0036.

∴ result is  $-.004$ .

Exercise ii.

Find the value of

- $x^4 - 11x^3 - 11x^2 - 13x + 11$ , for  $x = 12$ .
- $x^4 + 50x^3 - 16x^2 - 16x - 61$ , for  $x = -17$ .
- $2x^4 + 249x^3 - 125x^2 + 100$ , for  $x = -125$ .
- $2x^3 - 473x^2 - 234x - 711$ , for  $x = 200$ .
- $x^5 - 3x^2 - 8$ , for  $x = 4$ .
- $x^6 - 515x^5 - 3127x^4 + 525x^3 - 2090x^2 + 3156x - 15792$ , for  $x = 521$ .
- $2x^5 + 401x^4 - 199x^3 + 399x^2 - 602x + 211$ , for  $x = -201$ .
- $1000x^4 - 81x$ , for  $x = .1$ .
- $99x^4 + 117x^3 - 257x^2 - 325x - 50$ , for  $x = 1\frac{2}{3}$ .
- $5x^5 + 497x^4 + 200x^3 + 196x^2 - 218x - 2000$ , for  $x = -99$ .
- $5x^5 - 620x^4 - 1030x^3 + 1045x^2 - 4120x + 9000$ , for  $x = 205$ .

Calculate, correct to three places of decimals,—

- $x^3 + 3x^2 - 13x - 38$  for  $x = 3.58443$ , for  $x = -3.77931$ , and for  $x = -2.80512$ .
- $y^4 - 14y^2 + y + 38$  for  $y = 3.13131$ , for  $y = -1.84813$ , and for  $y = -3.28319$ .

Exercise iii.

What do the following expressions become (1) when  $x = a$ , (2) when  $x = -a$ ?

- $x^4 - 4ax^3 + 6a^2x^2 - 4a^3x + a^4$ .
- $\sqrt{(x^2 - ax + a^2)}$ .
- $\sqrt{(x^2 + 2ax + a^2)}$ .
- $(x^2 + ax + a^2)^3 - (x^2 - ax + a^2)^3$ .

If  $x = y = z = a$ , find the value of the following expressions:

5.  $(x-y)(y-z)(z-x)$ .

6.  $(x+y)^2(y+z-a)(x+z-a)$ .

7.  $x(y+z)(y^2+z^2-x^2)+y(z+x)(z^2+x^2-y^2)+z(x+y)(x^2+y^2-z^2)$ .

8.  $\frac{x}{y+z} + \frac{y}{x+z} + \frac{z}{x+y}$ .

Find the value of

9.  $\frac{x}{a} + \frac{x}{b}$  when  $x = \frac{abc}{a+b}$ .

10.  $\frac{1}{a(b-x)} + \frac{1}{b(c-x)} + \frac{1}{a(x-c)}$ , when  $x = \frac{b}{a}(a-b+c)$ .

11.  $\frac{x}{a} + \frac{x}{b-a}$ , when  $x = \frac{a^2(b-a)}{b(b+a)}$ .

12.  $(a+x)(b+x) - a(b+c) + x^2$ , when  $x = \frac{ac}{b}$ .

13.  $bx+cy+az$ , when  $x=b+c-a$ ,  $y=c+a-b$ ,  $z=a+b-c$ .

14.  $\frac{a(1+b)+bx}{a(1+b)-bx} - \frac{a}{a-2bx}$ , when  $x=-a$ .

15.  $\left(\frac{x+a}{x+b}\right)^3 - \frac{x+2a+b}{x-a-2b}$ , when  $x = \frac{1}{2}(b-a)$ .

16.  $(p-q)(x+2r) + (r-x)(p+q)$ , when  $x = \frac{r(3p-q)}{2q}$ .

17.  $a^2(b-c)+b^2(c-a)+c^2(a-b)$ , when  $a-b=0$ .

18.  $(a+b+c)(bc+ca+ab) - (a+b)(b+c)(c+a)$ , when  $a=-b$ .

19.  $(a+b+c)^3 - (a^3+b^3+c^3)$ , when  $a+b=0$ .

20.  $(x+y+z)^4 - (x+y)^4 - (y+z)^4 - (z+x)^4 + x^4 + y^4 + z^4$ , when  $x+y+z=0$ .

21.  $a^3(c-b^2)+b^3(a-c^2)+c^3(b-a^2)+abc(abc-1)$ , when  $b-a^2=0$ .

22.  $a^5 \left(\frac{a^5+5b^5}{a^5-b^5}\right)^5 + b^5 \left(\frac{5a^5+b^5}{b^5-a^5}\right)^5$ , when  $a^5+b^5=0$ .

23. Express in words the fact that

$$(a-b)^2 = a^2 - 2ab + b^2.$$

24. Express algebraically the fact "that the sum of two quantities multiplied by their difference is equal to the difference of the squares of the numbers."

25. The area of the walls of a room is equal to the height multiplied by twice the sum of the length and breadth: what are the areas of the walls in the following cases:

(1) length  $l$ , height  $h$ , breadth  $b$ .

(2) height  $x$ , length  $b$  feet more than the height, and breadth  $b$  feet less than the height.

26. Express in words the statement that

$$(x+a)(x+b) = x^2 + (a+b)x + ab.$$

27. Express in symbols the statement that "the square of the sum of two quantities exceeds the sum of their squares by twice their product."

28. Express in words the algebraic statement.

$$(x+y)^3 = x^3 + y^3 + 3xy(x+y).$$

29. Express algebraically the fact that "the cube of the difference of two quantities is equal to the difference of the cubes of the quantities diminished by three times the product of the quantities multiplied by their difference."

30. If the sum of the cubes of two quantities be divided by the sum of the quantities, the quotient is equal to the square of their difference increased by their product; express this algebraically.

31. Express in words the following algebraic statement:

$$\frac{x^3 - y^3}{x - y} = (x+y)^2 - xy.$$

32. The square on the diagonal of a cube is equal to three times the square on the edge; express this in symbols, using  $l$  for length of the edge, and  $d$  for length of the diagonal.

33. Express in symbols that "the length of the edge of the greatest cube that can be cut from a sphere is equal to the square root of one-third the square of the diameter."

34. Express in symbols that any "rectangle is half the rectangle contained by the diagonals of the squares upon two adjacent sides." [The square on the diagonal of a square is double the square on a side.]

35. The area of a circle is equal to  $\pi$  multiplied into the square

of the radius; express this in symbols. Also express in symbols the area of the ring between two concentric circles.

36. The volume of a cylinder is equal to product of its height into the area of the base, that of a cone is one-third of this, and that of a sphere is two-thirds of the volume of the circumscribing cylinder; express these facts in symbols, using  $h$  for the height of the cylinder, and  $r$  for the radius of its base.

Exercise iv.

Perform the additions in the following cases :

1.  $(b - a)x + (c - b)y$ , and  $(a + b)x + (b + c)y$ .
2.  $ax - by$ ,  $(a - b)x - (a + b)y$ , and  $(a + b)x - (b - a)y$ .
3.  $(y - z)a^2 + (z - x)ab + (x - y)b^2$ , and  $(x - y)a^2 - (z - y)ab - (x - z)b^2$ .
4.  $ax + by + cz$ ,  $bx + cy + az$ , and  $cx + ay + bz$ .
5.  $(a + b)x^2 + (b + c)y^2 + (a + c)z^2$ ,  $(b + c)x^2 + (a + c)y^2 + (a + b)z^2$ ,  $(a + c)x^2 + (a + b)y^2 + (b + c)z^2$ , and  $-(a + b + c)(x^2 + y^2 + z^2)$ .
6.  $x(a - b)^2 + y(b - c)^2 + z(c - a)^2$ ,  $y(a - b)^2 + z(b - c)^2 + x(c - a)^2$ , and  $z(a - b)^2 + x(b - c)^2 + y(c - a)^2$ .
7.  $(a - b)x^2 + (b - c)y^2 + (c - a)z^2$ ,  $(b - c)x^2 + (c - a)y^2 + (a - b)z^2$ , and  $(c - a)x^2 + (a - b)y^2 + (b - c)z^2$ .
8.  $(a + b)x + (b + c)y - (c + a)z$ ,  $(b + c)z + (c + a)x - (a + b)y$ , and  $(a + c)y + (a + b)z - (b + c)x$ .
9.  $a^2 - 3ab - \frac{1}{2}b^2$ ,  $2b^2 - \frac{2}{3}b^3 + c^2$ ,  $ab - \frac{1}{3}b^2 + b^3$ , and  $2ab - \frac{1}{3}b^3$ .
10.  $ax^n - 3bx^n$ ,  $-9ax^n + 7bx^n$ , and  $-8bx^n + 10ax^n$ .
11. What will  $(ax - by + cz) + (bx + cy - az) - (cx + ay + bz)$  become when  $x - y - z = 1$ ?

SECTION II.—FUNDAMENTAL FORMULAS AND THEIR APPLICATION.

4. By Multiplication we get

$$(x + r)(x + s) = x^2 + (r + s)x + rs \dots \dots \dots A.$$

$$(x + r)(x + s)(x + t) = x^3 + (r + s + t)x^2 + (rs + st + tr)x + rst \dots \dots B.$$

From A we immediately get

$$(x \pm y)^2 = x^2 \pm 2xy + y^2 \dots \dots \dots [1]$$

$$(x + y + z)^2 = x^2 + 2xy + 2xz + y^2 + 2yz + z^2 \quad [2]$$

$$(\Sigma a)^2 = \Sigma a^2 + 2 \Sigma ab \dots \dots \dots [3]$$

$$(x + y)(x - y) = x^2 - y^2 \dots \dots \dots [4]$$

From B we derive

$$(x \pm y)^3 = x^3 \pm 3x^2y + 3xy^2 \pm y^3 \dots \dots \dots [5]$$

$$= x^3 \pm y^3 \pm 3xy(x \pm y) \dots \dots \dots [6]$$

$$(x + y + z)^3 = x^3 + y^3 + z^3 + 3x^2(y + z) + 3y^2(z + x) + 3z^2(x + y) + 6xyz \dots \dots \dots [7]$$

$$= x^3 + y^3 + z^3 + 3(x + y)(y + z)(z + x) \dots \dots \dots [8]$$

$$= x^3 + y^3 + z^3 + 3(x + y + z)(xy + yz + zx) - 3xyz \dots [9]$$

$$(\Sigma a)^3 = \Sigma a^3 + 3\Sigma a^2b + 6\Sigma abc \dots \dots \dots [10]$$

[The symbol  $\Sigma$  means THE SUM OF ALL SUCH TERMS AS]

FORMULA [1].—EXAMPLES.

1. We have at once  $(x + y)^2 + (x - y)^2 = 2(x^2 + y^2)$ , and  $(x + y)^2 - (x - y)^2 = 4xy$ .

2.  $(a + b + c + d)^2 + (a - b - c + d)^2$  may be written  $\{(a + d) + (b + c)\}^2 + \{(a + d) - (b + c)\}^2$ , which (Ex. 1) =  $2\{(a + d)^2 + (b + c)^2\}$ ; similarly

$$(a - b + c - d)^2 + (a + b - c - d)^2 = \{(a - d) - (b - c)\}^2 + \{(a - d) + (b - c)\}^2 = 2\{(a - d)^2 + (b - c)^2\};$$

$$\therefore (a + b + c + d)^2 + (a - b - c + d)^2 + (a - b + c - d)^2 + (a + b - c - d)^2 = 2\{(a + d)^2 + (b + c)^2 + (a - d)^2 + (b - c)^2\} = \text{(again by Ex. 1) } 4(a^2 + b^2 + c^2 + d^2).$$

3. Simplify  $(a + b + c)^2 - 2(a + b + c)c + c^2$ ;

This is the square of a binomial of which the first term is  $(a + b + c)$  and the second  $-c$ ; the given quantity  $\therefore = \{(a + b + c) - c\}^2 = (a$

4. Simplify  $(a+b)^4 - 2(a^2 + b^2)(a+b)^2 + 2(a^4 + b^4)$ .

By Ex. 1.  $2(a^4 + b^4) = (a^2 + b^2)^2 + (a^2 - b^2)^2$ ;  $\therefore$  given quantity  $= (a+b)^4 - 2(a^2 + b^2)(a+b)^2 + (a^2 + b^2)^2 + (a^2 - b^2)^2 = \{(a+b)^2 - (a^2 + b^2)\}^2 + (a^2 - b^2)^2 = a^4 + 2a^2b^2 + b^4 = (a^2 + b^2)^2$ .

### Exercise v.

1.  $(x+3y^2)^2 + (x-3y^2)^2$ ,  $(\frac{1}{3}a^2 + 3b^2)^2 - (\frac{1}{3}a^2 - 3b^2)^2$ .
2. Shew that  $(mx+ny)^2 + (nx-my)^2 = (m^2+n^2)(x^2+y^2)$ .
3. " "  $(mx-ny)^2 - (nx-my)^2 = (m^2-n^2)(x^2-y^2)$ .
4. Simplify  $\{a+3b\}^2 + 2(a+3b)(a-b) + (a-b)^2 \{a-b\}^2$ .
5. "  $(x+3)^2 + (x+4)^2 - (x+5)^2$ , and  $(\frac{1}{2}x^2 - 2y^2)^2 - (\frac{1}{2}y^2 + 2x^2)^2$ .
6. Simplify  $(a+b+c)^2 + (b+c)^2 - 2(b+c)(a+b+c)$ .
7. Shew that  $(ax+by)^2 + (cx+dy)^2 + (ay-bx)^2 + (cy-dx)^2 = (a^2+b^2+c^2+d^2)(x^2+y^2)$ .
8. Simplify  $(x-3y^2)^2 + (3x^2-y)^2 - 2(3x^2-y)(x-3y^2)$ .
9. "  $(x^2+xy-y^2)^2 - (x^2-xy-y^2)^2$ , and  $(1+2x+4x^2)^2 + (1-2x+4x^2)^2$ .
10. If  $a+b = -\frac{3}{4}c$ , shew that  $(2a-b)^2 + (2b-c)^2 + (2c-a)^2 + 2(2a-b)(2b-c) + 2(2b-c)(2c-a) + 2(2c-a)(2a-b) = \frac{1}{16}c^2$ .
11. Simplify  $2(a-b)^2 - (a-2b)^2$ ;  $(a^2+4ab+b^2)^2 - (a^2+b^2)^2$ .
12. "  $(a+b)^2 - (b+c)^2 + (c+d)^2 - (d+a)^2$ .
13. "  $(\frac{1}{2}x-y)^2 + (\frac{1}{2}y-z)^2 + (\frac{1}{2}z-x)^2 + 2(\frac{1}{2}x-y)(\frac{1}{2}z-x) + 2(\frac{1}{2}y-z)(\frac{1}{2}z-x) + 2(\frac{1}{2}x-y)(\frac{1}{2}y-z)$ .
14. Prove that  $(x-y)^2 + (y-z)^2 + (z-x)^2 = 2(x-y)(z-y) + 2(y-x)(z-x) + 2(z-y)(z-x)$ .
15. Simplify  $(1+x)^4 - 2(1+x^2)(1+x)^2 + 2(1+x^4)$ .
16. "  $(x+y+z)^2 - (x+y-z)^2 - (y+z-x)^2 - (z+x-y)^2$ .
17. "  $(x-2y+3z)^2 + (3z-2y)^2 + 2(x-2y+3z)(2y-3z)$ .
18. "  $(a^2+b^2-c^2)^2 + (c^2-b^2)^2 + 2(b^2-c^2)(a^2+b^2-c^2)$ .
19. "  $(x+y)^4 + (x-y)^4 - 2(x-y)^2(x+y)^2$ .

20. "  $(5a + 3b)^2 + 16(3a + b)^2 - (13a + 5b)^2$ .

21. Shew that  $(3a - b)^2 + (3b - c)^2 + (3c - a)^2 - 2(b - 3a)(3b - c) + 2(3b - c)(3c - a) - 2(a - 3c)(3a - b) - 4(a + b + c)^2 = 0$ .

22. If  $z^2 = 2xy$ , prove that  $(2x^2 - y^2)^2 + (z^2 - 2y^2)^2 + (x^2 - 2z^2)^2 - 2(2x^2 - y^2)(z^2 - 2y^2) + 2(x^2 - 2z^2)(z^2 - 2y^2) - 2(x^2 - 2z^2)(2x^2 - y^2) = (x + y)^4$ .

23. Simplify  $(1 + x + x^2 + x^3)^2 + (1 - x - x^2 + x^3)^2 + (1 - x + x^2 - x^3)^2 + (1 + x - x^2 - x^3)^2$ .

24. Simplify  $(ax + by)^4 - 2(a^2x^2 + b^2y^2)(ax + by)^2 + 2(a^4x^4 + b^4y^4)$ .

FORMULAS [2] AND [3].—EXAMPLES.

1.  $(1 - 2x + 3x^2)^2 = 1 - 4x + 6x^2 + 4x^2 - 12x^3 + 9x^4$

$= 1 - 4x + 10x^2 - 12x^3 + 9x^4$ .

2.  $(ab + bc + ca)^2 = a^2b^2 + 2ab^2c + 2a^2bc + b^2c^2 + 2abc^2 + c^2a^2 = a^2b^2 + b^2c^2 + c^2a^2 + 2abc(a + b + c)$ .

3.  $\{(x + y)^2 + x^2 + y^2\}^2 = (x + y)^4 + 2(x + y)^2(x^2 + y^2) + x^4 + 2x^2y^2 + y^4 = (x + y)^4 + (x + y)^2\{(x + y)^2 + (x - y)^2\} + x^4 + 2x^2y^2 + y^4 = 2(x + y)^4 + (x^2 - y^2)^2 + x^4 + 2x^2y^2 + y^4 = 2\{(x + y)^4 + x^4 + y^4\}$ .

4.  $(x^2 + xy + y^2)^2 = x^4 + 2x^3y + 2x^2y^2 + x^2y^2 + 2xy^3 + y^4 = (x + y)^2x^2 + x^2y^2 + y^2(x + y)^2$ .

5. In Ex. 3, substitute  $b - c$  for  $x$ ,  $c - a$  for  $y$ , and consequently  $b - a$  for  $x + y$ , then since  $(b - a)^2 = (a - b)^2$ , Ex. 3 gives  $\{(a - b)^2 + (b - c)^2 + (c - a)^2\}^2 = 2\{(a - b)^4 + (b - c)^4 + (c - a)^4\}$ .

6. Making the same substitutions in Ex. 4, we have  $(x^2 + b^2 + c^2 - ab - bc - ca)^2 = (a - b)^2(b - c)^2 + (b - c)^2(c - a)^2 + (c - a)^2(a - b)^2$ , or, multiplying both sides by 4,

$\{(a - b)^2 + (b - c)^2 + (c - a)^2\}^2 = 4(a - b)^2(b - c)^2 + 4(b - c)^2(c - a)^2 + 4(c - a)^2(a - b)^2$ , and  $\therefore$  from Ex. 5,  $(a - b)^4 + (b - c)^4 + (c - a)^4 = 2(a - b)^2(b - c)^2 + 2(b - c)^2(c - a)^2 + 2(c - a)^2(a - b)^2$ .



## Exercise vi.

1.  $(1 - 2x + 3x^2 - 4x^3)^2$ ,  $(1 - x + x^2 - x^3)^2$ .
2.  $(1 - 2x + 2x^2 - 3x^3 - x^4)^2$ ,  $(1 + 3x + 3x^2 + x^3)^2$ .
3.  $(2a - b - c^2 - 1)^2$ ,  $(1 - x + y + z)^2$ ,  $(\frac{1}{2}x - \frac{1}{3}y + 6z)^2$ .
4.  $(x^3 - x^2y + xy^2 - y^3)^2$ ,  $(ax + bx^2 + cx^3 + dx^4)^2$ .
5. Shew that  $(a^2 + b^2 + c^2)(x^2 + y^2 + z^2) - (ax + by + cz)^2 = (ay - bx)^2 + (cx - az)^2 + (bz - cy)^2$ .
6. Prove that  $(a + b)x + (b + c)y + (c + a)z$  multiplied by  $(a - b)x + (b - c)y + (c - a)z$ , is equal to the difference of the squares of two trinomials.
7. Shew that  $(a - b)(a - c) + (b - c)(b - a) + (c - a)(c - b) - \frac{1}{2}\{(a - b)^2 + (b - c)^2 + (c - a)^2\} = 0$ .
8. Simplify  $\{a - (b - c)\}^2 + \{b - (c - a)\}^2 + \{c - (a - b)\}^2$ .
9. Shew that  $(a^2 + b^2 - x^2)^2 + (a_1^2 + b_1^2 - x^2)^2 + 2(aa_1 + bb_1)^2 = (a^2 + a_1^2 - x^2)^2 + (b^2 + b_1^2 - x^2)^2 + 2(ab + a_1b_1)^2$ .
10. Prove that  $\{(a - b)(b - c) + (b - c)(c - a) + (c - a)(a - b)\}^2 = (a - b)^2(b - c)^2 + (b - c)^2(c - a)^2 + (c - a)^2(a - b)^2$ .
11. Square  $2a - \frac{1}{2}bx - \frac{1}{3}cx + 2dx$ .
12. If  $x + y + z = 0$ , shew that  $x^4 + y^4 + z^4 = (x^2 - y^2)^2 + (y^2 - z^2)^2 + (z^2 - x^2)^2$ .
13. Prove that  $a^2(b + c)^2 + b^2(c + a)^2 + c^2(a + b)^2 + 2abc(a + b + c) = 2(ab + bc + ca)^2$ .

**Art. V.** To apply formula [4] to obtain the product of two factors which differ only in the signs of some of their terms:—group together all the terms whose signs are the same in one factor as they are in the other, and then form into a second group all the other terms.

## EXAMPLES.

1. Multiply  $a + b - c + d$  by  $a - b - c - d$ ; here the first group is  $a - c$ , the second  $b + d$ ;  $\therefore$  we have

$$\{(a - c) + (b + d)\} \{(a - c) - (b + d)\} = (a - c)^2 - (b + d)^2.$$

2.  $(1 + 3x + 3x^2 + x^3)(1 - 3x + 3x^2 - x^3) = \{(1 + 3x^2) + (3x + x^3)\} \{(1 + 3x^2) - (3x + x^3)\} = (1 + 3x^2)^2 - (3x + x^3)^2 = 1 - 3x^2 + 3x^4 - x^6.$

3. Find the continued product of  $a + b + c$ ,  $b + c - a$ ,  $c + a - b$  and  $a + b - c$ .

The first pair of factors gives  $\{(b + c) + a\} \{(b + c) - a\} = (b + c)^2 - a^2 = b^2 + 2bc + c^2 - a^2.$

The second pair gives  $\{a - (b - c)\} \{a + (b - c)\} = a^2 - b^2 + 2bc - c^2$ ; the only term whose sign is the same in both these results is  $2bc$ ; hence, grouping the other terms, we have

$\{2bc + (b^2 + c^2 - a^2)\} \{2bc - (b^2 + c^2 - a^2)\} = (2bc)^2 - (b^2 + c^2 - a^2)^2 = 2a^2b^2 + 2b^2c^2 + 2c^2a^2 - a^4 - b^4 - c^4.$

4. Prove  $(a^2 + ab + b)^2 - a^2b^2 = (a^2 + ab)^2 + (ab + b^2)^2.$

The expression  $= (a^2 + b^2)(a^2 + 2ab + b^2) = (a^2 + b^2)(a + b)^2 = a^2(a + b)^2 + b^2(a + b)^2 = (a^2 + ab)^2 + (ab + b^2)^2.$

Exercise vii.

1.  $(a^2 + 2ab + b^2)(a^2 - 2ab + b^2).$

2.  $(\frac{1}{2}x^2 - xy + y^2)(\frac{1}{2}x^2 + y^2 + xy).$

3.  $(a^2 - ab + 2b^2)(a^2 + ab + 2b^2); (x^4 + 4xy)(x^4 - 4xy).$

4.  $\{(x + y)x - y(x - y)\} \{(x - y)x - y(y - x)\}.$

5. Simplify:  $(x + 3)(x - 3) + (x + 4)(x - 4) - (x + 5)(x - 5).$

6.  $(1 + x)^4 + (1 - x)^4 - 2(1 - x^2)^2.$

7.  $(x^2 + y^2)^2 - (2xy)^2 - (x^2 - y^2)^2.$

8.  $(2a^2 - 3b^2 + 4c^2)(2a^2 + 3b^2 - 4c^2).$

9.  $(2a + b - 3c)(b + 3c - 2a); (2a - b - 3c)(b - 3c - 2a).$

10.  $(x^4 + y^4)(x^2 + y^2)(x + y)(x - y).$

11.  $(x^2 + xy + y^2)(x^2 - xy + y^2)(x^4 - x^2y^2 + y^4).$

12.  $(a + b - ab - 1)(a + b + ab + 1).$

13. Prove  $(a^2 + b^2 + c^2)(b^2 + c^2 - a^2)(c^2 + a^2 - b^2)(a^2 + b^2 - c^2) = 4b^4c^4$  when  $a^4 = b^4 + c^4.$

14.  $(x^2 + y^2 - \frac{5}{4}xy)(x^2 + y^2 + \frac{5}{4}xy).$

15.  $(x^4 - 2x^3 + 3x^2 - 2x + 1)(x^4 + 2x^3 + 3x^2 + 2x + 1).$

16. Multiply  $(2x-y)a^2 - (x+y)ax + x^3$  by  $(2x-y)a^2 + (x+y)ax - x^3$ .

Prove the following:

$$17. (a^2 + b^2 + c^2 + ab + bc + ca)^2 - (ab + bc + ca)^2 = (a + b + c)^2 \times (a^2 + b^2 + c^2).$$

$$18. (a^2 + b^2 + c^2 + ab + bc + ca)^2 - (a^2 + ab + ca - bc)^2 = \{(a+b)(b+c)\}^2 + \{(b+c)(c+a)\}^2.$$

$$19. 4(ab+cd)^2 - (a^2 + b^2 - c^2 - d^2)^2 = (a+b+c-d)(a+b-c+d)(c+d+a-b)(c+d-a+b).$$

20. Find the product of  $x^3 - y^3 + z^3 - 2xy + 2xz - 2yz$  and  $x^3 + y^3 + z^3 - 2xy - 2xz + 2yz$ .

$$21. (x^2 + y^2 + xy\sqrt{2})(x^2 - xy\sqrt{2} + y^2)(x^4 - y^4).$$

$$22. (1 - 6a + 9a^2)(\frac{1}{3} + 2a + 3a^2).$$

$$23. \{(m+n) + (p+q)\} (m - q + p - n).$$

24. Obtain the product of  $1+x+x^2$ ,  $x^2+x-1$ ,  $x^2-x+1$ , and  $1+x-x^2$ .

$$25. (a-b^2)^2 (a+b^2)^2 (a^2+b^4)^2 (a^4+b^8)^2.$$

26. Shew that  $(x^3 + xy + y^2)^2 (x^2 - xy + y^2)^2 - (x^2 y^2)^2 = (x^4 + x^2 y^2)^2 + (x^2 y^2 + y^4)^2$ .

#### FORMULA A.—EXAMPLES.

1. Multiply  $x^2 - x + 5$  by  $x^2 - x - 7$ : here the common term is  $x^2 - x$ , the other terms  $+5$ , and  $-7$ , hence the product  $= (x^2 - x)^2 + (-7 + 5)(x^2 - x) + (-7 \times 5) = (x^2 - x)^2 - 2(x^2 - x) - 35 = x^4 - 2x^3 - x^2 + 2x - 35$ .

2.  $(x-a)(x-3a)(x+4a)(x+6a)$ : taking the first and third factors together, and the second and fourth, we have the product  $= (x^2 + 3ax - 4a^2)(x^2 + 3ax - 18a^2) = (x^2 + 3ax)^2 - (4a^2 + 18a^2) \times (x^2 + 3ax) - 72a^4 = \&c$ .

#### Exercise viii.

$$1. (x^2 + 2x + 3)(x^2 + 2x - 4); (x - y + 3z)(x - y + 5z).$$

$$2. (x+1)(x+5)(x+2)(x+4); (x^3 + a - b)(x^3 + 2b - a).$$

$$3. (a^2 - 3)(a^2 - 7)(a^2 + 5)(a^2 + 7); (x^4 + x^2 + 1)(x^4 + x^2 - 2).$$

$$4. \{(x+y)^2 - 4xy\} \{(x+y)^2 + 5xy\}.$$

5.  $(x^n + a + 7)(x^n - a - 9) ; \left(\frac{x}{y} + \frac{y}{x} - 1\right) \left(\frac{x}{y} + \frac{y}{x} + 3\right)$ .
6.  $(nx + y + 3)(nx + y + 7)$ .
7.  $(x + a - y)(x + a + 3y)$ .
8.  $(x^{2n} + x^n - a)(x^{2n} + x^n - b)$ .
9.  $(\frac{1}{2}x^4 - y^2 + 2)(\frac{1}{2}x^4 - y^2 - 4)$ .
10.  $\left(\frac{1}{x} + \frac{1}{y} - \frac{1}{2}\right) \left(\frac{1}{x} + \frac{1}{y} + 2\frac{1}{2}\right)$ .

11. Multiply together  $x - 2 + \sqrt{2}$ ,  $x - 2 + \sqrt{3}$ ,  $x - 2 - \sqrt{2}$ , and  $x - 2 - \sqrt{3}$ .

12.  $(x + a + b)(x + b - c)(x - a + b)(x + b + c)$ .

13.  $(a + b + c)(a + b + d) + (a + c + d)(b + c + d) - (a + b + c + d)^2$ .

14. Prove that

$$(2a + 2b - c)(2b + 2c - a) + (2c + 2a - b)(2a + 2b - c) + (2b + 2c - a)(2c + 2a - b) = 9(ab + bc + ca)$$

FORMULAE [5] AND [6].—EXAMPLES.

1. We get at once

$$(x + y)^3 + (x - y)^3 = 2x(x^2 + 3y^2)$$

$$(x + y)^3 - (x - y)^3 = 2y(3x^2 + y^2)$$

2. Simplify  $(a + b + c)^3 - 3(a + b + c)^2c + 3(a + b + c)c^2 - c^3$ .

This plainly comes under formula [5], the first term being  $a + b + c$ , the second  $-c$ ; hence the expression is  $\{(a + b + c) - c\}^3 = (a + b)^3$ .

3. Shew that  $(x^2 + xy + y^2)^3 + (xy - x^2 - y^2)^3 - 6xy(x^4 + x^2y^2 + y^4) = 8x^3y^3$ .

This comes under formula [6], the first term being  $(x^2 + xy + y^2)$ , and the second  $-(x^2 - xy + y^2)$ ; we have therefore  $\{(x^2 + xy + y^2) - (x^2 - xy + y^2)\}^3 = (2xy)^3 = 8x^3y^3$ .

Exercise ix.

Simplify

1.  $(1 - x^2)^3 + (1 + x^2)^3$ ,  $(x^2 + xy^3)^3 - (x^2 - xy^3)^3$ .

2.  $(a + 2b)^3 - (a - b)^3$ ,  $(3a - b)^3 - (3a - 2b)^3$ .

3.  $(x+y-z)^3 + 3(x+y-z)^2z + 3(x+y-z)z^2 + 3(x+y-z)z^2$ .
4.  $(a-b)^3 + (a+b)^3 + 6a(a^2 - b^2)$ .
5.  $(x-y)^3 + (x+y)^3 + 3(x-y)^2(x+y) - 3(y-x)(x+y)^2$ .
6.  $(1+x+x^2)^3 - (1-x+x^2)^3 - 6x(1+x^2+x^4)$ .
7.  $(a-b-c)^3 + (b+c)^3 + 3(b+c)^2(a-b-c) + 3(c-b-c)^2(b+c)$ .
8.  $(3x-4y+5z)^3 - (5z-4y)^3 + 3(5z-4y)^2(3x-4y+5z) - 3(3x-4y+5z)^2(5z-4y)$ .
9.  $(1+x+x^2)^3 + 3(1-x^3)(2+x^2) + (1-x)^3$ .
10. Show that  $a(a-2b)^3 - b(b-2a)^3 = (a-b)(a+b)^3$ .
11. Show that  $a^3(a^3-2b^2)^3 + b^3(2a^3-b^3)^3 = (a^3-b^3)(a^3+b^3)^3$ .
12.  $(x^2+xy+y^2)^3 + 6(x^2+y^2)(x^4+xy^2+y^4) + (x^2-xy+y^2)^3$ .
13. Show that  $a^3(a^3+2b^3)^3 + b^3(2a^3+b^3)^3 + (3a^2b^2)^3 = (a^6+7a^3b^3+b^6)^2$ .
14. Simplify  $(ax+by)^3 + a^3y^3 + b^3x^3 - 3abxy(ax+by)$ .
15. What will  $a^3+b^3+c^3-3abc$  become when  $a+b+c=0$ ?
16. Find the value of  $x^6-y^6+z^6+3x^2y^2z^2$  when  $x^2-y^2+z^2=0$ .

FORMULAS [7], [8] AND [9].—EXAMPLES.

1. Simplify  $(2x-3y)^3 + (4y-5x)^3 + (3x-y)^3 - 3(2x-3y)(4y-5x)(3x-y)$ .

By [8] this is seen to be  $\{(2x-3y) + (4y-5x) + (3x-y)\}^3 - (0)^3 = 0$ .

2. Prove that  $(a-b)^3 + (b-c)^3 + (c-a)^3 = 3(a-b)(b-c)(c-a)$ .

In [8] substitute  $a-b$  for  $x$ ,  $b-c$  for  $y$ , and  $c-a$  for  $z$ ; for these values  $x+y+z=0$ , and the identity appears at once.

3. Prove  $(a+b+c)^3 - (b+c-a)^3 - (a+c-b)^3 - (a+b-c)^3 = 24abc$ .

In [7] let  $x=b+c-a$ ,  $y=c+a-b$ ,  $z=a+b-c$ , and therefore  $x+y+z=2c$ ,  $y+z=2a$ ,  $z+x=2b$ , and this identity at once appears.

Exercise x.

1. Cube the following:  $1-x+x^2$ ,  $a-b-c$ ,  $1-2x+3x^2-4x^3$ .
2. Simplify  $(x^2+2x-1)^3+(2x-1)(x^2+2x-2) - (x^3+3x^2-1)^3$ .
3. \*Prove that  $(x+y)(y+z)(z+x)+xyz=(x+y+z)(xy+yz+zx)$
4. Prove that  $(ax-by)^3+a^3y^3-b^3x^3+3abxy(ax-by) = (a^3-b^3)(x^3+y^3)$ .
5. Simplify  $(x-2y)^3+(y-2z)^3+(z-2x)^3+3(x-y-2z)(y-z-2x)(z-x-2y)+(x+y+z)^3$ .
6. Simplify  $(2x^2-3y^2+4z^2)^3+(2y^2-3z^2+4x^2)^3+(2z^2-3x^2+4y^2)^3$ .
7. Simplify  $(2ax-by)^3+(2by-cz)^3+(2cz-ax)^3+3(2ax+by-cz)(2by+cz-ax)(2cz+ax-by)$ .
8. Prove  $(x^3+3x^2y-y^3)^3+\{3xy(x+y)\}^3=\{(x-y)^3+9x^2y\} \times \{x^2+xy+y^2\}^3$ .
9. Prove  $9(x^3+y^3+z^3)-(x+y+z)^3=(4x+4y+z)(x-y)^2+(4y+4z+x)(y-z)^2+(4z+4x+y)(z-x)^2$ .
10. If  $x+y+z=0$ , shew that  $x^3+y^3+z^3=3xyz$ .
11. If  $x=2y+3z$  shew that  $x^3-8y^3-27z^3-18xyz=0$ .
12. Shew that  $(x^2+xy+y^2)^3+(x^2-xy+y^2)^3+8z^3-6z^2(x^4+x^2y^2+y^4)=0$ , if  $x^2+y^2+z^2=0$ .
13. Prove that  $8(a+b+c)^3-(a+b)^3-(b+c)^3-(c+a)^3=3(2a+b+c)(a+2b+c)(a+b+2c)$ .

Prove the following :

14  $(ax-by)^3+b^3y^3=a^3z^3+3abxy(by-ax)$ .

---

\*Note that the right-hand member is formed from the left-hand one by changing additions into multiplications, and multiplications into additions; hence in  $(x+y+z)(xy+yz+zx)$  the signs + and . may be interchanged throughout without altering the value of the expression.

$$15. a^3 + b^3 + c^3 - 3abc = \frac{1}{2} \{ (a-b)^2 + (b-c)^2 + (c-a)^2 \} \times (a+b+c).$$

$$16. (x+b+c) \{ (a+b-c)(b+c-a) + (b+c-a)(c+a-b) + (c+a-b)(a+b-c) \} = (a+b-c)(b+c-a)(c+a-b) + 8abc.$$

$$17. a^3 + b^3 + c^3 + 2abc = (a+b+c)^3 - 3 \{ a(b-c)^2 + b(c-a)^2 + c(a-b)^2 \}.$$

$$18. (a+b+7c)(a-b)^2 + (b+c+7a)(b-c)^2 + (c+a+7b)(c-a)^2 = 2(a+b+c)^3 - 54abc.$$

$$19. (a+b+c) \{ (2a-b)(2b-c) + (2b-c)(2c-a) + (2c-a)(2a-b) \} = (2a-b)(2b-c)(2c-a) + (2a+b-c)(2b+c-a)(2c+a-b).$$

$$20. \text{ If } x^2(y+z) = a^3, y^2(z+x) = b^3, z^2(x+y) = c^3, \text{ and } xyz = abc, \text{ shew that } a^3 + b^3 + c^3 + 2abc = (x+y)(y+z)(z+x)$$

## EXPANSION OF BINOMIALS.

We have from formula [5]

$$(a+b)^3 = a^3 + 3a^2b + 3ab^2 + b^3; \text{ multiplying by } a+b \text{ we get}$$

$$(a+b)^4 = a^4 + 4a^3b + 6a^2b^2 + 4ab^3 + b^4; \text{ multiplying this by } a+b \text{ we get}$$

$$(a+b)^5 = a^5 + 5a^4b + 10a^3b^2 + 10a^2b^3 + 5ab^4 + b^5.$$

From these examples we derive the following law for the formation of the terms in the expansion of  $a+b$  to any required power:—

(1). The *index* of  $a$ , in the *first* term, is that of the given power, and *decreases* by unity in each succeeding term; the index of  $b$  begins with unity in the *second* term and *increases* by unity in each succeeding term.

(2). The coefficient of the first term is unity, and the coefficient of any other term is found by multiplying the coefficient of the immediately preceding term by the index of  $a$  in that term, and dividing the product by the number of that preceding term. It will be observed that the coefficients equally distant from the extremes of the expansion, are equal.

Exercise xi.

1. Expand  $(x+y)^6$ ,  $(x+y)^7$ ,  $(x+y)^8$ ,  $(x+y)^{12}$ .
2. What will be the law of signs if  $-y$  be written for  $y$  in (1) ?
3. Expand  $(a-b)^5$ ,  $(a-2b)^4$ ,  $(2b-a)^4$ .
4. Expand  $(1+m)^6$ ,  $(m+1)^5$ ,  $(2m+1)^6$ .
5. What is the coefficient of the 4th term in  $(a-b)^{10}$  ?
6. Expand  $(x^2-y)^4$ ,  $(a-2b^2)^5$ ,  $(a^3-2b^3)^6$ .
7. In the expansion of  $(a-b)^{12}$ , the third term is  $66a^{10}b^2$ , find the 5th and 6th terms.
8. Shew that  $(x+y)^5 - x^5 - y^5 = 5xy(x+y)(x^2+xy+y^2)$ .
9. From (8) shew that  $2\{(a-b)^5 + (b-c)^5 + (c-a)^5\} = 5(a-b)(b-c)(c-a)\{(a-b)^2 + (b-c)^2 + (c-a)^2\}$ .

SECTION III.—HORNER'S METHODS OF MULTIPLICATION AND DIVISION.

EXAMPLES.

1. Find the product of  $kx^2+lx^2+mx+n$  and  $ax^2+bx+c$ . Write the multiplier in a column to the left of the multiplicand, placing each term in the same horizontal line with the partial product it gives :

	$kx^3$	$+lx^2$	$+mx$	$+n$	.....Q
$ax^2$	$akx^5$	$+alx^4$	$+amx^3$	$+anx^2$	..... $p_1$
$+bx$		$+bklx^4$	$+blmx^3$	$+bmx^2 + bnx$	..... $p_2$
$+c$			$+ckx^3$	$+clx^2 + cmx + cn$	..... $p_3$
	$akx^5 + (al+bk)x^4 + (am+bl+ck)x^3 + (an+bm+cl)x^2 + (bn+cm)x + cn$				.....P.

Art. VI. The above example has been given in full, the powers of  $x$  being inserted ; in the following example detached coefficients are used. It is evident that if the coefficient of the first term of the multiplier be unity, the coefficients of the multiplicand will be the same as those of the first partial product, and may be used for them, thus saving the repetition of a line.



2. Multiply  $3x^4 - 2x^3 - 2x + 3$  by  $x^2 + 3x - 2$ .

$$\begin{array}{r}
 1 \quad 3 \quad -2 \quad +0 \quad -2 \quad +3 \\
 +3 \quad \quad \quad +9 \quad -6 \quad +0 \quad -6 \quad +9 \\
 -2 \quad \quad \quad \quad \quad -6 \quad +4 \quad -0 \quad +4 \quad -6 \\
 \hline
 3x^6 + 7x^5 - 12x^4 + 2x^3 - 3x^2 + 13x - 6.
 \end{array}$$

3. Find the product of  $(x-3)(x+4)(x-2)(x-5)$ .

$$\begin{array}{r}
 +4 \quad \left| \begin{array}{r} 1 \quad -3 \\ \hline \quad +4 \quad -12 \end{array} \right. \\
 -2 \quad \left| \begin{array}{r} 1 \quad +1 \quad -12 \\ \hline \quad -2 \quad -2 \quad +24 \end{array} \right. \\
 -5 \quad \left| \begin{array}{r} 1 \quad -1 \quad -14 \quad +24 \\ \hline \quad -5 \quad +5 \quad +70 \quad -120 \end{array} \right. \\
 \hline
 x^4 - 6x^3 - 9x^2 + 94x - 120.
 \end{array}$$

4. Multiply  $x^3 - 4x^2 + 2x - 3$  by  $2x^3 - 3$

$$\begin{array}{r}
 1 \quad -4 \quad +2 \quad -3 \\
 \hline
 2 \quad 2 \quad -8 \quad +4 \quad -6 \quad \quad \quad [x^3 \times x^3 = x^6] \\
 0 \quad \quad \quad 0 \quad \quad 0 \quad \quad 0 \\
 0 \quad \quad \quad \quad \quad 0 \quad \quad 0 \quad \quad 0 \\
 -3 \quad \quad \quad \quad \quad -3 \quad +12 \quad -6 \quad +9 \\
 \hline
 2x^6 - 8x^5 + 4x^4 - 9x^3 + 12x^2 - 6x + 9
 \end{array}$$

In this example the missing terms of the multiplier are supplied by zeros; but instead of writing the zeros as in the example, we may, as in ordinary arithmetical multiplication, "skip a line" for every missing term.

5. Multiply  $x^4 - 2x^2 + 1$  by  $x^4 - x^2 + 3$ .

$$\begin{array}{r}
 1 \quad 1 \quad +0 \quad -2 \quad +0 \quad +1 \\
 -1 \quad \quad \quad -1 \quad -0 \quad +2 \quad -0 \quad -1 \quad \quad \quad [x^4 \times x^4 = x^8] \\
 +3 \quad \quad \quad \quad \quad +3 \quad +0 \quad -6 \quad +0 \quad +3 \\
 \hline
 x^8 \quad -3x^6 \quad +6x^4 \quad -7x^2 \quad +3
 \end{array}$$

6. Find the value of  $(x+2)(x+3)(x+4)(x+5) - 9(x+2)(x+3) \times (x+4) + 3(x+2)(x+3) + 77(x+2) - 85$ .

$$\begin{array}{r}
 \phantom{+4} \phantom{+3} \phantom{+2} \phantom{+} \\
 \phantom{+4} \phantom{+3} \phantom{+2} \phantom{+} \phantom{+} \\
 +4 \phantom{+3} \phantom{+2} \phantom{+} \phantom{+} \phantom{+} \\
 +3 \phantom{+2} \phantom{+} \phantom{+} \phantom{+} \phantom{+} \\
 +2 \phantom{+} \phantom{+} \phantom{+} \phantom{+} \phantom{+} \\
 \hline
 x^4 + 5x^3 - 7x^2 + 12x - 9
 \end{array}$$

7. Find the coefficient of  $x^4$  in the product of  $x - ax^3 + bx^2 - cx + d$  and  $x^2 + px + q$ .

$$\begin{array}{r}
 \phantom{+p} \phantom{+q} \\
 +p \phantom{+q} \\
 +q \\
 \hline
 +(b-ap+q)
 \end{array}$$

Exercise xii.

Find the product of

1.  $(1+x+x^2+x^3+x^4)(1-x+x^3-x^7+x^8-x^{12}+x^{13})$ .
2.  $(1+x^5)(1-x^5+x^6)(1+x+x^2+x^3+x^4)$ .
3.  $(x-5)(x+6)(x-7)(x+8); (2x^5-x^2+1)(x^4-x+2)$ .
4.  $(x^3+5x^2-16x-1)(x^3-5x^2-16x+1)$ .
5.  $(6x^6-x^5+2x^4-2x^3+2x^2+19x+6)(3x^2+4x+1)$ .

Obtain the coefficients of  $x^4$  and lower powers in

6.  $(1+\frac{1}{2}x-\frac{1}{8}x^2+\frac{1}{16}x^3-\frac{5}{128}x^4)(1-\frac{1}{2}x-\frac{1}{8}x^2-\frac{1}{16}x^3-\frac{5}{128}x^4)$ .
7. Multiply  $2x^7-x^3+3x-4$  by  $3x^5-2x^2-x-1$ .

Simplify the following :

$$8. (x+1)(x+2)(x+3) + 3(x+1)(x+2) - 10(x+1) + 9.$$

$$9. x(x+1)(x-2)(x+3) - 3x(x+1)(x+2) - 2x(x+1) + 2x.$$

$$10. x(x-1)(x-2)(x-3) + 3x(x-1)(x-2) - 2x(x-1) - 2x.$$

$$11. (x-1)(x+1)(x+3)(x+5) - 14(x-1)(x+1) + 1.$$

12. Given that the sum of the four following factors is  $-1$ , find (1) the product of the first pair; (2) the product of the second pair; and (3) the product of the sum of the first pair by the sum of the second pair.

$$(1) \quad x + x^4 + x^{13} + x^{16}$$

$$(2) \quad x^2 + x^5 + x^9 + x^{15}$$

$$(3) \quad x^3 + x^6 + x^{12} + x^{14}$$

$$(4) \quad x^6 + x^7 + x^{10} + x^{11}.$$

13. Given that the sum of the three following factors is equal to  $-1$ , find their product.

$$(1) \quad x + x^5 + x^8 + x^{13}$$

$$(2) \quad x^2 + x^3 + x^{10} + x^{11}$$

$$(3) \quad x^4 + x^6 + x^7 + x^9.$$

Art. VII. Were it required to *divide* the product  $P$  in the first of the above examples by  $ax^2 + bx + c$ , it is evident that could we find and *subtract* from  $P$  the partial products  $p_2, p_3$ , (or what would give the same result, could we *add* them with the sign of each term changed), there would remain the partial product  $p_1$ , which, divided by the monomial  $ax^2$ , would give the quotient  $Q$ . This is what Horner's method does, the change of sign being secured by changing the signs of  $b$  and  $c$ , which are factors in each term of  $p_2, p_3$ , respectively.

$$\begin{array}{r}
 1. \quad \begin{array}{r}
 ax^5 + (al + bl)ax^4 + (am + bm + cl)x^3 + (bn + cm)x + cn \dots\dots\dots P \\
 - bx \phantom{+ (al + bl)ax^4} \phantom{+ (am + bm + cl)x^3} \phantom{+ (bn + cm)x} \phantom{+ cn} \phantom{\dots\dots\dots} \\
 - c \phantom{+ (al + bl)ax^4} \phantom{+ (am + bm + cl)x^3} \phantom{+ (bn + cm)x} \phantom{+ cn} \phantom{\dots\dots\dots} \\
 \hline
 ax^2 \quad \begin{array}{r}
 alx^5 \phantom{+ (al + bl)ax^4} \phantom{+ (am + bm + cl)x^3} \phantom{+ (bn + cm)x} \phantom{+ cn} \phantom{\dots\dots\dots} P_1 \\
 \phantom{alx^5} + alx^4 \phantom{+ (am + bm + cl)x^3} \phantom{+ (bn + cm)x} \phantom{+ cn} \phantom{\dots\dots\dots} \\
 \phantom{alx^5} \phantom{+ alx^4} + amx^3 \phantom{+ (bn + cm)x} \phantom{+ cn} \phantom{\dots\dots\dots} \\
 \phantom{alx^5} \phantom{+ alx^4} \phantom{+ amx^3} + aux^2 \phantom{+ cn} \phantom{\dots\dots\dots} P_2 \\
 \phantom{alx^5} \phantom{+ alx^4} \phantom{+ amx^3} \phantom{+ aux^2} + mx \phantom{+ cn} \phantom{\dots\dots\dots} P_3 \\
 \phantom{alx^5} \phantom{+ alx^4} \phantom{+ amx^3} \phantom{+ aux^2} \phantom{+ mx} + n \phantom{+ cn} \phantom{\dots\dots\dots} Q
 \end{array}
 \end{array}
 \end{array}$$

The dividend and divisor are arranged as in the example, the sign of every term in the divisor, except the first, being changed in order to turn the subtractions into additions. The first term of the dividend ( $alx^5$ ) is brought down into the line of  $p_1$ ; dividing this by  $ax^2$ , the first term of the divisor, we get  $lx^3$ , the first term of the quotient. Multiplying this term,  $lx^3$ , by  $-bx$  and  $-c$ , respectively, and writing the products in the proper columns and rows, makes all ready to give the second term of  $p_1$ , which is got by simply adding up the second column of the work, giving  $alx^4$ . Dividing this second term of  $p_1$  by  $ax^2$ , gives  $lx^2$ , in the same way as was done in getting the second term of the quotient, and the third will be obtained. Repeating the steps, the complete quotient and the remainder will finally be obtained.

Should the coefficient of the first term of the divisor be unity, the coefficients of the line  $Q$  will be the same as those of  $p_1$ , and the line  $Q$  need not be written down, since one line does for both.

2x.  
x.  
-1, find  
second  
the sum  
  
is equal  
  
P in the  
that could  
(or what  
the sign of  
product  $p_1$ ,  
quotient  $Q$ .  
sign being  
factors in

2. Divide  $3x^6 + 7x^5 - 12x^4 + 2x^3 - 3x^2 + 13x - 6$  by  $x^2 + 3x - 2$ .

$$\begin{array}{r|rrrrrrrr}
 -3 & 3 & +7 & -12 & +2 & -3 & +13 & -6 & \{x^6 \div x^2 = x^4\} \\
 +2 & & -9 & +6 & -0 & +6 & -9 & & \\
 \hline
 & & & +6 & -4 & +0 & -4 & +6 & \\
 \hline
 & & & & 3x^4 & -2x^3 & +0 & -2x & +3
 \end{array}$$

Compare this example with the second example of Horner's Multiplication, performing a step in multiplication, then the corresponding step in division; then another step in multiplication and the second (corresponding) step in division, and so on.

3. Divide  $x^7 - 3x^6 + 4x^4 + 18x^3 - 7x^2 + 12$  by  $x^3 - 3x^2 + 3x - 1$ .

$$\begin{array}{r|rrrrrrrr}
 +3 & 1 & -3 & +0 & -4 & +18 & +0 & -7 & +12 \\
 -3 & & +3 & +0 & -9 & -36 & -27 & & \\
 +1 & & & -3 & -0 & +9 & +36 & +27 & [x^7 \div x^3 = x^4]. \\
 \hline
 & & & & +1 & +0 & -3 & -12 & -9 \\
 \hline
 & & & & & x^4 & +0 & -3x^2 & -12x & -9; & 6x^2 & +8x & +3
 \end{array}$$

The quotient is therefore  $x^4 - 3x^2 - 12x - 9$ , and the remainder  $6x^2 + 8x + 3$ .

4. Divide  $x^8 - 3x^7 - 5x^5 + 2x^4 + 5x^3 + 4x^2 + 1$  by  $x^3 + 2x - 1$ . The zero coefficient in the divisor may be inserted, or it may be omitted and allowance made for it in the  $2x$ -line. See examples 4 and 5 in multiplication.

$$\begin{array}{r|rrrrrrrrr}
 -2 & 1 & -3 & +0 & -5 & +2 & +5 & +4 & +0 & +1 \\
 +1 & & & -2 & +6 & +4 & -4 & -6 & +2 & \\
 \hline
 & & & & & 1 & -3 & -2 & +2 & +3 & -1 \\
 \hline
 & & & & & & 1 & -3 & -2 & +2 & +3 & -1; & 0 & +5 & +0
 \end{array}$$

$[x^8 \div x^3 = x^5]$ . The quotient is therefore  $x^5 - 3x^4 - 2x^3 + 2x^2 + 3x - 1$ , and the remainder  $5x$ .

5. Divide  $10x^6 - 11x^5 - 3x^4 + 20x^3 + 10x^2 + 2$  by  $5x^3 - 3x^2 + 2x - 2$ .

Arranging as in the ordinary method, we have

$$\begin{array}{r|rrrr|rrr}
 & 10 & -11 & -8 & +20 & +10 & +0 & +2 \\
 +3 & & 6 & -3 & -6 & +12 & & \\
 -2 & & & -4 & +2 & +4 & -8 & \\
 +2 & & & & +4 & -2 & -4 & +8 \\
 \hline
 5 & 2 & -1 & -2 & +4 & 24 & -12 & +10
 \end{array}$$

$$\text{Quotient} = 2x^3 - x^2 - 2x + 4 + \frac{24x^2 - 12x + 10}{5x^3 - 3x^2 + 2x - 2}$$

We first draw a vertical line with as many vertical columns to the right as are less by unity than the number of terms in the divisor. This will mark the point at which the remainder begins to be formed. We then divide 5 into 10, and thus obtain the first coefficient of the dividend. We next multiply the remaining terms of the divisor by the 2 thus obtained. Adding the second vertical column and dividing by 5, we obtain -1; we multiply by the -1, add the next column and divide the sum by 5, and so on for the others.

This method is not, however, always convenient. If the first term of the dividend be not divisible by the first term of the divisor, the work would be embarrassed with fractions. We may then proceed as in the following examples :

6. Divide  $x^5 - 3x^4 + x^3 + 3x^2 - x + 3$  by  $2x^3 + x^2 - 3x + 1$ .

Let  $2x = y$ , or  $x = \frac{y}{2}$ .

Substitute  $\frac{y}{2}$  for  $x$  in the dividend and divisor, and we have

$$\begin{aligned}
 \frac{y^5}{2^5} - \frac{3y^4}{2^4} + \frac{y^3}{2^3} + \frac{3y^2}{2^2} - \frac{y}{2} + 3 & \div \frac{2y^3}{2^3} + \frac{y^2}{2^2} - \frac{3y}{2} + 1 \\
 & = \frac{y^5 - 2 \times 3y^4 + 2^2y^3 + 2^3 \times 3y^2 - 2^4y + 2^5 \times 3}{2^5} \div
 \end{aligned}$$

$$\frac{y^3 + y^2 - 2 \times 3y + 2^2}{2^2}$$

$$\therefore \frac{y^5 - 6y^4 + 4y^3 + 24y^2 - 16y + 96}{2^3} \div y^3 + y^2 - 6y + 4 \dots A$$

Dividing  $y^5 - 6y^4 + 4y^3 + 24y^2 - 16y + 96$  by  $y^3 + y^2 - 6y + 4$  by the ordinary method, and the quotient by  $2^3$  we have

$$\frac{y^2 - 7y + 17}{2^3} - \frac{1}{2^3} \cdot \frac{39y^2 - 114y - 28}{y^3 + y^2 - 6y + 4}.$$

Substituting for  $y$  its value  $2x$ , and simplifying we get

$$\frac{x^2}{2} - \frac{7x}{4} + \frac{17}{8} - \frac{1}{8} \cdot \frac{39x^2 - 57x - 7}{2x^3 + x^2 - 3x + 1}. \quad \dots\dots\dots B.$$

By comparing the dividend of  $A$  with the original question, we find that we have multiplied the successive coefficients of the dividend by  $2^0, 2^1, 2^2, \&c.$ , and, omitting the first term, we have multiplied the successive coefficients of the divisor by the same numbers. Dividing then by Horner's division we get the coefficients  $1, -7, 17$ , and for coefficients of remainder,  $-39, 114$ , and  $28$ . The first of these divided by  $2, 2^2, 2^3$  are the coefficients of  $x^2 \&c.$ ; and,  $-39, \&c.$ , are divided by  $1, 2, 2^2$ . Hence the work will stand as follows:—

	$x^5 - 3x^4 + x^3 + 3x^2 - x + 3 \div 2x^3 + x^2 - 3x + 1$										
	<table style="border-collapse: collapse; margin: 0 auto;"> <tr> <td style="padding: 0 5px;">1</td><td style="padding: 0 5px;">2</td><td style="padding: 0 5px;">4</td><td style="padding: 0 5px;">8</td><td style="padding: 0 5px;">16</td><td style="padding: 0 5px;">32</td><td style="padding: 0 5px;">1</td><td style="padding: 0 5px;">2</td><td style="padding: 0 5px;">4</td> </tr> </table>	1	2	4	8	16	32	1	2	4	
1	2	4	8	16	32	1	2	4			
-1	1	-6	+4	+24	-16	+96	1	-6	+4		
+6		-1	+7	-17							
-4			+6	-42	+102						
				-4	+28	-68					

$$\begin{aligned} \text{*Quotient} &= \frac{x^2}{2} - \frac{7x}{4} + \frac{17}{8} - \frac{1}{8} \cdot \frac{39x^2 - \frac{114x}{2} - \frac{28}{4}}{2x^3 + x^2 - 3x + 1} \\ &= \frac{x^2}{2} - \frac{7x}{4} + \frac{17}{8} - \frac{1}{8} \cdot \frac{39x^2 - 57x - 7}{2x^3 + x^2 - 3x + 1}. \end{aligned}$$

\*It will, in general, be as convenient to multiply the dividend by such a number as will make its first term exactly divisible by the first term of the divisor, and afterwards divide the quotient by this multiplier.

7. Divide  $5x^5 + 2$  by  $3x^2 - 2x + 3$ .

$$\begin{array}{r}
 5x^5 \quad 0 \quad 0 \quad 0 \quad 0 \quad +2 + 3x^2 - 2x + 3 \\
 1 \quad 3 \quad 9 \quad 27 \quad 81 \quad 243 \quad \underline{1 \quad 3} \\
 \hline
 +2 \left| \begin{array}{ccc|cc}
 5 & 0 & 0 & 0 & 0 + 486 \\
 & 10 & +20 & -50 & -280 \\
 -9 & & -45 & -90 & +225 +1260
 \end{array} \right. \quad \underline{-2 \quad +9} \\
 \hline
 5 \quad +10 \quad -25 \quad -140 \quad \left| \begin{array}{cc}
 -55 & +1746.
 \end{array} \right.
 \end{array}$$

Coeffs. of Quotient =  $\frac{5}{3} + \frac{10}{3^2} - \frac{25}{3^3} - \frac{140}{3^4} - \frac{1}{3^4} \cdot \frac{55 - 1746}{3 - 2 + 3}$ .

Quotient =  $\frac{5x^3}{3} + \frac{10x^2}{9} - \frac{25x}{27} - \frac{140}{81} - \frac{1}{81} \cdot \frac{55x - 582}{3x^2 - 2x + 3}$ .

Exercise xiii.

1. Divide  $6x^5 + 5x^4 - 17x^3 - 6x^2 + 10x - 2$  by  $2x^2 + 3x - 1$ .

2.  $(5x^6 + 6x^5 + 1) \div (x^2 + 2x + 1)$ .

3.  $(a^6 - 6a + 5) \div (a^2 - 2a + 1)$ .

4.  $(x^5 - 4x^3y^2 - 8x^2y^3 - 17xy^4 - 12y^5) \div (x^2 - 2xy - 3y^2)$ .

5.  $(a^6 - 3a^4x^2 + 3a^2x^4 - x^6) \div (a^3 - 3a^2x + 3ax^2 - x^3)$ .

Divide

6.  $4x^4 + 3x^2 - 3x + 1$  by  $x^2 - 2x + 3$ .

7.  $10x^5 + 5x^4 - 90x^3 - 44x^2 + 10x + 1$  by  $x^2 - 9$ .

8.  $x^5 - x^4y + x^3y^2 - x^2y^3 + xy^4 - y^5$  by  $x^3 - y^3$ .

9. Multiply  $x^4 - 4x^3a + 6x^2a^2 - 4xa^3 + a^4$  by  $x^2 + 2xa + a^2$ , and divide the product by  $x^4 - 2x^3a + 2xa^3 - a^4$ .

Divide

10.  $x^5 - ax^4 + bx^3 - bx^2 + ax - 1$  by  $x - 1$ .

11.  $6x^5 + 7x^4 + 7x^3 + 6x^2 + 6x + 5$  by  $2x^2 + x + 1$ .

12.  $60(x^4 + y^4) + 91xy(x^2 - y^2)$  by  $12x^2 - 13xy + 5y^2$ .



13.  $6x^6 - 481x^5 + 79x^4 + 81x^3 - 81x^2 + 86x - 481$  by  $x - 80$ .  
 14.  $6x^6 - x^5 + 2x^4 - 2x^3 + 2x^2 + 19x + 6$  by  $3x^2 + 4x + 1$ .  
 15.  $a(a+2b)^3 - b(2a+b)^3$  by  $(a-b)^3$ .  
 16.  $(x+y)^3 + 3(x+y)^2z + 3(x+y)z^2 + z^3$  by  $(x+y)^2 + 2(x+y)z + z^2$ .  
 17.  $10x^{10} + 10x^9 + 10x^8 - 200$  by  $x^7 + x^3 - x + 1$ .  
 18.  $lmx^4 + (lm+cn)x^3 + cnx^2 + abx + ac$  by  $bx+c$ .  
 19. Multiply  $1 + \frac{2}{13}x - 18x^3$  by  $1 - \frac{1}{4}x^2 + \frac{3}{2}x^3$  and divide the product by  $1 + \frac{1}{2}x - 3x^2$ .

Find the remainders in the following cases :

20.  $(x^3 + 3x^2 + 4x + 5) \div (x - 2)$ .  
 21.  $(x^4 - 3x^2 + x - 3) \div (x - 1)$ .  
 22.  $(x^4 + 4x^3 + 6x + 8) \div (x + 2)$ .  
 23.  $(27x^4 - y^4) \div (3x - 2y)$ .  
 24.  $(9x^5 + 5x^4 - 3x^3 + 7x^2 - 5x + 8) \div (x^2 - 2x)$ .  
 25.  $(5x^4 + 90x^3 + 80x^2 - 100x + 500) \div (x + 17)$ .

**Art. viii.** The following are examples of an important use of Horner's Division :

1. Arrange  $x^3 - 6x^2 + 7x - 5$  in powers of  $x - 2$ .

2	1	-6	7	-5
		2	-8	-2
2	1	-4	-1	-7
		2	-4	
2	1	-2	-5	
		+2		
	1	0		

Hence,  $x^3 - 6x^2 + 7x - 5 = (x - 2)^3 - 5(x - 2) - 7$ , or as it is generally expressed,  $x^3 - 6x^2 + 7x - 5 = y^3 - 5y - 7$  if  $y = x - 2$ .

2. Express  $x^4 + 12x^3 + 47x^2 + 66x + 28$  in powers of  $x+3$ .

$$\begin{array}{r|rrrrr}
 -3 & 1 & 12 & 47 & 66 & 28 \\
 & & -3 & -27 & -60 & -18 \\
 \hline
 & 1 & 9 & 20 & 6 & 10 \\
 & & -3 & -18 & -6 & \\
 \hline
 & 1 & 6 & 2 & 0 & \\
 & & -3 & -9 & & \\
 \hline
 & 1 & 3 & -7 & & \\
 & & -3 & & & \\
 \hline
 & 1 & & & & \\
 & & & & & 0.
 \end{array}$$

Hence  $x^4 + 12x^3 + 47x^2 + 66x + 28 = y^4 - 7y^2 + 10$  if  $y = x+3$ .

After a few solutions have been written out in full, as in the above examples, the writing may be lessened by omitting the lines opposite the increments ( $-2$  in Ex. 1, and  $3$  in Ex. 2), the multiplication and addition being performed mentally. The last example written in this way would appear as follows:

$$\begin{array}{r|rrrrr}
 -3 & 1 & 12 & 47 & 66 & 28 \\
 & 1 & 9 & 20 & 6 & (10) \\
 & 1 & 6 & 2 & (0) & \\
 & 1 & 3 & (-7) & & \\
 & 1 & & (0) & & 
 \end{array}$$

Exercise xiv.

- 1 Express  $x^3 - 5x^2 + 3x - 8$  in powers of  $x-1$ .
2. "  $x^3 + 3x^2 + 6x + 9$  "  $x+1$ .
3. "  $x^4 - 8x^3 + 24x^2 - 32x + 97$  in powers of  $x-2$ .
4. "  $x^4 + 12x^3 + 5x^2 - 7$  "  $x+2$ .
5. "  $3x^5 - x^3 + 4x^2 + 5x - 8$  "  $x-2$ .
6. "  $x^4 - 7x^3 + 11x^2 - 7x + 10$  "  $x-1\frac{3}{4}$ .
7. "  $x^3 - 2x^2 - 4x + 9$  "  $x-\frac{2}{3}$ .
8. "  $x^3 - 9x^2y + 6xy^2 - 8y^3$  "  $x-2y$ .
9. "  $x^5 - 5x^4y + 5xy^4 - y^5$  "  $x-y$ .

- |     |   |  |   |                                |
|-----|---|--|---|--------------------------------|
| 10. | “ | $8x^3 + 12x^2y + 10xy^2 + 8y^3$                      | “ | $2x + y.$                      |
| 11. | “ | $x^3 - \frac{3}{2}x^2 + \frac{3}{8}x - \frac{5}{72}$ | “ | $\frac{1}{3}x - \frac{1}{18}.$ |
| 12. | “ | $x^4 + 8x^3 - 15x - 10$                              | “ | $x + 2.$                       |

---

## CHAPTER II

---

### SECTION I.—THE PRINCIPLE OF SYMMETRY.

---

**Art. ix.** An expression is said to be symmetrical with respect to two of its letters when these can be interchanged without altering the expression :

Thus if in  $a^3 + a^2x + ax^2 + x^3$ , we write  $x$  for  $a$ , and  $a$  for  $x$ , we get  $x^3 + x^2a + xa^2 + a^3$ , which is identical with the given expression. So, in  $x^2 + b^2x + ba + a^2x$ , if we interchange  $a$  and  $b$ , there results  $x^2 + a^2x + ab + b^2x$  which is identical with the given expression ; but it will be seen that the expression is *not* symmetrical with respect to  $x$  and  $b$ , or  $x$  and  $a$ .

An expression is symmetrical with respect to *three* of its letters  $a, b, c$ , when  $a$  can be changed into  $b$ ,  $b$  into  $c$ , and  $c$  into  $a$ , without altering the expression.

Thus  $a^3 + b^3 + c^3 - 3abc$  remains unaltered by changing  $a$  into  $b$ ,  $b$  into  $c$ , and  $c$  into  $a$ , and is therefore symmetrical with respect to these letters. So,  $a^2b + b^2a + a^2c + c^2a + b^2c + bc^2$ , and  $(a-b)^3 + (b-c)^3 + (c-a)^3$ , are each symmetrical with respect to  $a, b, c$ .

Again  $(x-a)(a-b)^2 + (x-b)(b-c)^2 + (x-c)(c-a)^2$  is symmetrical with respect to  $a, b, c$ , but not with respect to  $x$  and any of the other letters.

*Generally*, an expression is symmetrical with respect to any number of its letters  $a, b, c, \dots, h, k$ , when  $a$  can be changed into  $b$ ,  $b$  into  $c$ ,  $c$  into  $d$ , . . . .  $h$  into  $k$ , and  $k$  into  $a$ , without altering the expression.

A symmetric function of several letters is frequently represented by writing each *type-term* once, preceded by the letter  $\Sigma$ ; thus for  $a+b+c+\dots+l$  we write  $\Sigma a$ , and for  $ab+ac+ad+\dots+bc+bd+\dots$  (*i. e.* the sum of the products of every pair of the letters considered) we write  $\Sigma ab$ .

Exercise xv.

Write the following in full :

1.  $\Sigma a^2b$ ,  $\Sigma (a-b)^2$ ,  $\Sigma a(b-c)$ ,  $\Sigma ab(x-c)$ ,  $\Sigma a^3b^2c$ ,  $\Sigma (a+b) \times (c-a)(c-b)$ ,  $\Sigma \{(a+c)^2 - b^2\}$ , and  $\Sigma a(b+c)^2$ , each for  $a, b, c$ .

2.  $\Sigma abc$ ,  $\Sigma a^2b$ ,  $\Sigma a^2bc$ ,  $\Sigma (a-b)$ , and  $\Sigma a^2(a-b)$ , each with respect to  $a, b, c, d$ .

Show that the following are symmetrical :

3.  $(x+a)(a+b)(b+x) + abx$ , with respect to  $a$  and  $b$ .

4.  $(a+b)^2 + (a-b)^2$  with respect to  $a$  and  $b$ , and also with respect to  $a$  and  $-b$ .

5.  $(ab-xy)^2 - (a+b-x-y)\{ab(x+y) - xy(a+b)\}$  with respect to  $a$  and  $b$ , and also with respect to  $x$  and  $y$ .

6.  $a^2(b-c) - b^2(a-c) - c^2(b-a)$  with respect to  $a, b, c$ .

7.  $(ac+bd)^2 + (bc-ad)^2$  with respect to  $a^2$  and  $b^2$ , and also with respect to  $c^2$  and  $d^2$ .

8.  $x^6 + y^6 + 3xy(x^2 + xy + y^2)$  with respect to  $x$  and  $y$ .

9.  $\{x^3 - y^3 + 3xy(2x+y)\}^3 + \{y^3 - x^3 + 3xy(2y+x)\}^3$  with respect to  $x$  and  $y$ .

10.  $a(a+2b)^3 + b(b+2a)^3$  with respect to  $a$  and  $b$ , and also with respect to  $a$  and  $-b$ .

11.  $ab[\{(a+c)(b+c) + 2c(a+b)\}^2 - (a-c)^2(b-c)^2]$  with respect to  $a, b, c$ .

12.  $a^2b^2 + b^2c^2 + c^2a^2 + 2abc(a+b+c)$  with respect to  $ab, bc, ca$

With respect to what letters are the following symmetrical ?

13.  $xyz + 5xy + 2(x^2 + y^2)$ .

14.  $2(a^2x^2 + b^2y^2) - 2ab(xy + yz + zx)$ .

15.  $(f^2 - h^2)^2 + 4g^2(f+h)^2 + (2fh - 2g^2)^2$ .

16.  $(x+y)(x-z)(y-z) - xyz$ .  
 17.  $a^2b^2 + b^2c^2 + c^2a^2 - 2abc(a+b-c)$ .  
 18.  $x^6 - y^6 + z^6 - 3(x^2 - y^2)(y^2 - z^2)(z^2 + x^2)$ .  
 19.  $(a+b)^2 + (a+c)^2 + (b-c)^4$ .  
 20.  $(a+b)^4 + (a-c)^4 + (b+c)^4 + (a+c)^4$ .  
 21.  $(a+b)^4 + (a-c)^4 + (b+c)^4 + (a+c)^4 + (c-b)^4$ .

Select the type-terms in :

22.  $a^2 + 2ab + b^2 + 2bc + c^2 + 2ca$   
 23.  $a(b^2 - c^2) + b(c^2 - a^2) + c(a^2 - b^2) + (a+b)(b+c)(c+a)$ .  
 24.  $a(b+c)^2 + b(c+a)^2 + c(a+b)^2 - 12abc$ .

Write down the *type-terms* in :

25.  $(x+y)^5, (x-y)^5, (x+y)^5 - x^5 - y^5$ .  
 26.  $(x+y)^7 + (x-y)^7, (x+y)^7 - (x-y)^7$ .  
 27.  $(x+y+z)^4, (x-y-z)^4$ .  
 28.  $(a+b+c+d)^4, (a^2 + b^2 + c^2 + d^2)^2$ .  
 29.  $(a+b)^3 + (b+c)^3 + (c+a)^3$ .

Art. x. In reducing an algebraic expression from one form to another, advantage may be taken of the principle of symmetry : for, it will be necessary to calculate only the *type-terms*, and the others may be written down from these.

#### EXAMPLES.

1. Find the expansion of  $(a+b+c+d+e+\&c.)^2$

This expression is symmetrical with respect to  $a, b, c, \&c.$  ; hence the expansion also must be symmetrical, and as it is a product of *two* factors, it can contain only the squares  $a^2, b^2, c^2, \&c.$ , and the products in pairs,  $ab, ac, ad . . . bc, bd, \&c.$  ; so that  $a^2$  and  $ab$  are *type-terms*.

Now  $(a+b)^2 = a^2 + 2ab + b^2$  ; and the addition of terms involving  $a, b, c, \&c.$ , will not alter the terms  $a^2 + 2ab$ , but will merely give additional terms of the same type. Hence from symmetry we get

$$\begin{aligned}
 (a+b+c+d+e+\&c.)^2 = a^2 + 2ab + 2ac + 2ad + 2ae + \dots \\
 + b^2 + 2bc + 2bd + 2be + \dots \\
 + c^2 + 2cd + 2ce + \dots \\
 + d^2 + 2de + \dots \\
 + e^2 + \dots
 \end{aligned}$$

This may be compactly written

$$(\Sigma a)^2 = \Sigma a^2 + 2\Sigma ab.$$

2. Expand  $(a+b)^3$ .

This has been found by actual multiplication—see formula [5]—but we may also proceed as follows :

(1) The expression is of *three* dimensions, and is symmetrical with respect to  $a$  and  $b$ .

(2) The type-terms are  $a^3, a^2b$ .

Hence  $(a+b)^3 = a^3 + b^3 + n(a^2b + b^2a)$ , where  $n$  is *numerical*.

To find the value of  $n$ , put  $a = b = 1$ , and we have  $(1+1)^3 = 1+1+n(1+1)$ ;  $\therefore n=3$ .

3. Expand  $(x+y+z)^3$ .

This is of three dimensions, and is symmetrical with respect to  $x, y, z$ . We have

$$\begin{aligned}
 (x+y+z)^3 &= \{(x+y)+z\}^3 = (x+y)^3 + \&c. \\
 &= x^3 + 3x^2y + \&c., \text{ which are type-terms, the only other possible} \\
 &\text{type-term being } xyz.
 \end{aligned}$$

Now, since the expression contains  $3x^2y$ , it must also contain  $3x^2z$ , that is, it must contain  $3x^2(y+z)$ . Hence

$$\begin{aligned}
 (x+y+z)^3 &= x^3 + 3x^2(y+z) \\
 &\quad + y^3 + 3y^2(z+x) \\
 &\quad + z^3 + 3z^2(y+x) \\
 &\quad + n(xyz), \text{ where } n \text{ is numerical, and}
 \end{aligned}$$

may be found by putting  $x=y=z=1$  in the last result, giving

$$\begin{aligned}
 (1+1+1)^3 &= 1+1+1 + 3(1+1) + 3(1+1) + 3(1+1) + n; \\
 \therefore n &= 6.
 \end{aligned}$$

4. Similarly we may shew that

$$\begin{aligned}(a+b+c+d)^3 = & a^3 + 3a^2(b+c+d) + 6bcd \\ & + b^3 + 3b^2(c+d+a) + 6cda \\ & + c^3 + 3c^2(d+a+b) + 6dab \\ & + d^3 + 3d^2(a+b+c) + 6abc.\end{aligned}$$

5. Expand  $(a+b+c+\&c.)^3$ .

The type-terms are  $a^3, a^2b, abc$ .

Expanding  $(a+b+c)^3$ , we get  $a^3 + 3a^2b + 6abc + \&c$ .

Hence by symmetry we have

$$(\Sigma a)^3 = \Sigma a^3 + 3\Sigma a^2b + 6\Sigma abc.$$

6. Simplify  $(a+b-2c)^2 + (b+c-2a)^2 + (c+a-2b)^2$ .

This expression is symmetrical, involving terms of the types  $a^2$  and  $ab$ . Now  $a^2$  occurs with 1 as a coefficient in the first square, with 4 as a coefficient in the second square, and with 1 as a coefficient in the third square, and hence  $6a^2$  is one type-term of the result:  $ab$  occurs with 2 as a coefficient in the first square, with  $-4$  as a coefficient in the second square, and with  $-4$  as a coefficient in the third square, and hence  $-6ab$  is the second type-term in the result: hence the total result is  $6(a^2 + b^2 + c^2 - ab - bc - ca)$ .

7. Simplify  $(x+y+z)^3 + (x-y-z)^3 + (y-z-x)^3 + (z-x-y)^3$ .

This is symmetrical with respect to  $x, y, z$ ; and the type-terms are  $x^3, 3x^2y, 6xyz$ :

(1)  $x^3$  occurs in each of the first two cubes, and  $-x^3$  in each of the second two cubes.  $\therefore$  there are no terms of the type  $x^3$  in the result.

(2)  $3x^2y$  occurs in the *first* and *third* cubes, and  $-3x^2y$  in the second and fourth,  $\therefore$  there are no terms of this type in the result.

(3)  $6xyz$  occurs in each of the four cubes,  $\therefore 24xyz$  is the total result.

8. Prove  $(a^2 + b^2 + c^2 + d^2)(w^2 + x^2 + y^2 + z^2) - (aw + bx + cy + dz)^2 = (ax - bw)^2 + (ay - cw)^2 + (az - dw)^2 + (by - cx)^2 + (bz - dx)^2 + (cz - dy)^2$ .

The left hand member (considered as given) is symmetrical with respect to the pairs of letters,  $a$  and  $w$ ,  $b$  and  $x$ ,  $c$  and  $y$ ,  $d$  and  $z$ , that is, any two pairs may be interchanged without affecting the expression. As the expression is only of the second degree in these pairs, no term can involve three pairs as factors; hence the type-terms may be obtained by considering all the terms involving  $a, b, w, x$ ; these are  $a^2w^2, a^2x^2, b^2w^2, b^2x^2, -a^2w^2, -b^2x^2, -2abwx$ , and are the terms of  $(ax-bw)^2$  which is consequently a type-term. From  $(ax-bw)^2$  we derive the five other terms of the second member by merely changing the letters.

9. Prove that

$(x^2 - yz)^3 + (y^2 - zx)^3 + (z^2 - xy)^3 - 3(x^2 - yz)(y^2 - zx)(z^2 - xy)$  is a complete square.

The expression will remain symmetrical if  $(x^2 - yz)(y^2 - zx)(z^2 - xy)$ , instead of being multiplied by  $-3$ , be subtracted from each of the preceding terms, thus giving

$$\begin{aligned} & (x^2 - yz) \{ (x^2 - yz)^2 - (y^2 - zx)(z^2 - xy) \} \\ & + (y^2 - zx) \{ (y^2 - zx)^2 - (z^2 - xy)(x^2 - yz) \} \\ & + (z^2 - xy) \{ (z^2 - xy)^2 - (x^2 - yz)(y^2 - zx) \} \\ & = (x^2 - yz)x(x^3 + y^3 + z^3 - 3xyz) \\ & + \&c. \\ & + \&c. \\ & = (x^3 + y^3 + z^3 - 3xyz)(x^3 + y^3 + z^3 - 3xyz). \end{aligned}$$

Exercise xvi.

Simplify the following:

1.  $(a+b+c)^2 + (a+b-c)^2 + (b+c-a)^2 + (c+a-b)^2$
2.  $(a-b-c)^2 + (b-a-c)^2 + (c-a-b)^2$ .
3.  $(a+b+c-d)^2 + (b+c+d-a)^2 + (c+d+a-b)^2 + (d+a+b-c)^2$ .
4.  $(a+b+c)^3 - a(b+c-a) - b(a+c-b) - c(a+b-c)$ .
5.  $(x+y+z+n)^2 + (x-y-z+n)^2 + (x-y+z-n)^2 + (x+y-z-n)^2$ .
6.  $(a+b+c)^3 + (a+b-c)^3 + (b+c-a)^3 + (c+a-b)^3$ .



$$7. (x-2y-3z)^2 + (y-2z-3x)^2 + (z-2x-3y)^2.$$

$$8. (ma+nb+rc)^3 - (ma+nb-rc)^3 - (nb+rc-ma)^3 - (rc+ma-nb)^3.$$

$$9. a(b+c)(b^2+c^2-a^2) + b(c+a)(c^2+a^2-b^2) + c(a+b)(a^2+b^2-c^2).$$

$$10. (ab+bc+ca)^2 - 2abc(a+b+c).$$

Prove the following :

$$11. (ax+by+cz)^2 + (bx+cy+az)^2 + (cx+ay+bz)^2 + (ax+cy+bz)^2 + (cx+by+az)^2 + (bx+ay+cz)^2 \\ = 2(a^2+b^2+c^2)(x^2+y^2+z^2) + 1(ab+bc+ca)(xy+yz+zx).$$

$$12. (a+b+c)^4 + (b+c-a)^4 + (c+a-b)^4 + (a+b-c)^4 \\ = 4(a^4+b^4+c^4) + 24(a^2b^2+b^2c^2+c^2a^2).$$

$$13. (a+b+c)^4 = \Sigma a^4 + 4\Sigma a^3b + 6\Sigma a^2b^2 + 12\Sigma a^2bc.$$

$$14. (\Sigma a)^4 = \Sigma a^4 + 4\Sigma a^3b + 6\Sigma a^2b^2 + 12\Sigma a^2bc + 24\Sigma abcd.$$

$$15. (a^2+b^2+c^2)^3 + 2(ab+bc+ca)^3 - 3(a^2+b^2+c^2) \times (ab+bc+ca)^2 = (a^3+b^3+c^3-3abc)^2.$$

$$16. (a-b)^2(b-c)^2 + (b-c)^2(c-a)^2 + (c-a)^2(a-b)^2 = (a^2+b^2+c^2-ab-ac-bc)^2.$$

$$17. (2a-b-c)^2(2b-c-a)^2 + (2b-c-a)^2(2c-a-b)^2 + (2c-a-b)^2(2a-b-c)^2 = 9(a^2+b^2+c^2-ab-bc-ca)^2$$

$$18. (ar^2+2brs+cs^2)(ax^2+2bxy+cy^2) - \{arx+b(ry+sx)+csy\}^2 = (ac-b^2)(ry-sx)^2.$$

$$19. (a^2+ab+b^2)(c^2+cd+d^2) = (ac+ad+bd)^2 + (ac+ad+bd)(bc-ad) + (bc-ad)^2.$$

20. Shew that there are two ways in which the given product in the last example can be expressed in the form  $p^2+pq+q^2$ , and two ways in which it can be expressed in the form  $p^2-pq+q^2$ .

$$21. 6(w^2+x^2+y^2+z^2)^2 = (w+x)^4 + (w-x)^4 + (w+y)^4 + (w-y)^4 + (w+z)^4 + (w-z)^4 + (x+y)^4 + (x-y)^4 + (x+z)^4 + (x-z)^4 + (y+z)^4 + (y-z)^4.$$

$$22. \frac{1}{2}\{(a+b+c)^5 + (a-b-c)^5 + (b-c-a)^5 + (c-a-b)^5\} = \frac{1}{2}\{(a+b+c)^3 + (a-b-c)^3 + (b-c-a)^3 + (c-a-b)^3\} \times \frac{1}{2}\{(a+b+c)^2 + (a-b-c)^2 + (b-c-a)^2 + (c-a-b)^2\}.$$

SECTION II.—THEORY OF DIVISORS.

Any expression which can be reduced to the form  $ax^n + bx^{n-1} + cx^{n-2} + \dots + hx + k$ , in which  $n$  is a positive integer and  $a, b, c, \dots, h, k$  are independent of  $x$ , is called a **POLYNOME** in  $x$  of degree  $n$ .

The expressions  $f(x)^n, F(x)^n, \phi(x)^n$ , are used as general symbols for polynomes; the index  $n, m$ , indicates the degree of the polynome.

**Theorem I.** If the polynome  $f(x)^n$  be divided by  $x-a$ , the remainder will be  $f(a)^n$ .

Cor. 1.  $f(x)^n - f(a)^n$  is always exactly divisible by  $x-a$ .

(Particular case:  $x^n - a^n$  is always exactly divisible by  $x-a$ ).

Cor. 2. If  $f(a)^n = 0, f(x)^n$  is exactly divisible by  $x-a$ , i.e.,  $f(x)^n$  is an algebraic multiple of  $x-a$ .

Cor. 3. If the polynome  $f(x)^n$  on division by the polynome  $\phi(x)^m$  leave a remainder independent of  $x$ , such remainder will be the value of  $f(x)^n$  when  $\phi(x)^m = 0$ .

EXAMPLES.—THEOREM I.

1. Find the remainder when  $x^5 - 7x^4 + 13x^3 - 16x^2 + 9x - 12$  is divided by  $x-5$ .

The remainder will be the value of the given polynome when 5 is substituted for  $x$ . (See Art. III.).

$$\begin{array}{r|rrrrrr} & 1 & -7 & +13 & 16 & +9 & -12 \\ 5 & & 5 & -10 & 15 & -5 & 20 \\ \hline & 1 & -2 & 3 & -1 & 4 & 8 \end{array}$$

Hence the remainder is 8.

2. Find the remainder when  $(x-a)^3 + (x-b)^3 + (a+b)^3$  is divided by  $x+a$ .

For  $x$  substitute  $-a$ , then  $(-2a)^3 + (-a-b)^3 + (a+b)^3 = -8a^3$ .

3. Find the remainder when  $x^3 + a^3 + b^3 + (x+a)(x+b)(a+b)$  is divided by  $x+a+b$ .

For  $x$  substitute  $-(a+b)$  and we get

$$-(a+b)^3 + a^3 + b^3 + ab(a+b) = -2ab(a+b). \quad \text{See Formula [6].}$$

4. Find the remainder when  $(x^2 + 2ax - 2a^2)^3(x^3 - 2ax - 2a^2) + 32(x-a)^4(x+a)^4$  is divided by  $x^2 - 2a^2$ .

$x^2 - 2a^2$  may be struck out wherever it appears.

This reduces the dividend to

$$(2ax)^3(-2ax) + 32(x-a)^4(x+a)^4 = -16a^4x^4 + 32(x^3 - a^2)^4.$$

In this substitute  $2a^2$  for  $x^2$  and it becomes

$$-64a^8 + 32a^8 = -32a^8,$$

which is the required remainder.

### Exercise xvii.

1. Find the remainder when  $3x^4 + 60x^3 + 54x^2 - 60x + 58$  is divided by  $x + 19$ .

2. Find the remainder when  $px^3 - 3qx^2 + 3rx - s$  is divided by  $x - a$ .

3. What number added to  $4x^5 + 34x^4 + 58x^3 + 21x^2 - 123x - 41$  will give a sum exactly divisible by  $2x + 13$ ?

4. What number taken from  $10x^{10} - 20x^8 - 10x^6 - 89x^4 - 89x^2 + 20$  will leave a remainder exactly divisible by  $10x^2 - 11$ ?

Find the remainders from the following divisions:

5.  $(x+1)^5 - x^5 \div x+1$ , and  $(x+a+3)^3 - (x+a+1)^3 \div x+2$ .

6.  $x^n + y^n \div x-y$ ;  $x^{2n} + y^{2n} \div x+y$ ;  $x^{2n+1} + y^{2n+1} \div x+y$ .

7.  $(x+1)^3 + x^3 + (x-1)^3 \div x-2$ .

8.  $(x-a)^3(x+a)^3 + (x^2 - 2b^2)^3 \div x^2 + b^2$ .

9.  $(x^2 + ax + a^2)(x^2 - ax + a^2) - (x^2 - 3ax + 2a^2)(x^2 + 3ax + 2a^2) \div x^2 + 2a^2$ .

10.  $(9a^2 + 6ab + 4b^2)(9a^2 - 6ab + 4b^2)(81a^4 - 36a^2b^2 + 16b^4) \div (3a - 2b)^2$ .

11.  $a^2(x-a)^3 + b^2(x-b)^3 \div x-a-b$ .

12.  $(ax+by)^3 + a^3y^3 + b^3x^3 - 3abxy(ax+by) \div (a+b)(x+y)$ .

13.  $x^3 + a^3 + b^3 - 3abx \div x-a+b$ ; also  $\div x+a-b$  also  $\div x-a-b$ .

14. Any polynome divided by  $x-1$  gives for remainder the sum of the coefficients of the terms.

## EXAMPLES.—COR. 1.

1.  $x^5+y^5$  is exactly divisible by  $x+y$ .

In " $x^5-a^5$  is exactly divisible by  $x-a$ ," substitute  $-y$  for  $a$ .

2.  $mx^3-px^2+qx+m+p+q$  is exactly divisible by  $x+1$ .

This may be written

$\{nx^3-px^2+qx\} - \{m(-1)^3 - p(-1)^2 + q(-1)\}$  is exactly divisible by  $x - (-1)$ .

3.  $(x^2+6xy+4y^2)^5 + (x^2+2xy+4y^2)^5$  is exactly divisible by  $(x+2y)^2$ . For  $(x^2+6xy+4y^2)^5 - (-x^2-2xy-4y^2)^5$  is exactly divisible by  $(x^2+6xy+4y^2) - (-x^2-2xy-4y^2)$ , which is  $2(x^2+4xy+4y^2) = 2(x+2y)^2$ .

## Exercise xviii.

Prove that the following are cases of exact division :

1.  $x^{2n+1}+y^{2n+1} \div x+y$ ;  $x^{2n}-y^{2n} \div x+y$ .
2.  $x^{12}+y^{12} \div x^4+y^4$ ;  $x^{30}+y^{30} \div x^6+y^6$ ; also  $\div x^{10}+y^{10}$ ; also  $\div x^2+y^2$ .
3.  $(ax+by)^5 + (bx+ay)^5 \div (a+b)(x+y)$ .
4.  $(ax+by+cz)^3 - (bx+cy+az)^3 \div (a-b)x + (b-c)y + (c-a)z$ .
5.  $(2y-x)^n - (2x-y)^n \div 3(y-x)$ .
6.  $(2y-x)^{2n+1} + (2x-y)^{2n+1} \div y+x$ .
7.  $(my-nx)^5 - (mx-ny)^5 \div (m+n)(y-x)$ .
8.  $(x+y)^6 + (x-y)^6 \div 2(x^2+y^2)$ .
9.  $(x^2+xy+y^2)^3 + (x^2-xy+y^2)^3 \div 2(x^2+y^2)$ .
10.  $(a+b)^9 - (a-b)^9 \div 2b(3a^2+b^2)$ .
11.  $(x^2+5bx+b^2)^7 + (x^2-bx+b^2)^7 \div 2(x+b)^2$ .
12.  $(a+b)^{4n+2} + (a-b)^{4n+2} \div 2(a^2+b^2)$ .
13.  $\{x^3+3xy(x-y)-y^3\}^3 + \{x^3-3xy(x-y)-y^3\}^3 \div 2(x-y)^3$ .
14.  $3x^3-5x^2+4x-2 \div x-1$ .

15. Any polynome in  $x$  is divisible by  $x-1$  when the sum of the coefficients of the terms is zero.

16. Any polynome in  $x$  is divisible by  $x+1$ , when the sum of the coefficients of the even powers of  $x$  is equal to the sum of the coefficients of the odd powers. (The constant term is included among the coefficients of the even powers).

EXAMPLES.—COR. 2.

1. Show that  $a(a+2b)^3 - b(2a+b)^3$  is exactly divisible by  $a+b$ . By Cor. 2, the substitution of  $-b$  for  $a$  must cause the polynome to vanish.

$$\text{Substituting; } a(a-2a)^3 + a(2a-a)^3 = -a^4 + a^4 = 0.$$

2. Show that  $(ab-xy)^2 - (a+b-x-y)\{ab(x+y) - xy(a+b)\}$  is exactly divisible by  $(x-a)(y-a)$ , also by  $(x-b)(y-b)$ .

For  $x$  substitute  $a$  and the expression becomes

$$(ab-ay)^2 - (b-y)\{ab(a+y) - ay(a+b)\} = \\ a^2(b-y)^2 - (b-y)\{a^2(b-y)\} = 0.$$

The expression is, therefore, exactly divisible by  $x-a$ . But it is symmetrical with respect to  $x$  and  $y$ , hence it is divisible by  $y-a$ , and as  $x-a$  and  $y-a$  are independent factors, the expression is exactly divisible by  $(x-a)(y-a)$ . Again, the given expression is symmetrical with respect to  $a$  and  $b$ , hence, making the interchange of  $a$  and  $b$ , the expression is seen to be divisible by  $(x-b)(y-b)$ .

3. Show that  $6(a^5 + b^5 + c^5) - 5(a^3 + b^3 + c^3)(a^2 + b^2 + c^2)$  is exactly divisible by  $a+b+c$ .

For  $a$  substitute  $-(b+c)$  and the result which would be the remainder were the division actually performed, must vanish.

$$6\{- (b+c)^5 + b^5 + c^5\} - 5\{- (b+c)^3 + b^3 + c^3\}\{(b+c)^2 + b^2 + c^2\} \\ = 6\{- (b+c)^5 + b^5 + c^5\} + 30bc(b+c)(b^2 + bc + c^2). \text{ See [1] and [6].}$$

The expansion being of the 5th degree, and symmetrical in  $b$  and  $c$ , it will be sufficient to show that the coefficients of  $b^5$ ,  $b^4c$ ,  $b^3c^2$  vanish, the coefficients of  $b^2c^3$ ,  $bc^4$ ,  $c^5$  being the coefficients

of the former terms in reverse order. Calculating the coefficients of these type-terms we get

$$6\{-5b^4c - 10b^3c^2 - \dots\} + 30(b^4c + 2b^3c^2 + \dots),$$

which evidently vanishes. Hence the truth of the proposition.

$$4. \text{ If } a+b+c=0, \frac{1}{3}(a^5+b^5+c^5) = \frac{1}{3}(a^2+b^2+c^2) \cdot \frac{1}{3}(a^3+b^3+c^3).$$

In the last example it has been proved that the *difference* of the quantities here declared to be equal, is a multiple of  $a+b+c$ , *i.e.*, in this case, a multiple of zero. Hence under the given condition they are equal.

### Exercise xix.

Prove that the following are cases of exact division :

1.  $(ax-by)^3 + (bx-ay)^3 - (a^3+b^3)(x^3-y^3) \div a, b, x, y, a+b, x-y.$
2.  $ax^3 - (a^2+b)x^2 + b^2 \div ax-b.$  (Substitute  $ax$  for  $b$ .)
3.  $\begin{cases} (ax+by)^2 - (a-b)(x+z)(ax+by) + (a-b)^2xz \div x+y. \\ (ax-by)^2 - (a+b)(x+z)(ax-by) + (a+b)^2xz \div x+y. \end{cases}$
4.  $6a^3x^2 - 4ax^3 - 10axy - 3a^2xy + 2x^2y + 5y^2 \div 2ax-y.$
5.  $1 \cdot 2a^4x - 16 \cdot 32a^3x^2 + 4 \cdot 8a^2x^3 + 9ax^4 - x^5 \div 6ax - 2x^2.$
6.  $x^8 + x^6y^2 + x^2y + y^3 \div x^6 + y.$
7.  $(c-d)a^2 + 6(bc-bd)a + 9(b^2c-b^2d) \div a+3b.$
8.  $x(x - \frac{1}{12}y)^5 + y(\frac{1}{12}x - y)^5 \div x-y.$
9.  $a(a+2b)^3 - b(b+2a)^3 \div a-b, \text{ also } \div a+b.$
10.  $a^5 - 2a^4b + a^3b^2 + a^2x^3 - 2abx^3 + 3^2x^3 \div (a-b)(x+a).$
11.  $a(b-c)^3 + b(c-a)^3 + c(a-b)^3 \div (a-b), (b-c), (c-a).$
12.  $a^3(b-c) + b^3(c-a) + c^3(a-b) \div (a-b), (b-c), (c-a).$
13.  $a^4(b-c) + b^4(c-a) + c^4(a-b) \div (a-b), (b-c), (c-a).$
14.  $(a-b)^2(c-d)^2 + (b-c)^2(d-a)^2 - (d-b)^2(a-c)^2 \div (a-b), (b-c), (c-d), (d-a).$
15.  $\{(a-b)^2 + (b-c)^2 + (c-a)^2\} \{(a-b)^2c^2 + (b-c)^2a^2 + (c-a)^2b^2\} - \{(a-b)^2c + (b-c)^2a + (c-a)^2b\}^2 \div (a-b), (b-c), (c-a).$
16.  $(x+y)(y+z)(z+x) + xyz \div x+y+z.$

17.  $ab(a^2 - b^2) + bc(b^2 - c^2) + ca(c^2 - a^2) \div a + b + c.$   
 18.  $(ab - bc - ca)^2 - a^2b^2 - b^2c^2 - c^2a^2 \div a + b - c.$   
 19.  $(a + 2b)^3 + (2b - 3c)^3 - (3c - a)^3 + a^3 + 8b^3 - 27c^3 +$   
 $a + 2b - 3c.$   
 20.  $a^3b^3 + b^3c^3 + c^3a^3 - 3a^2b^2c^2 \div ab + bc + ca.$

EXAMPLES.—COR. 3 AND 2.

1. Find the value of  $4x^5 + 9x^3 - 5x^2 + 23x + 6$  when  $2x^2 = 3x + 4$ .

Since  $2x^2 - 3x + 4 = 0$ , we have simply to find the remainder on division by  $2x^2 - 3x + 4$ , and if it is independent of  $x$ , it is the value sought, Cor. 3.

$$\begin{array}{r|rrrrrr} & 4 & 0 & 9 & -5 & 23 & 6 \\ 3 & & 6 & 9 & 15 & -3 & \\ -4 & & & -8 & -12 & -20 & 4 \\ \hline 2 & 2 & 3 & 5 & -1 & 0 & 10 \end{array}$$

Hence the required value is 10.

2. What value of  $c$  will make  $x^3 - 5x^2 + 7x - c$  exactly divisible by  $x - 2$ .

If 2 be substituted for  $x$ , the remainder must vanish, Cor. 2.

$$\begin{array}{r|rrrr} 2 & 1 & -5 & 7 & -c \\ & & 2 & -6 & 2 \\ \hline & 1 & -3 & 1 & 2-c \end{array}$$

Hence  $2 - c = 0$ , or  $c = 2$ .

3. What value of  $c$  will make  $6x^5 - 5x^4 + cx^3 - 20x^2 + 19x - 5$  vanish when  $2x^2 = 3x - 1$ ?

By Cor. 3, the remainder must vanish when the given poly-  
 nome is divide by  $2x^2 - 3x + 1$ . We may divide at once and find,  
 if possible, a value of  $c$  that will make both terms of the remainder  
 vanish, or we may first express  $cx^3$  in lower terms in  $x$ , and  
 then divide and find the required value of  $c$  from the remainder.

1st. Method, (see page 28),

$$\begin{array}{r|rrrrrr} & 6 & -10 & 4c & -160 & 304 & -160 \\ 3 & & 18 & 24 & 12c+36 & 36c-420 & \\ -2 & & & -12 & -16 & -8c-24 & -24c+280 \\ \hline & 6 & 8 & 4c+12 & 12c-140 & 28c-140 & -24c+120 \end{array}$$

Hence  $28c = 140$  and  $24c = 120$ . Both of these are satisfied by  $c = 5$ .

2nd Method.  $x^3 = \frac{1}{4}x(3x-1) = \frac{3}{4}x^2 - \frac{1}{4}x = \frac{3}{4}(3x-1) - \frac{1}{4}x = 2\frac{1}{4}x - \frac{3}{4} - \frac{1}{4}x = 1\frac{3}{4}x - \frac{3}{4}$ ;  $\therefore cx^3 = 1\frac{3}{4}cx - \frac{3}{4}c$ .

Substituting for  $cx^3$  in the given polynome it becomes

$$6x^5 - 5x^4 - 20x^2 + (1\frac{3}{4}c + 19)x - \frac{3}{4}c - 5.$$

Divide and apply Cor. 3.

3	6	-10	0	-160	$28c + 304$	$-24c - 160$
-2		18	24	36	-420	
			-12	-16	-24	280
	6	8	12	-140;	$28c - 140$	$-24c + 120$

We thus obtain the same remainder as by the former method, and consequently the same result. A comparison of the two methods shews that they are but slightly different in form, but the second method shows rather more clearly that  $c$  need not be introduced into the dividend at all, but the proper multiples of it found by the preliminary reduction can be added to or taken from the numerical remainder, and the "true remainder" be thus found, and  $c$  determined from it.

### Exercise xx.

Find the value of

1.  $x^4 - 3x^3 + 4x^2 - 3x + 4$ , given  $x^2 = x - 1$ .
2.  $x^5 - 2x^4 - 4x^3 + 13x^2 - 11x - 10$ , given  $(x-1)^2 = 2$ .
3.  $2x^5 - 7x^4 + 12x^3 - 11x^2 + 2x - 5$  given  $(x-1)^2 + 2 = 0$ .
4.  $3x^6 + 11x^5 + 10x^3 + 7x^2 + 2x + 3$  given  $x^2 + 3x^2 - 2x + 5 = 0$ .
5.  $6x^7 + 9x^6 - 16x^4 - 5x^3 - 12x^2 - 6x + 60$  given  $3x^4 + x - 4 = 0$ .

What values of  $c$  will make the following polynomes vanish under the given conditions.

6.  $x^4 + 13x^3 + 26x^2 + 52x + 8c$ , given  $x + 11 = 0$ .
7.  $x^4 - 2x^3 - 9x^2 + 2cx - 14$ , given  $3x + 7 = 0$ .
8.  $x^4 - 4x^3 - x^2 + 16x + 6c$ , given  $x^2 = x + 6$ .
9.  $2x^4 - 10x^2 + 4cx + C$ , given  $x^2 + 3 = 3x$ .
10.  $2x^4 + x^3 - 7cx^2 + 11x + 10$ , given  $2c = 5$ .



11.  $4x^4 + cx^2 + 110x - 105$ , given  $2x^2 - 5x + 15 = 0$ .  
 12.  $3x^5 - 16x^4 + cx^3 - 5x^2 - 114x + 200$ , given  $x^2 = 3x - 4$ .  
 13. What values of  $p$  and  $q$  will make  $x^4 + 2x^3 - 10x^2 - px + q$  vanish, given  $x^2 = 3(x - 1)$ ?  
 14. What values of  $p$  and  $q$  will make  $a^{12} - 5a^{10} + 10a^8 - 15a^6 + 29a^4 - pa^2 + q$  vanish, given  $(a^2 - 2)^2 = a^2 - 3$ ?

**Theorem II.** If the polynome  $f(x)^n$  vanish on substituting for  $x$  each of the  $n$  (different) values  $a_1, a_2, a_3 \dots a_n$

$$f(x)^n = A(x - a_1)(x - a_2)(x - a_3) \dots (x - a_n)$$

in which  $A$  is independent of  $x$  and consequently is the coefficient of  $x^n$  in  $f(x)^n$ .

Cor. If  $f(x)^n$  and  $\varphi(x)^m$  both vanish for the same  $m$  different values of  $x$ ,  $f(x)^n$  is algebraically divisible by  $\varphi(x)^m$ .

#### EXAMPLES.

1.  $x^3 + ax^2 + bx + c$  will vanish if 2, or 3, or  $-4$  be substituted for  $x$ , determine  $a, b, c$ .

The coefficient of the highest power of  $x$  is 1;

$$\therefore x^3 + ax^2 + bx + c = (x - 2)(x - 3)(x + 4) = x^3 - x^2 - 14x + 24.$$

$$\therefore a = -1; b = -14; c = 24.$$

2.  $x^3 + bx^2 + cx + d$  will vanish if  $-3$  or 2, or 5 be substituted for  $x$ ; determine its value if 3 be substituted for  $x$ .

The given polynome  $= (x + 3)(x - 2)(x - 5)$ ;

$$\therefore \text{the required value is } (3 + 3)(3 - 2)(3 - 5) = -12.$$

3.  $ax^3 + 3bx^2 + 3cx + d$  will vanish if for  $x$  be substituted  $-3$ , or  $\frac{1}{3}$ , or  $1\frac{1}{2}$ , but it becomes 45 if for  $x$  there be substituted 8; determine the values of  $a, b, c, d$ .

The coefficient of the highest power of  $x$  is  $a$ ;

$$\therefore ax^3 + 3bx^2 + 3cx + d = a(x + 3)(x - \frac{1}{3})(x - 1\frac{1}{2})$$

$$\therefore a(3 + 3)(3 - \frac{1}{3})(3 - 1\frac{1}{2}) = 45; \therefore a = 2.$$

$$\therefore 2x^3 + 3bx^2 + 3cx + d = 2(x + 3)(x - \frac{1}{3})(x - 1\frac{1}{2})$$

$$\therefore b = \frac{2}{3}, c = -3\frac{1}{2}, d = 4\frac{1}{2}$$

4. If  $x^3 + px^2 + qx + r$  vanish for  $x = a$  or  $b$ , or  $c$ , determine  $p$ ,  $q$ , and  $r$  in terms of  $a$ ,  $b$ ,  $c$ .

$$\begin{aligned} x^3 + px^2 + qx + r &= (x-a)(x-b)(x-c) \\ &= x^3 - (a+b+c)x^2 + (ab+bc+ca)x - abc \\ \therefore p &= -(a+b+c) \quad \text{or} \quad -\Sigma a. \\ q &= ab+bc+ca \quad \text{or} \quad \Sigma ab \\ r &= -abc \quad \text{or} \quad -\Sigma abc. \end{aligned}$$

5. If  $x^3 + px^2 + qx + r$  vanish for  $x = a$ , or  $b$ , or  $c$ , determine the polynome that will vanish for  $x = b+c$ , or  $c+a$ , or  $a+b$ .

Since  $x^3 + px^2 + qx + r$  vanishes for  $x = a$  or  $b$  or  $c$ ,  
 $x^3 - px^2 + qx - r$  will vanish for  $x = -a$  or  $-b$  or  $-c$ ,  
 and  $-p = a+b+c$ ;

But the required polynome will vanish for

$$x = -p - a, \text{ or } -p - b, \text{ or } -p - c;$$

that is, for  $x+p = -a$ , or  $-b$ , or  $-c$ .

$$\begin{aligned} \text{Hence it is } (x+p)^3 - p(x+p)^2 + q(x+p) - r &= \\ x^3 + 2px^2 + (p^2 + q)x + pq - r. \end{aligned}$$

The following is the calculation in the last reduction. (See page 31).

	1	-p	q	-r
p	1	0	q ;	pq - r
p	1	p ;	p <sup>2</sup> + q	
p	1 ;	2p		
	1			

6. In any triangle, the square of the area expressed in terms of the lengths of the sides, is a polynome of four dimensions; and the area of the triangle, the lengths of whose sides are 3, 4, and 5, respectively, is 6. Find the polynome expressing the square of the area.

Let  $a$ ,  $b$ , and  $c$  be the lengths of the sides, and  $A$  the required polynome.

1st. The area vanishes if any two of the sides become together equal to the third side, hence if  $a+b=c$ ,  $A=0$ , and consequently  $A$  is divisible by  $a+b-c$ . Similarly it is divisible by  $b+c-a$  and by  $c+a-b$ .

2nd. The area vanishes if the three sides vanish together, hence if  $a+b+c=0$ ,  $A=0$ , and consequently  $A$  is divisible by  $a+b+c$ .

We have thus found four linear factors, but  $A$  is of only four dimensions.

$$\therefore A = m(a+b+c)(b+c-a)(c+a-b)(a+b-c),$$

in which  $m$  is a numerical constant.

$$\text{But } 6^2 \text{ or } 36 = m(3+4+5)(4+5-3)(5+3-4)(3+4-5) \\ = 576m; \therefore m = \frac{1}{16}.$$

(The above includes all the ways in which the area of a triangle can vanish, for the vanishing of only one side involves the equality of the other two, or if  $a=0$ ,  $b=c$ , and  $\therefore a+b=c$ , which is included in 1st.; if two sides vanish simultaneously, the three must vanish).

#### EXAMPLES ON THE COROLLARY.

7. Prove that  $(x+1)^{12} - x^{12} - 2x - 1$  is divisible by  $2x^3 + 3x^2 + x$ .

Factoring the latter expression we find it vanishes for  $x=0$ , or  $-1$  or  $-\frac{1}{2}$ . Substituting these values in the former polynome, it also vanishes. But these are different values of  $x$ , hence the truth of the proposition.

8.  $(x+y+z)^5 - x^5 - y^5 - z^5$  is divisible by  $(x+y+z)^3 - x^3 - y^3 - z^3$ .

The latter expression vanishes if  $x = -y$ , so also does the former.

By symmetry they both vanish if  $y = -z$  and if  $z = -x$ . Hence they are both divisible by  $(x+y)(y+z)(z+x)$ . But this expression is of three dimensions, as also is the latter of the given polynomes, hence it is a divisor of the former.

9. Prove that  $\{(a+b)^5 + (c+d)^5\}(a-b)(c-d) + \{(b+c)^5 + (a+d)^5\}(b-c)(a-d) + \{(b+d)^5 + (c+a)^5\}(b-d)(c-a)$  is algebraically divisible by  $(a-b)(c-d)(b-c)(a-d)(b-d)(c-a) \times (a+b+c+d)$ , and find the quotient.

Let  $a=b$  and the former polynome reduces to

$$\{(a+c)^5 + (a+d)^5\}(a-c)(a-d) + \{(a+d)^5 + (c+a)^5\}(a-d)(c-a)$$

which vanishes, the second complex term differing from the first only in the sign of one factor, having  $(c-a)$  instead of  $(a-c)$ .

Hence the former polynome is divisible by  $a-b$ , and by symmetry it is also divisible by  $a-c$ , by  $a-d$ , by  $b-c$ , by  $b-d$ , by  $c-d$ .

Again,  $(a+b)^5 + (c+d)^5$  is divisible by  $(a+b) + (c+d)$ ; for, on putting  $a+b = -(c+d)$ , it becomes  $\{-(c+d)\}^5 + (c+d)^5$  which  $= 0$ .

Similarly the other terms of the former of the given polnomes are each divisible by  $a+b+c+d$ , and consequently the whole is so divisible.

Now all these factors are different from each other, hence the former of the given polnomes is divisible by the product of these factors, *i.e.*, by the latter of the given polnomes.

Both of these polnomes are of seven dimensions, hence their quotient must be a number, the same for all values of  $a, b, c, d$ .

Put  $a=2, b=1, c=0, d=-1$ , and divide. The quotient will be found to be  $-5$ .

$$\begin{aligned} \therefore \{ & (a+b)^5 + (c+d)^5 \} (a-b)(c-d) + \{ (b+c)^5 + (a+d)^5 \} \times \\ & (b-c)(a-d) + \{ (b+d)^5 + (c+a)^5 \} (b-d)(c-a) = -5(a-b)(c-d) \\ & \times (b-c)(a-d)(b-d)(c-a)(a+b+c+d). \end{aligned}$$

• N.B.—It is not always necessary to find the factors of the divisor, as the following examples show.

10. Prove that  $x^2+x+1$  is a factor of  $x^{14}+x^7+1$ .

$x^2+x+1$  will be a factor of  $x^{14}+x^7+1$  provided

$$x^{14}+x^7+1=0 \text{ if } x^2+x+1=0.$$

$$\text{If } x^2+x+1=0$$

$$\therefore x^3+x^2+x=0$$

$$\therefore x^3+x^2+x+1=1$$

$$\therefore x^3=1$$

$$\therefore x^6=1 \text{ and } x^{12}=1$$

$$x^7=x \text{ and } x^{14}=x^2$$

$$\therefore x^{14}+x^7+1=x^2+x+1=0$$

$$\therefore x^2+x+1 \text{ is a factor of } x^{14}+x^7+1.$$

Art. XII. Two other methods of proving this proposition are worthy of notice.

1st.  $x^2+x+1$  will be a factor of  $x^{14}+x^7+1$  provided it is a factor of  $\{(x^{14}+x^7+1) \pm \text{a multiple of } (x^2+x+1)\}$ .

$$\begin{aligned} & x^{14}+x^7+1 \text{ differs by a multiple of } x^2+x+1 \text{ from} \\ & x^{14}+x^{11}(x^2+x+1)+x^8(x^2+x+1)+x^7+x^4(x^2+x+1)+ \\ & x(x^2+x+1)+1 \\ = & x^{12}(x^2+x+1)+x^9(x^2+x+1)+x^6(x^2+x+1)+x^3(x^2+x+1)+ \\ & (x^2+x+1) \\ = & (x^{12}+x^9+x^6+x^3+1)(x^2+x+1). \end{aligned}$$

Hence  $x^2+x+1$  is a factor of  $x^{14}+x^7+1$ .

$$\begin{aligned} \text{2nd. } \frac{x^{14}+x^7+1}{x^2+x+1} &= \frac{x^{21}-1}{x^7-1} \cdot \frac{x-1}{x^3-1} = \\ & \frac{(x^{21}-1)\{(x^{15}-1)-x(x^{14}-1)\}}{(x^7-1)(x^3-1)} = \\ & \frac{(x^{21}-1)(x^{15}-1)}{(x^7-1)(x^3-1)} - \frac{x(x^{21}-1)(x^{14}-1)}{(x^3-1)(x^7-1)}. \end{aligned}$$

But we see at once that on reduction both of these fractions give an integral quotient, hence  $(x^{14}+x^7+1) \div x^2+x+1$  gives an integral quotient.

11.  $x^2+x+1$  is a factor of  $(x+1)^7-x^7-1$ .

If  $x^2+x+1=0$ ,  $(x+1)^7-x^7-1$  will vanish also, for in such case  $x+1=-x^2$ .

$$\therefore (x+1)^7-x^7-1 = (-x^2)^7-x^7-1 = -x^{14}-x^7-1,$$

which by the last example vanishes if  $x^2+x+1=0$ ;

$$\therefore x^2+x+1 \text{ is a factor of } (x+1)^7-x^7-1.$$

For  $x$  substitute  $\frac{x}{y}$  and multiply by  $y^2$  and  $y^7$  respectively, and this example becomes

$$x^2+xy+y^2 \text{ is a factor of } (x+y)^7-x^7-y^7.$$

## Exercise xxi.

Determine the values of  $a$ ,  $b$ ,  $c$ ,  $d$ ,  $e$ , in the following cases:—

1.  $x^3 + 3bx^2 + 3cx + d$  vanishes for  $x=2$ , or  $3$ , or  $4$ .
2.  $x^4 + cx^2 + dx + e$  “ “  $x=1\frac{1}{3}$  or  $-3$  or  $4\frac{1}{3}$ .
3.  $x^3 + bx^2 + cx + 24$  “ “  $x=2$  or  $-3$ .
4.  $ax^3 + bx^2 + cx + 90$  “ “  $x=3$  or  $-5$  or  $2$ .
5.  $ax^4 + cx^2 - 30x + e$  “ “  $x=1\frac{1}{2}$  or  $-4$ , or  $2\frac{1}{2}$ .
6.  $81x^4 + 6cx^2 + 4dx + e$  “ “  $x=1\frac{2}{3}$  or  $-3\frac{1}{3}$  or  $1\frac{1}{3}$ .
7.  $ax^4 + bx^3 + cx^2 - 81$  “ “  $x=\frac{3}{2}$  or  $\frac{3}{4}$  or  $3$ .
8.  $ax^4 + cx^2 + dx + e$  “ “  $x=2$  or  $1\frac{1}{2}$  or  $-1$  and becomes  $14$  for  $x=1$ .

9.  $ax^3 + cx + d$  vanishes for  $x=1\frac{1}{4}$ , or  $2\frac{3}{4}$ , and becomes  $49$  for  $x=3$ , determine its value for  $x=-3$ .

Given that  $x^3 - px^2 + qx - r$  vanishes for  $x=a$ , or  $b$ , or  $c$ , determine the polynome that vanishes for

10.  $x = a+1$ , or  $b+1$ , or  $c+1$ .

11.  $x = a-1$ , or  $b-1$ , or  $c-1$ .

12.  $x = 1 - \frac{1}{a}$ , or  $1 - \frac{1}{b}$ , or  $1 - \frac{1}{c}$ .

13.  $x = ab$ , or  $bc$ , or  $ca$ .

14.  $x = a^2$ , or  $b^2$ , or  $c^2$ .

15.  $x = a(b+c)$ , or  $b(c+a)$ , or  $c(a+b)$ .  $\left\{ \begin{array}{l} a(b+c) = q - \frac{r}{a} \end{array} \right\}$ .

16.  $x = \frac{a+b}{c}$  or  $\frac{b+c}{a}$  or  $\frac{c+a}{b}$ .  $\left\{ \begin{array}{l} \frac{a+b}{c} = \frac{p}{c} - 1 \end{array} \right\}$ .

Prove that the following are cases of exact division:

17.  $(x-1)^{12} - x^6 + (x^2 - x + 1)^2 \div x^3 - 2x^2 + 2x - 1$ .

18.  $(x-1)^{16} - x^8 + (x^2 - x + 1)^3 \div x^3 - 2x^2 + 2x - 1$ .

19.  $(x-2)^{10}(2x-5)^{10} - x^{10} + 2^{10}(x^2 - 4x + 5)^5 \div x^3 - 6x^2 + 13x - 10$ .

20.  $(x^2 + 4x + 3)^{18} - x^{18} - x^2 - 5x - 3 \div x^3 + 6x^2 + 8x + 3$ .

21.  $(9x-4)^{21}(x-1)^{21} - x^{21} - (9x^2 - 14x + 4)^{21} \div (x-1) \times (9x-4)(9x^2 - 14x + 4)$ .

22.  $\{6(x-1)\}^{13} - (2x^2 + 3x - 4)^{13} + (2x^2 - 3x + 2)^{13} \div (2x^2 + 3x - 4)(2x^2 - 3x + 2)(x-1)$ .

$$23. \{2(x+1)(x-2)\}^{17} + (x^2 - 3x + 3)^{17} - (3x^2 - 5x - 1)^{17} \div (x+1)(x-2)(x^2 - 3x + 3)(3x^2 - 5x - 1).$$

$$24. \{6(x-1)\}^{16} - (2x^2 + 3x - 4)^{16} - (2x^2 - 3x + 2)^{16} + 2(2x^2 + 3x - 4)^8(2x^2 - 3x + 2)^4 \div (x-1)(2x^2 + 3x - 4)(2x^2 - 3x + 2)$$

$$25. \{2(x+1)(x-2)\}^{20} - (x^2 - 3x + 3)^{20} - (3x^2 - 5x - 1)^{20} + 2(x^2 - 3x + 3)^9(3x^2 - 5x - 1)^{11} \div (x+1)(x-2)(x^2 - 3x + 3) \times (3x^2 - 5x - 1).$$

$$26. 1 + x^4 + x^8 \div 1 + x + x^2.$$

$$27. x^{10} + x^5y^5 + y^{10} \div x^2 + xy + y^2.$$

$$28. 1 + x^3 + x^6 + x^9 + x^{12} \div 1 + x + x^2 + x^3 + x^4.$$

$$29. 1 + x^4 + x^8 + x^{12} + x^{16} \div 1 + x + x^2 + x^3 + x^4.$$

$$30. x^{15} + x^{10}y^5 + x^5y^{10} + y^{15} \div x^3 + x^2y + xy^2 + y^3.$$

$$31. x^{17} + x^4 + x^3 + x + 1 \div x^4 + x^3 + x^2 + x + 1.$$

$$32. 1 + x + x^2 + x^3 + x^5 + x^6 + x^5 \div 1 + x + x^2 + x^3 + x^4 + x^5 + x^6.$$

Find the quotient of the following divisions in which  $D$  denotes the product

$$(b-c)(c-a)(a-b)(a-d)(b-d)(c-d);$$

$$33. (b^2c^2 + a^2d^2)(b-c)(a-d) + (c^2a^2 + b^2d^2)(c-a)(b-d) + (a^2b^2 + c^2d^2)(a-b)(c-d) \div D.$$

$$34. (bc + ad)(b^2 - c^2)(a^2 - d^2) + (ca + bd)(c^2 - a^2)(b^2 - d^2) + (ab + cd)(a^2 - b^2)(c^2 - d^2) \div D.$$

$$35. (b+c)(a+d)(b^2 - c^2)(a^2 - d^2) + \text{the two similar terms} \div D.$$

$$36. (b^2 + c^2)(a^2 + d^2)(b-c)(a-d) + \quad \quad \quad \div D.$$

$$37. \{bc(b+c)^2 + ad(a+d)^2\}(b-c)(a-d) + \quad \quad \quad \div D.$$

$$38. \{bc(b+c) + ad(a+d)\}(b^2 - c^2)(a^2 - d^2) + \quad \quad \quad \div D.$$

$$39. \{bc(b^3 + c^3) + ad(a^3 + d^3)\}(b-c)(a-d) + \quad \quad \quad \div D.$$

$$40. (b+c-a-d)^4(b-c)(a-d) + \quad \quad \quad \div D.$$

41. The sum of the fractions  $\frac{1}{1}, \frac{1}{2}, \frac{1}{3}, \dots, \frac{1}{n}$ , increased by the sum of their products two by two, increased by the sum of their products three by three, ..... increased by their product is equal to  $n$ .

42. In any trapezium the square of the area expressed in terms of the lengths of the parallel sides and the diagonals, is a polynome of four dimensions, determine that polynome.

43. In any quadrilateral inscribed in a circle, the square of the area expressed in terms of the lengths of the sides, is a polynome of four dimensions, find that polynome.

**Theorem III.** If the polynome  $f(x)^n$  vanish for more than  $n$  different values of  $x$ , it vanishes identically, the coefficient of every term being zero.

Cor. If a rational integral expression of  $n$  dimensions be divisible by more than  $n$  linear factors, the expression is identically zero.

## EXAMPLES.

1.  $\frac{(x-a)(x-b)}{(c-a)(c-b)} + \frac{(x-b)(x-c)}{(a-b)(a-c)} + \frac{(x-c)(x-a)}{(b-c)(b-a)} - 1 = 0$ , if  $a$ ,  $b$ , and  $c$  are unequal; for this is a polynome of two dimensions in  $x$ , but it vanishes for  $x=a$ , and, therefore, by symmetry for  $x=b$ , and for  $x=c$ , that is, for three different values of  $x$ , hence it vanishes identically.

2.  $\{(a+b)^2 + (c+d)^2\}(a-b)(c-d) + \{(c+b)^2 + (b+d)^2\}(b-c)(a-d) + \{(c+a)^2 + (b+d)^2\}(c-a)(b-d) = 0$ .

Substitute  $b$  for  $a$  and the expression becomes  $\{(b+c)^2 + (b+d)^2\}(b-c)(b-d) + \{(c+b)^2 + (b+d)^2\}(c-b)(b-d)$  which vanishes, hence the given expression is divisible by  $a-b$ , and consequently by symmetry it is divisible by  $(a-b)$ ,  $(b-c)$ ,  $(c-d)$ ,  $(a-c)$ ,  $(b-d)$ , and  $(a-d)$ . But the given expression is of only four dimensions, while it appears to have six linear factors, hence it vanishes identically.

## Exercise xxii.

Verify the following:

$$1. \frac{x^2 y^2 z^2}{b^2 c^2} + \frac{(x^2 - b^2)(y^2 - b^2)(z^2 - b^2)}{b^2(b^2 - c^2)} + \frac{(x^2 - c^2)(y^2 - c^2)(z^2 - c^2)}{c^2(c^2 - b^2)} = x^2 + y^2 + z^2 - b^2 - c^2.$$



$$2. \frac{y^2 z^3}{b^2 c^2} + \frac{(y^2 - b^2)(z^2 - b^2)}{b^2(b^2 - c^2)} + \frac{(y^2 - c^2)(z^2 - c^2)}{c^2(c^2 - b^2)} = 1.$$

$$3. \frac{x^2 y^2}{a^2 b^2} + \frac{(x^2 - a^2)(a^2 - y^2)z^2}{(z^2 - a^2)(b^2 - a^2)a^2} + \frac{(x^2 - b^2)(b^2 - y^2)z^2}{(b^2 - z^2)(b^2 - a^2)b^2} + \frac{(z^2 - x^2)(z^2 - y^2)}{(z^2 - a^2)(b^2 - z^2)} = 0.$$

$$4. \frac{1}{(x+a)(a-b)(a-c)} + \frac{1}{(x+b)(b-c)(b-a)} + \frac{1}{(x+c)(c-a)(c-b)} = \frac{1}{(x+a)(x+b)(x+c)}.$$

$$5. bc(b^2 - c^2) + ca(c^2 - a^2) + ab(a^2 - b^2) = (a+b+c)\{a^2(b-c) + b^2(c-a) + c^2(a-b)\}.$$

$$6. \frac{a+x}{x(x-y)(x-z)} + \frac{a+y}{y(y-x)(y-z)} + \frac{a+z}{z(z-x)(z-y)} = \frac{a}{xyz}.$$

$$7. \frac{a^4(b^2 - c^2) + b^4(c^2 - a^2) + c^4(a^2 - b^2)}{a^2(b-c) + b^2(c-a) + c^2(a-b)} = \frac{1}{2}\{(a+b+c)^3 - a^3 - b^3 - c^3\}.$$

$$8. (adf + bcf + bed - ace)^2 + (bec + aed + acf - bdf)^2 = (a^2 + b^2)(c^2 + d^2)(e^2 + f^2).$$

$$9. \frac{(a-b)^5 + (b-c)^5 + (c-a)^5}{(a-b)(b-c)(c-a)} = \frac{5}{2}\{(a-b)^2 + (b-c)^2 + (c-a)^2\}.$$

$$10. (-x+y+z)(x-y+z)(x+y-z) + x(x-y+z)(x+y-z) + y(x+y-z)(-x+y+z) + z(-x+y+z)(x-y+z) = 4xyz.$$

$$11. \frac{(a^2 - b^2)^3 + (b^2 - c^2)^3 + (c^2 - a^2)^3}{(a+b)(b+c)(c+a)} = (a-b)^3 + (b-c)^3 + (c-a)^3.$$

$$12. x^2(y+z)^2 + y^2(z+x)^2 + z^2(x+y)^2 + 2xyz(x+y+z) = 2(xy+yz+zx)^2.$$

**Theorem IV.** If the polynomials  $f(x)^n$ ,  $\phi(x)^m$  ( $n$  not less than  $m$ ) are equal for more than  $n$  different values of  $x$ , they are equal for all values, and the coefficients of equal powers of  $x$  in each are equal to one another.

(This is called the Principle of Indeterminate Coefficients. The full use of it cannot be exhibited till the student is able to work simultaneous equations.)

## EXAMPLES.

$$1. \frac{a^2}{(a-b)(a-c)(a-d)} + \frac{b^2}{(b-a)(b-c)(b-d)} + \frac{c^2}{(c-a)(c-b)(c-d)} + \frac{d^2}{(d-a)(d-b)(d-c)} = 0.$$

Assume  $\frac{x^3}{(x-a)(x-b)(x-c)(x-d)} = \frac{A}{x-a} + \frac{B}{x-b} + \frac{C}{x-c} + \frac{D}{x-d}$  (α)

in which  $A, B, C, D$  are independent of  $x$ .

Multiply by  $(x-a)(x-b)(x-c)(x-d)$ .

$$\therefore x^3 = (A+B+C+D)x^3 + \text{terms in lower powers of } x.$$

Now this equality holds for more than three values of  $x$ , holding in fact for *all* finite values of  $x$ .

$$\therefore A+B+C+D=0$$
 (β)

Again multiply both sides of (α) by  $x-a$

$$\frac{x^2}{(x-b)(x-c)(x-d)} = A + \left( \frac{B}{x-b} + \frac{C}{x-c} + \frac{D}{x-d} \right) (x-a).$$

Put  $x=a$

$$\therefore \frac{a^2}{(a-b)(a-c)(a-d)} = A.$$

By symmetry  $\frac{b^2}{(b-a)(b-c)(b-d)} = B$ , &c.

Adding

$$\frac{a^2}{(a-b)(a-c)(a-d)} + \frac{b^2}{(b-a)(b-c)(b-d)} + \frac{c^2}{(c-a)(c-b)(c-d)} + \frac{d^2}{(d-a)(d-b)(d-c)} = A+B+C+D=0 \text{ by } (\beta).$$

$$2. \frac{a^2(a+b)(a+c)}{(a-b)(a-c)} + \frac{b^2(b+c)(b+a)}{(b-c)(b-a)} + \frac{c^2(c+a)(c+b)}{(c-a)(c-b)} = (a+b+c)^2.$$

$$\text{Assume } x^3 - px^2 + qx - r = (x-a)(x-b)(x-c). \quad (\alpha).$$

$$\therefore x^3 + px^2 + qx + r = (x+a)(x+b)(x+c). \quad (\beta).$$

$$\frac{x^4 + px^3 + qx^2 + rx}{x^3 - px^2 + qx - r} = x + 2p + \frac{A}{x-a} + \frac{B}{x-b} + \frac{C}{x-c} \quad (\gamma).$$

Multiply by  $x^3 - px^2 + qx - r$  and equate the coefficients of the terms in  $x^2$ . {In multiplying the fractions in the right-hand member of ( $\gamma$ ), use the factor side of ( $\alpha$ ).}

$$q = q - 2p^2 + A + B + C$$

$$\therefore A + B + C = 2p^2.$$

Multiply both members of ( $\gamma$ ) by  $x-a$

$$\frac{x(x+a)(x+b)(x+c)}{(x-b)(x-c)} = A + \left\{ x + 2p + \frac{B}{x-b} + \frac{C}{x-c} \right\} (x-a).$$

Put  $x = a$ ,

$$\frac{2a^2(a+b)(a+c)}{(a-b)(a-c)} = A.$$

By symmetry

$$\frac{2b^2(b+c)(b+a)}{(b-c)(b-a)} = B \text{ and } \frac{2c^2(c+a)(c+b)}{(c-a)(c-b)} = C;$$

$$\therefore \frac{a^2(a+b)(a+c)}{(a-b)(a-c)} + \frac{b^2(b+c)(b+a)}{(b-c)(b-a)} + \frac{c^2(c+a)(c+b)}{(c-a)(c-b)}$$

$$= \frac{1}{2}(A+B+C) = p^2.$$

$$= (a+b+c)^2.$$

3. Extract the square root of  $1+x+x^2+x^3+x^4+\&c.$

Assume the square root to be  $1+ax+bx^2+cx^3+dx^4+\&c.$

$$\therefore 1+x+x^2+x^3+x^4+\&c. = (1+ax+bx^2+cx^3+dx^4+\&c.)^2$$

$$= 1+2ax+(a^2+2b)x^2+2(ab+c)x^3+(2d+2ac+b^2)x^4+\&c.$$

$$\therefore 2a = 1$$

$$\therefore a = \frac{1}{2}$$

$$2b+a^2 = 1$$

$$\therefore b = \frac{1}{2}\left(1 - \frac{1}{4}\right) = \frac{3}{8}$$

$$2(c+ab) = 1$$

$$\therefore c = \frac{1}{2} - \left(\frac{1}{2} \times \frac{3}{8}\right) = \frac{5}{16}$$

$$2d+2ac+b^2 = 1$$

$$\therefore d = \frac{1}{2} \left\{ 1 - \frac{5}{16} - \frac{9}{64} \right\} = \frac{35}{256}.$$

$$\therefore \sqrt{1+x+x^2+\&c.} = 1 + \frac{1}{2}x + \frac{3}{8}x^2 + \frac{5}{16}x^3 + \frac{35}{256}x^4 + \&c.$$

(NOTE.—As it is frequently necessary to determine the coefficient of a particular power of  $x$ , a few preliminary exercises are given on this subject.)

## Exercise xxiii.

Determine the coefficient of

1.  $x^4$  in  $(1+ax)^3 + (1+bx)^5 + (1-cx)^6$ .
2.  $x^5$  in  $(1+x+2x^2+3x^3)(1-x+3x^2+x^3-5x^4)$ .
3.  $x^4$  in  $(1+x+2x^2+3x^3+4x^4+\&c)(1-x-x^2-x^3+x^4-\&c)$ .
4.  $x^3$  in  $A(x-b)(x-c)(x-d) + B(x-a)(x-c)(x-d) + C(x-a)(x-b)(x-d) + D(x-a)(x-b)(x-c)$ .
5.  $x^4$  in  $(1-ax)^3(1+ax)^5$ .
6.  $x^4$  in  $(1+ax)^3(1-bx)^5$ .
7. In the product

$$(1+ax+bx^2+cx^3+\&c)(1-ax+bx^2-cx^3+\&c)$$

prove that the coefficients of the odd powers of  $x$  must be all zeros.

Determine the value of the following expressions :

$$8. \frac{1}{(a-b)(a-c)(a-d)} + \frac{1}{(b-a)(b-c)(b-d)} + \frac{1}{(c-a)(c-b)(c-d)} + \frac{1}{(d-a)(d-b)(d-c)}$$

$$9. \frac{a}{(a-b)(a-c)(a-d)} + \frac{b}{(b-a)(b-c)(b-d)} + \&c$$

$$10. \frac{a^2}{(a-b)(a-c)(a-d)} + \text{three similar terms.}$$

$$11. \frac{a^3}{(a-b)(a-c)(a-d)} + \text{“ “}$$

$$12. \frac{a^4}{(a-b)(a-c)(a-d)} + \text{three similar terms.}$$

$$13. \frac{bcd}{(a-b)(a-c)(a-d)} + \text{“ “}$$

$$14. \frac{a(a+b)(a+c)}{(a-b)(a-c)} + \text{two “}$$

$$15. \frac{a^3(a+b)(a+c)}{(a-b)(a-c)} + \text{“ “}$$

$$16. \frac{a^4(a+b)(a+c)}{(a-b)(a-c)} + \text{“ “}$$

$$17. \frac{a(a+b)(a+c)(a+d)}{(a-b)(a-c)(a-d)} + \text{three similar terms.}$$

$$18. \frac{a^2(a+b)(a+c)(a+d)}{(a-b)(a-c)(a-d)} + \text{ " " "}$$

$$19. \frac{a^3(a+b)(a+c)(a+d)}{(a-b)(a-c)(a-d)} + \text{ " " "}$$

$$20. \frac{bc(b+c)}{(a-b)(a-c)} + \text{two similar terms.}$$

[For numerator use  $x^3 + 2px^2 + (p^2 + q)x + (pq - r)$ .]

$$21. \frac{(2a+b)(2a+c)}{(a-b)(a-c)} + \text{two similar terms.}$$

[For numerator use  $x^3 - 2px^2 + (p^2 + q)x - (pq - r)$ .]

$$22. \frac{a(b+c)}{(a-b)(a-c)} + \text{two similar terms.}$$

[For numerator use  $x(x+p)$ .]

$$23. \frac{b+c+d}{(a-b)(a-c)(a-d)} + \text{three similar terms.}$$

$$24. \frac{a^3(bc+cd+db)}{(a-b)(a-c)(a-d)} + \text{ " " "}$$

$$25. \frac{bc+cd+db}{(a-b)(a-c)(a-d)} + \text{ " " "}$$

Extract the square-root of (to 4 terms):

$$26. 1+x. \quad | \quad 27. 1-x. \quad | \quad 28. 1+2x+3x^2+4x^3 + \&c.$$

$$29. 1-4x+10x^2-20x^3+35x^4-56x^5+84x^6.$$

$$30. \text{Extract the cube-root of } 1+x. \quad (\text{To 4 terms}).$$

**Art. XI.** 1. Find the condition that  $px^2+2qx+r$  and  $p'x^2+2q'x+r'$  shall have a common factor.

Multiply the polynomes by  $p'$  and  $p$  respectively, and take the difference of the products, also by  $r'$  and  $r$  respectively, and divide the difference of the products by  $x$ .

$$\begin{array}{r|l} \begin{array}{l} p'px^2 + 2p'qx + p'r \\ pp'x^2 + 2pq'x + pr' \\ \hline 2(pq' - p'q)x + (pr' - p'r) \end{array} & \begin{array}{l} pr'x^2 + 2qr'x + rr' \\ p'r x^2 + 2q'r x + r'r \\ \hline (pr' - p'r)x + 2(qr' - r'q) \end{array} \end{array}$$

Multiply the former of these remainders by  $(pr' - p'r)$  and the latter by  $2(pq' - p'q)$ , and the difference of the products is

$$(pr' - p'r)^2 - 4(pq' - p'q)(qr' - r'q).$$

But if the given polynomes have a linear factor this remainder must vanish, or

$$(pr' - p'r)^2 = 4(pq' - p'q)(qr' - r'q).$$

If the given polynomes have a quadratic factor, the *linear* remainders must vanish identically, or (Th. III.)

$$pq' - p'q = 0, pr' - p'r = 0, \text{ and } qr' - r'q = 0,$$

$$\text{or, } \frac{p}{p'} = \frac{q}{q'} = \frac{r}{r'}.$$

2. Find the condition that  $px^3 + 3qx^2 + 3rx + s$  shall have a square factor.

Assume the square factor to be  $(x-a)^2$ . On division, the remainder must be zero for every finite value of  $x$ , and consequently (Th. III.) the co-efficient of each term of the remainder must be zero. Divide by  $(x-a)^2$ , neglecting the first remainder.

$a$	$p$	$3q$	$3r$	$s$
		$pa$	$pa^2 + 3qa$	
$a$	$p$	$pa + 3q$	$pa^2 + 3qa + 3r$	$R$
		$pa$	$2pa^2 + 3qa$	
		$p$	$2pa + 3q$	$3(pa^2 + 2qa + r)$

$$\therefore pa^2 + 2qa + r = 0;$$

$\therefore px^2 + 2qx + r$  is divisible by  $x-a$  (Th. I. Cor. 2),

or,  $px^3 + 3qx^2 + 3rx + s$  and  $px^2 + 2qx + r$  have a common divisor. Multiply the latter polynome by  $x$  and subtract the product from the former, and the proposition reduces to

If  $px^3 + 3qx^2 + 3rx + s$  have a square factor,  $px^2 + 2qx + r$  and  $qx^2 + 2rx + s$  will have the square-root of that factor for a common divisor.

3. If  $px^3 + 3qx^2 + 3rx + s$  vanish for  $x=a$ , or  $b$ , or  $c$ , find in terms of  $x, p, q, r$  the value of

$$\frac{1}{x-a} + \frac{1}{x-b} + \frac{1}{x-c}.$$

Reduce to a common denominator and add the numerators

$$\frac{3x^2 - 2(a+b+c)x + (ab+bc+ca)}{(x-a)(x-b)(x-c)}$$

Multiply both numerator and denominator by  $p$  and reduce by Th. II., and Ex. 4 of Th. II.

$$\frac{\mathfrak{B}(px^2 + 2qx + r)}{px^3 + 3qx^2 + 3rx + s}$$

$$\therefore \frac{x^{m+1}}{x-a} + \frac{x^{m+1}}{x-b} + \frac{x^{m+1}}{x-c} = \frac{\mathfrak{B}(px^{m+3} + 2qx^{m+2} + rx^{m+1})}{px^3 + 3qx^2 + 3rx + s}$$

4. If  $px^3 + 3qx^2 + 3rx + s$  vanish for  $x = a$ , or  $b$ , or  $c$ , express in terms of  $p, q, r, s$ , the following,  $a + b + c, a^2 + b^2 + c^2, a^3 + b^3 + c^3$   
 . . . . .,  $a^m + b^m + c^m$ .

Divide  $x^{m+1}$  by  $x - a$ .

$$a \left| \begin{array}{cccccccc} 1 & & & & & & & \\ & a & a^2 & a^3 & & & a^m & a^{m+1} \\ \hline & 1 & a & a^2 & a^3 & . . . . . & a^m & a^{m+1} \end{array} \right.$$

Similarly divide  $x^{m+1}$  by  $x - b$  and also by  $x - c$ .

add together the quotients

$$\frac{x^{m+1}}{x-a} + \frac{x^{m+1}}{x-b} + \frac{x^{m+1}}{x-c} = \mathfrak{B}x^m + (a+b+c)x^{m-1} + (a^2 + b^2 + c^2)x^{m-2} + (a^3 + b^3 + c^3)x^{m-3} + \&c.$$

Hence, by the last example, the required expressions are the coefficients taken in order, beginning with the second, of the terms in the quotient of  $\mathfrak{B}(px^{m+3} + 2qx^{m+2} + rx^{m+1}) \div (px^3 + 3qx^2 + 3rx + s)$ . These may now be found by Horner's Division.

5. Writing  $s_1$  for  $a + b + c, s_2$  for  $a^2 + b^2 + c^2, \&c.$ , express  $(a-b)^4 + (b-c)^4 + (c-a)^4$  in terms of  $s_1, s_2, s_3, s_4$ .

By actual expansion

$$(x-a)^4 + (x-b)^4 + (x-c)^4 =$$

$$3x^4 - 4(a+b+c)x^3 + 6(a^2 + b^2 + c^2)x^2 - 4(a^3 + b^3 + c^3)x + a^4 + b^4 + c^4 = \mathfrak{B}x^4 - 4s_1x^3 + 6s_2x^2 - 4s_3x + s_4.$$

Put  $x = a, = b, = c$  in succession.

$$(a-b)^4 + (c-a)^4 = 3a^4 - 4s_1a^3 + 6s_2a^2 - 4s_3a + s_4$$

$$(a-b)^4 + (b-c)^4 = 3b^4 - 4s_1b^3 + 6s_2b^2 - 4s_3b + s_4$$

$$(b-c)^4 + (c-a)^4 = 3c^4 - 4s_1c^3 + 6s_2c^2 - 4s_3c + s_4$$

$$\therefore 2\{(a-b)^4 + (b-c)^4 + (c-a)^4\} = 3s_4 - 4s_1s_3 + 6s_2^2 - 4s_3s_1 + 3s_4$$

$$\therefore (a-b)^4 + (b-c)^4 + (c-a)^4 = s_0s_4 - 4s_1s_3 + 3s_2^2$$

in which  $s_0$  is written for  $\mathfrak{B}$  or  $1 + 1 + 1, i.e., a^0 + b^0 + c^0$ .

## Exercise xxiii. (a).

1. Determine the condition necessary in order that  $x^2 + px + q$  and  $x^3 + p'x + q$  may have a common divisor.

2. The expression  $x^6 + 3a^2x^5 + 3bx^4 + cx^3 + 3dx^2 + 3e^2x + f^3$  will be a complete cube if

$$f = \frac{e}{a} = \frac{d}{b} = \frac{c - a^6}{6a^2} = b - a^4.$$

3. Prove that  $ax^5 + bx + c$  and  $a + bx^4 + cx^5$  will have a common quadratic factor if

$$b^2c^2 = (c^2 - a^2 + b^2)(c^2 - a^2 + ab).$$

4. Prove that  $ax^5 + bx^2 + c$  and  $a + bx^3 + cx^5$  will have a common quadratic factor if

$$a^2b^2 = (a^2 - c^2)(a^2 - c^2 + bc).$$

5. Prove that  $ax^4 + bx^3 + cx + d$  and  $a + bx + cx^3 + dx^4$  will have a common quadratic factor if

$$(a + d)^2 = (b - c)(bd - ac).$$

6.  $x^3 + px^2 + qx + r$  will be divisible by  $x^2 + ax + b$  if

$$a^3 - 2pa^2 + (p^2 + q)a + r - pq = 0, \text{ and } b^3 - qb^2 + rpb - r^2 = 0.$$

7.  $x^4 + px + q$  will be divisible by  $x^2 + ax + b$  if

$$a^6 - 4qa^2 = p^2 \text{ and } (b^2 + q)(b^2 - q) = p^2b^3.$$

8. Determine the condition necessary in order that  $x^4 + 4px^3 + 6qx^2 + 4rx + t$  may have a square factor.

If  $x^4 + 4px^3 + 6qx^2 + 4rx + t$  vanish for  $x = a$ , or  $b$ , or  $c$ , or  $d$ , find in terms of  $x, p, q, r, t$ , the value of

$$9. \frac{x^n}{x-a} + \frac{x^n}{x-b} + \frac{x^n}{x-c} + \frac{x^n}{x-d}.$$

$$10. \Sigma a, \Sigma a^2, \Sigma a^3, \Sigma a^4, \Sigma a^5, \Sigma a^6.$$

$$11. \Sigma (a-b)^4, \Sigma (a-b)^6.$$

12. Determine the values of the expressions in Ex. 9, 10, 11, for the polynome  $x^4 - 14x^2 + x - 38$ .



## CHAPTER III.

## SECTION I.—DIRECT APPLICATION OF THE FUNDAMENTAL FORMULAS

FORMULAS [1] AND [2].  $(x \pm y)^2 = x^2 \pm 2xy + y^2$ , &c.

**Art. XII.** From this it appears that a trinomial of which the extremes are squares, is itself a square if four times the product of the extremes is equal to the square of the mean, and that to factor such a trinomial, we have simply to connect the square root of each of the squares by the sign of the other term, and write the result twice as a factor.

## EXAMPLES.

1.  $4x^4 - 80x^2y^2 + 400y^4 = (2x^2 - 20y^2)(2x^2 - 20y^2)$
2.  $1 - 12x^2y^2 + 36x^4y^4 = (1 - 6x^2y^2)(1 - 6x^2y^2)$ .
3.  $(a-b)^2 + (b-c)^2 + 2(a-b)(b-c)$ . This equals  $(a-b+b-c) \times (a-b+b-c) = (a-c)(a-c)$ .
4.  $x^2 + y^2 + z^2 + 2xy - 2xz - 2yz$ .

Here the three squares and the three double products suggest that the expression is the square of a linear *trinomial* in  $x, y, z$ .

An inspection of the signs of the double products enables us to determine the signs which are to connect  $x, y, z$ : we see that

- 1st. The signs of  $x$  and  $y$  must be *alike*.
- 2nd. The signs of  $x$  and  $z$  must be *different*.
- 3rd. The signs of  $y$  and  $z$  must be *different*. Hence we have  $x+y-z$ , or  $-x-y+z = -(x+y-z)$ , and the factors are  $(x+y-z)(x+y-z)$ .

## Exercise xxiv.

1.  $9m^2 + 12m + 4$ ;  $c^{2m} - 2c^m + 1$ .
2.  $y^6 - 2y^3z^3 + z^6$ ;  $16x^2y^2 + 16xy^3 + 4y^4$ .
3.  $9a^2b^2 + 12abc + 4c^2$ ;  $36x^2y^2 - 24xy^3 + 4y^4$

4.  $\frac{1}{4}x^4 + 16y^2z^2 - 4x^2yz$ ;  $\frac{1}{4}a^4 - \frac{1}{3}a^2b^2c^2 + \frac{1}{9}b^4c^4$ .
5.  $(a+b)^2 + c^2 - 2c(a+b)$ ;  $9x^8 - \frac{3}{2}x^4y^2 + \frac{1}{16}y^4$ .
6.  $z^2 + (x-y)^2 - 2z(x-y)$ ;  $\left(\frac{a}{b}\right)^{2m} + \left(\frac{b}{a}\right)^{2m} - 2$ .
7.  $(x^2-y)^2 + 2(x^2-y)(y-z^2) + (y-z^2)^2$ .
8.  $(x^2-xy)^2 - 2(x^2-xy)(xy-y^2) + (xy-y^2)^2$ .
9.  $(a+b+c)^2 - 2c(a+b+c) + c^2$ ;  $\frac{1}{16}p^6 - 2p^3q^2 + \frac{1}{6}q^4$ .
10.  $(3x-4y)^2 + (2x-3y)^2 - 2(3x-4y)(2x-3y)$ .
11.  $(x^2-xy+y^2)^2 + (x^2+xy+y^2)^2 + 2(x^4+x^2y^2+y^4)$ .
12.  $(5x^2+2xy+7y^2)^2 + (4x^2+6y^2)^2 - 2(4x^2+6y^2) \times (5x^2+2xy+7y^2)$ .
13.  $\left(\frac{a}{b}\right)^{2m} + \left(\frac{b}{a}\right)^{2n} - 2\left(\frac{a}{b}\right)^{m-n}$ .
14.  $a^2 + b^2 + c^2 - 2ab - 2bc + 2ac$ .
15.  $a^4 + b^4 + c^4 - 2a^2b^2 - 2a^2c^2 + 2b^2c^2$ .
16.  $(a-b)^2 + (b-c)^2 + (c-a)^2 + 2(a-b)(b-c) - 2(a-b)(c-a) + 2(b-c)(a-c)$ .
17.  $4a^4 - 12a^2b + 9b^2 + 16a^2c + 16c^2 - 24bc$ .

FORMULA [4].  $x^2 - y^2 = (x+y)(x-y)$ .

Art. XIII. In this case we have merely to take the square-root of each of the squares, and connect the results with the sign + for one of the factors, and with the sign - for the other.

EXAMPLES.

1.  $(a+b)^2 - (c+d)^2$ .  
 This =  $\{(a+b) + (c+d)\} \{(a+b) - (c+d)\}$   
 =  $(a+b+c+d)(a+b-c-d)$ .
2. Factor  $(x^2 + 5xy + y^2)^2 - (x^2 - xy + y^2)^2$ .

Here we have

$$\{(x^2 + 5xy + y^2) + (x^2 - xy + y^2)\} \{(x^2 + 5xy + y^2) - (x^2 - xy + y^2)\}$$

$$= 2(x^2 + 2xy + y^2)(6xy) = 12xy(x+y)^2.$$

3.  $a^2 - b^2 - c^2 + 2bc$ .

$$\text{This} = a^2 - (b-c)^2 = (a+b-c)(a-b+c).$$

$$4. \text{ Resolve } (a^2 + b^2)^2 - (a^2 - b^2)^2 - (a^2 + b^2 - c^2)^2.$$

$$\begin{aligned} \text{This} &= 4a^2b^2 - (a^2 + b^2 - c^2)^2 \\ &= (2ab + a^2 + b^2 - c^2)(2ab - a^2 - b^2 + c^2). \end{aligned}$$

The former of these factors  $= (a+b)^2 - c^2 = (a+b+c)(a+b-c)$ ;  
and the latter  $= c^2 - (a-b)^2 = (c+a-b)(c-a+b)$ .

$\therefore$  the given expression

$$= (a+b+c)(a+b-c)(c+a-b)(c-a+b).$$

### Exercise xxv.

$$1. 49a^2 - 4b^2.$$

$$2. 9a^2 - \frac{1}{4}b^2.$$

$$3. 81a^4 - 16b^4.$$

$$4. 100x^2 - 36y^2.$$

$$5. 5a^2b - 20bx^2y^2.$$

$$6. 9x^6 - 16y^4.$$

$$7. \frac{1}{16}c^2 - 1.$$

$$8. 4y^4 - \frac{1}{9}x^2z^2.$$

$$9. 81a^4 - 1.$$

$$10. a^4 - 16b^4.$$

$$11. a^{16} - b^{16}.$$

$$12. a^2 - b^2 + 2bc - c^2.$$

$$13. (a+2b)^2 - (3x-4y)^2.$$

$$14. (x^2+y^2)^2 - 4x^2y^2.$$

$$15. (x+y)^2 - 4z^2.$$

$$16. (3x+5)^2 - (5x+2)^2.$$

$$17. 4x^2y^2 - (x^2 + y^2 - z^2)^2.$$

$$18. (x^2 + xy - y^2)^2 - (x^2 - xy - y^2)^2.$$

$$19. (x^2 - y^2 + z^2)^2 - 4x^2z^2.$$

$$20. (a+b+c+d)^2 - (a-b+c-d)^2.$$

$$21. (2+3x+4x^2)^2 - (2-3x+4x^2)^2.$$

$$22. (a^2 + b^2 + 4ab)^2 - (a^2 + b^2)^2.$$

$$23. (a^2 - b^2 + c^2 - d^2)^2 - (2ac - 2bd)^2.$$

$$24. (x^2 - y^2 - z^2)^2 - 4y^2z^2.$$

$$25. (a^6 - a^3b^3 + b^6)^2 - (a^6 - 5a^3b^3 + b^6)^2.$$

$$26. a^{12} - b^{12} + 6a^9b^3 - 6b^9a^3 + 8b^9a^3 - 8a^9b^3.$$

$$27. (x^2 + y^2 + z^2 - xy - yz - zx)^2 - (xy + yz + zx)^2.$$

$$28. (x^2 + y^2 + z^2 - 2xy + 2xz - 2yz) - (y+z)^2.$$

$$29. 2a^2b^2 + 2b^2c^2 + 2c^2a^2 - a^4 - b^4 - c^4.$$

$$30. x^4 + y^4 + z^4 - 2x^2y^2 - 2y^2z^2 - 2z^2x^2.$$

1.  
2.  
3.  
4.  
5.  
= (ax

Art  
the re  
the fi  
are de

(1)  
pressio

(2)  
already

sion.  
an exp

1st.

2nd.

sum is

if the s

3rd.

whose

and bef

fore b i

1. x^2

10, 4 a

whose s

hence t

$$\text{FORMULA A. } (x+r)(x+s) = x^2 + (r+s)x + rs.$$

## EXAMPLES

1.  $x^3 - 9x^2 + 20 = (x^3 - 5)(x^2 + 4).$
2.  $(x-y)^2 + x - y - 110 = (x-y+11)(x-y-10).$
3.  $(a^2 - ab + b^2)^2 + 6b(a^2 - ab + b^2) - 4a^2 + 9b^2 =$   
 $\{(a^2 - ab + b^2) + (2a + 3b)\} \{(a^2 - ab + b^2) - (2a - 3b)\}.$
4.  $(x^2 - 5x)^2 - 6(x^2 - 5x) - 40 = (x^2 - 5x + 4)(x^2 - 5x - 10).$
5.  $(ax + by + c)^2 - (m - n)(ax + by + c) - mn$   
 $= (ax + by + c - m)(ax + by + c + n).$

**Art. XIV.** It will be seen that the first (or *common*) term of the required factors, is obtained by extracting the square root of the first term of the given expression, and that the other terms are determined by observing two conditions :

(1) Their product must equal the *third* term of the given expression.

(2) Their sum (*algebraic*) multiplied into the *common* term already found, must equal the middle term of the given expression. Hence, to make a systematic search for integral factors of an expression of the form  $x^2 \pm bx \pm c$ , we may proceed as follows :

1st. Write down every pair of factors whose product is  $c$ .

2nd. If the sign before  $c$  is  $+$ , select the pair of factors whose *sum* is  $b$ , and write *both* factors  $x+$ , if the sign before  $b$  is  $+$ ;  $x-$ , if the sign before  $b$  is  $-$ .

3rd. But if the sign before  $c$  is  $-$ , select the pair of factors whose *difference* is  $b$ , and write before the *larger* factor  $x+$  or  $x-$ , and before the other factor  $x-$  or  $x+$ , according as the sign before  $b$  is  $+$  or  $-$ .

## EXAMPLES.

1.  $x^2 + 9x + 20$ . The factors of 20 in pairs are 1 and 20, 2 and 10, 4 and 5. The sign before 20 is  $+$ , hence select the factors whose *sum* is 9. These are 4 and 5. The sign before 9 is  $+$ , hence the required factors are  $(x+4)(x+5)$ .

2.  $x^2 - 8x + 12$ . Pairs of factors of 12 are 1 and 12, 2 and 6, 3 and 4. Sign before 12 is +, therefore take pair whose sum is 8. These are 2 and 6. Sign before 8 is -, hence the factors are  $(x-2)(x-6)$ .

3.  $x^2 - 21x - 100$ . Pairs of factors of 100 are 1 and 100, 2 and 50, 4 and 25, 5 and 20, 10 and 10. Sign before 100 is -, therefore take the pair whose difference is 21. These are 4 and 25. The sign before 21 is -, therefore  $x-$  goes before 25, the larger factor, and the factors are  $(x+4)(x-25)$ .

4.  $x^2 + 12x - 108$ . Pairs of factors of 108 are 1 and 108, 2 and 54, 3 and 36, 4 and 27, 6 and 18, 9 and 12. Sign before 108 is -, therefore take the pair whose difference is 12. These are 6 and 18. Sign before 12 is +, therefore  $x+$  goes before 18, the larger factor, and  $x-$  before 6, the other factor; hence the factors are  $(x-6)(x+18)$ .

NOTE.—It will be found convenient to write the factors in two columns, separated by a short space. Taking Ex. 2 above, proceed thus: Since the sign of the third term is +, write the sign of the second term (in this case -) above both columns.

$$\begin{array}{cc} \overline{1} & \overline{12} \\ (x-2) & (x-6) \end{array}$$

Ex. 3 above. Since the sign of the third term is -, write the sign of the 2nd term (in this case -) above the column of larger factors, and the other sign of the pair  $\pm$ , above the other column.

$$\begin{array}{cc} + & - \\ \overline{1} & \overline{100} \\ \overline{2} & \overline{50} \\ (x+4) & (x-25) \end{array}$$

5.  $x^2 - 34x + 64$ .

Here we have the factors

$$\begin{array}{cc} \overline{1} & \overline{64} \\ x-2, & x-32 \\ \overline{4}, & \overline{16} \end{array}$$

and since the last term has the sign +, and the middle term has the sign -, we write - over both columns.

6.  $x^2 + 12x - 64$ .

-	+
1,	64
2,	32
$x-4, x+16$ .	

Here, since the last term has the sign -, we write the sign (+) of the middle term, over the column of larger factors, and the sign - over the other column.

7.  $x^2 - 10x^2 - 144$ .

Here we have the pairs of factors :

+	-
1,	144
2,	72
4,	36
$x+8, x-18$ .	

And since the sign of the third term is -, we write the sign of the second term (in this case -) above the column of larger factors, and the other sign (of the pair ±) above the other column.

Exercise xxvi.

1.  $x^2 - 5x - 14$  ;  $x^2 - 9x + 14$  ;  $x^2 + 7x + 12$ .
2.  $x^2 - 8x + 15$  ;  $x^2 - 19x + 84$  ;  $x^2 - 7x - 60$ .
3.  $4x^2 - 2x - 20$  ;  $9x^2 - 150x + 600$ .
4.  $\frac{1}{4}x^2 + 4\frac{1}{4}x - 36$  ;  $25x^2 + 40x + 15$  ;  $9x^6 - 27x^3 + 20$ .
5.  $\frac{1}{16}x^2 + 1\frac{3}{4}x + 12$  ;  $16x^4 - 4x^2 - 20$ .
6.  $x^4 - (a^2 + b^2)x^2 + a^2b^2$  ;  $4(x+y)^2 - 4(x+y) - 99$ .
7.  $(x^2 + y^2)^2 - (a^2 - b^2)(x^2 + y^2) - a^2b^2$ .
8.  $(a+b)^2 - 2c(a+b) - 3c^2$ .
9.  $(x+y)^2 + 2(x^2 + y^2)(x+y) + (x^2 - y^2)^2$ .
10.  $(a+b)^2 - 4ab(a+b) - (a^2 - b^2)^2$ .
11.  $(x^2 + xy + y^2)^2 + x^3 - y^3 - 5xy - 2y^2 - 2x^2$ .
12.  $a^2 - 2a(b-c) - 3(b-c)^2$ .

13.  $(x^3 + y^3)^2 + 2a^3(x^3 + y^3) + a^6 - b^6$ .  
 14.  $(x^2 - 10x)^2 - 4(x^2 - 10x) - 96$ .  
 15.  $(x^2 - 14x + 40)^2 - 25(x^2 - 14x + 40) - 150$ .  
 16.  $(x^2 - xy + y^2)^2 + 2xy(x^2 - xy + y^2) - 3x^2y^2$ .  
 17.  $z^4 - 2z^2 + 2$ ;  $x^4 - 2x^2 - 3$ ;  $9x^8 + 9x^4y^2 - 10y^4$ .  
 18.  $c^{2m} + c^m - 2$ ;  $x^6 - x^3 - 2$ ;  $x^{2m} - 2x^m y^n - 8y^{2n}$ .  
 19.  $x^{2m} - (a-b)x^m y^n - aby^{2n}$ .

**Art. XV.** Trinomials of the form  $ax^2 + bx + c$  ( $a$  not a square) may sometimes be easily factored from the following considerations:—

The product of two binomials consists of

1st. The product of the *first* terms.

2nd. “ “ *second* “

3rd. The sum (*algebraic*) of the products of the terms taken diagonally.

These three conditions guide us in the converse process of resolving a trinomial into its binomial factors.

#### EXAMPLES.

1. Resolve  $6x^2 - 13xy + 6y^2$ .

Here the factors of the first term are  $x$  and  $6x$ , or  $2x$  and  $3x$ ; those of the third term are  $y$  and  $6y$ , or  $2y$  and  $3y$ . These pairs of factors may be arranged

(1)	(2)	(3)	(4)
$x$	$y$	$2x$	$2y$
$6x$	$3x$	$6y$	$3y$

Now, we may take (1) with (3) or (4), or (2) with (3) or (4); but none of these combinations will satisfy the third condition. If, however, in (4) we interchange the coefficients 2 and 3, then (2) and (4) give

$2x$      $3y$ , and

$3x$      $2y$ , where we can combine the “diagonal”

products to make 13, and the factors are

$$2x - 3y, \text{ and}$$

$$3x - 2y.$$

The coefficients of (2), instead of those of (4), might have been interchanged, giving the same result.

$$2. 6x^2 - 15xy + 6y^2.$$

Here, comparing (2) and (3), Ex. 1, we see that their diagonal products may be combined to give 15, and the factors are

$$2x - y, \text{ and } 3x - 6y.$$

$$3. 6x^2 - 20xy + 6y^2.$$

Here, again referring to Ex. 1, we see at once that it is useless to try *both* (2) and (4), since the diagonal products cannot be combined in any way to give a higher result than  $13xy$ . But comparing (1) and (4), we obtain by interchanging the coefficients in (4)

$$x - 3y, \text{ and}$$

$$6x - 2y, \text{ which satisfy the third condition.}$$

Or, we might interchange the coefficients of (3), and take the resulting terms with (2), getting  $2x - 6y$ , and

$$3x - y.$$

$$4. 6x^2 + 35xy - 6y^2.$$

Here the large coefficient of the middle term shows at once that we must take (1) and (3) together. Interchanging the coefficients of (1) we have

$$6x - y, \text{ and}$$

$x + 6y$ . The same result will be obtained by interchanging the coefficients of (3).

#### Exercise xxvii.

$$1. 6x^2 - 37xy + 6y^2.$$

$$2. 6x^2 + 9xy - 6y^2.$$

$$3. 56x^2 - 76xy + 20y^2.$$

$$4. 56x^2 - 36xy - 20y^2.$$

$$5. 56x^2 - 1121xy + 20y^2.$$

$$6. 56x^2 - 68xy + 20y^2.$$

$$7. 56x^2 - 558xy - 20y^2.$$

$$8. 56x^2 + 36xy - 20y^2.$$

$$9. 56x^2 - 67xy + 20y^2.$$

$$10. 56x^2 + 3xy - 20y^2.$$

$$11. 6x^2 - 16xy - 6y^2.$$

$$12. 6x^2 + 5xy - 6y^2.$$

$$13. 56x^2 + 562xy + 20y^2.$$

$$14. 56x^2 - 122xy + 20y^2.$$

$$15. 56x^2 - 102xy - 20y^2.$$

$$16. 56x^2 - 229xy + 20y^2.$$

$$17. 56x^2 - 94xy + 20y^2.$$

$$18. 56x^2 - 276xy - 20y^2.$$

$$19. 36x^2 - 33xy - 15y^2.$$

$$20. 72x^2 - 19xy - 40y^2.$$



**Art. XVI.** More generally, trinomials of the form  $ax^2+bx+c$  ( $a$  not a square) may be resolved by Formula A, thus

Multiplying by  $a$  we get  $a^2x^2+baax+ac$ . Writing  $z$  for  $ax$  this becomes  $z^2+bz+ac$ . Factor this trinomial, restore the value of  $z$  and divide the result by  $a$ .

## EXAMPLES.

1.  $6x^2+5x-4$ . Multiplying by 6, we get  $(6x)^2+5(6x)-24$  or  $z^2+5z-24$ . Factoring, we get  $(z-3)(z+8)$ , hence the required factors are  $\frac{1}{6}(6x-3)(6x+8) = (2x-1)(3x+4)$ .

2.  $6x^2-13xy+6y^2$ . Factoring  $z^2-13zy+36y^2$  we get  $(z-4y)(z-9y)$ , hence the required factors are  $\frac{1}{6}(6x-4y)(6x-9y) = (3x-2y)(2x-3y)$ .

3.  $33-14x-40x^2$ . Factoring  $1320-14z-z^2$  we get  $(30-z)(44+z)$ , hence the required factors are  $\frac{1}{40}(30-40x)(44+40x) = (3-4x)(11+10x)$ .

**NOTE.**—The factors may conveniently be arranged in two columns, each with its appropriate sign above it.

	—	+
Ex. 1, above	1	24
	2	13

$$\frac{1}{6}(6x-3)(6x+8) = (2x-1)(3x+4).$$

	—	—
Ex. 2, above	1	36
	2	18
	3	12

$$\frac{1}{6}(6x-4)(6x-9) = (3x-2)(2x-3).$$

[Another method of factoring trinomials of the form  $ax^2+bx+c$  is as follows :

Multiply by  $4a$ , thus obtaining  $4a^2x^2+4abx+4ac$ . Add  $b^2-b^2$ , which will not change the value,  $4a^2x^2+4abx+b^2-b^2+4ac$ ; by [1] this may be written  $(2ax+b)^2-(b^2-4ac)$ . Factor this by [4] and divide the result by  $4a$ .

*Ex.* Factor  $56x^2 + 137x - 27885$ . Multiply by  $4 \times 56$  or  $2 \times 112$ ,  $112^2x^2 + 2.137.112x - 6246240$ . Add  $137^2 - 137^2$ , then  $112^2x^2 + 2.137.112x + 137^2 - (137^2 + 6246240) = (112x + 137)^2 - 6265009 = \{(112x + 137) + 2503\} \{(112x + 137) - 2503\} = (112x + 2640)(112x - 2366)$ .

We multiplied by  $4 \times 56$ , we must, therefore, now divide by that number. Doing so, we obtain as factors  $(7x + 165)(8x - 169)$ .]

## Exercise xxviii.

- |                               |                                      |
|-------------------------------|--------------------------------------|
| 1. $10x^2 + x - 21$           | 9. $12x^2 - x - 1$                   |
| 2. $10x^2 - 29x - 21$         | 10. $9x^2y^2 - 3xy^2 - 6y^4$         |
| 3. $10x^2 + 29x - 21$         | 11. $4x^2 + 8xy + 3y^2$              |
| 4. $6x^2 - 37x + 55$          | 12. $6b^2x^2 - 7bx^2 - 3x^4$         |
| 5. $12a^2 - 5a - 2$           | 13. $6x^4 - x^2y^2 - 35y^4$          |
| 6. $12x^2 - 37x + 21$         | 14. $2x^4 + x^2 - 45$                |
| 7. $12x^2 + 37x + 21$         | 15. $4x^4 - 37x^2y^2 + 9y^4$         |
| 8. $15a^6 + 13a^3b^2 - 20b^4$ | 16. $4(x+2)^4 - 37x^2(x+2)^2 + 9x^4$ |
17.  $6(2x+3y)^2 + 5(6x^2+5xy-6y^2) - 6(3x-2y)^2$ .  
 18.  $6(2x+3y)^4 + 5(6x^2+5xy-6y^2)^2 - 6(3x-2y)^4$ .  
 19.  $6(x^2+xy+y^2)^2 + 13(x^4+x^2y^2+y^4) - 385(x^2-xy+y^2)^2$ .  
 20.  $21(x^2+2xy+2y^2)^2 - 6(x^2-2xy+2y^2)^2 - 5(x^4+4y^4)$ .

## SECTION II.—EXTENDED APPLICATION OF THE FORMULAS.

**Art. XVII.** The methods of factoring just explained may be applied to find the rational factors, where such exist, of quadratic multinomials.

## EXAMPLES.

1. Resolve  $12x^2 - xy - 20y^2 + 8x + 41y - 20$ .

In the first place we find the factors of the *first three terms*, which are

$$4x + 5y, \text{ and} \\ 3x - 4y.$$

Now, to find the *remaining terms* of the required factors, we must observe the following conditions:

1st. Their product must = -20.

2nd. The sum (*algebraic*) of the products obtained by multiplying them diagonally into the  $y$ 's, must =  $41y$ .

3rd. The sum of the products obtained by multiplying them diagonally into the  $x$ 's, must =  $8x$ .

We see at once that -4 with the first pair already found, and +5 with the second pair, satisfy the required conditions, and  $\therefore$  the factors are

$$4x + 5y - 4, \text{ and} \\ 3x - 4y + 5.$$

2.  $p^2 + 2pr - 2q^2 + 7qr - 3r^2 + pq$ .

Here the factors of  $p^2 + pq - 2q^2$ , are

$$p + 2q, \text{ and} \\ p - q.$$

Now find two factors which will give  $-3r^2$ , and which multiplied diagonally into the  $p$ 's and  $q$ 's respectively, will give  $2pr$ , and  $7qr$ ; these are found to be  $-r$  taken with the *first* pair, and  $+3r$  taken with the second pair. Hence the required factors are

$$p + 2q - r, \text{ and} \\ p - q + 3r.$$

**Art. XVIII.** But the following examples illustrate a surer method.

3.  $x^2 + xy - 2y^2 + 2xz + 7yz - 3z^2$ .

Reject 1st the terms involving  $z$ ,

2nd. " "  $y$ ,

3rd. " "  $x$ .

and factor the expression that remains in each case.

1st.  $x^2 + xy - 2y^2 = (x - y)(x + 2y)$ .

2nd.  $x^2 + 2xz - 3z^2 = (x + 3z)(x - z)$ .

3rd.  $-2y^2 + 7yz - 3z^2 = (-y + 3z)(2y - z)$ .

Arrange these three pair of factors in two sets of three factors each, by so selecting one factor from each pair that two of each set of three may have the same coefficient of  $x$ , two may have the

same coefficient of  $y$ , and two the same coefficient of  $z$  (coefficients including sign). In this example there are

$$x - y, x + 2z, -y + 2z,$$

$$\text{and } x + 2y, x - z, 2y - z.$$

From the first set select the common terms (including signs) and form therewith a trinomial,  $x - y + 2z$ .

Repeat with the second set, and we get  $x + 2y - z$ .

$$\therefore x^2 + xy - 2y^2 + 2xz + 7yz - 3z^2 = (x - y + 2z)(x + 2y - z).$$

4.  $3x^2 - 8xy - 3y^2 + 30x + 27$ .

1st.  $3x^2 - 8xy - 3y^2 = (3x + y)(x - 3y)$ .

2nd.  $3x^2 + 30x + 27 = (3x + 3)(x + 9)$ .

3rd.  $-3y^2 + 27 = (y + 3)(-3y + 9)$ .

$\therefore$  the factors are  $(3x + y + 3)(x - 3y + 3)$ .

5.  $6a^2 - 7ab + 2ac - 20b^2 + 64bc - 48c^2$ .

1st.  $6a^2 - 7ab - 20b^2 = (2a - 5b)(3a + 4b)$ .

2nd.  $6a^2 + 2ac - 48c^2 = (2a + 6c)(3a - 8c)$ .

3rd.  $-20b^2 + 64bc - 48c^2 = (-5b + 6c)(4b - 8c)$ .

$\therefore$  the factors are  $(2a - 5b + 6c)(3a + 4b - 8c)$ .

### Exercise xxix.

1.  $7x^2 - xy - 6y^2 - 6x - 20y - 16$ .

2.  $20x^2 - 15xy - 5y^2 - 68x - 42y - 88$ .

3.  $3x^4 + x^2y^2 - 4y^4 + 10x^2 - 17y^2 - 13$ .

4.  $20x^2 - 20y^2 + 9xy + 28x + 35y$ .

5.  $72x^2 - 8y^2 + 55xy + 12y - 169x + 20$ .

6.  $x^2 - xy - 12y^2 - 5x - 15y$ .

7.  $8x^2 + 18xy + 9y^2 + 2xz - z^2$ .

8.  $6x^2 + 6y^2 - 13xy - 8z^2 - 2yz + 8xz$ .

9.  $6x^4 - 10y^4 + 11x^2y^2 - 25z^2 + 10y^2 + 25y^2z^2 - 15x^2 + 10x^2z^2$ .

10.  $15x^4 - 16y^4 - 22x^2y^2 + 15z^4 + 14y^2z^2 + 50x^2z^2$ .

11.  $4a^2 - 15b^2 - 4ab - 21c^2 - 36bc - 8ac$ .

12.  $a^4 + b^4 + c^4 - 2a^2b^2 - 2b^2c^2 - 2c^2a^2$ .

**Art. XIX.** Trinomials of the form  $ax^4 + bx^2 + c$  can always be broken up into real factors.

If  $a$  and  $c$  have different signs, the expression may be factored by Art. XVI.

If  $a$  and  $c$  are of the same sign, three cases have to be considered: i.  $b = 2\sqrt{ac}$ , ii.  $b > 2\sqrt{ac}$ , iii.  $b < 2\sqrt{ac}$

**Case I.**  $b = 2\sqrt{ac}$ . This case falls under Art XII., formula [1]. where examples will be found.

**Case II.**  $b > 2\sqrt{ac}$ . This case falls under Art XVI., where examples will be found. The following additional examples are resolved by the second method of that article.

#### EXAMPLES.

1.  $4x^4 + 5x^2y^2 + y^4$ .

Here we see that  $(\frac{5}{4}y^2)^2$  will make, with the first two terms, a perfect square, and we therefore add to the given expression  $(\frac{5}{4}y^2)^2 - (\frac{5}{4}y^2)^2$ . The expression then becomes

$$\begin{aligned} & 4x^4 + 5x^2y^2 + (\frac{5}{4}y^2)^2 + y^4 - (\frac{5}{4}y^2)^2 \\ &= (2x^2 + \frac{5}{4}y^2)^2 - \frac{9}{16}y^4 \\ &= (2x^2 + \frac{5}{4}y^2 + \frac{3}{4}y^2)(2x^2 + \frac{5}{4}y^2 - \frac{3}{4}y^2) \\ &= (2x^2 + 2y^2)(2x^2 + \frac{1}{2}y^2) = (x^2 + y^2)(4x^2 + y^2). \end{aligned}$$

2.  $3x^4 + 6x^2 + 2$ .

Here multiplying by  $4 \times 3$ , and completing the square as in Ex. 1, we have

$$\begin{aligned} & 36x^4 + 72x^2 + 6^2 + 24 - 6^2 = (6x^2 + 6)^2 - 12 \\ &= (6x^2 + 6 - \sqrt{12})(6x^2 + 6 + \sqrt{12}), \text{ which divided by } 4 \times 3 \text{ give} \\ & \text{the required factors.} \end{aligned}$$

3.  $ax^4 + bx^2 + c$ .

Proceeding as in Ex. 2 we have, by multiplying by  $4a$ ,

$$\begin{aligned} & ax^4 + bx^2 + c = \{4a^2x^4 + 4abx^2 + b^2 - b^2 + 4ac\} \div 4a \\ &= \{2ax^2 + b + \sqrt{(b^2 - 4ac)}\} \{2ax^2 + b - \sqrt{(b^2 - 4ac)}\} \div 4a. \end{aligned}$$

## Exercise. xxx.

1.  $x^4 + 7x^2 + 1$ ;  $4x^4 + 14x^2 + 1$ .
2.  $x^4 + 7x^2y^2 + y^4$ ;  $3x^4 + 5x^2y^2 + y^4$ .
3.  $4x^4 + 10x^2 + 3$ ;  $3(x+y)^4 + 5z^2(x+y)^2 + z^4$ .
4.  $x^4 + 7x^2y^2 + 3\frac{1}{4}y^4$ ;  $x^4 + 7x^2y^2 + 8\frac{1}{4}y^4$ .
5.  $4x^4 + 9x^2y^2 + \frac{1}{16}y^4$ ;  $4(a+b)^4 + 10c^2(a+b)^2 + 3c^4$ .
6.  $3x^4 + 8x^2y^2 + 4\frac{7}{12}y^4$ ;  $36x^4 + 96x^2 + 55$ .
7.  $5x^4 + 20x^2 + 2$ ;  $4a^4 + 12a^2 + 1$ .
8.  $4(x+y)^4 + 12(x+y)^2z^2 + z^4$ ;  $5x^4 + 20x^2y^2 + 2y^4$ .
9.  $9x^4 + 14x^2 + 4$ ;  $2x^4 + 12x^2(y+z)^2 + 15(y+z)^4$ .
10.  $2x^4 + 12x^2 + 15$ ;  $7x^4 + 40x^2 + 45$ .
11.  $8x^4 + 36x^2y^2 + 29y^4$ ;  $7x^4 + 20x^2y^2 - 20y^4$ .
12.  $7(a-b)^4 + 16(a-b)^2c^2 + 5c^4$ ;  $\frac{3}{2}a^4 + 3a^2b^2 + b^4$ .
13.  $3x^4 + 6x^2y^2 + 2y^4$ ;  $3(a+b)^4 + 6(a^2 - b^2)^2 + 2(a-b)^4$ .
14.  $49a^4 - 84a^2b^2 + 22b^4$ ;  $25m^4 + 60m^2n^2 + 27n^4$ .
15.  $49(m+n)^4 - 84(m^2 - n^2)^2 + 22(m-n)^4$ .

**Case III.**  $b < 2\sqrt{ac}$ . This case may be brought under Art. XIII. The following examples illustrate the process of reduction and resolution.

## EXAMPLES.

$$1. x^4 - 7x^2 + 1.$$

We have to throw this into the form  $a^2 - b^2$ :

$$x^4 - 7x^2 + 1 = (x^2 + 1)^2 - 9x^2 = (x^2 + 1 + 3x)(x^2 + 1 - 3x).$$

$$2. 9x^4 + 3x^2y^2 + 4y^4 = (3x^2 + 2y^2)^2 - 9x^2y^2 \\ = (3x^2 + 2y^2 - 3xy)(3x^2 + 2y^2 + 3xy).$$

$$3. x^4 + y^4 = (x^2 + y^2)^2 - 2x^2y^2 \\ = (x^2 + y^2 + xy\sqrt{2})(x^2 + y^2 - xy\sqrt{2}).$$

$$4. x^4 - \frac{1}{4}x^2y^2 + y^4 = (x^2 + y^2)^2 - \frac{3}{4}x^2y^2 \\ = (x^2 + y^2 + \frac{3}{2}xy)(x^2 + y^2 - \frac{3}{2}xy).$$

$$5. ax^4 + bx^2 + c = (\sqrt{a} \cdot x^2 + \sqrt{c})^2 - \{2\sqrt{ac} - b\}x^2 \\ = \{\sqrt{a} \cdot x^2 + \sqrt{c} - \sqrt{(2\sqrt{ac} - b)}x\} \times \\ \{\sqrt{a} \cdot x^2 + \sqrt{c} + \sqrt{(2\sqrt{ac} - b)}x\}.$$

**Art. XX.** It is seen from these examples that we have merely to add to the given expression what will make with the *first and last* terms (arranged as in Ex. 5) a perfect square, and to subtract the same quantity. In Ex. 2, *e. g.*, the square root of  $9x^4 = 3x^2$ , the square root of  $4y^4 = 2y^2$ ,  $\therefore 3x^2 + 2y^2$  is the binomial whose square is required; we need  $\therefore 12x^2y^2$ ; but the expression contains  $3x^2y^2$ ;  $\therefore$  we have to *add and subtract*  $12x^2y^2 - 3x^2y^2 = 9x^2y^2$ .

Hence we derive a practical rule for factoring such expressions.

(1). Take the square roots of the two extreme terms and connect them by the proper sign; this gives the first two terms of the required factors.

(2) Subtract the middle term of the given expression from twice the product of these two roots, and the square roots of the difference will be the third terms of the required factors.

6.  $x^4 + \frac{7}{6}x^2y^2 + y^4$ . Here  $\sqrt{x^4} = x^2$ ,  $\sqrt{y^4} = y^2$ , and the first two terms of the required factors are  $x^2 + y^2$ ; twice the product of these is  $+2x^2y^2$ , from which subtracting the middle term,  $\frac{7}{6}x^2y^2$ , we get  $\frac{5}{6}x^2y^2$ ; the square roots of this are  $\pm\frac{5}{6}xy$ . Hence the factors are  $x^2 + y^2 \pm \frac{5}{6}xy$ .

Note that since  $\sqrt{y^4} = +y^2$ , or  $-y^2$ , it may sometimes happen that while the former sign will give irrational factors, the latter will give rational factors, and conversely.

7.  $x^4 - 11x^2y^2 + y^4$ . Here, taking  $+y^2$ , we have

$$x^2 + y^2 + xy\sqrt{13}, \text{ and } x^2 + y^2 - xy\sqrt{13}.$$

But taking  $-y^2$ , we have

$$x^2 - y^2 + 3xy, \text{ and } x^2 - y^2 - 3xy.$$

Sometimes *both* signs will give rational factors.

8.  $16x^4 - 17x^2y^2 + y^4$ . Here we have

$$(4x^2 + y^2 + 3xy)(4x^2 + y^2 - 3xy), \text{ and also} \\ (4x^2 - y^2 + 5xy)(4x^2 - y^2 - 5xy).$$

## Exercise xxxi.

1.  $x^4 + 2x^2y^2 + 9y^4$ ,  $x^4 - x^2y^2 + y^4$ ,  $x^4 + x^2y^2 + y^4$ .
2.  $x^4 + 4y^4$ ,  $16x^4 + y^4 - x^2y^2$ ,  $\frac{1}{4}x^4 + y^4$ .
3.  $x^4 + 1$ ,  $x^4 + 9y^4$ ,  $1 - 12y^2 + 16y^4$ .
4.  $x^4 - 7x^2 + 1$ ,  $x^4 + 9$ ,  $\frac{1}{4}x^4 + y^4 - 3x^2y^2$ .
5.  $y^4 - x^4 + 11x^2y^2$ ,  $x^8 + 4y^8$ ,  $x^4 + 4x^2 + 16$ .
6.  $4x^4 + y^4 - 8\frac{1}{4}x^2y^2$ ,  $x^4 + y^4 - \frac{7}{16}x^2y^2$ ,  $4x^4 + 1$ .
7.  $x^{4m} + 64y^{4m}$ ,  $x^{4m} + 4y^{4m}$ ,  $\frac{1}{4}x^4 + \frac{9}{16}y^4 - 5\frac{3}{4}x^2y^2$ .
8.  $4x^4 - 8x^2 + 1$ ,  $7x^2y^2 - \frac{1}{4}x^4 - 36y^4$ ,  $x^4 + a^4y^4$ .
9.  $m^2x^4 + n^2y^4 - (2mi + p)x^2y^2$ ,  $x^{4m} + 2^{4m-2}y^{4m}$ .
10.  $16x^4 - 25x^2 + 9$ ,  $4x^4 - 16x^2 + 4$ ,  $13x^2y^2 - 9x^4 - 4y^4$ .
11.  $4x^4 - 12\frac{1}{2}x^2y^2 + 9y^4$ ,  $x^4 + 6x^2 + 25$ .
12.  $a^4 + b^4 + (a+b)^4$ ,  $1 + a^4 + (1+a)^4$ .
13.  $(x+y)^4 - 7z^2(x+y)^2 + z^4$ .
14.  $(a+b)^4 + 7c^2(a+b)^2 + c^4$ .
15.  $16a^4 + 4(b-c)^4 - 9a^2(b-c)^2$ .
16.  $4(a+b)^4 + 9(a-b)^4 - 21(a^2 - b^2)^2$ .
17.  $(x^2 + y^2 - xy)^4 - 7(x^3 + y^3)^2 + (x+y)^4$ .
18.  $(a^2 + ab + b^2)^4 + 7(a^3 - b^3)^2 + (a-b)^4$ .
19.  $16a^4 + 4a^2 + 1$ ,  $x^4 - 41x^2 + 16$ .
20.  $x^4 + 81y^8 - 63x^2y^4$ ,  $1 + z^4 + 25z^8$ .
21.  $(a^2 + 1)^4 + 4(a^2 + 1)^2a^2 + 16a^4$ ,  $(x+1)^4 + 2(x^2 - 1)^2 + 9(x-1)^4$ .

Art. XXI. We can apply [4], Art. XIII., to factor expressions of the form  $ax^4 + bx^3 + rbx - r^2a$ . This may be written

$$a(x^4 - r^2) + bx(x^2 + r) = \{a(x^2 - r) + bx\}(x^2 + r).$$

## EXAMPLES.

1.  $6x^4 + 4x^3 + 12x - 54$ . This  
 $= 6(x^4 - 9) + 4x(x^2 + 3) = (x^2 + 3)\{6(x^2 - 3) + 4x\}$   
 $= (x^2 + 3)(6x^2 + 4x - 18)$ .



2.  $11x^4 + 10x^3 - 10x - 176$ . This  
 $= 11(x^4 - 16) + 10x(x^2 - 4) = (x^2 - 4)\{11(x^2 + 4) + 10x\}$   
 $= (x^2 - 4)(11x^2 + 10x + 4)$ .
3.  $40x^4 + 30x^3 + 60x - 160$ . This  
 $= 10(4x^4 - 16) + 15x(2x^2 + 4) = (2x^2 + 4)\{10(2x^2 - 4) + 15x\}$   
 $= (2x^2 + 4)(20x^2 + 15x - 40)$ .

NOTE.—To determine  $r$ , take the ratio of the coefficient of  $x^3$  to the coefficient of  $x$ .

### Exercise xxxii.

Resolve into factors

- |   |   |
|---|---|
| 1. $x^4 + 2x^3 + 6x - 9$ .                                    | 13. $\frac{1}{4}x^4 + \frac{1}{10}x^3 - \frac{2}{5}x - \frac{4}{5}$ . |
| 2. $2x^4 + 2x^3 + 6x - 18$ .                                  | 14. $80x^4 - 32x^3y + 64xy - 320y^4$ .                                |
| 3. $x^4 + 3x^3 + 12x - 16$ .                                  | 15. $24x^4 - 12x^3y + 30xy^3 - 150y^4$ .                              |
| 4. $3x^4 + x^3 - 4x - 48$ .                                   | 16. $2x^4 + \frac{1}{2}x^3y - 8xy^3 - 512y^4$ .                       |
| 5. $5x^4 + 4x^3 - 12x - 45$ .                                 | 17. $11x^4 + 10x^3 - 12x - 15\frac{1}{2}$ .                           |
| 6. $10x^4 + 5x^3 + 30x - 360$ .                               | 18. $40x^4 + 30x^3 + 60x - 160$ .                                     |
| 7. $\frac{1}{4}x^4 + 20x^3 + 4x - 10\frac{1}{5}$ .            | 19. $13x^4 - 12x^3y + 72xy^3 - 468y^4$ .                              |
| 8. $25x^4 - 40x^3 + 8x - 1$ .                                 | 20. $3x^4 + 3x^3y + 12xy^3 - 48y^4$ .                                 |
| 9. $37\frac{1}{2}x^4 - 30x^3 + 48x - 96$ .                    | 21. $5x^4 + 4x^3y - 12xy^3 - 45y^4$ .                                 |
| 10. $63x^4 - 39x^3 + 52x - 112$ .                             | 22. $4x^4 - 14x^3y + 28xy^3 - 16y^4$ .                                |
| 11. $810x^4 + \frac{8}{4}x^3 + \frac{2}{5}x - 2\frac{1}{2}$ . | 23. $x^4 + 80x^3y + 16xy^3 - \frac{1}{2}y^4$ .                        |
| 12. $242x^4 - 33x^3 - 3x - 2$ .                               | 24. $2x^4 - x^3y + 6xy^3 - 72y^4$ .                                   |

Art. XXII. Formulas [1] and [4] may sometimes be applied to factor expressions of the form

$$ax^4 + bx^3 + cx^2 + rbx + r^2a.$$

This may be put under the form

$$a(x^4 + r^2) + bx(x^2 + r) + cx^2 = a(x^2 + r)^2 + bx(x^2 + r) + (c - 2ar)x^2, \text{ which can sometimes be factored.}$$

#### EXAMPLES.

1.  $x^4 + 6x^3 + 27x^2 + 162x + 729$ .

We have  $x^4 + 729 + 6x(x^2 + 27) + 27x^2$ .

$$= (x^2 + 27)^2 + 6x(x^2 + 27) + 9x^2 - 36x^2$$

$$= \{x^2 + 27 + 3x\}^2 - 36x^2, \text{ which gives the factors}$$

$$x^2 - 3x + 27, \text{ and } x^2 + 9x + 27.$$

$$\begin{aligned}
 2. \quad & x^4 + 4x^3 + 4x^2 + 20x + 25. \quad \text{This} \\
 & = (x^2 + 5)^2 + 4x(x^2 + 5) - 6x^2 \\
 & = (x^2 + 5)^2 + 4x(x^2 + 5) + 4x^2 - 10x^2 \\
 & = \{x^2 + 5 + 2x - x\sqrt{10}\} \{x^2 + 5 + 2x + x\sqrt{10}\}.
 \end{aligned}$$

## Exercise xxxiii.

Resolve into factors.

1.  $x^4 - 6x^3 + 27x^2 - 162x + 729.$
2.  $x^4 + 2x^3 + 3x^2 + 8x + 16.$
3.  $x^4 + x^3 + x^2 + x + 1.$
4.  $x^4 - 4x^3 + x^2 - 4x + 1.$
5.  $4x^4 - 12x^3 - 6x^2 - 12x + 4.$
6.  $x^4 + 14x^3 - 25x^2 - 70x + 25.$
7.  $16x^4 - 24x^3 - 16x^2 + 12x + 4.$
8.  $x^4 + 5x^3 - 16x^2 + 20x + 16.$
9.  $x^4 + 6x^3 - 11x^2 - 12x + 4.$
10.  $x^4 + 4x^3y + x^2y^2 + 12xy^3 + 9y^4.$
11.  $x^4 + 6x^3 - 9x^2 - 6x + 1.$
12.  $x^4 + 4x^3y - 19x^2y^2 + 4xy^3 + y^4.$
13.  $4x^4 + 4x^3y - 65x^2y^2 - 10xy^3 + 25y^4.$
14.  $x^4 + 6x^3y - 9x^2y^2 - 6xy^3 + y^4.$
15.  $x^4 + 6x^3y + 10x^2y^2 + 12xy^3 + 4y^4.$
16.  $9x^4 + 18x^3y - 52x^2y^2 - 12xy^3 + 4y^4.$
17.  $11x^4 + 10x^3y + 39x^2y^2 + 20xy^3 + 44y^4.$

## SECTION III.—FACTORIZING BY PARTS.

**Art. XXIII.** To factor an expression which can be reduced to the form  $a.F(x) + b.f(x).$

When the expression is thus arranged, any factor common to  $a$  and  $b$ , or to  $F(x)$  and  $f(x)$ , will be a factor of the whole expression. The method about to be illustrated will be found useful in cases where only *one* power of some letter is found.

## EXAMPLES.

1. Factor
- $acx^2 - abx - bc^2x + b^2c$
- .

Here we see that only one power of  $a$  occurs, and we therefore group together the terms involving this letter, and those not involving it, getting

$$\begin{aligned} & a(cx^2 - bx) - bc^2x + b^2c \\ &= ax(cx - b) - bc(cx - b) = (ax - bc)(cx - b). \end{aligned}$$

2. Factor
- $m^2x^2 - mna^2x - mn^2x + n^2a^2$
- .

Here we observe that  $a$  occurs in only one power ( $a^2$ ).

Therefore we have

$$\begin{aligned} & -a^2(mnx - n^2) + m^2x^2 - mn^2x \\ &= -na^2(mx - n) + mx(mx - n) \\ &= (mx - n)(mx - na^2). \end{aligned}$$

- 3.
- $2x^2 + 4ax + 3bx + 6ab$
- .

Here we observe that the expression contains only one power of both  $a$  and  $b$ . We may, therefore, collect the coefficients in either of the following ways :

$$\begin{aligned} & a(4x + 6b) + (2x^2 + 3bx), \\ \text{or, } & b(3x + 6a) + (2x^2 + 4ax). \end{aligned}$$

Now the expressions in the brackets ought to have a common factor, and we see that this is the case. Hence,

$$\begin{aligned} & a(4x + 6b) + (2x^2 + 3bx) \\ &= 2a(2x + 3b) + x(2x + 3b) = (2x + 3b)(x + 2a). \end{aligned}$$

- 4.
- $abxy + b^2y^2 + acx - c^2$
- 
- $= a(bxy + cx) + b^2y^2 - c^2$
- 
- $= ax(by + c) + (by + c)(by - c) = (by + c)(ax + by - c).$

- 5.
- $y^3 - (2a + b)y^2 + (2ab + a^2)y - a^2b$
- 
- $= -b(y^2 - 2ay + a^2) + y^3 - 2ay^2 + a^2y$
- 
- $= -b(y^2 - 2ay + a^2) + y(y^2 - 2ay + a^2)$
- 
- $= (y - b)(y - a)^2.$

- 6.
- $2x^3y + 2bx^4 - bx^5y + 4abx^2y - x^2y^2 + 4axy^2 - 2abxy^2 - 2ay^3$
- .
- 
- $= b(2x^4 - x^3y + 4ax^2y - 2axy^2) + 2x^3y - x^2y^2 + 4axy^2 - 2ay^3$
- 
- $= bx(2x^3 - x^2y + 4axy - 2ay^2) + y(2x^3 - x^2y + 4axy - 2ay^2)$
- 
- $= (y + bx)(2x^3 - x^2y + 4axy - 2ay^2).$

And  $2x^3 - x^2y + 4axy - 2ay^2$   
 $= a(4xy - 2y^2) + 2x^3 - x^2y$   
 $= 2ay(2x - y) + x^2(2x - y) = (2ay + x^2)(2x - y).$

7.  $x^3 + (2a - b)x^2 - (2ab - a^2)x - a^2b$   
 $= b(-x^2 - 2ax - a^2) + x^3 + 2ax^2 + a^2x$   
 $= -b(x + a)^2 + x(x + a)^2 = (x - b)(x + a)^2.$

8.  $px^3 - (p - q)x^2 + (p - q)x + q$   
 $= q(x^2 - x + 1) + px^3 - px^2 + px$   
 $= q(x^2 - x + 1) + px(x^2 - x + 1) = (px + q)(x^2 - x + 1).$

Exercise xxxiv.

- |  |                                      |
|--|--------------------------------------|
| 1. $x^2y - x^2z - y^2 + yz.$   | 6. $x^3 - b^2x^2 - a^2x + a^2b^2.$   |
| 2. $abxy + b^2y^2 + acx - c^2.$  | 7. $x^5 - a^3x^2 - b^2x^3 + a^3b^2.$ |
| 8. $x^2z^2 + ax^2 - a^2z^2 - a^3.$   | 8. $8x^2 + 12ax + 10/x + 15ab.$      |
| 4. $2x^2 - ax - 4bx + 2ab.$  | 9. $a^2 + (ac - b^2)x^2 + bcx^3.$    |
| 6. $x^2 + 2bx + 3ac + 6ab.$  | 10. $a^2 + (ac - b^2)x^2 - bcx^3.$   |
| 11. $abx^3 + (ac - bd)x^2 - (af + cd)x + df.$  |                                      |
| 12. $px^3 - (p + q)x^2 + (p + q)x - q.$  |                                      |
| 13. $a^2 + ab + 2ac - 2b^2 + 7bc - 3c^2.$  |                                      |
| 14. $x^3 + (a + 1)x^2 + (a + 1)x + a.$   |                                      |
| 15. $mpx^3 + (mq - np)x^2 - (mr + nq)x + nr.$  |                                      |
| 16. $x^3 - (a + b + c)x^2 + (ab + bc + ac)x - abc.$  |                                      |
| 17. $x^3 + (a - b - c)x^2 - (ab - bc + ca)x + abc.$  |                                      |
| 18. $x^3 + (a + b - c)x^2 - (bc - ca - ab)x - abc.$  |                                      |
| 19. $a^2x^3 - a^3x^2y - a^2xy + a^3y^2 - ax^2yz - a^2z - xyz + ay^2z.$                         |                                      |
| 20. $a^2bx^2 + ab^2xy + acdxy + bcdy^2 - aefxz - bcfyz.$                                       |                                      |
| 21. $a^2x^3 - a(b - c)x^2 + c(a - b)x + c^2.$  |                                      |
| 22. $mx^3 - nx^2y + rx^2z - mxy^2 + ny^3 - ry^2z.$   |                                      |
| 23. $amx^2 + (mby - nay + mcz)x - nby^2 - ncyz.$   |                                      |
| 24. $(am - bcm)x^2 + (am - bcn)x + an + nax.$  |                                      |
| 25. $a^2b^2c^2 - b^2c^2xy - a^2c^2yz + c^2xy^2z - a^2b^2zx + b^2x^2yz + a^2z^2xy - x^2y^2z^2.$ |                                      |
| 26. $x^5 - m^2x^4 - (n - n^2)x^3 + (m^2n - m^2n^2)x^2 - a(x^2 + n^2 - n).$                     |                                      |
| 27. $1 - (a - 1)x - (a - b + 1)x^2 + (a + b - c)x^3 - (b + c)x^4 + cx^5.$                      |                                      |
| 28. $a^3x^3 - a^2(b - c + d)x^2y - (abc - abd + acd)xy^2 + b^2ay^2.$                           |                                      |
| 29. $m^2npix^3 - (n^2p - m^2n^2 - m^2py)x^2 - (n^3 + npq - m^2nq)x - n^2q.$                    |                                      |

$$30. m^2p^2x^5 + m^2p^2x^4 - (p^2n^2 - q^2m^2)x^3y^2 - (p^2n^2 - q^2m^2)x^2y^2 - (n^2q^2 + n^2q^2x)y^4.$$

**Art. XXIV.** Sometimes an expression which does not come directly under the preceding form, may be resolved by first finding the factors of its parts.

## EXAMPLES.

$$1. abx^2 + aby^2 - a^2xy - b^2xy.$$

Here, taking  $ax$  out of the first and third terms, and  $by$  out of the second and fourth terms, we have

$$ax(bx - ay) - by(bx - ay), \text{ and hence} \\ (ax - by)(bx - ay).$$

$$2. x^4 - (a + b)x^3 + (a^2b + ab^2)x - a^2b^2.$$

Here, taking the first and last terms together, and the two middle terms together, we have

$$(x^2 + ab)(x^2 - ab) - (a + b)x^3 + ab(a + b)x \\ = (x^2 + ab)(x^2 - ab) - (a + b)x\{x^2 - ab\} \\ = (x^2 - ab)\{x^2 + ab - (a + b)x\} = (x^2 - ab)(x - a)(x - b).$$

$$3. x^{3m} - 4x^m + 3. \text{ This equals}$$

$$x^{3m} - x^m - 3(x^m - 1) = x^m(x^{2m} - 1) - 3(x^m - 1) \\ = x^m(x^m + 1)(x^m - 1) - 3(x^m - 1) \\ = (x^m - 1)\{x^m(x^m + 1) - 3\}.$$

## Exercise xxxv.

- |                                     |   |
|-------------------------------------|---|
| 1. $a^2 - ab + ax - bx.$            | 14. $a^3 + a^2 - 2.$                              |
| 2. $abx^2 + b^2xy - a^2xy - aby^2.$ | 15. $3a^2b^4 - 2ab^2 \quad 1.$                    |
| 3. $x^4 + ax^3 - a^2x - a^4.$       | 16. $y^3 - 3y + 2.$                               |
| 4. $a^3x + 2a^2x^2 + 2ax^3 + x^4.$  | 17. $2a^3 - a^2b - ab^2 + 2b^3.$                  |
| 5. $acx^2 + (ad - bc)x - bd.$       | 18. $b^{3m} + b^{2m} - 2.$                        |
| 6. $25x^4 - 5x^3 + x - 1.$          | 19. $y^{3n} - 2y^{2n}z^n - 2y^n z^{2n} + z^{3n}.$ |
| 7. $a^2 - b^2 + ax - ac - bx + bc.$ | 20. $a^3 - 4ab^2 + 3b^3.$                         |
| 8. $a^3 + (1 + a)ab + b^2.$         | 21. $a^{2m} - 3a^m c^n + 2c^{2n}.$                |
| 9. $x^4 + 2xy(x^2 - y^2) - y^4.$    | 22. $ax^3 - (a^2 + b)x^2 + b^2.$                  |
| 10. $x^3 - y^3 + x^2 + xy + y^2.$   | 23. $35x^{2n} - 6a^2x^n - 9a^4.$                  |
| 11. $2b + (b^2 - 4)x - 2bx^2.$      | 24. $a^2b^2 + 2abc^2 - a^2c^2 - b^2c^2.$          |
| 12. $x^3 + 3x^2 - 4.$               | 25. $am^2 - ab^2 + b^2m - m^3.$                   |
| 13. $p^3 - p^2q - 2pq^2 + 2q^3.$    | 26. $\frac{1}{3} - 6a^2 + 27a^4.$                 |

- 27.  $(x-y)^3 + (1-x+y)(x-y)z - z^2$ .
- 28.  $24m^3 - 28m^2n + 6mn^2 - 7n^3$ .
- 29.  $x^{m+n} + x^ny^n + x^my^m + y^{m+n}$ .
- 30.  $x^4 + 2x^3y - a^2x^2 + x^2y^2 - 2axy^2 - y^4$ .

SECTION IV.—APPLICATION OF THE THEORY OF DIVISORS.

Art. XXV. By Theorem I. we prove that

- $x^n - a^n$  is divisible by  $x - a$  always
- $x^n - a^n$  " " "  $x + a$  when  $n$  is even
- $x^n + a^n$  " " "  $x + a$  when  $n$  is odd.

By actual division we find, in the above cases;—

$$\frac{x^n - a^n}{x - a} = x^{n-1} + x^{n-2}a + \dots + xa^{n-2} + a^{n-1} \dots \dots \dots (1).$$

$$\frac{x^n - a^n}{x + a} = x^{n-1} - x^{n-2}a + \dots + xa^{n-3} - a^{n-1} \dots \dots \dots (2).$$

$$\frac{x^n + a^n}{x + a} = x^{n-1} - x^{n-2}a + \dots - xa^{n-2} + a^{n-1} \dots \dots \dots (3).$$

EXAMPLES.

1. Resolve into factors  $x^3 - y^3$ ; here  $x - y$  is one factor and by (1) the other is  $x^2 + xy + y^2$ .

2. Resolve  $a^3 + (b - c)^3$ ; here  $a + (b - c)$  is one factor; and by (3) the other is  $a^2 - a(b - c) + (b - c)^2$ .

3. Resolve  $x^{15} + 1024y^{10}$ . This  $= (x^3)^5 + \{(2y)^2\}^5$ , one factor of which is  $x^3 + (2y)^2$ , and by (3) the other factor is

$$(x^3)^4 - (x^3)^3(4y^2) + (x^3)^2(4y^2)^2 - x^3(4y^2)^3 + (4y^2)^4 \\ = x^{12} - 4x^9y^2 + 16x^6y^4 - 64x^3y^6 + 256y^8.$$

4. Resolve  $(x - 2y)^3 + (2x - y)^3$  into factors.

Here by (3) we have

$$\frac{(x - 2y)^3 + (2x - y)^3}{x - 2y + 2x - y} = (x - 2y)^2 - (x - 2y)(2x - y) + (2x - y)^2$$

∴ the factors are

$$3(x - y)(7x^2 - 13xy + 7y^2),$$

5. Resolve  $x^5 + x^4y + x^3y^2 + x^2y^3 + xy^4 + y^5$  :

$$\text{By (1) we see that this} = \frac{x^6 - y^6}{x - y} = \frac{(x^3 + y^3)(x^3 - y^3)}{x - y}$$

$$= (x + y)(x^2 - xy + y^2)(x^2 + xy + y^2).$$

6. Resolve  $x^{11} - x^{10}a + x^9a^2 - x^8a^3 + x^7a^4 - x^6a^5 + x^5a^6$   
 $- x^4a^7 + x^3a^8 - x^2a^9 + xa^{10} - a^{11}$ . This =  $\frac{x^{12} - a^{12}}{x + a}$

$$= \frac{(x^6 + a^6)(x^6 - a^6)}{x + a} = \frac{(x^6 + a^6)(x^3 - a^3)(x^3 + a^3)}{x + a}$$

$$= (x^2 + a^2)(x^4 - x^2a^2 + a^4)(x - a)(x^2 + xa + a^2)(x^2 - xa + a^2).$$

### Exercise xxxvi.

Factor the following :—

1.  $x^6 - y^6$ ,  $x^3 - 1$ ,  $x^3 + 8$ ,  $8a^3 - 27x^3$ ,  $8 + a^3x^3$ .
2.  $x^5 - a^{10}$ ,  $27a^3 - 64$ ,  $a^{12} - b^8$ ,  $x^{10} - 32y^5$ .
3. Find a factor which, multiplied into  $a^4 + a^3b + a^2b^2 - 3 + b^4$ , will give  $a^5 - b^5$ .
4. By what factor must  $x^4x^2y + 16xy^2 - 64y^3$  be multiplied to give  $x^4 - 256y^4$ ?
5. Factor  $x^7 + x^6y + x^5y^2 + x^4y^3 + x^3y^4 + x^2y^5 + xy^6 + y^7$ .  
Find the factors of the following :
  6.  $(3y^2 - 2x^2)^3 - (3x^2 - 2y^2)^3$ ,  $a^8 - 16b^4$ .
  7.  $x^3 - y^3 - x(x^2 - y^2) + y(x - y)$ .
  8.  $b(x^3 - a^3) + ax(x^2 - a^2) + a^3(x - a)$ .
  9.  $b(m^3 + a^3) + am(m^2 - a^2) + a^3(m + a)$ .
  10.  $x^6 - y^6 + 2xy(x^4 + x^2y^2 + y^4)$ .
  11.  $(a^2 - bc)^3 + 8b^3c^3$ ,  $x^{4m} - a^{4n}$ .
  12.  $x^3 - 3ax^2 + 3a^2x - a^3 + b^3$ .
  13.  $x^3 + 8y^3 + 4xy(x^2 - 2xy + 4y^2)$ .
  14.  $8x^3 - 6xy(2x + 3y) + 27y^3$ .
  15.  $1 - 2x + 4x^2 - 8x^3$ .
  16.  $a^5 + a^4bc + a^3b^2c^2 + a^2b^3c^3 + ab^4c^4 + b^5c^5$ .

Art. XXVI. The principles illustrated in Section II., chap. II., may be applied to factor various algebraic expressions, as in the following cases :

## EXAMPLES.

1. Find the factors of

$$(a+b+c)(ab+bc+ca) - (a+b)(b+c)(c+a).$$

1st. Observe that the expression is *symmetrical* with respect to  $a, b, c$ .

2nd. If there be any *monomial* factor  $a$  must be one. Putting  $a=0$ , the expression vanishes.  $\therefore a$  is a factor, and, by symmetry,  $b$  and  $c$  are also factors.  $\therefore abc$  is a factor.

3rd. There can be no other *literal* factor, because the given expression is of only *three* dimensions, and  $abc$  is of three dimensions.

4th. But there may be a *numerical* factor,  $m$  suppose, so that we have

$$(a+b+c)(ab+bc+ca) - (a+b)(b+c)(c+a) = mabc.$$

To find  $m$ , put  $a=b=c=1$  in this equation, and  $m=1$ .

$\therefore$  the expression  $= abc$ .

2. Resolve  $a^2(b-c) + b^2(c-a) + c^2(a-b)$ .

1st. For  $a=0$  this does not vanish.  $\therefore a$  is not a factor, and by symmetry neither is  $b$  nor  $c$ .

2nd. Try a *binomial* factor; this will likely be of the form  $b-c$ ; put  $b-c=0$ , i.e.,  $b=c$  in the given expression, and there results

$$a^2(c-c) + c^2(c-a) + c^2(a-c), \text{ which } = 0,$$



$\therefore b-c$  is a factor, and by symmetry  $c-a$  and  $a-b$  are factors. Since the given expression is only of *three* dimensions, there can be no other *literal* factor; but there may be a *numerical* factor,  $m$  (say), so that

$$a^2(b-c) + b^2(c-a) + c^2(a-b) = m(a-b)(b-c)(c-a).$$





To find the value of  $m$ , give  $a, b, c$ , in this equation, any values which will not reduce either side to zero; let  $a=1, b=2, c=0$ ,







**IMAGE EVALUATION  
TEST TARGET (MT-3)**



**Photographic  
Sciences  
Corporation**

23 WEST MAIN STREET  
WEBSTER, N.Y. 14580  
(716) 872-4503

15  
16  
18  
20  
22  
25  
28  
32  
36  
40  
45  
50  
56  
63

11  
12  
15  
18  
22  
27  
32  
36  
40  
45  
50  
56  
63

and we have  $2 = m(-2)$ , or  $m = -1$ ; so that the given expression =  $-(a-b)(b-c)(c-a)$ , or  $(a-b)(b-c)(a-c)$ .

3. Resolve  $a^3(b+c^2) + b^3(c+a^2) + c^3(a+b^2) + abc(abc+1)$ .

Here we see at once that there is no *monomial* factor: put  $b+c^2=0$ , *i.e.*,  $b = -c^2$ , and the expression becomes  $a^3(-c^2+c^2) - c^6(c+a^2) + c^3(a+c^4) - c^3a(-c^3a+1)$  which = 0;  $\therefore b+c^2$  is a factor, and by symmetry  $c+a^2$  and  $a+b^2$  are also factors; and proceeding as in former examples we find  $m=1$ ;  $\therefore$  the expression =  $(b+c^2)(c+a^2)(a+b^2)$ .

4. Resolve into factors the expression

$$(a-b)^3 + (b-c)^3 + (c-a)^3.$$

As before, we find that there are no *monomial* factors.

Let  $a-b=0$ , or  $a=b$ , and substituting  $b$  for  $a$  the expression becomes zero; hence

$a-b$  is a factor.

By symmetry  $b-c$  “

and  $c-a$  “

Hence the factors are

$$m(a-b)(b-c)(c-a).$$

To find  $m$  let  $a=0$ ,  $b=1$ ,  $c=2$ , and we have

$$6 = 2m, \text{ or } m=3.$$

The factors are, therefore,

$$3(a-b)(b-c)(c-a).$$

5. Resolve into factors

$$a^3(b-c) + b^3(c-a) + c^3(a-b).$$

As before, we find that there are no *monomial* factors.

Let  $a-b=0$ , or  $a=b$ ; substituting  $b$  for  $a$ , the expression becomes zero;

therefore  $a-b$  is a factor.

By symmetry  $b-c$  “

and  $c-a$  “

Now the product of these three factors is of *three* dimensions, while the expression itself is of *four* dimensions. There must, therefore, be another factor of *one* dimension. It cannot be a

monomial factor, for the expression has no such factors. It cannot be a binomial factor, such as  $a+b$ , for then, by symmetry,  $b+c$  and  $c+a$  would also be factors, which would give an expression of *six* dimensions. It cannot be a trinomial factor, unless  $a$ ,  $b$ , and  $c$  are similarly involved. For instance, if  $a-b+c$  were a factor, then, by symmetry,  $b-c+a$  and  $c-a+b$  would also be factors, and the dimensions would be *six* instead of *four*. The other factor must, therefore, be  $a+b+c$ . Hence,

$$a^3(b-c) + b^3(c-a) + c^3(a-b) = m(a-b)(b-c)(c-a)(a+b+c).$$

To find  $m$ , put  $a=0$ ,  $b=1$ , and  $c=2$ , and we have

$$-6 = 5m;$$

$$\therefore m = -1.$$

Hence the factors are

$$-(a-b)(b-c)(c-a)(a+b+c),$$

or,  $(a-b)(a-c)(b-c)(a+b+c).$

6. Prove that

$$a^3 + b^3 + c^3 + 3(a+b)(b+c)(c+a)$$

is exactly divisible by  $a+b+c$ , and find all the factors.

Let  $a+b+c=0$ , or  $a=-(b+c)$ ; substituting this value for  $a$ , we have

$$-(b+c)^3 + b^3 + c^3 + 3bc(b+c), \text{ or}$$

$$-(b+c)^3 + (b+c)^3 \text{ which } = 0, \text{ and}$$

therefore  $a+b+c$  is a factor.

As before, we find that there are no monomial factors. Since  $a+b+c$ , the factor already obtained, is of *one* dimension, the other factor must be of *two* dimensions, and cannot, therefore, be a binomial; for if  $a+b$  were a factor, by symmetry  $b+c$ , and  $c+a$  must also be factors. The factors in that case would give a quantity of *four* dimensions, while the expression itself is only of three dimensions. Nor can  $a^2+b^2+c^2$  be a factor. For if so, the other factor must involve a numerical multiple of the first power of  $a$ , and, therefore, on taking the first power of  $a$  out of terms involving first and third powers, we should have left some numerical multiple of  $a^2+b^2+c^2$ , instead of which we get

$a^2 + 3(b+c)^2$ . Nor can  $a^2 + (b+c)^2$  be a factor, for symmetry would require two other factors, viz.:  $b^2 + (c+a)^2$ , and  $c^2 + (a+b)^2$ , thus giving a quantity of *seven* dimensions.

The only factor admissible is, therefore,  $(a+b+c)^2$ .

Hence

$$a^3 + b^3 + c^3 + 3(a+b)(b+c)(c+a) = m(a+b+c)(a+b+c)^2 \\ = m(a+b+c)^3.$$

To find  $m$ , let  $a=1$ ,  $b=0$ , and  $c=0$ , and we have  $1=m$ .

Hence the factors are

$$(a+b+c)(a+b+c)(a+b+c).$$

7. Simplify

$$a(b+c)^2 + b(a+c)^2 + c(a+b)^2 - (a+b)(a-c)(b-c) \\ - (a-b)(a-c)(b+c) + (a-b)(b-c)(a+c).$$

Let  $a=0$ , and the expression becomes  $bc^2 + cb^2 + bc(b-c) - bc(b+c) - bc(b-c)$ , which equals zero; therefore  $a$  is a factor; by symmetry  $b$  and  $c$  are also factors.

The expression is of three dimensions, and  $abc$  is of three dimensions, there cannot therefore be any other literal factor.

Hence the expression  $= abc$ .

To find  $m$ , let  $a=b=c=1$ , and we have

$$4+4+4=m;$$

$$m=12.$$

$\therefore$  the expression  $= 12abc$ .

In the preceding examples the factors have been *linear*, but the principle applies equally well to those of higher dimensions. (See Th. ii. Cor.)

8. Examine whether  $x^n + 1$  is a factor of  $x^{3n} + 2x^{2n} + 3x^n + 2$ .

Let  $x^n + 1 = 0$ , or  $x^n = -1$ , and substituting, the expression vanishes, therefore,  $x^n + 1$  is a factor.

9. Examine whether  $a^2 + b^2$  is a factor of

$$2a^4 + a^3b + 2a^2b^2 + ab^3.$$

Let  $a^2 + b^2 = 0$ , or  $a^2 = -b^2$ , substituting, we have

$$2b^4 - ab^3 - 2b^4 + ab^3 \text{ which } = 0, \text{ and}$$

therefore  $a^2 + b^2$  is a factor.

10. Prove that  $a^3 + b^3$  is a factor of

$$a^5 + a^4b + a^3b^2 + a^2b^3 + ab^4 + b^5.$$

Let  $a^3 + b^3 = 0$ , or  $a^3 = -b^3$ ; substituting, we have

$$-a^2b^3 - ab^4 - b^5 + a^2b^3 + ab^4 + b^5, \text{ which } = 0, \text{ and}$$

therefore  $a^3 + b^3$  is a factor.

### Exercise xxxvii.

Resolve into factors

1.  $(x+y+z)^3 - (x^3 + y^3 + z^3)$ .
2.  $bc(b-c) - ca(a-c) - ab(b-a)$ .
3.  $(a^2 - b^2)^3 + (b^2 - c^2)^3 + (c^2 - a^2)^3$ .
4.  $x(y+z)^2 + y(z+x)^2 + z(x+y)^2 - 4xyz$ .
5.  $(a+b)^3 - (b+c)^3 + (c-a)^3$ .
6.  $a(b-c)^3 + b(c-a)^3 + c(a-b)^3$ .
7.  $(a+b+c)(ab+bc+ca) - abc$ .
8.  $a^3(c-b^2) + b^3(a-c^2) + c^3(b-a^2) + abc(abc-1)$ .
9.  $a^2(b+c) + b^2(c+a) + c^2(a+b) + 2abc$ .
10.  $(a-b)(c-h)(c-k) + (b-c)(a-h)(a-k) + (c-a)(b-h)(b-k)$ .
11.  $x^4y^2 + x^2y^4 + x^4z^2 + x^2z^4 + y^4z^2 + y^2z^4 + 2x^2y^2z^2$ .
12.  $(a-b)^5 + (b-c)^5 + (c-a)^5$ .
13.  $ab(a+b) + bc(b+c) + ca(c+a) + (a^3 + b^3 + c^3)$ .
14.  $a^4(c-b^3) + b^4(a-c^3) + c^4(b-a^3) + abc(a^2b^2c^2 - 1)$ .
15.  $x^4(y^2 - z^2) + y^4(z^2 - x^2) + z^4(x^2 - y^2)$ .
16.  $x^4 + y^4 + z^4 - 2x^2y^2 - 2y^2z^2 - 2z^2x^2$ .
17.  $(b-c)(x-b)(x-c) + (c-a)(x-c)(x-a) + (a-b)(x-a)(x-b)$ .
18.  $(a+b)^3 + (b+c)^3 + (c+a)^3 + 3(a+2b+c)(b+2c+a)(c+2a+b)$ .
19. Shew that  $a^5 + a^2b^2 - ab^2 - b^3$  has  $a^2 - b$  for a factor.
20. Shew that  $(x+y)^7 - x^7 - y^7 = 7xy(x+y)(x^2 + xy + y^2)^2$ .
21. Examine whether  $x^2 - 5x + 6$  is a factor of  $x^3 - 9x^2 + 26x - 24$ .

22. Shew that  $a - b + c$  is a factor of  
 $a^2(b+c) - b^2(c+a) + c^2(a+b) + abc.$
23. Shew that  $a^2 + 3b$  is a factor of  
 $a^4 - 4a^3b^3 + 3a^2b^4 + 3a^2b - 12ab^4 + 9b^5,$   
 and find the other factor .
24. Find the factors of  $a^4(b-c) + b^4(c-a) + c^4(a-b).$

SECTION V.—FACTORING A POLYNOME BY TRIAL DIVISORS.

Art. XXVII. To find, if possible, a rational linear factor of the polynome.

$$x^n + bx^{n-1} + cx^{n-2} + \dots + hx + k.$$

Substitute successively for  $x$  every measure (both positive and negative) of the term  $k$ , till one is found, say  $m$ , that makes the polynome vanish, then  $x - m$  will be a factor of the polynome.

EXAMPLES.

1. Factor  $x^3 + 9x^2 + 16x + 4.$

The measures of 4 are  $\pm 1$ ,  $\pm 2$  and  $\pm 4$ . Since every coefficient of the given polynome is positive, the positive measures of 4 need not be tried. Using the others, it will be found that  $-2$  makes the polynome vanish; thus

$$-2 \left| \begin{array}{cccc} 1 & 9 & 16 & 4 \\ & -2 & -14 & -4 \\ \hline 1 & 7 & 2 & 0 \end{array} \right.$$

Hence the factors are  $(x+2)(x^2+7x+2).$

The labour of substitution may often be lessened by arranging the polynome in ascending powers of  $x$ , and using  $1 \div$  (measure of  $k$ ) instead of the measures of  $k$ . (This is really substituting  $1 \div$  measure of  $k$ , for  $1 \div x$ ). Should a fraction occur during the course of the work, further trial of that measure of  $k$  will be needless.



EXAMPLES.

2. Factor  $x^3 - 10x^2 - 63x + 60$ .

The measures of 60 are  $\pm 1, \pm 2, \pm 3, \pm 4, \pm 5, \&c.$  Neither  $+1$  nor  $-1$  will make the polynome vanish. Try 2; thus

$$\begin{array}{r|rrrr} 1 & 60 & -63 & -10 & 1 \\ & & 30 & & \\ \hline 2 & 30 & -16\frac{1}{2} & & \end{array}$$

A fraction occurring we need go no further.  $-2$  will also give a fraction, as may easily be seen. Next try 3; thus

$$\begin{array}{r|rrrr} 1 & 60 & -63 & -10 & 1 \\ & & 20 & & \\ \hline 3 & 20 & -14\frac{1}{3} & & \end{array}$$

A fraction again occurring, we may stop.  $-3$  will also give a fraction. Next try 4; thus

$$\begin{array}{r|rrrr} 1 & 60 & -63 & -10 & 1 \\ & & 15 & -12 & \\ \hline 4 & 15 & -12 & -5\frac{1}{4} & \end{array}$$

Next try  $-4$ .

$$\begin{array}{r|rrrr} -1 & 60 & -63 & -10 & 1 \\ & & -15 & & \\ \hline 4 & 15 & -19\frac{1}{4} & & \end{array}$$

Next trying 5 we find it fails, then try  $-5$ , thus

$$\begin{array}{r|rrrrr} -1 & 60 & -63 & -10 & 1 \\ & & -12 & 15 & -1 \\ \hline 5 & 12 & -15 & 1 & 0 \end{array}$$

The remainder vanishes as required; the factors are, therefore,  $(x+5)(x^2 - 15x + 12)$ .

**Art. XXVIII.** When  $k$  has a large number of factors, the number that need actually be tried can often be considerably lessened by the following means.

Add together all the coefficients of  $x$  (including the constant term  $k$ ); let the sum be called  $k_1$ .

From the sum of the coefficients of the even powers of  $x$  (including  $k$ ) take the sum of the coefficients of the odd powers of  $x$ ; let the remainder be called  $k_2$ . (In the coefficients are included the signs of the terms).

1st. If  $k_1$  vanish,  $x-1$  will be a factor of the polynome.

2nd. If  $k_2$  vanish,  $x+1$  will be a factor of the polynome.

3rd. If both  $k_1$  and  $k_2$  vanish,  $x^2-1$  will be a factor of the polynome.

4th. If neither  $k_1$  nor  $k_2$  vanish, (writing  $p$  for "a positive measure of  $k$  greater than 1");

( $\alpha$ ) We need not try the substitution of  $p$  for  $x$  unless  $p-1$  be a measure of  $k_1$ , and  $p+1$  a measure of  $k_2$ .

( $\beta$ ) Nor need we try the substitution of  $-p$  for  $x$  unless  $p+1$  be a measure of  $k_1$ , and  $p-1$  a measure of  $k_2$ .

(In trying for measures, the signs of  $k$ ,  $k_1$ , and  $k_2$  may be neglected.

#### EXAMPLES.

1. Find the factors of  $x^3-10x^2-63x+60$ . (See Ex. 2 above).

$$\text{Here } k=60; k_1=1-10-63+60=-12,$$

$$k_2=-1-10+63+60=112.$$

Tabulating the trial-measures we get

12	1,	2,	3,	4,			
60	2,	3,	4,	5,	6,	10,	12,
112		4,			7,		

12	3,	4,		6,			
60	2,	3,	4,	5,	6,	10,	
112	1,	2,		4,			

(It is evident that 12 is the highest measure of 60 we need try in the upper table, for the next measure, 15, would give 14 as a trial-measure of 12, and higher measures of 60 would give higher trial-measures. Similarly, 10 is the highest measure that need be tried in the lower table.)

In the upper table, 3 is the only measure of 60 that gives a full column; hence of the positive measures of 60 we need try only the substitution of 3 for  $x$ .

In the lower table, 2, 3, and 5 give full columns, hence we must try the substitutions  $-2$ ,  $-3$ ,  $-5$  for  $x$ .

On trying the four substitutions to which we are thus restricted we find  $-5$  is the only one for which the polynome vanishes. (See Ex. 2 above).

2. Find the factors of  $x^4 + 12x^3 - 40x^2 + 67x - 120$ .

$$k = -120; k_1 = 1 + 12 - 40 + 67 - 120 = -80;$$

$$k_2 = 1 - 12 - 40 - 67 - 120 = -238.$$

80	1,	2,	4,	5,					
120	2,	3,	4,	5,	6,	8,	10,	12,	15, &c.
238				7,					

80		4,	5,					16,			
120	2,	3,	4,	5,	6,	8,	10,	15,	20,	24, &c.	
238	1,	2,					7,	14,	21,		

The upper table gives us 6 as a trial-measure, and the lower gives us  $-3$  and  $-15$ .

Trying these we get

1	-120	67	-40	12	1
		-20			
6	-20	$7\frac{5}{6}$			

-1	-120	67	-40	12	1
		40			
3	-40	$35\frac{2}{3}$			

-1	-120	67	-40	12	1
		8	-5	3	-1
15	-8	5	-3	1:	0

Hence  $x+15$  and  $x^3-3x^2+5x-8$  are the factors. The latter cannot be resolved, for our tables above tell us we need try only  $x-6$ ,  $x+3$ , and  $x+15$ . The first two have been found not to be factors, and 15 will not measure 8.

4. Factor  $x^4-27x^2+14x+120$ .

$$k=120; k_1=1-27+14+120=108$$

$$k_2=1-27-14+120=80.$$

108	1,	2,	3,	4,	9					
120	2,	3,	4,	5,	6,	8,	10,	12,	15,	&c.
80	4,	5,							16,	

108	3,	4,	6,	9						
120	2,	3,	4,	5,	6,	8,	10,	12,	15,	&c.
80	1,	2,	4,	5.						

The upper table gives us 3 and 4, the lower table gives us -2, -3, and -5. Using these in order we get

1	120	14	-27	0	1	
3	40	18	-3	-1	-1	
1	40	18	-3	-1	0	Hence $x-3$ is a factor.
4	10	7	1	1		
4	10	7	1	1	0	Hence $x-4$ is a factor.
-1	-5	-1				
2	5	1	0			Hence $x+2$ is a factor.

and there remains  $x+5$ , a factor.

Hence the factors are  $(x-3)(x-4)(x+2)(x+5)$ .

5. Factor  $x^4-px^3+(q-1)x^2+px-q$ .

$$k=-q; k_1=1-p+(q-1)+p-q=0;$$

$$k_2=1+p+(q-1)-p-q=0.$$

Since both  $k_1$  and  $k_2$  vanish, the polynome is divisible by both  $x-1$  and  $x+1$ .

1	1	-p	q-1	p	-q
1	1	-p+1	q-p	q-p	q
-1	1	-p+1	q-p	q	0
-1	-1	+p	-q		
1	1	-p	q	0	

Hence the other factor is  $x^2 - px + q$ .

6. Factor  $x^4 + 2ax^3 + (a^2 + a)x^2 + 2a^2x + a^3$ .

$$k = a^3; k_1 = 1 + 2a + (a^2 + a) + 2a^2 + a^3 = (a + 1)^3;$$

$$k_2 = 1 - 2a + (a^2 + a) - 2a^2 + a^3 = a^3 - a^2 - a - 1.$$

The positive measures of  $k$  are 1,  $a$ ,  $a^2$ ,  $a^3$ . Of these 1 may be rejected at once, since neither  $k_1$  nor  $k_2$  vanish, and  $a^2$  and  $a^3$  may also be rejected since  $k_1$  or  $(a + 1)^3$  is not divisible by either  $a^2 \pm 1$  or  $a^3 \pm 1$ . But  $k_1$  is divisible by  $a + 1$ , and  $k_2$  is divisible by  $a - 1$ ; thus we need only try the substitution of  $-a$  for  $x$ . (See 4  $\beta$ , page 92)

$$\begin{array}{r|rrrrr}
 -a & 1 & 2a & a^2 + a & 2a^2 & a^3 \\
 & & -a & -a^2 & -a^2 & -a^3 \\
 \hline
 -a & 1 & a & a & a^2 & 0 \\
 & & -a & 0 & -a^2 & \\
 \hline
 & 1 & 0 & a & 0 & 
 \end{array}$$

Hence the factors are  $(x + a)^2(x^2 + a)$ .

7. Factor  $x^3 - (a + c)x^2 + (b + ac)x - bc$ .

$$k = -bc;$$

$$k_1 = 1 - (a + c) + (b + ac) - bc = 1 - a + b - c + ac - bc$$

$$k_2 = -1 - (a + c) + (b + ac) - bc = -(1 + a + b + c + ac + bc).$$

The factors of  $k_1$ , other than 1, are  $b$  and  $c$ .  $k_1$  is not divisible by either  $b \pm 1$  nor by  $c + 1$ . However,  $k_1$  is divisible by  $c - 1$ , and  $k_2$  is at the same time divisible by  $c + 1$ ,  $\therefore$  we need only try the substitution of  $c$  for  $x$ . (See 4  $\alpha$ , page 86).

$$\begin{array}{r|rrrr}
 c & 1 & -(a + c) & (b + ac) & -bc \\
 & & c & -ac & bc \\
 \hline
 & 1 & -a & b & 
 \end{array}$$

Hence the factors are  $(x - c)(x^2 - ax + b)$ .

Exercise xxxviii.

- |                                     |  |
|-------------------------------------|--|
| 1. $a^3 - 9a^2 + 16a - 4.$          | 15. $x^3 - 11x^2 + 39x - 45.$          |
| 2. $x^3 - 9x^2 + 26x - 24.$         | 16. $x^3 + 5x^2 + 7x + 2.$             |
| 3. $x^3 - 7x^2 + 16x - 12.$         | 17. $a^3 - 3a^2 - 193a + 195.$         |
| 4. $x^3 - 12x + 16.$                | 18. $p^3 - 3p^2 - 6p - 8.$             |
| 5. $x^3 + 3x^2 + 5x + 3.$           | 19. $a^4 + 5a^3 - 3a^2 - 7a + 6.$      |
| 6. $x^4 + 4x^3 + 10x^2 + 12x + 9.$  | 20. $a^{6n} - 6a^{4n} + 11a^{2n} - 6.$ |
| 7. $x^3 - 3x + 2.$                  | 21. $a^4 - 41a^2b^2 + 16b^4.$          |
| 8. $x^4 + 2x^2 + 9.$                | 22. $a^4 - a^2b^2 - 2ab^3 + 2b^4.$     |
| 9. $m^3 - 3m^2n + 4mn^2 - 2n^3$     | 23. $p^3 - 4p^2 + 6p - 4.$             |
| 10. $x^3 + 2x^2 + 2.$               | 24. $x^n + 4x^{2n} - 5.$               |
| 11. $m^3 - 5m^2n + 8mn^2 - 4n^3.$   | 25. $y^4 - 5y^3 + 8y^2 - 8.$           |
| 12. $b^3 + b^2c + 7bc^2 + 39c^3.$   | 26. $a^4 - 2a^3 + 3a^2 - 2a + 1.$      |
| 13. $m^4 - 4mn^3 + 3n^4.$           | 27. $a^3 + a^2b^2 + ab^2 - 3b^3.$      |
| 14. $a^4 - 7a^2b + 28ab^3 - 16b^4.$ | 28. $2a^{3n} - a^{2n} - a^n + 2.$      |
29.  $x^4 - 18x^3 + 113x^2 - 288x + 252.$   
 30.  $x^4 - 9x^3y + 29x^2y^2 - 39xy^3 + 18y^4.$

Art XXIX. To find, if possible, a rational linear factor of the polynome

$$ax^n + bx^{n-1} + cx^{n-2} + \dots + hx + k.$$

*First Method.* Multiply the polynome by  $a^{n-1}$ .

$$(ax)^n + b(ax)^{n-1} + ac(ax)^{n-2} + \dots + a^{n-2}h(ax) + a^{n-1}k;$$

or writing  $y$  for  $ax$ ,

$$y^n + by^{n-1} + acy^{n-2} + \dots + a^{n-2}hy + a^{n-1}k.$$

Factor this polynome by the method of the last article, replace  $y$  by  $ax$ , and divide the result by  $a^{n-1}$ .

EXAMPLE.

Factor  $3x^4 + 5x^3 - 33x^2 + 43x - 20.$

Multiply by  $3^3$  and express in terms of  $3x$ .

$$(3x)^4 + 5(3x)^3 - 99(3x)^2 + 387(3x) - 540;$$

or,  $y^4 + 5y^3 - 99y^2 + 387y - 540.$

Here  $k = -540$ ;  $k_1 = 1 + 5 - 99 + 387 - 540 = -246$ ;  
 $k_2 = 1 - 5 - 99 - 387 - 540 = -1030$ .

246	1, 2, 3, 6, 41, 82, 123, 246.
540	2, 3, 4,
1030	5,

246	3, 6, 41, &c.	(Trying by factors of 246 instead of by factors of 540, for convenience).
540	2, 5,	
1030	1,	

The only factors of 540 in full columns are 4 in the upper table and 2 in the lower one; hence we need try only the substitutions 4 and -2.

1	- 540	387	- 99	5	1
2	- 135	- 135	63	- 9	- 1
4	- 135	63	- 9	- 1;	0

Hence  $y - 4$  is a factor. The substitution  $-2$  need not now be tried, since we see that 135 is not a multiple of 2. The other factor is therefore  $y^3 + 9y^2 - 63y + 135$ .

Replacing  $y$  by  $3x$  and dividing by 27;

$$\begin{aligned} & \frac{1}{27}(3x-4)(27x^3+81x^2-189x+135) \\ & = (3x-4)(x^3+3x^2-7x+5), \end{aligned}$$

which are the factors.

**Art. XXX. Second Method.** Writing  $m$  for "a measure of  $a$ ," and  $p$  for a "measure of  $k$ , positive or negative;"

For  $x$  substitute every value of  $p \div m$  till one, say  $p' \div m'$  be found which makes the polynome vanish; then  $m'x - p'$  will be a factor. Should a fraction be met with in the course of substitution, further trial of that value of  $p \div m$  will be useless.

Should  $k$  have more factors than  $a$ , it will generally be better to arrange the polynome in ascending powers of  $x$  and use values of  $m \div p$  instead of  $p \div m$ , making  $p$  positive and  $m$  positive or negative.

To reduce the number of trial-measures, calculate  $k_1$  and  $k_2$ , as directed on page 92, then 1, 2, 3 hold as on that page, but in 4 read  $p-m$  for  $p-1$  and  $i+m$  for  $p+1$ .

## EXAMPLES.

1. Factor  $36x^3 + 171x^2 - 22x + 480$ .

$$k = 480, k_1 = 36 + 171 - 22 + 480 = 665$$

$$k^2 = -36 + 171 + 22 + 480 = 637.$$

$m$  may have any of the values  $\pm 1, \pm 2, \pm 3, \pm 4, \pm 6, \pm 9, \pm 12, \pm 18, \pm 36$ .

In forming the table write out the measures of  $k_1$ ; take each measure in succession and add to it each value of  $m$  separately, should the sum measure 480, *i.e.*,  $k$ , add to it the same value of  $m$ , and should the new sum measure 637, *i.e.*,  $k_2$ , keep the measure of 480, writing above it the value of  $m$  used. Should the sum in either case not be a measure, another value of  $m$  must be tried; when all the values of  $m$  have been tried, another measure of 665, *i.e.*,  $k_1$  must be tried till all have been tested. (Measures of  $k_1$  or 665 have been used in this instance because they are much fewer than those of 480; measures of  $k_2$  or 637 would have done equally well).

$m = +3,$	$+1,$	$+3$	$-2$	$-3$	$-9$	$-3$	
665	1,	5,	7	5	7	19	19
480	4,	6,	10	3	4	10	13
637	7,	7,	13	1	1	1	13

Hence the only substitutions that need be tried are

$$\frac{3}{4}, \frac{1}{6}, \frac{3}{10}, \frac{-2}{3}, \frac{-3}{4}, \frac{-9}{10}, \frac{-3}{16}, \text{ for } \frac{1}{x}.$$

Arrangement in ascending powers of  $x$ .

By actual trial, as below, we find  $\frac{-3}{16}$  is the only one of these giving a zero remainder.



	480	- 22	171	36
3		360		
4	120	84½		
1		80		
6	80	9⅔		
3		144		
10	48	12·2		
-2		-320	223	-266
3	160	-114	133;	-230
-3		360		
4	120	- 95½		
-9		-432		
10	48	-45·4		
-3		- 90	21	-36
16	30	- 7	12;	0

(The coefficients are written only once, and understood for the other lines of substitution.)

Hence the factors are  $3x+16$  and  $12x^2 - 7x+30$ .

The latter factor cannot be resolved, for 16 will not measure 30, and all the other factors left for trial by the tables above, have been tried and have failed.

2. Factor  $10x^4 - x^3(15y+4z) - x^2(40y^2 - 6yz) \cdot$

$x(60y^3 + 16y^2z) - 24y^3z.$

Here  $m = \pm 1, \pm 2, \pm 5, \text{ or } \pm 10.$   $k = -24y^3z.$

$$\begin{aligned} k_1 &= 10 - (15y+4z) - (40y^2 - 6yz) + (60y^3 + 16y^2z) - 24y^3z \\ &= 10 - 15y - 40y^2 + 6yz - 2z(2 - 3y - 8y^2 + 12y^3) \\ &= (5 - 2z)(2 - 3y - 8y^2 + 12y^3). \end{aligned}$$

$k_2 = (5 + 2z)(2 + 3y - 8y^2 - 12y^3),$  as may easily be found by making the calculation.

We get at a glance  $2z$  a factor of  $k,$   $2z-5$  a factor of  $k_1,$  and  $2z+5$  a factor of  $k_2;$  hence taking  $m=5,$  we are directed to try the substitution  $\frac{2z}{5}$  for  $x.$

	10	$-(15y+4z)$	$-(40y^2 - 6yz)$	$(60y^3 + 16y^2z)$	$-24y^3z$
$2z$		$4z$	$-6yz$	$-16y^2z$	$24y^3z$
$5$	$2$	$-3y$	$-8y^2$	$12y^3;$	$0$

and  $k_2,$  as  
out in 4

$-6, \pm 9,$

like each  
separately,  
value of  
the mea-  
sured the  
must be  
measure  
measures  
they are  
uld have

3  
9  
6  
13

of these

Hence  $5x - 2z$  is a factor, the other being

$$2x^3 - 3x^2y - 8xy^2 + 12y^3.$$

The latter factor being homogeneous, the method of this article may be applied to it.

$$m = \pm 1 \text{ or } \pm 2, \quad k = 12, \quad k_1 = 3, \quad k_2 = 15.$$

$m = 1,$	2,	1,	-1		
3	1,	1,	3,	3	The other columns
12	2,	3,	4,	2	are not full.
15	3,	5,	5,	1	

Hence the trial-substitutions (arrangement in ascending powers of  $x$ ) are  $\frac{1}{2}, \frac{2}{3}, \frac{1}{4}, \frac{-1}{2}$ .

1	12	-8	-3	2	
2	6	-1	-2;	0	
2	4	2			
3	2	1;	0		Final factor is $2y + x$ .

Hence the factors are  $(x - 2y)(2x - 3y)(x + 2y)$ , and these, with the factor  $5x - 2z$  already found, give the complete resolution of the polynome proposed.

(The factor  $5x - 2z$ , might easily have been got by the method of Art. XXIII., page 79, but the present solution shows we are independent of that article. It may also be obtained by rearranging the polynome in terms of  $y$ ).

#### Exercise xxxix.

Factor

1.  $2x^3 - 20x^2 + 38x - 20$ ;  $x^3 - 7x^2y + 16xy^2 - 12y^3$ .
2.  $12x^3 + 5x^2y + xy^2 + 3y^3$ ;  $8x^3 - 14x + 6$ .
3.  $3x^3 - 15ax + a^2x - 5a^3$ ;  $2x^3 + 9x^2y + 7xy^2 - 3y^3$ .
4.  $2b^4 - 7b^3c - 4b^2c^2 + bc^3 - 4c^4$ ;  $15a^3 + 47a^2b + 13ab^2 - 12b^3$ .
5.  $4p^4 + 8p^3q + 7p^2q^2 + 8pq^3 + 3q^4$ .
6.  $150x^4 - 725x^3y + 931x^2y^2 + 920xy^3 - 1152y^4$ .
7.  $36x^4 - 6(9 - 7y)x^3 - 7(9 + 14y)x^2y + 3(49 - 40y)xy^2 + 180y^3$ .
8.  $10x^4 - x^3(15y + 4z) + x^2(40y^2 + 6yz) + x(60y^3 - 16y^2z) - 24y^3z$

CHAPTER IV.

SECTION I.—DIVISION. MEASURES AND MULTIPLES.

Art. XXXI. When one quantity is to be divided by another the quotient can often be readily obtained by resolving the divisor or dividend, or both, into factors.

EXAMPLES.

1. Divide  $a^2 - 2ab + b^2 - c^2 + 2cd - d^2$  by  $a - b + c - d$ . Here we see at once that the dividend  $= (a - b)^2 - (c - d)^2$ , and  $\therefore$  quotient  $= a - b - (c - d) = a - b - c + d$ .

2. Divide the product of  $a^2 + ax + x^2$  and  $a^3 + x^3$  by  $a^4 + a^2x^2 + x^4$ . Here  $a^3 + x^3 = (a + x)(a^2 - ax + x^2)$ , and the divisor  $= (a^2 + ax + x^2)(a^2 - ax + x^2) \therefore$  the quotient is  $a + x$ .

3. Divide  $a^3 + a^2b + a^2c - abc - b^2c - bc^2$  by  $a^2 - bc$ . The dividend is  $a(a^2 - bc) + b(a^2 - bc) + c(a^2 - bc) \therefore$  the quotient  $= a + b + c$ .

4.  $(a^3 + b^3 - c^3 + 3abc) \div (a + b - c)$ .

Dividend  $= a^3 + b^3 + 3ab(a + b) - c^3 - 3ab(a + b) + 3c(a + b - c) = (a + b)^3 - c^3 - 3ab(a + b - c)$  which is exactly divisible by  $a + b - c$ ; quotient  $= a^2 + b^2 + c^2 - ab + bc + ca$ .

5. Divide  $x^5 - x^4y + x^3y^2 - x^2y^3 + xy^4 - y^5$  by  $x^3 - y^3$ .

The dividend is (Art. XXV.) evidently  $(x^6 - y^6) \div (x + y)$ , and this divided by  $x^3 - y^3 = (x^3 + y^3) \div (x + y) = x^2 - xy + y^2$ .

6. Divide  $b(x^3 + a^3) + ax(x^2 - a^2) + a^3(x + a)$  by  $(a + b)(x + a)$ . Striking the factor  $x + a$  out of dividend and divisor we have  $b(x^2 - ax + a^2) + ax(x - a) + a^3 = b(x^2 - ax + a^2) + a(x^2 - ax + a^2) = (a + b)(x^2 - ax + a^2) \therefore$  quotient  $= x^2 - ax + a^2$ .

7. Divide  $apx^4 + x^3(aq + bp) + x^2(ar + bq + pc) + x(qc + br) + cr$  by  $ax^2 + bx + c$ .

Factoring the dividend (Art. XXIII.) we have

$$(ax^2 + bx + c)(px^2 + qx + r).$$

$\therefore$  the quotient = the latter factor.

8. Divide  $6x^4 - 13ax^3 + 13a^2x^2 - 13a^3x - 5a^4$  by  $2x^3 - 3ax - a^2$ .

This can be done by Art. XVII. The divisor is  $2x^3 - a^2 - 3ax$ , and we see at once that  $3x^2 + 5a^2$  must be two terms of the quotient.

Multiplying diagonally into the first two terms of the divisor, and adding the products, we get  $+7a^2x^2$ ; but  $+13a^2x^2$  is required.  $\therefore +6a^2x^2$  is still required, and as this must come from the third term multiplied into  $-3ax$ , that third term must be  $-2ax$ ;  $\therefore$  the quotient is  $3x^2 + 5a^2 - 2ax$ .

NOTE.—By multiplying the terms  $-2ax$ ,  $-3ax$ , diagonally into the  $x^2$ 's and  $a^2$ 's respectively, we get the remaining terms of the dividend; it is, of course, necessary to test whether the division is exact.

9. Divide  $2a^4 - a^3b - 12a^2b^2 - 5ab^3 + 4b^4$  by  $a^2 - b^2 - 2ab$ .

Here, as before, one factor is  $a^2 - b^2 - 2ab$ ;  $\therefore$  two terms of the other factor are  $2a^2 - 4b^2$ . Multiplying, as in the last example, we get  $-6a^2b^2$ ; but  $-12a^2b^2$  is required.  $\therefore -6a^2b^2$  is still needed, and  $+3ab$  is the third term of the required quotient, which is therefore  $2a^2 - 4b^2 + 3ab$ .

Prove that

$$10. (1 + x + x^2 + \dots + x^{n-1})(1 - x + x^2 - \dots + x^{n-1}) \\ = 1 + x^2 + x^4 + \dots + x^{2n-2}.$$

$$\text{Product} = \frac{1 - x^n}{1 - x} \cdot \frac{1 + x^n}{1 + x} \\ = \frac{1 - x^{2n}}{1 - x^2} = 1 + x^2 + x^4 + \dots + x^{2n-2}.$$

$$11. \text{ Divide } (a^2 - bc)^3 + 3b^3c^3 \text{ by } a^2 + bc. \\ = (a^2 - bc)^3 + (2bc)^3 \text{ by } (a^2 - bc) + 2bc \\ = (a^2 - bc)^2 - (a^2 - bc) \times 2bc + (2bc)^2 \\ = a^4 - 4a^2bc + 7b^2c^2.$$

12.

Find

1.

2.

3.

4.

5.

6.

7.

8.

9.

10.

11.

12.

13.

14.

15.

16.

17.

18.

19.

20.

21.

22.

12. Divide  $1+2357947691x^9$  by  $1-11x+121x^2$

$$\text{Dividend} = 1 + (11x)^9$$

$$= \{1 - (11x)^3 + (11x)^6\} \{1 + (11x)^3\}$$

$$\text{Divisor} = \{1 + (11x)^3\} - (1+11x).$$

$$\therefore \text{quotient} = \{1 - (11x)^3 + (11x)^6\} (1+11x).$$

### Exercise xl.

Find the quotients in the following cases :

1.  $1 - x + x^2 - x^3 \div 1 - x.$
2.  $1 - 2x^4 + x^8 \div x^4 + 2x^2 + 1.$
3.  $x^{16} + a^8x^8 + a^{16} \div x^4 - a^2x^2 + a^4.$
4.  $x^4 + 4x^2y^2 - 32y^4 \div x - 2y.$
5.  $1 - 4x^2 + 12x^3 - 9x^4 \div 1 + 2x - 3x^2.$
6.  $(a^2 - 2ax + x^2)(a^3 + 3a^2x + 3ax^2 + x^3) \div a^3 - x^3.$
7.  $x^3 - y^3 + z^3 + 3xyz \div x - y + z.$
8.  $6a^4 - a^3b + 2a^2b^2 + 13ab^3 + 4b^4 \div 2a^2 - 3ab + 4b^2.$
9.  $4x^4 - x^2y^2 + 6xy^3 - 9y^4 \div 2x^2 + 3y^2 - xy.$
10.  $a^4 + b^4 - c^4 - 2a^2b^2 \div a^2 - b^2 - c^2.$
11.  $21a^4 - 16a^3b + 16a^2b^2 - 5ab^3 + 2b^4 \div 3a^2 - ab + b^2.$
12.  $2a^3 - 7a^2 - 46a - 21 \div 2a^2 + 7a + 3.$
13.  $\{a^3(b-c) + b^3(c-a) + c^3(a-b)\} \div a + b + c.$
14.  $x^3 - 3ax^2 + 3a^2x - a^3 + b^3 \div x - a + b.$
15.  $x^4 - y^4 + z^4 + 2x^2z^2 - 2y^2 - 1 \div x^2 - y^2 + z^2 - 1.$
16.  $x^4 - (a+c)x^2 + (b+ac)x^2 - bcx \div x - c.$
17.  $x^3 + x^2y + xy^2 + y^3 \div x + y.$
18.  $x^7 - x^6y + x^5y^2 - x^4y^3 + x^3y^4 - x^2y^5 + xy^6 - y^7 \div x^4 + y^4.$
19.  $a^4 + b^4 - c^4 - 2a^2b^2 - 2c^2 - 1 \div a^2 - b^2 - c^2 - 1.$
20.  $a^4 - ab^3 - ac^3 - 2a^3b + 2b^4 + 2bc^3 - 3a^3c - 3b^3c - 3c^4 \div a + 3c - 2b.$
21.  $a^2b - bx^2 + a^2x - x^3 \div (x+b)(a-x).$
22.  $a(b-c)^3 + b(c-a)^3 + c(a-b)^3 \div a^2 - ab - ac + bc.$

$$23. a^2b^2 + 2abc^2 - a^2c^2 - b^2c^2 \div ab + ac - bc.$$

$$24. x^3 + y^3 + 3xy - 1 \div x + y - 1.$$

$$25. x^6 - x^2 - 2 \div x^2 - x + 1.$$

$$26. a^4 - 29a^2 - 50a - 21 \div a^2 - 5a - 7.$$

$$27. (2x - y)^2 a^4 - (x + y)^2 a^2 x^2 + 2(x + y) a x^4 - x^6 + \\ (2x - y) a^2 + (x + y) a x - x^3.$$

$$28. (x^3 - 1) a^3 - (x^3 + x^2 - 2) a^2 + (4x^2 + 3x + 2) a - 3(x + 1) \\ \div (x - 1) a^2 - (x - 1) a + 3.$$

**Art. XXXII.** The HIGHEST COMMON FACTOR of two algebraic quantities may, in general, be readily found by factoring. The H. C. F. is often discovered by taking the sum or difference (or sum *and* difference) of the given expressions, or of some multiples of them.

#### EXAMPLES.

1. Find the H. C. F. of  $(b - c)x^2 + (2ab - 2ac)x + a^2b - a^2c$ , and  $(ab - ac + b^2 - bc)x + a^2c + ab^2 - a^2b - abc$ .

Taking out the common factor  $b - c$  we get  $(b - c)(x^2 + 2ax + ab)$  and  $(b - c)\frac{1}{2}\{(a - b)x - a^2 + ab\}$ ;

$\therefore b - c$  is the H. C. F. of the given expressions.

2. Find the H. C. F. of

$$1 - x + y + z - xy + yz - zx - xyz, \text{ and}$$

$$1 - x - y - z + xy + yz + zx - xyz.$$

Their difference is  $2y + 2z - 2xy - 2zx = 2(1 - x)(y + z)$ .

Their sum is  $2 - 2x + 2yz - 2xyz = 2(1 - x)(1 + yz)$ .

$\therefore$  the H. C. F. is  $(1 - x)$ .

3. Find the H. C. F. of  $x^5 + 3x^4 - 8x^2 - 9x - 3$ , and

$$x^5 - 2x^4 - 6x^3 + 4x^2 + 13x + 6.$$

The annexed method of finding the H. C. F. depends on the principle, that if a quantity measures two other quantities, it will measure any multiple of their sum or difference.

The  
are th  
multi  
(d) ha  
of th  
with  
do no  
to fin  
same  
contai

4. ]

If (a  
8 and

MEASURES AND MULTIPLES.

<u>1 + 3</u>	<u>0 - 3 - 9 - 3</u>	<i>(a)</i>	
<u>1 - 2 - 6 + 4 + 13 + 6</u>		<i>(b)</i>	
<hr style="border-top: 1px solid black;"/>			
<u>5 + 6 - 12 - 22 - 9</u>		<i>(c)</i>	[ = <i>(a)</i> - <i>(b)</i> ]
<hr style="border-top: 1px solid black;"/>			
<u>2 + 6</u>	<u>0 - 16 - 18 - 6</u>		<i>(a)</i> × 2
<u>1 - 2 - 6 + 4 + 13 + 6</u>			<i>(b)</i>
<hr style="border-top: 1px solid black;"/>			
<u>3 + 4 - 6 - 12 - 5</u>		<i>(d)</i>	
<hr style="border-top: 1px solid black;"/>			
<u>15 + 18 - 36 - 66 - 27</u>			<i>(c)</i> × 3
<u>15 + 20 - 30 - 60 - 25</u>			<i>(d)</i> × 5
<hr style="border-top: 1px solid black;"/>			
<u>- 2 - 6 - 6 - 2</u>			
<hr style="border-top: 1px solid black;"/>			
<u>1 + 3 + 3 + 1</u>		<i>(f)</i>	
<hr style="border-top: 1px solid black;"/>			
<u>25 + 30 - 60 - 110 - 45</u>			<i>(c)</i> × 5
<u>27 + 36 - 54 - 108 - 45</u>			<i>(d)</i> × 9
<hr style="border-top: 1px solid black;"/>			
<u>- 2 - 6 - 6 - 2</u>			
<hr style="border-top: 1px solid black;"/>			
<u>1 + 3 + 3 + 1</u>		<i>(g)</i>	

H. C. F. =  $(x+1)^3$ .

The coefficients are written in two lines, *(a)* and *(b)*. They are then subtracted so as to cancel the first terms. *(a)* is next multiplied by 2, and added to cancel the last terms. If *(c)* and *(d)* had been the same their terms would have been the coefficients of the H. C. F. Since they are not, we proceed with them as with *(a)* and *(b)* till they become the same. When *(a)* and *(b)* do not contain the same number of terms it is more convenient to find only *(c)*, and then use this with the quantity containing the same number of terms. The general rule is to operate on lines containing the same, or nearly the same number of terms.

4. Find the H. C. F. of  $3x^3 + 2x^2 - 14x + 8$ , and  $6x^3 - 11x^2 + 13x - 12$ .

<u>3 + 2 - 14 + 8</u>	<i>(a)</i>	
<u>6 - 11 + 13 - 12</u>	<i>(b)</i>	
<hr style="border-top: 1px solid black;"/>		
<u>6 + 4 - 28 + 16</u>		<i>(c)</i> × 2
<u>15 - 41 + 28</u>	<i>(c)</i>	<i>(b)</i> - <i>(a)</i> ,
<u>(5-7)(3-4)</u>		

H. C. F. =  $3x - 4$ . *(d)*

If *(a)* and *(b)* have a common factor its first term must measure 3 and 6, and its last term must measure 8 and 12. *(c)* is not

therefore, the H. C. F. Resolve (c) into factors.  $5x-7$  is not a factor of (a) and (b). If, therefore, (a) and (b) have a common factor it is  $3x-4$ . On trial  $3x-4$  is found to be a factor of (a) and  $\therefore$  it is the H. C. F. of (a) and (b).

5. If  $x^2+px+q$ , and  $x^2+rx+s$  have a common factor, prove that this factor is

$x + \frac{q-s}{p-r}$ . If  $x-a$  be the common factor then the remainders on dividing the given expressions by  $x-a$ , must be zero, i. e.,  
 $a^2+pa+q=0$ , and  $a^2+ra+s=0$ , or

$$(p-r)a=s-q, \therefore a = \frac{s-q}{p-r}, \text{ and}$$

$$x-a = x - \frac{s-q}{p-r} = x + \frac{q-s}{p-r}.$$

6. What value of  $a$  will make  $a^2x^2+(a+2)x+1$ , and  $a^2x^2+a^2-5$ , have a common measure.

They cannot have a monomial factor. Neither can they have one of two dimensions unless  $(a+2)$  vanishes, i. e., unless  $a=-2$ , in which case the expressions become  $4x^2+1$ , and  $4x^2-1$ , which have no C. F. Hence if the given quantities have a C. F., it must be of the form  $x+m$ ; dividing  $a^2x^2+a^2-5$  by  $x+m$ , we have for remainder,

$a^2m^2+a^2-5=0$ , or  $m^2 = \frac{5-a^2}{a^2}$ ;  $\therefore m = \frac{1}{a} \sqrt{5-a^2}$ , in which  $\sqrt{5-a^2}$  must be possible and integral,  $\therefore a^2=4$ , ( $a^2=1$  gives values to  $m$  which on trial fail) and  $a = \pm 2$ , of which the positive value must be taken, and  $\therefore 2x+1$  is the C. F.

7. If the H. C. F. of  $a$  and  $b$  be  $c$ , the L. C. M. of  $(a+b)(a^3-b^3)$ , and  $(a-b)(a^3+b^3)$  is  $\frac{a^6-b^6}{c^2}$ .

Let  $a=mc$ ,  $b=nc$ , and  $\therefore a^3=m^3c^3$ ,  $b^3=n^3c^3$ . Thus  
 $(a+b) = c(m+n)$ ;  $(a-b) = c(m-n)$ , and  
 $(a^3+b^3) = c^3(m^3+n^3)$ ;  $(a^3-b^3) = c^3(m^3-n^3)$ .  
 $\therefore (a+b)(a^3-b^3) = c^4(m+n)(m^3-n^3)$ , and  
 $(a-b)(a^3+b^3) = c^4(m-n)(m^3+n^3)$ .

Th  
 L. C.  
 8. I  
 and r.  
 Let  
 $x^3+qa$   
 equatin

$\therefore$   
 Or  
 Divid

and as t  
 ing coeff

Find

1.  $2x^4$
2.  $x^3 +$
3.  $px^3 -$
4.  $ax^3 -$
5.  $1-3$
6.  $ac^{2a} +$



The H. C. F. of the last expressions is  $c^4(m^2 - n^2)$ ,  $\therefore$  the  
 L. C. M. =  $c^4(m^6 - n^6) = \frac{c^6(m^6 - n^6)}{c^2} = \frac{a^6 - b^6}{c^2}$ .

8. If  $(x-a)^2$  measures  $x^3 + qx + r$ , find the relation between  $q$  and  $r$ .

Let  $x+m$  be the other factor, then  
 $x^3 + qx + r = (x-a)^2(x+m) = x^3 + (m-2a)x^2 + (a^2 - 2am)x + ma^2$   
 equating coefficients,

$$m - 2a = 0, \quad a^2 - 2am = q, \quad ma^2 = r$$

$$\therefore m = 2a, \text{ and } \therefore a^2 - 4a^2 = q, \quad 2a^3 = r, \text{ and}$$

$$a^2 = -\frac{q}{3}, \text{ or } a^6 = -\frac{q^3}{27}; \text{ and } a^3 = \frac{r}{2} \text{ or } a^6 = \frac{r^2}{4}$$

$$\therefore \frac{r^2}{4} = -\frac{q^3}{27}, \text{ or } \frac{r^2}{4} + \frac{q^3}{27} = 0.$$

Or thus:—

Dividing  $x^3 + qx + r$  by  $(x-a)^2$  we find the remainder  
 $(q + 3a^2)x + r - 2a^3$   
 and as this will be the same for all values of  $x$ , we have, by equat-  
 ing coefficients,

$$q + 3a^2 = 0,$$

$$\text{and } r - 2a^3 = 0,$$

$$\text{or } q^3 = -27a^6$$

$$\text{and } r^2 = 4a^6;$$

$$\text{therefore } \frac{r^2}{4} + \frac{q^3}{27} = 0, \text{ as before.}$$

### Exercise xli.

Find the H. C. F. of the following :

1.  $2x^4 + 3x^3 + 5x^2 + 9x - 3, 3x^4 - 2x^3 + 10x^2 - 6x + 3.$
2.  $x^3 + (a+1)x^2 + (a+1)x + a, x^3 + (a-1)x^2 - (a-1)x + a.$
3.  $px^3 - (p+q)x^2 + (p-q)x + q, px^3 - (p+q)x^2 + (p+q)x + q.$
4.  $ax^3 - (a-b)x^2 - (b-c)x - c, 2ax^3 + (a+2b)x^2 + (b+2c)x + c.$
5.  $1 - 3\frac{2}{3}x - 3\frac{1}{3}x^2 + \frac{1}{3}x^3 - x^4, 1 - 1\frac{1}{6}x - 3x^2 + 1\frac{1}{6}x^3 + x^4.$
6.  $ac^{2a} + bc^{2b} + (a+b)c^{a+b}, a^c c^a + a^c c^b + a^a b^c + b^c c^b.$

7.  $a^2x^3 + a^5 - 2abx^3 + b^2x^3 + a^3b^2 - 2a^4b$ , and  
 $2a^2x^4 - 5a^4x^2 + 3a^6 - 2b^2x^4 + 5a^2b^2x^2 - 3a^4b^2$ .
8.  $(ax+by)^2 - (a-b)(x+z)(ax+by) + (a-b)^2xz$ , and  
 $(ax-by)^2 - (a+b)(x+z)(ax-by) + (a+b)^2xz$ .
9.  $a(b^2 - c^2) + b(c^2 - a^2) + c(a^2 - b^2)$  and  
 $a(b^3 - c^3) + b(c^3 - a^3) + c(a^3 - b^3)$ .
10.  $a^{3m} + a^{2m} + a^m + 1$ , and  $a^{3m} - a^{2m} + a^m - 1$ .
11. If  $x^3 + ax^2 + bx + c$ , and  $x^2 + a'x + b'$ , have a common factor of one dimension in  $x$ , it must be one the factors of  
 $(a-a')x^2 + (b-b')x + c$ .
12. Determine the H. C. F. of  $(a-b)^5 + (b-c)^5 + (c-a)^5$ , and  
 $(a^2 - b^2)^5 + (b^2 - c^2)^5 + (c^2 - a^2)^5$ .
13. Find the H. C. F. of  
 $2(y^3 - 2y^2 - y + 2)x^3 + 3(y^2 - 1)x^3 - (2y^3 - y^2 - 2y + 1)$ , and  
 $3(y^3 - 4y^2 + 5y - 2)x^3 + 7(y^2 - 2y + 1)x - (3y^3 - 5y^2 + y + 1)$ .
14. If  $x^2 + px + q$ , and  $x^2 + mx + n$  have a common linear factor, shew that  
 $(n-q)^2 + n(m-p)^2 = m(m-p)(n-q)$ .
15. Find the L. C. M. of  $x^3 - 3x^2 + 3x - 1$ ,  $x^3 - x^2 - x + 1$ ,  
 $x^4 - 2x^3 + 2x - 1$ , and  $x^4 - 2x^3 + 2x^2 - 2x + 1$ .
16. Find the L. C. M. of  
 $x^3 + 6x^2 + 11x + 6$ ,  $x^3 + 7x^2 + 14x + 8$ .  
 $x^3 + 8x^2 + 19x + 12$ , and  $x^3 + 9x^2 + 26x + 24$ .
17. Find the value of  $y$  which will make  
 $2(y^2 + y)x^2 + (11y - 2)x + 4$  and  
 $2(y^3 + y^2)x^3 + (11y^2 - 2y)x^2 + (y^2 + 5y)x + 5y - 1$ , have a common measure.
18. The product of the H. C. F. and L. C. M. of two quantities is equal to half the sum of their squares, one of them is  
 $2x^3 - 11x^2 + 17x - 6$ ; find the other.
19. If  $x+a$  and  $x-a$  are both measures of  $x^3 + px^2 + qx + r$ , shew that  $pq = r$ .

20.  
(x-a)21.  
L. C.22.  
of the23.  
F., sh  
 $x^4 +$ 24.  
mon fa25.  
and  $a^5$ 26.  
y the H

the L.

27.  
L. C. M28. S  
(above) v

29. F

Art.  
lowest te  
of factor  
nominato

20. If  $x^3 + qx^2 + r$  and  $x^3 + mx + n$  have a common measure  $(x-a)^2$ , show that  $q^2n^3 = m^2r^3$ .

21. If the H. C. F. of  $x^3 + px + q$  and  $x^2 + mx + n$ , be  $x+a$ , their L. C. M. is

$$x^4 + (m-a)x^3 + px^2 + (a^3 + mp)x + a(m-a)(a^2 + p).$$

22. If  $x^2 + qx + 1$ , and  $x^3 + px^2 + qx + 1$ , have a common factor of the form  $x+a$ , shew that  $(p-1)^2 - q(p-1) + 1 = 0$ .

23. If  $x^3 + px^2 + q$ , and  $x^2 + mx + n$ , have  $x+a$  for their H. C. F., shew that their L. C. M. is

$$x^4 + (m-a+px)x^3 + p(m-a)x^2 + a^2(a-p)x + a^2(a-p)(m-a).$$

24. If  $x^2 + px + 1$ , and  $x^3 + px^2 + qx + 1$ , have  $x-a$  for a common factor, shew that  $a = \frac{1}{1-q}$ .

25. Find the H. C. F. of  $(a^2 - b^2)^3 + (b^2 - c^2)^3 + (c^2 - a^2)^3$ , and  $a^5(b-c) + b^5(c-a) + c^5(a-b)$ .

26. If  $\alpha$  be the H. C. F. of  $b$  and  $c$ ,  $\beta$  the H. C. F. of  $c$  and  $a$ ,  $\gamma$  the H. C. F. of  $a$  and  $b$ , and  $\delta$  the H. C. F. of  $a$ ,  $b$ , and  $c$ , then

the L. C. M. of  $a$ ,  $b$ ,  $c$ , is  $\frac{abcd}{\alpha\beta\gamma}$ .

27. If  $x+c$  be the H. C. F. of  $x^2 + ax + b$ , and  $x^2 + a'x + b'$ , their L. C. M. will be  $x^3 + (a+a'-c)x^2 + (aa' - c^2)x + (a-c)(a'-c)c$ .

28. Shew that the L. C. M. of the quantities in Ex. 2 (solved above) will be a complete square if  $x = y^2 + z^2 - y^2z^2$ .

29. Find the H. C. F. of  $x^8 + 2x^6 + 3x^4 - 2x^2 + 1$ , and  $6x^8 + x^7 + 17x^5 - 7x^3 - 9$ .

---

## SECTION II.—FRACTIONS.

---

Art. XXXIII. When required to reduce a fraction to its lowest terms, we can often apply some of the preceding methods of factoring to discover the H. C. F. of the numerator and denominator.

## EXAMPLES.

$$1. \frac{ac+by+ay+bc}{aj+2bx+2ax+bf} = \frac{c(a+b)+y(a+b)}{j(a+b)+2x(a+b)} = \frac{c+y}{j+2x}.$$

$$2. \frac{a^4-ba^3-a^2b^2+ab^3}{a^5-ba^4-ab^4+b^5} = \frac{a\{a^3+b^3-ab(a+b)\}}{a(a^4-b^4)-b(a^4-b^4)}$$

$$= \frac{a(a+b)(a-b)^2}{(a-b)(a^4-b^4)} = \frac{a}{a^2+b^2}.$$

$$3. \frac{x^5+x^4y+x^3y^2+x^2y^3+xy^4+y^5}{x^5-x^4y+x^3y^2-x^2y^3+xy^4-y^5}.$$

Here the numerator is evidently  $(x^6-y^6) \div (x-y)$ , and the denominator is  $\frac{x^6-y^6}{x+y}$ . The result is  $\therefore \frac{x+y}{x-y}$ .

$$4. \frac{(x+y)^5-x^5-y^5}{(x+y)^4+x^4+y^4} = \frac{5x^4y+10x^3y^2+10x^2y^3+5xy^4}{(x+y)^4-x^2y^2+(x^2+y^2)^2-x^2y^2}$$

$$= \frac{5xy\{x^3+y^3+2xy(x+y)\}}{(x^2+y^2+xy)\{(x+y)^2+xy+x^2+y^2-xy\}}$$

$$= \frac{5xy(x+y)(x^2+xy+y^2)}{2(x^2+xy+y^2)^2} = \frac{5xy(x+y)}{2(x^2+xy+y^2)}$$

$$5. \frac{x^2-12x+35}{x^3-10x^2+31x-30}.$$

Here we see at once that the numerator  $= (x-5)(x-7)$ ; and it is plain that  $x-7$  is not a factor of the denominator; we  $\therefore$  try  $x-5$  (Horner's division), and find the quotient to be  $x^2-5x+6$ .

$$\therefore \text{the result} = \frac{x-7}{x^2-5x+6}.$$

$$6. \frac{x^4+2x^2+9}{x^4-x^3+8x-21}.$$

The factors of the numerator are at once seen to be  $x^2+2x+3$ , and  $x^2-2x+3$ , of which the latter is one factor of the denominator, the other being (Horner's division)  $x^2-2x-7$ :  $\therefore$  the result

$$\text{is } \frac{x^2+2x+3}{x^2-2x-7}.$$

## Exercise xlii

Reduce the following to their lowest terms:

$$1. \frac{x^2 - 7x + 6}{x^3 - 2x^2 - 8x - 96}, \quad \frac{3xy^2 - 19xy + 14x}{7y^3 - 17y^2 + 6y}$$

$$2. \frac{x^4 + a^2x^2 + a^4}{x^4 + ax^3 - a^3x - a^4}, \quad \frac{x^2 + x - 12}{x^3 - 5x^2 + 7x - 3}$$

$$3. \frac{x^3 - 3x + 2}{x^3 + 4x^2 - 5}, \quad \frac{x^4 + 2x^2 + 9}{x^4 - 4x^3 + 4x^2 - 9}$$

$$4. \frac{2 + bx}{2b + (b^2 - 4)x - 2bx^2}, \quad \frac{x^3 + 2x^2 + 12x}{x^3 + 4x}$$

$$5. \frac{5a^5 + 10a^4x + 5a^3x^2}{a^3x + 2a^2x^2 + 2ax^3 + x^4}, \quad \frac{20x^4 + x^2 - 1}{25x^4 + 5x^3 - x - 1}$$

$$6. \frac{x^7 - x^6y + x^5y^2 - x^4y^3 + x^3y^4 - x^2y^5 + xy^6 - y^7}{x^7 + x^5y + x^5y^2 + x^4y^3 + x^3y^4 + x^2y^5 + xy^6 + y^7}$$

$$7. \frac{3a^2x^4 - 2ax^2 - 1}{4a^3x^6 - 2a^2x^4 - 3ax^2 + 1}, \quad \frac{x^2 + \left(\frac{a}{b} + \frac{b}{a}\right)xy + y^2}{x^2 + \left(\frac{a}{b} - \frac{b}{a}\right)xy - y^2}$$

$$8. \frac{a^2(b-c) + b^2(c-a) + c^2(a-b)}{abc(a-b)(b-c)(c-a)}$$

$$9. \frac{(a+b+c)^2}{a^3(b-c) + b^3(c-a) + c^3(a-b)}$$

10. From Ex. 4 (solved above) show that

$$\frac{(a-b)^4 + (b-c)^4 + (c-a)^4}{(a-b)^5 + (b-c)^5 + (c-a)^5} = \frac{(a-b)^2 + (b-c)^2 + (c-a)^2}{5(a-b)(b-c)(c-a)}$$

$$11. \frac{(x+y)^5 - x^5 - y^5}{(x+y)^7 - x^7 - y^7}$$

12. Shew that

$$\frac{(a-b)^7 + (b-c)^7 + (c-a)^7}{(a-b)^5 + (b-c)^5 + (c-a)^5} = \frac{7}{14} \{(a-b)^2 + (b-c)^2 + (c-a)^2\}$$

Art. XXXIV. In reducing complex fractions it is often convenient to multiply both terms of the complex fraction by the L. C. M. of all the *denominators* involved.

## EXAMPLES.

1. Simplify  $\frac{\frac{1}{2}(x+1\frac{1}{2})-\frac{2}{3}(1-\frac{3}{4}x)}{1\frac{1}{2}-\frac{1}{2}(x+4\frac{1}{4})}$ .

Here the L. C. M. of all the denominators involved is 12;  $\therefore$  multiplying both terms of the complex fraction by 12, and removing brackets, we have

$$\frac{6x+8-8+6x}{21-4x-17} = \frac{12x}{4-4x} = \frac{3x}{1-x}.$$

2.  $a - \frac{a-b}{1+ab}$   
 $1 + \frac{a(a-b)}{1+ab}$

Here multiplying both terms by  $1+ab$ , we get

$$\frac{a(1+ab)-a+b}{1+ab+a(a-b)} = \frac{b(a^2+1)}{a^2+1} = b.$$

3.  $\frac{1}{x-1 + \frac{1}{1 + \frac{x}{4-x}}}$

Here multiplying both terms of the fraction which follows  $x-1$  by  $4-x$ , the given fraction becomes at

once  $\frac{1}{x-1 + \frac{4-x}{4}}$ , and now multiplying both terms by 4, we

have  $\frac{4}{4x-4+4-x} = \frac{4}{3x}$ .

It may be observed that when the fraction is reduced to the form  $\frac{a}{b} \div \frac{c}{d}$ , we may strike out any factor common to the two *denominators*, and also any factor common to the two *numerators*; it is sometimes more convenient to do this than to multiply directly by the L. C. M. of all the denominators.

4. Simplify  $\left(\frac{a+b}{a-b} + \frac{a-b}{a+b}\right) \div \left(\frac{a^2+b^2}{a^2-b^2} - \frac{a^2-b^2}{a^2+b^2}\right)$ .

Here the numerator of the first fraction is  $(a+b)^2 + (a-b)^2$  and the denominator is  $a^2 - b^2$ ; the numerator of second fraction is  $(a^2+b^2)^2 - (a^2-b^2)^2$ , and the denominator is  $a^4 - b^4$ ; the former denominator cancels this to  $a^2 + b^2$ , which, of course, becomes a multiplier of the first numerator:

$\therefore$  we have  $\frac{(a^2+b^2)\{(a+b)^2+(a-b)^2\}}{(a^2+b^2)^2-(a^2-b^2)^2} = \frac{(a^2+b^2)^2}{2a^2b^2}$ .

Occasionally, we at once discover a common complex factor, strike this out, and simplify the result.

5.  $\frac{1}{a} + \frac{1}{b} + \frac{1}{c}$   
 $\frac{1}{a^2} + \frac{1}{b^2} - \frac{1}{c^2} + \frac{2}{ab}$ : here the den. =  $\left(\frac{1}{a} + \frac{1}{b}\right)^2 - \frac{1}{c^2}$   
 $= \left(\frac{1}{a} + \frac{1}{b} + \frac{1}{c}\right) \left(\frac{1}{a} + \frac{1}{b} - \frac{1}{c}\right)$ , and cancelling the common factor we have

$\frac{1}{\frac{1}{a} + \frac{1}{b} - \frac{1}{c}}$ , and multiplying by  $abc$ , this =  $\frac{abc}{bc+ca-ab}$ .

Exercise xliii.

Simplify the following:

1.  $\frac{1 - \frac{1}{2}\{1 - \frac{1}{3}(1-x)\}}{1 - \frac{1}{3}\{1 - \frac{1}{2}(1-x)\}}$ ,  $\frac{\frac{a+b}{a-b} + \frac{a-b}{a+b}}{\frac{a+b}{a-b} - \frac{a-b}{a+b}}$ .

2.  $\frac{\frac{x}{x+y} + \frac{x}{x-y}}{\frac{2x}{x^2-y^2}}$ ,  $\frac{\frac{1}{1-a} - \frac{1}{1+a}}{\frac{a}{1-a} + \frac{1}{1+a}}$ .

3.  $1 + \frac{a}{1+a + \frac{2a^2}{1+a}}$ ,  $x - \frac{x-y}{(x-a)(x-z)} \cdot \frac{x-y}{x+y}$

$$4. \frac{\frac{a^2+b^2}{2a^2} - \frac{2b^2}{a^2+b^2}}{\frac{a^2+b^2}{2b^2} - \frac{2a^2}{a^2+b^2}}; \quad \frac{\frac{1}{a} + \frac{1}{ab^3}}{b-1 + \frac{1}{b}}$$

$$5. \frac{\frac{a+b}{c+d} + \frac{a-b}{c-d}}{\frac{a+b}{c-d} + \frac{a-b}{c+d}}; \quad \frac{a+b + \frac{b^3}{a}}{a+b + \frac{1}{b}}$$

$$6. \frac{3xyz}{yz-zx-xy} - \frac{\frac{x-1}{x} + \frac{y-1}{y} + \frac{z-1}{z}}{\frac{1}{x} + \frac{1}{y} + \frac{1}{z}}$$

$$7. \frac{\frac{2}{a^2} + \frac{2}{b^2} + \frac{2}{c^2} + \frac{a^4+b^4+c^4}{a^2b^2c^2}}{\frac{a}{bc} + \frac{b}{ac} + \frac{c}{ab}}$$

$$8. \frac{a^3+a^2b+ab^2+b^3}{a^3-a^2b+ab^2-b^3} \div \frac{a^2+2ab+b^2}{a^2-b^2}$$

$$9. \left( \frac{a+b}{a-b} + \frac{a^2+b^2}{a^2-b^2} \right) \div \left( \frac{a-b}{a+b} - \frac{a^3-b^3}{a^3+b^3} \right)$$

$$10. \frac{\frac{1}{a} + \frac{1}{b+c}}{\frac{1}{a} - \frac{1}{b+c}} \left\{ 1 + \frac{b^2+c^2-a^2}{2bc} \right\}$$

$$11. \frac{\frac{2(1-x)}{1+x} + \frac{(1-x)^2}{(1+x)^2} + 1}{\frac{2(1+x)}{1-x} + \frac{(1+x)^2}{(1-x)^2} + 1}; \quad \frac{\left(\frac{x-a}{x+a}\right)^2 + \left(\frac{x+a}{x-a}\right)^2 - 2}{\left(\frac{x-a}{x+a}\right)^2 + 2 + \left(\frac{x+a}{x-a}\right)^2}$$

$$12. \frac{\frac{x}{y} + 1 + \frac{y}{x}}{\frac{x}{y} - 1 + \frac{y}{x}} \div \frac{1 - \frac{y^3}{x^5}}{1 + \frac{y^2}{x}}$$

13.

14.

15.

16.

17.

18.

Art.  
found,  
fraction  
them to

1. FI

Here

$$\frac{(x+y)}{2}$$

a



$$13. \frac{3\left(\frac{a-b}{a+b}\right)^3 - \left(\frac{a-b}{a+b}\right)^3 - 3\left(\frac{a-b}{a+b}\right) + 1}{3\left(\frac{a+b}{a-b}\right) - 3\left(\frac{a+b}{a-b}\right)^2 + \left(\frac{a+b}{a-b}\right)^3 - 1}$$

$$14. \frac{x^5 - x^4y + x^3y^2 - x^2y^3 + xy^4 - y^5}{x^5 + x^4y + x^3y^2 + x^2y^3 + xy^4 + y^5} \div \left(\frac{x-y}{x+y}\right)^3.$$

$$15. \left(\frac{1-x^2}{1-x^3} + \frac{1-x}{1-x+x^2}\right) \div \left(\frac{1+x}{1+x+x^2} - \frac{1-x^2}{1+x^3}\right).$$

16. Find the value of

$$\frac{a}{2na - 2nx} + \frac{b}{2nb - 2nx} \text{ when } x = \frac{1}{2}(a+b).$$

17. Find the value of  $\sqrt{1 - \sqrt{1-x}}$

$$\text{when } x = 2\left(\frac{1-b}{1+b}\right)^2 - \left(\frac{1-b}{1+b}\right)^4.$$

18. Find value of

$$\frac{\sqrt{a+bx} + \sqrt{a-bx}}{\sqrt{a+bx} - \sqrt{a-bx}} \text{ when } x = \frac{2ac}{b(1+c^2)}.$$

Art. XXXV. When the sum of several fractions is to be found, it is generally best, instead of reducing at once all the fractions to a common denominator, to take two (or more) of them together, and combine the results.

#### EXAMPLES.

1. Find the sum of

$$\frac{x+y}{2x-2y} - \frac{y-x}{2x+2y} - \frac{x^2-y^2}{x^2+y^2}.$$

Here taking the first two together we have

$$\frac{(x+y)^2 + (x-y)^2}{2(x^2 - y^2)} = \frac{x^2 + y^2}{x^2 - y^2}; \text{ now add this to } -\frac{x^2 - y^2}{x^2 + y^2}.$$

$$\text{and we get } \frac{(x^2 + y^2)^2 - (x^2 - y^2)^2}{x^4 - y^4} = \frac{4x^2y^2}{x^4 - y^4}.$$

2. Find the sum of

$$\frac{1+x}{1-x} + \frac{4x}{1+x^2} + \frac{8x}{1+x^4} - \frac{1-x}{1+x}$$

Here, taking the first and the last together, we have

$$\frac{(1+x)^2 - (1-x)^2}{1-x^2} = \frac{4x}{1-x^2};$$

taking this result with the second fraction, we have

$$4x \left( \frac{1}{1+x^2} + \frac{1}{1-x^2} \right) = \frac{8x}{1-x^4};$$

now take this result with the remaining fraction and we get

$$8x \left( \frac{1}{1-x^4} + \frac{1}{1+x^4} \right) = \frac{16x}{1-x^8}.$$

3.  $\frac{x^{2n}}{x^n-1} - \frac{x^{2n}}{x^n+1} - \frac{1}{x^n-1} + \frac{1}{x^n+1}$ . Taking in pairs those whose denominators are alike, we have

$$\frac{x^{2n}-1}{x^n-1} - \frac{x^{2n}-1}{x^n+1} = x^{2n} + x^n + 1 - (x^n - 1) = x^{2n} + 2.$$

The work is often made easier by *completing the divisions* represented by the fractions.

4. Find the sum of  $1 + \frac{2x+1}{2(x-1)} - \frac{4x+5}{2x+2}$ . By dividing numerators into denominators, this

$$\begin{aligned} &= 1 + 1 + \frac{3}{2x-2} - 2 - \frac{1}{2x+2} = \frac{3}{2x-2} - \frac{1}{2x+2} \\ &= \frac{3x+3-x+1}{2x^2-2} = \frac{x+2}{x^2-1}. \end{aligned}$$

5.  $\frac{x}{x-2} + \frac{x-9}{x-7} - \frac{x+1}{x-1} - \frac{x-8}{x-6}$ : we have, by division

$$\begin{aligned} &1 + \frac{2}{x-2} + 1 - \frac{2}{x-7} - 1 - \frac{2}{x-1} - 1 + \frac{2}{x-6}, \text{ or} \\ &\frac{2}{x-2} + \frac{2}{x-6} - \frac{2}{x-7} - \frac{2}{x-1} = \frac{2(2x-8)}{(x-2)(x-6)} - \frac{2(2x-8)}{(x-1)(x-7)} \\ &= (4x-16) \left\{ \frac{1}{x^2-8x+12} - \frac{1}{x^2-8x+7} \right\} \\ &= (80-20x) \div (x^4-16x^3+88x^2-152x+84). \\ &\quad [\text{denominator} = (x^2-8x)^2 + 19(x^2-8x) + 84]. \end{aligned}$$

6. Find the value of

$$\frac{x+2a}{x-2a} + \frac{x+2b}{x-2b} \text{ when } x = \frac{4ab}{a+b}$$

By division,  $1 + \frac{4a}{x-2a} + 1 + \frac{4b}{x-2b}$

$$= 2 + 4 \left( \frac{a}{x-2a} + \frac{b}{x-2b} \right); \text{ but the quantity in the brackets}$$

$$= \frac{(a+b)x-4ab}{(x-2a)(x-2b)} = 0 \text{ since } (a+b)x = 4ab$$

$\therefore$  the value of the given expression is 2.

### Exercise xliv.

Simplify the following :

1.  $\frac{x-a}{5} + \frac{x^2+ax+a^2}{x+a} - \frac{x^3-a^3}{x^2-a^2}$

2.  $\frac{a^3+b^3}{a^2-ab+b^2} + \frac{a^3-3a^2b+3ab^2-b^3}{a^3-b^3} - \frac{a(a-b)-b(a-b)}{a^2+ab+b^2}$

3.  $\left( \frac{1}{a+x} + \frac{1}{a-x} + \frac{2a}{a^2+x^2} \right) \times$   
 $\left( \frac{1}{a+x} - \frac{1}{a-x} - \frac{2x}{a^2+x^2} \right)$

4.  $\frac{a}{a+b} + \frac{b}{a-b} - \frac{ab}{ab-b^2} + \frac{ab}{a^2+ab}$

5.  $\frac{3+2x}{2-x} - \frac{2-3x}{2+x} + \frac{16x-x^2}{x^2-4}$

6.  $\frac{1}{4a^3(a+x)} + \frac{1}{4a^3(a-x)} + \frac{1}{2a^2(a^2+x^2)}$

7.  $\frac{1}{2} \left( \frac{3x+2y}{3x-2y} \right) - \frac{1}{2} \left( \frac{3x-2y}{3x+2y} \right)$

8.  $\frac{x+1}{2x-1} - \frac{x-1}{2x+1} - \frac{1-3x}{x(1-2x)} + \frac{x}{x(4x^2-1)} + \frac{1}{x(16x^4-1)}$

9.  $\frac{1}{2x+2} - \frac{4}{x+2} + \frac{9}{2(x+3)} - \frac{x-1}{(x+2)(x+3)}$

$$10. \frac{2(x+y)}{x-y} - \frac{2(y-x)}{x+y} - \frac{4(x^2-y^2)}{x^2+y^2} + \frac{4(x^4+y^4)}{x^4-y^4}.$$

$$11. (a-b) \left\{ \frac{1}{(x+a)^2} + \frac{1}{(x+b)^2} \right\} + 2 \left\{ \frac{1}{x+a} - \frac{1}{x+b} \right\}$$

$$12. \left\{ \frac{a+x}{a-x} + \frac{4ax}{a^2+x^2} + \frac{8a^3x}{a^4+x^4} - \frac{a-x}{a+x} \right\} + \left\{ \frac{a^2+x^2}{a^2-x^2} + \frac{4a^2x^2}{a^4+x^4} - \frac{a^2-x^2}{a^2+x^2} \right\}.$$

$$13. \frac{5x-4}{9} + \frac{12x+2}{11x-8} - \frac{10x+17}{18}.$$

$$14. \frac{a}{a^2+b^2} + \frac{a}{a^2-b^2} + \frac{a^2}{(a-b)(a^2+b^2)} - \frac{2a^3-b^3-ab^3}{a^4-b^4}$$

$$15. \frac{12x+10a}{3x+a} + \frac{117a+28x}{9a+2x} - 18.$$

$$16. \frac{4x-17}{x-4} - \frac{8x-30}{2x-7} + \frac{10x-3}{2x-5} - \frac{5x-4}{x-1}.$$

$$17. \text{Find the value of } \frac{a+b+2c}{a+b-2c} + \frac{a+b+2d}{a+b-2d}$$

$$\text{when } a+b = \frac{4cd}{c+d}.$$

$$18. \frac{x^{3n}}{x^n-y^n} - \frac{y^n x^{2n}}{x^n+y^n} - \frac{x^{3n}}{x^n-y^n} + \frac{y^{3n}}{x^n+y^n}$$

$$19. \frac{(a-b)^{3n}}{(a-b)^n-1} - \frac{(a-b)^{2n}}{(a-b)^n+1} - \frac{1}{(a-b)^n-1} + \frac{1}{(a-b)^n+1}.$$

$$20. \frac{1}{(a^2-b^2)(x^2+b^2)} + \frac{1}{(b^2-a^2)(x^2+a^2)} - \frac{1}{(x^2+a^2)(x^2+b^2)}.$$

$$21. \frac{1+x}{1-x^3} + \frac{1-x}{1+x^3} - \frac{2}{1-x^2} - \frac{2x^2}{x^6+1}.$$

$$22. \frac{a^3+a^2b+ab^2+b^3}{a^3-a^2b-ab^2+b^3} \times \frac{(a+b)^2-3ab}{(a-b)^2+3ab} \times \frac{(a-b)^3-a^3+b^3}{(a+b)^3-a^3-b^3}.$$

Art. XXXVI. The following are additional examples in which a knowledge of factoring and of the principle of symmetry is of advantage.

## EXAMPLES.

$$1. \frac{x^2 - (y-z)^2}{(x+z)^2 - y^2} + \frac{y^2 - (z-x)^2}{(y+x)^2 - z^2} + \frac{z^2 - (x-y)^2}{(z+y)^2 - x^2}.$$

Cancelling the common factor  $x-y+z$  in the two terms of the first fraction, there results  $\frac{x+y-z}{x+y+z}$ , hence by symmetry, the denominators of the other two fractions will be  $x+y+z$ , and the numerators will be  $y+z-x$ ,  $z+x-y$ ;  $\therefore$  sum of the three numerators  $= x+y+z$ , and the result  $= 1$ .

$$2. \text{Simplify } \frac{ab}{(c-a)(c-b)} + \frac{bc}{(a-b)(a-c)} + \frac{ca}{(b-c)(b-a)}.$$

The L. C. M. of denominators is evidently  $(a-b)(b-c)(c-a)$ . This gives for numerator of first fraction  $-ab(a-b)$ , and by symmetry the other numerators are  $-bc(b-c)$ ,  $-ca(c-a)$ .

$$\begin{aligned} \therefore \text{ we have } & - \frac{ab(a-b) + bc(b-c) + ca(c-a)}{(a-b)(b-c)(c-a)} \\ & = - \frac{(a-b)(b-c)(a-c)}{(a-b)(b-c)(c-a)} = 1. \end{aligned}$$

2. Reduce the following to a single fraction :

$$\frac{a}{(a-b)(a-c)(x-a)} + \frac{b}{(b-a)(b-c)(x-b)} + \frac{c}{(c-a)(c-b)(x-c)}.$$

Here the L. C. M. is  $(a-b)(b-c)(c-a)(x-a)(x-b)(x-c)$ ; the numerator of the first fraction is

$$\begin{aligned} & -a(b-c)(x-b)(x-c), \text{ and } \therefore \text{ by symmetry that of} \\ \text{second is } & -b(c-a)(x-c)(x-a), \text{ and that of third is} \\ & -c(a-b)(x-a)(x-b); \text{ and their sum is} \\ & -\{a(b-c)(x-b)(x-c) + b(c-a)(x-c)(x-a) + \\ & \quad c(a-b)(x-a)(x-b)\}. \end{aligned}$$

This vanishes if  $a=b$ , hence  $a-b$  is a factor, and  $\therefore$  by symmetry  $b-c$  and  $c-a$  are also factors. Now the product of these

is of the third degree, while the whole expression rises only to the fourth, hence  $x^2$  cannot be involved. The other factor must therefore be of the form  $mx+n$ , in which  $m$  is a number.

To determine  $n$  put  $x=0$ , and the expression becomes  $abc\{a-b+b-c+c-a\}=0$ ;  $\therefore n=0$ , or the other factor is  $mx$ .

To determine  $m$  put  $a=0$ ,  $b=1$ ,  $c=-1$ , and  $m$  will be found to be 1. The numerator is  $\therefore x(a-b)(b-c)(c-a)$ , and the result is

$$\frac{x}{(x-a)(x-b)(x-c)}$$

3. Simplify  $\frac{a+b}{(b-c)(c-a)} + \frac{b+c}{(c-a)(a-b)} + \frac{c+a}{(a-b)(b-c)}$ .

L. C. M. of denominators is  $(a-b)(b-c)(c-a)$ ;

$\therefore$  first numerator is  $a^2 - b^2$ , and by symmetry

second "  $b^2 - c^2$ , and

third "  $c^2 - a^2$ ;

the sum of these = 0, which is the required result.

4. Reduce

$$\frac{2}{x-y} + \frac{2}{y-z} + \frac{2}{z-x} + \frac{(x-y)^2 + (y-z)^2 + (z-x)^2}{(x-y)(y-z)(z-x)}$$

Here the numerator becomes

$$2(y-z)(z-x) + 2(x-y)(z-x) + 2(x-y)(y-z) + (x-y)^2 + (y-z)^2 + (z-x)^2, \text{ which is evidently } \{(x-y) + (y-z) + (z-x)\}^2 = 0.$$

5.  $a^3 \left\{ \frac{a^3 + 2b^3}{a^3 - b^3} \right\}^3 + b^3 \left\{ \frac{2a^3 + b^3}{b^3 - a^3} \right\}^3$ .

Observe that the denominators become the same by changing the sign between the fractions, and that the expression is symmetrical with respect to  $a$  and  $b$ . The numerator of the first fraction is  $a^{12} + 6a^9b^3 + 12a^6b^6 + 8a^3b^9$ , and by symmetry that of the other is  $-b^{12} - 6b^9a^3 - 12b^6a^6 - 8b^3a^9$ . Their sum is  $\therefore a^{12} - b^{12} + 6a^3b^3(a^6 - b^6) - 8a^3b^3(a^6 - b^6)$   
 $= (a^6 - b^6)\{a^6 + b^6 + 6a^3b^3 - 8a^3b^3\} = (a^6 - b^6)(a^3 - b^3)^2$   
 $= (a^3 + b^3)(a^3 - b^3)^3$ , and since the denominator of the given expression is  $(a^3 - b^3)^3$   $\therefore$  the result is  $a^3 + b^3$ .

## Exercise xlv.

Simplify the following :

1.  $x \left( \frac{x-2y}{x+y} \right)^3 + y \left( \frac{2x-y}{x+y} \right)^3.$

2.  $a \left( \frac{a+2b}{a-b} \right)^3 + b \left( \frac{2a+b}{b-a} \right)^3.$

3.  $\frac{a+b}{(b-c)(c-a)} + \frac{b+c}{(c-a)(a-b)} + \frac{c+a}{(a-b)(b-c)}.$

4.  $\frac{1}{(a-b)(a-c)} + \frac{1}{(b-a)(b-c)} + \frac{1}{(c-a)(c-b)}.$

5.  $\frac{a-b}{a+b} + \frac{b-c}{b+c} + \frac{c-a}{c+a} + \frac{(a-b)(b-c)(c-a)}{(a+b)(b+c)(c+a)}.$

6.  $\frac{a^2}{(a+b)(a+c)(x+a)} + \frac{b^2}{(a+b)(b-c)(x+b)} - \frac{c^2}{(a+c)(b-c)(x+c)}.$

7.  $\frac{x^3}{(x-y)(x-z)} + \frac{y^3}{(y-x)(y-z)} + \frac{z^3}{(z-x)(z-y)}.$

8.  $\frac{a^3}{(a-b)(a-c)} + \frac{b^3}{(b-a)(b-c)} + \frac{c^3}{(c-a)(c-b)}.$

9.  $\frac{1}{\left(\frac{a}{b}-1\right)\left(\frac{a}{c}-1\right)} + \frac{1}{\left(\frac{b}{a}-1\right)\left(\frac{b}{c}-1\right)} + \frac{1}{\left(\frac{c}{a}-1\right)\left(\frac{c}{b}-1\right)}.$

10.  $\left( \frac{x^3 - 2y^3}{x^2} \right)^3 + y^3 \left( \frac{2x^3 - y^3}{x^3 + y^3} \right)^3.$

11.  $\frac{1}{(b+c-a)(c+a-2b)} + \frac{1}{(c+a-2b)(a+b-2c)} + \frac{1}{(a+b-2c)(b+c-2a)}.$

12.  $\frac{b^2-c^2}{(b+c)^2} + \frac{c^2-a^2}{(c+a)^2} + \frac{a^2-b^2}{(a+b)^2}.$

13.  $\frac{a^3}{(a-b)(a-c)(x-a)} + \frac{b^3}{(b-a)(b-c)(x-b)} + \frac{c^3}{(c-a)(c-b)(x-c)}.$

$$14. \frac{x(y+z)}{(x-y)(z-x)} + \frac{y(z+x)}{(y-z)(x-y)} + \frac{z(x+y)}{(z-x)(y-z)}$$

$$15. \frac{(a+b)^2 + (b-c)^2 + (a+c)^2}{(a+b)(b-c)(a+c)} = \frac{2}{a+c} - \frac{2}{b-c} + \frac{2}{a+b}$$

$$16. \frac{1}{x(x-a)(x-b)} + \frac{1}{a(b-a)(x-a)} + \frac{1}{b(b-a)(x-b)}$$

SECTION III.—RATIOS.

Art. XXXVII. If  $\frac{a}{b} = \frac{c}{d} \therefore ad = bc$ . Now,

dividing  $ad = bc$  by  $ca$  we have  $\frac{b}{a} = \frac{d}{c} \dots (1)$ .

“  $ad = bc$  by  $cd$  “  $\frac{a}{c} = \frac{b}{d} \dots (2)$ .

“  $ad = bc$  by  $ab$  “  $\frac{d}{b} = \frac{c}{a} \dots (3)$ .

Also  $\frac{ma + nc}{mb + nd} =$  each of the given fractions  $\dots (4)$ .

For  $\frac{ma + nc}{mb + nd} = \frac{mb\left(\frac{a}{b}\right) + nd\left(\frac{c}{d}\right)}{mb + nd} = \frac{(mb + nd) \frac{a}{b}}{mb + nd} = \frac{a}{b}$  or  $\frac{c}{d}$ .

A very important case of this is  $m = 1, n = \pm 1$ , hence

$$\frac{a}{b} = \frac{c}{d} = \frac{a+c}{b+d} = \frac{a-c}{b-d} \dots (5)$$

Also  $\frac{a-b}{a+b} = \frac{c-d}{c+d} \dots (6)$ .

For by (2) and (5)

$$\frac{a}{c} = \frac{b}{d} = \frac{a-b}{c-d} = \frac{a+b}{c+d} \therefore \frac{a-b}{a+b} = \frac{c-d}{c+d}$$

Or thus:  $\frac{a-b}{a+b} = \frac{\frac{a}{b} - 1}{\frac{a}{b} + 1} = \frac{\frac{c}{d} - 1}{\frac{c}{d} + 1} = \frac{c-d}{c+d}$



Generally, to prove that if  $\frac{a}{b} = \frac{c}{d}$ , any fraction whose numerator and denominator are homogeneous functions of  $a$  and  $b$ , and are of the same degree, will be equal to a similar fraction formed with  $c$  instead of  $a$  and  $d$  instead of  $b$ :—Express the first fraction in terms of  $\frac{a}{b}$ , and for  $\frac{a}{b}$  substitute its equivalent  $\frac{c}{d}$ , and reduce the result.

By (2) the fractions may be formed of  $a$  and  $c$ , and  $b$  and  $d$ .

$$\text{If } \frac{a}{b} = \frac{c}{d} = \frac{e}{f}, \quad \frac{ma+nc+pe}{mb+nd+pf} = \frac{a}{b} \text{ or } \frac{c}{d} \text{ or } \frac{e}{f} \dots\dots (7)$$

$$\frac{ma+nc+pe}{mb+nd+pf} = \frac{mb\left(\frac{a}{b}\right) + nd\left(\frac{c}{d}\right) + pf\left(\frac{e}{f}\right)}{mb+nd+pf}$$

(1).

(2).

$$= \frac{(mb+nd+pf)\frac{a}{b}}{mb+nd+pf} = \frac{a}{b}$$

(3).

$$\text{If } \frac{a}{b} = \frac{c}{d} \text{ and } \frac{m}{n} = \frac{p}{q}$$

(4).

$$\frac{ma \pm pc}{nb \pm qd} = \frac{pa \pm mc}{qb \pm nd} = \frac{ma}{nb} \text{ or } \frac{pa}{qb}, \text{ or } \&c \dots\dots (8)$$

$$\text{For } \frac{ma}{nb} = \frac{pc}{qd} = \frac{ma \pm pc}{nb \pm qd} \text{ by (5)}$$

$$\frac{pa}{qb} = \frac{mc}{nd} = \frac{pa \pm mc}{qb \pm nd}$$

(5).

$$\text{But } \frac{ma}{nb} = \frac{pa}{qb}, \text{ hence the equality stated in (8).}$$

(6).

$$\text{If } \frac{a}{b} = \frac{c}{d} = \frac{e}{f} \text{ and } \frac{m}{n} = \frac{p}{q} = \frac{r}{s},$$

$$\frac{ma \pm pc \pm re}{nb \pm qd \pm sf} = \frac{pa \pm rc \pm me}{qb \pm sd \pm nf} = \&c., = \frac{ma}{nb} = \&c. \dots (9)$$

If an upper sign be taken in a numerator, the corresponding upper sign must be taken in the denominator; if a lower sign, the corresponding lower sign, otherwise all the signs are independent of each other.

## EXAMPLES.

1. If  $\frac{a}{b} = \frac{c}{d}$ , show that  $\frac{5a-4b}{7a+5b} = \frac{5c-4d}{7c+5d}$

The given fraction =  $\frac{5\frac{a}{b} - 4}{7\frac{a}{b} + 5} = \frac{5\frac{c}{d} - 4}{7\frac{c}{d} + 5} = \frac{5c-4d}{7c+5d}$

2. If  $\frac{a}{b} = \frac{c}{d}$  shew that  $\frac{2a^3+3a^2b}{3a^2b-4b^3} = \frac{2c^3+3c^2d}{3c^2d-4d^3}$ .

Dividing the given fraction by  $b^3$  we have

$$\frac{2\frac{a^3}{b^3} + 3\frac{a^2}{b^2}}{\frac{a^2}{b^2} - 4}, \text{ and this becomes, on substituting for } \frac{a}{b} \text{ its equal } \frac{c}{d},$$

$$\frac{2\frac{c^3}{d^3} + 3\frac{c^2}{d^2}}{\frac{c^2}{d^2} - 4} = \frac{2c^3 + 3c^2d}{3c^2d - 4d^3}.$$

3. If  $3a=2b$ , find the value of  $\frac{a^3+b^3}{a^2b-ab^2}$ . This =  $\left(\frac{a^3}{b^3} + 1\right) \div \left(\frac{a^2}{b^2} - \frac{a}{b}\right)$  [by dividing both numerator and denominator by  $b^3$ ]. But from the given relation  $\frac{a}{b} = \frac{2}{3}$  we have, by substituting for  $\frac{a}{b}$ ,

$$\left(\frac{8}{27} + 1\right) \div \left(\frac{4}{9} - \frac{2}{3}\right) = 35 \div (-6) = -5\frac{5}{6}.$$

4. If  $\frac{a}{b} = \frac{c}{d}$ . Prove that  $\frac{a^3+b^3}{c^3+d^3} \times \frac{b}{d} = \left(\frac{a+b}{c+d}\right)^4$ .

We have  $\frac{a}{c} = \frac{b}{d} = \frac{a+b}{c+d}$ . Also

$$\frac{a^3+b^3}{c^3+d^3} = \frac{b^3}{d^3} \left(\frac{a^3}{b^3} + 1\right) \div \left(\frac{c^3}{d^3} + 1\right) = \frac{b^3}{d^3}, \text{ and this multiplied}$$

by  $\frac{b}{d}$  gives  $\frac{b^4}{d^4} = \left(\frac{a+b}{c+d}\right)^4$ .

5. If  $\frac{x^3+ax^2-bx+c}{x^3-ax^2+bx+c} = \frac{x^2+ax-b}{x^2-ax+b}$ , shew that  $x = \frac{b}{a}$ .

Multiplying both terms of second fraction by  $x$ , it becomes

$$\frac{x^3+ax^2-bx}{x^3-ax^2+bx}; \text{ now each of the given fractions =}$$

$\frac{\text{difference of numerators}}{\text{difference of denominators}};$

$$= \frac{c}{c} = 1 \therefore x^2+ax-b=x^2-ax+b$$

$$\text{or } 2ax = 2b \therefore x = \frac{b}{a}.$$

6. If  $\frac{a}{b} = \frac{c}{d} = \frac{e}{f}$ , shew that  $\frac{ac+ce+ea}{bd+df+fb} = \frac{a^2+c^2+e^2}{b^2+d^2+f^2}$ .

For  $\frac{ac}{bd} = \frac{ce}{df} = \frac{ea}{fb} = \frac{ac+ce+ea}{bd+df+fb}$ . By (7) making  $m=n=p=1$ .

Also  $\frac{a^2}{b^2} = \frac{c^2}{d^2} = \frac{e^2}{f^2} = \frac{a^2+c^2+e^2}{b^2+d^2+f^2}$ . By (7).

But  $\frac{ac}{bd} = \frac{a^2}{b^2}$  hence the required equality.

The problem is a particular case of (9), with all the signs + and  $a$  for  $m$ ,  $b$  for  $n$ ,  $c$  for  $p$ , &c.

(If the fractions given equal to one another have not monomial terms, instead of seeking to express the proposed quantity in terms of one fraction and then substituting an equivalent fraction, it is often better to assume a single letter to represent the common value of the fractions given equal, and to work in terms of this assumed letter.)

7. If  $\frac{a+b}{3(a-b)} = \frac{b+c}{4(b-c)} = \frac{c+a}{5(c-a)}$ ,

prove that  $32a+35b+27c=0$ .

Assume each of the given fractions =  $x$ , so that  $a+b=3(a-b)x$ ,  
 $b+c=4(b-c)x$ ,  $c+a=5(c-a)x$ ,

$$\text{or } \frac{a+b}{3} + \frac{b+c}{4} + \frac{c+a}{5} = x(a-b+b-c+c-a) = 0.$$

$\therefore$  adding these fractions we have  $32a+35b+27c=0$ .

This example might also be worked as a particular case of (7), thus

$$\begin{aligned} \frac{a+b}{3(a-b)} &= \frac{b+c}{4(b-c)} = \frac{c+a}{5(c-a)} \\ &= \frac{20(a+b) + 15(b+c) + 12(c+a)}{60(a-b) + 60(b-c) + 60(c-a)} = \frac{32a+35b+27c}{0}, \end{aligned}$$

$$\therefore 32a+35b+27c = 0 \times \frac{a+b}{3(a-b)} = 0.$$

8. If  $\frac{a^2}{b^2} + \frac{c^2}{f^2} = \frac{2c}{d} \left\{ \frac{a}{b} - \frac{c}{d} + \frac{e}{f} \right\}$ , prove that

$$\left( \frac{a+c+e}{b+d+f} \right)^2 = \frac{a^2+c^2+e^2}{b^2+d^2+f^2}.$$

Transposing terms, &c., we have

$$\frac{a^2}{b^2} - \frac{2ac}{bd} + \frac{c^2}{d^2} + \frac{e^2}{f^2} - \frac{2ce}{df} + \frac{c^2}{d^2} = 0,$$

$$\text{or } \left( \frac{a}{b} - \frac{c}{d} \right)^2 + \left( \frac{e}{f} - \frac{c}{d} \right)^2 = 0;$$

that is, the sum of two essentially positive quantities = 0;

$\therefore$  each of them must = 0; hence we have

$$\frac{a}{b} - \frac{c}{d} = 0, \text{ and } \frac{e}{f} - \frac{c}{d} = 0;$$

$$\therefore \frac{a}{b} = \frac{c}{d} = \frac{e}{f}; \therefore \frac{a^2}{b^2} = \frac{a^2+c^2+e^2}{b^2+d^2+f^2}.$$

$$\text{Also } \frac{a}{b} = \frac{a+c+e}{b+d+f}; \therefore \frac{a^2}{b^2} = \left( \frac{a+c+e}{b+d+f} \right)^2;$$

$$\therefore \left( \frac{a+c+e}{b+d+f} \right)^2 = \frac{a^2+c^2+e^2}{b^2+d^2+f^2}.$$

## Exercise xlvi.

1. If  $\frac{a}{b} = \frac{c}{d}$ , prove  $\frac{a^2 - ab + b^2}{ab - 4b^2} = \frac{c^2 - cd + d^2}{cd - 4d^2}$ .
2. If  $\frac{a}{b} = \frac{c}{d}$ , prove  $\frac{a^2 - c^2}{b^2 - d^2} = \left(\frac{a-c}{b-d}\right)^2 = \left(\frac{a+c}{b+d}\right)^2$ .
3. Given the same, shew that each of these fractions
- $$= \sqrt{\left(\frac{a^2 + c^2}{b^2 + d^2}\right)}.$$

4. If  $2x = 3y$ , write down the value of

$$\frac{2x^3 - x^2y + y^3}{x^2y + xy^2 + 2y^3}, \text{ and of } \frac{x^4 - 3x^3y + 2y^4}{(x^2 - y^2)^2}.$$

5. If  $\frac{a}{b} = \frac{c}{d} = \frac{e}{f}$ , shew that  $\frac{a}{b} = \frac{ma - nc - pe}{mb - nd - pf}$ .
6. From the same relations prove that  $\frac{a^3}{b^3} = \left(\frac{a - mc - ne}{b - md - nf}\right)^3$ .
7. If  $\frac{1+x}{1-x} = \frac{b}{a} \left(\frac{1+x+x^2}{1-x+x^2}\right)$ , then  $x^3 = (b-a) \div (b+a)$ .
8. If  $\frac{\sqrt{a+x} + \sqrt{a-x}}{\sqrt{a+x} - \sqrt{a-x}} = a$ , prove that  $x = \frac{2a^2}{1+a^2}$ .
9. If  $\frac{mx+a+b}{nx+a+c} = \frac{mx-c-d}{nx-b-d}$ , prove  $x = \frac{b-c}{n-m}$ .
10. If  $\frac{a-b}{ay+bx} = \frac{b-c}{bz+cx} = \frac{c-a}{cy+az} = \frac{a+b+c}{ax+by+cz}$ ,

then each of these fractions =  $\frac{1}{x+y+z}$ ,  $a+b+c$  not being zero.

11. If  $\frac{a+b}{a-b} = \frac{b+c}{2(b-c)} = \frac{c+a}{3(c-a)}$ , then  $8a+9b+5c=0$ .
12. If  $\frac{\sqrt{a} + \sqrt{a-x}}{\sqrt{a} - \sqrt{a-x}} = \frac{1}{a}$ , shew that  $\frac{a-x}{a} = \left(\frac{1-a}{1+a}\right)^2$ .
13. If  $\frac{x^2 - yz}{x(1-yz)} = \frac{y^2 - xz}{y(1-xz)}$ , and  $x, y, z$  be unequal, shew that each of these fractions is equal to  $x+y+z$ .

14. If  $\frac{x^2+2x+1}{x^2-2x+3} = \frac{y^2+2y+1}{y^2-2y+3}$ , show that each of these fractions  $= (xy-1) \div (xy-3)$ .

15. If  $\frac{25x^2-16}{10x+8} = \frac{3(x^2-4)}{2x-4}$ , shew that  $\frac{x-4}{x+5} = \frac{3}{5}$ .

16. If  $y = \frac{4bc}{b+c}$  shew that  $\frac{y+2b}{y-2b} + \frac{y+2c}{y-2c} = 2$ .

17. If  $\frac{1}{4} \left( \frac{a^2+b^2}{a^2-b^2} \right) = \frac{1}{5} \left( \frac{b^2+c^2}{b^2-c^2} \right) = \frac{1}{6} \left( \frac{c^2+a^2}{c^2-a^2} \right)$ , prove that  $25a^2+27b^2+22c^2=0$ .

18. If  $\frac{a^2}{x^2-yz} = \frac{b^2}{y^2-zx} = \frac{c^2}{z^2-xy}$ , shew that  $a^2x+b^2y+c^2z = (a^2+b^2+c^2)(x+y+z)$ .

19. If  $\frac{x}{a+b-c} = \frac{y}{b+c-a} = \frac{z}{c+a-b}$ , then will  $(a-b)x + (b-c)y + (c-a)z = 0$ .

20. If  $\frac{a}{b} = \frac{c}{d} = \frac{e}{f}$  then  $\left( \frac{a^2+c^2+e^2}{b^2+d^2+f^2} \right)^2 = \frac{a^4+c^4+e^4}{b^4+d^4+f^4}$ .

21. If  $\frac{bx+ay}{a-b} = \frac{cy+bz}{b-c} = \frac{az+cx}{c-a}$ , shew that  $(a+b+c)(x+y+z) = ax+by+cz$ .

22. If  $\frac{x^3-5x^2a-a^3+5xa^2}{x^3+x^2a+xa^2+a^3} = \frac{x-a}{x+a}$ , shew that each of these expressions = 1.

23. If  $\frac{1}{6} \left( \frac{a-b}{a+b} \right) = \frac{1}{5} \left( \frac{b-c}{b+c} \right) = \frac{1}{10} \left( \frac{c-a}{c+a} \right)$ , and  $a, b, c$  be different, shew that  $16a+11b+15c=0$ .

24. If  $\left( \frac{x+yz}{y+zx} \right)^2 = \frac{1-y^2}{1-x^2}$ , prove that  $x^2+y^2+z^2+2xyz=1$ .

25. If  $\frac{a}{x-y} = \frac{b}{y-z} = \frac{c}{z-x}$ , shew that  $a+b+c=0$ .

26. If  $\frac{a}{b} = \frac{c}{a}$ , prove that  $\frac{a+b}{a-b} = \frac{\sqrt{ac} + \sqrt{bd}}{\sqrt{ac} - \sqrt{bd}}$ .

27. If  $\frac{a}{b} = \frac{c}{d} = \frac{e}{f}$ , then each is equivalent to

$$\frac{la+mc+ne}{lb+md+nf}, \text{ hence shew that}$$

$$\frac{a}{2z+2x-y} = \frac{b}{2c+2y-z} = \frac{c}{2y+2z-x}, \text{ when}$$

$$\frac{x}{2a+2b-c} = \frac{y}{2b+2c-a} = \frac{z}{2c+2a-b}.$$

28. If  $\frac{a}{b} = \frac{c}{d}$ , prove that  $\left(\frac{a-b}{c-d}\right)^n = \sqrt[n]{\frac{a^{2n}+b^{2n}}{c^{2n}+d^{2n}}}$ .

29. If  $\frac{x}{a(y+z)} = \frac{y}{b(x+z)} = \frac{z}{c(x+y)}$ , prove that

$$\frac{x}{a}(y-z) + \frac{y}{b}(z-x) + \frac{z}{c}(x-y) = 0.$$

30. If  $\frac{a}{lx(ny-mz)} = \frac{b}{my(lz-nx)} = \frac{c}{nz(mx-ly)}$ , then will

$$\frac{a}{lx}(l-x) + \frac{b}{my}(m-y) + \frac{c}{nz}(n-z) = 0.$$

31. If  $z = \frac{\sqrt{(ay^2-a^2)}}{y}$ , and  $y = \frac{\sqrt{(ax^2-a^2)}}{x}$ , shew that

$$x = \frac{\sqrt{(az^2-a^2)}}{z}.$$

32. If  $\frac{x^2-yz}{a^2} = \frac{y^2-xz}{b^2} = \frac{z^2-xy}{c^2} = 1$ , shew that

$$x+y+z = \frac{a^2x+b^2y+c^2z}{a^2+b^2+c^2}.$$

33. If  $\frac{m}{x} = \frac{n}{y} = \frac{r}{z}$ , and  $\frac{x^2}{a^2} = \frac{y^2}{b^2} = \frac{z^2}{c^2} = 1$ ,

prove that  $\frac{m^2}{a^2} + \frac{n^2}{b^2} + \frac{r^2}{c^2} = 3 \frac{m^2+n^2+r^2}{x^2+y^2+z^2}$ .

34. If  $\frac{a}{b} = \frac{c}{d} = \frac{e}{f} = \&c.$ , then

$$\frac{a^{3n}-e^{3n}}{b^{3n}-d^{3n}} = \frac{a^n c^n e^n - (a^n - c^n + e^n)^3}{b^n d^n f^n - (b^n - d^n + f^n)^3}$$

35. If  $\frac{a_1}{b_1} = \frac{a_2}{b_2} = \frac{a_3}{b_3} = \dots = \frac{a_n}{b_n}$ , then

$$\frac{a_1 a_2 - a_2 a_3 + \dots + (-1)^{n-1} a_{n-1} a_n}{b_1 b_2 - b_2 b_3 + \dots + (-1)^{n-1} b_{n-1} b_n} = \frac{a_1 \sqrt{a_2 a_3} + a_2 \sqrt{a_3 a_4} + \dots + a_{n-1} \sqrt{a_n a_1}}{b_1 \sqrt{b_2 b_3} + b_2 \sqrt{b_3 b_4} + \dots + b_{n-1} \sqrt{b_n b_1}}$$

36. If  $\frac{A+B+C}{abc} = \frac{A}{a} + \frac{B}{b} + \frac{C}{c}$ ,

$$\text{and } (A+B+C)(a+b+c) = Aa + Bb + Cc,$$

$$\text{then will } \frac{A}{1+a^2} + \frac{B}{1+b^2} + \frac{C}{1+c^2} = 0.$$

$$\text{and also } \frac{A}{a + \frac{1}{a}} + \frac{B}{b + \frac{1}{b}} + \frac{C}{c + \frac{1}{c}} = 0.$$

37. If  $\frac{xh}{a^2} = \frac{yk}{b^2} = \frac{zl}{c^2}$ , and  $\frac{x^2}{a^2} = \frac{y^2}{b^2} = \frac{z^2}{c^2} = 1$ ,

$$\text{then will } \left( \frac{x}{\frac{1}{3}h} + \frac{y}{k} + \frac{z}{c} \right)^2 = \frac{a^2}{h^2} + \frac{b^2}{k^2} + \frac{c^2}{l^2}.$$

#### SECTION IV.—COMPLETE SQUARES, &c.

1. What quantity must be added to  $x^2 + px$  to make it a complete square?

Let  $r$  be the quantity.

$$\begin{aligned} \text{Then } x^2 + px + r &= \text{complete square} = (x + \sqrt{r})^2 \\ &= x^2 + 2x\sqrt{r} + r. \end{aligned}$$

Equating coefficients we have

$$\begin{aligned} 2\sqrt{r} &= p \\ \therefore r &= \frac{p^2}{4} = \left(\frac{p}{2}\right)^2 \end{aligned}$$

Or thus: Since  $(a+x)^2 = a^2 + 2ax + x^2$ ; we observe, (See Art. XII), that *four times the product of the extremes is equal to the square of the mean*,

$$\therefore 4x^2 r = p^2 x^2;$$

$$\therefore r = \left(\frac{p}{2}\right)^2, \text{ as before.}$$



Or we may extract the square root and equate the remainder to zero. thus

$$\begin{array}{r}
 x^2 + px + r(x + \frac{p}{2}) \\
 \hline
 x^2 \\
 \hline
 2x + \frac{p}{2} \quad px + r \\
 \hline
 px + \frac{p^2}{4} \\
 \hline
 r - \frac{p^2}{4}
 \end{array}$$

Now, if the expression be a complete square, this remainder must vanish ; hence we have

$$r = \frac{p^2}{4} = \left(\frac{p}{2}\right)^2$$

2. Find the relation connecting  $a, b, c$ , if  $ax^2 + bx + c$  is a complete square.

$$\text{Assume } ax^2 + bx + c = (\sqrt{a}x + \sqrt{c})^2 = ax^2 + 2\sqrt{ac}x + c.$$

Now, since this holds for all values of  $x$ , we have  $2\sqrt{ac} = b$ , or  $b^2 = 4ac$ , the relation required.

3. Determine the relation amongst  $a, b, c$ , in order that

$$a^2x^2 + bx + bc + b^2 \text{ may be a perfect square.}$$

As in Ex. 1, we have  $4a^2x^2(bc + b^2) = b^2x^2$  ;

$$\therefore \frac{1}{4a^2} - \frac{c}{b} = 1.$$

Or thus :

$$\begin{aligned}
 \text{Assume } a^2x^2 + bx + bc + b^2 &= (ax + \sqrt{bc + b^2})^2 \\
 &= a^2x^2 + 2a\sqrt{bc + b^2}x + bc + b^2.
 \end{aligned}$$

Equating coefficients, we have  $b = 2a\sqrt{bc + b^2}$  ;

$$\therefore \frac{1}{4a^2} - \frac{c}{b} = 1, \text{ as before.}$$

The same result may also be obtained by extracting the square root and equating the remainder to zero.

4. Show that if  $x^4 + ax^3 + bx^2 + cx + d$  be a complete square, the coefficients satisfy the equation  $c^2 - a^2d = 0$ .

Is it necessary that the coefficients satisfy any other equation?

Extracting the square root of  $x^4 + ax^3 + bx^2 + cx + d$  in the usual manner, we have for the final remainder

$$\left\{ c - \frac{a}{2} \left( b - \frac{a^2}{4} \right) \right\} x + d - \frac{1}{4} \left( b - \frac{a^2}{4} \right)^2$$

Now, if the expression be a complete square, this remainder must vanish; and, that it may vanish for general values of  $x$ , we must have

$$c - \frac{a}{2} \left( b - \frac{a^2}{4} \right) = 0 \quad \dots \dots \dots (1),$$

$$d - \frac{1}{4} \left( b - \frac{a^2}{4} \right)^2 = 0 \quad \dots \dots \dots (2);$$

$$\text{Eliminating } b - \frac{a^2}{4}, \text{ we have } c^2 - a^2d = 0 \quad \dots \dots (3).$$

The coefficients must satisfy the equations (1) and (2), and therefore either of these equations, together with the equation (3), which results from them.

The same result may be obtained by assuming

$$\begin{aligned} x^4 + ax^3 + bx^2 + cx + d &= (x^2 + \frac{1}{2}ax + \sqrt{d})^2 \\ &= x^4 + ax^3 + 2x^2\sqrt{d} \\ &\quad + \frac{1}{4}a^2x^2 + ax\sqrt{d} + d. \end{aligned}$$

$$\text{Equating coefficients, we have } 2\sqrt{d} + \frac{1}{4}a^2 = b \quad \dots \dots (1)$$

$$\text{and } a\sqrt{d} = c \quad \dots \dots (2).$$

From (2) we have  $c^2 - a^2d = 0$ , as before.

5. What must be the value of  $m$  and  $n$  if  $4x^4 - 12x^3 + 25x^2 - 4mx + 8n$  is a perfect square?

$$\begin{aligned} \text{Assume the expression} &= \{(2x^2 - 3x + \sqrt{(8n)})\}^2 \\ &= 4x^4 - 12x^3 + 4x^2\sqrt{(8n)} + 9x^2 - 6x\sqrt{(8n)} + 8n. \end{aligned}$$

$$\text{Equating coefficients, we have } 6\sqrt{(8n)} = 4m \quad \dots \dots (1),$$

$$\text{and } 4\sqrt{(8n)} + 9 = 25 \quad \dots \dots (2);$$

$$\therefore n = 2,$$

$$m = 6.$$

Or remain have

6. I and b Assu

Equ

dividin

dividing

7. F

Assu

Equa

8. Sh rational equal to

Or thus: Extracting the square root in the ordinary way, the remainder is found to be  $(-4m+24)x+8n-16$ ;  $\therefore$  we must have

$$4m+24=0, \text{ or } m=6, \\ \text{and } 8n-16=0, \text{ or } n=2.$$

6. If  $ax^3+bx^2+cx+d$  be a complete cube, shew that  $ac^3=db^3$ , and  $b^2=3ac$ .

$$\text{Assume } ax^3+bx^2+cx+d=(x+d^{\frac{1}{3}})^3 \\ = ax^3+3a^{\frac{2}{3}}d^{\frac{1}{3}}x^2+3a^{\frac{1}{3}}d^{\frac{2}{3}}x+d$$

Equating coefficients,

$$b=3a^{\frac{2}{3}}d^{\frac{1}{3}} \dots \dots \dots (1)$$

$$c=3a^{\frac{1}{3}}d^{\frac{2}{3}} \dots \dots \dots (2);$$

dividing (1) by (2),  $\frac{b}{c} = \frac{a^{\frac{2}{3}}}{d^{\frac{1}{3}}};$   
 $ac^3=db^3.$

Also,  $b^2=9a^{\frac{4}{3}}d^{\frac{2}{3}} \dots \dots \dots (3);$

dividing (3) by (2),  $\frac{b^2}{c} = 3a;$

$$\therefore b^2=3ac.$$

7. Find the relations subsisting between  $a, b, c, d, e$ , when  $ax^4+bx^3+cx^2+dx+e$  is a complete fourth power.

Assume  $ax^4+bx^3+cx^2+dx+e=(a^{\frac{1}{4}}x+e^{\frac{1}{4}})^4$   
 $= ax^4+4a^{\frac{3}{4}}e^{\frac{1}{4}}x^3+6a^{\frac{1}{2}}e^{\frac{1}{2}}x^2+4a^{\frac{1}{4}}e^{\frac{3}{4}}x+e.$

Equating coefficients, we have

$$b=4a^{\frac{3}{4}}e^{\frac{1}{4}},$$

$$c=6a^{\frac{1}{2}}e^{\frac{1}{2}},$$

$$d=4a^{\frac{1}{4}}e^{\frac{3}{4}};$$

whence  $bd=16ac. \dots \dots \dots (1).$

$$bc=24a^{\frac{5}{4}}e^{\frac{3}{4}}=6a \cdot 4a^{\frac{1}{4}}e^{\frac{3}{4}}=6ad. \dots \dots \dots (2).$$

$$cd=24a^{\frac{3}{4}}e^{\frac{5}{4}}=6e \cdot 4a^{\frac{3}{4}}e^{\frac{1}{4}}=6be. \dots \dots \dots (3).$$

8. Shew that  $x^4+px^3+qx^2+rx+s$  can be so resolved into two rational quadratic factors if  $s$  be a perfect square, negative, and

equal to  $\frac{r^2}{p^2-4q}$ .

Since  $-s$  is a perfect square, let it be  $n^2$ .

$$\begin{aligned} \text{Assume } x^4 + px^3 + qx^2 + rx - n^2 \\ &= (x^2 + mx + n)(x^2 + m'x - n) \\ &= x^4 + (m + m')x^3 + mm'x^2 - n(m - m')x - n^2. \end{aligned}$$

Equating coefficients, we have

$$\begin{aligned} m + m' &= p \\ mm' &= q \\ m - m' &= \frac{r}{n} \\ m^2 + 2mm' + m'^2 &= p^2 \\ 4mm' &= 4q ; \end{aligned}$$

$$\therefore (m - m')^2 = p^2 - 4q = \frac{r^2}{n^2}$$

$$\therefore \frac{r^2}{p^2 - 4q} = n^2 = -s.$$

#### Exercise xlvii.

1. What is the condition that  $(a-x)(b-x) - c^2$ , may be a perfect square.

2. Find the value of  $n$  which will make  $2x^2 + 8x + n$ , a perfect square.

3. Find a value of  $x$  which will make  $x^4 + 6x^3 + 11x^2 + 3x + 31$ , a perfect square.

4. Extract the square root of

$$(a-b)^4 - 2(a^2 + b^2)(a-b)^2 + 2(a^4 + b^4)$$

5. Find the values of  $m$  and  $n$  which will make  $4x^4 - 4x^3 + 5x^2 - mx + n$ , a perfect square.

6. What must be added to  $x^4 - \sqrt{4x^4 - 16x^2 + 16} - 4x^2$  in order to make it a complete square?

7. The expression  $x^4 + x^3 - 16x^2 - 4x + 48$ , is resolvable into two factors of the form  $x^2 + mx + 6$ , and  $x^2 + nx + 8$ ; determine the factors.

8. Find the value of  $c$  which will make  $4x^4 - cx^3 + 5x^2 + \frac{cx}{2} + 1$ , a complete square.

9. Obtain the square root of  $4\{(a^2 - b^2)cd + ab(c^2 - d^2)\}^2 + \{(a^2 - b^2)(c^2 - d^2) - 4abcd\}^2$ .

10. If  $(a - b)x^2 + (a + b)^2x + (a^2 - b^2)(a + b)$ , is a complete square, then  $a = 3b$ , or  $b = 3a$ .

11. Find the simplest quantity which, subtracted from  $a^2x^2 + 4abx + 4acx + 5bc + b^2c^2$ , will give for remainder an exact square.

12.  $x^4 - 4x^3 - x^2 + 16x - 12$  is resolvable into quadratic factors of the form  $x^2 + mx + p$ , and  $x^2 + nx + q$ : find them.

13. Find the values of  $m$  which will make  $x^2 + max + a^2$  a factor of  $x^4 - ax^3 + a^2x^2 - a^3x + a^4$ .

14. Shew that if  $x^4 + ax^3 + bx^2 + cx + d$  be a perfect square, the coefficients satisfy the relations

$$8c = a(4b - a^2), \text{ and}$$

$$64d = (4b - a^2)^2.$$

15. Investigate the relations between the coefficients in order that  $ax^2 + by^2 + cz^2 + dxy + eyz + fax$  may be a complete square.

16. If  $x^3 + ax^2 + bx + c$  is exactly divisible by  $(x + d)^2$ , shew that

$$\frac{1}{2}(b - d^2) = \frac{c}{d} = d(a - 2d)$$

17. Determine the relations among  $a, b, c, d$ , when  $ax^3 - bx^2 + cx - d$ , is a complete cube.

18. The polynome  $ax^3 + 3bx^2 + 3cx + d$  is exactly divisible by  $(a - x)^2$ ; shew that  $(ad - bc)^2 = 4(ac - b^2)(bd - c^2)$ .

19. Find the relation between  $p$  and  $q$ , when  $x^3 + px^2 + q$ , is exactly divisible by  $(x - a)^2$ .

20. If  $x^2 + nax + a^2$  is a factor of  $x^4 + ax^3 + a^2x^2 + a^3x + a^4$ , shew that  $n^2 - n - 1 = 0$ .

21. If  $x^4 + ax^3 + bx^2 + cx + d$ , be the product of two complete squares, shew that

$$(4b - a^2)^2 = 64d, (4b - a^2)a = 8c, a\sqrt{(3a^2 - 2b)} = 3b.$$

22. Prove that  $x^4 + px^3 + qx^2 + rx + s$  is a perfect square, if

$$p^2s = r^2, \text{ and } q = \frac{p^2}{4} + 2\sqrt{s}.$$

23. If  $ax^3 + 3bx^2 + 3cx + d$  contain  $ax^2 + 2bx + c$  as a factor, the former will be a complete cube, and the latter a complete square.

24. If  $m^2x^2 + px + pq + q^2$  be a perfect square, find  $p$  in terms of  $m$ ,  $q$ , and  $x$ .

25. Find the relation between  $p$  and  $q$  in order that  $x^3 + px^2 + qx + r$  may contain  $(x+2)^2$  as a factor.

26. If  $x^3 + px^2 + qx + r$  be algebraically divisible by  $3x^2 + 2px + q$ , show that the quotient is  $x + \frac{p}{3}$ .

---

RELATION IN INVOLUTION.

**Art XXXVIII.** If  $aa' = bb' = cc'$ , prove that

$$\begin{aligned} 1. (a+b')(b+c')(c+a') &= (a'+b)(b'+c)(c'+a) \\ (a+b') \times a' &= aa' + b'a' = bb' + b'a' = (b+a') \times b' \\ (b+c') \times b' &= bb' + c'b' = cc' + c'b' = (c+b') \times c' \\ (c+a') \times c' &= cc' + a'c' = aa' + a'c' = (a+c') \times a' \end{aligned}$$

$$\begin{aligned} \therefore (a+b')(b+c')(c+a') \times a'b'c' &= \\ (a'+b)(b'+c)(c'+a) \times b'c'a' & \\ \therefore (a+b')(b+c')(c+a') &= (a'+b)(b'+c)(c'+a). \end{aligned}$$

$$\begin{aligned} 2. (a+b)(a+b')(a'-c)(a'-c') &= (a'+b)(a'+b')(a-c)(a-c'). \\ (a+b) \times a' &= aa' + a'b = bb' + a'b = (b'+a') \times b \\ (a+b') \times a' &= aa' + a'b' = bb' + a'b' = (b+a') \times b' \\ (a'-c) \times a &= aa' - ac = cc' - ac = (c'-a) \times c \\ (a'-c') \times a &= aa' - ac' = cc' - ac' = (c-a) \times c' \end{aligned}$$

$$\begin{aligned} \therefore (a+b)(a+b')(a'-c)(a'-c') \times (aa')^2 &= \\ (b'+a')(b+a')(c'-a)(c-a) \times bb'.cc' & \end{aligned}$$

$$\text{But } bb'.cc' = (aa')^2,$$

$$\text{and } (c'-a)(c-a) = (a-c)(a-c')$$

$$\therefore (a+b)(a+b')(a'-c)(a'-c') = (a'+b)(a'+b')(a-c)(a-c').$$

## Exercise xlviii.

If  $aa' = bb' = cc'$  prove that

$$1. (a-b')(b-c)(c'-a') = (b-a')(a-c)(c'-b').$$

$$2. (b-c')(c-a)(a'-b') = (c-b')(b-a)(a'-c').$$

$$3. (c-a')(a-b)(b'-c') = (a-c')(c-b)(b'-a').$$

$$4. (a-b')(b-c')(c-a') = (a-c')(b-a')(c'-b').$$

$$5. \frac{(a-b)(a-b')}{(a'-b)(a'-b')} = \frac{(a-c)(a-c')}{(a'-c)(a'-c')},$$

$$6. \frac{(b-c)(b-c')}{(b'-c)(b'-c')} = \frac{(b-a)(b-a')}{(b'-a)(b'-a')},$$

$$7. \frac{(c-a)(c-a')}{(c'-a)(c'-a')} = \frac{(c-b)(c-b')}{(c'-b)(c'-b')}.$$

8. Shew that the seven preceding relations may be derived from the single relation

$$(a+a')(bb'-cc') + (b+b')(cc'-aa') + (c+c')(aa'-bb') = 0.$$


---

## CHAPTER V.

## SIMPLE EQUATIONS OF ONE UNKNOWN QUANTITY.

Art. XXXIX. **Preliminary Equations.** Although the following exercise belongs in theory to this chapter, in practice the *numerical* examples should immediately follow Exercise I., and the *literal* examples Exercise III. Like those exercises, this one is merely a specimen of what the teacher should give till his pupils have thoroughly mastered this preliminary work. But few numerical examples are given, it being left to the teacher to supply these.

## Exercise xlix.

What values must  $x$  have that the following equations may be true?

1.  $x-5=0$ .  $x-3\frac{1}{2}=0$ .  $x-a=0$ .  $x+3=0$ .
2.  $x+4\frac{1}{2}=0$ .  $x+a=0$ .  $x+3=5$ .  $x-4=6$ .
3.  $x-a=b$ .  $x+a=c$ .  $x-b=-c$ .  $6-x=3$ .
4.  $8-x=10$ .  $5+x=11$ .  $9+x=4$ .  $7-x=-5$ .
5.  $8+x=-6$ .  $a-x=3b$ .  $2a=x+3b$ .  $3a=5b-x$ .
6.  $2x-6=8$ .  $3x+8=20$ .  $ax=a^2$ .  $mx=bm$ .
7.  $3x=c$ .  $ax=5$ .  $ax=0$ .  $(a+b)x=b+a$ .
8.  $(a-b)x=b-a$ .  $(a+bx)=(a+b)^2$ .  $(a-b)x=a^2-b^2$ .
9.  $(a+b)x=b^2-a^2$ .  $(a^2-ab+b^2)x=a^3+b^3$ .
10.  $(a^2-b^2)x=a-b$ .  $(a^2-b^2)x=a+b$ .  $(a^2+b^2)x=1$ .
11.  $(a+x-b)=(a+b)$ .  $x-a+b=b-x+a$ .
12.  $2a-x=x-2b$ .  $ax+bx=c$ .  $ax-b=cx$ .
13.  $ax-b=bx-c$ .  $ax-ab=ac$ .
14.  $ax-a^2=bx-b^2$ .  $ax-a^3=bx-b^3$ .



15.  $ax - a^3 = b^3 - bx$ ;  $ax + b + c = a + bx + cx$ .
16.  $a - bx - c = b - ax + cx$ ;  $a + bx + cx^2 = ax - b + cx^3$ .
17.  $bx - cx^2 + e = ex - b - cx^3$ ;  $3x = \frac{2}{3}$ ;  $4x = \frac{2}{7}$ .
18.  $10x = \frac{1}{6} - 1$ ;  $ax = \frac{b}{c}$ ;  $ax = \frac{a^2}{b}$ .
19.  $abx = \frac{a}{b} + \frac{b}{a}$ ;  $bcx = \frac{ac^2}{b} + \frac{ab^2}{c}$ .
20.  $\frac{1}{2}x = 5$ ;  $\frac{2}{3}x = 8$ ;  $\cdot 5x = 2$ ;  $\cdot 3x = \cdot 06$ .
21.  $\cdot 02x = 20$ ;  $\cdot 3x = \cdot 2$ ;  $\cdot 4x = \cdot 6$ .
22.  $\cdot 18x = 1\cdot 8$ ;  $\frac{x}{a} = b$ ;  $\frac{ax}{b} = c$ .
23.  $\frac{ax}{b} = \frac{b}{c}$ ;  $\frac{x}{a+b} = c$ ;  $\frac{ax}{a+b} = b$ .
24.  $\frac{a+b}{a-b}x = \frac{a}{b}$ ;  $\frac{a-b}{a+b}x = \frac{a+b}{b-a}$ .
25.  $\frac{a}{b-a}x = \frac{a}{a-b}$ ;  $\frac{b-a}{a+b}x = \frac{a-b}{b+a}$ .
26.  $\frac{a+b}{a+c}x = \frac{a-c}{a+b}$ ;  $\frac{1}{x} = \frac{1}{2}$ ;  $\frac{2}{x} = \frac{3}{5}$ .
27.  $\frac{1}{x} = \frac{1}{ab}$ ;  $\frac{1}{x} = \frac{a}{b}$ ;  $\frac{a}{x} = \frac{b}{c}$ ;  $\frac{7}{x} = \frac{1}{3} + \frac{1}{4}$ .
28.  $\frac{3}{20} + \frac{4}{5x} = \frac{33}{5x} - \frac{1}{3}$ ;  $\frac{a}{x} + \frac{b}{c} = 0$ .
29.  $\frac{5}{x-7} = 6 - \frac{7}{x-7}$ ;  $\frac{5}{3x-4} = 7 + \frac{9}{4-3x}$ .
30.  $(x-4) - (x+5) + x = 3$ ;  $2x - (x-5) - (4-3x) = 5$ .
31.  $2(3-x) + 3(x-3) = 0$ ;  $2(3x-4) - 3(3-4x) + 9(2-x) = 10$ .
32.  $a(1-2x) - (2x-a) = 1$ ;  $x-5(a-x) = bx-5a$ .
33.  $mx(3a-4) + 3mx - 3a + 1 = 0$ .
34.  $a(bx-c) + b(cx-a) + c(ax-b) = 0$ .
35.  $a(ax-b) + b(cx-c) + c(cx-a) = 0$ .

36.  $a(bx - a) + b(cx - b) + c(ax - c) = 0.$   
 37.  $a(x - 2b) + b(x - 2c) + c(x - 2a) = a^2 + b^2 + c^2.$   
 38.  $3\{3\{3(3x - 2) - 2\} - 2\} - 2 = 1.$   
 39.  $9\{7\{5(3x - 2) - 4\} - 6\} - 8 = 1.$   
 40.  $\frac{1}{3}\{\frac{1}{3}\{\frac{1}{3}\{\frac{1}{3}(x + 2) + 2\} + 2\} + 2\} + 2 = 1.$   
 41.  $\frac{1}{9}\{\frac{1}{7}\{\frac{1}{5}\{\frac{1}{3}\{x + 2\} + 4\} + 6\} + 8\} = 1.$   
 42.  $\frac{1}{2}\{\frac{1}{2}\{\frac{1}{2}\{\frac{1}{2}x - \frac{1}{2}\} - \frac{1}{2}\} - \frac{1}{2}\} - \frac{1}{2} = 0.$   
 43.  $\frac{2}{3}\{\frac{2}{3}\{\frac{2}{3}\{3x - 1\frac{1}{3}\} - 1\frac{1}{3}\} - 1\frac{1}{3}\} - 1\frac{1}{3} = 0.$   
 44.  $\frac{1}{8}\{\frac{7}{11}\{\frac{5}{4}\{\frac{3}{2}\{x + 4\} + 8\} + 12\} + 20\} + 32 = 58.$   
 45.  $\frac{4}{5}\{\frac{3}{4}\{\frac{2}{3}\{\frac{1}{2}\{x + 7\} - 3\} + 6\} - 1\} = 4.$   
 46.  $r\{q\{p\{n(mx - a) - b\} - c\} - d\} - e = 0.$   
 47.  $(1 + 6x)^2 + (2 + 8x)^2 = (1 + 10x)^2.$   
 48.  $9(2x - 7)^2 + (4x - 27)^2 = 13(4x + 15)(x + 6).$   
 49.  $(3 - 4x)^2 + (4 - 4x)^2 = 2(5 + 4x)^2.$   
 50.  $(9 - 4x)(9 - 5x) + 4(5 - x)(5 - 4x) = 36(2 - x)^2.$

**Art. XL.** In order that the product of two or more factors may vanish, it is necessary, and it is sufficient, that one of the factors should vanish. Thus, in order that  $(x - a)(x - b)$  may = 0, either  $x - a$  must = 0, or  $x - b$  must = 0, and it is sufficient that one of them should do so.

Hence the single equation  $(x - a)(x - b) = 0$  is really equivalent to the two disjunctive equations, either  $x - a = 0$  or  $x - b = 0$ , for either of these will fulfil the condition of the given equation, and that is all that is required.

Similarly, were it required to find what values of  $x$  would make the product  $(x - a)(x - b)(x - c)$  vanish, they would be given by

$$x - a = 0, \text{ or } x - b = 0, \text{ or } x - c = 0 \therefore x = a \text{ or } b \text{ or } c.$$

Hence the single equation

$$(x - a)(x - b)(x - c) = 0$$

is equivalent to the three disjunctive equations

$$x - a = 0, \text{ or } x - b = 0, \text{ or } x - c = 0.$$

## EXAMPLES.

1. Solve  $x^2 - x - 20 = 0$ .

The expression  $= (x-5)(x+4)$ , which will vanish if either of its factors does, that is, if  $x-5=0$ , or  $x+4=0$ ,

$$\therefore x=5, \quad \text{or } x=-4.$$

2. Solve  $x^4 - x^3 - x^2 + x = 0$ .

This gives  $x^3(x-1) - x(x-1) = x(x-1)(x^2-1)$

$= x(x-1)(x+1)(x-1)$ , which vanishes for

$$x=0, x=1, x=-1.$$

3. Solve  $x^3 + a^2x^2 - ax - a^3 = 0$ .

This  $= x(x^2 - a) + a^2(x^2 - a)$

$= (x+a^2)(x^2 - a)$ , which vanishes for

$$x+a^2=0, \text{ and } x^2 - a=0, \text{ or}$$

$$x = -a^2, \text{ and } x^2 = a.$$

4. Solve  $x^2(a-b) + a^2(b-x) + b^2(x-a) = 0$ .

The factors of the expression are (Ex. 2, page 79)

$x-a, x-b, a-b$ ; hence the expression vanishes if  $x-a=0$ , or  $x-b=0$ .

5. Solve  $221x^2 - 5x - 6 = 0$ .

Here we have the factors  $17x-3$  and  $13x+2$ ;

$\therefore$  the equation is satisfied by  $17x-3=0$ , or  $x = \frac{3}{17}$ ,  
and  $13x+2=0$ , or  $x = -\frac{2}{13}$ .

6. Solve  $2x^4 + 2x^3 + 6x - 18 = 0$ .

In this case we have  $2(x^4 - 9) + 2x(x^2 + 3)$

$= 2(x^2 + 3)\{x^2 - 3 + x\}$ , which vanishes for  $x^2 + 3 = 0$ , or  $x^2 + x - 3 = 0$ .

7. Solve  $(x-a)^3 + (a-b)^3 + (b-x)^3 = 0$ .

The expression is equal to  $3(x-a)(a-b)(b-x)$ ,

and therefore vanishes for  $x-a=0$ , or  $x=a$ ;

and for  $x-b=0$ , or  $x=b$ .

## Exercise 1.

1. If an equation in  $x$  has the factors  $2x-4$  and  $2x-6$ , find the corresponding values of  $x$ .

2. If an equation gives the factors  $2x-1$  and  $3x-1$ , what are the corresponding values of  $x$ ?

3. If an equation gives the factors  $3x^2-12$  and  $4x-5$ , find the corresponding values of  $x$ .

Find the values of  $x$  for which the following expressions will vanish;

4.  $x^2-2x+1$ ;  $4x^2-12x+9$ .

5.  $9x^2-4$ ;  $x^2-(a+b)^2$ ;  $x^2-2ax+a^2$ .

6.  $x^2-9x+20$ ;  $4x^2-18x+20$ .

7.  $x^2+x-6$ ;  $x^2-x-12$ ;  $9x^2-9x-28$ .

8.  $6x^2-12x+6$ ;  $6x^2-13x+6$ ;  $6x^2-20x+6$ .

9.  $6x^2-5x-6$ ;  $6x^2-37x+6$ ;  $6x^2+x-12$ .

10. A certain equation of the fourth degree gives the factors  $x^2-x-2$ , and  $4x^2-2x-2$ , find all the values of  $x$ .

Find values of  $x$  in the following cases :

11.  $x^3-2bx^2-3b^2x=0$ .

12.  $x^3-ax^2+a^2x-a^3=0$ .

13.  $x^3-2x+1=0$ ;  $x^3-3x+2=0$ .

14.  $x^4-2ax^3+2a^3x-a^4=0$ .

15.  $x^3+(b+c)x^2-bcx-b^2c-bc^2=0$ .

16.  $\frac{x-a}{x-b} + \frac{x-b}{x-a} - \frac{(a-b)^2}{(x-a)(x-b)} = \frac{x^2-a^2}{(x-a)(x-b)}$ .

17.  $x^3-bx^2-a^2x-a^2b=0$ .

18.  $3x^3+4abx^2-6a^2b^2x-4a^3b^3=0$ .

19.  $x^3(a-b)+a^3(b-x)+b^3(x-a)=0$ .

20.  $\frac{(x-b)(x-c)}{(a-b)(a-c)} + \frac{(x-c)(x-a)}{(b-c)(b-a)} = 1$ .

21.  $x \left( \frac{x-2a}{x+a} \right)^3 + a \left( \frac{2x-a}{x+a} \right)^3 = x^2-a^2$ .

22.

23.

24.

-6, or

25.

3a, or

26.

27.

and 1-

Art.

illustra

with ad

When

or more

1. So

Here,

denomin

at once

7

6

2. 2a

In th

numeric

8

2

1

:

$$22. (x+a+b)^3 - x^3 - a^3 - b^3 = (x+a)(a^2 - b^2).$$

$$23. \frac{ab}{(b-a)(x-a)} + \frac{bx}{(x-a)(a-b)} + \frac{ax}{(a-b)(b-x)} = \frac{1}{a-b}.$$

24. Form the polynome which will vanish for  $x$  equal 5, or -6, or 7.

25. Form the polynome which will vanish for  $x=a$ , or  $4a$ , or  $3a$ , or  $-4a$ .

26. Form the equation whose roots are 0, 1, -2, and 4.

27. Form the equation whose roots are  $1+\sqrt{2}$ ,  $1-\sqrt{2}$ ,  $1-\sqrt{3}$ , and  $1+\sqrt{3}$ .

**Art. XLI.** In solving fractional equations, the principles illustrated in the section on fractions may frequently be applied with advantage, as in the following cases.

When an equation involves several fractions, we may take two or more of them together.

#### EXAMPLES.

$$1. \text{ Solve } \frac{8x+5}{14} + \frac{7x-3}{6x+2} = \frac{4x+6}{7}.$$

Here, instead of multiplying through by the L. C. M. of the denominators, we combine the first fraction with the last, getting at once

$$\frac{7x-3}{6x+2} = \frac{7}{14} = \frac{1}{2}. \quad \therefore 7x-3=3x+1, \text{ and } x=1.$$

$$2. \frac{2x+8\frac{1}{2}}{9} - \frac{13x-2}{17x-32} + \frac{x}{3} = \frac{7x}{12} - \frac{x+16}{36}.$$

In this case, taking together all the fractions having only numerical denominators, we get

$$\frac{8x+34+12x-21x+x+16}{36} = \frac{13x-2}{17x-32}; \text{ or}$$

$$\frac{25}{18} = \frac{13x-2}{17x-32};$$

$$\therefore 425x-800=234x-36, \text{ hence } x=4.$$

It is often advantageous to *complete the divisions* represented by the fractions.

$$3. \frac{4x-17}{9} - \frac{3\frac{2}{3}-22x}{33} = x - \frac{6}{x} \left(1 - \frac{x^2}{54}\right).$$

Here, completing the divisions, we have

$$\frac{4x}{9} - \frac{17}{9} - \frac{1}{9} + \frac{2x}{3} = x - \frac{6}{x} + \frac{x}{9},$$

$$\frac{10x}{9} - 2 = x + \frac{x}{9} - \frac{6}{x} \therefore -2 = -\frac{6}{x}, \text{ or } x=3.$$

$$4. \frac{ax+b}{x-m} + \frac{cx+d}{x-n} = a+c$$

$$\therefore a + \frac{am+b}{x-m} + c + \frac{cn+d}{x-n} = a+c$$

$$\therefore (am+b)(x-n) + (cn+d)(x-m) = 0$$

$$\therefore (am+b+cn+d)x = (a+c)mn + bn + dm.$$

5. Similarly may be solved

$$\frac{ax+b}{x-m} + \frac{cx+d}{x-n} + \frac{ex^2+fx-g}{(x-m)(x-n)} = a+c+e$$

$$\therefore \frac{am+b}{x-m} + \frac{cn+d}{x-n} + \frac{\{e(m+n)+f\}x - emn - g}{(x-m)(x-n)} = 0.$$

$$\therefore (am+b)(x-n) + (cn+d)(x-m) + \{e(m+n)+f\}x - emn - g = 0.$$

$$\therefore \{(a+c)m+b+(c+e)n+d+f\}x = (a+b+c)mn + bn + dm + g.$$

$$6. \frac{132x+1}{3x+1} + \frac{8x+5}{x-1} = 52;$$

$$\therefore 44 - \frac{43}{3x+1} + 8 + \frac{13}{x-1} = 52, \text{ or}$$

$$\frac{13}{x-1} = \frac{43}{3x+1}; \therefore 39x+13=43x-43, \text{ and } x=14.$$

$$7. \frac{25-\frac{1}{3}x}{x+1} + \frac{16x+4\frac{1}{5}}{3x+2} = 5 + \frac{23}{x+1}.$$

Taking the last fraction with the first, and multiplying the resulting equation by 15, we have

$$\frac{240x+63}{3x+2} = 75 + \frac{5x-30}{x+1};$$

$$\therefore 80 - \frac{97}{3x+2} = 75 + 5 - \frac{35}{x+1}, \text{ or}$$

$$\frac{97}{3x+2} = \frac{35}{x+1}; \therefore 8x = 27, \text{ and } x = 3\frac{3}{8}.$$

$$8. \frac{x-a}{b+c} + \frac{x-b}{a+c} + \frac{x-c}{b+c} = 3.$$

$$\therefore \frac{x-a}{b+c} - 1 + \frac{x-b}{a+c} - 1 + \frac{x-c}{b+c} - 1 = 0;$$

$$\therefore \frac{x-(a+b+c)}{b+c} + \frac{x-(a+b+c)}{a+c} + \frac{x-(a+b+c)}{b+c} = 0,$$

which is satisfied by  $x-(a+b+c)=0$ ;  $\therefore x=a+b+c$ .

$$9. \frac{m}{x+a} + \frac{n}{x-b} = \frac{m+n}{x-c};$$

$$\therefore \frac{m(x-c)}{x-a} + \frac{n(x-c)}{x-b} = m+n,$$

which may be solved as in Ex. 1.

$$10. \frac{5x+5}{x+1} - \frac{4x+9}{2x+4} = \frac{15x+7}{3x+1} - \frac{12x+17}{3x+4}.$$

$$\therefore 3 + \frac{2}{x+1} - 2 - \frac{1}{2x+4} = 5 + \frac{2}{3x+1} - 4 - \frac{1}{3x+4}, \text{ or}$$

$$\frac{2}{x+1} - \frac{1}{2x+4} = \frac{2}{3x+1} - \frac{1}{3x+4};$$

$$\therefore \frac{3x+7}{2x^2+6x+4} = \frac{3x+7}{9x^2+15x+4}.$$

This can be divided by  $3x+7$ , giving  $3x+7=0$ , or  $x=-\frac{7}{3}$ .

The result of the division is

$$\frac{1}{2x^2+6x+4} = \frac{1}{9x^2+15x+4}, \text{ or}$$

$9x^2+15x+4=2x^2+6x+4$ , or  $7x^2=-9x$ , which we can divide by  $x$ ,  $\therefore x=0$ ; the result of the division is  $7x=-9$ , or  $x=-\frac{9}{7}$ .

## Exercise II.

$$1. \frac{10x+17}{18} - \frac{12x+2}{13x-16} = \frac{5x-4}{9}.$$

$$2. \frac{6x+13}{15} - \frac{9x+15}{5x-25} + 3 = \frac{2x+15}{5}.$$

$$3. \frac{7x+1}{x-1} = \frac{35}{9} \times \frac{x+4}{x+2} + 3\frac{1}{9}.$$

$$4. \frac{4x-7}{2x-9} + \frac{2-14x}{7} + \frac{3\frac{1}{3}+x}{14} = \frac{10-3\frac{2}{3}x}{2} - \frac{19}{21}.$$

$$5. \frac{2x+a}{3(x-a)} + \frac{3x-a}{2(x+a)} = 2\frac{1}{6}.$$

$$6. \frac{x-4}{6x+5} + \frac{3x-13}{18x-6} = \frac{1}{3}.$$

$$7. \frac{3x+1}{2x-15} - \frac{x-11}{2x-10} = 1; \quad \frac{x-9}{x-5} + \frac{x-5}{x-8} = 2.$$

$$8. \frac{x-12}{x-7} + \frac{x-4}{x-12} = 2 + \frac{7}{x-7}; \quad \frac{3x-19}{x-13} + \frac{3x-11}{x+7} = 6.$$

$$9. \frac{x-2}{2x+1} + \frac{x-1}{3(x-3)} = \frac{5}{6}; \quad \frac{x+1}{4(x+2)} + \frac{x+4}{5x+13} = \frac{9}{20}.$$

$$10. \frac{5(2x^2+3)}{2x+1} + \frac{5-7x}{2x-5} = 5x-6; \quad \frac{3}{x-7} + \frac{1}{x-9} = \frac{4}{x-8}.$$

$$11. \frac{7x+55}{2x+5} - \frac{3x}{2} = 9 - \frac{3x^2+8}{2x-4}; \quad \frac{17}{x-16} + \frac{15}{x-18} = \frac{32}{x-17}.$$

$$12. \frac{1-25x}{15} - \frac{3-2\frac{1}{2}x}{14(x-1)} = \frac{28-5x}{3} - \frac{10x-11}{30} + \frac{x}{3}.$$

$$13. \frac{1}{x-2} - \frac{2+2\frac{1}{2}x^2-\frac{1}{2}x^3}{6-5x+x^2} - \frac{1}{2}x = \frac{5}{x-5}.$$

$$14. \frac{30+6x}{x+1} + \frac{60+8x}{x+3} = \frac{48}{x+1} + 14.$$

$$15. \frac{5x^2+x-3}{5x-4} = \frac{7x^2-3x-9}{7x-10}.$$

$$16. \frac{x}{x-2} + \frac{x-9}{x-7} = \frac{x+1}{x-1} + \frac{x-8}{x-6}.$$

17.

18.

19.

20.

21.

22.

23.

24.

25.

26.

27.

28.

29.

30.

31.



17.  $\frac{x^2-3x-9}{x-5} + \frac{x^2-7x-17}{x-9} = \frac{x^2-6x-15}{x-8} \cdot 2$
18.  $\frac{4x+7}{4x+5} + \frac{4x+9}{4x+7} = \frac{4x+6}{4x+4} + \frac{4x+10}{4x+8}$ .
19.  $\frac{2x-3}{2x-4} - \frac{2x-4}{2x-5} = \frac{2x-7}{2x-8} - \frac{2x-8}{2x-9}$ .
20.  $\frac{7x+6}{28} - \frac{2x+4\frac{2}{3}}{23x-6} + \frac{x}{4} = \frac{11x}{21} - \frac{x-3}{42}$ .
21.  $\frac{x^2-5}{x^2-6} + \frac{x^2-11}{x^2-12} = \frac{x^2-7}{x^2-8} + \frac{x^2-9}{x^2-10}$ .
22.  $\frac{x-1\frac{2}{3}\frac{5}{6}}{2} - \frac{2-6x}{13} = x - \frac{5x-\frac{1}{4}(10-3x)}{39}$ .
23.  $\frac{1-2x}{3(x^2-x+1)} + \frac{1+x}{2(x^2+1)} + \frac{1}{6(x+1)} = \frac{1}{9(x^2+1)}$ .
24.  $\frac{2x^2+x-30}{2x-7} + \frac{x^2+4x-4}{x-1} = \frac{x^2-17}{x-4} + \frac{2x^2+7x-13}{2x-3}$ .
25.  $\frac{x-a}{x-b} + \frac{x-b}{x-a} - \frac{(a-b)^2}{(x-a)(x-b)} = \frac{2(a-x)}{a+x}$ .
26.  $\frac{12x+10a}{3x+a} + \frac{28x+117a}{2x+9a} = 18$ .
27.  $\frac{13\frac{1}{2}x-5}{13\frac{1}{2}x-6} + \frac{13\frac{1}{2}x-11}{13\frac{1}{2}x-12} = \frac{13\frac{1}{2}x-7}{13\frac{1}{2}x-8} + \frac{13\frac{1}{2}x-9}{13\frac{1}{2}x-10}$ .
28.  $\frac{1}{2(x-1)^2} + \frac{1}{2(x-1)} - \frac{x}{2(x^2+1)} = \frac{16x}{(x-1)(x^2+1)}$ .
29.  $\frac{1}{2}(\frac{2}{3}x+4) - \frac{7\frac{1}{2}-x}{3} = \frac{x}{2} \left( \frac{6}{x} - 1 \right)$ .
30.  $\frac{3x}{2} - \frac{81x^2-9}{(3x-1)(x+3)} = 3x - \frac{3}{2} \cdot \frac{2x^2-1}{x+3} - \frac{57-3x}{2}$ .
31.  $1 + \frac{2x+1}{2(x-1)} - \frac{4x+5}{2(x+1)} = \frac{x^2+3x+2}{x^2-2x+1} - 1$ .

$$32. \frac{7x-30}{10\frac{1}{2}} - \frac{5x-7}{\frac{1}{2}x-3} - \frac{2-21x}{21} =$$

$$\frac{42x-171}{63} - 10 + \frac{2x-9}{63-14x} - \frac{1}{7}(4 -$$

$$33. \frac{18x-22}{13-2x} + 6x + \frac{1+6x}{8} = 13\frac{1}{4} - \frac{101-64x}{8}$$

$$34. \frac{4-9x}{1-3x} - \frac{5-12x}{7-4x} = 2 - \frac{24x^2-5}{7-25x+12x^2}$$

$$35. \frac{8x+25}{2x+5} + \frac{16x+93}{2x+11} = \frac{18x+86}{2x+9} + \frac{6x+26}{2x+7}$$

$$36. \frac{1}{x+a+b} + \frac{1}{x-a+b} + \frac{1}{x+a-b} + \frac{1}{x-a-b} = 0.$$

Art. XLII. The results deduced in Section III., Chapter IV., may often be applied with advantage.

#### EXAMPLES.

$$1. \frac{ax+b}{cx+d} = \frac{m}{n}$$

$$\therefore \frac{(ax+b)d - (cx+d)b}{(cx+d)a - (ax+b)c} = \frac{md-nb}{na-mc} \quad (\text{page 123}).$$

$$\therefore x = \frac{md-nb}{na-mc}$$

$$2. \frac{ax^2+bx+c}{mx^2+nx+p} = \frac{a}{m}$$

$$\therefore \frac{(ax^2+bx+c) - ax^2}{(mx^2+nx+p) - mx^2} = \frac{a}{m} \quad (\text{page 122}).$$

$$\therefore \frac{bx+c}{nx+p} = \frac{a}{m} \quad \&c.$$

$$3. \frac{3x+7}{x+4} = \frac{3x-13}{x-4}$$

By (5) each of these fractions =

$$\frac{\text{difference of numerators}}{\text{difference of denominators}}, \quad \therefore \frac{20}{8} = \frac{3x+7}{x+4} = 3 - \frac{5}{x+4},$$

$$\text{or } \frac{1}{2} = \frac{5}{x+4}, \quad \therefore x=6.$$

$$4. \quad \frac{mx+a+b}{nx-c-d} = \frac{mx+a+c}{nx-b-d},$$

$$\therefore \frac{mx+a+b}{mx+a+c} = \frac{nx-c-d}{nx-b-d}; \text{ or By 4, page 122,}$$

$$\frac{mx+a+b}{b-c} = \frac{nx-c-d}{b-c}; \text{ or } (n-m)x$$

$$= a+b+c+d, \quad \therefore x = \&c.$$

$$5. \quad \frac{\sqrt{(a+x)} + \sqrt{(a-x)}}{\sqrt{(a+x)} - \sqrt{(a-x)}} = a.$$

Here by (6), page 122, we have

$$\frac{2\sqrt{(a+x)}}{2\sqrt{(a-x)}} = \frac{a+1}{a-1}; \text{ or, cancelling the 2 in left hand mem-}$$

ber, and squaring,

$$\frac{a+x}{a-x} = \frac{(a+1)^2}{(a-1)^2}, \text{ whence, again by (6),}$$

$$\frac{2x}{2a} = \frac{(a+1)^2 - (a-1)^2}{(a+1)^2 + (a-1)^2} = \frac{4a}{2(a^2+1)} = \frac{2a}{a^2+1};$$

$$\therefore x = \frac{2a^2}{a^2+1}.$$

$$6. \quad \frac{\sqrt{(x-a+b)} - \sqrt{(x+a-b)}}{\sqrt{(x-a+b)} + \sqrt{(x+a-b)}} = \frac{a-b}{a+b}.$$

$$\therefore \frac{\sqrt{(x-a+b)}}{\sqrt{(x+a-b)}} = \frac{a}{b};$$

squaring and again applying (6),

$$\therefore -\frac{2x}{2(a-b)} = \frac{a^2+b^2}{a^2-b^2}, \text{ and } x = -\frac{a^2+b^2}{a+b}.$$

## Exercise lii.

1.  $\frac{1+x}{1-x} = \frac{1}{a}$ ;  $\frac{x+a}{x-a} = m$ ;  $\frac{ax+b}{ax-b} = \frac{m}{n}$ .
2.  $\frac{a+x}{b+2x} = 1$ ;  $\frac{a(b+x)}{a-x} = b$ ;  $\frac{a}{a-x} = \frac{b}{b-x}$ .
3.  $\frac{a+x}{a-x} = \frac{a+b}{a-b}$ ;  $\frac{x+m}{x-m} = \frac{a+b}{a-b}$ ;  $\frac{a+b}{1+cx} = \frac{a-b}{1-cx}$ .
4.  $\frac{a+bx}{a+b} = \frac{c+dx}{c+d}$ ;  $\frac{a+bx}{a-b} = \frac{c+dx}{c-d}$ ;  $\frac{a-x}{b-x} = \frac{a+x}{b+x}$ .
5.  $\frac{2x^2-5x+6}{2x^2-7x+3} = \frac{x^2-7x+5}{x^2-9x+2}$ .
6.  $\frac{ax+b-c}{ax-b+c} = \frac{(b-c)^2}{(b+c)^2}$ .
7. If  $\sqrt{x+y} + \sqrt{x-y} = \frac{x}{y}$ , shew that  $\frac{x+y}{x-y} = 1$ .
8.  $\frac{2x-7}{2x-3} = \frac{x+7}{x+11}$ ;  $\frac{4x-5}{2x+10} = \frac{10x-32}{5x-8\frac{1}{2}}$ .
9.  $\frac{57x-43}{19x+13} = \frac{39x-7}{13x+25}$ ;  $\frac{23x+5\frac{1}{2}}{115x-29} = \frac{36x-7}{180x+23}$ .
10.  $\frac{210x-73}{310x-66} = \frac{21x+7\cdot3}{31x+8}$ ;  $\frac{mx-a-b}{nx-c-d} = \frac{mx-a-c}{nx-b-d}$ .
11.  $\frac{3x + \sqrt{4x-x^2}}{3x - \sqrt{4x-x^2}} = 2$ ;  $\frac{\sqrt{12x+1} + \sqrt{12x}}{\sqrt{12x+1} - \sqrt{12x}} = 18$ .
12.  $\frac{x^3+ax^2-bx+c}{x^3-ax^2+bx+c} = \frac{x^2+ax-b}{x^2-ax+b}$ .
13.  $\frac{\sqrt{2a^2-x^2} + b\sqrt{2a-x}}{\sqrt{2a^2-x^2} - b\sqrt{2a-x}} = \frac{\sqrt{a+b}}{\sqrt{a-b}}$ .
14.  $\frac{\sqrt{x^2+a^2} + \sqrt{x^2-a^2}}{\sqrt{x^2+a^2} - \sqrt{x^2-a^2}} = a^2$ .
15.  $\frac{8x^3+12x^2-8x+5}{8x^3-12x^2+8x+5} = \frac{4x^2+6x-4}{4x^2-6x+4}$ .

16.

17.

18.

18.

20.

21.

22.

23.

24.

25.

26.

27.

28.

29.

30.

31.

Art. 2  
simplify

16.  $\frac{\sqrt[3]{x+1} - \sqrt[3]{x-1}}{\sqrt[3]{x+1} + \sqrt[3]{x-1}} = \frac{1}{2}$
17.  $\frac{28 + \sqrt{x}}{28 - \sqrt{x}} = \frac{9 + 3\sqrt{x}}{9 + 2\sqrt{x}}$
18.  $\frac{a^3x^3 + a^2bx^2 - acx + d}{a^3x^3 - a^2bx^2 + acx + d} = \frac{a^2x^2 + abx - c}{a^2x^2 - abx + c}$
18.  $\frac{5\sqrt{2x-1} + 2\sqrt{3x-3}}{4\sqrt{2x-1} - 2\sqrt{3x-3}} = 2\frac{1}{3}$
20.  $\frac{\sqrt{2x} + \sqrt{3-2x}}{\sqrt{2x} - \sqrt{3-2x}} = \frac{3}{2}$
21.  $\frac{2\sqrt[3]{3x+3} + \sqrt[3]{7x+8}}{2\sqrt[3]{3x+3} - \sqrt[3]{7x+8}} = 5$
22.  $33\{13 - 2\sqrt{x-5}\} = 3\{13 + 2\sqrt{x-5}\}$
23.  $(\sqrt{n+1})\{\sqrt{nx+1} - \sqrt{nx}\} = (\sqrt{n-1})\{\sqrt{nx+1} + \sqrt{nx}\}$
24.  $\frac{\sqrt{x+c} + \sqrt{b}}{\sqrt{x+c} - \sqrt{b}} = \frac{\sqrt{x} + \sqrt{a}}{\sqrt{x} - \sqrt{a}}$
25.  $\frac{\sqrt{x+28}}{\sqrt{x+4}} = \frac{\sqrt{x+38}}{\sqrt{x+6}}; \quad \frac{\sqrt[3]{2x+17}}{\sqrt[3]{2x+9}} = \frac{\sqrt[3]{2x+27}}{\sqrt[3]{2x+15}}$
26.  $\frac{\sqrt{x+2a}}{\sqrt{x+b}} = \frac{\sqrt{x+4a}}{\sqrt{x+3b}}; \quad \frac{3x-1}{\sqrt{3x+1}} = \frac{1 + \sqrt{3x}}{2}$
27.  $\frac{\sqrt{a} - \sqrt{a-x}}{\sqrt{a} + \sqrt{a-x}} = a; \quad \frac{\sqrt{x} + \sqrt{b}}{\sqrt{x} - \sqrt{b}} = \frac{a}{b}$
28.  $\frac{ax+1 + \sqrt{a^2x^2-1}}{ax+1 - \sqrt{a^2x^2-1}} = b$
29.  $\frac{1 - \sqrt{1 - \sqrt{1-x}}}{1 + \sqrt{1 - \sqrt{1-x}}} = a$
30.  $\frac{1+x}{\sqrt{(2ax+x^2)}} = \frac{b+1}{b-1}; \quad \frac{1+x+x^2}{1-x+x^2} = \frac{62}{63} \times \frac{1+x}{1-x}$
31.  $\frac{5x^4 + 10x^2 + 1}{x^5 + 10x^3 + 5x} = \frac{a^5 + 10a^3 + 1}{5a^4 + 10a^2 + 1}$

Art. XLIII. Various other artifices may be employed to simplify the solution of equations.

## EXAMPLES.

1. Solve  $2 + \sqrt{4x^2 - 9x + 8} - 2x = 0$ : here there is but one surd, and it is convenient to make that surd one side of the equation and transpose all the rational terms to the other; this gives

$$\sqrt{4x^2 - 9x + 8} = 2x - 2; \text{ squaring both sides,}$$

$$4x^2 - 9x + 8 = 4x^2 - 8x + 4, \therefore x = 4.$$

2.  $\sqrt{a+x} + \sqrt{a-x} = 2\sqrt{x}$ . We might square this as it stands, but the work will be simplified if we first transpose, thus

$$\sqrt{a+x} = 2\sqrt{x} - \sqrt{a-x}; \text{ now squaring,}$$

$$a+x = 4x + a - x - 4\sqrt{(ax-x^2)}, \text{ or}$$

$$x = 2\sqrt{(ax-x^2)}. \text{ Again squaring,}$$

$$x^2 = 4ax - 4x^2, \text{ whence } x = 0, \text{ or } \frac{4a}{5}.$$

3. Clear of radicals

$$\sqrt[3]{x} + \sqrt[3]{y} + \sqrt[3]{z} = 0. \text{ Transposing,}$$

$$\sqrt[3]{x} + \sqrt[3]{y} = -\sqrt[3]{z}; \text{ cube by formula [6],}$$

$$x+y+3\sqrt[3]{xy}(\sqrt[3]{x}+\sqrt[3]{y}) = -z; \text{ and substituting for}$$

$$\sqrt[3]{x} + \sqrt[3]{y} \text{ its value } -\sqrt[3]{z}, \text{ this becomes}$$

$$x+y-3\sqrt[3]{xyz} = -z, \text{ or}$$

$$x+y+z = 3\sqrt[3]{xyz}; \therefore \text{ cubing again,}$$

$$(x+y+z)^3 = 27xyz.$$

$$4. \frac{a+x+\sqrt{(2ax+x^2)}}{a+x} = b.$$

Dividing and transposing, we have

$$\frac{\sqrt{(2ax+x^2)}}{a+x} = b-1, \therefore \frac{2ax+x^2}{a^2+2ax+x^2} = (b-1)^2; \text{ again by}$$

division in left-hand member,

$$-\frac{a^2}{(a+x)^2} + 1 = (b-1) \therefore \frac{a}{a+x} = \sqrt{\{1-(b-1)^2\}}, \text{ or}$$

$$\frac{x+a}{a} = 1 \therefore \sqrt{\{1+(b-1)^2\}}, \text{ or}$$

$$\frac{x}{a} + 1 = \&c.$$

5. Solve  $\sqrt{4x^2+19} + \sqrt{4x^2-19} = \sqrt{47+3}$ .

We have the identity

$$(4x^2+19) - (4x^2-19) = 38 = 47 - 9.$$

Now dividing the members of this identity by those of the given equation, we have

$\sqrt{4x^2+19} - \sqrt{4x^2-19} = \sqrt{47-9}$ . Adding this to the given equation, then

$$2\sqrt{4x^2+19} = 2\sqrt{47}, \therefore 4x^2+19 = 47, \text{ and } x = \pm\sqrt{7}.$$

6.  $\sqrt[3]{25+x} + \sqrt[3]{25-x} = 2$ .

Cubing by formula [6]. (See Ex. 3), we have

$$25+x+25-x+6\sqrt[3]{(25^2-x^2)} = 8, \text{ or}$$

$$\sqrt[3]{(625-x^2)} = -7, \text{ or } (625-x^2) = -343;$$

$$\therefore x^2 - 525 + 453 = 968, \text{ and } x = \pm 22\sqrt{2}.$$

### Exercise liii.

1.  $\sqrt{x+4} + \sqrt{x-3} = 7$ .

2.  $\sqrt{3x+1} + \sqrt{4x+4} = 1$ .

3.  $\sqrt{2x+10} + \sqrt{2x-2} = 6$ .

4.  $\sqrt{mx} - \sqrt{nx} = m - n$ .

5.  $\sqrt{bx} + \sqrt{ab+bx} = \sqrt{x}$ .

6.  $\sqrt{x} + \sqrt{x+3} = \frac{5}{\sqrt{x+3}}$ .

7.  $\sqrt{ax+x^2} = (1+x)$ .

8.  $\sqrt[3]{17x-26} = \frac{2}{9}$ .

9.  $\sqrt{x} - \sqrt{a+x} = \sqrt{\frac{a}{x}}$ .

10.  $b+x - \sqrt{b^2+x^2} = c^2$ .

11.  $\sqrt{8+x} - \sqrt{x} = 2\sqrt{1+x}$ .

12.  $\sqrt{2x-27a} = 9\sqrt{a} - \sqrt{2x}$ .

13.  $\sqrt[3]{1-x} + \sqrt[3]{1+x} = \sqrt[3]{3}$ .
14.  $\sqrt[3]{3+x} + \sqrt[3]{3-x} = \sqrt[3]{7}$ .
15.  $\sqrt[3]{x+1} - \sqrt[3]{x-1} = \sqrt[3]{11}$ .
16.  $\sqrt[3]{a+x} + \sqrt[3]{a-x} = \sqrt[3]{b}$ .
17.  $\sqrt[3]{1+\sqrt{x}} + \sqrt[3]{1-\sqrt{x}} = 2$ .
18.  $\sqrt{x} - \sqrt{a - \sqrt{ax+x^2}} = \frac{1}{2}\sqrt{a}$ .
19. Clear of radicals  $\sqrt[3]{a} + \sqrt[3]{b} - \sqrt[3]{c}$ .
20. Solve  $x + \sqrt{a^2 + x^2} = \frac{na^2}{\sqrt{a^2 + x^2}}$ .
21. Clear of radicals  $\sqrt{x} + \sqrt{y} + \sqrt{x} - \sqrt{y}$ .  
Solve the following equations :
22.  $\sqrt{1+x} + \sqrt{1+x + \sqrt{1-x}} = \sqrt{1-x}$ .
23.  $\sqrt{x + \sqrt{x}} - \sqrt{x - \sqrt{x}} = a\sqrt{\frac{x}{x + \sqrt{x}}}$ .
24.  $\sqrt{1+x+x^2} + \sqrt{1-x+x^2} = mx$ .
25.  $\sqrt{a^2 - x^2} + x\sqrt{a^2 - 1} = a^2(1 - x^2)$ .
26.  $\frac{bx - c^2}{\sqrt{bx} + c} = \frac{\sqrt{bx} + c}{n} - a$ .
27.  $\sqrt{2x^2 + 5} + \sqrt{2x^2 - 5} = \sqrt{15} + \sqrt{5}$ .
28.  $\sqrt{3x^2 + 10} + \sqrt{3x^2 - 10} = \sqrt{17} + \sqrt{3}$ .
29.  $\sqrt{3x^2 + 9} - \sqrt{3x^2 - 9} = \sqrt{34} + 4$ .
30.  $\sqrt{3a - 3b + x^2} + \sqrt{2a - 2b + x^2} = \sqrt{a} + \sqrt{x}$ .
31.  $\sqrt{4a^2 - 3b^2 - 2x^2} + \sqrt{3a^2 - 3b^2 - x^2} = a + x$ .
32. Clear of radicals,  $\sqrt[3]{2x} - \sqrt[3]{2y} - \sqrt[3]{2z}$ .
33.  $\sqrt{a+x} + \sqrt{a-x} = 2x \div \sqrt{a + \sqrt{a^2 + x^2}}$ .
34.  $\sqrt{x^2 + 2ax} + \sqrt{x^2 - 2ax} = \frac{na^2}{\sqrt{x^2 + 2ax}}$ .
35.  $\sqrt{\left(\frac{\sqrt{x+a}}{\sqrt{x-a}}\right)} - \sqrt{\left(\frac{\sqrt{x-a}}{\sqrt{x+a}}\right)} = \sqrt{x-a^2}$ .
36.  $\sqrt{\{(2a+x)^2 + b^2\}} + \sqrt{\{(2a-x)^2 + b^2\}} = 2a$ .

Art  
princip1.  $\frac{x}{a}$ 

Fact

 $(x)$ 

or

2.  $\frac{3}{a}$ 

Tran

 $\frac{ab}{a+b} \left\{ \frac{3}{a} \right\}$ 3.  $\frac{3}{(a)}$ 

Add

 $(a)$  $\therefore (a)$  $\therefore x =$



**Art. XLIV.** Sometimes a factor can be discovered, and the principle of Art. XL. applied.

## EXAMPLES.

$$1. \frac{x^4 + a^2x^2 + a^4}{x-a} = x^3 + (a-b)x^2 + (a^2-ab)x - a^2b.$$

Factoring we have

$$\frac{(x^2 + ax + a^2)(x^2 - ax + a^2)}{x-a} = (x-b)(x^2 + ax + a^2),$$

$$\text{or } x^2 - ax + a^2 = (x-a)(x-b);$$

$$\therefore (a+b-a)x = ab - a^2, \text{ and } x = a - \frac{a^2}{b}.$$

$$2. \frac{3abc}{a+b} - \frac{bx}{a} + \frac{a^2b^2}{(a+b)^3} = 3cx - \frac{b^2x}{a} \cdot \frac{2a+b}{(a+b)^2}.$$

Transpose  $\frac{bx}{a}$  and factor, then

$$\begin{aligned} \frac{ab}{a+b} \left\{ 3c + \frac{ab}{(a+b)^2} \right\} &= x \left\{ 3c + \frac{b}{a} \left( 1 - \frac{2ab+b^2}{(a+b)^2} \right) \right\} \\ &= x \left\{ 3c + \frac{b}{a} \cdot \frac{a^2}{(a+b)^2} \right\} \\ &= x \left\{ 3c + \frac{ab}{(a+b)^2} \right\} \\ \therefore \frac{ab}{a+b} &= x. \end{aligned}$$

$$3. \frac{x+a}{(x-b)(c-a)} - \frac{x-b}{(a-b)(b-c)} - \frac{x-c}{(b-c)(c-a)} = \frac{b+c}{(a-b)(b-c)(c-a)}$$

Add term by term the identity (Th. iii., page 54).

$$\frac{x-a}{(a-b)(c-a)} + \frac{x-b}{(a-b)(b-c)} + \frac{x-c}{(b-c)(c-a)} = 0.$$

$$\therefore \frac{2x}{(a-b)(c-a)} = \frac{b+c}{(a-b)(b-c)(c-a)}.$$

$$\therefore x = \frac{1}{2} \cdot \frac{b+c}{b-c}.$$

$$4. (x+a+b)^3 + (a+b)^3 - (x+b)^3 - (x+a)^3 + x^3 + a^3 + b^3 = abc.$$

The left hand member vanishes for  $x=0$ , and  $\therefore$  by symmetry for  $a=0$  and  $b=0$ ;  $\therefore$  it is of the form  $mabx$  in which  $m$  is numerical.

Put  $x=a=b$ , and  $m$  is found to be 6,

$\therefore$  the equation reduces to

$$6abx = abc, \therefore \text{and } x = \frac{1}{6}c.$$

$$5. \left(\frac{x-a}{x-b}\right)^3 = \frac{x-2a+b}{x-2b+a}; \text{ let } x-b=m, x-a=n, \text{ and } \therefore$$

$m-n=a-b$ , then we have

$$\frac{m^3}{n^3} = \frac{n-(m-n)}{m+(m-n)} = \frac{2n-m}{2m-n}$$

$$\therefore 2m^4 - nm^3 = 2n^4 - n^3m, \text{ and}$$

$$2(m^4 - n^4) - mn(m^2 - n^2) = 0, \text{ which is divisible by } m^2 - n^2,$$

$$\therefore m^2 - n^2 = 0, \text{ or } m+n=0;$$

But  $m+n=2x-a-b=0$ ,  $\therefore x = \frac{1}{2}(a+b)$ .

$$6. \frac{1}{3} \cdot \frac{x^2-4x+2}{x^2-4x-1} + \frac{1}{6} \cdot \frac{x^2-4x+3}{x^2-4x-3} - \frac{2}{9} \cdot \frac{x^2-4x+3}{x^2-4x-6} = \frac{5}{18}$$

Let  $y = x^2 - 4x$ , then this equation becomes

$$\frac{1}{3} \cdot \frac{y+2}{y-1} + \frac{1}{6} \cdot \frac{y+3}{y-3} - \frac{2}{9} \cdot \frac{y+3}{y-6} = \frac{5}{18}, \text{ or by division,}$$

$$\frac{1}{3} + \frac{1}{y-1} + \frac{1}{6} + \frac{1}{y-3} - \frac{2}{9} - \frac{2}{y-6} = \frac{5}{18}, \text{ or}$$

$$\frac{1}{y-1} + \frac{1}{y-3} - \frac{2}{y-6} = 0; \text{ this may be written}$$

$$\frac{1}{y-1} - \frac{1}{y-6} + \frac{1}{y-3} - \frac{1}{y-6} = 0, \text{ or}$$

$$\frac{5}{y-1} + \frac{3}{y-3} = 0, \therefore 5y-15+3y-3=0, \text{ or}$$

$$y = 2\frac{1}{4} \therefore x^2 - 4x = 2\frac{1}{4}, \text{ or } x^2 - 4x + 4 = 4 + 2\frac{1}{4},$$

and  $x-2 = \pm \frac{5}{4}$ . We might assume  $(x-2)^2 = y$ , when the given equation would take the form

$$\frac{1}{3} \cdot \frac{y-2}{y-5} + \frac{1}{6} \cdot \frac{y-1}{y-7} - \frac{2}{9} \cdot \frac{y-1}{y-10} = \frac{5}{18}.$$

And reducing as before, we should find

$$y = 6\frac{1}{4} = (x-2)^2, \therefore x-2 = \pm \frac{5}{2}, \text{ as before.}$$

### Exercise liv.

1.  $\frac{x^4 + a^2x^2 + a^4}{x-b} = x^3 - (a-b)x^2 + (a^2-ab)x + a^2b.$
2.  $\frac{x^4 + 4a^4}{x+b} = x^3 + 2a(a-b)x + (2a-b)x^2 - 2a^2b.$
3.  $\frac{a^2}{a^2+ab+b^2} - \frac{a^2x}{a^3-b^3} = \frac{2c}{a-b} - 2cx.$
4.  $\frac{1}{a+b+x} - \frac{1}{a} - \frac{1}{b} - \frac{1}{x} = 2ab(x+b)x^2.$
5.  $\frac{1}{(x-b)(x-c)} + \frac{1}{(a+c)(a+b)} = \frac{1}{(a+c)(x-c)} + \frac{1}{(a+b)(x-b)}.$
6.  $\frac{bx}{a} - \frac{3ab}{a-b} + \frac{a^2b^2}{(a-b)^3} = 3x - \frac{b^2x}{a} \cdot \frac{2a-b}{(a-b)^2}$
7.  $\frac{x^3 + 2ax}{x^4 - 11x^2a^2 + a^4} = \frac{x-a}{x^2 - 3ax - a^2}.$
8.  $\frac{1}{2} \left( \frac{x-a}{x+a} \right)^2 - \frac{1}{2}(-x+a) = \frac{x-a}{x+a}.$
9.  $x^3 - (a+b+c)x^2 + (a^2+b^2+c^2)x - \frac{1}{3}(a^3+b^3+c^3)$   
 $= (x-a)(x-b)(x-c).$
10.  $\frac{1}{ax} + \frac{1}{bx} + \frac{1}{cx} = \frac{1}{2}(a+b+c)^2 - \frac{1}{2} \left( \frac{a}{bcx} + \frac{b}{acx} + \frac{c}{abx} \right).$
11.  $\frac{1-ax}{bc} + \frac{1-bx}{ac} + \frac{1-cx}{ab} = \left( \frac{2}{a} + \frac{2}{b} + \frac{2}{c} \right) x.$
12.  $\frac{(a-b)^2}{abc} - 1 + \frac{a}{b} = \frac{a^2-b^2}{abc} + \left( 1 + \frac{a}{b} \right) x.$
13.  $x^3 + (b+c)^3 + 3b(b+c)x = b^3.$

$$14. x - a - 3\sqrt{abx} = b.$$

$$15. 11x^4 + 10x^3 - 40x = 176.$$

$$16. \frac{x}{(a+b)^2} + \frac{ac}{(a-b)^3} - \frac{c}{(a^2-b^2)(a+b)} = \frac{ax}{(a-b)^2}.$$

$$17. x^3 - \frac{a-b}{a+b} \cdot x^3 + \frac{2cx^2}{1+cx} = x - \frac{a-b}{a+b} \cdot \frac{1-cx}{1+cx}.$$

$$18. \frac{4x^4 + 4a^4 - 33x^2a^2}{2x+a} = \frac{1}{2}(4x^3 - 8x^2a - 9xa^2 - 2a^3).$$

$$19. \frac{7}{x^2 - 11x + 28} + \frac{7}{x^2 - 17x + 70} = \frac{3\frac{1}{2}x^2}{x^2 - 14x + 40}.$$

$$20. \frac{8}{x^3 - 6x + 5} + \frac{8}{x^2 - 14x + 45} = \frac{x^4}{x^2 - 10x + 9}.$$

$$21. \frac{x+a}{(a-b)(c-a)} - \frac{x-b}{(a-b)(b-c)} + \frac{x+c}{(b-c)(c-a)}$$

$$= \frac{a+c}{(a-b)(b-c)(c-a)}.$$

$$22. (x-a)^3 + (a-b)^3 + (b-x)^3 = x^3 - a^3.$$

$$23. x \left( \frac{x+2a}{x-a} \right)^3 + a \left( \frac{a+2x}{a-x} \right)^3 = 2a.$$

$$24. (x+a)^3 - (a+b)^3 + (b-x)^3 = (x+a)(x+b)(a+b).$$

$$25. x^3 - (x-b)^3 - (x-a+b)^3 - a^3 + (x-a)^3 + (a-b)^3 + b^3 = (a-b)c^2$$

$$26. (x+a)^3 - (x+b)^3 - (x-b)^3 - (2a)^3 + (x-a)^3 + (a+b)^3 + (a-b)^3 = (a^2 - b^2)c.$$

$$27. \frac{x+a}{x^2+ax+a^2} - \frac{x-a}{x^2-ax+a^2} = \frac{a^4}{a(x^4+a^2x^2+a^4)}.$$

$$28. (x+a+b)^4 - (x+a)^4 - (x+b)^4 + x^4 - (a+b)^4 + a^4 + b^4 = 12ab\{x^2 + (a+b)^2\}.$$

$$29. \frac{a-x}{a^2-bc} + \frac{b-x}{b^2-ca} + \frac{c-x}{c^2-ab} = \frac{3x}{ab+bc+ca}.$$

$$30. x^3(b-a^2) + a^3(x-b^2) + b^3(a-x^2) + abx(abx-1) = (a-x^2)(b^3-a^4).$$

31.

32.

33.

34.

35.

36.

37.

38.

39.

40.

41.

42.

43.

$$31. (1+x+x^2)^2 = \frac{ax+1}{ax-1} \cdot (1+x^2+x^4).$$

$$32. \sqrt{\left(\frac{x+a}{x+b} - \frac{a-b}{2(x+c)}\right)} = 1.$$

$$33. \frac{x+a}{x+b} = \left(\frac{2c+a+c}{2c+b+c}\right)^2.$$

$$34. \sqrt{(x^2+27x+180)} - \sqrt{(x^2+26x+168)} = \sqrt{\left(\frac{x+15}{x+12}\right)}$$

$$35. \{(x+a + \sqrt{(x^2+2ax+b^2)}\}^3 + \{x+a - \sqrt{(x^2+2ax+b^2)}\}^3 = 14(x+a)^3. \quad (\text{See page 17, Ex. 1}).$$

$$36. \{x+a + \sqrt{(x^2-2ax-2b^2)}\}^2 + \{x+a - \sqrt{(x^2-2ax-2b^2)}\}^2 = x^2 - b^2 + 2a(a-b).$$

$$37. \left(\frac{x+a}{x-b}\right)^3 = \frac{x+2a+b}{x-a-2b}.$$

$$38. (5x-7)^3 - (2x-4)^3 = 27(x^3-1).$$

$$39. \frac{1}{3} \cdot \frac{x^2-6x-1}{x^2-6x-4} + \frac{1}{5} \cdot \frac{x^2-6x-4}{x^2-6x-9} - \frac{2}{9} \cdot \frac{x^2-6x-7}{x^2-6x-16} \\ = \frac{14}{45} + \frac{4}{x^2-6x-9}.$$

$$40. \frac{1}{5} \cdot \frac{x^2-2x-3}{x^2-2x-8} + \frac{1}{9} \cdot \frac{x^2-2x-15}{x^2-2x-24} - \frac{2}{13} \cdot \frac{x^2-2x-35}{x^2-2x-48} \\ = \frac{2}{585}.$$

$$41. \{(x+a-b) + \sqrt{(x^2+a^2-b^2)}\}^3 - \{x+a-b - \sqrt{(x^2+a^2-b^2)}\}^3 = 8(x+a-b)^3.$$

$$42. \frac{1}{(x+a)^2 - a^2} + \frac{1}{(x+b)^2 - a^2} = \\ \frac{1}{x^2 - (a+b)^2} + \frac{1}{x^2 - (a-b)^2}.$$

$$43. 41 \left( \frac{6x-45}{x+1} + \frac{7x+67}{x-4} \right) + 130 = \\ 39 \left( \frac{8x+57}{x+2} + \frac{9x+68}{x+3} \right).$$

$$44. \quad 51 \left( \frac{16x+45}{x-1} - \frac{9x+25}{x-4} \right) + 863 =$$

$$61 \left( \frac{27x-3}{x-2} - \frac{7x+30}{x-3} \right).$$

$$45. \quad (x+a)(x+3a)(x+4a)(x+6a) = x^4 + 6a^2(x^2 + 7ax + 6a^2).$$

$$46. \quad \frac{1}{x+6a} + \frac{2}{x-3a} + \frac{3}{x+2a} = \frac{6}{x+a}.$$

## Exercise IV.

$$1. \quad a(b-x) + b(c-x) = b(a-x) + cx.$$

$$2. \quad (a+bx)(a-b) - (ax-b) = ab(x+1).$$

$$3. \quad (a-b)(x-c) + (a+b)(x+c) = 2(ax+ad).$$

$$4. \quad (a-b)(x-c) - (a+b)(x+c) + 2a(b+c) = 0.$$

$$5. \quad (a-b)(a-c)(a+x) + (a+b)(a+c)(a-x) = 0.$$

$$6. \quad (a-b)(a-c+x) + (a+b)(a+c-x) = 2a^2.$$

(solve in  $\{x-c\}$ ).

$$7. \quad (m+a)(a+b-x) + (a-m)(b-x) = a(m+b).$$

$$8. \quad m(a+b-x) = n(x-a-b).$$

$$9. \quad (m+n)(m-n-x) + m(x-n) - n(x-m) = m^2 - n^2.$$

$$10. \quad \frac{m-x}{m} + \frac{n-x}{n} + \frac{p-x}{p} = 3.$$

$$11. \quad \frac{a^2b-x}{a} + \frac{b^2c-x}{b} + \frac{c^2a-x}{c} = 0.$$

$$12. \quad \frac{a-x}{bc} + \frac{b-x}{ca} + \frac{c-x}{ab} = 0.$$

$$13. \quad \frac{1-ax}{bc} + \frac{1-bx}{ca} + \frac{1-cx}{ab} = 0.$$

(Deduce the solution from that of No. 12).

$$14. \quad \frac{a-bx}{bc} + \frac{b-cx}{ca} + \frac{c-ax}{ab} = 0.$$

$$15. \quad (a+b+c)x - \frac{a^2+b^2}{a-b} = \frac{2abx}{a+b} + \frac{a+b}{a-b} \cdot c$$

$$16. \frac{3abc}{a+b} + \frac{a^2b^2}{(a+b)^2} + \frac{(2a+b)b^2x}{a(a+b)^2} = \frac{(b+3ac)x}{a}.$$

$$17. \frac{10}{x} + \frac{4}{9} = \frac{9}{x} + \frac{2}{3}. \quad \left( \text{Solve in } \frac{1}{x} \right).$$

$$18. \frac{7}{x} + \frac{1}{3} = \frac{23-x}{3x} + \frac{7}{12} - \frac{1}{4x}.$$

$$19. \frac{7}{3} + \frac{13}{5x} = \frac{2(5x-12)}{3x} - \frac{17}{20} + \frac{10}{x}.$$

$$20. \frac{10-x}{3} + \frac{13-x}{7} = \frac{7x+266}{x+21} - \frac{4x+17}{2x}.$$

$$21. \frac{5}{x+3} + \frac{3}{2(x+3)} = \frac{1}{2} - \frac{7}{2(x+3)}.$$

$$22. \frac{6x+5}{8x-15} - \frac{1+8x}{15} = \frac{1-x}{3} - \frac{x-3}{5}.$$

$$23. \frac{a - \frac{1}{x}}{a + \frac{1}{x}} - \frac{1}{x} = \frac{x - \frac{1}{a}}{x + \frac{1}{a}} - \frac{1}{a}$$

$$24. \frac{a^2}{b - \frac{c^2}{d - \frac{e^2}{x}}} = 1.$$

$$25. (x-1)(x-2) - (x-3)(x-4) = 3;$$

$$(x-3)(x-4) = (x-2)(x-6).$$

$$26. 2(x-4)(3x+4) + (2x-3)(3x+2) - 6(x-2)(2x-3) = 0.$$

$$27. (a-x)(b-x) = x^2; \quad (a-x)(x-b) = x^2 - c^2.$$

$$28. (a-x)(b+x) = b^2 - x^2; \quad (x-a)(x-b) = x^2 - a^2.$$

$$29. (u+x)(b+x) = (a-x)(b-x);$$

$$(ax+b)(bx+a) = (b-ax)(a-bx).$$

$$30. (a-x)(b-x) + (a-c-x)(x-b+c) = 0.$$

$$31. (a-x)(b-x) - (c-x)(d-x) = (c+d)x - cd.$$

$$32. (x-a)(x-b) - (x-c)(x-d) = (d-a)(d-b).$$

$$33. \{(a^2 - b^2)x - ab\} \{a - (a+b)x\} + 2ab^2x = \\ \{(a+b)^2x + ab\} \{b - (a-b)x\}.$$

$$34. (x+1)(x+2)(x+3) = (x-3)(x+4)(x+5).$$

$$35. (x+1)(x+2)(x+3) = (x-1)(x-2)(x-3) + 3(x+1)(4x+1)$$

$$36. (x+1)(x+4)(x+7) = (x+2)(x+5)^2.$$

$$37. (x+2)(x+5)^2 = (x+3)^2(x+6).$$

$$38. (x-1)(x-4)(x-6) - x(x-2)(x-9) = 136$$

$$39. (a+x)(b+x)(c+x) - (a-x)(b-x)(c-x) = 2(x^3 + abc).$$

$$40. \frac{(x-a)(x-b)(x-c) - (d-a)(d-b)(d-c)}{x-d} = (x-d)^2.$$

$$41. x(x-a)^2 - (x-a+b)(x-a+c)(x-b-c) = (a^2+bc)(b+c).$$

$$42. (x-a+b)(x-b+c)(x-c+d) - x^2(x-a+d) = bc(d-a).$$

$$43. (x-a+b)(x-b+c)(x-c+d) - x(x-a+c)(x-c+d) \\ = bc(d-a).$$

$$44. (x-2a)(x-2b)(x-2c) - (x-a-b)(x-b-c)(x-c-a) \\ = (a+b+c)(a^2+b^2+c^2) - 9abc.$$

$$45. x^3 - (x-a+b)(x-b+c)(x-c+a) \\ = (a+b+c)(a^2+b^2+c^2) - 2(a^2b+b^2c+c^2a) - 3abc.$$

$$46. x \left( a - \frac{1}{x} \right) \left( b - \frac{1}{x} \right) \left( c - \frac{1}{x} \right) + \frac{1}{x^3} = \frac{a+b+c}{x}.$$

$$47. (x+a)(x+b) + (x+c)(x+a) = (x+b)(x+d) + (x+d)(x+c).$$

$$48. (ax+b)(ax-c) - a(b-x)(ax+b) = a^2(x-c)(x-b) - \\ a(ax-c)(c-x).$$

$$49. \frac{2x-3}{x-4} + \frac{3x-2}{x-8} = \frac{5x^2-29x-4}{x^2-12x+32}.$$

$$50. \frac{5x-1}{3(x+1)} - \frac{3x+2}{2(x-1)} = \frac{x^2-30x+2}{6x^2-6}.$$

51.

52.

53.

54.

55.

56.

57.

58.

59.

60.

61.

62.



$$51. \frac{3x-7}{2x-9} - \frac{3(x+1)}{2(x+3)} = \frac{11x+3}{2x^2-3x-27}.$$

$$52. \frac{7x-5}{3x-2} + \frac{8x-7}{3x-1} + \frac{10x+7}{9x^2-9x+2} = 5.$$

$$53. \frac{2x+7}{3x-7} + \frac{3x-6}{2x-5} + \frac{5(x-1)}{9x-25} = \frac{3x-2}{2x-5} + \frac{5x-8}{9x-25} + \frac{2x+2}{3x-7}.$$

$$54. \frac{4x^2-3x}{1+x} - \frac{3x}{1-x} = \frac{4x^3+2x}{x^2-1}, \quad \frac{x-a}{x-m} - \frac{x-b}{x-n} = 0.$$

$$55. \frac{\frac{1}{4} - x}{\frac{1}{4} + x} + \frac{1}{4} = \frac{x}{\frac{1}{4} - 2x} - \frac{1}{4}; \quad \frac{a}{c} + \frac{cx}{ax-b} = \frac{c}{a} + \frac{cx}{c-b}.$$

$$56. \frac{\frac{3}{2} - \frac{1}{x}}{\frac{3}{2} + \frac{1}{x}} - \frac{\frac{2}{3} - \frac{1}{x}}{\frac{2}{3} + \frac{1}{x}} = \frac{\frac{3}{2} - \frac{2}{3}}{\frac{2}{3} \cdot \frac{1}{x} + 1}.$$

$$57. \frac{2(x-1)}{x-7} + \frac{x+8}{x-4} = \frac{3(5x+16)}{5x-28}.$$

$$58. \frac{ax}{mx-p} + \frac{cx}{nx-q} = \frac{a}{m} + \frac{c}{n};$$

$$\frac{ax+b}{mx-p} + \frac{cx+d}{nx-q} = \frac{a}{m} + \frac{c}{n}.$$

$$59. \frac{b-x}{a+x} + \frac{c-x}{a-x} = \frac{a(c-2x)}{a^2-x^2};$$

$$\frac{a+b}{x-a} + \frac{b+c}{x-b} = \frac{a+c+2b}{x-c}.$$

$$60. \frac{ax+b}{ax-b} - \frac{bx}{ax+b} = \frac{ax}{ax-b} - \frac{(ax^2-2b)b}{a^2x^2-b^2}.$$

$$61. \frac{ax-b}{mx-p} + \frac{cx-d}{nx-q} + \frac{(bn+dm)x-(bq+dp)}{(mx-p)(nx-q)} = \frac{a}{m} + \frac{c}{n}.$$

$$62. \frac{m}{x-a} + \frac{n}{x-b} + \frac{p}{x-c} = \frac{m}{x-c} + \frac{n}{x-a} + \frac{p}{x-b}.$$

$$63. \frac{ax-2a}{ax-2b} = \frac{ax-2b}{ax+2a}; \frac{\frac{1}{a} - \frac{1}{x}}{\frac{1}{a} + \frac{1}{x}} = \frac{a - \frac{1}{x}}{a + \frac{1}{x}};$$

$$\frac{2x^2 - 3x + 5}{7x^2 - 4x + 2} = \frac{2}{7}.$$

$$64. \frac{ax^2 - bx + c}{mx^2 - nx + p} = \frac{a}{m}; \frac{ax^3 - bx^2 + ax - d}{mx^3 - nx^2 + mx - q} = \frac{ax - b}{mx - n}.$$

$$65. \frac{\frac{1}{4} - x}{\frac{1}{4} + x} + \frac{1}{4} = \frac{x}{\frac{1}{4} + x} - \frac{1}{4};$$

$$\frac{\frac{2}{3}x - \frac{2}{3}}{\frac{2}{3} - x} - \frac{2}{3} = \frac{2}{3} + \frac{\frac{2}{3}x + \frac{2}{3}}{x - \frac{2}{3}}.$$

$$66. \frac{21}{x-98} - \frac{71}{x-94} = \frac{21}{x+44} - \frac{71}{x-52}.$$

$$67. \frac{7}{x-6} + \frac{3}{x-11} = \frac{9}{x-7} + \frac{1}{x-12};$$

$$\frac{9}{x-51} - \frac{9}{x-15} = \frac{2}{x-81} - \frac{2}{x+81}.$$

$$68. \frac{5}{x-6} + \frac{4}{x-9} = \frac{8}{x-7} + \frac{1}{x-10};$$

$$\frac{1}{x-6} + \frac{8}{x-3} = \frac{5}{x-2} + \frac{4}{x-5}.$$

$$69. \frac{m-n}{x-a} - \frac{a-b}{x-m} = \frac{m-n}{x-b} - \frac{a-b}{x-n}.$$

$$70. \frac{a+b}{x-b} - \frac{a+c}{x-c} = \frac{b+d}{x-(a+b+2c+d)} - \frac{c+d}{x-(a+2b+c+d)}.$$

$$71. (x-a+b)^3 - (x-a)^3 + (x-b)^3 - x^3 + a^3 - (a-b)^3 - b^3 \\ = (a-b)c^2.$$

$$72. (x+a+b)^5 - (a+b)^5 - (x+b)^5 - (x+a)^5 + x^5 + a^5 + b^5 \\ = 10abx(2x+a+b)(x+a+b).$$

$$73. \frac{(m-n)(x-a)}{b+c} + \frac{(n-p)(x-b)}{c+a} + \frac{(p-m)(x-c)}{a+b} = 0.$$

74.

75.

76.

77.

78.

79.

80.

81.

Art  
ple illusDEFIN  
expressio  
Thus, ifBy TI  
 $f(a)^n = 0$ ,  
solving t  
 $x$  which  
pose  $f(x)$

$$74. \frac{ax-1}{a^2(c+b)} + \frac{bx-1}{b^2(c+a)} + \frac{cx-1}{c^2(a+b)} = \frac{3x}{ab+bc+ca}.$$

$$75. \frac{x-2a}{b+c-a} + \frac{x-2b}{c+a-b} + \frac{x-2c}{a+b-c} = 3.$$

$$76. \frac{x-2a}{b+c-a} + \frac{x-2b}{c+a-b} + \frac{x-2c}{a+b-c} = \frac{3x}{a+b+c}.$$

$$77. \frac{a-x}{a^2-bc} + \frac{b-x}{b^2-ac} + \frac{c-x}{c^2-ab} = \frac{3}{a+b+c}.$$

$$78. \frac{x+2ab}{a+b+c} + \frac{2ab-x}{b+c-a} = \frac{x-2ab}{a-b+c} + \frac{x+2ab}{a+b-c}.$$

$$79. \frac{a}{x+b-c} + \frac{b}{x+a-c} = \frac{a-c}{x+b} - \frac{b+c}{x+a}.$$

$$80. \frac{m^2(a-b)}{x-m} + \frac{n^2(b-c)}{x-n} + \frac{p^2(c-d)}{x-p} +$$

$$\frac{q \{pd+(n-p)c+(m-n)b-ma\}}{x-q} = 0.$$

$$81. \frac{(x-2)(x-5)(x-6)(x-9) + (a+2)(a-4)(a-5)(a-11)}{x} \\ + \frac{(b+1)(b+5)(b+8)(b+12)}{x} = (x-4)(x-7)(x-11) + \\ \frac{(a^2-1)(a-8)(a-10) + (b+2)(b+3)(b+10)(b+11)}{x}.$$

Art XLV. Employing the language of algebra, the principle illustrated in Art. XL. may be stated as follows:

DEFINITION.—Any quantity which substituted for  $x$  makes the expression  $f(x)$  vanish, is said to be a root of the equation  $f(x) = 0$ . Thus, if  $a$  is a root of the equation  $f(x) = 0$ , then  $f(a) = 0$ .

By Th. I., if  $x-a$  is a factor of the polynome  $f(x)^n$ , then  $f(a)^n = 0$ , and  $a$  must be a root of the equation  $f(x)^n = 0$ ; hence in solving the equation we are merely finding a value, or values, of  $x$  which will make the corresponding polynome vanish. Suppose  $f(x)^n = (x-a)\varphi(x)^{n-1} = 0$ , we are required to find a value, or

values, of  $x$  which will make  $(x-a)\varphi(x)^{n-1}$  vanish. The polynome will certainly vanish if *one* of its factors vanishes, whether the other does or not, and will not vanish unless at least *one* of its factors vanishes. Hence  $(x-a)\varphi(x)^{n-1}$  will vanish if  $x-a=0$ , quite irrespective of the value of  $\varphi(x)^{n-1}$ . Also, if  $\varphi(x)^{n-1}=0$ , the polynome will vanish, irrespective of the value of  $x-a$ . It follows, therefore, that if  $f(x)^n$  can be resolved into two or more factors, each of these factors equated to zero will give one or more roots of the equation  $f(x)^n=0$ .

When there can be found two or more values of  $x$  which satisfy the conditions of given equations, they are sometimes distinguished thus:  $x_1, x_2, x_3$ , &c., to be read "one value of  $x$ ," "a second value of  $x$ ," "a third value of  $x$ ," &c. Thus, if

$$(x-a)(x-b)(x-c)=0,$$

$$\therefore x_1 = a, x_2 = b, x_3 = c.$$

## EXAMPLES.

1. Solve  $2x^3 - 13x^2 + 27x - 18 = 0$ .

Factoring,

$$(x-2)(x-3)(2x-3)=0,$$

$$\therefore x_1 = 2, x_2 = 3, x_3 = 1\frac{1}{2}.$$

2.  $x^2 - (a+b)x + (a+c)b = (a+c)c$ ,

$$\therefore x^2 - (a+b)x + (a+c)(b-c) = 0,$$

$$\therefore x^2 - \{(a+c) + (b-c)\}x + (a+c)(b-c) = 0,$$

$$\therefore \{x - (a+c)\} \{x - (b-c)\} = 0,$$

$$\therefore x_1 = a+c, x_2 = b-c.$$

3.  $x^2(a-b) + a^2(b-x) + b^2(x-a) = 0$ .

$$\therefore x^2(a-b) - x(a^2 - b^2) + ab(a-b) = 0,$$

$$\therefore (x-a)(x-b)(a-b) = 0.$$

If  $a-b=0$ , the given equation holds irrespective of the values of  $x-a$  and  $x-b$ , and therefore of the values of  $x$ ; but if  $a-b$  is not zero,  $x_1 = a, x_2 = b$ .

$$4. \quad x = \frac{(a^2 + b^2)x - (a^2 - b^2)}{(a^2 - b^2)x - (a^2 + b^2)};$$

$$\therefore \frac{x+1}{x-1} = \frac{a^2(x-1)}{b^2(x+1)} \quad \therefore \left(\frac{x+1}{x-1}\right)^2 - \frac{a^2}{b^2} = 0,$$

$$\therefore \frac{x_1+1}{x_1-1} - \frac{a}{b} = 0 \quad \therefore x_1 = \frac{a+b}{a-b},$$

$$\frac{x_2+1}{x_2-1} + \frac{a}{b} = 0 \quad \therefore x_2 = \frac{a-b}{a+b}.$$

$$5. \quad \frac{(a-x)^2 + (b-x)^2}{(a-x)^2 + (a-x)(b-x) + (b-x)^2} = \frac{34}{49}.$$

$$\therefore \frac{(a-x)^2 + 2(a-x)(b-x) + (b-x)^2}{(a-x)^2 - 2(a-x)(b-x) + (b-x)^2} = \frac{2(49) - 34}{3(34) - 2(49)} = 16,$$

$$\therefore \frac{\{(a-x) + (b-x)\}^2}{\{(a-x) - (b-x)\}^2} - 16 = 0,$$

$$\therefore \frac{(a-x_1) + (b-x_1)}{a-b} - 4 = 0, \quad \therefore x_1 = \frac{1}{2}(5b-3a);$$

$$\frac{(a-x_2) + (b-x_2)}{a-b} + 4 = 0, \quad \therefore x_2 = \frac{1}{2}(5a-3b).$$

$$6. \quad \frac{(x-a)(x-b)}{(c-a)(c-b)} + \frac{(x-b)(x-c)}{(a-b)(a-c)} = 1.$$

Subtract term by term from the identity (See page 53),

$$\frac{(x-a)(x-b)}{(c-a)(c-b)} - \frac{(x-b)(x-c)}{(a-b)(a-c)} + \frac{(x-c)(x-a)}{(b-c)(b-a)} = 1$$

$$\therefore (x-c)(x-a) = 0, \quad \therefore x_1 = c, \quad x_2 = a.$$

7. Find the *rational* roots of  $x^4 - 12x^3 + 51x^2 - 90x + 56 = 0$ .

Factoring the left-hand member by the method of Art. xxviii.,

$$(x-2)(x-4)(x^2 - 6x + 7) = 0$$

$$\therefore x_1 = 2, \quad x_2 = 4, \quad \text{or } x^2 - 6x + 7 = 0.$$

Since  $x^2 - 6x + 7$  cannot be resolved into rational factors we know that it will not give rational roots,  $\therefore x_1 = 2, x_2 = 4$  are the only values that meet the condition of the problem.

Any literal equation of the second, third, or fourth degree, and many equations of the higher degree can be resolved into a series of disjunctive equations. A full analysis for the first four degrees will be given in Part II., meanwhile the following special forms of the Theorem in Art. XLV., will enable the student to solve nearly all the equations commonly proposed.

(A). In order that two expressions having a common factor may be equal, it is necessary either that the common factor should vanish, or else that the product of the remaining factors of one of the expressions should be equal to the product of the remaining factors of the other expression, and it is sufficient if one of these conditions be fulfilled. In symbols this is

$$\text{If } (x-a)f(x) = (x-a)\phi(x), \therefore x_1 = a \text{ or } f(x) = \phi(x).$$

(B). If an equation reduces to the form  $(mx+n)^2 = c^2$

$$\therefore (mx+n)^2 - c^2 = 0,$$

$$\therefore (mx_1+n) - c = 0 \text{ and } \therefore x_1 = \frac{c-n}{m},$$

$$\text{or } (mx_2+n) + c = 0 \text{ and } \therefore x_2 = \frac{-c-n}{m}.$$

(C). If an equation reduces to the form

$$\left\{ \frac{mx+n}{px+q} \right\}^2 = \frac{a^2}{b^2},$$

$$\text{then } x_1 = \frac{qa-nb}{mb-pa}, \quad x_2 = \frac{-qa-nb}{mb+pa}. \quad (\text{See Exs. 4 and 5 above}).$$

(D). If an equation appears under the form

$$(a-x)(x-b) = c, \tag{1}$$

$$\text{then } x_1 = \frac{1}{2}(a+b+r), \quad x_2 = \frac{1}{2}(a+b-r),$$

$$\text{in which } r^2 = (a-b)^2 - 4c.$$

From the identity  $(a-x) + (x-b) = a-b$

$$\text{we get } (a-x)^2 + 2(a-x)(x-b) + (x-b)^2 = (a-b)^2 \tag{2}$$

$$(2) - 4(1) \quad \therefore (a-x)^2 - 2(a-x)(x-b) + (x-b)^2 = (a-b)^2 - 4c = r^2 \text{ say}$$

$$\therefore \{(a-x) - (x-b)\}^2 - r^2 = 0,$$

$$\therefore \{(a-x_1) - (x_1-b)\} + r = 0, \text{ and } \therefore x_1 = \frac{1}{2}(a+b+r);$$

$$\text{or } \{(a-x_2) - (x_2-b)\} - r = 0, \text{ and } \therefore x_2 = \frac{1}{2}(a+b-r).$$

$$8. \quad x + \frac{1}{x} = a + \frac{1}{a} \quad \therefore x - a = \frac{1}{a} - \frac{1}{x},$$

$$\therefore \frac{x-a}{1} = \frac{x-a}{ax} \quad \text{Applying (A),}$$

$$\therefore x-a=0, \text{ or } ax=1,$$

$$\therefore x_1 = a, \quad x_2 = \frac{1}{a}.$$

$$9. \quad (x+a+b)(x+b+c) = (x-3a+b)(2x-3a+2b-c);$$

$$\therefore \frac{x+a+b}{x-3a+b} = \frac{2x-3a+2b-c}{x+b+c}$$

$$= \frac{x-4a+b-c}{3a+c}$$

Page 122. (5).

$$\therefore \frac{2(x-a+b)}{x-3a+b} = \frac{x-a+b}{3a+c},$$

$$\therefore (A) \quad x_1 = a - b$$

$$\frac{1}{2}(x_2 - 3a + b) = 3a + c \quad \therefore x_2 = 9a - b + 2c.$$

$$10. \quad \frac{(x+2)^2}{x^2-2x} = \frac{a}{b}, \quad \therefore \frac{(x+2)^2}{m(x+2)^2+n(x^2-2x)} = \frac{a}{ma+nb} \quad (1)$$

But (C) can be applied if  $m$  and  $n$  are so determined that  $m(x+2)^2+n(x^2-2x)$  is a square.

This requires that  $4m(m+n) = (2m-n)^2$ ,

$$\therefore 4m^2 + 4mn = 4n^2 - 4mn + n^2,$$

$$\therefore 8m = n.$$

Assume  $m=1$ , then  $n=8$ , and (1) becomes, on substitution and reduction,

$$\frac{(x+2)^2}{(3x-2)^2} = \frac{a}{a+8b} = r^2, \text{ say}$$

$$\therefore x_1 = \frac{2(1+r)}{3r-1}, \quad x_2 = \frac{2(r-1)}{1+3r}.$$

$$11. \quad \frac{(x+1)^4}{(x^2+1)(x-1)^2} = \frac{a}{b} \quad \therefore \frac{(x^2+2x+1)^2}{(x^2+1)(x^2-2x+1)} = \frac{a}{b}.$$

For  $x^2+1$  write  $xz$

$$\therefore \frac{(xz+2x)^2}{xz(xz-2x)} = \frac{a}{b} \quad \therefore \frac{(z+2)^2}{z(z-2)} = \frac{a}{b}.$$

This equation was solved in Ex. 10, hence  $z$  may be treated as known.

$$\text{But } \frac{x^2+1}{x} = z, \quad \therefore \frac{x^2+2x+1}{x^2-2x+1} = \frac{z+2}{z-2}.$$

$$\therefore \left(\frac{x+1}{x-1}\right)^2 = \frac{z+2}{z-2}, \text{ a formed solved in (C).}$$

$$12. (a-x)^4 + (b-x)^4 = c.$$

In the identity

$$(u+v)^4 = u^4 + v^4 + 4(u+v)^2 uv - 2u^2 v^2,$$

Let  $u = a-x$ ,  $v = x-b$ ,  $\therefore u+v = a-b$  and  $u^4 + v^4 = c$ ,

$$\therefore (a-b)^4 = c + 4(a-b)^2(a-x)(x-b) - 2(a-x)^2(x-b)^2$$

Write  $z$  for  $(a-x)(x-b)$

$$\therefore z^2 - 2(a-b)^2 z + (a-b)^4 = \frac{1}{2}\{c + (a-b)^4\} = t^2, \text{ say,}$$

$$\therefore \{z - (a-b)^2\}^2 = t^2$$

$$\therefore \text{by (B)} \quad z_1 = (a-b)^2 - t; \quad z_2 = (a-b)^2 + t, \quad \therefore z \text{ is known;}$$

But  $(a-x)(x-b) = z$

$$\therefore \text{by (D)} \quad x_1 = \frac{1}{2}(a+b+r); \quad x_2 = \frac{1}{2}(a+b-r) \quad (1).$$

$$\text{in which } r^2 = (a-b)^2 - 4z,$$

$$\begin{aligned} &= (a-b)^2 - 4\{(a-b)^2 - t\} = 4t - 3(a-b)^2 \\ \text{or } &(a-b)^2 - 4\{(a-b)^2 + t\} = -4t - 3(a-b)^2 \end{aligned} \quad (2)$$

$$\text{and } t^2 = \frac{1}{2}\{c + (a-b)^4\}. \quad (3)$$

Hence  $x$  is expressed in terms of  $a$ ,  $b$ , and  $r$ ,

$r$  is expressed in terms of  $a$ ,  $b$ , and  $t$ ,

$t$  is expressed in terms of  $a$ ,  $b$ , and  $c$ ,

and the expressions for  $r$  and  $t$  are cases of (B).

$$13. (a-x)(b+x)^4 + (a-x)^4(b+x) = ab(a^3 + b^3)$$

$$\text{Let } a-x = n-z \text{ and } b+x = n+z \quad \therefore n = \frac{1}{2}(a+b) \quad (1).$$

The equation reduces to

$$(n^2 - z^2)\{(n+z)^3 + (n-z)^3\} = ab(a^3 + b^3)$$

$$\therefore (n^2 - z^2)(2n^3 + 6nz^2) = ab(a^3 + b^3)$$

$$\therefore (n^2 - z^2)(n^2 + 3z^2) = ab(a^2 - ab + b^2)$$



$z^2$  may now be found by (D), and from the result  $z$  may be found by (B), and from (1)  $x = \frac{1}{2}(a-b) + z$ ;

$$3z^2 = \frac{3}{4}(a-b)^2 \text{ or } \frac{1}{4}(10ab - a^2 - b^2)$$

$$\therefore x = 0, \text{ or } a-b, \text{ or } \frac{1}{2}(a-b) + \frac{1}{6}\sqrt{(30ab - 3a^2 - 3b^2)}.$$

$$14. \{ \sqrt[4]{(a+x)} + \sqrt[4]{(a-x)} \}^2 \{ \sqrt{(a+x)} + \sqrt{(a-x)} \} = 2cx.$$

Divide the terms of the identity

$$\sqrt[4]{(a+x)}^4 - \sqrt[4]{(a-x)}^4 = 2x$$

by the corresponding terms of the equation,

$$\therefore \sqrt[4]{\frac{(a+x)}{(a-x)}} = \frac{c+1}{c-1}, \therefore \frac{a+x}{a-x} = \left(\frac{c+1}{c-1}\right)^4;$$

$$\therefore x = a \cdot \frac{(c+1)^4 - (c-1)^4}{(c+1)^4 + (c-1)^4}.$$

$$15. \sqrt[3]{(a-x)^2} + \sqrt[3]{\{(a-x)(b-x)\}} + \sqrt[3]{(b-x)^2} = \sqrt[3]{(a^2 + ab + b^2)}$$

Divide the terms of the identity

$$\sqrt[3]{(a-x)^3} - \sqrt[3]{(b-x)^3} = a-b$$

by the corresponding terms of the equation.

$$(1). \quad \therefore \sqrt[3]{(a-x)} - \sqrt[3]{(b-x)} = \frac{a-b}{\sqrt[3]{(a^2 + ab + b^2)}}.$$

Cube, using the form  $(u-v)^3 = u^3 - v^3 - 3uv(u-v)$ .

$$(a-x) - (b-x) - 3\sqrt[3]{\{(a-x)(b-x)\}} \cdot \frac{a-b}{\sqrt[3]{(a^2 + ab + b^2)}}$$

$$(3) \quad = \frac{(a-b)^3}{a^2 + ab + b^2} = a - b - \frac{3ab(a-b)}{a^2 + ab + b^2},$$

$$\therefore \sqrt[3]{\{(a-x)(b-x)\}} = \frac{ab}{\sqrt[3]{(a^2 + ab + b^2)^2}}$$

$$\therefore (a-x)(b-x) = \frac{a^2 b^3}{(a^2 + ab + b^2)^2}$$

a form solved in (D).

(1).

$$16. \frac{\{\sqrt{(a-x)} + \sqrt{(x-b)}\}^2}{\sqrt{(a-x)} - \sqrt{(x-b)}} = \sqrt{c}$$

Assume  $\sqrt{(a-x)} = z\sqrt{(x-b)}$

$$\therefore (a-x) + (x-b) = (z^2 + 1)(x-b).$$

$$\therefore x - b = \frac{a - b}{z^2 + 1}.$$

The proposed equation now becomes

$$\frac{\sqrt{(x-b)(z+1)^2}}{z-1} = \sqrt{c}$$

$$\therefore \frac{(x-b)(z-1)^4}{(z-1)^2} = c.$$

$$\therefore \frac{(z+1)^4}{(z^2+1)(z-1)^2} = \frac{c}{a-b}, \text{ a form solved in Ex. 11.}$$

$$17. (x-2)(x-5)(x-6)(x-9) + (y+2)(y-4)(y-5)(y-11) + \\ (z+1)(z+5)(z+8)(z+12) = x(x-4)(x-7)(x-11) + \\ (y+1)(y-1)(y-8)(y-10) + (z+2)(z+3)(z+10)(z+11).$$

Let  $x' = x^2 - 11x$ ,  $y' = y^2 - 9y$  and  $z' = z^2 + 13z$ ,

$$\therefore (x'+18)(x'+30) + (y'-22)(y'+20) + (z'+12)(z'+40) = \\ x'(x'+28) + (y'-10)(y'+8) + (z'+22)(z'+30)$$

$$\therefore x'^2 + 48x' + 540 + y'^2 - 2y' - 440 + z'^2 + 52z' + 480 = \\ x'^2 + 28x' + y'^2 - 2y' - 80 + z'^2 + 52z' + 660,$$

$$\therefore 20x' = 0, \therefore x^2 - 11x = 0, \therefore x_1 = 0, x_2 = 11.$$

### Exercise lvi.

What can you deduce from the following statements ?

1.  $A \cdot B = 0$ . 2.  $A \cdot B \cdot C = 0$ . 3.  $(a-b)x = 0$ . 4.  $12xy = 0$ .

5. What is the difference between the equation

$$(x-5y)(x-4y+3) = 0$$

and the simultaneous equations

$$x-5y = 0 \text{ and } x-4y+3 = 0.$$

What values of  $x$  will satisfy the following equations ?

6.  $x(x-a) = 0$ . 7.  $ax(x+b) = 0$ . 8.  $(x-a)(bx-c) = 0$ .

9.  $ax^2 = 3ax$ . 10.  $x^2 = (a+b)x$ . 11.  $x(x^2 - a^2) = 0$ .

12.  $a^2x^3 = b^2x$ . 13.  $x^2 + (a-x)^2 = a^2$ .

14.  $x^2 + (a-x)^2 = (a-2x)^2$ . 15.  $(a-x)^2 + (x-b)^2 = a^2 + b^2$ .

16.  $(a-x)(x-b) + ab = 0$ .

17.  $(a-x)^2 - (a-x)(x-b) + (x-b)^2 = a^2 + ab + b^2$ .

18.  $x^2 - (a-b)x - ab = 0$ .

19.  $x^3 - (a+b+c)x^2 - (ab+bc+ca)x - abc = 0$ .

If  $x$  must be positive, what value or values of  $x$  will satisfy the following equations ?

20.  $(x-5)(x+4)=0$ . 21.  $x^2+29x-30=0$ .

22.  $x^2-17x-84=0$ . 23.  $3x^2+10x+3=0$ .

24.  $x^4-13x^2+36=0$ . 25.  $x^3-2x^2-5x+6=0$ .

Solve the following equations :

26.  $(a-x)^2+(x-b)^2=(a-b)^2$ .

27.  $(a-x)^2-(a-x)(x-b)+(x-b)^2=(a-b)^2$

28.  $a^2(a-x)^2=b^2(b-x)^2$ . 29.  $a^2(b-x)^2=b^2(a-x)^2$ .

30.  $(x-a)^3+(a-b)^3+(b-x)^3=0$ . 31.  $(x-1)^2=a(x^2-1)$ .

32.  $\frac{a-x}{x-b} = \frac{x-a}{c+x}$ . 33.  $\frac{a+b-x}{a-c-x} = \frac{a-c+x}{a+c-x}$ .

34.  $(x-a+b)(x-a+c)=(a-b)^2-x^2$ .

35.  $(x-a)^2-b^2+(a+b-x)(b+c-x)=0$ .

36.  $(a+b+c)x^2-(2a+b+c)x+a=0$ .

37.  $\frac{a+b-x}{c} = \frac{a+b-c}{x}$ .

38.  $(a-x)^2+(a-b)^2=(a+b-2x)^2$ .

39.  $x(a+b-x)+(a+b+c)c=0$ .

40.  $(n-p)x^2+(p-m)x+m-n=0$ .

41.  $\frac{ax^2-bx+c}{mx^2-nx+p} = \frac{c}{p}$ . 42.  $\frac{ax^2-bx+c}{mx^2-nx+p} = \frac{a-b+c}{m-n+p}$

43.  $4x^2+a^2-b^2-2(a+b)x=(a-x)(b+x)-(a+x)(b-x)$ .

44.  $(2a-b-x)^2+9(a-b)^2=(a+b-2x)^2$ .

45.  $(2a+2c-x)^2=(2b+x)(3a-b+3c-2x)$ .

46.  $(3a-5b+x)(5a-5b-x)=(7a-b-2x)^2$ .

47.  $(3a-b+x)(3a+b-x)=(5a+3b-3x)^2$ .

48.  $a(a-b)-b(a-c)+c(b-c)x^2=0$ .

$$49. (ab+bc+ca)(x^2+x+1)+(a-b)^2=(2ac+b^2)(x^2+x+1) \\ + (a-c)^2x. \quad 74.$$

$$50. (x+1)(x+3)(x-4)(x-7)+(x-1)(x-3)(x+4)(x+7)=96. \quad 75.$$

$$51. (x-1)(x+3)(x-5)(x+9)+(x+1)(x-3)(x+5)(x+9)+18 \\ =0. \quad 76.$$

$$52. x + \frac{1}{x} = 3\frac{1}{2}. \quad 77.$$

$$53. x + \frac{1}{x} = \frac{a+b}{a-b} + \frac{a-b}{a+b}. \quad 78.$$

$$54. x - \frac{1}{x} = \frac{a}{b} - \frac{b}{a}. \quad 79.$$

$$55. \frac{a+x}{b+x} + \frac{b+x}{a+x} = 2\frac{1}{2}. \quad 80.$$

$$56. \frac{a-x}{x-b} + \frac{x-b}{a-x} = \frac{13}{6}. \quad 81.$$

$$57. \frac{a-x}{b+x} - \frac{b+x}{a-x} = \frac{m}{n} - \frac{n}{m}. \quad 82.$$

$$58. \frac{a}{x} + \frac{x}{a} = \frac{m}{n}. \quad 83.$$

$$59. \frac{x^2+ax+a^2}{x^2-ax+a^2} = c. \quad 84.$$

$$60. \frac{x^2+a^2}{x^2-ax+a^2} = c. \quad 85.$$

$$61. \frac{x^2+a^2}{(x+a)^2} = c. \quad 86.$$

$$62. \frac{(a-x)^2+(x-b)^2}{(a-x)(x-b)} = \frac{5}{2}. \quad 87.$$

$$63. \frac{a-x}{x-b} + \frac{x-b}{a-x} = \frac{m}{n}. \quad 88.$$

$$64. \frac{(x+a)^2+(x-b)^2}{(x+a)^2-(x-b)^2} = \frac{a^2+b^2}{2ab}. \quad 89.$$

$$65. \frac{(a-x)^2-(x-b)^2}{(a-x)(x-b)} = \frac{4ab}{(a^2-b^2)}. \quad 90.$$

$$66. \frac{(a-x)^2+(a-x)(x-b)+(x-b)^2}{(a-x)^2-(a-x)(x-b)+(x-b)^2} = \frac{49}{19}. \quad 91.$$

$$67. \frac{2a^2+a(a-x)+(a+x)^2}{2a^2+a(a+x)+(a-x)^2} = \frac{c+1}{c-1}. \quad (\text{Also for } c=5). \quad 92.$$

$$68. (5-x)^4+(2-x)^4=17. \quad 93.$$

$$69. x^4+(a-x)^4=c; \quad x^4+(x-4)^4=82. \quad 94.$$

$$70. (a-x)^4+(x-b)^4=(a-b)^4. \quad 71. (a-x)^5+(x-b)^5=c. \quad 95.$$

$$72. x^5+(a-x)^5=a^5; \quad x^5+(6-x)^5=1056. \quad 96.$$

$$73. (a-x)^2(x-b)^2+(a-x)^2(x-b)^3=a^2b^2(a-b). \quad 97.$$

$$74. (a-x)(b+x)^4 + (a-x)^2(b+x)^3 + (a-x)^3(b+x)^2 + (a-x)^4(b+x) = (a+b)c.$$

$$75. \frac{(a-x)^4 + (x-b)^4}{(a-x)^2 + (x-b)^2} = \frac{41}{20} (a-b)^2.$$

$$76. \frac{(a-x)^5 + (x-b)^5}{(a-x)^4 + (x-b)^4} = \frac{211}{97} (a-b).$$

$$77. \frac{(a-x)^4 + (x-b)^4}{(a-x)^2 + (x-b)^2} = \frac{a^4 + b^4}{a^2 + b^2}.$$

$$78. \frac{(a-x)^4 + (x-b)^4}{(a-x)^3 + (x-b)^3} = \frac{a^4 + b^4}{a^3 + b^3}.$$

$$79. \frac{(a-x)^5 + (x-b)^5}{(a-x)^3 + (x-b)^3} = \frac{a^5 - b^5}{a^3 - b^3}.$$

$$80. \frac{(a-x)^3}{b-x} + \frac{(b-x)^3}{a-x} = \frac{a^3}{b} + \frac{b^3}{a}.$$

$$81. \frac{a-x}{(x-b)^2} + \frac{x-b}{(a-x)^2} = \frac{a}{b^2} - \frac{b}{a^2}.$$

$$82. \frac{(a-x)^4 + (x-b)^4}{(a+b-2x)^2} = \frac{a^4 + b^4}{(a+b)^2}.$$

$$83. \frac{(a-x)^5 + (x-b)^5}{(a+b-2x)^2} = \frac{a^5 - b^5}{(a+b)^2}.$$

$$84. \frac{(a-x)^5 + (x-b)^5}{(a-x)^2 + (x-b)^2} = (a-b)^3.$$

$$85. \frac{(a-x)^4 - (x-b)^4}{(a-x) - (x-b)} = \frac{(a-b)c}{(a-x)(x-b)}.$$

$$86. \frac{(a-x)^5 + (x-b)^5}{(a-x)^2 + (x-b)^2} = c(a-x)(x-b).$$

$$87. \frac{(a-x)^3 + (x-b)^3}{(a-x)^4 + (x-b)^4} = \frac{c}{(a-x)(x-b)}.$$

$$88. (1+x^2)^3 = (x^3-3)^2.$$

$$89. \frac{x^4+1}{2x(x^2+1)} = \frac{a}{b} \quad 90. \frac{(x+1)^2(x^2+1)}{(x-1)^2(x^2-x+1)} = \frac{a}{b}$$

$$91. \frac{(x-1)^2x}{(x^2-x+1)^2} = \frac{a}{b} \quad 92. \frac{(x^2+x+1)^2}{(x+1)^2(x^2+1)} = \frac{a}{b}$$

$$93. \frac{(x^2+1)^2}{x(x+1)^2} = \frac{a}{b} \quad 94. \frac{(x+1)^4}{x(x^2+1)} = \frac{a}{b}$$

$$95. \frac{x(x+1)^2}{(x-1)^4} = \frac{a}{b} \quad 96. \frac{x^2+x+1}{(x+1)^2} \cdot \frac{x^2+x-1}{(x-1)^2} = \frac{a}{b}$$

$$97. \frac{x^4-x^2+1}{(x^2-1)^2} = \frac{a}{b} \quad 98. \frac{x(x^2+1)}{(x^2-1)^2} = \frac{a}{b}$$

$$99. \frac{(x+1)(x^3+1)}{(x-1)(x^3-1)} = \frac{a}{b} \quad 100. \frac{(x+1)(x^5-1)}{(x-1)(x^5+1)} = \frac{a}{b}$$

$$101. \frac{(x+1)^4}{x^4+1} = \frac{a}{b} \quad 102. \frac{(x+1)^5}{x^5+1} = \frac{a}{b}$$

$$103. 2(a-x)^4 - 9(a-x)^3(x-b) + 14(a-x)^2(x-b)^2 - 9(a-x)(x-b)^3 + 2(x-b)^4 = 0.$$

$$104. 4(a-x)^4 - 17(a-x)^2(x-b)^2 + 4(x-b)^4 = 0.$$

Find the rational roots in the following equations :

$$105. x^4 - 12x^3 + 49x^2 - 78x + 40 = 0. \quad [\text{Let } z = x^2 - 6x].$$

$$106. x^4 - 6x^3 + 7x^2 + 6x - 8.$$

$$107. x^4 - 10x^3 + 35x^2 - 50x + 24 = 0.$$

$$108. 32x^4 - 48x^3 - 10x^2 + 21x + 5 = 0.$$

$$109. x^3 - 6x^2 + 5x + 12 = 0.$$

$$110. \frac{5}{x} - \frac{4}{x-a} - \frac{9}{x-2a} - \frac{4}{x-3a} + \frac{5}{x-4a} = 0.$$

$$111. \frac{14}{x+20} + \frac{5}{x+5} - \frac{4}{x-4} = \frac{14}{x-55} + \frac{5}{x-40} - \frac{4}{x-25}$$

$$112. \frac{2x+5a}{x} - \frac{x+8a}{x-a} + \frac{x}{x-2a} =$$

$$\frac{x-a}{a-3a} - \frac{x+5a}{x-4a} + \frac{2x-5a}{x-5}$$

$$113. \frac{x+4}{x+2} + \frac{x+2}{x} + \frac{x+4}{x-1} = \frac{x+3}{x-2} + \frac{x-1}{x-3} + \frac{x-3}{x-5}$$

$$114. \frac{7}{x} - \frac{31}{x-1} + \frac{20}{x-2} + \frac{8}{x-3} + \frac{20}{x-4} - \frac{31}{x-5} + \frac{7}{x-6} = 0.$$

$$115. \sqrt{(x^2 - a^2 - b^2)} + \sqrt{(x^2 - b^2 - c^2)} - \sqrt{(x^2 - c^2 - a^2)} = x.$$

$$116. \frac{\sqrt{(a^2 + 2x)} + \sqrt{(a^2 - 2x)}}{\sqrt{(a^2 + 2x)} - \sqrt{(a^2 - 2x)}} = \frac{m^2 x^2}{a^2} \cdot \frac{\sqrt{(m^2 x + 2)} + \sqrt{(m^2 x - 2)}}{\sqrt{(m^2 x + 2)} - \sqrt{(m^2 x - 2)}}$$

$$117. \sqrt{(x^2 - a^2)} + \sqrt{(x^2 - b^2)} + \sqrt{(x^2 - c^2)} = x.$$

$$118. \{\sqrt{(a-x)} + \sqrt{(b-x)}\} \{\sqrt{(a-x)} - \sqrt{(b-x)}\} = a.$$

$$119. \frac{\sqrt[3]{(a-x)} - \sqrt[3]{(x-b)}}{\sqrt[3]{(a-x)} + \sqrt[3]{(x-b)}} = \frac{a+b-2x}{a-b}.$$

$$120. \sqrt[5]{(a+x)} + \sqrt[5]{(a-x)} = \sqrt[5]{(2a)}.$$

$$121. \frac{\{\sqrt[3]{(a-x)^2} + \sqrt[3]{(x-b)^2}\}^2}{\sqrt[3]{(a-x)} + \sqrt[3]{(x-b)}} = a-b.$$

[Write  $u$  for  $\sqrt[3]{(a-x)}$ , and  $v$  for  $\sqrt[3]{(x-b)}$ ].

 $\frac{a}{b}$  $\frac{a}{b}$  $\frac{a}{b}$ 

c].

 $\frac{1}{x-25}$

CHAPTER VI.

SIMULTANEOUS EQUATIONS.

Art. XLVI. There are three general methods of resolving simultaneous linear equations, 1° by substitution, 2° by comparison, 3° by elimination. The last is often subdivided into the method by cross-multipliers, and the method by arbitrary multipliers.

In applying the elimination-method the work should be done with detached coefficients, each equation should be numbered, and a register of the operations performed should be kept.

Ex. Resolve  $u + v + x + y + z = 15.$   
 $u + 2v + 4x + 8y + 16z = 57.$   
 $u + 3v + 9x + 27y + 81z = 179.$   
 $u + 4v + 16x + 64y + 256z = 453.$   
 $u + 5v + 25x + 125y + 625z = 975.$

Register	<i>u</i>	<i>v</i>	<i>x</i>	<i>y</i>	<i>z</i>	
	1	1	1	1	1	= 15 (1)
	1	2	4	8	16	57 (2)
	1	3	9	27	81	179 (3)
	1	4	16	64	256	453 (4)
	1	5	25	125	625	975 (5)
(2) - (1).		1	3	7	15	42 (6)
(3) - (2).		1	5	19	65	122 (7)
(4) - (3).		1	7	37	175	274 (8)
(5) - (4).		1	9	61	369	522 (9)
(7) - (6).			2	12	50	80 (10)
(8) - (7).			2	18	110	152 (11)
(9) - (8).			2	24	194	248 (12)
(11) - (10).				6	60	72 (13)
(12) - (11).				6	84	96 (14)
(14) - (13).					24	24 (15)
(15) ÷ 24.					1	1 (16)
$\frac{1}{24} \{ (13) - 60(16) \}.$						2 (17)
$\frac{1}{2} [ (10) - \{ 12(17) + 50(16) \} ].$				1		3 (18)
(6) - $\{ 3(18) + 7(17) + 15(16) \}.$				1		4 (19)
(1) - $\{ (19) + (18) + (17) + (16) \}.$						5 (20)

An ex  
 have be  
 (3) + (1)  
 3(2) - 3(

A gen  
 will be g  
 tions it

Form  
 thus—  
 Form  
 thus—  
 Subtra

Divide  
 quotient  
 value of  
 [Writin  
 the gener  
 Ex. 1.



An examination of the Register will show how easy it would have been to shorten the process, thus (10) is (7) - (6) which is (3) + (1) - 2(2); similarly (11) is (4) + (2) - 2(3);  $\therefore$  (13) is (4) + 3(2) - 3(3) - (1), &c.

A general systematic arrangement of the elimination-method will be given in Part II. For two or three simultaneous equations it may be stated as follows.

$$a_1x + b_1y + c_1 = 0$$

$$a_2x + b_2y + c_2 = 0.$$

Arrange the coefficients thus—

$$\begin{array}{cccc} a_1 & b_1 & c_1 & a_1 \\ a_2 & b_2 & c_2 & a_2 \end{array}$$

Form their products diagonally from left to right downwards, thus—

$$a_1b_2 \quad b_1c_2 \quad c_1a_2.$$

Form their products diagonally from right to left downwards, thus—

$$b_1a_2 \quad c_1b_2 \quad a_1c_2.$$

Subtract the latter products in order from the former, thus—

$$a_1b_2 - b_1a_2, \quad b_1c_2 - c_1b_2, \quad c_1a_2 - a_1c_2.$$

Divide the 2<sup>o</sup> and 3<sup>o</sup> remainders by the 1<sup>o</sup> remainder, the first quotient will be the value of  $x$ , the second quotient will be the value of  $y$ .

[Writing  $R_1, R_2, R_3$  for the three 'remainders' respectively, the general result is  $(mx + ny)R_1 = mR_2 + nR_3$ ].

Ex. 1. Solve

$$11x + 5y - 68 = 0$$

$$6x - 7y + 31 = 0$$

$$\begin{array}{cccc} 11 & 5 & -68 & 11 \\ 0 & -7 & 31 & 6 \end{array}$$

$$\hline -77 \quad 155 \quad -408$$

$$30 \quad 476 \quad 341$$

$$\hline -107 \quad -321 \quad -749$$

$$3 \quad 7$$


$$\parallel \quad \parallel$$



$$x \quad y.$$

olving  
compar-  
into the  
y multi-





be done  
numbered,  
t.

15 (1)  
57 (2)  
179 (3)  
453 (4)  
975 (5)  
42 (6)  
122 (7)  
274 (8)  
522 (9)  
80 (10)  
152 (11)  
248 (12)  
72 (13)  
96 (14)  
24 (15)  
1 (16)  
2 (17)  
3 (18)  
4 (19)  
5 (20)





**IMAGE EVALUATION  
TEST TARGET (MT-3)**



**Photographic  
Sciences  
Corporation**

23 WEST MAIN STREET  
WEBSTER, N.Y. 14580  
(716) 872-4503

15  
16  
17  
18  
19  
20  
22  
25

10  
11  
12  
13  
14

$$\text{Ex. 2. } \frac{12}{x} - \frac{25}{y} = 1.$$

$$\frac{22}{x} + \frac{30}{y} = 17.$$

$$\begin{array}{r} 12 \quad -25 \quad -1 \quad 12 \\ 22 \quad 30 \quad -17 \quad 22 \\ \hline 360 \quad 425 \quad -22 \\ -550 \quad -30 \quad -204 \\ \hline 910) \quad 455 \quad 182 \\ \hline \frac{1}{2} \quad \frac{1}{5} \\ \parallel \quad \parallel \\ \frac{1}{x} \quad \frac{1}{y} \end{array}$$

$$\therefore x=2 \text{ and } y=5.$$

2° Let the equations be

$$a_1x + b_1y + c_1z + d_1 = 0$$

$$a_2x + b_2y + c_2z + d_2 = 0$$

$$a_3x + b_3y + c_3z + d_3 = 0.$$

Arrange the coefficients thus

$$\begin{array}{cccccc} a_1 & b_1 & c_1 & -d_1 & -a_1 & -b_1 \\ a_2 & b_2 & c_2 & -d_2 & -a_2 & -b_2 \\ a_3 & b_3 & c_3 & -d_3 & -a_3 & -b_3 \\ a_1 & b_1 & c_1 & -d_1 & -a_1 & -b_1 \\ a_2 & b_2 & c_2 & -d_2 & -a_2 & -b_2. \end{array}$$

Selecting the first three columns form the diagonal products from left to right downwards, thus :

$$\begin{array}{ccc}
 a_1 & b_1 & c_1 \\
 & \diagdown & / \\
 a_2 & b_2 & c_2 \\
 & \diagdown & / \\
 a_3 & b_3 & c_3 \\
 & \diagdown & / \\
 a_1 & b_1 & c_1 \\
 & \diagdown & / \\
 a_2 & b_2 & c_2
 \end{array}
 \quad \text{giving } a_1 b_2 c_3$$

$$\begin{array}{ccc}
 & & a_2 b_3 c_1 \\
 & & a_3 b_1 c_2
 \end{array}$$

Form the diagonal products from right to left downwards, thus:

$$\begin{array}{ccc}
 a_1 & b_1 & c_1 \\
 & / & \diagdown \\
 a_2 & b_2 & c_2 \\
 & / & \diagdown \\
 a_3 & b_3 & c_3 \\
 & / & \diagdown \\
 a_1 & b_1 & c_1 \\
 & / & \diagdown \\
 a_2 & b_2 & c_2
 \end{array}
 \quad \text{giving } c_1 b_2 a_3$$

$$\begin{array}{ccc}
 & & c_2 b_3 a_1 \\
 & & c_3 b_1 a_2
 \end{array}$$

From the sum of the former products take the sum of the latter products obtaining a remainder, which call  $R_1$ .

Similarly form a 2<sup>o</sup> remainder,  $R_2$  from the 2<sup>o</sup>, 3<sup>o</sup> and 4<sup>o</sup> columns

$$\begin{array}{ccccccc}
 a \ 3^\circ & & " & R_3 & & " & 3^\circ, 4^\circ \text{ and } 5^\circ & " \\
 a \ 4^\circ & & " & R_4 & & " & 4^\circ, 5^\circ \text{ and } 6^\circ & "
 \end{array}$$

Then  $x = R_2 \div R_1$ ,  $y = R_3 \div R_1$ ,  $z = R_4 \div R_1$ ,  
and generally

$$(mx + ny + pz)R_1 = mR_2 + nR_3 + pR_4.$$

Ex. 3.  $3x + 2y - 4z + 20 = 0$

$5x - 7y - 6z - 1 = 0$

$7x + 5y + 5z - 24 = 0.$

3	2	-4	-20	-3	-2	
5	-7	-6	1	-5	7	
7	5	5	24	-7	-5	
3	2	-4	-20	-3	-2	
5	-7	-6	1	-5	7	
-105	-288	28	-500	$(3 \times -7 \times 5 = -105$		17.
-100	700	432	14	$5 \times 5 \times -4 = -100$		
-84	-20	500	-504	$7 \times 2 \times -6 = -84, \&c.)$		19.
-289	392	960	-990			
196	600	-15	240	$(-4 \times -7 \times 7 = 196$		20.
-90	10	480	980	$-6 \times 5 \times 3 = -90$		
50	672	-840	15	$5 \times 2 \times 5 = 50, \&c.)$		
156	1282	-375	1235			
-445)	-890	1335	-2225			22.
	2	-3	5			
	x	y	z			

Exercise Ivii.

Solve the following systems of equations :

- |   |   |
|---|---|
| <p>1. <math>2x + 3y = 41</math><br/><math>3x + 2y = 39</math></p> <p>3. <math>11x + 12y = 100</math><br/><math>9x + 8y = 80.</math></p> <p>5. <math>3x + 7y = 7</math><br/><math>5x + 3y = -36.</math></p> <p>7. <math>5x + 3y + 2 = 0</math><br/><math>3x + 2y + 1 = 0</math></p> <p>9. <math>10x + 7y + 4 = 0</math><br/><math>6x + 5y + 2 = 0.</math></p> <p>11. <math>\frac{1}{3}x + \frac{1}{4}y = 6.</math><br/><math>3x - 4y = 4.</math></p> | <p>2. <math>5x + 7y = 17</math><br/><math>7x - 5y = 9.</math></p> <p>4. <math>18x - 35y + 13 = 0</math><br/><math>15x + 28y - 275 = 0.</math></p> <p>6. <math>3x + 16y - 5 = 0</math><br/><math>28y = 5x + 19.</math></p> <p>8. <math>21x + 8y + 66 = 0</math><br/><math>23y - 28x + 13 = 0.</math></p> <p>10. <math>23x + 15y - 4\frac{1}{2} = 0</math><br/><math>32x + 21y - 6 = 0.</math></p> <p>12. <math>\frac{1}{2}x - \frac{1}{3}y = 1.</math><br/><math>\frac{1}{4}x - \frac{2}{3}y + 5 = 0.</math></p> |
|---|---|

13.  $\frac{1}{3}y = \frac{1}{2}x - 1.$

$\frac{1}{4}y = \frac{2}{5}x - 1.$

15.  $1.5x - 2y = 1.$

$2.5x - 3y = 6.$

17.  $5x - 4y + 1 = 0.$

$1.7x - 2.2y + 7.9 = 0.$

19.  $3.5x + 2\frac{1}{3}y = 13 + 4\frac{1}{2}x - 3.5y.$

$2\frac{1}{2}x + .8y = 22\frac{1}{2} + .7x - 3\frac{1}{3}y.$

20.  $\frac{1}{x} + \frac{1}{y} = \frac{5}{6}.$

$\frac{1}{x} - \frac{1}{y} = \frac{1}{6}.$

22.  $\frac{1.6}{x} = \frac{2.7}{y} - 1.$

$\frac{.8}{x} + \frac{3.6}{y} = 5.$

24.  $\frac{x}{3} + \frac{5}{y} = 4\frac{1}{3}.$

$\frac{x}{6} + \frac{10}{y} = 2\frac{2}{3}.$

26.  $\frac{3}{4}x - \frac{1}{2}(y+1) = 1.$

$\frac{1}{2}(x+1) + \frac{3}{4}(y-1) = 9.$

28.  $\frac{1}{3x+1} = \frac{2}{5y+4}.$

$\frac{1}{4x-3} = \frac{2}{7y-6}.$

30.  $\frac{15x+1}{45-y} = 8.$

$\frac{12y+19}{x-10} = 25.$

14.  $\frac{2}{3}x + \frac{3}{2}y = 17.$

$\frac{3}{4}x + \frac{2}{3}y = 19.$

16.  $7x = 10y + .1.$

$11x = 16y + .1.$

18.  $.16x - .04y = 1.$

$.19x - .11y = 1.$

21.  $\frac{3}{x} + \frac{8}{y} = 3.$

$\frac{15}{x} - \frac{4}{y} = 4.$

23.  $17x - \frac{.3}{y} = 3.$

$16x - \frac{.4}{y} = 2.$

25.  $\frac{5x}{.7} + \frac{.3}{y} = 6.$

$\frac{10x}{7} + \frac{9}{y} = 31.$

27.  $\frac{5}{x+2y} = \frac{7}{2x+y}.$

$\frac{7}{3x-2} = \frac{5}{6-y}.$

29.  $\frac{x+3y}{x-y} = 8.$

$\frac{7x-13}{3y-5} = 4.$

31.  $\frac{3x+1}{4-2y} = \frac{4}{3}.$

$x+y=1.$

-105

-100

-81, &amp;c)

= 196

-90

50, &amp;c.)

0.

).

)



$$\frac{7-2x}{5-3y} = \frac{3}{2}.$$

$$\frac{7-2y}{5-3x} = \frac{2}{3}.$$

$$34. \frac{x+3y+13}{4x+5y-2\cdot5} = 30.$$

$$\frac{8x+1y+6}{5x+3y-23} = \frac{1}{2}.$$

$$36. \frac{2x-y+3}{3} - \frac{x-2y+3}{4} = 4.$$

$$\frac{3x-4y+3}{4} + \frac{4x-2y-9}{3} = 4.$$

$$37. 20(x+1) = 15(y+1) = 12(x+y).$$

$$38. (x-2) : (y+1) : (x+y-3) :: 3 : 4 : 5.$$

$$39. (x-5) : (y+9) : (x+y+4) :: 1 : 2 : 3.$$

$$40. \frac{x+3}{x+1} = \frac{y+8}{y+5}.$$

$$\frac{2x-3}{2(y+1)} = \frac{5x-6}{5y+7}.$$

$$42. (x-1)(5y-3) = 3(3x+1).$$

$$(x-1)(4y+3) = 3(7x-1).$$

$$44. (3x-2)(5y+1) = (5x-1)(y+2).$$

$$(3x-1)(y+5) = (x+5)(7y-1).$$

$$45. x+y=37.$$

$$y+z=25.$$

$$z+x=22.$$

$$47. 1\cdot3x-1\cdot9y=1.$$

$$1\cdot7y-1\cdot1z=2.$$

$$2\cdot9z-2\cdot1x=3.$$

$$49. \frac{1}{5}x - \frac{1}{2}y = 0.$$

$$\frac{1}{3}x - \frac{1}{2}z = 1.$$

$$\frac{1}{2}z - \frac{1}{3}y = 2.$$

$$33. \frac{x+2y+1}{2x-y+1} = 2.$$

$$\frac{3x-y+1}{x-y+3} = 5.$$

$$35. \frac{x+1}{3} - \frac{y+2}{4} = \frac{2(x-y)}{5}$$

$$\frac{x-3}{4} - \frac{y-3}{3} = 2y-x.$$

$$41. (x-4)(y+7) = (x-3)(y+4).$$

$$(x+5)(y-2) = (x+2)(y-1).$$

$$43. (x+1)(2y+1) = 5x+9y+1.$$

$$(x+2)(3y+1) = 9x+13y+2.$$

$$46. 2x+2y=7.$$

$$7x+9z=29.$$

$$y+8z=17.$$

$$48. 5x+3y+2z=217.$$

$$5x-3y=39.$$

$$3y-2z=20.$$

$$50. 1\frac{1}{3}x+1\frac{1}{2}y=10.$$

$$2\frac{2}{3}x+2\frac{2}{3}z=20.$$

$$3\frac{1}{4}y+3\frac{3}{5}z=30.$$

51.

53.

55.

57.

59.

61.

63.

65.

67.

51.  $x+y-z=17.$   
 $y+z-x=13.$   
 $z+x-y=7.$
52.  $x+y+z=9,$   
 $x+2y+4z=15.$   
 $x+3y+9z=23.$
53.  $x+y+z=3.$   
 $2x+4y+8z=13.$   
 $3x+9y+27z=34.$
54.  $7x+6y+7z=100,$   
 $x-2y+z=6.$   
 $3x+y-2z=0.$
55.  $3x+2y+3z=110.$   
 $5x+y-4z=0.$   
 $2x-3y+z=0.$
56.  $x+y+z=9.$   
 $x+2y+3z=14.$   
 $x+3y+6z=20.$
57.  $x+2y+3z=32.$   
 $2x+3y+z=42.$   
 $3x+y+2z=40.$
58.  $x+y+2z=34.$   
 $x+2y+z=33.$   
 $2x+y+z=32.$
59.  $3x+3y+z=17.$   
 $3x+y+3z=15.$   
 $x+3y+3z=13.$
60.  $x+2y-z=4\cdot6.$   
 $y+2z-x=10\cdot1.$   
 $z+2x-y=5\cdot7.$
61.  $x+2y-7z=21.$   
 $3x+2y-z=24.$   
 $9x+7y-2z=27.$
62.  $x+y=1\frac{1}{2}z+8.$   
 $y+z=2\frac{2}{3}y-14.$   
 $z+x=3\frac{3}{4}x-32.$
63.  $\frac{1}{2}x+\frac{1}{3}y+\frac{1}{4}z=36\frac{1}{2}.$   
 $\frac{1}{3}x+\frac{1}{4}y+\frac{1}{5}z=27.$   
 $\frac{1}{5}x+\frac{1}{6}y+\frac{1}{7}z=18.$
64.  $2\frac{1}{2}x+3\frac{1}{3}y+4\frac{1}{4}z=140,$   
 $3\frac{1}{3}x+4\frac{1}{4}y+5\frac{1}{5}z=175.$   
 $2\frac{2}{3}x+3\frac{3}{4}y+4\frac{4}{5}z=157.$
65.  $\frac{x+1}{y+1} = 2.$   
 $\frac{y+2}{z+1} = 4.$   
 $\frac{z+3}{x+1} = \frac{1}{2}.$
66.  $\frac{3x+y}{z+1} = 2.$   
 $\frac{3y+z}{x+1} = 2.$   
 $\frac{3z+x}{y+1} = 2.$
67.  $\frac{x+y}{y-z} = 10.$   
 $\frac{x+z}{x-y} = 9.$   
 $\frac{y+z}{x+5} = 1.$
68.  $\frac{x+3}{y+z} = 2.$   
 $\frac{y+3}{x+z} = 1.$   
 $\frac{z+3}{x+y} = \frac{1}{2}.$

$$\frac{2(x-y)}{5}$$

$$2y-x.$$

$$-3)(y+4).$$

$$+2)(y-1).$$

$$x+9y+1.$$

$$x+13y+2.$$

7.

9.

0.

$$69. \quad \frac{4}{x} - \frac{3}{y} = 1.$$

$$\frac{2}{x} + \frac{3}{z} = 4.$$

$$\frac{3}{y} - \frac{1}{z} = 0.$$

$$71. \quad \frac{xy}{x+y} = \frac{1}{5}.$$

$$\frac{yz}{y+z} = \frac{1}{6}.$$

$$\frac{zx}{z+x} = \frac{1}{7}.$$

$$73. \quad (x+2)(2y+1) = (2x+7)y.$$

$$(x-2)(3z-1) = (x+3)(3z-1).$$

$$(y+1)(z+2) = (y+3)(z+1).$$

$$74. \quad (2x-1)(y+1) = 2(x+1)(y-1).$$

$$(x+4)(z+1) = (x+2)(z+2).$$

$$(y-2)(z+3) = (y-1)(z+1).$$

$$75. \quad (x+1)(5y-3) = (7x+1)(2y-3).$$

$$(4x-1)(z+1) = (x+1)(2z-1).$$

$$(y+3)(z+2) = (5y-6)(3z-1).$$

$$76. \quad 21x+31y+42z = 115.$$

$$6(2x+y) = 3(3x+z) = 2(y+z)$$

$$77. \quad 15(x-2y) = 5(2x-3z) = 3(y+z).$$

$$21x+31y+41z = 135.$$

$$78. \quad 6x(y+z) = 4y(z+x) = 3z(x+y).$$

$$\frac{1}{x} + \frac{1}{y} + \frac{1}{z} = 9.$$

$$79. \quad 3x+y+z = 20.$$

$$3u+x+4y = 30.$$

$$3u+6x+z = 40.$$

$$5u+8y+3z = 50$$

$$70. \quad \frac{6}{x} + \frac{4}{y} + \frac{5}{z} = 4.$$

$$\frac{3}{x} + \frac{8}{y} + \frac{5}{z} = 4.$$

$$\frac{9}{x} + \frac{12}{y} - \frac{10}{z} = 4.$$

$$72. \quad \frac{xy}{4y-3x} = 20.$$

$$\frac{xz}{2x-3z} = 15.$$

$$\frac{yz}{4y-5z} = 12.$$

81.

38.

85.

A  
the s  
whic  
invol  
wher  
mine

1.

Mu

Div

He

$$\begin{aligned} 81. \quad & u+x+y+z=144. \\ & u+2x+2y+2z=267. \\ & u+2x+3y+3z=359. \\ & u+2x+3y+4z=410. \end{aligned}$$

$$\begin{aligned} 82. \quad & u+x+y+z=24. \\ & u+2x+3y-9z=0. \\ & 3u-x-5y+z=0. \\ & 2u+3x-4y-5z=0. \end{aligned}$$

$$\begin{aligned} 83. \quad & u+x+y+z=60. \\ & u+2x+3y+4z=100. \\ & u+3x+6y+10z=150. \\ & u+4x+10y+20z=210. \end{aligned}$$

$$\begin{aligned} 84. \quad & u+x+y+z=1. \\ & 2u+4x+8y+16z=5. \\ & 3u+9x+27y+81z=15. \\ & 4u+16x+64y+256z=35. \end{aligned}$$

$$\begin{aligned} 85. \quad & \frac{1}{2}x + \frac{1}{4}y - \frac{1}{3}z = 1. \\ & \frac{1}{3}x - \frac{1}{4}y - \frac{1}{9}u = 1. \\ & \frac{1}{6}x + \frac{2}{3}z - \frac{1}{2}u = 1. \\ & \frac{3}{4}y - \frac{1}{2}z - \frac{1}{3}u = 0. \end{aligned}$$

$$\begin{aligned} 86. \quad & \frac{1}{2}u - \frac{1}{3}x + \frac{1}{5}y - \frac{1}{7}z = 47. \\ & \frac{1}{3}u + \frac{1}{2}x + \frac{1}{7}y - \frac{1}{2}z = 37. \\ & \frac{1}{5}u - \frac{2}{7}x + \frac{1}{2}y - \frac{1}{3}z = 17. \\ & \frac{5}{7}u - \frac{1}{2}x - \frac{1}{3}y + \frac{1}{2}z = 17. \end{aligned}$$

Art. XLVII. The principle of symmetry is often of use in the solution of symmetrical equations. For from one relation which may be found to exist between two or more of the letters involved, other relations may be derived by symmetry; also, when the value of one of the unknown quantities has been determined, the values of the others can be at once written down, &c.

$$\begin{aligned} 1. \quad & (x+y)(x+z) = a. \\ & (x+y)(y+z) = b. \\ & (x+z)(y+z) = c. \end{aligned}$$

Multiply the equations together and extract the square root.

$$\therefore (x+y)(y+z)(z+x) = \sqrt{(abc)}.$$

Divide this equation by the third.

$$\therefore x+y = \frac{\sqrt{(abc)}}{c}, \text{ and therefore, by symmetry,}$$

$$\therefore y+z = \frac{\sqrt{(abc)}}{a},$$

$$\therefore z+x = \frac{\sqrt{(abc)}}{b}.$$

Hence we get

$$x = \frac{ab - bc + ca}{2\sqrt{(abc)}},$$

whence  $y$  and  $z$  may be derived by symmetry.

$$2. \quad \begin{aligned} x+y+z &= 0 \dots\dots\dots (1). \\ ax+by+cz &= 0 \dots\dots\dots (2). \\ bx+cy+az+(a-b)(b-c)(c-a) &= 0 \dots\dots\dots (3). \end{aligned}$$

$$c \times (1) - (2) \text{ gives } (c-a)x + (c-b)y = 0.$$

$$\therefore y = \frac{(c-a)x}{b-c}, \text{ and similarly,}$$

$$z = \frac{(a-b)x}{b-c}.$$

Substitute in (3) these values of  $y$  and  $z$ , and reduce,

$$\therefore x(a-b)(c-a) = (a-b)(b-c)(c-a),$$

$$\therefore \text{ or } x = (b-c), \therefore y = c-a, z = a-b.$$

$$8. \quad a(yz - zx - xy) = b(zx - xy - yz) = c(xy - yz - zx) = xyz.$$

Divide the first and the last equations by  $axyz$ ;

$$\therefore \frac{1}{a} = \frac{1}{x} - \frac{1}{y} - \frac{1}{z}, \text{ and hence, by symmetry,}$$

$$\frac{1}{b} = \frac{1}{y} - \frac{1}{z} - \frac{1}{x},$$

$$\frac{1}{c} = \frac{1}{z} - \frac{1}{x} - \frac{1}{y}.$$

$$\therefore \frac{1}{b} + \frac{1}{c} = -\frac{2}{x}, \text{ and by symmetry,}$$

$$\frac{1}{c} + \frac{1}{a} = -\frac{2}{y},$$

$$\frac{1}{a} + \frac{1}{b} = -\frac{2}{z}.$$

$$4. \quad \begin{aligned} ax+by+cz &= 1 \dots\dots\dots (1) \\ a^2x+b^2y+c^2z &= 1 \dots\dots\dots (2) \\ a^3x+b^3y+c^3z &= 1 \dots\dots\dots (3) \end{aligned}$$

$$c \times (1) - (2) \text{ gives } a(c-a)x + b(c-b)y = c-1 \dots\dots\dots (4).$$

$$c \times (2) - (3) \text{ " } a^2(c-a)x + b^2(c-b)y = c-1 \dots\dots\dots (5).$$

$$b \times (4) - (5) \text{ " } ab(c-a)x - a^2(c-a)x = b(c-1) - (c-1),$$

$$\text{ or } a(a-b)(a-c)x = (c-1)(b-1),$$

$$\therefore x = \frac{(1-b)(1-c)}{a(a-b)(a-c)};$$

whence  $y$  and  $z$  may be derived by symmetry.

5. Eliminate  $x, y, z, u$  (which are supposed all different) from the following equations :

$$x = by + cz + du.$$

$$y = cz + du + ax.$$

$$z = du + ax + by.$$

$$u = ax + by + cz.$$

Subtracting the second equation from the first,

$$\therefore x - y = by - ax, \text{ or}$$

$$(1+a)x = (1+b)y = (\text{by symmetry}) (1+c)z = (1+d)u.$$

These relations may be also obtained by adding  $ax$  to both members of the first equation,  $by$ , to both members of the second equation, &c.

Now divide the first equation by these equals.

$$\therefore \frac{1}{1+a} = \frac{b}{1+b} + \frac{c}{1+c} + \frac{d}{1+d}.$$

And since  $\frac{1}{1+a} = 1 - \frac{a}{1+a}$ , we have

$$1 = \frac{a}{1+a} + \frac{b}{1+b} + \frac{c}{1+c} + \frac{d}{1+d}.$$

### Exercise Iviii.

- Given  $ax + by = c$  and that  $x = \frac{b'c - bc'}{b'a - b'a'}$   
 $a'x + b'y = c'$  derive the value of  $y$
- Given  $bx = ay$  and that  $x = \frac{a(dm - cn)}{bc - ad}$   
 $dx + md = cy + nc$  derive the value of  $y$ .
- Given  $ax + by + cz = d$  and that  $x =$   
 $a^2x + b^2y + c^2z = d^2$   $\frac{a(d-b)(d-c)}{d(a-b)(a-c)}$ , write down  
 $a^3x + b^3y + c^3z = d^3$  the values of  $y$  and  $z$ .

4. There is a set of equations in  $x, y, z, u,$  and  $w,$  with corresponding coefficients ( $a$  to  $x,$  &c.),  $a, b, c, d,$  and  $e;$  one of the equations is

$$x = by + cz + du + ew, \text{ write down the others.}$$

Solve the following equations :

$$5. \frac{x}{m} + \frac{y}{n} = a, \frac{y}{n} + \frac{z}{p} = b, \frac{x}{m} + \frac{z}{p} = c.$$

$$6. x + ay + bz = m, y + az + bc = n, z + ax + by = p.$$

$$7. x + ay = l, y + bz = m, z + cu = n, u + dw = p, w + ex = r.$$

8. Eliminate  $x, y, z,$  (supposed to be all different) from the following equations :

$$x = by + cz, y = cz + ax, z = ax + by.$$

9. Eliminate  $x, y, z,$  from

$$\frac{x}{y+z} = a, \frac{y}{z+x} = b, \frac{z}{x+y} = c.$$

10. Having given

$$x = by + cz + du + ew,$$

$$y = cz + du + ew + ax,$$

$$z = du + ew + ax + by,$$

$$u = ew + ax + by + cz,$$

$$w = ax + by + cz + du,$$

$$\text{Shew that } \frac{a}{1+a} + \frac{b}{1+b} + \frac{c}{1+c} + \frac{d}{1+d} + \frac{e}{1+e} = 1.$$

Art. XLVIII. Resolution of Particular Systems of Linear Equations.

Ex. 1.

$$x + y + z = a \quad (1)$$

$$y + z + u = b \quad (2)$$

$$z + u + x = c \quad (3)$$

$$u + x + y = d \quad (4)$$

$$(1) + (2) + (3) + (4) \quad 3(u + x + y + z) = a + b + c + d \quad (5')$$

$$3(1) \quad 3(x + y + z) = 3a \quad (6')$$

$$\frac{1}{3}\{(5') - (6')\} \quad u = \frac{1}{3}(-2a + b + c + d.)$$

The values of  $x$ ,  $y$  and  $z$  may now be written down by symmetry.

The following is a variation of the above method, applicable to a much more general system.

Assume the auxiliary equation

$$u + x + y + z = s, \quad (5)$$

$$\therefore (1) \text{ becomes } s - u = a, \quad (6)$$

$$(2) \quad " \quad s - x = b, \quad (7)$$

$$(3) \quad " \quad s - y = c, \quad (8)$$

$$(4) \quad " \quad s - z = d, \quad (9)$$

$$(5) + (6) + (7) + (8) + (9) \quad 4s = s + a + b + c + d.$$

$$\therefore s = \frac{1}{3}(a + b + c + d).$$

$s$  is now a known quantity, and may be treated as such,

$$\text{in (6) giving } u = s - a$$

$$" (7) \quad " \quad x = s - b$$

$$" (8) \quad " \quad y = s - c$$

$$" (9) \quad " \quad z = s - d.$$

$$\text{Ex. 2. } \quad yz = a(y + z), \quad (1)$$

$$zcx = b(z + x), \quad (2)$$

$$xy = c(x + y), \quad (3)$$

$$(1) \div ayz, \quad \frac{1}{y} + \frac{1}{z} = \frac{1}{a},$$

$$(2) \div bzx, \quad \frac{1}{z} + \frac{1}{x} = \frac{1}{b},$$

$$(3) \div cxy, \quad \frac{1}{x} + \frac{1}{y} = \frac{1}{c}.$$

This may now be solved like Ex. 1, using the reciprocals of  $a$ ,  $b$ ,  $c$ ,  $x$ ,  $y$  and  $z$  instead of these quantities themselves.

$$\text{Ex. 3. } \quad a_1 u + b_1(x + y + z) = c_1 \quad (1)$$

$$a_2 x + b_2(y + z + u) = c_2 \quad (2)$$

$$a_3 y + b_3(z + u + x) = c_3 \quad (3)$$

$$a_4 z + b_4(u + x + y) = c_4 \quad (4)$$

Assume the auxiliary equation

$$u + x + y + z = s. \quad (5)$$



(1) becomes  $b_1s - (b_1 - a_1)u = c_1$

$$\therefore \frac{b_1}{b_1 - a_1}s - u = \frac{c_1}{b_1 - a_1} \quad (6)$$

Similarly from (2)  $\frac{b_2}{b_2 - a_2}s - x = \frac{c_2}{b_2 - a_2} \quad (7)$

“ “ (3)  $\frac{b_3}{b_3 - a_3}s - y = \frac{c_3}{b_3 - a_3} \quad (8)$

“ “ (4)  $\frac{b_4}{b_4 - a_4}s - z = \frac{c_4}{b_4 - a_4} \quad (9)$

$$(5) + (6) + (7) + (8) + (9) \left( \frac{b_1}{b_1 - a_1} + \frac{b_2}{b_2 - a_2} + \frac{b_3}{b_3 - a_3} + \frac{b_4}{b_4 - a_4} \right) s \\ = s + \frac{c_1}{b_1 - a_1} + \frac{c_2}{b_2 - a_2} + \frac{c_3}{b_3 - a_3} + \frac{c_4}{b_4 - a_4} \quad (10)$$

From (10) we can at once get the value of  $s$ , which may therefore be treated as a known quantity.

in (6) giving  $u = \frac{b_1s - c_1}{b_1 - a_1}$

and the value of  $x$ ,  $y$ , and  $z$  may be obtained from (7), (8) and (9), or they may be written down by symmetry.

Ex. 4.  $ax + b(y + z) = c \quad (1)$

$$ay + b(z + u) = d \quad (2)$$

$$az + b(u + x) = e \quad (3)$$

$$au + b(x + y) = f \quad (4)$$

Assume  $u + x + y + z = s \quad (5)$

(1) + (2) + (3) + (4)  $(a + 2b)s = c + d + e + f \quad (6)$

Hence  $s$  is a known quantity and may be treated as such.

From (1) and (5)  $bs - bu + (a - b)x = c,$

$$\therefore bu - (a - b)x = bs - c, \quad (7)$$

Similarly from (2) and (5)  $bx - (a - b)y = bs - d, \quad (8)$

“ “ (3) “ “  $by - (a - b)z = bs - e, \quad (9)$

“ “ (4) “ “  $bz - (a - b)u = bs - f, \quad (10)$

$b(7) + (a - b)(8) \quad b^2u - (a - b)^2y = abs - bz - (a - b)d, \quad (11)$

$b(9) + (a - b)(10) \quad b^2y - (a - b)^2u = abs - bz - (a - b)f, \quad (12)$

$$b^2(11) + (a-b)^2(12) \quad \{b^4 - (a-b)^4\}u = abs\{b^2 + (a-b)^2\} \\ - a\{b^2d + (a-b)^2f\} - b\{b^2(c-d) + (a-b)^2(e-f)\} \quad (13)$$

(6) The values of  $x$ ,  $y$ , and  $z$  may now be written down by symmetry.

(7) Ex. 5.  $a^3 + a^2x + ay + z = 0,$

$$b^3 + b^2x + by + z = 0,$$

(8)  $c^3 + c^2x + cy + z = 0.$

The polynome  $t^3 + xt^2 + yt + z$  vanishes for  $t=a$ ,  $t=b$ ,  $t=c$ ,

(9)  $\therefore$  by Th. II., p. 46, for all values of  $t$ .

$$t^3 + xt^2 + yt + z = (t-a)(t-b)(t-c)$$

$$= t^3 - (a+b+c)t^2 + (ab+bc+ca)t - abc.$$

$\therefore$  Th. III., p. 53,  $x = -(a+b+c),$

$$y = ab + bc + ca,$$

$$z = -abc.$$

Ex. 6.  $x + y + z + u = 1,$  (1)

$$cx + by + cz + du = 0, \quad (2)$$

$$a^2x + b^2y + c^2z + d^2u = 0, \quad (3)$$

$$a^3x + b^3y + c^3z + d^3u = 0. \quad (4)$$

Employing the method of arbitrary multipliers,

$$(4) + l(3) + m(2) + n(1) \quad \begin{array}{cccc} a^3x + b^3y + c^3z + d^3u = n \\ +la^2x + lb^2y + lc^2z + ld^2u \\ +max + mby + mcz + md \\ +nx + ny + nz + nu \end{array} \quad (5)$$

$$\begin{array}{cccc} +la^2 & +lb^2 & +lc^2 & +ld^2 \\ +ma & +mb & +mc & +md \\ +n & +n & +n & +n \end{array}$$

To determine  $x$  assume

$$b^3 + lb^2 + mb + n = 0, \quad (6)$$

$$c^3 + lc^2 + mc + n = 0, \quad (7)$$

$$d^3 + ld^2 + md + n = 0, \quad (8)$$

$$\therefore x = \frac{n}{a^3 + la^2 + ma + n} \quad (9)$$

But the system (6), (7), (8) has been solved in Ex. 5, from which it is seen that

$$l = -(b+c+d), \quad m = bc + cd + db, \quad n = -bcd,$$

and  $a^3 + a^2l + am + n = (a-b)(a-c)(a-d):$

∴ using these values in (9)

$$x = \frac{-bcd}{(a-b)(a-c)(a-d)}$$

The values of  $y$ ,  $z$  and  $u$  may now be written down by symmetry.

Ex. 7. 
$$\frac{x}{m-a} + \frac{y}{m-b} + \frac{z}{m-c} = 1. \quad (1)$$

$$\frac{x}{n-a} + \frac{y}{n-b} + \frac{z}{n-c} = 1. \quad (2)$$

$$\frac{x}{p-a} + \frac{y}{p-b} + \frac{z}{p-c} = 1. \quad (3)$$

Assume 1 - 
$$\frac{x}{t-a} - \frac{y}{t-b} - \frac{z}{t-c} = \frac{t^3 + Bt^2 + Ct + D}{(t-a)(t-b)(t-c)} \quad (4)$$

But in virtue of equations (1), (2) and (3), the first member of (4) vanishes for  $t=m$ ,  $t=n$ , and  $t=p$ , and ∴  $t^3 + Bt^2 + Ct + D$  vanishes for the same values of  $t$ , and ∴ Th. II. p. 46,

$$t^3 + Bt^2 + Ct + D = (t-m)(t-n)(t-p),$$

∴ (4) becomes 
$$1 - \frac{x}{t-a} - \frac{y}{t-b} - \frac{z}{t-c} = \frac{(t-m)(t-n)(t-p)}{(t-a)(t-b)(t-c)}$$

To obtain the value of  $x$  multiply both sides of this equation by  $(t-a)$ ,

$$t-a-x - \frac{y(t-a)}{t-b} - \frac{z(t-a)}{t-c} = \frac{(t-m)(t-n)(t-p)}{(t-b)(t-c)}$$

Now  $t$  may have any value in this equation; let  $t=a$ .

$$\therefore x = \frac{(a-m)(a-n)(a-p)}{(a-b)(a-c)}$$

The substitution  $(xyz|abc)$  will give the values of  $y$  and  $z$ .

Ex. 8. 
$$\frac{x+a}{p} = \frac{y+b}{q} = \frac{z+c}{r} \quad (1)$$

$$lx + my + nz = s^2 \quad (2)$$

By Art. XXXVII.,

$$\frac{x+a}{v} = \frac{y+b}{q} = \frac{z+c}{r} = \frac{lx+my+nz+la+mb+nc}{lp+mq+nr}$$

$$(2) \quad = \frac{s^2+la+mb+nc}{lp+mq+nr} = R, \text{ say}$$

$$\therefore x = pR - a, \quad y = qR - b, \quad z = rR - c.$$

$$\text{Ex. 9.} \quad yz + zx + xy = (a+b+c)xyz \quad (1)$$

$$\frac{yz+zx}{a} = \frac{zx+xy}{b} = \frac{xy+yz}{c} \quad (2)$$

$$(1) \div xyz \quad \frac{1}{x} + \frac{1}{y} + \frac{1}{z} = a+b+c \quad (3)$$

$$(2) \div xyz \quad \frac{\frac{1}{x} + \frac{1}{y}}{a} = \frac{\frac{1}{y} + \frac{1}{z}}{b} = \frac{\frac{1}{z} + \frac{1}{x}}{c} \quad (4)$$

$$\text{Page 122 and (3)} \quad = \frac{\frac{2}{x} + \frac{2}{y} + \frac{2}{z}}{a+b+c} = 2 \quad (5)$$

$$(4) \text{ and } (5) \quad \therefore \frac{1}{x} + \frac{1}{y} = 2a, \quad \frac{1}{y} + \frac{1}{z} = 2b, \\ \frac{1}{z} + \frac{1}{x} = 2c. \quad (6)$$

$$(3) - (6) \quad \frac{1}{x} = a-b+c, \quad \frac{1}{y} = a+b-c, \quad \frac{1}{z} = -a+b+c.$$

$$\text{Ex. 10.} \quad \frac{x+c}{a+b} + \frac{y+b}{a+c} = 2. \quad (1)$$

$$\frac{x-b}{a-c} + \frac{y-c}{a-b} = 2. \quad (2)$$

$$(1) \quad \therefore \frac{x+c}{a+b} - 1 = 1 - \frac{y+b}{a+c}$$

$$\therefore \frac{x-a-b+c}{a+b} = \frac{a+c-b-y}{a+c} \quad (3)$$

Similarly from (2)  $\frac{x-a-b+c}{a-c} = \frac{a-b+c-y}{a-b}$  (4)

(3) and (4)  $\therefore x-a-b+c = \frac{a+b}{a+c}(a-b+c-y)$   
 $= \frac{a-c}{a-b}(a-b+c-y).$

But unless  $\frac{a+b}{a+c} = \frac{a-c}{a-b}$ , this cannot be the case except for

$$a-b+c-y=0,$$

in which case  $x-a-b+c=0$  also,

giving  $x=a+b-c$  and  $y=a-b+c$ . (5)

If  $\frac{a+b}{a+c} = \frac{a-c}{a-b} \therefore a^2-b^2=a^2-c^2$  (6)  
 $b^2-c^2=0$ , or  $(b+c)(b-c)=0$ ,

$$\therefore b=c, \text{ or } b=-c.$$

But if  $b=+c$  or  $-c$ , (1) and (2) are one and the same equation; hence if (1) and (2) are independent, (6) cannot be true, thus leaving only the alternative (5).

Ex. 11.  $2ax=(b+c-a)(y+z)$ , (1)

$$2by=(c+a-b)(z+x),$$
 (2)

$$(x+y+z)^2+x^2+y^2+z^2=4(a^2+b^2+c^2)$$
 (3)

(1) and page 122 (5)  $\frac{x}{b+c-a} = \frac{y+z}{2a} = \frac{x+y+z}{b+c+a}$  (4)

(2) " "  $\frac{y}{c+a-b} = \frac{x+z}{2b} = \frac{x+y+z}{c+a+b}$  (5)

(4), (5) and "  $\therefore \frac{x+y+z}{a+b+c} = \frac{x}{b+c-a} = \frac{y}{c+a-b} = \frac{z}{a+b-c}$ .

$$\therefore \frac{x^2}{(b+c-a)^2} = \frac{(x+y+z)^2+x^2+y^2+z^2}{(a+b+c)^2+(b+c-a)^2+(c+a-b)^2+(a+b-c)^2}$$

Reduction and (3)  $= \frac{(x+y+z)^2+x^2+y^2+z^2}{4(a^2+b^2+c^2)} = 1$ .

$$\therefore x^2=(b+c-a)^2.$$

(4) Ex. 12.  $ax = by = cz = \frac{1}{x} + \frac{1}{y} + \frac{1}{z}$  (1)

(1)  $\div xyz \quad \therefore \frac{a}{yz} = \frac{b}{zx} = \frac{c}{xy} \therefore = \frac{a+b+c}{xy+yz+zx}$  (2)

Also from (1)  $\div xyz \quad \frac{a}{yz} = \frac{1}{xyz} \left( \frac{1}{x} + \frac{1}{y} + \frac{1}{z} \right) = \frac{xy+yz+zx}{x^2y^2z^2}$  (3)

(2)  $\times$  (3)  $\frac{a^2}{y^2z^2} = \frac{a+b+c}{x^2y^2z^2}$

$\therefore a^2x^2 = a+b+c.$

(5) Ex. 13.  $\frac{y+z-x}{a} = \frac{z+x-y}{b} = \frac{x+y-z}{c}$  (1)

$xyz = m^3$  (2)

(1)  $\frac{z}{a+b} = \frac{x}{b+c} = \frac{y}{c+a} = \frac{m}{r}$  suppose (3)

then  $\frac{xyz}{(a+b)(b+c)(c+a)} = \frac{m^3}{r^3}$

$\therefore r^3 = (a+b)(b+c)(c+a)$

Hence the value of  $r$  is known and from (3)

$rx = m(b+c).$

Ex. 14.  $y+z = 2axyz$  (1)

$z+x = 2bxyz$  (2)

$x+y = 2cxyz$  (3)

$\therefore xyz = \frac{y+z}{2a} = \frac{z+x}{2b} = \frac{x+y}{2c} = \frac{x+y+z}{a+b+c}$

$= \frac{x}{b+c-a} = \frac{y}{c+a-b} = \frac{z}{a+b-c}$  (4)

$\therefore x^3y^3z^3 = \frac{xyz}{(b+c-a)(c+a-b)(a+b-c)}$

$\therefore x^2y^2z^2 = \frac{1}{(b+c-a)(c+a-b)(a+b-c)}$

Hence the value of  $x^2y^2z^2$  is known, call it  $\frac{r}{r^2}$  and substitute in (4)

$$\frac{1}{r} = \frac{x}{b+c-a}$$

$$\therefore rx = b+c-a,$$

in which  $r^2 = (b+c-a)(c+a-b)(a+b-c)$ .

$$\text{Ex. 15.} \quad y^2+z^2-x(y+z) = a \quad (1)$$

$$z^2+x^2-y(z+x) = b \quad (2)$$

$$x^2+y^2-z(x+y) = c \quad (3)$$

$$(1)+(2)+(3) \quad 2(x^2+y^2+z^2-xy-yz-zx) = a+b+c \quad (4)$$

$$(1) \text{ may be written } \quad x^2+y^2+z^2-x(x+y+z) = a \quad (5)$$

$$(2) \quad \text{“} \quad \text{“} \quad x^2+y^2+z^2-y(x+y+z) = b \quad (6)$$

$$(3) \quad \text{“} \quad \text{“} \quad x^2+y^2+z^2-z(x+y+z) = c \quad (7)$$

$$\therefore x+y+z = \frac{a-b}{y-x} = \frac{b-c}{z-y} = \frac{c-a}{x-z}$$

$$\therefore (x+y+z)^2 = \frac{(a-b)^2 + (b-c)^2 + (c-a)^2}{(y-x)^2 + (z-y)^2 + (x-z)^2}$$

$$= \frac{a^2+b^2+c^2-ab-bc-ca}{x^2+y^2+z^2-xy-yz-zx}$$

$$(4) \quad = \frac{2(a^2+b^2+c^2-ab-bc-ca)}{a+b+c} \quad (8)$$

$$= \frac{2(a^3+b^3+c^3-3abc)}{(a+b+c)^2} \quad (9)$$

Write  $r^2$  for  $2(a^3+b^3+c^3-3abc)$ .

$$(9) \quad \therefore x+y+z = \frac{r}{a+b+c} \quad (10)$$

$$\text{Returning to (8)} \quad (x+y+z)^2 = \frac{2(a^2+b^2+c^2-ab-bc-ca)}{a+b+c} \quad (8)$$

$$(4) \quad 2(x^2+y^2+z^2-xy-yz-zx) = \frac{(a+b+c)^2}{a+b+c} \quad (11)$$

$$\frac{1}{2}\{(8)+(11)\} \quad x^2+y^2+z^2 = \frac{a^2+b^2+c^2}{a+b+c} \quad (12)$$

$$(5) \text{ and } (10) \quad x^2+y^2+z^2 - \frac{rx}{a+b+c} = a$$

$$\begin{aligned} \therefore rx &= (a+b+c)(x^2+y^2+z^2) - a(a+b+c) \\ (12) \quad &= a^2+b^2+c^2 - a(a+b+c) \\ &= b^2+c^2 - a(b+c). \end{aligned}$$

(5), (6), (7) are symmetrical with respect to  $(xyz|abc)$ ; (10) shows this substitution does not affect  $r$ , and consequently the values of  $y$  and  $z$  may be written down at once from that of  $x$ .

## Exercise lix.

$$1. \quad ax+by=c, \\ mx+ny=d.$$

$$2. \quad ax+by=c, \\ mx-ny=d.$$

$$3. \quad ax+by=c, \\ mx+ny=c.$$

$$4. \quad \frac{x}{a} + \frac{y}{b} = 1, \\ x+y=c.$$

$$5. \quad \frac{x}{a} + \frac{y}{b} = 1,$$

$$6. \quad \frac{x}{a} + \frac{y}{b} = 1,$$

$$\frac{x}{b} + \frac{y}{a} = 1,$$

$$\frac{x}{b} = \frac{y}{a}.$$

$$7. \quad ax+bc=by+ac, \\ x+y=c.$$

$$8. \quad \frac{a}{x} + \frac{b}{y} = m,$$

$$\frac{b}{x} + \frac{a}{y} = n.$$

$$9. \quad (a+c)x - (a-c)y = 2ab, \\ (a+b)y - (a-b)x = 2ac.$$

$$10. \quad \frac{x-c}{y-c} = \frac{a}{b}, \\ x-y=a-b.$$

$$11. \quad \frac{x}{y} = \frac{a}{b},$$

$$12. \quad \frac{x+y}{y+1} = \frac{a+b+c}{a-b+c},$$

$$\frac{x+m}{y+n} = \frac{c}{d}.$$

$$\frac{y-1}{x+1} = \frac{a-b-c}{a+b-c}.$$



13. 
$$\frac{x-a+c}{y-a+b} = \frac{b}{c}.$$

$$\frac{y+b}{x+c} = \frac{c+a}{b+a}.$$

15. 
$$\frac{x}{m-a} + \frac{y}{m-b} = 1,$$

$$\frac{x}{n-a} + \frac{y}{n-b} = 1.$$

17. 
$$x+y+z=l,$$
  
$$ax+by+cz=m,$$

$$\frac{x}{l-a} + \frac{y}{l-b} + \frac{z}{l-c} = 1.$$

19. 
$$\frac{x-a}{p} = \frac{y-b}{q} = \frac{z-c}{r},$$

$$lx+my+nz=1.$$

21. 
$$x+y+z=a+b+c,$$
  
$$bx+cy+az=a^2+b^2+c^2,$$
  
$$cx+ay+bz=a^2+b^2+c^2.$$

23. 
$$x+y+z=m,$$
  
$$x:y:z=a:b:c.$$

25. 
$$xy+yz+zx=0, \quad ayz+bxz+cxy=0,$$
  
$$bcyz+acxz+abxy+(a-b)(b-c)(c-a)xyz=0.$$

26. 
$$(a+b)x+(b+c)y+(c+a)z=ab+bc+ca,$$
  
$$(a+c)x+(a+b)y+(b+c)z=ab+ac+bc,$$
  
$$(b+c)x+(a+c)y+(a+b)z=a^2+b^2+c^2.$$

27. 
$$mx+ny+pz+qu=r,$$

$$\frac{x}{a} = \frac{y}{b} = \frac{z}{c} = \frac{u}{d}.$$

14. 
$$\frac{x+c}{a+b} + \frac{y+b}{a+c} = 2,$$

$$\frac{x-b}{a-c} + \frac{y-c}{a-b} = 2,$$

16. 
$$x+y+z=0,$$
  
$$(b+c)x+(a+c)y+(a+b)z=0,$$
  
$$bcx+acy+abz=1.$$

18. 
$$\frac{x-a}{p} = \frac{y-b}{q} = \frac{z-c}{r},$$

$$l(x-a)+m(y-b)+n(z-c)=1.$$

20. 
$$a(x-a)=b(y-b)=c(z-c),$$
  
$$ax+by+cz=m^2.$$

22. 
$$x+y+z=0,$$
  
$$ax+by+cz=ab+bc+ca,$$
  
$$(b-c)x+(c-a)y+(a-b)z=0.$$

24. 
$$ax+by+cz=r,$$
  
$$mx=ny, \quad qy=pz$$

28

29.

30.

32.

34.

36.

37.

$$28. \frac{x(y+z)}{a} = \frac{y(x+z)}{b} = \frac{z(x+y)}{c}, \quad z = \frac{1}{a+b+c}$$

$$\frac{1}{x} + \frac{1}{y} + \frac{1}{z} = a+b+c.$$

$$29. (a-b)(x+c) - ay + bz = (c-a)(y+b) - cz + ax = 0, \\ x+y+z = 2(a+b+c).$$

$$30. \begin{aligned} ax+by &= 1, \\ by+cz &= 1, \\ cz+ax &= 1, \end{aligned}$$

$$31. \begin{aligned} ly+mx &= n, \\ nx+lz &= m, \\ mz+ny &= l. \end{aligned}$$

$$32. \begin{aligned} x+y &= a, \\ y+z &= b, \\ x+z &= c. \end{aligned}$$

$$33. \begin{aligned} y+z-x &= \frac{mn}{l}, \\ z+x-y &= \frac{ln}{m}, \\ x+y-z &= \frac{lm}{n}. \end{aligned}$$

$$34. \begin{aligned} \frac{1}{y} + \frac{1}{z} &= 2a, \\ \frac{1}{z} + \frac{1}{x} &= 2b, \\ \frac{1}{x} + \frac{1}{y} &= 2c. \end{aligned}$$

$$35. \begin{aligned} \frac{1}{y} + \frac{1}{z} - \frac{1}{x} &= \frac{2}{a}, \\ \frac{1}{z} + \frac{1}{x} - \frac{1}{y} &= \frac{2}{b}, \\ \frac{1}{x} + \frac{1}{y} - \frac{1}{z} &= \frac{2}{c}. \end{aligned}$$

$$36. \begin{aligned} (a+b)x + (a-b)z &= 2bc, \\ (b+c)y + (b-c)x &= 2ac, \\ (c+a)z + (c-a)y &= 2ab. \end{aligned}$$

$$37. \begin{aligned} x + \frac{y}{b} - \frac{z}{c} &= a, \\ y + \frac{z}{c} - \frac{x}{a} &= b, \\ z + \frac{x}{a} - \frac{y}{b} &= c. \end{aligned} \quad 38. \begin{aligned} \frac{x}{b+c} + \frac{y}{c+a} &= b-a, \\ \frac{x}{b+c} + \frac{z}{a+b} &= a-c, \\ \frac{y}{c+a} + \frac{z}{a+b} &= c-b. \end{aligned}$$

$$\begin{aligned}
 39. \quad x+y-z &= a, \\
 y+z-v &= b, \\
 z+v-x &= c, \\
 v+x-y &= d.
 \end{aligned}$$

$$\begin{aligned}
 40. \quad u+v-x &= a, \\
 v+x-y &= b, \\
 x+y-z &= c, \\
 y+x-u &= d, \\
 z+u-v &= e.
 \end{aligned}$$

## Exercise lx.

Resolve

1.  $(a+b)x + (a-b)y = 2(a^2 + b^2)$     2.  $x+y = a,$   
 $(a-b)x + (a+b)y = 2(a^2 - b^2)$      $x^2 - y^2 = b.$
3.  $2x - 3y = m,$     4.  $(a-b)x + (a+b)y = a+b.$   
 $2x^2 - 3y^2 = n^2 + xy.$      $\frac{x}{a+b} - \frac{y}{a-b} = \frac{1}{a+b}.$
5.  $(a-b)x + y = \frac{a+b+1}{a+b},$     6.  $(a+b-c)x - (a-b+c)y$   
 $x + (a+b)y = \frac{a-b+1}{a-b}.$      $= 4a(b-c).$   
 $\frac{x}{y} = \frac{a+b-c}{a-b+c}.$
7.  $\frac{x+y}{x-y} = \frac{a}{b-c},$     8.  $\frac{x-a}{y-a} = \frac{a-b}{a+b}.$   
 $\frac{x+c}{a+b} = \frac{y+b}{a+c}.$      $\frac{x}{y} = \frac{a^3 - b^3}{a^3 + b^3}.$
9.  $\frac{x-y+1}{x-y-1} = a,$     10.  $\frac{x+y+1}{x-y+1} = \frac{a+1}{a-1},$   
 $\frac{x+y+1}{x+y-1} = b,$      $\frac{x+y+1}{x-y-1} = \frac{1+b}{1-b}.$
11.  $\frac{x-y+1}{x+y-1} = a,$     12.  $\frac{x}{a+b} + \frac{y}{a-b} = a+b,$   
 $\frac{x+y+1}{x-y-1} = b.$      $\frac{x}{a} + \frac{y}{b} = 2a.$
13.  $(a+c)x + (a-c)y = 2ab,$     14.  $a^2 + ax + y = 0,$   
 $(a+b)y - (a-b)x = 2ac.$      $b^2 + bx + y = 0.$

$$15. \begin{aligned} y+z-x &= a, \\ z+x-y &= b, \\ x+y-z &= c. \end{aligned}$$

$$16. \begin{aligned} 7x+11y+z &= a, \\ 7y+11z+x &= b, \\ 7z+11x+y &= c. \end{aligned}$$

$$17. \begin{aligned} \frac{a}{x} + \frac{b}{y} - \frac{c}{z} &= 2ab, \\ \frac{b}{y} + \frac{c}{z} - \frac{a}{x} &= 2bc, \\ \frac{c}{z} + \frac{a}{x} - \frac{b}{y} &= 2ca. \end{aligned}$$

$$18. \begin{aligned} (a-b)(x+c) - ay + bz &= 0, \\ (c-a)(y+b) - cz + ax &= 0, \\ x+y+z &= 2(a+b+c). \end{aligned}$$

$$19. \begin{aligned} \frac{x}{b+c} + \frac{y}{c-a} &= a+b, \\ \frac{y}{c+a} + \frac{z}{a-b} &= b+c, \\ \frac{z}{a+b} + \frac{x}{b-c} &= c+a. \end{aligned}$$

$$20. \begin{aligned} \frac{x}{b+c} + \frac{y}{c-a} - \frac{z}{a-b} &= 0, \\ \frac{x}{b-c} - \frac{y}{c-a} + \frac{z}{a-b} &= 0, \\ \frac{x}{b+c} + \frac{y}{c-a} + \frac{z}{a+b} &= 2a. \end{aligned}$$

$$21. \begin{aligned} \frac{x}{a} + \frac{y}{a-1} + \frac{z}{a-2} &= 1, \\ \frac{x}{b} + \frac{y}{b-1} + \frac{z}{b-2} &= 1, \\ \frac{x}{c} + \frac{y}{c-1} + \frac{z}{c-2} &= 1. \end{aligned}$$

$$22. \begin{aligned} \frac{xy}{x+y} &= a, \\ \frac{yz}{y+z} &= b, \\ \frac{zx}{z+x} &= c. \end{aligned}$$

$$23. \begin{aligned} \frac{x}{a} + \frac{y}{b} + \frac{z}{c} &= \frac{x}{b} + \frac{y}{c} + \frac{z}{a} = \\ \frac{x}{c} + \frac{y}{a} + \frac{z}{b} &= \frac{1}{a} + \frac{1}{b} + \frac{1}{c}. \end{aligned}$$

$$24. \begin{aligned} \frac{x}{a} = \frac{y}{b} = \frac{z}{c} = \frac{u}{d}, \\ mx+ny+pz+qu &= r \end{aligned}$$

$$25. \begin{aligned} ax=by=cz=du, \\ y^2-z^2 &= x-u. \end{aligned}$$

$$26. \begin{aligned} y+z &= au, \\ x+z &= bu, \\ x+y &= cu, \\ \frac{1-x}{1-y} &= \frac{a}{b} \end{aligned}$$

$$27. \begin{aligned} x+y &= m, \\ y+z &= n, \\ z+u &= a, \\ u-x &= b. \end{aligned}$$

28.  $11x + 9y + z - u = a,$   
 $11y + 9z + u - x = b,$   
 $11z + 9u + x - y = c,$   
 $11u + 9x + y - z = d.$
29.  $x + ay + a^2z + a^3u + a^4 = 0,$   
 $x + by + b^2z + b^3u + b^4 = 0,$   
 $x + cy + c^2z + c^3u + c^4 = 0,$   
 $x + dy + d^2z + d^3u + d^4 = 0.$
30.  $x + y = a,$   
 $y + z = b,$   
 $z + u = c,$   
 $u + v = d,$   
 $v + x = e.$
31.  $x + ly = a,$   
 $y + mz = b,$   
 $z + nu = c,$   
 $u + pv = d,$   
 $v + qx = e.$
32.  $x + y + z = a,$   
 $y + z + u = b,$   
 $z + u + v = c,$   
 $u + v + x = d,$   
 $v + x + y = e.$
33.  $x - y + z = a,$   
 $y - z + u = b,$   
 $z - u + v = c,$   
 $u - v + x = d,$   
 $v - x + y = e.$
34.  $x + y + z - u = a,$   
 $y + z + u - v = b,$   
 $z + u + v - x = c,$   
 $u + v + x - y = d,$   
 $v + x + y - z = e.$
35.  $x + y + z - u - v = a,$   
 $y + z + u - v - x = b,$   
 $z + u + v - x - y = c,$   
 $u + v + x - y - z = d,$   
 $v + x + y - z - u = e.$
36.  $2x - y - z + 2u - v = 3a,$   
 $2y - z - u + 2v - x = 3b,$   
 $2z - u - v + 2x - y = 3c,$   
 $2u - v - x + 2y - z = 3d,$   
 $2v - x - y + 2z - u = 3e.$
37.  $v - 2x + 3u - 2y + z = a,$   
 $x - 2y + 3v - 2z + u = b,$   
 $y - 2z + 3x - 2u + v = c,$   
 $z - 2u + 3y - 2v + x = d,$   
 $u - 2v + 3z - 2x + y = e.$

## Exercise lxi.

Resolve the following systems of equations :

1.  $\frac{1+x+x^2}{1+y+y^2} = a,$   
 $\frac{1+y+y^2}{1+x+x^2} = b.$
2.  $\frac{x+1}{y+1} = a \left( \frac{x-1}{y-1} \right),$   
 $\frac{x^2+x+1}{y^2+y+1} = b^2 \left( \frac{x-1}{y-1} \right)^2.$
3.  $\frac{(1+x)(1+y)}{(1-x)(1-y)} = \frac{1+a}{1-a},$   
 4.  $\frac{x+y}{1+xy} = \frac{a^2 - \alpha^2}{a^2 + \alpha^2},$

- $\frac{(1+x)(1-y)}{(1-x)(1+y)} = \frac{1+b}{1-b}$        $\frac{x-y}{1-xy} = \frac{b^2-\beta^2}{b^2+\beta^2}$
5.  $\frac{x+y}{1+xy} = \frac{a}{b+c}$ ,      6.  $\frac{x+y}{1-xy} = \frac{2a}{1-a^2}$   
 $\frac{x-y}{1-xy} = \frac{b-c}{a}$ ,       $\frac{x-y}{1+xy} = \frac{2b}{1-b^2}$
7.  $\frac{x+y}{1-xy} = \frac{2a\alpha}{a^2-\alpha^2}$ ,      8.  $\frac{1+xy}{x+y} + \frac{x+y}{1+xy} = \frac{2a}{m}$ ,  
 $\frac{x-y}{1+xy} = \frac{2b\beta}{b^2-\beta^2}$ ,       $\frac{1-xy}{x-y} + \frac{x-y}{1-xy} = \frac{2b}{n}$
9.  $\frac{y(1+x^2)}{x(1+y^2)} = a$ ,      10.  $y+z = 2axyz$ ,  
 $\frac{y(1-x^2)}{x(1-y^2)} = b$ ,       $x+z = 2bxyz$ ,  
 $x+y = 2cxyz$
11.  $\frac{y+z-x}{a} = \frac{z+x-y}{b} = \frac{x+y-z}{c}$ ,      12.  $ax = by = cz$ ,  
 $xyz = m^3$ .       $= \frac{1}{x} + \frac{1}{y} + \frac{1}{z}$
13.  $y^2 + z^2 - x(y+z) = a$ ,  
 $x^2 + z^2 - y(x+z) = b$ ,  
 $x^2 + y^2 - z(x+y) = c$
14.  $2ax = (b+c-a)(y+z)$ ,  
 $2by = (c+a-b)(x+z)$ ,  
 $(x+y+z)^2 + x^2 + y^2 + z^2 = 4(a^2 + b^2 + c^2)$
15.  $\frac{x-1}{y-1} = \frac{a-1}{b-1}$ ,      16.  $\frac{x^2+xy+y^2}{x^2-xy+y^2} = \frac{x^2+y^2}{a} = \frac{xy}{b}$   
 $\frac{x^3-1}{y^3-1} = \frac{a^3-1}{b^3-1}$
17.  $x^4 + x^2y^2 + y^4 = a$ ,      18.  $x^3 + y^3 = \frac{a}{x-y}$ ,  
 $x^2 + xy + y^2 = b$ .       $x^2y - xy^2 = \frac{d}{x+y}$
19.  $xy + \frac{x}{y} = a(x^2 + y^2)$       20.  $x^3 = a(x^2 + y^2) - bxy$

$$xy - \frac{x}{y} = b(x^2 + y^2) \quad y^3 = b(x^2 + y^2) - axy.$$

$$21. \quad 4c(x^2 + 1) = (a+b)(x-y)^2, \\ 4c(y^2 - 1) = (a-b)(x-y)^2.$$

$$22. \quad x^3 - u^3 = \frac{b+c}{2a} (x^2 + xy + y^2)(x+y),$$

$$y^3 - u^3 = \frac{b-c}{2a} (x^2 + xy + y^2)(x+y).$$

$$23. \quad \frac{x+x^2}{y+y^2} = a,$$

$$\frac{y+x^2}{x+y^2} = b.$$

$$24. \quad \frac{x^2 + y^2}{xy} = a,$$

$$\frac{1 + x^2 y^2}{xy} = b.$$

$$25. \quad x(y+z) = a,$$

$$y(z+x) = b,$$

$$z(x+y) = c.$$

$$26. \quad (x+y)(x+z) = a,$$

$$(y+z)(y+x) = b,$$

$$(z+x)(z+y) = c.$$

$$27. \quad x(x+y+z) = a - yz,$$

$$y(x+y+z) = b - zx,$$

$$z(x+y+z) = c - xy.$$

$$28. \quad x^2 - (y-z)^2 = a,$$

$$y^2 - (z-x)^2 = b,$$

$$z^2 - (x-y)^2 = c.$$

$$29. \quad x^2 + y^2 = az,$$

$$x+y = bz,$$

$$x-y = cz.$$

$$30. \quad \frac{1}{x^2} + \frac{1}{y^2} = \frac{2a}{z^2},$$

$$\frac{1}{x^2} - \frac{1}{y^2} = \frac{2b}{z^2},$$

$$\frac{1}{x} + \frac{1}{y} = \frac{1}{c}.$$

$$31. \quad x^2 - y^2 = az,$$

$$x+y = bz,$$

$$x-y = cz.$$

$$32. \quad xy = \frac{z-1}{z+1},$$

$$(x-y)(z+1) = 2a,$$

$$(x^2 - y^2)(z+1)^2 = 4bz.$$

## CHAPTER VII.

EXAMINATION PAPERS : EDUCATION DEPARTMENT AND UNIVERSITY  
OF TORONTO.

## I.

1. State the rules for the addition and subtraction of Algebraic quantities. Express in the simplest form

$$(b+c-a)x+(c+a-b)y+(a+b-c)z$$

$$(c+a-b)x+(a+b-c)y+(b+c-a)z$$

$$(a+b-c)x+(b+c-a)y+(c+a-b)z$$

2. State and prove the Index Laws. Assuming these to be general, interpret  $x^{-n}$ .

Find the products in the following cases :

$$(1) (x^3+6x^2y+12xy^2+8y^3)(x^5-6x^2y+12xy^2-8y^3).$$

$$(2) (a+b+c)(b+c-a)(c+a-b)(a+b-c).$$

3. Prove the rule of signs in Division.

Divide : [Apply Horner's Method to (1)]

$$(1) x^6-22x^4+60x^3-55x^2+12x+4 \text{ by } x^2+6x+1.$$

$$(2) x^4+9+81x^{-4} \text{ by } x^2-3+9x^{-2}. \quad (3) x^m-1 \text{ by } x-1.$$

4. Find the square roots of

$$(1) 4x^{4m}-\frac{4}{3}x^{5m}+\frac{1}{9}x^{6m}$$

$$(2) \frac{a^2}{b^2}+\frac{b^2}{c^2}+\frac{c^2}{a^2}-2\frac{a}{c}-2\frac{c}{b}+2\frac{b}{a}.$$

5. Distinguish between an algebraic equation and an identity.

Solve

$$(1) \sqrt[3]{(1-2x)}+\sqrt[3]{(1+2x)}=3.$$



$$(2) \frac{x-2}{x+2} + \frac{x+2}{x-2} = 2 \cdot \frac{x+3}{x-3}$$

6. A person bought a certain number of oxen for \$320. If he had been able to purchase four more for the same sum, each would have cost him \$4 less. Find the number [of oxen. Explain the negative result.

7. (1) If  $\frac{a}{b} = \frac{c}{d}$  show that  $\frac{a^2+2ab+3b^2}{c^2+2cd+3d^2} = \frac{b(a-3b)}{d(c-3d)}$ .

(2) Find the value of  $x^6 - 200x^5 + 198x^4 + 200x^3 - 197x^2 - 397x$  when  $x = 199$ .

8. Three towns,  $A$ ,  $B$ ,  $C$ , are at the angles of a triangle. From  $A$  to  $C$ , through  $B$ , the distance is 82 miles; from  $B$  to  $A$ , through  $C$ , is 97 miles; and from  $C$  to  $B$ , through  $A$ , is 89 miles. Find the direct distances through the towns.

## II.

1. Prove  $x^m \div x^n = x^{m-n}$ .

Simplify  $(a+b+c)^3 - 3(a+b+c)^2c + 3(a+b+c)c^2 - c^3$ .

2. Prove the rule for finding the L. C. M. of two quantities. Find the L. C. M. of

$$a^3 + b^3 + c^3 - 3abc, \text{ and } (a+b)^2 + 2(a+b)c + c^2.$$

3. Prove  $\frac{a}{b} \times \frac{c}{d} = \frac{ac}{bd}$ .

Simplify  $\left( \frac{1-x^2}{1-x^3} + \frac{1-x}{1-x+x^2} \right) \div \left( \frac{1+x}{1+x+x^2} - \frac{1-x^2}{1+x^3} \right)$ .

4. Reduce to their lowest terms  $\frac{a^{3m} + a^{2m} - 2}{a^{2m} + a^m - 2}$ , and

$$\frac{a(a+2b) + b(b+2c) + c(c+2a)}{a^2 - b^2 - c^2 - 2bc}$$

5. (1.) If  $a^3 - pa^2 + qa - r = 0$ , then  $x^3 - px^2 + qx - r$  is exactly divisible by  $x - a$ .

(2.) Prove that  $(a+b+c)(bc+ca+ab) - (b+c)(c+a)(a+b)$  is divisible by  $abc$ . Is there any other divisor?

6. If  $x = \left(\frac{a+b}{a-b}\right)^{\frac{2mn}{n-m}}$ , then  $\frac{1}{2} \frac{a^2-b^2}{a^2+b^2} (\sqrt[n]{x} + \sqrt[n]{x}) = \left(\frac{a+b}{a-b}\right)^{\frac{m+n}{n-m}}$

7. Solve the equations—

(1)  $\frac{3-2x}{1-2x} - \frac{5-2x}{7-2x} = 1 - \frac{4x^2-2}{7-16x+4x^2}$

(3.)  $\frac{x+3}{x+4} - \frac{x+1}{x+2} = \frac{4x+9}{2x+7} - \frac{12x+17}{6x+16}$

8. A person going at the rate of  $p$  miles an hour, and desiring to reach home by a certain time, finds, when he has still  $r$  miles to go, that, if he were continuing to travel at the same rate, he would be  $q$  hours too late. How much must he increase his speed to reach home in time?

9. Of the three digits comprising a number, the second is double of the third; the sum of the first and third is 9, and the sum of the three digits is 17. Find the number.

10.  $A$  owes  $B$  \$ $a$  due  $m$  months hence, and also \$ $b$  due  $n$  months hence. Find the equation which determines the time at which both sums could be paid at once, reckoning interest at 5 per cent. per annum.

III.

1. If  $x=10$ ,  $y=11$ ,  $z=12$ , find the value of

$\left\{ x^2 - (y+z)^2 \right\} \times \frac{x+y-z}{x+y+z}$ ; and subtract

$(y-z)a^2 + (z-x)ab + (x-y)b^2$  from

$(y-x)a^2 - (y-z)ab - (z-x)b^2$ .

2. Divide  $a + (a+b)x + (a+b+c)x^2 + (a+b+c)x^3 + (b+c)x^4 + cx^5$  by  $1+x+x^2+x^3$ ; and find the square root of

$9 - 24x + 58x^2 - 116x^3 + 129x^4 - 140x^5 + 100x^6$ .

3. Solve (1)  $\frac{4x+5}{x+1} + \frac{x+5}{x+4} = \frac{2x+5}{x+2} - \frac{x^2-10}{x+3} + x$ .

$$(2) \frac{1}{2}x + \frac{1}{3}y + \frac{1}{4}z = 9, \quad \frac{1}{3}x + \frac{1}{4}y - \frac{1}{2}z = -1\frac{1}{2}, \\ \frac{1}{4}x - \frac{1}{2}y + \frac{1}{3}z = 1.$$

4. A boy bought a number of oranges at the rate of 45 cents a dozen; if he had received 20 oranges more for the same money the whole would have cost him only 40 cents a dozen. How many did he buy?

5. A farmer took to market two loads of wheat, amounting together to 75 bushels; he sold them at different prices per bushel, but received on the whole the same amount for each load; had he sold the whole quantity at the lower price he would have received \$78.75; but had he sold it at the higher price he would have received \$90. Find the number of bushels in each load.

6. Show how to find the square root of  $a + \sqrt{b}$ .

Find the square root of  $1 + \sqrt{1 - a^2}$

7. Solve  $\frac{6x+5}{2x-7} + \frac{4x-1}{x-2} = \frac{7x+1}{x-3}$ ; and find the value of  $x$

when  $ax^2 - 36x + 81 = 0$ , has equal roots.

8. If  $\frac{a}{b} = \frac{c}{d}$ , prove that  $\frac{a+c}{b+d} = \sqrt[3]{\frac{(a^3+c^3)}{(b^3+d^3)}}$ ; and

$$\text{that } \frac{a+b}{a-b} = \frac{\sqrt{(ac)} + \sqrt{(bd)}}{\sqrt{(ac)} - \sqrt{(bd)}}.$$

9. Show that  $a^3(b-c) + b^3(c-a) + c^3(a-b)$  is exactly divisible by  $a+b+c$ ; and resolve the expression into its factors.

#### IV.

1. Multiply  $a^2 + b^2 - c^2 + 2ab$  by  $a^2 - b^2 + c^2 + 2ac$ , and divide the product by  $a^2 - b^2 - c^2 + 2bc$ .

2. Simplify

$$\frac{18a^2b^2}{x+y} \div \left\{ \frac{3ab(x-y)}{7(c+d)} \div \left( \frac{4(c-d)}{21ab^2} \div \frac{8(c^2-d^2)}{a(x^2-y^2)} \right) \right\}$$

3. Find the L.C.M. of  $4x^2 - 9y^2$ ,  $4x^2 - 10xy + 6y^2$ , and  $6x^2 - 13xy + 6y^2$ , and the G.C.M. of  $1 + x^{\frac{1}{2}} + x + x^{\frac{3}{2}}$  and  $2x + 2x^{\frac{1}{2}} + 3x^{\frac{3}{2}} + 3x^{\frac{5}{2}}$

4. Obtain the square root of  $\frac{1}{2} - \frac{2}{3}\sqrt{\frac{1}{2}}$ , and find the value of  $c$  when  $4x^4 - 12x^3y + cx^2y^2 - 12xy^3 + 4y^4$  is a perfect square.

5. Distinguish between an *equation* and an *identity*. Give an example of each. What value of  $m$  makes  $(x-3)^2 - (x-1)(x-5) = m$  an identity? Can any value of  $m$  make it an equation?

6. Reduce to its simplest form

$$\frac{\sqrt{2+x} - \sqrt{1+x}}{\sqrt{1+x} - \sqrt{x}} + \frac{1 + \sqrt{1-1 \div (1+x)}}{1 + \sqrt{1+1 \div (1+x)}}$$

7. Solve the equations

$$(1) \frac{2x+5}{x+2} + \frac{2x-5}{x-2} - \frac{4x-5}{x-1} = 0.$$

$$(2) 73y - 5x = (x-5y)(x+3y),$$

$$\frac{2}{x-5y} - \frac{5}{x+3y} = \frac{7}{33}.$$

8. A person performed a journey of  $22\frac{1}{2}$  miles, partly by carriage, at 10 miles an hour, and partly by train, at 36 miles an hour, and the remainder by walking, at 4 miles an hour. He did the whole in 1 hour 50 minutes. Had he walked the first portion, and performed the last by carriage, it would have taken him 2 hours  $30\frac{1}{2}$  minutes. Find the respective distances by carriage, train and walking.

9. Solve

$$\frac{x+3}{x+4} - \frac{x+1}{x+2} = \frac{4x+9}{2x+7} - \frac{12x+17}{6x+16}$$

10. What value of  $y$  will make  $2x^2 + 3xy + 6y^2$  exactly divisible by  $x-3$ ?

If  $a$  and  $b$  are the roots of the equation  $x^2 + x + 1 = 0$ , show that  $a^3 - b^3 = 0$ .

## V.

1. Multiply

$$4x^2 - \frac{2}{3}x + \frac{1}{25} \text{ by } 2x + \frac{1}{5}.$$

Prove that

$$\left(\frac{1}{2}x - y\right)^3 - \left(x - \frac{1}{4}y\right)^3 \text{ is exactly divisible by } x + y.$$

2. Express in words the meaning of the formula

$$(x+a)(x+b) = x^2 + (a+b)x + ab.$$

Retaining the order of the terms, how will the right-hand member of this expression be affected by changing, in the left-hand member (1) the sign of  $b$  only, (2) the sign of  $a$  only, (3) the signs of both  $a$  and  $b$ ?

3. Simplify  $(a+b)^4 + (a-b)^4 - 2(a^2 - b^2)^2$ ; and show that

$$(a+b+c)(b+c-a)(a+c-b)(a+b-c) = 4a^2b^2$$

when  $a^2 + b^2 = c^2$ .4. Prove that  $\frac{a}{b} \div \frac{c}{d} = \frac{ad}{bc}$ .

Simplify

$$\left(\frac{a^2 + b^2}{2ab} + 1\right) \left(\frac{ab^2}{a^3 + b^3}\right) \div \frac{4a(a+b)}{a^2 - ab + b^2}.$$

5. I went from Toronto to Niagara, 35 miles, in the steamer "City of Toronto" and returned in the "Rothsay," making the round trip in 5 hours and 15 minutes; on another occasion I went in the "Rothsay" (whose speed on this occasion was 1 mile an hour less than usual), from Toronto to Lewiston, 42 miles, and returned in the "City of Toronto," making the round trip in 6 hours and 30 minutes; find the usual rates per hour which these steamers make.

6. Solve

$$(1) \frac{3}{x} - \frac{2}{y} = \frac{1}{a}, \quad \frac{2}{x} - \frac{1}{y} = \frac{2}{a}$$

$$(2) x^2 + 5x = 5\sqrt{(x^2 + 5x + 28)} - 4.$$

7. Find three consecutive numbers whose product is 48 times the middle number.

8. If  $m$  and  $n$  are the roots of  $ax^2 + bx + c = 0$ , then

$$ax^2 + bx + c = a(x-m)(x-n).$$

Show that if  $ax^2 + bx + c = 0$  has equal roots, one of them is given by the equation

$$(2a^2 - 2ab)x + ab - b^2 = 0.$$

9. If  $\frac{m}{x} = \frac{n}{y}$  and  $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ , prove that

$$\frac{m^2}{a^2} + \frac{n^2}{b^2} = \frac{m^2 + n^2}{x^2 + y^2}.$$

---

### VI.

1. Simplify

$$\left( \frac{ax^2 - ay^2 + 2axy}{x^2 + y^2} \right)^2 + \left( \frac{by^2 - bx^2 + 2axy}{x^2 + y^2} \right)^2.$$

2. Divide  $a^3 - b^3 - c^3 - 3abc$  by  $a - b - c$ , and show, without expansion, that

$$(1 + x + x^2)^3 - (1 - x + x^2)^3 - 6x(x^4 + x^2 + 1) - 8x^3 = 0.$$

3. Resolve into factors  $x^4 - \frac{1}{4}x^2y^2 + y^4$ , and

$$7x^2 - 6y^2 - xy + 19x + 33y - 36; \text{ and prove that}$$

$$b^2(c+a) + c^2(a+b) - a^2(b+c) + abc \text{ is exactly divisible by } b+c-a.$$

4. Apply Horner's method of division to find the value of  $5x^5 + 497x^4 + 200x^3 + 196x^2 - 218x - 2000$  when  $x = -99$ , and the value of  $6x^5 + 5x^4 - 17x^3 - 6x^2 + 10x - 2$  when  $2x^2 = -3x + 1$ .

5. Find what

$$\frac{\sqrt{a+x} + \sqrt{a-x}}{\sqrt{a+x} - \sqrt{a-x}} \text{ becomes when } x = \frac{2ab}{1+b^2}.$$

6. If  $a$  and  $b$  be any positive numbers, prove that

$$\frac{1}{a} + \frac{a}{1+a} > 1, \quad \frac{a}{b} + \frac{b}{a} > 2.$$

7. Solve the equations—

$$(1) \quad x^{\frac{1}{2}} + y^{\frac{1}{2}} = 5,$$

$$x^{-\frac{1}{2}} + y^{-\frac{1}{2}} = \frac{5}{6}.$$

$$(2) \quad x + 2y + 3z = 14,$$

$$2x + 3y + z = 11,$$

$$3x + y + 2z = 11.$$

$$(3) \quad (x+1)(x+3)(x+4)(x+6) = 13.$$

8. There are three consecutive numbers such that the sum of their cubes is equal to  $16\frac{2}{3}$  times the product of the two higher numbers: find the numbers.

9. (1) Form an equation three of whose roots are 0,  $\sqrt{-3}$ , and  $1 - \sqrt{2}$ .

(2) If one of the roots of the equation  $x^2 + px + q = 0$ , is a mean proportional between  $p$  and  $q$ , prove that  $p^3 = q(1+p)^2$ .

10. Two trains start at the same instant, the one from  $B$  to  $A$ , the other from  $A$  to  $B$ ; they meet in  $1\frac{1}{2}$  hours; and the train for  $A$  reaches its destination  $52\frac{1}{2}$  minutes before the other train reaches  $B$ : compare the rates of the trains.

## VII.

1. Give some application of the "rule of signs" in Algebraic Multiplication and Division.

2. Find the numerical value of the quantity

$$bc(c-a)(a-b) - ca(a-b)(b-c) + ab(b-c)(c-a),$$

when  $a = 10$ ,  $b = 01$ ,  $c = 0$ ; and prove that if

$$x = \frac{c^2 - ab}{a+b}, \text{ then will } (a+b) \cdot \frac{a^2 + b^2 + c^2 + x^2}{a+b-c+x} \\ = b^2 + c^2 + (a+b)(a+c).$$

3. Investigate a method of finding by inspection the remainder after dividing any rational and integral function of  $x$  by  $x+a$ .

Show that the quantity

$$a^2b^2 - ab^2x - (a^2 + 2b^2)x^2 + ax^3 + 2x^4$$

is divisible by each of the quantities  $x+a$ ,  $x+b$ ,  $a-2x$ ,  $b-x$ .

4. Investigate the rule for finding the H.C.F. of two algebraic quantities, showing under what limitations factors may be introduced or suppressed at any step.

Find the H.C.F. of

(1)  $6x^4 - 7x^3 - 13x^2 + 19x - 6$  and  $x^3 + 2x^2 - 1$ .

(2)  $(x+y)(ax^2 - by^2) - xy(a-b)(x+y)$ , and  
 $(x-y)(ax^2 - by^2) + xy(a-b)(x-y)$ .

5. Prove, by general reasoning, that the value of a fraction is not altered by multiplying or dividing both the numerator and denominator by the same quantity.

Simplify (1)  $\frac{13}{12(2x-3)} - \frac{7}{12(2x+3)} - \frac{x-4}{4x^2+9}$ .

(2)  $\left\{ \frac{1}{(x+a)(x-b)} + \frac{1}{(x-a)(x+b)} \right\} \div$   
 $\left\{ \frac{1}{(x+a)(x+b)} + \frac{1}{(x-a)(x-b)} \right\}$ .

6. With respect to  $x$ , the equations

$$\frac{18}{4} + \frac{2x-24}{11} + \frac{11x-34}{22} = \frac{7}{44}$$

(2)  $\frac{5x^2+x-3}{5x-4} - \frac{7x^2-3x-9}{7x-10} = \frac{x-3}{35x^2-78x+40}$ .

(3)  $x^2 = ax + by$ , and  $y^2 = bx + ay$ .

### VIII.

1. Define the terms "power," "root," "index," and "coefficient"; explain also the reasoning by which it is shown that

$$a - (b - c) = a - b + c.$$



2. Multiply  $(x^2 + xy + y^2)^2$  by  $(x - y)^3$ .

Find the values of  $a$  and  $b$  which will make

$x^2 + ax + b$  divisible by  $x + p$ , and also by  $x + q$ .

3. Divide  $x^6 + y^6 + 2x^3y^3$  by  $(x + y)^2$ , and

$a^8 + a^4b^4 + b^8$  by  $(a^2 - ab + b^2)(a^2 + ab + b^2)$ .

4. Investigate a rule for the extraction of the square root of any algebraic quantity, and deduce the rule for the extraction of the square root of a number.

If to any square number be added the square of half the number immediately preceding it, the sum will be a complete square: viz., the square of half the number immediately following it.

5. Find the square root of

$$(1) a^2x^6 + 2abcx^4 + (b^2 + 2ac)x^2 + c^2x^{-2} + 2bc.$$

$$(2) \frac{1}{4}x^{\frac{1}{2}} - \frac{1}{3}x^{\frac{1}{3}} + \frac{1}{4}x^{\frac{2}{3}} + \frac{1}{5}x^{\frac{2}{5}} - \frac{1}{6}x^{\frac{2}{6}} + \frac{1}{7}x^{\frac{2}{7}}.$$

6. If  $x^2 + ax + b$  and  $x^2 + a'x - b$  have a common measure, it will be  $x + \frac{a + a'}{2}$ , and the condition that they may have a common measure is  $4b = a^2 - a'^2$ .

Find the H. C. F. of  $x^4 + p^2x^2 + p^4$  and  $x^4 + 2px^3 + p^2x^2 - p^4$ .

Find the L. C. M. of  $2\frac{1}{2}(x^2 + x - 20)$ ,  $3\frac{1}{3}(x^2 - x - 30)$ , and  $4\frac{1}{6}(x^2 - 10x + 24)$ .

7. Find values of  $a$  and  $b$  which will render the fraction

$$\frac{3x^2 - (4a + b)x + a + 2b^2}{5x^2 - (8a + b)x - a + 4b^2}$$

the same, for all values of  $x$ .

8. Solve the equation  $2 + \sqrt{(x+1)(x+6)} - \sqrt{(x-1)(x+5)} = 0$ , and account for the circumstance, that the values of  $x$ , determined from it, apparently do not satisfy the equation.

## IX.

1. Prove that  $a(2n+1)(a^2+n\overline{a+1}) - n(2a+1)(n^2+a\overline{a+1}) = (a-n)^3$ .

2. If  $a$ ,  $b$ , and  $c$  are positive quantities, and if  $a > b$  and  $c > a - b$ , prove that

$$c - (a - b) = c - a + b.$$

Assuming this equation to hold good when  $a$ ,  $b$  and  $c$  are unrestricted, prove that the expression  $-(-a)$ , occurring in an algebraic operation, is equivalent to  $+a$ .

3. If  $x^3 + ax^2 + b$  and  $x^3 + px + q$  have a common measure of the form of  $x^2 + mx + n$ , then  $a^3bq = (b - q)^3$

4. Find the H. C. F. of

$$a^2 - b^2 - abxy + abx^{-1}y^{-1}, \text{ and } a^2x^3 - b^2y^{-1} + a^2bx^2y - b^2xy^{-2}.$$

5.  $A$  and  $B$  are two numbers, each of two digits. The left-hand digit of  $A$  exceeds that of  $B$  by  $x$ ; the excess of  $A$  above  $B$  is  $y$ ; but the sum of the digits of  $B$  exceeds the sum of the digits of  $A$  by  $z$ . Prove that  $y + z = 9x$ ; and give an example of two such numbers as  $A$  and  $B$ .

6. If  $\frac{a}{b} = \frac{b}{c} = \frac{c}{d}$ , prove that each of these ratios

$$= \sqrt[3]{\frac{a}{d}} \text{ and also } = \frac{a + b + c}{b + c + d}.$$

7. Solve the equations

$$(1) \frac{x+a}{x-a} - \frac{x-a}{x+a} = \frac{b+x}{b-x} - \frac{b-x}{b+x}$$

$$(2) a(x^2 + y^2) - b(x^2 - y^2) = 2a \\ (a^2 - b^2)(x^2 - y^2) = 4ab.$$

8. A farmer buys  $a$  sheep for  $\$P$  and sells  $b$  of them at a gain of 5 per cent.; at what price ought he to sell the remainder to gain 10 per cent. on the whole?

9. The sum of three numbers is 70; and if the second is divided by the first, the quotient is 2, and the remainder 1; but if the third is divided by the second, the quotient is 3, and the remainder is 3; what are the numbers.

## X.

1. Divide  $ax^3 + 2cxyz + by^3 + ax^2(y+z) + by^2(z+x) + 2cxy(x+y)$  by  $x+y+z$ .

2. Prove that if  $x^4 + px^2 + qx + a^2$  be divisible by  $x^2 - 1$ , it is also divisible by  $x^2 - a^2$ .

3. Explain the reason for introducing or suppressing factors in the process of finding the H.C.F. of two algebraical quantities.

Why is the name "Greatest Common Measure" objectionable?

Find the H.C.F. of  $x^4 - x^3 - x^2 - x - 2$  and  $3x^3 - 7x^2 + 3x - 2$ .

4. A traveller leaves  $A$  for  $B$  at the same time that another leaves  $B$  for  $A$ ; the former walks at the rate of 3 miles an hour till he has performed half the distance; he then rests for an hour; after which he resumes his journey, walking now at the rate of 4 miles an hour; the second traveller goes at the rate of 4 miles an hour till he has got over one-third of the distance between  $B$  and  $A$ ; he then rests for 40 minutes; after which he resumes his journey, walking now at the rate of 3 miles an hour. The travellers reach  $A$  and  $B$  respectively at the same time. Find the distance between  $A$  and  $B$ .

5. Show by examining the square of  $a+b$  how the square root of an algebraical quantity may be found.

Find the square roots of

$$(1) 25x^4 - 30ax^3 + 49a^2x^2 - 24a^3x + 16a^4, \text{ and}$$

$$(2) \frac{x^2}{y^2} + \frac{y^2}{x^2} - \left( \frac{x}{y} + \frac{y}{x} \right) \sqrt{2} + \frac{5}{2}.$$

6. Show that  $a^{\frac{m}{n}} = \sqrt[n]{a^m}$ , when  $m$  and  $n$  are integers, and  $m$  is divisible by  $n$ ; and state the principle on which you would maintain the truth of the equation for all values of  $m$  and  $n$ .

7. Solve the equations

$$(1) \frac{5x^2 + x - 3}{5x - 4} = \frac{7x^2 - 3x - 9}{7x - 10}.$$

$$(2) (3x-1)^2 + (4x-2)^2 = (5x-3)^2.$$

8. Two regular polygons are so related that the number of their sides is as 2 to 3, and the magnitude of their angles as 3 to 4; find the figures.

---

XI.

1. State in words the several operations to be performed in order to obtain the result expressed by the following algebraical expression :

$$\sqrt[4]{\frac{ma^2 + nb^2}{m+n}}$$

Also find its value when  $a = b = 4$ .

2. Two men,  $A$  and  $B$ , dig a trench in  $3\frac{2}{3}$  days. If  $A$  were to do more work by one-third than he does, and  $B$  more work by one-half than he does, they would dig the trench in  $2\frac{2}{3}$  days. In what time would each dig it alone, at his present rate of work ?

3. Perform the multiplications in

(1)

$$\left(2x^{\frac{1}{2}} + 3y^{\frac{1}{3}}\right) \left(2x^{\frac{1}{3}} - 2y^{\frac{1}{4}}\right) \left(4x^{\frac{1}{4}} + 6x^{\frac{1}{2}}y^{\frac{1}{3}} + 9y^{\frac{1}{2}}\right) \left(4x^{\frac{1}{2}} - 6x^{\frac{1}{3}}y^{\frac{1}{4}} + 9y^{\frac{1}{3}}\right)$$

(2)  $\left(\frac{1}{4}x^2 + \frac{1}{3}xy + \frac{2}{3}y^2\right) \left(\frac{1}{4}x^2 - \frac{1}{3}xy + \frac{2}{3}y^2\right)$ .

4. Divide

(1)  $x^4 + 9 + 81x^{-4}$  by  $x^2 - 3 + 9x^{-2}$ .

(2)  $x^4 - (a+b+p)x^3 + (ap+bp-c+q)x^2 - (aq+bq-cp)x - qc$  by  $x^2 - px + q$ .

5. Show that  $x^{2m+1} - x^{2n-1}$  is always divisible by  $x \pm 1$ ,  $m$  and  $n$  being any positive integers.

6. Define a fraction; and from your definition prove a rule for adding together two fractions with different denominators.

Add together the fractions,

$$\frac{a^2 - bc}{(a+b)(a+c)}, \quad \frac{b^2 - ca}{(b+c)(b+a)}, \quad \frac{c^2 - ab}{(c+a)(c+b)}$$

7. Solve the following equations :

$$(1) \frac{x^2 + 2x + 2}{x + 1} + \frac{x^2 + 8x + 20}{x + 4} = \frac{x^2 + 4x + 6}{x + 2} + \frac{x^2 + 6x + 12}{x + 3}.$$

$$(2) (x^2 + y^2) \frac{x}{y} = \frac{100}{3}, \quad (x^2 - y^2) \frac{y}{x} = \frac{21}{4}.$$

## XII.

1. When  $m$  and  $n$  are whole numbers, and  $m$  greater than  $n$ , show that  $\frac{a^m}{a^n} = a^{m-n}$  and that  $\frac{1}{a^n}$  is correctly symbolized by  $a^{-n}$ .

2. Multiply  $(a-b)(a+b)(a^2+b^2)(a^4+b^4) \dots$  to  $(n+1)$  factors.

3. Divide  $1-x$  by  $1-2x$ , to 5 terms, and write down the  $(r+1)$ th term, and the remainder after  $(r+1)$  terms.

4. If the number three be divided into any two parts, show that the difference of the squares is three times the difference of the numbers.

5. Find the L. C. M. of  $1-8x+17x^2+2x^3-24x^4$ , and  $1-2x-13x^2+38x^3-24x^4$ .

6. What relation must there be between the coefficients  $m$ ,  $n$ ,  $p$  and  $q$ , in order that

$$(x^2 + mx + n)^2 + px^2 + qx$$

may be an exact square for all values of  $x$ ?

7. Solve the following equations :

$$(1) \frac{1+x^3}{(1+x)^2} + \frac{1-x^3}{(1-x)^2} = a.$$

$$(2) \frac{ax-b^2}{\sqrt{(ax)+b}} = \frac{\sqrt{(ax)-b}}{n} - c.$$

$$(3) \frac{xy}{x+y} = 1, \quad \frac{xz}{x+z} = 2, \quad \text{and} \quad \frac{yz}{y+z} = 3.$$

8. Given  $x+y+z=ax=by$ , find  $(x+y+z) - z$ .

9. Find a number expressed in the decimal notation by two digits, whose sum is 10; and such, that if 1 be taken from its double, the remainder will be expressed by the same digits in a reversed order.

XIII.

1. Find the value, when  $a = 2\frac{1}{2}$ ,  $b = 3\frac{1}{2}$ ,  $c = 4\frac{1}{2}$  of

$$\frac{b^2c^2(c^2 - b^2) + c^2a^2(a^2 - c^2) + a^2b^2(b^2 - a^2)}{(b+c)(c+a)(a+b)}$$

2. Show that the value of the expression, in the preceding question, is not altered by changing  $a$  into  $a+x$ ,  $b$  into  $b+x$ , and  $c$  into  $c+x$ .

3. Multiply  $(1+a_1x)(1+a_2x)(1+a_3x) \dots (1+a_nx)$  to 3 terms.

4. A speculator borrows a sum of money at the yearly interest of 7 per cent.; part of the amount he invests at  $8\frac{1}{2}$  per cent., and the remainder at 9; and, at the end of the year, he finds that he has made a profit of \$75; but, had the former part been invested at 9 per cent., and the latter at  $8\frac{1}{2}$ , his profit at the end of the year would have been only \$65. Find the whole sum borrowed.

5. Given  $ax+by=c$ ,  $a'x+b'y=c'$ , determine the value of  $mx+ny$ , and find the conditions under which the value becomes indeterminate

6. If  $\frac{a_1}{a_2} = \frac{a_2}{a_3} = \dots = \frac{a_{n-1}}{a_n}$ ,

then will  $a_1 + a_2 + a_3 + \dots + a_n = \frac{a_1^2 - a_2 a_n}{a_1 - a_2}$ .

7. Eliminate  $x$  and  $y$  from the equations

$$x^{\frac{2}{3}} + y^{\frac{2}{3}} = a^{\frac{2}{3}}$$

$$\alpha = x + 3x^{\frac{1}{3}}y^{\frac{2}{3}}$$

$$\beta = y + 3x^{\frac{2}{3}}y^{\frac{1}{3}}$$

8. If  $ax^2+bx+c=0$  and  $a_1x^2+b_1x+c_1=0$ , then will

$$(ab_1 - a_1b)(bc_1 - b_1c) = (ac_1 - a_1c)^2.$$

9. Find that number of two figures to which if the number formed by changing the places of the digits be added, the sum is 121; and if the same two numbers be subtracted, the remainder is 9.

## XIV.

1. Simplify

$$a(b+c)^2 + b(c+a)^2 + c(a+b)^2 - \{(a-b)(a-c)(b+c) + (b-c)(b-a)(c+a) + (c-a)(c-b)(a+b)\}.$$

2. State the law of Indices, and prove it for positive integral indices; and assuming it to be general, interpret the expressions

$$x^{-m}, \quad x^{\frac{m}{n}},$$

where  $m$  and  $n$  are positive integers.

3. Having given the equations,

$$\begin{aligned} x + y + z &= 0, & x' + y' + z' &= 0, \\ a^2 &= x^2 + x'^2, & b^2 &= y^2 + y'^2, \\ c^2 &= z^2 + z'^2; \end{aligned}$$

$$\text{prove that } a^2(yz - y'z') + b^2(zc - z'c') + c^2(xy - x'y') = 0.$$

4. A traveller  $P$  sets out to walk from  $A$  to  $B$ , proceeding at the rate of 3 miles an hour; and, 32 minutes afterwards, another traveller  $Q$  sets out to walk from  $B$  to  $A$ , proceeding at a uniform rate. They meet half way betwixt  $A$  and  $B$ .  $P$  then quickens his pace by 1 mile an hour; and  $Q$  slackens his 1 mile an hour.  $Q$  reaches  $A$  at the same time that  $P$  reaches  $B$ . Find the distance between  $A$  and  $B$ .

5. How are equations classified?

Solve the equations—

$$(1) \quad mx + am = n^2x + an^2.$$

$$(2) \quad x^4 - x^2 + y^4 - y^2 = 84, \\ x^2 + x^2y^2 + y^2 = 49.$$

6. What two numbers are those whose difference, sum and product are to each other as the three numbers 2, 3, 5?

## XV.

1. What is the meaning of the symbols  $a, a^2, a^3 \dots$ ?Show *a priori* that  $a^0 = 1$ ; how do you know that  $ab = ba$ ?

How is it proved that the multiplication of like signs gives a positive, and that of unlike signs, a negative result.

2. Find the value of

$$(b-c)^3 + 2(c-a)^3 + (a-b)^3 - 3(b-c)(c-a)(a-b)$$

when  $a = 1$ ,  $b = -\frac{1}{2}$ ,  $c = \frac{3}{2}$ .

3. Simplify the following expression :

$$(ac - b^2)(ce - d^2) + (ae - c^2)(bd - c^2) - (ad - bc)(be - ca)$$

4.  $P$  and  $Q$  are travelling along the same road in the same direction. At noon  $P$ , who goes at the rate of  $m$  miles an hour, is at a point  $A$ ; while  $Q$  who goes at the rate of  $n$  miles in the hour, is at a point  $B$ , two miles in advance of  $A$ . When are they together?

Has the answer a meaning, when  $m - n$  is negative? Has it a meaning when  $m = n$ ? If so, state what interpretation it must receive in these cases.

5. Show how to find the Least Common Multiply of two or more algebraic quantities.

$$(1) \quad x^2 - ax - 2a^2, \quad x^3 + ax^2 \quad \text{and} \quad ax^2 - x^3.$$

$$(2) \quad x^3 - x^2y - a^2x + a^2y \quad \text{and} \quad x^3 + ax^2 - xy^2 - ay^3.$$

In what algebraic operations is the Lowest Common Multiple of two or more quantities required?

6. State and prove the principle upon which the rules of Addition and Subtraction of fractions are founded.

Simplify the following expressions :

$$(1) \quad \frac{(a+b-c)^2 - d^2}{(a+b)^2 - (c+d)^2} + \frac{(b+c-a)^2 - d^2}{(b+c)^2 - (a+d)^2} + \frac{(c+a-b)^2 - d^2}{(c+a)^2 - (b+d)^2}.$$

$$(2) \quad \frac{x^2 + y^2 - z^2 + 2xy}{x^2 - y^2 - z^2 + 2yz}, \quad \frac{a^3 + a^2b}{a^2b - b^3} - \frac{a(a-b)}{(a+b)b} - \frac{2ab}{a^2 - b^2}.$$

$$7. \quad \text{If } ax - by + c(x-y) = (a-b)(a+b-c),$$

$$by - cz + a(y-z) = (b-c)(b+c-a),$$

$$cz - ax + b(z-x) = (c-a)(c+a-b)$$

$$\text{then will } a^2(b-c) + b^2(c-a) + c^2(a-b) = 0.$$

8.  $P$  is a number, of two digits,  $x$  being the left hand digit, and  $y$  the right. By inverting the digits, the number  $Q$  is obtained. Prove that  $11(x+y)(P-Q) = 9(x-y)(P+Q)$ .



## XVI.

1. Show that

$$\{(ax+by)^2+(ay-bx)^2\} \{(ax+by)^2-(ay+bx)^2\} = (a^4-b^4)(x^4-y^4); \text{ and that}$$

$$2(a-b)(a-c)+2(b-c)(b-a)+2(c-b)(c-a)$$

is the sum of three squares.

2. If  $s = a + b + c + \&c.$  to  $n$  terms, then

$$\frac{s-a}{s} + \frac{s-b}{s} + \frac{s-c}{s} + \&c. = n - 1.$$

3. Show that  $a-b$ ,  $b-c$ , and  $c-a$  cannot be all three positive or all three negative.

4. Extract the square root of

$$4x^8 + 9x^6 - 12x^4 + 16x^2 + 9 - 2x(6x^6 - 8x^4 + 9x^2 - 12).$$

5. Given

$$ab - \frac{1}{2}(a+b)(p+q) + pq = 0,$$

$$cd - \frac{1}{2}(c+d)(p+q) + pq = 0,$$

find the value of  $p-q$ , and show that if either  $a$  or  $b$  is equal to  $c$  or  $d$ , then  $p$  is equal to  $q$ , unless  $a+b=c+d$ .

6. Find the value of  $\frac{x}{y}$ , having given

$$\frac{x^{2n} - ay^{2n}}{x^{2n} + ay^{2n}} = \frac{x^n - b(x-y)^n}{x^n + b(x-y)^n}.$$

7. Prove that  $(a-b)(b-c)(c-a)$  is a common measure of the quantities

$$(a^2 - b^2)^5 + (b^2 - c^2)^5 + (c^2 - a^2)^5,$$

$$c^4(a-b) + a^4(b-c) + b^4(c-a).$$

8. Find the conditions that  $a_1x + b_1y = c_1$ ,  $a_2x + b_2y = c_2$ , and  $a_3x + b_3y = c_3$  may be satisfied by the same values of  $x$  and  $y$ .

9. Two persons,  $A$  and  $B$ , start at the same instant from two stations ( $c$ ) miles apart, and proceed in the same direction along the line joining the stations with velocities ( $a$ ) and ( $b$ ) miles per hour. Find the distance ( $x$ ) from the stations where  $A$  overtakes  $B$ , and interpret the result when  $a < b$ .

## XVII.

1. Express in symbols the result of subtracting from unity the quotient obtained by dividing the sum of  $a$  and  $b$  by their product.

2. Multiply together  $x + \sqrt{a+b}$ ,  $x - \sqrt{a+b}$ ,  $x + \sqrt{a-b}$  and  $x - \sqrt{a-b}$ ; and divide  $24a^3 - 22a^2b + 2a^2c - 5ab^2 + 27abc - 34ac^2 + 6b^3 - 22b^2c + 16bc^2 + 8c^3$  by  $3a - 2b + 4c$ .

3. If  $x+a$  be the H. C. F. of  $x^2+px+q$  and  $x^2+p'x+q'$ , their L. C. M. will be  $(x+a)(x+p-a)(x+p'-a)$ .

Show that the difference between

$$\frac{x}{x-a} + \frac{x}{x-b} + \frac{x}{x-c} \text{ and } \frac{a}{x-a} + \frac{b}{x-b} + \frac{c}{x-c}$$

is the same whatever values be given to  $x$ .

4. Prove, if the four fractions

$$\frac{bx+cy+dz}{b+c+d-a}, \quad \frac{cx+dy+az}{c+d+a-b}, \quad \frac{dx+ay+bz}{d+a+b-c}, \quad \frac{ax+by+cz}{a+b+c-d}$$

are equal to one another, their common value will be equal to  $\frac{x+y+z}{2}$  as long as  $a+b+c+d$  does not vanish.

5. What do you mean by *solving an equation*. Show that 3 is a root of the equation

$$\sqrt{(x^2-3x+4)} = \frac{3+\sqrt{3}(x-2)}{x^2-7}.$$

6. Eliminate  $x$  between the equations

$$x^3 + \frac{1}{x^3} + 3\left(x + \frac{1}{x}\right) = m, \text{ and}$$

$$x^3 - \frac{1}{x^3} - 3\left(x - \frac{1}{x}\right) = n.$$

7. If  $\frac{1}{a} + \frac{1}{b} - \frac{1}{c} = \frac{1}{a+b-c}$ ,  $a, b, c$  are not all different.

8. A cask,  $A$ , contains  $m$  gallons of wine and  $n$  gallons of water; and another cask,  $B$ , contains  $p$  gallons of wine and  $q$  gallons of water, how many gallons must be drawn from each cask so as to produce by their mixture  $b$  gallons of wine and  $c$  gallons of water?

## XVIII.

1. Multiply together the factors

$$1-x, 1+x, 1+x^2, 1+x^4, \text{ and } 1+x^8,$$

and show that if  $n$  is any uneven number, the sum of the  $n$ th powers of any two numbers is always divisible by the sum of the numbers.

2. Find the numerical value of the expression

$$\frac{c}{b} \cdot \frac{\sqrt[3]{a+\sqrt{c}}}{\sqrt[3]{a-\sqrt{c}}}$$

where  $a, b, c$  are connected by the equation  $a^2(b-c)^2 - c(b+c)^2 = 0$ .

3.  $A$  has a younger brother,  $B$ . The difference between their ages is  $\frac{2}{3}$  of the sum of their ages. By adding twice  $B$ 's age to 5 times  $A$ 's, we obtain the age of the father; and by subtracting twice  $B$ 's age from 5 times  $A$ 's, we obtain the age of the mother. Show that the age of the mother is  $\frac{1}{17}$  that of the father.

4. Find the H.C.F. of

$$x^3 - (2a+b)x^2 + a(2a+b)x - a^2(a+b), \text{ and}$$

$$x^3 - (2b+a)x^2 + b(2b+a)x - b^2(b+a).$$

5. If
- $\frac{1}{b} + \frac{1}{c} = \frac{4}{a}$
- , shew that

$$(a+b-c)^3 + 2(b+c-a)^3 + (c+a-b)^3 = 2(b+c)^3.$$

6. Show fully how the rule for finding the square root of a given number is obtained. If  $n+1$  figures of the square root of a number have been obtained, prove that the remaining  $n$  may be obtained by division.

Extract the square root of

$$x^2(x^2+y^2+z^2) + y^2z^2 + 2x(y+z)(yz-x^2).$$

7. Find the value of the expression

$$\frac{x-y}{1+xy} \text{ when } x = \frac{a+b}{a-b}, y = \frac{b}{a}.$$

8. Solve the equations :

$$(1) \frac{1}{2}(x-2a) - \frac{1}{3}(x+3a) + \frac{1}{6}(x-6a) = 0.$$

$$(2) \sqrt{(2x^2+1)} + \sqrt{(2x^2+3)} = 2(1-x).$$

9. Divide 21 into two parts, so that ten times one of them may exceed nine times the other by 1.

---

XIX.

1. Multiply together

$$x^2 - \frac{9}{4}ax + \frac{1}{2}a^2 + \frac{5}{4}x - \frac{3}{4}a + \frac{1}{4};$$

$$x^2 + \frac{1}{4}ax - a^2 - \frac{7}{4}x + \frac{3}{2}a - \frac{1}{2}.$$

Divide this product by

$$\frac{1}{4}x^2 + \frac{1}{2}ax - 2a^2 - \frac{1}{4}x + 2a - \frac{1}{2};$$

and extract the square root of the quotient.

2. If  $x+y+z = \frac{1}{x} + \frac{1}{y} + \frac{1}{z} = 0$ , shew that

$$(x^6 + y^6 + z^6) \div (x^3 + y^3 + z^3) = xyz.$$

3. Find the H. C. D. of  $20x^4 + x^2 - 1$  and  $75x^4 + 15x^3 - 3x - 3$ ; also of  $(x+y)^7 - x^7 - y^7$  and  $(x^3 - y^3)^2$ .

4. Given that  $ab - (a+b)(x+y) + 4xy = 0$ ,

$$cd - (c+d)(x-y) + 4xy = 0,$$

find the value of  $(x-y)^2$ .

5. Having given

$$x^2 = y^2 + z^2 - 2ayz$$

$$y^2 = z^2 + x^2 - 2bzx$$

$$z^2 = x^2 + y^2 - 2cxy,$$

Show that  $\frac{x^2}{1-a^2} = \frac{y^2}{1-b^2} = \frac{z^2}{1-c^2}$ .

6.  $\frac{1+x+\sqrt{(2x+x^2)}}{1-x+\sqrt{(2x+x^2)}} = 1-ax$ .

7. Determine  $x$  in terms of  $a$  and  $b$  in order that  $x^4 + 2ax^3 + 3b^2x^2 - 4a^3x + 4b^4$  may be a perfect square.

8. A company of 90 persons consists of men, women, and children; the men are 4 in number more than the women, and the children exceed the number of men and women by 10. How many men, women, and children are there in the company.

## XX.

1. Divide  $(1+m)x^3 - (m+n)xy(x-y) - (n-1)y^3$  by  $x^2 - xy + y^2$ .

2. If  $x^3 + px^2 + qx + r$  is exactly divisible by  $x^2 + mx + n$ , then  $nq - n^2 = rm$ .

3. Prove that if  $m$  be a common measure of  $p$  and  $q$ , it will also measure the difference of any multiples of  $p$  and  $q$ .

Find the G.C.M. of  $x^4 - px^3 + (q-1)x^2 + px - q$  and  $x^4 - qx^3 + (p-1)x^2 + qx - p$ .

4. Prove the rule for multiplication of fractions.

Simplify  $\frac{x^2 - (y-z)^2}{(y+z)^2 - x^2} \times \frac{y^2 - (z-x)^2}{(z+x)^2 - y^2} \times \frac{z^2 - (x-y)^2}{(x+y)^2 - z^2}$ ;

and  $\frac{a}{a^2 + b^2} - \frac{a}{a^2 - b^2} + \frac{a^2}{(a-b)(a^2 + b^2)} - \frac{2a^3 - b^3 - ab^3}{a^4 - b^4}$ .

5. What is the distinction between an *identity* and an *equation*?  
If  $x - a = y + b$ , prove  $x - b = y + a$ .

Solve the equation

$$\frac{16x-13}{4x-3} + \frac{40x-43}{8x-9} = \frac{32x-30}{8x-7} + \frac{20x-24}{4x-5}.$$

6. What are *simultaneous equations*? Explain why there must be given as many independent equations as there are unknown quantities involved. If there is a greater number of equations than unknown quantities, what is the inference?

Eliminate  $x$  and  $y$  from the equations  $ax + by = c$ ,  $a'x + b'y = c'$ ,  $a''x + b''y = c''$ .

7. Solve the equations—

$$(1) \quad \sqrt[3]{(n+x)} + \sqrt[3]{(n-x)} = m.$$

$$(2) \quad 3x + y + z = 13, \quad 3y + z + x = 15, \quad 3z + x + y = 17.$$

8. A person has two kinds of foreign money; it takes  $a$  pieces of the first kind to make one £, and  $b$  pieces of the second kind: he is offered one £ for  $c$  pieces, how many pieces of each kind must he take?

9. A person starts to walk to a railway station four and a-half miles off, intending to arrive at a certain time; but after walking a mile and a half he is detained twenty minutes, in consequence of which he is obliged to walk a mile and a half an hour faster in order to reach the station at the appointed time. Find at what pace he started.

10. (a) If  $\frac{a}{b} = \frac{c}{d}$  then will  $\frac{a^4+c^4}{b^4+d^4} = \frac{a^2c^2}{b^2d^2}$ .

(b) Find by Horner's method of division the value of  $x^5 + 290x^4 + 279x^2 - 2892x^2 - 586x - 312$  when  $x = -289$ .

(c) Show without actual multiplication that  $(a+b+c)^3 - (a+b+c)(a^2+ab+b^2-bc+c^2-ac) - 3abc = 3(a+b)(b+c)(c+a)$ .

NOTE.—In Ex. 6, p. 87, after proving that  $a+b+c$  is a factor, we may proceed as follows to discover the remaining quadratic factor:

The quadratic factor must be of the form

$$m(a^2+b^2+c^2) + n(ab+bc+ca),$$

in which  $m$  and  $n$  are independent, being either zero, or a positive or negative number. To determine them put  $c=0$ , then the given expression gives

$$\{a^3+b^3+3ab(a+b)\} \div (a+b) = a^2+b^2+2ab.$$

$$\text{but also} = m(a^2+b^2) + nab. \quad \therefore m=1 \text{ and } n=2.$$

$$\therefore a^3+b^3+c^3+3(a+b)(b+c)(c+a) \div (a+b+c) =$$

$$a^2+b^2+c^2+2(ab+bc+ca) = (a+b+c)^2.$$

+n, then

t will also

2  
2

$$\frac{3-ab^3}{b^4}$$

equation?

here must  
unknown  
equations

$$x+by=c.$$

17.

es a pieces  
cond kind:  
each kind

## XXI.

1. Find the value of  $x^3 - \left(\frac{1}{a} + \frac{1}{b}\right)x^2 + \left(\frac{1}{b} - \frac{1}{a}\right)x + \frac{2}{b^2}$  when  $a = \frac{1}{2}$ ,  $b = \frac{1}{3}$ ,  $x = 2$ . Simplify

$$a^3 - [x^3 - \{(3a^2x - 3ax^2) + a^3\} + 2ax^3] + x^3.$$

2. Find, by symmetry, the sum of  $(a+b+c)^3 - (a+b-c)^3 - (a-b+c)^3 - (b-a+c)^3$ , and of  $(a^4 - 4a^3x + 3a^2x^2 - 2ax^3 + 3x^4)^2$  and  $(a^4 + 4a^3x + 3a^2x^2 + 2ax^3 + 3x^4)^2$ .

3. Explain and illustrate the signs  $>$ ,  $<$

Prove:  $x^2 + y^2 > 2xy$ ,  $(x+y+z)^2 > 3(xy+yz+zx)$ , and

$$x^3 + y^3 + z^3 > 3xyz.$$

4. Determine the value of  $x+y-z+3x^{\frac{1}{3}}y^{\frac{1}{3}}z^{\frac{1}{3}}$ , when  $x^{\frac{1}{3}}+y^{\frac{1}{3}}-z^{\frac{1}{3}}=0$ , &c.; of  $a^7 + 7ax^3 + 8x^2 - 3a^2 - (x^4 + 7ax^3 - 8x^2 - 3a^2)$ , when  $x = -1$ .

5. Show that  $(a^m)^{\frac{p}{q}} = a^{\frac{mp}{q}}$ .

Simplify  $\left\{\left(\frac{1}{16}\right)^{\frac{1}{2}}\right\}^{-\frac{1}{2}} \times \left(\frac{1}{8}\right)^{-\frac{1}{3}} \times \sqrt[4]{256}$ , and divide

$$x^{\frac{5}{2}} - 6ax^{\frac{3}{2}} + 5a^2x^{\frac{1}{2}} + 2a^2x^{\frac{1}{2}} - 2a^{\frac{5}{2}} \text{ by } x^{\frac{3}{2}} - 2a^{\frac{1}{2}}x + a^{\frac{3}{2}}.$$

6. If  $u = \frac{1}{2}\left(x + \frac{1}{x}\right)$  and  $v = \frac{1}{2}\left(y + \frac{1}{y}\right)$  prove that

$$uv - \sqrt{(1-u^2)}\sqrt{(1-v^2)} = \frac{1}{2}\left(xy + \frac{1}{xy}\right).$$

7. Gold is  $19\frac{1}{2}$  times as heavy as water, and silver  $10\frac{1}{2}$  times. A mixed mass weighs 4,160 ounces, and displaces 250 ounces of water. What proportion of gold and silver does the mass contain?

8. Shew that  $1 + px + qx^2 + rx^3$  is a perfect cube if  $p^2 = 3q$ , and  $q^2 = 3pr$ .

9. Solve the equations:

$$(1) \sqrt{\frac{x-2}{x+2}} + \sqrt{\frac{x+2}{x-2}} = 4.$$

$$(2) (x^4 + y^4)^2 + x^2y^2(x^2 - y^2)^2 + x^2 - y^2 = 328, \quad x^2 - y^2 = 3.$$

$$(3) \frac{x^2}{y^2} + \frac{2x+y}{\sqrt{y}} = 20 - \frac{y^2+x}{u}, \quad x+8=4y.$$

10. A person buys two bales of cloth, each containing 80 yards, for \$240. By selling the first at a gain of as much per cent. as the second cost him, and the second at a loss of as much per cent., he makes a profit of \$16 on the whole. Find the cost price per yard of each bale.

SECOND CLASS TEACHERS, 1880.

XXII.

1. Find the value of  $x^5 + x^4 - 166x^3 - 166x^2 + 81x + 81$  when  $x = -7$ ; and the value of  $x^3 - 3px^2 + (3p^2 + q)x - pq$  when  $x = a + p$ . (Arrange the latter result according to powers of  $a$ ).

2. What is the condition that  $x + b$  shall be a factor of  $ax^2 + bx + c$ ?

Find the factors of

(a).  $(a^2 - ab) + 2(b^2 - ab) + 3(c^2 - b^2) + 4(a - b)^2$ ; and

(b).  $(ax + b)(bx + c)(cx + a) - (ax + c)(bx + a)(cx + b)$ .

3. What must be the relation among  $a, b, c$ , that  $ax^2 + bx + c$  may be a perfect square?

(a). Extract the square root of

$$(a - b)^4 - 4(a^2 + b^2)(a - b)^2 + 4(a^4 + b^4) + 8a^2b^2.$$

(b). If 5 be subtracted from the sum of the squares of any four consecutive numbers, the remainder will be a perfect square. (Prove this.)

4. If  $\frac{a}{b} = \frac{c}{d} = \frac{e}{f}$  and  $\frac{h}{k} = \frac{l}{m} = \frac{n}{p}$

prove that  $\frac{(a+c+e)(h+l+n)}{(b+d+f)(k+m+p)} = \frac{ah+cl+en}{bk+dm+fp}$ .

5. (a). Reduce  $\frac{ab(x^2 - y^2) + xy(a^2 - b^2)}{ab(x^2 + y^2) + xy(a^2 + b^2)}$  to its lowest terms.

(b). If  $xy + yz + zx = 1$  prove that

$$1 - x^2 + \frac{y}{1 - y^2} + \frac{z}{1 - z^2} = \frac{4xyz}{(1 - x^2)(1 - y^2)(1 - z^2)}.$$



6. Prove that

$$(a) \frac{2\{x+2+\sqrt{(x^2-4)}\}}{x+2-\sqrt{(x^2-4)}} = x+\sqrt{(x^2-4)}.$$

$$(b) (b+c-a)a^{\frac{1}{2}}+(c+a-b)b^{\frac{1}{2}}+(a+b-c)c^{\frac{1}{2}} = (a+b+c)(a^{\frac{1}{2}}+b^{\frac{1}{2}}+c^{\frac{1}{2}})-2(a^{\frac{3}{2}}+b^{\frac{3}{2}}+c^{\frac{3}{2}}).$$

7. Solve the equations—

$$(a). (b-c)(x-a)^3+(c-a)(x-b)^3+(a-b)(x-c)^3=0.$$

$$(b). x+y=4xy; y+z=2yz; z+x=3zx.$$

$$(c). x+y+z=0.$$

$$ax+by+cz=0.$$

$$bcx+cay+abz+(a-b)(b-c)(c-a)=0.$$

$$(d) \frac{x-1}{x+3} + \frac{x-3}{x+1} + 2=0.$$

### FIRST-CLASS TEACHERS, 1876.

#### XXIII.

1. Investigate Horner's method of division.

Divide  $x^9 - 3x^8 - 31x^7 + 25x^6 + 3x^5 - 8x^3 + 19x^2 + 8x + 10$  by  $3x^4 - 21x^3 + 9x - 6$ , showing the "final remainder."

Find the value of  $2x^5 + 803x^4 - 398x^3 + 1605x^2 - 1204x + 122$ , when  $x = -402$ .

2. If  $f(x)$ , a rational and integral function of  $x$  is divided by  $x^2 + px + q$ , the remainder is  $\frac{\{f(a) - f(\beta)\}x + a f(\beta) - \beta f(a)}{a - \beta}$ ,

where  $a, \beta$  are the roots of  $x^2 + px + q = 0$ .

Examine the case where  $p^2 = 4q$ .

3. Show without actual expansion that

$$\frac{a^4(b^2 - c^2) + b^4(c^2 - a^2) + c^4(a^2 - b^2)}{a^2(b - c) + b^2(c - a) + c^2(a - b)} =$$

$$\frac{(a^2 - b^2)^3 + (b^2 - c^2)^3 + (c^2 - a^2)^3}{(a - b)^3 + (b - c)^3 + (c - a)^3}.$$

4. Find the value of  $x$  and  $y$  that will render the fraction  $\frac{2z^2 + (x - a)z + 2b(x - 2c)}{3z^2 + (y - b)z + 2a(y - 3c)}$  the same for all values of  $z$ .

5. Show how to find the sum of  $n$  terms of a series in Geometric progression.

(1) Show that the sum of  $n$  terms of the series  $1 + r + (1 + 2r)(1 + r) + (1 + 3r)(1 + r)^2 + \dots$ , is  $n(1 + r)^n$ .

(2) Sum to infinity the series  $\frac{1}{2 \cdot 4 \cdot 6} + \frac{1}{4 \cdot 6 \cdot 8} + \frac{1}{6 \cdot 8 \cdot 10} + \dots$

6. Explain the notation of functions: prove that if

$f(m) = 1 + mx + \frac{m(m-1)}{1 \cdot 2}x^2 + \dots$ , then  $f(m) \times f(n) = f(m+n)$ .

Show that in the expansion of  $(1+x)^n$  the sum of the squares of the co-efficients =  $\frac{1 \cdot 2 \cdot 3 \dots 2n}{(1 \cdot 2 \cdot 3 \dots n)^2}$ .

7. Solve the equations—

(1)  $\frac{x-a}{b+c} + \frac{x-b}{a+c} + \frac{x-c}{a+b} = 3$ .

(2)  $x^4 - 10x^3 + 35x^2 - 50x + 24 = 0$ .

(3)  $\frac{1}{21x^2 - 13x + 2} + \frac{1}{28x^2 - 15x + 2} = 12x^2 - 7x + 1$ .

8. Give a brief account of mathematical induction, and show that a square of a multinomial is equal to the square of each term together with twice the product of each term into the sum of all that follow it.

Find the sum of the products of the first  $n$  natural numbers taken two and two together?

9. If  $\frac{x}{a} = y + z$ ,  $\frac{y}{b} = z + x$ ,  $\frac{z}{c} = x + y$ , prove

(1)  $\frac{1}{a} \cdot \frac{1}{b} \cdot \frac{1}{c} = \frac{1+a}{1-ab} \cdot \frac{1+b}{1-bc} \cdot \frac{1+c}{1-ca}$ .

(2)  $\frac{x^2}{a(1-bc)} = \frac{y^2}{b(1-ca)} = \frac{z^2}{c(1-ab)}$ .

(3)  $\frac{\sqrt{1-bc}}{a} + \frac{\sqrt{1-ca}}{b} + \frac{\sqrt{1-ab}}{c} = \frac{\sqrt{1-bc}}{a} \cdot \frac{\sqrt{1-ca}}{b} \cdot \frac{\sqrt{1-ab}}{c}$

10.  $AB$  is divided in  $C$ , so that  $AB, BC = AC^2$ ; from  $CA$  is cut off a part  $CD$  equal to  $CB$ ; from  $DC$  is cut off a part  $DE$  equal to  $DA$ ; from  $ED$  is cut off a part equal to  $EC$ , and so on *ad inf.* Show that the points of section continually approach a point  $C'$  such that  $AC' = BC$ .

14. Eliminate  $x, y, z$  and  $u$  from the equations

$$a_1x + b_1y + c_1z + d_1u = 0.$$

$$a_2x + b_2y + c_2z + d_2u = 0.$$

$$a_3x + b_3y + c_3z + d_3u = 0.$$

$$a_4x + b_4y + c_4z + d_4u = 0.$$

12. A railway train travels from Toronto to Collingwood. At Newmarket it stops 7 minutes for water, and two minutes after leaving the latter place it meets a special express that left Collingwood when the former was 28 miles on the other side of Newmarket; the express travels at double the rate of the other, and runs the distance from Collingwood to Newmarket in  $1\frac{1}{2}$  hour; and if on reaching Toronto it returned at once to Collingwood, it would arrive there three minutes after the first train; find the distance between Toronto, Newmarket and Collingwood.

### FIRST CLASS TEACHERS, 1877.

#### XXIV.

1. Simplify  $\left\{ \left( \frac{x+y}{x-y} \right)^2 + 1 \right\} \left\{ \left( \frac{x+z}{x-z} \right)^2 + 1 \right\} \left\{ \left( \frac{y+z}{y-z} \right)^2 + 1 \right\} \times$   

$$\frac{x^2(y-z) + y^2(z-x) + z^2(x-y)}{x^4y^2 + x^2y^4 + x^4z^2 + x^2z^4 + y^4z^2 + y^2z^4 + 2x^2y^2z^2}$$

2. Solve (1.)  $\frac{ax+m+1}{ax+m-1} + \frac{ax+n}{ax+n-2} = \frac{ax+m}{ax+m-2} + \frac{ax+n+1}{ax+n-1}$ .

(2.)  $\sqrt[3]{1+\sqrt{x}} + \sqrt[3]{1-\sqrt{x}} = 2$ .

3.  $A, B,$  and  $C$  start from the same place;  $B,$  after a quarter of an hour, doubles his rate, and  $C,$  after walking 10 minutes, diminishes his rate one-sixth; at the end of half an hour,  $A$  is a quarter of a mile before  $B,$  and half a mile before  $C,$  and it is

observed that the total distance walked by the three, had they continued to walk uniformly from the first, is  $6\frac{1}{4}$  miles. Find the original rate of each.

4. (1) Investigate the relations that must exist between the constants in order that  $Ax^2 + By^2 + Cz^2 + ayz + bxz + cxy$  shall be a perfect square.

(2) Find the conditions that the values of  $x$  and  $y$  derived from the equations  $ax + by = \frac{a^3}{x} + \frac{b^3}{y} = c^2$  may be rational.

5. If  $x^2 + px + q$  and  $x^2 + mx + n$  have a common factor, then will  $(n - q)^2 + n(n - p)^2 = m(m - p)(n - q)$ .

6. Prove  $(a^n)^m = a^{mn}$ , whether  $m$  and  $n$  be positive or negative, integral or fractional.

Show that  $(x^{2m} + x^{2n})^{\frac{1}{mn}} = x^{\frac{1}{m}} + \frac{1}{n} \times (x^{m-n} + x^{n-m})^{\frac{1}{mn}}$

7. (1.) If  $\frac{a}{b} = \frac{c}{d}$  then  $\sqrt{\frac{a^{2n} + b^{2n}}{c^{2n} + d^{2n}}} = \left(\frac{a - b}{c - d}\right)^{\frac{n}{2}}$

(2.) If  $\frac{a^n d^n - b^n c^n}{\frac{1}{2}n(a^n - b^n - c^n + d^n)} = \frac{a^n c^n - b^n d^n}{\frac{1}{2}n(a^n - b^n - d^n + c^n)}$ , then each

of these fractions =  $\frac{1}{n} (a^n + b^n + c^n + d^n)$ .

8. If  $x$  be very small, show that—

$$\frac{(1 + 2x)^{\frac{1}{2}} + (1 + 3x)^{\frac{1}{3}}}{2 + 5x - (1 + 4x)^{\frac{1}{2}}} = 2 - 4x, \text{ very nearly.}$$

9. Prove that  $1 - n^2 + \frac{n^2(n^2 - 1^2)}{1^2 \cdot 2^2} - \frac{n^2(n^2 - 1^2)(n^2 - 2^2)}{1^2 \cdot 2^2 \cdot 3^2} + \dots = 0$

10. If a debt \$ $a$  at compound interest be discharged in  $n$  years by annual payment of \$ $\frac{a}{m}$ , show that  $(1 + r)^n(1 - mr) = 1$ , where  $r$  is the interest on \$1 for a year.

11. Solve—(1.)  $3x^2 - 2xy = 55$ .

$$x^2 - 5xy + 8y^2 = 7.$$

$$(2) \frac{5}{x^2 - 7x + 10} + \frac{5}{x^2 - 13x + 40} = x^2 - 10x + 19$$

$$(3) a^2 b^2 x^{\frac{1}{2}} - 4a^{\frac{3}{2}} b^{\frac{3}{2}} x^{\frac{n+q}{2}} = (a-b)^2 x^{\frac{1}{2}}$$

FIRST CLASS TEACHERS, 1878.

XXV.

1. Simplify  $\left(\sqrt{\frac{a+x}{x}} - \sqrt{\frac{x}{a+x}}\right)^2 - \left(\sqrt{\frac{x}{a}} - \sqrt{\frac{a}{x}}\right)^2 - \frac{x^2}{a(a-x)}$

and  $\frac{x^2 - (y-z)^2}{(x+z)^2 - y^2} + \frac{y^2 - (z-x)^2}{(x+y)^2 - z^2} + \frac{z^2 - (x-y)^2}{(y+z)^2 - x^2}$

2. Divide  $\frac{x}{a} - 1 - \frac{b}{a} - \frac{b^2}{a^2} + \frac{b}{x} + \frac{b^2}{x^2}$  by  $x-a$ ;

shew that  $(-9a^2)^{\frac{1}{2}} = \frac{1}{2} \{ \sqrt{(6a)} + \sqrt{(-6a)} \}$ .

3. If  $\frac{m}{x} = \frac{n}{y} = \frac{r}{z}$  and  $\frac{x^2}{a^2} + \frac{y^2}{b^2} + \frac{z^2}{c^2} = 1$ , prove that

$$\frac{m^2}{a^2} + \frac{n^2}{b^2} + \frac{r^2}{c^2} = \frac{m^2 + n^2 + r^2}{x^2 + y^2 + z^2}$$

4. Find the relations between the roots and coefficients of the equation  $ax^2 + bx + c = 0$ .

If  $m$  and  $n$  are the roots of the equation  $ax^2 + bx + c = 0$ , show that the roots of the equation  $acx^2 + (2ac - b^2)x + ac = 0$  are

$$\frac{m}{n} \text{ and } \frac{n}{m}.$$

5. Solve the equations:

(1)  $x^2 + 2\sqrt{x^2 - 2x} = 2x + 8$ .

(2)  $\frac{x^3}{y} - \frac{y^3}{x} = 10\frac{5}{8}, \frac{x}{y} - \frac{y}{x} = \frac{5}{8}$ .

(3)  $xz = y^2, x + y + z = 12, x^2 + y^2 + z^2 = 91$ .

6. Two men start at the same time to meet each other from towns which are 28 miles apart; one takes five minutes longer than the other to walk a mile, and they meet in four hours. Find each man's rate per hour.

7. If  $P$ ,  $Q$ ,  $R$  be respectively the  $p$ th,  $q$ th,  $r$ th terms of a G.P., show that

$$P^{q-r} \times Q^{r-p} \times R^{p-q} = 1.$$

Sum to infinity the series  $\frac{1}{x} + \frac{2}{x^2} + \frac{3}{x^3} + \dots$  &c.

8. Find the amount of \$ $P$  at compound interest for  $n$  years,  $r$  being the interest on \$1 for one year.

Supposing \$ $p$  to be withdrawn at the end of each year, what will be the amount at the end of  $n$  years?

9. Determine the number of combinations of  $n$  things taken  $r$  together.

The number of combinations of  $n$  things taken two together exceeds by 6 the number of combinations of  $n-1$  things taken two together: find  $n$ .

10. (1) Find the limit of  $(1 + \frac{1}{x})^x$  when  $x$  increases without limit.

(2) Find the  $(r+1)$ th term in the expansion of  $(3-5x)^{-\frac{1}{2}}$ .

11. Determine the limits between which lies  $\frac{x^2 - 3x - 3}{2x^2 + 2x + 1}$  for all possible values of  $x$ .

---

FIRST CLASS TEACHERS, 1879.

---

XXVI.

7. Prove that  $2\{(a-b)^2 + (b-c)^2 + (c-a)^2\} = 7(a-b)(b-c)(c-a)\{(a-b)^4 + (b-c)^4 + (c-a)^4\}$ .

2. Extract the square root of  $ab - 2a\sqrt{(ab-a^2)}$ , and find the simplest real forms of the expression

$$\sqrt{3+4\sqrt{-1}} + \sqrt{3-4\sqrt{-1}}.$$

3. Solve the equations :

$$(1). 2x^4 + x^3 - 11x^2 + x + 2 = 0.$$

$$(2). x^2 + y^2 + z^2 = a^2$$

$$yz + zx + xy = b^2$$

$$z + y - x = c.$$

$$(3). \sqrt{(x^2 + 5x + 4)} + \sqrt{(x^2 + 3x - 4)} = x + 4.$$

4. Prove that the number of positive integral solutions of the equation  $ax + by = c$  cannot exceed  $\frac{c}{ab} + 1$ .

In how many ways may £11 15s. be paid in half-guineas and half-crowns?

5. If  $xy = ab(a + b)$ , and  $x^2 - xy + y^2 = a^3 + b^3$ , shew that

$$\left(\frac{x}{a} - \frac{y}{b}\right) \left(\frac{x}{b} - \frac{y}{a}\right) = 0.$$

6. Given the sum of an arithmetical series, the first term, and the common difference, shew how to find the number of terms. Explain the negative result. *Ex.* How many terms of the series 6, 10, 14, &c., amount to 96?

7. Find the relation between  $p$  and  $q$ , when  $x^3 + px + q = 0$  has two equal roots, and determine the values of  $m$  which will make  $a^2 + max + a^2$  a factor of  $x^4 - ax^3 + a^2x^2 - a^3x + a^4$ .

8. In the scale of relation in which the radix is  $r$ , shew that the sum of the digits divided by  $r - 1$  gives the same remainder as the number itself divided by  $r - 1$ .

9. Assuming the Binomial Theorem for a positive integral index, prove it in the case of the index being a positive fraction.

Shew that the sum of the squares of the co-efficients in the expansion of  $(1 + x)^n$  is  $\frac{2n}{3} + \frac{1}{3}n^2$ ,  $n$  being a positive integer.

10. Sum the following series :—

$$(1.) 1 + 3x + 5x^2 + 7x^3 + \&c. \text{ to } n \text{ terms.}$$

$$(2.) \frac{1}{3 \times 8} + \frac{1}{8 \times 13} + \&c. \text{ to } n \text{ terms, and to infinity.}$$

11. Shew that  $\begin{vmatrix} bc, & -ac, & -ab \\ b^2 - c^2, & a^2 + 2ac, & -a^2 - 2ab \\ c^2, & c^2, & (a+b)^2 \end{vmatrix}$  is divisible by  $abc(a+b+c)$ .

FIRST CLASS TEACHERS, 1880—GRADE C.

XXVII.

1. If in  $ax^2 + 2bxy + cy^2$ ,  $ku + lv$  be substituted for  $x$  and  $mu + nv$  for  $y$ , the result takes the form  $Au^2 + 2Buv + Cv^2$ . Find the value of  $(B^2 - AC) \div (b^2 - ac)$  in terms of  $k, l, m, n$ .

2. Resolve  $a(b-c)^3 + b(c-a)^3 + c(a-b)^3$  into factors.

Prove that  $\frac{Au^3 + Bv^3 + Cw^3}{uvw} = \frac{Ax^3 + By^3 + Cz^3}{xyz}$

if  $u = x(By^3 - Cz^3)$ ,  $v = y(Cz^3 - Ax^3)$ ,  $w = z(Ax^3 - By^3)$ .

3. Extract the square root of

$$(a-b)^2(b-c)^2 + (b-c)^2(c-a)^2 + (c-a)^2(a-b)^2,$$

and the cube root of

$$A(a-b)^6 + (b-c)^6 + (c-a)^6 - 3(a-b)^2(b-c)^2(c-a)^2.$$

4. Eliminate  $x, y, z$  from

$$ax + by + cz = 1 \quad \frac{a}{x} = \frac{b}{y} = \frac{c}{z}$$

$$k(x^2 + y^2 + z^2) + 2(lx + my + nz) + h = 0.$$

5. Simplify  $\frac{a\sqrt{b} + b\sqrt{a}}{\sqrt{a} + \sqrt{b}}$ ,  $\{\sqrt{(4+3j)} + \sqrt{(4-3j)}\}^2$ ,

$$\text{and } \left(\frac{-1+j\sqrt{3}}{2}\right)^2 + \frac{-1+j\sqrt{3}}{2} + 1,$$

in which  $j = \sqrt{-1}$ .

6. Given the first term, the common difference and the number of terms of an arithmetical progression, find (i.) the sum of the terms, (ii.) the sum of the squares of the terms.



7. Solve the equations

$$(i.) (a-x)^3 = (x-b)^3.$$

$$(ii.) ax + by = \frac{a}{x} + \frac{b}{y} = 1.$$

$$(iii.) x(y+z^{-1}) = a, y(z+x^{-1}) = b, z(x+y^{-1}) = c.$$

8. What value (other than 1) must be given to  $q$  that one of the roots of  $x^2 - 2x + q = 0$ , may be the square of the other.

If  $a, b, c$  are the roots of  $x^3 - px^2 + qx - r$ , express

$$\frac{2a^2b^2 + 2b^2c^2 + 2c^2a^2 - a^4 - b^4 - c^4}{2ab + 2bc + 2ca - a^2 - b^2 - c^2}$$

in terms of  $p, q$  and  $r$ .

9. A vessel makes two runs on a measured mile, one with the tide in  $m$  minutes and one against the tide in  $n$  minutes. Find the speed of the vessel through the water, and the rate the tide was running at, assuming both to be uniform.

10. Five points,  $A, B, C, O$  and  $P$  lie on a straight line. The distances of  $A, B$ , and  $C$ , measured from the point  $O$ , are  $a, b$ , and  $c$ ; their distances measured from the point  $P$  are  $x, y, z$ . Prove that whatever be the positions of the points  $O$  and  $P$ ,

$$x^2(b-c) + y^2(c-a) + z^2(a-b) + (b-c)(c-a)(a-b) = 0.$$

The remain

$D'A$

let  $f_1(x)$  degree duct of produce

But  $R$  merely Take the does so dividen

It ha formed ing  $R$  to not equ values of that it such ca divisor althoug not equ objectio braic or sion by proof de not ope

## APPENDIX.

### SECTION I.—ELEMENTARY THEOREMS ON POLYNOMES.

(See page 39, *et seq.*)

**Theorem I.** If the polynome  $f(x)^n$  be divided by  $x-a$ , the remainder will be  $f(a)^n$ .

*D'Alembert's Proof.*  $f(x)^n$  is the dividend,  $x-a$  is the divisor; let  $f_1(x)^{n-1}$  be the quotient, which is necessarily a polynome of degree  $n-1$ , and let  $R$  be the remainder. Then, since the product of the quotient and the divisor added to the remainder reproduces the dividend,

$$f(x)^n = (x-a)f_1(x)^{n-1} + R.$$

But  $R$  does not contain  $x$ , hence it will remain the same, not merely in form but in actual value, whatever value be given to  $x$ . Take the case  $x=a$ , then  $(x-a)f_1(x)^{n-1}$  vanishes for its factor  $x-a$  does so, hence  $R=f(a)^n$ . Thus the remainder is the value of the dividend when  $x$  has the value which makes the divisor vanish.

It has been objected to the above proof "Division can be performed only when there is an actual divisor, therefore in assuming  $R$  to be the remainder of  $f(x)^n \div (x-a)$  it is assumed that  $x$  is not equal to  $a$ , and although  $R$  will remain unchanged for all values of  $x$  that fulfil this assumption, it cannot thence be inferred that it will do so if the contradictory assumption be made. In such case the only legitimate conclusion is that there being no divisor there is neither quotient nor remainder. Therefore, although  $f(a)^n$  may be the remainder in the case in which  $x$  is not equal to  $a$ , yet the above argument does not prove it." This objection confuses arithmetical or numerical division with algebraic or formal division, division by a definite quantity with division by an undetermined or variable quantity. The following proof does not involve the assumption  $x=a$ , and consequently is not open to the foregoing objection.

*Lagrange's Proof. Lemma.*  $x^n - a^n$  is divisible by  $x - a$ , if  $n$  be a positive integer.

$$\text{By actual division } \frac{x^n - a^n}{x - a} = x^{n-1} - a, \frac{x^{n-1} - a^{n-1}}{x - a};$$

$\therefore x^n - a^n$  is divisible by  $x - a$  if  $x^{n-1} - a^{n-1}$  is so divisible, hence  $x^{n-1} - a^{n-1}$  " "  $x - a$  "  $x^{n-2} - a^{n-2}$  . . . .

Thus we can reduce the exponent unit by unit until at last we arrive at,  $x^2 - a^2$  is divisible by  $x - a$  if  $x - a$  is so divisible. But  $x - a$  is certainly divisible by itself,  $\therefore x^2 - a^2$  is divisible by  $x - a$ ,  $x^3 - a^3$  is also divisible by  $x - a$ ,  $\therefore$  so also is  $x^4 - a^4$  and thus we may go on to any positive integral exponent whatsoever.

**Theorem.** Writing  $f(x)^n$  in polynomial form arranged in ascending powers of  $x$ ,

$$\begin{aligned} f(x)^n &= A_0 + A_1x + A_2x^2 + A_3x^3 + \dots + A_nx^n, \\ \therefore f(a)^n &= A_0 + A_1a + A_2a^2 + A_3a^3 + \dots + A_na^n, \\ \therefore f(x)^n - f(a)^n &= A_1(x-a) + A_2(x^2 - a^2) + A_3(x^3 - a^3) + \dots \\ &\quad + A_n(x^n - a^n). \end{aligned}$$

But every term of this polynomial is divisible by  $x - a$ , and the highest power of  $x$  in the quotient is  $x^{n-1}$  got from the term  $A_n(x^n - a^n)$ , so the quotient may be represented by  $f_1(x)^{n-1}$ ,

$$\begin{aligned} \therefore \{f(x)^n - f(a)^n\} \div (x - a) &= f_1(x)^{n-1} \\ \text{or } \frac{f(x)^n}{x - a} &= f_1(x)^{n-1} + \frac{f(a)}{x - a} \end{aligned}$$

**Theorem II.** If the polynome  $f(x)^n$  vanish on substituting for  $x$  each of the  $n$  different values  $a_1, a_2, a_3, \dots, a_n$ , then  $f(x)^n = A(x - a_1)(x - a_2) \dots (x - a_n)$ , in which  $A$  is independent of  $x$  and consequently is the coefficient of  $x^n$  in  $f(x)^n$ .

Since  $f(a_1) = 0$ ,  $\therefore f(x)^n = (x - a_1)f(x)^{n-1}$ . In this substitute  $a_2$  for  $x$ ,  $\therefore$  since  $f(a_2)^n = 0$ , it becomes  $0 = (a_2 - a_1)f_1(a_2)^{n-1}$ . Of this product the factor  $a_2 - a_1$  does not vanish since by hypothesis  $a_2$  is not equal to  $a_1$ , therefore the other factor  $f_1(a_2)^{n-1}$  must vanish that the product may vanish, and consequently  $f_1(x)^{n-1}$  is

divisible by  $x - a_2$ . Let the quotient be denoted by  $f_2(x)^{n-2}$ ,  $\therefore f(x)^n = (x - a_1)(x - a_2)f_2(x)^{n-2}$ . Substitute  $a_3$  for  $x$  and proceed as before, and it will be proved that  $x - a_3$  is a factor of  $f(x)^n$ . Continuing to  $n$  factors we get a quotient independent of  $x$ , since each division reduces the exponent of  $x$  by unity,  $\therefore$  finally

$$f(x)^n = A(x - a_1)(x - a_2) \dots (x - a_n).$$

Cor. If  $f(x)^n$  and  $\varphi(x)^m$  both vanish for the same  $n$  different values of  $x$ ,  $f(x)^n$  is algebraically divisible by  $\varphi(x)^m$ .

Let  $a_1, a_2, a_3, \dots, a_m$  be the  $m$  different values of  $x$  for which the polynomes vanish,

$$\therefore f(x)^n = (x - a_1)(x - a_2) \dots (x - a_m) F(x)^{n-m}$$

$$\text{and } \varphi(x)^m = A(x - a_1)(x - a_2) \dots (x - a_m)$$

$$\therefore f(x)^n \div \varphi(x)^m = F(x)^{n-m} \div A,$$

which is an integral function of  $x$  since  $A$  does not contain  $x$ .

**Theorem III.** If the polynome  $f(x)^n$  vanish for more than  $n$  different values of  $x$  it will vanish identically, the coefficient of every term being zero.

Let  $a_1, a_2, a_3, \dots, a_n, a_{n+1}$  be  $n+1$  different values of  $x$  for which  $f(x)^n$  vanishes,

$$\therefore f(x)^n = A(x - a_1)(x - a_2)(x - a_3) \dots (x - a_n)$$

Substitute  $a_{n+1}$  for  $x$ , and since  $f(a_{n+1})^n = 0$ ,

$$\therefore 0 = A(a_{n+1} - a_1)(a_{n+1} - a_2)(a_{n+1} - a_3) \dots (a_{n+1} - a_n),$$

But none of the factors  $a_{n+1} - a_1, a_{n+1} - a_2$ , &c. vanishes,

$\therefore A$  must be zero, or

$$f(x)^n = 0(x - a_1)(x - a_2)(x - a_3) \dots (x - a_n)$$

and the factor, zero, will be a factor in the coefficients of every term.

**Theorem IV.** If the polynomes  $f(x)^n, \varphi(x)^m$  ( $n$  not less than  $m$ ) are equal for more than  $n$  different values of  $x$ , they are equal for all values, and the coefficients of equal powers of  $x$  in each are equal to one another.

$$\begin{aligned}
 f(x)^n &= A_0 + A_1x + A_2x^2 + A_3x^3 + \dots + A_nx^n \\
 \varphi(x)^n &= B_0 + B_1x + B_2x^2 + B_3x^3 + \dots + B_mx^m, \\
 \therefore f(x)^n - \varphi(x)^n &= A_0 - B_0 + (A_1 - B_1)x + (A_2 - B_2)x^2 + \\
 &\quad (A_3 - B_3)x^3 + \dots + (A_m - B_m)x^m \\
 &\quad + A_{m+1}x^{m+1} + A_{m+2}x^{m+2} \dots + A_nx^n,
 \end{aligned}$$

and this is a polynome of degree  $n$  at most. But  $f(x)^n = \varphi(x)^n$  for more than  $n$  different values of  $x$ , that is  $f(x)^n - \varphi(x)^n$  vanishes for these values,  $\therefore$  by Theorem III.  $f(x)^n - \varphi(x)^n$  vanishes identically, and the coefficients  $A_0 - B_0, A_1 - B_1, A_2 - B_2, \dots, A_m - B_m, A_{m+2}, A_{m+2}, \dots, A_n$  are all equal to zero,

$$\therefore A_0 = B_0, A_1 = B_1, A_2 = B_2, \dots, A_m = B_m, A_{m+1} = 0, A_{m+2} = 0 \dots$$

*Note to Art. XVII.* To find, where such exist, the factors of

$$ax^2 + bxy + cax + ey^2 + gyz + hz^2.$$

Multiply by  $4a$

$$4a^2x^2 + 4abxy + 4acax + 4aey^2 + 4agyz + 4ahz^2.$$

Select the terms containing  $x$  and complete the square, thus

$$\begin{aligned}
 &4a^2x^2 + 4abxy + 4acax + b^2y^2 + 2bcax + c^2z^2 \\
 &\quad - (b^2 - 4ae)y^2 - 2(bc - 2ag)yz - (c^2 - 4ah)z^2 = \\
 &(2ax + by + cz)^2 - \{(b^2 - 4ae)y^2 + 2(bc - 2ag)yz + (c^2 - 4ah)z^2\}
 \end{aligned}$$

If the part within the double bracket is a square say  $(my + nz)^2$  the given expression can be written

$$(2ax + by + cz)^2 - (my + nz)^2$$

which can be factored by [4]. Factor and divide the result by  $4a$ . If the part within the double bracket is not a square, the given expression cannot be factored. If  $b$  and  $c$  are both even, multiply by  $a$  instead of by  $4a$  and the square can be completed without introducing fractions. If  $e$  is less than  $a$  it will be easier to multiply by  $4e$  instead of by  $4a$  and select the terms containing  $y$ . A similar remark applies to  $h$ .

This method can evidently be extended to quadratic multinomials of any number of terms.

## EXAMPLES.

1. Resolve
- $x^2 + xy + 2xz - 2y^2 + 7yz - 3z^2$
- into factors.

Multiply by 4

$$4x^2 + 4xy + 8xz - 8y^2 + 28yz - 12z^2$$

Complete the square selecting terms in  $x$ ,

$$4x^2 + 4xy + 8xz + y^2 + 4yz + 4z^2 - 9y^2 + 24yz - 11z^2 =$$

$$(2x + y + 2z)^2 - (3y - 4z)^2 =$$

$$\{(2x + y + 2z) + (3y - 4z)\} \{(2x + y + 2z) - (3y - 4z)\} =$$

$$(2x + 4y - 2z)(2x - 2y + 6z) = 4(x + 2y - z)(x - y + 3z)$$

 $\therefore$  the factors are  $(x + 2y - z)(x - y + 3z)$ .

- 2.
- $6a^2 - 7ab + 2ac - 20b^2 + 61bc - 48c^2$
- .

Multiply by  $4 \times 6 = 24$ 

$$144a^2 - 168ab + 48ac - 480b^2 + 1536bc - 1152c^2 =$$

$$(12a - 7b + 2c)^2 - 520b^2 + 1564bc - 1156c^2 =$$

$$(12a - 7b + 2c)^2 - (23b - 31c)^2 =$$

$$(12a + 16b - 32c)(12a - 30b + 36c) =$$

$$24(3a + 4b - 8c)(2a - 5b + 6c),$$

 $\therefore$  the factors are  $3a + 4b - 8c$  and  $2a - 5b + 6c$ .

- 3.
- $x^2 + 12xy + 2xz + 26y^2 - 8yz - 9z^2 =$

$$(x^2 + 12xy + 2xz + 36y^2 + 12yz + z^2) - 10y^2 - 20yz - 10z^2 =$$

$$(x + 6y + z)^2 - \{(y + z)\sqrt{10}\}^2 =$$

$$\{x + (6 + \sqrt{10})y + (\sqrt{10} + 1)z\} \times$$

$$\{x + (6 - \sqrt{10})y - (\sqrt{10} - 1)z\}$$

- 4.
- $3a^2 + 10ab - 14ac + 12ad - 8b^2 - 8bd + 8c^2 - 8cd$
- .

Multiply by 3, not  $4 \times 3$ , since the coefficients of the other terms in  $a$ , are all even,

$$9a^2 + 30ab - 42ac + 36ad - 24b^2 - 24bd + 24c^2 - 24cd.$$

Select the terms containing  $a$  and complete the square

$$\begin{aligned} & (3a+5b-7c+6d)^2 - 49b^2 + \\ & \quad 70bc - 84bd - 25c^2 + 60cd - 36d^2 = \\ & (3a+5b-7c+6d)^2 - (7b-5c+6d)^2 = \\ & (3a+12b-12c+12d)(3a-2b-2c) = \\ & 3(a+4b-4c+4d)(3a-2b-2c), \end{aligned}$$

$\therefore$  the factors are  $a+4b-4c+4d$  and  $3a-2b-2c$ .

Work Exercise XXIX by this method.

## SECTION II.—INDICES AND SURDS.

The general Index-laws are

$$a^{\frac{m}{n}} \cdot a^{\frac{p}{q}} = a^{\frac{m}{n} + \frac{p}{q}} \quad (1)$$

$$a^{\frac{m}{n}} \div a^{\frac{p}{q}} = a^{\frac{m}{n} - \frac{p}{q}} \quad (2)$$

$$(ab)^{\frac{m}{n}} = a^{\frac{m}{n}} \cdot b^{\frac{m}{n}} \quad (3)$$

$$(a \div b)^{\frac{m}{n}} = a^{\frac{m}{n}} \div b^{\frac{m}{n}} \quad (4)$$

$$\left(a^{\frac{m}{n}}\right)^{\frac{p}{q}} = a^{\frac{mp}{nq}} \quad (5)$$

The law connecting the Index and the Surd symbols is

$$a^{\frac{m}{n}} = \sqrt[n]{a^m} \quad (6)$$

[The indices  $\frac{1}{2}$ ,  $\frac{1}{3}$ ,  $\frac{1}{4}$ , &c., are generally used to denote 'either square-root,' 'any of the cube-roots,' 'any one of the fourth-roots,' &c.

The surd symbols  $\sqrt{\quad}$ ,  $\sqrt[3]{\quad}$ ,  $\sqrt[4]{\quad}$ , &c., are by some writers restricted to indicate the arithmetical or absolute roots, sometimes called the positive roots. Thus

$$\sqrt{4} = 2, \text{ but } 4^{\frac{1}{2}} = \pm 2, \therefore 4^{\frac{1}{2}} = \pm \sqrt{4}$$

Also,  $\sqrt{\{(-2)^2\}} = \sqrt{4} = 2$ .

$$\sqrt[3]{27} = 3, \text{ but } 27^{\frac{1}{3}} = 3 \text{ or } 3\left(\frac{-1 \pm j\sqrt{3}}{2}\right) \therefore 8^{\frac{1}{3}} = (1^{\frac{1}{3}}) \cdot 3^{\frac{2}{3}}$$

$$\sqrt[4]{16} = 2, \text{ but } 16^{\frac{1}{4}} = \pm 2 \text{ or } \pm 2j, \therefore 16^{\frac{1}{4}} = (1^{\frac{1}{4}}) \sqrt[4]{16}$$

With this restriction the general connecting formula would be

$$a^{\frac{m}{n}} = (1^{\frac{m}{n}})^{\sqrt[n]{a^m}}$$

In the following exercises this restriction need not be observed.]

## EXERCISE.

1. What is the arithmetical value of each of the following :

$$36^{\frac{1}{2}}, 27^{\frac{1}{3}}, 16^{\frac{1}{4}}, 32^{\frac{1}{5}}, 4^{\frac{1}{6}}, 8^{\frac{1}{7}}, 27^{\frac{1}{8}}, 64^{\frac{1}{9}}, 32^{\frac{1}{10}}, 64^{\frac{1}{11}}, 81^{\frac{1}{12}}, (3\frac{3}{8})^{\frac{1}{3}},$$

$$(5\frac{1}{16})^{\frac{1}{4}}, (1\frac{9}{16})^{\frac{1}{4}}, (.25)^{\frac{1}{2}}, (.027)^{\frac{1}{3}}, 49^{\cdot 5}, 32^{\cdot 2}, 81^{\cdot 75}$$

2. Interpret  $a^{-2}$ ,  $a^0$ ,  $a^2$ ,  $(a^2)^{-2}$ ,  $a^2^{-2}$ ,  $a^{-4}$ ,  $(a^{-4})^{-4}$ ,  $a^{\frac{1}{2}}$ ,  $a^{-\frac{1}{2}}$ .

3. What is the arithmetical value of

$$36^{-\frac{1}{2}}, 27^{-\frac{1}{3}}, (.16)^{-\frac{1}{2}}, (.0016)^{-\frac{1}{4}}, (\frac{1}{4})^{-\frac{1}{2}}, (\frac{4}{25})^{-\frac{1}{2}}, (\frac{9}{16})^{-\frac{1}{2}}, (5\frac{1}{16})^{-\frac{1}{4}}$$

4. Prove  $(a^m)^n = (a^n)^m$ ;  $(a^m)^{\frac{1}{n}} = (a^{\frac{1}{n}})^m$ ;  $a^{-m} = (a^{-1})^m$ ;  
and express these theorems in words.

5. Simplify  $a^{\frac{1}{2}} \cdot a^{\frac{1}{3}}, c^{\frac{1}{2}} \cdot c^{\frac{1}{5}}, m^{\frac{1}{2}} \cdot m^{-\frac{1}{3}}, n^{\frac{3}{4}} \cdot n^{-\frac{1}{2}}, (7\frac{1}{2})^{\frac{1}{3}} \cdot (2\frac{2}{3})^{\frac{1}{5}} \cdot (3\frac{1}{5})^{\frac{1}{3}}$

$$\frac{a^{\frac{1}{2}}}{a^{\frac{1}{3}}}, \frac{c^{\frac{5}{8}}}{c^{\frac{1}{2}}}, \frac{d^{\frac{1}{4}}}{d^{\frac{1}{2}}}, \frac{e^{-\frac{1}{2}}}{e^{\frac{1}{3}}}, \frac{e^{\frac{1}{2}}}{e^{-\frac{1}{3}}}, (2\frac{2}{5})^{\frac{3}{4}} \cdot (6\frac{3}{5})^{\frac{3}{4}} \div (\frac{1}{4})^{-1\frac{1}{2}}$$

6. Remove the brackets from

$$(a^6)^{\frac{1}{2}}, (b)^{-\frac{1}{3}}, (c^{\frac{1}{2}})^{-\frac{1}{3}}, (d^{\frac{2}{3}})^{\frac{1}{4}}, (e^{-\frac{1}{3}})^{\frac{2}{3}}, (f^{-\frac{2}{3}})^{-\frac{2}{3}}$$

$$(a^2b^2)^{\frac{1}{2}}, (a^{\frac{3}{4}}b^{\frac{6}{5}})^{\frac{5}{2}}, (a^2c^{-1})^{-\frac{1}{2}}, (a^{-5}c^{\frac{2}{5}})^{-\frac{1}{2}}, (x^{\frac{2}{3}}y^{-\frac{4}{5}})^{-6}$$

7. Remove the brackets and simplify

$$(x^{\frac{1}{2}-\frac{1}{3}})^{\frac{2}{3}} (x^{\frac{1}{2}-\frac{1}{3}})^{\frac{2}{3}} (x^{\frac{1}{2}-\frac{1}{3}})^{\frac{2}{3}}; x^{(\frac{1}{2}-\frac{1}{3})^{\frac{2}{3}}} x^{(\frac{1}{2}-\frac{1}{3})^{\frac{2}{3}}} x^{(\frac{1}{2}-\frac{1}{3})^{\frac{2}{3}}};$$

$$x^{\frac{1}{2} + \frac{1}{16}} \cdot x^{\frac{1}{2} - \frac{1}{16}}; x^{\frac{1}{2} + \frac{1}{16}} \cdot x^{\frac{1}{2} - \frac{1}{16}};$$

$$\{x^2^{-2} \div x^{-2}^{-2}\} \{x^{(-2)^2} \div x^{-2}^2\}.$$



$$8. \text{ Simplify } -x\{x^{-\frac{1}{2}}(-x)^{-1}\}^{\frac{1}{3}}, x\{(-x)^{-\frac{2}{3}}(-x)^{-2}\}^{-\frac{5}{3}}, \\ (-x)^{-1}\{x^{-3}x^{-\frac{3}{4}}\}^{\frac{3}{4}}$$

$$9. \text{ Determine the commensurable and the surd factors of } \\ 12^{\frac{1}{2}}, 24^{\frac{1}{3}}, 18^{-\frac{1}{2}}, (-81)^{\frac{1}{3}}, 12^{\frac{2}{3}}, 64^{\frac{3}{4}}, (\sqrt[4]{\pi})^{\frac{2}{3}}, (6\frac{3}{4})^{-\frac{2}{3}}.$$

(The surd factor must be the incommensurable root of an integer.)

$$10. \text{ Simplify } 8^{\frac{1}{3}} + 18^{\frac{1}{3}} - 50^{\frac{1}{3}}; 72^{\frac{1}{3}} + (\sqrt[4]{25})^{\frac{3}{4}} - (\sqrt[3]{25})^{-\frac{1}{3}};$$

$$\{(6+2^{\frac{1}{3}})(6-2^{\frac{1}{3}})\}^{\frac{1}{3}}; (2^{\frac{1}{3}}+3^{\frac{1}{3}})^2 + (2^{\frac{1}{3}}-3^{\frac{1}{3}})^2;$$

$$(2^{\frac{1}{3}}+3^{\frac{1}{3}})(4^{\frac{1}{3}}+9^{\frac{1}{3}}-6^{\frac{1}{3}}); (7^{\frac{1}{3}}-3^{\frac{1}{3}})^{\frac{1}{2}}(7^{\frac{1}{3}}+3^{\frac{1}{3}})^{\frac{1}{2}}.$$

$$[\{(a+x)(x+b)\}^{\frac{1}{2}} - \{(a-x)(x-b)\}^{\frac{1}{2}}]^2;$$

$$\{a^{\frac{2}{3}} + (a^3 - x^3)^{\frac{1}{2}}\}^{\frac{1}{3}} \cdot \{a^{\frac{2}{3}} - (a^3 - x^3)^{\frac{1}{2}}\}^{\frac{1}{3}}$$

Express as surds,

$$11. a^{\frac{2}{3}}, x^{\frac{5}{3}}, p^{\frac{3}{2}}, c^{-\frac{1}{2}}, h^{-\frac{2}{3}}.$$

$$12. x^{n+\frac{1}{2}}, y^{-n+\frac{2}{3}}, a^{.25}, b^{-n+\frac{1}{m}}$$

$$13. (ax-b)^{\frac{a}{b}}, (x^2-4x+1)^{\frac{m-3}{4}}, (p-qx)^{n-\frac{3}{2}}$$

Express with indices,

$$14. \sqrt[3]{a^2}, \sqrt[4]{c^3}, \sqrt[n]{x^m}, \sqrt[3]{y^{m-n}}, \sqrt[a]{(ax)}, \sqrt{a^{-b}}.$$

$$15. \sqrt[3]{(a^3+b^3)}, \sqrt[3]{(a^3+b^3)^2}, \{\sqrt[3]{(a^3+b^3)}\}^2, \sqrt{x}\{(a-b)x\}, \\ \sqrt[2]{(a-bx)^{n-1}}, \sqrt[n]{(a^n-b^n)^{m-3n}}$$

$$16. (a^{\frac{2}{3}})^{\frac{2}{3}}, (b^{-\frac{1}{3}})^{\frac{2}{3}}, (c^{-\frac{2}{3}})^{-\frac{3}{4}}, (x^{\frac{2}{3}})^{-\frac{3}{4}}, (a^2x)^{-\frac{1}{2}}, (a^{-3}x^{-\frac{1}{2}})^{-\frac{1}{2}}, \\ (x^{\frac{1}{2}}y^{-\frac{1}{3}})^{14}.$$

Simplify the following, expressing the results by both notations.

17.  $a \cdot a^{-\frac{1}{2}}, a^0 \cdot a^{-\frac{1}{2}}, a^{\frac{1}{3}} \cdot a^{-\frac{1}{2}}, a \cdot a^{-\frac{1}{4}}, a^{-\frac{1}{2}} \cdot \sqrt{a}, a^{\frac{2}{3}} \sqrt[3]{a^2}, a^{\frac{1}{4}} \sqrt[4]{a}$   
 $a^{\frac{2}{3}} \sqrt[3]{a^3}, a^{\frac{2}{5}} \sqrt[5]{a^{-3}}, a^{\frac{5}{8}} \sqrt[8]{a^{-1}}, a^{\frac{2}{3}} b^{\frac{1}{2}} c^{-\frac{1}{4}}, a^{\frac{1}{3}} b^{-\frac{1}{2}} c^{\frac{1}{4}},$   
 $a^{\frac{1}{3}} b^{\frac{2}{3}} c^{\frac{1}{3}} \cdot a^{-\frac{2}{3}} b^{-\frac{1}{3}} c^{-\frac{1}{3}}$

18.  $\frac{a^{\frac{1}{3}}}{\sqrt{a}}, \frac{a^{\frac{2}{3}}}{\sqrt[3]{a}}, \frac{\sqrt[3]{c^5}}{\sqrt[3]{c}}, \frac{\sqrt[4]{x^3}}{x^{\frac{1}{3}}}, \frac{\sqrt[3]{y^{-2}}}{y^{-\frac{1}{2}}}, \frac{\sqrt[3]{(24a^{-2})}}{2^6 \sqrt[3]{a}}, \frac{c(ab)^{\frac{1}{2}} - ac}{bc - c(ab)}$

19.  $\frac{a^{\frac{1}{2}} + a^{-\frac{1}{2}}}{a^{\frac{1}{2}} - a^{-\frac{1}{2}}}, \frac{a^{\frac{3}{2}} - a^{-\frac{3}{2}}}{a^{\frac{1}{2}} - a^{-\frac{1}{2}}}, \frac{a^{-\frac{3n}{2}} - a^{\frac{6n}{2}}}{a^{-\frac{3n}{2}} + a^{\frac{3n}{2}}}, \frac{a^2 + 1 + a^{-2}}{a + 1 + a^{-1}}$

20. Divide  $x - y$  by  $x^{\frac{1}{n}} - y^{\frac{1}{n}}$ ;  $x^{\frac{1}{3}} + a^{\frac{2}{3}} x^{\frac{2}{3}} + a^{\frac{1}{3}}$  by  $x^{\frac{2}{3}} + a^{\frac{1}{3}} x + a^{\frac{1}{3}}$ ;  
 $x + y + z - 3x^{\frac{1}{3}} y^{\frac{1}{3}} z^{\frac{1}{3}}$  by  $x^{\frac{1}{3}} + y^{\frac{1}{3}} + z^{\frac{1}{3}}$ ;  
 $2ab + 2bc + 2ca - a^2 - b^2 - c^2$  by  $a^{\frac{1}{2}} + b^{\frac{1}{2}} + c^{\frac{1}{2}}$

EXERCISE.

1. Express the following quantities i. as quadratic surds, ii as cubic surds, iii. as quartic surds.

$a, 3a, 2a^2, a^2x, x^n, y^{\frac{1}{2}}, a^{-n}, \frac{x}{y}, mx^{-\frac{n}{p}}, \cdot 1, \cdot 01, 1 \cdot 1x^2.$

2. Reduce to entire surds,

$x \sqrt{x}, a \sqrt[3]{a}, b^2 \sqrt[3]{b^2}, 3 \sqrt[3]{3}, 4 \sqrt[3]{2}, \frac{1}{2} \sqrt{2}, \frac{1}{3} \sqrt[3]{4}, \frac{1}{5} \sqrt[5]{9}, 3 \sqrt[3]{\frac{1}{2}}.$

$a \sqrt{\left(\frac{b}{a}\right)}, \frac{a}{b} \sqrt{b}, \frac{a}{b} \sqrt{\left(\frac{b}{a}\right)}, \frac{a}{b} \sqrt{\left(\frac{a}{b}\right)},$

$\frac{x^3}{y} \sqrt{\left(\frac{y}{x}\right)}, \frac{x^3}{y} \sqrt{\left(\frac{y}{x}\right)^2}, \frac{x^2}{y} \sqrt[3]{(x^2 y^{-1})},$

$a^n/b, a^n \sqrt[3]{(a^m)}, (a+x)^n \sqrt[3]{(a+x)^n}, (a+x)^n \sqrt[3]{(a-x)^{n+1}},$

$\frac{x+y}{m} \sqrt{\left(\frac{x+y}{m}\right)}, (x+y) \sqrt{\left(\frac{x-y}{x+y}\right)}, \frac{a-b}{a+b} \sqrt[3]{\left(\frac{a+b}{a-b}\right)^2},$

$(x-y)^{-2} \sqrt[5]{(x^2 + 2xy + y^2)^{-4}}, (x-x^{-1}) \sqrt[3]{(x^2 + 1)^2}.$

3. Reduce to their simplest form

$$\begin{aligned} & \sqrt{12}, \sqrt[3]{8}, \sqrt{50}, \sqrt[3]{16}, \sqrt[4]{\sqrt[3]{250}}, \sqrt{\frac{1}{2}}, \sqrt[3]{\frac{1}{4}}, \sqrt{\frac{8}{3}}, 5\sqrt[3]{(-320)}, \\ & \sqrt[4]{(1-\frac{8}{11})}, \sqrt{a^3}, \sqrt{(a^3b^7)}, \sqrt[3]{a^5}, 3\frac{1}{2}\sqrt[3]{(54x^9)}, \sqrt[4]{(x^5y^7z^9)}, \\ & \sqrt{\{a^3(1-x^2)\}}, \sqrt[3]{\{a^2(a^2-1)^4\}}, \sqrt{(ab)}, \sqrt[n]{a^{n+1}}, \sqrt[n]{a^{m+n}}, \\ & \sqrt[n]{a^{2n+3}}, \sqrt[n]{a^{3m-2}}, \sqrt{(a^2x+a^3)}, \sqrt[3]{(a^3+2a^4x+a^5x^2)}, \\ & \sqrt{\{(x-1)(x^2-1)\}}, \sqrt[3]{\{(a^2+2ax+x^2)(a^3+x^3)\}}, \\ & \sqrt[3]{\{(x^2-a^2)^2(x-a)\}}, \sqrt{(4x^3-8x^2+4x)}, \\ & \sqrt{(8x^2-16x+8)}, \sqrt[3]{\{(x^2-2+x^{-2})(x^4-2x^2+1)\}}, \\ & \sqrt{\left(\frac{2x-2+2x^{-1}}{x+2+x^{-1}}\right)} \sqrt{\left(\frac{3x^3-6x^2+3x}{27x^2+18x+3}\right)} \sqrt{\left(\frac{(a^2-ab)^2+4a^3b}{a-b}\right)} \end{aligned}$$

4. Compare the following quantities by reducing them to the same surd index :

$$\begin{aligned} & 2 : \sqrt{3} ; 2 : \sqrt[3]{9} ; \sqrt{2} : \sqrt[3]{3} ; \sqrt{10} : \sqrt[3]{30} ; 2\sqrt{2} : \sqrt[3]{22} ; \\ & a^2 : \sqrt{a^3} ; \sqrt[6]{x} : \sqrt[3]{y} ; \sqrt[4]{x} : \sqrt[2]{y} ; \sqrt[3]{x^n} \cdot \sqrt[2]{x^m} ; \sqrt[4]{a} : \sqrt[6]{b} : \sqrt[8]{c} ; \\ & \sqrt[n]{a} : \sqrt[n]{b} : \sqrt[n]{c} ; \sqrt[m]{a^n} : \sqrt[n]{b^p} : \sqrt[p]{c^m}. \end{aligned}$$

5. Reduce to simple surds with lowest integral surd index

$$\begin{aligned} & \sqrt{\sqrt[3]{a}}, \sqrt[3]{\sqrt[4]{b}}, \sqrt[3]{\sqrt{c}}, \sqrt[3]{\sqrt{x^3}}, \sqrt[4]{\sqrt[3]{x^2}}, \sqrt[5]{\sqrt[3]{x^{10}}}, \\ & \sqrt[3]{\sqrt[4]{x^{15}}}, \sqrt[3]{\sqrt{2^7}}, \sqrt{(\sqrt[3]{81})}, \sqrt[4]{(\sqrt[3]{81})}, \sqrt{(a\sqrt{a})}, \\ & \sqrt[3]{(a\sqrt{a})}, \sqrt{(x\sqrt[3]{x})}, \sqrt[3]{(x^2\sqrt[4]{x})}, \sqrt[3]{(5\sqrt{5})}, \sqrt{(3\sqrt[3]{3})}, \\ & \sqrt[4]{(3\sqrt[3]{3})}, \sqrt[6]{(x\sqrt[5]{x})}, \sqrt[n]{\{a\sqrt[n]{b\sqrt[n]{c}}\}}, x\sqrt{(x^{-1}\sqrt{x^{-1}})}, \\ & y\sqrt[3]{(y^{-2}\sqrt[3]{y^{-2}})}, z\sqrt[4]{(z^{-3}\sqrt[3]{z^{-2}})}, \sqrt[m]{\{x\sqrt[n]{y\sqrt[r]{z}}\}}, x^{-1}\sqrt[3]{(x^2\sqrt{x^3})} \end{aligned}$$

6. In the following quantities, combine the terms involving the same radical ;

$$\begin{aligned} & 3\sqrt{2}+5\sqrt{2}-7\sqrt{2} ; \sqrt{8}-\sqrt{2} ; \sqrt[3]{16}+3\sqrt[3]{2} ; \\ & \sqrt[3]{16}+\sqrt{2} ; a\sqrt{x}-\sqrt{x} ; a^3\sqrt{x}-b^3\sqrt{x} ; \\ & 8\sqrt{a}+5\sqrt{x}-7\sqrt{a}+\sqrt{(4a)}-3\sqrt{(4x)}+4\sqrt{(9x)} ; \\ & \sqrt{x}+3\sqrt{(2x)}-2\sqrt{(3x)}+\sqrt{(4x)}-\sqrt{(8x)}+\sqrt{(12x)} ; \end{aligned}$$

$$\begin{aligned}
& 7x - 3\sqrt{x} + 5\sqrt[3]{x} - 2\sqrt{x^2} + \sqrt[5]{x^2}; \\
& 4\sqrt{(a^2x)} + 2\sqrt{(b^2x)} - 3\sqrt{\{(a+b)^2x\}}; \\
& \sqrt{\{(a-b)^2x\}} + \sqrt{\{(a+b)^2x\}} - \sqrt{(a^2x)} + \sqrt{\{(1-a)^2x\}} - \sqrt{x}; \\
& \sqrt{(a-b)} + \sqrt{(16a-16b)} + \sqrt{(ax^2-bx^2)} - \sqrt{\{9(a-b)\}}; \\
& \sqrt{(a^3+a^2b)} - \sqrt{(b^3+ab^2)}; \\
& \sqrt{(a^3+2a^2b+ab^2)} - \sqrt{(a^3-2a^2b+ab^2)} - \sqrt{(4ab^2)}.
\end{aligned}$$

7. In the following quantities, perform, as far as possible, the indicated multiplications and divisions, expressing the results in their simplest forms:

$$\begin{aligned}
& \sqrt{2}\sqrt{6}; \sqrt{3}\sqrt{12}; \sqrt{14}\sqrt{35}\sqrt{10}; \sqrt{a}\sqrt{(3a)}; \\
& \sqrt{c}\sqrt{(12c)}; \sqrt{(6x)}\sqrt{(8x)}; \sqrt{y^3}\sqrt{y^3}; \sqrt[3]{y^5}\sqrt[3]{y^7}; \\
& \sqrt[3]{a}\sqrt[3]{a^2}\sqrt{b}; \sqrt{a}\sqrt{\left(\frac{x}{a}\right)}; \sqrt{a}\sqrt{\left(\frac{a}{x}\right)}; \sqrt{3a}\sqrt{\left(\frac{5c}{6a}\right)}; \\
& \sqrt{a^{n+1}}\sqrt{a^{n+1}}; \sqrt[3]{b^{n+1}}\sqrt[3]{b^{2n+1}}; \sqrt{12} \div \sqrt{3}; \sqrt{(6x)} \div \sqrt{(2x)}; \\
& a \div \sqrt[3]{a}; a^2 \div \sqrt[4]{a^3}; a \div \sqrt[2]{a^{n-1}}; a^p \div \sqrt[2]{a^{n-m}}; \\
& (a+x) \div \sqrt{(a+x)}; (a^2-x^2) \div \sqrt{(a-x)}; (x^2-1) \div \sqrt[3]{(x+1)^2}; \\
& (3\sqrt{8}-5\sqrt{2}+\sqrt{18}+\sqrt{32}+\sqrt{72}-2\sqrt{50})\sqrt{2}; \\
& (7\sqrt{2}-5\sqrt{6}-3\sqrt{8}+4\sqrt{20})\sqrt{18}; (\sqrt{5}+\sqrt{3})(\sqrt{5}-\sqrt{3}); \\
& (\sqrt{2}+1)(\sqrt{6}-\sqrt{3}); (3-\sqrt{2})(2+3\sqrt{2}); \\
& (5\sqrt{3}+\sqrt{6})(5\sqrt{2}-2); (\sqrt{a}-\sqrt{b})(\sqrt{a}+\sqrt{b}); \\
& (a\sqrt{b}+b\sqrt{a})(b\sqrt{a}-a\sqrt{b}); \\
& \{\sqrt{(x+1)}+\sqrt{(x-1)}\}\{\sqrt{(x+1)}-\sqrt{(x-1)}\}; \\
& \{\sqrt{(3a-b)}+\sqrt{(3b-a)}\}\{\sqrt{(3a-b)}-\sqrt{(3b-a)}\}; \\
& \sqrt{(a+\sqrt{b})}\sqrt{(a-\sqrt{b})}; \sqrt{(\sqrt{x}+\sqrt{y})}\sqrt{(\sqrt{x}-\sqrt{y})}; \\
& \sqrt{a+\sqrt{(a^2-x^2)}}\sqrt{a-\sqrt{(a^2-x^2)}}; \\
& \sqrt[3]{x-\sqrt{(x^2-1)}}\sqrt[3]{x+\sqrt{(x^2-1)}}; \\
& \sqrt[3]{a\sqrt{a}-\sqrt{(a^3-x^3)}}\sqrt[3]{\sqrt{(a^3-x^3)}+a\sqrt{a}};
\end{aligned}$$

$$\begin{aligned}
& \sqrt[3]{8+3\sqrt{7}}, \sqrt[3]{8-3\sqrt{7}}; (\sqrt{a+\sqrt{b}})^2; (\sqrt{a+\sqrt{b}})^3; \\
& (a-\sqrt{cx})^2; (\sqrt{x+\sqrt{x-1}})^2; \\
& \{a+\sqrt{(1-a^2)}\}^2; \{\sqrt{(a+b-x)}-\sqrt{(a-b+x)}\}^2; \\
& [\{\sqrt{(a+x)(x-b)}\}+\sqrt{(a-x)(x+b)}]^2; \left\{\sqrt{\frac{2a}{b}}-\sqrt{\frac{3b}{2a}}\right\}^2 \\
& [\sqrt{(a+x)(x+b)}+\sqrt{(a-x)(x-b)}]^2; \\
& \left\{\sqrt{\frac{a-x}{x-b}}-\sqrt{\frac{x-b}{a-x}}\right\}^2; \sqrt{(x^2-1)}\cdot\sqrt{\frac{x+1}{x-1}}; \\
& (\sqrt[3]{a}+\sqrt[3]{b})^3; \{\sqrt{(1/a+\sqrt{b})}+\sqrt{(a-\sqrt{b})}\}^2; \\
& \{\sqrt{(\sqrt{10+1})}-\sqrt{(\sqrt{10-1})}\}^2; \\
& [\sqrt{a+\sqrt{(a^2-x^2)}}+\sqrt{a-\sqrt{(a^2-x^2)}}]^2; \\
& (\sqrt{x+\sqrt{y}})^4+(\sqrt{x-\sqrt{y}})^4; (a^2+ab\sqrt{2+b^2})(a^2-ab\sqrt{2+b^2}); \\
& (\sqrt[3]{a}+\sqrt[3]{c})(\sqrt[3]{a^2}-\sqrt[3]{ac}+\sqrt[3]{c^2}); \\
& (\sqrt{a+\sqrt{b}}+\sqrt{c})(\sqrt{b+\sqrt{c}}-\sqrt{a})(\sqrt{c+\sqrt{a}}-\sqrt{b})(\sqrt{a+\sqrt{b}}-\sqrt{c}); \\
& \left(\sqrt{\frac{x}{a}}+\sqrt{\frac{y}{b}}+1\right)\left(\sqrt{\frac{x}{a}}+\sqrt{\frac{y}{b}}-1\right) \times \\
& \left(\frac{x}{a}-\sqrt{\frac{y}{b}}+1\right)\left(-\sqrt{\frac{x}{a}}+\sqrt{\frac{y}{b}}+1\right) \\
& (\sqrt[3]{a}+\sqrt[3]{b}+\sqrt[3]{c})\{\sqrt[3]{a^2}+\sqrt[3]{b^2}+\sqrt[3]{c^2}-2\sqrt[3]{bc}-2\sqrt[3]{ca}-2\sqrt[3]{ab}\}.
\end{aligned}$$

8. Find rationalizing multipliers for the following expressions, and also the products of multiplication by these :

$$\begin{aligned}
& a+\sqrt{b}, \sqrt{a+b\sqrt{c}}, a\sqrt{b}-b\sqrt{a}, a+\sqrt{(a^2-x^2)}, \\
& \sqrt{(a-x)}-\sqrt{(a+x)}, \sqrt{(a^2+\sqrt{c})}+\sqrt{(a^2-\sqrt{c})}, \\
& \sqrt{8+\sqrt{(24+\sqrt{5})}}-\sqrt{8+\sqrt{(24-\sqrt{5})}}, \sqrt{a}+\sqrt{b}+\sqrt{c}, \\
& 3+\sqrt{2}+\sqrt{7}, \sqrt{6}+\sqrt{5}-\sqrt{3}-\sqrt{2}, \sqrt{a}+\sqrt{b}+\sqrt{c}+\sqrt{d}, \\
& \sqrt{(1+a)}-\sqrt{(1-a)}+\sqrt{(1+b)}-\sqrt{(1-c)}, \sqrt[3]{a}+\sqrt[3]{c}, \\
& \sqrt[3]{a^2}-\sqrt[3]{c^2}, \sqrt[4]{a}+\sqrt[4]{c}, \sqrt{a}-\sqrt[3]{b}, \sqrt[4]{a}+\sqrt{a}, \sqrt{x}+\sqrt[4]{y^3}, \\
& \sqrt{x+1}+\sqrt{x^{-1}}, \sqrt{(ab^{-1})}-\sqrt{(a^{-1}b)}, \sqrt[3]{2}+\sqrt[3]{3}-\sqrt[3]{5}, \\
& \sqrt[3]{a}+\sqrt[3]{b}+\sqrt[3]{c}, a+\sqrt{b}+\sqrt[3]{c}.
\end{aligned}$$

9. Rationalize the divisors and the denominators in the following, and reduce the results to their simplest form:

$$\begin{aligned}
 & 1 \div (2 - \sqrt{3}), \quad 3 \div (3 + \sqrt{6}), \quad 5 \div (\sqrt{2} + \sqrt{7}), \\
 & (\sqrt{3} + \sqrt{2}) \div (\sqrt{3} - \sqrt{2}), \quad (7\sqrt{5} + 5\sqrt{7}) \div (\sqrt{5} + \sqrt{7}), \\
 & a \div (\sqrt{a} + a), \quad (x - a) \div (\sqrt{x} - \sqrt{a}), \\
 & (a^2 + ab + b^2) \div \{a + \sqrt{(ab) + b}\}, \quad (x + a) \div (\sqrt[3]{x} + \sqrt[3]{a}), \\
 & \frac{a\sqrt{x} + b\sqrt{y}}{c\sqrt{x} - e\sqrt{y}}, \quad \frac{2\sqrt{6}}{\sqrt{2} + \sqrt{3} - \sqrt{5}}, \quad \frac{1 + 3\sqrt{2} - 2\sqrt{3}}{\sqrt{2} + \sqrt{3} + \sqrt{6}}, \\
 & \frac{\sqrt{6} - \sqrt{5} - \sqrt{3} + \sqrt{2}}{\sqrt{6} + \sqrt{5} - \sqrt{3} - \sqrt{2}}, \quad \frac{2}{\sqrt{(a+1)} - \sqrt{(a-1)}}, \\
 & \frac{2c}{\sqrt{(a+c)} + \sqrt{(a-c)}}, \quad \frac{a+x + \sqrt{(a^2+x^2)}}{a+x - \sqrt{(a^2+x^2)}}, \\
 & \frac{\sqrt{(a+x)} + \sqrt{(a-x)}}{\sqrt{(a+x)} - \sqrt{(a-x)}}, \quad \frac{1}{a\sqrt{(1+b^2)} + b\sqrt{(1+a^2)}}, \\
 & \frac{a\sqrt{(1-b^2)} - b\sqrt{(1-a^2)}}{\sqrt{(1-b^2)} + \sqrt{(1-a^2)}}, \quad \frac{a\sqrt{(1-a^2)} + c\sqrt{(1-c^2)}}{a\sqrt{(1-c^2)} + c\sqrt{(1-a^2)}}, \\
 & \frac{\sqrt{\{(1+a)(1+b)\}} - \sqrt{\{(1-a)(1-b)\}}}{\sqrt{\{(1+a)(1+b)\}} + \sqrt{\{(1-a)(1-b)\}}}, \\
 & \frac{(a-x)\sqrt{(b^2+y^2)} - (b-y)\sqrt{(a^2+x^2)}}{(a+x)\sqrt{(b^2+y^2)} + (b+y)\sqrt{(a^2+x^2)}}, \\
 & \frac{\sqrt{(1+a)} - \sqrt{(1-a)} + \sqrt{(1+b)} - \sqrt{(1-b)}}{\sqrt{(1+a)} + \sqrt{(1-a)} + \sqrt{(1+b)} + \sqrt{(1-b)}}, \\
 & \frac{\sqrt{(x+a)} - \sqrt{(x-a)} - \sqrt{(x+b)} + \sqrt{(x-b)}}{\sqrt{(x+a)} + \sqrt{(x-a)} + \sqrt{(x+b)} + \sqrt{(x-b)}}, \\
 & \frac{\sqrt{a}}{\sqrt{b}} + \frac{\sqrt{b}}{\sqrt{a}}, \quad \sqrt{\frac{(a+x)}{(a-x)}} - \sqrt{\frac{(a-x)}{(a+x)}}, \quad \sqrt{\frac{a+\sqrt{x}}{a-\sqrt{x}}}, \quad \sqrt{\frac{\sqrt{x}-\sqrt{y}}{\sqrt{x}+\sqrt{y}}}, \\
 & \frac{\sqrt{a+\sqrt{(a^2-1)}}}{\sqrt{a-\sqrt{(a^2-1)}}}, \quad \frac{1}{\sqrt{x}} - \frac{1}{\sqrt{y}}, \quad \frac{\sqrt{a}}{\sqrt{x}} - \frac{\sqrt{x}}{\sqrt{a}}, \\
 & \frac{1}{\sqrt{x} + \sqrt{y}}, \quad \frac{1}{\sqrt{x}} + \frac{1}{\sqrt{y}}, \quad \frac{\sqrt{a}}{\sqrt{x}} + \frac{\sqrt{x}}{\sqrt{a}}.
 \end{aligned}$$

10. Find the values of the following expressions for  $n = 1, 2, 3, 4, 5$ , respectively.

$$\frac{1}{\sqrt{5}} \left\{ \left( \frac{1+\sqrt{5}}{2} \right)^n - \left( \frac{1-\sqrt{5}}{2} \right)^n \right\}.$$

$$\frac{1}{2\sqrt{6}} \left\{ \frac{(2+\sqrt{6})^{n+1} - (2+\sqrt{6})}{1+\sqrt{6}} - \frac{(2-\sqrt{6})^{n+1} - (2-\sqrt{6})}{1-\sqrt{6}} \right\}$$

11. Show that

$$\frac{1}{2(x-1)} \left[ \{x + \sqrt{(x^2-1)}\}^{4n+1} + \{x - \sqrt{(x^2-1)}\}^{4n+1} \mp 2 \right]$$

is a square for  $n = 1, 2$ , or  $3$  respectively.

12. Extract the square roots of

$$x+y-2\sqrt{(xy)}, \quad a+c+e+2\sqrt{(ac+ce)},$$

$$a+2c+e+2\sqrt{\{(a+c)(c+e)\}}, \quad 2a+2\sqrt{(a^2-c^2)},$$

$$2\{a^2+b^2-\sqrt{(a^4+a^2b^2+b^4)}\}, \quad x-2+x^{-1},$$

$$\sqrt{x+2}+\sqrt{x^{-1}}, \quad x+3x^2+x^3+2x\sqrt{x}+2x^2\sqrt{x},$$

$$x^2-xy+\frac{1}{4}y^2+\sqrt{(4x^3y-8x^2y^2+xy^3)}, \quad 2x+\sqrt{(9x^2-y^2)},$$

$$5-2\sqrt{6}, \quad 10+2\sqrt{21}, \quad 9+4\sqrt{5}, \quad 4-\sqrt{15}, \quad 7+4\sqrt{3},$$

$$12-5\sqrt{6}, \quad 70+3\sqrt{451}, \quad 4-\sqrt{15},$$

$$9+2\sqrt{6}+4(\sqrt{2}+\sqrt{3}), \quad 15.25-5\sqrt{6}.$$

13. Find the value of

$$\frac{(a+b)xy}{ay^2+bx^2}, \text{ given } x = \frac{a\sqrt{a}}{\sqrt{(a+b)}} \text{ and } y = \frac{b\sqrt{b}}{\sqrt{(a+b)}};$$

$$\sqrt{(x^2+y^2)}, \text{ given } x = \sqrt[3]{(a^2c)} \quad y = \sqrt[3]{(a^2e)};$$

$$\frac{x+\sqrt{(x^2+1)}}{x-\sqrt{(x^2+1)}}, \text{ given } x = \frac{1}{2} \left\{ \sqrt{\frac{a}{c}} - \sqrt{\frac{c}{a}} \right\}$$

$$\frac{\sqrt{(1+x)}-\sqrt{(1-x)}}{\sqrt{(1+x)}+\sqrt{(1-x)}}, \text{ given } x = \frac{2ab}{a^2+b^2};$$

$$\frac{2a\sqrt{(1+x^2)}}{x+\sqrt{(1+x^2)}}, \text{ given } x = \frac{1}{2} \left\{ \sqrt{\frac{a}{e}} - \sqrt{\frac{e}{a}} \right\}.$$

14. If  $\sqrt{x+a+b} + \sqrt{x+c+d} = \sqrt{x+a-c} + \sqrt{x-b+d}$ ,  
 $\therefore b+c=0$ .

15. Simplify  $\frac{\frac{1}{2}(1+\sqrt{5})x-2}{x^2-\frac{1}{2}(1+\sqrt{5})x+1} + \frac{\frac{1}{2}(1-\sqrt{5})x-2}{x^2-\frac{1}{2}(1-\sqrt{5})x+1}$ .

COMPLEX QUANTITIES.

Quantities of the form  $a+b\sqrt{-1}$  in which neither  $a$  nor  $b$  involves  $\sqrt{-1}$ , are called complex quantities. The letter  $j$  (or  $i$ ) is frequently used as the symbol of the ditensive unit  $\sqrt{-1}$ , so that  $a+b\sqrt{-1}$  would be written  $a+bj$ . So also  $\sqrt{-x}=j\sqrt{x}$ ,  $\sqrt{-x}\sqrt{-y}=j^2\sqrt{xy}=-\sqrt{xy}$ , and  $j^3=-j$ .

EXERCISE.

Simplify the following, writing  $j$  for  $\sqrt{-1}$  in any result in which the latter occurs:

1.  $\sqrt{-4}$ ,  $\sqrt{-36}$ ,  $\sqrt{-81}$ ,  $\sqrt{-8}$ ,  $\sqrt{-12}$ ,  $\sqrt{-72}$ ,  $\sqrt[3]{-8}$ ,  
 $\sqrt{-5}\sqrt{-6}$ ,  $\sqrt{-6}\sqrt{-8}$ ,  $\sqrt{-8}\sqrt{12}$ ,  $\sqrt{-8}\sqrt[3]{-8}$ ,  
 $\sqrt{-5}\sqrt{-20}$ .

2.  $\sqrt{-x}$ ,  $\sqrt{-x^2}$ ,  $\sqrt{-a^3}$ ,  $\sqrt{-a^{2n}}$ ,  $\sqrt{(-a)^2}$ ,  $\sqrt{(-a)^3}$ ,  
 $\sqrt{(-3ax^3)}$ ,  $\sqrt{-a}\sqrt{a^3}$ ,  $\sqrt{-x^2}\sqrt{-y^3}$ ,  $\sqrt{-a}\sqrt{-1}$ ,  
 $\sqrt{5}\sqrt{-a}$ .

3.  $j^2$ ,  $j^3$ ,  $j^4$ ,  $j^5$ ,  $j^9$ ,  $j^{15}$ ,  $j^{16}$ ,  $j^{17}$ ,  $j^{18}$ ,  $j^{4n}$ ,  $j^{4n+1}$ ,  $j^{4n+2}$ ,  $j^{4n+3}$ ,

4.  $aj$ ,  $bj$ ,  $j\sqrt{x}j\sqrt{y}$ ,  $5j$ ,  $j^2\sqrt{5}$ ,  $j\sqrt{-a}$ ,  $j\sqrt{-a^2}$ ,  $j\sqrt{a}\sqrt{-a}$ .

5.  $\sqrt{-j^2}$ ,  $\sqrt{-j^3}$ ,  $\sqrt{-j^4}$ ,  $\sqrt{-j^5}$ ,  $\sqrt{-j^{2n}}$ ,  $\sqrt{-j^{4n}}$ .

6.  $\frac{\sqrt{-6}}{\sqrt{3}}$ ,  $\frac{\sqrt{-6}}{\sqrt{-3}}$ ,  $\frac{\sqrt{6}}{\sqrt{-3}}$ ,  $\frac{\sqrt{a}}{\sqrt{-b}}$ ,  $\frac{\sqrt{a}}{\sqrt{-b}}$ ,  $\frac{1}{\sqrt{-1}}$ ,  
 $\frac{a}{\sqrt{-a}}$ ,  $\frac{a^2}{\sqrt{-a^2}}$ ,  $\frac{\sqrt{(-ax)}}{\sqrt{-x}}$ ,  $\frac{-\sqrt{-1}}{\sqrt{-a}}$ ,  $\frac{a^3}{\sqrt[3]{-a^3}}$ ,  
 $\frac{-c}{\sqrt{-c^3}}$ ,  $\frac{\sqrt{(-a)^{2n+1}}}{\sqrt{(-a)^{2n-1}}}$ .



$$7. \frac{1}{j}, \frac{1}{j^2}, \frac{1}{j^3}, \frac{-1}{j}, \frac{1}{j^6}, \frac{1}{j^{4n+1}}, \frac{-1}{j^{4n+1}}, \frac{1}{j^{4n-1}},$$

$$\frac{a^2j}{\sqrt{-a^3}}, \frac{x}{j\sqrt{x}}, \frac{-y}{j\sqrt{-y^2}}, -\frac{cj^3}{\sqrt{-c^2}}.$$

$$8. \sqrt{(a-b)} \cdot \sqrt{(b-a)}, \sqrt{(3x-4y)} \cdot \sqrt{(4y-3x)}, (3+5j)(7+4j),$$

$$(8-9j)(8-7j), (7-j\sqrt{5})(7+j\sqrt{10}), (\sqrt{3}-j\sqrt{6})(\sqrt{2}-j\sqrt{6}),$$

$$(a+bj)(c+ej), \{a+(a-1j)\} \{a+(a+1j)\},$$

$$(\sqrt{a+j\sqrt{b}})(\sqrt{a-j\sqrt{c}}) (a+bj)(a-bj), (aj+b)(aj-b),$$

$$(\sqrt{a+j\sqrt{b}})(\sqrt{a-j\sqrt{b}}), (a\sqrt{b+cj\sqrt{x}})(a\sqrt{b-cj\sqrt{x}}),$$

$$\sqrt{(1+j)} \cdot \sqrt{(1-j)}, \sqrt{(3+4j)} \cdot \sqrt{(3-4j)},$$

$$\sqrt{(12+5j)} \cdot \sqrt{(12-5j)}, (1+j)^2, (\sqrt{a-j\sqrt{b}})^2, (5-2j\sqrt{6})^2,$$

$$(a+bj)^2+(a-bj)^2, (a+bj)^2-(a-bj)^2, (a+bj)^2+(aj-b)^2,$$

$$\{\sqrt{(4+3j)}+\sqrt{(4-3j)}\}^2, \{\sqrt{(3-4j)}-\sqrt{(3+4j)}\}^2,$$

$$\{\sqrt{(1+j)}+\sqrt{(1-j)}\}^2, (1+j)^3, (1+j)^4,$$

$$(a+bj)^4, (a+bj)^3+(a-bj)^3, (a+bj)^3-(a-bj)^3,$$

$$\left(\frac{1+j\sqrt{3}}{2}\right)^3, \left(\frac{-1+j\sqrt{3}}{2}\right)^3, \left(\frac{-1-j\sqrt{3}}{2}\right)^3, \left(\frac{1+j}{\sqrt{2}}\right)^4, \left(\frac{1-j}{\sqrt{2}}\right)^4,$$

$$(x+jy)^4+(x-jy)^4, (x+jy)^4-(x-jy)^4,$$

$$(1+j\sqrt{5})^4+(1-j\sqrt{5})^4, (a+bj)^5+(a-bj)^5,$$

$$(a+bj)^5-(a-bj)^5, (1+j)^5+(1-j)^5, (1+j\sqrt{2})^5+(1-j\sqrt{2})^5$$

$$\left(\frac{j+\sqrt{3}}{2}\right)^6, \left(\frac{j-\sqrt{3}}{2}\right)^6,$$

$\left[\frac{1}{8}\{\sqrt{(30-6\sqrt{5})}-1-\sqrt{5}\}+\frac{1}{8}j\{\sqrt{15}+\sqrt{3}+\sqrt{(10-2\sqrt{5})}\}\right]^n$   
for all positive integral values of  $n$ .

$$9. \frac{4}{1+j\sqrt{3}}, \frac{64}{1-j\sqrt{7}}, \frac{21}{4+3j\sqrt{6}}, \frac{5}{\sqrt{2}+j\sqrt{3}}, \frac{1-20j\sqrt{5}}{7-2j\sqrt{5}},$$

$$\frac{1-j\sqrt{3}}{1+j\sqrt{3}}, \frac{1+j}{1-j}, \frac{1+j^3}{1+j}, \frac{1-j^3}{1-j}, \frac{1-j^3}{(1+j)^3}, \frac{1+j^3}{1-j}, \frac{x+yj}{x-yj},$$

$$\frac{a+j\sqrt{x}}{a-j\sqrt{x}}, \frac{j\sqrt{a+\sqrt{-b}}}{\sqrt{-a-j\sqrt{b}}}, \frac{a-bj}{aj+b}, \frac{a+j\sqrt{(1-x^2)}}{a-j\sqrt{(1-x^2)}}$$

$$\frac{\sqrt{(x-y)} - \sqrt{(y-x)}}{\sqrt{(y-x)} + \sqrt{(x-y)}} \cdot \frac{1}{1+j} + \frac{1}{1-j} \cdot \frac{1+j}{1-j} + \frac{1-j}{1+j},$$

$$\frac{1}{(1+j)^2} + \frac{1}{(1-j)^2}, \quad \frac{1}{(1+j)^4} - \frac{1}{(1-j)^4}, \quad \frac{x+yj}{a+bj} + \frac{x-yj}{a-bj},$$

$$\frac{x+yj}{a+bj} - \frac{x-yj}{a-bj}, \quad \frac{\sqrt{x+j\sqrt{y}}}{\sqrt{x-j\sqrt{y}}} - \frac{\sqrt{y+j\sqrt{x}}}{\sqrt{y-j\sqrt{x}}},$$

$$\frac{\sqrt{(1+a)} + j\sqrt{(1-a)}}{\sqrt{(1+a)} - j\sqrt{(1-a)}} - \frac{\sqrt{(1-a)} + j\sqrt{(1+a)}}{\sqrt{(1-a)} - j\sqrt{(1+a)}}.$$

10.  $\sqrt{(3+4j)} + \sqrt{(3-4j)}, \quad \sqrt{(3+4j)} - \sqrt{(3-4j)},$   
 $\sqrt{(4+3j)} \pm \sqrt{(4-3j)}, \quad \sqrt{(1+2j\sqrt{6})} \pm \sqrt{(1-2j\sqrt{6})}$   
 $\sqrt{(5+2j\sqrt{6})} \pm \sqrt{(5-2j\sqrt{6})},$   
 $\sqrt{(2\sqrt{15}+30j)} \pm \sqrt{(2\sqrt{15}-30j)},$   
 $\sqrt{(\sqrt{3}+j\sqrt{105})} \pm \sqrt{(\sqrt{3}-j\sqrt{105})},$   
 $\sqrt{\{a+j\sqrt{(x^2-a^2)}\}} \pm \sqrt{\{a-j\sqrt{(x^2-a^2)}\}},$   
 $\sqrt{\{a^2+jx\sqrt{(x^2+2a^2)}\}} \pm \sqrt{\{a^2-jx\sqrt{(x^2+2a^2)}\}}.$

11. Prove that both  $\frac{1}{2}(-1+j\sqrt{3})$  and  $\frac{1}{2}(-1-j\sqrt{3})$  satisfy the equation  $\frac{x^3-1}{x-1} = 0$ ,

that  $(x+wy+w^2z)^3 = x^3+y^3+z^3+3(x+wy)(y+wz)(z+wx)$   
 and that  $(x+y+z)(x+wy+w^2z)(x+w^2y+wz) = x^3+y^3+z^3-3xyz$ , in which  $w$  represents either of the preceding complex quantities.

Hence, prove that

$$(i) \{2a-b-c+(b-c)j\sqrt{3}\}^3 = \{2b-c-a+(c-a)j\sqrt{3}\}^3 = \{2c-a-b+(a-b)j\sqrt{3}\}^3;$$

$$(ii) u^3+v^3+w^3-3uvw = (a^3+b^3+c^3-3abc)x$$

$(x^3+y^3+z^3-3xyz)$ , if  $u = ax+by+cz, \quad v = ay+bz+cx,$   
 $w = az+bx+cy$ , or if  $u = ax+cy+bz,$   
 $v = cx+by+az, \quad w = bx+ay+cz.$

12. Prove that  $\frac{1}{2} \{ \sqrt{5} + 1 + j \sqrt{(10 - 2\sqrt{5})} \}$  satisfies the equation

$$\frac{x^5 + 1}{x + 1} = 0.$$

Writing  $w$  for the preceding complex quantity, prove that

$$(7 + w + w^2 + 3w^3)(7 - w^4 - w^3 - 3w^2) = 71,$$

$$\text{and } (x + y + z)(x + w^2y - w^3z)(x - w^3y - wz)(x - wz + w^4z)$$

$$(x + w^4y + w^2z) = x^5 + y^5 + z^5 - 5x^3yz + 5xy^2z^2.$$

Prove that  $\{4a + (b - c)(\sqrt{5} - 1) + (b + c)j\sqrt{(10 + 2\sqrt{5})}\}^5 =$   
 $\{[(a + b)\{-1 + j\sqrt{(\sqrt{5} + 2)}\} + (a - b)\{\sqrt{5} + j\sqrt{(\sqrt{5} - 2)}\}]\}$   
 $\times \sqrt[4]{5 - 4c}\}^5.$

### SECTION III.—PURE QUADRATICS.

#### EXAMPLES.

$$1. \quad \frac{x + 3(a - b)}{x - 3(a - b)} = \frac{a(3x + 9a - 7b)}{b(3x - 7a + 9b)}.$$

Apply, if  $\frac{m}{n} = \frac{p}{q}$ ,  $\therefore \frac{m + n}{m - n} = \frac{p + q}{p - q}$ ;

$$\therefore \frac{x}{3(a - b)} = \frac{3x(a + b) + 9a^2 - 14ab + 9b^2}{3x(a - b) + 9(a^2 - b^2)}.$$

Dividing the denominators by  $3(a - b)$

$$\therefore x\{1 + 3(a + b)\} = 3x(a - b) + 9a^2 - 14ab + 9b^2,$$

$$\therefore x^2 = 9a^2 - 14ab + 9b^2$$

$$2. \quad \left(\frac{x - 2a + 4b}{x + 4a - 2b}\right)^2 = \frac{5x - 9a + 3b}{5x + 3a - 9b}.$$

Apply, if  $\frac{m}{n} = \frac{p}{q}$ ,  $\therefore \frac{n - m}{n} = \frac{q - p}{p}$ , and factor the numerator

$$(x + 4a - 2b)^2 - (x - 2a + 4b)^2,$$

$$\therefore \frac{12(x+a+b)(a-b)}{(x+4a-2b)^2} = \frac{12(a-b)}{5x+3a-9b},$$

$$\therefore \frac{x+a+b}{x+4a-2b} = \frac{x+4a-2b}{5x+3a-9b} = \frac{3(a-b)}{4x-a-7b}, \text{ by taking differ-}$$

ence of numerators and difference of denominators. To the first and third of these fractions, apply if

$$\frac{m}{n} = \frac{p}{q}, \therefore \frac{m}{n-m} = \frac{p}{q-p},$$

$$\therefore \frac{x+a+b}{3(a-b)} = \frac{3(a-b)}{4x-4a-4b},$$

$$\therefore 4\{x^2 - (a+b)^2\} = 9(a-b)^2,$$

$$\therefore x^2 = \frac{1}{4}\{4(a+b)^2 + 9(a-b)^2\}.$$

$$3. \frac{\sqrt{3x^2-1} + \sqrt{3-x^2}}{\sqrt{3x^2-1} - \sqrt{3-x^2}} = \frac{a}{b},$$

$$\therefore \sqrt{\frac{3x^2-1}{3-x^2}} = \frac{a+b}{a-b},$$

$$\therefore \frac{3x^2-1}{3-x^2} = \frac{(a+b)^2}{(a-b)^2},$$

$$\therefore x^2 = \frac{3(a+b)^2 + (a-b)^2}{(a+b)^2 + 3(a-b)^2} = \frac{7^2 + ab + b^2}{a^2 - ab + b^2}.$$

$$4. m\sqrt{1+x} - n\sqrt{1-x} = \sqrt{m^2+n^2} \quad (1)$$

Square both members and reduce

$$\therefore (m^2 - n^2)x - 2mn\sqrt{1-x^2} = 0. \quad (2)$$

Transfer the radical term and square both members,

$$\therefore (m^2 - n^2)^2 x^2 = 4m^2 n^2 (1 - x^2) \quad (3)$$

$$\therefore (m^2 + n^2)^2 x^2 = 4m^2 n^2 \quad (4)$$

$$\therefore x = \frac{\pm 2mn}{m^2 + n^2}. \quad (4)$$

The above follows the usual mode of solving equations involving radicals, viz., make a radical term the right-hand member gathering all the other terms into the left-hand member, square each

member, repeat, if necessary, until all radicals are rationalized. This method is convenient but it does not explain the difficulty

that only one of the values of  $x$  in (4) satisfies (1) viz.  $\frac{+2mn}{m^2+n^2}$

The other value,  $\frac{-2mn}{m^2+n^2}$  satisfies the equation

$$m\sqrt{1+x} + n\sqrt{1-x} = \sqrt{m^2+n^2}.$$

The explanation is simple. Squaring both members of (1) is really equivalent to substituting for (1) the conjoint equation

$$\begin{aligned} &\{m\sqrt{1+x} - n\sqrt{1-x} - \sqrt{m^2+n^2}\} \\ &\{m\sqrt{1+x} + n\sqrt{1-x} - \sqrt{m^2+n^2}\} = 0 \end{aligned} \quad (5)$$

which reduces to (2) above.

Treating (5) or (2) by transferring and squaring is equivalent to substituting for it, the equation

$$\begin{aligned} &\{m\sqrt{1+x} - n\sqrt{1-x} - \sqrt{m^2+n^2}\} \times \\ &\{m\sqrt{1+x} - n\sqrt{1-x} + \sqrt{m^2+n^2}\} \times \\ &\{m\sqrt{1+x} + n\sqrt{1-x} - \sqrt{m^2+n^2}\} \times \\ &\{m\sqrt{1+x} + n\sqrt{1-x} + \sqrt{m^2+n^2}\} = 0 \end{aligned} \quad (6)$$

which reduces to

$$\{(m^2-n^2)x - 2mn\sqrt{1-x^2}\} \{m^2-n^2\}x + 2mn\sqrt{1-x^2}\} = 0 \quad (7)$$

which further reduces to (3)

Thus the whole process of solving (1) is equivalent to reducing it to an equation of the type  $A=0$  and then multiplying the member  $A$  by rationalizing factors. Thus instead of solving (1) we really solve (6), *i.e.*, a conjoint equation equivalent to *four* disjunctive equations. (See page 140, Art xl) Now the values given in (4) will satisfy (6), the positive value making the first factor vanish, the negative value making the third factor vanish, while no values can be found that will make either the second or the fourth factor vanish.

Hence, if one of such a set of disjunctive equations is proposed for solution, the conjoint equation must be solved, and if there be a value of  $x$  which satisfies the particular equation proposed, that value must be retained and the others rejected.

(This process is the opposite to that given in Arts. XL. and XLV. : there a conjoint equation is solved by resolving it into its equivalent disjunctive equations. The two processes are related somewhat as involution and evolution are).

Further, it should be noticed that just as there are four factors in (6) while there are only two values in (4), it will in general be possible to form more disjunctive equations than there are values of  $x$  that satisfy the conjoint equation, and consequently it will be possible to select disjunctive equations that are not satisfied by any value of  $x$ , or, in other words, whose solution is impossible.

This will perhaps be better understood by considering the following problem.

Find a number such that if it be increased by 4 and also diminished by 4 the difference of the square-roots of the results shall be 4.

Reduced to an equation this is

$$(6) \quad \sqrt{x+4} - \sqrt{x-4} = 4 \quad (8)$$

Rationalizing this becomes

$$\begin{aligned} & \{4 - \sqrt{x+4} + \sqrt{x-4}\} \{4 - \sqrt{x+4} - \sqrt{x-4}\} \times \\ & \{4 + \sqrt{x+4} + \sqrt{x-4}\} \{4 + \sqrt{x+4} - \sqrt{x-4}\} = 0 \quad (9) \end{aligned}$$

which reduces to

$$\{24 - 8\sqrt{x+4}\} \{24 + 8\sqrt{x+4}\} = 0$$

$$i.e. 9 - (x+4) = 0, \text{ or } x = 5.$$

Now  $x = 5$  satisfies (9) because it makes the factor

$$4 - \sqrt{x+4} - \sqrt{x-4}$$

vanish and it is the only finite value of  $x$  that does satisfy (9), or, in other words, there are no values of  $x$  which will make any of the factors

$$4 - \sqrt{(x+4)} + \sqrt{(x-4)}, \quad 4 + \sqrt{(x+4)} + \sqrt{(x-4)},$$

$$\text{or } 4 + \sqrt{(x+4)} - \sqrt{(x-4)}$$

vanish. There is, therefore, no number that will satisfy the conditions of the problem.

[It will be found that as  $x$  increases,  $\sqrt{(x+4)} - \sqrt{(x-4)}$  decreases, hence as 4 is the least value that can be given to  $x$  without involving the square-root of a negative, the greatest real value of  $\sqrt{(x+4)} - \sqrt{(x-4)}$  is  $\sqrt{8}$  which is less than 4. We see by this that our method of solution fails for (8) simply because (8) is *impossible*].

$$5. \quad \sqrt{\{(a+x)(b+x)\}} - \sqrt{\{(a-x)(b-x)\}} =$$

$$\sqrt{\{(a-x)(b+x)\}} - \sqrt{\{(a+x)(b-x)\}} \quad (1)$$

Collecting the terms involving  $\sqrt{(a+x)}$  and  $\sqrt{(a-x)}$  respectively the equation becomes

$$\{\sqrt{(a+x)} - \sqrt{(a-x)}\} \{\sqrt{(b+x)} + \sqrt{(b-x)}\} = 0 \quad (2)$$

This is satisfied if either

$$\sqrt{(a+x)} - \sqrt{(a-x)} = 0 \quad (3)$$

$$\text{or } \sqrt{(b+x)} + \sqrt{(b-x)} = 0 \quad (4)$$

The rational form of (3) is  $(a+x) - (a-x) = 0$  which is satisfied by  $x=0$  and this also satisfies (3).

The rational form of (4) is  $(b+x) - (b-x) = 0$  which requires  $x=0$ , but this does not satisfy (4). Hence the second factor of the left-hand member of (2) cannot vanish.

Therefore the only solution of (2) and  $\therefore$  of (1) is  $x=0$ , derived from (3).

$$6. \quad \sqrt[3]{(a+x)} + \sqrt[3]{(a-x)} = \sqrt[3]{(2a)}$$

$$\text{Cube by the formula } (u+v)^3 = u^3 + v^3 + 3uv(u+v)$$

$$\therefore (a+x) + (a-x) + 3\sqrt[3]{\{2a(a^2 - x^2)\}} = 2a.$$

$$\therefore 2a(a^2 - x^2) = 0,$$

$$\therefore x = \pm a.$$

Both these values belong to the proposed equation.

The rationalizing factors of

$$\sqrt[3]{a+x} + \sqrt[3]{a-x} - \sqrt[3]{2a} = 0$$

$$\text{are } \sqrt[3]{a+x} + \omega \sqrt[3]{a-x} - \omega^2 \sqrt[3]{2a},$$

$$\text{and } \sqrt[3]{a+x} + \omega^2 \sqrt[3]{a-x} - \omega \sqrt[3]{2a}. \text{ See page 257.}$$

The remarks on Ex. 4, will apply *mutatis mutandis* to equations of this type.

$$7. \frac{\sqrt[3]{(a+x)^2} + \sqrt[3]{(a^2-x^2)} + \sqrt[3]{(a-x)^2}}{\sqrt[3]{(a+x)^2} - \sqrt[3]{(a^2-x^2)} + \sqrt[3]{(a-x)^2}} = c. \quad (1)$$

Assume  $\sqrt[3]{a+x} = u$  and  $\sqrt[3]{a-x} = v$

$$\therefore u^3 + v^3 = 2a \text{ and } u^3 - v^3 = 2x,$$

$$\text{and } \therefore \frac{u^3 - v^3}{u^3 + v^3} = \frac{x}{a} \quad (2)$$

Also (1) becomes

$$\frac{u^2 + uv + v^2}{u^2 - uv + v^2} = c \quad (3)$$

Multiply both members by  $\frac{u-v}{u+v}$

$$\therefore \frac{u^3 - v^3}{u^3 + v^3} = c \frac{u-v}{u+v}, \therefore \text{ by (2) } \frac{x}{a} = c \frac{u-v}{u+v} \quad (4)$$

Again adding and subtracting denominators and numerators in (3)

$$\frac{u^2 + v^2}{uv} = \frac{c+1}{c-1}.$$

Adding and subtracting 2 (denominators) and numerators in this

$$\frac{u^2 - 2uv + v^2}{u^2 + 2uv + v^2} = \frac{3-c}{3c-1}, \text{ or } \left(\frac{u-v}{u+v}\right)^2 = \frac{3-c}{3c-1}.$$

$$\therefore \text{ substituting by (4), } \frac{x^2}{a^2} = c^2 \frac{3-c}{3c-1},$$

$$\therefore x = ac \sqrt{\frac{3-c}{3c-1}}.$$



$$8. \{ \sqrt[4]{x+a} + \sqrt[4]{x-a} \}^3 \{ \sqrt[4]{x+a} - \sqrt[4]{x-a} \} = 2c \quad (1)$$

Assume  $u = \sqrt[4]{x+a}$  and  $v = \sqrt[4]{x-a}$ , and (1) becomes

$$(u+v)^3(u-v) = 2c \text{ or } (u+v)^2(u^2-v^2) = 2c \quad (2)$$

$$\text{Also } u^4 - v^4 = 2a \quad \text{or } (u^2+v^2)(u^2-v^2) = 2a \quad (3)$$

$$\text{and } u^4 + v^4 = 2x. \quad (4)$$

$$\text{From (2) and (3), } (u-v)^2(u^2-v^2) = 4a-2c \quad (5)$$

$$\therefore (2) \times (5), (u^2-v^2)^2(u^2-v^2)^2 \text{ or } (u^2-v^2)^4 = 4c(2a-c) \quad (6)$$

Also (3)<sup>2</sup> + (6),

$$\{ (u^2+v^2)^2 + (u^2-v^2)^2 \} (u^2-v^2)^2 = 4(a^2+2ac-c^2)$$

$$\text{or } (u^4+v^4)(u^2-v^2)^2 = 2(a^2+2ac-c^2)$$

Substituting by (4) and (6)

$$2x\sqrt[4]{2ac-c^2} = a^2+2ac-c^2.$$

#### EXERCISE.

$$1. (x+a+b)(x-a+b) + (x+a-b)(x-a-b) = 0.$$

$$2. (a+bx)(b-ax) + (b+cx)(c-bx) + (c+ax)(a-cx) = 0.$$

$$3. (a+bx)(ax-b) + (b+cx)(bx-c) + (c+ax)(cx-a) \\ = \frac{1}{2}(a^2+b^2+c^2).$$

$$4. (a+x)(b-x) + (1+ax)(1-bx) = (a+b)(1+x^2).$$

$$5. (a+x)(b+x)(c-x) + (a+x)(b-x)(c+x) + (a-x)(b+x)(c+x) \\ + (a-x)(b-x)(c+x) + (a-x)(b+x)(c-x) + \\ (a+x)(b-x)(c-x) = 5abc.$$

$$6. (a+x)(b+x)(c+x) + (a+x)(b+x)(c-x) + (a-x)(b-x)(c+x) \\ + (a-x)(b+x)(c+x) + (a+x)(b-x)(c-x) + (a-x)(b+x)(c-x) \\ + (a-x)(b-x)(c+x) + (a-x)(b-x)(c-x) = 8x^2$$

$$7. (a+5b+x)(5a+b+x) = 3(a+b+x)^2.$$

$$8. (a+17b+x)(17a+b+x) = 9(a+b+x)^2.$$

$$9. (9a-7b+3x)(9b-7a+3x) = (3a+3b+x)^2.$$

- (1) 10.  $\frac{ab}{a^2 - b^2x^2} + \frac{cd}{c^2 - d^2x^2} = 0.$  11.  $\frac{x-a}{x+1} + \frac{x+a}{x-1} = 2c.$
- (2) 12.  $\frac{a+x}{a-x} = \frac{x+b}{x-b}.$  13.  $\frac{ax+b}{a+bx} = \frac{cx+d}{c+dx}.$
- (3) 14.  $\frac{a-x}{1-ax} = \frac{1-bx}{b-x}.$  15.  $\frac{a-x}{1-ax} = \frac{4-x}{1-bx}.$
- (4) 16.  $\frac{x+a+2b}{x+a-2b} = \frac{b-2a+2x}{b+2a-2x}.$  17.  $\frac{a+4b+x}{a-4b+x} = \frac{3b-a+x}{3b+a-x}.$
- (5) 18.  $\frac{x+5a+b}{x-3a+b} = \frac{x-a+b}{a-x+3b}.$  19.  $\frac{a-7b+x}{7a-b-x} = \frac{a+5b+x}{5a+b+x}.$
- (6) 20.  $\frac{3a-b-x}{a-3b+x} = \frac{5b-3a+x}{5a-3b+x}.$  21.  $\frac{3a-2b+3x}{a-2b+x} = \frac{x-a+2b}{3x-3a+2b}.$
22.  $\frac{3a-2b+3x}{a-2b+x} = \frac{x-7a+8b}{3x-5a+4b}.$
23.  $\frac{5a-6b+x}{a+x} = \frac{3a-5b+3x}{a+b+x}.$  24.  $\frac{a+b-x}{3a-b-3x} = \frac{3(a-b+x)}{a-5b+x}.$
25.  $\frac{7a+b-x}{5a+3b-3x} = \frac{3(a-b+x)}{a-17b+x}.$
26.  $\frac{5a-b+x}{2(a+2b-x)} = \frac{2(2a-b+x)}{a+11b-x}.$
27.  $\frac{7a-b+x}{7b-a+x} = \frac{a(a+5b+x)}{b(5a+b+x)}.$  28.  $\frac{x+a-b}{x-a+b} = \frac{a(x+a+5b)}{b(x+5a+b)}.$
29.  $\left(\frac{5a-3b+x}{5b-3a+x}\right)^2 = \frac{7a-9b+3x}{7b-9a+3x}.$
30.  $\left(\frac{a+5b+x}{5a+b+x}\right)^2 = \frac{a+17b+x}{17a+b+x}.$
31.  $\left(\frac{7a-b+x}{7b-a+x}\right)^2 = \frac{17a+b-x}{17b+a-x}.$
32.  $\frac{17a+b-x}{a+17b-x} = \frac{a^2(a+17b+x)}{b^2(17a+b+x)}.$

$$33. \frac{(x+7a+b)(x-a+b)}{(5x+3a-11b)(x-a+17b)} = \frac{x-5a+b}{5x+7a-59b}$$

$$34. \frac{(1+3x+5x^2)(x^2+3x+5)}{(1+2x+3x^2)(x^2+2x+3)} = \frac{9}{4}$$

$$35. \frac{\sqrt{1+x^2} + \sqrt{1-x^2}}{\sqrt{1+x^2} - \sqrt{1-x^2}} = \frac{a}{b}$$

$$36. \frac{\sqrt[3]{1+x^2} + \sqrt[3]{1-x^2}}{\sqrt[3]{1+x^2} - \sqrt[3]{1-x^2}} = \frac{a}{b}$$

$$37. \frac{\sqrt[4]{1+x^2} + \sqrt[4]{1-x^2}}{\sqrt[4]{1+x^2} - \sqrt[4]{1-x^2}} = \frac{a}{b}$$

$$38. \frac{\sqrt[5]{1+x^2} + \sqrt[5]{1-x^2}}{\sqrt[5]{1+x^2} - \sqrt[5]{1-x^2}} = \frac{a}{b}$$

$$39. \frac{\sqrt[6]{1+x^2} + \sqrt[6]{x^2-1}}{\sqrt[6]{1+x^2} - \sqrt[6]{x^2-1}} = \frac{a}{b}$$

$$40. \frac{\sqrt[n]{x^2+1} + \sqrt[n]{x^2-1}}{\sqrt[n]{x^2+1} - \sqrt[n]{x^2-1}} = \frac{a}{b}$$

$$41. \sqrt{4a+b-4x} - 2\sqrt{a+b-2x} = \sqrt{b}$$

$$42. \sqrt{3a-2b+2x} - \sqrt{3a-2b-2x} = 2\sqrt{a}$$

$$43. \sqrt{2a-b+2x} - \sqrt{10a-9b-6x} = 4\sqrt{a-b}$$

$$44. \sqrt{3a-4b+5x} + \sqrt{x-a} = 2\sqrt{x+a}$$

$$45. \sqrt{3a-4b+5x} + \sqrt{x-a} = 2\sqrt{2x-2b}$$

$$46. \sqrt{5x-3a+4b} + \sqrt{5x-3a-4b} = 2\sqrt{x+a}$$

$$47. \sqrt{2a+b+2x} + \sqrt{10a+9b-6x} = 2\sqrt{2a+b-2x}$$

$$48. 2\sqrt{2a+b+2x} + \sqrt{10a+b-6x} = \sqrt{10a+9b-6x}$$

$$49. \sqrt{2a-13b+14x} + \sqrt{3(b-2a+2x)} = 2\sqrt{2a-b+2x}$$

$$50. \sqrt{3(7a+b+x)} - \sqrt{a+7b-x} = 2\sqrt{7a+b-x}$$

$$51. \sqrt{(a+x)(x+b)} + \sqrt{(a-x)(x-b)} = 2\sqrt{ax}$$

$$52. \sqrt{(a+x)(x+b)} - \sqrt{(a-x)(x-b)} = 2\sqrt{bx}$$

$$53. \sqrt{ax+x^2} - \sqrt{ax-x^2} = \sqrt{2ax-a^2}$$

$$54. \sqrt{a}$$

$$55. \frac{1}{1+}$$

$$56. \frac{x+}{a-}$$

$$57. \sqrt{\frac{1}{}}$$

$$58. \sqrt{\frac{1}{}}$$

$$59. \sqrt[3]{(}$$

$$60. \sqrt[3]{(}$$

$$61. \sqrt[3]{(}$$

$$62. \sqrt[3]{(}$$

$$63. \sqrt[3]{(}$$

$$64. \sqrt[3]{(}$$

$$65. \sqrt[3]{(}$$

$$66. \sqrt[3]{(}$$

$$67. \sqrt[3]{(}$$

$$68. \sqrt[3]{(}$$

$$69. 5\sqrt{4}$$

$$70. \sqrt{4}$$

$$71. \sqrt{4}$$

$$26\sqrt{}$$

$$72. \sqrt[3]{\frac{1}{}}$$

$$73. 2\sqrt[3]{\frac{1}{}}$$

54.  $\sqrt{ax-x^2} + \sqrt{ax+x^2} = \sqrt{2ax+a^2}$ .
55.  $\frac{1}{1+\sqrt{1-x}} + \frac{1}{1-\sqrt{1-x}} = \frac{2}{9}x$ .
56.  $\frac{x+\sqrt{ax}}{a-\sqrt{ax}} + \frac{a+\sqrt{ax}}{x-\sqrt{ax}} = \frac{x-a}{a}$ .
57.  $\frac{\sqrt{\{(a+x)(x+b)\}} + \sqrt{\{(a-x)(x-b)\}}}{\sqrt{\{(a+x)(x+b)\}} - \sqrt{\{(a-x)(x-b)\}}} = \sqrt{\frac{a}{b}}$ .
58.  $\sqrt{\frac{3a-2b+2x}{3a-2b-2x}} = \frac{\{\sqrt{a} + \sqrt{2a-2b}\}^2}{2b-a}$ .
59.  $\sqrt[3]{a+x} + \sqrt[3]{a-x} = 2\sqrt[3]{a}$ .
60.  $\sqrt[3]{(a+x)^2} - \sqrt[3]{(a^2-x^2)} + \sqrt[3]{(a-x)^2} = \sqrt[3]{a^2}$ .
61.  $\frac{\sqrt[3]{(1+x)^2} + \sqrt[3]{(1-x^2)} + \sqrt[3]{(1-x)^2}}{\sqrt[3]{(1+x)^2} - \sqrt[3]{(1-x^2)} + \sqrt[3]{(1-x)^2}} = 2\frac{1}{3}$ .
62.  $\sqrt[3]{(1+x)^2} + \sqrt[3]{(1-x)^2} = 2\frac{1}{3}\sqrt[3]{(1-x^2)}$ .
63.  $\sqrt[3]{(3+x)} + \sqrt[3]{(3-x)} = \sqrt[3]{6}$ .
64.  $\sqrt[3]{(1+x)^2} + \sqrt[3]{(1-x)^2} = 5\{\sqrt[3]{(1+x)} + \sqrt[3]{(1-x)}\}^2$ .
65.  $\sqrt[3]{(14+x)^2} - \sqrt[3]{(196-x^2)} + \sqrt[3]{(14-x)^2} = 7$ .
66.  $\{\sqrt[3]{(9+x)} + \sqrt[3]{(9-x)}\} \sqrt[3]{(81-x^2)} = 12$ .
67.  $\{\sqrt[3]{(14+x)^2} - \sqrt[3]{(14-x)^2}\} \{\sqrt[3]{(14+x)} - \sqrt[3]{(14-x)}\} = 16$ .
68.  $\{\sqrt[3]{(57+x)^2} + \sqrt[3]{(57-x)^2}\} \{\sqrt[3]{(57-x)} + \sqrt[3]{(57+x)}\} = 100$ .
69.  $5\{\sqrt[4]{(41+x)} + \sqrt[4]{(41-x)}\}^2 = 8\{\sqrt{(41+x)} + \sqrt{(41-x)}\}$ .
70.  $\{\sqrt[4]{(x+5)} + \sqrt[4]{(x-5)}\}^3 \{\sqrt[4]{(x+5)} - \sqrt[4]{(x-5)}\} = 2$ .
71.  $\{\sqrt[4]{(x+1)} + \sqrt[4]{(x-1)}\} \{\sqrt{(x+1)} + \sqrt{(x-1)}\} =$   
 $26\{\sqrt[4]{(x+1)} - \sqrt[4]{(x-1)}\}$ .
72.  $\sqrt[3]{\frac{1+x}{1-x}} + \sqrt[3]{\frac{-x}{1+x}} = a. \quad [y+y^{-1}=a]$ .
73.  $2\{\sqrt[3]{(1+x)^2} + \sqrt[3]{(1-x^2)}\} = (c^2+1)\{\sqrt[3]{(1+x)} + \sqrt[3]{(1-x)}\}^2$ .

74.  $\sqrt[3]{a+x} + \sqrt[3]{a-x} = \sqrt[3]{c}$ .
75.  $\{\sqrt[3]{a+x} + \sqrt[3]{a-x}\} \sqrt[3]{a^2 - x^2} = c$ .
76.  $\sqrt[3]{a+x}^2 = \sqrt[3]{a^2 - x^2} + \sqrt[3]{a-x}^2 = \sqrt[3]{c^2}$ .
77.  $\{\sqrt[3]{a+x}^2 - \sqrt[3]{a-x}^2\} \{\sqrt[3]{a+x} - \sqrt[3]{a-x}\} = c$ .
78.  $\{\sqrt[3]{a+x}^2 + \sqrt[3]{a-x}^2\} \{\sqrt[3]{a+x} + \sqrt[3]{a-x}\} = c$ .
79.  $(a+x)\sqrt[3]{a-x} - (a-x)\sqrt[3]{a+x} = c\{\sqrt[3]{a+x} - \sqrt[3]{a-x}\}$ .
80.  $(a+x)\sqrt[3]{a+x} - (a-x)\sqrt[3]{a-x} = c\{\sqrt[3]{a+x} - \sqrt[3]{a-x}\}$ .
81.  $\{\sqrt[3]{a+x}^2 - \sqrt[3]{a^2 - x^2} + \sqrt[3]{a-x}^2\}^2 =$   
 $c\{\sqrt[3]{a+x} + \sqrt[3]{a-x}\}$ .
82.  $\{\sqrt[4]{a+x} + \sqrt[4]{a-x}\}^2 = (c+1)\{\sqrt{a+x} + \sqrt{a-x}\}$ .
83.  $\{\sqrt[4]{x+a} - \sqrt[4]{x-a}\} \{\sqrt{x+a} + \sqrt{x-a}\}^2 =$   
 $c\{\sqrt[4]{x+a} + \sqrt[4]{x-a}\}$ .

SECTION IV.—QUADRATIC EQUATIONS AND EQUATIONS THAT  
CAN BE RESOLVED AS QUADRATICS.

EXAMPLES.

1.  $x^4 + (ab+1)^2 = (a^2 + b^2)(x^2 + 1) + 2(a^2 - b^2)x + 1,$   
 $\therefore x^4 + a^2b^2 = (a^2 + b^2)x^2 + 2(a^2 - b^2)x + (a-b)^2$   
 $\therefore x^4 + 2abx^2 + a^2b^2 = (a+b)^2x^2 + 2(a^2 - b^2)x + (a-b)^2$   
 $\therefore x^2 + ab = \pm \{(a+b)x + (a-b)\},$   
 or  $x^2 \mp (a+b)x + ab = \pm(a-b),$   
 $\therefore x^2 \mp (a+b)x + \frac{1}{4}(a+b)^2 = \frac{1}{4}(a-b)^2 \pm (a-b),$   
 $\therefore x \mp \frac{1}{2}(a+b) = \frac{1}{2}\sqrt{\{(a-b)^2 \pm 4(a-b)\}}.$

$$2. \frac{(a-x)^2 \sqrt{a-x} + (x-b)^2 \sqrt{x-b}}{(a-x)\sqrt{a-x} + (x-b)\sqrt{x-b}} = a-b.$$

Write  $a-b$  in the form  $(a-x) + (x-b)$  and multiply by the denominator of the left-hand member,

$$\begin{aligned} \therefore (a-x)^2 \sqrt{a-x} + (x-b)^2 \sqrt{x-b} &= \\ (a-x)^2 \sqrt{a-x} + (a-x)(x-b) \{ \sqrt{a-x} + \sqrt{x-b} \} + & \\ (x-b)^2 \sqrt{x-b}, & \end{aligned}$$

$$\therefore (a-x)(x-b) \{ \sqrt{a-x} + \sqrt{x-b} \} = 0,$$

$$\therefore (a-x) = 0, \text{ or } x-b = 0,$$

$$\text{or } \sqrt{a-x} + \sqrt{x-b} = 0.$$

$$x_1 = a, \quad x_2 = b.$$

The equation  $\sqrt{a-x} + \sqrt{x-b} = 0$  has no solution for the sum of two *positive* square-roots, cannot vanish.

The solution  $x = \frac{1}{2}(a+b)$  belongs to the equation

$$\sqrt{a-x} - \sqrt{x-b} = 0.$$

$$3. \frac{ax+b}{bx+a} = \frac{mx+n}{nx+m}.$$

Add and subtract Numerators and Denominators

$$\frac{(a+b)(x+1)}{(a-b)(x-1)} = \frac{(m+n)(x-1)}{(m-n)(x+1)},$$

$$\therefore \left( \frac{x+1}{x-1} \right)^2 = \frac{(a-b)(m+n)}{(a+b)(m-n)} = s^2 \text{ say,}$$

$$\therefore x_1 = \frac{s+1}{s-1}, \quad x_2 = \frac{s-1}{s+1}.$$

$$4. \sqrt[3]{\frac{a-x}{b+x}} + \sqrt[3]{\frac{b+x}{a-x}} = c.$$

Square both members, subtract 4 and extract the square-root.

$$\therefore \sqrt[3]{\frac{a-x}{b+x}} - \sqrt[3]{\frac{b+x}{a-x}} = \pm \sqrt{c^2 - 4}$$

$$\therefore \sqrt[3]{\frac{a-x}{b+x}} = \frac{1}{2}\{c \pm \sqrt{c^2-4}\} = e \text{ say,}$$

$$\therefore \frac{a-x}{b+x} = e^3 \quad \therefore \frac{2x-(a-b)}{a+b} = \frac{1-e^3}{1+e^3},$$

$$\therefore x = \frac{1}{2}\{(a-b) + (a+b) \frac{1-e^3}{1+e^3}\}.$$

Or thus, cube both members,

$$\therefore \frac{a-x}{b+x} + 3c + \frac{b+x}{a-x} = c^3$$

$$\therefore \frac{(a-x)^2 + (b+x)^2}{2(a-x)(b+x)} = c^3 - 3c$$

$$\therefore \frac{\{(b+x)-(a-x)\}^2}{\{(b+x)+(a-x)\}^2} = \frac{c^3 - 3c - 2}{c^3 - 3c + 2} = \frac{(c+1)^2(c-2)}{(c-1)^2(c+2)}$$

$$\therefore \frac{2x-(a-b)}{a+b} = \frac{c+1}{c-1} \sqrt{\frac{c-2}{c+2}}.$$

(Prove that  $\frac{1-e^3}{1+e^3} = \frac{c+1}{c-1} \sqrt{\frac{c-2}{c+2}}$  if  $2e = c \pm \sqrt{c^2-4}$ .)

5.  $\frac{\sqrt{a-x} - \sqrt{b-x}}{\sqrt{a-x} + \sqrt{b-x}} = \frac{\sqrt{\{(a-x)(b-x)\}}}{c}$ . Rationalize Denom.

$$\therefore \frac{\{\sqrt{a-x} - \sqrt{b-x}\}^2}{(a-b)} = \frac{\sqrt{\{(a-x)(b-x)\}}}{c},$$

or  $\frac{\{\sqrt{a-x} - \sqrt{b-x}\}^2}{\sqrt{\{(a-x)(b-x)\}}} = \frac{a-b}{c}$ , (A)

$$\therefore \frac{\{\sqrt{a-x} + \sqrt{b-x}\}^2}{\sqrt{\{(a-x)(b-x)\}}} = \frac{a-b}{a-b+4c};$$

$$\therefore \frac{\sqrt{\{(a-x)(b-x)\}}}{c} = \sqrt{\frac{a-b}{a-b+4c}}. \quad (B)$$

Also from (A),

$$\frac{a+b-2x}{\sqrt{\{(a-x)(b-x)\}}} = \frac{a-b+2c}{c},$$

Multiply (B) and (C) member by member

$$\therefore a+b-2x = (a-b+2c) \sqrt{\frac{a-b}{a-b+4c}}$$

$$\therefore x = \frac{1}{2} \left\{ a+b - (a-b+2c) \sqrt{\frac{a-b}{a-b+4c}} \right\}$$

$$6. \quad x^4 - 4 = \frac{x^2 + 20}{x^2 - 2}; \quad x^6 - 2x^4 - 5x^2 - 12 = 0.$$

Find the rational linear factors of the left-hand member by the method of Art. XXVII., page 90.

$$\therefore (x-2)(x+2)(x^4 + 2x^2 + 3) = 0,$$

$$\therefore x-2=0, \text{ or } x+2=0, \text{ or } x^4 + 2x^2 + 3 = 0.$$

The last of these equations may be solved as a quadratic giving

$$x^2 = -1 \pm 2\sqrt{-2}, \quad \therefore x = \pm 1 \pm \sqrt{-2},$$

$$\therefore x_1 = 2, \quad x_2 = -2, \quad x_3 = 1 + \sqrt{-2}, \quad x_4 = 1 - \sqrt{-2},$$

$$x_5 = -1 + \sqrt{-2}, \quad x_6 = -1 - \sqrt{-2}.$$

N.B.—In solving numerical equations of the higher orders, *the rational linear factors should always be found and separated as disjunctive equations*, before other methods of reduction are applied. Such separation may always be effected by the methods of Arts. XXVII. to XXX., and unless it is done the application of the higher methods may actually fail. Thus, if it be attempted to solve as a cubic the equation,

$$x^3 - 9x - 10 = 0$$

the result is  $x = \{5 + \sqrt{-2}\}^{\frac{1}{3}} + \{5 - \sqrt{-2}\}^{\frac{1}{3}}$ , which can be reduced only by trial. The left-hand member can however be easily factored by the method of Art. XXVII. and the equation reduces to

$$(x+2)(x^2 - 2x - 5) = 0,$$

which gives  $x = 2$  or  $1 \pm \sqrt{6}$ .

Denom.

(A)

(B)



$$7. (x-2)^7 - x^7 + 2^7 = 0.$$

Factor, (See No. 20, p. 89), rejecting constant factors,

$$\therefore (x-2)(x^2-2x+1)^2 = 0$$

$$\therefore x=0, \text{ or } x-2=0, \text{ or } x^2-2x+4=0.$$

The last equation gives  $x = 1 \pm \sqrt{-3}$ .

## EXERCISE.

Solve the following equations :

$$1. (x+a+b)^3 = x^3 + a^3 + b^3. \quad 2. (x+a+b)^5 = x^5 + a^5 + b^5.$$

$$3. (a-b)x^3 + (b-x)a^3 + (x-a)b^3 = 0.$$

$$4. (a-b)x^2 + (x-b)a^2 + (x+a)b^2 = 2abx.$$

$$5. (x-a)^5 + (a-b)^5 + (b-x)^5 = 0.$$

$$6. (x-a)^7 + (a-b)^7 + (b-x)^7 = 0.$$

$$7. (a^3-b)x^4 + (x^3-a)b^4 + (x-a)a^4 = abx(a^2b^2x^2-1).$$

$$8. (x-a)(x-b)(a-b) + (x-b)(x-c)(b-c) + (x-c)(x-a)(c-a) = 0.$$

$$9. \frac{x^5-1}{x-1} = 0. \quad 10. \frac{x^{12}-1}{x^4-1} = 0.$$

$$11. \frac{x^{16}-1}{x^4-1} = 0. \quad 12. \frac{x^{20}-1}{x^4-1} = 0.$$

$$13. x^4 + 5x^3 - 16x^2 + 20x - 16 = 0. \quad (\text{See Art. XXII.})$$

$$14. x^4 - 3x^3 + 5x^2 + 6x + 4 = 0.$$

$$15. (x-a)^4 + x^4 + a^4 = 0. \quad 16. 2x^3 = (x-6)^2.$$

$$17. x(x-2)^2(x+2) = 2. \quad 18. (4x^3-17)x+12=0.$$

$$19. x^4 + (ab+1)^2 = (a^2+b^2)(x^2+1) + 2(a^2-b^2)x+1.$$

$$20. x^2(x-169)^2 + 17x = x^2 - 3540.$$

21.

22.

24.

26.

27.

29.

31.

33.

34.

35.

36.

37.

39.

41.

43.

45.

46.

$$21. \quad 6x(x^2+1)^2 + (2x^2+5)^3 = 150x+1.$$

$$22. \quad 2x(x-1)^2 + 2 = (x+1)^2. \quad 23. \quad x^4 = 12x+5.$$

$$24. \quad 3x^4 = 12x^3 + 1. \quad 25. \quad (x+4)^3 = 3(2x-1)^2.$$

$$26. \quad \sqrt{x^2+m^2} + \sqrt{(n-x)^2+m^2} = \sqrt{(x-\frac{1}{2}n)^2 + (\frac{1}{2}n\sqrt{3}-m)^2}.$$

$$27. \quad \frac{(x+1)^4}{(x^2+1)(x-1)^2} = \frac{m}{n}. \quad 28. \quad \frac{(x+1)^5}{x(x^3+1)} = \frac{m}{n}.$$

$$29. \quad \frac{(x^2+1)(x^3+1)}{(x+1)(x^4+1)} = \frac{m}{n}. \quad 30. \quad \frac{(x^2+1)(x^3+1)}{(x^2-1)(x^3-1)} = \frac{m}{n}.$$

$$31. \quad \frac{(x^2+1)(x^3+1)}{x^2(x+1)} = \frac{m}{n}. \quad 32. \quad \frac{(x^3-1)^2}{x(x^2+1)(x-1)^2} = \frac{m}{n}.$$

$$33. \quad \frac{x(x+1)^2}{(x^2+1)(x-1)^2} = \frac{n(n-m)}{2m(2m-n)}.$$

$$34. \quad \frac{(x^3+1)^2}{x(x^2-1)^2} = \frac{4m^2}{m^2-n^2}.$$

$$35. \quad \frac{(x-1)(x^2+1)^2}{(x^3-1)(x+1)^2} = \frac{2(m-n)^2}{mn}.$$

$$36. \quad \frac{x^6-1}{(x+1)(x^5-1)} = \frac{2m}{2m-n}.$$

$$37. \quad \frac{(x^3-1)(x+1)^3}{(x^3+1)(x-1)^3} = \frac{m+n}{m-n}. \quad 38. \quad \frac{(x+1)(x^4+1)}{(x-1)(x^4-1)} = \frac{m+n}{m-n}.$$

$$39. \quad x^3 = \frac{ax-b}{bx-a}. \quad 40. \quad x^4 = \frac{ax-b}{bx-a}.$$



$$41. \quad x^5 = \frac{ax-b}{bx-a}. \quad 42. \quad x^4 = \frac{ax^2+bx+c}{a+bx+cx^2}.$$

$$43. \quad x^2 = (x-1)^2(x^2+1). \quad 44. \quad a^2x^2 = (a-x)^2(a^2-x^2).$$





$$45. \quad x^2 = (x-a)^2(x^2-1).$$

$$46. \quad a\sqrt{x^2+1} - x\sqrt{x^2+a^2} = cx.$$





**IMAGE EVALUATION  
TEST TARGET (MT-3)**



**Photographic  
Sciences  
Corporation**

23 WEST MAIN STREET  
WEBSTER, N.Y. 14580  
(716) 872-4503



47.  $\sqrt[3]{(a^3+x^3)} + \sqrt[3]{(a^3-x^3)} = \sqrt[3]{(a^6-x^6)^2}$ .
48.  $m(x+m-n)(x-m+7n)^2 = n(x-m+n)(x+7m-n)^2$ .
49.  $m^2(x+m+17n)(x-m-5n)^2 = n^2(x+17m+n)(x-5m+n)^2$ .
50.  $m^2(x+m+17n)(x-m+7n)^2 = n^2(x+17m+n)(x+7m-n)^2$ .
51.  $\frac{\sqrt{(x-a)} + \sqrt{(x-b)}}{\sqrt{(x-a)} - \sqrt{(x-b)}} = \sqrt{\frac{x-a}{x-b}}$ .
52.  $\frac{\sqrt{(x-a)} + \sqrt{(x-b)}}{\sqrt{(x-a)} - \sqrt{(x-b)}} = \sqrt{\frac{a-x}{x-b}}$ .
53.  $\sqrt{\frac{a-x}{b+x}} - \sqrt{\frac{b+x}{a-x}} = c$ .
54.  $\sqrt{\frac{a-x}{b-x}} + \sqrt{\frac{b-x}{a-x}} = c$ .
55.  $\sqrt[3]{\frac{a-x}{b+x}} - \sqrt[3]{\frac{b+x}{a-x}} = c$ .
56.  $\sqrt[3]{\left(\frac{a-x}{b-x}\right)^2} - \sqrt[3]{\left(\frac{b-x}{a-x}\right)^2} = c$ .
57.  $\sqrt[4]{\frac{a-x}{b+x}} + \sqrt[4]{\frac{b+x}{a-x}} = c$ .
58.  $\sqrt[4]{\frac{a-x}{b-x}} - \sqrt[4]{\frac{b-x}{a-x}} = c$ .
59.  $\sqrt[5]{\frac{a-x}{b+x}} + \sqrt[5]{\frac{b+x}{a-x}} = c$ .
60.  $\sqrt[5]{\frac{a-x}{b-x}} - \sqrt[5]{\frac{b-x}{a-x}} = c$ .
61.  $\sqrt[6]{\frac{a-x}{b+x}} + \sqrt[6]{\frac{b+x}{a-x}} = c$ .
62.  $\sqrt[6]{\frac{a-x}{b-x}} - \sqrt[6]{\frac{b-x}{a-x}} = c$ .
63.  $\frac{\sqrt{(a-x)^3} + \sqrt{(b-x)^3}}{\sqrt{(a-x)} + \sqrt{(b-x)}} = c$ .
64.  $\frac{\sqrt{(a-x)^3} + \sqrt{(b-x)^3}}{\{\sqrt{(a-x)} + \sqrt{(b-x)}\}^3} = c$ .
65.  $\frac{\sqrt{(a-x)^3} + \sqrt{(b-x)^3}}{\sqrt{(a-x)} - \sqrt{(b-x)}} = c$ .
66.  $\frac{\{\sqrt{(a-x)} + \sqrt{(b-x)}\}^3}{\sqrt{(a-x)} - \sqrt{(b-x)}} = c$ .
67.  $\frac{\sqrt{(a-x)^5} + \sqrt{(x-b)^5}}{\sqrt{(a-x)} + \sqrt{(x-b)}} = c$ .

$$68. \frac{\sqrt{(a-x)^5} - \sqrt{(x-b)^5}}{\{\sqrt{(a-x)} - \sqrt{(x-b)}\}^5} = c.$$

$$69. \frac{\sqrt{(a-x)^5} + \sqrt{(x+b)^5}}{\sqrt{(a-x)^3} + \sqrt{(x+b)^3}} = \sqrt{\{(a-x)(x+b)\}}.$$

$$70. \frac{\sqrt{(a-x)^3} + \sqrt{(x+b)^3}}{\sqrt{(a-x)} + \sqrt{(x+b)}} = \frac{(a+b)^2}{4\sqrt{\{(a-x)(x+b)\}}}.$$

$$71. \frac{x^3 + (a-x^2)\sqrt{(a-x^2)}}{x + \sqrt{(a-x^2)}} = c.$$

$$72. \frac{x^3 + (a^2-x^2)\sqrt{(a^2-x^2)}}{x + \sqrt{(a^2-x^2)}} = cx\sqrt{(a^2-x^2)}.$$

$$73. \sqrt[3]{(a-x)^2} - \sqrt[3]{\{(a-x)(x-b)\}} + \sqrt[3]{(x+b)^2} = \sqrt[3]{(a^2-ab+b^2)}$$

$$74. \frac{b\sqrt{(a-x)} + c\sqrt{(x-b)}}{\sqrt{(a-x)} + \sqrt{(x-b)}} = x.$$

$$75. \frac{a\sqrt{(a-x)} + b\sqrt{(x-b)}}{\sqrt{(a-x)} + \sqrt{(x-b)}} = x.$$

$$76. \frac{\sqrt{(x-a)} + \sqrt{(x+a)} - \sqrt{(2a)}}{\sqrt{(x-a)} - \sqrt{(x+a)} + \sqrt{(2a)}} = \sqrt[4]{\frac{x+c}{x-c}}.$$

$$77. \frac{\sqrt{(a-x)} + \sqrt{c}}{\sqrt{(x-b)} + \sqrt{c}} = \sqrt[4]{\frac{a-x}{x-b}}.$$

$$78. \sqrt[3]{(a-x)^2} - \sqrt[3]{\{(a-x)(x+b)\}} + \sqrt[3]{(x+b)^2} = \sqrt[3]{(a^2-ab+b^2)}.$$

$$79. \{\sqrt[3]{(a-x)^2} - \sqrt[3]{\{(a-x)(x-b)\}} + \sqrt[3]{(x-b)^2}\}^2 = (a-b)\{\sqrt[3]{(a-x)} + \sqrt[3]{(x-b)}\}.$$

$$80. \{\sqrt[3]{(a-x)^2} + \sqrt[3]{(b+x)^2}\}^2 = (a+b)\{\sqrt[3]{(a-x)} + \sqrt[3]{(b+x)}\}.$$

$$81. \sqrt[3]{(a-x)} + \sqrt[3]{(x-b)} = \sqrt[3]{c}.$$

$$82. \sqrt[3]{(a+x)^2} - \sqrt[3]{(a-x)^2} = \sqrt[3]{(2cx)}.$$

$$83. \sqrt[3]{(a-x)^2} + \sqrt[3]{\{(a-x)(b-x)\}} + \sqrt[3]{(b-x)^2} = \sqrt[3]{c^2}.$$

$$84. \sqrt[3]{(a-x)^2} - \sqrt[3]{\{(a-x)(x+b)\}} + \sqrt[3]{(x+b)^2} = c\{\sqrt[3]{(a-x)} + \sqrt[3]{(x+b)}\}.$$

$$85. \{ \sqrt[3]{a-x} + \sqrt[3]{x+b} \} \sqrt[3]{(a-x)(x+b)} = c.$$

$$86. \sqrt[3]{(a-x)^2} + \sqrt[3]{(x-b)^2} = c \{ \sqrt[3]{a-x} + \sqrt[3]{x-b} \}^2.$$

$$87. x + \sqrt[3]{(a^3-x^3)} = \frac{c^3}{x \sqrt[3]{(a^3-x^3)}}.$$

$$88. \frac{a^3}{x^3-b^3} = \frac{x + \sqrt[3]{(2b^3-x^3)}}{x - \sqrt[3]{(2b^3-x^3)}}.$$

$$89. (a+x)\sqrt[4]{a+x} + (a-x)\sqrt[4]{a-x} = a \{ \sqrt[4]{a+x} + \sqrt[4]{a-x} \}.$$

$$90. (a+x)\sqrt[4]{a-x} + (a-x)\sqrt[4]{a+x} = a \{ \sqrt[4]{a+x} + \sqrt[4]{a-x} \}.$$

$$91. \sqrt[4]{26-x} + \sqrt[4]{x-10} = 2.$$

$$92. \{ \sqrt[4]{a-x} + \sqrt[4]{x-b} \}^2 = c \{ \sqrt{a-x} + \sqrt{x-b} \}.$$

$$93. (a-x)\sqrt[4]{a-x} + (x-b)\sqrt[4]{x-b} = (a-b) \{ \sqrt[4]{a-x} + \sqrt[4]{x-b} \}.$$

$$94. \{ \sqrt[4]{a-x} + \sqrt[4]{x-b} \}^2 \{ \sqrt{a-x} + \sqrt{x-b} \} = c(a+b-2x).$$

$$95. \{ \sqrt[4]{a-x} + \sqrt[4]{b-x} \} \{ \sqrt{a-x} + \sqrt{b-x} \}^2 = c \{ \sqrt[4]{a-x} - \sqrt[4]{b-x} \}.$$

$$96. a \sqrt{1+x^2} - x \sqrt{x^2+a^2} = e.$$

$$97. (a-x)\sqrt[3]{x-b} + (x-b)\sqrt[3]{a-x} = c \{ \sqrt[3]{a-x} + \sqrt[3]{x-b} \}^3.$$

$$98. \{ \sqrt[3]{a-x} + \sqrt[3]{b+x} \}^5 = c \{ \sqrt[3]{a-x}^2 + \sqrt[3]{b+x}^2 \}.$$

$$99. \{ \sqrt[3]{a-x} + \sqrt[3]{b+x} \}^5 = c \sqrt[3]{(x-x)(b+x)}.$$

$$100. \sqrt[3]{(a-x)^2} - \sqrt[3]{(b-x)^2} = c \sqrt[3]{(a+b-2x)}.$$

$$101. \sqrt[4]{a-x} + \sqrt[4]{x-b} = \sqrt[4]{c}.$$

$$102. \sqrt[5]{a-x} + \sqrt[5]{x-b} = \sqrt[5]{c}.$$

$$103. \frac{(a-x)\sqrt[4]{a-x} + (x-b)\sqrt[4]{x-b}}{(a-x)\sqrt[4]{x-b} + (x-b)\sqrt[4]{a-x}} = c.$$

$$104. \frac{(a-x)\sqrt[4]{b-x} + (b-x)\sqrt[4]{a-x}}{\sqrt[4]{a-x} - \sqrt[4]{b-x}} = c.$$

$$105. \frac{\sqrt[4]{a-x} + \sqrt[4]{x-b}}{\sqrt[4]{a-x} - \sqrt[4]{x-b}} = \frac{c}{a+b-2x}.$$

106.

107.

108.

109.

110.

111.

112.

SECT

1.

I +

∴ x-

I ÷

By II

Also

(Not an  
ently in  
square-



106.  $\frac{\{\sqrt[4]{a-x} + \sqrt[4]{b-x}\}^5}{\sqrt[4]{a-x} - \sqrt[4]{b-x}} = c.$
107.  $\{a-x\}^5 \sqrt[5]{a-x} - \{x-b\}^5 \sqrt[5]{x-b} = c \{ \sqrt[5]{a-x} - \sqrt[5]{x-b} \}.$
108.  $\{a-x\}^5 \sqrt[5]{x+b} - \{x+b\}^5 \sqrt[5]{x-a} = c \{ \sqrt[5]{a-x} - \sqrt[5]{x+b} \}.$
109.  $\{ \sqrt[5]{a-x}^3 + \sqrt[5]{x-b}^3 \} \sqrt[5]{(a-x)(x-b)} = c.$
110.  $\{ \sqrt[5]{a-x} - \sqrt[5]{x-b} \}^3 \{ \sqrt[5]{a-x}^2 - \sqrt[5]{x-b}^2 \} = c.$
111.  $\{ \sqrt[5]{a-x}^2 - \sqrt[5]{x-b}^2 \}^2 \{ \sqrt[5]{a-x} + \sqrt[5]{x-b} \} = c.$
112.  $\{ \sqrt[5]{a-x}^3 + \sqrt[5]{x+b}^3 \}^2 = c \{ \sqrt[5]{a-x} + \sqrt[5]{x+b} \}.$

---

SECTION V.—QUADRATIC EQUATIONS INVOLVING TWO OR MORE VARIABLES.

---

1.  $(x+y)(x^2+y^2) = a,$  I.  
 $x^2y + xy^2 = c.$  II.

I+2II.  $\therefore (x+y)^3 = a+2c$

$\therefore x+y = \sqrt[3]{a+2c}.$  (Any one of the three cube-roots). III.

I  $\div$  II  $\frac{x^2+y^2}{xy} = \frac{a}{c}; \therefore \left(\frac{x-y}{x+y}\right)^2 = \frac{a-2c}{a+2c}.$

By III.  $x-y = \frac{\sqrt{a-2c}}{\sqrt[6]{a+2c}}.$

Also  $x+y = \frac{\sqrt{a+2c}}{\sqrt[6]{a+2c}},$

$$\therefore x = \frac{\sqrt{a+2c} + \sqrt{a-2c}}{2\sqrt[6]{a+2c}}$$

$$y = \frac{\sqrt{a+2c} - \sqrt{a-2c}}{2\sqrt[6]{a+2c}}.$$

(Not any one of the six sixth-roots of  $a+2c$  may be used indifferently in the denominator, but only any cube-root of whichever square-root of  $a+2c$  is used in the numerator. Thus if the radi-

cal sign be restricted to denote merely the arithmetical root, if  $k$  be defined by the equation  $k^2 - k + 1 = 0$ , and if  $m$  and  $n$  indicate any integers whatever, equal or unequal, the value of  $x$  may be written

$$\{k^{2m} \sqrt{(a+2c)} + k^{2n-m} \sqrt{(a-2c)}\} \div 2^m \sqrt{(a+2c)}.$$

$$2. \quad 8x^2 - 5xy + 3y^2 = 9(x+y) \quad \text{I.}$$

$$11x^2 - 8xy + 5y^2 = 13(x+y) \quad \text{II.}$$

1st Method. Eliminate  $(x+y)$ .

$$\therefore 104x^2 - 65xy + 39y^2 = 99x^2 - 72xy + 45y^2,$$

$$\therefore 5x^2 + 7xy - 6y^2 = 0,$$

$$\therefore (5x - 3y)(x + 2y) = 0,$$

$$\therefore x = \frac{3}{5}y \text{ or } -2y.$$

Substitute these values for  $x$  in I.

$$\therefore 72y^2 = 360y \text{ or } 45y^2 = -9y$$

$$\therefore y = 0, \text{ or } 5, \text{ or } -\frac{1}{5},$$

$$\text{and } x = 0, \text{ or } 3, \text{ or } \frac{3}{5}.$$

2nd Method. Take the sum of the products of I. and II. by arbitrary multipliers  $k$  and  $l$ ,

$$k(8x^2 - 5xy + 3y^2) + l(11x^2 - 8xy + 5y^2) = (9k + 13l)(x + y). \quad \text{III.}$$

Determine  $k$  and  $l$  so that the left-hand member of III. may, like the right-hand member, be a multiple of  $x + y$ . This may be done by putting  $x = -y$  in III. from which

$$16k + 24l = 0, \quad \therefore 2k = -3l$$

$$\therefore \text{if } k = 3, \quad l = -2.$$

Substituting these values in III., it becomes

$$2x^2 + xy - y^2 = x + y$$

$$\therefore (x+y)(2x-y) = x+y, \text{ or } (x+y)(2x-y-1) = 0.$$

$$\therefore \text{either } x+y = 0, \text{ or } 2x-y-1 = 0,$$

$$\therefore y = -x, \text{ or } 2x-1.$$

Substituting these values for  $y$  in I., it becomes

$$16x^2 = 0, \text{ or } 10x^2 - 7x + 3 = 27x - 9,$$

$$\therefore x = 0, \text{ or } 3, \text{ or } \frac{2}{3};$$

$$\text{and } y = 0, \text{ or } 5, \text{ or } -\frac{1}{5}.$$

$$8. \quad \frac{x^5 + y^5}{x^3 + y^3} = \frac{a^5 + b^5}{a^3 + b^3} \quad \text{I.}$$

$$x^2 + xy + y^2 = a^2 + ab + b^2 \quad \text{II.}$$

$$\text{I.} \div \text{II.}, \therefore \frac{x^4 - x^3y + x^2y^2 - xy^3 + y^4}{(x^2 + y^2)^2 + x^2y^2} =$$

$$\frac{a^4 - a^3b + a^2b^2 - ab^3 + b^4}{(a^2 + b^2)^2 - a^2b^2},$$

$$\therefore \frac{x^3y + xy^3}{(x^2 + y^2)^2 - x^2y^2} = \frac{a^3b + ab^3}{(a^2 + b^2)^2 - a^2b^2} \quad \text{III.}$$

Write  $z$  for  $\frac{xy}{x^2 + y^2}$  and  $k$  for  $\frac{ab}{a^2 + b^2}$

$$\text{III.}, \therefore \frac{z}{1 - z^2} = \frac{k}{1 - k^2}, \therefore z = k \text{ or } -\frac{1}{k},$$

$$\therefore \frac{xy}{x^2 + y^2} = \frac{ab}{a^2 + b^2} \text{ or } \frac{a^2 + b^2}{-ab},$$

$$\therefore \frac{xy}{x^2 + xy + y^2} = \frac{ab}{a^2 + ab + b^2}, \text{ or } \frac{a^2 + b^2}{a^2 - ab + b^2}.$$

$$\text{II.}, \therefore xy = ab, \text{ or } (a^2 + b^2) \frac{a^2 + ab + b^2}{a^2 - ab + b^2} \quad \text{IV.}$$

$$\sqrt{(\text{II.} + \text{IV.})}, \therefore x + y = \pm(a + b)$$

$$\text{or } \sqrt{(2a^2 - ab + 2b^2)} \sqrt{\frac{a^2 + ab + b^2}{a^2 - ab + b^2}};$$

$$\sqrt{(\text{II.} - 3\text{IV.})} \text{ and } x - y = \pm(a - b),$$

$$j\sqrt{(2a^2 + ab + 2b^2)} \sqrt{\frac{a^2 + ab + b^2}{a^2 - ab + b^2}}.$$

$$\therefore x = \pm a, \pm b \text{ or}$$

$$\frac{1}{2} \{ \sqrt{(2a^2 - ab + b^2)} + j \sqrt{(2a^2 + ab + 2b^2)} \} \sqrt{\frac{a^2 + ab + b^2}{a^2 - ab + b^2}}.$$

$$y = \pm b; \pm a \text{ or}$$

$$\frac{1}{2} \{ \sqrt{(2a^2 - ab + b^2)} - j \sqrt{(2a^2 + ab + 2b^2)} \} \sqrt{\frac{a^2 + ab + b^2}{a^2 - ab + b^2}}.$$

$$4. \quad (x^2 + y^2)(x^3 + y^3) = a, \quad \text{I.}$$

$$(x + y)(x^4 + y^4) = b. \quad \text{II.}$$

$$\text{Put } z = \frac{xy}{x^2 + y^2}, \therefore \frac{1 - z}{1 - 2z^2} = \frac{a}{b}$$

$$\therefore 2az^2 - bz - (a - b) = 0$$

$$\therefore 4az^2 = b \pm \sqrt{(8a^2 - 8ab + b^2)} = b + r \text{ say.}$$

$$\therefore \frac{xy}{x^2 + y^2} = \frac{b + r}{4a} \quad \text{III.}$$

$$\therefore \frac{x + y}{x - y} = \pm \sqrt{\frac{2a + b + r}{2a - b - r}}$$

$$\therefore \frac{x}{y} = \frac{\sqrt{(2a + b + r)} + \sqrt{(2a - b - r)}}{\sqrt{(2a + b + r)} - \sqrt{(2a - b - r)}}$$

$$= \frac{\{\sqrt{(2a + b + r)} + \sqrt{(2a - b - r)}\}^2}{2(b + r)} \quad \text{IV.}$$

$$\text{I}^2. \quad (x^2 + y^2 + 2xy)(x^2 + y^2)^2 \{(x^2 + y^2) - xy\}^2 = a^2.$$

$$\text{III.} \quad \therefore (xy)^5 \left\{ \frac{4a + 2b + 2r}{b + r} \right\} \left\{ \frac{4a}{b + r} \right\}^2 \left\{ \frac{4a - b - r}{b + r} \right\}^2 - a^2,$$

$$\therefore x^{10} \left( \frac{y}{x} \right)^5 \left\{ \frac{32a^2(2a + b + r)(4a - b - r)^2}{(b + r)^5} \right\} = a$$

$$\therefore x^{10} = \left( \frac{x}{y} \right)^5 \left\{ \frac{(b + r)^5}{32(2a + b + r)(4a - b - r)^2} \right\}$$

$$\text{IV} \quad = \frac{\{\sqrt{(2a + b + r)} + \sqrt{(2a - b - r)}\}^{10}}{1024(2a + b + r)(4a - b - r)^2} \quad \text{X.}$$

$$\therefore x = \frac{\sqrt{(2a+b+r)} + \sqrt{(2a-b-r)}}{2\sqrt{\{(2a+b+r)(4a-b-r)^2\}}}$$

in which  $r = \pm \sqrt{(8a^2 - 8ab + b^2)}$ .

The value of  $y$  may be derived from that of  $x$  by the first form in IV.

I.	5. $x^4 = ax - by,$	I.
II.	$y^4 = ay - bx.$	II.
	$x.I. - y.II. \quad x^5 - y^5 = a(x^2 - y^2)$	
	$y.I. - x.II. \quad xy(x^3 - y^3) = b(x^2 - y^2),$	
	$\therefore$ either $x - y = 0$ from which $x = y = 0$ , or $\sqrt[3]{(a-b)}$	III.
	or $x^4 + x^3y + x^2y^2 + xy^3 + y^4 = a(x+y)$	IV.
III.	and $xy(x^2 + xy + y^2) = b(x+y)$	V.
	(IV. + V.) $(x+y)^2(x^2 + y^2) = a + b$	VI.
	V. $(x+y)^4 - (x^2 + y^2)^2 = 4b(x+y)$	VII.
	$\sqrt{(VII^2 + 4.VI.)} \quad (x+y)^4 + (x^2 + y^2)^2 = 2t(x+y)$	VIII.
	in which $t = \sqrt{\{(a+b)^2 + 4b^2\}}$ .	IX.
IV.	$\frac{1}{2}(VII. + VIII.), \therefore (x+y)^4 = (2b+t)(x+y)$	
	$\therefore (x+y)^3 = 2b+t$	
	$\therefore (x+y) = \sqrt[3]{(2b+t)}$	X.
	VI. $\div X^2 \quad \therefore x^2 + y^2 = \frac{a+b}{\sqrt[3]{(2b+t)}}$	XI.
	2.XI. $- X^2 \quad \therefore (x-y)^2 = \frac{2(a+b)}{\sqrt[3]{(2b+t)}} - \sqrt[3]{(2b+t)^3} = \frac{2a-t}{\sqrt[3]{(2b+t)}}$	
	$\therefore x-y = \frac{\sqrt{(2a-t)}}{\sqrt[6]{(2b+t)}}$	
X.	and $x+y = \frac{\sqrt{(2b+t)}}{\sqrt[6]{(2b+t)}}$	

$$\therefore x = \frac{\sqrt{(2b+t)} + \sqrt{(2a-t)}}{\sqrt[4]{(2b+t)}}$$

$$\text{and } y = \frac{\sqrt{(2b+t)} - \sqrt{(2a-t)}}{\sqrt[4]{(2b+t)}};$$

in which  $t = \sqrt{(a^2 + 2ab + 5b^2)}$ .

$$6. \quad x^4 - c^4 = m(x+y)^4;$$

$$y^4 + c^4 = n(x-y)^4.$$

II.

$$\text{Let } z = \frac{x+y}{x-y}, \therefore z+1 = \frac{2x}{x-y} \text{ and } z-1 = \frac{2y}{x-y}$$

III.

$$\text{I. + II.} \quad x^4 + y^4 = m(x+y)^4 + n(x-y)^4$$

$$\therefore (z+1)^4 + (z-1)^4 = 16(mz^4 + n)$$

$$\therefore (8m-1)z^4 - 6z^2 + (8n-1) = 0,$$

$$\therefore z = \sqrt{\frac{3 + \sqrt{9 - (8m-1)(8n-1)}}{8m-1}}$$

IV.

$$\text{II. \& III.} \quad (z-1)^4(x-y)^4 + 16c^4 = 16n(x-y)^4$$

$$\therefore x-y = \frac{2c}{\sqrt[4]{16n - (z-1)^4}}$$

V.

$$\text{and } x+y = \frac{2cz}{\sqrt[4]{16n - (z-1)^4}}$$

$$\therefore x = \frac{c(z+1)}{\sqrt[4]{16n - (z-1)^4}} = \frac{c(z+1)}{\sqrt[4]{(z+1)^4 - 16mz^4}}$$

and  $y = \frac{c(z-1)}{\sqrt[4]{16n - (z-1)^4}}$ , and the value of  $z$  is given by IV.

$$7. \quad x^2 + y^2 = \frac{1}{3}(2m + n^2),$$

$$x^3 + y^3 = mn.$$

$$\therefore (x+y)^2 - 2xy = \frac{1}{3}(2m + n^2)$$

$$\text{and } (x+y)^3 - 3xy(x+y) = mn.$$

Let  $u = x + y$  and  $v = xy$ , and the equations become

$$u^2 - 2v = \frac{1}{2}(2m + n^2);$$

$$u^3 - 3uv = mn.$$

Eliminate  $v$ ,  $\therefore u^3 - (2m + n^2)u + 2mn = 0$ ,

$$\therefore u^4 - (2m + n^2)u^2 + 2mn u = 0,$$

$$\therefore u^4 - 2mu^2 + m^2 = n^2u^2 - 2mnu + m^2.$$

$$\therefore u^2 - m = \pm(nu - m),$$

$\therefore u = n$ , (the value  $u = 0$  was introduced by the multiplication by  $u$ ),

$$\text{or } u^2 + nu - 2m = 0,$$

$$\therefore u = \frac{1}{2}\{-n \pm \sqrt{(n^2 + 8m)}\}$$

$$\therefore v = \frac{1}{2}(n^2 - m) \text{ or } \frac{1}{8}\{n^2 + 8m \mp 3n\sqrt{(n^2 + 8m)}\}$$

$\therefore u$  and  $v$  are completely determined.

Also  $x + y = u$ ,  $x - y = \sqrt{(u^2 - 4v)}$

$$\therefore x = \frac{1}{2}\{u + \sqrt{(u^2 - 4v)}\};$$

$$\therefore y = \frac{1}{2}\{u - \sqrt{(u^2 - 4v)}\}.$$

If  $m = 7$  and  $n = 5$ , the above equations become

$$x^2 + y^2 = 13, \text{ and } x^3 + y^3 = 35.$$

Solving, as above, gives

$$u = 5, \text{ or } 2, \text{ or } -7,$$

$$2v = 12, \text{ or } -9, \text{ or } 36,$$

$$\therefore x + y = 5, \text{ or } 2, \text{ or } -7,$$

$$x - y = \pm 1, \text{ or } \pm \sqrt{22}, \text{ or } \pm j\sqrt{23}.$$

$$\therefore x = 3, 2, \frac{1}{2}(2 \pm \sqrt{22}) \text{ or } \frac{1}{2}(-7 \pm j\sqrt{23});$$

$$y = 2, 3, \frac{1}{2}(2 \mp \sqrt{22}) \text{ or } \frac{1}{2}(-7 \mp j\sqrt{23}).$$

$$8. \quad x^2 + y = \frac{17}{8};$$

$$x + y^2 = \frac{5}{4}.$$

$$\therefore \frac{17}{8} - y = (\frac{5}{4} - y^2)^2,$$

$$\therefore y^4 - \frac{5}{2}y^2 + y + \frac{1}{2} = 0.$$

Testing this for rational linear factors it is easily reduced to

$$(y-1)^2(y^2 + 2y + \frac{1}{2}) = 0,$$

$$\therefore y = 1 \text{ or } \frac{1}{2}(-2 \pm \sqrt{2});$$

$$x = \frac{1}{4} \text{ or } \frac{1}{4}(-1 \pm 4\sqrt{2}).$$

$$9. \quad (2x - y + z)(x + y + z) = 9;$$

I.

$$(x + 2y - z)(x + y + z) = 1;$$

II.

$$(x + y - 2z)(x + y + z) = 4.$$

III.

Let  $s = x + y + z$  and the equations may be written

$$(s + x - 2y)s = 9$$

IV.

$$(s + y - 2z)s = 1$$

V.

$$(s - 3z)s = 4.$$

VI.

$$IV. + 3.V. \quad (4s + x + y - 6z)s = 12, \text{ or } (5s - 7z)s = 12$$

VII.

$$8.VII - 7.VI. \quad \{(15s - 21z) - (7s - 21z)\}s = 8,$$

$$\therefore 8s^2 = 8, \quad \therefore s = \pm 1.$$

Substituting in I, II. and III. they become

$$2x - y + z = \pm 9, \quad x + 2y - z = \pm 1, \quad x + y - 2z = \pm 4,$$

$$\therefore x = \pm 4, \quad y = \mp 2, \quad z = \mp 1.$$

$$10 \quad x^2 + y^2 = a;$$

$$u^2 + v = b;$$

$$xy + uv = c;$$

$$xu + yv = e.$$

Let  $t = xy - uv.$

$$\therefore (x+y)^2 = a+c+t, \quad \therefore x = \frac{1}{2}\{\sqrt{(a+c+t)} + \sqrt{(a+c-t)}\}$$

$$(x-y)^2 = a-c-t, \quad y = \frac{1}{2}\{\sqrt{(a+c+t)} - \sqrt{(a-c-t)}\}$$



$$(u+v)^2 = b+c-t, \quad u = \frac{1}{2} \{ \sqrt{(b+c-t)} + \sqrt{(b-c+t)} \}$$

$$(u-v)^2 = b-c+t, \quad v = \frac{1}{2} \{ \sqrt{(b+c-t)} - \sqrt{(b-c+t)} \}$$

Also  $2(xu+yv) = (x+y)(u+v) + (x-y)(u-v) = 2e,$

$$\therefore \sqrt{(a+c+t)(b+c-t)} + \sqrt{(a-c-t)(b-c+t)} = 2e,$$

$$\therefore \{ 4e^2 + (a-c-t)(b-c+t) - (a+c+t)(b+c-t) \}^2 = 16e^2(a-c-t)(b-c+t).$$

$$\therefore \{ (a-b)^2 + 4e^2 \} t^2 - 2(a^2 - b^2)ct + (a+b)^2 e^2 - 4e^2(ab+c^2) + 4e^4 = 0,$$

$$\therefore t = \frac{(a^2 - b^2)c \pm 2c \sqrt{[(ab - e^2) \{ (a-b)^2 - 4(c^2 - e^2) \} ]}}{(a-b)^2 + 4e^2}.$$

11.  $xy = uv$

$$x+y+u+v = a$$

$$x^3 + y^3 + u^3 + v^3 = b^3$$

$$x^5 + y^5 + u^5 + v^5 = c^5$$

Let  $x+y = \frac{1}{2}(a+z).$   $\therefore u+v = \frac{1}{2}(a-z).$

Also let  $r = xy = uv$

$$(x+y)^3 = x^3 + y^3 + 3xy(x+y)$$

$$(u+v)^3 = u^3 + v^3 + 3uv(u+v)$$

$$\therefore a(3z^2 + a^2) = 4(b^3 + 3ar)$$

Also  $(x+y)^5 = x^5 + y^5 + 5xy(x^3 + y^3) + 10x^2y^2(x+y)$

$$(u+v)^5 = u^5 + v^5 + 5uv(u^3 + v^3) + 10u^2v^2(u+v)$$

$$\therefore a(5z^4 + 10a^2z^2 + a^4) = 16\{c^5 + 5^3r + 10ar^2\}$$

Eliminating  $r$  between VII. and VIII,

$$45a^2z^4 - 30a(a^3 + 2b^3)z^2 + a^6 - 20a^3b^3 - 80b^6 + 144ac^5 = 0$$

$$\therefore 15az^2 - 5(a^3 + 2b^3) = \pm 2\sqrt{\{5(a^3 + 5b^3)^2 - 180ac^5\}}$$

$$\therefore z = \sqrt{\frac{a^3 + 2b^3 \pm 2\sqrt{[\frac{1}{5}\{(a^3 + 5b^3)^2 - 36ac^5\}]} }{3a}}$$

uced to

I.  
II.  
III.

IV.  
V.  
VI.  
VII.

±4,

+c-t}  
-c-t}

I.  
II.  
III  
IV.  
V.  
VI.  
VII.  
VIII.  
IX.  
X.

$$\text{VII. \& IX. } 12ar = a^3 - 4b^3 + 3az^2$$

$$= 2a^3 - 2b^3 \pm 2\sqrt{\frac{1}{5}\{(a^3 + 5b^3)^2 - 36ac^5\}}$$

$$\therefore r = \frac{5(a^3 - b^3) \pm \sqrt{5\{(a^3 + 5b^3)^2 - 180ac^5\}}}{30a}$$

XI.

X. and XI. give the values of  $z$  and  $r$  which may now be treated as known in  $\sqrt{\quad}$  and  $\sqrt{\quad}$ .

$$x + y = \frac{1}{2}(a + z), \text{ and } xy = r$$

$$\therefore x - y = \frac{1}{2}\sqrt{\{(a + z)^2 - 16r\}}$$

$$\therefore x = \frac{1}{4}(a + z \pm \sqrt{\{(a + z)^2 - 16r\}});$$

$$y = \frac{1}{4}(a + z \mp \sqrt{\{(a + z)^2 - 16r\}}).$$

The values of  $u$  and  $v$  may be obtained from those of  $x$  and  $y$  respectively by changing  $z$  into  $-z$ .

## EXERCISE.

1.  $6\{(7-x)^2 + y^2\} = 13(7-x)y, \quad x^2 + 4y = y^2 + 4.$
2.  $10x^2 - 9y^2 = 2x^3, \quad 8x^2 - 6y^2 = 13x.$
3.  $xy = (3-x)^2 = (2-y)^2. \quad 4. \quad x^2 + y^2 = 8x + 9y = 144.$
5.  $x^2 + y^2 = x + y + 12, \quad xy + 8 = 2(x + y).$
6.  $x + xy + y = 5, \quad x^2 + xy + y^2 = 7.$
7.  $x^3 + y^3 = 7xy = 28(x + y). \quad 8. \quad x^2 + xy + y^2 = \frac{35}{x^2 + y^2} = \frac{28}{xy}.$
9.  $x^4 + x^2y^2 + y^4 = 133, \quad x^3y + x^2y^2 + xy^3 = 114,$
10.  $(x + y)(x^2 + y^2) = 17xy, \quad (x - y)(x^2 - y^2) = 9xy.$
11.  $25(x^3 + y^3) = 7(x + y)^3 = 175xy.$
12.  $2x^2 - y^2 = 14(x^2 - 2y^2) = 14(x - y).$
13.  $2x^2 - 3xy = 9(x - 3y), \quad 3(x^2 - 3y^2) = 2(2x^2 - 3xy).$
14.  $2x^2 - xy + 5y^2 = 10(x + y), \quad x^2 + 4xy + 3y^2 = 14(x + y).$
15.  $(2x - 3y)(3x + 4y) = 39(x - 2y), \quad (3x + 2y)(4x - 3y) = (99(x - 2y))$

16.  $(x+2y)(x+3y) = 3(x+y)$ ,  $(2x+y)(3x+y) = 28(x+y)$ .
17.  $x+y=8$ ,  $x^4+y^4=706$ . 18.  $x+y=5$ ,  $x^5+y^5=275$ .
19.  $x+y=2$ ,  $13(x^5+y^5) = 121(x^3+y^3)$ .
20.  $x+y=4$ ,  $41(x^5+y^5) = 122(x^4+y^4)$ .
21.  $x^2-5xy+y^2+5=0$ ,  $xy=x+y-1$ .
22.  $x^2+y=5(x-y)$ ,  $x+y^2=2(x-y)$ .
23.  $3(x^2+y) = 3(x+y^2) = 13xy$ .
24.  $10(x^2+y) = 10(x+y^2) = 13(x^2+y^2)$ .
25.  $x^2+y = \frac{1}{9}$ ,  $x+y^2 = \frac{2}{9}$ . 26.  $9(x^2+y) = 3(x+y^2) = 7$ .
27.  $x+xy+y=5$ ,  $x^3+xy+y^3=17$ .
28.  $x+y=2$ ,  $(x+1)^5 + (y-2)^5 = 211$ .
29.  $3(x-1)(y+1) = 4(x+1)(y-1)$ ,  $\frac{x^2+x+1}{y^2+y+1} = \frac{31}{39} \left( \frac{x^2-x+1}{y^2-y+1} \right)$
30.  $x+y = \frac{1}{xy}$ ,  $x-y = xy$ .
31.  $x+y+1=0$ ,  $x^6+y^6+2=0$ .
32.  $x+y=1$ ,  $3(x^8+y^8) = 7$ .
33.  $4xy^2 = 5(5-x)$ ,  $2(x^2+y^2) = 5$ .
34.  $27xy = 17$ ,  $9(x^3+y^3) = -8$ .
35.  $(x^2+y^2)^2 + 4x^2y^2 = 5 - 12y$ ,  $y(x^2+y^2) + 3 = 0$ .
36.  $x+y=xy$ ,  $x^2+y^2 = x^3+y^3$ .
37.  $x^4 - 6x^2 \sqrt{(y^2-x^2)^{\frac{1}{2}}} - 16y^2 = 9x^2$ ,  
 $(x^2+2)^2 = 4\{2+x^2\sqrt{(y^2+x^2)-y^2}\}$ .
38.  $x(y^2+3y-1) = 2y^2+2y+3$ ,  $y(x^2+3x-1) = 2x^2+2x+3$ .

XI.

be treated

f x and y

= 144.

 $\frac{28}{y^2} = \frac{xy}{xy}$ 

-y).

99(x-2y)

$$39. \frac{x^3}{a^3} + \frac{y^3}{b^3} = 2c^3, \quad \frac{x}{a} + \frac{y}{b} = c \left( \frac{x}{a} - \frac{y}{b} \right).$$

$$40. x^3 + xy^2 = a, \quad y^3 + x^2y = b.$$

$$41. x + y = a, \quad \frac{x}{b-y} + \frac{b-y}{x} = c.$$

$$42. x^2 + ay^2 = \frac{a+1}{a-1}, \quad ax^2 + y^2 = (a^2 - 1)y.$$

$$43. x + y^2 = ax, \quad x^3 + y = by. \quad 44. x + y^3 = ay^2, \quad x^2 + y = bx^2.$$

$$45. x^4 - y^4 = a^2(x-y)^2, \quad x^3 - x^2y + xy^2 - y^3 = b^2(x+y).$$

$$46. (x+y)(x^2 + 3y^2) = m, \quad (x-y)(x^2 + 3y^2) = n.$$

$$47. x^2y^2 = y(a-x)^3 = x(b-y)^3.$$

$$48. x^3(b-y) = y^3(a-x) = (a-x)^2(b-y)^2.$$

$$49. a^2(x^2 + e^2) = b^2(x+y)^2, \quad a^2(y^2 + 3z^2) = c^2(x+y)^2.$$

$$50. x^3 - y^3 = a(x^2 - y^2), \quad x^2 + y^2 = b(x+y).$$

$$51. x + y = a, \quad x^3 + y^3 = bxy.$$

$$52. \sqrt{\frac{x}{y}} - \sqrt{\frac{y}{x}} = \frac{x-y}{a}, \quad \frac{x(c^2 + xy)}{y(c^2 - xy)} = b^2.$$

$$53. x + y = xy = x^2 + y^2. \quad 54. x - y = \frac{x}{y} = x^2 - y^2.$$

$$55. x^3(1+y^2)(1+y^4) = a, \quad x^3(1-y^2)(1-y^4) = b.$$

$$56. \frac{x^2 + xy + y^2}{x^2 - xy + y^2} = \frac{x^2 + y^2}{a} = \frac{xy}{b}.$$

$$57. x^2y + xy^2 = \frac{a}{x^2 + y^2}, \quad x^4y + xy^4 = b.$$

$$58. x^2y + xy^2 = a(x^2 + y^2), \quad x^2y - xy^2 = b(x^2 - y^2).$$

$$59. \left( \frac{x}{y} + \frac{y}{x} \right) (x+y) = a, \quad \frac{x^2}{y} + \frac{y^2}{x} = b.$$

60. x

61. a

62. x

63. (

64. x

65. a

66. x

67. x

68. :

69. (

70. :

71. :

73. :

74. :

75. :

76. :

77. :

78. :

79. :

80. :

81. :

82. :

60.  $x^2 + y^2 = ax^2y^2 = xy(x+y)$ .
61.  $abxy = a(x^3 + y^3) = b(x+y)^3$ .
62.  $xy(x+y) = a, \quad x^3y^3(x^3 + y^3) = b$ .
63.  $\left(\frac{1}{x} + \frac{1}{y}\right)(x^3 - y^3) = a, \quad \left(\frac{1}{x} - \frac{1}{y}\right)(x^3 + y^3) = b$ .
64.  $x^4 + y^4 = m(x^2 + y^2), \quad x^2 + xy + y^2 = n$ .
65.  $ab(x+y) = xy(a+b), \quad x^2 + y^2 = a^2 + b^2$ .
66.  $x^3 + y^3 = a(x+y), \quad x^4 + y^4 = b(x+y)^4$ .
67.  $x^2 + y^2 = a, \quad x^5 + y^5 = b(x^3 + y^3)$ .
68.  $xy = a, \quad x^5 + y^5 = b(x^3 + y^3)$ .
69.  $(x-y)(x^3 + y^3) = (a-b)(a^3 + b^3), \quad x^2 - y^2 = a^2 - b^2$ .
70.  $x^2 - y^2 = a, \quad x^3 + y^3 = b(x-y)$ .
71.  $x+y = a, \quad x^4 + y^4 = b$ .
72.  $x+y = a, \quad x^5 + y^5 = b$ .
73.  $x+y = a, \quad x^2 + y^2 = b^2x^2y^2$ .
74.  $x+y = a+b, \quad (a-b)^2(x^4 + y^4) = (x-y)^2(a^4 + b^4)$ .
75.  $x+y = a, \quad c(x^4 + y^4) = xy(x^3 + y^3)$ .
76.  $(x+y)^3 = a(x^2 + y^2), \quad xy = c(x+y)$ .
77.  $x^2y + xy^2 = a^3, \quad c^3(x^3 + y^3) = x^3y^3$ .
78.  $x^3 = a(x^2 + y^2) - cxy, \quad y^3 = c(x^2 + y^2) - axy$ .
79.  $x^3 - y^2 = a^2, \quad x^3 - y^3 = c^4 \left(\frac{1}{x} - \frac{1}{y}\right)$ .
80.  $x^4 - y^4 = a^2xy, \quad (x^2 + y^2)^2 = b^2(x^2 - y^2)$ .
81.  $(x+y)x^2y^2 = a, \quad x^5 + y^5 = b$ .
82.  $(x+y)xy = a, \quad x^5 + y^5 = bxy$ .

83.  $x^4 + y^4 = a(x+y)^2$ ,  $x^5 + y^5 = b(x+y)^3$ . 100. (
84.  $x^4 + x^2y^2 + y^4 = a$ ,  $x^2 - xy + y^2 = 1$ . 101. (
85.  $(x^2 + y^2)xy = x^2 - y^2$ ,  $\frac{x^4(1 + x^2y^2)}{y^4(1 + xy)^2} = \frac{a}{b} \cdot \frac{1 + xy}{1 - xy}$ . 102. (
86.  $x + y = (x - y) \sqrt{(xy)}$ ,  $\frac{\sqrt{(x^7y^5)} - x}{\sqrt{(x^5y^7)} + y} = a$ . 103. (
87.  $\frac{1}{x} + \frac{1}{y} = \frac{1}{a}$ ,  $\sqrt{(1-x)} - \sqrt{(1-y)} = b$ . 104.
88.  $x^2 + y^2 = a(x+y)$ ,  $x^4 + y^4 = b(x^3 + y^3)$ . 105.
89.  $x^3 + y^3 = a$ ,  $(x+y)(x^4 + y^4) = b(x^2 + y^2)$ . 106.
90.  $(x^2 + y^2)(x^3 + y^3) = axy$ ,  $(x+y)(x^4 + y^4) = bxy$ . 107.
91.  $(x+y)^2(x^2 + y^2) = a$ ,  $(x^2 + y^2)^2(x^4 + y^4) = b$ . 108.
92.  $(x-y)(x^2 - y^2)(x^4 - y^4) = 4axy$ ,  
 $(x+y)(x^2 + y^2)(x^3 + y^3) = b(x-y)$ .
93.  $x^4y + xy^4 = a(x^3y + xy^3) = b(x^4 + y^4)$ . 109.
94.  $a(x^5 + y^5) = ab(x+y) = bxy(x^3 + y^3)$ . 110.
95.  $\frac{x^3 - y^3}{x^2 - y^2} = \frac{a^3 - b^3}{a^2 - b^2}$ ,  $\frac{x^5 - y^5}{x^4 - y^4} = \frac{a^5 - b^5}{a^4 - b^4}$ . 111.
96.  $\frac{x^3 + y^3}{x^2 - y^2} = \frac{a^3 + b^3}{a^2 - b^2}$ ,  $\frac{x^4 + y^4}{x^3 - y^3} = \frac{a^4 + b^4}{a^3 - b^3}$ . 112.
97.  $x^5 = 2ax - by$ ,  $y^5 = 2ay - bx$ . 113.
98.  $(x+y)(x^3 + y^3) = a$ ,  $(x-y)(x^3 - y^3) = b$ . 114.
99.  $\frac{(x+y)^4(x^2 + xy + y^2)}{(x^2 + y^2)(x^2 - xy + y^2)} = 3m^2$ ,  
 $\frac{(x-y)^4(x^2 - xy + y^2)}{(x^2 + y^2)(x^2 + xy + y^2)} = 3n^2$ . 115.

$$100. (x+y)(x^3+y^3) = axy, \quad (x-y)(x^3-y^3) = bxy.$$

$$101. (x+y)(x^3+y^3) = a(x^2+y^2), \quad (x-y)(x^3-y^3) = b(x^2+y^2).$$

$$102. \frac{(x+y)^3(x^3+y^3)}{(x^2+xy+y^2)(x^2+y^2)} = a^2,$$

$$\frac{(x-y)^3(x^3-y^3)}{(x^2-xy+y^2)(x^2+y^2)} = b^2.$$

$$103. \frac{(x^3+y^3)(x+y)^3}{x^2+xy+y^2} = 2a^2, \quad \frac{(x^3-y^3)(x-y)^3}{x^2-xy+y^2} = 2b^2.$$

$$104. \frac{(x^3+y^3)(x+y)^5}{(x^2+xy+y^2)^2} = 8a^2, \quad \frac{(x^3-y^3)(x-y)^5}{(x^2-xy+y^2)^2} = 8b^2.$$

$$105. xy(x+y)(x^3+y^3) = a, \quad xy(x-y)(x^3-y^3) = b.$$

$$106. x(x+y)(x+2y)(x+3y) = a^2, \quad (x+y)^2 + (x+2y)^2 = b.$$

$$107. \sqrt{(x-xy)} + \sqrt{(y-xy)} = a, \quad \sqrt{(x-x^2)} + \sqrt{(y-y^2)} = b.$$

$$108. (x+1)(y-1) = a(x-1)(y+1),$$

$$(x^5+1)(y-1)^5 = b^3(x-1)^5(y^5+1).$$

$$109. x+y = a, \quad \sqrt[4]{(x-k)} + \sqrt[4]{(y-k)} = c.$$

$$110. x+y = a(1+xy), \quad (x+y)^4 = b^4(1+x^4y^4).$$

$$111. x+y = a(1+xy), \quad x^5+y^5 = b^3(1+x^5y^5).$$

$$112. (x+1)(y-1) = a(x-1)(y+1),$$

$$(x^5-1)(y-1) = b^2(y^5-1)(x-1).$$

$$113. \frac{(1+x)(1+y)}{(1-x)(1-y)} = a, \quad \frac{(1+x)^3(1+y)^3}{(1-x^3)(1-y^3)} = b.$$

$$114. \frac{(c+x)(c+y)}{(c-x)(c-y)} = a, \quad \frac{(c^4+x^4)(c^4+y^4)}{(c^4-x^4)(c^4-y^4)} = b.$$

$$115. \frac{(x+m)(y+n)}{(x-m)(y-n)} = a, \quad \frac{(x^4+m^4)(y^4+n^4)}{(x-m)^4(y-n)^4} = b.$$

116.  $\frac{(x+1)(y+1)}{(x-1)(y-1)} = \frac{a}{c}, \quad \frac{(x^5+1)(y^5+1)}{(x^5-1)(y^5-1)} = \frac{b}{d}$
117.  $\frac{y(1+x^2)}{x(1+y^2)} = a, \quad \frac{y^2(1+x^4)}{x^2(1+y^4)} = b.$
118.  $\frac{1+x}{1+y} = a\sqrt{\frac{x}{y}}, \quad \frac{y(1+x+x^2)}{x(1+y+y^2)} = b.$
119.  $\frac{y(1+x^2)}{x(1+y^2)} = a, \quad \frac{y^3(1+x^4)}{x^3(1+y^6)} = b.$
120.  $\frac{y(1+x^2)}{x(1+y^2)} = a, \quad \frac{y^4(1+x^8)}{x^4(1+y^8)} = b.$
121.  $\frac{y(1+x^2)}{x(1+y^2)} = a, \quad \frac{y^5(1+x^{10})}{x^5(1+y^{10})} = b.$
122.  $\frac{(x+y)(xy+1)}{(x-y)(xy-1)} = \frac{a^2+b^2}{2ab}, \quad \frac{x(y^2+1)}{y(x^2-1)} = \frac{a-b}{a+b}$
123.  $\frac{(x+y)(1+xy)}{(x-y)(1-xy)} = a^2(b^2-1), \quad \frac{x(1-y^2)}{y(1-x^2)} = b,$
124.  $\frac{(x+y)(1+xy)}{(x-y)(1-xy)} = a, \quad \frac{(x^2+y^2)(1+x^2y^2)}{(x^2-y^2)(1-x^2y^2)} = b.$
125.  $\frac{(x+y)(1+xy)}{(x-y)(1-xy)} = a, \quad \frac{(x^3+y^3)(1+x^3y^3)}{(x^3-y^3)(1-x^3y^3)} = b.$
126.  $\frac{(x^4+y^4)(1+x^4y^4)}{(x^4-y^4)(1-x^4y^4)} = a, \quad \frac{(x+y)(1+xy)}{(x-y)(1-xy)} = b.$
127.  $\frac{(x+y)(1+xy)}{(x-y)(1-xy)} = a, \quad \frac{(x^5+y^5)(1+x^5y^5)}{(x^5-y^5)(1-x^5y^5)} = b.$
128.  $\frac{(x^2+xy+y^2)(1+xy+x^2y^2)}{(x+y)^2(1+xy)^2} = a.$   
 $\frac{(x^2-xy+y^2)(1-xy+x^2y^2)}{(x-y)^2(1-xy)^2} = b.$
129.  $x^4 - 3x^2y + 5a^3x + y^2 = 0, \quad y^3 - x^2y^2 - 2a^3x = 0.$

130.

(Her

131.

132.

1.

3.

4.

6.

8.

10.

12.



$$130. \quad 2x(y^2 - 2x)^2 = a, \quad y(y^2 - 2x)^2 \sqrt{(y^2 - 4x)} = b.$$

(Hence deduce the solution of  $x^5 - 5x^2 + 2 = 0$ ).

$$131. \quad 2xy(x^2 + y^2)^2 = a, \quad (x^2 - y^2)(x^2 + y^2)^2 = b.$$

$$132. \quad \sqrt{(x^2 + y^2)} + \sqrt{(a - x)^2 + y^2} = \sqrt{\left\{\frac{1}{2}a\sqrt{3} - y\right\}^2 + \left(\frac{1}{2}a - x\right)^2},$$

$$6(x^2 - y^2) = a(6x - 2y\sqrt{3} + a).$$

## EXERCISE.

- |  |  |
|--|--|
| 1. $(2x + y - 4z)(x + y + z) = 24,$                    | 2. $x^2 - yz = 1,$                           |
| $(x + 2y - 2z)(x + y + z) = 6,$                        | $y^2 - xz = 2,$                              |
| $(-2x + 3y + 5z)(x + y + z) = 30,$                     | $z^2 - xy = 3.$                              |
| 3. $(x + 2y - 3z)(x + y + z) - 2(xy + yz + zx) = -12,$ |  |
| $(2x - 3y + z)(x + y + z) + (xy + yz + zx) = 61,$      |  |
| $(3x - y + 2z)(x + y + z) - 5(xy + yz + zx) = 5.$      |  |
| 4. $x^2 - yz = 0,$                                     | 5. $(x^5 + y^5 + z^5)^3 + (x + y)^2 = 31.$   |
| $x + y + z = 7,$                                       | $(x^5 + y^5 + z^5)^3 + (x + y + z)^3 = 729.$ |
| $x^2 + y^2 + z^2 = 21.$                                | $(x + y)^2 + (x + y + z)^3 = 31.$            |
| 6. $x^2 - yz = 0,$                                     | 7. $x + yz = 14,$                            |
| $x + y + z = 21,$                                      | $y + zx = 11,$                               |
| $(x - y)^2 + (y - z)^2 + (z - x)^2 = 126.$             | $z + xy = 10,$                               |
| 8. $x + y = 8z,$                                       | 9. $x + y = 5z,$                             |
| $x^3 + y^3 = 134z^3,$                                  | $x^2 + y^2 = 39z,$                           |
| $x^2 + y^2 + z^2 = 134.$                               | $x^3 + y^3 = 105z^2.$                        |
| 10. $x + y = 7z,$                                      | 11. $x + y = 7z,$                            |
| $x^2 + y^2 = 25z^2,$                                   | $x^3 + y^3 = 25z^2,$                         |
| $x^4 + y^4 = 674z^3.$                                  | $x^5 + y^5 = 20272z.$                        |
| 12. $x + y : y + z : z + x :: a : b : c,$              | 13. $x + y : y + z : z + x :: a : b : c,$    |
| $(a + b + c)xyz = 2,$                                  | $(a + b + c)xyz = 2(x + y + z.)$             |

14.  $ax = by = cz = \frac{1}{x} + \frac{1}{y} + \frac{1}{z}$       16.  $z \left( \frac{x}{y} + \frac{y}{x} \right) = a,$
15.  $(x+y-z)x = a,$   
 $(x-y+z)y = b,$   
 $(-x+y+z)z = c.$        $y \left( \frac{x}{z} + \frac{z}{x} \right) = b,$   
 $x \left( \frac{y}{z} + \frac{z}{y} \right) = c.$
17.  $(y+z)(2x+y+z) = a,$       18.  $x(y+z):y(z+x):z(x+y) =$   
 $(z+x)(x+2y+z) = b,$        $b+c:c+a:a+b,$   
 $(x+y)(x+y+2z) = c,$        $xy+yz+zx = (a+b+c)(x+y+z).$
19.  $(a+b)x + (b+c)y + (c+a)z = (a+b+c)(x+y+z),$   
 $a(x+y) = c(y+z),$   
 $(x+y)^2 + (y+z)^2 + (z+x)^2 = 4(a^2 + b^2 + c^2).$
20.  $c(x+y) + b(x-z) - a(y+z) = 0,$   
 $b(x-z) = (a-c)y,$   
 $x^2 + y^2 + z^2 = a^2 + b^2 + c^2.$
21.  $x+y-az = x-by+z = -cx+y+z = xyz.$
22.  $(a+b+c)(x-y) + a(x+z) - b(y+z) = 0,$   
 $(a+b+c)(x-z) + a(x+y) - c(y+z) = 0,$   
 $\frac{ax^2}{(b+c)^2} + \frac{by^2}{(c+a)^2} + \frac{cz^2}{(a+b)^2} = 1.$
23.  $xy + \frac{x}{z} = a; \quad yz + \frac{y}{x} = b, \quad zx + \frac{z}{y} = c.$
24.  $y+z:z+x:x+y :: b+c:c+a:a+b,$   
 $(x+y+z)(xyz) = (a+b+c)(xy+yz+zx).$
25.  $n^2 - yz = a, \quad y^2 - xz = b, \quad z^2 - xy = c.$
26.  $x^2 + (y-z)^2 = a^2, \quad y^2 + (z-x)^2 = b^2, \quad z^2 + (x-y)^2 = c^2.$

$$27. \quad x^2 + xy + y^2 = a^2, \quad y^2 + yz + z^2 = b^2, \quad z^2 + zx + x^2 = c^2.$$

$$28. \quad x^3 + y^3 - z^3 + 3xyz = a(x + y - z) \\ x^3 - y^3 + z^3 + 3xyz = b(x - y + z), \\ -x^3 + y^3 + z^3 + 3xyz = c(-x + y + z).$$

$$29. \quad x + y + 2az = 0, \\ x^2 + y^2 - 2b^2z^2 = 0, \\ x^n + y^n + z^n = c^n.$$

$$30. \quad x + y - az = 0, \\ y(x^2 + y^2) = b^2, \\ x^3 + y^3 = c^3.$$

$$31. \quad x(y-1)(z-1) = 2a, \\ x^2(y^2-1)(z^2-1) = 4bx, \\ x^3(y^3-1)(z^3-1) = 6cx^2$$

$$32. \quad x(y-1) = a(z-1), \\ x^2(y^2-1) = b^2(z^2-1), \\ x^3(y^3-1) = c^3(z^3-1).$$

$$33. \quad x(y-1) = a(z-1), \\ x^2(y^2-1) = b^2(z^2-1), \\ x^4(y^4-1) = c^4(z^4-1).$$

$$34. \quad x(y-1) = a(z-1), \\ x^2(y^2-1) = b^2(z^2-1), \\ x^4(y^4+1) = c^4(z^4+1).$$

$$35. \quad x(y-1) = a(z-1), \\ x^2(y^2-1) = b^2(z^2-1), \\ x^5(y^5-1) = c^5(z^5-1).$$

$$36. \quad (x-y)^2 = az(x+y), \\ (x^3-y^3) = bz(x+y)^3, \\ (x-y)^3 = cz(x^3+y^3)$$

$$37. \quad x - y = a, \\ u - v = b, \\ xy = uv, \\ x^5 - y^5 + u^5 - v^5 = c(a + b).$$

$$38. \quad x + y = a, \\ u + v = b, \\ x^2 + u^2 = c^2, \\ y^2 + v^2 = c^2.$$

$$39. \quad xy = uv = a^2, \\ x + y + u + v = b, \\ x^3 + y^3 + u^3 + v^3 = c^3.$$

$$40. \quad xy = uv = a^2, \\ x + y + u + v = b, \\ x^4 + y^4 + u^4 + v^4 = c^4.$$

$$41. \quad xy = uv = a^2,$$

$$42. \quad xy = uv = a^2,$$

- $x + y + u + v = b,$   
 $x^5 + y^5 + u^5 + v^5 = c^5.$
43.  $xy = uv = a^2, \quad x + y + u + v = b, \quad (x + u)^4 + (y + v)^4 = c^4.$
44.  $xy = uv,$   
 $x + y + u + v = a,$   
 $x^2 + y^2 + u^2 + v^2 = b^2,$   
 $x^3 + y^3 + u^3 + v^3 = c^3.$
45.  $xy = uv,$   
 $x + y + u + v = a,$   
 $x^2 + y^2 + u^2 + v^2 = b^2,$   
 $x^4 + y^4 + u^4 + v^4 = c^4.$
46.  $xy = uv,$   
 $x + y + u + v = a,$   
 $x^2 + y^2 + u^2 + v^2 = b^2,$   
 $x^5 + y^5 + u^5 + v^5 = c^5.$
47.  $xy = uv,$   
 $x + y + u + v = a,$   
 $x^3 + y^3 + u^3 + v^3 = b^3,$   
 $x^4 + y^4 + u^4 + v^4 = c^4.$
48.  $xy - uv = 0,$   
 $xu + yv = a^2,$   
 $x + y + u + v = b,$   
 $x^3 + y^3 + u^3 + v^3 = c^3.$
49.  $x^2 + y^2 = a^2,$   
 $u^2 + v^2 = m^2,$   
 $ux + vy = c^2,$   
 $vx + uy = n^2.$
50.  $x + y + u + v = a,$   
 $xy + uv = b^2,$   
 $x^2 + y^2 = m^2,$   
 $u^2 + v^2 = n^2.$
51.  $y(1 + x^2) = 2x,$   
 $u(1 + y^2) = 2y,$   
 $v(1 + u^2) = 2u,$   
 $x(1 + v^2) = 2v.$
52.  $x + y + u + v = a, \quad (x + y)^2 + (u + v)^2 = b^2,$   
 $(x + u)^2 + (y + v)^2 = c^2, \quad (x + v)^2 + (y + u)^2 = e^2.$
53.  $\frac{x}{y + z} = \frac{2a - u}{a - 2u}, \quad \frac{y}{z + x} = \frac{2b - u}{b - 2u}, \quad \frac{z}{x + y} = \frac{2c - u}{c - 2u},$   
 $x^2 + y^2 + z^2 = e^2.$

1.  
8. —  
7. —  
7. 14

1.  
5. 96  
10. —

1. C  
5. 0.  
11. c  
14. C  
17. C  
25. 9  
85.

1.  
3.  
4.  
6.  
8.  
10.

# ANSWERS.

## EXERCISE i.

1. 9, -69, 1, 0, 1206, -29,  $1\frac{3}{8}$ .      2. -160, 106, 41, 108.  
3.  $-\frac{5}{19}$ ,  $\frac{13}{8}$ , -25, 125,  $\frac{7}{19}$ , -81,  $-4\frac{1}{8}$ , 0, -1.      4. 9, 8,  
7,  $-\frac{1}{2}$ .      5. 176, 82,  $254\frac{2}{9}$ ,  $-37 \div 7\frac{2}{3}$ .      6. 18 each.  
7. 146, 14, -72, -270, 396.      8. Each=0.

## EXERCISE ii.

1. -1.      2. -166542.      3. 100.      4. -2967511.  
5. 968.      6. -162.      7. 10.      8. -8.      9. 0.  
10. -20.      11. 706440254900.      12. 0 each.      13. Each 0.

## EXERCISE iii.

1. 0,  $16a^4$ .      2.  $a$ ,  $a\sqrt{3}$ .      3.  $2a$ , 0.      4.  $26a^6$ ,  $-26a^6$ .  
5. 0.      6.  $4a^4$ .      7.  $6a^4$ .      8.  $\frac{3}{2}$ .      9.  $c$ .      10. 0.  
11.  $a \div (a+b)$ .      12.  $a^2c(b+2c) \div b^2$ .      13.  $a^2+b^2+c^2$ .  
14. 0.      15.  $(12a^2b - 24ab^2 + 28b^3) \div (3b-a)^3$ .      16. 0.  
17. 0.      18.  $-b^2c$ .      19, 20, 21 and 22, each 0.  
25.  $2(b+l)h$ ,  $4x^2$ .      32.  $d^2 = 3l^2$ .      33.  $l = \sqrt{(\frac{1}{3}d^2)}$ .  
35.  $\pi r^2$ ,  $\pi(r+r')(r-r')$ .

## EXERCISE iv.

1.  $2(bx+cy)$ .      2.  $3(ax-by)$ .  
3.  $a^2(x-z) - ab(x-y) - b^2(y-z)$ .  
4.  $(x+y+z)(a+b+c)$ .      5.  $(a+b+c)(x^2+y^2+z^2)$ .  
6.  $2(x+y+z) \times (a^2+b^2+c^2 - ab - bc - ca)$ .      7. 0.  
8.  $2(rx+by+cz)$ .      9.  $a^2+b^2+c^2$ .  
10.  $2x^2(a-2b)$ .      11.  $a+b-c$ .

## EXERCISE v.

1.  $2(x^2 + 9y^4)$ ,  $4a^2b^2$ .
2.  $x^2 + 4x$ ,  $-3\frac{1}{4}x^4 - 4x^2y^2 + 3\frac{1}{4}y^4$ .
3.  $x^3 - 6x^3 + 9x^4 + 2xy - 6xy^2 - 6x^2y + 18x^2y^2 + y^2 - 6y^3 + 9y^4$ .
4.  $4(x^2 - b^2)^2$ .
5.  $6a^3$ .
6.  $4x^2y(x^2 - y^2)$ ,  $2(1 + 12x^3 + 16x^4)$ .
7.  $\frac{1}{16}c^3$ .
8.  $a^2 - 2b^2$ ,  $8ab(a + b)^2$ .
9.  $2(a - c)(b - d)$ .
10.  $\frac{1}{4}x^2 + \frac{1}{4}y^2 + \frac{1}{4}z^2 + \frac{1}{4}(xy + yz + zx)$ .
11.  $(1 + x^2)^2$ .
12.  $4(xy + yz + zx) - 2(x^2 + y^2 + z^2)$ .
13.  $x^2$ .
14.  $(a^2 + 2b^2 - 2c^2)^2$ .
15.  $16x^2y^3$ .
16.  $-4ab$ .
17.  $4(a + b + c)^2$ .
18.  $4(1 + x^2 + x^4 + x^6)$ .
19.  $(a^2x^2 + b^2y^2)^2$ .

## EXERCISE vi.

1.  $1 - 4x + 10x^2 - 20x^3 + 25x^4 - 24x^5 + 16x^6$ ,  
 $1 - 2x + 3x^2 - 4x^3 + 3x^4 - 2x^5 + x^6$ .
2.  $1 - 4x + 8x^2 - 14x^3 + 14x^4 - 8x^5 + 5x^6 + 6x^7 + x^8$ ,  
 $1 + 6x + 15x^2 + 20x^3 + 15x^4 + 6x^5 + x^6$ .
3.  $4a^2 + b^2 + c^4 + 1 - 4ab - 4ac^2 - 4a + 2bc^2 + 2b + 2c^2$ ,  
 $1 + x^2 + y^2 + z^2 - 2x + 2y + 2z - 2xy - 2xz + 2yz$ ,  
 $\frac{1}{4}x^2 + \frac{1}{9}y^2 + 36z^2 - \frac{1}{3}xy + 6xz - 4yz$ .
4.  $x^6 - 2x^5y + 3x^4y^2 - 4x^3y^3 + 3x^2y^4 - 2xy^5 + y^6$ ,  
 $a^2x^2 + 2abx^3 + (2ac + b^2)x^4 + 2(ad + bc)x^5 + (2bd + c^2)x^6 + 2cdx^7 + d^2x^8$ .
5.  $3(a^2 + b^2 + c^2) - 2(ab + bc + ca)$ .
6.  $4a^2 + \frac{1}{4}b^2x^2 + \frac{1}{16}c^2x^2 + 4d^2x^2 - 2abx - acx + 3adx + \frac{1}{4}bcx^2 - 2bdx^2 - cdx^2$ .

## EXERCISE vii.

1.  $(a^2 - b^2)^2$ .
2.  $\frac{1}{2}x^4 + y^4$ .
3.  $a^4 + 3a^2b^2 + 4b^4$ .
4.  $x^4 - y^4$ .
5.  $x^2$ .
6.  $16x^2$ .
7.  $0$ .
8.  $4a^4 - 9b^4 - 16c^4 + 24b^2c^2$ .
9.  $b^2 - 9c^2 - 4a^2 + 12ac$ ,  
 $9c^2 - 4a^2 - b^2 + 4ab$ .
10.  $x^8 - y^8$ .
11.  $x^8 + x^4y^4 + y^8$ .

12.  $a^2 - a^2b^2 + b^2 - 1$ . 14.  $x^4 + y^4 + \frac{7}{16}x^2y^2$ .  
 15.  $x^8 + 2x^6 + 3x^4 + 2x^2 + 1$ .  
 16.  $4a^4x^2 - 4a^4xy + a^4y^2 - a^2x^4 - 2a^2y^3y + 2ax^5 + 2ax^4y - x^6$ .  
 20.  $(x^2 + y^2 - 2xy - z^2)^2$ . 21.  $x^8 - y^8$ .  
 22.  $\frac{1}{3} - 6a^2 + 27a^4$ . 23.  $(m+p)^2 - (n+q)^2$ .  
 24.  $2x^2 + x^4 + 2x^6 - x^8 - 1$ . 25.  $a^8 - b^{16}$ .

## EXERCISE viii.

1.  $x^4 + 4x^3 + 3x^2 - 2x - 12$ ,  $x^2 + y^2 - 2xy + 8xz - 8yz + 15z^2$ .  
 2.  $x^4 + 12x^3 + 49x^2 + 78x + 40$ ,  $x^6 + bx^3 - a^2 + 3ab - 2b^2$ .  
 3.  $a^8 + 8a^6 - 10a^4 - 104a^2 + 105$ ,  $x^8 + 2x^6 - x^2 - 2$ .  
 4.  $x^4 + 5x^3y^2 - 12x^2y^2 + 5xy^3 + y^4$ .  
 5.  $x^{2n} - 2x^n - a^2 - 16a - 63$ ,  $\frac{x^2}{y^2} + \frac{y^2}{x^2} + \frac{2x}{y} + \frac{2y}{x} - 1$ .  
 6.  $n^2x^3 + 2nxy + y^2 + 10nx + 10y + 21$ .  
 7.  $(x+a)^2 + 2y(x+a) - 3y^2$ . 8.  $x^{4n} + 2x^{3n} + x^{2n}(1-a-b) - x^n(a+b) + ab$ . 9.  $\frac{1}{4}x^8 - x^4y^2 + y^4 - x^4 + 2y^2 - 8$ .  
 10.  $\left(\frac{1}{x} + \frac{1}{y}\right)^2 = 2\left(\frac{1}{x} + \frac{1}{y}\right) - \frac{5}{4}$ .  
 11.  $x^4 - 8x^3 + 19x^2 - 12x + 2$ .  
 12.  $(x+b)^4 - (a^2+c^2)(x+b)^2 + a^2c^2$ . 13.  $ab + cd$ .

## EXERCISE ix.

1.  $2(1+3x^4)$ ,  $2xy^3(3x^4+x^2y^0)$ . 2.  $9b(a^2+b^2+ab^2)$ ,  
 $b(27a^2-27ab+7b^2)$ . 3.  $(x+y)^3$ . 4.  $8a^3$ . 5.  $8x^3$ .  
 6.  $8x^3$ . 7.  $a^3$ . 8.  $27x^3$ . 9.  $(2+x)^3$ . 12.  $8(x^2+y^2)^3$ .  
 14.  $(a^3+b^3)(x^3+y^3)$ . 15. 0. 16. 0.

## EXERCISE x.

1.  $1 - 3x + 6x^2 - 7x^3 + 6x^4 - 3x^5 + x^6$ ,  $a^3 - b^3 - c^3 - 3a^2(b+c)$   
 $+ 3b^2(a-c) + 3c^2(a-b) + 6abc$ ,  $1 - 6x + 21x^2 - 56x^3 +$   
 $111x^4 - 174x^5 + 219x^6 - 204x^7 + 144x^8 - 64x^9$ .

2.  $-(x^9 + 18x^8 + 27x^7 + 29x^6 - 24x^5 - 36x^4 + 5x^3 - 3x^2 - 2)$ .  
 5. 0. 6.  $45x^6 + 168\Sigma x^4 y^2 - 432x^2 y^2 z^2$ . 7.  $(ax + by + cz)^3$ .

## EXERCISE xi.

1.  $x^6 + 6x^5 y + 15x^4 y^2 + 20x^3 y^3 + 15x^2 y^4 + 6xy^5 + y^6$ ,  
 $x^7 + 7x^6 y + 21x^5 y^2 + 35x^4 y^3 + 35x^3 y^4 + 21x^2 y^5 + 7xy^6 + y^7$ ,  
 $x^8 + 8x^7 y + 28x^6 y^2 + 56x^5 y^3 + 70x^4 y^4 + 56x^3 y^5 + 28x^2 y^6 +$   
 $8xy^7 + y^8$ ,  $x^{12} + 12x^{11} y + 66x^{10} y^2 + 220x^9 y^3 + 495x^8 y^4 +$   
 $792x^7 y^5 + 924x^6 y^6 + 792x^5 y^7 + \&c$ .  
 2. The signs will be alternately positive and negative.  
 3.  $a^5 - 5a^4 b + 10a^3 b^2 - 10a^2 b^3 + 5ab^4 - b^5$ ,  
 $a^4 - 8a^3 b + 24a^2 b^2 - 32ab^3 + 16b^4$ , same as last, terms in  
 inverse order. 4.  $1 + 6m + 15m^2 + 20m^3 + 15m^4 + 6m^5 +$   
 $m^6$ ,  $m^5 + 5m^4 + 10m^3 + 10m^2 + 5m + 1$ ,  $64m^5 + 192m^5 +$   
 $240m^4 + 160m^3 + 60m^2 + 12m + 1$ . 5. 120.  
 6.  $x^8 - 4x^6 y + 6x^4 y^2 - 4x^2 y^3 + y^4$ ,  $a^5 - 10a^4 b^2 + 40a^3 b^4 -$   
 $80a^2 b^6 + 80ab^8 - 32b^{10}$ ,  $a^{18} - 12a^{15} b^3 + 60a^{12} b^6 -$   
 $160a^9 b^9 + 240a^6 b^{12} - 192a^3 b^{15} + 64b^{18}$ .  
 7.  $495a^8 b^4 - 792a^7 b^5$ .

## EXERCISE xii.

1.  $1 - x^3 + x^4 + x^6 + x^{17}$ . 2.  $1 + x + x^2 + x^3 + x^4 + x^6 + x^7 +$   
 $x^8 + x^9 + x^{15}$ . 3.  $x^4 + 2x^3 - 85x^2 - 86x + 1680$ ,  
 $2x^9 - 3x^6 + 4x^5 + x^4 + x^3 - 2x^2 - x + 2$ . 4.  $x^6 - 57x^4 +$   
 $266x^2 - 1$ . 5.  $18x^8 + 21x^7 + 8x^6 + x^5 + 63x^3 + 96x^2 +$   
 $43x + 3$ . 6.  $1 - \frac{1}{2}x^2 - \frac{1}{8}x^4$ . 7.  $6x^{12} - 4x^9 - 5x_0^8 - 2x^7 +$   
 $9x^6 - 10x^5 + x^4 - 5x^3 + 5x^2 + x + 4$ . 8.  $x^3 + 9x^2 + 10x + 11$ .  
 9.  $x^4 + 3x^3$ . 10.  $x^4 - 3x^3$ . 11.  $x^4 + 8x^3 - 8x$ .  
 12. (1), -1, (2), -1, (3) -4. 13. -1.



## EXERCISE xiii.

1.  $3x^3 - 2x^2 - 4x + 2$ .
2.  $5x^4 - 4x^3 + 3x^2 - 2x + 1$ .
3.  $a^4 + 2a^3 + 3a^2 + 4a + 5$ .
4.  $x^3 + 2x^2y - 3xy^2 + 4y^3$ .
5.  $a^3 + 3a^2x + 3ax^2 + x^3$ .
6.  $4x^2 + 8x + 7, -13x - 20$ .
7.  $10x^2 + 5x^2 + 1, 10x + 10$ .
8.  $x^2 - xy + y^2$ .
9.  $x^2 - a^2$ .
10.  $x^4 + (1-a)x^3 + (1-a+b)x^2 + (1-a)x + 1$ .
11.  $3x^3 + 2x^2 + x + 1\frac{1}{2}, 3\frac{1}{2}(x+1)$ .
12.  $5x^2 + 13xy + 12y^2$ .
13.  $6x^5 - x^4 - x^3 + x^2 - x + 6, -1$ .
14.  $2x^4 - 3x^3 + 4x^2 - 5x + 6$ .
15.  $a + b$ .
16.  $x + y + z$ .
17.  $10x^3, 10(x^4 - 20)$ .
18.  $mx^3 + nx^2 + a$ .
19.  $1 + x - 5\frac{3}{4}x^2 - 3x^3 + 9x^4$ .
20. 38.
21. -4.
22. -20.
23.  $15y^4$ .
24.  $85x + 8$ .
25. 755.

## EXERCISE xiv.

1.  $y^3 - 2y^2 - 4y - 9$ , if  $y = x - 1$ .
2.  $y^3 + 3y + 5$ , if  $y = x + 1$ .
3.  $y^4 + 81$ , if  $y = x - 2$ .
4.  $y^4 + 4y^3 - 43y^2 + 92y - 67$ , if  $y = x + 2$ .
5.  $3y^5 + 30y^4 + 119y^3 + 238y^2 + 249y + 106$ , if  $y = x - 2$ .
6.  $y^4 - \frac{59}{8}y^2 - \frac{91}{8}y + \frac{845}{56}$ , if  $y = x - 1\frac{3}{4}$ .
7.  $y^3 - \frac{16}{3}y + \frac{155}{7}$ , if  $y = x - \frac{2}{3}$ .
8.  $(x - 2y^3) - 3y(x - 2y)^2 - 18y^2(x - 2y) - 24y^3$ .
9.  $(x - y)^5 - 10y^2(x - y)^3 - 20y^3(x - y)^2 - 10y^4(x - y)$ .
10.  $(2x + y)^3 + 2y^2(2x + y) + 5y^3$ .
11.  $512y^3 - 3y - \frac{19}{4}$ , if  $y = \frac{1}{4}x - \frac{1}{16}$ .
12.  $y^4 - 24y^2 + 49y - 28$ , if  $y = x + 2$ .

## EXERCISE xv.

1.  $a^2b, +ab^2 + a^2c + b^2c + bc^2 + ac^2,$   
 $(a-b)^2 + (b-c)^2 + (c-a)^2, a(b-c) + b(c-a) + c(a-b),$   
 $ab(x-c) + bc(x-a) + ac(x-b),$   
 $abc(a^2b + a^2c + b^2c + ab^2 + ac^2 + bc^2),$

$$(a+b)(c-a)(c-b) + (b+c)(a-b)(a-c) + (c+a)(b-c)(b-a).$$

$$(a+c)^2 - b^2 + (b+a)^2 - c^2 + (c+b)^2 - a^2,$$

$$a(b+c)^2 + b(c+a)^2 + c(a+b)^2.$$

2.  $abc + bcd + cda + dab,$

$$a^2(b+c+d) + b^2(c+d+a) + c^2(d+a+b) + d^2(a+b+c),$$

$$(a-b) + (a-c) + (a-d) + (b-c) + (b-d) + (c-d),$$

$$a^2(a-b) + b^2(b-c) + c^2(c-d) + d^2(d-a).$$

13.  $x$  and  $y$ . 14.  $ax$  and  $by$ ,  $x$ ,  $y$ ,  $z$ . 15.  $f$  and  $h$ .

16.  $x$  and  $y$ , also  $x$  and  $-z$ , and  $y$  and  $-z$ .

17.  $a$ ,  $b$  and  $-c$ . 18.  $x^2$ ,  $-y^2$  and  $z^2$ . 19.  $b$  and  $c$ .

20.  $a$  and  $c$ . 21.  $a$  and  $b$ . 22.  $a^2$  and  $2ab$ .

23.  $a^2b$  and  $abc$ . 24.  $a^2b$ ,  $abc$ . 25.  $x^5$ ,  $x^4y$ , and  $x^3y^2$ ;  
same;  $x^4y$ ,  $x^3y^2$ . 26. Not symmetrical.

28.  $a^4$ ,  $a^3b$ ,  $a^2bc$ ,  $abcd$ ;  $a^4$ ,  $a^2b^2$ . 29.  $a^3$ ,  $a^2b$ .

## EXERCISE XVI.

1.  $4(a^2 + b^2 + c^2)$ . 2.  $3(a^2 + b^2 + c^2) + 2(ab + bc + ca)$ .

3.  $4(a^2 + b^2 + c^2 + d^2)$ . 4.  $2(a^2 + b^2 + c^2)$ .

5.  $4(x^2 + y^2 + z^2 + n^2)$ . 6.  $2(a^3 + b^3 + c^3) + 6\Sigma a^2b - 12abc$ .

7.  $14(x^2 + y^2 + z^2) + 2(xy + yz + zx)$ . 8.  $24abcnmr$ .

9.  $2abc(a + b + c)$ . 10.  $a^2b^2 + b^2c^2 + c^2a^2$ .

## EXERCISE XVII.

1. 115. 2.  $pa^3 - 3qa^2 + 3ra - s$ . 3. 2. 4.  $-17\cdot3538$ .

5. 1,  $2(3a^2 + 1)$ . 6. 0 or  $2y^n$ ,  $2y^n$ , 0. 7. 36.

8.  $-(b^2 + a^2)^3 - (3b^2)^3$ . 9.  $-15a^4$ . 10.  $3888a^4b^4$ .

11.  $a^2b^2(a+b)$ , 12. 0. 13.  $2a^3 - 3ab(a-b)$ ,

$2b^3 + 6ab(a+b)$ ,  $2(a^3 + b^3)$ .

## EXERCISE XX.

1. 3. 2. 1. 3.  $-1 \pm 2\sqrt{-2}$ . 4. 2. 5. 36. 6. 11.

7.  $-1\frac{9}{27}$  13.  $p = -q$ ,  $q = 6$ . 14.  $p = -46$ ,  $q = 14$ .

## EXERCISE XXI.

1.  $b = -3, c = 8\frac{3}{4}, d = -24.$  2.  $c = -20\frac{1}{2}, d = -13\frac{1}{2}, e = 60\frac{1}{4}.$   
 3.  $b = -3, c = -10.$  4.  $a = 3, b = 0, c = -57.$  5.  $a = -2,$   
 $c = 24\frac{1}{2}, e = 0.$  6.  $c = -106\frac{1}{2}, d = 202\frac{1}{2}.$  7.  $a = 200,$   
 $b = -810, c = 639.$  8.  $a = 4, c = -27, d = 7, e = 30.$   
 9. 399. 10.  $x^3 - (p+3)x^2 + (2p+q+3)x - (p+q+r+1).$   
 11.  $x^3 - (p-3)x^2 - (2p-q-3)x - (p-q+r-1).$   
 12.  $rx^3 - (3r-q)x^2 + (3r-2q+p)x - (r-q+p-1).$   
 13.  $x^3 - qx^2 + prx - r^2.$  14.  $x^3 - (p^2-2q)x^2 + (q^2-2pr)x - r^2.$   
 15.  $x^3 - 2qx^2 + (pr+q^2)x + r^2 - pqr.$  16.  $rx^3 - (pq+3r)x^2 +$   
 $(p^3-2pq+3r)x - (pq-r),$  33. -1. 34. 1. 35. -1.  
 36. 1. 37. -1. 38 and 39.  $a+b+c+d.$  40. -1.

## EXERCISE xxiii.

1.  $5b^4+15c^4.$  2. 6. 3. 3. 4.  $- \{A(b+c+d) + \dots + \dots + \dots\}.$   
 5. 0. 6.  $5b^4 - 30ab^3 + 30a^2b^2 - 5a^3b.$  8. 0. 9. 0. 10. 0. 11. 1.  
 12.  $(a+b+c+d).$  13. -1. 14.  $a+b+c+d.$   
 15.  $(a+b+c)(a^2+b^2+c^2+ab+bc+ca)+abc.$   
 16.  $(a+b+c)^2(a^2+b^2+c^2)+2abc(a+b+c).$  17.  $a+b+c+d.$   
 18.  $(a+b+c+d)^2.$  19.  $(a+b+c+d)\{(a+b+c+d)^2 -$   
 $(ab+ad+ac+bc+bd+cd)\} + abcd.$  20.  $a+b+c.$  21. 3.  
 22. -1. 23. 0. 24. 0. 25. 0. 26.  $1 + \frac{1}{2}x - \frac{1}{8}x^2 + \frac{1}{16}x^3.$   
 27.  $1 - \frac{1}{2}x - \frac{1}{8}x^2 - \frac{1}{16}x^3.$  28.  $1+x+x^2+x^3.$   
 29.  $1-2x+3x^2-4x^3.$  30.  $1 + \frac{1}{2}x - \frac{1}{8}x^2 + \frac{5}{81}x^3.$

## EXERCISE xxiii. (a)

1.  $(p-p'+q)^2 = (p+1)(p^2-pp'-q).$   
 8.  $9(p^2-q)(r^2-qt) - (pr-t)^2 = 9\{3(p^2-q)(qr-pt) -$   
 $(pq-r)(pr-t)\} \times \{3(pq-r)(r^2-qt) - (pr-t)(qr-pt)\}.$   
 9.  $x^2(4x^3+3px^2+3qx+r) \div (x^4+4px^3+6qx^2+4rx+t).$

6. 11.

q = 14.

10.  $-4p, (4p)^2 - 2(6q), -(4p)^3 + 3(4p)(6q) - 3(4r),$   
 $(4p)^4 - 4(4p)^2(6q) + 4(4p)(4r) + 2(6q)^2 - 4t,$   
 $-(4p)^5 + 5(4p)^3(6q) - 5(4p)^2(4r) - 5(4p)(6q)^2 + 5(4p)t +$   
 $5(6q)(4r), (4p)^6 - 6(4p)^4(6q) + 6(4p)^3(4r) + 9(4p)^2(6q)^2 -$   
 $6(4p)^2t - 12(4p)(6q)(4r) - 2(6q)^3 + 6(6q)t + 3(4r)^2.$
11.  $s_0s_4 - 4s_1s_3 + 3s_2^2, s_0s_6 - 6s_1s_5 + 15s_2s_4 - 10s_3^2,$  where  $s_0, s_1,$   
 &c., are the coefficients of the terms (taken in order) of  
 the quotient in No. 10.
12.  $x^n(4x^3 - 28x + 1) \div (x^4 - 14x^2 + x - 38); s_1 = 0, s_2 = 28, s_3 =$   
 $= -8, s_4 = 544, s_5 = -70, s_6 = 8688; \Sigma(a-b)^4 = 4526,$   
 $\Sigma(a-b)^6 = 264122.$

## EXERCISE xxiv.

1.  $(3m+2)^2, (c^m-1)^2.$  2.  $(y^3-z^3)^2, 4y^2(2x+y)^2.$
3.  $(3ab+2c)^2, 4y^2(3x-y)^2$  4.  $(\frac{1}{2}x^2-4yz)^2, (\frac{1}{2}a^2-\frac{1}{3}b^2c^2)^2.$
5.  $(a+b+c)^2, (3x^4-\frac{1}{2}y^2)^2.$  6.  $(z-x+y)^2, \left\{ \left( \frac{a}{b} \right)^m - \left( \frac{b}{a} \right)^m \right\}^2.$
7.  $(x^2-z^2)^2.$  8.  $(x-y)^4.$  9.  $(a+b)^2, (\frac{3}{4}p^3-\frac{4}{3}q^3)^2.$
10.  $(x-y)^2.$  11.  $4(x^2+y^2)^2.$  12.  $(x+y)^4.$
13.  $\left\{ \left( \frac{a}{b} \right)^m - \left( \frac{b}{a} \right)^n \right\}^2$  14.  $(a-b+c)^2.$
15.  $(a^2-b^2-c^2)^2.$  16.  $(2a-2c)^2.$  17.  $(2a^2-3b+4c)^2.$

## EXERCISE xxv.

1.  $(7a+2b)(7a-2b).$  2.  $(3a+\frac{1}{2}b)(3a-\frac{1}{2}b).$
3.  $(3a-2b)(9a^2+4b^2)(3a+2b).$  4.  $(10x-6y)(10x+6y).$
5.  $5b(a+2xy)(a-2xy)$  6.  $(3x^3-4y^2)(3x^3+4y^2).$
7.  $(\frac{3}{4}c+1)(\frac{3}{4}c-1).$  8.  $(2y^2-\frac{2}{3}xz)(2y^2+\frac{2}{3}xz).$
9.  $(3a-1)(3a+1)(9a^2+1).$  10.  $(a-2b)(a+2b)(a^2+4b^2).$
11.  $(a-b)(a+b)(a^2+b^2)(a^4+b^4)(a^8+b^8).$

12.  $(a+b-c)(a-b+c)$ . 13.  $(a+2b-3x+4y)(a+2b-3x+4y)$ .  
 14.  $(x^2-y^2)^2$  15.  $(x+y+2z)(x+y-2z)$ . 16.  $16(x+1)(1-x)$ .  
 17.  $(x+y+z)(x+y-z)(z-x+y)(z+x-y)$ .  
 18.  $4xy(x+y)(x-y)$ . 19.  $(x-z+y)(x-z-y)(x+z+y)(x+z-y)$ .  
 20.  $4(a+c)(b+d)$ . 21.  $24x(1+2x^2)$ . 22.  $8ab(a+b)^2$ .  
 23.  $(a+b+c+d)(a+c-b-d)(a-b-c+d)(a+b-c-d)$ .  
 24.  $(x+y+z)(x-y-z)(x+y-z)(x-y+z)$ .  
 25.  $8a^3b^3(a^6-3a^3b^3+b^6)$ . 26.  $(a^3+b^3)(a^3-b^3)^3$ .  
 27.  $(x^2+y^2+z^2)(x^2+y^2+z^2-2xy-2yz-2zx)$ .  
 28.  $(x+2z)(x-2y)$ . 29.  $(a+b-c)(a-b+c)(b+c+a)(b+c-a)$ .  
 30.  $(x-y+z)(x+y-z)(x+y+z)(x-y-z)$ .

## EXERCISE xxvi.

1.  $(x-7)(x+2)$ ,  $(x-7)(x-2)$ ,  $(x+4)(x+3)$ .  
 2.  $(x-3)(x-5)$ ,  $(x-7)(x-12)$ ,  $(x-12)(x+5)$ .  
 3.  $2(2x-5)(x+2)$ ,  $3(3x-20)(x-10)$ .  
 4.  $\frac{1}{2}(x+12)(\frac{1}{2}x-3)$ ,  $5(x+1)(5x+3)$ ,  $(3x^3-4)(3x^3-5)$ .  
 5.  $(\frac{1}{2}x+4)(\frac{1}{2}x+3)$ ,  $4(4x-5)(x+1)$ .  
 6.  $(x-a)(x+a)(x-b)(x+b)$ ,  $\{2(x+y)-11\}\{2(x+y)+9\}$ .  
 7.  $(x^2+y^2-a^2)(x^2+y^2+b^2)$ . 8.  $(a+b-3c)(a+b+c)$ .  
 9.  $(x+y)(1+x+y)\{x+y+(x-y)^2\}$ .  
 10.  $(a+b)(1-a-b)\{a+b+(a-b)^2\}$ .  
 11.  $(x^2+xy+y^2+2x+y) \times \{x^2+xy+y^2-(x+2y)\}$ .  
 12.  $(a-5b+3c)(a+b-c)$ . 13.  $(x^3+y^3+a^3)^2-b^6 = \&c$ .  
 14.  $(x^2-10x-12)(x^2-10x+8)$ .  
 15.  $(x^2-14x+10)(x-9)(x-5)$ . 16.  $(x^2-y^2)^2$ .  
 17.  $(z+1)(z-1)(z^2-2)$ ,  $(x^2-3)(x^2+1)$ ,

$$(3x^4 + 5y^2)(3x^4 - 2y^2), \quad 18. (c^m + 2)(c^m - 1),$$

$$(x^3 - 2)(x + 1)(x^2 - x + 1), (x^m - 4y^n)(x^m + 2y^n),$$

$$19. (x^m - ay^n)(x^m + by^n).$$

## EXERCISE xxvii.

$$1. (x - by)(bx - y).$$

$$2. 3(x + 2y)(2x - y).$$

$$3. 4(14x - 5y)(x - y).$$

$$4. 4(14x + 5y)(x - y).$$

$$5. (14x - y)(x - 20y).$$

$$6. 4(7x - 5y)(2x - y).$$

$$7. 2(28x + y)(x - 10y).$$

$$8. 4(14x - 5y)(x + y).$$

$$9. (8x - 5y)(7x - 4y).$$

$$10. (8x + 5y)(7x - 4y).$$

$$11. 2(3x + y)(x - 3y).$$

$$12. (3x - 2y)(2x + 3y).$$

$$13. 2(28x + y)(x + 10y).$$

$$14. 2(28x - 5y)(x - 2y).$$

$$15. 2(28x + 5y)(x - 2y).$$

$$16. (56x - 5y)(x - 4y).$$

$$17. 2(4x - y)(7x - 10y).$$

$$18. 4(14x + y)(x - 5y).$$

$$19. 3(3x + y)(4x - 5y).$$

$$20. (8x + 5y)(9x - 8y).$$

## EXERCISE xxviii.

$$1. (5x - 7)(2x + 3). \quad 2. (5x + 3)(2x - 7). \quad 3. (5x - 3)(2x + 7).$$

$$4. (2x - 5)(3x - 11). \quad 5. (4a + 1)(3a - 2). \quad 6. (3x - 7)(4x - 3).$$

$$7. (3x + 7)(4x + 3).$$

$$8. (5a^3 - 4b^2)(3a^2 + 5b^3).$$

$$9. (4x + 1)(3x - 1).$$

$$10. 3y^2(x - y)(3x + 2y).$$

$$11. (2x + 3y)(2x + y).$$

$$12. x^2(3b + x)(2b - 3x),$$

$$13. (3x^2 + 7y^2)(2x^2 - 5y^2). \quad 14. (2x^2 - 9)(x^2 + 5),$$

$$15. (2x + y)(2x - y)(x - 3y)(x + 3y).$$

$$16. (2x + 4 + y)(2x + 4 - y)(x + 2 - 3y)(x + 2 + 3y).$$

$$17. 169xy. \quad 18. (19y^2 + 60xy - 6x^2)(35x^2 - 12xy + 30y^2).$$

$$19. 2(4xy - 3x^2 - 3y^2)(61x^2 - 49xy + 61y^2).$$

$$20. 2(5x^2 + 4xy + 10y^2)(x^2 + 10xy + 2y^2).$$

## EXERCISE XXIX.

1.  $(7x+6y+8)(x-y-z)$ .      2.  $(5x-5y-22)(4x+y+4)$ .  
 3.  $(3x^2+4y^2+13)(x^2-y^2-1)$ .      4.  $(4x+5y)(5x-4y+7)$ .  
 5.  $(9x+8y-20)(8x-y-1)$ .      6.  $(x+3y)(x-4y-5)$ .  
 7.  $(4x+3y-z)(2x+3y+z)$ .      8.  $(3x-2y-2z)(2x-3y+4z)$ .  
 9.  $(3x^2-2y^2+5z^2)(2x^2+5y^2-5)$ .  
 10.  $(15x^2+8y^2+5z^2)(x^2-2y^2+3z^2)$ .  
 11.  $(2a-5b-7c)(2a+3b+3c)$ .  
 12.  $(a-b+c)(a+b-c)(a+b+c)(a-b-c)$ .

## EXERCISE XXX.

1.  $x^2 + \frac{7}{2} \pm \frac{3}{2}\sqrt{5}$ ,  $2x^2 + \frac{7}{2} \pm \frac{3}{2}\sqrt{5}$ .      2.  $x^2 + \frac{7}{2}y^2 \pm \frac{3}{2}y^2\sqrt{5}$ ,  
 $\frac{1}{12}(6x^2+5y^2 \pm y^2\sqrt{13})$ .      3.  $\frac{1}{4}(4x^2+5 \pm \sqrt{13})$ ,  
 $\frac{1}{12}\{6(x+y)^2+5z^2 \pm z^2\sqrt{13}\}$ .  
 4.  $(x^2 + \frac{1}{2}y^2)(x^2 + 6\frac{1}{2}y^2)$ ,  $(x^2 + \frac{1}{2}y^2)(x^2 + \frac{3}{2}y^2)$ .  
 5.  $(2x^2+4\frac{1}{2}y^2)(2x^2+\frac{1}{2}y^2)$ ,  $\frac{1}{4}\{4(a+b)^2+5c \pm \sqrt{13}\}$ .  
 6.  $\frac{1}{12}(6x^2+5y^2)(6x^2+11y^2)$ ,  $(6x^2+5)(6x^2+11)$ .  
 7.  $\frac{1}{5}(5x^2+10 \pm 3\sqrt{10})$ ,  $(2a^2+3 \pm 2\sqrt{2})$ .  
 8.  $\{2(x+y)^2+(3 \pm 2\sqrt{2})z^2\}$ ;  
 $\frac{1}{5}\{10x^2+(10 \pm 3\sqrt{10})y^2\}\{10x^2+(20-6\sqrt{10})y^2\}$ .  
 9.  $\frac{1}{9}(9x^2+7 \pm \sqrt{13})$ ,  $\frac{1}{2}\{2x^2+(6 \pm \sqrt{16})(y+z)^2\}$ .  
 10.  $\frac{1}{2}(2x^2+6 \pm \sqrt{6})$ ,  $\frac{1}{7}(7x^2+20 \pm \sqrt{85})$ .  
 11.  $\frac{1}{2}\{4x^2+(9 \pm \sqrt{23})y^2\}$ .  
 12.  $\frac{1}{7}\{7(a-b)^2+8c^2 \pm c\sqrt{29}\}$ ,  $\frac{1}{6}\{3a^2 \pm b^2\sqrt{3}\}$ .  
 13.  $\frac{1}{3}\{3x^2+(3 \pm \sqrt{3})y^2\}$ ,  $\frac{1}{3}\{3(a+b)^2+(3 \pm \sqrt{3})(a-b)^2\}$ .  
 14.  $\{7a^2+(6 \pm \sqrt{14})b^2\}$ ,  $(5m^2+9n^2)(5m^2+3n^2)$ .  
 15.  $\{7(m+n)^2+(6 \pm \sqrt{14})(m-n)^2\}$ .

## EXERCISE XXXI.

1.  $(x^2 \pm 2xy + 3y^2)$ ,  $(x^2 \pm xy - y^2)$ ,  $(x^2 \pm xy + y^2)$ .
2.  $(x^2 \pm 2xy + 2y^2)$ ,  $(4x^2 \pm 3xy + y^2)$ ,  $(\frac{1}{2}x^2 \pm xy + y^2)$ .
3.  $(x^2 \pm \sqrt{2}x + 1)$ ,  $(x^2 \pm \sqrt{6}xy + 3y^2)$ ,  $(1 \pm 2y - 4y^2)$ .
4.  $(x^2 \pm 3x + 1)$ ,  $x^2 \pm \sqrt{6}x + 3$ ,  $\frac{1}{2}x^2 \pm 2xy + y^2$ .
5.  $(y^2 \pm \frac{1}{2}x^2 \pm \frac{5}{2}x^2 \sqrt{5})$ ,  $(x^4 + 2y^4 \pm 2x^2y^2)$ ,  $x^2 + 4 \pm 2x$ .
6.  $(2x^2 + y^2 \pm \frac{7}{2}xy)$ ,  $(x^2 + y^2 \pm \frac{1}{4}xy \sqrt{39})$ ,  $(2x^2 + 1 \pm 2x)$ .
7.  $(x^{2m} + 8y^{2m} \pm 4x^m y^m)$ ,  $x^{2m} + 2y^{2m} \pm 2x^m y^m$ ,  
 $(\frac{1}{2}x^2 - \frac{3}{4}y^2 \pm xy \sqrt{5})$ .
8.  $(2x^2 - 1 \pm 2x)$ ,  $-(\frac{1}{2}x^2 - 6y^2 + xy)(\frac{1}{2}x^2 - 6y^2 - xy)$ ,  
 $(x^2 + a^2 y^2 \pm axy \sqrt{2})$ .
9.  $mx^2 - ny^2 \pm xy \sqrt{p}$ ,  $x^{2m} + 2^{m-1}y^{2m} \pm 2^m y^m$ .
10.  $4x^2 - 3 \pm x$ ,  $2x^2 - 2 \pm 2x \sqrt{2}$ ,  
 $-(3x^2 - 2y^2 + xy)(3x^2 - 2y^2 - xy)$ .
11.  $2x^2 \pm \frac{4}{3}xy - 3y^2$ ,  $x^2 \pm 2x + 5$ .
12.  $2(a^2 + ab + b^2)^2$ ,  $(2a^2 + a + 1)^2$ .
13.  $\{(x+y)^2 + 3(x+y)z + z^2\} \{(x+y)^2 - 3(x+y)z + z^2\}$
14.  $(a+b)^2 + \frac{7}{2}c^2 \pm \frac{3}{2}c^2 \sqrt{5}$ .
15.  $\{4a^2 + 5a(b-c) + 2(b-c)^2\} \{4a^2 - 5a(b-c) + 2(b-c)^2\}$ .
16.  $4(a^2 + 5ab - 2b^2)(b^2 + 5ab - 2a^2)$ .
17.  $\{(x^2 + y^2 - xy)^2 \pm 3(x^2 + y^2 - xy)(x+y) + (x+y)^2\}$ .
18.  $\{(a^2 + ab + b^2) + \frac{7}{2}(a-b)^2 \pm \frac{3}{2}(a-b)^2 \sqrt{5}\}$ .
19.  $(4a^2 \pm 2a + 1)$ ,  $x^2 \pm 7x + 4$ .
20.  $(x^2 \pm 9xy + 9y^2)$ ,  $(1 \pm 3z + 5z^2)$ .
21.  $4(3x^2 - 2x + 1)(x^2 - 2x + 3)$ .

## EXERCISE XXXII.

1.  $(x^2 + 3)(x + 3)(x - 1)$ .
2.  $2(x^2 + 3)(x^2 + x - 3)$
3.  $(x^2 + 4)(x + 4)(x - 1)$ .
4.  $(x + 2)(x - 2)(3x^2 + x + 12)$ .

5.  $(x^2$
7.  $(\frac{1}{2}x$
9.  $(5x$
10.  $(3x$
12.  $(11$
14.  $8(x$
15.  $(2x$
16.  $(x^2$
17.  $(x^2$
19.  $(x^2$
20.  $(x^2$
21.  $(x^2$
22.  $2(x$
23.  $(x^2$
1.  $(x^2$
3.  $\{x$
5.  $2x$
7.  $(4$
9.  $(x$
10.  $(x$
11.  $(x$
12.  $(x$
13.  $2x$
15.  $x^2$
16.  $(3$
17.  $\frac{1}{2}$



5.  $(x^2 - 3)(5x^2 + 4x + 15)$ .      6.  $(x^2 + 6)(10x^2 + 5x - 60)$ .  
 7.  $(\frac{1}{2}x^2 + \frac{1}{10})(\frac{1}{2}x^2 + 40x - \frac{1}{10})$ .      8.  $(5x^2 - 1)(5x^2 - 8x + 1)$ .  
 9.  $(5x^2 - 8)(7\frac{1}{2}x^2 - 6x - 12)$ .  
 10.  $(3x^2 - 4)(21x^2 - 13x - 28)$ .      11.  $(18x^2 + 1)(45x^3 + \frac{9}{3}x + \frac{5}{2})$ .  
 12.  $(11x^2 + 1)(22x^2 - 3x - 2)$ .      13.  $(\frac{1}{2}x^2 - \frac{2}{5})(\frac{1}{2}x^2 + \frac{1}{5}x + \frac{2}{5})$ .  
 14.  $8(x^2 - 2y^2)(10x^2 - 4xy + 20y^2)$ .  
 15.  $(2x^2 - 5y^2)(12x^2 - 6xy + 30y^2)$ .  
 16.  $(x^2 - 16y^2)(2x^2 + \frac{1}{2}xy + 32y^2)$ .  
 17.  $(x^2 - \frac{9}{5})(11x^2 + 10x + \frac{9}{5})$ .      18.  $10(x^2 + 2)(4x^2 + 3x - 8)$ .  
 19.  $(x^2 - 6y)^2(13x^2 - 12xy + 78y^2)$ .  
 20.  $(x^2 + 4y^2)(3x^2 + 3xy - 12y^2)$ .  
 21.  $(x^2 - 3y^2)(5x^2 + 4xy + 15y^2)$ .  
 22.  $2(x^2 - 2y^2)(2x^2 - 7xy + 2y^2)$ .  
 23.  $(x^2 + \frac{1}{5}y^2)(x^2 + 80xy - \frac{1}{5}y^2)$ .      24.  $(x^2 - 6y^2)(2x^2 - xy + 12y^2)$ .

## EXERCISE xxxiii.

1.  $(x^2 + 3x + 27)(x^2 - 9x + 27)$ .      2.  $x^2 + x(1 \pm \sqrt{3}) + 4$ .  
 3.  $\{x^2 + 1 + \frac{1}{2}(1 \pm \sqrt{5})x\}$ .      4.  $x^2 + 1 - x(2 \pm \sqrt{5})$ .  
 5.  $2x^2 + 2 - 3x \pm x\sqrt{23}$ .      6.  $(x^2 + 15x - 5)(x^2 - x - 5)$ .  
 7.  $(4x^2 - 2)(4x^2 - 6x - 2)$ .      8.  $(x^2 + 8x + 4)(x^2 - 3x + 4)$ .  
 9.  $(x^2 + 7x - 2)(x^2 - x - 2)$ .  
 10.  $(x^2 + 5xy + 3y^2)(x^2 - xy + 3y^2)$ .  
 11.  $(x^2 + 10x - 1)(x^2 + 2x - 1)$ .  
 12.  $(x^2 + 7xy + y^2)(x^2 - 3xy + y^2)$ .  
 13.  $2x^2 + xy - 5y^2 \pm xy\sqrt{46}$ .      14.  $(x^2 + 7xy - y^2)(x^2 - xy - y^2)$ .  
 15.  $x^2 + 2y^2 + 3xy \pm xy\sqrt{3}$ .  
 16.  $(3x^2 + 10xy - 2y^2)(3x^2 - 4xy - 2y^2)$ .  
 17.  $\frac{1}{11}\{11x^2 + 22y^2 + 5xy \pm \frac{2}{11}xy\sqrt{11}\}$ .

## EXERCISE XXXIV.

1.  $(y-z)(x^2-y)$ .
2.  $(by+c)(ax+by-c)$ .
3.  $(z^2+a)(x+a)(x-a)$ .
4.  $(2x-a)(x-2b)$ .
5.  $(x+3a)(x+2b)$ .
6.  $(x-b^2)(x-a)(x+a)$ .
7.  $(x-b)(x+b)(x-a)(x^2+ax+a^2)$ .
8.  $(2x+3a)(4x+5b)$ .
9.  $(a+bx)(a-bx+cx^2)$ .
10.  $(a-bz)(a+bx+cz^2)$ .
11.  $(ax-d)(bx^2+cx-f)$ .
12.  $(px-q)(x^2-x-1)$ .
13.  $(a-b-c)(a+2b+3c)$ .
14.  $(x+a)(x^2+x+1)$ .
15.  $(mx-n)(px^2+qx-r)$ .
16.  $(x-a)(x-b)(x-c)$ .
17.  $(x+a)(x-b)(x-c)$ .
18.  $(x+a)(x+b)(x-c)$ .
19.  $(a^2+z)(x-ay)(x^2-y)$ .
20.  $(abx+cly-cfz)(ax+by)$ .
21.  $(ax+c)(ax^2-bx+c)$ .
22.  $(x-y)(x+y)(mx-ny+rz)$ .
23.  $(mx-ny)(ax+by+cz)$ .
24.  $(mx+n)(ax-bcx+a)$ .
25.  $(c^2-xx)(b^2-yy)(a^2-xy)$ .
26.  $(x^3-m^2x^2-a)(x^2-n+n^2)$ .
27.  $(1+x-x^2)(1-ax+bx^2-cx^3)$ .
28.  $(ax-dy)(ax-by)(ax+cy)$ .
29.  $(mx+q)(px+n)(m^2x-n)$ .
30.  $(mx+ny)(mx-ny)(p^2x^2+q^2y^2)(x+1)$ .

## EXERCISE XXXV.

1.  $(a+x)(a-b)$ .
2.  $(ax+by)(bx-ay)$ .
3.  $(x-a)(x+a)(x^2+ax+a^2)$ .
4.  $x(a+x)(a^2+ax+x^2)$ .
5.  $(ax-b)(cx+d)$ .
6.  $(5x^2-1)(5x^2-x+1)$ .
7.  $(a-b)(a+b+x-c)$ .
8.  $(a^2+b)(a+b)$ .
9.  $(x-y)(x+y)^3$ .
10.  $(x-y+1)(x^2+xy+y^2)$ .
11.  $(b-2x)(2+bx)$ .
12.  $(x-1)(x+2)^2$ .
13.  $(p-q)(p^2-2q^2)$ .
14.  $(a-1)(a^2+2a+2)$ .
15.  $(ab^2-1)(3ab^2+1)$ .
16.  $(y-1)^2(y+2)$ .

17.  $(a+b)(2a^2-3ab+2b^2)$ . 18.  $(b^m-1)(b^{2m}+2b^m+2)$ .  
 19.  $(y^m+z^n)(y^{2n}-3y^nz^n+z^{2n})$ . 20.  $(a-b)(a^2+ab-3b^2)$ .  
 21.  $(a^m-c^n)(a^m-2c^n)$ . 22.  $(ax-b)(x^2-ax-b)$ .  
 23.  $(5x^n-3a^2)(7x^n+3a^2)$ . 24.  $(ab+bc-ca)(ab-bc+ca)$ .  
 25.  $(m-b)(m+b)(a-m)$ . 26.  $(\frac{1}{3}-3a^2)(1-3a)(1+3a)$ .  
 27.  $(x-y-z)(x^2-2xy+y^2+z)$ . 28.  $(6m-7n)(4m^2+n^2)$ .  
 29.  $(x^m+y^n)(x^n+y^m)$ . 30.  $(x^2+xy+ax+y^2)(x^2+xy-ax-y)^2$ .

## EXERCISE XXXVI.

1.  $(x-y)(x+y)(x^2+xy+y^2)(x^2-xy+y^2)$ ,  $(x-1)(x^2+x+1)$ ,  
 $(x+2)(x^2-2x+4)$ ,  $(2a-3x)(4a^2+6ax+9x^2)$ ,  
 $(2+ax)(4-2ax+a^2x^2)$ .  
 2.  $(x-a^2)(x^4+x^3a^2+x^2a^4+xa^6+a^8)$ ,  
 $(3a-4)(9a^2+12a+16)$ ,  $(a^3-b^2)(a^3+b^2)(a^6+3^4)$ ,  
 $(x^2-2y)(x^8+2x^6y+4x^4y^2+8x^2y^3+16y^4)$ .  
 3.  $(a-b)$ . 4.  $x+4y$ . 5.  $(x+y)(x^2+y^2)(x^4+y^4)$   
 6.  $5(y^2-x^2)(7x^4-11x^2y^2+7y^4)$ ,  $(a^2-2b)(a^2+2b)(a^4+4b^2)$ .  
 7.  $y(x-y)(y+1)$ . 8.  $(x-a)(x^2+ax+a^2)(a+b)$ .  
 9.  $(a+b)(m+a)(m^2-am+a^2)$ .  
 10.  $(x^2+xy+y^2)(x^2-xy+y^2)(x^2+2xy-y^2)$ .  
 11.  $(a^2+bc)(a^4-4a^2bc+7b^2c^2)$ .  
 12.  $(x-a+b)\{(x-a)^2-(x-a)b+b^2\}$ .  
 13.  $(x^2-2xy+4y^2)(x+2y+4xy)$ .  
 14.  $(2x+3y)(2x-3y)^2$ . 15.  $(1-2x)(1+4x^2)$ .  
 16.  $(a^2+abc+b^2c^2)(a+bc)(a^2-abc+b^2c^2)$ .

## EXERCISE XXXVII.

1.  $3(x+y)(y+z)(z+x)$ . 2.  $(a-b)(b-c)(a-c)$ .  
 3.  $3(a^2-b^2)(b^2-c^2)(c^2-a^2)$ . 4.  $(x+y)(y+z)(z+x)$ .

5.  $3(a+b)(b+c)(c+a)$ .      6.  $(a+b+c)(a-b)(b-c)(c-a)$ .  
 7.  $(a+b)(b+c)(c+a)$ .      8.  $(a^2-b)b^2-c)(c^2-a)$ .  
 9.  $(a+b)(b+c)(c+a)$ .      10.  $(a-b)(b-c)(c-a)$ .  
 11.  $(x^2+y^2)(y^2+z^2)(z^2+x^2)$ .  
 12.  $(a^2+b^2+c^2-ab-bc-ca)(a-b)(b-c)(c-a)$ .  
 13.  $(a^2+b^2+c^2)(a+b+c)$ .  
 14.  $(c-b^3)(a-c^3)(b-a^3)$ .      15.  $(x^2-y^2)(y^2-z^2)(x^2-z^2)$ .  
 16.  $(x+y+z)(x-y+z)(y-z+x)(z+y-x)$ .  
 17.  $(a-b)(b-c)(a-c)$ .      18.  $8(a+b+c)^3$ .  
 24.  $(a-b)(b-c)(a-c)(a^2+b^2+c^2+ab+bc+ca)$ .

## EXERCISE XXXVIII.

1.  $(a-2)(a^2-7a+2)$ .      2.  $(x-2)(x-3)(x-4)$ .  
 3.  $(x-3)(x-2)^2$ .      4.  $(x-2)^2(x+4)$ .  
 5.  $(x+1)(x^2+2x+3)$ .      6.  $(x^2+2x+3)(x^2+2x+3)$ .  
 7.  $(x+2)(x-1)^2$ .      8.  $(x^2+2x+3)(x^2-2x+3)$ .  
 9.  $(m-n)(m^2-2mn-2n^2)$ .      10. None.  
 11.  $(m-n)(m-2n)^2$ .      12.  $(b+3c)(b^2-2bc+13c^2)$ .  
 13.  $-(m-n)^2(m^2-mn+n)$ .      14.  $(a+2b)(a-2b)(a^2-7ab+4b^2)$ .  
 15.  $(x-5)(x-3)^2$ .      16.  $(x+2)(x^2+3x+1)$ .  
 17.  $(a-1)(a^2-2a-195)$ .      18.  $(p+2)(p-1)(p+4)$ .  
 19.  $(a-1)^2(a+2)(a+3)$ .      20.  $(a^{2n}-1)(a^{2n}-2)(a^{2n}-3)$ .  
 21.  $a^2+4b^2 \pm 7ab$ .      22.  $(a-b)^2(a^2+2ab+2b^2)$ .  
 23.  $(p-2)(p^2-2p+2)$ .      24.  $(x^n-1)(x^{2n}+5x^n+5)$ .  
 25.  $(y-2)(y^3-3y^2+2y+4)$ .      26. None.  
 27.  $(a-b)(a^2+2ab+3b^2)$ .      28.  $(x^n+1)(2a^{2n}-3a^n+2)$ .  
 29.  $(x-2)(x-3)(x-6)(x-7)$ .      30.  $(x-y)(x-2y)(x-3y)^2$ .

## EXERCISE XXXIX.

1.  $2(x-1)(x^2-9x+10)$ ,  $(x-2y)^2(x-3y)$ .
2.  $(4x+3y)(3x^2-xy+y^2)$ ,  $(x-1)(4x-2)(2x+3)$ .
3.  $(x-5a)(3x^2+a^2)$ ,  $(2x+3y)(x^2+3xy-y^2)$ .
4.  $(b+c)(b-4c)(2b^2-bc+c^2)$ ,  $(5a+4b)(3a^2+7ab-3b^2)$ .
5.  $(2p+q)(2p+3q)(p^2+q^2)$ .
6.  $(10x-9y)(15x+16y)(x^3-5xy+8y^2)$ .
7.  $(2x-3y)(2c+3y)(3x+4y)(3x-5y)$ .
8.  $(5x-2z)(2x^3-3x^2y+8xy^2+12y^3)$ .

## EXERCISE XL.

1.  $1+x^2$ .
2.  $(x^2-1)^2$ .
3.  $(x^4+a^2x^2+a^4)(x^8-a^4x^4+x^8)$ .
4.  $(x+2y)(x^2+8y^2)$ .
5.  $1-2x+3x^2$ .
6.  $(a-x)(a+x)^2$ .
7.  $x^2+y^2+z^2+xy+yz-zx$ .
8.  $(a+b)(3a+b)$ .
9.  $(x-y)(2x+3y)$ .
10.  $a^2-b^2+c^2$ .
11.  $7a^2-3ab+2b^2$ .
12.  $a-7$ .
13.  $(a-b)(b-c)(a-c)$ .
14.  $(x-a)^2-b(x-a)+b^2$ .
15.  $x^2+y^2+z^2+1$ .
16.  $x(x^2-ax+b)$ .
17.  $x^2+y^2$ .
18.  $(x-y)(x^2+y^2)$ .
19.  $a^2-b^2+c^2+1$ .
20.  $a^3-b^3-c^3$ .
21.  $a+x$ .
22.  $(c-b)(a+b+c)$ .
23.  $ab-ca-bc$ .
24.  $x^2+y^2+1-xy+x+y$ .
25.  $(x^3-2)(x+1)$ .
26.  $a^2+5a+3$ .
27.  $(2x-y)a^2-(x+y)ax+x^3$ .
28.  $a(x^2+x+1)-(x+1)$ .

## EXERCISE XLI.

1.  $x^2-3$ .
2.  $x+a$ .
3.  $x^2-x+1$ .
4.  $ax^2+bx+c$ .
5. None.
6.  $c^a+c^b$ .
7.  $(a-b)(x+a)$ .
8.  $b(x+y)$ .

9.  $(a-b)(b-c)(c-a)$ . 10.  $a^{2m}+1$ .  
 12.  $5(a-b)(b-c)(c-a)$ . 13.  $(y-1)(x-1)$ .  
 15.  $(x+1)(x^2+1)(x-1)^3$ . 16.  $(x+1)(x+2)(x+3)(x+4)$ . 17. 5  
 18. Same as given quantity. 25.  $(a-b)(b-c)(c-a)$ .  
 29.  $x^4+x^2+2x+1$ .

## EXERCISE xlii.

1.  $(x-1) \div (x^2+4x+16)$ ,  $x(3y-7) \div y(7y-3)$ .  
 2.  $(x^2-ax+a^2) \div (x^2-a^2)$ ,  $(x+4) \div (x-1)^2$ .  
 3.  $(x-1)(x+2) \div (x^2+5x+5)$ ,  $(x^2+2x+3) \div (x^2-2x-3)$ .  
 4.  $1 \div (b-2x)$ ,  $1 \div (x^2-2x+2)$ .  
 5.  $5a^3(a+x) \div x(a^2+ax+x^2)$ ,  $(4x^2+1) \div (5x^2+x+1)$ .  
 6.  $(x-y) \div (c+y)$ . 7.  $(3ax^2+1) \div (4a^2x^4+2ax^2-1)$ ,  
 $(ax+by) \div (ax-by)$ . 8.  $-1 \div abc$ .  
 9.  $-(a+b+c) \div (a-b)(b-c)(c-a)$ . 11.  $5 \div 7(x^2+xy+y^2)$

## EXERCISE xliii.

1.  $(4-x) \div (5-x)$ ,  $(a^2+b^2) \div 2ab$ .  
 2.  $x$ ,  $2a \div (a^2+1)$ . 3.  $a(1+a) \div (1+2a+3a^2)$ ,  $x$ .  
 4.  $b^2 \div a^2$ ,  $(b+1) \div ab^2$ . 5.  $(ac-bd) \div (ac+bd)$ ,  $b \div a$ .  
 6.  $1+6x^2yz(y+z) \div \{y^2z^2-x^2(y+z)^2\}$ . 7.  $(a^2+b^2+c^2) \div abc$ .  
 8. 1. 9.  $-(a^4+a^2b^2+b^4) \div ab(a-b)^2$ .  
 10.  $(a+b+c)^2 \div 2bc$ . 11.  $\left(\frac{1-x}{1+x}\right)^2$ ,  $4a^2x^2 \div (a^2+x^2)$ .  
 12.  $(x+y) \div (x-y)$ . 13.  $(a-b)^3 \div (a+b)^3$ .  
 14.  $(x+y) \div (x-y)$ . 15.  $1 \div x^3$ .  
 16.  $1 \div n$ . 17.  $\pm(1-b) \div (1+b)$ . 18.  $1 \div c$ .

## EXERCISE xlv.

1.  $(x-a) \div 5$ .      2.  $a+b$ .      3.  $16a^5x \div (a^4-x^4)^2$ .  
 4. 0.      5.  $1 \div (x+2)$ .      6.  $1 \div (a^4-x^4)$ .  
 7.  $12xy \div (9x^2-4y^2)$ .      8.  $(4x^2+2) \div x(16x^4-1)$ .  
 9.  $1 \div (x+1)(x+2)(x+3)$ .      10.  $4(x^4+4x^2y^2+y^4) \div (x^4-y^4)$ .  
 11.  $(a-b)^3 \div (x+a)^2(x+b)^2$ .      12.  $2a \div x$ .  
 13.  $(236-77x) \div 18(11x-8)$ .      14.  $1 \div (a-b)$ .  
 15.  $15a(3a-x) \div (9a+2x)(a+3x)$ .  
 16.  $(10x-7) \div (x-1)(2x-5) - 1 \div (2x-7)(x-4)$ .      17. 2.  
 18.  $y^n(y^n-x^n)$ .      19.  $(a-b)^{2n}+2$ .  
 20. 0.      21.  $4x^2 \div (x^{12}-1)$ .  
 22.  $-(a^2+b^2)(a^2-ab+b^2) \div (a^2-b^2)(a^2+ab+b^2)$ .

## EXERCISE xlv.

1.  $x-y$ .      2.  $a+b$ .      3. 0.      4. 0.      5. 0.  
 6.  $\{(a+b)(c+a)x^2+2(ab+bc+ca)ax-2a^2bc\} \div$   
 $(a+b)(a+c)(x+a) \times (x+b)(x+c)$       7. 1.  
 8.  $a+b+c$ .      9. 1.      10.  $x^3-y^3$       11. 0.  
 12.  $(a-b)(b-c)(a-c) \div (a+b)(b+c)(c+a)$ .  
 13.  $x^2 \div (x-a)(x-b)(x-c)$ .      14. 1.      15. 0.  
 16.  $\{b(x+a-b)+ax\} \div \{ab+(b-a)(x-b)\}$ .

## EXERCISE xlvii.

1.  $(a-b)^2+4c^2=0$ .      2. 8.      3. 10.      4.  $a^2+b^2$ .  
 5.  $m=2, n=1$ .      6.  $2x^2$ , or 5.      7.  $m=-5, n=6$ .  
 8.  $\pm 12$ .      9.  $(a^2+b^2)(c^2+d^2)$ .      11.  $-3bc-4c^2+b^2c^2-4b^2$ .  
 12.  $(x^2-4x+3)(x^2-4)$ , also  $(x^2-3x+2)(x^2-x-6)$ .  
 13.  $\frac{1}{2}(-1 \pm \sqrt{5})$ .      15.  $a \div c = d^2 \div e^2, a \div b = f^2 \div e^2$ ,

- $b \div c = d^2 \div f^2$ .      17.  $ac^3 = b^3d$  and  $9ad = bc$ .  
 19.  $4p^3 + 27q = 0$ .      24.  $p = 2m^2q \pm 2mq\sqrt{(m^2 + 1)}$ .  
 25.  $4(p - 3) = q$ .

## EXERCISES xlix.

1.  $5, 3\frac{1}{4}, a, -3$ .      2.  $-4\frac{1}{2}, -a, 2, 10$ .  
 3.  $a + b, c - a, b - c, 3$ .      4.  $-2, 6, -5, 12$ .  
 5.  $-14, a - 3b, 2a - 3b, 5b - 3a$ .      6.  $7, 4, a, b$ .  
 7.  $\frac{1}{3}c, 5 \div a, 0, 1$ .      8.  $-1, \{(a + b)^2 - a\} \div b; a + b$ .  
 9.  $(b - a), a + b$ .      10.  $1 \div a - b, 1 \div (a - b), 1 \div (a^2 + b^2)$ .  
 11.  $2b, a$ .      12.  $a + b, c \div (a + b), b \div (a - c)$ .  
 13.  $(b - c) \div (a - b), b + c$ .      14.  $a + b, a^2 + ab + b^2$ .  
 15.  $a^2 - ab + b^2, 1$ .      16.  $-1, (a + b) \div (a - b)$ .  
 17.  $(e + b)(e - b), 2 \div 15, 3 \div 14$ .      18.  $-1 \div 12, b \div ac, a \div b$ .  
 19.  $(a^2 + b^2) \div a^2b^2 \div a^2b^2, a(b^3 + c^3) \div bc$ .  
 20.  $10, 12, 4, \frac{1}{3}$ .      21.  $1000, \frac{3}{5}, \frac{3}{2}$ .      22.  $9\frac{9}{10}, ab, bc \div a$ .  
 23.  $b^2 \div ac, c(a + b), b(a + b) \div a$ .  
 24.  $a \div b, (a - b) \div (a + b), -(a + b)^2 \div (a - b)^2$ .  
 25.  $-1, -1$ .      26.  $(a^2 - c^2) \div (a + b)^2, 2, 3\frac{1}{3}$ .  
 27.  $ab, b \div a, ac \div b, 12$ .      28.  $12, -ac \div b$ .  
 29.  $9, 2$ .      30.  $12, 1$ .      31.  $3, 1$ .      32.  $(2a - 1)(2a + 2), 0$ .  
 33.  $1 \div m$ .      34.  $1$ .      35.  $(ab + bc + ca) \div (a^2 + bc + c^2)$ .  
 36.  $(a^2 + b^2 + c^2) \div (ab + bc + ca)$ .      37.  $a + b + c$ .      38.  $1$ .  
 39.  $1$ .      40.  $1$ .      41.  $1$ .      42.  $15$ .      43.  $16\frac{1}{2}$ .      44.  $6$ .  
 45.  $5$ .      46.  $(npqa + pqb + qc + d) \div mnpq$ .      47.  $-\frac{1}{8}$ .  
 48.  $0$ .      49.  $-25 \div 136$ .      50.  $1$ .



## EXERCISE I.

1. 2, 3.    2.  $\frac{1}{2}, \frac{1}{3}$ .    3.  $\pm 2, 1\frac{1}{2}$ .    4. 1,  $1\frac{1}{2}$ .    5.  $\pm \frac{2}{3}$ ,  
 $\pm(a+b), a$ .    6. 4, 5, 2,  $2\frac{1}{2}$ .    7. -3 or 2; 4, -3;  
 $2\frac{1}{3}, -1\frac{1}{3}$ .    8. 1;  $\frac{2}{3}$  or  $\frac{5}{3}$ ;  $\frac{1}{3}$  or 3.    9.  $-\frac{2}{3}$  or  $\frac{2}{3}, \frac{1}{6}$  or 6;  
 $\frac{4}{3}$  or  $-\frac{2}{3}$ .    10. -1, 2;  $-\frac{1}{2}, 1$ .    11. 0,  $-b, 3b$ .
12.  $a, \pm a\sqrt{-1}$ .    13. 1;  $\frac{1}{2}(-1+\sqrt{5})$ .    14.  $\pm a$ .
15.  $\pm bc, -(b+c)$ .    16.  $a+2b$ .    17.  $b$  or  $\pm a$ .
18.  $-2ab, \frac{1}{3}ab(1\pm\sqrt{7})$ .    19.  $a, b, -(a+b)$ .    20.  $a, b$ .
21.  $a$  or  $1-a$ .    22.  $-a, -b, a-2b$ .
23.  $a, b, b(1-b)\div(1+a-b)$ .    24.  $x^3-6x^2-37x+210$ .
25.  $x^4-4ax^3-13a^2x^2+64a^3x-48a^4$ .
26.  $x(x-1)(x+2)(x-4)=0$ .    27.  $x^4-4x^3+x^2+6x+2=0$ .

## EXERCISE II.

1. 4.    2.  $-7\frac{6}{7}$ .    3. -107.    4. 8.    5.  $3a$ .    6.  $\frac{31}{13\frac{1}{3}}$ .
7.  $50\frac{30}{9}, 17$ .    8. 22,  $46\frac{1}{3}$ .    9. 7, 3.    10, 10, 10, 11.
11. 0 or 11; 33.    12.  $3956\div 3971$ .    13.  $\frac{1}{7}(15\pm\sqrt{190})$ .
14. 3.    15. 3.    16. 4.    17.  $1\frac{3}{8}$ .    18.  $1\frac{1}{2}$ .    19.  $3\frac{1}{4}$ .
20. 4.    21.  $\pm 3$ .    22. 11.    23. 2 and  $-1\pm\sqrt{-3}$ .
24.  $2\frac{1}{2}$ .    25. 0.    26.  $3a$ .    27.  $\frac{2}{3}$ .    28.  $1\frac{6}{5}$ .    29. 3.
30. 10.    31. 0, 1, or  $(-5\pm\sqrt{-23})\div 8$ .    32.  $102\frac{3}{5}$ .
33.  $(-11\pm\sqrt{4681})\div 20$ .    34. 2,  $\frac{1}{3}, \frac{7}{4}$ .    35. -4
36. 0 or  $\pm\sqrt{(a^2+b^2)}$ .

## EXERCISE III.

1.  $(1-a)\div(1+a), a(m+1)\div(m-1), b(m+1)\div a(m-1)$ .
2.  $a-b, 0, 0$ .    3.  $b, ma\div b, b\div ca$ .    4. 1, -1, 0.
5.  $-\frac{3}{2}$  or -1.    6.  $(c-b)(b^2+c^2)\div 2abc$ .    8. 14,  $4\frac{1}{2}$ .
9. 2, 6-295.    10.  $73\div 210, (a+b+c+d)\div(m+n)$ .

11.  $b \div a$ .      12.  $b \div a$ .      13.  $a$  or  $0$ .  
 14.  $\pm \sqrt{a^2 + 1} \div 2$ .      15.  $\frac{1}{3}$ .      16.  $\frac{1}{3}$ .      17.  $0$  or  $4$ .  
 18.  $c \div ab$ .      19.  $83\sqrt{(2x-1)} = 100\sqrt{(3x-3)}$ .  
 20.  $75 \div 52$ .      21.  $8$ .      22.  $34\frac{49}{4}$ .  
 23.  $1 \div n(n-1)$ .      24.  $ac \div (b-a)$ .      25.  $4, 3\frac{1}{2}$  or  $13\frac{1}{2}$ .  
 26.  $a^2b^2 \div (a-b)^2, 3$       27.  $4a^2 \div (1+a)^2, b(a+b)^2 \div (a-b)^2$ .  
 28.  $(1+b^2) \div 2ab$ .      29.  $\sqrt{(1-x)} = 2 \div (a+1)^2$ .  
 30.  $-a \pm a\sqrt{(1+b+b^2) \div 2b}$ .  
 31.  $\left(\frac{x+1}{x-1}\right)^5 = \left(\frac{a+1}{a-1}\right)^5$ .

## EXERCISE liii.

1.  $8$ .      2.  $0$ .      3.  $3$ .      4.  $(\sqrt{m} \div \sqrt{n})^2$ .  
 5.  $ab \div (1-2\sqrt{b})$ .      6.  $4 \div 7$ .      7.  $1 \div (a-2)$ .  
 8.  $18962 \div 12393$ .      9.  $\sqrt{a} \div (\sqrt{a+2})$ .  
 10.  $(c^4 - 2bc^2) \div (2c^2 - 2b)$ .      11.  $\frac{1}{2}$ .      12.  $18a$ .  
 13.  $x^2 = 80 \div 81$ .      14.  $\pm \frac{1}{3} \sqrt{\frac{1}{2}}$ .      15.  $\pm \frac{1}{11} \sqrt{-11}$ .  
 16.  $\pm \sqrt{\left\{a^2 - \frac{(b-2a)^2}{27b}\right\}}$ .      17.  $0$ .      18.  $\frac{9}{16}a$ .  
 19.  $(c-a-b)^3 = 27abc$ .      20.  $x^2 = a^2(n-1)^2 \div (2n-1)$   
 21.  $16xy = (n-4x-y)^2$ .      22.  $0, -\frac{2}{5}$ .  
 23.  $\left(\frac{a^2}{2a-2} - 1\right)^2, 0$ .      24.  $2\sqrt{(1-m^2)} \div m\sqrt{(4-m^2)}$ .  
 25.  $(a^2-1)\{a^2+2 \pm \sqrt{(a^2+1)}\} \div a^2$ .  
 26.  $(cn-an+c)^2 \div b(n-1)^2$ .      27.  $\pm 5$ .  
 28.  $2\sqrt{(3x^2+10)} = (17\sqrt{17} - 3\sqrt{3}) \div 7$ .      29.  $\pm 5$ .  
 30.  $\pm \frac{1}{4} \sqrt{(3b-2a)}$ .      31.  $\frac{1}{2} \sqrt{(a^2-b^2)}$ .

32.  $(2y + 2z - 2x)^3 + 216xyz = 0$ .      33.  $\frac{2}{3}a\sqrt{6}$ .  
 34.  $a(n^2 - 4n + 8) \div (2n - 4)$ .      35.  $a^2 + 2a$ .  
 36.  $\pm \sqrt{(3a^2 + b^2)} \div \sqrt{3}$ .

## EXERCISE liv.

1.  $-(a^2 + b^2) \div a$ .      2.  $(2a^2 + b^2) \div 2a$ .  
 3.  $\{(a-b)a^2 - 2c(a^2 + ab + b^2)\} \div \{a^2 - 2c(a^3 - b^3)\}$ .  
 4.  $-b$ .      5.  $a + b + c$ .      6.  $ab \div (b - a)$ .  
 7.  $x^2 - 3ax - a^2 = 0$ , &c.      8.  $a$ .      9.  $\frac{1}{3}(a + b + c)$       10.  $1 \div abc$ .  
 11.  $1 \div (a + b + c)$ .      12.  $(a - b)(ac - 2b) \div (a + b)ac$ .      13.  $-c$ .  
 14.  $(\sqrt[3]{a} + \sqrt[3]{b})^3$ .      15.  $\pm 2$ .      16.  $c \div (a - b)$ .  
 17.  $(a - b) \div (a + b)$ .      18.  $\frac{5}{6}a$ .      19.  $\pm 2$ .      20.  $\pm 2$ , &c.  
 21.  $\frac{1}{2}(a + c) \div (a - c)$ .      22.  $a, (3ab - 3b^2 - a) \div (1 + 3a - 3b)$ .  
 23.  $a$ .      24.  $a, b, 2b$ .      25.  $a, (c^2 + 6ab) \div 6b$ .      26.  $\frac{1}{6}(c + 6a)$ .  
 27.  $\frac{1}{2}a$ .      28.  $a + b$ .      29.  $(ab + bc + ca) \div (a + b + c)$ .  
 30.  $\pm b, \pm a$ .      31.  $\sqrt{1 \div (a - 1)}$ .  
 32.  $\{6(a - b) - 4c(c - b)\} \div \{4c - 3b - a\}$ .  
 33.  $(c^2 - ab) \div (a + b - 2c)$       34.  $\frac{1}{2}(-29 \pm \sqrt{37})$ .  
 35.  $(x + a)^2 = 2b^2 - a^2$ .      36.  $\sqrt{(b^2 - \frac{2}{3}ab)}$ .      37.  $\frac{1}{2}(b - a)$ .  
 38.  $3\frac{1}{2}, \frac{5}{4}$ .      39.  $x^2 - 6x = a$ .      40.  $1 \pm \sqrt{19}$ .      41.  $b, b - a$ .  
 42.  $(a^2 + b^2) \div (a + b)$ .      43.  $x = -5 \div 2$ .      44.  $\frac{1}{2}(5 \pm \sqrt{3})$ .  
 45.  $-2a, \frac{5}{7}a, \frac{3}{2}a$ .      46.  $-3a$ .

## EXERCISE lv.

1.  $bc \div (a + c)$ .      2.  $(a^2 + b - 2ab) \div (a + b^2)$ .  
 3.  $(ad - bc) \div (a - b)$ .      5.  $\frac{a^2 + bc}{b + c}$ .      6.  $c$ .      7.  $\frac{1}{2}(a + b)$   
 8.  $a + b$ .      9.  $0$ .      10.  $0$ .      11.  $abc$ .  
 12.  $(a^2 + b^2 + c^2) \div (a + b + c)$ .      13.  $(a + b + c) \div (a^2 + b^2 + c^2)$ .

14.  $(a^2 + b^2 + c^2) \div (ab + bc + ca)$  15.  $\frac{a+b}{a-b}$  16.  $\frac{ab}{a+b}$   
 17.  $4\frac{1}{2}$ . 19. 4. 20. -140. 21. 17. 22. 10. 23.  $a$ .  
 24.  $\frac{e^2(a^2 - b)}{a^2d + c^2 - bd}$ . 25.  $3\frac{1}{4}$ , 0. 26.  $3\frac{1}{2}$ . 27.  $(ab - c^2) \div (a + b)$ .  
 28.  $-b, a$ , 29.  $\infty, 0$ . 30.  $\frac{1}{2}(a + b - c)$ . 31.  $\frac{ab}{a+b}$ .  
 32.  $d$ . 33.  $at - b^2$ . 34.  $-3\frac{2}{3}$ . 35.  $\frac{2}{3}$ . 36.  $-3\frac{2}{3}$   
 37. Infinity. 38. 10. 39.  $abc \div (ab + bc + ca)$ .  
 40.  $(ab + bc + ca - ad - bd - cd) \div (a + b + c - 3d)$ .  
 41.  $a(b + c)^2 \div (b^2 + c^2 - ab + bc - ca)$ .  
 42.  $bc(d - a) - (a - b)(b - c)(c - d) \div (ab + bc + cd - ad - b^2 - c^2)$ .  
 43.  $bc^2 - b^2c - ac^2 + b^2d - abd + acd \div (ab + bc - ac - b^2)$ .  
 44.  $-(a + b + c)$ . 45.  $a + b + c$ . 46.  $(ab + bc + ca) \div abc$ .  
 47.  $-\frac{1}{2}(b + c)$ . 48.  $(ab + c) \div 2a$ . 49. 9. 50. 2. 51. 7.  
 52. 4. 53.  $\frac{1}{2}(5 \pm \sqrt{785})$ . 54. 4,  $(am - nb) \div (n - m + a - b)$ .  
 55.  $\frac{1}{12}$ ,  $b(a + c) \div (a^2 + ab + b^2)$ . 56. 0,  $-\frac{2}{3}$ ,  $\frac{1}{2}$ . 57. 10.  
 58.  $(apnq - cmpq) \div (apn^2 + cqm^2)$ ;  $\frac{nq(ap + mb) - mp(cq + nd)}{apn^2 + mn^2b - m^2cq - m^2nd}$ .  
 59.  $ab \div (b - c)$ ,  $c\{a^2 + (b - c)a - bc\} + a(b^2 - c^2) \div$   
 $(a^2 + b^2 - c^2 + ab - bc - ac)$ . 60.  $b \div (a + b)$ .  
 61.  $mpeq + apmq \div (apn^2 - cqm^2)$ .  
 62.  $\{bm(a - c) + cn(b - a) + ap(c - b)\} \div$   
 $\{m((a - c) + n(b - a) + p(c - b))\}$ . 63.  $(a^2 + b^2) \div ab$ , 0,  $3\frac{1}{3}$ .  
 64.  $(ap - cm) \div (an - bm)$ ,  $\frac{d(n - q) - q(b - d)}{a(n - q) - m(b - d)}$ . 55.  $\frac{1}{4}$ ,  $\frac{1}{8}$ .  
 66. 100. 67. 13, 111. 68. 11, 7.  
 69.  $(a + b - m - n)$ . 70.  $\frac{a^2 + 2ac + aa + 2bc + 2ab}{a - d}$ .

71.  $\frac{1}{2}(a+b) \pm \sqrt{\frac{1}{4}(a-b)^2 - \frac{1}{3}c^2}$ ,  $a$  or  $b$ . 72. 0. 73.  $a+b+c$ .  
 74.  $\frac{ab+bc+ca}{abc}$ . 75.  $a+b+c$ . 76.  $a+b+c$ .  
 77.  $(ab+bc+ca) \div (a+b+c)$ . 78.  $b^2+a^2-c^2$ .  
 79.  $c-a-b$ . 80. 0. 81. 0 or 11.

## EXERCISE lvi.

1.  $A=0$ , or  $B=0$ . 2.  $A=0$ , or  $B=0$ , or  $C=0$ .  
 3.  $x=0$ , or  $a-b=0$ . 4.  $x=0$ , or  $y=0$ .  
 5. In the first case either  $x-5y=0$ , or  $x-4y+3=0$ , in the second case both conditions hold. 6.  $x=0$ , or  $x=a$ .  
 7.  $x=0$ , or  $x=-b$ . 8.  $x=a$ , or  $x=c \div b$ .  
 9.  $x=0$ , or  $x=3$ . 10.  $x=0$ , or  $x=a+b$ .  
 11.  $x=0$ , or  $x=\pm a$ . 12.  $x=0$ , or  $x=b \div a$ .  
 13.  $x=0$ , or  $x=a$ . 14.  $x=0$ , or  $a$ .  
 15.  $x=0$ , or  $x=a+b$ . 16.  $x=0$ , or  $a+b$ .  
 17.  $-(2ab) \div (a+b)$ . 18.  $x=a$ , or  $b$ .  
 19.  $x=a$ , or  $b$ , or  $c$ . 20. 5. 21. 1. 22. 21.  
 23.  $x=\frac{1}{3}$ ,  $x=3$ . 24.  $x=9$ ,  $x=4$ .  
 25.  $x=1$ , or 3. 26.  $(ab) \div (a+b)$ .  
 27.  $x=a$ , or  $b$ . 28.  $x=(a^2+b^2) \div (a+b)$ ,  $x=b+a$ .  
 29.  $(2ab) \div (a+b)$ . 30.  $x=a$ , or  $b$ .  
 1.  $x=1$ , or  $(1+a) \div (1-a)$ . 32.  $x=a$ .  
 34.  $x=a-b$ , or  $\frac{1}{2}(b+c)$ . 35.  $x=a+b$ , or  $\frac{1}{2}(a+c)$ .  
 36.  $x = \frac{a}{a+b+c}$ , or 1. 37.  $a+b-c$ .  
 38.  $x=a$ , or  $\frac{1}{3}(4b-a)$ . 39.  $x=-c$ , or  $a+b+c$ .

40.  $x = 1$ , or  $\frac{m-n}{n-p}$ .
41.  $x = \frac{nc-pb}{mc-ap}$ .
42.  $\frac{p(a-b)-c(m-n)}{m(c-b)-a(n-p)}$ .
43.  $x = \frac{1}{2}(a+b)$ , or  $\frac{1}{2}(b-a)$ .
44.  $x = 2a - b$ , or  $3b - 2a$ .
45.  $x = a + c - b$ , or  $x = \frac{4a+4c-2b}{8}$ .
46.  $x = a + b$ , or  $\frac{17a-7b}{5}$ .
47.  $x = 4a + b$ , or  $a + b$ .
48.  $x = \frac{a-b}{b-c}$ , or  $\frac{a}{c}$ .
49.  $(a-b)(b-c)x^2 - (a^2 + b^2 + c^2 - ab - bc - ca)x + (a-c)(a-b) = 0$ .
50.  $x = \pm 3$ , or  $\pm 2$ .
51.  $x = \pm 6$ , or  $\pm 2$ .
52.  $x = 3$ , or  $\frac{1}{3}$ . 53.  $x = \frac{a+b}{a-b}$ , or  $\frac{a-b}{a+b}$ .
54.  $x = \frac{a}{b}$ , or  $-\frac{b}{a}$ .
55.  $x = b - 2a$ , or  $a - 2b$ .
56.  $x = \frac{2a+3b}{5}$ , or  $\frac{3a+2b}{5}$ .
57.  $x = (mb+na) \div (m+n)$ , or  $(ma-nb) \div (m+n)$ .
58.  $x = \sqrt{\{(m+2n) - \sqrt{(m-2n)}\} \div \sqrt{\{(m+2n) + \sqrt{(m-2n)}\}}}$ .
59.  $a \left\{ \frac{\sqrt{(c+1)} + \sqrt{(c-1)}}{\sqrt{(b+1)} - \sqrt{(c-1)}} \right\}$ .
60.  $a \{ \sqrt{(3c-2)} + \sqrt{(2-c)} \} \div \{ \sqrt{(3c-2)} - \sqrt{(2-c)} \}$ .
61.  $a \{ \sqrt{(2c-1)} + 1 \} \div \{ 1 - \sqrt{(2c-1)} \}$ .
62.  $\frac{1}{2}(a+2b)$ . 63.  $\frac{1}{2}(a+b) - (a-b) \sqrt{(m-2n)} \div \sqrt{(m+2n)}$ .
64.  $\frac{-a^2+b^2}{2b}$ , or  $\frac{a^2+b^2}{2a}$ .
65.  $2ab \div (a+b)$ .
66.  $x = \frac{3a+5b}{8}$ , or  $\frac{3b-5a}{8}$ .
67.  $x = 2a \{ \sqrt{(c+4)} - \sqrt{(c-4)} \} \div \{ \sqrt{(c+4)} + \sqrt{(c-4)} \}$ .

68.

69.

70.

72.

73.

74.

76.

77.

78.

81.

84.

85.

87.

89-

89.

90.

91.

92.

94.

95.

96.

93.

99.

68.  $x = 4$ , or  $3$ .

69.  $\frac{1}{2}\{a \pm \sqrt{a^2 - 4m}\}$  where  $m = a^2 \pm \frac{1}{2}\sqrt{(c+a^4)}$ ;  $3$  or  $1$ .

70.  $a, b$ . 71.  $x = \frac{1}{2}[a+b \pm \sqrt{\{(a+b)^2 - 4(ab+t)\}}]$ ,  
where  $t = \frac{1}{2}(a-b)^2 \pm \frac{1}{2}\sqrt{\{(a-b)^4 + 4r\}}$ .

72.  $x = 0$ , or  $a$ , or  $\frac{1}{2}a(1 \pm \sqrt{-3})$ ;  $x = 4$ , or  $2$ .

73.  $x = 0$ , or  $a+b$ , or  $\frac{1}{2}\{(a+b) \pm \frac{1}{2}\sqrt{(a-b)^2 - 4ab}\}$ .

74.  $x^2 - (a-b)x + ab = 0$ . 75.  $x = \frac{1}{2}(3a-b)$ , or  $\frac{1}{2}(3b-a)$ .

76.  $x = 3a - 2b$ , or  $3b - 2a$ .

77.  $y^2 - m^2 = 0$ , where  $y - m - x$  and  $2m = a + b$ . See Key.

78.  $y^2 - m^2 = 0$ . 79.  $y^2 - m^2 = 0$ . 80.  $y^2 - m^2 = 0$ .

81.  $y^2 - m^2 = 0$ . 82.  $y^2 - m^2 = 0$ . 83.  $y^2 - m^2 = 0$ .

84.  $(y^2 - k^2)(5y^2 + 7k^2) = 0$ , (where also  $k = \frac{1}{2}(a-b)$ ).

85.  $k^4 - y^4 = c$ . 86.  $k^5 + 10k^3y^2 + 5ky^4 = c(k^4 - y^4)$  &  $c$ .

87.  $sy \pm k\sqrt{(k-3c \pm r)} = 0$ , where  $s^2 = 3k + c$ , and  $r^2 = (k-3c)^2$   
 $+ (k-c)(3k+c)$ . 88.  $-3 \pm \sqrt{(9 \pm 12\sqrt{24})}$ .

89—102. Work with a variable  $w$  such that  $wx = x^2 + 1$ .

89.  $w = (a \pm s) \div b$ , where  $s = a^2 + 2b^2$ .

90.  $w = (3a + 2b \pm s) \div 2(a-b)$  where  $s = \pm 5\sqrt{(a^2 + 2ab + 4b^2)}$ .

91.  $w = (3 \pm s) \div (1 \pm s)$  where  $s = (b-4a) \div b$ .

92.  $(w+1)^2 = a \div (a-b)$ . 93.  $w^2 = 2a \div (b-a)$ .

94.  $(x+1) \div (x-1) = a \div (a-8b)$ .

95.  $(w+2) \div (w-2) = \frac{1}{2}(1 \pm s)$  where  $s = (16a+b) \div b$ .

96.  $w^2(4a-b) \div (a-b)$ . 97.  $w^2 = (4a-3b) \div (a-b)$ .

98.  $w = (b \pm s) \div 2a$  where  $s^2 = b^2 + 16a^2$ .

99.  $w = (a+b \pm s) \div 2(a-b)$  where  $s = (a+b)^2 + 8(a-b)$ .

100.  $w = (a+b \pm s) \div 2(a-b)$  where  $s^2 \div (a-b)^2 = \{(a+b)^2 + 4(a-b)^2\} \div (a-b)^2$ .
101.  $(w+2) \pm (w-2) = \pm s \div (4 \pm 3s)$  where  $s^2 = 2a \div (a+b)$ .
102.  $(w+2) \div w = \pm \sqrt{\{5a \div (a+4b)\}}$ .
103.  $\frac{1}{2}(2a+b)$ ,  $\frac{1}{3}(a+2b)$ .      104.  $2a-b$ ,  $\frac{1}{2}(a+b)$ , &c.
105. 1, 2, 4, 5.      106.  $\pm 1$ , 2, 4.      107. 1, 2, 3, 4.
108.  $-\frac{1}{2}$ ,  $-\frac{1}{4}$ , 1,  $\frac{5}{4}$ .      109. -1, 3, 4.      110.  $-a$ ,  $5a$ ,  $5a$ .
111. 15, 20.      112.  $2\frac{1}{4}a$ .      113. 4, -1.
114. 7, -1.      115.  $\frac{1}{2}(bc \div a + ca \div b + ab \div c)$ .      116.  $\pm x \div m$ , &c.
117.  $2s(s-a)(s-b)(s-c) \div \sqrt{\{s'^2 - a^2\}(s'^2 - b^2)(s'^2 - c^2)\}$  where  $2s = a+b+c$ ,  $2s^1 = a^2 + b^2 + c^2$ .
118.  $(2ab + 2ac^2 + 2bc^2 - a^3 - b^3 - c^3) \div 4c^2$ .      119.  $a$ ,  $b$ ,  $\frac{1}{2}(a+b)$ .
120.  $\pm a$  or  $\pm ja \sqrt{3}$ .      121.  $a$ ,  $b$ ,  $\frac{1}{2}(a+b)$ .

## EXERCISE lvii.

- |   |  |  |     |
|---|--|--|-----|
| 1. $x, 7; y, 9$ .                         | 2. $x, 2; y, 1$ .                            | 3. $x, 8; y, 1$ .                      | 37. |
| 4. $x, 9; y, 5$ .                         | 5. $x, -10\frac{1}{2}; y, 5\frac{1}{2}$ .    | 6. $x, -2; y, \frac{1}{2}$ .           | 40. |
| 7. $x, -1; y, 1$ .                        | 8. $x, -2; y, -3$ .                          | 9. $x, -\frac{3}{4}; y, \frac{1}{2}$ . | 42. |
| 10. $x, -\frac{1}{4}; y, \frac{3}{4}$ .   | 11. $x, 12; y, 8$ .                          | 12. $x, 8; y, -9$ .                    | 45. |
| 13. $x, 10; y, 12$ .                      | 14. $x, 12; y, 15$ .                         | 15. $x, 18; y, 13$ .                   | 47. |
| 16. $x, 3; y, 2$ .                        | 17. $x, 7; y, 9$ .                           | 18. $x, 7; y, 3$ .                     | 50. |
| 19. $x, 7; y, 3$ .                        | 20. $x, 2; y, 3$ .                           | 21. $x, 3; y, 4$ .                     | 53. |
| 22. $x, \frac{4}{5}; y, \frac{9}{10}$ .   | 23. $x, 3; y, \frac{1}{4}$ .                 | 24. $x, 12; y, 15$ .                   | 56. |
| 25. $x, \frac{7}{10}; y, \frac{3}{10}$ .  | 26. $x, 8; y, 9$ .                           | 27. $x, 3; y, 1$ .                     | 59. |
| 28. $x, 7; y, 8$ .                        | 29. $x, 11; y, 7$ .                          | 30. $x, 17; y, 13$ .                   | 62. |
| 31. $x, 5; y, -4$ .                       | 32. $x, -\frac{31}{10}; y, -\frac{13}{10}$ . | 33. $x, 13; y, 10$ .                   | 64. |
| 34. $x, 4\frac{2}{5}; y, 3\frac{3}{10}$ . | 35. $x, 11; y, 6$ .                          | 36. $x, 7; y, 5$ .                     | 67. |
|   |  |  | 70. |
|   |  |  | 73. |
|   |  |  | 76. |
|   |  |  | 78. |
|   |  |  | 81. |
|   |  |  | 83. |
|   |  |  | 85. |
|   |  |  | 86. |
|   |  |  | 1.  |
|   |  |  | 3.  |
|   |  |  | 4.  |
|   |  |  | 5.  |



37.  $x, 2; y, 3.$       38.  $x, 5; y, 3.$       39. Equations  
40.  $x, 3; y, 1.$       41.  $x, 7; y, 5.$       not independent.  
42.  $x=0=y=0.$       43.  $0, 0.$       44.  $x=0$  or  $13; y=0$  or  $\frac{2}{11}.$   
45.  $x, 17; y, 20; z, 5.$       46.  $x, \frac{2}{13}, y, \frac{2}{13}, z, \frac{2}{13}.$   
47.  $11, 7, 9.$       48.  $21, 22, 23.$       49.  $-15, -6, -8.$   
50.  $3, 4, 5.$       51.  $12, 15, 10.$       52.  $5, 3, 1.$   
53.  $\frac{5}{8}, 1\frac{1}{2}, \frac{2}{3}.$       54.  $3, 5, 7.$       55.  $11, 13, 17.$   
56.  $5, 3, 1.$       57.  $9, 7, 3.$       58.  $7\frac{1}{2}, 8\frac{1}{2}, 9\frac{1}{2}.$   
59.  $3\frac{1}{2}, 2\frac{1}{2}, 1\frac{1}{2}.$       60.  $2\cdot3, 3\cdot4, 4\cdot5.$       61.  $30, 20, 70.$   
62.  $88 \div 59, 1098 \div 59, 1004 \div 59.$       63.  $30, 12, 70.$   
64.  $6, 12, 20.$       65.  $5, 2, 0.$       66.  $1, 1, 1.$   
67.  $11, 9, 7.$       68.  $5, 3, 1.$       69.  $2, 3, 1.$   
70.  $3, 4, 5.$       71.  $\frac{1}{3}, \frac{1}{3}, \frac{1}{3}.$       72.  $5, 4, 3.$   
73.  $7, 3, 1.$       74.  $2, 3, 1.$       75.  $1, 3, 5.$   
76.  $0, 1, 2.$       77.  $1755 \div 698, 360 \div 349, -15705 \div 698.$   
78.  $\frac{1}{2}, \frac{1}{3}, 1.$       79.  $5, 4, 1, 3.$       80.  $4\frac{2}{3}, 3\frac{3}{10}, 2\frac{2}{15}, 1\frac{1}{10}.$   
81.  $31, 41, 51, 21.$       82.  $7, 4\frac{1}{2}, 4, 8\frac{1}{2}.$   
83.  $20, 10, 0, 30.$       84.  $11 \div 24, \frac{1}{4}, 1 \div 24, \frac{1}{4}.$   
85.  $270 \div 117, -52 \div 117, 15 \div 117, -126 \div 117.$   
86. Each 210.

## EXERCISE lviii.

1.  $(a'c-ac') \div (a'b-ab').$       2.  $b(cn-dm) \div (ad-bc).$   
3.  $b(d-c)(d-a) \div d(b-c)(b-a), c(d-a)(d-b) \div d(c-a)(c-b).$   
4.  $y = cz + du + ew + ax, z = du + ew + ax + by,$   
 $u = ew + ax + by + cz, w = ax + by + cz + du.$   
5.  $x = \frac{1}{2}m(a-b+c), \&c.$       6.  $x = \{p(a^2-b) - m(ab-1)\} +$

$$n(b^2 - a) \div \{a^3 + b^3 - 3ab + 1\}, \text{ \&c.}$$

7.  $x = (l - am + abn - abcp + abcd r) \div (1 + abcde), \text{ \&c.}$

8.  $1 = a \div (1 + a) + b \div (1 + b) + c \div (1 + c).$

9.  $1 = ab + bc + ca + 2abc.$

EXERCISE lix.

1.  $(nc - bd) \div (na - bm), (mc - ad) \div (mb - na).$

2.  $(no + bd) \div (an + bm), (mc - ad) \div (bm + an).$

3.  $c(n - b) \div (an - mb), c(m - a) \div (bm - am).$

4.  $(b - c)a \div (b - a), b(a - c) \div (a - b).$  5.  $ab \div (a + b), y, \text{ same.}$

6.  $ab^2 \div (a^2 + b^2), a^2b \div (a^2 + b^2).$  7.  $ac \div (a + b), bc \div (a + b)$

8.  $(a^2 - b^2) \div (am - bn), (b^2 - a^2) \div (bm - an).$

9.  $a + b - c, c + a - b.$  10.  $a + c, b + c.$

11.  $a(cn - dm) \div (bd - ac), b(en - dm) \div (ad - bc).$

12.  $y = \{9\delta(a^2 - c^2) - b(b + 2a)\} \div \{(a - b)^2 - c^2 + 4bc\}.$

13.  $a + b - c, a - b + c.$  14.  $a + b - c, c + a - b.$

15.  $(m - c)(n - a) \div (b - a), \text{ \&c.}$  16.  $1 + (a - b)(a - c), \text{ \&c.}$

17.  $(m - bc)(l - a) \div (c - a)(a - b), \text{ \&c.}$

18.  $x = p \div (pl + mq + nr) + a, \text{ so } y \text{ and } z.$

19.  $p\{1 - (la + mb + nc)\} \div (pl + mq + nr) + a, \text{ \&c.}$

20.  $(m^2 + 2a^2 - b^2 - c^2) \div 3a, \text{ \&c.}$  21.  $y = a - b + c, \text{ \&c.}$

22.  $x = (ab + bc + ca)(b + c - 2a)(2b - a - c) \div \{(a - c)(b + c - 2a) + (b - c)(2b - a - c)\}.$  Corrected equation,  $x = \frac{1}{2}(b + c), \text{ \&c.}$

23.  $ma \div (a + b + c), \text{ \&c.}$  24.  $npr \div (anp + bmp + cmq).$

25.  $1 \div (b - c), \text{ \&c.}$  26.  $\frac{1}{2}(b + c - a), \text{ \&c.}$

27.  $au \div d, \text{ \&c.}$  28.  $z = 1 \div (a + b - c).$

29.  $a + b, \text{ \&c.}$  30.  $1 \div 2a, \text{ \&c.}$

31.  $(m^2 + n^2 - l^2) \div 2mn$ , &c.      32.  $\frac{1}{2}(a+c-b)$ , &c.  
 33.  $l(m^2 + n^2) \div 2mn$ , &c.      34.  $1 \div (b+c-a)$ , &c.,  
 35.  $bc \div (b+c)$ , &c.      36.  $b+c-a$ , &c.  
 37.  $a, b, c$ .      38.  $b^2 - c^2$ , &c.  
 39.  $\frac{1}{2}(a+2b-c+3d)$ , &c.      40.  $\frac{1}{4}(4a+b+3c-2d+5e)$ .

## EXERCISE 1X.

1.  $a+b, a-b$ .      2.  $\frac{1}{2}(a^2+b), \frac{1}{2}(a^2-b)$ .  
 3.  $(3n^2+m^2) \div 5m, (2n^2-m^2) \div 5m$ .  
 4.  $a \div (a-b), b \div (a+b)$ .      5.  $1 \div (a-b), 1 \div (a+b)$ .  
 6.  $a+b-c, a-b+c$ .      7.  $a+b-c, a-b+c$ .  
 8.  $(a^2+ab+b^2) \div (a+b), (a^2-ab+b^2) \div (a-b)$ .  
 9.  $(ab-1) \div (a-1)(b-1), (a-b) \div (a-1)(b-1)$ .  
 10.  $(1+a) \div (ab-1), (1+b) \div (ab-1)$ .  
 11.  $(a+1)(b+1) \div (ab-1), (a-b) \div (ab-1)$ .  
 12.  $a(a+b), b(a-b)$ .      13.  $a\{b(a+b)-c(a-c)\} \div (a^2-bc)$ ,  
 $a\{b(a-b)+c(a+c)\} \div (a^2-bc)$ .      14.  $-(a+b). ab$ .  
 15.  $\frac{1}{2}(b+c)$ , &c.      16.  $(a-2b+3c) \div 38$ , &c.  
 17.  $2 \div (b+c)$ , &c.      18.  $a+b$ , &c. (by symmetry).  
 19.  $b^2 - c^2$ , &c.      20.  $b^2 - c^2$ , &c.  
 21.  $\frac{1}{2}abc, (1-a)(1-b)(1-c), (2-a)(2-b)(2-c)$ .  
 22.  $2abc \div (ab+bc-ca)$ .      23.  $1, 1, 1$ .  
 24.  $ar \div (ma+nb+pc+qd)$ , &c.  
 25.  $u=0$ , or  $\left(\frac{d}{a}-1\right) \div \left(\frac{d^2}{b^2}-\frac{d^2}{c^2}\right)$ .  
 26.  $(b+c-a) \div (a+b+c), y = (b-c-a) \div (a-b-c)$ .  
 27.  $\frac{1}{2}(a-b+m-n)$ , &c.

28.  $(4a+2c-d-3b)$ ,  $y+z$  by symmetry. 16.  
 29.  $-(a-b+c+d)$ ,  $(ab+bc, \&c.)$ ,  $-(abd+\&c.)$ ,  $abcd$ . 17.  
 30.  $\frac{1}{2}(a-b+c-d+e)$ , others by symmetry. 18.  
 31.  $x = (a-lb+lmc-lmnd+lmnp) \div (1+lmnpq)$ , the others by 19.  
 symmetry. 32.  $x = b+c-e$ , &c. 18.  
 34.  $y = (a+5b+3c-7d+9e) \div 22$ , &c. 19.  
 35.  $z = \frac{1}{2}(a+c)$ , then symmetry. 36.  $z = c+d+e$ , &c.  
 37.  $x = a-2b+3c-2d+e$ , then by symmetry. 20.

## EXERCISE lxi.

1.  $x = (2ab+a+b+r) \div 2(a-b)$  where  $r = 4a(b^2+b+1) + (3a-b)(3b-a)$ . 24.  
 2.  $x = (ar+1) \div (ar-1)$  where  $r^2 = (b^2-1) \div 3(a^2-b^2)$ .  
 3.  $x = \{\sqrt{(1+a)(1+b)} - \sqrt{(1-a)(1-b)}\} \div \{\sqrt{(1+a)(1+b)} + \sqrt{(1-a)(1-b)}\}$ . 25.  
 4.  $(ab-a\beta) \div (ab+\alpha\beta)$ ,  $(a\beta+b\alpha) \div (a\beta-b\alpha)$ . 26.  
 5.  $x = \{\sqrt{(a+b+c)(a+b-c)} + \sqrt{(b+c-a)(a+c-b)}\} \div \{\sqrt{(a+b+c)(a+b-c)} - \sqrt{\dots}\}$ .  
 6.  $(a+b) \div (1-ab)$ . 7.  $x = (\alpha\beta-ab) \div (a\beta+b\alpha)$  28.  
 8.  $\left(\frac{x+1}{x-1}\right)^4 = \frac{(a+m)(b+n)}{(a-m)(b-n)}$ . 29.  
 9.  $x = \{a\sqrt{(1-b^2)} - b\sqrt{(1-a^2)}\} \div \sqrt{(a^2-b^2)}$ . 30.  
 10.  $x = (b+c-a) \div \{\sqrt{(b+c-a)(c+a-b)(a+b-c)}\}$ . 31.  
 11.  $x = (b+c) \div \sqrt[3]{(a+b)(b+c)(c+a)}$ . 32.  
 12.  $x = \sqrt{(a+b+c)} \div a$ , &c.  
 13.  $x = \{b^2+c^2-a(b+c)\} \div \sqrt[3]{2(a^3+b^3+c^3-3abc)}$ .  
 14.  $(b+c-a)$ , &c. 15.  $a$  or  $(a^2-b) \div (1-ab)$ .

16.  $x+y = \sqrt{(a+b)(a+2b)} \div \sqrt{(a-b)}$ ,  
 $x-y = \sqrt{(a+b)(a-2b)} \div \sqrt{(a-b)}$ , &c.
17.  $(x+y)^2 = \frac{1}{2} \left( 3b - \frac{a}{b} \right)$ ,  $(x-y)^2 = \frac{1}{2} \left( \frac{3a}{b} - b \right)$ .
18.  $(x+y) \div (x-y) = \sqrt{(a+3b)} \div \sqrt{(a-b)} = m$  suppose.
19.  $y^2 = m \div (am^2 - m + 1)$  where  $m =$   
 $1 \div a + b \pm \sqrt{(a^2 - b^2 + 1)} \div (a+b)$ .
20.  $a, b$ .      21.  $x = \{ \sqrt{(a-c)} + \sqrt{c} \} y \div \{ \sqrt{(a-c)} - \sqrt{c} \}$ .
22.  $x = (a+c)y \div (a-c)$ , &c.
23.  $x+y = (ab-1) \div (a-b)$ , &c.
24.  $xy = \{ \sqrt{(b+2)} - \sqrt{(b-2)} \} \div \sqrt{\{ (b+2) - \sqrt{(b-2)} \}} =$   
 $p$  suppose,  $x \div y = \{ \sqrt{a+2} + \sqrt{(a-2)} \} \div$   
 $\{ \sqrt{(a+2)} - \sqrt{(a-2)} \}$ .
25.  $x^2 y^2 z^2 = \frac{1}{2} (a+b-c)(b+c-a)(c+a-b)$ , &c.
26.  $x = (ab-bc-ca) \div 2\sqrt{abc}$ .      27.  $a-x^2 = \pm m$ , where  $m$  is  
the value of  $v$  in the equation  $4ca - 4(c+a)v + 4v^2 =$   
 $(ca-ab-bc)^2 + 4b(ca-ab-bc)v + 4b^2v^2$ .
28.  $x = \frac{1}{2} \sqrt{(abc)} \left( \frac{1}{b} + \frac{1}{c} \right)$ ,  $y$  and  $z$  by symmetry.
29.  $a(b^2+c^2) \div (b^2+c^2) \div (b^2+c^2)$ , &c.
30.  $c \{ \sqrt{(a+b)} + \sqrt{(a-b)} \} \div \sqrt{(a+b)}$ .
31. 0 or  $a(b+c) \div 2bc$ , &c.
32.  $z = -1$  or  $a\sqrt{(a^2-1)} \div \sqrt{(a^2-b^2)}$ , &c.

## EXAMINATION PAPERS.

## I.

1.  $(a+b+c)(x+y+z)$ .
2.  $(x^2-4y^2)^3$ ,  $2(a^2b^2+b^2c^2+c^2a^2)-a^4-b^4-c^4$ .
3.  $x^4-6x^3+13x^2-12x+4$ ,  $x^2+9x^{-2}+3$ ,  
 $x^{n(n-1)}+x^{n(n-2)}+x^{n(n-3)} \dots\dots$
4.  $2x^{2m}-\frac{1}{3}x^{3m}$ ,  $\frac{a}{b}-\frac{b}{c}-\frac{c}{a}$ .      5.  $\pm 14\sqrt{-19} \div 27$ , 0 or  $\frac{4}{3}$ .
6. 16, 7. 199, 8. AB 37, CA 52, BC 45.

## II.

1.  $(a+b)^3$ .      2.  $(a+b+c)(a^3+b^3+c^3-3abc)$ .      3.  $1 \div 2^3$ .
4.  $(a^{2m}+2a^m+2) \div (a^m+2)$ ,  $(a+b+c) \div (a-b-c)$ .
7.  $-\frac{7}{8}$ , 4 or 6.      8.  $p^2q \div (r-pq)$ .
9. 584.      10.  $(am+bn) \div (a+b)$ , (by common rule).

## III.

1.  $-117$ ,  $a^2(z-x)+(x-y)ab+(y-z)b^2$ .
2.  $a+bx+cx^2$ ,  $3-4x+7x^2-10x^3$
3.  $2\frac{1}{2}$ ; 6, 9, 12.      4. 160 eggs.
5. 40, 35.      6.  $\sqrt{\frac{1}{2}(1+a)}+\sqrt{\frac{1}{2}(1-a)}$ .
7. 5 or  $\frac{1}{4}$ ; 4.      9.  $(a+b+c)(a-b)(b-c)(a-c)$ .

## IV.

1.  $(a+b+c)^2$ .      2.  $a \div b$ .      3.  $(4x^2-9y^2)(4x^2-4y^2)$ ,  $1+\sqrt{x}$ .
4.  $\frac{1}{2}\sqrt{3}-\frac{1}{6}\sqrt{6}$ , 17.      5. 4.      6. 1.      7. 0 or 4; 8, 1.
8.  $7\frac{1}{2}$ , 12.      9. 4 or 6.      10.  $\frac{1}{2}(-3 \pm \sqrt{-39})$ .

## V.

1.  $8x^3+1 \div 125$ .      3.  $16a^2b^2$ .      4.  $b \div 8a$ .      5. 15, 12.
6.  $\frac{1}{2}a$ ,  $\frac{1}{4}a$ ; 4 or -9,      7. 6, 7, 8, or -6, -7, -8.

## VI.

1.  $(a^2 + b^2)$ .      2.  $a^2 + b^2 + c^2 + ab + ac - bc$ .  
 3.  $x^2 + y^2 \pm \frac{3}{2}xy$ ,  $(7x + 6y - 9)(x - y + 4)$ .      4.  $-20, 0$ .  
 5.  $b$  or  $1 \div l$ .      7.  $x = 4$  or  $9$ ,  $y = 9$  or  $4$ ;  $1, 2, 3$ ;  
 $x = \frac{1}{2}(-7 \pm \sqrt{33})$ .      8.  $-1, 0, 1$ , or  $5, 6, 7$ , or  $-\frac{1}{7}, -\frac{4}{7}, \frac{3}{7}$ .  
 9.  $x(x^2 + 3)(x^2 - 2x - 1) = 0$ .      10.  $x \div y = 3 \div 4$ .

## VII.

2.  $-.01$ .      4.  $x^2 + x - 1$ .  
 5.  $(36x^2 + 18x + 9) \div (16x^4 - 81)$ ,  $(x^2 - ab) \div (x^2 + ab)$ .  
 6.  $9; 3; x = \frac{1}{2}(a - b)$ .

## VIII.

2.  $(x^3 - y^3)^2(x - y)$ ;  $a = p + q$ ,  $b = pq$ .  
 3.  $(x^2 - xy + y^2)^2$ ;  $a^4 + b^4 - a^2b^2$ .  
 5.  $ax^3 + bx + cx^{-1}$ ;  $\frac{1}{2}x^{\frac{3}{2}} + \frac{1}{3}x^{\frac{5}{3}} - \frac{1}{4}x^{\frac{7}{2}}$   
 6.  $x^2 + px + p^2$ ;  $50(x + 5)(x - 4)(x - 6)$ .  
 7.  $a = 0 = b$  or  $a = 1$ ,  $b = 2$ .  
 8.  $3$  or  $-43 \div 7$  satisfies the equation  $2 - \sqrt{\dots}$  &c.

## IX.

4.  $axy + b$ .      7.  $\pm \sqrt{ab}$ ;  $\sqrt{(a+b)} \div \sqrt{(a-b)}$ , and  $y = \text{reciprocal}$   
of this.      8.  $P(22a - 21b) \div 20a(a - b)$ .      9.  $7, 15, 48$ .

## X.

1.  $ax^2 + by^2 + 2cxy$ .      3.  $x - 2$ .      4.  $24$ .  
 5.  $5x^2 - 3ax + 4a^2$ ;  $\frac{x}{y} + \frac{y}{x} - \sqrt{\frac{1}{2}}$ .      7.  $3; \frac{1}{2}$ .      8.  $4, 6$ .

## XI.

1. 2.      2.  $6, 8$ .      3.  $(4x^{\frac{1}{2}} - 9y^{\frac{1}{2}})^3, \frac{1}{19}x^4 + \frac{4}{81}y^4$ .  
 4.  $x^2 + 3 + 9x^{-2}$ ,  $x^2 - (a + b)x - c$ .      6.  $0$ .  
 7.  $0$  or  $-2\frac{1}{2}$ ;  $y = \pm 3$  or  $\pm \sqrt{-9}$ , &c.

## XII.

2.  $a^{2^n} - b^{2^n}$ .      3.  $1+x+2x^2+4x^3+8x^4, 2^rx^{r+1}$ .  
 5.  $(1-3x-4x^2)(1-2x-13x^2+38x^3-24x^4)$ .  
 6.  $m^2 - m = p, q = 0$ .  
 7.  $x^2 = (a-2) \div (a+4), \sqrt{x} = \{b(n-1) - cn\} \div \sqrt{a(n-1)}$ .  
 8.  $ab \div \{(a-1)b - 1\} - 1\}$ .      9. 37.

## XIII.

1. 2. 3.  $1+c_1x+c_2x^2+\&c.$ , where  $c_1, c_2, \&c.$ , represent the combinations of  $a_1, a_2, \dots$  taken one, two, &c. at a time.  
 4. \$4000.      5.  $\{m(b'c-bc') - n(ca'-c'a)\} \div (ab' - a'b)$ ,  
 $\frac{a}{a'} = \frac{b}{b'} = \frac{c}{c'}$ .      7.  $(\alpha+\beta)^{\frac{3}{2}} + (\alpha-\beta)^{\frac{3}{2}} - 2\alpha^{\frac{3}{2}} = 0$ .      9. 65.

## XIV.

1. 12abc.      4. 8 miles.      5.  $am \div n; x+y = \pm 5$  or  $\pm 1$ ,  
 $x-y = \pm 1$  or  $\pm 5$ .      6. 20.

## XV.

2.  $\frac{1}{8}$ .      3.  $c^2(c^2 - bd) + d^2(b^2 - ac) + ac(ad - bc)$ .      4.  $2 \div (m-n)$ .  
 If  $m-n$  is negative  $x$  is neg. which shows that they were together before noon. If  $m-n=0$ ,  $x$  is infinite, i.e., they are never together.  
 5.  $x^2(x^2 - a^2)(x-2a); (x^2 - a^2)(x^2 - y^2)$ .  
 6.  $(a+b+c+3d) \div (a+b+c+d); (x+y+z) \div (x-y+z);$   
 $3a \div (a+b)$ .

## XVI.

4.  $2x^4 - 3x^3 + 4x + 3$ .      5. See paper XIX., prob. 4.  
 6.  $\frac{1}{2}\{1 \pm \sqrt{(a^{\frac{1}{n}} + 4b^{\frac{1}{n}}) \div \sqrt{4b^{\frac{1}{n}}}}\}$ .      8.  $(b_2c_1 - b_1c_2) \div (a_1b_2 - a_2b_1)$   
 $= (b_3c_2 - b_2c_3) \div (a_2b_3 - a_3b_2)$ .      9.  $ac \div (a-b)$ ,



## XVII.

1.  $\{2ab - (a+b)\} \div ab$ .      2.  $x^4 - 2x^2(a+b^2) + (a-b^2)^2$ ;  
 $8a^2 - 2ab - 10ac - 3b^2 + 2c^2 + 5bc$ .      6.  $m^{\frac{3}{2}} - n^{\frac{3}{2}} = 4$ .  
 8.  $(m+n)(bq-pc) \div (mq-pn)$ ,  $(p+q)(mc-bn) \div (mq-pn)$ .

## XVIII.

1.  $1 - x^{16}$ .      2. 1.      4.  $x - a - b$ .      6.  $(x-y)(x-z)$ .  
 7. 1.      8.  $9a$ ;  $\pm 1 \div \sqrt{2}$  or  $\pm \frac{1}{2} \sqrt{6}$ .      9. 10, 11.

## XIX.

1.  $\frac{1}{4}(4x-a+1)^2$ .      3.  $5x^2 - 1$ ;  $(x^2 + xy + y^2)^2$ .  
 4.  $(a-c)(a-d)(b-c)(b-d) \div (a+b-c-d)^2$ .  
 6.  $(a+2)^2 \div 4(a^2+a)$ .      7.  $(7b^2 - a^2) \div 12a$ .      8. 18, 22, 50.

## XX.

1.  $(1+m)x + (1-n)y$ .      3.  $x^2 - 1$ .  
 4.  $(x+y-z)(y+z-x)(z+x-y) \div (x+y+z)^2$ ;  
 $(a^2b - 2ab^2 - a^3 + ab^3 + b^3) \div (a^4 - b^4)$ .  
 5. 1.      6.  $a''(cb' - bc') + b''(ac' - a'c) + c''(a'b - ab') = 0$ .  
 7.  $\sqrt{\left\{n^2 - \left(\frac{m^3 - 2n}{3m}\right)^3\right\}}$ ; 2, 3, 4.  
 8.  $a(b-c) \div (b-a)$ ,  $b(c-a) \div (b-a)$ .      9. 3.      10. 2000.

## XXI.

1. 8.      2.  $24abc$ ;  $2a^8 + 43a^6x^2 + 62a^4x^4 + 44a^2x^6 + 18x^8$ .  
 4. 0;  $a^7 + 16$ .      5. 16;  $x + 2a^{\frac{1}{2}}x^{\frac{1}{2}} - 2a$ .      7. 3377 oz. of gold,  
 783 oz. of silver.      9.  $-(8 \pm 4\sqrt{3}) \div (3 \pm 2\sqrt{3})$ ;  
 $x = \pm 2$  or  $\sqrt{-1}$ ,  $y = \mp 1$  or  $\mp 2\sqrt{-1}$ ; 8, 4.  
 10.  $y = \text{cost of 2nd bale} = 60 \pm 20\sqrt{7}$ .

## XXII.

1.  $\cdot 02997$ ,  $a^3 + aq + p^3$ .      2.  $ab^2 - b^2 + c = 0$ ,  
 (a).  $(a-b)(8a-3b)$ . (b).  $x(x-1)(a-b)(b-c)(a-c)$ .  
 3.  $b^2 = 4ac$ . (a).  $(a+b)^2$ .      5. (a)  $(ax-by) \div (ax+by)$   
 7. (a)  $\frac{1}{3}(a+b+c)$ , (b),  $\frac{2}{3}$ ,  $\frac{2}{3}$ , 2.  
 (c).  $b-c$ ,  $c-a$ ,  $a-b$ . (d).  $-1 \pm \sqrt{2}$ .

## XXIII.

1.  $\frac{1}{3}x^5 + \frac{4}{3}x^4 - x^3 + \frac{1}{3}x^2 + \frac{1}{3} + (95x^3 + 21x^2 - 40x + 42) \div$   
 $(3x^4 - 21x^3 + 9x - 6)$ ;  $-382$ .      4.  $x = a + 2c$ ,  $y = b + 3c$ .  
 5. (2).  $\frac{1}{3}$ .      7. (1).  $a+b+c$ . (2), 1, 2, 3, 4. (3), 0 or  $\frac{7}{3}$ .  
 8.  $\frac{1}{2^4}(u^2 - 1)(3n+2)$ .      11.  $\begin{Bmatrix} a_1 b_1 c_1 d_1 \\ a_2 b_2 c_2 d_2 \\ a_3 b_3 c_3 d_3 \\ a_4 b_4 c_4 d_4 \end{Bmatrix} = 0$ .  
 12. Coll. to Newmarket 63 miles.

## XXIV.

1.  $1 \div \{(x-y)(y-z)(z-x)\}$ .      2.  $3 - m - n$ ; 0.      3. 5, 3,  $4\frac{1}{2}$ .  
 4. (1).  $A = bc \div 2a$ ,  $B = ac \div 2b$ ,  $C = ab \div 2c$ ; (2).  $a^2 + b^2 = c^2$ .  
 11.  $y = \pm 2$ ,  $x = \pm 5$ , &c.;  $x^2 - 10x = -19$  or  $-16$ ,  
 $\therefore x = 8$  or  $3$ , &c.;  $(b^{-\frac{1}{2}} \pm a^{-\frac{1}{2}})^{\frac{4pq}{p-a}}$

## XXV.

1.  $(a^2 - 3x) \div (a^2 - x^2)$ ; 1.      2.  $\frac{1}{a} - \frac{b}{ax} - b^2(x+a) \div a^2x^2$ .  
 5.  $x^2 - 2x = 2$ ,  $x = 3$ ,  $y = 2$ ;  $y = 53 \div 24$ , &c.      6. 4 miles, 3 do.  
 7.  $(x^n - 1) \div (x^2 - 1)x^{n-1}$ .      8.  $R^n \left( P - \frac{p}{r} \right) + \frac{p}{r}$ .      9. 7.

10  $1 + \frac{1}{1} + \frac{1}{1 \cdot 2} + \frac{1}{1 \cdot 2 \cdot 3} + \&c. = 2.71828$  approximately;

$\{(5x)^r \div 3^{r+1}\} \{1 \cdot 6 \cdot 11 \dots (5r-4)\} \div r.$  11.  $\frac{2}{3}$  and  $-\frac{7}{3}$ .

## XXVI.

2.  $a - \sqrt{ab - a^2}$ ; 4. 3.  $2, \frac{1}{2},$  or  $\frac{1}{2}(-3 \pm \sqrt{5})$ ;  $x + y + z = \sqrt{a^2 + 2b^2}$ ,  $\therefore z = c + \sqrt{a^2 + 2b^2}$ , &c.;  $\frac{1}{3}(-4 \pm \sqrt{76})$ .

4. 3. 7.  $p^2 \div 4 + r^3 \div 27 = 0.$  10.  $(1+x) \div (1-x)^2 - x^{n-1} \{3x - 1 + 2n(1-x)\} \div (1-x)^2$ ;  $n \div (15n+9), \frac{1}{15}$ .

## XXVII.

1.  $(lm - nk)^2.$  2.  $(a+b+c)(a-b)(b-c)(a-c).$

3.  $(a-b)(b-c) + (b-c)(c-a) + (c-a)(a-b)$ ;  $(a-b)^2 + (b-c)^2 + (c-a)^2.$  4.  $x = a \div (a^2 + b^2 + c^2),$  &c.

5.  $\sqrt{ab}$ ; 18 or -2;  $\frac{3}{4} + 3j.$  7.  $\frac{1}{2}(a+b)$  &c.;

$x = \{1 + a^2 - b^2 \pm \sqrt{(1-a-b)(1-a+b)(1+a-b)(1+a+b)}\} \div 2a$ ;  $x \div y = (1+a)(1-b) \div (1-ac),$  &c.

8. -8;  $(p^4 - 4p^2q + 8pr) \div (p^2 - 4q).$

9.  $(m+n) \div 2mn, (n-m) \div 2mn$

3c.

r  $\frac{7}{13}$ .4  $\frac{1}{2}$ .=  $c^2$ . $x^2$ .

3 do.

7.

**W. J. GAGE & CO'S,**  
**'List of Educational Publications,**  
**FOR SCHOOLS AND COLLEGES.**

---

**MATHEMATICS.**

**HAMBLIN SMITH'S ARITHMETIC.**—An Advanced treatise, on the Unitary System, by J. Hamblin Smith, M.A., of Gonville and Caius Colleges, and late lecturer of St. Peter's College, Cambridge. Adapted to Canadian Schools, by Thomas Kirkland, M.A., Science Master, Normal School, Toronto and William Scott, B. A., Head Master Model School for Ontario, 6th Edition..... PRICE  
\$0 75

**KEY.**—A complete Key to the above Arithmetic, by the Authors ..... 2 00

“I consider Hamblin Smith's Arithmetic by Kirkland and Scott, and Hamblin Smith's Algebra, with appendix by Mr. Baker, admirable works, far better adapted for use in our schools and for private study than any other similar works that I know of. They will soon, I have no doubt, supersede textbooks hitherto used in our schools, as they have already done in the Galt Collegiate Institute.”—*Alex. Murray, M.A., Mathematical Master Galt Collegiate Institute.*

**KIRKLAND & SCOTT'S ELEMENTARY ARITHMETIC.**—An Elementary treatise, on the Unitary System, intended as an introductory textbook to Hamblin Smith's Arithmetic, by Thomas Kirkland, M.A., Science Master Normal School, Toronto, and William Scott, B.A., Head Master Model School, for Ontario; 40th thousand *within first year of its issue* ..... 0 25

“Introducing Fractions immediately after the ‘Simple Rules’ will be hailed by all practical teachers as a step in the right direction. I shall advocate the exclusive use of your work in all elementary schools, as my past experience enables me to estimate its value.”—*John Macoun, M.A., F.L.S., Rector of Albert College Grammar School, Belleville.*

W. J. GAGE & Co's Educational Series.

	PRICE.
<b>McLELLAN &amp; KIRKLAND'S EXAMINATION PAPERS IN ARITHMETIC.</b> —A complete series of Problems, designed for use in Schools and Colleges, and especially adapted for the preparation of candidates for Teachers' Certificates, by J. A. McLellan, M.A., LL.D., Inspector of High Schools and Thomas Kirkland, M.A., Science Master Normal School, Toronto. 4th Edition.....	1 00
<b>McLELLAN &amp; KIRKLAND'S EXAMINATION PAPERS.—PART I.</b> —Containing the Examination Papers for admission to High Schools, and for Candidates for Third-Class Teachers' Certificates .....	0 50
<b>HINTS AND ANSWERS TO McLELLAN &amp; KIRKLAND'S EXAMINATION PAPERS,</b> containing answers to Problems and Solutions to all difficult questions. Prepared by the authors. 2nd Edition .....	1 00
<p>The leading American Educational Journal (<i>National Teachers' Monthly</i>) says of McLellan and Kirkland's Examination Papers:—"In our opinion, the best collection of problems on the American Continent."</p>	
<b>SMITH &amp; McMURCHY'S ADVANCED ARITHMETIC.</b> .....	0 50
<b>SMITH &amp; McMURCHY'S ELEMENTARY ARITHMETIC.</b> .....	0 25
<b>McLELLAN'S MENTAL ARITHMETIC.—PART I.</b> Containing the Fundamental Rules, Fractions, and Analysis. By J. A. McLellan, M.A., LL.D., Inspector of High Schools, Ontario. 2nd Edition.....	0 30
<b>McLELLAN'S MENTAL ARITHMETIC.—PART II.</b> By the same author, fully treats Percentage in its various applications, General Analysis, Stocks and Shares, Interest, Discount, &c., &c., and gives practical solutions of almost every type of question likely to be met with in any treatise on Arithmetic. 2nd Edition .....	0 45
<p>"His treatment of the subject has been so practical and skillful that teachers have frequently expressed the wish that he would prepare a text-book on Mental Arithmetic. The volume before us, Part I. of the work, treats systematically and comprehensively of the fundamental rules, fractions, analysis, reduction, &amp;c. It contains about 1,200 well graded practical problems. We can recommend the book to all teachers of arithmetic."—<i>London Advertiser</i>.</p>	
<b>JUVENILE MENTAL ARITHMETIC.</b> —By John F. Stoddard, M.A.....	0 15

**W. J. GAGE & Co's Educational Series**

	PRICE.
<b>HAMBLIN SMITH'S ALGEBRA.</b> —An Elementary Algebra, by J. Hamblin Smith, M.A., of Gonville and Caius Colleges, and late Lecturer at St. Peter's College, Cambridge, with Appendix by Alfred Baker, B.A., Mathematical Tutor, University College, Toronto .....	0 90
<b>KEY.</b> —A complete Key to Hamblin Smith's Algebra. 2 75 "Arrangements of subjects good; explanations and proofs exhaustive, concise and clear; examples for the most part from University and College Examination papers are numerous, easy and progressive. There is no better Algebra in use in our High Schools and Collegiate Institutes." —George Dickson, B.A., Head Master Collegiate Institute, Hamilton.	2 75
<b>HAMBLIN SMITH'S EXERCISE IN ALGEBRA. PART I.</b> .....	0 75
<b>GROSS' ALGEBRA.</b> —PART II. By E. J. Gross, M.A., Fellow of Gonville and Caius Colleges, and Mathematical Lecturer at Gerton College, Cambridge .....	2 50
<b>HAMBLIN SMITH'S ELEMENTS OF GEOMETRY,</b> containing Books I. to VI., and portions of Books XI. and XII., of Euclid with Exercises and Notes, by J. Hamblin Smith, M.A., &c., and Examination Papers, from the Toronto and McGill Universities, and Normal School, Toronto .....	0 90
<b>HAMBLIN SMITH'S GEOMETRY.</b> —BOOKS I. and II., with Exercises, &c.....	0 30
<b>HAMBLIN SMITH'S GEOMETRY.</b> —BOOKS II. and III., with exercises, &c.....	0 30
<b>POTTS' EUCLID.</b> —Containing the first six books with explanatory notes; a series of questions on each book; and a selection of Geometrical Exercises from the Senate, House, and College Examination Papers; with hints &c., by Robert Potts, M.A., Trinity College, with Appendix by Thomas Kirkland, M.A., Science Master Normal School. 500th thousand.....	0 50
<b>POTTS' EUCLID.</b> —BOOKS I. and II., with Exercises .....	0 30
<b>POTTS' EUCLID.</b> —BOOKS II. AND III. with Exercises .....	0 30
I shall recommend Pott's Euclid to the teachers in training as a book of invaluable use.— <i>W. Crockett, A.M., Principal Normal Training School, New Brunswick.</i>	

## W. J. GAGE & Co's Educational Series.

	PRICE.
<b>KIRKLAND'S STATICS.</b> —An Elementary Treatise on Statics. By Thos. Kirkland, M. A., Science Master Normal School, Toronto, with numerous examples and exercises; principally designed for the use of candidates for first and second class certificates, and for the Intermediate Examination. 4th edition .....	1 00
<p>“It supplies a great want felt by those preparing for Teachers' Certificates. This—did it possess no other merits—should make it a great success. It is by far the best text book on the subject for the schools of Ontario I have seen.” <i>Geo. Baytie, M. A., M. D., Science Master Normal School, Ottawa.</i></p>	
<b>HAMBLIN SMITH'S STATICS.</b> —ELEMENTARY STATICS. By J. Hamblin Smith, M. A., Gonville and Caius College, and late lecturer at St. Peter's College, Cambridge, with appendix by Thomas Kirkland, M.A., Science Master, Normal School, Toronto .....	0 90
<b>HYDROSTATICS.</b> —ELEMENTARY HYDROSTATICS. By J. Hamblin Smith, M.A., Gonville and Caius College, late lecturer at St. Peter's College, Cambridge. ....	0 75
<b>KEY.</b> —A Key to Hamblin Smith's Statics and Hydrostatics in one volume .....	2 00
<b>TRIGONOMETRY.</b> —ELEMENTARY TRIGONOMETRY. By J. Hamblin Smith, M. A. ....	1 25
<b>KEY.</b> A Key to Hamblin Smith's Elementary Trigonometry .....	2 50

## ENGLISH.

<b>MASON'S ADVANCED GRAMMAR.</b> —Including the principles of Grammatical Analysis. By C. P. Mason. B.A., F.C.P., fellow of University College, London. Enlarged and thoroughly revised, with Examination Papers added by W. Houston, M.A. 27th edition.....	0 75
<p>“I asked a grammar school inspector in the old country to send me the best grammar published there. He immediately sent Mason's. The chapters on the analysis of difficult sentences is itself sufficient to place the work far beyond any English Grammar hitherto before the Canadian public.”—<i>Alex. Sims, M.A., H. M. H. S., Oakville.</i></p>	

**W. J. GAGE & Co's Educational Series.**

	PRICE.
<b>MASON'S ENGLISH GRAMMAR.</b> —(Common School edition) with copious and carefully graded exercises, 243 pages .....	0 60
<b>MASON'S OUTLINES OF ENGLISH GRAMMAR,</b> for the use of junior classes.....	0 50
<b>ENGLISH GRAMMAR EXERCISES.</b> —By C. P. Mason. Reprinted from Com. Sch. Edition ..	0 36
<b>MILLER'S SWINTON'S LANGUAGE LESSONS,</b> (revised edition), adapted as an introductory text-book to Mason's Grammar, by J. A. Macmillan, B.A., Ottawa Collegiate Institute. It contains the Examination Papers for admission to High Schools, and teaches grammar and composition simultaneously. 5th edition, 40th thousand... 0 25	0 25
<p>"In accordance with a motion passed at the last regular meeting of the County of Elgin Teachers' Association, appointing the undersigned a Committee to consider the respective merits of different English Grammars, with a view to suggest the most suitable one for Public Schools, we beg leave to report, that, after fully comparing the various editions that have been recommended we believe that 'Miller's Swinton's Language Lessons' is the best adapted to the wants of junior pupils, and would urge its authorization on the Government, and its introduction into our Public Schools." (Signed),</p> <p>A. F. BUTLER, Inspector.            J. McLEAN, Town Inspector.            J. MILLAR, M. A., Head Master St. Thomas High School.            A. STEELE, B.A., Head Master Aylmer High School.            N. M. CAMPBELL, Head Master Co. of Elgin Model School.</p> <p>It was moved and seconded that the report be received and adopted. Carried unanimously.</p>	
<b>NEW ENGLISH GRAMMAR.</b> —In three parts: Etymology, Syntax and Analysis. By William Swinton. A. M. Revised by J. B. Calkin, M. A., Principal of the Normal School, Truro, N. S.....	0 50
<b>DAVIES' INTRODUCTORY ENGLISH GRAMMAR</b> .....	0 25
<b>DAVIES' NOTES ON 5TH READER.</b> —Literary Extracts selected from Book V of the authorized series of Readers, for "Examination in English Literature," of candidates for third class certificates, with notes original and selected. By H. W. Davies, D. D., Principal Normal School, Toronto. 5th edition .....	0 25
<b>MILLER'S ANALYTICAL AND PRACTICAL GRAMMAR</b> .....	0 36



**W. J. GAGE & Co's Educational Series.**

	PRICE.
<b>EPOCH SERIES.—PART I.</b> Containing first four of Series .....	\$0 50
<b>EPOCH SERIES.—PART II.</b> Containing last four of the Series .....	0 30
<b>EPOCH SERIES.—COMPLETE</b> , in one volume..	1 00
<b>CREIGHTON'S EPOCH PRIMER OF ENGLISH HISTORY.</b> —An introductory volume to "Epochs of English History." A complete summary of the history of England, in 140 pages. By Mandell Creighton, M. A., late Fellow and Tutor of Merton College, Oxford.....	0 30
"The work is admirably done, and it will no doubt obtain a very considerable sale."— <i>Athenæum</i> .	
"This volume, taken with the eight small volumes, containing the accounts of the different epochs, presents what may be regarded as the most thorough course of elementary English History ever published.— <i>Berdeen Journal</i> .	
<b>PINNOCK'S CATECHISM OF ENGLISH HISTORY</b> .....	0 10
<b>A SCHOOL MANUAL OF ENGLISH COMPOSITION.</b> For advanced classes in Academies, High and Public Schools. By William Swinton.....	0 45
<b>REID'S ENGLISH DICTIONARY</b> of the English language, containing the Pronunciation, Etymology, and Explanation of all words authorized by eminent writers; to which are added a vocabulary of the roots of English words and an accented list of Greek, Latin and Scripturo proper names. By Alexander Reid, A.M., Rector of the Circus-place School, Edinburgh; author of "Kudiments of English Composition," &c.; with an introduction by Henry Reid, Professor of English Literature in the University of Pennsylvania; and an appendix showing the pronunciation of nearly 3,000 of the most important geographical names. 3rd Canadian and 23rd English edition.....	1 00
"It is a very complete and useful work."— <i>Montreal Daily Witness</i> .	
<b>UNIVERSAL PRONOUNCING POCKET DICTIONARY OF THE ENGLISH LANGUAGE.</b> —Founded on the principles of Walker, Webster, Worcester, Johnston, Goodrich and Porter.....	0 20
<b>NATIONAL PRONOUNCING DICTIONARY</b> .....	0 40

**W. J. GAGE & Co's Educational Series.**

**GOLDSMITH'S TRAVELLER, AND GRAY'S ELEGY.**—In one volume; edited by the Rev. E. T. Stevens, M. A., Oxon, joint editor of "The Grade Lesson-books," "The Useful Knowledge Series, etc.;" and the Rev. D. Morris, B. A., London, author of "The Class-Book History of England," etc. Interleaved edition ..... 0 40

**SCOTT'S LADY OF THE LAKE.**—With introduction, notes, and glossarial index; by R. W. Taylor, M. A., Assistant Master at Rugby School, and formerly Fellow of St. John's College, Cambridge. Interleaved edition ..... 0 40


**MORRISON'S ENGLISH COMPOSITION.**—For the use of schools. By Thomas Morrison, M. A., Rector of the Free Church Normal School, Glasgow ..... 0 45


**CREIGHTON'S EPOCHS OF ENGLISH HISTORY.**—Edited by the Rev. M. Creighton, M. A., late Fellow and Tutor of Merton College, Oxford. Eight volumes in convenient and cheap form, adapted to Public and High Schools. Price 20c each.


**THE SERIES CONSIST OF:**

- I. Early England up to the Norman Conquest. By Frederick York-Powell, M. A. With four maps.
- II. England a Continental Power from the Conquest to Magna Charta, 1066-1216. By Louise Creighton. With a coloured map of the Dominion of the Angevin Kings.
- III. The Rise of the People, and Growth of Parliament, from the Great Charter to the Accession of Henry VII, 1215-1485. By James Rowley, M. A., Professor of Mod. Hist. and Lit., Univ. Coll. Bristol. With four maps.
- IV. The Tudors and the Reformation, 1485-1603. By the Rev. Mandell Creighton, M. A., late Fellow and Tutor of Merton College, Oxford, Editor of the series. With three maps.
- V. Struggle against Absolute Monarchy, from 1603 to 1688. By Bertha M. Cordery.
- VI. The Settlement of the Constitution from 1689-1788. By James Rowley, M. A., Professor of Modern History and Literature, University College, Bristol.
- VII. England during the American and European Wars, from 1789-1820. By O. W. Tancock, M. A., Assistant Master King's School, Sherborne, Dorset.
- VIII. Modern England, from 1820-1875. By Oscar Browning, M. A., Fellow of King's College, Cambridge.

"Amongst manuals in English History the Epoch Series is sure to take high rank."—*Daily Globe*.







MASON'S GRADUATED SERIES OF ENGLISH GRAMMARS.

**Mason's Outlines of English Grammar.**

By C. P. MASON, B. A., F. C. P., Fellow of University College London.  
Authorized for use of Schools in Ontario. For the use of junior classes.

Price, 45 Cents.

**Mason's Shorter English Grammar.**

With copious and carefully graded exercises, 243 pages.

Price, 60 Cents.

**Mason's Advanced Grammar.**

Including the principles of Grammatical Analysis. Enlarged and thoroughly revised, with Examination Papers added by W. Houston, M.A.

27th Edition, Price, 75 Cents.

"I asked a grammar school inspector in the old country to send me the best grammar published there. He immediately sent Mason's. The chapters on the analysis of difficult sentences is of itself sufficient to place the work far beyond any English Grammar hitherto before the Canadian public."—Alex. Sims, M. A., H. M. H. S., Oakville.

**English Grammar Practice.**

This work consists of the Exercises appended to the "Shorter English Grammar," published in a separate form. They are arranged in progressive lessons in such a manner as to be available with almost any text book of English Grammar, and take the learner by easy stages from the simplest English work to the most difficult constructions in the language.

Price, 30 Cents.

**Outlines of English Grammar.**

These elementary ideas are reduced to regular form by means of careful definitions and plain rules, illustrated by abundant and varied examples for practice. The learner is made acquainted, in modern measure, with the most important of the older forms of English, with the way in which words are constructed, and with the elements of which modern English is made up. Analysis is treated so far as to give the power of dealing with sentences of plain construction and moderate difficulty. In the

**English Grammar,**

the same subjects are presented with much greater fulness, and carried to a more advanced and difficult stage. The work contains ample materials for the requirements of Competitive Examinations reaching at least the standard of the Matriculation Examination of the University of London.

**The Shorter English Grammar**

is intended for learners who have but a limited amount of time at their disposal for English studies; but the experience of schools in which it has been the only English Grammar used has shown that, when well mastered, this work also is sufficient for the London Matriculation Examination.

## Examination Primer in Canadian History.

(History Taught by Topical Method.)

By JAMES L. HUGHES, Inspector of Public Schools, Toronto.  
A Primer for Schools, and Students preparing for Examinations. Price 25c.

### NEW AND SPECIAL FEATURES.

1. The History is divided into periods in accordance with the great national changes that have taken place.
2. The history of each period is given topically instead of in chronological order.
3. Examination Questions are given at the end of each chapter.
4. Examination Papers, selected from the official examinations of the different provinces, are given in the Appendix.
5. Student's Review Outlines, to enable a Student to thoroughly test his own progress, are inserted at the end of each chapter.
6. Special attention is paid to the Educational, Social, and Commercial progress of the country.
7. Constitutional Growth is treated in a brief but comprehensive exercise.

*By the aid of this work Students can prepare and review for Examinations in Canadian History more quickly than by the use of any other work.*

## Gage's Practical Speller.

A Manual of Spelling and Dictation. Price 30c.

### PROMINENT FEATURES.

The book is divided into five parts as follows:

PART I. Contains *the words in common use in daily life, together with Abbreviations, Forms, etc.* If a boy has to leave school early, he should at least know how to spell the words of common occurrence in connection with his business.

PART II. Gives *words liable to be spelled incorrectly* because the same sounds are spelled in various ways in them.

PART III. Contains *words pronounced alike but spelled differently* with different meanings.

PART IV. Contains a large collection of the *most difficult words in common use*, and is intended to supply material for a general review, and for spelling matches and tests.

PART V. Contains *Literary Selections* which are to be memorized and recited as well as used for Dictation Lessons, and lessons in Morals.

DICTIONATION LESSONS.—All the lessons are suitable for Dictation Lessons on the slate or in dictation book.

RENEWALS.—These will be found throughout the book.

