

THE
CANADIAN JOURNAL
OF
MEDICINE AND SURGERY

A JOURNAL PUBLISHED MONTHLY IN THE INTEREST OF
MEDICINE AND SURGERY

J. J. CASSIDY, M.D., EDITOR.

VOL. XX.
JULY TO DECEMBER, 1906

BUSINESS MANAGER
W. A. YOUNG, M.D., L.R.G.P. (LOND.),
145 COLLEGE ST., TORONTO, CAN.

1906

INDEX TO VOLUME XX.

CORRESPONDENCE.

Physicians' Fees for Life Insurance Examinations in the Province of Quebec	426
The Recent Fire at Queen's University, Kingston	277

EDITORIALS.

A Few Hours with the British Medical . . .	232
A Symposium of French Physicians on the Dangers from the Hypodermic Injection of Insoluble Salts of Mercury.	153
Editorial Notes	161, 270, 345, 420
Enthusiasm as a Cure	418
"Epistaxis, Prophylaxis"	351
Fees Paid to Physicians for Life Insurance Examinations	416
"Good Morrow to You.—Here, the Street Is Narrow"	419
Inaugural Address by Sir A. E. Wright, at the Opening of the Twentieth Session of the Medical Faculty of the University of Toronto	346
Open Meeting of the Toronto Medical Society	348
Our British Medical Association Number.	57
Personals	166, 237, 276, 358, 425
Relative to the History of the Constitution of the British Medical Association	58
Seventy-Fourth Annual Meeting of The British Medical Association at Toronto.	227
Some of the Leading Facts in the History of the British Medical Association	72
The <i>British Medical Journal</i>	70
The Canadian North-West, Its Climatology and Opportunities	90
The Committees of the British Medical Association	66
The Fifteenth International Medical Congress at Lisbon	157
The Nature and Treatment of Hysteria	264
The Recent University Senate Elections	350
The Report of the Registrar General of Ontario for 1904—Tuberculosis in Ontario	267
The Results of Examining Bologna Sausages and Potted Meats	413
The Treatment of Hemiplegia	261
"To See Ourselves as Others See Us"	352
Whole Milk, Skim Milk, Buttermilk and Cream, 1906	343

LARYNGOLOGY, RHINOLOGY AND OTOTOLOGY

Electrolysis in Eustachian Stenosis	409
Ludwig's Angina	410

NEWS OF THE MONTHS.

A Creditable Record	278
British Medical Association Exhibits	280
Final Examinations Ontario College Physicians and Surgeons	282
Items of Interest	170, 238
Medical Council Elections	420
New Staff for the Hospital for Sick Children	281
Ontario Medical Council in Annual Session	167
Ontario Medical Library Association	279
Queen's University Visited by \$70,000 Fire.	169
The New Ontario Provincial Board of Health	360
The Work of the New York Post-Graduate Medical School and Hospital	427
Trinity Meds. Dine	279

OBITUARY

Death of Dr. Minerva M. Greenaway, Toronto	359
Death of Dr. John Matthew Lefevre	359
Death of Dr. Jas. Stewart, of Montreal	359

ORIGINAL ARTICLES.

Acute Ascending Paralysis—Landry's Paralysis. By A McPhedran, M.B.	113
Address in Medicine. By Sir James Barr, M.D., F.R.C.P., F.R.S.E.	239, 305
Addresses in Obstetrics. By W. S. A. Griffith, M.D., F.R.C.P., F.R.C.S.	173
A Series of Intestinal Anastomoses. By Thomas S. Cullen, M.B. (Tor.), Associate Professor of Gynecology, Johns Hopkins University	1
Combination Operation for the Cure of Inguinal Hernia. By F. N. G. Starr, M.B. (Tor.)	330
Penetrating Stab-Wounds of the Abdomen. By N. A. Powell, M.D., Toronto	238
Race Suicide and Christian Science. By James H. Richardson, M.D., Toronto	255
The British Medical Association—Toronto Meeting, August 21-25, 1906	36

The Employment of Physical Measures in the Treatment of Disease. By Charles R. Dickson, M.D., Toronto 322

The Hygiene of the Home. By J. J. Cassidy, M.D. 373

PROCEEDINGS OF SOCIETIES.

Exhibit Hall 21P

Preliminary Programme of the Seventy-Fourth Annual Meeting of the British Medical Association 122

The British Medical Association, Toronto Meeting 186

SCHOOL HYGIENE.

Children's Teeth 408

Medical Inspection of Schools in Montreal 407

Mental Overwork in Schools 404

School Nurses 408

SELECTED ARTICLES.

Abstracts 266

Antagonistic Action of Veronal and Morphine 340

Bromopin and Its Therapeutic Importance 395

Celestins—Grand-Grille—Hopital Springs. 333

Citrophen in Malaria 335

Horace Wells, Humanity's Greatest Benefactor, the Discoverer of Anesthesia.. 378

"Manna" from Harvard 400

Something About the World's Great Fraternal Assurance Society 336

The Treatment of Bright's Disease 339

The Treatment of Pains in Locomotor Ataxia 398

Uric Acid Diathesis—Report of a Successful Case. (Abstract) 397

THE PHYSICIAN'S LIBRARY.

A Compend of Operative Gynecology. By William Seaman Bainbridge, M.D. 295

A Guide to Urine Testing. By Mark Robinson, L.R.C.P., L.R.C.S.Ed. 435

A Manual of Bacteriology. By Herbert U. Williams, M.D 366

A Manual of Medicine. By Thomas Kirkpatrick Monro, M.A., M.D 361

A Manual of Midwifery. By Thos. Watts Eden 434

American Practice of Surgery. By Jos. Bryant, M.D., Albert H. Buck, M.D.. 432

Anesthetics. By J. Bumfield, M.D. 362

A New Edition of Morris' Anatomy 366

A Non-Surgical Treatise on the Diseases of the Prostrate Gland and Adnexa. By George Whitefield Overall, A.B., M.D.. 297

A Text-Book of Genito-Urinary Diseases. By Dr. Leopold Casper 431

A Text-book of Human Physiology. By Dr. Robert Tigerstedt 367

A Text-Book of Obstetrics. By Barton Cook Hirst, M.D. 430

A Text-Book on the Practice of Gynecology. By W. E. early Ashton, M.D., LL.D. 437

A Treatise on Surgery. By George R. Fowler, M.D 291

Beverly of Graustark. By George Barr McCutcheon 295

Caro-Hua Lee. By Lillian Bell 293

Clinical Lectures on Neurasthenia. By Thomas D. Savill, M.D.Lond 435

Comparative Otology—Surgical Pathology and Treatment of Diseases of the Ear. By Clarence John Blarc, M.D. 299

Eczema. By Samuel Horton Brown, M.D. 365

Elements of General Chemistry with Experiments. By John H. Long, M.S., Sc.D 370

Ellis's Demonstration of Anatomy. Revised and edited by Christopher Addison, M.D., B.S. (Lond.), F.R.C.S 298

Gastric Surgery. By Herbert J. Patterson, M.A., M.B., B.C. (Cantab) 363

Genito-Urinary Diseases and Syphilis. By Henry H. Morton, M.D 433

Green's Encyclopedia and Dictionary of Medicine and Surgery 287, 438

Handbook of Meat Inspection. By Dr. Robert Ostertag 290

Heart Disease and Aneurysm of the Aorta. By Sir William H. Broadbent, Bart., K.C.V.O., and John F. H. Broadbent, M.D., F.R.C.P 298

Indications for Operation in Disease of the Internal Organs. By Prof. Hermann Schlesinger, M.D 361

International Clinics 430

In the Van. By Price-Brown 433

Kiepe's Materia Medica and Therapeutics. By Edward J. Kiepe. 438

Lectures on Clinical Psychiery. By Dr. Emil Krapelin 288

Lectures on Midwifery for Midwives. By A. B. Calder, M.D 367

Lury of the Stars. By Frederick Palmer.. 294

Manual of Diseases of the Ear, Nose and Throat. By John Johnson Kyle, M.D. 304

Medical Jurisprudence, Forensic Medicine and Toxicology. By R. A. Witthaus, A.M., M.D 284

New Serum Therapy. By D. Montgomery Paton, L.R.C.S. and L.R.C.P. (Ed.) . . .	302	The Delineator for October	237
On Leprosy and Fish-Eating. By Jonathan Hutchinson, F.R.S., F.R.C.S., etc.	292	The Diseases of the Nose and its Accessory Sinuses. By H. Lambert Lack, M.D., (London). F.R.C.S.	304
Osborne's Introduction to Materia Medica and Pharmacology. By Oliver T. Osborne, A.M., M.D.	302	The Ear and its Diseases. By Seth Scott Bishop, B.S., M.D., LL.D.	365
Phlebitis and Thrombosis. By Warrington Haward, F.R.C.S. (Eng.)	369	The Healers. By Maartens Maartens.	369
Physiology of the Nervous System. By J. P. Morat	291	The Health-Care of the Baby—A Handbook for Mothers and Nurses. By Louis - Fischer, M.D.	300, 362
Portfolio of Demochromes. By Professor Jacobi, of Freiburg, in Breisgau	285	The Influence of the Mind on the Body. By Paul Dubois, M.D.	363
Progressive Medicine. Edited by Hobart Amory Hare, M.D.	289, 262, 363	The International Medical Annual	296
Prophylaxis and Treatment of Internal Diseases. By Frederick Forchheimer, M.D.	289	The Medical Annual	286
Rational Hydrotherapy. By J. H. Kellogg, M.D.	285	The Medical Annual Synoptical Index to Remedies and Diseases.	366
Some of H. K. Lewis' Publications	368	The Nature and Treatment of Cancer. By John A. Shaw-Mackenzie, M.D. Lond.	288
Student's Handbook of Operative Surgery. By William Ireland de C. Wheeler (Mod. B.A., M.D., (Dun. Univ.), F.R.C.S.	296	The Operating Room and the Patient. By Russell S. Fowler, M.D.	291
Study, Treatment and Prevention of Tuberculosis	303	The Practice of Pediatrics. By Walter Lester Carr, A.M., M.D.	436
Surgery, Its Theory and Practice. By William Johnson Walsham, F.R.C.S. (Eng.), M.B. and C.M. (Aberdeen)	368	The Practical Medicine Series. Edited by Gustavus P. Head, M.D.	293, 361
The American Illustrated Dictionary. By W. A. Newman Dorland, M.D.	436	The Ship-Surgeon Hand-book. By A. Vavasour Elder, M.R.C.S., L.R.C.P.	369
The Autotoxicoses: Their Theory, Pathology and Treatment. By Heinrich Stern, Ph., M.D.	299	Treatise on Diseases of the Skin. By Henry W. Stelwagon, M.D., Ph.D.	300
The Chemistry, Physiology and Pathology of Uric Acid, and the Physiologically Important Purin Bodies. By Francis H. McCrudden	286	Treatment of Gonorrhoea in the Male. By C. Leedham-Green, M.B.	301
		Year-Book of Pediatrics and Orthopedic Surgery By Isaac A. Abt, M.D., and John Ridlon, A.M., M.D.	431

The Canadian Journal of Medicine and Surgery

A JOURNAL PUBLISHED MONTHLY IN THE INTERESTS OF
MEDICINE AND SURGERY

VOL. XX.

TORONTO, JULY, 1906.

NO 1.

Original Contributions.

A SERIES OF INTestinal ANASTOMOSES.

BY THOMAS S. CULLEN, M.B. (TOR.),
Associate Professor of Gynecology, Johns Hopkins University.

In going over my cases of the last few years I thought it might be advisable to describe in detail those in which it was necessary to remove portions of the bowel. The number is somewhat limited, but each case offers several points of interest:

Secondary carcinoma of the small bowel, 1 case.

Primary carcinoma of the cecum, 2 cases.

Tuberculosis of the cecum with perforation, 1 case.

Carcinoma of the sigmoid flexure, 1 case.

Carcinoma of the sigmoid flexure, complicated by a large uterine myoma, 1 case.

Carcinoma of the rectum secondary to a primary growth in the right Fallopian tube, 1 case.

Rectal diverticula, with perforation and abscess, 1 case.

SECONDARY CARCINOMA OF THE SMALL BOWEL.

In the following case a loop of the small bowel had become adherent to a friable carcinoma of the ovary. The growth had invaded the intestinal wall and the slightest traction was sufficient to rupture it. The Connell interrupted suture was employed except for the last few sutures, where we used mattress sutures penetrating the peritoneal and muscular coats but not piercing the mucosa. To make doubly sure we reinforced with a running suture entirely around the bowel. As it was impossible to completely remove the carcinoma of the ovary, a large gangrenous

area being left behind and requiring drainage, we found it necessary to push the loop containing the anastomoses far over to the left among healthy loops; otherwise it would certainly have been infected by the necrotic and gangrenous tissue. As noted in the history the bowel gave no further trouble.

*Tentative diagnosis: Subperitoneal and intraligamentary myomata. Actual condition: Hydrosalpinx, adeno-carcinoma of the right ovary, involvement of the small bowel and marked extension to the bladder. Hysterectomy, partial removal of the cancerous growth, resection of a portion of the small bowel; temporary recovery.**

History.—On Jan. 25, 1904, I saw the patient, who was 48 years of age. Her menstrual periods had continued regularly until she was 44. Since then the flow had appeared every three or four months, and there had been a slight vaginal discharge. Two years previously she had passed a calculus, apparently from the left kidney.

Examination.—On vaginal examination I found the uterus half as large again as normal. Projecting from the fundus on the right side, and very prominent, was what appeared to be a subperitoneal myoma about 5 cm. in diameter. The right side of the pelvis was filled by a growth which apparently sprang from the uterus and filled the broad ligament. This growth in contour and consistence resembled a myoma.

Operation.—On opening the abdomen (Feb. 2) I found the uterus moderately enlarged. The supposed subperitoneal myoma proved to be a very tense hydrosalpinx, which was kinked forward, thus accounting for its prominence. The growth on the right side was a carcinoma of the ovary. It filled the broad ligament and had infiltrated the bladder wall. Attached to the cancerous mass was the omentum with a loop of small gut. As the gut at this point was markedly constricted, I attempted by gentle dissection to release it, but the bowel was so infiltrated by cancer that it commenced to tear and resection of a portion was imperative. It was decided that the only hope of even temporary relief would be hysterectomy with as thorough removal of the growth as possible. This was done, but a raw, green, offensive, cancerous area, fully 6 cm. in diameter, remained attached to the surface of the bladder. Three inches of the bowel were then resected and the ends united by means of the Connell suture, supplemented by the Lembert suture. The anastomosed bowel was then placed among healthy loops of gut as far removed from the necrotic area as feasible. The pelvis was drained through the vagina and abdomen. The patient recovered

*Extracted from the *Journal of the American Medical Association*, November 19, 1904.

promptly, but naturally still has a small abdominal sinus. We have employed a retention catheter continuously, as even its temporary removal was promptly followed by the signs of ascending renal infection. In November, 1904, the patient was in fairly good condition and had been entirely relieved of abdominal distension and cramps, to which she had been subject for some time prior to the operation.

In this case the clearly outlined subperitoneal nodule associated with the growth on the right side gave us a clinical picture very characteristic of multiple myomata, and this diagnosis was further strengthened by the healthy appearance of the patient. Some may doubt the wisdom of attempting any operative procedure in these cases, but in the liberation of the constricted and friable intestinal loop the bowel was opened, and then the more




FIGURE I.—SCHEME OF OPERATION IN GROWTHS OF THE CECUM.

1. Lateral anastomosis between the transverse colon and small bowel. 2. Section of ileum and closure of end. 3. Removal of the growth and closure of the end of the transverse colon. If the patient should suddenly collapse the operation may be abandoned at any one of these three steps.

radical procedure seemed to offer the best chance of temporarily relieving the patient. In this case an absolute diagnosis would have been impossible without opening the abdomen.

PRIMARY CARCINOMA OF THE CECUM.

We have operated upon two cases of this variety. One patient was 55 years of age, the other 56. In Gyn. No. 12197 the patient was greatly emaciated, had complained for months of straining in the lower abdomen and later had passed much blood. The tumor was easily palpable in the cecal region and operation was at first deemed out of the question. After a week's rest in the hospital, however, she had improved and at her earnest sollicita-

tion the operation was undertaken. The subsequent months of comparative comfort were certainly well worth the ordeal of the operation, and up to the last she never had the unpleasant and racking symptoms that had been present before the operation.

In case No. 12016 the patient had lost some weight but was still in fairly good condition. She had never had any bloody stools and complained of very little discomfort. It is sometimes difficult to understand why in the one case there was so much hemorrhage while in the second, apparently equally far advanced, there was never any loss of blood. In the latter the character of the growth may afford the explanation. It was a colloid carcinoma. The greater part of the growth had been converted into colloid material. Near the surface few blood vessels were present. Case 12197, on the other hand, was a typical instance of adenocarcinoma with small glands.

In one case we left a fistulous opening, in the other we closed without drainage. The latter method is, I think, the better procedure. In cases of carcinoma of the cecum it seems wiser to make the lateral anastomosis with Robson's or Moynihan's clamps first. If the patient be too weak, the subsequent steps of the operation can be omitted (Fig. 1). Is she be still in fair condition the growth is removed and the ends of the ileum and ascending colon can be closed.

Adeno-carcinoma of the cecum; great emaciation; lateral anastomosis between the ileum and transverse colon; resection of the diseased bowel: temporary recovery.

Gyn. No. 12197. Mrs. J. R., white, aged 56. Admitted to the Johns Hopkins Hospital, June 21, 1905. Discharged Aug. 2, 1905.

The patient's chief complaint is of weakness and exhaustion. She has never been strong. Six years ago she had general dropsy. Has been married 37 years. Has had five children, the youngest 25 years old. The menopause occurred five years ago. Two years ago the patient began to pass much mucus by the rectum and had a good deal of straining in the lower abdomen. She passed no blood. This condition persisted until four weeks ago when the movements became very dark and foul-smelling; there was never any bright blood in the stools. There has been rapid loss of weight and strength and a tender lump has recently been noticed in the right iliac fossa just above the crest. This has become increasingly tender and for the past week the exhaustion has been extreme. There have been no nausea, vomiting, or stomach symptoms of any kind. On examination I found the patient very much emaciated, of a sallow tint, the mucous membranes were pale and it was with great difficulty that she could walk. Just

above the crest of the ilium on the right side, extending into the right iliac fossa, a firm, irregular and very tender mass can be felt. This is apparently situated in the cecum or in the abdominal wall directly over the cecum. Extending upward from this is a tumor mass. When the patient came to me I told her husband, who is a physician, that it was useless to perform any operation, but that we could send her to the hospital for a week's rest prior to her going away. During the week she gained considerably but then had an intestinal hemorrhage and lost ground. She again improved to some extent and wished to have something




FIGURE II.—CARCINOMA OF THE CECUM.

The greater part of the picture is occupied by the crater-like growth with undulating walls. Its line of advancement in the ascending colon is indicated by *a*. Its encroachment on the ilium by *a'*. The pericecal fat is infiltrated by discrete nodules as seen by *b*.

done. It was only after a great deal of persuasion that we decided to do an exploratory operation, not for a moment deeming that it would be feasible to remove the growth. On June 18, 1905, she was very much improved in color and strength and her hemoglobin had increased to 60 per cent. The mass in the right iliac fossa was not nearly so tender as on admission.

Operation, July 5.—The tumor mass involving the cecum was found freely movable. No enlarged glands in the mesentery or in the omentum could be detected, nor was there evidence of peritoneal metastases. On account of the apparent limitation of the growth we decided to remove it. The mass was freed from the

peritoneum of the lateral wall to which it was adherent. Not knowing just how long the patient could stand the operation we divided the procedure into three steps. First we made a lateral anastomosis between the lower end of the ileum and the transverse colon. Next the ileum was cut across at a safe distance from the growth and the end turned in and closed. The third step consisted in loosening up the growth, severing the ascending colon above the growth and closing the colon. In this way we could have hurriedly concluded at any one of the three steps. The abdomen was closed without drainage. In freeing the tumor we had to be exceedingly careful, as the ureter lay directly beneath the tumor. The right kidney had been prolapsed and the edge of it also lay beneath the tumor. The kidney was in close contact with the tumor and helped to make the growth seem so large.

July 8.—The patient has done well since the operation; she has had no nausea nor vomiting since the first day. No distension. She is taking her nourishment well.

Several days after this she became exceedingly weak and it was thought that she could not recover, but she speedily regained ground and was discharged apparently well on Aug. 2.

Sept. 6.—The doctor wrote me: "I am glad to say that the patient has been home from the hospital five weeks to-day and has increased one pound a week in weight. Her appetite is good, in fact, better than for two years. Her complexion is fairer than for years. She is on her feet the greater part of the day. Takes breakfast in her room, but the other two meals she enjoys at the table with the family. Her bowels are all right. At times she has some abdominal soreness and swelling."

I saw the patient in November. Her general condition was good, but she had some soreness in the right side. On careful palpation we could still detect the sensitive and prolapsed right kidney, but there was no evidence of metastases at any point.

She grew a good deal weaker and died on Jan. 8, 1906, free from pain and perfectly conscious.

Path. No. 8823. The specimen consists of the cecum, of the surrounding fat and of the appendix. The entire mass is board-like in consistency. The appendix is practically normal in size and is glued down to the cecum and to the neighboring fat. The hollow cup of the cecum is surrounded by a dense wall varying from 1 to 3 cm. in thickness. The cavity presents a crater-like appearance and is 3 cm. in depth (Fig. 2). The tissue is dark and crumbly. The mucosa, where present, is dark in color. Projecting from the mucous membrane are large and small nodules of the growth. On one end of the section is normal mucosa belonging to the ascending colon, on the other a considerable flap of normal ileum.

Histological Examination.—The cavity is found to be lined by many glands which present a tree-like arrangement, the epithelium being one layer in thickness. This tissue shows a great deal of round-celled infiltration. The well-advanced parts of the growth are composed of quantities of glands closely packed together. These glands are small and in many places the epithelium has proliferated to such an extent that the gland cavity is completely filled. The cell nuclei are remarkable for their uniformity in size. The growth is a typical adeno-carcinoma which has extended far beyond the contour of the wall of the bowel. The outlook, of course, is unfavorable.

Adeno-carcinoma of the cecum with extensive involvement of the lymph glands; resection of the diseased bowel. Patient apparently well.

Gyn. No. 12016. Mrs. F. H., admitted to the Johns Hopkins Hospital, April 2, 1905. Discharged, June 1. The patient is a widow 55 years of age, white. Her family and previous histories are not important. She has had two children. Her present trouble began about three years ago with an attack of diarrhoea, loss of weight, and general ill health. During the past two years she has had several attacks of colitis. Repeated examinations of the stools have been negative. Abdominal palpation from time to time did not reveal anything. She has lost about 30 pounds in weight during the last year, but recently has gained some. She is quite anemic; red corpuscles 2,700,000, leucocytes 7,000, hemoglobin 40 per cent. She has had little or no pain but a general sense of soreness at short intervals. In the right iliac fossa Dr. Nathan R. Gorter noticed a slight thickening about three weeks ago. This has been growing since that time. Appetite poor, bowels regular, no bleeding from the bowel at any time. On careful palpation I was able to detect a distinct area of induration in the region of the cecum. This appeared to be 4 cm. in diameter, but was no index to the actual size of the growth.

April 3.—A long incision was made through the right rectus. A carcinoma was found involving the cecum and a small portion of the ileum and about half of the ascending colon. The bowel was freed and clamped above and below. A lateral anastomosis was then done by means of the Moynihan forceps. The free end of the ascending colon was closed, the end of the ileum brought out through the lower angle of the abdominal incision and the abdomen closed.

April 6.—The patient has been unable to retain any nourishment. The nausea continues. The bowels have moved, per rectum, several times. The free end of the ileum that was brought

out through the lower angle of the wound is sloughing off to some extent. There is no escape of fecal matter through it.

Nausea and vomiting continued at intervals for a week and there was at times free fecal discharge from the enterostomy wound. The patient gradually improved, and several attempts were made to close the fistulous opening, but the bowel was so much indurated as a result of fecal matter coming over it, that the sutures did not hold. The patient made a very satisfactory re-



FIGURE III.—PRIMARY COLLOID CARCINOMA OF THE CECUM.

Gyn. Path. 8490. In the lower part of the section healthy ileum is seen. In the upper part unaltered mucosa of the ascending colon. The lower margin of the growth is indicated by *a*. The extension in the ascending colon by *b*. The growth is very thick and projects in places fully 1.5 cm. into the lumen of the bowel. It presents a translucent appearance and shows very little breaking down except in the vicinity of *b*. This accounts for the absence of hemorrhage. *c* is a very large mesenteric gland. It was fairly riddled with the adeno-carcinomatous growth.

covery and was discharged from the hospital on June 1. There was, however, a slight fecal fistula.

Feb. 28, 1906.—The fistulous tract closed fully three months ago. The patient is in excellent condition and is able to go everywhere. She is in better health than for years. Of course the outlook is very unsatisfactory, considering the histological findings.

Gyn.-Path. No. 8490.—The specimen consists of the cecum, appendix and a small part of the ileum, also of several mesenteric lymph glands. The growth itself is approximately 10 cm. in length, 9 cm. in breadth and about 8 cm. in thickness. The outer surface is nodular and at several points rather friable. It looks waxy or gelatinous and at first sight would make one think that it was somewhat edematous. The enlargement, on careful examination, is found to be due to infiltration of the fat, especially in the vicinity of the appendix, by the nodular growth which here and there is granular. The walls of the cecum vary from 5 mm. to 1.5 cm. in thickness. The tissue has a gelatinous appearance and is somewhat transparent. In some places the growth is dirty and necrotic-looking. The line of junction between the growth and the ascending colon is sharply defined, the growth projecting about 8 mm. from the surface. The line of demarcation between the growth and the ileum is also sharply defined, but here the mucosa of the ileum is undermined. The largest lymph gland in the mesentery reaches 2.5 cm. in diameter.

On *histological examination* the mucosa at the edge of the growth is seen to be normal. As we approach the growth, however, it shows considerable small round-celled infiltration. It then ends abruptly and is replaced by the new growth, which also consists of glands. These glands, however, are large and small and not regular as we find in the normal mucosa. Their epithelium in many places has so proliferated that the gland lumen is obliterated. In other places large and small colonies of glands are seen. The nuclei of the gland epithelium are fairly uniform in size; some, however, are larger than usual and stain deeply. From the gland grouping one would not hesitate to make an immediate diagnosis of carcinoma. In other places the glands are exceedingly small and closely packed together. This is especially evident where the tissue is dense and surrounded by much small round-celled infiltration. At other points the glands are separated from the stroma by a colloid secretion, and in the outlying portions of the growth where the cancer has run wild this colloid material is so pronounced that the epithelium has almost entirely disappeared, apparently being converted into this colloid material. The growth has extended to the outer surface of the bowel and, as was noted at the operation, extended to the adjoining mesentery. Far out in the adipose tissue is a lymph nodule 4 mm. in diameter. Along its margin at two points are large areas of carcinomatous infiltration where the gland type is perfectly preserved. The large lymph gland has been given over almost entirely to the new growth and few if any lymphoid elements are to be detected except just along the margin of the nodule. The case is one of adeno-carcinoma of the cecum, in which the colloid-producing cells predominate.

TUBERCULOUS STRICTURE OF THE ASCENDING COLON.

The careful and exhaustive articles dealing on lesions of this character that have already appeared render it superfluous for me to enter into a detailed consideration of the subject. Those wishing to study the subject fully are referred to the interesting articles of Henri Hartmann and Pilliet,* and Reclus,† in the French; of Hofmeister,‡ Adolf Hartmann,§ and Gross,|| in the German, and of Lartigau,** in this country. Hofmeister has tabulated all the cases he could find in the literature, and his consideration of the subject is most thorough, while Baumgarten, through his students, Hartmann and Gross, has contributed not a little to the pathological aspect of this disease. The works of Lartigau and Hofmeister should be carefully read by all particularly interested in this class of cases.

Tuberculous ulceration of the intestine is relatively frequent, as evidenced by the findings at autopsy, but stricture of the lumen of the bowel following as a result of this condition is somewhat rare. Hofmeister says that Eisenhardt, in 1,000 autopsies on tuberculous patients, found intestinal lesions 566 times. In only 9, however, was there a more or less definite stricture of the bowel.

Tuberculous strictures of the bowel are usually single and situated at the ileocecal valve. The cecum is converted into a sausage-shaped mass, which is adherent, as a rule, posteriorly and occasionally laterally. The omentum, although at times adherent to the growth, is not as prone to engraft itself on the tumor as in cases in which appendicitis exists. The outer surface, while relatively smooth, may be studded by a few tubercles. At one point the gut shows a constriction, and usually around this the adipose tissue is very dense. Where the cecum is cut into, the mucosa frequently shows considerable alteration. It is sometimes studded with irregular or serpiginous tuberculous ulcers, while the intervening mucous membrane is the seat of a chronic inflammatory process. At the point of stricture the lumen of the gut is so narrow that the tip of the finger can hardly be introduced. In some cases so small is the calibre of the bowel that a sound is passed with difficulty, and in our case a small bird-shot was sufficient to completely occlude the canal. The degree of alteration in the cecum varies with the individual case, and it is only necessary for the reader to picture the tuberculous process advancing until the cecum becomes matted and densely adherent to all the neighboring structures, and, in rare instances, the process gradually involves the abdom-

* Note sur une variété de typhlité tuberculeuse simulant les cancers de la région. Bull. de la Soc. anat. de Paris, 1891, t. lxxvi, p. 471.

† Typhlité et appendicite tuberculeuses. Cliniques Chirurgicales de la Pitié, 1894, p. 317.

‡ Ueber multiple Darmstenosen, tuberkulösen Ursprungs, Beiträge zur klinischen Chirurgie, 1896, Bd. xvii, S. 577.

§ Ein Fall von tuberkulöser Darmstenose. Inaug. Diss., Tübingen, 1897.

|| Ueber Stricturirrende Darmtuberkulose. Inaug. Diss., Tübingen, 1901.

** Journal of Experimental Medicine, 1901, vol. vi., p. 23.

inal wall until finally there is a fistulous opening on the surface. Even in the early stages the mesenteric glands are enlarged and already involved in the tuberculous process, and where the cecal invasion is apparently in its incipency there may be caseation of these glands.

Tuberculous stenoses of the gut, when multiple, are almost invariably situated in the ileum. Anywhere from one to twelve strictures have been noted in the same patient. In one case Hofmeister found twelve strictures scattered over a distance of about seven feet of gut. The bowel between the strictures is frequently distended, and in rare cases has been known to reach 17 cm. in circumference. Lartigau draws especial attention to a group of these cases in which, associated with the tuberculous process, there is a marked diffuse thickening of the bowel wall, which occasionally reaches 1 cm. or more in thickness.

The appendix is usually adherent, but, except where the tuberculosis of the cecum is far advanced, shows no implication in the specific process. Our case proved no exception to the rule. Although bound down by adhesions, the appendix was otherwise normal.

Histological Picture.—In sections from the cecum the edges of the ulcers may show tuberculous tissue, but, as a rule, epithelioid cells or typical tubercles are wanting, and nothing but granulation tissue can be made out. In the vicinity of the muscle, however, groups of epithelioid cells, and now and then tubercles, are seen. The peritoneal surface is usually free from tuberculous nodules until the disease is far advanced or unless the cecal lesion has been associated with tuberculous peritonitis. Sections from the stricture are composed entirely of connective tissue; sometimes with, at other times without, areas even slightly suggestive of tuberculosis. The adipose tissue surrounding the gut at the point of stricture is much infiltrated with small round cells, rendering the fat exceedingly hard and firm. Sections from the lymph glands in the region of the cecum almost invariably yield typical tubercles.

Naturally the tuberculosis gradually extends to the muscle and outer coats of the bowel. The farther away the process extends from the lumen of the bowel, the more characteristic will be the specific lesions, since the inflammatory changes produced by the intestinal bacteria have less opportunity of masking the tubercles. The diffuse thickening or "chronic hyperplastic tuberculosis" of the intestine yields a picture very different from that of simple tuberculosis, as has been clearly pointed out by Henri Hartmann, Lartigau, and others. In these cases the tuberculous process has been relegated entirely to the background, while the mucosa and muscle have been overrun with round cells. Intestinal bacteria

have doubtless gained entrance to the walls through the tuberculous lesions and have continually kept up a chronic inflammation of the bowel wall so widespread in character that the tuberculosis is entirely overshadowed. At a few points, however, it will still be demonstrable, and can be detected with certainty in the mesenteric lymph glands. Even in the cecal wall, when the typical lesions are totally wanting, tubercle bacilli can still be readily demonstrated.

Clinical History.—Patients presenting tuberculosis of the cecum are usually between twenty and thirty years of age. The condition, however, may be found in the very young, and has been noted in persons fairly advanced in years. Quite commonly the patient has suffered from an old tuberculous process in the lungs or has a suspicious family history. In many of the cases which have come to autopsy healed lesions in the lungs have been demonstrated, while in a few instances there has been swelling of the cervical, axillary, or other lymph glands coincident with the cecal lesion. One of the first symptoms is constipation. After a time dull or sharp pain is felt in the appendiceal region. As the constriction develops there may be an intermittent diarrhea, with the gradual narrowing of the bowel, and fulness may be noted over the cecum. Where there is much infiltration of the intestinal wall the gut becomes very firm and feels like a sausage-shaped tumor. With the gradual growth of tuberculous tissue and narrowing of the bowel symptoms of obstruction manifest themselves, as evidenced by abdominal distension, colicky pain, marked peristalsis, vomiting, and rapid loss in weight.

But although these symptoms may be present, in some instances definite indications of the presence of the lesions may be entirely absent. In our case the patient felt well until the day before operation, complaining only of slight discomfort near the appendix.

Diagnosis.—With the increased attention paid to cecal tuberculosis the possibilities of overlooking these lesions will be lessened. It was only a few days after our case was operated upon that Dr. Finney saw a patient giving symptoms sufficiently suggestive of a tuberculous lesion in the cecum to render such a diagnosis justifiable. At operation the cecum was found to be the seat of a most extensive tuberculous ulceration. Fortunately, it was found possible to excise the whole of the diseased area.

Given a tumor in the right iliac fossa, of slow growth, a clinical history pointing to a previous pulmonary tuberculosis, and a comparative absence of temperature, it is highly probable that tuberculosis is present. If a patient be fairly well advanced in years, of course, the possibility of a malignant growth must be considered. As pointed out by Hartmann, Lartigan, and other

authorities, tuberculosis of the cecum, especially of the hyperplastic form, has often been taken for sarcoma. This has been due to the massive infiltration with small round cells. But provided that we remember that they form a definite infiltration, instead of one or more large foci, and further, that the cells are uniform in size instead of being large and small and actively dividing, confusion is not likely to occur.

The gross diagnosis between tuberculosis and carcinoma of the cecum may offer numerous difficulties, but on microscopic examination no confusion can exist, as in the tuberculous process the epithelial elements play an entirely passive role or have disappeared. Moreover, the demonstration of the tubercle bacilli is generally easy.

The diagnosis between cecal tuberculosis and appendicitis is usually dependent on the tuberculous history and the slow growth of the tumor, together with the absence of a temperature suggestive of a pus accumulation. Of course, in a case similar to the present one, a differential diagnosis would be absolutely impossible.

Treatment.—If tuberculosis of the cecum be diagnosed early operation is indicated. Resection of the entire diseased area is, of course, necessary for an absolute cure. Lateral anastomosis between the ileum and ascending colon is the ideal operation. If after resection of the diseased portion of the gut very little mobility be obtainable, in order to avoid tension an end-to-end anastomosis is the only alternative. Where there are numerous strictures scattered over an area of several feet of gut, the question arises as to whether the entire diseased area should be excised or several anastomoses be made, removing only the diseased segments and leaving the intervening normal gut. If the span of gut involved by the tuberculous process be not over three or four feet, it is wiser to remove this portion in its entirety. In one of the cases reported six or seven feet were removed, and the patient recovered. With the diseased cecum it is always necessary to carefully examine the glands of the mesentery, and if they be involved, they too should be excised. The results from resection have been very gratifying, Hofmeister in his table of 83 operative cases showing a recovery of 62 per cent.

*Tuberculous stricture of the ascending colon, with sudden total obstruction of the bowel; perforation of the intestine; removal of the cecum and half the ascending colon. Recovery.**

The following is taken from my case-book, November 29, 1902: At 11 p.m. I saw, in consultation with Dr. Charles E. Simon, Miss K. G., aged twenty-four years. The day before she

had had indefinite pains in the region of the appendix. They were, however, not very severe and lasted but a short time. To-day she did her work as usual and prepared supper, but shortly afterward was taken with severe pain in right side and was forced




FIGURE IV.—TUBERCULOSIS OF THE CECUM WITH PERFORATION.

Above is a cross-section of the ascending colon. Below and to the right the ileum. At a point directly opposite the ileum is a perforation of the cecum, and just above the perforation the adipose tissue is thickened and there is a constriction of the gut. At *a* are two enlarged and tuberculous lymph glands. (For the interior view of the specimen, see Figures 5 and 6.)

to go to bed. At 9 p.m. Dr. Simon saw her. There was marked rigidity of the right rectus over the appendiceal region. There was little temperature. On examination of the blood Dr. Simon noted that all eosinophiles had disappeared and that there was

an evident leukocytosis.* When I saw her two hours later the rigidity of the right side had in part disappeared, probably as she was slightly under the influence of morphine. The general condition was good; pulse full and regular. Nevertheless, I advised immediate operation.

At 1.30 a.m. the abdomen was opened and a thin, watery pus immediately escaped from the peritoneal cavity, and the pelvis was found to be completely filled with pus. The intestinal loops, however, on the whole, presented a fairly normal appearance. Here and there they were covered by a few flakes of fibrin. The appendix was easily recognized and was bound down by adhesions. It was tied off from tip to base. As the distal extremity appeared to be normal, we expected to find a perforation near the cecum, but on complete removal of the appendix it was found that, apart from adhesions, no alteration was present. After removing the pus from the abdomen a sponge was passed into the right renal pocket to see if any pus was there, and, to our surprise, some dark fluid escaped. This was entirely different from that found in the pelvis. The abdominal incision was continued upward to the ribs, and we immediately saw a perforation, about 4 mm. in diameter, in the ascending colon. As there was a good deal of fluid escaping, I temporarily closed this fistulous opening with a purse-string suture. I then drew the ascending colon out and made a longitudinal incision, and on introducing the finger into the colon found total obstruction a short distance above the ileocecal valve. The lower third of the ascending colon, the cecum, and a small portion of the ileum were tied off and removed, together with some enlarged glands in the mesocolon. The ascending colon and ileum were then united by end-to-end anastomosis. Lateral union would have been preferable, but we had no choice, as the tissues would have been on too great a tension. A Connell suture was employed for two-thirds the circumference of the gut, the remaining third being turned in with rectangular mattress sutures. The entire line of suture was reinforced by running mattress sutures. The pelvis was carefully sponged out, the intestinal loops were brought up into the abdomen, and the entire pelvis was loosely packed with iodoform gauze.†

A gauze drain was left at the site of the anastomosis. The patient stood the operation well. Her pulse did not rise above 100. The outlook, however, was not particularly flattering, con-

* Simon lays much stress on the frequent absence of eosinophiles where pus is accumulating and thinks that this sign is of more practical value than the degree of leukocytosis.

† For several years, where the pelvis has been filled with free pus, I have made it a practice, after having wiped the pelvis and intestines off, to place the patient for a moment in the Trendelenburg posture. The pelvis has then been loosely but fully packed with gauze, the ends of which are brought out through the appendix incision. My object has been to prevent the intestinal loops from dropping down and becoming adherent or kinked in the pelvis. In my hands this procedure has yielded very gratifying results. The loops, although still liable to become adherent, are on a level and are not nearly so prone to become obstructed.

sidering the fact that there was a commencing peritonitis and also considerable edema of the intestinal wall. Eight days after operation, on removal of the last of the gauze, some fecal matter was found on the dressing. The fistula gradually closed, and the patient made an excellent recovery.

February 12, 1904.—The patient has been at work for several months, performing general household duties without the slightest




FIGURE V.—COMPLETE OBSTRUCTION OF THE ASCENDING COLON WITH PERFORATION ON THE PROXIMAL SIDE OF THE STRICTURE.

The walls of the colon are greatly thickened and the narrowed lumen is completely obstructed by a small bird-shot. Opposite the ileo-cecal valve is the small perforation and at the lower end of the picture the inverted appendix stump is seen.

inconvenience. Her general condition is excellent. From her I learned that she had had typhoid (?) fever six years previously and was in bed for two weeks. For the last year she has had cramp-like pains throughout the abdomen two or three times a month, and recently the bowels have been more constipated than usual. She gives no history whatever of injury or bruising of the abdomen. For about a week before her admission to the hospital

she had had intermittent abdominal pain. From the family history we were unable to get any data suggestive of hereditary tuberculosis.

March 1, 1906. The patient is now in excellent health.

Pathological Report.—Gynecological-Pathological No. 6316.) The specimen consists of a small portion of the ileum, of the cecum, and of about one-half of the ascending colon. The mucosa of the ileum is unaltered, that of the cecum in most places is normal, but at a point directly opposite the ileocecal valve is a perforation 5 mm. in diameter (Fig. 4). The walls of the perforation are rather smooth and the surrounding mucosa, over an area 1 cm. in diameter, is somewhat thickened. The ascending colon, about 5 cm. above the perforation, shows a marked constriction. At this point the lumen narrows down until it is not more than 2 mm. in diameter. Indeed, so small is it that a fine bird-shot would lodge and completely plug the canal at this point (Figs. 5 and 6). The intestinal wall at the point of constriction varies from 5 mm. to 8 mm. in thickness and is exceedingly firm in consistency. The constriction is 1 cm. in length and the ascending colon above this point is unaltered.

Histological Examination.—The appendix, beyond showing a few adhesions on its outer surface, is normal. The cecum in the vicinity of the perforation has entirely lost its glandular elements, the specimen consisting almost entirely of granulation tissue. The underlying muscle shows a varying amount of small round-celled infiltration. This is especially abundant in the vicinity of the peritoneal covering.

Along the margin of the perforation there is also much granulation tissue, and the underlying muscle is everywhere infiltrated by small round cells. The ulceration is evidently an old process, as nowhere is a very acute inflammatory reaction present. The walls of the stricture are, to a great extent, composed of fibrous tissue. Here and there we have some light areas somewhat suggestive of tuberculosis. No giant cells are, however, demonstrable. Several mesenteric glands were removed with the intestine. Some of these reached 1.5 cm. in diameter. On histological examination they show typical tubercles, some sections of which contain four or five giant cells. The tuberculous process in the lymph glands has here and there advanced to caseation.

The following points merit attention in this case:

1. The total absence of definite symptoms until a few hours before operation.
2. The presence of symptoms identical with those of acute appendicitis.
3. Marked contraction of the stricture.

4. The advisability of always exploring the right renal pocket in all cases in which there is free purulent fluid in the pelvis.

As seen from the history, the patient had practically no symptoms until about five hours before operation, and then there was moderate pain over the appendix, accompanied by rigidity of the right rectus.

Examination of the blood showed a total absence of eosinophiles. The only way in which we can account for the lack of symptoms is that for some reason there occurred an acute contraction of the stricture, which, up to that time, had permitted the




FIGURE VI.--TUBERCULOUS STRICTURE OF THE ASCENDING COLON WITH PERFORATION OF THE CECUM.

Directly opposite the ileocecal valve is a small perforation with slightly ragged edges. A short distance above this point the intestinal walls grow thicker and then form an annular constriction. The lumen of the ascending colon at the stricture has been so narrowed that a small bird-shot, when introduced, lodged therein and completely plugged the gut.

free passage of feces. The possible existence of such a condition supplies another indication for early operation whenever trouble exists in the appendiceal region. Already peritonitis had developed, although the symptoms had existed for so short a time; and had we delayed until morning there would have been little chance of saving the patient.

After having removed the appendix and wiped the pus from the pelvis, the abdominal cavity appeared normal, and I probably should not have explored the right renal pocket had I not been familiar with the renal work of Max Broedel, who has shown

clearly that where there is a free accumulation of fluid in the region of the appendix, by gravity it will travel down into the right renal fossa.

I should have preferred a lateral anastomosis, but we were forced to make an end-to-end union on account of tension.

CARCINOMA OF THE SIGMOID FLEXURE.

We have had one uncomplicated carcinoma of the sigmoid flexure. The patient was 53 years of age and, when I saw him, was very weak. He had lost much in weight and toward the last had had copious hemorrhages several times a week. We had no alternative but to make an end-to-end anastomosis. As noted in the history he succumbed on the tenth day, but there was no evidence of peritonitis. He was too weak to stand the strain and developed cardiac depression similar to the attack of a year previous, at which time his life had been despaired of.

Our second case of carcinoma of the sigmoid flexure was accidentally discovered during a hysterectomy for a large myomatous uterus which was firmly wedged in the pelvis. The intestinal obstruction was thought to be caused by the myoma. There had been no symptoms on which one could definitely base a diagnosis of carcinoma. In this case the patient returned after several weeks and died from peritonitis in the right upper abdominal quadrant, a point far removed from the site of the anastomosis. The autopsy also clearly demonstrated that metastases were freely scattered throughout the abdomen, and further that we had not entirely removed the original growth. A more extensive operation would not, however, have been feasible.

Adeno-carcinoma of the sigmoid flexure; resection of the diseased area, end-to-end anastomosis; death on the tenth day.

Dr. A. G. W. This patient has been failing for nearly two years. First he noticed that he was losing in weight, but was able to go around and do his work fairly well. He was very easily fatigued; could not do as much night work as before; had been under treatment for some time with apparently no relief. When I saw him there had been rectal hemorrhages for over a year. On careful palpation we were unable to detect any growth in the abdomen, and on using a short proctoscope could make out nothing. As he continued to lose greatly in strength we sent him to the mountains, deeming that he could not stand an operation. He improved slightly, but soon again lost ground as a result of the frequent hemorrhages. Finally he was admitted to the hospital, and we decided to make an exploratory operation, remembering, however, that his heart a year previous had given rise to

such alarming symptoms that it was thought he would succumb. After entrance the patient improved slightly, but the hemorrhages continued.

Sept. 22, 1904.—We made a median incision and examined the appendix. This we found twice its natural size and partly filled with fecal matter. It was removed. We then carefully examined the intestines and found hard masses throughout the transverse colon. These proved to be fecal concretions. In the pelvis was a hard mass which on pressure proved to be a malignant growth involving the bowel just below the brim. No glandular enlargement could be detected, but here and there fine white lines—evidently dilated lymphatics—were seen passing down the meso-sigmoid. We carefully walled off the abdominal cavity and also the abdominal incision, clamped above and below the growth, and after removing the growth did an end-to-end anastomosis with a Connell suture for three-fourths the circumference of the bowel. The remaining portion was closed in with mattress sutures. Posteriorly the lower portion of the sigmoid on the right was rather thin and there was just the faintest possibility that there might be a subsequent leakage. Everything, however, looked perfectly solid. On account of the fecal concretions, we brought up a loop of the descending colon into the left inguinal region. This loop was opened the same night.

Sept. 29.—The patient since operation has had a practically normal temperature, but on one or two occasions it ran up to 100° F. His pulse has been fairly good. During the entire time there has been a good deal of nausea, but no vomiting. He has had an ice bag over his stomach, which has been exceedingly sensitive. At no time has there been any distension. The bowels moved thoroughly after calomel and magnesia. This morning at 12.30 he woke up in a profuse perspiration. His pulse was almost imperceptible, although an hour before he had been in excellent condition. I examined him between two and three o'clock. The pulse was not demonstrable either in the facial or in the radial region. He was given strychnin and digitalis. He soon lost consciousness, was very restless, and died at 4 a.m. We had here a definite cardiac syncope. He retained his nourishment from the beginning. It may be noted that a year ago he had a similar attack, and on that occasion his heart's action became so weak that he was not expected to rally.

Gyn.-Path. No. 7786. The specimen consists of six inches of sigmoid flexure (Fig. 7). Outer surface of bowel looks fairly normal except for a slight bulging. On palpation it is found to be very firm and gristle-like. On examination two distinct and separate growths can be detected. One is 4 cm. in diameter, the other 5 cm. Each one has raised edges and is sharply circum-

scribed. The growth extends on an average about 5 mm. from the surface, but in some places projects at least a centimetre into the cavity. While the edges are markedly raised the central portions present depressions. The growths are rather porous in




FIGURE VII.—CARCINOMA OF THE SIGMOID FLEXURE.

Path. No. 7786. The specimen shows two distinct foci of carcinoma. *a* and *b* each has a depression in the centre with wavy elevated margins sharply circumscribed from the surrounding healthy bowel. They are separated from one another by an interval of at least 1 cm. of healthy mucosa.

appearance. The surrounding mucosa looks perfectly normal. Sections through the growths show that they are typical adenocarcinomata. All resemblance to the bowel mucosa has, however, entirely disappeared. The growths show irregular invasion of

the submucosa and of the muscular layer of the bowel. There is considerable small round-celled infiltration.

Diagnosis.—Adeno-carcinoma of the sigmoid flexure.

Acute intestinal obstruction; large myoma wedged in the pelvis; non-suspected adeno-carcinoma of the sigmoid flexure; hysterectomy; resection of the diseased bowel; end-to-end anastomosis. Temporary recovery.

Gyn. No. 12000. E. S., colored, aged 40. Admitted to the Johns Hopkins Hospital, March 26, 1905; discharged, June 9. I saw this patient in consultation with Dr. Clement A. Penrose. On admission she was suffering from intestinal obstruction. This was thought to be caused by a myoma which had been known to exist for fifteen years. The family and previous history was negative. The menses began at 16, were always regular, and caused a great deal of pain; the flow was excessive. Twelve years ago she had a severe attack of abdominal pain. This was sharp and shooting in character, but there was no intestinal obstruction. For the past six weeks, beginning at the time of a menstrual period, she again noticed sharp, shooting pains in the abdomen. These were intermittent and practically limited to the left side. The bowels have not moved for several days, and the pains have been spasmodic, occurring at intervals of four to five minutes. She does not think that she has had any fever. There had been no blood in the stools before the obstruction. On the morning of her admission she vomited a small amount. On admission the lower abdomen was found distended by a mass. The tumor reached as high as the umbilicus on the left side, presenting a large nodule which pressed down in the left iliac fossa. A similar and smaller nodule was present in the right iliac fossa. Around the umbilicus peristaltic movements were marked, and were accompanied by loud gurgling in the intestines. Tympany was marked everywhere except just over the nodules and above the symphysis. The abdomen was opened at once. A myomatous uterus was found reaching as high as the umbilicus. Springing from the region of the right cornu was a pedunculated tumor about 10 cm. in diameter. There were no adhesions above and the appendages appeared to be normal. As the tumor was free above, but could not be easily lifted, we suspected an intra-ligamentary growth or inflammatory adhesions in the pelvis. The round ligaments on both sides were tied and the ovarian vessels controlled. The enucleation was begun from left to right. The uterus was amputated through the cervix, but its removal was accomplished with a great deal of difficulty owing to the broad cervical attachment.

After removal of the uterus the rectum was found to contain

a growth which seemed to be malignant in character and was adherent to the cervix posteriorly (Fig. 8). The rectal tumor was loosened as carefully as possible from the cervix; it lay entirely below the brim of the pelvis. There was a nodule apparently about 7 cm. in diameter and the intestines for a length of 8 cm. were involved. The general peritoneal cavity was again carefully walled off and the bowel clamped above and below the diseased area. After removal of the growth an end-to-end anastomosis was done. Three-fourths of the bowel was closed by




FIGURE VIII. BROAD-BASED MYOMATOUS UTERUS FILLING THE PELVIS
CARCINOMA OF THE SIGMOID FLEXURE.

The multi-nodular uterus is very broad-based, rendering the hysterectomy difficult. *a* indicates the uterine cavity. The myoma *b* contained many cancerous areas. Occupying the sigmoid flexure is the carcinoma *c*. This almost completely occluded the bowel. Its upper limits are indicated by *d*, its lower by *d'*.

Connell sutures, the remaining one-fourth by mattress sutures. The entire suture line was reinforced by continuous sutures. The posterior vaginal fornix was punctured and the pelvis packed with one strip of iodoform gauze. The anastomosis was very satisfactory and the condition of the bowel good. The growth was very low down, thus rendering anastomosis difficult. It was, however, too high up to permit removal through the anus. In order

to give the anastomosis a complete rest a left inguinal colostomy was done, the descending colon being sutured to the peritoneum, and opened later on in the evening. As we found it very difficult to get a good exposure for the anastomosis we incised the abdominal wall transversely, making an incision three inches long extending through the left rectus. We were thus enabled to greatly facilitate the operation and save much time. The liver and the omentum were free from nodules. The patient was returned to the ward in a very weak state, but in fairly good condition considering the severity of the operation. Her temperature at that time was 101.5° F. For several days after operation the patient was very restless and it was difficult to keep her quiet. She was continually trying to remove the binder. She gradually improved, however, and on May 13th an attempt was made to close the fecal fistula. In this, however, we were not successful, as when she left the hospital, on June 9, there was still a slight fecal discharge from the fistulous tract. She seemed to be in very good condition. The bowels moved well; there was little pain, but some tenderness over the region of the anastomosis. She was gaining in weight and strength.

Gyn.-Path., No. 8447. E. S. The specimen consists of a large myomatous uterus, of both tubes and ovaries and of a portion of the sigmoid flexure. The myomatous uterus has been amputated through the cervix. It is 16 cm. in length, 12 cm. in breadth and 11 cm. in its antero-posterior diameter. Attached to the surface are several interstitial and one pedunculated myomata. The pedunculated nodule is rough, oblong in shape, 7 cm. in its longest diameter. The undercut surface is 10 cm. in diameter (which would account for the difficulty encountered in the enucleation). On section many myomata are seen scattered throughout the walls.

Our chief interest is centred in the section of the sigmoid flexure. This is 9 cm. in length. The outer covering of the bowel looks fairly normal except for some slight whitish elevations. Occupying the entire thickness of the bowel near the centre is a hard, light-colored growth (Fig. 8). This is 4 cm. in length and extends throughout the entire thickness of the bowel. The growth itself with the indurated adipose tissue surrounding it is fully 3 cm. in thickness.

Histological examination shows in some places perfectly normal mucosa surrounded on either side by colonies of small glands. In some of these colonies the epithelium is so proliferated that the gland arrangement is lost. The epithelial cells of the new growth are much smaller than those of the normal epithelium. They stain more deeply and some of them are rather large. On the whole, however, they are of uniform size. The muscular coat is involved and here the nests of cells are much denser. They are,

however, in many places surrounded by fibrillated tissue that takes the hemotoxylin stain and resembles mucin. The growth has extended to the outer surface of the bowel, but the chief thickening here is due to new connective-tissue formation in the fat.

The picture is one of typical adeno-carcinoma of the rectum.

Gyn. No. 12204. The patient was again admitted on June 24, 1905. At the seat of the former colostomy was a small sinus just admitting a probe. There had been no fecal discharge from this for several days. Until a week previously she had been in good condition. The bowels became constipated, there were frequent attacks of pain in the abdomen and during the last seven days there had been no movement. For the last two or three days the pains had increased in severity, but there had been no vomiting. Her temperature and pulse were normal. The abdomen was slightly distended. No peristaltic movements were visible. Enemata were ineffectual.

On June 25 the bowels moved spontaneously. On June 27 considerable vomiting occurred and distension was noted; there was great tenderness on palpation in the right upper quadrant. On June 28 enemata were given and there was some fecal discharge through the wound. The distension, however, continued and the vomiting persisted. On July 3 patient was taken to the operating room, as the condition had become alarming. No operation, however, was performed. She died the same day.

Autopsy No. 2558. Autopsy July 4, 1905, by Dr. W. Francis. Anatomical diagnosis. Old abdominal operation wound, hysterectomy and resection of the sigmoid for carcinoma, anastomosis of the colon, recurrence of carcinoma in anastomosis with stricture of the lumen, metastases in the peritoneum, small fecal fistula in the left inguinal region communicating with the descending colon. Fibrino-purulent peritonitis, source not determined. Atelectasis in the lower lobes of both lungs.

There is a small opening in the left inguinal region. The abdomen is slightly distended and on opening it a quantity of foul gas escapes. In the right upper quadrant in the region of the liver and extending over to the left upper quadrant is a little fibrino-purulent peritonitis. Large masses of fibrin cover the intestinal walls, surface of the liver, etc. This is walled off above the umbilicus by recent adhesions. Elsewhere about the abdomen there are adhesions which for the most part can be broken down. At the seat of the operation wound the structures are closely adherent to the abdominal wall. The intestines are everywhere bound down by adhesions which are of three varieties, fibrinous, fibrous, and nodular. In other places throughout the small intestines adherent coils are found to be strongly bound together,

but there are also localized small areas of adhesions consisting of round, hard nodules varying in size from a pea to a walnut and on examination consisting of dense, hard, more or less fibrous tissue studded with yellowish, opaque points. These new growths in many places project into the lumen, but in no way seem to have destroyed the mucosa. The nodules of new growth in the peritoneum are few in number, but each is apparently of considerable size. Except for these points the peritoneum seems to be free from new growth. The lower nine inches of the colon were removed three months previous to the autopsy and an end-to-end anastomosis was made between the colon and the rectum. This line of junction runs behind the stump of the cervix uteri. The lumen of the bowel at this point is greatly contracted, admitting only the tip of the little finger. On section through this line of junction, it is found to consist of dense fibrous tissue with very fine, yellowish, opaque points through it. The mucosa of the rectum is injected. The source of the peritonitis in the right upper quadrant is not determined. The appendix is perfectly normal. The vagina and the stump of the cervix appear normal, but the scar tissue around the cervix and along the line of the peritoneum is suggestive of a new growth. There is a large amount of carcinomatous-looking tissue between the cervix and the rectum.

On histological examination sections from the region of the anastomosis show normal mucosa and a thickened muscular coat. There is a thick mass of fibrous tissue with carcinomatous alveoli scattered throughout it. These show the type of the original tumor. The growth is a typical adeno-carcinoma. Sections from the large fibrous nodules in the peritoneum which bound the intestines together at several points show that they also consist of fibrous tissue with abundant areas of adeno-carcinoma scattered throughout them.

The mesentery consists chiefly of fat. It also shows alveoli. Sections from the scar tissue in the region of the intestinal anastomoses also contain masses of cancer cells. The original growth was evidently not entirely removed, and there had also been metastases before the operation was undertaken.

CARCINOMA OF THE RECTUM, SECONDARY TO A PRIMARY GROWTH IN THE RIGHT FALLOPIAN TUBE.

This case is of interest on account of the extent of the operation. Complete removal of the uterus by Wertheim's method is usually sufficiently severe to tax the patient's strength without any attempt to remove a large segment of the bowel. In this case the rectum was fortunately very lax, and after freeing it without in

any way disturbing the blood supply, we were able to do an exaggerated Whitehead operation, bringing down and cutting off the necessary amount of bowel, while still preserving the sphincter. Although in the end a hopeless case the patient was absolutely relieved of the distressing bowel symptoms, and to the day of her death, months later, never suffered from the slightest intestinal obstruction.

*Primary carcinoma of the right Fallopian tube (Fig. 9) with secondary involvement of the uterus, both ovaries, pelvic peritoneum, omentum, and rectum. Removal of omentum, uterus and appendages, one-third of the pelvic peritoneum, and six inches of the bowel. The patient was comfortable and considered herself well, five months after operation. The respite was, of course, only temporary.**

Mrs. Z. was seen in consultation with Dr. J. Milton Linthicum, Jan. 5, 1905. The patient was 55 years of age. She was sparely built, fairly well nourished, but slightly anemic. For months she had had some hemorrhage from the uterus and later great pain on defecation; in fact, her discomfort had been so great that she said she could not endure it much longer. On examination, under anesthesia, I found the uterus slightly enlarged and on the right side a firm mass about 6 cm. in diameter. I thought it to be a myoma.

Jan. 7.—On opening the abdomen I found the omentum everywhere studded with nodules, some of them being very small, others 1 cm. or more in diameter, and umbilicated. I questioned the advisability of operating, but Dr. Linthicum thought it wiser to operate, as the patient said "she would rather die than go through the torture that she had been experiencing for several weeks." The omentum was separated close to the transverse colon, as in the vicinity of the colon no metastases were to be found. The right tube was much enlarged and apparently involved in a malignant growth. It was attached to the pelvic floor and the peritoneum at this point, over an area fully 5 by 6 cm., was involved in the process. On the right side the ureter ran directly beneath the thickened peritoneum. On the left side the ovary, although small, was glued down to the pelvic floor directly over the ureter. Posteriorly the uterus was firmly attached to the rectum. It was found necessary to carefully dissect out the ureters first, as it was evident that much of the pelvic peritoneum must be removed. The hysterectomy was carried out practically along the lines laid down by Wertheim's operation. Fully one-third of the pelvic peritoneum, however, was removed. I had hoped to remove part of the rectum with the uterus in one piece,

*From the *Johns Hopkins Hospital Bulletin*, Vol. XVI, No. 177, December, 1905.

but found that it was impossible. Consequently it was necessary to separate the uterus from the rectal growth. The rectum was freed on all sides, care being taken, however, not to interfere with the blood supply. The patient was then placed in the perineal position and the skin separated from the rectal mucosa, just as is done in a Whitehead operation. Six inches of the rectum were drawn down through the sphincter and cut off and the upper edge of the rectum was sutured to the skin. The ureters were covered over as far as possible with the remaining peritoneum. A medium-sized gauze drain was introduced into the pelvis and brought out through the vagina. The entire operation took a little more than three hours. The patient had a very feeble pulse when she left the table, which was not surprising, as she was in a weak condition at the commencement of the operation.

Jan. 8.—The patient is improving greatly. Her pulse is 126, temperature normal, respirations about 30. This evening there has been considerable vomiting. Sixteen ounces of water were ordered with the hope of washing the stomach out. She vomited four ounces, fortunately retaining the twelve. There have only been about 70 c.c. of urine in twenty-four hours, but the general condition does not seem to indicate any uræmia.

Feb. 1.—The patient has steadily improved since operation. There has been a great deal of discharge from the pelvis, but that is rapidly diminishing. She occasionally has a temperature of 101° F. The sphincter action at first was rather tardy, but is now much better.

March 1.—The patient is up and around and suffering little or no discomfort. Of course, a complete recovery is out of the question. The operation was performed merely to relieve her intense suffering.

Subsequent History.—During the spring and part of the summer she was free from pain, journeyed to distant points, and looked very well. About the middle of August she became rather weak; after seven days' rest in bed she suddenly grew worse and died in half an hour. From the symptoms it is possible that death was due to embolism. The operation relieved her of great suffering and gave her over six months of comparative comfort.

Gyn.-Path. No. 8114. The specimen comprises the uterus and enlarged right tube, both ovaries, the small left tube, and a cuff of pelvic peritoneum, the greater part of the omentum, and several inches of the rectum.

The uterus has been removed entirely. It is 7 by 5 by 3.5 cm. and is covered with numerous adhesions. The uterine cavity is of the normal size. The mucosa is thinner than usual and shows nothing of interest.

The right tube at the uterus is 3 mm. in diameter. After passing outward 1.5 cm. it suddenly increases in size, reaching a diameter of 1.8 cm. It gradually increases until near the fimbriated extremity it is 4 cm. in diameter. The entire length of the tube is approximately 12 cm. It is for the most part smooth, but at two points on its inner aspect the muscular coats have given way and we have hernial spaces .8 by 1.8 cm. in diameter covered only by peritoneum (Fig. 9). The under sur-




FIGURE IX.—PRIMARY CARCINOMA OF THE RIGHT TUBE.

face of the extremity of the tube is roughened where it has been attached to the peritoneum of the pelvic floor. The tube was not opened until hardened. Sections near the uterus show that the lumen is fully 1 cm. in diameter, and that it is filled with a friable, porous, granular-looking growth which is free on the under side, but intimately blends with the upper or convex side of the tube. Sections near the outer end of the tube show that the walls are not over 1 mm. in thickness. Here also the tube lumen is filled with a similar friable growth which is whitish

yellow or mottled, evidently as a result of old hemorrhages. The tube itself is nearly as large as the uterus.

The left tube is about 5 cm. in length, is slightly beaded, and varies from 3 to 5 mm. in diameter. The fimbriated end is patent and the tube has grown fast to the lower and outer end of the right tube. The right ovary is very small, is approximately 1.5 by 1 by 1 cm. The left ovary is also atrophic, being 2 by 1.5 by .6 cm. Attached to the right side of the cervix is an irregular area of peritoneum which was approximately 7 by 6 cm. The central portion of this is hard and indurated, the outlying portions are smooth.

The rectum is atrophied to a considerable extent. The length of the portion removed, in its fresh state, is about six inches. The rectal mucosa is smooth and apparently normal. The constriction was due to infiltration of the adipose tissue surrounding the rectum. The nodules in the omentum, as noted in the clinical description, are firm. Some of them measure fully 3 cm. in length.

Histological Examination.—Sections from the uterus show that the surface epithelium is intact. The glands are normal. At numerous points the muscle is becoming active and growing up into the stroma of the mucosa. It shows us fairly well how an adeno-myoma may develop from an in-growth of the muscle fibres.

Sections from the tube near the uterus show that springing from the upper wall of the tube is a new growth, as indicated in the gross description. The lower part of the tube is free. Projecting from the side of the tube where the lumen is free are little finger-like outgrowths covered by a single layer of delicate epithelium. The nuclei are oval and vesicular. Some of the nuclei stain very deeply and are rather increased in size. They immediately remind one of a malignant growth. Springing from the wall of the tube and filling almost the entire cavity is a papillary growth. The stroma of the out-growth consists of spindle-shaped connective tissue cells. The epithelial covering is one or many layers in thickness. In numerous places the epithelial covering has greatly proliferated, and we have solid masses of cells many layers thick. Here also there are large and deeply staining nuclei. In the deeper portion of the growth the connective tissue predominates and we have solid nests of cells. The epithelium tends to retract from the connective tissue. At numerous points large areas of the growth have undergone coagulation necrosis and we have fragmentation of the nuclei. The outer muscular wall in most places is still preserved. At some points, however, the entire thickness of the tube has been involved by the growth. Sections from the outer portion of the tube yield practically the

same picture. The papillary arrangement is particularly well marked and many of the nuclei are spindle-shaped or irregular, very large and deeply staining. In some sections fully three-fourths of the field have undergone coagulation necrosis. In such areas only a few of the cells around the larger blood vessels still retain their vitality. Scattered throughout the muscular walls of the tube are definite masses of growth chiefly in the form of cell nests or penetrating glands and on the outer surface of the tube are little bunches of new growths. We have undoubtedly a primary carcinoma of the Fallopian tube with a penetration of the entire tubal wall at numerous points.

Sections from the right ovary, which was very small, show that the organ in some places is normal, but at many points it has been penetrated by masses of carcinoma which send out branches in all directions. The cells present exactly the same characteristics and are manifest in the depth as well as on the surface. Here also there is some breaking down. The left tube near the uterus is practically normal. The left ovary, although also very small, shows diffuse infiltration by the growth. The structure is recognized as that of typical, carcinomatous glands or as isolated, large, irregular cells with irregular and deeply staining nuclei; in fact the ovary is riddled by the growth.

Sections from the omentum show a most typical picture. In such areas the fat of the omentum is to a great extent replaced by young and old connective tissue, and lying in the connective tissue are masses of epithelial cells, very solid, with a definite glandular-like arrangement. The same large, deeply staining and irregular nuclei are also here in evidence. The nuclear figures are particularly well formed. We find considerable hemorrhage and also breaking down of the carcinomatous elements. The only extension to the rectum is by continuity from the outer surface. The rectal mucosa has not been involved.

Diagnosis.—Primary carcinoma of the right Fallopian tube with extension to the peritoneum of the pelvic floor, to both ovaries, and also to the rectum by continuity, general pelvic adhesions; extensive metastases into the omentum.

For a further discussion of the various forms of cancer of the tube, we would refer the reader to Dr. Elizabeth Hurdon's article, published in the *Johns Hopkins Hospital Bulletin*, Vol. XII., p. 315, 1901, and to the recent article by G. J. Tomson, published in *La Gynécologie* in February, 1905.

RECTAL DIVERTICULA.

Rectal diverticula are not common. They are usually encountered at autopsy, but rarely detected during life. Had it not been for the perforation of two of these with subsequent develop-

ment of an abscess between the indurated bowel and the uterus, the surgeon's aid would hardly have been required. As noted in the pathological report the greater part of the tumor is made up of indurated fat surrounding the diverticula. Had no microscopic examination been made this would have been classed as a brilliant and permanent recovery after removal of carcinoma of the rectum.

Operations on the sigmoid or lower rectum are much more easily handled in women than in men, as we can so readily drain through the vagina if need be. Gauze coming in contact with the point of anastomosis is, however, very prone to cause suppuration and then leakage from the bowel.

As the contents of the lower bowel are solid and usually rather hard, we have erred on the safe side and in each case brought out a loop of the descending colon and attached it to the skin, being prepared to open the bowel with the cautery if the slightest unfavorable symptoms should present themselves.

Diagnosis.—*Pelvic abscess, with retroverted myomatous uterus. Actual condition: Rectal diverticula, with rupture into the surrounding rectal fat, producing a definite tumor. Small abscess between the tumor and the pelvic floor** (Fig. 10).

History.—This patient was seen early in February, 1904, in consultation with Dr. S. T. Haffner. She was 60 years of age. For some time she had experienced slight difficulty in defecation, and for a few days had been running a temperature varying from 100 to 103° F.

Examination.—On vaginal examination, I found the uterus somewhat enlarged. Posterior to it, and apparently continuous with it, was a globular mass. This was very hard and resembled a myoma in contour. There was, however, a hard ridge over its lower portion, as is so often noted where pelvic abscess exists.

Operation.—On February 13 I made a small incision in the vaginal vault just posterior to the cervix, and after peeling back the mucosa entered Douglas' pouch with a pair of blunt artery forceps. A very small amount of pus and a few flakes of fibrin escaped, but the mass was in no way diminished in size. Realizing the presence of an unusual condition, I packed the opening in the vault and immediately entered the abdomen from above. Filling Douglas' sac almost completely was a tumor mass evidently springing from the sigmoid flexure, which had rotated 90 deg. and had become firmly embedded in the pelvis. It closely resembled a rectal cancer. On careful manipulation it was brought out of the pelvis, and on inspection no lymph glands were demonstrable. The diseased segment of gut was removed and an

* Reprinted from the *Journal of the American Medical Association*, November 1, 1901.

end-to-end anastomosis done with Connell and Lambert sutures, the former being employed at the mesenteric junction and for about two-thirds the circumference of the gut. A portion of the descending colon was brought up into a small incision in the left inguinal region and made fast, so that if occasion demanded it




FIGURE X.—DIVERTICULA OF RECTUM WITH ABSCESS BETWEEN BOWEL AND UTERUS.

could be opened with a thermo-cautery at a moment's notice. Drains were then introduced into the vagina and also through the lower angle of the abdominal incision. At the end of the fourth day there was considerable abdominal distension and the patient was very weak. We accordingly opened the descending colon at its point of attachment to the abdominal wall and at the same time forced the patient's nourishment. She promptly recov-

ered. The small fistulous opening was a few weeks later readily closed under local anesthesia, and the patient is now (March 1, 1906) perfectly well.

Examination of Tumor.—On laying the tumor open we found that there were two rectal diverticula passing out into the adipose tissues, and communicating with the lumen of the gut by openings not more than 1 mm. in diameter (Fig. 10). The larger diverticulum was 1 cm. in diameter and filled with a fecal mass. The floor of this diverticulum had given way, and the surrounding fat was everywhere infiltrated by inflammatory products. The excessive hardness of the tumor was due to replacement of the fat in many places by recent connective tissue. The small abscess between the tumor and the pelvic floor was due to the extension of the inflammatory process to the peritoneum of Douglas' pouch. The diverticula were lined by atrophic mucosa.

A rectal examination of this case would have yielded little information beyond the detection of some narrowing of the lumen of the bowel, which is often present in cases of pelvic abscess. In this case cancer of the bowel might very readily have been diagnosed and a colostomy performed.



MR. GEORGE COOPER FRANKLIN, F.R.C.S., LEICESTER, ENGLAND,
Retiring President of the British Medical Association.

The Canadian Journal of Medicine and Surgery.

**THE BRITISH MEDICAL ASSOCIATION—TORONTO
MEETING, AUGUST 21-25, 1906.**

CONSIDERABLE progress has been made with the arrangements for that notable event, the meeting of the British Medical Association in this city in the closing part of August. From the inquiries that are being received from every part of the continent, as well as from the British Isles, it is evident that a very large attendance will be recorded at this meeting. Over 200 members resident in the British Isles have already asked for accommodation, and in many cases they will be accompanied by members of their families. The Association will be convened under thirteen sections, which will meet daily from 9.20 to 1 o'clock. The afternoons and evenings will be devoted to general meetings, public addresses and various entertainments. There will be three public addresses delivered. Sir James Barr will present the address in Medicine, his topic being, "The Circulation Viewed from the Peripheral Standpoint." Dr. W. S. A. Griffith will deliver the address in Obstetrics, Sir Victor Horsley the address in Surgery, "The Technique of Operations on the Central Nervous System." and it is just possible that a public address will be delivered by Dr. Marie, of Paris. It is intended that clinics shall be held each morning at 8.30, when interesting cases will be reviewed by some of the prominent physicians and surgeons in attendance. Considerable advance has already been made in arranging for the work of the sections.

Anatomy.—The section of Anatomy will be under the presidency of Dr. Arthur Robinson, of Birmingham. Papers have been promised by the following: Dr. C. R. Bardeen, University of Wisconsin, Madison, Wis.; Prof. G. C. Huber, University of Michigan, Ann Arbor, Mich.; Prof. J. P. McMurrich, University of Michigan, Ann Arbor, Mich.; Dr. Ross E. Harrison, Johns Hopkins, Baltimore, Md.; Dr. H. Knowler, Johns Hopkins, Baltimore, Md.; Dr. G. L. Streeter, Johns Hopkins, Baltimore, Md.

It is also possible that Prof. Mall, of Johns Hopkins, Baltimore; Prof. C. S. Minot, Harvard Medical School, Boston; Dr. E. A. Spitzka, New York, and Dr. R. R. Bensley, of Chicago, may communicate papers.

Laryngology and Otology.—The section of Laryngology and Otology will be under the presidency of Dr. J. Dundas Grant, of London, and will have three or four principal topics for discussion:

1. "Operations for the Correction of Deviations of the



DR. R. A. REEVE, TORONTO,
President-Elect of the British Medical Association.

The Canadian Journal of Medicine and Surgery.

Nasal Septum." (Discussion to be opened by Dr. St. Clair Thompson, of London.)

2. "Laryngeal Disturbances Produced by Voice Use."
3. "The Indication for Ligation of the Jugular Vein in Otitic Pyemia."
4. "The Diagnosis and Treatment of Ethmoidal Disease."

Each discussion will occupy about two and a half hours, the remainder of the day being devoted to papers. It is just possible that Dr. Logan Turner will open the discussion on "Ethmoidal Disease."



DR. ALEXANDER MCPHETRAN, TORONTO,
President Canadian Medical Association.

Medicine.—Tuesday, Aug. 21st: "Blood Pressure in Its Relation to Disease." (a) Physiological Introduction (Dawson, of Baltimore). (b) Clinical Methods of Determining Blood Pressure; Their Uses and Limitations (Geo. Gibson, Edinburgh). (c) Pathology and Therapeutics of Blood Pressure (Sir Wm. Broadbent). Also possibly a paper on the subject by Clifford Allbutt, and one or two others, including one Canadian.

Wednesday, Aug. 22nd: Discussion in conjunction with the section of Physiology upon "Over and Under Nutrition, with



THE UNIVERSITY OF TORONTO (FROM THE SOUTH-EAST).
The Meeting Place of the British Medical Association, August, 1906.

The Canadian Journal of Medicine and Surgery.

Special Reference to Proteid Metabolism." Introduced by Chrittenden. Other special speakers: Herter, Starling, Hutchison, Francis Hare, A. Haig, and others.

Thursday, Aug. 23rd: Papers from William Osler, J. Mackenzie, and Erlanger on "Heart Block." Other papers: L. F. Barker, A. Stengel, A. McPhedran.

Friday, Aug. 24th: Papers devoted to neurological subjects. W. G. Spiller, "Syringomyelia." J. J. Putman.

The following gentlemen have signified their intention to contribute to the section: Dr. J. J. Putman, Boston, Mass.; Dr. W. G. Spiller, Philadelphia, Pa.; Dr. Alfred Stengel, Philadelphia, Pa.; Dr. Barber, Baltimore, Md.

Obstetrics and Gynecology.—The section of Obstetrics and Gynecology is under the presidency of Dr. A. H. Freeland Barbour, of Edinburgh. The following is the programme suggested.

Tuesday—Discussion on "Hyperemesis Gravidarum." Opened by Dr. J. C. Cameron, of Montreal.

Wednesday—"The Changes in Uterine Fibroids after the Menopause, with Special Reference to Operations."

Thursday—Subject for discussion and opener to be selected by Dr. Barbour.

Papers—"Uterine Myomata and Their Degenerative Changes," T. S. Cullen; "Sectional Anatomy of Labor" (lantern demonstration), A. H. F. Barbour; "Condition of Ovaries in Normal and Abnormal Pregnancy," C. Lockyer (lantern demonstration).

Surgery.—The section of Surgery is under the presidency of Sir Hector Clare Cameron, M.D., Glasgow. The following is the programme suggested:

Tuesday—"Enucleation of the Prostate Gland." Reader, Dr. Bingham, Toronto.

Wednesday—"Treatment of Ascites Secondary to Chronic Hepatitis."


Thursday—"Surgical Treatment of Ulcer of the Duodenum." Reader, Dr. W. J. Mayo, Rochester, Minn.

Friday—"Treatment of Acute Septic Peritonitis."

Pediatrics.—The section of Pediatrics is under the presidency of George A. Sutherland, M.D., London. The following is the programme suggested:

Tuesday—Discussion on "Congenital Pyloric Stenosis." The medical aspect of the subject will be introduced by Dr. Edmund Cautley, London, and the surgical aspect by Dr. Harold Stiles, Edin.

Wednesday—Discussion on "Pneumococcal Infection." The



DR. DAWSON WILLIAMS,
Editor *British Medical Journal*.

The Canadian Journal of Medicine and Surgery.

medical aspect will be introduced by Dr. Henry Ashby, Manchester.

Thursday—A Symposium on "Enterocolitis." The subject will be taken up under the following headings: (a) Etiology, (b) Pathology, (c) Symptoms, (d) Diagnosis and Prognosis, (c) Medical Treatment, (f) Dietetic Treatment.

Friday—A Discussion on "Rheumatism."

Psychology.—The section of Psychology is under the presidency of Wm. Julius Mickle, M.D., London. It has been arranged to have four discussions, one each day of the sectional meetings. The subjects are:

Tuesday—"General Paresis."

Wednesday—"Classification of Insanity."

Thursday—"So-called Mental Degeneracy."

Friday—"Dementia Precox."

The leaders and those chosen to discuss these subjects will be eminent British, American and Canadian psychologists, and the President, Dr. Mickle, is expected to present the first paper, as he is a recognized authority on general paresis. The second subject chosen will be one of great interest to both countries, as it is a question now under general discussion.

A series of papers will also be presented by eminent men, and the following Canadians have already signified their intention to take part: Dr. C. K. Clarke, Toronto; Dr. Ryan, Kingston; Dr. Moher, Brockville; Dr. Sherris, Montreal; Dr. Daniel Clark, Toronto.

State Medicine.—The section of State Medicine is under the presidency of Dr. F. Montizambert, of Ottawa. The following programme has been arranged:

Tuesday—"The Prevention of Tuberculosis."

Wednesday—"Water Supplies."

Thursday—"The Hygiene of Homes and Educational and Industrial Institutions."

Friday—"International Sanitary Protection."

Guests.—Prof. Brouardel, member of the Institute and the Academy of Medicine, France; Dr. Mattin, City Health Officer of Paris, France; Dr. Letulle, Professor of the Medical Faculty of Paris; Dr. Liccaga, Sanitary Adviser of the Government of Mexico, Mexico; Dr. Wyman, Surgeon-General of the United States Public Health and Marine Hospital Service, Washington.

Therapeutics.—The section of Therapeutics is under the presidency of Donald MacAlister, M.D., Cambridge. The following is the programme arranged:

Tuesday—The Study of the Kidney: (a) Its Physiology and Pharmacology; (b) The Therapeutics of Acute Nephritis;



MR. GUY ELLISTON,
Secretary of the British Medical Association.

(c) The Treatment of Chronic Nephritis; (d) The Treatment of Uremia.

Wednesday—"Serum Therapy."

Thursday—"The Place of Materia Medica and Therapeutics in the Medical Curriculum."

Friday—"The Value of Alcohol in Therapeutics," Dr. A. D. Blackader, Montreal; "The Teaching of Pharmacology"; "The Teaching of Therapeutics."

Pathology and Bacteriology.—The section of Pathology and Bacteriology, under the presidency of Professor J. G. Adami, M.D., F.R.S., Montreal, have made the following preliminary arrangements:

Tuesday—"Nuclear Physiology and Pathology." To be opened by Professor Adami and Dr. Macallum.

Wednesday—"Etiology and Life-History of Malignant New Growths."

Thursday—"The Forms of Arteriosclerosis, Their Classification and Experimental Production."

Friday—Papers upon "Pathogenic Protozoa" by various workers. Papers have been promised by Prof. Aschoff, Marburg, Germany; Prof. Novy, Ann Arbor; Dr. Pearce, Bender Laboratory, Albany; Dr. Bushnell; Prof. Grunbaum; Prof. Calder Leith; Dr. Oskar Klotz, Montreal; Prof. J. J. Mackenzie, Toronto.


The American Association of Pathologists and Bacteriologists have been formally invited to be present, and a number of the members will likely attend.

Dermatology.—This section will meet under the presidency of Dr. Norman Walker, of Edinburgh, who will open the section by an address on "The Teaching of Dermatology." During one of the days of the meeting there will be a discussion on the subject of "Eczema," to be opened by Dr. A. J. Hall, of Sheffield, Eng. A paper on "Psoriasis and Light" has been promised by Dr. J. N. Hyde, of Chicago. Papers will also be given by Dr. Gilchrist, Baltimore; Dr. A. R. Robinson, New York; Dr. Elliott, New York.

Physiology.—The section of Physiology will meet under the presidency of Professor W. D. Halliburton, M.D., F.R.S., London. The following programme has been arranged:

Discussions: (1) Discussions in junction with the section of Medicine on, "Over Nutrition and Under Nutrition, with Special Reference to Proteid Metabolism in Health and Disease"; (2) Discussion in junction with the section of Pathology on "The Role of the Nucleus in Nutrition."

Papers: Dr. S. P. Bebe, New York, on "Serum under the Influence of Injected Nucleo-proteid"; Prof. T. G. Bordie,



OFFICES OF THE BRITISH MEDICAL ASSOCIATION,
STRAND, LONDON, ENGLAND.

F.R.S., London, on "The Functions of the Renal Tubules and Glomeruli"; Prof. F. Gotch, F.R.S., Oxford, on "Demonstration of the Sphintharoscope"; Prof. W. B. Hall, Chicago, on "New Apparatus"; Prof. W. D. Halliburton, F.R.S., London, on "Proteid Nomenclature"; Prof. C. F. Hodge, Worcester, Mass., on "Structures and Physiological Functions of Amoeba Proteus"; Profs. C. F. Hodge and M. F. Duncan, Worcester, Mass., on "Differentiation of Contractile Protoplasm"; Prof. W. H. Howell, New York, on "Physiology of Heart"; Prof. G. C. Huber, Ann Arbor, on "Physiology of Renal Tubules"; Dr. G. T. Kemp, Champaign, Ill., on "Blood-platelets"; Dr. Louis Lapicque, Paris, on "Electrical Excitation of Nerves and Muscles"; Prof. J. S. Macdonald, Sheffield, on "Structure and Functions of Nerve Fibres"; Prof. J. J. R. MacLeod, Cleveland, on "Experimental Glycosuria"; Dr. Gustav Mann, Oxford, on "A Plea for Micro-physiology"; Prof. B. Moore, Dr. M. Edie, Dr. Spence, and Dr. H. E. Roaf, Liverpool, on "Experimental Glycosuria"; Prof. B. Moore, E. Whitley, and Dr. H. E. Roaf, Liverpool, on "Effects of Ions on Growth and Cell Division"; Dr. F. W. Mott, F.R.S., London, on "The Functional Significance of the Convolutional Pattern in the Primates"; Dr. Maurice Nicloux, Paris, on "Chloroform Anesthesia and a Simple Method of Estimating Chloroform"; Prof. C. S. Sherrington, F.R.S., and Dr. H. E. Roaf, Liverpool, on "Lock-jaw"; Prof. F. S. Lee, New York, on "The Causes of Fatigue in Certain Pathological States."

Papers are also promised by the following: Dr. Harvey Cushing, Baltimore; Dr. P. T. Herring, Edinburgh; Dr. F. G. Hopkins, F.R.S., Cambridge; Prof. Waldemar Koch, Columbia, Mo.; Dr. S. J. Meltzer, New York; Dr. Sutherland Simpson, Edinburgh; Prof. L. B. Mendel, New Haven; Prof. Porter, Boston; Prof. Jacques Loeb, Berkeley, Cal.

Ophthalmology.—The section of Ophthalmology will meet under the presidency of Robert Marcus Gunn, F.R.C.S., London. The following provisional programme has been arranged:

Tuesday—"Rare Forms of Choroiditis."

Wednesday—"Sympathetic Ophthalmia."

Thursday—"Affections of the Lachrymal Passages."

Friday—"Visual Tests for Marine and Railroad Service."

Most of the buildings of the University will be utilized in connection with the meeting. On the ground floor of the Main Building, in addition to the Post Office and Reception Rooms, there will be rooms for the regular meetings of some six or eight sections. The second floor, in addition to special offices for the Secretariat and the Editor of the *British Medical*




DR. RADCLIFF CROCKER,
Treasurer of the British Medical Association.

Journal, will be devoted almost entirely to the Museum, which will afford some 12,000 square feet for exhibitors. Already a large amount of this space has been disposed of to leading manufacturers of instruments and drugs in Great Britain, the United States and Canada. This alone will be one of the most interesting parts of the Association to Canadian visitors. Accommodation for other sections will be provided in rooms closely adjoining the Main Building. It is expected that the new Convocation Hall will be completed in sufficient time to enable the ceremonies of the official reception, on the evening of the 21st of August, and the public addresses, to take place there. Already the Committee is actively engaged in providing accommodation for the host of visitors that is expected. Queen's Hall, Wycliffe College, Annesley Hall, the Fraternity houses, and other buildings adjacent to the University will probably be utilized, and many of the citizens are already offering their hospitality. The Committee on Entertainment have a most excellent programme prepared, one of the interesting features of which will be an excursion to Niagara Falls at the invitation of Sir Henry Pellatt. Owing to the exceptionally favorable travelling rates which have been obtained over the Canadian lines of steam and rail, the attendance will be made very easy, and physicians wishing to avail themselves of the privileges of this meeting should communicate with the Secretaries at an early date, in order to obtain accommodation.

EASTERN CANADIAN PASSENGER ASSOCIATION.

(Including the following Lines.)

Algoma Central and Hudson Bay Railway, Bay of Quinte Railway, Boston and Maine R.R., Canadian Pacific Railway, Central Ontario Railway, Central Vermont Railway, Grand Trunk Railway System, Great Northern Railway of Canada, Huntsville, Lake of Bays and Lake Simcoe Nav. Co., Kingston and Pembroke Ry., Muskoka Lakes Nav. and Hotel Co., New York Central and Hudson River R.R., Niagara Navigation Co., Northern Navigation Co., Ontario and Quebec Navigation Co., Oxford Mountain Railway, Ottawa and New York Railway, Ottawa River Navigation Co., Pembroke Navigation Co., Pere Marquette, C.H. and D. System, Quebec Central Railway, Quebec Railway Light and Power Co., Quebec Southern Railway, Richelieu and Ontario Navigation Co., Rideau Lakes Navigation Co., Rutland R.R., Temiscouata Railway, Temiskaming and Northern Ontario Railway, Toronto, Hamilton and Buffalo Railway, Trent Valley Navigation Co., Dominion Atlantic Railway, Intercolonial Railway.



DR. LANGLEY BROWNE,
Chairman of Council of the British Medical Association.

1. *Fares, Going Dates and Limits.*—(a) Domestic Business, Certificate, Plan Arrangements; free return regardless of number in attendance. Passengers going rail, returning R. and C. Navigation Co., or *vice versa*, rate to be one-half fare.

(b) European Business.—On presentation of certificate, to be prepared and signed by the Secretary of the Eastern Canadian Passenger Association, and countersigned by the Secretary of the Canadian Committee, or the Secretary of the British Medical Association, one-way tickets to be issued at one-half lowest one-way first-class rail fare; round trip tickets at lowest one-way first-class rail fare between all points in Canada. Rates to the Pacific Coast subject to concurrence of Transcontinental Passenger Association. Steamship lines to advise Secretary what, if any, additional arbitraries are required. Dates of sale, July 1st to September 30th, 1906, inclusive. Final return date, September 30th, 1906.

2. *Extension of time Limit.*—On deposit with Joint Agent of Standard Convention certificates issued from points in the Maritime Provinces, points west of Port Arthur and from points in the United States, on or before August 28th, 1906, and on payment of fee of \$1.00 at time of deposit, an extension of time until September 30th to be granted. Joint Agency to be conducted in the name of G. H. Webster, Secretary, Eastern Canadian Passenger Association, will be kept open from August 21st to September 15th, 1906.

3. *Side Trips.*—(a) Side trip tickets to be sold from Toronto to delegates from the Maritime Provinces, from points west of Port Arthur, and from the United States on presentation of validated certificate, or deposit receipt, at lowest one-way first-class fare for the round trip, to all points in Canada. Dates of sale, August 23rd to September 1st, 1896, inclusive. Return limit, September 30th, 1906.

(b) Side trip tickets also to be sold to delegates from Ontario and Quebec to stations west of and including Sudbury, and east of and including Montreal, on presentation of validated certificate or deposit receipt, at lowest one-rate first-class fare for the round trip. It being understood, also, that the arrangements authorized for the extension of time limit from points in the Maritime Provinces, from points west of Port Arthur and from points in the United States will also apply for delegates from Ontario and Quebec.

Usual additional arbitraries *via* Upper Lake Steamships to apply, viz., going lake returning same, \$8.50 additional to be collected. Going lake, returning rail, or going rail returning lake, \$4.25 additional to be collected. Also usual arbitraries *via* St. Lawrence route, for delegates desiring to return by



SIR VICTOR HORSLEY, F.R.S.,
Who will deliver the Address in Surgery.

The Canadian Journal of Medicine and Surgery.

steamer, on presentation of tickets to purser, viz., \$6.50 Toronto to Montreal; \$3.50 Kingston to Montreal.

Via Northern Navigation Company on lines where meals and berth are included, rate to be single fare plus meal and berth arbitrary.

Ocean Transportation.—The "Lines" will grant the minimum rates named in the circulars published by the respective lines.

The Trunk Line Association covering New York State, Pennsylvania, West Virginia, Maryland, and Washington, including the following railways: Grand Trunk Railway, New York Central and Hudson River R.R., West Shore Railroad, New York, Ontario and Western Railway, Erie Railroad, Delaware, Lackawanna and Western R.R., Lehigh Valley R.R., Central Railroad of New Jersey, Philadelphia and Reading Railway, Pennsylvania R.R., Baltimore and Ohio R.R., Chesapeake and Ohio Railway.

Persons from points in the territory of the Trunk Line Association, who pay full first-class fare going to the meeting, shall be returned at *one-third* the highest limited fare by the route travelled, *on the certificate of the Trunk Line Association*. This certificate must be obtained from the ticket agent at starting points, and when endorsed by the Secretary of the Canadian Committee or the General Secretary of the British Medical Association, and vised by the special agent of the railway companies, may be presented by the holder at the place of meeting to obtain the concession returning. A fee of 25 cents will be charged for each certificate vised.

The returning journey must be made by the line over which the going journey was made.

Guests attending and making use of these certificates should give the ticket agents *timely notice* of their intention, in order that through ticket and certificates may be ready when required.

Going tickets and certificates will be issued August 17th to 23rd. Certificates will be vised August 21st to 25th, inclusive, and then honored for return tickets to and including August 29th. Extension of the return limit may be obtained to leave Toronto up to and including September 30th, by depositing validated certificates with Mr. G. H. Webster, Joint Agent at Toronto on or before August 28th, and making a payment to him of \$1.00 at the time of deposit.

The New England Passenger Association covering Maine, New Hampshire, Vermont, Massachusetts, Connecticut and Rhode Island, including the following railways: Bangor and Aroostook R.R., Boston and Albany R.R., Boston and Maine R.R., Canadian Pacific Railway, Central Vermont Railway, Eastern S.S. Co., Grand Trunk Railway, Maine Central R.R.



SIR JAMES BARR,
Who will deliver the Address in Medicine.

The Canadian Journal of Medicine and Surgery.

New York and New Haven and Hartford R.R., Portland and Rumford Falls Railway, Quebec Central Railway, Rutland R.R.

Certificate Plan Arrangements.—Free return regardless of the number in attendance.

Passengers going by rail, returning by Richelieu and Ontario Navigation Co., or *vice versa*, rate to be one and one-half fare.

On deposit with Joint Agent of Standard Certificates on or before August 28th, 1906, and on payment of fee of one dollar at time of deposit, an extension of time until September 30th will be granted. Joint Agency to be conducted in the name of G. H. Webster, Secretary, E.C.P. Association, will be kept open from August 21st to September 15th, 1906.

Side-trip tickets to be sold from Toronto to delegates from the Maritime Provinces, from points west of Port Arthur and from the United States, on presentation of validated certificates, or deposit receipt, at lowest one-rate way first-class fare for the round trip to all points in Canada.

Dates of sale, August 23rd to September 1st, 1906, inclusive; return limit, September 30th, 1906.

Usual additional arbitraries *via* Upper Lake Steamships to apply, viz., going lake, returning same, \$8.50 additional to be collected. Going lake, returning rail, or going rail, returning lake, \$4.25 additional to be collected. Also usual arbitraries *via* St. Lawrence route, for delegates desiring to return by steamer, on presentation of tickets to purser, viz., \$6.50, Toronto to Montreal; \$3.50, Kingston to Montreal.

Via Northern Navigation Company on lines where meals and berth are not included, the rail rate will apply; on lines where meals and berth are included, rate to be single fare, plus meal and berth arbitrary.

Excursions and Side Trips being Arranged.—A large number of excursions and side trips are being arranged for, some that can be accomplished in a day, and others taking several days, so that there is no question that our visitors from across the Briny will be afforded every opportunity of seeing the beauty spots of this fair Dominion. We append herewith a list of a few of the trips which the Committee contemplate arranging:

1. A trip to Algonquin National Park—Distant from Toronto, 205 miles. Cost for round trip (Association rate), \$5.65 (about £1 2s. 3d.).

2. To Lake Nipissing and French River—Toronto to North Bay, 227 miles. Cost for round trip (Association rate), \$6.85 (about £1 7s. 3d.).

3. A trip to the Temagami Region—300 miles. Cost of round trip (Association rate), \$9.95 (about £1 19s. 10d.).

4. A trip to Lake Simcoe and Couchiching, Bass Lake and the Severn River.



DR. W. S. A. GRIFFITH,
Who will deliver the Address in Obstetrics.

5. To Muskoka Lakes—Round trip, distance, 112.

6. A trip to the Kawartha Lakes—Round trip fare (Association rate), \$4.30 (about 17s. 2d.).

7. A trip to the Lake of Bays—Distance from Toronto, 146 miles; cost of round trip (Association rate), \$4.45 (about 17s. 10d.).

HOTELS AND LODGINGS COMMITTEE.

Name of Hotel, and Address.	Number of Guests	Rates per Day.
The New Russell House, 217 Yonge St.	40	\$1.50 (special rate).
The Tremont, 163 Yonge St.	100	\$2.00 (American plan).
The Queen's, Front St.	200	\$3.00 up, without bath; \$4.00, with bath (American plan).
The Arlington, King St. W.	50	\$2.00 to \$2.50 (American plan).
The King Edward, King St. E.	700	EUROPEAN—Single room without bath \$1.50 up, with bath \$2.00 up; double room without bath \$2.50 up, with bath \$3.50 up. AMERICAN—Single room without bath \$3.50 up, with bath \$4.00 up; double room without bath \$6.00 up, with bath \$7.00 up.
The Iroquois, King St. W.	60	\$1.50 up.
Rossin House, King and York Sts.	500	\$2.50 (American plan).
Hotel Gl.ystone, 1204 Queen St. W.	75	\$1.50 (\$1.00 for bed and breakfast).
Walker House, Front and York Sts.	75	\$1.50 up.
Name of Boarding House.	Number of Guests	Rates per Day.
The Waverley, 484 Spadina Ave.	10	\$1.00 to \$1.50.
Matheson Hall, 599 Yonge St.	25	\$1.00 to \$1.50.
The Avonmore, 276 Jarvis St.	25	\$1.25 (single meals 25 cents).
The Abberley, 258 Sherbourne St.	100	\$1.00 (special rate).
Mrs. Snell, 39 Grosvenor St.	10	\$1.25 to \$1.50.
Conservatory Residence, 47 St. George St.	20	\$1.50 to \$2.00.
Mrs. Ray, 70 Murray St.	6	\$1.00 (room and breakfast).
Queen's Hall, 7 Queen's Park.	48	\$2.00 (including meals).
Annesley Hall, Queen's Park.	65	\$1.25 (gentlemen preferred to ladies).
Nurses' Home, Children's Hospital.	75	

The Canadian Journal of Medicine and Surgery

J. J. CASSIDY, M.D.,
EDITOR,

43 BLOOR STREET EAST, TORONTO.

Surgery—F. N. G. STARR, M.B., Toronto, Associate Professor of Clinical Surgery, Toronto University; Surgeon to the Out-Door Department Toronto General Hospital and Hospital for Sick Children; N. A. POWELL, M.D., C.M., Prof. of Medical Jurisprudence, Toronto University, Surgeon Toronto General Hospital etc.

Clinical Surgery—ALEX. PRIMROSE, M.B., C.M. Edinburgh University; Professor of Anatomy and Director of the Anatomical Department, Toronto University; Associate Professor of Clinical Surgery, Toronto University; Secretary Medical Faculty, Toronto University.

Orthopedic Surgery—B. E. MCKENZIE, B.A., M.D., Toronto, Surgeon to the Toronto Orthopedic Hospital; Surgeon to the Out-Patient Department, Toronto General Hospital; Assistant Professor of Clinical Surgery, Ontario Medical College for Women; Member of the American Orthopedic Association; and H. F. H. GALLOWAY, M.D., Toronto, Surgeon to the Toronto Orthopedic Hospital; Orthopedic Surgeon, Toronto Western Hospital; Member of the American Orthopedic Association.

Gynecology and Obstetrics—GEO. T. MCKENUGH, M.D., M.R.C.S. Eng., Chatham, Ont.; and J. H. LOWE, M.D., Toronto.

Medical Jurisprudence and Toxicology—ARTHUR JUKES JOHNSON, M.B., M.R.C.S. Eng.; Coroner for the City of Toronto; Surgeon Toronto Railway Co., Toronto; W. A. YOUNG, M.D., L.R.C.P. Lond.; Associate Coroner, City of Toronto.

Physiology—CHAS. F. DICKSON, M.D., C.M., Queen's University, M.D., University of the City of New York; Electrologist Toronto General Hospital; Hospital for Sick Children and St. Michael's Hospital.

Pharmacology and Therapeutics—A. J. HARRINGTON, M.D., M.R.C.S. Eng., Toronto.

Pediatrics—A. R. GORDON, M.D., Toronto; HELEN MACMURCHY, M.D., Toronto.

W. A. YOUNG, M.D., L.R.C.P. Lond.,
MANAGING EDITOR,

145 COLLEGE STREET, TORONTO.

Medicine—J. J. CASSIDY, M.D., Toronto, Member Ontario Provincial Board of Health; Consulting Surgeon, Toronto General Hospital; and W. J. WILSON, M.D., Toronto, Physician Toronto Western Hospital.

Oral Surgery—E. H. ADAMS, M.D., D.D.S., Toronto.
Clinical Medicine—ALEXANDER McPHERAN, M.D., Professor of Medicine and Clinical Medicine Toronto University; Physician Toronto General Hospital, St. Michael's Hospital, and Victoria Hospital for Sick Children.

Mental and Nervous Diseases—N. H. BLSHER, M.D., Minko Insane Asylum; CAMPBELL MEYERS, M.D., M.R.C.S. L.R.C.P. (London, Eng.), Private Hospital, Deer Park, Toronto.

Public Health and Hygiene—J. J. CASSIDY, M.D., Toronto, Member Ontario Provincial Board of Health; Consulting Surgeon Toronto General Hospital; and E. H. ADAMS, M.D., Toronto.

Physiology—A. B. EADIE, M.D., Toronto, Professor of Physiology Woman's Medical College, Toronto.

Pathology—W. H. PEPLER, M.D., C.M., Trinity University; Pathologist Hospital for Sick Children, Toronto; Associate Demonstrator of Pathology Toronto University; Physician to Out-Door Department Toronto General Hospital; Surgeon Canadian Pacific R.R., Toronto; and J. J. MACKENZIE, B.A., M.B., Professor of Pathology and Bacteriology Toronto University Medical Faculty.

Ophthalmology and Otolaryngology—J. M. MACCALLUM, M.D., Toronto, Professor of Materia Medica Toronto University; Assistant Physician Toronto General Hospital; Oculist and Aurist Victoria Hospital for Sick Children, Toronto.

Laryngology and Rhinology—J. D. THORBURN, M.D., Toronto, Laryngologist and Rhinologist, Toronto General Hospital.

Dermatology—D. KING SMITH, M.B. Tor., Toronto

Address all Communications, Correspondence, Books, Matter Regarding Advertising, and make all Cheques, Drafts and Post-office Orders payable to "The Canadian Journal of Medicine and Surgery," 145 College St., Toronto, Canada.

Doctors will confer a favor by sending news, reports and papers of interest from any section of the country. Individual experience and theories are also solicited. Contributors must kindly remember that all papers, reports, correspondence, etc., must be in our hands by the first of the month previous to publication.

Advertisements, to insure insertion in the issue of any month, should be sent not later than the fifth of the preceding month. London, Eng. Reprint at 4 ve. W. Hamilton Mill, Thanet House, 231 Strand, W.C. Agents for Germany, Saitzsch's News Exchange, Mainz, Germany.

VOL. XX.

TORONTO, JULY, 1906.

NO. 1.

Editorials.

OUR BRITISH MEDICAL ASSOCIATION NUMBER.

As yet we hear only the fog horn, and we are straining our eyes for the lights of the good ship with its human freight of British medicos, so soon due on this side of the herring pond.

A welcome awaits our expected guests, and so we have unfurled a few flags and a bit of bunting in this issue of our journal, and hung a few portraits of faces that will greet each other and

perchance give us occasion to raise the shout, "For he's a jolly good fellow!"

It's the second time during the last decade that this honored Association has graced Canada with its presence, and the promise of meeting place, upon the former occasion in Montreal, and now in Toronto, giving this journal its first opportunity to wear court dress, and we hope the bow will be accepted as graciously as it is made. Many difficulties had to be surmounted and several disappointments suffered, including the circumstance that, at the last moment, our esteemed Canadian confrere, Dr. William Osler, now of Oxford, was unable to finish his promised paper in time for this issue. Also, the corrected galley proofs of the paper of our collaborator, Dr. Alexander McPhedran, who is travelling abroad, have not reached us, and it is now the eleventh hour. Early summer is with us in this lovely land, and we let the old inhabitant, Hiawatha, of song and story, extend his greeting:

"Beautiful is the sun, O strangers,
When you come so far to see us!
All our town in peace awaits you,
All our doors stand open for you;
You shall enter all our wigwams,
For the heart's right hand we give you.

"Never bloomed the earth so gaily,
Never shone the sun so brightly,
As to-day they shine and blossom
When you come so far to see us!

"Never was our lake so tranquil,
Nor so free from rocks and sand-bars;
For your birch canoe in passing
Has removed both rock and sand-bar!

"Never before had our tobacco
Such a sweet and pleasant flavor,
Never the broad leaves of our corn-fields
Were so beautiful to look on,
As they seem to us this morning,
When you come so far to see us!"


W. A. Y.

RELATIVE TO THE HISTORY OF THE CONSTITUTION OF THE BRITISH MEDICAL ASSOCIATION.

THE British Medical Association, under its present constitution, is a federation of local medical societies, called Divisions. The Divisions are grouped for certain purposes in local bodies



A. W. MAYO ROBSON, F.R.C.S.



H. MACNAUGHTON-JONES, M.D.,
M.Ch., R.U.I.




W. ROSE, M.S., F.R.C.S.



LENNOX BROWNE, F.R.C.S.



A. CARLESS, M.S., F.R.C.S.



H. W. ALLINGHAM, F.R.C.S.

called Branches. The aggregate of Branches composes the Association.

Each Division and Branch has its own local administration and rules, subject only to such restrictions upon each as are deemed needful for the harmonious co-operation of all.

The Divisions and Branches are linked together in the Association by the following means:

(a) Every member of a Division is a member of the Association, and conversely every member of the Association is a member of the Division in whose area he resides.

(b) The whole Association is subject to certain general regulations, defined in the Memorandum of Association, Articles of Association and By-laws.

(c) The governing bodies of the whole Association are: For certain specific purposes, the General Meeting of the Members, in which every member is entitled to take part; and for all other purposes, the Representative Meeting, in which every member through his Division is entitled to be represented.

(d) The Association has a Central Executive, consisting of a Representative Council and Committees.

(e) The whole Association has an official organ, the *British Medical Journal*, published under control of the Council.

(f) A uniform subscription of 25s. is paid by every member, and entitles to all ordinary privileges of membership, both of the Association, and of any Division and Branch to which a member may at any time belong.

In the early years of its life, the constitution of the British Medical Association was simple—a small Council by which all its affairs were governed, and, until the establishment of the Branches began, this sufficed for all its requirements.

But on this Council the Branches soon desired to have some representation, and as they grew in number, so the Council also gradually increased in size, until it became unwieldy and greatly hampered the conduct of business. It was decided that a Special Committee, to be known as the "Committee of the Council," should be brought into existence, and, subject to the veto of the Council at the annual meeting, was to be the real managing "cabinet" of the Association.

The Committee was to consist of Past Officers, Presidents,



PROF. WILLIAM OSLER, OXFORD, ENGLAND.

The Canadian Journal of Medicine and Surgery.

Presidents of Council, Treasurers, and so on, who were made "Vice-Presidents for life," and twenty members of the Council to be elected by ballot by the Council, was to meet four times a year, and at the end of each year was to be accountable to the Council for its actions and policy, and to stand or fall thereby.

The members of the Committee of Council paid their own expenses, the honor of the position being regarded as ample recompense for any sacrifice made.

After a time, however, a demand for representative government was made. At the annual meeting, held in Worcester in 1882, the questions of the constitution and government of the Association were raised by the following motion, of which due notice had been previously given: That the Committee of Council be requested to consider in which way direct representation of the Branches can best be secured.


This resolution was carried, and in response to it on October 18th, 1882, a sub-committee was appointed to consider the question and to report to the Council. This sub-committee came to the conclusion, that its first duty was to ascertain the mind of the Association generally concerning the matter in question.

With this object in view, it was determined to issue through the President of the Council a series of questions, together with a covering circular letter, addressed to the President and Secretary of every home Branch, asking its views on the subject.

This letter set forth the question under consideration in its fullest aspect, sought the counsel and advice of the Presidents and Secretaries of every home Branch concerning it, and appended to it was a series of questions, the answers to which, it was thought, would be decisive. Up to this time every Branch had been represented on the Committee of Council by its Honorary Secretary, *ex-officio*, and the first question was: Is your Branch satisfied with its present method of representation on the Committee of Council by its Honorary Secretary, *ex-officio*?

Sixteen questions were added to enable each Branch to make clear its objections to the above, if it had any, and aimed at getting fairly at the root of the matter. They ran as follows:

Has your Branch, having the power to elect a special Honorary Secretary to represent it on the Committee of the Council, availed itself of that power?



VIEW OF THE OLD RESIDENTIAL WING, UNIVERSITY OF TORONTO.
The Canadian Journal of Medicine and Surgery.

What is the total number of the members of your Branch, and what is the income of your Branch from Branch subscriptions?

Are the travelling expenses of your Honorary Secretary to the meetings of the Committee of Council defrayed by the Branch?

Is there any feeling in your Branch of inadequate representation?

Is the attendance of your Honorary Secretary at the meetings of Committee of Council influenced by the payment or non-payment of his travelling expenses?

Have you any other suggestions to offer to the sub-committee with regard to the representation of your Branch in the Committee of Council?

The answers received in reply to these questions were difficult to summarize, but the sub-committee, acting on what appeared to be their general sense, drew up an exhaustive report, and this report, which, for convenience of debate, was drawn up in two parts, was discussed at a special meeting of the Council held at Birmingham, May 17th, 1883.

The report with some slight modifications was adopted, and it was resolved: That the Committee of Council be requested to give due notice before the annual meeting of such alterations in the laws and by-laws of the Association as may be necessary for carrying into effect the change in the constitution of the government of the Association, embodied in the report of the sub-committee on the representation of the Branches in the Committee of the Council, as amended by the Council this day.

And this resolution was followed by another: That, as the Council, as proposed to be elected, will in future be the executive or governing body, and the Committee of Council, thus being no longer required, will cease to exist, the required alterations of the laws and by-laws be made by the Committee of Council, with the assistance of the solicitor of the Association.

The new Council of the Association was to consist of the Officers, as before, the Vice-Presidents, who had been, or should in future, be elected for life, of one representative for every Branch as heretofore, the larger Branches having the right to elect out of their members additional representatives on the fol-



TOWER, WEST END, MAIN BUILDING, UNIVERSITY OF TORONTO.

lowing scale, namely: That in Branches numbering over two hundred members there should be one additional representative for every two hundred members.

The question of the payment of the expenses of representatives was, after a good deal of discussion, finally decided at a general meeting, held at Leeds in 1889, a resolution being carried, that first-class railway fares, to and fro, were to be allowed to such representatives when travelling in the United Kingdom.

At the annual meeting of 1900, at Ipswich, a Committee was appointed to consider and report, first to the Branches, and subsequently to the Annual General Meeting, on the changes required in the constitution. The scheme prepared by this Committee was approved, subject to a few minor changes, by the Annual General Meeting of 1901, at Cheltenham, and a further Committee appointed to supervise the preparation of new articles of Association and By-laws to give effect to the scheme.

The articles and by-laws so prepared were adopted by special resolution on July 9th, 1902, and are those now in force, subject to certain amendments of details of the by-laws, which have been subsequently adopted in the manner provided in the constitution.*


J. J. C.

THE COMMITTEES OF THE BRITISH MEDICAL ASSOCIATION.

In dealing with the many and important subjects cropping up and claiming consideration or settlement at its hands, the Council of the Association commits, in the first instance, nearly everything to the scrutiny of a small committee, by whom it may be examined in all its bearings and reported on, in print, to the Council.


Several of the committees are permanent bodies, only a certain number of whose members are changed from time to time, and they have thus a thorough knowledge of the matters each takes in hand. The Journal and Finance Committee, the Parliamentary Bills Committee, the Scientific Grants Committee, the Trust Funds Committee, and the Library and Premises Committee may be taken as examples of permanent committees, with-


* For further details of the constitution of the British Medical Association see British Medical Association Year Book, 1906, p. 69.



SOUTHEASTERN VIEW FROM UNIVERSITY COLLEGE TOWER.

The Canadian Journal of Medicine and Surgery.





BIOLOGICAL BUILDING, QUEEN'S PARK.

The Canadian Journal of Medicine and Surgery.

out whose special attention in their several directions the Council could not act intelligently. As special questions arise from time to time, concerning which some members are known to possess special knowledge, they, in like manner, are generally placed in the hands of such members—are thrashed out by them first and then reported on to the Council for ultimate decision or dismissal. The Committee of Anesthetics, the General Practitioners' Committee, the Committee on the Eye-Sight of Public Servants, that on Weak-Minded Children, are examples of such Committees.

J. J. C.


THE "BRITISH MEDICAL JOURNAL."

THE *British Medical Journal* is, as we would say in Canada, the linch-pin of the Association, for, were it other than this—one of the first medical journals in the world—the British Medical Association could not have reached its present position of influence and numerical strength.

For many years after the formation of the Association, in the hands of editors resident in various provincial centres, the *Journal* remained provincial in character and of very ordinary reputation. As a record of the work and proceedings of the Association it was welcome to members, and in its pages are preserved addresses, lectures, papers, records of cases and other matters of the greatest value and importance.

Mr. Ernest Hart was appointed editor in 1867, and, from that time, with a slight interregnum in 1869 and 1870, up to the time of his death in 1900, he continued to occupy the editorial chair. It can be fairly said that much of the success of the *British Medical Journal* depended on his skill and untiring industry.

It is pleasing to be able to bear witness to the fact that the achievements of Mr. Dawson Williams, the present editor of the *British Medical Journal*, show no recession from the high mark left on the sands of editorial fame by his gifted predecessor. Devoted as the *British Medical Journal* has been to the service of the members of the Association of every class—to the public services, military, naval, poor law, and representative as it has been of every advance in medicine, surgery and midwifery in all



GYMNASIUM, UNIVERSITY OF TORONTO.

The Canadian Journal of Medicine and Surgery.

countries, it has become one of the leading medical journals of the world. It is likewise acknowledged everywhere to be the type of what a reputable medical journal should be.

In the editorial columns topics of current medical interest are discussed, and great efforts are made to supply members with early, accurate and complete information on all questions scientific, political or social affecting the profession. The editor is ably assisted by a large staff of experienced writers, experts and specialists. Special correspondents supply news from the principal foreign countries and the British colonies.

A great part of the space of the *British Medical Journal* is devoted to articles on medicine, surgery and pathology. In it are published each autumn the addresses delivered before the annual meeting, and reports of the discussions which take place in the sections. Clinical lectures and papers, shorter notes from private practice are published at other seasons of the year. The clinical and scientific work done by the Divisions and Branches of the Association, as well as by the principal medical sections in the United Kingdom, are reported in the body of the *Journal*.

The appearance of the *Journal* is fairly indicative of its worth and importance in the field of medical journalism. With the first number of 1906 improvements were introduced in the style of printing, which add to the ease with which it is read.

The issue of the *Journal* is now 23,000 copies. It is published weekly in London, and is sent post free to all members of the Association, whether resident in the United Kingdom, in British colonies, or in foreign countries. J. J. C.

SOME OF THE LEADING FACTS IN THE HISTORY OF THE BRITISH MEDICAL ASSOCIATION.*

THE British Medical Association, which will hold its seventy-fourth annual meeting at Toronto, August 21-26, 1906, is considered to be the largest and most influential medical association in the world. It seems opportune, therefore, to present, in the current number of THE CANADIAN JOURNAL OF MEDICINE

*For the facts and data which appear in these articles we are indebted to an abstract from the *British Medical Journal* dated June 19th, 1897, entitled "A Cursory Survey of the history of the British Medical Association from its Institution to the Present Time," and also to the *British Medical Association Year Book, 1906*.



SCHOOL OF PRACTICAL SCIENCE, CHEMISTRY AND MINING BUILDING.


The Canadian Journal of Medicine and Surgery.

AND SURGERY, some of the main facts in the history of this celebrated association, including some of the ups and downs it encountered before reaching its present influential position.


The British Medical Association, originally designated "The Provincial Medical and Surgical Association," was founded at Worcester (Eng.), on July 19th, 1832. On that day, fifty medical men attended a meeting at the Worcester Infirmary, convened by Dr. (afterwards Sir) Charles Hastings, to consider the formation of the proposed Association. After pointing to the avidity with which the proposal for such a society had been received as an omen favorable to the undertaking, Dr. Hastings outlined the work, which it was hoped that the Association might accomplish, under the following general heads:

(1) Advancement of Medical Science; generally by the collection of useful information, by means of original essays and reports of hospitals and similar institutions, and of private practice, (a) medical topography, (b) the investigation of endemic and epidemic disease, and (c) medico-legal science.


(2) The maintenance of the honor and respectability of the profession by promoting friendly intercourse and free communication among its members. Under this head with regard to medical ethics, attention was drawn to the fact "that with the exception of a few essays, no attempt had been made to establish a code for the guidance of those needing such direction. In a well-organized profession there could be no difficulty in adopting to its exigencies the doctrines of general ethics." With reference to medical politics attention was called to the fact "that the organization of the profession which obtained was not what it should be; the whole system of medical polity in England was both defective and erroneous. Opinions differed widely as to the evils and remedies, but few commended the existing state of things. The subject was closely connected with the advancement of science, for, if the profession were constituted as it ought to be, the harmony that would be established could not fail to be a direct means of more cordial and efficient co-operation in extending the science and improving the practice of medicine. During the first decade of its existence annual meetings were held at Bristol (1833), Birmingham (1834), Oxford (1835), Manchester (1836), Cheltenham (1837), Bath (1838), Liverpool



The Canadian Journal of Medicine and Surgery.



VICTORIA UNIVERSITY, QUEEN'S PARK.



ANNESLEY HALL—THE VICTORIA UNIVERSITY WOMEN'S RESIDENCE.

The Canadian Journal of Medicine and Surgery.


(1839), Southampton (1840), York (1841), Exeter (1842), Leeds (1843), and so on. From this time on towns in different parts of England began to vie with each other for the honor, as it had soon come to be considered, of entertaining the Association.

These annual gatherings served to bring together the best men, old and young, of each district, and as year after year, district was joined to district, so in direct proportion grew the acquaintance, good fellowship and friendship of practitioners from all parts of England.

It was not very long before the Association had become too large to be able to continue in its original lines, and it became necessary to divide it into "Branches," each to be in itself a fair representation of the parent Society—each to have its President, Secretary, Council, and its own by-laws, which were to be subject to the approval of the Council of the Association; and each to hold its own separate, ordinary and annual meetings, and to have its own representative or representatives, according to numerical strength on the Council of the parent Association. Of these Branches, the East Anglian and the Bath and Bristol seem to have been the first formed; Bath taking the lead in 1836, followed by Bristol in 1840, and these two becoming united, with a combined membership of 286 in 1842.

By the end of 1878, not only had the Association spread into Wales, Scotland and Ireland, but by the formation of the Jamaica Branch, the Colonies had put in their claim for representation. In the eighties thirteen colonial applications were received and granted. The Adelaide and South Australia in 1880; the Melbourne and Victoria in 1880; the Sydney and New South Wales in 1880; the British Guiana in 1883; the South Indian and Madras in 1884; the Bermuda in 1886; the Halifax and Nova Scotia in 1887; the Colombo and Ceylon in 1887; the Malta and Mediterranean in 1888; the Griqualand West in 1888; the Barbadoes in 1889; the Cape of Good Hope in 1889; the Bombay in 1889; the Punjaub in 1889.

During the nineties the following Branches were formed: The Londonderry and North-West of Ireland in 1890; the Leeward Islands in 1890; the Burmah in 1891; the Hong Kong in 1891; the Montreal in 1891; the Manitoba and West of Canada in 1891; the Trinidad and Tobago in 1892; the Cork and South



KNOX COLLEGE, TORONTO.

The Canadian Journal of Medicine and Surgery.

of Ireland in 1893; the Dundee and District in 1893; the South-east District of Ireland in 1893; the Grahamstown and Eastern Province in 1893; the Brisbane and Queensland in 1894; the British Columbia in 1894; the Deccan in 1894; the Gibraltar in 1894; the Malaya in 1894.

In 1875	the membership of the Association	was	6,112
" 1885	" "	" "	11,249
" 1895	" "	" "	16,000
" 1905	" "	" "	19,582

During all the early periods of its existence, far, indeed, into its life, a want of organization in the conduct of the affairs of the Association and divided administration tended to impede what might otherwise have been more rapid progress.

With editors in one place, treasurers in another, and secretaries, again, elsewhere, with offices in yet another, and with no settled abode anywhere, with no one to give business attention to the collection of subscriptions, it cannot be wondered at if the financial condition of the Association was a source of great anxiety and of much discussion at the annual meetings.

A Committee of Inquiry was appointed by the Association to examine into these matters. After several meetings the result of the deliberations of this Special Committee was handed to Council, and its judgment was summed up in the following recommendations:

That the office of the Association in Birmingham be closed and that the office be removed to London.


That the Secretary be replaced by a paid "General Manager," whose whole time should be devoted to the duties of the office.

That these should be the collection of the subscriptions from either the local Secretaries of the Branches, or from such subscribers, attached or unattached, to the Branches, as preferred to pay them through him.

To overlook and arrange for procuring advertisements for the *Journal*.

To be responsible for the keeping of the books and correspondence of the Association.

To overlook the office work, the printing of the *Journal*, and generally to manage the business of the Association, as distinct



UPPER CANADA COLLEGE, DEER PARK

The Canadian Journal of Medicine and Surgery.

from the literary and editorial work of the *Journal*, with which, of course, he was to have nothing to do.


These recommendations were carried out, and the entire management of the business of the Association was transferred to London. Mr. Francis Fowke was appointed General Manager, and he entered upon his duties, January 1st, 1872. His was a difficult task. Nevertheless, he began his work with the determination to bring it to a successful issue, and, though he had to labor long and arduously before any great result was apparent, he had the satisfaction of seeing a tendency towards improvement from the beginning, and of knowing that he had the support of every member of the Council, and the good-will of all in the furtherance of his efforts.

At the close of 1871 advertisements produced £1,992; by 1881 they had risen to £6,089; by 1891 to £14,568. The issues of the *Journal* and the subscriptions for the same periods showed equally marked and encouraging results. In 1871 subscriptions amounted to £4,677; in 1881, £9,147; in 1891, £14,759, and up to the present time the increase is continuing in similar proportions, and the number of copies of the *Journal* of the British Medical Association, issued weekly, is now 23,000. It must be remembered, however, that the number of *Journals* issued does represent the number of members of the Association, as it includes, besides those devoted to the members, the sales in the office to private individuals, clubs and such like.

After a few years residence in Great Queen Street, the offices of the Association were moved to 16 L.A. Strand, where increased accommodation was secured for the General Manager and his clerks, and the more public business of the Association. These premises also provided a large room in which the Committee of Council could meet.


The printing of the *Journal* was taken over by the Association in 1879.

This last step necessitated another change of premises, and the establishment of the Association in its present home, in what had hitherto been the offices of the Briton Life Insurance Co., still in the Strand, with a second facade in Agar Street. With necessary alterations, the new premises afforded the space




TORONTO GENERAL HOSPITAL, GERRARD STREET EAST.

The Canadian Journal of Medicine and Surgery.



DR. MEYERS' PRIVATE HOSPITAL FOR NERVOUS DISEASES - DEER PARK, TORONTO.

The Canadian Journal of Medicine and Surgery




HOSPITAL FOR SICK CHILDREN, COLLEGE STREET, TORONTO.
The Canadian Journal of Medicine and Surgery.

required, not only for commodious offices, but for a printing establishment as well.

Over these offices is the library of the Association, which contains some 9,000 volumes; an appointed librarian is in constant attendance. These handsome premises, a photograph of which appears at p. 45 of this issue, are as yet only leased, but it is likely that the Council will not relax its efforts to accumulate a surplus until the British Medical Association is in a position to purchase the freehold of the present property or some equally eligible one.

J. J. C.



SUMMER QUARTERS OF THE HOSPITAL FOR SICK CHILDREN, HASLAN'S POINT, TORONTO.



ST. VINCENT DE PAUL HOSPITAL, BROCKVILLE.
The Canadian Journal of Medicine and Surgery.



GRACE HOSPITAL, COLLEGE STREET, TORONTO.

The Canadian Journal of Medicine and Surgery.



ST. JOSEPH'S HOSPITAL, GUELPH, ONT.



ROSS MEMORIAL HOSPITAL, LINDSAY, ONT.

THE CANADIAN NORTH-WEST, ITS CLIMATOLOGY AND OPPORTUNITIES.


It was suggested by an esteemed Canadian physician, now resident in Oxford, that this journal gather some reliable facts as to the climatology and farming facilities of the great West, and publish them in the interest of those Old Country physicians who are so frequently appealed to by parents in regard to sending sons, as settlers, to this new country. Life in Western Canada is strenuous, health-giving and inspiring, to-day a wilderness, to-morrow a wheat field, the next day a thriving settlement. A




HOTEL DIEU HOSPITAL, WINDSOR, ONT.

perfect network of a railroad system has superseded the long, tedious jaunts by stage or cart. Provided the man is young, strong, willing to work and has sufficient means to establish himself in a cattle ranch or farm, his prospects are good. A man nervous with the strain of a city life, cumbered with a wife and sickly young children, had better far stay in the Old World, where others have ploughed for him and home comforts surround him. Such a settler is unsuited to the life of the West, and a nuisance to the Canadian people, who are too busy themselves building their own "camp fires" to listen to his tale of disappointment.

When we consider that the three Provinces of Manitoba, Saskatchewan and Alberta, and incidentally New Ontario, cover




A FEW OF THE TORONTO CHURCHES.



OSGOODE HALL, TORONTO—LAW COURTS.

The Canadian Journal of Medicine and Surgery.



LIBRARY, OSGOODE HALL, TORONTO.

The Canadian Journal of Medicine and Surgery.

in round numbers 600,000 square miles and embrace within their combined limits the available agricultural and ranching regions of Western Canada, it is easy to grasp the full significance of the exodus to Canada from European countries, the sister Colonies and the United States.


Wrong impressions frequently prevail regarding the climate and physical features of the great Canadian West, but a glance at any map having climatic lines will show that Edmonton has as high an average temperature as St. Paul—that Northern Michigan and Manitoba have similar temperatures—that as we go north-west the climate is modified by the influence of the winds from the Pacific. The mean temperature for July in Winnipeg




ROYAL VICTORIA HOSPITAL, BARRIE, ONT.

is 66 deg., which is higher than in any part of England. The average diurnal range is also much greater than in England, being from a maximum of 78 degrees to a minimum of 53 degrees. This high daily temperature during the growing months, with the long hours of sunshine, matures the crops quickly. In Alberta the climate is dominated by the warm chinook winds. Little snow falls, and both cattle and horses can remain outside the entire winter, living on the sun-cured buffalo grass which covers the plain like a carpet.


The surface of Western Canada slopes to the east with a slight tilt to the north—the rivers finding their way to Hudson Bay. From the mountains on the west to the granite country of New Ontario on the east stretches one vast, alluvial plain. This plain is watered and drained by three great river systems—the Red and



MAIN CORRIDOR, CITY HALL, TORONTO.
The Canadian Journal of Medicine and Surgery.



TORONTO STREET—THE WALL STREET OF TORONTO.
The Canadian Journal of Medicine and Surgery.



NEAR CORNER OF KING AND YORK STREETS, TORONTO.

the Assiniboine in Manitoba, the Saskatchewan in Southern Alberta and Saskatchewan, and the Peace and Athabasca in Northern Alberta. These river systems make of this inland empire one vast network of intersecting valleys. To these topographical features and the mild climate is due the remarkable productivity of the soil.

As regards the soil of North-western Canada, Prof. Shaw, an eminent agriculturist writer, says: "The first foot of soil in the three provinces of Manitoba, Saskatchewan and Alberta is its greatest natural heritage. It is worth more than all the mines in the mountains from Alaska to Mexico and more than all the forests from the United States boundary to the Arctic Sea, vast as these are. And next in value to this heritage is the three feet of soil which lies underneath the first. The subsoil is only secondary in value to the soil, for without a good subsoil the value of a good surface soil is neutralized in proportion as the subsoil is inferior. The worth of a soil and subsoil cannot be measured in acres. The measure of its value is the amount of nitrogen, phosphoric acid and potash which it contains, in other words, its producing power. Viewed from this standpoint, these lands are a heritage of untold value. One acre of average soil in the North-west is worth more than twenty acres of average soil along the Atlantic sea-board. The man who tills the former can grow twenty successive crops without much diminution in the yields, whereas the person who tills the latter must pay the vendor of fertilizers half as much for materials to fertilize an acre as would buy the same in the Canadian North-west in order to grow a single remunerative crop."


Of the three Provinces referred to, Manitoba is the oldest. It is the smallest of the Western Provinces, measuring but 65,000 square miles, yet is as large as England, Scotland and Ireland, and has 27,000,000 acres of arable land, less than one-sixth of which is under plough. The natural resources of the country are as great as those of any other part of the North American continent. The soil is a rich black loam of great strength and depth, that of the Red River Valley being particularly well adapted for the growth of wheat.

The two new Provinces of Alberta and Saskatchewan contain




MURAL DECORATION, CITY HALL, TORONTO.

The Canadian Journal of Medicine and Surgery.



HORTICULTURAL GARDENS, TORONTO.

1. *The Canadian Journal of Medicine and Surgery.*



YONGE STREET WHARF, TORONTO.

The Canadian Journal of Medicine and Surgery.

between them the largest unbroken tract of wheat growing land to be found on the American continent. A large area of desirable free grant land has been opened up for homesteading. The Province of Alberta is situated immediately east of the Rocky Mountains and north of the State of Montana, covering an area of about 253,000 square miles. It is characterized by a mild climate in winter and cool breezes in summer. Its location gives it the benefit in winter of the chinook winds, which follow a north-easterly direction from the current in the southern Pacific Ocean, whence they receive their warmth. The snow in winter rarely lies longer than four or five days at a time, when it is melted by this wind, thus making the winters mild and filling the creeks and ponds with water for the stock on the ranches. In the summer these creeks are constantly supplied with water from the melting snow in the mountains, so that during the summer and winter there is always to be found an abundance of water for grazing and all other purposes. The grain raised in the Edmonton district does but little more than supply local requirements. The cool temperature in summer, with the grasses and pure cool mountain streams mentioned, make Alberta one of the best countries to be found for cheese and butter-making, and it is rapidly becoming as noted for such industries as for its ranches. There are countless herds of fat cattle on the ranges of Southern Alberta, which at any season are neither fed nor sheltered. The ranching industry in Southern Alberta seems, however, to be undergoing a radical alteration. The rancher is giving way to the mixed farmer. Some of the larger men are realizing on their property, and are being replaced by farmers who have some of their land under crop, but keep a herd of cattle as well. The Alberta horse has already become noted for endurance, lung power and perfect freedom from hereditary and other diseases. Thoroughbreds from Great Britain and Kentucky, Clydesdales from Scotland, Percherons from France, and trotting stock from the United States have been imported at great expense, with the result that the young horse of Alberta will compare with any in Canada.

The valley of the Province of Saskatchewan, which extends from the Rocky Mountains to Manitoba, contains some of the most fertile soil in the world. This newly-formed Province



HUMBER RIVER IN WINTER (NEAR TORONTO).
The Canadian Journal of Medicine and Surgery.



A BIT OF ROSEDALE, TORONTO.

The Canadian Journal of Medicine and Surgery.




ROSEDALE RAVINE, TORONTO.

The Canadian Journal of Medicine and Surgery.

embraces a large portion of the valley and extends south to the international boundary. The total area of the Province is about 251,000 square miles. Throughout this district are many thriving settlements, in which are located a large number of prosperous settlers.

The new-comer to the Canadian North-west has the choice of three ways of securing a farm. He may homestead; he may buy land from the Canadian Pacific Railway or other holders; he may rent an already established farm. Good land may be purchased




QUEEN'S HOTEL, TORONTO.

on easy terms and at reasonable prices, ranging from \$5.00 to \$20.00 per acre. Homesteads can still be secured on the outskirts of settlements in districts adapted for mixed farming and stock raising, where hay and water are abundant and timber for building purposes is conveniently obtainable.

The following figures are eloquent of the crop areas and total yields for 1905 in Manitoba, Saskatchewan and Alberta:—

	Area in crop	Average yield per acre.	Total yield bushels.
Wheat	4,019,000 acres	@ 21.6	86,810,400
Oats	1,423,000	" 46.6	66,311,800
Barley	433,800	" 31.	13,447,800
Flax	34,900	" 13.7	478,130



ROYAL MUSKOKA HOTEL.

The Canadian Journal of Medicine and Surgery.

In addition, considerable quantities of peas and rye were produced, the yields being satisfactory, also the usual large crops of potatoes and other roots.

So bountiful are the harvests in the Canadian West that it is now necessary to bring in from Eastern Canada and elsewhere 10,000 to 20,000 farm laborers to work in the wheat fields. These earn good wages and many remain and become actual settlers themselves.

It must not be understood from the above that the Canadian North-west is a wheat country and nothing else. That is far from the case, for most that is produced in a temperate climate grows here abundantly. It is the natural home of the cereal and the garden root; for dairying it has no superior, and it is the stockman's paradise.

It has been well said that, "Next in value to the soil is the heritage of climate." No citizen of North-western Canada should be anxious to apologize for the climate of his country. Good as the soil is it would never have brought supremacy in grain production in this country had it not been for the climate. The blessing of the climate is threefold. It consists in the purity of the air, in the temperature of the same, and in the happy equilibrium in the precipitation. Every one knows the value of the pure air in this country, viewed from the standpoint of health. But does every one know as to the inestimable character of the blessing which pure air proves to the agriculture of the country? It prevents the rapid decay and transformation of the vegetable matter in the soil, and also the too rapid transformation of inert fertility, thus virtually preventing waste in the hand of nature. In this fact is found our explanation of the extraordinary fertility of the soil. The cool temperature of the summer nights is responsible for the large relative yields of the grain. Raise the temperature of the summer days and nights, and the yield of grain will be proportionately reduced. The relatively cool temperature is one of the agricultural glories of this land. The relatively light precipitation is also a great boon to the North-western farmer. It grows his crops and does not destroy them when grown. Nearly every portion of these three Provinces has a rainfall of 15 or 20 inches, enough to grow good crops of grain on



HARVEST SCENE IN THE NORTH-WEST.




REAPING—WESTERN CANADA.



CUTTING GRAIN IN MANITOBA.



"THE STOCKMAN'S PARADISE."



HERDING CATTLE IN ALBERTA.



ON THE SASKATCHEWAN.

farms that are properly tilled, and not enough to waste the fertility of the soil through cracking. In this another reason is found for the wonderful producing power of these lands.

From a consideration of facts relating to the Canadian Northwest, and that during the past year upwards of 150,000 people have entered the great region as settlers, one is inclined to believe that the opening up of the vast territories was, and continues to be, a boon to humanity, since it has enabled man to assert his manhood, in ownership and cultivation of the soil, and has placed within the reach of all opportunities for obtaining health and wealth, a condition of life unknown beneath some other skies.

World-weariness has no place in the great prairie land. The heart must be young, the step buoyant, and the soul of the youth must listen and hear the call of the wild ere it answers.