

**CIHM
Microfiche
Series
(Monographs)**

**ICMH
Collection de
microfiches
(monographies)**



Canadian Institute for Historical Microreproductions / Institut canadien de microreproductions historiques

© 1997

The copy filmed here has been reproduced thanks to the generosity of:




Library
Agriculture Canada

The images appearing here are the best quality possible considering the condition and legibility of the original copy and in keeping with the filming contract specifications.

Original copies in printed paper covers are filmed beginning with the front cover and ending on the last page with a printed or illustrated impression, or the back cover when appropriate. All other original copies are filmed beginning on the first page with a printed or illustrated impression, and ending on the last page with a printed or illustrated impression.

The last recorded frame on each microfiche shall contain the symbol → (meaning "CONTINUED"), or the symbol ∇ (meaning "END"), whichever applies.

Maps, plates, charts, etc., may be filmed at different reduction ratios. Those too large to be entirely included in one exposure are filmed beginning in the upper left hand corner, left to right and top to bottom, as many frames as required. The following diagrams illustrate the method:



L'exemplaire filmé fut reproduit grâce à la générosité de:

Bibliothèque
Agriculture Canada

Les images suivantes ont été reproduites avec le plus grand soin, compte tenu de la condition et de la netteté de l'exemplaire filmé, et en conformité avec les conditions du contrat de filmage.

Les exemplaires originaux dont la couverture en papier est imprimée sont filmés en commençant par le premier plat et en terminant soit par la dernière page qui comporte une empreinte d'impression ou d'illustration, soit par le second plat, selon le cas. Tous les autres exemplaires originaux sont filmés en commençant par la première page qui comporte une empreinte d'impression ou d'illustration et en terminant par la dernière page qui comporte une telle empreinte.

Un des symboles suivants apparaîtra sur la dernière image de chaque microfiche, selon le cas: le symbole → signifie "A SUIVRE", le symbole ∇ signifie "FIN".

Les cartes, planches, tableaux, etc., peuvent être filmés à des taux de réduction différents. Lorsque le document est trop grand pour être reproduit en un seul cliché, il est filmé à partir de l'angle supérieur gauche, de gauche à droite, et de haut en bas, en prenant le nombre d'images nécessaire. Les diagrammes suivants illustrent la méthode.

MICROCOPY RESOLUTION TEST CHART

(ANSI and ISO TEST CHART No. 2)



1.45

1.50

1.56

1.63

1.71

1.80

1.90

2.00


2.25

2.50



APPLIED IMAGE Inc

1653 East Main Street
Rochester, New York 14609 USA
(716) 482 - 0300 - Phone
(716) 288 - 5989 - Fax



DEPARTMENT OF AGRICULTURE

CENTRAL EXPERIMENTAL FARM
OTTAWA, CANADA

PLUM CULTURE

AND

DISTRICT LISTS OF PLUMS SUITABLE FOR ONTARIO AND QUEBEC

WITH

DESCRIPTIONS OF VARIETIES

BY

W. T. MACCUN

Horticulturist, Central Experimental Farm.

BULLETIN No. 43.


JULY, 1903.

PUBLISHED BY DIRECTION OF THE HON. SYDNEY A. FISHER, MINISTER OF AGRICULTURE



TABLE OF CONTENTS.

	PAGE.
Classes and groups of plums.....	7
Plum culture in the provinces of Ontario and Quebec.....	11
Experiments with plums at the Central Experimental Farm.....	11
Seedling varieties.....	12
Cross-breeding.....	13
Propagation by grafting and budding.....	13
Tools and appliances used in pruning and grafting.....	18
The nursery.....	19
The orchard.....	19
Soil and exposure.....	19
Preparation of the land.....	20
Laying out the orchard.....	20
Windbreaks.....	20
Kind of trees to plant.....	21
Planting.....	21
Varieties.....	22
A district list of varieties for the provinces of Ontario and Quebec.....	23
Descriptions of varieties.....	28
American varieties.....	28
European varieties.....	33
Japanese varieties.....	37
List of varieties of plums being tested at the Centr. Experimental Farm.....	33
Pollination of plums.....	40
Pruning.....	41
Fertilizers for the plum orchard.....	41
Cultivation.....	43
Cover crops.....	43
Picking, packing, storing and marketing the fruit.....	44
Canning and preserving American plums.....	45
Spraying.....	46
Diseases and other injuries to plums.....	47
Plum insects.....	51



To the Honourable
The Minister of Agriculture.

Sir,—I beg to submit for your approval Bulletin No. 43 of the Experimental Farm series which has been prepared under my direction by Mr. W. T. Macoun, Horticulturist of the Central Experimental Farm.

The subject treated of is Plum Culture and the information presented has been largely derived from the experience gained in connection with plum growing at the Central Experimental Farm during the past fifteen years. In this bulletin information is given as to the best methods of preparing the soil for a plum orchard, with particulars also as to the planting and subsequent care of the trees. Instructions are also given as to methods of pruning, grafting and on many other topics relating to this branch of fruit industry. Lists of varieties of plums suitable for planting in different parts of Ontario and Quebec are given with descriptions as to the character, quality and time of ripening of each sort. Some of the diseases to which the plum is subject are also referred to and methods of treatment suggested.

A brief account is given by Dr. James Fletcher, Entomologist and Botanist to the Experimental Farms, of some of the more important insects which are injurious to the plum. A chapter has also been prepared by Mr. F. T. Shutt, Chemist to the Farms, on fertilizers for the plum orchard.

It is hoped that the information submitted will be useful to those interested in the cultivation of this fruit in Canada, that it will prove a stimulus to the further planting of plums and that it may aid in making the growing of this useful fruit more successful and more profitable in this country.

I have the honour to be,
Your obedient servant,

WM. SAUNDERS,
Director Experimental Farms.

Ottawa, July 24, 1903.

PLUM CULTURE:

BY

W. T. MACOUN,

Horticulturist, Central Experimental Farm, Ottawa.

If every member of every family in Canada had a plentiful supply of plums of the best quality for home use, and if enough of this fruit were exported profitably from Canada to supply the demands of the people of other lands, there would be less necessity for writing a bulletin on plum culture. But unfortunately this is far from being the reality. Comparatively few people get good plums to eat, especially in the fresh condition, and the export trade in plums, whatever it may be in the future, is still in its infancy. The past experience has been that the consumption of fruit increases almost or quite as rapidly as the supply, providing the latter is properly distributed, and it is not too much to expect that it will do so in the future, at least for some time. The desire for fruit grows on the consumer, and the better the quality of the fruit supplied the greater will be the desire for it.

In order to make this bulletin on plum culture as complete as possible, it has been necessary, in addition to using the information obtained from experimental work here, to correspond freely with fruit growers and to consult many publications and thus get the benefit of the work of others. In this connection, special mention should be made of the recent work "Plum and Plum Culture," by Prof. F. A. Waugh, from which practical and useful book many suggestions have been obtained and several descriptions copied. To Canadian fruit growers I am especially indebted for information regarding varieties and for specimens, as well as assistance in preparing the district lists. In this respect I wish to refer particularly to Mr. Auguste Dupuis, Director of the Quebec Fruit Experiment Stations, Village des Aulnais, Que., who very kindly sent me a large collection of European plums grown in his district, and who gave me very full information regarding the culture of this fruit there and elsewhere in the province of Quebec. Horticultural workers in the United States have also been very courteous and have given me much information, especially in reference to the newer American plums.

I desire to gratefully acknowledge the assistance of Dr. James Fletcher, Entomologist and Botanist of the Dominion Experimental Farms, who prepared the article on "Plum Insects," and I am much indebted also to Mr. F. T. Shutt, Chemist, for the chapter on "Fertilizers for the Plum Orchard."

Some of the matter relating to propagation and culture in this bulletin has been taken from bulletin No. 37 on apple culture by the writer, but changes have been made in this where necessary.

The plum has been cultivated for a very long time, and the origin of the older varieties, like the apple, is unknown.

There are three great classes of plums from which are derived most of the cultivated varieties of to-day, namely, the European, derived from *Prunus domestica*, the Japanese, and the American.

EUROPEAN PLUMS.

This class of plums has reached a higher stage of development than either the Japanese or American, which is due to the fact that they have been under cultivation from very early times and that more systematic and intelligent labour has been spent on their improvement. It is thought that many of the European plums have gradually developed from the Damson, and that the latter originated from the European sloe, *Prunus spinosa*. The Damsons, however, are so distinct in tree and fruit from other European plums that some authorities make a separate group of them. There is such a wide difference between the Damson and the Reine Claude or Green Gage groups that it seems reasonable to suppose that the origin of the two was different, and indeed they have been regarded as different types, at least since the Middle Ages. Both the Damsons and the Green Gages come fairly true from seed. In his book on "Plums and Plum Culture," Prof. F. A. Waugh, who has made a thorough study of the European plums, after making a separate class of the Damsons and a group of the Reine Claude or Green Gages, divides the other varieties into several types. This division is valuable and is adopted here. The varieties described in this bulletin being mostly given as representatives of these types.

Dame Aubert type.—This includes the largest European plums and is represented by Yellow Egg, Golden Drop, and Grand Duke.

Prunes.—Represented by Fellenberg (Italian prune), German Pruno and Raynes.

Perdrigons.—The only variety of this type which is grown in Ontario and Quebec, to the writer's knowledge, is the French Red Perdrigon, introduced by the Trappist Fathers, Oka, Que.

Diamond Type.—This includes some of the most productive, firmest and best shipping varieties and usually those of only medium quality. Varieties of this type are Diamond, Kingston, Quackenboss, Glass, Shipper, Mount Royal and Arctic.

Bradshaw Type.—Under this type Prof. Waugh includes Bradshaw, Victoria, Pond, Field, Duane Purple, Oswego and Giant Prune. These varieties, however, have not as many characteristics in common as those of the previous types.

Lombard Type.—Represented by Lombard, Communia, Voronesh 20 of Budd, Leipsic of Budd, Prince of Wales and Merunka. The Early Red Russian and White Nicholas of Budd would also be included in this type.

Damsons.—The best known varieties of this class are Shropshire, French, Frogmore, and Cluster.

Reine Claude or Green Gage Group.—There are a large number of fine dessert varieties in this group, the principal being Bavay, Green Gage, Lawrence, Imperial Gage, McLaughlin, Jefferson, Washington, General Hand, Bleeker, Peter's Gage and Queen May.

Plums of the European class succeed best in Ontario in the counties bordering on the great waterway southward and eastward from the Georgian Bay to the Thousand Islands. They may be grown very successfully all through south-western Ontario and the Niagara peninsula, and through the central counties to those bordering on Lake Huron. East of Toronto the best varieties are grown within twenty five or thirty miles of Lake Ontario. North and east of these areas only the hardiest kinds succeed well, and but very few are hardy north of latitude 45°.

In the province of Quebec a few varieties give fair satisfaction in the Eastern Townships and a few on the Island of Montreal, and along the Ottawa above Montreal, past the Lake of the Two Mountains. Along the south shore of the St. Lawrence, below the city of Quebec, in the counties of Lévis, Bellechasse, Montmagny, L'Islet, Beauce, Kamouraska and Temiscouata, and along the north shore in the counties of Portneuf, Quebec (Island of Orleans, especially), Charlevoix, most of the finest varieties of European plums can be grown with good success. It is, however, near the river, and where its influence is felt, that these plums succeed best.

The hardiness of the European plums appears to be governed partly by the moisture of the atmosphere, and partly by the temperature. At Ottawa very few varieties will fruit at all, and most of these only bear a good crop once in three or four years. It is not the wood which is killed by winter, although winter killing of the tips does

occur, but the flower buds are destroyed. This killing of the buds appears to be due partly to the dry, cold weather of winter, and partly to spring frosts. Good crops of European plums have been produced at Ottawa after the temperature had been lower than 20° below zero the previous winter, showing that temperature alone is not the cause of failure. Along the south shore of the St. Lawrence, below the city of Quebec, where the temperature falls 30 degrees Fahr. below zero, good crops of the best varieties of European plums are regularly produced. The moisture of the air in this case appears favourable to the preservation of the flower buds, even though the temperature is low. There is a great difference, however, in the hardiness of the different varieties, and in time seedlings may be produced which will have flower buds capable of surviving the coldest temperatures which occur in these districts in any winter.

JAPANESE PLUMS.

Owing partly to their novelty, partly to their vigour, productiveness and good shipping qualities, and partly to the way they have been boomed, the Japanese plums have been widely planted on this continent. Although introduced into America as recently as 1870, this class of plum is now almost as well known as the European. The origin of the Japanese plum is unknown, but it is thought to be a native of China. The Japanese were not very systematic in their nomenclature, and when the plums were introduced into America there was great confusion of names. They have, however, been pretty well worked out, and the best known and most profitable kinds can now be obtained true to name. Many seedlings have been grown in the United States and a large number of hybrids originated, some of which have great merit.

The fruit of most of the varieties is only medium in quality: a few, however, are good, and a still smaller number very good. The productiveness, firmness, appearance and good keeping qualities are what make these plums valuable. The majority of the Japanese plums are not quite as hardy in the flower bud as the hardiest of the European, but a few seem to be quite as hardy. They bloom very early, and on this account are more likely to be injured by spring frosts.

The Simon or apricot plum *Prunus Simonii*, Carr., though not a Japanese plum has some of its characteristics and may be classed with the Japanese plum here. This plum is thought to be a native of China and was introduced into America from France about twenty years ago. It has not been planted to any extent in Canada, nor is it widely grown in the United States. The tree is productive and the fruit is large and handsome, though inferior in quality. The tree is quite productive, and the fruit keeps and ships well. It does not appear to be any hardier than the Japanese plums. It has been grown with success in south-western Ontario, but is not a desirable plum to plant.

AMERICAN PLUMS.

These plums have a wide range on the American continent, being found wild from Mexico north to the province of Manitoba, and from the Atlantic to the Pacific oceans. They are represented over this great area by seven distinct species and six recognized groups or types. Of these, the cultivated varieties originated from the American wild plum *Prunus americana*, Marsh, and the Canada plum *P. nigra*, Ait. include nearly all the American plums that are profitably grown in the provinces of Ontario and Quebec. The more southern groups, of which a few varieties are partially successful, are: the Miner group, *Prunus hortulana mineri*, Bailey, which is closely related to *Prunus americana*, but has some resemblance to the Wildgoose group; and the Wildgoose group *Prunus hortulana*, Bailey. It is, however, only in the milder parts of the province of Ontario that they are even fairly satisfactory. At Ottawa, the flower buds are injured and the crop light. The varieties of the Wayland and Chicasaw groups of plums,—two other important groups—are too tender for most districts in eastern Canada.

Americana Group.—This group has, up to the present time, furnished the best varieties of American plums. The range of *Prunus americana* is given by Waugh as being New Jersey and Ohio to Minnesota, Montana and Colorado. The tree is a

spreading grower, sometimes reaching a height of 20 feet. On account of its spreading habit the trunk and large branches are often split or broken by winds when laden with fruit or by heavy snow storms in winter, and this is a serious drawback to some varieties especially. The varieties of this species bloom later than the Canada plum *P. nigra*, and sometimes escape frost from which the native species suffers, an example of which occurred in Ottawa in 1902. The trees are very productive and the fruit is much improved in size by thinning. The fruit varies greatly in size and the colour ranges from yellow to red. The skin is generally tough, and often thick and astringent, but the flesh is usually firm or moderately firm, very juicy, sweet, and sometimes rich and high flavoured. The stone usually clings, but occasionally is almost or quite free. When growing wild, this tree is found thriving best in rich and rather moist soil, but under cultivation it succeeds very well even where the soil is comparatively poor and not very moist. It is only about 50 years since the Americana plums were thought worthy of cultivation, but during the past few years the cultivated and named varieties have increased so fast that there are now over 200 of them. The size, appearance, and quality is also very much improved. Indeed, the improvement is remarkable considering the short time in which it has been made, and we may hope for still greater improvement yet. The appearance of the plums is all that could be desired, and the best varieties are almost large enough, but there is great room for improvement in the character of the skin and flavour of the fruit, although the latter is good. Earlier varieties are also wanted. The present fruiting season of the Americana plums at Ottawa is from the fourth week of August until about the last of September. Some varieties of this group are Bixby, Wolf, Hawkeye, Stoddard and De Soto.

Nigra Group.—From this group of plums, of which *Prunus nigra* is the species, will probably be originated the varieties which will be of greatest value in the northern parts of Ontario, and the coldest parts of the province of Quebec. This is the wild plum of Canada, having a range from the Maritime Provinces westward and northward to Assiniboia. Although in some places this species can scarcely be distinguished from *P. americana*, into which it seems to merge in some districts, in the colder parts of the provinces of Ontario and Quebec the tree is very distinct, and for this reason we prefer considering it a separate species although some good authorities make it merely a variety of *P. americana*. It is a more upright grower than *P. americana* and the wood is darker and tougher. The tree seldom breaks down like *P. americana*, which is a great advantage. It blooms earlier and has larger flowers than the Americana. The flowers also have a pink tinge, especially when opening. The fruit ripens early, and on this account is sometimes more profitable, as it can be marketed when there is little competition with other plums. Although the named varieties which are now upon the market are not as high in quality as the best of the Americana group, some of the wild seedlings are fully as good in quality, though not as large. The skin of the Canada plum is thinner than the other species and breaks up easier when cooked. The colour of the fruit varies almost as much as *P. americana*, but is more often entirely red with little or no bloom. The shape varies, but the fruit is more regularly oblong and oval than *P. americana* and is not flattened like that species. The Canada plums begin to ripen early in the second week of August and the season extends to September. Very little has yet been done either in the United States or Canada to improve *P. nigra*, but as good results are likely to be obtained as with *P. americana*. If a plum can be originated which will ripen by August 1, the season of the hardy American plums will cover two months. Some of the varieties of this group are, Arkin, Odegard, Cheney.

HYBRID PLUMS.

During the past ten years a number of hybrid plums have been placed upon the market. Most of the more prominent of these were originated by Luther Burbank, of California, who has devoted much time to this work. Few of his hybrids, however, which are now on the market will be likely to prove valuable in the colder parts of Ontario and Quebec, as the parents of most of them are varieties which do not prove successful where the climate is severe. Some of them may be of value in the warmer parts of Ontario where the Japanese plums do well.

There is a wide field for work in plum hybridization. If blood of the European and Japanese varieties can be introduced in the Americanas and Nigras and the hardiness of the latter maintained, plum culture in the north will receive a great impetus. It is not too much to hope that this will soon be accomplished.

PLUM CULTURE IN THE PROVINCES OF ONTARIO AND QUEBEC.

The plum has been most largely planted in the Niagara peninsula, where it grows to perfection. From Hamilton to Niagara, orchard after orchard of this fruit may be seen, and in the spring when in full bloom and in the autumn when laden with fruit the trees are a sight to behold. But plum growing is by no means confined to the Niagara district. All through South-western Ontario and along Lake Huron and the Georgian Bay, plums succeed well and are grown in large quantities. Indeed, the Georgian Bay district appears quite equal to the Niagara peninsula for plum growing. Plums are also grown very successfully from Hamilton eastward within thirty miles of Lake Ontario and the St. Lawrence River, to about the city of Kingston. Some of the best plums also succeed in the midland counties of Ontario. North and east of the districts mentioned, plum growing is confined mainly to the Americana and Nigra varieties, though a few of the hardest European kinds will prove fairly satisfactory in some of these northern sections.

In the province of Quebec plums are not grown to a very large extent, although in some parts the best European plums can be grown very successfully. On the Island of Montreal a few of the best kinds are grown, but they are not thoroughly satisfactory, as the flower buds are frequently killed by winter. There are, however, some European seedlings originated on the island which give good satisfaction. The best district for growing the European plums in Quebec is probably along the south shore of the St. Lawrence, below the city of Quebec in the counties already mentioned.

Here even the more tender European kinds appear to do well, and Mr. Dupuis informs the writer that the season begins in the first half of August with Favorite and Mirabelle and that Grand Duke and Coe's Golden Drop will keep in good condition until December. In the Eastern Townships, a few of the hardest European kinds are fairly satisfactory, but the climate is too severe to grow them profitably on a large scale. In the province of Quebec, with the exception of favoured districts along the St. Lawrence River, and perhaps along the Bay of Chaleur, the Americana and Nigra plums will, as a rule, be the most profitable kinds to plant, unless hardier European or hybrid kinds are originated.

The profits from plum growing have been good in the past, but such great numbers of trees have been planted in recent years that unless a large part of this fruit can be safely exported the market is likely to be soon glutted. Very early, very late, and good shipping varieties will probably be the most profitable kinds to plant in the future. When a market is glutted, fruit of the best quality will command the highest prices, provided it is in good condition and of fine appearance, and this fact should also be taken into consideration when planting.

EXPERIMENTS WITH PLUMS AT THE CENTRAL EXPERIMENTAL FARM, OTTAWA.

Experiments with plums were begun at the Central Experimental Farm in 1888 and have been continued ever since. The testing of varieties to determine their hardiness, productiveness, and other qualities has been one of the most important experiments, and a large number of varieties have been tested. Experiments have also been carried on with various stocks, to learn which was the most satisfactory for the various classes of plums, and different methods of grafting have also been tried. The spraying of the trees with various mixtures and solutions to control fungous diseases and insect pests has also been an important part of the work. Careful records have been kept of the dates of blooming of the different varieties, and the information thus obtained is

very useful, as it has been proven that few varieties of American plums are self-fertile, and it is thus necessary to have, in most cases, some other kind blooming at the same time in order that fruit may set well. A table showing the blossoming period of the different varieties will be found in this bulletin. The yields from each individual plum tree are kept separate, and it is thus possible to tell whether one tree is bearing better than another. Much work has been done in testing and originating seedling plums, especially of the American varieties, and a few very promising sorts have been produced. Some experimental work in cross-breeding has also been carried on.

The soil in the plum orchard is not as suitable for plum growing as it might be, being a light, sandy loam, but good plums are raised. The trees were originally 20 by 20 feet apart, but interplanting has been carried on and the trees are now 20 by 10 feet apart. There are now 569 trees in the orchard, consisting of 459 trees of named varieties and 110 seedlings. In addition there are a large number of seedlings planted 10 by 10 feet apart.

Following are the number of varieties of each group or class in the orchard and nursery:—


Group or Class.	Number of Varieties.
European.....	68
Japanese.....	11
Americana.....	100
Nigra.....	11
Miner.....	13
Wayland-like.....	3
Wildgoose.....	10
Hybrid.....	31
Total varieties.....	247

Seedling Varieties.

While there are many fine named varieties of plums which succeed in the more favourable parts of the provinces of Ontario and Quebec, there is always the possibility of getting something better. In the colder parts of the provinces where few of the better plums succeed, there is a fine opportunity for developing hardier and better kinds. One of the easiest and best methods of obtaining new varieties is by growing seedlings from the stones of the best plums which have ripened in the district where new kinds are desired, and if no plums have yet been grown there stones should be procured from the nearest place where they can be obtained. The European plums have been improved for so many centuries that it will be difficult to obtain a seedling plum of this group which will be better in quality than the best now in cultivation, but the prospects for obtaining trees with hardier fruit buds are very bright. The native and Americana plums have been, relatively speaking, little improved as yet, and there is a wide field for development here. Stones should be saved from the largest plums from the most productive tree of the variety of which seedlings are desired. These stones should, when possible, be planted immediately after the fruit is ripe, as if the stones become dry they will not germinate as well, and sometimes will not grow at all. If they are planted at the time the fruit is ripe it is not necessary to remove the pulp. If the stones cannot be conveniently planted at this time they should not be allowed to become dry. They may be kept over winter in boxes mixed with sand, which should be moist but not wet. A layer of sand about one inch in thickness is put in the bottom of the box and is merely covered with a layer of stones, the latter are then lightly covered with sand and another layer of stones spread on top, and so on until the box is filled. This is called stratification. The box should be buried outside where there is good drainage and no danger from small animals, or kept in a cellar. If the stones freeze when they are thus moist in sand they will crack and seedlings grow quicker in the spring, but there is a danger of their drying out when exposed to the frost unless the box is buried. Stones spread outside in the autumn in a well drained place and lightly covered with soil will often come through the winter in fine condition.

The stones should be planted not more than one inch deep, (often they are planted too deep,) in good loamy soil, in rows three feet apart and dropped from one to two

inches apart in the rows. These usually germinate in the spring, after which the soil should be kept thoroughly cultivated during the summer to induce a thrifty growth. The next spring or the one following, the young trees should be planted out about ten feet apart or ten by five feet and left to fruit, which they will do in from three to six years. Scions may then be taken from the promising varieties and grafted. Stones planted from the best of these will be likely to produce something still better. As many of the seedling plums, especially of the Americana and native varieties are as good as their parents, the ground on which they are growing is not lost as the fruit can be sold to advantage.



European plum top grafted on Nigra plum, three feet from ground.

CROSS-BREEDING.

New varieties of plums may be obtained by cross-breeding, and by this method one is even more likely to obtain the kind of plum desired, but the time for this work is so limited, being only a few days when the flowers are opening, that only specialists can very well undertake it. The method of crossing is explained in Bulletin No. 37 on Apple Culture.

Propagation.

The plum is propagated principally by budding and grafting, although a few varieties will strike more or less readily from cuttings, and some kinds when on their own roots are increased from suckers.

STOCKS.

While it has not yet been clearly proven that the stock on which a scion is grafted will materially change the flavour or season of the fruit, it does affect the vigour and fruitfulness of the tree in a greater or less degree. If a scion is grafted on a dwarf stock the tree will be dwarfed and will come into bearing sooner than if grafted on a thrifty stock as anything which checks the growth of the tree promotes early fruitfulness. It is possible, however, especially in top grafting, to have such a slow growing stock that the graft outgrows it too much and the tree becomes top heavy and if it does not die owing to a poor circulation of sap, the graft is liable to be broken off by wind. The tree in the accompanying cut was killed because there was not a free circulation of sap. The stock also, if it is tender, may be winter killed and a tree which may be perfectly hardy

above ground is ruined by being grafted on such a stock. Nurserymen find that the scion influences the root system of the stock, but it is not so clear that the scion or graft makes the stock any hardier.

If a tree is planted deep enough, roots may be thrown from the scion and the tree will eventually be on its own roots but this should not be depended on as a rule, and a stock should be used that will be hardy, give a good union, and make a thrifty tree.

The two main stocks used in grafting and budding the plum are the Myrobolan and Marianna.

Myrobolan.—This is a European plum and is imported principally from France. It unites readily with the scion and throws few suckers, which make it a very desirable stock, where the winters are not very severe. It is used very generally in America, but more in the north than in the south where the Marianna is chiefly used. The Myrobolan plum is not a satisfactory stock for the colder parts of Ontario and Quebec, as it is liable to winter kill.

Marianna.—The Marianna plum is used largely in the Southern States as stock, where it propagates freely from cuttings. It is thought to be a hybrid between the Myrobolan and the Chickasaw plums. This, also, is not a desirable stock for the colder parts of Ontario and Quebec.

Peach.—The peach unites readily with the plum and is used in the United States as a stock for it. It can be grown cheaply, and strong young stocks are readily obtained for grafting and budding. It has, however, the disadvantage of not being hardy enough in many parts of Canada.

St. Julien.—The St. Julien is a European stock that is used to a limited extent in America for propagating the European plums, but the Myrobolan and Marianna stocks can be obtained so cheaply that it is not used much now. The St. Julien is, however, the safest stock for European plums in the north.

Americana and Native.—Americana and Native plum seedlings furnish the best stocks for the colder parts of Canada. The young trees make strong growth and are very suitable as stocks for grafting and budding. This stock is not usually satisfactory when the European plums are top grafted on it, as the top outgrows the stock and either breaks off from being top heavy or dies from lack of nourishment, as the native varieties, especially, grow much slower. No bad results, however, have followed from root grafting the European plums on Americana stock, and good thrifty trees have been obtained but the trees are still young.

Sand Cherry (Prunus pumila).—The Americana plums have been successfully root grafted on the Sand Cherry at the Experimental Farm, and trees which have been grafted 10 years are still in good condition, with a perfect union and bearing well. The trees are considerably dwarfed by this stock. The Sand Cherry may prove very useful where close planting is adopted, as many more trees could be planted on an acre when dwarfed by this stock. Trees grafted on this stock, however, for ten years are not as firm in the ground as we should like, and strong winds have a tendency to loosen them. This may be a serious disadvantage when the tree gets older.

BUDDING.

The favourite method of propagating plums is by Shield budding, and the best season for doing the work is in late summer, some time during August being the best time in Ontario and Quebec. At Ottawa the trees have been found in good condition during the second week of August. Young stocks one or two years old are the most satisfactory.

Budding is best performed when there is still sufficient sap beneath the bark to permit of the latter being easily raised with a knife. On the other hand, if the work is done when the tree is still growing vigorously the bud is liable to be 'drowned out,' or, in other words, forced out by reason of too much sap and growth of the stock.

The stock which is to receive the bud should be at least three-eighths of an inch in diameter near the ground. The lower leaves are rubbed off to a height of five or six inches to enable the bud to work more freely. A perpendicular cut is now made in the stock as near the ground as possible from an inch to an inch and a-half long and preferably on the north side of the tree, as the bud will not be so readily dried out by the sun on that side. The cut should only extend through the bark. Another cut should now be made across the top of the perpendicular one. The two cuts when made will appear thus: T

The buds are cut from well developed and ripened shoots of the current season's growth of the variety it is desired to propagate. Before the buds are removed the leaves should be cut off the shoots; a piece of the petiole or leaf stem is left, however, by which the bud may be handled after it has been removed. A very sharp, thin-bladed knife is necessary in removing the bud. Knives are specially made for this purpose. The bud is cut off the shoot downwards or upwards, whichever is more convenient, the general practice, however, is to cut upwards. The length of the piece removed with the bud should be about one inch long, and the cut surface smooth. It should be quite thin, as but little of the wood is taken with the bud. The buds or

twigs should be kept where they will not dry out while the work of budding is going on. The bud is inserted under the bark by raising the latter with the blade of the knife or the part of the budding knife made for that purpose. The bud is then pushed down and under the bark with the fingers, and finally the piece of leaf stalk which was left when it was removed from the twig is pressed with the blade of the knife to bring the bud into the proper position. The bark on each side of the bud, which should now be under the bark of the stock, will hold it in position. In order to bring the bud and stock into close contact and prevent the former from drying up before the union takes place, they should be tied tightly



Example of Shield-budding.

together with raffia or some soft string, taking care not to cover the bud with it. The bud should unite with the stock in two or three weeks, and after that time the string should be cut, as otherwise the bud may be injured. If the proper season has been chosen for the work the bud should remain dormant until the spring. If it starts in the autumn it may be killed during the winter. In the following spring the stock should be cut off just above the bud which will cause all the strength of the stock to be directed into the bud and produce rapid growth, three feet not being an exceptional growth for the first season.

Budding is now a very popular method of propagating plums. The first season's growth is greater than from root-grafted trees and there is a larger proportion of straight trunked trees by this method. If it is desired also to prevent trees from growing on their own roots, budding is preferable, as trees propagated in this way may be planted so that the stock is just at the surface of the soil and all roots are thrown from it.

Buds may also be inserted in the branches of trees with good results. When the buds have united and grown the top may be shaped up as if top grafted.

GRAFTING.

Scions.—As much of the success in grafting depends on the condition and quality of the scions, too much stress cannot be laid on the importance of having them of the best quality and in the best condition at the time of grafting.


Scions may be cut any time after the wood is well ripened in the autumn and before the buds begin to swell in the spring. The best time, however, is in the autumn, as they may then be kept in the condition desired. If they are cut in cold weather, in winter, trees from which they are taken may be injured if large numbers are removed from them, as the bark is liable to split, there is less sap also in the scions at that time and thus the chance of their drying up is greater than if they were cut in the autumn. One cannot tell very well, either, in winter whether the young wood has been injured or not. Scions should be cut from healthy, bearing trees. The wood of old trees is liable to be diseased, and if diseased wood is used it is likely to produce a diseased tree when grafted. Scions should also be cut from the most productive trees. Occasionally, one or more trees of a variety will produce more and heavier crops than the others. If scions are taken from these trees, the probability is that a larger proportion of the grafted trees will produce crops like the trees from which the scions were taken than they otherwise would. The scions should be cut from the wood of the current season's growth, as older wood is not satisfactory. The buds should be well developed and the wood thoroughly ripened. It is not wise to use the water sprouts or young shoots which spring from the main branches or trunk for this purpose. They may not be thoroughly ripened, and it is also possible that sprouting propensities may be thus more developed in the grafted trees. The entire season's growth may be cut off and packed away until required for grafting, when it should be cut into pieces from four to six inches in length having three well developed buds.

Scions may be kept in good condition in moss, saw-dust, sand, or forest leaves. The last named are found very satisfactory at Ottawa. These materials should be slightly moist, but not wet; the object being to keep the scions fresh and plump without there being any danger of their rotting. They should be kept in a cool cellar which is not too dry, and should remain dormant until ready for use.

Root Grafting.—Plums are propagated successfully by root grafting, although budding is more general and gives, as a rule, better results. Strong one year old or two year old stocks are heeled in during the autumn in a cool cellar in moist sand. Grafting may be done any time during the winter, but it is usually not started until January or February. Whip or tongue grafting is the method usually employed. As only the root is required, the trunk and branches are cut off and thrown away. As there is but little advantage in using the whole root, it may be divided into several pieces, much depending on its size. Each piece should be at least four inches long. A smooth, sloping cut upwards, about two inches long, is made across the main part of the root most suitable to receive the scion. The scion is prepared by cutting off a piece of the wood procured for this purpose in the autumn from four to six inches long and with about three well



Example of Root-Grafting.



COTTBELL PLUM IN BLOOM—Typical Tree of *Prunus Americana*.

184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000

developed buds on it; a smooth, sloping cut downwards and across it is now made of about the same length as that already made on the stock. Clefts are now made in the sloping surface of both scion and stock, in the former, upwards; and in the latter downwards. They are then joined together by forcing the tongue of the scion into the cleft of the stock. The inner bark, or cambium, of both scion and stock, should be in contact with one another on at least one side of the graft, as it is at this point of contact where the union begins to take place. In order to ensure a speedy and successful union, waxed cotton thread is wound tightly around to hold the parts together. Amateurs are also advised to rub grafting wax all over where the two parts are joined, as with this treatment, success is likely to be more certain.

The operation having been completed, the grafts are packed away in moss or saw-dust until spring. They are then planted out in nursery rows about three feet apart and one foot apart in the rows, the point of union being about three inches below the surface of the soil. The ground should then be kept thoroughly cultivated throughout the season.

Crown Grafting.—Crown grafting is usually done on young stocks in the nursery row in the spring. The trees are cut at or just beneath the surface of the soil at the crown or collar. A sloping cleft is then made in the side of the crown and a scion, cut wedge-shape at the lower end, is inserted in the cleft. The same precautions should be observed as in root-grafting, of having the inner bark of both stock and scion touching on at least one side. The grafted part should then be well covered with grafting wax, in order to exclude the air. The trees usually make a strong growth when grafted in this way, but as the work has to be done in April before growth begins it is often inconvenient to do it at that busy season of the year.

Top Grafting.—Plum trees are not top grafted as frequently as apple trees, but they can be very successfully grafted in this way. When there are trees which produce poor or unprofitable fruit they may be made to bear good fruit by top grafting other varieties upon them. An unsymmetrical top may also be improved by top grafting. European or Japanese varieties should not be top grafted on Americana or Nigra stock. In our experience at the Central Experimental Farm it has been found that although a good union is made the European will outgrow the Americana so much that the top will die a few years after grafting, the trunk of the stock expanding too slowly. It may be mentioned here that it is a good practice to slit the bark of the stock in top-grafted plum trees if there is an indication of its getting hidebound. In top grafting plums it is best to have both stock and scion as nearly related botanically as possible. Top grafting is done in the spring before growth begins, and early grafting is more important with the plum than with the apple. As the shock to a large tree would be very great if all the branches, on which leaves develop, were cut off the first season, about three years should be devoted to changing the top of the tree. Cleft grafting is the method usually adopted in top working plum trees, it being simple and satisfactory.

The branches to be grafted should not exceed one inch and a half or two inches in diameter. If they are larger, it is so long before the stub heals over, that disease may set in. It is possible, however, to graft larger branches by putting in more scions. The top grafting of a large tree should be done with a view to having the new top as symmetrical as possible, and great care should be taken in selecting the branches to be grafted upon. After the branch is sawn off it is cleft by means of a mallet and strong knife to the depth of an inch and a half to two inches. It is held open to receive the scion by driving a wedge in it. Scions for use in top grafting are cut from dormant wood which has been kept in good condition in the manner already described. They should have about three strong buds and be cut wedge shape at the base, one side, however, being a little thicker than the other. Two scions are now inserted in the cleft of the stub, with the wide side of the wedge on the outside, and thrust down until the lowest bud is almost on a line with the edge of the stub. The inner bark of both scion and stub should meet at some point, so that the union will take place readily, and this is more easily effected if the scion is given a slightly outward slope when inserted. When the wedge has been withdrawn from the cleft the advantage of having the wedge-shaped end of the scion thicker on one side will be apparent, as it will be held much

more tightly than if both sides were the same. If the scion is not a tight fit all along, there is something wrong in the way it has been cut or the stub has been cleft. The cut parts should now be covered with grafting wax to exclude the air and hold the scion in place. Cotton is also sometimes wrapped around the wax in order to more effectively hold the scion in place. If both of the scions grafted on a stub should grow, the weaker one should be removed after the other is well united and the surface of the stub at least partially healed over.

It is often desirable to top graft young trees, and this may be done very readily. The main branches are cut back to within a short distance of the trunk, and the scions grafted on, either by cleft or whip grafting. The closer the grafted part is to the trunk, the better, as the tree will be stronger than if the union occurred further out on the limb, since the growth of graft and scion may not be equal. It is possible to cut off the whole top of the tree and graft successfully on the main trunk, when the tree is young, but unless one is sure that the union will be perfect and the top not outgrow the stock, it is better not to run the risk of losing the tree. Furthermore, if the whole top is cut off there will be such a growth the first season that the scions are liable to get broken off. In top grafting a young tree that has been planted from three to five years, it is better to take two seasons to do the work, as the results will usually be a rule, more satisfactory.

It is necessary to examine the grafted trees during the summer and remove any young shoots from the stocks which are interfering with the scions. It is not wise, however, especially when the tree has been cut back severely for grafting, to remove all the shoots until the grafts have grown considerably and furnish a good leaf surface.

While grafting implements and appliances are numerous, the work can be done with a few, and as it is not often convenient for the farmer or fruit

grower to get a large outfit, only the really necessary things are mentioned. These are:—A sharp, fine-toothed hand saw, to be used for sawing off large limbs, or for making the stubs on trees to be top-grafted where the limbs are too large to be cut with the pruning knife.

A strong pruning knife for cutting the smaller limbs; for smoothing the wounds made by the saw or pruning shears; for trimming off torn edges of branches, and for pruning roots of young trees when planting.

A budding knife, with a thin steel blade, for removing buds, having an ivory handle which is made thin at the end and is used for raising the bark.


A grafting knife, which is used in top-grafting trees. Home-made grafting knives can be easily made. A strong, sharp blade is the chief requisite.

Pruning shears, which are intermediate in their uses between the saw and the pruning knife. They are used for cutting off branches which are too large for the latter and too small to need the saw; for rough pruning and for cutting scions.

A wedge and mallet are also necessary in top grafting large trees.

Raffia, which is one of the best tying materials. It is very strong and very pliable and is particularly useful for bandaging when budding.

Cotton yarn, which is used for tying root grafts and is one of the most satisfactory materials for the purpose. The size known as No. 18 knitting cotton is the best. It is bought in balls, which should be soaked for a few minutes in melted grafting wax before using. The yarn may also be drawn through melted wax, which ensures its all being thoroughly soaked, and is, perhaps, on this account preferable to soaking the ball.



GRAFTING WAX.

There are many kinds of grafting wax recommended, but it is unnecessary to enumerate them all. One of the cheapest and best is that recommended in *The Horticulturist's Rule Book*, under the name 'Reliable Wax,' the receipt of which is as follows:—

'Reliable Wax.—Resin, 4 parts, by weight; beeswax, 2 parts; tallow, 1 part. Melt together and pour into a pail of cold water. Then grease the hands and pull the wax until it is nearly white. One of the best waxes for either indoor or outdoor use.' This should be heated before using if too hard.

The principal value of grafting wax is to exclude air from the wound, and thus prevent the wood from drying before a union takes place. A good grafting wax should not crack when on the tree, else the air will reach the wound and the wax prove of little value. Many materials may be used instead of grafting wax for this purpose, one of the simplest being a mixture of clay and cow dung, but grafting wax is much to be preferred. Strips of cotton are often used, especially in top-grafting and crown-grafting, for wrapping around the wound after the wax has been applied for the purpose of helping to exclude the air, and also to assist in holding the scion in position until the union takes place. This cotton is unnecessary if good grafting wax is used; but if a very valuable variety is grafted it is safer to use the cotton, as when the growth of the scion is rapid there is a chance of its getting broken off during the first season before it is thoroughly united with the stock. Large wounds on trees should be covered with some material that will protect the cut surface from the weather, prevent disease from setting in, and which will not peel off easily. A good dressing of lead paint is probably the best material to use for this purpose. Grafting wax may be used on smaller branches.

THE NURSERY.

Although, as a rule, it will be the most convenient plan to buy trees from the professional nurseryman, yet he who propagates plum trees by root-grafting, crown-grafting, or budding, for his own use, should have a nursery in which to grow them until they are ready for the orchard. A good sandy loam soil, which does not bake and is well drained, is best suited for this purpose, and will grow the strong, healthy trees which are desired. The ground should be thoroughly prepared and the young trees planted about 12 inches apart, in rows from 2½ to 3 feet apart. Cultivation should be thorough up to about the middle of July, when it should cease, as in colder climates, especially, it is very desirable that the wood ripen well, and late cultivation would encourage late growth. It will be necessary the first year the grafted or budded trees are growing in the nursery to go over them carefully and cut out any shoots which may be coming from the stocks, and also to reduce the graft to one stem should more develop. If any side branches grow, however, they should be left intact. In small nurseries it is sometimes advisable to tie the young trees to stakes the first season. This will make them straighter and will help to keep them from being broken. These trees may be planted in the orchard the following spring if one-year old trees are to be used. By the end of the second year or the beginning of the third, after the branches have been pruned to the proper height and the tops shaped, the trees will be in the best condition for planting in the orchard.

THE ORCHARD.

Soil and Exposure.—Plums will succeed well on a great many kinds of soils, but some groups appear to succeed better on certain soils than on others, the best soil depending somewhat on the climate in which the plums are grown. In those sections of Ontario where the European plums succeed best, well drained clay loam has given the most satisfactory results. Along the south shore of the St. Lawrence, below the city of Quebec, where the European plums succeed well, these plums do better on sandy loam.

soils. The Japanese plums on the whole give better results on warm loamy soils than on clay loam. The Americanas and Nigra plums succeed best on clay loam soil, but also do well on sandy loam. All soils should be well drained or success need not be expected. The more severe the climate in which the plums are grown the warmer the soil should be.

If there is danger from spring frosts a northern or north-eastern exposure would be likely to give best results, as the flower buds would not develop as soon as on a southerly exposure. The flower buds of the European and Japanese plums suffer badly in the north and there is no doubt that a northerly exposure would be best for these plums.

Preparation of the Land.—It very often happens that the farmer or fruit grower suddenly decides to plant an orchard. No previous thought had been given to the matter, or if there had, nothing was done to get the land into better condition for the young trees. The trees are bought, the land hastily, and not very well, prepared and the trees set out to take their chances. No after cultivation will fully make up for neglect of the thorough preparation of the land. Trees should begin to grow thriftily from the time they are planted if they are to obtain a good size before they begin to bear heavily, and if the land is not thoroughly prepared and in good condition when they are planted, growth is likely to be slow. It is much better, if one has no land in good condition, to delay planting a year, and give the soil the necessary attention. The time will not be lost, as the trees will do much better. Land which has been well manured for root crops, ploughed in the autumn, and again ploughed in the spring and thoroughly levelled and pulverized with the harrow should be in good condition for planting the trees. If the subsoil is near the surface the subsoil plough should be used after the ordinary one, loosening the soil from four to six inches deeper than the former.

Sod land ploughed in the autumn, top dressed in the spring with a good coating of barn-yard manure and then ploughed again and thoroughly pulverized with the harrow, should also bring the soil into good condition. A green crop, such as clover ploughed under in the spring and the land thoroughly harrowed, would also be a very good method.

Laying Out of the Orchard.—Plum trees require thorough spraying, and this should be taken into consideration when planting so that the trees will not be set too close. Trees should also have abundance of sunlight to thrive best and produce fruit of good colour, and they cannot obtain this if they are too crowded. There are several good methods which may be adopted in laying out the orchard.

If the trees are planted the same distance apart each way with a view to leaving them all as permanent trees, they should be planted from 15 to 20 feet apart, depending on the varieties chosen. If, however, the branching and upright growing varieties were mixed as might be necessary for good pollination, 18 feet apart would be a very satisfactory distance to plant.

Another good method is to plant the trees a greater distance apart one way than the other. This is a satisfactory system when properly carried out. By this plan trees may be planted successfully 10 feet apart in the rows with the rows 15 feet apart, the latter distance leaving ample room for spraying. When the trees become crowded, every other one may be taken out, thus leaving the permanent trees 20 by 15 feet apart.

A third method is to plant the trees in an apple orchard with the object of getting some profit from the land before the apple trees come into full bearing.

If the permanent apple trees were 35 by 35 feet apart, a row of plum trees 17½ feet apart in the row could be planted between the rows of apple trees. Plum trees could also be planted between the apple trees in the rows. This would leave the plum and the apple trees 17½ feet apart. Planted in this way, large crops of plums can be produced before it is necessary to cut the trees out.

Windbreaks.—If the orchard is not naturally protected from the wind by trees or by rising ground, a windbreak may be planted with good effect along the north and west sides, or any other side from which the greatest injury comes, the object being not to stop the wind altogether, but simply to check its velocity, as if a windbreak is high and very dense it stops the circulation of air in the orchard to a large extent, and this gives

very favourable conditions for the spread of both insect pests and plant diseases. On the other hand, a proper windbreak lessens the force of the wind and thus protects the trees, which will grow straighter and shapelier; it will also very materially lessen the amount of windfalls, and it will permit of growing varieties which will not succeed under ordinary exposure. Wind is one of the most important factors in drying out the land and causing drought. If its force is checked by a windbreak the evaporation of moisture from the soil will not be so great.

One of the best trees to plant for a windbreak is the Norway spruce (*Picea excelsa*). It is a rapid growing evergreen and is hardy almost everywhere where plums can be grown successfully. A single row of these trees planted from 8 to 10 feet apart is quite sufficient. They should grow, if properly cared for, at the rate of from 2 to 3 feet a year until they reach a height of 50 to 60 feet. In very exposed places it may be desirable to plant two rows of trees, the trees forming the second row being planted between and 8 or 10 feet behind the trees in the first row. The first row may be composed of arbor-vite, which are rather slow growing, and the row behind made of Norway spruce, if desirable. White pine and European larch are rapid growing trees which may be used for this purpose. Scotch pine is inclined to be irregular in growth, and is, on this account, sometimes not satisfactory. If the trees already mentioned cannot be obtained there are other native trees which will give good satisfaction.

Kind of Trees to Plant.—Plum trees one or two years of age will give the best satisfaction. If the planter does all the cultivation himself, small trees will be more satisfactory, as they start more readily than larger ones, but if hired help is employed a good sized tree is important, as small sized trees are liable to be tramped down or otherwise injured. The paragraph on stocks should be read carefully, as the stock of the plum plays an important part in the growth of the tree.

Planting.—The spring is the best time to plant plum trees, and the earlier it is done the better, providing the soil is dry enough to work without puddling. Plum trees suffer more from late planting than apple trees. The trees may be planted with success in the autumn if the work is done early, as they will throw out roots before winter, but if planted late they are very likely to be killed by drying out. As it is of the greatest importance to get the trees planted early in the spring and as when ordered from nursermen in the spring, it is difficult to get them as early as required, a good plan is to order them to be delivered in autumn and when received heel them in well drained soil until spring. After the trees are taken out of the soil great care should be taken to prevent the roots from becoming dry before planting, as if they do the tree is almost sure to die. Dipping the roots in a thin mixture of clay loam and water will protect them somewhat, but wet burlap, old bags, or wet straw should also be used. Before exposing the roots of the trees, however, the holes should be made. Many planters seem to have the idea that if they dig a hole barely large enough for the roots to be crowded into they will have good results. Sometimes they do; much oftener they do not. If the whole field has been subsoiled and is in a thorough state of tillage it would not matter so much, as the soil all over would be in the same state of friability, but this is very rarely the case. So that, as a rule, it is necessary to make the hole somewhat larger than will accommodate the roots, spread out to their full extent. It should be made about 18 inches deep, after which the subsoil should be loosened a few inches more, but not removed. In digging the hole, the surface soil should be kept separate from the subsoil or that of poorer quality. Sufficient surface soil should now be thrown back in the hole to make the tree, when planted, about an inch deeper in the ground than it was before. If a tree is not planted deep enough, the roots may become exposed and the tree die. On the other hand, it should not be planted too deep. Before it is planted permanently in the hole, the soil which has been thrown in should be raised and rounded off in the centre. If this is done, the roots of the tree can be spread out much more readily and placed more in their natural position. Roots of plum trees have not many fibres and it is necessary to spread what are left on the tree, carefully, in order to get the best results. Broken or bruised roots should be cut off before planting the tree.

The tree being now placed upright in the hole and the roots carefully spread out, the surface soil is gently thrown in and worked in among them, by the hand, if necessary. It is very important to have the soil come in close contact with the root fibres, in order that the best conditions may be afforded the tree to begin growth promptly. When the roots are well covered, more good soil should be thrown in and when the hole is about half full it should be well tramped with the feet, after which the hole should be filled level with the surface of the soil, tramping being done while it is being filled. The surface of the soil should be left loose, as this will help to prevent evaporation of moisture from the soil which has been thrown in. It is not necessary to water any tree if planting is done at the proper season and the soil fairly moist and well compacted about the roots.

If one year old trees are used all side branches should be removed and the trunk pruned back to a height of two to three feet above the ground, leaving the tree a mere whip. The branches of two year old trees should be pruned back so as to leave only about four buds on each, but from four to six branches are all that are necessary to make a good and symmetrical top and others should be cut back to the trunk. The Stringfellow method of planting, which consists in cutting back the roots to a stub and the top to about eighteen inches from the ground and planting in a small hole, should be practised with caution in the provinces of Ontario and Quebec.

If the orchard is in an exposed position and the trees large and with high trunks, it will pay to tie stakes to them to keep them from getting loose.

In districts where drought is liable to occur, or even in places where the soil is likely to become rather dry, it will be wise to mulch the trees to a depth of from 4 to 6 inches with manure, straw, sawdust, or anything of that nature which will not become a compact mass. If this is placed about the base of the tree and left during the summer it will keep the surface soil loose and prevent evaporation of moisture and the growth of the trees will be much more rapid. A good mulch may be the means of preventing a tree from dying if the season is very unfavourable or the tree in poor condition. If the mulch is loose when winter sets in there may be danger from mice, and this should be guarded against.

Varieties.

Although a large number of varieties of plums is now offered for sale by nursery-men, the number which can be recommended and suggested as worthy of trial is comparatively limited. Few additions have been made of late years to the list of best European plums, the greatest number of new named varieties having come either from the Japanese or native plums. A considerable number of hybrid varieties have been introduced quite recently, but most of these have not yet been sufficiently tested to judge of their merits. Great improvement has been made in the Americana and other North American plums, and a great many named varieties have been introduced, no less than 137 of which are being tested at the Central Experimental Farm.

The plum season extends over a period of about three months, beginning about the 1st of August, and ending late in October or early in November, although along the River St. Lawrence below the city of Quebec some European varieties will keep until December.

By a judicious selection, varieties may be planted which will give an unbroken succession of ripe fruit during this period. The market to which the plums are to be sent should also be carefully considered as if they are to be sent long distances the firmer varieties will be the most satisfactory.

In Bulletin No. 37, on Apple Culture, published by the writer, the provinces of Ontario and Quebec were divided into thirteen different districts, representing in a greater or less degree different conditions of soil and climate. These same divisions are adopted in this Bulletin.

Although the varieties recommended are likely to be the most suitable, the intending planter should learn what varieties are proving the most profitable in his vicinity. This is important, as the districts are large and conditions will vary somewhat from one end of a district to the other. The lines dividing the districts are not arbitrary. It is not possible to make an exact dividing line on one side of which a variety will do well

and on the other side of which it will prove a failure. The boundary lines are suggestive only. It is often the case that there will be especially unfavourable locations for orchards in a milder district in which it would be safer to plant the varieties recommended for a colder one. The planter should use his judgment in the matter.

In Bulletin No. 37, on Apple Culture, a map was published showing the different districts in the provinces of Ontario and Quebec. Unfortunately this map was destroyed by fire and could not be reproduced here, but anyone having Bulletin No. 37 may refer to it as the districts are practically the same.

The varieties recommended in the following list are arranged, as far as possible, in order of ripening, beginning with the earliest. It was only through the kindness of a large number of Canadian fruit growers that it has been possible to prepare a list of the best varieties for the different districts. These men have given the results of their experiences most willingly, and I take this opportunity of again thanking them publicly for their assistance.

A DISTRICT LIST FOR THE PROVINCES OF ONTARIO AND QUEBEC.

DISTRICTS 1 AND 2.

(Counties of Essex, Kent, Bothwell, Elgin, Norfolk, Haldimand, Welland and Lincoln and the southern part of the counties of Lambton, Middlesex, Oxford and Wentworth).

VARIETIES RECOMMENDED FOR MARKET.

European.—Bradshaw, Imperial Gage, Lombard, Bavay, Italian Prune, Shropshire Damson, Grand Duke, Monarch.

Japanese.—Red June, Abundance, Burbank, Chabot.

ADDITIONAL VARIETIES SUGGESTED.

European.—Gueii, Quackenboss, Yellow Egg, Diamond, Golden Drop (Coe's).

VARIETIES RECOMMENDED FOR HOME USE.

Emerald, Washington, Fairbank, Bradshaw, Imperial Gage, Bavay, Italian Prune, Shropshire Damson.

DISTRICT 3.

(Counties of Wellington, Dufferin, Waterloo, Halton, Peel and Brant and the eastern part of Oxford, the northern part of Wentworth, the western part of York and the southern part of Simcoe).

VARIETIES RECOMMENDED FOR MARKET.

European.—Bradshaw, Gueii, Imperial Gage, Glass, Lombard, Pond, Yellow Egg, Bavay.

Japanese.—Red June, Abundance, Burbank.

ADDITIONAL VARIETIES SUGGESTED.

Mount Royal, Raynes, Starton, Chabot.

VARIETIES RECOMMENDED FOR HOME USE.

Washington, Burbank, Bradshaw, Imperial Gage, Lombard, Bavay.

DISTRICT 4.

(The northern parts of Lambton, Middlesex and Oxford; the counties of Perth, Huron, Bruce and Grey and the county of Simcoe with the exception of the extreme southern portion).

VARIETIES RECOMMENDED FOR MARKET.

European.—Bradshaw, Imperial Gage, Glass, Lombard, Yellow Egg, Bavay, Italian Prune, Monarch.

Japanese.—Red June, Abundance, Burbank, Chabot.

ADDITIONAL VARIETIES SUGGESTED.

Grand Duke, Diamond.

VARIETIES RECOMMENDED FOR HOME USE.

Washington, Burbank, Bradshaw, Imperial Gage, Bavay, Italian Prune.

DISTRICT 5.

(The county of York with the exception of the extreme western portion and the counties of Ontario, Durham, Northumberland, Prince Edward, Lennox, Frontenac to Kingston, and Hastings and Addington, within thirty miles of the St. Lawrence River; also the southern portion of Victoria and Peterborough).

VARIETIES RECOMMENDED FOR MARKET.

European.—Bradshaw, Gueii, Imperial Gage, Glass, Lombard, Yellow Egg, Bavay Italian Prune.

Japanese.—Red June, Abundance, Burbank.

ADDITIONAL VARIETIES SUGGESTED.

Grand Duke, Diamond, Monarch, Chabot.

VARIETIES RECOMMENDED FOR HOME USE.

Washington, McLaughlin, Burbank, Bradshaw, Imperial Gage, Bavay, Italian Prune, Shropshire Damson.

DISTRICT 6.

(County of Leeds, Grenville, Dundas, Stormont).

VARIETIES RECOMMENDED FOR MARKET.

Americana and Nigra.—Aitkin, Bixby, Mankato, Cheney, Wolf, U.S., Hawkeye, Stoddard.

ADDITIONAL VARIETIES SUGGESTED.

Americana.—Terry, Smith, Atkins, Bouncer.
Wildgoose.—Whitaker, Milton.

VARIETIES RECOMMENDED FOR HOME USE.

Americana, Nigra and Wildgoose.—Bixby, Mankato, Cheney, Whitaker, Cottrell, Hawkeye.

EUROPEAN SUGGESTED FOR TRIAL.

Early Red Russian, Lunn, Mount Royal, Raynes, Richland, Gueii, Glass, Arctic, Lombard, White Nicholas, Yellow Egg.

None of the European plums are very satisfactory in this district, as the fruit buds of most varieties are usually killed by winter.

JAPANESE SUGGESTED FOR TRIAL.

Red June, Burbank.

DISTRICT 7.

(The counties of Victoria, Peterborough, Hastings and Addington, except the southern portions; Manitoulin and St. Joseph Islands, and the counties of Renfrew, Lanark, Carleton, Russell, Prescott, Glengarry; the counties of Pontiac, Wright, and Ottawa south of latitude 46; also the counties of Argenteuil, Two Mountains, Terrebonne, L'Assomption; and Montcalm, Joliette, Berthier, Maskinonge, and St. Maurice, within 25 miles of the St. Lawrence River.)

VARIETIES RECOMMENDED FOR MARKET.

Americana and Nigra.—Earliest and best native seedlings, Aitkin, Bixby, Mankato, Cheney, Wolf, U.S., Hawkeye, Stoddard.

ADDITIONAL VARIETIES SUGGESTED.

Americana.—Terry, Smith, Atkins, Bender, Queen, Kieth, Etta, Bouncer. These are newer varieties, highly recommended by experimenters.

VARIETIES RECOMMENDED FOR HOME USE.

Americana and Nigra.—Best native seedlings, Bixby, Mankato, Cheney, Cottrell, Hawkeye, Bouncer.

EUROPEAN SUGGESTED FOR TRIAL.

Early Red Russian, Rowley, Lunn, Mount Royal, Raynes, Richland, Glass, Montmorency, White Nicholas, Perignon, Ungarish. None of the European plums are very satisfactory in this district, as the fruit buds of most varieties are usually killed by winter.

JAPANESE SUGGESTED FOR TRIAL.

Red June, Burbank.

DISTRICT 8.

(Counties of Huntington, Beauharnois, Chateaugay, Jacques Cartier, Laval, Hochelaga, Chambly, Laprairie, Napierville, St. Johns, and the western part of Iberville and Missisquoi).

VARIETIES RECOMMENDED FOR MARKET.

Americana and Nigra.—Earliest and best native seedlings, Aitkin, Bixby, Mankato, Cheney, Wolf, U.S., Hawkeye, Stoddard.

European.—Mount Royal, Raynes, Montmorency.

ADDITIONAL VARIETIES SUGGESTED.

Americana.—Terry, Smith, Atkins,

European.—Arctic, Glass, Yellow Egg, Richland, Early Red Russian, White Nicholas, Lombard, Damson.

VARIETIES RECOMMENDED FOR HOME USE.

Americana and Nigra.—Bixby, Mankato, Cheney, Cottrell, Hawkeye.

European.—Lunn, Raynes, Mount Royal, Brodie, McLaughlin, Montmorency, Green Gage, Queen May, Perdrigon, Ungarish.

JAPANESE SUGGESTED FOR TRIAL.

Red June, Burbank.

DISTRICT 9.

(Counties of Verchères, Richelieu, Yamaska, St. Hyacinthe, Rouville, Bagot, Drummond, Richmond, Shefford, Sherbrooke, Brome, Stanstead, and the eastern part of Iberville and Missisquoi and the western part of Compton).

VARIETIES RECOMMENDED FOR MARKET.

Americana and Nigra.—Best early native seedlings, Bixby, Mankato, Cheney, Wolf, U.S., Hawkeye, Stoddard.

ADDITIONAL VARIETIES SUGGESTED.

Americana and Nigra.—Terry, Smith, Atkins, Bender, Queen, Kieth, Etta, Bouncer.

European.—Mount Royal, Raynes, Glass, Richland, Early Red Russian, White Nicholas.

VARIETIES RECOMMENDED FOR HOME USE.

Americana and Nigra.—Bixby, Mankato, Cheney, Cottrell, Hawkeye.

EUROPEAN SUGGESTED.

Lunn, Mount Royal, Brodie, Montmorency, Glass, Richland, Early Red Russian, White Nicholas, Arctic, Damson, Ungarish.

JAPANESE SUGGESTED FOR TRIAL.

Red June, Burbank.

DISTRICT 10.

(Counties of Nicolet, Arthabaska, Wolfe, the eastern part of Compton and the counties of Beauce, Megantic, Dorchester, Lotbinière, Lévis and Bellechasse).

VARIETIES RECOMMENDED.

Americana and Nigra.—Best early native seedlings, Aitkin, Bixby, Mankato, Cheney, Wolf, U.S., Hawkeye, Stoddard.

ADDITIONAL VARIETIES SUGGESTED.

Americana.—Terry, Smith, Atkins, Bender, Queen, Kieth, Etta, Bouncer.

VARIETIES RECOMMENDED FOR HOME USE.

Americana and Nigra.—Bixby, Mankato, Cheney, Cottrell, Hawkeye.

EUROPEAN SUGGESTED FOR TRIAL.

Lunn, Mount Royal, Raynes, Brodie, Queen May, Early Red Russian, White Nicholas, Arctic, Ungarish.

JAPANESE SUGGESTED FOR TRIAL.

Red June, Burbank.

DISTRICT 11.

(Counties of Montmagny, L'Islet, Kamouraska, most of Temiscouata, Bonaventure, and Gaspé on the Bay of Chaleur side to Gaspé Basin).

VARIETIES RECOMMENDED FOR GROWING NEAR THE ST. LAWRENCE RIVER.

European.—Washington, Green Gage, Arctic, Montmorency, Lombard, Damson, Grand Duke.

ADDITIONAL VARIETIES SUGGESTED.

Mount Royal, Raynes.

VARIETIES RECOMMENDED FOR HOME USE.

Washington, Imperial Gage, Green Gage, Arctic, Lombard, Damson.

VARIETIES RECOMMENDED FOR GROWING INLAND.

Americana and Nigra.—Best early native seedlings, Aitkin, Bixby, Mankato, Cheney, Wolf, U.S.

ADDITIONAL VARIETIES SUGGESTED.

Americana.—Terry, Smith, Bender, Etta.

VARIETIES RECOMMENDED FOR HOME USE.

Bixby, Mankato, Cheney, Cottrell.

DISTRICT 12.

(Counties of Champlain, Portneuf, Quebec, Montmorency, Charlevoix, and Chieoutimi, east of the St. Maurice River and south-west of Lake St. John and to the St. Lawrence River.)

VARIETIES RECOMMENDED.

Best early native seedlings, Aitkin, Cheney, Bixby, Wolf, Cottrell, U.S.

ADDITIONAL VARIETIES SUGGESTED.

Terry, Smith, Bender, Etta.

Near the St. Lawrence River, especially in the vicinity of Quebec and below and on the Island of Orleans, the following European varieties would give more or less satisfaction:—Washington, Green Gage, Aretie, Montmorency, Mount Royal, Raynes, Lombard, Danson.

DISTRICT 13.

(North of latitude 46° in Ontario, and in Quebec north of districts 7 and 12 as far as plums will grow; also the north-eastern part of Temiseouata, Rimouski, and Matane.)

VARIETIES SUGGESTED.

Best early native seedlings, Aitkin, Odegard, Bixby, Mankato, Cheney.

DESCRIPTIONS OF VARIETIES.

The following descriptions were, most of them, made by the author from specimens either grown at the Central Experimental Farm or in other parts of the provinces of Ontario and Quebec. In some cases, however, especially among the European plums, the descriptions were obtained from other sources which are considered reliable. Where these descriptions are used the author's name is given. The varieties which are described are divided into the various groups to which they belong. They are limited to those mentioned in the district lists with the exception of a few new kinds considered promising but which have not been tested long enough to recommend, and a few of the older varieties.

AMERICANA VARIETIES.

(Where not otherwise noted.)

American Eagle.—Fruit above medium size, roundish; cavity narrow, medium depth; suture a fairly distinct line; apex rounded; colour deep purplish red; dots numerous, small, yellow; bloom moderate, pale blue; skin thick and tough; flesh deep yellow, juicy; stone medium size, oval, slightly flattened, cling; sweet, rich flavour; quality good. Season mid-September. Would be more promising if colour were brighter.

Aitkin.—Fruit large, oval; cavity narrow, medium depth; suture obscure, apex pointed; colour uniformly deep red all over; dots none; bloom none; skin thin, flesh deep yellow, juicy, moderately sweet, not rich or high flavoured; stone large, flat, oval, semi-cling, no astringency; quality medium to above. Season, last week of August. Tree only fairly productive. Nigra group. The earliness of this plum is the principal point for recommendation.

Bender.—Fruit large, oval, slightly compressed; colour dark red; dots, very many, dull yellow; bloom thick, blue; cavity shallow, suture obsolete, skin thick, tough, flesh, yellow; stone large, oval, flat, quite free; quality, good; season rather early; tree very vigorous, with very large, fine healthy foliage; very productive. (Ward.)

Planted at the Central Experimental Farm in 1899. Has not fruited yet, but is highly praised by some growers.

Birby.—Fruit above medium to large, roundish; cavity narrow, medium depth, suture rather indistinct, slightly depressed, apex rounded, colour yellow, more or less covered with bright red; dots numerous, small, yellow; bloom fairly heavy; skin moderately thick, rather tender; flesh deep yellow, juicy; stone medium size, oval in outline, considerably flattened, cling; sweet but not rich in flavour, no astringency; quality good; season late August to early September.

A very handsome, early plum. Chief fault is unevenness of ripening. Makes good preserves.

Buencer.—Fruit very large to large, roundish, somewhat heart-shaped; cavity medium width, shallow; suture a distinct line; apex pointed; colour uniformly deep purplish red all over; dots numerous, yellow, distinct; bloom moderate; skin thick, tough; flesh deep yellow, juicy; stone large, flat, oval, cling; sweet, rich, very good flavour; quality very good; season mid to late September.

A seedling of Yosemite Purple, originated at the Central Experimental Farm.

A very promising variety.

Caro.—Fruit large, roundish; cavity narrow, medium depth, suture fairly distinct; colour bright red, showing yellow in patches; dots numerous, yellow, distinct; bloom light; skin thick, moderately tender; flesh deep yellow, juicy; stone large, oval in outline, considerably flattened; sweet, rich, good flavour; quality good; season early to mid September.

A seedling of Wolf, originated at the Central Experimental Farm.

A promising seedling. More attractive than Wolf and better in quality.

Carstensen.—Fruit medium size, roundish, somewhat uneven; cavity narrow, medium depth; suture obscure; apex rounded; colour yellow, nearly covered with deep red, dots obscure; bloom none; skin thin, tender; flesh yellow, juicy, sweet, good flavour; stone flat, roundish, semi-cling almost free; quality good; season early to mid August. Tree vigorous, productive. The earliest native plum tested and valuable on the account. A Nigra seedling, originated by H. P. Carstensen, Billings Bridge, Ont., last season.

Cheney.—Fruit large, round to somewhat oval, uneven; cavity narrow, medium depth; suture merely an indistinct line; apex rounded; colour uniformly deep red all over, sometimes paler on one side; dots none; bloom none; skin moderately thick, tough, not astringent; flesh deep yellow, juicy; stone medium size, flat to a slight curve, sweet, moderately rich flavour; quality good; season late August to mid September. One of the best. It soon gets soft, however, after ripening. Tree a strong grower moderately productive. Nigra group.

Clyde.—Fruit above medium size, roundish, somewhat heart-shaped; cavity medium width, deeper than most; suture a distinct line, slightly compressed; colour yellow, almost covered with deep red; dots numerous, small, yellow, distinct; bloom moderate; skin thin, moderately tender, slightly astringent; flesh deep yellow, juicy; stone medium size, oval, considerably flattened, semi-cling; quality good; season mid September. Among the good kinds, but quite a number are better.

Confort.—Fruit medium size, roundish; cavity narrow, shallow; suture merely a distinct line; apex rounded; colour uniformly deep red all over; dots indistinct; bloom moderate; skin very thick, tough; flesh deep yellow, juicy; stone medium size, oval, considerably flattened, cling; sweet, good flavour; quality good; season, mid-September.

to October. A firm plum, but not large enough to be one of the best. Keeps better than most.

Consul.—Fruit large, roundish; cavity narrow, medium depth; suture a distinct line; apex rounded; colour deep red; dots moderately numerous, yellow, distinct; bloom light; skin rather thick, tough; flesh deep yellow, juicy, sweet; stone medium size, oval, considerably flattened, almost free; quality good. Season late September to early October. A seedling of Wolf originated at the Central Experimental Farm. Will probably prove a useful late plum.

Cottrell.—Fruit above medium to large, oblong and roundish to heart-shaped; cavity narrow, medium depth; suture a distinct line; apex rounded; colour yellow, almost covered with bright red; dots rather numerous, small, yellow; bloom medium; skin moderately thick, tender; flesh pale yellow, juicy; stone medium size, oval, much flattened, cling; sweet, good flavour; quality good. Season early to mid-September. Promising, makes a good preserving plum and is attractive-looking.

Dr Soto.—Fruit medium to above medium in size, roundish, somewhat heart-shaped, slightly flattened; cavity narrow, medium depth; suture a distinct line; colour deep yellow, well washed with deep red or dark red; dots obscure; bloom slight; skin moderately thick, fairly tender; flesh deep yellow, juicy; stone medium size, oval, considerably flattened, cling; sweet, good flavour; quality good. Season mid to late September. A good plum on account of its quality and great productiveness, but is not as large as it should be to be one of the best.

Don.—Fruit large, roundish; cavity narrow, medium depth; suture a distinct line; colour uniformly deep, lively red all over; dots numerous, small, distinct; bloom, moderate; skin thick, tough; flesh deep yellow, juicy, firm; stone medium size, oval, somewhat flattened, cling; sweet, rich, good flavour; quality very good. Season late September to October. A seedling of Wolf, originated at the Experimental Farm. A very promising plum. One of the best late plums fruited here.

Dr. Dennis.—Fruit above medium to large, somewhat heart-shaped, flattened; cavity narrow, medium depth; suture a distinct line; apex rounded; colour deep red; dots small, numerous, distinct; bloom moderate; skin thick, rather tough; flesh deep yellow, juicy; stone large, flat, broad, cling; moderately sweet, slightly astringent; quality above medium. Season mid to late September. A good variety but not as promising as some.

Etha.—Fruit large, nearly round, yellow, striped and splashed with pale red; suture distinct; skin medium, tender; flesh sweet and rich; stone oval, smooth, grooved on back; ripe August 29, 1900. Originated by H. A. Terry, Crescent, Ia; parentage unknown, fruiting first in 1895; tree described as a slow grower. A fine plum, attractive in appearance, high in quality, and promises to be very productive. Worthy of trial for market. (Goff.)

Forest Garden.—Fruit medium size, roundish; cavity narrow, medium depth; suture a distinct line; apex rounded; colour yellow, almost entirely covered with dark purplish red; dots small, numerous, yellow; bloom moderate; skin thick, tough; flesh deep yellow, juicy, sweet; stone medium size to small, oval, considerably flattened, cling; quality good. Season early to mid September. Plum not attractive enough to be promising.

Gayford.—Fruit above medium to large, roundish, somewhat heart-shaped; cavity narrow, shallow; suture a distinct line, very slightly depressed; apex pointed; colour deep, dull red on yellow ground; dots obscure; bloom moderate; skin thick, rather tough, slightly astringent; flesh deep yellow, juicy; stone medium size, oval, considerably flattened, semi-cling; sweet, good flavour; quality good. Season mid September. Would be promising if colour were more attractive.

Hammer.—Fruit large, roundish to oval; cavity narrow, medium depth; suture a line, rather indistinct; colour uniformly deep red all over; dots numerous, yellow, distinct, prominent; bloom heavy; skin thick and tough; flesh deep yellow, juicy, meaty, sweet; stone below medium size, oval, considerably flattened, cling; quality good. Season late September. A very handsome plum having more the flavour of Miner than Amerianna. This variety cracks badly on the trees, otherwise it would be one of the most promising.

Huckle.—Fruit large, roundish; cavity shallow, narrow; suture merely a distinct line; apex rounded; colour yellow, more or less covered with purplish red; dots small, indistinct; bloom medium; skin thick, moderately tough, flesh deep yellow, juicy; stone large, broad, much flattened, cling, sweet, good flavour. Quality good. Season mid to late September. One of the best.

Kieth.—Fruit large to very large, long oval; colour, orange overlaid with crimson, dots many, very minute, white; bloom blue; cavity shallow, stem short and stout, suture a faint line; skin thick; flesh yellow, firm, stone large, oval, somewhat flattened, cling; quality good to best. Season of De Soto and Wolf. One of the best Americanas (Waugh.) Planted at Central Experimental Farm in 1903.

Mankato.—Fruit above medium to large, roundish, cavity narrow, medium depth, suture a distinct line; apex rounded; colour deep, dull red with a moderately heavy bloom; dots numerous, small, yellow; bloom rather heavy, skin thick, tough, flesh deep yellow, juicy, sweet, good flavour, not astringent, stone large, flat, somewhat, quality good. Season late August to early September. Better in quality than Bixby, but not as handsome. A good early plum. Promising.

Milton.—Fruit medium to large, oval, bright to rather deep red, dots numerous, small, yellow, prominent; suture merely a distinct line, skin thin but tough, flesh yellow, juicy, sweet; stone medium size, cling, quality medium to good. Season last week of August. Tree a strong grower and very productive, when fruit buds are not injured by winter. Wildgoose group.

New Elm.—Fruit large roundish, pointed or somewhat heart-shaped, cavity narrow, shallow; suture merely a distinct line, apex rounded almost pointed, colour yellow, more or less covered with bright purplish red, dots numerous, small, yellow, bloom moderate; skin thick, tough, flesh deep yellow, juicy, sweet, stone medium size, oval, considerably flattened, cling, quality good. Season early to mid September. A firm plum and should make a good slipper. Too thick and large for the can for home use.

Ocheeda.—Fruit medium to above medium size, roundish to heart-shaped, cavity narrow, shallow; suture a distinct line; apex almost pointed, colour deep red almost black, dots numerous, small, yellow; bloom rather heavy, skin thick, moderately tough, flesh deep yellow, juicy; stone medium size, rounded, considerably flattened, cling, sweet, good flavour; quality good. Season early to mid September. A firm plum and should ship well.

Queen (Golden Queen).—Fruit very large, roundish, orange or bright golden yellow and of the most delicious flavour; quite unequalled for canning and very fit for eating out of hand or for slicing and serving with sugar and cream, as for pastries. Ripens latter part of August to September 10. Tree remarkable for its unusually fine up-right growth. (Introducer's description.) Originated with H. A. Terry, Canada. It has given good satisfaction elsewhere. Planted at Central Experimental Farm in 1901.

Silas Wilson.—Fruit large, roundish, cavity narrow, shallow, suture an indistinct line; apex rounded; colour yellow, more or less marked with purplish red, dots very small, yellow, sparse; bloom medium, skin rather thick, moderately tough, flesh deep yellow, juicy, stone medium to above medium, broad, considerably flattened, semi-cling; sweet, rich, good flavour; quality very good. Season mid to late September. One of the best Americana plums.

Smith.—Fruit very large, globose, with slight indentation to bottom, semi-cling, medium season. This variety originated here and is the given to it by many others. The largest plum I have seen of this group was of this variety, fine quality, too. Tree vigorous, healthy and productive. W. L. H. Benton, No. 1 Highway spoken of by growers. Planted at Central Experimental Farm in 1901.

Standard.—Fruit large to very large, roundish, cavity shallow, suture a distinct line, apex rounded, colour deep yellow, almost entirely covered with deep purplish red, dots fairly numerous, small, yellow, bloom light, skin thin, tough, slightly astringent, flesh deep yellow, juicy, stone medium size, large, flat, sweet, good rich flavour; quality very good. Season mid September. One of the largest and best flavoured Americana plums.

Sunrise.—Fruit large, oval; cavity narrow, shallow; suture a distinct line, not depressed; apex rounded; colour yellow, more or less covered with bright red; dots few, yellow, distinct; bloom medium, skin thick, moderately tough; flesh deep yellow, juicy, sweet; stone large, flat, oval; practically free; quality good. Season mid-September. A seedling of De Soto, originated at Central Experimental Farm. Promising owing to freeness of stone.

Terry (Free Silver).—Fruit large, regular, oval; surface smooth but not shiny; colour a clear dark red; dots small, grey; bloom thin lilac; cavity small, shallow; stem, $\frac{3}{8}$ to $\frac{1}{2}$ inch; suture indistinct; apex rounded; skin thin, tough, acid but not astringent; flesh firm but melting; stone large, ovate, pointed; flattened, acid next to stone, adherent; flavour a mingling of Americana and Augustifolia; quality good. Season end of August. One of the largest and handsomest native plums yet produced. A very promising plum. (Craig.) Planted at Central Experimental Farm in 1902.

U. S. (Brittlewood No. 2).—Quite large, spherical; dark purplish red; cling; fine quality; medium season; does not rot much here. Tree thrifty and clean, good bearer. This I regard as one of the very best of all the Americans in my collection. (J. W. Kerr, Denton, Md.)

Planted at Central Experimental Farm in 1901.

This plum is very highly recommended by several experimenters, and is said to be the largest Americana plum on the market.

Van Buren.—Fruit medium to above medium size, almost round; cavity narrow, shallow; suture only a fairly distinct line; apex rounded; colour yellow, more or less mottled and washed with bright red; dots numerous, small, yellow; bloom light; skin thick, tough; flesh deep yellow, juicy, firm; stone above medium size, roundish, broad, much flattened, cling; sweet, good flavour; quality good. Season mid-September to October.

A handsome plum. One of the latest and best keepers.

Weaver.—Fruit above medium size, roundish, somewhat heart-shaped; cavity narrow, medium depth; suture a distinct line; colour yellow, nearly entirely overspread with bright red; dots numerous, small, purple; bloom light; skin moderately thick, tough; flesh deep yellow, juicy, sweet; stone medium size, oval, considerably flattened, almost free; quality good. Season mid-September.

An attractive plum, but not large enough to be promising.

Whitaker.—Fruit large, oval, somewhat heart-shaped; colour bright red; suture merely a distinct line; dots numerous, yellow prominent; bloom thin, bluish; skin thin, tough; flesh yellow, juicy, moderately firm, sweet, good flavour; stone medium size, long oval, cling; quality good. Season first and second weeks in September, does not ripen evenly. A handsome plum. Tree vigorous, spreading and productive where fruit buds are not injured by winter. Wildgoose group.

Wolf.—Fruit large, roundish; cavity narrow, shallow; suture shallow, fairly distinct, not depressed; apex rounded; colour deep red; dots fairly numerous, small, yellow, distinct; bloom moderate; skin thick, tough; flesh deep yellow, juicy, sweet, rich, good flavour; stone above medium size, outline oval, considerably flattened, cling; quality good. Season early to mid-September. One of the best. This does not answer the description of Wolf given by some authorities. Both, however, are good plums.

Wyant.—Fruit large, oblong, flattened; cavity narrow, deep; suture a distinct line; apex almost pointed; colour deep red; dots numerous, small, purple; bloom medium; skin rather thick, somewhat tough, astringent; flesh deep yellow, moderately juicy, fairly sweet; stone large, much flattened, oval, semi-cling, almost free; quality medium. Season mid-September. Quality not good enough. This plum is highly recommended in the Western States, but has not proven as good as some others here.

Yosemite Purple.—Fruit large, roundish, somewhat flattened; cavity medium depth and width; suture an indistinct line; colour deep, dull purplish red; dots numerous, small, yellow, distinct; bloom moderate; skin thick, but tender; flesh deep yellow, juicy; stone medium size, oval, considerably flattened, semi-cling; sweet, rich flavour, but somewhat astringent; quality above medium. Season mid to late September.

EUROPEAN VARIETIES.

Amargillis (Seedling of Mirabelle).—Fruit above medium to large, roundish to heart-shaped, cavity medium depth and width, abrupt; stem medium to long, moderately stout; suture distinct, slightly depressed; apex rounded; colour greenish yellow; dots moderately numerous, indistinct; skin moderately thick, moderately tender; flesh yellow, juicy; stone medium size, oval, cling; sweet, rich flavour; quality very good. Grown from seed of Mirabelle in 1899. Began to bear in 1896. Tested September 30, 1902. Originated by Aug. Dupuis, Village des Aulnoies, P.Q.

Arctic (Moore's Arctic).—Fruit medium to below medium in size, roundish or somewhat oval; colour dark purple, almost black; bloom thin, blue; suture indistinct; flesh greenish yellow, juicy, moderately sweet; quality medium; season early September. Tree vigorous and a good cropper. Hardier than some European plums, but not desirable where the best varieties succeed, as it is too small and not good enough in quality.

Baroy (Reine Claude de Bayay).—Fruit large, roundish, slightly flattened at ends; colour greenish yellow with green splashes; bloom thin, pale; suture medium depth; stem short; flesh yellow, juicy, melting, sweet, rich, very good flavour; stone free; quality very good; season late September to early October. Tree vigorous, very productive. One of the best both for home use and for market.

Bradshaw (Niagara).—Fruit above medium to large, obovate; dark purplish red with a bluish bloom; dots few; cavity narrow, shallow; stem medium length, moderately stout; suture distinct but shallow; apex rounded; skin rather thick, tough; flesh greenish yellow, juicy, moderately firm, sweet, with a rich flavour, stone semi-cling; quality good. Season middle of August to first week of September. Tree a strong upright grower and very productive.

Brodie.—Fruit below medium size, almost round; colour dark purple with a blue bloom; dots obscure; suture merely a distinct line; skin thin, tender; flesh greenish yellow, juicy, moderately firm, sweet, rich flavour; stone small, roundish, semi-cling; quality good to very good; season second and third weeks of September. A good dessert plum, but rather small for market. Specimens received from R. Brodie, Montreal, Que. Tree has been on Mr. Brodie's place since his grandfather's time. Thought to be a seedling.

Diana.—Fruit medium to large, oval; colour dark blue, with a heavy blue bloom; dots obscure; cavity narrow, abrupt; stem short to medium, rather stout; suture merely a distinct line, not depressed; flesh yellow, moderately juicy; quality medium. Season medium. Tree a strong grower and very productive.

Early Red Russian.—Fruit medium size, oval; cavity narrow, shallow, abrupt; stem medium length, slender; suture an indistinct line, no depression; apex rounded; colour dull purplish red; dots moderately numerous, yellow, distinct; bloom thin, blue; skin fairly thick, moderately tender; flesh yellowish green, juicy; stone medium size, long, oval, cling; moderately sweet with an acid after-taste; quality medium. Season late September. Of the Lombard type. Imported from Russia by Prof. Budd from Dr. Regel, St. Petersburg, during the winter of 1881-2. Prof. Budd writing in 1890 said of this plum, 'This was sent out quite extensively eight years ago marked "Mixed Arab." The sorts mixed were Early Red Russian, White Nicholas and Black Arab.' Most of the trees proved to be Early Red Russian No. 3. There is still some doubt regarding this plum which may be the variety sent out by Prof. Budd as White Nicholas. Another variety, called Late Red, somewhat like this one, which may be the true Early Red, ripens at Ottawa during the last week of August.

Emerald.—Fruit above medium size, oval; colour yellow; suture distinct, flesh yellow, juicy, sweet, good, rich flavour; stone free; quality good to very good. Specimens received on August 4th., from E. D. Smith, Winona, Ont. Said to ripen by the end of July. Originated by the late Warren Holton, Hamilton, Ont.

Promising on account of earliness and quality.

German Prune.—'Fruit small to medium; long oval; cavity very shallow; stem rather slender, medium long; suture hardly more than a line; apex somewhat pointed; colour blue; dots few, scattered; bloom blue; flesh greenish or slightly yellow; stone small, oval, pointed, moderately flattened, very free; quality hardly more than fair; season medium; tree strong, tall grower, productive,' (Waugh). This has long been a popular plum.

Glass (Glass Seedling).—'Fruit large, roundish, deep purple with a blue bloom; suture very shallow, indistinct; stem medium length, slender; flesh yellow with a shade of green, juicy, moderately sweet; skin medium in thickness, tender; stone medium size, cling; quality medium. Season second and third weeks of September. Tree a strong grower and productive where it succeeds well. Very similar, if not identical with Quackenboss.

Golden Drop (Coe's Golden Drop).—'Fruit large to very large; oval with a short neck, the two halves unequal; cavity very shallow and abrupt; stem medium length, stout; suture deep; apex somewhat depressed; colour golden yellow; dots very many, yellow; bloom yellow; flesh firm, meaty; stone medium large, long, pointed, somewhat flattened, ribbed at the edge, half free; quality good; season medium late. Tree a good grower with large, coarse, rough foliage.' (Waugh).

Grand Duke.—'Fruit large to very large obovate; cavity narrow, shallow; stem an inch long; suture rather deep; colour very dark blue; bloom heavy, blue; flesh yellow, firm; stone oval, hardly flattened, cling; quality good; season late. Tree moderately vigorous with a spreading open head. Regarded by many as one of the very best late shipping plums' (Waugh). This is a favourite plum in some parts of the best plum districts of Ontario.

Guéni.—'Fruit medium size; oval, cordate; cavity shallow; stem an inch long, pubescent; suture shallow; apex somewhat pointed; colour blue; dots not visible; bloom blue; flesh greenish yellow; stone medium size, round oval, oblique pointed, cling; quality fair; season medium' (Waugh).

Imperial Gage.—'Fruit medium to above medium in size, roundish; colour yellowish green; dots indistinct; cavity narrow, medium depth; stem medium to long, moderately stout; suture distinct but very slightly depressed; skin fairly thick, rather tough; flesh yellowish green, firm, juicy, sweet, rich flavour; stone medium size, oval, semi-cling to almost free; quality very good. Season early September. Tree a strong grower and very productive.

Italian Prune (Fellenberg).—'Fruit medium to large, elliptical, straighter on one side and longer on the other; cavity very shallow; stem nearly as long as the fruit; suture shallow; colour dark blue; dots not many, dull yellow; bloom blue; skin thin; flesh greenish yellow; stone medium size, oval, pointed, rough, ridged at edge, quite free; quality good to extra; season late; tree rather spreading' (Waugh).

This is one of the most satisfactory European plums both for home use and for market.

Jefferson.—'Fruit medium to large, round or round oval; cavity very shallow; stem medium short; suture, hardly any; apex very slightly depressed; colour greenish yellow; dots many, greenish; bloom white; skin thin and tender; flesh yellow; stone medium size, blunt, with a short neck, slightly flattened, rough, free; flavour, rich and sugary; quality good to best; season medium late; a good tree' (Waugh). One of the finest varieties for home use.

Lombard.—'Fruit medium size, oval, slightly flattened at ends; colour purplish red with a thin blue bloom; dots fairly numerous, yellowish, distinct; stem short, slender; suture shallow, indistinct; skin rather thin, tender; flesh yellow, juicy, sweet, but not rich, firm; stone medium size, cling; quality medium; season second and third weeks of September. Tree vigorous and a very heavy bearer. One of the hardiest of the European plums.

Lunn (Montreal No. 60).—Fruit received from W. W. Dunlop, Outremont, Que.:

Fruit large, oval, broad (round oval); cavity shallow, medium width, slightly flaring; stem medium length, $\frac{1}{2}$ inch, stout; suture a distinct line, very little if any depression; apex rounded, very slightly flattened; colour dark purple; dots fairly numerous, irregular, indistinct, brownish; bloom moderate, blue; skin moderately thick, tough; flesh yellowish green, very juicy, fairly firm; stone large, oval, cling; sweet, rich; quality very good. Season early to middle of September. A fine dessert plum.

Monarch.—'Fruit large, roundish oval; cavity deep, broad, rounded; stem short and stout; suture hardly visible; colour dark purplish; bloom heavy, bluish; flesh yellowish; stone free; quality good; season late. An English variety lately introduced to this country and thought to be a valuable late shipping plum' (Waugh). This plum is well worthy of trial.

Montmorency (Reine Claude de Montmorency).—Fruit medium size, almost round; cavity narrow, abrupt, rather shallow; stem short to medium, moderately stout; suture indistinct; sometimes very slightly depressed; apex rounded or slightly flattened; colour yellow and greenish yellow before quite ripe with a light orange blush or dots of orange on sunny side; dots obscure; bloom thin, white; skin moderately thick, tough; flesh yellow, very juicy, moderately firm, sweet, rich; stone small, oval, almost free; quality very good.

Mountain.—Fruit received from W. W. Dunlop, Outremont, Que.:

Fruit medium to above medium size, roundish, flattened slightly at ends; cavity medium depth and width, slightly flaring; stem medium to long, moderately stout; suture distinct, usually slightly depressed, apex slightly flattened; colour, greenish yellow, more or less overspread with dull coppery red; dots numerous, yellow, distinct; bloom thin, bluish; skin moderately thick, tough; flesh yellowish green; stone above medium, broad, roundish, cling; sweet, rich; quality very good. Season early to middle September. An excellent dessert plum. Well worth propagating.

Mount Royal (Dunlop 54).—Fruit received from W. W. Dunlop, Outremont, Que.

Fruit medium size, roundish, flattened at stem end; cavity medium to open, medium depth, somewhat flaring; stem short to medium, moderately stout; suture distinct, very slightly depressed; apex rounded, slightly flattened; colour dark purple; dots numerous, irregular, distinct; bloom blue, moderate; skin moderately thick, fairly tender; flesh greenish yellow, juicy, firm, sweet, moderately rich flavour; stone below medium, roundish, cling; quality good. Season early to mid-September. Should be a good shipping plum.

McLaughlin.—'Fruit medium size, round or even oblate; cavity shallow, with a ridge around the stem; stem strong, rather long; suture very shallow; apex very slightly depressed; colour greenish yellow with a pink blush; dots many, greenish; bloom white; skin thin; flesh yellow; stone medium size, oblique oval, slightly flattened, rough, cling; flavour rich, sugary; quality extra. Season medium. Tree hardy and a fairly good grower.' (Waugh).

One of the best varieties for home use, the quality being exceptionally good. It is of the Green Gage type.

Peters (Peters' Yellow Gage).—'Fruit medium to large; round oval; cavity medium, shallow, abrupt; stem long, pubescent; suture shallow; apex slightly depressed; colour greenish yellow, sometimes with a slight blush; dots many, yellow; bloom white; skin thin; flesh greenish yellow; stone medium large, oval, pointed, hardly flattened, cling; quality good to best. Season early. Tree moderately vigorous and upright. A good amateur variety of the Green Gage type.' (Waugh).

This has succeeded well in District No. 3.

Pond (Pond's Seedling).—Fruit very large; nearly oval, but tapers slightly towards cavity; colour purplish red; bloom purplish; dots numerous, dull yellow, distinct but not prominent; cavity narrow, shallow; stem medium length, fairly stout; suture distinct and but slightly depressed; skin thick, rather tough; flesh yellow, juicy, sweet,

good flavour; stone large, rough, cling; quality good. Season early September. Tree a strong grower and quite productive.

Quackenboss.—Fruit medium size or larger; round oval; cavity shallow, flaring; stem rather long; suture a line; colour blue; dots blue; bloom blue; skin thin; flesh greenish; stone oval, pointed, flattened, cling; quality fair to good. Season medium. A good rapid growing tree and fairly productive. (Waugh).

Ripens in Ontario in second and third weeks of September.

Queen May.—Fruit large, roundish, almost perfectly round; colour greenish yellow splashed with pale green when not ripe; bloom thin, pale bluish; dots small, pale, indistinct; cavity narrow, medium depth; stem medium length to rather long, fairly stout; suture indistinct; apex rounded; skin moderately thick, tough; flesh greenish yellow, very juicy, moderately firm, sweet, rich flavour; stone medium size, almost oval, cling; quality very good; season September. Tree a strong, moderately spreading grower, fruiting heavily when young, but does not live very long. A very promising dessert plum. Specimens received from N. E. Jack, Chateaugay Basin, Que., who is growing trees received from Thos. Clark, Chateaugay. Thought to be a seedling.

Raynes (Dunlop 53).—Fruit received from W. W. Dunlop, Outremont, Que.:—

Fruit above medium to large; oval, long, flattened on side of suture; cavity medium depth and width, abrupt; stem medium length, moderately stout; suture distinct, slightly depressed; apex rounded; colour dark reddish purple; dots small, numerous, indistinct; bloom moderate, blue; skin thin, tender; flesh yellowish green, firm, fairly juicy; stone above medium to large, long, oval, free; moderately sweet; quality above medium. Season early to middle of September. A prolific bearer and should be a good shipper. A prune plum.

Rickland.—Fruit medium to above medium size, oval; cavity narrow, medium depth, abrupt; stem medium length, $\frac{3}{4}$ -inch, slender; suture a distinct line, no depression; apex rounded; colour deep purplish red; dots fairly numerous, yellow, indistinct; bloom moderate, blue; skin thick, fairly tender; flesh greenish yellow, juicy, moderately firm; stone medium size, oval, flat, cling; sweet but not rich; quality above medium. Season middle of September. Hardier than most European sorts. Originated on the farm of Randall Elden, Rickland, Pennsylvania.

Rowley.—Fruit above medium size, round, dark purplish red with a bluish bloom; dots obscure; suture an indistinct line; apex rounded; skin thin, moderately tough; flesh yellow, moderately juicy, firm, sweet, rich flavour; stone medium size, oval, slightly flattened, cling; quality good to very good. Season end of August and first week of September. A promising seedling originated by Jos. Rowley, Cummings Bridge, Ont. (near Ottawa). Said to fruit well nearly every year.

Shropshire (Damson).—Fruit small, oval; cavity, hardly any; stem about one-half inch long; suture none; colour dark blue; dots none visible; bloom blue; skin firm; flesh greenish, sour; stone small, oval, turgid, cling; quality fair. Tree a good grower and enormously productive. (Waugh).

This is a popular damson in Canada.

Ungarish.—Fruit above medium to large; long oval; cavity narrow, shallow, abrupt; suture distinct, very slightly if at all depressed; apex round; colour dark purple; dots moderately numerous, indistinct, brown; bloom moderate, blue; skin fairly thick, tender; flesh greenish yellow, firm, fairly juicy; stone large, long, oval, free; moderately sweet; quality above medium. Season middle of September. Introduced by Prof. Budd from C. H. Wagner, Riga, Russia.

This plum is somewhat like the Raynes (Dunlop 53) A prune plum. Promising on account of hardiness.

Washington.—Fruit large, roundish, slightly flattened at ends; colour greenish yellow with a pink blush on sunny side; dots obscure; cavity narrow, shallow; stem short, stout; suture distinct and slightly depressed; skin tough; flesh greenish yellow, firm, juicy, sweet, rich, stone medium size, roundish, almost or quite free; quality very good. Season early to mid-September. Tree a strong grower with a roundish top.

Yellow Egg.—Fruit large to very large, oval; colour deep yellow; bloom white; dots small, numerous, indistinct; cavity shallow, ridged; stem long, moderately stout; suture distinct, slightly depressed; skin thick, rather tough; flesh yellow, juicy, sweet; stone large, oval, cling; quality good. Season end of August to early September. Tree vigorous and productive.

JAPANESE VARIETIES.

Abundance.—Fruit large, roundish; bright to deep red with a yellow ground; dots numerous, yellow, prominent; cavity narrow, abrupt; stem medium length, rather stout; suture distinct; apex pointed; skin thin, moderately tender; flesh yellow, juicy, firm, sweet, rich; stone oval, cling. Quality good to very good. Season August 10 to 25. Tree a strong upright grower, an early bearer and productive.

Burbank.—Fruit large to very large, roundish; colour deep red with dark red on sunny side and about cavity, on a yellow ground; dots numerous, small, distinct, yellow; stem medium length; suture merely a distinct line; apex sometimes pointed; skin thin, moderately tender; flesh yellow, firm, juicy, sweet, good flavour; stone roundish, cling; quality good. Season latter part of August, a few days after *Abundance*. Tree an exceptionally vigorous grower, very branching and bears early and heavily. One of the most, if not the most, satisfactory of the Japanese plums. The flower buds appear hardier than most European varieties.

Chobot.—Fruit medium to large, roundish, heart-shaped; colour deep, rather dull red; dots numerous, small, yellow; cavity narrow; stem short, stout; suture fairly distinct; apex usually rounded; skin moderately thick, tough; flesh yellow, juicy, firm, sweet; stone below medium size, oval, cling; quality good. Season late. Tree a strong upright grower and bears well. This variety is one of the later introductions and has proven one of the best of the Japanese plums.

Red June.—Fruit medium to below in size, roundish, somewhat flattened; colour deep to dark red; dots small, yellow, numerous; cavity deep; stem short, moderately stout; suture distinct but shallow; skin thin, tender; flesh pale yellow, firm, juicy, briskly sub-acid with little richness; stone small, roundish, cling; quality medium. Season last week of July to first week of August. Tree moderately spreading. A medium bearer. This plum is valuable on account of its extreme earliness. This variety was received at the Central Experimental Farm under the name of *Botan* and *Shiro momo*. The flower buds are hardier than most European varieties but although there is usually much bloom comparatively little fruit sets, probably because blossoms are self-sterile.

Varieties of Plums which are being tested at the Central Experimental Farm,
Ottawa.

List of Varieties.

Names of Varieties.	Date of Planting Oldest Trees.	Names of Varieties—Continued.	Date of Planting Oldest Trees.
EUROPEAN PLUMS.		EUROPEAN PLUMS—Continued.	
Amaryllis	1902	Rowley	1903
Arctic (<i>Moore's Arctic</i>)	1899	Saratoga	1901
Baker Prune	1899	Sharp (<i>Victoria</i>)	1895
<i>Beauty of Naples</i> = <i>Naples</i>		Shropshire (<i>Shropshire Dunsdon</i>)	1898
Bonnie	1901	Smith's Early	1897
Bohemian	1895	Tatze	1903
Pomme Ste. Anne	1895	Smith's October	1897
Bradshaw	1899	Ungarish	1888
Brodie	1903	<i>Victoria</i> = Sharp	
Chataouqua	1899	Voronesh Blue	1903
Columbia	1898	Voronesh Yellow	1893
Cochet père	1901	Waggenheim	1901
Czar	1900	White Nicholas	1895
Denniston Superb	1902	<i>Yellow Moldavka</i> = <i>Moldavka</i>	
Diamond	90		
Diane (<i>Duane's Purple</i>)	95	JAPANESE PLUMS.	
Early Red Russian	1895	Abundance	1900
Emerald	1903	Bereckman's	1903
Englebert (<i>Prince Englebert</i>)	1898	<i>Botan</i> = Red June	1900
<i>Fellenberg</i> = Italian Prune		Burbank	1903
Field	1900	Chabot	1903
<i>General Hand</i> = Hand		Engre	1903
German Prune	1895	Hale	1903
Glass (Glass Seedling)	1893	Kerr	1903
Grand Duke	1898	October (<i>October Purple</i>)	1902
Gueli	1898	Red June	1894
Hand (General Hand)	1898	Satsuma	1903
Hanszweitsche	1900	<i>Shiro-Sinomo</i> = Red June	
Horrigan	1903	Willard	1900
Ickworth (<i>Ickworth Imperatrice</i>)	1901		
Imperial Gage	1900	AMERICANA PLUMS.	
Italian Prune (<i>Fellenberg</i>)	1898	Admiral Dewey	1901
John A.	1895	Admiral Schley	1901
Kingston	1900	Advance	1903
Lachino	1903	American Eagle	1895
Lapsic	1893	Bailley	1903
Lingoh	1900	Bender	1900
Lombard	1900	Bixby	1893
Lunn	1903	Blackhawk	1893
McLaughlin	1900	Bomberger	1901
Mallard	1903	Bouquet	1895
Moldavka (<i>Yellow Moldavka</i>)	1897	Brackett	1903
Monarch	1900	Brittlewood No. 1	1901
Monroe	1901	<i>Brittlewood No. 2</i> = U. S.	
Montmorency (<i>Reine Claude de Montmorency</i>)	1899	Brittlewood No. 3	1903
<i>Moore's Arctic</i> = Arctic		Brooklyn	1903
Mount Royal	1903	Bryan (<i>W. J.</i>)	1903
Mountam	1903	Budd (<i>Prof.</i>)	1903
<i>Naples</i> (<i>Beauty of Naples</i>)	1890	Cero	1895
Niagara (<i>Bradshaw I.</i>)	1898	Champion	1895
Outremont	1903	City	1895
Perdigan	1903	Coinage	1903
Pond (<i>Pond's Seedling</i>)	1000	Colorado (<i>Colorado Queen</i>)	1895
<i>Prince Englebert</i> = Englebert		Collman	1901
Quackenboss	1898	Comfort	1893
Queen May	1903	Consul	1895
Quebec	1902	Cottrell	1895
Raynes	1903	Craig (<i>Prof.</i>)	1903
Red Egg	1898	Cyclone	1901
<i>Reine Claude de Montmorency</i> Montmorency		Deepereck	1895
Richard Trotter	1895	Dennis (<i>Dr.</i>)	1895
Richland	1888	De Soto	1888

List of Varieties of Plums.

Names of Varieties—Continued.	Date of Planting Oldest Trees.	Names of Varieties—Continued.	Date of Planting Oldest Trees.
AMERICANA PLUMS—Continued.		AMERICANA PLUMS—Concluded.	
Diana.....	1901	Value.....	1903
Don.....	1895	Van Buren.....	1890
Dunlap (<i>Dunlap's No. 1</i>).....	1895	Van Deman.....	1895
Eldorado.....	1901	Warren.....	1901
Emma.....	1901	Weaver.....	1895
Etta.....	1901	Wolf.....	1888
Forest Garden.....	1890	Wyant.....	1890
Freestone.....	1901	Yellow Sweet.....	1895
<i>Free Silver</i> = Terry.....		<i>Yosemite Purple</i> = Purple Yosemite.....	
Galena.....	1895		
Gaylord.....	1895	NIGRA PLUMS.	
Golden.....	1903	Aitkin.....	1897
<i>Golden Queen</i> = Queen.....		August.....	1899
Hammer.....	1895	Brandon Ruby.....	1903
Hanson.....	1901	Cheney.....	1890
Hawkeye.....	1898	Manitoba No. 4.....	1895
Holt.....	1901	Manitoba No. 5.....	1895
Hunt.....	1893	Mills Seedling.....	1890
Irene.....	1895	Odegard.....	1901
Ironclad.....	1895	Smith Red.....	1900
Jessie.....	1894	Suelling.....	1894
Julia.....	1901	Whyte.....	1893
Kennedy Red.....	1897		
Kieth.....	1903	MINER-LIKE PLUMS.	
Labert (<i>Labert's Red</i>).....	1897	Carver.....	1895
<i>Large Red Sweet</i> = Plunk.....		Clinton.....	1895
Legal Tender.....	1901	<i>Col. Wilder</i> = Wilder.....	
Leonard.....	1897	Esther.....	1895
Lillie.....	1901	Forest Rose.....	1898
Lottie.....	1901	Idall (<i>Idol</i>).....	1895
Louisa.....	1901	Iroquois.....	1895
Mankato.....	1895	Miner.....	1895
Marellus.....	1901	Nebraska.....	1895
Marjorie.....	1903	Oren.....	1900
Mary.....	1901	Prairie Flower.....	1895
Maude Lacey.....	1903	Rachel.....	1897
Mollie.....	1891	Surprise.....	1900
Moon.....	1899	Weir (<i>Weir's Large Red</i>).....	1895
Nellie.....	1895		
Nellie Blauche.....	1901	WAYLAND-LIKE PLUMS.	
Newton Egg.....	1897	Golden Beauty.....	1895
New Uln.....	1895	Moreman.....	1888
Ocheeda.....	1895	Red.....	1895
Old Gold.....	1897		
Omega.....	1901	WILDGOOSE PLUMS.	
Pearl.....	1901	Downing, (<i>Charles</i>).....	1893
Peffer Premium.....	1895	Dunlop, (<i>No. 2</i>).....	1895
Plunk (<i>Large Red Sweet</i>).....	1897	James Vick.....	1897
Purple Yosemite (<i>Yosemite Purple</i>).....		Milton.....	1893
Quaker.....	1895	Roulette.....	1899
Queen (<i>Golden Queen</i>).....	1901	Sophie.....	1895
Reel.....	1901	Van Houten.....	1901
Rockford.....	1892	Whitaker.....	1895
Rollingstone.....	1888	Wilder (<i>Col.</i>).....	1901
Ruby.....	1903	Wildgoose.....	1898
Sada.....	1901		
Silas Wilson.....	1895	HYBRID PLUMS.	
Smith.....	1901	America (<i>Botan X Robinson</i>).....	1901
Spier.....	1888	Ames (<i>P. Americana X P. triflora</i>).....	1901
Stella.....	1903	Apple (<i>Parentage unknown</i>).....	1901
Stoddard.....	1891		
Sunrise.....	1895		
Terry's De Soto.....	1903		
Terry (<i>Free Silver</i>).....	1902		
U. S. (<i>Brittlewood No. 2</i>).....	1901		

List of Varieties—Concluded.

Names of Varieties—Continued.	Date of Planting Oldest Trees.	Names of Varieties—Concluded.	Date of Planting Oldest Trees.
HYBRID PLUMS—Continued.		HYBRID PLUMS—Concluded.	
Bartlett (<i>Delaware X P. Simoni</i>).....	1901	Occident (<i>Parentage unknown</i>).....	1903
Chalco, (<i>P. Simoni X Burbank</i>).....	1903	Pendent (<i>Pottawattamic X Forest Garden</i>)	1901
Climax, (<i>Botan X P. Simoni</i>).....	1901	Preserver (<i>Kelsey X Early Red</i>).....	1903
Combination.....	1903	Ragland (<i>Parentage unknown</i>).....	1903
Compass Cherry (<i>Prunus pumila Besseyi X Minor</i>).....	1901	Red May (<i>Abundance X Wildgoose</i>)....	1903
Doris.....	1903	Rupert (<i>Prunus pumila X P. americana</i>)	1901
Duke.....	1903	Shiro (<i>Robinson X Myrobalan X Wickson</i>).....	1901
Excelsior (<i>Kelsey X Wildgoose</i>).....	1903	Six Weeks.....	1903
First.....	1903	Sultan = Occident.....	1901
Golden (Gold) (<i>Robinson X Botan</i>).....	1903	Watson.....	1903
Goosedyc.....	1903	Waugh.....	1903
Gonzales (<i>Parentage unknown</i>).....	1903	Wickson (<i>P. triflora X P. Simoni</i>).....	1903
Holland (<i>Kelsey X Lone Star</i>).....	1903	Yates.....	1903
Kelbalan.....	1903		
Kelmyro.....	1903		
Nona (<i>P. triflora X P. angustifolia?</i>) ..	1903		

POLLINATION OF PLUMS.

Until quite recent years the pollination of fruits was given comparatively little study, but during the past ten years a number of experimenters have been at work investigating the causes of failure in the setting of fruit and have shown the importance of having the blossoms fertilized by pollen which will ensure the setting of the fruit. The plum has received more careful study in regard to pollination than any other fruit. For five years or more Prof. F. A. Waugh gave this question his special attention, and the results which he obtained and the facts which he published have been of great assistance to fruit growers.

As a result of his experiments he found that of all the varieties of plums of American origin which he studied (and he studied most of those on the market), only one variety, the Robinson was self-fertile. In other words, if a tree of any other variety of American origin than the Robinson were planted where its flowers could not be fertilized by the pollen from a tree of another variety no fruit or practically no fruit would set. The Japanese plums were almost as self-sterile as the American, but the European plums were more or less self-fertile. It will be seen from the above that many varieties of plums are not fertilized at all or only partly fertilized by their own pollen. It is necessary, therefore, that varieties be so mixed in the orchard that proper pollination and a good setting of the fruit will be ensured. Americana varieties of plums should be planted to pollinize Americana varieties, Nigra to pollinize Nigra, although Americana will do, Japanese to pollinize Japanese and European to pollinize European. Varieties should be planted near each other which bloom at the same time. This is very important, as if the varieties do not bloom at the same time pollination cannot take place. The nurserymen do not as a rule give in their catalogues the relative time of blooming of the different varieties bought from them, but it is necessary for the fruit grower to know this before planting if he is to obtain the best results. The value of bees and other insects in the orchard for assisting in pollinizing the flowers cannot be over-estimated, and where possible, colonies should be kept for this purpose.

The dates of blooming of plums in different parts of Canada were recorded by fruit growers for the Horticultural division of the Central Experimental Farm for five years. These dates have been compiled and the average of each variety thus obtained.



FRUITING BRANCH OF DON PLUMS.



CHENEY PLUM IN BLOOM—Typical Tree of *Prunus Nigra*.

The following is a table of varieties of Americana and Nigra plums mentioned in this bulletin, giving their season of blooming. This will be of assistance to fruit growers when planting. A table of the European varieties is not given, as the information obtained was not full enough to be perfectly reliable, and furthermore the intermixing of European kinds is not so important as with American, as most of the varieties appear to be self fertile. The Japanese varieties described in this Bulletin are all extra early and early bloomers, and will thus pollinize one another.

There is ten days' difference between the time of blooming of the earliest and latest varieties in the following table, so that it would be impossible for the early blooming varieties to pollinize the late ones, but the early might pollinize the medium, the medium the medium late, and the latter the late.

AMERICANA AND NIGRA PLUMS—SEASON OF BLOOMING.

Extra Early.—Aitkin.

Early.—Cheney, Mankato, Odegard.

Medium.—Bixby, Bouncer, City, Comfort, Gaylord, New Ulm, Ocheeda, Queen, U.S., Van Buren, Weaver.

Medium Late.—Bender, Cottrell, Smith, Wolf, Wyant.

Late.—American Eagle, De Soto, Forest Garden, Hammer, Hawkeye, Quaker, Silas Wilson, Stoddard.

PRUNING.

The pruning of the trees is not nearly as important in the culture of the plum as of the apple, and there is more danger of injury by over-pruning. While an apple tree will, as a rule, recover from severe pruning, plum trees often never recover from it. In plum culture it is much better to err on the side of under-pruning than over-pruning. The tree should be pruned when young to a symmetrical top with the main branches so disposed that there will be no bad croches, after which very little pruning is necessary beyond cutting out dead and broken branches and thinning out where the top is very thick. Some varieties will require more pruning than others, some of the Japanese being especially rampant growers. There is a difference of opinion and a difference of results in regard to the heading back of plum trees. As a rule, very good results will be obtained without heading back. A few strong growing varieties, such as Burbank, however, have to be kept cut back to keep them within reasonable limits. When trees are headed in, the work should be done early in the spring, at which time the ordinary pruning can be done to best advantage. Wounds should be covered with lead paint or grafting wax.

The following note on useful fertilizers for the plum orchard has been prepared for this Bulletin by the Chemist of the Experimental Farms.

FERTILIZERS FOR THE PLUM ORCHARD.

By FRANK T. SHULTZ, M.A., *Chemist, Dom. Exptl. Farms.*

In common with other orchard crops, the requirements of the plum tree, as regards plant food, are chiefly nitrogen, phosphoric acid, and potash, and these essential elements must be present not only in fair quantities but in more or less readily available forms if vigorous growth and an abundance of fruit are to be expected. In addition, lime is frequently of value, since all stone fruits make a considerable demand upon the available lime of the soil, and this, owing to natural causes or exhaustive cropping, may be reduced to mere traces.

Further, all rational systems of fertilization must include the renewal, from time to time, of the vegetable organic matter of the soil, and this may best be attained when the supply of barnyard manure is insufficient, by the growth and turning under of green crops.

Nitrogen and Organic Matter.—Since these, from an agricultural point of view, are intimately associated the one with the other, the latter being the natural conservator of the former, and economy, as a rule, dictating their use in orchards in such an associated form, we may conveniently consider their application under the same heading.

Barnyard manure undoubtedly stands first on the majority of farms for enriching the soil in these constituents and improving its physical condition. In composition—and hence in value—it is extremely variable, but good average samples of fresh manure may be considered to contain: nitrogen 0.5%, phosphoric acid 0.25%, potash 0.45%. The organic matter in fresh manure is usually in the neighbourhood of 25%.

With many orchardists, however, the available supply of manure is insufficient and inadequate for the area to be fertilized, and it is for such that the system of green manuring—as by the growth and turning under of a cover crop—is especially valuable. The important role of the cover crop in the modern systems of orchard soil management is set forth in another place in this bulletin and the whole subject of increasing fertility by means of clover has been very fully explained in Bulletin No. 40 of the Experimental Farms series. It will, therefore, only be necessary here to state one or two of the principal reasons why clover or some other of the legumes should be employed for this purpose of enriching and improving the soil.

A vigorous crop of clover will contain at a moderate estimate, in its foliage and roots:—

Nitrogen.....	from 100 to 150 lbs	per acre.
Phosphoric acid....	" 30 to 45	"
Potash.....	" 85 to 115	"

It is evident, therefore, that by this use of clover we can with a single crop furnish the soil with as much nitrogen as would be supplied by a dressing of 10 tons of manure per acre. The greater part of this nitrogen is taken by the clover from the atmosphere, and hence is a distinct addition to the soil. The phosphoric acid, potash and lime, it is true, have been obtained from the soil, but have largely been drawn from a considerable depth and hence increase the stores of these elements in the upper layers of the soil. Moreover, the decay of the clover sets free all these important elements of plant food in forms readily utilizable by trees.

One or two words should also be added regarding the value of the organic matter so supplied. This eventually is converted into humus, the importance of which as a soil constituent it is difficult to over-estimate. It not only liberates slowly and continuously its plant food, but vastly improves the soil texture, whether it be a clay or a sandy loam. It increases the soil's power for absorbing and retaining moisture and it furnishes the best material for the development of microbial life, which, as now known, plays so important a part in increasing a soil's fertility.

There is, of course, a possibility of too great an enrichment with nitrogen by this means; this would be indicated by luxurious and excessive growth, dark-green foliage and poor fruiting. Under such circumstances, the use of organic and nitrogenous fertilizers should be discontinued.

To furnish immediately available nitrogen to young trees showing a deficiency of this element (indicated by a sparse development of foliage of a yellowish-green colour), nitrate of soda may be used as a top dressing, at the rate of 100 to 200 lbs. per acre.

Phosphoric Acid, Potash and Lime.—There is probably no better medium for supplying these elements to orchard soils than unleached hardwood ashes. These should contain from 5 to 6 per cent of potash, in the neighbourhood of 2 to 3 per cent of phosphoric acid, and from 30 to 35 per cent of lime. Not only are these constituents in wood ashes in a condition that renders them readily assimilable, but in many parts of Canada they are cheaper than in any other form of commercial fertilizer. Fifty to eighty bushels per acre is the usual application.

Bone meal should contain from 3 to 4 per cent of nitrogen, and 20 to 24 per cent of phosphoric acid. Since its plant food is only slowly liberated in the soil, it is considered a 'lasting' fertilizer and one well adapted to orchard use. The application is usually about 300 lbs. per acre. Superphosphate will contain from 15 to 20 per cent of phos-

phoric acid, the greater portion of which should be in a soluble (available) condition. The application is from 200 to 400 lbs. per acre.

Potash can be obtained in the form of muriate (50 per cent actual potash) or as kainit (12 per cent actual potash); 100 to 150 lbs. of the former and 200 to 500 lbs. of the latter are the limits for ordinary application.

Voorhees, in his work on fertilizers, suggests the following mixtures for orchards: (a) one part, or 100 lbs. each, of ground bone, superphosphate, and muriate of potash; and, (b) a mixture of one and a half parts, or 150 lbs., of ground bone and one part, or 100 lbs., of muriate of potash.

Soils differ so greatly as to their fertility that it is impossible to state definitely the amounts of these fertilizers that could in all cases be used with profit. From 300 to 500 lbs. of such mixtures per acre on fairly good soils would, no doubt, be ample, but on very poor soils these amounts, according to the experience of many orchardists, might be considerably increased.

Lime.—Where lime only is required, by reason of natural deficiency in this element, of excess of humus, sourness, or the refractory character of the soil, it may be applied at the rate of 1 to 2 tons per acre."

CULTIVATION.

Plum orchards should be kept under a high state of cultivation or otherwise the fruit is likely to be small. The surface soil should be stirred about once a week or after every heavy rain until July in order to conserve moisture and to open the soil so that the air can get in freely and assist in nitrification and help promote a thrifty growth of the trees. Trees sometimes produce good crops of plums when they are growing in sod, but this is not the rule. It has been found that there is more than twice as much moisture when the ground is cultivated than where there is sod. Furthermore, the curculio becomes very troublesome in sod orchards, and for this reason, if for no other, the orchard should be kept cultivated. Cultivation should be discontinued towards the middle of July, in order that the wood may ripen in good time and that a good cover crop may be established. Where trees are planted close and cultivation is difficult it will be preferable to mulch the trees with grass or straw than to leave them in sod as the fruit will be larger and the trees more thrifty.

COVER CROPS.

A cover crop is a crop of some kind the seed for which is planted for the purpose of having a growing crop in the orchard after cultivation has ceased, in order to help use up plant food which has been made available during the summer and which might leach away if the soil were bare; and a more important purpose still of this crop, especially in some localities, is to have a covering which will protect the roots of the trees and help to hold the snow in winter. A third important use of the cover crop is to have some vegetable matter to plough under in the spring to improve the soil both by adding humus and by adding nitrogen where leguminous plants are used.

Experiments with cover crops have been carried on rather extensively at the Central Experimental Farm for the past eight years, and it has been found that the most satisfactory plant for this purpose is the common red clover, sown not later than the middle of July at the rate of 10 to 12 lbs. per acre, no nurse crop being used as a rule. The hairy vetch (*Vicia villosa*) has given good satisfaction in some places, especially in the Niagara peninsula, and has the advantage of growing very late in the autumn. This may be sown at the rate of 40 or 50 lbs. per acre with good results. It is not necessary that the plant used as a cover crop should live over winter.

Sometimes it is not possible, owing to dry weather, to get a good cover crop by sowing about the middle or latter half of July. Experiments are now in progress at the Central Experimental Farm to learn how satisfactory plants in drills two feet or more apart will prove as cover crops. The object of planting this way is that the seed may be sown early enough to ensure a good crop and yet the soil may be cultivated

until the usual time. Soja beans, horse beans, and hairy vetch are being used for this purpose. This method is said to have given good satisfaction where it has been tried. If the soil suffers from lack of moisture in a dry time, the cover crop should be ploughed under as early in the spring as possible and cultivation begun. By doing this, much of the moisture which would otherwise be transpired through the leaves of the growing plants would be conserved. If, however, there is always sufficient moisture in the soil it is a good practice to let the crop grow until about the third week of May, as there would be more humus and nitrogen obtained by this method.

PICKING, PACKING, STORING AND MARKETING THE FRUIT.

Experience, only, will teach the fruit grower the proper time to pick each variety of plum for storing or shipping, as some may be picked greener and some riper than others. As a rule, however, the European plums should be picked when they are well coloured but still quite firm. The Americana varieties are so juicy that they also have to be picked before fully ripe if they are to be shipped far. The Japanese varieties may be picked earlier than either of the preceding as the colour and flavour will develop well even though harvested when rather on the green side. Plums should be picked with the stems on when it is possible to do so and on no account should they be shaken from the tree when intended for shipment. A step-ladder to stand on and a strong basket to put the fruit in are necessary. Plums are usually sold in medium sized fruit baskets, the larger sized grape baskets being preferred. It is sometimes advisable when the fruit is especially fine to face the basket, thus giving the fruit a still more attractive appearance, but if this is done the face should represent the kind of fruit to be found lower down. Some varieties of plums will keep for several weeks in cool or cold storage. A temperature of from 35 to 42 degrees Fahrenheit has been found the best for storing this fruit.

The marketing of the plums will depend very much on the location of the orchard. The greatest care should be observed to have the fruit reach the market in prime condition, and if it is consigned to a commission merchant he should be one of recognized integrity.

Thinning the Fruit.—Until quite recent years the thinning of plums on the trees had not been practiced very extensively. Plum orchards have increased so much, however, in size and number that the competition has been keener and the prices lower, and in consequence the most advanced growers are now thinning their fruit, and find it profitable to do so, as the prices obtained for the larger fruit more than compensate for the labour required in thinning. Furthermore, if part of the fruit is picked when green it does not have to be picked when ripe, so there is little extra handling. Some varieties of plum trees bear very heavily, and this is particularly true of the Americana varieties. In consequence, the fruit when unthinned is much smaller than it would be if there were less of it, and the drain upon the vitality of the tree from the production of so much seed shows itself before long, and it frequently happens, especially in poorly tilled and poorly fertilized orchards, that trees literally bear themselves to death.

The time to thin is after the fruit is well set and when it is fairly certain what the crop is going to be. There is always a dropping of plums during the month of June, caused principally by improper pollination, natural thinning and injury from curculio, and as soon as possible after these have thinned the crop, hand thinning should be done. The Americana plums fruit so heavily that, in an experiment conducted at the Wisconsin Experiment Station, it was found that nearly four-fifths of the crop should be removed in order to get really satisfactory results. When Americana plums were thinned as heavily as this the fruit was left about two inches apart, which was found a good distance in the experiment, but a greater distance was suggested. In 1897 an experiment in thinning plums was conducted at the Central Experimental Farm with native plums, when the thinned fruit when ripe was considerably larger than that unthinned. An experiment was also conducted for the horticultural division at St. Catharines the same year with a European variety, when a distinct advantage was obtained by thinning.

Some varieties of European and Japanese plums are left as much as six inches apart by fruit growers, and at this distance profitable crops are said to be obtained of fruit of the best quality. The most profitable distance apart to leave the plums will be largely governed by the variety. Some varieties will not need thinning at all, and even where trees are bearing heavily the scarcity and cost of labour may prevent the profitable thinning of the fruit.

CANNING AND PRESERVING AMERICAN PLUMS.

All good housewives are familiar with the way to can and preserve European plums, but the American varieties require different treatment in order to get the most satisfactory results. Some varieties are thick in the skin, while others are more or less astringent, and for these reasons special methods are employed to make the fruit more palatable. The skin of many varieties may be readily removed by pouring boiling water on the fruit and then peeling it. Preserves made from fruit thus treated is very good. The skin of some varieties will not break down in cooking, and peeling is especially desirable in such cases.

American plums are not as good for canning as the European, and, if they are canned, are best for making pies.

In the autumn of 1902 the following eight varieties of American plums were preserved, in order to learn what differences there were in these kinds for this purpose:—Bixby, Cheney, New Ulm, Mankato, Cottrell, Bouncer, American Eagle, Silas Wilson. These were preserved with and without the skin. In nearly every case the peeled fruit made the best preserves. The Bixby, however, cooked with the skin on was the best of all those tested, having a better flavour than any of the others both peeled and unpeeled. With some varieties 1 lb. of sugar to 1 lb. of fruit was found to make the preserves too thick; on the other hand, $\frac{3}{4}$ lb. sugar to 1 lb. of fruit, in some cases did not make them quite sweet enough. None of the varieties tested were found markedly astringent, though most of those cooked with the skins had a flavour, not unpleasant, but peculiar to the American plums. The proper proportions to be used in preserving each variety will have to be learned by experience. The following are some of the notes made on the preserves, arranged in descending order of merit:—

Bixby.—1-lb. sugar to 1-lb. fruit, unpeeled; good colour, good flavour, skin tender.

Cheney.—1-lb. sugar to 1-lb. fruit, peeled; attractive amber colour, good flavour.

Cottrell.—1 lb. sugar to 1-lb. fruit, peeled; amber colour, sweet, rich, good.

New Ulm.—1-lb. sugar to 1-lb. fruit, peeled; attractive, pale, good flavour.

Mankato.—1-lb. sugar to 1-lb. fruit, peeled; pale, clear amber, good flavour, but too sweet.

Bouncer.—1-lb. sugar to 1-lb. fruit, peeled; attractive, but too thick, good flavour.

Bixby.—1-lb. sugar to 1-lb. fruit, peeled; too sweet, not as good as unpeeled.

Cottrell.— $\frac{3}{4}$ -lb. sugar to 1-lb. unpeeled fruit; attractive colour, not as good as peeled.

Silas Wilson.—1-lb. sugar to 1-lb. fruit, peeled; dull amber colour, good, but too rich.

Bouncer.— $\frac{3}{4}$ -lb. sugar to 1-lb. fruit, unpeeled; attractive colour, good flavour, but skin tough.

Mankato.—1-lb. sugar to 1-lb. fruit, unpeeled; fairly attractive colour, but too sweet.

New Ulm.—1-lb. sugar to 1-lb. fruit, unpeeled; attractive colour, good flavour, but tough skin.

Silas Wilson.—1-lb. sugar to 1-lb. fruit, unpeeled; good flavour, but skin tough.

American Eagle.— $\frac{3}{4}$ -lb. sugar to 1-lb. fruit, deep red, rather tough skin, medium quality.

Cheney.—1-lb. sugar to 1-lb. fruit, unpeeled; unattractive colour, but skin breaks well.

The following recipes for canning and preserving American plums, published by the late Prof. Goff, of the Wisconsin Experiment Station, were consulted and in part adopted in making the preserves:—

"The native plums, especially those with firm pulp after being treated by any of the methods mentioned below, are well adapted to all purposes for which the foreign plums are used. As a rule, more sugar is required for the native plums, but the preparations are rich in proportion. The harshness in the skin and stone of some native plums is readily removed by steaming them in an ordinary cooking steamer until the skin cracks; or pour over them boiling water to which has been added common baking soda in the proportion of half a teaspoonful to a quart. The thicker skinned varieties may be readily peeled by placing them in boiling water two or three minutes. The recipes follow:—

Canning.—Pick the fruit when well coloured but a little hard, steam or cook in a porcelain-lined kettle until tender, put in cans that have first been treated to boiling water, and cover with boiling syrup made of equal parts of granulated sugar and water, filling the can to the top; then run a silver knife around the can inside and let out the air, and seal at once. Plums cooked in the syrup are likely to be tough. Canned plums may be used for pies and for mixing with or flavouring other fruit. Plums are often canned without sugar to be used in winter for making fresh plum butter. The juice of canned plums makes excellent jelly. One lady recommends splitting native plums to the stone on one side before cooking, to avoid crumbling.

Drying.—De Soto, Wyant, and doubtless other varieties, may be pared, pitted and spread on plates, lightly sprinkled with sugar and dried, first in the oven and later in the sun. Cook like dried peaches.

Plum Jelly.—The fruit should be gathered when only part ripe—about half coloured. This point is very essential. Put plums in a large granite or porcelain kettle—the latter is best—with barely enough water to cover them. Cook until tender, but not until they are in a pulpy mass. Having previously covered a large jar with a cloth, strain the fruit in and let the juice drop through, but do not squeeze. When all has drained through, strain once or twice more through another cloth, until the juice is perfectly clear. To one measure of juice provide one measure of granulated sugar, but do not put together at once. A very important point in the making of all jelly is that only a small quantity should be cooked at one time. Into a medium sized kettle put, say, four tumblers of juice, let it boil briskly fifteen or twenty minutes, then add the four tumblers of sugar, and in a very short time, usually from three to ten minutes, the jelly will be finished, light, clear, and delicious. To test the jelly, dip a spoon into the boiling juice and sugar and hold it up; when the jelly clings to the spoon in thick drops, take it off quickly and put into jelly glasses. The plum pulp which is left can be put through a cullender and used for plum butter.

Another recipe—Plum preserves.—Take equal weights of fruit and sugar, place in stone jar a layer of fruit, then a layer of sugar, alternating thus until quantity desired is reached. Let stand over night; in the morning drain off the syrup that will have formed, into a porcelain kettle, place same over a fire and let syrup come to a boil, then pour it over fruit in jar again, repeat this every day until the fourth heating, when fruit and syrup are both put in kettle and boiled for a few minutes. Place same in glass jars while hot, seal and put away in some cool and preferably dark place.

Still another recipe.—To each pound of plums add a pound of sugar, put the fruit into boiling water until the skins will slip, peel and sprinkle sugar upon each layer of fruit in a bowl, allowing them to stand over night, then pour off the juice, bring quickly to a boil, skim and add the plums, cook very slowly till tender and clear, which will take about one-half hour, take them out carefully and put into a pan, boil the syrup for a few minutes longer until it thickens, pour it over the fruit, seal or tie them up.

SPRAYING.

It should not now be necessary to point out the advantages of spraying to control insect enemies and fungus diseases, as so much has been said and written on this sub-

ject. The beneficial results where spraying has been thoroughly done and the conditions have been favourable have also been so marked that the advantages of spraying need no further proof. It is, however, found necessary to continually advocate this practice. Fruit growers become discouraged after an unfavourable season for spraying, when the conditions are favourable for the development of fungi and rainy weather prevents the application of mixtures altogether or which if applied are washed off by the rain almost as soon as the work is done. With an experience of this kind fruit growers are often inclined to stop spraying and let nature take her course. It sometimes happens, however, that the mixtures and solutions have been improperly made or the wrong mixture has been used to destroy a certain insect or disease and the fruit grower wonders why he does not get good results. It is more frequently the case, however, that the spraying is not thoroughly done. The object of spraying is to cover the leaves, fruit and bark with the fungicides and insecticides in order that the latter may destroy the insects or diseases which come in contact with them, and in order that the materials may be distributed evenly over the tree it should be forced out of the pump in a fine mist-like spray, to accomplish which it is necessary to have a good nozzle. If the trees are not covered with the mixtures and solutions at all times when the insects and diseases may be affected by them, the trees and fruit may be injured just in proportion to the thoroughness and continuity with which the work is done. It is, therefore, important that every fruit grower should know the life history of every insect and disease which injures his trees or fruit, in order that he may know the best time to spray for each enemy. A spraying calendar has been published at intervals at the Central Experimental Farm by the Entomologist and the Horticulturist, in which are given full directions for the preparation of the different formulæ recommended and the time when each spraying should be made. These calendars are of great value to fruit growers and should be in the hands of everyone. Although a certain number of applications are recommended for the prevention and control of the different insects and diseases referred to, it may be necessary to make more. If, for instance, a heavy rain came on, say, within 24 hours after a spraying had been made, which washed most of the material off, another application should be made as this might be the very time when the disease or insect which it is desirable to control is making the greatest headway. It is often too late, also, when a spraying for a certain purpose is made, and labour and material are thus practically lost. Spraying is rather expensive when there is a large orchard to cover. It is, therefore, very important that the right mixtures and solutions are used; that they are prepared properly and applied thoroughly, constantly and at the right time, and that the trees be kept covered with the mixtures and solutions during all the time when injury is likely to occur.

As the mixtures and solutions may have very injurious effects on the trees if improperly made, and as they may prove of little or no value if not applied at the right time, the formulæ recommended in this Bulletin should be closely followed.

DISEASES OF THE PLUM.

RIPE ROT, BROWN ROT (*Monilia fructigena*).—The ripe rot causes serious injury to the plum crop every year, especially in the province of Ontario. Its spread is so rapid that a fine crop of plums is soon rendered almost worthless. The disease is usually first noticed on the ripening fruit by a discoloration of the skin, which becomes brown or black and is soon covered by small pustules or clusters of spores, the fruit rotting and falling to the ground. If the weather is damp and sultry the conditions are most favourable to the rotting of the fruit, and a large proportion of the crop will be destroyed in one day. Sometimes the disease does not show when the fruit is picked, and infected fruit is shipped and rots before reaching its destination, causing serious loss to shipper or buyer. This disease is often not recognized in the spring, at which time it attacks the twigs, fruit spurs and blossoms, causing them to blacken and wither. At this time, also, there is often an exudation of gum from the twigs and spurs, brought about by this disease.

Remedies.—The ripe rot fungus spreads by means of spores which germinate early in the spring and penetrate the twigs from the leaf and flower buds on which they alight.

In order to destroy as many of these spores as possible all diseased fruit should be gathered and burned whether it is on the ground or on the tree. This fruit harbours myriads of spores, which endure the winter and are capable of infecting the trees the following spring.

The trees should be thoroughly sprayed in time to destroy the spores before they germinate in the spring. The first spraying should be made with poisoned Bordeaux mixture, or a sulphate of copper solution, 1 lb. sulphate of copper to 25 gallons of water, shortly before the buds start to develop, and with poisoned Bordeaux mixture just before the blossoms open. These sprayings are very important and should never be neglected. After the trees have bloomed they should be thoroughly sprayed again with ordinary poisoned Bordeaux mixture, and also ten days or two weeks before the fruit begins to colour. The trees should also be sprayed with ammoniacal copper carbonate when the fruit is beginning to ripen. This will destroy the spores which appear in great numbers on the mature plums, and not discolour the fruit. Plums which touch one another on the tree give very favourable conditions for the spread of the disease from one fruit to another. Being close together, moisture is retained on the skin and the spores which may be on one fruit germinate readily and soon infect the next, and thus the disease spreads rapidly. Thinning the fruit would make the conditions much less favourable for the development of the disease. All discoloured or dead wood should be cut out and burned in the meantime. If spraying is thoroughly done the injury from this disease will be much lessened.

BLACK KNOT (*Plowrightia morbosa*).—The black knot is more feared by the average fruit grower than almost any other disease. Its development appears to him mysterious and its control impossible. Many orchards have been completely ruined by the black knot, which spreads very rapidly once it has gained a foothold in the orchard. Trees which have been cared for from the beginning are, however, not so subject to this disease, and if treated properly from the outset it may be kept in check and even eradicated altogether, if infection does not take place from neighbouring orchards or from planting trees from infected nurseries.

The black knot is a fungus which spreads by means of spores. A spore is blown through the air and alights in the axil of a leaf, on a spur, or on some other part of the tree where it can get a resting place. When conditions are favourable the spore germinates, penetrates the tree and grows in it. In the spring yellowish swellings appear on the branches, the first visible indication of the disease, and during the months of May and June become darker in colour with a velvety surface which is caused by innumerable spores which cover the exposed part of the lump or knot. These spores are soon blown away and alight on other branches and trees. These germinate and penetrate the tree as already described and new knots develop in time from them. A second crop of spores is produced in autumn or early winter, but these are not liberated from the knot until the latter part of the winter or early in the spring when they are distributed and alighting on the trees germinate as the others. The mycelium is not all exhausted by the production of the knots in spring, but is capable of producing more from the same knot the following year. The disease is also liable to be carried from one orchard to another, which makes it difficult to control sometimes. To fight a disease which is provided with such sure means of reproduction and dissemination requires watchfulness, thoroughness, and continuous effort until it is eradicated.

Remedy.—If trees are already badly affected with this disease the best plan is to cut them down and burn them. If only moderately affected, the knots should be removed and burned and the orchard thoroughly cultivated and sprayed. Vigorous trees are not so subject to the disease as those making little growth, and vigorous trees will recover more quickly from wounds made by removing the knots.

As the early spores appear in the latter part of spring or early in summer, the knots should be removed as soon as possible after they appear. When they are on small branches these should be cut off three to six inches below the affected part and burned immediately. When this is not practicable without badly injuring the tree the knot should be removed with a sharp pruning knife and the wound given a thorough painting with pure kerosene, after which it should be covered with grafting wax or lead

paint. Great care should be taken to prevent the kerosene running on the branch as it may injure it. Old knots which cannot be removed with the knife should get a good painting with kerosene. By putting some colouring material in the kerosene, one can more readily see when a wound has been painted well. All knots should be burned. The treatment already described is only for the knots. Unless the life history of this disease is known a fruit grower might imagine that if the knots were removed as they appeared that it was all that was necessary. This might be all, if the knots were removed before the spores were formed and distributed, but if one knot were missed it would be capable of producing myriads of spores which would soon re-infect the orchard.

The advantages of spraying with Bordeaux mixture were clearly demonstrated in an experiment conducted at the Cornell Experiment Station, the results of which were published in Bulletin No. 81 of that station. It was found that the number of knots was reduced from 2,000 to 165 by spraying.

A large proportion of the late spores which are exposed in late winter or early spring and are ready to germinate when conditions are favourable, would probably be destroyed by an application of sulphate of copper solution at the rate of 1 lb. of sulphate of copper to 25 gallons of water when the trees are still dormant in the spring. A second spraying of poisoned Bordeaux mixture should be made just as or before the buds are breaking, which would also be a good time to spray for other diseases of the plum and for insects, and a third spraying should be made at the time when the knots are at the velvety or spore producing stage in order to destroy these spores. A fourth application would also probably be well worth the expense. Every precaution should be taken to prevent the spread of this disease which has destroyed so many plum trees in Canada.

SPOT OR BLIGHT OF THE NATIVE PLUM (*Cladosporium carpophilum*, V. Thumen).—The almost complete absence of native plums during recent years in the Ottawa district and elsewhere in eastern Ontario and the province of Quebec, is due in a large measure or almost entirely to the disease known as blight. The fruit forms and reaches more than half its size, but colours prematurely. When affected by the disease it shrivels and falls to the ground without ripening. If the fruit is examined when half grown or later, small pale green or yellow patches will be noticed. These gradually enlarge until finally they are about half an inch in diameter, at which time the blotches are darker in colour, of more irregular outline and are raised on the skin. The Americana plums are not, as a rule, seriously affected with this disease, which is principally confined to the Nigra varieties.

Remedy.—This fungus is nearly related to the apple spot and can be satisfactorily treated in much the same way. The trees should be sprayed with Bordeaux mixture just after the blossoms fall, again two weeks later, and a third time two weeks after the second application. It is also advisable to spray a fourth time with ammoniacal copper carbonate just when the fruit is beginning to colour. The native varieties ripen early, and if the ordinary Bordeaux mixture were applied the last time, the fruit might remain stained. The ammoniacal copper carbonate does not leave a noticeable stain on the fruit. This remedy has been very satisfactorily used by one grower in particular near Ottawa who has thus been able to grow native plums very profitably. The Americana varieties may be top grafted on the native ones with the result that there will be less disease as the latter are not as much affected as the native. All other plum trees not looked after or bearing poor fruit should be burned; also all fruit which is diseased.

SHOT-HOLE OR LEAF-SPOT (*Cylindrosporium padi*).—The first indications of this disease are small, yellowish spots with reddish margins, which appear on the young leaves. These spots increase in size and finally reach a diameter averaging about one-eighth to one-sixth of an inch. When fully grown, the central part dries up and drops out, leaving a clean cut around the margin, very suggestive of a shot-hole, after which the disease is named. When these spots occur in large numbers, as they frequently do, so much of the leaf is destroyed that it drops prematurely. The early dropping of the leaves pre-

vents the fruit, twigs and buds reaching their full development, and on this account causes serious injury where the disease is troublesome. Prof. Beach, Geneva, N.Y., who conducted experiments to control this disease on European plums, found the Bordeaux mixture satisfactory and recommends three applications. The first about ten days after the blossoms fall; the second, three weeks later, and the third, three or four weeks after the second. At the Central Experimental Farm, Americana and Nigra plums have been thoroughly sprayed with Bordeaux mixture from three to five times, without satisfactory results. Some varieties are more susceptible to the disease than others.

PLUM POCKETS—*Exoascus (Taphrina)*.—The disease of plum known as 'plum pockets' does not cause widespread injury, but in some places does occasion considerable loss. The mycelium of the fungus which causes the pockets lives year after year in the same tree, and thus it is not necessary for a tree to be infected by the spores every year in order to perpetuate the disease. The fruit is affected soon after the trees have blossomed and is indicated by the unnatural swelling and bladder-like appearance of the fruit and by its unusual yellow colour. When the spores of the disease which has been working inside the fruit appear on the surface they give the pockets a gray appearance. Later on, the pockets turn almost black and fall to the ground. The leaves and twigs are also noticeably affected with this disease, the former becoming curled and unhealthy looking and the twigs swelling unnaturally. There is no known effective remedy for this disease, but it will probably be much lessened if the affected parts with the pockets are cut off and burned.

GUMMOSIS.—The exudation of gum from plum trees, which is known as gumming or gummosis, is quite common in plum orchards. It has been given careful study and is not attributed to any one disease. The conclusion reached is that any weakening of the tree by severe pruning, by winter, by injury to the trunk or branches, or by diseases of different kinds will cause gumming. It is often very noticeable when trees are affected with the *Monilia* or Ripe Rot fungus. It appears to follow any breaking or injury to the wood tissue.

MICE.—Mice often girdle the trees in the orchard in winter, especially when it is in sod or when there is rubbish lying about in which they like to harbour. Everything in the way of rubbish should, however, be removed before winter. Their depredations may be prevented either by wrapping the trunks with building paper in autumn and banking up the earth about the base to the height of about a foot; by encircling the trunk with fine galvanized iron netting; or by using the veneer protectors used to prevent scalding. Where the latter are used the earth should be banked up a little at the base to prevent the mice from going underneath.

If a tree is girdled by mice it usually dies. If, however, as soon as the injury is noticed, the wound is cleaned and covered with grafting wax and wrapped with cloth so that the air is excluded and the wood prevented from drying out, the sap which rises through the soft wood will continue to do so and returning through the inner bark, growth will be made all around the upper part of the wound, and if the latter be not too large there is a chance of its healing over. If, however, the wood becomes dry before the bandage is put on it will almost certainly die. When the wax and bandage are applied the tree should be headed back considerably to lessen the amount of transpiration of moisture, as there will not be as much sap rise as if the tree were uninjured. Girdled trees are sometimes saved by connecting the upper and lower edges of the girdle with scions, which are inserted all around the trunk. Mice may be destroyed in the orchard by using a mixture of one part by weight of arsenic with three parts of corn meal. To use this safely nail two pieces of board each six feet long and six inches wide together so as to make a trough. Invert this near the trees to be protected and place about a tablespoonful of the poison on a shingle and put it near the middle of the run, renewing the poison as often as is necessary.

PLUM INSECTS.

By JAMES FLETCHER, *Entomologist and Botanist.*

Insects may be divided into two classes according to the nature of their mouth parts. Those of the first or larger division, Biting Insects, are furnished with mandibles, or biting jaws, by means of which they consume the substance of their food, as caterpillars, beetles, &c. The second class, Sucking Insects, have, instead of mandibles, a beak or tube, by means of which they suck up their food in a liquid form from beneath the surface, as the true bugs, plant-lice, &c. As regards the insects of the first class, all that it is necessary to do is to place on the food plant some poisonous material which will be eaten with the food. With the second class, however, this would be useless, for they would push their beaks through the poisonous covering on the outside of the food plant and extract from the interior the juices upon which they live. For this class, therefore, some substance must be used which will kill by mere contact with their bodies.

INSECTICIDES AND FUNGICIDES.

The following are the formulæ of standard remedies which are recommended by the Entomologist and Botanist:—

I. PARIS GREEN.

For Biting Insects.

Paris green.....	1 lb.
Lime (fresh).....	1 lb.
Water.....	200 gallons.

For dry application.—1 lb. Paris green with 50 lbs. flour, land plaster, slaked lime or any other perfectly dry powder.

II. KEROSENE EMULSION.

(RILEY-HUBBARD FORMULA.)

For Scale insects and Plant-lice.

Kerosene (coal oil).....	2 gallons.
Rain water.....	1 gallon.
Soap.....	$\frac{1}{2}$ lb.

Dissolve soap in water by boiling; take from fire, and, while hot, turn in kerosene and churn briskly for five minutes. To be diluted before use with nine parts of water.

III. WHALE-OIL SOAP.

For scale-insects (young).....	1 lb. in 5 gallons water.
For aphid.....	1 lb. in 8 "
For San José scale (in winter).....	$2\frac{1}{2}$ lbs. in 1 "

IV. TOBACCO AND SOAP WASH.

For Plant-lice or Aphid.

Soak in hot water for a few hours 10 lbs. of tobacco leaves (home grown will do); strain off, and add 2 lbs. of whale-oil soap. Stir until all is dissolved, and dilute to 40 gallons. Apply early and two or three times at short intervals.

V. POISONED BORDEAUX MIXTURE.

For Fungi and Insects on Fruit-trees.

Copper sulphate (bluestone).....	4 lbs.
Lime (fresh)	4 "
Paris green.....	4 ozs.
Water (1 barrel).....	40 gallons.

Dissolve the copper sulphate (by suspending it inside a wooden or earthen vessel containing 4 or 5 or more gallons of water). Slake the lime in another vessel. If the lime, when slaked, is lumpy or granular, it should be strained through coarse sacking or a fine sieve. Pour the copper sulphate solution into a barrel, or it may be dissolved in this in the first place; half fill the barrel with water, add the slaked lime, fill the barrel with water and stir thoroughly. It is then ready for use.

Stock solutions of dissolved copper sulphate and of lime may be prepared and kept in separate covered barrels throughout the spraying season. The quantities of bluestone, lime and water should be carefully noted.

VI. COPPER SULPHATE SOLUTION.

For Fungi.

Copper sulphate (bluestone).....	1 lb.
Water	25 gallons.

As soon as dissolved it is ready for use. *For use only before the buds open.*

VII. AMMONIACAL COPPER CARBONATE.

For Fungi.

Copper carbonate.....	5 oz.
Ammonia	2 quarts.
Water (1 barrel).....	40 gallons.

Dissolve the copper carbonate in the ammonia. The ammonia and concentrated solution should be kept in glass or stone jars, tightly corked. It is ready for use as soon as diluted with the 40 gallons water. To be used when Bordeaux cannot be applied on account of staining the fruit.

THE WORST ENEMIES OF THE PLUM TREE.

ATTACKING THE FOLIAGE.

1. The EYE-SPOTTED BUD-MOTH (*Tmetocera ocellana*).—Small, dark brown caterpillars, $\frac{1}{4}$ of an inch in length, with black heads and collars, destroying the buds when just unfolding and later attacking the leaves, two or three of which they attach together, feeding inside. They also sometimes bore down the centre of the twig. *Remedy*:—Spray early with a strong Paris green wash (Paris green 1 pound, fresh lime 1 pound, water 100 gallons).

2. The CIGAR CASE-BEARER (*Coleophora fletcherella*).—Small orange-coloured caterpillars with black heads, encased in brown leathery cigar-shaped cases, which they carry about with them. They pass the winter as caterpillars on the twigs, and cluster around the opening buds, injuring the foliage and flowers. *Remedy*:—Spray early with the wash mentioned under No. 1 above, or with kerosene emulsion (Formula II).

3. **TENT CATERPILLARS** (*Clisiocampa*).—Two kinds attack the foliage of the plum as well as many other trees. The Apple-tree Tent Caterpillar forms a tent in the fork of two twigs; the Forest Tent Caterpillar does not make a tent but spins a flat mat of silk on the side of a branch or on the trunk; to these resting places the young caterpillars resort when not feeding. The mature insects are thick-bodied, reddish-brown moths expanding about $1\frac{1}{2}$ inches across the wings, which are crossed obliquely by two bands. These bands are pale in the first named, but dark in the moth of the Forest Tent Caterpillar. During July the females lay rings of about 200 eggs on the twigs of the trees, in which state the insect passes the winter. *Remedies*:—Collect and destroy the egg clusters during the winter. Spray the trees with poison (Formula I. or V.) directly the young caterpillars are noticed. All tents should also be cut off and destroyed early before the leaves hide them.

4. **PLANT-LICE OR APHIDS** (*Aphis prunifoliae*, *Hyalopterus pruni*).—In appearance and habits much the same as the Apple Aphid, clustering on the underside of leaves, sucking out the juices and causing the leaves to become wrinkled and twisted. *Remedies*:—Whale-oil soap 1 lb. in 6 gallons of water, tobacco and soap wash (Formula IV) or kerosene emulsion (Formula II). Begin to spray when the plant-lice are first noticed.

5. **THE PEAR-TREE SLUG OR CHERRY-TREE SLUG** (*Eriocampa cerasi*).—In June and August, slimy, greenish-brown, slug-like caterpillars, $\frac{1}{2}$ inch long, feeding on the upper surface of the leaves, often doing considerable damage to plum trees. *Remedies*:—Spray with a weak solution of Paris green or dust with freshly slaked lime, or Paris green diluted with 50 times its weight of some dry powder.

6. **PLUM LEAF CATERPILLARS**.—There are other kinds of leaf-eating caterpillars which occasionally occur in sufficient numbers on the foliage of plum trees to do serious injury. *Remedy*:—Spraying regularly with Formula V. will prevent injury from these insects.

ATTACKING THE WOOD.

7. **THE SHOT-BORER** (*Xyleborus dispar*).—Small blackish beetles, which bore into the trunks and limbs, causing serious damage in apple and plum orchards. *Remedy*:—Wash the trees liable to attack three times, early and late in June and once in July, with the following:—Soft soap, 1 gallon; water, 3 gallons; carbolic acid, $\frac{1}{2}$ pint.

OCCURRING ON THE BARK.

8. **THE SAN JOSÉ SCALE** (*Aspidiotus perniciosus*).—Minute, almost circular scale-insects, one-thirtieth of an inch wide, shaped like an inverted saucer, with a depressed ring around a central point; inside this ring, black or dark coloured. This very inconspicuous insect when in small numbers is easily overlooked, but when abundant gives to the bark a dirty, scurfy and grayish colour, as though dusted with ashes. *Remedies*:—This is by far the most difficult insect to eradicate which the fruit growers have ever had to deal with. The treatments which have so far given the best results are: (1) the ordinary kerosene emulsion (Formula II.) two treatments during the summer (an extra one may advantageously be applied in May, just before the foliage is so thick that it is difficult to reach all parts of the tree), the first summer spraying in the middle of June, and the second one after the fruit is picked; this to be followed in winter or spring with a thorough general spraying with lime and sulphur, 1 lb. lime and $\frac{1}{2}$ lb. of sulphur to one gallon of water boiled together until dissolved. (2) Whale-oil soap $2\frac{1}{2}$ lbs. to the imperial gallon of water, the best time to spray being just before the buds burst in spring. The soap should be dissolved in hot water and applied as hot as is practicable.

9. **THE NEW YORK PLUM SCALE** (*Lecanium cerasifera*).—Conspicuous, dark brown, hemispherical scales, about $\frac{1}{8}$ of an inch long by $\frac{1}{8}$ of an inch wide, occurring at all times of the year, clustered along the small branches, particularly along the lower sides. The presence of this enemy upon a plum tree may be detected, especially in July and August and also in the spring, by the filthy black condition of the bark due to the growth of a fungus upon the copious deposit of honey-dew which is emitted by the young scale insects during the time of their growth. *Remedies*:—Spray the trees during the winter with kerosene emulsion (Formula II.) diluted with four parts of water, or with the whale-oil soap solution (Formula III.)

ATTACKING THE FRUIT.

10. **THE PLUM CURCULIO** (*Conotrachelus nenuphar*).—Small, rough, grayish beetles about $\frac{1}{8}$ of an inch long. The females, in the operation of egg-laying, make upon the sides of plums small crescent-shaped marks, with a single hole in the centre of each. An egg is laid in the central spot, from which hatches a white grub; this soon destroys the fruit. *Remedies*:—(1) The mature beetles feed in early spring upon the unopened buds and afterwards upon the young leaves and may be reached by spraying the trees before the buds open with Paris green (Formula I), repeating this as soon as the fruit has formed and spraying ten days later with the poisoned Bordeaux mixture (Formula V). (2) The beetles are sluggish in the early morning and drop from the trees if a sudden jar be given to the trunk. This jarring, if repeated every day or two over a sheet or other receptacle, will be the means of greatly lessening the numbers of the mature insects. The beetles drop readily and lie quietly for some time, when they can be easily collected and destroyed.

NOTE.

The operation of 'spraying' consists of applying liquids by means of a force pump and spraying nozzle, with such force as to break up the liquid so thoroughly that it falls upon the plants treated as an actual mist or spray. The word 'spraying,' however, to many who endeavour to practise this operation, has still little more meaning than doing something in any sort of way, to fruit trees with a spraying pump. Such terms as 'sprinkling' and 'showering' are inaccurate for the operation here intended. Unfortunately, much of the so-called spraying as usually carried out, could more accurately be designated as sprinkling or showering, which describe a much less careful and less even distribution of liquids.

The Entomologist and Botanist will be glad to give further information concerning attacks upon crops by insects if inquiry be made by letter.

