



# IMAGE EVALUATION TEST TARGET (MT-3)



Photographic  
Sciences  
Corporation

23 WEST MAIN STREET  
WEBSTER, N.Y. 14580  
(716) 872-4303

**CIHM/ICMH  
Microfiche  
Series.**

**CIHM/ICMH  
Collection de  
microfiches.**



**Canadian Institute for Historical Microreproductions / Institut canadien de microreproductions historiques**

**© 1985**

# Technical and Bibliographic Notes/Notes techniques et bibliographiques

The Institute has attempted to obtain the best original copy available for filming. Features of this copy which may be bibliographically unique, which may alter any of the images in the reproduction, or which may significantly change the usual method of filming, are checked below.

L'Institut a microfilmé le meilleur exemplaire qu'il lui a été possible de se procurer. Les détails de cet exemplaire qui sont peut-être uniques du point de vue bibliographique, qui peuvent modifier une image reproduite, ou qui peuvent exiger une modification dans la méthode normale de filmage sont indiqués ci-dessous.

☐ Coloured covers/  
Couverture de couleur

☐ Coloured pages/  
Pages de couleur

☐ Covers damaged/  
Couverture endommagée

☒ Pages damaged/  
Pages endommagées

☐ Covers restored and/or laminated/  
Couverture restaurée et/ou pelliculée

☐ Pages restored and/or laminated/  
Pages restaurées et/ou pelliculées

☐ Cover title missing/  
Le titre de couverture manque

☒ Pages discoloured, stained or foxed/  
Pages décolorées, tachetées ou piquées

☐ Coloured maps/  
Cartes géographiques en couleur

☐ Pages detached/  
Pages détachées

☐ Coloured ink (i.e. other than blue or black)/  
Encre de couleur (i.e. autre que bleue ou noire)

☒ Showthrough/  
Transparence

☐ Coloured plates and/or illustrations/  
Planches et/ou illustrations en couleur

☐ Quality of print varies/  
Qualité inégale de l'impression

☒ Bound with other material/  
Relié avec d'autres documents

☐ Includes supplementary material/  
Comprend du matériel supplémentaire

☐ Tight binding may cause shadows or distortion along interior margin/  
La reliure serrée peut causer de l'ombre ou de la distortion le long de la marge intérieure

☐ Only edition available/  
Seule édition disponible

☐ Blank leaves added during restoration may appear within the text. Whenever possible, these have been omitted from filming/  
Il se peut que certaines pages blanches ajoutées lors d'une restauration apparaissent dans le texte, mais, lorsque cela était possible, ces pages n'ont pas été filmées.

☐ Pages wholly or partially obscured by errata slips, tissues, etc., have been refilmed to ensure the best possible image/  
Les pages totalement ou partiellement obscurcies par un feuillet d'errata, une pelure, etc., ont été filmées à nouveau de façon à obtenir la meilleure image possible.

☒ Additional comments:/  
Commentaires supplémentaires:

Pagination as follows : p. [368]-372.

This item is filmed at the reduction ratio checked below/  
Ce document est filmé au taux de réduction indiqué ci-dessous.

10X	14X	18X	22X	26X	30X
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
12X	16X	20X	24X	28X	32X

The copy filmed here has been reproduced thanks to the generosity of:



Medical Library  
McGill University  
Montreal

The images appearing here are the best quality possible considering the condition and legibility of the original copy and in keeping with the filming contract specifications.

Original copies in printed paper covers are filmed beginning with the front cover and ending on the last page with a printed or illustrated impression, or the back cover when appropriate. All other original copies are filmed beginning on the first page with a printed or illustrated impression, and ending on the last page with a printed or illustrated impression.

The last recorded frame on each microfiche shall contain the symbol ➡ (meaning "CONTINUED"), or the symbol ▼ (meaning "END"), whichever applies.

Maps, plates, charts, etc., may be filmed at different reduction ratios. Those too large to be entirely included in one exposure are filmed beginning in the upper left hand corner, left to right and top to bottom, as many frames as required. The following diagrams illustrate the method:



L'exemplaire filmé fut reproduit grâce à la générosité de:

Medical Library  
McGill University  
Montreal

Les images suivantes ont été reproduites avec le plus grand soin, compte tenu de la condition et de la netteté de l'exemplaire filmé, et en conformité avec les conditions du contrat de filmage.

Les exemplaires originaux dont la couverture en papier est imprimée sont filmés en commençant par le premier plat et en terminant soit par la dernière page qui comporte une empreinte d'impression ou d'illustration, soit par le second plat, selon le cas. Tous les autres exemplaires originaux sont filmés en commençant par la première page qui comporte une empreinte d'impression ou d'illustration et en terminant par la dernière page qui comporte une telle empreinte.

Un des symboles suivants apparaîtra sur la dernière image de chaque microfiche, selon le cas: le symbole ➡ signifie "A SUIVRE", le symbole ▼ signifie "FIN".

Les cartes, planches, tableaux, etc., peuvent être filmés à des taux de réduction différents. Lorsque le document est trop grand pour être reproduit en un seul cliché, il est filmé à partir de l'angle supérieur gauche, de gauche à droite, et de haut en bas, en prenant le nombre d'images nécessaire. Les diagrammes suivants illustrent la méthode.

rrata  
to

pelure,  
n à

32X

NOTES ON THE DISSECTION OF A CASE OF CON-  
GENITAL DISLOCATION OF THE HEAD OF THE  
FEMUR. By FRANCIS J. SHEPHERD, M.D., C.M., M.R.C.S.,  
Eng., *Demonstrator of Anatomy, McGill University, Montreal.*

A FEMALE subject about 50 years of age was brought to the dissecting-room of McGill University in the autumn of 1879 in whom it was noticed that the right leg was smaller and shorter than the left, and everted. The woman was very stout, and the head of the femur could not be distinctly made out, but it was noticed that the trochanter was higher than usual; abduction was limited, but otherwise the joint was freely movable. The tuberosity of right ischium was quite subcutaneous, the muscles appearing as if carried up with the trochanter-major. Amount of shortening, 2 inches. The exact nature of the lesion was not made out from an external examination, and it was only on dissection of the part that it was discovered to be a congenital dislocation.

*Dissection.*—There are no scars to be seen on the skin of gluteal or femoral region, indicating that old sinuses had existed, the skin is perfectly smooth. On removing the skin and fascia of the gluteal region, the fibres of the gluteus maximus, and gluteus medius muscles are found to be shorter than normal, their points of origin and insertion being approximated; the muscular tissue is much altered, in some places being completely changed into fatty tissue. I may here remark that all the muscles of the body have become fatty degenerated to a greater or less extent, but this condition is much more marked in some of the muscles about the affected hip. On removing the two superficial gluteal muscles (between which a lot of fat is seen), the quadratus femoris, obturator externus, gemelli, and obturator internus are brought into view. These muscles, instead of passing transversely outwards to get to their insertion, pass upwards; the pyriformis is found to pass outwards and slightly upwards, instead of downwards. The gluteus medius, obturator internus, gemelli, and pyriformis in passing to their insertions go over the head of the femur. More of the obturator externus is

exposed to view than usual; this muscle passes upwards to the great trochanter close to the old capsule, which is seen lying between it and the gluteus minimus; the obturator externus is much longer than normal, and is quite a strong muscle, the tissue of which is healthy and without any trace of fatty degeneration. The gluteus minimus is shortened, and nearly all composed of fibrous tissue, it is pushed upwards by the ascent of the great trochanter, and has a smaller origin than usual, owing to the space on the dorsum of the ilium between the middle and inferior curved lines being partly occupied by the new socket for the head of the femur. This muscle covers the head of the femur, and is so closely united to the new capsule that it can only in places be separated from it.

The adductors and gracilis muscles are shortened and tense, and have participated but slightly in the general degeneration; the pectineus is smaller than usual, its outer portion being fibrous; it is attached as usual to the shaft of the femur. The iliacus internus muscle has nearly all atrophied away, and now consists of a very thin flat layer of muscular fibres, streaked with fat, which become united to the tendon of the psoas magnus, an inch above Poupart's ligament, so that none of its muscular fibres are seen below this ligament; above and below this muscle in the iliac fossa is a large deposit of fat. The psoas magnus has the usual origin, but consists almost entirely of fatty tissue, streaked here and there with muscular fibres, it ends in a flat tendon a little higher up than usual, and after being joined by the wasted iliacus muscle passes as a narrow, thin, ribbon-like tendon in the groove below the anterior inferior spine of the ilium under Poupart's ligament, in a direction outwards and upwards, and becomes blended with the anterior part of the old capsule.

*Ligamentous Structures.*—Two capsular ligaments are seen, the old and the new. The *old capsule* is still attached to the margin of the acetabulum, and is much thickened; the ilio-femoral ligament is well marked, being strengthened by the blending with it of the tendon of the psoas and iliacus muscles. The capsule is stretched upwards and outwards, following, of course, the ascent of the head of the femur; in its upper and posterior part is seen the head of the femur protruding through

a slit; this slit embraces the neck of the femur, and it appears as if the head of the femur has worn away the capsule at this point by pressing it against the dorsum ilii. The *new capsule* is attached at its pelvic extremity around the socket which has been formed on the dorsum ilii by the wearing away of the bone, —at its femoral end it is attached internally to the slit in the old capsule through which the head of the femur protrudes, and externally is continuous with the inner surface of the gluteus minimus. It is ligamentous in structure, its inner surface is smooth, and a number of small villous processes are seen hanging from it. No synovial fluid can be seen. The *ligamentum teres* is entirely absent, no trace of it being seen either in the old acetabulum or in the head of the femur; it has probably been worn through and the two ends absorbed.

*Osseous Structures.—Femur.* The whole bone is perceptibly smaller than that of the opposite side. The head is remarkably altered in appearance, it is of less size than normal, and where the ligamentum teres should be attached it is flat and devoid of cartilage, as if it had been worn away or had lost its epiphysis; on the upper part of this flattened surface a shallow groove is seen, crescentic in form. The remaining part of the head is covered with cartilage. The compact bony tissue covering the head is remarkably thin, and on breaking it through the cancellated structure is found to be very soft. The neck of the bone is much shortened, and forms a right angle with the shaft; the upper part of the shaft is arched outwards; where the gluteus maximus was attached a rather prominent crest is seen. The *lesser trochanter* is absent, its place being occupied by a shallow groove.

*Pelvis.*—The whole right half is smaller than the left. The wing of the right ilium is much thinner, more upright, more curled inwards, and altogether smaller than the left; the anterior inferior spine is prominent, and in consequence the groove below it in which lay the tendon of the psoas and iliacus is deeper. The rami of the pubis and ischium on the right side are flatter, thinner, and seem to spread out more than those of the opposite side, that is if a perpendicular be dropped from the symphysis pubis, the angle formed by it with the rami of the right side is much more obtuse than that formed by it with the

rami of the left. The right obturator foramen has lost its triangular shape, and is now broadest in its transverse diameter. The *acetabulum* is much altered in appearance, being a mere triangular depression in the bone; the apex of the triangle is upwards and to the right, and the base corresponds to the cotyloid notch. The edges of this triangular depression are smooth and curled inwards, and but slightly covered with fibrocartilage; the transverse ligament has almost disappeared. Around the edges of the depression is attached the old capsular ligament. The bottom of the cavity is completely ossified, and shows no trace of any disease having existed, the depression for the Haversian gland is well seen, it is small, being about the size of a sixpence. The acetabulum measures two inches in length, three-quarters of an inch in breadth, and half an inch in depth, it was filled with fatty tissue. The *new socket* for the head of the femur is an oval depression on the dorsum ilii between the middle and inferior curved lines, and on a level with the great sciatic notch, the inner edge of this new socket corresponding to the bony edge of the notch. It has not been deepened by ossific deposit, but appears to have been made by the wearing action of the head of the femur. It is two inches in diameter and a quarter of an inch in depth. Around its edge the new capsule is attached, and its floor is covered by a sort of periosteum, which receives fibres from the capsule. The ilium is not very thin at this point.

The measurements of the pelvis are altered, the diameters of the inlet being increased and outlet slightly diminished:—

Diameters.	Inlet.	Outlet.
Conjugate, . . . .	4½ inches.	4 inches.
Right oblique, . . .	5½ "	—
Left oblique, . . .	5½ "	—
Transverse, . . . .	6 "	4½ "

The distance between the anterior superior spines of the ilia measured nine inches.

The sacrum is slightly turned to the affected side, but otherwise is not much altered in appearance. There is no great anterior convexity of the lower dorsal and lumbar vertebrae, but the vertebrae incline laterally to the affected side.

*Remarks.*—(1.) *The absence of anterior convexity* in the lower



dorsal and lumbar regions was probably due to the atrophied condition of the psoas and iliacus muscles and their abnormal attachment to the old capsule. Probably during life, when the woman assumed the erect position, this anterior convexity existed, though not to the great degree that is usual.

(2.) *The absence of the lesser trochanter* was due no doubt to its having been torn away from its attachment to the femur at the time the dislocation occurred, and to its afterwards having been absorbed. This would account for the insertion of the psoas and iliacus tendon into the old capsule, the muscles thus having a very limited action would atrophy. This condition would also favour the opinion held by some that this form of dislocation is due to violence (at birth).

(3.) There was no twisting forwards of the head and neck of the femur, a condition which is described by some, and which, had it existed, would have accounted for the eversion. The eversion of the limb may have been only a *post-mortem* symptom due to the relaxed state of the muscle, the large capsule, and small head of femur.

---

*Note.*—The subject in which the above described dislocation occurred, owing to the unfortunate state of affairs in Canada, had been illegally obtained, so that no history could at the time be procured. I have since heard that this woman had suffered from "lameness" all her life, that she walked with a waddling motion, and also that she had borne a large family of children. This information I have fair reasons for supposing is authentic.

UR.

rophied  
normal  
en the  
nvexity

t to its  
at the  
g been  
psoas  
naving  
d also  
cation

eck of  
which,  
The  
*ortem*  
large

tion  
ada,  
time  
ered  
ling  
ren.  
ic.

