





**IMAGE EVALUATION
TEST TARGET (MT-3)**



**Photographic
Sciences
Corporation**

23 WEST MAIN STREET
WEBSTER, N.Y. 14580
(716) 872-4503

**CIHM/ICMH
Microfiche
Series.**

**CIHM/ICMH
Collection de
microfiches.**



Canadian Institute for Historical Microreproductions / Institut canadien de microreproductions historiques

© 1983

Technical and Bibliographic Notes/Notes techniques et bibliographiques

The Institute has attempted to obtain the best original copy available for filming. Features of this copy which may be bibliographically unique, which may alter any of the images in the reproduction, or which may significantly change the usual method of filming, are checked below.

L'institut a microfilmé le meilleur exemplaire qu'il lui a été possible de se procurer. Les détails de cet exemplaire qui sont peut-être uniques du point de vue bibliographique, qui peuvent modifier une image reproduite, ou qui peuvent exiger une modification dans la méthode normale de filmage sont indiqués ci-dessous.

- Coloured covers/
Couverture de couleur
- Covers damaged/
Couverture endommagée
- Covers restored and/or laminated/
Couverture restaurée et/ou pelliculée
- Cover title missing/
Le titre de couverture manque
- Coloured maps/
Cartes géographiques en couleur
- Coloured ink (i.e. other than blue or black)/
Encre de couleur (i.e. autre que bleue ou noire)
- Coloured plates and/or illustrations/
Planches et/ou illustrations en couleur
- Bound with other material/
Relié avec d'autres documents
- Tight binding may cause shadows or distortion along interior margin/
La reliure serrée peut causer de l'ombre ou de la distortion le long de la marge intérieure
- Blank leaves added during restoration may appear within the text. Whenever possible, these have been omitted from filming/
Il se peut que certaines pages blanches ajoutées lors d'une restauration apparaissent dans le texte, mais, lorsque cela était possible, ces pages n'ont pas été filmées.
- Additional comments:
Commentaires supplémentaires:

Pagination as follows : [i] - xii, [1] - 12, 13* - 15*, 13 - 18, 19* - 24*, 23 - 134 p.

- Coloured pages/
Pages de couleur
- Pages damaged/
Pages endommagées
- Pages restored and/or laminated/
Pages restaurées et/ou pelliculées
- Pages discoloured, stained or foxed/
Pages décolorées, tachetées ou piquées
- Pages detached/
Pages détachées
- Showthrough/
Transparence
- Quality of print varies/
Qualité inégale de l'impression
- Includes supplementary material/
Comprend du matériel supplémentaire
- Only edition available/
Seule édition disponible
- Pages wholly or partially obscured by errata slips, tissues, etc., have been refilmed to ensure the best possible image/
Les pages totalement ou partiellement obscurcies par un feuillet d'errata, une pelure, etc., ont été filmées à nouveau de façon à obtenir la meilleure image possible.

This item is filmed at the reduction ratio checked below/
Ce document est filmé au taux de réduction indiqué ci-dessous.

10X	14X	18X	22X	26X	30X
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
12X	16X	20X	24X	28X	32X

The copy filmed here has been reproduced thanks to the generosity of:




National Library of Canada

The images appearing here are the best quality possible considering the condition and legibility of the original copy and in keeping with the filming contract specifications.

Original copies in printed paper covers are filmed beginning with the front cover and ending on the last page with a printed or illustrated impression, or the back cover when appropriate. All other original copies are filmed beginning on the first page with a printed or illustrated impression, and ending on the last page with a printed or illustrated impression.

The last recorded frame on each microfiche shall contain the symbol \rightarrow (meaning "CONTINUED"), or the symbol ∇ (meaning "END"), whichever applies.

Maps, plates, charts, etc., may be filmed at different reduction ratios. Those too large to be entirely included in one exposure are filmed beginning in the upper left hand corner, left to right and top to bottom, as many frames as required. The following diagrams illustrate the method:



L'exemplaire filmé fut reproduit grâce à la générosité de:

Bibliothèque nationale du Canada

Les images suivantes ont été reproduites avec le plus grand soin, compte tenu de la condition et de la netteté de l'exemplaire filmé, et en conformité avec les conditions du contrat de filmage.

Les exemplaires originaux dont la couverture en papier est imprimée sont filmés en commençant par le premier plat et en terminant soit par la dernière page qui comporte une empreinte d'impression ou d'illustration, soit par le second plat, selon le cas. Tous les autres exemplaires originaux sont filmés en commençant par la première page qui comporte une empreinte d'impression ou d'illustration et en terminant par la dernière page qui comporte une telle empreinte.

Un des symboles suivants apparaîtra sur la dernière image de chaque microfiche, selon le cas: le symbole \rightarrow signifie "A SUIVRE", le symbole ∇ signifie "FIN".

Les cartes, planches, tableaux, etc., peuvent être filmés à des taux de réduction différents. Lorsque le document est trop grand pour être reproduit en un seul cliché, il est filmé à partir de l'angle supérieur gauche, de gauche à droite, et de haut en bas, en prenant le nombre d'images nécessaire. Les diagrammes suivants illustrent la méthode.

THE
NORTH AMERICAN
SYLVA;
OR,
A DESCRIPTION OF THE FOREST TREES
OF THE
United States, Canada and Nova Scotia,
NOT DESCRIBED IN THE WORK

OF
F. ANDREW MICHAUX,

AND CONTAINING ALL THE

FOREST TREES DISCOVERED IN THE ROCKY MOUNTAINS, THE TERRITORY
OF OREGON, DOWN TO THE SHORES OF THE PACIFIC, AND INTO
THE CONFINES OF CALIFORNIA, AS WELL AS IN VA-
RIOUS PARTS OF THE UNITED STATES.

ILLUSTRATED BY 121 FINE PLATES.

BY THOMAS NUTTALL, F. L. S.,

Member of the American Philosophical Society, and of the Academy of Natural
Sciences of Philadelphia, &c. &c. &c.

IN THREE VOLUMES.—VOL. I.

BEING THE FOURTH VOLUME OF MICHAUX AND NUTTALL'S
NORTH AMERICAN SYLVA.

PHILADELPHIA:

ROBERT P. SMITH, 15 MINOR STREET,

R. BALDWIN, PATERNOSTER ROW, LONDON; H. BOSSANGE, NO. 11
QUAI VOLTAIRE, PARIS; PERTHES, BESSER & MAUKE,
NO. 22 JUNGFERNSTIEG, HAMBURG.

1853.

QK481

244039

119

1953

fol.

v. 1

KITE & WALTON, PRINTERS,
No. 3, Ranstead Place.
T. R. JONES, Colorist.

WI

TO THE LATE

WILLIAM MACLURE, Esq.

President of the Academy of Natural Sciences in Philadelphia, &c. &c.

AS A MEMENTO OF HIS ATTACHMENT TO, AND LIBERAL ENCOURAGEMENT
OF, NATURAL SCIENCES IN NORTH AMERICA.

ALSO TO

F. ANDREW MICHAUX,

*Member of the American Philosophical Society, Correspondent of the Institute
of France, &c. &c.*

WHOSE NAME IS IDENTIFIED WITH THE HISTORY AND IMPORTANCE OF
THE PRODUCTIONS OF THE NORTH AMERICAN FOREST,

THIS WORK IS MOST RESPECTFULLY DEDICATED

BY

THE AUTHOR OF THIS SUPPLEMENT.

s
v
b
cr
S
w
of
in
ca
pe
on
Be
tat
pro
da
ext
tre
hav
ma
Bal
for
thro
cre
Her

P R E F A C E .

THE FOREST TREES OF AMERICA, being a subject of such great extent and importance, I felt, consequently, very diffident of undertaking their study, after what has been already done so well by my predecessor M. MICHAUX. Yet, in offering a new edition of the AMERICAN SYLVA in English, it appeared requisite, in keeping pace with the progress of discovery, that all the forest trees of the extended dominion of the United States, should, in some way or other, be included in the present publication; and, I confess, the magnitude of the task appeared, at first, sufficiently appalling, when we reflect on the vast territory now claimed by the United States. Beginning with the arctic limits of all arborescent vegetation, in the wilds of Canada, which we cannot, with propriety, exclude, forming as it does the boreal boundary of the North American forest; we then follow the extended shores of the Atlantic, until, towards the extremity of East Florida, and its Keys or Islands, we have attained the very confines of the tropical circle and make a near approach to the island of Cuba and the Bahamas: turning westward, we pass over the wide forests of the Mississippi, pursue the western streams, through vast woodless plains, until we attain the long crests of the "Rocky Mountains" or Northern Andes. Here, in these alpine regions, we meet with a total

change in the features of the forest; resiniferous evergreens, of the family of the Pines, now predominate, and attain the most gigantic dimensions. All the species, and they are numerous, have peculiar traits, and form so many curious and distinct species, of which little is yet known more than their botanical designation. Other remarkable forest trees, also imperfectly known, inhabit this great range of mountains, which continues uninterruptedly into the interior of Mexico in its southern course; while on the north, following the sources of the Missouri and the Oregon, and after thus dividing the waters which flow into the Atlantic and Pacific, it is, at length, merged in the "Shining Mountain" which send off their distant tributaries to the Arctic ocean.

The plains of the Upper Platte, those of the Oregon and of Northern California, a region bereft of summer rains, forming extensive barren steppes, like those of Siberia, present no forests, scarcely an alluvial belt along the larger streams of sufficient magnitude to afford even fuel for the camp fire of the wandering hunter or the erratic savage. The scanty drift wood, borne down from the mountains, the low bitter bushes of the arid plain, even the dry ordure of the bison, is collected for fuel, and barely suffices to prepare a hasty meal for the passing traveller; who, urged by hunger and thirst, hurries over the desert, a region doomed to desolation, and, amidst privations the most appalling, lives in the hope of again seeing forests and green fields in lieu of arid plains and bitter weeds, which tantalized our famished animals with the fallacious appearance of food, like the cast-away mariner raging with thirst, though surrounded with water as fatal to the longing appetite as poison.

Towards the shores of the Pacific, and on the banks of the Oregon, we again meet with the agreeable features of the forest.

"Majestic woods, of every vigorous green,
 Stage above stage, high waving o'er the hills;
 Or to the far horizon wide diffus'd,
 A boundless deep immensity of shade."

Transported in idea to the borders of the Hudson or the Delaware, we recline beneath the shade of venerable Oaks and spreading Maples; we see, as it were, fringing the streams, the familiar Cotton-wood and spreading Willows. On the higher plains, and ascending the hills and mountains to their summits, we see a dark forest of lofty pines; we hear the light breeze sigh and murmur through their branches as it did to the poets of old. But the botanist, in all this array, fails to recognise one solitary acquaintance of his former scenes: he is emphatically in a strange land; a new creation, even of forest trees, is spread around him, and the tall Andes and wide deserts rise as a barrier betwixt him and his distant home.

My indulgent reader will then excuse me, if I, on this occasion, appear before him only as a botanist; culling those objects which have given him so much delight, he wishes to present them to the curious public, alive to the beauties and symmetry of Nature's works. Whatever is yet known of their uses and history, is also given; and that the task might be more complete, we have rambled a little beyond, rather than fallen short of, the exact limits of the republic. We have thus added, as our friends TORREY and GRAY have done, or intend to do, in their general Flora, a collection of the trees of Upper California, extending our ramble as far as the vicinity of Sta. Barbara, in about the 34th degree of north latitude. We here met with several Oaks, Pines, a Plane-tree, a Horse-chestnut, and a Box Elder, which have not yet been found within the limits of the territory of Oregon.

While the work was in progress, Professor TORREY informed me of the arrival of a large collection of dried plants from Key West, in East Florida, made by Doctor BLODGETT, of the United States army. All the trees in this herbarium, at least forty species, were in the most generous manner given up to me for publication by the Professor. Most of them form distinguishing features in the tropical landscape of the West India Islands. Among them were the Mahogany, *Simaruba*, the Guaiacum or Lignum Vitæ, the poisonous Manchineel, several trees of the family of the Myrtles, (*Eugenia*,) three or four species of Fig trees, the Calabash, and Papaw or Melon-tree, the Mangrove, two species of Cordia, the West India Birch (*Bursera gummifera*), and many other arborescent plants which are now for the first time added to the Flora of the United States, and thus in a measure resolving the problem of the geographical limits of the Caribbæan Flora. The island of Key West lies about eighty-five miles from East Florida, and is the same distance from Cuba. It is about nine miles long and three broad, containing a population of about 400 people, chiefly engaged as wreckers.

Besides the trees we have noticed, I have been recently informed of the existence of thickets of *Cactuses* on the island, one of which with an erect, cylindric and divided stem, attains the height of 30 or more feet.

In the islands of the Everglades, considerably inland in East Florida, we have been informed that a Palm about 90 feet high, forming a magnificent tree, has been seen, but of this plant we have been unable to obtain, as yet, any further account.

The haste with which I have been obliged to proceed with the publication, has prevented me from receiving much advantage from correspondents. Such as have honoured me with their remarks are mentioned under

the appropriate articles as they occur in the work; and I take this opportunity of tendering them my sincere thanks for all such assistance.

As fast as new materials may be discovered, we intend to give them to the world in the form of a supplement, and we shall then also have an additional opportunity for correcting any errors which may have occurred either in regard to information or in the progress of printing, as well as of making such additions as a more thorough examination of the subject may suggest, particularly the characters of the different kinds of wood indigenous to the most extended limits of the republic.

Thirty-four years ago, I left England to explore the natural history of the United States. In the ship *Halcyon* I arrived at the shores of the New World; and after a boisterous and dangerous passage, our dismayed vessel entered the Capes of the Delaware in the month of April. The beautiful robing of forest scenery, now bursting into vernal life, was exchanged for the monotony of the dreary ocean, and the sad sickness of the sea. As we sailed up the Delaware my eyes were rivetted on the landscape with intense admiration. All was new!—and life, like that season, was then full of hope and enthusiasm. The forests, apparently unbroken, in their primeval solitude and repose, spread themselves on either hand as we passed placidly along. The extending vista of dark pines gave an air of deep sadness to the wilderness.

“—— these lonely regions, where, retired
From little scenes of art, great Nature dwells
In awful solitude, and nought is seen
But the wild herds that own no master's stall.”

The deer brought to bay, or plunging into the flood

*

from the pursuit of the Indian, armed with bow and arrow, alone seemed wanting to realize the savage landscape as it appeared to the first settlers of the country.

Scenes like these have little attraction for ordinary life, but to the naturalist it is far otherwise; privations to him are cheaply purchased, if he may but roam over the wild domain of primeval nature, and behold

“Another *Flora* there, of bolder hues,
And richer sweets, beyond our garden’s pride.”

How often have I realized the poet’s buoyant hopes amidst these solitary rambles through interminable forests. For thousands of miles my chief converse has been in the wilderness with the spontaneous productions of Nature; and the study of these objects and their contemplation has been to me a source of constant delight.

This fervid curiosity led me to the banks of the Ohio, through the dark forests and brakes of the Mississippi, to the distant lakes of the northern frontier; through the wilds of Florida; far up the Red River and the Missouri, and through the territory of Arkansas; at last over the

“Vast savannahs, where the wandering eye,
Unfixt, is in a verdant ocean lost.”

And now across the arid plains of the far west, beyond the steppes of the Rocky Mountains, down the Oregon to the extended shores of the Pacific, across the distant ocean to that famous group of islands* where Cook at length fell a sacrifice to his temerity. And here for the first time, I beheld the beauties of a tropical vegetation; a season that knows no change; but that of perpetual spring and summer: an elysian land, where Nature offers spontaneous food to man. The region of

* Sandwich islands.

the Bread fruit; the Tarrow (*Colocasia esculenta*) which feeds the indigent mass of the population; the Broussonetia, a kind of Mulberry tree, whose inner rind, called *tapa*, affords an universal clothing. The low groves produce the Banana, the Ginger, the Turmeric, the inebriating *Kava*, (*Piper methysticum*.) a kind of Arrow root, resembling the potato, (*Tacca*.) and the Saccharine Tee root, (*Dracæna terminalis*.) at the same time the best of portable fodder. The common timber for constructing houses, boats, various implements, and the best of fuel, is here the produce of a Mimosa, (*Acacia heterophylla*.) For lights and oil, the *too too* kernels (*Alcurites triloba*) produce an excellent and inexhaustible supply; the cocoa-nut and the fragrant *Pandanus* afford delicious food, cordage and mats, and the very reeds, reduced in size, which border the rivulets, are no other than the precious sugar-cane of commerce.

Leaving this favoured region of perpetual mildness, I now arrived on the shores of California, at Monterey. The early spring (March) had already spread out its varied carpet of flowers; all of them had to me the charm of novelty, and many were adorned with the most brilliant and varied hues. The forest trees were new to my view. A magpie, almost like that of Europe, (but with a yellow bill,) chattered from the branches of an Oak, with leaves like those of the Holly, (*Quercus agrifolia*.) A thorny Gooseberry, forming a small tree, appeared clad with pendulous flowers as brilliant as those of a Fuchsia. A new Plane tree spread its wide arms over the dried up rivulets. A Ceanothus, attaining the magnitude of a small tree, loaded with sky-blue withered flowers, lay on the rude wood-pile, consigned to the menial office of affording fuel. Already the cheerful mocking-bird sent forth his varied melody, with rapture imitating the novel notes of his neighbouring songsters.

The scenery was mountainous and varied, one vast wilderness, neglected and uncultivated; the very cattle appeared as wild as the bison of the prairies, and the prowling wolves (*Coyotes*) well fed, were as tame as dogs, and every night yelled familiarly through the village. In this region the Olive and the Vine thrived with luxuriance and teemed with fruit; the Prickly Pears (*Cactus*) became small trees, and the rare blooming Aloe (*Agave americana*) appeared consigned without care to the hedge row of the garden.

After a perilous passage around Cape Horn, the dreary extremity of South America, amidst mountains of ice which opposed our progress in unusual array, we arrived again at the shores of the Atlantic. Once more I hailed those delightful scenes of nature with which I had been so long associated. I rambled again through the shade of the Atlantic forests, or culled some rare productions of Flora in their native wilds. But the 'oft told tale' approaches to its close, and I must now bid a long adieu to the 'new world,' its sylvan scenes, its mountains, wilds and plains, and henceforth, in the evening of my career, I return, almost an exile, to the land of my nativity!

l, one vast
very cattle
ies, and the
as tame as
through the
Vine throve
the Prickly
rare bloom-
ned without

Horn, the
mountains of
l array, we
Once more
with which I
ain through
some rare
But the 'oft
ust now bid
scenes, its
rth, in the
exile, to the



Quercus Garryana.

Western Oak.

Chêne occidental



Mo
r
v
FE
s
s
C
N
al

T
alter
appe
allie

QUE
va
tis
bi
sq

QUE

IN
of th

O A K S .

Natural Order, CUPULIFERÆ. *Linnean Classification*,
MONECIA, POLYANDRIA.

QUERCUS. (TOURNEFORT.)

MONOICOUS. MALE flowers in loose catkins or racemes. *Calyx* monophyllous more or less deeply 5-cleft. *Stamens*, 5 to 10 with short filaments, the anthers oval and 2-celled.

FEMALE flower solitary, with a cup-shaped undivided hemispherical involucre formed of agglutinated imbricate scales, sometimes free at the summit. *Perianth* minute, superior. *Ovary* terminated by 2 to 3 stigmas, 3-celled, with 2 ovules. *Nut or gland* ovate-cylindric, coriaceous and smooth, 1-celled; albumen none, germ erect, with thick and fleshy cotyledones.

Trees or shrubs, principally of temperate regions. Leaves alternate, stipulate, simple. Flowers green and inconspicuous, appearing before the complete expansion of the leaves. Nearly allied to the Chestnuts, (*Castanea*.)

WESTERN OAK.

QUERCUS GARRYANA, (Dougl. Mss.), *foliis petiolatis, obovatis, utrinque obtusis sinuatis subtus pubescentibus, adultis subglabris, lobis obtusis subæqualibus superioribus subbilobis, fructibus sessilibus, cupula subhemisphærica dense squamosa, squamis acuminatis pubescentibus, glande ovata.*

QUERCUS *Garryana*, HOOKER, Flor. Bor. Amer. vol. 2. p. 159.

IN our western tour across the continent, no feature of the landscape appears more remarkable, after passing

the Mississippi to the wide alluvial borders of the Platte, than the almost total absence of our most characteristic forest trees, the Oaks. When at length we approached the Rocky Mountains or Northern Andes, we looked in vain for any species of this important genus, and as far as the eye could trace we commonly saw nothing but a dark unbroken mass of gigantic Firs and Pines. It was not till we had nearly reached the shores of the Pacific, that we again beheld any of the familiar features of the Atlantic forest. At the confluence of the Columbia and the Wahlamet we pitched our tents and moored our vessel, which had passed Cape Horn, beneath the spreading shade of majestic Oaks. With the first appearance of extended alluvial plains, immediately below the singular falls of the Oregon, called the Dalles, or Dykes, we observed, for the first time, this western oak loaded with its fruit.

The strong resemblance of the leaf of this species to that of the Post Oak (*Quercus stellata*) is almost a libel upon our gigantic plant, which may well rank among the largest of its species. It attains the height of 90 or 100 feet, if not more, with a diameter of 3 to 6; indeed, amidst a forest the most remarkable in the world for its extreme elevation, our Oak still bore a strict comparison with the rest. Its character, in nearly all respects, equals the famous Oak of northern Europe, (*Q. pedunculata*); its lofty summit and enormous branches spread out far and wide, affording the most perfect shade; and, as a picturesque tree, it is much the most striking in the western landscape. As an object of economy we found it of the last importance, useful timber trees being scarce along the Oregon; indeed, no other Oak exists along the coast of the Pacific but the present till we arrive at the absolute boundary of California. How far it extends to the north, I am unable to

say, but probably as far as Nootka Sound. In Upper California it is scarcely found beyond Monterey; its limit is probably somewhere between the 38th and 50th degree.

The wood is remarkably white for an Oak, hard and fine grained, and well suited for almost every kind of construction for which the White Oak or English Oak is employed. It was used by our trading party as barrel staves, and was found no way inferior to White Oak. Logs of it brought a good price at the Sandwich Islands, and, in short, there is scarcely any thing in which strength or durability are requisite, for which this timber is not suited. The acorns being sweet and agreeable, form an excellent mast for hogs, and even the aborigines of this region, who never cultivate the soil, employed them for food, first preparing them by stoving and afterwards laying them away under ground for future use.

The acorns are much larger than those of the Post Oak, as well as rounder. The leaf bears a considerable resemblance to that species, but is smaller, and, in fact, intermediate in form between it and the European species (*Q. pedunculata*.) It differs from both, in the whiteness of its wood. The bark is whitish and scaly, almost similar to that of the White Oak. The leaves from the first are not pubescent above, or only slightly so along the midrib, the hairs more numerous beneath, are, as in many other species, collected into stellated clusters; the young leaves of the Post Oak, previous to expansion, appear brownish-yellow, and like a mass of velvet, with the copious pubescence by which they are clad; in ours this appearance never occurs, and the old leaves become nearly smooth; the lobes have narrow sinuous openings, which scarcely pass half way down through the leaf; the lobes are usually 4 on a side, and

possess no great inequality with each other; the upper pair mostly present a notch or small division on the lower side, but nothing analogous to the singular obtuse dilatation which that part of the leaf exhibits in the Post Oak. The acorns, besides being larger, are not striated, and the scales of the cup are acuminate, and the upper ones free.

PLATE I.

A young branch, with the leaves not fully expanded, with barren aments.

; the upper
sion on the
gular obtuse
ibits in the
er, are not
minate, and

banded, with



Quercus Agrifolia

Holly leaved Oak

Chêne à feuilles de Houx



Her de Hour

Q

Qu

An

ma

dis

situ

pre

and

spa

the

atta

ter

rou,

rede

coa

A

war

spe

little

wou

gree

almo

HOLLY-LEAVED OAK.

QUERCUS AGRIFOLIA; *foliis lato-ovatis subcordatis dentato-spinosis glabris, fructibus axillaribus sessilibus.* NEE in Annal. Scienc. Nat. vol. 3. p. 271. Annals of Botany, No. 4. p. 106.

QUERCUS AGRIFOLIA, *foliis perennantibus subrotundo-ovatis subcordatis utrinque glabris remote spinoso-dentatis, cupula hemisphærica; squamis adpressis obtusiusculis, glunde ovata acuta.*—PURSH, Flor. Bor. Am. 2. p. 657. WILLD. Sp. pl. 4. p. 431.

An Ilex folio agrifolii americana, forte agria, vel aquifolium glandifera. PLUKENET, tab. 196. fig. 3?

THIS species, almost the only one which attains the magnitude of a tree in Upper California, is abundantly dispersed over the plain on which Sta. Barbara is situated, and, being evergreen, forms a conspicuous and predominant feature in the vegetation of this remote and singular part of the western world. It appears more sparingly around Monterey, and scarcely extends on the north as far as the line of the Oregon territory. It attains the height of about 40 or 50 feet, with a diameter rarely exceeding 18 inches. The bark is nearly as rough as in the Red Oak; the wood hard, brittle, and reddish, is used only for the purposes of fuel or the coarse construction of a log cabin.

As an ornamental tree for the south of Europe or the warmer states of the Union, we may recommend this species. It forms a roundish summit, and spreads but little till it attains a considerable age; as a hedge it would form a very close shelter, and the leaves evergreen and nearly as prickly as a holly, would render it almost impervious to most animals.

The leaves vary from roundish ovate to elliptic, and are of a thick rigid consistence, the serratures quite sharp; the young shoots are covered more or less with stellate hairs, and, for some time, tufts of this kind of down remain on the under side of the midrib of the leaves, which are, however, at length perfectly smooth and of a dark green above, often tinged with brownish-yellow beneath. The stamiferous flowers are very abundant and rather conspicuous, the racemes the length of 3 or 4 inches, the flowers with a conspicuous calyx and 8 to 10 stamens. The female or fruit-bearing flowers, are usually in pairs in the axills or juncture of the leaf with the stem and sessile, or without stalks. The cup of the acorn is hemispherical, and furnished with loose brownish scales, the acorn much longer than the cup, is ovate and pointed.

We do not recollect to have seen this tree properly associated with any other, except occasionally the *Platanus racemosus*; their shade is also hostile to almost every kind of undergrowth.

By Persoon this species is said to have been found on the eastern coast of North America, while Pursh attributes it to the north-west coast, about Nootka Sound. It does not, however, extend even to the territory of Oregon, as far as my observations go. Nee says "I have only seen branches collected at Monterey and Nootka." The leaves of the young plants (if I am not mistaken) are perfectly smooth when first developed, of a thin consistence, with numerous slender sharp dentures; beneath they are of a brownish-yellow colour, and appear smooth and shining.

PLATE II.

A young branch with barren aments. *a.* A branch with acorns.

SMALL LEAVED OAK.

QUERCUS *DUMOSA; ramis gracilibus pubescentibus; foliis rotundato-ovalibus subsessilibus spinoso-dentatis glabriusculis, subtus villosis concoloribus.

I observed this species to form entangled thickets over the base of the hills which flank the village of Sta. Barbara, in Upper California. It attains the height of 4 to 6 feet, is of a very unsightly appearance, forming what we should call scrub-oak thickets, of considerable extent, over a barren and rocky soil, which denies sustenance to almost everything else; the branches divide into many irregular straggling and almost naked slender twigs, clothed with a whitish smooth bark. The leaves are evergreen, small, and wholly resemble those of the *Quercus coccifera*, but are somewhat pubescent above and softly so beneath; the young twigs are also hairy, with a persisting pubescence. Being unable to discover upon it at the season I visited that country (in the month of April) either flowers or fruit, I am unable to give a figure of it that would be at all interesting.

with acorns.

ROCKY-MOUNTAIN OAK.

QUERCUS UNULATA; *fruticosa ramosissima; foliis perennantibus brevipetiolatis oblongis acutis sinuato-dentatis dentibus acutis, basi cuneatis, subtus pulverulento-tomentosis, supra nitidis; fructibus subsolitariis sessilibus, cupula hemisphaerica squamis appressis, glande ovata acuta.* TORREY in the Annals of the Lyceum of New York, vol. 2, p. 248.

THIS dwarf Oak, considerably allied to our small leaved preceding species, was discovered by Dr. James, in Long's Expedition, towards the sources of the Canadian, a branch of the Arkansa, and likewise in the Rocky Mountains. It is said to be a small straggling shrub, with the under surface of the leaves clothed with a close whitish tomentum or down, more or less spread, though more thinly, also on the upper surface, with the hairs stellated. The leaves are small, and somewhat resemble those of the Holly, about an inch and a half or two inches long, rather narrowed at the base, of a thick and rigid consistence, as in all the sempervirent Oaks, reticulately veined beneath, with the margin sinuately toothed, but not that I can perceive waved, as the specific name implies; the teeth sharp and acute at the points; above somewhat shining and minutely pubescent. The acorns are large and strongly resemble those of the Live Oak; they are, however, without stalks, and grow alone or in pairs; the cup is deep and hemispherical, with the scales pointed.

It is so nearly allied to the Holly Oak of the South of Europe (*Quercus Ilex*), that it is necessary to distinguish them. In our plant the base of the leaf is wedge-formed;

foliis peren-
uato-dentatis
mento-tomento-
vilibus, cupula
acuta. Tor-
ork, vol. 2, p.

o our small
Dr. James, in
the Canadian,
the Rocky
gling shrub,
with a close
read, though
th the hairs
at resemble
half or two
a thick and
Oaks, reti-
n sinuately
as the spe-
cute at the
y pubescent.
those of the
, and grow
mispherical,

he South of
distinguish
dge-formed;



Quercus Undulata.

Rocky Mountain Oak

Chino undulata

in
to
ac
and

in the *Ilex* it is usually rounded, the border less deeply toothed, and not in the least sinuated. The cup and acorn are wholly similar, but in our plant a little larger and less pointed.

PLATE III.

A branch of the natural size, with the acorn.

DOUGLAS' OAK.

QUERCUS DOUGLASSI, *foliis membranaceis oblongo-ovalibus basi acutis petiolatis sinuato-pinnatifidis siccitate haud nigrescentibus, supra glabris, subtus puberulis, lobis brevibus acutiusculis, petioliis ramulisque junioribus dense fulvo-pubescentibus; fructibus sessilibus solitariis binisve, cupula hemisphaerica dense squamosa squamis ovalis convexis in appendicem submembranaceam fulvum appressam linearem obtusam productis pubescentibus; glande ovata cupulam triplo superante obtusa cum umbone conico.*—HOOK. Ic. ined. HOOK. and ARNOT, Botany of Beechy's Voyage, p. 391.

THIS curious species, of which we have seen only a dried specimen, was collected in Upper California, and bears some affinity to the *Q. Garryana*. According to Hooker and Arnot, the leaves and whole appearance of the plant closely resemble *Q. sessiliflora*, but with different scales to the cup of the acorn. The leaves appear to be smaller, narrower, and less deeply divided than in *Q. Garryana*. The young leaves are covered with down on both sides, and the lobes tipped with short, soft, acute points.

To us, the branch which we have seen bears some resemblance, though vague, to the Post Oak, (*Q. stellata*.) The cup and acorn is also somewhat similar, but larger, while the leaf is smaller and scarcely dilated above. The under surface is covered with the same stellated pubescence. The stigmas of the fertile flowers are from 3 to 5 in number.

PLATE IV.

A branch of the natural size, with acorns. *a.* The male catkin and young leaf. *b.* The stamiferous flower magnified.

longo-ovalibus
eccitate haud
s, lobis brevi-
oribus dense
ariis binisve,
is ovatis con-
m appressam
glande ovatu
me conico.—
of Beechy's

seen only a
ifornia, and
ccording to
pearance of
with differ-
aves appear
ded than in
vered with
with short,

bears some
ak, (*Q. stel-*
similar, but
eely dilated
n the same
tile flowers

he male cat-
agnified.

1850



Dougl.




Quercus Douglasii

Douglas's Oak

Chêne de Douglas







Quercus Densiflora.

Dense flowered Oak.

Chêne à fleurs denses.

§ *C
la
lo
gu
ou

Tr
India
nerve
cong
rathe
cerne
Wall

QUE
lat
par
bus
mu
bus
fen
mi
lux
Pl.

Tr
gene
India
the a
alone
green
comm
entir

§ *CASTANOPSIS. *Aments elongated and persistent, perianth lanuginous, divided to the base; scales of the spreading cup loose and squarrose; stamens exerted; nut somewhat angular and downy; stigmas several, filiform and deciduous.*

Trees of Oregon, California, and the Himmalay Mountains in India, with the aspect of the Chestnut. Leaves entire, pennately nerved, sempervirent. Aments elongated, erect, the flowers conglomerated. Fertile flowers . . . ? To this section, or rather genus, belong also, as far as the male specimens are concerned, the *Quercus glomerata* and *Quercus spicata* of Dr. Wallich.

DENSE-FLOWERED OAK.

QUERCUS DENSIFLORA; *foliis perennantibus coriaceis petiolatis oblongo-lanceolatis basi obtusis breviter acuminatis parulle nervosis integerrimis margine revolutis junioribus fulvo-furfuraceo-tomentosis subtus pallidioribus demum glabris, amentis musculis elongatis folia superantibus densifloris valde tomentosus nunc ad basin flores paucos femineos gerentibus, fructibus sessilibus, cupula brevi hemisphærica dense squamosa, squamibus elongato-linearibus laxis sericeis, glande ovato-globosa sericea.*—Hook. Icon. Pl. ined. Hook. and ARNOT, Bot. Beechy, p. 391.

THIS remarkable tree, scarcely a true Oak, but congeneric with species in the Himmalaya Mountains, in India, is a native of Upper California. It has so much the appearance of a Chestnut, that the cup of the fruit alone attests what it really is. The leaves are evergreen and of the same lanceolate outline with the common Chestnut, having similar pennate nerves, but entire, or nearly so, on the margin; at first they are

softly clothed beneath with dense stellate brownish hairs, but at length become smooth; they are about 4 inches long and 1 to $1\frac{1}{4}$ wide. The catkins are erect, about 4 inches long, presenting the appearance of cylindrical, woolly spikes, beset with numerous exerted stamens with long slender filaments, as in the Chestnut. The cup is shallow and patulous, within and without softly sericeous, the scales numerous and acuminate, very loose, somewhat spreading and $2\frac{1}{2}$ to 3 lines long. The acorn is large, evidently angular, and more convex on one side, covered with whitish down, and terminated with several filiform, lanuginous and deciduous stigmas.

The *Castanea chrysophylla* of Douglas, if not the same plant, appears to be another species of this section or genus.

PLATE V.

A branch of the natural size. a. The acorn.

wish
out 4
erect,
of cy-
erted
stnut.
thout
nate,
long.
nvex
ated
mas.
ame
n or



Quercus leana.

Leas Oak

Chêne de leas



QU
la
an
an
cu
gt

O
spok
in co
sider
anal
the m
a dis
am b
taned
libert
disco
also
Lea's
speci

"T
phyllo
the s
are o
corda
from
The
pedun
of Q
fruit
The

LEA'S OAK.

QUERCUS LEANA, *foliis membranaceis, longissime petiolatis, oblongo-ovalibus, basi rotundatis, subcordatis, sinuato-pinnatifidis, demum glabris, lobis latis integris setaceo-acuminatis; fructibus brevi-pedicellatis; solitariis binisve, cupula hemispherica, squamis ovalis obtusis, glande subglobosa vittata subsemi-immensa, cum umbone brevi conico.*

OF this remarkably ambiguous Oak I have already spoken, in a note on *Q. heterophylla*, having at that time, in concert with Mr. Thomas G. Lea, its discoverer, considered it as a variety of that rare species or some analogous hybrid. Other specimens, accompanied with the ripe glands, have now convinced me that it is either a distinct species or another strange hybrid; but as I am by no means satisfied of the existence of such spontaneous mixed races among our Oaks, I have taken the liberty of giving it as a species, and dedicating it to its discoverer, an ardent and successful botanist. I shall also take the liberty of adding a quotation from Mr. Lea's notes, made on this plant, and sent to me with the specimens.

"The fruit resembles Michaux's figure of *Q. heterophylla*, but differs in being more depressed and obtuse at the summit. The cups, I think, are alike. The leaves are on longer petioles, but accord in being inclined to be cordate at base. If it is a hybrid, it may have come from the *Q. imbricaria* or *Q. tinctoria* or *Q. coccinea*. The fruit is too widely different from *Q. rubra*. The peduncles are about the same length as in my specimens of *Q. imbricaria*; in Michaux's figure of that species the fruit is represented as *sessile*, which I think is wrong. The petioles are much longer than in *Q. imbricaria*, the

leaves larger and more obtuse at base. These modifications (if it is a hybrid) may be derived from the long petioles and larger leaves of the *Black* and *Scarlet Oaks*. I think it does not partake of *Quercus phellos*, (Willow Oak,) a species that does not grow, to my knowledge, within several hundred miles of this place (Cincinnati)."

"I saw two individuals of *Q. phellos* in the Bartram garden, which Colonel Carr assured me were propagated from the seed of the original Bartram Oak. Certainly our plant is very like Michaux's figure; but as that appears to be a hybrid of *Q. phellos*, I think they must be considered distinct. If ours be a hybrid, it most likely comes from *Q. imbricaria* and *Q. tinctoria*, or *coccinea*."

"I have found but a single stock of this (about five years ago). It grows three miles north of Cincinnati."

I confess I see too little resemblance in our plant with *Q. imbricaria*, to agree with my friend Mr. T. Lea, as to any hybrid connection with that remotely allied species. Betwixt the Grey Oak, (*Q. ambigua*, Michx.) and *Q. tinctoria*, I perceive a nearer resemblance. The fruit appears to be wholly that of the Grey Oak. The gland in both is striated, and with a small conic projection. In our plant, however, the base of the gland and that of the cup are yellow, indicating its alliance to *Q. tinctoria*. The leaf differs wholly from both in its simple undivided lobes, though the long petiole and rounded base is that of *tinctoria*. Scarce as this species yet appears to be, under the present circumstances, I am inclined to believe it of a distinct race, with features as distinct as any species in the genus. For the Grey Oak being, I believe, unknown in Ohio, is again out of the question. I suspect it is in all physical respects allied to *tinctoria*, and would equally afford a yellow dying material.

The full grown leaves are from 5 to 5½ inches long, by 3 to 3½ wide, smooth and shining above, with a small quantity of deciduous stellate pubescence beneath. The lobes are about a single pair on a side. The central lobe only sometimes again subdivided into three lesser lobes, all of them ending in bristles. The base is rounded, and often hollowed out, or somewhat sinuated. The buds are small and brown. The fertile flower often by threes, on a short, thick, common pedicel, the middle flower abortive. Male flowers . . . not seen. Cups rather deep, as in *Q. tinctoria*, with the scales ovate, obtuse, and closely imbricated. The acorn roundish, somewhat ovate, broadly striate, with a short roundish conic point or umbo about half way, or nearly so, immersed in the cup.

PLATE V.(bis.)

A branch of the natural size with fruit. *a.* The cup. *b.* The gland.

this
W
So
ven
the

ele
de

TH
An
de
ac
I
ce
la
s
g
r
C
i

OBSERVATIONS ON THE OAKS.

QUERCUS MARITIMA, (*the Maritime Oak.*) The fruit of this species, and sometimes the leaves approach to the Willow Oak, but this is a low shrubby plant of the Southern States, with sempervirent leaves, which are very often deeply and distinctly sinuated, rigid, with the lobes often obtuse and mucronate.

QUERCUS MYRTIFOLIA, (*Myrtle-leaved Oak.*) Of this elegant and curious species, we have yet no materials deserving of a figure.

QUERCUS STELLATA. Q. OBTUSILOBA, MICH. (*Post Oak.*) The variety which I mentioned in the Genera of North American plants, vol. 2, p. 215, under the name of *β. depressa*, rarely exceeds 3 feet in height, and bears acorns at the height of 12 to 18 inches from the ground. I first observed it on the hills of the Missouri, up to its confluence with the river Platte, and it is also almost the last species which we find to the westward. I have since met with apparently the same low variety on the gravelly poor hills of the island of Martha's Vineyard, near Massachusetts Bay; it is this scrubby growth of Oak which still affords shelter to the grouse on that island. In some parts of Massachusetts, (according to Dr. Emerson,) the usual large growth of this tree is occasionally met with. The species of Quercus which I called *Q. Michauxii*, is, I now believe, nothing more than a mere variety of *Q. prinus*.

SWAMP WHITE OAK, (*Quercus bicolor.*) Of this spe-

cies I first observed a curious variety, which I called β . *mollis* or *Soft Leaved Swamp Oak*, in the swampy elevated forests of the Hudson, near New York; it occurs likewise near Philadelphia and Boston. The leaves, I find, are of the same form as in *Q. bicolor*, but the under side is not white but partly ferruginous or green and softly pubescent, the quantity of this clothing, however, varies, and in large leaves it becomes very thin. It forms a somewhat pyramidal tree, 60 or 70 feet high, branched nearly from the base, the branches deflected and intricately ramified. The leaves are narrowed at the base, and abruptly dilated towards the summit, the dentures are few and sometimes almost wanting, the breadth is about two-thirds of the length; the fruit stalk or peduncle filiform, 2 or 3 inches long, bearing about 1 to 3 acorns on each. It may perhaps be *Quercus filiformis* of Muhlenberg's Catalogue, page 87.

MOSSY-CUP OAK, (*Quercus oliviformis*.) This rare Oak, (which Michaux found only above Albany and in Genessee,) or at least a variety of it with less attenuated cups, is met with in Orange county, New York, where it was observed by Dr. Horton, and it also grows near Vernon in Sussex county, New Jersey. It has much the aspect of the Water White Oak, (*Q. discolor*,) but the leaves are sinuated.

WHITE OAK, (*Quercus alba*.) According to Mr. Emerson, the roots of the White Oak make very beautiful furniture. In England five pounds sterling have been given for the roots of a White Oak. The pieces have been taken out, and, when sawed and planed, present a wood of extraordinary beauty. A cabinet and table, made from the forked branches of this Oak, now in the possession of Mr. C. Wistar, in Germantown, may well

vie with the finest woods known; it is of a clear pale yellow, inclining to olive, and feathered in the most beautiful manner; the polish is also equal to that of the finest mahogany.

BARTRAM'S OAK, (*Quercus heterophylla*, MICH. vol. 1. pl. 16.) This curious tree, which, in 1837, had attained the height of 50 feet and a circumference of 3 feet 9 inches, was inadvertently cut down, and with it the species, if such it was, appeared to be annihilated; but Thomas G. Lea, Esq., of Cincinnati, informs me "that several years ago he discovered an Oak between two and three miles north of that city, the leaves and fruit of which accord with Michaux's figure. The leaves are sometimes larger than those represented, but with the same outline, irregularly and coarsely toothed, or sublobed, and longish petioles, the margin is very rarely entire. The tree is about 25 feet high, and in a vigorous state of growth. Some scattering Oaks of other species are in its immediate neighbourhood. I think it is not a variety of *Q. imbricaria*, many trees of which I have examined, but never found them with leaves the least indented. The *Q. phellos*" [to which it might be allied] "does not grow in the vicinity of Cincinnati, nor, that I know of, in any part of Ohio; this tree, therefore, cannot be a variety of that species." Its nearest affinity appears to me to be to the *Quercus ambigua* of Michaux, jr., from which it is principally distinguished by the narrower and more simple divisions of its leaves.

The WILLOW OAK appears to be very nearly allied to the Cluster-Leaved Oak of New Spain, (*Quercus confertiflora*), figured and described by Humboldt and Bonpland, but in that, though otherwise so very similar, the leaves are hairy beneath, while ours are perfectly smooth.

The **WILLOW OAK** is found as far west as the banks of the Arkansa and several of its branches.

LIVE OAK, (*Quercus virgens.*) Trees near Magnolia, in West Florida, occur of 8 to 9 feet diameter; it consequently affords large timber. Great quantities of this wood are now brought from the coast of West Florida. According to Wm. Bartram, the Live Oaks on the St. John's in East Florida, are from 12 to 18 feet in circumference; the trunk there rises only from 12 to 20 feet, when it throws out 3 to 5 large limbs, which continue to grow in nearly a horizontal direction, each limb forming a gentle curve from its base to its extremity, (*Bartram's Travels*, p. 85;) and he adds, I have stepped above 50 paces, on a line, from the trunk of one of these trees to the extremity of the branches. The wood is almost incorruptible, even in the open air. The acorn is small, agreeable to the taste when roasted, and in this state they are eaten by the aborigines as we do Chestnuts.

Stately avenues are formed of the Live Oak in South Carolina and Georgia, which, robed in Long Moss, put on an air of sombre grandeur and wildness.

In addition to the geographical limits of the Oaks, I may add, that according to the observations of Mr. G. B. Emerson, the *Rock Chestnut Oak* (*Quercus montana*, WILLD.), occurs in many parts of Massachusetts; he has also found the *Yellow Oak* (*Q. castanea*, WILLD.), about Agamenticus Mountain in York, (Maine.) "It is also found at Saco, in Maine, 25 miles further north." The *Black Oak*, (*Q. tinctoria*), "is found in York county, Maine, *Q. palustris*, (Pin Oak,) is very rare in Massachusetts." Mr. Emerson also corroborates my own observations concerning the prevalence of the *Post Oak* on the Island of Martha's Vineyard, and adds, that "it

hardly exceeds 20 inches in diameter, and 30 feet in height," which is a circumstance I had overlooked, its prevailing character there being that of a shrub.

The OAKS, though a very extensive genus, are confined to the Northern hemisphere. Besides the numerous species which pervade the United States, 16 were discovered by Nee in Mexico and New Spain, one of which, the *Q. agrifolia* is found in Upper California; 21 species were added to the Flora of North America by Humboldt and Bonpland found also in New Spain; 4 species were discovered in Japan by Thunberg; 2 in China by Bunge; 1 in Cochinchina, and 1 in the island of Formosa; 2 very remarkable species with lanceolate entire leaves and very long spikes of flowers, like those of a Chestnut, were met with in Nepal by Wallich; 6 other species likewise exist in that portion of India; Europe, chiefly the southern part, Northern Africa and Armenia afford about 28 species and several varieties; Java, Sumatra and the Molucca Islands also produce 19 species. Thus it appears, of the whole number, (according to the enumeration of Willdenow and more recent discoveries,) the Old World contains 63 species, and North America, including New Spain, about 74. Of these the United States possess about 37, and New Spain the same number. To these I may also add an additional species from the island of Cuba, nearly allied to our Southern Grey Oak, (*Q. cinerea*;) this I propose to call after its discoverer, M. La Sagra,

QUERCUS SAGRÆANA, foliis perennantibus oblongo-ellipticis obovatisque integris s. sublobatis brevi petiolatis obtusis nitidis margine revolutis subtus tomentosissimis nervosis subalbiddis, fructibus binis pedicellis incrassatis, cupula hemisphærica, squamis appressis, nuce ovata.

This species apparently forms a tree. The leaves are broader than those of the Grey Oak, of a thick and rigid texture, and are strongly veined both above and beneath; they are about $2\frac{1}{2}$ inches long and about 1 inch wide.

In
every
first
almo
brow
brow
a bro
subje
green
faint
densit
from
are m
of thi
dollar
faint
forty
the bu
that p
densit
magni
the Un
sured
probab
60 fee
Oak v
foot.
98 cer
timber
been c
and si

ADDITIONAL OBSERVATIONS.

In density and hardness the Live Oak much exceeds every other species of the genus hitherto examined. At first glance, and aided by its great weight, it appears almost like *Lignum Vitæ*. The sap-wood is of a pale brownish-yellow, the perfect wood of a pale chestnut-brown, and the extremely fine saw-dust almost as bright a brown as that from Mahogany. Growing in a climate subject to small changes of temperature, and being ever-green, the woody circles of annual increment are very faint and obscurely marked, which adds to the common density of the fibres. These rings, on young trees, vary from 1 to 2 lines in width, but in the older wood they are much narrower. One of the most striking features of this wood, however, is the distinctness of the medullary rays which traverse in strong and pale lines the faint waves of the annual increments. For the first forty or fifty years, the Live Oak appears to increase in the bulk of its trunk, as fast as our White Oak; but after that period the growth is much more slow; still the density of its wood is so great, that, through a strong magnifier, the pores and vessels are barely visible. In the United States Navy Yard, in this place, I have measured a squared log of Live Oak, 32 feet long, which probably formed the trunk of a tree not less than 50 to 60 feet in height. The present value of moulded Live Oak varies from \$1.20 to \$1.30 and \$1.45 per cubic foot. Promiscuous unprepared logs sell from \$1.20 to 98 cents and \$1 the cubic foot. Some very choice timber sells as high as \$1.65. This valuable timber has been employed in the United States navy between fifty and sixty years.

Little is yet known respecting the southern limits of this species of Oak, though there can be little doubt that it continues along the borders of the Mexican Gulf to Yucatan. Dr. Burroughs informs me that it is said to be found growing on the banks of the Alvarado river, about seventy-five miles south of Vera Cruz. I am also informed of the existence of the Live Oak near Matagorda in Texas.

It is stated in a late Texian paper that an English company have recently landed on the Brassos, in the neighbourhood of Brazoria, for the purpose of getting out Live Oak. They are said to have contracted with the English government to deliver two millions of cubic feet. The country about Brazoria is loaded with enormous trees, some of them casting a shade of 150 feet in diameter. The Live Oak extends into Texas at least one hundred and fifty miles, according to the observations of Mr. Caspar Wistar, jr., of Germantown.

John Lenthall, Esq., United States Naval Constructor, has favoured me with the following remarks concerning the timber used in the United States Navy.

The frames and principal pieces are all of Live Oak, and the frames of several of our ships that were cut from the islands of Georgia and on the coast, thirty years since, are still in an excellent condition, though in some ships, in which the timber was cut inland, the result is not so favourable. The weight of a cubic foot varies from 73 to 78 pounds. This timber is peculiarly adapted to ship building, and is scarcely fit for anything else, being short and crooked, so that the timbers are rarely grain cut.

The White Oak, used almost exclusively for plank, is cut from the seaboard of the middle states, and is equal to the best English or foreign timber. The Red Oak is never used. The Oak from Canada, is that which has

gen
ver
the
tim
and
Oak
pou
lake
uppe
Bay
This
F
quan
East
built
them

Th
freed
yellow
and v
the a
tanni
a gr
colou
with
obtai
ened
the p
infusi
tanni
glass.
Ac
calls
obtain

generally been introduced into England, and from it a very erroneous opinion has been formed with regard to the Oak timber of the United States, for the northern timber is much inferior to that from the southern states, and is never used. A cubic foot of unseasoned White Oak is from 58 to 60 pounds, and seasoned, 47 to 49 pounds. White Oak timber is often brought from the lakes and used for keels and bottom planks; but for upper works that from the Delaware and Chesapeake Bay is preferred, being much stronger and more durable. This lake timber is principally to be found at Newport.

From the Delaware river and Chesapeake Bay large quantities of White Oak are likewise shipped for the Eastern States, of which the better class of ships are built. A great deal of Pine timber is also shipped from thence for the same purpose.

The Quercitron is the bark of the *Quercus tinctoria*, freed from the epidermis. Besides tannin, it contains a yellow colouring matter, which may be extracted by water, and which, on evaporation, yields a peculiar extract to the amount of 8 per cent. of the bark employed. The tannin belongs to that variety which precipitates iron of a green colour. This tannin is very injurious to the colour, because it is precipitated by the same reagents with the colour, and imparts to it a brownish tint. To obtain the colouring matter free from it, a bladder softened in water, and cut into small pieces, freed from all the parts which are soluble in water, is applied to the infusion of the Quercitron bark, which takes up the tannin; or it may be precipitated by a solution of isinglass.

According to Chevreul, the colouring matter which he calls Quercitrin, although not a simple substance, is obtained by cautiously concentrating an infusion of

Quercitron. A crystalline substance then precipitates, which, while yet in suspension in the liquid, imparts to it a pearly appearance. It exhibits a slight acid reaction by curcuma-paper. It is slightly soluble in ether, but more completely so in alcohol. Water dissolves it; and the solution becomes orange-yellow by the addition of alkali. The acetate of lead and of copper, as well as the protochloride of tin, precipitate it in yellow flakes. Sulphate of the peroxide of iron colours it at first olive-green, and then causes a precipitate. Sulphuric acid dissolves Quercitrin, and the greenish orange coloured solution becomes cloudy by the addition of water. By dry distillation it yields, among other products, a liquid which soon crystallizes, the crystals possessing all the properties of Quercitrin.

In the dyeing establishments the clear yellow colour is obtained by precipitating the tannin by means of a solution of glue or buttermilk; the colouring matter then remaining in the solution is mixed with the solution of alum and carbonate of potash, by which it is precipitated of a yellow colour in combination with the alumina. Protochloride of tin also produces with it a strong yellow precipitate.

QUERCUS RUBRA? The largest Red Oak in North America, says a correspondent of the Natchitoches Herald, can be seen on the plantation of W. Smith, Esq., eighteen miles from Natchitoches, on the road leading to Opelousas. This majestic Oak stands in the midst of a rich and heavy bottom, on the Bayou St. Barb. Two feet from the ground it measures *forty-four* feet in circumference, and at six feet, *thirty-two* feet. The trunk appears sound and healthy, and its height, *to the branches*, is from fifty to sixty feet.

From Dr. G. Engelmann, of St. Louis, I learn that the White Oak, (*Q. alba*), and the Rock Chestnut Oak, (*Q. montana*, WILLD.), grow in that vicinity, where there are two varieties of each with sessile and with pedunculated fruit, in this respect agreeing with the two varieties of the English Oak, (*Q. robur*), which have been considered as two species. He also informs me that the Chinquepin Oak, (*Q. prinoides*, WILLD.; *Q. prinus chinicapin*, MICH. Sylva, t. 11,) grows commonly in South-western Missouri. He also adds, that the Spanish Oak, (*Q. falcata*), he has only seen in the southern extremity of Missouri; and that the Water Oak, (*Q. aquatica*), grows no nearer to him than the banks of the Arkansa.

In regard to the western range of our forest trees, Doctor Engelmann informs me, by letter, that though the Chestnut (*Castanea americana*) does not grow in the immediate valley of the Mississippi, it still reappears again in south-western Missouri and the north-western portion of Arkansas, where is also found the Locust tree, (*Robinia pseudacacia*.)

The Sweet Gum tree, (*Liquidambar styraciflua*), he saw on the borders of the Wabash; it grows also in Southern Missouri and all through Arkansas to the province of Texas, but he has not seen it through the greater part of Missouri and Illinois.

The Black Gum tree, (*Nyssa multiflora*, Walt. N. *Sylvatica*, MICH. Sylva, t. 110,) according to Dr. Engelmann, is common in the southern parts of Missouri.

The Wahoo Elm, (*Ulmus alata*), Dr. Engelmann finds as far north in Missouri as the vicinity of Herculaneum.

Around Cape Girardeau, one hundred and fifty miles south of St. Louis, he also observes the Tulip tree, (*Liriodendron*.)

Becch trees, the Doctor informs me, he has not seen

west of the Wabash, except near Capo Girardeau; they grow associated with Pines in Western Louisiana, and I have seen them in the forests which border the Arkansa.

CHESTNUT TREE, (*Castanea americana*.) The wood of this tree is capable of receiving a fine polish, and well selected pieces present waves and feathered figures of considerable beauty and variety, the more striking as they are seen with great distinctness through a pale and light ground. Furniture of this kind may be seen at Mr. Crout's cabinet warehouse in Sixth street, in this city.

u;
a,
he

of
all
of
s
d
t
s

100



Castanea alnifolia.

Dwarf Chestnut

Chataignier à feuilles d'aulne



Na

Por
o
a
m
cl
in
it
er

T
Am
and

CAS
m
b
β. P
Cas
Cas
Fag

A

†
rive

C H E S T N U T .

Natural Order, AMENTACEÆ, (JUSS.) *Linnaean Classification*, MONOECIA, POLYANDRIA.

CASTANEA.† (TOURNEFORT.)

POLYGAMUS. The *male* ament elongated, composed of numerous interrupted clusters of flowers, with a 5 or 6 parted perianth. Stamens 10 to 20. *Female* flowers about 3 in an ovoid muricate valvular involucre. Perianth urceolate, 5 or 6-cleft, having rudiments of abortive stamens. The ovary incorporated with the perianth, the stigma pencillate, exerted, its divisions rigid and pungent. Nuts 1 to 3, included in the enlarging echinate, 4-cleft involucre.

These are trees or shrubs of temperate Europe and North America, with alternate, stipulate, mucronately serrated leaves, and very long axillary aments. Nuts farinaceous, edible.

DWARF CHESTNUT.

CASTANEA ALNIFOLIA, *depressa, foliis obovatis subacutis, mucronato-serratis subciliatis junioribus subtus pubescentibus, amentis filiformibus solitariis tomentosis.*

β. PUBESCENS, *foliis brevioribus, adultis subtus pubescens.*

Castanea alnifolia, NUTT. Gen. Am., vol. 2. p. 217.

Castanea nana, ELLIOTT, Sk., vol. 2. p. 615. (not of Muhl.)

Fagus pumila, var. *præcox*, Walter, Carolin., p. 233.

A SPECIES remarkable for its dwarf growth, and in-

† So named from *Castanea*, a town of Thessaly, near the river Peneus, where large Chestnut trees are still found.

sorted only to complete the history of the genus. It rarely exceeds a foot in height, growing in small patches, with creeping roots. I first met with the variety β in the vicinity of Charleston, South Carolina, afterwards the smoother kind, much more abundant, and in flower in the month of March, around Tallahassee, in West Florida.

The Floridian plant is scarcely a foot in height, with smooth purplish-grey branchlets; the leaves obovate, on very short petioles, deeply serrate, obtuse or acute, elliptic-obovate, when young whitish pubescent; the adult almost perfectly smooth on both surfaces; about 3 inches long by 1 inch or more wide. Stipules subulate, rather persistent. Male aments solitary, long, and filiform, tomentose. The fruit I have not seen.

The Charleston plant grows in sandy pine barrens, and the nut, which is solitary, is said by Elliott to be much larger, but less abundant than in the other native species. This plant rarely exceeds 2 feet in height. Its leaves are glossy above, pubescent but not tomentose beneath. Fertile flowers 1 to 3 in an involucre, only one perfected.

The wood of the Chinquepin, (*C. pumila*), whenever it can be obtained large enough for posts, is much valued, as it is supposed to be more durable when exposed to the weather than any other timber, except the Red Cedar. (*Elliott*.)

PLATE VI.

A branch of the natural size.

GOLDEN LEAVED CHESTNUT.

CASTANEA CHRYSOPHYLLA, (Dougl. Mss.,) *foliis sempervirentibus lato-lanceolatis acuminatis coriaceis integerrimis glabris subtus aureo-farinosis.* Hook. Flor. Bor. Am., vol. 2. p. 159.

According to Douglas, this is a splendid evergreen tree, varying in height from 20 to 70 feet, with leaves 4 to 5 inches long, deep green above, and below of a rich golden-yellow. These leaves are, also, (very different from all the rest of the genus,) quite entire. The spikes or catkins of flowers scarcely exceed an inch in length, including the peduncle, and they are solitary in the axills of the upper leaves. Sometimes all the flowers on a catkin are male; sometimes the 2 or 3 lower flowers are female. The fruits are 2 or 3, crowded, or densely covered with acicular prickles. Said to be common at the Grand Rapids of the Columbia, Cape Orford, and near Mount Hood; constantly affecting the hills. This species rests wholly on the authority of Douglas. I did not meet with it, nor does it appear that any specimens were sent to England. It will probably prove to be some very different genus to that of the present.

B I R C H .

Natural Order, BETULINEÆ, (Richard.) *Linnæan Classification*, MONŒCIA, POLYANDRIA.

BETULA.† (LINN.)

Male flowers in long cylindric aments. Scales in a double series, the inner by 3's, 1-flowered; stamens 6 to 12. *Female* flowers with ovoid or oval aments; the scales trifid, 1 to 3-flowered. *Styles* 2. *Nuts* minute, compressed, 1-seeded, edged with an alated thin margin.

Trees or shrubs of the colder parts of the northern hemisphere on both continents with the bark often exfoliating in thin circular plates. Leaves alternate, ovate, or deltoid, serrated; producing stipules; aments axillary.

WESTERN BIRCH.

BETULA OCCIDENTALIS; *ramis resinoso-verrucosis, foliis lato rhombico-ovatis sublobatis inciso-serratis hirsutulis, subtus ; allidioribus punctatis, nervis remotis, amentis femineis lato-cylindraceis squamis lobis lateralibus ovatis intermedio longiore.*

BETULA *occidentalis.* Hook. Flor. Bor. Amer., vol. 2. p. 155.

THIS low species of Birch, only 6 to 10 feet high, was first observed westward, near the sources of the Sweet-Water, a northern branch of the Platte, and where it penetrates into the first range of the Rocky Mountains. On the borders of this clear stream, diminished to a small purling brook, and accompanied by clumps of willows, we first saw it growing. According to Drum-

† Supposed to be derived from *Betu*, the Celtic name for the Birch.

sifi-

ble
ale
3-
ed,

ere
cir-
ro-

is
is,
ix-
is

5.

as

t-

it

s.

a

f

-

-

e



Western



Betula occidentalis.

Western Birch

Bouleau occidental

me
tal
sp
De
of
co
hea
vir
spr
bra
del
my
son
abo
ner
with
on,
cim
(Th
2 to
stan
scal
leng
at t
atec
line
bro
sum
T
dian
The

A

mond, it occurs on the east side of the Rocky Mountains down to Edmonton House. Douglas found it near springs on the west side of the Rocky Mountains; and Doctor Scouler met with it in Oregon, near to the straits of Juan de Fuca; it also grows near Walla-Walla, and continues up the Oregon to the country of the Flat-heads.

The principal branches are erect and somewhat virgate, clothed with a bright brown bark, copiously sprinkled with small resinous warts, so as to render the branches rough to the touch. The leaves are somewhat deltoid, or rhomboidly-ovate, on shortish petioles (in my specimens), acute, but not acuminate, sharply and somewhat unequally serrated, and very slightly lobed, above somewhat glutinous, with very few pinnated nerves, below paler; the midrib and nerves sprinkled with a few long hirsute hairs, which are also seen above, on, and near the petiole. The leaves, in flowering specimens, are only about $1\frac{1}{2}$ inches long by an inch wide. (The adult leaves described by Hooker, are much larger, 2 to $2\frac{1}{2}$ inches long.) The aments are cylindric, in the stamiferous plant, composed of a double series of scales. Female aments pedunculated, cylindric, at length drooping, often accompanied by a very small leaf at the base; the scales trifid and dilated, strongly ciliated, the lateral lobes ovate; the central one nearly linear and longer; 3 germs beneath each scale. Nuts broadly winged. Styles 2, very long and subulate; summit of the germ pubescent.

The trunk of this species is only a few inches in diameter, so that it scarcely ranks with proper trees. The leaves are bitter to the taste.

PLATE VII.

A branch of the natural size. *a.* The seed vessel.

OVAL-LEAVED BIRCH.

BETULA RHOMBIFOLIA, ramis resinosis gracilibus, foliis subrhomboideo-ovalibus, vix acutis, grosse serratis, subtus pallidioribus epunctatis; venis pilosiusculis amentis femineis cylindraccis, squamis tripartitis glabriusculis lobis ovatis lateralibus brevibus.

THIS is a still more humble shrub than the preceding, which it somewhat resembles. It grows in the central Rocky Mountain range, and continues more or less to the banks of the Oregon. It is spreading and somewhat decumbent, with slender brown twigs, which, when young, are more or less covered with resinous atoms. The leaves, with their petioles, which are 2 or 3 lines, are not more than an inch long by $\frac{1}{2}$ an inch wide, oval, and somewhat rhombic, deeply, sharply, and almost equally serrate, rounded, but still generally acute, smooth above, paler beneath, with a very few distant nerves, somewhat hairy along their margins beneath. External scales of the male aments ovate, and ciliate. Stamens about 6. Female aments with nearly smooth, deeply 3-parted scales, of which the central division is the longest. I have not seen the ripe fruit.

PLATE VIII.

A branch of the natural size. a. The seed vessel.

Obs. On the summit of the White Mountains of New Hampshire grows the *Betula nana* of Europe, found there by Mr. Oakes as well as myself.

Mr. Charles Pickering also collected a specimen on

liis
tus
mi-
bis

ng,
ral
to
at
en
as.
es,
e,
ad
e,
nt
h.
e.
l,
s

r
l



Oval leaf



Betula rhombifolia.

Oval leaved Birch

Bouquet à feuilles ovales

thos
cosa
In
now
is a
Stat

C
This
the
Dr. I

W
justly
to the
aspec
have
kins.
comp
middl
lobes,
nearly

those mountains, which appears to be the *Betula fruticosa* of Pallas, first found in Siberia.

In the Rocky Mountains, besides the two species now described, we met with the *Betula glandulosa*, which is also found on the high mountains of the United States.

CANOE BIRCH, or PAPER BIRCH, (*Betula papyracea*.) This very useful species of Birch to the aborigines of the north, is found, according to the observations of Dr. Richardson, as far as the 65th degree of latitude.

WHITE BIRCH, (*Betula populifolia*.) Hooker very justly remarks the near affinity which this species bears to the common European Birch, (*B. alba*.) The general aspect is the same. In our plant, however, the leaves have longer acuminate points, and smaller fertile catkins. The scale of the same catkin in ours, is also comparatively smaller and shorter clawed, with the middle lobe acute and much smaller than the lateral lobes, whereas, in the European Birch, the lobes are nearly all equal and obtuse.

A L D E R S .

Natural Order, AMENTACEÆ, (JUSSIEU.) *Linnæan Classification*, MONECIA, TETRANDRIA.

GENUS ALNUS. (TOURNEFORT, DECANDOLLE.)

CHARACTER. The flowers are *monœcious*, (or of two different kinds on the same plant,) disposed in catkins, (or cylindric spikes of short duration,) those producing the stamens are long and cylindric; those of the fruit or seed are ovoid or globular, produced upon branching peduncles. The scales of the *male* flower are pedicellated, and in the form of an inverted heart, bearing beneath each 3 lesser scales; the proper flowers are situated at the base of each of these, and are composed of a cup with 4 lobes and 4 stamens. The scales of the fruiting catkins are wedge-shaped, hard, and persistent. The ovary is compressed, and bears 2 long stigmas. The envelope of the seed is hard, with a border which is either thick or membranaceous, and presents 2 cells with 2 seeds; the ovules in the germ are about 4, or 2 in a cell, 3 of them usually abortive.

The plants of this small genus, confined to the temperate or colder parts of Europe and North America, are either shrubs or trees, with deciduous leaves, generally growing by streams, or in cool and humid places. As trees they seldom attain a greater elevation than 30 to 40 feet; the wood is hard and yellowish, becoming of a brownish-red, nearly like mahogany, when exposed to the air, and capable of acquiring a fine polish. When stained black it resembles ebony, and it is capable of enduring moisture for a great length of time.

The Alders may be divided into the two following sections; in both the peduncles are subdivided.

§ 1. *The seed-vessel furnished with a membranaceous winged margin, and with the scales of the fertile ament retuse or obscurely lobed.*

The White Alder, (*Alnus incana.*)

The Oregon Alder, (*Alnus Oregona.*)

The Heart-leaved Alder, (*Alnus cordata.*)

Mountain Alder, (*Alnus viridis.*)

§ II. *The margin of the seed-vessel thick and opaque, and with the scales of the fruiting ament distinctly lobed.*

Common Alder, (*Alnus glutinosa.*)

Fine-toothed Alder, (*Alnus serrulata.*)

Sea-side Alder, (*Alnus maritima.*)

Oblong-leaved Alder, (*Alnus oblongata.*)

Short-leaved Alder, (*Alnus brevifolia.*)

Rhombic-leaved Alder, (*Alnus rhombifolia.*)

§ I. *Fruit alated.*

OREGON ALDER.

ALNUS *OREGONA; *foliis lato-ovatis utrinque acutis, duplicato-serratis junioribus glutinosis, venis subtus pubescentibus pallidis; stipulis oblongis deciduis glutinosis, ramulis glabris.*


ALNUS glutinosa. PURSH, Flor. Bor. Am., vol. 2. p. 623, (in part.)

THIS tree, like the common Alder of Europe, attains the height of 30 or 40 feet, with an erect smooth trunk of small diameter, the wood of which is very similar to that of the European species, and might, no doubt, be employed for the same purposes; it is, however, lighter in colour, but of a close grain.

As an ornamental tree, it is well worth attention, producing an elegant erect top, and affording considerable shade by the largeness of its leaves, which are about 3 inches long by $2\frac{1}{2}$ wide. We found it, as usual with the plants of this genus, growing along the borders of small clear brooks, near the confluence of the Wahlamet, but seldom, if ever, on the banks of the larger streams which are subject to inundation. In our progress to the West, we first observed this tree on the borders of the rivers Boisée and Bruléc, which pass into the Shoshonee, not far from Walla-Walla, and at intervals it continues more or less common to Point Chinook, near the shores of the Pacific.

The twigs are smooth and of a brown colour, and the young buds of every kind resinous, as well as the upper surface of the younger leaves; beneath, the leaves are more or less pubescent, particularly along the veins,

i-
u-
t-
)
s
k
o
o
t
-
s
.





Alnus Oregona.

Oregon Alder

Aune de l'Oregon

and
are
teet
at t
the
are
bud
simi
T
nus
lets,
as i
It a
A. a
disc
are n

Be
wood
beau
have
hoga
wood
plove
clogs
its du
peras
been
on th

A B


and paler and often somewhat ferruginous. The veins are very strongly marked and prominent beneath, the teeth large, but the denticulations minute and glandular at the points. Sometimes the leaves are elliptic-ovate, the stalks about the third of an inch long. The stipules are resinous, and disappear with the evolution of the bud. The fruiting ament is roundish-ovoid, and very similar to that of the common Alder.



This species is nearly allied to the White Alder, (*Alnus incana*), but differs sufficiently in its buds, branchlets, stipules, and leaves; in both the fruit is provided, as in the Birch, with a translucent membranous wing. It appears, likewise, to have a considerable affinity to *A. acuminata* of Humboldt and Bonpland, a tree of Peru, discovered by Dombey, but in that species the leaves are more lanceolate than ovate and acuminate.

Besides the other economical uses for which the wood of the Alder is employed, the knots furnish a beautifully veined wood for cabinets; handsome chairs have been made of it, which acquire the colour of mahogany. In France it is used in making sabots, or wooden shoes, and in the north of England it is employed for the thick soles of a kind of shoes called clogs, and is preferred to these uses in consideration of its durability and lightness. The chips, boiled with copperas, give a black dye to wool; and the leaves have been used in tanning; sheep will browse on them and on the smaller branches.





PLATE IX.

A branch of the natural size. *a.* The seed-vessel.





**IMAGE EVALUATION
TEST TARGET (MT-3)**



**Photographic
Sciences
Corporation**

23 WEST MAIN STREET
WEBSTER, N.Y. 14580
(716) 872-4503

0
11
E 28
E 32
E 25
E 22
E 20
E 18
6

11
11
10
51

WHITE ALDER.

ALNUS INCANA; *foliis oblongis acutis subtus pubescentibus, axillis venarum nudis, stipulis lanceolatis.* WILLD. Sp. pl.

Alnus undulata, WILLD. Sp. pl., vol. 4. p. 336.

Black Alder, (*Alnus glauca*,) Mich. Sylva, vol. 1. p. 378.

Betula Alnus, crisper, MICH. Flor. Bor. Am., vol. 2. p. 181.

Betula crisper, ARTON. Kew., vol. 3. p. 339.

Betula alnus, β. LINN. Sp. pl.

THIS species forms a much smaller tree than the common Alder, being only 12 to 18 feet high, and sometimes indeed a mere shrub, as in the Alleghany Mountains in Pennsylvania. In Massachusetts and Maine it attains its greatest size. Its bark is grey or cinereous; the leaves are sometimes villous beneath, and the stipules persistent after the development of the leaves, which are no way glutinous; those of the young plants are smooth and glaucous beneath. It is common to the mountainous parts of Europe nor less than to the northern parts of the United States. It occurs likewise in this vicinity.

MOUNTAIN ALDER.

ALNUS VIRIDIS, (DECANDOLLE;) *foliis rotundato-ovatis irregulariter argute serratis glabriusculis, stipulis ovatis membranaceis deciduis; fructibus late alatis.*

Alnus viridis, DECANDOLLE, Flore Française, vol. 3. p. 304.

Betula viridis, VILLARS, Dauphin, vol. 4. p. 789.

Betula ovata, SCHRANK, Salisb. p. 25.

Betula incana, *β.* LAMARCK, Dict., vol. 1. p. 455.

Alnus Alpina minor, BAUHIN'S Pinax, p. 428.

LABRADOR, the elevated summits of the White Mountains of New Hampshire,† and the tops of the high mountains of North Carolina‡ are the only localities on this continent where the Mountain Alder has yet been found. It occurs, likewise, in the Alps of Switzerland, at an elevation of between 4 and 5000 feet above the level of the sea, where it frequently forms a small tree about 6 feet in height. In the White Mountains its stature is much more depressed, and it is of rare occurrence. It grows likewise in the barren and cold climate of Kamtschatka.

The wood is white, and the branches are covered with a cinereous smooth bark. The leaf is near 2 inches long and $1\frac{1}{2}$ wide, nearly smooth on both sides, but generally somewhat hairy along the veins beneath, rather acute, with numerous sharp small and irregular serratures, but not doubly serrate. The male catkins are long, and grow 2 or 3 together at the extremities of the twigs; each scale contains 3 tetrandrous flowers, as usual in the genus. The fertile aments are roundish and elliptic, about 3 together and terminal, the scales are truncated and obscurely lobed at the extremities; the fruit, like that of the Birch, is furnished with a broad, thin, conspicuous winged margin.

† A specimen in the herbarium of the Academy of Natural Sciences in Philadelphia, was discovered on the White Mountains by my friend Charles Pickering, at an elevation of 4000 feet above the level of the sea.

‡ Recently discovered by Dr. Gray and Mr. Curtis.

§ II. *Fruit not alated; the margin opaque.*

THIN LEAVED ALDER.

ALNUS TENUIFOLIA; *foliis lato-ovatis subacutis duplicato-crenatis glabris basi rotundatis longe petiolatis, stipulis deciduis, pedunculis femineis duplicato-ramosis.*

This very distinct species of Alder which arranges with our common species, (*A. serrulata*), was met with on the borders of small streams within the range of the Rocky Mountains, and afterwards in the vallies of the Blue Mountains of Oregon, a chain which may be called, as it were, in comparison of their elevation, the Alleghanies of the West.

This species falls short of the character of a tree, but yet it is scarcely inferior in size with our common species, growing to about the height of a man, with numerous short branches, covered with a smooth grey bark. The leaves are about 2 inches long by $1\frac{1}{2}$ wide, with slender petioles, from a half to $\frac{3}{4}$ of an inch in length; they are of a thin consistence, and usually smooth, with obtuse denticulations. The fruiting branches are often subdivided, each branch bearing from 3 to 5 small roundish ovate aments, of which the scales are very distinctly lobed. The fruit is unusually small and elliptic, terminated by the 2 remaining styles, and having a thin opaque margin.

PLATE X.

A branch of the natural size. *a.* The fruit.


to-
lis

es
th
ne
ne
d,
e-

nt
e-
e-
s.
h
;
h
n
l
y
-
a



Thu



Alnus tenuifolia.

Thin leaved Alder

Alune menue feuille

S
a
t
h
A

RHOMBIC LEAVED ALDER.

ALNUS RHOMBIFOLIA; *foliis subrhomboido-ovatis obtusiusculis glutinosis basi acutis, subduplicato-serrulatis serraturis crebris acutis, subtus puberulis axillis venarum nudis, stipulis oblongis membranaceis deciduis.*

I OBSERVED this species, a large shrub, in the vicinity of Monterey, in Upper California. Its nearest relation appears to be to the European Alder, (*A. glutinosa*), from which, however, it is abundantly distinct. The fruit I have not seen.

The leaves are about 2 inches long and $1\frac{1}{2}$ wide, glutinous, beneath nearly the same colour as above, and pubescent along the veins; the petioles are not more than about 2 lines long. The twigs are smooth and brownish. It appears to be allied to the oblong-leaved Alder (*A. oblongata*) of the South of Europe.

SEA SIDE ALDER.

ALNUS MARITIMA; *foliis ovalibus glabris serratis obtusis vel acuminatis, basi acutis, subtus ferrugineis; amentis femineis maximis, squamis duplicato-lobatis.*


ALNUS MARITIMA; *foliis ovatis serratis, basi acutis.* Muhl. Mss. *Observationes Botanicae de Plantis Am. Septent.* p. 193. (in the Library of the Acad. Nat. Sc. Philad.) and Herbarium.

A SPECIMEN of this very distinct species of Alder was collected on the Eastern shore of Maryland, by my friend Charles Pickering. It has the appearance of being a low shrub, with slender smooth branches. The leaves are $2\frac{1}{2}$ to 3 inches long by $1\frac{1}{2}$ or more wide, of an elegant well defined oval outline, and supported upon longish petioles; the young buds and leaves, after the manner of the genus, are slightly glutinous; the nervings very slender, serratures shallow, and in the larger leaves rather remote; the uppermost leaves on the infertile shoots are acuminate, and, at first glance, look almost like the leaves of a Camellia. The male catkins are unknown, as are the stipules, which are probably small. The fertile ament in size and general appearance might be taken for the strobile or cone of a Spruce; it is about the size of a Hop cluster, nearly black, with the scales very thick and deeply and obviously lobed. The carpel is small in proportion, and with a thick opaque and obscure margin, as in *A. serrulata*.

PLATE X.

A branch of the natural size. *a.* The seed vessel.

el
i-
.
.
s
r
f
e
.







Alnus maritima.

Sea-side Alder

Alnus maritima







Elmus opaca.

Opaque leaved Elm.

Orme opaque



N

UL
s
A

L
rior
Ree
110
of c
spee
the
oper
prov
for
ally
only
brea
scab
and
marg
leaf

E L M S .

Natural Order, ULMACEÆ. (Mirbel.) *Linnaean Classification*, PENTANDRIA, DIGYNIA.

ULMUS. (LINN.)

OPAQUE LEAVED ELM.

ULMUS *OPACA; *foliis parvis oblongo-ovatis obtusis scubris, subduplicato denticulatis, basi cuneatis obliquis subtus pubescentibus, floribus fusciculatis, fructibus hirsutis.*

In the summer of 1818, on my journey into the interior of the territory of Arkansa and on the plains of Red River, near its confluence with the Kiamesha, 1100 miles up the former stream I had the satisfaction of discovering this curious Elm, which, like our other species, forms a majestic and spreading forest tree of the dimensions of an ordinary Oak. In those dry and open savannahs, the shade of this densely verdant tree proved more than usually acceptable. It is remarkable for the smallness and thickness of its oblique and usually blunt leaves, which, with their short stalks, are only about an inch in length by half that dimension in breadth; they are also very numerous, close together, scabrous, with minute papillæ, of a deep green above, and somewhat shining, oblong-ovate, mostly obtuse, the margin with shallow double denticulations; beneath, the leaf is paler, a little brownish, with strong pennate,

simple, or forked nerves; the base of the leaf is oblique, as well as the whole outline, and one half of the leaf is much narrower than the other; the nerves are pubescent. The young branches are smooth and brownish. The leaves, before complete development, are canescently tomentose and attended by large oblong membranous brown stipules. The taste of the plant is astringent, but no way mucilaginous.

This remarkable species appears to be nearly allied to *Ulmus chinensis*, judging from the short description in Persoon and Duhamel. The flowers are fasciculated in small numbers and on short peduncles. The samara is elliptic, rather deeply bifid at the summit, covered with a dense and somewhat ferruginous pubescence even when ripe.

Of the uses and quality of the timber of this species, I am unable to speak from experience, as it grew remote from the settlements at that time established in the territory. The density of shade produced by it, so crowded with rigid leaves, and the peculiarity of its appearance, entitle it to a place in the nurseries of the curious, and it is probably quite hardy enough for all temperate climates. To this species Virgil's epithet—

“Fœcundæ frondibus ulmi”—

might more justly be applied than to any other.

PLATE XI.

A branch of the natural size.

c,
is
s-
l.
s-
t-
t-
d
n
n
s
n
n

,
-
n
o
s
e
l
.



Elmus racemosa.

Thomas's Elm

Orme à grappe



U

U

ce
w
th
ve
Y
E
sp
th

pr
Lo
on
ab
ce
in
to
of
be
tin
10
ve
th
of

w

THOMAS' ELM.

ULMUS RACEMOSA, (THOMAS;) *foliis ovatis acuminatis duplicato-serratis glabris subtus pubescentibus; floribus racemosis fasciculatis.*

ULMUS RACEMOSA, flowers in racemes; pedicels in distinct fascicles, united at their bases; leaves ovate, acuminate, doubly serrate, glabrous above, [minutely] pubescent beneath; stigmas recurved. EATON'S North Am. Bot. (ed. 8.) p. 464. THOMAS, in Silliman's Journ. Sci., vol. 19. p. 170, with a Plate.

THIS species, confounded with our other Elms, is, according to Professor Torrey, an abundant species in the western part of the State of New York, and probably of the Western States generally. Mr. Thomas, its discoverer, found it in Cayuga county, in the State of New York, and in the adjacent country. According to G. B. Emerson, Esq., Mr. Oakes, he believes, has obtained specimens from Vermont, collected by Dr. Robbins, so that it is probably a northern and western species.

The lower stout branches, according to Mr. Thomas, produce corky excrescences like the Wahoo Elm. Leaves broad-ovate acuminate, obliquely auriculated on one side, doubly serrate, smooth, and somewhat shining above, with the under surface and ribs minutely pubescent. The flowers, unlike any other Elm, are disposed in racemes, composed of several clusters of 2 to 4 together and extending to the length of 1 to 2½ inches, often furnished with one or two small but perfect leaves before the terminal buds are open; the flowers distinctly pedicellate. Calyx 7 to 8 cleft. Stamens 7 to 10. Stigmas 2, recurved. Samara elliptic, large and very pubescent, with the margin thickly fringed, and the membrane more extended on one side as indicative of a second but abortive cell.

PLATE XII.

A branch of the natural size. *a.* The flower. *b.* A branch with the corky bark.

HICKORY.

Natural Order JUGLANDÆE, (Decand.) *Linnean Classification*, MONŒCIA, POLYANDRIA.

CARYA† (NUTT. Gen. Am.)

Staminate flowers in very long and loose ternate aments, scales imbricated, 3-parted. Stamens 3 to 6, with pilose anthers. *Fertile* flower with a single 4-lobed superior herbaceous perianth. Style none; stigma partly discoid, 2-lobed, the segments bifid. Pericarp woody, 4-valved. Nut mostly somewhat quadrangular, with an even surface.

Large trees of North America, confined to the eastern side of the Rocky Mountains, and extending from Upper Canada to Florida. Leaves alternate, unequally pinnate, without stipules. Flowers polygamous, in compound pendulous pedunculated aments appearing with the leaves; female flowers terminal; the pericarp opening by 4 valves. Nuts edible or bitter, usually more or less quadrangular; in the *Pakan* even. Pubescence tufted or stellate. (Persoon, as far back as 1807, divided the genus *Juglans* into the two natural sections which it presented; his second division included the Hickories only. “ * * * *Amentis masculis compositis, tetrandris.*”)

† From *καρυα*, the ancient Greek name of the Walnut. *Hickory* is an Indian name for *some* of the species of this genus; one of them was known to the Indians by the name of *Pecan* or *Pakan*; Rafinesque applied the barbarous name *Hickoria* to this genus, without describing or limiting it; in so doing he has no higher claims for the adoption of the name than our woodsmen and the aborigines.

si-

es
s.
i-
ts
at

f
o
.
l
e
.



Carya Microcarpa.

Small fruited Hickory.

Noyer à petit fruit.

PLXIII



§

CA
l
s
r
Can
T
Jug
F
Jug

T
Hic
high
with
the
frien
mois
nut
tosa,
small
grow
seen
and
recon
rema
that
they a
3 pair
and 2

§ I. *Nuts more or less quadrangular.* HICKORY, properly so called.

SMALL FRUITED HICKORY.

CARYA MICROCARPA; *foliis quinis ad septenis, oblongo-lanceolatis serratis promisse acuminatis glabris subtus glandulosis; amentis glabris, nuce subglobosa subquadrangulata, testa tenui.*

Carya microcarpa. NUTT. Gen. Am., vol. 2. p. 221. DARLINGTON, Flora Cestrica. [Ed. alt.] p. 545.

Juglans compressa. a. microcarpa. MUHL. Catal., p. 88. BART. Flor. Philad., vol. 2. p. 179.

Juglans alba odorata. Balsam Hickory. MARSHALL, p. 68.

This species, allied to *C. tomentosa*, or the common Hickory, becomes a fine lofty spreading tree 60 to 80 feet high, having a diameter of 18 inches to 2 feet or more, with an even bark. I first observed it on the banks of the Schuylkill, in the vicinity of Philadelphia, and my friend Dr. Darlington remarks that it is frequent in moist woodlands in the vicinity of West Chester. The nut is of the same form nearly as that of *C. tomentosa*, of a pleasant taste, with a thin shell, but usually small, not much exceeding the size of a nutmeg. It grows, I believe, also in Massachusetts, where I have seen these peculiar nuts. The wood is white and tough, and possessed of most of the good qualities which recommend the ordinary Hickory. This species is remarkable for the smoothness of its leaflets, which, in that respect, approach *C. glabra* or the Pig Nut, but they are every way larger and less deeply serrate; 2 or 3 pairs with a terminal odd one, 4 to 8 or 9 inches long and 2 to 3½ inches wide, oblong-lanceolate, with shallow

serrulations, smooth on both sides when fully expanded, except a slight tuft in the axills of the nerves beneath; the under surface sprinkled with minute resinous particles; the lateral leaflets subsessile and rather obtuse at base, the terminal one with a short petiole and attenuated below. Aments 3 together, upon a common peduncle, slender, nearly quite smooth, scales trifid, the lateral segments ovate, the middle one long and linear; anthers hairy, mostly 4, sometimes 3 or 5. Female flowers 2 or 3 together, sessile, on a common peduncle; segments of the perianth very long and somewhat foliaceous. Stigma discoid, 4-lobed; fruit globose-ovoid, about $\frac{3}{4}$ of an inch in diameter; the pericarp thin, with the sutures rather prominent. Nut somewhat quadrangular with the shell thin.

By the leaves it appears to be allied to *C. glabra*, but the nut, on a small scale, is that of *C. tomentosa* or the common Hickory.

PLATE XIII.

A small branch, reduced about a third. *a.* The nut.

COMMON HICKORY, (*Carya tomentosa*, β . *maxima*.) This is a remarkable variety for the great size of its fruit, which are as large as a moderate apple. It grows a few miles from Philadelphia. Mr. Elliott also observed it on the sea islands of South Carolina.

Carya glabra, (*Juglans glabra*, Du Roi, Harbk., vol. 1. p. 335. *J. porcina*, MICH. North Am. Sylva, vol. 1. pl. 38,) of this there are two varieties, one with globose, and the other with turbinate fruit: intermediate forms are also met with, proving them to be no more than varieties.

§ II. *Nuts even, without prominent angles. Leaflets often numerous.*—PECANS.

Carya angustifolia. Juglans angustifolia, AIT. Kew. vol. 3. p. 361. *J. Pecan*, MUHL. in Nov. Act. Soc. Nat. Scrut. Berlin., vol. 3. p. 392. *J. olivæformis*, WILLD. Sp. Pl. 4. p. 457. A fine stately tree, formerly cut down for the sake of obtaining a single crop of nuts; remarkable for its numerous leaflets and their almost falcate form. In Massachusetts, where it has been submitted to cultivation, it never grows beyond the size of a shrub, being every year more or less cut down by the effects of the severe frosts.

Carya Pecan. Juglans Pecan, WALTER. *J. myristiciformis?* MICH. Sylva, vol. 1. pl. 39. This obscure plant of Walter, may, perhaps, be nothing more than *C. glabra*. Michaux's plant was unknown to Elliott.

Carya amara. Juglans amara, MICH. Sylv., vol. 1. pl. 33.

Obs. According to an experiment published in the Massachusetts Agricultural Journal, the sap of the Butternut tree (*Juglans cinerea*), is capable of producing as much sugar as that of the maple. Four of the trees yielded in one day nine quarts of sap, which produced 14 pounds of sugar.

THE BLACK WALNUT, (*Juglans nigra*), is met with as far north as Massachusetts, particularly in the western part of the State, as around Northampton. Mr. Emerson says, "*Juglans nigra* I have found repeatedly as far north as Boston. It is in Middlesex, Worcester, and Norfolk counties" (Massachusetts).

On the banks of the Sciota, in Ohio, I have seen a tree of 6 feet diameter.

CANDLEBERRY MYRTLE.

Natural Order, MYRICACEÆ, (Richard.) Linnæan Classification, DIOECIA, TETRANDRIA to OCTANDRIA.

MYRICA, LINN., (in part.)

Flowers unisexual; those of the two sexes upon the same or more commonly upon different plants. *Male* flowers in cylindrical sessile catkins; each flower with 4 to 8 stamens, with the filaments elongated and more or less united at the base; the stamens exserted beyond the borders of the dilated short scale, many stamens in branching clusters nearly without scales at the summit of the catkin; bracteoles *none* in either sex. *Female* flowers in loose, sometimes filiform catkins, with many of the lower scales abortive; scales 1-flowered, the germ naked. Styles 2, very long, linear, and acuminate, ovary villous. Drupe 1-seeded, spherical, coated with a grumose waxy pulp. Nut very hard; seed erect; embryo without albumen, the radicle superior. Cotyledons thick and oily.

A genus wholly distinct from *Myrica Gale*, which is common to northern Europe and North America. The character of lunate scales given to *Myrica* by Linnæus applies only to the *GALE*, which therefore constitutes a genus by that name. The rest of our species belong to *Myrica*. In the *Gale*, the fruit is a small ovate dry nut, with an indurated bracte on either side of it, giving it the appearance of being 3-lobed.


The species of this genus are few; natives of the warmer and colder zones of both hemispheres, growing generally near the sea coast, and are chiefly shrubs, with alternate persistent, or annual simple leaves, usually more or less serrated, or pinnatifid, and besprinkled with aromatic resinous scales, as are also the scales of the buds. Catkins axillary, expanding early in the

ifi-

or
in-
ith
se;
ort
ut
er
is,
d,
e,
u-
ut

n
te
e,
r
e
it

d
e
r
l,
e
e



Myrica inodora.

Inodorous Simile Tree

Crice inodore



y
C
a
d

M

M
M

E
co
sc
T
A
cu
ve
(M
wh
in
ris
of
ma
fig
ber
wa
frag
for
it a
it i
Tra
V

year. There are several species in Nepal in India and at the Cape of Good Hope. The *M. Faya*, with a 4-celled drupe, and about 10 stamens in loose catkins, will, no doubt, constitute a different genus, which I propose to call *FAYA AZORICA*.

INODOROUS CANDLE TREE.

MYRICA INODORA; *arborea, foliis lanceolato-ellipticis integris obtusis margine revolutis basi cuneatis subtus vix squamosis plerisque nudis, baccis majusculis albidis.*

MYRICA inodora, BARTRAM'S Travels in Florida, &c. p. 405.

MYRICA obovata, CHAPMAN, MSS.

IN the month of March, 1830, on the borders of the Escambia, in West Florida, I had the fortune to re-discover this fine species of *Myrica*, so long since described by the amiable and excellent Wm. Bartram, near Taensa Bluff, on the Mobile river, where he remarks, August the 5th, 1776, (67 years ago:) "In my excursions about this place, I observed many curious vegetable productions, particularly a species of *Myrica*, (*Myrica inodora*.) This very beautiful evergreen shrub, which the French inhabitants call the Wax tree, grows in wet sandy ground, about the edges of swamps; it rises erect 9 or 10 feet, dividing itself into a multitude of nearly erect branches, which are garnished with many shining deep green entire leaves of a lanceolate figure. The branches produce abundance of large round berries, which are covered with a scale or coat of white wax; no part of this plant possesses any degree of fragrance. It is in high estimation with the inhabitants for the production of wax for candles, for which purpose it answers equally well with bees-wax, or preferable, as it is harder and more lasting in burning." BARTRAM'S Travels, p. 405 and 406.

We found it exactly in similar situations as those

described by Burtram, and it has also been found in Alabama by Dr. Juet, from whom I have been favoured with specimens. It may with propriety be called a tree, though never so large a one as the *Myrica Faya*, or Fayal Myrtle. The stem sometimes attains the thickness of a man's arm, and, like the rest of the genus, it is gregarious and forms stout thickets on the margins of small streams and swamps. The berries are twice as large as those of the common Wax Myrtle. Though the leaves have no perceptible scent, they are not always entirely without the usual scaly resinous glands; they have no serratures, and are about 3 to 3½ inches long, by 1 to 1½ wide. The bark is of a grey colour, inclining to brown. The male catkins are unusually large, as well as the berries, and the leaves, when old, are as stiff as in the laurel. The stamens beneath each scale of the ament are 8, with distinct filaments and monadelphous at base, the summit of the catkin is nearly without scales, and terminates in monadelphous branchlets of stamens, each bearing 3 or 4 anthers. The female catkin is loose, and the lower scales empty; the germ is pilose. The wood appears compact, fine grained, and nearly white. The candles formed of the myrtle wax burn long, yield a grateful smell, and are destitute of the disagreeable scent produced on extinguishing tallow candles. In Carolina a kind of sealing wax has been made of it, and the root has been accounted a specific in toothache. In Prussia it has been cultivated for the wax.

The Fayal Myrtle, (*M. Faya*,) is in Fayal the principal article of fuel; it there attains the ordinary height of a peach tree, with a more erect stem; it produces a considerable quantity of compact reddish wood. It is also cultivated in rows between and around the orange

trees for the purpose of shelter from the cutting sea breezes, which would otherwise abridge their height and retard their growth. All the gardens of the island require the same shelter for which the evergreen Faya is so well calculated, being perfectly hardy and indigenous to the Azores.

PLATE XIV.

A branch of the natural size. *a.* The berry.

PLANE TREE.

Natural Order, PLATANÆ. *Linnæan Classification*, MONŒCIA, POLYANDRIA.

PLATANUS.† (TOURNEFORT.)

Flowers of one sex, those of the two kinds situated upon the same plant, and each of them disposed in spherical aments on pendulous stalks, producing from 2 to 5 upon each. MALE flowers formed of minute thickish bractes; the filaments very short, situated between the bractes. *Anthers* 2-celled, attached to a connectivum broader than the filament, with a peltate summit FEMALE. Pistils numerous, in pairs. *Ovary* of 1-cell, including 1 to 2 pendulous ovules. *Stigmas* 2, long and filiform. *Fruit* a carpel seated in a tuft of articulated hairs, including 1 pendulous oblong seed, destitute of albumen.

Lofty deciduous leaved trees, with widely spreading branches and a dense broad foliage having a pentangular outline. Natives of Europe, Asia, Northern Africa, and the temperate parts of North America. The species about 3. Leaves alternate, palmate; the bud concealed beneath a conical envelope and immersed in the base of the petiole. The young shoots and leaves covered with a deciduous down. The old bark at length scaling off in extensive patches, leaving the trunk smooth.

† The name is from the Greek word *platys*, broad, in allusion to its wide spread leaves and branches.

0-

nc
n
E
y
r-
e
-
d
,


g
-
,
e
l



Platanus racemosa.

California Sycamore

Platanus de Californie.



v
f
r
v
s
a
f
h
c
s
t
s
a
c
v

CALIFORNIA BUTTONWOOD, OR PLANE.

PLATANUS RACEMOSUS; *foliis quinquelobo-palmatis basi truncatis subsinuatis subtus lanuginosis pallidis, laciniis lanceolatis acuminatis integris, stipulis angulatis, fructibus racemosis.* NUTTALL, Mss. in Audubon's Birds of America, tab. 362.

THIS remarkably distinct species of *Platanus* is a native of Upper California, in the vicinity of Sta. Barbara, where it puts on very much the appearance of our common Buttonwood, (*Platanus occidentalis.*) As far as I yet know, it is the only species on the western coast of America. It grows probably farther north, but I did not meet with it in the territory of Oregon. It does not appear in this unfriendly climate to arrive at the gigantic magnitude of its eastern prototype, though it equally affects rich bottom lands and the borders of streams, but the scarcity of rain, in this climate, which had not for three years been sufficient to encourage the raising of crops, and the consequent disappearance of water in most of the brooks, prevented, no doubt, this subaquatic tree from assuming its proper character in a more favourable soil. At first view it would be taken for the ordinary species, spreading out the same serpentine picturesque limbs, occasionally denuded of their old coat of bark, and producing the same wide and gigantic trunk, but a glance at the leaves no less than the fruit, would remind the eastern traveller that he sojourned in a new region of vegetation, and objects apparently the most familiar he met around him, associate them as he would, were still wholly strangers.

The leaves not fully expanded were about 4 inches wide and the same in length, divided more than half

way down into five sharp pointed, lanceolate portions, of which the two lower are the smallest; all the divisions are quite entire, two of them in small leaves are suppressed, thus producing a leaf of only three parts. Above, as usual, the surface is at first clad with a yellowish copious down, formed of ramified hairs, which quickly falls off and spreads itself in the atmosphere. The under surface of the leaves are, however, always copiously clad with a coat of whitish wool, which remains. The young leaves, clad in their brown pilose clothing, have a very uncommon appearance, and feel exactly like a piece of stout thick woollen cloth. The branchlets, petioles, and peduncles are equally villous. The *male* catkins are small, less in size than peas, full of long haired scales, and with unusually small anthers. The *female* catkins are in racemes of 3 to 5 in number, with remarkably long styles, being between 2 and 3 tenths of an inch in length, and persistent on the ripe balls. The raceme with the full grown balls measures 9 inches. The tree has, therefore, a very unusual appearance, filled with these very long pendulous racemes, each bearing from 3 to 4 or even 5 balls, at the distance of about an inch from each other. The stigmas are at first of a deep and bright brown.

The wood of this species, as far as I could learn from the American residents at Sta. Barbara, is far preferable to that of the common Buttonwood, being much harder, more durable, less liable to warp, and capable of receiving a good polish; it is of a pale yellowish colour, like the young wood of the Oriental Plane, and bears some resemblance to beech wood in its texture. In the radiation of its medullary vessels, it resembles the wood of the common species.

PLATE XV.

A branch of the natural size. *a.* The carpel.

THE ORIENTAL PLANE (*Platanus Orientalis*) deserves to be planted in the United States as an ornamental tree. It grows to the height of from 70 to 90 feet, with widely spreading branches and a massive trunk, forming altogether a majestic object. The leaves are more deeply divided and indented than in our common species. A native of the East, where shady trees are not so abundant as in North America, it was celebrated in the earliest records of Grecian history. Xerxes, it seems, (according to Herodotus,) was so fascinated with a beautiful Plane tree which he found growing in Lycia, that he encircled it with a ring of gold, and confided the charge of it to one of the Ten Thousand. He passed an entire day under its shade, encamping with his whole army in its vicinity, and the delay so occasioned was believed to be one of the causes of his defeat. Pausanius (A. D. 170) mentions a Plane tree of extraordinary size and beauty in Arcadia, which was said to have been planted by Menelaus, the husband of Helen, and to have been at the time he saw it 1300 years old.

Plane trees were planted near all the public schools in Athens. The groves of Epicurus, in which Aristotle taught his peripatetic disciples; the shady walks planted near the Gymnasia and other public buildings of Athens; and the groves of Academus, in which Plato delivered his celebrated discourses, were all formed of this tree.

The remarkable Plane tree at Buyukdere, or the Great Valley, mentioned by Olivier, the naturalist, and after him by Poucqueville, Hobhouse, and various other writers, has a trunk that presents the appearance of 7 or 8 trees, having a common origin, which Olivier supposes to be the stool of a decayed tree, and which were all connected at their base. Dr. Walsh, who measured the tree in 1831, found the trunk 141 feet in circumference

at the base, and its branches covered a space of 130 feet in diameter. The trunk divides into 14 branches, some of which issue from below the present surface of the soil, and some do not divide till they rise 7 or 8 feet above it; one of the largest is hollowed out by fire, and affords a cabin to shelter a husbandman. The tree, if it can be considered a single plant, is certainly the largest in the world. But what renders it an object of more than usual interest, is that M. Decandolle conjectures it must be more than 2000 years old.

The wood of the Oriental Plane in the Levant and in Asia is used in carpentry, joinery, and cabinet-making. It is said to make beautiful furniture, on account of the smoothness of its grain, and its susceptibility of receiving a high polish.

Concerning our common Plane tree or Buttonwood, (*Platanus occidentalis*.) Dr. Darlington remarks in his *Flora Cestrica*, page 542: "It makes a noble shade in front of houses where it has room to develop itself;" and he further remarks that "the *wood* is not much esteemed; but is occasionally sawed into joists and other lumber." It is beginning to be considerably planted as a shade tree on the side walks of the streets in several of the large towns of the United States, and being seldom attacked by insects, and rarely elevating the pavements, it is exceedingly well calculated for this useful purpose in a climate subject to such ardent summer heats; but, if the Oriental Plane would answer the same purpose, and it is easily propagated, we should not only possess an ornamental but also a *useful tree*, as it regards the wood. The finest specimens of trees of this species in the vicinity of Philadelphia, are those round the Pennsylvania Hospital, which were planted about the year 1760.

POPLARS.

Natural Order, AMENTACEÆ, (JUSSICU.) *Suborder*, SALICINEÆ. *Linneæan Class and Order*, DICECIA, OCTANDRIA.

POPULUS† LINN.

DICÆIOUS.—*Aments* cylindric, with the scales deeply cleft. *Perianth* cup-shaped, oblique, and entire. *Stamens* about 8, (or from 30 to 100 or more,) inserted on the scale or perianth. **FERTILE FLORETS** with the scales and perianth as in the male. *Stigmas* 3 or 4. *Capsule* 1-celled, 2 to 4-valved. *Seeds* numerous, comose, with long soft hairs like wool.

Trees of the temperate and colder parts of Europe and North America, with one species in Asia. The leaves are alternate, roundish, or deltoidly cordate, the petiole, for the most part, vertically compressed towards the summit, and often glandular at the base; the flowers, (as in the Willows, to which they are intimately allied,) appearing before the leaves.

The Poplars are divisible into two sections or subgenera.

§ 1. Those properly so called, with about 8 stamens, and, usually, filiform stigmas.

§ 2. Those with from 30 to 100 or more stamens, and with broad dilated reniform stigmas.—**POTAMERIA.** These are mostly large trees which affect the banks of rivers, and includes *Populus lævigata*, (*P. canadensis*, MICH. fil.,) *P. angulata*. *P. monilifera*, *P. heterophylla*, (*P. argentea*, MICH. fil.,) and probably *P. candicans*, *P. balsamifera*, and our *P. angustifolia*.

† An old Latin name of uncertain derivation.

NARROW LEAVED BALSAM POPLAR.

POPULUS ANGUSTIFOLIA; *foliis ovato-lanceolatis lanceolatisve acutis, superne attenuatis penninerviis concoloribus glabris adpresso-serratis; ramulis teretibus glabris, gemmis resinosis.*

P. angustifolia, TORREY, *Lycum Nat. Hist. N. York*, vol. 2. p. 249.

Narrow Leaved Cotton-wood, of LEWIS and CLARKE.

As we ascended the banks of the river Platte, in our extended journey to the West, about Larimic's Fork, a northern branch of that extensive stream, we observed scarcely any other tree along the alluvial plains but the present and the Cotton-wood, and those were chiefly confined to the islands, a circumstance accounted for by the annual burning of the prairies, which wholly strips the streams of their margin of forest, so that we behold far and wide nothing but a vast plain, a sea of grass undulating before the breeze, and the illusion appears more sensible by the fact, that the only variation to the scene is produced by the scattered islands of the lofty Poplar which gives life and variety to the wild and boundless landscape.


The height of this species, which so nearly resembles the Balsam Poplar, may be about 60 to 100 feet, having a trunk of proportionate diameter, clad like the Cotton-wood with a rough greyish bark. Although a brittle and poor wood for almost every purpose, it will, like the Cotton-wood of the Mississippi, (*Populus angulata*), become of necessity important for fence and fuel, whenever this country shall become settled, as scarcely any other timber exists in sufficient quantity for economical

la-
ous
m-

2.

ur
a
ed
e
y
or
y
e
f
n
-
s
e

s
y
.






Populus Angustifolia.

Peuplier baumier à feuilles étroites

Narrow-leaved Balsam Poplar



purposes. When dry it burns well, but is quickly reduced to ashes.

Whatever may be the immediate uses of the Narrow Leaved Poplar, we must say, that in a country so exposed and arid as the Rocky Mountain region, we felt grateful for the shade and shelter it so often exclusively afforded us, with the exception of a few insignificant willows, that frequently associate with it. In short, we rarely lost sight of this tree, which accompanied us to the northern sources of the Platte, presented itself, as usual, on the alluvial banks of the Colorado of the West, along Lewis river, of the Shoshonee, the banks of the Oregon from the Walla-Walla to its estuary, attaining along the banks of this noble stream, and its southern tributary the Wahlamet, an augmentation so great as to vie in magnitude with the tall Cottonwood of the Mississippi, and to pass amidst the mighty forests of the West, as one of the largest deciduous leaved trees of the country. We find this species of Poplar also on the banks of the Missouri, on the upper part of the river, from whence it continues uninterruptedly to the vallies of the Rocky Mountains. In the severity of winter the boughs are collected by the aborigines to support their horses; the beaver likewise feeds upon them by choice.

Early in the spring the balsamic odour of its resinous buds may be perceived far and wide, and the shade of its tall, round, and spreading summit invited hosts of birds to its branches, particularly the large and fine Band-Tailed Pigeon, which feeds with avidity on its seed buds in the spring, affording them an abundant article of food.

The lesser branches are rather tough than brittle, covered with a smooth yellowish bark. The leaves vary on the same branch, in this respect, that the earli-

est brought out are the widest; these are generally ovate and acute, (not acuminate as in the Balsam Poplar,) rounded below; afterwards come out a set of lanceolate leaves, much longer than the others, often three inches by only one inch wide; these are acute at both ends, and would pass very well for those of a Willow; they are all margined with fine bluntish appressed serratures, and beneath they are nearly the same colour as above. The footstalks of the leaves or petioles are about $\frac{3}{4}$ to an inch in length. The nerves are all faint beneath and pennate, with no appearance of being 3-nerved at the base; the number of these nerves or lateral vessels are 12 to 15 on a side, at least double the number they are in the Balsam Poplar. The rachis of the female ament is smooth, the germs much corrugated in drying, and the scaly cup of the germ very shallow, not more than half the size of that of the species to which it is so closely allied. Upon the whole comparison, we are satisfied that this is as distinct a species as any in the genus. In the FLORA ROSSICA, vol. 1. p. 21. t. 41. B. a leaf is given as a variety of the Balsam Poplar from Altai, very similar to our present species.

PLATE XVI.

A branch of the natural size. a. The capsule.

COTTON-WOOD, *POPULUS LÆVIGATA*, Willd. (*P. canadensis*, Mich. Sylva, vol. 2. pl. 95.)

This species, more hardy than the Mississippi Cottonwood, is found on the banks of most of the western rivers for a considerable distance towards the Rocky Mountains. It is abundant on the borders of the Arkansas, but is no where met with in Oregon Territory.

BALSAM POPLAR, (*Populus balsamifera*.) In *Flora Rossica*, vol. 1. p. 67 and 41, is given a figure of a Poplar from Daouria, called the Balsam Poplar, which does not appear to be our plant. The leaves are wider, with shorter petioles. This species stretches into eastern Siberia, along the river banks to the peninsula of Kamtschatka.

AMERICAN ASPEN, (*Populus tremuloides*, MICH. Fl. Am., vol. 2. p. 243. *P. tremula*, WILLD.) This elegant species, of which the leaves, like the European Aspen, tremble and vibrate in the faintest breeze, is met with west of the Mississippi, in scattering groups, chiefly in the ravines of the mountains, where springs issue out, and in narrow vallies, where considerable groups of this tree may be seen into the centre of the Rocky Mountain chain, but they here attain a very inconsiderable magnitude, being seldom thicker than a man's arm.

WHITE POPLAR, (*P. Alba*.) now commonly cultivated with us for ornament, soon attains the magnitude of an Oak. This wood, remarkable for its whiteness, is generally used in the South of France for wainscoting and flooring their houses, and in England it is employed by turners for a host of small articles and utensils of housekeeping.

The wood of various species of Poplar has been employed for dyeing; that of the Lombardy or Italian Poplar boiled in water with wool communicates a very permanent golden yellow colour; other species, as the Black, Virginia Poplar and Cotton-wood give, according to the length of time employed, and the quantity of the wood boiled, various shades of colour, as that of nankin, musc, the colour of the Vicuna, with other shades, and answer conveniently for the ground of other colours.

W I L L O W S .

Natural Order, AMENTACEÆ, (JUSSIEU.) *Suborder*, SALICINEÆ. *Linnaean Classification*, DICEIA, DIANDRIA, &c.

SALIX.† (TOURNEFORT, LINN.)

The flowers are DICEIOUS‡ or very rarely MONÆCIOUS, disposed in cvoid or cylindric catkins, composed of undivided scales, which are imbricated over each other, and each with the pistils or stamens form a flower; at the base of these scales exists a small glandular body, which is either simple or bifid, and surrounds the interior organs of reproduction. In the *male* flowers there exists from 1 to 5, or even 7 to 9 stamens, (ordinarily there are only 2.) In the *female* flowers the ovary is single, terminated by a bifid style, having usually 4 stigmas. The capsule or follicle consists of 1 cell with 2 valves. The seeds are very numerous and minute, each terminated by a long tuft of hairs or pappus; the radicle is inferior, or in an inverse position to that of the Poplars, to which they are so intimately allied.

THE Willows, numerous in species, are all (with two exceptions in the straits of Magellan and Peru) natives of the northern hemisphere, and all of them shrubs or trees, some not more than an inch in height above the ground, confined to the highest summits of lofty moun-

† Said to be derived from the Celtic *sal*, near, and *lis*, water.

‡ A term used by Linnæus to designate a class of plants which have flowers of different sexes on two different individuals: *monœcia*, with two sorts of flowers on different parts of the same plant.

tains; others attain an elevation of 50 or 60 feet. The wood is usually light and close-grained; the twigs of several species, used by basket-makers, are remarkable for their pliability. The bark of most of the species contains a peculiar vegetable principle, called *salicine*, which, for intermittent fevers, is nearly as serviceable as Peruvian bark, and the down of the seeds has been manufactured into a coarse paper. Most of the species affect wet or humid situations, being common on the immediate borders of brooks and rivers, which they fringe with a luxuriant and agreeable vegetation. The flowers generally precede the appearance of the leaves, and though not beautiful, they are seen with delight, as the earliest harbingers of our northern spring. The species, more than 200 in number, present nearly the same general form of foliage, with the margin entire or serrated; the catkins are lateral or terminal, and the stamens are too variable in number to admit of a classification by them; there are some with only 1, a good many with 2, others with 3, 4, 5, or even 9 or 10. The capsules, though small, afford the best traits of specific distinction. The leaves put on various appearances, and even outlines, in the progress of their growth, and the Willows justly rank among botanists as the most protean and difficult family of plants to discriminate in the northern hemisphere. In our tour across the continent we have met with some remarkable species, four or five of which become trees; we shall offer descriptions of the whole, but only give figures of those which rank among arborescent species.

LONG-LEAVED WILLOW.

SALIX SPECIOSA; *foliis longissimis lanceolatis serrulatis promissa acuminatis glabris subtus glaucis junioribus pilosis; amentis serotinis 6—9-andris, germinibus pedicellatis lanceolatis acuminatis glabris, squamis lanceolato-oblongis sericeis, stigmatibus lobis bifidis.*

SALIX LUCIDA, HOOK. (as it regards the Oregon plant.) Flor. Bor. Am., vol. 2. p. 148.

No Willow on the American continent presents so remarkable and splendid an appearance as the present; the effect of which is produced no less by its magnitude than the size and beauty of its foliage. Its aspect is that of a large Peach tree, with the leaves and their stalks from five to eight inches long by an inch to an inch and a half wide; beneath, when adult, they are glaucous, like those of the River Maple. The summit is tufted and spreading; and the tree attains the elevation of about 20 to 30 feet, with a trunk of 12 to 18 inches in diameter. When in full bloom, which is with the first expansion of the leaves, in May, the numerous and large bright yellow catkins, loading the branches, emulate the finest Acacia of New Holland; they are also agreeably fragrant, and attract swarms of wild bees and other insects, in continual motion among their waving branches. We have seen this noble species nowhere in such perfection as along the banks of the deep Wahlamet and the wide Oregon, whose numerous islands are almost exclusively decked with this imposing Willow, which continues to the Blue Mountains, and along the neighbouring streams as far east as the river Boisé. As we sailed along the smooth bosom of these

serrulatis
junioribus
us pedicel-
lanceolato-

) Flor. Bor.

resents so
e present;
magnitude
aspect is
and their
nch to an
they are
summit is
elevation
18 inches
with the
erous and
ies, emu-
are also
wild bees
ng their
ecies no-
the deep
numerous
imposing
ins, and
the river
of these

Handwritten text on the right margin, possibly a page number or reference, including the number 110.



Salix speciosa.

Long leaved Willow.

Saule gracieux.



extensive streams, for many miles we never lost sight of the Long-leaved Willow, which seemed to dispute the domain of the sweeping flood, fringing the banks of the streams and concealing the marshes entirely from view; at every instant, when touched by the breeze, displaying the contrasted surface of their leaves; above of a deep and lucid green, beneath the bluish-white of silver; the whole scene, reflected by the water and in constant motion, presented a silent picture of exquisite beauty. Immediately behind this foreground of spreading Willows, arose in the first rank of the legitimate forest, the lofty Poplars we have already described, succeeded by the majestic Oaks and Maples, while the distant hills to their summits were impenetrably hid by the vast towering Pines and Firs, which mingling as it were with the clouds, close in the rest of the landscape with funereal grandeur.

This species is related to the Lucid and Bay Willows, and the buds have something of the same aromatic exudation, the serrulations and the base of the leaf are also equally glandular. The bark of the trunk is rough and divided, the twigs smooth and shining, of a yellowish-brown. The leaves, at first green on both surfaces, are, before expansion, clothed with long brown loose parallel hairs, which disappear with the progress of their growth, at length they become silvery and glaucous beneath; they are finely serrulated, acute at both extremities, with the points very much attenuated. The stipules are semicircular and broad, serrulated on the margin. The flowers come out with the expanding leaves from lateral buds, containing, in the male, also two or three leaves, so that the catkin appears as a pedunculated spike. The stamens are very long, from 5 to 9 in a scale, with filaments which are hairy towards the

base, and as well as the hairy, broad, cuncate, serrated scales, are of a bright golden-yellow.

The female plant, at the time of flowering, appears to have smaller leaves than the male, and those on the branches which bear the catkin, are green on both surfaces for a considerable time; they are also but little acuminate: 5 or 6 leaves with their appropriate stipules grow out on the same branch, which terminates in a female spike. The scales of the catkin or spike are oblong or lance-oblong, and less hairy than in the stamiferous catkin. The germ is lanceolate, pedicellate and smooth, acuminate and terminated by a short bifid style, with 2 pubescent bifid stigmas. The capsule is likewise smooth, and contains seeds with a very long pappus, as abundant almost as on a seed of cotton.

The wood is whitish and close-grained, and might probably be employed for the same purposes as that of the White Poplar, but the nearly uninhabited state of the country in Oregon, prevents the possibility of making any useful experiments. As an ornamental and hardy tree, however, it stands pre-eminent among all its fraternity, and well deserves to be introduced into pleasure grounds, where it would be perfectly hardy as far north as New York, or in any part of Great Britain.

PLATE XVII.

A twig and leaf of the natural size. *a.* The female catkin. *b.* The male catkin. *c.* The male flower and scale of the catkin. *d.* The open capsule.

e, serrated

g, appears
ose on the
both sur-
but little
oriate sti-
erminates
or spike
an in the
ate, pedi-
ted by a
as. The
ls with a
a seed of

nd might
as that of
ate of the
f making
nd hardy
l its fra-
pleasure
far north

le catkin.
he catkin.




Salix pentandra.

Long leaved Bay Willow.

Sauze laurier.





LONG-LEAVED BAY WILLOW.

SALIX PENTANDRA; *foliis ellipticis acuminatis serratis glabris, petiolis superne glandulosis, amentis scrotinis pentandris, germinibus lanceolatis glabris*. WILLD. Sp. pl. l. c. VAHL. in Flora Danica, tab. 943. HOST. Sal. Austr. 1. t. 1. f. 2. Eng. Bot. t. 1805.

SALIX PENTANDRA, *foliis serratis glabris, floribus pentandris*. LINN. Hort. Cliffort. p. 454. et Sp. pl. p. 144. Flora Lapponica, p. 370. t. 8. fig. 3. GMELIN, Flora Sibirica, vol. 1. p. 153. t. 34. fig. 1.

Salix foliis glabris, ovato-lanceolatis; petiolis glandulosis; floribus hexastemonibus. HALLER, Flora Helvetica, No. 1639.

Salix montana major; foliis Laurinis. TOURNEFORT, Institutes Rei. Herb. p. 591.

Salix foliis laureo sive lato glabro odorato. RAII. Hist. p. 1420.

SALIX PENTANDRA, β . CAUDATA, *foliis longissime acuminatis attenuatis; ramulis junioribus hirsutis; capsulis rugulosis opacis*.

WE met with this species, hitherto wholly European, in the very centre of the North American continent, by streams in the valleys of the Rocky Mountains, towards their western slope, in Oregon, and also the Blue Mountains of the same territory. It is true, the present variety is a remarkable one. The uppermost leaves on the adult branches are about 5 or 6 inches long, one-half of which distance, or about $2\frac{1}{2}$ to 3 inches, may be considered as the slope of the summit and the point of the leaf; the buds and last developed branches are also hirsute. This variety, like its prototype, becomes a tree 15 to 20 feet high, of a bright, but not deep,

and rich green. The flowers are fragrant, produced after the appearance of the leaves, and the capsules in our plant are slightly corrugated, even when ripe and open, and do not shine as in the common Bay Willow. The leaves have the same odorous glands, and the bark of the branches is smooth, shining, and of a brownish-yellow colour. This tree is met with in mountainous situations, by streams, in all the northern parts of Europe, in Britain, France, Switzerland, Sweden, Lapland, and throughout Siberia and Russia. Its branches are too fragile to be employed for any economical purpose, and the wood decrepitates in the fire. The leaves, which are fragrant from the resinous glands of their margin, however, furnish a yellow dye, and the abundant down of its seeds in some of the northern countries, is used with success as a substitute for cotton, mixed with a third part of the true material.

According to Loudon, it is one of the most desirable species of the genus for planting in pleasure-grounds, on account of the fine display made by the blossoms, their abundant fragrance, the shining rich deep green of the leaves, and the comparatively slow growth and compact habit of the tree. It is also one of the latest flowering Willows, the blossoms seldom expanding till the beginning of June.

PLATE XVIII.

A branch of the natural size. *a.* The capsule.

produced
capsules in
ripe and
y Willow.
the bark
brownish-
mountainous
parts of
den, Lap-
branches
nical pur-
ne leaves,
of their
abundant
untries, is
ixed with

desirable
unds, on
ms, their
n of the
compact
owering
the be-




Western yellow Willow.

Salix lutea.

Saule jaune

PL. XIX.





WESTERN YELLOW WILLOW.

SALIX LUTEA, *foliis ovato-lanceolatis acutis leviter serrulatis glabris, stipulis lunatis, amentis præcocibus brevibus cylindraceis, germinibus pedicellatis glabris, ramis luteis.*

THIS is a smallish arborescent species of Willow, which inhabits the Rocky Mountain region, and proceeds westward to the banks of the Oregon, but is no where common. It attains to the magnitude of a small tree, and at first glance appears nearly allied to the common yellow twigged Willow (*S. vitellina*), but it never grows so large, and is after the very first evolution of the leaf buds perfectly smooth, with none of those tufts of hairs which in that species appear at the interior base of the bud; it likewise possesses stipules, and bears short close catkins, with smooth capsules, and the serrulations of the leaves are not glandular, indeed scarcely visible, the leaf appearing to have a thin and often almost entire diaphanous margin.

It is at the same time a very elegant and distinct species, remarkable for its smooth bright yellow branches and pale green, rather small, lucid leaves. It is well deserving of cultivation for the sake of variety, and is undoubtedly hardy from the climate it inhabits. The leaves are rather narrow, more lanceolate than ovate, acute, or somewhat acuminate, about an inch and a quarter long, and less than half an inch wide. The minute serratures or serrulations are rather crowded, but very minute and shallow, and not in the least cartilaginous. The young leaves before evolution are silky and white beneath, but the pubescence wholly disappears with their expansion.

PLATE XIX.

A branch of the natural size with fertile catkins. *a.* The capsule.

HOOKER'S WILLOW.

SALIX HOOKERIANA. (Barratt, MSS); *ramis valde robustis pubescentibus, junioribus dense tomentoso-lanatis, foliis late obovatis fere subrotundatis rigidiusculis serratis margine planis supra (adultis) nudiusculis subtus tomentoso-lanatis, stipulis [semicordatis], amentis cylindraccis crassis, squamis longe densissime lanatis, ovarii longe stipitatis lanceolatis glaberrimis, stylo breviusculo, stigmatibus lobis integris.* Hook. Flor. Bor. Am., vol. 2. p. 145. t. 180.

THIS small tree, nearly allied to *S. cinerea*, we found of common occurrence in the lower part of the territory of Oregon towards the sea, particularly frequent on the margins of ponds and in wet places near the outlet of the Wahlamet, where it attained the height of eight or ten feet. It was also found in the territory of Hudson's Bay, near the Grand Rapids of the Saskatchewan, by Douglas. The leaves are remarkably protean in their form, sometimes wholly soft and villous on both sides, more commonly so beneath, the nature of the pubescence also varying till it at times resembles in appearance and to the touch the most brilliant velvet; the old leaves are generally obovate, smooth and shining above, often nearly so beneath, and then of a bluish-green or glaucous hue; they are, moreover, both acute and obtuse, sometimes even rounded at the extremity, and are generally among the largest, or at least widest, of Willow leaves. The stipules on some branches are very conspicuous, circular and serrated. The male aments are extremely pilose in an early stage, as in our *S. conifera*, to which this species makes some approach; the stamens are two to each scale. The capsule is ovate-lanceolate

and very smooth, the style elongated, and the stigmas two and undivided.

The twigs of this species are flexible, but the wood is too small for almost any economical employment. The old wood of the trunk of *S. caprea* or the Sallow, so much like the present species, is much esteemed by wood turners.

BLUNT LEAVED WILLOW.

SALIX FLAVESCENS, foliis obovatis sublanceolatis flavescente tomentosis integris demum glabris, stipulis parvulis subsemicordatis denticulatis, amentis præcocibus, capsulis lanceolatis sericeo-pubescentibus, stigmatibus villosis.

WE met with this shrubby species in the range of the Rocky Mountains, and we are doubtful if it is not also an inhabitant of Europe. It agrees very nearly with the obtuse leaved Willow of Willdenow, (*S. obtusifolia*), of which we have seen no specimens, and which is apparently a very obscure species, said to be a native of the Lapland Alps. Ours is a large shrub, with much of the aspect and general character of the Grey Willow, which is indeed the type of a groupe of kindred Willows. For a good while the leaves still remain downy, particularly on the under surface, which is tinged also with pale yellow. The leaves, when old, are about $1\frac{1}{2}$ inches long, $\frac{3}{4}$ of an inch wide or more, wedge-shaped at the base, obovate or oblanceolate, quite smooth, entire apparently, yet the stipules are denticulate. The branches are brown or dark purple. The catkins are short and cylindric. The scales blackish, hairy, and obtuse, the capsules white and silky; the style is distinct, and the four stigmas long and pubescent.

POND WILLOW.

SALIX STAGNALIS, foliis oblongo-lanceolatis obtusis integerrimis basi cuneatis subtus pubescentibus, stipulis nullis, amentis coætaneis, capsulis lanceolatis acuminatis sericeis, squamis sublanceolatis.

THIS is another species indigenous to the banks of the Oregon, related to the same section with the last. The twigs are dark brown and slender, and the leaves scattered beneath with a minute brown pubescence, which communicates a somewhat rusty appearance to the leaves; they are about $1\frac{1}{2}$ inches long, and $\frac{1}{4}$ to $\frac{3}{4}$ of an inch wide, in the bud they are covered with long silky hairs. The capsules are remarkable for their great attenuation and length, the scales of the ament are oblong-lanceolate and hairy. We have not seen the male plant, and our account is therefore imperfect.

 VELVET WILLOW.

SALIX CUNEATA, foliis lanceolatis oblongisve apice subserulatis acutis, basi cuneatis glabris subtus holosericeis, stipulis minutis plerisque nullis, amentis coætaneis elongatis, capsulis ovatis sericeis, stigmatibus brevibus glabris.

THIS beautiful Willow we found growing in clumps near the rocky margin of the Oregon at its confluence with the Wahlamet, attaining the height of six or eight feet. The branches are slender, and, according to their age, vary much in appearance, at first villous and

downy, but at a later period brown, and sometimes quite blue, with a glaucous bloom. The leaves, with their short petioles, measure about two inches, and are about half an inch wide, pubescent above, at length nearly smooth and deep green, but always clad beneath with a whitish close tomentum, producing all the brilliant display of the finest velvet. The male flowers we have not seen. The female catkins are rather long, loose, and subcylindric, often accompanied at the base by two or three leaves, and come out when the leaves are considerably grown. The capsules are silky, short, ovate, and acute. The style short, and the stigmas four and smooth. The scales of the catkin are brown and oval, somewhat hairy, and much shorter than the capsules. In the narrow leaved varieties, the leaves appear almost wholly entire. The broader leaved plants bear some resemblance to the Grey Willow, but the serrulations are minute and the stipules very small, or wholly wanting.

WESTERN POND WILLOW.

SALIX MACROCARPA, foliis lanceolatis angustatis subintegerrimis utrinque acutis subacuminatis demum glabris subtus glaucis, stipulis obsoletis, amentis coetaneis dian-drus, capsulis ventricosus caudatis glabriusculis pedicellatis, stigmatibus subsessilibus quadrifidis.

This species, like our Pond Willow (*S. grisea*), to which it is closely related, is found forming clumps in wet places where the water is stagnant, situations which it always seems to prefer to the banks of running streams. It attains the height of 3 or 4 feet. The

branches are smooth and brownish-black, sometimes glaucous or whitish. The leaves (about 2 to 3 inches long and half an inch wide) are at first covered with a brownish silky down which disappears with age, when the leaves become dark green and bluish-white, or glaucous beneath, they are usually very acute, and mostly entire. The catkins are small and oblong, with 2 or 3 leaves at the base of the pedicel; the scales of the stamens are small and blackish, oval and obtuse, somewhat hairy. The female aments have very short pedicels, and produce at the base about 2 leaves; the scales are narrow and linear; the capsules pedicellated, somewhat villous, but at length nearly smooth, ventricose-lanceolate, with long points, and nearly sessile stigmas.

SOFT LEAVED WILLOW.

SALIX SESSILIFOLIA, foliis lanceolatis sessilibus acutissimis apice suberrulatis villosis mollibus, stipulis nullis, amentis scrotinis diandris elongatis terminalibus! germinibus lanatis, capsulis lanceolatis demum subglabris, stylo profunde bipartita stigmatibus bifidis.

This beautiful and very distinct species of Willow, formed dense tufts on the rocky borders of the Oregon, at the confluence of the Wahlamet, attaining to the height of about 6 to 8 feet, and when in flower appeared as showy as a *Mimosa*. It is remarkably leafy, and the leaves are hoary, with a rather long and somewhat copious pubescence, which communicates a softness to the touch equal to that of velvet; whether the leaves afterwards become more smooth or not I cannot say, but think it probable. Different from almost every other Willow I have seen, the catkins each terminate so

many small leafy branches about 2 inches long, exclusive of the catkin or spike, which is itself about 1½ inches—in both sexes attenuated at the base; these branchlets as well as the leaves are whitish, with soft hairs, particularly the former, but still the green colour of the leaf predominates; the points of the leaves are somewhat rigid, sharply acute, and, unlike most other species, they are destitute of petioles or footstalks; the scales of the ament are oval and unusually conspicuous, more hairy in the stamiferous flower. The stamens are 2 to a scale. The capsule is pubescent and lanceolate, at length nearly smooth. The stigmas 4, and rather long. No vestiges of stipules appear in any of our numerous specimens: the older branches are dark brown.

It is difficult to decide on the affinities of this very distinct species, which at a little distance resembles a *Protea* or *Leucodendron*, the leaves being equally grey and silvery, with soft hairs, which are so equally distributed on either surface as to obliterate the presence of the veins, and render both surfaces almost similar. It appears, in some respects, to resemble *S. arenaria*, the Sand Willow, but the late appearance of the aments and their remarkable disproportion, are almost without a parallel.

PROSTRATE WILLOW.

SALIX BRACHYCARPA, foliis ovatis lanceolatisve acutis subsessilibus integerrimis cinereo-pubescentibus subtus incanovillosis, stipulis nullis, amentis coætaneis brevibus glomeratis, capsulis ovatis abbreviatis tomentosis, stigmatibus subsessilibus.

THIS singular prostrate and dwarf Willow we met

with in the Rocky Mountain range, on the borders of Bear river, a clear rapid brook cutting its way through basaltic dykes to the curious lake of Timpanagos, in New Mexico. This locality is likewise famous for the numerous seltzer springs so strongly impregnated with carbonic acid as to sparkle and effervesce like champagne. Our hunters called them the "Beer Springs," and for a day and a half that we spent at this memorable place, the waters afforded us a most delicious treat during the warm weather, in those arid plains. In an open marshy situation, on the margin of the river forest, grew an abundance of this curious, depressed and hoary shrub, which has somewhat the aspect of the European Sand Willow, (*S. arenaria*,) but it is much more dwarf, with the leaves smaller, and always hoary with pubescence. The stem branches from the base only rising four or five inches above the surface of the earth, but with many diffuse, tough, woody branches, which spread out into a circle of a foot or more. The root stock is woody and thick; the branches full of cicatrices, pubescent, but brown beneath, thickly covered with small leaves, which in some plants are elliptic-ovate, in others oblong-lanceolate, all very entire, nearly sessile and acute, from half an inch to an inch long, about three lines wide; above always grey with pubescence, but beneath rather whitely villous; some of the lowest small leaves are smooth on the upper surface. There are no stipules in any of my specimens. The male flowers I have not seen. The fertile catkins are short and somewhat clustered, not cylindric, few flowered, the capsules oblong-lanceolate and short, villous, with appressed hairs, not densely lanuginous, as in *S. arenaria*, terminated by a short slender style and four short stigmas.

orders of
through
gos, in
for the
ed with
e cham-
brings,"
memo-
delicious
plains.
e river
sed and
of the
much
hoary
se only
earth,
which
e root
trices,
with
ate, in
sessile
t three
e, but
small
are no
vers I
some-
psules
ressed
termi-
nas.



Salix argophylla.

Silver leaved Willow

Saule à feuilles argentées



villos argenteas

SILVER LEAVED WILLOW.

SALIX ARGOPHYLLA, *foliis lineari-sublanccolatis acutis sessilibus integerrimis utrinque argenteo-sericeis, stipulis obsolete, amentis serotinis diandris, capsulis villosis lanccolatis.*

IN our devious progress to the West, we at length approached one of the branches of the Oregon, the river Boiséc, towards its junction with the Shoshonee; its banks were not fringed with a belt of forest, but so stripped of every character of an alluvial nature, that when we suddenly approached it, there appeared no break in the plain, and the clear and rapid flood shot through a deep perpendicular chasm of columnar basaltic rocks. We descended towards its brink and pursued our path along its obstructed banks, climbing over fearful rocks and along the margins of impending precipices, night approached without any cessation of our incessant toil. At length we hailed with satisfaction a small portion of the river bottom, almost covered by tall bushes of a remarkable silvery appearance, which proved to be the subject of our present examination, a very curious species of Willow, which, with a kind of Mock Orange, (*Philadelphus*,) composed nearly all the conspicuous vegetation of this sterile chasm. At length our new *Alnus*, (*A. Oregona*,) the Narrow-leaved Balsam Poplar, and the Long-leaved tree Willow, gave evidence of an ample alluvial plain and the proximity of the great Shoshonee.

This species becomes a small tree from 12 to 15 feet in height, as silvery and white as the *Leucodendron*

argenteum, the branches are brown, but the twigs are hoary with villous hairs. The leaves are very much crowded, soft, with whitish shining silky down, so abundant on either side as wholly to hide the veins, and nearly the midrib; they are also nearly without foot-stalks, entire on the margin, of a narrow linear outline and sharply acute, with a distinct bristly point, $1\frac{1}{2}$ to 2 inches long, and only about three lines wide. Stipules small and linear, seldom seen. The aments come out late with the leaves, and the flower branches produce from four to seven leaves. The male ament is small and narrow, with the scales lanceolate and villous; the female aments are oblong, the capsules lanceolate and villous.

The wood of this species, though small, is very white, smooth, and close-grained; the species must also be hardy, well worth cultivating, and would then probably grow to a considerable size. We perceive no affinity that this species bears, except perhaps to the *S. angustifolia* of the borders of the Caspian, from which at the same time it is probably very distinct.

PLATE XX.

A branch of the natural size. *a.* The male catkin. *b.* The capsule.

LONG SPIKED WILLOW.

SALIX MACROSTACHYA, *foliis lineari-sublanceolatis elongatis remote serrulatis acutis utrinque argento-sericeis, stipulis semiovatis deciduis, amentis longissimis præcocibus, capsulis lanceolatis demum glabriusculis.*

WE met with this species sparingly on the banks of

the Oregon. It bears a strong resemblance to the last, without however being any way intimately allied to it. It forms a slender bush, remarkable for its white and silvery pubescence, which appears as soft and glossy as velvet, the twigs are also pubescent. The leaves are two to three inches long, two or three lines wide, distinctly serrulated and nearly sessile, with the veins conspicuous through the pubescence. The female catkins, with their peduncles, are three or four inches in length, the capsules are nearly sessile, and at length but slightly pubescent. In this species there are distinct stipules on the young branches.

RIVER WILLOW.

SALIX FLUVIATILIS; foliis linearibus utrinque acuminatis, sublancoelatis spinuloso-serratis demum glabris concoloribus, stipulis nullis, amentis serotinis pedunculatis villosis diandris, squamis oblongis, fructibus lanceolatis glabris pedicellatis, stigmatibus sessilibus.

THIS species lines the immediate border of the Oregon, a little below its confluence with the Wahlamet, attaining the height of about six feet or more. We believe this is also the same Willow that we mistook for the Long-leaved species of Pursh and Muhlenberg, (*Salix longifolia*,) which so commonly lines the banks of the Missouri and Mississippi, and which often forms the exclusive growth of the small islands and sandbars, preparing these wastes, recovered from the flood, for a superior growth of trees, and they are also accompanied and succeeded commonly by the Cotton-wood, (*Populus canadensis*.) We met with this species like-

wise on the banks of Lewis River of the Shoshonee, accompanying our Long-leaved Willow, both of which continue almost uninterruptedly to occupy the banks and bars of all the Western streams to the Oregon, and proceed along that river to the borders of the Pacific.

In this remarkably fluviatile species, the leaves of the very young plants are somewhat pinnatifid, and at all times the serratures, rather distant, are sinuated and very sharp or spinulosely acute. The branches are brownish and very full of leaves. The leaves two to three inches long, are seldom more than two lines wide, except in the young shoots, when they are twice that breadth. The branches producing the male flowers are as short as usual, bearing only three or four small leaves: the catkins are narrow, solitary, and rather short, and come out with the opening of the leaves. In the female plant the inflorescence is similar with that of the male, but there is also produced a later growth of catkins, which terminate proper divided branchlets. The scales of the female catkin are oblong and densely bearded below, the germ is smooth, with four sessile stigmas as in *S. longifolia*. The young leaves are at first somewhat hoary and pubescent, with minute hairs; the young plants have also often pubescent foliage.

We have met with the *Salix longifolia* on the banks of the Arkansa, and it greatly resembles the present species, but differs in producing distinct stipules, in the minute serratures, and above all in the pubescent capsule and elongation of the catkin. The leaves are also generally broader, and it is said to grow only about two feet high.

SLENDER WILLOW.

SALIX EXIGUA, *foliis linearibus utrinque acutis subintegerrimis sericeis, stipulis nullis, amentis serotinis elongatis, capsulis lanceolatis sessilibus, demum nudiusculis.*

THIS species is also a native of the territory of Oregon, and grew with the preceding, which it strongly resembles: it is, however, a smaller species, with still narrower leaves, at all times more or less grey and silky, the serrulations are mostly wanting, though very minute ones are sometimes seen: the capsules are smaller, and not pedicellated. The male plant I have not seen. The branches are reddish brown and smooth.



ROUND LEAVED WILLOW.

SALIX ROTUNDIFOLIA, *foliis subrotundis ovatisve minutissime serrulatis vel integris utrinque glabris concoloribus, petiolis ramulisque pilosis, stipulis maximis dilatato-cordatis membranaceis glanduloso-serratis, amentis coetaneis diandris lanuginosis, capsulis glabris lanceolatis, stylis elongatis.*




β. *OVATA*, *foliis ovatis acutis dense serrulatis.*

THIS dwarf and remarkable species of Willow, attaining about two feet in height, was obtained in the gorge of a lofty alpine ravine, through which we fruitlessly endeavoured to pass. It was in the month of July, and the perpetual snows which still covered the mountains, mingled their outline with the skies. On the





**IMAGE EVALUATION
TEST TARGET (MT-3)**



**Photographic
Sciences
Corporation**

23 WEST MAIN STREET
WEBSTER, N.Y. 14580
(716) 872-4503

0
E 128
E 132
E 136
E 140
E 144
E 148

10
01

skirt of this gelid region grew our present subject. We named this scene of toil and disappointment, Thornburgh's Pass, or rather ravine, as no passage was practicable, from the man who undertook to be our guide. It was in the central chain of the Rocky Mountains, and near to the pass of the Shoshonees, which the following day we attained.

We know of no species with which we can compare this Willow. The older branches are brown, smooth, and full of cicatrices left by leaves that have grown near together, giving the plant a stunted appearance; the petioles are about $\frac{1}{2}$ an inch long, with the younger branches hairy, the younger leaves are also somewhat so on the mid-rib. The leaves are nearly round, from $1\frac{1}{2}$ to 3 inches wide, and about the same in length, though some of the later produced leaves are ovate and sometimes even acute, both sides are equally green, the margin in the ovate leaves elegantly and very closely serrulated, but in the round leaves the serrulations are often nearly obliterated. The stipules are very large, wide, and heart-shaped, finely serrated with glandular points, at length they become membranaceous and deciduous. The male aments are oblong, large and sessile, the scales blackish and ovate, producing copious white hairs longer than their whole length, the filaments are very long. The female aments grow on thick stalks, and have the scales also very woolly; the capsules are smooth and ovate, acuminate; the style is long, terminated by 4 stigmas. The *ovate leaved* variety was collected by the late Dr. Gairdener, on the hills of the Wahlamet.

MINUTE WILLOW.

SALIX NIVALIS, *foliis ovalibus subspathulatis integerrimis glabris obtusis, subtus concoloribus reticulatis, amentis serotinis pedicellatis paucifloris, germinibus ovatis sericeis squamæ glabræ retusæ longioribus, stigmatibus subsessilibus.*

SALIX nivalis. HOOKER, Flor. Bor. Am., vol. 2. p.

THIS elegant and very diminutive shrub, merely about 2 inches high, was obtained in the same alpine ravine as the former, but on a plain elevated to the very line of perpetual snow, at the height of about 10,000 feet above the ocean level. The stems are smooth, brown and woody, sending out very few short branches terminated by scanty tufts of smooth coriaceous leaves, 2 to 3 lines wide and about 4 lines long. The male plant we have not seen; the female catkin is produced upon a pubescent pedicel nearly its length, the flowers are about 6, sessile, and the subtending scales smooth, retuse and nerved. The germ is short ovate, and silky, terminated by a sessile or almost sessile quadrifid stigma.

This species is allied to the *Salix myrtilloides*, but at the same time perfectly distinct.

PLATE XIX.

A fertile plant of the natural magnitude. *a.* A leaf. *b.* The young capsule and its scale.


DUSKY WILLOW.


SALIX MELANOPSIS, foliis lineari-lanceolatis serrulatis glabris basi attenuatis vix petiolatis, stipulis nullis, amentis coetaneis diandris, squamis striatis obtusis, capsulis glabris lanceolatis subsessilibus.

This species, which I have called Dusky Willow from the dark appearance it assumes in drying, we met with at our station called Fort Hall, in the plains of the Rocky Mountains, on the alluvial lands of Lewis river of the Shoshonee, not growing in masses but scattered over the banks of the river in the more elevated situations, and there attaining the magnitude of a small tree 12 to 15 feet high, with a spreading summit, and when in flower forming a very elegant object. It is closely related to the *Triandrous Willow* of Europe, (*Salix triandra*,) but still sufficiently distinct. It never, like that species, becomes a considerable tree, but more resembles in its magnitude and mode of growth our common Black Willow (*S. nigra*). The wood is white and close grained as in that species: the young branches appear blackish-brown in a dead state, and the young leaves appear also very dark, they are about an inch and a half long, and about three lines wide, attenuated at the base, so as to present no distinct petiole, acute above, with minute serratures; there are no hairs at the base of the bud, as in *S. triandra*. The male flowering branchlets are provided with five or six leaves; the catkins linear and elongated, with dusky pubescent scales, marked each with about five striatures. The female catkins are rather short, and the fruit smooth,

glatis gla-
s, amentis
salis gla-

low from
met with
s of the
wis river
scattered
ed situa-
small tree
nd when
s closely
, (*Salix*
ver, like
at more
wth our
is white
branches
e young
an inch
enuated
, acute
airs at
le flow-
ves; the
bescent
. The
smooth,






Dusky Willow

Salix melanopsis.

Saule noirâtre



with sessile stigmas, as in other species of this particular groupe.

PLATE XXI.

A branch of the natural size. *a.* The scale. *b.* The stamen. *c.* The germ.

The *Salix triandra*, so nearly allied to the present species, becomes a tree 30 feet high, and is frequently planted in osier grounds for the basket-maker; but the best kind planted for this purpose is the *Osier*, properly so called, (*S. viminalis*,) which might be propagated in almost every part of the United States.

BLACK WILLOW.

SALIX NIGRA, *Mich. Arb.*, vol. 2. plate 125. fig. 1. A variety of this tree occurs in South Carolina and Florida, in which the leaves are villous and the scales of the ament densely lanuginous. In the herbarium of Mr. Schweinitz it was marked, on the authority of Elliott, as a species *S. subvillosa*.

This tree, a native of all the States from New England to Florida, and west nearly to the foot of the Rocky Mountains, is one of the few native species which becomes a tree, attaining the height of 15 to 25 feet, with a diameter of 10 to 15 inches. It affects the banks of rivers and lakes, and generally grows near to the water, in moist, occasionally overflowed situations. In the warmer parts of the States it puts on a handsome appearance, particularly when in flower, but has the defect of branching almost from the base, and leaving

in a posture more wild and picturesque than beautiful. It has a dark coloured rough bark, and the branches are brittle at the base; the male catkins coming out with the leaves are very elegant, and attractive and useful to bees. In the southern parts of the Union, according to Mr. Elliott, the stems when found sufficiently large are employed for the timbers of boats, and are found to be light and durable.

The **WHITE WILLOW**, (*Salix alba*, LINN.) appears almost naturalized in many parts of the United States. It is particularly abundant in the western interior of New Jersey along the borders of the Walkill. It grows rapidly, becoming a stout tree in a few years, and in Europe it is considered the most valuable timber tree of the genus; it produces a white close-grained wood, capable of receiving a polish, it also affords fence-wood, fuel, and bark for the tanner, nearly as good as that of the Oak.

The uses of Willows and Willow-wood in Europe are very numerous. It is generally a close fine-grained white wood, capable of taking a smooth and equal polish, and remarkably light withal. The osiers are very extensively used for all kinds of basket-work, and, as Virgil remarks, the shepherd sits beneath its shade, while it affords fence for his field, browsing for his flock, and honey for the bees,—

“Salices, humilesque genistæ,
Aut illæ pecori frondem, aut pastoribus umbram
Sufficiunt, sepemque satis et pabula melli.”

GEORGICON, II.

MAGNOLIA.†

Natural Order, MAGNOLIACEÆ, (JUSSIEU.) Linnæan Classification, POLYANDRIA, POLYGYNIA.

Calyx of 3 deciduous petaloid sepals. *Corolla* of from 6 to 12 petals. *Stamens* numerous as well as the pistils. *Carpels* disposed in an imbricated cone, 1 to 2-seeded, opening by the dorsal suture. *Seeds* pulpy, red, suspended, when ripe hanging out of the carpel by a long umbilical thread composed of spiral vessels.

Trees and shrubs with large entire alternate deciduous leaves, and solitary terminal large and usually odoriferous flowers. Chiefly natives of North America, China, and Japan.

LARGE FLOWERED MAGNOLIA.

MAGNOLIA GRANDIFLORA, *Linn.* (MICH. Sylva, plate 51.)

IN the neighbourhood of Savannah in Georgia, and near New Orleans, this splendid tree often presents an almost equal, smooth, columnar shaft of 60 to 80 feet elevation, and attains the height of 100 feet or upwards, with a graceful high and spreading summit. On the trunk of this species, near Savannah, I observed large quantities of the parasitic air plant, *Epidendrum conopseum*, and it appeared there to grow on no other tree. According to Wm. Bartram, who saw the species so

† Named by Linnæus in honour of Pierre *Magnol*, a botanist of Montpellier.

abundant in his tour in Florida, its summit forms a perfect cone, rising from a straight clear trunk, resembling a beautiful column, and, from its dark foliage "silvered over with milk-white flowers," it is seen at a great distance. The succession of flowers is also long continued, in favourable situations from May to August. Though confined very much to the neighbourhood of the sea-coast, it extends westward in Georgia as far as Milledgeville; and I met with it in Alabama, on the banks of Utchee creek, about 12 miles from Columbus in Georgia, and afterwards in other parts of that State down to West Florida. It is known to the Creeks by the native name of *Toola*.

In the new edition of Duhamel, we have the following account of its introduction into France. There is at Maillardière, about five miles from Nantes, a fine Magnolia, which was brought from the banks of the Mississippi, in 1732, and planted in a poor soil. It grew there neglected for more than 30 years, till M. Bonami, a physician of Nantes and professor of botany there, recognised this beautiful tree to be the *Magnolia grandiflora*; and at the meeting of the States of Bretagne in September, 1760, in Nantes, he presented to the Princess of Rohan-Chabet, a fine branch of this Magnolia in flower, which became a subject of conversation and interest to all assembled. Louis XV. possessed several small plants of this species in his garden at the Petit Trianon, but they did not thrive; and, having heard of a Magnolia 35 to 40 feet high, which every year was covered with fine flowers of a delicious perfume, he sent two of his gardeners to ascertain if it was possible to transport this tree to Versailles; and, above all, should they do so, if it would be certain to grow. They saw the tree, and, being of opinion that it would not survive removal, it was suffered to remain in its place. It was at that

time from 35 to 40 feet high; but, during the troubles of the civil war of La Vendee, it was mutilated, and lost most of its branches. Afterwards, the burning of the house near which it was planted having damaged its head, the branches were cropped down to the trunk, and it again shot out with vigour, but the young shoots, not having had time to ripen, were destroyed by the frost; notwithstanding this severe check, it again recovered, and afterwards became a fine tree, between 25 and 30 feet high, with a large well proportioned head, and a trunk of four feet in circumference, the lower branches sweeping the ground, and the whole tree producing annually from 350 to 400 large, elegant, and fragrant flowers. The seeds, however, never arrive at perfect maturity, although the fruit attains its full size, and remains upon the tree till the following spring. This tree still exists, and is now upwards of 30 feet high, and more than 100 years of age.

At Caserta, in the neighbourhood of Naples, this tree has attained the height of nearly 60 feet. In this climate they also ripen seeds freely.

LONG-LEAVED MAGNOLIA.

MAGNOLIA MACROPHYLLA, *Michaux*, Flor. Bor. Am., vol. 1. p. 327. MICH. Sylva, pl. 57.

THE principal locality of this fine species has been for many years confined to the vicinity of Lincolnton, North Carolina, 10 miles S. E. of the town, near or on the estate of a man named Smith. The trees occupy the banks of a small stream, and are chiefly dispersed

over its acclivities, in a rich loamy soil. In Tennessee, near the Cumberland river, I afterwards saw a few small trees of this species; but in the winter of 1830, in an extensive tour which I made through the interior of the Southern States, I met with abundance of the *Magnolia macrophylla*, 70 miles from Tuscaloosa, on the banks of the Coosa in Bibb county, Alabama, growing often in the same bottom lands as the *Illicium floridanum*. To give me some idea of the magnitude of the Magnolias growing in this vicinity, a farmer told me some trunks produced 16 rails to the cut, and that the trees were two feet in diameter; but as timber it was little esteemed, not enduring long in the air. To the town of Cahaba, I still saw the Magnolia, and afterwards, in the immediate vicinity of Tuscaloosa, on the road down the banks of the Black Warrior, towards Florida, I observed this species in the greatest abundance, often as much as 60 feet in height; but being the depth of winter, I, of course, could form no adequate conception of the splendour of its appearance when in vigorous vegetation.

EAR-LEAVED MAGNOLIA, (*magnolia auriculata*, LAM. *Long-leaved Cucumber Tree*, MICH. Sylva, pl. 56.) In Bartram's Garden, at Kingsessing, in this vicinity, there is a tree of this species, 70 or more feet high, and with a trunk of the diameter of two to three feet.

UMBRELLA TREE, (*Magnolia tripetala*.) According to Professor Torrey, no person since Michaux has found this tree in any part of the State of New York.

TULIP TREE, (*Lyriodendron tulipifera*, LINN.) Respecting the northern limits of this tree, G. B. Emerson, Esq., informs me, "I have found a single tree of *Lyriodendron* in Norfolk county, Massachusetts. It is plentiful on Westfield river."

Tennessee,
saw a few
of 1830, in
interior of
of the *Mag-*
a, on the
a, growing
floridanum.
of the *Mag-*
I saw some
at the trees
was little
the town of
wards, in the
oad down
rida, I ob-
e, often as
of winter,
ion of the
is vegeta-

lata, LAM.
(p. 56.) In
s vicinity,
feet high,
ree feet.
According to
has found
k.

(ENN.) Re-
Emerson,
of Lyrio-
is plenti-




Drimophyllum pauciflorum.

California Bay tree

Drimophyllum pauciflorum.



paniciflora.



DRIMOPHYLLUM.†

Natural Order, LAURINEÆ. (Ventenat.) *Linnean Classification*, ENNEANDRIA, MONOGYNIA.

Flowers HERMAPHRODITE.—*Perigonium* 6-cleft, subcampanulate, the segments spatulate-linear, equal, deciduous, the base persistent. *Stamina* 9, in 3 series, all fertile, filaments short, the 3 of the inner series with a pair of roundish large sessile glands, near the base of each. *Anthers* elliptic-ovate and similar, all opening on the inner side, 4-celled, the cells equal and parallel, with all the valves ascending. *Ovarium* 1-celled, with one ovule. *Style* short; *stigma* somewhat depressed, capitate. *Berry* 1 seeded

A sempervirent, small tree of Upper California, with alternate, lanceolate, minutely reticulated, pungently aromatic, smooth leaves. Flowers small, yellow, smooth, in small terminal naked clusters. (Nearly allied to *Ocotea* of Aublet, but with hermaphrodite flowers, similar and uniformly introrse anthers, and a deciduous perianth. The leaves are also naked, without pennate nerves, and the inflorescence not in panicles. It is also nearly allied to *Aperiphracta* of Nees, but with flowers of a very different habit, and, with those genera, belongs to the tribe OREODAPHNEÆ.)

CALIFORNIAN BAY TREE.

DRIMOPHYLLUM PAUCIFLORUM.

OCOTEA SALICIFOLIA? Kunth, Synops. vol. 1. p. 458.

THIS is a very elegant evergreen tree of Upper California, growing round Sta. Barbara, 20 to 25, or 30

† From *δρμος*, acrid to the taste, and *φυλλον*, a leaf.

feet high, with erect, terete, and smooth branches. The wood is white and rather soft. The leaves are alternate, evergreen, coriaceous, perfectly smooth, three to four inches long, and $\frac{3}{4}$ to $1\frac{1}{4}$ inches wide, lanceolate pointed, but obtuse, entire, with very indistinct slender lateral nerves, and strongly but minutely reticulated above; the footstalks are about two to three lines long. The odour and taste of the leaves are very aromatic, the latter so much so as to be quite pungent, even more so than the leaves of the Bay, and they are employed as condiments by the inhabitants. The flowers are in small contracted clusters, at first surrounded with bud scales, which are caducous, but not in the form of an involucre. The flowers are about four or five together, on pedicels nearly as long as themselves. The perianth is yellowish, funnel-formed, and somewhat spreading, deeply 6-cleft, the segments linear-spathulate and smooth, a little pubescent within towards the base. Stamens 9, with short and broad filaments, the anthers oblong, 4-celled, all opening from within, the cells parallel and nearly all equal, with the valves ascending, the three innermost each furnished towards the base of the filament with two large reniform, sessile glands. The perianth is deciduous, the base alone being persistent, and enlarging with the 1-seeded berry. The perfect fruit I have not seen.

PLATE XXII.

A branch of the natural size. *a.* The flower enlarged.

UMBELLULARIA,

(NEES ab Esenb.)

Natural Order, LAURINEÆ. Linnæan Classification, ENNEANDRIA, MONOGYNIA.

Flowers HERMAPHRODITE.—The *perianth* deeply 6-parted and shortly companulate, the segments equal and deciduous. *Stamina* 12, the 9 exterior fertile, interior sterile, the 3 fertile innermost ones each with a pair of large glands covering the back of the filament. *Anthers* 4-celled, those of the 3d series with the two upper cells introrsely opening, the lower extrorsely and lateral; the sterile filaments thread-shaped. *Stigma* peltate, subrepand. *Berry* 1-seeded, seated on the cup-shaped base of the perianth.

A tree of Upper California, with alternate pennately nerved leaves. The flowers aggregated in axillary umbellated clusters, inclosed by the involucre, the scales of the involucre broadish, alternate and approximate, after the opening of the flower, caducous.

CALIFORNIAN UMBELLULARIA.

UMBELLULARIA CALIFORNICA, *hermaphrodita, foliis perennantibus oblongo-lanceolatis vix acutis penninerviis reticulata-venosis glabris, pedunculis axillaribus simplicibus, floribus pluribus, umbella subcapitata, pedunculo folio brevioribus.*

TETRANTHERA? CALIFORNICA, HOOKER. et ARN. in Botan. Beech. Voy., p. 159. HOOKER. Flor. Bor. Amer., 2. p. 137.
LAURUS REGIA? DOUGLAS. Journal, in Hook. Compan. Bot. Mag., vol. 2.

THIS splendid evergreen tree was discovered on the

north-west coast, in Upper California, by Mr. Menzies, who first made known to botanists the vegetable treasures of that interesting and then unexplored region. Douglas afterwards found it in nearly the same country, south of the Columbia or Oregon, and adds, that it attains the height of from 40 to 120 feet, with a diameter of from 2 to 4 feet. It commences at the southern limit of the prevailing pine and fir forests, which line the wastes of Oregon. The foliage gives out, when bruised, a most powerful camphorated odour, which from its pungency is capable of exciting sneezing. Flowering specimens of this interesting tree were in Douglas's collection from California. It is to be regretted, however, that no detailed description nor figure is given; and I had not the good fortune to meet with it myself.

SASSAFRAS (*Laurus Sassafras*. LINN). The inhabitants of North and South Carolina distinguish two kinds of Sassafras, the Red and the White. The Red or true *L. Sassafras* I referred (in the Genera of North American plants, vol. 1. p. 259, 260.) to a sub-genus *Euosmus*, embracing also the following variety, which I then considered as a species, by the name of *L. (Euosmus) albida*. It is distinguishable from the Red by having the buds and twigs smooth and glaucous; its leaves are also smooth and thin, and the veins almost obsolete beneath, the petiole is apparently longer. The root is much more strongly camphorated than that of the Red sort, and is nearly white. It is better calculated to answer as a substitute for ochra (*Hibiscus esculentus*) than the common kind, as the buds and young branches are much more mucilaginous. It is abundant in North and South Carolina, from the Catawba Mountains to the east bank of the Santee, growing with the common kinds.

From the present order of plants we derive the Cinnamon, Cassia and the Camphor. Several species afford the Cinnamon of commerce, and the *Laurus Quiros* produces that of Peru. The Cinnamon of Santa Fe de Bogota is afforded by *Laurus Cinnamomoides*. A great deal of the finest camphor of India, however, is the product of the *Dryobalanops Camphora*. The volatile oil obtained from some species of *Laurus* found in the vast forests between the Oronoko and the Parime, is produced in great abundance by merely making an incision into the bark with an axe, as deep as the liber or young wood. It gushes out in such quantities that several quarts may be obtained by a single incision. It has the reputation of being a powerful discutient.

THE LINDEN, OR LIME TREE.

Natural Order TILIACEÆ (*Jussieu*). Linnæan Classification,
POLYANDRIA, MONOGYNIA.

TILIA.† (LINN.)

Sepals 5. *Petals* 5. *Stamens* numerous, disposed more or less in 5 clusters, the central tuft (chiefly in the native species) transformed into a petal. The *ovary* globular, villous, and 5-celled, each of the cells bearing 2 ovules. *Capsule* ligneous, globular, by abortion only 1-celled, with 1 or 2 seeds. *Cotyledones* sinuate.

Trees of Europe and North America with alternate dilated or cordate leaves, oblique at the base, serrated on the margin, and with a tough and fibrous bark; stipules caducous. The flowers disposed in flattish pedunculated clusters (or eymes), and with the peduncle curiously adnate for a great part of its length to a large membranaceous, linear bracte. The rest of this family of plants are nearly all tropical productions.

LARGE LEAVED LINDEN, OR LIME.

TILIA HETEROPHYLLA, *foliis ovalis, argute serratis, basi nunc cordatis, nunc oblique aut æqualiter truncatis, subtus tomentosis; nucis pisiformi*. VENTENAT, Mem. de l'Institut., tom. 4. p. 16. pl. 5. PURSH, Flor. Bor. Amer., vol. 2. p. 363. NOUVEAU DUHAMEL, vol. 1. p. 229. DECANDOLLE, Prod. vol. 1. p. 513.

TILIA HETEROPHYLLA, leaves glabrous and deep green above,

† An ancient Latin name, probably from the Greek *τιλιάς*, the ELM.

REE.

Classification,

more or less
(five species)
villous, and
Capsule lig-
or 2 seeds.

te dilated or
margin, and
The flowers
) and with
s length to a
his family of


ME.

erratis, basi
utis, subtus
e l'Institut.,
l. 2. p. 363.
e, Prod. vol.

green above,

ek $\pi/\delta\alpha\epsilon\zeta$, the






Tilia heterophylla.

Large leaved Linden

Tilial heterophylla



very white and velvety-tomentose beneath, the veins dark coloured and nearly glabrous, with coarse mucronate serratures; petals obtuse, crenulate; staminodia [inner petals] spatulate, entire, style hairy at the base. TORREY and GRAY, Flor. North Amer., vol. 1. p. 239.

TILIA ALBA. *Smith's Insects of Georgia*, vol. 1. p. 21. t. 11?

This is one of the rarest and most ornamental trees of the whole genus, and as far as my own observations go, it is almost wholly confined to the shady forests of the Ohio and its tributary streams, to which Pursh also adds the banks of the Mississippi; Torrey and Gray received it likewise from the neighbourhood of Macon, in Georgia, where it was collected by our late mutual friend and excellent observer Doctor Loomis. In descending the Ohio, late in autumn, (about the year 1816,) I got out of the boat in which I was descending to walk round Le Tart's Rapids above Cincinnati, here I observed almost an exclusive forest of this fine Linden, on a rather elevated alluvial platform, in a light, rich calcareous soil. Most of the trees were tall and rather slender, 60 to 80 feet in height, and the ground was thickly strewn with their large and singular leaves, almost as white as snow beneath. According to the herbarium of Mr. Schweinitz, it exists also in Virginia, probably on the borders of the streams which flow into the Ohio, near Pittsburgh; and according to Doctor Short, of Lexington, Kentucky, it forms in his vicinity one of the largest forest trees in the rich lands there. Decandolle speaks of having received a specimen of some very similar species from Mexico. It does not yet appear to have been introduced into Europe, though it is properly described in the new Duhamel, probably from Ventenat's essay, as the leaves are said to be snow-white beneath.

The young branches are purplish and somewhat glau-

cous. The largest leaves I have seen are about 6 or 7 inches long, and 3 to 5 broad. In the young state, the white pubescence beneath is most conspicuous when the leaves are thinly covered, the hairs are stellate, the serratures are strong and sharp, with acuminate rigid points, the upper surface is dark green; the base of the leaf varies considerably, sometimes it is sinuated, at other times perfectly flat and truncated; the leaves are always very oblique at the base. The flowers are somewhat larger than those of *T. americana*, and the fruit is villous, nearly spherical, and certainly always without any ribs.

The *TILIA ALBA*, White Lime of Michaux, plate 132, not being the *T. alba* of Kitaibel and Aiton, (Hort. Kew. l. c.) which is a native of Hungary, it is necessary to change his name, and we propose to call it *TILIA MICHAXII*, (Michaux's Lime,) if his plant should indeed prove to be any thing more than a smoother variety of our *T. heterophylla*.

PLATE XXIII.

A branch of the natural size. *a.* The fruit. *b.* The flower.

GENERAL OBSERVATIONS.

The Lime has long been a favourite tree for avenues and public walks; it is planted in the streets of some of the principal towns of France, Holland and Germany, and it is used for forming avenues both on the continent of Europe and in Great Britain. It has of late years been much planted along the streets in several towns and cities of the United States, but in Philadelphia it is so much attacked by insects that it probably will not long survive. The species employed for this purpose appears

to be principally the European, while the native kinds being more hardy and vigorous, ought to have the preference, particularly the present species, (*T. heterophylla*), which in a good soil becomes a large tree, and is at the same time splendidly ornamental. The insect that devours the leaves of the Linden appears to be a moth,† which suspends its cocoons at the ends of the twigs of the trees it has stripped; these ought carefully to be removed and destroyed, by which means the evil, if not wholly cured, would be decidedly mitigated.

The Dutch plant the Lime in towns, along their widest streets, and by the sides of their canals; and the whole country is thus perfumed by their flowers during the months of July and August; they likewise afford an ample repast for the bees.

The wood of the European Lime tree is of a pale yellow or white, close-grained, soft, light, and smooth, and not liable to be attacked by insects. It is used by pianoforte-makers for sounding-boards, and by cabinet-makers for a variety of purposes. It is turned into domestic utensils of various kinds, carved into toys, &c. The most elegant use to which it has been applied is for carving, for which it is superior to any other wood. Many of the fine carvings in Windsor Castle, St. Paul's, Trinity College Library at Cambridge, and in the Duke of Devonshire's mansion at Chatsworth, from the hand of the celebrated Gibbons, are of this wood. It makes excellent charcoal for gunpowder. Baskets and cradles were formerly made from the twigs. The leaves are also employed as fodder for cattle in Europe. It is in Russia and some parts of Sweden that the well-known bass mats are formed from the inner bark of this tree. The bark stripped from young trees of six


† A species of *Oiketicus*.

inches to a foot in diameter are selected for this purpose. These strips are steeped in water till the bark separates freely into layers, it is then taken out and separated into strands, which are dried in the shade, and afterwards manufactured into the mats so much used by gardeners and upholsterers, and for covering packages. The fishermen of Sweden make fishing-nets of the fibres of the inner bark, formed into a kind of flax; and the shepherds of Carniola even weave a coarse cloth of it, which serves them for their ordinary clothing. The whole plant abounds with mucilage, the sap, like that of the Maple, affords a considerable quantity of sugar, and the honey produced by the flowers is considered superior to all other kinds for its delicacy, selling at three or four times the price of common honey; in Europe it is used exclusively in medicine, and for making some particular kinds of *liqueurs*, especially Rosolio. This Lime tree honey is only to be procured at the little town of Kowno, on the river Nieman, in Lithuania, which is surrounded by an extensive forest of Lime trees. The triturated fruit produces also a paste very similar to that of Cocoa. During the taste for grotesque decorations, the Lime, like the Yew, was cut into various imitative forms, and in some of the public gardens of recreation round Paris and Amsterdam there are very imposing colonnades, arcades, walls, pyramids, and other architectural looking masses formed of this tree.

The European Linden attains a height of upwards of 100 feet, and grows with vigour for several centuries. In Switzerland there are some very large and ancient Lime trees: one mentioned by Decandolle the younger, near Morges, has a trunk of 24 feet 4 inches in circumference; another near the great church at Berne, which was planted before the year 1410, is 36 feet in girth.

s purpose.
separates
rated into
fterwards
gardeners
es. The
fibres of
and the
oth of it,
ng. The
ke that of
ngar, and
superior
three or
rope it is
ng some
o. This
ttle town
which is
s. The
r to that
orations,
mitative
creation
mposing
r archi-

wards of
enturies.
ancient
ounger,
in cir-
Berne,
fect in



J. Berria

Rhizophora Americana.


American Mangrove

Rhizophore d'Amerique



Thoria

phore d'Amérique



M A N G L E .

Natural Order, RHIZOPHOREÆ, (R. BROWN.) *Linnean Classification*, DODECANDRIA, MONOGYNIA.

RHIZOPHORA.† (LINN.)

Tube of the *calyx* obovate, coherent with the ovary, the border divided into 4 oblong, persistent segments. *Petals* 4, oblong, emarginate, coriaceous, conduplicate, before expansion embracing the alternate stamens, the margins each with a double row of long woolly hairs. *Stamens* twice as many as the petals; anthers nearly sessile, large, linear-oblong. *Ovary* 2-celled, with 2 ovules in each cell. *Style* conical, short, 2-furrowed; *stigma* 2-toothed or bifid. *Fruit* ovate or oblong, crowned near the base with the persistent segments of the calyx, longer than the tube, at length perforated at the apex by the radicle of the germinating embryo.

Maritime trees of the tropics with entire opposite leaves, and axillary flowers.

AMERICAN MANGLE, OR MANGROVE.

RHIZOPHORA AMERICANA, *foliis obovato-oblongis obtusis; pedunculis trichotomis petiolo longioribus, stylis subulatis bifidis, fructibus subulato-clavatis obtusis.*

RHIZOPHORA *mangle*. JACQUIN, Amer., p. 141. t. 89. BROWN, Jam. p. 211. DECAND. prod. vol. 3. p. 32. NUTT. Florid. pl. Sill. 5. p. 295. TOR. and GRAY, vol. 1. p. 484. (not of LINN.)

† The name, from *ρίζα*, a *root* *φέρω*, *to bear*, in allusion to the seed germinating before it falls from the branches.

Candela americana foliis laurinis. CATESBY'S Carol., vol. 2. p. 63. t. 63.

Mangle aquatica, foliis subrotundis et punctatis. PLUMIER, Gen. p. 13. SLOANE, Jam. p. 155. Hist. vol. 2. p. 63.

Margue Guapariba. PISO'S Brazil, l. 4. c. 87. R.

THIS tree is found in the maritime swamps of Louisiana and East Florida, and along the coast of Texas, not uncommon. The Mangrove, like the famous Banyan Fig, sends out innumerable roots from the fusiform fruits which terminate its branches into the surrounding marshes, so that after a while a single tree becomes, as it were, the parent of a whole forest of several miles in extent; and growing well even into the salt water, it is not unfrequent to see their branches loaded with oysters (the *Ostrea folium*) of an exquisite flavour. Those thickets likewise afford a resort for various kinds of sea-fowl, and fringing the margin of the ocean and the salt pools with their spreading summits, they give a peculiar feature to the tropical landscape, but at the same time afford shelter to clouds of musquetoos. The bark and fruit are useful for tanning; the flower, according to Loureiro, dyes a very durable black, and, according to Sloane, affords a material for ink.

The Mangrove of the West Indies and tropical America becomes a tree about 40 to 50 feet high, and 2 to 3 feet in diameter, with a ferruginous bark and white wood of no great value, except for fuel, yet, according to Sloane, the wood is good for building and shingles. The wood of that of India, as described by Roxburgh in his *Flora Indica*, is of a dark reddish colour, hard, and durable.

The Mangrove is not very tall, but very branching, the branches almost always opposite, elongated and pendant. When touching the soil, they strike root and become new trees, which re-multiply themselves in the

same manner, thus forming an almost impenetrable barrier on the borders of the sea.

The leaves are opposite, entire, coriaceous, at first folded inwards, with caducous stipules between the petioles. The flowers are pale yellow, the segments of the calyx lanceolate. The anthers are subulate; the margin of the petals pilose. The style bifid, with the divisions rather long and subulate. The verdure of the Mangrove is dark and gloomy, and the whole tree, inhabiting a region of desolation, presents an aspect of sadness.

The most extraordinary plant of this, or rather a nearly allied genus, is the *Rhizophora gymnorhiza* of Linnæus, (now *Brugiera*.) This tree grows commonly in the maritime marshes of India, and the branches of its numerous roots ascending into the air produce the appearance of a large umbrageous tree, as it were, on stilts, or as Roxburgh says, supported in the air on a circle of converging hop-poles. The fruit, the leaves, and the bark of this species are also said to afford food to the native inhabitants. A figure of it is given by Rumphius, vol. 3. t. 68, and by Rheede, in the Flora Malabarica, vol. 6. tab. 31, 32.

PLATE XXIV.

A branch of the natural size. a. The fruit.

G U A V A .

*Natural Order, MYRTACEÆ. Linnæan Classification,
ICOSANDRIA, MONOGYNIA.*

PSIDIUM,† (LINN.)

Calyx-tube (or external germ) ellipsoid or obovate, often contracted at the summit; the border at first undivided and ovate while in flower, afterwards 1 to 5-cleft. *Petals* 5. *Stamens* very numerous, distinct. *Style* filiform: *stigma* capitate. The *ovary* with from 5 to 20 cells, some of them abortive, each cell subdivided by the interposition of a placenta resembling a dissepiment. *Ovules* numerous, horizontal. *Fruit* a many-seeded berry, coated with the adhering tube of the calyx and crowned by its persisting lobes. The *seeds* scattered through the pulp in the ripe berry, having a bony or hard shell. The *embryo* curved in a half circle round the protruded base of the testa. *Cotyledones* minute, the radicle rather long.

Trees or shrubs chiefly indigenous to the intertropical regions of America, with opposite, entire, impunctate, feather-nerved leaves. Peduncles axillary, 1 to 3 flowered, each flower with a pair of bractes. The flowers white.

FLORIDA GUAVA.

PSIDIUM BUXIFOLIUM, *glabrum, ramulis teretibus, foliis parvulis coriaceis cuneato-obovatis obtusis subsessivous margine revolutis, pedunculis solitariis brevissimis unifloris, fructu pyriformi.*

For a knowledge of this interesting tree or shrub we

† One of the Greek names for the Pomegranate. *Guava* is a corruption of the American aboriginal name of *Guayaba*.

fication,

, often con-
d and ovate

Stamens
a capitate.

m abortive,
enta resem-

tal. *Fruit*

ube of the

eds scatter-

a bony or

round the

the radicle

ical regions

her-nerved

wer with a

bus, foliis

bsessitibus

nis uniflo-

shrub we

Guava is a

uba.






Pisidium buxifolium.

Florida Guava.

Coyavter de la Floride



are indebted to the late indefatigable Dr. Baldwin, who met with it in some part of East Florida, near the river St. Johns. To show how very unlike this species is to all the others known, it was hastily marked by Mr. Schweinitz in his herbarium, (of which the specimen forms a part,) "*Quercus virens*," and, at the first hasty glance some resemblance may be traced with the Live Oak in the leaf and twig, but, of course, the presence of the fruit at once dispels the illusion.

I have seen but the single specimen now figured, and would recommend its examination to some future traveller. The twig is round, covered with a grey bark, and at near distances marked with the cicatrices of opposite fallen leaves. The leaves on the upper branchlets are crowded together in opposite pairs, of a very thick, opaque, rigid consistence, and appear to be sempervirent, they are perfectly smooth on both sides, paler beneath, dark green above, cuncate-obovate, obtuse, sometimes with an attempt at a very short and blunt acumination, with the margin reflected, and beneath marked with numerous approximating feathered nerves; they are from 1 inch to $1\frac{1}{2}$ inches long by $\frac{1}{2}$ to $\frac{3}{4}$ of an inch wide. The peduncles are axillary and solitary, very thick in the fruit-stalk and scarcely two lines long. The flowers I have not seen. The berry is blackish-purple, pear-shaped, about the size of a cherry, and appears to have been succulent, as usual; internally it is filled with horizontal rows of flat, subreniform, pale brownish bony seeds, with a narrow embryo curved into the form of a horse-shoe. The cotyledones are very small, and in the seed of a bright waxy-yellow. This species is very nearly allied to the Purple-fruited Guava, (*P. Cattleianum*,) scarcely differing in any thing but the smallness of the leaves and the pyriform fruit, though the leaves of the Purple Guava, besides being much

larger, are also pubescent when young. Most of the species of this genus are cultivated in the tropics for their fruit. The *P. pyriferum*, or common Guava, bears a fruit about the size of a hen's egg, yellowish, with a peculiar odour; the pulp is rather firm, flesh-coloured, sweet, agreeable, and aromatic. In the West Indies it is highly esteemed by all classes, being eaten raw, as a dessert, or formed into an excellent sweetmeat and jelly.

Of the fruit of the Purple Guava, to which ours is so closely related, Lindley remarks, "The excellent flavour of its fruit, which is very like that of strawberries and cream, is far superior to either *P. pyriferum*, *pomiferum*, or *polycarpon*." Mr. Sabine remarks of the fruit of this species, "that it is juicy, of a consistence much like that of a strawberry, to which it bears some resemblance in flavour."

What the present species may become, when cultivated, remains to be proved, but in a genus so generally interesting for their fruit, the experiment is worth making, when an opportunity may offer. Probably, Dr. Baldwin found it growing near or above New Smyrna, as he did not go much farther into the interior of East Florida.


PLATE XXV.

A branch of the natural size in fruit.

Most of the
tropics for
guava, bears
ish, with a
h-coloured,
st Indies it
n raw, as a
tmeat and

ours is so
ent flavour
berries and
pomiferum,
fruit of this
ch like that
mblance in

when culti-
o generally
th making,
Dr. Bald-
Smyrna, as
or of East



Calyptroanthus chytroculia


Forked Calyptroanthus

Calyptroanthus chytroculia

PLATE



Aranthe chytocula



CALYPTRANTHES,†

(SWARTZ.)

Natural Order, MYRTACEÆ. Linnæan Classification,
ICOSANDRIA, MONOGYNIA.

Tube of the *calyx* obovate, with the border entire, when flowering bursting circularly in the form of a lateral, and at length, deciduous lid. *Petals* none, or 2 or 3 and minute. *Stamens* many. *Style* 1; stigma simple. *Ovary* 2 to 3-celled, the cells 2-seeded. The *berry* by abortion 1-celled, 1 to 4-seeded.

Small trees of the West India Islands and of Brazil, the leaves with pinnated veins. Flowers small and numerous, usually in axillary or terminal panicles.

FORKED CALYPTRANTHES.

CALYPTRANTHES CHYTRACULIA, *arborea, foliis ovatis apice attenuatis rigidiusculis demum glabris, pedunculis axillari-terminalibus trichotomis paniculatis floribusque rufo-velutinis.*—DECAND. Prod. vol. 3. p. 257.

C. CHYTRACULIA, *arborea, pedunculis terminalibus trichotomis tomentosis, foliis ovatis apice attenuatis.* SWARTZ, Prod. p. 79. Flor. Ind. Occid., vol. 2. p. 921.

MYRTUS CHYTRACULIA, *pedunculis dichotomis paniculatis tomentosis, foliis geminis subovatis terminalibus.* LINN. Amœn. Academ., vol. 5. p. 398. SWARTZ, Observ. p. 202.

Chytraculia arborea, foliis ovatis glabris oppositis, racemis terminalibus. BROWN, Jamaic. p. 239. t. 37. fig. 2.

EUGENIA PALLENS? Poiret. Suppl., vol. 3. p. 122.

THIS plant forms an elegant and curious small tree,

† The name from *καλυπτρα*, a veil, and *αρβου*, a flower, in allusion to the operculid form of the calyx.

with hard wood, and in Jamaica is accounted an excellent timber, but the trunks seldom exceed 14 or 15 inches in diameter. In Jamaica it is found in the dry mountain lands; it is also indigenous to the islands of St. Thomas and Guadaloupe, and it has now also been found on Key West by Dr. Blodgett.

The branches appear to be covered with a grey and smooth bark. The leaves, when in bud, as well as the young branches, flower-stalks and calyx are clad with a short, soft, ferruginous down, which from the leaves, as they advance in their development, wholly disappears, they are of a lanceolate-ovate form, narrowed into a short petiole below; above acuminate but obtuse; beneath they are distinctly pennate-nerved, and too opaque to admit the light through the resinous glands with which they are nevertheless provided; they are about two inches long by an inch in width. The flowering panicles are trichotomous, usually terminal, and considerably ramified. The flowers are small and whitish, from the colour of the stamens. The calyx is ferruginous and tomentose, formed of a small obovate even cup, the whole border separating in a circular manner flies over to one side, in the form of a rounded petal, from whence issues the numerous filiform stamens with small whitish anthers. The germinal fruit appears small, dry, and tomentose, but I am unacquainted with it in a ripe state.

PLATE XXVI.

A branch of the natural size. *a.* A flower magnified, showing the lateral adherence of the lid of the calyx.

an excel-
14 or 15
in the dry
islands of
also been

grey and
ell as the
ad with a
leaves, as
disappears,
ed into a
; beneath
opaque to
ith which
out two
; panicles
siderably
from the
ous and
cup, the
flies over
n whence
l whitish
dry, and
n a ripe

, showing



E.D. Long del.


Sinclair's lab.

Small-leaved Eugenia. **Eugenia dichotoma.** *Tambosier dichotome.*



Sinclair's Lith.

hosier dichotome.



EUGENIA.†

(MICHELI, LINN.)

*Natural Order, MYRTACEÆ. Linnæan Classification,
ICOSANDRIA, MONOGYNIA.*

The tube of the *calyx* roundish, with the border deeply 4-parted. *Petals* 4. *Stamens* many, free. Ovary 2 to 3-celled, the cells containing several ovules. *Berry* sub-globose, crowned with the persisting calyx, when mature 1 or rarely 2-celled. *Seeds* 1 or 2, roundish and large. The *embryo* pseudo-mono-cotyledonous, the *cotyledones* very thick and wholly blended together, the radicle more or less distinct and very short.

These are trees or shrubs, mostly indigenous to the Caribbean Islands, or the warmer parts of America. The leaves and inflorescence are very similar to those of the Myrtles.

SMALL-LEAVED EUGENIA.

EUGENIA DICHOTOMA, *pedunculis axillaribus oppositis et subterminalibus folio longioribus bifidis aut bis bifidis, floribus in dichotomiis sessilibus cæteris pedicellatis, foliis elliptico-lanceolatis basi attenuatis pellucido-punctatis, adultis glabris, junioribus utrinque ramulis calycibusque pubescentibus.*—DECAND. Prod., vol. 3. p. 278.

MYRTUS dichotoma, Vahl! MSS. POIRET, Supplem., vol. 4. p. 53? *β. FRAGRANS, foliis ovatis glabris.* *EUGENIA FRAGRANS.*

WILLD. Sp. Pl., vol. 2. p. 964. Bot. Magaz., t. 1242. *E. montana*, AUBL. Guian., vol. 1. p. 495. t. 195?

EUGENIA divaricatu, Lam. Encyc., p. 202.

THIS elegant and fragrant species of *Eugenia* resem-

† So named in honour of Prince Eugene of Savoy, who was a protector and encourager of botany, and possessed a botanic garden.

bling a Myrtle, becomes, at Key West, according to Dr. Blodgett, a tree. It is also indigenous to the islands of St. Domingo and Cuba, where specimens have been collected by Poitcau and La Sagra. The variety *fragrans*, for such I must consider it, is a native of the high mountains in the southern part of Jamaica and Martinique, and if the same with Aublet's *E. montana* it is also a native of Guiana. The *E. fragrans* has many years since been collected by Dr. Baldwin, in the vicinity of New Smyrna in East Florida.

The wood of *E. divaricata*, according to Lamarck, is hard, close grained and reddish, and is much esteemed for articles of furniture. The wood of the Florida tree is exactly similar, while that of *E. montana*, according to Aublet, is hard, compact and white.

The branches of the plant now figured are covered with a smooth light grey or silvery bark, and at the summits are crowded with small shining almost opaque leaves, but yet interspersed with the usual resinous vesicles of the genus; they are from an inch to an inch and a half in length, and about three-quarters of an inch in breadth, mostly elliptic, or elliptic-oblong, and always narrowed below; sometimes they are nearly lanceolate and obtuse at the point; scarcely any veins are visible on either side, but the mid-rib is prominent beneath. The young leaves, buds, peduncles and calyx are clothed with a close, short, hoary pubescence, which in the variety *fragrans* is much less distinct or almost wanting. The peduncles are axillary, coming out towards the summits of the branches, and are of various lengths, sometimes only a little longer than the leaves, at other times crowded into trichotomous branchlets, two or three times longer than the leaves; in their most simple form, except by the abortion of the lateral buds, they terminate in three flowers, the central one sessile in the fork,

and the lateral ones are longish diverging pedicels; at other times the peduncles are twice trifid, or even more ramified, and lengthened out very much in the progressive ripening of the fruit. The segments of the calyx are always four, broad and rounded, covered with resinous cists or vesicles, and pubescent or ciliate on the margins. The petals are likewise rounded or concave, whitish, with a tinge of red. The stamens are numerous. Style simple and subulate. The berry at length only 1-seeded. There are a pair of minute subulate bractes under the base of each flower bud, but so deciduous that they are seldom to be seen.

One of the specimens of the variety *fragrans* from New Smyrna has very slender twigs; and on the same specimen there are obtuse and very *sharply acute* leaves. In this also the peduncles are chiefly axillary. This plant is nearly as fragrant as the common Myrtle.

PLATE XXVII.

A branch of the natural size *a*. The flower a little enlarged.
b. The berry, of the natural size.

TALL EUGENIA.

EUGENIA PROCERA, *pedicellis unifloris axillaribus 2-4 confertis folio brevioribus sub flore bibracteolatis, foliis ovatis obtuse acuminatis ramisque glabris.* POIRET, Suppl. Encycl., vol. 2. p. 129. DECAND. Prod., vol. 3. p. 268.

MYRTUS PROCERA, *pedunculis confertis axillaribus unifloris, foliis ovatis acuminatis planis glabris, ramis virgatis, caule arboreo.* SWARTZ, Prodr. p. 77. Flor. Ind. Occident., vol. 2. p. 387. WILLD. Sp. pl., vol. 4. p. 968.

THIS is another plant with the aspect of a Myrtle, which becomes a tree and attains an elevation of 20 to 30 feet. It was discovered by Swartz in the forests of the interior of Hispaniola. It is likewise indigenous to the islands of Martinique, Sta. Cruz, and has now been found common on Key West by the same gentleman who met with the preceding species.


The wood appears to be white and close-grained. The twigs are clothed with a light grey, almost white and silvery bark, and are spreading and sometimes zig-zag. The leaves are on short petioles $1\frac{1}{2}$ to $2\frac{1}{2}$ inches long, by an inch to an inch and a half wide, ovate, acuminate, and obtuse, rather opaque, nearly scentless, though provided with the usual resinous vesicles, and from the bud they are perfectly smooth. The flowers are said to be fragrant, and come out on separate axillary peduncles, from 2 to 4 together; the peduncles at first are not more than 3 or 4 lines long, but grow out at length to the extent of half an inch. The segments of the calyx are 4, rounded and broad, rough, with aromatic vesicles, but smooth. The petals 4, are rounded and concave, slightly ciliated, and appear to

bus 2-4 *con-*
foliis ovatis
appl. Encycl.,

us unifloris,
is virgatis,
d. Occident.,

a Myrtle,
n of 20 to
e forests of
igenous to
now been
gentleman

se-grained.
most white
etimes zig-
2½ inches
de, ovate,
scentless,
sicles, and
ne flowers
separate
peduncles
but grow
The seg-
ad, rough,
tals 4, are
appear to






W. W. Wood del.

Eugenia procera.

Tall Eugenia.

lambosier elevé



have been reddish-white. Stamens numerous, the anthers whitish. The berry spherical, brownish-yellow, about the size of a grain of black pepper, studded over with numerous glands or aromatic cists, and crowned with the broad persisting border of the calyx: at first 2-celled, with several ovules, at length the berry is only 1-seeded, the seed large, with no distinct cotyledones. It flowers in April. The size of the leaves appear to vary, so that in some specimens they are uniformly only about half the dimensions we have given.

PLATE XXVIII.

A branch of the natural size and of the large leaved kind. *a*.
A cluster of the berries.

BOX-LEAVED EUGENIA.

EUGENIA BUXIFOLIA, *pedunculis axillaribus ramosis multifloris brevissimis, pedicellis sub flore bibracteolatis, foliis obovato-oblongis obtusis basi attenuatis opucis subtus punctatis margine subrevolutis.*—DECAND. Prod., vol. 3. p. 275. WILLD. Sp. pl., vol. 2. p. 960.

MYRTUS BUXIFOLIA, *racemulis brevissimis confertis axillaribus, foliis cuneatis oblongis obtusis convexiusculis.* SWARTZ, Prod. p. 78. Flor. Ind. Occident., vol. 2. p. 899. *M. monticola?* SWARTZ, Flor. Ind. Occid., vol. 2. p. 898.

MYRTUS AXILLALIS, *Poiret*, Dict. vol. 4. p. 412, (non Swartz.) *M. Poireti*, Spreng. Syst., vol. 2. p. 483.

THIS plant, also a native of Cuba, St. Domingo, and Jamaica, has been observed at Key West by Dr. Blodgett, where it is very common in sterile places, affecting the vicinity of the sea, and becoming a tree of about 20 feet in height, with a hard, white, close-grained wood. The bark is whitish-grey and even, the twigs are slender and chiefly clothed with leaves towards their summits, they are wedge-oblong, sometimes almost lanceolate, obtuse, and always narrowed below into a minute petiole, so that they appear to be nearly sessile, above of a darkish green and somewhat shining, beneath dull and paler, slenderly nerved beneath, somewhat opaque, punctate and slightly revolute on the margin, they are about $1\frac{1}{2}$ inches long by $\frac{1}{2}$ to $\frac{3}{4}$ of an inch wide. The flowers are very small, in axillary branching clusters of 3 to 7 together on the minute and very short bracteate raceme; there are 2 minute bracteoles under each flower; the calyx as well as the petals are studded with resinous glands, and the latter are more than twice the

mosis multi-
lutis, foliis
obtus punc-
l. 3. p. 275.

is axillari-
s. SWARTZ,
M. monti-

(on Swartz.)

ingo, and
Dr. Blod-
, affecting
about 20
ned wood.
s are slen-
their sum-
st lanceo-
a minute
ile, above
neath dull
at opaque,
, they are
ide. The
clusters of
bracteate
der each
dded with
twice the








Eugenia buxifolia

Box-leaved Eugenia.

Lambowiera feuilles de buis.



length of the calyx. The calyx, racemes, and minute branchlets are covered with a close brownish pubescence. The flowers are *polygamous*, on many specimens sterile, though furnished with the pistillum, and many of the flowering clusters are produced on the naked branches where they have been preceded by the former leaves. The berry is dark brown, covered with resinous glands or cists, about the size of a grain of black pepper, and when mature contains one or more (rarely 2) large seeds in 1 or 2 cells, with blended, inseparable cotyledones.

PLATE XXIX.

A branch of the natural size. *a.* A flower enlarged. *b.* The berry. *c.* A berry with 2-cells.

INDIAN ALMOND.

*Natural Order, COMBRETACEÆ, (R. BROWN.) Linnæan
Classification, DECANDRIA, MONOGYNIA.*

TERMINALIA, (Linn. DECAND.)

Flowers often POLYGAMOUS from abortion.—Border of the *calyx* deciduous, campanulate, 5-cleft, the divisions acute. *Petals* none. *Stamens* 10, in a double row, longer than the calyx. *Ovary* with 2 or 3 ovules. *Style* filiform, somewhat acute. *Drupe* not crowned by the calyx, often dry, indehiscent, 1-seeded. The *seed* resembling an almond. *Cotyledones* spirally convolute.

Trees of the largest size or shrubs, with alternate or rarely opposite leaves, crowded towards the extremities of the branches, and hence the generic name. Flowers in spikes, the spikes in racemes or panicles, bisexual in the lower part, and male in the upper.

§ I. CATAPPA, (Gærtner.) *The drupe compressed, with the margin winged or much attenuated.*

CATAPPA, OR INDIAN ALMOND.

TERMINALIA CATAPPA, *foliis obovatis basi attenuatis subtus molliter pubescentibus, glandulis minimis subtus in basi folii ad latus nervi medii.*—DECAND. Prod., vol. 3. p. 11. LINN. Mantis. p. 519.

TERMINALIA CATAPPA, leaves about the extremities of the branchlets on short petioles, obovate, cuneate and attenuated, at the same time slightly cordate at the base, a little repand, with a large depressed gland beneath on each side the midrib near the base: racemes axillary, solitary, simple, shorter than

D.

) *Linnaean*

A.

of the *calyx*
ute. *Petals*
an the calyx.
ewhat acute.
dehiscent, 1-
yledones spi-


ate or rarely
the branches,
the spikes in
l male in the


sed, with the

D.

enuatis sub-
is subtus in
od., vol. 3. p.

ities of the
d attenuated,
little repand,
le the midrib
, shorter than





Terminalia catappa.

Indian Almond

Botanicus de Malabar

11

the leaves; drupe oval, compressed, glabrous, with elevated navicular margins, convex on both sides. ARNOT, Prodr. Ind. Orient., vol. 1. p. 313. JACQUIN'S Ic. rar., vol. 1. tab. 197. LAM. Illust. tab. 848. fig. 1. *Adamarum*, RHEED, Flora Malabarica, vol. 4. tab. 3 and 4. TORREY and GRAY, Flor. N. Amer., vol. 1. p. 485.

ACCORDING to Torrey and Gray, Dr. Hasler has discovered this splendid tree in Southern Florida. A variety of it is known to exist in the Caribbean Islands, which Humboldt and Kunth imagined to be introduced, but for this supposition there is probably no sufficient ground, as Poiteau collected it in the forests of St. Domingo, of which I have a specimen now before me. A near congener, if not the same thing, was found in Guiana by Aublet, his *Tanibouca*; yet the favourite region of its existence is in the tropical forests of India, on the sandy and gravelly coasts of Malabar, and in the island of Java; it there becomes, according to Rheed, a very large and splendid tree of a pyramidal form, like that of a lofty spruce, the leafy summit being composed of almost horizontal branches disposed in circular stages. Its wood is white, very hard, covered with a smooth grey bark which is red within. The leaves, situated near to the extremities of the branchlets, 6 or 7 together, at intervals, form circular clusters of great regularity; they are about 6 to 9 inches long, by 3 to 5 wide, of an inversely ovoid or cuneate oval figure, widening towards the summit, where they become almost round, with a short, abrupt, slanting point in the centre, narrowed and somewhat cordate at the base, nearly entire, or obscurely, though sometimes very distinctly crenulated on the border, green and smooth above, slightly pubescent beneath; the young leaves and shoots as well as the petioles, clothed with a brown and close tomentum. The flowers are small, without scent, of a whitish-green, and disposed in great numbers in

several almost terminal axillary slender spikes, they are nearly sessile, with caducous, concave, oval, pointed bractes. The calyx contains a small, very hairy, 5-toothed cup. The spikes are not as long as the leaves. The fruit is an elliptic shell, a little compressed, glabrous, surrounded with an elevated margin, convex on both sides, and reddish-brown when mature. This dry drupe includes an oblong very hard nut, of one cell, containing a white kernel, of a taste approaching to that of the filbert nut, but more oily and soluble.

In India it is also cultivated in gardens. The large almond-like kernels of its nuts are eaten and served at the best tables. An oil is obtained from the kernels by expression similar to that of the olive, which is said never to become rancid. It is made also into emulsions like almonds. The Indians employ the leaves medicinally for indigestion, bilious affections, and other maladies.

PLATE XXXII.

A small branch of the natural size. *a.* The flower. *b.* The nut.

Terminalia Benzoin has a milky sap, and was believed to produce the Benzoic acid, which, however, is now doubted.

Another of the species, *Terminalia vernix*, is said to afford the celebrated Chinese and Japanese varnish used in their lacquer ware. This tree grows on the mountains of several of the southern provinces of China, and in the Moluccas. It possesses a lactescent juice, which as well as its exhalations even, are said to be deleterious, but the kernels of its fruits, like those of the Catappa, are perfectly harmless and agreeable. At Batavia regular plantations are made of the *Terminalia Moluccana* in the gardens and places of public resort, for the sake of its agreeable shade.

es, they are
ral, pointed
y hairy, 5-
the leaves.
d, glabrous,
ex on both
s dry drupe
containing
that of the

The large
d served at
the kernels
hich is said
o emulsions
ves medici-
other mala-

ver. *b.* The

was believed
ver, is now

, is said to
arnish used
e mountains
, and in the
, which as
deleterious,
e Catappa,
atavia regu-
Moluccana in
the sake of




Button tree.

Conocarpus erecta.

Conocarpe droit.



urpe droit.



CONOCARPUS.†

(GÆRTNER.)

*Natural Order, COMBRETACEÆ. Linnæan Classification,
PENTANDRIA, MONOGYNIA.*

The flowers densely aggregated in globular or oblong spikes like aments.—Tube of the *calyx* about the length of the ovary, persistent; the border 5-cleft. *Petals* none. *Stamens* 5 to 10, exserted; the anthers heart-shaped. *Ovary* compressed, containing 2 ovules. The *fruits* coriaceous, corky and scale-like, closely imbricated, and indehiscent. *Cotyledones* spirally convolute.

Small maritime trees or shrubs, with alternate, entire, somewhat coriaceous leaves. Heads of flowers pedunculated, axillary or terminal, solitary or in panicles.

BUTTON-TREE.

CONOCARPUS *ERECTA*, *foliis oblongis utrinque acuminatis sæpius basi biglandulosis, capitulis paniculatis*.—DECAND. Prod., vol. 2. p. 16. JACQ. Amer., p. 78. t. 52. CATESBY'S Carolina, tab. 33.

a. arborea. DECAND. l. c.

Conocarpus erecta, fruits retrorsely imbricated in a subglobose head, somewhat boat-shaped, scarcely winged; tube of the calyx not produced beyond the ovary; leaves oval-lanceolate, mostly acute or acuminate at each end, usually with 2 glands at the base; heads panicled.—TORREY and GRAY, Flor. N. Amer., vol. 1. p. 485.

† From *κωνος*, a *cone*, and *καρπος*, a *fruit*, its fruit resembling the cone of an alder.

Manghala arbor Curassavica foliis salignis. HERM. Parad.

Bat. COMMELIN, Hort. Amst., p. 115. cum. ic.

Alnus maritima myrtifolia coriariorum. PLUK. Almag. 18.
t. 240. f. 8.

Alni fructu laurifolia arbor maritima. SLOANE, Jam. Hist.,
2. p. 18. t. 161. f. 2.

Innominata. PLUM. ic. 135. t. 144. f. 2.

THIS is another tropical West Indian tree which the southern extremity of East Florida has afforded. It has been observed on the shore of Key West, Southern Florida, and around Tampa Bay. In the West Indies, like the Mangrove with which it grows, and for a kind of which it is taken by the Spaniards, who call it *Mangle Saragoza*, it affects the low sandy and muddy shores near the sea, where it becomes an erect tree about 30 feet high, with the trunk a foot in diameter, having a smooth whitish-grey bark and angular branchlets. In South America it also exists on the coast of Guayaquil, and in Chili, near Valparaiso. In a country where the finest kinds of wood are so common, that of the "Button Tree" is little esteemed, and it is, therefore, only used for fuel; it is, however, fine and close-grained, in the branches brownish-white, capable of a high polish, with scarcely any visible annual layers, and made up almost wholly of dotted medullary rays. The general aspect of its inflorescence, and, indeed, its closely imbricated inelegant heads of flowers lead us almost to compare it with some of the *amentaceæ*, particularly the Alder, while its real relations are to the present family, which includes in the *Combretum* itself, and the singularly splendid *Cacoucia* of Aublet, some of the most elegant and beautiful of plants.

The bark is grey, bitterish and astringent, and no doubt medicinal. The leaves of a yellowish-green, are from 2 to 3 inches long, $\frac{3}{4}$ to an inch broad, acute at

each end, very smooth, and on short petioles, which have frequently two glands at the base. The flowers, for which butterflies have a great predilection, are very inconspicuous, greenish-yellow, small, and collected into globose heads, in axillary and terminal few-flowered panicles on pedicels about the length of the capituli. The heads at length become reddish; the capsules are small and scale-like, corky, dilated elliptic, internally concave, with broad, thin, carinated margins, and are very often abortive, never more than 1-seeded, and pubescent at the summit.

The island of Cuba affords another allied but very distinct species, which may be the *C. procumbens* of Jacquin, put down as a variety of the present by Decandolle; the calyx, however, is almost entirely smooth, with very acute segments, and the leaves are sharply apiculated, and sometimes obtuse with a short point. In this the wood appears to be very hard and as close-grained as Mahogany, of a dull white, inclining to grey, with a delicate feathered appearance, and a thick bark, grey externally and blackish within.

According to Ponce Maximilian, the bark of the *Conocarpus racemosa* (one of the plants called Mangrove in Brazil) is much used at Rio Janeiro for tanning.

PLATE XXXIII.

A branch of the natural size. *a.* The flower. *b.* The fruit, of the natural size. *c.* The same magnified.

SILKY BUTTON-TREE.

CONOCARPUS ERECTA. γ. SERICEA, (Forster in herb. P'Herit.)
*foliis oblongis utrinque acuminatis ad utramque paginam
 etiam adultis adpressè villososericcis.* DECAND. Prod., vol.
 3. p. 16.

*Mangle foliis oblongis integris utrinque molli lanugine holo-
 sericea obductis.* SLOANE, Hist. Jam., vol. 2. p. 67. tab. 187.
 fig. 2.

ACCORDING to Sloane, this tree is known in Jamaica by the name of the White Mangrove, and attains the height of 20 feet, having white wood with a very small pith; the bark is also smooth and whitish. This variety, or species, has also been found, with the above, at Key West, in East Florida, by Dr. Blodgett. We do not see any thing to distinguish it as a separate species from the *erecta* except the peculiar silky, shining pubescence with which the leaves continue to be clothed even in the adult state.

erb. P'Herit.)
ue paginam
. Prod., vol.

ugine holo-
67. tab. 187.

n Jamaica
attains the
very small
his variety,
ve, at Key
We do not
te species
ing pubes-
othed even



Laguncularia racemosa

White Mangrove

Manglieri's grapes.



licia grappes.



LAGUNCULARIA,†

(GÆRTNER.)

Natural Order, COMBRETACEÆ. *Linnean Classification*,
DECANDRIA, MONOGYNIA.

Calyx border persisting, subcampanulate, 5-lobed. *Petals* 5, minute, spreading and caducous. *Stamens* 5 or 10, in two series, included. *Style* subulate, stigma capitate. *Nut* margined, coriaceous, valveless, 1 or 2-seeded, crowned with the calyx. *Cotyledones* convolute; the radicle very long.

A tree of the Caribbee islands with opposite, elliptic, smooth leaves. Racemes opposite, many-flowered, the flowers sessile, furnished with deciduous bractes; the calyx bilobed at the summit. The seed germinating within the nut. A genus nearly allied to LUMNITZERA of India.

WHITE MANGROVE.

LAGUNCULARIA RACEMOSA, *Gærtner*, fil. *Carpol.*, vol. 3. p. 209. t. 217. DECAND. *Prodr.*, vol. 3. p. 17.

CONOCARPUS RACEMOSA, *foliis lanceolato-ovatis obtusiusculis, fructibus segregatis*. LINN. *Sp. pl.* WILLD., vol. 2. p. 995. SWARTZ, *Obs.* p. 79. JACQ. *Amer.* p. 80. t. 53.

Conocarpus, foliis elliptico-ovatis, petiolis biglandulosis, racemis laxis, fructibus sejunctis. BROWNE, *Jam.* p. 159.

Mangle julifera, foliis ellipticis ex adverso nuscentibus. SLOANE, *Jam.* p. 156. *Hist.* vol. 2. p. 66. tab. 187. f. 1. RAII, *Dendr.* p. 115.

SPHENOCARPUS. *Richard*, *Anal. Fr.* p. 92.

SCHOUSBOA *commutata*. SPRENGEL, *Syst. Veget.*, vol. 2. p. 332.

THIS plant is a native of the sandy and muddy shores

† From *laguncula*, a little bottle, in allusion to the form of the nut.

of the Caribbean islands and the neighbouring continent, where it becomes a lofty branching tree, sometimes dividing into three or four trunks close to the ground: it is called White Mangrove by the English inhabitants of these islands. Dr. Blodgett has sent specimens of this tree also from Key West, in East Florida.

The branches are cylindrical and brownish, the twigs ferruginous; the leaves are opposite and smooth, about three inches long and an inch and a half wide, upon short petioles, quite entire, thick, and somewhat coriaceous, elliptic or ovate, obtuse, and sometimes emarginate, with a pair of glands near the summit of the petiole, and, in most of the leaves, towards the edge appear a number of raised glandular points, which are closed or open. The flowers are disposed in axillary and terminal elongated racemes, the racemes sometimes trifid. Flowers small, sessile, greenish-white; the germ pyriform, and, as well as the short border of the calyx, covered with a short whitish, silky pubescence. Petals 5, very small and caducous. Stamens 5, not exerted. The germ at its summit with 2 small dentiform bracteoles, the bractes themselves short, broad ovate and caducous. Style, at length somewhat exerted, with a small capitate stigma. Nut 1-seeded.

PLATE XXXIV.

A branch of the natural size. *a.* The nut in an early stage.

g continent,
sometimes
the ground:
inhabitants
specimens of
la.

, the twigs
ooth, about
wide, upon
what coria-
es emargi-
mit of the
s the edge
, which are
in axillary
sometimes
; the germ
the calyx,
ce. Petals
ot exerted.
form brac-
ovate and
ted, with a

arly stage.





Shepherdia argentea.

Rabbit Berry.




Argousier argente.







**IMAGE EVALUATION
TEST TARGET (MT-3)**




**Photographic
Sciences
Corporation**

23 WEST MAIN STREET
WEBSTER, N.Y. 14580
(716) 872-4503

0
LE 128 125
LE 132
LE 122
LE 120
LE 118
6

10
LE 130
LE 128



R A B B I T B E R R Y .

Natural Order, ELÆAGNEÆ, (JUSSIEU.) *Linnæan Classification*, DIÆCIA, OCTANDRIA.

SHEPHERDIA,† (NUTTALL.) HIPPOPHÆ, (PURSH.)

Flowers DIÆCIOUS.—MALE *calyx* 4-cleft, much larger than that of the female. *Corolla*, none. *Stamina* 8, alternating with a torus of 8 glands.—FEMALE flower with a small 4-cleft, superior, campanulate *calyx*, and 8 glands. *Style* 1; *stigma* oblique, subcapitate. *Berry* juicy, 1-seeded, globose, invested with the fleshy *calyx*.

Small trees, spinescent or unarmed, with the general aspect of *Eleagnus*. Leaves entire, opposite, clothed with silvery and ferruginous scales. Flowers small, in axillary clusters, or in spikes. Berries pulpy, diaphanous, scarlet, subacid.

R A B B I T B E R R Y, OR WESTERN SHEPHERDIA.

SHEPHERDIA ARGENTEA, *foliis oblongo-ovatis, obtusis, glabris, utrinque argenteo-lepidotis, floribus glomeratis*.—NUTT. Gen. Amer., vol. 2. p. 240. LONDON'S ENCYC. PLANTS, p. 836. Arboretum et Frutic. p. 1321. fig. 1208. HOOKER, Flor. Bor. Amer., vol. 2. p. 138. tab. 178, (well illustrated.) HIPPOPHÆ *argentea*, PURSH, Flor. Bor. Am. p. 115.

This very useful, hardy, and ornamental tree, is

† Named in honour of the late *Mr. Wm. Shepherd*, then curator of the Liverpool Botanic Garden. A most scientific gardener and skillful cultivator.

wholly a native of the northern and western regions of North America. Dr. Richardson observed it on the banks of the Saskatchewan, between Carlton and Edmonton House Forts, in the latitude of 51, and Major Long's party found it growing on the borders of Rainy Lake, about latitude 49. On the banks of the Missouri, the limit of its southern range is the borders of the Platte, but it appeared to be most abundant and fertile around Fort Mandan, or the Great Northern bend of the Missouri, in about the latitude of 48; here it becomes a small tree 12 to 18 feet in height, and when adorned with its brilliant scarlet berries, produced in thick clusters so as almost to conceal the branches, few objects are more ornamental, contrasted also with the silvery hue of the leaves, which reminds one of the useful olive; it presents at once an appearance both striking and novel.

Amongst the natives and Canadian voyageurs it has several different names. According to Lewis and Clarke, it was known on the Missouri, to the natives, by the name of the Rabbit Berry, from being fed on probably by those animals, and it was met with by their party in the gorges of the Rocky Mountains. Richardson says the Creek Indians give it a name which signifies Bloody Berry, (*Methcoo-meeva*,) from the singular redness and transparency of its fruit. The Canadian voyageurs called it *Graisse de bœuf*, or Buffalo-fat, from the imaginary relish of the berries, or from the practice of mixing them with their fat pounded meat or pemican.

In 1815 I introduced a plant into the Liverpool botanic garden, but being kept in the greenhouse, it was, I presume, killed with kindness, and was soon lost. About 12 or 15 years ago, my friends Messrs. Windships, of Brighton, in Massachusetts, in the vicinity of

Boston, obtained a few seeds of the *Shepherdia* from the banks of the Missouri, which growing, gave origin to all the plants now in the United States and in Europe. A standard tree, at this time growing in their nursery, is quite as large as an ordinary Apple-tree of the same age, and it is yet very vigorous and increasing in size. It is, I believe, about 20 feet high, with a handsome erect trunk, clothed with a somewhat smooth bark, and presents an appearance of the most vigorous health. Its resemblance to the Olive is very striking; it has a rounded uniform elegant summit, and when in fruit, which is at the close of summer, scarcely any thing can be more brilliant, from the load of berries with which it is every where clad; these are about the size of small red currants, juicy, but not watery, of a pleasant sub-acid taste, mixed with a sweetness, which renders them generally agreeable. Made into sweet jelly, in the manner of currants, they are thought preferable by most who have tasted them. But the great use of the *Shepherdia* will be for constructing hedges or live fences, at least in the Northern States where it thrives well. Kept down by cutting it becomes sufficiently close, and has also the advantage of being thorny, green, or rather silvery, till late in autumn, and it is attacked by no insect, nor subject to any disease or blight.

The berries are greedily devoured by all the autumnal birds, particularly Robins and Blue-birds, who flock round the tree in throngs while any thing remains to be had.

In its native state it is a small, rather narrow topped tree, with the branches ending in stout spines. The leaves are oblong-ovate, obtuse, shortly petiolate, on both sides free from hairs, but covered with peltate or rounded scales, which (through a lens) appear to be ciliated. The flowers, which come out as early as in

March, are in clusters. The calyx of the *male* flower is considerably larger than that of the female, and divided down to the base into four subovate, obtuse divisions, internally yellowish, but outside scaly like the leaves. The stamens are eight, with oftentimes rather short pubescent filaments; the anthers are oblong and 2-celled. The *female* flowers are smaller and shortly pedunculate, without any vestiges of stamens. There is one style, and a thickish, oblique, subelliptic stigma. The germ appears inferior, but is, in fact, only invested by the tube of the calyx. The berries are collected into clusters, and are sparingly scattered with scales, but bright and pellucid. The seed, or rather nut, with a cartilaginous shell, is subovate and shining, much like that of *Hippophae*, it is also scored externally as if partly 2-lobed, with a small projection at the base. The embryo is straight and flat, without albumen, and the radicle inferior. The cotyledons are large, thick and oval. Mr. Wyeth, in the Rocky Mountains, observed a variety of this species with yellow berries.

PLATE XXXV.

A branch of the natural size.

A second species of this genus is the

SHEPHERDIA CANADENSIS, or **CANADIAN SHEPHERDIA**, with elliptic-ovate leaves, nearly smooth above and naked beneath, clothed with stellate hairs and ferruginous scales, the flowers are also in axillary spikes. This species, it appears from Hooker, ranges far to the north, throughout Canada to Fort Franklin, on Mackenzie River, and from Newfoundland and Hudson's Bay to the Rocky Mountains. In the State of New York I have

met with it on the borders of the smaller inland lakes, as well as on the shores of Lake Erie; but this species scarcely becomes a tree, is without thorns, and the berries, though equally brilliant with the preceding, are rather unpleasant to the taste. On the authority of Menzies it is said also to exist on the North West coast, but I have not seen it there.

ale flower is
and divided
e divisions,
the leaves.
rather short
ong and 2-
and shortly
ns. There
otic stigma.
ly invested
ollected into
scales, but
nut. with a
, much like
nally as if
base. The
en, and the
thick and
s, observed

SHEPHERDIA,
above and
nd ferrugi-
ikes. This
o the north,
Mackenzie
Bay to the
ork I have

XIMENIA,†

(PLUMIER.)

Natural Order, OLACINÆE. Linnæan Classification,
OCTANDRIA, MONOGYNIA.

Calyx 4-toothed, minute, persistent, not enlarging. *Petals* 4, densely pilose within, connivent, above revolute. *Stamens* 8, the filaments capillary, anthers adnate, long and linear, not exserted. *Ovary* 4-celled, 4-seeded. *Style* 1. *Drupe* ovate, 1-seeded.—DECAND. 1. p. 533.

Tropical trees or shrubs, with smooth, alternate, entire, elliptical or ovate, exstipulate leaves; flowers mostly in small axillary umbels.

MOUNTAIN PLUM.

XIMENIA AMERICANA, *spinosa, foliis oblongis, pedunculis multifloris*.—LINN. Sp. Pl. Hort. Cliff. 1193. Swartz, Obs. p. 149. DECAND. Prod., vol. 1. p. 533.

a. ovata, foliis ovatis. *X. multiflora*. JACQUIN, Amer. p. 106. t. 277. fig. 31. LAMARCK, Illust. tab. 297. fig. 1.

XIMENIA *montana*. MACFADYEN, Flora of Jamaica, p. 121, a variety, however, without thorns.

XIMENIA *aculeata, flore villosa, fructu luteo*. PLUMIER, Gener. p. 6. Ic. 261. fig. 1.

THIS plant forms a small tree with an erect stem and spreading, grey, verrucose and somewhat angular branches. It is indigenous to the mountains of Jamaica,

† Named in honour of Francis Ximenes, a Spanish naturalist and missionary.

sification,

g. *Petals* 4,
Stamens 8,
d linear, not
Drupe ovate,

entire, ellip-
small axillary

, *pedunculis*
wartz, Obs. p.


mer. p. 106. t.

ica, p. 121, a

PLUMIER,

et stem and
at angular
of Jamaica,

ish naturalist






J.W. MacDougal

Ximelia Americana

Mountain Plum

Ximelia Americana



Key West in Florida, where it was found by Dr. Blodgett, and is also met with in the neighbourhood of Carthagera, in Hispaniola, and many years ago it was collected in the interior of East Florida by John Bartram, as Mr. A. Gray saw specimens of it in his collection still extant. According to Drs. Wight and Roxburgh it is also indigenous to the coast of India.

It bears a drupe the size of the Plum of Europe, or of a pigeon's egg, yellow, smooth, and shining, 1-seeded, with a thin rind and watery pulp of a pleasant sweet subacid taste. The seed is large and white. This Plum is of an agreeable flavour, and not inferior to the common varieties of that of Europe; it has a slight degree of astringency with a pleasant acidity. The flowers have a fragrant odour said to be like that of frankincense. The wood is as yellow as that of the Sandal, and, in India, its powder is often substituted for it by the Brahmins in their religious ceremonies.

The leaves grow 2 or 3 together, on short, lateral, tuberculoid branchlets; they are petiolate, oblong-lanceolate, obtuse and narrowed below, smooth, obscurely veined, about 2 or 2½ inches long, and less than an inch broad. The flowers are disposed in small pedunculated axillary and subterminal umbels, the umbels 3 or 4-flowered. The calyx is minute and 4-toothed. Petals 4, linear-oblong, conniving into a tube below, recurved at the apex, and covered with rather long and dense brownish-yellow hairs within. Stamens 8, as long as the petals, the filaments like the most delicate threads, the anthers long and linear, ovary 8-angled at the base, conical and subulate, with the style as long as the stamens.

PLATE XXXVI.

A branch of the natural size. *a.* The fruit.

MACLURA, †

(NUTTALL.)

*Natural Order, ARTOCARPEÆ, (R. BROWN.) Linnæan
Classification, DIÆCIA, TETRANDRIA.*

Flowers **DIÆCIOUS**.—**MALE** flowers with the *calyx* 4-cleft. *Stamens* 4, exserted. **FEMALE** flowers in globular aments; the *calyx* fleshy, 4-parted, with the bases ingrafted together. *Corolla* none. *Style* 1, filiform, villous. *Germs* numerous, each 1 seeded, coalescing into a compound globular berry of many cells, the cells 1-seeded. *Seed* obovate.

A lactescent tree, related to the Fustic, and with nearly similar yellow wood. Leaves alternate, entire, without stipules, producing superaxillary simple spines. Male flowers in pedunculated umbels. Aments axillary. Berry verrucose and large, resembling an orange, at first lactescent, greenish-yellow or yellow. Nearly allied to *Broussonetia*, but distinguished by the coalescence of the germs, and a peculiar habit.

OSAGE ORANGE, OR YELLOW WOOD.

MACLURA AURANTIACA. *Nutt. Gen. Amer.*, vol. 2. p. 234.
LOUDON'S Encycl. of Plants, p. 784. fig. 13256. *Arboretum et Frutic.*, vol. 3. p. 1342. 1362. figures 1226, 1227, 1228.
LAMBERT'S Supplement to Pinus, l. c. *EATON'S Manual*, (Ed. 8.) p. 311.

THIS remarkable tree, though perfectly thriving and

† In honour of the celebrated geologist, philosopher, and patron of natural science, Wm. Maclure.

Linnaean

calyx 4-cleft.
lobular aments;
fused together.
stamens numerous,
lobular berry of


nearly simi-
lar stipules,
scales in pedun-
cle and large,
yellow or yel-
lowish by the

WOOD.

Vol. 2. p. 234.
Arboretum
1227, 1228.
Gordon's Manual,

thriving and

philosopher, and







A. S. W. 1844


Maelua Aurantiaca.


Osage Orange

Bois d'Arc










Maclura Aurantiaca.

Osage Orange

Bos d'Arc



hardy in the climate of Philadelphia, raised from seeds, does not naturally extend to the north of the Arkansa river; it is even there of rare occurrence, and generally destitute of fruit. We saw a few old ill-grown trees on the banks of the Pottoc, a few miles from Fort Smith. It was only on the rich low bottom lands of Red River, near the confluence of the Kiamasha, that we beheld the *Maclura* in perfection, forming a great part of the prevailing umbrageous forest, and attaining an elevation of 50 or 60 feet, by a diameter of between 2 and 3 feet. It is, however, much inclined, when full grown, to throw out irregular spreading limbs of considerable extent, though at first it presents a very elegant roundish spreading summit. But at all times it strikes the beholder as something remarkable in the northern forest by the beauty and splendour of its dark and shining foliage, which in appearance strongly resembles that of the Orange, and the numerous spines which the branches present seem to confirm the comparison. The fruit is alike singular, resembling full grown yellowish-green Oranges, but filled with a milky juice, and, as they ripen at intervals, or become abortive, the ground beneath, like an orchard, is strewn with this curious fruit, which, when first discovered, lying neglected beneath the tree, led the voyagers to fear and report it as a poison; but the family to which it appertains and its relation to the Mulberry, affords a presumptive sanction for its harmless qualities.

The *Maclura* was first noticed by Hunter and Dunbar in their voyage up Red River, on the banks of the Little Missouri, of the Washita River, also near Natchitoches, and upon the banks of the Arkansa. It was likewise observed by Dr. James, in Major Long's Expedition, along the banks of the Arkansa and the Canadian. I first saw living plants, bearing fruit, about the year 1810,

in the garden of Mr. Chouteau, at St. Louis, which were raised from seeds collected in the country of the Osages. It was afterwards introduced into the garden of the late Mr. M^cMahon of Philadelphia, from seeds collected by Governor Lewis. The largest tree I have seen in cultivation is in the garden of Mr. Thomas (now David) Landreth, in Federal Street, Philadelphia. It is about 20 feet high and 5½ in circumference. This tree has for many years borne fruit, but the seeds have only of late been perfected, in consequence of the absence of the stamiferous individual, on the accession of which, however, it produced abundance of seeds nearly as large as those of the Orange, and much of the same form; these were indicated in the fruit by an unusual projection of the external tubercles.

The wood of the *Maclura* is solid, heavy, and elastic, of a fine yellow colour, which, like the Fustic, it readily communicates to water, and it might consequently be used as a dye. It is also capable of receiving a fine polish, and resembles satin-wood by its brilliant gloss. The elasticity and durability of its wood has long been celebrated by the Indians, who, from its use, have bestowed upon it the name of Bow-wood, and the Canadians, who traversed these regions in quest of furs, knew it long by the name of the *Bois d'arc*. The bark, as in *Broussonetia*, affords a fine white flax. Another important use of the *Maclura*, in this climate, is that of forming live fences or hedges, for which purpose it is well adapted, as it bears cutting, grows close, and is very thorny, as well as free from the attacks of blight and insects; it has all the advantage of keeping for a long time green, and appears, in all respects, as elegant a fence as that of the Wild Orange in the south. It has, besides, an additional recommendation in its use

for feeding silk worms, for which purpose it is scarcely inferior to the *famous Morus Multicaulis*.†

The branches are flexuous and round, clothed with a smooth grey bark. The leaves are alternate, upon long foot-stalks, and are usually oval and acuminate; on the bearing branches they are, however, often considerably larger and heart-shaped at the base, very entire, with the point mucronated and a little pungent, the upper surface is smooth and shining, but the petiole and nerves on the under side of the leaf are somewhat hirsutely pubescent. The petiole is often an inch or more long; the leaf itself 2 to 4 inches, and $1\frac{1}{2}$ to 3 inches wide. The *staminiferous* plant appears uniformly weaker, more delicate, and smaller leaved than the fertile plant. The flowers in it are axillary, in pedunculated small umbels, each umbel containing about 15 to 20 flowers, consisting merely of a small 4-cleft calyx, with oval hairy segments, and 4 stamens, on lengthened and exerted filaments. The anthers are 2-celled, large and oval, opening lengthwise. In both plants, single, undivided thorns come out in the upper axills of the leaves. The *female* capitulum consists of a congeries of flowers united into a globular form, about the size of a cherry; these consist also in a calyx of 4 divisions, but less regular than in the male. The styles and stigmas, one to each germ, are $\frac{3}{4}$ of an inch long, giving to the ament the appearance of a tuft of long pubescent threads. The berry filled with a milky juice, becomes about the size of a moderate but not large Orange, having an irregular tessellated appearance, almost like that of the Bread-fruit; these tessellations are the unduly enlarged

† Different opinions are now entertained of the value of the leaves of the *Maclura* as a food for silk-worms, some approving and others discouraging their use.

fleshy summits of the segments of the calyx. The whole of the calyces, at a short distance below the immediate surface, become partly engrafted together into one mass, and a transverse section of the fruit, therefore, presents a series of radiating and woody fibres, amongst which are immersed the one-seeded germs. The seeds are obovate, compressed, and nearly as large as those of the Orange. The testa is yellowish-white and cartilaginous; there is no albumen, and the seed is of a yellowish-brown, pendulous, with the radicle inverted from the axis of the fruit, and curved partly over the margin of the thick and fleshy cotyledones towards the point occupied by the hylum. The fruit, when ripe, is succulent, has a sweetish but insipid taste, and is somewhat acrid. As far as we know, it is not eaten by any animal.

It is readily propagated by sowing the seeds, and also by cuttings of the root, which grow much more readily than cuttings of the branches. Although several male plants were raised in this vicinity, it is singular to remark, that while the fertile plant is in its utmost vigour, very few of the former survive; and as their presence is necessary for fertility, it probably would be the best way to ingraft the male on some one of the branches of the bearing plant.

PLATE XXXVII.

A branch of the male plant of the natural size *a*. The male flower a little enlarged. *b*. The female capitulum.

PLATE XXXVIII.

The fruit of the natural size. *a*. A transverse section of the same. *b*. The appearance of the abortive germs in the section. *c*. The seed with the testa. *d*. The same divested of the testa. *e*. The embryo. *f*. The back view of the same.

Besides the Fig, we have, in this interesting family of plants, the famous Bread-fruit of India and the islands of the Pacific. The Cow Tree, or Palo de Vaca of South America, which is tapped like a fountain, and yields a copious supply of rich and wholesome milk. The nuts of the *Brosimum alceastrum*, when roasted, are used in the place of bread, in the West Indies, and have a taste similar to that of Hazel-nuts: the juice of all the plants of this family also contains more or less of caoutchouc. Amidst this generally harmless group of plants, it is singular to find the deadly Upas of Java, (*Antiaris toxicaria*,) whose inspissated juice is found to contain that most virulent of all poisonous principles, strychnia. It is, however, somewhat doubtful what the real affinities of this plant are, as it is acknowledged to be an anomaly in the family.

the whole
mediate
into one
therefore,
amongst
the seeds
as those
and car-
d is of a
inverted
over the
wards the
n ripe, is
l is some-
en by any

, and also
are readily
veral male
ular to re-
st vigour,
presence is
e the best
branches of

The male

tion of the
the section.
of the testa.

NETTLE TREE.

Natural Order, ULMACEÆ, (Mirbel.) *Linnæan Classification*, POLYGAMIA, DICEIA.

CELTIS,† (LINN.)

Flowers POLYGAMOUS.—*Staminate* flower with the calyx 5 or 6-parted. *Corolla* none. *Stamens* 4 to 7. *Perfect flower* with the calyx deeply 5-parted. No *corolla*. *Ovarium* 1-celled; the *ovules* solitary and pendulous. *Style* very short; *stigmas* 2, thickish, subulate, and spreading. *Drupe* globular, 1-seeded, thinly coated with a sweetish pulp. *Embryo* inverted. *Cotyledones* folded.

The genus within its proper bounds includes deciduous leaved trees of South Europe, the Levant, the mountains of Nepal and Cuba, and the forests of the United States. The true Celtides have alternate, entire, deciduous, and mostly cordate leaves, generally oblique at the base, 3-nerved, entire, but mostly serrated on the margin. The stipules are membranaceous and deciduous. The flowers are precocious, or appear before the expansion of the leaves, with a film-like irregularly torn membranous perianth, the stamiferous ones near the base of the branchlet pedicellate, and 3 or 4 together. The fertile flowers are solitary and axillary, on short peduncles. The drupes brownish-yellow, rather sweet, insipid, and nearly juiceless.

For the tropical species with axillary cymes coeval with the leaves, 2 distinct styles and an ovary with 2 ovules, I propose the name of TRACHYDENDRON, (in allusion to their rough pubes-

† The ancient name of the *Lotus*, applied to this genus by Linnæus.

Classifica-

the calyx 5 or
perfect flower
Ovarium 1-
very short;
frupe globu-
s. *Embryo*

luous leaved
of Nepal and
true Celtides
date leaves,
mostly ser-
ous and deci-
e the expan-
membranous
the branchlet
are solitary
nish-yellow,

val with the
es, I propose
rough pubes-

his genus by



Celtis reticulata.

Small leaved Nettle tree.

Wormwood tree.





cent leaves and twigs.) Most of these species have a tough fibrous bark of the nature of hemp.

SMALL LEAVED NETTLE TREE.

CELTIS RETICULATA, foliis brevibus, lato-cordatis, subcoriaceis, vix et irregulariter serratis acutis basi obliquis scabris, subtus subglabris venis elevatis reticulatis, pedunculis fructiferis unifloris.

CELTIS reticulata. TORREY, in *Annals of Lyceum*, N. Y., vol. 2. p. 247.

THIS low growing species of Nettle tree was discovered by Dr. James near the base of the Rocky Mountains; I likewise met with it in the same mountain range, by small streams, and also along the borders of the Oregon, towards the Blue Mountains, particularly along the banks of the Brulée, a small stream falling into that river. It does not, in the situations where we observed it, become a timber tree, but rather a tall shrub, full of slender, and, at length, smooth branches. The leaves become thick and rigid, and are about an inch and a half long, by less than an inch wide, acute, but scarcely acuminate, with a few irregular serratures towards the point of the leaf, though a number of the leaves may be observed possessing no serratures at all; the upper surface is shining and scabrous, beneath the leaves are pubescent along the nerves, though at length nearly quite smooth; the petioles are 1 or 2 lines long and pubescent; the base of the leaf is very oblique, rounded and slightly sinuated. The drupe is globose, solitary, brownish-yellow, on a short peduncle. Of the wood of this species nothing is yet known.

PLATE XXXIX.

A branch of the natural size.

LONG LEAVED NETTLE TREE.

CELTIS LONGIFOLIA, *foliis ovato-lanceolatis promissa acuminatis integerrimis levigatis demum glabris basi rotundatis obliquis subcuneatis, pedunculis fructiferis unifloris, cortice subtevi.*

CELTIS occidentalis, *β. integrifolia*, NUTT. Gen. Am., vol. 1. p. 202. (not of LAMARCK.)

THIS tree, growing to the height of 60 or 70 feet, inhabits the deep shady forests which border the Mississippi from St. Louis to the vicinity of the sea. Its even and not deeply cleft bark, in the absence of its aspiring summit, at once distinguishes this species from the Common Hack-Berry. Like all the rest of the genus, (confined within its proper limits,) the insignificant filmy flowers appear early in the spring, before the expansion of the leaves. The small branches are smooth and yellowish-brown. The leaves are smooth, of a thin consistence, and remarkable for the great length of their acuminate points, of an ovate or ovate-lanceolate form, subtended by deciduous stipules, and at first pubescent beneath, particularly along the large vessels or veins; at length almost absolutely smooth, at no time scabrous either above or beneath, and wholly entire on the margin. The length is about from 3 to 3½ inches by 1 to 1½ wide. The obliquity of the base varies according to the position of the leaf on the branch; those first developed are nearly equal at base, and of a lanceolate form; the later ones are larger, wider, and more oblique. The flowers are as usual; the males about 3 together towards the base of the branch, the females


E.

omisse acu-
basi rotun-
iferis uni-

Am., vol. 1.

70 feet, in-
the Missis-

Its even
s aspiring
from the
the genus,
cant filmy
expansion
ooth and
of a thin
th of their
lanceolate
l at first
ge vessels
at no time
entire on
3½ inches
varies ac-
ch; those
of a lan-
and more
s about 3
e females





Celtis longifolia.

Long leaved Nitte tree

Micoquier à longues feuilles

