





**IMAGE EVALUATION
TEST TARGET (MT-3)**



**Photographic
Sciences
Corporation**

23 WEST MAIN STREET
WEBSTER, N.Y. 14580
(716) 872-4503

14
18
28
32
25
22
20
9

**CIHM/ICMH
Microfiche
Series.**

**CIHM/ICMH
Collection de
microfiches.**

oi



Canadian Institute for Historical Microreproductions

Institut canadien de microreproductions historiques

1980

Technical and Bibliographic Notes/Notes techniques et bibliographiques

The Institute has attempted to obtain the best original copy available for filming. Features of this copy which may be bibliographically unique, which may alter any of the images in the reproduction, or which may significantly change the usual method of filming, are checked below.

L'Institut a microfilmé le meilleur exemplaire qu'il lui a été possible de se procurer. Les détails de cet exemplaire qui sont peut-être uniques du point de vue bibliographique, qui peuvent modifier une image reproduite, ou qui peuvent exiger une modification dans la méthode normale de filmage sont indiqués ci-dessous.

- Coloured covers/
Couverture de couleur
- Covers damaged/
Couverture endommagée
- Covers restored and/or laminated/
Couverture restaurée et/ou pelliculée
- Cover title missing/
Le titre de couverture manque
- Coloured maps/
Cartes géographiques en couleur
- Coloured ink (i.e. other than blue or black)/
Encre de couleur (i.e. autre que bleue ou noire)
- Coloured plates and/or illustrations/
Planches et/ou illustrations en couleur
- Bound with other material/
Relié avec d'autres documents
- Tight binding may cause shadows or distortion
along interior margin/
La reliure serrée peut causer de l'ombre ou de la
distortion le long de la marge intérieure
- Blank leaves added during restoration may
appear within the text. Whenever possible, these
have been omitted from filming/
Il se peut que certaines pages blanches ajoutées
lors d'une restauration apparaissent dans le texte,
mais, lorsque cela était possible, ces pages n'ont
pas été filmées.
- Additional comments:/
Commentaires supplémentaires:

- Coloured pages/
Pages de couleur
- Pages damaged/
Pages endommagées
- Pages restored and/or laminated/
Pages restaurées et/ou pelliculées
- Pages discoloured, stained or foxed/
Pages décolorées, tachetées ou piquées
- Pages detached/
Pages détachées
- Showthrough/
Transparence
- Quality of print varies/
Qualité inégale de l'impression
- Includes supplementary material/
Comprend du matériel supplémentaire
- Only edition available/
Seule édition disponible
- Pages wholly or partially obscured by errata
slips, tissues, etc., have been refilmed to
ensure the best possible image/
Les pages totalement ou partiellement
obscurcies par un feuillet d'errata, une pelure,
etc., ont été filmées à nouveau de façon à
obtenir la meilleure image possible.

This item is filmed at the reduction ratio checked below/
Ce document est filmé au taux de réduction indiqué ci-dessous.

10X	12X	14X	16X	18X	20X	22X	24X	26X	28X	30X	32X
						✓					

The copy filmed here has been reproduced thanks to the generosity of:




Library of the Public
Archives of Canada

The images appearing here are the best quality possible considering the condition and legibility of the original copy and in keeping with the filming contract specifications.

Original copies in printed paper covers are filmed beginning with the front cover and ending on the last page with a printed or illustrated impression, or the back cover when appropriate. All other original copies are filmed beginning on the first page with a printed or illustrated impression, and ending on the last page with a printed or illustrated impression.

The last recorded frame on each microfiche shall contain the symbol \rightarrow (meaning "CONTINUED"), or the symbol ∇ (meaning "END"), whichever applies.

Maps, plates, charts, etc., may be filmed at different reduction ratios. Those too large to be entirely included in one exposure are filmed beginning in the upper left hand corner, left to right and top to bottom, as many frames as required. The following diagrams illustrate the method:



L'exemplaire filmé fut reproduit grâce à la générosité de:

La bibliothèque des Archives
publiques du Canada

Les images suivantes ont été reproduites avec le plus grand soin, compte tenu de la condition et de la netteté de l'exemplaire filmé, et en conformité avec les conditions du contrat de filmage.

Les exemplaires originaux dont la couverture en papier est imprimée sont filmés en commençant par le premier plat et en terminant soit par la dernière page qui comporte une empreinte d'impression ou d'illustration, soit par le second plat, selon le cas. Tous les autres exemplaires originaux sont filmés en commençant par la première page qui comporte une empreinte d'impression ou d'illustration et en terminant par la dernière page qui comporte une telle empreinte.

Un des symboles suivants apparaîtra sur la dernière image de chaque microfiche, selon le cas: le symbole \rightarrow signifie "A SUIVRE", le symbole ∇ signifie "FIN".

Les cartes, planches, tableaux, etc., peuvent être filmés à des taux de réduction différents. Lorsque le document est trop grand pour être reproduit en un seul cliché, il est filmé à partir de l'angle supérieur gauche, de gauche à droite, et de haut en bas, en prenant le nombre d'images nécessaire. Les diagrammes suivants illustrent la méthode.

NOTE ON EXCISION OF THE KNEE JOINT.¹

BY

SIR JAMES GRANT M.D.

The knee-joint, the most complicated anatomically in the human system, has been subjected to excision, chiefly for three conditions

- 1st. Chronic disease of the articulation.
- 2nd. Ankylosis, whether fibrous or bony.
- 3rd. Compound fracture or dislocation.

The first excision of the "knee-joint" successfully, was performed by Filkin of Norwich, in 1762, and again by Park, in 1781, 40 years ago, it has been performed in all less than 25 times, owing chiefly to its receiving but little favour. This operation has been very considerably revived by Textor of Germany, and the late Sir Wm. Ferguson, of London. The first excision of this joint in the United States, 1856 was performed by Dr. Kinloch of Charleston; and in March, 1865, in the Protestant Hospital, by myself, this being one of the first knee excisions in Canada.² The term excision has quite a different signification now, from what it had thirty years ago. Then it meant somewhat free removal of much of the structures essential to the functions of the articulation. At present it means, in many cases, very partial interference with the structures essential to the joint, while it aims in every case, at the complete removal of the diseased tissues. To dig out sequestra from the surfaces of bones, and scrape the latter, and thus not shorten the limb, is the order of the day. Quite a change from former times, when the ends of the bones would be sawn off, in order to coapt parts and form a mechanical joint without much consideration as to future usefulness. Such, however, was more the result of a want of knowledge of our present advanced pathology than otherwise. The term "arthrectomy" was for a time spoken of as meaning partial excision of the joint, where the diseased synovial membrane is chiefly or solely removed and "excision" to removal of the ends of the bones completely. These terms arthrectomy and excision so overlap each other, and fail to define the exact meaning, that the terms partial and complete excision are now generally adopted, and chiefly owing to the fact that it is exceedingly rare to meet with a case of knee-joint disease in which the whole of the synovial membrane requires extirpation and yet none of the bone.

¹ Read before the Ottawa Medical Society, Feb., 1866.

² Reported in Medical Times and Gazette, London, Eng., March 18, 1865.

and vice versa, it is just as rare to meet with a case requiring the free removal of the ends of the bones, and not calling for the removal of the synovial structures. To-day complete excision of the knee-joint is a novelty, that is, opening the joint, sawing off the ends, and coapting the parts as in the days of Professor Geo. Fenwick, of McGill, whose name is handed down as one of the most celebrated surgeons Canada has produced. Why these changes or modifications as to treatment of such cases? First, the more accurate recognition of the vast importance of early and vigorous treatment of tuberculous disease, has arrested much, which formerly progressed to complete destruction of the ends of both the tibia and femur; and on the other hand, when operation is decided upon, it is at a stage when disease of the bones can be removed without complete excision of the ends, although the bone tissue is conserved as much as possible, the removal of the synovial tissue is now particularly free. In fact, Treves states it is not an uncommon procedure to dissect out the entire synovial membrane, whether visibly affected or not. In general terms, in a case of tuberculous joint, the first principle is to remove all tissues in which we believe the bacilli are deposited and to spare all structures, except the synovial membrane, not charged with the tubercle bacillus. It must be admitted that this is after all, a complex problem, not only in theory, but also in practice. In adults exsection is truly the shortest and safest way of eliminating the tedious morbid process and substituting ankylosis for a comparatively useless joint. To adult cases, orthopaedic treatment is rarely applicable, while in children, mechanical and general treatment frequently are attended with the most practical results. In the child, the growth of the thigh and tibia, depends so much on the epiphyses adjoining the knee, that exsection is liable to be followed by very considerable shortening, and consequently is to be avoided as much as possible. The antiseptic treatment in such operations, as defined by Gerster, requires no comment. To be truly progressive is to be truly antiseptic in treatment, the results of which certainly mark the present as a progressive era in surgery. In the older operations on the knee-joint, in which an absolutely stiff knee was the object in view, it was not looked upon as important, whether the ligamentum patellæ was divided or not, in such a case the action of the extensor muscles not being considered. More recent operations, however, tend to the preservation intact of the ligamentum patellæ, in order to preserve, as far as possible, some of the movements. Treves recommends that if the patellar ligament has been divided, it should be stitched together again with silk, which remains buried. He considers it better not to include the capsular

the joint in the stitches which unite the skin wound, as the skin is liable to be dragged in and the edges displaced, thus interfering with primary union. It is also important that the edges of the capsule and the fibrous expansion of the muscle should not be united, as any effusion into the joint cavity after the operation would thus not be pent up, but escape into the areolar tissue around and become more rapidly absorbed, and should even any pus be formed, it would be more easily evacuated. Sayre, in his recent work on "Joint Diseases," states as his opinion that exsection of the knee-joint is attended with considerable danger, and in many instances you may justly hesitate before resorting to the operation. Mr. Bryant, in a lecture at Guy's Hospital, published in the *Lancet*, "On the least sacrifice of parts as a principle in operative surgery," states: "I trust that this series of cases is enough to demonstrate with sufficient clearness, the value of the practice I am now inculcating, and to show that in a large number of cases of disease of the joints, a cure may be secured by a simple incision into the affected joint and the removal of the necrosed bone. The series includes examples of disease of the shoulder and elbow, hip, knee and ankle and great toe joints, and I do not think I should be far wrong if I were to express my belief that in many cases, if not in all, many surgeons, more particularly those who are advocates for excision, would have excised the joints, and some few, would have amputated. In the treatment I am now advocating, the surgical proceedings are simple and are attended with a minimum of danger. The success of the practice I have recorded was great." The opinion of Mr. Bryant is one founded on large practical experience and is certainly worthy of the fullest consideration, when such trying cases present for surgical interference and treatment.

