

CIHM/ICMH Microfiche Series.

CIHM/ICMH Collection de microfiches.



Canadian Institute for Historical Microreproductions / Institut canadian de microreproductions historiques



Technical and Bibliographic Notes/Notes techniques et bibliographiques

The Institute has attempted to obtain the best original copy available for filming. Features of this copy which may be bibliographically unique, which may alter any of the images in the reproduction, or which may significantly change the usual method of filming, are checked below.

L'Institut a microfilmé le meilleur exemplaire qu'il lui a été possible de se procurer. Les détails de cet exemplaire qui sont peut-être uniques du point de vue bibliographique, qui peuvent modifier une image reproduite, ou qui peuvent exiger une modification dans la méthode normale de filmage sont indiqués ci-dessous.

	Coloured covers/ Couverture de couleur		Coloured pages/ Pages de couleur	Or
	Covers damaged/ Couverture endommagés		Pages damaged/ Pages endommagées	the
	Covers restored and/or laminated/ Couverture restaurée et/ou pelliculée		Pages restored and/or laminated/ Pages restaurées et/ou pelliculées	firs sig
	Cover title missing/ Le titre de couverture manque		Pages discoloured, stained or foxed/ Pages décolorées, tachetées ou piquées	
	Coloured maps/ Cartes géographiques en couleur		Pages detached/ Pages détachées	Th
	Coloured ink (i.e. other than blue or black)/ Encre de couleur (i.e. autre que bleue ou noire)	•	Showthrough/ Transparence	TII wt
1	Coloured plates and/or illustrations/ Planches et/ou illustrations en couleur		Quelity of print varies/ Qualité inégale de l'impression	Ma dif en:
	Bound with other material/ Relié avec d'autres documents		Includes supplementary material/ Comprend du matériel supplémentaire	rig rec
v	Tight binding may cause shadows or distortion along interior margin/ La re liure serrée peut causer de l'ombre ou de la		Only edition available/ Seule édition disponible	
	distortion le long de la marge intérieure		Pages wholly or partially obscured by errata slips, tissues, etc., have been refilmed to	
	appear within the text. Whenever possible, these have been omitted from filming/ Il se peut que certaines pages blanches ajoutées lors d'une restauration apparaissent dans le texte, mais, lorsque cela était possible, ces pages n'ont pas été filmées.		ensure the best possible image/ Les pages totalement ou partiellement obscurcies par un feuillet d'errata, une pelure, etc., ont été filmées à nouveau de façon à obtenir la meilleure image possible.	
	Additional comments:/ Commentaires supplémentaires;			

This item is filmed at the reduction ratio checked below/ Ce document est filmé au taux de réduction indiqué ci-dessous.



The to th

> The poss of th filmi

gi gi n, ne st m, 111

e 11 VL lic

pp ife tir giu ĮU itt The copy filmed here has been reproduced thanks to the generosity of:

Library of the Public Archives of Canada

The images appearing here are the best quality possible considering the condition and legibility of the original copy and in keeping with the filming contract specifications.

Original copies in printed paper covers are filmed beginning with the front cover and ending on the last page with a printed or illustrated impression, or the back cover when appropriate. All other original copies are filmed beginning on the first page with a printed or illustrated impression, and ending on the last page with a printed or illustrated impression.

The last recorded frame on each microfiche shall contain the symbol \longrightarrow (meaning "CON-TINUED"), or the symbol ∇ (meaning "END"), whichever applies.

Meps, plates, charts, etc., may be filmed at different reduction ratios. Those too large to be entirely included in one exposure are filmed beginning in the upper left hand corner, left to right and top to bottom, as meny frames as required. The following diagrams illustrate the method:

1 2 3

L'exemplaire filmé fut reproduit grâce à la générosité de:

Le bibliothèque des Archives publiques du Cenade

Les images suivantes ont été reproduites svec le plus grand soin, compte tenu de la condition et de la natteté de l'exemplaire filmé, et en conformité evec les conditions du contrat de filmage.

Les exemplaires originaux dont la couverture en papier est imprimée sont filmés en commençant par le premier plat et en terminent soit par la dernière page qui comporte une empreinte d'impression ou d'illustration, soit par le second plat, selon le cas. Tous les autres exemplaires originaux sont filmés en commençant par la première page qui comporte une empreinte d'impression ou d'illustration et en terminant par la dernière page qui comporte une telle empreinte.

Un des symboles suivants apparaîtra sur la dernière image de cheque microfiche, selon la cas: le symbole → signifie "A SUIVRE", le symbole ▼ signifie "FIN".

Les cartes, planches, tablesux, etc., peuvent être filmés à des taux de réduction différents. Loreque le document est trop grand pour être reproduit en un seul cliché, il est filmé à partir de l'angle supérieur gauche, de gauche à droite, et de haut en bas, en prenent le nombre d'Images nécessaire. Les diagrammes suivants lilustrent la méthode.



1	2	3
4	5	6 ·

étails étails is du nodifier ir une iimage

errata i to

t e pelure, :on à

S] T BY

ARCTIC

SEARCHING EXPEDITION:

JOURNAL OF A BOAT-VOYAGE

THROUGH RUPERT'S LAND AND THE ARCTIC SEA,

IN SEARCH OF THE DISCOVERY SHIPS UNDER COMMAND OF

SIR JOHN FRANKLIN.

WITH AN APPENDIX ON THE PHYSICAL GEOGRAPHY OF NORTH AMERICA.

BY SIR JOHN RICHARDSON, C.B., F.R.S.

INSPECTOR OF NAVAL HOSPITALS AND FLEETS, ETC. ETC. ETC.

IN TWO VOLUMES.

VOL. II.

PUBLISHED BY AUTHORITY.

LONDON: LONGMAN, BROWN, GREEN, AND LONGMANS. 1851.



CHAPTER XIII.

OF THE 'TINNE OR CHEPEWYANS.

Page

1

Geographical Position. National Name. Tribes. Hare Indians and Dog-Ribs. Personal Appearance. Women. Dispositions. Wars. Socialism. Improvi-Dress. dence. Suffering. Affection for their Children. Hospitality feeble. Falsehood. Honesty. Religious Belief. Volatility. Marriages. Wrestling for a Wife. Dogs. Moose hunting. Public Opinion the only Rule of Conduct. Chiefs. Introduction of Christianity. Horses. Houses. Dawnings of Civilization. Members of the 'Tinnè People west of the Rocky Mountains. Southern Athabascans . . .

CHAP. XIV.

EYTIHNYUWUK, OR CREES AND CHIPPEWAYS.

National Names. Division. Tribes. Territory. Wars with the Mengwe. Conventional Character not true. Persons. Gait. Crimes. Wabunsi. Wigwams. Religious Belief. Vapour Baths. Everlasting Fire. Its rites. Used in Sickness. Its Priests. Its Origin. Chief Sun. Policy. Calumet. Maize. Food. Rein-White-fish. deer. Bison. Earth-works. Pottery. Language. Half-breeds. Colony of Red River or Osnaboya. Spirituous Liquors .

CHAP. XV.

OCCURRENCES IN WINTER.

Page

M

M

Fort Confidence. Situation. Silurian Limestone. Lake basin. Trees. Dwelling-house. Occupations. Letters. Galena Newspaper. Oregon "Spectator." Extent of the Hudson's Bay Company's Territory. Fisheries. Venison. Wolverines. Native Socialism. Provisions collected at Fort Confidence. Fetes. Winter Fishery. Eskimo Sleds. Reindeer. Wolverine. Wolves. Honesty of the Dog-Ribs. Their Indolence. Provisions not individual Property. Indians move off. An Accouchement. Colebs in search of a Wife. Might makes Right. None but the brave deserve the fair. Progress of the Seasons. Temperature. Arrival of Summer Birds. At Fort Confidence. At Fort Franklin. On the Yukon -- 61

CHAP. XVI.

Mr. Rae's Expedition in the Summer of 1849. Instructions. He crosses to the Coppermine. Descends that River. Sea covered with Ice. Surveys Rae River. Cape Kendal. Cape Hearne. Basil Hall Eskimos. Bay. Cape Krusenstern. Douglas Island. Detention. Dangerous Situation. August 23., return. Author and Mr. Bell leave Fort Confidence. Cross Great Bear Lake. Descend Bear Lake River. David Brodie lost in the Woods. His Adventures. Fort Simpson. Methy Portage. Receive English Letters. Norway House. Part from the Seamen and Sappers and Miners. Continue the Voyage to Canada. Boston. Land at Liverpool. Summary of the present State of the Search for Sir John Franklin -- 113

APPENDIX.

No. I.

PHYSICAL GEOGRAPHY.

GENERAL VIEW. Rocky Mountains. Their Length. Their Height. Glaciers. Parallelism to the Pacific Coast. Continental Slopes. Russian America. Eastern Slope. Prairies. Mississippi Valley. Its Slope. Fundamental Rocks of the Basin. Silurian Strata. Carboniferous Series. Tertiary Beds. Lignite Formations. Series of Lake Basins. Transverse Valleys. Intermediate Belt of primitive, hypogenous, or metamorphic Rocks. Its Rivers mere Chains of Lakes. Its Breadth. Altitude. Sources of three great River Systems. Great Fish River. The Yukon or Kwichpack. Basins of Excavation. Glacial Actiou. Active Volcanoes - - - 161

- VALLEY OF THE ST. LAWRENCE. Altitudes of the Lakes above the Sca. Lake Superior. Lake Michigan. Lake Huron. Lake Erie. Lake Ontario. Lake Chaplain. Northern Brim of the St. Lawrence Basin. Its geological Structure. North Shore of Lake Superior. Structure of the Country at the Sources of the Mississippi - 177 WINIPEG OR SASKATCHEWAN VALLEY. Height of Lake

Elk, or Red-deer River. Lesser Slave Lake. Peace

Page Lake eters. nt of eries. sions hery. dives. Prooff. light fair. d of klin. - 61

tructhat iver. Hall tion. nthor Great rodie pson. rway ners. d at earch - 113 v

Page

								Page
River.	Slave	River.	River of	the M	lounta	ins.	Noh'-	
hanne l	Bute.	Great 1	Bear Lake		•	-		1 9 9
YUKON V	ALLEY.	Yuko	on or Kwiel	hpack.	Vol	canic	Chain	
of Alas	ka. C	oal. F	ossil Bones					205

No. II.

CLIMATOLOGY.

Snow Line. Ground Ice. Thermometrical Observations in the Valley of the St. Lawrence. Comparative Temperature of the Two Sides of the Continent. Phenomena of the Seasons at Penetanguishene: At Fort William: At Fort Vancouver. Thermometrical Observations in the Valley of the Saskatchewan. On the East and West Sides of the Continent in that Parallel. Phenomena of the Seasons at Cumberland House : At Carlton House: At Marten's Falls on Albany River. Thermometrical Observations on the Missinipi and in the same Parallels on the East and West Sides of the Continent. Thermometrical Observations in the Valleys of the Mackenzie, Yukon, and Pelly. Progress of the Seasons at Fort Franklin. The mometrical Observations on the Arctic Seas. General Remarks. Nocturnal Radiation - 212

E

E

Cr

No. III.

ON THE GEOGRAPHICAL DISTRIBUTION OF PLANTS NORTH OF THE 49TH FARALLEL OF LATITUDE.

Generic and Specific Forms of Plants decrease in Number as the Latitude increases. Analogy between Altitude and Increase of Latitude. Culture of the Vine. Of the *Cerealia*. Maize. Wheat. Oats. Barley. Potatoes. Botanieal Districts. Their Physiognomy. Woodland District. Barren Grounds. Prairies. Rocky Mountains.

Page h'-- 199 ain - 205

0118 mnoort serast hearlerthe onevs the ons nal - 212

DRTH

ber ide ihe es.

nd ns.

No. IV.

LIST OF INSECTS.

Note on Hymenoptera in Arctic North America. List of Coleoptera. Orthoptera. Neuroptera. Hymenoptera. Hemiptera. Homoptera. Lepidoptera. Diptera - 354

No. V.

VOCABULARIES.

EXPLANATION OF PLATES I. AND II. - - 403

PLATE.

Cree Wigwams. (Vide p. 43.) - - - - Frontispiece

vii

Page



ARCTIC

SEARCHING EXPEDITION.

CHAPTER XIII.

OF THE 'TINNE OR CHEPEWYANS.

GEOGRAPHICAL POSITION. — NATIONAL NAME. — TRIBES. — HARE INDIANS AND DOG-RIBS. — PERSONAL APPEARANCE. — WOMEN. — DRESS. — DISPOSITIONS. — WARS. — SOCIALISM. — IMPROVI-DENCE. — SUFFERING. — AFFECTION FOR THEIR CHILDREN. — HOSPITALITY FEEBLE. — FALSEHOOD. — HONESTY. — RELIGIOUS BELIEF. — VOLATILITY. — MARRIAGES. — WRESTLING FOR A WIFE. — DOGS. — MOOSE-HUNTING. — PUBLIC OPINION THE ONLY RULE OF CONDUCT. — CHIEFS. — INTRODUCTION OF CHRISTI-ANITY. — HORSES. — HOUSES. — DAWNINGS OF CIVILISATION. — MEMBERS OF THE 'TINNÈ PEOPLE WEST OF THE ROCKY MOUNTAINS. — SOUTHERN ATHABASCANS.

'TINNE' or 'Dtinne', Athabascans, or Chepewyans.
Under these national appellations I have to speak of a people whose southern border is the Churchill River, or the Missinipi, as it is termed by the Eythinyuwuk, to whom it is also a boundary line.
Every where, in the country lying east of the VOL. II.

Mackenzie, the 'Tinnè lands are conterminous with the Eskimo coast, and, to the westward of the Rocky Mountains, with the Kutchin grounds, though the precise geographical limits of the two nations in that direction have not yet been correctly ascertained. The 'Tinnè, however, extend across the continent, since the $T\bar{a}$ -kuli and almost the entire population of New Caledonia have been referred by ethnologists to their nation.

The name by which the 'Tinnè designate themselves has, as is usual with the native Americans, the signification of "people," or "the people," and its proper application, when ascertained with care, would seem, at first sight, to be a good test for fixing the nationality of some tribes whose position in the ethnological scale is still uncertain. But as our acquaintance with the various American languages extends, and the way in which the pronunciation of the same word in the mouths of different tribes is gradually modified becomes known, doubts arise as to the value of such a test, or, rather, the opinion of the intimate connection between the various tongues is strengthened, though it may be difficult to trace their links in vocabularies compiled by Europeans. Thus, though no two languages can be apparently more dissimilar than the harsh, guttural, unpronounceable, and unwritable 'Tinnè speech, and the flowing, harmonious, and

g

iı

ir

W

fe

w

u

bι

gu

· A

Br cur

for

TRIBES.

3

s with Rocky gh the ons in asceross the entire eferred

themericans, e," and th care, test for position But as an lanpronunifferent doubts her, the een the may be es comwo lanhan the vritable us, and easily acquired tongue of the Eythinyuwuk, yet the 'Thinyu (man) of the latter may be resolved into the 'Tinnè, 'Tinye, or 'Dunnè, of the former, and the Ting-i of the Kutchin, without much philological artifice.*

Various tribes have been distinguished by peculiar names, but there is little variety in their general appearance, and few discrepancies in their dress, customs, or moral character. The Hare Indians (Kā-cho-'dtinnè) inhabit the banks of the Mackenzie, from Slave Lale downwards, and the Dog-ribs (Thling-è-ha-'dtinnè) the inland country on the cast, from Martin Lake to the Coppermine. There is no perceptible difference in the aspect of these two tribes. They meet in the same huntinggrounds at the north end of Great Bear Lake, intermarry, and their speech scarcely differs even in accent. The Hare Indians, frequenting a thickly wooded district in which the American hare abounds, feed much on that animal, and clothe themselves with its skins, while the Dog-ribs depend more upon the rein-deer for a supply of winter dresses, but in all essential respects they are the same

* Mr. Isbister says the Chepewyan tongue is "harsh and guttural, difficult of enunciation, and unpleasant to the ear." "As a language it is exceedingly meagre and imperfect."—Rep. Brit. Ass. for 1847. Mr. M. Pherson pointed out to me, as a curious coincidence, the similarity in sound of the Gaelic word for people, with the 'Dunnè of the Dog-rib Indians.

people. To the eastward of the Dog-ribs are the Red-knives, named by their southern neighbours the Tantsa-ut-'dtinne (Birch-rind people). Thev inhabit a stripe of country running northwards from Great Slave Lake, and in breadth from the Great Fish River to the Coppermine. They were also formerly in the habit of resorting to the north end of Great Bear Lake, to kill musk-oxen and rein-deer; but many of their influential men being cut off by treachery in a feud with the Dog-ribs, they have lately kept more towards the east end of Great Slave Lake. These three tribes roam northwards to the Eskimo boundary line, but mutual fears cause the two people to leave an ample neutral ground, on which neither party are willing to venture.

Other members of the 'Tinnè nation inhabit the country at the mouth of the Missinipi, and carry their furs to Fort Churchill, where they meet the Eskimos that come from the northward, and, through the influence of the traders, carry on an amicable intercourse with them, so that 'Tinnè families occasionally accompany the Eskimos to their hunting-grounds. A wide tract of barren lands intervenes between the Churchill 'Tinnè and the Red-knives, and the tribes on the Slave and Elk Rivers which resort to Fort Chepewyan. These "barren grounds" are very thinly peopled,

a

TRIBES.

the ours hey ards the were orth and being they dreat vards fears sutral og to

habit and they ward, ry on Finnè os to arren è and e and vyan. opled, and rather by isolated families who resort thither for a year or two to hunt the rein-deer than by parties associated in such numbers as to deserve the name of a tribe. Part of these wandering, solitary people resort at intervals of two or three years to Churchill for supplies, and part to Fort Chepewyan, where, from the direction in which they came, they are named Sa-i-sa-'dtinne' (Eastern or Rising Sun folks). The Athabasca 'Tinnè, named also Chepewyans, frequent the Elk and Slave Rivers, and the country westward to Hay River, which falls into Great Slave Lake. There is some difference between their dialect and that of the tribes on the Mackenzie, but not so much as to occasion any difficulty to an interpreter, versed in either tongue. The name Chipewyan has no relation to the word Ojibbeway or Chippeway, which designates an Eythinyuwuk people frequenting the coasts of Lake Superior, but has rather, I believe, its origin in the contempt felt by the warlike Crees for the less manly 'Tinné, whom they oppressed by their inroads, before commerce introduced peace between them. Chi-pai-uk-'tim (you dead dog) is a most opprobrious epithet. The appellation of "slave," given to the Dog-ribs by the same people, whose war-parties penetrated even to the banks of the Mackenzie, has a similar origin; and it has been stated in a preceding page, that

в 3

the Kolushes also called the Eskimo Kadyakers with whom they warred "slaves." To the south of the Athabascans, a number of 'Tinnè frequent the upper part of the Missinipi, where they mingle with the Crees, and in common with them trade with the posts on Lac la Ronge and Isle à la Crosse. (See Vol. I. p. 91.) The Sarsis or Circees, who live near the Rocky Mountains, between the sources of the Athabasca and Saskatchewan Rivers, are said to be likewise of the 'Tinnè stock.

Between the Peace River and the west branch of the Mackenzie are the Beaver Indians, who take their name from an affluent of the latter. Their dialect is reported to be softer than that of the other 'Tinnè, having probably been modified by their intercourse with the Crees of the prairies. Other tribes on the mountain branch of the Mackenzie differ somewhat either in language or manners from the eastern part of the nation, and have peculiar designations. The Noh'hannè inhabit the angle between that branch and the great bend of the trunk of the river, and are neighbours of the Beaver Indians. Higher up are the 'Dtcha-ta-ut-'tinne, "Mountain Indians" or "Strong-bows," who keep to the ranges of the Rocky Mountains, and the Tsilla-ta-ut-'tinne, or "Brushwood-people."

Between the trunk of the Mackenzie, on the 65th parallel, and the Rocky Mountain ranges,

t 1 C tltla ta A h T. is dy all do of the pa res bas kne fur det

TRIBES.

kers h of the ngle rade à la *cees*, the vers, ranch take Their other their

take Their other their Other enzie nners have it the nd of of the *ta-ut-*' who nd the

> n the inges,

dwell a tribe named Dahā-'dtinnè by the Dog-rib Indians, and Noh'hai-è by the Kutchin. They descend the Gravel River to come to Fort Norman, and are ill understood by the Dog-rib interpreters there. In the first volume (p. 180.) I have mentioned, on the authority of Mr. M'Kenzie, that the Dahā-'dtinnès name themselves in their own tongue Cheta-ut-tdinnè, which indicates their identity with the Strong-bows, both being mountaineers. Further down the Mackenzie, near the 65th parallel, another small tribe also descends from the mountains to visit Fort Good Hope, and is named Amba-ta-ut-'tinne', or "Sheep-people," because they hunt the Ovis montana on the mountain-tops. These people speak a dialect of the 'Tinnè, which is well understood by the Hare Indians.

This enumeration of the various 'Tinnè tribes dwelling on the east side of the Rocky Mountains, all of whom believe that they are sprung from a dog, will give some idea of the geographical extent of the nation. It is not my intention to speak of them severally, as my personal acquaintance is too partial to enable me to state correctly in what respects they differ from each other. The Athabascans or Chepewyans proper have been so long known, and so often mentioned by writers on the fur countries, and Hearne has given so many details of the habits of the 'Tinnè of Churchill, and

of the tribes he encountered in his journey over the barren grounds, that I could add little of importance; I shall, therefore, restrict my remarks to the Dog-ribs and Hare Indians, who resorted to Fort Franklin and Fort Confidence during my residences on Great Bear Lake.

These people possess more regular features than the Eskimos, with, at the same time, a greater variety among individuals, many of whom have good profiles. Taken as a whole, they exhibit all the characteristics which we observe in the red races dwelling further south; but their inattention to personal appearance, want of cleanliness, and their abject behaviour, give them a very inferior aspect, particularly when in the company of white people. For they possess the whine and air of accomplished beggars, and their solicitations are constant as long as they have any hope of gain. The women are inferior to the men in height, features, and care of their dress; for, dirty as the men generally are, they do paint their faces and wear ornaments on festive occasions, while few of the women take so much trouble. Most of the latter, however, are tatooed on the chin, or at the angles of the mouth.

1

9

1

n

N

16

iı

n

ee

tł

 \mathbf{sl}

tl

cl

lo

The clothing of the men in summer is rein-deer leather, dressed like shammy, and is beautifully white and soft when newly made. A shirt of this

DRESS.

y over ttle of emarks rted to ng my

es than greater n have exhibit in the ieir incleanlia very mpany ine and itations of gain. ht, feaas the ces and few of of the at the

in-deer atifully of this

material, cut evenly below, reaches to the middle; the ends of a piece of cloth secured to a waistband, hang down before and behind; hose or Indian stockings descend from the top of the thigh to the ankle; and a pair of mokassins or shoes of the same soft leather, with tops which fold round the ankle, complete the costume. When the hunter is equipped for the chase, he wears, in addition, a stripe of white hare-skin, or of the belly part of a deer-skin, in a bandeau round the head, with his lank, black elf-locks streaming from beneath; a shot pouch, suspended by an embroidered belt, which crosses the shoulder; a fire-bag or tobacco-pouch tucked into the girdle; a pair of mittens; and a long fowling-piece in its coat thrown carelessly ecross the arm or balanced on the back of the neck. The several articles here enumerated are ornamented at the seams and hems with leathern thongs wound round with porcupine quills, or are more or less embroidered with bead-work, according to the industry of the wife or wives. One of the young men even of the slovenly Dog-rib tribe, when newly equipped from top to toc, and tripping jauntily over the mossy ground with an elastic step, displays his slim and not ungraceful figure to advantage. But this fine dress, once donned, is neither laid aside nor cleaned while it lasts, and soon acquires a dingy look, and an odour which can be perceived from

10

some distance. In the camp a smoky, greasy blanket of English manufacture is worn over the shoulders by day, and forms, with the clothes, the bedding by night.

In winter the skins of fawn rein-deer, retaining the hair, are substituted for the shammy leather, and a large robe of the same material is thrown over the shoulders, and hangs down to the feet, in place of the blanket. As the preparation of so much leather and dressed fur keeps the women busy, they are glad to use English cloth, of blue, red, or green colours, or Canadian capots of white or blue cloth, which they acquire at the trading posts in exchange for venison or furs. But with regard to the winter dress especially, the substitution of the produce of the English loom for their native leather is a loss both of comfort and of appearance.

The women's dress resembles the men's, except that the shirt is somewhat longer, and, for the most part, is accompanied by a petticoat which reaches nearly to the knee.

The form of the dress here described is common to the whole 'Tinnè nation, and also to the Crees and Dakotas, though the material varies with the district; moose deer, red deer, or bison leather, being used in the south and west, where those animals abound; and the Hare Indians make their

WOMEN.

greasy ver the nes, the

taining leather, thrown feet, in of so women of blue, f white trading ut with substiom for ort and

except for the which

ommon c Crees ith the eather, those e their shirts of the skin of the hare. This, being too tender to be used in the ordinary way, is torn into narrow strips, which are then twisted slightly, and plaited or worked into the required shape. I have noticed no process among the northern Indians that approaches so nearly to weaving as the manufacture of the white hare-skin shirts.* Such is the closeness and fineness of the fur, that they are exceedingly warm, notwithstanding the looseness of their texture. Though the dress of the southern Indians is after the same pattern with that of the 'Tinnè, the Kutchin, both in the interior and on the coast, form, as has been already mentioned, the hose and shoes of the same piece; thus initating the Eskimo boot, though with a different material.

The Dog-rib men and women leave their hair without other dressing than simply wiping their greasy hands on the matted locks, when they have been rubbing their bodies with marrow, which they occasionally do.

The Hare Indian and Dog-rib women are certainly at the bottom of the scale of humanity in North America. Not that they are treated with cruelty, for the 'Tinnè are not a cruel people, but that they are looked upon as inferior beings, and in this belief they themselves acquiesce. In early

* The Kenaiyer of Cook's Inlet are said to weave the wool of the mountain goat (*Capra americana*) into a stuff used for clothing.

infancy the boy discovers that he may show any amount of arrogance towards his sisters, who, as soon as they can walk, are harnessed to a sledge, and inured betimes to t' labours which are their inevitable lot through ' ;; while the future hunter struts in his tiny snow shoes after the men, and apes their contempt of the women. The women drag the sledges alone or aided by dogs, clear the ground for the tent, cut poles to extend the lodge or tent-skins upon, collect fire-wood, bring water, make all the dresses and shoes, clean the fish, and smoke or jerk the venison for its preservation. They also cook both for themselves and their husbands, the 'Tinnè not holding the opinion of the Kutchin that a man ought not to eat meat prepared by a woman. Neither are the 'Tinnè women altogether precluded from eating with the men; though in times of scarcity the man would expect to be first fed, as it is a maxim with them that the woman who cooks can be well sustained by licking her fingers. The women are not, however, generally discontented with their lot, and better days are certainly dawning upon them, as the opinions of the traders are beginning to tell visibly on the Notwithstanding their servile conwhole nation. dition they are not without influence over the stronger sex; and they seldom permit provisions or other articles to be disposed of without ex-

pro car

to shr fea or dre all, are wh are of the the out on and the sinl son nea ram grea any hav and

Ί

DISPOSITIONS.

how any who, as a sledge, are their e hunter nen, and e women clear the the lodge ng water, fish, and servation. heir huson of the neat prenè women the men; ld expect that the y licking er, genetter days opinions ly on the vile conover the rovisions hout expressing their thoughts on the matter with much earnestness and volubility.

Few traces of the stoicism popularly attributed to the red races exist among the Dog-ribs: they shrink from pain, show little daring, express their fears without disguise on all occasions, imaginary or real, shed tears readily, and live in constant dread of enemies, bodied and disembodied. Yet all, young and old, enjoy a joke heartily. They are not a morose people, but, on the contrary, when young and in a situation of security, they are remarkably lively and cheerful. The infirmities of age, which press heavily on the savage, render them querulous. They are fond of dancing, but their dance, which is performed in a circle, is without the least pretensions to grace, and is carried on laboriously with the knees and body half bent and a heavy stamping, having the effect of causing the dancers to appear as if they were desirous of sinking into the ground. It is accompanied by a song resembling a chorus of groans, or pretty nearly the deep sigh of a pavier as he brings his rammer down upon the pavement. They are great mimics, and readily ape the peculiarities of any white man; and many of the young men have caught the tunes of the Canadian voyagers, and hum them correctly.

They are an unwarlike people, and averse to

a

p

C

SC

a1

rd

k

m

m we

co

co

qı ot

ci

a av

an

di

in

m

 $^{\text{th}}$

ste

shedding blood; yet, as they do not meet their foes in open warfare or man to man, their very timidity impels them to treachery or a violation of the laws of hospitality, when, by long-continued oppression and the loss of relatives, they have been driven to retaliate upon the few individuals or families of the domineering tribe who were living in confidence among them. This remark applies directly to their feud with the Redknives, who for many years resorted to the huntinggrounds of the Dog-ribs, tyrannised over them, and carried away their women. This was long borne, but at length, some lives having been lost in the contests which occasionally ensued, the Dog-ribs, watching their opportunity, cut off several leading Red-knives and their families, who, not dreading any thing at the time, were scattered among the Dog-rib encampments. The details of these reprisals give a curious insight into the character of the people. Some of the victims, deprived of the means of resistance, and aware of their intended fate, travelled for a whole day with the hostile party; but the latter required to have their passions roused by altercation before they acquired sufficient boldness to perpetrate the deed, and were finally incited to its commission by the sufferers demanding to be killed at once if their death was intended, for they would go no further. When

SOCIALISM.

eet their eir very violation ong-cones, they few indiibe who This . the Redhuntinger them, vas long been lost ied, the off sevees, who, scattered letails of the chaleprived of their with the ve their acquired ed, and the sufir death When the husbands and grown men were killed, the Dogribs argued that pity impelled them to slaughter also the wives and children, who would be unhappy and perish for want, having lost their means of support. To a people who could no longer support the tyranny of their bolder neighbours, nor combine so as to repel aggression by force, treachery seemed to be the only mode of obtaining redress; and in fact the extent to which they carried their reprisals effectually broke the spirit of the Redknives, and drove them to a distance.

The Dog-ribs are practical socialists; and, as much of the misery they occasionally experience may be traced to this cause, the study of the working of such a system may be instructive in a community like this, whose members owe their condition in the social scale solely to their personal qualities, and not to inheritance, favour, or the other accidents which complicate the results in civilised life. Custom has established among them a practice universally acted upon, - that all may avail themselves of the produce of a hunter's energy and skill; and they do not even leave to him the distribution of his own game. When it is known in the camp that deer have been killed, the old men and women of each family sally forth with their sledges, and, tracing up the hunter's footsteps to the carcases of the animals he has slain,

proceed to divide them among themselves, leaving to the proper owner the ribs, which is all that he can claim to himself of right. He has also the tongue, which he takes care to cut out on killing the deer. It is not in the power of these people to restrain their appetites when they have abundance; and the consequence is, that when the chase is successful, all the community feast and grow fat, however little many of the men-and there are not a few idle ones - may have contributed to the common good. The hunter's wife dries the rib-pieces, after cutting out the bone, in the smoke, or over a fire, to carry to a fort for the purposes of trade; but, unless there is a superabundance, little provision is made by the party for a time of scarcity, which is sure to arrive before long; since the deer, when much hunted, move to some other district. Taught by their frequent sufferings on such occasions, the more active hunters frequently withdraw themselves and their families from the knowledge of the drones of the community, leaving them at some fishing station, where, with proper industry, they may subsist comfortably. A fish diet is not, however, agreeable to the palates of these people for any length of time; and, as soon as rumours of a hunter's success reach them,-which they do generally much exaggerated by the way, -a longing for the fleshpots is instantly excited, especially among the old,

a el di hı \mathbf{th} re m un ma the spe the Su car du and the opi of dre dur mos full unq sho[.] plia sona loss V

AFFECTION FOR PARENTS.

aving 1 that s also out on f these y have en the st and -and ibuted ies the smoke, oses of e, little carcity, e deer, listrict. h occathdraw edge of it some y, they t, howfor any unter's y much e fleshhe old,

and a general movement to the hunting-ground ensues. If, on their march, the craving multitude discover a hoard of meat stored up by any of the hunting parties, it is devoured on the spot; but they are not always so fortunate. Before they reach the scene of anticipated abundance, the deer may have gone off, followed by the hunters, with uncertain hopes of overtaking them, and nothing remains for the hungry throng, including the old and the lame, but to retrace their steps, with the prospect of many of them perishing by the way, should their stock of food have been quite exhausted. Such occurrences are by no means rare; they came several times under our immediate notice during our winter residence at Fort Confidence, and similar facts are recorded by Mr. Simpson of This gentleman expresses his the same tribe. opinion that the charge made against this nation, of abandoning their infirm aged people and children, had its origin in the sauve qui peut cry raised during a forced retreat from some one of these most injudicious excursions; and I am inclined fully to agree with him; for I witnessed several unquestionable instances of tenderness and affection shown by children to their parents, and of compliance with their whims, much to their own personal inconvenience. The grief they show on the loss of a parent is often great and of long conti-VOL. II.

18

nuance, and it is the custom both for men and women to lament the death of relations for years, by nightly wailings.

Hospitality is not a virtue which is conspicuous among the Dog-ribs, who differ in this respect from the Eythinyuwuk, in whose encampments a stranger meets a welcome and a proffer of food. It is not customary, however, for the Dog-rib to receive the traveller who enters his tent with the same show of kindness. If he is hungry, and meat hangs up, he may help himself without eliciting a remark, for the 'Tinnè hold it to be mean to say much about a piece of meat; or he may exert his patience until some cookery goes on, and then join in the meal; and should there be venison at hand, he will not have long to wait, for every now and then some one is prompted to hang a kettle on the fire, or to place a joint or steak to roast before it.

Another habit which darkens the shade in the character of these Indians is that of lying, which they carry to such an extent, even among themselves, that they can scarcely be said to esteem truth a virtue. If a young man has been successful in his morning's hunt in a time of famine, he does not rush into his family circle with joy beaming on his countenance, to tell that there is food, but, assuming an aspect of sadness, squats

de ho th

v

0

A

p

b

it

it

sa

ar

HONESTY.

nen and or years,

spicuous respect oments a of food. Dog-rib cent with gry, and out elicitmean to nay exert and then enison at very now a kettle to roast

le in the g, which ng themo esteem een sucf famine, with joy there is s, squats

himself in silence beside the fire. The women with doubt and anxiety examine his shoes and dress for spots of blood, that may betoken the death of an animal, but discovering none, put the question, "Did you see no deer ?" "Not one, the deer are all gone, not a single footstep was to be seen." When the colloquy has continued for a time, and hope seems to be extinct, he then draws out from beneath his shirt two or three tongues, as the case may be, and says with an air of the utmost indifference, "You may go for the meat." It is not, however, merely at such times, and to enhance the pleasure by previous disappointment, that truth is violated, but on almost every occasion: and the skill of an Old Bailey practitioner would find exercise in eliciting facts from the mass of contradictions with which they overload them. A story which was at first a pure invention, or perhaps, a perversion of some simple occurrence, becomes so changed by the additions it receives in its transmission from individual to individual, that it deceives the originators, and if it bears on the safety of the community, may spread consternation among them, and occasion a hasty flight.

It is pleasant, instead of dwelling longer on this defect, to turn to another feature—their strict honesty; the practice of the 'Tinnè with regard to the property of white people differing remarkably

c 2

CHEPEWY. NS.

from their northern neighbours, the Eskimos, and their southern ones, the Crees, though the temptations to which they are exposed are equally great. No precautions for the safety of our property at Fort Confidence were required. The natives carefully avoided touching the magnetic instruments, thermometers, and other things placed outside the house, and could be trusted in any of the rooms without our finding a single article dis-Our dining-hall was open to all comers; placed. and though the smallness of our separate apartments caused us to exclude hangers on, new comers were permitted to satisfy their curiosity respecting our occupations, and they always squatted themselves down at the door, and looked on in silence, wondering, as we were told, at our constant writing. From M. La Flèche, the intelligent missionary at Isle à la Crosse, I received a similar character of the southern part of the nation, who, if they find any article left by the voyagers on the portages, are sure to bring it in to be claimed at the forts.

n

p

t

S

p

0

tł

Ce

b

tł

a

aı

pe

Of the peculiarities of their religious belief I could gain no certain information. The interpreters to whom I applied for assistance disliked the task, and invariably replied, "As for these savages, they know nothing; they are ignorant people." The majority of the nation recognise a

RELIGIOUS BELIEF.

21

os, and tempequally ir pro-'he naetic inplaced any of cle disomers; apartcomers pecting themsilence, onstant nt missimilar h, who, rers on claimed

belief I interlisliked r these morant gnise a

"Great Spirit," at least by name, but some doubt his existence, assigning, as a reason for their atheism, their miserable condition; or they say, "If there be such a being, he dwells on the lands of the white people, where so many useful and valuable articles are produced." With respect to evil spirits, their name in the Dog-rib country is The 'Tinnè recognise them in the Bear, legion. Wolf, and Wolverene, in the woods, waters, and desert places; often hear them howling in the winds, or moaning by the graves of the dead. Their dread of these disembodied beings, of whom they spoke to us under the general name of "enemies" is such, that few of the hunters will sleep out alone. They never make any offerings to the Great Spirit, or pay him an act of adoration; but they deprecate the wrath of an evil being by prayer, and the sacrifice of some article, generally of little value, perhaps simply by scattering a handful of deer-hair or a few feathers.

The dead are not burnt, after the manner of the Kolushes, but are buried. In lamenting for deceased relatives the mourners sometimes gash their bodies or limbs with knives, but more rarely now than in old times. It was formerly the custom, on a death occurring, for the family to abandon every article they possessed, and betake themselves in a perfectly destitute condition to the nearest body

c 3

of their own people, or to the trading post. The advice of the traders is gradually breaking down this practice.

Shamanism does not seem to exert the important influence upon the 'Tinnè that it does among the Asiatic Tchukche, the Kutchin, or the Eskimos. There are men in the nation, with the reputation of sorcerers, who profess to have power over spirits; but they have but little personal influence, and are generally of small repute, to which, perhaps, the contempt of the white people for their arts con-A belief, however, in the power of the tributes. Eskimos and of strange Indians to hurt them by incantations, or "bad medicine," prevails. White people are said to be exempt from such dangers, their "medicine" being the most powerful. The "conjurers" are occasionally employed to cure the sick, and I suppose on such occasions receive some reward; but I heard of no instance of their being beat and coerced to influence the spirits favourably in the manner that the Asiatic Tchukche are reported in Baron Wrangell's work to deal with their shamans.

Among the Crees the conjurers perform a much more prominent part than with the 'Tinnè, and their practices come frequently under the observation of residents on the lands of that people; but I never saw one exhibit among the Hare Indians,

0 e: as th di ho or de th im \mathbf{th} nu rat vei ski rea gai fro son Th wit

c

C

MARRIAGES.

The down

ortant ng the kimos. tation pirits; nd are os, the s conof the em by White angers, The are the e some being urab!y are rewith

much è, and pservae; but ndians, Dog ribs, or Red-knives, in the course of four or five years passed among them, though I have many times seen some of the old men throw trifling articles into the water, to procure a fair wind, or secure a safe passage across a lake or down a rapid.

From a people so liable to be actuated by fears of imaginary evils no steady line of action can be expected, and the Dog-ribs are in reality as volatile as children. When accompanied by a white man, they will perform a long journey carefully to a distant post; but we found, by experience, that however high the reward they expected to receive on reaching their destination, they could not be depended upon to carry letters. A slight difficulty, the prospect of a banquet on venison, or a sudden impulse to visit some friend, were sufficient to turn them aside for an indefinite length of time.

In general, the 'Tinnè have only one wife, the numbers of the sexes being equal, or the males rather predominating. The women are married very young, but the man must have shown some skill in hunting before he obtains a helpmate readily. The consent of the parents is usually gained by the suitor, and is seldom withheld from a man whose activity promises the old folks some addition to their comforts or consequence. The woman's wishes have, perhaps, some weight with her parents, but I could not ascertain that

c 4
CHEPEWYANS.

r

t

t

П b

a

la

c

C

to

m

W

m

b

bo

cc

st

w. ev

tr

pa

ig

ke

th

any show of courtship* was made, or that her disinclination was allowed to interfere with the man's determination to take her, if the parents did not oppose. No ceremony attends the union. Hearne says, that it is the established etiquette among the Eastern 'Tinnè for the woman to affect unwillingness to change her condition, and for the man to rush into her father's tent, and drag her off by the hair of the head. We witnessed no scene of this kind among the Dog-ribs, but more than once saw a stronger man assert his right to take the wife of a weaker countryman. Any one may challenge another to wrestle, and, if he overcomes, may carry off his wife as the prize. The younger children generally follow the fortunes of the mother, but the father may retain them if he chooses. In such contests, it is suspected that the wife sometimes prompts the aggressor; but I have been told-for I never actually witnessed one of these wrestling matches - that she looks on with composure and impartiality, and does not insult her late master with a display of pride on being the object of such a struggle, the causa teterrima belli. The bereaved

* The term "dear," or "beloved," is said to be unknown in the language; and Captain Lefroy, who tried to ascertain if it was so, says, "I endeavoured to put this intelligibly to Nannette, by supposing such an expression as ma chère femme; ma chère fille. When at length she understood it, her reply was (with great emphasis), "I disent jamais ça; i disent ma femme; ma fille."

 $\mathbf{24}$

MOOSE-HUNTING.

25

hat her ith the parents e union. tiquette o affect for the g her off scene of an once the wife hallenge ay carry children ier, but In such metimes ld_for restling ure and master of such ereaved

unknown aseertain ligibly to e femme ; her reply lisent ma

husband meets his loss with the resignation which custom prescribes in such a case, and seeks his revenge by taking the wife of another man weaker than himself. From a passage in one of Mr. Murray's letters, I infer that this practice extends to the Kutchin, but it is unknown among the Cree tribes, and does not exist among the Eskimos. The 'Tinne' are said to be jealous of their wives; but rather, I believe, lest they should be enticed away, than from any nice sense of honour. The laxity of morals, however, with respect to female chastity, which prevails in the Eskimo tribes is not conspicuous in the 'Tinnè, and is, perhaps, contrary to the national character, though some corruption may have crept in through their acquaintance with white people.

Before the introduction of articles of European manufacture, the 'Tinnè caught fish with hooks of bone, or speared them with weapons pointed with bone or copper. Some of their fish harpoons were constructed very artistically. They also used, and still continue to use, nets made of lines of twisted willow bark, or thin stripes of deer-hide cut very evenly. Nets are unknown among the northern tribes west of the Mackenzie, and some of the parties of the Eskimos that we saw declared their ignorance of their use. On the banks of the Mackenzie and other rivers frequented by moose-deer, these animals are hunted in spring by a small

CHEPEWYANS.

26

breed of dogs, which run lightly over the crusted snow, and hold the animal at bay until the Indian comes up in his snow shoes. At other times of the year, the success of the Hare Indians and Dogribs in killing the moose is small, as they have not the skill of approaching so wary an animal which the Athabascans and Crees possess. Rein-deer are captured in pounds and by nooses, but are in the present day more generally killed with the fowlingpiece, which is also the weapon used against the musk-ox. The pounds are formed on the verge of the woods, and are made with much less labour than those of the Kutchin; yet, as they need the exertions of all the community for their construction, the indolence of the major part causes them to be rarely made. The black bear is snared or shot, but few of the Dog-ribs will venture to attack the "brown barren-ground bear," whose fierceness, or, as they say, " potent medicine," appals them. It is killed by them, however, without risk when it is detected hybernating under the snow in spring.

Order is maintained in the tribe solely by public opinion. It is no one's duty to repress immorality or a breach of the laws of society which custom has established among them, but each opposes violence as he best may by his own arm or the assistance of his relations. A man's conduct must be bad indeed, and threaten the general peace,

CHIEFS.

crusted Indian imes of nd Dogave not l which leer are in the fowlinginst the verge of labour the exruction, hem to or shot, ack the ness, or, em. It ien it is ring. public horality custom opposes or the et must peace,

before he would be expelled from the society; no amount of idleness, nor selfishness, entuils such a punishment. Superior powers of mind, combined with skill in hunting, raise a few into chiefs, under whose guidance a greater or smaller number of families place themselves; and a chief is great or little, according to the length of his tail. Ilis clients and he are bound together only by mutual advantage, and may and do separate as inclination The chief does not assume the power of prompts. punishing crimes, but regulates the movements of his band, chooses the hunting-ground, collects provisions for the purchase of ammunition, becomes the medium of communication with the traders, and extends his sway by a liberal distribution of tobacco and ammunition among his dependents. At present, the rank of a chief is not fully established among his own people until it is recognised at the fort to which he resorts. The Company send in annually a number of red coats, ornamented with lace, for presents to the chiefs, which are worn as badges of office on great occasions. The power of a chief varies with his personal character. Some have acquired an almost absolute rule, by attaching to themselves in the first instance an active band of robust young men, and using them to keep in order any refractory person by claiming his wife after the custom of the tribe. It is in vain in

CHEPEWYANS.

such cases that the poor husband, dreading to be deprived of his most valuable property, retires to a remote hunting-ground; for he is sure to receive a message, from some passing Indian, expressive of the chief's intentions; and he generally comes to the conclusion that submission is the best He is certain to fall in with the chief and policy. his band sooner or later, either as he goes to the fort for supplies of ammunition or elsewhere. A free expenditure by the chief of the presents he receives from the traders, and even of the produce of his furs, is a main bulwark of his authority, in addition to the skill which he must possess in the management of the various tempers with which he has to do.

The sounds of the 'Tinnè language can scarcely be expressed by the English alphabet, and several of them are absolutely unpronounceable by an Englishman. In my attempts to form a vocabulary I had great difficulty '- distinguishing several words from one another which had dissimilar sounds to the native ear, and were widely different in their signification. A Dog-rib or Athabascan appears, to one unaccustomed to hear the language, to be stuttering. Some of the sounds must have a strong resemblance to the Hottentot cluck, and palatal and guttural syllables abound in the language. Vocabularies of this tongue cannot be greatly de-

or 'Ti md the ert \mathbf{As} ent a l yer we the gre out occ con bei son for to reli don

184

Mo

men

and

beli

pe

INTRODUCTION OF CHRISTIANITY.

g to be retires sure to lian, exgenerally the best hief and s to the here. A sents he produce ority, in s in the which he

scarcely everal of Englishry I had rds from to the r signifi-, to one be stutstrong palatal nguage. atly depended upon, as no two people will agree on the orthography.

With respect to the future prospects of the 'Tinnè, the nation in general may be said to be more docile and confiding, and more directly under the influence of the traders, or of missionary exertions, than their southern neighbours, the Crees. As yet, Roman Catholic missionaries alone have entered the 'Tinnè country, and they have already a large number of nominal converts. For some years Canadian priests from the Red River colony went annually to Methy Portage, where many of the Athabascans and Churchill River 'Tinnè congregate at the usual season of transporting the outgoing furs and incoming supplies. On these occasions, numbers of the Indians were baptized, a considerable inducement to submit to the rite being the present of a piece of tobacco, or perhaps some vague notion of the protection thereby afforded against evil influences. There was no time to instruct them in the truths of the Christian religion, and this could be but very imperfectly done through the medium of interpreters. In 1846, however, the Roman Catholic mission under Monsieur La Flèche was established, as has been mentioned in a preceding chapter. This gentleman and his associate, Monsieur Taschè, members, I believe, of the Society of Jesus, applied themselves

CHEPEWYANS.

to the study of the 'Tinnè language, and were soon enabled to teach many of their converts to read and write.

By sympathising with their people in all their distresses, taking a strong interest in every thing that concerns them, by acting as their physicians when sick, and advisers on all occasions, the priests of the mission have gained their entire confidence. It is not likely that Protestant missionaries, coming later into the field, will succeed in introducing their more spiritual but less imposing form of worship among a people whose first teachers have been so successful.

When the fur traders first penetrated to the Elk River, the Athabascans had only a small breed of dogs useful for the chase, but unfitted for draught; and the women did the laborious work of dragging the sledges. Now the cultivation of a stouter race of dogs has in some respects ameliorated the lot of the females, and within a few years the acquisition of horses by many of the natives on that river has introduced a still greater improvement. Houses are beginning to be built, and the more provident and staid of the people have fixed homes to retire to. With the means of securing their property and provisions, new ideas respecting them spring up, and a revolution in the opinions of the nation is evidently in progress. Recently, also, it has been

the ma nav to bro hab wel res A Ind of ti wya 0 inha the New thei The serv com the. trial is at simi Tsit Sim said thei

WESTERN TRIBES.

soon read

their thing icians riests lence. oming ucing m of have

e Elk eed of aght; gging race lot of ition has s are and e to. and up, n is been the policy of the Hudson's Bay Company to employ many of the young natives, during the summer, in navigating their boats to the depôts, and back again to the outposts. By these trips prejudices are broken down; the youth acquire information and habits of labour and steady industry, and, being well paid, the clothing they purchase gives them respectability in the eyes of their countrymen. A generation has passed away since the whole Indian country was demoralised by the opposition of trading companies, and the present race of Chepewyans are ignorant of the use of spirituous liquors.

Of the nations belonging to the 'Tinnè stock who inhabit the country west of the Rocky Mountains, the *Tā-kuli* or Carriers occupy the greater part of New Caledonia. They subsist chiefly on fish, and their name denotes people employed on the waters. They burn their dead; the widow becomes the servant of the relations, is harshly treated, and is compelled to carry about with her for several years the ashes of the deceased. When the time of her trial ends, a feast is made by the kindred, and she is at liberty to marry again. A custom somewhat similar prevails among the Chippeways. The Tsitka-ni, who dwell between the Stikeen and Simpson's Rivers to the north of the Carriers, are They bury said also to be of the same stock. their dead, and are hunters.

31

in

CHEPEWYANS.

In addition to these tribes, a detached portion of the 'Tinnè people is mentioned by Dr. Latham, under the name of Southern Athabascans. They occupy the sea-coast from the north bank of the Oregon southwards, to the River Umqua, in $43\frac{1}{2}^{\circ}$ of lat. For an account of these, I must refer the reader to the works of the author just named, and to the Transactions of the American Ethnological Society from which he quotes.

Dr. Latham may, also, be consulted for an account of four or five isolated languages, spoken by tribes that interpose between the North and South Athabascans to the west of the Rocky Mountains, and for notices of the inhabitants of the Archipelago skirting that coast. The Kolush language ends, he thinks, at the north end of King George's Archipelago.

The Chenooks, one of these isolated people, are noted for their habit of flattening the foreheads of their infants artificially, a custom which crosses the continent southwards to the coast of Florida, and was practised, though not exactly in the same way, by the extinct Peruvian races of Lake Titicaca.*

* Among some good examples of flattened skulls from the west coast of America, in the Museum at Haslar, there is the remarkable one of Comeomly, the hero of Washington Irving's Astoria.

G R CI D Тн or l the ing of t thei witl ider lena cou lina of t to a terr gon the v

NAT

CHAP. XIV.

EYTHINYUWUK, OR CREES AND CHIPPEWAYS.

NATIONAL NAMES. -- DIVISION. -- TRIBES. -- TERRITORY. -- WARS WITH THE MENGWE. -- CONVENTIONAL CHARACTER NOT TRUE. -- PERSONS. -- GAIT. -- CRIMES. -- WABUNSI. -- WIGWAMS. -- RELI-GIOUS BELIEF. -- VAPOUR BATHS. -- EVERLASTING FIRE. -- ITS RITES. -- USED IN SICKNESS. -- ITS PRIESTS. -- ITS ORIGIN. --CHIEF SUN. -- POLICY. -- CALUMET. -- MAIZE. -- FOOD. -- REIN-DEER. -- BISON. --- WHITE-FISH. -- EARTH-WORKS. -- FOTTERY. --LANGUAGE. -- HALF-BREEDS. -- COLONY OF RED RIVER OR OSNABOYA. -- SPIRITUOUS LIQUORS.

THE people who designate themselves Eythinyuwuk or Ininyu-wë-u, occupy the country lying between the Rocky Mountains and Hudson's Bay, and reaching from the 'Tinnè boundary down to the plains of the Saskatchewan and valley of the St. Lawrence; their hunting-grounds on the plains interlocking with those of the Dakotas or Sioux. They are identified as a nation with the Algonkins and Lennilenape or Delawares, who once owned the whole country east of the Mississippi as far south as Carolina, but who, blighted by the precocious expansion of the Anglo-Saxon colonists, have dwindled down to a few remnants of mixed blood. The generic term Algic, taken from the root of the word Algonkin, has been employed by the philologists of the United States to comprehend all the tribes who

D

VOL. II.

ortion tham, They of the $43\frac{1}{2}^{\circ}$ er the l, and ogical

poken n and founof the olush King

e, are ds of osses orida, same Titi-

m the is the ving's $\mathbf{33}$

speak dialects of the Algonkin tongue, and whose southern limits are stated by Schoolcraft to be conterminous with the *Catawbas*, *Creeks*, *Cherokees*, *Chactas*, and *Chickasas*. The tract which they occupied in the year 1600 includes the whole area of the United States east of the Mississippi, north of these nations, excepting the grounds of several tribes of the Iroquois race, north and south.* In 1603, when the French settled in Canada, the Algonkins, according to Colden, were "the most warlike and polite nation in all North America."

The national name of this people is derived, according to the custom of the Americans, from the word "man," which is in different dialects *Ethinyu*, *Ethin-u*, *Inin-yu*, or *Ininè.* † According to Schoolcraft, they do not call themselves Unischauba ‡, or "aborigines," but, on the contrary, have a tradition current among the southern

* The Abenakis, Etchemins, and some kindred tribes located to the south of the Gulf of St. Lawrence, and the Hochungarras, or Winnebagoes, and Wyandots, to the north-west, belonged to the Iroquois stock.

† Dr. Latham traces affinities between the terms for "people," in several languages. The similarity of the terms *Inuk* or *Inuit*, and '*Tinnè*, or '*Tinyè*, to some of the above is obvious. Mr. Howes makes *Ethin'u* = $\epsilon \theta \nu - o c$. From *Ethinu* comes *ethiniseu*, "manly;" "wise," indicating the opinion the Crees have of their own nation.

[‡] From unisha, "common" or "general," ininè, "a man," and aub, a generic particle denoting "light," "virility," or "life." Schoolcraft.

m ho of of Hu or nat rite \mathbf{Th} \mathbf{the} the not cou not Hu Am feuc and trib Red

the a cordi † so-ca his f south

 $\mathbf{34}$

NATIONAL NAME.

members of the nation, that the country they now hold was previously possessed by the *Alligèwi*, of whom the name only remains in the appellation of the Alleghany Mountains.

Before the European invasion, the Dakota, Huron, Oneida, Mohawk, and Iroquois association. or Mengwe'*, generally known as the "Five nations," had penetrated into the Eythinyuwuk territory by way of the Missouri and St. Lawrence. The contests by which the Mengwè established themselves in a district, surrounded on all sides by their enemies, must have been severe; and they are not even now ended, but are carried on in the country between the Saskatchewan and Missouri, notwithstanding the persevering efforts of the Hudson's Bay Company and the officers of the American outposts to suppress them. Deadly feuds exist between the Blackfeet Eythinyuwuk, and the Mandans, Minetares†, and other Dakota tribes which frequent the bison plains; and on the Red River of Lake Winipeg, fatal conflicts took

† Called also Absoroka. These and the Mandans are the so-called Welch Indians, said to be descended from Madoc and his followers. The same origin has been attributed to the southern *Tuscarora*, of the Iroquois stock.

D 2

whose to be rokees, they e area north everal Īn a, the most ca." erived. from ialects ing to Unistrary, athern

located garras, nged to

beople," inuk or bvious. ethinis have

man," y," or

^{*} This confederacy assumed the appellation of *Mengwè* from the ancient Iroquois title, *Ongwè-honwe*, which signifies, according to Colden, "men surpassing all others."

place in the last year, between the Chippeways and the Sioux, who are Dakotas.

Dr. Latham states that the *Shyennes*, who dwell on the head waters of the Yellow Stone and Platte Rivers, are of the Algonkin race, though insulated by other people from the rest of their nation; and that there is in like manner a southern detachment of Iroquois (*Tuscaroras*, &c.), between whom and their countrymen the Delawares interpose; but as I mean to restrict myself to the St. Lawrence Valley, and the country lying north of it, I must refer the reader who wishes for a general classification of the native American races, to the "Natural History of the Varieties of Man," by the learned author just named.

The various tribes of Eythinyuwuk assume local designations from the rivers or other remarkable features of the districts they inhabit, and they have also names of more general import. Thus the northern ones, who border on the 'Tinnè, call themselves Nathèwy-withinyu, Nehethè-wuk, or Nithè-wuk*, "Exact or complete men." These are the Crees of the fur traders; and Mr. Howse, though he does not publish the grounds of his opinion, considers them to be the stem of the Algonkin race. On the south of these, in the country extending from Lake Winipeg to the south side

* From Ni, "exactly."

of (C or na de

Ch Ad Ag † the trib

Na Mus Ath Omi Pege Miss Wus Po-i Pem Mas

Ne n Chuk Pene Was Weu Kà-s

DIVISIONS.

eways

dwell Platte ulated ; and etachwhom rpose; . Lawof it, I general to the by the

assume emarkid they Thus nè, call or *Ni*ese arc Howse, of his he Alcounth side of the basin of Lake Superior, dwell the Odchipewa (Chippeways or Ojibbeways, called also Sauteurs * or Sotoos). A third great division of the nation name themselves Lenni-lenape (Delawares), which denotes "Uncommon men."[†]

* Spelt by some Canadian writers Saulteaux. A populous Chippeway tribe frequent the Saut Ste. Marie to feed on the Adikumayg or Attikamaig (White-fish, Coregonus sapidissimus, Agassiz), whence the name of "Cascade people" (sauteurs).

[†] The following list, drawn up in 1770 by Mr. Hutchins of the Hudson's Bay Service, gives the names of the Eythinyuwuk tribes then trading with the Hudson's Bay Company :---

KEISCATCHEWAN NATION.

Names of Tribes or Places.		Districts they inhabit.		
Muska-siskow	-	Saskatchewan prairies.		
Athăpèskow	-			
Omiska-sipi	-	Beaver River and Lake.		
Pegogĕ-mè-u nipi -	-	Muddy Lake, Moose Lake.		
Misi-nipi	-	Churchill or English River.		
Wuskèsew-sipi	-	Red-deer River.		
Po-i-thinnè-kaw-sipi -	-	Nelson or North River.		
Pemmichi-ke-mè-u -	-	Cross Lakc (Nelson River).		
Maskègowuk	-	Swampy or low grounds near		
M		Hudson's Day.		
Ive me-u sipi	-	Sturgeon River.		
Chuki-tanu sipi	-	Hill River.		
Penesay-: ichewan sipi	-	Hay's River.		
Washè-u-sipi	-	Severn River.		
Wewanito-wuk	-			
Kà-stitchewanuk -	-	Albany River. Here Hudson		

had his first interview with the natives, among whom traditions of the circumstances attending it were current in 1770.

D 3

The Iroquois name the Algonkin race Adirondak, and Mr. Schoolcraft thinks that this appellation,

NAKA-WE-WUK, OR NORTHERN UTAWAWA.

This people inhabit the country lying between Christianux Lake (Lake Winipeg) and James's Bay, approaching within one hundred miles of the latter. They speak the Odchipewa tongue.

Names of Tribes or Places.		Districts they inhabit.
Namèkusi-sipi	•	Trout River.
Wà-pusi-sipi	-	Hare River.
Christianux	-	Lake Winipeg.
Weniska-sipi	-	Badger River.
Odchipewe-sipi	-	River Winipeg.
Mistĕhè-saka-hegen -	-	Great Lake Winipeg.
Mith-kwa-ga mè-u-sipi	-	Red or Bloody River.
Shama-tawa		Henly House River.

UPE-SHI-POW.

This people resort to the eastern coast of Hudson's Bay, between Rupert's and Whale Rivers. Their language differs in some words both from the Keiskatchewan and Nakawawa. (They border on the Eskimos of the Labrador peninsula.)

Muswà-sipi -	-	-	Moose River
Winne-peskowuk	-	-	East Main.

La Hontan enumerates the tribes speaking the Algonkin language in 1700 as follows: ---

In Acadia (Nova Scotia) :- Abenakis*; Mickemac; Canibas;

be of Ma of sev the pin gon 0 kan war C Ilin Kik min С touc gua С kan Ten cow 0 on I bato Т Esk to of posi who

> and sound *

^{*} This tribe are of the Iroquois stock, according to Schoolcraft, who says that *Abenaki* is a derivative from *Wabanung*, "the east," and *akkè*, "earth," and signifies "eastlanders." The *Abenakis* were called Tarrenteens by the early English colonists, and formerly inhabited part of the present states of Maine and New Hampshire. They were divided into several sub-tribes, of whom the best known are the *Penobscots*, *Norridgewocks*, and *Ameriscoggins*. About the year 1754, all but the Penobscots withdrew into Canada. The fullest vocabulary of the Abenaki langnage is furnished by the manuscripts of Father Rale, and has been published by the American Academy of Arts

TRIBES.

39

ondak, lation,

istianux thin one tongue. bit.

n's Bay, e differs kawawa. a.)

Algonkin

Canibas;

who says , "earth," ens by the it states of o-tribes, of *riscoggins*. ida. The anuscripts y of Arts being still retained for the highlands at the source of the Hudson, countenances the traditions current

Muhingaus; Openangos; Soccokis (Sokokies, living eastward of Boston, New England—Colden); Etechemins.* These seven tribes are brave warriors, more expert and less cruel than the Iroquois. Their language differs little from the Algonkin.

On the St. Lawrence, from the Sea up to Montreal:—Papinachois; Montagnois; Gaspesiens; Abenakis of Sciller; Algonkins.

On Lake Huron :— Outaouas; Nockes; Missisagues; Attekamek; Outchipoues (Odchipewa), called Sauteurs, b ave warriors.

On the borders of Lake Ilinois (Lake Michigan):-Some Ilinois of Chegakou; Oumamis, brave warriors; Maskoutens; Kikapous, brave warriors; Outagamis, brave warriors; Malomimis; Pouteouatamis; Ojatinons, brave warriors; Sakis.

On the borders of Lake Frontenac (Lake Ontario): -- Tsonontouans; Goyoguans; Onontagues; all of whom speak a language differing from the Algonkin; Onnoyoutes and Agnies.

On the Outaouas (Uttawa):— Tabitibi; Monzoni; Machakandibi; Nopemin of Achirini; Nepisirini; Temiskamnik (Lake Temiscamaing). These six tribes speak Algonkin, and are all cowards.

On the north of the Mississippi, and in the country bordering on Lake Superior and Hudson's Bay: -- Sonkaskitons; Ouadbatons; Atintons; Clistinos, brave and skilful warriors.

The Assimpouals (Assinipoytuk, or Stone Indians) and Eskimaux are struck out from La Hontan's list, as they belong to other nations. The chart appended to his book gives the positions he assigns to these several tribes; but Schoolcraft whose authority is of the greatest weight, says that the list

and Sciences. The language is peculiar, from the frequent use of the rolling sound of r, or a burr.

^{*} The Etchemins are of the Iroquois race, according to Schoolcraft.

among the western Algonkins, that their ancestors came from the eastern coast. It is probable that the Mengwè, on the contrary, advanced from the west, if we may judge by the way in which their tribes are distributed. The aggressive movements, however, of the two nations, would throw little light on the primary peopling of the continent, even were they ascertained, since there are traces of their respective districts having been previously occupied by a people of small stature but superior in the arts, who have left memorials of their existence, in numerous and extensive earth-works, and mounds of ancient date, wherein copper bosses overlaid with silver have been found. The shafts or galleries lately discovered at the copper mines on the south side of Lake Superior, containing immense quantities of stone chisels, betoken a people more advanced than the Canadians were on the first arrival of the French; and are said to be now followed by the American miners, as guides to the most valuable deposits of native copper.

The wars of the *Eythinyuwuk* and the *Mengwè* with each other, or with Europeans, have been recorded by many pens, and have supplied incidents

contains many errors. The chart, of course, has all the imperfections which attached to the geography of the great lakes and more northern country then, and for more than a century afterwards. for ha qu val wit rea by cor sav bee Th pri var tur me taiı and rica exa ten ria We deı be a res the

WARS WITH THE MENGWE.

cestors

le that

m the their

ments,

little

traces iously

perior

exists, and

bosses shafts

mines

aining

ken a

ere on

to be ruides

engwè been

dents

imper-

es and

after-

for numerous works of fiction, in which the writers have ascribed a loftiness of soul and other noble qualities to these people, of which it would be in vain to seek traces in the present day; and we may without much scepticism assert, that they never really possessed them. Actions prompted only by the caprice of a barbarous people, have been considered as the results of refined sentiment : and savage cunning, seen through a false medium, has been elevated to the promptings of far-seeing policy. The revolting cruelty with which they tortured prisoners of war, and the stoicism with which, when vanquished, they endured such treatment in their turn, are more certain traits of character. A few men, remarkable for their powers of mind, have certainly appeared among the Eythinyuwuk nations, and from them the abstract idea of a North American Indian has been formed by Europeans.

Among this people there are to be found finer examples of the human figure, handsomer countenances, and a more manly and independent carriage, than among the Eskimos and 'Tinnè; and West's exclamation on seeing the Apollo Belvidere, that he was a young Mohawk warrior, may be adduced as evidence of the natural grace which a ranger of the woods, unfettered by artificial restraints, may possess. In fact, the attitudes of the Eythinyuwuk are occasionally, and especially

when actuated by strong passion, striking, and sometimes elegant; yet the habitual gait of the Red man is not a graceful one. The toes are turned in; the step, though elastic, has an appearance of insecurity, and is by no means majestic, nor even pleasing, to one unaccustomed to see the centre of gravity thrown so much forwards. Even though the palm of personal appearance be given to the Eythinyuwuk, in moral conduct I hold them to be decidedly inferior to the Eskimos and 'Tinnè. They are less honest, and though perhaps not so much given to falsehood as ties 'Tinnè, are more turbulent and more prompt to invade the rights of their countrymen, as well as of neighbouring na-Their wars are carried on by ambuscade tions. and treachery, seldom in open field: they spare neither infants nor women in their forays; and instances of personal bravery, such as the Eskimos often exhibit, are rare indeed among them. The worst of the vices of which St. Paul accuses the heathen world are said to exist among the Crees of the plains, and gambling is practised to excess by the whole nation.* One game in which the

* A society named *Wabuno* is said to have been formed arong the Chippeways, for the practice of certain nocturnal orgies called *Wabunsi*, — an appellation signifying "not yet light," from *wauben*, "daylight," and the negative suffix.—See Schoolcraft. The appellation of *Wahunsenacawh*, by which





WIGWAMS.

odd or even number of pebbles, and the hand in which they are held, are to be guessed, is constantly resorted to whenever two or three meet together, and it is accompanied by singing and gestures, indicating some kind of divination. They will pass a whole day so occupied, and will stake all that they hold most valuable on the result.

As the Narrative of Sir John Franklin's First overland Expedition in 1819-21 contains all the particulars of the manners and religious belief of the Crees, or *Nethewuk*, that I had then collected, I shall not here repeat them; but shall merely allude in a brief way to such of their habits and usages as have not been noticed among the 'Tinnè.

The ordinary wigwams*, skin tents, or "lodges" as they are called, of the two people are exactly alike as to form, being extended on poles set up in a conical manner; but, as a general rule, the tents of the Crees are more commodious, and more carefully

Powhattan's subjects were accustomed to address him, had probably reference to his being chief of a society of this kind. (See Virginia by Strachey.)

* Wiggè or Wigwap signifies a dwelling. Most of the Indian words and names of places adopted into the languages of the United States and Canada are of Algic origin. The Mandans and other Dakotas of the Missouri build more substantial huts, with dome-shaped roofs, covered with earth, to which, as to look-out places, the men resort.

C

۲C

RTLI

& N HANAHT

1 p +3 /

44

and frequently supplied with a fresh lining of the spray of the balsam fir. This people also occasionally erect a larger dwelling of lattice work, covered with birch bark, in which forty men or more can assemble for feasting, debating, or performing some of their religious ceremonies. These erections are chiefly made on the skirts of the bison plains, or in localize s where a large number of the nation are accustomed to assemble together. The entire nation of the Eythinyuwuk cultivate oratory more than their northern neighbours, who express themselves much more simply and at the same time with much less readiness.

Neither among the Eskimos nor 'Tinnè did I observe any image or visible object of worship; but most of the Crees carry with them one or more small wooden figures rudely carved, some of which they state to be representatives of a malicious or at least capricious being named Kepuchikan*, to whom they make offerings. They acknowledge other spirits or Manito-wuk, and demons or vampires called Witako; but I could not ascertain that prayer was ever made to the Kitche-manito, the "Great Spirit" or "Master of Life." The vapour-bath,

* Or Gepuchikan. The propensities as well as designation of this being resemble those of "Puck." Dr. Johnson derives the latter word and Pug, from the Icelandic and Gothic word Puke, signifying "a hobgoblin."

wh is i or the jur shu hou a ki out, hot mov 0 dest Gre trib eter craf duti for such whe of. " the Sup the

corre

EVERLASTING FIRE.

g of the occasioncovered nore can ing some tions are ns, or in ation are ntire nary more express he same

C.

did I obnip; but or more of which ous or at to whom ge other vampires at prayer "Great ur-bath,

esignation on derives othic word which is comparatively seldom used by the 'Tinnè, is in frequent request with the Crees, and is more or less connected with religious observances. It is the great medium by which the shamans or conjurors cure the sick. The operator in this case shuts himself up with his patient in the sweatinghouse, where he shampoes him, singing all the time a kind of hymn. As long as the shaman can hold out, so long must the sick man endure the intensely hot atmosphere, and then, if the invalid be able to move, they both plunge into the river.

One custom of the Chippeways which fell into desuetude after the arrival of the French on the Great Lakes, is still preserved by tradition in the tribe. It was an institution for preserving an eternal fire named Kagagish'koda. Mr. Schoolcraft gives the following account of the rites and duties connected with it, which I make no apology for quoting, as so singular a custom, related on such good authority, deserves to be mentioned, when the peculiar habits of the race are spoken of.

"The Chippewa tribe had its council-house, and the seat of eternal fire, on the south side of Lake Superior, west of Keeweenau Point. Here lived the principal chief, called the *Mutchèkewis*, who ex-

* From Ka-gi-gi, "everlasting," and iskōda, "fire." The corresponding words in Cree are $k\bar{a}$ -ki-ki and iscu-teyu.

ercised more authority, and assumed more state, than would be compatible with the present feelings of the Indians. The designation was official, and not personal, and the office was hereditary in the direct male line. He was supported by voluntary contributions, his *mushinawa*, or provider, making known from time to time his necessities by public proclamation. Whatever was required on these occasions, whether food or clothing, was immediately furnished. He appears to have been the chief priest, and could neither engage in war nor hunting."

"In the village where he resided, and near his cabin, the eternal fire was kept burning. The altar was a rude kind of oven, over which no building was erected. Four guardians were selected by the Mutchèkewis, to take care of the fire. Two of these were men, and two women. They were all married; but the wives of the men employed on this service, were required to cook and do the necessary domestic work, while the husbands of the women destined to the sacred duty were always engaged in hunting, and in providing whatever else was wanted. The four persons devoted to the altar were thus left without any secular cares to divert their attention from the holy trust committed to them. A perpetual succession was kept up in the priesthood, by a prerogative of the

Mu forn for sons and Dea it w 66 and the t caref Whe enou kewi was l which then addeo "(assem seaso morn fire, a one si these mann This morni

FIRE USED IN SICKNESS.

47

e state, feelings al, and in the luntary making y public n these s immeeen the war nor

id near The g. hich no s were e of the women. the men ook and usbands ty were g whatroted to ar cares st comas kept of the Mutchèkewis, and the principal head woman; the former selecting a husband, and the latter a wife for the survivor, whenever one of these eight persons died. The chain was thus always unbroken, and the traditionary rites transmitted unimpaired. Death was the penalty for any neglect of duty, and it was inflicted without delay and without mercy."

"The council fires were lighted at the great fire, and carried wherever the council was held. After the transaction of the business, a portion of i⁺ was carefully returned and the remainder extinguished. Whenever a person became dangerously ill, if near enough, he was taken to the house of the Mutchèkewis, where his fire was extinguished, and a brand was brought from the altar and a fire kindled, at which a feast was prepared. A great dance was then held, and the viands consumed. And it is added that the patient seldom failed to recover.

"Once in eight years, the whole Chippewa tribe assembled at their principal village, about the season of the swelling of the buds. Farly in the morning the great pipe was lighted at the sacred fire, and delivered to the Mutchèkewis. He took one smoke, and then handed it to the women, and these to the men, by all of whom it was in like manner smoked. It was then passed to the children. This ceremony consumed the day, and early next morning a feast was held, at which the men and

women and children sat in separate groups. This feast was partaken silently, and without singing or dancing. In the evening they departed to their different villages."

"The principal male attendant on the Kaugagiskoda was the Kauga-gizhek, or "Everlasting Sun;" and his assistant was named Kanawaudenk-shkuda, or the "Fire Keeper." The principal female was called Gaubewekwa, or the "Everlasting standing woman;" and her assistant Kabagaubewekwa, "The woman who stands all the time."

"The Chippeways assert that they received this custom from the Shawnees, who are the most southern of the western Algonkins, their country being in the present state of Kentucky. Traces of its prevalence at a former period among other North American nations exist. The Natchez and most of the Louisiana tribes are represented by Charlevoix as having had a perpetual fire in their temples. Both he and Du Pratz were eye-witnesses of the rite. The hereditary ruler, or "Chief Sun," whose title was equivalent to that of Inca or Emperor, exercised a more despotic power than appears to have been permitted in any other nation north of Mexico. This power and this worship were kept up with an oriental display of honor and ceremony long after the French had settled in the valley of the Mississippi, and indeed up to the destruction

of eate befo 'the \mathbf{F} that the than than knov was coun beyo M carry her h lation long it. year, again longe \mathbf{Th} so im Eythi 'Tinne ropear have VOL

THE CALUMET.

. This ging or to their

ugagis-Sun;" shkuda, ale was tanding a, "The

zed this e most country races of g other hez and nted by in their itnesses ef Sun," or Em. han apnation hip were nd ceree valley ruction of the nation by them in 1729. 'The Sun has eaten,' proclaimed an official functionary daily, before the Ruling Sun, after his morning's repast; 'the rest of the princes of the earth may now eat.'"

From this interesting extract we may gather, that the Algic race were much more advanced in the forms of government and association of tribes than the more northern nations, and especially than the 'Tinnè, who had no villages when first known to Europeans. Cultivation of the earth was not carried on to the north of the Chippeway country, since maize does not prosper in America beyond the 52nd parallel.

M'Kenney relates that a Chippeway widow must carry a bundle of rags, or a doll, which is called her husband, constantly in her arms, until the relations of the deceased think that she has mourned long enough, when one of them releases her from it. This occurs generally at the expiration of a year, and the widow is then allowed to marry again; but the probation may be extended much longer, if her husband's relations choose.

The use of the Uspogan, or Calumet, which forms so important a part of every ceremony among the Eythinyuwuk, was not an original practice of the 'Tinnè, but was introduced to that people by Europeans along with tobacco, whereas this weed must have been grown from the most ancient times by

Е

VOL. II.

the Chippeways, if the traditions which Mr. Schoolcraft collected during his long residence with that people are to be trusted. Maize is more used on the Missouri than in the proper Chippeway country, its cultivation forming a part of the regular economy of the Dakota tribes; the Chippeways, however, do not admit that they received it from that quarter; but, in a legend related by Mr. Schoolcraft, ascribe its origin to one of their own chiefs, who received it as the prize of a victory he obtained over a spirit. Hence its name of Mondamin, or the Spirit's Grain. The Delawares had extensive fields of maize at the time of the discovery of America, and to them the early Virginian colonists were indebted on their first landing for food, which being afterwards withheld, produced extreme misery and famine.

From some of the details of Mr. Schoolcraft's account of the rites of Kagagishkoda, we may infer that the national polity and social condition of the Chippeways have greatly deteriorated since their acquaintance with Europeans. The contact with civilised man has induced among them an incontrollable desire for intoxication, unaccompanied by any real benefit. For though missionaries have made a number of nominal converts, the blessings of vital Christianity are confined, as far as I could ascertain, to only a few Chippeway communities

on Hu Sar pen and on T cour rein dant appi Eski prain popu and of bi deer speci

* Ac ric word (School

the

figur

Adika

chey'

nor t

of the

being

REINDEER.

choolh that sed on untry, egular eways, t from Schoolchiefs, he ob-Monres had e discorginian ing for oduced

blcraft's ny infer of the te their ct with inconnied by s have lessings I could hunities on Lake Huron, and to some of the Crees in the Hudson's Bay Company's territory. The well-fed Sauteurs of the river Winipeg, who are independent of the traders, repel the missionaries; and the same is the case with the bison-hunters on the prairies.

Throughout the whole eastern wooded and barren country, down to the 42nd parallel of latitude, the reindeer was, three centuries ago, the most abundant of the deer kind, and, being the most easily approached, furnished the staple provision for the Eskimo, 'Tinnè, and Eythinyuwuk. On the wide prairies of the Missouri and Saskatchewan, the populous Sioux, Stone-Indians, or Assini-poytŭk, and other Dakota tribes, fed on the countless herds of bison which pasture there. Next to the reindeer in importance in the eastern districts, is the species of Coregonus, named "white-fish," to which the Chippeways and Nithè-wuk have given the figurative appellation of " reindeer of the waters," Adikumaig or Atih-hameg.* On referring to Strachey's account of Virginia, I do not find this word, nor the name of the reindeer, in his vocabulary of of the Delaware tongue; the white-fish indeed not being an inhabitant of the southern waters. The

* Adihumaig, from adih, a "reindeer," and guma, a generic word for "water" in composition, and the animate plural ig. (Schoolcraft.) Athik or atik, "a reindeer," in Cree.

Chippeways have a legend, which relates that the white-fish sprung first into existence at the outlet of Lake Superior, being produced from the scattered brains of a woman, whose head, for some very guilty conduct, was doomed to wander through the country, but, coming in its travels to the falls of St. Mary, was there dashed in pieces. A crane, by virtue of that inherent power so frequently attributed to birds and beasts by the aborigines of America, instantly transformed the particles of brain into the roe of a white-fish, to the widespread benefit of the Indian nations.*

Though the earth-works already alluded to are supposed to have been raised by a people more ancient than the Eythinyuwuk, yet the fact of their northern limits being within the Chippeway lands is worthy of note; and vestiges of pottery works, apparently of a rude kind, have been found on the south branch of the Saskatchewan within the Nithè-wuk bounds, but not further north †, the substitute for earthenware among the Eskimos being vessels of potstone, and among the 'Tinnè water-tight baskets, in which the fluid was warmed by hot stones dropped into it.

* Schoolcraft.

The Eskimos † On the east side of the Rocky Mountains. on the western coast of Russian America manufactured a very rude pottery when first visited by the Russians.

mo tior pro par ract of t l an plai the East is d omis befo: tatio affec lang v, no peop usua ferre throa little, quent The . vocab * M

has bee

LANGUAGE.

at the attered e very ogh the falls of ane, by y attriines of cles of e wide-

to are e more fact of ppeway pottery n found within north †, Eskimos 'Tinnè varmed

> Eskimos d a very

I have already alluded to the softness and harmony of the Cree language. It differs in construction from the Eskimo tongue, in the personal pronot as being prefixes, not suffixes, and in other particulars; but both have the polysynthetic character of the other American idioms. The sounds of the English f and v do not occur in the Cree; l and r are also wanting in the pure Cree of the plains. Other Algic tribes substitute y, n, or l for the Cree th, and instead of k, the inhabitants of East Maine use the sound of tch. The Chippeway is distinguished from the Cree by the frequent omission of s before k and t, and the insertion of mbefore b, and of n before d and g. The permutations of the Cree and its cognate dialects chiefly affect the linguals; but the Mohawk and Huron languages have none of the labials, neither b, p, f,v, nor m. When conversing, the teeth of these people are always visible; the auxiliary office usually performed by the lips being by them transferred, or superadded, to that of the tongue and throat.* Of the grammar of the 'Tinnè I know little, but the nouns seem to be much more frequently monosyllabic than in the Algonkin dialects. The Appendix contains some portions of a Cree vocabulary, which I formed in 1819-20.

* Mr. Howse, from whose grammar much of this paragraph has been borrowed.

е З

It is from among the Eythinyuwuk that most of the servants of the Fur Companies, who have married native women, have selected their wives; few of them having chosen Chepewyan females, and no one, I believe, an Eskimo maiden. From these marriages a large half-breed population has arisen, which will ere long work a change in the fur trade, and in the condition of the whole native popula-In character, the half-breeds vary according tion. to their paternity; the descendants of the Orkney labourers, in the employ of the Hudson's Bay Company, being generally steady, provident agriculturists of the Protestant faith; while the children of the Roman Catholic Canadian voyagers have much of the levity and thoughtlessness of their fathers, combined with that inability to resist temptation, which is common to the two races from whence they are sprung. Most of the half-breeds have been settled by the Hudson's Bay Company in the colony of Osnaboya, which extends for fifty miles along the banks of the Red River of Lake Winipeg. Of the six thousand souls, to which the mixed population of this settlement is said to amount, three fifths are stated by Mr. Simpson to be Roman Catholics; while the valuable property is mostly in the hands of the remaining two fifths, who own sixteen out of eighteen wind and water mills, erected within the precincts of the colony.

sca and Coi jud just Cor I over som the bish Hud the adva ploy Rom tions and both are : color being the y son's have factor as wi

 $\mathbf{54}$

EDUCATION.

t most o have wives; es, and n these arisen, r. trade, populacording Orkney 's Bay nt agrihildren s have of their resist es from -breeds mpany or fifty f Lake ich the said to bson to roperty fifths, water ony.

The settlement is under the government (it can scarcely be said the control) of a governor, council, and recorder, all nominated by the Hudson's Bay Company. The recorder is the civil and criminal judge, presides at jury trials, and is aided by justices of the peace, and a constabulary in the Company's pay.

In 1849 a bishop was sent from England to oversee the Episcopal church. There are also some ministers of the Wesleyan persuasion; and the Roman Catholic worship is maintained by two bishops, a staff of priests, and a nunnery. The Hudson's Bay Company aid the clergymen of all the persuasions by free passages, rations, and other advantages, besides granting salaries to those employed at their fur posts, whether Protestants or Roman Catholics. There are also various educational establishments in the colony for the settlers and native population; and most of the children, both male and female, of the Company's officers are now instructed in a boarding-school in the colony of a high character, a few of them only being sent to Great Britain or Canada. Many of the young men so educated have entered the Hudson's Bay Company's service as clerks, and some have attained the rank of chief traders and chief factors; while the young women, in their vocations as wives of the officers and clerks, diffuse a know-

Е 4

56

ledge of Christianity and a taste for domestic comfort and decorum to the remotest posts. The present state of society in the fur countries contrasts most favourably with the almost general heathenism which prevailed during the murderous contests between the trading companies by which the country was demoralised when I first traversed it thirty years ago.

The half-breeds, as a class, show great quickness in acquiring a knowledge of letters, as well as skill in the mechanical arts. As joiners, workers in iron, and boat-builders, many of them would rank high among European craftsmen; and, taught by necessity, they have generally the advantage of being able to work at all the several branches of the carpenter's and blacksmith's arts, even to the forging of their tools.

At the Wesleyan Missionary establishment of Rossville, near Norway House, and round the Episcopal church at the Pas on the Saskatchewan, native villages have sprung up, and agriculture to a small extent is practised. Though the cerealia and leguminous vegetables thrive well at Red River, and horses, cattle, hogs, poultry, and sheep flourish, agriculture is eschewed by the large section of the population, who are descendants of the Canadian voyagers. The pleasures of the precarious chase are preferred by this part of the community to steady

indu exte for v Α ingi the s quai can tribe inter are f yet, often to wh posse tors o haug at on are s virtu Chip their Mons speak gener Go both keepe

SPIRITUOUS LIQUORS.

c com-The cs congeneral derous which aversed

ickness as skill in iron, ik high necesbeing of the to the

ent of e Episnative a small a and River, ourish, of the nadian ase are steady industry, and every summer there is accordingly an extensive movement to the plains to dry bison meat for winter use.

As to the effect of the colony on the neighbouring natives, Mr. Simpson, who from his residence in the settlement had an opportunity of becoming acquainted with the facts, speaks as follows. "Nothing can overcome the insatiable desire of the Indian tribes for intoxicating liquors; and though they are interdicted from the use of spirits, and the settlers are fined when detected in supplying them with ale, yet, from the great extent of the colony, they too often contrive to gratify that debasing inclination, to which they are ready to sacrifice everything they They feel no gratitude to their benefacpossess. tors or spiritual teachers; and while they lose the haughty independence of savage life, they acquire at once all the bad qualities of the white man, but are slow indeed in imitating his industry and virtues." It appears from this testimony that the Chippeways have not the friendly feelings towards their instructors which the 'Tinnè, according to Monsieur La Flèche, manifest; but Mr. Simpson speaks more favourably of the Crees, who are in general better disposed than the Chippeways.

Goods for the use of the colonists are imported both by the Company and by individual storekeepers in the ships that come annually to York
CREES.

58

Factory; but the distance is too great, and the inland navigation too difficult, to admit of agricultural produce being carried down profitably in return. Hence most of the half-breed settlers, encouraged by some of the colonial merchants and Roman Catholic priests, have made strenuous attempts to share the fur trade with the Hudson's Bay Company, who at present have the monopoly of that traffic; and the Company do not seem to possess a force adequate to prevent their eventually succeeding in their object.

Of late years, a communication has been formed between the colony and the United States by way of the plains and St. Peter's River. This furnishes a channel for the disposal of peltry without detection; and through the relationship existing between the half-breeds of the colony and the various tribes of Indians as far north as Methy Portage, no great difficulty is experienced by them in withdrawing a considerable quantity of the most valuable furs from the Company's trade.

In the winter of 1848 a half-breed was summoned before the Recorder of Osnaboya for a breach of the Company's regulations in this respect, and on the day of trial, five hundred of his class, armed to the teeth, surrounded the court-house. The Recorder was obliged to secrete himself, and the matter was finally compromised by the Company's

HUDSON'S BAY COMPANY.

and the agricultably in lers, ennts and uous atludson's onopoly seem to r even-

formed by way urnishes t detecbetween s tribes to great wing a ble furs

nmoned each of and on med to he Rcnd the upany's

agent purchasing the furs from the delinquent. Secretly or openly, this contravention of the right of exclusive trade in fur claimed by the Company is sure to proceed, and, emboldened by success, the young half-breeds are not likely to acknowledge any law that is contrary to their own will. They hold that the territorial right derived from their Indian ancestry is theirs, and not the Company's; and their claims have been supported by a philanthropic body in England, and advocated in parlia-Without entering into the question of the ment. chartered rights of the Hudson's Bay Company, or the propriety of maintaining a monopoly of the fur trade, it is my firm conviction, founded on the wide-spread disorder I witnessed in times of competition, that the admission of rival companies or independent traders into these northern districts would accelerate the downfall of the native races. This has been rapid on the confines of the settled parts of the United States and of Canada, and has been stayed only by the extinction of the fur-bearing animals, by which the power of the Indians to purchase spirits has been cramped. Even the benevolence of the English government in making annual presents of clothing and blankets to the Indians of Canada is converted into an injury by a set of unscrupulous petty dealers, who hang about the encampments to purchase these articles as soon

CREES.

as they come into the possession of the Indians, by supplying them with the baneful liquid they so This is punishable by the colonial ardently covet. laws; but when crimes are committed beyond the pale of civilisation, conviction is difficult. By the laws of the United States, also, it is penal to supply Indians with spirits; but according to general report this benevolent enactment is extensively violated by their fur traders; and it is greatly to be regretted that competition for the Indian trade in that quarter should induce the Hudson's Bay Company to follow so bad an example, after having abolished the use of spirits with so much advantage in the north, where they have no rivals.

I was informed that in 1848 the natives at the Red River colony of Osnaboya were paid a high money price for their furs by the Company's agent, and that they immediately crossed the boundaryline to purchase rum at the American post with their money; but it would be better to seek for the redress of such an abuse by a representation to the United States' government, than resort to retaliatory measures of the same nature.

FORT O BASI TER TEN FISH PRO WIN — W LENO MOVI A W DESE RATU DENO

THE about on a which the la feet. from yards towar above

* Tł 1848 by

FISHERY ISLAND.

61

CHAP. XV.

OCCURRENCES IN WINTER.

FORT CONFIDENCE. — SITUATION. — SILURIAN LIMESTONE. — LAKE BASIN. — TREES. — DWELLING-HOUSE. — OCCUPATIONS. — LET-TERS. — GALENA NEWSPAPER. — OREGON "SPECTATOR." — EX-TENT OF THE HUDSO.'S BAY COMPANY'S TERRITORY. — FISHERIES. — VENISON. — WOLVERINES. — NATIVE SOCIALISM. — PROVISIONS COLLECTED AT FORT CONFIDENCE. — FETES. — WINTER FISHERY. — ESKIMO SLEDS. — REINDEER. — WOLVERINE. — WOLVES. — HONESTY OF THE DOG-RIBS. — THEIR INDO-LENCE. — PROVISIONS NOT INDIVIDUAL PROPERTY. — INDIANS MOVE OFF. — AN ACCOUCHEMENT. — CŒLEBS IN SEARCH OF A WIFE. — MIGHT MAKES RIGHT. — NONE BUT THE BRAVE DESERVE THE FAIR. — PROGRESS OF THE SEASONS. — TEMPE-RATURE. — ARRIVAL OF SUMMER BIRDS. — AT FORT CONFI-DENCE. — AT FORT FRANKLIN. — ON THE YUKON.

THE site selected for our winter residence was about three miles from the mouth of Dease River, on a peninsula having an undulating surface, which, at the distance of three or four miles from the lake, attained a height of about three hundred feet. In front, or to the south, and separated from the main by a strait five or six hundred yards in width, lies Fishery Island, elevated towards its centre two hundred and forty-five feet above the water.*

* This altitude was ascertained by Mr. Rae in the spring of 1848 by the aneroid barometer.

ians, by they so colonial ond the By the o supply general ensively eatly to an trade n's Bay c having vantage

es at the a high s agent, undaryst with t for the h to the to reta-

SILURIAN LIMESTONE.

The peninsula is composed of limestone, which forms low precipices at the edge of the water, as well as in various places of the interior; and the same rock appears in higher cliffs on the borders of the lake, about eight miles to the westward, at Limestone Point. Six or seven miles back, on the banks of Dease River, red sandstone is the prevailing rock. The soil generally is a mixture of gravel and loam; and boulders of granite and trap rocks are scattered over the surface of both hill and valley.

Ten miles to the eastward, a range of primitive rocks rises gradually from the borders of the lake, to the height of, perhaps, six hundred or seven hundred feet, and separates Dease's Bay from the northern arm of M'Tavish's Bay. This rising ground is a continuation of the "intermediate primitive belt" mentioned in p. 316, and many other parts of the preceding journal, and which will be described more fully in the Appendix. The nearest pyrogenous or metamorphic rocks to Fort Confidence that we observed are about four miles off, in a bay on the south-east side of Fishery Island.

The limestone is probably the remains of the silurian strata, which were removed when the basin of the lake was excavated. On the south side of the lake, about ninety miles distant in a

direc Grass consi extre const Athal been shale The naked a thin wet p Little is very sian p few yo been o three from t trees circum ground and fif The

place 1 and 11 pretty mouth

63

, which ater, as and the borders rard. at , on the he preture of nd trap oth hill

imitive ne lake, om the rising nediate many which pendix. ocks to it four 'ishery

of the en the south t in a direct line from Fort Confidence, stands the Scented Grass Hill, between Smith's and Keith's Bays. It consists of bituminous shale, and is one of the extreme points of that shaly formation, which constitutes so large a part of the banks of the Athabasca and Mackenzie Rivers, and which has been thought to be equivalent to the Marcellus shale of the New York system of rocks.

The summits of the higher eminences are mostly naked, but on the edges of streams and small lakes a thin forest of spruce fir covers the ground. In wet places there is a tolerable growth of willows. Little underwood of any other kind exists. Birch is very scarce; neither the balsam spruce nor banksian pine were observed on the lake, and only a few young aspens. Except where the forest has been destroyed by fire, the spruce firs are from three to four hundred years old, as ascertained from their annual rings. One of the best-grown trees that I saw, measured fifty-seven inches in circumference, at the height of four feet from the ground. The tallest of them are between forty and fifty feet high.

The observations of Mr. Simpson in 1837-8 place Fort Confidence in 66° 54' of north latitude, and 118° 49' of west longitude, which corresponds pretty closely with the position I assigned to the mouth of Dease River on the chart constructed in

DWELLING-HOUSE.

1825. The mean of Mr. Rae's observations for latitude gave about a quarter of a mile more northing than Mr. Simpson's.

Our winter dwelling, though dignified, according to custom, by the title of "the fort," had no defensive works whatever, not even the stockade which usually surrounds a trading post. It was a simple log-house, built of trunks of trees laid over one another, and morticed into the upright posts of the corners, doorways, and windows. The roof had considerable slope: it was formed of slender trees laid closely side by side, resting at the top on a ridge pole, and covered with loam to the depth of six or eight inches. A man, standing on the outside, could touch the eaves with his hand. Well-tempered loam or clay was beat into the spaces left in the walls by the roundness of the logs, both on the outside and inside, and as this cracked in drying, it was repeatedly coated over, for the space of two months, with a thin mixture of clay and water, until the walls became nearly impervious to the air. The rooms were floored and ceiled with deal. Massive structures of boulder stones and loam formed the chimney-stacks, and the capacious fire-places required three or four armfuls of fire-wood, cut into billets three feet long, to fill them.

The building was forty feet long by fourteen

wide sixte vided ment house room purp tory. skin Two storeh sides tallest withir being for a was b Of erected Augus stack throug stroye most o what y in fron were c others VOL.

DWELLING-HOUSE.

65

ons for north-

cording no detockade It was ees laid upright s. The med of sting at h loam A man, e eaves lay was by the de and was remonths, htil the The •. Massive hed the ces reut into

burteen

wide, having a dining-hall in the centre, measuring sixteen by fourteen, and the remaining space divided into a store-room and three sleeping apartments. A kitchen was added to the back of the house, and a small porch to the front. Mr. Rae's room and mine had glazed windows, glass for the purpose having been brought up from York Fac-The other windows were closed with deertory. skin parchment, which admitted a subdeed light. Two houses for the men stood on the east, and a storehouse on the west, the whole forming three sides of a square, which opened to the south. The tallest and straightest tree that could be discovered within a circuit of three miles was brought in, and, being properly dressed, was planted in the square for a flag-post; and near it a small observatory was built, for holding magnetic instruments.

Of the buildings which Dease and Simpson erected, Mr. Bell, on his arrival in the middle of August, found only part of the men's house and a stack of chimneys standing; the others having, through the carelessness of the Indians, been destroyed by fire. Our predecessors had cut down most of the timber within a mile of the house, and what we needed had consequently to be brought in from a wider circle. A part of Mr. Bell's people were constantly engaged with the fisheries, but the others had worked so diligently, that the buildings VOL. II.

were all covered in on our arrival, and the flooring, ceiling, and partitions were shortly afterwards completed. Two of the sappers and miners, Mackay and Brodie, carpenters by trade, were employed to make tables and chairs; and Bruce, the guide, acted as general architect, and was able and willing to execute any kind of joiner's work that was needed. Two men were constantly employed as sawyers; four as cutters of fire-wood, each of them having an allotted task of providing a cord of wood daily; others were occupied in drawing it home on sledges; and four men were continually engaged in fishing. On the Sunday no labour was performed, the fishing party came in, and all were dressed in their best clothes. Prayers were said in the hall, and a sermon read to all that understood English; and some of the Canadians, though they were Roman Catholics, usually attended. James and Thomas Hope, who were Cree Indians, having been educated at Norway House as Protestants, and taught to read and write, were regular attendants; and James Hope's eldest son, a boy about seven years of age, who had already begun to read the Scriptures, frequently recognised passages in the lessons that he had previously read.

During the winter Mr. Rae and I recorded the temperatures hourly, sixteen or seventeen times a day; also the height of the mercury in Delcro's barome declin mont was l were vario certai neton dippin the t servat winds was c Fro

seen t no lei we en ably e ceeded little and re produce The letters

* The at Wool intendan of the m

OCCUPATIONS.

ooring, ds comcay and o make cted as to exed. Two s; four ving an daily; me on aged in formed, ssed in he hall, nglish; y were nes and ng been ts, and ndants; t seven ead the in the

led the times a ro's barometer; the degrees of the aneroid barometer, the declinometer, and dipping-needle. Once in the month a term day, extending to thirty-six hours, was kept, in which the fluctuations of the magnets were noted every two and a half minutes, and various series of observations were made for ascertaining the magnetic intensity with the magnetometer, the vibration apparatus, and Lloyd's dipping-needle. Mr. Rae ascertained frequently the time and rates of the chronometers by observations of the fixed stars; and a register of the winds and weather and appearances of the aurora was constantly kept.*

From this sketch of our occupations, it will be seen that our time was filled up, and that we had no leisure for ennui in the long winter. In fact, we enjoyed as much comfort as we could reasonably expect, and had our postal arrangements succeeded as well as the others, we should have had little more to desire. Our schemes for sending and receiving letters were, however, failures, and productive of much subsequent disappointment.

The packet of Admiralty despatches and private letters sent off on the 18th of September, 1848, on

* The magnetic observations are now in process of reduction at Woolwich, and will soon be published under the superintendance of Lieutenant Colonel Sabine, along with an abstract of the meteorological øbservations.

г 2

the third morning after our arrival from the coast, was placed in charge of François Chartier and Louis la Ronde, with directions for them to proceed with all speed to Isle à la Crosse, at which place Chartier's wife was residing. I wrote to Mr. M'Pherson, requesting him to forward the party without delay; and Mr. Rae, who put up the packet, enclosed, I believe, a circular, soliciting the gentlemen at the several posts to send the packet on as quickly as possible. Mr. Rae himself was of opinion that he enclosed such a document, though he does not perfectly recollect that he did But whether the circular was enclosed or not 80. in the first instance, or afterwards left out, the circumstance of a packet being sent express for fifteen hundred miles ought to have ensured its being forwarded from the further posts. No delay occurred at Fort Simpson, Mr. M'Pherson sending the party on as soon as their provisions could be prepared. Chartier and his companion reached Fort Chepewyan by open water, and were despatched to Isle à la Crosse as soon as the ice was strong enough for travelling over. At Isle à la Crosse the letters were put en route again after a fortnight's detention, and at Carlton House they were kept two months. This last delay was unaccountable. When they did reach Red River they were sent on; but instead of reaching England

in A when desir the " of Jul witho tunat howev chanc Land. On guide Fort S for an the so lost hi astray, hunter failed; tions, v They d after th were no until tl with se Fort C the firs finding

LETTERS.

coast, r and o prowhich ote to rd the out up liciting nd the himself ument, he did or not ut, the ess for red its o delay sending ould be reached ere deice was le à la after a se they ras un-River ngland

in April or May, as we had a right to expect, and when a knowledge of our proceedings was much desired by the Admiralty previous to the sailing of the "North Star," they did not arrive till the middle of July, and our families were nearly twelve months without intelligence from us. We were also unfortunate with our subsequent letters, which were not, however, sent by special express, but were left to the chance of the ordinary conveyance through Rupert's Land.

On the 31st of October, two men and an Indian guide were sent with a second packet of letters to Fort Simpson, hoping that they would be in time for an express which leaves that post annually for the south on the 1st of December. The Indian lost himself, or rather, I believe, went wilfully astray, for the purpose of falling in with some hunters that he expected to find. In this he failed; and the party, after suffering some privations, were saved from starvation by killing a deer. They did not reach Fort Simpson till some time after the winter express had left; and as the letters were not of public importance they remained there until the spring, when they were forwarded along with some others that we subsequently sent to Fort Chepewyan, that they might go down with the first boats. On my way out in the summer, finding part of these letters at one of the posts,

I took them on with me; the others reached England by the same mail packet that I crossed the Atlantic in, and were delivered on the day after my arrival at home.

The only letter-bag from England that we received during our stay at Fort Confidence came in on the 12th of April, 1849, and brought us home news up to the 22nd of June, 1848, ten months This came by the usual canoe route, and was old. brought up from Canada with the Red River mail; but at the same time we received a single newspaper, which gave us some English intelligence as late as the 15th of September. The history of this newspaper is that of the triumph of the electric telegraph. While the English mail packet was steaming up the sound of New York, on the 30th of September, a summary of European news having been carried on shore by an express steamvessel, was in the act of being transmitted by telegraph to the banks of the Mississippi. Within a few hours, it was published there in the "Galena Advertiser," of which it filled one entire folio. This paper, being carried over the plains to Red River, by a party which set out on the day following its publication, was sent to Great Bear Lake, and gave us the first intimation of a rebellion in Ireland. The other newspapers that we received at the same time were of very old date, but every paragraph of them, as well as of our letters, was read again

and a only b tions in Ru extend up his down that h ceding patience examp By Simpso " Spect 1848, v takes i for so dearth tempera insight a com play a world. war, w hunted tility w the vici race.

OREGON "SPECTATOR."

71

and again with a keenness that can be understood only by those who have undergone similar privations of intelligence. We heard of an old resident in Rupert's Land, who was philosophic enough to extend this pleasure over the whole year, by laying up his annual file of newspapers, and taking one down daily for perusal according to its date, so that he had just mastered the news of the preceding year when a new file arrived. Our impatience was too great to permit us to follow an example so systematic.

By the return of our packet men from Fort Simpson in January, we received the Oregon "Spectator," dated Oregon city, February 10th, 1848, with the motto "Westward the star of Empire takes its way." It was a creditable production for so young a state, remarkable for the extreme dearth of "news," but a strenuous advocate of temperance and morality, and curious for the insight which it gave of the first movements of a community destined at no distant period to play a conspicuous part among the nations of the The state is already involved in an Indian world. war, which will not cease until the Red Men are hunted from their native soil. The cause of hostility was one of those unavoidable accidents which the vicinity of white people entails on the Indian race. A large body of emigrants brought small-

F 4

ched ossed day

e rene in home onths 1 was mail; newsgence story f the acket n the news teamtelehin a alena This er, by pubve us The same graph again

72

pox and measles with them, which, spreading among the populous and warlike Kaiyuses or Black-feet, cut off many of the tribe. By the Indian moral code, the death of their brethren was to be revenged by the slaughter of people from whom the injury came, and as it was sufficient if the victims were of the same nation with the offenders, the Kaiyuses fell upon the nearest and most defenceless. A missionary and his family, to the number of twelve persons, were cut off, and their property and some women and children seized. Through the interference of the Hudson's Bay Company's Governor at Fort Vancouver, the captives were redeemed, but five hundred of the inhabitants of Oregon marched to chastise the Indians. The paper says, "the thunders of war have commenced; let them be continued until American property and American life shall be secure upon American soil." We afterwards learnt that the demonstration had little effect upon the Indians, who, being well provided with horses, shunned the encounter, or returned to harass the Oregon army at their pleasure. Sooner or later, however, the Kaiyuses will feel the strong arm of the white man, and be compelled to cede their native lands to the emigrant hordes that are pressing westward.

As the crow flies the distance between Fort Vancouver on the Oregon and Fort Confidence

excee betwe coast, on tl Throu comm and the forwa regula hazard Beside suppoi active furnisk nition, credits to sev object. which success reporte we we sufficie A few a and ou time th woman and a l VENISON.

73

long feet, ioral nged jury were uses A velve some nterernor med, egon says, them rican fterffect with arass r or rong cede are

> Fort ence

exceeds 1350 geographical miles, and the space between the Company's posts on the Labrador coast, or on Lake Huron, and their advanced station on the Porcupine, measures about 2500 miles. Throughout this vast extent of territory, a regular communication is kept up between the Governor and the numerous scattered posts, and supplies are forwarded to all the districts annually with a regularity which cannot be interrupted without hazarding the lives of both traders and natives. Besides the establishment of fisheries for our winter support, Mr. Bell employed several of the most active Dog-ribs in the capacity of fort hunters, furnishing them with clothing, guns, and ammunition, to be repaid in venison. He also gave large credits of ammunition and other articles of trade to several leaders of small bands for the same object. In the end of September and in October, which is the best hunting season, we heard of great Two hundred carcases of rein-deer were success. reported as having been put en cache for us, which we were to send for as soon as the snow was sufficiently deep to permit the dog-sledges to run. A few animals killed near the fort were brought in, and our prospects looked flourishing. In the mean time the Indians sent a sick man and a very aged woman to be nourished by us through the winter, and a large body of old men, elderly widows, and

children settled down near us at a fishing station behind Fishery Island. From this body visitors came to us almost daily, begging for a meal or two of dried meat to vary their diet, or bringing in a trout or two for sale. The fish were always purchased, and then the seller invariably asked for a bit of meat, as he could not walk back without eating. The simple cunning by which these poor folks endeavoured to accomplish their ends, and to move Mr. Bell to be liberal, was amusing, and generally in the end successful, for his habitual good-nature was not long proof against their varied entreaties.

In the first two weeks of October, the ice driving about compelled the fishermen to take the nets out of the water, and during that time the Indian party subsisted mostly on rations from us, being supplied with both meat and fish. As soon as the straits separating Fishery Island from the main froze completely over, which occurred on the 16th of the same month, the old men were well supplied with trouthooks to set under the ice, and they caught, I believe, fish enough for their wants, but they concealed their success that they might continue to draw aid from our store. By dint of much talking, occasionally withholding supplies, and threatening to do so entirely, Mr. Bell at length persuaded most of the party to move towards Cape M'Donald, where fish were reported to be more plentiful. Our own

fisher cline scarc fisher Thsmoo cover out been receiv The friend quant cursic the st to us. made on be could they heavil the la tinuar white the bo suffici Chepe firm. tendir

 $\mathbf{74}$

NATIVE SOCIALISM.

fisheries, however, as well as the Indian ones, declined as the winter wore on, and in February scarcely sufficed to furnish a meal daily to the fishermen themselves.

The snow having by the middle of October smoothed the inequalities of the surface, and covered the stones and stumps, Mr. Bell sent out parties to bring in the venison that had been stored up for us; but instead thereof we received a very beggarly account of empty eaches. The wolverenes had destroyed some; our Indian friends at the fishery had eaten up a greater quantity, having, unknown to us made several excursions for the purpose; and we did not take into the storehouse a tithe of what had been reported. to us. The hunters by whom the caches had been made came in for fresh supplies of amaunition, and on being remonstrated with, merely said, what could they do if hungry Indians came their way? they must eat. This socialist | actice presses heavily on the industrious heater, and encourses the lazy individuals in their idleness; but its comtinuance in force after so long an intercourse with white men is a proof of a fund of good-nature at the bottom of the national character. It is of itself sufficient evidence against the imputation that the Chepewyan tribes habitually desert the old and infirm. We saw on several occasions children attending their sick or aged parents with tenderness

ation sitors two in a purfor a chout poor ad to and situal aried

iving s out party plied traits comsame routlieve, ealed v aid bccag to most here own

76

and solicitude. Instances of desertion, which have undoubtedly occurred, are to be ascribed to the pressure of famine, which has urged the ablebodied to hurry on in quest of relief, disregarding those who were unable to keep up with the line of march.

Our intercourse with the Indians continued throughout the winter in the way that has been stated. The more industrious among them resorted to good hunting stations, generally in parties of two or three families together, and also in two more numerous bands, under the direction of two Most of them resorted to the confines of chiefs. M'Tavish's Bay, where the animals are plentiful in winter. From them we received occasional supplies of venison, and two or four of our men were employed for a considerable part of the winter in bringing it in with the dog-sledges. From two steady old men, who had been furnished with nets, we purchased some hundreds of fine trout, together with a quantity of white-fish and freshwater herrings. Some of the Martin Lake Dog-ribs also, though not fitted out by us with ammunition, found it convenient to bring their meat to Fort Confidence, instead of going to Fort Simpson. In this way we obtained more than we required for our present and future wants, including the eleemosynary demands of the Indians, which were,

howe mana Th journ receiv April

1848

In in au Simps of rou bags c a full exped return So any t Our n veniso two fr

PROVISIONS.

however, kept within bounds by Mr. Bell's careful management.

The following table, extracted from Mr. Bell's journal, gives a summary of all the provisions received into our storehouse up to the middle of April, 1849:—

Months.	No. of Fish.	Fresh Venison.	Half dry Venison.	Pound- ed Meat.	Rein- deer Fat.	Reindeer Tongues.
		lbs.	lbs.	lbs.	k lbs.	No.
1848 September	420	500	1500	200	150	120
October	2370	130	2100		170	105
November	1163	4330	570		18	65
December	560	1830	140			10
1849 January	279	2005				
February	223	1850	560			40
March	176	3165	1680	105	10	125
April			2670	55	5	160
Totals	5191	13,810	9,220	360	353	625

In addition to the above, Mr. Bell brought up, in autumn, 1200 lbs. of dried meat from Fort Simpson, 6 cwt. of barley-meal, and three kegs of rough barley, several 90 lb. bags of flour, some bags of potatoes, with tea and sugar, together with a full supply of pemican for Mr. Rae's summer expedition, and for the provisioning of the men returning to England.

So well provided, we had no dread of want at any time, and passed the winter in abundance. Our men had each a daily ration of 8 lbs. of venison on five days in the week, and on the other two from 10 lbs. to 15 lbs. of fish. The women also

have b the ablecding line

nued been orted es of two two es of ul in supwere er in two nets, ether vater also. tion, Fort son. ired the ere,

received rations, and the children smaller allowances. Barley and potatoes were issued in addition as long as they lasted, and flour occasionally. All the men preferred barley-meal to wheaten flour, as it answered better for thickening the soup, and they thought that it was a more substantial article of dict. The rough barley was beaten in a wooden trough until the husks separated, and then boiled whole along with venison, in which way it made a nourishing soup, that was much relished by all the party. Few of the Europeans consumed the whole of their provisions, and the Indians were generally in attendance at their meals to receive the surplus. Several feasts varied the monotony of our winter life: one was given as a house-warming when the buildings were finished; another, as is customary at all the posts, on the first day of the new year; and two others when the winter was further advanced. On these occasions, the fishermen and wood-cutters were called in, and the whole establishment, man, woman, and child, supped at long tables placed temporarily in the hall. Preparations for the feasts were made by a great baking of bread, pies, and tarts* for two days previously; and tea was served liberally as long as any of the

* We had large supplies of cranberries, bleaberries, and the fruit of the amelanchier, the produce of the country; which, with a few pounds of Zante currants, served for tarts and pies all the winter.

party were up 😗 early morn the m the h under and t A which suppo The r fisher and ty eighty line a varies emplo ture o four p and w to six the sn vary f In ope fishery or fir f

WINTER FISHERY.

allowdition All ur, as , and article rooden boiled nade a all the whole ierally irplus. winter en the omary year; er adn and e estat long ations ng of ously; of the and the h, with all the

party felt an inclination to drink. The tables were then cleared away, and the dance was kept up with vigour to a late hour, or rather to an early one, for the party did not separate till the morning was advanced. Mr. Bell and Bruce were the musicians. The latter, with that aptness which the half-breeds show to learn any thing that comes under their observation, had made his own fiddle, and taught himself to play upon it.

A short description of the modes of fishery by which most of the fur posts in Rupert's Land are supported may not be inappropriate in this place. The nets, formed like those used in the herring fishery, measure, before mounting, one hundred and twenty yards in length, but are gathered in to eighty yards by the introduction of the backingline along the upper edge. The depth of the net varies with that of the waters in which it is to be employed, from two to four yards. For the capture of white-fish, of the ordinary size of three or four pounds, the mesh is five and a half inches long, and where these fish are very large it is increased to six. For taking the Bear Lake herring, and the small coregoni of other localities, the meshes vary from two inches to two and three quarters. In open waters the nets are shot, as in the herring fishery; the upper margin being buoyed with cedar or fir floats, and the lower one depressed by stones.

The fish hang themselves in the meshes, being unable,' from the form of the gill-plates, to withdraw their heads after having once passed them through. Trout of 15 lbs. weight may be taken in the white-fish nets, and also *inconnu* (Salmo mackenzii) weighing 25 lbs.; but the meshes will not admit the heads of the larger trouts (namaycush), which weigh from 30 lbs. to 50 lbs. These are caught with cod-hooks.

In winter the nets are set under the ice. The first step is to make a series of holes, about fifteen feet apart. A pole is then introduced, and conducted along the surface of the water from hole to hole, carrying with it a line, which serves to haul in a string of nets, properly buoyed and loaded, but seldom exceeding five in number. The rope is then detached, and each end of the net is fastened to a piece of wood, laid across its respective hole, or to a stake driven into the ice. On visiting the nets next day only the extreme holes are opened, the rope is attached anew at one end, and is veered away as the nets are withdrawn by the opposite hole. The fish that have been caught being removed, the nets are drawn back to their places by the line. A line of nets reaches about 400 yards, and the fisherman generally endeavours to carry it entirely across a strait or pass in the lake which fish are known to frequent.

Ev care à and the th meshe not of erects shelter snow h will no to turn accoun In t

neatly supplie through to the easily : hole re conveni ration thrown snow-di that the in the : procure the imp there to VOL. I

WINTER FISHERY.

81

being withthem taken Salmo s will may-These

The ifteen conole to haul baded, rope stened hole, g the bened, nd is y the aught their about vours h the Every second or third day, fishermen who are careful take their nets out of the water to dry and repair them. If this be not attended to, the threads swell and rot and few fish enter the meshes; the floats also become water-logged if not often dried. In severe weather, the fisherman erects a canvass or skin screen to windward, to shelter him while he overhauls his nets. Eskimo snow barricades are much more effective, but pride will not permit the Orkney or Canadian fisherman to turn the useful expedients of the Eskimos to account.

In the winter, Albert built a snow wall very neatly round the water-hole by which the fort was supplied, to keep off the snow drifts, cut steps through the ice down to the water, and then fitted to the aperture a light snow-lid, that could be easily removed. By this contrivance the waterhole required little clearing for a week, and the convenience was great; but after the first admiration of his ingenuity subsided, the cover was thrown aside, and the hole allowed to fill up with snow-drift. The consequence of this neglect was, that the first man or woman who went for water in the morning had half an hour's hard work to procure it, and then it was necessary to remove all the impurities left by the dogs which had resorted there to drink on the preceding day.

VOL. II.

G







Albert found more ready imitators in another practice which he taught the men. He was appointed to attend to the officers' fires, and immediately set about preparing his wood sledge according to his own fashion. He first coated the runners with earth or clay tempered with water, coat after coat freezing as rapidly as it was applied. Hot water was used in this operation, otherwise it would have frozen too quickly for him to give it the convex form and smoothness that were necessary. He next washed the runners with water, polishing the ice with his naked hand as it formed. Canadians and Europeans looked on carelessly, merely saying to one another, "What can the savage be about ?" but none of them having the most distant idea that they would follow his example next day. The four sledges employed by the woodmen were of equal size, and each was drawn by two men. The drawers of wood went out together, were equally loaded, and, to their extreme surprise, Albert and his companion outstripped them all on the journey home; their emulation was excited, they laboured hard the whole day, and at night confessed that they were fairly beaten. Then they tried Albert's sledge, and found it run so easily, that forthwith they requested him to prepare their sledges in the same way; and during the winter the sledge-men invariably dressed their runners in the Eskimo mode.

T the its y This in se clum the latte seldo are r were ing 1 Arcti with with south ing p not b institu nerall and y scarce defect compa elicit s Hudso Bay in them

REIN-DEER.

83

other s ap-1 imledge d the vater. plied. vise it rive it neccswater, ormed. elessly, in the ng the his exved by h was ent out streme ripped ulation e day, beaten. it run im to during their

The venison that we obtained was the flesh of the small or barren ground rein-deer, which drops its young on the coasts or islands of the Arctic Sea. This kind does not penetrate far into the forest even in severe seasons, but prefers keeping in the isolated clumps or thin woods that grow on the skirts of the barren grounds, making excursions into the latter in fine weather. A full grown well-fed buck seldom weighs more than 150 lbs. after the intestines The bucks of the larger kind, which are removed. were mentioned in a preceding chapter as frequenting the spurs of the Rocky Mountains near the Arctic Circle, weigh from 200 lbs. to 300 lbs., also without the intestines. Whether these be the same with the woodland rein-deer, which inhabit the southern districts of Rupert's Land and the adjoining parts of Canada, and of the United States, has not been determined, no comparisons having been The small barren ground deer are geinstituted. nerally in excellent condition in the proper season, and yield the very finest venison, hence they can scarcely be supposed to have been dwarfed through defective pasture; and it is probable that a rigid comparison of examples of the several kinds would elicit specific differences. The rein-deer that visit Hudson's Bay travel southwards towards James' Bay in spring. In the year 1833, vast numbers of them were killed by the Cree Indians at a noted

G 2

pass three or four days march above York Factory. They were on their return northwards, and were crossing Hayes River in incredible multitudes. The Indians, excited by the view of so many animals thronging into the river, committed the most unwarrantable slaughter; man, woman, and child rushed into the water and stabbed the poor deer wantonly, letting most of the carcases float down the stream or putrefy on the beach, for they could use only a small number of those they slew. From that date the deer did not use the pass until last year, when a few resumed their old route and were suffered to go unmolested, the Indians not being prepared for their coming. Mr. Rae made two successful excursions in search of deer in the winter, and the Hopes likewise went a short way into the barren grounds on the same errand, but though the latter also killed animals, their caches being carelessly made, were invaded by the wolverenes, and our store reaped no advantage from their efforts. The wolverene is extremely wary, and shows extraordinary sagacity and perseverance in accomplishing its ends. The Indians believe that it is inspired with a spirit of mischief, and endowed with preternatural powers. Though more destructive to their hoards of provision than the wolf, or even the bcar, and able to penetrate fences that resist their powerful efforts, it is only about

thir one com be I a log gnav it sh

* I depths exact of our follows

Lengtl betw Length

Total l Height "Breadt

"

The leg when s portionquadrup matted exist an each sid secretion A young it bring inhabiti

WOLVERENES.

thirty inches long, a foot high at the shoulder, and one foot six inches at the rump, but it is very compactly made.* With teeth that do not seem to be peculiarly fitted for cutting wood, it will sever a log equal to a man's thigh in thickness by constant gnawing. In selecting the spot it intends to breach it shows as much skill as the beaver, generally

* Its Dog-rib name is Noh-gaiye, pronounced from the depths of the throat with a strong aspirate on the h. The exact dimensions of a wolverene, which was surprised in one of our *caches* and killed in the month of March, were as follows:—

Length from root of tail, measured along the b	ack	and	Inch.
between the ears to the point of the nose -	-	•	30.5
Length of stump of tail	-	-	6.2
" of long hairs at top of ditto	-	-	5.5
Total length of animal, including the entire tail	-	•	42.5
Height at fore shoulder	•	-	12.0
"at rump	-	-	12.6
Breadth of fore paw, the toes moderately spread	-	-	2.5
" of hind paw	-	•	2.7

The legs are remarkably muscular, the fore ones in particular, when skinned, having a strong resemblance to a finely proportioned, muscular human arm, rather than to the limb of a quadruped. The paws are covered underneath with long matted fur in winter, so that the small callosities that do exist are discovered with difficulty. The anal glands, one on each side, of the size of an olive, are filled with a very fetid secretion of a yellowish colour, and the consistence of cream. A young female produces two young; an older one four; and it brings forth later in the summer than other ferine animals inhabiting the same districts.

G 3

tory. were The imals st unchild deer down could From il last were being e two n the t way l, but caches wolfrom wary, rance elieve , and more n the ences bout

contriving to cut a log near one end, so that it may fall down into some void space, and thus open an entrance into the hoard. The animal works so hard in carrying on this operation that it causes its mouth to bleed, as the ends of the logs and the snow often testify. Once admitted into the hoard, it has to gnaw the pieces of meat asunder, as they are generally frozen together, and then it proceeds to drag them out one by one, and to bury them in the snow, each in a separate place. As it travels backwards and forwards over the meat, it smears it with a peculiarly fetid glandular secretion, after which no other animal will touch it. In this way one of these beasts will spoil a large cache in an hour or two, and wholly empty it in a few nights. The pieces which are carried off are so carefully concealed in the snow, and the wolverene makes so many tracks in the neighbourhood, that it is difficult to trace out the deposits, and they are seldom found.

Where there are trees, the meat *caches* are generally made with logs let into each other at the corners by notches, as in building a log-house. This, as we have seen, can be invaded by the wolverene. Mr. Rae, however, made a safe cellar by cutting a hole in the ice, covering it thickly with snow, and then pouring water over all, until the frost had rendered the whole a solid mass.

nei The into the dam of d for an pick are fero bret abje sidin but : log, He and sisted behin to b The made satisf drive mont who

that it s open orks so causes nd the hoard, is they roceeds nem in travels smears ı, after iis way e in an nights. refully akes so is diffiseldom

es are at the house. he wollar by y with til the

WOLVES.

Wolves also follow the hunter, and lurk in his neighbourhood, to share in the produce of his gun. Their strength enables them to break occasionally into a cache, but they have neither the skill nor the tenacity of purpose of the wolverene, and the damage they commit is generally on the carcases of deer recently slain, while the hunter has gone for a sledge to bring them in. On his return in an hour or two he often finds only the well picked bones. These wolves, though of large size, are a timid race, and seldom or never exhibit the ferocity and ravenous boldness of their Pyrenean brethren. When reduced by famine they are very abject and unresisting. Mr. Bell once, while residing on Mackenzie's River, caught a full-grown but famished wolf in a marten-trap tied to a small log, which it had not strength to carry away. He went to the fort for a line to lead it home, and the children who accompanied him back assisted in bringing it in, by pushing it on from behind. It made no resistance, and suffered itself to be tied quietly to the stockades of the fort. The experiment of taming it was not, however, made, and after the curiosity of the people was satisfied, it was killed. At another time, a wolf, driven by hunger, was prowling about Fort Edmonton, when, being scared by some of the people who were passing, it took shelter in the kitchen.

G 4

The cook, an old Canadian, who was busily engaged in frying pancakes, was frightened by the aspect of his visitor, and oversetting the frying-pan in the fire, and leaping into bed, he hid himself beneath the blankets. The poor wolf, astonished at the novelty of the scene, and amazed by the blaze of the flaming grease and the screams that issued from the bed, retreated into the square of the fort, and was there killed by the people who had rushed from their several houses on the alarm being raised.

One of Mr. Rae's hunting and exploring excursions was made in the month of December, in the coldest period of the winter; and he informed me that at that time the vapour which rose from the rein-deer completely hid the individuals of a herd; so that, unless he could approach a detached one, or get into the midst of them, he could not take an aim.

Notwithstanding the indolence of a large part of the Dog-rib and Hare Indians who resorted to Fort Confidence, and their total disregard of truth, they had the merit of being strictly honest with respect to property, and also of being quiet and unwilling to offend. No precautions were taken to guard knives and other articles used in the house to which they had easy access, but they meddled with nothing, and we missed none of our effects.

The con and mag that then dini them part they come ment main gazin view, In

presse a low had b en ca their distar young From ration with huntin any th

INDOLENCE OF THE DOG-RIBS.

89

gaged ect of n the meath t the aze of issued e fort, ushed being

g exer, in ormed from of a ached d not

art of Fort they spect illing tuard se to Idled fects. The thermometers, of which at least a dozen were constantly hanging up outside, were never touched, and none of the natives ever intruded into the magnetic observatory, after a general intimation that they were not to do so. When parties of them came in with venison, they slept in our dining-hall, and their friends from the fishery joined them to hear the news, and to talk for a great part of the night, yet, though the place was crowded, they gave us very little trouble. Some of the new comers would frequently enter the sleeping apartments, and crouching down against the wall, remain in perfect quietness for an hour together, gazing at the books and other things exposed to view, and watching Mr. Rae and myself writing.

In December, January, and February, the Indians pressed heavily on our store, as the fishery was at a low ebb, and they had either consumed, or we had brought in, all the venison that had been put en cache for us. The more active hunters, with their families, had followed the game to a greater distance, and several detachments of stout but idle young men had joined the fishery encampment. From these able-bodied fellows we steadily withheld rations, though they were more than once furnished with provisions and ammunition to go out on a hunting excursion. The efforts to get them to do any thing were, however, ineffectual. They gene-

rally returned when their food was exhausted, saying that the animals were all gone. They were afraid, I believe, to venture far in the proper direction, lest they should meet enemies of whom they are in constant dread, and especially of their spiritual foes. Their complaints of the want of game were, however, proved to be unfounded, by the excursions of Mr. Rae and the two Hopes.

In January, intelligence came that the hunters had stored up a number of deer at a considerable distance from the fort, but the news reached Cape Macdonald as soon as Fort Confidence, and a party from the fishery set off in quest of it. They were, nevertheless, too late, some of their wandering countrymen having discovered and consumed the store before their arrival, and they were compelled to return, in a famishing condition, to the fishery. Our men had also a fruitless journey to the empty The lies that were told on these occasions cache. were innumerable, and every one was ready to clear himself and inculpate some other party. We generally, however, succeeded after a time in finding out the real delinquents, who consequently were coolly received, and had short rations when they came to the fort.

As the months of February and March rolled away, and the days lengthened, cheering reports from the hunting parties came in, and some of the mor Mr. who prov stati sum their they so, t leng hunt comi party they sledg fit ou on th took duce of th the c foiled Mr. prom secui At brou had
PROVISIONS NOT INDIVIDUAL PROPERTY. 91

where ber diwhom f their rant of ed, by s.

unters lerable d Cape a party were, dering ed the npelled ishery. empty casions ady to We n finduently when

rolled eports of the more active of the fishery party went to join them. Mr. Bell, on two several occasions, fitted out the whole of the residents at Cape Macdonald with provisions enough to take them to the hunting stations; but their hearts failing them, they consumed what was given to them, and came with their daily petitions as before. During this time they were taking trout, and denying that they did so, though they occasionally sold us a few. At length, towards the end of March, two of our fort hunters. who had left their wives at the fishery, coming to fetch them, the desire of the whole party to eat venison became uncontrollable, and they came in en masse to the fort, with their sledges and all their moveables, to receive another They remained encamped near the house fit out. on this occasion for about three weeks before they took their departure, subsisting chiefly on the produce of their trout lines. During this time some of the young men made two excursions to plunder the caches of the hunters employed by us, but were foiled in the attempt, and came back fasting; for Mr. Bell, taught by former losses, had been very prompt in sending men for the venison, and had secured it before the marauders reached the spot.

At this time some considerable supplies were brought in by the Martin Lake Indians, and we had several opportunities of observing the way in

FORT CONFIDENCE.

which these people act towards each in regard to The venison intended for us was provisions. neatly packed on the sledges, but each Indian generally carried a kind of knapsack on his shoulders, containing some choice pieces of meat to be consumed on the journey back to his tent. The sharp eyes of the hungry party at the encampment discovered the approach of strangers while yet at a great distance, and the rumour of an arrival spreading with rapidity, men, women and children crowded into the square of the fort. The first act of the new-comers was to run the loaded sledges at once into the store-house, which was opened to receive them; but as they arrived in succession, the women from the camp generally pressed in, and throwing their arms round a young hunter, with much kindness of manner would say to him, "It is long since we have seen you, my relation; how have you fared since we met? You are a generous man!" and so on. While his attention was thus engaged, and before he could free himself from the unwashed sirens, whose unwonted softness of speech led him to suspect either ridicule or plunder, one of the females having cut the strings of his knapsack would carry it off, amidst the laughter of the crowd. The young fellow, thus despoiled of his provisions, however much he might be vexed in secret, was

obli pres upor dist mate part he d the exce Marc a ve pains the f infan skin a was b This two l and 1 shoul suffer pursu before to it task, hunti an Ir footst

AN ACCOUCHEMENT.

gard to us was Indian on his of meat is tent. the enrangers nour of women he fort. run the , which arrived enerally a young uld say ou, my ? You ile his could whose suspect females would crowd. isions, t, was

obliged to join openly in the mirth; and the expression of face of some of the youths thus preyed upon, as they endeavoured to force a smile in their distress, was irresistibly comic. The loss fell ultimately on the store, as Mr. Bell had to furnish the party with food for their return, though in that case he did not certainly select titbits. When at length the great move was made, and the fishery party, exceeding forty in number, went off, in the end of March news came that the wife of one of the hunters, a very small woman, had been taken with the pains of labour. The three females belonging to the fort went to her aid, and found the new-born infant, the mother's first child, wrapped in a deer's skin and stuck into a hole made in the snow. It was brought into the house, and dressed by the fire. This event delayed the mother's departure about two hours. She then set out dragging a sledge, and having her first-born suspended between her shoulders, in a bag or Indian cradle. She was not suffered, however, to profit by the well-beaten path pursued by the rest of the party, who had gone before her, but had to make a new track parallel to it through the leose snow, always a laborious task, even to a stout man. Want of success in hunting, or some other calamity, was sure to befal an Indian who should incautiously tread in her footsteps. This was the custom; no slight or

FORT CONFIDENCE.

unkindness was shown to her; her husband was, I believe, really fond of her; and her sledge was a light one, being loaded only with things belonging to herself.

Soon after what may be considered a great event in the simple annals of Fort Confidence-the departure of so numerous a body of hangers-on-we had a visit from a Dog-rib who had been residing on the other side of the lake, at the plentiful fishery opposite Fort Franklin. Our visitor, a stout, ablebodied fellow, came empty-handed, and introduced himself as a very serviceable man, who had been of great use to our two fishermen stationed at Fort We readily understood that the truth Franklin. concealed under this self-praise was, that he had been living by their labours, perhaps bringing in a little wood for them occasionally, but really sharing in the produce of their nets. On questioning him more closely as to the object of his coming, he at length said that he came to look for a wife. Had he no wife? Yes, he had had one, but an Indian had taken her from him. Had he an aversion to his wife that he had parted from her without a struggle? No; his wife was a very good wife; she suited him very well. Then why did he not fight for her? "You see," said he, "it was a big Indian that took her! I am a little fellow, what could I do !" This hero met with little countenance from mati more and, helpi TH being upon blind lines, of th gener who quest the re of de in th sprin tunes Af rein-d India activ quest them He d dried quan

MIGHT MAKES RIGHT.

95

nd was, e was a longing

it event the dem-we ding on fishery it, ableoduced been of at Fort e truth he had ging in really n quesof his ook for ad one, Had he om her y good did he is a big , what enance

from us, and after being fed for two days, an intimation was made to him that he would receive no more rations; on which he followed the others, and, as we afterwards learnt, soon procured another helpmate.

The singular national custom of the women being the property of the strongest, had been acted upon a short time previously at the fishery. A blind man, who was more assiduous in setting trout lines, and more successful as a fisherman than most of the others, was deprived of his wife on the general movement of the party, by an old fellow who wanted her to drag his sledge. Mr. Bell questioned the ancient Paris about the truth of the report when he came to the fort, and instead of denying the fact, or seeming ashamed, he gloried in the deed as a manly action. A child, the offspring of her former connection, followed the fortuncs of the mother.

After Mr. Bell and I quitted the fort in May, the rein-deer migrating towards the sea-coast left the Indians again in straitened circumstances. The active hunters moved towards the open country in quest of musk-oxen, and the more helpless threw themselves on Mr. Rae's compassion at the fort. He distributed among them the remains of the dried meat brought from Fort Simpson, and a quantity of musk-bull meat, which was too strong

FORT CONFIDENCE.

for his own men, but which the natives relished greatly. Thus furnished they betook themselves reluctantly to various fisheries. Such is the life which these poor creatures lead. Occasional feasting and rejoicing, with intervening periods of want, sometimes of absolute famine. With proper management, the natural resources of the country would support a population ten times as great; but as long as all the drones of the community claim a right to appropriate to their own wants the produce of the exertions of an industrious hunter or fisherman, no certain provision for the future will be made. The first step in advance will be the formation of fishing villages, and the culture of barley and potatoes; and, under the guidance of intelligent missionaries, this might be effected without much difficulty; while, at the same time, the truths of Christianity might be brought to bear on the heathenism and moral defects of the 'Tinnè nation.

The preceding details may lead the reader to imagine that the Dog-ribs or Hare Indians are an unhappy race; but such is not the case. They are timid, and assume the attitude and solicitations of beggars in their intercourse with white people; but among themselves they are lively, volatile, and full of fun and mirth, which even an empty stomach cannot suppress.

"In weel teml open Octo As tl the a in cd am in trude reside Th rigoro down forma eight

with t

enoug

they a

or by

each c

circles

firmly

is at f.

of a

of a t

mains vor

PROGRESS OF THE SEASONS.

97

relished mselves the life al feastiods of proper country great; munity wants istrious for the idvance and the der the ight be at the ght be moral

der to are an hey are tations cople; le, and omach With regard to the progress of the seasons, the "Indian summer," as it is called, brought us three weeks of fine weather after our arrival in September. The centre of Bear Lake usually remains open till late in December, but by the middle of October the bays and straits are frozen across. As the structure of ice has of late years attracted the attention of speculative geologists, principally in connection with the movements of glaciers, I am induced to mention here a few facts which intruded themselves on my observation during my residences in the fur countries.

The first step in the freezing of rivers in this rigorous climate, after the water has been cooled down to 32° by a succession of cold weather, is the formation of somewhat circular plates of ice, six or eight inches in diameter. These drift for a time with the current, until they have become numerous enough to cover the surface of the water, when they are arrested in a narrow part of the river, or by any slight obstacle, and speedily adhere to each other, after which the interstices between the circles fill rapidly with crystals that bind all firmly together. The sheet of ice thus produced is at first nearly opaque; but when, in the course of a day or two, it has acquired the thickness of a few inches, it becomes transparent, and remains so until a fall of snow has obscured the VOL. II. Iľ

FORT CONFIDENCE.

In unsheltered lakes the wind drifts the surface. snow to the beach, and would perhaps keep the ice clean for great part of the winter, were it not that in certain hygrometric conditions of the atmosphere small starry tufts of most beautiful tabular and latticed crystals are deposited at short intervals on the ice, and freeze firmly to it. In a dry atmosphere these crystals evaporate again, but should a fall take place of the fine, dust-like snow, which is the most common kind in the high latitudes, they serve to detain it until it consolidates, so as to resist the wind. It is rare, however, for the snow to lie more than a foot deep on any of the large lakes, unless where it has drifted under the lee of piled-up slabs of ice, or of rocks, islands, or other shelter.

During winter the ice receives an increase of thickness from beneath, and at the same time evaporates above; the latter process going on with a rapidity that would scarcely be credible to one ignorant of the extreme dryness of the air in an arctic winter. The ice acquires a thickness of from four to eight feet, according to the severity of the season, the depth of the lake, and other modifying circumstances; and I desire here to advert especially to the fact, that although it is constructed of successive horizontal additions beneath, when it decays in spring it consists of

ICE AND SNOW.

99

fts the the ice ot that atmotabular ort in-In a in, but e snow, gh latiblidates, ver, for any of d under islands,

ease of le time on with to one ir in an ness of severity l other nere to ugh it lditions sists of

vertical prisms, penetrating its whole thickness, and standing side by side like the columns of a basaltic cliff; which, in their mode of formation, have, I imagine, a close analogy. Dr. Slagintweit informed me that neither the ice nor the basalt forms exact prisms, the angles never having the precise measurements of true crystals. In this condition the ice may be strong enough to support a considerable weight; and I have travelled over it with a large party on several occasions when the prisms on which the foot rested were depressed at every step, and a pointed stick could be driven through the whole thickness into the water beneath, with as much ease as into a bank The ice then, in fact, presents the phyof snow. sical characters of a semi-fluid mass, as pointed out by Professor Forbes, its parts being moveable on each other, not only vertically, but as in the case of travelling glaciers, capable of gliding past one another horizontally.

In spring, when the action of the sun-light is very powerful, an incipient thaw takes place at mid-day on the surface of the snow, which, on freezing again, acquires a glassy crust. As the season advances, but while the temperature of the air is still even at noon far below the freezing point, the crust in clear weather becomes penetrated in the direction in which it is struck by the

н 2

FORT CONFIDENCE.

100

sun's rays at mid-day by innumerable canals, and finally crumbles into a granular mass like the *firn* of the high Switz glaciers, that crackles under the feet as soon as the sun sinks towards the horizon. This firn is not universal; it is more common within the arctic circle, and in situations where there seems to have been originally a certain looseness in the texture of the snow, and where its surface is so much inclined that the sun's rays do not fall on it obliquely about noon. I did not notice it in any quantity on the level surface of a lake.

The rapid evaporation of both snow and ice in the winter and spring, long before the action of the sun has produced the slightest thaw or appearance of moisture, is made evident to residents in the high latitudes by many facts of daily occurrence; and I may mention that the drying of linen furnishes a familiar one. When a shirt, after being washed, is exposed in the open air to a temperature of 40° or 50° below zero, it is instantly rigidly frozen, and may be broken if violently If agitated when in this condition by a bent. strong wind, it makes a rustling noise like theatrical thunder. In an hour or two, however, or nearly as quickly as it would do if exposed to the sun in the moist climate of England, it dries and becomes limber.

Mr. Rae mentioned to me another example of

the bou mer Bay part froz lifte The and boul on it In atmo man to R The scale room body ness bed, out cloth appro door, the s the e In

TEMPERATURE.

101

, and e firn er the rizon. mmon where loosere its ays do notice ke. ice in ion of ppearnts in occurf linen being empetantly lently by a thea-

er, or

to the

es and

ple of

the same fact, which bears on the transportation of boulders, and may interest geologists. During his memorable residence on the shores of Repulse Bay, he noticed several large boulders which were partially exposed at low water. When the sea froze they became engaged in the ice, and were lifted with it from the bottom by the flood-tides. The ice gaining at each tide in thickness beneath, and losing above by superficial evaporation, the boulders in process of time came to rest in pits on its surface.

In consequence of the extreme dryness of the atmosphere in winter, most articles of English manufacture made of wood, horn, or ivory, brought to Rupert's Land, are shrivelled, bent, and broken. The handles of razors and knives, combs, ivory scales, and various other things kept in the warm rooms, are damaged in this way. The human body also becomes visibly electric from the dryness of the skin. One cold night I rose from my bed, and, having lighted a lantern, was going out to observe the thermometer, with no other clothing than my flannel night-dress, when, on approaching my hand to the iron latch of the door, a distinct spark was elicited. Friction of the skin at almost all times in winter produced the electric odour.

In November the snow was deep enough for

н З

FORT CONFIDENCE.

sledges to run without receiving much injury. On the 1st of December the sun was just visible for an instant at noon, from an eminence behind This month was a cold one, and the the house. coldest days in the year were the 17th and 18th, when the average temperature for forty-eight hours was 551° below zero of Fahrenheit. The lowest observation made was at seven in the afternoon of the 17th, at which time the temperature was 58.9° F. The thermometer, one of Adie's making, was hung by the side of a dozen by the same artist, none of which differed a degree from When tested by freezing mercury, this one it. stood at 36°, which is considered to be from 4° to 6° too high; so that, making the smallest of these corrections, the actual minimum temperature we observed was 65° F.* This is one of the greatest colds on record, and pains were taken to ascertain it correctly.†

* Mr. Saunders, commander of the North Star, records $64\frac{1}{2}^{\circ}$ F. as the lowest temperature observed in Wolstenholme Sound in the winter of 1850.

† In two thermometers made by a London artist, and hung up beside the others, the spirit retreated into the bulb, though the scales were graduated down to 73°. In freezing mercury, these thermometers indicated 55°, being about 15° too low. Mr. Adie's thermometers were constructed under the superintendence of Professor Forbes of Edinburgh. The precautions used to insure their correctness will be detailed when the meteorological observations at Fort Confidence are published.

half requ the days zon limb emin of F howe been sky a Janu On 9 o'c rapid withc 2 P. M light. beaut above in th shone a sky weath beam In (or 6

TEMPERATURE.

njury. visible pehind id the 18th, -eight The afterrature Adie's by the from is one 4° to these re we reatest ertain

ds 64½° Sound

l hung though ercury, 7. Mr. ndence ised to logical Even at mid-winter we had three hours and a half of daylight. On the 20th of December I required a candle to write at the window at ten in the morning. On the 29th the sun, after ten days' absence, rose at the fishery, where the horizon was open; and on the 8th of January, both limbs of that luminary were seen from a gentle eminence behind the fort, rising above the centre of Fishery Island. For several days previously, however, its place in the heavens at noon had been denoted by rays of light shooting into the sky above the woods. The lowest temperature in January was 50° F.

On the 1st of February the sun rose to us at 9 o'clock and set at 3, and the days lengthened rapidly. On the 23d I could write in my room without artificial light from 10 A.M. to half-past 2 P.M., making four hours and a half of bright daylight. The moon in the long nights was a most beautiful object; that satellite being constantly above the horizon for nearly a fortnight together in the middle of the lunar month. Venus also shone with a brilliancy which is never witnessed in a sky loaded with vapours, and unless in snowy weather, our nights were always enlivened by the beams of the Aurora.

In February the lowest temperature was 56° F. (or 62° corrected), and in March, 44° F. (48°)

11 4

FORT CONFIDENCE.

was the lowest observed. On the 20th of April, signs of the snow softening on the south side of the house, contiguous to the walls, were perceived. The day by th' ime had lengthened so far, that I was able to read off the degrees of the thermometer by daylight at nine in the evening. Snow birds arrived in small flocks, and on the 27th the snow began to melt in sheltered places, exposed to the direct rays of the sun. Ospreys, gerfalcons, eagles, and gulls appeared on the 17th of May; and, from the 19th to the 23d, melted snow stood deep on the surface of the ice. On the prior of these two dates, the first goose was seen. The geese approach the high latitudes as soon as the swamps are uncovered, when they feed on the undeveloped flowering stalks of the cotton-grass and other Cyperaceae. Their arrival is thus an indication of the progress of spring, and frosty weather will sometimes drive them back for a week or so to a milder district. The impulse, however, by which they are urged to their breeding stations is so uncontrollable, that in backward seasons they are driven to the sea coast before the snow is gone, and then, from want of food, they are in a very lean condition. Their incubation, the fledging of the goslings, and the moulting of the parents, has to take place before the end of August, when old and young pass southwards to spend the winter

in 1 lead suin war The Indi enca run in ea TI breed arriv then north latter On 24th, flocks north longand r earlie day. shove fur o \mathbf{Th} subse Mr. I

ARRIVAL OF SUMMER BIRDS.

105

April, side of ceived. r, that hermo-Snow th the osed to alcons, May; v stood rior of . The as the he unass and ication er will to to a which so uney are gone, a very ing of ts, has en old vinter

in more genial climes. An indelible attachment leads them back to their natal places, and the ensuing summer sees them winging their way northwards in cuneal bands, with unerring instinct. Their arrival in a district enlivens white man and Indian: during their passage, plenty reigns in every encampment; and the dingy, pot-bellied children run about with smiling, greasy faces, brandishing in each hand the leg or wing of a goose.

The Canada geese come in the van, and remain breeding in the woody country; snow geese next arrive, and pass onwards to Wollaston's Land; then the laughing geese come and go, holding a north-west course; and at the same time with the latter, the Hutchin's geese speed to the sea coast.

On the 22d, pin-tail ducks were seen; on the 24th, swans; and, on the 30th and 31st, large flocks of snow geese and brown cranes passed northwards. On the 1st of June, bees, sandpipers, long-tailed ducks or cacawees, eider and king ducks, and northern divers were seen: the catkins of the earliest willows also burst their envelopes on this day. On the 5th, teal, widgeon, scaup-ducks, shovellers, and jagers arrived; but, on the 8th, the fur of the polar hare was still white.

The progress of spring at Fort Confidence, subsequent to the 7th of May, is recorded from Mr. Rae's notes, as on that date Mr. Bell and I

FORT FRANKLIN.

commenced our journey southwards. To contrast with the above the dates of the arrivals of the migratory birds at Fort Franklin in the same season are here added, the difference of latitude between the two places being a degree and three quarters.

On the 11th of May, under a hot sun, a pool of water had formed on the ice near the bay of the Deer Pass. We bivouacked on the shore beside it, and had not yet arranged our sleeping-places, when a Canada goose alighted in the pool. It was scarcely allowed to settle before it was shot, and, with a celerity unknown in civilised lands, stripped of its feathers and committed to the cooking-kettle. This was evidently a straggler, and must have seen the small pond in which it alighted from a great distance; for, on our arrival at Fort Franklin on the following day, we learnt that neither our two fishermen employed there, nor an Indian residing near them, had as yet seen any of the spring birds. On the 14th the Indian saw gulls; on the 18th, snow geese and various small birds came, together with the pretty little gull named Xema bonapartii, which in large flocks sought for insects in the open water now forming along the shores of the smaller lakes. On the 22d, bands of snow geese passed to the north-west, flying high. They evidently found the country about Fort Franklin still too closely wrapped in its winter

garb the I Ge at Pe north varyi date time and c on w Canad two. high believ avails goose, flight. small notice confir any of it is g Wh was e out to snow

* Th Land.

ARRIVAL OF SUMMER BIRDS. 107

ontrast the miseason etween rters.

pool of of the beside places, ol. It s shot, lands, to the aggler, hich it arrival learnt there, et seen Indian rarious little flocks rming e 22d, flying about vinter garb, and were winging their way to the valley of the Mackenzie, where the season is earlier.

Geese, according to Mr. Bell's information, arrive at Peel River Fort, upwards of two degrees farther north, from the 12th to the 15th of May, rarely varying above a day or two, the 15th being the date of their coming in backward seasons. At that time they find the marshy places bare of snow, and can procure the roots of bents and other plants on which they feed. There, as clsewhere, the Canada geese precede the snow geese a day or two. The Hutchin's geese* come later, and pass high overhead towards the north. The Indians believe that a small finch (Plectrophanes lapponica) avails itself of the strength of wing of the Hutchin's goose, and nestles among its feathers during its flight. When a goose is shot, they often see the small bird flying from it. Neither Mr. Rae nor I noticed such an occurrence, nor did I obtain a confirmation of it from the personal observation of any of the gentlemen resident in the country, but it is generally affirmed by the Indians.

While we were at Fort Franklin, Mr. Bell, who was employed all day in shooting geese, pointed out to me a fact in the natural history of the snow and laughing geese. Though they migrated

* These are commonly called "Eskimo geese" in Rupert's Land.

FORT FRANKLIN.

in large flocks, each had already selected its mate; and if the female was shot, the male bird instantly separated from the rest, and descended to look after her. In this way, he often fell a victim to his conjugal fidelity; but if he escaped the shots aimed at him, and became shy, he would still continue for hours, and even days, searching the neighbourhood for his lost mate. The case was different if the male bird fell first. The female, it is true, also left the flock, but she kept more aloof, and generally, after making a circle or two round the spot where the body of her partner lay, went off with the next flock that came up.

The laughing geese passed Fort Franklin a few days later than the snow geese, but a single individual was often seen some days before the arrival of the main body, associated with a flock of snow geese, and generally acting as leader by assuming a station in the apex of the angle in which they fly. About the same time, the American robin, or migratory thrush, came with the yellowpoll and black and yellow warblers (*Sylvia æstiva et maculosa*). The latter fed on the berries of the Alpine arbutus, as did likewise the golden plovers, whose stomachs also contained the juicy fruit of the *Empetrum nigrum*. The Eskimo curlew at this time fed on large ants. It would appear that these insects descend to the stomachs of the curlews | fast 1 mand death huntir malis) were a began was a The 1 was ol ground one w of the pellets owl. River easily (C. glassifier)grey r before merate I had

* I h which b press it I could two syll two last

ARRIVAL OF SUMMER BIRDS.

109

stantly stantly to look etim to e shots d still ng the se was female, t more or two her lay,

a few single ore the a flock der by gle in herican vellowæstiva of the lovers, uit of ew at r that e cur-

lews alive, since I found that several, having taken fast hold of the lining of the gullet with their mandibles, remained sticking there, and even after death required some force to detach them. The tree hunting (Fringilla canadensis), black-finch (F. husmalis), and white-crowned finch (F. leucophrys)* were also early visitors, and soon after their arrival began to construct their nests. The Lapland finch was also seen, but only on its passage to the coast. The Lestris richardsonii flew about in pairs, and was observed to have the habit of quartering the ground like the hen-harrier. In the stomach of one which I killed, there were the skin and some of the bones of a mouse, rolled into a ball, like the pellets that are rejected from the stomach of an The purple-throated diver visited Bear Lake owl. River in considerable numbers. This species is easily distinguished from the great northern diver (C. glacialis), while flying, by its swollen, bluishgrey neck. Almost all the summer birds arrived before we left that neighbourhood; but I have enumerated only the earliest comers, or those which I had not previously seen in so high a latitude,

* I have already mentioned the nocturnal song of this bird, which breeds throughout Rupert's Land. In attempting to express its clear, loud notes by syllables, the nearest approach I could make was *cheet-cheet*, *tareet*, *cheet*. The first two syllables are loud and high, the next two short, and the two last lower and softer.

VALLEY OF THE YUKON.

110

and whose range is, therefore, not correctly given in the Fauna boreali-Americana.*

· From Mr. Murray, I have received the following account of the arrivals of the water-fowl in the Valley of the Yukon. "Of the two kinds of swan, only the largest sort (Cygnus buccinator) are seen here; they pass on to the northward of the Porcupine River, to breed among the lakes. Bustards (i.e. Canada geese) are plentiful, and breed everywhere, from Council Bluffs on the Missouri to the vicinity of the Polar Sea. On the ramparts of Porcupine River they frequently build high up among the rocks, where one would suppose only hawks and ravens would have their nests. How they take their young down is unknown to me, but they must be carried somehow. Ravens and large gulls are very destructive to young geese. With respect to the breeding-quarters of the laughing geese (Anser albifrons), I am able to inform you correctly, having myself seen a few of their nests; and, since the receipt of your letter, made further inquiry among the northern Indians. Their nests are built on the edges of swamps and lakes, throughout most of the country north of the Porcupine, where the ground is marshy. It is only near the most northerly bends of that river that any are seen in the breeding season, and these are male birds. They pass to their breeding-places in the beginning of June, and make their nests among long grass or small bushes, where they are not easily seen. They are shy birds when hatching; and, when any one comes near the nest, manage to escape unperceived, and then show themselves at a distance, and manœuvre like grouse to lead the intruder away from the place. Notwithstanding our ruthless habit of collecting eggs of all kinds to vary our diet, I have often felt for a laughing goose, whose anxiety for the safety of its eggs was frequently the means of revealing to us the situation of its nest. When the bird was swimming some hundreds of yards off, immediately that any person in walking round the lake came near its treasure, the poor bird began to make short, impatient turns in the water, resuming her calm demeanour if

Gi extei over

the int eggs a the he These than ar are per Gens d larly to any fly (snow there a seen. abunda siderabl or brow head an descript nicla). that ar velocity do, som water; very fa taste. their ap geese a other k end of t in June shores of in Octo

ARRIVAL OF SUMMER BIRDS. 111

Great Bear Lake is navigable for its whole extent for only fifty days in the year. It is frozen over later than shallower pieces of water in its

the intruder passed the nest without seeing it. As soon as the eggs are taken, the goose rises out of the water and flies elose to the head of the eaptor, uttering a frightened and pitiful ery. These geese are more numerous in the Valley of the Yukon than any other kind; and the numbers that pass northwards there are perhaps equal to that of all the other species together. The Gens du large (Neyetse-kutchin), who visit the north coast remlarly to traffie with the Eskimos, say that they have never seen any flying northwards over the sea in that quarter. White geese (snow geese, Chen hyperboreus) are also passengers here, and there are likewise black geese, which I presume you have never seen. A few of them pass down Peel's River, but they are more abundant on the Yukon. They are very handsome birds, considerably smaller than the white geese, and have a dark brown or brownish black colour, with a white ring round the neck, the head and bill having the shape of that of the bustard." (This description applies pretty well to the brent goose (Anser bernicla). The black geese are the least numerous and the latest that arrive here. They fly in large flocks with remarkable velocity, and generally pass on without remaining, as the others do, some days to feed. When they alight, it is always in the water; and, if they wish to land, they swim ashore. They are very fat, and their flesh has an oily and rather disagreeable taste. Bustards, laughing geese, ducks, and large gulls make their appearance here from the 27th to the 29th of April. Snow geese and black geese about the 15th or 16th of May, when the other kinds become plentiful. They have mostly passed by the end of the month, though some, especially the bustards, are seen in June. The white geese and black geese breed only on the shores of the Arctic Sea. They return in September and early in October, flying high, and seldom halting."

y given

g account e Yukon. gnus bucrd of the ards (*i. e.* a Council

On the high up nd ravens Jown is vens and espect to lbifrons), a few of e further ilt on the try north nly near n in the to their eir nests ily seen. e comes en show lead the ruthless I have afety of ituation reds of and the e short, nour if

GREAT BEAR LAKE.

vicinity, but the ice remains longer unbroken, and drift ice continues on its surface till the middle of August. On the 7th of May, the day on which Mr. Bell and I set out to travel to Fort Franklin, the ice was as firm as in mid-winter; and, though the snow softened in fine weather, it still covered the ground deeply.

MR. RA TION RIVEI ESKID BAY.-— DA AND A LAKE. IN TH PORTA PART TINUE POOL.-FOR S

Having occurre Confide is the account to reach in that boats u Eskimo plan of been re three bo would vol.

MR. RAE'S EXPEDITION.

n, and dle of which nklin, hough overed

CHAP. XVI.

MR. RAF'S EXPEDITION IN THE SUMMER OF 1849. — INSTRUC-TIONS. — HE CROSSES TO THE COPPERMINE. — DESCENDS THAT RIVER. — SEA COVERED WITH ICE. — SURVEYS RAE RIVER. — ESKIMOS. — CAPE KENDALL. — CAPE HEARNE. — BASIL HALL BAY. — CAPE KRUSENSTERN. — DOUGLAS ISLAND. — DETENTION. — DANGEROUS SITUATION. — AUGUST 23., RETURN. — AUTHOR AND MR. BELL LEAVE FORT CONFIDENCE. — CROSS GREAT BEAR LAKE. — DESCEND BEAR LAKE RIVER. — DAVID BRODIE LOST IN THE WOODS. — HIS ADVENTURES. — FORT SIMPSON. — METHY PORTAGE. — RECEIVE ENGLISH LETTERS. — NORWAY HOUSU. — PART FROM THE SEAMEN AND SAPPERS AND MINERS. — CON-TINUE THE VOYAGE TO CANADA. — BOSTON. — LAND AT LIVER-POOL. — SUMMARY OF THE PRESENT STATE OF THE SEARCH FOR SIR JOHN FRANKLIN.

HAVING in the preceding chapter mentioned such occurrences during our winter's residence at Fort Confidence as seemed most worthy of notice, this is the most appropriate place for introducing an account of Mr. Rae's endeavours, in the summer, to reach Wollaston Land, and complete the search in that quarter. Had we succeeded in taking our boats up the Coppermine, beyond the reach of the Eskimos, according to our expectations when the plan of search was formed, the voyage might have been resumed in the summer of 1849 with two or three boats; and, in that case, both Mr. Rae and I would have gone, that we might aid each other VOL. II. I

114

MR. RAE'S EXPEDITION.

among the ice. But, having been compelled to leave our craft in September without the smallest hope of their being found again in a seaworthy condition, and having only one boat remaining that could be employed on the service, it became necessary to determine which of us should take charge of that vessel and of the small party it could contain. Setting all personal considerations aside, and looking solely to the means of providing for the examination of as large a portion of the Arctic Sea as could be accomplished, I had no hesitation in deciding in favour of Mr. Rae. His ability and zeal were unquestionable; he was in the prime of life, and his personal activity and his skill as a hunter fitted him peculiarly for such an The arrangement I made for withenterprise. drawing the European party, and employing volunteers from the men engaged in the country, was a considerable pecuniary saving, which I was bound to consider, as far as I could, without cramping the means of search.

It has been already mentioned that Mr. Rae explored the country between Fort Confidence and the Coppermine River in winter, to select the best route for dragging the boat over in spring. In April he conveyed provisions, boat stores, and other necessaries across to the Kendall on dog-sledges, and left two men at Flett's station in charge of

MAY,

the the tain dryi proc men nam

1849

Ne Jan Th Ha All Lou

Bap Lou

(Memo "A Disco the co Coppe part o aminat toria J Union these t lying l almost flood t

INSTRUCTIONS.

MAY,

1849.

elled to smallest aworthy maining became ild take it could ns aside, ding for e Arctic esitation ility and e prime his skill such an or withig voluny, was a as bound ramping

Rae exence and the best ing. In nd other -sledges, harge of

them, together with two Indian hunters. Before the end of the month we learnt that they had obtained both musk-ox and rein-deer meat, and were drying part of it for summer use. The subsequent proceedings are told in the following official documents, which I quote entire, premising that the names of the men composing his party were-

Boat's crew.

Neil M'Leod	-	-	-	-	Orkney man.
James Hope	-	-	-	-	Cree Indian.
Thomas Hope	•	-	-	-	"
Halcro Humphrey	-	-	-	-	Orkney man.
Albert One-eye	-	-	-	-	Eskimo.
Louis Olivier	-	-	-	-	Canadian.

Left in charge of Fort Confidence.

Baptiste Paul	-	-	-	••	Canadian.
Louis Dubrill	-	-	-	-	,,

"Fort Confidence, Great Bear Lake,

(Memorandum.)

May 1. 1849.

"As in the prosecution of the search for traces of the Discovery Ships under command of Sir John Franklin, the continental coast line between the Mackenzie and Coppermine Rivers has been carefully examined, the only part of my instructions not yet complied with, is the examination of the adjoining shores of Wollaston and Victoria Lands, which the state of the ice in Dolphin and Union Straits rendered inaccessible last autumn. That these two islands are separated from each other by a strait lying between the 111th and 113th meridians, is rendered almost certain by a consideration of the direction of the flood tide, which, on the west side of these parallels, sets

MR. RAE'S EXPEDITION.

to the westward through Dolphin and Union Straits, and to the eastward of them, sets to the eastward towards Cape Alexander; coming, we must conclude, from the northward between the lands in question: for the survey by Messrs. Dease and Simpson has shown that the coast of Victoria Land is continuous up to the 111th parallel; and the latter gentleman records his opinion, that much of the heavy drift ice that encumbers Coronation Gulf descends between these lands from the north.

116

"The exploration of the shores of this strait is of much importance in the search for the Discovery Ships, for the following reasons:-Sir John Franklin having been directed to steer to the south-west after he had passed Barrow's Straits, would be led directly into it, and he would be deterred from attempting a more westerly course by the circumstance of Sir Edward Parry having found that route impracticable for two successive seasons. Should there be several islands between Wollaston and Banks's Lands, and the channels between them be intricate, it is not unlikely that the ships may have been shut up therein by ice. It was the intention of Sir James Ross, in the event of his reaching Winter Harbour last year, to send a party across the ice this spring to pass between Victoria and Wollaston Lands towards Cape Krusenstern and the Coppermine River. To co-operate with that party and to aid it with provisions, or to supply its place, should circumstances have prevented its being sent, it is expedient that a party should go from hence, and as you and a sufficient number of men have volunteered for this service, I hereby, in virtue of the clause of my instructions which authorises me to detach you and a party of volunteers under your command, appoint you to this duty. You are therefore to descend the Coppermine River; and as soon as the sea opens in July, are to proceed to explore the strait in

1849

MAY.

ques of I neigl coast stone ccedi that paint. the sa the gu you c Shoul party, that th place, you ar from t or obt the shi quarte "H perienc

perience advance safety your reinstance commen " Ha as prace

* The

prudent of Dolpł private 1 Mar,

1849.

aits, and cds Cape orthward Messrs. Victoria and the h of the descends

of much s, for the n directed Barrow's ıld be dey the eirhat route d there be 's Lands, s not unherein by the event d a party toria and and the ty and to d circumlient that sufficient I hereby, huthorises der your therefore s the sea strait in

INSTRUCTIONS.

117

question, endeavouring to communicate with any parties of Eskimos you may meet with on Wollaston or the neighbouring islands. Should you reach the northern coast of Banks's Land, you are there to erect a pile of stones, and deposit a memorandum of your object and procccdings at the distance of 10 feet from its base, marking that side of the pile with a broad arrow in red or white paint. You are also to erect similar piles, and deposit in the same manner on conspicuous headlands, memoranda for the guidance of the party detached by Sir James Ross, when you can do so without materially delaying your progress. Should you discover any signal posts erected by that party, and learn from the memoranda deposited near them that the strait has been sufficiently explored down to that place, you are to proceed no further in that direction; and you are at liberty to use your own judgment in deviating from this route, if, from information given by the Eskimos, or obtained from other sources, you are of opinion that the ships, or part of their crews, may be found in another quarter.

"Having the fullest confidence in your judgment, experience and prudence, I shall not name a period to your advance, further than by requesting you not to hazard the safety of the party entrusted to your care, by delaying your return too long. The last season furnishes a strong instance of the early date at which the winter occasionally commences in these seas.*

"Having performed this service, or prosecuted it as far as practicable, with a due regard to the safety of your

* The 25th of August was the date at which I considered it prudent that Mr. Rae should endeavour to be on the south side of Dolphin and Union Straits, and I expressed this opinion in a private note to him.

MR. RAE'S EXPEDITION.

party, you are to return with all speed to Fort Confidence, and, embarking without delay the instruments and stores remaining at that post, to proceed forthwith to Fort Simpson. Such of the stores as are useful to the Company are to be valued and handed over to them, and the instruments are to be forwarded to England, addressed to the Secretary of the Admiralty. The men are to be sent to whater at some fishing station sufficiently to the southward to ensure their early arrival at Norway House next spring; and you are to direct them to be furnished with nets, that they may provide for their own sustenance during the winter, with as little expense to the Government as possible.

"Immediately on your return from the coast, you are to communicate an account of your proceedings to the Secretary of the Admiralty, for the information of their Lordships; and you are also to transmit to him a chart of any hitherto unexplored coasts or straits you may discover, as soon as you have had leisure to construct it.

"Given under my hand, at Fort Confidence, 1st May, 1849.

(Signed) "JOHN RICHARDSON, Commanding Aretic Searching Expedition. "John Rae, Esq."

Copy of a Despatch from Mr. Rae to the Secretary of the Admiralty, narrating the Proceedings of the Expedition under his Command to the shores of the Arctic Sca, in the Summer of 1849.

> "Fort Confidence, Great Bear Lake. September 1. 1849.

" Sir,

"I have the honour to acquaint you, for the information of my Lords Commissioners of the Admiralty, that the 1849 expe

MAY,

perm exam searce place and was unfor "J fidem Copp

ice on 7th o of the I star a coup were and s haule "C

the m long) the 12 it was and S ascence obstruc coldnee high g almost snow o it was sufficie

ASCENDS THE DEASE.

MAY,

1849.

et Confients and n to Fort he Comand the ressed to be sent ne southbuse next shed with ustenance Govern-

you are is to the i of their chart of discover,

1st May,

on, cpedition.

ry of the Expedition c Sea, in

r Lake.

ormation that the expedition under my command, which descended the Coppermine this summer to the Arctic Sea, for the purpose of examining the shores of Wollaston and Victoria Lands, in search of Sir John Franklin and party, returned to this place to-day, having been quite unsuccessful in its object, and with the loss of Albert, the Eskimo interpreter, who was drowned at the Bloody Fall, the particulars of which unfortunate accident I shall mention hereafter.

"Having made every requisite arrangement at Fort Confidence for facilitating our progress across land to the Coppermine, I waited impatiently for the disruption of the iee on Dease River, to which our boat was hauled on the 7th of June. Next day we learned that the upper parts of the river were clear of ice; and on the following morning I started in company with four men and two Indians, and a couple of sledges on which our baggage and provisions were stowed. The Dease was still covered with strong and solid ice for two miles up its course, over which we hauled the boat before getting to open water.

"Our ascent of the stream was extremely slow, owing to the many barriers of ice (some of them six or eight miles long) over which we had to launch the boat, and it was the 15th before we arrived at the forks of the river, where it was my intention to diverge from the route of Dease and Simpson. They followed the north branch, whilst we ascended the south-east one. This stream was also much obstructed by ice, and so very shallow, consequent on the coldness of the weather, which prevented the snow on the high grounds from thawing, that the whole party were almost continually up to their knees among water and snow engaged in launching the boat. In ordinary seasons it was evident, by marks along the banks, that there is sufficient water for a boat drawing some inches more than

MR. RAE'S EXPEDITION.

ours throughout its whole length, which is little more than 15 miles, including its various curves. On the 17th we passed over the ice on the lake from which the stream flows. It contains many islands, and its breadth, where we crossed it in a nearly south direction, is 31 miles.

"Indian report had led me to believe that there was a creek, having sufficient depth of water for the boat, flowing from this lake into the south branch of the Kendall, which we were to descend to the Dismal Lakes; but in this we were disappointed, and consequently had a portage of six geographical miles to make overland nearly due east. I had examined this place in the winter, but the ground being then covered with snow and ice, I could not form a correct opinion on the subject. The west end of the portage is situate in latitude 67° 10′ 48″ N.; longitude by account * 117° 18′ W.; and the variation of the compass 50° 49′ E.

"Crossing the portage occupied us two days; the ice had not yet broken up in many places on the river on its east side, and the water was 10 inches lower than when Sir John Richardson and party forded it last autumn; we consequently found some difficulty in descending it. Its general course is north-east by east, and length from where we entered it, to its influx at the west end of the most easterly of the Dismal Lakes, $17\frac{1}{2}$ miles. On the 21st we arrived at the station on the banks of the Kendall River, to which provisions for the sea voyage had been hauled on dog sleds in April, and found the two men who had been left in charge quite well. We descended the Kendall next day to the Coppermine, which was still covered with ice, so firm and solid that a person might

* The rates of the chronometers had become so irregular, or had altered so much, that they were of no use until rated anew. have in wi 66 occuj ing d when merid 07' 2 azimu and 5 birch willow day th to des but af blocke terrup we arr in doin of our men), shippir " H easily double

more a

by the

could

remain

circum

of fish Halket

some 1

1849.

JUNE,

JUNE,

1849.

e than 7th we stream where 3.

was a lowing which this we of six ast. I ground form a of the oude by ompass

the ice r on its n when in; we it. Its where e most e 21st Cendall d been n who ed the s still might

> lar, or anew.

DESCENDS THE COPPERMINE.

121

have crossed the river without being more than ankle deep in water.

"During five days that we were detained here, we were occupied repairing the injuries received by the boat, shooting deer to save our pemican, and making observations when the weather would permit. The result of three meridian observations of the sun gave mean latitude 67° 07' 20" N., and the mean variation from five sets of azimuths on different days (the extremes being 49° 38', and 51° 55') was 50° 37' 48" E. On the 28th the dwarf birch was observed to be in leaf, and the leaf-buds of the willows began to develope. In the afternoon of the same day the river was thought sufficiently open to permit us to descend it for some distance among the driving ice; but after proceeding six miles, we found the stream again blocked up. We were so often and so long detained by interruptions of this kind, that it was the 11th of July before we arrived at the Bloody Fall, having been fourteen days in doing the work of one. Notwithstanding the inefficiency of our steersman James Hope (one of Dease and Simpson's men), we ran all the rapids, including the Escape, without shipping much water, and with all the cargo in the boat.

"Hitherto deer had been so numerous that we could easily have shot enough for the maintenance of a party double or treble our numbers. Here they had become more scarce and shy, which could be only accounted for by the proximity of the Eskimos, no recent traces of whom could however be seen. From the fall to the sea the ice remained fixed until the 13th, when it cleared away, a circumstance that was very soon indicated by the numbers of fish which appeared below the fall. With the aid of Halkett's air-boat, which had been brought from a hill some miles distant, where it had been left last season, a net was set in one of the eddies, and before the men had finished arranging it, seven fine salmon and two white fish were caught, and we afterwards obtained a supply for several days' consumption.

"On the 14th we entered the sea, and found a narrow and very shallow channel along the shore of Richardson Bay, until we came to its north side, where the ice lay against the rocks. Here the latitude 67° 51' 19" N. was observed; and two azimuths of the sun, the one on the meridian and the other when on the prime vertical, gave variation 57° 04' and 56° 25' E. Whilst encamped at this place, we were visited by seven Eskimos, one of whom I at once recognised as the active, intelligent man who had afforded Sir John Richardson's party such efficient assistance last season, when crossing the river at the head of Back's Inlet. On inquiry I learnt that they had been well supplied with provisions in the early part of winter and in spring; but that in the interval they had nearly starved, owing to the scarcity of seals, having had to subsist for some time on the skins of the larger species of these animals, which they had reserved for making boots. In the winter they had communicated, either directly or indirectly, with the natives of Wollaston Land, none of whom had ever seen whites, large boats, or ships. They were all made happy by some small presents, and a supply of fish, which they ate raw, and appeared to relish much. They left us near midnight, promising to return next day with some boots and shoes for sale.

"On the 16th, by making a number of portages over the ice, we rounded Point Mackenzie and entered Back's Inlet, which was partially open. Having a fine breeze from the cast, we set sail and soon ran to its head, when we entered Rae River (discovered and named by Sir John

184 Rid up VISI seci run " alor the the assi At is a stre rock and fine and two, grap and vary stroi draw and or le 200 as th " party

JULY.

party that as th days knew

SURVEYS RAE RIVER.

July,

1849.

nen had hite fish oply for

narrow hardson ice lay N. was e on the al, gave d at this whom I who had t assisthead of id been f winter d nearly had to species making , either n Land, or ships. s, and a to relish return

bver the Back's breeze I, when ir John Richardson last autumn); and on proceeding three miles up it, came to the lodges, six in number, of our Eskimo visitors, who said that they had been so much alarmed at seeing the boat under sail that they were on the point of running away.

"As there was no possibility of our making much progress along shore until the ice wasted a little more, I devoted the two following days to an examination of this river; the Eskimo whom I have already mentioned as our active assistant last year willingly agreeing to accompany us. At the distance of $9\frac{1}{2}$ miles from the river's mouth, there is a perpendicular fall of 10 feet, which extends across the stream, except a few yards on the north side, where the rock slopes so much that, during the spring-floods, salmon and white fish are able to ascend, affording the natives a fine opportunity of spearing them. Here I left the boat and four of the men, whilst in company with the other two, and our Eskimo guide, I traced the river 191 geographical miles further. Its course is nearly due west, and very straight; about the size of the Dease River, and varying in breadth from 80 to 200 yards, with a very strong current, and sufficient depth of water for a boat drawing 14 or 15 inches. It flows over a bed of limestone, and is bounded on the north, at the distance of half a mile or less from its banks, by precipices of basalt from 100 to 200 feet high, superimposed on limestone of the same kind as that which forms the bed of the river.

"At the extreme west point of our journey, we found a party of ten Eskimos with their families, who informed us that the stream maintained the same course and size as far as they had seen it, which was somewhat more than three days' march, or about sixty miles; how much further they knew not, as they had never been to its source. Two of

MR. RAE'S EXPEDITION.

JULY,

this party returned with us to the boat, where they received presents, which I had some difficulty in getting them to accept, as they said they had nothing to give in exchange. On the evening of the 18th we ran down stream, and landed our guide among his friends, who seemed very glad to see him return safe. They were now much less timid than when we first met them, and we pitched our tent close to theirs, without causing any apparent alarm, although it was afterwards observed that two of the men kept watch during the night.

"Early on the 19th we took a friendly leave of these simple and inoffensive people, and pulled down to the mouth of the river, where I stayed until noon, when the latitude $67^{\circ} 55' 20''$ N. was observed. The cloudy state of the weather during the two preceding days had prevented any observations being made. New moon occurring to-day, it was high water a few minutes before 1 P. M.; the rise being 10 inches. In the afternoon, when on our way to Cape Kendall, we experienced a severe thunder-storm from north-north-west, which obliged us to land for shalter.

"Our advance along the coast was to slow, the we did not arrive at the place where the boats were left last autumn until the 24th. We found that they had been much broken up by the Eskimos to obtain the ironwork. The tents, oil-cloths, and part of the sails, still remained uninjured, and were of much value to us, as we were illprovided with the two first of these articles. The 'cache' of pemican and ammunition was also untouched, having apparently escaped notice from being covered with snow. The latitude 68° 10′ 44″ N., and variation 56° 8′ E., were observed here.

"On the 27th a west-north-west breeze having cleared away the ice for a short distance from the shore, we con-

1849 tinu befo 68° side a lig wide of th wate 3 P. when from again haste save i seasor evide where come, white, " V neares passin for ou been a high r hand, the ot in the served change

"A and on deer, o

July,

1849.

cceived lem to hange. landed to see d than close to tit was during

f these mouth e latiof the ed any day, it he rise way to m from er. we did ft last l been hwork. nained ere illcache ' aving snow.

> learcd con-

were

CAPE KRUSENSTERN.

tinued our course towards Cape Hearne, which we reached before noon, and found its extreme point to be in latitude 68° 11' 17" N. Basil Hall Bay being filled from side to side with unbroken ice, we encamped here. Next forenoon a light south-south-east breeze opened a crack in the ice, wide enough for the boat to cross to an island in the middle of the bay, on the north side of which we found some open water, which enabled us to get two miles beyond it. At 3 P. M. on the 30th, we arrived at Cape Krusenstern, and when opposite its high cliffs a strong breeze sprung up from north-north-east, which drove the ice so forcibly against the rocks, that we were obliged to unload with all haste, and haul the beat up on a drift-bank of snow to save it from being squeezed. Here for the first time this season we found the ice broken up in the offing, caused evidently by the strong currents of the ebb and flood tides; whereas on looking in the direction from which we had come, all, except immediately along shore was smooth, white, unbroken, and apparently as firm as in winter.

"We were now at the most convenient though not the nearest point for making the traverse to Wollaston Land, passing close to Douglas Island, and there was no neccessity for our proceeding further along the shore, even had we been able to do so, which at present was impossible; the high rocks presenting an insurmountable barrier on the one hand, and the ice by its roughness equally impassable on the other. We pitched our tents on the top of the cliffs, in the ascent of which the before-mentioned snow-bank served as a ladder, and waited for the first favourable change in the ice.

"A few days afterwards, Albert (the Eskimo interpreter) and one of the men, when some distance inland looking for deer, overtook five Eskimos travelling to the interior with

MR. RAE'S EXPEDITION.

loads of salmon, which they had speared in a rivilet that falls into Pasley Cove. From these the interpreter learnt that the sea ice had commenced breaking up only one day before our arrival, and that they had been in company with the natives of Wollaston Land during the winter, none of whom had ever seen Europeans, large ships, or boats.

"During our long and tedious detention here, several gales of wind occurred, principally from the northward, but the space of open water was so small that they produced little effect upon the ice. Our situation was most tantalising to all the party : occasionally at turn of tide a pool of water, a mile or more in extent, would appear near us, and every thing would be prepared for embarkation at a minute's notice, in expectation of the opening increasing and permitting us to cross to Douglas Island, but our hopes were always disappointed. A number of observations were obtained, which placed our encampment in latitude 68° 24' 35'' N., the longitude being very nearly the same as that assigned to it by Sir John Richardson and Mr. Kendall. The mean variation of the compass, from eight sets of azimuths, on different days, and at different hours, was 59° 8' 08" E., the extremes being 57° 42' and 61° 25'.

"The ice continued driving to and fro with the tides, without separating sufficiently to allow of the practicability of passing among it until the 19th August, when there was more open water to seaward than we had yet observed, caused by a moderate southerly breeze that had been blowing for the last two days. After waiting some hours for a favourable opportunity of forcing our way through a close-packed stream of ice that was grinding along the rocks as it drove to the northward, we at last pushed off, and after more than once narrowly escaping being squeezed, we reached comparatively open water, where we had room

to u and we c that fog h the so visab early south west away we to some We I Point and th scarce and in visible "T

1849

AUG.

for the was in by lau the 22 view v scope i white east ar betwee breadt Hearn "As as ther
SEA COVERED WITH ICE.

127

to use our oars. We had pulled more than seven miles and werc still three miles from the island (Douglas), when we came to a stream of ice, so close packed and so rough that we could neither pass over nor through it; a thick fog had come on, and the ebb tide was carrying us fast to the south-east. Under these circumstances I thought it advisable to return towards the main shore, on which we landed early on the morning of the 20th, a short distance to the south of the place from whence we had started. A northwest breeze came on some hours after landing, and cleared away the ice a few yards' distance from the beach, of which we took advantage to shift our quarters, which being under some crumbling cliffs, were neither safe nor convenient. We poled along shore for three miles or more towards Point Lockyer (the only direction in which we could go), and then took shelter in a small bay, into which we had scarcely entered when the wind changed to east-north-east, and in a very short time left not a spot of open water visible, either near shore or at a distance.

"The wind continued to blow from east and north-east for the two following days, during which, when the tide was in, we advanced a few miles to the south, principally by launching the boat over the ice. On the evening of the 22d I ascended a hill near the shore, from which a fine view was obtained. As far as I could see with the telescope in the direction of Wollaston Land, nothing but the white ice forced up into heaps was visible, whilst to the east and south-east there was a large space of open water, between which and the shore a stream of ice, some miles in breadth, was driving with great rapidity towards Cape Hearne and its vicinity.

"As the fine weather had now evidently broken up, and as there was every appearance of an early winter, I thought

Aug.

1849.

ilet that one day any with none of ats.

ral gales but the ed little lising to f water, nd every minute's and perpes were were ob-68° 24' e as that Kendall. sets of ars, was 25'. he tides, icability

he tides, icability here was bserved, ad been he hours rough a ong the hed off,

ueezed,

nd room

MR. RAE'S EXPEDITION.

it would be a useless waste of time to wait any longer in hopes of being able to cross to Wollaston Land; I therefore gave orders for our return towards the Coppermine, at which I did not expect to arrive in less than a week, as the ice wore as unfavourable an aspect as it did last year.

"At an early hour on the 23d the men commenced carrying the baggage to Point Lockyer, still more than a league distant, and afterwards hauled the boat to the same place; doing this gave all the party ten hours' hard work; but our fatigues were soon forgotten on finding some open water on the south side of the point, in which we were speedily afloat and sailing before the fast-decreasing breeze. It fell calm in half an hour; but we plied our oars to such good purpose, that we pitched our tents late at night on Cape Hearne. Here I had expected to find the ice close packed on the shore, and quite impassable, but I was agreeably surprised to discover as we advanced, that there was a lane of open water between the beach and the pack, wide enough for us to pass through. The only way that I can account for this is, by supposing that the gale of wind that had been blowing from north-east and eastnorth-east had shifted to the north and north-west as it approached the land, and carried the ice along with it.

"By working seventeen hours next day we came to the mouth of the river, and on the following morning ascended to the Bloody Fall. Here fish were still very numerous, and whilst some of the party were cooking, others set a couple of nets to obtain a supply for some days. I may here mention, that when on the coast, we obtained as many salmon and herring as we could consume, wherever there was a piece of open water large enough for setting a net to be found.

"Dease and Simpson, after hauling their boats over the

18 ro ha bo un tin hay the of app was smo how was the suffi on s to s the eddi ran, whic was, came and eithe be of befor dang small ever, the la a lea nor d VC

Aug.

AUG.

longer in

1849.

FATAL ACCIDENT

I thereermine, at ek, as the year. ommenced re than a the same ard work; some open h we were ng breeze. irs to such night on e ice close out I was that there l the pack, y way that he gale of and eastwest as it ith it. me to the ascended numerous, hers set a I may tained as wherever setting a

over the

rocks opposite the lower and strongest part of the fall, had them towed up the remainder by water; and as our boat was now much weakened by the rough usage it had unavoidably been exposed to, for the purpose of saving time, and the tear and wear that would be caused by having it dragged over the portage, I was desirous of doing the same. I was the more led to do this, as the men were of opinion that it might be effected with safety. All that appeared in any way difficult was easily done, and there was only one short place to be ascended, which was so smooth that a loaded boat might have passed it; here, however, from some unaccountable cause, the steersman was seized with a sudden panic, and called to those towing the boat to slack the line. This was no sooner done sufficiently to allow him to get firm footing, than he leaped on shore, followed by the bowman, and allowed the boat to sheer out into the current, where the line broke, and the boat soon oversetting, was carried into one of the eddies some distance down stream; to which Albert and I ran, and stationed ourselves at two points of rock near which the wreck would pass. It drove to where Albert was, and he hooked it by the keel with an oar until I came to his aid, when I fixed a pole in a broken plank and called my companion to assist me in holding on; he either did not hear me correctly, or thinking that he would be of more use on the bottom of the boat, sprung to it, and before I had time to call him off, or even think of his danger, they were carried by a turn of the current into a small bay, where I believed both were safe; not so, however, for the next minute they were swept out again, and the last I saw of our excellent interpreter was his making a leap towards the rocks; he missed them and disappeared, nor did he rise again to the surface. VOL. II. K

MR. RAE'S EXPEDITION.

"This melancholy accident grieved me much, as the brave lad was universally liked for his activity, lively and amiable disposition, and extreme good nature. On James Hope, the steersman, rests all the blame of the loss of the boat; his carelessness in using a small towing-line when there was a much stronger one in the boat, and his cowardice where there was no danger, can admit of no excuse.

"On taking up our nets (which we laid carefully on the rocks for the Eskimos), they were found full of herringsalmon, in fine condition. We commenced our journey, across land, towards Great Bear Lake, on the 26th, each of the men carrying about 90 lbs., and my own bundle being nearly 50 lbs. Three days' easy march brought us to that part of the Coppermine (lat. 67° 12' N.) from which it was my intention to make a straight course to Fort Confidence. Here we expected to meet with some Indians, but we did not fall in with them until the following evening, when our loads were much lightened, and we arrived here early this afternoon (being our seventh day from the Bloody Fall), accompanied by upwards of thirty of the natives, who had joined us at different parts of the route.

"The two men who had passed the summer at Fort Confidence were well; and having all the stores ready packed, I shall leave this to-morrow, after supplying the Indians with ammunition to enable them to hunt their way to the trading-posts.

> "I have, &c. (Signed) "JOHN RAE.

"Fort Simpson, September 26. 1849. "P.S. I arrived here this afternoon, and intend sending off the expedition men to Big Island, Slave Lake, on the 28th enab prov yern

M is to with letter resul prope prese Coron where expec the m these flectio Eve of the

of the indebt throug approp sense of co-ope cheereabsence a grea

ICE IMPASSABLE.

28th, with an ample supply of nets and twine, so as to enable them to procure sufficient fish for their winter provisions, without being any additional expense to Government.

(Signed) "J. RAE."

131

Mr. Rae's failure in crossing to Wollaston Land is to be attributed solely to the strait being filled with impracticable ice. I know from his private letters that the mortification he experienced in the result is much more severe than he has thought proper to express in his official despatch. The presence of ice in Dolphin and Union Straits and in Coronation Gulf for two or more successive seasons, where the experience of former years had led us to expect a comparatively open sea, is suggestive of the manner in which a party may be shut up in these regions, and leads to many melancholy reflections.

Every reader of my narrative of the proceedings of the expedition will be aware of how much I was indebted to Mr. Rae's activity and intelligence throughout its progress; and this seems to be the appropriate place for me to express formally my sense of obligation to him for his sound advice and co-operation on every emergency. His scciety cheered the long hours of an arctic winter's absence from my wife and family, and it was in a great measure owing to his skill and assiduity

к 2

, as the ively and on James ass of the ine when and his it of no

ly on the herringjourney, 6th, each n bundle cought us N.) from course to with some until the ightened, r seventh pwards of ent parts

at Fort es ready lying the ant their

RAE.

. 1849. sending , on the

LETTER TO MR. RAE.

in observing, that our experiments on magnetism during our stay at Fort Confidence were carried on so as to be productive of scientific results.

In consequence of intelligence obtained from newspapers on my journey home, I addressed the following letter to Mr. Rae:—

"Lake Winipeg, August 19. 1849.

"My Dear Rae,

132

"As I learn from the newspapers which I have just read, and shall forward for your perusal, that Sir James C. Ross did not reach Barrow's Strait till after the 28th of August, and that it is probable that he may have been arrested short of his intended wintering quarters at Melville Island or Banks's Land, and could not consequently send off his proposed spring party to the Coppermine River this season, I consider it likely that he may determine on sending that party next spring; and if so, by the present arrangements they will, on their arrival at Great Bear Lake, find Fort Confidence deserted.

"I therefore think it important that you should engage either the chief of some band, or two expert hunters, to pass the months of June and July, 1850, on the portage between Bear Lake and the Coppermine River, promising them a handsome reward if they render any assistance to the expected white party, and paying them such moderate sums, in addition to a full supply of ammunition, as may content them for spending the summer on such excellent hunting grounds.

"You will have no difficulty in engaging either Martin Lake or Bear Lake Indians for this service; and there is abundance of time, after the arrival of the March packet, by which you will receive this letter, for them to reach Fort will rands consp Statio "T Ross's some left, y and th pense, will sa the lor

"P. this ser

Extrac

"W

party of ne-betai engagin the rou received Europea requeste Indians, them by

LETTER FROM MR. RAE.

133

gnetism rried on

ed from ssed the

9. 1849.

have just Sir James after the may have uarters at not consete Copperthe may and if so, arrival at

ild engage iunters, to he portage promising istance to moderate h, as may excellent

er Martin l there is h packet, to reach Fort Confidence long before the snow begins to melt. I will thank you to furnish them with five or six memoranda in waterproof cases, with directions to plant them in conspicuous places at the mouth of the Kendall, Flett's Station, Fort Confidence, and elsewhere.

"These precautions may prove to be unnecessary, as Ross's party will most likely, early in their march, discover some of your landmarks, and learn, by the notes you have left, your intention of quitting Fort Confidence this season, and thereupon turn back to the ship. But, at a small expense, if the Indians carry their instructions out fully, they will save the party, should it come on, from having to make the long journey round Bear Lake without assistance.

"I remain, &c.

(Signed) "JOHN RICHARDSON.

"P.S. Mr. M'Pherson recommends Tecon-ne-betah for this service."

Extract of a Letter from Mr. Rae to Sir John Richardson.

"Portage La Loche (Methy P.) July 30. 1850.

"When the winter packet arrived and I received your instructions respecting the establishment of an Indian party on the Coppermine, the Martin Lake Chief Teconne-betah was at Fort Simpson, and I had no difficulty in engaging him to pass the summer at certain stations on the route between the Kendall and Bear Lake. He received three notes which were to be delivered to any Europeans he might meet, in which the strangers were requested to put themselves wholly in the hands of the Indians, who would guide them by the best ron' and feed them by the way. In the event of his not seeing any

к З

parties, other notes, with a rough chart of the best route, were supplied well wrapped in oil cloths, which were to be placed on a pole in conspicuous stations.

(Signed) "J. RAE."

MAY

Having by the preceding quotations brought the narrative of the search made by the expedition to a close, I now revert to the period at which Mr. Bell and I left Fort Confidence. A party of men preceded us by a week, taking with them the baggage, which they were directed to carry on to Fort Norman along with some pemican and stores left at Fort Franklin. This was intended as a precautionary measure to avoid the delay that might be caused by Bear Lake River being late in ' breaking up.

On the 7th of May we took leave of Mr. Rae, and left the fort. Bruce, Mitchell, Brodie, M'Leod, and Mastegon accompanied us in our journey over the ice, which was completed on the 12th, having occupied five days and a half. On the northern shores of the lake, the snow had lost little of its depth, and we had to clear it away to the thickness of five feet in making our encampment on Cape M'Donald; but on the southern shores we found some exposed sandy spots of ground bare, and pools of water forming at midday. Mastegon shot the first goose of the season on the 10th, a straggler that was tempted to cross

1849 our the wea befd T near ther pem Rive the the r Acco with dog-On praye Bruc Dore. and I them the a they ' which quick cipate lofty · the t part o

FORT FRANKLIN.

est route, ere to be

MAY

1849.

RAE."

brought spedition hich Mr. of men hem the ry on to nd stores led as a lay that ng late in

Mr. Rac, M'Lcod, journey he 12th, On the had lost away to encampsouthern spots of at mide season to cross our pathway by a pool of water produced under the influence of a powerful sun (p. 105.). Cloudy weather followed, and five or six days elapsed before the geese began to arrive in earnest.

The men who preceded us were at the fishing hut, near the site of Fort Franklin, when we arrived there. They had transported three sledge loads of pemican about seventeen miles down Bear Lake River to the usual winter crossing-place, but, finding the stream open, they had put them *en cache* on the right bank, and returned to wait further orders. According to their report the rivulets were swollen with melting snow, and travelling by land with dog-sledges was at an end for this season.

On the 14th, being Sunday, we assembled to read prayers; and, early on the following morning, Bruce set out for Fort Norman, taking with him Dore, Cousins, Thomas Hope, Mastegon, Plante, and M'Leod. They travelled light, carrying with them merely their blankets and provisions. By the arrangement I had made with Mr. M'Pherson they were to find at Fort Norman a small barge, which Bruce was directed to bring up to us as quickly as he could. The only difficulty we anticipated was at the rapid in Bear Lake River, where lofty walls of ice remain to a late date, covering the tracking-ground. Hope and M'Leod, being part of Mr. Rae's boat's crew, were to return over-

к 4

BEAR LAKE RIVER.

land immediately with some articles from the store at Fort Norman. They did come back nine days afterwards, and passed onwards to Fort Confidence; and we learnt in the sequel that the Mackenzie did not break up at Fort Simpson till the 23d of May, being fifteen days later than Mr. M'Pherson had known it to do during twenty years' residence on its banks.

We remained waiting nearly a month for the barge, having with us Mitchell and Brodie, with the two Fort Franklin fishermen, Hector Morrison and Narcisse Tremblè. Our diet consisted of trout, white fish, Bear-lake herring and geese, the latter being Mr. Bell's contribution to the common stock. Of Fort Franklin the only vestige remaining was the foundation of a chimney stack; and the fishing hut not being large enough to hold us all, we bivouacked under the shelter of a boat's sail, as a substitute for a tent. When the water had run off the surface of the ice on the lake, so that we could transport our effects across it without wetting them, we moved to the banks of Bear Lake River; being glad to quit the vicinity of the hut, which, like all fishing quarters, became extremely disagreeable as soon as the accumulated impurities of the winter were revealed to view by the wasting of the snow. The marshy places or dry sandy banks first became bare, but many wreaths of

the n Lake that conje Bruc infor still barge four d in ex bridge ceivin to the in the Mr. M These

and le

transfe

being

withou

a large

ice cel

precau the do

On

barked

Mitche

1849.

drift

JUNE,

store days

JUNE,

ence; ie did May, n had ce on

or the with rrison trout, latter stock. g was fishing we bi-, as a un off could etting River; which, y disarities asting sandy hs of 1849.

DESCENT OF BEAR LAKE RIVER.

137

drifted snow continued unmelted till the end of the month. We stayed at the encampment on Bear Lake River till the 8th of June, much surprised that the boat did not arrive, and forming various conjectures to account for the delay. On that day, Bruce and Mastegon came to us on foot, bringing information of the lower part of the river being still covered with ice, and that they had left the barge a few miles within its mouth. They had been four days on the march, one of which was occupied in examining the ice, which they stated formed a bridge at the rapid many feet in thickness. On receiving this intelligence, I resolved on descending to the mouth of the river on foot, and after going in the barge to Fort Norman, where I could join Mr. M'Pherson, to send her back for the stores. These were accordingly placed in a secure cache, and left under the charge of Narcisse, to whom we transferred our surplus stock of fish and geese, being sufficient to maintain him eight or ten days without touching the pemican, of which there was a large reserve. The snow drifts formed excellent ice cellars for preserving fresh provisions, the only precaution necessary being to protect them from the dogs by branches of trees.

On the 9th, Mr. Bell, Bruce, and Mastegon, embarked with me in the fishing coble, and Morrison, Mitchell, and Brodie, were directed to walk along

138 DESCENT OF BEAR LAKE RIVER. JUNE,

the bank of the river, each of them carrying his own bedding and clothing. I cautioned them against going inland, and promised that we would wait from time to time at particular points for their coming up. Half an hour after setting out, finding the river smoother than we expected, and Bruce being of opinion that we could embark all the party, we put ashore, and in a short time Morrison and Mitchell joined us, but David Brodie, having struck into the woods with the view of making a straighter course, did not arrive in the hour that we remained waiting for him. Supposing then that he had gone past, we resumed our voyage, taking into the coble Morrison and Mitchell.

The rate of our descent of the stream rather exceeded four miles an hour, and at half-past six A. M. we reached the *cache* situated fourteen miles from the lake. Brodie not arriving in the course of the day, I became convinced that he had lost himself in the woods, and therefore sent Morrison and Mitchell back to the lake to acquaint Narcisse with what had happened, and to endeavour to engage an Indian who was residing at the fishery to go in quest of Brodie. In the mean time we fired our fowling-pieces at intervals, and set fire to some trees that the smoke might be seen from a distance.

inf rive the ken join the that and folld Ν havi the a land in t shor whic pape to th Narc incid easin suffe woul perso and fault lost,

DAVID BRODIE LOST.

139

JUNE,

1849.

them would ts for g out, ected, nbark time frodie, ew of in the Supsumed n and

er exx A. M. from of the imself n and creisse ur to shery ne we t fire from Mastegon, who knew that part of the country, informed me, that ten or twelve miles back from the river, there was an extensive swamp, from which there flowed one stream that fell into the Mackenzie, and another, named the Black River, which joined Bear Lake River about four miles below the *cache*. The latter stream was so rapid, he said, that Brodie would be unable to cross it on a raft, and it was not fordable; he would therefore, by following it, be certainly led to Bear Lake River.

Next day our two men came back from the lake, having themselves gone astray for some hours in the attempt to make a short cut across a neck of land. After placing written directions for Brodie in the cache, we all embarked again, and in a short time came to the influx of the Black River, which was evidently flooded. Here I left another paper of instructions for Brodie, directing him to the cache for provisions, and to remain with Narcisse until the barge came for them. The incident of Brodie's straying gave me much uneasiness, as I feared he would experience some suffering, though I did not apprehend that he He was a man of much would lose his life. personal activity and considerable intelligence, and though his judgment would be probably at fault when he first became conscious that he was lost, I knew that as soon as he was in a condition

BRODIE'S ADVENTURES.

coolly to consider his position, he would be enabled to shape a course for the river by the sun, and following its bank to vturn to the lake. And so it eventually happened. When he discovered that he was walking in a wrong direction, he began to mend his pace and to run as is usual in such cases, but took an inland course, and at length came to the borders of the swamp above mentioned. Here the woods being more open he obtained a distant view of the "Hill at the Rapid," which he recognised from having seen it on his former journey to the cache, and as he knew that we must pass it in descending the river, he resolved on walking straight for it in the hope of arriving there before After this he came to the Black River, and us. being a fearless swimmer, swam across it, carrying his clothes on his head. The stream, being very tortuous, came again in his way, when he crossed it a second and a third time in the same manner, but on the last occasion, his bundle slipping off floated away, and he regained the bank with difficulty in a state of perfect nudity. After a moment's reflection, he came to the conclusion that without clothes he must perish, and that he might as well be drowned in trying to recover them as to attempt proceeding naked. On which he plunged in again, and fortunately landed this time safely with his habiliments. He now refreshed himself with

184 par anx wit froi Jun visi use, that He by l T happ of th were havi in Co of le that learn in th tente horiz mista The fortu suit e many Ał

JUNE

BRODIE'S ADVENTURES.

141

V

JUNE

1849.

abled and nd so l that an to cases, ne to Here istant e reurney ass it lking before , and rying very ossed nner, g off diffimothat \mathbf{night} as to ed in with with part of a small piece of dried meat, which, in his anxiety he had hitherto left untouched, and forthwith decided on finding the *cache*, and returning from thence to the lake. On the third day (11th June), he found my note together with some provisions which we had suspended to a pole for his use, but he had so husbanded his own small supply that he had still a morsel of dried meat remaining. He had no difficulty afterwards in joining Narcisse, by keeping sight of the river the whole way.

This adventure is recorded as an example of what happened to all the seamen and sappers and miners of the expedition each in their turn. Four of them were lost in the winter of 1848-49, for three days, having mistaken their way to the boat encampment The straggling of the others was in Cedar Lake. of less account, but none of them could be taught that they were liable to such accidents until they learnt it by experience. One man who strayed in the winter on Cedar Lake, when found, was contentedly steering for the moon, which, being near the horizon and gleaming red through the forest, was mistaken by him for the fire of the men's bivouack. The snow which covered the ground at the time fortunately enabled the Indian who went in pursuit of him, to trace his steps before he had gone many miles.

About twelve miles below the mouth of Black-

FORT SIMPSON.

water River we came to the commencement of the Rapid, and hauling the coble on the beach there, proceeded on foot to the "Hill," immediately below "the Rapid," where we encamped. While on our march we perceived that the bridge of ice was giving way, and the river directly fell some feet. Early next morning Bruce went back with the men for the coble, and brought it down by the time that Mr. Bell and I had prepared breakfast for the party. After concluding that meal, we crossed the river in her obliquely, among high walls of ice; and a mile or two below came to the barge which the crew had brought up so far two days previously. A short time sufficed to launch her, embark the tents kindly provided for us by Mr. M'Pherson, and descend to the mouth of Bear Lake River, where we passed the night. Next day we went on to Fort Norman, where I purposed to wait for Mr. M'Pherson, who had gone to Fort Good Hope to bring up the year's returns of furs from that post, and from the Peel and Porcupine Rivers.

On his arrival on the 14th I sent back the small barge, to bring down Narcisse, Brodie, and the stores, and embarking with Messrs. Bell and M'Pherson proceeded to Fort Simpson. The rest of the journey homewards, being by the same route as the outward voyage, need not be mentioned in detail.

0 viou Gree brin Lake end did the the 2 the 1 from encar had broug canoe at the to Ma affair since, that chang societ from leave ligatio forwar high Indian

1849

JULY.

JULY,

1849.

of the there, below on our ce was e feet. ie men e time for the crossed alls of barge o days ch her, us by uth of night. here I d gone eturns el and

ek the e, and ll and e rest route hed in METHY PORTAGE.

On the 25th we left Fort Simpson, having previously been joined by the men who wintered at Great Slave Lake, and also by the small barge, bringing Brodie and Narcisse Tremblè from Bear Lake. We were detained by drift ice at the west end of Great Slave Lake till the 6th of July, and did not reach Fort Resolution till the 11th. On the 19th we arrived at Fort Chepewyan, and on the 26th at Methy Portage, which we crossed on the 27th with all our baggage, on horses hired from the Indians. From L'Esperance, who was encamped with his brigade on Methy Lake, I had the pleasure of receiving English letters, brought up from Canada by the governor's light canoe, which leaves La Chine in May. Mr. Bell at the same time received instructions to return to Mackenzie's River, to conduct the Company's affairs there. This was unpleasant tidings to him, since, having spent the greater part of his life in that northern region, he had been soliciting a change, but the mortification was softened by the society of his two daughters who had been sent from Isle à la Crosse to meet him. In taking leave of this gentleman, I must express my obligations to him for his assiduous endeavours to forward the interests of the expedition, and my high sense of his excellent management of the Indians at Fort Confidence, to which we owed a

NORWAY HOUSE.

winter of abundance, and the excellent condition in which the store was left in spring. I had enjoyed much pleasure in his society, and parted from him with regret.

The remainder of the voyage down was performed in company with Mr. M'Pherson, who was going down to Canada on furlough with his family. At Norway House, where we arrived on the 13th of August, the men of my party who had been engaged there in 1847 were discharged, and the Europeans were sent down to York Factory to go to England in the Company's ship. During the time these men were under my command not a single act of disobedience occurred. Crews better fitted for heavy portage work, and for the ordinary duties of a winter's residence in the north, might doubtless have been selected in the country, but none that I could have depended upon with so much confidence in adverse circumstances. Dore and Cousins, the two younger seamen, who were extremely serviceable from their activity, intelligence, and willingness to perform any duty that was required, have gone again on the same errand to Beering's Straits.

From Norway House I proceeded to Canada in a brigade of three light canoes, manned by voyagers who were returning thither at the close of their engagements in the country. On the 14th of Sep-

temb Saut vesse a sta coe i throu about the st and M Aft Georg counts I wen mail s at Live an abs passed Wit miralty narrati afterwa approb Here

pedition dition the man absent Sir J VOL.

LAND AT LIVERPOOL.

145

ndition I had parted

as perho was family. e 13th d been ind the y to go ing the not a s better rdinary , might ry, but vith so Dore o were intellity that errand

> da in a yagers eir enof Sep

tember we came to Fort William, and on the 25th to Saut Ste. Marie. From thence we went in a steamvessel to the lower end of Lake Huron, and taking a stage coach there for Orillia, crossed Lake Simcoe in a steamboat. Then we travelled by coach through Young Street to Toronto, a distance of about forty miles, and there embarked in one of the steam-packets that ply daily between that port and Montreal.

After a few days passed at La Chine with Sir George Simpson in revising the outstanding accounts between the Company and the Expedition, I went to Boston, and embarking in the British mail steam-packet, crossed the Atlantic, and landed at Liverpool on the 6th of November, 1849, after an absence of nineteen months, twelve of them passed in incessant travelling.

Without delay I presented myself at the Admiralty, and, having laid before their Lordships a narrative of my proceedings, had the honour soon afterwards to receive a letter announcing their approbation of my conduct.

Here the journal of the transactions of the expedition ends, but a summary of the present condition of the search may not be unacceptable to the many who take an interest in the fate of our absent countrymen.

Sir James C. Ross, with the "Enterprise" and VOL. II.

146 PRESENT STATE OF THE SEARCH.

"Investigator," reached the three Islands of Baffin in lat. 74° N., on the 26th of July, 1848, but was not able to cross the "middle ice" till the 20th of August, on which day he attained open water in lat. 751° N., and long. 68° W. He then steered for Pond's Bay, and examined the coast carefully from thence to Possession Bay, in which he landed on the There he found a memorandum left by Sir 26th. Edward Parry in 1819, but no trace of Sir John On the 1st of September, the ships Franklin. arrived off Cape York, where a conspicuous landmark was erected. Sir James next examined Maxwell Bay, and some smaller indentations of the north coast of Barrow's Strait, but was prevented by a firm barrier of ice from approaching Cape Riley at the entrance of Wellington Channel. Neither could he get near Cape Rennell, because of compact, heavy ice extending from Wellington Channel to Leopold Island. Not being able to penetrate to the west, the ships were run into Port Leopold on the 11th, and on the following morning the main pack of ice closed in with the land and shut them in for the season. On the 12th of October the ships were hove into their winter quarters. During the winter many white foxes were taken in traps; and copper collars, on which were inscribed notices of the situation of the vessels, and of the depots of provision, having

beer liber In M'Cl coast and separ Cress retur In the w Beach Brown Lieute to the reach state conjur of 184 the Gu ceptior Fury I were n any pa certain passage house t with he

SIR JAMES C. ROSS.

f Baffin but was he 20th water in ered for lly from d on the ft by Sir Sir John he ships ous landexamined ations of was preoroaching Channel. , because Wellingeing able run into e followl in with On the nto their ny white llars, on nation of h, having

been secured round their necks, they were set at liberty again.

In May, 1849, Sir James Ross and Lieutenant M'Clintoch thoroughly explored on foot the west coast of North Somerset down to lat. 72° 38', N., and long. $95\frac{3}{4}^{\circ}$ W., where a very narrow isthmus separates Brentford Bay of the Western Sea, from Cresswell Bay of Prince Regent's Inlet. They returned to the ship on the 23rd of June.

In the mean time, Lieutenant Robinson examined the western side of Regent's Inlet down to Fury Beach, and several miles beyond it. Lieutenant Brown had crossed the inlet to Port Bowen, and Lieutenant Barnard had traversed Barrow's Strait to the vicinity of Cape Hurd, but was unable to reach Cape Riley on account of the hummocky state of the ice. By these excursions, taken in conjunction with Mr. Rae's expedition in the spring of 1847, the whole of Prince Regent's Inlet and the Gulf of Boothia was examined, with the exception of one hundred and sixty miles between Fury Beach and Lord Mayor's Bay; and as there were no indications of the ships having touched on any part of the coast so narrowly traced, it is certain that they had not attempted to find a passage in that direction. Sir James caused a house to be buil. at Port Leopold, and covered with housing cloths, in which he left provisions

г 2

148 PRESENT STATE OF THE SEARCH.

and fuel for twelve months, together with the Investigator's launch and steam-engine. He then proceeded to cut a way out for the ships through the ice, which was not effected until the 28th of August, 1849. On leaving the harbour he crossed over towards Wellington Channel, where he found the land-ice still fast and preventing his approach. While contending with the loose packs, and struggling to advance to the westward, a strong gale of wind on the 1st of September suddenly closed the ice around the ships, wherein they remained helplessly beset until the 25th, by which time they had drifted out of Lancaster Strait, and were off Pond's Bay. As the season was now far advanced, further search that year was thus frustrated by an accident, often experienced in the navigation of the Arctic Seas; and all harbours in that vicinity being closed for the winter, Sir James reluctantly gave the signal to bear up for England.

While Sir James C. Ross was still engaged in the ice on the west side of Baffin's Bay, Mr. James Saunders, Master and Commander of the "North Star," having been sent out with supplies in the spring of 1849, was working up on the east side, with imminent danger to his ship. Owing to the unusual quantity of ice in the bay that summer, and the frosts which glued the floes into one impenetrable mass, he was unable to cross of coming that the caught Septemil was prowhere the length effective 76° 33' northering known the coldest in casions rebelow zettion

hauled or ten mont Possessio ders next Jackson's from land by the he Pond's Ba on Wollas Meanwl quarter of the "Hera Wainwrig

"NORTH STAR."

cross over to Lancaster Sound, and his ship becoming involved in the ice about the same date that the "Enterprise" and "Investigator" were caught in the pack, drifted with it the whole of September, until on the last day of that month she was providentially driven into Wolstenholme Sound, where there being a pool of open water she was at length extricated. There the ship wintered in lat. 76° 33' N., long. 68° 56¹/₄ W., being the most northerly position in which any vessel has been known to have been laid up. February was the coldest month, and the thermometer on two occasions marked $63\frac{1}{2}^{\circ}$, and once $64\frac{1}{2}^{\circ}$, of Fahrenheit below zero.

On the 1st of August, 1850, the "North Star" was hauled out of the cove in which she had remained ten months, and on the 8th she had crossed over to Possession Bay, which was examined. Mr. Saunders next proceeded to Whaler Point, Port Bowen, Jackson's Inlet, and Port Neill; but being prevented from landing his provisions at any of these places by the heavy landfloes of old icc, he bore up for Pond's Bay, and succeeded in depositing his cargo on Wollaston Island.

Meanwhile the search was proceeding from the quarter of Beering's Straits. Captain Kellett in the "Herald," on July 25th, 1849, after examining Wainwright's Inlet, despatched Lieutenant Pullen

the then ough h of ossed ound oach. trugle of d the helpy had 'ond's irther acciof the cinity tantly

ed in Mr. of the pplies n the ship. e bay e floes ple to

150 PRESENT STATE OF THE SEARCH.

to the Mackenzie; and afterwards, in standing along the margin of the ice, discovered a group of high islands on the Asiatic coast in lat. 71° 20' N., long. 175° 16' W., with extensive and very high land to the north of them deeply seated in the ice.* Commander Moore, also, in the "Plover," made several attempts to penetrate to the eastward at this time, and not succeeding, returned first to Kotzebue Sound, and subsequently to Norton Sound, where he wintered.

Lieutenant (now Commander) Pullen, accompanied by Mr. Hooper, mate, and twelve men, performed the coasting voyage to the Mackenzie in two 27-foot whale-boats. He was convoyed past Point Barrow by the "Herald's" pinnace named the "Owen," and the Royal Thames Yacht-Club schooner, the "Nancy Dawson." The latter was owned and commanded by Mr. Shedden, a mate of the Royal Navy, who had come thus far with his small craft, solely at his own expense, to prosecute the search for the discovery ships, and who, though he was in the last stage of consumption, was not prevented by the languor of the disease, which carried him off two months afterwards, from giving most efficient aid to Lieutenant Pullen.

* Captain Smyth suggests that this land may be that reached by Sergeant Andreyer in 1762, which he reported to be inhabited by a people named Kraïhaï.

C 184 vigo ingl agai the son, The " In thro Strai and The the l the v sump Th Lance "Res T. Aı Omm the " the ty perien search Frank expedi furnisl

BEERING STRAITS.

anding coup of 20' N., cy high ne ice.* ' made vard at first to Norton

accomen, perenzie in ved past e named cht-Club tter was mate of with his prosecute , though was not e, which m giving

> hat reached to be in-

On Sir James C. Ross's return to England in 1849, the Admiralty resolved that a still more vigorous search should be organised, and accordingly the "Enterprise" and "Investigator" were again fitted out and despatched to Beering's Straits, the former under the command of Captain Collinson, C.B., and the latter of Commander McClure. These ships having separated in the Pacific, the "Investigator," which was the dullest sailer, through a fortunate choice of route, reached the Straits first, succeeded in passing Point Barrow, and was last seen on the 4th of August, 1850. The "Enterprise," having been unable to penetrate the barrier of ice, went to Hong Kong to refit in the winter, and is to make another attempt this summer of 1851.

The preparations for the search on the side of Lancaster Sound were on a large scale. The "Resolute" was commissioned by Captain Horatio T. Austen, and the "Assistance," Captain Erasmus Ommaney, was put under his orders, together with the "Pioneer" and "Intrepid," steam tenders to the two vessels. Captain William Penny, an experienced whale-fisher, was also engaged for the search, and placed in command of the "Lady Franklin" and "Sophia." In addition to these expeditions fitted out by the Admiralty, others furnished from private sources showed the interest

L 4

152 PRESENT STATE OF THE SEARCH.

that was widely and deeply felt in the cause. Captain Sir John Ross, notwithstanding his advanced years, sailed in the "Felix" schooner; and by the munificence of Mr. Henry Grinnell, a New York merchant, the United States sent forth the "Advance" and "Rescue" on the same humane quest, under the command of Lieutenant De Haven, U. S. N., and Mr. S. P. Griffin. Lady Franklin likewise, with that untiring energy and conjugal devotion which has marked her conduct throughout, despatched the "Prince Albert" under the orders of Commander Forsyth of the Royal Navy.

This squadron was assembled in Lancaster Sound in the month of August, 1850, at which time the "North Star" was also there, forming in all a fleet of ten vessels. On the return of the last-named ship and of the "Prince Albert," we received intelligence from Barrow's Straits as late as the 25th of August, 1850. By that time both sides of Lancaster Sound had been thoroughly searched as far as Cape Riley on the north side, and Port Leopold on the south, also Prince Regent's Inlet down to Port Neill and Fury Beach. On the above-mentioned date, Mr. Snow, of the "Prince Albert," went ashore at Point Riley to examine a flag-post, which had been erected by Captain Ommaney; and found a note from that officer, stating that he had landed with the officers of the "As-

sist had rem som deta onann Cap trac men colle note Fran De I on th signa away piece with chip (amina sion t ordina they a atmos snow proved been 1

BARROW STRAITS.

ause. s adand New h the mane aven, unklin njugal ghout, orders

caster which rming urn of lbert," uits as t time ughly side, gent's **)n** the Prince nine a h Omtating " Assistance" and "Intrepid" on the cape on the 23rd, had found traces of an encampment and collected the remains of materials, which evidently proved that some party belonging to Her Majesty's ships had been detained there. Traces of the same party were found on Beechey Island. The note concludes by the announcement that Captair. Ommaney proceeds to Capes Hotham and Walker, in search of further traces of Sir John Franklin's Expedition. No mention is made of the nature of the materials collected; but the tenor of Captain Ommaney's note indicates that he had no doubt of Captain Franklin having been off Cape Riley. Lieutenant De Haven, of the "Advance," landed on the cape on the morning of the 25th, and erected a second signal post, but seems to have carried nothing away. Mr. Snow gathered and brought off five pieces of beef, mutton, and pork bones, together with a bit of rope, a small rag of canvass, and a chip of wood cut by an axe. From a careful examination of the beef bones, I came to the conclusion that they had belonged to pieces of salt-beef ordinarily supplied to the Navy, and that probably they and the other bones had been exposed to the atmosphere and to friction in rivulets of melted snow for four or five summers. The rope was proved by the ropemaker who examined it to have been made at Chatham of Hungarian hemp, subse-

154 PRESENT STATE OF THE SEARCH.

quent to 1841. The fragment of canvass which seemed to have been part of a boat's swab, had the Queen's broad arrow painted on it; and the chip of wood was of ash, a tree which does not grow on the banks of any river that falls into the Arctic Sea. It had, however, been long exposed to the weather, and was likely to have been cut from a piece of drift-timber found lying on the spot, as the mark of the axe was recent compared to the surface of the wood, which might have been exposed to the weather for a century.* Mr. Snow counted five rings of stones with two or three slabs in the centre of each circle, which he took for fire-places, but on which there were no traces of smoke nor any remains of burnt wood. As tent-pegs could not be driven into the shingly beach, the stones had been evidently used in the erection of as many tents as there were circles, and the slabs in the centre were likely to have served as stands for magnetic instruments. Colonel Sabine remarked that four tents would be needed in using the instruments supplied to Sir John Franklin's expedition, and a fifth for the protection of the observers. If the ships were

* The grounds of these conclusions were fully stated in a report made to the Admiralty by Sir W. Edward Parry, myself, and other officers, which has been published with other parliamentary returns.

sto m tai tw sel day cor sen aft off and shi Str tere othe hay pora Fra Isla best Wit evid cair the Т ceed infer Had

TRACES OF THE "EREBUS" AND "TERROR." 155

stopped in that locality about the time of the monthly term-day, the officers would almost certainly make the term observations, which last for twenty-four hours, and in that case each ship would select a separate place of observation. The termday in August, 1845, was the 29th; and we may conclude, from the information which we at present possess, that on that day, or about a month after they were last seen, the discovery ships were off Cape Riley. It is ascertained that the bones and rope were not left by any party from other ships of the Royal Navy that have visited Barrow's Straits; and had the "Erebus" and "Terror" wintered there, a cairn, with memoranda, and many other evidences of that fact, would undoubtedly have been found. The ships must have been temporarily arrested by a barrier of ice; and Sir John Franklin, having good anchorage between Beechey Island and Cape Riley, turned the delay to the best account by making the term observations. With the prospect of soon passing the Straits, he evidently had not thought it necessary to erect a cairn, or signal-post, at the threshold, as it were, of the enterprise.

These first traces of the Expedition are exceedingly interesting, and they lead directly to the inference that it pursued its course to Cape Walker. Had Sir John, finding the Strait barred, gone up

which d the chip grow Arctic o the rom a ot, as o the n ex-Snow three e took traces l. As hingly in the e cir. ely to nents. uld be to Sir r the were

> d in a myself, parlia-

> > -

156 PRESENT STATE OF THE SEARCH.

Wellington Inlet, he would undoubtedly have left a memorandum at Cape Riley, assigning reasons for departing from his instructions.

g

in

of

to dis

at

lad

on to

the

und

is e vill

is s

the

that

tha: Sou

boa

On the 25th of August, 1850, the "Assistance" and "Intrepid" were well over towards Cornwallis Island, and a little to the north of Cape Hotham. Captain Penny with his consort were standing up the west side of Wellington Channel, Sir John Ross was making for Cape Hotham, the "Rescue" was near the head of Wellington Sound on the east side, land being visible from Cape Riley, crossing the Sound just beyond her, and the "Advance" was lying under Beechey Island, having grounded on a shoal; but Lieutenant De Haven, expecting to get his vessel off without difficulty, declined the assistance from the "Albert" which was offered. Captain Austin, with the "Resolute" and "Pioneer," was at that time examining the south side of Barrow's Straits. He had visited Possession Bay on the 18th, and then intended to look into Pond's Bay. He has not been seen since, and it is probable that he passed the "North Star" and "Albert" without being descried, in the thick weather that prevailed when they were coming out of Lancaster Sound.

Such was the state of the search in August, 1850. As Sir John Ross intended to return in 1851, after

EXPECTED RETURN OF SIR JOHN ROSS. 157

landing his stores on Melville Island*, we may expect that he at least will bring further intelligence in October or November next.

* Sir John Ross took with him four carrier pigeons belonging to a lady residing in Ayrshire, intending to liberate two of them when the state of the ice rendered it necessary for him to lay his vessel up for the winter, and the other two when he discovered Sir John Franklin. A pigeon made its appearance at the dovecot in Ayrshire on the 13th of October, which the lady recognised by marks and circumstances that left no doubt on her mind of its being one of the younger pair presented by her to Sir John. It carried no billet; but there were indications, in the loss of feathers on the breast, of one having been torn from under its wing. Though it is known that the speed of pigeons is equal to one hundred miles an hour, the distance from Melville Island to Ayrshire being in a direct line about 2,400 miles, is so great, that evidence of the bird having been sent off about the 10th of October must be had, before that we can well believe that no mistake was made in the identification of the individual that came to the dovecot. Sir John's letters from Lancaster Sound mention that when he wrote he had the pigeons on board.

e left is for

nce" vallis ham. g up Ross ' was side, g the ' was ed on ng to d the fered. 'Pioside ssion into nd it ' and thick ming 850. after



APPENDIX.



APPENDIX.

No. I.

PHYSICAL GEOGRAPHY.

CHAPTER I.

- GENERAL VIEW.—Rocky Mountains. Their Length. Their Height. — Glaciers. — Parallelism to the Pacific Coast. — Continental Slopes. — Russian America.—Eastern Slope.— Prairies.—Mississippi Valley. — Its Slope — Fundamental Rocks of the Basin. — Silurian Strata. — Carboniferous Series.—Tertiary Beds.—Lignite Formations.—Series of Lake Basins. — Transverse Valleys. — Intermediate Belt of primitive, hypogenous, or metamorphic Rocks. — Its Rivers mere Chains of Lakes. — Its Breadth.— Altitude. — Sources of three great River Systems. — Great Fish River. — The Yukon or Kwichpack. — Basins of Excavation. — Glacial Action. — Active Volcanoes.
- VALLEY OF THE ST. LAWBENCE. Altitudes of the Lakes above the Sea. — Lake Superior. — Lake Michigan. — Lake Huron. — Lake Erie. — Lake Ontario. — Lake Champlain. — Northern Brim of the St. Lawrence Basin. — Its geological Structure. — North Shore of Lake Superior. — Structure of the Country at the Sources of the Mississippi.
- WINIPEG OR SASKATCHEWAN VALLEY.— Height of Lake Winipeg.— Sea River.—Katchewan River.—Thousand Lakes.—Portages.— River Winipeg.—Red River.—Saskatchewan River.

MISSINIPI VALLEY. -- Its Lakes. -- Frog Portage.

- MACKENZIE RIVER VALLEY. Methy Portage. Athabasca, Elk, or Red-deer River. — Lesser Slave Lake. — Peace River. — Slave River. — River of the Mountains. — Noh'hanne Bute. — Great Bear Lake.
- YUKON VALLEY. Yukon or Kwichpack. Volcanic Chain of Alaska. — Coal. — Fossil Boncs.

In drawing up the following Appendix, my object has been to record facts in the light in which they appeared

М

VOL. II.

PHYSICAL GEOGRAPHY.

vie

able

flow

for

who

doni

Cop

of A

to is

trave

the s

cross

The

to th

explo

from

parall

from i

rapid

noted

Conti

the lo

corner

westwa

charact

volcano

at pres

lying e

hundred

consist (

The

Up

162

5

to me. When treating of districts which I did not visit, I have borrowed from every work concerning them to which I had access. The most important sources of information are generally named or expressly alluded to; but I thought that it would give too much formality to so slight a sketch were I to parade every authority for the statements it contains. Where practised geologists have examined the country, their report has been chosen in preference to my own observations; and this is the case on the route of the expedition up to the 49th parallel. Beyond Lake Winipeg no geologist has yet penetrated, and the descriptions of the rocks occurring within the space of twenty degrees of latitude that lie to the north of that sheet of water are, with all their imperfections, entirely my own. It would be true economy in the Imperial Government, or in the Hudson's Bay Company, who are the virtual sovereigns of the vast territory which spreads northwards from Lake Superior, to ascertain without delay the mineral treasures it contains. I have little doubt of many of the accessible districts abounding in metallic wealth of far greater value than all the returns which the fur trade can ever yield.

The Rocky Mountain chain, which is the northern prolongation of the Andes, has a general course of north 26° W. for 2,700 geographical miles, from the 30th parallel of latitude up to the shores of the Arctic Sea. Its higher peaks rise from 12,000 to 15,000 feet above the ocean, and enter the region of perpetual snow; but the northern part of the chain, which touches on the Mackenzie, is so much lower, that even its summits * are denuded during the short summer of that district, and perennial patches of snow exist there only in shady cre-

* Supposed to be at least 3,000 feet high, in the 62nd parallel.
ROCKY MOUNTAINS.

ot visit, hem to s of inded to; ality to y for the sts have nosen in the case parallel. netrated, ithin the north of , entirely erial Gowho are h spreads without ave little inding in e returns

northern of north the 30th retic Sea. above the but the the Macbits * are rict, and hady cre-

arallel.

vices which have a northern aspect. I have not been able to discover, after many enquiries, that glaciers which flow through mountain gorges into the lower country are formed in any part of North America, though travellers who have crossed to California, Oregon, and New Caledonia, speak of hills clothed with perpetual snow; and the Copper River, which joins the sea opposite to the peninsula of Alaska, is said by Bäer, on the authority of Klimowskij, to issue from a solid mass of ice. Several passes which traverse the chain do not rise more than 6,000 feet above the sea level, and being free from snow in summer may be crossed in that season by pack-horses and even by waggons. The more northern of these passes have long been known to the fur-traders; the southern ones have lately been explored and used by the multitudes who have hurried from the United States to California in search of gold.

Up to the 60th degree of latitude the chain runs nearly parallel to the coast of t^1 Pacific, and not far distant from it; the descent to the level of the sea is consequently rapid on the west, — a configuration which M. Guyot* has* noted as peculiar to the New World, while in the Old Continent the short slopes are turned to the south, and the long ones towards the north. A large triangular corner, which belongs to the empire of Russia, and extends westward to Beering's Straits, has a different physical character, in the existence of a transverse series of active volcanocs, as we shall hereafter have occasion to notice; at present my remarks will be confined to the continent lying eastward of the mountains.

The width of the chain is stated at from forty tc one hundred miles, and the central parts and peaks are said to consist of granite and other igneous rocks.

> * Phys. Geogr., р. 50. м 2

The eastern slope towards the Atlantic commences by a belt, formed mostly of sandstone, 150 miles in width, which rests on the shoulder of the chain, with an inclination of about 37 feet in the mile, in its descent from 8,000 feet above the sea, to 2,500. The more gradual slope of the great prairies, beginning at the last-named elevation, has a breadth of 700 or 800 miles, and retains in its descent the prairie character of a treeless, sandy, and moderately undulated, or, as it is locally named, "rolling" plain.

Most of the streams which cross the prairie flow through deep furrows, sunk abruptly below the general level; nevertheless the Mississippi, Missouri, and some of their larger tributaries have wider valleys, skirted by successive terraces and alluvial deposits. On the banks of all the rivers there are belts of woodland, and clumps of trees that encroach on the prairie, intercepting grassy lawns, and producing remarkably fine park scenery, which is often enlivened by small lakes. In the interior of the prairie, however, water is scarce, and there is such a total want of wood, that for [•]days together the traveller can find no other fuel than the dung of the bison. Near the mountains the soil is coarsely sandy, strewn with boulders, and sterile; farther eastward the sand is finer, and the boulders disappear, but they recur in numbers on the lower border of the prairie: they are also scarce or wholly absent over very extensive tracts of the rich alluvial deposits of the valley of the Mississippi, south of the Ohio.

The Mississippi drains the entire space between the Rocky Mountains and the Alleghanies or Apalachian chain, embracing thirty degrees of longitude. The whole of this vast water-shed may be considered as one valley, whose bottom, indicated by the channel of the river, has a southerly course, inclining slightly to the eastward. The length of the river, from its source in Itasca Lake, at an

alti of stat rou line whi of t para valle mile of t inche A geogr sumn 2,500 valley inche might and c drain The 1 from than t ordina consid

* Th in Tenn history the chan an area and the one mil

MISSISSIPPI VALLEY.

165

width, lination 000 feet of the n, has a cent the tely un-

through ; neverir larger terraces ers there roach on icing reened by er, water that for than the coarsely eastward but they ie: they ve tracts ssissippi,

veen the palachian he whole e valley, er, has a d. The ie, at an altitude of 1,490 feet above the sea, to Balize, on the Gulf of Mexico, has been estimated by Schoolcraft at 3,160 statute miles, following its windings, and may be stated in round numbers at 2,400 geographical miles in a straight line. The moderate hills and eminences of the country in which the river has its origin do not rise above the plane of the general eastern slope of the continent in the same parallel; and the general longitudinal descent of the great valley from Lake Itasca is at the rate of 10 inches in the mile for the upper half of the river, and for the other half of the way, or from St. Louis downwards, of only two inches and a half per mile.

As St. Louis is 500 feet above the sea, and about 600 geographical miles distant transversely from the western summit of the prairie slope (which has been taken at 2,500 feet above the sea), the lateral descent of the valley to the channel of the river in that parallel is 40 inches in the mile; but the sinuosities of the Missouri, a mightier stream than the one in which it loses its name, and of the other grand affluents of the Mississippi that drain the prairies, give a gentler inclination to their beds. The Illinois, Ohio, and the minor streams which come in from the other side of the valley, flow at a lower level than the western feeders; their upper branches being subordinate to the general slope, which has there descended considerably.*

* The following facts, ascertained at the Navy Yard of Memphis, in Tennessee, by R. A. Marr, Esq., are interesting points in the history of the Mississippi. The quantity of water passing through the channel of the river at that place in 1849 was sufficient to cover an area of 100,000 square miles to the depth of seven feet and a half, and the quantity of silt it carried down would make a bed of earth one mile square and seventy-six feet deep. The current in the

166

From the Apalachian chain to the Rocky Mountains the fundamental rocks would appear to be silurian* overlaid in large tracts by the old red sandstone and the car-

central area of the river generally exceeded three miles an hour, and was less towards the sides.—*Proceed. Amer. Assoc.*, p. 340.

* As the silurian rocks are most extensively developed in North America, and are of constant recurrence throughout the route of the Expedition, I shall introduce here, for reference, a tabular enumeration in the order of their superposition, as ascertained by the United States geologists, beginning with the lowest.

- A. CHAMPLAIN DIVISION; supposed to underlie three-fourths of the territory of the United States, and to occupy the surface of one half.
 - 1. Potsdam sandstone; a quartzose rock, generally grey, often striped, sometimes partially or wholly red: is supposed to be the lowest sedimentary fossiliferous rock. The Taconic rocks of Dr. Emmons, consisting of lamellar white limestone, with specular iron ore, are supposed to be the earliest deposits of this period, modified by metamorphic agency. Footmarks of reptiles have been found by Mr. Hunt in this sandstone.
 - 2. Calciferous sandrock; a deposit of calcareous and earthy matter variously mixed.
 - 3. Black river or Chazy and Birds-eye limestone; a brittle limestone, having a smooth, flat-conchoidal fracture. 'This limestone, characterised by its peculiar fossils, is of frequent recurrence northwards, up to the islands of the Arctic Sea.
 - 4. Trenton limestone; a dark-coloured limestone, interlaminated with dark shaly matter. Often metalliferous.
 - 5. Utica slate; dark-coloured carbonaceous slates, which readily disintegrate.
 - 6. Hudson River group, Loraine shales, Frankfort slate, and Rubblestone; mostly dark slates and shales, and grey thick-bedded grit stones.
- B. ONTARIO DIVISION, includes a series of limestones, shales, and sandstones, which pass insensibly into each other.
 - 7. Grey sandstone of Oswego.
 - 8. Medina sandstone; red or slightly variegated, and of every degree of coherence. Originates many brine springs.
 - 9. Oneida conglomerate; a variable intermixture of sand and quartz pebbles.
 - 10. Clinton group; consisting of deposits of various characters,

bor of t the

C. I

13

15.

16.

17.

a

a

ť

le 19,

t

is

tı

of

in cl

VC

D. ER

20.

18.

SUJURIAN STRATA.

boniferous series. The sandstone, which rises on the flanks of the Rocky Mountains to the height of 8,000 feet above the sea, is referred by some geologists of the United States

such as thin shaly sandstones, shales, conglomerates, thin-bedded impure limestones, with iron ores.

- 11. Niagara group; consisting of dark-bluish shale and dark limestone, and taking its name from the cataract, where a section of both its members is exposed.
- C. HELDERBERG DIVISION.
 - 12. Oxondago salt group, is an immense development of argillaceous shales and marks with shaly limestones, veins and beds of gypsum, giving origin to copious and very rich salt springs. This formation re-appears near Slave River, on the 60th parallel of latitude, and also, I believe, on the shores of the Arctic Sea.
 - 13. Water-line group, consists generally of two layers of drabcoloured water-linestone, with an intervening layer of blue lime-rock.
 - 14. Pentamerus Emestone; named from its characteristic fossil, and rarely a pure limestone, being more or less mixed with black shale.
 - 15. Delthyris or Catshill shaly limestone; composed of beds of subcrystalline grey limestone, slaty limestone, and slaty argillaceosiliceous limestone.
 - 16. Oriskany sandstone; a whitish sandstone, composed of sand derived from granitic rocks or mica schist.
 - 17. Caudagalli grit; named from the feathery forms in which it abounds. It is a drab-coloured or brownish calcareous and argillaceous sandstone.
 - 18. Shoharie grit; a fine-grained calcareous sandstone, from which the calcareous matter may be washed away by long exposure, leaving the siliceous skeleton of the rock.
 - 19. Onondago limestone; recognisable by its crystalline structure, toughness, and numerous organic remains.
 - 20. Corniferous limestone; a fine-grained compact limestone, which is very durable, and produces cascades where the smaller streams traverse it. It is the uppermost of the important limestone beds of the New York or silurian system, being succeeded by shales in which the limestone fossils give place to others of a different character.
- D. ERIE DIVISION, referred by most English geologists to the Devonian or to the carboniferous series.

intains * overhc car-

our, and

n North te of the enumee United

hs of the ce of one

ey, often to be the *rocks* of specular is period, tiles have

hy matter

imestone, one, chace north-

aminated

h readily

d Rubbledded grit

ales, and

of every

ad quartz

aracters,

to the triassic system; but its exact geological position is not yet satisfactorily determined. Cretaceous beds, known by their organic remains (but not containing white chalk with flints), occur extensively along the Missouri, and spread widely on both sides of the Mississippi, below the influx of the Ohio. They exist also near the Rocky Mountains, high above the sea, in the vicinity of the sources of the Arkansas, Platte, and Gila*; while towards

- 21. Marcellus shales; black, slaty, bituminous shales, containing septaria, with occasional thin bands of limestone. Similar in lithological characters to Genesee slate.
- 22. Hamilton group; an immense deposit of dull-olive calcarcous shales, which change to light ash-grey in weathering. It contains septaria.
- 23. Tully limestone; usually thick-bedded, blue, or nearly black, limestone, often divided by seams into irregular fragments.
- 24. Genesee slate; is a great mass of argillaceous, black, fissile slate, which rapidly exfoliates and falls down. Fluid bitumen is of common occurrence. Either this deposit or No. 21. exists on the north side of Methy Portage, on the Elk or Athabasca river.
- 25. Portage group; a vast deposit of shale, flag-stones, and thickbedded sandstone, rising with a slope or abruptly from the shales on which it reposes.

t

N

c

S

0

b

C

of

S

st

aı

ac

sti

88

TI

A

D

ne

18

26. Chemung or Ithaca group; a highly fossiliferous series of shales and thin-bedded sandstones.

To the last succeed the OLD RED SANDSTONE and the COAL MEASURES.

The lower part of the Champlain division has been considered as the equivalent of the Cambrian system, the Utica slate being parallel to the Landeilo flags; the Hudson River group and Ontario division, up to the Niagara group, is thought to be equivalent to the Caradoc sandstone; the Niagara group and the whole Helderberg division is supposed to be co-equal with the Wenlock rocks; and the Erie division as equivalent to the Upper and Lower Ludlow rocks, including the Devonian system of Phillips.—Vide Hall, Geol. Report of New York, p. 518.

* "The cretaceous formations terminate in the Atlantic regions of New York, before they have reached the city of New York, so that their limit scarcely touches the 40th degree of latitude, or 16° lower than in Europe. In Kentucky and Tennessee it remains below 37°,

TERTIARY BEDS.

position is s, known ite chalk ouri, and elow the e Rocky y of the e towards

containing Similar in

caleareous g. It con-

early black, nents. black, fissile bitumen is 21. exists on basea river. and thickn the shales

ies of shales

COAL MEA-

nsidered as ing parallel rio division, he *Caradoc* ; division is t the Erie *rocks*, in-*L. Report of*

> regions of rk, so that 16° lower below 37°,

the mouth of the Mississippi there are very extensive recent tertiary and alluvial deposits, which, skirting the Gulf of Mexico, run into Texas on the one side, and along the Florida coast on the other.

Tertiary coal, containing dicotyledonous leaves, exists in the Raton Pass, between the sources of the Red River and the Arkansas, at an elevation of 4,600 feet. Coal of the same description, associated with similar leaves, occurs on the Mackenzie in latitude 65° N.; and at various intermediate parts on the flanks of the Rocky Mountains

but it is very different far up the Missouri ; this great river flows uninterruptedly from the foot of the Rocky Mountains for 1,400 miles through strata of chalk, at least as far as the Sioux river. This is the result of the researches of the Prince of Neuwied, and of the reports of the celebrated astronomer, Nicollet. In these western parts of America the chalk rises to 50° of latitude. There, also, it nows a continuous extension greater than that of any other formation on the globe. Captain Fremant saw chelk strata and fields covered with Inoceramus crispii, on the River Plane; Lieutenant Libert sound them on the Arkansas, and Dr. Winteren also beyond the Rio del Norte, nea. Munterey and Larendo. The Rocky Monotains, and heir continuation beyond Santa Fe, have entirely cut off this erelace bus sea. No trace of chalk was discovered either by Contain Fremont on the Commisia River or on the Humboldt, in that - more fully great basin which dips to the Pacific; or yet by the observant Captains Cooke and Johnstone along the River Gila, in Sonora, or California." "The whole of this vastly externation consists only of the upper beds. After very areful and accurate investigation, Sir Charles Lyell decided, that in the whole of Light America, chalk strata, from the Maestricht beds down to the said, done occurred; and Mr. Ferdinand Römer, as the result of his highly valuable and accurate researches in Texas goes the length of estimating all the strata in that region, already o far removed from the Mantic coast, as entirely of the upper division, and not even touching on the gault. This peculiarity is, however, singularly enough, limited to North America alone. Even in Mexico deeper chalk beds occur; and Darwin saw cretaceous shells in abundance 2,000 feet above the sea, near Port Famine, in 53° south latitude."-Silliman's Journal, Sept. 1850.

beds of lignite are known to exist, which are probably also of the tertiary era.*

In the whole width of the Mississippi basin, from the falls of St. Anthony downwards, no primitive (or hypogenous) rocks appear, except in the low ridges of the Ozark Mountains, which have a hilly prominence, owing to the excavation of the valley; but their summits scarcely rise above the plane of the general slope, supposing that it were extended with an even descent to the Atlantic. They range from Red River to St. Louis, parallel to the Alleghanies, and consequently make an angle with the Rocky Mountain chain. Coal measures crop out on their flanks.

Of the forty degrees of latitude which intervene between the Gulf of Mexico and the Arctic Sea, the valley of the Mississippi occupies, as we have seen, about one half; and the whole drainage of that portion west of the Alleghanies is accumulated in one great channel, which is directed southwards and a little eastwards. From the head of Lake Superior northwards, there is a series of great transverse excavations, occurring in succession, on to Great Bear Lake, which lies under the Arctic circle; and it is remarkable that nearly all the lake basins † of these valleys commence in the silurian strata, and are continued into or entirely across a belt of primitive, or hypogenous, and metamorphic rocks ‡, which extends from Lake Superior

* Vide Journ. p. 194.

† This grand series of lakes forms the line of canoe navigation from Canada northwards; and the fact of its position in the fracture between limestone and granite, was perceived and recorded by Sir Alexander Mackenzie.

[†] My imperfect acquaintance with the science of geology renders me incompetent fully to appreciate the worth of the several systems that profess to explain the mode in which the beds forming the crust of the earth have been formed; neither have I exclusively adopted

to lim mo of slig the par tern of nort an syst acco recti T posit inter nowl Its h assoc slope

any of the ter nation theorie

vated

shape

rated

these occuj

of wa

excee

Th

PRIMITIVE ROCKS.

to the shores and islands of the Arctic Sea. The western limit or strike of this formation, which I have traced for more than 1,400 geographical miles, has a general course of north 30° W., its rhumb line, consequently, inclines slightly towards the axis of the Rocky Mountains, and the intercepted space grows narrower towards the north.

On the other hand, the Apalachian chain, running parallel to the Atlantic coast for a thousand miles up to its termination in the Shickshock and Notre Dame Mountains of the promontory of Gaspé, and having a direction of north 46° E., diverges from the Rocky Mountain axis at an angle of 72°. These are the three chief pyrogenous systems of North America, the Ozark Hills being of lesser account, and coinciding, as has been said above, in direction with the Alleghanies.

The middle belt of primitive rocks may, both from its position and diagonal direction, be distinguished as the *intermediate primitive or hypogenous* formation. Its altitude nowhere entitles it to the appellation of a mountain chain. Its hypogenous rocks, which are chiefly granite and gneiss associated with trap, scarcely rise above the mean eastern slope, and do not present acute peaks or continuous elevated ridges. They exhibit generally rounded or domeshaped summits, or form oblong eminences, which are separated from one another by narrow inclined valleys. Most of these valleys, and the larger ones without exception, are occupied by lakes, which are often deep; and the proportion of water in the district is very great, probably considerably exceeding that of the dry land.

The rivers that traverse the intermediate primitive belt

bably

m the hypoof the owing arcely g that lantic. to the th the n their

etween of the f; and ghanics irected f Lake nsverse : Bear is revalleys ed into is, and aperior

> vigation Fracture by Sir

renders systems e crust dopted

any of the current opinions: I would therefore be understood to use the terms "primitive," "hypogenous," and "metamorphic," as designations of the rocks so called by geologists, and not as exponents of theories.

(and for ten degrees of latitude all the rivers that originate high on the prairie slope do so) form, on entering it, lakelike expansions, which are studded with rocky islets, and send long winding arms into all the neighbouring valleys. These dilatations have little or no current, but they are connected with each other by one or more straits, in which the stream is turbulent and rapid, and the overfall frequently great enough to produce a cascade. The tortuous arms of such expansions often wind for many miles through the country; and the Indian, by making short portages from one string of lakes to another, may travel with his canoe in every direction, as far as the formation extends. Sometimes a river forks in this rocky district, and its branches running far apart, just as they would in an alluvial delta, unite again, intercepting a considerable tract of country of the prevailing character, that is, having a predominance of water surface. Examples of this peculiarity occur in the River Winipeg, which conveys the waters of the Lake of the Woods to Lake Winipeg; also in the discharge of the latter sheet of water by Nelson River and Play-Green Lake; as well as in some of the expansions of the Missinipi or Churchill River. Instances also of lakes having more than one outlet are not rare in this formation; and now and then, though comparatively very seldom, these outlets lead to distinct river systems.

On the east side of Lake Winipeg the width of the primitive belt is about two hundred geographical miles; and from the summit of an eminence which rises only a few hundred feet above the general level, but yet is sufficiently conspicuous to have obtained the distinctive appellation of "the Hill," thirty-six lakes may be counted. In many localities, where the knolls are denuded of soil, the surfaces of the rocks are evenly ground down, and are sometimes smooth, polished, and striated.

on and the bra Sco we sur heig con equ cour the issu is b hilly des t does rise sum thou river nate Be whic state but * S

Missis Long, reckor from observ tween 1,460

PRIMITIVE ROCKS.

ginate

lake-

s, and

alleys.

ey are

its, in

verfall

e tor-

miles

short

travel mation

istrict, ould in

lerable

having 3 pecu-

ys the

; also

Nelson

of the

stances

rare in

ntively

miles;

only a

suffi-

e ap-

unted.

f soil,

hd are

ms. of the

By a rough measurement, the centre of this formation on the 53rd parallel of latitude is found to be between 700 and 800 geographical miles from the Rocky Mountains; the Great Canadian Lake district is of equal width; and Labrador on to Newfoundland and the eastern shores of Nova Scotia occupy a similar space in the map. Now assuming, as we have done, and as the observations of the topographical surveyors of the United States entitle us to do, that the height at which the gentle eastern slope of the continent commences is 2,500 feet, and supposing the descent to be equable, we should have an altitude above the sea in the country from whence the sources of the Mississippi proper, the St. Lawrence, and the Red River of Lake Winipeg issue, of 1,800 feet. The actual elevation of that district is between 1,400 and 1,500 feet*, and the only marked hilly eminence in the district, which is named the hauteur des terres, and is said to consist of drift-sand and boulders, does not appear, from the descriptions we have of it, to rise more than 300 feet beyond the general level. The summits, therefore, of this tract of land, distinguished though it be by shedding its waters into three separate river systems and as many different seas, are also subordinate to the general eastern slope of the continent.

Before naming more particularly the transverse basins which cross the intermediate belt of primitive rocks, I may state that the Mackenzie, inferior indeed to the Mississippi, but yet a river of the first class, running in an opposite

* Schoolcraft estimates the height of Itasea Lake, from which the Mississippi issues, at 1,490 feet above the Gulf of Mexico. Major Long, who ascended the St. Peter's, a head branch of the Mississippi, reckons the altitude of the short portage, which separates its sources from those of Red River, at 1,400 feet; and my own barometrical observations and estimates place the summit of the water-route between Lakes Superior and Winipeg, traversed by the Expedition, at 1,460 feet above the tide level of the St. Lawrence.

174

direction, drains seventeen degrees of latitude into the Arctic Sea, taking its course through a valley which differs in its character from that of the Mississippi, as the details of progress of the expedition through it have already shown. In this place it will be sufficient to recall to mind that from Methy Portage (*Portage la Loche*) to the sea, a distance of 1,400 geographical miles, the fall is about 900 feet, the successive portions of the river being designated the Washacummow, Elk or Athabasca, Slave, and Mackenzie Rivers.

Two other rivers of magnitude cross the Arctic circle, viz., Back's Great Fish River, which, originating near Great Slave Lake at an altitude of 150 feet above its surface, runs cast-north-east into the Arctic Sea, draining the north-eastern corner of the continent; and the Yukon, which, rising to the westward of the Rocky Mountains, not far from the union of the Francis and Lewis, which form the Pelly, flows first to the north, and after receiving a large tributary named the Porcupine, to the westward, falling into Beering's Sea, where it is known to the Russians by the name of the Kwichpack.

A glance at the map will show, that on the eastern side of the continent the water basins generally maintain the north-casterly inclination of the Alleghanies, while further to the westward the basins of the two great rivers assume a parallelism to the Rocky Mountains; and that the influence of the intermediate hypogenous formation has been of a different character, the rivers winding their way across it, sometimes with a southerly, sometimes with a northerly inclination, seemingly indicating the obstruction offered by the harder rocks to the agent by which the river channels were excavated. On emerging from the belt, the lower parts of the rivers generally incline towards the north-cast,

* A St. La to wh or less haps n sive d rocks, of the

SOURCES OF THREE GREAT RIVER SYSTEMS. 175

with a considerable degree of parallelism to each other before they fall into Hudson's Bay.*

The peculiar configuration of the continent which I have endeavoured to sketch must be duly considered by any one who endeavours to detect the agency by which the river valleys and lake basins were excavated. It is not, however, my purpose to enter upon the discussion of this question, or to speak of the partial and often repeated elevations and depressions by which the lacustrine and fluviatile terraces have been accounted for ; nor would a summary of this kind admit of the necessary elucidations. I shall merely say that, adopting the opinion of the United States' geologists, that they are basins of excavation, I consider them all to be of the same epoch, and that the currents or waves of translation, if such they were, must have had an easterly direction in the middle latitudes, and gained strength as they rolled towards the Atlantic, when they swept away wholly or partially the fossiliferous deposits that once covered the primitive rocks of Hudson's Bay, Canada, and the castern parts of the United States; the former extent of the newer rocks being indicated by the patches which remain. By a singular coincidence of a political with a natural limit, the northern boundary of the United States, or the 49th parallel, marks the line on the great prairie slope, where the current took a southerly direction, to excavate the wide and magnificent valley of the Mississippi. A similar diversion of the excavating

* A study of the map will show, that the lake basins north of the St. Lawrence have generally their long axes across the river courses to which they respectively belong, and that many assume a greater or less degree of parallelism to the intermediate primitive belt. Perhaps movements of elevation or depression had occasioned an extensive disruption of strata along the western border of the hypogene rocks, previous to the removal of the silurian beds on the excavation of the lake valleys.

to the which as the have ent to *Loche*) ne fall being Slave,

circle, g near ove its caining Zukon, ntains, which ceiving tward, to the

n side in the iurther issume he ins been across therly red by innels lower i-cast,







force northwards would produce the basin of the Maekenzie, commencing on the 53rd parallel, but in a district narrowed and disturbed by the approximation of the intermediate primitive rocks to the Rocky Mountains.

Supposing the continent to have retained its present form since the era of these excavations, it seems searcely possible to reconcile the existence of extensive glacial action with any modification of climate; yet the smoothened surfaces, streaks, and furrows referred to that action, whether in the form of glaciers or of drift ice, are of no rare occurrence, wherever durable rocks show themselves, between the St. Lawrence and the Arctic Sea.

In connection with the excavations of the North American continent, the fact may be mentioned of the great indentations of the coast line, including Hudson's Bay, the Gulfs of St. Lawrence and Mexico, and the Caribbean Sea being on the east side; while breaks of the west shore, to the south of the peninsula of Alaska, are comparatively small, and both coasts of South America are nearly entire.

The geologists of the New York Survey consider that the present continent of North America was constructed from the debris of land lying more to the eastward. Mr. Hall, speaking of the strata exposed between the Hudson and the Mississippi, states that they contain organisms which must have lived in the bed of the ocean, and that the chief source of the sedimentary deposits lay to the east and south-east. To the westward, the sedimentary rocks are of a finer grain, and at the same time diminish in quantity, while the carbonate of lime increases, indicating, in conjunction with the contained fossils, the bottom of an ocean of greater depth and more quiet condition. The erctaceous and tertiary deposits of the western prairies show, according to the geologist just quoted, that the eastern part of the continent was first elevated, and that

the the of pos the mov the coas chai T othe and 75°, theor earth in th of m exist serva cases,

The gr

a leng

contain the wo prairie the Mi on the aided 1 from th * Spel

* Spel French a VOL.

VALLEY OF THE ST. LAWRENCE. 177

Macstrict e in-

esent rcely ection surether rare s, be-

Ame · great Bay, bbean shore, atively entire. hat the l from Hall, n and which e chief t and rocks ish in ating, of an The rairics t the l that the older rocks in the west were subsequently overlaid by the new deposits. However that may be, the occurrence of the chalk fossils and tertiary deposits in their present position and altitude clearly indicates that the elevation of the Rocky Mountain chain was one of the latest of the great movements that have occurred in this continent. It is to the westward of these mountains only, along the Pacific coast, and in the peninsula of Alaska*, and the Aleutian chain of islands, that recent volcanic action can be traced.

The existence of coal measures, containing ferns and other plants of a tropical character, in Jameson's Land and Melville Island, in the high latitudes of 71° and 75°, is a curious fact, to be accounted for by those who theorise on the ancient condition of the surface of the earth; and the vast accumulations formed at a later epoch in the Siberian Sea and Kotzebue's Sound, of fossil bones of mammoths, rhinoceri, and other animals which do not exist in arctic regions at the present day, and the preservation to this date of some of their undecomposed carcases, are equally interesting facts, which need explanation.

VALLEY OF THE ST. LAWRENCE.

The first in order of the great transverse excavations, and the grandest, is the basin of the St. Lawrence, which has a length equal to the whole course of the Mackenzie, and contains by far the greatest accumulation of fresh water in the world. It has no connection with the drainage of the prairie slopes in the same parallels, which is performed by the Missouri and its numerous affluents on one side, and on the other by the Saskatchewan and its tributaries, aided by the upper feeders of the Mackenzie. It differs from the three great lacustrine basins which succeed it, in

* Spelt thus, and also *Alashka* in Cook's Third Voyage. The French and Russian authors write the word *Aliaschka*.

VOL. II.

N

178

Ś

its head lying within or to the eastward of the termination or elbow of the "intermediate primitive rocks." The position of this head, or as it is well named from its elevation above the other members of the basin, Superior Lake, is midway between the Gulf of Mexico and the Arctic Sea, and its water-surface is 641 feet above the level of the tide.* The other great lakes descend successively in the following order of their heights above the sea: Lakes Huron and Michigan, 600 feet; Erie, 565 feet; Ontario, 492 feet; and Lake Champlain, 93 feet.

Ν

b tl

0

m

m of

th

pr of

th

18

tor

dep

me

mo]

seri

of

.

t

west

sum

Ham

York

primi

over

At the west end of Lake Superior, and on its northern shores, several promontories, having an altitude of from 800 to 1000 feet above the water, give a mountainous character to the coast when seen from the surface of the lake, but which it is far from possessing when viewed in relation to the country lying behind it. It is such as would result from the excavation of the basin by the removal of the softer rocks which have covered the granites, porphyrics, and traps of these eminences. The silurian beds, not having been so extensively broken up to the westward of the *Fon du lac*, envelop the pyrogene nucleus so as, in conjunction with recent arenaceous deposits and drift, to cover it on that flank almost to the summit. From this locality, which is rather a plateau than a mountainous district, issue the feeding streams of the three several river

* Some discrepancy exists between the heights assigned to this lake by different authors. We have taken that deduced by Captain Lefroy from barometrical measurements made in connection with the observatory at Toronto. Dr. Houghton, the Michigan geological surveyor, estimates its height at 641 feet; but he makes the descent from it to Lake Huron 45 feet, while Professor Henry reckons this descent at only 18 feet, which must be under-estimated. Mr. Logan, in 1847, sets down the height of Lake Superior as 597 feet, having adopted for the height of Lake Michigan 578 feet, from Professor Henry. In the height of Erie and the inferior lakes authors are generally agreed.

LAKES MICHIGAN, HURON, AND ERIE. 179

systems, — of the Mississippi, St. Lawrence, and Saskatchewan, — as has been mentioned in a preceding page.

If we trace the south side of the St. Lawrence basin from this quarter, we find that already on the upper lake it assumes a different aspect, being composed chiefly of sandstone, and having less elevation than the north bank. Lake Michigan runs far to the southward among the silurian strata, entering as it were into the valley of the So small is the elevation of the ground Mississippi. between the Fox River which falls into Green Bay on the west side of that lake, and the Wisconsin, a tributary of the Upper Mississippi, that in times of flood a barge may float readily from one stream into the other.* A very moderate elevation in like manner separates the south end of Lake Michigan from the Illinois, another affluent of the Mississippi, so that a depression of 600 feet would produce a communication between the waters of the Gulf of the St. Lawrence and those of the Gulf of Mexico, through the Illinois valley.+ The basin of Lake Huron is excavated in the silurian strata, and the great promontory which divides it from Lake Michigan is said to be a deposit of old red sandstone enclosing the extensive coal measures of Saginaw. The lake shores are bold, but not mountainous.

Lake Erie has lower shores, and is the shallowest of the series. Its bed and much of its northern margin is formed of the corniferous limestone, one of the upper members of

* Featherstonhaugh.

[†] A depression of thrice that amount would carry the ocean to the western border of the prairie islands, leaving, as insular ranges, the summits of the Alleghanies and their continuation in Vermont, New Hampshire, and Gaspé, with a few peaks in the hilly region of New York, which lies between Lakes Ontario and Champlain; while the primitive masses on the north of Lake Superior would be mere reefs, over the highest of which an agitated sea would dash its spray.

ninaecks." om its perior d the re the l sucve the feet;

rthern f from ainous of the wed in uch as the reranites, silurian e westleus so d drift, om this tainous al river

to this Captain with the cical surdescent cons this r. Logan, t, having Professor thors are

the silurian rocks. Mr. Hall remarks that, had the eroding agency removed this bed, and penetrated to the soft strata of the Onondago salt-group, Eric would probably have been the deepest of the lakes.* The southern brim of its basin is so low, that an easy canal communication has been opened to the Ohio, a tributary of the Mississippi; and other water connections might be made with facility. Lake Ontario occupies a hollow in the silurian rocks inferior to those of the Erie basin+; and the country which lies to the south of it has in general a level character, though the Clinton and Niagara groups of rocks rise in places in high escarpments.

For a full account of the heights of the southern border, the whole valley of the great lakes and the St. Lawrence, and for detailed descriptions of the rock formations, the reader may consult the several geological surveys published under the authority of the legislatures of New York, Michigan, and Ohio: the preceding brief notices have been extracted therefrom for the purpose of showing the general character of the country, and the lowness of the barrier which separates the valleys of the Mississippi and St. Lawrence. I may add, as a further exemplification of the passes on the south side of the St. Lawrence, that a subsidence of 400 feet would cause the waters of Lake Ontario to flow through the valleys of the Mohawk and Hudson into the Atlantic, and at the same time convert Lake Champlain into a maritime strait, thereby forming islands of the states of New York, New England, and Maine, and of the British colonies of New Brunswick and Nova Scotia.

tl n

ab

St

the

des par

La

the

the Kir

a d

cons the

Before we proceed to trace the northern bank of the valley of the St. Lawrence, it will be convenient to notice

* Hall, p. 408.

 \dagger In the Medina sandstone, grey sandstone, the Hudson River group of shales, and, towards its eastern extremity, in the Trenton limestone, -Hall.

LAKE CHAMPLAIN.

more particularly Lake Champlain, since it is there that we have an approximation of the Apalachian chain to the primitive rocks which form the northern brim of the St. Lawrence basin, and unite with the intermediate belt on the north shore of Lake Superior.

Dr. Emmons* estimates the length of the valley of Lake Champlain at 180 miles, and its average width at about 20. Its bed is most depressed between West Point, Burlington, and Port Kent, where its soundings reach 600 feet, or 500 below the surface of the ocean.† It is in fact a deep chasm with a very abrupt slope on the western side, and a more gradual one on the eastern bank. The direction of the lake is north and south; it opens into the St. Lawrence basin on the north, and the valley of the Mohawk crosses its axis at some distance on the south. The summit level of the canal which connects it to the Hudson is only 147 feet above the tide; and a depression to that amount would cause the waters of the ocean to flow through it from New York Sound into the Gulf of St. Lawrence.

The New York highlands, bounded by Lake Champlain, the St. Lawrence, and Lake Ontario, are formed of primitive granite, hypersthene, and limestone rocks, which

* New York Geol. Survey.

[†] The lake lies, as has been stated in a preceding page, 93 feet above the tidal waters of the St. Lawrence: hence the fall of the St. John or Richelieu River, which discharges its waters just above the tidal level, may be estimated at 10 inches in the mile; and the descent of the St. Lawrence from Lake Ontario, excluding the comparatively currentless expansions of the Thousand Islands' Lake and Lake St. Francis, is nearly in the same ratio. These facts may aid in the calculation of the inclination of the beds of similar rivers. With the same view, I may add that the fall of the St. Lawrence between Kingston, at the outlet of Lake Ontario, and Montreal is 220 feet, in a distance of 160 geographical miles in a straight line, of which a considerable part is lake-way, which gives an average of 16 inches in the mile.

N 3

ding trata been basin pened water atario ose of uth of n and nents. order, rence, is, the blished York, s have ing the of the ppi and fication that a f Lake vk and e conthereby ngland, inswick

> of the notice

n River Trenton

constitute many striking and picturesque groups of conical peaks. Among these Mount Marcy rises 5,467 feet above the tide; Mount M'Intyre, 5,183; Mount Seward, 5,100; Mounts Martin and Santanoni, each 5,000; and Whiteface, Taylor's Mountain, and Nipple-top are a few hundred feet lower. These highlands extend into Canada, where they form a mountain belt twenty-five or thirty miles wide, along the sources of the Chaudiere and St. Francis.* They are separated from the neighbouring districts of the state of New York by the river and lake valleys named above; and, being impassable for an army, Lake Champlain was the only route by which the Atlantic states could be assailed from Canada East, or vice versa; — hence the celebrity of its defiles in the annals of colonial and revolutionary warfare.

g

re

si

sc si

th

C

ro

7,0

Po

thi bal

lan

Ma

the

wes

lime

met

epid rock

red

mor

of willong

bold

from

Acroc

The valley of the lake is excavated in the lowest group of the silurian rocks. The Potsdam sandstone, together with a greenish-white marble, and a trap rock, are quarried extensively at Whitehall, situated at the upper extremity of the lake, for the double purpose of clearing sites for houses, and procuring building stones. In various parts of the lake shores, the silurian rocks are covered with beds of clay and sand, in which there have been detected about twenty species of marine shells, which exist in a living state on the coast of the Atlantic at the present time. The general character of the scenery, and especially of the upper half of the lake, is bold, hilly, and picturesque, often rocky, but occasionally cliffs of clay and sand 100 feet high border on the water. The shores are low and shelving only in the bays, which are formed by short lateral valleys.

• Orford and Sutton Mountains are each reckoned at more than 4,000 feet high. The latter is the summit of a wide hilly tract, composed of chloritic and micaceous schists and gneiss. - Hunt.

GREEN MOUNTAINS.

183

The state of Vermont lies along the east side of the Champlain Valley; and the country, as it recedes from the lake, rises gradually into the acclivities of the Green Mountains, which are a continuation of the Alleghanics, and are prolonged into the Shick-Shock and Nôtre Dame Mountains of the promontory of Gaspé on the Gulf of St. Lawrence.*

The predominating rocks on this slope are an alternation of argillaceous slates, with slaty and fine-grained sandstones, and shaley grits belonging to the Hudson's River

* Mr. Hunt, of the Canada geological survey, says that "the whole of the Green Mountain Rocks belong to the Hudson River group, with the possible addition of a part of the Shawagunk conglomerates. The fossiliferous rocks of the St. Francis valley are referrible to the Niagara limestones of the upper silurian beds: a similar formation exists at Gaspé, and has been traced 150 miles south-west (in the direction of the Green Mountains); and from the similarity of Nôtre Dame (Gaspé) to the Green Mountains, and the fact that the Hudson River rocks flank the St. Lawrence to Cape Rosiere (Gaspé), we may conclude that the upper silurian rocks will be found to be nearly continuous throughout. Resting upon this formation, in Gaspé, is a body of arenaceous rocks, 7,000 feet thick, which apparently correspond to the Chemung and Portage group of New York, with the old red sandstones. As this formation is found extending quite to the Mississippi, it is probuble that it will accompany the silurian rocks through New England, surrounding the coal-fields of New Brunswick, and of eastern Massachusetts, and Rhode Island. To this may be referred, in part, the rocks of the White Mountains, which may sweep around the western border of Massachusetts' anthracite formation, until lost under the super-carboniferous rocks of the Connecticut River. The limestones of western New England seem to be no other than the metamorphic Trenton limestones of Phillipsburg; while the chloritoepidotic rocks and serpentines of Sutton valley appear again in the rocks of southern Connecticut, between these limestones and the new red sandstone. With such a key to the structure of the metamorphic rocks of New England, and of the great Apalachian chain of which these form a part, we may regard the difficulties that have long environed the subject as in a great degree removed, and the hold conjectures as to their metamorphic origin, which have been from time to time put forth, fully vindicated."-Hunt, Proceed. Am. Acroc. at Cambr., 1849, p. 333.

N 4

ical oove 00; ince, feet they wide, neis.* i the anned plain id be e the wyolu-

group gether arried emity es for parts about living time. lly of esque, sand re low short

e than

184

group. In breaking down, the shales produce a cold, clayey, retentive soil, much less favourable for agricultural purposes than the "birds'-eye" and other limestones which crop out nearer the shores of the lake. *

With respect to the northern slope of the St. Lawrence valley, a reference to the map will show that the brim of the basin, where its feeding streams have their source, is generally about 150 geographical miles from the centre of the lake-way from Ontario upwards, and considerably further off from the river channel lower down. The bays of the great lakes, in many places, curtail the breadth of the slope; and it is every where exceeded in breadth by the northern slope towards Lake Winipeg, James's Bay, and Hudson's Straits. It will also be perceived that the valley, with reference to its northern bank, makes an acute bend, of which the west end of Lake Erie forms the elbow; the lower or eastern arm being parallel to the Alleghany range, while the upper one takes more the direction of the axis of the Rocky Mountains. The angular form of the basin is in conformity with the course of the primitive rocks from Labrador to Lake Superior, where they blend with the "intermediate belt." † The general level of the

0

n

B

sh

ne

pe

da

the isla

and

of

sla

isca

the

which

612

and

acco

taine

the t

* During our voyage through Lake Champlain, I was informed by a fellow-passenger that the agriculture of Vermont was very superior, especially near Burlington, where there are many large orchards, and sheep farming is extensively pursued. Cleared land, he told me, sells currently at forty dollars (8*l*.) an acre, and fifty dollars are thought to be a high price. Two hundred acres form a good-sized farm in the opinion of the neighbourhood, and eighteen hundred acres a very large property. Many steamers are constantly employed in summer in visiting the various bays of the lake, and carrying the produce to Whitehall, whence it is transferred by canal to the Hudson.

+ The whole belt from the Labrador coast, along the valley of the St. Lawrence, and northwards to the Arctic Sea, seems to be a segment of the border of a great basin of which Hudson's Bay is the centre, and fragments of its eastern brim may be found on the shores of Hudson's Straits, and in the islands to the north.

ST. LA /RENCE BASIN — GEOLOGICAL STRUCTURE, 185

ridge formed by these rocks does not exceed the height of 700 feet above the river or lake surface; but, partaking of the general eastern slope of the continent, it attains a height of about 1,400 feet above the sea on the north side of Lake Superior. The course of the St. Lawrence through Canada East is conformable to the general strike of the beds in which the channel is excavated.

The geological structure of the north side of the St. Lawrence basin, as ascertained by the Canadian state survey under Mr. Logan, has been summed up as follows by his assistant, Mr. Hunt : - "A formation of syenitic gneiss, often passing into mica-schist, and interstratified with crystalline limestone, forms a ridge of high land extending from the coast of Labrador along the north side of the St. Lawrence, at a distance of from twelve to twenty miles from the shore, until it crosses the Ottawa, near Bytown, and thence is traced across Lake Simcoe to the shores of Lake Huron, where its northern limit is observed near the mouth of the French River, while it again appears at the south-eastern extremity of the lake in Matchedash Bay. Resting upon this is a series of rocks forming the whole north coast of the lake, and numerous small islands. It is made up of sandstones, often coarse-grained, and sometimes becoming conglomerate from the presence of red jasper pebbles. These beds are associated with slates and one or more bands of limestone. . . . The

• The following ascertained points may be noticed :- Lake Temiseaming, which is high up on the Ottawa, and near the line dividing the water-shed of that river from the Abitibbe and Moose River, which falls into Hudson's Bay, was ascertained by Mr. Logan to be 612 feet above the tide. The dividing portage between the Ottawa and Lake Nipissing is 696 feet above the sea, the lake itself being, according to Mr. Murray, 647 feet high. Lake Simcoe was ascertained, by the same observer, to have an altitude of 704 feet above the tide. - Logan's Geol. Rep. for 1848.

cold, tural /hich

rence im of ce, is tre of rably bays lth of th by Bay, at the acute lbow; ghany of the of the mitive blend of the

med by aperior, ds, and ne, sells thought farm in a very summer luce to

> of the egment centre, ores of

formation is much cut by greenstone dikes, and exhibits very frequently interstratified beds of greenstone, often of great thickness. Both these and the sedimentary rocks contain metalliferous quartz veins, of which the copper mines of this region are examples. Resting uncomformably on the tilted edges of this formation, and in other places directly upon the southern limit of the sycnitic gneiss, appear the silurian rocks, identical with those which are found in New York, and covering the peninsula between Lake Huron and Lake Ontario. Beginning with the Potsdam sandstone, we have, upon the Manatoulin Islands and the coast between Matchedash Bay and Sarnia, a complete exposure of those formations known as the Trenton limestone, Utica slates, Loraine shales, Medina sandstones, and the Niagara limestones, with the rocks of the Clinton group. All these are well characterised by their respective fossils, and are spread out quite undisturbed at a very gentle dip of about thirty-five feet in a mile.

"Passing to the east, we find that the synitic rocks have divided where they cross the Ottawa, and, taking a southward course, are spread over a considerable extent of country between the Ottawa and the St. Lawrence. Crossing this river below Kingston, they constitute the greater part of the Thousand Isles *, and are extensively developed in the northern counties of New York."

Mr. Logan gives a more particular account of the north

* In the Lake of the Thousand Isles, as the funnel-shaped outlet of Lake Ontario is denominated, many of the round-backed hummocks of granite which form the innumerable islets exhibit the parallei furrows, streaks, and smooth surfaces attributed by some geologists to glacial action. This expansion, in fact, has exactly the aspect of many of the dilatations of the northern rivers which flow through the "intermediate primitive district."

NORTH SHORE OF LAKE HURON.

187

shore of Lake Huron, of which I have made the following abstract : - " The north shore of Lake Huron, on which twenty-two mining locations have been claimed of Government, presents an undulating country rising into hills which sometimes attain the height of from 400 to 700 feet above the lake. These occasionally exhibit rugged escarpments and naked rocky surfaces; but, in general, their summits are rather rounded, and their flanks, with the valleys separating one ringe from another, are most frequently well clothed with hard and soft wood, often of large growth, and of such species as are valuable in commerce, in many places giving promise of a good arable soil. Many of the slopes are gentle, and many of the valleys wide. The Thessalon, of the reported length of 200 miles, and the Spanish river of 120 miles, flow through this country, with three other rivers of from 50 to 60 miles' length each." With respect to the rocks occupying this country, he says, that "for 120 miles from the upper end of the lake one great formation, having a breadth in some places of ten and in others exceeding twenty miles, exists. It is composed of sandstones, conglomerates, slates, and limestones; the sandstones often vitreous, and presenting the character of a perfect quartz rock. These are associated with greenstone trap and other igneous rocks, in the form of overflows, dykes, and veins, and with amygdaloidal trap in layers. The whole reposes on granite, which is the metalliferous rock of the district, and lics beneath the lowest known fossiliferous beds. The Potsdam sandstone, Trenton limestone, Utica slates, and Loraine shales are exposed in successive deposits, resting on the tilted beds of the quartz rock in a nearly horizontal, unconformable position in the Grand Manatoulin, La Cloche, Snake, Thessalon, Sulphur, and other islands, and at the east end of the lake. Medina sandstone and Niagara limestones

hibits en of rocks opper formother enitic which la bewith toulin Sarnia, as the **I**edina cks of sed by undis. et in a

e rocks king a extent vrence. ite the nsively

e north

d outlet mmocks parallei cologists spect of ngh the

188

exist in certain localities of the promontory of Cabot's Head."

is

pa

po

an

3.

pel

abl

wi

rat

Tra

to

silv

rocl

mar

veir

show

witl

six (

the

cons

of th

dalo

shap

capp

The

The

Supe

A

Of the mining places above alluded to by Mr. Logan, the Bruce mines, situated on the main shore behind the island of St. Joseph, are the principal; and, in 1849, when I was returning to Canada in the month of September, and had an opportunity of visiting them, 160 people were there employed, forming, with their families, a considerable village population. Mr. Logan calculated that 250 tons of dressed ore might be raised monthly, yielding, at an average, about 15 per cent. of copper. At present, fuel for mining purposes is obtained from the Pennsylvania coal-field, through the port of Cleveland, on Lake Erie; but, ere long, the coal district in Saginaw Bay, on the south side of Lake Huron, will become the smelting place for both the Huron and Lake Superior minerals.

To complete the outline of the north bank of the St. Lawrence valley, we may state that the primitive rocks continued from Lake Huron to the outlet of Lake Superior, form there the bold promontory of Big Cape (Gros Cap), which is a mass of flesh-coloured granite and Point Iroporphyry rising 700 feet above the water. quois, on the opposite side, three miles distant, is 600 feet high, has a more table-shaped summit, and from its base a line of low sandy beach stretches away on the south side, sending out a projecting tongue named "White-Fish Point." The north shore, as seen from Big Cape, presents a grand and finely varied coast line, deeply indented by Goulais and Batchewaung Bays, with the promontory of Mamainse, composed of rugged and crumbling amygdaloid, dipping into the wide expanse of waters on the north-west. The granitic ridge which skirts the bottoms of these bays comes out in bold cliffs on both sides of

NORTH SHORE OF LAKE SUPERIOR. 189

Michipicoten Bay, at the Otter's Head, and at the bottoms of Nipigon and Thunder Bays, projecting also in various capes between them. From the last named bay it stretches across the northern bend of the Kamenistikwoya River, by which stream the canoe navigation to the interior is carried on.

Mr. Logan considers the granite, which frequently passes into gneiss, as the base of the series of rocks composing this bank of the lake. 2. To this succeeds gneiss, and both are traversed by dykes and veins of granite. 3. The next in order are dark green talcose slates, and a pebbly and slaty conglomerate. 4. Resting unconformably on these, is a series of bluish shales, interstratified with trap. 5. Lastly, white or spotted sandstones, indurated marls, and conglomerates interstratified with trap. Trap dykes in vast numbers traverse all the beds down to the granite. Veins containing copper, lead, zinc, and silver, belonging to two systems,—cne coincident with the rock masses, the other parallel to them, - occur in very many places on the north shore. The courses of the veins vary in different bays, and Professor Agassiz has shown that the outline of the lake has a close connection with the directions of the trap dykes, of which he describes six different systems, each of them associated with one of the great curvatures or bays.

A granite porphyry, which is very durable, forms a considerable portion of the boldest and most barren parts of the north shore of the lake; the projections of amygdaloid, being more perishable, assume the most picturesque shapes; and some of the loftiest headlands are thickly capped with greenstone, basalt, and other trappean rocks. The low, flat, well-wooded islands are mostly sandstone. The general elevation of the northern brim of the Lake Superior basin may be stated at between 800 and 900

abot's

ogan, d the when ember, e were asiderat 250 ing, at resent, ennsylt Lake Bay, smeltuperior

of the imitive f Lake cape ite and ht Iro-**DO** feet s base south e-Fish resents ted by ory of amygon the ottoms des of

feet above the water surface, and the distance of its crest from the margin of the lake at from 20 to 50 miles.

Many mining stations have been granted to adventurers by the Canadian legislature, but workings are carried on in three only; viz. one in Pigeon Bay, one among the islands of Nipigon Bay, and the third in Mica Bay. At the last named 100 miners were employed in 1849, when the establishment was broken up by a foray of Chippeways, who thought that their territorial rights were invaded.^{*} But little ore has as yet been shipped from the Canadian side of the lake. The export of native copper from the United States shore is considerable. In 1849 1,114 tons of pure copper were transshipped at Saut Ste. Mary.[†]

Our limits will not allow us to dwell longer on the St. Lawrence basin[‡]; but with respect to the general character of the ridge which divides it from the Winipeg excavation, the aspect of the country traversed in pur-

* Narrative, Vol. I. p. 73.

- + The Detroit Free Press states that, in 1850, the shipments exceeded 4,000 tons, and it is calculated that they will equal the whole consumption of the United States, which is 6,000 tons in 1851. There are twenty-two copper companies in operation on the Michigan shores, employing 800 men. The masses of native copper on Kewawoonan Point are enormous; but, from their very purity, they can neither be blasted nor hewn out in the ordinary way. The method adopted at present is to use long iron chisels, which are turned slightly by the hand at each blow of the hammer. In this way large slabs weighing two tons or more are cut out, to which not above five per cent. of quartz rock and other impurities adhere. On this point there are said to be parallel ridges of trap rising through beds of sandstone. Among these the native copper lies in walls or veins which have two directions, one running across the trap ridges in N.W. 6 N. 1 N. direction, the other parallel to the ridges and strike of the sandstone.

[‡] Professor Agassiz has devoted a large volume to the natural history of Lake Superior alone, which is full of interesting facts and comprehensive general views.

LAKE SUPERIOR.

191

s crest

turers ied on og the r. At , when Chippeere incom the copper n 1849 ut Ste.

on the general Vinipeg in pur-

nents exthe whole 1. There Michigan on Kewathey can be method d slightly urge slabs five per his point beds of or veins ridges in nd strike

acts and

suing the canoe route may be considered as a type of the whole. The surface of that tract is hilly, the granite rising in rounded and sometimes in rugged knolls abruptly from lakes or swamps, but only to small heights above the general level. Here and there, but particularly towards the summit of the ridge, there are considerable deposits of sand, gravel, and loam, with many boulders. The term ridge is used with reference to its being a height separating two depressions; but its summit is a marshy plateau of some extent, across which narrow winding lakes afford a canoe navigation in a variety of directions. *

This summit of the water-shed, which, level as it is with respect to its water communications, is rendered very uneven by the protrusion of numerous granitic masses to various but moderate heights, lies much nearer to Lake Superior than to Lake Winipeg; and in descending towards the north, the same rocks appear in succession which have been noticed as forming the bank of the St. Lawrence valley, the "birds'-eye limestone" of the Champlain division of the New York system being the newest deposit on the Winipeg Lake.

If we trace the water-shed to the south-west, beyond the head of Lake Superior, by the sources of the St. Louis and head waters of the Mississippi, the uneven marshy surface gradually merges in the sandy prairie lands of the Red River, the Saskatchewan, and Missouri, where the naked rocks disappear, or are to be found only in the deep river channels.

The manner in which an elbow of the "intermediate primitive rocks," which form the nucleus of the watershed here spoken of, encloses the head of Lake Superior,

* Vide map of the district in Franklin's Second Overland Journey.

and extends along part of its south coast until lost under newer deposits, may be learned from the subjoined extract of a "Report by Dr. Owen, geologist to the States of Iowa and Wisconsin."*

* "The protozoic strata form sections on the Mississippi for an average distance, in a direct line, into the interior of Wisconsin, of from fifty to seventy-five miles, or up to the low falls of the principal castern tributaries of that river, where the crystalline rocks first appear. In this part of the country the igneous ranges do not rise into elevated mountains, but, on the contrary, they are seldom seen except in the immediate cuts of the streams, being, for the most part, covered with drift. The character of the country generally, towards the summit levels leading to Lake Superior, is a succession of terraces of moderate elevation, chiefly composed of drift, having a nucleus, no doubt, of granite, syenite, or hornblende rocks; but these protrude only occasionally. At intervals the streams are ruffled into rapids, being filled with boulders which materially obstruct their navigation. A portion of these boulders may have been transported from great distances, but the greater part appear not to be far removed from their parent rock. It is a matter of surprise that so large an area of the interior of this district, and indeed of the sources of the Mississippi generally, should be level tamarack and cedar swamps, since, on approaching a great water-shed that gives rise to one of the greatest rivers in the world, one is led to anticipate a country with physical features of quite a different character. Interposed between the crystalline and igneous rocks of the interior of the district and the lowest sandstones, some green and red schistose beds have been observed at different localities. These appear to have been derived from the decomposition and detritus of the more easily disintegrating felspathic granites. The lower bcds of sandstone adjacent to the igneous outburst are not unfrequently changed to hard quartzite.".... "The highest ranges of the Wisconsin side of Lake Superior, situated from ten to sixteen miles from its south shore, are estimated to be near 1,000 feet above the lake, and are formed of hornblende rocks, metamorphic slates, syenite, and trap. No organic remains have been detected in the great mass of sandstone bounding this part of the lake whereby its geological era may be determined. On the west side of the Mississippi, north of the Winebago reserve (Minesota), and as far north as St. Peter's river, limestone, with underlying sandstone, prevails to the extent of half a degree of longitude."

e p ri M su Т L L cos Li abo mil Ma and wer the a w tarie or A of tl prain bank the **H** V

VALLEY OF THE WINIPEG.

193

under extract ates of

for an onsin, of orincipal eks first not rise om seen ost part, towards 1 of terhaving a cks; but re ruffled ruct their ansported be far ret so large ces of the swamps, ne of the ntry with between trict and have been n derived tegrating t to the te." situated ted to be de rocks, ave been t of the the west ota), and andstone,

VALLEY OF THE WINIPEG OR SASKATCHEWAN.

The next great transverse excavation includes the whole valley of the Saskatchewan and Nelson Rivers; and from the sources of the former near Mount Hooker, one of the Rocky Mountain peaks said to be 15,700 feet high, to the mouth of the latter in Hudson's Bay, the axis of the valley runs in a direct line about east-northeast for 660 geographical miles. Lake Winipeg, the principal lake basin in connection with the river, lies at right angles to that axis, is nearly parallel to the Rocky Mountain chain, and forms one of a series of great lakes succeeding one another in a north-north-west direction. Their names are Lake Superior, Lake Winipeg, Deer Lake, Wollaston Lake, Athabasca Lake, Great Slave Lake, Marten Lake, and Great Bear Lake; the northern coast line being moreover indented on the same bearing by Liverpool and Franklin Bays.

Lake Winipeg itself is 230 geographical miles long and about 40 wide, but its width would be increased to 120 miles, if Moose Lake, Muddy Lake, Winepegoos and Manitoba Lakes, which differ very slightly from it in level, and are evidently component parts of the same lake basin, were included in the measurement. Into this great lake the Saskatchewan, by its two diverging branches, gathers a wide extent of prairie drainage, its northern tributarics being conterminous with the affluents of the Elk or Athabasca River, and its southern ones with those of the Missouri. The Assinaboyn also traverses much prairie land, one of its branches originating on the banks of the great southerly bend of the Missouri; and the Red River, which the Assinaboyn joins, rises, as has

VOL. II.

been already mentioned, on the same level and in the close neighbourhood of the sources of the Mississippi and St. Lawrence. The waters of the lake wash, on the east side of the basin, the "intermediate primitive rocks," and find their way through them by anastomosing channels named Sea River, Katchewan, and in its lower part Nelson River. After it has passed through the intermediate belt, it takes its way over silurian linestones, and finally enters Hudson's Bay through alluvial deposits of some extent.

The surface of the lake has been calculated to be 853 feet * above the sea, and its basin is excavated in the silurian beds. Along the whole eastern shore the granite, gneiss, and trap rocks are every where exposed, the first named rock being the most extensive; and nowhere do these masses rise to the altitude of hills. On the north and west the birds' eye limestone is the prevailing rock, and forms low cliffs, in a country otherwise every where flat; and towards the south end of the lake, and in the narrows, arenaceous deposits appear in the immediate vicinity of granite, trap-rocks, and chlorite slates, having a close resemblance to those of Pigeon Bay of Lake Superior, where argentiferous veins occur. It is, therefore, an interesting quarter for exploration by the practical miner.

In ascending from Lake Superior by the Kamenistikwoya, and its upper branch named the Dog River,

* Captain Lefroy's observations :----" In the Geological Appendix to Sir John Franklin's Second Journey, I estimated the height of Lake Winipeg above Hudson's Bay at 800 feet, which I considered to be a rough approximation. Major Long places it at the same level with Lake Superior, and the dividing ridge between the two basins at 600 feet higher; but, by the best estimates I have been able to form, he makes the summit of the ridge 230 feet too low." a

FU

A

U

Dog Asc Port Wes

Lake Heig

Th was Majo Scott the S Surve *kabikk* found differe way al by bar In 1 by me

HEIGHT OF LAKE WINIPEG.

195

to Thousand Lakes* (Milles Lacs), forty portages are made, in which the whole or part of the cargo is carried,

* The height of this lake above Lake Superior was ascertained approximately as follows :---

Foot of Mountain Portage estimated by Captain Lefroy	-	Feet. 48	
Upper end of ditto (bar. meas. Rich.)	-	127	
Ascent of river to Dog Portage (estimated, Rich.) -	-	150	
Upper end of Dog Portage above the lower (bar. meas. Ric	h.)	332	
Ascent of Dog River to Thousand Lakes (estimate)	-	160	
		817	
Lake Superior above the sea	-	641	
Thousand Lakes above the sea	-	1458	

Owing to the obstructions in the upper part of the Dog River, the canoe route diverges from that stream at Cold-water Lake, and passes by the Prairic, Middle, and Savannah Portages, and a small stream which flows from the latter into Thousand Lakes. The heights of this part of the route are as follows :---

Dog Lake, above Lake Super	ior		-	-	Feel 657
Ascent of Dog Rive.	-	-	-	-	14
Portage to Cold-water Lake	-	-	-	-	2
West end of Prairie Portage a	and Mi	dle Portage	-	-	161
					834
Lake Superior above the sea	•	-	-	-	641
II. LA CONTRACTOR	Dente				1470

Height of Prairie or of Middle Portage above the sea

The difference of height between the ends of the Mountain Portage was carefully ascertained by me with Delcro's barometer in 1849. Major Delafield estimates the difference at 125 feet, and Lieutenants Scott and Denny, who accompanied Major Long in his voyage up the St. Peter's, at 130 reet. Mr. Murray, of the Canadian Geological Survey, measured the actual height of the falls here (named Kakkabikka) from the smooth water at their summit to the base, and found it to be 119 feet; but at least 8 feet must be added to give the difference between the ends of the portage, which terminates some way above the brow of the cascade. Captain Lefroy makes the height, by barometrical measurement, only 115 feet.

In 1849 the height of the upper end of Dog Portage was ascertained by me with Delcro's barometer; in the previous season the aneroid 02

n the pi and ie cast ," and annels rt Nelnediate finally of some

be 853 in the granite, the first here do ie north ng rock, y where d in the mmediate s, having Lake Sutherefore, practical

Kamenisg River,

Appendix e height of considered same level two basins een able to
PHYSICAL GEOGRAPHY.

196

there being besides some long rapids where it is not necessary to unload; and in descending to Lake Winipeg the portages are about fifty, their number and even the route varying with the height of water. Thousand Lakes (Journ. p. 62.) is an extremely irregular piece of water, having many extensive arms, some of which are very Multitudes of islands well covered with birch, shallow. aspen, arbor-vitæ (Thuya occidentalis), and the various pine trees of the region, render the scenery pleasing. A few granite knolls and mural precipices show among the trees; but many of the islets appear to be formed of sand, of which sections twenty feet high occur. The lake not only branches into the neighbourhood, but water communications diverge from it in every direction, as is customary in the "middle primitive belt."

From the Thousand Lakes the canoe route keeps on the border of the primitive rocks, touching on silurian deposits, when it bends to the southward or westward. At first it is flanked on both sides by granite. In Rainy Lake there is much mica-slate; and at its outlet the stream, falling over gneiss rocks, produces the cascade of *La Chaudière*. The greater fertility of the country about Nemican Lake and Rainy River show the vicinity of newer formations. In the Lake of the

barometer gave 328 feet as the height, which was a greater degree of accordance between the instruments than I generally found. Major Long estimates the water-shed between Lake Winipeg and Superior at 1,200 feet above the tide; Major Delafield calculates the height of Cold-water Lake at 505 feet, to which, if 161 be added for the Prairie Portage, and 641 for Lake Superior, we have 1,307 feet for the height of Prairie Portage above the sea. Captain Lefroy, by barometrical measurements made in connection with the observatory at Toronto, makes the west end of Prairie Portage 1,361 feet above the sea; but the distance between the two places of observation renders the result liable to some error.

V R cc ni ro 18 asc the we wh to . be the Lak ridg sepa the the Pete part and i conce

t

l

(

I

3

prairi

COTEAU DES PRAIRIES.

is not inipeg en the Lakes water, e very birch, various ng. A ong the of sand, lake not er comis cusceeps on silurian restward. In te. ts outlet the casr of the er show ke of the

r degree of id. Major 1 Superior e height of the Prairie the height arometrical t Toronto, ie sea; but s the result Woods Dr. Bigsby found the Pentamerus Knightii, a fossil characteristic of the upper silurian rocks; but granite is the chief constituent of the islands and shores of the lake. The River Winipeg flows wholly within the granite district, and has the lake-like dilatations and other characteristics of the streams which traverse the "intermediate primitive rocks." The wide extent of land which its reuniting arms enclose is remarkable. One of its affluents, named English River, issues from Lake Sal, which lies near the water-shed, dividing the Winipeg basin from that of James's Bay. The Berens River which falls into Lake Winipeg, issuing from the common brim of the same basins further north, in the vicinity of the sources of the Severn River, affords a canoc route to Hudson's Bay. From my own observations in 1819, and Mr. Barnston's at a later period, it has been ascertained that limestones of the silurian epoch occur on the northern flank of the "intermediate primitive belt," as well as in the basin of Lake Winipeg. The Red River, which has been already repeatedly referred to, and which to prevent confusion with its southern namesake it would be well to call Osnaboyna, lies wholly to the westward of the "intermediate belt," and has a direct course from Lake Travers of 300 geographical miles. The sandy ridge named Côteau des Prairies, or Hauteur des Terres, separates its upper part from the Missouri valley; but the metamorphic rocks which present themselves around the sources of the Mississippi and its tributary the St. Peter, are also visible near Lake Travers. In the lower part of Red River limestone crops out in one place only, and is quarried by the settlers. Elsewhere the rocks are concealed by the sandy deposit forming the soil of the prairies, along whose eastern border the river flows. Major

o 3

PHYSICAL GEOGRAPHY.

Long enumerates thirty affluents of the Red River, and its western branch, the Assinaboyn.

The Saskatchewan, which is to be considered as the main feeder of the Winipeg basin flows from a considerable distance above Pine Islar . Lake down to Lake Winipeg over horizontal beds limestone, through so flat a country that the river forks as it would in an alluvial delta. A rich mud is deposited in parts, particularly between Pine Island Lake and the main stream, and round Moose and Muddy Lakes. In Beaver Lake, which lies immediately to the north of Pine Island Lake, the silurian strata are again seen covering the flanks of the primitive rocks; while to the southward an eminence named Basquiau*, lying at the distance of nearly a degree, separates the river valley from the Red Deer Lake and Swan River. As powerful salt springs exist on this eminence, we may conjecture that it belongs to the Onondago salt group.

I have been more particular in the topographical and geological remarks on the St. Lawrence and Saskatchewan basins, because economy of space made it expedient to omit the details of the voyage of the Expedition through them. The remaining districts will be more briefly mentioned here, since the narrative included many facts relating to them.

VALLEY OF THE MISSINIPI.

The next river basin that we have to notice is that of the Churchill, English River, or Missinipi[†], the latter, or

* See Journal, I. p. 68.

† Misi or mitchi, in Cree, significs "much or great," and nipi "water," while sipi means "river."

_1

ha ka pra tw few pou thr para

C

(

t

g

ea

I

th

F kenz subo south borde

VALLEY OF THE MACKENZIE.

199

r, and

as the asider-Lake agh so a allucularly l round ch lics silurian imitive ed Base, sepad Swan ainence, ago salt

raphical Saskatit expepedition be more ed many

that of tter, or

and nipi

Cree appellation, being nearly synonymous with Mississippi. This basin, in crossing the intermediate primitive rocks, lies nearly parallel to the Nelson River or Katchewan, which is the lower part of the Saskatchewan. Deer and Wollaston Lakes, which discharge their waters into the Churchill River, lie in the line of lake basins mentioned above as running northwards from Lake Superior; but their axes do not cross the river valley so nearly at a right angle as Lake Winipeg does the valley with which it is connected. Further up the Missinipi the Methy, Buffalo, Clear, and Isle à la Crosse Lakes, which are situated just to the westward of the primitive rocks, taken in the aggregate, lie more in the plane of Lake Winipeg. On the eastern flank of the intermediate primitive ridge lie Big Indian and Waskayow-washgow Lakes, belonging also to the Missinipi river system.

The Missinipi, by its principal feeder, the Beaver River, has its source lower down the eastern slope than the Saskatchewan, and drains a comparatively small extent of prairie lands. At the Frog Portage and elsewhere the two basins are divided from each other by rocks only a few feet high, over which, in times of flood, the waters pour; so that the two may be viewed as one great valley through which two large rivers flow, their trunks running parallel to each other.

VALLEY OF THE MACKENZIE.

Further to the north lies the great valley of the Mackenzie, extending to the Arctic Sea, but having also its subordinate transverse lake basins, which differ from the southern ones in their heads merely entering the western border of the intermediate primitive rocks, and their dis-

PHYSICAL GEOGRAPHY.

charging streams taking an opposite direction through the newer deposits.

Methy Portage forms the dividing brim between the Missinipi basin and the Mackenzie River valley, at the place where it is crossed in the usual canoe route; and, though the country be wooded, it may be considered as a partial extension of the prairie slope. The strata are bituminous shale * resting on silurian limestone, and covered with a thick arenaceous deposit. This is deeply furrowed by the channels of the Elk River, and its tributary the Washacummow; but the lake basins which mark out the border of the intermediate primitive rocks must be sought for further to the eastward. It is probable that this border touches a straight line drawn from Knee Lake, across the outlet of Athabasca Lake, to the deep northern arm of Great Slave Lake, and onwards by Marten Lake, across the two eastern arms of Great Bear Lake, to the Copper Mountains. That portion of the line which lies Letween Athabasca Lake and Methy Portage is little known, because the water route lies to the westward of it. Wollaston and Deer Lakes already mentioned, will, if this line be correctly drawn, be situated considerably within the border of the primitive rocks; and an illustration of the manner in which the waters occupying the minor valleys of that district communicate with each other is afforded by Wollaston Lake sending a stream from its north end into Athabasca Lake, and one from its south end into the different river system of the Missinipi. This fact, which, as reported on the authority of the fur-traders, is expressed in Arrowsmith's map, may be considered as proving that Wollaston Lake is considerably elevated.

* It is doubtful whether this shale be referable to the Marcellus shale or not.

* heig

i

tl

tł

to

p

W

of

be

se

pr

of

fou

un

ATHABASCA RIVER.

h the

1 the t the and, ed as a are , and leeply tribumark must le that Lake, orthern Lake, to the ch lies s little d of it. if this within tion of minor ther is om its south This traders, ered as evated.

larcellus

Indeed, it seems to be high up on the water-shed which separates the Mackenzie valley from the basin of Hudson's Bay.

According to Captain Lefroy's measurements and estimates, Methy Lake is about 1,540 feet above the sea; and I ascertained the summit of Methy Portage road to be 188 feet higher than the lake; while the Washacummow, or Clear-water River, on the north side of the portage, is 590 feet below it,' and by my calculations 910 feet above the sea. From Methy Portage westwards, the country, though deeply furrowed by river courses and ravines, and more or less thickly wooded, partakes so much of a prairie character that horsemen may travel over it to Lesser Slave Lake and the Saskatchewan. At the junction of the Washacummow with the Elk, the channel of the river is sunk 925 feet below the summit of the portage, which may be assumed as near the mean level of the district, and is 1,688 feet above the sca. *

The Athabasca, Red Deer (La Biche), or Elk River, the most southerly feeder of the Mackenzie, originates in the Rocky Mountains near Mount Browne, which is said to rise 15,000 feet above the sea, and, flowing through prairie lands, receives the waters of Lesser Slave Lake, whose axis crosses its general course, and afterwards those of Red Deer Lake. Its bed is in many places deeply cut beneath the level of the prairie plateau, which is not separated by any marked ridge from the Saskatchewan prairie country. Further north, the wide-spreading sources of the Peace River drain the Rocky Mountain chain for four degrees of latitude, and the trunk formed by their union curves across the slope to join the Athabasca a

* See Narrative, Vol. I. p. 114. Captain Lefroy's estimate of the height of Methy Lake exceeds mine by 40 feet.

PHYSICAL GEOGRAPHY.

little below the transverse basin of the Lake of the Hills. The channel of the Peace River is cut into silurian or Devonian strata; but the Lake of the Hills, or Athabasca Lake, runs eastward among the "intermediate primitive rocks," and, as has been said above, receives a tributary stream from Wollaston Lake, situated near the watershed which divides the basin from Hudson's Bay. The conjoined stream of the Peace and Athabasca Rivers assumes the name of Slave River, which flows in the fracture between the silurian and primitive rocks. The junction of the western arm of the river with Great Slave Lake marks the western border of the primitive rocks, which is also indicated on the northern side of the basin by the Fort Providence Inlet. Aylmer and Artillery Lakes lie 150 feet higher at the east end of the lake, into which they send their waters; and in their immediate vicinity, on the same plateau, are the sources of the Great Fish or Back's river, which falls into the Arctic Sea, as has been already stated. The streams that run from the westward into Chesterfield Inlet must come from near the same place; and this inlet, from its transverse direction, and east-north-east bearing from Great Slave Lake, has an evident connection with that excavation, their axes being nearly at right angles to the Rocky Mountain chain.

In Great Slave Lake the Mackenzie is deflected from the intermediate primitive belt, and flows first westwards then northwards, in a channel scooped out of the upper silurian strata and still newer deposits, for 1,000 statute miles of river course, or nearly 600 geographical miles in a direct line; neither granite, gneiss, nor mica slate are seen on its banks, and even trap rocks are rare, if any actually occur.

d

fi

0

li

n

h

of

The River of the Mountains, embracing by its feeders a more northern part of the Rocky Mountain chain, after

202

NOH'HANNÈ BUTE.

203

disengaging itself from the rugged hills from which it draws its supplies, makes a northerly bend nearly parallel to Slave River, and then joins the Mackenzie at Fort Simpson. Of this mountain stream I have already given a slight notice in the Narrative (Vol. I. pp. 167-170.); and I may add here that for twenty-five miles upwards from its mouth it flows through sand and shale, with limestone occasionally cropping out. At the end of that distance there is a rapid, above which low wooded points exist, with at intervals mountain bluffs coming down to the banks of the river. The most remarkable of these stands at the influx of the Noh'hannè River, and is named the Noh'hanne bute. It is the highest hill in that quarter, and is about seventy-five miles from Fort Simpson. Perhaps it is a member of a range whose prolongation is seen indistinctly in descending the Mackenzie about halfway between Hare Skin River and Fort Simpson. Messieurs M'Pherson and Bell ascended it, and the latter gentleman was seized with nausca and vertigo before reaching its summit; so that its altitude is probably considerable, but the snow disappears from it in summer. On its top there is a salt spring, having a basin fifteen feet in diameter, which is never dry. For this notice I am indebted to Mr. M'Pherson, who brought from it some fragments of limestone that were similar in lithological character to those procured at the rock by the river's side, described in page 182. of the first volume of the Narrative.

Great Bear Lake, the most northerly of the transverse freshwater lakes, lies about 150 feet above the channel of the Mackenzic, and crosses the arctic circle on the line where the hypogenous and silurian rocks meet. Coronation Gulf is also excavated on the same line, which has a general parallelism to the Great Slave Lake series of excavations enumerated above.

[ills. n or asca itive itary ater-The s asfrac-The Great nitive of the tillery e, into ediate Great Sea, as om the ear the ection, e, has r axes chain. l from twards upper statute es in a re seen tually

ecders

after

PHYSICAL GEOGRAPHY.

The northerly trending of the coast line west of the Mackenzic is evidently due to the prolongation of the Rocky Mountains, whose successive ridges or spurs come out *en échelon*, diminishing in altitude as they approach the shore.

The short western slope of the continent from the Rocky Mountains to the Pacific differs from the eastern one in its configuration, its river valleys being all more or less transverse. The peculiar wing-like projection in the north, towards Asia, is evidently due to the volcanic chain of Alaska, which runs at right angles to the Rocky Mountains. The great transverse river valley of the Yukon or Kwichpack lies to the north of it. The western seacoast from Cook's Inlet to Behring's Straits is delineated in a map attached to Baer's "Nachrichten über den Nordwestküste von Amerika," published in 1839, though it seems to have been hitherto neglected by our topographers; and the interior of Russian America has remained a blank in our maps.*

For the following information respecting the Hudson's Bay Company's land to the west of the Mackenzie, and for some account of Russian America gathered from the natives, I am indebted to Chief Factor M'Pherson, Chief Trader Bell, and to the letters of Mr. Alex. H. Murray, alluded to in a preceding page. By ascending the River of the Mountains, and tracing its north-west branch to Lake Frances, a very elevated mountainous country is reached. In this the Lewis, which flews towards the Pacific, takes its rise, its sources springing to the eastward of those of the River of the Mountains; so that here,

* The newest work on this district of America is, "Beitrage zur Kenntniss der orographischen und geognostischen Beschaffenheit der Nordwestkiiste Amerikas, von Dr. C. Grewingk, St. Petersburg, 1850; but this I have not been able to procure.

 $\mathbf{204}$

THE YUKON.

as well as in other parts of the Rocky Mountain chain, the rivers falling into opposite seas interlock at their origin. In lat. 61° 30' N., long. 130° W., the Lewis is joined by the River Francis, which has no connection with the lake of a similar name, but comes from Russian Lake, a sheet of water lying more to the south. At the junction of the Francis and Lewis, Mr. Roderick Campbell has built a trading post, named " Pelly Banks," which, according to his experiments with the boiling water thermometer, is elevated 1,400 feet above the ocean. The united streams take the name of the Pelly, which falls into the Pacific, probably into Tchilikat or Lynn Canal, but the exact situation of its mouth has not been ascertained. Native traders come from the head of Tchilikat to Pelly Banks in a fortnight.

From the same elevated district in which the Lewis or Pelly and the north-west branch of the River of the Mountains take their source, the Yukon, a river of great magnitude, issues; and for a considerable part of its course flows to the north, through a country which, as far as I can judge from the descriptive notices of it which I have collected, closely resembles the valley of the Mackenzie. Mr. Murray was disposed at one time to identify this river with the Colville, which falls into the Arctic Sea about 120 miles to the east of Point Barrow; but more recent and full native information leads him now to conclude that it flows towards Norton Sound; and one of the officers of the Enterprise, in a private letter which has been published, states that the Russians have established its identity with the Kwichpack, which falls into Beering's Sea between Cape Stephens and Cape Romanzoff.*

* The following is an extract of a letter to me from Mr. Murray, dated May 1850, on the Yukon :--- "My account of the course of this river, also a sort of chart I made of it from the description given by

the the come oach

the stern re or a the chain Iouncon or scaneated r den hough topoas re-

dson's e, and m the Chief urray, River nch to try is Is the eastthere,

> ige zur ieit der , 1850;

PHYSICAL GEOGR. PHY.

206

In 1847, Mr. Bell, having heard of the Yukon from the Kutchin who visisted the fort on Peel's River, set out in quest of it accompanied by a native guide. He first crossed the mountains to a stream termed the Rat River. on which an outpost named La Pierre's House has since been built. This post is about sixty miles distant from Bell's Fort on the Peel, and is about ten miles to the southward of it. Shortly after embarking in a canoe on the Rat River, Mr. Bell came to one of much larger size to which it is tributary, and which is named the Porcupine. Three days' descent * of this carried him into the Yukon, which it enters at right angles in the 66th parallel, and in the supposed longitude of $147\frac{1}{4}^{\circ}$ west. † At this place the Yukon is 14 mile wide, and is full of wellwooded islands, with a very strong current in the channels which separate them. After issuing from the mountainous

the Indians, might, perhaps, lead you to have a wrong impression respecting the mouth of the river. I am now convinced that it is not the same with the Colville, and I have for some years suspected that its mouth lay to the west. The Russians have come up the lower part of the river regularly for some seasons : I was at first informed that they entered it from another river, but I am now told positively by Indians who went down and met them last summer that they come into it direct from the sea. By one of these Indians I received a letter from the Russians, which, being in their own language, is unintelligible to me. Salmon and hooked-nosed trout (Salmo scouleri) ascend the river, but are not found in the Mackenzie, or rivers falling into the Arctic Sea. Again, I have made frequent inquiries of the 'Gens du large,' or the northern Indians, who visit the Arctic Seacoast, and find that they are unacquainted with the mouth of the river. For two winter days' walking below the Porcupine, the Yukon trends to the west and south-west, and the natives say that it flows on in the same direction. I am therefore inclined to believe that the Colville is a smaller river, and that the Yukon empties its waters into Norton Sound."

p

E

80

p

te

rig

po

* In returning from the Yukon against the stream, by tracking the canoe, nine days were required to reach Rat River.

† By Mr. Murray's courses and distances.

COURSE OF THE YUKON.

district north of Pelly Banks, the course of the Yukon is, according to the Indians, to the westward of north; and in one place where it cuts a spur of the Big Beaver Mountains, it passes between high limestone cliffs resembling the "ramparts" on the Mackenzie. Its current is every where more rapid than that of the river just The first important tributary which it receives is named. the Red Island River, that flows in from the mountains on its eastern bank, and is divided at its source from the head of the Peel by a single ridge of land. Between it and the Lewis there is a barren plateau, which the Indians cross in four days, but on which they find no water.* Another tributary from the east comes in lower down, and below that Deep River enters from the west. The Russians appeared on this stream with a skin boat (baidar), coming overland, it was supposed, from the Copper River or Atna, which joins the Pacific in Comptroller's Bay. From below the "ramparts," which are reckoned to be about seventy miles above the influx of the Porcupine, the Yukon flows to the north-west through a flat country; but twenty or thirty miles below the mouth of that tributary it again cuts a spur of the Beaver Mountains, trending at the same time to the south-westward and westward, and finally issuing in Beering's Sea, under the Eskimo appellation of Kwichpack, as has been mentioned above. Below the Big Beaver Mountains it receives from the south the "River of the Mountain Men," which runs parallel to the main stream for some distance before entering it. From thence to the sea the Kwichpack flows,

* There is a district in Siberia on which neither snow nor rain is said to fall. This may be a similar one; for it can scarcely be, in so rigorous a climate, that melting snow, if it exists, should not leave pools of water all the summer.

the out first ver, ince rom the e on size Poro the allel, this wellnnels inous

ression t is not ed that er part ed that ely by r come ived a age, is ouleri) falling of the c Seaof the Yukon t flows e that waters

ng the

PHYSICAL GEOGRAPHY.

according to the natives, through a low, marshy, and very sparingly-wooded country.

Mr. Murray, in his letter to me quoted above, mentions his intention of exploring the Yukon as far as he had leisure to do so in the summer of 1850; and it was probably the report of his party having been seen which induced Captain Collinson to land Lieutenant Barnard and Mr. Adams at Fort Michaelowsky, that they might ascertain who the white men were.

On the Pacific coast only of North America does volcanic action exist at the present time. Many peaks throwing out fire and smoke are mentioned by Bäer. They lie in a line running west-south-west from the north side of Kanai or Cook's Inlet, through the peninsula of Alaska and the Aleutian Islands. Many of their summits rise into the region of perpetual snow. Among these Ilämän has a height of 13,151 feet, and Winkelhöhe of 11,270, both lying on the north bank of Cook's Inlet. In Alaska there is the Peak of St. Paul, and in the islands Schischaldin which is 8,953 feet high, besides Schimaldin, Makuschkin, Korowenskische, and many others. The line in which these volcanic peaks lie when prolonged to the eastward, strikes the Big Beaver Mountains on the Yukon.

Mounts Edgecumbe, Fairweather, and St. Elias are, I believe, extinct volcanoes, which form, with those of Alaska, nearly the segment of a circle.

On the side of the Atlantic modern volcanic rocks occur in Jan Mayen's Island only, whose principal mountain, Becrenberg, rises 6,870 feet above the sea. In the coal of Jameson's Land, lying in north latitude 71° on the east side of Greenland, and in that of Melville Island, in latitude 75° north, Professor Jameson found plants resembling fossils of the coal fields of Britain. This fact is sufficient of itself to raise a world of conjecture respecting the conan ria co Be bo: bei the nor 73c

n

0

* the g V

moi

208

N

FOSSIL BONES.

very

ntions e had s proch inrd and ascer-

oes volpeaks Bäer. e north asula of ummits g these lhöhe of 's Inlet. e islands imaldin, The line d to the Yukon. as are, I those of

> ks occur ountain, e coal of the east in latiembling ufficient the con

dition of the earth when these ancient fossils were living plants. If the great coal measures, containing similar fossil vegetable forms, were deposited at the same epoch in distant localities, there must have existed, when that deposition took place, a similarity of condition of the North American continent from latitude 75° down to 40°.*

In concluding this sketch I shall advert to the distribution of the fossil remains of mammalia, and especially to those of the mammoth. Teeth of this animal have been discovered on the banks of several rivers in Russian America, north of Mount St. Elias; and there is a celebrated locality in Kotzebue Sound where the thawing and waste of frozen cliffs is continually exposing the bones and tusks of mammoths and other quadrupeds. Dr. Buckland, in his interesting account of the specimens collected at this place, on Captain Beechey's voyage, enumerates fragments of bones of mammoths, and of the urus, the leg-bone of a large deer, and a cervical vertebra of some unknown animal, different from any that now inhabit Arctic America. Along with these there were found also the skull of a musk ox and some bones of the reindeer in a more recent condition than the others. In Asia, from the Don to Beering's Straits, on the banks of all the great rivers, bones and teeth are still more plentiful, the local deposits being richer and more extensive the more we advance to the north. They are especially abundant on the islands north of Sviatoi Noss (Sacred Cape), lying between the 73d and 76th parallels; and some of these islands seem to consist in a great part of organic remains, which fill more space than the matrix in which they lie. The soil

* The coal of Vancouver's Island and Oregon belongs, it is said, to the great coal measures. P

VOL. II.

PHYSICAL GEOGRAPHY.

containing them is frozen as hard as a rock; and, as it thaws annually, the bones drop out or are quarried by the natives. From time immemorial this has been proceeding, and the ivory hunters have been obtaining their annual supply without any sensible diminution of the store they draw from. Some tusks found in New Siberia weigh 480 lbs., and from that island alone the merchants of Yakutzk, in 1821, obtained 20,000 lbs. weight of this fossil ivory. Though the tusks of the mammoth (Elephas primigenius) are chiefly sought for, bones and teeth of other animals are found, and among the rest those of the Rhinoceros tichorinus, of which the skulls, flesh, and skin, with its hair, have been procured. The remarkable discovery in 1799, by a Tungusian, at the mouth of the Lena, of the entire carcase of a mammoth, in excellent preservation, is well known. The existence of these numerous testimonials of an ancient fauna is suggestive of many curious speculations, and geologists seem hitherto to have failed in explaining the circumstances under which accumulations so vast could occur in such high latitudes. The difficulty is increased when we consider that these bones have not been detected to the east of the Rocky Mountains in the northern latitudes. None have hitherto been found in Rupert's Land, though the annual waste of the banks of the large rivers and the frequent land slips would have revealed them to the natives or fur traders had they existed even in small numbers. They are rare also, or altogether wanting, in Canada, but in the valley of the Mississippi the bone-licks are well known as most extensive and furnishing the remains of a different series of extinct quadrupeds.

The existence of the skull of a musk ox in Kotzebuc Sound is of much interest. This relic is preserved in the

BOULDERS.

as it y the protheir f the Siberia chants of this Elephas eth of of the id skin, ble dise Lena, reservaumerous of many to have ch aceu-The 29. se bones v Mounerto been waste of and slips r traders are rare he valley h as most ent series

> Kotzebue ed in the

British Museum, and the naturalists who have compared it with recent skulls of the animal brought from Melville Island or Churchill, perceive no difference. But, as the species is not known to frequent any district to the westward of the Mackenzie, the transport of any of its bones in modern times to Kotzebue Sound cannot be readily accounted for.

In a preceding page I have alluded, in a general way, to the distribution of boulders on the eastern prairie slope and in the valley of the Mississippi. The journal of the voyage shows that these are everywhere present in the north. The surface soil, the beds of rivers, and sea-shore abound in them. I noticed them also in various places accumulated in clusters, forming small eminences of from 10 to 100 yards in diameter, and from 8 to 20 feet high. These may be ice-borne boulders. The usually circular form of the heaps militates against their being glacier moraines. Such collections are of frequent occurrence on the borders of Great Bear Lake, and in the valleys that separate the spurs of the Rocky Mountains, - several hundred feet above the present levels of the lakes and rivers. They also occur in more southern localities.

No. II.

CLIMATOLOGY.

Snow Line.—Ground Ice.—Thermometrical Observations in the Valley of the St. Lawrence.—Comparative Temperature of the Two Sides of the Continent.—Phenchiena of the Seasons at Penetanguishene: At Fort William: At Fort Vancouver.—Thermometrical Observations in the Valley of the Saskatchewan.—On the East and West Sides of the Continent in that Parallel.—Phenomena of the Seasons at Cumberland House: At Carlton House: At Marten's Falls on Albany River.—Thermometrical Observations on the Missinipi and in the same Arallels on the East and West Sides of the Continent.—Thermometrical Observations in the Valleys of the Mackenzic, Yukon, and Pelly.—Progress of the Seasons at Fort Franklin.— Thermometrical Observations on the Arctic Seas.— General Remarks.—Nocturnal Radiation.

THE remarks which follow on the climate of North America have reference especially to the districts through which the Expedition travelled, some general facts being, however, illustrated by observations made in other quarters.

The course of the snow line has always engaged the attention of travellers, who have ascended high mountain ranges, or penetrated into Polar regions. Nowhere on the route of the Expedition is the snow permanent; not even on the summits of that part of the Rocky Mountains which skirts the Mackenzie. Snow may indeed be occasionally found in summer which has drifted into some narrow ravine, or under a high cliff with a northern exposure; but these patches are of small extent, and have no general influence on the temperature of the district in which they occur.

p

lo

th

ce

T

(E

SNOW LINE.

In lat. 65° N., the snow remains continually on the ground from the middle of October till the beginning of May, at which time the soil begins to appear after it has been covered up for 200 days. The thickness of the snowy covering materially affects the depth to which the low winter temperature can be traced into the subsoil. In places of small extent, sheltered from cold winds, and having a good drainage, with a southern aspect, the vernal rays of the sun assist in removing the snow early; while a retentive clay, even in lower and more southern localities, produces a tardy summer and late frosts. Such are the hypogenous or primitive districts, which from their cold climate and ungenial soil have been justly named "barren grounds," and are analogous to the terra damnata of Lapland, described by Linnæus; or the tundras of Arctic Siberia; in such tracts the snow falls early and remains long. The active vegetation of forest lands reacts on the soil; excited by the sun's rays the trees are roused from their winter's sleep, while the soil is still as hard as a rock, and the snow disappears sooner from over their roots than elsewhere. At the Equator the permanent snow line is said to vary from 15,000 feet of altitude to 20,000, to sink to 3,800 on the 60th parallel of latitude, and to be one foot high on the 75th parallel.* The latter assumption does not, however, accord with observations in the northern hemisphere. In no arctic district to which man has yet penetrated, is there a permanent covering of snow through any wide extent of low country. Even at Spitzbergen, only nine degrees from the Pole, there is a summer in which vegetation proceeds, of which we have witnesses in the flora and fauna. The well-fed herds of reindeer, which that hyperborean

* Von Buch states the height of the snow line at the North Cape (Europe) at 2,275 Prussian feet.—Meyen, Geog. of Plants.

р З

Valley) Sides shene :)bserv-1 West ne Sea-'s Falls lissinipi he Conne Macat Fort Seas. —

North hrough facts n other

ed the ountain here on ht; not Mounleed be to some ern exhave no trict in

land maintains, must find grass and lichens, whereon they fatten.*

On the north side of Lake Superior, the duration of the winter's snow is less by at least sixty days than it is in lat. 65° N., that being the difference due to about twenty-seven degrees of latitude. In favourable seasons, at Melville Island, in lat. 743° N. and long. 110° W., the snow towards the end of June lies only in the valleys where it had drifted deeply, and the level meadows remain uncovered for seventy days, or till the beginning of September. In Regent's Inlet, which is more to the eastward, the snow dissolves less rapidly; and at Igloolik, still further east, notwithstanding a difference of 5¹/₄° of southing, the soil is uncovered for a still shorter period, furnishing an illustration of the depression of temperature on the eastern coasts, on which some observations will be made in a subsequent page. The subjoined tables will furnish the dates of the disappearance of the snow in the spring, of various more southern localities.

The opinion held at no distant date by eminent meteorologists, that a mean annual temperature of the freezing point of water coincides with the snow line, has been satisfactorily disproved by observation. Everywhere to the eastward of the Rocky Mountains, the isothermal of 32° F. is below the fifty-seventh parallel of latitude.

When the rigour of the climate of Arctic America is

* The Herald Island, discovered by Captain Kellett in lat. 71° 19' N., long. 175° 23' W., to the north of Beering's Straits, is 1200 feet high, and consists of granite precipices to the height of 900 feet, and then a succession of terraces, on which there grows a turfy vegetation. Eight species of plants were gathered from these terraces, viz. a *Hepatica*, *Poa arctica* and another grass, *Artemisia borealis*, *Cochlearia fenestrata*, *Saxifraga laurentiniana*, a moss and a red lichen, which covered the rocks.

zo lin ha the me " r sub tere wai pro year less vani decr of the of th

limit

the a

the t

p

81

81

a

at

ca

GROUND ICE.

y

of

it

ut

18,

1.,

he

en-

the

is

ly;

g a

for

the

hich

age.

dis-

nore

hete-

ecz-

been

e to

l of

a is

° 19′ feet

and

ation.

atica, trata,

d the

considered, the under limit of permanent snow on the hills appears to be very elevated. The true laws of the altitude of this line have not yet been ascertained. Correct measurements have shown that within the tropics the snow line varies several thousand feet, even in the same range of mountains, according to its different aspects. The active radiation of the sun in the continuous day of an arctic summer, in conjunction with the comparatively small winter deposit, must tend to elevate the snow line; while within the tropics, the effect of a vertical sun is compensated by nocturnal terrestrial radiation, and the deposition that attends the sudden cooling of an atmosphere charged with moisture. The same or similar causes must tend to vary the breadth of the mountain zone, comprehended between the summer and winter snow The east coast climate everywhere north of Canada line. has an analogy to this zone, and not to its upper limit or the permanent snow line.

Another phenomenon intimately connected with the mean temperature of a district, is the "ground ice" or " permanently frozen subsoil." The lateral extent of this substratum, its southern limits, and its thickness, are interesting subjects of inquiry. It is well known that the warmth excited by the sun's rays is conducted slowly and progressively into the earth, the effects of seasons and years following like tides, but becoming less sensible and less distinct until they are blended, and at a certain depth vanish altogether. Professor Forbes says, that "the decrease of the annual range is common to the strata of the air above the surface of the carth and to those of the soil beneath; both ultimately, no doubt, exhibit a limit, first where the diurnal variations disappear, then the annual." The cause, however, he states is different in the two cases; the one being chiefly the result of the

P 4

radiation, and the other of the conduction of heat. The limit here spoken of, or the depth to which the periodical changes of summer and winter are felt, is influenced by a variety of circumstances, and differs in different localities. The temperature of that limit would be, it is supposed, that of the mean of a number of years (fourteen or fifteen perhaps) forming a complete cycle of the annual variations.

The limit or commencement of permanently frozen soil, or ground ice, is coincident, according to Bäer, with the isothermal line of 32° F.; and its thickness increases in proportion as the mean temperature of the locality falls below that degree, its unlimited descent being checked by the interior heat of the earth.

Observations of the temperatures of mines and of artesian wells have established the fact that the temperature rises as we penetrate into the crust of the earth; but the rate of increment has not yet been satisfactorily determined. The temperatures of mines in the same district, and of different parts of the same mine at equal depths, vary greatly. Some authors fix the increment at one degree of Fahrenheit for every forty-five feet of descent, after the superficial stratum which is directly influenced by the solar heat has been passed.

At Yakutzk, in Siberia, on the sixty-second parallel, with a mean annual temperature of 14° F., a well dug to the depth of 382 feet just penetrated the frozen earth, and the resulting increment of heat there is one degree for twenty-eight feet of descent. At Fort Simpson, on the Mackenzie, very nearly in the parallel of Yakutzk, but having a mean annual temperature of 25° F., the frozen substratum was found to terminate at the depth of seventeen feet from the surface, the underlying bed being loose and sandy. The surface soil there was thawed at the close of summer (19th October, 1837), to the depth of nearly

5

d

ir

e

of

W

tił

ca

GROUND ICE.

eleven feet, so that the ground ice was only six feet thick.

At York Factory, on Hudson's Bay, in lat. 57°, in October, 1835, recent frosts had penetrated eight inches into the soil; the thaw due to the summer heat extended twenty-eight inches beyond this, beneath which a frozen bed seventeen and a half feet thick reposed on thawed mud which had a temperature of 33° F. The mean annual heat of this place is $25\frac{1}{2}$ F., being equal to that of Fort Simpson, which lies five degrees further north.

At Severn outpost, exactly one degree of latitude to the South of York Factory, and on the same coast of Hudson's Bay, the surface had thawed at the close of the summer of 1835 nearly to the depth of six feet, and the frozen substratum was dug into seven and a quarter feet, being thirteen feet in all, but not passed through, so that its thickness was not ascertained.

I have no information respecting the ground ice of the Peace River or Saskatchewan prairie districts.

At Rupert's House, on James's Bay, near the level of the sea, in lat. $51^{\circ} 26'$ N., long. $78^{\circ} 40'$ W., the soil in an exposed situation in the month of April was frozen to the depth of seven feet; but under a snow-drift the frost had penetrated only thirteen inches into the earth. At East Main, situated on the opposite side of the same bay, in lat. $52^{\circ} 15'$ N., long. $78^{\circ} 40'$ W., the ground under a snowdrift eight feet thick was frozen to the depth of only ten inches. The pits at these places were not dug deep enough to give any information respecting the existence of a permanently frozen substratum.

On Michiskum Lake, which lies in lat. 49° N., long. 78° W., on the north side of the watershed that divides the Abitibbe, or eastern branch of Moose River, from Lake Temiscaming and the Ottawa, and which is about 700 feet

The odical by a lities. posed, ifteen ations. n soil, th the uses in cy falls hecked

of arteerature but the deterdistrict, depths, one dedescent, fuenced

parallel, ell dug n earth, gree for on the zk, but frozen f sevenng loose he close ' nearly

above the sea, the frost towards the close of winter had penetrated three and a half feet in a cultivated field, while in the woods not above six inches of soil were frozen. No search was made for a frozen substratum. These are the chief observations * I have been able to collect, having reference to this subject. It is plain that much more extensive researches are required to enable us to form any general conclusions which can be relied on.

Professor Bäer informs us that in Europe and Siberia, the farther we go east, the more southerly do we find the limit of perpetual ground ice to be. In the environs of Lake Baikal, ice remains at least in one locality all the year. No permanent ice was found at Tobolsk in lat. 58° N., but Humboldt discovered small pieces at the depth of six feet in summer in the elevated district of Boguslowsk, near the foot of the Ural Mountains, in lat. 59° 45' N. It would appear from these instance: that the ground ice has a more southerly limit on the shores of Hudson's Bay than in Siberia; and that in America, as far as observations go, that limit follows the course of the isothermal lines which dip to the south as they proceed to the eastward.

t

t

W

W

0

se

of

tr co

co

of ap

im

alti

equ

If

tha

on

THERMOMETRICAL OBSERVATIONS IN THE VALLEY OF THE ST. LAWRENCE.

The subsequent tables of temperature and notices of the progress of the seasons are arranged so as to convey, as far as my materials go, a view of the climates of the successive lake basins from the St. Lawrence northwards. *Table II.* is intended also to illustrate the fact that the climate, on the west side of the continent, is milder than

* All the experiments made in Rupert's Land, in 1835-36, are detailed in the Ed. New Philos. Journ. for January 1841.

VALLEY OF THE ST. LAWRENCE. 219

in the eastern states. Fort Vancouver being upwards of forty miles from the coast in a direct line, its climate cannot be called maritime, and its temperatures may be conveniently contrasted with those of Montreal, which is also situated on the banks of a first-rate river, and not under the influence of sea-breezes.

There is very little difference in the latitudes of the two places, or in their altitudes above the sea; but the direct distance between them exceeds 2,200 geographical miles. The difference of their mean annual temperatures is, at least, eight degrees of Fahrenheit in favour of the Pacific coast, and is equal to what would be produced by a diminution of four or five degrees of latitude on the same meridian. The greater mean heat of the Pacific side of the continent has been long known, and a reference to the table will show that it is mainly due to the milder winters; the mean difference between the summer and winter temperatures being twice as great at Montreal as on the Oregon. On the other hand, the summers are sensibly warmer at Montreal. The notices of the progress of the seasons on the two sides of the continent will illustrate these facts equally well. The early spring at Vancouver, with the two inches of snow and a rainy winter, contrasts strongly with the long dry winter and three feet of snow at Penetanguishene, where the ice does not disappear from the lake till past the middle of April.

Columns 1. and 2. of the same table refer to places immediately on the coast, and Franklin Malone is at an altitude above Fort Vancouver, which is considered to be equivalent to two degrees of mean annual temperature. If we compare these columns with column 4., we perceive that with little change of mean annual temperature, places on the coast have a more equable climate, the three winter

had while No e the aving e exa any

beria, id the ons of ill the at. 58° pth of lowsk, 45' N. ind ice udson's obserhermal e east-

EY OF

ices of onvey, of the wards. hat the er than

-36, are

months being comparatively milder, and the three summer ones less warm.

Table I. shows that as we advance into the interior, the heat of the summer is kept up, or even augments, notwithstanding the elevation above the sea attained by ascending the successive stages of the St. Lawrence basin.

The nature of the rock formations has a considerable influence on the elimate of a district. In the primitive country, such as has been described in a preceding page as abounding in lakes and swamps, the climate is extreme, the winters being not only longer, but also more severe, the dissolution of the ice in such districts absorbing much A marked difference occurs when we pass from the heat. "intermediate primitive range" to the prairie districts lying to the westward; for, notwithstanding the greater elevation of the latter, the winters are milder, the snow less deep and less durable, the rivers break up earlier, and the sap flows sooner in the trees. A corresponding difference in the vegetation occurs; the prairie plants have much less of an arctic aspect than those of the primitive districts. Professor Agassiz, in his work on Lake Superior, has instituted a very interesting comparison between the vegetation of that basin and the lower and middle subalpine zones of the higher tracts of the Jura, proving their very great similarity. With the prairie districts, the analogy of the Jura is very much less strong. Many of the plants which give the peculiar character to the prairies south of the Missouri, range northwards to the branches of the Peace River, a main affluent of the Mackenzie; and several prairie plants enter the silurian wooded tracts which lie to the westward of the "intermediate primitive range," though they have not been discovered in the more eastern parts of the continent.

Januar Februa March April May June July August Septemil October Novemit Decemb Year

Winter Spring Summer Autumn

Difference hottest coldest Difference summe winter

> sei lin Re

 $\mathbf{220}$

TEMPERATURE OF VARIOUS PLACES.

TABLE I.

TEMPERATURE OF VARIOUS PLACES IN LAKES ONTARIO, ERIE, HURON, AND MICHIGAN.

	Lake Ontario, North Side Toronto. Lat. 43° 40' N. Long. 79° 22' W. Alt. 340 Feet.	Lake Huron, East Side Penetangulsinene. Lat. 44° 48' N. Long. 80° 40' W. Alt. 626 Feet.	Lake Michigan. Green Bay, Fort Howard. Lat. 44° 40' N. Long. 87° 00' W. Alt. 600 Feet.	Lake Erie, East End Buffalo. Lat. 42° 55′ N. Long. 78° 55′ W. Alt. 575 Feet.	Lake Erie and R. S. Clair, Detroit. Lat. 42° 19' N. Long. 82° 58' W. Alt. 575 Feet.
Periods.	Hourly Obs. ior 5 Years at Magn. Obs.	8 and 8. by C. Todd, Surgeon R.N., corrected by Dove's Form.	7, 2, 9, corrected by Dove's Form.	Computed by the Form. N.Y. Dove. 2 Yrs. Obs.	8 and 2, 31 Years, corrected by Dove's Form.
	(1.)	(2.)	(3.)	(4.)	(5.)
January February	25·34 23·71	23·42 22·60	17·76 20·05	23·41 21·14	20·26 31·32
March	30.37	31.90	30.87	35.49	39.88
April	40.00	55.36	56.57	55.29	57.07
Juno	60.88	67.99	67.86	67.45	68.02
July	66.28	73.31	71.71	71.55	70.83
Anoust	66.73	69.40	68.67	69.99	68.87
Sentember -	58.71	56.03	56.91	59.89	61.68
October	45.69	50.11	46.74	48.75	54.28
November -	36.03	38.86	33.97	37.22	38.02
December -	27.24	25.24	20.95	27.80	29.32
Year	44.81	45.74	44.57	45.54	49.10
Winter	25.43	23.75	19.59	24.12	26.97
Spring	42.34	35.02	43.43	43.83	48.87
Summer	64.43	70.24	69.41	69.66	69.24
Autumn	46.81	48.34	45.87	48.62	51.34
Difference of hottest and coldest months	48.81	49.99	53.95	50 [.] 41	50 [.] 57
summer and winter -	39.20	46.49	49.82	45.54	42.27

No. 1. is taken from the volume of observations made at the observatory at Toronto, published by Government. No. 2. from Franklin's Second Journey. Nos. 3., 4., and 5. from Dove's tables in the Report of the British Association for 1847.

ımmer

iterior, ments, ned by basin. derable rimitive ng page xtreme, severe, g much from the districts greater he snow earlier, sponding nts have primitive ke Supebetween ldle subproving ricts, the Many of e prairies branches zie; and ed tracts orimitive the more

TABLE

0 y si w di

3

	EAST SIDE.						
	Atlantic	c Coast.	St. Lawrence Valley.				
Periods.	Mance, Eastport, Lat. 44° 54' N. Long. 66° 56' W. Alt. 15 Feet.	Nova Scotia, Halifax. Lat. 44° 39' N. Long. 63° 38' W. Alt. 15? Feet.	North N. York, Fr. Malone, Lat, 44 ⁰ 50' N. Long, 74 ⁰ 23' W. Ait, 645 Feet.	Canada, East, Montreal, Lat. 45 ^o 31' N. Long. 73 ^o 35' W. Ait. 60 Feet.			
	(1.)*	(2.)	(3.)	(4.)			
January	19.18	20.00	28.24	13.03			
February	22.71	18.00	26.12	17.68			
March	29.85	25.00	31.42	30.89			
April	38.72	30.00	45.08	46.80			
May	48.99	40.00	52.67	53.97			
June	56.31	50.00	60.22	59.73			
July	63.28	63·00	66.90	68.90			
August	63.24	70.00	65·45	68·04			
September -	57.15	51.00	55.17	57.61			
October	46.33	51.00	46.92	46.20			
November -	36 18	38.00	32.85	31.28			
December -	24.55	25.00	21.22	19.66			
Year	42.41	40.08	43.52	42.12			
3 Winter months	22.15	21.00	21.87	16.95			
3 Spring months	39.19	31.67	43.06	39.03			
3 Summer months	60.94	61.00	64.19	67.26			
3 Autumn months	46.55	46.67	44.98	45.18			
Difference of hottest and coldest months	} 44.10	52 ·00	48.66	55.87			
Difference of summer and winter -	38.79	40.00	42.32	50·31			

COMPARATIVE TABLE OF TEMPERATURES ON THE EAST AND

* Columns 1., 2., and 3. are extracted from Dove's paper, published in the Report of the British Association for 1847, being corrected for the diurnal variation computed for Toronto by Dove, in the same paper. No. 4. is abstracted from a paper by J. S. M'Cord, Esquire, printed at Montreal, and discussing observations of temperature for 1839 and 1840, and 1840-41, 12 times daily at the even hours. Column 5. is also from Dove's paper, and is reduced by his formula. Columns 6. and 8. ner abstracted from a year's observations, by G. B. Roberts, Esquire, of the Hudsoo's Bay Company, with a thermometer of Newman's construction. Column 6. is reduced for the hours of observation, according to Dove's formula for Toronto. Column 8. is transcribed without correction. Column 7. is an abstract of observations by the Rev. S. Parker, also corrected by Dove's formula. The subjoined is an abstract of observations for seven years, made by Harry Poole, Eaquire, at Albion Mines, near Pictou, in Nova Scotia, and on the coast of the Gulph of st. Lawrence. Mr. Poole ascertained the minima in the night by a self-registering ther-mumeter, and the maxima between 1 and 2 P. M., by a thermometer hung on the north side

COMPARATIVE TABLE OF TEMPERATURES. 223

0

ABLE

ST AND

cy.	
ada, East, ontreal, 45° 31' N. . 73° 35' W. . 60 Fcet.	
(4.)	
13.03 17.68 30.89 46.80 53.97 59.73 68.90 68.04 57.61 46.50 31.58 19.66	
42·1 2	
16-95 39-03 67-26 45-18	
55.87	
50.31	

eport of the for Toronto rd, Esquire, and 1840-41,

ts, Esquire, Column 6. r Toronto. rvations by

enry Poole, he Gulph of ering there north side

Π.

WEST SIDES OF THE CONTINENT NEAR THE 45TH PARALLEL.

•				
	Oregon	State.		-
Fort Van	Periods.			
11ours 7, 2, 9.	Hours 8 and 8.	Hours 7-1.	Hour 2 p.m.	
(5.)	(6.)	(7.)	(8.)	-
37.62	36.58	37.10	40.68	Jannary.
42.89	35.37	38.98	44.07	February.
43.68	44.05	47.47	55.06	March.
45.55	50.22		62.50	April.
53.44	58.43		74.61	May.
62.48	58.72		73.17	June.
65.46	61.76		75.71	July.
65.84	63.02		79.74	August.
60.30	61.10		74.27	September,
53.23	52.62	48.27	68.16	October.
42.68	38.83	39.23	48.27	November,
42.95	34.49	38.60	42.06	December.
51.34	49.73		61.91	Year.
41.15	35.49	38.23	42.21	3 Winter months.
47.55	50.90		64.07	3 Spring months.
64.67	61.30		76.14	3 Summer months
52.00	50.92		64 71	3 Autumn months
28.28	28.56		39.06	Difference of hottest and coldest months
23.52	25.81		33.93	Summer and winter.

of the house, and sheltered from the north winds by a plantation of firs at the distance of a few yards. The observations are corrected for the Toronto diurnal variation for an hour before suntise, and 2 P.M. As there appears to have been some local cause, producing higher winter temperatures than in other parts of the coast, I did not think it expedient to introduce these observations into the comparative table.

Abstract of Temperatures at Albion Mines, 120 Feet above the Sea, Lat. 45° 344' N., Long. 62° 42' W. Means of Maxima.

Ì	Jan.	50.58 F.	March	52·53	June	68·86	Sept.	62.42	Year	-		56·03
	Dec.	51-11	May	58.64	Angust	70°61	Nov.	49 98	Diff. h	ot and co	old mths.	20.66
	Wint.	51.18	Spring	53-23	Summer	69.04	Autumn	55.83	Diff. o	f sum. a	nd wint.	17.86

TABLE III.

FABLE	OF	TEMPERATURES	ON	LAKE	SUPERIOR,	AND	ON	THE
		UPPER PART O	F TH	IE OTT.	AWA RIVER			

		I Transienses			
Poleir	Fort Brady, Saut St. Mary. Lat. 46° 31' N. Long. 84° 25' W. Alt. 660 feet.	North-east angle, Michipicoten. Lat. 47° 56' N. Long. 85°(6' W. Alt. 660 feet.	Thunder Bay, Fort William. Lat. 48° 234' N. Long. 89° 22' W. Alt. 660 feet.	H. Bay Post, on the Ottawa. Lat. 470 19' N. Long. 790 31' W Ait. 630 feet.	
Perious.	7. 2. 9. Dove corrected.	8 and 8, by G. Kelth. Esq , corrected by Dove's form.	8 and 8, corrected by Dove's form,	Sunrise, sunset, and noen, by - Severight, Esq., corrected by Dove's form.	
	(1.)	(2.)	(3.)	(4.)	
January	⊥18.30	+10.63	+5.70	+0.93	
February -	19.69	16.66	8.22	18.44	
March	27.04	26.09	22.72	24.41	
April	38.05	34.66	31.42	39.04	
May	52.00	51.88	48.87	49.35	
June -	58.61	55.00	58.73	62.75	
July	65.36	57.03	62.19	67 28	
August	64.36	60.04	58.84	65.58	
September -	55.55	49.67	48.16	53.39	
October	44.75	44.92	41.88	40.83	
November -	33.29	29·01	23.43	25.97	
December -	22.23	22.38	18.16	17.68	
Year	41.63	38.25	35.90	38.58	
3 Winter months	20.07	16.68	10.75	15.02	
3 Spring months	39.04	37.57	39.67	37.58	
3 Summer months	62.77	57.39	59.94	65.23	
3 Autumn months	44.63	41.24	37.80	40.02	
Difference of hottest and coldest months Difference of	46.06	49:41	56.49	58.05	
summer and winter months	42.70	40.71	49.19	50.21	

Column 1. is extracted from Dove's paper, already referred to; columns 2 and 3. are compiled from documents obtained from the Hudson's Bay Company; and column 4. is extracted from Mr. Logan's Geological Report for 1845-6.

224

vo

PROGRESS OF THE SEASONS AT MICHIPICOTE. 225

Phenomena illustrative of the Climate of Penetanguishene, on Lake Huron.*

The spring sets in very suddenly. Snow continues until the latter end of April, and longer in the forest than in cleared lands. The weather in March is clear and cloudless, and the ice, which in winter has attained a thickness of sixteen inches, begins to dissolve. Towards the end of the month the sap of the maple flows, and the sugar harvest commences. Flocks of Canada geese and various ducks appear about the same time, and are the harbingers of fine weather. The ice disappears on an average on the 24th of April. Alders and various willows flower about the middle of the month, and the Hepatica triloba blossoms on the 25th. Potatoes are planted between the 1st and 20th of May, and encumbers and melons are usually sown between the 25th and the end of the month. Viola blanda, Xylosteum, Leontice, Erythronium, and many other plants, blossom in this month; the forest trees come all into leaf about the 16th, and, about the 19th, the moschetoes begin to be troublesome.

In the month of June the temperature rises to 90° F. in the day, and heavy dews fall in the night. Barley and oats are sown about the 15th, and towards the end of the month garden peas are fully podded, and the male flowers of maize spring up. The *Lilium philadelphicum* blossoms at this time.

In July and August the weather is usually dry and sultry. About the beginning of the former month *Penstemon pubes*cens, *Rhus typhinum*, garden melons, and cucumbers blossom; and, towards the middle of August, melons grown without artificial warmth are ripe, and the wheat and oat harvest commences. Maize is fit for pulling about the end of the month.

In September numerous flocks of *Turdus migratorius* and other birds arrive from the north, and remain for a time feeding on the berries of various rasp bushes. Maize ripens about the first of the month, and near its end frost destroys the

> * By the late C. C. Todd, Esq., Surgeon, R. N. 11. Q

VOL. 11.

THE

iscaming.

y Post, Ottawa 70 19' 90 31' W. 530 feet. se, sunset, noen, ght, Esq., ected by e's form. (4.) -9.2318.44 24.41 39.0449.35 62.75 67 28 65.58 53.39 40.83 25.97 17.68

> 37·58 65·23 40·07

38.58

15.02

58·05

50.21

red to; rom the Logan's

cucumber and melon vines; forest trees change their hue, and potatoes are dug and stored for winter use.

The forest assumes a variety of autumnal hues in the beginning of October; about the middle of the month, many flocks of geene and ducks pass to the southward, and their appearance precedes a series of cold weather, which strips the leaves from the trees. A fall of snow usually occurs about the 25th.

November is generally calm and pleasant, and about three weeks of peculiar weather, named the "Indian summer," occurs. It is characterised by a fog or haze rising from the earth, through which the sun is scen obscurely, and there is little or no wind. In December, the thermometer sinks a few degrees below zero, and much snow falls. The harbour freezes in the beginning of the month. In January, the thermometer sinks 20 degrees below zero, and rarely to -32° . The snow attains a depth of three feet in the woods, but the ground it covers is not frozen. A great fall of snow takes place in February, and there is usually a temporary thaw about the end of the month, accompanied by heavy rain, and occasionally by thunder.

Phenomena indicating the Progress of the Seasons at Michipicoten, Lake Superior*, in the year 1840.

Jan. 19. Open water in the bay. Mergansers frequenting it. Feb. 14. Bay again closed by ice.

- March 8. Snow birds departed for the north. 26th. The snow birds returned again, the weather having been severe. Domestic hens began to lay eggs. Two ducks seen.
- April 10. Lake clear of ice in Michipicoten Bay. 12th. Turdus migratorius came.
- May 2. All the snow gone. 6th. Swallows came. 22nd. Potatoes planted.
- Sept. 4. Small trout collecting in the rivulets to spawn. On

* By chief factor George Keith, Esq.

PROGRESS OF THE SEASONS AT FORT WILLIAM. 227

the 23rd they ceased spawning. Frosty nights. Potatoes not hurt. 26th. Corregonus lucidus spawning in the river.

- Nov. 10. Ice beginning to drift in the river. 18th. The large trout and *Corregonus albus* cease to spawn here at this time, though they carry on this operation later in other parts of the lake.
- Dec. 3. Ice broken up in the river. Little snow lying on the ground. 13th. River again frozen over. Season mild.

Phenomena indicating the Progress of the Seasons at Fort William, Lake Superior, in the year 1840.

Feb. 29. Thermometer at noon rose to 39° F.

- March 1. Temperature 61° in the middle of the day. On the 27th a grey hawk, and on the 31st a barking erow, Corvus americanus, were seen.
- April 2. The sap of the sugar maple began to run. On the 4th small holes began to perforate the ice. On the 9th the first wild ducks of the season came, and on the 10th butterflies, blue flies, and gulls were noticed. 2toh. The general thaw commences at this period. Ground frozen to the depth of 3 feet 9 inches. 21st. Anser canadensis, Anos boschas, and mergansers frequenting the neighbourhood. 28th. Heard a nightingale (Turdus?). 30th. River partially open.
- May 2. River free of ice. Bay of the lake full of drift
 ice. 6th. Anser hyperboreus passing in flocks.
 8th. Moschetoes seen. 10th. The birch tree and
 maple budding.
- June 15. Swallows building in the outhouses. 17th. Sturgeons spawning in the rapids of the river. 19th. *Catastomi* beginning to descend the river from the rapids. 21st. *Corregonus lucidus* comes to the entrance of the river in shoals.

Q 2

and

e bemany ir aps the about

three r," ocearth, ittle or legrces in the r sinks ttains a overs is ary, and month, r.

ns at 40.

ting it.

he snow ng been s. Two

Turdus

22nd.

wn. On

- July 3. The Corregoni have left the mouth of the river. 15th. Barley just coming into ear. Potatoes in flower. The Lepus americanus having its second litter of young. 31st. Raspberries ripening.
- Aug. 8. Red currants and blue berries (Vaccinium) perfectly ripe. 10th. Reindeer begin to rut. 19th. Barley ripening. 29th. Peas quite ripe. 31st. The swallows have disappeared.
- Sept. 2. Reindeer rutting senson ends. On the 7th the leaves of the birch and aspen change colour. 10th. Small trout begin to spawn. 13th. Potatoes, cabbages, turnips, and cauliflowers nipped by the frost. 14th. A few ducks arriving from the north. 16th. The first stock-ducks arrived from the north this autumn. 20th. Small trou. spawning abundantly on the shoals. 23rd. The orioles have departed for the south. Canada geese arriving from the north, and going southwards. 30th. Corregonus lucidus begins to spawn in the rapids of the river.
- Oct. 6. The large trout begin to spawn in the lake at the Shaguinah Islands; they cease on the 18th. Thunder. 7th. Leaves of the birch and aspen falling. 10th. The Corregonus lucidus has ceased spawning on the rapids. 14th. Thunder. Anser hyperboreus arriving from the north. 15th. Passing in large flocks. 20th. Hail, thunder, and lightning. Plovers, divers, snipes, orioles, geese, and ducks in the neighbourhood. On the 31st, snow-birds began to arrive from the north.
- Nov. 3. The small lakes frozen over. On the 9th, the river (Kamenistikwoya) covered by a sheet of ice, which broke up again. 21st. The spawning season of the Corregonus albus terminates.
- Dec. 1. Ice driving about in the lake with the wind. On the 17th, the bay was frozen across to the Welcome Islands.

PROGRESS OF THE SEASONS AT FORT VANCOUVER. 229

The following Notices of the Progress of the Seasons at Fort Vancouver in 1838*, may be contrasted with the state of vegetation at the same times of the year in the valley of the St. Lawrence : —

- Jan. 2d. Short young grass, affording good pasturage for sheep in places that were flooded in summer. 8th. Berry-bearing bushes budding, such as the wild cherry, black currant, &c. Swans abundant, ducks, geese, and cranes scarce. 10th-12th. Snow fell to the depth of an inch and a half, and vegetation was retarded by unfavourable weather till
- Feb. 17. when wild gooseberry bushes were observed budding. Between the 26th of this month and
- Mar. 16. Thunder and hail showers occurred. Ribes sanguineum blossoming. Trillium grandiflorum, having the local name in Oregon of "Herb Paris," in full flower. 21st. Apple and pear-trees budding. The wild gooseberry in full leaf, and further advanced than bushes from England cultivated in the garden. 24th. The swallow first noticed. 30th. Hummingbirds appearing ; strawberries flowering.
- April 3. Mahonia in blossom, and, on the 5th, peach-trees flowering. 8th. Potatoes that have lain in the ground all winter, beginning to show. 11th. Dogwood and elder in blossom. 17th. Several species of violet in flower. 20th. Field iris in flower. 23d. Brambles flowering. On the 25th, clover in bloom; and, on the 26th, wild tares flowered. Hail and thunder storms. 28th. Blossoms of the fruit-trees falling.
- May 1. Lupins in flower. 7th. Wild rose and eglantine in flower. 12th. Strawberries ripening. 28th. Fieldpeas in blossom. 30th. Garden-peas brought to table. Thunder.
 - * By G. B. Roberts, Esq., of the Hudson's Bay Company.

Q 3

river. es in econd

fectly Barley swal-

leaves Small bbages, 14th. th. The this auindantly leparted rom the rregonus e river. e at the . Thunh falling. pawning perboreus in large Plovers, s in the began to

> the river ce, which season of

ind. On Welcome

- June 1. Spring barley in ear. 5th. New potatoes raised from tubers, left all winter in the ground, fit for table.
 7th. Oats in flower. 10th. Spring wheat coming into ear. 17th. Bramble-berries ripening. 23rd. Gooseberries, currants, and raspberries ripe. Thunder. 26th. Blackberries (Vaccinium) ripe.
- July 4. Humming-birds scarce this season. 19th. Barley fit for the scythe. 22nd. Winter wheat ripe. 28th. House-flies numerous.
- Aug. 3. Oats ripe. 7th. Field-peas harvested. 9th. Foggy mornings, followed by a clear sky and excessive heat. Dewless mornings succeeded by rain. 12th. Salmon season usually ends. Several cases of ague occurring. 27th. Moschetoes very troublesome. Geese arriving from the north.
- Sept. 7. Hail and thunder. 12th. Buckwheat harvested. 13th. Ague prevalent. 19th. Mowed barley that was sowed on the 16th of June—a fair crop. 22nd. Peas sown on the 19th of June, on alluvial land, that was flooded in winter, now ripe.
- Oct. 27. The Columbia or Oregon river unusually low, the greatest depth opposite the fort being less than fifteen feet.
- Nov. 8. Potatoes killed by the frost. 18th. A little drift ice in the river. 20th. Rain and sleet.
- Dec. 26. Snow at this time two inches deep, being the greatest fall this season, and less than usual.

TEMPERATURES IN SASKATCHEWAN VALLEY. 231

TABLE IV.

THERMOMETRICAL OBSERVATIONS. — VALLEY OF THE SASKAT-CHEWAN.

Months and Seasons.	Cumberland House. Lat. 53° 57' N. Long. 102° 20' W. A 900 Feet. Mean Temp., 8 A. M., 8 P. M., corrected, 1839.	Oxford House. Lat. 54° 55' N. Long. 96° 28' W. Alt. 400 Feet? Mean Temp., 7 A. M., Noon, and 8 P. M., corrected, 1833.	York Factory. Lat. 57° 0' N. Long. 92° 26' W. Alt. 20 Feet. Mean Temp., Morn., Noon, and Evening, corrected, 1830.	Rupert House. Lat. 51° 21' N. Long. 83° 40' W. Alt. 20 Feet ? Mean Temp. corrected, Sunrise, 13 P. M., Sunset, 1839.
June			47.67	
July			59.99	
August	62.84		54.85	
September -	44.20		41.90	
October	33.12	17.53	33.43	34.80
November -	21.48	13.29	25.17	23.33
December	7.94	-2306	3.73	15.59
January	-0.89	-22.06	-5.12	-4.09
February	-8.06	- 1.90	-6.60	-0.68
March -	18.30	8.57	4.77	7.64
April	27.01	28.62	19.21	21.05
May	52.59	38.01	33.23	41.51
Year	33.20		25.63	
Summer	62.62		52.07	
Autumn	33.04		33.20	
Winter	-0.12	-0.85	-2.23	0.14
Spring	32.70	7.51	19.17	0.78
Difference of hottest and coldest months	72.06		66•59	
Difference of summer and winter -	61•79		54 .60	

The observations recorded in the above table were made at Cumberland House, by chief factor John Lee Lewis, between the beginning of August 1839 and the end of September 1840, the register being deficient in June and July, when Mr. Lewis was travelling on the business of the company. The temperatures were measured by a thermometer made by Newman, and sent out at that time with many others, by the Geographical Society, but at the expense of the Hudson's Bay Company. The monthly means of the combined temperatures at 8 A. M. and at 8 p. M. were corrected by Dove's table, calculated for Toronto.—*Rep. Br. Ass.* 1847.

The observations at Oxford House were made between October

Q 4

rom ble. aing 3rd. hun-

ey fit 28th.

Foggy essive 12th. f ague esome.

vested. ey that 22nd. al land,

ow, the ss than

le drift

greatest
TABLE V.

Periods.	Nain. Lat. 57° 10' N. Long. 61° 50' W. Alt. 20 Feet. 3 Years. 8. 12. 4. 8. Dove.	Okak. Lat. 57° 30'. Long. 66°. Alt. 20 Feet. 2 Years. 8. 12. 4. 8. Dove.	Sitka, Lat. 57° 3' N. Long. 135° 18' W. Alt. 20 Feet. 24 Years. Reduced. Bäer.
June	42.53	44.65	53.83
July	50.18	51.65	57.11
August	50.99	52.00	57.79
September	44.98	44.45	55.96
October	33.98	31.15	46.63
November	26.51	22.40	42.89
December	6.51	8.45	36.35
January	0.95	2.15	34.18
February	3.51	1.95	33.60
March	7.52	8.25	38.01
April	29.97	29.00	40.64
May	36.23	38.25	48.18
Year	27.82	27.86	45.44
Summer	47.90	49.43	56.24
Autumn	35.16	32.67	48.49
Winter	3.66	4.18	34.74
Spring	24.57	25.17	42.28
Difference of hot- test and coldest months}	50.04	50.05	24.19
Difference of sum- mer and winter }	44 ·24	45.25	21.50

COMPARATIVE TABLE OF TEMPERATURE ON EAST AND WEST SIDES OF THE CONTINENT.

1833 to May 1834, at 7 A.M., noon, and 8 P.M., and these are also corrected by Dove's formula for Toronto.

The register for York Factory, Hudson's Bay, was kept by chief factor Joseph Charles, from June 1830 to end of May 1831, the temperatures being recorded in the morning, noon, and afternoon, but the exact hours of the morning and evening are not specified. They have been corrected by Dove's table, on the supposition that the hours were sunrise and sunset.

The observations at Rupert House, on the east side of James's Bay, were made at sunrise, an hour and a half after noon, and at sunset, and Dove's corrections have also been applied to them.

An examination of the several columns will show clearly the effect of open water in Hudson and James's Bays, in tempering the atmosphere in the months of October, November, and even December, and of the presence of ice in those seas, keeping down the summer heat.

EDMONTON HOUSE.-CARLTON HOUSE.

233

İ

TABLE VI.

Abstract of a Record of Temperatures of the Air in the Shade, kept by Mr. Drummond, at Edmonton House, Lat. 53° 40' N., Long. 113° W., Alt. 1,800 Feet.

		Monthly Means	Extreme Temperatures.		
Months.	Of Maxima. Of Minima.		Of these.	Highest,	Lowest.
1827. Jan Feb	18·68 29·96	3·42 3·68	11·05 14·32	+42·0 +47·0	27·0 25·0

TABLE VII.

Abstract of a Journal of Temperatures kept at Carlton House by the Author, Lat. 52° 51'N., Long. 106° 13 W., Alt. 1,100 Feet.

Months.		Monthly Means	Extreme Temperatures.		
	Of Maxima.	Of Minima.	Of these.	Highest.	Lowest.
Feb., last	} 12.50	-1.20	5.65	31.0	-29.0
March - April - May -	23·10 40·97 61·90	+0·74 18·53 33·95	11·92 29·75 47·92	42:0 59:0 75:0	-26.0 + 2.0 +22.0

EST

N. s' W. et.

•

494

8

9

60

are also

by chief ne temon, but They e hours James's at sun-

> e effect e atmober, and heat.

TABLE VIII.

TABLE OF THE EXTREME TEMPERATURES OCCURRING IN EACH MONTH AT SEVERAL PLACES IN THE SASKATCHEWAN VALLEY, AND IN HUDSON'S AND JAMES'S BAYS.

	Cumberland House. Lat. 53° 57' N. Long. 102° 20' W. Alt. 900 Feet.		Oxford House. Lat. 54° 55' N. Long. 96° 28' W. Alt. 400 Feet.		York Factory. Lat. 57° 00' N. Long. 92° 26' W. Alt. 20 Feet.		Rupert House. Lat. 51° 21' N. Long. 83° 40' W. Alt. 26 Feet.	
	Highest Temp.	Lowest Temp.	Highest Temp.	Lowest Temp.	Highest Temp.	Lowest Temp.	Highest Temp.	Lowest Temp.
June -	87	42	-	-	83	25		
July -	98	47	-	_	96	35	1	
August -	90	49	-	-	79	40		
September	73	30	-	-	70	28		
October -	68	5	49	- 3	58	19	61	13
November	38	-10	39	- 6	42	8	36	-10
December	25	-26	32	-29	30	-30	37	- 5
January -	25	-32	0	-44	19	-34	16	32
February	34	-35	30	33	37	35	39	-36
March -	50	- 9	34	-27	47	-25	34	-25
April -	55	- 3	64	- 8	55	- 9	53	-16
May -	93	27	65	15	62	15	71	19

This table shows that tropical temperatures occur in the Saskatchewan basin, for a day or two, or it may be only for a few hours at a time in summer, yet that the three summer months seldom pass without night frosts. These destroy tender plants, and in untoward seasons injure the growth of *cerealia*. Wheat, however, ripens well in the drier limestone districts, and still better in the prairie country; but it is there subject to periodical ravages of the larvæ of caterpillars, which come like an army of locusts, and eat up all that is green. Were the country more generally cultivated, and rooks and domestic poultry encouraged, this plague might be lessened. Maize ripens well at the Red River and Carlton, and, I believe, at Cumberland House also, though I did not see it in cultivation there. The 54th parallel may be considered as its northern limit, but its profitable culture does not, perhaps, extend beyond the 49th or 50th degree on the east side of the Rocky Mountains.

PROGRESS OF SEASONS AT CUMBERLAND HOUSE. 235

In the following Table of Phenomena, indicating the Progress of the Seasons at Cumberland House, I have combined my own observations in the spring of 1820 with those of chief factor John Lee Lewis, in 1839 and 1840, distinguishing the remarks by the years. The supposed altitude of Cumberland House above the sea is 900 feet, according to Captain Lefroy's calculations.

- March 4. Water collecting in pools round the establishment. 1840.
 - 7. Much bare ground visible.

N EACH

VALLEY,

rt House

51° 21' N. 83° 40' W. 26 Feet,

> Lowest Temp.

> > 13

-10

 $-5 \\ -32$

---36

 $-25 \\ -16$

Saskat-

hours at

lom pass

intoward

is well in country ;

erpillars,

is green.

domestic

e ripens

nberland The 54th

rofitable

egree on

- 8. The snow which covered the ground t the depth of three feet, was observed to moisten in the sun for the first time this season. 1820.
- 12. Temperature in the shade rose for the first time to $+ 30^{\circ}$ F. The melting snow began to drop from the eaves of the houses.
- 21. Patches of earth became visible, the season being in respect to the melting of the snow fourteen days later than that of 1840. The river Saskatchewan broke up partially, the melting snow covered with *Poduræ*, as it is also frequently in the autumn.
- 24. A white-headed eagle was seen, this being almost always the first of the summer birds which arrives; it comes as soon as it can obtain fish. In 1840, the first eagle was seen on the 26th.
- April 2. The river Saskatchewan froze over again, after some very cold days.
 - 7. Barking crows (Corvus americanus) seen. They were not observed till the 19th in 1840.
 - 8. First snow bunting seen (Emberiza nivalis). 1840.
 - 9. A merganser seen. 1820.
 - 10. Willow catkins beginning to burst.
 - 12. Geese and swans seen in 1820. In 1840 they were not seen till the 20th; and pelicans and ducks were observed that year on the 21st.

April 13. Buds of Populus balsamifera bursting. 1820.

- 17. Plovers, grakles, and orioles seen, and, on the following day, Canadian jays and fly-catchers. Frogs croaking.
- 20. Coltsfoot, Nardosmia palmata flowering.
- 26. Alder flowering. The sugar harvest, which is collected in this district from the Negundo fraxinifolium, commenced in 1820, on the 20th of this month, and lasted till the 10th of May. The flow of the sap is greatly influenced by the direct action of the sun, and is greatest when a smart night's frost is succeeded by a warm sun-shining day. The flow ceases in a cold night.
- 28. The Saskatchewan thoroughly broken up. The ice on Pine Island Lake did not disappear until nearly a month afterwards. Wahlenberg observes that the mean temperature of the air in Lapland must rise to 40° F. before the rivers are completely free. The Saskatchewan opens in this district before the mean heat for ten days rises so high; but its upper part flows from a more southerly and warmer, though a more elevated, country.
- 30. Commenced ploughing 1840.
- May 1. Anemon's patens, or wind flower, in blossom, its leaves not yet expanded. 1820.
 - 2. A fall of snow to the depth of two feet. 1840.
 - 13. Planting potatoes.
 - 14. Sowing barley. 1820. Negunio fraxinifolium and gooseberry bushes in flower.
 - 17. Willows, gooseberries, aspens (*Populus tremuloides* in leaf. Various *Drabæ* in flower. 1820. In 1840 the trees were bursting their buds at this time.
 - 17. Wheat sown on the 8th of this month, above ground to-day, having germinated in nine days. 1840.

~~

21. Barley sown on the 14th above ground, having taken seven days to germinate.

 $\mathbf{236}$

PROGRESS OF SEASONS AT CUMBERLAND HOUSE. 237

May 22. Leave of the trees expanding rapidly.

- 24. Ulmus americana flowered. 1820.
 - 25. Pine Island Lake clear of ice. 28th. Prunus pennsylvanica, P. virginiana, and Amelanchier in flower. 30th. From the 23rd to the 30th of this month in 1840, the temperature in the shade at 2 P.M. varied between 78° F. and 93° F. On the 30th, potatoes planted on the 13th appeared above the ground. 1840.
- June 12. All the forest trees in full leaf. 1820.
- Aug. 1. Commenced reaping barley. On the 15th, 18th, 19th, and September 1., the thermometer at noon ranged between 80° and 90°, being the hottest days in the month. There was much thunder and hail on these days. 1839.
- Sept. 2. Flocks of water-fowl beginning to arrive from the north. 3rd. The first fall of snow this autumn. 4th. Vast numbers of water-fowl flying southwards. A severe fall of snow and frost in the north causes these birds to hurry to the south. 11th. First hoar-frost. Birch and aspen leaves turning yellow. 14th. Wild-fowl numerous. 20th. Snow; 21st, ditto very heavy. 24th. Thunder and lightning.
- Oct. 1. Taking up potatoes. 5th. Leaves all fallen from the deciduous trees. On the 11th, thermometer at 2 P.M., in the shade, 68° F., being unusually high.
 - 14. Water-fowl passing southwards in large flocks. 1839.
 - 15. Bays of the lake frozen over. 16th. The ground frozen hard. 17th. Last water-fowl seen this season. 18th. Lake entirely frozen over. In 1839, the Little River was frozen over on the 24th of this month, but broke up again in part, and remained partially open all the winter.
 - 31. Waveys (Anas hyperborea) passing. Lake partially opened.

follow-Frogs

is col*raxini*of this he flow t action night's ng day.

The ice l nearly that the nust rise ly free. t before but its warmer,

ts leaves

0.

ium and

muloides 20. In at this

ground 840. 1g taken

The following are the *Phenomena of the Spring of* 1827 at Carlton House, in lat. 52° 51′ N., long. 106° 13′ W., on the eastern limits of the Saskatchewan prairie lands, and at an elevation above the sea of about 1,100 feet.*

- Feb. 15. Snow thawing in the sunshine, and on the 17th many sandy hummocks on the plains were bare. This is at least three weeks earlier than the thaw commences in an early season at Cumberland House, which is a degree further north, but is 200 feet lower.
- March 6. Trees thawed in fine days, and on the 8th the black earth on the immediate banks of the river was softened to the depth of two inches by the power of the sun's rays. At this place the westerly winds bring mild weather, and the easterly ones are attended by fog and snow.
 - 13. Sparrow-hawks (*Falco sparverius*) arrived from the south, and on the 17th several migratory small birds were noticed.
 - 29. Large flocks of snow-birds (*Emberiza nivalis*) came about the establishment; and, by the 31st, steep banks, which had a southern aspect, were clear of snow.
- April 1. Many Fringillidæ (birds of the sparrow tribe) were seen. On the 2nd, swans arrived, and, by the 3rd, much snow had disappeared from the plains.
 - 4. The snow at this time was melting in the shade, and the sap of the maple trees (*Negundo fraxinifolium*) began to flow.

* This was estimated in 1827 at 1,000 feet from the length of the river course between Carlton House and York Factory, making a smaller allowance per mile for the descent, as far as Lake Winipeg, and a considerably greater one for the falls and rapids of Nelson River. It accords sufficiently with Captain Lefroy's observations; and its error does not probably exceed 200 feet at most.

PHENOMENA OF SPRING AT CARLTON HOUSE. 239

- April 6. Geese arrived. Stormy weather, about the middle of the month, retarded the arrival of the summer birds; but the plants continued to grow fast. On the 20th, the Telltale plover (*Charadrius vociferus*) and several small birds came.
 - 22. Turdus migratorius, Pyrrhula ludoviciana, and Lanius excubitor were seen, and the flowers of Anemone patens expanded.
 - 27. Ice in the river Saskatchewan gave way. Frogs began to croak.
 - 28. Canada cranes (Grus canadensis) arrived.
- May 1. Sturnus ludovicianus arrived, and the last flocks of Emberiza nivalis departed for the north.
 - 2. On this day, Icterus phæniceus and Scolecophagus ferrugineus were seen, and most of the water-fowl had by this time arrived. On the 4th, Phlox hoodii flowered.
 - 5. Ranunculus rhomboideus, Viola debilis, Nardosmia palmata, and several carices flowered.
 - 6. Hirundo viridis and many gulls arrived.
 - 7. On this day the sap of the ash-leaved maple, which had flowed scantily for ten days, ceased to run altogether, and the sugar harvest closed. Avocetta americana arrived. Populus tremuloides in flower.
 - 9. Crow-blackbirds were first seen. Corydalis aurea, Corylus americana and rostrata, Hippophäe canadensis, Thermopsis rhombifolia, Vesicaria arctica, and Alnus viridis flowered. 12th. Potentilla concinna, Townsendia sericea flowered. 14th. Gooseberry bushes coming into leaf. Ash-leaved maple flowering, seven days after the sap had ceased to flow from wounds in the stem. 16th. The Picus varius arrived in considerable numbers, and on the 19th the Viola nuttalliana flowered.

The average antecedence of spring phenomena at Carlton

1827 ., on ., and

17th bare. thaw House, O feet

th the e river by the vesterly cly ones

rom the y small

is) came it, steep clear of

be) were the 3rd,

e shade, fraxini-

th of the • naking a Winipeg, f Nelson rvations;

House to their occurrence at Cumberland House is between a fortnight and three weeks. 'The difference of latitude, which is only one degree, is nearly counterbalanced by 200 feet of greater altitude; but the dry sandy soil of the plains, which are early denuded of snow, gives the spring there a great superiority over that of the lower country, where the ground is almost submerged, and the greater part of it icebound for a month after the river is open.

I obtained no other register of temperatures at Edmonton House, or from the country near the base of the Rocky Mountains in that parallel, than the daily maxima and minima for two winter months observed by Mr. Drummond in 1827. These are included in the subjoined table; and it will be observed that the winter, as far as we can judge from a few isolated observations, is even milder at Edmonton than at Cumberland House, though it is on the same parallel and at a much greater altitude. Edmonton House is in latitude 54° N., and longitude 113° W., and its elevation above the sea is estimated by Captain Lefroy, from his observations on the boiling point of water, at 1,800 feet.

Neither have I been able to procure registers of temperatures kept at any post in the southern parts of the Saskatchewan basin. The Red River Colony extends to the boundary line of the United States, or the 49th parallel; and I have been informed that the Fagus ferruginea, or American beech, grows within the limits of the settlement, though it does not exist on Lake Superior, but terminates in that direction at Michilimackinac, on the 46th parallel. In the interesting account of the Alps of New Hampshire by Professor Agassiz, the trees which grow in the zone comprised between the elevations of 830 and 1,500 feet above the sea are the same kinds which grow on the Red River and forks of the Saskatchewan at nearly as great altitudes, and from five to ten degrees further north; with the addition of the oaks, which find their northern limit on Lake Winipeg, though they do not enter the corresponding New Hampshire zone. The same trees, however, if I understand the passage in the

ล

iı

MARTIN'S FALLS.

241

which feet of which great ground and for

Mounima for 1827. will be a a few at Cuma much N., and stimated ang point

eratures tchewan y line of nformed thin the e Supe-, on the of New v in the 500 feet d River des, and n of the though e zone. e in the work on Lake Superior (p. 186.) to which I here refer, continue to ascend the New Hampshire Mountains to the height of 2,080 feet; above which, to the height of 4,350 feet, the vegetation consists almost entirely of Abies alba and balsamea, and Betulr excelsa and papyracea. Betula excelsa scarcely reaches the Saskatchewan basin, or, if it does, is rare even in the southern parts; but the other trees here named go northwards on the Mackenzie till the 69th parallel, and terminate there on coming within the direct influence of the winds blowing from the Arctic Sea. The summits of the New Hampshire Alps 6,280 feet high, present a truly arctic flora, much more so, I believe, than the slope of the Rocky Mountains at that elevation on the 52nd parallel. On this point, however, I cannot speak with any confidence, as I have no measurements of the heights at which Mr. Drummond gathered his plants, and he is the only authority for the botany of that district.

That I may carry on as complete a view of the climate of the country as I can produce from the observations of others as well as from my own, I introduce here a table drawn up by a very intelligent officer of the Hudson's Bay Company, chief factor George Barnston*, giving the progress of the seasons at Martin's Falls. This post is situated in lat. 51° 32' N., long. 86° 39' W., on Albany River, about 200 geographical miles in a direct line from the coast of Hudson's Bay, an equal distance in a westerly direction from the water-shed between the Albany and Winipeg Rivers, and somewhat less from the beach of Lake Superior bearing south. According to Mr. Barnston's remark which follows, the country about the falls is similar in its geological character to the west side of Lake Winipeg, and referrible to the silurian epoch. "Our geological

* Mr. Barnston having sent me this paper in 1840, I published it in the Edinburgh Philosophical Journal for April, 1841.

VOL. II.

position," says that gentleman, " is upon the confines of the great basin of James's Bay, an immense extension of the older calcareous strata. Between the 'falls' and the coast the bed of the river is composed of limestones and clays, both containing extinct genera of shells; while above, towards the interior, little is to be seen but gneiss and greenstone schist, with a mixture here and there of less fissile granite rocks. The fossils I have been able to procure in this neighbourhood are principally spirifers, producta, terebratula, and impressions of trilobites. Although in winter we have the cold of Russia, in the months of July and August we enjoy the climate of Germany and the north of France." (Barnston, l. c.) A reference to the map will show that there is a canoe route from Winipeg River through Lake Sal, and by a portage over the water-shed to Lake Saint Joseph, and Albany River, and also a shorter one from the Peek River on Lake Superior to the southern tributaries of the Albany. In short, as has been already frequently stated, the primitive rocks forming the brims of these several river basins are traversed in every direction by sheets of water occupying more surface than the rocks themselves.

On the Progress of the Seasons at Martin's Falls.

Dec. Jan. Feb. We are frequently visited in these dead winter months by the white owl (Strix nyctea) from James's Bay, but the hawk owl (Strix funerea) is our most common bird of prey. Tetrao umbellus, T. canadensis, and T. phasianellus are residents in this district the whole year. The Tetrao saliceti, or willow ptarmigan, is a winter visitor which comes from the north.

SEASONS AT MARTIN'S FALLS.

March. Martens (Mustela martes) pair, and soon afterwards rabbits (Lepus americanus).

- 15. In the middle of the month the snow often melts in the height of the day, and by the 20th a snowbird (*Emberiza nivalis*) may be seen occasionally if the season be early.
- 20. Tops of the higher grasses, which have been concealed beneath the snow, begin to show. A few brown feathers clothe the necks of the willow-grouse, and these birds leave us.
- April.
- A slight crust now forms on the snow, produced by night frosts after thaw in the day. In mild weather, a few insects show in the sunshine.
- 8. Two species of *Perla* and one of *Nemoura* come up through the crevices of the ice and porous snow, and all proceed straight to the nearest beach. The cold renders them too weak to fly, though most of them have got rid of their nymphine investments before emerging from the ice.
- 15. Snow-birds have become plentiful, and are now joined by the *Emberiza lapponica* and *Alauda alpestris.* 20th. The flesh-fly is still scarce. The small owl (*Scops*) calls in the warm nights, and the common woodpecker (*Picus pileatus*) drums on the hollow trees. 22nd. The Canada goose (*Anser canadensis*) and stock-ducks (*Anas boschas*) sometimes arrive at this date, but are frequently forced to return for want of water, and by northerly blasts.
- 25. A few spots of ground bare; (more than a month later than the earth begins to appear at Carlton House.) 28th. The red-breasted thrush (*Turdus migratorius*) and cattle blackbird (*Scolecophagus ferrugineus*), are now arriving, and pick up the benumbed grubs and caterpillars. Goshawks arrive.

Falls.

ines of

sion of

and the

ies and

while

t gneiss

here of

able to

pirifers,

8. Al-

in the

of Ger-

) A re-

oe route

portage

Albany

River on

Albany.

he primi-

er basins

vater oc-

ead winter ea) from unerea) is umbellus, residents to saliceti, or which

May 1. Snow melting rapidly. Ground getting barer. 5th. Wild geese and ducks passing to the northward. Hawks still arriving. 10th. Every fine day brings an accession to the small bush birds, fly-catchers, &c. Food for these is still scarce, and they approach the houses in quest of Dipteræ, which rise from the manure and rich earth round the place. Snow-birds have left us, and ermines and hares (Lepus americanus) are becoming altogether brown. The ice is now shingly and dangerous, and strong currents and rapids are open. Wavies (Anser hyperboreus) and brents (Anser bernicla) passing towards James's Bay in large flocks. No weather stops them after this time.

- 12. The northern diver (Colymbus glacialis) and several black ducks (Anas nigra, fusca, and perspicillata) are still scarce, but are sometimes seen. The buds of the balsam poplar, aspen, and of various willows, swell. On the latter may be found the earliest Tenthredines, the larvæ being nursed in the tender bud. Two species of butterfly (Vanessa and Argynnis) sport over the ice and snow, when these are not gone.
- 15. The larger rivers break up. (The Saskatchewan opens about twenty days earlier at Carlton House). Fish ascend the small streams to spawn. The pike (Esox lucius) and dorè (Lucioperca) spawn. Sucking-carp (Catastomi) soon follow them. Trout take bait greedily. The cliff-swallow (Hirundo fulva, Vieillot) is seen. (On May 28th, 1849, Mr. Rae found this swallow on the banks of the Coppermine, having constructed its clustered nest against the cliffs at the mouth of Kendall River, lat. 67° N., but not yet laid its eggs.) Swamps and stagnant pools are thawed. Frogs begin to croak, and moschetoes to bite.

SEASONS AT MARTIN'S FALLS.

- May 20. Shells (Limnæi) begin to move in the pools along the river. Snails (Limax, Helix, Bulimus, &c.) remove from under stones and fallen timber. The end of the month discloses some species of moths (Noctualites).
 - 25. Our only goatsucker (*Chordeiles virginianus*), and the golden-winged woodpecker (*Colaptes auratus*), the last of the spring birds in this district, arrive. Beavers, otters, and musk-rats have their young.
 - 28. The balsam poplar, and aspen expand their leaves. (Not so early as on the Saskatchewan, two or three degrees further north.) The Hudson's Bay reindeer has young.
- June 1. Sturgeon begin to frequent falls and rapids, and to spawn. 5th. Insects are busy on warm days; the *Tenthredinidæ* on bushes, the *Sphingides, Andre*netæ, and Pangoniæ on the ground, all attended by a variety of parasite ichneumons. The first flowers blow, and those of the willow are surrounded by Sylphides and flower-flies (Anthomoyæ).
 - 10. A night-frost will sometimes intervene, even at this late period; and, in the woods, the ground is still frozen solidly at the depth a foot from the surface. Vegetation, nevertheless, still goes forward. Moschetoes become a torment; the swamps and puddles swarm with their larvæ. Small talpoles abound in the pools. 13th. The country is now covered with verdure. Birds are nestling; geese and ducks hatching. The natives are occupied with the sturgeon fishery.
 - 15. The latest shrubs are in leaf, and the majority of moths and butterflies are disclosing themselves. The large Ephemeræ (Perlæ and Phryganeæ) issue from the water. 20th. Trout take the fly-hook. White-fish (Coregonus albus) rise to the surface. Cattle seek the houses to get rid of the tormenting

. 5th. hward. brings atchers, ney apich rise e place. d hares r brown. l strong *mser hy*ssing toweather

d several picillata) The buds s willows, e earliest he tender *tessa* and then these

atchewan n House). The pike) spawn. m. Trout (*Hirundo* th, 1849, ks of the tered nest all River, Swamps begin to

Tabani. In dry seasons rivulets become low, though rivers retain their strength.

Our warmest month. The river unusally gets low. Sturgeon fishing continues. Oattle become lean, being tormented by flies in the day, and feeding in the night only. 10th-20th. Many genera of Coleoptera appear, some of them of forms more characteristic of warmer climes; Cicindelæ, Necrophori, many Buprestes, and a species allied to Lucanus. Of those whose larvæ live on wood, the Serropalpus, a very fine Dorcacenis, Cerambyx, Callidium, Lamia, and numerous species of Lepturetæ. Neuroptera are abundant on the banks of the river -Libellulæ, Agrion, &c.; and on the leaves Hemerobius, Panorpes, Sialis. In the other orders, also, there are many genera to keep up the character of the month. It ends with bringing in strawberries, and in sending off sturgeon, which return to the depths.

Aug.

- The raspberries begin to ripen. Young ducks are well feathered. We have sultry weather for a few days, and then thunder-storms followed by chilly nights. 10th. Pigeons (*Ectopistes migratorius*) are numerous. Young geese (*Anser canadensis*) can fly. Gnats decrease, and sand-flies (*Similium*) replace them.
- 15. The raspberries, and red and black currants ripen. Grasshoppers are full grown. Trout move about, ascending the river. Sturgeon are very scarce. Grass becomes brown in dry situations. 20th. The noisy lesser tell-tale (*Totanus flavipes*) appears, and if we have much rain we are visited by a species of snipe. The golden and ring-necked plovers are not uncommon.

Sept.

The air is generally cooler and the wind stronger, and frosty nights may be expected.

July.

SEASONS AT MARTIN'S FALLS.

Sept. 10. Trout spawn. Many insectivorous birds depart. The passenger pigeon disappears. Hawks and the large horned owl (Strix virginianus) are common. Night frosts frequent.

- 15. Tops of potatoes always blackened. Caterpillars nearly all cased. Trout refuse the fly-hook, but still take bait; they are now poor fish. Canada geese pass to the southward. Ducks abound in the grassy lakes. Leaves turning ye low rapidly. 20th. Pleasant weather at mid-day but cold in the night. The fall-moth (*Phalæna autumnalis*) is now to be seen. Sand-flies bite only in the height of the warmer days. 'The moscheto is utterly defunct. Diving ducks common; the others gone.
- Oct. 1. Pools and swamps crusted with ice. White-fsh (Coregonus albus) begin to spawn. 5th. Suckingcarp (Catastomi) and trout desert the small streams. Foliage is yellow, and falls. Deer rut. Instead of rain we have snow, which generally melts, the earth being warmer than the atmosphere.
 - 10. A single blast of northerly wind will now suffice to bare the trees, strip the shrubs, and send all waterfowl to the south. The last of these are weak or lingering flocks of wavies (Anser hyperboreus), which may be observed passing. They seldom alight unless when met by adverse winds.
 - 20. The small lakes and rivers sometimes frozen. (Between the 21st and 28th is the usual period for the freezing of the lake at Cumberland House.) Tullibee (*Coregonus artedi*) spawns. Quadrupeds acquire thic'. fur. The willow-grouse (*Tetrao saliceti*) arrives from the north. There is usually a little snow on the ground; and the American hare and ermine are becoming white.
- Nov. 1. The ground is covered with snow, which in mild weather harbours multitudes of *Podura*. 10th. I
 - **r** 4

e low,

ts low. e lean, ling in of Colecharacophori, ucanus. erropal-Ilidium, . Neuriver s Hemeers, also, racter of wberries, n to the

ucks are for a few by chilly orius) arc ensis) can ilium) re-

nts ripen. ve about, y scarce. 20th. The appears, ited by a ng-necked

stronger,

have frequently observed at this time, and later, a wingless tipula (Chionea hiemalis) crawling about. 20th. Large rivers and the lakes often solidly frozen; strong rapids filling with ice, and setting fast. American hares and ermines entirely white. Swamps passable, and winter fairly set in.

(Geo. Barnston.)

The following abstract of temperatures, ascertained by John Rae, Esq. in 1846-7, at Fort Hope, Repulse Bay, 62° 32' lat., 86° 56' long., which was accidentally omitted to be inserted in Table XII., is placed here, the space being convenient for its reception.

The observations were made eight times in the twentyfour hours, at equal intervals; and the house was situated only a few feet above the sea level.

Sept.	28·57	Dec	March —28·10	June 31·38
Oct.	12·56		April —3·95	July 41·46
Nov.	0·68		May 17·88	August 46·32
Aut.	13.81	Winter-25.04	Spring -4.73	Summer 3959

Mean heat of the year 1846-7 Difference of bottest and coldest months - 75.64 summer and winter -22

6.14° Fah.

- 64.63

COMPARATIVE TABLE OF TEMPERATURES. 249

TABLE IX.

TEMPERATURES BETWEEN THE PARALLELS OF 52° AND $61\frac{1}{2}^{\circ}$ North Latitude.

Periods.	Churchill. Lat. 59° 02' N. Long. 93° 10' W. Ait. 20 ft. Year, 1838-39. Morning, Aftern., Evening. Corrected by Dove's formula. (1.)	Greenland. Frederick sthal. Lat. 60° N. Long. 45° W. Alt. 20 ft. Dove. (2.)	Pelly Banks. Lat. 610 30' N. Long. 130° W. Alt. 1,400 ft. 5 Months, 1848, 1849. Sunrise, 3 P. M., Dusk. Corrected. (3.)
January February March April June July September October November December Year Year Summer Summer Spring Autumn - Difference of hot. and cold. months Difference of	$\begin{array}{r}21 \cdot 21 \\7 \cdot 31 \\ -4 \cdot 63 \\ 16 \cdot 29 \\ 28 \cdot 42 \\ 44 \cdot 69 \\ 56 \cdot 80 \\ 53 \cdot 39 \\ 36 \cdot 03 \\ 26 \cdot 50 \\ 3 \cdot 32 \\ -14 \cdot 00 \\ \hline 18 \cdot 19 \\ \hline -14 \cdot 17 \\ 51 \cdot 63 \\ 13 \cdot 36 \\ 21 \cdot 95 \\ \hline 78 \cdot 01 \\ \end{array}$	19.62 18.72 22.10 27.50 32.45 35.15 29.75 30.00 ? 22.70	
Difference of summer and winter	65.80		

No. 1. The register at Churchill was kept by Mr. Harding, of the Hudson's Bay Company's Service, from February, 1838, to May, 1839, the observations being made daily in the morning, afternoon, and evening. Column 2. is extracted from Dove's table; and for column 3. I am indebted to Murdock M'Pherson, Esq., who sent me a register kept at Pclly Banks by R. Campbell, Esq.

about. solidly setting white.

nston.)

ed by e Bay, mitted space

wentyituated

	31·38 41·46 46·32	
•	39 59	

ah.

TABLE

Periods.	Godhaab. Lat. 64º 10' N. Long. 52º 24' W. Sea level.	Fort Chepewyan. Lat. 58° 43' N. Long.:118° 20' W. Alt. 700 ft.	Fort Resolution. Lat. 61° 10' N. Long.113° 51'W. Alt. 500 ft.	Fort Simpson. Lat. 61° 51′ N. Long.121° 51′ W. Alt. 400 ft.
	(1.)	(2.)	(3.)	(4.)
January	12.38	-8.76	0.42	-12.54
February	12.56	-4.01	-25.60	-9.06
March	15.60	3.08	9.95	5.55
April	22.01	19.80	12.88	26.28
May	32.16	45.40	40.14	48.16
June	39.09	55.00		63.64
July	41.92	63.00		60.97
August	40.84	58.10		53.84
Sentember -	35.65	43.53		49.10
October	29.84	33.00	26.06	24.28
November -	21.94	19.13	12.041	8.53
December	17.49	2.76	-2.29	-8.36
Year	26.79	27.52	21.00?	25:75
Winter	14.14	-3.34	-8.09	-10.00
Summer	40.62	58.70		59.48
Spring	23.26	22.76	20.99	26.66
Autumn	29.14	31.89		27.34
Difference of hot. and cold. months -	29.54	71.76	. .	76·18
Difference of summer and winter -}	26.48	62.04		69.48

THERMOMETRICAL OBSERVATIONS IN THE VALLEY OF THE

No. 1., extracted from Pove's table, Seventeenth Rep. Br. Ass. p. 376., gives the temperature of the west coast of Greenland in that parallel, and is inserted for comparison.

paramet, and is inserted for comparison. No. 2. is compiled from $3\frac{1}{2}$ years' observation by Messrs. Keith and Stewart in 1825 and 1826, and by another officer in 1838 and 1839. In the latter case, the times of observation were eight and eight, and the correction for these hours, calculated by Dove for Toronto, has been applied. The summer months were not entered in the register on the second occasion. Observations were also made at 2 p. M., and are recorded in another table.

2 P. M., and are recorded in another table. No. 3. is abstracted from a register of observations made at eight and eight, in the winter and spring : Dove's correction has been used.

and eight, in the winter and spring: Dove's correction has been used. No. 4. is an abstract of 2½ years' observation by Murdock M'Pherson, Esq. in 1837, 1838, 1839, and 1840, at eight and eight, with Dove's correction for Toronto applied.

COMPARATIVE TABLE OF TEMPERATURES. 251

BLE

THE

npson. 51' N. 951'W. 90 ft.	
.)	
54 06 55 28 16 64 97 84 10 28 53 -53 -36	
:75	
•00 •48 •66 •34	
5·18	
•48	

r. Ass. n that

Keith 8 and 1t and ve for red in ade at

> eight used. rdock eight,

X.

MACKENZIE; IN GREENLAND; AND ON THE YUKON.

Fort Rellance. Lat. 62° 46′ N. Long. 109° 00′ W. Alt. 650 ft.	Foit Enterprise. Lat. 64° 28' N. Long. 113° 06' W. Alt. 850 ft.	Fort Franklin. Lat. 65° 12' N. Long. 123° 13' W. Alt. 500 ft.	Fort Confidence. Lat. 66° 54' N. Long.118° 49' W. Alt. 500 ft.	Yukon. Lat. 66° N. Lon. 147°W, Alt. 200 ft. ?	Periods.
(5.)	(6.)	(7.)	(8.)	(9.)	
-25.00	-15.57	-23.33	-21.57	-26.85	January.
-18.84	-25.88	-16.75	-21.52	-26.44	February.
-6.14	-13.48	-5.38	-20.21	-11.16	March.
8.23	5.78	12.35	-4.71	12.66	April.
36.03	31.20	35.18		41.24	May.
		48.02		53.49	June.
		52.10		65.75	July.
		50.56		59.90	August.
	31.59	41.00		38.66	September.
20.70	21.75	22.47	19.43	21.60	October.
13.44	-1.70	-0.11		-8.28	November.
-17.07	-30.54	-10.88		-18.43	December.
16.00 %	17.94	17.75	9.00 \$	14.58	Year.
-16.97	-24.00	-17.00	-27.26	-23.80	Winter.
	55.00?	50.41		56.73	Summer.
12.21	23.20	12.69		14.60	Spring.
	17.21	21.15		17.37	Autumn.
	86.88 ?	74.43		92.60	{Difference of hot. and cold. months.
	79-00 ?	67.41		80.23	Difference of summer and winter.

No. 5. is compiled from Sir George Back's observations in 1834, (fifteen daily.)

No. 6. is abstracted from a register kept at Fort Enterprise in 1820, 1821, and 1822. The summer observations are defective; but the mean summer heats are inferred from observations made in travelling over the barren grounds further north.

No. 7. Two years' observations at Fort Franklin; first year nineteen observations daily, except in summer, during which, and in the second year, six observations daily at equal intervals.

No. 8. Fifteen to seventeen observations, daily.

No. 9. Observations at six and six in summer; and in winter as early as the thermometer could be read in the morning, and as late in the evening. Dove's correction for these hours at Toronto has been applied.

TABLE

TAPLE OF THE SUMMITS OF THE DAILY OR MONTHLY CURVES OF BETWEEN THE PARALLELS OF $55\frac{3}{4}^{\circ}$ and

An a second design of the seco								
Periods.	Fort Churchill. Lat. 59° 02'. Long. 93° 10'. Sea Level. (1.)		Fort Chepewyan. Lat. 58° 43' N. Long. 118° 20 W. Alt. 700 ft. (2.)		Fort I Lat. Long. Alt	Resolution. 61° 10' N. 113° 51' W. 2. 500 ft. (3.)	Fort Simpson. Lat. 61° 51' N. Long. 121° 51' W Alt. 400 ft. (4.)	
	2 p. m. Mean .	Monthly Maxima.	2 P. M. Mean.	Monthly Maxima.	M M	onthly axlma.	2 P. M. Mean.	
Junuary -	-16.48	-2	-7.04	22		20°	-17.55	
I ebruary	-1.18	42	3.21	44		2	-1.68	
March -	8.51	37	16.72	47		37	12.68	
April -	31.66	58	39.12	65		32	47.22	
May -	37.94	65	33.33	76		64	56.98	
June -	55.17	82	62.00	97	-	-	73.40	
July -	66.81	86	69.00	78	-	-	70.61	
August -	62.97	83	66.10	72	-	-	65.87	
Sentember	54.09	84	46.00	65	-	-	60.00	
October -	33.32	46	41.06	56		47	30.08	
November	8.00	23	16.71	39		31	12.51	
December	-9.77	5	4.95	40		20	-7.98	
Year -	27.59	86	32.61	97	-	•	33.51	
Winter -	-9.14	-	0.38	44		20	-9.37	
Summer -	61.65	-	65.70	97	-	-	69.96	
Spring -	26.04	-	29.72	76		64	38.89	
Autumn -	31.80	-	34.59	65	-	-	34.15	

XI.

Темрі 66° 54

. 0

COMPARATIVE TABLE OF TEMPERATURES. 253

BLE

XI.

VES OF

mpson. 5 bi N. 10 bi W. 100 ft. 7.555 1.68 2.68 7.22 6.98 3.40 0.61 5.87 0.00 0.08 2.51 7.98 3.51 9.37 9.96 8.89 4.15 TEMPERATURE IN THE VALLEYS OF THE MISSINIPI, MACKENZIE, YUKON, AND PFLLY, 66° 54' N., EXPRESSED IN DEGREES OF FAHRENHEIT.

Fort Enterprise. 1 at. 64° 28' N. Long. 113° 06' W. Alt. 850 ft. (5.)		terprise. 28' N. 3° 66' W. 50 ft.	Fort Franklin. Lat. 65º 12' N. Long. 123º 13' W. Alt. 500 ft. (6.)		Fort Confidence. Lat. 66° 54' N. Long. 118° 49' W. Alt. 500 ft. (7.)		Yukon, 1 st. 66° N. Lot & 147° W. M. 200 ft.? (S.)		Pelly Banks. Lat. 61° 30' N. Long. 130° W. Ait. 1,200 ft. (9.)		Periods.
Mea Max	ins of tima.	Monthiy Maxima.	Mean Temp. at 2 P. M.	2 Years, Mnthiy Max.	Mean Temp. at 2 P. M.	Mor Maxima	lean at 1 P. M.	Monthly Maxima.	Mean at 3 p. m.	Monthly Maxima	
-	9.68	20	21.53	11.8	-17.83	2.0	27.58	17	-16.61	16	Jan.
-1	9.10	1	-8.24	27.8	—16·49	2.0	-23.55	10	-00.35	29	Feb.
-	0.90	24	1.71	31.8	-7.45	12.8	-0.94	28	10.35	34	Mar.
1	6.40	40	23.21	49.0	6.20	24.5	19.43	52	36.20	50	Apr.
4	2.80	68	41.65	69.0	-	-	48.81	70			May
	•	-	51.00	75.0	-	-	62.00?				June
		-	59.02	80.0	-	-	74.84	76			July
		-	56.88	74.0	•	-	70.94	86			Aug.
3	19.30	53	44.82	66.0	-	-	52.73	69			Sept.
2	7.90	37	23.75	47.0	22.42	32.0	9.49	50			Oct.
1	2.80	25	4.81	32.5	0.41	19.0	-5.40	28			Nov.
-2	25.80	6	-12.07	27.5	-32.23	-7.0	-5.39	22	-0.34	4	Dec.
	•		22.09	80	•	-	23.89	86			Year
-1	17.15	20	-14.14	27.8	-22.58	27	-22.70	22	-5.72	29	Wint.
	•	-	55.64	80	-	-	69.28	86			Sum.
	•	-	22.18	69	-	-	22.71	70			Spr.
1 5	23.33	-	24.45	66.2	-	-	26.03	69			Aut.

Progress of the Seasons at Fort Franklin, on Great Bear Lake, in lat. 65° 12' N., long. 123° 12' N.

The mean temperature of the three winter months varies comparatively little in different years; but the relative temperatures of these months differ greatly among themselves, so that in one year December is the coldest month, in another February, and in a third January. In some years the temperature of places exposed to the sun rises for a day or two in winter above the freezing point, and the snow moistens on the surface; but in other winters no thaw whatever occurs.

In March the snow is deepest, and averages about three feet, being, however, often drifted to a much greater thickness under cliffs and on the borders of lakes. In the end of March or beginning of April trees begin to thaw, the mean temperature in the shade being about zero Fah.; but the effect of the sun's rays on the blackened bulb of a thermometer being sufficient to raise the mercury to $+90^{\circ}$ Fah.

About the 10th of April the snow begins to thaw decidedly in the sunshine, and myriads of *Poduræ* are seen at such times moving actively in its cavities. Ptarmigan begin to assume their summer plumage towards the end of the month.

From the 1st to the 6th of May, according to the season, water-fowl arrive. The *Colymbus glacialis* and *arcticus*? arrive occasionally earlier, and frequent a piece of water at the efflux of Bear Lake River from the lake, which remains open all the year.

Swans (Cygnus buccinator and americanus) are among the early arrivals, the larger species coming first. The Anas acuta, A. crecca, Clangula histrionica, and Oidemia perspicillata make their appearance within the first eight days. Gulls come about the 9th or 10th. Singing birds, orioles, and swifts arrive about the middle of the month; the latter varying their time of appearance to a week later, if the spring is tardy. Pools of water and swamps must have been thawed long enough to release a sufficiency of winged insects for the support of the swallow tribe, before they show themselves in a

SEASONS AT FORT FRANKLIN.

district. About the 10th or 12th of the month small streams break up, the mean temperature of the ten preceding days having risen to 37° Fah. Bear Lake River, which is fed from the depths of the lake with warmer water, breaks its bonds at its efflux earlier. Lower down, this river remains fast till the first or second week of June. Mackenzie River usually opens at Fort Simpson about the 7th o. May, and in the more northern quarters in the course of a week later; the boats which follow the flood in its descent taking about that time to go to Fort Good Hope. In 1849 the river broke up on the unprecedentedly late date of the 23d of May.

At this latter date there is bright light at midnight on Great Bear Lake, and the *Fringilla leucophrys* is employed with other songsters in singing at that hour.

Snow-geese arrive about this time, or a week earlier, and are followed in a fortnight by the laughing-geese. Both kinds wing their way northwards in bands of from fifteen to forty individuals, which are passing every few minutes day and night for about three weeks. Many go on without halting; others alight in the marshes to feed on the nascent stems of the early Cyperaceæ, which are developed with marvellous rapidity after the commencement of the thaw, and, though still wrapped in the dead leaves of last season, have acquired juiciness and a sweet taste by the time that the snow has mostly gone. Towards the end of May, or in the first week in June, according to the earliness of the spring, Chrysosplenium alternifolium, Arbutus alpinus, Eriophorum vaginatum begin to flower, and the Betula glandulosa and some willows show their tender foliage and catkins. Early in June the Potentilla fruticosa, the Rhododendron lapponicum, and several anemones flower. Frogs at this time croak loudly; and by the middle of the month, summer may be considered as fairly established. About the 24th or 26th of July, ripc bleaberries (Vaccinium uliginosum) may be gathered. Strawberries are generally a week earlier, and the Arbutus alpina and Rubus chamamorus, or cloudberry, are somewhat later.

In the beginning of August stars may be seen at midnight;

lear

tems, so other temtemvo in n the

feet, under ch or ature sun's ent to

idedly such gin to nth. eason, ticus ? iter at emains

> ig the acuta, cillata come swifts ng is nawed or the s in a

and in the last week of this month the van flocks of snow-geese are seen going southwards, having spent between eighty and ninety days at their breeding stations. The laughing-geese follow in a day or two; but they pass on in autumn without any of the delays that characterise their spring flights, which a.e necessarily checked as often as a few cold days arrest the melting of the snow on the sea-coast. Drift ice obstructs the navigation of the lake in some seasons till the first or second week of August.

Sept

Octo

Nove Dece

Janu

Febr Marc

April

May .

June July

Augu

Year

Spring

Autun

Summ Winter

Differe

hot.

mont Differen

sumn

winte

H

T

lis

th Ro ev

In the last week of August, or in the beginning of September, snow falls, and by the 10th of the month the deciduous leaves begin to drop. By the 18th, most of the birds which breed in the district have migrated southwards, a few waterfowl and the winter residents alone remaining. Between the first appearance of vegetation, till the falling of the leaves of deciduous trees, about a hundred days elapse; but although this may be taken as the length of the season for the growth of plants. some of the grasses continue to ripen their seeds till the beginning of October, notwithstanding much severe frost before that date. In ordinary seasons the frost sets in severely before the end of September, and the seeds of carices and grasses, instead of dropping off, are frozen hard in their glumes, and remain hanging on the culm till next spring, when they drop off into a soil prepared by the thaw for their reception. It is on these grass seeds of the preceding year that the graminivorous birds feed on their first arrival from the south. In October, when the soil begins to freeze again, the summer thaw has penetrated about twenty-one inches in the neighbourhood of Fort Franklin. The small lakes are covered with ice by the 10th or 12th of the month; and, when that occurs, the last of the water-fowl depart. By the 20th of the month the smaller trees are frozen through, the larger ones remaining soft and moist in the centre. By the end of the month, or early in November, the young ice, filling the bays, puts an end to the navigation of the lake, after it has continued open about sixty days. The centre of the lake does not freeze over till late in December.

COMPARATIVE TABLE OF TEMPERATURES. 257

TABLE XII.

THERMOMETRICAL OBSERVATIONS IN THE ARCTIC SEAS.

Periods.	Melville Island. Lat. 744° N. September, 1819- August, 1820.	Port Bowen. Lat. 7340 N. September, 1824- August, 1825.	Igloolik. Lat. 6940 N. August, 1822- July, 1823.	Winter Island. Lat. 664º N. August, 1821- June, 1822.	Wolstenholme Sound. Lat. 764 N. Six times daily. August. 1849- July, 1850.
	(1.)	(2.)	(3.)	(4.)	(5.)
September -	+22.66	+25.90 +10.84	+25.10 +13.72	+31.62 +13.15	+26.76 +11.32
November -	-20.85	-5.00	-18.66	+7.80	-18.60
December	-21.80	-19.05	-28.25	-14.24	-27.05
January	-30.64	-28.91	-16.13	-23.13	-25.07
February	-32.31	-27.31	-19.58	24.01	-34.02
March	-18.10	-28.36	-19.01	-10.78	-17.47
April	-8.30	-6.20	-0.82	+6.20	3.74
May	+16.74	+17.62	+25.14	+23.31	+25.82
June	+36.22	+36.12	+32.16	+33.16	+39.73
July	+42.43	+38.87	+38.28	+35.33	+40.52
August	+32.63	+35.77	+33.88	+36.88	+33.67
Year	+1.43	+4.33	+5.76	+9.80	+4.54
Spring	-3.13	-5.74	+1.	+6.35	+1.29
Autumn	-0.43	+10.28	+ 80	+17.53	+6.55
Summer	-+ 37.09	+36.93	+35.07	+35.15	+37.97
Winter	-28.36	-25.02	-21.38	-20.35	-28.53
Difference of hot. and cold. months	74.74	67•78	66•83	60.89	74.54
Difference of summer and winter	65.45	61.95	56.45	55.20	66.20

SPITZBERGEN.

	June.	July.	August.	Sum. Quarter.
Hecla Cove, lat. 80° N.	+35.86	+40.17	+38.40	+38.15
m. lat. 80° N., long. 10° E.	33.73	35.98	33.80	34.52

Columns 1, 2, 3, and 4. are extracted from a paper which I published in the Ed. Phil. Journ. for 1841. Column 5. is an abstract of a thermometrical register kept by James Rae, Esq., surgeon of the Royal Navy, on board the North Star. An observation was made every four hours.

VOL. II.

1

S

geese y and geese at any ch a.'e meltnaviweek

f Sepiduous which waterhe first iduous is may plants, the bebefore before ses, inremain f into a n these s birds hen the etrated anklin. h of the depart. hrough, By the filling r it has

ke does

The preceding pages contain the temperatures of the districts through which the Expedition travelled, wherever I have been able to ascertain them, and also data for extending the lines of mean annual heat (isothermal), mean summer heat (isothæral), and mean winter heat (isocheimenal) across the continent. By comparing the sca-coast temperatures in Table II. and those of the shores of the great lakes in Table I. with those of places in France and Italy lying between the same parallels of 42°---45° north latitude, we perceive that the mean annual heat of Europe is from 8° to 15° Fah. greater than that of America at the same distance from the equator, while the summer heats differ only from 2° to 6°.* The inferior mean heat of America is therefore due principally to excessive winter colds, and this is decidedly the case in the interior. As the summer heats, however, regulate the culture of the cerealia and the growth of deciduous plants generally, the severe winters of America do not cause a scanty vegetation. From the 50th parallel northwards the trees are frozen to their centres in winter; and, consequently, the development of buds and other vital processes which go on in the temperate climate of England, even in the coldest months, are completely arrested. This hybernation of plants increases in length with the severity and duration of winter which, generally speaking, augment in the interior of America with the latitude. The summer heats

* Dove's table in the Report of the Brit. Association for 1847 has furnished the means of making this comparison. The places compared were Alais, Arles, Bourdeaux, Dax, Manosque, Marseilles, Montpellier, Pau, Puy, Tarascon, Toulon, Toulouse, Perpignan, Alba, Bologna, Cascina, and Lucca. Oleron of Bearn has the same altitude with Lake Superior, and Mount Louis is 4,900 feet high. All of them lie within the parallels of 43°-45° except the last, which is in 42° 50′ N. The maps of isothermal lines of this author express the general results of the study of the table referred to.

d

81

pe

th

COMPARISON OF TEMPERATURES.

do not, however, decrease in the same ratio as we go to the north; on the contrary, the isothæral lines nearly follow the canoe route, and run to the northward and westward. The elevation of the prairie slopes has less influence in depressing the summer heat, than the nature of the soil and other causes have in raising it.

Experiments are still wanting whereby we may ascertain the ratio of the decrease of mean heat in America with the increase of altitude. In Table II. we find that, notwithstanding the elevation of Franklin Malone above Eastport of 645 feet, its mean temperature is greater; its interior position giving it an advantage in summer heat over the sea-coast, which its greater altitude does not destroy. If we refer to Dovc's table, and contrast the temperatures of Mont Louis, which lies near the 43d parallel, with those of low country situations enumerated at the foot of the preceding page, we find a mean difference of temperature of one degree of Fahrenheit for 350 feet of altitude. A similar allowance for the elevation of the successive steps of the St. Lawrence basin would place in still greater prominence the rise of the isothermal lines, and more still that of the isotheral ones, as they recede from the Atlantic coast. There is, however, this difference in the climate of the summit of a mountain and of an elevated plateau, that in the former case we approach near the line of invariable temperature, and the summer heat therefore differs less from the mean of the year, and more from that of the plains, than on a plateau where the depression of mean temperature produced by elevation is due chiefly to winter colds, and in a small degree only to defect of summer heat.

From Table II. also we may learn that the mean temperature of the coast districts of the Pacific is greater than that of the Atlantic countries, and, at the same time, more

\$ 2

of the erever stendimmer across atures akes in lying de, we rom 8° e same s differ merica ds, and summer alia and severe getation. e frozen the dech go on e coldest ation of luration the inor heats

> for 1847 he places farseilles, han, Alba, e altitude ll of them hich is in spress the

equable; the difference between the hot and cold months being less. We find in it an expression of the general fact, that the west coasts of continents are warmer than the east ones; and as Montreal and Fort Vancouver lie nearly in the same latitudes and at the same altitudes above the sca, and both are far enough removed from the coast to be beyond the direct influence of the sca breezes, columns 4. and 6. furnish the means of eliciting many of the peculiarities of climate on the two sides of the continent. Instead of four or five months of continuous snow and ice which Canada may be said to enjoy, for it is the season of general enjoyment, Oregon has an open, rainy winter, with little frost or snow; but, at the same time, a summer of less power.

Table V. exhibits even greater differences in the Pacific and Atlantic climates in a higher parallel. The course of the occan currents, and the interposition of the peninsula of Alaska and its prolongation by the Aleutian chain of islands, protect the west coast of America from the masses of drift ice which in the same latitudes encumber and chill the Labrador coast for most of the year. Even in the polar regions the west coasts have milder climates. Table X. shows, as far as it goes, that the mean temperature of the west coast of Greenland exceeds that of places on the continent, up to the 150th meridian, though the summer on the coast is greatly colder than that of the interior. By the study of Table XII. we learn that in the polar seas the summer heats vary little, as we might expect from the constant presence of ice; but the annual mean seems to decrease generally with the latitude, the only exception being that of Wolstenholme Sound, in which we have a confirmation of the greater mildness of the west Greenland coast. In the high latitudes the mean heat of the three winter months does not differ greatly in different

years; but in some sensons one of these months, in others another, is the coldest, the temperature being ruled greatly by the prevailing winds.

Generally speaking, the mean annual temperature of places in the interior of North America falls within a degree or two of the mean heat of the two months of April and October. The mean temperature of the whole surface of the earth is, according to Dove's calculations, 58.2° Fah., being 54° in January and 62° in July. For such a mean annual heat in America we must descend to the 34th parallel of latitude; but the July heat of 62° Fah. extends northwards to the Mackenzie.

The stanse winter colds in the high latitudes are apparently in a great measure owing to the active nocturnal radiation into the clear blue sky. The observatory, which was a small log building without a fire-place, furnished us with the means of judging how much greater the depression of temperature in the night was in places exposed to the sky, than in those covered in.

The daily curve of atmospheric temperature for the three winter months was a bold and nearly regular hyperbolic curve, of which the mean was $-25 \cdot 2^{\circ}$ Fah., the maximum $-22 \cdot 2^{\circ}$ Fah., and the minimum $-26 \cdot 7^{\circ}$ Fah.* The maximum occurs at 1h. 18m. P. M., and the mean line is crossed by the curve at 9h. 19m. A. M., and 6h. 28m. P. M.; the lowest temperature being reached at 7 A. M. The ascending branch of the curve, therefore, corresponds to an interval of 6h. 18m., and the descending one to 17h. 42m. During 14h. 51m. the temperature is below the mean, and it is only 9h. 9m. above it, which indicates a tolerably bold curve in the day, and a nearly horizontal course in the night.

* The correction for the error of the thermometer at low temperatures used in Tab. X. col. 8. was not applied to these numbers.

8 3

nonths general r than over lie titudes om the preczes, g many of the tinuous for it is n, rainy ne time,

Pacific ourse of eninsula chain of e masses ber and Even in climates. an temthat of though that of arn that ve might e annual ade, the in which the west heat of different

In the observatory the mean for the same period was only -15.91° Fah., and the range no more than 0.97° . The maximum occurred at 6 P.M., being retarded 43 hours; and the minimum at 10 A.M., being delayed 3 hours. For most of the night the temperature was above the mean, such being the effect of the interposition of the building between the thermometer and the blue atmosphere. The walls of the observatory, it is necessary to remark, were by no means air-tight, and the door was opened at least once an hour in the day, and sometimes, especially on term days, much oftener. There was, consequently, a considerable and frequent admission of the external air; and, on the other hand, during the experiments on magnetic intensity, the heat of the observer's body had an evident effect in raising the temperature of the room.

I had intended to have instituted a series of observations, with Sir John Herschell's actinometer, on the nocturnal radiation, and also on the momentary intensity of the direct rays of the sun; but the instrument was unfortunately broken on the voyage. 'The Edinburgh New Philosophical Journal for 1841 contains the results of observations made at Fort Franklin, with the black bulb thermometer, on the heating power of the sun's rays, and I renewed these observations at Fort Confidence; but, as they were not carried on later than April, they furnish no information respecting the power of the sun in the months in which the processes of vegetation are active. As the black bulb thermometer indicates the accumulative effect of the sun's rays, it seems to be an useful instrument for ascertaining the heating power of the sun on the stems and larger branches of trees, at least, if not also on their leaves and on herbaceous plants. The hybernation of trees ceases long before the temperature of the atmesphere is sufficient to restore activity to the vegetative

EFFECTS OF SUN-LIGHT ON VEGETATION. 263

processes, and before the earth, still enveloped in its snowy eovering, has felt the influence of returning spring. This is evidently mainly or wholly due to the sun's light direct or reflected; and perhaps its rays as reflected from the pure and glassy surface of the snow, after the days have increased considerably in length, may have the same powerful effect on the forest that, according to Professor Forbes, they have on the black-build thermometer. For some time after the trees have begun to thaw by day, they freeze again in the night; and in more southern localitics, where the sugar-maple grows, the sugar makers are well acquainted with the fact that a hard frost arrests the flow of the sap in the night. Should a hot day, however, follow such an occurrence, the flow is more abundant than ever, the short rest seemingly mereasing the irritability of the organs by which the sup is eliminated and circulated.*

* As I was revising this sheet, Sir William Tooker favoured me with an extract from the join all of Mr. Bertonic Seeman, botanist of the Herald, part of which follows: -- "Doing our stuy at fort Clarence, in September, 1850, I made several experiments to ascertathe depth to which the naw penetrated the sole the result varied in some places it did not descend above two feet into the earth, while in sandy places the ground was free from frost to the depth of four or five feet. The season was much colder the in 1849, the sca more loaded with ice, and the terrestrial vegetation ess vigorous"

od was 2. The hours; s. For ean, uilding . The vere by st once m days, uble and he other ity, the raising

observathe nocnsity of was ungh New sults of ck bulb nys, and e; but, furnish in the active. nulative instruon the also on ernation the atgetative

No. III.

ON THE GEOGRAPHICAL DISTRIBUTION OF PLANTS IN THE COUNTRY NORTH OF THE 49TH PARALLEL OF LATITUDE.

Generic and Specific Forms of Plants' Decrease in Number as the Latitude increases. — Analogy between Altitude and Increase of Latitude. — Culture of the Vine. — Of the Cerealia. — Maize. — Wheat. — Oats. — Barley. — Potatoes. — Botanical Districts. — Their Physiognomy. — Woodland District. — Barren Grounds. — Prairies. — Rocky Mountains. — Sitka. — Polar Plants. — Arctic Zone. — Trees and Shrubs. — Table of Distribution of Species in three scveral Zones. — Carices.

THOUGH the isotharal lines, when the term is restricted to the mean temperatures of the three summer months of June, July, and August, run from Lake Superior northwards to the Mackenzie, yet the short duration of the summer on the banks of that river, and the occasional frosts in June and August, and in some years even in July, render the climate unsuitable for numbers of vegetables which flourish in the northern districts of the United States. Many trees, shrubs, and perennial roots can be frozen without injury if the frost be continuous throughout the winter; and they acquire so much irritability in their hybernation, that the stimulus of perpetual though less fervid day within the arctic circle causes them to perform the functions of foliation and fructification with a rapidity unknown in more temperate regions. Other plants which need longer time to perfect their fruit or woody fibre, terminate in succession according to their several constitutions as the latitude increases. Their place is only partially

LAKE TITICACA.

10

supplied by other speciet, which have in like manner their equatorial and polar limits. These are not, however, so numerous as those which die out, there being no rule more general than the decrease of generic and specific forms in passing from temperate zones to arctic or polar ones.

There is a similarity in many respects between the vegetation of alpine tracts and that of high latitudes, but not an identity, the condition of the two regions differing in some essential particulars. No more apt illustration of this fact is needed than that adduced by Meyen, of Titicaca. This alpine lake, situated on the plateau of Chuquito in southern Peru, at the elevation of 12,700 feet, is surrounded by a rich and beautiful vegetation, which flourishes under a perpetual spring. On its banks a populous community, inhabiting magnificent cities, is supported by a fertile soil, yet trees are wanting in the country; whereas we have seen that the pine forest extends in North America to the 69th parallel beyond the limits of the cerealia. On the shores of Lake Titicaca barley and oats grow, but wheat does not succeed, and maize is raised only by artificial heat. In respect of these products, therefore, its climate agrees with that of Fort Simpson on the 62nd parallel. Its summer heat, which, according to Meyen, ranges between 52° and 66° Fah., is more equable than that of Fort Simpson which has a mean daily summer temperature of about 59° Fah., with a mean at noon of 70° Fah.; and a range of from 90° Fah. down to below the freezing point.

It is necessary to remark, however, that the decrease of vegetable forms with an increase of latitude has more analogy to that which is observed on a lofty isolated mountain than on an elevated plateau; and plants actually grow on the summits of the White Mountains of New Hamp-

LANTS

r as the rease of Iaize. — — Their Prairies. Zone. in three

stricted months uperior ation of casional n July, retables United can be ughout in their gh less perform apidity s which re, teritutions artially

266 GEOGRAPHICAL DISTRIBUTION OF PLANTS.

shire which are not met with again until we reach the shores of the Arctic Sea.

The peculiarities of the climate of Canada and Rupert's Land may be in part shown by reference to a few of the plants usually cultivated for food. The *vine* would thrive with the summer heat of Fort Simpson were the season long enough; but the September and October heats, which are required to ripen its fruit, do not occur in any district of Rupert's Land; and the grape is destroyed by the severe night frosts which are frequent in autumn even in so low a parallel as the north shore of Lake Superior. The conditions essential to the due growth of the vine, mentioned by Meyen*, do not extend in the basin of the St. Lawrence beyond the 43rd parallel, while on the Rhine wine is a profitable production up to the 51st.

Maize is a plant which thrives best in the dampest and hottest tropical climates, where it brings forth eight hundred-fold. Its culture extends into temperate regions, but with a greatly diminished yield; and it is cultivated near its northern limit only as a green vegetable, the grain seldom ripening, and being eaten in its milky state. This is its condition in most parts of Great Britain, when reared in the open field. On the western shore of Europe it is not cultivated beyond 46°, though in the valley of the Rhine it extends to 49° north lat. In South America, on the Chili coast, it is planted as low as 40° south lat.; and on

• This author states that the culture of the vine is regulated more by the length of summer than by its high temperature, though the latter is also an element in the proper ripening of the grape. It will succeed, he says, under every tropical heat, provided the atmosphere be not too moist. It thrives well under a mean heat of 60° Fah.; it ripens with a lower mean heat of 48° Fah., and a summer heat of 67° Fah., but the juice contains less sugar and yields less alchohol. (Meyen, Geogr. of Plants.)

WHEAT.

1 the

pert's of the thrive season which istrict oy the ven in . The ationed wrence ne is a

est and nt hunons, but ed near seldom is is its ared in t is not Rhine on the and on

> ed more ugh the It will hosphere Fah.; heat of lchohol.

the Peruvian plateau, at the height of 12,000 feet, above which it requires artificial shelter and warmth. A profitable return can be obtained from it in Rupert's Land between the 49th and 51st parallels, where, however, the vine does not accompany it as on the banks of the Rhine. Garden cultivation and shelter from spring frosts would extend its cultivation in Rupert's Land even higher than in On the fertile acclivity of Young Street which England. leads from Toronto to Lake Simcoe, and crosses the 44th parallel of latitude, we may behold heavy crops of maize, and cucumbers and gourds, ripening, in the same field, with but little expenditure of care or labour, though the mean annual heat, being 41° Fah., is inferior to that of the Orkney and Shetland Islands, where barley, one of the most northern of the cereals, grows imperfectly. The summer heat of Young Street, however, exceeds that of any part of the British Isles.

Wheat is the cereal which requires most heat of those usually cultivated in England. Its culture is said to ascend to 62° or 64° north lat. on the west side of the Scandinavian peninsula, but not to be of importance beyond the 60th. On the route of the Expedition it is raised with profit at Fort Liard in lat. 60° 5' north, long. 122° 31' west, and having an altitude of between 400 and 500 feet above the sea. This locality, however, being in the vicinity of the Rocky Mountains, is subject to summer frosts; and the grain does not ripen perfectly every year, though in favourable seasons it gives a good return. At Dunvegan, on Peace River, lying in lat. 56° 6' north, long. 117° 45' west, and at an altitude of 778 feet, the culture of this grain is said to be equally precarious. It grows, however, freely on the banks of the Saskatchewan, except near Hudson's Bay, where the summer temperature is too low. From Mr. M' Pherson I learnt, that on the west side of

C
the Rocky Mountains good crops of wheat are raised with facility at Alexandria, on Frazer's River, in lat. 52° 30' north, long. 122° 40' west, and 300 or 400 feet above the sea; also at Fort George, on the same river, more than a degree further north, and 100 feet higher.

At Fort James, on the borders of Stuart's Lake, in lat. $54\frac{1}{2}^{\circ}$ north, in a mountainous region near the source of Frazer's River, wheat continues to grow, but often suffers from the summer frosts. In these quarters the grain comes to maturity in about four months. In the colony of Red River its growth is luxuriant, though the upper part of that country, which touches the 49th parallel of latitude, is elevated about 1,000 feet above the sea. Periodical ravages of grasshoppers, however, frequently destroy the hopes of the husbandman.

At Fort Francis, situated on the banks of Rainy River in lat. 48° 36' north, long. 93° $28\frac{1}{2}$ ' west, wheat is generally sown about the 1st of May, and is reaped in the latter end of August, after an interval of about 120 days. In 1847 multitudes of caterpillars spread like locusts over the neighbourhood. They travelled in a straight line, crawling over houses, across rivers, and into large fires kindled to arrest them. Throughout the whole length of Rainy River, on the Lake of the Woods, and on the River Winipeg, they stripped the leaves from the trees, and ate up the herbage. They destroyed the folle avoine on Rainy Lake, but left untouched some wheat that was just coming This was the first time that Fort Francis had into ear. experienced such a visitation. When we passed that way in 1848, the still leafless trees were covered with the coccoons of last year, in each of which there remained the hairy skin of a caterpillar.

On the island of Sitka, lying in 57°-58' north latitudes, though the forest, nourished by a comparatively high mean l with 2° 30' ve the than a

in lat. rce of suffers grain colony upper allel of Periodlestroy

liver in nerally e latter vs. In ver the crawlkindled Rainy River and ate Rainy coming cis had at way th the hed the

> itudes, 1 mean

temperature and a very moist atmosphere, is equal to that of the richest woodlands of the northern United States, yet corn does not grow.

In the middle temperate zone of France wheat is cultivated to the height of 5,400 feet only. In Mexico its culture commences at the altitude of 2,500 or 3,000 feet, and ascends to more than 9,000 feet. On the plateau of southern Peru, 8,000 feet above the sea, its yield is extraordinary; and on the foot of the volcano of Arequipa it succeeds as high as 10,000; but it will not grow in the equable temperature of Lake Titicaca, the heat there not being sufficient to ripen either it or rye. It requires for its growth, says Meyen, the mean annual heat of 39° Fah., combined with a summer heat of 56° Fah. : a much inferior mean heat is, however, sufficient in the extreme climate of subarctic America, provided the summer heat for 100 or 120 days be great enough.

Oats are little cultivated in Rupert's Land; they require longer time than barley to ripen, and are therefore not likely to grow so far north. They have not been tried at Fort Norman, however, which is the most advanced post in that direction where barley is cultivated. Mr. M⁴ Pherson finding some grains of oats accidentally in a barley field, propagated them, and raised some good crops on the River of the Mountains, and I believe also at Fort Simpson. On the Scandinavian peninsula this grain is said to extend to $62\frac{1}{2}^{\circ}$ N. and 65° N., but, even on the latter parallel, falling five degrees short of the latitude which barley reaches. Meyen saw ripe oats at Lake Titicaea.

In good seasons *barley* ripens well at Fort Norman on the 65th parallel, as has been mentioned in the narrative (l. p. 165.). All Mr. Bell's attempts to raise it at Fort Good Hope, two degrees further north, failed. It reaches, as we have just observed, the 70th degree of latitude on







the Scandinavian peninsula, and it is cultivated for green fodder in Peru up to the height of 13,800 feet, but seldom ripens its grain higher than 10,000 feet. (Meyen.)

Potatoes, which have been cultivated from time immemorial on the banks of Lake Titicaca, yield abundantly at Fort Liard, and grow, though inferior in quality, at Fort Simpson and Fort Norman. They have not succeeded at Fort Good Hope, near the 67th parallel. At the latter place turnips in favourable seasons attained a weight of from two to three pounds, and were generally sown in the last week of May. At Peel's River the trials made to grow culinary vegetables had no success. Nothing grew except a few cresses. Turnips and cabbages came up about an inch above the ground, but withered in the sun, and were blighted by early August frosts.

In the preceding narrative, as well as in the geographical sketch, we have had frequent occasion to allude to several great divisions of North America, each of which has a peculiar physiognomical character in its vegetation.

1st. The eastern woodland country constitutes the first division, in which the forest extends from the Atlantic westward till it meets the great prairies.

2nd. The second division lies to the north of the forest lands, and is appropriately named the "Barren Grounds.' This tract has its greatest north and south extension on the eastern coast. On the shores of Hudson's Bay and the Welcome it reaches from the 60th or 61st parallel to the extremity of the continent, but narrows to the westward; since the boundary line of the woods takes a diagonal or north-east direction from the 91st meridian, and, before reaching the 120th, has risen to the 67th parallel. Further to the west the Barrens Ground form a border to the Arctic Sea of greater or less breadth according to the northerly prolongation of the continental pro-

DISTRICTS OF VEGETATION.

montories, since the southern limit is nearly coincident throughout with the arctic circle, on which it approaches Beering's Straits — clumps of spruce fir*, the usu~l outliers of the forest, having been observed on Buckland or Noatak River which falls into Eschscholtz Bay. The fertile alluvial deposits of the well-sheltered valley of the Mackenzic interrupt the continental continuity of the Barren Grounds by carrying the woods nearly to the sea shore; but there seems to be no other material indentation of the barren district; and even on the Mackenzie the valley is bridged, as it were, by the naked summits of the alpine ridges.

3rd. The prairie slope forms a third physiognomical district of vegetation, which has the greatest transverse expansion on the Missouri, and, narrowing as it goes north, runs out on the 60th parallel, having, after passing the Saskatchewan, been much indented by the woods which feather the numerous rivers that drain the declivity. These prairies have much analogy with chalk downs in aspect as well as in mineral constitution.

4th. The Rocky Mountain chain, and the alpine ranges and isolated peaks which rise to the westward of it, may be considered as a fourth district which nourishes some peculiar species of plants.

5th. And the lower woodland country on the Pacific side of the range forms a fifth.

If we trace any one of these districts northwards, making due allowance for the varying altitude of the country above the sea, we may ascertain the effect of increase of latitude on the vegetation of that meridian; but, if we compare one district with another, we must keep in view the climatological fact of the rise of the isothermal lines in proceeding westward. The course of the forest boundary is one illus-

* The species is doubtful.

green ldom

emoly at Fort led at latter ght of n the grow except out an l were

phical several has a

e first tlantic

forest unds.', ension Bay st paows to takes ridian, 67th orm a ccordl pro-

tration of this phenomenon; and we have another in the range of certain species or forms constituting that forest. Thus the *Cupressus thyoides* is rare beyond the 49th parallel in the eastern district, and terminates altogether along with the *Thuya occidentalis* on the 53rd, while the magnificent *Cupressus* or *Thuya nutkatensis* adorns the forests of Norfolk Sound on the 58th parallel of the Pacific coast. The distribution of the *Pinus inops*, *Abies canadensis*, *Rubus nutkanus*, and of some other conspicuous trees and shrubs, show that the vegetation of the district of Sitka on the north-west coast is equal and similar to that of the eastern states of Wisconsin or Minesota eight degrees further south.

The physiognomy of the woodland district through which the canoe route lies has been incidentally touched upon in the descriptions of several localities that have been introduced into the narrative, yet it will not be out of place to recall its general features here. Of this district, which has a breadth of about 600 geographical miles between the 50th and 55th parallels, the white spruce is the most abundant and characteristic tree; yet up to the 54th parallel it is conjoined, and especially on the banks of rivers, with other trees which break the monctony of the dark evergreen forest. Beyond the banks of the Saskatchewan the oaks, elms, ashes, maples, bass-wood, white thorns, Virginian clematis, and various other trees and shrubs cease to grow; and the white spruce may be said to cover the face of the country, except on the alluvial borders of rivers and lakes, where the aspen, balsam poplar, balsam fir, alder, and multitudes of willows usurp its place, or on the edges of swamps where the black spruce leads a lingering, unhealthy existence. With the black spruce the larch is often associated; though it is not confined to morasses, yet it is too much isolated in its distribution to produce a

WOODY DISTRICT.

in the forest. 49th ogether ile the rns the Pacific *cana*picuous district milar to ta eight

h which upon in n introplace to , which veen the he most 4th paf rivers, he dark tchewan thorns, shrubs o cover borders balsam e, or on lingerhe larch orasses, oduce a

difference of tint sufficiently massive to please the eye, except in a very few localities. The Banksian pine is more frequently seen in considerable patches, and its appearance is agreeable to the voyager; for, independent of the fact that its spreading branches and general form, resembling that of the Scotch fir, is a rest to the eye wearied with the tapering stiffness of the spruce, it offers the prospect of a dry and comfortable encampment. It always grows in a sandy soil, and is remarkable among the Rupert's Land trees for its freedom from underwood. Not so the white spruce, which admits of a thick undergrowth of willows, cornel bushes, viburnum, roses, brambles, and gooseberries; and, in the country south of Lake Winipeg, of maples, American yews, and many other shrubs and trees. The willows, especially when conjoined with the falling or inclined stems of forest trees — the growth of bygone centuries, form a barrier to the progress of a white man in the forest; but the slim and agile native glides through the tangled thicket with a noiseless and ghostlike case, impassive to the annoyance of the moscheto clouds that darken the air. The prickly twining Panax horridum, which interlaces and arms the brushwood on the north-west coast up to the 58th parallel, has no representative on the east side of the continent, except perhaps the Aralia hispida, which, though of the same family, has feeble defences, and is not a climber. The Cratagi have the most offensive weapons of any of the shrubs in Rupert's Land.

Even beyond the Saskatchewan, where the maples, ampelopsis, and some other trees and shrubs whose leaves assume the orange and red tints before they fall, cease to grow, the river banks are enlivened by the bright purplish shoots of the white cornel berry (osier rouge) and the gay spires of the *Epilobium angustifolium*, which rise above a VOL. II.

man's height in the alluvial deposits, and are varied also by other shrubs that have been noticed in the descriptions interspersed through the preceding pages. These are merely the foreground incidents, however; the sombre spruce everywhere forms the back ground.

The agency of man is working a change in the aspect of the forest even in the thinly peopled north. The woods are wasted by extensive fires, kindled accidentally or intentionally, which spread with rapidity over a wide extent of country, and continue to burn until they are extinguished by heavy rains. These conflagrations consume even the soil of the drier tracts; and the bare and whitened rocks testify for centuries to the havoc that has been made. A new growth of timber, however, sooner or later springs up; and the soil, when not wholly consumed, being generally saturated with alkali, gives birth to a thicket of aspens instead of the aboriginal spruce.

The frozen subsoil of the northern portions of the woodland country does not prevent the timber from attaining a good size, for the roots of the white spruce spread over the icy substratum as they would over smooth rock. As may be expected, however, the growth of trees is slow in the high latitudes. On the borders of Great Bear Lake, four hundred years are required to bring the stem of the white spruce to the thickness of a man's waist. When the tree is exposed to high winds, the fibres of the wood are spirally twisted; but in sheltered places, or in the midst of the forest, the grain is straight, and the wood splits freely.

At the limit of the woods the white spruce is everywhere the most advanced tree, growing either solitarily, with its branches clinging to the ground and its dwarfed top bent from the blast, or in small clumps in some favourable locality. The *Salix speciosa* may indeed be said to pass

WOODY DISTRICT.

l also by ations ine mercly e spruce

rs.

aspect of he woods ly or inde extent are extinconsume whitened een made. er springs ing genethicket of

the wooduttaining a d over the As may be h the high four hunthe white h the tree d are spithe midst wood splits

verywhere y, with its d top bent favourable id to pass beyond the spruce; but it does so only on the alluvial points of rivers, and not in its tree form.

Though the species of plants become less numerous as we advance northwards through the woody region, there is no falling off in the number of individuals of the species that remain. For not only is the forest crowded, and often almost impenetrably so, when the trees are young, but on the margins of rivers, and other open places, there is a dense herbaceous vegetation, which clothes the ground in Rupert's Land as perfectly as it is covered in a lower latitude, though the vegetation be less rank. On the inundated alluvial flats tall carices grow as closely as they can stand, and furnish an abundance of nutritious hay. There is, however, a total absence in the north of the Lianas, Tillandsia, and parasitic Orchidea, which impart so peculiar an aspect to the forests in some of the warmer districts of the earth. The great hedge bindweed, (Calystegia,) the Virginia creeper, the hop plant, and the twining herbaceous Smilacina, with its grape-like clusters of blackberries, disappear on the south side of Lake Winipeg, and the only aerial parasite in the north is the leafless Arceuthobium oxycedri, which seats itself on the branches of the Banksian pinc. The graceful Usneæ which hang from the branches of the ancient black spruces in long, thread-like hanks, have, it is true, some resemblance to the Tillandsiæ which forms an elegant drapery to the ever-green oaks of Georgia and Florida.

In the castern woodland district, from the St. Lawrence to the Saskatchewan, the *Composita* are the most numerous family of plants, and they form between the sixth and seventh of the whole phænogamous vegetation. Next to them come the *Cyperacea*, which, owing to the great development of the genus *Carex*, constitute more than oneninth of the *Phanerogamia* of the district.

т 2

In the second, or barren ground district, in places where the soil is formed of the coarse sandy débris of granite, and is moderately dry, the surface is covered by a dense carpet of the Corniculariæ tristis, divergens, ochroleuca, and pubescens, mixed in damper spots with Cctrariæ cucullata and islandica. In more tenacious soils other plants flourish; not, however, to the exclusion of lichens, except in tracts of meadow ground. The Rhododendron lapponicum, Kalmia glauca, Vaccinium uliginosum, Empetrum nigrum, Ledum palustre, Arbutus uva ursi, Andromeda tetragona, and several depressed or creeping willows, lie close to the soil, their stems short, twisted, and concealed, with only the summits of the branches showing among mosses or lichens. Here and there, on the moister sides of the hills, there is a gay display of saxifrages, pediculares, or primroses; and a few of the sandy spots on the coast are enlivened by a beautiful dwarf phlox or a handsome dodecathcon. On the alluvial banks of rivers only are willows of erect growth to be found, and of these the Salix speciosa is the most robust and the handsomest. On the sandy shore of the sea the Pisum (or Lathyrus) maritimum, the Polemonium cæruleum, various blue and yellow Astragali, and several Artemisiæ flourish. Most of these plants also occur, though more sparingly, in the interior. One circumstance which came under my observation, and has been cursorily alluded to in the Narrative (I. p. 322.), is the existence of very ancient stumps of trees, either solitary or grouped, in various places of the barren grounds, seemingly the vestiges of the forest, which had spread more widely over the country some centuries ago than in the present day. Further evidence that such was the case may be obtained in the extension of Pyrolæ, and some other woodland plants, to the coasts of the Arctic Sca. On the sheltered banks of rivers, even in the barren grounds, clumps of living trees occasionally occur; but the

BARREN GROUND DISTRICT.

277

cs.

es where te, and is carpet of pubescens, nd islanish; not, tracts of 1, Kalmia Ledum paad several soil, their e summits ns. Here e is a gay id a few of beautiful he alluvial wth to be robust and the Pisum m, various æ flourish. aringly, in under my the Narrastumps of accs of the rest, which nturies ago t such was Pyrolæ, and Arctic Sea. the barren ur; but the stumps I speak of stand often on the exposed side of a hill, and indicate a deterioration of climate, however that may have been produced. We saw no young firs growing up in such situations to leave similar vestiges in a future age.

In many sheltered valleys on the sea-coast, and even in the more elevated interior, especially where a fertile soil has been formed by the decomposition of trap rocks, there is a good growth of grasses, several of which flourish well on lands that are occasionally inundated by the sea. Among these are Elymus mollis, Spartina cynosuroides, Calamagrostis stricta, Carices stans, compacta, glarcosa, membranacea, and livida, Colpodium, Deschampsia, Festuca, and several Poa. In some of the maritime meadows to which the reindeer resort to bring forth their young, there are treacherous mud-banks, which are soft enough and deep enough to swallow up a deer or musk ox, that may rush heedlessly into them when chased by a wolf; but in general the frozen subsoil is so near the surface as to preclude any such accident. The existence of these boggy places, which were seen only on the sca-coast, scarcely affords a satisfactory solution to the problem of the entombment of a living elephant or rhinoceros, and the subsequent preservation of the entire carcass in the frozen soil. But in whatever manner this may have taken place, I should infer, from the economy of the arctic regions, that these animals were migratory, like the reindeer of the present day, and wintered in milder climates.

On approaching the arctic circle the relative proportion of the *Compositæ* greatly decreases, and that of the *Cyperaceæ* increases within the woody tracts, though it falls off on the barren grounds. Taking the whole zone between the arctic circle and the extremity of the continent, which includes much woodland, the *Cyperaceæ* are the most

т З

numerous family of plants, and are more than double the Gramineæ. The Cruciferæ come next to the Cyperaceæ in this zone. In the polar regions beyond the continent, the Cruciferæ take the first place in respect of number of species, then come the Gramineæ, which are closely followed by the Saxifrageæ.

The third, or prairie district, has the prevailing aspect of a grassy plain, the herbage, however, having a considerable intermixture of carices among the true grasses. The herbage grows up rather wiry in the dry summers of that region; but, in consequence of the fires that frequently spread over vast tracts, a young growth takes place, to which the bison and deer resort. On the Arkansas, the "buffalo or bison grass" is the Sesleria dactyloides. Whether this species extends to the Saskatchewan or not. I am unable to say: we certainly did not gather it there; but at the time that Mr. Drummond and I visited that part of the prairie, recent fires had made flowering specimens of grasses very rare. Of the phænogamous prairie plants actually collected, the Gramineæ form about the eleventh, and the Cyperaccæ the sixteenth. On the plains the Compositæ are numerous and showy; there is a considerable variety of handsome Leguminosæ, with some pretty Boragineæ; and the Artemisiæ, owing to the quantity of surface they cover, though the species are not numerous, contribute greatly to the hoary aspect of the prairie vegeta-The Rosuceae vie with the Cyperaceae in number of tion. species; but many of them are fruit-bearing shrubs, growing on the banks of the rivers that serpentine among alluvial points, in channels sunk deeply below the general surface of the prairie.

Between the 32nd and 33rd parallels, on the Gila and Rio del Norte, west of the Rocky Mountain ridge, Colonel Emory gathered many examples of *Cacti*, of which Dr. Engelman

PRAIRIE DISTRICT.

279

.

ble the peraceæ ntinent, mber of ely fol-

spect of iderable The of that quently lace, to isas, the ctyloides. n or not. it there; ted that ng specis prairie bout the he plains is a conne pretty antity of umerous, e vegetaumber of growing g alluvial al surface

> ı and Rio el Emory Ingelman

has described fifteen species belonging to the genera Mammillaria, Echinocactus, Opuntia, and Cereus. Among these, the Pitahaya, or Cereus giganteus, is the most remarkable, as it grows in the shape of a candelabra, or Titanic tuningfork, with three or four prongs, to the height of sixty feet. Cacti are numerous also on the eastern side of the mountains in the same parallel; and the smaller kinds, chiefly Opuntia, range northwards over the prairies to the 49th parallel, and perhaps still further north. We gathered Opuntia glomerata, or the crapaud verd of the voyagers, on the Lake of the Woods; and a species of the same section of the genus attains an equally high parallel on the Pacific coast.

With the physiognomy of the vegetation on the Rocky Mountains, and of the district to the west of that range, I have no personal acquaintance, and borrow the following notice of the vegetation of Sitka from Bongard. Sitka is situated in the entrance of Norfolk Sound, on the 57th parallel, near an extinct volcano, named Mount Edgecumbe, which marks the entrance of the sound. The most remarkable mountain in the immediate vicinity of the settlement is Westerwöi, which is 3,000 Parisian feet in height, and is clothed to its summit by a dense forest of pines and spruces, some of which acquire a diameter of seven feet, and the prodigious length of 160 feet. The hollow trunk of one of these trees formed into a canoe, is able to contain thirty men, with all their household effects. The climate of Sitka is very much milder than that of Europe on the same parallel. The cold of winter is neither severe nor of long continuance; but the atmosphere is charged with vapours, whose condensation occasion almost constant rains. In the month of July the sun is seldom visible on more than three or four days, and then only for

т 4

an instant. This humidity gives astonishing vigour to the vegetation, yet corn does not grow there; and, in fact, the want of level surface is an impediment to cultivation. In six weeks the botanists collected 222 species of plants, of which thirty-five were new to science.

Of the Polar plants, amounting to ninety-one species, which inhabit Melville Island, the shores of Barrow's Straits and Lancaster Sound, and the north coasts of Greenland, between the 73rd and 75th parallels of latitude, about seven-ninths range to Greenland, Lapland or Northern Asia. Of the remainder, some have been gathered on the shores of the Arctic Sea from Baffin's Bay to Beering's Straits; and it is probable that if these high latitudes were fully explored, the flora of the entire zone would be found to be uniform. Some of the more local plants will perhaps be ascertained, on further acquaintance, to be mere varieties altered by peculiarities of climate. That the flora as well as the fauna in the high northern latitudes is nearly alike in the several meridians of Europe, Asia, and America, has long been known. And even when we descend to some distance south of the arctic circle, we find that this law is superior to the intrusion of high mountain chains, and is but partially infringed upon. In taking the St. Lawrence basin for instance, if we allow for the rise of the isothermal lines on the west coast, and make our comparisons in an oblique zone, including Sitka and Wisconsin, we shall find that there is much similarity in the floras on the two sides of the continent. The Rocky Mountain ridge is not by any means a boundary to the peculiar vegetable forms of the Pacific coast; on the contrary, many of them cross the ridge to its eastern declivity, though they do not descend into the low country; and there is actually more similarity between the vegetation of

POLAR PLANTS.

ir to the fact, the ion. In lants, of

rs.

species, Barrow's f Greenlatitude, land or ve been Baffin's that if rn of the e of the n further uliarities the high neridians n. And the arctic rusion of ed upon. we allow oast, and ing Sitka similarity he Rocky ry to the the condeclivity, try; and etation of the prairies of Oregon, and those of the Missouri and Saskatchewan, than there is between the latter and the eastern parts of the United States and Canada. In still more southern latitudes the case may be different; and Ehrenberg has found a totally different group of *Infusoria* in California from that which exists on the east side of the continent; the Rocky Mountains, in his opinion, proving a complete barrier to these organisms.

The families of Polar plants which are most rich in species are the Crucifera, Graminea, Saxifragea, Caryophyllea, and Compositæ. Of these the Saxifrageæ are most characteristic of extreme northern vegetation. All of them that inhabit the 74th parallel in America are found also in Spitzbergen, Lapland, or Siberia; and even the polar species are twice as numerous as those which exist in the wide district which Gray's "Flora of the Northern States" comprehends. If we reckon all that enter the arctic circle, we shall find them to be four times as many as those which Dr. Gray enumerates; and we may add that the plant which Humboldt traced highest on the Andes was a saxifrage. The Caryophylleæ and Cruciferæ, which vie with the saxifrages in number on the 74th parallel, include many of the doubtful local species above alluded to. Of the most northern Gramineæ, about one half are, as far as we yet know, exclusively American; the few species which the other families contain have as extensive a lateral range as the saxifrages.

Arctic zone.—On descending to the main land from the 71st parallel down to the arctic circle, including a zone of four degrees of latitude, we find that the species have increased eight-fold in number, and there is a large addition of generic forms, as might be expected on entering within the limits of the forest.

281

Ranunculaccæ Papaveraccæ Cruciferæ Caryophylleæ Leguminosæ Rosaceæ Onagrariæ Saxifrageæ

The Polar familics are-

Compositæ Cichoraceæ Eupatoriaceæ Senecionideæ Campanulaceæ Ericeæ Polemoniaceæ Scrophularineæ Polygoneæ Salicaccæ Junceæ Cyperaceæ Gramineæ Lycopodineæ Equistaceæ Cryptogamia

In addition to the above the following enter the Arctic

Sarracenieæ Fumariacea Violarieæ Droseraceæ Polygaleæ Lineæ Balsamineæ Celastrineæ Halorageæ Ceratophylleæ Portulaceæ Crassulaceæ Grossularieæ Umbelliferæ Araliaceæ Corneæ

A

Circle :---

Caprifoliaceæ Valerianeæ Compositæ Asteroideæ Vaccinicæ Monotropeæ Gentianeæ Diapensiaceæ Hydrophylleæ Boragineæ Orobanchem Labiatæ Verbenaceæ Primulaceæ Plumbagineæ Plantagineæ

Chenopodicæ Eleagneæ Santalacese Empetrem Urticeæ Betulaceæ Coniferæ Juncagineæ Aroideæ Naiades Smilaceæ Melanthaceæ Asphodeleæ Orchideæ Irideæ Filices

I made a pretty full collection of lichens and mosses within the arctic circle; but since so many of them are almost cosmopolites, and a still greater number are common to both the temperate and frigid zones, under similar conditions of moisture and exposure, I have avoided swelling the lists with their names. *Fungi* are not wanting in the northern regions, but the difficulty of preserving them prevented me from gathering many. All the families in the above two lists are represented in England, except *Diapensiaceæ*, which is a Lapland form; and *Sarracenieæ* and *Araliaceæ* which are more purely American.

Between the arctic circle and the south side of the Winipeg or Saskatchewan basin on the 50th parallel, embracing

ST. LAWRENCE PLANTS.

the entire width of the continent, the following families make their appearance : --

Berberideæ Nymphæaceæ Capparideæ Cistineæ Malvaceæ Tiliaceæ Hypericineæ Accrineæ Accrineæ Ampelideæ Geraniaceæ Oxalideæ Rhamneæ Terebinthaceæ Cucurbitaceæ Loranthaceæ Paronychieæ Jasmineæ Apocyneæ Asclepiaceæ Convolvulaceæ Solaneæ Amaranthaceæ Aristolochieæ

Euphorbiaceæ Ulmaceæ Cupuliferæ Myriceæ Liliaceæ Alismaceæ Pontederiaceæ Restiaceæ Hydrocharideæ Marsiliaceæ Salvinaceæ

The families which reach the St. Lawrence basin, but do not extend northwards to the Winipeg valley, or enter the western prolongation of that zone, are—

Menispermaceæ Podophylleæ Limnanthaceæ Oxalideæ Rutaceæ Lythrarieæ Cacteæ Hamamelideæ Compositæ Vernoniaceæ Acanthaceæ Nyctagineæ Phytolacceæ Laurineæ Saurureæ Juglandaceæ Platanaceæ Camelineæ Hypoxideæ Dioscoreæ

To give a further view of the accession of families in going southwards, the following are added from Dr. Gray's "Botany of the Northern States": —

Magnoliaceæ Anonaceæ Cabombaceæ Resedaceæ Elatinaceæ Anacardiaceæ Hippocastanaccæ Melastomaceæ Hydrangeæ Aquifoliaccæ Ebenaceæ Bignoniaceæ Nyssaccæ Podostomcæ Balsamifluæ Amaryllidaceæ Hæmodoraceæ Xyridaceæ

In tracing individual species to their northern limits, we did not discover in any one instance that the crest of a water-shed between successive transverse river basins was a boundary to the plant. Many of the more remarkable trees, oaks, &c. flourish in the neighbourhood of Rainy Lake and on the upper part of Red River, but die

Arctic

mosses m are mmon r convelling in the m prein the ; *Diaea* and

Winiracing 283

out on approaching the south end of Lake Winipeg. Others go a degree or two further north to the banks of the Saskatchewan, about Cumberland House, and there make their last appearance: among these are the ashes, elins, and maples. Some which are not seen beyond that locality on the canoe route, go three or four degrees further north on the western side of the prairies, in the sheltered valleys of the Rocky Mountains. In these valleys also the lamented Drummond found a considerable number of the species of the Pacific coast, their range not being cut short by the dividing ridge, but being seemingly more effectually limited by the dry prairies. It is unfortunate that the vertical limits of the species gathered by Drummond in the mountains were not noted, as a careful list containing that element, and which no one was more able than he to make. would have conveyed much information with respect to the distribution of plants. The statistical enumeration of the mountain species, collected between 52° and 57°, in the subjoined table evidently contains a mixed flora; some families having an arctic, almost a polar character; others a subarctic, or almost temperate one.

LIST OF TREES AND SHRUBS.*

RANUNCULACE. — Clematis vi yiniana is common to Oregon, the eastern United States, and Canada, and extends northwards to the Saskatchewan.

BERBERIDEÆ. - Berberis vulgaris has been found in Canada,

* I am indebted to Dr. Asa Gray for some valuable information respecting the range in the United States of some of the trees in the following list. The Northern States referred to in the list extend from New England to Wisconsin, and south to Ohio and Pennsylvania, inclusive.

Newfoundland, and New England, and is considered as having been introduced from England. The pinnate-leaved barberries or *Mahoniæ* are natives of Oregon, and perhaps extend northwards into New Caledonia or Vancouver's Island.

CISTINE E. — Hudsonia tomentosa grows in New Jersey and Canada, cz the borders of all the great lakes, and onwards on the canoe route to Clear-water River on the 57th parallel, beyond which it was not observed.

TILIACE. — *Tilia glabra*, the lime tree, white wood, or bass wood, is a familiar ornamental and useful tree in the United States and Canada. We observed it as far north as Lake Winipeg, but only as underwood, sending out long flexible branches, which the natives convert into temporary cordage.

ACERINE A. — Acer montanum (vel spicatum), the mountain maple, has a range from Maine, Pictou, Wisconsin, and Minesota, to the River Winipeg, and, from the beautiful orange and red tints which its leaves assume in decay, is a great ornament to the woods in autumn. A. circinatum is confined to the west coast, is common in Oregon, and extends to the British territory on that side of the mountains. It grows in the woody country only, and chiefly in the pine forests, where its pendulous branches, taking root, form almost impenetrable thickets. The close-grained tough wood is used by the natives for making hoops.

A. saccharinum, sugar maple, with the variety, or perhaps species, named A. nigrum by Michaux, has been traced by Dr. Asa Gray along the Alleghany Mountains to Gcorgia. In the low country it scarcely passes to the south of Pennsylvania, but on the west side of the valley of the Mississippi is found as far south as Arkansas. Its northern limit is a short way beyond the 49th parallel on the elevated southern water-shed of Lake Winipeg; but it may, perhaps, attain a greater northern latitude in the lower country of Canada. A little to the south of Rainy Lake it yields abundance of good sugar. The variety named bird's-eye maple grows on one of the islands of the Lake of the Woods, and has been employed for making gun-stocks. Goat Island, at the Falls of Niagara, according to Mr. David

Others : Sasmake elins, cality north alleys iented cies of by the imited ertical moung that make, to the of the in the some hers a

regon, wards

anada,

mation in the extend Penn285

Douglas, nourishes some of the largest sugar maples in North America.

A. rubrum, red or swamp maple, ranges southwards, according to Dr. Gray, to Florida, and round the whole Gulf of Mexico to northern Texas; but some of the southern forms, he says, would probably be considered by European botanists to be specifically distinct from the northern tree. It grows in Nova Scotia, throughout Canada, westward to Lake Winipeg and the Rocky Mountains on the 52nd or 53rd parallel, and also crosses that chain to the head waters of the Columbia.

A. pennsylvanicum, striped maple, or moose wood, comes down along the coast to Boston, and follows the mountains from Pennsylvania to the borders of Georgia, to which Dr. Gray has traced it. It grows also in Kentucky, and was seen by us on the banks of the Winipeg, where it has more the character of a flexible willow than of a tree. A. macrophyllum is confined to the mountainous country on the Pacific up to the 50th parallel, and is one of the most graceful trees, rising to the height of ninety feet, with a circumference of sixteen. A. dasycarpum, white or silver maple, is a fine large ornamental tree, well known in the United States. It is found on Lake Huron, but does not appear to rise northwards out of the St. Lawrence basin. Good sugar is made from the juice of this tree.

Negundo fraxinifolium or aceroides, ash-leaved maple, does not, to Dr. Gray's knowledge, grow wild in New England. It abounds in Pennsylvania, and extends westwards to western Texas and the Rocky Mountains, growing at the high elevation of 6,000 or 7,000 feet near Santa Fé and the Pawnee Fork, according to Lieut. Alert. It terminates northwards about the 54th parallel on the banks of the Saskatchewan, and is the tree which yields most of the sugar made in Rupert's Land. Though this product varies much with the skill of the operator, the kind obtained from the juice of this tree is generally of a darker colour than that which the true sugar maple yields.

AMPELIDE *Ampelopsis quinquefolia*, Virginia creeper, or American ivy, extends northwards to Lake Winipeg, and is a great ornament to the protruding rocks over which it creeps.

287

It is a familiar shrub in the Northern States, but I have not been able to ascertain its southern limit. The Vitis cordifolia or riparia, frost grape, grows, on the evidence of collections made on my former journeys, as far north as the south end of Lake Winipeg, on the 50th parallel. I did not observe it on my late voyage, in which, indeed, I had very little leisure to search for plants; and if it actually grows in so high a latitude, it does not produce edible fruit so as to attract the attention of the residents, who could give me no information concerning it. Together with the Vitis astivalis, or summer grape, it is common in Wisconsin and Minesota. Some of the native American vines are cultivated in the eastern states; and the Isabella grape, a variety of the Vitis labrusca, has an agreeable though peculiar flavour.

ZANTHOLACE ... Zanthoxylum americanum, northern prickly ash, and Ptelea trifoliata, the shrubby trefoil, grow in Canada and Wisconsin, where they seem to find their northern limit.

CELASTRINE E. — Staphylea trifolia, bladder nut; Euonymus atropurpureus, burning bush; E. americanus, strawberry bush; and Celastrus scandens, inhabit Canada and Wisconsin, but were not observed to the north of Lake Superior. E. atropurpureus crosses the continent to Oregon.

RHAMNEÆ. — Rhamnus alnifolius, alder-leaved buckthorn, grows from Maine and Michigan northwards, to about the 58th parallel. It is a low shrub, and is applied to no economical purpose. R. purshianus is an Oregon plant, which extends to Vancouver's Island and New Caledonia. Ceanothus americanus, New Jersey tea, ranges from Maine, Michigan, and Wisconsin, to Canada West, but was not gathered by us to the north of Lake Superior. The C. lævigatus, a west-coast species, which extends from Oregon to Vancouver's Island, seems to be the only member of the genus that enters the British territory. C. sanguineus is common in the valley of the Columbia, and crosses the mountains to the upper tributaries of the Missouri, forming one of the many instances of west-coast plants traversing the dividing ridge to the eastern prairies, but not extending to the eastern woodland districts.

North

to be not not the crosses

comes untains r. Gray n by us aracter confined parallel, sight of *carpum*, known ut does e basin.

le, does nd. It western evation e Fork, s about d is the s Land. perator, lly of a ds. creeper, and is creeps.

TEREBINTHACEÆ, or ANACARDIACEÆ. — Rhus radicans or toxicodendron, the poison oak; R. aromatica, the fragrant sumach; and R. glabra, the smooth sumach, reach the banks of the Saskatchewan, or latitude 45°. R. typhina and R. venenata extend to Canada, but have not been discovered north of Lake Superior.

LEGUMINOS.E. — Amorpha fruticosa, false indigo; A. canescens, lead plant; and A. nana, grow abundantly on the prairies of Osnaboya, and are the only shrubby leguminous plants which extend to Rupert's Land. Robinia pseudacacia is plentiful in Canada East.

CÆSALPINEÆ. — Gymnocladus canadensis, the Kentucky coffee-bean tree; Cercis canadensis, red bud; and Cassia chamæcrista, partridge pea, have their northern limits in Canada or Wisconsin.

ROSACE E. - Prunus americana, the wild yellow plum, seems to reach its northern boundary on the River Winipeg, not having been observed by me beyond the 50th parallel. Lieut. Abert gathered its fruit as far south as the banks of the Canadian and Pawnee forks of the Arkansas, on the 40th parallel. It is a common bush on the river banks in the Northern States. The American plums and cherrics require further investigation, as the number of the species and their distinctive characters are imperfectly known. This one grows to the height of ten or fifteen feet on the Winipeg, producing in the woods long flexible branches, armed with a few slender sharp thorns. Its ripe fruit is fleshy and well flavoured, but rather mealy, of a yellowish colour inclining to orange. It has an ovoid shape, with a shallow groove on one side like a peach, is nearly an inch in diameter, and its stone is so much compressed that its thickness is less than half its width; while its length, being 0.63 inch, exceeds the width by a fourth part. The nut is oblique, with convex valves, being circumscribed by two unequal curves. One edge is acute, with a groove on each side of it; the other edge is occupied by a narrow groove. I have been thus particular in the description of the northern fruit, that it may be compared with plums growing in other districts. The fruit is

289

cans or fragrant banks d *R. ve*ed north

A. caneon the uminous udacacia

Centucky Cassia limits in

seems to ot having ut. Abert idian and . It is a tes. The gation, as acters are of ten or g flexible ripe fruit yellowish e, with a h inch in thickness ·63 inch, ue, with curves. the other thus part may be e fruit is the Puckësāminan of the Crees; and La Prune, or the plum, of the white residents.

The Nekā-u-mina of the Chippeways, or Thekā-u-mina of the Crees, and sand cherry of the residents on Rainy River and Lake Winipeg, is a bush or small shrubby tree a foot and a half high, which grows on sand hills. The bark of its annotinous and biennial shoots is reddish, and the older twigs are brownish, with small warty speeks. When in fruit (in which state only I examined it in September, 1849) the fruitstalks are solitary, and spring from the base of the summer's growth; they are rather more than half an inch long, or about equal to the diameter of the fruit, which is black and rather austere, but edible. The stone is 0.38 inch long, almost regularly elliptical and acute at the ends, but more so at one end than the other. Its valves are very convex, so that its width exceeds its thickness very little. The sides are not acute-edged : one suture is depressed, forming a shallow groove; on the other side, which is very obtuse and almost flattened, there is a furrow above and below the suture, and rather remote from it. The annotinous shoots are smooth, angular, and generally flexuose, with the leaves springing alternately at the curves. The leaves measure two in thes and a half, the footstalk forming about one fifth part of this length. The lamina is lanceolato-elliptical; that is, nearly regularly elliptical, with an acute end, and a gradual tapering into the footstalk; it is serrated by acute, appressed teeth at the upper end, and is entire towards the footstalk; its under surface is pale and somewhat glaucous, the upper one dark green, and both sides are perfectly smooth. The footstalks are edged by the decurrent lamina for more than half their length, and the deciduous linear-lanceolate stipulæ are inciso-pinnate inferiorly. This cherry is probably the Cerasus pumila of Michaux and later American botanists. It was not traced by me beyond the 50th parallel.

Another small shrubby cherry grows, on moist sandy soil, by the banks of rivers and lakes, from Lake Superior to the Elk River on the 57th parallel. Its fruit is scarcely half the size of the preceding, but is, like it, black, and hangs generally on

VOL. II.

U

solitary footstalks, though the flowers grow by two or threes in short racemes. This shrub lies close to the ground; makes no approach to the tree form; seldom exceeds a foot in height; and so much resembles the *Salix myrsinites* and some other depressed willows, that, on looking for catkins, I have not been undeceived until I found the footstalks of the last year's cherries. The fruit of this sand cherry is sweeter than that of the preceding one. Whether it be the *Cerasus depressa* of some botanists I cannot determine; nor do I pretend to clear up the confusion that exists in botanical works respecting *C. pumila* and *depressa*.

C. pennsylvanica, wild red cherry, the Pasis-so-wey-minan of the Chippeways, and Pasi-a-wey-minan of the Crees, produces a small sour red fruit, which grows in a many-flowered raceme on long slender footstalks. Its equatorial limit, according to the United States' botanists, is the New England States and Pennsylvania, where it is a slender tree 20 or 25 feet high. Its polar limit is within the Saskatchewan basin, which it ascends towards the base of the Rocky Mountains, nearly to the height of 2,000 feet above the sea. C. virginiana, choke cherry, is named by the Crees Ta-hwoy-minan, and by the Dog-ribs Ki-e-dunnè-yerrè. It was found by Licut. Abert on the Kansas and Arkansas, and on Purgatory Creek; and is, in northern latitudes, a shrub with long branches. At Fort Liard, on the 61st parallel, it is 20 feet high, and on the confines of the arctic circle, where it terminates, it does not exceed four or five feet. The fruit can scarcely be said to be edible by itself; but it is often pounded, stones and all, and mixed with pemican. C. serotina, wild black cherry, is a general inhabitant of Rupert's Land, extending westward to the valleys of the Rocky Mountains on the Pacific side, where, however, it is generally dwarfed; and northwards to near Great Slave Lake. It is said by Dr. Gray to be a fine large tree in the Northern States, with purplish-black fruit, having a pleasant vinous flavour. Besides these I gathered specimens of a cherry-tree, not in flower, on Athabasca and Slave Rivers, which Sir William Hooker is inclined to consider as the C. mollis, discovered by the unfortunate

291

.

r threes makes height; he other not been cherries. the preof some or up the *t*, pumila

y-minan ees, pro--flowcred t, accordnd States 25 feet in, which is, nearly na, choke d by the Abert on and is, in ort Liard, onfines of ed four or by itself; pemican. bitant of he Rocky generally It is said ates, with Besides flower, on oker is infortunate

David Douglas, on the banks of the Columbia, growing on subalpine hills to the height of from 12 to 25 feet.

Purshia tridentata inhabits the Rocky Mountain prairies near the head waters of the Missouri and Columbia, and extends northwards to the 49th parallel. Spiraa opulifolia, nine bark, ranges from Maine, Canada, and Wisconsin, westwards to the valley of Oregon, in which it is found from the sources of the Columbia downwards. It is common on the low islands of Lake Superior, and has its polar limit in the colony of Red River. S. chamædrifolia inhabits the north-west coast up to Sledge Island in Beering's Straits, and Chamisso Island in Kotzebue Sound. It does not cross the Rocky Mountains, nor does any other Spiræa go so far north on the east side of the continent. S. betulifolia is another western species which inhabits the Blue Mountains and Mount Hood, and crosses the Rocky Mountains to their eastern valleys between latitudes 52° and 54°, but docs not descend to the lower eastern country. S. tomentosa, hardhack or steeple-bush, is common in the meadows and low grounds of New England, and spreads through Nova Scotia, Canada, and Rupert's Land to Lake Winipeg. S. douglasii is an Oregon species resembling the preceding, which extends to the Straits of Da Fuea. S. ariæfolia forms part of the underwood in forests on the Pacific coast, on the Kooskoosky, Spokan, Flathhead, Salmon, and M'Gillivray Rivers up to the 49th parallel. S. salicifolia is very abundant on the banks of every lake and river in the St. Lawrence and Saskatchewan basins, and northwards to Slave River. It is often associated with the Myrica gale, growing in the water. In its northern range it approaches the S. chamadrifolia of the west coast, but does not attain so high a latitude, owing to the greater severity of the climate on the east side of the mountains.

Rubus occidentalis, black raspberry or thimbleberry, extends from the Northern States to the Saskatchewan basin, and also to the Pacific coast. *R. strigosus*, wild red raspberry, is also found on both sides of the continent; on the east side it inhabits the United States, Newfoundland, and Canada, and may be traced in the interior canoe route throughout the Sas-

katchewan basin. R. nutkanus, white flowering raspberry, was discovered in Queen Charlotte's Sound by Mr. Menzies in lat. 51° on the Pacific coast, since which time it has been found in Norfolk Sound, lat. F , and traced down to Cape Orford in lat. 43°, and to the be waters of the Columbia in 52°. On the eastern declivity o. he Rocky Mountains it grows between latitudes 52° and 54°, and on the River Winipeg, Lake Superior, and Upper Michigan. Near the Pacific it is ten feet high, and forms the underwood on the island of Sitka; but in the passes of the Rocky Mountains dwindles down to a foot or eighteen inches. In thickets on the Winipeg its leaves attain remarkable dimensions : the fruit is inedible. R. odoratus, purple flowering raspberry, is a native of the Northern States, Canada, and the country between Lake Superior and the Saskatchewan. R. spectabilis is a prickly shrub, ten feet high, inhabiting the Pacific coast from Oregon to Unalashka. R. suberectus, bram-This species, which is also European, is an inhabitant of ble. Newfoundland, and of the country between Lake Superior and the Saskatchewan, where it was found both in 1825 and 1848. R. villosus is common in the Northern States, and is found also in Nova Scotia and Canada-West up to Lake Huron and Wisconsin. It is included in Elliott's "Flora Carolina," but I have not ascertained its equatorial limit. R. hispidus vel obovalis, running swamp blackberry, is common in the Northern States, and extends through Canada to Lake Superior. R. canadensis (L.), vel trivialis (Pursh), low blackberry or dewberry, has a similar range with the preceding species. R. nivalis (Douglas), is an alpine shrub, found on the snowy ridges of the Rocky Mountains, and not growing more than six inches high. There are also several herbaceous species of this genus; as R. triflorus, dwarf raspberry, which is common in the Northern States and throughout Rupert's Land, northward to Slave and Mackenzie Rivers. The Dog-ribs name it Tāsillè-ki-ch. Its northern limit is about lat. 68°. R. chamæmorus, cloudberry, is found on the White Mountains of New Hampshire near the limit of trees; also in Maine and Nova Scotia, Newfoundland and Labrador. In lat. 54°, and more to the north, it crosses the

continent, and is found on the summits of the Rocky Mountains between latitudes 52° and 56°; in Unalashka, on the shores of Beering's Straits, and on the most northern promontories of the continent. Near the Arctic Sea it is a common plant on mossy plains, but produces fruit there only in fine seasons. The fruit, which has a rich honey flavour, perishes with the early frosts. It is perhaps the most delicious of the arctic berries when in perfection, but cloys if eaten in quantity. R. stellatus, resembling the preceding, has been found only at Foggy Harbour on the north-west coast. I. arcticus and R. acaulis inhabit the shores of Hudson's Bay, Labrador, and the country westward to Kotzebue Sound. Their southern limit seems to be in the Saskatchewan basin, in about lat. 53°. On many parts of the flat beaches of Slave and Mackenzie Rivers the lively red flowers of R. acaulis cover large patches of ground which are partially flooded by small rivulets. In woods the last-mentioned species has a stouter growth, and emits long flagelli which run among the mosses.

Potentilla fruticosa, the shrubby einquefoil, grows abundantly from the northern states to the Arctic Sea, by river banks as well as in the most exposed and elevated situations. It occurs in the high valleys of the Rocky Mountains, at Pelly Banks on the west side of that range, and in Kotzebue Sound. On the Coppermine River near the sea it is almost herbaceous, the woody stem being extremely short and subterranean.

Rosa woodsii, R. carolina, R. blanda, R. cinnamomea, R. majalis, and R. stricta grow in the wooded districts; but, from their similarity to each other, their respective limits have not been ascertained. R. blanda was found flowering freely near the mouth of the Mackenzie on the 69th parallel. This species and R. cinnamomea cross to the Pacific coast. R. woodsii and majalis have been traced as far north as the Mackenzie. R. nitida and lucida grow in Newfoundland and in the New England States. R. fraxinifolia is confined to the Pacific coast; and R. lævigata has not been found beyond Lake Huron. R. seligera, a fine climbing rose, grows from Ohio to Wisconsin, but has not been detected to the north of the great lakes.

s.

spberry,

enzies in en found)rford in 52°. On between Superior, high, and he passes eighteen emarkable ole flowermada, and atchewan. biting the tus, bramabitant of perior and and 1848. and also in d Wiscon-I have not valis, run-States, and densis (L.), as a similar ouglas), is cky Moun-There are R. triflorus, States and Mackenzie s northern ry, is found the limit of ndiand and crosses the

υ 3

293

Cratagus punctata, dotted thorn, is found everywhere in the Northern States, extends northwards to Wisconsin, and crosses the continent to the coast of the Pacific; but has not, so far as I have learnt, been found within the British territory. C. glandulosa occurs in Canada, and northwards to the south side of the Saskatchewan basin; probably also on the Pacific coast. C. coccineu, scarlet-fruited thorn, is a common low tree in the Northern States; was found by Lieut. Abert, as far south as Stranger Creek, in lat. 39°; and extends to Wisconsin and the great lakes. C. cordata, Washington thorn, is supposed by Dr. Asa Gray to have been introduced into New England, but to grow wild in Pennsylvania and the more southern states. It is found from Canada to the Saskatchewan and the valleys of the Rocky Mountains, and about the sources of the Columbia, in between lat. 52° and 54°. The Cratægi flourish on the banks of Rainy and Winipeg Rivers; but are scarce further north. Mertens found several in the forest lands of Sitka. Amelanchier canadensis (botryapium et ovalis), shad-bush and serviceberry, is La Poire of the voyagers, the Misass-ku-tu-mina of the Crees, and the Tche-ki-eh of the Dog-ribs. This shrub extends along the banks of rivers nearly as far northwards as the woods go, and produces fruit up to the 65th parallel on the Mackenzic. It is common in the Northern States, in Nova Scotia, Newfoundland, and Labrador, and westward to the Pacific. The black fruit is about the size of a pea, is well tasted, drics well, and in that state is mixed with pemican, or used for making puddings: for which purpose it nearly equals the Zante currant. Its wood, being tough, is used by the natives for making arrows and pipe-stems, and has obtained on that account the name of bois de flèche from the voyagers; but in the United States the name of arrow wood is given to a different tree. The variety or species named A. sanguinea was traced up to the 60th parallel.

Pyrus rivularis, Powitch tree, inhabits Oregon and Vancouver's Island. Its fruit is edible, and its wood, which is hard enough to take a fine polish, is used for wedges. *P. ameri*cana, the American mountain ash, is found on the southern parts of the Alleghanics, and more commonly in the swamps

295

3.

e in the l crosses so far as ory. C. outh side fic const. e in the south as and the posed by land, but rn states. valleys of Columbia, the banks her north. elanchier serviceu-mina of shrub exirds as the the Macva Scotia, e Pacific. sted, dries used for the Zante atives for at account he United erent tree. l up to the

> and Vanch is hard *P. ameri*southern he swamps

and mountain woods of the Northern States. It is frequent on the shores of Lakes Huron and Superior; but is seldom seen on the cance route beyond Lake Winipeg. On the acclivities and in the valleys of the Rocky Mountains, however, it ranges northwards to Fort Liard, near the 60th parallel. It has been observed as high as the 56th degree of north latitude on the Pacific coast, from whence it extends southwards through the subalpine regions of Oregon. *P. arbutifolia*, choke-berry, is common in the damp thickets of the northern states, in Newfoundland, in Canada, and onwards to the Saskatchewan basin; but was not observed so far north as the immediate banks of that stream.

GROSSULARIEE. — The species of this family seem to attain their maximum number to the north of the United States. Ribes oxyacunthoides, sharp-thorned gooseberry, inhabits Newfoundland, Canada, and the canoe route northwards to the 62nd parallel, or perhaps further. R. cynosbati, prickly gooseberry, is common in the rocky woods of the Northern States, and accompanies the preceding species northwards to Slave Lake. R. saxosum inhabits New England, the shores of Lake Huron, and the valley of the Saskatchewan, extending also probably to Oregon. R. hirtcllum, short-stalked wild gooseberry, the most common species in New England, extends to Canada, Wisconsin, and northwards to Great Slave Lake. The preceding one seems to be considered by Dr. Asa Gray to be a variety of this species. R. lacustre, swamp gooseberry, is common in the most northern parts of the United States, and Nova Scotia; crosses the mountains to North California and Oregon; and extends northwards along the Mackenzie, nearly or quite to its delta. It is the Tagossay-ki-eh of the Hare Indians, which name is common to several kinds of gooseberry in the Dog-rib country, where there is a greater variety of species. R. divaricatum is common near Indian villages on the north-west coast, from 45° to 52° north lat. R. rotundifolium, Michx. (triflorum, Willd.), is a rare inhabitant of the mountainous districts of Oregon, and inhabits the Northern States from Massachusetts to Michigan and Wisconsin, but has not been found on the north side of Lakes Superior or Huron. R. rubrum, the common red

u 4

15

currant, native both of Europe and America, extends from the northern states very nearly to the shores of the Arctic Sea, having been gathered beyond the 69th parallel; and it ranges westward to Kotzebue's Sound. It is the Ki-eh-eth-lule-āzè of the Dog-ribs and Hare Indians. R. prostratum, fetid currant, inhabits cold damp woods from Nova Scotia and the Northern States northwards to the Athabasca, and westward to the Rocky Mountains and Oregon. The fruit is produced in copious racemes; but, in common with the foliage, it has an unpleasant odour, and a strong taste of turpentine. R. hudsonianum. This is the Nut-sinne of the Dog-ribs, and is a common gooseberry from Hudson's Bay to the Rocky Mountains and subalpine districts of Columbia; also in a northerly direction on the Mackenzie to lat. 67°. R. floridum, wild black currant, resembles the preceding, and is a common species in the Northern States, westward to Wisconsin, and ranges northwards to lat. 54°. R. sanguineum, which has become so common an ornament of our gardens, is a native of the Pacific coast only, where it ranges from 38° north lat. to 52°. There are several other very handsome species in Oregon, and, among others, the rich P. aureum; but they have not been traced beyond the 49th parallel.

ARALIACE. — Panax horridum, prickly ash-leaved panax, a twining shrub common in California, Oregon, and New Caledonia, as far north as 57° or 58°, crosses the Rocky Mountain ridge to the upper tributaries of the Saskatchewan, but does not descend to the eastward. Aralia hispida, bristly sarsaparilla, may be considered as the eastern representative of the preceding, though it is scarcely shrubby, having merely a very short, tough stem, almost buried in the crevices of the rocks from which it springs.

CORNEZ.—Cornus alba vel stolonifera, red osier cornel. This willow-like shrub, which is the osier rouge of the voyagers, ornaments the river strands from the Northern States, Nova Scotia, and Newfoundland, northwards to near the mouth of the Mackenzie, and westward to the shores of the Pacific. It is named by the Crees, on account of the bright red colour of its

297

twigs, Mithkwa-pè-min-àhtik (red stick), and its fruit Mushwamina (bear-berry), because the bears eat it. The Dog-ribs call this berry, Kai-gossai-ki-ch. A warm decoction of the bark and twigs is used by the natives for bathing their limbs when swelled by fatigue. C. alternifolia, C. paniculata, C. sericea, and C. circinata, which are inhabitants of the Northern States, are said to extend to Canada; but except the last named, which occurs on Lake Superior, none of them were gathered by us on the canoe route. C. sericea and C. florida, also Canadian species, cross the continent to Oregon, but do not occur north of the great lakes. The herbaceous C. canadensis reaches the shores of the Arctic Sea, crossing the continent from east to west; and the P. suecica, a European plant, is found in the Gulf of St. Lawrence, and on the west coasts of arctic America as high as Kotzebue Sound, and southwards to Oregon; but has not been detected in the interior districts.

LORANTHACE *E.* — Arccuthobium oxyccdri, this leafless parasitical shrub, is common to Europe, Central Asia, and North America, where it grows on cedars and pine trees. On the eastern declivities of the Rocky Mountains it ranges from lat. 52° to 57° north, and also eastward to Hudson's Bay, growing on the *Pinus* banksiana. On the western side of the mountains, from the Spokan River in 47° north lat. to near the sources of the Columbia, it infests the *Pinus ponderosa*.

CAPRIFOLIACE \mathcal{E} .— Sambucus canadensis, black-fruited elder, was gathered by Lieut. Abert on the Cottonwood Creek of the Neosha, in lat. $38\frac{1}{5}^{\circ}$ north, at an altitude of about 1,400 feet. It has its northern limit in the Saskatchewan basin, and ranges westward from Nova Scotia across the prairies. S. racemosa vel pubens, the red-fruited elder, is common in the Northern States on the shores of Lake Superior, going northward to the Saskatchewan, and westward to Oregon. Its polar limit, as far as ascertained, is on the eastern declivity of the Rocky Mountains between 52° and 59° north lat. V. prunifolium, black haw, or sloe-leaved viburnum, reaches the north shore of Lake Huron; but is more common in New York and Ohio. V. ientago, sweet viburnum, is a handsome tree in the Northern States, grows in

s.

rom the tic Sea, t ranges *le-āz*è of currant, Northern ie Rocky copious ipleasant mianum. n goosesubalpine 1 on the rant. re-Northern ls to lat. an ornaly, where ral other the rich the 49th

d panax, nd New e Rocky itchewan, a, bristly sentative g merely es of the

hel. This voyagers, es, Nova th of the ic. It is our of its

Nova Scotia and Wisconsin, and extends northwards to the south side of the Saskatchewan basin. V. nudum, withe-rod, is more common from New Jersey southwards than towards the great lakes; but occurs as far northwards as the last-named V. dentatum, arrow wood, is common in the low species. grounds of the Northern States up to Wisconsin, and is said by Pursh to extend to Canada; but it seems to be rare in that country. V. acerifolium, maple-leaved arrow wood, is a more northern species; and, in common with other shrubs that approach the arctic circle, it crosses the Rocky Mountains to the valley of Oregon, and also to Sitka. It has been traced as far north as Great Slave Lake, occurs also in Newfoundland, and is common in the rocky woods of the Northern States. V. opulus vel oxycoccus. Sir William Hooker is inclined to consider the European and American shrubs known by these names-and of which the handsome snow-ball tree, or guelder rose, is a cultivated variety-to be one species, and Dr Asa Gray unites them. In America the shrub extends from the Northern States and Nova Scotia to lat. 68° on the Mackenzie, and perhaps very nearly to the verge of the woods. It also crosses to the Pacific coast, having been found in the valley of Oregon. Its fruit, of a bright pinkish red colour, has a sharp acid taste, and is the Mongsö-a minā (moose-berry) of the Crees, and the Dunnèki-e or Indian-berry of the Dog-ribs and Hare Indians. The fruit being sometimes used as a poor substitute for cranberries, has obtained for the bush the name of cranberry tree in the Northern V. edule, the pembina of the voyagers, was traced by States. us northwards to the Elk River. It is much less common than the preceding, and has a more fleshy and less acid fruit, of an orange-red colour. The voyagers relish this fruit; and it has given name to many of the rivers of Rupert's Land. It is the Nipi-minan (water-berry) of the Crees. Michaux, Dr. Asa Gray, and other authors consider it to be scarcely a variety of V. opulus: I have found, however, its foliage retaining pretty constantly its peculiar character. In V. oxycoccus the lobes of the leaves are separated by acute sinuses, and have long, tapering, jagged, or deeply serrated points. In V. edule the sinuses

between the lobes are rounded, and the lobes themselves are shorter, though the lamina of the leaf is cut to within a short distance of its base. The European V. opulus has generally obtuse sinuses, and a less deeply cut lamina, but the lobes also short. Pursh, who separated V. oxycoccus from edule, described the bases of the leaves of the one as acute and of the other obtuse; but, as there seems to be no difference in that respect, it is probable that he meant the sinuses.

Diervilla trifida ve! canadensis, bush honeysuckle, has a herbaccous aspect, and is one of the most common underwoods on the portages. It occurs in all the woody districts of the Saskatchewan basin up to the acclivities of the Roc!- Mountains, but is rare to the north of Cumberland House. It grows also on Lake Superior, in Wisconsin, Nova Scotia, and the Northern States. Lonicera parviflora, small honeysuckle, has a conterminous range with the preceding. L. douglasii gathered on Saskatchewan is considered by Dr. A. Gray to be merely a variety produced by cultivation. L. hirsuta, hairy honeysuckle, is a coarse-leaved climber, common in moist rocky woods of the Northern States, and Canada as far as Lake Huron. L. ciliata, fly honeysuckle, grows at Pictou, on the Catskill Mountains, in Ohio, Wisconsin, generally throughout the rocky woods of the Northern States; also on Lake Superior, and northwards along the whole Saskatchewan basin. L. cærulea, the mountain fly honeysuckle, extends northwards to the arctic circle ; it likewise ranges from the Labrador coast and Newfoundland to the Rocky Mountains, and we should suppose also to the Pacific coast, since it is both a European and a Siberian species; but it is not named by Mertens or Bongard among the Sitka plants, nor does it appear to have been found by Douglas, Tolmie, or Scouler in Oregon. It grows in Wisconsin, New Hampshire, Massachusetts, and New York.

Symphoricarpus racemosus, snow-berry, and S. occidentalis, wolf-berry, range from Vermont, Michigan, and Wisconsin, over the St. Lawrence and Saskatchewan basins, to the 60th parallel on the Mackenzie. They also occur in the Oregon valley, Vancouver's Island, and doubtless much further along that coast.

ITS.

ds to the withe-rod, n towards ast-named the low is said by are in that is a more s that apains to the traced as Newfoundern Stales. nclined to a by these or guelder · Asa Gray • Northern nd perhaps ses to the egon. Its taste, and he Dunnè-The fruit erries, has Northern traced by nmon than ruit, of an and it has It is the Dr. Asa variety of ng pretty e lobcs of ng, taperie sinuses

299

A

RUBIACE *E*. — Cephalanthus occidentalis, button bush. This shrub, which belongs to the sub-family of Cinchone *æ*, occurs in thickets of the Northern States and Canada, but does not extend to Lake Superior.

COMPOSITE. — Of this large family no shrub has been detected in the canoe route north of Lake Superior; though the *Crinitaria viscidiflora* grows as high as the 55th parallel on the banks of the Salmon River, west of the Rocky Mountains, and on the upper branches of the Columbia above the Kettle Falls. A small annual herb was found on the Saskatchewan, which Sir William Hooker placed next this species; but, from the imperfect specimens, he could not ascertain its genus satisfactorily.

VACCINE. — Gaylussacia resinosa, black huckleberry, is common in the Northern States westward to Wisconsin, and extends northwards to the Saskatchewan. Vaccinium corymbosum, common swamp blueberry, extends from the Northern States to Newfoundland and Canada, as far north as Quebce, but has not been gathered to the westward of Lake Superior. V. pennsylvanicum, low shining-leaved blueberry, is very common in the dry rocky woods of the Northern States, Canada, and the country between Lakes Superior and Winipeg. V. canadense, downy-leaved blueberry, is the most abundant species by the sides of streams and in thickets, from Maine and Michigan to the shores of Hudson's Bay, and northwards in the woody districts to the arctic circle. It extends also westward across the mountains to the upper feeders of the Columbia. V. uliginosum, bog bilberry, occurs on the summits of the New Hampshire Alps; on the Green Mountains of Vermont, and on Essex county mountains of New York; on the Newfoundland, Labrador, and Greenland coasts ; also from Lake Superior northwards to the Arctic Sea. On the west side of the Rocky Mountains it has been gathered on Sitka, Unalashka, and Kotzebue Sound. In Europe it grows in the forests of the higher Jura, in England, and the Scandinavian peninsula. Beyond the arctic circle its fruit is not abundant every year; but in good seasons it is plentical to an extraordinary degree, and is of
a finer quality than in more southern localities. It then affords food to the bears and large flocks of geese, which fatten on it, and acquire a fine flavour. The berries, when frozen by the autumnal frosts, remain hanging on the bushes until the snow melts in the following June, and may then be gathered in a very juicy but tender condition. V. salicinum, willow-leaved bilberry, is an inhabitant of Unalashka. V. myrtillus, myrtleleaved bilberry, was gathered by Mr. Drummond on the summit of the pass between the head waters of the Saskatchewan and Columbia, but has not been detected further to the east, though it is a European plant. Chamisso found a Vaccinium on Unalashka, which he was inclined to refer to this species; but his specimens were imperfect. Bongard, however, enumerates it as existing among the plants gathered by Mertens on Sitka. V. myrtilloides vel angustifolium is found in Canada, and from Hudson's Bay to the woody declivities of the Rocky Mountains, between the 52nd and 54th parallels. It crosses the dividing ridge also to the alpine valleys of Oregon, and to the sca-coast further north, where the purplish-brown fruit is eaten with relish by the natives. V. cæspitosum, dwarf bilberry, grows on the alps of New Hampshire, the shores of Lake Superior near James' Bay, and northward to the valley of the Saskatchewan and the Rocky Mountains, between 52° and 57° north; also in the Oregon valley. V. ovalifolium grows in Oregon from the mouth of the Columbia up to the Portage River, near the crest of the Rocky Mountains, on the 50th parallel, and also on the island of Sitka at lat. 57° north. V. vitis-idæa, eow-berry, or alpine cranberry, is the Wi-sā-gù-minā of the Crees, and the cranberry most plentiful and most used throughout Rupert's Land. This berry is excellent for every purpose to which a cranberry can be applied; and though inferior to the V. oxycoccus in flavour in autumn, is far superior to it after the frosts; and, as it may be gathered in abundance in a most juicy condition when the snow melts in June, it is then a great resource to the Dog-ribs and Hare Indians, as well as to the immense flocks of waterfowl that are migrating to their breeding places at that date. It

TS.

sh. This eæ, occurs does not

has been r; though th parallel he Rocky abia above n the Sasnis species; certain its

leberry, is consin, and um corym-Northern as Quebec, Buperior. y, is very ern States, d Winipeg. t abundant Maine and ards in the westward Columbia. f the New mont, and Newfounde Superior the Rocky , and Kotthe higher Beyond r; but in , and is of

grows in perfection, in the most exposed situations, round a boulder or granite rock, over whose face its branches may spread, and where it can have at one time both moisture and the reflected heat of the sun's rays. It is found at Danvers, in Massachusetts, in Maine, and the higher mountains of New England, where its fruit is reported by Dr. Asa Gray to be barely edible, bitter, and mealy. In the parallel of Lake Superior it spreads from the Atlantic to the Pacific (being absent, however, on the prairies). In a higher latitude it crosses the continent also from Churchill Fort to Sitka and Kotzebue Sound, and it extends in the middle districts to the Arctic Sea in latitude 71°. In Sitka its leaves are said to be small. In Rupert's Land they vary in size, according as the plant is exposed or under shade. V. ovatum is common in Oregon and rocky places of the west coast northwards to the 49th parallel. V. oxycoccus, dotted cranberry, is, like the preceding, common to the New and Old World. It grows in peat bogs from New England and Wisconsin, northwards to the aretic circle, and from Newfoundland and Labrador to the Rocky Mountains, between 52° and 57° north lat.; in Sitka on the latter parallel, and in Kotzebue Sound. V. macrocarpum, American cranberry, is common in the peat bogs of the Northern States, and has its limit in the Saskatchewan basin. It crosses the continent from Newfoundland to Oregon; and the natives near the mouth of the Columbia eat its fruit, when boiled, under the name of Su-labich. Chiogenes hispidula, creeping snowberry, is common in the Northern States, where it grows under evergreens in turfy places. It extends across the continent from Newfoundland to the sources of the Columbia, and northwards along the Rocky Mountains to the 55th parallel.

ERICE E. — Gaultheria procumbens, creeping winter-green tea-berry, chequer-berry, partridge-berry, or box-berry. This fragrant creeping shrub is a great ornament of the woods north of Lake Superior. It inhabits moist woods in the Northern States, grows at Pictou and on Lakes Huron and Superior, and was traced by us northwards to the Lake of the Woods, or near

the 50th parallel. G.? myrsinites has hitherto been found on the declivities of the Rocky Mountains only between the 52nd and 57th parallels. Mr. Drummond says, that its small berries have a delicious pine-apple flavour. The plant was cultivated in the Botanic Garden at Glesgow, but I have not heard that it produced fruit there. G. shallon is an Oregon plant growing between Cape Mendocino and Puget Sound, but not extending inland more than a hundred miles from the sea-coast. Epigæa repens, ground laurel, or trailing Arbutus, inhabits sandy and rocky woods in the northern states, Canada, Nova Scotia, Newfoundland, and Rupert's Land, as far north as the Saskatchewan. Arbutus menziesii and A. tomentosa inhabit Oregon northwards to Puget Sound; but no true Arbutus has been detected on the east side of the Rocky Mountains. Arctostuphylos uva-ursi, bear-berry, is common to Europe and America, and descends from the Arctic Sea-coast to Rainy Lake and the rocks and hills of the Northern Sates. It crosses the continent to the valley of Oregon, where the Chenook Indians mix its dried leaves with tobacco. It is used for the same purpose by the Crees, who call it Tchakashè-pukk ; by the Chepewyans, who name it Kleh; and by the Eskimos north of Churchill, by whom it is termed Attung-ā-wi-at. On account of the Hudson's Bay officers carrying it in bags for a like use, the voyagers gave it the appellation of Sac-a-commis. On the north-west coast, Mertens found it at Sitka, and it doubtless extends along the whole coast. Its dry farinaceous berry is utterly inedible. A. alpina, alpine bearberry, though a herbaceous plant, may be mentioned with the others: it is also European. In the United States the only habitat given is the Alps of New Hampshire; but it grows at a much lower altitude in Newfoundland and Canada. It was found by Drummond on the Rocky Mountain ridge, and is very common on the barren grounds beyond the woody district, and along the whole arctic coast to Kotzebue Sound. There are two varieties, one with bright red and more juicy fruit; the other, having a dark purplish-black berry, of more fleshy consistence, and a stronger peculiar flavour. Both are eaten in the autumn; and, though not equal to some of the

rs.

round a

ches may sture and nvers, in of New ray to be of Lake fic (being atitude it Sitka and icts to the said to be ng as the ommon in rds to the e the prews in peat the arctic he Rocky ka on the rocarpum, gs of the van basin. n; and the ruit, when hispidula, s, where it hcross the Columbia, the 55th

nter-green ry. This ods north Northern erior, and s, or near

other native fruits, are not unpleasant. The two kinds are exactly alike in foliage. Andromeda hypnoides, moss-like andromeda, an inhabitant of the Alps of New Hampshire, Mount Marcy in New York, Labrador, and the north-west coast, was not detected by us on the interior canoe route. A. lycopodioides, a Kamtschatka plant was found by Chamisso on Unalashka. A. cupressina inhabits the Rocky Mountains in lat. 56° north. A. mertensiana and A. stelleriana, so named by Bongard, were discovered on Sitka by Mertens. A. tetragona is one of the most northern plants, being an inhabitant of the north end of Spitzbergen. It occurs on all the islands and coasts of the Arctic Sea, from Greenland to Kotzebue Sound, at Sitka, and as far south as Mount Hood on the 45th parallel. It is also a Lapland and Siberian plant. Like the two preceding species, it is rather a wiry herb than a shrub. The withered leaves of past years remain attached to the thread-like stem, and may be used as fuel, a fact which Mr. Rae so fully demonstrated, as we have mentioned, in a preceding page. A. polifolia, rosemary andromeda, inhabits the Alps of New Hampshire and New York, Wisconsin, Lake Superior, and the country northwards to the Arctic Sea; also the whole breadth of the continent from Newfoundland and Labrador to Sitka and Kotzebue Sound, with the exception of the prairies. It is an inhabitant also of the higher Jura. A. calyculata, rustyleaved andromeda, grows in sphagnous bogs and on the flooded strands of clear streams in the Northern States, and Rupert's Land as far as the upper part of the Mackenzie, and also on the shores of Beering's Sea. A. racemosa, cluster-bearing andromeda, grows in the moist copses of Canada, Massachusetts, and New Jersey near the coast, extending from thence southwards. A. ligustrina, privet andromeda, a common shrub of the Northern States, extends northwards to the Saskatchewan basin. In this genus and in Arbutus it may be noticed, that the more herbaceous species have generally the highest range.

Phyllodoce taxifolia, or Menziesia cærulea. This English plant grows on the New Hampshire Alps, and has been found on the Labrador coast. Steller is also said to have gathered it on the

American coast and islands opposite Kamtschatka. M. ferruginea and M. aleutica were found by Mertens at Sitka; the former, which is one of Menzics's discoveries, has since been gathered by Seeman on the coast of Beering's Sea, and the latter was previously found by Chamisso on Unalashka. Menziesia glanduliflora is one of Mr. Drummond's discoveries on the Smoking River, an elevated tributary of Peace River, on the 55th parallel. It is remarkable for its gracefully drooping yellow flowers. M. empetriformis inhabits Vancouver's Island and the alpine districts of Oregon. M. grahamii and M. interinedia grow on the Rocky Mountains, in lat. 55° eastward of their crest. M. globularis inhabits the same districts on the Smoking River northwards to 56°; and, according to Pursh, it occurs also on the high mountains of Carolina, and on the Cacapon Mountains, near Winchester, in Virginia. None of the Menziesiæ are mentioned by Dr. Asa Gray as existing in the Northern States; and it would appear that many of the species are very local, particularly the alpine ones.

Kalmia latifolia, calico bush, mountain laurel, or spoon wood, forms dense thickets on the mountains of Carolina and Pennsylvania, and is common northwards from Maine to Ohio and Canada, where it is a much humbler shrub. It was not observed by us on the north side of the St. Lawrence basin. K. glauca inhabits moorish places from the Northern States to the Arctic Sea, and crosses the continent to Sitka. K. angustifolia, sheep laurel, is common in the Northern States and Canada, to James's Bay and Newfoundland. We did not observe it on the canoe route north of Lake Superior. Azalea viscosa inhabits the Northern and Eastern States and Canada, but was not seen by us beyond the St. Lawrence basin. A. nudiflora, purple azalea, or pinxter flower, a common shrub in the Northern States, extends to Canada. The Rhododendron maximum, which is common on the mountains of Carolina and Pennsylvania, and is more rare in the Northern States and Canada, grows also in Oregon, on the subalpine range of Mount Hood, and more to the north on the high mountains near the "Rapids" of the Columbia. R. lapponicum, Lapland rose-bay,

VOL. II.

х

TS.

kinds are ss-like anire, Mount rest coast, oute. A. **Chamisso** Mountains , so named A. tetrahabitant of islands and bue Sound, 5th parallel. he two prehrub. The thread-like Rae so fully eding page. lps of New ior, and the ole breadth or to Sitka iries. It is lata, rustythe flooded nd Rupert's also on the ring androssachusetts, ence southon shrub of skatchewan ed, that the range. nglish plant ound on the

ed it on the

is another arctic plant which is found isolated on the peaks of the White Mountains of New Hampshire, and on Mount Marcy in the north corner of New York. It has been gathered as far south on the coast as the Labrador peninsula and the shore between York Factory and Churchill River, and grows on the summits of the Rocky Mountains on the 56th parallel, and throughout the whole extent of the barron grounds from Repulse Bay to Norton Sound, and northwards to the Arctic Sea. An infusion of the leaves and flowering tops was drunk by us instead of tea, but it makes a less grateful beverage than the Ledum palustre. It is a Scandinavian plant. R. hamtschaticum is an inhabitant of the north-west coast in lat. 53°, and of Unalaschka, as well as of the Asiatic shore. R. albiflorum, an elegant and ornamental plant, was discovered by Drummond on the Rocky Mountains between 52° and 57° of north latitude, where alone it has been found.

Loiseleuria vel azalea procumbens inhabits the alps of New Hampshire, and the coasts of Newfoundland, Labrador, Hudson's Bay, and the Arctic Sea; also the north-west coast at Mount Edgecumbe, Sitka, and Kotzebue Sound. Ledum palustre, narrowleaved Labrador tea, the Kā-ki-ki-pukwā (perennial leaves), or the maskego-pukwā (medicine leaves), of the Crees, is an inhabitant of the colder parts of Canada, the coasts of Newfoundland and Labrador, and the whole of Rupert's Land to the Arctic Sea, on whose shores it grows from Repulse Bay and the mouth of the Thlewee-choh to Kotzebue Sound. It is also found at Sitka; but Dr. Asa Gray has seen no specimens gathered south of the United States' boundary line. It is frequently used as a substitute for tea. L. latifolium grows in the woody districts of Rupert's Land, often in the immediate vicinity of the other species; but extends further south, being common in cold boggy grounds in the Northern States.

MONOTROPEE. — Cladothamnus pyrolifolius (Tolmiea, Hook.) inhabits Norfolk Sound on the Pacific coast in lat. 57°, and the country southwards to Puget Sound. Chimaphila umbellata, Prince's pinc, Pipsissewa, goes northward to 53° on the Rocky Mountains, but does not pass the 50th parallel in the much

NTS.

he peaks of at Marcy in as far south re between summits of oughout the lse Bay to , An infuy us instead the Ledum aticum is an and of Unabiflorum, an rummond on orth latitude,

alps of New dor, Hudson's Mount Edgeestre, narrowal leaves), or ees, is an inof Newfound-Land to the ulse Bay and id. It is also nens gathered is frequently in the woody te vicinity of g common in

> *lmiea*, Hook.) 57°, and the *la umbellata*, on the Rocky in the much

lower country through which the cance route lies. It crosses the dividing range, descends to the mouth of the Columbia, and is common in the Northern States. The Chippeways, in whose country it grows abundantly, do not appear to have discovered its admirable diurctic qualities. *C. maculata* is a more southern species: it was gathered by us on the great lakes, but is not common north of the Middle States.

JASMINE. — Fraxinus sembucifolia, black ash, is said to grow in Virginia, and by Dr. Asa Gray to range from Maine to Wisconsin. It also inhabits New Brunswick, Nova Scotia, and Canada. F. americana, white ash, was found by Lieut. Abert on the Arkansas, high up on the western slope of the Mississippi valley. It is a large forest tree in the Northern States. It grows at Pictou, also on Lake Superior, Rainy Lake, the River Winipeg, and the banks of the Saskatchewan, to latitude 54° north, where it is still a tree. F. pubescens, red ash, does not grow thicker than a man's thigh on Rainy River, where it terminates near the 49th parallel. It extends southwards to the Middle States, and also across the mountains to Oregon.

ELEAGNACE .- Eleagnus argentea, silver berry, is a very common shrub on the banks of rivers throughout the basins of the Saskatchewan and Mackenzie, up to the 68th parallel of latitude. Its dry husky berries are covered with the same silvery epidermis that gives the hoary appearance to the leaves, and are used by the Kutchin to ornament their dresses. This apparently sapless fruit is often found in the stomachs of geese on their northerly migrations. It is the wapow-muskwaminan, or "white-bear berry," of the Crees; and the branches, which harden in drying, are used by the natives for making pipe-stems. It ascends the Saskatchewan, and occurs in Canada, but does not find a place in Gray's "Flora of the Northern States." Shepherdia canadensis grows from Vermont and Wisconsin northwards to beyond the arctic circle, and is very common on the Mackenzie. Its small, red, juicy, very bitter, and slightly acid, berry is useful for making an extempore beer, which ferments in twenty-four hours, and is

an agreeable beverage in hot weather. S. argentea is a prairie shrub common to the plains of the Missouri and Saskatchewan, but which does not grow in the eastern districts. It is the Mith-y \ddot{u} min \ddot{a} or blood-red berry of the Crees.

THYMELER. — Dirca palustris, leather-wood, is common in the Northern States, and extends to the north side of Lake Superior, disappearing about the Lake of the Woods.

EMPETREE. — Empetrum nigrum occurs on the alps of New Hampshire and New York, and is found throughout the whole extent of Rupert's Land up to the Aretic Sea, along which it ranges to Kotzebue Sound, descending the western coast to Sitka, and perhaps lower. It is absent only on the prairies. In the more sandy tracts of the barren grounds it covers the surface with its prostrate branches, that are loaded with fruit in favourable scasons. The snow-geese feed and fatten on the berries, which, after the fresh frosts, become very juicy, and are highly refreshing to the weary and thirsty traveller.

ULMACER. - Ulmus americana, white elm, was found by Lieut. Abert on the Pawnee Fork in latitude 38° 10' N., at an elevation of 1.658 feet; and Dr. Asa Gray informs me that it descends to Texas. It is a majestic tree in the Northern States, much prized for its rapid growth and the beauty of its form. Its wood is in requisition there for the use of wheelwrights. On the north banks of the Saskatchewan, in about latitude 54°, which is its polar limit, it grows only in rich alluvial soil, and, being crowded among balsam poplars and other trees which inhabit such places, does not exhibit its handsome outline so as to strike the eye. Its timber there is often decaved at heart, and, even when sound, is so porous that we found it to be unfit for planking boats. It is probable that the U. fulva, slippery or red elm, known by the corky and angled bark of its branches, has an equal northerly range; but we did not trace it, though two kinds extend to the Saskatchewan, and we gathered It is perhaps an elm of which I heard, but did their flowers. not see in leaf, which inhabits wet places on the banks of Rainy River, and produces a wood that is considered there to

809

TS.

s a prairie atchewan, It is the

common in e of Lake

e alps of ughout the Sea, along he western nly on the grounds it are loaded e feed and ecome very and thirsty

s found by 0' N., at an me that it hern States, of its form. heelwrights. out latitude ich alluvial other trees some outline decayed at bund it to be e U. fulva, l bark of its not trace it, we gathered ard, but did he banks of ed there to be of no value. Dr. Asa Gray has not traced U. fulva on the Atlantic side of the Alleghanies further south than Maryland. Celtis occidentalis, sugarberry or hackberry, is common in the Northern States, and extends to Wisconsin and to the Oregon, but was not seen by us on the north side of Lake Superior.

JUGLANDINEE.—Juglans cinerca, butter-nut; J. nigra, black walnut; Carya alba, shell bark or shag bark hickory; C. amara, bitter nut or swamp hickory; and C. glabra, pig nut or broom hickory, reach Wisconsin and the basin of the St. Lawrence, but were not seen by us north of Lake Superior.

CUPULIFERÆ. - Quercus obtusiloba vel stellata, post oak, abounds in Texas as far as San Antonio de Bexar, wherever any hard wood grows. It was traced with Q. rubra, red oak, as far as the River Winipeg. Of the latter, Dr. Asa Gray says that he does not know whether it extends to Texas or not. It is a good-sized tree in the Northern States, and common in rocky woods. Q. alba, white oak. This, a most valuable forest tree in the Northern States, ranges northwards to Lake Winipeg, where it has a crooked and rather unsightly growth of 20 feet. Michaux states that its southern limit is in Florida; but I do not find it in the lists of plants gathered by Messrs. Emory and Abert in their journeys from Fort Leavenworth on the Missouri to North Mexico and California. Q. bannisteri; Q. tinctoria, quercitron or black oak; Q. macrocarpa, burr oak; Q. bicolor, swamp oak; Q. prinos, swamp chesnut oak; and Q. palustris, pin oak, grow in Canada or Wisconsin, but were not detected by us to the north of the great lakes, with the exception perhaps of Q. macrocarpa, since I gathered the immature acorns of an oak resembling this on Rainy River. Q. garryana inhabits Oregon northwards to Puget Sound, is a tree which reaches 80 feet in height, and is well adapted for shipbuilding.

Fagus ferruginea et sylvestris, American becch: is said by Pursh to range southwards to Florida; it ceases at Mackinac on Lake Huron, and does not grow on Lake Superior, but reappears further to the north-west on the Red River of Lake Winipeg, beyond which it was not seen. Carpinus americana, hornbeam, blue or water beach, called also iron wood, inhabits

Canada, Wisconsin, and the other Northern States, but was not seen by us on the canoe route. Ostrya virginica, hop hornbeam. This tree, which has also the trivial name of iron wood, grows as far north as the River Winipeg, and is plentiful on Rainy and Red Rivers. Its southern range, according to authors, is to Carolina and Georgia.

Corylus americana, hazel nut, and C. rostrata, beaked hazel nut, range northwards to the Saskatchewan, the former also crossing the continent to the Pacific coast.

MYRICACE.E. — Myrica gale, sweet gale or Dutch myrtle, is also a European shrub. It is common in North America, on the stony margins of lakes, and in peat bogs, from Virginia to the arctic circle, and in the island of Sitka. The native population of Rupert's Land use the buds as a material for dycing: *Comptonia asplenifolia*, sweet fern, is common in the Northern States, and terminates on the northern slope of the Saskatchewan basin; it ranges southwards along the mountains of Carolina and Georgia.

BETULACE .-- Betula papyracea, paper or canoe birch, is an invaluable tree to the population of Rupert's Land. Its bark is indispensable for the construction of their canoes, and also serves for the covering of tents in localities where the skins of large animals are scarce. Neatly sewed and ornamented with porcupine quills, it is moulded into baskets, bags, dishes, plates, and drinking vessels; in short it is the material of which most of the light and easily transported household furniture of the Crces is formed. The ruder Tinnè use it, but dispense with many of the forms into which it is worked by their southern neighbours. The wood serves for paddles, the framework of snow shoes, sledges, hatchet helves, and occasionally for gun-stocks; and in spring the sap forms a pleasant sweet drink, from which a syrup may be manufactured by boiling. Beyond the arctic circle it is a scarce and crooked tree, but occurs of a small size as high as the 69th parallel. It grows in perfection on the north shore of Lake Superior, in the neighbourhood of Fort William, where, owing to the ample supply of good bark, a manufactory of canoes or the use of the Hudson's Bay Company has been established.

As the Kolushes north of Sitka use birch-bark canoes, I infer that this tree extends to the Pacific; but I have not seen it in the lists of plants of that coast. Pursh mentions Hudson River as its southern limit; and Gray states its range as extending from New England to Wisconsin, but chiefly through the northern parts of that district. It grows in Nova Scotia, Newfoundland, and Labrador. B. occidentalis occurs in Oregon, along the Rocky Mountains, northwards to the straits of Da Fuca, and crosses the ridge to the vicinity of Edmonton House, on the 54th parellel. **B.** excelsa, yellow birch, was not traced by us beyond the banks of the Kamenistikwoya, which falls into Lake Superior. In the Northern States it is a stately tree, 60 feet in height. lenta, cherry or sweet birch, is a rather large tree, which is common in the Northern States, Nova Scotia, Canada, and Newfoundland; but does not appear to go far westward, as it was not found by us nor by Agassiz on Lake Superior. B. pumila vel glandulosa, little birch, is rare in New England, but grows in bogs of the northern parts of Pennsylvania, Ohio, Michigan, and Wisconsin; also in Nova Scotia, Canada, Newfoundland, and Labrador. It goes considerably beyond the arctic circle, being found on the banks of the Thlewee-choh, the Coppermine river, and other arctic streams, and also on the Mackenzie to about the 68th parallel. It is very like the following, but has a more crect and slender growth, which may be perhaps owing to locality. The leaves are generally longer. B. nana, dwarf birch, exists on the summits of the White Mountains of New Hampshire, and of the Essex Mountains of New York. It grows also in the higher parts of Labrador and Canada, along the shores of the Arctic Sea, from Davies' Straits to Kotzebue Sound; and generally throughout the Barren Grounds. Alnus viridis, green or mountain alder, and A. incana, speckled or hoary alder, range northwards to the delta of the Mackenzie on the 68th parallel, and from Newfoundland and Labrador to Kotzebuc Sound; the first species being also found on Sitka. They are common bushes in New England and Wisconsin; but I have not seen their southern limits mentioned. A. rubra, rcd alder, is an inhabitant of

TS.

t was not ornbeam. od, grows on Rainy uthors, is

aked haze] ormer also

myrtle, is merica, on Virginia to ative popufor dycing. e Northern skatchewan of Carolina

birch, is an l. Its bark es, and also the skins of nented with shes, plates, hich most of of the Crees ith many of neighbours. snow shoes, cks; and in lich a syrup c circle it is e as high as rth shore of iam, where, ry of canoes established.

x 4

311

В.

Sitka and Norfolk Sound, and extends southwards to Oregon. It seems to be the western representative of the *A. serrulata* of the southern parts of New England.

SALICACE. - Willows are numerous on the east side of the Rocky Mountains, and seem to attain the maximum developement of species in the southern parts of Rupert's Land, but to be less abundant on the Pacific coast, except to the north of the peninsula of Alaska. From Lake Superior to the Arctic Sea they form dense thickets on the shores of every river and lake. It is scarcely possible to note the range of willows, or to collect satisfactory specimens of the species on a rapid journey, as many of them which perfect their catkins before the evolution of their leaves, remain undetermined in the herbarium. Of twenty-two species described by Dr. Asa Gray as inhabitants of the Northern States, only S. tristis, Aiton; S. humilis, Marshall; S. sericea, Marshall; S. alba, L.; and S. angustata, Pursh, were not collected by us on our northern voyages. .S. uva ursi, S. repens vel fusca, and S. herbacea, which grow on the alps of New Hampshire, extend beyond the arctic circle on both sides of the continent, the latter being one of the most northern plants, as it grows on the north end of Spitzbergen. The other fourteen named by Dr. Gray reach one or more of the northern basins.

The following were traced from Lake Superior to the arctic circle, or beyond it: S. villosa, S. rostrata, S. discolor, S. viminalis, S. lucida, S. longifolia, S. corduta, S. rigida, S. planifolia, and S. pedicellaris. Some others were not gathered higher than the valley of the Saskatchewan, but their southward range included the St. Lawrence basin: as S. candida; S. petiolaris, S. rosmarinifolia, S. purpurea, and S. fragilis.

S. drummondii, S. barattiana, and S. cordifolia were gathered near the elevated sources of the Saskatchewan only, though the last named has been detected by other collectors on the Labrador coast. Salix sitchensis is known only as an inhabitant of the island from whence it derives its name. Salix richardsonii and S. acutifolia are common to the Saskatchewan and Mackenzie River basins, the former being also an inhabitant of the coast between York Factory and Churchill.

TS.

o Oregon. rrulata of

ide of the developeand, but to orth of the Arctic Sea and lake. r to collect ourney, as the evoherbarium. inhabitants S. humilis, angustata, yages. .S. ch grow on ic circle on of the most pitzbergen. or more of

o the arctic discolor, S. rigida, S. ot gathered heir south-S. candida; fragilis. re gathered though the e Labrador tant of the ichardsonii h and Mactant of the

The following are specially arctic in their habitats: S. myrsinites, S. vestita, S. speciosa, S. reticulata, and S. nivalis, which grow on the peaks of the Rocky Mountains between 52° and 57° north, and within the arctic circle, some of them reaching very high latitudes. S. reticulata grows on the coast between York Factory and Churchill. S. speciosa inhabits the Arctic Sea coasts from Coronation Gulf to Kotzebue Sound, and ranges southwards on the Mackenzie to about the 60th or 61st parallel. It is perhaps the nandsomest of the genus, having an agreeable growth, and very large leaves, which are of a silvery whiteness beneath, and when bruised have a rather pleasant odour. On the Mackenzie it grows to the height of 15 feet, in form of a bush, with very stout and long yearly shoots, which distinguish it from all the other willows of the same localities On the coasts of the Arctic Sea, wherever the rivers afford a suitable point of alluvial soil, a thicket of this willow may be expected as tall as a man. Mr. Seeman observed it in the tree form on the north-west coast, where it is from 18 to 20 feet in height, and having a stem five inches in diameter. It resembles S. lapponum in its habit. S. stuartiana and S. retusa grow on the more northern banks of the Mackenzie and in Kotzebue Sound, and have not as yet been detected south of the arctic circle. S. rostrata, S. speciosa, S. lucida, S. longifolia, S. depressa, S. reticulata, S. arctica, and S. polaris, have been enumerated by authors as crossing to the Pacific side of the Rocky Mountains in their respective zones. Salix glauca was found by Seeman on the shores of Beering's Sea.

The following are not confined to the American continent, but range to either Europe or Asia, as well as to Rupert's Land, or the arctic coasts: S. petiolaris, S. rosmarinifolia, S. viminalis, S. purpurea, S. fragilis, S. acutifolia, S. fusca, S. myrsinites, S. stuartiana, S. reticulata, S. herbacea, S. polaris and S. ammaniana.

The most common in Rupert's Land are the *S. rostrata*, which extends southwards to New England, and in the north forms almost impenetrable thickets 20 feet or more in height, in which the old twisted and sordid grey stems spread in all directions. *S. longifolia*, which has the growth of an osier, covers the new-

formed sandbanks of the rivers up to the 68th parallel, its flexible, densely growing young stems serving to arrest the mud, and speedily to raise the bank above the ordinary level of the water. In drier spots, by river banks in the Saskatchewan basin, it forms bushes from 20 to 25 feet in height. Even in the passes of the Rocky Mountains, as at Jasper Lake, it grows freely on drifting sands. Lieut. Abert found it growing at Council Grove, and Hundred-and-ten-mile Creek, at the height of 1,200 feet above the sea, between the 38th and 39th parallels. It inhabits also the banks of the Susquehannah, and all the Northern States. The soft pliable twigs are a favourite food of the moose-deer, and might, indeed, as they grow on the flooded sandbanks, be mowed like hay. Populus balsamifera, balsam poplar, or tacamahac, was found growing on the banks of the Mackenzie up to lat. 59°, where it makes a very slender tree. In the southern part of the delta of that river it forms groups of healthy young trees, and from thence to the United States it flourishes on rich alluvial and occasionally flooded banks of rivers to the exclusion, on such spots, of most other trees : its trunk attains a greater circumference than any other member of the northern forest, but its wood is of no value, except for fuel; and, when old, the tree is unsightly from having very generally lost its top. Its growth is rapid, and its decay apparently equally so. I measured some drift logs of this tree which were floating down the Mackenzie, and found them to be about 15 feet in circumference, with a very moderate tapering upwards. The Crees name it Mathèh-mètus, or ugly poplar. Dr. Asa Gray gives as its southern limit New England, Wisconsin, and perhaps Pennsylvania, but not further. It crosses Beering's Sea to Kamtschatka; and on the rivers of Oregon it grows, according to Douglas, to the height of 140 feet, and 20 feet in diameter. P. candicans, balm of Gilead, which greatly resembles the preceding, has not been detected north of Wisconsin: it is the common balsam poplar of Pennsylvania, New York, and New England. P. monilifera, lævigata vel cunadensis (Mx.), is a more southern species, being rare in New England, but taking the place of *P. candicans* in Western Pennsylvania,

Ohio, and Kentucky. It grows on the banks of the Arkansas and other southern tributaries of the Missouri. P. tremuloides, aspen. Dr. Gray believes that, south of Pennsylvania and Kentucky, this tree is confined to the Alleghanies, and even on these mountains it is rare. In the Northern States it is common, and varies in height from 20 to 50 fect. It abounds in Rupert's Land in the more fertile soils, and very generally springs up in place of the white spruce, when that tree has been destroyed by fire. Its range is co-extensive with the forest land; but towards the Arctic Sea, and in lat. 69° on the Mackenzie, it is a slender willow-like tree. It is the best firewood in the country, but is applied to no other economical purpose, except that its ashes are collected on account of the abundance of potash they contain. It do not know whether it inhabits the Pacific coasts or not. P. grandidenta, big-toothed aspen, is common in the Northern States, and reaches New Brunswick and Canada, but did not come under our notice on the canoe route.

PLATANACE*E.* — *Platanus occidentalis*, American Plane-tree, resembles the well-known *P. orientalis* in the way that its exterior bark falls off in thin plates. It extends northwards to Canada, but does not appear north of Lake Superior.

CONIFERÆ. — Pinus banksiana, gray pine, the Cyprès of the voyagers, grows from the arctic circle on the Mackenzie, down to the great Canada lakes, south of which, Dr. Gray has scarcely seen it, but has heard that it is found in the northern districts of Maine; and it occurs in the list of Wisconsin plants published by the American Association. It crosses the Rocky Mountains to the Spokan River in latitude 47° north. This would be an ornamental tree on many sandy and otherwise unproductive wastes. *P. resinosa*, red pine, has its southern limit, according to Emerson, at Wilkesbarre, in Pennsylvania (latitude $41\frac{1}{2}^{\circ}$ north). I have traced it to $56\frac{1}{2}^{\circ}$ of latitude on Methy River, and it crosses the Rocky Mountains to latitude 43° in Oregon. Dr. Gray says that its height in the northern states is from 60 to 80 feet, and Emerson relates that a few years ago it was not uncommon to find trees

s.

allel, its rest the v level of atchewan t. Even Lake, it growing he height parallels. nd all the te food of he flooded a, balsam nks of the ender tree. ms groups d States it s of rivers : its trunk ber of the fuel; and, herally lost equally so. re floating 15 feet in rds. The Asa Gray h, and perering's Sca grows, ac-20 feet in tly resem-Visconsin : New York, cunadensis v England, nsylvania,

of this species in the southern parts of Maine exceeding 100 feet in height, with a stem four feet in diameter. *P. inops*, which was not seen by us on the canoe route to the north of the United States' boundary, extends on the northwest coast from Oregon to Sitka, and ascends Mount Rainier to rear the snow limit. *P. strobus*, white or Weymouth pine, has its equatorial limit on the Alleghanies of Virginia or North Carolina, and it ranges northwards to the south end of Lake Winipeg. In the Middle States this tree has a shaft of 100 feet; and Emerson has collected instances of trees formerly existing which had the extraordinary length of from 220 to 260 feet. Even near its northern termination it is still a stately tree.

Abies balsamea, balsam fir, was not traced beyond the 62nd parallel on the canoe route. It is Le Sapin of the voyagers, who prefer its spray to that of any other tree for laying the floor of a tent or winter bivouack. Dr. Gray traced it on the Alleghanies only to Pennsylvania. In the latitude of Norfolk Sound (57°) it crosses the Rocky Mountains to the Pacific. In Virginia, North Carolina, and Georgia, Pinus fraseri, or the small fruited balsam fir, occupies the Alleghanies to the exclusion of the preceding. It does not reach the great lakes. A. canadensis, hemlock spruce, was observed on the Kamenistikwoya, but not further north than the 49th parallel; though Mr. Tolmie traced it up to the 57th degree of latitude on the shores of the Pacific, and it was observed by Mertens on Sitka. In Maryland this species is found on the Alleghanies only; and Dr. Gray thinks that it ceases to grow in North Carolina and Tennessee. A. alba, white spruce. Of this species we have had frequent occasion to speak in the preceding pages, as it is especially the forest tree in Rupert's Land. It is L'epinette blanche of the voyagers, and the Mina-hik of the Crees. Within the arctic circle it seldom exceeds 40 or 50 feet in height; though in ravines, where it is well-sheltered, and has a suitable soil, it attains twice that altitude. Its age in these high latitudes exceeds 400 years before it shows signs of decay. It most probably has a range from one side of the continent to the other, but has not yet been detected on the

west coast. From the 69th parallel on the Mackenzie, it crosses obliquely to the 61st or 60th on the coast of Hudson's Bay; and it is the common spruce in Canada, Nova Scotia, New Brunswick, and New England, but its southern limits are unknown to Dr. Gray. In Canada the sweet cedar is much used for the thin hoops (varandes) and lining of the bark canoes, being a straight-grained light wood; but in more northern districts the white spruce supplies its place. It is also exclusively used north of Lake Winipeg, for building purposes, sawing into deals, and boat-building. With its tough roots split to a convenient thickness, and used under the Cree appellation of Watap, the pieces of canoe bark are sewn together; and, in districts where birch bark is scarce, a rude canoe is formed of the bark of a spruce fir. A well grown tree, with 30 feet or so free from branches, is chosen ; an incision made down to the wood along one side; and the bark, being skilfully raised in one piece, receives the canoe shape by the two ends being skewered together and stuffed with a few branches to add stiffness. The cargo is then placed in the middle, and two or three Indians will descend a rapid river in this extempore vessel. Before many days, however, it becomes water-logged, and, losing its stiffness, spreads out flatly almost to the level of the water, so as to be nearly useless as well as dangerous. Pieces of the bark are sometimes used for covering the roofs of houses.

A. nigra, black spruce, falls little short in its northern range of the preceding, but in the higher latitudes it is a much inferior tree in numbers, beauty, and utility, and is almost confined to swamps and bogs. According to Emerson, it is in perfection in the northern parts of Maine, or about the 46th parallel, and is less flourishing in more southern localities. It is found on the higher mountains of North Carolina and Tennessee. Up to the Saskatchewan it retains a vigorous growth, beyond which it becomes visibly inferior to the white spruce, its branches being short, irregular, and overgrown with usneæ and other parasitic lichens. A. mertensiana and A. sitchensis grow in the forests of Norfolk Sound on the rorth-west coast in lat. 57° ; Mr. Seeman found the latter extending northwards

TS.

exceeding eter. P. ite to the he northt Rainier outh pine, or North d of Lake ft of 100 rmerly ex-20 to 260 tately tree. the 62nd voyagers, laving the d it on the of Norfolk acific. In seri, or the to the exreat lakes. e Kameniel; though ude on the s on Sitka. nies only; th Carolina species we ling pages, nd. It is hik of the s 40 or 50 -sheltered, ude. Its e it shows one side of ted on the

to the coasts of Beering's Sea; and a spruce which grows on the banks of the Niatok or Buckland River is thought by Sir William J. Hooker to be one of these Sitka species, and decidedly different from A. *alba*, to which Mr. Seeman at first referred it.

Larix americana, American larch, tamarack or hackmatack, L'epinette rouge of the voyagers, and the Wagginā-gan or "tree that bends" of the Crecs, ranges northwards to the arctic circle, and from Newfoundland and Labrador across the continent to the Pacific. It grows in the swamps of the Northern States, and extends southwards to Virginia, where it is confined to the mountains. In high latitudes this tree yields a very heavy wood, so much twisted in the grain as not to be readily worked, but it is tough and very durable. It is a tree of no great importance, and is generally thinly scattered through the forest, and if it is any where grouped in numbers it is on the borders of swamps, where it never attains much height.

Cupressus thyoides, white or sweet cedar, extends from North Carolina to the south side of the Saskatchewan basin. Clumps of it grow on the west side of Rainy Lake, and solitary trees range northwards to the vicinity of Cumberland House in latitude 54°, where a specimen was gathered by Mr. Drummond. C. nutkatensis vel Thuja excelsa inhabits the Pacific coast from Norfolk Sound down to Observatory Inlet and Van-Thuja occidentalis, American arbor vitæ, also couver's Island. called white cedar, has its northern limit on the east side of the Rocky Mountains at Lac Bourbon or Cedar Lake, a dilatation of the Saskatchewan lying between the 53rd and 54th parallels. Michaux mentions the mountains of Virginia as its southern limit. It is a handsome ornament to the banks of Rainy River and the River Winipeg, where it overhangs the water in a picturesque manner; but, as it commonly grows on the occasionally inundated points of lakes and in swamps unmixed with other trees, it has a sombre aspect; and its stems are generally inclined, crooked, and even contracted. T. gigantea, the Wyeth of the Wallamet Indians, grows in the valley of Oregon from the Rocky Mountains to the sea, and northwards to Vancouver's

٧v

319

rs.

ws on the at by Sir s, and den at first

ekmatack, *nā-gan* or the arctic a the con-Northern is confined ds a very be readily tree of no hrough the t is on the ht. from North

i. Clumps litary trees use in lati-Drummond. cific coast and Vanr vitæ, also side of the dilatation h parallels. ts southern ainy River water in a h the occamixed with e generally the Wyeth egon from ancouver's Island. Mr. Douglas found it growing to the height of 170 feet, with a trunk 40 feet in circumference.

Juniperus communis extends from the vicinity of the Arctie Sea to the New England States and Newfoundland. It produces berries freely on elevated grounds within the arctic circle. J. prostrata (Persoon), repens, vel humilis aliorum, is considered by Sir William Hooker to be a variety of J. sabina, which includes the J. virginiana. It has always the prostrate form in Rupert's Land, and was observed within the arctic circle, 1,000 feet above the sea, associated with the preceding, and bearing fruit. Dr. Gray informs me, that it is not found in this prostrate flagelliform condition south of New York and Northern Pennsylvania. The ordinary J. virginiana, red ccdar or savin, ranges to the furthest limits of Texas, and to the country about Santa Fé and Tampas Creek, which is elevated from 3,000 to 5,000 feet above the sea. Col. Emory found it on the 35th parallel, at an altitude of from 6,000 to 7,000 feet, in form of a large tree. Taxus canadensis, American yew or ground hemlock, grows in Massachusetts and Newfoundland, and on the borders of the great lakes northwards to the southern slope of the Saskatchewan basin. It is a very pleasing underwood, with an almost herbaceous aspect, which grows thickly under the shade of many kinds of trees. On the canoe route it never assumes the tree form; but in the valley of Oregon Mr. Douglas found yew trees as large as those of Europe.

The following table is founded on Sir William J. Hooker's Flora Boreali-Americana. Since the publication of that work, Sir George Back's voyage down the Thlewee-choh, Messrs. Dease and Simpson's through the Arctic Sea, Mr. Rae's from York Factory to Repulse Bay, the voyage detailed in the preceding pages, a list of Nova Scotia plants contributed by Mr. Dawson of Pictou, a collection

of plants gathered by Mr. Campbell at Pelly Banks, and Dr. Asa Gray's important Botany of the Northern States, have contributed to our knowledge of the distribution of species, and have been severally had recourse to.

The first zone extends on the eastern side of the continent from latitude 45° to 55°, or it comprehends the St. Lawrence and Saskatchewan basins: it rises obliquely, in accordance with the course of the isothermal lines, in going westward, and on the Pacific coast it includes the 49th and 58th parallels, or Vancouver's and Sitka Islands. It is subdivided into three districts; viz., the eastern forest country, the eastern prairies, and the country west of the crest of the Rocky Mountains.

The second zone comprehends all the country lying between the arctic circle and the extremities of the continent in latitude 72°. It was not found practicable, owing to the way in which the herbaria were formed, to separate the barren ground species from those growing in the woody country, and the zone has been made to include three districts, one of which is Kotzebue Sound, where 273 species have been collected by Chamisso and others.*

The Rocky Mountain ridge, between the 52nd and 57th parallels, has been made a second district of this zone, as many arctic species go southwards along the elevated crest of the ridge. I have not been able, however, to separate the species collected by Drummond in the lower valleys of the ridge from those gathered high up on the peaks. From the untiring diligence and unrivalled quickness of eye of this celebrated collector, we may consider the district as well explored; and had the vertical ranges of the species

* Mr. Seeman, who was employed as botanist in the Herald's late voyages to that quarter, has made a much more ample herbarium of the north-west coast; but as he has just arrived in England as these pages are passing through the press, I can avail myself only partially of his researches for the improvement of the table.

SECOND AND THIRD ZONES.

INTS.

Banks, and ern States, ribution of

the continds the St. obliquely, nal lines, in les the 49th Islands. It stern forest west of the

intry lying of the conticable, owing to separate n the woody le three dis-273 species

nd and 57th this zone, as evated crest to separate er valleys of aks. From as of eye of district as the species

Herald's late herbarium of gland as these only partially been noted, there would be nothing more required for the present investigation.

The third district of the zone comprises the entire arctic country, from Barrow's Point to Davis' Straits.

The third zone lies to the north of the 73rd parallel, and extends from Melville Island to Spitzbergen, including a few species gathered above that parallel in Baffin's Bay, and some collected by Dr. Scoresby on the east coast of Greenland. No American land to the westward of Melville Island in so high a latitude has been discovered; and I have met with no flora of the polar Asiatic islands; but it is probable that, so far north, a nearly uniform vegetation encircles the earth.

Y

STATISTICAL ENUMERATION OF THE NUMBERS OF SPECIES BY THEIR FAMILIES AND GENERA IN DIFFERENT ZONES.

	ones.	Bet to 55 and	First ween o on Lat. West	Zone Lat. East 49° t side.	450 side, o 580	Si F Circ	econd rom le no to 72	Arct rthw N.	ie. le ards	Po Ly of	Thio biar ing 730	d o Zo: No La	r ne. rth it.	
Familles.	Tr'-1 Specles in Three Z	Species in First Zoac.	On Pacific Coast.	On Eastern Prairies.	In East. Woody Distr.	Species in Second Zone.	Kotzebue Sound.	Rocky Mts. 52º to 57º.	Eastern Arctic Distr.	Species in Third Zone.	Lancaster Sound, &c.	North Greenland	Spitzbergen.	Europe, Asia, and Africa.
DICOTYLEDONE/	E 1718	1499	498	391	1130	551	290	340	376	70	57	40	29	403
I. RANUNCULACEA	- 72	59	24	19	47	42	12	31	23	5	5	1	1	22
Clematis -	- 3	3	3	1	1	1		1	1.00					
Thalietrum -	- 5	4	2	2	3	2	1		2					1
Anemone -	- 11	9	4	4	8	8	2	7	4					6
Hepatica -	- 1	1	1	1	1									1
Hydrastis -	- 1	1			1		. 1							
Adonis -	- 1	1			1									
Ranunculus	- 30	23	6	8	20	21	7	17	13	4	4	1	1	13
Caltha -	- 4	3	2	2	2	4		3	2	1	1			1
Trollius -	- 1					1		1						
Isopyrum -	- 1	1			1									
Coptis -	- 2	2	2		1									
Aquilegia -	- 2	2	1		2	2			2					
Delphinium	- 9	2	1		1	2	1	1						
Aconitum -	- 2	2	1		1	1	1	1						
Actæa -	- 2	2		1	2						1			
Cimicifuga -	- 1	1			1									
Hydrastis -	- 1	1			1									
Pæonia brownii	- 1	1	1											
II. MENISPERMACEÆ	- 1	1			1		ŀ							
Menispermum	- 1	1			1									
III. BERBERIDE	- 5	5	8	1	2									
Berberis -	. 3	3	2	l î	1									1
Leontice -	- 1	i	-		l î									
Epimedium	- i	i	1											
IV. PODOPHYLLE #	- 3	9			3									
Jeffersonia -	- 1 ĭ	Ĭ			i									
Podophyllum		1			i									
Undersaltin					1						1	1	1	

ANTS.

SPECIES BY ZONES.

No.26 Eastern Arctic Distr.	HL C C C C C C C C C C C C C C C C C C C	Toyle mor nin T il selonds 0 5	1 I Tancaster Sound, &c.	1 25 North Greenland	retht. Spltzbergen.	No O Europe, Asia, and Africa.
1:	322	4				

NUMBERS OF SPECIES IN DIFFERENT ZONES. 323

				Lones.	F Be 45° E. L 58	twe to sic at.	Zo en 55 le, 1 490 V. 8	ne. Lat. on ind to ide.	Se Fi Cli	con om reie war 72 ⁰	dZc Ar noi ds I	one. etic rth- io	Po L	Thi blar ying 730	rd c Zo 7 N. Lat	ne. of	
Fami	illes.			Total Species in Three	Species in First Zone.	On Pacific Coast.	On Eastern Prairies.	In East. Woody Distr.	Species in Second Zone.	Kotzebue Sound.	Rocky Mts. 52º to 57º.	Eastern Arctic Distr.	Species in Third Zone.	Lancaster Sound, &c.	North Greeniand.	Spitzbergen.	Europe, Asia, and Africa
V. NYMPHAACER	-			4	4	1		4									1
Nymphæa	•	•	-	1	1			1							1		
Nuphar Nelumbium	:	:	:	2 1	2	1		2 1									1
VI. SARRACENIEÆ				1	1			1	Ι,		1	1					
Sarracenia	•	-	-	1	1		i	i	li		i	1					
VII. PAPAVERACE.	/K		-	3	2			2	1	1	1	1	1	1	1	1	1
Papaver	-	-	-	1		1			1	1	1	1	1	1	1	1	1
Stylophorum	-	-	•	1	1			1									
Sanguinaria	-	•	•	1	1			1									
VIII. FUMARIACE.	Æ	-		9	7	3	1	5	4	2	1	2					2
Dielytra	-	-	-	4	3	3	1	1	1	1							
Adlumia	-	-	-	1	1			1									
Corydalis	-	-	-	3	2			2	3	1	1	2					1
Fumaria	-	-	-	1	1			1									1
IX. CRUCIFER 28	-	-	-	104	65	31	21	48	66	16	34	53	16	14	7	5	50
Cheiranthus	-	-	-	1	1	1					1						
Nasturtium	-	-	-	4	4	2	1	3	1	1	1	1					3
Baibarea	-	-	-	3	3	2	2	2	1		1	1					2
Turritis	-	•	-	7	5	2	3	4	6		5	5					1
Arabis -	-	-	-	7	7	3	2	6	2	1	2	2					4
Cardamine	-	-	-	7	5	2	2	5	6	4	3	5	1	1	1	1	4
Dentaria	-	-	-	2	2			2									
Parrya -	•	-	-	2		_			2	1		2	1	1		1	1
Vesicaria	-	-	-	3	3	1	3	1	2		2	1					
Draba -	-	-	-	25	8	5	1	5	22	5	9	16	7	6	2	2	15
Cochlorait	•	•	•	1	1	1		1				-					I
Thleani	•	-	-	8	4	3		1	3	-1		5	4	3	3		4
Intaspi	•	•	-	5	3			8	1			1					3
Caltila	•	•	-	- 4					14	1	-1						1
Hosporie	-	•	-	1	1			1	Ι,								
Sieumhainna	•	-	-	10	4		اے	-		4	_						2
Campling	-	-	-	10	9	P	9	:		4	5	1				,	3
Brava	-	-	-	2	1			1									z
Platynotalum	-	-	-	7					4		귀	4			,		
Eutromo	-	-	-	2					1		-1	4	2	2	1	1	
Orone		•	-	7		, I			z		Ì	z	1	1			
Sinapis	•		-	2	2	4		2									2

y 2

-

324 GEOGRAPHIC

GEOGRAPHICAL DISTRIBUTION OF PLANTS.

				Cones.	F Be 450 E. L 58	twe to sid at.	Zoi en 1 550 e, a 490 . si	ne. Lat. on nd to de.	Sec Fro Cir	cle var 72	dzo Arc nor ds t	ne. tic th-	Po	Thin dar ying y30	d o Zor N. Lat	or of	
Famil	lies.			Total Species in Three 2	Species in First Zone.	On Pacific Coast.	On Eastern Prairies.	In East. Woody Distr.	Species in Second Zone.	Kotzebue Sound.	Rocky Mts. 52º to 57º.	Eastern Arctic Distr.	Species in Third Zone.	Lancaster Sound, &c.	North Greenland.	Spitzbergen.	Europe, Asia, and Africa
CRUCIFERÆ — (cont Lepidium Capsella	tinued.) -	-	-	3 1	3	2	2	2 1	2 1		21	1					1
X. CAPPARIDEÆ Cleome Polanisia	-	:	- -	2 1 1	9 1 1	2 1 1				•							
XI. CISTINEÆ Helianthemum Lechea - Hudsonia	-	-		5 1 3 1	5 1 3 1			5 1 9									
XII. VIOLARIER Viola -	-	-	-	18 18	18 18	4	5 5	15 15	2 2		22	22					1
XIII. DROSERACE. Drosera Parnassia	Æ -	-	-	9 4 5	7 4 3	9 2 1	3 2 1	6 4 2	6 2 4	3 1 2	6 2 4	9 1 2					4 3 1
XIV. Polvoalez Polygala	-	:	-	77	7 7			777	1			1 1					
XV. CARYOPHYLLI L.anthus Saponaria	- -	:	-	66 1 1	48 1	26	13	33 1	35 1	15 1	19	25	11	7	7	7	29 1 1
Silene - Lychnis Agrostemma Moliugo	-	-	-	10 3 1	10 2 1	3	1	7 2 1	2 1	1	1 1	21	1 2	1	1 2	1	3 2 1
Spergula Larbræa Stellaria	-	-	-	4 1 14	3 1 10	2 1 7	2 3	2	3 1 9	1 4	2 1 5	1 1 8	1 3	1	2	1 2	4 1 3
Arenariæ Merckia Cerastium	-		-	19 1 10	11 8	7 5	5 2	9 5	18 1 4	4 1 2	6 9	11 1	3 1	3 1	1 1	1	9 4
XVI. LINEÆ Linum	:	- -	-	3 3	9 3	1 1	22	3 3	1	1	1 1	1 1					1
XVII. MALVACEÆ Malva - Hibiscus Sida -	-	-	•	5 3 1 1	5 3 1 1	1 1	2 1 1	9 2 1									1

)

X

x

X

XX

XX

xx

XXV F

ANTS.

NUMBERS OF SPECIES IN DIFFERENT ZONES.

r	
dZone. Arctic north- rds to P N.	Third or Polar Zone. Lying N. of 73° Lat.
Rocky Mts. 52° to 57°. Eastern Arctic Distr.	Species in Third Zone. Lancaster Sound, &c. North Greenland. Spitzbergen. Europe, Asia, and Afric
2 1 1 2 2 2 2 2 2 2 2 4 5 3 6 1 2 2 4 5 15 19 1 1 1 1 1 1 1 1	2 1 2 1 2 1 2 1 3 4 1 2 1 1 2 1 1 1 2 1 1 1 2 1 1 1 2 1 1 1 2 1 1 1 2 1 1 1 2 1 1 1 2 1 1 1 2 1 1 2 2 1 1 2 2 1 1 1 1 2 1 1 1 1 1 1 1 1 1 2 2 1
4 1 2 1 1 1 1	6 11 3 3 1 1 3 1 1 1 1 1 1 1 1 1 1

			Cones.	F Be 450 E. L 58	irst twe to sid at.	Zor en 1 550 e, a 190 7. si	ne. on nd to de.	Sec Fr Cin	com om cie war 72 ⁰	dZo Arc noi ds t	ne. tic th-	Po	Thin lar ring 73 ⁰	d o Zon N. Lat	r of	
Families.			Total Species in Three 2	Species in First Zone.	On Pacific Coast.	On Eastern Prairies.	In East. Woody Distr.	Species in Second Zone.	Kotzebue Sound.	Rocky Mts. 520 to 570	Eastern Arctic Distr.	Species in Third Zone.	Lancaster Sound, &c.	North Greenland.	Spitzbergen.	Europe, Asia, and Africa
XVIII. Tiliaceæ - Tilia	-	-	2 2	2 2			2 2				-			-		
XIX. Hypericine. Hypericum -	:	-	8 8	8 8			8 8									1
XX. ACERINEÆ - Acer Negundo -	-	:	8 7 1	8 7 1	2 2	1 1	6 5 1									
XXI. Амрелідеж Ampelopsis - Vitis	-	:	4 1 3	4 1 3			4 1 3									
XXII. GERANIACEÆ Geranium - Erodium -	-	-	6 5 1	6 5 1	3 2 1	1	3 3									4 9 1
XXIII. BALSAMINEÆ Impatiens -	:	-	2 2	2 2	1 1		2 2	1 1		11	1 1					
XXIV. Limnanthaceæ Floërkia -	:	-	1 1	1 1			1 1									
XXV. Oxalideze Oxalis		-	5 5	5 5	3 5	1 1	3 3									1
XXVI. RUTACEÆ - Xanthoxylum - Ptelea	:	-	4 3 1	4 9 1			4 3 1									
XXVII. CELASTRINEÆ Staphylea - Euonymus - Celastrus - Myginda - Ilex Prince	-		8 1 2 1 1 1	8 1 2 1 1	2 1 1	1	7 1 2 1 1	1		1						
Nemopanthes -	-	-	1	1			1									
Rhamnus - Ceanothus -	:	-	0 2 4	2 4	3 1 2		3 1 2									

¥ 9

$\mathbf{326}$

GEOGRAPHICAL DISTRIBUTION OF PLANTS.

			ones.	F Bel 450 E. L 58	irat twee to sid t. 4	Zon en I 550 e, an 190 f. sie	ne. .at. on nd to de.	Sec Fre Cir	om cle var 720	a Zo Arc nor ds t	ne. tic th-	T P. Ly	Thin ing 730	d o Zoi N. La	r of	
Families			Total Species in Three Z	Species in First Zone.	On Pacific Coast.	On Eastern Prairies.	In East. Woody Distr.	Species in Second Zane.	Kotzebue Sound.	Rocky Mts. 52º to 57º.	Eastern Arctic Distr.	Species in Third Zone.	Lancaster Sound, &c.	North Greenland.	Spitzbergen.	Europe, Asia, and Airica.
XXIX. TEREBINTHA Rhus	CEÆ -	•	6	6 6	1	_	6			-		_				-
XXX. LEGUMINOS		-	98	88	27	94	51	10	11	11	14	9	2	,		10
Thormonsis -		_	0	9	1	1	51	13			• •	-	1	1		1
Banticia -	-	-	4	a a	1	1	4									'
Medicago -		-	2	9			2									2
Trifolium	-		7	7	6	9	ą									3
Psoralea -		_	5	5	3	ã	Ŭ			1						Ĭ
Potalostemum -	-	-	2	9	ľ	0	9									
Gluoinshiza	-	-	1	1	١.	ĩ	1						1			
Tophrosia	-	-	1	1	•	1	1									
Amorpha	-	-	9	a		9	0									
Robinia -	-	•	3	1		3	2									١,
Dhees	-	-	10	10	0	-	1		6			١,	١,			
Phaca	-	-	12	10	2		2	4	3		1	11		1.		
Oxytropis -	-	-	10	1,4		4	2		4	4	4	ľ '	1	1		3
Astragalus -	•	-	12	10	4	1	2	2		1	1					2
Desinodium -	-	-	9	3			5		Ι.	Ι.	۱.				ł	
Fledysarum -	-	-	2					2	1.	1.	1	ĺ.				
Lespedeza -	-	-	5	2			5	Ι.	١.	١.	Ι.					
Vicia	-	-	5	5	3		4	11	1	1	1		1			2
Ervum -	-	-								Ι.			1			
Lathyrus -	-	-		1	2	3	0	1 2	1	1	2					14
Amphicarpea -	-	-	1	1 3						1						
Apios	-	-					1						1			
Phaseolus -	-	-				١.	1				Ι.					
Lupinus -	-	-	0	0	5		1	2	12	1	11	1				
Gymnocladus -	-	-										1				
Cercis	-		1				1									
														.		
XXXI. ROSACER	-	-	124	106	37	33	78	49	20	25	527	5	4	4	1	19
Cerasus -	-	-	8	8	1	4	8	1			1	1		1		
Prunus -	-	-	1	1	1	1	1	1	1			1	1			
Purshia -	-	-	1	1	1			1				1				
Lütkea -	-	-	1	1	1				1							
Spiræa -	-	-	8	8	6	3	3	2	1	1						1
Gillenia -	-	-	1	1			1	1	1		1	1	1		1	E.I
Dryas	-	-	3	2	1	1	2	3	2	1 2	2 3	1	1	1	1	11
Geum	-	-	4	4	1	2	3	1	1	1		1	1			1
Sieversia -	•	-	5	2	1	1	1	3	3	1	1	1	1			
Comaropsis -	-	-	2	2	1		1	1			1	1	1			1
Rubus		-	18	16	8	3	12	5	3	3	4			1		12

LANTS.

NUMBERS OF SPECIES IN DIFFERENT ZONES.

				ones.	Fi Bet: 45° E. La 58°	rst wee to i side	Zon n L 550 e, ai 190 f	nd nd to de.	Sec Fre Cire V	ond om . cie vare 720	Zo Arc nor is t	ne. tic th-	T Po Ly	hin iar ing	d o Zor N. Lat	r ie. of	
Famil	ies.			Total Species in Three Z	Species in First Zone.	On Pacific Coast.	On Eastern Prairies.	In East. Woody Distr.	Species in Second Zone.	Kctzebue Sound.	Rocky Mts. 52º to 57º.	Eastern Arctic Distr.	Species in Third Zone.	Lancaster Sound, &c.	North Greenland.	Sp tzbergen.	Europe, Asia, and Africa.
Rosace	ued.)					-	-	_	-				-		-	-	
Dalibarda	-	-	-	1	1			1					1		ļ		
Fragaria	-	-	-	3	3	1		2	1		1	1					li
1'otentilla	-	-	-	36	24	6	14	16	19	10	13	12	3	2	3		7
Sibbaldia	•	-	-	1	1			1	1		1						
Chamærhodos	-	-	-	1	1		1		1		1						1
Agrimonia	-	-	-	1	1			1									1
Alchemilla	-	-	-	1	1			i									1 1
Sanguisorba	-	-	_	2	2	9		9	1	1	1	۱,			ł		1
Poterium	-	_	_	ĩ	Ĩ	1 ~		Ĩ	1.	1.		۱ *	1				,
Rosa	-	-	-		6		0				Ι.	1.					
Cratmana	-	-	-	9								10					2
Amulandian	•	•	•	0		1.	Z	0	1:		11	Ι.					
Amelanchier	-	•	-	2		1		2	1			11					
Fyrus -	-	-	-	6	6	4		1 3									
VVVII O				0	0			I	۱.,								
Talla UNAGBA	RIAS	•	•	28	25	9	11		12	5	0	4	12	I	2	Ì.	10
Ephoblum	-	-	-	12	9	5	2	4	17	5	6	4	1 2	1	2	1	6
Gaura -	-	-	-	4	4		14										
Enothera	-	-	-	7	7	2	5	1 3	5					1			1
Clarkia	-	-	-	1	1	1											
Isnardia	•	•	•	2	2			2									1
Circæa -	-	-	-	2	2	1		2	2			1		1			2
																1	
XXXIII. HALOR	AGL	•	-	10	8	2	4	7	5	3	2	4	ł			i	5
Proserpinaca	-	-	-	1	1			1					1	1			
Myriophyllum	•	-	-	3	9		1	1 3	2		1	1 2	2	1	1		2
Callitriche	-	-	-	3	3	1	2	1 5	2					1			2
Hippuris	-	-	-	3	1	1	1	1	3	3	1	1 2	2			1	1
								1									
XXXIV. CERATO	PHYLLE2	E	-	1	1		1	1	1		1	1			1		1
Ceratophyllum	-	-	-	1	1		1	1		1					1		1
VVVV I															1		
AAAV. LYTHRAF	u AS	-	-	3	3	1			5			1			1	1	11
Lythrum	-	-	-	2	2	1		1 2	1	1		1					11
Decodon	•	-	•	1	1			1 1	4								1
VVVVI C	-						.	.				1					
AAAVI. CUCURE	TACER	-	-		2		11	1	1						1		1
Sicyos	-	-	-					1 1							1		1
Echinocistus	-	•	-	1	11	1	1										
VVVVII D				110			Ι.										١.
Maut	LACEAS	•	-	13	11	9	' '		រ ដ	۲ <u>۱</u>		4			1		
Montia	•	-	-			1		1	ł			1			1		11
Portulaca	•	-	-		1 1			1 1	4	1	1		1		1	1	
Lewisia	-	-	-	1 1	6 1	1 1	1	1		1	1	1		F	1	1	

¥ 4

328

GEOGRAPHICAL DISTRIBUTION OF PLANTS.

				Zones.	F Bet 45% E. 1. 58	irst twe to sic at.	Zo en 1 55 le, 1 490 V. si	ne. Lat. on .nd to ide.	Se Fr Ci	con om rcle war ?2	dZd Arc noi ds f	one. ctic rth- to	Pe Lj	Thl plar ring 73 ⁰	rd c Zo N. La	ne. of t.	
Fan	nilies.			Total Species in Three	Species in First Zone.	On Pacific Coast.	On Eastern Prairies.	In East. Woody Distr.	Species in Second Zone.	Kotzebue Sound.	Rocky Mts. 52º to 57º.	Eastern Arctic Distr.	Species in Third Zone.	Lancaster Sound, &c.	North Greenland.	Spitzbergen.	Europe, Asia, and Africa
PORTULACEA (c	ontinued.)					-			-	-			-	-	_	-
Claytonia	:	-	-	19	1 7	7	1	1	3	3	1						
XXXVIII. PAR	ONYCHIE!	E-		2	2		1	1									
Anychia		-	-	1	1			1	1								
Paronychia	-	-	•	1	1		1										
XXXIX. CRASSU	LACEA	-	-	3	3		1	2	1	1		1					1
Sedum -	-	-	-	2	2		1	1	1	1		1					1
Penthorum	-	-	-	1	1			1									
XL. CACTER	-	-	-	2	2	1	1	1									
Cactus -	-	-	-	2	2	1	1	1									
XLI. GROSSULAR	EÆ	-	-	16	15	8	3	10	4	1	2	4					1
Ribes -	-	•	-	16	16	8	3	10	4	1	2	4					1
XLII. SAXIFRAGE	A	-		56	32	13	4	19	35	23	21	18	12	11	7	7	18
Heuchera	-	-	-	6	6	3	1	3	1		1	1					
Tiarella	-	-	-	3	3	2		1									
Tellims.	-	-	-	1	1	1		_									
Mitella	-	-	-	4	3	-	1	2	3		3	1					
Chrysospleniu	m	-	_	2	ï		-	1	ĭ	1	ĩ	i	1	1			1
Saxifraga		-		38	17	7	2	11	28	20	15	15	n	10	7	7	17
Eriogyna	-	_		1	1	· '	1	1	1	1	ĩ	10		10	'	1	•••
Leptarrhena	-	-	•	1	Î			•	î	î	1						
XLIII. UMBELLI	FERÆ	-	-	39	36	13	6	29	10	6	2	5					10
Hydrocotyle	-	-	-	2	2			2			1						1
Eryngium	-	-	-	1	1		li	1									
Sanicula	-	-		1	1	1		1									
Cicuta -	-	-	-	3	3	1		3	2		1	2					1
Zizia -	-	-	-	3	3		1	3	1		1	1					-
Ammi -	-	-		1	1		-	1			-	1					
Carum	-	-	-	1	ī			1									1
Cryptotania	-	-		1	j			1									
Sium -	-	-		2	9	2	1	0								Į	
Bunleurum			_	2	ĩ	~	1	1		1							1
Seseli -	-		,	ĩ	i		1	1		1		1					
Cnidium				1	1		-1		'			1					
Theenium		_		- 1				1									
Liquetioum		-	1	- 1				1	,	,							
Conioselinum	-	-	[2	2	1		1		-	1	1					
e.omoscumum			- 1	~	-	- A I			- A I	- ÷ i	- 1						•

TS.

ne. tic th-	T Pol Ly 7	hir ar 30	d or Zon N. Lat	e. 51		
Eastern Arctic Distr.	Species in Third Zone.	Lancaster Sound, &c.	North Greeniand.	Spitzbergen.	Europe, Asia, and Africa	
	1 1 1 1 1 5 2 1 1	221	1 10	7 7 7	7	1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1

NUMPERS OF SPECIES IN DIFFERENT ZONES.

First Zone. econd Zone Third or Polar Zone. Lying N. of 73° Lat. Between Lat. 45° to 55° on From Arctic Circle north-wards to 72° N. E. side, and Lat. 49° to 58° W. side. Total Species in Three Zones. Europe, Asia, and Africa. Rocky Mts. 52º to 57º. Species in First Zone. In East. Woody Distr. Species in Second Zone Species in Third Zone Lancaster Sound, &c. Eastern Arctic Distr. Familles. **On Eastern Prairies.** North Greenland. Kotzebue Sound. On Pacific Coast. Spitzbergen. UMBELLIFER .- (continued.) Angelica 2 2 • 2 Archangelica -2 ï . Pleurospermum 1 • 1 1 Færula 2 2 2 1 -. • 1 1 Imperatoria --• 1 Pastinaca 1 1 -• • 1 1 Heracleum -• • 1 1 1 1 Laserpitium -• -1 1 1 Polytænia 1 . • -1 1 1 1 Daucus --1 Osmorrhiza 2 1 2 2 2 --ī Conium -1 1 Erigeneia 1 -. 1 -XLIV. ABALIACEÆ 7 7 1 1 6 1 1 • -Adoxa -1 1 1 1 1 -. -1 1 3 Panax -1 23 • • 3 1 3 3 Aralia -_ • . XLV. HAMAMELIDEÆ 1 1 1 . -Hamamelis 1 1 . 7 7 2 2 XLVI. CORNER -2 2 7 $\overline{7}$ 4 22 1 • • 1 7 7 4 Cornus • 1 1 1 XLVII. LORANTHACEÆ 1 1 1 1 -1 1 Arceuthobium -1 1 1 . 1 1 1 XLVIII. CAPRIFOLIACE .. 24 24 7 4 29 6 2 6 3 2 -12 Sambucus 2 1 2 9 1 -2 9 1 8 1 2 1 . . 9 2 Viburnum -1 1 . 1 Diervilla 1 1 . • 1 Lonicera • 8 1 7 3 1 3 1 Triosteum 1 . • • 1 2 1 2 2 1 Symphoricarpus -• 1 -Linnæa . 1 1 1 1 XLIX. RUBIACEAS . 15 15 5 2 14 3 3 2 2 -Hedyotis 3 1 1 3 3 -• • Mitchella 1 • 11 • Cephalanthus -1 -. 9 1 9 1 8 1 Galium • 5 2 3 2 • 3 -Spigelia •

GEOGRAPHICAL DISTRIBUTION OF PLANTS.

		Contes.	Fi Bet 450 E. L 58	rst wee to sid at.	Zon 550 e, a 490 7. si	e. .at. on nd to de.	See Fr Cir	con om clo varo 72°	d Zo Arc nor is to N.	ne. ctic th-	Po Ly 7	Thin biar ying	d o Zo N.	r ne. of	
Families.		Total Specles in Three 2	Species in First Zone.	Or: Pacific Coast.	On Eastern Prairies.	In East. Woody Distr.	Species in Second Zone.	Kotzebue Sound.	Rocky Mts. 52º to 57º.	Eastern Arctic Distr.	Species in Third Zone.	Lancaster Sound, &c.	North Greenland.	Spitzbergen.	Europe, Asia, and Africa
L. VALERIANEJE - Valeriana - Fedia	:	- 6 - 5 - 1	5 4 1	22	1	3 2 1	3 3	1	2 2	1		-	_		1 1
I. Compositation 1. Cichoraceae Sonchus -	:	- 321 - 41 - 7	285 36 6	66 7 3	80 9 2	218 31 5	85 11 1	31 2	60 10 1	57 6 1	7 1	6 1	3 1	1 1	46 12 4
Nabalus - Lygodesma Leontodon Anaruja -	:	- 2	6 1 2 2	1 2 1	2	5 1 2 1	1 2	2	12	2	1	1	1	1	2
Cynthia - Lapsana - Cichorium	:	- 1 - 1	1111			1 1 1									1
Crepis - Hieracium Troximon Krigia -	-	- 3 - 12 - 3 - 1	3 10 2 1		1 3 1	2 10 1 1	2 3 2		222	111					1 2
2. Cynareæ - Centaurea Arctium -	:	- 13	10 1	1	1	10 1 1	5	1	4	2					5 1 1
Carduus - Cirsium - Saussurea	:	- 6	5	1	1	5 3	3 2	1	3 1	2					2 1
3. Vernoniaceæ Vernonia	:	- 2	2 2 2		1 1	2 2									
4. Eupatoriaceæ Kuhnia – Eupatorium	:	- 15 - 1 - 4	14 1 4	2	4	14 1 4	4	2	2	3	2	2			2 1
Liatris - Liatris - Nardosmia Adenocaulon	:		1 5 2 1	111	2 1 1	1 5 2 1	3 1	1	1	3	2	2			1
5. Senecionideæ Silphium - Xanthium	:	- 116	104 6 2	36	29 1	74 6 2	35	21	24	25	4	3	2		23
Ambrosia Parthenium Lea	:	- E - 1	5 5	2	1	4 1									

NTS.

NUMBERS OF SPECIES IN DIFFERENT ZONES.

		Zones.	Fi Bel 450 E. L 580	sid at. 4	Zon 550 e, a 190 . slo	e. .at. on nd lo le.	Sec Fr Cir	concom cle vard 720	Zo Arc nor ls to N.	ne. tic th-	Po Ly	This lar ing	rd o Zoi N. Lat	nc. of	-
Families.		Total Species in Three	Species in First Zone.	On Pacific Coast.	On Eastern Prairies.	In East. Woody Distr.	Species in Second Zone.	Kotzebue Sound.	Rocky Mts. 52º to 57º.	Eastern Arctic Distr.	Species in Third Zone.	Lancaster Sound, &c.	Notth Greenland.	Spitzbergen.	Europe, Asia, and Africa
COMPOSITE, Senecionida-(continued.)					_	-	-		-	-	-	-	
Heliopsis -	• •	1	1		1	1				i					
Echinacea		1	1			1									
Rudbeckia		3	3		2	2									
Lepachys		1	1			1							1		
Coreopsis		3	3			3									
Helianthus		12	12	2	3	9			1						
Bidens -		6	6	1		6					l I				
Gaillardia		2	2	2	2	1		1		ŀ					
Trichophyllum		2	2	2											
Hymenopappus		1	1	1											
Picradenia		1	1	ļ	1							ļ			
Helenium	• •	1	1	1	1	1	1	1	1	1					
Anthemis		1	1			1						1			1
Marruta -		1	1			1								i i	
Achillea -	- •	3	3	2	1	2	2	1	2	2					1
Chrysanthemum		4	3	2		1	3	3	1	2	1	1			2
Pyrethrum	• •	1	1	١.		1	1	1	1	1					1
Cotula -	• •	1	1	1						ί.	1				1
Omalanthus		1	1	1		1	1			1					
Tanacetum		3	9	1		2									11
Artemisia		17	12	6	8	5	րո	7	7	8					5
Gnaphalium		6	6	1		5									4
Antennaria		6	5	3	3	5	4	1	4	2	1	1	1		4
Arnica -		5	4	4			3	1	3	1	1	1	1		
Senecio -		16	12	3	3	10	9	6	5	7	11				1 2
Cacalia -		2	2		1	2									
6. Asteroideæ		134	119	20	36	87	30	5	20	21				1	4
Solidago -	• •	31	31	4	5	30	3	1	3	2		1			li
Aster -		59	49	7	10	37	11	3	5	9				1	l i
Eurybia -		2	2			2			1		1				Ľ
Seriocarpus		1	1			1							1	1	
Tripolium		2	2		2		1			1	1				
Galatella -		2	2			2			[1		1		L
Townsendia		1	1	1	1		1		1	1	1	1			L
Erigeron -		17	12	6	6	9	12	2 1	10	8	3		1		1 9
Dipplopappus		9	9	1	6	4			1		1				Г
Boltonia -		1	1			1	1	1							1
Brachyris		1	1		1			1						1	
Polymnia		1	1	1	1	1		1	1		1				
Madia -		1	1		1		I	1		1		1		1	1
Crinitaria			2 2	1	1		1		1		1				
Donia -		. 4	4		1 9		1 .		1 1	1		1	1	1	

				ones.	Fi Bei 450 E. L 58	rst wee to sid at.	Zor en 1 550 e, a 490 . si	ne. Lat. on ind to de.	Sec Fr Cir	om cie varc 72 ⁰	d Zo Arc nor is to N.	ine, tic th-	Po	This lar ying 30	rd o Zoi t N. Lat	ne. of	
Famil	les.			Total Species in Three Z	Species in First Zone.	On Pacific Coast.	On Eastern Prairies.	In East. Woody Distr.	Species in Second Zone.	Kotzebue Sound.	Rocky Mts. 52º to 57º.	Eastern Arctic Distr.	Species in Third Zone.	Lancaster Sound, &c.	North Greenland.	Spitzbergen.	Europe, Asia, and Africa.
LII. CAMPANULACI Campanula Lobelia	- -	-	-	14 8 6	11 5 6	- - 	1	9 3 6	4 4	99	22	22	1		1	1	3 2 1
LIII. VACCINEAS Vaccinium	-	-	:	16 16	16 16	13 13	5 5	10 10	6 6	22	5 5	4					4
LIV. ERICEAE Gaultheria	-	:	:	40	33	19	9 1	23	18	8	13	10	2	1	1	2	10
Arbutus Andromeda Menziesia Kalmia	-	-	-	4 10 8 9	4 9 4 9	9 6 3 1	1	2 6 1 3	2441	2 2	2 2 4 1	2 3 1	2	1	1	2	2 2 1
Epigæa Rhodora Rhododendron Azalca - Ledum -	-	-	-	1 1 6 1 2	1 1 5 1 2	112		1 1 4 1 2	219	112	2	1					2 1 9
LV. MONOTROPER		-	-	16	15	8	4	10	6	10	5	5					5
Cladothamnus Pyrola - Pterospora Monotropa Chimaphila	- - -			1 10 1 2 2	1 9 1 2 2	1 6 1	3	5 1 2 2	6	2	5	5					5
LVI. JASMINEÆ Fraxinus	<u> </u>	-	-	3 3	3 3	1	1 • 1	3									
LVII. APOCYNEÆ Apocynum	-	:	-	4	4 4		1 1	3 5									1
LVIII. Asclepiad Asclepias	EÆ	:	:	11 11	11 11		2 2	11 11									1 1
LIX. GENTIANEÆ Gentiana Pleurogyne Swertia		-	-	34 23 1	27 16 1	10 6 1	2	19 11 1	19 10 1	8 6 1	9 7	8 6 1					5 2 1
Halenia Sabattia Menyanthes		-		3 2 1	9 2 1	1	1 1	9 2 1	1 1	1	1	3					1
Villarsia	-	-	-	2	2	2		1									

Lancaster Sound, &c.	North Greenland.	Spitzbergen.	Principal very series and the series of the
2 1	1111	1 1 2 2	3 2 1 4 4 10 2 2 1
			2 1 2 5 5
			1 1 1 1 5 2 2 1 1 1 1

NUMBERS OF SPECIES IN DIFFERENT ZONES. 333

Families. Junctifies. Junctifies. <thjunctifies.< th=""> <thjunctifies.< th=""></thjunctifies.<></thjunctifies.<>			,	Zones.	Fi Bet 45° E. La 58°	side	Zon n L 550 e, ar 90 t . sid	e. at. on id o le.	Seci Fro Circ	ond on a cie ard 720	Zor Arc nor s to N.	ne. tic th-	Po Ly	Thir lar ing 30	d or Zon N. Lat.	e. of	
LX. POLEMONIACE	Families.			Total Species in Three 2	Species in First Zone.	On Pacific Coast.	On Eastern Prairies.	In East. Woody Distr.	Species in Second Zone.	Kotzebue Sound.	Rocky Mts. 52º to 57º.	Eastern Arctic Distr.	Species in Third Zone.	Lancaster Sound, &c.	North Greenland.	Spitzbeigen.	Europe, Asia, and Africi
Polemonium 2 2 2 1 1 2 1 1 1 1 1 1 1 1 1 1 1	LX. POLEMONIACER	-		13	11	4	2	8	4	2	2	2	1	_	-	1	2
Phiox - - - 9 7 2 5 3 1 1 1 Collomia - - 2 2 2 1	Polemonium -	-	-	2	2		1	2	1	1	1	1	1			1	1
Collomia - - 2 2 2 1<	Phlox	-	-	9	7	2		5	-3	1	1	1					1
LXI. DIAPENSLACE - - 1	Collomia -	-	-	2	2	2	1	1									
Diapensia - - 1	LXI. DIAPENSIACEA		-	1					1			1					1
LXII. CONVOLVULACE.R	Diapensia -	-	-	1	ł				1			1					1
Convolvulus - - 2 1 <t< td=""><td>LXII. CONVOLVULACE &</td><td>-</td><td>_</td><td>6</td><td>6</td><td>2</td><td>2</td><td>6</td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td>2</td></t<>	LXII. CONVOLVULACE &	-	_	6	6	2	2	6									2
Calystegia - - 2 1 1 1 2 1	Convolvulus -	-		2	2	-		2		İ I							ī
Cuscuta - - 2 2 2 2 2 1 1 4 2 1 2 1 1 4 2 1 </td <td>Calvstegia -</td> <td>-</td> <td>-</td> <td>2</td> <td>2</td> <td></td> <td></td> <td>2</td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td>ī</td>	Calvstegia -	-	-	2	2			2									ī
LXIII. HYDROPHYLLEE 5 4 1 4 2 1 2 1 Hydrophyllum 3 3 3 Eutoca 2 1 1 1 2 1 2 1 LXIV. BORAGINEE 27 25 7 6 20 9 4 5 6 Myosotis 4 3 2 1 1 1 1 1 1 Echinospermum 7 7 1 7 2 1 1 Lithospermum 1 1 1 1 Echium 1 1 1 1 LXV. SOLANEE 2 2 2 1 1 LXV. SOLANEE 8 8 4 2 3 Solanum 1 1 1 1 Datura 1 1 1 1 Niceadra 1 1 1 Niceadra 2 2 2 1 LXVI. OROSANCHEE - 6 6 3 3 4 1 1 Niceotiana - 2 2 2 2 Physalis 2 2 2 1 LXVI. OROSANCHEE - 6 6 3 3 4 1 1 LXVI. OROSANCHEE - 1 1 LXVII. ACANTHACEE - 1 1 Dianthera 1 1 1 LXVII. SCROPHULARINEE - 74 69 27 22 46 20 10 13 14 2 2 1 12 Verbascum 2 2 2 Verbascum 2 2 2 LING AND A A A A A A A A A A A A A A A A A A	Cuscuta -	•	-	2	2	2	2	2									
Hydrophyllum - - 3 3 1 3 2 1 Lutoca - - - 2 1 1 1 2 1 1 LXIV. BonAGINEÆ - - 2 1 1 1 2 1 1 1 2 1	LXIII HYDROPHYLLES		_	5	4		1	4		١,	0	١,					
Eutoca - - - 2 1 1 1 2 1 2 1 1 1 2 1 1 1 2 1<	Hydrophyllum -		_	9	9		1	3	-	•	_	1					
LXIV. BORAGINEE 27 25 7 6 20 9 4 5 6 Myosotis 4 3 2 1 1 1 1 1 1 1 Echinospermum 7 7 1 7 2 1 1 1 Lithospermum 14 13 4 3 11 6 3 3 4 Onosmodium 1 1 1 1 1 Echium 1 1 1 1 1 UXV. SOLANEE 8 8 4 2 3 Solanum 2 2 1 1 Datura 1 1 1 1 Nicatiana 2 2 2 1 Nicatiana 2 2 2 1 Nicatiana 2 2 2 1 Nicatiana 2 2 2 1 LXVI. OROSANCHEE 6 6 3 3 4 1 1 1 Nicatiana 2 2 2 1 LXVI. OROSANCHEE 6 6 3 3 4 1 1 Dipteracanthus 1 1 1 LXVII. ACANTHACEE 1 1 1 Dianthera 1 1 1 LXVII. SCROTHULABINEE 1 1 LXVII. SCROTHULABINEE 2 2 2 Verbascum 3 3 Linaria 2 2 2 2 Verbascum 2 2 2 Linaria 2 2 2 LXVI. SCROTHULABINEE 2 2 2 Linaria 2 2 2 LXVII. SCROTHULABINEE 2 2 2 Linaria	Eutoca -	•	-	2	i		1	1	2	1	2	1					
MY. DOMAGNER: - - 2 2 9 4 5 6 Myosotis - - 4 3 2 1 1 1 1 1 Echinospermum - - 7 7 1 7 2 1	IXIV BODIENT			07	0.5								l	1			
Lithospermum - - - - - 1 <t< td=""><td>Mucrotic</td><td>-</td><td>-</td><td>21</td><td>23</td><td></td><td>0</td><td>20</td><td>1 ?</td><td>1</td><td>10</td><td>0</td><td></td><td></td><td></td><td></td><td></td></t<>	Mucrotic	-	-	21	23		0	20	1 ?	1	10	0					
Lithospernum	Echipotros -	-	-		3	Z				1							3
Dnosmodium - - 1 <td< td=""><td>Lithospermum -</td><td>-</td><td>-</td><td>114</td><td>1.6</td><td></td><td></td><td>.,</td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td></td<>	Lithospermum -	-	-	114	1.6			.,									
Echium - - 1 1 1 1 LXV. SOLANEÆ - - 1 1 1 1 1 LXV. SOLANEÆ - - 2 1 1 1 1 1 Datura - - - 2 2 1 1 1 1 Nicandra - - 2 2 1 1 1 1 Nicotiana - - 2 2 2 1 1 1 1 1 Nicotiana - - 2 2 2 1	Opermedium -	-	-	1	1.	7	10	11	۱°	3	1 3	17	l I	1			14
LXV. SOLANEÆ 8 8 8 4 2 3 Solanum 2 2 1 1 Datura 1 1 1 1 Nicandra 1 1 1 1 Nicotiana 2 2 2 2 Physalis 2 2 2 2 1 LXVI. OROSANCHEÆ 6 6 3 3 4 1 1 1 1 Orobanche 4 4 3 3 2 1 1 1 Epiphegus 1 1 1 1 LXVII. ACANTHACEÆ 2 2 2 Dipteracanthus 1 1 1 Dianthera 1 1 1 LXVII. ScrophularineÆ - 74 69 27 22 46 20 10 13 14 2 2 1 12 Verbascum 3 3 Linaria 2 2 2 2	Echium -	-	-	1	i i	1	1										
LA V. SOLANEA: - - - - 8 8 4 2 3 Solanum - - - 2 1 1 1 Datura - - - 1 1 1 1 Nicoandra - - - 1 1 1 Nicotiana - - 2 2 2 1 LXVI. OROSANCHEÆ - - 2 2 2 1 LXVI. OROSANCHEÆ - - 2 2 2 1 1 Drobanche - - - 4 4 3 3 2 1 1 Conophilus - - 1 <td>VV Carrier</td> <td></td>	VV Carrier																
Solanum - - - 2 2 1 1 Datura - - - 1 1 1 1 1 Nicatdra - - 1 1 1 1 1 Nicotiana - - 2 2 2 1 1 1 Nicotiana - - 2 2 2 1 1 1 1 LXVI. OROSANCHEÆ - - 2 2 2 1 <t< td=""><td>AV. SOLANEA -</td><td>-</td><td>•</td><td>8</td><td>8</td><td>4</td><td>2</td><td>9</td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td></t<>	AV. SOLANEA -	-	•	8	8	4	2	9									
Datura - - - 1 1 1 Nicatdra - - 1 1 1 1 Nicotiana - - 2 2 2 1 1 Nicotiana - - 2 2 2 1 1 1 Nicotiana - - 2 2 2 1 1 1 LXVI. OROSANCHEÆ - - 2 2 2 1 1 1 1 Orobanche - - - 4 4 3 2 1 1 1 Conophilus - - 1 1 1 1 1 1 1 1 LXVII. ACANTHACEÆ1 - 2 2 2 2 2 1 1 Dipteracanthus - - 1 1 1 1 1 1 LXVIII. SCROFHULARINEÆ - 74 69 27 22 46 20 10 14 2	Solanum -	-	-	2			11								1		
Nicatara - - 1 1 1 Nicotiana - - 2 2 2 1 Physalis - - 2 2 2 1 1 1 LXVI. OROSANCHEÆ - - 2 2 2 1 1 1 1 Orobanche - - 4 4 3 3 2 1 1 1 Conophilus - - 1	Datura	-	•							1							
Nicoluma - - - 2 2 2 1 Physalis - - 2 2 2 1	Nicandra -	-	-					1					I	L.			
LXVI. OROSANCHEÆ 6 6 3 3 4 1 1 1 1 Orobanche 4 4 3 3 2 1 1 1 Conophilus 1 1 1 Epiphegus 1 1 1 LXVII. ACANTHACEÆ1 2 2 Dipteracanthus - 1 1 1 Dianthera - 1 1 1 LXVIII. SCROFHULARINEÆ - 74 69 27 22 46 20 10 13 14 2 2 1 12 Verbascum 3 3 Linaria 2 2 2 2 2	Physalis -	-		2			1										
LXVI. OROBANCHEÆ - - 6 6 3 3 4 1																	
Orobanche - - 4 4 3 3 2 1	LXVI. OROSANCHEÆ	-	-	6	6	3	3	4	1	1	1	1		1	1	1	
Conophilus 1 1 1 1 Epiphegus 1 1 1 1 LXVII. ACANTHACEÆ1 2 2 2 Dipteracanthus - 1 1 1 Dianthera 1 1 1 LXVIII. SCROPHULARINEÆ - 74 69 27 22 46 20 10 13 14 2 2 1 12 Verbascum 3 3 Linaria 2 2 2 2	Orobanche -	-	-	4	4	9	3	2	11	1		11					1
Epiphegus 1 1 1 1 LXVII. ACANTHACEÆI 2 2 2 Dipteracanthus - 1 1 1 Dianthera 1 1 1 LXVIII. SCROFHULARINEÆ - 74 69 27 22 46 20 10 13 14 2 2 1 12 Verbascum 3 3 3 Linaria 2 2 2 2	Conophilus -	-	-													1	
LXVII. ACANTHACEÆI 2 2 2 Dipteracanthus 1 1 1 Dianthera 1 1 1 LXVIII. SCROFHULARINEÆ - 74 69 27 22 46 20 10 13 14 2 2 1 12 Verbascum 3 3 Linaria 2 2 2 2	Epiphegus -	-	-	1				1									
Dipteracanthus - 1 1 1 Dianthera - - 1 1 1 LXVIII. Scrophularing - - 74 69 27 22 46 20 10 13 14 2 2 1 12 Verbascum - - - 3	LXVII. ACANTHACEÆL		-	1 2	2 2			5									
Dianthera - - 1	Dipteracanthus	-	-	1	1		1						1		1		
LXVIII. SCROPHULARINE - 74 69 27 22 46 20 10 13 14 2 2 1 12 Verbascum 3 3 Linaria 2 2 2 2	Dianthera -	-	-	1	1	4		1									
Verbascum 3 3 3 Linaria 2 2 2 2	LXVIII. SCROPHULARI	NEÆ	-	74	69	27	22	40	520	10	119	314	5		2 1	1	21
Linaria 2 2 2	Verbascum -		-	1 5	3	3			3	1	1	1	1.	1	1	Ι.	2
	Linaria -	-		9		2	1	1	2							1	ī
Collinsia 4 4 1 3	Collinsia -	-		4		I.	1		3								

334

GEOGRAPHICAL DISTRIBUTION OF PLANTS.

ION OF PLANTS.

				Cones.	F Bel 45 E. L 58	twee to aid at. 4	Zon 6, a 190 . al	ne. .at. on nd to de.	Sec Fr Cli	om om cle varc 720	Arc non is to N.	tic th-	Po	Thin iar ying '30	d o Zoi N. Lat	r ne. of	
Fami	lies.			Total Species in Three 2	Species in First Zone.	On Pacific Coast.	On Eastern Prairies.	In East. Woody Distr.	Species in Second Zone.	Kotzebue Sonnd.	Rocky Mts. 52° to 57°.	Eastern Arctic Distr.	Species in Third Zone.	Lancaster Sound, &c.	North Greenland.	Spitzbergen.	Europe, Asia, and Africa
SCROPHULARINE -	-(conti	nued.)									-						
Chelone	-	-	-	1	1			1									
Penstemon	-	-	-	8	8	5	4	1									1
Mimulus	-	-	-	4	4	2	1	2									
Gratiola		-	-	1	1	1	1	1									
Synthyris	-	-	-	1	1			1									
Limosella	-	•	-	1	1			1									1
Veronica		-	-	11	11	6	4	8	3	1	3	1					6
Gymnandra			-	2	1	Ĭ		Ŭ	1	ī		1					
Romanzoffia		-	_	9	2	0			lî	1						i 1	
Gorardia	-		-	i c	e e	-	1	6	1	•						11	
Orthousanna	-	-	-			0											
Castillain	-	-	-	3	3	3	1										
Castineja	-	-	-	5	5	3	3	4	э	z	3	3					
Euphrasia	•	•	-	1	1			1									
Bartsia -	•	-	-	1	1			1									1.
Rhinanthus	-	-	-	1	1	1	1	1	1		1	נן					1
Melampyrum	-	•	-	1	1		1	1									1
Pedicularis	•	-	-	16	12	3	4	8	11	5	6	8	2	2	1	1	6
LXIX. LABIAT M.			_	40	40	8	10	97	2		9	9					8
Montha			-	1	1	1	1	1	~		-	-					Ũ
Traonus	-	-	-				6	4						•			
Saluia	-	•	-	1		•	4	1									
Manuala	•	-	-	1				1									
Monarda	-	-	-	3	3			3			1						
Blephila	•	-	-	1	1			1									
Pycnanthemun	n	-	-	9	3			3									
Collinsonia	•	-	-	1	1			1									
Cunila -	-	-	~	1	1			1									
Hedeoma	-	-	-	1	1			1									
Micromeria	-	-	-	1	.1			1									
Melissa	-	-	-	1	1			1									
Prunella	-	-	-	1	1	1	1	1									1
Scutellaria	-	-		6	6	3	2	5									1
Lophanthus	-	-	-	9	.9		11	9									
Nepeta	-	-	-	Ĩ	ï			ĭ									1
Dracocenhalun	1	-	_	1	i		1	1	1			1					1
Physostumia	-	-	_	1	;	1			1 1		^	1					
I missostegia	-	-	-			1	1	1									,
T	•	-	-					1									
Leonurus	-	-	-		1			1									1 I
Galeopsis	•	-	-	1	1			1									1
Stachys	-	-	-	3	3	1	1	3	1		1	1					11
Marrubium	-	-	-	1	1			1									1
Teucrium	-	-	-	1	1			1									
Trichostema	-	-	-	1	1			1									
Trichostema	-	-	-	i	i			1									

ľ

Thola yir 73		l or Zon N.	e. of		
	Lancaster Sound, ac.	North Greenland.	Spitzbergen.	Europe, Asia, and Afric	
					1
2		2	1	2	1 1 1 6 8
					1

NUMBERS OF SPECIES IN DIFFERENT ZONES.

			cones.	E. 1	at.	6, 8 490 7, 81	on ind to de.	Ċĥ	cie varo 72 ⁰	noi is to N.	th-	Ľ	ving 730	Zo N La	ne, of	
. Families.			Total Species in Three Z	Species in First Zone.	On Pacific Coast.	On Eastern Prairies.	In East. Woody Distr.	Species in Second Zone.	Kotzebue Sound.	Rocky Mts. 520 to 570.	Eastern Arctic Distr.	Species in Third Zone.	Lancaster Sound, &c.	North Greenland.	Spitzbergen.	Europe, Asia, and Africa.
LXX. VERBENACER	-	•	15	15	5	5	13	4	-	4	2		-		-	6
Verbena -	•	•	6	6	2	2	5									1
Iltricularia -		•	6	1	1	1	6			0						
Pinguicula -	-	-	2	2	2	2	1	2		2	1					3 2
LXXI. PRIMULACE &	-	-	23	17	9	7	13	11	7	4	8					10
Dodecatheon -	-	-	2	2	2	1	1	2	2		1					
Androsace -	-	-	2					2	2	2	2					2
Douglasia -	-	-	2	1		1		l			1					
Primula -	-	-	6	3	1	1	3	5	3	2	3					4
I rientalis -	-	-	3	3	2	_	1									1
Glaux	-	-	0	0	2	1	0	1			- 1					
Samolus -	-	-	1	1	1	1	1									1
LXXII Prowercowe			0	0	,		9	,			,					,
Statice	-	-	2	2	1		2	i	1		1					1
LXXIII. PLANTAGINE &		_	5	5	3	1	4	9			9					9
Plantago -	-	-	5	5	3	i	4	2			2					3
LXXIV. NYCTAGINER	-	-	3	3	1		2									
Oxybaphus -	-	-	1	1	1		-			- 1						
Allionia -	-	-	2	2			2									
LXXV. AMARANTHACEÆ	-	-	6	6	1	2	6									
Amaranthus -	•	-	6	6	1	2	6									
LXXVI. CHENOPODEÆ	-	-	20	19	6	11	12	4	3		1					14
Salicornia -	-	-	2	2	1	1	1									2
Salsola	-	-	1	1			1									1
Corispermum -	-	-	1	1	1	1	1									1
Eurotia -	-	-	1	1		1	,									1
Chanapadium	-	-	2	2	0	1	1	,			1					
Ambring -	1		1	1	3	°	1	-			-					5
Atriplex -	-	-	4	4	1	1	3	3	3							3
LXXVII. PHYTOLACE			1	1			,									
Phytolacca -				i			i									

336

GEOGRAPHICAL DISTRIBUTION OF PLANTS.

		Cones.	Fi Bet 450 E. L. 58	to shi	Zor n I 550 e, a 190	ie, .al. on nd to de.	Sec Fri Cir	om cle arc 72º	Arc nor is to N.	tic th-	T Pc Ly	Thir Diar May 30	d o Zo N. Lat	r ne. of	
Families.		Total Species in Three 2	Species in First Zone.	On Pacific Coast.	On Eastern Prairies.	In East. Woody Distr.	Species in Second Zone.	Kotzebue Sound.	Rocky Mts. 52º to 57º.	Eastern Arctic Distr.	Species in Third Zone.	Lancaster Sound, &c.	North Greenland.	Spitzbergen.	Europe, Asia, and Africa
LXXVIII. POLYGONER -	-	34	32	16	8	22	11	6	6	9	2	2	2	2	17
Konigia	-	1					1	1	1			,			1
Rumer	-	10	10	5	4	9	5	1	2	4	l '	1	•	1	8
Polygonum	-	18	17	7	2	19	4	3	1	4	1	1	1	1	7
Eriogonum	-	4	4	3	2										
LXXIX. LAURINES	-	2	2			2									
Benzoin	-	1	ī			1									
Sassafras	-	1	1			1									
XXX ELEMANTE		q	q		2	9	9		0	9					
Eleagnus	-	ĭ	1		ĩ	1	1		ĩ	ĵ					
Shepherdia	-	2	2		1	1	1		1	1					
XXXI Turners		1				1									
· Direa		i	li			1									
			-												
LXXXII, SANTALACER -	-	2	2		2	2	1		1	1					
Comandra	-	2			Z	2	1		1	1					
LXXXIII. ARISTOLOCHIAE	-	1	1			1									
Asarum	-	1	1			1									
LXXXIV. EMPETERS		1	1			1	1	1	١,						1
Empetrum	-	1	i			i	1	i	i	i					1
EXXXV. EUPHORBIACER	•	87	87	20	20	8									1
Acalypha		i	i			i									
LXXXVI. URTICEAE -	-	10	10			10									4
Pilea		1	1	1	1	3	1			1					ſ
Parietaria	-	1	i			1									
Boehmeria	-	1	1			1									
Humulus	-	1	1			1									1
Cannabis	-	1	1			1									1
LXXXVII. ULMACER -	-	3	3	1		8		-							
Ulmus	-	2	2			2									
Celtis	-	1	1	1		1									

xc

2

х

M X C

v
NUMBERS OF SPECIES IN DIFFERENT ZONES.



 Х

		ones.	B 4 1 La	First etwee 5° to 2. sid t. 49° W.	Zon- en L. 55° c e, an b to l side.	e at. on id 580	Se F Cl	ron rcie wai 72	d Zo Arc nor rds to N.	ne. tic th-	Pely	This plar ying 30	d or Zo N. Lat	ne. of	
· Families.		Total Species in Three Z	Species in First Zone.	On Pacific Coast.	Oo Eastern Prairies.	In East. Woody Distr.	Species in Second Zone.	Kotzebue Sound.	Rocky Mts. 52° to 57°.	Eastern Arctic Distr.	Species in Third Zone.	Lancaster Sound, &c.	North Greenland.	Spitzbergen.	Europe, Asia, and Africa.
LXXXVIII. SAURUREÆ Saururus -	•	1	1			1	_				-	-		-	
LXXXIX. JUQLANDINEA Juglans Carya		4 2 2	4 2 2			4 9 2									
XC. SALICACEÆ - Salix Populus		48 44 4	39 35 4	8 7 1	6 5 1	36 32 4	27 25 2	8 7 1	14 12 2	24 22 2	3 3	1	3 3]]	14 13 1
XCI. BETULACEÆ - Betula Alnus	• • •	11 8 3	11 8 3	4 2 2	4 2 2	9 7 2	6 4 2	3 1 2	6 4 2	5 3 2					2 1 1
XCII. PLATANACEÆ Platanus -	•	1	1 1			1 1									
XCIII. CUPULIFERÆ Quercus - Fagus Carpinus - Ostrya Corylus	•••••	15 10 1 1 1 2	15 10 1 1 1 2	2 1 1	2	14 9 1 1 1 2									
XCIV. MYRICACEÆ Myrica Comptonia -	•	3 2 1	3 2 1	2 2	1 1	2 1 1	1 1		1 1	1 1					1 1
XCV. CONIFERÆ - Pinus Abies Larix Cupressus - Thuja		20 5 6 1 2 2	20 5 6 1 2 2	9 2 3 1 1	6 1 2 1	14 4 1 1	7 1 3 1	1	4 2	6 1 2 1					2
Juniperus - Taxus	-	3 1	9 1	2	2	2	2	50	2	2		-			2
XCVI. COMELINEZE Tradescantia -	-	554 1 1	193 1 1	102	1 22	399 1 1	198	53	120	140	21	20	9	9	199
VOL. II.				Z											

GEOGRAPHICAL DISTRIBUTION OF PLANTS.

<u> </u>				Cones.	Fir B La 55' side 49 V	st 2 letw t. 4 o or , an o to V. s	Zone seen 5° t d La 58° ide.	o at.	Seco Fro Circ W	m d le i ard	Zon Arc nor s to N.	ne. tic th-	T Po Ly 7	hir iar ing 3°	d o Zor N. Lat.	r ie. of	
Familles.				Total Species in Three 2	Species in First Zone.	On Pacific Coast.	On Eastern Prairies.	In East. Woody Distr.	Species in Second Zone.	Kotzebue Sound.	Rocky Mts. 52º to 57º.	Eastern Arctic Distr.	Species in Third Zone.	Lancaster Sound, &c.	North Greenland.	Spitzbergen.	Europe, Asia, and Africa
XCVII. ALISMACEÆ Sagittaria - Alisma -	-	-	-	3 1 2	9 1 2	2 1 1		3 1 2									3 1 2
XCVIII. JUNCAGINEÆ Scheuchzeria	-	-	:	4	4	1	3 1 9	4	2 1		2	1					3 1 2
XCIX. AROIDEÆ	-	<i>a</i> .	-	19	12	2	3	11	3	1	3	2					10
Orontium - Arum -	:	-	:	1				1									
Calla - Symplocarpus Lemna -	-	-	-	1 2 9		2		1									2 3
Sparganium Typha -	-	:	:	2	2		1	21	2 1	1	2						3 1
C. NAIADES - Zannichellia	:	:	:	14	14	1 2	3	14	1		1	1					9 1
Naias - Ruppia - Potamogeton	-	-	-	1					1								17
CI. SMILACEÆ - Smilax	-	-	-	19			5 8	16	5 3		:	3	2				4
Streptopus - Uvularia -	-	-	-			3 5	3 2	2	5								2
Polygonatum	-	-	-				2 4					3					i
CII. MELANTHACEÆ Limnanthemum	:	-	-	1	610 1	6 ' 1	7 4	1:	3 4		2 :	3	3				3
Xerophyllum Helonias -	-	-	-		1	1			1		1	1	1				
Veratrum - Tofieldia - Medeola -	-	-			1 9 1	1 3	1 1 2 9	2	1 1 3 9	2	1	1	2				2
Trillium -	-	-			7	7	1		6								1
Lillum	-			:	9 3	9 3	6 2	1	5 3								1

NTS.

 $\{ l \}$

.

cone. rctic orth-to N. Third or Polar Zone. Lying N. of 73° Lat. Spitzbergen. 8 Lurope, Asia, and Africa. Rocky Mts. 52º to 57º. Species in Third Zone. Lancaster Sound, &c. Eastern Arctic Distr. North Greenland. 1 2 1 1 1 1 1

1

NUMBERS OF SPECIES IN DIFFERENT ZONES.

				Zones.	Fi La 5 side 49	rst Betw at. 4 5° c , an 0° to V. 8	Zon veer 5° t on E id L o 58 side.	n to at.	Sec Fre Cir	ond om cie /arc 72 ⁰	Zo Arc nor ls to N.	ne. tic th-	Po Ly	Thin lar ying 730	d o Zou N. Lat	r ne. of	
Familie				Total Species in Three	Species in First Zone.	On Pacific Coast.	On Eastern Prairies.	In East, Woody Distr.	Species in Second Zone.	Kotzehue Sound.	Rocky Mts. 52º to 57º.	Eastern Arctic Distr.	Species in Third Zone.	Lancaster Sound, &c.	North Greenland.	Spitzbergen.	Europe, Asia, and Africa
LILIACE - (continued, Fritillaria -)			0	0	0											1
Erythronium Calochortus	-	-	-	3	$\frac{2}{3}$ 1	1 1 1		2									1
CIV. ASPHODELEÆ	-	-	-	12	11	7	3	4	2	1		2					1
Anthericum	•	-	-	1					1	1		1					
Allium -	•	-	-	3	8	4	3	4	i			1					1
Camassia -	-	-	-	1	1	1											
Brodiæa -	•	•	-	2	2	2											
CV. PONTEDERIACEÆ	-	-	-	2	2			2									
Pontederia -	-	-	-	1	1			1	1	1	ł						
Leptanthus -	-	-	-	1	1			1									
CVI. RESTIACEÆ	•	-	•	1	1			1									
Eriocaulon -	•	-	-	1	1		!	1									1
CVII. JUNCEÆ	-	-		09	90	10	9	15	17	8	14	19	۱.	0	0	1	
Luzula -	-	-		7	6	4	2	4	5	4	5	10	0	1	1		17
Juncus -	-	-	-	16	14	6	7	11	12	4	9	10	ĩ	1	i		10
CVIII. HYDROCHARIDE	Æ	-	-	2	2			2									Ι.
Valisneria -	-	-	-	1	1	1		1									I î
Udora -	-	-	-	1	1			1									ſ
CIX. ORCHIDEÆ		-	-	54	52	217	9	43	10	9	9	8					1
Microstylic -	-	-	-	3	3	1		2					1	1			
Liparis -	-	-	-	2	2 2	2		2			1	1	1	1			
Corallorrhiza	-	-	-	3	8 8	3 2	2 1	2	1	1			1		1		
Amplectrum	-	-	-	1	1	4		1			1			1	1		
Catypso -	-	-	-	1	1		1	11	1		1	1			1		1
Guernadania	-	-	-	12	1	1		11						1	1		1
Platanthana	-	-	-	1				1,1					1				1
Paristylus -	-	-	-	20	ľ	1	15	115	2 3	1	1 3	1 8	5				
Arothusa -	-	-	-				1	1									I
Pogonia -		-				5			5				1			1	
Calopagon -		-	-								1		1		1		1
Spiranthes -	-	-	_				Ι.				1					1	
Goodvera -	-	-	_	6		1	۱		1					1			
Listera -	-	-	_			2 0			,	1	1	1	1			1	
Cypringdium		-	_							1.			1		1	1	

(A)

						Zones.	F Be 45 E 1 5	'irst etwe o to . sic Lat.	Zo en 559 de, a 499 V. s	one. Lat. on and to side.	Sec Fre Cin	ond m cle var 720	ds to N.	ne. tic th-	Pe	This plar ying 73°	rd o Zo N. Lat	ne. of	
	Fami	lies.		•		Total Species in Three 2	Species in First Zone.	On Pacific Coast.	On Eastern Prairies.	In East. Woody Distr.	Species in Second Zone.	Kotzebue Sound.	Rocky Mts. 520 to 570.	Eastern Arctic Distr.	Species in Third Zone.	Lancaster Sound, &c.	North Greenland.	Spitzhergen.	Europe, Asia, and Airica
C	X. IRIDEÆ Iris - Sisyrinchium	-	:		•	8 7 1	7 6 1	2 1 1	1	7 6 1	3 2 1	22		1		-			
С	XI. Нурохідел Hypoxis	s -	-		-	1	1			1									
С	XII. Dioscore. Dioscorea	æ	-		-	1 1	1 1			1 1									
	KIII. CYPERACE Carex - Elyna - Eleocharis Scirpus Eriophorum Dulichium Cyperus Rhynchospora	Æ - - - - -				218 183 2 7 10 8 1 5 2	184 153 6 10 7 1 5 2	51 39 4 3 4	33 24 3 5 1	160 131 5 10 7 1 4 2	102 91 2 3 6	12 9 3	54 45 2 2 5	77 68 1 2 6	5 3 2	5 3 2	1	2 2	81 65 5 4 6
C	XIV. GRAMINE. Lecrsia Hydropyrum	лЕ -	-		-	153 2 1	134 2 1	48	46	96 2 1	51	24	32	36	13	13	5	6	56
	Phleum Phalaris Hierochloe Anthoxanthum Milium	-			-	4 2 1 3 1 1	3 2 1 2 1 1 1	2 2 1	1 1 1 1	3 1 1 2 1 1	2 1 1 3	1 2	2 1 2	2 1 1 2	1	1 2	1	1	3 2 1 2 1 1
	Panicum Holcus Oplismenus Setaria Cenehrus	-				8 1 1 1 1	8 1 1 1 1	1	3	8 1 1 1									1
	Oryzopsis Stipa - Muhlenbergia Phippsia Colpodium	- - -	••••			9 4 6 2 3	3 4 6 1	2	2 3 1	3 1 5 1	22	22	1	2	1	1		1	1
	Vilfa - Agrostis Calamagrostis Ammophila	-	-		-	1 6 10 1	1 6 9	6 3	1 2 2	3 7	9 9	2	9 9	1 9					421

NTS.

on cl rt to	ic h-	T Pol Lyi 7	hird lar 3° I	l or Zon N. o Lat.	e. of	а.	
Rocky Mts. 52 to 31.	Eastern Arctic Distr.	Species in Third Zone.	Lancaster Sound, &c.	North Greenland.	Spitzbergen.	Europe, Asia, and Afric	
54	4755622255	77812 6 361 21 1 2 1 1 2 3 3 1 3 3		5 3 1 2 1 2 1	5511	2282	

NUMBERS OF SPECIES IN DIFFERENT ZONES.

1

		Zones.	l Be 45 F La	First etwee o to f t. sid at. 49 W. s	Zone n La 55° o e, an c to t side.	n d 38°	Se F C wal	cond rom ircie rds to	Zon Arcti north 72°	e. c 1- N.	Po Ly 7	Thin iar ing 3 ⁰	nd o Zoi N. Lat	r ne. of	÷
Familles.		Total Species in Three	Species in First Zone.	On Pacific Cuast.	Cn vastern Prairies.	In East. Woody Distr.	Species in Second Zone.	Kotzebue Sound.	Rocky Mts 520 to 570	Eastern Arctic Distr.	Species in Third Zone.	Lancaster Sound, &c.	North Creenland.	Spitzbergen.	Europe, Asia, and Afric
GRAMINEE - (continue	ed.)								-		-	-		-	
Graphephorum	-	1	1			1									
Phragmites -	-	1	1	1	1	1									1
Spartina -	-	1	1			1	1			1			ļ		
Eutriana -	-	1	1		1		1		1				Ì		
Deschampsia	-1	2	1	1	1	1	1	1		1	1	1			1
Dupontia -	-	1	1			1	1	1		1	[1	1	i i	1	1
Boutelouia -	-	2	2			2			i						
Aira -	-	3	2	1		2	1		1		١.	3			1
1 risetum -	-	2	2	1		1	1		1	1	11		1		
Avena -	-	2	1		1	1	2		- 1	1		1			-
Dantnoma -	•	00	1	10	6	10		_	_				0		10
Foa	•	20	23	10	0	15	11	1	5	9	1 '			2	15
Glucaria		0	8	0	q	5	1			1					0
Blouropogon		9	°	3	0	5	1			1	۱.	1	1		-
Rahoules -	_	9	9		1	9	1		,		ľ	1			
Catubrosa -		1	ĩ			Ű	1		1						1
Koeleria -		i	i	1	1	i									i
Festuca -		9	6	9	3	2	5	3	3	4	1 :	1		1	4
Bromus -		3	3	ĩ	Ŭ	2	ĭ	ĩ	ĭ	i	1	-		1	
Ceratochloa -	_	1	1	1		-	•	-	-	-					
Briżopyrum	-	1	1	i	1										
Triticum -	-	3	3	2	2	1	2		2	1					3
Elymus -	-	8	8	3	2	6	3	1	3	2					5
Asprella -	-	1	1			1									
Hordeum -	-	2	2	2	1	1	1			1					1
Andropogon	-	4	3		3	1	1		1						
ACROGENES	-	71	67	26	18	57	30	6	21	24					26
CXV. EQUISETACE &	-	9	9	3	2	8	6	2	4	5					8
Equisetum -	-	9	9	3	2	8	6	2	4	5					8
CXVI. FILICES -	-	47	43	16	11	35	20	3	13	16	1				20
Polypodium -	-	3	3	3		3	3		1	3					3
Woodsia -	-	3	S		1	3	2		1	2					2
Cistopteris -	-	3	2	_	1	2	2	1	2	1					2
Aspidium -	-	11	10	5	2	9	4	2	3	4		1			6
Unoclea -	-	1	1			1									
Struthiopteris	-	1	1			1									
Athyrium -	-	1	1	1	1	1	1		1	1					
Rischnum -	-		5	,		3	2		1	1					3
meennum •	-		- 4	- 4			1 1				1	1		1	1

z 3

	Zones.	l Be 45 H La	First etwee 0 to 0. sid at. 49 W.	Zone 55° o e, an o to f side.	n d 80	Se F Ci wai	rcie i rcie i rcis to	Zon Arcti north 72°	e. c N.	T Pol Ly 7	'hir lar ing 30 J	d oi Zon N. Lat.	r ie. of	
Families.	Total Species in Three	Species in First Zone.	On Pacific Coast.	On Eastern Prairies.	In East. Woody Distr.	Species in Second Zone.	Kotzebue Sound.	Rocky Mts. 52º to 57º.	Eastern Arctic Distr.	Species in Third Zone.	Lancaster Sound, &c.	North Greenland.	Spitzbergen.	Europe, Asia, and Africe
FILICEF - (continued). Pteris - Cryptogramma - Adiantum - Cheilanthes - Dicksonia - Osmunda - Schizæa -	3 1 1 1 1 3 1	3 1 1 1 1 3 1	1 1 1 1	1	3 1 1 3 1	I 1 I		1	1		-			I I I
CXVII. Lycopodium - Lycopodium - Selaginella -	5 12 11 1	5 12 11 1	2 7 7	4 5 5	1 11 10 1	2 4 4	2 2	2 4 4	1 3 3					1 7 7
CXVIII. Hydropterines Isöetes Salvinia Azolla	3 1 1 1	3 1 1 1			3 1 1 1									1 1
Dicotyledones - Monocotyledones	1725 554	1499 493	498 162	391 122	1130 399	568 198	271 53	340 120	377 146	69 21	57 20	40 9	29 9	403 188
Acrogenes -	2279 71	1992 67	660 26	513 18	1529 57	766 30	324 6	460 21	523 24	90	77	49	38	591 26

Obs. In the preceding table, and in that which follows, species that range to several zones are enumerated in each. The proportionate numbers of the second table are found by dividing the whole *Phanerogamæ* of a district by the numbers of each family in that district, and they may, therefore, be considered as denominators of fractions having 1 for a numerator. The proportions vary remarkably in different districts. The predominance

The proportions vary remarkably in different districts. The predominance of Compound Flowers, Leguninous and Rosaccous plants in the Prairies, combined with the paucity of Saxifrages, Gentians, and Ericaceous plants, affect the proportions of the other families materially. The Grasses, as might b = expected, are more numerous in the Prairies than elsewhere, with the remarkable exception of the Polar Zonc, in which the Gramineæ form one-seventh of the species, and in conjunction with the Cruciferæ, Carnophylleæ, and Saxifrageæ, constitute more than half the Phanerogamæ. The small numbers of Asters, Willows, and Carices, on the Pacific coast, modify the numbers of that district. NTS.

_							
	T Pol Ly 7	hird ar 2 ing 3° I	i or Ion N.	e. of			
Eastern Arctic Disu-	Species in Third Zone	Lancaster Sound, &c.	North Greenland.	Spitzbergen.	Europe, Asia, and Africa		
	1 1 3 3 3 77	69 21	57	40 9	29.4 91	1 1 7 7 1 1 1 1003 88	
	523 24	90	77	49	38	26	

species that range ate numbers of the α of a district by therefore, be conator.

The predominance the in the Prairies, Ericaceous plants, The Grasses, as an elsewhere, with he Gramineæ form Cruciferæ, Caryohanerogamæ. The cific coast, modify

PROPORTIONS OF PHANEROGAMOUS VEGETATION. 343

PROPORTIONATE NUMBERS OF SOME OF THE PRINCIPAL FAMI-LIES TO THE WHOLE PHANEROGAMOUS VEGETATION OF THE SEVERAL DISTRICTS OF THE THREE ZONES.

		First Zon	ne.	Se	cond Zon	ne.	at.
Families.	Pacific Coast.	Estern Prairies.	East. Woody Dist.	Kotzebue Sound.	Rocky Mts. 52° to 57°.	Arctic Region.	Polar Zone. North of 73 ⁰ Li
Monocotyledones	- 4.1	4.2	3.8	6.1	3.8	3.6	4.3
Compositae -	- 10.0	6.4	7.0	9.0	7.7	25.1	12.9
Cyperaceæ -	- 12.9	15.9	9.5	27.0	8.5	6.8	18.2
Gramineæ .	- 13.8	11.2	15.9	13.1	14.4	14.5	7.0
Rosaceæ -	- 17.8	15.6	19.5	12.0	18.4	19.4	18.0
Cruciferæ -	- 21.3	24.4	31.9	17.1	13.9	9.8	6.1
Leguminosæ -	- 24.5	15.1	30.0	23.1	41.8	37.3	45.5
Scrophularineæ	- 24.4	23.3	32.2	27.0	35.4	37,•1	45.5
Caryophylleæ	- 25.4	39.5	46.3	18.0	24.2	20.9	8.3
Ranunculaceæ	- 27.5	27.0	32.5	23.1	14.8	22.7	18.2
Ericeæ	- 34.7	171.0	66.5	36.0	35.4	52.2	45.5
Oreliideæ -	- 38.8	57.0	35.6	108.0	57.6	65·3	
Saxifrageæ -	- 50.8	128.2	80.5	12.5	21.9	29.0	7.5
Umbelliferæ -	- 50.8	85.5	54.5	54.0	230.0	104.4	
Gentianeæ -	- 66.0	256.5	80.2	40.5	51-1	65.2	
Coniferæ -	- 73.3	85.5	106.4	324 •0	125.0	87.0	
Labiatæ -	- 82.5	51.3	41.3		23.0	26.2	
Salieaceæ -	- 82.5	85.5	42.5	40.5	32.9	21.7	30.3
Boragineæ -	- 94.3	85.5	76.4	64.8	11.5	87.0	

A list of the plants collected by Mr. Seeman on the coasts of Beering's Sea having been received subsequent to the first part of the preceding tables having passed through the press, some emendations are requisite. The numbers of Dycotyledones, in p. 342., are to be substituted for those in p. 322., and the following changes made in the column headed Kotzebue Sound: viz. Ranunculaceæ, 14; Cruciferæ, 19; Caryophylleæ, 18; Leguminosæ, 14; Rosaceæ, 27; Portulacea, 5; Saxifrageæ, 26; Composite, 36; Ericeæ, 9; Boragineæ, 5; Scrop iularineæ, 12; Verbenaceæ, 1; Diapensiaceæ, 1. These occasion some slight alterations in the first column of the second zone, and in the total number of plants.

The following Table of the Distribution of the Carices was drawn up by Dr. Boott, whose intimate acquaintance with that genus of *Cyperaceæ* renders it of the highest value to the student of the Geography of Plants: —

My dear Sir John,

London, May 6th, 1850.

I have examined the Carices you brought from your last excursion to America. They are —

C. scirpoidea Michx. Arctic Sea coast.

ursina Dewey. Arctic Sea coast.

glareosa Wahlg. Arctic Sea coast.

stans Drejer. Arctic Sea coast.

saxatilis L. Arctic Sea coast.

compacta Brown. Arctic Sea coast.

fuliginosa St. & Hoppe. Arctic Sea coast.

livida Willd. Arctic Sea coast.

Novæ Angliæ Schwz. Arctic Sea coast and Meiny Portage.

canescens L. (& b.) Arctic Sea coast and Methy Portage.

var. polystachya. Lakes Superior, Rainy, and of the Woods.

adusta Boott. Methy Portage.

,,

siccata Dewey. Methy Portage, Saskatschawan.

lanuginosa Michx. Methy Portage, Saskatschawan.

lenticularis Michx. Methy Portage.

Houghtonii Torrey. Methy Portage.

Racana Boott. Methy Portage.

utriculata Boott. Methy Portage, Lakes Superior, Huron.

aqua^eilis Wahlg. Methy Portage, Lakes Superior, Huron, Fort Simpson and Chipewyan.

umbellata Schk. Methy Portage, Lakes Superior, Huron, Fort Simpson and Chipewyan.

oligosperma Michx. Lake Superior.

aristata Br. Lakes Superior and Huron.

scoparia Schk. (§ b.). Lakes Superior and Huron, Winipeg, Athabasca.

vulgaris Fries. Lakes Superior and Huron, Winipeg.

retrorsa Schwz. Winipeg River, Lake of Woods.

pedunculata Muhlg. Winipeg River, Rainy Lake.

intumescens Rudge. Lake of Woods, Rainy Lake.

Ederi Ehrh. Rainy River and Lake.

TABLE OF DISTRIBUTION OF CARICES. 345

Pennsylvanica Lam. Saskatschawan, Winipeg, and Cumberland Lakes.

incurva Light. Valleys of the Sask. and Mack.

Of the above, C. Racana is new, and C. stars uew to British America.

I find from my notes that the number of Carices in North America is 250; of which 178 are found in all Arctic America, including 97 common to Arctic America and the States, leaving 81 Arctic species.

Of these 81, there are 36 common to Europe, leaving 45 peculiar to Arctic America.

Of the 97 found in Arctic America and the States, 28 are common to Europe, leaving 69 exclusively American.

There are in the States, besides, 72, of which 4 only are European, leaving 68 exclusively American.

The exclusively American species are therefore 182, and 68 common to America and Europe.

American species in Arctic America-EuropeandoAmerican species in Arctic America and the StatesEuropeandoAmerican species in the StatesEuropeandoAmerican species in the StatesEuropeando	45 69 68	36 28 4	<pre>} 81 } 97 } 72</pre>
	182	68	250

You will find that the large proportion of Carices in the Northern part of America, common to it and to Europe, is in accordance with the observations of Agassiz, made in his late interesting excursion to Lake Superior. He remarks that the farther north we proceed the greater is the uniformity of the plants common to the two continents; and it is remarkable that Leconte, in his list of the Coleoptera of Lake Superior, was struck with the absence of all the groups peculiar to the American continent, the large increase of the species of genera feebly represented in the more temperate regions, and the existence of many genera heretofore regarded as confined to the southern parts of Europe and Asia.

Yours sincerely,

F. BOOTT.

NTS.

ne Carices luaintance he highest s:—

6th, 1850.

m your last

Portage. ortage. y, and of the

Huron. Huron, Fort

Huron, Fort

ron, Winipeg,

eg.

The 97 species found in Arctic America and in the States, are -

C. anceps Muhlg. arida Tor. aristata Brown. aurea Nutt. angustata Boott. arctata Boott. adusta Boott. aperta Boott. aquatilis Wahl. atrata L. blanda Dewey. bromoides Schk. Backii Boott. bullata Schk. Buxbaumii Wahl. cephalophora Muhl. conoidea Schk. cristata Schwz. crinita Lam. commutata Gay. capillaris L. canescens L. chordorrhiza Ehrh. capitata L. debilis Michx. Deweyana Schwz. digitalis Willd. eburnea Boott. Ehrhartiana Hoppe. festucacea Schk. flexilis Rudge. filiformis L. flava L.

C. fulva Good. grisca Wahlg. gracillima Schwz. granularis Muhlg. gynocrates Worm. hystericina Muhlg. intumescens Rudge. irrigua Willd. lupulina Muhlg. lagopodioides Schk. Liddoni Boott. longirostris Tor. lanuginosa Michr. lacustris Willd. lenticularis Michx. limosa L. livida Willd. Muhlenbergii Schk. miliacen Muhlg. monile Tuckn. muricata L. NovæAngliæ Schwz oligosperma Michx. Ederi Ehrh. polytrichoides Muhl pubescens Muhlg. PennsylvanicaLam. pedunculata Muhlg. plantaginea Lam. pseudocyperus L. pallescens L. pauciflora Light. rosea Schk.

C. Richardsoni Brown. retrorsa Schwz. rostrata Michx. retroflexa Muhlg. rigida Good. subulata Michx. squarrosa L. striata Michx. stipata Muhlg. scoparia Schk. straminea Schk. scabrata Schwz. Schweinitzii Dewey siccata Dewey. scirpoidea Michx. stellulata Good. triceps Michx. tentaculata Muhlg. trisperma Dewey. teretiuscula Good. tenuiflora Wahl. tenella Schk. umbellata Schk. utriculata Boott. varia Muhlg. verticillata Boott. vulpinoidea Michx. vesicaria L. vulgaris Fries. vitilis Fries. Willdenowii Schk.

Of the 97 in Arctic America and in the States, 28 are European.

In England, C. Buxbaumii Wg. canescens L. filiformis L. flava L. fulva Good. irrigua Willd. limosa L. muricata L. Œderi Ehrh. pseudocyperus L. pallescens L. stellulata Good. teretiuscula Good. vesicaria L. vulgaris Fries.

6

In Scotland. C. capillaris L. aquatilis Wahl. atrata L. pauciflora Light. rigida Good. In North of Europe. C. chordorrhiza Ehrh. capitata L. gynocrates Worm. tenuiflora Wahlg. livida Willd. tenella Schk. vitilis Fries. Ehrhartiana Hoppe.

TABLE OF DISTRIBUTION OF CARICES. 347

Of these 28 species of Europe, 12 are Alpine, or found in high northern latitudes.

C. Ehrhartiana is found in Germany, and is probably a form of C. teretiuseula Good.

C. fulva was originally established upon a Newfoundland specimen, and has only been found once near Boston, U.S.A.

The 72 found in the United States are -

C. alopecoidea Tuckn. | C. folliculata L. æstivalis Curtis. alveata Boott. Boottiann Benth. Barrattii Tor. Baltzellii Chapman. Buckleyi Dewey. crus-corvi Shutt. Careyana Dewey. Cherokeensis Schwz. Crawei Dewey. Cooleyi Dewey. Caroliniana Buckley comosa Boott. decomposita Dewey. Davisii Schwz. dasyearpa Muhlg. exilis Dewey. Elliottii Tor. Floridana T'or. Fraseri Sims. fænea Willd. formosa Dewey. flaccosperma Dewey

glaucescens Eh. Grayii Carey. gigantea Rudge. hyalina Boott. Halsevana Dewey. Hite Rockiana Dew. imbil.cats doott. juncea W %d. Kaarkernin Dewey. i orum Willd. 10pt Hormis Sartw. iavighta Smith. virabilis Dewey. Mitchelliana Curtis microdonta Tor. Meadii Dewey. mirata Dewey. oligocarpa Schk. oxylepis Tor. præcox Jacq. panicea L. platyphylla Carey. planostachys Kunz.

C. polymorpha Muhlg. refracta Schk. retrocurva Dewey. sterilis Willd. sparganioides Muhl. Sartwellii Dewey. setacea Dewey. Shortii Tor. Steudelii Kunth. styloflexa Buckley. stenolepis Tor. Sullivantii Boott. sychnocephala Car. strictior Dewey. tenax Chapman. tetanica Schk. torta Boott. turgescens Tor. Tuckermani Boott. vestita Willd. venusta Dewey virescens Muhl. vulpina L. Woodii Dewey.

Of these 72 species, 4 only are common to Europe (England).

C. præcox Jacq. (introduced), found only in Salem, Massachusetts. lævigata Smith (introduced), found once near Boston, Massachusetts.

panicea L.

vulpina L. Doubtful, probably a form of C. stipata. (Ohio, Illinois.)

I can offer you little that is satisfactory to myself as to the geographical range of the 97 species that are common to Arctic America and the States, for want of precise data as to the Carices of the Southern and Western States.

A.-I find, from such data as I have, that from lat. 30° to 35°, that is, from N. Orleans through the Carolinas, there are 33 species extending into Arctic America, one of which, C. Novæ Angliæ Schwz., ranging from N. Orleans to the Arctic Sea, maintains an equally vigorous development through 40° of latitude.

rs.

s, are --

soni Brown. Schwz. Michx. a Muhlg. food. Michx. a L. Michx. Muhlg. Schk. ea Schk. Schwz. nitzii Dewey Dewey. ea Michx. ta Good. Michx. lata Muhlg. na Dewey. scula Good. ora Wahl. Schk. ata Schk. ata Boott. Muhlg. llata Boott. oidea Michx. ia L. is Fries. Fries. nowii Schk.

European.

of Europe. rrhiza Ehrh. h L. ates Worm. ora Wahlg. Willd. Schk. Fries. tiana Hoppe.

CFre	om lat. 42° to 45°, Massa	achusetts to Wisconsin, there are 4
S	pecies extending northwa	ras.
Of th	ie 72 species found in t	he States, 27 are southern species
ranging	from Florida to Kentuck	y.
Nowform	adland	l C remoto I
Labrado	r,	 C. recta B., nigra All., ustulat. We.
Greenla	nd,	 C. duriuscula Mr., hæmatolepi Dr., reducta Dr., rufina Dr., holostoma Dr., hyperbore Dr., microglochin Wg., pe data Wg., microstachya Ehrh
Canada,		1. C. miliaris Mx.
Rocky N	Iountains	 C. petasata Dy., petricosa Dy., filiforia Nutt., Geyeri B., Ly. oni B., Jamesii T., dioica L. Pyrenaica Wg.
Rocky N	Iountains and Altai,	1. C. Franklinii B. (C. macrogyna Turczon).
North-V	Vest coast,	 C. anthoxantha Pr., antheri coides Pr., Hoodii B., ampli folia B., Gmelini H., circinnata Mr., leiocarpa Mr., marcida B., micropoda Mr., macro cephala W., Mertensii Pres. nigella B., Sitchensis Pres. Tolmiei B., elongata L., le porina L., stricta G., physocarpa Presl. C. oveta Budga
Greenlar	d to Laka Superior	1. C. bigolog All
meeniai "	and Newfoundland to Arctic Sea.	1. C. glareosa Wg.
"	to Cumberland House,	1. C. subspathacea Worm.
,,	to Arctic Sea,	3. C. stans Dr., vahlii S., ursina Dy
>>	Slave Lake, and Ft. Enterprise,	1. C. rotundata Wg.
**	and Mackenzie River,	1. C. rariflora Sm.
**	Bear Lake, and Rocky Mts.,	2. C. supina Wg., vaginata Tausch
,	Bear Lake, Church R. and Sask.	1. C. ampullacea G.
	and Danulas Dar	1 C fuliningen St to Han

NTS.

there are 21

there are 43

hern species,

All., ustulata

, hæmatolepis ., rufina Dr., hyperborea in Wg., pestachya Ehrh.

petricosa Dy., eyeii B., Ly-T., dioica L.,

(C. macrogyna

Pr., antheriodii B., ampli-H., circinnata Mr., marcida Mr., macroertensii Pres., chensis Pres., ongata L., leta G., physo-

Vorm. S., ursina Dy.

cinata Tausch.

Hop.

	Mountains, and North-West coast,		
99 1	to Rocky Mountains,	7.	C. festiva Dy., incurva Light. lagopina Wg., nardina Fr., obtusata Lil., rupestris All., saxatilis L.
" t	o North-West coast (net Rocky Mts.),	3.	C. salina Wg., cryptocarpa Mr., stylosa Mr.
Arctic Sea,		1.	C. marina Dy.
Hudson's B	ay to Methy Portage,	1.	C. Houghtonii Tor.
"	to Cumberland H.,	2.	C. heleonastes .Eh., maritima Mull.
Cumberland River, an	House, Mackenzie d Rocky Mts.,	1.	C. concinna Br.
Carlton Ho	use,	2.	C. Torreyi Tuck., Hookeriana Dy.
,,	to North-West coast,	1.	C. Parryana Dy.
Methy Port	age,	1.	C. Racana B.
Wooded Co	untry,	2.	C. affinis Br., podocarpa Br.
Rocky Mo West coa	ountains to North- st,	5.	C. Douglassii B., Rossii B., nigricans Mr., macrochæta

Greenland, Arctic Sea, Rocky 1. C. compacta Br.

nigricans Mr., macrochœta Mr., stenophylla Wg.

Of the above 81 species, 36 are European !

	England !	
3.	remota L.	
	dioica L.	
	elongata L.	
	leporina L.	
	stricta G.	
	ampullacea G.	

Scotland ! C. ustulata Wg. vahlii Schk. rariflora Sm. vaginata Tausch. incurva Light. lagopina Wg. rupestris All. saxatilis L.

North of Europe ! C. bicolor All. subspathacea Worm. fuliginosa S. & Hop. nigra All. microglochin Wg. microstachya Ehrh. rufina Dr. Pyrenaica Wg. (Pyrenees!) obtusata *Ĺil*. supina Wahl. salina Wahl. maritima Muller. stenophylla Wahlg. heleonastes Ehrh. glareosa Wahl. festiva Dy. nardina Fries. pedata Wahlg. rotundata Wahlg. hyperborea Dr. (Lapland). holostoma Dr. (Lapland). cryptocarpa Mr. (Iceland).

349 TABLE OF DISTRIBUTION OF CARICES.

Of these 36 European species found in Arctic America, 30 are Alpine or belonging to high northern latitudes.

C. Pyrenaica Wahlg. is confined in Europe to the Pyrenees!

C. remota L., common in England, has been found only in Newfoundland. Royle has found it also on the Himalayas and at Ku nawur in the East Indies!

C. festiva Dewey, found in Norway, Finmark, and Lapland, extends in America from Greenland to Unalaschka, and along the **Rocky Mountains to the Straits of Magellan!**

The 45 American species in Arctic America are :-

C. duriuscula Meyer. circinnata Meyer. leiocarpa Meyer. micropoda Meyer. stylosa Meyer. nigricans Meyer. macrochæta Meyer. anthoxantha Presl. anthericoides Presl. physocarpa Presl. Mertensii Prescott. Sitchensis Prescott. miliaris Michaux. filifolia Nuttall. Jamesii Terrey.

,,

C. Houghtonii Terrey. C. podocarpa Brown. reducta Drejer. hæmatolepis Drejer. stans Drejer. petasata Dewey. petricosa Dewey. Hookeriana Dewey. marina Dewey. ursina Dewey. Parryana Dewey. Torreyi Tuckerman. ovata Rudge. Gmelini Hooker. macrocephala Willd. affinis Brown.

compacta Brown. concinna Brown. Geyeri Boott. Lyoni Boott. Hoodii Boott. Rossii Boott. Tolmiei Boott. marcida Boott. nigella Boott. Douglassii Boott. Franklinii Boott. Raeana Boott. amplifolia Boott. recta Boott.

A .- New Orleans, Cumberland House, Rocky Mountains, C. umbellata, debilis.

> Greenland, Rocky Mountains, Arctic Sea, C. Novæ Angliæ.

Texas to Canada, C. retroflexa, grisea, blanda, triceps.

Hudson's Bay, C. Muhlenbergii. ,,

Carlton House, North-West coast, C. anceps. ,,

Georgia to Canada, C. lupulina, commutata, squarrosa, tentaculata, striata, hystericina, miliacea.

South Carolina to Canada, C. cephalophora, varia, granularis, vulpinoidea.

**	,,	,, a)	nd Rock	y Mountains, C. bromoides.
,,	,,	Cumberla	nd Hous	se, C. crinita, intumescens.
"	"	"	"	Rocky Mountains,
				North-West coast,
				C. stellulata.
"	"	"	"	North-West coast, C.
				lagonodioides.
"	"	Hudson's trichoid	Bay, I	Norway House, C. poly-

NTS.

erica, 30 are

ences only in Newand at Ku

nd Lapland, nd along the

arpa Brown. acta Brown. una Brown. ri Boott. i Boott. ii Boott. i Boatt. ici Booti. ida Boott. a Boott. glassii Boott. klinii Boott. na Boott. ifolia Boott. Boott.

ains, C. umbel-

retic Sea, C.

ceps.

nceps. arrosa, tentacu-

ria, granularis,

s, C. bromoides. , intumescens. Mountains, -West coast, lulata. Vest coast, C. scoparia, dioides. use, C. poly-

TABLE OF DISTRIBUTION OF CARICES.

South Carolina to Lake Winnipeg, C. lacustris, North-West coast, C. rosea. 99

North Carolina Mountains of, Observatory Inlet, Cumberland House, North-West coast, C. Buxbaumii. 11

to Cunada, C. conoidea. 11 Cumberland House, Rocky Mountains. .,, C. Pennsylvanica.

B. - Kentucky to Canada, C. pubescens, digitalis.

11

,,

Mackenzie River, Rocky Mountains, C. eburnea. Illinois, Cumberland House, Rocky Mountains, North-West coast, C. Richardsoni.

New Jersey to Canada, C. Schweinitzii.

North-West coast, C. aperta. "

to Hudson's Bay, Arctic Sea, Recky Mountains, 99 North-West coast, C. livida.

Ohio to Cumberland House, C. arida.

Carlton House and Rocky Mountains, C. Backii. •• Pennsylvania to Canada, C. subulata, scabrata, bullatu.

> Cumberland House, C. gracillima, cristata, plantaginea.

Rocky Mountains, C. pe-,, dunculata, utriculata.

Greenland, North-West coast, C. vesicaria. "

North-West const, C. angustata.

Rhode Island to Bear Lake, C. monile.

Greenland, Cumberland House, Rocky Moun-• • tains, North-West coast, C. adusta.

C.-Massachusetts and Newfoundland, C. fulva.

North-West coast, C. muricata, verticillata.

Michigan to Canada, C. festucacea.

Cumberland House, C. Ehrhartiana. 17

- Carlton House, Rocky Mountains, C. teretiuscula, ,, trisperma.
- Canada, North-West coast, C. straminea. ,,

and North-West coast, C. Liddoni.

New York to Canada, C. arctata.

Rocky Mountains, C. Willdenowii.

Cumberland House, C. pallescens, pseudocyperus, •• tenuiflora, vitilis, flava, aristata, filiformis, irrigua. Rocky Mountains, C. sic-11 ,, •• cata, longirostris.

New	York	to Cumberla	nd House	, North-W	est coast, C. re-
				Greenland	i. Hudson's Bay.
	"	,,	"	C. Œa	leri.
	,,	Mackenzie	River, R	locky Mou	ntains, C. aquatilis.
	"	"	N	Vorth-West ginosa.	coast, C. lanu-
	"	Bear Lake C. limos	, Rocky I z.	Mountains,	North-West coast,
	"	Hudson's I North-V	Bay, Carl Vest coast	ton House, t, <i>C. aurca</i> .	Rocky Mountains,
	,,	Newfound	and, C. f	lexilis.	
	"	>>	Gree M	enland, A Iountains, . vulgaris,	rctic Sea, Rocky North-West coast, canescens.
	,,	33	Roc	ky Mount	ains, North-West
			co	ast, C. par	uciflora.
	"	Greenland,	C. gynoe	crates.	
New	York	Mountains of,	to Mack	enzie River	, C. lenticularis.
	"	**	Greenl	and, Arc	tic Sea, Rocky
				Moun	tains, C. scirpoidea.
	"	**	"	Labra	dor, Arctic Sea,
		White Mts.	of New 1	C. 7 Tampshire	Bear Lake, C. oli-
	"		01 1101 1	ramponno,	gosperma.
	,,	,,	"	"	North-West coast,
					C. rostrata.
New	Hamp	oshire, White	Mts. of,	Canada, Mounts	Greenland, Rocky
				Greenland	l. Hudson's Bay.
"		- "	"		Rocky Moun-
					tains, C. capi-
					tata.
"		,,	,,	"	Bear Lake,
					tains, C. capil- laris.
Wisco	nsin t	o Canada, C. a	enella.		
,		Cumberland	House, E	Iudson's Ba	y, C. chordorrhiza.
,	,	,,	"]	Rocky Mo	untains, C. Dew-

eyana.

The geographical range, as far as I know it, of the 72 species found in the United States is as follows: --

ITS.

ast, C. re-

dson's Bay,

C. aquatilis. t, C. lanu-

-West coast,

y Mountains,

Sea, Rocky -West coast, cens. North-West

a.

lenticularis. Sea, Rocky , C. scirpoidea. Arctic Sea, r. r. Lake, C. oliosperma. th-West coast, f. rostrata. mland, Rocky

C. atrata. Iudson's Bay, locky Moun-

nins, C. capi-

ar Lake, Rocky Moun-

ains, C. capillaris.

chordorrhiza. ns, C. Dew-

he 72 species

TABLE OF DISTRIBUTION OF CARICES.

Florida, C. tenax, Baltzellii, oxylepis.

- " to New Orleans, C. gigantea, Floridana.
- " Georgia, C. dasycarpa, turgescens.
- " Carolinas, C. glaucescens, venusta, Elliottii.
- " New England, C. folliculata, polymorpha.

Texas, C. alveata, hyalina, imbricata, microdonta, planostuchys.

- " to Alabama, C. Cherokeensis.
- " Kentucky, C. stenolepis.
- " New Jersey, C. flaccosperma.
- " Rhode Island, C. fænea.

New Orleans, C. Boottiana.

" to Wisconsin, C. Meadii,

Carolina, South, C. Buckleyi, Caroliniana, Mitchelliana, Fraseri, juncea, styloplexa, lucorum.

- " to Virginia, C. æstivalis.
- " Massachusetts, C. comosa.
- Carolina, North, to Ohio, C. Sullivantii.

Virginia to Kentucky, C. Shortii.

Kentucky to New York, C. oligocarpa, Hitchcockiana.

" Connecticut, C. Davisii, virescens.

Ohio, C. tetanica, crus-corvi.

- " to Pennsylvania, C. strictior.
- " New Jersey, C. Tuckermani.
- " New York, C. Careyana.
- " Illinois, C. vulpina.
- Pennsylvania, C. refracta.
 - " to Cherokee, C. sterilis.
 - Connecticut, C. torta.
- New Jersey, C. Barrattii, Kneiskernii.

" to Rhode Island, C. Halseyana, platyphylla.

" Connecticut, C. vestita.

New York, C. alopecoidea, formosa, lupuliformis, mirata, sychnocephala, Woodii.

- " Michigan, C. crawei, Sartwellii, decomposita, Steudelii.
- " Rhode Island, C. retrocurva.
- " Massachusetts, C. exilis.
- " Connecticut, C. Grayii.

Massachusetts, C. setacea, præcox, panicea, lævigata.

to Michigan, C. mirabilis.

Michigan, C. Cooleyi.

Wisconsin to New England, C. sparganioides.

VOL. II.

ΛΛ

354 INSECTS OF ARCTIC NORTH AMERICA.

For the following notice and list of insects collected on the Expedition, I am indebted to Adam White, Esq., F.L.S., &c., of the British Museum. With respect to the extent of the collection, it is to be observed that no time was devoted to the capture of insects. Such as presented themselves at convenient times were taken, but none were sought for; and the numbers of the list are not, therefore, to be considered as a criterion of the richness of the country in that division of the animal kingdom.

NOTE ON HYMENOPTERA IN ARCTIC NORTH AMERICA.

"Otho Fabricius first, perhaps, recorded any of the Hymenoptera of Arctic North America. Doubtless Baffin, Frobisher, and other manly navigators recognised humble bees and other bees during their summer voyages, and may have, in print or in manuscript, with sailor-like earnestness, made mention of every such occurrence in their journals. It is delightful to read the notices of flowers and verdure in their accounts of the hurried spring, summer, and autumn two months of a Greenland year, of five-sixths winter. Where flowers and verdure abound, even for six weeks or a shorter time, there insects must be found;—there insects of the order Hymenoptera, the order to which this notice is limited, must occur. Flowers and Hymenoptera must be together.

"O. Fabricius records two species of Hymenoptera as being brought by him from Greenland. His book, so admirable a model of a local fauna as to be even now one of the standards of excellence, was published in 1780. The next considerable accession to our acquaintance with the Hymenoptera of British America was made by Redman, who collected in Nova Scotia many fine species now in the British Museum. Some of these, such as Pelecinus, Sirices, Ichneumonidæ, &c., were very prominent species, and are now being worked out in the vast collections of the National Museum.

"Sir John Richardson and his brave comrades collected many

NOTE ON THE HYMENOPTERA, ETC. 355

lected on ite, Esq., respect to ed that no ach as preaken, but ist are not, richness of om.

MERICA.

the Hymeni, Frobisher, is and other is in print or is mention of the of a Greenand verdure there insects enoptera, the Flowers and

tera as being admirable a standards of considerable ra of British Nova Scotia ome of these, re very prothe vast col-

llected many

species, which were lost during their disastrous journey. They still, however, brought many insects to England, and in the 'Fauna Boreali-Americana' these insects are described by the venerable Kirby. The species of Hymenoptera are very few; there are only thirty-two altogether; the circumstances attending the journey not admitting of their collection and preservation.

"An eminent man, reasoning on such data as he had, has recorded his belief that it will be found that *Hymenoptera* do not abound in British North America; now it may be remarked in making generalisations on the distribution of animals, especially those of the lower orders, 'that, before generalising on a collection from any place not often visited or not often explored, attention be paid to the taste or tastes, or, in other words, to the bias or direction of the eye, hand, and mind of the person or prisons who collect, supposing such reasoning is recorded as on authentic data.'

"Mr. George Barnston, to whose researches Sir John Richardson directed public attention in the 'Edinburgh New Philosophical Journal' for April, 1841, has published a very admirable summary of the Progress of the Seasons as affecting Animals and Vegetables at Martin's Falls, Albany River, James's Bay, about lat. 51° 30' N., and in long. 86° 20' W. In this fresh and refreshing journal, there are more than indications that Hymenoptera, Diptera, and Neuroptera abound. In a year or two afterwards Mr. Barnston came to London and presented his collection to the British Museum.

"As one instance of his excellence as a collector, I may mention that Mr. Walker, who named and described the species of *Diptera* in the Cabinet of the British Museum, has alluded to or has described nearly 250 species of his dipterous insects from the single station mentioned above; there being only 14 species of these insects recorded in the 'Fauna Boreali-Americana' of the Rev. Wm. Kirby. Mr. Barnston's researches among the Neuroptera also were considerable and very valuable. One insect brought by him, the Pteronarcys regalis (although previously found in Canada), afforded Mr. Newport a fit sub-

A A 2

356 INSECTS OF ARCTIC NORTH AMERICA.

ject for his genius as an accurate anatomist and recorder of facts and reasonings on the insect economy. This gentleman discovered persistent *branchiæ* in the *imago* or perfect state of the *Pteronarcys*, and has recorded his discovery and quoted some observations of Mr. Barnston's in a paper read at the Linnæan Society. As Mr. Gray's Catalogues of the collections in the British Museum, (mines of information to the reasoner and writer on geographical distribution,) are published, it will be seen how valuable are Mr. Barnston's and Sir John Richardson's collections to our acquaintance with the articulated animals of British North America, especially in its more northerly parts.

"I have mentioned that Kirby describes or alludes to only thirty-two species of Hymenoptera in his 'Insects of North America;' while Mr. Barnston in one spot found 192 distinct species, exclusive of Chalcididæ. "I subjoin a comparative list of the families of Hymenoptera, the comparison being made with the British species existing in the Museum collection at the time of this record. Mr. B. and myself worked out the Tenthredinidæ; my friend and coadjutor Mr. Frederick Smith, an able hymenopterist, determined the other species; so the list may be deemed as correct as the circumstances will admit.

"It must be borne in mind that our British collection of Hymenopfera has been accumulating for at least thirty years, was a favourite part of Dr. Leach's collection, and has been made over a wide and variegated country; while Mr. Barnston's was formed in three months, on one spot and under almost unheard-of disadvantages, counterbalanced, however, by an enthusiasm not easily deterred by difficulties.

		5	Br in 1	itish Colle British Mu	Collected at Martin's Falls.				
Cimbicidæ	-	-	-	10	-	-	-	4	
Tenthredinidæ		-	-	157	-	-	-	76	
Siricidæ, &c.	-	-	-	7	-	-	-	2	
Ichneumonidæ		-	-	200	-	-	-	47	
Chalcididæ	-	-	-	2	-	-	-	2	
Chrysididæ	-	-	-	22	-	-	-	1	
Formicida	-	-	-	11	-	-	-	7	

COLEOPTERA.

			British Collection in British Museum.				Collected at Martin's Fai			
Mutillidæ	•	-	-	5		-	-	0		
Sapygidæ	•	-	-	2	-	•	-	0		
Pompilidæ, &c.		-	-	38	-	-	-	2		
Crabronidæ	-	•	-	57	-	-	-	16 .		
Vespidæ	-	-	-	17	-	-	-	4		
Apidæ -	•	-		170	-	-	-	33		

"A striking proof that the time has not yet come to reason correctly on the distribution of Hymenopterous insects, — at least in British North America."

LIST OF INSECTS

TAKEN BY SIR JOHN RECEIPTING AND JOHN RAE, ESC. IN ARCTIC NORTH AMERICA, DRAWF UP BY ALAM WITTE, ESQ., F.L.S., ETC.

Consorticka.

Cieindela	longilabris.	, Say.	(C. zibilabris,	Kirby).	Shores of Ar tic	
Sea, lat	. 70° N.; a	and at	Fort Supson,	lat. 62°	12	

Cicindela micollis, Say. Borders of Mackenzie and Blave Rivers, lat. 59°-62° N.

Dromius nigrinus, Eschsch. Great Bear Lake, lat. 66°--67° N.

- Carabus ? n. s. (C. gladiator, 7 arnston MS.). Borders of Mackenzie and Slave Rivers, lat. 5 -65° N.
- Carabus Chamissonis, Eschsch. Borders of Mackenzie and Slave Rivers; and Cape Krusentern, lat. 58°-68° N.
- Carabus —? n. s. (C. Hudsonicus?) Borders of Mackenzie and Slave Rivers, lat. 58°-65° N.
- Calosoma calidum, Auct. B rders of Mackenzie and Slave Rivers, lat. 58°-65° N.

Loricera pilicornis, Auct. Great Bear Lake, lat. 66°-67° N.

- Elaphrus intermedius, Kirby. Great Bear Lake, lat. 66°-67° N.
- Notiophilus sibiricus, Motchoulsky. Great Bear Lake, lat. 66°--67° N.
- Dicelus ? n. s. (D. sculptilis?) Borders of the Mackenzic and Slave Rivers, lat. 58°-65° N.
- Agonum melanarium, Dej. Great Bear Lake, and district to the south of Lake Winipeg, lat. 49°-68° N.

AA 3

1.

recorder of s gentleman erfect state and quoted read at the e collections the reasoner published, it ad Sir John the articuy in its more

udes to only ets of North 192 distinct comparative n being made n collection at orked out the derick Smith, eccies; so the will admit.

ection of Hyrty years, was as been made Ir. Barnston's under almost rever, by an

Collected at artin's Falls. 4 76 2 47 ? 1

7

358 INSECTS OF ARCTIC NORTH AMERICA.

Argutor brevicornis, Kirby. Borders of Mackenzie and Slave Rivers. Omaseus orinomum, Leach. District south of Lake Winipeg, lat. 50°-54° N.

Platysma vitrea, Eschsch. Borders of Mackenzie and Slave Rivers.

Pæcilus lucublandus, Say. Borders of Mackenzie and Slave Rivers.

Harpalus — ? n. s. (near H. obtusus). Borders of Mackenzie and Slave Rivers.

Stenolophus — ? n. s. Great Bear Lake, lat. 66°-67° N.

Amara —? sp. (near A. trivialis). South of Lake Winipeg, lat. 49° N. Amara — ? sp. Great Bear Lake.

Bembidium conicolle, Motchoulsky (B. impressum, Kirby). Great Bear Lake, and north of Lake Winipeg, lat. 49°-67° N.

Acupalpus --- ? n. s. Great Bear Lake, lat. 66°--67° N.

Peryphus - ? sp. Great Bear Lake.

Platytrachelus -? n. s. Great Bear Lake.

Notaphus nigripes, Kirby. Great Bear Lake.

Notaphus variegatus, Kirby. South of Lake Winipeg, lat. 49° N.

Dytiscus Harrisii, Kirby. South of Lake Winipeg.

Agabus — ? n. s. (near A. arcticus). Cape Krusenstern, lat. 68° N. Colymbetes — ? sp. Borders of Mackenzie and Slave Rivers.

Hydrophilus picipes, Auct. South of Lake Winipeg, lat. 49° N.

Heterocerus -? n. s. (near H. fossor). Great Bear Lake.

Staphylinus villosus, Grav. South of Lake Winipeg, lat. 49° N.
Quedius, n.s. (near Q. molochinus). Great Bear Lake, lat. 66°-67° N.
Omalium, n. s. (near O. rivulare). Shore of Arctic Sea, near mouth of Mackenzie River, lat. 70° N.

Anthophagus, sp. Borders of Mackenzie and Slave Rivers.

Silpha Lapponica, Auct. Fort Simpson; Borders of Mackenzie and Slave Rivers.

Silpha opaca, Auci. Borders of Mackenzie and Slave Rivers.

Silpha, n. s. (near S. Baikalica), *Motchoulsky*. Borders of Mackenzie and Slave Rivers.

Ptinus fur, Auct. South of Lake Winipeg. Throughout Rupert's Land.

Byrrhus —? n. s. Borders of Mackenzie and Slave Rivers, lat. 58° --65° N.

Byrrhus -? n. s. South of Lake Winipeg.

Rhisotrogus fervens, Gyll. South of Lake Winipeg, lat. 49° N.

Platycerus piccus, Web. Fort Simpson, on the Mackenzie River, lat. 62° N.

Cyphon fuscipes, Kirby. Great Bear Lake.

Elater æripennis, Kirby. Shore of Arctic Sca, near Mackenzie River, lat. 70° N.

Elater ancus ? Borders of Mackenzie and Slave Rivers.

۸.

Slave Rivers. Winipeg, lat.

Slave Rivers. Slave Rivers. lackenzie and

N. 900 N. 49° N.

ïrby). Great '° N. N.

, lat. 49° N.

ern, lat. 68° N. Rivers. lat. 49° N. Lake. , lat. 49° N. at. 66°—67° N. Sea, near mouth

livers. Mackenzie and

Rivers. rs of Mackenzie

ghout Rupert's

Rivers, lat. 58°

at. 49° N. Ickenzie River,

ear Mackenzie

ers.

COLEOPTERA.

Elater,	n. s.	(near	Е.	melancholicus).	Borders	of	Mackenzie	and
Slave	Rive	ers.						

- Elater, n. s. (near E. sanguineus). Borders of Mackenzie and Slave Rivers.
- Ludius, n. s. (near L. sibiricus). Great Bear Lake, lat. 65°-67° N. Ampedus, n. s. Great Bear Lake.
- Buprestis tenebrica, Kirby. Borders of Mackenzie and Slave Rivers. Chrysobothris, n. s. Fort Simpson, on the Mackenzie River, lat. 62° N.
- Trachypteris Drummondi, Kirby, var. Fort Simpson, on Mackenzie River, lat. 62° N.
 - Trachypteris decolorata (Bupr. appendiculata, Kirby). Fort Simpson, on Mackenzie River, lat. 62° N.
 - Ellychnia corrusca, Auct. South of Lake Winipeg.
 - Ragonycha cembricola, Eschsch. Great Bear Lake, lat. 65°-67° N.
 - Thanasimus abdominalis, Kirby. Great Bear Lake.
- Hydnocera, n. s. Great Bear Lake.
- Blapstinus æneus, Deg. South of Lake Winipeg.
- Upis ceramboides, Auct. Fort Simpson, and Borders of Mackenzie and Slave Rivers.
- Anthicus -? n. s. Great Bear Lake.
- Formicoma -? n. s. Great Bear Lake.
- Stenotrachelus Roulieri, *Motch.* var. Shores of Arctic Sea, near Mackenzie River, lat. 70° N.
- Serropalpus ? sp. Fort Simpson, on Mackenzie River.
- Hylobius -? sp. Borders of Mackenzie and Slave Rivers.
- Alophus ? sp. South of Lake Winipeg.
- Alophus ? sp. Cape Krusenstern, lat. 68° N.
- Erirhinus, sp. (near E. tremulæ). South of Lake Winipeg.
- Tomicus -? sp. South of Lake Winipeg.
- Asemum striatum, Auct. Fort Simpson on the Mackenzie River.
- Callidium bifoveolatum. Cape Krusenstern and Arctic Coast, between 671° and 68°.
- Callidium Proteus, *Kirby*; and C. simile, *Kirby*, var. Arctic Coast, between 67¹/₂° and 68°; Fort Simpson on the Mackenzie River, lat. 62° N.
- Clytus undulatus, Say. Shore of Arctic Sea; Mouth of Mackenzie River.
- Clytus —? sp. Shore of Arctic Sea; Mouth of Mackenzie River, lat. 70° N.
- Acanthocinus pusillus, Kirby. Great Bear Lake, lat. 66°-67° N.
- Monochamus resutor, Kirby. Fort Simpson, on the Mackenzie River.
- Monochamus confusor, Kirby. Borders of Mackenzie and Slave Rivers.

A A 4

360 INSECTS OF ARCTIC NORTH AMERICA.

Acmæops Proteus (Kirby), Leconte; Leptura strigilata, var.? Fort Simpson, on the Mackenzie River.

Acmæops strigilata (Fabr.), Lec. Shore of Arctic Sea (Mouth of Mackenzie).

Pachyta liturata, Kirby. Fort Simpson, on the Mackenzie.

Pachyta, n. s. Fort Simpson, on the Mackenzie.

Rhagium lineatum, Auct. Fort Simpson, on the Mackenzie.

Syneta carinata, Eschsch. About Great Bear Lake.

Galleruca marginella, Kirby. Borders of Mackenzie and Slave Rivers; and about Great Bear Lake.

Chrysomela multipunctata, Say. Borders of Mackenzie and Slave Rivers.

Phædon Adonidis, Pall. Shore of Arctic Sea (Mouth of Mackenzie); Fort Simpson.

Adoxus vitis, Fabr. District about Great Bear Lake, and Borders of Mackenzie and Slave Rivers.

Coccinella 13-punctata, Auct. Great Bear Lake, lat. 65°-67° N.

Coccinella 5-notata, Kirby. Shore of the Arctic Sca; Mouth of Mackenzie, lat. 70° N.

Coccinella ocellata, Auct. Borders of Mackenzie and Siave Rivers.

ORTHOPTERA.

Locusta tuberculata, *Pal de Beauv.* ? Borders of Mackenzie and Slave Rivers; Fort Simpson.

Locusta, four species. Borders of Mackenzie and Slave Rivers.

Acrydium granulatum, Kirby. Borders of Mackenzie and Slave Rivers; Fort Simpson.

NEUROPTERA.

Æschna borealis, Zetterst. ? Borders of Mackenzie and Slave Rivers. Libellula — ? sp. Fort Simpson on Mackenzie.

Libellula scotica, Donov.? Between Lake Winipeg and Lake Superior.

Agrion cyathigerum, Charp. var. Borders of Mackenzie and Slave Rivers.

Ephemera viridescens — ? n. s., Barnston, Between Lake Winipeg and Lake Superior, lat. 47°-52°.

Ephemera -? n. s. South of Lake Winipeg, lat. 49° N.

Pteronarcys regalis, Newman. Borders of Mackenzie and Slave Rivers.

Pteronarcys Proteus, Newman? Shores of Arctic Sea (Mouth of Mackenzie River), lat. 70°.

NEUROPTERA, HYMENOPTERA.

361

ar.? Fort

(Mouth of

ie.

ie.

and Slave

and Slave

Mackenzie);

and Borders

-67° N. ; Mouth of

ave Rivers.

ackenzie and Rivers. e and Slave

Slave Rivers.

g and Lake

ie and Slave

ake Winipeg

and Slave

(Mouth of

Perla — ? (sp. near P. abnormalis, Newman). Borders of Mackenzie and Slave Rivers.

Perla — ? (sp. near P. sonans, *Barnston*). Borders of Mackenzie and Slave Rivers.

Semblis - ? n. s. Borders of Mackenzie and Slave Rivers.

Phryganea striata, n. s., Barnston. Borders of Mackenzie and Slave Rivers.

Phryganea variegata, n. s., *Barnston*, and two or three other species. Borders of Mackenzie and Slave Rivers.

HYMENOPTERA.

Trichiosoma lucorum, Auct. Fort Simpson, on Mackenzie River. Tenthredo (Nematus). Great Bear Lake

Tenthredo (Nematus) - ? n. s. South of Lake Winipeg.

Tenthredo integra ? About Great Bear Lake.

Tenthredo (Dolerus). South of Lake Winipeg.

Sirex flavicornis, Fabr. Cape Krusenstern; Fort Simpson on Mackenzie, and country south of Lake Winipeg.

Ephialtes - ? sp. Fort Simpson, on Mackenzie River.

Aspizonus - ? sp. Arctic Sea (Mouth of Mackenzie River).

Ichneumon - ? sp. Fort Simpson, on the Mackenzie River.

Ichneumon — ? sp. Cape Krusenstern, and Fort Simpson on the Mackenzie River.

Cryptus -? sp. South of Lake Winipeg.

Chrysis - ? sp. Cape Krusenstern.

Mutilla - ? sp. About Great Bear Lake.

- Formica herculeana. About Great Bear Lake; Borders of Mackenzie and Slave Rivers; Fort Simpson.
- Formica sanguinea. South of Lake Winipeg, and Fort Simpson.

Pompilus - ? n. s. Borders of Mackenzie and Slave Rivers.

Odynerus — ? n. s. Shore of Arctic Sea (Mouth of Mackenzie River).

Vespa maculata, var. Borders of Mackenzie.

Vespa vulgaris, Auct. Borders of Mackenzie and Slave Rivers and Fort Simpson.

Vespa marginata, Kirby. Cape Krusenstern.

Halictus — ? (n. s. near H. quadricinctus). South of Lake Winipeg. Halictus, three black species. South of Lake Winipeg.

- Megachile Willughbiella ? Fort Simpson, on the Mackenzie River.
- Bombus arcticus? Arctic Coast between 67¹/₂° and 68°; Borders of the Mackenzie and Slave Rivers.

Bombus (sp. near B. lapponicus). Arctic Coast between $67\frac{1}{2}^{\circ}$ and 68° . Bombus — ? sp. Shore of Arctic Sea (Mouth of Mackenzie River).

362 INSECTS OF ARCTIC NORTH AMERICA.

Bombus pratorum. Borders of Mackenzie and Slave Rivers. Bombus, n. s. (near B. lucorum). Arctic Coast between $67\frac{1}{2}^{\circ}$ and 68° Bombus, n. s. Arctic Coast.

HEMIPTERA.

Acanthosoma boreale, *Hope.* Great Bear Lake. Acanthosoma nebulosum, *Kirby.* South of Lake Winipeg. Miris — ? sp. Great Bear Lake.

Rhyparochromus, two species. South of Lake Winipeg. Salda — ? sp. (near S. riparia). Cape Krusenstern, lat. 68° N.

HOMOPTERA.

Aphrophora, sp. Great Bear Lake, and to the south of Lake Winipeg.

LEPIDOPTERA.

Papilio Turnus, L. Fort Simpson, on Mackenzie River.

Pontia casta, Kirby. Arctic Coast between 67¹/₄° and 68°.

Pontia, sp. Fort Simpson, on the Mackenzie.

Anthocharis — ? n. s. (near A. Simplonia). Arctic Coast between 67¹⁰/₂ and 68°.

Colias Palæno, L. Fort Simpson, on the Mackenzie River.

Colias Boothii, Curtis. Arctic Coast between 671° and 68°.

Colias Chione, var. ? C. Arctie Coast; Cape Krusenstern.

Argynnis Freija (*Thunb.*), var. Melitæa Tarquinius, *Curtis*. Arctic Coast between $67\frac{1}{2}^{\circ}$ and 68° .

Argynnis - ? n. s. Arctic Coast.

Vanessa Milberti, Godart. (V. furcillata, Say). Fort Simpson, on the Mackenzie River.

Vanessa Progne, Godart. (V. C. argenteum, Kirby). Fort Simpson, on the Mackenzie River; Arctic Coast between $67\frac{1}{2}^{\circ}$ and 68° .

Nymphalis Artemis, Auct. Fort Simpson on Mackenzie River, and Borders of Mackenzie and Slave Rivers.

Chionobas Bore, Boisd. ? Arctic Coast between 67¹° and 68°.

Hipparchia, n. s. ? (near H. discoidalis), Kirby. Arctic Coast between $67_{\pm}^{\circ}^{\circ}$ and 68° .

Hipparchia Rossii, Curtis. Arctic Coast between 671° and 68°.

Polyommatus Franklinii, Curtis. Arctic Coast.

Arctia Americana, Harris, var. Borders of Mackenzie and Slave Rivers.

Hadena Richardsonii, Curtis. Arctic Coast between 67¹/₂° and 68°.

Anarta -? sp. Arctic Coast between 671° and 68°.

Geometridæ, two species. Arctic Coast between $67\frac{1}{2}^{\circ}$ and 68° . Tincidæ, three species. Arctic Coast between 67^{1} and 68° . **A**.

ivers. 671° and 68°

eg.

;. . 68° N.

Lake Winipeg.

er. 18°.

Coast between

liver. 168°. Itern. *Curtis*, Arctic

impson, on the

ort Simpson, on nd 68°. zie River, and

and 68°. ctic Coast be-

and 68°.

zie and Slave

71° and 68°.

and 68°. 68°.

VOCABULARIES OF NORTH AMERICA.

DIPTERA.

Culex — ? sp. Borders of Mackenzie and Slave Rivers.
Chironomus, sp. Borders of Mackenzie and Slave Rivers.
Tipula, sp. Arctic Coast between 67¹/₂° and 68°.
Tabanus, three species. Borders of Mackenzie and Slave Rivers.
Tabanus, two species. Arctic Sea, Mouth of Mackenzie River.
Eristalis flavipes, Walker. Borders of Mackenzie and Slave River, and district to the South of Lake Winipeg, lat. 49°-65° N.
Syrphus, sp. Borders of Mackenzie and Slave Rivers.
Musca, five species. Borders of Mackenzie and Slave Rivers.
Musca — ? sp. South of Lake Winipeg.
(Estrus Tarandi ? Arctic Coast between 67¹/₂° and 68°.

No. V.

VOCABULARIES.

A. Eskimo Vocabulary.

THE Kuskuchewak column of the following vocabulary is extracted from Bäer's work.* To draw up an effective comparative table would require a thorough acquaintance with both dialects, since the names of articles of dress, and implements of art, change with the materials of which they are formed; natural objects are differently designated,

* BAER, Statische und ethnographische Nachrichten über die Russischen Besitzungen an der Nordwestkuste von Amerika. St. Petersburg, 1839, p. 259. The Kuskuchewak words being written in this work in Russian characters, with which I am unacquainted, J. F. von Bach, Esq., of the British Museum, had the kindness to furnish me with a translation. This gentleman drew up carefully columns representing the conventional English equivalents of the Russian characters for each word, and added also the French pronunciation, which want of space compels me reluctantly to omit. I have made some small alterations in words written by him according to the English pronunciation, to suit the plan of orthography which I have followed in the other vocabularies.







364 VOCABULARIES OF NORTH AMERICA.

according to the circumstances under which they are viewed; and the terms for actions are altered, as the agents, time, place, and other circumstances, vary. Unless, therefore, these facts be known and attended to by one who forms a vocabulary, there may appear to be no resemblance between the dialects of two tribes who mutually understand each other, and converse together with ease. The introduction of the syllabic characters used by the late Rev. Mr. Evans in teaching the Cree Indians would, I believe, remove the difficulties which orthography throws in the way of a European, who endeavours to reduce the native languages of North America to writing.

The column containing Eskimo spoken on the Labrador coast, is extracted from a pretty large vocabulary and grammar, which the Rev. Peter Latrobe had the kindness to procure for my use on the expedition. I have reason to believe that some errors may have crept into this vocabulary, from the similarity of the German written hto s not being always adverted to by the transcriber, and also from the uncertainty of the proper English equivalent of the German v. These are not, however, I trust numerous among the examples I have used. Where the Labrador dictionary was defective, the excellent English and Eskimo vocabulary, drawn up by Captain Washington, and published by the Admiralty for the use of the Searching Expeditions, has been referred to. The dialects spoken by the intermediate Eskimo tribes inhabiting the north shores of the continent are seldom quoted, my object having been to identify the language spoken by members of the nation occupying geographical positions the most remote from each other.

In writing out the table it was obvious to me that the Labrador dialect is in general the softer of the two.

ESKIMO VOCABULARY.

they are ad, as the ary. Untended to appear to tribes who to together characters g the Cree tics which h, who enth America

e Labrador bulary and he kindness have reason t into this n written h scriber, and ish equivaver, I trust Where the ent English Vashington, the Searchlects spoken the north ject having bers of the lost remote

ne that the f the two.

Instead of the hard tch so frequent in the Kuskuchewak tongue, the east coast tribes generally use s, and in Coronation Gulf h is substituted. The strongly aspirated sound which is heard in the Scottish word "loch" is of frequent occurrence in the Kuskuchewak column of the vocabulary, where it is denoted by kh. An Englishman in attempting this sound lets the k be heard, which he ought not to do. The difficulty of constructing a correct Eskimo vocabulary is increased by the necessity of previously mastering the exceedingly numerous inflections of the nouns, adjectives, pronouns, and verbs, which supply the place occupied by auxiliary verbs, possessive pronouns, prepositions, and adverbs in the European languages. These inflections are briefly noticed in the introduction to Capt. Washington's vocabulary, and I shall merely add here, that in the Labrador grammar, obtained for us by Mr. Latrobe, there are examples of thirty different terminations of the dual and plural numbers of nouns, which have evidently had their origin in euphuism.

Each noun has six cases in each number, distinguished by their terminations, the vocative being, however, absent in some. The cases are formed by affixes having the power of prepositions, as mut, mik, mit, me, and hut in the singular, and nut, nih, nit, ne, and gut in the plural. The nominative is also varied by affixes which perform the functions of possessive pronouns, as ga, go, ne, ait, anga, ara, &c.; as kivgah, a servant, kivganga, my servant, kivgane, his servant; nuna, land, nunaga, my land; nelegah, a master, nelegara, my master; tunnusuga, my nation, &c. Pagit, panga, or parma are affixes employed when the noun is connected with a verb signifying action or suffering. The noun, when changed by a qualifying affix, is declined in its new form, in the usual way. Besides the ordinary active nominative, each noun has also an intransitive one, which

366 VOCABULARIES OF NORTH AMERICA.

ends in b, and is differently declined. Examples are subjoined. The power of the affixes varies according as the noun is used with a transitive or intransitive verb. Nouns may also be varied by affixes expressive of augmentation, diminution, affection, ridicule, humility, or multitude. Some of these terminations are arsuk, arsuit, diminutives; and soak, sudset, sudsek, augmentatives; vak placed before any of them increases their power; and the adverbial aluk denoting "very" may be put after them, and is applicable to either good or bad. Vavak, signifying an extraordinary number, is placed before söareluit.*

Adjectives have also their declensions; and likewise comparisons made by the addition of the syllable *nek*, or by verbs. The adjective generally follows the noun, and must agree with it in case and number. If the substantive have an affix, so must the adjective. Nouns may be changed into verbs by the affix *evok* or *ovok*, and the adjective then must take the same termination.

Pronouns are declined like the nouns by affixes, which require much nicety in their due employment. Affixes supply the place of possessive pronouns.

The third person singular of the indicative is considered to be the root of the verb, and may be used as a noun with a change in the termination, "a hunter" being equivalent to "he hunts." The inflections of the verb are extremely numerous, and are expressive of affirmation, negation, interrogation, and of the various circumstances in which the agent or object can be placed with respect to time, place, mood, or possession. The infinitive, formed by the termination *nek*, is used when things are spoken of indefinitely, or when two verbs come together, and is conjugated in the

* Akkatu is employed by the Eskimo of Churchill in the same way as söareluit.

ESKIMO VOCABULARY.

es are subas the noun Nouns may ation, dimide. Some tives; and before any erbial *aluk* s applicable traordinary

nd likewise e nek, or by n, and must antive have be changed jective then

fixes, which ht. Affixes

s considered a noun with equivalent e extremely egation, inh which the time, place, y the termindefinitely, gated in the

the same way

same way as the other moods and tenses, there being a past and future infinitive. Generally the verb in the Labrador Eskimo agrees in its inflections with the Greenland dialeet; but there are some special differences, and particularly with regard to the future, which has a threefold construction in the Greenland tongue, but is more simple in the Labrador speceh. With the ample means which the regular verb possesses of expressing every mood and tense, the Eskimo has little occasion for auxiliaries, and in fact the structure of the language is very regular and exact.

There are, however, one or two auxiliaries which have an affinity to adverbs — such as *pi-wok*, which is used in a variety of ways, sometimes in immediate relation to a noun, sometimes only as an adjunct to a verb: it occasionally seems to be equivalent to the English "get" or "do." When placed after participles, which is its most common position, it signifies the action of a thing. *Ipsok*, another auxiliary, seems to be equivalent to the Latin *est*; it often increases the meaning of the verb with which it is connected. "To be," or "to have," is denoted by the syllables *gi* or *vi* in composition, as *nunagiva*, "it is his land."

The adverbs are numerous, and have relation to time, place, equality, size, number, order, union, separation, &c.; and also to questioning, denying, affirming, negativing, including, excluding, desiring, admonishing, and distinguishing.

An example of the inflections of a single verb would occupy many pages, and cannot be given here; but the precoding short notices will suffice to show that vocabularies of the same language, formed by different people, may have little similarity, and that much care is requisite before we can venture to affirm the distinct origin of two tribes upon such evidence. In a language which is transmitted orally

368 VOCABULARIES OF NORTH AMERICA.

alone, and which is not preserved in its integrity by an appeal to the eye, alterations for the sake of euphony are frequent, and these, which are not uncommon with the Eskimo, vary with the delicacy of the car of the speaker. Thus when the termination *wangha* does not blend pleasingly with the preceding syllable, *langha* is substituted, and the general pronunciation is more nasal with some small communities, more guttural with others.

EXAMPLES OF NOUNS DECLINED TRANSITIVELY AND INTRANSITIVELY.

TUPEK, a tent.

	Sing.		Dual.						
Nom. tr. intr.	Tupek turkib	•	tuppak	•	-	-	turket.		
Gen.	turkib	-	tuppak	-	-	-	turket.		
Dat.	tuppek	-	tuppak	-	-	-	turket.		
	tuppermut	-	tuppang	nut	-	-	tuppernut.		
Acc.	tuppak	-	tuppak	-	-	•	turkinut.		
	tuppernik	-	tuppang	nit	-	-	turkit.		
Voc.	caret.								
Abl.	tuppermit	-	tuppang	nit	-	-	tuppermit.		
	tuppermut	-	tuppang	nut	•	-	turkinnut.		

NELEGARA, my master.

Nom. tr intr	. Nelegara - . nelekama -	nelegakka nelekang-ma	-	-	nelekakka. nelekama.
Gen.	nelekama -	nelekangma	-	-	nelekama.
Dat.	nelegara - nelekamnut	nelegakka nelegamnut	:	:	nelekakka. nelekamnut.
Acc.	nelegara - nelekamnik	nelegakka nelegamnik	•	-	nelekakka. nelekamnik.
Voc.	nelegara -	nelegakka	-	•	nelekakka.
Abl.	nelekamnit - nelekamnut	nelegamnit nelegamnut	:	-	nelekamnit. nelekamnut.

Nelegane, his master; neleganga, another person's master; and similar variations of the noun, have, in like manner, their various inflections.

ESKIMO VOCABULARY.

grity by an ony are frehe Eskimo, Thus ker. l pleasingly ted, and the small com-

١.

LY AND

Plural.

ırket.

arket. urket. uppernut. urkinut. urkit.

uppermit. urkinnut.

elekakka. elekama. elekama. elckakka. elekamnut. elekakka. elekamnik. elekakka. elekamnit. elekamnut.

> master ; and their various

COMPARATIVE TABLE OF THE DIALECTS SPOKEN BY THE BEERING'S SEA AND LABRADOR ESKIMOS.

Obs. S. denotes "singular;" D. "dual;" P. "plural." W. points out words taken from Captain Washington's vocabulary. a is sounded as in "father;" \bar{a} as in "law;" e as in "there;" i as ee in "see;" t as in "ink," "pin;" u as oo in "good;" kh or ch in the Scottish word "loch," or Irish "lough;" h after g signifies that the latter has the soft pronunciation as in "give;" the hyphen following g, gives nearly the same sound.

English.				Ku	skutch		Labrador Eskimo.			
The only	Crea	tor	•	•		-		Ping-ortitsi-o-wok.		
God (the	Crea	tor)	-	nuna-li	shta	-	-	Nuna, country.		
Heaven ; ment.	the	firm	a-	ki-il-ya	k	•	-	killēk.		
Earth. la	nd. a	count	rv	nuni	-	-	-	nuna.		
Air	-	-	1	u-i-ueb	u-vuo	hi-ak	-	u-i-a-wak, west wind.		
Air, wir	nd, a	lso t	he	-	-	-	•	silla (sillata, in the open air).		
Wind	-	-	-	-	•	-	•	anorre; sullu-ar-nèk, also		
The sun	-	-	-	-	-	•	-	sekkinek; (Perngit);		
Sun	-	-	•	akkta;	pukli	-anok	-	akki-suk-pok, the sun breaks		
Moon	-		-	tang-ek			-	takkek.		
Month		_	-	tang_ok	· . ino	Link	-			
Stars	-	•	-	mittit	-	-	-	ubluri-ak; ubloriak; (D.		
Comet	-	-	-	ar-i-ak	hn-ak	htak.				
A star su	rrour	ded	by	-	-		•	agsuk; (P. aguthet).		
Water		-		musek	-			immek, fresh water		
River	_	-		kvak	-			ku: kok: kogguk: koggut		
A larga	ivor	-	_		_	-	_	kokso-ak		
Son		-		immakl	nik		_	immak		
The wide	0000	- -		man	-Pres	-	-	immarbikeo-ak		
Laka	ocea			nonvik	-	-	-	akkor-oktok a lake uhene		
Lake		•	-	nanvik	•	-	-	deer are speared.		
A pond,	tresh	-	-			-	•	tessek; (P. tessit).		
Brook	-	-	-	kitchik	li-ak	•	-	kõgak.		
A tear	-	•	-		-	-	-	kogve; (P. kogvit).		
Straits	-	-	-	u-ikakh	-	-	-	ikkarasak, W.		
Cliff	-	•	•	-	-	-	-	ikkargok ; (P. ikkarut).		
Deeply c.	left	•	-	-	•	-	-	korok; (P. kor-kut).		
VOL.	II.				в	в				
370 VOCABULARIES OF NORTH AMERICA.

English.	Kuskutchewak.	Labrador Eskimo.
A long inlet		kang-erdluk.
Gulf or creek	nang-vagnak	(kang-ŭ-agnak ? u-agnak the west).
Current	tchag-vak	sag-vak; sarvak.
Current in the sea -		ingerarnek.
Current in fresh water		aksarnak.
Bottom of waters -	notu-ik	tung-a-wik, W.
He treads his boot down at the heel.		tung-mark-pok.
Shore	chna; agavnu-ik -	sīg-sak.
Mouth or source of rivers, a well.	pa-i	(pè, a court yard; pa, the round opening in a kai-yak).
A bank in the sea, a sunken rock	ithalh-nuk	siorak-so-ak-nuk, much sand; ipek-so-ak-nuk, much mud.
Stone	tkalhk-uk	ka-ertok, a rock.
A sea hird		tullik.
It is deep (valley or		ittiwok.
river).		
Day	ignu-ik	uvlok, or úblok, a day.
In the morning		uvlakut : uble-tillugo.
The day closes		uvlokliwok.
The morning, or the		kau; kauk; kaut.
The front or forehead		ke-uk : kauk : (P. karrut).
It is daylight		kau-ma-wok.
To-day		uvlut ; ubluk ; ovetsi-ak.
Night		u-nu-ak (P. unu-et).unuame
Marsh	mag-ik	(mannek. moss. W.).
Dry summer	ki-nu-ig-nu-ik.	
Summer		au-i-ak.
Mountain	ing-ik	kakkak; (P. kakket).
Lowland	tchu-iv-nu-ik -	(su-yuk, wet, dirty).
Shell	ammokt	amomio-yok, an oyster; (P. amomio-yut).
White shell fish -		ayöarnet.
Mussel		uvi-lok.
Snail shell		si-ut-terok, W.
The ear		si-ut.
A grain of sand -		si-orak (Dkek, Pket).
Sand	kag-u-i-ak.	
Clay (loam, chalk) -	magai-ak	machak.
Fire	knu-ik	ikoma, ignis.
One who fires a gun -		kukni-wok.
Wind	a-nuka	annore, W.; an-o-i, W.
Thunder	kalik.	
It thunders		kalukpok; kalludlarpok.
Lightning		kaumarlok (vide bright).
Rain	tchali-ali-ak; kitok	silla-luk; (silla, air).

ESKIMO VOCABULARY.

or Eskimo.

nak? u-agnak

rvak.

, W. -pok.

rt yard; pa, the opening in a

k-nuk, much ipek-so-ak-nuk, al. z rock.

úblok, a day. uble-tillugo. k. k; kaut. nuk; (P. karrut). ok. luk; ovetsi-ak. P. unu-et),unuame moss, W.).

(P. kakket). vet, dirty). ok, an oyster; mio-yut).

k, W.

. -kek, P. -ket).

pis.

.; an-o-i, W.

kalludlarpok. (vide bright). (silla, air).

English.	K	Kuskutchewak.			Labrador Eskimo.	
Hailstone		- kakhu	lat		-	(kakkulak, it is round, W.).
Anything sharp)		-	-	-	kakilaut; P. lautit.
It is round	-		-	-	•	angma-la-rik-pok.
Snow -	-	- kanikl	ı-chak	- 3	•	kannek, falling snow; (P. kang-it).
Ice -	-	- tchiko	-	-		siko.
Storm -	•	- anu-ge	vak	•	•	ani-gavak, an extraordinary quantity of snow; (anio, a snow storm).
Strong wind	-	- Canug-	wei. K	otz. S.	W.)	akkunak : akkunak-so-ak.
Wind or air	-		-	•	-	annorre ; annorer-ho-ak, great wind.
The wind is sti	11		-		-	annorre-karung-napok.
Calm -	•	- ku-nu	-ik	-		kunigok : (D. kunikuk).
Clear -	-	- tankik	h-teh	uk	-	alla-kak-pok.
A bright sky	-		-	-		alla-ki-wok.
Dark -	-	- telhk	-	-	-	tek. darkness.
Fog -	•		- \	-	•	tek-tuk; tartuk; (P. tar-tu-it).
It is foggy	-		-	-	-	niptai-pok. W.
The weather cl	ears		-		-	nipter-pok.
Vapour or fog	-		-	-	-	iseriak. (isse. severe cold).
A cold -	-		-	-	-	ikkě.
Clouded -	-	- tali-gu	ık	-	-	tali-pok, it is hidden; (P. -pot).
A cloud -	-		-	-	-	nu-vu-i-a.
Bright, or light		- noakh	tok	-		kauma-wok, it is bright.
Coals -		- khuma	wit	-	-	P. aumakut ; (S. aumako).
Ashes -	-	- acak	-			arsek. W.
Blue -	-	- vitok :	minu	kh-ka	t -	(minnu, a sea-weed, W.).
Bluish -	-		-	-		tungo-i-uktak.
Berry juice	-		-		-	tungo.
Red .	-	- kivago	k	-		auna-luk-tok, it is red.
Blood -	-	all all all all all all all all all all	-			ank : acout.
White -	-	- ngolh-	kak		_	kaggark-nok, it is white.
Night -	-	- unuk	-	-	•	u-nu-ak; (D. u-nu-ek; P. unu-et).
Smoke -	-	- punk	-	-	-	pu-i-ok, damp smoke, steam.
It smokes	-		-	-	-	pu-i-ok-pok.
Smell -	-	- nagnal	c	-	•	naimawa, or nai-wok, he smells something.
Man (homo)	-	- tatchu				
A shadow	-		-	-	-	tatchak.
A looking-glass			-	-	-	tatchartut.
Man -	-	- nukalh	ni-ak	-	•	(nukak, a brother).
Man (relation)	stock	-	•	-	-	ang-ut; (D. ang-u-tek; P. ang-ulit).
His father			-	-	-	ang-uta.
A helm -	-			-	-	ang-ut.
An adult -	-		-	-	-	ang-uti-marik.
People (Eskimo	(ac	- tagut:	VUOU	at	-	inu-it (S. i-nuk).

B B 2

VOCABULARIES OF NORTH AMERICA.

English, Kuskutchewak,		Labrador Eskimo.	
A name			taggisek.
My nation			tunnisuga.
Life			inusek.
A portrait of a man -			inu-i-ak.
White man, European			kablunak: (P. kablunet).
Evebrows			kublo: (D. kabluk: Pt).
Inland Indian stranger	allisaguk		allani-a-wok (S. allak).
A man's foot mark	uni-a-gux		alluk: (D alluk)
Stranger Felimon			si od lor min : ho-od-lor-
Stranger Hakimos -			
An unprotected man -			sole enviole
Husband	wi no		wi a manual man (P
nusband	vi-na -	• •	wi, a married man, (1. wi-nit).
One is with another -		• •	una.
She has a husband -		• •	wi-ghi-wok.
Wife	nuli-ga -		 nuli-a; (nuli-ang-a, my wife).
She or he has a brother's son or daughter.		• •	nu-a-karpok.
An unmarried woman			wi-ga-sok.
A bachelor			nule-tok.
Widower			nuler-tok
Old man	utobi nuk . a	nuli_uv	nulci-tox.
Male of man or boast	uten-nuk, a	unum-uv	ment
Hais old		-	itta-wak (itak W)
He is old		· ·	- Itth-WCK; (ItOK, W.)
Woman	aunals		uing i mole (al in D alti
woman	aganak -		ak, a mother).
Old woman	aganukli-uv	ak 🚽	- ning-i-vok.
The oldest of a family			- ang ai-i-uklek.
Brother's or son's wife		•	- uk-ang-a.
Grandfather	apnugli-uk.		•
Grandmother	annugli-u	-	- anenak-si-ak.
They have a mother -			 mikli-ak-attig-ekput.
Mother	ani -	-	- anenak ; akko-a.
Mother's milk			- ammak.
She is beautiful -			- enanau-wok.
A relative	tunka.		
Father	atti -	- ·	- attatak; (P. attatet).
The father		-	- attatu.
An adopted father -		-	- attatak-sak.
Son	igni-ak -		 ergnek: (P. ergnerit).
Daughter -	nanaga : nar	nik .	- nanik : (P. naniknit).
Brother	annak		(unenak, mother, heautu).
His elder brother			 anningna.
Elder brother or sister		-	- angosisuma.
Younger brother		_	- nukak + nukka my hrather
Sister -	agna-wu_il-		- nova · novence his sister ·
	agna-vu-ik	-	neyara, my sister.
Two uterine brothers		•	- angu-tauk-attigekpuk.
A LWIN		-	- likking-ui: karrisarek, huins.

ESKIMO VOCABULARY.

r Eskimo.
(P. kablunet). kabluk ; Pt). ; (S. allak). alluk). u ; he-ad-ler-
ried man; (P.
nuli-ang-a, <i>my</i> k.
(itok, W.)
; (aköa, D. akö- ther). klek.
nk. ttig-ekput. .kko-a.
ok. P. attatet).
c. P. ergnerit). . paniknit). nother, beauty).

n. akka, my brother. ango, his sister; my sister. -attigekpuk. karrisarek, twins.

English.	Kuskutchewal	4.	Labrador Eskimo.
A boy	tangoy-ali-uvak	-	nuka-pi.ak; (P. nuka-pitset)
A young woman -			ni-wi-ark-si-ak.
Agirl	nozi-atchak -		ni-wi-azi-ak.
Grandehild	tut-khih.		
A child belonging to the parents.	• • •	•	kittorng-ak.
A mother's only child		-	attung-ektak.
A woman's last child		-	mikki-erngo-a.
It is my child		•	kittom-yarivara.
An orphan		-	ananak-ang-ilak.
A fatherless child -		-	atatni-tok.
An orphan deprived of both parents.		•	illi-arksuk; (P. illi-arksu- it).
His sister's child -		-	u-i-orva.
A child		-	kittorng-ak; sorusek.
A little, or new born,		-	nutarak.
Uncle	anahkli-uvan.		
His mother's sister -	(root? aganak	, a	ai-ang-a.
His father's sister -			at-sang-a
Sister-in-law -			sak i-a. W.
A prudent woman			arnanda
A robust man -		_	atsu-ilik
A countryman -		-	nung-kut · (P nung-kattiget)
A friend, one of two		-	illek-sak.
A walking companion		-	tupperkat; (P. tupperka-
A travelling com-		•	ing-i-a-ket; (D. ingi-akattek
A comrade (house-		•	iglo-mokat; (P. iglo-mokat-
T	khvana		11-wing-a
Thou	lhou-it		ig-vit.
Thee acc.			illing.nik
He	ikum	_	toim-na.
We		_	u-vag-ut P. · (u-va-guk D)
Ye, you	ilhli-te-pik -	-	ilipsè, P.; (illiptik, D.; il-
They		-	okkö-a.
He demonstr.			taimna : tainsoma
This			tamanno.
He, or she, intransitive	1177.8	-	ung · tampo
He or she transitine -		_	oma : tansoma
They	unnt	-	okkös · tankön
Of mine		_	uvango: uwango
To me	kvinum -		uvamnut
To thee	lhninum -	-	iling-nut
Tous		_	the second second

373

ввЗ

VOCABULARIES OF NORTH AMERICA.

English. Kuskutchewak. Labrador Eskimo. To you -To him illipting-nut. umu-in ; ikumin omunga; tapsomunga. . To them okkoniunga; tapsomunga. Of himself . ing-me. -To themselves ingming-nut. kina? ki-a, who ? (P. kikut). Whose ? what ? what kind ? of whom ? kai-a; tchambi-a Who? what thing? suna? su-ub? (P. su-unt). • What do you say ? suva. • • • . ki-mit ; kikkunit. sumik ? Of or by whom With what thing? • • . • . . What company ? kikkut ? • . . Which ke kiput. . -His (vide of his). umnia • . Of his oma; tapsoma. kho-in-tchati-ka; Mine u-wanga. khvona. illi-pu-it; lipu-it-ik igvit, of thine. Thine • This, masc. unakh-wina una ; (inung-una, this man). -This thing oma; (oma-pung-a, his thing). That imna. -Self nang-ninek; (P. nang-. merngit). How ? what ? -• tcha-itun kannak. The same ingna. . Thus tava. Who are these people? kik-ut? who is the head of the family ? kamikuk; uksi-u; ni-akko; (P. ni-akkut). Head . niba-gun. Crown of the head kausek. ke-uk; ka-uk; (P. karrut). i-ye (ai-i-ga, W.); isse; (P. issit). Forehead tchughi-uk vi-tatu-ik Eyes • kablo, S.; (P. kablut). Eyebrow ka-i-ag-mi-ut -Evelashes kemerit-set na-i-utu-ik; tchu-usi-ut; (P. si-utik). Ears . tu-ik. Mouth • kanik kannerk ; kaurngit. --Face kènak. . • . king-ak; (P. king-et). naksuk; (P. naksu-it). nikh Nose • • A horn . -• -Cheek ulu-ak. -• -• katang-ak. Muzzle -• ki-u-tit; (S. kig-ut). umik; (P. umgit), also a Teeth khu-u-tu-ik . Beard unik curtain. Neck u-i-ak; (P. u-i-ait), fore u-i-anut . quarter of an animal; konge-sek.

ESKIMO VOCABULARY.

Eskimo. somunga. tapsomunga. ho ? (P. kikut). ? (P. su-unt). unit. . na. e. -una, this man). a-pung-a, his ; (P. nang-P. ni-akkut). uk;(P. karrut). ga, W.); isse; (P. kablut). i-utik). aurngit. P. king-et). . naksu-it). 8. kig-ut). umgit), also a .u-i-ait), fore f an animal;

Paglish.	Kuskutch	hewak.		Labrador Eskimo.
Hair (human) on the	nu-i-at -	•	•	nu-i-ak ; (P. nutset).
Hair, fur	mu-ilh-kut	•	•	merserpok, also a feather ((merkut, W.).
A needle		-		merkut.
A skin, general name	• •	•	•	amek; (D. amok; F
Hands	yagatchu-tu	-ik	-	aggait, hands and fingers;
Two, numer			-	aggait.
My hands	• •	•	•	aggakka; aggaktit, your
Foot	ig-uk -	-		ittigak : (P. ittiket).
Finger	si-evogat	-	-	
Thumb				kublo: (P. kublut).
Fore finger		•	-	tikkek ; (P. tikkerit), also a thimble.
A sign to indicate any-		-	-	tikkorut.
thing (finger post).				
Middle finger		-	-	kettert-lek, W.
The middle		-	-	kerka.
He is in the middle -		-	-	ketterpok.
Third finger		-	-	mikilirak, W.
Smallest		-	-	mikke; (mikkinek, the least
It decreases		-	-	mikki-orpek.
The first		-	-	mikkledklek.
Little finger		•	•	erkekok, W.; mikkillerak (P. mikkillaket).
Hands and feet to- gether.		•	•	igluktuk.
Belly	aksi-ak -	-	-	nek. also the body.
Tongue	ali-anuk	-		okak : (P. oket).
He licks with his		-	-	alluktorpok.
My tongue		-		okara.
A member of the body		-	-	nabgo-ak.
A leg		-	•	nabguk-pa; (ni-o, leg or thigh).
The trunk or body -		-	•	minumernet; time; (P.
A headless body -		-	-	kattik.
The back bone -		-		ku-i-a-pigak.
The rump		-	-	nullok.
Rlood	ka-i-unkak		-	auk : agout.
To speak	kalkhtu-ik	-	-	o-karlune.
He is spoken of				kalle-mayok.
To erv	vikhna-on-o	n.		
The waves roar -	The party of the state	-		kadlarpok.
He weeps much		-	-	kai-u-mi-wok-nudlarnok.
He distorts his face in		-	-	kakkerlu-arpok.
To laugh	nu-inhli-akh	ta.		

в в 4

376 VOCABULARIES OF NORTH AMERICA.

English.	Kuskutchewak.	Labrador Eskimo.
He laughs in mocker		- i-yorkpork.
He is in a laughing of weeping mood.		- illapsukpok.
To kill	tchikaliz-gi-u.	
Dead. (he is) -	tukumak -	- tokkowok.
A corpse -		- time (body) -tokkung-a-
•		yok.
Alive	u-nung-vak -	- innu-wok.
Life		- innusek.
A living man -		- innuk.
Bad	tchakii-uk -	- (sèg-lu-wok, he lies).
He is very bad		 a-yorpok; yudlarpok.
He becomes bad		- assilè-wok.
Not good -		- nama-lung-ilak.
Good	knu-ignag-kuk	- ai-ung-itok.
Very good, or great		- ang-i-yok.
He is a good man		- pillorik-pok ; ridlarpok.
Brave	tuvgak	- aksut, also strong.
Very brave or strong		- aksorso-ak.
Coward -	alantak -	- erksinadlarpok, afraid.
He is terror-struck		- sakko-arpok.
He is timid -		- innimi-wok.
He loscs courage		- kotso-alavok.
Thick	ukughelghi-a.	
Lean, not stout	kui-migu-ilhnagak	
He is stout -		- kuini-wok.
Thin, or lean, (he is)		- sallukpok; ludlarpok.
High	yukhtuli -	- portovok.
Low	yukh-kalhna-gak.	
Warm	kikh-tehatuk	- ki-ek-pok; onatomik.
Heat	kalhtok -	- kivek; onarsivok.
It boils		- kallapok.
To smell	nagne-chuk -	- nawok, he smells.
To spit	kchigu	- (sigguk, the beak).
He spits out -		- oviakpok.
To cough -	kuzgh-ga -	- ko-erlorpok, he coughs.
Pain	aknakhtu-a.	
Health	yuguntu-a.	
He is healthy -		- atsu-ili-wok, W.
Angry	wik-nu-i-chuk	- ning-akpok; gadlarpok.
Quarrel	agu-i-a-uk -	- akgiwok, he retaliates.
Terrible	alu-innakh-kuk	- adhei! (interjection).
Buy	kupuzg-u -	- akpang-erpok, he buys, W. pussi-wok, he buys it.
Taken	tkhwaka.	
Take	tkhu	- pi-uk, W.; pi-vok, or pi- wok, (auxiliary).
Give	ta-iz-ghu -	- ta-ug-si-lugo, let us barter. W.; pillata-wok, he gives in reward; pittipa, he gives it to him to hold

e

RICA.

Labrador Eskimo. pork. ukpok. (body) -tokkung-a-k. -wok. sck. k. -lu-wok, he lies). rpok; yudlarpok. 2-wok. a-lung-ilak. ng-itok. i-yok. rik-pok ; ridlarpok. it, also strong. rso-ak. inadlarpok, afraid. co-arpok. mi-wok. o-alavok. ni-wok. kpok; ludlarpok. ovok.

k-pok; onatomik. k; onarsivok. apok. ok, he smells. guk, the beak). kpok. rlorpok, he coughs.

-ili-wok, W. -akpok; gadlarpok. wok, he retaliates. i! (interjection). ing-erpok, he buys, W.; assi-wok, he buys it.

k, W.; pi-vok, or piok, (auxiliary). g-si-lugo, let us barter, .; pillata-wok, he gives reward; pittipa, he pes it to him to hold.

ESKIMO VOCABULARY.

English.			Kurkutche	ewak.		Labrador Eskimo.
Sell -	-	•	kiputna-waka	a		ni-u-werpok, he trades.
Lively, joyous	-	-		•	•	pio-ri-wok; ku-wi-a-sik- pok, W.
Merry -	-	-	nuna-nikh-k	uk		nunan-ghi-a-suk-pok, brisk.
Tedious -		-	nuna-ni-tu.			a 1 /
He is weak	-	-		-	-	nuneněpok : sanghe-pok.
Song -	-	-	i-vagun	-	-	iming-arkpok.
A song, hymn, o	or osa	lm		-	-	iming-erut-set.
Dance -		-	kazi-i-achi-k	ut	-	okkigenek, W.
Truth -	-	-	pachikh-pi-a	k	-	(padsitik-sak, an excuse).
Lie -	-	-	ikli-uk		-	seglu-wok, he lies.
Thief -	-	-	tu-joli-nak		-	tiglik-nok, he steals
Forest -	_	-	nu-i-ku-ig-w	akhtut		usportok trees comething
			nu-i-ku-ig-v	anutui		erect.
Place having r	io tre	es		-	-	nappartu-itok.
Grass -	-	-	tchangu-it.			
Straw or grass sea shore.	on t	he	• •	•	-	i-wik; ibgit.
Berries -	-	-	nangat -	-	•	paung-at.
Moss -	-	-	kumagu-i-tu	-it	-	ting-ang-yak, a bluish moss; marnek, W.; neka- gasek, W.
Sphagnum pal	ustre	-		-		orkso : (P. orksut).
Fir tree -	-	-	nu-ikvag-vak	h-tur	vek.	(1100)
Firewood	•	-		-	-	ikko-maksak, W.; (ikkoma,
Birch -	-	-	ilhgnuk	•	-	okpit; kai-volik; (P. kai-
Alder -	-	-	tohugyagyat			108.01
Poplar -	-	_	avant	,		
Willow -	-	_	tchagatu-it		-	okan-jak
Rock -	•	-	u-ipnat	-	-	u-i-arak, a stone; kai-
Ward a hash						ertok, w.
vessei, a bark	-		shunnak	-	-	umi-akso-ak.
A small boat o	of woo	od	· · · ·	-	-	umi-arak.
Baidare (skin	boat)) -	anh-1-ak		-	umi-avik.
Baidarka	•	-	pukhtan; ka	i-ak	•	kai-ak, (for one person); (P. kai-net).
Flat-bottomed	boat		anhi-akh-li-u	uk.		
Dog -	-	-	anna-kukta	-	•	kemmek; king-mek; (P.
Dog-sled	-	-	i-kam-chak	-	-	kam-utik, W.
Calls the do gether.	ogs	to-		•	-	kang-marpok.
Tanned sea-ec	w hie	le	amakh-kak.			
Arch -	-	•	ugli-vu-ik	-	-	korok, W., hollow.
A valley -	-	-		-	-	korkinek.
A bird arrow	-	-		-	-	nugit.
Arrow -	•	-	ikkh-uk ; tchagak.	pickh	-	karksok; (P. karksut).
Fish-hook				-	-	karsuk-sok.

VOCABULARIES OF NORTH AMERICA.

English.		Kuskutchewak.			Labrador Eskimo.	
Strap Hand drum, tambou	- r-	nuk-tchaklik tcha-ul.				
Shaman, sorcerer	•	tungalhkh ;	analb	kh-	anghe-kok.	
House .		tur.			iglo: (P. iglut).	
Hut, (abode of ma ried people).	r-	u-ina.			.B.o., (1B.u.).	
Tent	-		•	-	tuppek; (P. turkit).	
Snow house -	-		-	-	iglorigak.	
Indian tavern -	-	akumgavak.			5 5	
To take a vapour ba	th	mu-ichtak.				
He bathes, he dips i	t		-	-	missukpok.	
A.mour	-	annu-i-akh-o	chutu	-it.		
Guest	-	ali-anik	-	-	allak ; allani-a-wok.	
Give for a treat		tebaktchu.			,	
He gives a feast	-		-		nerri-marpok.	
Eat -		nuiga -	-	-	nerri-wok, W., he eats.	
Dining hall -	-		-	-	nerriving-me.	
To make a present	of	vaguzhah-ah	i ·		nilli-ta-wok.	
to make a present o		nikazhzah	-obi		I'm u woar	
To some	_	minka -	Brin.	_	moreornak he cause	
A noodlo	-	IIIIIKa -	-	-	workut: (P morkutit)	
Te heat	-	milli-alablen	-		anaumok W	
He oute it in pieces	-	phil-akiku	-	•	nillalmole	
Ded feb	-		-	•	pinakpok.	
Red fish	-	nu-l-ku-it.				
Salmo orientalis	-	tagni-akvak.				
Salmo sanguineus	-	Kak-ki-a.			1 11 1 (1) 1 11 1 (1)	
Salmo muksun -	-	ka-ukn-tut	-	-	ekanuk, (P. ekanuktut),	
					salmon trout.	
Salmo alpinus -	-	ankhli-u-gat	-	-	ekalli-et, trout.	
Salmo proteus -	•	atakak.				
Chaiko ?	•	nu-ik-ni-at.				
Syrka	-	imakh-ping-	at.			
Smelt	-	kpuka-chat.				
Eel pout	-	managnat.				
Pike	-	tehukvak.				
Fishing-net -	-	kughya.				
A bag, a poke	•		-	-	pok.	
Spawn	•	mass-i-uk.				
Cup	•	val-i-uk	-	-	korkok, wide-mouthed-cup; erngusok, drinking-cup.	
Spoon	-		-	•	alu-pa-ut.	
Pot	-	gant.				
Earthen pot, native	•	• •	-	-	illuterkut, W.; (illuli-wok, he hollows it out).	
Bladder	~	imangvik.				
Oar (boat's) -	-	anvagun		-	(pa-ut: P. pa-ntit): enat.	
Entrails					erchavit P. : (S. erchavik).	
Gut -	-	iggzh-n-joli-	nk		inelo: (P. inelu-it).	
Kamlaika (cloth)	-	iggmagna-tu	-ik.		, (

7.
ador Eskimo.
•
iglut).
P. turkit).
ani-a-wok.
, W., he eats.
-me. ok.
k. he sews.
(P. merkutit).
(P. ekalluktut),
trout. trout.
k, drinking-cup.
W.; (illuli-wok, ws it out).
. pa-utit); epat.
inelu-it).

ESKIMO VOCABULARY.

Engiish.		Kuskutchewak.		Labrador Eskimo.	
Woollen cloth -	-		•	ateg-ek-sai-ah, W.	
Parka	-	atkuk.			
Fur-boots -	-	kamu-ik-si-ak	-	kamikso-ak.	
Breeches	-	khulik -	-	karlik ; (P. karlit).	
Cap	-	nachak	•	ketsivak, akkordlek, also a	
Castor oil -	-	alli-ukit-khak.		Jucker.	
Beaver	-	kini-i-uli -	•	kig-i-ak; (P. kig-ilset).	
Otter	-	chvignil'nuk.			
Sable	-	kakhchichvak	-	karvi-ait-si-ak.	
Grey fox -	•	u-ikh-pu-ikhtuk	-	arvngasek.	
Red fox	-	kavhiatchak -	-	ka-i-ok.	
White fox -	-	ulhi-gu-ik -	-	teri-enniak ; P. teriennit- sek.	
Young fox -	-	pi-i-a-gak -	•	pei-a-raka, a young qua- druped or bird.	
Female bear -	-		-	akbik.	
Bear	-	unu-valh-jäkh	-	akhlak.	
Polar bear -	-		-	nennok: (P. nennut).	
Wolf	-	ku-isli-unu-ik		amarrok ; (P. amarkut) ;	
				ammarwok.	
Hare	-	ka-i-ukh-li -		ukalek : (Plit): ikkingna	
Wolverene -	-	Kab-tehak -		kablia-ri-u.	
Marmot -	-	kalb-ganakh-tuli		sik-sik : ullick	
Musk rat -	-	sig-vak -	-	kiy-galuk, W.	
Casan marmot (C	Citil-	kaninik -	•	ik-ik; sik-sik.	
Frmino	_	naculhkk-ak -	_	torrisis W	
Losser otter -		amagmi-utak -		nammi-oktok.	
Tail of an animal		anagini-utuk -	-	pammi-ok	
Mouso -		avilh-not -	-	owing-ok W	
Mouse	_	abuvot -	-	nivul-i-wol a lunge fu	
A spidor -	-	chuvat		assi-wok, (P assi-voit)	
A spluer -	-	in Aughi als . mi	1	historials (D historitant).	
Gnat, moschetoe	-	ghi-ak.	ĸu-	kiktoriak; (F. kiktoritset)	
Walrus tusk -	•	tul-1-ak -	-	togak, W.; (tok, a point).	
Walrus -	-	azgh-vu-ik	-	ai-wek; (P. ai-werit).	
Mammoth tooth	-	chagu-nu-ek.			
Back-fat of deer	-		-	tunnuk; (tunno, the back)	
Tallow	-	anu-ignak	-	kui-ni-wok, fat he is.	
Fat	-	u-ig-nu-ik.			
Rein-deer -	-	tun-tu	-	tuktu; (P. tuktut.)	
A large whale -	-		-	korchak; (P. korchetset).	
Delphinus leucas	-	chtvak	-	(sav-gak, a water-serpent).	
A seal, general na	ame		-	puese; (P. pue-sit).	
Seal, largest kind	-	izli-ugvak -	-	uksuk; oguk.	
Small seal -	-		-	netsi-arksuk; netsek.	
Middle seal -	-		-	kai-rolik.	
Young ditto -	-		-	pai-yarak.	
Seal lying on ice	-		-	otok.	
		1			

380 VOCABULARIES OF NORTH AMERICA.

English.	Kuskutchewak.	Labrador Eskimo.
A seal with a pointed		abba; (P. abbit).
Unborn seal		iblau.
The seal comes up to		pu-i-rook.
breathe.		P
I will	pivukh-tu-a	pivok.
I will not	piyuk-nak-tu-a -	piwak.
Gol	ai-i: ai-aghi	ailer-it. W.
Come!	ikh-tchika	kai-it!
He comes		kai-wok.
I come	ta-i-tchika.	
Bring	ta-iski-u	kaitsi-wok. to bring.
It is	pitankh-tok.	
Verily		ahammarik, (emphatic offir.)
Yes		katz : kassak : kaitsok :
		alieila ; ang-ernok.
Certainly not!	tchata-i-tok	se-i-ovut, aukai-lo.
He sava no		ang-ing-ilak.
Not		naukak, aukok.
Yet	tchali	sulle.
It is so (as you say) -		ahale.
Where ?	na.ni?	nane P
Where is it ?		nauk P
Whither?	nairt?	namut?
Which way? whereby?		naukut?
Here	khonikho	ovane mane : tamane.
Here about, around -		ovona.
There	vani: ung-napi -	mâne : tamâne.
Thither	vavu-it.	
Hither	akavu-it	ma-ungo: owunga.
There	khavana	ikkane.
Throw!	igazhghi-u	millorpok, he throws; egi- pok, he throws away.
So	khwatum	taimak : sorlo. as.
Now	khwatu-a	mâna.
In the present time -		manakut.
Later, afterwards -	atakh	king-urgane.
Aforetime		itsak.
Before	tehu-nu-imtpu -	sivurnga-gat.
Straight before -		miksane.
Before another -		ane-taima, also southward.
Behind	kanulhklimtiv.	
Backwards		knig-o-mut.
Above, upwards -	kulhma	kollanut; pa-ungo.
Below, beneath -	ochi-mi	kanna; sammand.
Num. 1	atu-u-chik	atou-sek.
. 2	a-i-nak : malhkhok	marruk : maggok, W.:
,, _	,	(agga, hands).
" 3	pa-i-na-i-vak	ping-a-sut ; ping-ahuk, W.; ping-a-nuk, W.
., 4	tchamik	sittamut.

CA.

ESKIMO VOCABULARY.

				_
dor Eskimo.	Englis	h.		
abbit).	Num. 5 -	•	-	t
	" 6 -	-	-	a
	"7-	-	-	a
	" 8 -	-	-	р
	" 9 -	-	-	cl
to bring.	" 10 -	-	-	te
	,, 11 -	•	•	
;, (emphatic affir.) ssak ; kaitsok ;	" 16 - " 20 -	-	-	t
ang-erpok.	21 -			
ukai-lo.	Whale -	-		2
k. ikok.	Bird in gene	ral	-	ti
	Eagle -	-	-	n
	Raven -	-	-	k
	Magpie -	-	-	k
	Hawk -	-	-	n
	Owl -	•	-	ig
ne; tamane.	Goose			
	Swan	-		1
nane.	Crane -	-		g
owunga.	Duck -	-	-	t
he throws . oris	Sinew -	-	-	ι
throws away.	Glass bead	-	-	t
	Blue -	-	-	te
ne.	White -	-	-	k
gat.	It is white	-	•	r
also southward.	Block	-	-	*
	DIACK .	-	•	
pa-ungo.	Krelle -	-	-	t
mmanu.	 Axe -	•	-	1
maggok, W.;	A. 2-0			
ands).	Pielesso	-	-	
; ping-ahuk, W.;	Knife -	-	-	
uk, W.	irme .	•		"
	Alantian av			1 1

E	nglish.			Kuskutchewak. Labrador Eskimo.
n. 5	-	•	•	tali-mik (tallek, the hand), tedlima;
6	-	-	-	akhvinok arvanget ; ping-a-su-yok-
7	-	-	-	a-i-na-akh-vanam - ping-a-sullo sitta-mello,
8	•	-	-	pi-na-i-vi-akh-vanam
9	-	-	-	chtami-akh-vanam - tellimella sitta-mello, W.,
10	_	_	_	tamini akh vanom - talling-voktut W
11	-	-	_	taninin-akii-vanitii - tenina-yoktut, w.
11	•	-	•	urkang-et.
10	-	-	-	arvertanget.
20	•	•	-	tzvinnak igluktut, hands and feet to- gether.
21	-	-	-	ungna.
ale	-	-	-	akh-vu-ik ar-wek.
l in g	enera	1	-	tu-in-mi-ak ting-mi-ak; (D. ting-mit- sck).
le	-	-	-	nu-itu-i-gavi-ak - nektoralik.
en	-	-	-	kolb-ka-gak
mie	-		-	kalla-ka-gai-ak
-	-	-	-	wantel
A	-	-	-	ingi alahan ali anili Wannigu ali anagi
	-	-	-	iggi-akutu-gan - upik, w.; upigu-ak, great
				owl, w.
se	-	-	-	nu-ikli-uk nerlek, (P. nerlit).
n	-	-	-	kughi-uk kog-uk, W.
ne	-	-	-	ghi-na-tali.
k	-	-	-	tu-in-mu-ik ting-mi-ak, (P. ting-mid- get), a bird, in general.
ew	-	-	-	ulhi-un; ivali-ut - uli-yut, W.
s bea	d	•	-	tu-ikh-lit (tuč, the shoulder); sang-
e	-	•	-	tchunaizi; tchu-a-gat tung-a-yuktak; (tungo,
ite	-	-	-	katu-ighi-agvak - kaud-luk-pok, W.; (kau,
				ady).
white	le	-	•	kaugarpok, w.
•	•	•	•	kivikh-tchitkkhiat - auk-paliki-tak; (auk, aggut, blood).
ek	•	•	-	tunulhgat kerngut; kernerpok; ker- nian-garvok.
lle	•	-	•	tchunaglat; anat - (sunak, W. polar-bear); (annak, refuse of animals).
•	•	•	•	kalhk-anak tukkingai-ok; (Put;
A			-	nollo_vok
0.80	-	-		
fa	-	•		ulbush (D alternation
ie	•	•	•	ulina-ut, (P. ulima-utit, <i>a hatchet</i>).
utian	axe	-	-	knun.

VOCABULARIES OF NORTH AMERICA.

English.		Kuskutchewak.		Labrador Bskimo.
Scissors He cuts something	e off	ku-ipli-a-unu-ik	•	kipsaut, W. kippiwok.
Needle	· -	minkuk: tchikuk		merkut: (sig-uk, a beak).
Button	-	nikht-ku-tu-it	-	sennero-ak; (P. sennerut- set).
Mirror	-	tangh-i-u-guk	-	takh-artut.
He sees him, or it	-		-	takko-wok.
Iron	-	nu-ilhkh-agak	•	kikki-ek, general name, also a wooden or ivory pin.
Copper	-	kanukh	-	kanu-vak. W.
Lead		khu-i-akak -	-	aggiktok, W.
Shirt	•	tulhpakhak -	•	uvincrok; (uvinite the flesh).
Linen waistcoat		alkuk	-	altighigha, under jacket, W.
Worsted ditto -		tunulbkhan-i-alku	k.	and and and factory it?
Kottle -		antohaval		nly-kn sik stone bottle
Dentelium shell	-	gantenavak -	-	uk-Au-Bik, stone neute.
Dentanum suen	-	numm-vagni-ut.		
Lar-rings -	•	aklatu-it -	-	ukla.
Long	-	tatkhli.		
Short -	-	nanilh-nuk -	-	nai-pok.
Broad	-	yu-gu-tuli -	-	silikpok, it is broad and thin.
Narrow	-	igu-kink-nuk	-	nerikipok, W.; amitok, W.
Fresh	-	milukapak.		_
Sweet	-	mi-iknik-kuk	-	mamakpok, it tastes nice.
Sugar	-		•	mamamak-sauk.

VOCABULARY OF THE KUTCHIN OF THE YUKON OR KUTCHI-KUTCHI, DRAWN UP BY MR. M'MURRAY; TO WHICH THE CHEPEWYAN SYNONYMS WERE ADDED BY MR. M'PHERSON.

	English.			1	Kutchi	n.		Chepewyan.
A	Animals bear - grizzly bear beaver	-		80 - si-i sè -	-	-	•	sasz. tlizè. tza: tsha.
	red fox black fox	:	-	na-kath nakath - iz- zè	- barha	- ita-ni	- il-	na-ghirhè-gossè. na-ghirhè-sin.
	cross fox white fox (a	- retie) -	nakath- etchi-a-	so thwi	:	-	na-ghirhè-netlizzè. na-ghirhè-gai.
	Canada lyn:	x	-	ni-itchi tsu-ko	:	:	-	ghisè.
	mink -	-	-	tchith-e	i	-	-	til-chusè; tekh-tusè.
	musquash	:	-	tzĕnn	-	-	-	tzěn; tshěn.

KUTCHIN VOCABULARY.

Eskimo.	
g-uk, a beak). (P. sennerut-	
general name, ooden or ivory	
w.	
V. (uvinite the	
under jacket, W.	
stone hettle.	
is broad and thin. W.; amitok, W.	
, <i>it tastes nice</i> . -sauk.	
Ситені-Китені	,
e Chepewyan	
hepewyan.	
gossè. sin.	
netlizzè. gai.	
tekh-tusè.	

n.

A wolf zo hare (American) - kà wolverene lekh-ethu-e - seal nīt-tchuk - tin-djukè - rein-deer bet-zey - goose krè swan ta-arr-zyne - crane che-a duck tet-sun - grouse akh-tail - tleukh-ko - tet-sun akh-tail - tleukh-ko - tleukh-ko - tleukh-ko - tleukh-ko-tak-hei gonus). pike alle-ti-in - blue-fish (grayling) methy (Lota) - tha An axe ta-è - Beads ta-è - bluto bluto bluto blue blue-fish (grayling) methy (Lota)	 yess; nuni-è. ka. nakh-ei. (nètsèk; netsi-arksuk, Esk). dunikh. bedzi. teha. kha-goss. dhell. yurrth-teho. dikh (pintailed gr.); kasbà (white gr). tlu-e-teho; tlu-e-zanè, trout thlu. uldai. thu-è-detla. tin-tellei. thuth, a spear.
hare (American) - $k\partial$ wolverene $lekh-ethu-e$ - $n\bar{n}t-tchuk$ - moose-deer $lich-tchuk$ - $tin-djukd$ - rein-deer $bet-zey$ goose krd $ta-arr-zyne$ - crane $che-a$ duck $tet-sun$ grouse $akh-tail$ fish, a salmon - white-fish (Core- gonus). pike $tet-sun$ blue-fish (grayling) methy (Lota) - tha A n awe $ta-de$ Beads $ta-de$ A tobace - box ta-de but - fish (grayling) methy (Lota) - tha A blanket $ta-de$ A tobace - box	 ka. nakh-ei. (nètsèk; netsi-arksuk, Esk). dunikh. bedzi. tcha. kha-goss. dhell. yurrth-tcho. dikh (pintailed gr.); kasbà (white gr). tlu-e-tcho; tlu-e-zanè, trout thlu. uldai. thu-è-detla. tin-tellei. thuth, a spear.
wolverene lekh-ethu-e - seal nīt-tehuk - moose-deer tin-djukè - rein-deer bet-zey - goose krè swan ta-arr-zyne - crane che-a duck tet-sun - grouse akh-tail - fish, a salmon - white-fish (Core- gonus). pike alle-ti-in - blue-fish (grayling) methy (Lota) - tha - A n axe ta-è - Beads ta-è - A belt ta-è - A tobace2-box A bonnet tsa-til-ek-ha. A cap tsa-til-ek-ha. A cap tsa-til-ek-ha. A cap tsa-til-ek-ha. A chisel A dagger A dagger	 nakh-ei. (nètsèk; netsi-arksuk, Esk). dunikh. bedzi. tcha. kha-goss. dhell. yurth-tcho. dikh (<i>pintailed gr.</i>); kasbà. (white gr). tlu-e-tcho; tlu-e-zanè, trout thlu. uldai. thlu-è-detla. tin-tellei. thuth, a spear.
seal	 (nètsèk; netsi-arksuk, Esk). dunikh. bedzi. tcha. kha-goss. dhell. yurrth-tcho. dikh (pintailed gr.); kasbà. (white gr). tlu-e-tcho; tlu-e-zanè, trout thlu. uldai. thu-è-detla. tin-tellei. thuth, a spear.
moose-deertin-djukèrein-deerbet-zeygoosekrèswanta-arr-zynecraneche-aduckche-agrousekrègrouseakh-tailfish, a salmontleukh-kogrouseakh-tailfish, a salmontleukh-kogrousealle-ti-ingrous).pikepike-blue-fish (grayling)methy (Lota)-Trading Goods.An awl-A naxe-A tobac-A tobac-bluton-tseltrow-ti-ak.Button-A bonnet-A cap-A duffle coat-A comb-a comb-a comb-theirzug.A chager-heirzug.A cheisel-A combA cheisel	 dunikh. bedzi. tcha. tcha. kha-goss. dhell. yurrth-tcho. dikh (pintailed gr.); kasbà (white gr). tlu-e-tcho; tlu-e-zanè, trout thlu. uldai. thlu-è-detla. tin-tellei. thuth, a spear.
rein-deerbet-zeygoosekrèswanta-arr-zynecraneche-aducktet-sungrousekh-tailfish, a salmontleukh-kogrouseakh-tailfish, a salmontleukh-kogrousealle-ti-inrwhite-fish(Core-gonus).alle-ti-inpikealle-ti-inblue-fish (grayling)rsi-tchamethy (Lota)che-tlukhTrading Goods.An awlthaAn awethathata-A belttseltaA tobacco-boxtseltaA bonnet-tseltaA capot or coatikA chiselso-itt-se.A combtha-ik.A duffle coatso-itt-se.A chelso-itt-se.A combtheir.A daggernil-ei-shoA fileso-itt-se.	 bedzi. tcha. kha-goss. dhell. yurth-tcho. dikh (pintailed gr.); kasbà. (white gr). tlu-e-tcho; tlu-c-zanè, trout thlu. uldai. thu-à-detla. tin-tellei.
goose -krè- $swan -$ ta-arr-zyne $crane -$ che-a $duck -$ tet-sun - $grouse -$ akh-tailfish, a salmonteukh-ko $white-fish$ (Core- $gonus)$.alle-ti-inpikeblue-fish (grayling)alle-ti-inmethy (Lota)-Trading Goods.An awlA blanket-A belttobacco-box-tseltatseltatseltatseltatseltatseltatseltoutotseltoutotsa-kol-u-tsa-kol-u-tselte coat-tselt coat-tselt coat-tselt coat-tselt coat-tselt coat-tselt coat-tselt <td> tcha. kha-goss. dhell. yurth-tcho. dikh (pintailed gr.); kasbà. (white gr). tlu-e-tcho; tlu-e-zanè, trout thlu. uldai. thlu-è-detla. tin-tellei. thuth, a spear. </td>	 tcha. kha-goss. dhell. yurth-tcho. dikh (pintailed gr.); kasbà. (white gr). tlu-e-tcho; tlu-e-zanè, trout thlu. uldai. thlu-è-detla. tin-tellei. thuth, a spear.
swan-ta-arr-zynecrane-che-a-duck-tet-sun-grouse-akh-tail-fish, a salmon-tleukh-kowhite-fish(Core-tleukh-ko-tak-heigonus).alle-ti-in-pikeblue-fish(grayling)alle-ti-inmethy(Lota)Trading GoodsAn axeAn axeBeads-nak-kai-c.A belttho.tselta-A tobacco-box-tseltrow-ti-ak.Button:A bonnettsa-kol-uA duffle coatA chiselA combA daggerA fileA bile	 kha-goss. dhell. yurth-tcho. dikh (<i>pintailed gr.</i>); kasbà (<i>white gr</i>). tlu-e-tcho; tlu-e-zanè, <i>trout</i> thlu. uldai. thu-è-detla. tin-tellei. thuth, a spear.
crane	 dhell. yurrth-tcho. dikh (pintailed gr.); kasbà. (white gr). tlu-e-tcho; tlu-e-zanè, trout thlu. uldai. thu-è-detla. tin-tellei. thuth, a spear.
duck	 yurrth-tcho. dikh (pintailed gr.); kasbà. (white gr). tlu-e-tcho; tlu-e-zanè, trout thu. uldai. thlu-è-detla. tin-tellei. thuth, a spear.
grouse	 dikh (pintailed gr.); kasbà. (white gr). tlu-e-tcho; tlu-c-zanè, trout thu. uldai. thlu-è-detla. thu-è-detla. thu-tellei.
fish, a salmon - tleukh-ko - tleukh-ko - tgonus). pike	 tlu-e-tcho; ilu-e-zanè, trout thlu, uldai. thu-è-detla. tin-tellei. thuth, a spear.
white-fish (Core- gonus). pike blue-fish (grayling) methy (Lota) - Trading Goods. An awl An axe beads A bluket - A tobacco-box - button: A tobacco-box - button: A cap A cap A duffle coat - A duffle coat - A dugger - A file - - box box box box che-tlukh - tha - - tha - - tha - - tha - - tha - - - - - - - - - - - - - -	 thlu, uldai. thlu-è-detla. tin-tellei. thuth, a spear.
pike alle-ti-in - blue-fish (grayling) methy (Lota) - rsi-tcha che-tlukh - Trading Goods. An awl tha An axe ta-è Beads ta-è Beads ta-è Beads ta-è A belt tho. A tobacco-box - tseltrow-ti-ak. Button: yei-kai-thit-le A cap tsa-kol-u - A bonnet tsa-til-ek-ha. A capot or coat - ik A duffle coat so-itt-se. A comb tseltrow-ti-ak. Button: so-itt-se. A comb their-zug. A file kuk-i	 uldai. thlu-è-detla. tin-tellei. thuth, a spear.
blue-fish (grayling) methy (Lota) - che-tlukh - Trading Goods. An awl tha An awe ta-è Beads ta-è Beads ta-è A belt tselta A tobacco-box - tseltrow-ti-ak. Buttol: yei-kai-thit-le A cap tsa-kol-u - A bonnet tsa-kol-u - A capot or coat - A duffle coat A chisel so-itt-se. A comb tseltrow-ti-ak. Bo-itt-k. A chisel so-itt-se. A chiger A file kuk-i	 thlu-è-detla. tin-tellei. thuth, a spear.
methy (Lota)-che-tlukhTrading Goods.An awl-An awl-An awe-banket-banket-tobacco-box-tseltrow-ti-ak.Button-bonnettsa-kol-u-A bonnettsa-kol-u-A cap-tsa-kol-u-A bonnetA capot or coat-A chiselA combA daggerA file <td< td=""><td> tin-tellei. thuth, a spear. </td></td<>	 tin-tellei. thuth, a spear.
Trading Goods.An awltha-An axeta-è-Beadsnak-kai-e.A blankettselta-A tobacco-box -tseltrow-ti-ak.Buttonyei-kai-thit-leA captsa-kol-u-A bonnettsa-kol-u-A cap ot or coat-tk-A duffle coat-chai-ikA combso-itt-se.teheir-zug.A daggernil-ei-sho-A filekuki -	- thuth, a spear.
An awl tha ta-è	- thuth, a spear.
An axe	
Beads	- thell; theith.
A belt	
A blanket tselta A tobacco-box - tseltrow-ti-ak. Button: yei-kai-thit-le A cap tsa-kol-u - A bonnet tsa-kol-u - A capot or coat - ik A duffle coat - chai-ik. A chisel so-itt-se. A comb tcheir-zug. A dagger - nil-ei-sho - A file kuk-i -	
A tobacco-box - tseltrow-ti-ak. Button yei-kai-thit-le A cap tsa-kol-u - A bonnet tsa-til-ek-ha. A capot or coat - ik A duffle coat - chai-ik. A chisel so-itt-se. A comb teheir-zug. A dagger - nil-ei-sho - A file kuk-i -	- tsurai.
Sutton::: - - yei-kai-thit-le A cap - tsa-kol-u - A bonnet - tsa-kol-u - A capot or coat - ik - - A duffle coat - chai-ik. - A chisel - - so-itt-se. A comb - - tehcir-zug. A dagger - nil-ei-sho - A file - - kuk-i -	
A cap tsa-kol-u - A bonnet tsa-kol-u - tsa-til-ek-ha. A capot or coat	- bun-eil-lay; pa-il-lay.
A connet tsa-til-ek-na. A capot or coat - ik A A duffle coat chai-ik. A chisel so-itt-se. A comb tcheir-zug. A dagger nil-ei-sho - A file kuk-i	- tsa-kulay.
A capot or coat	11
A dume coat challer. A chisel so-itt-se. A comb tcheir-zug. A dagger - nil-ei-sho - A file kuk-i -	- ekn.
A chisel so-ttt-sc. A comb tcheir-zug. A dagger nil-ei-sho - A file kuk-i	
A comb	
A dagger nil-cl-suo - A file kuk-i	la thuth
A life Kuk-1 -	- In-thuth.
	- nogunn; nok-keith.
A looking gloss - mutchei o i o	
A roun - to organ	tol-minthà
A gun fint hoch tai	- tel-gurthe.
A gun more - beca-to	- He-tell.
Gunnowden togge ton	- tolgumo koun nd
A powelow hown	- teigurre-koun-ne.
A rottle this	+113
A knife	- hòss
A ring	- 0035.
A shirt	tso tsi oh · thisitai
A small shot togga_atsil _	- tolistheis
A ball	- toll-gith-toho
A fractol	- ven-grun-veno.
Cloth thit.	U
Thread at a this is the	

VOCABULARIES OF NORTH AMERICA.

English. Kutchin. Chepewyan. sel-tu-i. Tobacco se' ei-i-ti-it -• Trowsers illei-ik -(karlik, Eskimo). _ Vermilion tingi-ta-tseikh. _ Miscellaneous. A tree -A willow tetch-hau tsu. kai-thsinnè. kai-i ---Grass tlo tlo. • . The ground • กบักก nih. • Water -. tehu tu; to. . A river -han dessh. A lake theu-tu-ĭ. van akh-tsin Rain dsha. konni-etha Warm -etu. Cold . konni-eka etdza. Hungry --. sei-ze-kwetsik seth-ithu. Fatigued -• kei-a-sethelth-krei ni-nitsau. ĕth-ill-seyk -Sick ai-a'. -A mountain • • tha sheth. A valley -The sun -kra-tannè shegussè. . sakh. r' sey-è --The stars thun thun. -A rock -. tchi thi 'tsunnè-cho. A house or fort izzè. ni-ti-a -A lodge or tent nèballè; nepalle. -A bow alt-heikh . elthi. --An arrow ki-e -. kah. A canoe tri tsi. • . neir-zi -Good nesu; neso; nazu. bets-hè-tè neso-ulla. Bad --Dav . • tzin tzinna. . tatha hetleghè. Night -• • Sleep nokh-tchi belkh. • -• Rest --tuggath-illa-è thĭlleh. . To sit • tchith-u-ĕtcha théda. To walk -• ka-whot-ĕl nathall. • sha-tocha thebakall. To run -. at-ĕl-ke thelguth. To shoot -• beshei-en-i-echa thega thul. To kili --A man • tenghi -'dunne; duneh; 'tinne. -A woman tren-djo tshēkwè. • A boy tse-a dunne-yazè. A girl A dog mitchet-ei tsekwe-azè. thling; thline. bet-tchinnai. . tleine -A sled latchan-vultl -. Numerals. nthlare, D., (en-clai, L.), (sthlagi, C.). nakkhe, D., (nakka, L.), (nakke, C.). 1 • tih-lagga • 2 nak-hei -

is D C.

so

KUTCHIN VOCABULARY.

epewyan.		Englis	h.		Kutchin.	Chepewyan.
	3	-	•	•	thi-eka	khtare, D., (ita-rgha, L.), (takkè, C.).
skimo).	4	•	•	•	tān-na	tinghe, D., (iting, L.), (tingee, C.).
	5	•	-	•	illa-kon-člei	zazunlare, D., (sa-soo-la,
ð.	6	•	•	-	neckhki-ět-hei -	eleathare, D., (ut-ke-tlai, L.), (alkitakhe, C.).
i	7	-	-	•	ataitsa-newk-he -	nthlazuntinghe, D., (kko- sing-ting, L.), (sthlasi- tingie, C.).
	8	•	-	•	nak-hei-etan-na -	alcatinghe, D., (elzenting, L.), alketingie, C.).
	9	-	-	•	nuntcha-niko	nthla-otta, D., (kkahooli, L.), katchine-onnuna, C.)
	10	-	•	-	tikh-lagga-chow-et- hi-en.	'nthla-una, D., (ito-nanna, L.), onnuna, C.).
	11	-	-	-	tikh - lagga - mik- ki- tagga.	(sthlagi-juthet, C.).
	12	-	-	-	nak-hei mikki-tagga	(nacke-juthet, C.).
	13	-	-	-	thi-eka-mikki-tagga	(takhe-juthet, C.).
	14	-	-	-	tanna-mikki-tagga -	(tingee-juthet, C.).
è-cho.	15	•	-	•	ilakon-ĕlei-mikki- tagga.	
nepalle.	20	•	•	•	nak-how-chow-ethi-en	(non-nanua, L.), (nackhe- onnuna, C.).
	21	•	-	•	nak-how-chow-ethi-in- unsla-tikh-lagga.	(nacke-onnuna, nathetsin sthlage, C.).
	30	-	•	-	thi-eka-chow-ethi-en	(tacke-onnuna, C.).
so; nazu.	40	-	•	-	tanna-ha-chow-ethi-en	(tingie-onnuna, C.).
	50	•	-	-	atla-konělei-chow-ethi- en.	(sasulagi-onnuna, C.).
	60	•	•	-	nikh-ki-at-hei-chow- ethi-en.	(alkitakhe-onnuna, C.).
	70		-	-	atait-sa.	
	80	•	-	-	nich-ki-etanna-chow- ethi-en.	
	90	-	-	-	muntcha-niko-chow- ethi-en.	
l. duneh ; 'tinnè.	100	-	-	-	tikh-lagga, chow-ethi- en-chow-ethi-en.	(onnuna-onnuna, C.), ten tens.
zè.	200	•	-	•	nak-kaggo-chow-ethi- en-chow-ethi-en.	(nacki-onnuna-onnuna, C.).
e. hline.	300	•	-	•	thi eka-chow-ethi-en- chow-ethi-en.	(takhi-onnuna-onnuna, C.).

Note.—The orthography of the names of numerals enclosed by crotchets is different from that of the other parts of the vocabulary. D. denotes Dog-rib words obtained by myself. L. is Dog-rib recorded by Capt. Lefroy. C. denotes Chepewyan words extracted from a list furnished by Mr. M'Pher-son, who has adopted the French orthography in part.

VOL. II.

D., (en-clai, L.), i, C.). D., (nakka, L.), . C.).

CC

CA.

386 VOCABULARIES OF NORTH AMERICA.

The fragment of a vocabulary of the Chepewyan dialect, which follows, was formed entirely from the diction of Mrs. M'Pherson, to whom the language has been familiar from her infancy. It was written to the following manner : Having at hand a pretty full be abulary of the Cree, drawn up at Carlton House i. 320, in which the words were arranged in alphabetical order, I propounded the Cree expressions to her in succession, assisting her with a French translation when she had any doubt of their meaning. The Chepewyan equivalent was pronounced by her again and again, until my ear caught the sound, and I was able to repeat it after her. I then wrote it down, and read it to her from the manuscript. Such words as I was unable to pronounce to her satisfaction, and they were not few, were left out. The nasal sounds resembling the French final nwere the most difficult, and they are of frequent occurrence in the language. The Chepewyan tongue also abounds in the burring sound of the letter r combined with an aspirate, which I know not how to express in English; and such words have consequently been left out of the vocabulary. The ordinary aspirate, similar to the och of the Scottish or Irish, is denoted in the vocabulary by kh. The vocabulary, short as it is, took some weeks to produce. It was interrupted by a change in our arrangements in travelling, canoes having been substituted for boats, which made it less convenient for me to receive lessons in Chepewyan. This difficulty would not, however, have prevented the prosecution of the task, especially as Mrs. M'Pherson with much kindness expressed her willingness to proceed until we had gone through the whole Cree vocabulary, of which about nine-tenths remained; but knowing that the language was becoming a written one, under the active superintendence of the Roman Catholic missionaries at Isle a la Crosse, I gave up my intention of endeavouring to ascertain its structure, and contented myself with the following specimen.

A A A A A A A A

y y M

Al

CHEPEWYAN VOCABULARY.

CA.

wyan dialect, ne diction of been familiar wing manner : of the Cree, ich the words inded the Cree with a French meaning. The her again and I was able to and read it to was unable to not few, were French final nent occurrence also abounds in ith an aspirate, lish; and such the vocabulary. the Scottish or

The vocabuoduce. It was ts in travelling, which made it in Chepewyan. prevented the Irs. M'Pherson ness to proceed vocabulary, of nowing that the nder the active missionaries at of endeavouring nyself with the

VOCABULARY OF THE CHEPEWYAN TONGUE, WITH CREE AND ENGLISH TRANSLATIONS.

A as in "father;" a as in "awe" "law;" è as in "theme;" e as in "dell" "bed;" i as in "ravine," or as ee in "see;" I as in "ill" "ink;" o as in "or" "for;" ŭ as in "us" "husk;" ai as in "aim" "nnaim;" an as ow in "how;" oy as in "hoy;" yu as "you;" eu as ew in "dew" or eau in "beauty;" kw as in "awkward;" ng as the French nasnl n; kh as ch in "loch," Scottice.

Cree, of Cariton House.	Chepewyan, of Athabasca.	English.
A.		
Abu-ye	tu-a-will	liquor, soup, or drink.
Agatha-shu, or aggai- a-shu.	the-ut-'tinne	an Englishman.
Aggiskow, or akkiskow	el-ka-ti	pin-tailed grouse.
Aggusk	sis-thère	a blunt arrow.
Atchak, or akhchak -	i-yu-nè	the soul.
Ai-ammi-hè-u	vu-alanè-pallè -	a flag.
Akop	tsirrè, or tchirrè -	a blanket or covering.
	Tsirrè-kai-cho -	a large blanket.
Ai-u-wannis	vu	all kinds of goods.
Akwa-napov-igan-askek	tille-arakai-ĭnka -	a covered kettle.
Akwatin	hātkin	frost.
Amĭsk	tza	a beaver.
Amis-kwa-wiste -	ekhkè: tza bèkong	a beaver house.
Āmu: amo	klize : ti-ranna -	a bee.
Annèk-kutchass -	tli-i: tchillè	a squirrel.
Annèk-kutchassis -	tillel-kuzè	small or ground-squirrel.
Apikh-tow-kishi-kow	'tchi-èn-tize; tchinnè-	mid-day, or half a day.
A	tan-m-se.	
Apikn-tow-tippiskow	thir-hize	mianight.
Apisi-mongsus	1 hours a salitat :	jumping deer.
Apiste-shipis	a-sè akhth.	a teal, or small duck.
Apistat-tchèkus -		prong-horned antelope.
Appakwa-sun	ni-pallé	a leathern tent-cover.
Appèk-kusis	kleunè	a mouse.
Appisk	tannonè-tcho, big bird	black or white-headed eagle.
Appistis-kis	kai-yazè; kai-gusè	a Hutchins's goose.
Appovè	toth; tö-a	paddle or oar.
Appoy-nask	kès	a spit.
Appüssuk, (Pl. appüss- ye-akhtik, or appussu-	thai-yc; nepalli-tetchun (<i>tent legs</i>).	tent poles.
yuk).		
Miskahtuk	nepalli	a man's legs.
Akhtai-yè (Pwuk)	thè	a fur skin.

VOCABULARIES OF NORTH AMERICA.

Cree, of Cariton House	Chepewyan, of Athabas	ca.	English.
Ammiskus tai và	trabatha a		hoaver skin
Aultile	Arillà	-	a kattla
Askin pasiski	kmotlda	-	a Kettle.
ASKI; HOSISKI	Kwottes -	•	innu.
Intaski	ni-tanninne -	•	my native land.
Kit-aski	na-minne -	•	your native land.
Int'askinan		•	our native land.
	Be-anninne -	•	his land.
Aski-tin-wè-as; aski- we-as.	bét; per-elind	•	raw or fresh meat; flesh.
Askow-i	ten-de-ila (ice, he not); ten-nailer.	ard,	holes in the icc.
	Kin-the leuk -		ice breaking up.
Assam (P. assam-uk)	akhè; akh; akhi		a snow-shoe.
Alikik	tīllè		a kettle, or copper kettle.
Assist	Sampas-tille -	•	a tin kettle.
	thekn.	1.3	strug ning on columnst
Assini-uspogan	tut-thékh -	ne-	stone pipe, or calumet.
Assini-watche-a -	sheth; thè-she	•	Rocky Mountains.
Assini-poyt, or, E-askab		•	a Stone Indian.
Assiske	otlès		mud or earth.
Assiske-pakwè-sigan-	thlès		wheaten flour.
Assiggan (P. assiga-	tel	•	a sock, foot stocking.
Assiss-wi	ètlè		an ice chisel (lit. a horn).
Astu-thèggum-ik -	tsi-yè	•	a shed in which canoes are
Astu-tin	tsá	•	ladies' cap or bonnet (beaver).
	'Tsa-kalló -		man's hat.
Astum-astaik	tsa-ne-tum -		sunshine.
Atchakht or atchak -	thin		a etar
Atchappi -	oltà	-	a bow
Atchappi-	blawlahd-olting		a fiddle
AtohYmmosia	thline word		a huure.
Atulth on ottols	at this	•	a puppy.
Attent	et-thin	•	rem-deer.
Athappi	ta-oith	•	a net.
Athabiskow -	the-minne-u-ye	•	a rocky country.
Atha-wak-kiska-mat- tinow	hokar-ritha -	•	a very steep bank.
Athā-wastin	tethi-èl	-	a calm.
Athik	tsai èllè	•	a frog (grenouille).
	Tsai-el-cho -		large frog (crapaud).
Athuskan	ta-kallè-chi-a		a raspherry.
Atchak-ash	til-chusè		a mink (mustela lutreola).
Attei-gan -	VII		trading stock
Attikh-hamer -	thlew	-	white fish (corregonue)
Attim	thling	-	a dog
Atueni	kaithlin einnd -	-	aldon
Aku-nusà-win	thaisi -	-	a plattar
4 2 08 08 1/ VLU W = 17 414 -	WARRAN -		

No.

CHEPEWYAN VOCABULARY.

۸.

English.

in.

e land. ve land. e land.

esh meat ; flesh.

he ice.

ing up. noe. or copper kettle. tle.

oe, or calumet.

Iountains. Indian. earth. flour. foot stocking.

hisel (*lit.* a horn). n which canoes are

cap or bonnet er). at. e.

er.

country. teep bauk.

grenouille). og (crapaud). erry. (mustela lutreola). stock. sh (coregonius).

Cree, of Carlton House.	Chepewyan, of Athabas	ca.	English.
Āpètte-kā-hīggan -	denti-lita-thil-tillè	- 1	a chest lock key (properly, but used for keys in ge-
Akhikhk (a guttural	hèkh	- 3	yes.
grunt).			
	nanne		across.
Akwo-kukhtin -	yuuna		on the other side.
Annutch a attà	An har -	1	it is moundy.
A ++>	kultund -	1 1	now; at the present time.
Annutch hak ky sikely	Killune		this name day.
Annutch-kak-ke-sikak	u-uzinne -	- L'	this very day.
Annuten ka-tippiskak	terri-kitina -	- 1	uns mgat.
Annu-watch-gal-as -	tunta)	-	rather long ago.
Apikn-tow	tanize	- !!	in the middle.
Apatishew	bet-arutha -	-	it is useful.
Appatun	bet-taritha -	-	useful.
Apputchiga			once on a time.
Askow	athke	- 1	sometimes.
Astum-uspi	ekku-aze -	-	since such a time.
Athi-mun	sutu-yè	-	it is difficult.
Athè-wak	ona-hadzūn -	-	more.
Athè-wak kishè-wak-	edzun-kuthè -	-	nearer; very near.
Athe-wak-pètsow -	hona-hedza-nitha	-	further; very far.
Eshunila	hulè-ho	-	he is troublesome; badly disposed.
Ai-ā	nitsa-ula -	•	keep it; have thou it.
Int'ai-an; or int'ai-a- wā-u.	se-itza-heila; huni zoni.	è-	I possess it; it is minc.
Kit'ai-an	netzè	-	it is yours.
Ai-akuski-tè-u	petothè-karth -	-	it has a broad bill.
Ai-āmi	yan-ilti	-	speak thou.
Ai-amew ; ai-atchi- mè-u.	yalti	-	he talks.
Ai-amihin	zedzun-yar-ilti	-	speak to me.
Ai-ami-hi-tu-tak -	althlai-yalthi (toge let us speak).	ther	let us talk together.
Ai-ami-hi-tu-wuk -	elthney-alti -	-	they talk to one another.
Ai-ami-hè-u	vedzonne-alti-	-	he spoke to him.
Int'ai-amin	è-āsti	-	I talk.
Int'ai-ami-ha-u -	bedze-asti -	-	I talk to him.
Int'ai-ami-hik	zedzun-alti -	-	he spoke to me.
Kit'ai-ami-hik	nedzunè-alta -	-	he talks to you.
Ekau-witha-atche-	zedzun-ye-innè-al hila-kula.	ti-	do not tell it.
Ai-atchčmow-akwa -	nu-hei-lund -	-	tell us the news; relat thou now.
Ai-anè-tika-	pevè-onla-honnà	-	it is full of partitions
Aikh-tu-kā-mik -	nu-anku -	- 1	another house.
Waska-iggan -	và -		a house.
Wiggi		. 1	a tent or dwelling
	1	-	a cont of anening.

15

VOCABULARIES OF NORTH AMERICA.

Cree, of Carlton House.		Chepweyan, of Athaba	sca.	(English.
Ai-ish-ku-shu; ai-isl ku-tè-u.	1 -	kalyè-ni-nan-idza	-	he is tired (with walking).
Ai-ish-kutan-nè-wu Int'ni ish-kuzin Int'ai-iskutchè-man	-	kalyè-ni-tan-idza kalyè-ne ninna-ch toth-ne-zin-alnilza		they are tired (ditto). I am tired (ditto). I am tired with paddling.
Kitai-iskutehè-man- nă?		teth-ne-ni-nan-ilz uza ?	a-	are you tired with paddling?
Ai-u -	•	nu-a-edzon-illa	-	he is there.
Akkushew -	-	ey-a-hilla -	-	he is sick or ill.
Int'akushin -	-	ey-a-hèzlè -	-	1 am sick.
Akkustemmu -	-	edzil -	-	he is wet.
Int'akkusktemmun	-	dzedzil	-	1 am wet.
Annĭskutāpan -	-	chās-inninne-ai	-	a knot.
Anniskutāpè! -	-	chās-nos-al! -	-	tie a knot !
Ing'annisku-tāpan	-		-	I will tie a knot.
Gè-annisku-tā-pè-u	-	chās-ninne-al	-	he has tied a knot.
Apikh-ku-pai-u	-	tey-kunnè-takh	•	it has become loose; it is loose.
Ge-appaha -	-	tey-kunnè-arlth; r ke-urth.	ney-	he has untied it.
Apith-kuna; appaha	ι-	pey-kè-urth -	-	loose it (a knot); open it.
Nè-gè-apith-kunain	-	kalthonna-pev-kè	urth	I have loosened it.
Ne-ge-ava-hain	-	ev-kè-urrth		I have untied it.
Int'anikh-tā-nă-hā-u		peino-harre-kluk	-	I gave him a blue eve.
Int'anikh-tā-pă-huk	_	zunno-arrè-kluck		he gave me a blue eve.
Apisal	-	per-il-thilth! -	-	warm it (as a garment at the fire).
Apisum	-	yi-èr-il-thilth	-	he warms it.
Sā-sev-int′apisain	-	kuda-ber-il-thilth	-	I have already warmed it.
Appil -	-	thein-'tă! -	-	sit down!
Iltè-anni!	_	ev-er-thein-ta!	-	sit here! (here sit!)
Annew -	- 1	nèltă	.	he sits.
Appè-wé-ŭk	_	hed-nilthi -		they sit.
Int/appin -	-	thistā		Teit
Kitannin nx 9 -	-	thin-to-wang?		are you sitting?
Kit appnin nx 9	-	unto uzong		do you wish to sit?
A shamin	-	unta-uzang	-	give me food to get!
Asnainin : -	-	bega-van-nenu :	-	T will give you food to eat!
Niga-asnami-tin	-	he-a-urchu -	-	I will give you lood to eat.
Michema; hughes	-	bet-ho	-	victuals.
Aspun-ishew -	-	ā-ă-ontzun -	-	he is niggardly (of his victuals).
Assitĭnă	-	èltan-nilè -	-	mingle them; add one to another.
A-sustatin -	-		-	it is hidden.
Kiga-kasustatin	-	necha-itus-'i -	- 1	I will hide myself from you.
As-swè-tè-u -	-	te-vè-thèlla -	- 1	it is in (a bag).
As-swè-tā-u -	-	te-yè-yèlla -	-	he puts it in.
Int'aswethn -	-	te-ve-ila -	-	I put it in.
Aswèthim -	-	hega-etu-u-elnè	-	be on your guard against
AN IT OFMINE				him.

CHEPEWYAN VOCABULARY.

l	l	L.	2	
ŀ	ł		4	,

glish.	Cree, of Carlton House.	Chepewyan, of Athabasca.	English.
ith walking).	Aswithi-min	zethè-sekor-u-elné -	be on your guard against
l (ditto).	Ing'aswithi-mow -	pa-us-o-èlnè	I will be on my guard
ith paddling.	Atchis-chapum-wè-u	na-seil-hitchè	he gives a side glance to a
with paddling?	Athag-uskow	ne-ctèl	it is broad.
	Athin-isew: evthiniseu	hung-va: huva -	he is wise or knowing
	Int/sin-gathi-uisen	hong-she-a	he is wise or prudent
111.	Athinew -		he is abstemious
	Atuske-n-	è-hul-ana	he labours.
	Int'atuskaim	e-walasna	Llabour
	Atta-wanna! (imn)		harter ! trada !
	Ki-wi-atta-wanna?	na-ukh-unouza?	will you barter?
	Atto-wa-gun		goods for trade
not.	Attathowskil	sel-honninnè!-	tell a story or fable
knot.	Atta-thow-Ki	Honnè	a story
ne loose; it is	Attà-michi-ka-tà-n	necha-ladi-nelthun -	ho grove bigger
	Attà-mi-shà-u -	neena-nau-nermun -	it grows bigger.
d it.	Atikh_tà_n	nouth-lurth -	it is ripo or wellow
	Ationus	neuth-furth	she dues on tingen it
not); open it.	Int/stigsin	uuedza	I due it
ned it.	Atima am	uriuza	La ouentaleas
d it.	Auma-ow	na ni osha	T overtakes.
a blue eye.	Int atima-ow	hen nicha lillà	I overtook nm.
a blue eye.	Ing atima-ow	ben-nisna-inie -	I will overtake nim.
a garment at	Int atomik		he overtook me.
	Int atimanuk	a a a a	ne overtook me (by water).
	Atimi-thowuk	nar-neitein	they ny from us (birds).
dy warmed it.	Attohu	tehirr-lite	he is choking.
	Int'attohun	tchirr-este	I am choking.
ere sit!)	Ki-wi-au-tote memi-	et-te-to-tin-in-uste-	I wish to be your friend.
	A-wuss; a-wussètè!-	nusè!	keep off! let me alone!
ng?	E		
to sit?			
d to eat!	E-atchi-inyu-wukk -	et-dunni-'tinne -	Indians of a strange nation
you food to eat.	Ek-kwa	yah	a louse.
drink; food;	Epètchè-kishi-wükk -	yelkon	dawn of day.
	Eskann-shi-ka-un -	edtè-thidzi	a horn comb.
ardly (of his	Eskwai-atch-tchi- tchan.	tinnè-la-dthaille-dzilla	the last or little finger.
m; add one to	Esputtinow	kokkarritha	high ground; a bank (une cote).
	Ethik-kwatin	ne-edia	hoar frost - hoar frost.
and from you	Ethiko-new -	ine caju	hoar frost - rimy.
nysen from you.	Ethikwuk	tcharti -	ants.
ag).	Etiskew	èkei-ghè	foot-mark or track of an
	×	ou 1	animal.

VOCABULARIES OF NORTH AMERICA.

Cree, of Carlton House.	Chepewyan, of Athabasca.	English.
E-a-kusin : thah-kusin	nedtha	light.
	Ned-tarrilla	not heavy.
E-anitch		still : quiet.
- aprica	Na teillà	it stirs not.
Èka	hila	not.
Enètebè-kishikak	inte	as the day was coming
Ekushi-kak	dzithà -	by day
Kishiskow : kisgow -	dzinà	the day
Ekospi · àg-mieni	klasing_tingd_voivd_	at that time
Fackwo-pukku	ashmoh	only that
Eskwai-atch	no-ontò -	lost
Eskwal-aten		a little above
Espinini-sik	l-yaze-beke	a intre above.
Espinink	the definit she	above.
Etakkusik	the-dzini-gne -	yesterday.
Ethipinne-ok-tapo-an	oti-a-eitne	truly.
Euppiskak	hedkleghe	by night.
* 1	Dza-kin	beaver lodge.
E-aske-u-	dza-kin-nannelya -	he breaks up a beaver lodge.
Int'e-askann		I break up a beaver lodge.
Ekau-witha!		do not!
Entau-wi		go and open it.
E-ukh-tinnè-gatè-u -	peta-harelta; peta-ha- elta.	it is opened.
E-ukhte-nammuk -	peta-klell	open ye it.
E-ukhte-num	· · · ·	he opened it.
Nè-ukhtè-nain		I opened it.
Ètapoy-ikhta	bethna-ilkis	mix it; stir it.
Ethepo-akwow; nepo- akwow; athin-ni-	huya	he is wise — knowing.
Ey-thin-akhtěk -	eln	small spruce fir (Abies bal-
Frythin atta abishin -	tabith_taba	stock duck (Anas houshas)
Ey-tum-actu-smiship-	vltei và	stock duck (Anas boschas).
Ey-thinni mine	ultai-ye	a pike or jack.
Ey-thinni nithov v		Canada grundo
Ey-thin m (P or	'dainno	on Indian of the anoshou's
thin_my_mille)	uunne	nation
Inter and hult		he mode my eve ashe hare
Intey-apa-nuk -		blow.
Tans-ey-sinikassort?	etla-hulyè?	what is his name?
Tans-ey-sini-kassu- yŭn ?	ey-la-hunlye? -	what is your name?
Tanna-si-te-kateg-		how do you call this?
oma?		-
Init. I. sounded as ce.		
I-ā-pit		
Kah-nup-ātè-i-a-pit -	nakith	he has an eye on one side.

ė

CHEPEWYAN VOCABULARY.

	·		
sh.	Cree, of Carlton House.	Chepewyan, of Athabasca.	English.
	Tann-ikè? Tann-ikh-tè-an ? Istè-kwa-nan; ustě- kwan; mistekwan.	etla-djah ? edthi	what is the matter? what is the matter with you? the head.
coming.	Int'istèkwanan Nistè-kwan - Uta-pètchè-itotè ! Nètè-itotè ! - Tanti-wy-i-tukh-tè- an ?	zedthi-cy-a zedthi è-o-kŭ-si! è-o-kŭ-si-nēk-iltkh! etla-se-nek-ăltkh? -	my head aches. my head. come hither ! go there or thither ! where are you going ?
	тсн.		
	Tchakkatinow - Tchi-kè-kum - Tchi-tchè -	shethi-azè shith 'tinnila-theyl lé (man's toe).	a knoll ; small hill. a wart. a finger.
beaver lodge. beaver lodge. t	Tchi-ka-ĕgan - Tchi-ka-ĕgan-akhtik Tcheŭk-sa-ĕgan -	thell; kong-kwi - thell-tchinnè klell-thelth; thléh- kon.	a hatchet. a hatchet helve. a gun flint.
	Tchè-măn Tchi-pai (Ppa-ŭkk) Tchi-pai-ŭktim	tsi; alle ethi-a	a canoe. a dead body; the deceased. you dead dog! an oppro- brious epithet.
t.	Tchi-pai-ŭkk (dance of the dead).	nè-èlkai	Aurora borealis.
knowing.	Petāpan O-wanni-wagan - Tchis-ā-wan	yel-kon pernatal	dawn of day. dusk of the evening. a hash, or haggis.
fir (Abies bal-	Tchis-kè-pi-son Tchis-ta-bà-sun	dza-thulth pan-neyla; luneylè-	a garter. a button; an anchor.
Anas boschas). k.	Tchista-ka-wè-sew Tchista-ka-nan-wi-	ther-onna el-karrè (pine-leaf	a tork. a wasp. a teal (Anas discors).
se. the speaker's	ship. Tehistèm-ow Tebimm-ashěn	<i>duck</i>) sel-tu-yè ned-tu-a	tobaceo. it is short.
eye ache by a	Tchuk-tchuk-athu ; tchuk-tchuk-ai-u.	tadzon-zellè	a blackbird (Scolephagus).
ame ? name ?	Tchika-wa-sis Tchi-kima Tchi-kima-numma?	yazè (few) ta-tu-ahaddè ta-tu?	not many. true, truly; verily.
call this?	Tchi-ka-ka-win 'Tchist! tchistè! Tchŭppasis	vicen i	close to the shore. hist! listen! look! below; underneath.
on one side	N'tchā-kā-pā-huk · N'tchā-kā-pi-chi-nin	dzè-noy-înkê	he poked it (a finger or stick) into my eye. it has run into my eye (a

ł

VOCABULARIES OF NORTH AMERICA.

Cree, of Carlton House.	Chepewyan, of A	thabasca.	English.
Tchèp-wow	kai-intchuthè	•	it tapers.
Tchès-kwä; tchès- kwa-nitta!	karrè! -	• •	wait! wait a little!
Tchitchei-mi-kŭskwċ-	pè-'kunne-nel	tu-yè	he or it has short nails.
su. Tchika-ai-gè-u -	thelth-ta-nai-i thelth.	lkh-	he hews with a hatchet.
Ne-tchika-iggan -	thelth-ta-nai-i	lkh-thell	I hew with a hatchet.
Tchi-kè-si-sè-u -	dzerè-hai-èllè	-	he plays at draughts.
Tchi-kè-si-sè-a-wukk			they play at draughts.
Tchi-kwa-ha-mè-u -	belekh-hered-	ye -	he crumbles the leaves
			(rubs them to powder).
Ingā-tchi-kwa-hain -		• •	I will crumble the leaves.
Tcbi-pè-tükk-wow	tell-klukk -	• •	it is light blue.
Ni-ghe-tchi-pusti-hà-u	thilk-tas -	• •	I put it with my arrow.
Wa-was-ki-shu -	tse-thil -	• •	the wapiti.
Oya-peyu-mus-tus -	ettirre-ya-ne -		bison bull.
Nosia-mustus	ettirre-su-ta-h	a -	bison cow.
Wā-pis-tànn	tha -	-	a marten.
Si-kāk	nult-si-ai -	-	a skunk.
Si-ku-sew; sigus -	del-kathlei	-	an ermine.
Winusk	tel-leh	• •	a marmot or spermophile.
Winustè-key		•	a Quebec marmot.
Wapusk	sass-dei-gai -	-	white bear.
Apek-ku-sis	tlunne -	• •	a mouse.
Shi-shi-pise	cigarre -	-	a teal.
Key-ask; kai-ask -	bess-gai-e -	-	a gull.
	Kallei -	-	a plover.
	Bekh-hulla -	-	Salmo mackenzii.
Okkau; uka	ettchu-e -	-	Dore.
Miki-sew	ded-donne-tch	ú -	an eagle.
Ahà-sew	dadsang -	•	an American crow.
Ottoni-bis	the-tchuthe -	-	Coregonus artedi (Tullibee)
Namay-pith	till-tulei -	-	Catastomus.
O-wi-pi-tchi-sis -	thlu-dathe -	-	Hiodon.
Nipe; nipi	tu; to	-	water.
The-kwus-kwun -		-	it is cloudy.
Kŭsku-wŭnŭsk-	kothé	-	clouds.
Kishi-kau; kis-gau;		-	day; day-light.
wa-pan.	hambi		morning
Ki-ki-ship	taindágani		noon
Apikh-tow-kishi-kau (middle-day)	tsindessai	-	noon.
Pakkisimu (oun-cot) -	tchilsin -		evening.
Tinnis-kak . tinniskau			night.
Mistiko-tohomon	tetsin-tsi -	.	a boat.
Wini-nègh	tu-tcho		the sea.
Thaka-stimmun-sigen	tsini-ball -		a sail.
Paska-stammun-algan.	telourthe-vaze		a pistol.
Kitchè-kuman	bèss-tcho -	-	a sword (big knife).

1. 1. S. . . .

DOG-RIB VOCABULARY.

little !

lish.

hort nails.

a hatchet.

hatchet. raughts. draughts. s the leaves n to powder). le the leaves. ne. my arrow.

r spermophile. armot.

enzii.

n crow. rtedi (Tullibce)

ght.

knife).

Cree, of Carlton House.	Chepewyan, of Athabasca	English.	
Tappis-kā-gan Mokasin ; muskesin - Tippiskā-wi 'peshim - Kesik ; kishik Pinasi-wuk Wa-waskhsta-punu ; owā-sāmusk. Kunu ; konă Miskwumi Piki-sè-u (<i>it is foggy</i>)	kothi-ghirrè ke eltsi yaha edihi tsinago-thethi- yath ti-enn etzil Hothin Nahalgi	a handkerchief. a shoe. the moon. the sky. thunder. lightning. snow. ice. fog. frost. thaw.	
Utin; thow-tin Atchimow Ni-ku-mun Mitzu Wappamow	niltsi yalthi Netghin tch`li etethi Urcltha-nelsi Su-sinnè	wind. to speak. to sing. a song. to eat. to see. to hear from you. a great happiness.	

The following words of Dog-rib were collected by myself at Fort Confidence. The want of a good interpreter caused me to discontinue the formation of a vocabulary of this dialect.

Dog-Rib	Vocabulary
---------	------------

English.	Dog-Rib, of Fort Confidence.	English.	Dog-Rib, of Fort Confidence.
A kettle	tillè.	A blanket -	zidda (tzud-di-e,
Little ditto -	tillè-yazè.	Indian hose -	thelth.
Fire wood -	sus.	The beard	tarra.
Gunpowder -	tel-kithé-ka.	A crooked knife	bèss-ha.
Shot		A knife	bèss.
Shot-pouch -	tel-ketha.	A knife sheath -	bess-thè.
Ball	tel-kethi-'tcho.	A fork	pakwa.
No meat ! -	par-ulla!	Snow	tzill ; tchill.
Dried ribs of rein-	átcharna; et-	Smoke	thlet.
[•] deer.	chanka.	A stone	thai.
Water	to.	A brisket	ana-rāne; ei yid-
A tin pan -	thai.	The shoulder -	da.
A coat, or capot	i.		ak-kànna.

VOCABULARIES OF NORTH AMERICA.

English.	Dog-Rib, of Fort Confidence.	English.	Dog-Rib, of Fort Confidence.
Leg bone or knuckle.	ak-kai-tchinna.	A file Pole for hanging	kokètha. telle-kaiza.
A firebrand -	halai-kun.	a kettle upon.	
A tent	nepàlle.	Buttons	pai-illa.
Tent poles -	thai-è.	Mittens	gis.
Transverse poles	tanè-ai.	The head	ta.
to hang meat		The nose	tinnetze.
upon.		The knee	et-thětha.
Tent door	ku-latche.	An encampment	zutès.
Leathern door for tent.	thidai-nepàlle.	The encampment is distant.	in-tu-è-zutès.
Dressed leathern	tel.	The encampment	thi-si-tè-zutès.
blanket.		is near.	
A spark from the	kantida.	A warm woollen	kow-i-tchitha.
fire.		collar; a com-	
Rein-deer tongue	et-thu.	forter.	1.41.5
Deer-skin hose -	et-thidda.	One	inthiare.
Breeches	tillai-1.	1wo	nakhke.
Deer nead -	et-thi.	Inree	kntarre.
A shoe	Ku.	Four	unge.
Cloth worn by	tnan.	rive	zazuniarre.
men round the		Six	elkaunarre.
midale.		Seven	ntniazintinge.
A bag = -	naitche; kleithe.	Light -	alkatinge.
A natchet	theith.	Nine	ntnia-otta.
A spoon	thius, or slus.	1en	nthia-una.

The vocabularies which follow were made by gentlemen whose system of orthography varies more or less from that adopted in the preceding pages. The dialects of the Dog-ribs who resort to Great Bear Lake, and of those who hunt on Marten Lake and visit Fort Simpson, differ little when spoken, and offer no difficulty to an interpreter who is acquainted with either; but many of the words have a very different aspect when written in English characters; and these tables may serve to illustrate a remark made in a preceding page respecting the difficulty which an English ear experiences in apprehending the sounds of the Tinnè languages. The Kutchin words collected by Mr. M'Murray, though not numerous, show a close affinity between the language spoken by that people and the

DOG-RIB VOCABULARY.

Tinnè, and will perhaps be considered as a proof of the common origin of the Tinnè and the Kolush tribes down to the 54th parallel of latitude.

English.	Dog.Rib.	English.	Dog-Rib.
English. Head - Neck - Tongue - Eyes - Ears - Nose - Cheeks, chin Shoulders - Thighs - Brisket - Brisket - Rump - Belly - Hands - Fingers - Nails - Teeth - Brain - Liver - Heart - Blood - Skull - Entrails - Udder; milk Butter - Flour - Sugar -	Dog.Rib. - bet-thi. - bdi-korh. - eth-thadu. - mendi. - bed-ze-gai. - mi-gou. - mi-ta. - ai-kon-nai. - ed-zaddai. - a-ethin. - etchin-nai. - be-tchuki. - mila. - mila-tchinnai. - mila-konnai. - bighu. - bet-the-ghu. - et-thut. - ed-zai. - et-tillai. - et-tillai. - et-tillai. - et-tuzai. - edgiddai-thlissai - hatai-kotliss. - suka.	English. Go! Come! Take! Cut! Large Srnall Large Short Short Short Short Short Short Short Y ! Laugh! Speak! or talk ! How many ? Heavy - Light High Low Good Bad Fat Eat!	Dog-Rib. aga ! ya-kusi ! hi-tcho ! bekan-nèthu ! si-nekai. no-sai. nai-tcha. ti-ula. nundeth. nundeth.helai. nitha. whā-yai. azel ! mena-thi-ukla ! betha ! tanna-itai ? addow-adlis ? tai-it. naisou. tlenai. tlaika. tlaika. tlaika.
Sugar - Tea Pepper - Medicine - Paper -	 suka. suka.tu (sugar water). tenni-tsi. na-diddu. eddithi. 	Eat! Drink! Smoke! Sleep! Give! Tell!	ath-uluston ! ath-uluston ! ustud ! notai ! mi-ne-kai ! adin-dai !

The above vocabulary was formed, I believe, at Fort Simpson, by one of the Hudson's Bay officers for his own use; but, having forgotten to note the circumstances under which it was drawn up, I can give no further information regarding it.

g-Rib, of Fort Confidence. tha. •kaiza.

lla.

etze. Ičtha. s. 1-è-zutès.

i-tè-zutès.

-i-tchitha.

larè. hkè. urre. ge. inlarrè. tharrè. azintinge. tingè. la-otta. la-otta.

y gentleless from ts of the of those on, differ iterpreter ords have aracters; made in an Enls of the by Mr. e affinity and the

VOCABULARIES OF NORTH AMERICA.

A VOCABULARY OF FORT SIMPSON DOG-RIB, BY MR. O'BRIAN, OF THE HUDSON'S BAY COMPANY.

Dog-Rib.7		English.	Dog-Rib.		English.
Edza-zinnè		Tetrao umbellus.	Noga -	-	wolverene.
Tih	-	Tetrao canadensis.	Kling -		dog.
Bet-theu -	-	owl.	Tzus -		wood fire.
Thlu-ai -	-	Coregonus albus.	Tai-tchin -	-	trees.
Samba -	-	trout.	Tzu -	-	pine-tree.
Kazè -	-	salmon.	Ki	-	birch.
Tsai-teu -	-	Back's gravling.	Sinnai -		I.
Tai-tellai -		Catasiomus.	Tlinnai -	-	thou.
Klogai -	-	squirrel.	Ottinai -	-	he.
Emmu-i-u-ai	-	Columba migra-	Ige		it.
		toria.	Edetata -	-	ves.
Khun -	-	fire.	Helai : Odelis	-	no.
Tu	-	water.	Id-zeunai -	-	to-day.
Tchon -	-	rain.	Kambai -	-	to-morrow.
Yah	-	snow.	Zeunai -	-	day.
Ten	-	ice.	Tethi -	-	night.
Sa	-	sun.	Yakh-kai -		winter.
Tethi-sa -	-	moon.	Klukai -	-	spring
Thi-u -	-	stars.	Senai -	-	summer.
Kose -	-	clouds.	Ai-tonkai -		autumn.
E-tu-ai -	_	girl.	Tai-chin-ala	-	boat.
Ab	_	snow-shoes.	Ki-ala -		canoe.
Kai	-	shoes.	Tami -	-	net.
Whoghi -	-	suare.	Tau-ai-on		full.
Thai -	_ 1	sinew.	Tu-tai -		empty
Do	-	now.	Tlon -		plenty.
Ye-won -	- 1	then.	Hulai -		none.
Tau-dezzei	-	half.	Tzuddi-è	-	a blanket.
Mal-lionai	_	rings.	Tai-si-ai -	-	a shirt.
Hai-ai -	-	trowsers.	Ed.geid-dai	-	a powder-horn.
Memba-ulai	_	waistcoat.	Mad-deli	-	buttons.
Tsi	_ 1	vermilion.	Thai-on-tithei		thread.
Sat-sn-wai		wire snare.	Et-thai-ai	-	scissors.
Sas		black bear.	Meni-di-e-dai		looking-glass.
Sa-tai-kuzè	_	brown bear.	Ai-tchusai	-	beads.
Tsa	_	beaver.	Ai-tai -	-	ice chisel.
Tsa-thu-ai		castoreum.	Bai-huch -		crooked knife
Tai-tchesi		mink.	Bai-chin-ai-i	_	clasn.
Tzin		musk-rat.	Bed-do-ai-du	-	not.
Tè.ki		wolf	Tha		pan.
TC-VI a	-		- 114	-	Lucare.

"MAUVAIS MONDE" VOCABULARY.

The following vocabulary of the language of a tribe dwelling near the sources of the River of the Mountains, and known to the voyagers by the name of "Mauvais Monde," and of the Dog-rib dialect, was drawn up by Mr. O'Brian, of the Hudson's Bay Company's service.

Mauvais	Monde	e.		Dog-	Rib, or S	lave.		English.
Thèlgai		•		thli-e		-		one.
Olki-e		•	-	olki-e	-	-	-	two.
Ta-dette	-	-	-	ti-e -	-	-	-	three.
Tinghi	•	•	-	tinghè	•	-	-	four.
Sazelli	-	-	-	sazelli ; la	akithe.	the ha	nd	five.
Et-seu-ti	•	-	-	et-seu-ti		-	-	six.
Thlad za-di-	e	-	-	han-die	-	-	-	seven.
Et-zan-di-e		-	-	et-zan-di	-e -	-	-	eight.
Et-thlei-hu-	lai	-	-	ethli-e-h	oulai	-	-	nine.
Ken-na-tai	-	•	-	o-nai-u-n	ion	-	-	ten.
El-lai-zai *		-	-	tel-kithi-	kun	-	-	gunpowder.
Bai-ka	•	-	-	tel-kithi-	teho	-	-	ball.
Ni-tai-ton	-	-	-	thai-thi	-	-	-	shot.
Ai-tai-kai	-	-		sel-tu-e	-	-	-	tobacco.
Et-ton-nai	-	-	-	tel-kithè	-	-	-	gun.
E-tha-thai-	m	-	-	hai-ko	-	-	-	gun-flint.
Utha -		-		tiu-ni-e		-	-	kettle.
Thei -	-	-	-	thei -	-	-		axe.
Ai-tchut	-	-	-	ai-tchut	-	-	-	awl.
Bèss -	-	-		bèss -	-	-	-	knife.
Ta-chill-ai		-	-	et-lev-na	ui -	-	-	cloth (strouds).
Kestu-ai		-	-	ai -		-	-	coat (capot).
Theth	-	-	-	theth -	-	-	-	leggings : also a belt.
				Edgidda	i -	-	-	powder-horn.
Kothegettai		, ,	-	ko-the-g	at -	-	-	handkerchief.
Set-tsa-tai	-	-		tsa -	-	-	-	bonnet-can.
Hai -		-		kun -		-	-	fire-steel.
A_tai-kai-te	nnev		-	seltu-ter	ne	-		tobacco-box.
Ta-ti-e	-		-	ta-ti -		-	-	needle
Thairka	-		_	ko-kasse		-	-	file
Etabaiai		-	_	haith-lai	ka -	-	-	seissors
E-kodzi	_	_	_	kud-dai		-	_	gup-worm
Ai-kathai-te		-	_	sa-kotha	i_tai	-	_	garters
Kloi si		-		sa-katha		-	_	gray haar
II thai	-	-	-	no-githi		-		for
Ustaidad	-	-	-	no-gitin	-	-	-	luny
Ustainge	-	-	-	no-thei		-	-	martan
Kocho	-	-	-	nom ha		-	-	ottor
Wallon	-		-	ton di o	ai =	-	-	mala moore deer
101101	•	-	-	l ren-ui-e	-	-	-	male moose-deer.

* This word is used also by the Beaver and Thekannè Indians.

BRIAN, OF

erene. fire. tree.

ıy. orrow.

er.

g aer. mn. e. y. y.

nket. rt. vder-horn. ns. d. rs. ng-glass. . isel.

ed knife.

VOCABULARIES OF NORTH AMERICA.

Mauvais Monde.	Dog-Rib,	or Slave.	English.
Intsei Wod-su-tchu - Wod-su-mon-bed-sai Kāg-kalai Ogha-tchai Ea-sai U-ta-dja Niton Setsa-on Te-sonnai Klu-chu-i-nai - Sunbaddei	 teu-di-etse bed-su bed-su-tsi kam'ba ogha - tai-tonna-tel uti - monalla tchikwe tchiltawe bai-tchinai dsheth 	10	female moose-deer. male rein-deer. female rein-deer. ptarunigan. goose. eagle. pike-fish. white man. woman. boy. sled. mittens.

A VOCABULARY OF CHEPEWYAN AND DOG-RIB WORDS.

The Chepewyan was taken down from the mouth of the interpreter at Great Slave Lake. The Dog-rib from that of the female interpreter (Nanette) at Fort Simpson. The whole in 1844. J. H. LEFROY.

Toronto, March, 1850.

English.Chepewyan.Dog-R b.Yes!No!he-liA mandennetarqui-ietaltitle girltaboytittle boytarqui-thetenai-utarqui-the-tenai-utarqui-the-tenai-utarqui-the-tenai-utarqui-the-tartab-<
Yes!e-h!- $No!$ he-li- A mandenne- A womantchel-a-qui. A little girl-tza-qui-ie- A boytchilla-qui-ie A boytchilla-qui-ie A little boytchilla-qui-ie A little boytchilla-qui-te A little boytza-qui-the A tractabtenai-u-azzè.Tothor
No1 - - he-li - - he-li. A man - - denne - - tchel-a-qui. A woman - tza-quie - - tzek-qui. A little girl - ett-er-e-ka - tzek-qui-azzè. A boy - - tchilla-qui-ie - tenai-u. A little boy - tza-qui-the - tenai-u. Fothor - tza-tab - tenai-u.
A man tza-qui-ie tchei-a-qui. A woman tza-qui-ie tzek-qui. A little girl ett-er-e-ka tzek-qui-azzè. A boy tchilla-qui-ie tenai-u-azzè. Father tza-qui-the tenai-u-azzè.
A wohan
A little girl ett-er-e-ka tzek-qui-azzè. A boy tchilla-qui-ie - tenai-u. A little boy tza-qui-the tenai-u-azzè. Fothor tza-tab
A boy tchilla-qui-ie tcnai-u. A little boy tza-qui-the tcnai-u-azzè.
A little boy tza-qui-the tenai-u-azzè.
Father
\mathbf{T} acher \mathbf{T} = \mathbf{T} (\mathbf{Z}) = \mathbf{U}
Mother en-nè en-nè.
Brother, elder tzoon-noi tzoon-noi
vounger - tzachilli
Sister elder
vouncer - starta
", younger - sa-tez-zan, or tza-tazze.
To-morrow koume.
<u>x</u> esterday ou-ah-ta-tzenke - ye-hho-a.
Tobacco tza-twe tza-twe.
A knife pa-as pa-as.
A gun thel-ki-the

' Broad, ' nasal, ' guttural, " nasal and guttural.

CHEPEWYAN AND DOG-RIB VOCABULARIES.

4.

English.	
moose-deer.	
ein-deer.	
rein-deer.	
gan.	
sh.	
man.	
1.	
9	

WORDS.

h of the interm that of the The whole in I. LEFROY.



E	nglis	h.		Chipewy	an.		Dog-Rib.
An axe Gunpowe Ball Air Fire Water Earth A fish A dog A fox A buffalo Rein-dec A moose Snow sho A sled A kettle Evening	lder 	n.		than-thye thel-ki-the-co thel-ke-the-cl nutze - kkon - tto - clou-a - čling - no-ki-ki-the et-cherre - eet-than - den-nee - ah-he - beth-chin-ne tille - ectzson - kompe -	an.		qua-a-qui. thi-ke-c-cond. the-ke-e-chou. e-at-ti-ge. kkon. two. ko-ecela. clou-a. clou-a. clou-a. cle. e-et-tha-thà. et-cherri. , ettzae. et-thun (ettzae, f.). denne-a. a-c. ba-chen-ne. than-ne. eya-kka-ezza. so-tho
Colours -	- bl	ack	-	tel-zonne	-	-	ta-zun.
**	re	hite d	:	tel-ka-ye tel kkosse	:	-	tel-ka. et-tel-kkos.
>> 1>	gr bl	een ue	:	ta-ečloze not distinguis black.	hed fr	om	ta-eck-cles. ten-è-ëlè.
// · · · · · · · · · · · · · · · · · ·	ye	llow	-	tel-thoi -	-	-	tel-thoi.
The moo	-	•	-	ssa	-	-	ssa.
A star	-			thun -			thun.
(The Gre	at I	Bear)	_	(va-ee-telli)	-	-	(va-tha).

English.	Dog-Rib.	English.	Dog-Rib.
Snow - Ice - Numerals	 yya. t-than. en-člai. nà-kka. tta-rgha. tting. sa-soo-la. ut-ke-ttai. kkosing-ting. etzenting. kka-hooli. ho-nanna. nou-nanna. 	Good-Bad-Beautiful-Ugly-Large-Small-Heavy-Light-Dark-Bright-	naa-zo. naa-zo-heli; dzoun-de. bur-a-oonde; tzoo-na-e-ti. pa-chi-ri. natza-konde; in- cha. tzoo-ta. net-ta; hinka. hin-ka-he-li. tel-zen. atz-za.

VOL. II.

D D

VOCABULARIES OF NORTH AMERICA.

Eng	lish.		Dog-Rib.	English.	Dog-Rib.
Low High Hard Soft New Old Denr (be	eloved)		ne-otzin-ik. tan-ne-e-tha. taa-y-eet. taa-yeet-heli. e-e-yes-e. e-e-ranna. The word un- known to the language.*	When-Which-What ?-To me-To him-To you-To us-Idon'tunder-	kkonde. w co. et-cloy ? tzen-ez-ctze. ne-ghon-em-etze. ne-nin-etze. c-e-cla-toon-nim- etze. nd-ad-'hear-des-
Wise	-	•	koo-rac-yon.	stand.	tha-helili.
Foolish	-	•	nà-a-ghal.	Idon't speak Cree	(same sentence
Strong	-	•	na-tz-ap.		taken down).
Weak	-	-	pa-a-ttha-to- rghelli.	I won't give it you.	na 'rha tchou-heli.
Right	•	-	nochnesse.	I will give it to	na 'rha ochou-
Left -	-	-	intzesse.	you.	eze.
My frien My com	nd panion	•	tza-teleg-ga.	What shall I give	than-etcha-na-rha- bela-hūze ?
Head	-	-	tzat-the.	Take care -	ča-re.
Eves	-		tzen-nhae.	Make haste -	aga-annitè.
Nose	-		tze-etze.	Get out (va t'en)	or-rhink-là.
Mouth	-		tze-tha.	Where is it? -	ye-in-kon-ecla?
Ears	-	-	setz-r-rgha.	Carry this for me	sse-ragh-di-ach.
Hair	-	-	setz-thè-rgha.	Don't touch that	perrone-te-sonna.
Tongue	-	•	tze-tthou.	What do you	na-nu-at-cloy?
Teeth	-	-	tze-w-who?	want?	
Neck	-	•	tze-e-e-cottle.	What do you	etcha-nette-ousa-
Arm	-	•	tze-int-chinne.	want for this?	nousa-ou-sinne?
Hand	-	•	ssa-la.	Give me a piece	tza-twc-tza-gan-
Foot	-	•	tzè-ka.	of tobacco.	a-two.
Legs		-	tze-thunna.	I have no tobacco	tza-twe-ta-oo-
Canoe pa	addle	-	ola; tho.	** ** **	twe.
Here	-	-	a-jahn.	Hold this	Ou-net-ton.
Where?	-		a-c-yà. djahn-tin ?	Whose is this?	me-etze-hande?

* I endeavoured to put this intelligibly to Nanette, by supposing such an expression as ma chère femme, ma chère fille. When at length she understood it, her reply was (with great emphasis) : "I' dit jamais ça. I' dit ma femme, ma fille."

Dog-Rib.

nde.

oy ? -ez-etze. hon-em-etze. in-etze. cla-toon-nimze. nd-'hear-desna-helili. ne sentence aken down). rha tchou-heli.

'rha ochourze. n-etcha-na-rhaocla-bāze? re. "a-annitē. -in-kon-ecla? g-ragh-di-ach. rrone-te-sonna. -nu-ät-cloy?

cha-nette-ousanousa-ou-sinne? a-twe-tza-gana-two. a-twe-ta-ootwe. u-net-ton. e-etze-hande?

posing such an agth she unders ça. I' dit ma

EXPLANATION OF PLATES I. & II.

No. 1.

Plates I. and II. represent impressions of fossil leaves found in a bed of potter's clay, belonging to a tertiary lignite deposit near the mouth of Bear Lake River, described in vol. i. p. 190. of the narrative. The leaves must have been very numerous, and were evidently deposited quietly from water turbid with fine potter's clay, which forms the matrix. By the spontaneous burning of the adjacent seams of lignite, the fossiliferous layer has been subjected to heat of varying intensity, so that some portions are semi-vitrified and rendered hard enough to resist a file, while the greater part is in the condition of moderately baked porcelain biscuit, and in some few specimens the clay is but slightly altered.

The impressions only, and none of the substance of the leaves remain; and owing to fusion of the leaves at their margins from pressure, and the cracking of the clay matrix from heat, none of the impressions of the larger leaves are perfect in their outlines, though portions of the surface are very delicately rendered so that the minute nervation is distinctly shewn, and the existence of pubescence may be made out.

Table 1. fig. 2. is a representation of the impression of a twig which has the character of *Taxites acicularis*, (Brongniart Prodr. 108, and "Descript. Geol. des Environs de Paris, p. 362., t. ii. f. 13." *Taxites foliis subdistichis*, *linearibus obtusis*). The leaf is scarcely half the length of that of *Taxus baccata*, and is decidedly smaller than

404 EXPLANATION OF PLATES I. AND II.

that of T. canadensis, being generally about 0.44 inch in length and about onc-fifth part as wide ; or, more precisely, the breadth of the lamina is 0.09 inch, which is somewhat broader in proportion than the leaf of the common yew. The outline of the leaf is linear with a slightly lanceolate narrowing near the apex, which is rounded without any perceptible projection of the mid-rib. Yet though the matrix has rendered the most delicate impressions of the surface, an actually existing minute projection of the midrib may have been obscured, owing to the convexity of the lamina; for on making casts of the common and American yew leaves in Paris plaster, the acute apices of the mid-ribs were not distinctly shewn. The footstalk is as short or shorter than in the common yew, and appears to have had the same kind of half twist which gives the distichous direction to the leaves. The surface of the lamina is slightly convex, with about as much recurvature of the edges as in the Canadian yew, and there is a regular fine undulation, or obtuse transverse wrinkling, which is perceptible in all the impressions when they are examined with a lens; but, except the straight, tapering, prominent mid-rib, there are no veins.

Owing to the distichous attachment of the leaves, the impressions of the adnate scales of the bark to which the footstalks are jointed are oblique, and the proper form of the scales is not easily determined. They do not appear, however, to have differed greatly from those of the common yew. The elevated triangular areas shewn in the stem of the figure were depressions in the plant between adjacent scales of the bark. In a few specimens the tops of the twigs are shewn to have had an arrangement similar to that of the yew. Buds are numerous in the axils of the leaves of the annotinous spray of yew, and a few impressed hollows in the fossils may have been caused by such bodies, but they are comparatively rare. More numerous small detached bodies in the matrix may have been produced by the berries or nuts of this plant. Five or six of the twigs

EXPLANATION OF PLATES I. AND II. 405

terminate inferiorly in ovate or irregularly oval expansions, which are such as a cluster of buds situated there would produce; in only one specimen is there any appearance of the branching of a spray, and in that the seeming branch may be merely the impression of one twig crossing another.

Out of upwards of fifty impressions of twigs of this Taxites, eight or ten have small round depressions on some of their leaves, disposed sometimes in a pretty regular row on each side of the mid-rib, but more often they are irregular both in distribution and size. Now and then one or more of the dots approaches nearer the margin of the leaf than the others, sometimes they are seated on the mid-rib, and occasionally one dot encroaches on another. Most of the dots have a little pit in the centre, and their circumferences more deeply impressed than the area, which is often convex, though not raised above the impression of the lamina. They must, therefore, have projected above the surface of the leaf, whose cast is all that remains. These dots bear some resemblance to the fructification of a fern; but on exhibiting the casts to Mr. Brown, he at once remarked the dissimilarity of the dots to sori, in their having no perceptible connection with veins, and in the appearance of a membranous expansion from the epidermis covering them, which his practised eye detected. On examining twigs of the Canadian and common yews I observed many sphacelated dots raised more or less above the surface of the leaves, which would make impressions very similar to those of the fossil. The dots in the recent plant occur more commonly on the under surface of the leaf than on the upper one, and are generally circular, though sometimes irregular. They are covered by the epidermis, which in the larger dots is always ruptured in the centre. I have not been able to discover their precise nature; they may possibly be caused by insects, or perhaps by the rupture of terebinthaceous collections. They do not appear to be fungi; and when the epidermis is removed, the minute cavity is found to be lined with indurated **DD** 3

inch in recisely, mewhat on yew. nceolate out any ugh the is of the the midty of the merican mid-ribs short or have had listichous amina is re of the ular fine h is perexamined rominent

aves, the which the r form of t appear, e common e stem of adjacent ps of the ar to that leaves of d hollows dies, but small deduced by the twigs
parenchyma, which, under the microscope, exhibits cells similar to those of the rest of the leaf.

Figures 2 and 3 are magnified, but the lithograph has failed in truly representing the delicate undulations of the lamina shewn in the fossil casts.

No. 2.

There are also eight or ten impressions of yew-like twigs differing from the preceding, but none of them sufficiently perfect to give precise characters. The leaves are narrower than in the former species, though generally of the same length: there is, however, more variety in this respect, the same plant containing leaves of very different lengths. They are narrowly lanceolate, tapering gradually from the base to the tip, which is acute. Instead of terminating in rounded lobes at the insertion of the footstalk, the leaf appears to be decurrent, with the mid-rib continued into the decurrent portion. The mid-rib is slender, but distinctly impressed throughout the whole length of the leaf; and the surface is less evidently wrinkled transversely than in the first species. The leaves are also more crowded and more erect, with less of the distichous appearance, and a 3 arrangement may be made out. Some of the sprigs are branched like those of the common yew. This plant possesses the characters of Taxites phlegethonteus of Unger (Plant. fossil. p. 390.). Round dots also exist in the impressions of this species, disposed as irregularly as in the preceding one, and some are visible on the decurrent base of the leaf. Part of these dots had in the original an elevated margin, a convex disk, not so high as the margin, and a pointed central point; others have left an uniformly concave impression.

Scattered through the matrix, and often in the close vicinity of the *Taxites* twigs, but only in one instance connected with them, there are impressions of a minute fruit, such as would be made by the nut-like seed of yew, deprived of its outer investment and of the coloured pulpy

calix. None of these impressions are, however, above onefourth of the size of a nearly ripe seed of the common yew. They are compressed, ovate, and acute, without a prominent point, and all of them shew a faint furrow descending on each side from the apex, more acute than the impression which would be made by the slight ridges on that part of the yew seed, which is only very slightly compressed. Several shew indications of an investment at the base, and in one impression the soft integument of the fruit scems to have been pressed aside so as to allow a cast to have been made of the nut within it. This pulp enveloped the nut entirely; or, if an opening like that of the calix of a yew-berry existed, it must have been obliterated by pressure. The solitary fruit attached to a twig of No. 1, is inclined downwards on a very short fruitstalk.

No. 3.

Some imperfect casts also exist of a plant, most probably belonging to the family *Ericaceæ*, and approaching nearly to Vaccinium. Some of the casts shew a five-parted, or five-leaved calix, composed of thick ovate acuminate sepals meeting at the apex. In two others a berry scems to have been crushed, leaving a flat floor of minute, very numerous seeds, partially covered with integument or There are also impressions which may have been pulp. produced by an urceolate corolla. These flowers grew on short fruitstalks, springing apparently solitarily from the axils of the leaves. As far as the form of the leaves can be made out, they are linear lanceolate, narrow, but scarcely acute at the point, with a concave surface and a not very prominent mid-rib. The leaves are approximated, apparently not in any regular order, applied to the stem at their bases, and curving outwards at the tips with a sigmoid flexure. They are rather more than a quarter of an inch long, and the height of the swollen calix or corolla is about as much.

рр 4

ts cells

iph has s of the

ke twigs ficiently arrower he same pect, the s. They e base to rounded pears to lecurrent mpressed urface is the first re erect, arrangebranched esses the nt. fossil. s of this one, and Part of a convex central ression. he close instance n minute of yew, ed pulpy

No. 4.

Plate II. is a representation of a segment of an impression of the upper surface of a palmately veined leaf, magnified to rather more than twice its linear dimensions. Though it bears a general resemblance in its nervation, and in the areolæ formed by the minor reticulating veins, to a leaf of the Maple, it differs at least as a species from the American maples with which it has been compared. Many small circular depressions of different sizes are irregularly distributed over the surface of the leaf; the more perfect of them are pitted by twenty or more minute points visible by the aid of a lens; and in some the central point is larger than the others, producing the appearance of an umbo, as in the dots of the Taxites figured These depressions were of course produced by in Plate I. bodies rising above the surface of the leaf and rough with little points; their unequal dispersion on the leaf, much of whose surface was smooth, is against their having been produced by hairy glands, and they were most probably moulded on fungi growing on the leaf. There are also some smaller and deeper depressions, most frequent towards the upper part of the leaf, but considerably less numerous than the larger shallow ones. Two other fragments of impressions, seemingly of the same kind of leaf, have footstalks not complete, but exceeding an inch in length. One of these, representing the upper surface of the leaf, has a few circular depressions of both kinds; on the other, which is an impression of the under surface of the leaf, there are no depressions.

None of the impressions are so complete as to give the whole outline of the leaf. The base runs at right angles with the footstalk, and is entire for nearly an inch, beyond which it is rounded off and crenated by almost semicircular, minutely apiculated teeth, separated from one another by very acute sinuses. The leaf appears at first

sight to be three-ribbed, the central nerve or mid-rib being straight, and the one on each side curved, so that if prolonged they would meet the central one at the point were the impression complete. There are, however, in fact seven nerves springing together from the footstalk, the lower pair having less prominence and more the character of the secondary veins; the next pair are but little smaller than the three principal ones. On the upper surface of the lamina the main voins were concave, while the impressions of their under surfaces are sharp and rectangular. The secondary veins and ultimate reticulations were prominent on both sides of the leaf. The minute areola are plain and smooth, and there is no indication of any pubescence in the axils of the veins.

Impressions of leaves from the coal beds of the Raton, in lat. 37¹/₂° N., long. 104¹/₂° W., (vide New Mexico by Emery, Abert, Cooke, and Johnston, p. 522., plate), resemble this species, but are too imperfect for identification.

No. 5.

An impression of a smaller leaf than the preceding is so like it in the character of the veins, ultimate reticulations, and general surface, that, but for a little difference in the crenatures of the margin, they might be pronounced without hesitation to be of the same species. The outline is subrotund, transverse at the base near the footstalk, and apparently entire there, crenated more irregularly and with generally smaller teeth than the preceding on the sides; and entire near the tip, which is deficient in the specimen. The diameter of the lamina of this specimen is an inch and a half. It exhibits none of the round dotted depressions, but there are some of the irregular clusters of little pits on different parts of the surface which exist in the specimen figured in Plate II. and in some other casts.

n impresleaf, magmensions. nervation, ing veins, cies from compared. s are irrethe more e minute he central e appeartes figured oduced by ough with eaf, much ving been probably are also nt towards numerous nts of imfootstalks One of has a few , which is ere are no

to give at right 7 an inch, y almost from one s at first

No. 6.

Another palmately-veined leaf differs from the preceding ones in the arcolæ being concave above or puckered from the tightness of the veins, and in its margin being sharply toothed and irregularly crenated. Only one impression of this leaf exists in the collection, and that is imperfect, there not being enough of it to indicate the form of the outline. It does not appear, however, to have been lobed. The äiameter of the lamina is scarcely two inches.

No. 7.

There is still another palmately-veined leaf very different in form from any of the others. It is one-third wider than it is long, and seems to have been rounded at the apex, which is, however, narrower than the widely The base is cut horizontally, and is very rounded sides. The sides are tooth-crenated, the teeth being segentire. ments of circles, and the crenatures acute, but not deep. The footstalk is slender, and the primary veins, none of which are straight, are still more so. They spring five together from the footstalk; the lower pair being smaller than the other three, and the mid-rib largest. They are all branched, and the ultimate reticulations are polygons of very irregular shape. The arcolæ are smooth and flat. The length of this leaf is rather more than an inch.

No. 8.

The fourth palmate leaf, of which there is a definite impression, is subrotund, and about an inch and a half long, with a very entire, or at the most slightly undulated, margin. More than an inch of slender footstalk remains. The base is horizontal, with a minute curvature downwards, or decurrence where it joins the footstalk, then it is widely

rounded into the slightly concave sides, and rounded again towards the point which is deficient. Five principal veins originate from the footstalk, the lower pair, which run near the margin of the lamina, being less conspicuous than the other three. The mid-rib is straight, the veir on each side of it curved in the segment of a circle; so that were the leaf complete, the three would be found to meet at the point, as in No. 4. Numerous curving and forking branches spring from all. The areolæ are of very various sizes, and their surfaces, when viewed with a microscope, are seen to be minutely granular, indicating minute pits in the upper side of the leaf.

No. 9.

Impressions of a penniform leaf resembling that of Morus alba in its general outline and the character of its veins, are pretty numerous. Both surfaces seem to have been quite smooth, there being no evidences of the existence of any pubescence. The central nerve or mid-rib, and the primary lateral ones, are very distinctly impressed in the matrix; and the connecting ones of the second order, which run from one lateral branch to another, are more conspicuous than those of the mulberry. They vary, from being almost straight and parallel to each other, to a greater or less degree of curvature, or even a pointed arch, in the middle of their length, and towards the margin of the leaf are branched and pass gradually into reticulations. The minor veins are much less prominent, and form minute meshes of no uniform shape, but generally oblong, and from four to eight-sided, having a strong resemblance to those of the mulberry. The minute areas are flat and smooth. The lateral veins spring from the mid-rib in pairs, but, except at the base of the lamina, not exactly opposite to each other. From the footstalk two lateral branches spring together with the mid-rib, forming, as in the mulberry, a tripleribbed leaf; but there is a greater fulness of the lamina

the prebuckered in being impresis imperthe form ave been o inches.

very difone-third unded at e widely d is very eing segnot deep. none of ring five g smaller They are polygons and flat.

> inite imalf long, ed, marins. The vards, or s widely

there, and the nervation differs from that of the mulberry in that part, in two or sometimes three veins of smaller size originating also from the mid-rib or footstalk to supply the base of the leaf. Five, or sometimes six veins may, therefore, be said to spring from the top of the footstalk, the lower ones being secondary as to size. In the leaf of the mulberry, these smaller basal veins are branches of the lateral veins. No entire impression exists. The largest fragment must have belonged to a leaf between seven and eight inches long. The general outline seems to have been cordate, with the lobes at the base full and perhaps overlapping, and the apex acute, but not acuminate as in the mulberry. The margin is serrato-dentate above, and simply undulated, or nearly even, at the base. The teeth are generally semi-ovate with a little point, or mammillate as in the mulberry, but are not so closely set. They are somewhat unequal in size, and occasionally denticulate, a larger tooth being notched by a single smaller one. In some specimens the sinuses between the teeth are acute, but more generally they are obtuse. This is one of the most common leaves in the deposit, and one impression of it often succeeds another in layers thinner than common writing paper, and so blended together at the margins of the leaves that the impressions cannot be obtained perfect.

Fig. 1. Plate I. is intended to represent a small fragment of this leaf, drawn of the natural size, but the distinction between the secondary transverse veins and their minor ramifications has not been maintained in the figure as to size, nor are the ultimate veins shown. The teeth are more obtuse in the side of the cast which has been drawn than they appear in the layer which was removed from it, owing to the way in which the matrix has adhered at that part; but it would appear that some of the leaves varied in having more obtuse teeth, if we may judge from two other casts very similar to this one in all other respects. Upwards of an inch of petiole remains in some of the specimens.

No. 10.

There are two good impressions of another cordate leaf, which, instead of the smooth lamina of the preceding, have the surface densely and equally covered with pores invisible to the naked eye, and which may have been produced by a close, stiffish pubescence. The outline of the leaf has been exactly cordate with a short acuminated point, and the base not so full as the preceding, but rather reniform. The point has a perfectly even edge, and the basal lobes are also quite entire; but the rest of the margin, of which only a small part is quite complete, seen s to have been undulated, the projections not amounting to teeth. A straight, tapering mid-rib gives off about seven lateral branches on each side, at an angle of about 45°, not exactly opposite, but so approximated as to form pairs. The lowest pair rise more nearly opposite, but are not so much more conspicuous than the others as to give a triple-ribbed appearance to the leaf. Some of the lateral branches fork near their tips; they are all joined by transverse veins similar to those of the preceding species, and the intervals are filled up by less obvious reticulations. The lower pair of lateral branches send ramifications downwards to the basal lobes stronger than the ordinary connecting veins. This leaf is about five inches in length, and of equal or superior breadth. A small part of the footstalk only remains.

No. 11.

Another impression of a pennately-veined leaf presents a distinctive character in the fine acute reticulating veins, which are prominent in the cast of the under as well as of the upper surface of the leaf. The arcolæ are flat, but traversed also by minute winding prominent veinlets, the ultimate reticulations being very minute. In the general character of the principal veins, this species strongly resembles No. 10. Of the outline, not much can be said,

mulberry of smaller to supply eins may, footstalk, n the leaf ranches of he largest seven and s to have id perhaps minate as ate above, The ase. , or mamset. They enticulate, rone. In are acute, one of the pression of n common nargins of ed perfect. mall fragat the disand their the figure The teeth has been removed as adhered the leaves udge from r respects. f the spe-

as only the lower half of the cast remains. About an inch and a half of broad footstalk is met very acutely by the cordate base of the lamina. The margin is cut by rather large and somewhat remote acute serratures, the first of which is about three quarters of an inch from the footstalk. The under side of each tooth of the serrature slopes gradually down to meet the much shorter upper side of the next tooth, which stands out horizontally. Each tooth is traversed by a vein of the third order, which ends in the acute point. From the fragments we may estimate the length of this leaf at five inches, and the breadth at three.

I forbear describing other fragments, which probably represent the upper end of this leaf, as there are some differences in the surfaces of the casts.

No. 12.

A very sharp cast occurs among the others of the upper part of a leaf, which in the acuteness of the veins and their form resembles Tilia europea, while in its general outline and rugose surface (but not in its margin) it is similar to some of the lower leaves of *Corylus avellana*. The veins meet the mid-rib in pairs, or alternately; the lower ones are nearly straight, the upper pairs are segments of a cordate curve, concave upwards. The veins of the next order pass directly between the branches in a straight line, or with a few anastomoses, and the ultimate reticulations are minute. The lamina is prominent above, from the tightness of the principal and secondary veins, and the cast of the upper surface shows also the prominence of the minute netting, corresponding of course to furrows in the original leaf, so close as to give it a somewhat woolly appearance to the naked eye. The cast of the under surface is convex where that of the upper one is hollow, shows the ultimate ramifications of the veins less distinctly, and is irregularly dotted by punctures visible by aid of a lens, and which probably had their origin in a stiff and scattered pubes-

cence. The axils of the veins do not appear to have been woolly. No punctures are perceptible on the cast of the upper surface. The general contour of the end of the leaf is very obtuse, and as the central tooth which ought to form the apex is broken away, we cannot determine whether it was prolonged into an acuminated point or not. The existing portion of the margin is tooth-crenated, the teeth being comparatively large and obtuse, with a minute point formed by the end of the vein which traverses each of them. A tooth corresponds to every vein that proceeds from the mid-rib, and the sinuses between are wide, shallow curves.

A second cast, which seems to be of the same kind of leaf, shews the apex rounded, without other projections than the wide and not very prominent teeth.

No. 13.

In contact with the apex of this leaf, and partly hidden by it, there is the cast of a slender seven or eight-ribbed fruitstalk with the upper part broken away. Globular depressions are situated alternately on the sides of this fruitstalk and in contact with it. They might have been produced by sessile woolly fruits of the same form with those of *Tilia europea*, and about half the size, or by little tufts of withered flowers on an interrupted spike like that of the male florets of *Castanea vesca*, but each tuft confined to one side of the fruitstalk, and not, as in *Castanea*, verticillate. The exactly globular form of the cavities seems to be more accordant with the casts of a fruit, while their rough interior must have been produced by pubescence or some other inequality of surface.

No. 14.

A rugose leaf differing from No. 12.; has pointed teeth, but the impressions are very imperfect.

VOL. II.

ЕΕ

at an inch by by the by rather a first of the footure slopes er side of Each tooth nds in the imate the at three. probably are some

the upper s and their ral outline similar to The veins ower ones of a cornext order t line, or ations are the tightbe cast of he minute e original opearance is convex ultimate regularly nd which d pubes-

No. 15.

There are two casts of a lanceolate penninerved leaf having the aspect of a leaf of one of the *Rhamneæ*. In the outline of the lamina, and the nervation, they closely resemble a leaf of the common *Alaternus*, with the exception of being only slightly crenated instead of serrated. The greatest breadth is a little below the middle, and the base is somewhat fuller than the apex, which is acute. The impressions are an inch and a half long. There is a fulness scarcely amounting to wrinkling, of the minor meshes.

No. 16.

Is the lower half of an oval leaf resembling that of a willow, with a margin entire, or indented only by small pits, which are probably the casts of minute marginal glands. The base is obtuse, the apex unknown, and the lamina flat and smooth, with a very slight prominence of the principal veins. The transverse diameter is half an inch.

There are also various fragments of impressions of ribbed grasses or carices.

The group of plants seems to be such as one would expect to flourish in the climate of Canada West, and belongs perhaps to the meiocene epoch.

11.

inerved leaf mamneæ. In they closely h the excepof serrated. Idle, and the ich is acute. There is a f the minor

ing that of a only by small ute marginal own, and the prominence of er is half an

pressions of

a West, and

POSTSCRIPT.

THE preparation of the illustrations, and other circumstances, having retarded the publication of these volumes for some months after the letterpress was ready, the delay has enabled us to learn the result of the last year's search for the lost Expedition. The first traces of the missing ships, discovered on the south side of Beechey Island and on Cape Riley, as mentioned in vol ii. p. 155., were followed up by the discovery of seven hundred empty meat-tins, and other remains, which furnish undoubted proof of Franklin's ships having wintered, in 1845-6, on the inside of the above-named Island. The tombs of three men, with headboards bearing their names and the dates of their deaths, were erected on the east side of the Island, not far from the site of the armourer's forge, an observatory or store-house, and other enclosures opposite to the an-One of these men belonged to the "Terror," chorage. and two to the "Erebus," which is sufficient evidence of the presence of both ships; and the latest death supplies us with the date of 3rd April, 1846. The mortality does not exceed that of previous expeditions; and we may therefore conclude, that the Expedition was in highly effective order when it left that anchorage, with only a moderate inroad into its stock of preserved meats, the seven hundred empty tins found on the island forming but a small proportion of the 24,000 canisters with which the ships were supplied.

Captain Penny and his officers, who examined Beechey

E E 2

Island and the neighbourhood very carefully and minutely, believe that the Expedition did not quit its winter anchorage till the end of August or beginning of September, 1846, founding their opinion mainly on the lateness at which the ice knows up; that much of the summer was passed there, they consider as proved by the deep sledge ruts in the shingle, which must have been made after the snow had partially disappeared, and by small patches of garden ground bordered with purple saxifrage, and planted in compartments with the native plants.

It is also the opinion of several officers of the searching party that Franklin's ships left their wintering station suddenly. The reasons assigned for this belief are, that several articles which might have been useful were left behind, and that at a look-out or fowling station, on Cape Spencer, a long day's journey from the anchorage, the lines for securing the covering of a circular enclosure, formed by a low wall of stones, had been cut, instead of having been deliberately untied, when the covering was removed, leaving the ends of line attached to the stones. The absence also of any memorandum of past efforts or future intentions, either at the stone cairn erected on the south side of Beechey Island, at the pile of canisters, or in the neighbourhood of the kitchen, forge, and other marked localities opposite the anchorage, is thought by some to be an indication of the sudden departure of the Expedition. The value of the articles left behind is too trifling to support such an inference* and the absence of the diligently-

* These were an armourer's wooden stand, used when laid on its side for the support of an anvil, and when standing on its end for the insertion of a vice; several coal bags, two of them containing coal dust mixed with a small proportion of small cinders and ashes, some pieces of rope, and scraps of old canvass, and a small piece of oaken fire wood, besides many fragments of worn clothing utterly worthless. An iron stove that had been made on board ship was also found at a fowling

l minutely, ter anchornber, 1846, t which the assed there, ruts in the ne snow had s of garden planted in

the searching g station sudief are, that ful were left tion, on Cape rage, the lines are, formed by f having been removed, leav-The absence

future intenthe south side in the neighr marked loby some to be e Expedition. rifling to supthe diligently-

en laid on its side end for the intaining coal dust shes, some pieces oaken fire wood, thless. An iron and at a fowling searched for memorandum does not seem to be sufficiently accounted for by such a supposition. The time required for calling in the parties from Cape Spencer, Caswall's Tower in Radstock Bay, and other points where they have been traced, and for embarking the instruments and utensils from the observatories and kitchen, might have sufficed for the planting of a copper cylinder or bottle, with a memorandum. That the ships drifted out unexpectedly in a floe of ice is not considered by the nautical men who have examined the anchorage to be possible. The north point of Beechey Island being connected to North Devon by a shingle beach, covered by only two or three feet of water, no pressure of ice can operate ou the harbour from that direction so as to drive out vessels by the south-eastern and only navigable entrance, and it is almost certain that Franklin's ships must have made their exit by the tedious and laborious operation of sawing out.

The absence of a memorandum at the wintering station is remarkable, and, in my opinion, wholly unexplained by any suggestion that has hitherto been given by the many writers who have made their opinions known, through the medium of the periodical press. From Sir John Franklin's well-known anxiety to act up to the tenor of his instructions, combined with the expressed desire of the Admiralty, that he should embrace every opportunity of forwarding

station near the east corner of the island, but it is stated to have been not worth carrying on board. The bird's bones remaining in the vicinity of the stone enclosure on Cape Spencer show that the sportsmen encamped there had been tolerably successful; and much small shot was found scattered among the stones with which the enclosure was paved. In the interstices of the stone wall there were many pieces of newspapers, also two bits of paper of much interest to the friends of two of the missing officers — one being inscribed with the name of Mr. M'Donald the surgeon; the other containing part of a memorandum in the handwriting of Captain Fitzjames, giving directions as to the times of recording certain meteorological observations.

accounts of the progress of the Expedition to England, I should have thought that he would certainly have left a record of his doings at a winter station, which he knew to be within reach of the whalers, before he commenced his voyage of the second season, in the hope of penetrating either to the south-west or northward, where he knew there would be little or no chance of finding a channel of communication, unless he succeeded in overcoming all obstacles, and pushing his way through that archipelago, which has hitherto proved a is vier to successive expeditions. And should he, as some suppose, but contrary as I think to all likelihood, have cut his way out of Beechey Harbour merely to turn his face to England, still I think he would have left some authentic record on the spot, mentioning his labours, and the cause of his return.

As there are no natives on the north side of Lancaster Strait to disturb any memorial or flag-post that may be erected, Sir John Franklin would probably not think it necessary to bury the copper cylinder or bottle containing his memorandum, but would rather suspend it in the most conspicuous way he could devise. Now, I have learnt, by experience, that the wolverene * will ascend trees to cut down a package hung to a branch; and that bears have similar habits was fully ascertained by Captain Austin's sledge parties. A depôt formed by Lieutenant McClintock on Griffith Island was entirely caten by bears, the tin cases proving to be but a poor defence against the tusks of these omnivorous animals, who expressed their approval of preserved potatoes by the way in which they cleared That they would relish the pemican out the canisters. which was part of their spoil, might have been predicted. They did not respect even the sign-post, but overthrew it,

* The wolverene inhabits the islands north of Lancaster Strait, and its recent footmarks were often seen by Lieutenant M'Clintock.

ingland, ave left knew to need his netrating ew there of comobstacles, hich has s. And nk to all Harbour he would entioning

ancaster may be think it ontaining the most earnt, by s to cut ars have Austin's Clintock the tin he tusks approval cleared pemican edicted. hrew it, trait, and

ck.

and bit off the end of the metal cylinder containing the record.

The want of this memorial leaves us totally in the dark as to Franklin's intended course, which would in all probability have been decided upon before he left the harbour; for, from his position, he had the means of ascertaining the state of the ice both in Barrow's Strait, and in Wellington Channel. If the former was open, his course would be to Cape Walker and the south-west, agreeable to his instructions; but if Barrow's Strait was closed, as he had found it to be the preceding year, and Wellington Channel open, then he would gladly follow the latter, which one at least of his intelligent officers considered to be the most promising route of all, and which the spirit of his instructions permitted him to take, if shut out from the west or southwest.

The well-planned and thoroughly organised travelling parties of the searching squadron, though they traced with extraordinary perseverance extensive portions of insular coast, failed in detecting any further decisive vestiges of Franklin's course. Captain Austin's two ships, with their tenders, wintered at the south-west end of Cornwallis Island, under the shelter of Griffith Island. From thence Lieutenant M'Clintock, who made the longest journey of all the pedestrian parties, setting out in spring, rounded the west end of Melville Island in longitude 114° W., and, passing over the extreme discoveries of Sir W. Edward Parry, saw distant land extending beyond the 116th meridian. The intermediate passages and bays were explored by Lieutenant Aldrich, Mr. Bradford, and Mr. M'Dougall. On the south side of Barrow's Strait, Cape Walker, and the adjoining coasts, were traced by Captain Ommaney and Lieutenants Osborne, Meecham, and Browne; Lieutenant Osborne having carried his re-

searches nearly to the 72d parallel on the 104th meridian. This was the most southerly point attained. It lies within 180 miles of the south shore of Victoria land, a 1 is perhaps part of the same island. Throughout the whole of the great extent of coast-line closely examined by these officers, on both sides of the strait, no traces whatever of Sir John Franklin's ships were discovered, though Lieutenant M'Clintock found the wheels of a cart used by Sir W. E. Parry in 1820, and other traces of that officer's travelling parties. The signal posts planted by the latter were thrown down by wind or animals.

Captains Penny and Stewart in the Lady Franklin and Sophia, wintered in Assistance Harbour, in company with Rear-Admiral Sir John Ross, of the Felix. The spring journeys of the two former, and of their officers, were directed to the examination of Wellington Sound. Captain Stewart and Dr. Sutherland explored the west and north sides of this inlet, their most northern points being in latitude 76° 24' N. Messrs. Goodsir and Marshall traced its south and west sides to the 99th meridian; and both parties, from their most westerly stations, saw a navigable sea extending northward and westward, to the utmost limits of their vision. Wellington Strait, closed to the eastward and northward, opens into this westerly passage by three channels, separated from one another by Baillie Hamilton's and Deans Dundas Islands. Baring's Island lies more to the westward, opposite the middle channel. Its shores, and those of the two other principal islands, were examined by Captain Penny, who crossed over to the point of Sir Robert Inglis Bay on the northern shore, which has been named Albert Land; and from whence he had the melancholy prospect of boundless open water, which he had not the means of navigating. A boat was transported over the ice towards it with much labour; but, the

th meridian. t lies within l is perhaps thole of the l by these whatever of ough Lieuart used by that officer's by the latter

y Franklin in company Felix. The neir officers, ton Sound. he west and points being d Marshall ridian; and saw a navito the utlosed to the rly passage by Baillie ng's Island annel. Its s, were exthe point which has e had the which he vas trans-; but, the

provisions of the crew running short, it was abandoned. Mr. Goodsir found a spar of American spruce, untrimmed, with its bark worn off, and broken at both ends, twelve feet long, and as thick as a man's ankle, on the shore facing the open water; also many smaller pieces of the same kind of drift wood, while none was picked up by Captain Stewart in Wellington Sound. From this fact these officers inferred, that the drift wood had come from the westward. The currents or tides among the islands at the western outlet of Wellington Strait, were at times, according to Captain Penny's judgment, not less than four knots; and the general opinion of his officers was that the principal set of the stream came from the westward, and the prevailing winds from the north-west.

Animal life was abundant in the open water, and on its Walruses were seen repeatedly in the several coasts. channels, north and south of Baillie Hamilton's Island; and polar bears were numerous and bold, so as to be dangerous to parties not well armed. Several of the bears were killed, and one of them contained an entire seal in its stomach, the practice of these voracious animals being to swallow their prey without mastication when it is not too large to pass their gullets. The walrus cannot exist except when it has access to open water; nor is the polar bear usually found at a distance from it, except in its passage from one sheet of water to another. The travellers also saw polar hares, wolves, foxes, herds of rein-deer, vast flocks of king and eider ducks, brent geese, and many gulls and other water-fowl of less utility to man. Musk oxen were seen only on Melville Island, where Licutenant M'Clintock killed four, and might have procured more had he wished to do so.

On the 5th of September, 1850, a floc of ice at least two years old, and upwards of thirty miles in width, filled the VOL. II. F F

lower part of Wellington Strait, and remained fast, though diminished in breadth, when last visited on the 24th of July, 1851. Captain Penny is of opinion that open water existed beyond it all the winter.

With respect to traces of Sir John Franklin's Expedition, beyond Cape Spencer none whatever were observed by Captain Penny's travelling parties, except a small piece of drift wood, which had been recently charred, and had been exposed to little or no friction subsequent to the operation of fire.* This was found by Mr. Goodsir in Disappointment Bay, in latitude 75° 36' N., longitude 96° W.; and I consider it to be certainly a relic of Sir John Franklin's Expedition, as these coasts are not now visited by natives, and this piece of charred wood could not have been waterborne from any great distance. It must have travelled, however, some short way subsequent to its having been exposed to the action of fire; for if it had been the remains of a fire kindled on the spot, other fragments of charcoal would have been found lying beside it. Franklin would, undoubtedly, during the spring passed in Beechey Bay, send out a party up Wellington Sound, as he would never let the opportunity escape of examining, as far as he was able to do, a route that might influence his future movements; and as the course to the westward within the reach of pedestrian parties was known, the resources of the two ships would be turned to the undiscovered way, commencing in their vicinity. That such exploring party went beyond the limits of Captain Penny's researches, I infer

* A piece of eln board that had been originally coated on one side with mineral pitch or tar, and after long exposure to the weather split oy an axe, was too much weathered even on the most recent surface to come within the date of Sir John Franklin's Expedition. It was found on Baillie Hamilton's Island, and must have drifted a very long way.

from neither post nor cairn marking the limit of its journey having been seen. If the same expanse of open water was visible, in 1846, from Baillie Hamilton's Island, which Captain Penny saw in 1851, we may readily conceive the efforts that would be made to carry the Erebus and Terror into it by any practicable extent of icc sawing, particularly if Barrow's Strait remained closed. The age of a floe of ice filling a strait does not indicate with certainty the length of time that the strait has been blocked up, for drift ice, loaded with the remains of several years' snow, may be carried into a narrow passage, so as to shut it up, and as suddenly removed again on a favourable concurrence of winds and tides. One navigator, therefore, may be able to sail, as Sir W. E. Parry did, nearly quite through that northern archipelago in one season, while his successors may find impassable barriers thrown across the path which he pursued, and new avenues opened. It would be unsafe, therefore, to argue that Wellington Strait is always closed, because it was choked by a floe of some age in 1850 and 1851.

By the efforts of the searching parties, which have just returned, combined with those of preceding years, all the accessible parts of the continental coast of America have been explored, and both sides of Barrow's Strait, to the further side of Melville Island, and the land beyond Cape Walker. Land has also been traced, though only by distant view, round the bottom of Jones's Sound. This has narrowed the lines of search to two distinct points—that is, to the south west of Cape Walker, which, from its being the direction in which Sir John was instructed to go, seemed to be especially the one in which he was to be sought; and the newly-found channel opening out to the westward from Wellington Strait. It is greatly to be desired that this one may be pursued by new efforts.

he 24th of open water

Expedition, bserved by all piece of nd had been e operation Disappoint-6° W.; and ı Franklin's by natives, been watere travelled, naving been the remains of charcoal klin would, echey Bay, would never r as he was ture movein the reach of the two y, commenparty went nes, I infer

d on one side the weather most recent Expedition. we drifted a

Mr. Rae, in April last, was on the eve of setting out from Great Bear Lake, in the hope of crossing on the ice to Victoria Land, and of continuing his search in a boat as soon as the navigation opened. Though he may not actually attain Lieutenant Osborn's furthest, he may, under favourable circumstances, approach so near to the scene of that officer's search, or of Lieutenant M'Clintock's, as to prove, should he find no traces of the ships, that the intervening space is too confined for the seclusion of living men. Captain M'Clure, who passed to the eastward of Point Barrow last season, if he found the sea as open as the more sanguine believe it to be, may have reached the west side of Parry's Archipelago, and have spent the winter not far from the supposed outlet of Victoria Channel; and this season Captain Collinson may be sailing eastward in the same direction. It is from Beering's Straits, then, that we are next to look for tidings of great interest to the civilised world, which sympathises so universally with the efforts made to trace and relieve so many gallant victims to science.*

20 October, 1851.

* With reference to Sir John Ross's pigeons, mentioned in a note on page 157. of Vol. II., it appears that he despatched the youngest pair on the 6th or 7th of October, 1850, in a basket suspended to a balloon, during a W.N.W. gale. By the contrivance of a slow-match the birds were to be liberated at the end of twenty-four hours.

THE END.

LONDON: Spottiswoodes and Shaw, New-Street-Square. etting out from ice to Victoria as soon as the ly attain Lienarable circumfficer's search, hould he find ace is too con-M'Clure, who season, if he lieve it to be, hipelago, and sed outlet of linson may be om Beering's ings of great es so univereve so many

oned in a note the youngest uspended to a f a slow-match hours.



