THE PRINCIPLES OF PATHOLOGY

VOLUME I

GENERAL PATHOLOGY

By J. George Adami, M.A., M.D., LL.D., F.R.S.

VOLUME II

SYSTEMIC PATHOLOGY

By J. George Adami, M.A., M.D., LL.D., F.R.S. and Albert G. Nicholls, M.A., M.D., D.Sc., F.R.S. (Can.)

THE

PRINCIPLES OF PATHOLOGY

BY

J. GEORGE ADAMI, M.A., M.D., LL.D., F.R.S.

PROFESSOR OF PATHOLOGY IN MCGILL UNIVERSITY, AND PATHOLOGIST-IN-CHEF TO THE ROYAL VICTORIA HOSPITAL, MONTREAL; LATE FELLOW OF JESUS COLLEGE, CAMBRIDGE, ENGLAND

VOLUME I

GENERAL PATHOLOGY

SECOND EDITION, REVISED AND ENLARGED WITH 329 ENGRAVINGS AND 18 PLATES



LEA & FEBIGER PHILADELPHIA AND NEW YORK



Entered according to the Act of Congress, in the year 1910, by LEA & FEBIGER,

in the Office of the Librarian of Congress. All rights reserved.

22092

IN ALL GRATITUDE

V GRVELEE OF V CENTREY THE FEIEND AND COUNSELLOR OF MORE THAN

Y E' S'

O.L



PREFACE.

In this work I have made the attempt to place before student and physician in an orderly and reasoned manner the principles of Pathology, the science, as distinct from the practice of medicine; the science upon which that practice is, or should be, based. Preëminently it is the duty of the pathologist as teacher to train the student in the habits of medical thought, to afford those data which bear upon disease in general, to show how such data are to be weighed, and what deductions may logically be drawn therefrom, so that later the student investigating a particular case may do so armed with a sound knowledge of general principles; that he may recognize individual symptoms not as isolated facts, but as indications of definite orders of disturbance affecting one or other organ, and, knowing what in general induces those disturbances, may form a judgment regarding the causation and meaning of the sum total of symptoms in a case. As Bacon laid down, "Vere scire est per causas scire"-To know truly is to know through causes-and he is the scientific physician or surgeon who seeks and determines causes; for only when the cause is deduced can treatment be rational.

I hold, therefore, that, whatever may be the case with other subjects, what is needed in a text-book of Pathology is not the mere record and description of phenomena, but the attempt to analyze those phenomena in an orderly manner. That text-book should be a training in medical thought. It was, however, one thing to hold these views, another thing to write a treatise embodying them. I will not state how many times most of the chapters of this work have been written and rewritten; nor how often the arrangement has been changed before the work has assumed its present shape in two volumes, this, the first of two, dealing with what is usually termed General Pathology; the second dealing with Systemic (including Special) Pathology. I will only say that constantly, in working over each section, I was forced, with Virchow, to recognize the cell and the changes undergone by it as the basis of all pathological sterdy, and thus, eventually, to guard against constant reversion to elementary but basal and all-important matters, was com-

PREFACE

pelled to write an introductory section upon the cell and its properties, more particularly in relationship to morbid changes.

The work has thus far assumed a novel, but what I am convinced is a logical, form. It begins not with a study of the blood and of circulatory disturbances, as has been usual with most works on General Pathology, but with a study of the properties of living matter. The study of circulatory disturbances is not, indeed, a part of General Pathology, and, accordingly, it is treated as the introduction to Systemic Pathology, that is, to the study of the diseases affecting individual systems and the effects of those diseased states upon the organism as a whole. As such it is treated in the second volume. It would be as appropriate, if not more so, to begin the study of General Pathology with the discussion of nervous disturbances and their effects upon the body at large.

There are different orders of minds, and no work can appeal to all. For myself, in beginning my studies, I found that I could easily remember the matter of such works as the larger Lyell's *Principles of Geology*, Foster's *Physiology*, and Fagge's *Medicine*, to cite examples in which there was a reasoned treatment of the subject, whereas, to attempt to commit to memory "cram books" laden with facts and names was mental agony. I saved time and gained knowledge by reading my subject at large. It is to those possessing a like order of mind that this work is addressed.

It would be false modesty on my part were I to pass by in silence the welcome with which the first edition of this work has been greeted. To put forth two very large volumes, in which the subject of Pathology was treated along unaccustomed lines, was a bold venture, and when, in addition, I had at hesitated to express individual views often at variance with time-honored teachings, and knew only too well that he who attempts to cover the whole subject could not treat each department with the sureness of the specialist in that department, I had expected very considerable criticism. Thus, I am deeply sensible of the warm words of commendation which have come from my colleagues and reviewers. In this edition I have endeavored to the best of my ability to remedy some at least of the defects of that first edition. More particularly in Section I the pages upon nuclear and nucleolar function have been largely rewritten; the lipoids, their relationships and chemistry, have been given fuller notice; the more recent views regarding the reduction process in the germ cells have been included, together with Godlewski's important observations upon the influence of the cytoplasm upon the developing individual, together with some of the more important recent studies upon hybridism and Mendelism. In Section II attention may be called

viii

PREFACE

to the fuller treatment of physical agencies-light, heat, cold, and electricity-and of nutritional disturbances as causes of disease. In Section III material additions have been made to the discussion of the febrile state; fuller attention is directed to the subject of non-specific immunity; while the diversion of complement and Wassermann's reaction, and the newer work upon the relationship of lipoid substances to the immune bodies are taken into consideration. The chapters upon Grafting and Metaplasia have been rearranged and, it is hoped, rendered clearer. In the chapters upon Neoplasia, what is perhaps the most material change is in connection with the tumors derived from neuroblastic elements. and more especially the establishment of the class of neurinomas to include the multiple cutaneous fibromas or neurofibromas, while Mallory's valuable studies upon intercellular fibrillation as an aid to diagnosis are given increased prominence. Among the regressive tissue changes, the most extensive alteration made is in the analysis of the Hyaline degenerations and the establishment of Elastoid degeneration as a distinct class. The pages upon Cholelithiasis have been largely rewritten and have undergone material alteration. The subject matter of the previous appendixes has been incorporated in the text, while the newer work upon the ultramicroscopic causes of disease which has appeared while this volume has been passing through the press has necessitated a brief appendix.

If in writing the original work the hopelessness of the attempt to master the subject in all its aspects impressed the writer, still more has his inability to keep abreast of the advances in all branches of medicine and cognate sciences during the last two years weighed upon him in revising this edition.

It is impossible for any individual to keep abreast of the manifold developments of all the sciences ancillary to medicine, physics and physical chemistry, biochemistry, biology and embryology, parasitology, histology, and physiology, and at the same time to master the literature of Pathology proper. In bacteriology alone and its one branch, the study of immunity, there is enough material being brought forth month by month to keep the reader fully engaged. Much that is of first-class importance is passed by in silence in these pages. At most, the writer has made the attempt to call attention to the intimate bearing of these other sciences upon medicine, and, in addition, to the important work now being accomplished by English-speaking workers. This last not through Chauvinism, or as a protest against the neglect that this work has too often received at the hands of Continental writers, but primarily to encourage the student in the habit of consulting authorities at first hand, of reading original articles and making his own deductions independently of the

PREFACE

opinion expressed by the writer of the text-book. There is no difficulty in obtaining the leading American and English medical journals, and when once the student appreciates the added interest and strength that come from first-hand reading, he will not be content until he masters the other languages of science—German, French, and Italian also. And, what is of like importance, it is sought to impress upon the student the opportunities that are before him in our university laboratories and well-equipped hospitals to undertake equally valuable investigations. If others—it may be of the same school, or known to him—have accomplished work of high order, why should not he also undertake research and seek to add to the sum of medical knowledge?

I cannot conclude without expressing my most hearty appreciation of the cordial cooperation I have received from my publishers in preparing this edition. For such new illustrations as are not my own I would especially thank Professor Reichert, of Philadelphia; Professors Mallory and J. H. Wright, of Boston; Dr. Fordyce, of New York, and Dr. Bashford, of London.

The heaviest debt of all I owe to my old teachers. Under the guidance of one near to me, the late Professor D. J. Leech, of Manchester, it was my good fortune and my privilege as a student to come under the influence, and that intimately, of not a few men who have been masters of their particular subjects, who, diverse it may be, in their gifts, have each possessed that intangible something that we recognize as greatness; men who have inevitably moulded my thoughts—Milnes Marshall and Francis Maitland Balfour, Michael Foster and Rudolf Heidenhain, Julius Dreschfeld and Charles Smart Roy, Emile Roux and Élie Metchnikoff. To them and to their teaching and inspiration is due whatever of virtue these pages may possess.

The cordial reception which this work has received leads me to ask the authors of researches on general pathological subjects to be kind enough to send me reprints, in order that it may be kept up to date.

331 PEEL STREET, MONTREAL.

J. G. A.

SECTION I.

PROLEGOMENA.

CHAPTER I.

CHAPTER II.

THE HISTOLOGY OF THE CELL.

CHAPTER III.

THE PHYSIOLOGY OF THE CELL.

CHAPTER IV.

THE CHEMISTRY OF THE CELL.

The Proteins, Amino-acids, and Polypeptids-	The Chemistry of the Nucleus-	
The Proteins and Metabolism-Life		54

CHAPTER V.

THE CHEMISTRY OF THE CELL (CONTINUED).

CHAPTER VI.

THE CHEMISTRY OF THE CELL (CONTINUED).

The	Non-proteid	Constituents-Water-Colloids and Crystalloids-Simple	
	Salts-Carl	pohydrates—Lipoids	85

CHAPTER VII.

GROWTH-RESERVE FORCE-STATES OF CELL ACTIVITY.

On Certain Functions of Living Matter-Growth-Physiological Inertia and 98 Habit-Reserve Force-The States of Cell Activity

CHAPTER VIII.

CELL MULTIPLICATION.

CHAPTER IX.

ADAPTATION.

Evolution and Adaptation-Its Physical Basis-Adaptation to Physical

CHAPTER X.

Cell and Tissue Differentiation-Individual Development.

Relationship	to	Ac	lapt	ati	on-	-E	oige	nesi	s	and	F	refe	orm	atio	n-	-Th	e)	Mos	aic	
Theory																				129

CHAPTER XL.

FERTILIZATION.

Germ Cells-Spermatogenesis-Oogenesis-The Part Played by the Chromo-

CHAPTER XII.

The Biophoric Hypothesis 156

CHAPTER XIII.

INHERITANCE.

The Different Forms-Raci	land Familial 15
--------------------------	------------------

CHAPTER XIV.

INHERITANCE (CONTINUED).

Parental ar	id Indivi	dual	Inh	erita	nce-1	Iendel	's Pr	incip	les-	Galt	ton'	s I	aw		
Mosa	ic, Atavis	stie,	and	Reve	ersion	ry In	herit	ance	-Fai	milia	al D	ege	ener	a-	
tion	and Diath	etic	Reve	ersion	-Sex	Limit	ed ar	nd Cu	mula	tive	In	er	itan	ce	
Mu	itation														167

CHAPTER XV.

INHERITANCE (CONTINUED).

The Theory of Inheritance-Inheritance of Acquired Character . . . 185

SECTION IL

THE CAUSES OF DISEASE.

CHAPTER I.

INTRODUCTORY.

Disease Inherited or Acquired-The Classification of Morbid State . . . 201

CHAPTER IL.

INHERITED MORBID CONDITIONS.

What is and What is Not Heritable-The Marriage of Consanguines . . 205

CHAPTER III.

THE CAUSATION OF MORBID CONDITIONS OF INTRAUTERINE AND PARTURIENT ACQUIREMENT.

Classification-Placental Disease and Its Influence-Diseases Peculiar to the

CHAPTER IV.

MONSTROSITIES AND ABNORMALITIES.

Abnormalities of Excess-Twins and Double Monsters-Monsters by Serial Deduplication: by Dichotomy; by Fusion-Deduplication of Organs. 226

CHAPTER V.

MONSTROSITIES AND ABNORMALITIES (CONTINUED).

Abnormalities of Defect-Dwarfism-Polar Hypogenesis-Defective Closure of Dorsal Groove of Thoracico-abdominal Fissure-Defects of Diaphragm. Face, Gill Clefts, Cloaca-Situs Inversus-Hermaphroditism . . .

CHAPTER VI.

POSTNATAL ACQUIREMENT OF DISEASE.

CHAPTER VIL.

EXOGENOUS INTOXICATIONS.

Non-parasitie-Poisons Acting upon and through the Nervous, Muscular, Circulating, and Other Systems 303

CHAPTER VIII.

EXOGENOUS INTOXICATIONS (CONTINUED).

Parasitic Causes-Bacteria as Causes of Disease-The Normal Defences of

CHAPTER IX.

EXOGENOUS INTOXICATION (CONTINUED).

rotozoan Parasites—Th	Various	Orders of	Pathogenic	Protozoa and Their	
-----------------------	---------	-----------	------------	--------------------	--

CHAPTER X.

EXOGENOUS INTOXICATIONS (CONTINUED).

CHAPTER XI.

Endogenous Intoxications.

CHAPTER XII.

Endogenous Intoxications (Continued).

Disintegrative Intoxications-Autolysis, Burns, etcImpaired Metabolism as	
a Cause of Disease-Gout and Allied Conditions-The Intermediate	
Intoxications - Gastro-intestinal "Auto-intoxication" - Obstructed	
Elimination as a Cause of Disease	370

CHAPTER XIII.

CHAPTER XIV.

PREDISPOSITION AND SUSCEPTIBILITY.

Life Periods and Incidence of Disease-Tissue Susceptibility-Idiosyncrasy . 404

SECTION III.

THE MORBID AND REACTIVE PROCESSES.

INTRODUCTORY.

PART I.

THE MORBID AND REACTIVE PROCESSES PROPER.

CHAPTER I.

THE LOCAL REACTION TO INJURY.

xiv

CHAPTER II.

	The	LOCAL	REACTION	TO	INJURY	(CONTINUE	D).		
ronic Infl	lammation	-Stage	s and Forn	ns—'l	he Mai	n Data Reg	arding	Inflam-	
mati	on-Fibro	sis and i	its Relation	iship	to Infl:	ammation			439

CHAPTER III.

THE SYSTEMIC REACTION TO MICROBIC INJURY.

The Process of	In	fection	-It:	: Ce	urse	<u> </u>	Type	s	-Exe	oge	nous	Ba	eter	ial	II	ito?	sica	÷	
tion .																			453

CHAPTER IV.

The Systemic Reaction (Continued).

The Febrile State .																		465
---------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	-----

CHAPTER V.

The Systemic Reaction (Continued).

Thermogenesis and Pyrexia—The Significance of the Febrile State—Relationship of the Nervous System to Pyrexia 478

CHAPTER VI.

IMMUNIZATION AND IMMUNITY.

CHAPTER VII.

IMMUNIZATION AND IMMUNITY (CONTINUED).

The Various Orders of Immunity—Indifferent or Non-specific Immunity— Immunity toward Substances of Known Constitution—Toward Phytotoxins and Enzymes—Toxins and Antitoxins—The Side-chain Theory 499

CHAPTER VIII.

IMMUNIZATION AND IMMUNITY (CONTINUED).

CHAPTER IX.

IMMUNIZATION AND IMMUNITY (CONTINUED).

Theories of Immunity-The Phagocytosis Theory-The Side-chain Theory 561

CHAPTER X.

SYSTEMIC REACTION THROUGH THE NERVOUS SYSTEM.

XV

PART II.

THE TISSUE CHANGES.

A. The Progressive Tissue Changes.

CHAPTER XI.

HYPERTROPHY.

CHAPTER XII.

REGENERATION.

Simple Regeneration-	-He	eter	om	orp	hosi	is-	Re	gene	erat	ion	of	the	Va	riou	sТ	issu	les	
in Man									*									601

CHAPTER XIII.

GRAFTING OR TRANSPLANTATION 631

CHAPTER XIV.

METAPLASIA AND HETEROPLASIA.

CHAPTER XV.

NEOPLASIA

CHAPTER XVI.

NEOPLASIA (CONTINUED).

CHAPTER XVII.

NEOPLASIA (CONTINUED).

CHAPTER XVIII.

NEOPLASIA (CONTINUED).

Typical Hylie Tumors of Mesenchymatous Origin (Connective-tissue Tumors) --Fibroma, Myxoma, Lipoma, Chondroma, Osteoma

XVI

CHAPTER XIX.

NEOPLASIA (CONTINUED).

Typical Hylic Tumors of Mesenchymatous Origin: Myelomas, Lymphomas, and Lymphomatosis. Leiomyomas-Of Mesothelial Origin : Rhabdomyomas-Of Epiblastic Origin: Neuromas, Gliomas, and Neurinomas-Of

CHAPTER XX.

NEOPLASIA (CONTINUED).

CHAPTER XXI.

NEOPLASIA (CONTINUED).

Primary Lepidic Tumors-Typical: Papilloma and Adenoma-Atypical: Car-

CHAPTER XXII.

NEOPLASIA (CONTINUED).

Transitional Lepidic Tumors-Mesotheliomas and Endotheliomas, including Hemangiomas-Tumors of Doubtful Relationship: Melanomas, Cho-

CHAPTER XXIII.

NEOPLASIA (CONTINUED). Theories of Neoplasia

CHAPTER XXIV.

CYSTS.

B. The Regressive Tissue Changes.

CHAPTER XXV.

NORMAL HISTOLYSIS AND CYTOLYSIS.

CHAPTER XXVI.

THE DEGENERATIONS AND INFILTRATIONS.

CHAPTER XXVII.

THE DEGENERATIONS AND INFILTRATIONS (CONTINUED). Conjugated Proteins and Sclero-proteins: Mucoid, Colloid, Amyloid, Elastoid,

xvii

CHAPTER XXVIII.

THE DEGENERATIONS AND INFILTRATIONS (CONTINUED).

Disturbances of the Aqueous Contents of the Cell: Of Fatty Contents—Of Carbohydrate Contents—Glycogenous Infiltration 903

CHAPTER XXIX.

CALCIFICATION AND CALCAREOUS DEPOSITS.

Calcareous Metastasis, Calcification Proper-Calcareous Concrements 924

CHAPTER XXX.

Deposits of Other Products: Calculi.

CHAPTER XXXI.

PIGMENTATION AND PIGMENTARY DEPOSITS.

Endogenous, from Hemoglobin and Its Derivatives: Hemochromatosis and Icterus—From Melanin and Lipochromes—Exogenous Pigmentation . 956

CHAPTER XXXII.

NECROSIS.

CHAPTER XXXIII.

DEATH.

APPENDIX.

xviii

THE PRINCIPLES OF PATHOLOGY.

SECTION I.

PROLEGOMENA.

CHAPTER I.

INTRODUCTORY.

Definition.—Just as physiology is the study of the functions of the body in health, so is pathology the study of the same functions in disease. To this extent we can accurately define the scope of our subject, but a moment's thought reveals that the definition is only superficially precise; further thought shows that, strive as we may, we cannot approach nearer to perfect definition. For everything depends upon what we understand by the terms "health" and "disease," and we are forced straightway to recognize that there is no boundary line—no definition—between the two—they merge insensibly the one into the other.


We all, it is true, have a general understanding of what health is, but when we come to attempt to express our understanding in words we find that, like "life" itself, it eludes definition. Why this is we may here briefly indicate.

It depends primarily upon the basal fact of variability. No two living beings, although belonging to the same species, and the same family, are structurally identical, nor even born identical, and if this be true of structure, it is true also of the outcome of structure—namely, function. There is thus no absolute standard of either structure or function in any one species. Every individual of the human or other species varies in every particular from every other individual; the dimensions of the different component parts, the proportional relationship between the parts, the action of the parts, present more or less evident divergence in any two individuals studied. At most, by the statistical method we can in some cases arrive at an approximate or theoretical standard. To give an example: There is no absolute standard of height for human beings; the average height varies in the different branches of our race, and differences, sometimes pronounced, occur

2

INTRODUCTORY

among the members of one and the same family. But if, as Quetelet' first noted, we take one thousand, or preferably ten thousand, fullgrown males—the larger the number the better—belonging to the same branch of the human race, and measure accurately their height, and plot out the results obtained, we find that the majority conform to, and are included in, a relatively short range. We can speak of that particular height to which the greatest number of individuals conform as the standard, or mode—i. e., the fashionable height; or again, taking the sum of the heights and dividing by the total number measured, we can obtain the type or arithmetical mean. ("Mode" and "mean," it may be added, do not necessarily correspond.) It will be found that the majority of individuals come within a relatively small range on either side of this mode, and we can on either side of the mean (or of the mode) determine the median or probable deviation (A A, and B B, in our first



Normal curve of frequency. This is a flowing curve symmetrical about the mean (which corresponds with the mode) and without limit at either end. XX, ordinate of mean or type (and of mode); A, A, B, ordinates of lower and upper median deviation.



A skew curve of frequency in which X X, the ordinate of mean or type, does not correspond to M M, the ordinate of the mode. The continuous then here represents the estimated (theoretical) curve of frequency, the interrupted line joins the numbers of the classes or grandes found in a particular case.


curve) above and below which fall 50 per cent. of the measurements. We can go farther and *assume* that the 50 per cent, within these limits constitute the normal class, those outside the abnormal. We, in short, obtain a curve which, in an uncomplicated case, corresponds curiously with that of the law of chance (Fig. 1). To illustrate: Taking the case of twelve dice, throwing an infinite number of times and counting the sum of the numbers turned uppermost (from 12–all "ones"–up to 72–all "sixes"), it can by this "law of chance" be determined mathematically what is the probable frequency of a given number—say 25—being exposed. Plotting out these probabilities, they are found to form an exact curve. This has been tested by Weldon, who selected a somewhat simpler case. Taking twelve dice, he threw them several

¹Anthropométrie ou Mésure des différentes Facultés de l'Homme, Brussels, 1871.

VARIABILITY

thousand times, and tabulated the number of dice at each throw which presented more than three points on their uppermost surface. Plotting out the number of dice—1, 2, 3 to 12—which fulfilled the conditions, he found that the curve giving the successive numbers followed with remarkable closeness the mathematical curve of chance.¹

In plotting out the measurements from a series of men or of other species, we obtain a like curve presenting a regular diminution at either side of the mode, the least frequent classes being those situated farther



Curve of variation in the head breadth in millimeters of 3000 criminals. Irregular continuous line = polygon of observed frequency; darker continuous line = normal curve of frequency as estimated by law of chance; dotted line = estimated skew curve of frequency of type IV (Pearson's table). (After MacDonnell.)

and farther away. While the majority of individuals approximate to the mode, few will exactly conform either to it or to the type (Figs. 2 and 3).

But in all such tabulations it has to be realized that the results obtained are not absolute; measurements of another thousand or ten thousand individuals *might*, nay, would surely, to some extent, move the mode or

¹ A concise and excellent account of the mathematics of variation is given by R. P. Bigelow, art. Variation, Buck's Ref. Handbook, second edition, 1904. See also Elderton, W. P., Frequency Curres and Variation, London, 1907; Davenport, C. B., Statistical Methods with Special Reference to Biological Variation, second edition, New York, 1904; and for yet fuller details, the various papers by Karl Pearson, Phil. Trans. Royal Soc. Lond., 1895 to 1905. The journal Biometrika has been established to deal with studies of this order.

INTRODUCTORY

20

mean in one or other direction; we only obtain an approximate standard. We need not here discuss *skewness*, or want of correspondence between mode and type (mean), or again, *multimodal curves*, in which two or more classes or modes are discovered grouped around certain dimensions and separated by intermediate classes containing fewer individuals. What we wish to emphasize at the present moment is that the normal gives place imperceptibly to the abnormal, every gradation being found between cases which approximate to the mode, and so may be regarded as normal, and those which show extreme departure from the same.

What is true of bodily dimensions and structure must inevitably be true also regarding function. From this it follows that, if health be regarded as the indication of perfect functional activity, and disease of imperfect function, the two merge imperceptibly the one into the other. For this must be accepted from the beginning, that conditions of disease are conditions of disturbed or imperfect function; they connote either the exaggeration or the diminution of processes which are of normal occurrence.¹

Local Disease, its Relationship to General Health .- If the difficulty in drawing a sharp line of distinction between the normal and the abnormal obtains in connection with inherent bodily states, it obtains also in connection with those that are acquired during the course of the individual existence. Considering man in the first place-but the same is true of all multicellular organisms—it is seen that, while the organism is in one sense a complete unit, in another sense it is a compound, formed of a great number of different structures, each made up of individual cells and the products of their activity. These structures and the cells composing them are at the same time-to an extent varying in the various organs-interdependent and independent. So far as they are interdependent, disturbance or disease of a given organ is liable to affect the other organs, and the body as a whole, causing constitutional disturbance; so far as they are independent, local disturbance may remain wholly local, or, in other words, the organism as a whole may be healthy while a part is diseased. For example: injury, disease, or loss of one tooth is liable to throw more work on the other teeth, and, rendering mastication not so perfect, may throw more work on the stomach; while again, through the disturbed innervation of the part there may be profound nervous irritation, sleeplessness, lack of rest, and in this and other ways marked constitutional disturbance may be set up. One knows full well, however, that any single tooth may be the seat of progressive caries or may be entirely lost, and no such train of constitutional disturbances be set up; there may be local disturbance and excellent general well-being. Thus our conception of what is health and what disease must be dependent upon whether we take into consideration the organism as a whole or the condition of the various constituent parts. To this paradoxical interdependence and independence of the cells we shall return.

¹ As was first faid down by Michael Servetus. See Osler, Michael Servetus, Baltimore, 1909;13. 13

In the forthcoming chapters we shall have to consider the more important features bearing upon the production of the conditions of disease, and the reaction of the organism to the same whereby a condition of health or relative health is brought about. In the meantime it will be well to afford a working definition. Bearing in mind that these terms are, and can only be, relative, it is well to consider health as a condition of metabolic equilibrium—a condition in which the organism or the part, is attuned, or in complete adaptation to its surroundings; disease as a condition in which equilibrium and adequate adaptation are wanting. In other words, to employ a metaphor encountered by chance in the works of a seventeenth century Italian theologian, "health is harmony, disease discord," a statement which can be applied to either general or local bodily condition, and which, to continue the metaphor, acknowledges or embraces the fact that the harmony may be in a minor key. That individual is regarded as enjoying good health, and in fact actually does enjoy good health, who nevertheless may for years have had extensive local disease of the heart valves, which in its turn has caused hypertrophy of the heart muscles in response to the increased work thrown on the organ. It is true that in such a one any sudden excitement or demand for increased work, which would have no deleterious effect upon the normal organs, suffices to bring on indications of heart failure. But within certain limits, employing ordinary caution, a subject of valvular disease of the heart may for long years enjoy life and carry out well all the ordinary duties without obvious bodily disturbance.

Diseases and Affections .- From the time of Galen onward there has been discussion and confusion regarding the relationship between local and general disturbances, so that as an aid to exact thinking and clear conception an increasing number of French and German writers emphasize the distinction between "maladie" and "affection," "Krankheit" and "Leiden." Disease is the process leading to affection of one or other organ. Pneumonia, as Roger aptly illustrates, is at the same time an infectious disease and an affection of the lungs. It is urged that to speak of "heart disease" or "kidney disease" leads to an incorrect mental imagery, to wrong ideas as to the nature of the malady, and very possibly to wrong treatment. In a case of thickening and contraction of the mitral valve in a young adult the morbid process that has led to the stenosis of the valve is acute rheumatism, and that is the disease which has resulted in the affection known as stenosis. And it is recommended that to indicate that the various resultant states in the different organs are not primary diseases but resultant states, we should designate these as "pathies"-cardiopathies, nephropathies, neuropathies, etc.

With these recommendations we are wholly in accord, provided that it be kept ever in mind that a cardiopathy, for example, may in itself

¹ For a résumé of French opinions upon this subject see Roger, in Bouchard's Pathologie Générale, 1: 1895; 18; and for the latest German pronouncement, Aschoff, Ucber den Krankheitsbegriff und verwandte Begriffe, Deut, med. Woch., 1909; 33.

INTRODUCTORY

be the causative agent of further morbid processes. Heart disease, that is, does exist. Rheumatism does not in itself induce congestion of the lungs, liver, and viscera in general, with cedema and ascites such as we see in mitral stenosis. Those are due directly to the cardiopathy, to the narrowed condition of the mitral valve and the consequent obstruction to the onward flow of blood. Acute rheumatism is here the remote cause, the mitral stenosis the direct. The above syndrome of vascular disturbances constitutes a definite morbid process, obstructive heart disease. Or, to sum up:

 Disease is a process or succession of disturbances induced by any agent which disturbs the normal activities of the organism as a whole, or of its constituent parts.

 Such disease may have purely a local seat, modifying the function and activities of an individual organ or portion of an organ, or may coincidently affect several organs.

3. In the latter case it frequently happens that the brunt of the disturbance falls upon one organ or tissue, so that we are apt, mistakenly, to regard the affection of that organ as the primary disease.

4. The affection of an organ or part may be of such a degree as not seriously to disturb the working of other parts of the organism; or, on the other hand, may be the direct cause of affections of other parts, and so is apt to be mistaken for the primary cause of disease, where, as a matter of fact, it is the secondary.

The recognition of these distinctions in practical medicine is of the highest importance. He who, for example, treats a case of obstructive heart disease as a primary condition and directs his efforts purely toward overcoming the disturbances due to cardiac inadequacy, may, it is true, bring about temporary alleviation of the condition, but only temporary. The primary cause may still be effective, be it the virus of acute rheumatism or factors inducing raised blood pressure. That is the rational medicine which seeks and discovers the primary disturbing agent, and, counteracting this, gives the fullest opportunity for the measures directed against the secondary effects to be permanently serviceable. It may not always be possible to determine the nature of the primary cause. Again, the local affection may have led to irreparable tissue changes; so also serious lesions inducing characteristic trains of symptoms may show themselves as remote effects long years after the acute incidence of the primary disease (e. g., the nervous lesions causing locomotor ataxia may only develop twenty years after the primary syphilitic infection, and when the causative agents are no longer present in the system). All this has to be admitted, but does not in any sense contradict the general principle. On the contrary, it expands it, indicating that it is the duty of the physician to determine more surely the primary causes of disease, to recognize at the earliest stage the effects of the same, and to institute treatment before those effects have proceeded to the stage of setting up irreparable injury.

Cellular Physiology and Cellular Pathology.—If, then, we take the position that for every structure and every function of the body, and for the

22

CELLULAR PATHOLOGY

body as a whole, there is a certain mode within the confessedly vague limits of which conditions are to be regarded as normal, it follows that anything outside the limits on either side is abnormal, and it is these conditions of excess or defect that for us must constitute disease; it is these we have to study. It follows also that constantly in our study of pathology we must base ourselves upon physiology and, so far as it throws light upon functions and functional disturbances, upon anatomy; for, obviously, pathological conditions are but extreme examples of physiological. Nav, more, it is useless to begin the study of our subject unprovided with a sound knowledge of physiology. Throughout this work, therefore, we would take for granted a knowledge of the main facts of physiology. Unfortunately, however, this is not always possible. More correctly, we should say that a knowledge of the physiology generally taught is taken for granted; certain branches of the subject must be considered with a fulness which is regrettable so far as, having regard to limitations in the size of our work, we thereby become compelled to condense our presentation of other matter, but is necessary, and in fact advantageous, so far as it tends to give the reader a more thorough comprehension of the meaning of morbid processes, and by laying down clearly the data upon which certain conclusions are based is indeed economical of space, since once laid down fully the briefest reference to general principles will suffice in later sections.

Why we have to dilate upon certain matters physiological becomes evident when we call to mind that physiology and pathology have for the last seventy-five years, at least, been divorced to this extent, that they have undergone development under separate influences. Under the influence more particularly of Ludwig and his pupils, physiological research has been directed to the study of organs and tissues. The organ as a whole has been taken into account. Results have been obtained by exact measurements—mechanical, electrical, or chemical of the work performed by one or other organ under varying conditions, with the result that nowadays we possess a rich store of data bearing upon what may be termed *mass effects*. Nor can it be denied that these methods have very materially advanced our knowledge of function and of the bodily processes.

But the oncometer, the pendulum myograph, the recording cylinder, and Kjeldahl's apparatus cannot be applied to the study of the individual cell units of which the tissue is composed. It is under the influence of another great master (Virchow) that modern pathology has been developed. His teaching was based upon exact study of diseased organs and the correlation between gross and microscopic appearances. It was largely histological, and, as a result, mass effects were followed back to the disturbances in the individual cells composing the tissue. In place of an organ, or tissue pathology, there was developed a "cellular pathology."

Thus, we owe it to Virchow that for now more than half a century pathology has held steadily before it the view that eventually the cell and the modifications undergone by it must be studied if we are to

INTRODUCTORY

understand aright the disturbances affecting the tissues and the organism as a whole. The cell, and not the tissue, is our unit.

We admit freely that for a long period after the publication of Virchow's great work comparatively little advance was made in our comprehension of cellular pathology. It is in science as in gold-mininga new "find" is announced, and workers from other fields forsake them and join in the rush. These other workers in pathology have, it is true, in the course of years contributed an enormous mass of facts, but the facts in the main have confirmed rather than advanced Virchow's observations. Virchow, indeed, employed his great influence in discouraging pathological speculation. He recognized that more facts must be accumulated before sure advance could be made. These facts preparatory to advance were in the meantime being accumulated, not by the pathologists, nor, again, by the physiologists, but by the zoologists, the botanists, and the embryologists-facts, namely, bearing upon the finer structure and the functions of the normal cell. It was left to a zoologist (Metchnikoff) to realize the bearing of these facts upon the cell in disease, and, by his studies upon the leukocytes, to emphasize the importance of the study and to develop new methods.

So strong was the influence of Virchów that through the last half of the nineteenth century pathology, as usually taught, consisted of little beyond the facts of gross and minute morbid anatomy. 'The text-books on our subject were devoted to the data of disease—to descriptions of the appearances (more particularly under the microscope) of the tissues and their component cells under various conditions of disease, and the abundant nomenclature in connection with the same. Attempts to explain and to generalize were reduced to a minimum.

It has taken many years—Metchnikoff's studies upon the leukocyte began in the early eighties—for a general realization of the bearing of these researches of Metchnikoff upon pathology and pathological research, and then only through the demonstration of their profound effect upon the related studies of the bacteriologist on infection and immunity. Now more than ever is pathology becoming truly cellular, even if, coincidently, it is widening itself in the opposite direction, and more particularly along chemical lines and through the study of disturbed metabolism is concerning itself with mass effects.

We admit freely also that the physiologists are, from other considerations, being simultaneously led back from tissue to cellular physiology. It is the natural course of events that, having established their science upon the reaction of organs and tissues as a whole, they should proceed to study the reaction of the component cells. As a matter of fact, we already possess at least one important work upon cell physiology, that, namely, of Verworn. But, excellent as it is, and suggestive, this is not yet generally read by the ordinary student; add to which, approaching the subject independently and from a different standpoint, we find ourselves not wholly in accord with Verworn over more than one matter of importance. Hence, since cell physiology is not given a proper place in the routine teaching of the student, and as this must be the basis of a

SCOPE OF THE WORK

cellular pathology, it is essential that we bring together and discuss in some little detail those facts bearing upon the life of the healthy cell, a knowledge of which, in our opinion, is necessary for an adequate comprehension of the changes which take place in the cell in disease. Associated with a study of the life of the cell we shall have to discuss certain phenomena—growth, adaptation, reserve force, heredity—which have an intimate bearing upon certain pathological processes, which again receive but the most summary treatment in the ordinary text-book of physiology.

Scope and Order of the Work.—These more physiological subjects we shall endeavor to deal with in an introductory section of the work. They will form the basis upon which we propose to develop our treatment of pathology proper.

As to this treatment: If, in its widest significance, pathology is the study of the functions of the body in disease (and conversely, we should add, of noxæ, or serious alterations in environment, as they bear upon the bodily functions), then clearly our subject embraces the whole field of scientific medicine, save and except prognosis and treatment. In other words, it has to deal with:

1. The causes of disease.

2. The course of disease (including the reactive processes on the part of the organism, whereby that course is guided.)

3. The results of disease.

Each of these main divisions can be approached and treated in at least two ways. Thus, on the one hand, forming or attempting to form a classification of diseases, we can discuss the *etiology*, or cause of each in turn; similarly we can describe the course of each separate disease, giving the symptomatology; and thirdly, we can note the results of each separate disease. On the other hand, studying all the conditions which cause disease, we can endeavor to *classify* the etiological factors, grouping together those influences which, acting on the organism, are seen to produce allied morbid conditions; similarly, from a knowledge of the course of various diseases, we can attempt to distinguish and describe certain morbid processes, one or more of which we recognize as underlying. and as common to, the course of individual forms of disease; and, coming to the results of the disease, instead of dealing with individual cases, we can discuss and classify the results of disturbed function upon individual organs and tissues, and attempt to gain a broad idea of how these local disturbances in one organ or tissue affect other organs and the organism in general.

The first of these methods is that of *special pathology*, so-called; it is the method employed in text-books of medicine and surgery; indeed, on the European continent, what we would term general works upon medicine and surgery are entitled text-books of internal and external pathology. Dealing, as is our purpose, with the broad principles of pathology, the second is the only possible procedure. We shall endeavor to arrange our matter in the order given—namely, to discuss first the causes, next the course (the morbid and reactive processes), and thirdly.

INTRODUCTORY

the results of disease. As will readily be understood, it is not always easy, nor indeed desirable, to discuss the causes of disease without at the same time indicating the processes which they originate; again, it is not possible to describe the morbid processes without indicating to some extent the results of the same. Nevertheless, this is, I feel assured, the only satisfactory and logical method of covering the vast territory before us. It is the means whereby the surest grasp is obtained of the principles underlying and governing those disturbances of vital activity which we recognize as disease.

This, we acknowledge, is not the course usually pursued in works upon general pathology, and to this extent is disadvantageous. By tradition starting with the morbid processes, inflammation and disturbances of the vascular system are first discussed. By our arrangement, inflammation is first considered in our third section, vascular disturbances not until the fourth. That more usual arrangement, we take it, is one of convenience coupled with prescription. When pathology is taught pari passu with medicine and surgery it is undoubtedly convenient that the student obtain a knowledge of such predominant processes as inflammation and infection as early as possible in the course, and when inflammation was regarded as essentially a vascular disturbance, it was natural that other vascular disturbances should be taken in close association. The result has been that the ordinary student has obtained an exaggerated idea of the importance of vascular disturbances, whereas what is of equal value for a general understanding of disease-namely, the pathology of the nervous system-has been relegated to the very end of the course, if, indeed, it has come in for treatment at all. The system has been imperfect. As regards convenience, we would point out that, conformably with our firm belief that the student should have a sound basis of physiology before entering upon the study of pathology, it is well that the matter contained in our first section should form a course in the "second vear"-whether given by pathologist or physiologist is a matter of indifference-and at the end of the second year a course on causation or etiology may well be introduced also. By this means the teaching upon the morbid processes comes to be given at the right momentnamely, when the student possesses his knowledge of physiology and is obtaining the first introduction to actual cases of disease in the hospital, and his first lectures upon "Internal and External Pathology."

We recognize, however, that the teaching and the periods of teaching of our subject vary greatly, and do not believe that the student should be a man of one book, mastering that book from alpha to omega. Hence we have arranged our material, so far as possible, in such a way that, while it forms a more or less progressive system, any one section or important subject becomes, by cross-references, etc., complete in itself.

We should add that, in order to afford illustrations of the different processes and their results, there are given, in the second volume of this work, that upon what may be termed *systemic* pathology, chapters dealing with the main outlines of special pathological histology. For

26

ON PRINCIPLES

these chapters, upon the special pathology of the different organs, the reader is indebted to our colleague, Professor A. G. Nicholls.

The Principles of Pathology.-One more word before embarking upon our course. What we have already said will have indicated that we have no narrow conception of the scope of our subject. The time has passed when morbid anatomy and morbid histology could be regarded as the sum and substance of pathological teaching, and when "to name his tools" was what was, in the main, demanded of the student. In the evolution of every science there are three stages to be recognized: The first, that of its dawn, when, from the observations of a few facts, the wide possibilities of the science impress themselves upon the worker and stimulate the imagination, so that forthwith he proceeds to indulge in wide hypotheses. These, in their turn, form the basis of further observations in order that they be tested, with the result that time and again they are found either erroneous or inadequate. With this, a second (reactive) stage becomes manifest, in which it is appreciated that before any sound generalizations or laws can be established the facts and data of the science must be carefully accumulated and marshalled. It is when this has been accomplished that the science can enter upon its third (complete) stage, in which an adequate knowledge of the factors involved permits the establishment of general laws. Needless to say these three stages are apt to overlap. At an early period sufficient facts may be at hand, or some clear-sighted observer may arise, to lay down with precision one or more secure generalizations, and so certain broad principles may be developed perfect at the very beginning, born complete, like Venus out of the sea foam. And, on the other hand, when, after much study and accumulation of facts, some other principle or law becomes established, its very establishment leads us to further possibilities, to the study and recording of yet other phenomena. For knowledge is progressive, and, while one department of science may have reached its third stage, other departments may scarcely have entered upon the first.

A study of the history of medical science-of pathology-amply bears out this statement. Galen, indeed, may be taken as the archetype of those who, conscientiously systematizing the known facts of a subject, proceed to develop a system which is unavoidably false, because the facts on which it is based are too few. He is at the same time a striking illustration of the value of even such necessarily imperfect work as he accomplished-for it cannot be denied that his efforts created a medical science, and that for a period of fifteen hundred years he and his system dominated the medical world. With the medical renaissance in the sixteenth century the gaining of new facts from experimental observation led to the downfall of the Galenistic philosophy-it was tested and found wanting-and led forthwith to a succession of hypotheses, as one series of observations followed another. The new development of mechanics and physics led to Borelli's investigations upon animal movements and the appearance of the *iatromechanical* school, which explained animal activities on a purely mechanical or physical basis. The remarkable experimental ability and genius of van Helmont led to

INTRODUCTORY

the recognition of ferments; his no less remarkable imagination led to extraordinary speculations, some wholly wild, others so prescient as only nowadays to be found to approximate to the truth. The more immediate outcome was the development of the *iatrochemical* school under Sylvius, in which, more particularly in connection with digestion and respiration, chemical processes were seen to be at the basis of animal activities. And so, to the middle of the nineteenth century, school succeeded school-mechanicodynamical, Brunonian, vitalistic, Cullen's system, and so on; the last of influence, more particularly in Germany, being that of Oken. System after system was overthrown as successive observations demonstrated their inadequacy. Virchow represented the revolt against all such. His teaching was that hitherto theories had been founded on insufficient data; the time had come to gather facts and cease theorizing; and so potent was his influence that we have had the remarkable spectacle of the workers in pathology among that most philosophical of peoples, the Germans, restraining themselves for fifty years from philosophizing, and sedulously bending themselves to accumulate facts and record details. And the same strong influence has told upon the pathologists of all countries.

During these fifty years the amount of material collected has been extraordinarily great—in fact, overwhelming—so much that no one individual can pretend to master it. Never has there been such a period; much more has been garnered than in all the preceding ages together. If in 1858⁴ the data were inadequate, today we experience the contrary danger of being overcome and blinded by excess of detail. The time has surely arrived to attempt to systematize our knowledge and so to order it that each new fact acquired is seen to have its place and to exemplify some general principle. Pathology, we hold, is now ripe to enter, and has entered, upon its third stage of development. It is with this opinion in mind that we have written the pages that follow.

¹ The year of publication of Virchow's Cellular Pathologie, and of Darwin's Origin of Species.

CHAPTER II.

THE HISTOLOGY OF THE CELL.

As already stated in our introductory chapter, we owe it to Virchow that, for more than half a century, pathology has held steadily before it the view that eventually the cell and the modifications undergone by it must be studied if we are to understand aright the disturbances affecting the tissues and the organism as a whole. The cell, and not the tissue, is our unit. Modern pathology, however, demands a much fuller study of the cell and its activities than is usually afforded in the physiological course. From this it follows that if we are to grasp the more recent advances in the study of disease and of the reactions to the same, some little space and time must be afforded to a consideration of the cell and its properties. Such facts as are matters of common knowledge may be passed over rapidly; others must be dwelt upon.

THE CONSTITUENTS OF THE CELL.

The animal cell, we recognize, consists of two main parts, the cell body and the nucleus; and this differentiation is clearly of great antiquity, for only the very simplest forms of life, whether animal or vegetable, do not show it. Even in these very simplest forms, if a sharply defined nucleus be not present, we have evidence that nuclear material exists. It used to be taught that the lowest "animal" forms-Haeckel's order of Monera-and the lowest plant forms-the Schizomycetes, or bacteriaare non-nucleated; but with the elaboration of more perfect staining methods, either a definite nucleus has been determined in those of the monera so far examined for this purpose; or again, in some, as shown by Gruber, granules of nuclear material are to be seen scattered through the cell body (Fig. 4). As regards the bacteria, there are parallel observations, observations which at first sight appear contradictory. Thus, Bütschli, from his observations, regards all the stainable substance of the bacterial body as equivalent to the nucleus, and studying some of the larger forms demonstrated the existence of a fine surrounding substance. apt to be gathered at the poles, which he holds to represent the cell body or cytoplasm. This view is confirmed by other workers. Several more recent observers, dealing with the smaller forms, figure minute granules of stainable material diffused through the bacterial body. These they regard as nuclear material, and call attention to a characteristic rearrangement of these granules prior to fusion, which suggests strongly the "quadrille" of the nuclear chromatin preceding cell division in the cells of higher forms of life.

A full study of the divergent observations upon this subject up to the year 1900 is given by Professor A. B. Macallum.¹ Macallum denies that these lowest forms have a nucleus—that is, a sharply defined and well-differentiated organ presenting characteristic changes in cell division. Herein we cannot but agree with him. He admits, however, the existence of substances allied to the nucleus and to chromatin, substances containing phosphorus and "masked" iron, whether present in a "central body," as in the *Cyanophycew*, or as scattered granules. There is nuclear material, but no proper nucleus. Other more recent studies favor this view.

Both series of observations would thus seem to be correct; in these lowliest forms there is no nucleus proper, although nuclear material



Absence of nucleus proper, with diffusion of scattered granules of nuclear material through body of a *monera*, (After Gruber.) exists. We may thus lay down that in all forms of life, animal and vegetable, from the lowest to the highest, nuclear and cytoplasmic matter co-exist in the cell. To the significance of their co-existence we shall revert later.

In the cells of man and of all save the lowest forms of life it can be seen that neither the nucleus nor the cell body is homogeneous. We are still very far from having resolved the minute anatomy of either. That anatomy, indeed, is beyond the power of the microscope to determine.

This much may here be laid down, that the nucleus is a well-defined body, most often approximating in shape to the spherical or oval; at times greatly elongated (as in plain, non-striated muscle tissue); at times irreg-

ular and lobate (as in the polymorphonuclear leukocyte); at times moniliform or beaded, as in *Stentor*; or, rarely, extensively branched, as in certain gland cells of insects, or possessing pseudopodia-like processes, as in the egg rays of the water scorpion.² In the higher animals, a more or less distinct nuclear membrane is to be made out, within which the substance exhibits an alveolar or netted arrangement. Of this nuclear matter, at least three constant constituents are to be distinguished: (1) The *linin* or achromatic (non-staining) network in which is deposited (2) the *chromatin*, or material which is rendered noticeable by nuclear dyes. In the spaces of the network is (3) the *nuclear fluid* or sap. Inconstant features are (a) the *nucleous*, a minute accumulation of chromatin-like substance, varying in amount, which nevertheless takes on certain differential stains, and hence would seem to have a somewhat different composition; (b) *vacuoles*—these are very rare; they are to be seen in the nuclei of fat cells; (c) definite crystals; these have been recog-

¹ On the Cytology of Non-nucleated Organisms, University of Toronto Studies; Physiological Series, No. 2, 1900.

² Vide Korschelt, Naturwissensch. Rundschau, 18; 1887; 409.

nized by Marchand, Podwyssozki, and others. These last two constituents indicate that the nucleus is not inert—that active metabolic processes occur within it.

Of these constituents, the nucleolus has of late received increased attention. Nucleoli are spherical bodies, apparently fluid or colloidal, exhibiting no structure. They vary greatly in size; in some orders of cells they are single, in others multiple. They may be relatively very large; thus, in the oocyte of Antedon bifida (a crinoid "starfish") this single nucleolus may be half the diameter of the nucleus (Chubb). Or, on the other hand, there may be multiple minute nucleoli situated at points of intersection of the chromatin network. Of these, there may be as many as 300 (Montgomery; unicellular glands of the leech, (Piscicola). All observers agree that they are intimately related to the chromatin. Thus, as M. Heidenhain points out, they are absent from the feebly staining nucleus of the young cell immediately after cell division, but grow pari passu with the development of the nucleus and appearance of increased chromatin. In general they take on a differential staining with nuclear (chromatin) dyes, and so are not chromatin proper; at times, however, they take on the same stain. The main opposing views: (1) That they are waste products of chromatin activity (Häcker); (2) that they supply nutriment to the chromatin (Flemming, Bambeke, Korschelt, Lubosch); (3) that they are the means through which the unorganized chromatin assimilated from the cytoplasm is converted into the organized chromatin of the nucleus (R. Hertwig); and (4) that they represent the dissociation product, rich in albumin but devoid of phosphorus, of the nucleo-albumins taken up from the cytoplasm to provide the nucleoproteins of the chromatin (M. Heidenhain¹)-all these may be harmonized by regarding the nucleolus as store or reserve material from which, on the one hand, assimilated material is withdrawn to provide (by dissociation) the chromatin constituents needed by the growing or regenerating nuclear network, and to which, on the other hand, are passed the products of chromatin disintegration. We must, that is, accept for the nucleus as for the cytoplasm the general principle, to which attention must be frequently drawn in these pages, that proteid matter tends to present like intermediate products in the opposed conditions of integration and disintegration. We shall have occasion shortly to discuss further nucleolar function (p. 47).

The type cell has but a single nucleus; this is the general rule, but it is not uncommon in active glandular tissues to find cells with two, and other instances occur in which cells are multinucleate. Two processes are possible: either with growth the nucleus may divide without subsequent division of the cell substance, or what had been separate uninucleate cells may fuse, forming a *plasmodium*.² We have evidence that

¹ For bibliography, see Chubb, Phil. Trans. Royal Soc. Biol., 98: 1906; 447, and M. Heidenhain, *Plasma und Zelle*, Fischer, Jena_r 1: 1907; 212.

^a Some writers draw a distinction between the syncytium, caused by the fusion of cells, and the *plasmodium*, caused by nuclear multiplication without cell division; others include both forms as plasmodia. both processes take place, and shall have to refer to these matters more fully in discussing *giant cells*.

As already indicated, in some of the simpler forms of life nuclear material is scattered through the cell substance. In higher forms, this nuclear material, as a rule, is strictly confined within the nucleus. But it has to be laid down that this is far from being absolutely constant. In certain cells and under certain conditions minute masses of chromatin are to be detected lying in the cell body away from the nucleus. These and other not so definitely *chromidial* substances have been shown by




Diagram of component parts of a cell illustrating the various theories of the structure of the cytoplasm. The lower segment represents the *ibrillar* or *sponge* theory; the upper, the granular theory; the left, the *loam* theory. At the right the protoplasmic threads (archoplasm) radiate from the centrosome. (Szymonowicz.)

numerous observers to be derived and discharged from the nucleus. Of late years there has been an active study of these nuclear products, which appear to play an important part in cell activities and to their relationship to nucleolar matter. We shall refer to them in fuller detail later.

The cell substance also exhibits indications of structure. Without there being in the animal cell the distinct membrane or wall so conspicuous in many vegetable cells, there is, nevertheless, often to be recognized a condensation of the cytoplasm or cell substance at the periphery, homogeneous, and constituting the *ectoplasm*, which passes almost imperceptibly into the main mass of cell substance, or *endoplasm*. This endoplasm is seen by careful study to possess, like the nucleus, an

32

PARAPLASM AND CELL GRANULES

alveolated arrangement, regarding the exact nature of which cytologists are still at variance, it being most difficult, in the first place, to translate optical appearances under high magnification; in the second place, to assure ourselves that what is seen represents the natural conditions of the living substance, and is not a secondary effect brought about either by the death of the cell or by the action of reagents (Fig. 5); thirdly, save for deposits within it, the cell substance is obviously fluid; we have to differentiate between portions that are freely fluid and those which, if of greater density and more viscid, are still of fluid nature. We shall not here enter into the controversy which has raged, and continues to rage, regarding this matter, but would lay down the general conclusions reached, which are that the cell substance consists of (1) a coarse or finer reticulum, which may be termed the cytoplasm proper; (2) the cell sap or fluid lying within the meshes of the reticulum; and (3) paraplasmic substances. Under the term paraplasm we include (a) granules of solid matter taken up by the cell by phagocytic action and not yet dissolved or discharged; (b) granules of solid or semisolid matter, crystalline or amorphous, which are the products of cell metabolism; (c) the fluid contents of secretory vacuoles; and (d) inactive substances laid down as a framework within the cell. Passing beyond purely histological appearances, we may say that the cytoplasm is the active cell substance (termed by some the *bioplasm*), though it must be kept in mind that this term also includes, if, indeed, it should not be confined to, the active substance of the nucleus; the paraplasm, all material, whether in a dissolved or precipitated form, which is within the cell, and represents matter resulting from the cell activity, whether products of disintegration of the cell substance or the unassimilated portions of absorbed material.

Regarding the granules constantly present within the cell substance, it deserves mention that some have given them a much more prominent position than is here ascribed to them. By special methods of staining, a fine granulation of the cell substance can be clearly demonstrated. and Altmann¹ has regarded these as the "elementary organisms, or bioblasts," as the units or fundamental elements in cell activities. Apart from the indications being, as we shall endeavor to show, that the primary cell activities are inherent in the nucleus, further study shows that Altmann's⁵ methods bring out granules of more than one order, and that there is no apparent relationship between the number of these granules (which varies greatly) and the active nature of the cell. All that Altmann has accomplished has been to call attention to the existence of these granules; he has not brought forward a single valid argument in favor of their possessing the attributes with which he endows them. While his facts are accepted, his interpretation of the same is now generally discredited.

There is another important structure in the cell substance, important in that it is seen to be actively engaged in the process of cell multipli-

¹ Die Elementärorganismen und ihre Beziehungen zu den Zellen, Leipzig, 1890.

- 3

THE HISTOLOGY OF THE CELL

cation. This is the *centrosome*, a minute dot or area of condensation surrounded by a fine areola, generally situated toward the centre of the cell in the neighborhood of the nucleus; in some rare cases it has been described as actually within the nucleus. It forms a centre from which, prior to cell division, the cytoplasmic substance becomes arranged in fine rays, and, even before the nucleus, it undergoes division. In the resting stage of the cell it is not constantly recognizable, and in some it has not yet been made out. Regarding its nature and relationship thère has been keen debate—whether it be an independent constituent, carried over by division from cell to cell, just as are the nucleus and the cytoplasm, or derived from the nucleus or from the cytoplasm. Martin Heidenhain,¹ for example, in a singularly full study, has suggested that it is the homologue of the *micronucleus*, of the infusorian cell. (The infusoria have two nuclei, of which the larger, the *macronucleus*, is most in

Ph. 0



Section of normal tubule of kidney, stained to show the regular arrangement of Altmann's granules in the cells, (After Altmann.)

Leukocytes from larva of salamander, showing centrosome with surrounding cytoplasmic radiation, Flemming.)

evidence in the functioning organism, but disintegrates and disappears during the process of conjugation and fertilization, the micronucleus then becoming active.) More recently, Yatsu,² confirming E. B. Wilson, has shown that if the eggs of *Cerebratulus* be cut up, and fragments, devoid of nuclei and centrosomes, be placed in sterilized calcium chloride solution and then in sea water, a centrosome with surrounding aster develops in them identical with those of whole eggs subjected to the same treatment. From which it is evident that the centriole or centrosome can be formed *de novo* from the cytoplasm—that it is of cytoplasmic origin.

¹ Arch. f. mikr. Anat., 43:1894:423; see also Arch. f. Entwickelungsmechanik, 1:1895:473.

² Proc. Soc. Exptl. Biol. and Med., 2: 1905, L;si, and Amer. Med., 1905;493.

CELL CONNECTIONS.

These are the main histological details regarding the constitution of the animal cell. But if our pathology is to be cellular, more is needed. The organism being multicellular, but derived from a single cell, it is necessary to have a definite conception regarding the histological relationship between the individual component cell units, and this because of the light this must throw upon the dependence of the cells one upon the other in disease as well as in health.

The usual conception of the organism, we take it, is that it is an accumulation of cells which are distinct separate entities, acting the one on the other, either by their products, or by physical influences, through conduction. The general idea is that the multicellular organism has developed primarily from the unicellular as an aggregation of separate unicellular units which have remained associated for mutual protection and benefit, the separate units undergoing differentiation as a result of relative position, and so of environment. Such a conception has induced a false view as regards what constitutes the individual, and to some extent as regards the relationship of the tissues one to the other.


We owe to the botanists, first among whom must be mentioned Walter Gardiner, the demonstration that in the multicellular plant the individual cells are not isolated and wholly detached, but are united to each other by fine bridges. It has been proved by them that by the intermediation of these bridges stimuli are directly conveyed from cell to cell. Now, in the animal body, it is becoming proved for most tissues that the cells are similarly connected. The cogwheel-like appearance of the cells of the epidermis was for long suspected, and has now been proved, to be the indication of a system of filaments passing from one cell to its neighbors, and Kolossow¹ has shown, and MacCallum, of Baltimore, has confirmed, that similar bridges pass between apposed cells of the endothelia. Ciliated and columnar epithelial cells are likewise joined together (Barfurth²) and similar direct connection has been described between the cells of both plain and striated muscle (Kultschitzki³). Even the neurons, which have been regarded so generally as independent cells, have now, by Apathy and others, been shown to communicate by extremely fine filaments. The difficulty in accepting Apathy's results has been due as much to the want of recognition of this general principle of cell connection as to the prevalent theories of nervous action.

These filamentous cell connections are evidently present from the very earliest period of individual existence. Mrs. Andrews,⁴ studying

¹ Arch. f. mikr. Anat., 42; 1893; 318, and Ztschr. f. wiss. Mikr., 9, Heft 1 and 3; see also W. B. MacCallum, Johns Hopkins Bull., 14: 1903; 105, and Muscatello, Virch. Arch., 142: 1895; 327.

² Anatomie, 1897; 79. ³ Quoted by Waldeyer, Areh. f. mikr. Anat., 57; 1901; 2. ⁴ Mrs. G. F. Andrews, Jour. of Morphol., 12; 1897, and E. A. Andrews, Zoöl. Bull., 2; 1898; 1; confirmed by C. Shearer, Proc. Roy. Soc., B., 77; 1906; 498, who gives recent literature. The fullest study of this subject is by Schüberg, Zeitschr. f. wiss. Zoöl., 74; 1903; 155. the recently laid eggs of echinus and the starfish, and employing very high powers, noted that in the process of cleavage, while momentarily the blastomeres become separate, there follows an active discharge of fine threads across the intervening space, resulting in the union of the

F16. 8



Cell bridges of "prickle cells" of epidermis. (From a photograph by Schridde.)

cells by protoplasmic processes. These could well be seen in the 8- and 16-cell stages, and, with their formation granules could be seen streaming from the one cell to the other.

In short, it may be laid down that the absolutely detached cell is the exception and not the rule. The leukocytes—the wandering cells of the organism—are wholly independent; but this, judging from MacBride's¹ observations, is an acquired and not a primary characteristic. In the larval echinoderm (starfishes and the like), whose wandering meso-

dermal cells are the earliest homologues of the leukocyte, these cells floating in the body cavity are found to be connected by a network of singularly fine processes.

F1G. 9



Cell bridges of vascular endothelium. (Kolossow.)



Cell bridges between cartilage cells,

The organism or individual, therefore, is not to be regarded as essentially a conjugation or colony of detached units, but rather as a connected whole, in which, for reasons to be immediately discussed, there has been partial and incomplete division of the living matter; the cells in general are not detached, only semidetached.

The Significance of the Cell.-This

conception leads up to a comprehension of the nature and significance of the cell. Studying this, whether among the unicellular individuals or in its many modifications among

¹ Proc. Camb. Phil. Soc., 9:1896:153,

THE SIGNIFICANCE OF THE CELL

the multicellular organisms, one is struck by the fact that it in general is of minute size. The exceptions—the cases in which single cells are large enough to be visible to the naked eye—are of three orders:

I. Where the cells contain a large amount of stored-up food material. This is notably the case with the ova of very many species. Here, on fuller study, it is found that the cytoplasm forms a delicate membrane, spreading over and limiting the yolk. In this superficial layer lies the nucleus, which thus, with the cytoplasm, is close to the exterior.

2. In certain of the infusoria the increase in size of the cell, until it becomes visible to the naked eye, is brought about by the development of the cell substance into a series of delicate radiating processes. By this means neither the nucleus nor any part of the cell is at a distance from the surrounding medium.

3. As in *Gromia* and sundry other protozoa, the enlargement of the cell may be associated with the presence of multiple nuclei.⁴

In all these cases we appear to have a mechanism whereby no portion of the cytoplasm is remote either from the external medium on the one hand or the nucleus on the other. There is clearly a relationship as regards size between the nucleus, the cytoplasm, and the surrounding medium, and this is determined primarily by the size of the nucleus. As we shall point out, the evidence is conclusive that the nucleus is the dominant controlling element in the cell; it governs the cell body and cytoplasm; the larger it becomes the greater the disproportion between its surface area and its mass, and, as interaction between it and the cytoplasm must in the main take place at the surface, the greater its size the less its relative efficiency. If the nucleus exceed a certain size, the more centrally situated nuclear material must be largely inactive, and so useless. Thus, it has come to pass that that living nuclear material has been most favorably circumstanced, or, in other words, best fitted to survive, which has undergone division and so increased its surface area at the same time as it has augmented its substance and mass. Thereby the maximum activity of the nuclear material has been insured. Hence the development in the first place of the multinucleated cell.

But if this be true of the interaction between nucleus and cytoplasm, it obtains also between the cytoplasm and the external medium. Here, again, we have to deal with surface action. It is obvious that, as in the radiolaria, the cell surface can be enormously increased by production into a large number of fine processes, but if, as already indicated, the cell activities are determined by the nucleus, by the reaction between nucleus and cytoplasm, such extension of the cytoplasm to a distance from the nucleus has its disadvantages. The most economical system is the spherical; and possibly as the result in the main of surface tension —by the same law as that causing the globular form of the rain drop or detached small mass of mercury, it is noticeable that free cells in general approximate in shape to the sphere. Here, again, we have the

¹ A stage leading up to this is seen in Stentor and other ciliated protozoa, in which the relatively large nucleus is moniliform, beaded, and elongate. same considerations of economy of action. Materials are absorbed and built into the cell substance from the external medium, and as the process of absorption and formation of new cytoplasm proceeds, the mass of the cell increases in a greater ratio than does the surface, until the point is reached at which the accumulation of inactive cytoplasm is subversive to proper action. The same processes that induced nuclear division have brought about cell division.

We thus recognize the following successive stages:

1. The cell or mass of living matter in which the nuclear matter is scattered through the cytoplasm.

2. The unicellular organism having the nuclear matter aggregated into a central mass, the nucleus.

3. The multinucleate unicellular organism.

4. The multicellular organism.

It follows, thus, that the multicellular organism is not to be regarded as the outcome of the fusion of a number of separate individuals for mutual advantage. Such fusion, it is true, does occur in nature; witness the myxomycetes. It is, however, the exception, and is not found along what we regard as the direct line of vertebrate ancestry. This communal, or, as the Germans term it, "Baustein" theory, must be replaced by one more directly in harmony with the facts of individual development and our knowledge of evolution—by what we may term the theory of decentralization, which regards the individual as the sum total of protoplasmic matter capable of existing as an entity under particular conditions of environment, the multicellular individual acquiring its greater size and more complete activities by means of nuclear division followed by cell division.

As regards the nuclei, this division is complete, and as the nucleus is we hold, the primary and controlling structure in the cell, to this extent each cell is an independent entity; as regards the cytoplasm, as stated (p. 35), the separation is incomplete, and to this extent the individual is a single connected whole. But, while making this statement, it must be borne in mind that the nucleus cannot persist without the cytoplasm; that there evidently is an intimate relationship between the two such that the nucleus is acted upon not directly by the external medium, but through the intermediation of the cytoplasm. From this it follows that cytoplasmic alterations, if conveyed from cell to cell, are capable of influencing the nuclei; these latter may control the individual cells, but are at the same time capable of being influenced by the cytoplasm. This conception of the relationship of the cells and tissues in the multicellular organism is fitted, we think, to throw light upon the otherwise somewhat paradoxical coincident independence and interdependence of the cells, to which we have already referred in discussing what is disease (p. 20), which must thus be regarded as a primal or necessary outcome of cell relationship.

Intercellular Substances: Are They Living Matter?—It has to be confessed that of late years the cell theory has from different quarters encountered searching criticism. To some of this, as, for example, to the

THE INTERCELLULAR SUBSTANCES

non-existence of a discrete nucleus in the very lowest forms of life, we have already incidentally referred. The Baustein theory above mentioned has introduced other difficulties: What is the need, and what the evolutionary process by which individual cells have become joined to form distinct organs when there exist sundry higher unicellular organisms among the ciliate infusoria which, possessing a single or two nuclei, possess also distinct organs, such as contractile internal fibrils (muscle), a peristome of cilia directing food to a distinct mouth region, excretory organs (contractile vacuoles), organs of defence (trichcoysts)? The answer to this question is difficult in terms of the communal, simple in terms of the decentralization hypothesis. But there are those whose opposition is directed against the very being of the cell theory. If it can be demonstrated that there is living matter in the multicellular organism which is external to cell boundaries, then this cell theory, if not wholly upset, demands very material modification.

What is the nature of the intercellular substances, of which there are several in the tissues of the higher animals—the fibrils of white connective tissue, the fibres and sheets of yellow elastic tissue, the matrix of cartilage, the lamella of bone? Are these alive, or are they to be regarded as dead matter? Recently a leading German histologist, Professor Martin Heidenhain, 'has claimed that they must be regarded as living matter—as extracellular living matter.

According to him, these exhibit metabolism, growth, formative energy, and perhaps, also, a definite grade of functional activity. The fibrils of connective tissue and the coarser deposits of elastin in yellow elastic tissue begin in general within the cell. This we freely admit. Later they come to be extracellular and freed from the cell body proper; they increase in length and also in bulk. A similar origin is upheld for the cartilaginous matrix; that, according to Schaffer, begins as a modification of the ectoplasm of the cartilage cells, and as the cells shrink and this modified ectoplasm, derived from one cell, fuses with that of neighboring cells, so does the matrix become a homogeneous mass, in which are embedded what are now sharply defined cells. Nay, in this matrix is also living matter of another order, since study shows running through it fine fibrils of connective-tissue type. The living matter of the bony matrix, he admits, has not been the subject of adequate study, but that it exists he has no doubt.

To this it may be replied with, I think, considerable force:

1. That, from a chemical and physical point of view, the albuminoids, or scleroproteins, which form the characteristic constituents of matricial matter—collagen, elastin, and chondrin—are of all the proteins of the body the most inert; in characters they are most nearly allied to the coagulated proteins, which are obviously "dead;" that, knowing their insolubility in various reagents, and the difficulty with which they are dissociated, it requires a severe stretch of the imagination to conceive these bodies as possessed of the power of recurrent satisfaction and dissatisfaction, which we regard as the prime attribute of living matter.

¹ Plasma und Zelle, Fischer, Jena, 1:1907.

2. That, physiologically, interstitial matter is strikingly inert, exhibiting nothing that we can regard as a direct reaction to irritation; what reaction is shown—dissolution, etc.—is obviously determined by the enclosed cells, *i. e.*, it is in the immediate neighborhood of cells exhibiting obvious reaction that changes are first to be noted in the matrix as the result of irritation.

3. That because the fibrils, the cartilaginous matrix, etc., show their earliest signs of development within the cell body, that does not, *ipso facto*, make them living matter, any more than intracellular fat globules are to be regarded as living matter.

4. That, as regards actual growth of the connective and yellow elastic fibrils upon which Heidenhain lays so much emphasis, we obtain, as Professor Leo Loeb has shown "growth" of a curiously similar type outside the body in lymph or blood subjected to strain. When a drop of uncoagulated lymph is placed between two glass slides, the mere act of pulling the one slide over the other leads to the appearance of fibrils, which grow in length and bulk; which, like those of connective tissue, are not only intracellular, but actually traverse cell bodies situated in their path; which show themselves first in immediate connection with these cells, the cells, as we now hold, liberating an enzyme that determines the modification. But the lines of this precipitation are evidently along the lines of strain. And so identically do we observe that the direction of the individual connective-tissue fibrils in a tendon, a fascia, etc., bears a direct relationship to the strain to which the tissue is habitually exposed.

5. That if cartilaginous matrix is to be regarded as living, then, also, the hyaline and amyloid deposits in pathological conditions are to be regarded as living. But among these we have every transition to conditions of deposit in successive layers, to conditions clearly of precipitation, and not of growth by intercalation or progressive building up of new molecules in immediate association with the old; the process is of a passive, not an active, type; nay, more, as Ophüls has shown, it may take on the type of deposit of successive layers of needle-like crystals of hyaline matter.

In his Cellular Pathology (1858, p. 23), Virchow laid it down that the intercellular substances had not life of their own, "but borrowed or obtained vital properties at second-hand from the (associated) cells." Heidenhain, with justice, points out that here is a basal mistake. The concept of life is that it is something inherent, something automatic. There is no such thing as one individual instilling or breathing life into inanimate matter outside itself. Yet, Virchow, despite his illogicality, was, it seems to me, nearer the truth than is Professor Martin Heidenhain. What is to be our primary postulate regarding the essential distinguishing property of living matter? If we lay down that it is growth—is the inherent power of assimilating other molecules, of so arranging them as to build up like proteidogenous molecules, possessing, therefore, like properties—then it must be held that intracellular substances have not

THE CELL THEORY

this property, and are not living. If, on the other hand, we declare that growth is not essential, and recognize three orders or grades of living matter, then, with Virchow, we can well admit that the cells discharge living molecules of the order of enzymes which act upon and modify the surrounding matrix. But in this case it is not the matrix itself that is living, as claimed by Heidenhain, but the discharged cell molecules, which act upon substances present in the surrounding lymph. And it has to be admitted that molecules of the same order acting within the cell can similarly act upon assimilated substances of the same order, and lead there to the production of the like precipitum. We fail to see, therefore, any necessity to alter our views regarding the intimate and essential relationship between the aggregation of matter into the form of cells and vital activities. The "cell theory" is still to be accepted as a foundation stone of our conceptions of living matter, even if we replace the older Baustein theory by one that regards the individual as the main unit, and the cells of which that individual is composed as means of bringing the living matter of that individual into an economical relationship with its environment.

CHAPTER III.

THE PHYSIOLOGY OF THE CELL.

WE have stated that we regard the nucleus as the controlling constituent of the cell. Here it will be well to indicate the grounds upon which this view is based, more particularly because this view is not universally accepted, but is apt to be propounded with some hesitation in works upon physiology, and because a correct appreciation of the influence of the nucleus is, as we shall repeatedly have to indicate, of primary importance in the study of morbid processes. It is only of late years that the attention of pathologists has been attracted to nuclear changes; only, in fact, after the cytologists had estab ished a basis of knowledge regarding the normal nucleus did it become possible to study departures from the normal.¹

It has, in the first place, been fully established that without a nucleus, growth and reproduction of the cell cannot occur. The cell, deprived of its nucleus, can exist for a time, can be the seat of certain metabolic activities, but its substance is progressively used up, and, judging from its complete incapacity to reproduce itself, it cannot form new living material, either cytoplasmic or nuclear. The red corpuscle, for example, the type of non-nucleated cell in the normal vertebrate organism, can act as a carrier of oxygen, but cannot perpetuate itself. The individual erythrocyte, once it is liberated into the blood stream, has but a limited period of life. Hunter² estimates that the red corpuscles of the rabbit live, at most, three or four weeks. Quincke³ and Worm Müller⁴ give a life of about fourteen days to the red corpuscles of the dog. Throughout life there is a constant development of new erythrocytes to take the place of those undergoing disorganization.

What is true with regard to the red corpuscles has been experimentally proved with regard to unicellular organisms. Brandt,⁵ in 1877, showed that pieces of *Actinospharium Eichhornii* containing a nucleus assume a characteristic form typical of the whole organism; those without a nucleus fail to do so. With *Siphonocladus* (another simple multicellular form), Schmidt,⁶ many years ago, showed that when broken up the protoplasm formed into spherical masses; those not having a nucleus failed

¹ For a fuller statement of these views regarding the dominance of the nucleus I would refer to my address at the meeting of the British Medical Association at Toronto, in 1906. Brit, Med. Jour., 2: 1906.

² Brit. Med. Jour., 1887; ii: 192.

³ Deut. Arch. f. klin. Med., 25: 1880: 567, and 27: 1880; 193.

⁴ Transfusion and Plethora, Christiania, 1875; see also von Ott, Virch. Arch., 93; 1883; 125.

⁵ Ueber Actinospharium Eichhornii, Dissert., Halle, 1877.

⁶ Festschr. d. naturforsch. Gesellsch., Halle, 1879; 3.

THE SIGNIFICANCE OF THE NUCLEUS

to produce a surrounding membrane, and soon disintegrated, while those containing one or more nuclei developed into the typical organisms. Fuller confirmatory results were gained by Nussbaum¹ with *Oxytricha*, and by Gruber² with *Stentor*. Klebs³ noted that enucleated cells of alge, like spirogyra, might live six weeks, and during that time might produce new starch granules—might, that is, synthesize starch from the carbon, oxygen, and water absorbed. This starch was formed in the sunlight and used up in the dark. Notwithstanding, unlike nucleated portions of such cells, these enucleated portions produced no cellulose wall, and disorganization and death were inevitable.

In studying the degenerations and necroses affecting the tissues we shall see that so long as the nucleus is intact, however grave may be the disturbances in the cell body, the indications are that cell regeneration is still possible; nuclear disintegration is the most obvious sign of cell death.

Enough has been said to indicate that the nucleus is essential for the continued growth of the cell. There is not, to our knowledge, a single observation to the contrary. It is, however, worthy of note that, as Boveri⁴ and Lillie⁵ have pointed out, there is a minimal limit to the size of the separate (nucleated) cell portions capable of undergoing further development.⁶ The nucleus, with the surrounding cytoplasm, is capable of regeneration and growth, provided that the amount of cytoplasm exceeds a certain minimal volume relative to the normal cell. For, as Verworn⁷ was the first to emphasize, the nucleus without the surrounding cytoplasm is as incapable of regenerating the cell as is the cytoplasm without nucleus. Nevertheless, Verworn takes a position which is untenable. He admits freely that cell growth and reproduction are not possible in the absence of the nucleus, and that the nucleus plays an essential part in such conditions as the formation of cellulose by the plant cell, the formation of chitin in the insect cell, sundry secreting processes in gland cells of higher animals, and that the remarkable change in the size of the nucleus during cell life can be brought about only by the nucleus receiving substances from the protoplasm and giving off others to it. He, however, denies wholly that the nucleus is the dominating portion of the cell, pointing out that, although the spermatozoon, in fertilization, introduces a minimal amount of cell substance into the ovum, and is composed, as regards its functional head, almost wholly of nuclear matter, nevertheless that minimal amount is introduced and cannot be neglected;⁸ that if the cell without nucleus cannot exist neither can the nucleus without cell substance, and demonstrates abso-

¹ Arch. f. mikr. Anat., 26: 1886: 485.

² Biol. Centralbl., 4:1885:717; 5:1885:253; and 6:1886:1.

³ Ibid., 7: 1887; No. 6, und Unters. a. d. botanisch. Inst., Tübingen, 1887.

⁴ Arch. f. Entwickelungsmech., 2:1895:394. ⁵ Jour. of Morphol., 12:1896:241.

⁶ In the cases studied, about one-twenty-seventh of the whole mass.

⁷ General Physiology, translated by F. S. Lee, Macmillan, 1899, 504 et seq.

⁸ But, as Strasburger points out, in the flowering plants the male sexual cells lose their cell-body when passing down the pollen tube and the nucleus only reaches the egg. (Durvien and Modern Science.) Cambridge, 1009; 104.

THE PHYSIOLOGY OF THE CELL

lutely from his studies upon the ciliated infusorian Lacrymaria olor that the nucleus does not control the motor apparatus of the cell—that non-nucleated sections of the organism move as actively as do nucleated sections, and this for a day, sometimes for several days. What he proves is not that the nucleus is not the dominating portion of the cell complex, only that the association of nucleus and cytoplasm is essential for full cell activity. He thus fails to grasp the significance of the nucleus, and his whole treatment of cell processes, if not vitiated, is, at least, greatly weakened.

All that Verworn's facts prove is that nucleus and cytoplasm are equally essential for the full function of the cell, not that they are of equal value. We might as well argue that in the community of bees the individual drone or worker is of importance equal to the queen bee, on the ground that, separate the queen bee from the rest of the community, and, being incapable of obtaining food for herself, she starves to death. Under no condition, that is, can the developed worker continue the race; this all-important function belongs to the queen bee, and to her alone. This simile, it is true, must not be pushed too far; advanced thus far, it will, however, illustrate our contention. The necessary association between nucleoplasm and cytoplasm does not contradict the evidence we possess that in the nucleus reside the controlling activities of the cell. Taking that evidence into account, it proves that the nucleus connot directly act upon the surrounding medium, and that so the function of the cytoplasm is to act as an intermediary between the nucleus and the environing substances.

The prominent part played by the nucleus in simple cell division; the series of processes insuring that each daughter cell obtains an equivalent amount of nuclear material; the remarkable part it plays in fertilization and the reproduction of the new individual—these matters need here but be referred to. They indicate, with a force that cannot be gainsaid, the controlling part played by the nucleus in the processes of cell and individual reproduction. The evidence that the nucleus is active in matters of cell metabolism is not so familiarly known, and deserves mention in a little more detail.

THE NUCLEUS AND METABOLISM.

Respiration, motility, the formation of contractile vacuoles, the seizing and destruction of minute living organisms, have all been seen to take place in cells deprived of their nucleus; even so elaborate a process as the formation of starch in the vegetable cell can occur under the same conditions. As already noted, enucleated cells having these properties cannot grow, but eventually become exhausted and disintegrate. The nucleus is necessary for the continued anabolism and katabolism of cytoplasmic matter, and it may be suggested that these higher activities manifested by enucleated cytoplasm are dependent upon the presence within it of matter previously elaborated by this nucleus from the cytoplasm and subsequently discharged.

METABOLISM AND SEMIPERMEABLE MEMBRANES

45

On the ground that in the course of his long and exact studies upon the cell he has never determined the passage of nucleolar or of nuclear matter out into the cytoplasm, Martin Heidenhain strenuously denies that the nucleus plays any part in the secretory activities of the cell. He considers the nucleus as the conservative agent in the cell, unconcerned with specific cell activities, but carrying on in the specific cell properties from one generation to another. This, I imagine, is the current conception, but it cannot be accepted. That it is the conservative agent I wholly agree; the part played by it in cell division (p. 115) and in fertilization (p. 144) would seem amply to support this view: and, as Professor A. B. Macallum¹ has suggestively brought forward, the chemistry of the nucleus and its structure indicate that the nuclear membrane is a mechanism whereby certain bodies only are permitted to pass into the nucleus, whereas others are prevented from entering, thereby acting as a mechanism to prevent gross change. Thus, although abundant chlorides and phosphates and potassium, sodium, and calcium salts may be present in the cytoplasm, his extensive studies show that the nucleus, whether vegetable or animal, is absolutely free from phosphates and chlorides; nor has he been able to detect either potassium or calcium. So also, as in the liver cell of pernicious anemia, the cytoplasm may be charged with *inorganic* iron, but the nucleus contains not a trace. On the other hand, both Browicz² and Sutherland Simpson and Herring³ have demonstrated the passage of hemoglobin (containing organic iron) into the nuclei of hepatic cells and the presence of the same there in a crystalline form. These are very significant facts, and can only mean that the envelope of the nucleus is what is known as a "semipermeable membrane," its structure being such that mechanically it permits the passage through it of certain bodies, whereas it is impervious There is evidence that the surface layer of the cytoplasm to others. constitutes similarly another semipermeable membrane having different properties. For this has been abundantly demonstrated from the long and complicated series of data which have accumulated bearing upon the subject of osmosis, namely, that the properties of these membranes vary greatly. It is not a matter of size of molecules that determines passage or arrest, nor the crystalloid or colloid nature of the substances in solution on one or other side, nor, again, their nature as electrolytes or non-electrolytes. A caoutchouc membrane, for example, will keep back the simple and highly soluble sodium chloride, and if ether be dissolved in methyl alcohol will permit the passage of the former but not of the latter; similarly, if the more colloidal copper oleate and the crystalloid cane sugar be dissolved in pyridine, the former will pass through such a rubber membrane; the latter will not. A pig's bladder membrane will permit the passage of a water-ether solution, but not of benzene equally dissolved in the mixture (Nernst); while Barlow⁴ has

¹ Proc. and Trans. Roy. Soc. Can., 3d ser. 2: 1908; sec. 4: 145.

² Bull. Internat. de l'Acad. des Sciences de Cracovie, July, 1899.

³ Proc. Roy. Soc., B., 78:1906:455.

⁴ Philosoph. Mag., 6: series 8: 1904: 1.

THE PHYSIOLOGY OF THE CELL

shown that a cupric ferrocyanide membrane is more readily permeable under pressure to the larger molecules of alcohol than it is to the smaller molecules of water. This is not the place to discuss the various theories of osmosis; suffice it say that it is not the sieve-like nature of a membrane and the size of the pores, nor, again, the surface tension of a solution (although this is a partial factor), that is the determining agent in osmosis, but, as Kahlenberg¹ has shown, the membrane permits the passage of those bodies for which it is a solvent, for which, therefore, it has an affinity of a particular order. The indications are that the outer layer of the cytoplasm-certainly in red corpuscles, and probably to a large extent in cells proper-is of a lipoid nature, formed of or containing bodies such as "lecithin," cholesterin, etc.; whereas the nuclear membrane (as shown by its staining properties) is of nucleoproteid nature. We are ready to admit with Macallum² that to the extent that this does not admit the passage of inorganic salts, and it may well be various other bodies, like glycogen, to that extent it exerts a conservative action; but the evidence, so far as it goes, suggests that proteids are somewhat widely solvents for other proteins; thus the passage of protein molecules through the nuclear membrane may be extensive, and as a conseouence there may be a wide range of variation in the eventual constitution of the nuclear material. While resembling each other in the main, distinct chemical differences have been determined between the nuclear constituents of the different tissues of the animal-between those isolated from hepatic, pancreatic, and renal nuclei; from cells, that is, having a common origin in the fertilized ovum. This in itself shows that nuclear matter is modified by its environment. As suggested on p. 31, it may well be that matter thus taken up but not vet differentiated or structurally worked up into the nuclear chromatin is to be recognized in the form of nucleoli.

If there are these indications that bodies of certain orders penetrate the nuclear membrane, and are built into the nuclear matter, we have, on the other hand, a steadily increasing amount of evidence that matter passes out through the same into the cytoplasm, and, indeed, that this matter is of singularly high importance in connection with the specific activities of various orders of cells.

There is, in the first place, what may be termed indirect evidence of such passage. Verworn noted that non-nucleated pieces of foraminifera did not show the slightest capacity to secrete the calcareous salts which form so characteristic a framework in these unicellular organisms. In another lowly form, *Thalassicola pelagica* (and the same is true in the amecba), while non-nucleated fragments can seize and kill living organisms, they cannot completely digest them, from which it would appear that the elaboration of the digestive fluids, the existence of which in unicellular forms has been demonstrated by Miss Greenwood,³

¹ Jour. Phys. Chem., 10: 1908: 41.

² Macallum is wrong in stating that fat is not found in normal nuclei.

³ Jour. of Physiol., 7:1886:254; 8:1887:263; 11:1890:576.

THE NUCLEUS AND METABOLISM

Le Dantec,¹ and others, is determined by the nucleus. Hofer² has shown that the excretion of slime by the amecha does not occur when the nucleus is absent; Korschelt,³ that the formation of chitin by insects is associated with change in the characters of the nucleus; Haberlandt⁴ has demonstrated that in the formation of the cell membrane in plants the nucleus becomes eccentric, passing to the immediate neighborhood of the site of deposit of the cellulose; a similar localization is seen in cells developing root hairs; Schniewind-Thies⁶ has called attention to the shrinkage and loss of staining power of the nucleus of nectar cells in flowers in the course of secretion; Lily Huie,⁶ to the marked changes that occur in the nuclei of the secretory cells of the leaves of the wellknown insectivorous plant, the Drosera, when fed with egg albumin.

These, it will be seen, are observations of many years' standing, which could be abundantly extended from more recent botanical and zoological literature. Coming nearer home, there are parallel observations by physiologists and histologists. Thus, I would recall R. Heidenhain's' observations, made long years ago, upon the differences in the appearances of the nuclei of salivary gland cells at rest and after stimulation, confirmed by Greenough's in his observations out the eells of the submaxillary gland, and to the interesting observations of Hodge'' (confirmed by Gustav Mann,'' Lugaro,'' and others) upon the nuclear alterations in the motor ganglion cells of bees, birds, cats, and other vertebrates, brought about by naturally and experimentally produced fatigue.

But now, in addition to these, are abundant observations upon the direct passage of material from the nucleus into the cytoplasm, and that in very immediate and significant relationship to the development of what we now recognize as the specific cytoplasmic constituents of particular orders of cells. All the observations have this in common, that they record the appearance of minute globules within the nucleus which, like the nucleoli, take on a differential stain with nuclear dyes, and these tend to accumulate immediately beneath the nuclear membrane. Following upon this, either (1) these become smaller and diminish in number, while coincidently the perinuclear cytoplasm takes on a diffuse stain of the same character, or (2) the actual passage of these globules out

¹ Ann. de l'Inst. Pasteur, 4: 1890; 273, and 5: 1891: 163.

² Jenaisch, Zeitschr, f. Wissensch., N. F., 17:1890;105.

³ Zool. Jahrbuch, Abth. f. Anat., 4; 1899; 1, and Naturwiss. Rundschau, 1887; 409.

⁴ Die Beziehungen zwischen Funktion u. Loge des Zellkernes bei den Pilanzen, Jena, 1887; also Sitzungsber, d. Kaiserl. Akad. d. Wiss., Vienna, Math. Naturwissenchs. Kl., 18; 1889; Abth. 1: 190.

⁵ Beitr. z. Kentniss der Septalnectarien, Jena, 1897.

⁶ Quart. Jour. Mier. Sci., N. S., 39: 1897: 387.

⁷ Hermann's Handb. der Physiol., 5: 1883.

⁸ Jour. of Med. Research, 7: 1902; 360.

¹⁰ Jour. of Anat., and Physiol. 29: 1894: 100.

¹¹ Lo Sperimentale, 44: 1895; Soc. Biol., 2.

⁹ Jour. of Morphol., 7: 1892: 95.

THE PHYSIOLOGY OF THE CELL

of the nucleus into the cytoplasm is described. With improved technique observations upon this second order of events are becoming more and more common, and we have different orders of workers giving to these nucleolar derivatives various names—plasmosomes, chromidia, mitochondria, etc.¹ Organized chromatin as such is not discharged from the nucleus, nor is it to be found free in the cytoplasm, save in the rare cases of aberrant mitosis, as in cancer cells, when sundry chromosomes may fail to be gathered into the daughter nuclei. What is dis-



A, resting nerve cell with large rounded nucleus, showing chromatin network, the Nisel bodies in the cytoplasm (derived from the nuclear material) also large and prominent; R, exhausted nerve cell of same order, with shrunken irregular nucleus, chromatin network indistinct, Nisel bodies diminished in size and pordy staining. (After Gustay Mann.) Charged is nucleolar matter, and this either by a process of diffusion or as discrete globules or granules. Once free in the cytoplasm, it is acted upon and interacts with the cytoplasm and its contents, and, according to the nature of the cell and of its cytoplasm, so does it take an active part in the formation of one or other specific cell substance. Modification and disappearance or conversion of the plasmosomes is seen to take place *pari passu* with the accumulation within the cell of these specific substances.

The process, in the first place, has been made out in the vegetable cell. Here may be recalled 'Torrey's² observations upon the secretion of diastase in maize seeds, etc. At the beginning of germination of the

seeds the nuclei of the columnar di^astase-producing cells contain darkstaining granules, with but a few in the cytoplasm. Small breaks are to be made out in the membrane of the heavily loaded nucleus, and through them the granules exude in small streams. These, at first spread through the cell, become later collected at the end next to the endosperm. Here they become ultimately dissolved. It is following upon their dissolution that the first action of a ferment upon the cell wall and matrix of the endosperm is observable.

It is described also throughout the animal kingdom from the protozoa upward. R. Heriwig⁸ and his pupils afford numerous instances of the development of such chromidia in the protozoa, in the form either of fine granules or of fine filaments or filamentous network. He has described their conversion into and relationship to the development of

⁴ As per example Archoplastic loops, centrophormia, trophopspongia, ergastoplasm!

³ Arch. f. Protistenkunde, 3; 1992; Sitzungsber, d. Gesellsch, f. Morphol, u. Physiol., 18; 1992; 77; Festschr, f. Ernst Hacekel, 1994; 301. See also Gerassimow, Beihefte z. Botan. Centralbl., 18; 1904; and Goldschmidt, Biol. Centralbl., 24; 1904; Arch. f. Protistenkunde, 5; 1904; and Zool. Jahrb., Anat. Abt., 21; 1905.

² Memoirs of the Torrey Botan, Club, New York, 29: 1902: 421.

NUCLEOLI AND METABOLISM

pigment granules of melanin-like nature in the cytoplasm in certain abnormal states of the individual *Actinospherium*. A. B. Macallum,¹ in 1890, studying the nuclei of developing ova in the ovaries of the lake lizard (Necturus) and of the frog, gave a very clear description of the formation and discharge of those nucleoli and their conversion into the yolk granules. Montgomery,² in *Piscicola* (a leech parasitic



Developing eggs of Astelan bifda, showing extrusion of nuclear matter. Young oncyte, the nuclear chromatin in the form of scattered branching threads. The deeply stained nucleolus is seen in the act of extruding spherules into the cytoplann, $\times 2000$,



Nucleolus from an adult egg, showing discharge of nucleolar spherules from the unstained area of the nucleolus into the nuclear substance. \times 1000,



A relatively young oncyte, showing discharge of spherales from the unstained area, with accumulation of spherales previously discharged to form first stage of the "yolk nucleus" -m; other spherales scattered through the eytoplasm, \times 1000. (Chubb.)

upon fresh water fish) found the discharge most active and abundant. We append certain figures from a very full study by Chubb³ of the ovarian ova of *Antedon*, a crinoid starfish, or, more accurately, brittle fish.

¹ Proc. Can. Instit., Toronto, 1:1896:11, and Trans. Can. Instit., Toronto, 1: 1891:247.

² Jour, of Morphol., 15; 1898. ² Phil. Trans. Roy. Soc. Biol., 98; 1906; 447.

Fig. 14 shows, in addition, how the plasmosomes may become swollen, fused, and modified into such intracytoplasmic accumulations as the "yolk nucleus" of the ovum, the "archoplasm" of spermatocytes, and other cells.

It is in gland cells, with their characteristic granules, that the stages of the process have been repeatedly observed. In mucous goblet cells, F. Hermann has noted that during active secretion minute globules taking the same stain as nucleoli appear immediately around the nucleus, these being absent in the resting stage.¹ Maximow gives a fuller account of like appearances in the serous salivary gland cells of the dog. The youngest, smallest, and most deeply staining granules are situated in the immediate neighborhood of the nucleus, and are indistinguishable from the intranuclear plasmosomes (nucleol). As these pass to a further distance from it their staining power diminishes



Relationship of nuclear plasmosomes to zymogen granules and secretory substances ysecreting cell: a, intranuclear plasmosome; b, granule (extranuclear plasmosome) in cytoplasm, near nucleus having same staining renation, and evidently discharged from the nucleus; c, conversion of same into more feebly staining secretory (prezymogen) granules; d, further stage; zymogen granules about to be discharged. (After Maximow). and they appear to give place to definite secretory granules. Steinhaus and Macallum give a very similar description of the mode of production of the granules in the pancreatic cell. Bensley,² Carlier,⁴ Mathews, all record the *prezymogens*, or granules becoming converted into the *zymogens* of the secreting cell, as of nuclear origin.

Mathews,⁴ of Chicago, studying the development of the well-known secretory (zymogen) granules of the pancreatic cell, noted first the appearance of a "Nebenkern" (or paranuclear body, similar to the

yolk nucleus figured in Fig. 14), within and in direct association with which were to be noted spiral twists or coils, which eventually became distributed through the cytoplasm. These coils proceed to decompose and a stage intervenes between their disappearance and the appearance of the zymogen granules in the cell. It is interesting to note that C. E. Walker,⁵ has described an almost identical mode of development of the granules in the leukocytes of the bone marrow of the rat and guinea-pig. The threads, it will be seen (Fig. 16), are characteristically beaded, and the beads become the eventual granules.

So, also, Macallum has shown that the hemoglobin of the red corpuscles of amphibia is derived from the nucleus, the hemoglobin, or its antecedent, diffusing through the nuclear membrane and becoming

¹See also Reichenow, Arch. f. Mikr. Anat., 72: 1908; 671.

² Proc. Can. Instit., Toronto, 1: 1896; 1, and Quart. Jour. Micros. Sci., N. S., 41: 1898; 301.

³ Brit. Med. Jour., 1900; ii; 740, and La Cellule, 16: 405.

⁴ Jour. of Morphol., 15: 1899, Supplement.

⁵ Proc. Roy. Soc., B., 79: 1906: 491.

THE NUCLEUS AND METABOLISM

en.

the

nd

he

я

bf ac

8

fixed in the cytoplasm. Arrigoni,¹ by a process of intravital staining of the nucleated red corpuscles seen in the blood of a case of pernicious anemia, has observed the presence of buds upon the nuclear membrane similar to those figured by Schmaus and Albrecht in karyolysis (see Fig. 18), the development of metachromatic granules in the cytoplasm, and their solution in the cell substance as hemoglobin. The nucleus, in short, is not, as used to be thought, cast out of the cythrocytes, but, as Pappenheim and others have taught, undergoes solution in the cell (intracellular karyolysis), giving rise to the hemoglobin



Cells (leukocytes) from bone marrow of guinea-pig showing: (a) development of thread in the archoplasm or chromidial mass in the cytoplasm apposed to the nucleus; (b) extension of thread through the cytoplasm, and commencing segmentation into cytoplasmic granules; (c) granules scattered through cell substance, their chain-like arrangement still noticeable. (Alter Walker)

Allied possibly to these last instances is the evidence that the nucleus of fat cells plays a part in the accumulation of fat. Study a section made from any collection of fat cells, and it is seen that the nuclei of those cells are characterized by the possession of isolated, sharply cut vacuoles. Shattock² has demonstrated that these take on the specific stain with Sudan III; in other words, that they contain fat. To our knowledge, these are the only nuclei in the body that are consistently vacuolated. The relationship is at least suggestive.

Lastly, Scott³ and others have clearly proved that the characteristic Nissl bodies so abundant in the body of the neurons or ganglion cells are nuclear products.

I give this evidence in such circumstantial detail because, in the first place, it is of the highest importance as throwing light upon the nature of specific cell activities and the part played by the nucleus in the same; in the second place, because I have not found it collected together in any work of reference known to me, certainly in no text-book available to the ordinary student. Data so abundant and harmonious, recorded by so large a number of independent observers in so many branches of biological science, can only point to the working of a law of wide application. Instead of the nucleus being an inert conservative body, it is seen to be the source of the specific activities of the cell, or, perhaps more correctly, of the substances which are necessary for the performance of those specific activities.

¹ Folia hæmatol., 6:1908:444. ² Trans. Path. Soc. London, 54:1903: 216.

³ Trans. Can. Instit., Toronto, 6:1899:405. See also Hatai, Jour. Comp. Neurol., 9: 1901.

THE PHYSIOLOGY OF THE CELL

It is deserving of note that both these processes, of what we may term *chromatolysis*, or diffusion outward and disappearance of the modified chromatins from the nucleus, and of *chromidiation*, or discharge of



Section from the liver of a child that died from acute sepis, to show various stages of knyolysis of the first order: 1. Unaffected nucleus; 2 and 3, paler staining nuclei, with some swelling and diminution of chromation; 4, nuclei still more swollen, the membrane only and an occasional nucleodar mass taking on the stain; 5, nuclei present as little more than unstrained vesicles. nucleolar globules from the cell, occur as, or as the result of, pathological conditions. The disappearance of the nucleus is a sign of the gravest lesions in the cell. It would seem that we must recognize three modes: (1) Karyolysis of the first order, in which the nucleus swells up, becomes vesicular or ædematous, and takes on the nuclear stains more and more feebly until eventually it becames invisible, indistinguishable from the surrounding cytoplasm. This is not infrequently encountered in areas of inflammation and in the condition of acute cloudy swelling (see Section III, Chapter XXV). It has still to be determined whether in all cases the modified chromatin has diffused out of the nucleus or whether (as would seem to be the

case in the temporary invisibility of chromatin in certain stages of cell division) the reaction and constitution of the chromatin still present has become altered, so that it no longer has an affinity for the basic nuclear dyes. (2) *Karyolysis of the second order*, or patho-.



Hyperchromatosis of nuclear wall: nucleus vesicular, with disappearance of chromatin network and accumulation of chromatin masses upon the nuclear membrane. (Schmaus and Albrecht.)



Karyolysis: extreme reduction or solution of nuclear chromatin, from a renal infarct. In b the nucleus is wholly disclosed save for some fine chromatin granules. (Lubarsch.)

logical chromidiation. In the liver cells in phosphorus poisoning Stolnikow has described the extensive passage out of minute bodies from the nucleus into the cell body, at first taking the nuclear stain, then later losing the power of staining, so that the cell body becomes filled with shell-like, scarce staining globules and smaller particles, while the

NUCLEAR DEGENERATION

nucleus becomes faded and scarce visible. Lukjanow, Galeotti, and Vigier have confirmed in connection with other tissues. It is common in many acute conditions to find a disappearance or great diminution of the nuclear network with coincident accumulation of nodules of modified chromatin immediately within the nuclear membrane, the condition of *hyperchromatosis*. This is apparently a preliminary stage. (3) There is, thirdly, an acute process of *karyorrhexis*, which is of a different order, the nuclear membrane breaking down and the nucleus disintegrating with discharge of masses of chromatin into the cell substance. Certain authorities make a distinction between this condition and *pyknosis*, in which the nucleus comes to be represented by shrunken but deeply staining irregular masses of chromatin collected together in the centre of the cell. We regard cells showing these conditions as dead.



Discharge of chromatin granules (plasmosomes) from the nuclear wall into the cytoplasm. (Schmaus and Albrecht.)

ified

e of

ell.

ho-

Par-

It

og-

1818

111-

he

les he

Leukocytes with disintegrating masses of nuclear material scattered through the cytoplasm (karyorrhexis).

Conclusions .- If, then, on the one hand, we regard the nucleus as the dominating portion of the cell, and, on the other, admit that this cannot act save in association with the cytoplasm, what must be our conception of the relationship of these two components of the cell and of the nature in general of cell activities? This question can, we think, best be answered after discussing the general principles of the chemistry of the cell. In the meantime the conclusion to be reached is that in the cell we have indications of the existence of living matter of two orders. There is in the nucleus matter which initiates growth, reproduction, and what we must regard as the very highest vital activities; matter which, moreover, can only react upon the cytoplasm, taking up substances from, and yielding other substances to this, and cannot react upon the external medium; in the cytoplasm, on the other hand, there is matter capable of taking up and acting upon other matter from without, from the external medium, but this is of a secondary order. It can manifest what may be termed the lower vital activities-absorption, respiration, mobility, and contractility, and these independently of nuclear control; it cannot initiate the higher activities of growth and reproduction.

Lastly, we may mention here, but will not discuss, a third order of matter that plays a most important part in cell activities; we refer to the organic ferments, substances produced by cell metabolism, capable of discharge from the cell and acting as a second group of intermediate bodies, this time between the external medium and the cytoplasm.

CHAPTER IV.

THE CHEMISTRY OF THE CELL.

THE PROTEINS.

IF we make a broad survey of all forms of life, animal and vegetable, we find that there is one order of substances common to and to be extracted from all (dead) cells, however simple or however highly differentiated, namely, proteids, or, as it is becoming now the custom to designate the wider group of related substances, proteins. Apart from these, with the exception of water and the phosphorus and iron which appear to be intimately associated with the proteins, we can recall no other constituent common to all cells. A very great variety of other components can be isolated from cell substance-salts of one or other order, fats, alcohols (ethyl alcohol,¹ cholesterin), and lipoids (the lecithins, so-called, cerebrin, etc.), carbohydrates (starch, glycogen, etc.), chlorophyll, and other complex bodies which we regard either as the results of disintegration of proteid matter or as stages leading up to the formation of the same. And some of these in certain cells may be accumulated in such abundance as to be the main constituents. But each of them may be wanting in one or other form of cell. The proteins alone-and water-are common to all cells.

It is true that the very analysis of living matter, whereby we isolate these proteins, renders that matter dead; that when isolated these proteins are inert substances, manifesting few of the phenomena which we recognize as proper to living matter. We must, nevertheless, conclude that "life" is bound up with the presence of proteins, even if at the same time we are compelled to admit that the chemical or physical constitution of living matter is something different from that of the inert substances we gam in the laboratory. To be more exact, life is bound up with the presence of *proteidogenous* matter, and if later, in order to prevent any possible confusion between the living substance and the dead proteins, we speak of *biophoric* molecules,² it must const ntly be kept in mind that we regard these as formed, in the main, if not essentially, of matter which by rearrangement or by satisfaction of its affinities, becomes converted into compound proteins.

¹ This has been distilled from normal brain and other tissues in recognizable amounts.

² Vide foot-note, p. 75.

PLATE I







FIG. 2

Oxyhemoglobin of Carp.



Oxyhemoglobin of Necturus.



Oxyhemoglobin of Squirrel,



Oxyhemoglobin of Guinea-pig



Oxyhemoglobin of Dog.

Protein Crystals.



THE CONSTITUTION OF PROTEINS

THE CONSTITUTION OF PROTEINS.

What, then, are these proteins?

They are singularly complex compounds of nitrogen, carbon, oxygen, hydrogen, and sulphur; the highest forms contain also iron and phosphorus.1 Their molecules are of such size and complexity that in general they are incapable of undergoing crystallization, so that they remain in a colloid state. This colloidal character renders it impossible for us to be sure that we are dealing with pure substances, and so makes analysis in most cases at the best approximate. But some of the simpler proteids are crystallizable; these can be obtained pure, and can be analyzed. Of such, the most familiar is *hemoglobin*, or, more accurately, are the hemoglobins, for the analysis of hemoglobin from different species of animals demonstrates that the composition is not identical; indeed, the fact that the crystals of this substance from different animals have widely different shapes is sufficient to indicate variation in composition. This difference is well shown in Plate I, which we owe to Professor Reicherb. Modern analyses vary between C₆₅₃H₁₁₈₁N₂₀₇O₂₁₀ FeS₂ and C₇₁₂H₁₁₃₀N₂₁₄O₂₄₅FeS₂. Whatever formula we take, we clearly have to deal with a molecule of enormous size, and this, to repeat, although we are dealing with one of the less complex proteins. It is, indeed, estimated that the average molecular weight of a protein is in the neighborhood of 15,000; there can be little wonder that the large proteid molecules are unable to make their way through the fine pores of an animal or vegetable membrane-that they do not diffuse. Even the simpler proteins, like egg albumin,2 have a molecular weight in the neighborhood of 5000.

Classification.—These proteins are of various orders. Material aid in their classification has been afforded recently by the labors of two committees of physiological chemists, the one appointed by the Physiological Society of Great Britain,³ the other appointed a little later by the American Society of Biological Chemists and American Physiological Society.⁴ Here the final report of the latter will in the main be followed as being somewhat the more precise, although the conclusions of the two are singularly harmonious. We shall have so frequently to refer to the different orders and classes of proteins that it will make for clearness if we give the classification in some detail.

We can divide them into two main groups—the simple and the combined or conjugated—(in a sense, as we shall show, all proteins are combined, wherefore conjugated, as the more uncommon, is the better word). A third group is to be found comprising the products of hydro-

¹ For a full and clear presentation of the chemistry of the proteins, see Gustav Mann, *Chemistry of the Proteils*, London and New York, 1906, or the *Practical Physiological Chemistry of Hawk*, 24 edition, Philadelphia, 1909.

² The formula of egg albumin is regarded as approximately C₂₂₀H₂₃₆N₄₈O₇₈S₂.

^a See Jour. of Physiol., 34: 1907: 17.

⁴See Proc. Amer. Soc. Biol. Chemists, 1 ; 1908 ; 142.

lytic dissociation of the protein molecules still affording proteid reactions, and, conversely, the products of synthetic building up of amino-acids into bodies of proteid affinities.

I. **The Simple Proteins.**—Among these are included those which yield only *a*-amino-acids or their derivatives on hydrolytic cleavage. Many of these are met with apparently in a free state in the fluids and cells of the body and are what we regard as type proteins—albumins, globulins, and the like. Although no means are available whereby the chemical individuality of any protein can be established, those isolated from animal and vegetable tissues are so well characterized by constancy of ultimate composition that, until we possess further knowledge, they may be treated as chemical individuals. These are:

1. **Albumins**.—Simple proteins soluble in pure water and coagulable by heat (*serum albumin*, *egg albumin*, *lactalbumin*, etc.).

2. Globulins.—Insoluble in pure water, but soluble in neutral solutions of strong bases with strong acids (*serum globulin*, *edestin*, *myosin of muscle*, *ovoglobulins*, *fibrinogen*, etc.).

3. **Glutelins**.—Insoluble in all neutral solvents, but readily soluble in very dilute acids and alkalies (simple proteins of this order—*glutelin*, etc. —occur in abundance in the seeds of cereals).

4. Gliadins (Br.), Prolamins (Osborne), or Alcohol-soluble Proteins (Am.).—Soluble in relatively strong alcohol (70 to 80 per cent.), but insoluble in water, absolute alcohol, and other neutral solvents (gliadin of wheat, *scin* of maize, *hordein* from barley, etc.).

5. Albuminoids (Am.) or Scleroproteins (Br.).—Simple proteins having essentially the same chemical structure as the preceding, but insoluble in all neutral solvents. Such compose the main organic constituents of the external coats of animals and of the skeletal structures and connective tissues (collagen, keratin, elastin, fibroin. Gelatin is not here included, being a derived protein, from collagen).⁴

6. **Histones.**—Soluble in water and insoluble in very dilute ammonia, insoluble also in the excess of ammonia if ammonium salts be absent; are basic proteins which yield precipitates with solutions of other proteins, and a coagulum on heating which is easily soluble in very dilute acid. On hydrolysis they yield a large number of amino-acids, the basic predominating (*qlobin*, *scombrone*, etc.).

7. **Protamins**.—The simplest proteins hitherto found in nature, approximating in simplicity of constitution to the synthetically derived polypeptids. Soluble in water, uncoagulable by heat, precipitate aqueous solutions of other proteins; possess strong basic properties, form stable salts with strong mineral acids. They yield comparatively few amino-acids, among which the basic amino-acids greatly predominate (*sturine, clupeine, sulmine*, etc., from fish sperm).

II. The Conjugated Proteins.—In these the protein moiety is united to some molecule or molecules of another order otherwise than as a salt.

¹ Although Emmett and Gies claim that gelatin is not a product of hydrolysis, but the outcome of intramolecular arrangement.

PEPTONES, AMINO-ACIDS, AND POLYPEPTIDS

1. Nucleoproteins.—Compounds of one or more protein molecules with nucleic acid (see p. 63).

2. Glycoproteins.—Compounds of the protein molecule or molecules with a substance or substances containing a carbohydrate group other than nucleic acid (mucins and mucoids, chondromucoid, amyloid, chondro-albuminoid, etc.).

3. **Phosphoproteins** (*Nucleo-albumins*).—Compounds of the protein molecule or molecules with an as yet undefined phosphorus-containing substance other than nucleic acid or lecithin (*cascin, vitellin, etc.*). The British Committee include these among the simple proteins on the ground that the cleavage products still contain phosphorus.

4. **Chromoproteins** (Br.), or **Hemoglobins** (Am.).—Compounds of the protein molecule with *hematin* or some similar iron-containing substance, or, in certain classes of invertebrates, with a substance containing copper, manganese, or zinc (*hemoglobin*, *hemocyanin*).

5. Lecithoproteins.—Compounds of the protein molecule with "lecithins" (*phosphatides*, *lecithans*). The British Committee objects that the existence of these as true compounds has not yet been determined, and it should be added that the existence of lecithins as actual chemical individuals is still a matter of controversy (see p. 96).

6. Lipoproteins.—The existence of compounds of the protein molecule with fatty acids is indicated by the synthetic formation of compounds between the amino-acids and the fatty acids.¹ On pathological grounds the existence of these compounds must be postulated, although as yet they have not been isolated.

III. **Derived Proteins.**—These are divided into two classes according to the extent of alteration of the protein molecule.

Primary Protein Derivatives.—Bodies exhibiting only slight alteration of the protein molecule—*proteans*, *metaproteins* (including "acid" and "alkali proteins," acid albumin, etc.), *computed proteins*.

Secondary Protein Derivatives.—Products of further hydrolysis, and, conversely, of the building up of amino-acids into molecules of protein type (*proteoses, peptones, polypeptids*).

PEPTONES, AMINO-ACIDS, AND POLYPEPTIDS.

To the opening years of this century belongs the solution of the problem as to the nature of the proteins. Valuable preparatory work had been accomplished in the preceding thirty years. Schaal (1871), Curtius (from 1881 onward), Grimeaux (1882), Schutzenberger, of Paris (1888-91), and Lilienfeld (1894) had each prepared synthetically bodies giving proteid reactions, and in 1892 Hofmeister had postulated, as we now know correctly, the mode of linkage of amino-acids in the

⁴ Thus Bondi has combined glycocoll and alanin with both laurie and palmitic acids, Biochem. Zeitschr., 17 (1909) 543 and 553. These lipoproteins are not included in either the American or British classifications.

THE CHEMISTRY OF THE CELL

protein molecule. It is the later brilliant researches of the great Berlin chemist, Emil Fischer, his isolation and preparation of a long series of amino-acids in addition to those already known, and his establishment of the general principles by which these may be linked together in series, that have at last placed upon a sure basis not merely the synthesis of bodies of proteid nature, but what is of equal importance, the constitution of the same. We can but indicate broadly the direction taken by these researches and their singular significance.

It is to be noted, in the first place, that if individual proteins be analyzed, whether these be only obtainable in the colloid state or be, like hemoglobin, crystallizable, successive analyses, while approximating, do not give identical proportions of C.H.N. and O. In other words, while these clearly represent distinct compounds having well-defined properties, the constitution of each of them is not absolutely fixed. An interesting and profoundly suggestive feature in all of these forms is that they may be broken down with relative ease into simpler bodies which still possess proteid characteristics. This we have already indicated in our description of the combined proteins. It is equally true in regard to the so-called free proteins. We need but recall the fact that proteolytic enzymes break down albumin, globulin, myosin, etc., into peptones and albumoses. In other words, hydrolysis, whether produced by the action of these enzymes or by boiling with dilute acids, or by the action of alkalies, splits up more elaborate proteins in the first place into bodies which are still proteins and give the characteristic reactions of the same, e. g., the biuret reaction, but which are evidently in the form of smaller molecules. That this is so is indicated by the fact that they diffuse through membranes. In this process of hydrolysis the proteins take up into their molecule one or more molecules of water, and the resulting *peptones* may be spoken of as degradation products. Thus the ordinary protein molecule is evidently a compound of like molecules, and is an example of polymerization, or the formation of a molecule of large size by the junction of a series of smaller similar molecules.

But now, by a continuation of this process, the peptones and albumoses afford still simpler degradation products, foremost among which are to be found members of the large group of *amino-acids*. About three-quarters of the albumin molecule is composed of such aminoacids. Thus, to give an example, the simplest bodies of proteid character thus far discovered in nature are the protamines (sturin, clupein, salmin, scombrin, etc.) obtained from the sperm of the sturgeon, herring, salmon, and mackerel. Compared with hemoglobin, the formulas of this class are relatively very simple, that of salmin, according to Kossel, being $C_{30}H_{s7}N_{17}O_{e}$. They all give the biuret test, and by hydrolysis give first bodies of the nature of peptones (protones), and by further action break up into still simpler nitrogenous bodies. Thus, by hydrolysis, sturin, $C_{30}H_{a0}N_{10}O_{2} + 5H_{2}O$, affords:

 $\begin{array}{rcl} C_6H_9N_3O_2 &+& 3C_6H_{14}N_4O_2 &+& 2C_6H_{14}N_2O_2 \\ Histidin & Arginin & Lysin \end{array}$

PEPTONES, AMINO-ACIDS, AND POLYPEPTIDS

Kossel, indeed, has determined that all proteids yield these nitrogencontaining amino-bodies: histidin, arginin, and lysin, called by him the hexone bases, since all contain six carbon atoms.⁴

The Amino-acids.—These amino-acids are intimately related to the fatty acid series; they are, indeed, fatty acids-aminated fatty acids, *i. e.*, fatty acids given partial basic properties by the addition of NH. molecules. Thus, glycocoll is amino-acetic acid, acetic acid being C.H.O., and glycocoll having the formula C.H.NO. Similarly, alanin is aminopropionic acid, C₃H₂NO₂, tyrosin is p.-oxyphenyl-a-amino-propionic acid, CaH11NO2, tryptophane is indol-amino-propionic acid, C11H12N2O2. By processes of hydration they are converted into the hydroxyl acids of that series. It is these amino-acids to which the chemists have especially directed their attention-Curtius, in the first place, and of late, more particularly, Emil Fischer and his pupils. Their constant presence as degradation products of proteins and their relative great abundance indicated not only that they are to be regarded as primary nuclei of the protein molecule, and that the proteid molecule is essentially built up by a linking together of amino-acid molecules, but also that, experimentally, by bringing about such a linkage, it might be possible to build up-synthesize-more complex molecules of the proteid type; or, in other words, to accomplish that most ambitious object of the chemist, the experimental production of proteins.

The Polypeptids.—As already noted, the first steps have been achieved toward this end.

Fischer and his pupils have devoted many years to an active study of the amino-acids. Some eight of these had already been produced synthetically (glycocoll, alanin, amidovalerianic acid, leucin, asparagin (aminosuccinic acid), aspartic acid (amino-succinamide), phenylalanin, glutamic acid). Others of the long series they themselves synthesized, so that now some thirty members of the group are known. Fischer devised a method of gaining the mono-amino-acids in a pure state by converting them into their esters, in which form they are volatile and can be distilled fractionally in vacuo. From these pure esters he gained the pure amino-acids and studied their compounds. The diamino acids (ornithin, lysin, etc.) he similarly purified (by the phosphomolybdic acid method). From Biot (1815) and Pasteur (1860) onward the optical activity of compounds which are products of vital activity had been taken as one of the particular manifestations of vital activity, as an evidence that what Moore would now term "biotic energy," or vitalism, is distinct from ordinary chemical processes, the corresponding products when gained by the chemist being optically inactive. Now, Fischer shows that he can gain both optically inactive and optically active forms of these amino-acids-that chemical methods outside the body can reproduce the products formed within, and that one of the last physical distinctions, if not the last, between "vital" and "laboratory" products has been swept away.²

¹ Deutsch Med. Woch. 24: 1898: 581.

² It deserves note that prior to Fischer, Alex. Mackenzie, of London, had achieved the production of optically active asymmetric carbon compounds (not, however, amino-acids) by direct synthesis.

THE CHEMISTRY OF THE CELL

A characteristic feature of these amino-acids is that they are amphoteric; they possess both acid and basic properties, being acid to a greater or less degree through the contained COOH group or groups, slightly basic through the NH₂ group or groups. It is this property that permits linkage. Thus, to take one of the simplest of the mono-amino acids glycocoll (NH₂.CH₂.COOH), or as, for convenience, we may write it, reversing the order of the NH₂ component, HNH.CH₂.COOH—by dehydration two molecules may become linked as follows:

$\begin{array}{c|c} \underline{OH} & \underline{H} \\ \underline{HNH.CH}_{2}CO - \underline{NH.CH}_{2}COOH; \\ \underline{Glycyl} & \underline{Glycin} \end{array}$

and in this way glycyl-glycin' be produced. If this be acted upon again by a halogen-containing acid chloride, and the halogen-salt be treated with ammonia, the di-glycyl-glycin can be obtained:

 $\begin{array}{c|c|c|c|c|} \underline{OH} & \underline{H} \\ \underline{H}_{t}N.CH_{2}.CO & -HN.CH_{2}.CO & -HN.CH_{2}COOH; \\ \underline{Glyeyl} & \underline{Glyeyl} & \underline{Glyein} \end{array}$

and this synthesis and process of polymerization can be continued until first the *pentapeptid* was obtained, with five nuclei linked in series; more recently the linkage of eighteen nuclei in series has been announced.

This linkage, it may be added, has been obtained not merely between identical nuclei, but between nuclei of different amino-acids. Thus, for example, Fischer has combined leucin and glycocoll into leucyltetraglycyl-glycin and leucyl-pentaglycyl-glycin. Leuchs and Suzuki have formed glycyl-phenylalanin and leucyl-phenylalanin; Fischer and Königs, glycyl-asparagin, and, among the di-amino acids, Fischer and Suzuki have formed lysyl-lysin and histidyl-histidin (dipeptids), etc.

These polymeric amino-acid compounds thus gained have been termed by Fischer *polypeptids*. They are bodies which, in appearance, certain color reactions (such as the biuret test), behavior toward alkalies and acids, and toward enzymes, so closely resemble the true peptones that, to quote Fischer, they must be regarded as their nearest relatives. What is more, bodies of this order have been recovered from organic substances. P. A. Levene has discovered glycylproline anhydride among the products of the digestion of gelatin. Fischer and Aberhalden have isolated a tetrapeptid from silk fibroin, etc.; and, lastly, Fischer² notes that *l*-leucyltriglycyl-*l*-tyrosin, prepared artificially, has all the properties of the albumoses.

Conclusion.—Let it be clearly understood that these polypeptids are not as yet demonstrated to be identical with any known peptone, and that the peptones themselves are degradation products of the higher proteins. Nevertheless, these observations go very far toward confirming us in a conception of these peptones and the higher proteins as

⁴ It will be observed that *glycocoll*, *glycyl*, and *glycin* differ according to the presence or absence of H and OH components.

² Faraday Lecture, 1907, Jour. Chem. Soc., 91: 1907: 1749.

PEPTONES, AMINO-ACIDS, AND POLYPEPTIDS

composed of polymerized molecules—as giant molecules, formed in the main of amino-acid molecules linked together by their otherwise unsatisfied NH and CO affinities. Thus, to modify Hofmeister's illustration,⁴ we may represent a portion, at least, of the protein molecule, as follows (Fig. 22):



So as not to confuse the student, we have in our diagram indicated the nuclei as leucin, tyrosin, lysin, etc.; it will be seen on study that the leucin is minus an H on the one hand and an OH on the other; thus, to employ Fischer's terminology, it is a leucyl. The same is true of the other bodies. It will be seen that we represent the protein molecule as composed of a main chain in serial repetition with a number of side-chains of varying constitution. Further study will show that in the chain as here indicated each separate link in its simplest form may be regarded as:

$$(H) - \text{NH.CH.CO} - (OH)$$

$$(H);$$

that is, as a glycocoll molecule, which the NH affinity on the one side and the CO affinity on the other have been satisfied by linkage with a like molecule, while one H of the CH₂ is substituted by butane, methylparaoxybenzene, acetic acid, and butylamin. In other words, we have a main glycocoll chain with a series of free swinging chains capable of being replaced or modified by processes of oxidation or by the action of bacteria and enzymes.

This is, perhaps, the simplest case that we can conceive: the constant presence of lysin and histidin as degradation products in the analysis of proteins suggests that the links of the main chains may not be so simple as here indicated, and the preponderance of nucleoproteids in the cell nucleus points to the conditions there as being of a more complex nature. But the studies of many different schools converge toward this conception of the structure of the protein molecule as a linking in series or repeated series of amino-acid nuclei.

⁴ Vide B. H. Buxton, American Medicine, 6:1903;581. A clear and concise presentation of the data supporting this conception of the nature of the proteid molecule.

THE CHEMISTRY OF THE CELL

It may be added that I have represented these links as portions of a circle in order to indicate that the complete molecule is of the nature of a ring.⁴ This idea, nevertheless I hold, best fits in with what we know regarding the proteins, namely, with their fixity to this extent; that we encounter in nature proteins of characteristic types and properties, bodies which cannot be conceived as capable of progressive linkage with an unending series of nuclei. The linkage in ring form best expresses the conception of completion and individualization of the compound molecules.




Diagram of suggested main ring of a protein molecule, without attached side-chains.

It deserves note also that there are indications that even in the simplest albumins yet other connections exist within the molecule. For instance, in the tryptic digestion of albumins, while certain amino-acids are split off with relative case, a certain proportion of the proteins are found most resistant, constituting the so-called *kyrins*, or, according to Fischer and Abderhalden, bodies of polypeptoid nature, yielding, on treatment with acid, certain amino-acids not present in the products of the tryptic digestion, along with othe s that are. Indeed, the existence of intermediate bodies like the peptones and albumoses, between the complete protein molecules and the amino-acids, suggests strongly that the amino-acids are built up into groups, of which the constituents have secondary, and therefore stronger, union.

THE CHEMISTRY OF THE NUCLEUS.

The dominant position which we have already indicated as taken by the nucleus in the cell economy renders it important to determine whether there are differences in the chemical composition of the nucleus

¹ Against this view may be advanced the amphoteric character of proteins which would seem to suggest a free amidogen radical balanced by a free carboxyl group.

as compared with the cell substance in general. As a matter of fact, there are pronounced differences.

In the first place, we find certain substances stored up in the nuclei which are present to but a slight extent, if, indeed, at times they are at all recognizable, in the cell body. Of these, as shown by Lilienfeld and Monti' and by Macallum,² phosphorus is most noticeable; another constant in nuclei, not so constant in the cell body, is "masked iron," *i. e.*, iron so united that in ionization no free Fe ions are dissociated, the Fe being present as a constituent of what is probably a very complex ion³ which has to undergo further dissociation before free Fe ions are liberated. On the other hand, as already noted (p. 45), certain substances commonly present in the cell body are characteristically absent from the nuclei.

When, now, we come to study more closely the proteid contents of the nuclei, these are found to exhibit certain pronounced features. Ordinary proteins, it is needless to say, are completely digested and dissolved by the gastric juice; but if a richly cellular tissue, or if free cells, such as are present in pus, be subjected to gastric digestion, as shown by Miescher,4 the nuclei are found largely unaffected; they show little decrease in volume, and on further study, as shown by Malfatti,5 the portions which are thus unaffected are the chromatin of the nuclear network, and the nucleoli. The linin or non-staining basis of the nuclear network is also undigested. We owe, especially to Kossel's⁶ investigations, the explanation of these peculiar features of nuclear material. Briefly, the nuclei are very largely constituted of a special group of proteins, the nucleoproteins, which split up into a simple protein, usually a histon, and nuclein, and it is the nucleins which are unacted upon by gastric juice, and, further, are characterized by a high phosphorus content, the amount of phosphorus varying in the different forms between 2 and 9 per cent. They are insoluble in water, dilute mineral acids, ether, and alcohol, but are soluble in alkalies. Like the nucleoproteins, they are of a proteid nature, affording another example of this combination of protein with protein to form complex molecules. On further decomposition they yield a protein and a nucleinic (or nucleic) acid. We say a nucleinic acid, for the different figures afforded by observers analyzing nuclear material from different sources would indicate that there are several of these acids. That obtained by Mathews' from herring sperm gave the formula C40H54N14O27P4.

¹ Zeitschr, f. Physiol. Chemie, 17: 1893; 410.

² Proc. Roy. Soc., 50; 1891; 277; Quart. Jour. Micr. Sci., N. S., 38; 1895; 175.

³ A relatively simple example to the point is that K_4Fe (CN)₆ dissociates into potassium and ferroeyanic (Fe(CN)₆) ions.

⁴ Verhandl. d. Naturforsch. Gesellsch., Basle, 1874.

⁵ Ber. d. Natur. med. Verein z. Innsbruck, 20: 1891-92.

⁶ Zeitschr, f. Physiol, Chem., 22: 1896; 172, 188; 26: 1899; 588; also Deutsch, med. Woeh., 24: 1898; 1581.

¹ Zeitschr. f. Physiol. Chemie, 23:1897:309. Other observers (Schmiedeberg, Herlaut, Bang, etc.) give figures fairly consonant as regards the phosphorus and nitrogen contents, not so consonant as regards the hydrogen and oxygen.

THE CHEMISTRY OF THE CELL

Miescher⁴ found that 96 per cent. of the proteins of the heads of fish spermatozoa consist of nucleoprotein—these heads, in fact, are formed of nuclear material and little else; according to Lilienfeld,² 77 per cent. of the dried substance of the leukocytes from a richly cellular organ (the thymus) consists of nucleoproteins. We give these figures in order to impress upon the reader the wealth of nucleoproteins in the organism.

The more generally accepted formula for the ordinary nucleic acid of higher animals is $C_{at}H_{zr}N_{15}P_{4}O_{30}$. And this nucleic acid can be dissociated into the purin bases—*adenin* $(C_5H_8N_5)^3$, *guanin* $(C_5H_8N_5O)$ *xanthin* $(C_5H_8N_4O_3)$, *hypoxinthin* $(C_5H_8N_4O_3)$. Of these, the first two would seem to be the primary bodies, the others being derivatives by the removal of an NH₂ group and oxidation. Along with these, other nitrogenous bodies (*pyrimidin* rather than the allied purin bases), are capable of being split off, namely, *cytosin*, *uracil*, and *thymin*, together with a hexose sugar,⁴ and phosphoric acid in notable amounts, namely, about twelve times as much as is obtainable from the phosphoproteins of the cytoplasm (casein, etc.). So that Steudel has suggested the following equation:

And the following provisional formula:

¹ Miescher and Schmiedeberg, Arch. f. Exp. Path. u. Pharm., 37:1896; 1.

² Arch. f. Anat. u. Physiol., Abt., 1892; 128.

³ The important part played by these bodies in the animal economy recalls Pflüger's classical article upon the nature of living matter and the suggestion that it is from the "half iiving molecule," eyanic acid, with its remarkable power of polymerization into the polycyanic acids HaCuNAOn, that living matter originated. Arch. f. d. Gesammt. Physiol., 10: 1875; 251.

⁴ The previous teaching that nucleic acid afforded a pentose was due, according to Haliburton, to its frequent admixture with the allied guanylic acid (of Bang), relatively abundant in pancreatic nuclei, which is dissociated into guanin, phosphoric acid, and a pentose. (Science Progress, No. 14, October, 1909; 197).

³ While the edition has been passing through the press there has appeared an important study by Levene and Jacobs (summarized by Levene in the Jour, Amer, Chem, Soc., 32: 1910: 231), establishing yet more definitely the components of the nucleoproteins and the constitution of the nucleic acids. They found (with Steudel and others) that the acids and bases are present in equivalent proportions, the number of molecules of acid, earbohydrate, and bases being equivalent, and they give the following classification of the nucleic acids: (1) Containing one purin base, no pyrimidin, phosphorie acid, and a pentose (guanylic and inosinic acids).

Thus, to epitomize, the nucleoproteins, as regards their protein moiety, give origin to the mono-amino and di-amino-acids, and as regards their nucleic acid moiety, to certain important purin bases, pyrimidin bases, a carbohydrate, and phosphoric acid.

Thus, so far as we can at present see, it is the existence of phosphorus and of these purin-base groups, or of compounds vielding these groups, that differentiate the nucleus from the cell body. How the iron is combined, which is also a feature of nuclear composition, is as yet undetermined. We would emphasize the coincident presence in the nucleus, as distinguished from the cell body, of iron and phosphorus. To quote Herter! "This masked iron, as it is sometimes called, is doubtless of the utmost importance in bringing about the oxidative processes in the body, and any considerable diminution of the organic iron of the cell is probably attended by a diminution in the intensity of these processes. As regards the phosphorus, this also appears to be closely associated with oxidative changes."

It is, indeed, these oxidative powers of the nuclear matter that are, perhaps, its most striking features. Spitzer² has pointed out that those cells which are characterized by most active metabolism—the cells, for example, of the liver, kidney, and thymus, along with blood corpuscles—exhibit the greatest oxidative powers, and, what is more, that the nucleoproteids derived from these cells exhibit these properties to a marked degree, even when isolated. By appropriate staining methods Lillie has demonstrated that in the living cell oxidation proceeds in immediate association with the nucleus. It may thus well be that, as suggested by Loeb, the explanation of the difference between the nucleated and the non-nucleated cell is, that in the absence of the nucleus and its nucleoproteids, those oxidative changes that are at the basis of growth and regeneration cannot proceed.

(2) Containing two purin bases (guanin and adenin), two pyrimidin bases (cytosin and uracil), and phosphorie acid (*phytomadeic acids*). (3) Containing two purin bases (guanin and adenin), two pyrimidin bases (thymin and cytosin), one hexose, and phosphorie acid (nucleic acids of animal tissues, *lhymonucleic acids*).

By a most ingenious series of analyses Levene and Jacobs showed the order of combination of the components of the simple nucleic acids (those of the first group). They found, as Steudel had surmised, that the pentose in these acids connects the phosphorie acid on the one side with the purin base on the other. Contrary to all previous osbervations, they discovered the phosphorie acid to be orthophosphorie, and that the pentose is *d* ribose and not xylose, as usually taught. Thus, Levene determines the following formula for guanylic acid:



¹ Chemical Pathology, Lea Bros. & Co., p. 75.
² Arch. f. d. gesammt. Physiol., 67: 1897: 615.

THE CHEMISTRY OF THE CELL

Attention has already been drawn to the histological evidence of the relationship between nuclear matter and the cell enzymes (p. 47); it will not, therefore, be surprising to find that, from a chemical standpoint, intimate relationships have been detected between the two. As pointed out by Gustav Mann, the nucleoproteins and many enzymes have like solubilities, and as a consequence are brought down together in the attempt to analyze the tissues and body fluids. Thus, Hammarsten obtained nucleoprotein along with trypsin from the pancreas, Pekelharing and many other workers have noted the association of pepsin and nucleoproteins in analyses of the gastric mneous membrane and gastric juice, and of nucleoprotein and fibrin ferment in analyses of the blood and of the thymus. It is difficult not to see some association between the active oxidative processes on the part of living nuclear matter and the enzyme action developed or derived from the same.

Conclusions.—Let us now sum up the conclusions that may be reasonably deduced from the above data. They are:

1. The one group of substances common to all dead cell material is the group of proteins, with which water holding certain simple salts in solution is constantly associated, apparently as a medium.

2. One particular group of proteins, the nucleoproteins, forms the main mass of the cell nuclei.

3. The presence of iron and phosphorus in these nucleoproteins differentiates them from the proteins of the cell body as a general group. While some of the cell body proteins contain iron (c. g., hemoglobin) and other phosphorus (c. g., casein and the nucleo-albumins), none contains both combined.

4. The oxidative properties of the cell are associated in a striking manner with the nucleoproteins. Their particular constituents suggest that the nuclear proteins are characterized by an "energy" superior to that of the cell body proteins.

5. Both histological and chemical considerations indicate an intimate relationship between the nucleoproteins and the cell enzymes.

These conclusions, it will be seen, support and strengthen the conclusions previously reached from histological and physiological considerations that the nucleus is the dominant portion of the cell economy.

METABOLISM IN RELATIONSHIP TO THE CHEMICAL COMPO-SITION OF THE PROTEIN MOLECULE.

Thus far we have dealt with dead cell substance. We can by chemical and physical means break down these proteins; we cannot obtain substances which, when isolated, exhibit the properties of living matter. This, at least, is the usual statement. We shall point out that this statement does not express the whole truth; or, rather, that its truth depends upon what we regard as life. How are we to correlate our chemical with our physiological findings?

It is obvious, in the first place, that life and vital phenomena in general are directly connected with the presence of radicals peculiar to the proteins. We cannot escape this conclusion. The great variety

PROTEIDOGENOUS MOLECULES AND LIFE

of these substances, the enormous complexity of their molecules, the fact that scarce two analyses of any protein of what we may term moderate complexity afford identical results, all indicate that these contain very labile groups, that in the body they undergo constant change. We are forced to conclude, that is, that in the living organism they do not exhibit a fixed composition, but that there is a continual taking up and giving off of atoms and radicals; that in the living organism the bodies which we isolate as proteins exist in a condition of "moving equilibrium," what may be termed their average composition over a long period of time remaining constant, their composition over a labile molecules as formed of a ring of nuclei after the type of the benzole ring, or, more accurately, of a ring of rings, each component primary ring being a primary protein of the first order (Fig. 23).

Each such ring, it will be seen, has some of its affinities satisfied by the adhesion of the components of the ring one to another, but others are unsatisfied, and in such a compound ring as that here suggested, remembering that we are dealing with carbon containing nuclei, and that the carbon atom is tetravalent, the number of satisfiable affinities will be very great.¹ Accepting, also, what we know is the condition under which the protein molecules exist in the living state, these rings must be conceived as present in a fluid medium containing free atoms, molecules and ions of electrolytes; and we must regard the difference between living protein matter and "dead" protein as this, that the "living" active protein molecule is so placed that, owing to the attraction and chemical activities of the surrounding atoms and molecules, the side-chains of the protein rings are in a state of continual change, an unsatisfied affinity now becoming satisfied by the adhesion of certain of these surrounding atoms or radicals; and, again, other sidechains of the protein ring becoming detached owing to the more powerful action exerted upon them by the surrounding molecules; so that once again certain affinities of the protein molecule are unsatisfied; whereas in the "dead" protein molecule in the very process of preparation the side-chains have all become satisfied, and, being satisfied, the molecule is inert-that is, unable to form other combinations.

Following this conception, life is to be regarded as a state of persistent and incomplete recurrent satisfaction and dissatisfaction of certain protein, or, as we have expressed it earlier, proteidogenous molecules,² and metabo-

¹ Here, in relationship to the prominent part played by carbon atoms in proteid and proteidogenous material, and to the constitution of the same, it is deserving of note that the earbon atom is not merely tetravalent, but, as shown by the organic chemists, these carbon atoms have a pronounced tendency to become linked to one another to form chains, and, thirdly, these chains tend to group themselves into ring formations.

² Herbert Spencer's celebrated definition of "life," given in his *First Principles*, is vitiated by his neglect to recognize this necessary association of the phenomenon with proteidogenous matter. It is, in short, not a definition, because, wanting this limitation, it includes a series of manifestations which are not generally accepted as manifestations of the state of "life,"

THE CHEMISTRY OF THE CELL

lism—the constant reaction and interaction between these molecules and the medium in which they exist—must be regarded as the primary and basal characteristic of living matter, whereby, on the one hand, certain of the constituents of the surrounding medium are acted upon by the proteidogenous molecule, are either attracted *in toto*, or some of their dissociated component ions become attracted, with the result that the molecule becomes enlarged; and, on the other hand, owing to the attraction of the surrounding molecules, certain of the side-chains of the molecule thus elaborated become split off and form new combinations with those other molecules.




Diagram of unsatisfied proteidogenous ring, formed of nuclei A to F, with side-chain (H) and unsatisfied affinities (M).

The above definition demands that life be regarded as a kinetic state of matter of a certain order. Certain recent observations by A. Macfadyen¹ and Dewar open up serious doubts as to whether this is necessarily the case-as to whether we are not forced to recognize what we may term potential life. Macfadyen found that many pathogenic bacteria can be immersed in liquid air for as long as six months with no impairment of vitality. The temperature to which these organisms have thus been exposed is one equal to about -190° C. On removal, the organisms were found still to retain unimpaired their pathogenic and agglutinative properties; the prococcus aureus still gave rise to active hemolysis. Dewar has obtained like results employing boiling liquid hydrogen, thereby subjecting the bacteria to a temperature which was surely within 20° C. of absolute zero (-273° C.). At these profound temperatures not merely is there a very g eat absence of heat, but also of what we have shown to be of the prime importance to living matter, namely, moisture; and we would imagine that intracellular metabolism must practically cease. A consideration further of the condition of

⁴ Proc. Roy. Soc., October 31, 1902. Some eighteen years earlier Pictet and Young had published similar observations, though the temperature they gained was not so low (~70° to -130° C). Compt. rend. Acad. 4. Sciences, 98 : 1884; 747.

PROTEIDOGENOUS MOLECULES AND LIFE

living matter in the spores of many bacteria seems to point in the same direction, i. e., that for life to continue it is not necessary that there be constant interaction with the surrounding medium. Such spores have been kept dry for twenty years and more, and when brought into favorable conditions have actively proliferated. Nevertheless, Macfadyen's remarkable observations suggest that (at -190° C.) there may still be some molecular change in his frozen organisms. He found, for example, that photogenic bacteria still gave luminosity at these low temperatures, whereas if, while freezing, he triturated the bacteria, the luminosity was abolished; he thus determined that the luminosity was a function of the living cells. The matter must thus be regarded as still sub judice, though the presumption is in favor of the existence of absolutely latent life. So far as we can see at the present time, the distinction between living and non-living matter is this: that certain complex proteidogenous compounds form systems in which this recurrent satisfaction and dissatisfaction of the constituent molecules proceed, in favorable circumstances, with an activity and rapidity unknown in connection with any other chemical compounds or systems of the same known to us.

CHAPTER V.

THE CHEMISTRY OF THE CELL-(Continued).

ENZYME ACTION.

It will be noted that in the above suggested definition of life we state "certain protein or proteidogenous molecules." Why, it may be asked, distinguish thus between proteid bodies? All are built up along the same lines, and all, we presume, in the living organism have affinities to be satisfied and side-chains which may be broken off. We introduce this word "certain" because it is still undetermined what is and what is not to be included in our conception of life. Let us for the moment leave it out and attempt to classify the proteins and potential proteins in the living organism according to the extent of their activities.

The Organic Ferments.-Attempting this, we discover that in the class of free organic ferments we encounter our simplest cases. We use the term "simple" relatively, for it may well be that the constitution of bodies exhibiting ferment, or enzyme action is very far from simple; we employ it because by it we would indicate that typical examples of this class exhibit a single metabolic activity, acting specifically upon one single order of bodies in the medium that surrounds them. Of these free organic ferments here referred to numerous examples immediately present themselves: the ptyalin of the saliva, pepsin, rennin, trypsin, and the extensive series of other ferments of the pancreatic juice and of other digestive secretins. It is all important to obtain a proper grasp of the nature of these bodies, or, as will be made evident, of ferment action rather than of ferments-for the more we investigate, the more it is brought home to us that metabolic activities are of the nature of ferment actions, or otherwise, that, if not all, at least the majority of the manifestations of change in the proteidogenous molecule are to be included under this term. Constantly in studying cell functions, whether normal or perverted, we find ourselves brought to recognize that at base we are dealing with ferment action. It is, therefore, all important to gain, if possible, a right conception of what we mean when we employ the term. The above-mentioned group of free organic ferments present certain features in common:

 Elaborated in certain cells they are discharged and act outside these cells.

2. Each acts upon a particular substance or series of substances in the external medium—ptyalin upon the starches, converting them into soluble sugar, but not upon proteins; pepsin upon proteins in an acid medium, converting them to peptones, but not acting upon starches;

FERMENTS AND ENZYMES

rennin upon casein, and upon casein only; trypsin upon proteins in an alkaline medium; steapsin upon fats, and so on.

3. It has, so far, been found impossible to obtain the ferments in what can be regarded as a pure state; they are constantly found "associated with" bodies giving proteid reactions.⁴ More particularly are they brought down along with bodies of the nature of globulins and of nucleoproteins.

4. Even though these apparent combinations of ferment and protein are present in extraordinarily minute quantities, given sufficient time they are capable of converting a maximum amount of the fermentescible substance, provided their action be not arrested by the accumulation of the products of fermentation. And in the action they are not themselves destroyed.

5. The attempts to obtain the ferment pure and isolated from the accompanying protein resemble those of the old woman who attempted to reduce daily the fodder of her horse. Just as the horse died when she succeeded in reducing that fodder to a wisp or two of hay daily, so with the reduction of material giving proteid reaction in the ferment solution to a minimum, and by repeated separations, it is found that the ferment is destroyed, or, more accurately, that the ferment action disappears.

What does this indicate? It may be asked, in the first place, do ferments as such really exist; is there a particular class of chemical compounds having this particular property of acting upon other matter and breaking it up, without themselves being altered; or, on the other hand, are ferments as a class non-existent; is ferment action simply the expression or outcome of the molecular arrangement of the compounds exhibiting this property; may there be several different classes of substances possessing ferment action? When we find that pure metals, like platinum, gold, iridium, and silver-or, again, metallic oxides-can exhibit properties identical in their nature with those possessed by the highly elaborate cell substance, and that the more we study chemical activity the greater the number of reactions we find which appear identical with this organic ferment action, the first of these alternatives cannot be wholly correct; at most we can lay down that there exist various inorganic bodies capable of manifesting ferment action, and with these a group of organic compounds manifesting like properties.

The Enzymes.—We may, that is, proceed to deal with the latter as a single class and give them a common name, that of *enzymes*. Even when we do this we recognize that under this term we include two orders,

⁴ That certain observers have recorded the existence of active enzyme-containing fluids which have given no biuret test or other proteid reaction is not in itself proof positive that enzymes are effective when present in such extraordinarily small amounts that the ordinary chemical tests may well be too gross to demonstrate their presence, and, in the second place, may presume that, like the amino-acids, their composition may be so simple as not to afford the protein tests. At the same time we frankly admit that the evidence that all enzymes are of this protein—or amino-acid—type is presumptive, and has still to be proved.

the extracellular and the intracellular enzymes, of which the former act when completely freed from the cell body, the latter only when in connection with the cell substance, and then in such intimate connection that we can only conclude that the enzyme action is part and parcel of the manifestation of the living (protein) molecule. To quote a familiar example, the yeast cell, growing actively, secretes and discharges something-invertin-which, when the yeast cells are wholly removed by filtration, is still capable of acting upon the malt sugar present in the solution, inverting it, that is to say, changing it into glucose, but this cannot proceed farther and break up the sugar into alcohol and carbonic acid. That process necessitates the presence of the living yeast cell, or, as Buchner demonstrated, the expressed living substance. Buchner found that if a mass of yeast cells be subjected to hydraulic pressure so great that the cell membranes were ruptured, the thick, glairy fluid so obtained was capable of effecting the conversion of the sugar into alcohol, but this only when the experiment was conducted with great care, the results obtained being far from constant. When, by any means, the yeast cells have been previously killed, the reaction does not occur. The only conclusion to be reached is that here the enzyme, or ferment action is a function of the living and active cell substance; that in the experiment, while the yeast cells are ruptured, the cell substance is not actually destroyed, but still is able to manifest certain properties. The further conclusion is that some so-called intracellular enzymes do not exist as free bodies; were this so, we could extract from the yeast cell or from this emulsion the specific alcohol-producing ferment, and that we cannot do. We are forced, then, to the conclusion that in this case enzyme action is a function of the unaltered cell substance.

Having determined that this is the only adequate explanation in the one case, it may well be asked whether this is not the explanation of the activity of the other enzymes (i. e., free organic ferments). These, as we have pointed out, are, with rare and possibly doubtful exceptions, found constantly associated with, or affording, the reaction of proteins. May not enzyme action be a function of active protein molecules, each particular enzyme action being due to the specific structure of a particular variety of these molecules? In other words, if we admit, as we have to admit, that metabolism throughout is determined primarily by enzyme action, and that metabolism is the property of living as distinguished from dead nitrogenous matter, may we not regard the free enzymesptyalin, pepsin, and so on-as free living protein molecules, divorced from cellular relationship, but continuing to manifest the one important function characteristic of living, as distinct from dead, protein, that, namely, of acting upon other molecules in their neighborhood, and bringing about a rearrangement of the atoms without at the same time being disintegrated? May we not, that is, regard the free enzymes as the simplest manifestation of life?

THE MODE OF ACTION OF ENZYMES

THE MODE OF ACTION OF ENZYMES.

This may, to most of our readers, be a novel, and, indeed, a revolutionary conception. Before attempting to answer it, we must endeavor to gain a more intimate knowledge of the nature of enzyme action and what this demands.

There are two possible modes by which ferment action in general and enzyme action in particular may be brought about. We can, on the one hand, imagine that the ferment has no chemical action, *i. e.*, that it does not even temporarily enter into combination with the fermentescible substance; that its influence is purely physical. This view necessitates that we regard the ferment as a body possessing very active molecular vibration, such that, in apposition to molecules of the fermentescible substance, it communicates its vibrations to these, so bringing about a rearrangement of the constituent atoms whereby the fermentescible is converted into the fermented substance. This "contact action" is held to be a not infrequent reaction in "inorganic" chemistry; it is by this means that the action of finely divided platinum, gold, or iridium, in converting hydrogen peroxide into water and oxygen, is held to be best explained. This we may speak of as *catalysis* proper.

The second mode may also be exemplified from inorganic chemistry by the manufacture of concentrated sulphuric acid from sulphurous anhydride by the agency of nitric acid. In this process the nitric acid is held to act as intermediary; a reaction occurs between it and the sulphurous anhydride, with the result that it gives up an atom of oxygen to the latter—sulphuric acid being formed and the nitric being converted into nitrous acid. This first stage affords the formula:


$$H_2SO_3 + HNO_3 = H_2SO_4 + HNO_3$$

In the second stage the nitrous acid so formed, exposed to the air, combines with oxygen and so is reconverted into nitric acid, which now can act upon another portion of the sulphurous anhydride:

$HNO_2 + O = HNO_3$.


Theoretically, therefore, a single molecule of nitric acid can in infinite time convert an infinite number of molecules of sulphurous anhydride into sulphuric acid, and at the completion of the reaction will still exist as a molecule of nitric acid.

In such a process, it will be seen, three members or factors enter: the sulphurous anhydride may be termed the fermentescible substance, the oxygen the fermentator or complement, and nitrous (not nitric) acid the intermediary or the ferment. It is the nitrous and not the nitric acid that is present throughout the series of alternate reactions. We can visualize the process as follows:



 F_{i} , the enzyme molecule or ferment; F, S_{i} , the fermentescible substance; F, R_{i} , the fermentator or body which, with the moiety detached from the fermentescible substance forms the resultant compound.

And we can along these lines represent the particular reaction here described as:



Many considerations lead to the conclusion that it is reactions of this order, and not simple catalysis, that take place in the living organism and in connection with the free enzymes, but more particularly the positive evidence that the enzymes become temporarily attached to the fermentescible substance. Thus, if fibrin be placed in an acidified solution of pepsin at 0° C. (at which temperature the pepsin is inactive), left there for a little time, and then be given successive washings in ice-cold water to remove all adherent pepsin solution, upon being brought to body temperature it undergoes solution without further addition of pepsin. The pepsin, that is, formed a preliminary union with the fibrin. The protein molecule, as we have pointed out, must be regarded as possessing unsatisfied affinities, satisfied by the junction of side-chains. Our conception of the whole process of metabolism, as already noted (p. 67), is along the lines here indicated. The whole proteid molecule may broadly be conceived as acting after the manner of the molecule of nitrous acid in the above reaction; constantly, that is, it attaches to itself atoms and radicals from the surrounding medium, either free in that medium or by its greater energy broken off from other molecules, and constantly it liberates these that they may enter into other combinations. So that, as already stated, the average composition of the molecule remains the same over long periods. Broadly stated, the condition of "moving equilibrium" is precisely that of the molecule of nitrous acid in the foregoing reaction. It may be serviceable to express this diagrammatically (Fig. 24).

The extracellular enzyme may be regarded as one of the side-chain molecules (II), or even as only one of the components of the same (M), still retaining on its part certain particular affinities only, to which it can attach itself and combine with specific substances, the act of combination causing a dissociation of those substances. The exact nature of such dissociations will be discussed later. In other words, we regard these free molecules as the ferments of common parlance, or more accurately, regard such free primary proteidogenous molecules as possessed each of specific enzyme action.

Enzyme Action and Metabolism in Relationship to Growth.— This generalization, broad as it is, does not include every property of the active protein molecule, or, as we may term it, for convenience in subsequent description, the *biophore*, using that term to indicate the ultimate molecule possessed of what we regard as the properties essential to life.¹ In its fully developed form that molecule, as stated (p. 67), is obviously polymeric, composed of a chain or ring of primary molecules, each of which has proteid properties. That very constitution renders it at the same time intermediary body and fermentescible substance. For another basal property of living matter, the result of metabolism has to be taken into account. While its average composition in the state of moving equilibrium remains the same, *the number of molecules increases*. In other words, there is *growth*, and growth demands that

⁴ This term was originally introduced by Weismann to indicate the ultimate collection of molecules of living matter endowed with specific properties. He had not apparently regarded the molecule as polymeric, and so demanded an accumulation of several molecules to carry out the requirements of his theory. The term is, however, so adapted to convey our present meaning that we employ it, believing that essentially we indicate the same conception as did he, and that the word has priority over Verworn's *biogen*, which has the same significance, but what we regard as a faulty root meaning—i.e., we postulate that these molecules do not *produce* life, but inherently *bear* the vital properties. Rubner, whose recent *Kraft und Stoffe im Haushalte der Natur* (Leipzig, 1909) well deserves perusal, terms them " bionts."

not all the atoms taken up in the form of side-chains become released to form metabolites; some at least must undergo rearrangement and become built up into new biophoric molecules (Fig. 27).

We shall have more to say regarding the intimate nature of growth. What we would here emphasize is that all matter endowed with properties which we term vital does not coincidently possess this property of growth.




Diagram of growth, i, e., formation of new biophoric molecules. A side-chain of the main (dark shaded) ring has built upon it a like series of (light shaded) nuclei.

We have already brought forward®the most pronounced confirmation of this statement, although its bearing may not have been immediately realized; we have shown that the non-nucleated cell may continue "alive" for days, and, it may be, weeks—can perform several activities which we regard as characteristically manifestations of vitality—and yet it cannot grow; it becomes more or less rapidly used up in the performance of function. Translating this into the terms of proteid activities, it is obvious that the ordinary proteins of the cytoplasm, while capable of sundry enzyme actions—of respiration, for instance (which now we recognize more and more clearly is associated with the presence of oxydases), of converting starch into sugar, and so on cannot grow, cannot build up new molecules of matter like unto themselves.

The nucleoproteids alone (in forms, at least, possessing nucleated cells) are associated with this capacity for growth, and, when present, they are associated not merely with multiplication of the number of molecules of nucleoproteid, but also with the increase in the amount and number of molecules of the cytoplasm, and, lastly, judging from the observations of the histologists, they are directly concerned in the growth, if we may venture so to term it, of the enzymes, or otherwise discharge of nuclear matter precedes the evidence of active enzyme action in the cell (p, 48).

THE ORDERS OF LIVING MATTER

THE ORDERS OF LIVING MATTER.

What, then, to come at last to our point, are we to regard as life? If we say that our conception must include the capacity for independent growth, it follows that we must regard the cytoplasm as not living as dead. If, on the contrary, we deny that growth is an essential part of our concept, then, comparing the free enzymes and their properties with the cytoplasm and its properties, we must recognize three orders or grades of living matter:

1. The nuclear matter, capable of both metabolism and growth in a medium of cell proteins.

 The cytoplasmic matter, capable of independent metabolism of several orders, but incapable of growth, save in relationship with the nuclear matter.

3. The free organic enzymes capable of causing "metabolism" of one or other order, but incapable of growth.

We confess that it is difficult to lay down positively which of these two views should be accepted as correct. Upon first consideration, we should be inclined to lay down that the property of growth is inherent in our conception of life, and therefore of living matter; but, on the other hand, it is scarcely possible to regard the non-nucleated cell the red corpusele, for example—with its active powers of metabolism, as non-living. For our purposes it is perhaps fortunate that we are not compelled to arrive at a positive conclusion. We have in general to deal with the cell, which, as a whole, manifests growth. We do not, however, think that the matter brought forward in the preceding paragraphs has merely an academic value; we shall, that is, have so frequently to deal with enzymes and ferment actions that at the outset it is important to possess an appreciation of the same and their relationship to the proteins and cell activities.

These considerations, it will be seen, taken alone, would lead us to regard these free molecules possessing enzyme action as the most elementary forms of life. Some would urge that, since this mode of action is common to these organic molecules and to other substances, therefore inorganic matter undergoing chemical change is also endowed with vitality; that logically, therefore, all matter, as urged by Haeckel, and yet earlier by the elder Lankester, is endowed with life. A little thought will show that we do not advance thus far. We do but lay down that enzyme action is a property of unsatisfied proteid—or proteidogenous—matter, and doing this we limit our conception of life to proteidogenous matter, exhibiting a particular order of changes.

Objections to the Above Hypothesis. We have here, perhaps, overboldly laid down one view regarding enzyme action. We have done this after not a little consideration, believing that for didactic purposes this is the better course. The matter, however, is very far from being settled, and it is but due to our readers to point out that at the present time many leading physicists and physical chemists

incline to the catalytic view of enzyme action, the view, namely, that enzymes act not by making temporary chemical combinations, but physically, and by propinquity, without combination. It has, indeed, been doubted whether the type example here afforded of what may be termed inorganic ferment action-the action, namely, of nitrous acid upon sulphurous anhydride (p. 73)-truly represents what happens, the suggestion being that here again what really occurs is a true catalysis.¹ Yet it has to be admitted that the objections brought forward are of a comparative nature and theoretical; no positive proof is adduced that the reactions indicated do not occur, and the actual detection of the intermediate bodies, even if only in relatively small amounts, definitely favors the occurrence of the stages we have indicated. The main objection to the chemical nature of enzyme action is, as Oswald has pointed out, that such theory of intermediate action fails entirely to account for the action of negative catalysts, in which, if there be a direct chemical action, it must proceed more slowly than the direct action which takes place in the absence of the (negative) catalyst. These negative catalysts are bodies which, instead of accelerating, delay reactions; 0.0000014 gram per c.c. of mannite, for example, reduces by onehalf the velocity of oxidation of 800 times its amount of sodium sulphite in solution. Here, however, we are once more confusing catalysts and enzymes. We are prepared to admit that catalysis does occur among inorganic substances; the enzymes we place in a different category. Now, as a matter of fact, no negative enzymes are known,² no proteid or "proteidogenous" bodies having such properties. The objection, therefore, does not hold. So far as we can see, the intermediation theory is adequate for all enzymes proper.

There is yet a third theory, that of *surface action*, first propounded by Faraday to explain the catalytic action of spongy platinum, powdered charcoal, etc. In the union of hydrogen and oxygen on the surface of spongy platinum, Faraday supposed a condensation of the gases on the surface of the metal, through which condensation there proceeds a more rapid action between the two gases. Applied to enzymes, this supposes an attraction of the fermentescible substances and what we may term the fermentator, and condensation of the same on the surface of the enzyme, so that under the altered conditions the two can act directly and more actively the one on the other. This conception of surface action would seem to promise most valuable results in explaining many vital phenomena. The conception, it will be seen, is for practical purposes not very far remote from that of the loose chemical union on the one side and on the other demanded by the intermediation theory. It does not, however, fit in so well with the facts of growth and increase in substance brought about by the intracellular enzymes, for growth demands actual chemical union. Further, if, as we shall point out

¹ Vide Moore, loc. cit., p. 126.

² These negative catalysts are not to be confused with *antienzymes*, bodies produced by the organism after injection of foreign enzymes which neutralize them.

REVERSIBILITY OF ENZYME ACTION

later, the amboceptors (in hemolysis, etc.) are best regarded as enzymes, then with them we obtain evidence of chemical union with the fermentescible substances (or, as German writers term it, the substrate), and that in the absence of the fermentator (complement). While, again, the fact that enzymes when in the presence of proteins or other body upon which they exert specific action are able to withstand a higher temperature without loss of properties than they can in the absence of these bodies, is also in favor of the view that they enter into combination with the same.

We speak here, however, with some diffidence. The increased study of the phenomena of surface action and of the allied adsorption during the last few years, of phenomena, that is, which are intermediate between chemical and physical activities, makes it possible that such adsorption is at the basis of enzyme action. Thus, Bayliss¹ points out that were this purely chemical in nature then concentration of an enzyme to twice its value should double the reaction velocity; but this is not the case, the increase, as in adsorption phenomena, is something less than this. The fact also that they are of colloidal nature, and as such are heterogeneous systems (Bredig), favors the view that they form adsorption compounds.2 We must, however, confess to some sympathy with the views expressed by Brailsford Robertson.3 "It certainly appears premature at the threshold of our physicochemical knowledge of proteins to declare a group of their compounds or reactions to be 'adsorption compounds' or 'adsorption reactions,' and to thereby . . . group them with phenomena which are not improbably essentially different in character and in mechanism, and to import into a field already sufficiently obscure a conception so misty as that of 'mechanical affinity.'" That adsorption very probably occurs in the cell in connection with its lipoid constituents we are very ready to accept, but are far from assured that enzyme action comes into the same category.

The Reversibility of Enzyme Action.—In this connection, as throwing a peculiarly strong light upon the nature of metabolism, attention must be called to a remarkable property of the enzymes first demonstrated within the last few years by Croft Hill,⁴ of Cambridge. The sugar, maltose ($C_{B}H_{22}O_{11}$), is split up by the enzyme maltase into two molecules of glucose ($C_{e}H_{22}O_{21}$), but in the flask or test-tube the reaction

³ Science Progress, 1906; see also his article upon "Adsorption Phenomena," Biochemical Jour., 1: 1906; 175.

² For fuller studies upon forments and enzyme action, consult Reynolds Green, The Soluble Ferments: Fermentation, Cambridge, 1899; Duclaux, Traité de Microbiologie, 1; 1800; 1; and A. E. Taylor, Fermentation, Univ. of California Publications, Pathology, No. 8, 1907. Other important studies are O'Sullivan and Tompson, Jour, of the Chem., Soc., 57; 1890; 834; Croft Hill, ibid., 73; 1898; 634; and Jacobson, Ztschr. f. Physiol. Chem., 16; 1899; 340. See also the very full article by B. Moore, already eited.

³ Jour, of Biol. Chem., 4:1908:35.

⁴ Trans. Chem. Soc., 73:1898;634; also Proc. Chem. Soc., 18:1901:184, and Trans. Chem. Soc., 83:1903:578.

is never complete; there remains a mixture of maltose and glucose. Hill found that if the enzyme maltase be added to a solution of glucose, he obtained a similar reaction—some molecules of glucose became dehydrated and combined to form maltose, and the solutions came to contain a mixture of the two sugars. Maltase, therefore, can, according to circumstances, either d'sintegrate maltose into glucose or accomplish the reverse process of synthesizing glucose into maltose, and having these properties, it is inevitable that, so long as the products of either the disintegration or the synthesis are not removed from solution, neither the glucose nor the maltose can be entirely used up in the reaction; there must result a stage of equilibrium at which the enzyme becomes inactive, the tendency to disintegrate the one substance balancing the tendency to synthesize the other.

This reversible reaction we may express as follows:

$\begin{array}{rcl} C_{12}H_{22}O_{11} &+ H_2O &+ Maltase \xrightarrow{\sim} C_6H_{12}O_8 &+ C_8H_{12}O_8 &+ Maltase, \\ Maltose && Glucose + Glucose \end{array}$

Here the resultant molecules, being of the same structure, it might seem at first sight that the enzyme splits the maltose into equal halves, A little consideration shows that this cannot be. From its constitution the simplest dissociation of maltose is into one molecule ($C_{a}H_{\mu}O_{s}$), requiring for its completion an hydroxyl ion, and into another $(C_{e}H_{u}O_{e})$, that is completed by a hydrogen ion. We might represent the action of the enzyme as being primarily upon the molecule of water, the resultant O and OH ions combining with the maltose and leading to reduplication; or, on the other hand, regard the primary action of the enzyme as being upon the maltose. The latter clearly is the reaction that occurs, and this because the enzymes as a body acts by hydrolysis, and thus, were the splitting up a molecule of water into free ions of H and OH their primary action, then a multiplicity of specific enzymes such as we find in nature would be unnecessary. The maltase must act upon the maltose primarily. The steps in the process must, therefore, be (1) attraction between enzyme molecule and either the C.H.O. moiety of the maltose, or the $C_6 \dot{H}_{\mu} O_6$; (2) this becomes detached from its fellow and joined to the enzyme; (3) a vet greater attraction between the moiety thus detached and joined on to the enzyme and an hydroxyl or hydrogen ion, respectively, free in the medium with (4) combination with the same and simultaneous detachment of the enzyme ion which now becomes free to repeat the process with another molecule of maltose. Where the reverse action occurs, there the surrounding conditions must be such that the enzyme ion is so placed as to exercise a greater attraction upon the glucose molecule, combining with it and leading to the simultaneous liberation of either an hydroxyl or a hydrogen ion.

Even in cases in which, through enzyme action, the fermentescible molecule is split up into still more dissimilar molecules, the same reversible action is found to occur. Wells' gives a good illustration of this


 $^{+}$ Jour, Amer, Med. Assoc., January 25, 1902, gives an excellent $r\acute{o}sum\acute{e}$ of the work accomplished in this direction.

REVERSIBILITY OF ENZYME ACTION

in a reaction noted by Schmiedeberg in 1881, although its significance was not comprehended by him. It is a well-known fact that benzoic acid and glycocoll introduced into the circulation are synthesized in the kidney into hippuric acid, and Schmiedeberg found that, similarly, hippuric acid can in this organ be split up into benzoic acid and glycocoll. What is more, he succeeded in extracting from renal tissue an enzyme *histosyme*—which brought about the splitting-up process. He did not advance so far as to establish the fact that the histozyme accomplished the reverse reaction, though with our present knowledge this is seen to be evident; it may be expressed thus:

```
C_sH_bCO.NH.CH_2COOH + H_2O + (E) \underset{\leftarrow}{\longrightarrow} C_sH_bCOOH + NH_2CH_2.COOH + (E).
Hippuric acid + water + enzyme \underset{\leftarrow}{\longrightarrow} Benzoic acid + glycocoll + enzyme
```

Here we deal with the same type of reaction as in the former case. An hydroxyl (OH) ion and a hydrogen ion, respectively, combine with separate moieties of the hippuric acid so as to form benzoic acid and glycocoll. We may express this graphically as follows (Fig. 28):



The evidence now accumulated indicates that all enzyme action is, at least potentially, reversible. Kastle and Loevenhart' have shown that this is the case with *lipuse* or *steapsin*, the fat-splitting ferment of

⁴ Amer. Chem. Jour., 24: 1900: 491; and Chemical News, 83: 1901.

the pancreatic juice, and Hanriot¹ has confirmed. Wroblewski² has studied the reversible action of invertase. E. Fischer and E. F. Armstrong³ have noted the like action of kephir lactase. Acree and Hinkins⁴ have found that the hydrolysis of triacetyl glucose by pancreatin is reversible, while Emmerling^b has shown that yeast extract will synthesize glucose and mandelonitrile glucoside into amygdalin. Quite recently and independently, A. E. Taylor⁶ has described the synthesis of a protein (protamin) through the action of a trypsin obtained from the liver of a soft-shelled Californian clam, and Brailsford Robertson⁷ has synthesized one of the first products of the peptic digestion of casein, gaining the substance or substances to which the name "paranuclein" has been applied. The latter subjected an alkaline suspension of casein to peptic digestion for several days, heated to 100° C., to destroy the ferment; filtered, gaining thus a solution free from either casein or paranuclein; treating this with a pepsin solution, he gained in two hours a thick white precipitate giving the reaction of paranuclein.

The observations of Kastle and Loevenhart are of peculiar interest for our present purpose. Lipase acts on all the fats proper, and the results obtained apply to all the fats of the food; they employed ethyl butyrate for greater ease in analysis, and found that the lipase not only split up this into alcohol and butyric acid, but could unite these two latter to form the fat; and they employed this second reaction to detect the existence of the enzyme in different tissues. By this means they discovered that the liver is peculiarly rich in lipase; that this exists also in fair amount in the mucosa of the small intestines and in the kidneys, and to some extent in nearly all tissues. We shall have occasion to revert to these matters when dealing with the cell fats.

It is, however, significant that several observers have called attention to the fact that the reversed action of the enzyme is of a different order from the direct. Brailsford Robertson⁸ has pointed out that with pepsin it proceeds actively at a temperature from 10° to 15° C, above that at which the ordinary direct action on proteins (casein) is wholly arrested. At this higher temperature the soluble peptones are reconverted by the pepsin into a precipitated protein. It is difficult to reach any other conclusion than that this difference connotes some change in chemical composition, and Robertson suggests that the direct action is induced by the hydrated form of the enzyme (*F*.OH), which, giving up its hydroxyl ion, brings about hydrolysis and peptonization of the proteins, and that heating brings about dehydration of the enzyme, thereby favoring a withdrawal of hydroxyl ions from the peptones and conversion into insoluble proteins. This suggestion, it will be seen, is wholly in line with the conception of enzyme action hald down by us.

⁴ Compt. rend. de l'Acad. de Sci., 132:1901:212.

² Bull, Acad. Sci., Cracow, 1901:94.

³ Ber. d. Chem. Gesell., 35: 1902: 3146.

- ⁸ Amer. Chem. Jour., 28:1902:370.
- ^{*} Jour. Biol. Chem., 3: 1907: 87.

4 Ibid., 34; 1901; 3810.

7 Ibid., 3: 1907: 95.

[#] Univ. of Calif. Publications, Physiol., 3: 1909; No. 16: 188.

ENZYME ACTION AND METABOLISM

Thanks to these studies, we have gained a much clearer conception of the course of enzyme action in the organism, and can understand why now it proceeds actively, now is arrested. Equilibrium and arrest of enzyme action, we see, occurs when the products of that action accumulate up to a certain point, while, on the other hand, if those products be removed as they are formed, the action may proceed until all the fermentescible substance is taken up. In the alimentary canal, for example, the products of the action of the various extracellular enzymes, being soluble, are absorbed, diffusing into the mucous membrane and thence passing into the blood and lymph; there is thus in health complete disintegration of the proteins, starches, fats, and other foodstuffs. Within the organism in the cells, and in connection with the intracellular enzymes, it is again a matter of diffusion. Take, for example, the glycogenic activity of the liver cell. If the cell, in its metabolic activities, has used or burnt up the sugar, glucose, brought to it by the blood, and so becomes deficient in carbohydrates, more sugar will diffuse into it from the blood, and, acting on this sugar, the enzyme will synthesize it into glycogen, and will continue to do this until there is a local equilibrium between the sugar and the glycogen in the cell. Glycogen, being insoluble, remains within the cell-becomes stored up. And there it must remain until one of two things happens: until either, owing to stimulation, the cell being called into activity dissociates the contained sugar-uses it up at a greater rate than other sugar can diffuse into the cell from the surrounding medium. So soon as this happens, the glycogen-sugar equilibrium is destroyed, and now the reverse enzyme action comes into effect and the glycogen is dissociated, until, through this formation of sugar, the equilibrium is restored, when the dissociation comes to a stop. Or, again, the amount of sugar in the circulating blood and lymph becomes lowered in consequence of dissociation and consumption by the tissues occurring at a greater rate than the absorption from the digestive tract. When this happens, so soon as the percentage of sugar in the surrounding lymph becomes lower than that in the cell, the cell sugar, being soluble, will tend to diffuse out into the lymph. Here, also, the glycogen-sugar equilibrium within the cell will be destroyed, and the enzyme will become active until it may be, if the sugar contents of the blood remains lowered for any considerable period (as it happens in prolonged starvation), with the steady passage out of the sugars as they are formed, all the glycogen of the cell is eventually used up.

These considerations open up a new vista regarding disturbances of metabolism. These are seen to be dependent primarily upon the enzyme activities of the cell; the cell equilibrium, in fact, depends upon the amount of enzymes of different orders present within and produced by it, and this to a greater extent than it does upon the material absorbed by the organism and acted upon by these enzymes. By which we mean that the cell and the organism by their constitution possess, within certain rather wide limits, a power of regulating the absorption of matter from without, but if enzyme production be interfered with, then the regulating

power is largely destroyed. If our conception be correct, that the enzymes are proteins, or, more exactly, that enzyme action is a property of the constitution of the protein molecule, we find ourselves brought back to the fundamental conception that the due and orderly carrying out of the cell activities is a function of the composition of the proteidogenous constituents of the cells. When, further, we recognize that the development of the free enzymes in the cell is associated with active discharge from the nuclei, we gain additional support for our view that the nucleoproteids are, or are immediately connected with the biophores, the primary molecules of living matter.

CHAPTER VI.

THE CHEMISTRY OF THE CELL-(CONTINUED).

NON-PROTEID CONSTITUENTS OF THE CELL.

WE have discussed in some detail the proteid groundwork of the cell, and must refer to the non-proteid constituents, if only to afford a due appreciation of the cell complex. As in our treatment of the proteins, so here our object is not to afford an elementary treatise upon physiological chemistry, but to bring together those facts which have a direct bearing upon cell function, and more especially upon perverted function, or disease.

Water.-First and foremost, it is well to take into consideration the water contained in the cell. That water is an essential constituent. It may be reduced to a minimum, as in the seeds of many plants and the spores of bacteria, but when this is the case we find that the living substance passes into a state of latency, or inaction. Cell activity is associated with the presence of water, or otherwise water is the medium in which occur the chemical processes constituting metabolism. So important a constituent is it that close upon 60 per cent, of the human body, as a whole, consists of water, and of certain organs, like the kidney, water forms rather more than 80 per cent. Leaving out of account matricial, extracellular, matter, such as bone substance, it is safe to state that the ordinary active cell of the human tissues contains not less than about 70 per cent. H₂O; i. e., seven-tenths are water, three-tenths proteins and other constituents; so large a proportion that it is still with some a matter of debate whether we should regard living matter as existing and acting in a state of solution or as solid undissolved molecules suspended in a fluid medium.

The discussion is of something more than academic importance, especially in the light of the more recent studies of the chemicophysicists upon the nature of solution and the molecular changes which take place in dissolved substances. These studies, in short, suggest that an intimate knowledge of the physics of solutions must be of the very highest importance for a correct understanding of metabolic processes. At the same time it must be admitted that we are only at the threshold. We know from these studies that when a simple salt like sodium chloride is suspended in water and sundry other media, a certain number of its molecules become dissociated into their constituents, and these free constituents (Na and Cl, for example), charged some with negative, others with positive, electricity, we term *ions—cathions* and *anions*, respectively; that, if the dilution be sufficient, all the molecules undergo

this ionization; that these free ions thus charged conduct themselves like independent molecules and are in a state in which they may readily be attracted by other ions having an opposite charge, and thus simply, but indirectly, pronounced chemical reactions may be brought about. It is, indeed, these dissociated molecules that are active in chemical processes.

Stable chemical compounds, according to modern views, are formed by the coming together of ions-electrolytes-having contrasted electrical charges, and the very act of combination neutralizes or liberates the energy represented by these charges. These combinations in the case of solid substances may be broken apart in two ways, either by heat or electricity (by means, that is, of imparting, if we may so express it, such violent action to the molecules that the constituent ions are dissociated), or by solution. The act of solution, provided the amount of water be adequate, will similarly bring about the dissociation of all the molecules of a salt; or, more correctly, at a given temperature each molecule requires a certain fixed amount of water to effect its dissolution. We say "at a given temperature," for here, also, in solution heat favors dissociation. It is when molecules are thus dissociated that fresh chemical combinations can occur. It is the liberated ions and not the compounds as such that react one with the other; considerations which indicate the importance of water to the cell. In its absence, metabolic processes could only take place at high temperatures. The fact that matter is assimilated by the cell in a state of solution permits the extensive disintegration, rearrangement, and combination of ions which are essential to metabolism and growth, and this without the cell or the economy being called upon to afford abundant energy in the form of heat, etc., to bring about the dissociation.

The presence of so large a proportion of water in the cell cannot but indicate that metabolic processes are brought about essentially by ionization, resulting from solution and not primarily from the dissociative effects of heat, although the bodily warmth promotes the process. What is more, judging from what has been gleaned concerning hydrolytic action and saponification, it would seem that the dissociated hydrogen (acid) and hydroxyl ions (alkaline) play dominant parts in metabolic processes in general. Enzyme action would seem to resolve itself largely into processes of hydrolysis-and dehydrolysis-either the breaking down of a molecule into two through the combination with an hydroxyl or a hydrogen ion, or the reverse process of the withdrawal of hydroxyl ions. But so far anything like adequate studies upon ionization in connection with the protein molecules are very largely wanting, although a distinct advance has been made by Brailsford Robertson in his studies upon "ion-proteids." We have to keep in mind the point upon which we have already laid stress, namely, that the chemical composition of the *dead* protein molecule is not identical with that of the living-a point demonstrated by the fact that, while living

⁴ Jour. of Phys. Chem., 10, 1906; 524, and 11; 1907; 437, etc.

SOLUTION : CRYSTALLOIDS AND COLLOIDS

cell substance is alkaline or neutral, with death it takes on an acid reaction; from which it follows that observations upon albumins, etc., outside the body do not by any means fully inform us as to the processes taking place within the cell.

Turning now to the question whether the cell is to be regarded as liquid or solid, it must, in the first place, be noted that a characteristic of liquids is that in them the constituent molecules can vary their position freely in relationship one to the other, whereas in solids the relationship of the molecules one to the other is fixed. The distinction, it is true, is only relative, and in the cell we encounter conditions which are, to say the least, ambiguous; on the one hand, by careful study of certain cases-the white corpuscle, for instance-we can observe a streaming motion of the cell substance, *i. e.*, a free change in the relationship of the constituent molecules. On the other hand, the nuclear material is, to a large extent, fixed both in relationship to the cytoplasm and in the relationship of its individual parts one to the other. The idea of structure, it may be laid down, involves relative solidity. The explanation of our difficulty depends largely upon the colloid constitution of living matter. The cell in general and proteids in particular are colloidal, i. e., are composed of molecules, or molecular aggregates, so large that they cannot enter into perfect solution.

On Solution: Crystalloids and Colloids.—Observations upon the nature and physical properties of colloid bodies have of late years been abundant and important, until now it may be said that at last we are beginning to realize their significance and to have an insight into the physics of the cell, composed as it is largely of colloid matter. They differ from crystalloids in not crystallizing readily, in diffusing very slowly through water, in not passing through animal membranes, although, paradoxically, they themselves constitute these membranes and give them their peculiar attributes.

Graham, who first investigated the class of bodies, held that crystalloids and colloids are like different worlds of matter. We now recognize that the two terms are only relative, that there exists a series of bodies exhibiting the above properties which, nevertheless, like certain lipoids (soaps and the simpler albumins and hemoglobin), are definitely crystallizable, as also that a colloid may enter into true solution, comporting itself thus like a crystalloid, although the tendency is for them not to enter into true solution.

Our conception of a true solution is that of a mixture of the "solute" in the "solvent" of such a nature that the molecules of the former are freely separated by the molecules of the latter, intermingled so homogeneously that the mixture is optically inactive, obeying the same laws as govern the mixture of gases. The colloid mixture is of a different type; characteristically the molecules tend to form looser or firmer aggregates, floating in the solvent—aggregates so large as to interfere with the passage of light through the mixture, which thus becomes optically active, so large also that the aggregates are capable of detection by the ultramicroscope. In such a mixture, as demonstrated by Hardy,

there exist two phases: one termed by him "water-solid," that of true solution of the molecules of the colloid in the water, the other, "solid-water," in which the molecules of the water are taken up between the aggregated molecules of the colloid. Popularly speaking, in the first case, the water is the solvent; in the second, is the solute; and these two states are largely interchangeable according to the relative proportion of the two substances.

What for our purposes is of interest is that by physical meansby alteration of temperature, mechanical agitation, etc.--the smaller colloidal aggregates are apt to "conglutinate," or join together into flocculi, threads, etc. This property would seem to have some bearing upon the formation of fibrils in the cells and in intercellular matter. More particularly on surfaces where colloidal "solutions" come into contact with air, gas, fluid of another order, or even with colloidal solutions of different nature, there is a marked tendency for the molecules of the colloid to collect and spread out into a surface membrane of greater or less resisting power. The cell membrane interposed between the nucleoproteins of the nucleus and the cytoplasm may be instanced as concentration films of this order.1 To the physical properties of such membranes and their great importance in the establishment and maintenance of the cell as an independent aggregate we have already referred (p. 45). The nature of these membranes determines very largely the constitution of the cell.

Simple Salts.—Certain salts, without being built up into the protein molecules, are obviously essential to the cell; the protein molecules, that is, do not manifest their activity in a pure watery medium, but in a dilute saline solution. More particularly we encounter chlorine salts, alkaline carbonates, phosphates and sulphates, and salts of the alkaline earths, notably those of sodium, potassium, ammonium, calcium, and magnesium. From the more recent studies upon electrolytic dissociation we have learned to be cautious in laying down how these are combined; many, indeed, are under ordinary conditions present in such minute quantities that they must exist largely dissociated into their constituent ions, and so must actively promote metabolism. That their presence is essential for cell activity was shown many years ago by Ringer. More recently, Jacques Loeb and Moore have called attention to their importance and to the profound effects upon cell activity of comparatively slight variations in their relative amounts.

Certain salts, on the other hand, even when present in extraordinarily small amounts, are most deleterious to different forms of life. Thus, years ago, Raulin² showed that while a minimal trace of zinc (0.07 gram to 1500 c.c. of the medium) favored markedly the growth of aspergillus niger, silver in amounts too small to be detected (the mere keeping the

¹ This, perhaps, is not an exact statement; the membranes in question are coarser than any simple concentration film; it has, however, been observed that there is a pronounced liability for molecules of various orders to become deposited upon such films; thus, the ectosare of the cell and the nuclear membrane may be regarded as primarily built up on a concentration film.

² Ann. des Sci. naturelles, Botanique, 1870.

SALTS AND OSMOSIS

10.710

water to be used for growth in a silver jug for a short time) absolutely arrested the growth of the same. A similar deleterious effect has been noted in connection with forms much higher in the chain of living forms: a strip of copper placed in a vessel containing tadpoles leads to arrest of activities in two or three hours; while, again, it is a matter of familiar knowledge that hydrocyanic acid can arrest enzyme action, and arrest cell and individual life when present in quantities wholly disproportionate to its effects as a mere acid. The observations of Rankin, carried out in our laboratory at the Royal Victoria Hospital,⁴ upon the influence of metals upon bacterial growth and destruction, lead to the conclusion that the liberation of ions of oxygen ("nascent oxygen") through the interaction of the water and the metal in presence of oxygen is the effective agent in these deleterious processes. We do but mention these to indicate the subtlety of cell activities.

Turning to the more ordinary salts, their existence within the cell, or, perhaps more exactly, their dissociation and the building up of 'certain of their ions into the biophores, and the combinations undergone in the cell sap, are accompanied by important phenomena of endosmosis and exosmosis, *i. e.*, alterations in the anounts of fluid of the cell brought about by diffusion. Salts of higher concentration within the cell tend to pass into the medium of lower concentration without the cell, and vice versa, and this process is accompanied by a reverse passage of water into and out of the cell, the colloidal cell substance, and more particularly the more condensed ectosare, acting as a semipermeable membrane, permitting this interchange of water and soluble salts, while at the same time preventing the escape of larger (colloidal) molecules.² Nay, more, it is found that in regard to simple salts there is with different colloids a marked difference in the rate of escape.

This matter of esmotic change is one of no little significance; it helps us to understand why it is that the biophores and the more complex protein molecules remain in the cell, and why, on the other hand, smaller and partly dissociated proteid molecules—for such, as already indicated, we must consider them to be—namely, the peptones, pass out or again are absorbed with relative ease. The same considerations apply to the discharge of the free organic ferments, or enzymes, if our contention be correct that these, also, are simple non-polymerized protein molecules. There are yet other proteins which apparently are on the borderline. Their retention within the cell body is largely dependent upon the state of the cell relationship to the external medium. The red corpuscle retains its hemoglobin only in fluids having more than a certain osmotic pressure. From the red corpuscles of man the hemoglobin is extruded at a concentration of or corresponding to that of 0.47 per cent. NaCl;

⁴ Ramsden has shown that solid or highly viscous films are rapidly formed at the surface of protein solutions, and probably, as Brailsford Robertson points out, it is such a concentration film, rather than the coarser ectosare, that is the agent mainly involved. Ztschr. f. phys. Chem., 47: 1904; 336; B. Robertson, Jour. of Biol. Chem., 4: 1908; 1; Hardy, Jour. of Physiol., 24: 1899; 158.

¹ Proc. Roy. Soc., B., 82;1910;78.

of the chicken, of 0.44 per cent.; of the frog, of 0.21 per cent. If the osmotic pressure of the medium be less than these amounts, the diffusion of salts out of the corpuscle is associated with so extensive a passage inward of water that the ectosarc is ruptured, and now the hemoglobin—which is one of the simple, crystallizable proteins—undergoes solution in the surrounding fluid, and colors it.

This limit of tonicity at which the corpuscles lose their hemoglobin is, we may point out, very different from the n.rmal "tone" of the blood serum. According to Hamburger's observations, the serum of man and many animals is isotonic with a 0.9 per cent. sodium chloride solution. We employ the term *hyperisotonic* for solutions having an osmotic pressure higher than this; those the osmotic pressure of which is lower, as *hypisotonic*. The blood serum must, therefore, be markedly hypisotonic to bring about a condition of *hemoglobinemia*, or passage of the hemoglobin out of the corpuscles, purely through reduction of the saline constituents of the serum.

We shall have to discuss later how far variations in the saline contents of the cell and difference between the percentage of these and the percentage of saline contents of the surrounding medium determine the watery contents of the cell, and are the basis of œdema of the cell, and how far, again, considerations of isotonicity helps in explaining general œdema. As a general rule, it would seem evident, from what we have said, that the greater the passage out of water from the cell and the less the amount of contained water, the less must be the amount of ionization occurring within that cell, the less its metabolic activity, and this as a matter of experience we find to be the case; while, on the contrary, extreme absorption or osmosis of water must favor dissociation and disintegration.

Carbohydrates.-From the fact that free carbohydrates are never found in association with the nucleus, notwithstanding the existence of a carbohydrate radical in the nucleoproteins, we may perhaps infer that such radical enters the nucleus not as a free carbohydrate but already combined, as a glycoprotein. When present in the cell body, they represent either material absorbed from the external medium (or lymph) and not yet dissociated, or material so absorbed and partly dissociated, or, lastly, material built up within the cell as the result of cytoplasmic activity; i. e., through the cell energy other substances have been dissociated, certain of their ions have become seized by the cytoplasm, while those not so seized have interacted between themselves to form carbohydrate molecules. This last process occurs notably in the vegetable cell, in the building up of the starch granules in the chlorophyll grains. There the chlorophyll, under the influence of sunlight, breaks up carbonic acid, and the carbon so dissociated enters into combination with oxygen and hydrogen. The starch so formed undergoes ultimate cleavage-through enzyme actionand by hydration is converted into soluble sugars, which, in their turn, are dissociated, and the carbon containing molecules, in the nucleated cell, are utilized for growth. The botanists have demonstrated clearly that these starches can give origin to fats.

CARBOHY DRATES

In the animal cell the direct synthesis of carbohydrates from carbonic acid and water, if it occurs at all, must be an unusual event, but, a priori, there would seem to be nothing opposed to the view that in the dissociation of the protein molecule carbohydrates may be evolved, and, as a matter of fact, the continued excretion of sugar by the urine in advanced cases of diabetes in which carbohydrate food has been wholly cut off can have no other explanation than this, while Lusk's observations on phloridzin diabetes, in which he found a definite relationship between the amounts of nitrogen and sugar in the urine, indicate that certain protein molecules under certain conditions become split up into a nitrogenous and a carbohydrate moiety. It seems, however, equally clear that the carbohydrates found in the cell are, in the main, absorbed from the food-assimilated; and, further, that they are rapidly "burned" or dissociated to provide energy, unless they are by them converted into storage compounds within the cell, into the more insoluble *glycogen*, into a form, that is, allied to the starches. This storage occurs particularly in certain cells, notably in the liver and the muscle and (in pathological states) the leukocytes. We see, in short, that the carbohydrates of the food undergo a remarkable series of alternate conversions into soluble and insoluble forms. In order to be absorbed they must be in a soluble state, and so we find that the starches are acted upon by extracellular enzymes-by the *ptyalin*, which converts them into the sugar (maltose), and by the *amylopsin* of the pancreatic juice, which is even more active, although, as Herter points out, it is doubtful whether by either process there is complete conversion into dextrose or glucose, the form in which the sugar is eventually utilized within the organism. This conversion is held to take place in the cells of the intestine, and from these the glucose is passed into the portal blood. From this it is taken up, particularly by the liver cells, in which, if not broken up and utilized in metabolism, with the coincident production of heat, it is stored up, becoming for this purpose acted upon by an intracellular ferment and converted into the more insoluble glycogen. Here, it would seem, it remains until the amount of glucose in the blood circulating through the liver falls below a certain percentage; when this happens the altered relationship between cell and surrounding medium favors a reversal of the enzyme action, and now the glycogen is reconverted into glucose and diffuses out. The loss of glucose from the systemic blood is evidently, in the main, due to absorption by the muscle cells.

F. S. Lee' has proved clearly that these utilize carbohydrates in their activity, that deprived of carbohydrates by phloridzin poisoning they pass into a state of fatigue, from which they recover rapidly when carbohydrates are given in the food. In the resting muscle, as in the liver, the soluble carbohydrates are converted into, and stored up as glycogen. With activity, this glycogen is broken down and disappears, an important cleavage product being lactic acid.

¹ Amer. Jour. Physiol., 4: 1900; 9.

It is clear from the above that in the animal cell the carbohydrates are characteristically metabolites in the broadest sense; they are taken up, dissociated into simple compounds, combined into more complicated bodies under the influence of the cell substance. Where energy is needed it is obtained from their dissociation; where there is excess energy it is stored up within the cell in the form of the built-up, more insoluble glycogen molecules.

Fatty Compounds (Lipoids) and Alcohols.1-These form another important group of metabolites-of substances appearing in the cell and utilized by it. We are at the present time uncertain how to designate them. Neither from a gross physical nor from a histologicalmicrochemical point of view is it possible to give a clear classification. Under the term "lipoid" it has of late years become the custom to include those bodies which resemble each other in being dissolved (like ordinary fats) in ether, alcohol, benzene, chloroform, and carbon bisulphide (Bang²), although Overton, who introduced the term, employed it conversely to designate those bodies which dissolve narcotics. He based the theory of narcosis on the absorption and solution of narcotic substances by the intracellular fatty substances. Either definition includes substances like cholesterin, which, although constantly found in association with fatty compounds, do not themselves possess a fatty moiety. The study of these bodies has been carried forward with great activity during the last few years; rapidly they have assumed a high importance in the consideration of cellular biology, in metabolism, as also in connection with immunity, so that some workers in their enthusiasm assign to them a place in the cell economy almost if not quite equal to that of the proteins. The following classification is modified from that given by Bang:

- I. Substances Containing Neither Phosphorus Nor Nitrogen.
 - 1. Fatty Acids, Neutral Fats (glycerin-fatty acid esters), and Soaps (simple salts of fatty acids).
 - Cholesterines and Phytosterines present, either free or as esters of the fatty acids.
- 11. Nitrogen and Phosphorus-containing Lipoids-Phosphatides.
 - Mono-amido-phosphatide or Lecithin (a glycerin phosphoric acid ester of two fatty acids plus cholin; one of the fatty acids is saturated, the other not). Associated with this are possibly to be included a compound of lecithin with carbohydrate (jecorin ? from autolyzed organs), and with protein (lecithalbumin ?)
 - Mono-amido-diphosphatide, e. g., kephalin, from the brain substance, having two fatty acids, one unsaturated (kephalic acid), and two bases, one being cholin. Other lecithin-like bodies having an acid reaction are here included.

⁴ For our treatment of this section we are largely indebted to Professor Aschoff's full and suggestive paper, Ziegler's Beitr. z. path. Anat., 47; 1909; 1.

² Biochemie der Zelllipoide, Ergebnisse d. Physiol., 6: 1907, and 8: 1909; also Ergeb. d. inner Med. u. Kinderheilk, 3: 1909: 447.

- Diamido-monophosphatide, c. g., sphingomyelin, the most important constituent of the so-called protagon of brain substance.
- 4. Triamidophosphatide, found by Fraenkel in the kidney.
- III. Nitrogen-containing, Phosphorus-free Bodies.
 - Here are included the cerebrosides, resembling glucosides in composition, and all containing galactose. Among these are phrenosin and kerasin, both constituents of brain "protagon," and
 - The mono-amino fatty acids, or lipoproteins, recently synthesized by Bondi.

IV. Bodies of Fatty Nature Not as Yet Analyzed, e. g.: Lipochromes and certain Antigens.

Fats, Fatty Acids, and Soaps.—Just as the carbohydrates are stored in the cell, as starch or glycogen, so we find certain cells containing stores of fatty substances in the form of insoluble neutral fats, and we recognize a curiously parallel series of processes in connection with the metamorphoses that these undergo. We recognize, that is, that their presence is governed by the action of intracellular enzymes-lipases. These neutral fats are triglycerides of the fatty acids-combinations of a triatomic alcohol (an alcohol containing three hydroxyl groups C.H. (OH), with one of the fatty acids, the fatty acid replacing the hydroxyl groups. The three most important of these are stearin (tristearin), $C_{3}H_{5}(C_{18}H_{25}O_{2})_{3}$ palmitin, $C_{3}H_{5}(C_{16}H_{31}O_{2})_{3}$; and olein, $C_{3}H_{5}(C_{18}H_{33}O_{2})_{3}$. Other fats occur in the economy, but more rarely. Thus, in milk, butyric and caproic acids and their glycerides are to be differentiated, and in the liver Leathes has demonstrated the existence of glycerides of higher unsaturated fatty acids of the linoleic acid series. It will be seen from their formulæ that these neutral fats contain "a ha'p'orth" of oxygen to an "intolerable quantity" of carbon; in other words, that their dissociation and combination with absorbed oxygen is capable of setting free a relatively enormous amount of energy. Hence their value to the cell. How, it must be asked, are they acquired?

While we must admit that the glycerin moiety may be in part derived from carbohydrate dissociation, and that from the same source, if not also from the amino-acids, the higher fatty acids may undergo a process of synthesis, undoubtedly the preponderating source is from the neutral fats of the food. Vegetable oils, it is true, contain fair amounts of free fatty acids, but these are little employed for food purposes. But at most we have obscure indications of how this fat becomes worked up in the intestine; how it is absorbed by the intestinal mucosa; in what form it is transported and built into the neutral fats found in the body cells.

We know that the fats of the food are emulsified by the action of the various digestive juices; that free fatty acids are to some extent split off; that soaps are formed, and glycerin liberated. But only a part of the neutral fats are thus affected. We have evidence from Heidenhain's observations of a certain amount of phagocytic absorption of fat drop-

lets by the cells of the intestinal epithelium, and by wandering leukocytes, but this, again, is not adequate to explain everything. Are the free fatty acids also absorbed, and that in the form of soaps, or are they taken up as colloidal solutions by the cell substance? These are still matters of discussion. There are indications that in the intestinal wall the fatty acids are reconverted into neutral fats, for as such, upon feeding with fatty acids, they are found in the thoracic duct, although, as Aschoff points out, this conversion might take place in the intestinal canal prior to absorption. In the next place, the condition of the fat in the blood is still a matter of open debate. In diabetes and other conditions the fat may be present in the form of an emulsion; but is this a normal state? Certainly the studies of Arnold¹ and Fischler,² in which cells placed in soap solutions showed the intracellular formation of fine fatty granules, show that fat may be taken up by the cells in other than a corpuscular form, even if it be converted into a corpuscular form within the cell.

100

As regards fatty acids, these are found in the cell as crystals, but only in degenerative or necrotic conditions, as in pancreatic fat necrosis (see Part III, Chap. XXXII), in necrotic areas of the lungs, and in degenerating tumor cells, although they are normally present in fecal matter. As regards soaps, it would seem that at times they may be present and mistaken for fatty acids, their reaction with Sudan III being somewhat similar. The studies of Dr. Klotz,3 in our laboratory, have more particularly indicated that sodium and potassium soaps may be present in recognizable amounts in degenerative states, and that what has been known as the large white fatty kidney is, in some instances at least, the large soapy kidney. Insoluble calcium soaps have been demonstrated in areas of old fat necrosis of the pancreas, in the atheromatous aorta, and other areas in which calcification is proceeding, and, according to Klotz, these bodies form a necessary phase in the process of calcification. To these observations we shall refer when discussing that process.

Cholesterin.⁴—Cholesterin, C₂₂H₄₆O, is, on the one hand, a monatomic unsaturated alcohol; on the other, a complex terpene; and in constitu-

¹ Virch, Arch., 171: 1903; 197; Münchener med. Woch., 1903; No. 43; and Anat., Hefte 24: 1904; 389.

² Virch. Arch., 174: 1903: 338.

³ Jour. of Med. Res., 20: 1909; 27. Windaus has confirmed the isolation of a sodium soap from the kidney. (See Aschoff, loc. cit.)

⁴ It is the fashion now on the part of certain chemists to speak of this as cholesterol. The law deserves to be laid down that when a name has been in use for fifty years, and has beene familiar not merely to experts in one particular department, but to all interested in several branches of science, so that there can be no mistake regarding what is meant by it, no attempt to alter it so as to make it fall in line with some newer and possibly transient code, shall be countenanced. This applies to the names of species of animals and plants as well as to those of chemical substances. An absolutely unnecessary confusion has been developed by the recent changes in the names of the commoner intestinal worms by the lack of common sense on the part of certain purists.

CHOLESTERIN

8.23

470

tion is wholly unlike the proteins, carbohydrates, and fats.¹ This notwithstanding it is almost a primary constituent of the cell, being found in greater or less amounts in all the tissues of the animal body (as was shown by F. W. Beneke, in the early sixties), and being represented among plants by the closely related *phytosterins*. Saturated combinations, possessing two molecules of hydrogen, are found in certain excretions-isocholesterin in the skin and hair fats (lanolin), and coprosterin in the intestinal contents, the latter evidently derived from biliary cholesterin. Part of the biliary cholesterin, like the bile pigment, is reabsorbed, part discharged as coprosterin. According to Windaus, cholesterin forms as much as 15 per cent. of the dried corpus callosum, 2.3 per cent. of the whole brain tissue (dried), 5.9 per cent. of dried human liver, 0.5 per cent. of dried muscle. Ordinarily, it is not to be recognized in the tissue cells or in the excretions, but under certain degenerative and necrotic conditions its characteristic rhomboid plates, with a corner "bitten out," are present in abundance in atheromatous areas of the aorta, for example, in gallstones, in certain tumors undergoing simple necrosis, in certain old cases of inflammation of the tunica vaginalis of the testis with hydrocele, and of hydrops cystidis felleæ, or obliteration of the cystic duct with distension of the gall-bladder. And as it is chemically a somewhat inert substance, the question has arisen as to the conditions under which it ordinarily exists in the tissues. Is it merely dissolved in the fats, soaps, or phosphatides of the cell, or does it, as Moore holds, enter into loose combination with the fats, the fats taking the place of "water of crystallization," such combinations having the power, in addition, of taking up relatively large quantities of water and assuming a colloid state; or, lastly, is it mainly present as firmer combinations with the fatty acids as cholesterin esters? Moore and White have thrown doubt upon the presence in the organism of the latter compounds; Rosenheim and Windaus have definitely demonstrated their presence. Hürthle also has demonstrated the existence of cholesterin esters in normal blood. Increased interest has been taken in this subject since the existence of myelin droplets has been recorded in a large number of conditions-droplets which under the ordinary lens are indistinguishable from fatty globules, but between the Nicolls' prisms are seen to be doubly refractive. As shown by Adami and Aschoff,² this doubly refractive quality indicates that they are of crystalline constitution-the "fluid crystals" of Lehmann. Cholesterin itself cannot assume the fluid crystal phase, and as Aschoff brings forward abundant evidence that in the main the doubly refractive myelin bodies contain cholesterin, the very fact of the crystalline constitution of these bodies indicates that they represent a compound rather than a mere solution or loose association between cholesterin and fats. It should be

¹ For a very full study of its constitution and properties, see Moore, Manchester Med. Chron., 47: 1907: 204.

⁹ Proc. Roy. Soc., B., 78: 1906; 359. See also Adami, Jeur. Amer. Med., Assoc., 48: 190: 463. added that another group of myelin bodies, seen in autolysis, exhibits no double refraction.

The widespread diffusion of the cholesterins in living matter suggests that they are of considerable importance to the economy. It must be confessed, however, that we know little regarding these functions, save that, combined with other lipoids, they promote the absorption of large amounts of fluid, and the assumption by those lipoids of a diffused rather than a globular discrete state. They are signs also that they play some part in the resistance of the organism to infectious disorders; that, whereas the phosphatides play the part of activators to cytolytic enzymes cholesterin plays that of a protective substance.

The Phosphatides.—Only slowly are we unravelling the confusion that has reigned regarding the *protagon* which Liebreich (1865) regarded as the main constituent of nervous matter, and the *lecithins*. Although there are those who, like Gamgee, and Wilson and Cramer' have valiantly upheld the chemical existence of protagon, the tendency is now to accept the conclusion of Rosenheim and Tebb,² and of Posner and Gies,³ that this is truly a mixture of phosphorus-containing and phosphorus-free bodies, more particularly of sphingomyelin, as representing the former group, and of phrenosin, representing the latter. Similar confusion has reigned regarding lecithin, owing to the difficulty in gaining it in a pure state free from cholesterin. For the sake of clear conception it would be better to banish both terms and speak of phosphatides and cerebrosides.

These phosphatides, as already indicated, are very widely distributed through the body; they are a most important constituent of the nervous tissue, form a constant and considerable constituent of the ervthrocyte: are to be obtained in abundance from egg yolk, and are a notable constituent of the adrenal. Chemically, their significance is their nature as compounds of amino-acids with fatty acids and one or more nitrogenous bases; physically, they are characteristically colloidal, and have the capacity of absorbing large amounts of water, of dissolving other colloids, and thus acting as carriers for fats, cholesterin, and other cell constituents. By Albrecht and others they have been regarded as playing an active part in the phenomenon of agglutination of the red corpuscles into rouleaux; Preston Keyes and others have indicated that they are of signal importance in the neutralization of toxins and the production of immunity, acting as complement (see "Immunity"). Overton has determined that those substances which are able to penetrate into the living cell have in common the property of being soluble in phosphatides (lecithin and protagon) and cholesterin, and he and Albrecht have independently suggested that by peripheral concentration these phosphatides constitute the semipermeable cell membrane, although, as we have pointed out, Brailsford Robertson demands a concentrated protein membrane, with underlying lipoids. Koeppe, Peskind,⁴ and others conclude that the hemolytic activity of substances such as chloroform, ether, bile salts, etc., probably depends upon their being solvents of lecithin.

¹ Quart. Jour. Exp. Physiol., 1: 1908; 101 and 102; 1909; 91.

² Jour. of Physiol., 37: 1908: 34, and Quart, Jour. Exp. Physiol., 1: 1908: 300.

³ Jour. Biol. Chem., 1: 1905: 59.

Cerebrosides.—We know less regarding these, save that phrenosin and kerasin are constant constituents of nerve matter. It is possible that they are constituents of the myelin bodies seen in autolysis.

Lipoproteins.—As regards the other group of protein-free nitrogenous bodies, the lipoproteins, whose existence is indicated by Bondi's synthesis of amino-acids with fatty acids, the study of autolysis has opened up a fertile field of conjecture. It is a commonly known fact that tissues which, like the kidney, may afford on section not a sign of fat upon histological examination with Sudan III and other fat-staining dyes, will, nevertheless, by thorough prolonged extraction, as by the method of Dormeyer, yield eventually a considerable percentage. Indeed, as much as 17.9 per cent. of the dry substance of the normal kidney has been found to consist of fats. The indications are that this fat is shielded, i. e., is in loose combination, under which it fails to give microchemical reactions and is not liberated by the simpler solvents. As pointed out by Hildesheim and Leathes,1 after autolysis or self-digestion of pieces of liver tissue under sterile conditions, these simpler methods give a definite increase in the percentage of fatty acids. This process of autolysis is brought about by the postmortem action of the enzymes present in the liver and other organs. When, now, Bondi and Fraenkel² demonstrate that the lipoproteins which they synthesized are unacted upon by trypsin or pepsin, but are dissociated by the ferments of the liver and kidney, it is at least suggestive that there exist in the cell compounds of fatty acids and proteins such as, from other considerations, have been postulated by Quincke, Klotz, Gies, and others. But it is deserving of note that the proteins themselves are compounds of amino-acids, that is, of aminated fatty acids; that the phosphatides, again, like the nucleins, are compounds of aminated fatty acids with much phosphorus; that, in short, we are dealing with bodies which show certain marked relationships one to the other, relationships which we may with confidence expect to find elucidated in the not distant future.

Lipochromes.—Pigmented intracellular granules or globules, giving more or less perfect microchemical tests for lipoid bodies, are usually present in such small quantity that it is impossible to isolate and analyze them. We, therefore, are ignorant of their exact constitution. Such bodies are found in the lutein cells of the corpora lutea, in the cardiac muscle fibres in brown atrophy, in senile nerve ganglion cells, in the cells of the vesiculæ seminales, testes, and epididymis, and in the liver cells in conditions of atrophy. There are some indications that fatty granules are laid down first, which later take up pigment. But evidently we deal with bodies of more than one nature, for some (like the pigment granules in heart and liver cells) are stained by Sudan III; others give (although weakly) the bluish color with sulphuric acid characteristic of plant lipochromes and the bluish-green color with iodine-potassium iodide solution. Regarding the lipoid bodies which have been found to act as (debatable) antignes we shall have something to say later when discussing the Wassermann reaction.

rafa.

See Leathes, Lancet, London, 1909; i; Feb. 27.
 Biochem. Zeitsch., 17: 1909, p. 555.

CHAPTER VII.

GROWTH-RESERVE FORCE-STATES OF CELL ACTIVITY.

GROWTH OF THE CELL.

IF, as has been indicated, we leave it an open question whether growth is to be regarded as an essential attribute of living matter; whether, that is, we can still, or cannot, regard certain forms of matter as living which do not in themselves possess the power of growth, this, nevertheless, is evident, that the cell as a whole, manifests the capacity to grow, and, studying the cell as a unit, growth must be regarded as an essential and primary attribute. And this being so, it is necessary to inquire more closely into the nature of the process, more particularly because we shall find that a correct appreciation of what is included in growth is all-important for a fuller understanding of certain most important and widespread pathological processes, notably tumor formation or neoplasia—processes in which, before everything, we encounter disturbances of this fundamental property of growth. Only a little less important in this connection are the processes of repair and regeneration.

Whatever our conception of the exact nature of living matter, the idea—and the facts—of growth demand an increase in the amount of living matter; whatever our conception of the biophore, or ultimate molecule of living matter, in the process of growth two biophores must develop where previously there was only one. How is this brought about? The general idea we have gained of the constitution of the protein molecule here comes to our aid; that molecule, we recognize, is polymeric—formed of a recurrence of simpler molecules, which become attached one to the other in series. In bodies, therefore, of a proteid nature growth must be dependent primarily upon conditions which favor the building up of the simpler molecules. Once these are formed, their inherent nature and constitution, and their surroundings, must be such as to lead them to combine in series. In this process there must (a) be eventually reached a point at which the series is complete, and the biophore, as such, becomes an integral whole; but (b) with the completion the biophore cannot be a satisfied compound molecule. All our data regarding metabolism show that this is far from being the case; the constituent primary molecules must possess affinities to be satisfied, and what happens in growth is that these are satisfied by the building up and attachment of other primary molecules which, in their turn, attract other molecules of like order; and so there is gradually built up a lateral compound molecule which, in its turn, becomes com-

GROWTH

plete, whereby now two biophores are present where previously there was only one.¹

We can best visualize this process by regarding the complete biophore as a ring, or, more exactly, a ring of rings, each constituent being constructed after the type of the benzole ring. It is not essential to conceive each primary ring (proteid molecule) as identical, although for diagrammatic purposes we so represent them. And that the biophore is actually built up in ring fashion we do not pretend to lay down categorically;




Diagram of growth, *i. e.*, formation of new biophoric molecules. A side-chain of the main (dark shaded) ring has built upon it a like series of (light shaded) nuclei.

but such ring formation best fulfils the requirements of the case, namely, that the polymerization of the primary molecules is determinate, leading to the formation of larger units.² To represent the primary molecules as arranged in linear series would fail to indicate rationally the need for the series to combine or break up into these larger units or the inevitability of this process (Fig. 29).

Such a conception of the building up of new biophores may seem at first sight to assume a selective capacity on the part of the preëxisting

¹ In an interesting and valuable criticism of the first edition of this work Dr. C. P. White (Med. Chron., Manchester, June, 1909) takes objection to this conception of growth, doubting whether it is in accordance with "biological principles," and laying down that "all living units grow by interealation and divide by fission." This I know is the ordinary routine statement, but I have never been able to in the least comprehend it or to understand how new living matter is formed by "intercalation." The statement is in my opinion meaningless, *i. e.*, inconceivable, and it has been in the endeavor to form a rational conception of what growth really means that I have arrived at the above conclusions as to the essential nature of growth.

² The use of the term polymerization is here convenient, but possibly, from a ehemical standpoint, not accurate. Your true polymer is formed of a series of nuclei of like constitution; our conception is more that we have here to deal with repetitions of *series* of nuclei of unlike constitution.

100 GROWTH-RESERVE FORCE-STATES OF CELL ACTIVITY

biophores and their affinities, which is unknown in inorganic nature. A little thought will show that this is not so. All that is demanded is the action of the same laws as govern crystallization. In a mixed solution of salts and colloids, for example, provided the concentration of the sodium and chlorine ions exceeds a certain degree, we know that these ions combine to form sodium chloride; and if, again, the concentration of the sodium chloride similarly exceeds a certain degree, its molecules combine to form solid crystals, independently (to a very large extent) of the other salts and ions in solution; just these particular molecules of NaCl are attracted one to the other, and so in this mixed solution of various salts there are built up the specific crystals of sodium chloride. We might have chosen for an illustration some salt of more complicated constitution, but this simplest example is adequate for our purpose. What is demanded in growth and multiplication of the biophores is a like attraction of ions and molecules, a like assumption by those ions and molecules of precise relationship the one to the other that we see in the process of crystallization.

Accepting this as a working scheme of the constitution of the biophore, it would demand too elaborate a discussion, and one perhaps somewhat out of place, to lay down the apparent stages in the process whereby the complete cell with its nucleus, cytoplasm, and excreted enzymes has become evolved. All we would say here is that if we accept reversible enzyme action as a primordial function of living matter, we can then figure the various steps in the evolution of the cell from a mass of undifferentiated bioplasm. We reach, however, the following conclusions:

 That the fundamental cell material is formed of complete biophores, and these alone have the capacity to reproduce themselves and so initiate growth.

2. These biophores, we must assume, at the beginning were able directly to assimilate and build up into primary protein molecules material presenting itself in the surrounding medium, and these again gave origin to new biophores; but with the evolution of the more complex biophoric molecule the stage was attained at which newly formed (compound) biophoric molecules could not act directly upon the surrounding medium, but required the intermediation of primary molecules of proteid type constituting a surrounding evtoplasm.

3. In the completely developed cell the first stages of assimilation are accomplished by these subsidiary cytoplasmic molecules; the material is taken up by them in the form of side-chains, and these act as carriers or "enzymes" for the biophores, which, in their turn (a) still further dissociate the matter so presented to them, thereby liberating energy (p. 86); (b) build up that matter into primary protein molecules, which are returned to the cytoplasm, and so bring about cytoplasmic growth; (c) build it up into or govern the formation of the constituent molecules of new biophores. In other words, the nuclear material must be regarded as initiating and controlling the growth both of the nucleus and the cytoplasm. The above conception, it will be seen upon consideration, is the only one possible and consonant with the facts known to us, *unless we accept a primary duality of living matter*.

It may seem to the reader that in discussing these matters we pass into the region of transcendentalism. But if we recognize so fully as we do the constant differentiation of the cell into nucleus and cytoplasm, and if, further, we have before us a series of facts bearing upon the relative activities of those two constituents, we surely are bound to ask what is the intimate meaning of this differentiation, and how we can best harmonize these facts and bring them into line. While it is true that our pathology is cellular, and that we take the cell as our unit, to understand the activities, and the disturbances in the activities of the cell, we must form for ourselves some working scheme which shall take into account the indications afforded by the cell structure. With the accumulation of new facts it is possible that the views here enunciated may have to be considerably modified. For that we are fully prepared. In the meantime the facts already possessed appear to us to give a satisfactory interpretation only along the lines here suggested. And this conception of the essential nature of the cell must be the basis of our subsequeat treatment of our subject.

The Relationship between Growth and the Other Cell Activities. —Before proceeding farther it will be well to say a few words regarding cell dynamics, and this in order that we may realize the relationship between growth and the other activities of the cell.

It is clear, in the first place, that, speaking broadly, in the performance of function the cell is to be regarded as a machine for the evolution of energy. Motion, for example, demands the liberation of energy; secretion and the discharge of various formed substances from the cell indicate similarly a loss of material, or, in other words, a discharge of potential energy from the cell; in the so-called warm-blooded animals the increase in temperature of the organism as a whole above that of its surroundings necessitates that individual cells liberate energy in the form of heat; the nerve cell liberates energy akin to electricity. On the other hand, growth and the accumulation of new molecules of living matter demand not the evolution, but the storage, of energy. Each complicated molecule of proteid type represents a relatively great store of potential energy. Speaking broadly, therefore, growth and the performance of other functions by the cell are widely contrasted. Both demand that energy be previously acquired by the cell, but in the one case this is stored up, in the other it is dissipated. It is useful, with Weigert, to speak of these two contrasted phases of cell activity as bioplastic and katabiotic, respectively.

Looking more closely into the matter, we recognize that in the cell, in general, the necessary energy is acquired through the assimilated foodstuffs. These main foodstuffs—the proteins, carbohydrates, and fats—even in the soluble, assimilable state, are complex carboncontaining bodies, which, when decomposed and burnt up outside the body, yield relatively abundant energy. The excreta of the organism —*i. e.*, of its component cells—the carbonic acid, water, urea, etc., as a class, have exactly the opposite characteristics; they are relatively simple in constitution, are dissociated with difficulty, and yield little energy in the process. In the breaking down of the foodstiffs by the

102 GROWTH-RESERVE FORCE-STATES OF CELL ACTIVITY

cell it follows that much energy is acquired by the cell, and this energy is utilized either in the performance of function or in growth.

These considerations are apt to lead the physiological student to conceive that energy is evolved in the dissociation of matter. Paradoxically, the reverse is the case. In the act of dissociation energy is used up and becomes potential; it is in the act of combination that energy is liberated. The explanation of the paradox is that the ultimate result has to be taken into account. In the dissociation of the molecules of the foodstuffs there is a primary loss of energy. But these foodstuffs are unstable, their elements loosely combined, and when in dissociation their ions become freed they combine among themselves or with other free ions to form more stable compounds, and in this combination of ions having greater affinities it is that an amount of energy is evolved, much greater than that set up in the primary act of dissociation. When the candle burns it is not the dissociation of its wax that causes the light and the heat; it is the combination of the dissociated carbon with the free oxygen of the air that is the cause. Heat and energy, in fact, are used up in the dissociation of the wax, but its carbon being relatively lightly combined, the loss is small compared with the evolution that occurs in the combination of the carbon and oxygen to form CO₂. Similarly, energy is required to bring about the dissociation of the cell molecules, and similarly, also, it is the oxygen absorbed that is the great source of cell energy-combining with liberated carbon ions to form CO₂, with hydroxyl ions to form H₂O. It is these ultimate combinations that are the great source of energy.

In short, the biophores and protein molecules in general are not to be compared with simple salts, but with such highly unstable bodies as nitroglycerin. It is a matter of familiar knowledge that dissociation of the molecules of this compound may be brought about by very slight stimuli—as by sharp vibration—and that the dissociation is accompanied not by a loss of energy, but by a rapid and abundant evolution of the same—by an explosion—with the production of light and heat, this being brought about by a reconstruction of the ions of O, C, H, and N into simpler, more stable compounds.

It is thus that dissociation of the cell molecules leads to liberation of energy, and that growth and the building up of the complex biophores represents, on the whole, a using up of energy, *i. e.*, a conversion of kinetic into potential energy. Or, in other words, the energy provided by the assimilated food may be:

(a) Dissipated in the performance of function (katabiotic).

(b) Stored up in the formation of the complex molecules of the cell substance, *i. e.*, in growth (bioplastic).

Growth and the performance of function other than growth (which in succeeding paragraphs we shall refer to briefly as the "performance of function") are two contrasted states of cell activity.

It must next be asked, If contrasted, can they proceed simultaneously in the cell? Can the cell simultaneously perform function and grow? All, it will be seen, depends upon the rate of assimilation of food and energy compared with the grade of functional activity of the cell. And here we require another distinction between the two processes now under discussion. The conditions which, for general purposes, we group together as functional are all the responses, or the preparations for responses, to external stimuli; it is the external stimuli that call them into activity, whether directly or through the intermediation of the nervous system. Growth, on the other hand, is an inherent property of living matter, and depends primarily upon intracellular conditions. If abundant food be presented to and assimilated by the cell, and if at the same time the cell be highly stimulated, all the energy so acquired may be utilized in the performance of function, and no growth may ensue. If the stimulation be still more intense, not only will all the foodstuffs be broken up to afford the requisite energy, but also the living cell substance will undergo dissociation in order to supply the energy needed. If, again, the external medium provides little food material, and the cell is called upon to perform work, then also, to afford the requisite energy, the molecules of cell substance proper must undergo disintegration. We have to recognize, however, that there is a certain stage, or grade, of adequate assimilation coupled with stimulation from without of medium intensity, in which the dissociation of the foodstuffs provides more energy than the cell is called upon to dissipate, and it is at this stage that the surplus energy may be rendered potential in the process of growth. We can, that is, conceive growth and the performance of function as occurring pari passu, and, what is more, this conception harmonizes with experience-that whereas excessive muscular activity results in exhaustion and shrinkage of a muscle, a moderate grade of muscular activity, if persevered in, and if accompanied by adequate nutrition, leads to the growth and enlargement of the individual fibres.

Indeed, while we recognize that growth is inherent and the performance of function a response to external conditions, we have at the same time to admit a certain relationship and interdependence of the two processes. To take again the case of muscle, it is a familiar fact that abundant food, unaccompanied by exercise, does not lead to muscular overgrowth, but rather to the reverse. A certain grade of functional activity appears to be essential, not merely for the maintenance of the status quo of the cell, but for its growth. When, however, we look more closely into the matter, we encounter difficulties which prevent us from making any more precise statements, for there are certain apparent exceptions. In the embryo, for example, growth is at its maximum, function at a minimum; throughout adult life, after a certain stage has been reached, the reverse may be said to obtain; by steady exercise the muscles may be brought to a certain bulk, but further exercise leads to no further increase; there is a maximum size induced by the due performance of function, and this cannot be exceeded. And, thirdly, as we shall see in studying tumors, at all life periods there are cells in the body which may take on excessive growth, and this, so far as we can determine, irrespective of functional activity. Only within certain narrow limits does this relationship obtain. We may say that the normal cell of the adolescent and adult individual exhibits a rela-

104 GROWTH-RESERVE FORCE-STATES OF CELL ACTIVITY

tionship between growth and function, but that under certain conditions growth is largely irrespective of function. What these conditions are we must attempt to determine.

Active assimilation and active growth with little apparent functional activity, characterize the embryonic and foctal stages of existence. The more the cells become differentiated and recognizable as the specific constituents of the different organs the less capacity they exhibit "for growth and the more their katabiotic activity.

Accumulation of energy, in short, characterizes the growing stage of the individual and of the growing cell, and we may regard this storage as continuing until the volume of the cell-and of the individualreaches the point beyond which further increase in mass becomes not merely uneconomical, but actually harmful to the system. To repeat, we have to recognize that there is a definite relationship between surface area and mass, and that when the mass of a cell exceeds a certain point its surface area, proportionately to the unit of mass, decreases at a rapid rate, and both assimilation and discharge are hindered. When this point is reached there are the alternatives: either of cell division (by which there is a rapid increase in surface area relative to mass), or of diminution of the mass through the performance of function; or, in other words, through dissociation of some of the cell substance and liberation of energy. The first of these methods obtains more particularly so long as the system as a whole (the individual) is below what we may express as the economical ratio between its mass and surface area relatively to external medium and environment; the second becomes more and more marked as this ratio is approached.

Carrying this chain of reasoning to its logical conclusion, it will be seen that the size of the adult individual of any species is a function of the constitution of its biophores; is the expression of the optimum economy of interaction between those biophores and the external medium. Growth of cell and individual continues until this optimum is reached. Function—katabiosis—with the liberation of kinetic energy is the means whereby the optimum is maintained.

But what is more, in the developed cell there must be a constant alternation between these two. So soon as the cell performs function, be it glandular and excretory, or motor, or what not, it liberates energy, and this it can only do by a dissociative process; in other words, by the disintegration of the cell substance. Thereby the cell falls below the limit of optimum efficiency, and here, again, it is in the position to take to itself new matter and grow; or, as Weigert' expresses it, "the katabiotic use of material in function removes the obstruction to growth."

The most highly differentiated of all cells, namely, the neurons, once they have attained to full development and differentiation, almost cease growing, save for diurnal variations in size and the making up of loss brought about by the performance of function; neither do they multiply, remaining fully differentiated and active throughout the whole period of existence of the normal organism. Cell differentiation, in short,

¹ Deut. med. Woch., 1896; 635.

PHYSIOLOGICAL INERTIA AND HABIT

manifesting itself mainly as cytoplasmic change, necessitates a modified metabolism. It may well be that the formation of more complex cytoplasmic molecules in itself is a hindrance to the formation of new biophores; that these cytoplasmic molecules have to be dissociated into molecules of a simpler type before their material can be utilized in the formation of the biophores. We shall have to revert to these matters in discussing the subject of cell multiplication.

PHYSIOLOGICAL INERTIA AND HABIT.

A factor of considerable importance in determining whether a given cell employs its acquired energy for growth or for the performance of function is clearly what Fraser Harris¹ has termed the "physiological inertia" of living matter. When a resting muscle is stimulated it does not immediately pass from the inert to the actively contracting state. There is, as is well known, a definite latent period before it responds to the stimulus. Contrariwise, when a gland cell is stimulated, it continues to secrete after the stimulus has been withdrawn. The cell acquires momentum, resembling the wheel set spinning, which continues to rotate after the hand has been withdrawn. Many other examples might be adduced showing that the inertia of the cell leads it to continue in the same state, whether of rest or of activity, after the conditions have changed. Prior to Harris, Weigert² had called attention to the same property of living matter, although, perhaps, in not such clear terms. And Ehrlich, basing himself on Weigert, makes this the basis of his theory of immunity. In fact, as we shall have occasion to point out, it affords the basis of our comprehension of all cases of individualad aptation.

For this physiological inertia is the starting point of what may be termed "habit." A cell stimulated to perform a certain act does not merely continue to perform that act for some little period after the stimulus has ceased, but, what is more, on a second occasion a slighter

¹ Brit. Med. Jour., 1900; ii: 741. See also the *Functional Inertia of Living Matter*, London, Churchill, 1908.

106 GROWTH-RESERVE FORCE-STATES OF CELL ACTIVITY

stimulus will induce the like series of molecular rearrangements, until a period is reached when the minimal stimulus produces an optimum reaction, and with this the cell energies become employed in one or more particular directions, and activities in other directions are distinctly lessened. Once a cell starts to grow, there is a tendency for it to continue to grow rather than to perform function, until the increasing size and increasing tension and other (external) stimuli attain such an extent that now functional activity is called into play. This, in turn, once started, is apt to continue.

It may be urged that here we assume for living matter properties not possessed by non-living matter and thus go counter to our main deduction or guiding principle that, essentially, vital properties are but particular instances of properties possessed by matter in general. Now, it is interesting to observe that pure physicists like Lord Kelvin have independently called attention to phenomena exhibited by inorganic matter (e. g., the "fatigue" exhibited by steel wire undergoing recurrent torsion), which curiously resemble phenomena seen in the tissues. Professor Bose has brought forward experiments of a very striking order to illustrate the parallelism.¹ From this it will be seen that in the nature of its responses to stimuli, living matter differs at most in degree, and not in kind from non-living.

RESERVE FORCE.

These considerations lead us to mention another prominent characteristic of living matter, namely, the possession of what is well termed "reserve force." It is a property which deserves the fullest recognition by the student of disease, for upon its existence depends the whole process of healing. Every living organism is so constituted that, under ordinary conditions, its cells and tissues do not work to their utmost limit. Just as, acting on sound mechanical principles, we construct, or ought to construct, a wall, a floor, or a bridge, so that it is capable of bearing a load several times greater than any to which in the ordinary course of events it is likely to be subjected, so are the cells of the organism

¹ He used a "sensitive cell"—a photo-electric cell—composed of brominated silver plates connected with a galvanometer to record changes in the electric eurrent through the system, and was able to show that every kind of response characteristic of a vital organ, such as the eye, could be obtained from the non-living mechanism. "In both we have under normal conditions 'a positive variation' (of the injury or resting current, or current of reference), in both the intensity of response, up to a certain limit, increases with the duration of the illumination, in both there is comparatively little fatigue, the increase of response with intensity of stimulus is similar in both, and, finally, even in abnormalities, such as reversal of response, preliminary megative twitch on cessation of illumination, and decline and reversal under continued action of light, parallel effects are noticed." He even obtained electric afteroscillation strictly comparable with the after-action (after-image) in the retina when light is shut off.—Response in the Living and Non-Living, Longmans, 1902.

FACTORS OF SAFETY

constructed. Engineers employ the term "factors of safety" to designate the margin of safety required in construction—and this margin is very considerable; as Professor Meltzer' points out, the ordinary boiler is constructed of steel plates capable of withstanding from six to ten times the pressure per square inch to which in use they will be subjected. There is a large reserve of force or energy within them over and above that which they exert under normal conditions.

Numerous illustrations of the fact immediately suggest themselves: The huge reserve of muscle power which there is in the ordinary individual, unsuspected by himself and by those around him until he is profoundly excited or the subject of delirium, when the patient, so weak a few hours previously as scarcely to be able to raise his arm from the bed, requires all the force of two or three strong men to hold him down. Similar to this is the sudden and great strain which the heart muscle can successfully withstand in violent exercise; it has been calculated that the heart can perform from three to four times as much work as it accomplishes under normal conditions, and this without producing the state of overstrain. Examples of another order are to be recognized in connection with the various glandular organs. Thus, three-quarters of the rabbit's liver may be removed and yet the animal continue to live in apparently sound health. In other words, onequarter of that organ suffices to satisfy the needs of the organism, or, what comes to the same thing, under normal conditions the liver cells are working only at one-quarter of their capacity. Nine-tenths of the dog's pancreas may be removed without glycosuria supervening; ninetenths, again, of the adrenals without fatal results ensuing. Almost all the thyroid gland may be successfully removed without apparent harm (provided the removal be not permitted at a single operation); or, again, the whole of the spleen of the dog may be excised and health be unaffected. The stomach, the greater part of the small intestine, and large portions of the colon may be removed in different individuals without serious disturbances. Meltzer, quoting Triepel's studies, points out that muscles, tendons, and elastic tissues have no factor of safety in themselves (a point of interest in connection with the development of arteriosclerosis), but are provided with some such by their connection with other tissues; bone and cartilage possess a very large margin of safety.

¹ There has been no such full study of this subject as that afforded by Meltzer's lecture (*The Factors of Sofety*, Harvey Lectures, 1906-1907: 139), which, although he had sent me a reprint, had escaped my notice prior to the issue of the first edition of this volume. Our point of view is so similar that it might well seem that I had "stolen his thunder.' Nevertheless, although, as he states, "the question of the body being provided with factors of safety (had) never been clearly raised," it had been in the air for many years, and pathologists the world over had very generally accepted the idea of reserve force. Ponfick's observations upon excision of the liver were well known, and the immunity with which one of a pair of organs could be put out of action, whether by disease or removal. I can recall an interesting article upon "Reservekraft" by Nothnagel, in the eighties. As a matter of fact, this section had been written by me some years before publication.

108 GROWTH-RESERVE FORCE-STATES OF CELL ACTIVITY

Where the spleen is removed other related tissues take on and perform its functions, exercising thus a vicarious activity, just as does the pituitary body (hypophysis cerebri) when the thyroid is atrophied or removed,1 or, as, according to others, do the Brunner's glands of the duodenum when the pancreas undergoes loss of function. Taking the organism as a whole into consideration, it is clear that these examples of vicarious activity are at the same time examples of reserve force. Yet other examples indicating the existence of abundant reserve force are met with in connection with the various paired and multiple structures of the body; a single lung is capable of carrying on satisfactorily the all-important process of respiration; a single kidney, urinary secretion; numerous lymph glands, teeth, or fingers may be removed without evident harm to the organism. Even the most highly specialized organ of all the brain, is largely composed of paired "centres," and when one of a pair is thrown out of action the other may often take up its functions, more especially when, as in the case of the heart, larvnx, and alimentary canal (organs of median line development), it governs muscles which normally contract simultaneously on both sides.

A striking example of yet another order of this protective affluence of nature is afforded by Meltzer. He points out that at puberty the woman's ovaries contain on the average 30,000 ova, so that, even were regular menstruation, and not pregnancy, the more normal function, sixty times more ova are provided than the individual could ever possibly employ.

It is thus very clear that the organism is so constructed as to possess in most of its functions an abundant margin of reserve force. This is capable of explaining how it is that the human body is at once so marvellously complex and delicate a mechanism—responding to a variety and extent of stimuli in a way that no machine constructed by man can nearly approach—and so able to withstand wide diversities of environment and extreme strains upon and injuries to individual organs without being destroyed; as also the fact that there may be extreme local disease or destruction of parts without of necessity constitutional disturbance. Until injury or disease of a tissue has reached a certain point the cells of that tissue are able to fulfil the needs of the organism, and in many cases, even after this point has been reached, other tissues may vicariously perform its functions.

In short, in the existence of this reserve force lies the secret of the continued existence of the individual, the explanation, as we shall point out, of immunity to disease, and of the healing of injuries of every order; if, indeed, it be not the keystone of adaptation, and, in brief, of the evolution of the race. This, however, may rightly be said: that if the existence of this reserve force be kept steadfastly in mind, we are saved from continual misconception of the mechanism and meaning of many processes.

If, for example, we appreciate the existence of this reserve force, there is no longer any inclination to suggest that the action of the leukoeytes

¹ Rogowicz, Ziegler's Beitrige, 4:1888;453; Boyce and Beadles, Jour. of Path., 1:1893;223 and 359; see also Schönemann, Virch. Arch., 129;1892;310, and Herring; Quart. Jour. Exp. Physiol, 1:1908;281 (with hibliography).

THE STATES OF CELL ACTIVITY

in inflammation is purposive. If, then, they take up and digest living bacteria, this is not the assumption by them of a new function, or the exercise of a new force to meet the exigencies of the case. We now know that in conditions of health bacteria constantly, if only to a slight extent, gain entrance to the tissues and are destroyed by endothelial and other cells; that if a drop of the blood of any normal individual be taken, the contained leukocytes can be shown to have these phagocytic properties; in short, that it is a normal property of sundry leukoevtes to ingest, and, when possible, digest, foreign substances. Or, if again, we find that in the majority of cases in which there is destruction of the pancreas, glycosuria shows itself, whereas in some rare cases an equal destruction of this organ is followed by no glycosuria, we must not forthwith conclude that severe injury to this organ can play no primary part in the production of diabetes mellitus. There is the possibility, that must not be ignored, that in the exceptional cases above cited so great a reserve force exists or is developed by the vicarious activity, it may be, of other organs that excessive exhibition and waste of sugar in the system is effectually prevented.

And underlying the development of this reserve force we must see the action of this same force of physiological inertia. Life is more than the continual precise adjustment of internal conditions to external changes of environment. Through inertia there is overadjustment; through it, when the cell assimilates, it continues to assimilate more than is actually needed; when it is stimulated to metabolize, it continues to form more paraplasmic matter than is necessary for immediate excertion; when it starts to grow, the extent of growth is over and above the extent of the initial stimulus. And, although these are the exceptions and not the rule, we are not without instances of cases, such as the cells of the mammary and sebaceous glands, in which, when once dissociative changes are initiated, they continue until the cell is completely disintegrated.

THE STATES OF CELL ACTIVITY.

These considerations prepare us to recognize certain states of the cell depending upon the ratio between assimilation, growth, and stimulation from without; states which it is well that we should recognize, for in conditions of disease we constantly encounter transitions from one to the other of these.

1. Subnormal Activity.—Under wholly normal conditions the process underlying the accumulation of reserve force leads to the presence in many tissues of redundant cells, cells which, from the accident of position, receive minimal stimulation and so pass into a latent, relatively inert state. Under abnormal conditions many cells may pass into this state. With lack of stimulation these cells undergo a very distinct alrophy: the paraplasmic matters are used up, the cell body becomes shrunken and inconsiderable, the stainable substance of the nucleus diminishes in amount, the nucleus as a whole becomes inconspicuous.

110 GROWTH-RESERVE FORCE-STATES OF CELL ACTIVITY

An excellent example of the passage into this state of subnormal activity and its results is to be seen in the muscles of a limb, whether of man or animal, kept forcibly at rest. If, for instance, through injury to the knee, or fracture, one lower limb of a muscular young adult be enclosed in plaster, it is a matter of familiar experience that within a week there is a marked diminution in the circumference around the middle of the thigh of the inmobilized as compared with the free limb. A similar and more extreme atrophy is to be encountered in the muscles of cases of hysterical paralysis. We mention these more particularly because in organic paralysis, due to recognizable injury to the nervous system, it is still a matter of debate to what extent injury to the "trophic" nerves is responsible for the atrophy.

If this condition of subnormal activity and latency be continued too long, the cells, or some at least of them, die and wholly disappear. That this is the case is well exemplified in the brain. In that organ, we need but remind the reader, nervous impulses are in many instances conducted through relays of cells; there are definite tracts along which specific impulses, and those impulses only, are conducted. If the neurons of the upper portions of such a tract be destroyed by disease or by removal, the second series of neurons with which they communicate can receive no impulses, and, as a consequence, are rendered largely inactive. As a matter of fact, in such cases we find that these secondary centres show pronounced atrophy of their constituent cells; these cells become greatly shrunken, and in the course of a few weeks or months their number is markedly reduced. We can proceed so far as to lay down with confidence that where the cells of basal nuclei do not exhibit this atrophy and disappearance, there the centres have not been connected with the destroyed area, and that where the atrophy is only partial or transient, there the affected centres have been, and are, in connection with more than one peripheral centre, thus continuing to receive stimuli, although not to the normal extent.

While we cannot follow Grawitz to the whole extent of his theory, or accept all the arguments upon which that theory is based, we have to admit the existence of certain orders of *Schlummerzellen*, or "sleeping cells"—cells atrophied through inaction, but capable under stimulus of returning to full vigor and full development. That these undergo such a grade of atrophy that, as he insists, while continuing to exist they become invisible, I cannot accept.

2. Vegetative Activity.—Cells in the process of active growth present certain well-marked characteristics. The nuclei are relatively large, rounded or oval, and deeply stained; paraplasmic granules and deposits absent, or, if present, in but slight amounts; the cell body unformed, tending to be rounded or oval, the cytoplasm exhibiting little differentiation. Such cells are apt to multiply, and we shall have more to say regarding them in our chapter upon Cell Multiplication. Here we would only give the warning that, from their general resemblance to the cells of the growing embryo, in which this type of cell predominates, it is customary to speak of these as *embryonic* cells. This

THE STATES OF CELL ACTIVITY

is a misnomer, leading to false conceptions, for cells of this type are to be encountered in normal tissues at all life periods. It is better to speak of them as vegetative cells.

3. Functional Activity.—Cells actively functioning as a general rule present signs of differentiation according to their specific function. Either, as in the case of muscle or nerve cells, the cytoplasm is highly elaborated, or, as in the case of gland cells (and the bodies of the neurons), there are paraplasmic deposits, granular, or in the form of minute globules. The nuclei are relatively not so large; the staining of the same varies with the stage of the cell activity. Here, however, distinctions are to be made between the cell action under a mean or normal stimulus and one that is more intense. We can thus further distinguish the following states:



To show the distinction between functional and vegetative cells. Two "clear cells" of regenerating liver, rounded, large, and clear, owing to relative absence of the paraplasmic granules seen in surrounding active cells. One of these exhibits mitosis. (Adler.)

4. Hyperactivity within the Limits of the Reserve Force of the Cell.—When, as already indicated, increased stimulation is accompanied by adequate nutrition, the functional activity of the cell is, up to a certain extent, accompanied by growth. In this way and to this extent increased work of the cell leads to what is termed hypertrophy. The dimensions of the cell are increased, the nucleus is of good size, the cell body presenting well-marked differentiation.

5. Excessive Functional Activity .-- When, on the other hand, the work that the cell is called upon to perform exceeds a certain point, the energy dissipated in the performance of function being greater than that acquired from the assimilated foodstuffs, provided that the stimulation be continued sufficiently long, then, first, the reserve force of the cell is used up, all the paraplasmic stores becoming exhausted, and next, the extra energy demanded has to be obtained from the disintegration of the active cell substance-cytoplasm and nucleoplasm. The result is cell exhaustion. The nucleus becomes poorly staining; the cell body exhibits a variety of changes, according to the specific activity of the cell. In some cases, i. e., cells of the convoluted tubules of the kidneys, there is actual breaking off and discharge of portions of the cytoplasm; in other cases, abnormal deposits occur within the meshes of the cytoplasm; in others again, owing to osmotic changes, the cytoplasm becomes vacuolated. These we shall have to describe more fully when dealing with the degenerations. If the stimulus be further continued, the result is cell death and complete disintegration.

CHAPTER VIII.

CELL MULTIPLICATION.

INCREASE in size of the tissue and of the multicellular organism as a whole is brought about by two processes: by enlargement of the individual units, the cells, and by the intercalation of new units derived from those already present. We speak commonly of increase in size of the individual as growth, by whichever of these two processes it be produced, and thus are apt to confuse the two, considering cell growth and cell multiplication as practically synonymous. While admitting that the second process is a sequence of the first, it is, nevertheless, well to keep the two perfectly distinct in our minds, and this we have endeavored to do thus far. In our treatment of the subject of growth we have taken into account purely the enlargement of the individual cell units by increase in the amount of living substance of the same. Following upon this, it is essential that we pass in review the subject of cell multiplication.

The rationale, if we may so express it, of cellular structure has already been touched upon (p. 100). It has been shown that the nucleus is to be regarded as the dynamic centre of the cell; that the nuclear biophores are to be regarded as requiring for their full activity a certain proportion of cytoplasmic matter, and that, in its turn, this cytoplasmic matter, for the due exercise of its functions, must be in intimate relationship with the external medium; that in a roughly spherical body, with increase in diameter, the surface increases at a far less rate than does the mass, so that growth of the individual cell beyond a certain limit is self-inhibitory, unless some means be employed to increase the surface in proportion to the mass of the cytoplasm; that when this is accompanied by coincident growth of the nuclear matter, means have also to be employed to increase the nuclear surface in relationship to its mass; that cell and nuclear multiplication, respectively, are the simplest means of accomplishing these objects.

Now, it is in place to describe the mechanism whereby this multiplication is brought about. The finer details and the variations in the process belong to the domain of the histologist and cytologist. Fortunately, throughout the cells of the higher animals, which more especially interest us, the same broad plan is to be recognized—the variations are very slight. Hence it is possible to describe in general terms the processes of cell division, knowing that what is said applies to individual cells, to the cells of the different tissues of the human body. We recognize two main types—direct, or *amitotic* division, and indirect, *mitotic* or *karyokinetic*.

AMITOSIS

DIRECT DIVISION: AMITOSIS.

This is the rarer form, although not so rare as it has been the custom to describe it. In text-books published within the last six years the statement will be found that among higher animals it is confined to the leukocytes. This is incorrect. There is an increasing number of observations indicating that it obtains also under certain conditions in cells that we regard as much higher in the scale. It used to be laid down that the cells of the metazoan (multicellular) embryo never exhibit it, but Patterson,¹ in the pigeon embryo, found frequent amitosis in certain regions of all those germ layers which were growing very rapidly, with indications that amitosis might be followed by mitosis. It is not thus, as Flemming² held, necessarily indicative of degeneration. In the fully developed organism, in tissues formed of aggregations of similar cells, in the liver,³ for example, we may encounter it, and it would appear to be particularly liable to occur in cells having the tendency to be multinucleate, cells exhibiting two or more nuclei without immediate separation of the cytoplasm into distinct cell bodies around each of the nuclei. It is in leukocytes and in cells of endothelial type that we encounter it most frequently, and in yet another group of cells, namely, those of pathological new-growths.

It is frequent also in the cells of the embryonic envelopes of insects, of the periblast of yolk nuclei, and in the syncytial (epiblastic) cells of the mammalian embryo. All the cells of this order are destined to but a temporary existence.

It is possible that further study will show that the cells in glandular and other organs already referred to which exhibit this direct division of the nuclei are also not wholly normal, or otherwise that amitosis is a sign of regressive change. As regards the leukocytes, it is worthy of note that in normal lymph glands, where the lymphocytes are continually being produced, we encounter frequent cases of mitosis or indirect division, while it is in the blood and in the tissues in conditions of inflammation that we meet with the amitotic division. Vom Rath,⁴ indeed, lays down that "when once a cell has undergone amitotic division it has received its death warrant; it may, indeed, continue for a time to divide by amitosis, but inevitably perishes in the end." This is, we think, too extreme a conclusion.

In addition to Patterson's observation noted above, Bashford⁵ records, in connection with the connective-tissue cells of the host in the neighborhood of transplanted portions of mouse cancer, that these cells at first divide by amitosis, and that eventually the products of amitotic

¹Anat. Anzeiger, 32:1908:117. In the frog and toad, Reichenow (Arch. f. mikr, Anat., 72:1908:671) could, however, find no instances in which it was not a precursor of cell degeneration, or occurred in cells very slightly differentiated.

² Areh. f. mikr. Anat., 37: 1891:249.


³ See Reinke, Verhandl. Deutsch. Anat. Gesell., Kiel, 1902.

⁴ Zool. Anzeiger, 14:1891:331.

⁶ Second Report of the Imperial Cancer Commission, London, Part 2, 1905.

-8

division undergo active mitosis, eventually giving rise to the stroma of the growing tumor. A somewhat similar case is the early amitotic proliferation of the cells in an inflamed area followed by later mitosis.⁴ The conclusion to be reached—provisionally—would seem to be that cells of low type like connective-tissue cells, when in an actively vegetative state, may exhibit amitosis which is not necessarily degenerative; whereas amitosis occurring in cells of higher type is, at least, suggestive of degeneration.




Amitosis. Stages of direct division in tumor cells: A, from an ovarian cancer; B, from an epithelioma of the hip: C, from a uterine sarcoma; D, from a metastatic cancer of the liver, showing the last stage of division of a cell into three equal parts. (Nedjelski,)

What happens in *amitosis* is that the nucleus divides without any apparent preliminary rearrangement of its structure. It becomes elongated, then dumb-bell shaped, and after a period in which (as can better be followed in the amecba) there is a certain amount of streaming of the nuclear material between the poles, the connecting neck becomes broken across and the two daughter nuclei pass apart, their separation being in some cases followed by division of the cell body, so that thus two complete daughter cells are developed; in other cases this further division is wanting, and the binucleate or multinucleate cell is produced (Fig. 31).

It deserves note that in this process, according to the majority of observers, the centrosome either plays no part or at most the attraction

¹ Similarly in the simple plant, *Spirogyra*, Pfeffer (Berichte Konigl, sachs, Gesell, d. Wissensch., 1899, July 3) has shown that in water containing 0.5 to 1.0 per cent. of ether the cells continue to multiply, but that by amitosis; transferred to ordinary water the cells now return to the normal mitotic division.





MITOSIS

sphere forms a ring around the equator of the dividing nucleus, *i. e.*, the part played is distinctly abnormal and unlike what is seen in mitotic division.

INDIRECT DIVISION: MITOSIS.

This is, par excellence, the natural mode of cell division, and that in animals and plants alike. That it should be so widely distributed indicates that the remarkable succession of changes seen in both nucleus and cytoplasm is not a matter of chance. The full significance of these changes we are still far from comprehending. One thing is obvious, namely, that they indicate a mechanism whereby the nuclear material is distributed with remarkable exactitude between the two daughter cells. And they indicate clearly something more. Were the biophores or essential constituents of the nuclear material all of the same nature and composition, the law of economy suggests that no such elaborate "quadrille" of the nuclear material would be indulged in; simple direct division into two equal halves would suffice, each daughter cell receiving approximately equal amounts of the nuclear material. That the nuclear material arranges itself in this remarkable manner prior to division may, in itself, be taken as proof positive that there is a differentiation of the biophores, and that mitosis is a mechanism whereby identical groups of biophores are conveyed into the daughter cells. Light will, we think, be shed upon the significance of the process when we come to consider the subject of heredity. For the present we shall merely detail the usual stages in the process of mitotic cell division. (See Plate II.)

The Stages of Mitosis.—For a full discussion of the phenomena of mitosis, as again of the part played by the cell in inheritance, the reader is referred to works upon histology, and more especially to Professor Wilson's valuable monograph.⁴ Here I can but in the briefest possible way recall the main features of the process.

1. Prophase or Preparatory Stage.—The nuclear chromatin which in the resting state of a cell is seen as an irregular and nodulated network, becomes modified into a continuous single (or, very rarely, a segmented) thread, having the appearance of a *skein* or *tangle*, and then proceeds to divide into a definite number of short lengths, the *chromosomes*. While these changes are proceeding the nuclear membrane disappears, so that the chromosomes come to lie naked in the cell. Every species of animal and plant has a fixed number of chromosomes, and in the mitosis of the cells this number regularly recurs. In man, more recent observations would indicate that the number is thirty-two. Side by side with these changes other changes take place outside the nucleus, in the cytoplasm or cell substance, leading to the development of the *amphiaster* or *spindle*. This arises under the influence of the *centrosome*; very frequently, while the nucleus is still at rest, this divides

¹ The Cell in Development and Inheritance, 2d edit., New York, Macmillan, 1906,

CELL MULTIPLICATION

into two similar halves; around each minute dot the protoplasmic network of the cytoplasm becomes concentrated, the fibrils radiating in all directions, so as to form star or *aster*, and, as the two separate, journeying toward opposite ends of the cell, a *spindle* of fine fibrils is seen to stretch between them.

2. Metaphase.—Each chromosome splits lengthwise into two exactly similar halves, the daughter chromosomes becoming apparently attached to certain mantle fibres of the spindle. This splitting of the chromosomes, discovered by Flemming in 1860, is the fundamental process of cell division.

3. **Anaphase**.—The daughter chromosomes diverge, the two members of each pair passing to opposite poles of the spindle. Here the chromosomes become closely crowded near the centre of the aster.

4. Telophase.—The cell body now divides into two in a plane passing through the equator of the spindle. Thus, each daughter cell contains half the daughter chromosomes, half the spindle, and one centrosome and aster. The two latter may persist or disappear; if they persist they form the *attraction sphere*. Before, during, or after the process of cell division there occurs the construction of the daughter nucleus. The commoner process is for the daughter chromosomes to fuse into a skein or tangle, which in its turn becomes irregularly swollen and dissociated into the chromatin network of the "resting" nucleus. Several recent observers claim that in the resting nucleus of certain species it is possible to distinguish the course of the original chromatin thread, and even of the different chromosomes that go to compose that thread.

CHAPTER IX.

ADAPTATION.

WE have suggested (p. 115) that the procession of changes seen in mitosis indicates that the biophores or specific ultimate molecules of living matter are not all identical, and in this and the succeeding chapters it will be seen that what we have to say practically centres around biophoric modification; around the extent to which the biophores, and through them the cells in general, become modified in their properties, and—as the properties of any substance depend upon the constitution of the same—in constitution. Here, again, it may at first seem a far cry from matters such as this to the needs of every-day pathology, but in reality, as we hope to demonstrate forthwith, a comprehension of these matters is essential for a proper grasp of the remarkable and superabundant facts elicited during the last few years in the study of immunity-a branch of pathology which has received of late more attention than has any other. And, although it is far from being generally recognized, it is through these studies that the pathologist and the bacteriologist are laying the foundation of an adequate theory of variation and inheritance.

Descent and variation are subjects which we have to dwell upon in future chapters as a foundation for our treatment of the inheritance of diathesis and disease and of the remarkable group of abnormal growths which we include under the heading of monstrosities and abnormalities. As a basis for our study of all these subjects, it is fitting that we first take up the subject of adaptation.

That living matter has adapted itself to its environment is a commonplace. Man and all other animals and plants exhibit countless evidences of the fact that each form of life is adapted to the particular environment in which it flourishes. But, admitting this, we are apt to ascribe the process of adaptation to chance. The zoologist and the botanist, recognizing that all living beings vary one from the other, that no two individuals are exactly alike, are apt to ascribe adaptation to the retention and descent of favorable variations; the individual, varying from "type" in a direction which gives it the advantage over other members of the species or tribe, is more liable to survive; if the variation be unfavorable, life is rendered more difficult and the individual and the stock descended from that individual tend to die out, they being at a disadvantage. There is seen to be a survival of the fittest, and it is by the summation and descent of favorable variations that the different species are adapted to their particular surroundings. This is the prevailing doctrine. Studying it, we see that adaptation

ADAPTATION

is regarded as based on chance; chance variations are at the bottom of the whole process.

If our studies in infection and immunity have any meaning, they teach us that this is not the truth—or at least not the whole truth. Adaptation is primarily an active process, or at least inevitable, and only subject to chance to this extent, that the individual may be impotent to choose or to control the changes in environment to which it becomes subjected, but once subjected to those changes the results are determined by inexorable law. It is not the mere fortuitous passive modification of living matter in a favorable direction, but a process whereby that living matter is able to a greater or less extent to change and suit itself to its surroundings, a given change in those surroundings leading to definite and corresponding alteration in that living matter. This we would emphasize.

For a comprehension of racial and species adaptation we have to begin with a study of individual adaptation. In connection with conjugation and amphimixis (the fusion of the germinal nuclear matter of the two parents in the fertilized ovum) chance undoubtedly enters, but only secondarily. We would, in the first place, afford the proof that adaptation is a regulated process affecting the individual, and, in the second, would seek to determine how a property apparently so wholly unlike those possessed by matter of all other orders has come to be developed.

It is a well-known fact that bacteria, like other living organisms, assimilate food through the action of enzymes, and these both extracellular and intracellular. Some bacteria, for example, living in media containing proteins and albuminoids possess active proteolytic ferments, whereby these bodies are reduced to soluble peptones, and may be still further dissociated, with indol as one of the ultimate products. Others more particularly act on sugars, splitting up these with the production of organic acids and gas (H and CO_2). On removal from their natural habitat, and growth upon the artificial media of the laboratory, the different bacteria exhibit these proteolytic and glycolytic properties in varying degrees: some have little or no proteolytic activity, others little or no glycolytic power; some ferment one particular sugar only, others a variety. We have, indeed, established our classification of the B. coli and allied forms largely upon these fermentative properties.¹ But now, as we believe was first pointed out by Peckham,² if the typhiol

¹ See Ford, The Florm of the Human Intestine, Studies from the Royal Victoria Hospital, 1: 1903: No. 5. In this most painstaking and elaborate study of the bacterial contents of the intestines of 50 cases Ford isolated as many as 50 different "species" of bacteria. Of these, 36 were non-spore bearing and non-pigment producing, and of these it will be seen that there are several groups containing three to seven members, each of which differs from its fellows only according to the fermentation or non-fermentation of one or other sugar. The recent studies upon the bacillus of epidemic dysentery recognize at least five "species" (some observers elaim very many more) indistinguishable morphologically, but each having a different action upon a series of sugars added to the medium of culture.

² Jour. of Exp. Med., 2: 1897; 549.

ADAPTATION IN BACTERIA

bacillus, which normally does not produce indol, be grown in a relatively strong proteid medium free from sugar, and be passed, over a considerable period, from tube to tube of this medium, there is eventual indol production. This cannot be said to be the result of chance-it is inevitable. Take any apparently normal culture of the B. typhosus and place it under one particular set of conditions, and the proteolytic indol producing function will manifest itself. Similarly, as pointed out some years ago by Sir Lauder Brunton and Macfadven," growth of certain bacteria in media containing particular sugars eventually results in those bacteria gaining the power to ferment the particular sugars. The property is not acquired immediately, but, with a given species, we can foretell absolutely that it will be acquired within the course of a few days, or at most weeks. Thus, in a recent paper, Klotz² has pointed out that the B, perturbans-a form intermediate between the B. coli and the B. typhosus-was able to ferment glucose when first isolated from water, but only gained the power of fermenting lactose and saccharose after growing in lactose and saccharose broths for some days.

The organism was then placed in a celloidin capsule and inserted into the peritoneal cavity of a rabbit. Left there for three days, it was found to have lost its power of fermenting the two latter sugars, regaining it, as regards saccharose, after forty-eight hours' sojourn (two passages) in saccharose broth; as regards lactose, after four days' incubation. The experiment was repeated by placing some of the stock culture in a celloidin capsule in the peritoneal cavity of a rabbit and leaving it there for one hundred and forty-four days. On removal, there was a slight fermentation of the glucose broth at the end of the first day; saccharose fermentation appeared on the fourth day; lactose fermentation on the sixth transfer, and then only at the end of seventy-two hours' growth; by the eighth transfer gas appeared in fair quantity. Work along these lines has recently been carried still farther by Twort,³ who, taking a series of members of the B. coli group which had been grown for a long period upon ordinary laboratory media, retaining fixed type characters, was able, by growing them now for long periods upon media containing unaccustomed sugars, to cause a certain number to eventually dissociate sugars which at first they did not ferment. He thus found that all members of the paratyphoid subgroup would ultimately ferment saccharose; the typhoid bacillus acquired the property of fermenting lactose and dulcite, and the dysentery bacilli of Shiga and Flexner ultimately fermented saccharose within twenty-four hours.

The same is true as regards resistant powers toward deleterious agents. Growing the Bacillus coli in broth made up with progressively increasing strengths of corrosive sublimate solution, beginning with 1 part in 100,000, von Hansemann⁶ was able to obtain active growth of the

- ¹ Proc. Roy. Soc., 46:1889:542.
- ² Jour. of Inf. Disease, Supplement, 2: 1906: 35.
- ^a Proe. Roy. Soc., Biol., 79: 1907: 329.
- * Descendenz und Pathologie, Berlin, 1908: 123.

ADAPTATION

microbes in strengths of the disinfectant which would kill the ordinary colon bacilli. Nor is this only true of vegetable forms; we need but recall the remarkable observations of Ehrlich, abundantly confirmed by other workers, upon the development of arsenic-resistant forms of different species of trypanosomes, produced by treating animals infected by these pathogenic protozoa with atoxyl and other arsenic-containing drugs.

The same is true also as regards pathogenic properties. As Vincent¹ has shown, it is possible to take absolutely non-pathogenic forms, like the B. megatherium and B. mesentericus vulgatus-forms which may be inoculated by the million into warm-blooded animals without the slightest disturbance being set up-and accustom or adapt them to growth within the warm-blooded animals by inserting celloidin capsules containing pure cultures of the same in the peritoneal cavities of these animals. These capsules, it may be explained, permit the diffusion of the body fluids, and so of nutritive material, and at the same time prevent the direct action of the body cells on the bacteria and the escape of the contained bacteria. After being grown thus for some months, upon removal of the capsules and making growths in culture media outside the body, it is found that the bacteria have become pathogenic, are capable of growing within the tissues when injected direct, and of causing the death of the inoculated animals. In other words, the bacteria now produce enzymes and other products capable of acting deleteriously upon or poisoning the animal tissues.

All these, it will be seen, are examples of the acquirements of new properties on the part of the lower organisms by adaptation. Within certain limits—at present by no means clearly defined—the simple forms of life are able to adapt themselves to their surroundings, and the adaptation cannot be ascribed to chance, for, with a given environment, the one particular alteration in properties surely results.

Let it be clearly understood that we do not pretend to lay down that these lower organisms can eventually enter into combination with and adapt themselves to every possible substance dissolved in the medium of growth or that every attempt to modify the properties of bacterial species is fraught with success. This is far from being the case. All we state is that the observations made so far indicate that there are certain substances with which living matter, or its metabolites, can enter into a more or less close combination, and toward which, therefore, it can adapt itself.

To these conclusions it has been objected that what, after all, we are dealing with is the survival of the fittest; that it is still a matter of chance; that among the thousands, not to say millions, of bacteria in a culture—owing to the *inherent tendency of living matter to vary*—it happens that some exhibit variation such that now these particular bacteria are able to ferment the unaccustomed sugars, etc.; that these having gained the new power, by chance, are at an advantage as compared with the others

¹ Ann. de l'Inst. Pasteur, 12: 1898: 785.

ADAPTATION AND IMMUNITY

which have not varied in this direction, and multiply at the greater rate, and their descendants, starting from this vantage ground, are even more likely to vary farther in the same direction, so that the particular property becomes exalted; so that, in short, in the process of time the descendants of the form exhibiting the favorable variation alone are represented. It is admitted that the new property is not gained at a bound; that a considerable number of "generations" of bacteria must pass before the acquired property is pronounced.⁴

So far as it carries, *i. e.*, as affording an alternative explanation of the phenomena, but not, it must be noted, as proof positive that adaptation is not active, this argument is quite valid. It is, however, demolished if we can show that adaptation can take place under conditions in which there can be no question of the survival of the fittest, in individual cells, and that with such certainty and in so short a period relatively to the life period of those cells that the process can only be of an active nature. And this we can do—at the very other end of the scale of living beings.

Acquired immunity in man, as in all animals, is adaptation, and this, again, is not a chance process; we can take germs-the cholera spirillum, for example—which, from their habit of life, must have at all times had a restricted local existence until man came on the scene and aided in their distribution, germs which, therefore, cannot at any time have affected certain of the lower animals in other regions, so that there can be no valid suspicion that at some remote period the ancestors of those animals had been subject to infection by, or had responded to, those particular species of microbes. Injecting these microbes into such lower animals, guinea-pigs, rabbits, and so on, we determine that they and their toxins are poisonous; so that with very considerable accuracy we can measure what fraction of a centigram of the toxin will cause the death of 100 grams of guinea-pig, rabbit, or other animal, within forty-eight hours. And, having determined this, we can by repeated injections of fractional portions of the lethal dose of the toxin so alter the constitution of the warm-blooded animal that now it can withstand ten or one hundred times the lethal dose without ill effect. Granted that we deal with healthy animals, animals having the normal powers of reaction, we can bring about this immunization with what, under the circumstances, is a marvellous precision. It is along these lines that Pasteur initiated the process of immunization against anthrax and other diseases, and upon these is based the now very considerable industry of antitoxin preparation. Here, again, is no matter of chance acquirement. Animals adapt themselves to, and combat, the toxins of disease according to very definite laws, the process varying somewhat, it is true, in connection with the different pathogenic microbes, nor is the animal body able with equal ease to gain immunity against each particular germ and its toxins. Against some, indeed, the immunity gained is either very

¹ It has been calculated that bacteria growing actively and under favorable conditions can divide, and so give rise to a new "generation" every fifteen minutes, and so afford close upon one hundred generations in the course of a single day.

ADAPTATION

feeble or is short-lived; but in any particular instance we realize that given amounts of toxin administered in a given way will in a given time result in the production of approximately the same grade of immunity in members of any one species of higher animal. And what is more, the adaptation is not merely temporary, existing only while the toxins are present and exerting their effects in the system. It is, in many cases, more or less permanent, so that in certain cases we see that it is in action for months, if not years. There must, that is, be impressed upon the cell substance an alteration in constitution which (remembering that the cells, as such, have most of them but a limited life period, becoming replaced by others of like nature) is conveyed from one cell generation to the other.

We shall, in our discussion upon immunity, adduce abundant instances affording proof of the statements here made and of the fact that the cell substance within certain limits can adapt itself adequately to alteration in the cell environment. And what is true of bacterial toxins is true also of not a few animal and vegetable poisons. For these, also, the system acquires a tolerance; or, expressed otherwise, while at first a certain quantity of each of these, absorbed and circulating in the blood and lymph, arrests cell activity either by breaking down the active cell substance, or by forming with it combinations which satisfy the biophoric molecules and so arrest metabolism, thus bringing about cell and systemic death; quantities less than the lethal act, and are reacted upon, in such a way by the cell substance, that this gains the property of dealing with quantities far in excess of what previously had been lethal.

Some of the most remarkable studies in this direction are those by Ehrlich and his pupils upon *abrin*, the active principle of the plant *Abrus precatorius*, and *ricin*, that of *Ricinus communis*, the castor oil plant. Both of these are intensely poisonous, are substances which, in the ordinary course of nature, are eminently unlikely to gain entrance into the systems of animals of the laboratory, and yet, with remarkable precision, those animals can, by repeated sublethal doses, be immunized so that they can stand doses several hundred times the ordinary lethal amount.

THE PHYSICAL BASIS OF ADAPTATION.

Along what lines can we explain this adaptation, so different from, or at least so far in advance of, anything we encounter in substances not endowed with life?

This we may safely say: that the capacity to adapt must be inherent in and depend upon the constitution of the molecules of living matter and upon the conditions under which those molecules carry out their ordinary activities. It is not so much that the tendency to vary is inherent, as that the labile nature of the biophores leads to their variation when subjected to modifications of environment; they vary according to circumstance, *i. e.*, according to law,

Let us try to reason out the simplest case first: that of the assumption

THE PHYSICAL BASIS OF ADAPTATION

by bacteria of new, or at least greatly exalted, powers of dissociating and fermenting unaccustomed foodstuffs-proteins or sugars.

We have already shown that substances in aqueous solution (and all the foodstuffs of the bacteria are assimilated in a state of aqueous solution) are liable to undergo ionization to a greater or less extent. We may, therefore, more than suspect that, either by direct ionization or by the secondary effect of free ions from other sources present in the cell sap, potential foodstuffs undergo dissociation—that the more complex bodies are broken down into others of a simpler type. This may well be a most important factor in the process we are discussing; the cytoplasmic molecules combining not with the molecules of the unaccustomed foodstuff as such, but with bodies of a simpler type, yielded by it, with bodies which are either ordinary constituents of the cytoplasmic and biophoric molecules, or which are so relatively simple that direct combination is possible between them and the molecules of living substance. An unaccustomed sugar, lactose, for example, may in this way be broken down and afford assimilable material to the bacterial cell.

This is, however, only one stage, the stage favored by the conditions under which the cell substance exists. It explains at most the assimilation of unusual foodstuffs, not the active adaptation to the same. For this latter we have to fall back upon the considerations already brought forward regarding the structure of the cytoplasmic and biophoric molecules-upon what, in brief, we may term the "side-chain theory" (p. 66). We are led, that is, to regard the molecules of living matter as a ring of subordinate radicals, each having numerous satisfiable affinities. If the environment remain unaltered, one constant series of "foodstuffs" diffuses or is absorbed into the cell; one regular order of dissociation products of the same is in solution in the cell sap, and the various affinities of the cytoplasmic and biophoric molecules are satisfied in one particular manner, associated with which growth proceeds. With a given environment, that is, these molecules build up side-chains which, having a particular composition, manifest particular properties.

But let the environment be altered; let a new potential foodstuff be introduced; through it and its dissociation products a new series of free ions is brought into the immediate sphere of action of the molecules of living matter. According to the strength of these ions, according also, it may be, to their number, these are attracted to the molecules of living matter and combined as side-chains, it may be replacing others in the process, others that on their part do not possess such strong affinities. If ions of a new type be thus taken up, new orders of sidechains will be developed and the molecular complex as a whole will acquire an altered composition—and altered properties.

At this stage we can figure to ourselves the central constituent rings as unaltered—merely the side-chains different. So long as the new foodstuff is presented, for so long will the cell molecules continue to form the new order of side-chains. And here let it be clearly under-

ADAPTATION '

stood we do not regard these side-chains as composed of the molecules of the foodstuffs combined in their entirety with the central cytoplasmic or biophoric molecules. The side-chains must, from every consideration, be regarded as tending toward the type of primary protein molecules. The new ions are built into them. If we regard the biophore as a polymeric molecule, and the simple protein molecules as of the same order, we cannot, as we have pointed out, regard growth as other than a process of development of new molecules by a process of accretion or building up of side-chains until these become united into new rings identical with the primary. And carrying out this idea, it is difficult to conceive side-chains in general as other than complete or partial polymerizations of the constituent nuclei of the biophoric molecule, Once a side-chain of a particular order is developed, it must, on its part, tend to polymerize, and if the radicals and ions identical with those that went to form it are present in the surrounding medium, a series or chain of like side-chain molecules will be developed within the cell.

Two, or it may be three, possibilities now present themselves:

1. The side-chain molecules may become detached in the cell sap or actually discharged into the surrounding medium, and being, as suggested, of the nature of primary protein molecules, may there present enzyme action. They may, in short, continue to dissociate the specific foodstuffs from which certain of their constituents were derived. As the whole molecule of cell substance was able to attract to itself certain of the constituents of that foodstuff, so, it may be, through side-chains formed, in the first place, from the products of disintegration of the foodstuffs, the cell now gains the power of acting directly on those foodstuffs. We shall encounter some very remarkable facts in our study of antitoxins, which can only be satisfactorily explained along the lines here laid down, namely, we have to assume that, in the first place, the cell gains its power to form antitoxins by combining with certain constituents of the toxins.

2. The second possibility is that these new side-chain molecules become utilized to form constituents of new cytoplasmic or biophoric rings—that they become utilized, in short, in growth. We conceive the biophore (p. 98) as being formed of a ring of primary (protein, amino-acid and nucleic acid) molecules, the constituent molecules not being necessarily identical in constitution. We can conceive the new side-chain molecules as replacing other molecules of simpler nature in the new biophoric rings that are in the process of being built up. If this should happen, then it is that we can regard the adaptation as not merely transient, but impressed upon the actual central living matter of the cell.

3. We mentioned above three possibilities; the third is the possibility that this combination takes place in three stages; that first, the constituents of the new foodstuff are incorporated in the side-chains; next, that they become constituents of the cytoplasmic molecules, and only in the third place become integral portions of the biophoric rings. This is likely to be the case if, as has been suggested, the biophores do not take up their specific constituents directly from the external medium, but

only from the cytoplasm, and through its intermediation. It is, indeed, possible to regard the cytoplasmic substance as of the nature of biophorie side-chain molecules.

The very fact that adaptation is in no case immediate, but requires some little period for its development, and this even in the simplest forms of life, favors this view of the existence of a succession of stages in its development.

But this is not all. Once the living matter of the cell becomes modified the modification is apt to persist, and apt, as we have said, to be carried on to later cell generations. A microbe that from the first moment of study has exhibited the power to ferment a given sugar, or that has acquired this power, is apt to retain that power if grown for a considerable period on a sugar-free medium. Under these conditions, no sugar being present, it cannot manifest this particular property, but, grown once more in the sugar-containing medium, it may immediately cause the fermentation. This must be said, that the power is apt to be weakened and not to show itself for a little time, and that the more recent the acquirement, the more rapidly is the power lost. Whether this last is a constant law we cannot say with absolute certainty. It is, however, a law of singularly wide application, this law that characters of more recent acquirement are those which are most easily lost, and its corollary that the older the character or property the more tenaciously is it retained. Specific properties are more firmly fixed than racial, racial than familial, and to this law we shall have frequently to refer. But, while admitting this, we are compelled to recognize that properties impressed upon the cell are retained for a longer or shorter period after the conditions which led to their acquirement have ceased to act. There is, as it were, a constitutional or truly a chemical inertia, and this is at the base of heredity.

We can only explain it by assuming that, whereas at first the modified constitution of the side-chains and primary molecules was due to the actual incorporation of dissociation products of the novel foodstuff, once these molecules become part and parcel of the biophores, these have the power to attract and combine not merely the already partly elaborated dissociation products of the foodstuff, but also simpler combinations of other origin, and to combine these in due proportions. We must admit that the biophores are capable of synthesizing (if the expression be permitted) the simplest hydroxyl ions, carbon compounds, etc., present in the cell sap, so that from them rings or primary molecules identical with the original continue to be produced.

In favor of this hypothesis, certain calculations of McFarland may here be quoted:¹

A horse may easily be so immunized against diphtheria that each cubic centimeter of its blood serum comes to contain 500 immunizing units of diphtheria antitoxin. Such a horse, it is calculated, has circulating sufficient blood to furnish 30 pounds—or 15,000 c.c.—of antitoxic serum, of which 1 c.c. will protect against, or neutralize, 225 c.c. of

¹ Text-book upon Pathogenic Bacteria, fourth edition, Philadelphia, 1903; 125.

ADAPTATION

the toxin. The amount of toxin injected to furnish such an immunity is 4200 c.c. As against this amount injected, the productive energy of the immunized horse is adequate to neutralize 3,375,000 c.c. of toxin; or, in other words, the blood drawn from his body is sufficient to protect 806 horses from doses of toxin as large as the total amount administered during the entire course of treatment, or against a very much greater amount than what, injected into an untreated horse would lead to its death (1 c.c. of strong diphtheria toxin administered to an untreated horse has, on more than one occasion, been followed by the death of the animal). It is obvious from these figures that the injection of a given amount of toxin leads to the development within the organism of not simply a corresponding, but a vastly increased, amount of antitoxin. What is more, if a treated animal be bled repeatedly, and the floating antitoxins be largely removed, the newly formed blood comes in a few days to contain amounts approaching those present previous to the bleeding.

Resume — Before proceeding farther, it will be well to sum up the successive stages in our argument:

1. All living matter exhibits obvious adaptation to the conditions under which it manifests its activity.

2. Specific and racial adaptation is best understood from a study of individual and cellular adaptation.

3. Study of individual and cellular adaptation demonstrates clearly that adaptation is a regulated process, and not the result of chance. Modify the conditions of life of one of the bacteria in certain particular directions, and, provided the modifications be not so severe as to arrest vital activities, the bacteria inevitably exhibit modifications in their properties, and these modifications are in direct relationship, or adaptation, to the particular alteration in environment.

 The study of immunity shows that what is true of the simplest unicellular organisms obtains also with individual cells in the highest animal forms.

5. Modifications in properties demand modification in the constitution of the cell substance; at base, therefore, adaptation indicates molecular alteration and rearrangement in the living matter of the cell. At base, therefore, we have to seek a chemical or physicochemical explanation for adaptation.

6. We find this according to the biophore theory, which regards the molecules of living matter as arranged as rings, and rings of rings, each ring being capable of attracting and affixing ions from the surrounding medium and building these up into side-chains.

 The rings of which the biophores are composed are, we hold, of proteid nature, and the tendency of protein molecules to undergo polymerization indicates that the side-chains are built up as polymers, *i. e.*, are also of proteid type.

8. What happens in adaptation, therefore, would seem to be this, that with modifications of environment new compounds are introduced into the cell sap; these undergo or have undergone dissociation into their constituent ions, and these new ions, either replacing other

ADAPTATION TO PHYSICAL CHANGES

groups of ions in the cell sap, or having greater affinities to the molecules of living matter, become fixed by those molecules and built up into side-chains. In this way we have the first alteration in the constitution of that living matter; they come to possess altered side-chains.

9. Such side-chains may (a) when complete become detached and free in the cell sap or be discharged into the surrounding medium, or (b) may become units in the building up of new cytoplasmic and biophoric (nuclear) molecules.

10. Once the living matter of the cell becomes modified to the extent that new biophores have been produced by reduplication, or, more exactly, growth, that modification is apt to persist and this long after the agent which caused the modification in the first place has ceased to act. The only valid explanation of these facts is that, while at first the specific dissociation products of the substance causing the modification were built into the side-chains and biophores, once these biophores or other molecules of living matter have assumed a particular constitution, they possess the power of attracting to themselves, and of building up into side-chains and new molecules, other and simpler ions in such proportion that from them they synthesize components of the side-chains and rings identical with the dissociation products of the substance which primarily brought about the modification.

Adaptation to Physical Alterations in Environment.-Thus far we have, for simplicity sake, taken into consideration only modifications in the cell produced by "foodstuffs." It will already have been determined by the reader that under this term is to be included everything capable of providing ions which can be seized upon by the living molecules and incorporated into side-chains or utilized for growth. The term is used as implying this idea, but it must be kept in mind that under it we include a large variety of substances-toxins and other poisons, for example-which ordinarily do not enter into our conception of "food." Our argument, in short, holds for all the adaptations in response to change of a chemical nature in the environment of the cell, with one possible exception, namely, that there may be substances absorbed or diffused into the cell which do not directly afford ions to be taken up by the molecules of living matter, but which break up matter already present in the cell, thus indirectly affording ions capable of utilization. This possible exception does not invalidate our main argument. It affords, indeed, a connecting link whereby to attach another series of phenomena, namely, the adaptations to physical, as distinguished from chemical, changes in environment. Changes in temperature, light, vibrations, do not introduce new ions into the cell from without; they tend, however, to modify the dissociation of the matter already within the cell, nuclear, cytoplasmic, and paraplasmic, and modifying the number and relative abundance of the different orders of free ions, they in a similar indirect manner must bring about change in the constitution of the biophoric molecular complex.

We possess, indeed, accurate observations upon the capacity of the lower forms of life to adapt themselves to temperature changes. The

ADAPTATION

earliest were those of Dallinger,¹ who by a very gradual increase in the temperature of the water in which they lived, extending over several months, accustomed infusoria, normally killed by a temperature of 25° C, to endure a temperature of 70° C. Davenport and Castle² have shown that tadpoles reared from the egg and kept at 15° C, for a month pass into heat rigor at a temperature of 40.3° C, whereas those reared at 24° to 25° C. do not manifest heat rigor until 43.5° C, is attained. Lastly, the experiments of Standfüss³ on the modification in the wing patterns and coloration in butterflies, caused by subjecting the eggs and caterpillars to different temperatures, have shown that forms so distinct as hitherto to have been regarded as different species are simply due to physical changes in environment, and that these diverse forms can be produced with exactitude.

From a general biological point of view these data regarding individual and cellular adaptation are of the very highest importance, and our conception of the means whereby it is brought about afford the necessary key to an understanding of variation, its origin and limitations, and through this to the process of evolution. We shall have to refer to these matters to some slight extent in later chapters. Here we would only lay stress upon the fact that cellular structure is the expression of the chemical constitution of the cells, that histological alteration presupposes modification in the arrangement and intimate constitution of the molecules of living matter, and lastly, that for the modifications to be more than merely transient the biophores or controlling molecules of living matter must have undergone alteration.

From a pathological point of view the data are of equal importance. We shall see that disease is two-sided. We have, on the one side, to regard the noxæ, or influences acting from without, setting up disturbances in cell activities; on the other side, the reactions on the part of the cells induced by such noxæ. And these reactions all come under the heading of adaptations to changed conditions. It is, perhaps, more correct to speak of these reactions as "tending to adapt," for time and again the adaptation is far from perfect. But in all the reactive processes we can recognize the existence and action of the same basal principles which are to be made out governing the microbe when its environment is altered-when a new sugar is introduced into its pabulum, and it proceeds to become modified, owing to the presence of that sugar and its dissociation products, with the result that the sugar becomes utilized as a foodstuff, and with this the microbe not merely accustoms itself to, but takes advantage of, the changed conditions. These considerations lead us to another possible definition of disease, *i. e.*, that "it is the expression of a reaction on the part of the cells to injurious agencies," just as the normal processes in the body are reactions to normal stimuli.

¹ Jour, Roy. Mier. Soc., 3: 1880; 1. ² Arch. f. Entwick, Mech., 2: 1895; 227.

^a Handb. der palüarktischen Grossschmetterlinge, 2 Aufl., Jena, 1896; Denkschr, d. Schweiz. Naturforsch. Gesellsch., 36: 1898, etc

CHAPTER X.

CELL AND TISSUE DIFFERENTIATION-INDIVIDUAL DEVELOPMENT.

MERELY to describe in outline the embryogeny of one of the higher vertebrates would demand more space than can here be afforded; we must take it for granted that the reader is familiar with the general details of the process. For our present purposes all that is necessary is to lay down that, by successive divisions and redivision, a single cellthe fertilized ovum-gives rise to all the cells which form the tissues and parts of the multicellular animal; that in the earlier periods of embryonic life the cells, the result of this division, show little sign of differentiation, but as development proceeds, differentiation becomes more and more marked in a larger and larger number of the cells, until at birth the separate organs, or almost all of them, are formed of constituent cells recognizably different from those of other organs even if the full differentiation of the same is not completed until some considerable time later. In other words, with progressive segmentation we pass gradually from the undifferentiated, or apparently undifferentiated, ovum to the most highly differentiated cells of the various tissues.

What we have now to consider is the means whereby this differentiation has been brought about, and this, again, not merely as an academic quest, but because in various states of disease we encounter extensive alterations in the characters and appearances of the cells of affected areas, and a knowledge of the laws governing the normal process of cell differentiation is essential for a comprehension of the abnormal processes. And here, at the outset, we would ask the reader for the time being to dismiss from his mind all thoughts of the modifications induced by the sexual fusion of the germ cells. These modifications are of a different order, and will be discussed in a subsequent chapter. The existence of parthenogenesis-of the development of individuals from non-fertilized ova-and the data gained from the abundant experiments on development initiated by physical and chemical means without spermatozoic fertilization, which we owe, in the first place, to Jacques Loeb,¹ prove that tissue differentiation is primarily independent of fertilization. For the present it will simplify matters to leave out of account the meaning and influence of this process.

Let us, in the first place, recall what we said (p. 36) regarding our

⁴ For a fuller study of these observations and of vital phenomena in general the reader may be recommended to Loeb's most interesting and suggestive lectures on the *Dynamics of Living Matter*, Columbia University Biological Series, New York, 1906.

conception of the multicellular individual, namely, that this is to be regarded not as a colony of individual unit cells, which have become and remained united for mutual benefit, but as a unit mass of living matter which, by increasing the surface presented to the external medium, has continued to remain a unit in spite of growth and increase in volume, and has preserved the due proportion between surface and mass through the agency of nuclear, followed by cell, division, the component cells in general being not wholly isolated, but remaining connected by cytoplasmic bridges.

1. In such a process, with continued nuclear division and distribution of the biophoric material into the constituent cells, inevitably that material is subjected to different influences. Just as in the freeswimming unicellular organs we note that a differentiation presents itself between the external and the internal cytoplasmic substance—

the former being directly acted upon by the surrounding medium and becoming modified into the denser ectoplasm—so, to take the simplest case that presents itself, in the even division of a spherical cell into a spherical cluster of cells, it must inevitably happen that those cells which are superficial are exposed to conditions distinct from the conditions acting upon the cells of the interior of the mass (Fig. 32). And, remembering what has been said in the preceding chapter regarding the capacity of living matter to adapt itself, it

is obvious that through adaptation the biophoric matter of the superficial cell layer will become modified, as compared with that of the deeper cell mass; and this modification in the constitution of the living cell substance will show itself in structural differentiation.

We gain, that is, our simplest and most natural explanation of cell differentiation by regarding it as *primarily* the result of adaptation to modified environment.

2. Accepting this as the primary cause of cell differentiation in the unicellular organism, it follows that, if there be two primordial cells possessing biophoric matter of identical constitution, and these each, under like conditions of environment, undergo conversion (growth and division) into a multicellular mass, then the component cells will undergo like differentiation. This, incidentally, is a basal law of heredity proper.

3. If, on the contrary, the composition of the biophoric material in two such primordial cells varies, then, although these be subjected to like environment, the cells resulting from their division will be effected diversely by that environment, and cell differentiation in the two resultant multicellular organisms, even if along the same lines, will nevertheless be distinct.

Granted the existence of living material after the order of biophores (as being at basis a chemical compound, however complex), and of adaptation, these must be our three primary postulates. And cell

EPIGENESIS AND PREFORMATION

differentiation in the multicellular organism is to be regarded as essentially the outcome of relative position in a complex of cells derived from one common biophoric material of particular constitution, subjected to the influence of a particular environment; that biophoric material becoming modified according to the influences brought to bear upon it in the different areas of the cell mass,

EPIGENESIS AND PREFORMATION.

This primarily. But a halt must be made. Are we justified in regarding the biophoric material of the ovum as "common," *i. e.*, as constituted of an aggregation of molecules of like order? There is the possibility that (even in the parthenogenetic ovum) the biophoric material is not homogeneous, but is composed of molecules of different orders, and that it is the mode of distribution of these diverse molecules that determines cell differentiation. Here, in short, we have to take sides in a controversy that has waged for close on a century and a half, now one party, now the other, appearing to gain the upper hand—the controversy between the upholders of *epigenesis* and *preformation*, respectively. Although with the progress of time and with fuller knowledge the field of battle has altered its position, the point at issue is essentially the same.

Before anything was known regarding the stages of development of the individual or of embryological histology, what may be termed the natural view held sway, and this was accepted by Aristotle and supported by Harvey as the result of his naked-eye studies of the developing hen's egg. The ovum in its earliest stage was seen to possess no internal structure that by the wildest imagination could be regarded as a minute edition of the future animal. No likeness could be made out between the germinal disk and primitive streak and the future chick. The natural view, therefore, was that the individual developed by the successive transformations of a germinal substance which originally was without form and without parts.

Only in the middle of the eighteenth century was this view called in question. Bonnet' recognized in the developing orum an unfolding or "evolution" of invisible small parts. These parts, he held, are present in the ovum from the first; are preformed. The ovum contains a "miniature model," as he unfortunately termed it, of the perfect animal—a model which he was careful to say is not exactly like the perfect animal, but consisted of "elementary parts" only. Bonnet had not observed the earlier stages of the chick. Caspar von Wolff² had, and saw clearly that the fertilized egg, as it proceeded to develop into the chick embryo, exhibited nothing that could be regarded as a "miniature model." He actively opposed Bonnet's doctrine of pre-

¹ Considerations sur les corps organisés, Amsterdam, 1762.

² Theoria Generationis, 1759.

formation.¹ The simple egg substance became modified under the action of its inherent formative power until, through continual new formations, and transformations more and more complex, the perfect animal was developed. And for a long period Wolff's "epigenesis" was the accepted doctrine, and this even after the development of studies of the cell showed that the ovum was by no means the simple substance which Wolff held it to be, and after the doctrine of vitalism rendered the conception of an inherent formative force unacceptable. For, on the whole, the successive discoveries of the embryologist favored Wolff's view. The morula and blastula stages of the embryo, the formation of the three germ layers, can only in an indirect way be dragged in to support the preformation theory.

Neither doctrine, as originally enunciated, is valid in the light of our present knowledge, but still the contest continues, and has, by Weismann,² been brought down to the biophores.

"Two fundamental assumptions," he states, "present themselves, and these can be related to every conception of germ plasm. . . . Either we may think of the id³ as made up of similar or of different kinds of parts, none of which has any constant relation to the parts of the perfect animal, or we think of it as composed of a mass of different parts, each of which bears a relation to a particular part of the perfect animal, and so, to some extent, represents its primary constituents. The assumption of a germ plasm composed of similar parts, which has been made, for instance, by Herbert Spencer, may be called the modern form of epigenesis, while the other assumption is the modern form of the (preformation) theory. The former theory can only explain development as induced by the influence of external conditions-temperature, air, water, gravity, position of parts-upon the chemical components of the germ plasm which are everywhere uniformly mingled, and it makes no difference whether this uniform germ plasm is thought of as composed of many different kinds of parts, so long as these parts are mingled uniformly to make a germ plasm and bear no relation to definite parts of the developing animal." We have quoted this in extenso because it states so accurately the conditions of the problem. Are we to regard the biophores present in the ovum of a given species as potentially of equal value, so that if in the process of cell division the biophore which finds itself in a nerve cell will have undergone those changes which convert it into a neuronic biophore; if, on the contrary, it has passed into a liver cell, the successive changes it has undergone in growth and multiplication have modified it into an hepatic biophore? Or, on the contrary, are we to suppose that the biophores present in the ovum are most varied in their constitution-that there preëxist in it biophores

¹We use this term in place of Bonnet's own "evolutio," so as to prevent a very possible confusion.

² The Evolution Theory. Translated (and that into clear and excellent English) by J. Arthur Thomson, London. Arnold, 1904, vol. 1, p. 350 et seq.

³ The unit mass of biophores, or, according to Weismann's terminology, of nuclear chromatin, capable of giving origin to the complete individual.

of the neuronic, hepatic, muscular, osseous, connective tissue, germ cell, and other types (the list could be lengthened prodigiously) which in the process of segmentation of the ovum are sorted out and distributed into the cells which form the *anlagen*, or basis of the different specific organs and tissue cells, and, entering these cells, control, or, more exactly, cause the differentiation of the same?

The point, it will be seen, is one of great importance, since our views not merely of tissue and cell differentiation, but of the broader subjects of evolution and heredity, materially depend upon which theory we accept. Weismann upholds strenuously the preformation theory, and as his views are widely quoted, it is necessary to inquire into his arguments.

Ontogeny (the development of the individual), he states, is not an isolated phenomenon, which can be interpreted without reference to the whole evolution of the living world, for it is most intimately associated with this, being, indeed, a piece of it. Ontogeng must be explained in harmony with phylogeny (the evolution of the race), and on the same principles. The assumption of a germ plasm without primary constituents, or of a completely homogeneous germ plasm, is irreconcilable with this, for it contradicts certain facts of inheritance and variation.

We take it that what Weismann means by this broad and rather vague pronouncement is that, to afford an example, if the lepidopterous insect, before attaining full development, has to pass through the caterpillar and chrysalis stages, this can only be explained by the preformation theory; that epigenesis is unable to explain the metamorphoses; that the effect of environment, merely, on the germ substance of the lepidopterous ovum would render the intermediate stages unnecessary, would cut them out, and would remove the manifold indications which individual development affords of the evolution of the race. We freely admit that, as a matter of fact, ontogeny affords most valuable indications as to phylogeny-that it is an abbreviated phylogeny, but how greatly abbreviated those who currently repeat this dictum are apt to slur over. The human embryo is at no period a pure worm, a perfect fish, a simple saurian; certain characteristic features only at certain stages are capable of explanation by the one theory alone-the theory that these features are reminiscences of the phylogeny. The retention of these features does not, however, demand the existence of determinants, *i. e.*, of biophores or groups of biophores of special constitution having the particular function of developing these particular features of special biophores which have descended unchanged from the annelid, fish, or saurian stage of existence; it can be explained more simply by the supposition that all the properties of the cells of the different tissues are the result of modifications of one common biophoric matter, these modifications being impressed upon that matter by the successive influences that have acted upon the cells in the course of development. From which it follows that we may regard it as essential that the cells which are ultimately to form certain organs shall have passed (or their progenitors shall have passed) through certain stages, in order that the

CELL AND TISSUE DIFFERENTIATION

contained biophores may undergo a particular line of modification. When the same result can be attained by a "short cut," this is done; whence it happens that the ontogeny does not by any means represent the full phylogeny. A very little knowledye of embryology furnishes abundant examples of these short cuts and of cases in which, in closely allied species, development is abbreviated by widely different "short cuts."

This argument, then, against epigenesis- if we understand Weismann's argument aright-is not unanswerable. His next appears, prima facie, to be more convincing. The existence, he urges, of a white lock of hair through several generations can only depend ultimately on a divergently constituted part of the germ plasm, which can only affect the one spot on the head and alter it, if it is itself different from what is usual. "On this account I call it the determinant of the relevant skin spot or hair group." In a germ plasm without primary constituents the variation could only depend on a uniform variation of all the parts, for the parts are either alike among themselves or, at any rate, have the same value for every part of the finished organism. How could an animal differing only in one minute part arise from a germ plasm which has varied in all its parts? There are five well-marked variations of the Indian species of butterfly, Kallima paralecta, in which the variation is in the markings on the under surface of the wing, while the upper surface is alike in all. How is this to be explained by the epigenetic theory? If each individual variation of the species depended on a variation of the whole germ plasm, the wood Kallima would soon bear no resemblance to its ancestral form, the meadow species. There must be primary constituents in the germ plasm, that is, vital units whose variation occasions the variation of definite parts of the organism, and of these alone.

As a consequence, Weismann has elaborated a scheme of inheritance in which the biophores (which he regards as supramolecular rather than molecular-as aggregates of molecules) are combined to form determinants or biophoric groups, each of which controls or determines the structure and function of one particular cell area of the body, and he assumes further that these determinants are combined into ids, each id containing the full complement of determinants necessary to give origin to the complete individual-numbers of these ids arranged serially are regarded as being present in the "idants" or "loops" of the wreath or aster of the nucleus of the ovum-the separate ids being conveyed to the ovum from different ancestors, and according to the ids which thus happen to pass into a particular ovum, so does one or other group of determinants derived from different ancestors come to gain control in the development of the individual. But of this more anon. We mention this here solely in order to give an idea of the relative size of these determinants as demanded by Weismann.

We could bring several arguments to bear against this chain of reasoning of Weismann's; could inquire, in the first place, whether Weismann is justified in assuming that where two varieties of a species exhibit to the naked ene only one single morphological point of differ-

THE PHYSICAL IMPOSSIBILITY OF WEISMANN'S THEORY 135

entiation, that is the only difference between them-whether more careful study would not demonstrate numerous concomitant variations not merely morphological, but functional also. We could quote the recent remarkable and extensive studies of Max Standfüss upon the experimental production of variation in butterflies, demonstrating that quite an extensive group of varieties which hitherto have been regarded as essentially due to difference in constitution of the germ substanceof the biophores of the germ cells-is due to the action of environment upon the germ substance, variation in the temperature to which the fertilized ova are subjected during the course of development sufficing to bring about an extraordinary variation in the coloration and marking of the eventual butterfly, a given temperature leading with striking constancy to a particular result. One single consideration, however, suffices to demolish the whole of Weismann's theorythe consideration, namely, that it is a physical impossibility that the id could contain all the requisite determinants; they could not be compressed into the space afforded, even were they atoms and not, as



Vanessa levana \Diamond to show influence of temperature during the larval period: A, winter form; B, summer form, which used to be considered a separate species (Vanessa prorsa). (After Grobben.)

he demands, collections of biophores, and these biophores not merely molecules of proteid nature and relatively great size, but collections of the same. We have already called attention to this reductio ad absurdum of Weismann's theory.1 Weismann2 freely admits, regarding determinants, that "in the higher multicellular organisms, as, for instance, in most arthropods, the number must be very high, reaching many thousands, if not hundreds of thousands, for in them almost everything in the body is specialized and must have varied through independent variations in the germ." And to make his image of these determinants quite clear, he adds: "In multicellular organisms I should be inclined to picture the determinants as a group of biophores which are bound together by internal forces to form a higher vital unity. This determinant must live as a whole, that is, assimilate, grow, and multiply by division, like every vital unit, and its biophores must be individually variable, so that the separate parts of a cell controlled by them may also be capable of transmissible variation.

¹ Adami, Inheritance and Disease, Osler's System of Medicine, vol. 1.

2 Loc. eit., p. 370,

CELL AND TISSUE DIFFERENTIATION

We employed previously Lord Kelvin's estimate of the size of a molecule of water, pointing out that, according to his figures, in the chromomeres or bead-like granules seen in certain chromosomes, which have been taken to represent Weismann's ids, there could be stretched across the diameter only about 150 molecules of water, and that when the highly complex molecules of the nucleoproteins have a molecular weight of not less than 15,000, the number of proteid molecules capable of occupying this diameter must be very much smaller. But in the opinion of many modern physicists this estimate of Lord Kelvin's represents, if anything, the maximum and not the minimum possible size of a molecule of water. Nernst,1 indeed, accepts van der Waals' calculation, based upon the molecular kinetic theory, that the magnitude of the molecule is one fivemillionth of a micromillimeter (0.000002μ) , or otherwise that along a line 0.5μ in length there could exist not 150, but 2500 molecules of water. Even taking this estimate, Weismann's conception is still outside the limits of the possible, when the huge size of the nucleoproteid molecule² is taken into account as compared with that of a simple molecule such as that of water.

If the biophores are, according to Weismann's conception, not simple molecules of proteid type, but aggregations of the same, the determinants composed of aggregations of biophores should be recognizable under the highest powers of the microscope, and the id formed in the higher animals, of thousands and tens of thousands of biophores, must inevitably be a body of from thirty to three hundred times the diameter of the determinant—so large, that is, that if it existed, it must have been recognized from the moment the nucleus of the cell was first observed—and if, as Weismann supposes, the nucleus of the ovum contains hundreds of ids derived from numerous ancestors, that nucleus would fill the whole field of the microscope! Needless to say, this is not the case; nor, we may add, does the coarseness of the nuclear structure vary materially according to the complexity of the animal. Physically, therefore, Weismann's conception is an impossibility, and, as Weismann has carried this conception of preformation to its logical outcome, it follows that, in demonstrating the impossibility of his theory, we simultaneously destroy all less fully developed theories of preformation.

Determinants, in Weismann's sense, cannot exist, and we must accept (with reservations, to be noted when we come to discuss the fertilized ovum) the alternative theory of epigenesis—the view that there exists primarily a single biophorie substance which in its growth and distribution to the various cells of the developing animal is subjected to varying influences whereby it becomes modified, and whereby the cells governed by it come to assume diverse functions and diverse structure. There is a preformation, but of the common biophoric substance alone; this must differ in the different species. And there

¹ Theoretical Chemistry, translated by Palmer, pp. 348 et seq.

[&]quot;As demonstrated inter alia by its non-filterable character, through a porcelain filter.

is an evolution, or unfolding, but the nature of this unfolding is of this order, that, given growth and cell division, the biophoric material subjected to a particular environment inevitably undergoes a definite series of transformations; and the different orders of cells, tissue, and organs are the result of the diversity of influences acting upon the common biophoric material of the ovum.

So far, let it be remembered, we have studiously kept out of consideration the facts of fertilization. It has seemed to us that we could make our statement of first principles clearer by neglecting them for the time being. Now we have reached the point at which they can no longer be neglected, for, obviously, in the gamogenetic individual the individual resulting from the union of the male and female germ cells—there is not a common biophoric substance; in them, at the moment of fertilization, at least, there are biophores of two orders, and it may be of many more, for, the parents being unlike, the biophores which controlled their growth must presumably have been unlike, and the same is true of all the ancestry. How, then, can we combine the conception of epigenesis from a common biophoric material (minus determinants, in Weismann's sense) with this necessary existence in the fertilized ovum of biophores of different constitutions? This we shall discuss in the next chapter.

THE MOSAIC THEORY.

But before leaving this portion of our subject there is a somewhat weighty objection to the theory of epigenesis which cannot be passed over in silence. The more carefully we study the earliest stages of segmentation of the ovum in the various forms of life the more clearly we recognize that, after the first or second division, the blastomeres or resultant segmentation cells begin to show signs of differentiation. In other words, cells apparently subjected to identical environment exhibit structural differentiation.

This point was emphasized strongly by Weismann in some of his earlier writings, and has been more particularly studied (1888) by Roux, of Breslau. The ovum in its earliest stages segments first along one median plane into two cells (or blastomeres), then each of these subdivides along a plane at right angles to the former, a four-cell stage being produced, and subsequently an eight, sixteen, thirty-two cell stage, etc. In certain most interesting observations upon the germinating frog's eggs, Roux' showed that by destroying one or other of the blastomeres in their earliest stages he could produce monsters of defect, one or other region of the body being undeveloped, according to the cell destroyed. If, in the four-cell stage, for example, one of the blastomeres be destroyed by means of a heated needle, a frog may develop wanting one entire quarter of the body. The conclusion appears obvious that in the

¹Virchow's Archiv, 114: 1888: 113; see also Anatom. Hefte, February, 1893.

CELL AND TISSUE DIFFERENTIATION

segmentation of the ovum, with the first division the determinants from one-half of the body pass into one of the primary blastomeres, those for the other half into the other; and that when these two divide, the determinants for the front half of the right side pass into the right anterior blastomere, for the hinder half of the left side into the left posterior blastomere, and so on. And more particularly from these observations he developed what has been termed the "mosaic theory" of development; that "the development of the frog gastrula and of the embryo formed from it is, from the second cleavage onward, a mosaic work, consisting of at least four vertical independently developing pieces;" organization, that is, precedes cell formation.

But in the course of these observations Roux himself was the first to note that, where he destroyed one of the cells in the two-cell stage, instead of gaining a half embryo (unilateral), he might gain a whole, though dwarfed, individual; and later, Driesch⁴ conducted a most suggestive series of experiments. Taking the eggs of the sea urchin in the two- and four-cell stages, he was able, by shaking, to separate the cells, each of which gave rise not to half and quarter embryos, but to entire, though dwarfed, larval forms. E. B. Wilson² obtained even more striking results with amphioxus eggs, while, not to mention several other confirmatory observations, Zoja,³ in certain jelly-fish (meduse), obtained perfect embryos, though correspondingly dwarfed, from the separated blastomeres of the sixteen-cell stage.

We shall have later to point out how these observations throw light upon the development of certain twins and double monsters. What we have to indicate here is that they absolutely contradict the mosaic theory. They show that in the earliest stages, and the same, we may presume, is the case in the later stages, the division of the cell—the ovum—and its nucleus is into similar parts. The daughter chromosomes are of equal value qualitatively and quantitatively.

But how are these facts to be reconciled with the opposed facts of Roux? This has been solved by Morgan.⁴ He has shown that in the frog's egg, if, after the destruction of one blastomere, the other be allowed to remain in its normal position, a half embryo develops, conformable with Roux's observations; if, on the other hand, following the action of gravity, it becomes inverted, it most frequently gives rise to a whole dwarf, although in some of his experiments, even under these conditions, the half embryo developed. Wilson has obtained similar results with amphioxus eggs. Through these and allied observations it has been determined that the different components of the ovum assume naturally particular relations, the one to the other. This is largely a mechanical matter. Thus, in the frog's egg the stored food material, yolk or deutoplasm, is heavier, and sinks, while the lighter nucleus and cytoplasm rise, and so far, it would seem, from purely mechanical causes there is developed a *polarity* in the orum. Similarly

¹ Zeitschr, f. wissensch. Zoologie, 53: 1892. ² Jour. of Morphology, 8: 1893.

³ Arch. f. Entwickelungsmechanik, 1 and 2:1895.

⁴ Anat. Anzeiger, 10:1895:623.

INFLUENCE OF CYTOPLASM ON NUCLEUS

the pigment in the frog's egg collects, or is developed, at the upper pole —the part exposed to the greater amount of light. Thus, we have indications that in the very earliest stages the fertilized egg obtains polarity; or, otherwise, that the different constituents—nucleus, cytoplasm, and paraplasm (deutoplasm)—take on a definite arrangement, which in itself determines to a large extent the subsequent course of cell division; if this arrangement be disturbed, then that subsequent course is liable to alteration. We can, that is, given these data, harmonize apparently contradictory facts, and, what is more, can from them gain an understanding of how it may come to pass that without determinants there may be potential cell differentiation in the very earliest stages of the segmenting eggs. Briefly, while the nuclear biophores are to be regarded as the controlling agents in the cell, their activity is determined by the surrounding cytoplasm and deutoplasm, and the relations of these three again are determined by physical agencies.

Observations of the foregoing order are brought forward by certain writers in support of this view that heredity does not reside in the nucleus solely, and that the cytoplasm exercises a pronounced determinative influence upon the future of the individual. Of this influence a most striking example has of late been adduced by Godlewski¹ in the course of an active controversy upon the respective influences of the paternal and maternal germ plasms upon the development of hybrid larvae. It is found in the first place that hybridization is possible between the most widely separated echinoderms-starfishes, stone lilies or crinoids, and sea urchins. Godlewski found that if he enucleated the egg of a sea urchin (Strongylocentrotus) and fertilized it with the sperm of a crinoid (Antedon) the resultant larva had sea urchin and not crinoid characters. There could be no more telling example of the influence of this cytoplasm upon development. But if the general conclusions we have reached regarding the relationship between cytoplasm and nucleoplasm be kept in mind, the case is seen to afford an exquisite example of adaptation. The nucleus for its growth depends upon what it can assimilate from the cytoplasm; the cytoplasm being of sea urchin type assimilates matter from the exterior which it builds up into "side-chains" of sea urchin type; it is this matter that has to be assimilated and built up by the nucleus whose new biophores inevitably take on to a very large extent sea urchin characters; or otherwise we have here an exquisite example of the means whereby the newly formed biophores are modified in accordance with their environment.

This controversy is still raging regarding the "mosaic theory," or, more definitely, regarding prearrangement of blastomeric constituents prior to segmentation and the meaning of the same. The reader will find a fuller discussion in Professor Wilson's work on *The Cell*. It will be seen that our conclusions very largely agree with his. Here we must call attention to the fact that there are certain fertilized eggs whose separated blastomeres cannot be brought to form complete

¹ Arch, f. Entwickelungsmechanik, 1906.

CELL AND TISSUE DIFFERENTIATION

dwarf individuals, but always-under the conditions of experimentdevelop into partial larva. Nevertheless, these cases cannot be adducd in favor of determinants. In the individual blastomere of the four-cell stage of the embryo of the gasteropod Ilyanassa, for example, there is present material which in the normal course of events would give origin to germ cells capable of developing into the complete individual, and yet such individual blastomere if separated from the rest is never found itself to originate a complete, if dwarfed, larva, but only a quarter larva. The matter capable of developing the whole individual is present; there must, however, be some arrangement, some mutual relationship of biophores, cytoplasm, and paraplasm which inhibits the full develop-Modifying Driesch's' conclusions, we may say: The relative ment. position of a blastomere in the whole agglomeration of blastomeres, coupled with the relation of the parts in that blastomere, determine in general what develops from it; if these relationships be changed, it gives rise to something different; to this extent "the prospective value of the blastomere is a function of position "-acting upon biophoric constitution.

This power of single cells to produce the entire body is, in general, limited to the earliest cleavage products, with the one prominent exception of the germinal blastomeres-cells that can be distinguished or followed back to a very early period in the embryo-which are destined to give rise to the germ cells. In certain of the lower multicellular animals there are indications that the body cells in general retain this property, as again in certain plants—the trite examples are the Hudra and the *Begonia*; but even in these it is at least questionable whether a single cell has this capacity. In these cases we are unable by experiment to isolate a single cell, and when the removed portion is below a certain size no results ensue. It is therefore probable that for the reproduction of the whole individual from the body cells there must be present representatives of the different germ lavers-a collection of cells rather than a single cell. In the higher animals, at least, a distinction between germ cells and somatic cells is very marked, and it may be laid down as a general principle that the more pronounced the differentiation of a cell, the less its capacity, not merely to reproduce the individual, but also to reproduce itself. In these higher animals, judging from the data regarding homologous twins and multiple births, and more particularly from Spemann's experiments upon the eggs of the newt, not beyond the gastrula period are we able to divide the embryo so that each half gives rise to the whole individual. Following upon this, with the development of the primary germ layers the constitution of the biophoric material has already become so modified that epiblast cells give rise to epiblastic structures only, hypoblast to hypoblastic. The apparent exceptions to this law we shall discuss in the chapter on Metaplasia. It is the existence of this law that permits us to classify the new growths or tumors. (See chapters on Neoplasia.) And when we study the fully developed tissues we find that the cells which are the most highly differentiated of all.

¹ Studien, IV, Zeitsch. f. wissensch. Zoologie, 55:1893:39.

CELL DIFFERENTIATION AND PROLIFERATIVE CAPACITY 141

namely, the neurons, or nerve cells proper, have completely lost the power of reproduction; once fully formed, they cannot multiply. Other welldifferentiated cells-muscle cells, gland cells, and even squamous epithelium-have retained the power of reproduction, but gland cells can only give origin to gland cells, muscle cells to muscle cells, epithelium to epithelium, and when fully developed they can only multiply after undergoing a preliminary "Entdifferenzierung," or undifferentiation, reverting to a simpler, less differentiated stage. The developed muscle cell, before it can multiply, loses its striation, reverts to a more embryonic type, its nuclei multiply, and each becomes surrounded by apparently undifferentiated protoplasm; the gland cells, to a large extent, lose their specific granules and paraplasmic matter, the cell body swells and stains poorly; the squamous epithelial cell becomes swollen and more rounded, its nucleus more prominent, its cogwheel-like processes unrecognizable. We shall describe these changes more fully when treating of the subject of tissue regeneration. The more fully we study the differentiated tissues of the body the more it is brought home to us that the fully developed and differentiated cell, as such, exhibits little active multiplication, and that to a very large extent under normal conditions the renewal of cells worn out by use is brought about by the presence and reproductive activity of "mother cells," of cells, that is, present in the tissue in a relatively undifferentiated form, or, as we are accustomed to term it, of embryonic type.

For instance, where the skin has been irritated, we find that even well out in the stratum corneum certain cells are swollen as above indicated, and show stages of mitosis. The normal skin does not present evidences of multiplication in these regions; in it the constant loss of surface cells is made up by the mother cells forming the deepest. Malpighian layer of the epidermis. This, as every student knows, is a palisade layer of small, simple cells with deep-staining nuclei. These are present throughout life; they never become converted into squamous cells; they exhibit mitotic figures and multiply, and it is their daughter cells given off toward the exterior which, as they pass farther and farther away from the nutrient basis, undergo successive modifications, until they become completely keratinized. There are similar mother cells for cartilage (perichondrium), bone (periosteum, osteoblasts), mucous membranes, and their grand follicles, lymph nodes, etc. In voluntary striated muscle it is probable that the so-called muscle spindles have a like function; in the heart muscle, as pointed out by MacCallum, there exists a layer immediately beneath the endocardium of this mothercell type;¹ in the brain and nerve centres we have indications that, with destruction of the neurons, certain of the less differentiated neuroglia cells can in early life undergo differentiation and development into neurons.

¹ The remarkable observations of Tawara in Aschoff's laboratory upon the "Reizleitung System" (Jena, Fischer, 1906) render it urgent that these observations be repeated to determine the distinction between such cells and those connected with the bundles of His and the conducting system of the heart. It is, in short, only the lowest and simplest of tissues that can impartially either perform function or multiply, and even here we note that *Entdifferenzierung* precedes multiplication. The simple connectivetissue cell, with attenuated nucleus and scarce visible cell body, swells prior to multiplication, its nucleus becomes spindle-shaped and deep staining, it gains a recognizable cytoplasmic body, it becomes identical with the spindle cell of developing connective tissue.

How are we to explain these facts in the terms of the biophore concept already laid down? Our conception of the biophores, it will be remembered, is that these primary molecules of living matter have, within relatively wide limits, the capacity of adaptation, i. e., with modified environment undergo structural modification. These facts indicate that, with active growth of the fertilized ovum and coincident multiplication of the biophores, the modifications undergone may be so inconsiderable that the first cleavage products possess biophores which retain all the properties possessed by the biophores of the ovum, and so, like the ovum, are capable of giving rise-if the blastomeres be separated-to complete individuals; if they be not separated, then, through the interaction of the cells and the polarity of the cell mass, to portions only of the simple individual. Rapidly as the cells multiply and the cell mass grows in extent, the biophores contained in each cell become modified. According to their environment, so do the foodstuffs assimilated vary, and the groups of jons seized upon and attached to the biophores exhibit variation, with resultant modification in the constitution of the new biophores, until these become so specialized that they give rise to biophores capable only of determining the characters of special orders of cells, incapable of giving origin to all the orders of cells present in the organism. And, as this process continues, eventually the elaboration of the biophores in adaptation to particular relationships and particular function becomes so extreme, their constitution so elaborate, that the capacity to deduplicate, *i*, *e*, to multiply, is lost: the cells containing these elaborated biophores cannot multiply; or it may be more correct to say that the biophores still possess the power of multiplication, but this is inhibited by the extreme differentiation of the cytoplasm.

Reverting to what we have written regarding cell energy (p. 101), it will be realized that the immediate reaction to external stimuli and the performance of function is exerted through the cytoplasm; that this is to be regarded as intermediary between the biophores and the external medium; that the performance of function demands discharge of energy on the part of the cell, while, contrariwise, growth demands storage of energy. From these considerations it follows that the more pronounced the differentiation of the cytoplasm the more is the cell prepared to expend the energy acquired from assimilated foodstuffs in the performance of function rather in growth. In other words, the greater the cytoplasmic differentiation the less the capacity of the biophores to initiate growth and cell multiplication. That this is so is strongly supported by the phenomenon of Entdifferenzierung above

described; or, concisely, the change from the functional to the vegetative type of cell is accompanied by a loss and using up of the cytoplasmic structures elaborated for the due performance of function in response to external stimuli.


We cannot sufficiently emphasize this antagonism between the katabiotic and bioplastic activities of the cell. It may not be absolute; within certain narrow limits, as already indicated (p. 103), the two must surely co-exist; but these limits appear soon to be overpast, and the more the cell prepares itself for the performance of special function the less becomes its vegetative activity.

In our study of tumors it will be seen how important is the bearing of these considerations upon our grasp of the essential nature of neoplasia.

CHAPTER XI.

FERTILIZATION.

Two facts in themselves indicate that sexual conjugation and fertilization, the result of that conjugation, essential as they have become for the continuation of the bulk of living species, are nevertheless of secondary import, or at least not primordial; the facts, namely, that growth, adaptation, and cell differentiation can proceed in animals



Development of the colonial flagellated Infusorian Pandorina morum, to show conjugation of like sexual cells: I_i ordinary colony formed of sixteen like cells (persistence of morula stage of embryo of higher forms); II_i , similar colony in which each cell has developed into a daughter colony of sixteen cells; III_i , colony like I_i in which the cells are escaping from the gelatinous envelope for purposes of conjugation; IV_i , V_i , conjugation of two like individuals cell; VI to X_i , subsequent stages leading up to development of a cell mass as in I_i . (After Pringsheim).)

developed parthenogenetically, and that in the lowest forms of life long successions of generations have been followed without signs of conjugation being detected; so that we may with security state that in these sexual conjugation does not occur. For these reasons, difficult as at times it has been, we have, to this point, studiously refrained from considering these processes and their results. Their study introduces a new

and complex order of phenomena, which best is taken into account after everything not directly due to sexual differentiation has been passed in review.

Here we shall not discuss the significance of that differentiation, nor the meaning of fertilization. We will provisionally accept the light afforded by Maupas' observations' upon long series of parthenogenetic generations of infusorian *Stylonychia pustulata*, confirmed as they have been by Calkins' recent most painstaking studies upon long generations of *Paramacium*, that fertilization is essentially a means of biophoric rejuvenation. Indeed, our treatment of the whole of what has now become a very



Egg of an Echinoderm with surrounding spermatozoa to show differentiation in size of invasive male, and yolk containing female germ cells. (Korschelt and Heider.)

considerable branch of biological research must be brief and eclectic. We can but select those data and general conclusions which lead us forward toward, and supply us with, a foundation for the study of heredity.


Thus, in passing, we may note that the simplest type of conjugation found among unicellular forms of life is that of fusion of two wholly similar individuals; that among the multicellular forms, whether of animals or plants, we find similarly, low down in the scale, that little differentiation is to be made out between the male and female germ cells (Fig. 34). Very soon this differentiation shows itself, so that the one cell —the male or invasive element—becomes motile, to the end that, being attracted, it may actively move toward and penetrate the more passive female element—passive, because it contains in its cytoplasmic meshes a store of foodstuff or yolk, necessary for the active growth which follows fertilization. Of such store material the male element, or spermatozoon, shows the veriest trace; it comes to consist of little beyond nucleus, centrosome, and actively motile tail or flagellum. The disproportion in size of the two elements involved in the act of fertilization becomes thus singularly great (Fig. 35).

From a very early period of development of the individual the germ cells, destined to give rise to either ova or spermatozoa, are marked off from the somatic cells, destined to give rise to the tissues of the body in general. In certain insects they have been traced back and recognized at the blastula stage; in the nematode worm, Ascaris, Boveri

¹ Archives de Zoologie, second series, 7:1889.

FERTILIZATION

has succeeded in tracing the differentiation back to the results of the first segmentation of the ovum—to the two-cell stage. Not to enter into details of modes of differentiation of the two orders of cells, which vary considerably in different forms of life, we may, with very slight alteration in wording, follow Professor Wilson, and lay down that the difference between the germ and the somatic cells is, that the former retain the sum total of egg chromatin elements handed down to them from the parents, whereas, by one or other process, the somatic cells



Two cells of segmenting egg of Ascaris megalocephala, A, destined to give rise to body cells, shows diminution and casting out of some of its chromatin. B, the germinal blastomere, shows no such reduction. (After Boveri.)


retain only a portion of the same. Following back the descent of cells destined to be germ cells, we find that the series is uniformly rich in chromatin—that there is no primary casting out or reduction; in somatic cells preliminary reduction does occur. "The original nuclear constitution of the fertilized cell is transmitted, as by a law of primogeniture, only to one daughter cell, and by this again to one, and so on, while in the other daughter cells the chromatin in part degenerates, in part is transformed, so that all the descendants of these side branches receive small reduced nuclei."¹ In conformity with what we have already stated regarding the nature of biophoric material, we would suggest that the somatic blastomeres receive, both a reduced amount of chromatin, and chromatin of modified constitution.

As already hinted, it is still an open question how far the nuclear chromatin is to be regarded as identical with biophoric matter; from the fact that chromatin (stainable material) may disappear from view entirely at certain stages of cell activity in certain of the lower forms, there are those who regard this not as the active living substance, but as a first product of the activity of the same. On the other hand, this loss of staining power has received what is at least a plausible explanation. The more acid the character of the chromatin the greater its affinity for basic dyes, such as hematoxylin. The greater the proportion of

¹ Boveri: Merkel and Bonnet's Ergebnisse, 1: 1891: 437.

GERM CELLS AND BODY CELLS

unsatisfied nucleic acid in the nucleus the more intense its staining properties. The intense stain taken on by nuclei prior to mitosis indicates thus a heaping up of bodies of the nature of nucleic acid; if later, as the nucleus breaks up during the process of mitosis the staining power diminishes greatly, this suggests that the nucleic acid becomes combined with (possibly) proteid matter to form nucleoproteids, the compound losing its strongly acid properties.¹ We are agreed that there is the closest possible relationship between the nuclear biophores and chromatin, and that a permanent reduction in the amount of the latter is the expression of a reduction in the amount of the former. Fig. 37 expresses graphically this relationship in descent of the germ cells to the rest of the organism.



Schema of germ and somatic cell differentiation. (After Klebs.)

We have pointed out that the primordial germ cells differ from the somatic cells in that they undergo no primary reduction in their chromatin. We have now to point out that a most remarkable feature of the adult germ cells, the immediate precursors of the ova and spermatozoa—the oocytes and spermatocytes, as we may term them—is that in the process of maturation they exhibit a *terminal* reduction.

If we study the mitotic figures in the growing tissues of multicellular individuals, we discover that, with one exception, which we shall have to deal with in studying tumor formation, these exhibit in the aster stage a number of chromosomes or loops of chromatin which is constant for the particular species, and is always even, always a multiple of two. Thus, it is 2 in one variety of *Ascaris*, 4 in certain worms, 18 in the sea

¹ It may, indeed, happen that at the period of mitosis when the nuclear membrane disappears, the nuclear matter, coming into more immediate contact with the cytoplasm, is peculiarly susceptible to modifications, from which at other periods of its existence it is, as we have pointed out (p. 45), largely protected.

FERTILIZATION

urchin, 32 in man, 168 in the phyllopod Artemia. That it is a multiple of two is due to the fact that one-half of the chromosomes are of paternal, one-half of maternal, origin. Nay, more, in not a few species the individual chromosomes vary in shape and this to such an extent that different types are clearly recognizable present in pairs. The convincing demonstration of this distribution has been afforded by Moenkhaus by crossing two fishes, *Fundulus* and *Menidia*. These happen to have the same number of chromosomes (36), but those of the former are three times as large as those of the latter. In the developing hybrid all the mitoses studied in the body cells exhibited half the chromosomes of the large (Fundulus) type, the other 18 of the small (Menidia) type.

In all cases so far studied the mature ovum and the spermatozoon contain or exhibit just one-half the number characteristic of the somatic cells of the particular species.¹ The mature ovum and the spermatozoon receive and contain half the number of chromosomes characteristic of the previous generations of germ cells and the somatic cells as a body.

The method of reduction varies in the two sexes, varies also to some extent in different species. The following account gives the stages common in all multicellular organisms, whether animal or plant, omitting details.

SPERMATOGENESIS: THE MATURATION OF THE SPERMATOZOON.

The primordial male germ cells give origin to *spermatogonia*, cells which divide and redivide, with the ordinary number of chromosomes, until, with adolescence, these cease dividing, attain a considerable size, and become known as *primary spermatocytes*. Each divides into two, giving rise to *secondary spermatocytes*; each of these again into two *spermatozoa*. Thus, each primary spermatocyte produces four spermatozoa.


Following now the chromosomes, this may be stated: In the ordinary cell, as has been indicated on p. 115, each chromosome splits longitudinally into two, and one of each pair thus formed passes into each daughter cell, which thus comes to possess the same number of chromosomes as did the mother cell, and as half the chromosomes of the mother cell are of paternal, half of maternal, origin, so does the daughter cell come to have also the like proportion of elements from both parents. Thus, if the parent cell in the monaster stage exhibits 8 chromosomes, each aster in the diaster stage has 8 members. We may, following Professor E. B. Wilson,² term these A, B, C, D, and a, b, c, d. But

¹ Here for the moment we leave out of consideration the "accessory chromosome" that has been determined in many species. To it and its significance we shall revert later.

² Harvey Lectures, 1906-1907: 212.

THE MATURATION OF THE SPERMATOZOON

now in the maturation process the primary spermatocyte exhibits a modification of this process, a modification known as *sunapsis*, the



The stages of Spermatogenesis in Ascuris megalocephala bivalens: I, spermatogonium, with four chromosomes (the normal number for the cells of this species); II, primary spermatocyte with two tetrads; III, primary spermatocyte undergoing division into two secondary spermatocytes, each with two dynds; IV, secondary spermatocyte; V, secondary spermatocyte undergoing division into two spermatozon, each with two monds; V, two young spermatozon, each bearing two monds—chromosomes (After A, Brauer.)

The stages of reduction in the Ovum and formation of polar bodies. Diagram based upon Boveri's observations on the maturation of the ovum of *Ascaris megalocephala biading*. I, and tranee of spermatozoon, sp., into ovum; 2, formation of two tetrads in place of four chromosome; 3, first division—formation of two pairs of dyads; 4, expulsion of first polar body (1 *P*, b), containing two dyads; 5, second division of nucleus of first polar body (1 *P*, b), containing two dyads; 5, second division of nucleus of ovum, and division of nucleus of first polar body —formation in each of two pairs of monads; 6, expulsion of second polar body and division of first polar body—ovum and each polar body provided with two monads. The arrows, pointing to Dv_c , indicate subsequent enlargement of the nucleus of ovum and conversion of the two monads into a chromatin network similar to that developed in Sp., the nucleus of the spermatozon.

FERTILIZATION

significance of which was first pointed out by the American cytologist Continuing to consider the case of 8 chromosomes, we Montgomery. find now that these undergo a coupling process, with apparent reduction; in place of 8 chromosomes there appear 4 "bivalents"-Aa, Bb, Cc, Dd (in Fig. 38, II, with 4 chromosomes, these are represented by tetrads), and in the subsequent mitosis to form the secondary spermatocyte, each group of bivalents divides so as to supply the complete series to each daughter cell. It will be seen that through the coupling process the number of elements is not actually, but only apparently, halved. This is followed by a true reduction process (the *meiosis* of Moore) in which the synapsis or fusion of the paternal and maternal elements is dissolved, the one portion, A, for example, passing into the one spermatozoon; the other, a, into another. It is still undetermined whether the complete paternal set-A, B, C, D-remains associated and passed over to the one spermatozoon, or whether there is distributed an admixture, e. g., A, b, c, D. The general opinion is in favor of the latter supposition. The result, however, is that the spermatozoon is constituted with half the number of chromosomes peculiar to the ordinary cells of the individual and of the species from whom it has been derived.

OOGENESIS: THE MATURATION OF THE OVUM.

A process, strictly parallel in intent, occurs in the maturation of the ovum, although so different in appearance are the stages that it was some little time before observers realized that they had to deal with like phenomena. We owe more particularly to Oscar Hertwig the demonstration of the identity of the two processes.

Like the spermatozoa, the ova are descended from the primordial germ cells, whose descendants in the ovary are termed oogonia, each germ cell dividing with the usual number of chromosomes, in general until the final stage. The differences now are these: that whereas, all the four cells derived from the primary spermatocyte become active spermatozoa, only one of the four cells descended from the corresponding oocvte becomes functional. The other three are degenerate and cast out as *polar bodies*—minute cells, with nucleus and cytoplasm, apparently functionless. Secondly, the period at which the process occurs is different; in general, the formation of these polar bodies either does not begin, or is not completed, until after the spermatozoon has entered the ovum (oocyte), and thirdly, and most perplexing of all, the process is conducted in such a way that all the stages happen within the body of the one cell, the oocyte. The polar bodies are intracellular formations. What was the oocyte undergoes nuclear changes and eventual reduction in its chromosomes and emerges as the mature ovum.

There is, however, and this prior to the formation of the first polar body, the same process of coupling or synapsis, whereby the number of chromosomes is *apparently* halved, although truly the full series of chromosomes is contributed, as dyads, to the first polar body, and to the




 Entry of spermatozoon into ovum.
 Loss of tail of spermatozoon; its mid-piece becomes the centrosome of the fertilized cell, 3. Division of centrosome. 4 and 5. Chromatin both of ovum and spermatozoon converted into a network; the two moleties gain approximately equal size.
 Chromatin of both becomes arranged into chromosomes (one-half of the number of each variety that is usual in the body cells of the species).
 Formation of spindle; division of chromosomes; partition of chromosomes derived from the two parents in equal number between the two future cells (blastomeres).



THE MATURATION OF THE GERM CELLS

nucleus of the oocyte. This first polar body now undergoes mitosis, its daughter nuclei receiving each one-half the normal number of chromosomes, and coincidently the nucleus of the oocyte undergoes a reducing mitosis, the third polar body and the resulting nucleus of the ovum being each provided similarly with one of each pair of chromosomes. The "monads" thus contributed to the nucleus of the ovum develop into typical chromosomes.



Schema of comparative descent of ovum and spermatozoa. The dotted lines indicate successive cell generation, the continuous lines connect successive stages of one cell. (Modified from Boveri.)

Conclusions.—Even leaving out the details (some of which may be of great significance), this is an extraordinary history. It is, nevertheless, one that has been confirmed by a large number of independent observers in connection with a great number of species both of animals and plants. What is the meaning of every step we have honestly to confess that we do not know. But certain conclusions are almost self-evident:

1. We are forced to see that what is in its essence the same process occurs in the maturation of both the male and the female elements. From the fact that the process obtains in all multicellular organisms examined, whether animal or plant, it is clearly of fundamental importance.

2. A feature that immediately arrests the attention is that in this process ovum and spermatozoon contribute to the fertilized ovum each one-half the number of chromosomes common to the cells of the particular species; so that the fertilized ovum, that is, the new individual, begins life with the normal number of chromosomes instead of double the number, which would be the case did spermatocyte and occyte give rise to sperm and ovum by the usual methods of mitosis.

3. What is the significance of the varying number of chromosomes in different forms of life we have no idea. We only know that in the same species varieties may exist, one of which has normally double the number of chromosomes possessed by the other.

FERTILIZATION

4. There are two facts which possibly throw light upon the need for reduction prior to fertilization. One of these we have already noted—namely, that if blastomeres be shaken apart in the early phases of segmentation, they give rise to dwarf larve, those from the four-cell stage being little more than half the size of those from the two-cell stage, and these in their turn little more than half as large as normal larve. The other is that similar dwarf larve have been produced by fertilizing the *enucleated* ova of certain species (such larve thus possess only the paternal chromosomes). Obviously, for due metabolism and growth to the normal size, there must exist a very precise quantitative relationship between chromatin and cytoplasm. This reduction on the part of both ovum and spermatozoon preserves that precise relationship.

5. An equally striking feature is that, leaving out of consideration the "accessory chromosome,"¹ these chromosomes are contributed to the new individual equally by both parents. These are, in fact, the one organ or component of the fertilized ovum and new individual, which is an identical contribution on the part of the two parents.² We have seen what curiously elaborate methods have been developed to insure their equality.

6. Equally striking and equally important is that, so far as we can determine, following the development of the new individual, it is evident that the same care is taken to insure that the chromosomes from both parents are distributed equally into each daughter cell, so that each cell of the various tissues of the adult is influenced by chromosomes derived from both parents. We do not say that chromosomes of paternal and maternal origin are of equal value. Everything indicates that is not the case; that now one, now the other, is the more potent; or, indeed, that in particular properties the one may be the more potent, in others the other. But histologically, or structurally, we cannot but recognize in these processes of fertilization and development a most marvellous mechanism tending to insure equal opportunities to both paternal and maternal chromosomes to influence the progeny.

7. Nay, more, we cannot but be led to the conclusion that in the chromosomes which become converted into the chromatin network of the functioning cell there must be contained the active controlling living matter of the cell—the biophoric matter. If we admit—as from everyday observation we are compelled to admit—that the properties or characteristics from both parents are conveyed to the offspring, and this variously in different members of the same family, we are equally compelled to admit that the chromosomes are the one component of the fertilized ovum capable of bringing about these results. The cyto-plasm of the beginning individual is derived in the main from the mother,

² Strassburger, indeed, points out that in the flowering plants the nucleus alone is contributed by the male sexual cell to the egg that is fertilized, its cell body being left behind during the passage down the pollen tube (*Darwin and Modern Thought*, Cambridge University Press, 1909; 104).

¹ Vide p. 153.

the centrosome from the father; the chromosomes are the one component contributed equally by both parents.

8. As we shall see in discussing the Mendelian principle (p. 168), the data bearing upon synapsis are of the highest significance. We are afforded clear indications that while each parent contributes one-half of the chromosomes present in the fertilized ovum and cells of the offspring, in that ovum and those cells only one-half of the chromosomes contributed to the one parent by the two grandparents can possibly be represented, the other half being cast out in the processes of oogenesis and of spermatogenesis, respectively.

In short, in these chromosomes of the spermatozoon and ovum and their subsequent distribution we have the anatomical basis of heredity. So far as it is determined and modified by sex, descent is dependent upon them, and no theory of descent can be considered adequate which does not take them into account. What is of direct interest to us, as students of disease, is that, following this line of thought to its legitimate conclusion, it is obvious that the inheritance of morbid states must be through the chromosomes, must be limited to conditions which are capable of affecting the biophores of the germ cells.

Numerous differences in detail have been recorded in connection with the successive stages of oogenesis and spermatogenesis in different species, but the broad principles are as here laid down. Studying the details of the process in various species, we are most strongly reminded of the slight differences in the technique of different assavers seeking to gain a perfect admixture of all parts of a given ore, in order that from a sample an ounce or less in weight they may estimate with accuracy the amount of precious metal in a ton of the material. To see such an assayer grind, and then shake, and roll, and mix the ground ore, smooth it out, cut into four parts, select two of the four diagonally opposite, and discard the other two, mix these thoroughly together and again divide, and continue the process until, finally, the amount is obtained just sufficient for melting in a small crucible, is curiously suggestive of what we see happening in these germ cells. In our opinion the most we can conclude is that here we have methods tending to insure that the ovum and the spermatozoon receive each a well-admixed sample of the biophores contained in the germ cells and, as a corollary, that hereby opportunity is afforded for every possible combination of the different orders of biophores contained in the parental germ cells. For, as we have already indicated (p. 137), and must now proceed to consider, while denving the existence of determinants, we have to recognize that biophores of more than one order are present in each germ cell, and it may well be in the different members of one set of chromosomes, and that inheritance is largely governed by the proportion of biophores of the different orders gaining entrance into the fertilized ovum.

The "Accessory Chromosome" and Sex.—For the sake of clearness and so as not to disturb our relation of events by continually noting an exception to the general principle, we have left out of account certain observations of the last few verrs which indicate that the number of chromosomes in a

FERTILIZATION

cell is not necessarily even, and that herein male and female individuals show a constant difference. McClung,1 of Kansas, was the first to investigate and draw conclusions from the appearances in question. Studying the spermatogenesis of certain insects, he noted that the cells could be divided into two approximately equal portions, of which the members of the one possessed a chromosome over and above those possessed by the other. And as sex is the only distinction which separates the fertilized cells into two approximately equal sets, he concluded that in the presence or absence of this accessory chromosome lies the physical and anatomical basis of sex. At first strongly contested, this view has of late gained an increasing number of adherents, more particularly through the studies and support of Professor E. B. Wilson, of New York, who has demonstrated the existence of the two orders of spermatozoa in close upon sixty species of insects, the difference involving the members of one, or, in a few cases, of two or three pairs of chromosomes. It is not necessary that the other member of a pair be entirely absent-either there is an unequal pair, one large member corresponding with a minute chromosome, or the latter has wholly disappeared. Now that Correns has discovered the same order of affairs governing the pollen and ova of directious (or ordinary flowering) plants, it is evident that we deal with a phenomenon of wide occurrence. The indications are that the cells of the female possess two complete series of chromosomes (Aa, Bb, Cc, etc.), those of the male the like series, minus one (A, Bb, Cc, etc., or Aa, Bb, Cc, etc.); that thus the mature ovum after reduction and prior to fertilization always has the full half set of chromosomes, the mature spermatozoon either the full half set, or the half set minus one. The hypothesis generally accepted by American workers is that the fusion of the ovum with a spermatozoon having the accessory chromosome gives rise to the female individual, with spermatozoon minus this chromosome to the male.² Nevertheless, in certain conditions this irregularity may appear in the ovum; thus, in insects producing a series of parthenogenetic generations (without fertilization) like the Aphides and Phylloxerans, and in which the parthenogenetic ova give origin to both males and females, while fertilized ova produce only females, T. H. Morgan^a has shown that the production of the one sex only is associated with the fact that the spermatozoon with the reduced number of chromosomes is so small, and contains so little cytoplasm, that it degenerates and does not attain to full development, there being thus only one order of functional spermatozoa. And as regards the males and females produced parthenogenetically, Morgan⁴ has made the remarkable discovery that the somatic or tissue cells of the males in the species studied contain only five chromosomes, whereas those of the

¹ Biolog. Bulletin, 3: 1902: 43.

² It will be seen that, strictly speaking, the term "accessory chromosome" is inaccurate and somewhat misleading. Compared with the number of chromosomes in the oocyte, neither order of spermatozoon possesses an additional chromosome, but one order is deficient in one or more chromosomes.

³ Proc. Soc. for Exp. Biol. and Med., 5; 1908; 55,

4 Ibid., p. 56.

female contain six. At some period, therefore, in the life of the parthenogenetic oocytes one chromosome disappears in cells destined to become males.

If this hypothesis be applicable to vertebrates, then it must shortly be shown from a study of ordinary somatic mitoses that the cells of the male contain one chromosome less than those of the female individuals of a given species. This has not to our knowledge been demonstrated. At the same time we admit that this hypothesis, based as it is upon our positive cytological data gained from widely different orders of living beings, is far in advance of any of the numerous and conflicting hypotheses of sex production of the past, based as they have been on controvertible and presumptive data, and incapable of exact verification hypotheses so crude and so unconvincing that it is useless to enumerate them.

It would take us too far afield to describe and criticise the opposing hypotheses of Castle,³ Bateson,² Berry Hart,³ and others, based upon these data. Not one of these has as yet established itself as satisfying all the conditions noted.

² Mendel's Principles of Heredity, Second Edition, 1909, and The Methods and Scape of Genetics, Cambridge, 1908,

^a Mendelian Action on Differentiated Sex, Edinburgh, Neill & Co., 1909.

¹ The Heredity of Sex, Bull. Mus. Comp. Zool., Harvard, 40: 1903: No. 4.

CHAPTER XII.

A RÉSUMÉ: THE BIOPHORIC HYPOTHESIS.

It will be well at the present point to pause and, looking backward, survey the road that we have travelled up to the present point, that we may see how far our quest has led us, and this because our present position gives entrance, to continue the metaphor, into the domain of heredity, and it is useful before entering this difficult territory to recall the path leading thereto. The present chapter will thus constitute a *résumé* of the conclusions reached in the preceding. We may characterize it as summing up (to this point) the biophoric theory, or, to be more modest, hypothesis, of living matter.

We have seen, in the first place, that all the phenomena that we regard as vital are manifestations of energy, and, as such, are bound up with matter, necessitating changes in the relationship of the molecules of the same. And what is more, that they are the attributes of a particular order of nitrogen and carbon containing bodies. We may speak of the molecules of this living matter as biophores, and, while still ignorant of their exact constitution, a study of metabolism leads us to regard them as "ring" formations, composed each of a ring of radicals, the component radicals having free affinities which are capable of being satisfied by the attachment of groups of free ions from the surrounding medium whereby "side-chains" are built up (synthesis).

These side-chains are capable of detachment (dissociation) when through alteration in the surrounding medium, ions are present in the same which exert greater attraction. By this means the biophores again become unsatisfied. Further, the side-chains themselves are to be regarded as unsatisfied bodies, able to build up like side-chain bodies in series. It is this unsatisfied nature or constitution of the side-chain bodies (and of the biophores) that permits them to act as *enzymes*. Therefore, it is this condition of persistent recurrent unsatisfaction that distinguishes living from dead matter, and that underlies all the processes of assimilation and dissimilation.

We have seen, further, that, given molecules of this nature, the sidechains attracted, and so the constitution of the biophores *in toto*, must very largely depend on the surrounding medium. The entrance of free ions of other nature into that medium, and alterations in the relative proportions of the ions there present, will lead to the building up of modified side-chains, and, as in the process of growth (polymerization) such side-chains, or some of them, must be utilized in the formation of new central molecular rings, so alteration in environment, if continued, is apt to lead to alteration in the essential constitution of the

A RESUME

biophores. So long as the environment remains unchanged, so long the basal constitution of the biophores remains unaltered; modify the environment, and, provided the modification be not so extreme as to arrest the series of ionic interchanges, the molecular constitution of the biophores becomes modified. It is this property of the biophores that is the basis of variation, adaptation, and evolution.

We have equally to recognize the co-existence of another property in the biophores, that of fixity of constitution, inherent in the relationship of the constituent atoms and radicals the one to the other. The biophores, that is, are to be regarded as built up according to a particular (geometrical) arrangement. It is this arrangement which prevents the indiscriminate linking of ions and grouping of radicals from the surrounding medium, and necessitates that, if unsuitable ions do become attached and the constitution thus interfered with, the compound loses its special properties; the biophore becomes "dead" matter. There are thus certain definite limits to modification, and these determine *heredity* in the narrower sense.

A study of the histology, physiology, and embryology of the cell renders it evident that the biophores are contained in the nucleus; that they are intimately associated with the chromatin. The fact that the main bulk of (dead) nuclear chromatin is formed of nucleoproteins, leads us to the conclusion that, although the biophores cannot be regarded as essentially nucleoproteins (for nucleoproteins, as such, have not by any means all the attributes of living matter), nevertheless the main component radicals of nucleoproteins must be contained in the biophores; or, otherwise, that the chemical changes which bring about the death of the biophores convert them largely into nucleo-proteins.

The extent of growth of the unit mass of biophores is determined by two factors: (1) The extent of surface of the same in contact with the intermediate medium (cytoplasm) in relation to the mass, and (2) the extent of the surface of the cytoplasm exposed to the external medium relative to its mass. These relationships (together with the nature of the cell reactions and activities) determine the size of the We must regard the multicellular organism not as a colony of cell. individual cells, which have primarily united for mutual protection, but as an adaptation, or means, whereby continued growth of the biophoric matter is most economically insured, with due retention of the above relationship of surface to mass. When this mass of biophoric matter passes the optimum the above relationship is preserved by nuclear division (with coincident increase in biophoric surface exposed to the cytoplasm), followed by cytoplasmic division (with like increase in cytoplasmic surface exposed to the external medium). In the multicellular organism the cytoplasmic division is in general incomplete, cytoplasmic bridges uniting related cells. With each segmentation of a cell, following upon biophoric growth, there is equal division of the biophores between the daughter cells.

The formation of multicellular masses renders it inevitable that all the cells of the mass are not exposed to an identical environment, and through altered constitution of the cytoplasm in the first place, brings about alterations in the biophores controlling cells subjected to different environment. It is to this effect of modified environment acting upon biophoric material of a common order, and to this adaptation of the biophores to such altered environment, that we must attribute tissue differentiation. Alteration in the biophores in its turn effects alteration in the cytoplasm and histological and functional modification of the entire cell. It is physically impossible for the ovum and its nucleus to contain determinants or specific biophores for the various tissues and the localized variations in those tissues which characterize the various species.

In the process of differentiation the constitution of the biophores in the different orders of cells undergoes progressive modification of such a nature that those controlling the main mass of cells of the body (the somatic cells), while capable of growth, i. e., of polymerizing identical biophores, and so of giving rise to cells of like nature, become incapable of giving rise under any conditions to the complete organism. They may, at most, exhibit an incomplete reversion to a simpler, less differentiated state. Those most highly differentiated lose even the power of reproducing cells of like nature. The power of reproducing the individual has become restricted to a group of cells (the germ cells), which, so long as they remain integral portions of the parent organism, present a minimum of differentiation. In these, presumably, the biophores undergo minimal modification. They are characterized histologically by showing no primary reduction in their chromatin in the process of successive division.

In all multicellular organisms (and in unicellular, save the lowest forms) the species is continued and the specific living matter propagated, if not in every successive generation, at least ultimately, by conjugation and fertilization. Sex has been developed, and fertilization is the process of fusion of the male (invasive) with the female (receptive) germ cell. The process is essentially one of combination of equal amounts (or numbers) of biophores from the two parental germ cells to constitute the nucleus of the new individual. Prior to the act a remarkable series of changes occurs in the germ cells involved, changes which would seem to have for their object to supply to the eventual ovum and spermatozoon an admixture or selection of the biophores present in the parental germ cells and to reduce the number of the same, so that the proportion present in the fertilized ovum is identical, relative to the cytoplasm of the same, with that of the unaltered germ cells of the parents, or, more exactly, in the ovum from which these gained development.

CHAPTER XIII.

INHERITANCE.

WE come now to consider the problems of heredity, by which, speaking strictly, we mean the conveyance to the offspring of the properties of the parents and of the parental stock, so that familial, racial, and specific characters are passed onward from generation to generation. Were inheritance pure, did the child absolutely reproduce the parent, were all the members of one family and stock identical in form and characters-as identical as are crystals of the same salt-or were they born identical, and did they owe individual differences purely to the diverse influences to which each individual becomes subject after birth, the problem before us would be relatively simple. But this is far from being the case. Interwoven with heredity there is variation, and this not of one but of two, if not of three, orders. We see, in the first place, that influences from without modify the individual. It is a matter of familiar knowledge that the soldier, the sailor, the professional musician, the undertaker, assume pronounced types; they have become modified by their course of life. In man and mammals we can subdivide these modifications (as one would term them, in contradistinction to inherited variations, or variations proper) into those acquired during intra-uterine and during postnatal existence. In the second place, from the fact that the individual is the result of amphimixis, of the fusion and intermixture of germ plasm from two parents who are not themselves identical, and whose germ plasms are not identical, it follows that the individual is not identical with either parent. This, again, is a matter of familiar knowledge; the child, while resembling the parents, now more the one, now more the other, is identical with neither. If the general tendency of successive acts of amphimixis is by the mingling of diverse germ plasms to produce intermediate grades, to maintain the mean, and so preserve the type, it clearly at the same time necessitates that each individual varies from either parent. And, thirdly, the study of the progeny of one pair of parents demonstrates to us that the interaction of the parental germ plasms-or, to preserve our terminology-of the germinal biophores in successive acts of fertilization, induces variation. No two children of the same parentage are born identical; no two fish, even of the same spawning (in which thousands of ova and millions of spermatozoa are matured at the same time), though these, admittedly, more nearly approach identity. As we shall have to point out, not only have we to recognize variation in the orders of biophores supplied by each parent to ovum and spermatozoon, respectively, but also we are forced to the conclusion that during the course of the individual life of the parent individual, biophores are liable to undergo a certain amount of modification of their constitution. Indeed, if we do not accept this capacity on the part of the biophores, to undergo modification, the facts of variation become incomprehensible.

Our study thus is not one of heredity pure and simple, but of the interaction of heredity and variation, and obviously, with so many factors to take into account and to analyze, the study is most complicated. Keeping in mind that here we are engaged upon an introduction to pathology, with particular reference to the principles underlying the same, the sense of proportion demands that our treatment be relatively brief. In almost every work upon general pathology with which we are acquainted the subject is dismissed in a brief paragraph or two. This we regard as a grave omission. It demands fuller treatment, and that because the physician, in his study of the individual case, is constantly brought to ask himself how far a given condition is an individual acquirement, how far it depends upon a vice of organization and is the outcome of inheritance; or, again, is himself asked to advise whether one or other condition of the parent, or would-be parent, is liable to influence the prospective offspring. And what is more, as we hope to show in later chapters, a broad grasp of the principles of heredity aids us materially in comprehending not a few pathological conditions, while, conversely, a knowledge of data acquired by the pathologist aids us equally in determining what are those principles.

While we are not blind to the virtue of according to the student a knowledge of diverse opinions and opposing theories, we believe that the better teaching is, where possible, to inculcate definite views, as also to afford a thoughful presentation of the data upon which one theory is based rather than a didactic epitome of many. And as we are already upon record as having definite views upon this subject of heredity.¹ and more particularly have based our theory upon pathological data; as, further, that theory conforms with and continues the considerations brought forward in the preceding chapters, we shall here limit ourselves largely to a presentation of that theory, although, at the same time, we shall endeavor, as already stated, to indicate to the reader in passing what are the other theories and what their salient points.

THE DIFFERENT FORMS OF INHERITANCE (*i. e.*, HEREDITY PLUS VARIATION).

We may, in the first place, make a classification of the properties exhibited by the individual. These are:

1. Individual, *i. e.*, properties peculiar to the individual, and not recognizably inherited from either parent or from any ancestor.

¹ British Medical Journal, 1901; i, June 1; and article "Inheritance and Disease," Osler's Modern Medicine, 1: 1906.

THE DIFFERENT FORMS OF INHERITANCE

2. Parental, *i. e.*, properties possessed by and peculiar to one or other parent and obviously inherited from that parent.

3. Familial, *i. e.*, properties possessed by and peculiar to the family of one or other parent.

4. Racial and stock properties; characters common to a particular stock.¹

5. Specific or ex-specie, *i. e.*, properties peculiar to the species, those, for instance, distinguishing the human individual from the ape.

6. Class, *i. e.*, properties peculiar to the class: those distinguishing man from animals that are not mammals.

And we can continue the classification yet farther to ordinal distinctions, peculiar to the particular order, *i. e.*, distinguishing man as a vertebrate from the invertebrates, and even to those distinguishing man as an animal from plant forms.

Studying, in man alone, the various departures from type, one prominent fact makes itself evident, namely, that the most basal features are those which are most commonly impressed on the individual-a fact which, in itself, is one of the strongest evidences in favor of evolution and against special acts of creation. Vertebrate characters are more firmly impressed and more constant than mammalian, mammalian than human, human than racial, racial than familial. In other words, a study of variation in its broadest aspects confirms the view, already indicated, of the progressive building up of the biophores in the course of the evolution of the different species. The facts are in harmony with the view that the original molecules of living matter were of relatively very simple constitution, becoming progressively more elaborate and complicated, not by loss of that original constitution, but by the addition of side-chains, those side-chains becoming successively integral portions of the more complex molecule. Under like conditions of environment, the same epigenetic series of transformations of these biophores must occur; polymerization, growth, and reproduction of these complex molecules is, with favorable changes in environment, still permissible, but new side-chains are either added to, or replace, those previously present. Following out this line of thought, it is seen that ontogeny, i. e., the course of development of the individual, becomes an epitome of phylogeny, *i.e.*, the evolution of the phylum or race,

¹ Our terminology is here a little indefinite; we speak indifferently, for instance, of the human race and of races of mankind, hereby meaning two different things. Of the human species we recognize several subspecies: Indo-European, Mongol, Australasian, etc., and of each of these several vares, of the Indo-European for instance, the Teutonic, the Celtic, etc., and of each of these races several stocks. Thus, among those of Teutonic descent, the Anglo-Saxon differs in certain recognizable particulars from the North German, the North German from the South German. Some of these differences are doubtless due to the fact that the stocks are not pure, that at one or other period there has been extensive intermarriage with individuals of other stocks, but some at least are independent of this cause, as witness the differences that already show themselves between the Anglo-Saxons who have remained in England and those whose ancestors migrated to North America or Australia.

INHERITANCE

not as a matter of mere historical survival, but as an epigenetic necessity. For the biophores controlling and giving rise to a given tissue or part of the body to be able to order the particular structure of that part, they must have a particular chemical or physical constitution. To gain that constitution the original common biophores of the fertilized ovum must in growth and distribution to the different organs pass through a particular series of chemical changes. To exhibit these successive changes they must be subjected to one particular series of alterations in environment. In ontogeny, therefore, and the development of the individual, the cells and the contained biophores reproduce the conditions undergone by the race in its evolution-this being essential for the due production of the specialized biophores of the different tissues. Where, in this process, conditions are such that the same reaction and end result may be attained by one step in place of two or more, the ontogenetic process is correspondingly abbreviated, and to this extent the ontogeny fails to reproduce the phylogeny. This, in brief, is what is known as the Recapitulation Theory, and our conception of its meaning.1

Racial Inheritance.- It is unnecessary to detail examples of hereditary properties older than racial, for the reader can easily call these to mind. And, as regards racial and stock inheritance, the same is largely true. Everyone is familiar with, for example, the differences in the form and structure of the individual hairs and the amount and nature of the cutaneous pigment in the different races of mankind, and has some knowledge of the differences in stature and in the conformation of the skull in the same. It may be of value from a pathological point of view to recall vet other differences, functional in nature and co-existent with these histological variation, which the morphologist is naturally apt to overlook. We refer to differences in reactive power toward various pathogenic agents, microbic and non-microbic. These we find well marked in the different races of domestic animals. Thus, the native cattle of Japan and the buffel, or native cattle of Austro-Hungary, are much more susceptible to tuberculosis than are ordinary European and American breeds, and one race of Algerian sheep is highly refractory to anthrax, a disease to which sheep in general are peculiarly susceptible. Between the races and stocks of mankind many such differences have been noted. Those of European descent and Malaysians subjected to the same conditions react very differently toward plague and beriberi. Both, it is true, are susceptible, but a far smaller proportion of Europeans exposed become victims to these diseases, and a far smaller proportion of those taking the diseases succumb. The reverse would seem to be the case as between those of European descent and negroes in regard to vellow fever; here it is the Europeans that are far more susceptible, while, contrariwise, negroes

¹ See more particularly Hurst, Natural Science, 6: 1895. It is also laid down at length by Reid, *The Principles of Heredity*, London, Chapman and Hall, 1905, without, however, any clear recognition of the underlying eauses.

THE DIFFERENT FORMS OF INHERITANCE

and American Indians and native races in general are more susceptible to tuberculosis than are "white men" under conditions of civilization.

So, also, with non-microbic diseases. Those of the Hebrew race are more prone to certain nervous affections and to diabetes than are the surrounding Gentile population; the French more subject to hysteroepilepsy; the English to gout; the inhabitants of North America of European descent to dyspensia and disorders of dentition.

And these finer distinctions even exhibit themselves in the almost indescribable *nuances* of character. It is not necessary to quote Shakespeare or Sir Thomas Browne on these matters; one has but to "walk the wards" of any large metropolitan hospital and observe how those of different nationalities bear themselves under conditions of disease, to become convinced of these pronounced underlying variations.

It may be argued that the differences are purely or in the main the result of differences in mode of life and general surroundings. But this is exactly our point: it is the various grades of variations in environment that are essential causes of these differences, from the pronounced distinctions in cutaneous pigment down to national gout and national dyspepsia.

This, however, may be said, that a study of these finer differences between the races and stocks of mankind throws doubt upon the argument of Weismann and other morphologists, based upon single morphological variations—that these demonstrate the existence of determinants. We find that such impalpable variations as occur between different stocks of the same race of mankind—morphological variations so slight that we can scarce measure them by instrumental means—are constantly accompanied by functional variation; or, otherwise, that a change in one slight direction affects not a single part, but the whole organism; while, conversely, change of any one part which produces no obvious morphological alteration has, nevertheless, an influence upon the body as a whole.

Familial Inheritance.--It is a well-known fact of the same order that not merely is there a certain likeness between the members of one family, but that certain traits peculiar to a family present themselves generation after generation with remarkable persistence-stature and length of limb, shape of some special part, as, for example, the "Hapsburg lip," which has characterized the present reigning House of Spain for so many centuries; actual malformations, like accessory digits, which have, in certain cases, been traced back for close upon three centuries; morbid conditions or tendencies to the same (diatheses), such as albinism (deficiency of cutaneous pigment), Daltonism (color blindness), hemophilia (liability to excessive hemorrhage as the result of relatively insignificant injury), nervous disorders; or, again, traits of character and tricks of expression. And all this we note in the absence of interbreeding-marriage of cousins, for example-which we might reasonably expect to intensify such family peculiarities, and in spite of the continual introduction of "new blood" into the family by marriage with members of other families not presenting these particular features;

INHERITANCE

so that we are forced to recognize the possession of what we may term *dominant* properties, by the germ plasm or biophores of particular strains, whereby, in amphimixis, these biophores assert themselves in one or other direction, overshadowing the effects of the conjoined biophores.

À study of family inheritance reveals other remarkable features. Thus, there are certain characteristics which affect male and female members of the family alike, others which appear in the one sex only, although they are conveyed by the other. The daughter of a family may, throughout her life, be free from gouty manifestations, and her daughter also, but her male children may exhibit gout at or before attaining middle age. This interrupted or sex-limited descent is most pronounced in the case of hemophilia; in this the female members of the family are rarely "bleeders," but they give birth to sons who are. Irrespective of sex, also, we encounter frequent cases in which one or other member of a family exhibits peculiarities not present in the parents, but well marked in one or other grandparent, or even in a great-grandparent. This reproduction of features, absent from the parents, but present in a preceding generation, is termed *atavism*. It has, as we shall note presently, to be distinguished from *reversion*.

In case after case the existence of this atavism renders it difficult, if not impossible, to trace a family peculiarity to its origin-a case of polydactylism (supernumerary digits), for instance, occurs in a family whose members are otherwise provided with the normal number of fingers and toes. To determine whether this is a wholly new appearance in the family, or merely the breaking out again of a dormant tendency, demands an extensive knowledge of lateral as well as of direct lines of descent. A man must have two parents, and may have -if there has been no inbreeding-four grandparents, eight great-grandparents, sixteen great-great-grandparents; thus, an intimate knowledge of at least sixteen family histories is requisite; such knowledge is notoriously hard to gain, especially where blemishes are concerned. Such difference must have made itself manifest in some one particular generation, must have had an origin. We have to turn to the smaller animals, with rapidly recurring generations, to throw light upon this point, and even with these have to breed for several generations and preserve exact records.

I owe to an old student a remarkable case of the hereditary transmission of family defects, which well illustrates many of the points here referred to. In 1620 two brothers landed from England and settled at Woburn, Massachusetts, and these, according to family history, married two sisters. The family and the descendants of the one brother have since then shown no abnormality; in the children of the other brother, N. X., himself, according to tradition, polydactylous, polydactylism presented itself. He had ten sons, whose descendants are now scattered through North America. For the next two generations it was dormant, or at least there are no records of its existence. In the direct line of my informant it has indeed been dormant for three generations.





PLATE IV




ato

nibe



FAMILIAL INHERITANCE

His great-grandfather was free, though other members of that generation were affected; his grandfather was also free, as was his father, although an uncle and aunt were affected. Of his father's twelve children, eight were affected, the condition being for the first time complete in himself. By complete is here meant that not only did the condition affect all his members, so that he had six fingers and six toes, but all the accessory digits were perfectly formed. What is more, his young son has them all perfectly complete. Another characteristic of the family history is that, whereas the daughters of the family may show the effect, it tends to die out with them; their children have normal digits. In this way, according to our patriarchal method of determining the family, the defect tends to remain familial, descending only through



the males. The potency of the "blood" of this family is, in other respects, strongly pronounced; there is a succession of large families, and the different members exhibit a great family likeness, so great that my informant could salute a stranger travelling in the West, "Good morning, Mr. X." and have him return the salute, "Good morning, Mr. X.," with the further remark: "I see you have the family sign (referring to his six fingers). I do not possess it, but my father did, and so does one of my eight children." And, on inquiry, the cousinship between the two was found to be at least beyond the third degree of removal. The following exbihits the genealogy of my informant's branch of the family:

A careful inquiry made by my friend, Dr. Brockbank, of Manchester,

INHERITANCE

has succeeded in bringing to light a family of the same name in England, two members of which migrated to America in or about 1620, and in this at least one member of the present generation exhibits poly-dactylism. It is evident, therefore, that the tendency to polydactylism *must* date back to the generation preceding the *emigres* of 1620, and *may* have originated much earlier. There are, indeed, indications that the tendency may be traceable back to earlier than 1500.

CHAPTER XIV.

PARENTAL AND INDIVIDUAL INHERITANCE.

STRICTLY speaking, every property possessed by the individual which is not and cannot be ascribed to intra-uterine and postnatal *acquirement* is the individual inheritance of the individual. It has reached him through the parental germ plasms. Specific (*ex-specie*), racial, and familial traits become the property of the individual. But here we would deal with those features which, peculiar to one or other parent, reappear in the offspring; and again, those which, not observable in either parent but present in the offspring, can only be ascribed to the interaction of the two parental plasms. What has once been inherited may be conveyed to a subsequent generation.

A little consideration shows that these two conditions as here defined are practically identical, and must be considered together. Considering these together and studying the various grades of inheritance, we find that these may be classified as follows:

A. Blended Inheritance.—Although the active interest taken of recent years in the next form to be mentioned is apt to obscure our view, as regards the majority of properties, this must be regarded as the most usual type. Of this the most familiar example is the skin color of the mulatto. The offspring, that is, tends in the main to exhibit a reblend of the paternal and maternal features, intermediate between those presented by either parent. In general, that is, we observe an equality of influence on the part of the respective germ plasms, and that the tendency of fertilization is to preserve the mean and perpetuate the type.

B. **Particulate Inheritance**,—'There are, however, many exceptions to this law of blended inheritance, nor have we as yet determined adequately what it is that determines these. All that we can say is that we are forced to recognize that, in respect to certain properties possessed by the parents, we have to note an incapacity to blend; they are antagomistic. First and foremost among these is the series of associated sexual properties; the offspring of most animals must be either male or female. Anything of the nature of coincident equal development of both sets of sexual organs is found incompatible with perfect function of either, and is an attribute of imperfect organization. We see the same in many properties which we may regard as of subsidiary importance. When one parent has brown eyes, the other blue, the children have either blue or brown eyes, rarely those of an intermediate color. And what is more, as between alternative antagonistic properties, in a given mating one of the two is apt to be *dominant*.

PARENTAL AND INDIVIDUAL INHERITANCE

MENDEL'S LAW.

This was well shown in the remarkable observations of the Austrian monk, Mendel,¹ upon hybridization, *i. e.*, upon the cross-fertilization of well-marked varieties of certain flowers, notably of the pea: observations which, neglected for long years, have of late been widely confirmed, by zoologists as well as by botanists. If two well-differentiated varieties of such a plant as the pea be selected and grown, varieties presenting, when grown pure, constant differences, differing in such particulars as position of the flowers on the stem (either axial or terminal), length of stem (whether markedly long-stemmed or markedly short-stemmed), character of the seed pods (either inflated or constricted between each seed and its fellow), seed coat (whether smooth or wrinkled), and if these be cross-fertilized, the pollen of the one variety being employed to fertilize the ova of the other, it is found that the hybrids which result manifest these antagonistic properties according to a very definite law. One of each pair of properties is dominant, i. e., present, in these "bastards" of the first generation, overwhelming in them wholly or mainly the corresponding character present in the other parent. Thus, if we hybridize a red-flowered pea with a white-flowering species, the first set of hybrids will have red flowers, and in this respect be indistinguishable from the red-flowered parent. In this respect they will show no relationship to the white-flowered parent. Red color of flower here is dominant, white recessive, and this irrespective² whether the pollen of the red-flowered variety be employed to fertilize the white-flowered, or vice versa. Almost always it is the progressive characteristic which is dominant. In this case, for example, red color of flowers is the positive acquirement; white color is latency, or loss of this acquirement. It is the positive acquirements-color, hairiness, presence of starch, in seeds etc.-characterizing the species as a whole that are dominant; absence of these, that are recessive.


But while this is the case, it does not mean that the characters of the other parent are completely thrown out; the recessive character is only latent. In other words, from what we know of nuclear division and distribution of parental chromosomes in the cells, biophores derived from both parents are conveyed to all the cells of the individual, only, as regards any particular feature, those from one source may be more active than those from the other. This is surely the case with the germ cells, as pointed out by Mendel.

³ Verhandl, d. Natur, Vereines in Brunn, 1866, reprinted by Göbel in "Flora," 89: 4001; 364, and translated by W. Bateson in Mendel's Principles of Heredity: A Defence, Cambridge Univ, Press, 1902, 2nd. Edit., 1909. De Vries, who rediscovered Mendel, gives a singularly clear account in his Species and Varieties, Chicago, 1906, lecture 10.

² In the main, Heider (Ueber Vererbungsgesetze, Berlin, Borntrager, 1905) gives some exceptions among plants, in which differences are noted. There are peculiarities in the development of the plant ovum—not seen in the animal—which help to explain this.

MENDEL'S LAW

Peas have this advantage, that they are capable, under ordinary conditions, of self-fertilization, and so the disturbing effects of cross-fertilization with pollen from other sources can be prevented. If, now, the offspring of this *bustard* first generation be grown and brought to flower, in their characteristics they are found to follow closely a numerical rule; the recessive character reappears in one quarter of this second generation, and if the flowers of this one-quarter be allowed to self-fertilize themselves, the subsequent generations show constantly the recessive character; the dominant is wholly cast out. As regards this one feature of flower color, for example, one can, after bastardization, regain a white-




Leaf characters of hybrids of Urtica pilalifera and U, dodariii (Correns): F^{i} , of first hybrid generation; F^{i} and F^{i} , or second and third self-fertilizations. The dentate character of the leaf edge is seen to be a dominant property.

colored variety. As regards the other three quarters, they it is found, divide themselves into two orders: One of the three quarters consists of purely dominant plants, and subsequent generations grown from these remain purely dominant—always, for example, have red flowers. The other two quarters, or one-half of the whole number of this second generation, have the properties of the first generation of hybrids; they are hybrids, and in each subsequent generation they give rise to the same proportion of pure dominants, hybrids, and pure recessives, namely, one-quarter that are found to be pure dominants, one-half hybrids, and one-quarter pure recessives.

The action of this law is, perhaps, best grasped from the accompanying

sive character as R, the proportions of the different types of hybrids evolved is represented by the formula n(D + 2DR + R); or, more exactly, n(DD + 2DR + RR). which has leaves with almost uninterrupted edges. Mathematically expressed, if we designate the dominant character as D, and the recesa species with strongly toothed or dentate leaves, and Urtica dodartii, figure, afforded by Correns, giving the character of the leaves in suc-cessive generations of hybrids of two species of nettles, *Urlica pillalifera*,

biophores as do the hybrid ova. of the pure dominant and recessive types receives the same number of We use the DD and the RR to indicate that each individual ovum



Schema of Mendel's have for a single pair of "mutagonishi," properties [A, the results of bybeich ration of a pure dominant (10) with a pure reseaves (10) form, 3), the decould of ensaine a hybrid with a reseave form. 20 per cert of progeny pure reseaves, 30 per cent, hybrid (hur apparently dominant); C, the result of crossing a hybrid with a dominant form; all apparently dominant (nucl) per cent, pure, 20 hybrid. (Massea)

If further generations be grown and self-fertilized, it is found that plants of the DD type always produce DD offspring; plants of the RR type always produce RR offspring; plants of the DR type constantly give the different types of offspring the proportion of n(DD + 2DR RR

mind that the law is not universal. The Limitations of Mendel's Law. It must, however, he borne in

acquirement on the part of one group of members of a species mated 1. It deals only with that class of cases in which we encounter an

PLATE V Mirabilis' Jalapa C-TO and the second alba + rosea round Carles to E EI П \overline{DD} RR DRDR



MENDEL'S LAW

with a *varietal* deficiency on the part of other members, and this varietal deficiency is not to be regarded so much as an absence of a particular property, as a lying latent of the same: the phenomenon, that is, is not strictly the result of interaction of opposed properties.

 Where in the hybrid the one property dominates, it does not drive the other out; the other is latent through the whole existence of the hybrid, and at times, under suitable conditions, may show itself.

Thus, as de Vries points out, the hybrid between the blue Veronica longifolia and its white variety has flowers with blue corolla; here and there among 1000 or more of the hybrids a completely white spike may be noted, or a side branch with white flowers, or one side of a spike may be blue, the other white. Bud varieties may show themselves, that is, with the recessive property in the ascendancy. It is there all the time, as shown by the subsequent generations, and now some influence has rendered it the more active in certain cells. Here possibly we have an insight into the nature of mosaic inheritance. (See p. 175.)

3. At times the coincident influences of both dominant and recessive unity is recognizable; in other words, we have indications of apparent blending rather than of particulate inheritance, indications that the one form passes into the other, as shown in the accompanying figure from Correns of the result of hybridizing two varieties of *Mirabilis jalapa*, *rosea* and *alba*. All the flowers of the first generation are of a lighter pink color than those of the dominant parent, although in the second generation the formula works out obviously into DD + 2DR + RR^{-1}

4. The law is of no avail in connection with hybridizations between distinct species. As is well known, it is only between species that are closely allied that such hybridization has any result, and then the offspring tends to be infertile. There are, nevertheless, cases in which a somewhat fertile progeny results, more particularly among plants. As de Vries points out, the conditions here are different; in the Mendelian cases we deal truly with pairs of conditions; the species and its variety differ in the presence or latency of a given property. Different species may, it is true, if allied, have many properties in common. Their difference lies in the fact that one has an acquirement in one or other direction (or, it may be, several) which is wholly unrepresented in the other; as a result the mating is incomplete and unbalanced, one or more units are introduced by the one germ substance for which the other introduces nothing that corresponds. If hybrids are produced under these circumstances, then, as de Vries has demonstrated experimentally:

¹ Bateson affords an equally striking instance. The *blue* Andalusian fowl is a hybrid of the DR type, produced by mating a dominant black Andalusian with a recessive white splashed with black. Breeders have striven in vanit or raise a pure strain of the blue variety; mating two blue birds gives progeny in the proportion 1 black, 2 blue, 1 white splashed with black; mating black with splashed white gives a *first* generation all blue.

(a) Different types are produced according as to whether the male element of the one parent, Λ, is used to fertilize the ovum of B, or vice versa.

(b) The hybrids yield very little seed, whereas Mendelian hybrids are as fertile as the parental stocks.

(c) Self-fertilized, they remain true to type and constant; there is no segregation or reversion to either parental form.

(d) All the individuals of each generation are alike.

Here, it will be seen, we have a wholly different condition of affairs. As de Vries well points out, up to the present there has been a great vagueness as to what constitutes a species, what a variety. Such hybridization would appear capable of affording an answer in any given case; while, conversely, cases which do not conform to the Mendelian law may be explained as demonstrating that the one or the other parent exhibits a specific and not a varietal difference from the other; in other words, that the point of difference between the two is not a retrogression. but a positive acquirement. Nevertheless, this cannot be affirmed with absolute precision, or, more accurately, there are cases in which it is impossible to lay down the limits between variety and species. Boulenger, of the British Museum, for example, studying one of the common European frogs, found a succession of varieties in different localities, neighboring and closely allied varieties freely interbreeding. The two extremes of the series were widely different in color and markings, so as to be regarded as wholly distinct species-and as such they comported themselves-for in one locality he found them inhabiting the same waters and incapable of interbreeding.

The Mendelian Law for Multiple Pairs of Features .-- When the individuals of the two Mendelian varieties thus hybridized differ in several pairs of features, the result, though following strictly the same law in respect to any one particular pair of features, becomes much more complicated. For two pairs of features, Ab and aB (Aa representing one pair of features, Bb the other, the capital letters representing the dominant feature of each pair), respectively, while in the first generation all the individuals will be of the AB type (the dominant features prevailing), in the second there will be sixteen possible combinations between germ cells carrying the different admixtures of these properties, and of these 16 possible combinations, externally, 12 will show the dominant A, 12 the dominant B character, 9 will show both dominant characters combined, 3 only will present the recessive a character, 3 the recessive b character, and 1 only will show both recessive characters (ab). For these and greater numbers of pairs of opposed characters (n), the law is that there will be 2^{n+n} possible combinations, 2^{n} forms which are externally different, and 3ⁿ which, from the internal constitution of their germ cells, will produce different orders of descendants. To show this in algebraic form, illustrating the different possible combinations, AB.AB, ABaB . . . to ab. ab, would occupy more space than perhaps the subject demands. Those who would follow it may

MENDEL'S LAW

be referred to Bateson's work.¹ Two important points are deducible from a mathematical study of these possible combinations between germ cells of different characters, and have been amply proved in practice: (1) That when the species crossed differ in two pairs of characters, one-sixteenth of the second and subsequent generations is liable to reproduce in a pure form the ancestral features of one or other of the original parent species, and by proper mating may be propagated, showing no trace of the *mésalliance* (as regards these two characteristics): and (2) that another one-sixteenth exhibits permanently the combination of the dominant character of the one species with the recessive character of the other-whereby, through proper self-fertilization or inbreeding, a new race or subspecies may be developed which retains this particular combination of features. This, for breeding purposes, is a matter of the very highest importance; already, by carrying out these indications and selecting advantageous features in two different species of plants, it has been found possible to develop species combining the two; as, for example, to obtain a new variety of wheat combining the rich yield of the English variety with the good flowering and early ripening qualities of the Manitoba wheat. It is along these lines of selection and appropriate mating that during the past centuries blindly, and with numerous failures, our breeders have developed the various pure breeds of domestic cattle; and that, in more recent years, Burbank has attained such marvellous developments in the practical production of improved varieties of fruits and flowers.

Unfortunately, with the human species we can so far hope to obtain little of practical benefit along these lines, nevertheless the abundant studies of the last few years have thrown some little light upon human pathology. In the first place, it has to be noted that this law, shown to be of wide application, among plants has, where tested, been found to apply also to animals. Coutagne² has proved it for silkworms; Lang² has shown that it holds in snails; Bateson⁴ with fowls; Cuenot,⁵ Darbishire,⁶ and G. M. Allen⁷ in mice; Castle,⁸ Woods,⁹ Hurst,¹⁰ and Schuster¹¹ in rabbits and guinea-pigs.

In man himself it is known that there occur such opposed pairs of characters; thus, where one parent has brown eyes, the other blue, the children do not have eyes of intermediate color, but either brown or blue. Davenport¹² and Hurst¹³ have independently shown that there

¹ Bateson, loc. cit. See also Brit. Med. Jour., 1906; ii; 61.

² Bull, Scient, d. I. France et I. Belgique, 37: 1902.

^a Festschr, f. E. Haeckel, 1904; 439.

⁴ Repts, to Evolution Committee, Roy. Soc., 1: 1902, and 2: 1905.

⁵ Arch, de Zool, Exp., Notes et Revues, 1902; 27, and 1903; 33.

⁶ Biometrika, 3:1904.

1000

7 Proc. Amer. Acad. of Arts and Sciences, 40: 1904.

⁸ Carnegie Institution Reports, Washington, No. 23: 1905.

⁹ Biometrika, 2:1903:299.

¹⁰ Jour. of Linnaen Soc., Zool., 29: 1905.

¹¹ Biometrika, 4:1905:1.

13 Proc. Roy. Soc. B., 80; 1908; 85.

¹² Science, 26:1907:589.

174

are two orders of eyes, the blue and hazel group, with pigment only in the back of the iris, the browns, greens, "ringed," and spotted group, with pigment also in front; this latter is the dominant, the former the recessive type. Study of long series of different matings in man shows a very close conformity to Mendel's law. Dark hair also, though not to such a marked extent, tends to be dominant over flaxen; but, confessedly, it is difficult to make accurate observations where each successive marriage introduces widely different "blood." We can recall one other definite observation made upon man, namely, that of Castle.¹ Albinos occasionally present themselves among negroes-individuals, that is, devoid of cutaneous pigment. As might be expected from what we have already said (p. 170), such loss of pigment, being a loss of a character, a retrogression, is recessive, and not dominant. Castle² and Bateson³ have, independently, suggested that sexual characters come under Mendel's law—that male and female properties are antagonistic. one or other presenting itself, while the other lies dormant, to show itself in a later generation. This cannot as yet be regarded as determined, but true it is that there are combinations of sets of characters which are combined in descent. The case already noted (p. 164) of polydactylism, conveyed along the male line," and the opposite conveyance of hemophilia through the female (although the female herself may show little tendency to manifest the condition), are instances in point.

GALTON'S LAW.

Before leaving the subject it is necessary to refer to a law which, for a time, was regarded as antagonistic to that of Mendel, but now has been shown by Darbishire⁵ and Correns to be another, though perhaps less satisfactory, expression of the same-less satisfactory in that it expresses the source of the effects in a given generation rather than an analysis of the same. It was held that blended and antagonistic inheritance were opposed, and that this law dealt only with the former. As we have pointed out, the two are not opposed, but various transitions are to be recognized, and, as a matter of fact, Francis Galton,6 who established his law upon a very extensive analysis of the coat color of Basset hounds, truly employed a pair of particular features. The law, as determined by him, is to the effect that in the composition of the individual the two parents contribute the half of the total (or each onequarter), the four grandparents one-quarter (or each one-sixteenth), and so on. According to this law, the series $\frac{1}{4} + \frac{1}{2} + \frac{1}{8} + \frac{1}{16}$, etc. = 1; *i. e.*, it equals the total inheritance. Karl Pearson, a strong adherent of Galton's

¹ Science, N. S., 17: 1903: 75.

² Contrib, from the Zool, Lab., Harvard, 40: 1903; No. 4.

³ Address to Zoological Section, Brit. Assoc., Cambridge, 1904.

⁴ This is by no means constantly the case with this particular abnormality.

⁵ Proc. Manchester Lit. and Phil. Soc., 49: 1905.

⁶ Proc. Roy. Soc., London, 1897.

GALTON'S LAW

methods, from a study of other sets of features (the color of the eves in man and the color of the hair in horses-both, again, Mendelian or antagonistic properties), arrived at slightly different results, giving the grandparents and earlier generations somewhat greater influence. Without discussing the matter in detail, it may be pointed out that if we cross a Mendelian bastard (first generation) with the recessive parent (see Fig. 43, B), we get 50 per cent, of the offspring taking purely after that parent, the offspring being constant in feature, and 50 per cent. presenting the dominant feature, but truly bastards, this offspring dividing up according to the Mendelian formula. If the cross be made with the dominant elder, all the offspring are of the dominant type; 50 per cent, of them are pure dominants, the other 50 per cent, hybrids (Fig. 43, C). These figures fit in, it will be seen, with Galton's law. The law, so far as we can see, fits a restricted series of cases, and does not, it seems to us, lead to practical application in the same way as do the Mendelian principles.

C. Mosaic Inheritance.-In this form, strictly speaking, in certain cells and cell groups the paternal influence is dominant; in others of the same order, the maternal. Such is best seen in connection with surface coloration, where the parents have been of different color, the result being a streaking or dabbling with the two colors without blending of the same; we obtain the effect (distantly, it is true) of a mosaic work. The condition is on the whole infrequent, and particularly so in man. The only examples that we can recall which would seem to come under this category are those rare ones in which the eyes are of a different color-and, possibly, the still rarer condition, in which the internal sexual organs on the one side are male, on the other female. It is a form of inheritance that has been little studied, which nevertheless has to be recognized, and may, indeed, color effects apart, be much more widespread than is generally held; for, passing beyond individual organs or tissues, we must realize that the individual is apt in certain organs and tissues to reproduce the properties and peculiarities of the one parent; in others, those characteristic of the other, and this is but mosaic inheritance on a larger scale. Regarded thus, recognizing that biophores from both parents pass equally into all the cells of the body multiplying as the cells multiply, it is not difficult to imagine that where those are relatively equally balanced, in certain environments the one order will have the upper hand, in others the other order.

D. Sex Limited Inheritance.—More than one reference has been made in the preceding pages to the appearance in members of the one sex of characters peculiar to that sex and recessive or absent in the other. That they are most often recessive is suggested (1) by the fact that rudimentary evidence of those features or parts are to be recognized in this other set (e. g., the breasts of the male, the clitoris of the female), and (2) by the way in which certain peculiarities, which in themselves are neither primary nor secondary sexual features, and which in the mother are unrecognizable and clearly recessive, are conveyed to her male and not to her female offspring. It is simplest to

suppose that in the germ plasm there is an intimate association between the elements governing the development of these peculiarities and those governing the attributes of maleness, and that if the former be recessive the latter are recessive also. We have noted, for instance, that the daughter of a "bleeder" most often exhibits no tendency to hemophilia, but transmits that tendency to her son, that the gouty diathesis is liable to be transmitted to the sons and not to the daughters, although in the sons of those daughters it may again show itself. Nettleship has recorded striking examples of the same sex-limited inheritance in connection with color blindness. This is relatively common in the male (4 per cent.), rare in the female (less than 0.5 per cent.). In these rarer cases in which the female is affected, all her progeny are color blind, whereas of the children of the color-blind father and normal mother none are color blind, although the daughters of such a family transmit color blindness to their sons; the sons do not. If the genealogy of the Mampel family given as Plate IV be studied, it will be seen that with a rare exception hemophilia follows the same law. I may add that in striving to determine the laws of human descent these occasional exceptions, or contradictions to the general scheme, are the despair of the systematists. Possibly, they are due to the crossing of widely distinct strains. Whatever the cause, we encounter them in connection with the transmission of all forms of inheritance so that unexceptional demonstrations of the action of Mendel's principles in man are rare.

E. Atavism.—As originally employed, all reversionary conditions —the appearance in a given generation of traits not present in the parent but characteristic of previous generation—were included under the one term atavism (*atavus*, a grandfather). It is needful, however, to make a distinction between familial reversion, or atavism proper, and racial or ex-specie reversion, with the appearance of properties characteristic of an earlier stage in the phylogeny. We shall use the term atavism for the former, phylogenetic reversion for the latter.

Such atavistic inheritance—the inheritance by a child of properties not manifest in either parent, but present in the grandfather or some relatively recent ancestor—is seen to fall in with Mendel's law. That law and the observations on which it is based demonstrate that a condition may remain latent through several generations, to reappear eventually in a definite proportion of the members of a stock. We gain also an understanding of *collateral* inheritance, *i. e.*, of possession of properties corresponding with those of members of collateral branches of the family—uncles, aunts, or distant cousins, and unlike those of the parents.

So, also, looking to the future, we see that all the progeny of an individual exhibiting one or other inherited taint need not of necessity exhibit that taint, although some are likely to do so, while, if both parents vary in the same direction, the probability becomes very great. Further, if either parent come from an unsound stock, though not individually presenting the stigmata of the family taint, these may reap-

GALTON'S LAW

pear in subsequent generations, being merely recessive. This may seem, and is, vague. It may be possible, when these principles become more firmly established and family histories more fully recorded, to state the probabilities of the inheritance of one or other condition.

F. Reversionary Inheritance.—A clear distinction is to be drawn between this and *alavistic inheritance*. In the latter we have the manifestation of properties present potentially in the germ plasm, though unable, owing to certain conditions, to manifest themselves fully in the body of the individual parent, while able to do this in the body of the offspring; those properties may be either favorable or unfavorable. In reversionary inheritance, we have *always* a return in the offspring to a *lower type*—a development which is incomplete, not reaching the standard of the type, but only attaining to a stage characteristic of an earlier period in the development of the species, whether affecting the body as a whole or more especially some particular system, such as the nervous system.

It is by no means easy in a large proportion of cases to determine whether reversionary traits present in an individual are truly inherited or merely acquired. A child is born microcephalic, for example, and with simian physiognomy, or possesses indications of persistent gill clefts, recalling earlier stages of evolution. Arrests of development of these orders may be due to no primary defect of the parental germ plasm, but to disturbance affecting the foctus *in utero*. Nor can we always determine with precision which order of events we have to deal with. Yet in certain cases we can have no doubt that we deal with undoubted reversion, brought about either by defect in the germ plasm of one or both parents or by the interaction of dissimilar germ plasms.

Darwin's Experiment.-The classical example of the latter method of inducing reversion is afforded by Charles Darwin's¹ well-known observations upon the effects of crossing a barb-fantail female pigeon with a barb-spot male, from which cross there developed a "bird hardly distinguishable from the wild Shetland species" (of blue-rock pigeon, Columba livia). Columba livia is the ordinary wild pigeon, common over a large portion of the northern hemisphere; everything indicates that from it during the course of long centuries, and by artificial selection, the extraordinarily divergent varieties of tame pigeons have been developed. Similar results have more recently been obtained by Ewart.² Crossing an absolutely white fantail, with thirty feathers in its tail, with an owl-archangel hybrid (the "owl" a powdered blue pigeon, with short beak, the "archangel" copper-colored, with well-developed crest), he obtained a bird almost identical in measurements with the bluerock, while in color and markings it showed complete reversion to the checkered blue-rock of India, and, like that, had only twelve tailfeathers. History indicates that the cult and culture of the domestic pigeon began in the East, and, like the crow-and like man-the wild pigeon shows variation in different regions of the earth's surface.

¹ Animals and Plants under Domestication, 1:204.

² The Penycuik Experiments, 1899:26.

Here, obviously, by crossing widely separated varieties of a species, we obtain not the domination of the characteristics of one or the other, but a reversion to an earlier type. We occasionally encounter in man instances apparently of the same nature, in which parents, each of sound constitution and each well developed, but coming of widely distinct stocks, have a series of children who in bodily constitution and mental growth are distinctly of lower type.

Familial Degeneration .- More often, however, in man we encounter cases in which reversion is to be ascribed not to antagonism of the germ plasms, but to defect in the same, defect which, as we shall point out, is to be regarded as primarily brought about by toxic influences telling upon and modifying the constitution of the parental germ cells. Such are the class which nowadays, though we do not wholly like the term, it is the custom to describe as *degenerate*—the progeny of those leading vicious lives. The typical degenerate is of poor bodily development, brain smaller than the normal, with convolutions less abundant and less fully formed, of degraded physiognomy, little capacity for sustained attention or for prolonged thought, cunning rather than intelligent, deficient in moral sense-in all these points resembling the lower, less-developed races of our species. No one studying a well-marked example of this order can fail to be impressed by the reversion to a lower type. The fortunate tendency is for families of this type to deteriorate in successive generations, for the latter members of this heritage of misery to exhibit idiocy, non-viable children, monstrous births, stillbirths, and so on, so that by the third or fourth generations the sins of the father have told so surely that the stock dies out.

G. Diathetic Reversion. - What must be regarded as a slighter grade of the reversionary process is still more frequently encounteredconditions in which one or other system does not attain full development, and in which the incomplete development descends from one generation to the other. We have already laid down that where the environment is unfavorable, conditions of latest development are those most easily lost; those of oldest phylogenetic acquirement are most firmly retained. The widest and latest evolved distinction between man and the other animals is the development of the higher nervous centres in the former. It is noteworthy how relatively unstable and various in their development are these centres; no other tissues of the organism vary so greatly in their functional capacity. This capacity, it is true, cannot easily be measured, but the variation is well indicated by certain figures from Cambridge University.¹ There they afford the students the option of taking either the ordinary or the honors course, and the better-trained men, who select the latter, have the choice of no less than eight triposes, or honor examinations, in different subjects, from mathematics and theology to Indian languages and agriculture, to work at for three years. The choice is thus singularly wide. The most celebrated of these tripos courses is that in mathematics; it draws

¹ For these I am indebted to my friend John Greaves, Esq., M.A., Christ's College.

MENTAL CAPACITY

the best men from all over Great Britain. The questions given range over a wide area, and the papers are set and marked not by one, but by several mathematicians of distinction, the candidates being in each paper given a choice of more questions than it is possible to cover in the time.

The examination, we repeat, is not compulsory; all who enter have had a mathematical training and a *penchant* for the subject. The following figures give (1) the maximum marks obtainable were every question answered and every problem worked out on each of the eight days the examination lasts, the average being struck for two successive recent years; (2) the means of the mark obtained by the highest candidates (Senior Wranglers) in two successive years; (3) the mean mark of the two lowest candidates to obtain first class honors; (4) the mean mark of the "Wooden Spoon" in those two years, *i. e.*, of the last candidate to be granted a "pass,"

AVERAGE OF TWO ORDINARY YEARS, CAMBRIDGE MATHEMATICAL TRIPOS.

Total examination		ol	daximum marks btainable, 4873	Maximum obtained Senior Wrangler. 1896	Lowest Wrangler, 942	Wooden Spoon, 190
First four days only			1486	787	477	170

Mr. Greaves has afforded also the figures for the first four days only, as the papers on these days cover what may be termed ordinary mathematics and the ordinary man often has little knowledge of the advanced mathematics of the last four days. The mean "Wooden Spoon" secured only 20 marks in the second, advanced part of the tripos, out of a possible 3400 or so. Even in these routine mathematical subjects the first man secures close upon five times as many marks as the last, whereas in the total examination he scores nearly ten times as many. There is, indeed, the memory of a tripos in which the Senior Wrangler (now a member of His Majesty's Ministry) secured twice as many marks as the second on the list (now a university professor of world-wide reputation).

It is difficult to imagine the difference in the development of the associated centres for calculation, figures, expression, and writing between the Senior Wrangler and the "Wooden Spoon" (the lowest on the pass list)—and also between the latter and the born idiot.

While, here again, mental development may be arrested by intrauterine or postnatal influences, and mental power is capable of great development by proper training, it is abundantly evident that mental capacity in the main is a matter of inheritance. We possess much evidence that imperfections of the higher centres and mental instability of various grades are markedly liable to descend; cases, indeed, are on record in which certain grades have shown themselves in the same family for two centuries and more. Marriage with those of good mental state reduces the liability, intermarriage of members of the same neurotic family is seen notably to increase the liability. In some

cases the descent is homeomorphic; in others, heteromorphic. The former type would seem to indicate the inheritance of some particular anatomical imperfection, or the subjection of the members of successive generations of the weak-headed to like strains, producing like results; the latter is nowadays accepted as indicating a general imperfection of development of the higher centres. For some years, so long as they are not subjected to strain, these reversionary individuals may show little departure from normal; subjected to some particular strain—alcohol, syphilis, or other infectious disease, anxiety, religious emotion, or intense mental activity of other nature—and the brain gives way in one or other direction, according as the strain falls on one or other centre. Thus, the mother may have religious mania, the son become an epileptic or victim of general paralysis, and his son be an imbecile.

Yet another recent and special acquirement of man, as distinct from nearly related animals, is his power of resistance against, and of susceptibility toward, sundry infectious diseases. These powers also vary considerably, and, like mental capacity, and deficiency in the same, are largely a matter of inheritance. Indeed, a lack of resistance toward infection frequently accompanies mental weakness-as it accompanies the more pronounced reversionary degenerations-rendering the children weakly and liable easily to succumb to childish ailments. Just as we noted that certain neuroses are distinctly inherited, so do we observe specific inheritance of susceptibility to one or another infectious agency. We have called attention to this in the different races of mankind; we observe it also in particular families: this family is particularly susceptible to tuberculosis, that to scarlet fever. What is still a matter of debate is whether an infectious disease affecting an individual under any conditions renders the offspring either more or less susceptible to that disease.

H. Cumulative Inheritance.—It occasionally happens that the blend, instead of showing a given property to an extent intermediate between what the two parents exhibit, shows this more pronounced than does either parent; there is, as it were, a summation rather than a mean product. Examples of this order are not frequent, nevertheless they occur. Thus, Mendel noted that the hybrids of certain peas, short-stemmed and long-stemmed, respectively, developed constantly a much greater length of stem than either parent form, or than the pure progeny of the longer stemmed subspecies. Nor can the cases properly belonging to this class be ascribed to ataxism. The conditions leading to cumulative inheritance are, so far, wholly undetermined, but its existence throws some light upon the conditions next to be studied.

I. Spontaneous Variations; Mutation.—So far we have had to deal with conditions present in the ancestors and conveyed to the individual, or, failing to be so conveyed; there is yet another condition to be taken into consideration, namely, the appearances in the offspring of conditions and relationships which are new to the stock. These we speak of as *spontaneous variations*. Of such, numerous examples in

SPONTANEOUS VARIATION OR MUTATION

animals and plants have been collected together by Bateson;¹ the most easily grasped examples of the conditions occur in flowering plants. The tulip, for example, is a plant having a three-partite arrangement of organs, well seen in the flower. Now, occasionally in a bed or field of these plants, flowers are to be encountered showing a four- or five-partite arrangement. The trefoil, or shanrock, and the clover have tripartite leaves; careful search in certain localities frequently affords leaves which are quadripartite, five-partite, and, very rarely, six-partite. Everything is opposed to the belief that the Liliacee, to which the tulip belongs, are descended from a four- or five-partite ancestry, or that the shanrock had originally cruciform leaves. Or, otherwise, in this and numerous similar instances, reversion is incapable of affording an explanation.

The frequent existence of these spontaneous variations has not been sufficiently recognized in man. By hook or by crook, anomalies of excess, supernumerary fingers, toes, vertebræ, ribs, teeth, breasts, hair, etc., have one and all been attributed to reversion or atavism, so-called, no matter how far back in, or how far off, the probable line of descent, it is necessary to travel to find a form possessed of the condition—of seven digits, for example. Thus, taking the case of supernumerary mammary glands: In mammals possessing several pairs, these are arranged in a row on either side of the ventral median line, either over the abdomen alone or stretching well over the thorax. The number of pairs is in relationship to the number of young brought forth—from one pair in man to six, seven, or even eight pairs in the sow.

Now in man it is not so very rare to have the presence of one or more accessory mamma—generally so small as to attract little notice, and less inconvenience, but sometimes large and well developed. These are situated, whether paired or unpaired, at some point or points along converging lines stretching, roughly, from the region of the axillary angles toward the symphysis pubis. Their position along these lines strongly suggests reversion toward the condition of a many-breasted ancestry. But, in the first place, what positive knowledge have we of such many-breasted ancestors in the direct line of human descent; and, in the second, how are we to account for those cases in which the supernumerary mamma is situated—as it may be—on the back of the shoulder, or the buttock—regions never the site of mammae in the lower animals?

And why, to continue this line of argument, do we not find the greatest number of mamma in the very lowest mammalian forms? The ornithorhynchus possesses only one pair. Atavism will not explain the progressive increase in number as we proceed upward along the line of ascent of certain species of mammal. Spontaneous variation, with subsequent inheritance of a favorable variation, must have occurred in them

¹ Materials for the Study of Variation, London, Maemillan, 1894.

² For literature on supernumerary mamme, see Bateson, *loc. cit.*, p. 81; Leichenstern, Virch, Arch., 73; 1878; 222; Puech, *Les Mammelles et leur anomalies*, Paris, 1876; and Roger Williams, J. Anet. and Physiol., 25; 1891; 225.

to produce the result, and if it has occurred in the one order of cases, it is simpler to invoke it to explain the other set, and this the more easily when we recall that the mammary glands are not totally new organs, but, strictly speaking, are collections of hypertrophied and modified sebaceous (or, some would say, sudoriparous) glands, of glands scattered all over the mammalian integument. It is true that more than one embryologist has called attention to the existence of a ridge the mammary ridge—extending bilaterally in the focus in the position of the mammary line already referred to; but in no mammal do mammae present themselves normally along the whole length of this ridge, and when variations do occur the existence of this ridge does not make them any less spontaneous.

Regarded thus, the more we consider the subject the greater appear to be the number of cases of spontaneous variation.

It may, indeed, in general be laid down that when a condition is progressive, or when, on the contrary, it appears to be regressive, but is isolated and unaccompanied by other indications of regression and incomplete development, then the indications are that we deal with spontaneous variation rather than with reversion. We would thus equally include in this class of cases of polydactylism and bradydactylism (shortness of certain phalanges), supernumerary vertebrae, as well as reduction in the number of the same, supernumerary and deficient ribs.

Once these occur, they have a tendency to be inherited. We have already referred to the case of inheritance of polydactylism.

Closely allied, and apparently due to the action of the same causes, are certain conditions of defect, which nevertheless cannot be attributed to reversion; they reproduce no stage in the previous history of the race, and these, too, are singularly linble to be inherited; such are articular laxity (with liability to specianeous dislocation of various joints), ichthyosis (with peculiar indication of the growth of the epidermis, so that the cells produced abundantly do not scale off, but accumulate to form thick masses or scales over the body), hemophilia (with its peculiar instability of the vascular system, rendering hemorrhage liable to occur upon the slightest provocation, and with inheritance not direct, but through the female who herself does not show the affection). Possibly in this class is to be included Daltonism, or color blindness, in which, while we have no evidence that the formation of the eve is modified, the individual is unable to recognize certain colors. This as noted on p. 175 would seem to be recessive in the female, dominant in the male, while *albinism*, another condition of this order, would appear to be purely recessive (see p. 174). So, also, several observers have noted that certain tumor formations, such as the development of multiple lipomas (fatty tumors), enchondromas (cartilaginous), and exostoses all tend to be inherited. All tumors proper, it may be noted, are progressive in type; as to how far they are to be regarded as primarily the result of spontaneous variation will be discussed later.

This may be noted, that the spontaneous variation, even when liable to be inherited, is not necessarily useful. There is some evidence that

MUTATIONS

if its development be correlated to that of some useful property, a useless or apparently useless, variation may become a specific character. Thus, Bland-Sutton' calls attention to the callosity on the inner side of the foreleg of the horse as a probable example of correlated inheritance. That area of cornification of the epithelium is not possessed by other species, and its existence, savc on this supposition, is an enigma.

Of late years the distinguished Belgian botanist de Vries² has thrown considerable light upon the appearance of this spontaneous variation. Cultivating the plant *Enothera lamarchiana* (one of the evening primroses) for some fifteen years, he noted the appearance from time to time of individuals which definitely varied from the parents. These appeared suddenly in growths of large numbers of young plants which, as a mass, did not depart from the parent stock; and, what is more, these "mutations," as he terms them, were true to seed. Thus in 1895-to quote an instance—there appeared the relatively huge *Enothera gigas*. There had been no gradual variation leading up to it; the appearance was sudden, and, subjected to self-fertilization, this single plant afforded seeds giving origin to several hundreds of the gigas type. Here, at a bound, a new species was seen to develop, and de Vries lays down very decidedly that new species are not brought about by the accumulation of small individual variations, but in nature and during the course of the artificial culture of plants there appear occasionally, if not periodically, individuals manifesting changes so pronounced from the beginning, and these so distinctly heritable, that one has to regard these individuals and their descendants as constituting a new species. Similarly, de Vilmorin^a points out that species which in the wild state are remarkably stable, are apt to become highly variable after several generations of culture. Under these conditions, Klebs⁴ has obtained mutants among the seedlings from Veronica chamaedrys. De Vries holds that in all cases evolution is of this discontinuous type. These mutations, it will be seen, are what we have considered as spontaneous variations. To quote Jacques Loeb:5 "If the determinants are comparable to a series of compounds, e. g., of alcohols, there is no more a transition possible between two species separated by a difference in only one determinant than there is a transition possible between two neighboring alcohols of the same series."

It will be seen from the following chapter that with this view of Loeb we must largely agree; although Boulenger's case, quoted on p. 172, would indicate that the difference between varieties and species is merely one of degree. There may, that is, in the terms of our theory, be modifications in the constitution of the biophore so slight that in fertilization the modified and the unmodified do not interfere, the one becoming

¹ Introduction to General Pathology, London, 1886; 166.

² Die Mutationstheorie, Versuche u. Beobachtungen, etc., Leipzig, 1; 1901.

³ Notices sur l'amélioration des plantes, Paris, 1886: 36.

⁴Künstliche Metamorphosen, Stuttgart, 1906: 152.

⁴ The Dynamics of Living Matter, New York, Macmillan, 1906: 225.

dominant; or, on the other hand, modifications so considerable as to bring about interference and sterility.

What is the cause of these spontaneous variations must be approached with caution. The fact already noted, that crossing of unlike races occasionally leads with certainty to offspring possessing a given property so developed as to suggest that the development represents the sum of, rather than the mean between, the individual possession of the two parents, would indicate that amphimixis is immediately concerned. Nevertheless, recent botanical observations would suggest that influences brought to bear upon the parent and its germ plasm are, in some cases at least, the prime cause. Thus, Macdougal¹ notes that, taking Raimannia odorata, another member of the evening primrose family, and subjecting its ovules to 10 per cent. sugar solutions and solutions of calcium nitrate (1 to 2000), definite mutants were obtained; several individuals developed from the seed were of a type wholly different from any previously seen-glabrous and not hairy and with altered shape of the leaves-and their appearance could only have a direct relation to the operation. By like methods he obtained also a mutant of Enothera biennis. These mutants bred true to the third generation. Similarly, Tower² reports that subjecting certain beetles to intense environmental change (cold and humidity) immediately before the maturation of the germ cells, ova, and sperm, he has obtained pronounced mutants. These mutants breed true and do not segregate the characters in subsequent generations.

¹ Report Dept. of Botan. Research, Fifth Year-book, Carnegie Institute, Washington, 1907; 119.

² Investigation of Evolution in Chrysomelid Beetles of the Genus Leptinotarsa, Carnegie Institute Publications, No. 48: 1906.

CHAPTER XV.

INHERITANCE-(Continued).

THE THEORY OF INHERITANCE.

This enumeration of the various forms of inheritance is apt to leave the impression that the possibilities are so varied and so haphazard that it is a hopeless matter from such data to construct any theory of inheritance capable of application to all or nearly all the cases. Certain facts, however, stand out, and utilizing these, we may advance to a certain extent:

1. In the first place, it is evident that, although considering any one feature, it may happen that that feature does not present itself in the immediate offspring, nevertheless the whole modern study of heredity proves convincingly that where the individual is the offspring of two members of the same species, each parent affords equivalent contributions to the offspring. These equivalent contributions of heritable material may, it is true, in one or other respect not be of equal potency; but there they are, and, contributed to the germ cells of that offspring, they may demonstrate their existence in the individuals developed from these germ cells.

This law, if we may so term it, is correlated with and evidently based upon the fact that in conjugation each parental germ cell supplies a like contribution of nuclear matter to the primordial cell of the new individual; half the chromosomes are of paternal, half of maternal, origin. No other conclusion is possible than that the heritable material resides in these chromosomes.

2. If this be so, the different forms of inheritance must be related to the properties of these chromosomes and to their interaction. Our theory of inheritance must, therefore, be essentially one which deals with the chromosomes and their constituents.

3. We have already laid down that the primordial living matter of the cell is contained in the nucleus; it is this matter that must be carried over in the chromosomes. From this it follows that our theory must be expressed in terms of the biophoric molecules, and that we have to endeavor to conceive a constitution of and mode of interaction between these biophores from the two parental germ cells which will satisfy the various conditions.

4. Coming now to analyze the different forms of inheritance, we make out that a particular feature showing itself in either parent may:

- (.4) Present itself also in the offspring:
 - Dominant, wholly replacing the corresponding but divergent feature seen in the other parent.
 - Blended, this particular feature in the offspring being intermediate in character between that exhibited in the two parents.
 - In mosaic form, in certain cells the paternal, in others the maternal, feature being dominant.
 - Blended and excessive, the feature being more pronounced than in either parent.
- (B) Be unrecognizable in the offspring:
 - Recessive, and replaced by corresponding feature derived from the other parent, but as such latent, capable of reappearing in later generations.
 - Absent, wholly wanting in subsequent generations, the absence being due either:
 - (a) To casting out of an inherited condition, or
 - (b) To the feature seen in the parent being an acquirement and not an inheritance.

5. Or, on the other hand, considering the individual, we note that as regards any particular feature or group of features, there may be:

- (.4) Normal Inheritance: The offspring not being in this respect advanced beyond either parent, but at the same time not fallen behind.
- (B) Progressive Inheritance: The offspring being advanced beyond the more advanced of the two parents and exhibiting either:
 - Excessive development of the condition or conditions already observable in one or both parents, or
 - Spontaneous variation (mutation), i. e., the appearance of conditions not previously noted in either parent or either parental stock.
- (C) Retrogressive or Reversionary Inheritance: The offspring reverting as regards any feature or group of features to a lower stage in the phylogeny of the species.⁴

(D) Non-inheritance: Apparent or actual.

From this analysis one thing at least is obvious, namely, that the biophores derived from either parent are liable to retain their identity for some generations. Or, to be more accurate, that qualities conveyed by the parental biophores may be retained even if in a recessive, latent condition.

That, indeed, is clearly proved by the Mendelian studies on hybridization: after six generations or more with self-fertilization the hybrid

¹ We are acquainted with no satisfactory evidence that this class should be redivided into *recessive* and *complete*. On the contrary, it is the experience of pigeon breeders that once, by cross-breeding, the old blue color and characters of the wild pigeon reassert themselves it is hopeless to use such birds for breeding purposes. Even when mated with birds of pure breed and dominant type it takes long to east out the signs of such reversion,

PROGRESSIVE VARIATION

can give origin to plants exhibiting the pure features of either the dominant or recessive ancestor. Conjugation cannot, therefore, be of the nature of a chemical union of the biophores from the two sources with resultant formation of a new biophoric substance. On the other hand, we cannot conclude that all the separate biophores contributed by and representing each ancestor are potentially present in the fertilized ovum. This would demand an infinite number. The existence of determinants such as Weismann conceived is, as we have pointed out, a physical impossibility, and this is equally so; were ten generations represented, that would demand the presence in each chromosome of more than one thousand separate orders of biophores.

Our way, then, lies between Scylla and Charybdis. Still, between those two the cautious mariner could advance his craft and, the gods helping, could achieve through the straits. And here we would urge that our conception of the constitution of the biophore affords us a proper equipment to achieve the passage. We have, it will be remembered, been led onward to regard the biophoric molecule as composed of a central body or ring of nuclei provided with side-chains which are dissociated with greater ease. As the environment has been modified, so have the side-chains undergone modification, and as these side-chains become utilized in the polymerization of the biophoric matter and the formation of new biophores, so has there been a progressive increase in complexity of the biophoric molecule. We have pointed out how, neglecting determinants, we must regard the biophores in the somatic cells as undergoing extensive modification when their environment has become altered, whereby they have given rise to or controlled the different orders of cells in the different tissues (p. 136). As regards the germ cells, their biophores must similarly be influenced, for it is upon their modification that the whole evolution of living forms has depended. Clearly the biophores of the human ovum are vastly more complicated than those of the amœba, or, again, than those of the lowest multicellular organism of the line of man's ascent, and yet the progressive elaboration of the soma or body throughout the course of the ascent has been the outcome of the germ plasm and the biophores of the same within ovary and testis.

6. There are two, or three, possible causes for the progressive variations of multicellular organisms: the mingling of germ plasms in conjugation (amphimixis), the effect of environment on the respective germ plasms, and the effect of both of these combined. The first of these was strenuously upheld for long by Weismann as the controlling cause, but he was compelled to admit that the second must also be in action. Here also it must be pointed out that Mendelism pure and simple is inadequate to throw light upon progressive evolution. It establishes the number and characters of individuals endowed with different properties capable of being produced by the amphimixis of germ cells possessing a given number of differing attributes; it accepts the divergent qualities of the germ plasms as already in existence without seeking to explain their origin; it deals with permutations, not with mutations. In regard to this second cause, we have demonstrated that it is clearly


in action in unicellular organisms which do not conjugate, as also in the somatic cells of the highest multicellular forms of life (p. 121); it is illogical to deny its action upon the germ cells of the same. Not to waste time by taking part in what has been an angry discussion, we are prepared to accept the third course—to admit that both the action of external agencies and amphimixis are factors in variation, retrogressive as well as progressive.

7. Granting this, and admitting that through the action of both causes it comes to pass that the germinal biophores in no two members of the same species are absolutely alike in constitution, what must we conceive to be their action upon each other when, through conjugation, biophores of two orders come together in the same cell, the fertilized orum?

The facts of inheritance, and what we know regarding its histological basis, entirely refute the hypothesis that the biophoric molecules as a whole undergo chemical union. We may, however, conceive these, in the first place, as lying side by side in a common cytoplasm or, to be, more exact, nuclear sap, in the process of assimilation attracting ions from the surrounding medium, building these up into side-chains of different orders. Of these side-chains, some of them are identical-common, that is, to the molecules of both sets of biophores-some, on the other hand, of unlike constitution, so that certain side-chains having corresponding position or attachments in the two sets of parental biophores are dissimilar. As demonstrated by studies upon immunity, we regard such side-chains as detachable and apt to be detached, that is, to be developed in excess, and then, becoming loose, passing into the surrounding cytoplasm. Again, as we have pointed out (p. 98), we must regard growth and increase in the number of biophores as brought about, in the first instance, by the building up of nuclei of side-chain matter, this matter attracting other matter in due order, so that gradually new rings are constituted-new biophores. If these views be correct, then, when molecules of closely allied constitution and properties are growing side by side, what is there in this process to determine that side-chain matter. which has been liberated under the influence of the one set of biophores and has become detached, does not become attracted to and built up into the substance of the "growing" biophores of the other set? I cannot but hold that under these conditions—that is, conditions under which we have compound molecules of very similar structure becoming built up side by side-this must inevitably occur in a common fluid medium. Whenever a greater affinity exists between the components of one growing biophore and certain side-chain nuclei developed under the influence of the molecules of the other set of biophores, then these nuclei will be apt to be built into, to become an integral part of, the new biophores, to the exclusion of the corresponding nuclei-those proper to the original molecules. In short, there will be, physically speaking, a contest between the two orders of growing biophores and, to a certain degree, a selection or rearrangement of constituent nuclei. This rearrangement in the simplest case will result in an interchange of constituent

INTERACTION OF PARENTAL BIOPHORES

parts; in other cases, may result in side-chain material derived from one parental biophore, and possessing powerful affinities to the growing biophores of both orders, becoming built up into both sets, to the exclusion of corresponding but weaker side-chains (so that these become wholly cast out), and with this the properties determined by their presence disappear in the next generation. In other cases, again, we can premise an interaction between certain side-chain groups derived from the two parental biophores, the resultants of this interaction becoming built up into the growing biophores, this interaction having as a result either an exaltation or a depression of parental character, or, again, leading to the production of mutation.



Schema of mode of interaction of two biophoric molecules in a common cell sap: A, of maternal; B, of paternal origin. 1, 2, 3, alledomorphic side-chains, which, when liberated into the cell sap, will be attracted to the biophore exercising the strongest affinity; 4, side-chains common to both molecules, built up indifferently into either.

Granted, that is, that in its broad lines we have come to realize the mode of constitution of the proteidogenous molecule, that we are justified in assuming that the biophoric or living molecules partake of similar constitution, and that our conception of growth is that which must be accepted, then under these conditions growth, in a common medium, of biophoric molecules of two orders, alke in general constitution but differing in certain of their component chemical nuclei, must result in a certain amount of interchange of those nuclei. Two sets of biophores may still be traced in the blastomeres, in germ cells, and other cells derived from the fertilized ovum; two sets each derived by direct physical descent from the original paternal and maternal biophores and chromosomes, respectively; but the members of each of these while building up into their structure material assimilated by their legitimate progenitors,

attract for purposes of growth allelomorphic1 matter formed similarly by the other. By this method, apart wholly from what may be regarded as external influence acting upon the germ cells during their existence within the organism of the individual, it must come to pass that through conjugation the biophores giving rise to a new individual are not identical with those of either parent, and that each comes to lose certain properties. which belonged to the biophores of the one, and gain some belonging to the biophores of the other. If this be so, then we can picture that in the process of reduction and casting out of biophoric material in the development of the oocyte and of the spermatocyte, while there are delivered to the orum molecules of living matter which in direct descent have been derived from one parent only, those molecules may convey to the ovum constituents and properties which have been derived from both parents. In this way, without any increase in the number of determinants or ids, by this chemical modification of biophores, a constant number of such biophoric molecules may become the bearers of properties derived from a long series of anecstors.

We purposely do not here consider all the different type of inheritance, for this is not a full treatise on the subject. We have taken up forms that are sufficiently wide apart to show that this biophoric theory is capable of elucidating their occurrence. It appears to us to have the great advantage of explaining how hereditary characters may be conveyed through a relatively small number of molecules of highly complex organization; how those molecules can in the course of amplimixis undergo modification through interaction; how they can become modified through the action both of amplimixis and of environment; how similarly they may undergo retrogressive changes and lose certain properties under the same influences.

S. With reference to the action of environment on the germinal biophores, it is still necessary that something be said, but our treatment of the subject of amphimixis will not be complete without reference to the remarkable reduction process which precedes fertilization. The mode of that reduction we have already described (p. 148 et seq.). We have seen that in the process of maturation of the ovum representatives of one-half of the chromosomes of the parental individual are cast out, and that similarly the spermatozoon contains only one-half of the chromosomes proper to the male parent. As shown by the abundant recent studies on Mendelism, the results of this reduction may be very remarkable; certain properties may at a single conjugation be thrown out so completely that they do not reappear in subsequent generations.

During the very first process of reduction in a hybrid a property or properties derived from the one parent may thus be thrown out; and yet when the parents had differed in several particulars, at this same moment properties derived from the other parent may likewise disappear.

¹ Bateson employs this term in connection with Mendelism, to indicate the corresponding or antagonistic property, either dominant or recessive, the two allelomorphs forming a "pair."

INTERACTION OF PARENTAL BIOPHORES


And as in such hybridization there may be as many as a score of properties in which the two parents had been contrasted—size, color of flower, position of flowers, shape of leaf, hairiness of leaves, shape of seed, etc.—the process of sorting prior to this casting out, if we regard these qualities as conveyed by distinct ids or determinantis, is beyond conception. It demands so exact a localization in each chromosome of the particular determinants, and at the same time so precise a distri-

10

18

1

ts



Schema to illustrate Mendel's law regarding the second hybrid generation as regards a single pair of features; as also to illustrate the effects of reduction of the chromosomes in oogenesis and spermatogenesis.

Each germ cell (first row) is originally provided with chromosomes of paternal (black) and of maternal origin (white). The existence of the law demands that in the process of reduction the ovum and the spermatozoon (second row) become provided with chromosomes (and biophores) that are of either paternal or of maternal descent, but not of both; although, as above noted, the biophores may in their growth and development have attracted side-chains formed primarily by the opposed order of biophores, to the exclusion of those originally belonging to them.

bution of the determinants for the various properties, that by no possible means have we been able to visualize what is supposed to happen. By the biophoric concept this casting-out process is, we think, comprehensible, namely, as already stated, we can imagine that during the sojourn together of the parental biophores in the germ cells of the new individual, from the moment of fusion of the parental germ plasms to give rise to that individual up to the maturation of his or her germ cells, there is an interaction and interchange between the side-chains to whose

presence is due these contrasted features, and this of such a nature that the newly developed biophores, descended, let us say, from the biophores of the female parent, have not the identical composition of those parental In the process of growth and formation there has been, biophores. as it were, a selective process. Owing to greater affinities, they have attracted and built unto themselves certain side-chains derived from the paternal biophores, and from merely attracting them in the first place, have come to form them actively. According to our conception, that is, a side-chain, to whatever central ring it is attached, tends to attract ions and radicals of a particular order to itself, so as to reproduce itself in series. This interchange depending on chemical affinities will not be universal, affecting all the side-chains of both paternal and maternal biophores; the newly formed biophores will present an admixture of the two orders; they will occupy definite positions in the nuclear thread and in the chromosomes derived from that thread.

Thus, it will happen that in the process of reduction, as indicated by the studies upon hybridization, the maturing ovum, or the spermatozoon, may come to contain biophores purely of paternal or purely of maternal origin.⁴ The accompanying diagram indicates what we conceive to be the process (Fig. 45).

Along these lines we believe it is possible to conceive the conveyance of a limited number of biophores in the germ cells from generation to generation, those biophores under favorable conditions gaining through amphimixis accretions to their properties, under unfavorable conditions becoming shorn of certain properties, and as a result the individuals developing from these germ cells may show either progressive evolution or devolution. To apply these considerations to the facts of hybridzation, etc., and thereby exemplify the mode of action of Mendel's law, would be altogether beyond the scope of the present work.

THE INHERITANCE OF ACQUIRED CHARACTER.

The above considerations upon amphimixis and its influence in causing the offspring to vary from either parent accept tacidy, it will be seen, the fact that there is variation between the two germ cells which enter into conjugation, but throw no light upon the primary cause of that variation. It is impossible to arrive at any other conclusion than that variation originates primarily in the action of modified environment upon the labile bioplasm. Nay, more, as we shall have to point out, such action of environment upon the germ cells during the course of their existence in the parent cannot be regarded as non-existent, though there are those

¹ So far as we can see, there are no indications that a given germ cell contains, for example, three-quarters of the grand paternal and one-quarter of the grand maternal. The rule appears to be that there is exclusive representation or it may also be equal, the one series lying latent; although there are difficulties in connection with this latter conception. This in itself indicates that the number of biophores gaining entrance is relatively small.

THE INHERITANCE OF ACQUIRED CHARACTER

who deny it; for upon its existence hangs the solution of the question whether any order of characters acquired by the parents in the course of their life can be conveyed to the offspring, and we cannot close our treatment of heredity without taking side in this ancient controversy. To what extent, if any, can acquired characters be inherited?

Before answering this it will be well to classify the characters which may be acquired; first, we divide them into the progressive acquirements and the regressive. Among the former come the increased use of parts, with improved functional activity of the same, swifter response to reflex or other nervous stimuli, and to these we must add acquired immunity to disease. Among the latter, mutilations and loss of parts; arrested development of parts, and abnormalities brought about by disturbances during development, whether the influence causing these have told upon the organism during intrauterine existence, or after birth during the period of postnatal growth; atrophy of tissues through disease, both in childhood and more particularly during adult life; retrogressive changes in the tissues brought about by disease, or, more broadly, by intoxications of various orders. For we recognize more and more clearly, not merely that bacteria and the larger parasites produce their deleterious effects upon the organism at large almost entirely through the agency of the toxins which they elaborate, but also that disturbances of very many orders which lead to continued depressed or perverted function of one organ lead thereby to either heaping up in the system of the deleterious substances which should be acted upon by that organ, or to modified internal secretions of the same, and so, secondarily, to poisoning of other tissues. These subjects will be dealt with more fully in the subsequent chapters upon Intoxication, Infection, and the Internal Secretions. Lastly, there may here, under protest, be included the legendary maternal impressions, because these are popularly and loosely held to come under the heading of acquired conditions.

1. Maternal Impressions.-We will deal with these first. If a mother while bearing her child has been frightened by a toad jumping toward her unexpectedly, and subsequently brings forth an anencephalous, toad-like monster, it must, imprimis, be advanced that the mother has not acquired that toad. None of the cases on record (and American medical literature a few decades back abounded in them) are in any sense instances of acquirement of a condition by the parent which is reproduced in the child. In the second place, if now the nervous theory be adopted and it be urged that a pronounced shock or stimulus referred by the parent to one area of her person is reproduced in a like area on the person of a child, it has to be pointed out that the child in utero is a separate individual, unconnected with the maternal nervous system; and, thirdly, with the rarest exceptions the fright or profound influence noted by the mother are stated to occur in the later months of pregnancy, when the different organs and parts of the foctus are already not merely laid down, but advanced in development, and we know that monstrosities and abnormalities date most often from the very earliest period of fortal

13

ores

een.

ave

first

ion.

ract

the

he

ree

gh

w,

n.

1P

life. All these tales are at the most examples of coincidence, where they are not the bizarre product of the female imagination.⁴

2. Use Acquirements.—Of acquirements in the strict sense of the word there is a complete lack of evidence that "use acquirements" are transmitted. The blacksmith's son has not larger biceps than has the ordinary individual, nor, with our knowledge of the relations of the germ cells to the rest of the organism, can we conceive why he should have. The utmost we can accept is that if the blacksmith has by exercise kept his system in excellent coordination, his germ cells will benefit thereby and his progeny be sound and generally well developed. But that one particular muscle or group of muscles should be picked out for progressive advance cannot be grasped. Confusion is here apt to arise between preëminent qualities that are inherited-the results, in the first place, of fortunate amphimixis-and such use acquirements. A great composer, for example, may have descendants with musicianly qualities above the normal: we have the Bach family, and, more recently, the three generations of Strauss, of waltz fame. In mental ability may be mentioned the Cecil and Sheridan families, the Darwin and Wedgwood group; but in no one of these cases have we the slightest evidence that the peculiar ability was acquired in the first place. More often than not the great genius has mediocre descendants; more often than this, none at all.

3. Acquired Immunity.- The evidence until recently has been definitely against this becoming inherited. The observations of Lustig² upon fowls rendered highly immune to abrin showed, indeed, indications of the reverse, the chickens in some cases being more susceptible to the special poison than were ordinary chickens of like age. More recently Conradi³ has reported upon the transmission of acquired immunity to rabies in the dog. He found that the offspring of a dog which had been immunized to this disease for three and one-half years, and a bitch immunized for five months showed very definite increased immunity, even to the most severe form of inoculation (intradural). Four of the six puppies survived doses of the virus which killed the controls in eight to ten days, and the two that succumbed did so only after lengthened periods of incubation. There have been not a few investigations along these lines with the infectious diseases, in general with negative or discordant results, or without adequate recognition of the conditions demanded for the proper carrying out of the experiments. Even here we deal with but one litter, and it might be objected that the immunity was of intra-uterine acquirement, from the blood of the immunized mother. To afford absolute proof, in mammals, of the transmission of acquired immunity it is essential to immunize the male parent alone, and that through a series of successive generations, and what is to be

¹ See McMurrich, The Physician and Surgeon (Ann Arbor, January, 1905), for an interesting study of the subject.

² Ctbl. f. Pathol., 15: 1904: 210.

^a Ctbl. f. Bakt., Abt. I, Originale, 46: 1908; 139.

THE INHERITANCE OF ACQUIRED CHARACTER

expected under these conditions is a Mendelian inheritance, certain of the progeny being immune, the others not. I would, however, recall here Ehrlich's convincing evidence that among protozoa—trypanosomes—the immunity to arsenic acquired by one generation becomes the property of successive generations, to indicate my firm belief that eventually a similar transmission will be found in higher animals. We shall, however, revert to this matter shortly.

4. Mutilations and Loss of Parts. It is a commonplace that the man who has lost an arm or a leg does not beget one-armed or one-legged children, but, on the contrary, offspring having the proper equipment of well-formed limbs. There is, it may be stated, no satisfactory recent case on record in which loss of a part by either parent has led to the offspring being minus that part. Weismann¹ cut the tails off successive generations of mice as soon as they were born; the twenty-second generation showed tails of perfect formation and normal length. We have the trite example of the Jews who have been circumcised religiously since the days of Abraham, and in whom the boy children have still to be circumcised.2 Here, again, as in connection with overdevelopment of one or other region of the body, it is not unlikely that the loss of a limb or important organ may have influence on the bodily health, and so tellin this case deleteriously-on the nutrition of the germ cells, but such influence must be general and not specific, leading to arrested development of one special organ or part in the embryo.

5. Arrested Development of Parts Due to Intra-uterine Disturbance.-We have been able to collect very little evidence under this heading, and the subject deserves fuller attention than has been given to it. Certain of these arrests are so extensive, as, for example, conditions of an encephaly due to amniotic pressure or adhesions, that life is arrested, and others not so severe must undoubtedly lead to general malnutrition, which must tell in a general way upon the germ cells. The difficulty before us consists, as we shall have further to point out in discussing abnormalities, in determining in very many instances whether a given arrest is inherent, due to the imperfect constitution of the embryo, or of external causation, brought about by intra-uterine conditions. But so far as we can see, a local arrest of development of definitely introuterine causation is not inherited. The earliest disturbance of development at all consonant with continued life-namely, the production of monochorial twins (from separation of the first two blastomeres)-does not lead to the offspring also producing twins, nor did the Siamese twins produce other than normal offspring; and cases are on record of indi-

⁺ The Evolution Theory, 2: 1904: 66.

² It has been objected that a definite percentage of modern Jews are born with a short foreskin, naturally circumcised. But so are "Gentiles," and it has not been shown that the percentage is greater. As I have pointed out elsewhere, although in different regions of the world circumcision is a religious rite, it must have been originated primarily from the observation that in hot climates those "naturally circumcised" were at a distinct advantage.

they word

ordi-

lave. reby one e. of · the ieramed oup: the not ione pon the ntly / to t to ned mg ons itv red ne, be

an

viduals born without limbs, apparently from intra-uterine amputation, being the sires of well-formed families.

6. **Disuse Atrophy.**—The same considerations must apply to this order of cases, so far as regards any individual organ. As regards the organism as a whole, it is the experience of breeders that the greatest fertility is associated with moderate exercise; that lack of exercise plus obesity tends toward sterility, and it is noticeable that the hard-worked wife of the poor curate has her quiver full and overflowing, whereas the millionaire's wife is apt to be childless. Clearly, therefore, there is an interaction between the soma and the germ cells, but that this is specialized, that atrophy of the muscles from disease influences the musculature of the offspring, has not been demonstrated.

7. Retrogressive Changes in the Tissue Due to Diverse Intoxications.—We here encounter conditions which we are inclined to think must to a greater or less extent tell upon the offspring. In fact, we have evidence that they do, though we would hasten to add that the influence would seem to be limited. It is in investigating these conditions that the pathologist can perform yeoman service to the study of heredity. Unfortunately, so far heredity has had little interest for medical men in general, and so far the observations are few and far between; they are, however, steadily increasing in number.

Man himself is difficult to deal with. As already stated, conditions of disease and intoxication in the female must be ruled out, for the maternal influence tells not on the germ cells only, prior to fertilization, but upon the developing foctus; and with regard to the male parent again, in most instances so many other factors have to be taken into consideration that to arrive at a sure conclusion is almost hopeless. Thus, take the commonest intoxication of all—the alcoholic. The general belief and we regard it as well founded—is that the children of the sot are, as a body, of lowered intelligence and vitality,' with unstable self-control. It is, however, next to impossible to prove this statistically, and this because:

1. If the mother be sound, her influence may be dominant upon the offspring; we must expect that a certain proportion will be of average development.

2. It is next to impossible in the majority of cases to determine whether already there be not hereditary taint in the father's family, and if there be, that this began from abuse of alcohol in a past generation—to show, in short, that alcoholism is the primary acquired condition and not the accompaniment (as apparently it often is) of retrogressive changes.

 Alcoholism in the father, as a general rule, carries in its train home misery and poverty. The poor development of the children may largely be due to neglect and malnutrition.

¹ Thus, for example, Imbault (Thèse de Paris, 1901) found that of 100 tuberculous ehildren, 36 per cent, were the offspring of alcoholies, 41 of tuberculous parentage. He quotes Arrivé's observations on 1506 cases of meningitis in children, that this occurs twice as often in those of alcoholies as in those of tuberculous parentage.

THE INHERITANCE OF ACQUIRED CHARACTER

tion,

order nism ty is ends ' the ends ' the ire's tion that ' the ticanink uave ence ence ence eral, ver, ions

ion, ain, eratake ef

as a trol.

this

the

'age

iere ow,

ome gely

age.

¹ Vide Legrain, Compt. rend. Soc. de Biol., 10 S., 2: 1895; 563.

² L'Obstetrique, January, 1900:63.

³ Arch. d. Méd. Exp., 12:1900:782.

⁴ Ctbl. f. Pathol., 15: 1904: 210 and 756. Lustig, while pointing out that his series of observations prove that immunity is not conveyed, curiously enough draws no conclusions from the frequency with which he encountered monstrosities and abnormalities of various orders.

Like difficulties present themselves if we attempt to study the heritage of tuberculous parentage. With syphilis, again, the fact that the father may infect the mother, and the further possibility of latent infection of the mother-and it may also be of the embryo-make it difficult to arrive at sure conclusions. Nevertheless, the frequency of stillbirths, monstrous births, and abnormalities in the children of syphilities as compared with other children, as has been noted by several observers,¹ cannot be merely a coincidence, while Mott has produced interesting data upon inherited parasyphilis, and notably upon the liability of the children of syphilitics to suffer from early general paralysis, and Georghiu,² studying the histories of a series of monstrous births, found in almost all cases the history either of syphilis or again of some acute infection of either parent shortly preceding the period of conjugation. It is when we make direct observations upon the lower animals that we gain the surest indications of these effects of parental intoxication; and here some of the most instructive figures are those of Carrière³ upon guinea-pigs. He inoculated his guinea-pigs over a period of several months with various soluble products of the tubercle bacillus, making altogether thirty separate matings in the course of two years. His results may be summed up in the following table:

	S	tillborn.	Dying before 16th day.		Surviving.		Total
	No.	Per cent.	No.	Per cent.	No.	Per cent.	born.
Male and female both inoculated	13	52.0	7	28.0	5	20.0	25
Female alone inoculated	7	26.9	9	34.6	10	38.4	26
Male alone inoculated	$\tilde{5}$	16.6	3	10.0	22	73.0	30

As might be expected, the influence of the intoxication was found greatest when both parents were subjected to the inoculation; least when the male alone was treated. But in this last category, although there were ten matings, the average litter was only three, whereas the average litter of the healthy guinea-pig is between four and five, and of those born, 16.6 per cent. were born dead, and of the 22 who survived beyond the sixteenth day, 7 are described as weaklings. There can be no doubt from this series that a bacterial poison such as the products of the tubercle bacillus has a distinct action on the paternal germ plasm—as, indeed, on the female. Lustig's' figures for the results of inoculations of fowls with abrin give parallel results; and both observers found as the result that the offspring were less resistant (and not more resistant) to inoculations of the tubercle bacilli and of abrin than were control animals of the same age.

With these figures may be compared those of Constantin Paul¹ upon the effects upon the offspring of saturnine poisoning in men working in lead, the wives not being subjected to the same influence. Of 32 pregnancies in which the husband alone was exposed to lead, he recorded 12 abortions; while of the 20 children born living, 8 died in the first year, 4 in the second, 5 in the third. Thus the 32 pregnancies yielded only 3 children living beyond the third year. Nor is premature death the only result; as Roque has pointed out, and as has been noted by others. there is a painful frequency of idiocy, imbecility, and epilepsy in the children of workers in lead. The figures are remarkable, but notwithstanding that we have brought them forward upon several occasions no one has submitted evidence in contradiction, and such additional evidence as we have obtained is in the same direction. Lize's² figures with regard to the family histories of those exposed to the fumes of nitrate of mercury are of the same order. (We have tabulated the results.)

	Number of cases.	Number of preg- nancies.	Abortions, pre- mature labor, and stillbirths.	Surviving infants,	Remarks.
Mothers alone exposed . Father and mother ex-	3	7	4	3	
posed	2	14	5	9	Of these, only 3 survived fifth year.
Father alone exposed .	?	12	4	8	Of these, 3 died before fourth year. One alone vigorous.

We possess, thus, clear evidence that substances circulating in the blood of the parent are capable of influencing the germ cells, and this not merely temporarily. In Lustig's cases the bad effects were noted months after the abrin had ceased to be given. It may be-it has been-objected that these are not cases of conditions acquired by the parent being conveyed to the offspring; the poisoned parent does not himself become a monster prior to begetting a monstrous progeny. This is quite true. It has been pointed out, again, that in all these cases we have regressive changes; the progenv tend to revert to a lower stage-as though, in the terms of our theory, the effect of the toxin had been to remove certain of the more recently acquired side-chains. This also would seem to be the case. The all-important point, however, is the demonstration that the germ cells within the ovary and testis are not inert, incapable of being acted upon by the rest of the parental organism. If we can demonstrate that retrogressive changes are possible, then under like influences progressive changes are equally so; if side-chains can be

¹ Arch. gén. de Méd., 15: 1860: 513.

² Union Méd., 25: 13: 1862: 106.

THE INHERITANCE OF ACQUIRED CHARACTER

removed from the biophores, other side-chains would seem capable of being added; so that here we have the first clear light thrown upon the mechanism whereby alteration in the environment of the individual, by telling upon his soma, may either:

1. Tell coincidently upon the germ cells.

pon

sing

reg-

ear.

only

the

ers,

the

ons

é's2

nes

dy

- 53

le.

CP

0-

or

10

2. Tell indirectly upon the germ cells, the modified internal secretions of one or other organ in the blood and lymph adding to or subtracting from it substances capable of acting upon the germ cells.

If the latter is demonstrable, then we are able to state definitely that the body cells themselves through their acquired conditions influence the germinal biophores.

All this is but just beginning to be realized⁴—as it is that different toxins and nutritive substances proper in the general circulation may have a specific action on the germ, causing modification in one or other direction. We have, it is true, indications that the offspring of the tuberculous, syphilitic, and alcoholic parentage differ somewhat in their degenerative stigmata, but these differences have not been determined experimentally.

Further researches along these lines may show that the acquired disturbances of a given organ may, through the consequent presence of abnormal cell products in the blood, influence the biophores specifically, so that under the action of different poisons the like organ in the offspring does not show the identical disturbance, but nevertheless exhibits departure from the normal. We would suggest that it is along these lines that Brown-Séquard's² remarkable observations upon guinea-pigs gain their explanation; observations which in our opinion have never been satisfactorily disproved, which further have been confirmed by Obersteiner³ Romanes, and others. * Brown-Séquard found that by section of the sciatic and other nervous lesions in guinea-pigs he could render the parents epileptic, and that the young were liable also to epilepsy and other nervous disturbances. Obersteiner found likewise that, of 32 young, the offspring of guinea-pigs in which he had cut the sciatic nerve of one or both parents, 13 were healthy, 19 showed disturbances, 11 weakly, 3 paretic, more particularly in the lower extremities; 2 had epileptic fits on irritating what he refers to as epileptogenous zones,

¹ Thus, Walter Heape (Phil. Trans. Roy. Soc., B., 200; 1909; 271), from a very full study of the proportion of the sexes produced by white and colored people in Cuba, can only explain the variations in these proportions by conditions other than heredity —by nutrition and physical conditions affecting the vitality and life of the ovarian ova. He affords evidence that privation and unfavorable conditions of life are correlated with an excess of males, prosperity with an excess of females, and suggests that extraneous forces may affect other than sex qualities of the ovarian ova, concluding with the remark that these deductions regarding the influence of environment on the ovar in the ovary are, to his knowledge, new.

² Researches on Epilepsy. Boston, 1857; also various papers in Jour. de Physiologie de l'homme, 1 and 3; 1858 and 1860; and in Arch. de physiol. normale et path., 1 to 4; 1868 to 1872.

³ Med. Jahrb., 1875. We quote Dietrich, *Die Bedeutung d. Vererburg f. d. Pathologie*, Tübingen, 1902; 14.

CL. BERNARD'S EXPERIMENTS

and were also paretic, soon dying; 3 showed corneal opacities and ulcers ascribed to atrophy of the fifth nerve.

If rodents are paretic, it is in the lower extremities that the paresis is most apt to show itself; thus, no stress is to be laid upon the relationship between section of the sciatic nerve in the parent and paralytic manifestations in the hind limbs of the offspring. There remain two possibilities: either that the operation, setting up irritation of the higher centres, induced a general malnutrition in the parent whereby the germ cells suffered, and the nervous instability of the offspring was but the manifestation of imperfect general development; or, secondly, that the irritation of the higher centres, by modifying the internal secretion of the nerve cells, led to the presence in the blood of substances exerting a specific action upon the biophores, in consequence of which the nerve cells of the offspring were imperfectly developed. We shall not attempt to decide between these two possibilities, but they deserve mention. We would only repeat that this study of the problems of heredity in this direction is but in its infancy, and although it promises to yield most important results, results which will determine definitely the extent to which conditions acquired by the parent influence the offspring, nevertheless years of patient study are requisite before this particular field of pathology is adequately worked over. Lastly, we would add the caution that too much must not be expected. The germ cells in the ovary and testis are characterized by the long period-extending in man over many years-in which they lie latent and inert. While thus inert it is unlikely that they present very active metabolism. This very latency would seem in itself to be a preservative against parental disturbances exerting too extensive an influence upon the constitution of the contained biophores. Nevertheless, to maintain life some metabolism must proceed, and, as our examples must demonstrate, they can be influenced by parental conditions. We have, therefore, not a little confidence that results of the highest value are to be expected, results of the highest value to us as medical men, for they will establish the limits of morbid heredity, and will afford us a sure basis for determining how far the frailties of the father, or the misfortunes of the mother, affect the progeny.

ilcers

resis tionlytic two igher t the t the m of rting the not lises the we endhile tabittle IOW the

SECTION II.

THE CAUSES OF DISEASE.

CHAPTER I.

INTRODUCTORY.

EVERY departure from the normal, whether in the cell, the organ, or the system in general, is a pathological condition, provided that, as indicated in the opening chapter (p. 18), we recognize that the "normal" is not an absolutely fixed point, but is the expression for the limits between which the majority of the individuals of a given species will be found to group themselves as regards any particular attribute. Such pathological conditions must, it will be seen, be of two orders: either primarily due to some constitutional defect transmitted from the parent or parents (included with which we must place the effects of imperfect interaction in the fusion of the male and female elements at the moment of fertilization. Such effects, being associated with the actual constitution of the individual, are of internal origin, inherent and inherited); or, in the second place, they may be the result of some influence which first affects the individual after his genesis. Such conditions are of external origin and acquired. Morbid conditions, then, are to be classified into inherited and acquired.

THE USE OF THE TERM "INHERITANCE."

Much confusion has been and continues to be introduced with the discussion of the inheritance of disease, as into that of heredity in general, by a lax comprehension and use of terms. By many "inherited" and "congenital" are employed as though they were interchangeable; by others, as conveying distinct ideas; disturbances to the fœtus, for instance, and conditions of intra-uterine origin being by them regarded as congenital, but not as inherited. We hold that the latter is the correct, or at least the more satisfactory, usage, but, owing to this confusion, would recommend that the term congenital be employed as little as possible, and then with a clear indication of what it is intended to imply.

INTRODUCTORY

A still greater confusion is introduced owing to the vulgar error, fostered by the legal profession, of regarding the individual as beginning existence with the moment of birth, and not until then, so that everything happening before that moment is grouped into one category; everything after, to another. The chick, so to speak, is not a chick until it breaks open the egg shell; its status, from the moment it ceases to be the expansible condition of "new-laid" egg until it emerges from the shell, is not recognized in law. But a very little reflection suffices to convince us that the individual existence of the chicken began even before the egg was laid; and what is true of the chick is equally true of the human being. The individual begins to be the moment that fecundation is accomplished-the moment the nuclear material of the spermatozoon fuses with the nuclear material of the ovum and "these twain become one." Compared with this event, birth is of secondary importance. The intra-uterine association of the embryo and foetus' with the maternal tissues is but one of the means employed by certain species only of the animal kingdom to insure the satisfactory nourishment of the young individual. The recognition of these facts is essential for any serious consideration of the causes of disease. To retain, in connection with man, the vulgar use of the term inheritance would be to employ a terminology having a different significance to that accorded to it by workers in other branches of biology. The biologist has no alternative but to define inheritance according to the principles here laid down, nor have we, dealing with a limited field of biology, the right to modify those terms for our own convenience. That alone, therefore, is inherited which is inherent, which is the property of the individual at the moment of his becoming an individual, which is part and parcel of the paternal and maternal "germ plasm" from which he originates, or is provided by the interaction of the same.

It is unnecessary to point out, save as a precaution, that what is a property of the individual from the moment of beginning existence need not show itself for long years—a family failing toward premature baldness not until years after puberty, or an inheritance of gouty tendencies not until after thirty-five. As the different organs and parts assume their particular conformation and properties at different periods, and do not develop *pari passu*, so must the various inherited peculiarities make their appearance at various times.

Similarly, morbid conditions—disease, injuries, malformations—may be acquired by the individual while still *in utero*, or in after life.

¹ The distinction here is that usually made, that the new individual is an embryo during the earlier stages in which the future conformation of the parts is unrecognizable; when these appear and the individual exhibits the various organs and parts laid down in the relative positions possessed in later life; when, in short, all important parts are recognizable in due position, it is a fartus. The distinction is not a sharp one, but is of some use. Thus, the human being is regarded as an embryo until the end of the second month. Ballantyne has usefully introduced a third period, the germinal, preceding the embryonic, and ending with the development of the neural groove.

THE CLASSIFICATION OF MORBID STATES

THE CLASSIFICATION OF MORBID STATES.

According to Period.—Inasmuch as birth is important as corresponding with the greatest change in the relationships of the individual to the external world, we may, therefore, proceed to make the following classifications of morbid conditions according to the incidence of their causation:

1. **Inherited**, due to influences affecting the ovum or the spermatozoon before or at the moment of fertilization.

2. Acquired.

e

1

(1) Antenatal, or of intra-uterine acquirement.

(2) Parturient, acquired at the time of birth before complete separation of the individual from the maternal organism.¹

(3) Postnatal, acquired after birth.

According to Cause.—This must be our primary classification, but we may approach our subject from another direction, that of causation —namely, by determining whether a given agent acts directly or indirectly in setting up morbid conditions.

The division so made is not so satisfactory as that just given, since the same agent may, according to circumstances, act in either way. Study, for example, the action of a single agent—cold. This may either lead immediately to systemic and local disturbances—to frostbite, and even death; or indirectly, whether by lowering the vitality of the tissues, or again, by reflex nervous action, may lead to such alteration in the conditions of the circulation in the respiratory tract that the tissues there become less resistant to external agencies and afford a nidus for the growth of microbes, whereby a pneumonia is set up. Cold is the cause of disease in both cases, but in very different ways; in the first, it is the definite *exciding* cause; in the second, the *predisposing* cause. It does not directly cause pneumonia; the direct exciting cause of that disease is the pneumococcus, or some other microorganism; and this, it may be pointed out, can set up the disease without of necessity the previous influence of cold upon the lungs.

The causes, then, may again be classified into predisposing and exciting, and these are in action in connection with both antenatal and postnatal acquirements. Imperfect development of the heart, whether brought about by intra-uterine disease or by inherent imperfection of development, just as well as imperfections in the organ, the result of disease after birth, may directly induce morbid conditions causing obstruction of the circulation—<u>marbus cœruleus</u> (in the former case), ordema, etc.; or indirectly, through the impaired nourishment of the tissues, may render them more vulnerable and easily acted on by external agencies.

When we come to consider them more closely, we observe that inher-

¹ This is a very minor class, but has to be included, there being a few conditions which are neither antenatal nor postnatal in their acquirement.

INTRODUCTORY

ited conditions act in the main as indirect causes of disease. This is particularly noticeable in the finer constitutional defects, which result in the individual being more susceptible to one or other diseased state. Such *diathesis*, or specific susceptiblity to a particular disease or group of diseases, is a predisposing cause. Tuberculosis, for example, is not inherited; it is a weakness of the tissues, rendering them incapable of resisting the tuberculous virus, that is inherited.

Here, however, we shall not attempt to enumerate the various predisposing causes of disease. We do but wish to emphasize the fact that, in studying individual cases, we must constantly keep before us the existence, and most often the co-existence, of the two orders of causation, and endeavor to distinguish clearly between them. The direct cause, we need not say, is all-important; nevertheless, due weight must be given to the predisposing.

At the present time it is well to emphasize this matter. Reading the works upon medicine and pathology of but a quarter of a century ago, it is impossible not to be impressed by the fulness with which, in connection with every morbid state, the possible predisposing causes are enumerated. This was inevitable. With lack of knowledge of the direct exciting cause, the known or apparent predisposing causes loom large.

With the remarkable series of discoveries of the direct causes of disease which characterized the end of the nineteenth century, it has been equally inevitable that our attention should be prominently directed to the part played by these direct causes and to the mode in which they act. It has been inevitable that, as a consequence, the study of predisposition has been relegated to a very inferior position, and, indeed, largely neglected. But already there are signs that the pendulum is swinging back-signs of a disposition to appraise these indirect causes at a higher and truer value. We see, for instance, more clearly nowadays than formerly, that the mere existence of pathogenic bacteria within the tissues is not the sole cause of infectious disease; such bacteria may pass into the lymph glands and there be destroyed. For such bacteria (in general) to be in a position to excite disease, there must coincidently be a lessened resistance on the part of the tissues, and the causes leading to this lessened resistance-the causes predisposing to infection-are being more fully studied and their importance more fully appreciated.

For the orderly consideration of the causation or etiology of disease it will be best to take up the subject according to our first scheme of classification. Following upon this, we shall, in a special section, deal with the subject of predisposition.

CHAPTER IL

ult te. up tot of

re

en

e-

d,

aia

ia

'e

INHERITED MORBID CONDITIONS.

WE have in the previous section laid down the general principles of inheritance, and we have there indicated what can and what cannot be inherited. It remains to apply those principles to the consideration of morbid states in man and the higher animals. And first, it will be well to emphasize what cannot be inherited.

1. No order of mutilations, as such, can be inherited, i. e., while some may have a deleterious effect upon the general well-being of the offspring (this must be rare), and some even may possibly influence the development of a particular system (which must be still rarer), in no case can the identical mutilation or anatomical disturbance in the parent reproduce itself in the child. We owe the establishment of this principle more especially to Weismann, though years previously it had been laid down clearly by Francis Galton (1872), who also, it may be added, was the first to enunciate the doctrine of the continuity of the germ material.

2. Infectious diseases in the parent cannot be inherited. There may be transmission of such from the parent to the embryo, or even in animals possessing abundant yolk and albuminous surrounding matter from the parent to the egg, but such transmission is not inheritance proper.

This is tacitly admitted by all modern writers in connection with tuberculosis, but in connection with the disease with which children are most often born infected it is still the usual custom to speak of inherited syphilis. At the most, we may be permitted to speak of congenital syphilis, using that term as indicated on page 179, and again of inherited *parasyphilitic* lesions.

For, in the first place—although this may seem to some a refinement of logic—if inheritance be as we have defined, and as it must be, through the bioplasm, another individual living being cannot be part and parcel of the heritable material. The microbe of an infectious disease cannot be a constituent of the biophore. At most, it can be an accidental inclusion in the surrounding non-heritable matter of the cell. And in the second place, among the mammalia even this accidental inclusion is so improbable that it must be dismissed.

Such transmissions can occur in lower forms of life having eggs provided with abundant food material, and we have positive evidence of its occurrence. Thus, in the disease of silkworms known as pébrine, which now we know to be due to a microsporidian parasite (*Nosema hombycis*), Maillot and Pasteur noted that the eggs are infected; they nevertheless develop, and only in the developing insect do the microbes so multiply

INHERITED MORBID CONDITIONS

as to cause death. Schaudinn¹ has shown the same to be the case with mosquitoes infected by *Trypanosoma nochuæ* (the "halteridium") of the stone owl). These parasites may pass into and be laid with the eggs, remain latent during the development of the young gnat, only becoming active when the latter is adult and begins to suck blood. Like conditions had previously been determined by Theolald Smith² in connection with the ticks which cause Texas fever. These, filling themselves with the blood of an infected ox, drop to the ground, there mature, and lay their eggs, and the young ticks can convey the *piroplasma* and the disease to other cattle. And in birds the same has been definitely proved to occur. Thus, Maffucci³ has demonstrated, and Baumgarten⁴ has confirmed, that in fowls the eggs frequently convey the tubercle bacilli, and the same latency of the microbes is noted. From the eggs normal, if weakly, chicks hatch out, which at first run about and eat just like the healthy chicks, and only after some weeks emaciate and exhibit tuberculosis.

Human ova are free, or almost free, from yolk, and are relatively very small, nor have we a single observation showing that the mammalian ovum is phagocytic—able to take up solid particles. That the minute spermatozoa should act as carriers is still more unlikely, and the possibility that they do so has been negatived by Gärtner's⁵ reductio ad absurdum.

According to Wyssokowicz, the minimal number of tubercle bacilli that will set up peritoneal infection in the guinea-pig is 8; in the rabbit, 24 to 30.⁶ Gärtner, obtaining the seminal ejaculations from tuberculous guinea-pigs, found that only five out of the thirty ejaculations contained a sufficient number of bacilli to cause tuberculosis. Rohlff, employing the semen of men suffering from phthisis, did not once succeed in rendering rabbits tuberculous by inoculation into the anterior chamber of the eye. From these and other observations Gärtner concluded that the semen emitted by a phthisical patient (not suffering from genital tuberculosis) does not on the average contain as many as 10 bacilli.

Now, on the average (Loeb), the human seminal ejaculations contain more than 226,000,000 spermatozoa. If the semen contained not 10 but 1000 bacilli, the chances that an individual spermatozoon, fertilizing the ovum, should bear with it a tubercle bacillus, and so lead to germinal infection, are as 1 to 226,000; if 1,000,000, 1 to 226. Only 1 out of about 85,000,000,000 spermatozoa has the chance of fertilizing an ovum. In short, the chance of a spermatozoon conveying tuberculosis from the father to the offspring is so absurdly minute that it may be neglected.

¹ Arbt. a. d. Kaiserl. Gesundheitsamt., 20; 1904; pt. 3.

² Smith and Kilborne, Bulletin of Bureau of Animal Industry, Washington, 1893.

³ Ztschr. f. Hyg., 11; 1892, 445. Arb. a. d. Path. Inst. zu Tübingen, 1; p. 322.
⁵ Ztschr. f. Hygiene, 13; 1893; 101.

* Ztsenr. 1. Hygiene, 15: 1895; 101.

⁶ These figures are in fair agreement with those of Webb and Williams, who found that 5 tuberele bacilli, followed later by 15, when inoculated subeutaneously, might produce tuberculosis in the guinea-pig (Sixth Intern. Congress for Tuberculosis, Washington, 1: 1908: 194).

THE NON-INHERITANCE OF SYPHILIS

tgs,

ler

us. in

me

elv

m-

it,

a

ıg

10

The same considerations may be brought to bear upon foctal syphilis, in which, as must now be accepted, the Spirochata pallida is the causative agent. That cases of syphilis in the newborn are most often of relatively late intra-uterine acquirement is rendered evident by the fact, to which Chiari has called attention—namely, that in more than 90 per cent. of infants presenting signs of syphilis, the liver is the seat of most extensive syphilitic disturbances. In the adult, in which the disease is acquired through some cutaneous infection, extensive hepatic syphilis is rare compared with the frequency of the disease. Infection through the placenta amply explains the condition in the infant; for all the blood on its way from the placenta passes through the liver, which thus is the organ first subjected to infection.

It may, in fact, be laid down that wherever there are active and specific manifestations of tuberculosis, syphilis, or other infectious disease in the newborn child, the condition is of intra-uterine acquirement, not inherited, and the conclusion is supported by the very various stages to which one may find the disease developed in the newborn, as also by the more recent evidence alforded by the Wassermann serum reaction, that the apparently healthy mothers of syphilitic offspring are, nevertheless, the subjects of syphilitic infection.⁴

Further support and illumination is given from Friedmann's² interesting series of observations. This worker injected healthy does, immediately after copulation, with a few drops of an emulsion of tubercle bacilli, and six to eight days later, killing the animals, made serial sections through the uterus, with its contained embryos, to observe the relationship of the bacilli. He discovered not a single bacillus in the mucous membrane of the vagina or uterus, but all the embryos showed within them numerous bacilli of characteristic form, and in clumps (growing). The bacilli can thus pass into the developing ovum or embrvo. Other observers have noted that bacilli introduced into the uterus outside the amnion may, some days later, be found in the amniotic fluid. Whether through the placenta (from maternal infection), through the walls of the foetal sac, or by passage into the developing ovum before that sac develops, the bacilli may infect the embryo. These various means are adequate to explain the phenomenon without calling upon the improbable infection of ovum or spermatozoon prior to fertilization.

But, if syphilis and tuberculosis themselves be not inherited, it is deserving of note that the children of syphilitic and tuberculous parentage may exhibit conditions which are derived from the infected state of the parent, and are strictly inheritances. Offspring themselves showing no signs of the active disease, may nevertheless exhibit certain stigmata—feetal cachexia, malnutrition, senile expression, even malformations, arrested development of the bony skeleton, of the teeth

¹ See, for example, Engelmann, Ztschr. f. Gynäk., 33; 1909; No. 3, and the full study by Baiseh, Münch. med. Woch., Sept. 21, 1909. Colles' and Profeta's laws gain their explanation on the ground that mother and child remain refractory to syphilis because they are already infected.

² Ztschr. f. klin. Med., 43: 1901: 11 (with bibliography).

INHERITED MORBID CONDITIONS

(<u>Hutchinson's teeth</u>), etc.; children of tuberculous parentage, delicate constitutions, precocious mentality, etc. These characteristics, presumably due to the action of the toxin on the germ cells, we may refer to as *inherited parasyphilitic* or *paratuberculous lesions*. In addition to the experimental observations of Lustig and Carrière, which we have given elsewhere, bearing on this subject, we may recall the observations of Charrin and Gley,⁴ that among the offspring of rabbits immunized against diphtheria they had noticed a particular liability to definite rickets: enlarged cartilages, enteritis, pot-belly, and delayed growth; whereas these appearances had been scarce noticed in their long series of other observations. Such rachitis in the rabbit may be spoken of as an inherited paradiphtherial condition.

Here, then, we have conditions in the offspring definitely inherited from the parent and due to accurited modification or disturbance of the parental germ plasm. We have not as yet determined absolutely the specific inheritance of a particular order of lesions directly associated with the action of a particular causative agent. That may be so, or it may not be. If the children of tuberculous parents manifest a liability to tuberculosis, it has still to be proved that this is something over and above the liability to infections in general brought about by their lowered vitality, and the same may be said with regard to Carrière's and Lustig's observations upon the higher susceptibility of the progeny of immunized animals to tuberculosis and abrin poison, respectively. More observations are requisite before anything definite can be laid down upon this point. At most, the indications favor the view that there exist specific paratoxic lesions.

What is true regarding infectious diseases must to some extent hold also regarding chronic intoxications of various orders, alcoholism, plumbism, etc.

From these and the other considerations which we have discussed elsewhere it will be made out that the results of constitutional disease in either parent may be the following, according, on the one hand, to the extent of the influence of that disease, or intoxication, upon the germ plasm in that parent, and on the other, to the activity or potency of the germinal matter contributed by the other parent:

1. **Sterility.**—The germ cells being so profoundly modified that either (a) they are destroyed, (b) their development is arrested, or (c) being developed, they (ova or spermatozoa) are imperfect and incapable of fusing with the germ cells of the other parent.

2: Imperfect Development of the Offspring.—(u) Of such extent as to lead to intra-uterine death and abortion; (b) of less extent, a viable individual being produced presenting either—

(1) Gross anatomical defects;

(2) No gross anatomical defects, but lowered vitality, presenting itself either in the form of weakened powers of resistance against disease in general, or (?) proneness to develop the same functional disease as the parent.

⁴ Compt. rend. de la Soc. de Biol., 10 S., 2; 1895; 705.

THE INHERITANCE OF ABNORMAL CONDITIONS

3. **Perfect development of the offspring**, with no appearance of functional disease or lessened power of resistance, (a) the offspring of these again being perfectly normal; (b) that offspring showing in subsequent generations constitutional weakness (recessive).

4. (?) Perfect development of the offspring, with increased power of resistance, the immunization of the parent having been accompanied by the development of an acquired tolerance to the particular toxin on the part of the germinal bioplasm.⁴

Spontaneous Variations; Mutations.—These show a marked tendency to be inherited (p. 180). In man, it is true, it is difficult to assure ourselves that a given departure from the normal has appeared for the first time in the history of any particular stock, so that we are apt to place all such conditions among those already inherited, descending from previous generations. There is no doubt, however, that albinism, Daltonism, hemophilia, and so on, have suddenly shown themselves at a bound in some one individual; no doubt, that is, because our experience with lower forms of life shows that this does happen.

THE INHERITANCE OF ABNORMAL CONDITIONS PASSED DOWN FROM PREVIOUS GENERATIONS.

If certain anomalies and constitutional defects are capable of being transmitted from the individual in whom they first arise to the descendants of the same, *a fortiori*, when a constitutional defect has shown itself in a family for several generations, there is increased likelihood of its being transmitted to further generations. While many doubt the inheritance of constitutional defects of any order that are acquired, the volume of facts at our disposal is too great, and the facts themselves too convincing, for any to deny that those already inherited—anomalies of certain orders, specific and general constitutional disturbances, or, more correctly, diatheses (predisposition to the same)—are frequently transmitted; it is to these that our attention as pathologists is most often called.

Inherited Anatomical Anomalies.—While it must be kept in mind that by no means all anomalies are heritable—many due to intrauterine disturbances are certainly not—nevertheless, there is a large and important class, that is, a class so large that only some of the best marked of those affecting man will here be noted, and that without description. Of these, we may mention polydactylism among anomalies of excess, bradydactyly and hypospadias among anomalies of defect.

Certain conditions it is difficult to classify. There must, for example, be an anatomical basis for color blindness (Daltonism), though that is

¹ We introduce this doubtfully, but because it is possible. Certainly, at the present day the bulk of the evidence is against the inheritance of acquired immunity. It is, however, proved that influences of a chemical nature acting upon the parental organism may coincidently modify the germ plasm in a retrogressive direction; on general principles, therefore, we should be prepared to admit that other substances exist capable of setting up progressive modifications. See page 198.

14

ve

te

PS

of

'd

le

it

y did

's

INHERITED MORBID CONDITIONS

beyond our present means to determine, and other conditions stand doubtfully between this class and the next, that of diatheses. Is hemophilia, for example, merely a predisposition to bleed, provided the individual be subjected to certain alteration in environment, or is there underlying this a definite anomaly, and weakness of the vessel walls? We suspect the latter, but what the exact anomaly is has not as yet been convincingly demonstrated. So, also, as regards many inherited nervous conditions; some undoubtedly are associated with an agenesia or aplasia (lack of development or imperfect development, respectively) of particular nerve centres, others are due to what Gowers would term an abiotrophy, or premature exhaustion and degeneration, of particular groups of nerve cells. Others, again, to no particular weakness of any particular set of nerve cells so much as to a general weakness and instability of the whole of the higher centres; it is not so much specific anatomical defects as strain, brought to bear upon particular centres in a weakened system, that originates the particular form of nervous breakdowns.

Diatheses.—From these borderline cases we pass to the diatheses proper. Some of these we have already discussed, notably those associated with sundry of the infectious diseases.

1. We have shown how vague and unsettled is our knowledge regarding acquirement of the specific lack of resisting power to particular infections. There is, on the contrary, a very definite conviction, based upon experience, regarding the existence of "racial diatheses," or, as it may be expressed, failure on the part of certain races of men to have developed resistance against particular infections-of the white races of mankind against yellow fever, the Polynesians against measles, the red Indians, and, to a little less extent, the negroes, against tuberculosis, etc.--and an equally clear recognition of the existence of familial diatheses toward such diseases as tuberculosis, scarlet fever, measles, and acute rheumatism. How far these diatheses express themselves as bodily configurations is still a matter of some doubt; this, however, is generally admitted, that, as regards tuberculosis, a fine skin and fine hair (of any color, not necessarily light), light bones, long chest with narrow sternal angle, small first rib, and expansion below the normal, indicate a predisposition to the disease, although it may attack, and attack acutely, those exhibiting none of those traits.

2. Other diatheses are of the metabolic type, exhibiting themselves as a predisposition to metabolic disturbances. Foremost among these is the gouty, or, as some would term it, and, unfortunately, the uric acid diathesis; others are the obese and the diabetic diatheses. With these, also, there is a certain racial liability, although to a large extent this would seem to be a matter of habits of life. Gout is frequent among the English, rare among the Scotch; diabetes common in Jewish communities, as also, among the females, is obesity; the Dutch are liable to greater corpulency than can be wholly ascribed to inaction; the Bushmen and Hottentots to such overdevelopment of the subcutaneous fatty tissue of the gluteal region (steatopygy), and, in some, also of the thighs, as to lead to a grotesque exaggeration of outline. More accurately, these states and the frequently associated *longinymph* condition

THE INHERITANCE OF ABNORMAL CONDITIONS

in the women of these races, or exaggerated size of the labia minora, are inherited anatomical variations of the soft tissues. From the earliest known reproductions of the "human form divine" found in the caves of southern France, which show this steatopygy, and the identical habit of the present Bushmen to live in caves, and make representations of himself and animals of the chase upon their walls, ethnologists conclude that this peculiarity has descended unchanged through long thousands of years from the paleolithic cave dwellers to the Bushmen of to-day.

m

T-

3. Others, again, are the nervous, and these form a very important group. Some of these we have already indicated in discussing whether here we have to deal with the transmission of anatomical defects. We would only repeat that in some cases clearly the recognition of anatomical changes is beyond our power to determine. Defects there must be, either in the relationship, or, more probably, in the functional capacity of the individual nerve cells or groups of the same. Hysteria, for instance, is notably diathetic, and, with the somewhat allied migraine. is noticeably transmissible, and these are typical examples of what, failing anatomical evidence of disturbance, we regard as functional conditions. Epilepsy also is without a definite anatomical basis, and is again transmissible. Thus, Echeveria¹ found that of the 553 children of 136 epileptics, only 105 were healthy, 78 were epileptic, 195 died of convulsions in childhood, 39 exhibited some form of paralysis, 51 were hysterical or choreic, 11 insane, 18 idiots. The remainder had died in childhood of various ailments. Virchow, indeed, would distinguish between a general and a systemic reversion. If the distinction is valid, then the nervous and infectious diatheses are examples of the latter. It is worthy of note that in the development of the nervous system and of resistance to specific infections, we have the latest group of acquirements by the human species, qua species; that, as a general law, conditions last acquired are those liable first to be lost. Thus, these two diatheses must be associated with failure to attain complete normal development. We would regard them not as special reversions so much as the first and slightest stages of general reversion. But a large number of nervous states appearing not at birth but in later life are clearly examples of hereditary weakness and premature exhaustion (abiotrophy) of particular nerve centres.2

¹ Quoted by von Hansemann, Descendenz und Pathologie, p. 269.

^a The following list of morbid states of nervous origin does not pretend to be complete, but is sufficient to give some indication of the extent and varied nature of either homeomorphous or heteromorphous transmission of nervous disease; *Mainly Homeomorphous*: Friedreich's ataxia (Marie's hereditary carebellar ataxia); Thomsen's disease; Huntingdon's chorea, as also convulsive tie and chorea minor; interstitial hypertrophic progressive neuritis of childhood (bejerine and Sottas); progressive muscular atrophy (Chareot-Marie-Tooth); progressive bulbar palsy (glosso-labio-laryngeal paralysis); progressive spinal muscular atrophy of childhood (type Werling-Hoffmann, this descends through the mother); bulbar paralytic facial type of muscular atrophy (Fazio-Londe); the Duchenne-Aran type of atrophy (Strümpell-Gowers); Bernhardt's combined spinal and bulbar type of muscular atrophy; hereditary spastic spinal paralysis (Strümpell-Bernhardt-

INHERITED MORBID CONDITIONS

Homeomorphous and Heteromorphous Inheritance.—While we observe that inheritance of liability to disease may in certain instances exhibit itself by the offspring succumbing to the identical influence as did the parent, and exhibiting the same symptomatology (homeomorphous inheritance), in other cases it manifests unlike disorders, although these may be along special lines. There are, as it were, families of diseases, the members of which may follow in direct descent, or, on the other hand, be interchangeable in inheritance (heteromorphous inheritance).

We have just given an admirable instance of these two types in connection with epileptic inheritance. An equally striking example of this grouping of diseases into families is to be seen in connection with the gouty diathesis. It is notorious that the gouty father tends to have the gouty son, but does not always. The majority of observations place the proportion of cases in which there is a history of gouty inheritance at 44 per cent. (Sir W. Gairdner places it as high as 90 per cent.), and the condition affecting the male (most often) transmission is mainly through the father. But gout is far from being the only condition affecting the gouty family more than the non-gouty. Bouchard¹ found the following percentages of incidence of disease in the ascendants and collateral members of the families of 33 gouty individuals.

										Per cent,
Gout .										44.0
Obesity										44.0
" Rheumat	isr	0. ¹³								25.0
Asthma										
Diabetes										12.5
Gravel .										
Eczema										
Biliary lith										

In only 12 per cent, of the cases studied was no heredity, either homeomorphous or heteromorphous, to be made out; or, in other words, in only 12 per cent, of the cases was the gouty condition acquired without antecedent diathesis. Similarly, data were obtained in studying a series of cases of obesity, and rheumatism (both chronic and acute). Everything nowadays points to acute articular rheumatism being an infective disorder; nevertheless, he found a history of heredity in 32 per cent, of the cases; or, in other words, certain constitutions are hereditarily prone to be affected. In migraine, biliary lithiasis, and diabetes one finds the same tendency for members of this same family of morbid conditions to preponderate in the personal and family histories.

It is clear that what we have to deal with here is not so much a special diathesis rendering the individual liable to succumb to one given disease, as a peculiar disturbance of nutrition rendering the individual liable

Krafft-Ebing, observed through five generations); amaurotic family idioey (Tay-Sachs disease); familiar form of lateral sclerosis and spastic paralysis (appearing in males of a family between the ages of twenty and thirty years); myoelonus epilepsy (of this 75 per cent. of the cases are hereditary). *Liable to be Heteromorphous*: Migraine, epilepsy, neurasthenia, hysteria.

¹ Quoted by Le Gendre in Bouchard's Pathologie Générale, 1: 1896; 348.

THE MARRIAGE OF CONSANGUINES

to be affected by one or other of a large group of diseases. There is, undoubtedly, the tendency in the one direction, e, g,, if 44 per cent, of gouty persons give a history of gout in the family, it cannot be denied that there is a special liability toward this one disease; but in another 44 per cent., according to Bouchard, in whom there is not the family history of gout, there is a history of one or more of these other conditions. It is very possible that environment may explain this interchange of disease; that members of a family exposed to the same influences and living in like surroundings will manifest the one special disease, whereas variations in environment lead to the manifestation of the disturbances characteristic of some other member of the group. This, however, does not detract from the remarkable fact that such a group or family of disease exists and that there is inherited diathesis.

N

For further discussion on the subject of the inheritance of disease the reader is referred to the introductory chapters (p. 159 et seq.), where also will be found reference to the subjects of atavism and reversionary inheritance, and the development of the so-called *degenerates*. Here, before closing, it is fitting that a little should be said upon the cognate and frequently discussed question of—

The Marriage of Consanguines.—Expressed in the terms of our theory, when the biophores of both parents possess particular sidechains, or groups of side-chains, of like composition, there is a much greater likelihood of those side-chains being potent in the offspring, and for that offspring to possess the traits brought about by the presence of those side-chains, than when, on the contrary, the corresponding sidechains in the two parents vary. If, again, certain side-chains characteristic of the fullest development of the species are wanting in the parental biophores, those side-chains cannot possibly be present in the offspring. In other words, we should expect the marriage of consanguines to reproduce with greater sureness family characteristics, whether progressive or regressive, than will marriage of an individual with the member of another family not possessing those characteristics.

And, as a matter of experience, this is what we find is the case, and not only that, but this principle of inbreeding is that depended upon by breeders to preserve, intensify, and, indeed, to fix and render constant variations which are regarded as favorable or advantageous.

The union of those belonging to the same family, when of nearly related degrees of consanguinity, is likely to have good or ill effects, according to the absence or presence of constitutional defects peculiar to the family. When such constitutional defects are present they tend to be intensified in the offspring of consanguineous union. Where they are wanting, the offspring is likely to be of good constitution. And, further, where the family is in any one respect above the normal, *i. e.*, when it exhibits a favorable variation, placing it at an advantage over ordinary mortals, the marriage of those nearly related is of actual advantage, by impressing and rendering more stable the favorable variation.

In stating this, the existence of latent or recessive properties in the germ plasm must be kept in mind. Not all the progeny of a consanguineous marriage will necessarily exhibit the family weakness or the

family beneficial property; atavistic, recessive characters may exhibit themselves here and there, but this, undoubtedly, is the tendency.

The vast body of facts accumulated upon the subject of intermarriage of relatives and its results can only be classified along the lines here laid down. It is by "in-and-in" breeding that, for example, the Hanover breed of pure white (albino) horses was established more than a century ago, the breed tracing back to a single albino horse, apparently a spontaneous variation. The Ancona breed of silk-haired sheep trace back similarly to a single animal.

A notable example of the effects of intermarriage as intensifying the inheritance of malformations in man is given by Poulton.¹ The village of Iseaux, in Isère (France), being remote from other villages, the inhabitants constantly intermarried. At the end of the last century the majority of these, male and female, presented an accessory finger and accessory toes on hands and feet; forty years ago the monstrosity was still very common, but now, with the improved means of communication and the introduction of new blood by marriage outside the village, it is tending to disappear.

Other village histories can only be explained by this law. Thus, to quote examples given by Le Gendre.² At Orthez (Basses Pyrenées), the Protestant families, being a class apart, were compelled to marry among themselves. The members of these were, in general, according to Réclus, poorly built and weakly, and among these were very numerous epileptics, so that each Protestant house possessed a special room reserved for the epileptic members. Since the advent of improved communication and increased facility for travel outside the previous narrow limits the state of affairs has disappeared.

The village (bourg) of Batz, studied by A. Voisin, shows the opposite condition of improvement by intermarriage. There intermarriage is the rule, and all are descendants of half a score of families—have names not found in the neighboring communes. Out of 2733 persons, Dally found that 870 had the same surname. But here the inhabitants are well built and of good physique; indeed, there is a smaller proportion of exemptions from military service than in the rest of the Department. Lavery cites a similar example in the commune Fort-Mardyck, near Dunkerque. Four families from Picardy were established there in the time of Louis XIV. At the time of Lavery's observations there were 1800 descendants, robust, without sign of any inherited defect, and their birth rate was higher and death rate lower than that of any of the neighboring communes. These had kept to themselves and intermarried among themselves.

The deduction here is obvious: marriage between consanguines may be recommended where the family is of sound physique, possessing no noticeable defect; it is powerfully contra-indicated when such defect exists. Inquiry must especially be made as to the existence of neuroses and constitutional predisposition to infectious disease.

¹I quote from Le Gendre: Bateson (*Materials for the Study of Variation*, p. 399) gives the name as Potton, but is unable to trace the original authority.

² Article Inheritance, in Bouchard's Pathologie Générale, 1:1896.

CHAPTER III.

THE CAUSATION OF MORBID CONDITIONS OF INTRAUTERINE AND PARTURIENT ACQUIREMENT.

THE subject of antenatal pathology is but yet in what may be termed a foetal state; much has been accomplished during the last twenty-five years to establish the framework of the subject, but much greater development is before it. Certain matters stand out clearly; they have been established with fair detail. Regarding others, only scattered observations exist among many theories. Were we to deal with the matter at all fully, this detail and the discussion of individual opinions would enter largely. Add to this, that dealing here with the causation of disease, it has to be confessed that, in general, we know more about the processes and conditions which appear to be due to intra-uterine disturbances than about the precise cause or causes of the individual con-This is not the place to enumerate those conditions in detail; we can but call attention to the more important. The most striking group-namely, the monstrosities and anomalies-will be treated in the following chapters (IV and V), and there it will be most satisfactory to note, at the same time, what we know concerning their causation. This chapter, therefore, will be relatively brief and will deal in generalities.

Briefly, under the action of various causes acting during intra-uterine existence, there may result:

1. Death with-

- (a) Absorption of the embryo: "blighted ovum," the embryo failing to develop and disappearing, the placenta and membrane still exhibiting growth for a time.
- (b) Abortion, the imperfect fortus being born dead or dying immediately after birth.
- (c) Premature labor.

2. Monstrosities.

- 3. **Malformations** (not so extreme as those causing monstrous births). These may be—
 - (a) Of defect.

(b) Of excess.

 Impaired vitality, with imperfect development, without gross anatomical change:

(a) General: infantilism, etc.

(b) Systemic: more particularly of the nervous system.


5. Cachexia.

6. Infection.

7. Traumatism (this distinct from the traumatism of parturition).

216 MORBID CONDITIONS OF INTRAUTERINE ACQUIREMENT

The causes of disturbances are, it will be noted, clearly, though not wholly, identical with those produced as a result of inheritance. As a matter of fact, we must admit, with Woodruff,¹ that causes acting on the ovum soon after fertilization—causes capable of affecting all the cells of the egg alike—must have results somewhat of the same order as are produced by those same causes acting on the bioplasm before fertilization; and, as experimental research is showing more and more definitely that the more pronounced anomalies and monstrosities, when set up by influences acting on the individual, date back to the earliest stages of development, we must not be surprised to find that an important group of disturbances may originate equally from influences acting before and after fertilization, and that no sharp line exists between



Early stages of marine gasterood Hyanasai: 1, the first two blastomeres; 2, the usual development of the eight-cell stage; 3 and 4, the first two blastomeres having been shaken apart, each proceeds to redivide independently, with the production of dwarf larve.

variations (due to the former set of conditions) and *modifications* (due to the latter). There are, however, many conditions which are only of intra-uterine origin.

Basing ourselves very largely upon the results of more recent studies upon the experimental pathology of the embryo, we may group the causes of these intra-uterine disturbances as follows:

- 1. Physical and mechanical, including injuries of various orders.
- 2. Malnutrition.
- 3. Intoxication.
- 4. Infection.

1. Mechanical and Physical Causes.—To purely mechanical and physical causes acting upon the earliest stages it appears evident that

¹ American Med., 10; 1905; 661 and 706.

some of the most remarkable monstrous growths are due. Experimentally, the eggs of lower forms of life, when in the two-, four-, and eight-cell stage, may be violently shaken, so that the cells fall apart, when each has been found capable of developing a separate individual, or, if they remain partially attached, double and other multiple monsters. That violent shaking is the cause of double monsters and of an order of twins in man is most doubtful. Here, Jacques Loeb's observations upon altering the tonicity of the surrounding medium affords a more probable cause. He found that placing echinus eggs in the early stages of division in diluted sea-water was followed by such osmotic swelling of the eggs that the membrane broke, and that part

of the egg which protruded underwent independent development. The same has been noted by Bataillon¹ in the case of teleostean (fish) eggs; by strong movements, the first blastomeres may be forced apart and separated by serous fluid to a greater or less extent, and undergo this independent growth.²

We believe, then, that mechanical and physical means are active in producing some of the most extreme forms of monstrosities. So, also, do they best account for one group of anomalies-those brought about by pressure effects, owing either (Dareste) to incomplete formation of the amnion with escape of fluid, or to deficient formation of the amniotic fluid; such may lead to adhesions between the amnion and foetus, the formation of amniotic bands, and consequent local retardation of growth. or produce effects of like nature without adhesions. Certain orders of anencephalic monsters, spina bifida, talipes equino-varus and yet severer forms of arrest of growth of



Amniotic threads on hand of seventh to eighth month fortus; loss of last phalanx of second finger; index finger constricted by band and terminal phalanges form a process adherent to the band from the second finger, (Marchand.)

the limbs, have been ascribed to this cause. Here, also, may be mentioned the occasional amputation of limbs, and, rarer, arrest of growth of other areas, brought about by loops of the unbilical cord—the fœtus in its movements tying itself in a knot. It must not be thought that all cases of imperfect limb development are due to this cause; in fact, the majority cannot be accounted for, but some undoubtedly must be so explained.

Cases occasionally occur of feetal fractures, and this usually unaccompanied by any history of maternal traumatism. Labor may have been

² For a fuller description of the subject of double growths see Chapter IV.

Arch. f. Entwickelungsmech., 11: 1901.

218 MORBID CONDITIONS OF INTRAUTERINE ACQUIREMENT

easy, and yet at birth, as in Linck's⁴ and in Chaussier's⁵ cases, and in Klotz's case,³ published from our laboratory, as many as thirty to one hundred separate fractures of long bones, ribs, stern im, etc., may be counted; obviously, the only satisfactory explanation in those extreme cases is an abnormal fragility, due to imperfect osteogenesis. In other instances, many bones are found presenting a sharp angle, with cicatrix above, strongly suggesting old compound fracture. Sparling⁴ regards those as not necessarily indicating fracture, but as produced by amniotic bands, which have deformed a part and subsequently have been torn away. To such tearing way of bands, Ballantyne attributes the occasional existence of wounds and skin defects in the newborn.

2. Malnutrition.—Simple malnutrition of the mother, lack of sufficient food, is a well-known cause of puny development and of weakly condition of the offspring. There is little evidence that in itself it leads to any definite anatomical defects. Where these are present it is likely that more than mere impaired nutrition is at fault, and that we have to deal with the third cause—namely, intoxication from placental absorption.

Malnutrition of the foctus may, however, be brought about in another way-namely, by placental disturbance. More particularly in heart disease, and again in the subjects of syphilis, there may be grave alterations of the placenta; so that, either from the fortal side the placental circulation is greatly lessened, or, on the maternal side, intra-uterine hemorrhages, etc., so reduce the interchange between the fœtal and the maternal blood that premature death is brought about. Yet another, and in this case extraordinarily advanced effect of placental circulatory disturbance is seen occasionally in monochorial twin pregnancies, where the twins, developing from a common ovum, have a common chorion and fused placenta. Here the more vigorous twin usurps more and more of the placental circulation, until, through anastomosis, it drives its blood (venous in quality, but arterial as regards the vessel carrying it) into the umbilical artery of the other twin, arrests the heart action and the circulation proper of that other twin, and so the development of the same; what does develop becoming extremely ædematous. In this way is developed a feetus acardiacus, or chorio-angiopagus (p. 231)

3. Intoxication.—If, as above stated, simple malnutrition of the mother mainly results in constitutional weakness of the child, without grave anatomical defects, maternal malnutrition, associated with constitutional disorders, has much more serious results—resulting, according to the nature of the disorder, in the whole gamut of disturbances mentioned in the opening paragraph of this chapter. Thus, at the one end of the scale obstructive cardiac disease, by the slowing of the uterine circulation and placental hemorrhages and infarcts, leads to death of

¹ Arch. f. Gynäk., 30; 1887; 264.

² Bull. Fac. de Méd. de Paris, 3: 1814: 301.

³ Journ. of Pathol., 13; 1909; 467.


⁴ Arch. f. Geburtsh. u. Gyn., 24: 1892: 225.

PLATE VI



Osteogenesis Imperfecta, with Multiple Antenatal Fractures of Ribs, Long Bones, etc.

The fractures show themselves as nodosities with abnormal curvatures of the bones.



INFLUENCE OF INTOXICATION

the foctus and stillbirth; at the other, infectious disorders may induce the gravest anomalies and definite monstrosities. More particularly it may be laid down that metabolic disturbances in the mother, when they do not induce actual sterility, as, for example, when they develop during the course of pregnancy, tell severely on the offspring; for example, kidney disease. Among these, obstructive liver diseases, with jaundice and renal incompetence, call for special mention, and may exert specific effects, although here more exact observation is requisite. In this connection the observation of Charrin¹ deserves remark. Taking gravid goats, he injected into them emulsions of liver tissue, and thereby developed in these animals a cytolytic action, *i. e.*, the blood of the goats gained the specific power of destroying foreign The kids born to these goats were apparently normal, liver cells. save that their livers showed grave degenerations. The conclusion to be drawn is that the cytolytic substance developed in the maternal organism, and, present in the blood, had diffused through the placenta or been taken up by the fortal tissue and had exerted its specific action upon the (foreign) liver cells of the foctus. Charrin is so experienced an observer that what is stated by him as a fact must be freely accepted.

It has been observed by Halsted,² and confirmed by others, that following partial removal of the thyroid in the bitch, the pups born are goitrous. Presumably, absence of adequate internal secretion from the maternal thyroid leads to heaping up of the bodies neutralized by that secretion, increased work and overgrowth of the fœtal thyroid glands.

With reference to infectious diseases, we have noted that toxins may tell upon the germ cells. Their influence upon the developing embryo and fœtus is very noticeable. Very often acute infectious diseases smallpox, scarlet fever, typhoid, etc.—lead to the expulsion of the dead child, and this without any indication that the child itself is infected; indeed, as a rule (although not constantly), with those acute infections in which the specific microbe is known, cultures from such fœtuses yield negative results. Intoxication, the effects of absorption of toxins from the maternal blood, is the simplest explanation of such. Where milder infections, la grippe, for example, occur in the early stages of pregnancy, if abortion is not brought about, the condition of blighted ovum may be encountered, or the development of anomalies.

This influence of the mother upon the fectus is well shown, more particularly in connection with lead and mercury poisoning. We have already quoted Lize's observations on the effect of poisoning by nitrate of mercury (p. 198), and have referred to those of Constantin Paul³ on lead. We here tabulate his figures:

¹ Semaine Méd., 1902; 413.

² Johns Hopkins Hosp. Rep., 1: 1896; 373; and Med. Record, New York, 34: 1888; 368; see also Marine and Williams, Arch. of Int. Med., May, 1908.

³ Union Méd., 2 S., 13 : 1862 : 106.

) MORBID CONDITIONS OF INTRAUTERINE ACQUIREMENT

	Number of cases,	Number of preg- nancies,	Abortions, pre- mature labor, and stillbirths.	Infants born living.	Remarks.
 Mother showing symp- toms of plumbism . Mother working in type foundry, all of 	4	15	13	2	infant died within twenty- four hours.
whose previous preg- nancies had been nor- mal	5	36	29	7	4 of these died in first year,
foundry had five preg- nancies	1	5	5	0	After ceasing to work had healthy child.
4. Mother working in- termittently in type foundry; while work- ing there	3	3	3	0	When away from work for
5. Mother in whom blue line on gum the only					some length of time gave birth to healthy children.
sign of lead poisoning . 6. Husband alone ex-	6	29	21	8	
posed to lead ,	2	32	12	20	Of these, 8 died in first year, 4 in second, 5 in third.

From this table it will be seen that the mothers who suffered from lead poisoning during pregnancy showed by far the most pronounced effects, though, as already noted, paternal poisoning had a very definite influence.

Nor is premature death the only result; in the children of workers in lead, there is a painful frequency of idiocy, imbecility, and epilepsy.

We would here recall the other figures given by us elsewhere. There can be no question that intoxication of the pregnant mother tends to exert a most deleterious effect upon the offspring; several observers, indeed, have proved experimentally that poisons, such as lead, mercury, arsenic, carbon monoxide, morphine, alcohol,¹ pass through the placenta, and may be detected in the foctal tissues.

As regards alcohol, Sullivan's² figures are especially striking. He investigated the histories of female chronic drunkards, choosing cases in which other degenerative features were wanting. In sober mothers the rate of stillbirths, abortions, and deaths of children before the third year he established as 23.9 per cent.; in these it was 55.8; squalor may account in part for this high figure, but a more careful study showed that the death rate increased progressively as the mother became longer

² Jour. Ment. Sci., 45: 1899; 489,

- 99

¹ This has been absolutely proved by Nicloux, L'Obstetriqué, 1900: 97.

PLACENTAL DISEASE

and longer a victim to alcohol, and when the history of the successive births was determined, thus:

								Per cent. dy-	Total	
						Cases.	born dead.	ing before 3.	per cent.	
First births						80	6.2	27.5	33.7	
Second "						80	11.2	40.8	50.0	
Third "						80	7.6	45.0	52.6	
Fourth to fi	ifth	bir	ths			111	10.8	54.9	65.7	
Sixth to ter	nth	- 6				93	17.2	54.8	72.0	

The infantile mortality, it will be seen, shows the same tendency to increase as do the stillbirths; if the foctus is impaired, so also is the vitality of the offspring.

4. Infection.-Normally, it is scarce necessary to say, there is no communication between the foetal and maternal circulation, the cells of the foetal villi acting as a barrier; and thus, as a rule, bacteria circulating in the maternal blood do not find their way into the foctus. Direct infection of the foctus from the mother is distinctly the exception and not the rule. Nevertheless, it does occur, and occasional cases have been reported in a considerable number of diseases, cases in which either the specific lesions of the infection, or the causative bacteria have been detected in the foctus at birth or so soon after birth that infection during parturition is wholly ruled out. Thus there are records of syphilis (the mother alone being infected), tuberculosis, variola (the child being born with the eruption fully pronounced), varicella (but not vaccinia, although there are several cases of at least temporary immunity conferred on the offspring when the mother has been vaccinated in the last three months of pregnancy), measles (some 20 cases), scarlatina, (likewise), erysipelas, and septic disorders, acute rheumatism, typhoid, anthrax (in the lower animals), cholera, epidemic cerebrospinal meningitis, influenza, mumps, relapsing fever, malaria, and yellow fever.1 The list is a long one, but the number of cases on record is small.

The simplest explanation of such cases is that the specific organism coming to rest in one of the maternal blood sinuses, or being taken up by the cells of the fœtal villi which have made their way into the sinuses, multiply there, lead to local tissue destruction, which, extending into the walls of the villi, eventually leads to the microbes being carried by the fœtal blood into the fœtal tissues.

PLACENTAL DISEASE AND ITS INFLUENCE UPON THE FOETUS.

The placenta, while strictly a part of the foctus, may, nevertheless, undergo primary disturbances, and, as upon it the foctus depends wholly

¹ The literature on this subject, as upon all branches of antenatal pathology, will be found indicated in Ballantyne's *Antenatal Pathology*, Edinburgh, Green & Sons, 1902. This is, to our knowledge, the only work in any language that attempts to deal with the whole subject of intra-uterine disorders, and is a mine of information. Another useful article upon placental transmission is that by Lynch, Johns Hopkins Hospital Bull, 10: 1902; 283.

222 MORBID CONDITIONS OF INTRAUTERINE ACQUIREMENT

for its nutrition and supply of oxygen, any extensive disturbance has a most serious effect. What we may term the active part of the placenta is essentially fœtal; our conception of this organ is simplified if we regard it as a series of finger-like processes of the chorino, or outer coat of the fœtus (*i. e.*, fœtal sac), which, containing vascular loops and covered by epithelium, make their way into the wall of the uterus until they penetrate and lie within the large blood sinuses of the uterus. This epithelium has extraordinary phagocytic powers: it absorbs the tissues before it until it gains entrance into the maternal vessels.

The extent of this process varies, or otherwise there is considerable variation in the dimensions of the placenta, and so of the nutrition of the foctus; the greater the number of the chorionic villi entering into its formation and the more active their plagocytic power, the greater the nutrition of the foctus and the more active its growth.

It follows, however, from this method of development that the invasion of the maternal sinuses is a precarious matter; the very act of invasion of the walls leads to weakening of the same; in fact, under wholly normal conditions hemorrhages occur on the maternal side, resulting in the formation of what we may regard as accessory sinuses, into which other villi make their way. The adaptation of the parasitic growth to the maternal vessels is such that with increased blood pressure (coupled, it may be, with impoverished state of the maternal blood and impaired nutrition of the villi) there may be very extensive and widespread hemorrhages. When this is the case—and this happens not infrequently in obstructive heart disease—the fortus is liable to become asphyxiated, the effused blood being stagnant, affording little nutrition and less oxygen to the foctus, and in itself obstructing the normal circulation. Such placental hemorrhages form one cause of premature labor and stillbirth.

Localized hemorrhages, again, may be followed by thrombosis and organization, and where the resulting fibroid areas are extensive, these also greatly reduce the area of nutrition of the fœtus, with resulting impoverished growth and impaired vitality. Fibroid areas of this nature in the placenta are far from uncommon and may be abundant in obstructive circulatory disturbances, as, again, in maternal syphilis; though here, as we shall point out, other factors are concerned in their production.

From causes that are little known—at times, it would seem, from inherent vices of development, at other times, as a consequence of impaired nutrition from the maternal blood (in maternal nephritis and cachexia, for example)—the fortal villi are liable to be diseased; to be œdematous or even cystic, or to undergo fibroid change, with consequent contraction and obliteration of the contained vessels. According to the extent of those disturbances so is there a greater or less amount of malnutrition of the fœtus, and of asphyxia, leading toward premature death.

So also there may be placental infection. The effects of this on the organ we see best in the more chronic infectious diseases, and here more particularly in syphilis. In tuberculosis the presence of actual tubercles has been recorded, but is rare, nor have the anatomical changes been so fully studied. In syphilis the disease, as in other organs, may show itself both by the production of actual gummata or by widespread vascular changes. This, at least, is the usual teaching. The more recent

work of Bondi,¹ and more particularly of Thomsen,² affords a very different picture of placental syphilis. The latter observer examined the placentas of 100 syphilitic women. He found, in the first place, that proliferation of the vascular intima in the foetal placenta is far from being a marked feature. What is more characteristic is the cellular overgrowth of certain villous processes, coupled with extensive œdema of others, both of them contributing to the greater size of the placenta, which is a pronounced characteristic of the condition. Whereas the weight of the normal placenta relative to that of the child is as 1 to 5 or 6, that of the syphilitic is usually given as 1 to 3 or 4. Thomsen found that it might be as high as 1 to 1.5. What is particularly characteristic is the co-existence of these changes in the villous processes with multiple small abscesses, for such are present rather than typical gummata. It is to be noted that in other conditions-in tuberculosis, for example -the abscesses may be present, or, again, proliferation; only in syphilis do both exist to a marked extent. Yet another feature is the extensive leukocytic (polynuclear) infiltration of the umbilical cord. Only in 5 cases out of 30 did he find these cord changes in cases not regarded as syphilitic, and of these 5, three of the infants were subsequently brought to the hospital with syphilitic lesions.



Girl, aged ten years, showing cicatricial grooves due to constriction of umbilied cord. At birth, according to the mother, the grooves in the abdominal wall and left thigh were occupied by the cord. (Hawthorne.)

As a result of these changes the placenta shows general enlargement, coupled with anemia; is pale, with yellowish whitish regions, indicating the more fibrous areas. It will be readily understood that there is obstruction to the circulation and malnutrition, and that the altered

¹ Arch. f. Gyn., 69; 1903; 223.

² Ziegler's Beitr., 38: 1905: 524.

224 MORBID CONDITIONS OF INTRAUTERINE ACQUIREMENT

relation of placental weight to that of the child is due in part to the imperfect growth of the latter.

It has been laid down by some observers that a distinction can be made out between fcctal syphilis (the mother being unaffected) and primary maternal syphilis, by the villi being more affected in the former, the sinuses in the latter. No such general rule can be laid down. The evidence afforded by the Wassermann reaction is that the older teaching is erroneous: where the fcctus is syphilized the mother also is infected, even if the disease remains latent or largely localized in the uterus.

The fœtal disturbance most frequently associated with the syphilitic disease of the placenta is *hydramnios*, excessive formation and accumulation of the amniotic fluid, associated frequently with small size of the fœtus. It has to be noted, however, that this association is not constant; on the one hand, the reverse conditions of *hypannios* has been recorded in syphilitic cases; on the other, hydramnios may occur in the absence of syphilis; thus cardiac defects in the fœtus may be a cause.

Other disorders of the fœtal adnexa may here be briefly noted. The cord may be abnormally long, and thus liable to be knotted, with more or less vascular obstruction; or, looped around the fœtus, it may be, causing compression, atrophy, and grooving of the body or limbs; it may be excessively short, arresting the movements of the fœtus and seriously interfering with labor. The *amnion* may be imperfectly developed and have undergone fusion with the fœtus, causing the development of bands, and by pressure arresting the development of one or other area, or, contrariwise, as indicated by Klaussner,¹ by setting up a certain grade of obstruction to the venules and lymphatic return, may induce localized giant growth; the *chorion* may exhibit abnormal vascularity.

DISEASES PECULIAR TO THE FOETUS.

There remain several disorders of the foctus, which Ballantyne would class together as "idiopathic," although, seeing that, in connection with almost all of the group, it is to be noted that instances occur of several members of the same family being affected and of the condition being transmitted indifferently along either parallel line, the term is, perhaps, unfortunate. Such states, in which there is this strong hereditary tendency, as, for example, elephantiasis congenita, ichthyosis, tylosis, hypertrichosis, achondroplasia (foctal rickets, osteopsathyrosis), congenital goitre, etc., we shall deal with best under the heading of Abnormalities of Development of Individual Tissues. At the same time, we freely admit the difficulty of exact classification of such disorders. Just as immunity may be inherited in some, acquired in others, so it is with these states; achondroplasia and micromelic (or short-limbed) dwarfism may crop out through a long series of generations; this is a definite inherited diathesis; it may, as indicated by Charrin and Gley's³

¹ Ueber Missbildungen der menschlichen Gliedmassen, Wiesbaden, Bergmann, 1905.
 ² Compt. rend. de la Soc. de Biol., 10 S., 2: 1895; 705, and 3: 1896; 22 and 1031,

MORBID CONDITIONS ACQUIRED DURING PARTURITION 225

results upon poisoning the male rabbit with the toxins of diphtheria or tubercle, or blue pus bacilli, be due to acquired modification of the parental germ plasm, and we are prepared to find that intrauterine (maternal) influences may produce like effects.

Of yet other conditions appearing sporadically, such as general foctal dropsy, ascites, hypertrophic stenosis of the pylorus, foctal endocarditis and peritonitis, foctal nephritis, congenital obliteration of the bile ducts, we know too little regarding the causation or causations to speak with any authority.

THE CAUSES OF MORBID CONDITIONS ACQUIRED DURING PARTURITION.

These may be briefly noted. They are either mechanical, traumatic, The mechanical causes are more particularly stranguor infectious. lation by short or shortened cord, abnormal shortness of cord obstructing descent, and undue narrowness of pelvic channel leading to the death of the child from exhaustion. Prominent among the traumatic disturbances are laceration and amputation from manual and instrumental aids to delivery; cephalhematoma, from rupture of vessels of the scalp through the intense compression of the head and the congestion produced by prolonged arrest of the partly delivered head at the external ring; distortions and partial fractures and dislocation of the skeletal parts; birth palsies, and hydrocephalus induced more particularly by instrumental injury. Among the infectious causes are the presence of pathogenic organisms in the vagina, notably the gonococcus (leading to gonorrheal ophthalmia) and the microorganisms of suppuration. Here, also, must be included improper treatment of the cord at the time of section, leading to local suppuration, infective icterus, and general pyemia.

CHAPTER IV.

MONSTROSITIES AND ABNORMALITIES.

HERE rather than elsewhere would seem most fitting to pass in review the subject of monstrosities and abnormal developments in general.

Definition.-The study of monstrosities and abnormalities, their structure, relationships, and mode of causation, is one that lies on the borderland between anatomy, embryology, and pathology. The pathologist is concerned, inasmuch as the conditions clearly represent departures from the normal, and certain of the minor grades of aberration lead to very definite disturbances in postnatal existence, as, for example, the long series of cardiac anomalies, and the morbus coeruleus (the "blue disease"), to which they give origin. But many again which are compatible with continued existence have no disturbing effect upon the health of the individual-as, for example, supernumerary digits; and many again, the greatest aberrations of all, are wholly incompatible with postnatal life. The subject is in itself fascinating; the departures from the normal are many of them so extraordinary that one seeks to know the cause; the more examples we encounter the more evident does it become that they arrange themselves into well-defined groups, the members of the groups showing orderly gradations, so that these monsters can be placed in classes and species as regularly as can the various forms of animals and plants. With this regularity it is obvious that underlying their development there must be a law, or laws. And the student is led onward to make himself familiar with the laws of normal development and the facts of human and comparative embryology in order to arrive at a satisfactory grasp of the meaning of these remarkable developments.

This is, it must be confessed, but a side path in the study of general pathology; it, however, occupies the same relationship to anatomy and embryology. We know no text-book on any of these subjects in which it receives due treatment.¹ And if we appear to accord it to space out of proportion to its importance, that is because to make ourselves intelligible, we have to enter into anatomical and embryological considerations. As it is, we do at most discuss the main varieties of monsters and abnormalities.

The terms monstrosity and abnormality are employed to denote grave anatomical departures from the normal, whether general or local (affecting but one section or part of the organism). There is no

⁴ In their recent *Text-baok of Embryology*, New York, 1009, Bailey and Miller devote a chapter to the subject, but this is based upon the older rather than the more recent studies on this subject.

ABNORMALITIES OF EXCESS

clear demarcation between the two. All monstrosities are anomalies, or, better, abnormalities (for, as we have already insisted, they are subject to $\nu\sigma\mu\sigma$, law); not all abnormalities are usually regarded as monstrosities; or, otherwise, the slighter aberrations are classed as abnormalities, the more pronounced as monstrosities. To the student who approaches the subject for the first time the number of forms seems as perplexing as the terminology is bewildering. But, as stated, each form finds its place in a general scheme. Two great divisions are to be made out: abnormalities of *excess* and those of *defect*, and each of these may be of one or other order—increase or decrease in *size* or in *number*. There are one or two other classes to which we shall refer briefly that do not come into this scheme, viz., those of *transposition* of parts and *hermaphroditism*; the vast majority are included in the two great divisions.

There is lacking a comprehensive and modern work on the whole subject in the English language. Hirst and Piersol's Monstrosities is valuable so far as it goes, and Ballantyne's pioneer work upon Antenutal Pathology is invaluable for the light it throws upon many conditions. The most masterly résumé of the subject is the article by Marchand in the last edition of Eulenburg's Real Encyclopedic. There is at the present time in the process of publication what promises to be the leading work upon the subject, viz., Schwalbe's Morphologie der Missbildungen (Jena, Fischer, 1906 and 1907). Taruffi's Storia di Teratologia, while most complete, has the disadvantage of introducing a complete new terminology, based upon a theory or theories of causation and relations that have already been found incorrect in several respects.

ABNORMALITIES OF EXCESS.

1. General Excess.—This may be (a) universal, as in true giantism; or (b) lateral, one-half of the body exhibiting greater development than the other (as though the first two blastomeres of the ovum had been disproportionate in their potentiality; or (c) local, one member or one organ being developed out of all proportion to the rest. These latter conditions will be referred to under Hypertrophy.

As to the causation of giantism we know little that is assured. While there exist families of which, through several generations, the members have been noted for their height, heredity is nevertheless infrequently observed in pronounced cases of giantism.

The minimum for true giantism is generally accepted as 6 feet 6 inches, or about 200 cm. Thoma places it at 188 cm. (or 6 feet 3 inches), but in Anglo-Saxon countries this is scarce considered giantism.

The giant exceeding 7 feet in height comes, as a rule, of a family whose members have been of medium height. In such giants, the excessive height is generally due largely to disproportionate length of the lower limbs. There is some question nowadays as to where the

MONSTROSITIES AND ABNORMALITIES

line is to be drawn—if it really exists—between such true giants and morbid giantism, a type of acromegaly. The more recent autopsies on giants have, in general, revealed enlarged pituitary bodies, while von Hansemann noted that in almost all of the many giants examined by him there has been evidence of rickets, and suggests that a grade of rickets insufficient to inhibit growth is followed by extreme bony development, especially of the limbs.

Regarding localized giantism, a distinction must be made between those cases that are apparently of inherent nature and those due to disturbed nutrition, even if these be often of congenital origin. In congenital elephantiasis and macroglossia, we have to deal with lymphatic obstruction, accompanied by secondary overgrowth of the connective tissues. Allied alterations in the relative vascular supply best explain, it may be, some conditions of the nature of macrodactuly. But where, as in some cases of gigantic hands, there is observable a tendency to duplication of fingers, the indications point strongly to a redundancy of vegetative matter in the anlage or growing point of the The same is clearly to be invoked in cases of premature and part. excessive development of the different components of the generative organs, although here, of late, more than one observer has called attention to an association between hypertrophy and new growths of the adrenals and precocious maturity and development of the external genitalia.1 Abnormal inheritance, associated with modifications, it would seem, of metabolism, best explains infantile goitre, hypertrichosis (or hairiness), and lipomatosis, or generalized obesity.

2. Numerical Excess.—A wide and varied range of conditions is to be included under this heading—from triplets on the one hand, to partial deduplication of a terminal phalanx on the other.



Heteroophal hen's egg with two embryos of sixth day), each with separate yolk. (Panum.)

Twins (Gemini Æquales).—If individual twins in themselves cannot be regarded as monstrosities, their appearance in a well-regulated human family is usually regarded as anomalous, and certainly the line of demarcation between twins of a certain order and some of the most *bizarre* of monsters is very slight.

We recognize two orders of twins: the *dichorial*, or heteroophal, and the *monochorial*, or monoophal. In the former, each child is born with a


separate "caul;" the sex may or may not be identical, and the features, configuration, and characters, while at times approximating, at others present wide differences. Such dichorial twins, obviously, originate

¹ An interesting study and record of cases of this and the allied association of these conditions with pituitary and pineal growths is afforded by Guthrie and Emery, Trans. Clinical Soc. Lond., 40: 1907: 175, 1907; 174; see also Bulloch and Sequiera, Trans. Path. Soc., 1905.

from two separate ova, fertilized at the same menstrual period, each of which, in its development, forms its own set of membranes, cord, and placenta, though ultimately the two placentas may fuse.

In the rare conditions of *superfactation* the ova are, it would seem, discharged and fertilized at different menstrual periods.

The rule is that each Graafian follicle contains a single ovum, but even in the young child, and in a large number of different species of animals, two, and sometimes three, ova have been observed in a single follicle. In those cases in which heteroophal twins are of the same sex and exhibit great similarity, the possibility must be borne in mind that they have been derived from a single follicle; nay, more, that they are due to early complete division of a single ovum into two independent ova after fertilization, but before implantation in the uterus.



Monoophal duck's egg with two embryos (of seventh day) upon a single yolk, (Panum.)

Monochorial twins are the less frequent. Ahlfeld's statistics are, that of 506 twin births, 444 were dichorial, 62 monochorial. They emerge from a single caul or chorionic sac, have a single placenta, are always of the same sex,¹ and, when equal in development, are remarkably alike. The condition is not hereditary, whereas in certain families there is a distinct tendency toward double, dichorial births.

The chorion, it will be remembered, is the outer wall of the ovum, and a single chorionic membrane enclosing two embryos can only mean that a single ovum has given rise to two individuals. Not infrequently the twins show identical abnormalities; right-sided hydrocele (Ahlfeld's case), spina bifda (d'Outrepont), hypospadias (Lehmann), etc. In cases of dichorial twins, there is never any indication of the separate chorions fusing into one (unless the rare and doubtful cases above noted of monochorial twins of different sex are to gain this explanation). The

¹ One or two exceptions to this rule are recorded, but these, if we mistake not, in the older literature only

MONSTROSITIES AND ABNORMALITIES

non-existence of haphazard fusion of the two halves of the double monster is against the idea of chorionic fusion, for did this occur we should expect to meet with occasional fusion of the embryos in various positions, the back of one to the side of the other, etc.

These cases, therefore, prove that a single ovum is capable of giving rise to two individuals, a conclusion wholly borne out by observations on the eggs of lower animals and by experimental diplogenesis.

Among the lower animals, multinucleated ova have been observed by several (Francque, Stockel, H. Rabe);⁴ on the other hand, numerous zoologists (Roux, Endres, Morgan, in the frog; Herlitzka, in the newt; Driesch, in sea urchins; Zoja, in the jelly-fish²) have proved that division of the ovum into its separate blastomeres at the two-, four-, eight-, and in the medusa even in the sixteen-cell stage, may give rise to separate,




Diagram of A, monochorial twins, each with separate amnion (dotted line) lying in a common chorionic sac; B, dichorial twins, each with its own chorionic sac.

though dwarfed, individuals from the separated blastomeres. In the newt, even as late as the gastrula stage, Spemann³ was able to gain two complete embryos by experimental division.

It is possible, therefore, that monochorial twins originate (1) from the separate fertilization of the two nuclei of one ovum, or (2) from the fertilization of an ovum with one nucleus, with subsequently (a) separation of the primitive blastomeres, or (b) cleavage, not of the ovum itself, but of the germinal area developing later upon that ovum; or, in other words, the formation of two primitive streaks upon a single germinal area. Without going into details, we may here say that the study of double monsters indicates that the period of origin is more

¹ For literature, see Schwartz, Anat. Anz., 18: 1900.

² For literature, see E. W. Wilson, *The Cell in Development and Inheritance*, second edition, New York, Macmillan, 1906.

³ Sitzungsber. d. Phys. med. Gesellsch., Würzburg, 1900.


PLATE VII

FIG. 1




Skiagraph of a Foetus Amorphus. (Charlton.) Showing presence of vertebral column and ribs, but absence of head and limbs.





Two Similar Examples of Fœtus Amorphus.





Fœtus Acardiacus Acormus, showing Vascular Relationship to the Umbilical Cord of the Healthy Fœtus. (Barkow.)

Photograph and Skiagraph of a Fœtus Acardiacus Acephalus. (Schwalbe.)



ACARDIAC MONSTERS

likely to be late than early, as does also the existence of a single chorion. There is more likelihood for identity where only one spermatozoon is concerned than where two.

Regarding these monochorial twins, it is to be noted that the amnions may be separated or fused, that the placenta, almost without exception, is common, that the umbilical cords may be well separated or inserted close together, or have for a varying distance a common amniotic sheath.

Unequal Monochorial Twins (Gemini Inæquales).—Fœtus Acardiacus.—Occasionally we encounter a curious abnormality. Instead of equal twins being born, there comes forth a single, well-formed individual, accompanied by, it may be, an amorphous lump of flesh, contained in the same sac and only recognizable as a product of conception

by possessing a distinctive small umbilical cord. Such is a fatus amorphus. Most often more complete organization can be made out in the mass; in general, the lower half of the body, with its limbs, is fairly developed (Acardiacus acephalus), or the head end, it may be little more than the cherub-like head itself (Acardiacus acormus), or, lastly, there may be a foetus fairly complete as regards its skeletal parts, and, unlike the other forms, this may possess a partly developed heart; but this and the other internal organs are imperfectly developed and the tissues are extremely cedematous (A. anceps).

A subgroup is to be recognized, including the cases of *A. acormus*, in which the foctus is implanted directly upon the placenta without the

intervention of an umbilical cord. It is still a matter of debate whether the mode of origin in these cases is the same as that of the others; all, it may be added, have separate amnions.

The causation of this type of monstrosity has been worked out by Meckel, Claudius, and Ahlfeld. In the first place, the course of the circulation is reversed. This explains the arrested development of the heart; the acardiac fœtus is nourished by the blood of its more developed and stronger brother. There is fusion and anastomosis of the allantoic and, later, placental vessels, so that the arterial blood' of the stronger fœtus (A), driven into the branches of the umbilical arteries of A, enters the branches of the umbilical artery of B, and, being under greater pressure, forces itself into B's aorta; passes into its branches and nourishes the various tissues of B; arrests the action, and leads to the atrophy B's heart; may be but sufficient in quantity, in quality, and in distri-

⁴ It must be remembered that this blood while in the umbilical artery of A is in quality venous, returning from the foctus to the placenta.




Developed twin and acephalic monster and their relationship to a common placenta. (Ahlfeld.)

MONSTROSITIES AND ABNORMALITIES

bution to nourish portions only of B's economy, whereby the other portions undergo aplasia and atrophy. 'The greater frequency of maldevelopment of the cephalic half of the body is apparently due to the fact that for a period the heart of B, repelling A's blood, continues the circulation of the upper half of the body with its own imperfectly aërated blood.

The above theory, worked out by Ahlfeld, whose diagram is here given, is that which has received wide acceptance; but, as pointed out by Marchand and Schwalbe, there are difficulties in accepting it in its entirety.



F16. 55



Schema of mode of development of acardiac monsters. (Ahlfeld.) "

In the first place, the human embryo would not seem to possess the free allantois assumed by Ahlfeld; the teratogenic period must, then, be placed at a later date; in the second, if all hemi- and holo-acardiaci originated at the same early period, it is difficult to u-derstand the singularly wide range of extent of development or retrogression. With Marchand, we must for many cases accept Dareste's view, that in a large proportion of cases (in the chick), where two primitive streaks develop upon one ovum, one of the two tends to be imperfect, and may presuppose that the imperfection extends to the heart and vascular system. The theory of

Schatz¹ is gaining adherents: It appears to me to be too elaborate and involved to be correct, though it may well be that my difficulty in following the lengthy exposition of the same


is at fault. The outcome of the arrest of the individual circulation of the weaker fortus may lead to its death long prior to full term. In this case, pressure, autolysis, and absorption may lead it to assume the form of a flattened mass within the membranes of the developing twin, and may result in a *fatus papuraceus*.

Triplets and Other Multiplex Births.—These may be mono-or polychorial, or a combination of monoand dichorial.

Wilder² gives careful notes of a case investigated by him of triplet sisters, evidently monochorial, so remarkable



Development of two embryos of the chick on a common germinal area, the lower of the two imperfect from the first (paracephalus,) (Dareste,)



Marchand's schema of mode of development of Acardiac monsters. A, the chorion ch is already developed; the yolk sac d has divided into two unequal halves, d⁰ and d², in consequence of which the one embryo a² receives through the yolk vessels more abundant nourishment than does the other embryo a². As a result, its allantoic circulation develops more actively, and the later developing allantoic artery of the smaller embryo anatomoses with it. He, later stage; the allantoic vessels of a¹ usurp the whole of the chorion, the smaller embryo gaining its blood entirely through the anatomosis of its allantoic artery with that of a¹, and this, therefore, in the reverse direction to the normal current.

¹ Archiv f. Gynäk., 1901.

² Amer. Jour. of Anat., 3: 1904.

in their identity that not even the mother (contrary to her conviction) could surely identify them; for on one occasion two of the sisters, in answer to inquiry, stated that they had not received their bath that morning, while the third gleefully announced that she had had three.


The largest number of simultaneous human births recorded by a reliable authority is seven; some few cases of quintuplets are on record, whether monochorial or not we have no adequate evidence. In the cat, the monochorial development of five kittens has been ascertained.

DOUBLE MONSTERS.

It will, we feel assured, materially facilitate a comprehension of the many and different forms of double monsters if, instead of first describing the more important forms, we first establish an adequate theory of causation and mode of development, and, laying this down, then proceed to classify them in terms of that theory. For close upon two centuries the explanation of diplogenesis has interested zoölogists and physicians alike, and each step forward in our knowledge of embryology has led to the propagation of new theories, which in general have, upon investigation, been found to be inadequate. Nevertheless, with increasing knowledge those theories have come to satisfy and embrace a larger and larger number of conditions, and, especially during the last quarter of a century, we have approached a rational classification.

The names that especially stand out in the study are those of Geoffroy St. Hilaire (1836), whose classification is still quoted; Forster (1861), who collected and classified what at the time was the fullest series of cases; Dareste (1877–1891), the pioneer in the study of experimental diplogenesis; Fol, whose observations on polyspermy were, nevertheless, in a wrong direction; and, among the more immediate moderns, Morgan, of Bryn Mawr; Roux, of Breslau; E. W. Wilson, Jacques Loeb, etc., upon the artificial separation of blastomeres; Rauber, on the "radiation theory;" Marchand, with his theory of fertilization of polar bodies; and Fischel, with his "head and trunk centres" theory. Many other names deserve quoting, but these are, perhaps, the foremost.


The difficulty thus far in propounding any adequate theory has lain in the fact that the more cases are studied the more surely it is seen that double monsters belong to two main categories. This has been a matter of contention since 1724. There have been those who would explain all examples as cases of *fusion* of two separate embryos lying in approximation; others equally convinced that *cleavage*, or dichotomy, of an originally single individual, is the essential cause. It is in the attempt to embrace both classes that, one after another, the various theories propounded have shown their weakness. While we realize ' that it is only just to the student, as to the older workers in this subject, to afford an account and criticism of previous theories, to set these forth adequately would take great space. For that reason, and because,



Diagrammatic Representation of Fischel's Theory.

The head centre H_t formed by the cells that first become invaginated, the trunk centre P_t formed by the fusion of the cells that undergo later invagination.





Mode of Origin of Different Orders of Double Monster According to Fischel's Theory.

A, monochorial twins without fusion; B, according to exact area of fusion (the head centre giving origin to the greater part of the body), various grades of craniopagus (head fusion), sternopagus, and xiphopagus (body fusion), passing on to conditions of anterior deduplication; C, the slighter cases of anterior deduplication by late dichotomy of the head centre: D, posterior deduplication by lack of fusion of the lateral components of the trunk centre.



in addition, we believe that it is less confusing for the student, we venture to put forward a single hypothesis, which we may term the "growing point" hypothesis,¹ it differs from those which have hitherto gained the fullest acceptance, viz., those of Rauber and Fischel—theories which likewise are based essentially upon a comparative study of embryology and teratology and experimental diplogenesis—in that it takes into consideration the mode of cell multiplication.

In the lower animals, as already noted, it is possible by various means to loosen and shake apart the earliest blastomeres of the fertilized ovum. Each then becomes, and develops as, a separate individual. Were this to happen in man, each separate blastomere would develop its own chorion and sac, whereby the contained embryos would be prevented from fusion. At most, as already suggested, such early separation would lead to identical dichorial twins. We have, therefore, as Schwalbe has most serviceably pointed out, to look to a later period; and, instead of attempting to ascertain what is the earliest must strive to establish what is the latest period at which double monsters can develop (Schwalbe's teratogenic termination period). To do this, it is best to consider the mode of development of the embryo of the higher vertebrates. This mode is not identical in the oviparous and viviparous forms, or, more accurately, in the lower vertebrates and the amniota (mammalia), respectively, but a common principle includes all. may thus first study a simple type, such as the fish. Here the whole ovum does not give rise to the embryo. Certain cells only at one pole form the germinal area; the rest, filled with yolk, afford food to the eventual embryo. Nor does the whole germinal area give rise to the embryo. This, as it grows, spreads over the yolk, and at one point, as pointed out first by His, that spreading growth becomes arrested, the edge elsewhere continuing to advance (Fig. 58). Hertwig has laid down that the sinus thus formed represents the cavity of the gastrula of holoblastic forms. It is the cells along the edge of this depression that give origin to the embryo, the primitive streak arising along the line of the same.

Rauber (1877) based the whole theory of diplogenesis upon this mode of development. He presumed the simultaneous appearance of two invaginations which, in the process of development underwent fusion, giving rise to a double monster. The accompanying figures from Kopsch gives the idea of the mode of development of superior deduplication (anadidymus) (Plate IX). Fischel, starting from the same basis, points out that Rauber's conception is false in so far as the embryo is not formed by a continuous infolding of the edge of the blastoderm behind the head end; on the contrary, at a relatively early period the cells of the two sides of the groove join to form the trunk or body centre, and that these two centres, head and trunk, between them constitute the whole embryo at this period, and between them form the whole

¹These earlier theories are given fully by Schwalbe, 'Die Missbildungen, Jena, B. Fischer, Pt. 2, 1907, and are carefully criticised by Fischel, Verh. d. Deutsch, Path. Gesell., Karlsbad, 5:1903; 272.


eventual individual. Starting from this, Fischel also ascribes the vast majority of double monsters to fusion, as indicated by the accompanying diagrams. He recognizes, however, that certain katadidymous forms, double below but single above, may be due to want of primary junction of the two halves of the trunk centre. This is not cleavage, but lack of



Diagrammatic representation of growth of germinal area of the fish egg as a cap spreading over the yolk, with arrest at one point, and infolding to form the primitive streak. union. As regards the slighter cases of anterior duplication—double-headed snakes, for example, and in cases in which merely the pituitary body is divided—he is at an *impasse*; he denies Gerlach's statement that the head centre grows forward, and yet he has to admit that in order to produce these there must be a cleavage of the head centre, by mechanical or other means. He thus is forced to the conclusion that the head centre gives origin to the head alone, the trunk centre being the great formative organ.

I am inclined to hold that had Fischel grasped the nature of growth

he would have escaped this difficulty and have afforded the adequate theory. What I mean is best illustrated by what we know concerning the mode of growth of the plant. Briefly, this exhibits two primary growing points, the superior giving origin to the stalk and its appendages, the inferior giving origin to the root, and it is from the cells of these growing points that eventually all the component cells and com-



Mode of origin of primitive streak and anlage of eventual embryo according to His,

ponent parts of the plant are derived. In the plant the cells of these growing points persist and are active throughout life, and their position is, in one sense, fixed; that is, the giant trees of California have attained their height, not by growth forward of, but by growth backward from, their growing points. The primary growing point forms the apex of the plant; its cells are vegetative; they undergo repeated division in such a way that, dividing, the outer daughter cell remains vegetative, retaining

THE GROWING-POINT HYPOTHESIS

the position of the mother cell, the inner or its descendants becoming converted into the eventual specialized tissue cells. In this way there persists a layer of vegetative cells ("cambium"), the cells given off from which become, in series, the primitive cells of successive regions or segments of the trunk. The growing point does not grow forward; it is *projected* forward by the intercalation of cells derived from its daughter cells, and it continues to occupy a fixed point in relationship to the rest of the economy.




Diagram of section of the growing point of a plant, A, A, the apical cells, which continually divide, giving off backward a series of cells; B, C, D, which cells divide as in B^0 , B^0 , to form the cells of the vegetative or cambial layer, these cells again, as in C, \mathcal{O} , and \mathcal{O}^0 , divide at right angles to the former plane to give origin eventually to the functional cells of the stem (or root); D, develoment of a secondary growing point.

Now, in plants, as all will have noted, we meet with cases in which the growing points have undergone cleavage at either superior or inferior pole; the monaxial trunk of a fir tree may, high up, divide into two or three equal trunks; the radish may fork, and so on. In many species it is natural for the growing point to divide early into several secondary growing points, with the production of many stems or numerous equal rootlets. The plant, it must be remembered, has not bilateral, but radial, symmetry; this notwithstanding, there is a tendency toward (bilateral) dichotomy, due, we would suggest, to the fact that cell division is binary; if there were a single apical growing point cell, each of the two products of division might assume equal properties and become the progenitor of independent growing points.

So, in animals, what Fischel terms the head and trunk centres are the superior and inferior growing points. These, once formed, do not

grow, but are projected apart; it is the successive daughter cells given off from these that give origin to the different segments of the body and their contained tissues, sundry of those daughter cells becoming secondary growing points for the lateral organs. Nor is this matter of the superior and inferior growing points a purely hypothetical conception on our part. Already Assheton' had demonstrated the existence of the same in the rabbit, frog, and chick, and incidentally and antecedently had proved the truth of Fischel's contention that the "trunk centre" gives origin to the greater part of the body. Thus, for example, he inserted a fine sable hair into the mid-point of the blastoderm of the unincubated hen's egg, and then, incubating the egg, found the hair to be situated at



Segittal section through head of individual with attached epignathus: a, epignathus; b, region of attachment at hypophysis; c, cerebrum of autosite; d, corpus callosum; e, cerebulum; f, vertebral colum; g, upper, and k, under lip, of autosite; i, torgue. (Schwabe).


the anterior end of the primitive streak; in fact, at the level of the first pair of somites. From this he concluded that from the superior growing point are developed the brain, medulla, heart, and foregut; from the inferior, all the rest of the body, *i. c.*, the primitive streak area. There is, however, this difference between the animal and the plant, that whereas in the latter the primary growing points may be active through the whole of existence, in the former they cease to function as such so soon as the anlage or matrix of the main axis is laid down. So soon as this occurs, further increase in length is only by intercalation.

Polar Hyperplasia and Serial Deduplication.—For, suppose this were not the case; that the medullary groove became laid down and

¹ Quart. Journ. Micros. Sci., 37: 1894; 113, and Proc. Roy. Soc., 60: 1896: 184.

POLAR HYPERPLASIA

the expansions of the neural canal had formed themselves, which are destined to give origin to the brain. Then any further production of daughter cells by the superior growing point could only give origin to a mass of tissue to which no eventual function could be assigned by the economy. We know, that is, that the site of the superior growing point in the developed organism is in the region of the sella turcica. The interesting point is that this "absurdity," in the Euclidean sense, does occasionally happen; or, at least, this conception best explains a very remarkable series of monstrosities. There are cases in which, adherent to the region of the sella turcica and projecting through the mouth of the newborn child, there is a large tumor. In the most advanced cases these masses contain bone, and may even show limbs, and are found to contain, when studied, representatives of organs



Section through a congenital sacral teratoma, (Nakayama,)

derived from epi-, meso-, and hypoblast. Such cases are known as *Epignathus*. The simplest conception of the origin is that they are due to redundant growth of the superior growing point after the anlagen of the tissues of the individual have been laid down, and that the "tera-togenic termination period" of the production of this form coincides with the development of the "neurula" stage of the embryo; the earlier the period and the more active the proliferative powers of the growing point, the more complete the series of tissues contained in the epignathus.

Similarly, springing from the site of the inferior growing point in the perineal region behind the rectum, we encounter another remarkable form of tumor, having the same general characters—the so-called *congenital sacral teratoma*. To this a like origin must be attributed.

These forms have commonly been regarded as examples of foctal inclusion, *i. e.*, of the inclusion of one embryo within the tissues of another. But no valid reason has been adduced why this inclusion should occur in these particular regions. Schwalbe concludes that

there must be "Versprengung," or splitting off of early formative cells, a view which approaches that here given, but, again, does not explain why this should be particularly apt to occur in these particular regions. I regard these forms as homologous with the cases among plants; in the garden rose, for example, where an imperfect flower stalk and head develop from the centre of another flower.

Polar Dichotomy.—Granting that up to the beginning of the neurula stage (when the medullary groove shows itself) the vertebrate embryo presents polar growing points, we now gain the teratogenic termination period for another group of monsters. It has, indeed, been shown experimentally by Spemann, in connection with the newt's egg, that up to this period it is possible to produce both the complete embryos and double monsters; later, this is not possible.

Spemann produced the latter by tying a fine hair around the developing ovum in the plane of the future long axis. He thus obtained various grades of deduplication. In other words, up to the welldeveloped gastrula stage it is possible that from a single embryo there may be produced dichotomy, or bifurcation, so that we may by this process gain:

1. Superior dichotomy, affecting the superior but not the inferior growing point, so that the upper end of the body is to a greater or less extent doubled' (Fig. 65, II and III).

2. Inferior dichotomy, affecting the inferior pole alone.

3. Both superior and inferior dichotomy, portions of the trunk region alone possessing a single axis (Fig. 65, VI).

4. Mesial deduplication, the superior and inferior growing points remaining single, but their cells projected backward from them on



Spemann's method of producing various grades of double monster from the newt's egg by tying a fine hair in the line of the longitudinal axis of the germinal area. either side, failing to unite, or becoming by mechanical means split asunder *mesodidymus* (Fig. 65, VIII).

5. Complete cleavage along the whole plane, giving rise thus to two embryos lying parallel, each provided with its individual longitudinal axis (Fig. 65, VII).

Regarding these figures, a few words are necessary with regard to the distinction marked between early and late dichotomy. By early dichotomy, I mean that the cleavage occurs so early² that, with separation between the grow-

ing point cells at one apex, all the cells which so far have been given off from one or other side of the growing point undergo cleavage,

¹The terms *anadidymus* and *katadidymus* have been in the past employed for conditions of superior and inferior deduplication, respectively, but this with so much confusion that there is now a definite movement to banish them.

² From Assheton's observations, already recorded, it is obvious that this cleavage must antedate the period of development of the primitive streak.

POLAR DICHOTOMY

which, indeed, must be held to extend into the area of cell proliferation from the growing point of the other pole, so that as growth continues





Diagram of various forms of dichotomy or cleavage: I, normal early primitive streak in germinal area with S. G. P. superior, and I. G. P., inferior growing point. II, result of early dichotomy of superior growing point. The separation affecting also the lateral rows of cells given off from the inferior growing point; III, late dichotomy of superior growing point, only the cells given off more recently from the two superior growing points affected. IV and V, similar results of early and late dichotomy of the inferior growing point. VI, relatively late dichotomy of both superior and inferior growing points—uankatadidymus; VI, early (complete) dichotomy involving both growing points—fused double monsters, lateral tusion, VIII, mesolidymus, the growing points remaining single, but the series of cells derived from them on either side undergoing separation, 16

two axes of symmetrical cell development diverge from this other pole —from which it results that when the anlagen of the primitive streak and structures are laid down they are doubled, and as the anlagen are doubled so from this time onward, the tissues derived from those anlagen are doubled. The result, it is true, is apparent fusion, but there are distinctions to be noted.

In what is referred to as late dichotomy, there, I hold, we have to deal with the origin of cleavage at a period when the proliferation behind either growing point has been so extensive and so long continued that between the two growing points a well-developed primitive streak has been laid down. According to our conception, already recorded, the

cells first derived from the growing points—the oldest—will become those farthest removed, while those immediately behind the growing points will be of most recent de-




Late dichotomy of superior growing point. Doubling of optic vesicles and forward part of head only of newt embryo in one of Spemann's experiments. (See Fig. 64.)



Early superior dichotomy in larval newt, from one of Spemann's experiments.

velopment. If this cleavage is brought about at a later stage it will affect these more labile and less fixed cells alone.


It follows thus, that I regard as probable examples of polar deduplication an important series which Rauber, Marchand, and Fischel and othern modern observers class among examples of fusion. Even granting what must be granted freely, that cases of superior and inferior cleavage show, upon study, a much more extensive doubling of their axes than external appearance suggests, and that in the former instances more particularly the chorda is often double as far as it can be traced inferiorly, even this does not prove the fusion of two primarily separate individuals. For the inferior growing point is not situated at the end of the tail region, but in front of that in the region of the anus (in man). Only where we have evidence throughout the length, of the existence of two longitudinal axes, are we absolutely assured that there have been





POLAR DICHOTOMY

two complete individuals which have fused. In some fish monsters we have, I think, evidence of this, but they are in the minority, and those conform in the lateral deviation of the fins, etc., to the principles we see in action in those cases in which there can be no question regarding fusion (Janus forms, etc., see p. 248).



Dicephalus tetrabrachius. (Ahlfeld's Atlas.) Skiagraph of a Dicephalus tribrachius, Museum, McGill University. (Dr. Girdwood.)

Tricephalic monster. (Ahlfeld's Atlas.)

In this class, therefore, I include all cases of double monstrosity in which, at any part of its course, even if it be only at the extreme end, there exists a single median axis. Of **superior deduplication** we possess a greater series of forms from mere bifurcation of the hypophysis cerebri through those showing more or less imperfect deduplication of the head (Figs. 68 and 69), complete double head (Fig. 70), dicephalus tribrachius (Fig. 72), to dicephalus tetrabrachius (Fig. 71).

Of inferior deduplication the slightest cases are (not those of double coccyx, but) of deduplication of the external organs of generation. Ballantyne and Skirving' have brought together some twenty cases of double penis. Next to this, we pass through more and more perfect examples of katadidymus tripus up to katadidymus tetrapus.

In both these series of cases there is a single umbilical cord, indicating a common yolk sac, *i. e.*, development from a single ovum.

Cases of anakatadidymus (Fig. 75) and mesodidymus are distinctly rare.

Fusional Deduplication.—In this order are to be included all those cases in which, whether the two elements are equal or unequal, each has



Katadidymus tripus, (Ahlfeld's Atlas.)

its individual main axis, not combining with that of the other, even if, in some cases, the two closely approximate. Keeping in mind the distinction between equality and symmetry, the latter term referring to the position relative to a common plane, we can make the redivision into (1) symmetrical diaxial double monsters, equal and unequal, and (2) non-symmetrical biaxial double monsters, these being always unequal. Of these, the first form is a very welldefined class.

Biaxial Symmetrical Double Monsters. — There are certain very striking features regarding these, which include the majority of double monsters: (1) First and foremost their symmetry: like part is applied to like. We never, for example, encounter monsters applied

side to side in opposite directions, or head to tail (the nearest approach to such is in the cases of epignathus and congenital sacral teratoma already discussed. (2) We may find them chest to chest (ventral fusion), head to head, rump to rump, side to side, but never wholly back to back with bony fusion.

These facts in themselves prove that they are derived from a single ovum and possess like polarity; had we to deal with two distinct ova that had fused, however early, such symmetry would be a chance occurrence, not a law.

We know that the ordinary egg has polarity. Take, for example, a

¹ Teratologia, 2; 1895, 92, 184 and 295.

hen's egg that has been incubated twenty-four to forty-eight hours, hold it with the blunt end to the left, the sharper end to the right; the developing chick is always found with the longitudinal axis at right angles to the main axis of the egg, and the head away from the observer.

In all the higher animals the embryo develops, if not on the surface of the ovum (fishes, birds), at least in a very definite relationship

to that surface (mammals); its ventral surface is always toward the yolk, its dorsal surface toward the surface of the ovum. Did two ova fuse, the commonest form of fusion should be one or other grade of back-to-back union. This is the rarest, and when it does occur, it is partial and easily explained. All symmetrical biaxial double monsters have had their ventral aspects directed toward a common yolk.

This conclusion again gives us an indication regarding the period teratogenesis. They have of originated from a common ger*minal area.* At what period it is difficult to state with precision, save that the termination of the period must coincide with that of "complete dichotomy," already noted, viz., the separation of germ matter to form two individuals cannot be later than the end of the gastrula stage; the probability is that it is frequently much earlier. The fusion of the two, resulting in monster formation, may, as we shall show, occur at a definitely later period. This conception, it will be seen, gives a common origin for all types of double monster-and thus harmonizes the two opposing theories. We have the three stages:

Anakatadidymus of sheep. (d'Alton.)

1. Partial dichotomy, giving origin to cleavage deduplication.

2. Complete dichotomy of germinal area, followed by subsequent fusion, giving origin to fusional deduplication.

3. Complete dichotomy of germinal area and yolk, giving rise to monochorial twins.

Granting these postulates, it will be seen that there may be the fol-





Diagram to demonstrate that two embryos developing upon a common ovum will, if they fuse, show a greater extent of fusion ventrally than dorsally.

F10. 77

lowing main varieties of fusion of two embryos developing on the surface of a sphere. As a matter of fact, we encounter all these, and can thus classify them into the following species:

1. Ventral Fusion.—This is relatively common, although frequently it is combined with a certain grade of—

2. Lateral Fusion. — The two individuals not being absolutely apposed, they may be regarded as not having accurately faced each other on the sphere, the radii joining the centre of the ovum to their mid-dorsal surfaces, forming an angle less than 180 degrees.

The greater size and development of the upper half of the body make it that where ventral fusion occurs, the fusion is above the region of the future umbilicus; this is so even in the very slightest cases, *i. e.*, such as xiphopagus, in which fusion is by the xiphoid region alone (the Siamese twins).

As regards lateral fusion, this is always more pronounced ventrally than dor-



Ventral fusion: Thoracopagus disymmetros. Heidelberg Pathological Institute. (Schwalbe,)



Ventral fusion. Skeleton of a thoracopagus double monster. (After Haller.)




Diagram of possible modes of fusion of two embryos developing on the surface of a single sphere (ovum): I, fusion by a common yolk sac (Xiphopauw); II, superior lateral fusion (various grades of Cepholdhoracopauw); II, median lateral (grades of Iekointherocopauw); IV, superior latera (grades of Pyuopagum monosymmetros); V, superior dorsipolar fusion (Craniopaguw); VI, upperior approximation to show asymmetric fusion in consequence of the head curvature; VII, superior apicopolar fusion (Janiceps); VIII, inferior dorsipolar fusion (Pyuopaguw disymmetros). sally: the conception of growth on the surface of a sphere explains why this must be so (Fig. 76).

3. Superior Polar Fusion.—Of this, two distinct subspecies are to be recognized:

(a) Superior Dorsipolar Fusion (Craniopagus).—Here the two individuals are joined by their cranial regions.

We have here a partial exception to the rule of symmetry, and one that throws light upon the period of origin of the monstrosity. The two heads are rarely in absolute conformity; they may be with their sagittal planes at angles of 90 degrees, or even more. Now the frontal, and still more the parietal region, is dorsal to the main axis of the body, and juncture by these regions can only mean that fusion has taken place *after the head fold or curvature of the embryo has been developed*. It is an interesting point that in all vertebrates the head of the embryo is not in the direct line of the main axis, but inclines to one or other side. This lateral curvature is well marked in the chick. Consequently, if two such heads fuse, the regions of fusion do not exactly correspond.



Superior dorsipolar fusion: Craniopagus. (Ahlfeld's Atlas.)

(b) Superior Apicopolar Fusion (Janiceps or Syncephalus).—The picture afforded by the Janiceps monsters is very different, and is best indicated by the illustrations (Figs. 81 to 87). Where the axes are in the same line there are two fully formed faces looking in opposite directions. Where they meet at an angle of less than 180 degrees the one face is imperfect; the more acute the angle, the greater the imperfection, until it may be represented only by a single median ear.

It is evident that here we deal with fusion that has occurred at an early period, before the ventral curve of the head has been developed, so that the growing points approximate. The accompanying diagrams (Figs. 85 and 86) indicate what has happened. In the first is represented the perfect Janus type. It will be seen (as universally admitted) that each face (represented by the lateral conical projection) is formed, as regards one-half, from the one individual, A, as regards the other, from the other, B.

We know that the site of what had been the superior growing point is in the immediate neighborhood of the infundibulum; in other words, that the cells derived from the growing point and those immediately



Apicopolar fusion: disymmetrical Janiceps (Cephalothoraeopagus disymmetros (Schwalbe's case). The two secondary front or facial aspects are absolutely similar.



Fro. 84



Apicopolar fusion at an angle less (or greater) than 180 degrees. Monosymmetrical Janiceps (Cephalothoracopagus monosymmetros) (Vrolik's case), Fig. 83, the perfect secondary front view; Fig. 84, the defective secondary front view, with synotic.

Fig. 83

behind it give rise by outward growth to the facial structures. If, then, cells in the immediate neighborhood of the two growing points become adherent, the only possibility on the part of their derivative cells is to develop in the line of least resistance. The growing points, as such, do not bifurcate, but their descendants grow out laterally on either side.




Diagram of mode of development of a disymmetrical Jamices monster. The longitudinal axes of the component embryos are in the same line. Fusion has taken place early in front of the growing points A and B respectively; as a result, instead of the earls of the head region derived from A growing to reach A', they are diverted by the opposing pressure of the cells derived from B; each face on either side is thus made up one-half from A (A'') the other from B (B'').


When the main axes of the two embryos are not in a straight line, then on the side in which the angle of juncture is less than 180 degrees a certain number of cells of the same embryo destined to develop parts of the face by growth in a laterosuperior direction (a) will, in their growth forward, find themselves opposed and arrested by the equal force with which the corresponding cells (b) of the other embryo are tending to grow in an exactly opposite direction. Where forces, equal and opposite, meet each other, as at a, b, c, d, in the diagram the result is that they neutralize each other; in other words, portions of the facial parts do not develop.

JANICEPS

Fig. 86

A constant feature of these Janiceps cases is that they exhibit also a well-marked grade of ventral thoracic fusion. Why this is so is illus-

Apicopolar fusion at an angle less (or greater) than 180 degrees, with resultant failure to develop parta of the secondary face on the side of the lesser angle.



Superior apicopolar fusion: a, approximation and fusion of the two embryos before the head curve has manifested itself; b, result of growth and development of head curve producing state of cephalothoracopagus.

trated in the diagram (Fig. 87). *After* fusion, each head, for its proper development, must form its ventral curve; this inevitably brings the bodies together ventrally.

Inferior Fusion.—It is noteworthy that we have identical subspecies at the inferior pole.



Superior spicopolar fusion: early stage of development of a Cephalothoracopagus monosymmetros in the chick. (Kaestner.)



Inferior dorsipolar fusion: Pygopagus monosymmetros. (Marchand.)

Fig. 90




Inferior dorsipolar fusion: Pygopagus monosymmetros. (As with superior dorsipolar fusion, the fusion in these cases is rarely absolutely symmetrical.) (Ahlfeld's Atlas.)

252

PYGOPAGUS AND ISCHIOPAGUS

(a) Inferior Dorsipolar Fusion (Pygopagus).—Where the fusion takes place after the ventral curvature of the tail end, it is parts that are dorsal to the main axis that become united.



Inferior apicopolar fusion: Ischiopagus disymmetros. (Ahlfeld's Atlas.)

In rare cases, as the bones form and become rigid the results of the sacral fusion is that the two individuals go through life not merely back to back, but with backs so close that the skin fuse; it is found, however, that the intimate bony union in them is only in the sacral region.




Diagram to show that in the amniota (mammalia), in which the embryo is formed not on the surface, but within the ovum and within the amniotic cavity, the same principles apply as in the case of the fish embryos developed on the surface of the ovum, the embryo developing parallel to the surface; I, superior apicopolar fusion; II, superior dorsipolar fusion; H, inferior apicopolar fusion; a, due amniotic eavities.

(b) Inferior Apicopolar Fusion (Ischiopagus).—This corresponds in every respect to the Janiceps fusion at the superior pole, only here it is the lower extremities and not the face that become forced laterally, and this in such a way that one of each lateral pair of legs is contributed by each individual. There may be the same union at an angle, so that there is a pair of legs on the one side, a single, compound leg on the other. There is the same, though slighter, tendency toward ventral

fusion as in Janiceps,¹ and similarly the development of a common umbilical cord.

Unequal or Parasitic Symmetrical Double Monsters.—Just as with monochorial twins, in which the fusion is in the placenta and one is the more vigorous, so with these double monsters, in which fusion is in the body; if there be unequal vigor the smaller comes to be nourished directly from the anastomosing bloodvessels of its stronger fellow, becomes acardiac and imperfect, and appears as a parasitic outgrowth. But the adaptation becomes more perfect in this latter case; the circulation is stronger, the parasite is not ordematous, although there is,



Thoracic parasite (Gastrothoracopagus parasiticus), (Wirtensohn,)

coincidently, a preliminary stage in which the distant parts are apt to have their circulation arrested, and, as a consequence, exhibit aplasia and lack of growth. We thus encounter parasitic thoracopagi, craniopagi, ischiopagi, and pygopagi. With Schwalbe, we may regard these as a further grade of the acardiac monsters, in which the vascular anastomosis has taken place, not in the placental region, nor in the umbilical, but into the body region of the stronger embryo.

1.00

Asymmetrical Double Monsters by Inclusion. —We occasionally encounter aberrant masses of tissue, sometimes (1) projecting; at others (2) definitely included within one or other body cavity. In the first series a weaker embryo has become adherent to the outer surface of the fully formed individual, and, it may be, becoming thus adherent, has become partially included as a result of the infolding and closing in of the body fissures. In the second, there are several possibilities, the simplest of which is that this same process has resulted in

complete inclusion and development of the state of "fœtus in fœtu." Time and again in examining a series of hen's eggs incubated for twenty-four hours, eggs are met with having two primitive streaks, one of them generally small or otherwise imperfect, not arranged symmetrically and with regard to a common polarity (see Fig. 56, p. 233). What is the cause of this want of polarity we do not know; it is possible that two unequal germinal disks have made their appearance in one ovum and undergone fusion. It is cases such as this that represent the early stage of these asymmetrical monsters.

But fœtal inclusions of this second order can only develop in relationship to the main fissures of the body, in relationship to the midline,

¹ For in the mammalian embryo the caudal curvature is not so pronounced as is the cranial.


FŒTAL INCLUSIONS

24

- 10

along the line of the thoracic or abdominal cleft. It is difficult to conceive of the process as operative in connection with the neural cleft at either extremity, and that because by the time that the cephalic and caudal curvatures become developed, the intervention of the amnion (as indicated diagrammatically in Fig. 94) must hinder any such inclusion. But in the thoracic and abdominal midline terata (*i. e.*, monsters) or teratomas are met with, and however rudimentary and chaotic the collection of tissues found, provided that representations of all three germ layers be present, the simplest explanation is this of foctal inclusion.

Teratomas.—There are, however, other cases which it is not so easy to explain by this inclusion theory, in which the position of the mass containing elements derived from all three germ layers is not in relationship to the primitive fissures. These demand another explanation, and a long series of hypotheses has been adduced to explain them, from that of aberrant snaring off of cells, or cell collections, during embryonic



Diagrammatic representations of development of fortal inclusions. With the more active development of a, the larger of two embryos lying in a common germinal area, as the yolk sae becomes exhausted, the smaller embryo (b) becomes drawn into the infolding body cavity of the former.

or fætal life, which cells eventually take on independent growth, forming an independent republic, or "free city," within the empire; through theories of aberrant blastomeres—single cells from the period of earliest division of the ovum becoming displaced and eventually taking up independent growth—down to the theory of parthenogenesis pure and simple, developed ova of the individual, or, it may be, spermatozoa, taking on spontaneous growth without due stimulus of fertilization. This last for those cases, alone, in which masses of this order develop in the ovary or testis.

We shall discuss these remarkable tumors in association with tumors in general. For many reasons they might be considered at this point; by mutual relationship it will be seen that they come very close to the double monsters; they are, in fact, one constituent of a double growth. It will, however, be more helpful to a grasp of the more important subject of tumor growth to consider them in the other connection. One group, indeed, we have already discussed—the cases of polar or serial deduplication, the epignathi, etc.—and have shown that we regard them

as due to the independent proliferation of aberrant growing point cells —cells of a later stage than the blastomeric, but still *totipotential*, capable of originating tissues of all three layers.

Here, as indicating what we believe to be their relationship, we would sum up by classifying the different forms of double growth in accordance with the conclusions arrived at in this discussion.



Ovarian teratoma (ovarian dermoid), to show development of hair, c_i and teeth, d_i^*

We may, in the first place, establish four main groups:

1. Twins.-Gemini.

2. Double Monsters Proper.-Terata.

3. Teratoid Growths.—Differing from the latter in that the weaker member of the partnership becomes subservient to and parasitic upon the other, still, however, exhibiting the development of certain recognizable organs, limbs, etc. This is an intermediate and often poorly defined group, grading into the preceding and the following.

Teratomas.—'The results of the independent growth within the tissues of one individual of a cell capable of giving rise to tissues representing all three germ layers, but incapable from its surroundings of developing completed organs and parts.

1. Twins. Gemini.

- (a) Heteroophal or dichorial, from separate ova.
- (b) Monochorial, from a single ovum (by complete cleavage). (1) Equal.
 - (2) Unequal. Acardiaci, or chorioangiopagi. Hemiacardiaci; possessing imperfect heart. Holoacardiaci; without heart.

REDUPLICATION OF ORGANS

2. Symmetrical Double Monsters; Double Terata.

(a) By cleavage: Dichotomous Deduplication.

- (1) Superior polar.
- (2) Inferior polar.
- (3) Superior and inferior polar: Anakatadidymus.
- (4) Mesial: Mesodidymus.
- (b) By fusion (following primary complete cleavage, as in 1 (b): Fusional Deduplication.
 - Ventral: Thoracopagus, various grades down to xiphopagus.
 - (2) Lateroventral: Sternopagus, various grades.
 - (3) Superior apicopolar: Syncephalus or Janiceps.
 - (4) Superior dorsipolar: Craniopagus.
 - (5) Inferior apicopolar: Ischiopagus.
 - (6) Inferior dorsipolar: Pygopagus.

All these may be (1) equal, or (2) unequal.

- 3. Asymmetrical Double Monsters.-Parasitic foetus, teratoid.
- 4. Teratomas.
 - (a) Twin teratomas, due to inclusion and imperfect development of a second embryo developing from the same ovum (and thus equally to be regarded as a second division of Class 3).
 - (b) Filial teratomas, the products of an aberrant totipotential cell of the host individual. (See Section III, Chapter XV.)


Multiplex births and triple monsters follow the same classification.

REDUPLICATION OF ORGANS; MERISTIC EXCESS.

The same principles which we have found at work in the growth of the individual as a whole are in action in connection with the development of the separate organs or parts. Just as in the plant subsidiary growth centres show themselves for the lateral branches, leaves, etc., so we may regard the different limbs and organs of the vertebrate as developed from, or through, the agency of growing centres, which may show aberrations in their growth just as do the primary growing points. Having made this statement, we may proceed rapidly.

We may divide the abnormalities of excess of individual parts into increase in the number of parts arising in (1) longitudinal series, and (2) in lateral series. Of the former, the most marked examples occur in connection with the longitudinal axis, the vertebre and their connections; we may have accessory vertebre, intercalated, as it were, in one or other region. There is the formation of an increased number of vertebral centres. Associated with this we may have a corresponding increase in the number of ribs. The two, however, do not necessarily correspond; the number of ribs may be increased without increase in the number of vertebra (e. g., cervical ribs). The ribs, indeed, arise from lateral-growing points, and may show tendency to deduplication of the lateral order, bifurcation of their sternal cartilages or ventral extremities, etc.

The examples of deduplication in lateral series are very numerous; the commonest are: (1) *polydactyly*, with all its various grades, from broadening of a terminal phalanx, through double nail, down to doubling of the whole phalanx and appearance of a complete accessory digit, or even the development of an incomplete double hand; (2) *polymastia*, and the development of accessory nipples and breasts. Elongation and partial or complete doubling of the kidneys, the adrenals, testes, ova, and other paired organs are usually instances of meristic excess in ongitudinal series.



Examples of complete and incomplete polydactyly. (Ahlfeld's Atlas.)

Accessory Organs.—Allied to, but different from, the above conditions are the frequent examples of the presence of small isolated accessory organs. The spleen and adrenal afford the most frequent examples, the liver and pancreas less frequent. What we deal with in these cases is evidently a snaring off, or segregation, of certain cells during the course of development, cells already so far differentiated that they have become unipotential, *i. e.*, capable of producing only one type of tissue. These sometimes may come to lodge in distant parts. Thus, the testes and the ovary originally lie in apposition to the developing adrenal; in their migration they may carry down with them some cells from the latter organ. As Marchand has pointed out, accessory adrenal nodules may be found in close connection with the developed testes and ovary.

CHAPTER V.

MONSTROSITIES AND ABNORMALITIES-(Continued).

ABNORMALITIES OF DEFECT.

General Defect.—**Dwarfism.**—In connection with anomalies of defect, we can proceed along the same lines as those followed in connection with anomalies of excess, and begin with general defect.

As with giantism, we recognize a dwarfism due to influences conveyed through the germ cells.

Thoma lays down that if 169 cm. in the male and 163 cm. in the female be taken as the mode, the mean embraces all those having heights within 3.8 cm. on either side of this figure; in other words, one-half the adult population is included within these figures, the other half lies outside. It is suggested that this number 3.8 be multiplied by 5 and subtracted from 169 and 163, respectively, to determine the heights below which dwarfism is present, e. g., 150 cm. (59 inches) for the male, 144 cm. (56 $\frac{1}{2}$ inches) for the female. It is found that only one individual in every thousand comes below these figures.

These figures we may regard as applying to the true dwarfs, those smaller than normal, in accordance with the law of chance. That the relative amount of "bioplasm" entering into the fertilized germ cell plays some part in their development is suggested by the fact that the division of the ovum into two in the two-celled stage has been found by all observers to result in the production of dwarfed individuals. There is another category, however, of those dwarfed in consequence of intra-uterine defect, affecting particularly the limbs, or of inherited diathesis toward osteogenesis imperfecta. With them should probably be included the cretinoid dwarfs (p. 353), in whom imperfect development is associated with deficient thyroid secretion. Undoubtedly, also, extra-uterine influences during childhood have their influence upon the stature of the adult. This has been well brought out recently in England by a study of primary school children in London and other But at Glasgow it was found that there must be something cities. determining the conformity of height and size to the "law of chance, and that something would seem to correspond with graded variation in the amount or quality of what we have termed the biophoric molecules primarily contributed to the individual. By constantly selecting the larger or best grown seed of any crop and sowing this only, the general average of that crop may be greatly improved within a few years; if, after this, the seeding be left to nature, there is progressive deterioration. Ultimately, it is environment that tells, as shown by

the above observation upon the stunting effects of town life and the fact, pointed out by Cantlie, that there is no fourth successive generation of dwellers in London.

Yet another cause of dwarfism, rare in the human race, is inbreeding, as all breeders know well. Calkins' observations upon the parthenogenetic development of the parameecium, and the eventual dying out or senescence of the stock, may be recalled in this relationship.



Various grades of cyclops formation. Fig. 97, arrhineephaly, with synotia, the fused orbits still retaining separate pupils. Fig. 98, cyclops prozer, with median single orbit and pupil. Fig. 99, more extreme grade, with complete absence of nasal passages, anotia, and microstomia. (After Ahlfeld.)

Regional and Organic Defective Development.—This is parallel with regional and organic excessive growth, and, like that, may be of more than one form. Thus, there may be:

 Hypoplasia.—Congenital or inherited hypoplasia, small size of an otherwise perfectly formed organ, due to relative deficiency of matricial matter—of one kidney, of one or both ovaries or testes, of a limb, etc.

 Polar Hypogenesis.—There are certain striking conditions which may best be explained as the converse of polar deduplication and dichotomy. These affect both the superior and inferior poles of the body, as, again, the secondary growing points.

Of such at the superior pole are the various grades of cyclops formation—monsters with the two eyes in a single fused orbit, the nasal passage being deflected upward into a proboscis; with an imperfect double eye in one orbit; with a single median eye; with no eye or medial facial parts. The figures opposite illustrate the cases. The slightest grade of all is that to which Welker gave the name *trigonocephaly*, showing peculiar smallness of the front of the skull, with approximation of the orbits. The other grades have been termed *arrhincephalus*, synotia, cyclops proper. Where, as in the advanced cases, the mouth also disappears, we have *astonia*.

Recalling our conception of the growing point, and how this gives off backward in succession the cells destined to be the mother cells of successive segments, the oldest becoming thus farthest away from the

POLAR HYPOGENESIS

growing point, the most recent in its immediate neighborhood, it is obvious that at the superior pole the anlagen for the extreme anterior parts are the last to be laid down. Along these lines cyclops formation are to be regarded as due to *premature exhaustion of the growing point*, or arrested growth of the same at a period when the mother cells for the superior portion of the body and most of the head have been given off.

It is not that the lateral parts undergo secondary fusion in the middle line, having crowded out and arrested the growth of the more median superior regions; those regions have never developed, and the lateral parts have never been other than in apposition. Certain interesting experiments of Stockard' deserve mention. He found that by subjecting the fertilized eggs of fish to the temporary action of dilute alcohol, as also, to a slighter extent, to the influence of other anesthetics, and mag-




Diagram to illustrate mode of production of polar hypogenesis: A_i the normal development of the apical portion of the organism, the daughter cells given off by the growing point controlling the development of the apical parts of the body; B_i premature exhaustion of the growing point cells, those controlling segments I and II not being developed. As a consequence, segments III meet in the middle line.

nesium salts, their growth was not absolutely arrested, but various grades of cyclopia showed themselves. With alcohol, from 90 to 98 per cent, of the developing fish showed either typical cyclopia, or symmetrical or asymmetrical microphthalmos, one or both eves being small, deeply buried, or absent. He concluded that these defects are produced by lessening the developmental energy at certain critical periods.

Siren Formation, or Symelia.—At the inferior pole we encounter a corresponding series of anomalies of defect: Sympus, or fusion of the two lower extremities into one; sympus apus, with still further fusion and a single foot; and apus, the fused limbs being represented by a single conical footless stump. These cases lack the external organs of

¹ Proc. Soc. for Exp. Biol. and Med., 7: 1909; 1.

generation, and only rarely, in the slightest grade, is the anus present. The ischia and acetabula are always fused, the pelvis greatly narrowed.

We would again emphasize that the end of the coccyx does not indicate the site of the inferior growing point, but that lies at a point above and anterior thereto. So that in polar hypogenesis it is not so much the coccyx that is involved as parts originating in front of this. Further, if the homology be correct of the lateral halves of the mammalian penis with the claspers of the ray or skate, and these claspers be the inferior rays or segments of the hind limb, it will be seen that these must be the segments of that limb to be first and most surely affected by premature



exhaustion of the inferior growing point. In other words, if there be arrest of formation of the inferior limb rays, with fusion of the representatives of the more cephalad portions of the same, there must be absence of external genitalia.

Hypogenesis of Secondary Growing Points. —Along the same lines are best explained cases of syndactyly and reduction in the number of phalanges, and, as we shall point out later, conditions of the order of agnathia.

3. Imperfect and Arrested Development of Parts Other than Polar.—In the course of our discussion of inheritance and in this last chapter there have been adduced more than one factor as effective in producing imperfect and arrested development, viz.: (1) Reversionary degeneration with defective constitution of the biophores, and so of parts of more recent evolution; (2) quantitative defect in

the matricial matter set apart for the development of a particular organ; (3) premature exhaustion of the matricial cells; and (4) intrauterine disturbance, (pressure, formation of bands, adhesions, etc.). While certain of the results produced by these agencies are distinctive and recognizable as being due to one order of cause, with many it is difficult, if not impossible, to assign a particular causation; aplasia, or arrested development of a part, however, produced in early embryonic life, leads to the same results, and, as Woodruff points out, there is no valid reason why one and the same cause, acting on the ovum prior to fertilization, may not exert the same chemical or physical action as it does immediately after fertilization. It is quite possible, therefore, that conditions of a certain order that are acquired may be identical with those due to influence acting on the germ cells prior to fertilization.

Thus, to attempt to analyze and determine in every case of defective growth what is the particular causation is beyond our power. It becomes necessary from this onward to pass in review the various forms of imperfect development of particular regions and systems. The latter

DEFECTIVE CLOSURE OF THE DORSAL GROOVE

we shall undertake in the special part of this work in which we deal systematically with the different systems. The regional malformations must be noted here; we refer more particularly to defects associated with the closure of the different fissures of the body—of the medullary groove behind, the great anterior thoracico-abdominal fissure, and the facial clefts. Following this, we shall conclude this section with a brief consideration of conditions which do not fall under any of the headings thus far noted.

MALFORMATIONS ASSOCIATED WITH DEFECTIVE CLOSURE OF THE DORSAL GROOVE.

This groove, it is scarcely necessary to remind the reader, originates as a longitudinal fold of the epiblast, extending from one end of the embryo to the other, and the epiblast lining it undergoes differentiation at a very early date to form the neuroblast or mother tissue for the whole nervous system, the folding in leading to the formation of the neural eanal. In the region of the head this neural canal distends into three bilateral pouches, the neuroblast lining the pouches giving rise to the fore-, mid-, and hind-brains, respectively.

There may be total failure of closure of the groove, or merely local failure, the necessary result in each case being that, instead of there being a neural canal (with expansions to form the ventricles of the brain), there persists exposed nerve matter, which at its edge passes and becomes transformed into the ordinary epiderm. We thus obtain the various grades of *Cranioschisis* and *Rachischisis*.

Anencephaly, Acrania, or Hemicephaly.—Anencephaly, acrania, or hemicephaly is a relatively frequent monstrosity. The frog-like appearance, due to absence of development of the frontals, is very characteristic. Owing to lack of closure of the neural groove in the cephalic region, there is lack of development of the vault of the skull and of the hairy scalp. In extreme examples even the orbital plates of the frontals are undeveloped. The freely exposed brain substance becomes extremely congested, so that little is to be made out beyond an amorphous mass of vascular membranes. Nevertheless, the basal portion of the brain has given origin to the optic and auditory vesicles and to the cranial nerves.

As to the direct cause of this condition opinion is still at variance. The view of Darcste still has its adherents, based upon his observation upon the anencephalic chick, that defective development of the amnion is at fault, leading to pressure upon the head at an early period, and, as a consequence, arrested development. Certainly, where there are amniotic adhesions of the foetal head there are accompanying grave developments of skull and brain; and the lordosis, or curvature of the cervical vertebrae, is difficult to explain on other grounds. But such adhesions are rare, nor is the general development of trunk and limbs of these monsters arrested to an extent corresponding to what

we should expect in defective development of the amnion and escape of anniotic fluid. Another view is that there has been early fortal hydrocephalus, with distension and rupture of the ventricles and arrested development of the cranial vault. There are, however, no transitional cases to support this hypothesis.

Neither view is adequate to explain the majority of cases. On the other hand, while the body as a whole is found, in general, well developed, there are certain commonly associated defects, viz., arrested development of the adrenals, while, as Shepherd and others have pointed out, the paired musculus sternalis is often markedly developed. These facts



Fig. 102

Anencephaly.

FIG, 103



Iniencephaly, (McGill College Museum,)


point to some general vice of development, and so commonly do we gain a history of parental infection in these cases, that we are inclined to give parental and germinal intoxication as the most frequent underlying cause. There is frequently an associated absence of closure of part or the whole extent of the vertebral canal and condition of *spina bifida*.

Exencephaly.—In these cases the cranial vault is in part developed, more often the frontal region. Where this is the case, the frontal bones are flattened and receding, owing to the escape of brain substances

SPINA BIFIDA

behind. The imperfectly formed brain substance, with its membranes, protrudes behind as a sac overhanging the back.

In most of these cases, however, as in meningocele, we deal with deficient formation of the bony vault of the skull rather than lack of closure of the neural canal. These conditions, along with that of hydrocephalus, will be discussed along with the regional malformations of the nervous system. The most extreme condition of this nature is *Iniencephaly*, in which, with spina bifida, there is imperfection of the occipital bone, through which part of the brain projects. What is most striking is that there is extreme flexion of the vertebral column, so that the occipit is approximated to the sacrum, the skin passing directly from one to the other.



A, schema of development of medullary groove; B, formation of neural canal by closure of the medullary groove; C, complete mehisebisis; the medullary groove remains open; Epi., epiderm; a, neural tract; p.a.s, pia-arachnoid space.

Rachischisis, or Spina Bifida.—Strictly speaking, every case in which the spine remains "bifid" in consequence of failure of the lamine of one or more vertebre to unite, is a case of spina bifida, or rachischisis. What is all important, from a diagnostic and surgical point of view, is the extent to which, and the mode in which, the spinal cord is involved in the defect. We thus distinguish the following series of cases:

 Spina bifida completa, in which the primary cause of lack of closure is failure of the medullary groove to close in and form the neural canal. As a consequence the ependyma, or superficial layer of nerve substance, remains in continuity with the skin on either side, the cord forming a flattened superficial plate, as indicated in the diagram (Fig. 104, C). Of this order we may distinguish the following grades:

(a) **Spina bifida completa totalis**, in association with anencephaly. The neutral matter forms a broad plate extending down the back to the coccyx, and fusing on either side with the skin of the back.

(b) **Spina bifida completa partialis**, also in association with an encephaly, but affecting the cervical region only, the neural canal becoming formed below this.

(c) Spina bifida completa restricta, local, affecting a restricted area in either cervical or, more often, lumbar region. Here, whether through abnormal curvature of the embryo, or lack of growth energy, one or other of the last regions of the medullary groove to close remains openas a flattened, exposed plate of nerve tissue, passing almost imperceptibly into the skin on either side. The cord in these cases, instead of becoming, with progressive growth, relatively short, compared with the vertebral column, remains long, and inevitably its canal opens on to the surface at the upper and lower extremities of the area. Very rarely this has been recorded in the dorsal region. Two results may ensue: (a) Either the area remains flattened, there being free discharge of fluid from the cerebrospinal canal, keeping the surface moist. Such cases, if born alive, inevitably exhibit infection, and, whether from this, or from the free loss of the fluid, die within a few days. Or (b) owing to the absence of pressure, fluid accumulates in the anterior pia-arachnoid space under the defect, causing the flattened cord to project backward, in which process the openings of the neural canal are apt to become occluded by pressure of the cord against the edge of the orifice. In this way originates the true *myelocele*.

In all these cases the neural epithelium covering the defect is apt to undergo extensive degeneration; the cord is represented by a flattened mass of greatly congested vessels, between which isolated nerve cell may be distinguished; the spinal nerves are given off from the anterior aspect and traverse the cyst (if present).

2. Spina Bifida Incompleta.—In all the remaining cases there has been due closure of the medullary canal, with junction of the cutaneous epithelium in the median dorsal line; the cord has become surrounded by its meninges, but there has been failure of the laminæ and associated tissues to develop adequately.

(a) The cord remains in almost immediate contact with the skin, and, with accumulation of fluid in the anterior arachnoid space, becomes represented by a flattened ribbon of nerve matter, from which pass the spinal nerves traversing the cyst—meningomyelocele of the first order. This form is relatively common.

(b) Lack of pressure leads to accumulation of fluid in the central canal of the cord in the area of defect—myelocystocele, or syringomyelocele (hydrorrachis interna). Here the spinal nerves lie outside the cyst wall.

(c) Through some defect in the walls of the vertebral canal there projects a cyst formed of the meninges—*meningocele*. Such defect may be either between the lamine, the cyst projecting backward, or, rarely,

DEFECTS OF ANTERIOR THORACICO-ABDOMINAL FISSURE 267

of the nature of a cleft between the bodies of the vertebra. There are no nerves in association with the cyst.

(d) Fluid accumulates in the *posterior* pia-arachnoid space, and at the same time a portion of the spinal cord also protrudes through the defect in the bony tube—*myelomeningocele* of the second order (very rare).

(e) Localized lack of junction of the laminæ, but no fluid projection, a pad of fatty and muscle tissue filling the space between the skin and the cord—*spina bifida occulta*. This form shows itself in either the cervical or lower lumbar region, and characteristically the area of defect is covered by a clump of long hairs.

DEFECTS OF THE ANTERIOR THORACICO-ABDOMINAL FISSURE.

We need but remind the reader that the embryo, at first a flattened elongated plate on the surface of the ovum, becomes gradually more cylindrical, the sides curving in to form the body cavity. For a considerable period these sides do not meet, and the developing viscera are exposed ventrally; for some time, indeed, a portion of the bowels actually protrudes, as does the allantois, with its vessels. Eventually, this great thoracico-abdominal fissure may fail to unite, and thus we encounter the following conditions:

Fissura Sterni.—Affecting the whole or part of the sternum, above or below. Either the sternal elements proper may fail to develop or the whole may be wanting, with resulting exposure and *ectopia* of the thoracic viscera. Where this is complete, the lungs cannot expand and extra-uterine life cannot be. Where incomplete, the heart alone may be exposed—*ectopia cordis*—at times with, at times without, development of the pericardial sac.

Complete Abdominal Fissure.—Eventration.—This is occasionally encountered, the viscera in general protruding through the median aperture.


Hernia Funiculi Abdominis.—Hernia funiculi abdominis is more common, or incomplete closure of the walls at the point of entry of the umbilical vessels. In this case a portion of the viscera lies within and distends the proximal portion of the umbilical cord.

Very rarely, the *omphalomesenteric duct*, the communication between the primitive small intestine and the yolk sac, may persist and remain patent, extending into the cord.

Fissura Vesicogenitalis.—Or the persistence of the fissure may be limited to the lower end of the abdomen. It will be remembered that the urachus represents the old communication between the bladder and the umblical cord, that originally from the region of the cloacal membrane and end of the gut the allantois was developed, passing to the region of the navel, and that in the dorsal end of this, at first freely communicating with the cloaca, the bladder develops, its ventral end being continued as the urachus. The bladder thus is, from the first,

MONSTROSITIES AND ABNORMALITIES

closely connected with the cloacal membrane. This latter eventually becomes perforated, the hind gut beyond it atrophies, and the rectum thus opens into the anus. Reichel, Enderlen, and the more recent workers on this subject hold that the various grades of *ectopia vesicw* are to be ascribed to a very early and abnormal division or fissure of the cloacal membrane extending navelward in the region of the bladder and abdominal wall, whereby the two sides of the abdominal wall do



Schema of development of rectum and urinary passages from cloaca,

not come together, and the bladder, also being associated in the fissure, is left open in front.¹ Associated with this condition we encounter that of *epispadias*, or patency of the urethral canal, on the upper aspect of the penis. The public arch, along with the abdominal wall, fails to gain complete formation; the two sides do not meet, and there is no symphysis; the anterior fissure of the bladder is continued along the urethra. The penis in these cases is short and imperfectly developed.

Another frequently associated condition is the persistence of the cloaca,

¹ For a full study of this complicated subject and criticism of the literature, reference may be made to Enderlen, *Ueber Blasenectopia*, Wiesbaden, Bergmann, 1904.


DEFECTS OF SPECIAL REGIONS

so that the gut, instead of opening into the anus, is connected with, and discharges into, the bladder, or opens into both.

In the most extreme cases, according to Marchand, the division through the genital eminence may be permanent; so that there is half a penis or half a clitoris on either side.

DEFECTS OF SPECIAL REGIONS.

Defective Development of the Diaphragm.—At this place, while discussing imperfect regional development, it is well to take into consideration the condition of defective closure between abdomen and thorax. This, the so-called *congenital diaphragmatic hernia*, is not



Congenital diaphragmatic hernia in man, aged fifty-seven years, partly false, there being at A and B direct communications between the peritoneal and pleural cavities, from the edges of which fatty omental folds of tissue passed into the thorax; partly true, the serous membrane forming a covering over the stomach and left kidney as they lay in the pleural cavity. The muscle of the diaphragm is shaded dark.

uncommon, and certain grades are compatible with continued existence. We have, indeed, encountered it in a man, aged fifty-seven years, who died from other causes.¹ As a result, the pleural and peritoneal cavities are in direct communication through an orifice frequently wide enough

¹ Fry, Montreal Medical Journal, 25: 1897.

MONSTROSITIES AND ABNORMALITIES

to allow a large part of the stomach, the spleen, the left lobe of the liver, and several coils of the intestine to lie in the pleural cavity, though sometimes the defect is trivial. It is much more common on the left than the right side, the presence of the liver appearing to favor the orderly development of the right half. When occurring on the right side, part of that organ passes into the pleural cavity.

While the condition is spoken of as diaphragmatic hernia, it must be remembered that, strictly speaking, a hernia is a protrusion of some of the abdominal contents through some abnormal opening, carrying the peritoneum before them, and as here most often the orifice is complete and there is no such covering of parietal peritoneum, most cases are properly those of false diaphragmatic hernia. At times, however, there is a membrane, peritoneal on the one side, pleural on the other, covering the defective area of the ligamentous or muscular portion of the diaphragm; then the viscera protruding into the pleura form a true hernia.

The commonest site of defect is in the ligamentous portion; smaller defects may occur behind in connection with the œsophageal foramen, or in front, from lack of development of the portio sternalis, when there may be *ectopia cordis abdominalis*. When the defect is large, the lung on that side is unable to develop fully, and, indeed, may remain in an atelectatic condition; in an infant dying at birth we have seen it scarcely recognizable. In the commoner, left-sided condition, the heart usually assumes a median position.

A more severe grade of this same condition of arrest of development is complete, or almost complete, absence of the diaphragm. This is incompatible with continued existence.

Defective Closure of the Facial Clefts.—Here, apart from the phenomena of polar reduction already noted, the abnormalities of defect show themselves specially in connection with the bilateral facial cleft extending from the orbit to the mouth. According to causation, Marchand divides them into primary and secondary—primary, due to inhibited local growth and fusion of the parts; secondary, due to amniotic adhesions and other causes of arrested junction; the former regular, or reproducing with fidelity an earlier developmental stage, whether unilateral or bilateral, the latter irregular, with more or less distortion of such earlier state. The inherent nature of the primary disturbances is indicated by the frequent inheritance of *harclip* in various grades.

In its very slightest grade, *harclip* affects only the upper lip, and that on one side; in severer grades there is associated lack of union between the maxillary process and the intermaxillary bone, so that there is an alveolar cleft; this may extend into and affect the hard and soft palate *cleft palate*—in which case the nasal cavity communicates with the mouth. In the severest cases of all we encounter either a cleft passing along the side of the nose into the orbit, or, again, though this is very rare, lack of formation of the intermaxillary bone, in which case there is a condition of median cleft of the lip.

To recite rapidly the various orders of defect in this region, they arefollowing Marchand's classification—as under:

Median cleft of the nose, with one nasal passage or with a simple tube on either side. Allied to this is the rare condition in which one nostril is fully formed, the other a mere conical snout, or proboscis.

Lateral nasal cleft, through lack of closure of the lateral frontal with the median nasal process.

Cheiloschisis. Simple harelip. See above.

Cheilognathoschisis. Cleft of lip and jaw, usually bilateral, with the intermaxillary forming an isolated median process (rare).

Cheilognathouranoschisis. Cleft of lip and jaw and palate.

(a) Median cleft of lip, with lack of development of intermaxillary bone and broad cleft of lip (seen in arrhinencephaly).

(b) Unilateral cleft of lip and jaw, due to defective union of the nasal and maxillary processes. Often associated with lateral cleft of the palate, lack of union of the alveolar process with that of the other side and the vomer. Most frequently left-sided.

(c) Bilateral cleft, with median or bilateral palatal cleft. Here the intermaxillary forms a snout-like projection connected with the nasal septum and the vomer.

(d) Unilateral or bilateral hardlip, with closure of the mouth through fusion of the upper and lower lips. In the first case only one-half of the mouth cleft is left, forming a common opening with the nose; in the latter the cleft and communication is bilateral.

(e) Primary lateral facial cleft. Cheilognathoprosoposchisis. Persistence of primary condition, or separation between the maxillary process of one side and the lateral and median frontal or nasal processes.

Makrostomia. Fissura buccalis. Lack of union of the sides of the mouth cleft, either unilateral or bilateral, the mouth thus reaching to the ear.

A prosopia. Complete lack of formation of the various processes forming the face, which thus is represented by an irregular cavity.

According to Marchand (who discusses the subject fully), the variation in the number of the teeth which may be present on the internaxillary is not an indication that the internaxillary on either side is formed of two halves, and that the lack of union is now on the one side, now on the other of the outermost of these two halves, but is to be ascribed to the fact that the anlagen for the teeth are not directly connected with the anlagen for the internaxillary and upper jaw, but are developed at a later period and liable to deduplication when, by a cleft or other defect, the row is interrupted.

Uranoschisis. Cleft palate. Affects the hinder portion of the hard palate, along with the soft palate at one side. Very rarely does it affect the anterior part of the hard palate only.


Staphyloschisis. Fissure limited to the soft palate, and in the slightest grade to the uvula only (bifid or double uvula).

A series of these cases of harelip and cleft palate is markedly hereditary, but not all. In other cases, as in micrognathia, the arrest is of

MONSTROSITIES AND ABNORMALITIES

mechanical origin. In the majority no clear primary cause is to be made out. As already stated, the irregular forms are obviously secondary to such mechanical disturbance as is produced by amniotic bands.

Other malformations of the face which are regional, affecting more than one organ, are best discussed here.



Development of the face of the human embryo (His): A, embryo of about twenty-nine days. The nasofrontal plate differentiating into processus globulares, toward which the maxillary processes of first visceral arch are extending; B, embryo of about thirty-four days; the globular, lateral frontal, and maxillary processes are in apposition; the primifive opening is now better defined; C, embryo of about the eighth week; immediate boundaries of mouth are more definite and the nasal orifices are partly formed, external ear appearing, D, embryo at end of second month. (Heisler.)

Malformations of the Lower Jaw.—Agnathia.—Absence or great aplasia of the lower jaw bilaterally. When this is the case the outer ears tend to approximate, and may, indeed, fuse in the median line of the neck below the upper jaw (synotia). There is accompanying microstomia. Winckel' ascribes the conditions as due to lack of growth owing to am-

¹ Münchener med. Wochenschr., 1896; No. 18.

ABNORMALITIES OF THE BRANCHIAL CLEFTS

niotic pressure on the back of the head, the lower jaw area being thus compressed against the neck. This, however, does not satisfactorily explain the synotia, and, with Dareste and L. Blanc,⁴ we would ascribe the condition to arrest of development of the third eranial vesicle (or, we would say, of the lateral growing point forming this region), with abnormal fusion of parts before and behind. The region of the third vesicle (fourth ventricle) remains a simple tube; the corresponding parts of the face are not developed. There is lack of development of the first branchial arch, in consequence of which the lower jaw and the basal portions of the superior maxilla are wanting. Certain bony portions of the internal and middle ear are still found, as also the external ears, but they come close together, and fuse in the midline.

This *otocephaly* is often accompanied by disturbed development of the region of the first vesicle, producing what Blanc terms *cyclotia*. Whereas in cyclopia the eyes remain in the upper part of the face, here, owing to the lack of development of the superior maxillaries, the eyes pass downward and join, and the new orbit is formed under the anterior portion of the sphenoid.

Microquathia, or imperfect development, with small size of lower jaw and conditions of chinlessness, may possibly be due to the compression suggested by Winckel.

Cleft tongue, median cleft of lower jaw, fistula and cyst of the lower lip are all rare conditions.

(Here, while discussing the malformations of the lower jaw, attention may be called to the opposite conditions of excessive growth, viz.: (1) Deduplication of the alveolar process with development more or less complete of a double row of teeth; and (2) dignathia, deduplication of the lower jaw, the converse of agnathia.)

Anomalies of the Branchial Clefts.-Incomplete closure of the second and lower branchial clefts leads to the formation of *congenital* fistulæ and certain congenital custs of the neck. These clefts pass from the exterior to what becomes ultimately the pharynx. The fistula may be complete, with a free passage of communication from without inward; may be incomplete internally or externally, resulting in a blind fistula; or, finally, the passage may be obliterated at either end, but persistent in its central part, in which case a cyst develops. According as this cyst originates from the more external or the more internal portion of a cleft, so may it be lined with squamous or with ciliated epithelium. Bland Sutton² more particularly has studied these conditions; he points out that the outer openings of the fistulæ, according as they are developed from one or other cleft, occur along a slightly curved line, with its concavity forward, extending from the external auditory meatus downward toward the sternoclavicular articulation on either side. Counting the Eustachian tubes which represent the first, there are potentially five

¹ Jour, de l'Anat, et Physiol., 1895. See also Le Gendre, Bouchard's Pathologie, 1: 260.


² Tumors, Innocent and Malignant, 1st edit., 1894; 323.

18

MONSTROSITIES AND ABNORMALITIES

pairs. It is the second pair which most commonly persists wholly or partially as an abnormality.

Cloacal Defects.—It will be remembered that, as at the front end of the body, so at the hinder, there occurs a most complicated series of developments, which, by arrest or imperfection, favor the production of numerous anomalies. To these we have already referred briefly in



Relationship of the sexual duets and their rudiments in the two sexes: I, the indifferent primary type; II, the differentiation in the female; III, the differentiation in the male; Ped, Hyd, stalked hydratist, the termination of the Wolffan duet; Hyd, Mor_{d} , hydratid of Morgania or terminal hydratid of Fallopian tube. The term organ of Rosennüller, which should be used to comprise the terminations of the Wolffan duet and the Wolffan body, is used so variously that it is better not to employ it, but rather to speak of Epoophoron (including the Wolffan tubules) and Paraoonboron.

discussing ectopia vesice. To understand the other anomalies, it will be well to review rapidly the main transformations. The gut, in its earliest stage, ends blindly in the coccygeal region. It comes close to the surface at the cloacal membrane and is continued a little beyond this as the "post-anal gut," regarding which all that is necessary to say is that it undergoes atrophy at a comparatively early date, though not without at times leaving rudiments which may be the seat of subsequent change. For the present we may neglect this and regard the

CLOACAL DEFECTS

original gut as ending beneath the cloacal membrane. From the front of this cloacal region there passes forward to the surface of the embryo the allantois, along what will eventually become the umbilical cord. Thus, at first, into the common terminal cloaca there pass two channels. the intestinal behind, the allantois in front. Next, by the formation of lateral folds, which meet in the middle line, the division between these two channels is carried down to the cloacal membrane, so that now two separate *cul-de-sacs* exist, the rectum behind, the allantois in front. In the normal course of events nothing occurs in relationship with the posterior channel save the absorption of its share of the cloacal membrane to form the anus. In connection with the anterior passage there are several changes. The proximal portion of the allantoic channel becomes converted into the medianly situated bladder; the passage itself becomes what is termed the urogenital sinus. Into this now open the Wolffian ducts, passing up to the kidney region. The anterior portion becomes closed off eventually to form the urethra (in the male its proximal part only, the distal part being contributed by the genital eminence, from which is developed the penis in the male, the clitoris in the female). Into the more posterior portion of the sinus opens the Müllerian duct. which undergo fusion at the lower end to form the vagina (in part, the anterior portion of this being contributed by the urogenital sinus) and the uterus, and remain distinct above as the Fallopian tubes. Thus it comes to pass that the Wolffian ducts, passing along the sides of the cervical end of the uterus, open into the vagina. But at the same time these Wolffian ducts split off the ureters, which gain entry into the base of the bladder, while the main duct on either side atrophies in the female, and, at most, at birth is represented by the rudimentary Gärtner's ducts. In the male they become converted into the vesiculæ seminales and the spermatic cords. The Müllerian ducts, on the other hand, atrophy in the male, and are, at the most, represented in the prostatic portion of the urethra, and, at the other extremity, the sessile hydatid (Fig. 108).

Here we shall deal with the rectal conditions only, taking up the genito-urinary defects later.

Atresia Ani.—Of this there are various grades: (1) A simple membranous septum closing off the rectum from the exterior, as the last remains of the cloacal membrane; (2) a thicker layer of tissue in the anal region, so that some little; or it may be considerable, distance intervenes between the anal site and the blind end of the rectum (Fig. 105, a); or (3) rarely the end of the rectum lies free in the pelvis. In the last case there has been primary lack of development of the lower end of the bowel. In cases other than the simplest, where there has been a cloaca formed, the connection of the intestinal tract with the urogenital sinus is apt to persist in the shape of fistulous or wider communication between the rectum and the urogenital organs (Fig. 109, B). Thus, there may be:

Atresia ani vulvovaginalis, the rectum communicating with the vulva or the vagina, and meconium being discharged through these passages.


Atresia ani uterina, communicating with the uterus; very rare.

MONSTROSITIES AND ABNORMALITIES

Atresia ani urethralis, with the urethra (pars membranacea).

Atresia ani vesicalis, with the bladder; rare.

Persistent cloaca. Here we have a still more complete arrest of development at an early stage; there is complete closure both of rectum and genito-urinary passages from without, and all open into a common



The various defects due to imperfect development in the cloacal region: A, imperformt anus atresia ani, the rectum closed off from the genito-urinary passage, which is patent; B, imperforate anus, the rectum opening into the bladder or urethrar; C, persistent cloacal membrane, imperforate anus and absence of urethra with persistent cloaca; D, the same, but with separation of rectum from the genito-urinary passage; E, pensistent cloaca, through lack of continuation downward of ridge separating the rectum from the genito-urinary passage,

cavity (as in C). More rarely, a later stage is indicated, the various channels being formed, but all being closed off from the exterior (D), or lastly, the cloaceal condition persists, but the membrane becomes absorbed (as in E).⁴

¹ For a fuller study of cloacal defects, see Keith, Brit. Med. Jour., 1908: ii; and of genito-urinary defects, with more particular reference to the ureters, Huntingdon's very thorough study may well be consulted. Harvey Soc. Lectures, 1906-07: 222.

TRANSPOSITION OF VISCERA

TRANSPOSITION OF VISCERA: SITUS INVERSUS.

There are yet other orders of anomalies, which only come under the heading of anomalies of defect, in so far as they do not represent the normal constitution. Such are transposition of viscera and hermaphroditism. Transposition of viscera naturally only shows itself in connection with viscera that are not paired or do not occupy the median line the heart and aorta, the stomach and intestines (organs which, originating in the median line, with development become diverted in one or other direction), the spleen and the liver. To this statement there is one slight exception, viz., the lungs, which, while paired, exhibit different lobation on the two sides.

There is also one known functional exception, viz., the speech centres in the island of Reil. Normally, it would seem that the left set of centres is functional, the right latent; this may be reversed.

The transposition may, on the one hand, affect only a single organ or group of organs; or, on the other hand, there may be complete situs inversus. Thus, the heart alone may be transposed, or the transposition may affect only the main arteries, the aorta passing from the right, the pulmonary artery from the left ventricle; or the thoracic organs may be normal, while the liver, spleen, and viscera exhibit transposition. Evidently, these partial cases can only be ascribed to local aberrations in development. With regard to complete situs inversus, it has been put forward that the individual presenting the condition has been one of a monochorial twin pregnancy; that, derived from the longitudinal division of a single ovum, he becomes a complete reflection, as it were, of his brother twin; where no history of twin birth can be obtained, it is suggested that the other brother become a foctus acardiacus, or papyraceus. Undoubtedly, there are facts telling in favor of this view; for example, in not a few cases of deduplication by cleavage and superior dichotomy the organs of one-half of the upper portion of the monster are transposed, as compared with those of the other. But, on the other hand, the general rule is that monochorial twins present no sign of such reflection. In the majority of cases no indication is afforded of the existence of situs inversus in one of the two, while, conversely, Kuchenmeister, studying 152 cases of transposition, found the history of twin birth in but a single case.¹ A more likely suggestion is that the main current of blood to or from the germinal area becomes diverted at an early stage of existence, and thus purely mechanical influences lead the vessels of one side of the organism to receive more blood, and therefore to grow more vigorously than those of the other. But it has to be confessed that we are still without any confidence regarding these hypotheses.

¹ Die angeb. Verlagerung d. Eingeweide d. Menschen. Leipzig, 1883: 268.

MONSTROSITIES AND ABNORMALITIES

HERMAPHRODITISM.

Sexual Differentiation .- The existence in the normal male and female of useless rudiments of parts characteristic of the opposite sex must not be taken as an indication that man is descended from an originally hermaphrodite ancestry. Of such in the line of descent there is no trace. Rather, such rudiments are, in Mendelian terminology, recessive features, due to the origin of the fertilized ovum from both male and female germ plasm. Elsewhere I have discussed somewhat fully the anatomical basis, as it may be termed, of fertilization (p. 144), as again of the determination of sex (p. 153). Whether as yet we are prepared to accept the doctrine of the accessory chromosome or not, the fact remains that as the result of conjugation there exist in every individual the anlagen for all the primary and secondary sexual characters of both sexes; whence it would seem that in every individual, hermaphroditism, or at least a blended, as distinct from a particulate, sexual inheritance should be potential or possible. As a matter of fact, such true hermaphroditism, or the presence in the one individual of both ovum- and sperm-producing organs, is one of the rarest conditions affecting the human being, and false hermaphroditism, or the assumption of the external organs of generation of the characters of the other sex, is far from common. This very rarity favors the view of Bateson, Castle, and others, that the properties in the germ cells determining sex are allelomorphic, or form an opposing pair or pairs, such that when the one is present the other must be inactive and recessive, equivalent to the recessive constituents of the first generation of a hybrid strain.

True Hermaphroditism.—The existence in one individual of both ovary and testis is among the very rarest of anomalies, and when it does show itself, one or both of the organs are sexually immature. With Klebs, we can recognize the following forms:

1. Lateral hermaphroditism, an ovary being developed on the one side, a testis on the other. This, in man, is the commonest form.

2. Unilateral hermaphroditism, there being on the one side both ovary and testis, on the other either ovary or testis or absence of both.

3. Bilateral hermaphroditism, there being on both sides both ovary and testis.

The terminology is perhaps confusing, but on consideration is found adequate to express the conditions.

In all these cases the general configuration of the body is of an intermediate type, now tending more to the male, now toward the other sexual type. In general, the external genitalia are of the intermediate type, *i. e.*, hypospadias is present, with small penis, separation of the two scrotal halves (or labia majora, for in general the testicle or testicles are undescended), small external orifice corresponding to the vagina, or vagina not recognizable externally, but opening into the urethra. Intermally, there is usually a uterus duplex, with tubes and ligaments.

The existence, however rare, of these cases of bilateral hermaphro-

HERMAPHRODITISM

ditism would seem opposed to the hypothesis of McClung and Wilson that sex is dependent on the accessory chromosome alone (p. 156); rather, it would seem to demand the existence of both male and female "determinants." To explain the cases of unilateral and lateral hermaphroditism it may be suggested that with the first cleavage of the ovum into the two blastomeres, which each gives rise to one lateral half of the body, these determinants become unequally distributed, even to the extent of both orders of determinants being absent from the one blastomere.

False Hermaphroditism.—As is well demonstrated in eunuchs and those castrated before puberty, the development of the secondary sexual characters, including that of the external genitalia, is largely governed by the development of the essential sexual organs, the ovaries or testes. Thus, it is where there is a congenital imperfection of the latter that we are particularly apt to encounter conditions in which imperfect formation of the external genitalia leads the individual to assume the external configuration of the other sex, or, more accurately, an intermediate type. We in this way distinguish:

1. Pseudohermaphroditismus masculinus. The individual being a male, *i. e.*, having testes, but the external genitalia and bodily habit approximating toward the feminine.

2. Pseudohermaphroditismus femininus. The individual being a female, with more masculine characteristics.

The first of these conditions is much more common. There is an insignificant, distorted penis, recalling the clitoris, perineal hypospadias (very often), the two scrotal halves without testes, which either are pelvic or in the upper part of the inguinal canal, and are immature. The outer margin of the urethral orifice may simulate labia minora, and its passage resembles a small vagina. Internally, the sacculus prostaticus (uterus masculinus) may be large, projecting behind the prostate proper as a bicornuate uterus of considerable proportions, provided with tubes; and where the testicles are undescended, they may lie in a broad ligament. Vesiculæ seminales and vasa deferentia and a small prostatic body are, however, present, and section of the testicles reveals the nature of the case. In such cases the growth of hair is more that of the female, with little or no development of beard or moustache, while the breasts may enlarge (gynacomastia) and approach the female type. In feminine pseudohermaphroditism, on the contrary, the clitoris tends to assume penile dimensions and the urogenital sinus (urethra and vagina) to be continued as far as the glans, whereby the labia majora become approximated and simulate the scrotum, the simulation being still greater when, as sometimes happens, one or both ovaries pass down the canal of Nuck. The urethra where it joins the vagina may be surrounded by a small prostate, the uterus is small, the tubes imperfect, the ovaries also small and imperfectly developed.

CHAPTER VI.

POSTNATAL ACQUIREMENT OF DISEASE.

FOLLOWING upon what has already been said, namely, that exciting causes of disease after birth must, of necessity, be of external origin, it is evident that these causes of acquired disease are either of the nature of alterations in the environment which tell directly upon one or other tissue, or are due to the entrance into the system from without of substances, either living or dead, which have a deleterious action upon the functions of the tissues. Thus, briefly, we may classify the agents producing disease acquired after birth into:

1. Mechanical-inducing "trauma."

2. **Physical**—under which can be included:

(a) Alterations in the pressure of the atmosphere, including both diminution and increase.

(b) Alterations in temperature, local and general, including both heightened and lowered temperature and freezing.

(c) Effects of electricity, both atmospheric and induced.

(d) Effects of light and of absence of same.

(e) Effects of soil and climate.

(f) Sociological effects, habitation, clothing, dwelling, occupation, and other environmental conditions.

3. Chemical Causes—under which, besides (a) the gross effects of caustic and other agents upon the tissues, we should include (b) the main effects of vitiation of the atmosphere by various gases, and (c) the main deleterious effects of improper food, as again, to some extent, the deleterious effect of certain occupations.

4. **Parasitic**—under which heading are to be included the deleterious effects of:

(a) Minute vegetable parasites—bacteria and fungi.

(b) Minute animal parasites—sporozoa, amœbæ, etc.

(c) The larger animal parasites, including worms (cestodes, trematodes, nematodes) and arthropods (arachnids and insects).

In the consideration of these as causes of disease, we have ever to keep in mind that vital activity co-exists with and depends upon physical and chemical changes in the living matter, and that "stimulation," with its resultant manifestations of increased vital activity in one or other direction, is to be recognized as primarily the action of physical or chemical agents, either category of which essentially induces alterations in the molecular condition and relationship of cell protoplasm. So long as these molecular arrangements are within certain limits, for

MECHANICAL CAUSES OF DISEASE

so long are they not merely not harmful, but actually beneficial to the organism. When these limits are overstepped in one or other direction, then it is that the molecular arrangements (or arrest of molecular arrangements) are harmful, that cellular and organic disturbances are set up, and conditions of local or general disease developed. In other words, it is purely a matter of degree whether a given agent, physical or chemical, which is capable of exerting an influence upon protoplasmic matter, acts as a physiological or a pathological agent.

It thus follows that everything capable of acting upon living matter comes under this heading of physical (and chemical) causes of disease. And as the causative agents of disease thus are so abundant, the most that we can do is to classify the causes and then to indicate those which are most frequently in action, and those which, in their action, induce certain special trains of phenomena. Here we shall say but little about most of these causes, and, more especially, we shall but glance at most of those which are spoken of as physical causes, because these are fully debated in the ordinary text-books of hygiene, in which a discussion of the effects of alterations in the composition of the atmosphere, effects of food, clothing, soil, habitation, climate, and offensive trades form so important a section.

Acting on the same principle, we shall not here describe in order the various microbic parasites; for nowadays, in every medical course, the subject of bacteriology, or, more accurately, of microbiology, has been elevated to a special subject. The case is somewhat different in connection with the animal parasites; these still are usually described, not in any special course, but in connection with the subject of pathology. But here, again, my treatment will be very brief, and that because in all schools of good standing, parasitology is being treated as a special subject, and text-books exist dealing wholly therewith, to which, rather than to a work on general pathology, the inquirer will naturally turn for detailed information.

While not enumerating and discussing the effects of individual mechanical and chemical noxe, as also the individual bacteria, and the part they play in the causation of disease, we shall discuss broadly the part played by these as a class in the production of disease. Thus, in the following pages we shall take up in order:

1. Traumatism and mechanical causes of disease.

2. The physical (and chemical) causation of disease, treated broadly.

3. Bacteria as causes of disease.

4. The animal parasites and their part in the production of disease.

MECHANICAL CAUSES OF DISEASE.

Although in most mechanical injuries we do not deal with a simple and single mode of action, we can usefully divide mechanical causes of disease and bodily disturbance into the following: 1. Concussion.

2. Puncture, with which may be included the effects of projectiles under high velocity.

3. Section.

4. Contusion, with which may be included lacerations and tearing.

5. Compression.

6. Distension.

7. Atmospheric pressure.

A little thought will show that in every one of the above we are dealing with pressure acting in various ways upon the tissues—how various will be seen if we briefly discuss the various forms. We may divide the effects even if broadly into those of *sharp* or incisive, and *blunt* or coarse traumatism.

1. Concussion (Commotio).—Here we are dealing with a brusque and momentary application of pressure to a soft, fluid or semifluid body. We see thus the effect of concussion best marked in parts in which a softer or more fluid constituent is in intimate association with tissues of greater density—in the brain and in hollow viscera having gaseous or fluid contents (the lungs, the urinary and gall-bladder, and the stomach) —as also in the bones with their marrow in contact with the unyielding osseous framework.

A sudden blow upon the skull, not fracturing the bone, and thus causing no direct laceration of the brain substance, is found to produce hemorrhages and extravasations of blood over the surface of the hemispheres and in the main tissues bordering upon the ventricles. Remembering that the brain is a soft viscus, floating, as it were, in a bath of surrounding fluid, the superficial hemorrhages may, in part, be explained by contrecoup. Following upon any such blow, the more solid contents of a cavity will partake of the motion imparted to that wall to a greater extent than will the more fluid contents. In this way the more solid brain may be driven violently against the brain case and its delicate vessels ruptured, more particularly on the side of the organ opposite to the region receiving the blow. Under similar conditions the semisolid tissues, bordering upon a cavity possessing fluid or gaseous contents, are apt to undergo rupture and exhibit hemorrhages on the side of the cavity nearest to the blow, and this because motion is imparted to these tissues by the blow, and in the absence of adequate support or restraint they tend to continue moving into the cavity, *i. e.*, the more superficial tend to separate from the underlying tissues and so to undergo rupture. In this way are to be explained the hemorrhages with rupture affecting the urinary and gall-bladder, the stomach, and, more rarely, the intestines, following upon concussion, when these viscera ase filled with fluid.

In the case of the thorax, it has frequently been observed that sudden violent blows, insufficient to rupture the skin or fracture the ribs, have been followed by rupture of the lung substance with pulmonary hemorrhage and pneumothorax, or escape of air into the pleural cavity. The same explanation holds here, namely, the different rate of movement of the lung tissue and the contained air. Other effects may show them

MECHANICAL CAUSES OF DISEASE

selves as a result of thoracic concussion, namely, profound disturbance of the heart beat and of the respiration. These appear to be in part due to the profound stimulation of the vagi, though, as G. W. Crile' has shown, the condition of collapse, and even of death, mainly results from the mechanical irritation of the heart muscle itself. With regard to the effects of concussion upon the bones, from the recorded cases of fat embolism, or blockage of certain arteries by fluid fat, it is evident that sudden blows and, still more, the "jar" of the sudden arrest of a falling body on a hard surface may lead to extensive rupture of the fat cells of the marrow.

2. Puncture.—The most familiar example of this form of traumatism is in wounds caused by stabbing. Here we have pressure applied locally by a fine instrument sufficient to cause local solution of continuity of the tissues. The results largely depend upon the region involved, and very largely also upon whether the instrument pierces any higher nerve centre or large bloodvessel or hollow viscus in its passage, as also upon whether the instrument introduces at the same time infective agents. On the one hand, we may have little or nothing beyond merely local disturbance; on the other, sudden death, or we may have profound hemorrhage or general infection, set up either by bacteria passing into the tissues from hollow viscera or by bacteria introduced into the wound from without. Usually projectiles, travelling at a high rate of velocity, produce wounds which can be compared to puncture wounds in general, but according to the velocity and the size of the projectile, so it must be kept in mind that practically every form of traumatism above discussed may be brought about by projectiles—concussion, contusion, laceration, and puncture wounds. Still further, the very force with which projectiles suddenly impinge upon the tissues leads to a more general and immediate disturbance than is seen in the case of ordinary puncture wounds. At certain rates of speed there may be laceration of soft parts. concussion, and multiple fracture of the bones extending over a very large area.

3. Section.—This, which, surgically speaking, is the commonest form of trauma, consists in the separation of tissues by a sharp-bladed instrument, whereby there is a minimal disturbance to the tissues which do not come into immediate contact with the instrument. So small are the individual cells and tissues, and such their shape and arrangement, that it is impossible to introduce any instrument, however fine, in such a way as to insinuate it between the cells without injuring them. Consequently, all along the surface of a cut there must inevitably be a layer of injured cells. But in pure section, pressure and tearing effect upon contiguous tissues are reduced to a minimum by the pressure being brought to bear in a shearing manner, namely, the instrument is not merely employed as a wedge, forcing the elements of the tissues apart, but, by the oblique movement of the wedge, a cutting action is brought about and the tissues severed rather than forced asunder.

¹ Philadelphia Medical Journal, March 31, 1900.

POSTNATAL ACQUIREMENT OF DISEASE

4. **Contusion**.—In contusion, as distinct from concussion, we have the pressure exerted directly upon the part, and have a pressure exerted such that the elements of the part are torn asunder to a greater or less extent—hemorrhage resulting. The separation of the part may be slight (contusion), or, on the other hand, may be such as to produce separation of the constituents visible to the naked eye (laceration), or even may be such as to cause complete separation of one portion of an organ from another, as where a limb, or the scalp, or portion of the integument is forcibly torn off. It follows that, in the first place, we have hemorrhage from the ruptured vessels of the part, and that, in addition, we have more or less profound alteration of function; while, again, the sudden profound disturbance brought about by laceration or rupture of the nerves may set up general disturbances and shock. Later, the solution of continuity and exposure of the parts to the atmosphere may be followed by the results of infection of the wound or wounds.

5. **Compression**.—With regard to compression, little need be said here. Continuous pressure tells especially upon the more fluid portions of a tissue, and so it is that the vessels, both blood and lymphatic, tend to be occluded as a result. We have especially to deal with disturbances of nutrition in the part and with the accumulation of fluid in the vessels and the lymphatic spaces beyond the area of compression. Such compression may be external, and one is familiar with its results where Esmarch or other bandages have been too firmly applied to a limb; or where, again, in consequence of low blood pressure, the capillaries of the back or other portion of the body are emptied by the mere weight of the body, and bedsores (*decubrus*) result. Or it may be internal, as where tumors and collections of fluid developing in one or other part of the economy press upon the neighboring organs. Its results are malnutrition of the affected parts, with atrophy, which may go on to necrosis and disintegration of the parts.

6. Distension.-The mode of action of a distending force upon the tissues is similar to that of compression, namely, the pressure tends to act more especially upon the more fluid portions of organs, driving the fluid away, so that here again we tend to have malnutrition of the part subjected to a distension, and subsequent atrophy. We recognize a well-marked example of the effects of such distension in hydronephrosis of the kidneys, in which condition, in consequence of obstruction of the lower urinary passages, and of the continued excretion, we have eventually practically the whole of the kidney tissue proper undergoing atrophy, and the organ may come to be represented by an enormous thin-walled cyst. The distension may be of intravascular nature, and produce results of like order, as in passive congestion, brought about by heart disease or venous obstruction. Where, as in hemorrhages and the escape of blood into the brain substance, the distending force acts rapidly, we may, by compression of the surrounding vessels and the arrest of nutrition of the cells of the higher nerve centres, have a rapidly supervening death.

PHYSICAL CAUSES

7. Atmospheric Pressure (Gaseous).-In addition to the chemical changes brought about by the reaction between the contained gases and the tissues, we must realize that there is a purely mechanical interaction between the organism and the atmosphere, dependent upon the pressure exerted by the latter. We may add that in animals living in water the same is true in regard to aqueous pressure. There are, that is to say, certain limits to pressure within which life can continue and beyond which the continued manifestation of vital processes become impossible. In the case of man, this variation in pressure especially influences the system by influencing the partial pressure, as it is termed, of the gases circulating in the blood. Blood and other fluids take up larger or smaller quantities of gases according to the pressure in the gaseous medium; hence, if the atmosphere becomes rarefied, although certain troubles are induced by reduction of pressure on the various surfaces and the consequent dilatation of the vessels of these surfaces, the main disturbance induced is that the amount of oxygen taken up from the air is materially reduced and a condition of partial asphyxia is brought about, or oxygen hunger in the tissues. When the atmospheric pressure is much increased, the vessels of superficial parts are compressed, and the blood in consequence is driven from them into the deeper organs. It may be added that the main symptoms of "caisson disease" do not show themselves while the individual is subjected to a greatly increased atmospheric pressure, but develop when the transition from the increased to the ordinary atmospheric pressure is too sudden. Partly this may be due to the sudden alteration in the distribution in the fluids of the body which thereby occurs, but it has been clearly demonstrated that when under heightened atmospheric pressure, the blood absorbs or dissolves increased quantities of air, then upon sudden transition to a lowered pressure it can no longer hold this gas. As a result, the oxygen of the air having been taken up by the tissues, the nitrogen presents itself within the vessels in the form of discrete bubbles, these seriously interfering with the circulation in the smaller vessels.

PHYSICAL CAUSES.1

Temperature.—Owing to the remarkably sensitive and effective mechanism whereby the heat of the body is controlled, the human organism can stand exposure to a remarkably wide limit of temperature—can, on the one hand, continue to exist when the surrounding air is as much as 100° F., or 62° C. below the freezing point (Back); or, on the other hand, as shown long years ago by Blagden and Fordyce, is as high as 260° F. (126.6° C.)—or, roughly, about 50° F. above the boling point of water; and if the body be not exposed to these temperature.

¹ In revising this section, and more particularly in the section upon light and electricity as etiological factors, I have been especially indebted to the admirable survey of the subject afforded by Professor Askanazy in the first volume of Aschoff's *Pathologische Anatomie*, Jena, Fischer, 1909.

POSTNATAL ACQUIREMENT OF DISEASE

tures for too long a period, the general effects, save for alteration in the distribution of blood, are singularly slight-or, more correctly, are within physiological limits. Nay, more, it is a matter of popular knowledge that the hand may be plunged momentarily into molten lead without damage. The body, that is, can be exposed to temperatures far beyond those at which, on the one hand, protoplasm is frozen, and on the other, undergoes heat coagulation.¹ The explanation is that, through the warming of the air immediately over the surface of the body when that body is subjected to intense cold, there is developed an insulating layer of warmer air in immediate contact with the surface cells. In the case of extreme heat, a similar protective layer of cooler air is developed by the abundant giving off of moisture on to the surface, which moisture, by its rapid evaporation, keeps that surface cool. In general, however, it may be laid down that the tissues are much less sensitive to loss of heat (cold) than they are to excess of heat. We may recall the observations of John Hunter and others upon the actual freezing of limbs and portions of the body of various animals which demonstrate that, due care being taken, the parts retain their vitality when thawed out.² Later years have accumulated yet more striking instances. Thus, as already noted (p. 68), bacteria may be exposed to the temperature of boiling liquid hydrogen for considerable periods and still retain vitality, while Carrel's remarkable studies have shown that organs such as the kidneys of warm-blooded animals may be kept in a refrigerator at about the freezing point for several days, and may then, upon being transplanted into another animal, exhibit apparently unimpaired function. Askanazy records that portions of mouse carcinoma have been kept for two years at a temperature varying between -8° and 12° C., and have then grown when transplanted into another mouse. The essential difference would seem to be that heat destroys the cell enzymes; cold merely arrests their activity.3 These enzymes are essential to vital processes, although, as elsewhere laid down, it is the continued existence of the biophoric molecules that determines the continuance of cell life.4 In the higher warm-blooded animals it is evident

 1 Isolated human cells, e.g., red and white corpuscles, undergo heat coagulation at or about 50° C.

² Dr. Tytler and I have repeated these observations, freezing the rabbit's ear by carbon dioxide snow (which is stated to have a temperature of --79° C.), and have found that the least disturbance ensues, and recovery is most complete, by slow thawing out in ice water, coupled with minimal handling of the frozen part (so as not to damage the delicate vessel walls and cause later intravascular clotting).

^a Thus the success of cold storage depends not only upon the arrest of bacterial growth, but also upon the arrested activity of the autolytic ferments which are only effective at or about the body temperature.

⁴ As pointed out by Miss J. White (Proc. Roy. Soc., B., 81: 1009; 417), the germination of seeds depends upon their being placed in such conditions that now their disastatic, proteolytic, and other enzymes can become active. Nevertheless, these ferments of the seeds of cereals retain their activity when the seed has been kept under favorable conditions for twenty years and more: for long years, that is, after the vitality and germinating power of such grain has been lost. Wheat can germinate at longest after sixteen years; barley after ten years.

that it is not the actual freezing of the cells, but the subsequent thawing that destroys those molecules, even although the intracellular enzymes (as in the parallel case of autolysis) still continue active.

Frostbite.-The degree of cold, the duration of its action, and the mode of its application are all factors in determining the grade of disturbance set up by the local action of cold upon a part. Effects are more slowly produced by radiation (i. e., under atmospheric influences) than by conduction. Damp clothes, for example, rapidly abstract heat from the body. Nevertheless, as those living in northern climes well know, atmospheric radiation is far from being without effect, and is especially effective upon peripheral parts-ears, fingers, cheeks, and nose-when unprotected. Conduction, as in the case of the toes, also tells more upon the peripheral parts-in fact, the heat of the body being preserved by the circulation, it is the peripheral parts which, under the contracting influence of cold upon the vessels, are the most likely to become anemic. The first effect, therefore, of frostbite is a dead whiteness of the involved area. This pallor gives place to hyperemia of moderate extent, if the return of the circulation be gradual-as by rubbing with snow out in the open-to a severe hyperemia, with abundant exudation and pain if the return be sudden, as by entrance into a warm atmosphere. Even when the frostbite is very extensive, e. g., where the whole foot becomes frozen, provided that the thawing-out process and reëstablishment of the circulation be very gradual, as by placing the extremity in ice-cold water, it is remarkable how little reaction there may be, and how little destruction of tissue. This leads to the conclusion that it is not the actual freezing of the tissues that induces cell death, but the subsequent too sudden reëstablishment of the circulation. with resultant paralytic dilatation of the vessels, intense exudation, and circulatory stasis leading to malnutrition. What our experiments lead us to regard as an important factor in the serious results is the loosening of the frozen vascular endothelium by manipulation of the part. This subsequent vascular dilatation may be extreme; the exudation may lead to vesication, as in the second degree of burns; stasis and thrombosis of the vessels may ensue, and eventually a condition of frost gangrene. That it is the vascular disturbance that is at fault, and not the primary death of the cells through the freezing process, is further indicated by the fact that similar gangrene may affect the extremities of those with enfeebled circulation, alcoholics, etc., from mere immersion in cold water—*i. e.*, above the freezing temperature. The contradictory statements made by various writers upon the subject of frostbite are due to a non-recognition of these general principles, which, it may be added. underlie successful treatment.

Chilblains.—Common in England, for example, where the cold is but moderate, these are unknown in those parts of Canada where the temperature may for weeks be in the neighborhood of 0° C. (-32° F.). They would appear to be associated with the conduction rather than the radiation of cold, to be intimately related to cold, damp shoes and gloves, and, it may be, skin rendered damp through the moist atmos-

POSTNATAL ACQUIREMENT OF DISEASE

phere. In countries prepared for cold, non-conducting overshoes and suitable gloves, or, as the writer can vouch from personal experience, the non-employment of gloves, even in zero weather, effectually prevents chilblains. The underlying process would seem to be a recurrent arterial contraction when the part is subjected to cold, followed by a mild grade of continued hyperemia and slight ordema when the part is again subjected to warmth. This hyperemia may lead to some hypertrophy of the skin and corium. As pointed out by Askanazy, such can be experimentally produced in the skin of the rabbit's ear by repeated momentary application of the ether spray.

Hypothermia Produced by Cooling .- That the Red Indian children of northern North America, in the days before civilization was introduced. were to be seen playing naked in the snow is a recorded fact; or otherwise, the circulation of the whole surface of the body can adapt itself to external cold equally as well as can that of the localized regions, such as the face of the ordinary mortal and the legs of the Highlander. Such adaptation presupposes increased heat production by the internal organs to make up for the increased superficial loss. If through lack of heat-producing food, and through muscular inaction or exhaustion, the production is not equal to the loss-or if, again, through conduction (as by immersion in cold water) the loss becomes extreme-then inevitably the general temperature of the body falls below the normal. A fall to the neighborhood of 20° C. (68° F.) is incompatible with continued existence. There are cases on record of those found unconscious in the snow who have recovered, although the temperature had fallen to 75° F. Such lowering of the body temperature is accompanied by intense bodily weakness and lassitude, an invincible somnolence, slowing and weakening of the pulse and respirations, complete unconsciousness and loss of reflexes. There is, as shown by experiments upon animals, a preliminary stage of increased warmth-production, indicated by muscle tremors, using up of glycogen in the cells, etc., evidently under the influence of the nervous system, but eventually the lowered temperature arrests cell activities; the enzyme activities, as already indicated, become arrested, and eventually the circulatory and respiratory centres cease to function; nor, once arrested, is there any mechanism, it would seem (in the higher animals), capable of renewing their activity.

On Cold as a Predisposing Agent.—The expression "to take a cold" indicates the universal belief that in certain conditions, at least, exposure to cold is followed by results which we recognize more and more are of infectious nature. Here we would briefly note that the researches of the last few years indicate that there is an assured basis for this belief. It has been shown experimentally that hens, rabbits, and other animals immersed in water, and so brought into a hypothermic condition are more easily infected than are normal animals. Their powers of resistance are reduced. In ordinary exposure to cold to which men are subjected there is no question of hypothermia. What does occur is a contraction of the superficial vessels, a congestion of the more internal organs. So, also, whether as a result of the latter state, or as a reflex

act, there is a very noticeable "running at the nose"-the discharge of increased quantities of a thin, watery mucus. If this be long continued or repeated it is apt to change its character, becoming mucopurulent, and indicating the development of an acute (infectious) catarrh. Or, otherwise, following upon this congestion and discharge from the nasal. mucous membrane, there has been active proliferation of bacteria, presumably already present in the respiratory tract. We must presume either that, reflexly, there has been a diminution of the antibodies of the nasal secretion, or that there has been an exhaustion of the same. The acute enteritis that may follow the swallowing of ice-cold water when the individual is heated, as also the pneumonia that may follow the sudden change from heated to cold air, must be of the same order. In all cases of this nature the "may" enters: two individuals may be subjected to identical conditions; the one is taken with pneumonia, the other not; and, confessedly, it is difficult to weigh the factors leading to the disturbance. We know, as a commonplace, that pathogenic organisms are normally present on the surface of our bodies. It would not, therefore, seem so much that these agents are present in the one case and absent in the other, as that the resisting power of the individual varies-that it is a question of antibodies and protective mechanisms, and the fact that the ill results of taking cold show themselves more particularly in the very young and the very old, where protective mechanisms are, from every consideration, seen to be less adequate than are those of the adult, as again, they are apt to be manifested in those of known weakness of constitution-alcoholics, convalescents from other illnesses, etc.--indicate that herein is the main explanation.

Heatstroke .- When the air is already saturated with moisture and evaporation cannot occur, the high temperature rapidly becomes dangerous, and morbid disturbances may develop; as, indeed, they may through prolonged exposure to excessive dry heat, when perspiration is arrested or the amount of fluid available for purposes of evaporation is exhausted. as in long hot marches without water. Such heatstroke-not to be confused with sunstroke-may occur in hot, sultry, overclouded weather. It is characterized by: (1) Rise of body temperature, both surface and rectal; it may be as much as 10° F., a rise not merely due to the external warmth, but, as indicated by experiments on animals, to disturbance of the heat-regulating apparatus, so that under the stimulus of increased heat there is increased cell activity, with increased discharge of CO. and absorption of O. (2) Headache and feeling of oppression, followed by loss of consciousness and convulsions. (3) Rapid respiration. (4) A definite but not corresponding increase in the pulse rate. (5) As shown by slowly fatal cases in man and by animal experimentation, dissociative changes occur in the tissues leading more particularly to fatty degeneration. (6) In rapidly fatal cases there is swift onset of rigor mortis and early onset of putrefaction.

In these cases we deal apparently with the direct effects of heightened temperature upon the nervous system and tissues generally, in which the increased extreme temperature, coupled with inadequate discharge of

POSTNATAL ACQUIREMENT OF DISEASE

heat, stimulates the tissues not to reduction, but to increased heat production.

Sunstroke (Insolation) .- In this the effects are largely, though not wholly, independent of the temperature of the atmosphere, wholly independent of the atmospheric moisture; they result directly from the action of the sun's rays upon the unprotected head or neck. The symptoms are: (1) Severe headache and pain in the neck; (2) nervous excitement and hallucinations, convulsions and coma. Death may occur within an hour or two after the onset of the symptoms; milder cases are followed by recovery, with, however, a residue of nervous disorder of one or other form. The disturbances, it will be seen, essentially influence the higher nerve centres in the brain and medulla. At autopsy there is found a condition of intense congestion of the brain, with cedema of the meninges. Sambon and others have seen here an acute infectious meningitis, and have ascribed these cerebral disorders to particular microbes. At most, it may be laid down that the meningeal disturbances predispose to local infection. The immediate heating effect of the light rays upon the central nervous system appears to be the prime cause.

Burns.—We may regard sunstroke as an outcome of radiant heat. Where there is direct conduction of extreme heat to any part of the body, burns are the result. Of these burns we are accustomed to recognize four grades:

 Erythema, or simple superficial inflammation, with hyperemia and pronounced reddening of the skin; slight swelling or inflammatory ordema, and marked pain from irritation of the superficial nerve endings.

2. Vesication.—The irritation is here more severe, and is accompanied by a certain slight grade of necrosis affecting the deeper vegetative cells of the Malpighian layer of the epidermis, together with a great accumulation of clear serous fluid, almost devoid of leukocytes, to form larger or smaller blisters, with separation of the epidermis from the corium. There is accompanying intense inflammatory congestion of the superficial vessels.

3. Necrosis.—Yet greater heat leads to generalized death of the epidermis, and it may be of the superficial portions of the corium, with coagulation of the blood in its contained vessels. The affected area, at first of a rich red color, soon becomes brown, leathery, and mumnified, although, through the accumulation of fluid and inflammatory products beneath (often becoming infected), this dry outer crust may become loosened, leaving a soft and purulent granulating surface. Usually the necrosed surface is not of any great depth; through the insulatory character of the human skin, even after exposure to boiling water for some little time, the underlying muscles may only exhibit a temperature of from 50° to 60° C. Prolonged action of boiling water will, however, result in the deeper tissues becoming "cooked."

4. Carbonization, or charring, results from the direct and prolonged action of flaming heat or molten metal. As a result, the affected parts may be almost unrecognizable; the blackened, dried skin may peel off from the underlying parts, the joints may be exposed, the intense con-

traction and shrinkage of the muscles lead to rupture of their tendons. The bones even may be charred and through the boiling of the soft brain the brain case may be burst open.

A fifth and final stage may here be noticed, namely, as seen in cremation, the carbon and all organic matter may be burnt off, leaving but the whitened inorganic ashes.

If more than half the surface of the body be burnt, death surely supervenes within twenty-four hours, while burns of the third degree, involving only one-sixth to one-eighth of the surface, are liable to lead to death in three or four days. The symptoms are primarily and predominantly nervous, and are those of *shock*. Broadly speaking, the larger the area involved, and the greater the number of nerve endings and neurons irritated, the greater the shock. This shock is a state of profound nervous exhaustion or inhibition, which may or may not be preceded by a stage of excitement and delirium. (See Chapter X, Section III.)

There continues to be debate as how far nervous conditions, how far changes in the blood, and the products of the necrosed tissues, as, again, of the decomposition of those tissues under bacterial agency, with entrance of bacteria from them into the unburnt tissues, are responsible for the subsequent symptoms. While through loss of the skin there is increased loss of heat, and the surface temperature is markedly reduced. the rectal temperature may be elevated. This cannot be ascribed to nervous agencies pure and simple, but is in harmony with what we know regarding the influence of dissolved cell and tissue products in producing a state of pyrexia or heightened temperature. So, also, the characteristic changes seen in the lymph nodes at a distance, with extreme enlargement, proliferation, and death of the central cells of those nodes described by J. McCrae and Bardeen, cannot be a nervous phenomenon, but must, like the similar condition found in diphtheria and typhoid, be due to the effect of circulating toxic substances. Nevertheless, the modified blood from cases of burns, as, again, the burnt tissues, have not been found to exhibit specific toxic effects when injected into the lower animals. Further observations, it may be urged, are requisite upon the results of injection of blood, as, again, of fresh tissues heated to various degrees.¹ Apart from local destruction of corpuscles in the burnt area the blood may show pronounced changes, more especially polycythemia, or apparent increase in the number of corpuscles as a consequence of the loss of fluid through drainage and exudation of the same into the burnt area. Secondary results of the blood changes are the appearance of hemoglobin in the urine and enlargement and deep coloration of the spleen-the organ in which damaged erythrocytes are removed from the circulation.

¹ The red corpuscles, for example, react differently—and must yield different products—at different temperatures. At 50° C, they show crenation with peripheral liberation of minute globules of their substances; at 60° C, they become laked, liberating their hemoglobin; at 70° C. they undergo immediate coagulation, without either of the previous changes. Nevertheless, as already stated, the profound nervous disturbance is the dominating feature, and in this connection, at autopsy, the brain is apt to be found profoundly hyperemic and œdematous.

In cases that recover, the healing of the destroyed skin surfaces is apt to be accompanied by extensive cicatrization, with contracture, leading to much maiming and disfigurement.

Light.—We rarely have to deal with light as a direct cause of bodily disturbance; nevertheless, we are yearly gaining increased evidence that it may have very definite pathological effects, or, more accurately, that the sun and other sources of light afford rays of very different wave lengths, of which those with longer waves toward the red end of the spectrum are relatively harmless, save for their heating effects, whereas those of short wave length toward the violet end, and the yet shorter ultraviolet rays beyond the visible portion of the spectrum, have both greater penetrating power and very material influence upon the tissues. It is these violet and ultraviolet rays which more particularly are involved in the destruction of bacteria and the lowest forms of life by the action of light; which, again, have been found capable of destroying or rendering useless various enzymes; which produce freckles, sunburns, and the severer skin lesions, with pain, hyperemia, and active inflammation, even to vesication, affecting those parts of the body exposed to intense sunlight, notably under conditions when the heat of the sun is not intense, as upon the snowclad Alps, or on the vast expanses of snow in Arctic and Antarctic regions. The hyperemia of the skin in these cases would seem to have a protective function. It has been shown that hemoglobin absorbs the light rays of shorter length. So, too, freckles and the bronzing of the skin of those much exposed to sunlight can only be regarded as adaptive, for this pigment also actively absorbs the light rays, and thus prevents them from injuring the deeper tissues. There is a basis of observed fact and of experiment for Woodruff's contention that prolonged exposure to the intense sunlight of the tropics is actually deleterious to those fair-skinned Northerners with little protective cutaneous pigment. It is evidently adaptation plus survival of the fittest that has led to the intense pigmentation of the negro.¹

The effects of ultraviolet light, as observed in the rabbit's ear, are not unlike those of burns of the first degree. Subjected for an hour to these rays, the epiderm cells show vacuolization and nuclear chromatolysis, passing to complete necrosis; changes are also seen in the vascular endothelium, with dilatation and congestion of the vessels of the cutis, exudation, and some escape of both red and white corpuscles into the tissue. Even thrombosis (intravascular coagulation) has been observed. This accute stage is followed by active and at times excessive regeneration of the damaged tissue. It is paradoxical that pigment-holding cells, instead of being damaged, are stimulated by these and other rays, as shown by the active expansion of the chromatophores of the skin of the frog and other animals, which thus act, to quote Askanazy, as

¹ Effects of Tropical Light on White Men, New York, 1905.

LIGHT, X-RAYS, RADIUM RAYS

"parasols." In forms like the squids (Cephalopods), which have different orders of pigment cells, while all the chromatophores (both yellow and red) are affected by ultraviolet rays, the red chromatophores become expanded by yellow light and not blue, the yellow cells by blue although not by yellow rays. It has been suggested that the anemia of moderate grade from which miners suffer may be due to the lack of action of light upon the red corpuscles, for it has been found that under the action of the shorter rays the erythrocytes absorb more oxygen.

A remarkable series of observations of recent date is that the tissues may be sensitized to light by actively fluorescent substances like erythrosin and eosin, so that now they permit a penetration of light of longer wave length. Paramoccia and other protozoa, which proliferate actively in diffuse light, are killed in a few minutes if eosin be added to the water, although when they are kept in the dark such eosin is without pronounced effect. Contrariwise, subpate of quinine and the smearing of the skin with black or dark grease acts as a protective against the ultraviolet raves by absorbing them.

Such a sensitization would seem effective in certain diseased conditions, notably in pellagra, which during the last two years has been found present over a wide area of North America. The subjects of the disease show an extraordinary grade of inflammation of the skin of the neck, wrists, and other exposed parts, coming on usually in spring. In occasional individuals, often as family idiosyncrasy, sunburn leads to a productive dermatitis (*Xeroderma pigmentosum*).¹

It is almost unnecessary to recall that these various observations have been applied therapeutically, as in the use of the Finsen light and of ultraviolet rays, to destroy superficial growths, lupus, etc. Here the underlying principle is to utilize the violet and ultraviolet rays for such a time and in such a way as to induce a limited necrosis of successive areas of the morbid growth, followed by regeneration and cicatrization.

X-Rays (Roentgen Rays) and Radium Rays.—These, although invisible to us, are, from their nature and effects, reasonably to be classed along with the light rays, although their effects, if of the same order, are far from identical. The more noticeable features regarding the *x*-rays are:

1. The hyperemia and inflammation of the skin are of much slower development than with light rays; the effect is markedly cumulative, and, as a result, a very obstinate dermatitis may be set up, of varying grades of severity, according to the nature of the tube employed, distance from which it acts, and duration of employment. There may be merely a somewhat purplish and painful erythema, leading to pronounced pigmentation, or a more chronic affection (generally after repeated treatments of moderate intensity), with thickening and fissuring of the skin, falling out of the hair, and loss of secretion through atrophy of the skin glands. This, as in Xeroderma pigmentosum, is apt to give place to malignant, epitheliomatous overgrowth, or, on the other hand, it may be followed by cutaneous atrophy. Or, without preceding overgrowth

¹ For fuller study, see Councilman and Magrath, 5th Rept. Harvard Cancer Commission, 1909; 5, and Jour. Med. Research, 21; 1909.

the inflammation may give place to necrosis and intractable ulceration, singularly slow in showing any tendency to heal.

2. They penetrate much more deeply than do the light rays: the stronger the vacuum in the tube, the greater the penetrative power. In this way pronounced and destructive effects may be wrought in the deeper organs. They specially affect—

(a) The lymph nodes and spleen, causing a rapid death of the lymphocytes, more particularly the vegetative cells in the germinal centres. The same occurs also in the red marrow: eventually regenerative changes occur, and with this there may develop a pronounced leukocytosis. These observations have led to the treatment of leukemia by the x-rays, with the result that a remarkable temporary improvement is brought about, the number of leukocytes falling to or below the normal; malignant lymphomas have also been greatly reduced in size. The betterment, however, is not permanent.

(b) Malignant Growths in General.—It may be laid down as a general principle that cells of a vegetative type, and in the process of active proliferation, are especially sensitive to the *x*-rays. It is for this reason that the Malpighian layer of the skin is particularly affected, and that skin ulcers are so long in healing over, as also that the germinal centres of the lymph nodes are picked out. Actively growing tumors thus easily exhibit cell necrosis and reduction in size under the *x*-rays. The destruction, however, is rarely complete, there is a pronounced tendency to be less influenced by the rays.

(c) The Essential Genital Glands.—It is now well established not only that those working with the *x*-rays, unless elaborate precautions be taken, are liable to become sterile, but that by the direct action of the *x*-rays upon the testicles and ovaries of man and of vertebrates in general a condition of *azospermia* (or absence of spermatozoa) and atrophy of the ovarian follicles, with menstrual disturbances and barrenness, respectively, may be brought about. So, also, abortion has been induced in the pregnant animal. The action of radium is similar, but more rapid and powerful; the effects upon deeply placed organs or tissues are even more pronounced.

(d) Several recent observers have called attention to the profound influence exerted by x-rays and radium upon the development of larvæ of different species. (See the discussion on Abiotrophy.)

What is the underlying cause of these destructive effects of the different orders of rays is still undecided. In the action of ultraviolet rays upon bacteria, increased oxygenation with formation of hydrogen peroxide, and briefly more active ionization with dissociation of the living molecules, has been suggested. An action of this nature is not unlikely and would explain the increased cell activity and subsequent proliferation brought about by slighter exposures, the cell disorganization and death due to more intense action. But also we have to recognize with all these agencies a tendency toward enzyme destruction, while observation upon the effects of radium upon the ovum show that the phosphatides become dissociated with liberation of cholin, a body of

ELECTRICITY

distinctly toxic properties. Comparable with this is the increased photoactivity of the tissues exposed to these rays. From what we know of the luminosity of the firefly and the photoactivity of living matter generally, this is intimately associated with the presence and oxidation of phosphatides. Possibly all these phenomena come under the one heading of accelerated and dissociative oxidation.

Electricity .- With its increasing employment in every-day life, electricity has rapidly assumed a more prominent position among the causes of disease and injury to the organism. But perhaps as a consequence of insecure knowledge of electrical phenomena possessed by most medical men, singularly little has so far been written upon this branch of our subject. Indeed, the results of the passage of electric currents through the body have been found so varied that it is difficult to make any but the most general statements. Like other agents, electricity depends for its effects upon the dose and the mode of administration. According to the method whereby the electric current acts upon and gains entrance into the tissues, according to the relationship of the body to other substances which are not conductors, and according to the intensity and nature of the current, so do we obtain very diverse results. When we call to mind that physicists are demonstrating more and more clearly that the electrical charges of ions, atoms, and molecules determine chemical action, it cannot but be realized that brusque changes induced in the electrical charges of the cell contents must profoundly affect metabolism—nay, may lead to dissociative changes of the severest grade. We may, in the first place, distinguish between the effects of electric discharge, or *fulguration*, and of the more ordinary current.

Fulguration.-We can reproduce this form of electricity by the discharge of a Leyden jar, whereby there is an immediate establishment of electrical equipoise. It occurs in nature as lightning, and as such the volume of the electricity passing between the positively charged cloud and the negatively charged earth in a single flash, has been calculated to be hundreds, if not thousands, of amperes. The effect upon the human body interposed in the passage of the "flash" is very remarkable. Comparatively little of the electricity gains entrance into the tissues; most passes over the surface of the body and through the clothing in such a way that the clothing is violently rent. More particularly, the nails may be drawn out of the soles of boots, the soles, and, it may be, the whole boots, violently wrenched off the feet. But at the regions where the electricity comes into contact with and leaves the tissues, there is intense local destruction and rending of the tissues, with effects as of burning, from the resistance offered by the tissues to the passage. Curious reddened arborescent "lightning figures" may radiate over the skin from the region of entrance. These are not burns, nor are they due to hemorrhages along the subcutaneous capillaries, since in nonfatal cases they disappear in the course of a few hours; their exact nature is undecided, although probably they are due to an intense hyperemia of superficial vessels. Nevertheless, lines of burning or electrolytic necrosis of the skin, as also superficial hemorrhages, are not uncommon.

POSTNATAL ACQUIREMENT OF DISEASE

296

Occasionally the electric current has been found to pass along the vessels to the heart, there causing rupture. Whether a lightning stroke is fatal or no depends largely upon the region struck; a fatal result would seem always to follow when the head is involved; where an extremity only, there, in general, recovery ensues. But in all cases there is immediate unconsciousness, and this, in cases that recover, may last for some hours, with gradual recovery of the pulse and respiration rate. The burns heal with extreme slowness. In fatal cases there is rapid, sometimes immediate, development of rigidity, and at autopsy, along with intense congestion of the internal organs, capillary hemorrhages may be found in the central nervous system and elsewhere. The treatment of malignant growths by local application of the electrical discharge has been introduced too recently to permit any assured statement as to its value.

The Constant and Alternating Currents.—Under pressure or voltage such as is used commercially, the constant and interrupted currents manifest very similar effects, save that, with like voltage, the effects of the latter are the more severe,¹ and tell more strongly upon the heart, producing more readily, as seen in animals, that condition of fibrillary contraction which most often is followed by complete arrest of heart beat. There is a remarkable variation in the recorded voltages which have been found fatal to man. These range from 100 to 2000 volts. Evidently very many factors are at work determining the fatal effect. Some, at least, of these are:

 Condition of the Skin.—The dry skin is a poor conductor of electricity; the moist or wetted skin permits the entrance of a larger volume of electricity, and this amperage is apparently a factor of very considerable importance in determining the extent of the lesions.

2. Area of Contact.—When this is extensive the effects are more diffuse; when small, more restricted, with more extensive local disturbance.

3. Position of Contact.—Extremely high voltage may pass between two neighboring points on the trunk or limbs, without fatal result, and with, in the main, local effects, which, however, may be severe and of the type seen in lightning strokes, with burning and necrosis of the tissues, muscular contraction, etc. Currents of moderate frequency and relatively low voltage may produce fatal effects when the heart is interposed between the points of entrance and exit.

4. Duration of Contact.—A current of 2000 volts and more, if its action be but for the fraction of a second, has little after effect; although the same repeated may be fatal, as it is if in action for from five to ten seconds. The amount, that is, of electricity that can gain entrance into the system through the dry skin from momentary contact is very small.

5. Frequency of Interruption.—It is a singular fact that a moderate frequency of interruption has far more serious effects upon the organism than has high frequency. In fact, as demonstrated by D'Arsonval and

¹ Whereas, the average total voltage of the interrupted current is in the neighborhood of 500, that of the constant current is about 1500.

ELECTRICITY

confirmed by Tesla, the human body may be placed in a circuit of from 100,000 to 200,000 volts, and if the interruptions be as many as 10,000 per second, all that is experienced is at most a comfortable glow with a feeling of warnth in the hand holding an incandescent lamp brilliantly lighted, being held in such a manner that the current passes from the body and through its carbon filament. The experience in the United States is that a frequency of 130, a voltage of 1500 to 2000, with one pole on a moistened surface over the brain and the other applied to one of the lower limbs, and a duration of contact of five to seven seconds, repeated several times, followed by a current of low voltage (200 to 300) for half a minute, brings about a sure and effective electrocution.

6. Voltage.—Experiments on dogs demonstrate that a voltage of 120 and below causes death by cardiac paralysis and fibrillary contraction. Above 120 the effects are more marked upon the respiratory centre. Indeed, as Oliver has shown, with high voltage (e. g., 4800) the heart is not paralyzed; with cardiac massage and artificial respiration, the circulation can be reëstablished and the respiratory centre gradually recover its function. It is to assure both respiratory and cardiac paralysis that the currents of high and low voltage are employed in electrocution. There are, indeed, indications that very high voltage (over 2000), like very high frequency, are less harmful than a moderate grade.

The studies upon electrocuted criminals have afforded more accurate information regarding the succession of events in severe electrical shock than have the numerous cases of industrial accidents. The following is seen to be the course of events: intense muscular contraction; syncope and unconsciousness; clonic convulsions; tonic convulsions; arrest of respiration; stoppage of heart. Recovery in industrial cases is followed by no permanent ill effects, save when the local passage of a current of high voltage has had intense effect upon the tissues. The burns heal very slowly. Whole muscles may necrose and slough away, and the resultant wounds take months rather than weeks in undergoing cicatrization.1 In fatal cases (electrocution), Spitzka2 notes that the maximum change is in the nervous system; there is no apparent change in the nerve cells themselves, but disruption and destruction, with capillary hemorrhages. In the pons, medulla, and cord, curious circular areas are encountered, with rarefied centres and peripheral zone of condensation, suggesting electrolytic liberation of gas in the perivascular spaces. There is evidence of some burning of the tissues at the region of application of the electrodes, brought about by the resistance by the tissues to the passage of the current. The blood is dark brown and rarely coagulates (? destruction of fibrin ferment). As in other cases in which, as in tetanus, and strychnine poisoning violent muscular contraction ushers in death, the temperature rises promptly after death, and this to as much as 120° F. to 129.5° F., within twenty minutes (Spitzka), the maximum being found at the site of the leg electrode, where the muscular rigor and coagulation are most extensive.

¹ Elder, Montreal Med. Jour.

² Proc. Amer. Philos. Sc., 47: 1908: 39.

POSTNATAL ACQUIREMENT OF DISEASE

The induction apparatus (Ruhmkorff's coil) has been found capable of causing death in the lower animals, and that by arrest of respiration through tetanus of the respiratory muscles. The amperage of this form of electricity is so low as scarce to be followed by serious effects in man.

Vibratory Motion.—The effects of rapid vibration upon the organism have not been fully worked out. As Meltzer' has indicated, vibration is essential to life. A certain minimum is indispensable, and must be regarded as associated with the molecular activities of living matter. Beyond a certain frequency in extent, vibration is, however, harmful. He and Welch have studied the effect of rapid vibration upon the blood corpuscles and have shown that, over a certain point, the cells become broken up when subjected to rapid vibration.² More recently, Meltzer and Shaklee³ have shown that rapid oscillation continued for a few hours renders pepsin, rennin, and trypsin completely inert. The same effect has been found by Abderhalden in connection with tyrosinase and zymase.

CHEMICAL CAUSES.

Under this heading we include all those causes in which there is a direct molecular interaction between the noxa and the constituents of one or other set of cells in the organism. We have, therefore, to consider not only those cases in which there are gross effects leading to immediate death of the cells, as by the action of caustic and other agents, but also those cases in which the cells, while not destroyed, have their functions arrested or disturbed without actual death of those cells being the immediate consequence.

Poisons.⁴—Chemical substances having a deleterious action upon the cells of this body may exert that in two ways: either they may immediately destroy or severely irritate the tissues with which they primarily come into contact, or, becoming absorbed, they may become diffused in the circulating fluids of the system, and thus have an action upon cells at a distance from the point of primary contact. The first order we speak of as caustic agents; the second as intoxicants. Both are poisons, if we accept the definition of Kobert, that "poisons are non-organized substances, organic or inorganic, existing within the organism or introduced from without, which, from their chemical constitution, are able under certain conditions to be harmful to living beings, by destroying or affecting their health or relative well-being." The series of changes occurring in the organism in general in consequence of the action of such poisons is known as *intoxication*, and may be of many orders.

The somewhat involved nature of the definition we have given be-

¹ Zeitschr. f. Biol., 1894. ² Jour. of Physiol., 5; 1884; 255,

³ Proc. Soc. Exp. Biol. and Med., 6: 1909: 48 and 103,

⁴ I have incorporated into the ensuing sections portions of a paper read by me at Washington, Trans. Assoc. Amer. Phys., 16:1901:38.

CHEMICAL CAUSES

comes justified when we realize that poisons are only such relatively; substances are harmful only when present in sufficient concentration either to set up molecular disturbance and chemical change in the protoplasm of the cells or to arrest normal molecular changes in the protoplasm. And, conversely, it follows that all substances capable of solution in the body fluids and absorption into the cells composing the body are also capable of acting as poisons, and this because the effect upon the cells depends upon the extent of their absorption, and above a certain point (the limit varying with each substance) their action is unfavorable to the continuance of the orderly cell processes. Thus, to give the most notorious example, water, which is absolutely essential to existence, and forms 70 per cent. of the body weight, and in some cells as much as 90 per cent.¹ if introduced into the tissues above a certain amount (60 c.cm.) per kilo of body weight, is found to be a poison, and, indeed, fatal. It causes the diffusion of hemoglobin out of the red corpuscles, preventing due oxygenation, and absorbed by the cells of certain tissues in excessive amounts, it deleteriously affects their activities, among other ways, by inducing undue ionization. It follows, thus, that the number of potential poisons is enormous.

How are we to classify then? The old familiar division into animal, vegetable, mineral, and gaseous is, for pathological purposes, absolutely useless; no particular sets of reactions follow upon the action of animal poisons, for example, as distinct from vegetable; the effect of a mineral may simulate that of a vegetable poison. Nor, for our purposes, useful as it would be, can we classify them according to the symptoms which they originate, and this because, with a given poison, the symptoms caused by one concentration may be widely different from those of another, and in one individual the effects produced may differ widely from those produced on another. It is more to the point to observe the changes in the individual tissues and to attempt to classify the toxic bodies in relationship to various forms of intoxication, *i. e.*, of the changes induced in the organism. This we do to a large extent in our study of the degenerations.

Ehrlich's Two Groups.—Ehrlich has introduced a broad division of pharmacological—and toxic—agents into two main groups, which must here be noted. Like that just suggested, it bears more particularly upon the reaction of the cell to the poison. The assimilation of foodstuffs, as we have pointed out in our chapter upon Growth (p. 98), is by him regarded as a linkage of side-chains to the nucleus of the living molecule, or, as we would express it, biophore. The members of Ehrlich's first class bear a close resemblance in this respect to the foodstuffs. He regards them as being linked to the living protoplasm. Their effects are not immediate; time is required, as he supposes, that they may become built into the living molecule, and so they exhibit what we may term a long latent period before developing these results. The members of this group are one and all the products of living matter.

¹ The gray matter of the fœtal brain.

POSTNATAL ACQUIREMENT OF DISEASE

and if not proteins, are of proteid affinities. Such are the bacterial toxins, the venoms of various animals, certain plant poisons, such as abrin, ricin, robin, certain definite poisonous proteins, and Ford would add certain glucosides. With the majority of these the result of union with the living molecule is to give rise to the discharge from the cells of *antibodies*, antitoxins, and the like.

The second group, while diffusing into the cell, and having very definite effects upon the cell activities, is regarded by him as not being built up into the living molecule; there is no latent period, the action is immediate, and there is no production of antibodies. Into this class come all the poisons not included in the first. How these act, Ehrlich finds it difficult to lay down, save that they do not become assimilated by the biophores. Some, like the anesthetics as a body, can be shown not to enter into chemical combination with the cell substance, since they can be recovered intact by simple processes. It is well within the bounds of possibility that the process affecting these bodies is of the nature of adsorption rather than true chemical combination;¹ or, again, it may be that some at least enter into chemical combination with the cytoplasmic constituents and act by blocking the intermediary substance and so arresting the activities of the biophores. The probability, indeed, is that this huge class will become divided into several divisions. Intermediate or contradictory data have been brought forward; thus some alkaloids (of the second group), like colchicine, exhibit a definite latent period, while some toxins (of the first), like those of the "Nasik" spirillum, can induce fatal results within half an hour.² Save for its suggestiveness, this classification of Ehrlich's is altogether too broad for our immediate purpose.

Exogenous and Endogenous Poisons .- Studying the degenerations, we determine that the poisons introduced from without are not the sole intoxicants. We can, indeed, proceed to establish two broad groups, the *exogenous* poisons arising outside the system, the *endogenous* arising within. But we have to be careful about our conception of what is endogenous and exogenous, and laxity in regard to these matters has caused great confusion, and has rendered particularly useless one term -auto-intoxication-capable of being of high value. According to vulgar parlance, the contents of the alimentary canal are a part, and an important part, of the "inside" of the individual; yet, strictly speaking, the food taken is not within the system until it has been absorbed by the cells of the organism. The following crude diagram (Fig. 110) of the simplified individual will make this point clear. It will be seen that only that which is within the shaded portion is really within the body. Intoxication arising from the absorption of substances from any point outside the shaded area must be regarded as exogenous. It was by an unfortunate lack of clearness regarding this—a lack unusual in

³ For a useful study of adsorption phenomena, see Bayliss, Biochemical Journal, 1; 1906; 175,

² Emery, Immunity and Specific Therapy, London, 1909; 41.

EXOGENOUS AND ENDOGENOUS POISONS

French writers—that Bouchard, who popularized, if he did not originate, the term *auto-intoacication*, classed all the toxic results, not only of perversion of cellular and tissue activity, but also of gastro-intestinal fermentation and the absorption of the products of bacterial activity in the digestive tract, under the one indiscriminate heading. It is true that the absorption of the products of bacterial activity, of fermented and altered foodstuffs, may, secondarily, affect the metabolism of the cells of sundry organs, and the products of perverted activity may be the direct cause of the general disturbances; indeed, in such cases we have a genuine auto-intoxication; but then the intestinal irritants become only the predisposing cause, not the direct. From this point of view every poison acting through the alimentary canal, from alcohol to mineral acids, is to a greater or less extent an auto-intoxicant, and this is clearly an absurdity.



-Intoxication by bacterial products is in no sense auto-intoxication, nor is it permissible to speak of gastro-intestinal auto-intoxication unless thereby is meant (and this the users of the expression rarely mean) that matter excreted into the bowels by the cells undergoes reabsorption. In consequence of this vague and illogical use the term has fallen into disrepute.

These considerations lead us to recognize that, from a pathological standpoint, we have to distinguish two distinct orders: (1) Those intoxications set up by substances actually derived from the cells of the organism, which alone are to be regarded as truly endogenous; and (2) all other intoxications set up by substances foreign to the economy: all these are exogenous. Having established this, we can proceed to divide these into their classes:

POSTNATAL ACQUIREMENT OF DISEASE

I. Exogenous Intoxications.

 Non-parasitic.—Intoxications due to the actions of poisons not produced in association with the organism, which gain an entrance into the system through the skin, digestive, respiratory, or urinary tracts.

2. Parasitic.—(a) Parasitic proper, due to the introduction into, and growth within the tissues of parasites of various orders, animal and vegetable, which, growing, give rise to toxic substances.

(b) Saprophytic, due to the growth of parasites of various orders on one or other surface communicating with the exterior of the organisms, the products of growth becoming absorbed and diffused into the tissues.

II. Endogenous Intoxications.—Of pure type; auto-intoxications proper.

1. Internal secretory, intoxications due to altered internal secretions on the part of the body cells affecting (a) the secretory cells and tissues themselves, and (b) the other tissues of the organism, through diffusion of the altered products of cell activity.

2. Disintegrative, due to the absorption of the products of disintegration of dead cells (e. q., in burns, internal hemorrhages, etc.).

3. *Metabolic*, the results of impaired metabolism and imperfect excretion.

III. Intermediate or Mixed.—Of impure type.

1. Obstructive, due to arrested elimination.

2. Gastro-intestinal.

One of the classes here formulated calls for remark, namely, that which I have termed the Intermediate. As will be noted more fully in the pages devoted to it, while (1) in certain cases it is quite clear that we deal with the effects of reabsorption of external secretions pure and simple, these cases are relatively rare. (2) In other cases there are indications that we deal with failure to secrete, with accumulation of metabolites that normally undergo discharge. (3) In yet others, with the deleterious effects of the absorption, not of the excretions themselves, but of toxic derivatives of the same, the products of bacterial disintegration. In such cases the poison itself originates outside the system, and is to that extent exogenous, although derived from a substratum of endogenous origin.

It is in this order that the various orders of intoxications and their relationship to the development of disease will be discussed.

CHAPTER VII.

EXOGENOUS INTOXICATIONS: NON-PARASITIC.

FOREIGN substances undergoing absorption or gaining entrance at the surface of the body have a twofold action: (1) Local, at the point of application, and (2) general. Only in rare cases is the first action wanting, as, for example, in that of hydrocyanic acid above a certain strength, when the general effects are so swiftly induced that there is no time for the local disturbances to show themselves.

For the local effects we have no special name; according to the nature of the poison, these are either degenerative or necrotic; and, if time be allowed, they are followed by evidences of inflammatory reaction of one or other grade.

Studying the general effects, such toxic agents either:

1. Cause arrested cell activity from the first.

Ce

n

2. Cause increased cell activity, followed by exhaustion and paralysis of function.

3. Cause increased cell activity, followed by disintegration.

Paradoxical as it may seem, the only satisfactory way for us to classify these exotic poisons as causes of disease is by a study of their effects. A given poison produces a particular chain of general disturbances; it is for us, as pathologists, to determine the basal lesion or lesions giving origin to these symptoms. Attempting this, we find that the different toxic agents have selective actions on one or other tissue. Thus it is these selective effects that must form the basis of our study, while secondarily in connection with each, it has to be determined whether the effect is characteristically inhibitive, irritative, or disintegrative.

It is the study of these exotic poisons that forms the basis of toxicology, and, to a considerable extent, of pharmacology. It is in works dealing with these subjects that extended observations upon the mode of action of any particular poison will be found. Nevertheless, we, as pathologists, have to consider these poisons, but approach them from a different standpoint; not from that of the individual drug or poison, but from that of the organism and the disturbances therein induced. It becomes necessary, therefore, that here we should at least call attention to this subject, if only in outline, keeping before us throughout that our endeavor at this point is not to discuss the general effects of poisons—the disturbances set up in a body at large. These constitute the *processes* of intoxication. It is the primary effects leading up to these general disturbances that here concern us. As already laid down, the poisons one and all exhibit a more or less obvious selective

304 EXOGENOUS INTOXICATIONS: NON-PARASITIC

action on the different tissues; we shall, therefore, pass in review the tissues most liable to be affected.

Poisons Acting through the Nervous System.—The system which most commonly appears to be affected in intoxication is the nervous system. This is, after all, what is but to be expected, for of all cells of the body the neurons are the most highly differentiated and most responsive to stimuli. At the same time, it has to be admitted that classification, according to their different effects upon the nervous system, is not easy, and this because the effects of a poison (1) vary very largely according to its concentration in the blood and rate of absorption by the neurons; (2) nor can we with any success apply data obtained from animal experimentation to the case of man. The higher we ascend in the scale of animals the more complex is the development of the nervous system. Thus, it may be laid down that man is, in general, more sensitive to cerebral manifestations of intoxication than are the lower animals. A very large number of substances which in man set up extreme cerebral disturbances, acute delirium, and convulsions have little or no effect on the lower animals, unless given in large quantities.

Admitting this, we may cautiously divide the poisons acting on the nervous system in the first place into:

1. Those causing immediate arrest of activity, e. g., hydrocyanic acid.

2. Those causing immediate diminution of activity, e. g., the hypnotics and sedatives.

 Those causing primary increased activity, followed by diminution of function, e. g., the intoxicant alcohol and the aldehydes, atropine and other alkaloids.

4. Those causing primary increased activity, followed by exhaustion and, it may be, disintegration—strychnine (also tetanus and rabies toxins).

In the second we can classify according to the part of the nervous system upon which the poison exercises its selective action, thus:

1. Higher cerebral centres: The hypnotics as a group, carbonic acid, santonin (in the frog).

2. The bulb: Picrotoxin (a convulsivant, the convulsions ceasing when the medulla is destroyed). Apomorphine (the centres which determine vomiting appear to be situated in the medulla).

3. Spinal cord: Here the type poison is strychnine, the higher centres and the cranial nerves not being affected by this poison. Brucin, quinine, thebaine, salts of potassium and ammonium to a less extent.

4. Peripheral nerves: Ether, chloroform, CO₂ (according to Waller's observations these induce changes in electromobility). Diphtheria toxin (which, by Sidney Martin and others, has been shown to cause active local destruction of axis cylinders or axones). Possibly, lead and alcohol act in a similar way.

 Nerve terminations: Curare (end-plates of motor nerve), cocaine, etc. (sensory nerve-endings). Veratrine, nicotine (stimulate nerve-endings).

Meyer and Overton have shown that the hypnotics as a group undergo an almost selective solution in fats and lipoid substances, and the indi-

ACTING ON THE MUSCULAR SYSTEM

cations are strong that the abundance of lipoids in nervous tissue explain why these are taken up by it. As Ehrlich and later writers have pointed out, adsorption rather than chemical combination proper is the process mainly involved. Nevertheless, whether by disturbing cell metabolism or by some slight chemical action, undoubtedly there is a direct effect on the nerve cells. As demonstrated by Hamilton Wright, working in my laboratory, where large doses of bromides are given, there is distinct histological alteration in the appearance of the nerve cells.

Of these, the most marked is the disappearance of the lateral gemmules of the cells, and he and other observers would suggest that the retraction and degeneration of these gemmules afford the anatomical explanation of the hypnotic action of the bromides and allied substances. If the communication and passage of stimuli from neighboring cells and processes is through the intermediation of these gemmules, and by their retraction the neuron becomes isolated, we obtain the explanation of the action of not only hypnotics, but of another set of drugs, like alcohol, which produce intoxication in the narrow sense, and may regard the incoördination in the action of these nerve cells, which is the essential characteristic of the condition, as being produced by the varying effects of the drugs upon different cell groups, leading at one period to excessive or irregular extension of the gemmules, at another to paralysis or retraction of the same. It cannot, however, be said that this function of the gemmules has been surely established, while disappearance of the gemmules has been recorded in many toxic conditions. To make it safe to accept this hypothesis, with some reservations, the alcohols and the aldehydes must be regarded as primarily exciting the nerve cells, the narcotics as primarily paralyzing the same. Among other primary excitants must be mentioned a series of alkaloids-atropine, hyocyanin, caffeine, and, to some extent, nicotine. All these are liable, under certain conditions, to produce delirium, followed by diminished excitability and coma. As pointed out by Langley, nicotine has the almost specific effect of stimulating, and finally paralyzing, the preganglionic nerve-endings of the sympathetic system.

On the Muscular System.—Substances especially affecting striated muscles act first and with most marked effect upon the heart, so that the circulatory overshadow all other effects, and in general very little has been observed in connection with these muscle poisons throwing light upon symptomatology. The most obvious muscular disturbances tonic and fibrillary contractions and paralyses—are, with rare exceptions, produced by poisons which act primarily on the nerve centres. There are, nevertheless, drugs acting directly on the muscle plates:

1. Irritative.—Causing increased contractility: <u>Veratrine</u> (in small doses), quinine, caffeine, <u>hypoxanthin</u>, <u>creatin</u>, <u>B. coli toxins</u> (Roger, leading to prolonged partial contracture).

2. Inhibitive.—Leading to enfeebled contraction: <u>Metallic salts</u>, of potassium, alkaline earths, copper, etc.

Still less can be said regarding non-striated muscles. There is a certain amount of evidence that ergot leads to contracture. Conversely,

20

he

ch

n,

)y

in

·t

EXOGENOUS INTOXICATIONS: NON-PARASITIC

306

norphine would appear to act directly on the intestinal muscles, and so arrest peristalsis; similarly, <u>atropine has been shown to relax the lower</u> end of the cat's coophagus (provided with plain muscle), while it has no effect upon the upper half (provided with striated muscle). How far the paralytic action of bacterial toxins is responsible for the intestinal dilatation of peritonitis is not absolutely decided. The direct action of certain drugs in producing vasodilatation and vasoconstriction, respectively, must be due to their influence upon the unstriated muscle of the arterioles. The recent observations of Josuć, among others, upon the experimental production of arteriosclerosis, <u>show that one group of</u> <u>substances—adrenalin, barium chloride, nicotine, etc.—acts directly upon</u> the muscular middle coats of arteries.

On the Blood.—There are very many substances which materially modify the condition of the blood, but of these the majority possess little action unless they be directly injected into the vessels. It would seem that in the act of being absorbed from one or other surface these become modified. Further, it has to be remembered (as preventing blood changes) that even when deleterious substances gain entrance into the blood there is a constant removal of the same by the liver and the tissues. Granted this, we recognize certain groups of poisons which act upon the blood in one or other way:

1. Hemolytics (or, better, Hemoclastics) .- Bodies acting upon the erythrocytes and leading to their dissolution, with liberation of the contained hemoglobin. This destruction may be brought about by physical means, as by altering the tonicity of the plasma (intravascular injections of H₂O, etc.) or by freezing and thawing, or by drugs, leading to disintegration of the red corpuscles, e. g., saponin (1 part in 125,000 of this in the blood sets up hemolysis), abrin, ricin, amanita extract, etc. One group of agents, e. g., bile salts and soaps, oleic and other acids, would seem to act by replacing the hemoglobin from its (loose) construction with the proteins of the stroma. The liability of certain bacterial products, of animal venoms, etc., to induce hemolysis, will be discussed in the section on Immunity. In the slightest cases such hemolysis produces hemoglobinemia (and hemoglobinuria); in more severe, methemoglobinemia, the liberated hemoglobin becoming acted upon. Methemoglobin would seem to be developed both in the presence of certain oxidizing agents (the chlorates, sodium hypochlorite, nitrites, etc.), and of reducing agents (pyrogallic acid, pyrocatechin, toluylenediamin, hydroquinone. Yet another group of bodies combine to form more stable compounds with hemoglobin without of necessity setting up hemolysis-carbon monoxide, sulphuretted, seleniuretted, and telluretted hydrogen, prussic acid, cyanogen, and the sulphocyanates, and, according to Liebreich, acetylene. All these combinations prevent the due inhalation of O and CO_a, and, as a consequence, tend to lead to asphyxia.¹

2. Leukolysis.—Two distinct processes may lead to leukopenia or reduction in the number of leukocytes in the circulating blood, nor is it

¹ The reader will find a useful review of data bearing upon hemolysis and leukolysis in Wells' "Chemical Pathology," p. 190.

z

ACTING ON THE ORGANS OF CIRCULATION

always easy to determine which is in action in a given case. These are (a) leukolysis proper, or destruction and dissolution of the leukocytes. and (b) local accumulation, the white corpuscles either accumulating within the vessels of internal organs, the pulmonary or abdominal capillaries, or undergoing actual migration. Such accumulation is apt to involve more particularly one or other form of leukocyte; thus, as pointed out by Opie, in peritoneal inflammation there is a remarkable accumulation of eosinophile leukocytes in the omental and mesenteric capillaries, with reduction in the number present in the general blood. But the indications are that, pathologically, acute leukolysis frequently occurs. Albertoni finds that pancreatin brings about a rapid and complete destruction of leukocytes; several of the bacterial toxins introduced into the blood appear to have a like effect. Bile salts present in excess possess this property. In many of these cases the leukopenia is followed by a pronounced leukocytosis, the increase in lymphocytes being apt to be especially marked. We deal here not so much with a stimulative multiplication of the leukocytes (this though may show itself later) as with a characteristic attraction of the cells in the lymph glands and bone marrow out of the tissue spaces into the capillaries.

The Organs of Circulation.—Poisons may affect primarily (1) the heart, (2) the vessels, more particularly the arteries, or (3) the nerve centres controlling the circulation, whereby the heart and vessels are secondarily affected. The effects of one or other of these actions upon the blood pressure and the circulation in general may be identical; thus, great care is necessary in order to come to a sure conclusion regarding the mode of action of any one poison. After the general circulatory effects have been determined, it is necessary to study the effects upon the isolated heart, or, still better, of transfusion through a vessel of the apical region of the left ventricle, after Townsend Porter's method, that region being devoid of nerve cells. Similarly, following the oncometric method of the late Professor Roy, the effects upon the blood flow must be studied through the isolated peripheral organs—kidneys, spleen, etc. In this way, per exclusionem, we can find the mechanism primarily influenced.

1. Acting Directly on the Heart.—Here we can redivide the poisons into (a) those causing stoppage in systole, and (b) those arresting the organ in diastole. To the former group belong the glucosides and allied bodies, digitalin, digitalein, digitoxin, strophanthin, convallamarin, antiarin, etc. Some of the animal venoms, such as that of the skin of the toad, have a like action, as have also the salts of barium. Causing diastolic stoppage are the metallic salts—of copper, arsenic, antimony, potash_ste. Alcohol (Roy and Adami) and chloroform (MeWilliam) in sufficiently large doses also act directly on the ventricular muscles, causing weakening and ventricular dilatation. Here we have the explanation of the acute dilatation of the heart seen in heavy drinkers (Steell and others).¹

¹ As pointed out by Ringer and Ford, in smaller doses nearly all these substances stimulate the heart and strengthen its beat; once again it is a matter of degree.

308 EXOGENOUS INTOXICATIONS: NON-PARASITIC

Other substances, like muscarin, act through the nervous system almost entirely; most of these nervous "diastolic" poisons act by paralyzing the augmentor or accelerator mechanism rather than by exciting the inhibitory vagus centre.

2. Action on the Vessels.—Here we gain similar groups of poisons: (a) causing contraction, (b) causing dilatation, and (c) acting secondarily through the nervous system.

1. After much contradictory evidence, it is now generally accepted that ergot acts directly on the vessels, for if dilatation be brought about by vasodilator stimulation, the passage of ergot or ergotin through the vessels of the part leads to constriction. At the same time there is direct action upon the heart. <u>Adrenalin and barium chloride</u> have, of late, been found to have even more profound local constricting effect.

 Nitrite of amyl and the nitrites in general, chloral hydrates, quinine, and atropine in small doses, lead to increased rapidity of flow through removed organs; hence, they directly induce dilatation; the same has been noted with regard to acids as a class (Gaskell).

It is interesting to note that these various drugs do not cause constriction or dilatation, respectively, of the vessels of all organs. They have, to a certain extent, a selective action. Quinine acts more especially upon the spleen, digitalein upon the vessels of the kidneys, amyl nitrite upon the vessels of the face and respiratory tract. Adrenin, as Herter shows, while it causes intense contraction and blanching of the vessels of most organs, when applied to the surface of the pancreas causes profound vasodilatation.

It may be noted that some at least of the bacterial toxins have a direct effect on the circulatory system. Thus, Bouchard found <u>tuberculin</u> to <u>act as a vasodilator</u>, Roger that those of the B. septicus putidus had a most powerful action on the heart, with slowing and prolongation of the contraction, and death in diastole. Kemp and Dewey, on the other hand, employing typhoid toxins on the terrapin's heart, gained no slowing but diminution in the size of the beats, and eventual death in systole.

The Digestive System.—Here again we have to distinguish the action upon the nervous, muscular, and secretory mechanisms of digestion, both direct and reflex. The full study of any individual poison, to determine how it affects the digestive system, demands (1) observations upon the results when introduced into the digestive channel: (a) when the nerves to a part (vagi and sympathetics) are intact, and (b) when one or other set is divided; and (2) study of the effect when it is introduced into the circulation simultaneously. Apomorphine, for example, has a direct effect upon the central nervous system, causing emesis when injected subcutaneously; ipecacuanha usually has no such effect; it causing vomiting only when introduced into the stomach, then acting reflexly by stimulating the vagus terminations. Cut the vagi, and even large doses are without effect. Magnesium sulphate and saline purgatives, as a class, at most cause increased peristalsis to a slight degree when introduced subcutaneously or into the blood; to produce abundant watery evacuations, they must act from within the gut. With anti-

ACTING ON THE DIGESTIVE TRACT

mony tartrate, the action may be both direct and reflex. And when definite anatomical lesions show themselves along the course of the alimentary canal, we cannot immediately conclude that these are due to the immediate action of the poison upon the tissues from without; even in the stomach they may be due not to absorption but to elimination of the already absorbed poison. Here, again, true conclusions can only be reached by comparing the effects of ingestion and of subcutaneous or intravenous infection. It is not surprising, therefore, that, contrary statements exist regarding the mode of action of many of the digestive poisons.

This being the case, it is best to pass in review the main orders of digestive disturbances in relation to their causes:

Salivation.—Drugs set up salivation and arrest of salivary secretions mainly by reflex nervous mechanism; the poisons must be absorbed before they tell upon the salivary gland. Of such reflex salivation, that set up by emetics affords a good example. To some extent the process may be regarded as eliminative, e, g, in mercurial salivation,

Vomiting.—While, as seen by study of the isolated stomach, several poisons can act directly upon the gastric musculature, setting up irregular peristaltic movements, contraction or relaxation, and paralysis, the process of vomiting is not due to the stomach alone. As Magendie showed, replace the stomach by a simple bladder, and vomiting still may be set up. Obviously, the stomach and alimentary tract in general play a secondary role in the process; to coördinate all the factors involved in the act, the nervous system must dominate, and, as Sir Lauder Brunton has shown, vomiting may be initiated in two ways: (1) Reflex, by gastric irritation of the branches of the vagus; (2) direct, by excitation of the nerve centres. We have already afforded examples, in ipecacuanha and apomorphine, of these two modes of action.

Diarrhœa and Dysentery.—'The causes of diarrhœa and the modes in which they act are manifold. Broadly speaking, under this heading discharges of two different types are included: (1) The premature removal of the liquid contents of the small intestine without due absorption and modification, and (2) the discharge of excessive secretion from the mucosa of the small and it may be of the large intestine. Dysentery, which, it may be recalled, is properly not a specific disease, is that form of diarrhœa characterized by straining and irritation of the lower bowel, accompanied by mucus, and, it may be, blood, derived from the inflamed mucosa of the colon and rectum.

The first process, that of premature removal, is brought about by increased *peristalsis*. This may be due to direct action on the nerve centres or to reflex irritation. Injections of rhubarb or senna will cause purgation when injected into the veins; croton oil only when introduced into the alimentary canal, and then, if the vagi be cut, no diarrhoea results. In other cases the action is even more indirect. Aloes, for example, acts only when injected, and then only when there is a free flow of bile; ligate the common bile duct, and no diarrhoea ensues.

As regards the second process, that of increased secretion, the saline

EXOGENOUS INTOXICATIONS: NON-PARASITIC

purgatives, as already noted, mainly act by this means, not causing purgation when injected intravenously.

When there are actual lesions—acute congestion, with or without ulceration—there is undoubtedly increased discharge, accompanied by diminished absorption. Regarding these lesions, it must be remembered that they are of two orders: (1) Those produced by direct irritative action of poisons acting upon the exposed surfaces, and (2) those due to elimination of an absorbed poison.

1. The former, naturally, are most apt to be seen in the upper part of the digestive tract. Bodies of the nature of acids or caustics setting up direct necrosis of the mucosa cause the greatest injury. This, however, often tends to be localized rather than generalized; it is regions of narrowing and compression—of arrest of the irritant—that are most liable to exhibit disturbance, as, for example, in the œsophagus, opposite to the larynx, the bifurcation of the trachea, the cardiac end of the œsophagus.

2. The eliminative lesions may occur from the stomach downward. Thus arsenious acid introduced subcutaneously will produce multiple hemorrhages and fatty degeneration of the mucous coats of the stomach, which pass on, as shown by Filehne, to multiple peptic ulcers, provided the gastric contents be acid. Whether the curious duodenal ulcers occasionally met with in extensive burns are eliminative is still an open question. The acute hemorrhagic, necrotic, and ulcerative condition of the colon seen after swallowing corrosive sublimate may be reproduced when the poison is introduced by other paths. We have similar evidence that mercury and antimony are eliminated through the colon. By analogy, the ulcerative origin.

The Liver.—The usual function of the liver is to neutralize or eliminate poisonous substances brought to it in the circulation; thus, on the one hand, its cells can excrete or modify relatively large amounts of toxic matter without being greatly affected; on the other, this very function, coupled with its position at the head of the portal circulation, renders them peculiarly liable to be damaged, and to exhibit either acute or chronic disturbances—cloudy, fatty, and other degenerations, with nuclear changes or even acute necrosis; or, again, atrophic changes coupled with fibrosis.

The rapidity with which many substances are absorbed from the intestinal canal and taken up by the liver cells is remarkable. Laffter, working under Heidenhain, found that rhubarb injected into the duodenum appeared in the bile in less than five minutes. Sulphindigotate of sodium introduced into the circulation began to enter the bile one minute after its injection.

In general, direct, as distinguished from reflex, toxic disturbances of the organ are more prominently in evidence, but the development of jaundice, as the result of strong emotion, in itself indicates the modifying influence of the central nervous system upon the organ.

Certain classes of poison stand out especially as being excreted or

acted upon by the liver, and as liable to set up disturbances in the process, notably:

 Among metals and metallic salts, lead, copper, mercury, arsenic, phosphorus. The last more particularly modifies profoundly the cellular actions, leading to nuclear and cytoplasmic disturbances. These metals, upon analysis, are found in greater quantities in this organ than in any other part of the body; they are excreted into the bile, a vicious circle being set up (*i. e.*, they may be reabsorbed again from the intestines).

2. The toxic products of digestion, indol, skatol, toxic albumoses, with which may be included the toxic products of bacterial growth in the intestines. Where these are found in excess, we gain evidence of *hepatic incompetence;* the overloaded cells permit the toxic bodies to pass through the organ into the general circulation, and, more particularly with non-neutralized digestive products, there are induced torpidity, slowness of pulse, dilatation of cutaneous vessels, muscular weakness—conditions which can be reproduced experimentally by injecting certain products of proteid and amylaceous digestion into the systemic circulation (albumoses, lactic acid, indol, etc.).

3. The toxins of pathogenic bacteria: One of the commonest changes in acute infections is a <u>state of cloudy swelling of the liver</u>, passing on in the more acute states to one of <u>fatty degeneration</u>. Certain toxins have an even severer action upon the liver cells, setting up localized areas of cell death (<u>focal necroses</u>). These we see in typhoid, diphtheria, the <u>plague</u>, and a number of other acute infections. There continues to be doubt as to the exact mode of causation of these necroses, but some at least are held to be due to the direct action of toxins.

4. The products of hemolysis: These and their effects will be discussed when we take up the subject of jaundice.

The Pancreas and Spleen, and Ductless Glands in General.—Our knowledge of the direct action of poisons upon these organs is not sufficiently extensive to permit of any general statements.

The Kidneys.—Just as the liver is the great organ for the elimination of toxic substances from the portal circulation, so the kidneys are the most prominent eliminating organs for toxic substances in the systemic circulation, and bear the brunt in cases of systemic intoxications. Thus the considerations laid down in connection with the former organ apply here very largely, mutatis mutandis. There may be nervous or direct irritation of the organs, leading to increased or decreased discharge of urine or of specific constituents of the same, cloudy and fatty degeneration of the tubular epithelium of particular areas, or disintegration and necrosis of the same; or, where the process of elimination is carried on over a long period, the gradual development of fibroid changes. Certain toxic substances, e. g., the metals above mentioned, act both on the hepatic and renal parenchyma; others, like cantharidin, act more particularly on the renal cells.

The Skin and its Constituent Glands.—Physiologically there is an interesting relationship between the sudoriparous and the salivary glands. Those substances which, like emetics, set up an increased

12 EXOGENOUS INTOXICATIONS: NON-PARASITIC

salivary flow, in general lead at the same time to diaphoresis. This is well marked in the case of <u>pilocarpine</u>. Other substances, like <u>atropine</u>, <u>arrest both salivation and diaphoresis</u>. Whether the action of diaphoretices is directly upon the gland cells, or through the terminal nerve filaments, is still a matter of debate.

Numerous disturbances, eliminative, vasomotor, trophic, may be set up in the skin by very numerous poisons; we are still far from understanding the causation of many of these conditions. One of the commonest is extensive localized or general erythema (active congestion of the superficial vessels). Mercury, the bromides, the iodides, iodoform, salicylic acid, etc., are on record as producing the condition. Closely allied to these are the erythemas produced by irregularities in diet or by eating certain foods. To a very large extent these erythemas and urticarias are idiosyncratic—only certain individuals are liable to manifest the lesions. Either some nervous disturbance must be at the bottom of these idiosyncrasies, or, as it is now suggested, some minute variation in the constitution of the cytoplasm of the endothelium of the vessels of a part, rendering it peculiarly susceptible to the actions of the toxins or poisons.

In the case of purpuric eruptions, it would seem that we have clearly to deal with this direct action of the poisons upon the endothelium of the smaller cutaneous vessels, for this often exhibits fatty degeneration. It is the localized weakening and necrosis of these cells that would seem to precede the rupture and hemorrhage, per rhexin and diapedesin, which set up the purpuric ecchymoses. This, it is true, is not the only cause of purpura, but would seem to be the commonest. We have encountered minute capillary emboli leading to minute hemorrhagic infarcts. In some cases of bacterial origin, as in the rose spots of typhoid, it is now fully demonstrated that there is local growth of bacteria, leading through the toxins to local degeneration and necrosis of the containing vessels, congestion, and, it may be, rupture of the same.

CHAPTER VIII.

EXOGENOUS INTOXICATIONS: PARASITIC CAUSES.

Or greater frequency and greater importance are the parasitic intoxications. Some of these, it is true, are due to absorption from without; there are parasites—using the term broadly—which live in the alimentary canal, for example, and do not themselves penetrate into the tissues; and living there, produce toxins which are absorbed. These are the exception. All that is necessary is to note that such exist, and that their products act after the same manner as those of the main mass of parasitic forms growing in the organism. All fall into one of the three groups: (1) Microparasites of vegetable nature; (2) microparasites of animal nature; (3) the larger animal parasites. In discussing how they act in setting up disease, we shall pass them in review in the above order.

BACTERIA AS CAUSES OF DISEASE.

Of all the various pathogenic microorganisms, the bacteria stand out as the largest class, and it has been by a study of their properties that the modern doctrine of infection has been developed. The realization of their existence and growth within the organism has brought about the greatest revolution in medicine and in surgery that our science has experienced. As already noted, we shall not deal with them in detail; that work is accomplished in the many special works upon bacteriology, wherein bacteria are dealt with from the point of view of the medical man, works in which the manner in which the individual pathogenic bacteria cause the different diseases is carefully discussed. We would but recall, and that rapidly, the characters of bacteria as a class, and deal generally with the methods whereby they may cause disease.

Briefly, then, bacteria are, as a class, characterized by their extreme minuteness; some, indeed, like the organism of contagious pleuropneumonia in cattle, are beyond the power of the strongest microscope to render visible—are less than one ten-thousandth of a millimeter in diameter; they possess no distinct nucleus; do not (so far as we can determine) conjugate, but multiply purely by fission; are, some of them, motile, by means of flagella, and some, again, but not all, exhibit sporulation, whether by internal development of such spores (endospores) or by conversion of the whole of the minute organism into an encysted resting stage, as occurs in some of the spherical forms, though whether

these so-called *arthrospores* are spores proper is gravely doubted. Morphologically, they can be separated into the three broad divisions of cocci, spherical or blunt oval forms, bacilli or rod-like forms, and spirilla, or forms exhibiting either complete spirals or segments of the same; transitional forms occur which it is difficult to range in either one or other division.

The temperature limits between which they grow are very various, as are also the media in which they grow, but of pathogenic bacteria as a class it may be said that they grow best at, or about, the temperature of the body of their hosts, and best also upon media containing organic matter, which, like the blood and tissues of their hosts, have a slightly alkaline reaction to litmus. The majority grow best in the presence of free oxygen (aërobes). A large number can exist in the complete, or almost complete, absence of free oxygen (facultative anaërobes); a small number (of pathogenic forms) can only grow when free oxygen is practically absent (obligatory anaërobes), obtaining the oxygen that is essential for existence by the breaking down and reduction of oxygencontaining foodstuffs. Possessing no mouths or organs, they can only live by absorption; in other words, in a fluid medium holding the necessary foodstuffs in solution. That they may act on potential foodstuffs. and bring about their dissociation, converting them into a soluble and nutritive form, they secrete enzymes-just as do the cells of the digestive tract of higher animals. These enzymes in the different species are of different orders-proteolytic, cellulose fermenting, diastatic and glycolytic, dissociating the various sugars, etc. Added to this, the pathogenic bacteria produce toxins, or substances having a poisonous action upon other living organisms.

With reference to their toxic powers, we may divide bacteria into three groups:

1. The non-toxic.

2. Those incapable of multiplying within the tissues, but grown outside the body, capable of producing toxic substances, which, being absorbed, set up disturbances. To this class belong many of the saprophytic and putrefactive bacteria, among them sundry organisms of so-called wound infection—microbes which will grow in pus of surface wounds, and there, through their products, set up irritation without gaining entry into the tissues. Here, also, are to be included many of the microorganisms of the digestive canal, which, growing in excess, set up local and general indications of intoxication. Many of these, it would seem, can under conditions become converted into members of the next group.

* 3. Bacteria capable of growing within the tissues and there setting up infection.

It is thus by their products of growth that bacteria cause disease, the difference between a bacterial intoxication pure and simple and an infection being that in the former case the products alone are absorbed; in the latter, the bacteria themselves gain entry and grow.

Toxins.—'The term "toxin" is now so generally and so vaguely used to embrace all the deleterious products of bacterial growth, that, doubtless, substances of very different orders are included under the term, including the direct excreta of the bacteria and the secondary products of the action of these discharges upon the medium of growth. That we have to deal, at least, with these two groups of bodies is, it seems to us, conclusively indicated by Sidney Martin's studies upon diphtheria. Martin has shown that from the spleen and other organs of those affected by this disease it is possible to isolate a highly toxic albumose, whereas from the false membrane in the throat (in which alone the bacilli have undergone multiplication, and are present in abundance) but little of the albumose is to be obtained. Nevertheless, an extract (sterile) of the false membrane has singularly toxic properties. From a study of the effects of this extract he concluded, with Roux and Yersin, that the primary product discharged by the diphtheria bacillus is an enzyme, that this diffusing, with some difficulty, it may be, and in very minute quantities from the region of growth of the bacilli, does not itself poison the tissues, but, acting upon certain proteid substances of the organism, converts them into highly toxic albumoses-and these it is that set up the symptoms of disease.

And there is much to be said in favor of this view of the nature of at least an important group of the primary products of bacterial growth. Like enzymes, extraordinarily small quantities suffice to produce eventually maximal disturbances; action is not immediate and is cumulative, herein differing from anything of the nature of a direct chemical process. They are brought down by the same substances which precipitate enzymes, are characteristically thermolabile, rendered inert by temperatures of 56° to 60° C. (the exceptions being no more marked than in the case of enzymes), and they diffuse either not at all or very slowly).

Not all pathogenic bacteria produce noticeable amounts of toxins discharged in the process of active metabolism. Some, like the typhoid bacillus and the Bacillus coli, afford culture fluids of very low toxic powers, but if they be frozen and triturated, as by Rowland's method, or subjected to great pressure, the body juices are found to be intensely toxic. The observations here are exactly parallel to the well-known experiment of Buchner. The fluid in which the yeast plant has been grown contains a ferment which will invert sugar, but will not convert that into alcohol. Express the active yeast cells under an hydraulic press, and the body juices, acting on the inverted sugar (glucose), produce alcohol. Here we have to deal with the existence of intracellular enzymes, bound up with the living cell substance, and the stimulus exerted by Buchner's observation has led to the discovery of abundant intracellular enzymes in the tissue cells. These "intracellular toxins" seem, obviously, to be bodies of the same order.

The views here put forth of the existence of primary and secondary toxins—the primary of the nature of enzymes, the secondary the active toxic substances—are, we must point out, not as yet universally accepted.

In this connection reference must be made to the remarkable studies of

Vaughan,¹ of Ann Arbor, and his pupils, upon the composition of the bacterial body in relationship to its pathogenicity. Making huge growths of various bacteria on solid media several square yards in extent, so as to obtain a sufficiency of material for analysis, Vaughan treats the extracted protein or proteins by digestion with an excess of a 2 per cent. solution of caustic soda in absolute alcohol. From these—as from egg albumin—he obtains a toxic moiety, soluble in alcohol, and a non-toxic, insoluble moiety. The toxic constituent is found constantly to possess a benzole ring, it affords tyrosin, and has the general properties of an albumose: to this ring-like constitution with side-chains to be satisfied he ascribes its toxic activities more particularly upon the cells of the respiratory centre.

THE NORMAL DEFENCES OF THE ORGANISM.

Recognizing, then, that we have these two ways in which bacteria can cause disturbances within the organism, namely, by the absorption of their products of growth and the substances produced thereby (simple intoxication), and by growth and production of toxins within the system (infection), it must next be asked, How do the bacteria gain entry into the tissues and thus cause the latter state?

The human body and, for the matter of that, the bodies of all multicellular organisms are to be regarded as closed corporations, in which corporations one of the special functions of the outer layer of cellular units is, for the benefit of the whole system, to hinder the entrance of individual organisms of other natures. And here it must be kept in mind that, contrary to first conceptions, these outer layers are not the external layers only, in the usual acceptation of the term, but are all layers bounding surfaces and channels which, however indirectly, communicate with the exterior. The mucous membrane of the stomach and intestines is thus strictly external, as will be grasped from the diagran given on p. 301.

There is in the higher animals only one direct channel of communication between the interior of the body and the exterior, and that only in the female, namely, the Fallopian tube, which has so fine a channel, and so protected, that to all intents and purposes it is closed; only under very exceptional circumstances—we have encountered one such case can acute peritonitis be brought about by suction of infective material from the uterus through the tube. The case in question was that of a thoroughly healthy girl, who, in the last days of her menstrual period, took part in a gymnastic competition, dying within eighteen hours. The viscera were found absolutely free from anything that could suggest

¹ Trans. Assoc. Amer. Phys., 20:1905:265; see also the Shattuck Lecture for 1906, Boston Medical and Surgical Journal, 1906; ii:215, 243, and 271. From the latter it will be seen that Professor Vaughan's conceptions of vital processes are of the same order as those set forth in this volume.

THE NORMAL DEFENCES OF THE ORGANISM

a primary focus; the peritoneal exudate contained a pure culture of streptococcus, and similar streptococci were present in the uterine cavity. It must be recalled that, save at the menstrual period and after parturition, the mucus of the cervical canal acts as an efficient plug against suction of vaginal or uterine contents through the Fallopian tubes. There are on record other cases of apparently primary peritonitis in which, *per exclusionem*, this mode of infection must be invoked.

Living outside this close corporation are countless other individual organisms. On the very surface of the human body, for instance, we know that there exist millions of microbes, mainly bacteria, many of them potentially pathogenic—pyococci, streptococci, B. pyocyaneus, B. coli, etc. The mouth contains them in abundance—pyococci, streptococci,¹ pneumococci. There may be countless millions in the intestinal canal, but these are outside the body, and, while they find nourishment in the cast-off dead cellular debris, in certain discharges from the surfaces and in the food material ingested, they are not taken into the tissues, or, as we shall point out, if they gain entrance, there are many mechanisms for arresting their growth and destroying them.

Of those mechanisms we recognize the following:

Surface discharges: Certain discharges or excretions either simply wash off the microbes, which, left at rest, might multiply, or, in addition, have definite bactericidal properties. But for the flow of the saliva, the mouth would be an admirable incubator. That flow carries them, with swallowing, to the stomach, where, if not already destroyed by other means, the acid gastric juice kills off the greater number. The mucus poured out from the various glands of the mouth and respiratory and other tracts, while it favors the arrest of bacteria, forms a layer through which microbes grow with difficulty to come into direct contact with the surface cells, and, in general, before this contact can be accomplished, either by the action of the cilia of those cells (respiratory tract, etc.), or by peristaltic or other movement, the mucus is liable to be carried away. Certain observers hold that mucus has, in addition, definite bactericidal properties.

The gastric juice is particularly active in destroying bacteria. The number taken at each meal with the food must be great. Milk, for instance, that has been kept—as most milk is kept—for twenty-four hours and more, may, on warm days, contain, it may be, 2,000,000 or more bacteria per cubic centimeter (15 minims). Nevertheless, a few hours after food the duodenum may be found quite sterile, and in general, as indicated by the slower development of the peritonitis following perforation of the stomach or upper portion of the intestine, as compared with that succeeding rupture of the lower portion of the small

¹ During our bacteriological course in 1905 the students, making cultures from each other's mouths, gained streptococci in 80 per cent. of the cultures. In view of the full studies of the last few years on the different strains of streptococci, more particularly by the English bacteriologists (Gordon, Houston, Dudgeon), it would be interesting to determine what proportion of these buccal streptococci belong to pathogenic and harmless races or species, respectively.

intestine, the bacteria which persist in the stomach are both reduced in number and, at least temporarily, inhibited in their activity. When there is gastritis, with arrest of secretion, or diminution of the hydrochloric acid, the same is no longer true. Then not only are the bacteria not destroyed, but, escaping into the small intestines, they find alkaline contents of the same or favorable medium of growth, and, proliferating, may by their products induce extensive irritation. Following Pettenkofer, Hankin found, for instance, that in normal health he could with impunity swallow billions of living virulent cholera spirilla without ill effects. Happening to repeat the experiment when he was suffering from transient gastric catarrh, there developed an acute diarrhoca, with abundant spirilla in the stools. It is interesting to note that what we may term are the commonest normal inhabitants of the intestines, namely, the B. coli and Bact. acidi lactici groups, are forms relatively tolerant of acids.

Even the excreta—the urine, milk, and, to a slight extent, the bile¹ have been demonstrated to exert a certain inhibitory and, in some cases, mild bactericidal action.

There are yet other protective external mechanisms. If a current of impure air, bearing dust, spores, etc., be caused to impinge upon a moistened surface, as it impinges it leaves behind it solid particles. The back of the pharynx, and, as an important auxiliary, the turbinated bones of the nose, act as such moistened plates, arresting the bacteria taken in with the inspired air. So effective is the mechanism that (in nose breathers) few bacteria gain entrance into the trachea. The further action of the moistened surfaces of the trachea and bronchi results in this, that the expired air of a healthy man is found absolutely sterile. As Arthur Ransome has shown, the same is true during quiet breathing, even where there is active tuberculous lung disease (although, as Flügge has demonstrated, the fine particles or globules of moisture discharged from the mouth of such a patient in speaking, coughing, sneezing, may contain the bacilli, and be infective). These globules, it is scarce necessary to say, are formed of saliva.

On such a surface as the back of the pharynx the mere presence of a frequently renovated layer of moisture would not seem to be a sufficient guard against lodgement of microbes and subsequent proliferation of the same. Ruffer² has called attention to the existence of a further mechanism. Even low down in the animal kingdom, as pointed out by Gaskell and Miss Allcock, Hardy, and others, we find that the surface discharges and mucinous coverings are not sufficient to arrest the growth of bacteria, and that certain leukocytes, passing out on to the surface, act as scavengers, and these, whether by actually taking up the microbes and digesting them, or, by their "explosion" and discharge of bactericidal contents, succeed in cleansing the surface and

¹ We have found that the bile, while not actively bactericidal, has a distinct inhibitory effect upon the growth of forms like the B. coli.

² British Medical Journal, 1890; ii: 491.

removing bacteria and low forms of life, which, growing in too large quantities, might poison the outer cells, and so break down the line of defence.

If we examine a scraping or swab from the surface of the pharynx or nose, we find that this contains fairly abundant leukocytes. In other words, leukocytes are constantly passing out between the epithelial cells to gain the surface. Many of these, properly stained, show within them bacteria and their remains. They are acting as scavengers and cleansers. It seems probable that the majority, having performed their functions, undergo dissolution or are swept away by currents of But some, at least, find their way between the lining cells into saliva. the subjacent tissues. This can be well determined if one takes, as did Ruffer, the rabbit's tonsil, kill the animal, remove and immediately harden and prepare the tonsil-staining sections for microorganisms. The tonsils are essentially lymph nodules, lying immediately beneath the surface epithelium. Such sections show, besides the lymphocytes forming the lymph nodes, two other orders of cells, the one polynuclear¹ --some of which contain bacteria-the other, larger cells of endothelial type and origin (Metchnikoff's macrophages), of which some contain polynuclear leukocytes and their remains. Such leukocytes, in short, as have wandered back from the surface find their way into the lymph channels, and so to the lymph nodes, and reaching them, are liable, if weakened, to be taken up by the larger hyaline endothelial cells lining the channels.

It is wrong, therefore, to imagine, as it is too often taught, that the hindrance to the entrance of bacteria into the tissues is, under all circumstances, complete in the healthy individual. A certain number of microbes is always gaining admission—nay, is being actively introduced by the cells of the organism. But under such circumstances they do not cause infection. In health they tend to be destroyed very soon after their reception. The evidence that this is the case is now overwhelming. It is true of the lower respiratory tract. Careful study of the peribronchial glands shows the same presence of intracellular bacteria, although not so abundantly as in the tonsils.

It is true also of the intestines. Here it can best be followed in the lower portion of the small intestines, the region in which the proliferation of bacteria attains its maximum. It is in this region that one notes that the lymph-glandular tissue of the submucosa is the most extensive (Peyer's patches and solitary follicles). Bizzozero' and Ruffer have shown that (in the rabbit) the lymphoid tissue presents appearances identical with those seen in the tonsil. Repeating the work in

¹ Polymorphonuclear is the more correct term, but we admit that it is too sesquipedalian for daily service. We employ polynuclear, with the reservation that it must be understood that by this we do not mean that the cells are multinuclear, only that they possess nuclei of many shapes.

² Centralbl, f. d. med. Wissensch., 23: 491; see also Ribbert, Deutsch. med. Woch., 1885: 197.

our laboratory, Dr. A. G. Nicholls¹ has shown that the amount of taking up of bacteria by these nodules in various animals is very considerable, though the rabbit, with its larger eccum, and arrest there of fermenting food, usually affords the most convincing demonstration.

Take a healthy, well-fed rabbit, kill it, open the abdomen, pull out the lower coils of the small intestine so as to stretch the mesentery; harden the mesentery with formalin in the stretched condition; with a sterile swab clean off the endothelial covering on either side; cut it out without opening the intestine. Now upon staining with carbol thionin there are to be observed along the fine vessels of the mesentery what, from their shape and relationship, can only be the more or less degenerated remains of bacteria. Occasionally a well-formed bacillus is to be recognized. Some are along the lymphatic vessels; some, judging from their nearness to irregularly lobed nuclei, are within polynuclear leukocytes; occasionally they are within small blood capillaries.

It is clear, then, that, taken into leukocytes upon the outer surfaces, bacteria may (1) be arrested in the first line of subcutaneous lymph nodes, or (2) may evade these and be arrested in the second line, the mesenteric and retroperitoneal lymph nodes, or (3) may pass into the minute radicles of the portal vein.

That such leukocytes as have escaped into the lumen of the gut and have returned can and do pass into the blood stream, has been demonstrated with admirable precision by Prof. A. B. Macallum, of Toronto,² In the course of his investigations upon the microchemistry of the cell, studying the fate of iron in the economy, he was led to feed lake lizards (Necturus) with peptonate and albuminate of iron. Taking them when they had fasted for thirty months (to make sure that their intestines were empty!), he fed them with the compounds above mentioned, and killed eight hours later. Employing Perl's (the Prussian blue) test, by which means the free iron, if present, takes on a pronounced blue color, he found that:

 Within the lumen of the intestine were leukocytes full of Prussianblue granules. These had, therefore, passed out and taken up the iron salts.

 Between the epithelial cells of the villi were cells of the same order (reëntering leukocytes), along with others quite free (wandering-out leukocytes).

3. Examining the other tissues, he found leukocytes containing the blue granules in the capillaries of the liver and in the spleen; or, in other words, the leukocytes containing the iron had found their way into both the portal and the systemic blood.

And, lastly, not to dwell upon the convincing results of Nocard, Ravenel, and Behring, upon the passage of tubercle bacilli and other organisms into the lymph of the thoracic duct of dogs and other animals

> ¹ Journal of Medical Research, N. S., 6: 1904; 485. ² Jour. of Physiol., 16: 1894; 268.

THE NORMAL DEFENCES OF THE ORGANISM

when fed more especially with fatty foods,¹ Dr. W. W. Ford,² in our laboratory at the Royal Victoria Hospital, has demonstrated most conclusively that immediately after death organs like the liver and kidneys of dogs, cats, rabbits, and guinea-pigs are not sterile, but contain some few bacteria, which are capable under favorable conditions of still growing.

Ford's cultures proved that over 70 per cent, of the livers and kidneys of these animals, if removed aseptically within a minute or two after death, and placed with every aseptic precaution in agar-agar, gelatin, or broth, yield cultures of various forms of pathogenic and nonpathogenic microbes, such as are found in the intestinal contents. The interesting point is that the growth of these forms is peculiarly slow. Ford, in general, obtained no growth within three days, but, keeping for several days, he obtained positive results. Evidently (1) the bacteria are attenuated, so that their growth is feeble, and (2) it is arrested until the bactericidal substances of the organs have become inert. There was a striking difference between the flora gained from the carnivorous animals and the rodents, and again between the flora of the different series of animals (dogs, cats, rabbits, guinea-pigs).

Wrosczek has made the further observation that, giving healthy animals cultures of non-pathogenic pigmented bacteria with their foods, these can be obtained in culture from the internal organs, without there being a sign of inflammatory or other lesions along the intestinal tract. More recently, our colleague, Dr. Wolbach,³ has called attention to the fact that in a large proportion of cases of what had hitherto been regarded as aseptic autolysis of the liver of the dog, cultures of an anaërobic bacillus are obtainable—so large a proportion that this must be regarded as habitually present in the healthy animal.

These observations, then, prove conclusively that *bacteria are constantly entering the organism*. But now to turn to another aspect of the subject: If bacteria, and these pathogenic, be injected directly into the blood stream, within fifteen minutes, and even within five minutes, although hundreds of thousands, not to say millions, have been thrown in, the circulating blood affords very few colonies. And, examining the tissues after these intervals, one finds that the bacteria have already been actively removed from the blood by the endothelium of the bloodvessels, more especially of the liver, kidneys, and spleen (splenic corpuscles). Within an hour the heart blood may be found sterile.

(At a later period, however, the blood may again be teeming with the bacteria.)

Bacteria, then, which gain entrance are taken up by the endothelial

¹ It is not an exaggeration to say that during the last few years there have appeared close upon fifty articles dealing more particularly with abdominal infection by the B. tuberculosis, and demonstrating experimentally the passage of solid particles through the unaltered mucosa of the intestine. Fuller data regarding the earlier work are given in an address by me. Jour. Am. Med. Assoc., 33:1899:1506 and 1572.

² Trans. Assoc. Amer. Phys., 15: 1900; 389, and Jour. of Hygiene, 1: 1901: 276.

³ Wolbach and Saiki, Jour. of Med. Research, 21: 1909: 267.

21


cells of the bloodvessels. Werigo¹ has observed and figured the endothelial cells in the liver sending out definite pseudopodia, whereby leukocytes and their contained bacteria are arrested. As these processes retract, the bacteria are to be seen lying within the endothelial cells.²

As we have pointed out,³ if one studies a series of livers removed at different periods from rabbits, into whose vessels B, coli have been injected, it is possible to recognize a series of stages of destruction of the bacteria.

At an early stage, large, well-formed bacilli are to be seen within the endothelial cells of the liver. Later, these break up into short stumpy segments; later, as the process continues, in place of bacteria, rows of two or three minute dots are observable; later again, the endothelial cells become free from any signs of germs. We are inclined to conclude



Swallen endothelial cell of capillary of rabbit's liver containing Bacillus cell in various stages of degeneration, within thirty minutes of injection of the bacilli into the blood stream. Part only of the nucleus is shown in the section.



Phagocytic cells from peritoneal cavity of guinea-pig, nine hours after intraperitoneal injection of Bacillus coli, to show stages of destruction of the bacilli: p, polynuclears; m, large mononuclear cell; l, lymphocytes (non-phagecytic).

that minute isolated double and treble dots seen within the liver cells represent the remains of the bacteria taken up from the overlying endothelium, for similar diplococcoid forms now appear in the bile.

Personally, we incline to the opinion, although we will not lay it down positively, that both the liver and the kidney actively excrete bacteria which have undergone preliminary action by the endothelium. Regarding this matter, it is extremely difficult to arrive at an absolute conclusion, and the many observations made since Columbian first pro-

¹ Ann. de l'Inst. Pasteur, 7: 1893: 593.

² A study of the phagocytic action of human leukocytes by Leishman's method shows that ingestion may occur without pseudopodial activity: mere contact, with, apparently, altered surface tension, results in the bacteria flowing into the cytoplasm.

^a Adami, Abbott, and Nicholson, Journal of Experimental Medicine, 4:1899:349.

THE MODES OF INFECTION

pounded the view, have been so conflicting—so positive in either direction —that we hesitate to express more than our belief.

Thus, the observations of Sherrington, Wyssokowicz, and not a few others are wholly opposed to this conclusion; those of Blachstein, Futterer, Wrosczek, our own observations, and those of Nicholls support it.

Whether the process continues so far that there is actual excretion of attenuated and destroyed bacteria, this is obvious, that while the tissues of the healthy body are not of necessity free from microorganisms, they are potentially sterile. While the mesenteric and other superficial lymph nodes may show abundant bacteria, the mass of these are clearly destroyed, or are undergoing destruction. While the removed liver may show abundant bacterial forms, in its endothelium, and even, as we hold, in its parenchyma, scarce any of these are normal, and cultures from the same organs made immediately after death are either sterile or give only delayed growths of a few forms; here and there throughout the organ a microbe not yet acted upon to the extent of rendering continuous growth impossible may yield a culture.

There are thus many means whereby the organism is prepared to arrest the entrance of microorganisms, and, in the event of such entrance, to inhibit their growth: (1) The physical and bactericidal action of the bodily discharges; (2) the structure of the surface layer; (3) the bactericidal activities of the wandering cells; (4) the phagocytic and bactericidal action of the lymphoid tissues; (5) the like action of the vascular endothelium; and (6) it may be, the bactericidal and excretory action of the cells of certain excreting glands. Nor is this all: (7) we have, in addition, ample proof that the circulatory fluids of the body have antibacterial properties. There is, however, still some want of accord between different observers as to how far these are in force under physiological conditions. We shall deal with this subject more fully at a later period.

THE MODES OF INFECTION.

We have purposely dealt with the subject of the normal defences of the organism at some little length, and this because an adequate appreciation of the relationship of the body toward bacteria in health is absolutely necessary for a full grasp of the conditions under which infection may originate. It will be seen that, instead of there being, as is so generally taught, one almost universal method whereby bacteria enter the body, namely, by some (traumatic) solution of continuity of the surface layers, the means are manifold. We may have:

1. Alterations of the surface discharges and secretions, either in amount or in quality, whereby microbes proliferate unduly on the surface, producing sufficiently concentrated toxic matter to affect the surface cells, lower their vitality, and destroy them, with the result that they now gain a focus of growth *within* the tissues.

The foul ulcerous condition of the mouth in certain fevers, accompanied by lessened salivation, is an instance to the point, as, again, is the

development of thrush. The accumulation of excreta by closure of the passages of discharge also favors the development of infection. This has been noted especially in the alimentary tract. Posner and Lewin' have proved that experimental closure of the rabbit's rectum leads to the presence of the B. coli in the various tissues and excreta in the course of a very few hours; Czaplewsky and Frazier,² that experimental closure of the rabbit's cecum rapidly leads to the supervention of peritonitis. Not only do bacteria proliferate excessively under these conditions, but their virulence is definitely exalted. B. coli isolated from the contents of the intestines before such closure may be found harmless for other animals; isolated some forty-eight hours later, is intensely virulent. The main cause of appendicitis is primarily, it would seem, not ulceration or erosion, but kinking or other obstruction of its narrow channel.

 Traumatic solution of continuity of surface layers: Here we have to deal not only with the destruction of the protective layers, but with the provision of a favorable nidus for bacterial growth in the necrosed cell tissue lining the injured surface.

When Pasteur fed a flock of sheep on a meadow which had been sprinkled abundantly with a virulent broth culture of the anthrax bacillus, scarce an animal succumbed to the disease; when, in addition, he scattered thorny particles and broken glass over the meadow, and then turned out the sheep to graze, the majority of the flock died of anthrax. It is, however, unnecessary to quote individual instances; the marvellous change which has come over surgical results since the application of Lister's method of keeping wounded surfaces free from exposure to contamination is our great object lesson. Little wonder that, with this before them, surgical pathologists regard trauma and solution of continuity as the essential causes of infection.

But here let us point out the significance of Welch's observations. It is not necessarily the destruction of surface layers which allows infection; the lowering of the vitality of the tissues is of almost equal importance. Despite the greatest care in the cleansing of cutaneous surfaces, and in the carrying out of aseptic or antiseptic treatment, suppuration may show itself in a wound. Welch has called attention to the almost universal presence in the lowest layers of the skin of the M. epidermidis-closely allied to, if not an attenuated form of, the pyococcus albus-these in health leading a harmless, saprophytic existence. This form is the common cause of, and is to be isolated from, "stitch abscesses," and may lead to extensive tissue destruction and general disturbances. If ligatures be made too tight, the included tissue is largely deprived of blood supply and nutrition, its vitality is lowered, and under these conditions it is that forms so feebly pathogenic as to be incapable, under ordinary conditions, of growth within the tissue, now proliferate, break down the tissues (by their products), with increased growth gain in virulence, and lead to abscess formation.

¹ Berlin, med. Gesell., February 6, 1895, abstr. in Med. Week., 1895; 82.

² Contributions from the William Pepper Laboratory, Philadelphia, 1900.

THE MODES OF INFECTION

3. Growth of bacteria and infection in an internal organ, with no recognizable solution of continuity of a surface—"cryptogenic infection." It has been usual to regard this as brought about by some local solution of continuity which has undergone healing, or is so small as to be passed over. Such, of course, may occur, but the facts brought forward above show clearly that, through unaltered surfaces, bacteria and other microbes may be introduced by the agency of the wandering cells of the organism, and, being so introduced, may be conveyed by the lymph or blood stream to various regions where, coming to rest, they may proliferate and set up infection.

We not infrequently encounter cases of tuberculous cervical glands in children without a sign of tuberculosis of the fauces, active tuberculosis of the mesenteric glands with the mucosa of the intestine showing not a single ulcer, or at times meet with acute localized osteomyelitis, due to streptococci, with no history of, and no sign of, local surface irritation anywhere. Similarly, an acute nephritis or cystitis may suddenly supervene, with no ulcerative lesions found anywhere at autopsy to explain its origin.

It may be asked why, if pathogenic organisms are so frequently present (as we know they are) on the surface of the body, in the mouth and the intestinal contents, and if the leukocytes are thus liable to carry them into the tissues, cryptogenic infections are not far more common; why, in short, we continue to live. The answer is (1) that leukocytes. in general, taking up very virulent microbes, tend to be destroyed or inhibited, so that they do not make their way back from the surface; or, indeed, through negative chemiotaxis (p. 416), do not take them up at all. We would not lay great stress upon this, though doubtless it is a factor, for occasionally we find them ingesting distinctly virulent forms (e. g., the gonococcus) and showing little obvious arrest of activity. Of more importance, we think, is (2) that, just as one swallow does not make the summer, so a single microbe cannot, according to numerous observations, produce infection (unless it be of extraordinary virulence). A certain minimal number must be at one spot in the tissues in order to produce enough toxic material to counterbalance the opposing cell activities.1 Ordinarily, therefore, a single microbe-bearing leukocyte coming to rest at any point does not produce disease. One or a few virulent germs introduced at one point are destroyed before they have time to proliferate. Thus isolated bacteria may simultaneously be introduced at various points and simultaneously be rendered harmless. Only when we have a special concatenation of circumstances is it likely that this method of infection shows itself: (a) the presence of an excessive number of virulent microbes at one surface region; (b) con-

¹ Thus, for example, employing a most ingenious method whereby to isolate individual bacteria, Webb, Williams, and Barber have demonstrated that it requires from 50 to 150 tuberele bacilli (according to the virulence of the culture) to set up tuberculosis in the guinea-pig by subcutaneous inoculation, in this confirming certain earlier observations of Wyssokowicz (Jour. of Med. Research, 20: 1909; 1).

gestion of a mucous surface, with passage out of an increased number of leukocytes; (c) reëntrance at one region of an undue number of the same bearing with them the microbes; (d) accumulation at one spot; or recurrent deposit of such numbers of microbes that the bactericidal powers of these cells then become exhausted; and lastly (e) temporary or habitual lowered vitality of the tissues of such a region antecedent to the introduction of the microbes. With all the protective mechanisms, it is unlikely that any one of these conditions alone is liable to set up infection. We must assume the concurrent working of several. Infection, indeed, must be regarded as the outcome of a contest between the protective mechanisms of the organism and the bacteria, in which, for a time at least, the latter gain the upper hand. Whether bacteria grow in the body and set up disease or not depends thus upon two main factors-the resisting power (or susceptibility) of the tissues and the virulence of the microorganism; and both of these are capable of great variation. The first of these we shall consider in a separate chapter, for it has a bearing not merely upon the causation of infections, but of all forms of disease; regarding the latter, it is appropriate that here we should call attention to the more important data.

The Channels of Entry in Relationship to the Modes of Growth of Bacteria.-All pathogenic bacteria, it is needless to say, do not have the same habits of growth. Some are strict parasites, growing in the animal body only, and at the temperature of the body, some, indeed, only in the human body. Such, while they may retain their vitality outside the organism, cannot proliferate there, whence it follows that communication of the bacteria-and of the disease-must be direct, or almost direct, either by immediate contact or by the conveyance of the virus in the form of *fomites*, in dust, scales of shed skin, etc., or in the discharges from the person from the one individual to the other. The tubercle bacillus and the microbe of gonorrhea are thus conveyed. Or, insects act as intermediaries; then, it would seem, only in a passive manner (we here refer to bacteria only); although, for example, the typhoid and the plague bacillus can proliferate within the intestinal canal of insects, it is doubtful whether these act as more than passive carriers; certainly they are not essential. Other microbes have a much wider range of growth, and here the process of infecting may be mediate: discharged from the body of the diseased individual, they proliferate in fluid outside the body at the ordinary temperature; and growing, they may exist for weeks, and through the contaminated fluid it is that the disease becomes carried to a second individual. But bacteria which commonly are conveyed by the one means may, with slight change of conditions, be transmitted by the other, and the old and still official usage of classifying certain diseases as contagious (i. e., conveyed by contact), others as infectious, is useless, save as a euphemism. It is better to classify all as infectious, and recognize those properties of growth which render infections more liable to occur mediately in certain diseases, immediately in others. So, too, the term miasmatic, as indicating that certain diseases are brought about by a miasm, influence, or effluvium emanating from the soil, belongs to a past generation and must be allowed to die a natural death. What is of more importance is to recognize how and why specific bacteria gain particular channels of entrance.

 Organisms floating in the air, whether capable of proliferating outside the body or not, are liable more particularly to gain entrance through the respiratory tract, and especially through the upper respiratory tract, the pharynx, and tonsils.

2. Those, like the typhoid and cholera microbes, which can proliferate in water, are particularly liable to gain entrance through the intestinal tract, although it has to be noted that the mouth and pharynx are common to both the digestive and respiratory systems, and they may here also be implicated. So, too, organisms discharged in the excreta of other animals, and not necessarily propagating in them, if these excreta be used as food (e. g., milk), or contaminate food; or, again, microbes growing in the more sterile tissues of diseased animals used as food, may gain entrance by this channel. In this way, for example, tuberculosis may be conveyed, more especially to young children.

3. Organisms capable of existing in the pores of the skin more particularly are liable to proliferate when there is solution of continuity of the skin. Thus, the pyococci, streptococcus pyogenes, and bacillus pyocyaneus more particularly gain entrance by this means, although, as the skin comes in direct contact with external objects, many other microorganisms, under particular circumstances, may gain entrance in this way; or wounds are inflicted by instruments already bearing pathogenic organisms, such as rusty instruments carrying the spores of the tetanus bacillus.

4. Organisms infecting the genital passages are liable to be conveyed directly to the other sex in conjugation, as also to the child in parturition.

5. Those infecting the placenta pass to the focus along the umbilical vein.

We must, however, repeat that the channel of entrance is not necessarily the seat of manifestation of primary growth. We have to note special tissue susceptibility (p. 408), whereby bacteria gaining entrance multiply in certain tissues and not in others.

Virulence.—In the early days of bacteriology it was held that virulence was not so much the expression of active antagonistic processes initiated by the bacteria, as of a disturbance of vital processes, more or less extensive, brought about by the mere mechanical presence of these organisms within the tissues and the abstraction by the same for the needs of their growth of substances equally necessary for the growth and due activities of the tissues. The anthrax bacillus thus was supposed to absorb the oxygen of the blood and to block the capillaries. Nowadays we recognize that it is the expression of the toxic properties of the substances excreted by the bacteria, either in the course of their growth, or, it must be added, of their disintegration. Just as the tissues have their protective mechanisms, so have the bacteria, and these latter protective mechanisms, so far as they affect us, may be

summed up in the one word—toxins; although, as we have already warned the reader under this heading, we clearly include bodies of more than one order, among them the aggressins. Or, in other words, the virulence of a microorganism is the expression of the toxicity of its products. Upon analyzing further, it will be seen that virulence depends upon two factors, namely, the intensity of the toxic action of these products (of a unit of the same) and the amount discharged in a given time. A third factor determines infection, namely, the number of bacteria present affording these toxins. The intensity of action of a given solution of enzyme depends, for example, not merely upon the existence of the enzyme in the solution, capable of converting a given amount of substance in a given time, but upon the amount of enzyme present relative to the amount of material to be converted. This, however, we can neglect for the moment.

It is found that:

1. This virulence is specific, and that in two ways: (a) The toxins produced by the different species (i. e., forms differing in morphologicaland cultural characteristics) are different, so that the results of inoculation constantly vary; and (b) the toxins are active for certain species of animals only, and not for others. An organism which will cause disease in one animal may be harmless for another of different species. Regarding the first of these properties, it is to be noted that of recent years facts have accumulated showing that closely allied species may elaborate common toxins, i. e., that they produce multiple toxic bodies, some of which are common to more than one species, but others are specific, and peculiar to the one species. Regarding the other, it is a matter of common knowledge that certain bacteria are specific for man and without effect on the lower animals (e. g., the gonococcus); whereas others, like the large group of species of organisms of hemorrhagic septicemia, in general, affect particular species of mammals and birds, but are without effect on man.1

2. The virulence of a given species is subject to great variation. No two strains isolated from different individuals are of identical virulence.

3. The virulence is exalted or increased for any species by "passage" through members of that species, *i. e.*, by inoculation of a culture into an individual of that species in sufficient amount to set up symptoms of severe infection, and when these symptoms present themselves, killing that animal and inoculating some of the body fluid of that animal into a second, and of that into a third animal, etc. To this increase a limit would seem to present itself after a certain number of passages, beyond which no further increase is found to occur, but the increase may be so great that a strain of an organism like the Streptococcus pyogenes, which, before passage, will only kill young individuals, and that after a period of three or four days and with the employment of 1 c.c. of a culture, will, by passage, be made so intensely virulent that

¹ It must, however, be remembered that one member of this group, the bacillus of bubonic plague, is strikingly virulent for man.

VIRULENCE

the thousandth part of a cubic centimeter, or even, it may be, the millionth part, inoculated into an animal will cause death in six hours.

4. While thus the virulence may, by passage, be exalted for the particular species employed, it may be considerably lessened for members of another species. This is not constantly the case, but some notable examples have been recorded. Thus, without exception, the virulence of pathogenic organisms is lessened by prolonged growth in or upon media of the laboratory, more particularly when this is accompanied by transfer to new media only at long intervals. It sometimes happens that the rapid transfer of a strain-every twenty-four hours-from one medium to another in rotation will cause a development of virulence in a weak stock up to a certain moderate amount; but if allowed to "stew in their own juice," bacteria tend to become attenuated. Various other agencies lead to lessening of the virulence outside the body, such as growth at a temperature bordering upon the maximum at which the particular species will retain its vitality; exposure to sunlight; action of small quantities of antiseptic or disinfectant substances; subjection to increased atmospheric pressure, etc. Broadly, it may be stated that bacteria exhibit the action of the law that has been made out for the higher forms of life, that within certain narrow limits the struggle for existence brings about the improvement of the race; above these limits, if the race gains too complete a mastery of its environment, it ceases to advance; below these limits the struggle is conducted at a loss -the race becomes enfeebled. So, also, it is in general to be observed that as growth becomes more active, the virulence increases, although as between any two species, or even between two distinct strains of the same species, relative luxuriance of growth is not by any means an absolute criterion of relative virulence.

It is needless to say that in connection with this subject of virulenceas with all others-there is much regarding which we are still ignorant. We cannot isolate the toxins and study their effects as pure chemical substances; we do not know, and can only infer, their nature. What is more, we are still at a loss to account for the mode of action in setting up infection of the two great groups of pathogenic organisms, namely, those which characteristically discharge extracellular toxins, so that the medium of growth becomes highly toxic, and those, on the other hand, which (grown outside the body, at least) produce inconsiderable amounts of diffusible toxins. Of the former group we have such organisms as those of diphtheria and tetanus, and that of blue pus (B. pyocyaneus); of the latter, the bacillus of typhoid, the B. coli, the anthrax, and tubercle bacilli. It seems evident, in the first place, that if outside the body they produce nothing which we can recognize as toxins, inside the body they discharge or afford something which affects the cells in their neighborhood, for, introduce attenuated anthrax bacilli into the tissues, and the leukocytes rapidly attack them; whereas, virulent anthrax bacilli introduced similarly are left severely alone. The same is true as regards colon bacilli inoculated into the peritoneal cavity; whether they are taken up and destroyed, or not, depends upon

their virulence. This group of organisms does not become pathogenic because its members are apparently innocent and inert and gain a footing in the tissues without irritating the cells and simulating them to employ their protective mechanism. There may be something in this; the unsuspecting scavenger in the mouth or intestines may take up an apparently innocent tubercle bacillus and convey it into the lymph gland, only to discover too late that it has swallowed more than it can digest; but, over and above this, we have to gain more knowledge of the exact nature of the products discharged by the group of bacteria while in active growth before we can feel that we have a satisfactory knowledge of what is the nature of virulence.⁴

Before we leave the subject it is necessary to say a word regarding a matter to which we have more than once referred, namely, the numerical relationship of bacteria to infection. With the possible exception of microbes, which, like the streptococcus already mentioned, have undergone by experimental means an extraordinary increase in virulence, it appears well established that in the individual mammal in normal health a single microorganism cannot cause disease. Even when the vitality is lowered it probably requires several in close proximity in order to produce so much discharge that the antibacterial substances of the enclosing or surrounding cells become neutralized. We have noted, for instance, the observations (p. 325, note) upon the minimal number of tubercle bacilli which will set up infection. If this be so, it is obvious that the number of bacteria gaining entry to one area at one time is a factor in the setting up of disease. A large number of bacilli of low virulence will as surely cause infection as will a few of exalted virulence. The number entering is very clearly a factor. Your physician-happily of the old days, for such unwise heroism is not called for now-who may have gone through an epidemic of diphtheria, and in so doing must time and again have breathed in the bacilli of the disease without taking the disease, succumbed surely when, sucking the tracheal tube of the little suffocating patient, in order to clear the passages, he introduced a mass of the microorganisms into his own throat.

It is well to keep in mind this influence of numbers in attempting to correlate experimental results with the natural course of infectious disease. In the laboratory, that is, we are accustomed to produce disease by the employment of immediate injections of millions of bacteria. It must be remembered that in nature there is rarely any such immediate overwhelming of the tissues. We may thus, at times, obtain positive results with microbes which under natural conditions would be relatively innocuous for the species under observation.


¹ Some light is thrown upon these matters by the recent studies upon the aggressins. (See Section III, Chapter VIII.)

CHAPTER IX.

PROTOZOAN PARASITES AS CAUSES OF DISEASE.

As already noted, active interest has been aroused within the last few years in sundry microbic unicellular forms of animal life as causes of disease, and at the present time scarce a month passes without some investigator announcing the discovery of a new form of protozoan parasite in one or other animal—at times apparently harmless, at times clearly associated with the appearance of definite symptoms of disease.

FIG, 113



Trypanosomes (T, gambiense) from the blood in sleeping sickness. \times 2000,

The study is relatively so new and incomplete that it may be the time is not ripe for the broadest generalizations regarding the mode of action of these protozoa as pathogenic agents. One is tempted, that is, from a study of forms which have been very fully worked out, such as the hematozoon malariæ, to see a broad distinction between the bacteria and these animal forms in the possession by the latter of an amoeboid stage, during which the microbes can actively attack the cells of the organism, and in a large number of instances can penetrate and grow within these

332 PROTOZOAN PARASITES AS CAUSES OF DISEASE

cells, that intracellular growth being a definite stage in the life cycle. With other forms, such as the trypanosomes, while the active stage of the disease they induce is characterized by the presence of extracellular forms, swimming freely in the body fluids, their complete disappearance in certain conditions (as after the administration of arsenical compounds) and subsequent reappearance suggests strongly a latent intracellular stage such as, indeed, has been demonstrated by Breinl,¹ while previously the studies of Rogers and other Indian observers had proved that the remarkable Leishman-Donovan bodies of kala-azar are the intracellular phase of existence of a trypanosome-like organism. We see, obscurely it is true, a difference between the nature of the diseases



Development of organism of kahazar in citrated sphero blood (Rogers' method): 1, Leishman-Donovan body in fresh sphero blood; 2, after three days' culture; 3, fifth day; 4, sixth day; 6, elongated fingeliated forms, sixth day; 6, group apparently developing from intracellular forms, with remains of cell. (After Christophers.) caused by these protozoan parasites and those set up by the vegetable bacteria, in that the latter give origin to powerful toxic bodies having a widespread action, whereas, while we have some evidence, and that definite, of the existence of toxins produced by the protozoan parasites, these would seem to be of a lower order of toxicity, so low that hitherto in most cases it has been impossible to recognize the development of antitoxins and passive immunity by experimental methods.² But if Guarnieri and Councilman³ be correct in their views regarding the protozoan causation of smallpox and vaccinia, and Mallory in his regarding scarlet fever, then this distinction must be given up, for the acute exanthemata are clinically the type examples of infections -of diffuse disturbance of all the tissues set up by toxins, resulting either in death or the production of well-marked immunity. Under these conditions, the most we can do is to pass in review the different orders

of protozoa that have been found causing disease, briefly noting the main features of their growth and distribution in the body. These have been found belonging to all the main orders of protozoa—the sarco-

¹ Reports, Liverpool Sch. of Tropical Med., 1909.

² In the previous edition reference was made to the acute exanthemata as very possibly due to protozoan parasites, and hence as opposed to the formulation of any such generalization. The trend of opinion at the present time is against this view and in favor of regarding smallpox, vaceinia, measles, etc., along with the pleuro-pneumonia of cattle, yellow fever, and acute anterior poliomyelitis (Flexner and Lewis) as induced by members of the group of ultramicroscopic organisms, which by analogy of their mode of action and the mode of reaction on the part of the system are more nearly related to the bacteria than to the protozoa. (See Appendix A.)


³See the series of articles by Councilman and his associates, Jour. of Med. Research, N. S., 6:1904:1.

SARCODINIÆ

diniæ, or rhizopoda, the mastigophora, or flagellata, the sporozoa, and the ciliate infusoria.

Sarcodiniæ.—Of these the type example is the amœba (or *ent-amœba*) of dysentery. This exists in a free state in water, and would seem to be capable of multiplying in the colon; it either attacks or passes through the mucous membrane to the submucosa, where it may be found in great numbers, containing ingested erythrocytes and cell debris, *i. e.*, it lives upon and ingests the cells of the part, setting up marked inflammatory swelling, ulceration, and necrosis. A later seat





Schematic life cycle of the Amste coli: 1, the adult ameba with nucleus (a) and contractile vacuole (r); 2, the same, multiplying by antiotic division; 3, appearance of chromidial granules in cytoplasm, which enlarge and become the sporces (μ). In 4; these sporces become discharged or liberated (5) and develop (6, 7, 8) into the adult ameba, or (9) under other conditions the ameba passes into an encysted starse. (After E. L. Walker.)

of election may be the liver, in which necrotic abscesses may be set up, the amoba being present in abundance in the tissue of the boundary zone of the abscess. Occasional rounded encysted amoba are to be made out more particularly in the walls of the colon. Whether the spores developed from these cysts developing into minute amoba remain throughout extracellular or exhibit an intracellular stage has not been determined.

It will be seen, therefore, that the indications are that these pathogenic amoeba live within the organism by attacking and digesting the cells and cell substance. Their action is essentially local, and whether

334 PROTOZOAN PARASITES AS CAUSES OF DISEASE

the remote effects seen in dysentery are to be regarded as, in part at least, due to the effusion of any toxic substances, discharged by the amœbe themselves, or wholly to the products of cell destruction along with the secondary infection of the ulcers in the colon, has not as yet been fully determined.¹

Mastigophora.-Of these, the trypanosomes may be taken as a type. These form a widely spread class of pathogenic protozoa, numerous species being found as blood parasites in vertebrates, both cold and warm blooded, from the fish and frog upward to man himself. The forms most fully studied have been the Trypanosoma evansii, causing a disease of horses in Assam, India, and the Philippine Islands known as surra (Evans, 1880); the Trypanosoma brucei, the cause of the n'gana, or tsetse-fly disease, affecting horses and cattle in Southeast Africa (Bruce), 1894; the allied form associated with dourine, or mal de coit (Rouget, 1896), in Algeria and Southern Europe, as also recently through importation in the United States and Canada; and with mal de caderas of South America, also affecting horses (Elmassian), It is now fully established that another trypanosome, the 1901. Trypanosoma gambiense, is the causal agent in the remarkable disease, sleeping sickness, which is spreading rapidly in Western and Central Africa, so rapidly that it is calculated that no less than half a million natives have died from the disease during the last ten years (Dutton and Ford, 1902; Castellani, 1903). Rogers and others have demonstrated that the minute Donovan-Leishman bodies found in the enlarged spleen in the Indian disease of man known as kala-azar, or dum-dum fever. are one stage in the life cycle of another trypanosome; and of the same order are the similar bodies found abundantly in the intractable Oriental sores which go by various names in various regions-Delhi boil, Aleppo button, etc. (J. H. Wright, 1903).

Without entering into the full details, for those belong to works devoted to the animal microparasites, it may be recalled that these minute organisms found in active motion in the removed blood possess an elongated, spindle-shaped body, with undulating membrane along one side, whose outer differentiated border, beginning within the head end of the organisms as an offset from a remarkable refractile granule, spoken of variously as centrosome or micronucleus, continue beyond the body as a flagellum (Fig. 113). There is a nucleus; at times a contractile vacuole may be made out. The length may be as much as 30 μ , or, in the largest forms, the *T. theileri* of cattle, 50 μ , or the huge *T. ingens*, recently discovered by Bruce² and his associates in Uganda, inhabiting the blood of the ox and reed buck, from 72 to 122 μ . The breadth ordinarily is from 2 to 3 μ . Multiplication is by a process of longitudinal fission, in which, in some cases, the nucleus, in others the micronucleus, first undergoes division.

¹ An admirable study of the amœbæ parasitic in animals has recently been published by Walker, of Boston, Jour. of Med. Research, 17: 1908; 379.

³ Bruce, Hamerton, and Bateman, Proc. Roy. Soc., B., 81: 1909: 323.


MASTIGOPHORA

This simple fission is, so far, the only mode of multiplication known; no sexual cycle has been made out, even though the organisms are transmitted through alternate hosts. It is this form also that alone is seen when, as first determined by Novy, the trypanosomes are grown

upon the media of the laboratory. In this they differ from the forms presently to be noted. Reference has already been made to the intracellular stage of these organisms. Further work is required to determine whether this be a necessary phase in the life cycle. although in addition it may be recalled that the brilliant protozoologist Schaudinn laid down that the Halteridium found parasitic in the erythrocytes of the owl, crow, and other birds is not a distinct and separate form, but is one stage in the life cycle of a trypanosome. It has been suggested that the *Piroplasma* or intracellular parasite of Texas fever in cattle has like relationships, a suggestion somewhat strengthened by the recent observation of



Piroplasma in red corpuscles of cattle suffering from Texas fever, (Smith and Kilborne.)



Glossina palpalis (\times 3%), the carrier of the trypanosome of sleeping sickness.

Nuttall and Hadwen,¹ that both *P. bovis* and *P. canis* disappear in animals treated with trypanroth and trypanblau.

¹ Proc. Roy. Soc., B., 81; 1909; 348.

BROTOZOAN PARASITES AS CAUSES OF DISEASE

The striking feature of all this group as disease producers is that they are conveyed to man and warm-blooded animals by the bites of insects. Specific insects act as intermediate hosts. In n'gana it is the testes effy (*Glossina morsiluns*); in sleeping sickness, a closely allied fly, *Glossina polpalis*. These act as intermediate hosts, the trypanosomes, succeed into the stomatchs, gaining cutrance thence into the issues of the fly and being discharged with the fluid lubricating the mouth parts—thus gaining entrance into the body of a larger animal when that is bitten by the fly.


Sether fresh blood, nor varmed, nor organic extracts, nor bile from affected animals have the slightest toxic effect; nor by centrifugalized

Inaterial, consisting almost wholly of trypanosomes, acted on by alternate freezing and thawing, nor by drying, could Laveran and Mesnil gain any indication of the presence of toxins,

The symptoms produced by trypanosomes are, as a class, essentially those of blood and circulatory disturbances — anemia, with a mild grade of

FIG. 119

Megastonne enterieum, Grassi, ventral and side views. (Schewinkoff.)



Trichomonas vaginalis,

Rever: statestica and accites, depression of cerebral activity, and coma. The indications are that, by the very abundance of these parasites in the blood, and their tendency to become agglutinated under certain conditions, they are apt to accumulate in and block capillaries of the brain and other organs, a phenomenon not encountered with the more minute bacteria, save as the result of local proliferation.

Other flagellate industriant *Tvichomonus* and *Megustoma* (*Lamblia*) intestimulis¹—have been rarely encountered in the intestinal discharges; most often in association with conditions of chronic diarrhora. Whether they have any casual relationship to the diarrhora is still a matter of debate. Recent observations indicate a close relationship between Trichomonas forms and the Trypanosomes.

Sporozoa.—The importance of this group of animal parasites as causes of disease in man may be estimated from the fact that in tropical

¹ The Cercomonus intestinulis used to be regarded as a separate form; Doffein regards it as identical with the Lamblia.



PLATE XII

Fig. 1.-Tertian Malarial Plasmodium.

2. Pigmented ring form.

7. Segmenting forms. 8. Flagellate form. (Microga9. Non-flagellate form, (Macro-

10. Segmenting form after de-struction of red corpusele.

Fig. 2.-Quartan Malarial Plasmodium.

 Segmenting forms after the destruction of red corpus-cle. 9. Flagellate form. (Microga-10. Non-flagellate form. (Macro-

Fig. 3.-Tertian Æstivo-autumnal Malarial Plasmodium.

10. Flagellate form. (Microga-

Fig. 4.-Quotidian Æstivo-autumnal Malarial Plasmodium.


5 to 7. Pigmented forms. In 6

9. Flagellate form. (Microga-


10, 11, 13 and 15. Crescentic forms.


14. Non-flagellate forms. (Ma-


PLATE XII



F1G, 3







SPOROZOA

and subtropical regions one disease of sporozoal origin—ague or malaria —occupies the position assumed by tuberculosis in the temperate zone. It has, indeed, been claimed that this disease brings about an even greater mortality.

All the organisms of this group are characteristically intracellular in their habits. In other words, the primary disturbance set up by them is that of cell parasitism, the microbes growing within and at the expense of individual cells, arresting their function and eventually leading to cell death-the cycle of the life history of the parasites being such that the maturation and spore formation of the intracellular individual coincides roughly with the exhaustion and death of the host cell. The spores, becoming free after a longer or shorter period of incubation, develop into minute amœboid forms, which penetrate other cells and repeat this process of asexual multiplication. This asexual cycle of forms may be repeated again and again. But now, in very many of these sporozoa, it has been determined that a second, sexual, cycle may be intercalated under certain conditions, more particularly (though not in every case) in connection with the transmission of the parasite from host to host. this second cycle being apt to occur in an intermediate host of another species. In ague, for example, the hematozoon malariæ exhibits the asexual cycle in the blood of man, the sexual cycle within the mosquito (various species of Anopheles), which, by feeding on human blood, acts as a transmitter of the disease to a second individual. when the sporozoites, the products of the sexual cycle, are introduced into a surface vessel of that individual along with the proboscis of the mosquito. It would seem, thus, that a large number of the sporozoa gain entrance into the systems of animals and are transmitted from individual to individual through the intermediation of biting and sucking insects. Not all, however; others, like the coccidiæ, gain entrance through the digestive tract.

Several suborders of the sporozoa contribute parasites to man and the higher warm-blooded animals. These we will rapidly note, calling particular attention to those data which throw light upon the mode of causation of disease.


(a) Hematosporidia.—Of these, the type example is the Hematozoon malaria. Other allied forms infesting the red corpuseles are met with in the blood of birds and other animals, and the study of these has elucidated the life history of the malarial parasite. Thus it was W. G. MacCallum's discovery of the process of conjugation between the forms present in the blood of Canadian crows that afforded the clue to the nature of the "flagellate" bodies of the malarial organisms. (See Plate XII, Fig. 1, No. 8; Fig. 2, No. 9, etc.) Some of these, like the halteridium of the owl, have been shown by Schaudinn to possess close affinities to the trypanosomes rather than to the sporozoa, or otherwise it may be suggested that the trypanosomes and sporozoa are closely allied. There is, however, some doubt regarding this observation of Schaudinn's. The main points to be noted regarding the hematozoon in relationship to ague are:

338 PROTOZOAN PARASITES AS CAUSES OF DISEASE

 The disease, being transmitted by particular species of mosquito, the anopheline, is only endemic where members of these species are present.

2. The anopheles, like all mosquitoes, lays its eggs in relatively still water, and the larvæ are aquatic. Save under the influence of strong winds, the mosquitoes do not travel any distance from their place of birth and from water. Malaria, therefore, is largely confined to lowlying, swampy, or badly drained regions and the neighborhood of stagnant water.

 The anopheles bites at night, not during the day; infection, therefore, occurs at night. It may be single or multiple, on different nights.



Anopheles maculipennis: solult male at left, female at right. (Howard.)

4. For its development the asexual cycle requires different periods in the different species of hematozoon—forty-eight hours for the organism of tertian fever; seventy-two for that of quartan; forty-eight hours (with irregular variations) for that of the estivo-autumnal type. The periodic attacks of ague are directly determined by these cycles, the chills and fever coinciding with the maturation of the hematozoa and their sporulation. Presumably, it is the breaking down of the corpuseles and liberation of the cell debris and pigment matter rather than any specific toxin (for this has not been determined) that is the cause of the febrile attacks.

5. The clusters of pigment and cell debris are apt to be separated from the blood in the spleen, there setting up those changes which lead to the enlargement of that organ. They may also accumulate in the

SPOROZOA

capillaries of the brain, of the kidney (Ewing), and other organs, setting up disturbances by arrest of the circulation.

6. The observations of Calkins on prolonged asexual multiplication of protozoa show that this leads to progressive weakening and degeneration of the later generations. Where, therefore, the affected individual removes himself to a region where he cannot be re-infected, it would seem that there is a natural tendency for the malarial organisms to become weaker and weaker, and so for this disease to pass off. Apart from this probability, there is evidence that more particularly young children (Koch) are relatively resistant to the disease, and despite repeated infection, gain immunity—*i. e.*, the power of destroying the

hematozoa—while in adults improvement in general health, brought about by removal to another climate, etc., would seem to favor the destruction of the hematozoa. This notwithstanding it is evident that, as with many bacteria, certain of the parasites may for long periods lead a latent existence within the body, and after the expiration of months and years take on active growth with the development of new crises of the disease.¹

The continuance of the disease over long periods with periodic exacerbations, but without reinfection would seem due to the occasional intermission of an imperfect, parthenogenetic, sexual cycle within the bloodvessels of the human host.² The other members of the sporozoa, while abundantly parasitic and pathogenic in worms, inseets, and



A. Λ " Rainey's corpuscle" or colony of sarcosporidia lying between the muscle fibers; B, the individual sarcosporidia (sarcoblasts) composing the same. (Perls.)

the lower forms of animal life, are rarely encountered in the higher animals, still more rarely in man. (b) The gregarines, for example, are not found in vertebrate forms: (c) the neosporidia (including the myxosporidia and the sarcosporidia), while occurring among vertebrates—the former common in fishes, the latter giving origin to Rainey's corpuscles within the muscle fibres of mammals—are almost unknown in man. (d) Of the coecidia, one form, the *Coecidium oriforme*, most common in the rabbit, where it affects more especially the upper part of the small intestine and


¹ Regarding this there is some debate. So high an authority as Sir Patrick Manson regards a new attack after months or years of freedom from malaria as evidence of reinfection.

² The existence of this type was first demonstrated by Schaudinn,

340 PROTOZOAN PARASITES AS CAUSES OF DISEASE

the liver, has been encountered not half a dozen times in man. As determined by Schaudinn and Siedleeki and Simond, the coccidium and other members of the order exhibit, like the hemosporidia, a sexual and an asexual cycle, though these occur in the one host.

Forms which curiously resemble certain stages in the sporozoan life cycles have been encountered by Pfeiffer, Guarnieri, and Councilman in




Life cycle of *Cocidium schulergi*. Sporonites penetrate epithelial cells, and grow into adult intracellular parasites (a). When mature, the nucleus divides repeatedly (b); and each of its subdivisions becomes the nucleus of a merozoite (c). These enter new epithelial cells, and the cycle is repeated many times. After five or six days of incubation, the merozoites develop into sexual differentiated gameters; some are large and well stored with yolk material (macrogrametes, d, e, f); others have nuclei which fragment into many small particles ("*Chromidia*"), each granule becoming the nucleus of a microgamete or male cell (d, h, i, j). The macrogamete is fertilized by one microgamete (g), and the syster immediately secretes a fertilization membrane which hardens into a cyst. The elsevage nucleus divides twice, and each of the four daughter nuclei forms a sporoblast (b) in which two sporonizes are produced (d). (After Schaudinn.)

the epithelial cells of the vaccinia and smallpox eruption (Fig. 123). In the former, Councilman and his associates determine but a single intracellular cycle; in the latter they found a second intranuclear cycle. Mallory, likewise, has described a remarkable stellate or rayed intracellular form in the epithelial cells in cases of scarlatina, recalling the daisy form assumed by the quartan malarial organism in the process of

CILIATE INFUSORIA

sporulation and others employing his method have confirmed his findings. The meaning of these forms is still *sub judice*.

During the last decade of the nineteenth century it was held by numerous observers (Sjöbring, Ruffer, Metchnikoff, Soudakewitch, Plimmer, etc.) that bodies seen within the cells of malignant growths—



a, Guarnieri's bodies in the cytoplasm epithelial cells in vaccinia (and smallpox); b, one stage of the presumed intranuclear cycle of these bodies in the epithelial cells in smallpox. (After Cakins.)

more especially of cancers—were of the nature of sporozoa. That such is their nature is now generally denied. While certain remarkable bodies are to be observed with fair frequency, sometimes in great abundance, the general opinion nowadays is that these are modified cell and nuclear products, and that they indicate peculiar forms of cell degeneration. In short, able and distinguished as are all the observers just



Balantidium coli.

mentioned, the trend of opinion now is opposed to regarding the vaccine and cancer bodies as other than allied to the chromidia or plasmosomes mentioned on p. 48.

Ciliate Infusoria.—These, the most highly specialized of the protozoa, are the least frequent of all the protozoon parasites. One form

342 PROTÒZOAN PARASITES AS CAUSES OF DISEASE

alone has been recorded —apparently with some justification—as possessing pathogenic properties, and this is not found within the tissues, but free in the alimentary canal. This is the *Balantidium coli*, a form not unlike the common paramecium of pond water, but more oval, and with the oral aperture more definitely at the one pole, the anus at the other; it is abundantly ciliated; this is said to be a normal parasite in the hog. Like the anœba coli, it is supposed to gain entrance into man through contaminated water. It has been found associated with extensive catarrhal inflammation of the colon, with dysenteric symptoms.¹ Observers have described a second species, the *Balantidium minutum*, of smaller size, found in association with other intestinal parasites.

Of the most highly differentiated suborder of the infusoria, the **Suctoria**, no parasitic examples are known in or upon the higher animals.

¹ In 1904 Strong (Pub. Phil, Isl. Bur. Sci., Biol. Lab., No. 26: 1904; 1) collected from the literature 127 cases of Balantidium infectius. Bowman has found the parasites in the mesocolic lymphatic glends as well as in the ulcerated walls of the large intestine (Philippine Jour. of Sci., 4: 1909; Section B; 417).

CHAPTER X.

METAZOAN PARASITES AS CAUSÉS OF DISEASE.

THE members of a limited number of classes of metazoan or multicellular animal organisms have adapted themselves to growth within the organisms, and, in certain cases, even within the tissues of other higher metazoa. Here I shall not attempt to describe these forms; such descriptions belong properly to works upon parasitology. At most it is necessary to note rapidly the classes capable of this parasitic existence. These are confined to members of the phyla, or groups **Platyhelminthes** and **Nemathelminthes** and rarer members of the **Insecta** and **Arachnidæ**. Among the platyhelminthes, or flat-worms, we encounter members of the *Trematodes*, or flukes, and the *Cestodes*, or tape-worms. The *Nematodes* are the chief representatives of the nemathelminthes, or round-worms.

CHARACTERISTICS OF METAZOAN PARASITES IN GENERAL.

Without exception, it may be stated that adaptation to a parasitic existence has been accompanied by simplification and retrogression. Forms that have not to hunt for their food, but receive it in a soluble, assimilable state, prepared by their host; that, further, are largely protected from the effects of external influences by their very mode of life, do not need elaborate organs of locomotion or an elaborate digestive apparatus; do not need organs of protection and defence beyond the means of neutralizing the digestive influence of the juices of their host. Nor do they need weapons of offence beyond those necessary to attach themselves to that host and penetrate its tissues in such a way as to gain therefrom the requisite pabulum. Thus, the organisms of these parasites are apt to become reduced to very simple terms; limbs and organs of locomotion may become rudimentary, sense organs atrophied, and, as in the cestodes, or tape-worms, the alimentary tract may wholly disappear, nourishment being gained purely by surface absorption. The prime necessity is the retention of life of the individual and preservation of the species, so that means have been developed to neutralize the harmful consequences of the host being but mortal and liable to die. Either the parasites are capable of existing for considerable periods outside the body of the host until fortuitously taken up by another, or are capable of living in different forms in a succession of different hosts, or lastly, and most commonly, have enormous reproductive capacity, becoming little more than animated masses of sexual glands, enormous

344 METAZOAN PARASITES AS CAUSES OF DISEASE

quantities of ova being produced and discharged, in preparation for the probability that, with rare exceptions, these will fall upon barren ground. More accurately, so precarious is this method of handing on the torch of life that only those species possessing an enormous reproductive capacity can possibly survive. For this retrogression and simplification of structure inevitably carries with it a lessened capacity on the part of the individual to adapt itself to other than a very narrow set of conditions; it has reduced its methods of offence and defence to a minimum, and thus we find, as a general rule, that a given species can only grow actively in a particular species of hosts. Or, conversely, each species of animal has its particular set of parasites, which differs from that of other species. At most, certain species of parasites may pass a particular stage of existence in closely allied species of hosts.

How, it may be asked, do these metazoan parasites cause disease?¹ It must be noted, in the first place, that on general principles we should not expect such parasites to set up severe disease. With forms that require a considerable period for the development of their life cycle, forms in which, further, as we have noted, the survival of the species by means of passage into other hosts is precarious, it would be a suicidal policy so to injure the host as to lessen its capacity to obtain nourishment, or to arrest its power of locomotion, and, indirectly, its opportunity to distribute the eggs of the parasite. And, as a matter of fact. we find that these larger parasites tend rather toward symbiosis-to harmonious living together, with minimal disturbance to the hostthan toward the production of states of severe disease. Nevertheless, the symbiosis is in no sense voluntary on the part of the host, nor have we any indication that the presence of parasites is an advantage to that host. We no longer hold, as did Jordens, in 1801, that intestinal worms are "the good angels and unfailing helpers of children." The indications are to the contrary. If, in general, the disturbance set up is slight, it is nevertheless there, and we have indications of definite reactive processes on the part of the host. The injury set up and the evidences of disturbance are of five orders:

1. That of actual presence in some organ, leading to displacement and pressure.

2. Disturbance due to migration of the parasites from one organ of the host to another.

3. Direct destruction of tissues.

4. Loss of foodstuffs diverted by the parasite and used up by it.

5. Disturbances induced by the excretions of the parasites.

These we may pass rapidly in review:

Injury Caused by the Mere Presence of the Parasite.—This, in general, is *nil*. There are, however, exceptions. A *Filaria nocturna*

¹ For the arrangement of the following paragraphs and the data therein contained we are largely indebted to a valuable summary of the subject by Shipley and Fearnsides, Journal of Economic Biology, 1:1906; No. 2,



EXPLANATION OF FIGURES IN PLATE XIII

VARIOUS STAGES OF MALLORY'S INTRACELLULAR PARASITE IN THE EPITHELIAL Cells in Scarlatina. (Mallory.)

The drawings were made with the Abbe camera lucida; projection on to table. Zeiss apochromatic homogeneous immersion 2.0 mm., apert. 130, compensation ocular 6.

Figs, 1 and 2 show numerous large and small searlet fever bodies (stained light blue) in and between the epithelial cells of the rete mucosum. In Fig. 1 is a large body in a lymph space of the corium just underneath the epidermis. Several of the bodies suggest fixation while in amorboid motion.

FIGS. 3, 5, and 6 are coarsely reticulated forms which may be degenerated forms of the scarlet fever bodies, or stages in sporogony.

FIGS. 4, 8, and 9 probably represent stages preceding the radiate bodies. In Fig. 9 the bodies lie in a lymph space. It shows also four small forms which have just got free from a rosette.

FIGS. 7, 10, 11, 12, 13, 14, and 15 show different stages in the development of the radiate bodies.

Fig. 10 is the earliest stage: there is a distinct central body and a definite, regular arrangement of granules at the periphery. Figs. 7, 11, and 12 show a little later stage of development; 11 and 12 are optical sections, while 7 is a surface view. Moreover, in Fig. 7 the body lies free in a lymph space in the corium. The segments begin to show a certain amount of lateral separation from each other. Fig. 13 is a still later stage; the segments are increasing in size and are more or less free from each other, although most of them are still attached to the central body. In Fig. 14 the segments are all free and enlarging, although still grouped around the central body. In Fig. 15 the bodies are still grouped around the central body, which is free and stains deeply with cosin.

(Mallory.)






PLATE XIV



The Kidney Worm, Dioctophyme Renale (Eustrongylus Gigas) of Man, from a Specimen in a Dog. Natural size. (Stiles.)



CHARACTERISTICS OF METAZOAN PARASITES IN GENERAL 345

in a lymph vessel or gland may set up little disturbance, but if it, or its eggs, blocks the vessel, elephantiasis or chyluria may ensue. A *Cysticercus cellulosa* in the muscle is without obvious effects; in the brain, by pressure upon important nerve centres, it may cause fatal results, and in the eye may lead to blindness. In this respect




Diagram of an Echinococcus hydatid: cu, thick external cuticle; pa, parenchymal (germinal) layer; c, d, e development of the heads according to Leuckar; f, a, h, i, k, d exployment of the heads according to Moniez; l, fully developed brood capsule with heads; m, the brood capsule has ruptured, and the heads hang in the lumen of the hydatid; n, liberated head floating in the hydatid; a, p, q, r, s, mode of formation of secondary excessions daughter cyst; l, daughter cyst, with one endogenous and one excessions granddaughter cyst; u, v, z, formation of exceenous cyst (after Kuhn and Davaine); y, s, formation of acquere cyst, u, v, z, formation of exceenous cyst (atter Kuhn and Davaine); y, z, formation z, form a brood capsule; erag, constricted portion of the mother cyst. (R. Blanchard, slightly modified.)

the larval form of the $Txnia\ echinococcus$ is the most dangerous human parasite, the size attained by the cysts in the liver and elsewhere being so considerable. In the dog, the *Eustrongylus gigas* (or *Dioetophyme renale*), growing slowly and attaining great size, may eventually, by pressure atrophy, replace the whole of the kidney substance.

346 METAZOAN PARASITES AS CAUSES OF DISEASE

Injury Caused by Migration.—This, again, may be infinitesimal. The minute larvæ of *Filaria nocturna* make a nightly migration from the deep-seated bloodvessels of the internal organs to the peripheral vessels



Larva of Filaria bancrofti in the blood of man, in Egypt: Nerv. nervous system; Ex., excretory; An., anus, \times 514, (Looss.)

without causing disturbance. The Filaria medinensis, the longest of the round-worms in man, may, without symptoms, make its way through the tissues of the body until it comes to lie under the skin of the leg, and then only sets up disturbance when it pierces the skin to allow escape of its ova, or if, in this situation, it becomes ruptured. The disturbance may, on the contrary, be very marked; most severe are the fever, myositis, and muscular pain, set up by the migrating Trichina larvae prior to encystment. A common cause of irritation and itching is the nocturnal passage outward at the anus of the Oxyuris vermicularis (Fig. 127). The observations of Looss and others of late years upon the life histories of the Ankylostoma duodenale and the Strongyloides intestinalis have demonstrated that the larvæ hatched from the eggs in water or moist earth gain entrance into the human host through the skin, where they set up a dermatitis, known by diverse names in different localities-"ground itch," "coolie itch," etc.

Injury Set Up by Active Destruction of Tissue.—According to Looss,¹ the ankylostomes feed upon the mucous membrane



FIG. 127.—Oxyuris vermicularis, the pin-worm, natural size; male smaller, female larger form, with sharply pointed tail.

Fig. 128.—Ankylostoma duodenale, the old-world hook-worm, natural size; the female is the larger and more curved.

FIG. 129.— Trichocephalus trichiurus, the whip-worm, natural size, male and female.

¹ Reports of the Egyptian Government School of Medicine, 3: 1905.

CHARACTERISTICS OF METAZOAN PARASITES IN GENERAL 347

of the small intestines, and only suck in blood when by chance they pierce an underlying capillary. Somewhat opposed to this view are the observations of Leo Loeb and Smith that there is to be obtained from the superior body region of these worms a discharge which has marked effects in arresting the coagulation of the blood. This suggests that the worms are prepared to pierce the intestinal capillaries, and that the continued minute hemorrhages may be a factor in the production of anemia. But it has been questioned whether the anemia of ankylostomiasis is not a secondary result of this erosion—not from loss of blood, but from absorption through the damaged mucosa of toxic intestinal contents, if

not from increased passage into the tissues of intestinal bacteria and the production of a state of sub-infection (p. 321).1 A similar and more severe anemia is set up by another parasite, the Dibothriocephalus latus, which attaches itself to the intestinal wall, and in so doing injures the mucous membrane. But, as will be shown shortly, another cause has been determined for the anemia in these cases. Of late, Metchnikoff² and Guiart³ have called attention to the fact that, by its long, whip-like anterior end the Trichocephalus trichiurus (T. dispar) (Fig. 129) can bore through the wall of the intestine, and have suggested that the escape of the intestinal bacteria along the fine passage thus made is a probable cause of some cases of peritonitis and appendicitis; nay, would look upon this as a common cause of appendicitis. The examination of thousands of sections from cases of appendicitis without once encountering in them anything corresponding to a trichocephalus must negative this latter view.

With these exceptions, the internal parasites cause singularly little destruction of tissue: to the passive dest



Larva of Strongyloides stereoralis as found in fresh feces: *Nerr.*, nervous system; (E.e., œsophagus; *Int.*, intestines; *Gen.*, genital primordium; *A.n.*, anus, × 228. (Looss.)

struction of tissue; to the passive destruction by pressure atrophy we have already referred.

Injury by Loss of Foodstuffs.—As pointed out by Shipley and Fearnsides, this, in general, is so slight as to be negligible. "A female roundworm, *Ascaris lumbricoides*, produces 42 grams of eggs every year, and must also extract from the host a certain amount of nutriment for herself

¹See also Section III, Chap. III.

² Bull. Acad. de Méd., Paris, 45: 1901: 301.

³ Ann. de l'Inst. Pasteur, 15; 1901; 440.

348 METAZOAN PARASITES AS CAUSES OF DISEASE

besides the amount that goes to build up the ova. When present in large numbers—and Franconneau Dufresne describes a case in which a boy got rid of 5000 worms in less than three years, and on one day evacuated 600—the loss is certainly serious. Strongyloides intestinalis (vel stercoralis) at one time thought to be the cause of Cochin China diarrhea, exists in such numbers that it is not uncommon for 100,000 to be expelled at one time. Such a number is said to weigh 200 grams." Others have described the evacuation of as many as a million at a time. Despite the small size, such numbers indicate a severe strain upon the host. Blanchard regards the anemia caused by the liver fluke, *Fasciola hepatica (Distoma hepaticum)*, as due to the fact that these nourish themselves in the blood which they suck from the small capillaries of the bile ducts inhabited by them. Shipley and Fearnsides doubt whether there can be any great loss of blood from this source, and ascribe the anemia to toxic action.

Morbid Conditions Caused by Excretions: Toxic Action of Metazoan Parasites .- The very definite symptoms which accompany the presence of the metazoan parasites, and the difficulty of explaining those symptoms by the extent of the lesions seen to accompany their presence, have led pathologists during recent years to surmise that, like bacteria, these discharge toxins which, diffused into the tissues and blood, are the essential cause of these symptoms. And they have been encouraged to hold this view by Weinland's' intensely interesting demonstration that cestodes defend themselves against the action of the digestive juices by the elaboration and excretion of an antibody (see Section III, Chapter VIII)-an antitrypsin, according to Weinland and Hamill, an antikinase, according to Dastre and Stessano-a body comparable with that elaborated by the cells of the intestinal mucous membrane, whereby it also prevents digestion. The demonstration of the existence of these defensive bodies favors the supposition that the parasites excrete also offensive substances. As a matter of fact, we have abundant proof that the tissues and body fluids of many of the parasites are themselves distinctly toxic. The cyst contents of T. echinococcus and other echinococci have been found toxic, setting up in the lower animals, when injected, peritonitis and urticaria.² Disturbances of a similar order have been noted to follow the rupture of echinococcus cysts in man. While there is little evidence of the existence of these body poisons in the trematodes, in the nematodes it is abundant and convincing. The body cavity fluid of Ascaris megalocephala, spurted accidentally into the eye, has set up violent corneal inflammation, and Charlton Bastian, Miram, and von Linstow have experienced uncomfortable, and even severe, effects from the mere emanation from this parasite when dissecting it-sneezing, conjunctival irritation, paroxysmal asthma, etc.; while 2 c.c. of the

¹ Zeitsch. d. Biol., 44: 1903: 1.

^a Blanchard, Traité de Zool. Méd., Paris, 1885–89; Debove, Compt. rend. Acad. des Sci., 105: 1887; 1285; Mourson et Schlagdenhaussen, Bull. et Mém. de Soc. Méd. PHóp., Paris, 5; 1888; 113.

TOXIC ACTION

expressed body fluid has been found to kill the guinea-pig within forty hours. Cattaneo' has gained like results with *Asearis lumbricoides*.

But the mere existence of toxic substances in the bodies of these animals is no evidence that they excrete toxins. It may, for instance be pointed out that the blood of one species of warm-blooded animal is toxic for other species; that is no proof that these excrete actively toxic substances. At most, the irritative effect of the emanations from the Ascaris megaloecphala suggests that toxic substances may be discharged, and in the case of the Dibothrioecphalus latus there is clear evidence that this is the case. This worns sets up a most pronounced form of pernicious anemia. Schauman and Tallquist² have produced a like anemia in the dog, by injecting extracts of the bothriocephalus.

Gravid segment (proglottis) of Dibothriocephalus latus, showing the rosette uterus in the median line, $\times 6$, (Leuckart.)

Of bodies which, perhaps, should not be included among the toxins, we possess some evidence. Thus, Leo Loeb and Smith have demonstrated that there is to be obtained from the anterior body region of the hook-worm (Ankylostoma) a discharge possessing pronounced effects in arresting the coagulation of blood. Their observations—contrary to those of Looss, already noted—would suggest that these parasites are prepared to pierce the intestinal capillaries; and secondly, that the continued minute hemorrhages are a factor in the production of anemia in those cases.

We possess, however, what is the clearest evidence from another quarter. I refer to the *cosinophilia*, or increase in the number of cosinophile leukocytes in the blood, which characterizes the presence of almost every vermiform parasite. That eosinophilia cannot be explained save on the assumption that there diffuses from the parasites into the tissues and so into the blood, some substance which, conveyed to the bonemarrow and other seats of origin of the eosinophile cells, there stimulates the proliferation and increased production of these cells. Müller


² Deutsche med. Woeh., 24: 1898: 312. Schaumann is the author of the classical work upon Bothriocephalus anemia (Z. Kenntniss der sog. Bothriocephalusanömie, Berlin, 1894). See also Askanazy, Zeitschr. f. klin. Med., 27: 1895: 492.

¹ Abstr. in Arch. ital. de Biol., 42: 1904: 496.

350 METAZOAN PARASITES AS CAUSES OF DISEASE

and Rieder' would seem to have been the first to call attention to this remarkable phenomenon, in the case of two patients infected with ankylostoma. Bucklers,² in 1894, first showed that eosinophila characterizes all forms of helminthiasis. All kinds of parasitic worms, from the harmless oxyuris upward, induce eosinophilia, and, accompanying this, as shown by Leichenstern, there is constantly the presence of Charcot-Leyden crystals in the feces (a frequent finding in cosinophilia in general).

Whereas, the normal percentage of cosinophiles, compared with the other leukocytes of the blood, is between 1 and 4 per cent. (25 to 500 per cmm.), in 50 cases of Bilharzia disease Douglas and Hardy found an average of 16.48 per cent., a maximum of 40 per cent. In trichinosis, T. R. Brown, of Johns Hopkins, has called attention to the marked cosinophilia, the proportion of cosinophiles in his first case rising from



Gravid segment of pork-measle tapeworm (Tania solium), showing the lateral branches of the uterus, enlarged, (Stiles.)

37 to 68.2 per cent.; in other cases he obtained percentages of 42.8, 45, and 48, respectively—observations which have since been abundantly confirmed, Harlow Brooks gaining a count of 84, and Kerr one of 86.6 per cent. In guinea-worm disease the same phenomenon is observed; in 6 cases Balfour found an average of 19.6, the figures varying from 6.4 to 36.6. So, also, with ankylostomiasis; here all recent observers call attention to cosinophilia as a constant feature, while percentages as high as 72 (Leichenstern), 66 (Boycott and Haldane), and 53.5 (Ashford) have been recorded. In cestode disease, various forms of filariasis, in ascariasis, and to a less extent in infection

Deutsch, Arch. f. klin, Med., 48; 1891; 96,
 Münch, med, Woch., 41; 1894; 21,

with Oxyuris vermicularis, eosinophilia shows itself in well-marked cases, though never to the same extreme extent as in the conditions already noted.

The intimate relationship between the parasites and the eosinophilia, as noted by Ward,¹ is shown by the following facts: (1) The increase and decrease of eosinophilies in the peripheral blood coincident with the appearance and disappearance of the Filaria bancrofti in the superficial capillaries; (2) Opie's experiments upon the induced trichinosis of the guinea-pig. It was found that the increase of eosinophiles dates from the beginning migration of the embryos, and that the eosinophilia reaches its maximum when the majority of the embryos are in the process of transmission from the intestinal mucosa by way of the lymphatics and the blood to the muscular system; (3) Sabrazes has also noted the accumulation of eosinophiles in the neighborhood of hydatid cysts. I am informed by Dr. Todd that at the site of attachment of ankylostomes to the mucosa of the small intestines a marked accumulation of eosinophiles is to be detected. All these observations indicate a positive chemiotactic influence leading these cells toward the source of the stimulus.

As we have said, it is difficult, when we consider the different modes of life of these different forms, to arrive at any other conclusion than that the parasites afford a diffusible irritant, or toxin, which stimulates the proliferation and increased entry into the blood of the eosinophile cells. If we accept this, then the frequent accompanying anemia is most rationally ascribed to a like cause.²

424

INSECT AND ARACHNID PARASITES.

Little need be said regarding the insect and arachnid parasites. These produce local disturbances (1) by puncture and the introduction of irritative salivary secretion, (2) by burrowing, as in scabies, or (3) by the deposit of eggs within the tissues and germination of the same, as in *Myiasis*, and (4) by affording means for secondary entrance of infective microbes. (5) Koch, I am informed, is of the opinion that the chigger, or sand fleas (*Sarcopsylla pewetrans*), gains entrance into the corium of the foot not through the action of its mouth parts, but through the digestive action of an excretion.

¹ Science, 25; 1907; 201. A thoughtful article on the effects of parasites on their host.

² A very full bibliography and analysis of cases of the cosinophilia of helminthiasis is afforded by Shipley and Fearnsides (loc. cit.). The observations to which we have more particularly referred are: Douglas and Hardy, Lancet, 1903; ii: 1540; T. R. Brown, Johns Hopkins Hospital Bulletin, 8; 1897; 79; Journal of Experimental Medicine, 3: 1898; 315, and Medical News, Philadelphia, 7: 1899; 12; H. Brooks, Medical Record, 59; 1900; 885; Kerr, Philadelphia Medical Journal, 6; 1900; 346; Balfour, Lancet, 1903; ii: 1649; Leichenstern, quoted by Ehrlich and Lazarus, Histology of the Blood, translated by Myers; Cambridge, 1900; 151; Boycott and Haldane, Journal, 71; 1900; 552, and American Medicine, 6; 1903; 391, New York Medical Journal, 71; 1900; 552, and American Medicine, 6; 1903; 391, New York

CHAPTER XI.

ad

wi

m

ac

on

or

"

an

br

ho

wj

an

it

as

ad

de

te

THE ENDOGENOUS INTOXICATIONS.

INTERNAL SECRETORY.

By Disturbance of the Internal Secretions.—An adequate recognition of the part played by the internal secretions of the economy has only come about during the present generation. Previous to this, attention had been almost wholly directed to the external secretions and the disturbances associated with or leading to alterations in excretory glands; this, although Claude Bernard, in his brilliant studies upon the liver and its glycogenic properties (1855–57), had demonstrated the existence and importance of the internal secretions, and, indeed, had given this name to substances which, formed through cell metabolism, become discharged, not externally, but into the blood and lymph; substances which, while from the point of view of the cells that form them they may be regarded as waste products, are by no means such for the rest of the economy, being essential to the proper carrying out of one or other function of one or other tissue, and affording most instructive examples of the mutual interdependence of the cells of the organism.

After Bernard's impressive demonstration that the liver cells convert into glycogen the sugar brought by the portal veins, and yield this glycogen (as sugar) to the blood, according to the needs of the muscle and other tissues, there came a long pause, broken in the late "eighties" by Brown-Séquard's bizarre campaign in support of injections of ovarian and testicular extracts as a cure for declining vigor. There is no doubt that these gland extracts have some tonic effect; but there is considerable doubt as to whether they have the specific effects claimed for them by Brown-Séquard and his followers. It has to be admitted that the use of organ extracts suffers even today from the suggestion of charlatanism, which inevitably accompanied the Brown-Séquard treatment. In the meantime the quiet work of other medical men and physiologists -Reverdin, Ord, Schiff, Horsley, Kocher, and others-was establishing the fact that the thyroid, a ductless gland, incapable, therefore, of affording an external secretion, played a very essential part in the organism, and that removal of this gland, or lack of development, or atrophy of the same, was followed by the appearance of a very remarkable train of symptoms-a train which might show itself as "cachexia strumipriva," cretinism, or myxœdema, respectively. The doctrine of the internal secretions may be said to have come into its own when George Murray demonstrated that injections of extracts of the healthy thyroid gland of the domestic animals causes the disappearance of all

INTERNAL SECRETORY

the distressing symptoms of myxcedema. Very soon it was shown that administration of the extract by the mouth is followed by equally good results; that the administration to healthy animals brings about the symptoms of hyperthyroidism, which in many respects resembles those of exophthalmic goitre, a condition which already, from histological considerations, Greenfield, of Edinburgh, had recognized as associated with overgrowth and overactivity of the gland. Next came the isolation by Baumann, of Freiburg, from the thyroid gland substance, of a compound protein, which he termed, first, thyroiodin, and later, and more appropriately, iodothyrin; he showed that this possessed the characteristic properties of the thyroid extract.

Here, then, has been afforded a full and scientific demonstration that disease is capable of being caused (1) by deficiency, (2) by excess of the specific internal secretion of a gland, or of particular constituents of the same.

While this is the most striking example, it is far from being the only one. We have evidence of one or other order, not merely of the development of internal secretions by glands, and those both ductless and affording external secretions; the indications, indeed, are now that the medulla of the adrenal bodies, which, although originating in connection with the sympathetic system, we are accustomed to regard as "glandular," and also the sympathetic ganglia in other regions afford an internal secretion. Nay, more, the valuable work initiated by Bayliss and Starling, upon the pancreatic secretion, is leading us to see that in a large number of cases certain portions of the mucous membrane of the alimentary tract and the cells of other organs afford hormones,¹ or internal secretions, which are necessary for the full activity of other glands at a distance. These hormones must not be confounded with enzymes; they are, for example, absolutely unaffected by boiling, and are soluble in alcohol.

Here we have to deal with the internal secretions as causes of disease; it would be out of place to describe *in extenso* the various morbid states associated with a disturbance in the internal secretions; at most, I can adduce briefly the evidence we possess associating these morbid conditions with such disturbances.

The Thyroid.—Myxœdema and Cretinism.—Myxœdema is a condition appearing in adult life in which there develops characteristically a loss of expression, associated with a thickening of the skin, or, more accurately, a subcutaneous infiltration of the face and body generally. At first this is due to a mucoid œdema (hence the name); later, this gives place to connective-tissue overgrowth. The skin is dry, the hair badly nourished, tending to drop out; the nose and lips become thick and bloated. With this the mental processes become slowed and undergo progressive failure, with a defective memory and, it may be, final dementia. Cretinism, on the other hand, is congenital; it is characterized by a striking retardation and imperfection of development.

¹ From δρμάω, I excite, or arouse.

The adult of forty is mentally an infant, often an imbecile, and his body retains infantile or childish features. Dentition is delayed; the sexual organs and functions do not attain maturity; the extremities are short and thick; the abdomen swollen; the features coarse and lacking expression. The evidence that these conditions are due to lack of thyroid secretion is:

1. Myxœdema, even after many years' duration, and cretinism, in the child, can be alleviated by administering thyroid extract. The cure



Cretin, male, aged twenty-one years, (Bourneville and Bricon.¹)

is not complete, *i. e.*, to preserve the normal state it is necessary to continue giving the extract from time to time.

 As first shown by Kocher,¹ symptoms identical with those of myxordema follow a complete removal of the human thyroid (cachexia strumipriva).

3. Histologically, in all cases of myxodema and cretinism we encounter either extreme atrophy or grave lesions of the thyroid.

It is interesting and, at first thought, paradoxical, that in a certain number of cases of both conditions we find enlarged thyroid, as also that upon operative interference in this latter order of cases symptoms of the very opposite condition—of exophthalmic goitre—are apt to show themselves. We have here, in my experience, examples of colloid goitre, *i. e.*, of great distension of the vesicles of the thyroid, with thickened, inspissated, firm, gelatinous secretion. The

normal thyroid is extremely vascular, a network of capillaries surrounding each vesicle, and these both hemic and lymphatic. I have suggested² that here, as in the expansion of the air-sacs in emphysema of the lungs, the distension of the vesicles results in such a flattening and compression of the vessels that circulation in and absorption of the thyroid secretion by these vessels is arrested, and symptoms of athyrea show themselves as a result. Acute congestion of the organ will thus lead to sudden absorption of large amounts of the secretion, and symptoms of hyperthyroidism manifest themselves. Similar considerations help to explain how overaction of the thyroid and exophthalmic goitre may give place to myxcedema.

> ¹ Arch. de Neurologie, 12: 1886: 137 and 292. ² The Practitioner, 64: 1900: 56.

Exophthalmic Goitre, or Graves' Disease (1835), **Basedow's Disease** (1840), **or Parry's Disease** (1825).—In this we have an unmistakable collection of symptoms: (1) Exophthalmos, or protrusion of the eyeballs; (2) tachycardia, or great rapidity of heart-beat and pulse; (3) enlargement of the thyroid; (4) tremor and nervousness. Flushing and abundant perspiration and increased pigmentation of the skin may also be present.

That this is caused by excessive secretion from the gland is shown by: (1) The cure of the disease by partial thyroidectomy; (2) the production of some of the most striking symptoms (tachycardia, tremors, and nervous irritability) by the administration of too large doses of thyroid extract to previously healthy men or animals (hyperthyroidism); (3) the increased nitrogenous output seen both in exophthalmic goitre and in hyperthyroidism; (4) the histological indications in typical cases of Graves' disease (as shown by Greenfield, and later by Halsted), of hyperactivity of the gland; the cells lining the vesicles are large; there are indications of overgrowth in the form of infoldings of the epithelium; the gland is found very vascular; the vesicular contents thin and fluid. The atypical form supervening upon colloid goitre has already been noted.

What is behind these states? What are the causes of athyrea and hyperthyrea? That is another matter. Here, for the time, we must be satisfied to recognize that these remarkable sets of symptoms are brought about essentially by defect and overproduction, respectively, of the internal secretion of the thyroid, without laying down what is the exciting cause of the defect or the overproduction. This statement holds in connection with all the other instances to be brought forward.

The Parathyroids.—Associated with the thyroid, either embedded in the lateral lobes or in their immediate neighborhood, are certain small bodies, the size of a pea or thereabouts. Of these there are usually a superior and an inferior pair. There is still debate regarding their functions. Some would regard them as rudimentary thyroid tissue,¹ but clearly they are functional, and their activity differs from that of the thyroid. Removal, in dogs, lends to muscular twitchings, giving place to tetany, exophthalmos, and rapid breathing, with death within a few days.² The symptoms are preëminently those of irritation of the nervous centres.

The brilliant observations of MacCallum and Voegtlin³ show that in the parathyroidectomized dog there is a rapid fall of the calcium salts to about half the normal amount, and that the intravenous injection of a calcium salt (the lactate or the acetate) almost instantly removes the violent symptoms produced by removal of these bodies—muscular twitchings, and rigidity, tachypnea, fibrillary tremors, increased rapidity of heart beat. It would seem thus that the parathyroids in some

¹ Kishi, Virch. Arch., 176: 1904: 260.

² MacCallum, Medical News, 1903: 820.

³ Johns Hopkins Hosp. Bull., 19:1908:91.

THE ENDOGENOUS INTOXICATIONS

way control the calcium metabolism, so that their removal is followed by a rapid excretion of the calcium salts. The observations further suggest that the toxic symptoms may be due to the unantagonized action of potassium salts upon the nerve centres, for in tetany the injection of potassium salts was found to intensify all the symptoms. It is doubtful, however, whether the parathyroids alone determine the calcium content of the tissues. Thus, in Kinnicutt's case¹ of tetany associated with gastric dilatation in an adult, parathyroid extracts had no influence, whereas the administration of calcium salts caused rapid and striking disappearance of the muscular twitchings and contractures. Here, also, after the death of the patient the parathyroids were found apparently normal.

Fro. 134

Normal skull.

Skull from case of acromegaly. (Osborne.)

It is still unsettled what is the exact causation of the *exophthalmos* of Graves' disease. It is not reduced by injections of parathyroid extract; nor, on the other hand, has it been satisfactorily reproduced by injection of thyroid extracts into normal animals. Here may be noted MacCallum and Cornell's observation² that where the dog's head is removed from the body (and so all influence removed of orbital congestion) stimulation of the cervical sympathetic results in an exophthalmos as pronounced as any produced in the living dog, and their demonstration that this is brought about by contraction of the so-called musculus orbitalis of Müller (1859), a case of smooth muscle and elastic fibrous tissue, enclosing the fatty bed of the eve, and having its apex posterior. They were unable to produce exophthalmos in the human being by sympathetic stimulation, so would leave the matter open; but Jonnesco,³ employing strong stimulation, had previously reported definite protrusion of the eveball. We are thus inclined to attribute the exophthalmos of Graves' disease to sympathetic irritation.

¹ Trans. Assoc. Am. Phys., 24: 1909: 475.

² Med. News, N. Y., October 15, 1904.

³ Thirteenth Internat. Med. Congr., Paris, 1900.

INTERNAL SECRETORY: PITUITARY

The Pituitary Body, or Hypophysis Cerebri.-Ever since the observations of Marie it has been known that there is a close relationship between disorders of the pituitary body and the development of the remarkable condition of overgrowth more particularly of the bones of the face and extremities known as aeromegaly. This condition develops in adult life, and is of slow development, over, it may be, twenty or more years. The face becomes enlarged, the superior and inferior maxillary bones being especially involved; the ears become of great size; the nostrils broaden; the eyelids thicken. The hands and feet are characteristically hypertrophied, becoming disproportionately large. Later, the spinal cord may be affected. In general *qiantism*, also, enlargement of the pituitary, has frequently been noted. Marie was of the opinion that loss of the active pituitary secretion induced the acromegaly, with replacement of the pituitary tissue by new growth. Schäfer (to whom, with his associates, Oliver and Herring, we owe the most recent advances in our knowledge) holds with Tamburini and Woods Hutchinson that the overgrowth is due to overgrowth of the anterior lobe; for in some cases a simple hyperplasia of this region has been noted; von Hansemann¹ points out that of tumors it is only adenomas (i. e., overgrowths of thisanterior lobe) and not sarcomas or colloid accumulations that are found associated with the disease. Schäfer² suggests that Marie's objection may be got over by supposing that, in the cases cited by Marie, the condition began by simple hyperplasia, the destructive new growth being a terminal event.

But if the indications are that this anterior glandular part supplies hormones or other products which stimulate the growth of bone and connective tissues, the observations of Herring demonstrate that the intermediate part, also glandular, develops colloid matter which finds its way through the posterior nervous portion of the organ into the infundibulum, and so into the ventricles of the brain. And this internal secretion, produced in these two portions, has different but equally striking properties, which, as Howell has shown, are not possessed by extract of the anterior part. Namely, as shown by Oliver and Schäfer, an extract of these portions causes a rise of blood pressure very similar to, but more prolonged than, that induced by adrenal extract, and, in addition, an increased discharge of urine-a polyuria which, further, is seen to be independent of the blood pressure and due to direct action on the renal tissue. Like adrenin this extract is unaffected by boiling. The indications are that the gland provides not one but several hormones.

Nor is this all. Like the adrenal, this minute gland (weighing in man only half a gram) is essential to life, and, as shown by Paulesco³ and by Reford and Harvey Cushing,⁴ its ablation in animals of the

² Croonian Lecture, Proc. Roy. Soc., B., 81: 1909: 442 (with bibliography).

⁴ Johns Hopkins Hosp. Bull., 20: 1909.

¹ Descendens und Pathologie, Berlin, 1909; 197.

³ Jour. de Physiol., 9: 1907.

THE ENDOGENOUS INTOXICATIONS

laboratory, is followed by death within forty-eight hours or so, with symptoms of acute inanition. What is the essential cause of this death is not known. It is, however, most significant that this minute mass of substance, mainly glandular, has this extraordinary influence over continued existence.

The Adrenals and the Chromaffin System .- Between the adrenals and the pituitary body an interesting parallelism is to be observed: both consist of a glandular portion and one of nervous origin, and, according to Gaskell,¹ these glandular portions are homologous, both derivatives of the primitive body cavity, and, therefore, mesothelial, being remnants of the primitive segmental excretory system. The medulla of the adrenal is essentially a derivative of the sympathetic system; the posterior portion of the pituitary, while not containing any neurons, is contributed by the infundibulum, and in the cat still exhibits the central canal of this part of the brain. Just as the glandular portion of the pituitary has been found associated with the growth activities of the skeletal tissues, so do we possess a series of observations that hyperplasia and simple tumors of the adrenal *cortex* may be accompanied by excessive and premature obesity, precocious muscularity, as in the rare cases of "Infant Hercules," and, more particularly, premature virility, with precocious development of the external genitalia.² So, also, just as the posterior nervous portion of the pituitary supplies an extract causing raised blood pressure, adrenin,³ extracted from the adrenal medulla, possesses to a striking degree the same properties.

The production of adrenin has been found to be associated with the presence of a particular order of cells—the *chromaflin* cells—so-called because of their affinity for chrome salts, these salts being retained, and the cells exhibiting a strong yellowish-brown color after immersion of the material in solutions of potassium bichromate.⁴ They are by no means confined to the medulla of the adrenal, but have been found in considerable numbers in the abdominal and other sympathetic ganglia, in the minute carotid body, situated on the anterior wall at the bifurcation of the common carotid, in the coccygeal gland, etc. We are brought, then, face to face with the fact that there exist cells of nervous origin with functions and characters widely distinct from those of ordinary nerve cells. The indications are that wherever these cells are present there adrenin, or a body having like effects upon the arterioles and the blood pressure, is also present.

As Addison, of Guy's, was the first to point out (1855) the significant syndrome of bronzing of the skin, progressive muscular and bodily

¹ Origin of Vertebrates, Longmans, 1908; 340.

² For a collection of cases and literature see Guthrie and Emery, Clinical Soc. Trans., 40: 1907. See also Bulloch and Sequeira, Trans. Path. Soc., Lond., 56: 1905; 189.

⁸ Schäfer urges the use of this term, rather than adrenalin, epinephrin, etc., the latter being the names of proprietary substances.

 4 Material that has been hardened in Orth's solution (Müller's fluid plus formalin) shows them well when cut in the ϵ rdinary manner and stained with hematoxylin and cosin,

INTERNAL SECRETORY: ADRENAL

weakness, with feeble pulse and heart action, nausea, vomiting, or other indications of gastric disturbances, is repeatedly found at postmortem to be associated with lesions involving the greater part of the adrenals, most often with replacement of the tissue by large fibrocaseous tubercular masses, less often with atrophy, or chronic interstitial fibrosis and atrophy, thrombosis of the adrenal veins, or with malignant growths in these organs. Nevertheless, long years before adrenin and chromaffin cells and their relationship to the sympathetic system had been discovered it was recognized that this peculiar disease might be present without there being any naked-eye or microscopic evidence of disease of the adrenal glands, and Rolleston and others had in one group of these cases called attention to the existence of chronic inflammatory and allied changes involving the abdominal sympathetics. Contrariwise, other cases are encountered in which, with practically complete destruction of both adrenals, as by secondary cancer, the symptoms of Addison's disease have been wholly wanting.¹ These apparently discordant findings have in the past been productive of much debate and uncertainty. It was soon discovered that removal of the pair of organs in the lower animals (here, again, as is the case with the pituitary) is almost always followed by rapid inanition and death. The rat alone frequently survives such removal. Marchand has supplied an important clue to these exceptions by his demonstrations of the existence of accessory adrenalsof adrenal tissue not merely embedded in the kidney (as is not infrequent), but carried down by the ovaries and testes in their development and descent. In the rat this accessory tissue is both constant and relatively abundant. This, however, is incapable of explaining all the cases, and the discovery of the chromaffin tissue and its relationships has elucidated matters to a very great degree. In Addison's disease we deal with disease of the chromaffin system rather than of the adrenals only.

I am very far from suggesting that all the problems in connection with the internal secretion of the adrenal and the pituitary are now solved. There are still many moot points. What, in the first place, is the significance of the intimate association between the glandular and nervous elements? In the case of the adrenals this is a secondary acquirement, for in certain lower vertebrates the glandular and nervous elements form distinct and separate organs. In the pituitary it is seen that the colloid secretion of the glandular part passes into and through the nervous portion. Does the adrenal cortex elaborate something from the blood which by the medulla is converted into adrenin; or, conversely, does it remove from the blood the substrate or material upon which the medullary substances have a specific action. It is at

¹ Lewin, in the study of 370 cases of typical Addison's disease, found:

Addison's disease with diseased adrenals, 88 per cent.

Addison's disease with sound adrenals, 12 per cent.

Diseased adrenals with bronzing, 72 per cent.; without bronzing, 28 per cent. He had also collected 44 cases in which the adrenals were found destroyed by various diseases in those who had shown no Addison's disease during life.

THE ENDOGENOUS INTOXICATIONS

least suggestive that, as pointed out by Pearce and other observers, a marked enlargement of the adrenals is frequently to be noted in cases such as those of chronic interstitial nephritis characterized by high blood pressure. Our own experience at the Royal Victoria Hospital is that in these the overgrowth of the cortex is even more marked than that of the medulla. This brings us to another point. It is a common mistake to regard the inner pigmented zone of the adrenal as medullary; it is nothing of the kind; it is the innermost zone of the cortex that thus contains pigment in its cells. And now adrenin has been isolated (Takamine, Abel, etc.¹), and even synthesized (Dakin), so that its chemical composition is fairly well ascertained. It is related to pyrocatechin and the aromatic products of proteid dissociation. Here, curiously, we find a series of independent observations converging suggestively upon one point. For, as will be noted more fully in the chapter upon pigmental changes, it is found, on the one hand, that the most important group of non-hematogenous pigments, the melanins, are allied to tyrosin and the aromatic group of proteid derivatives, and, on the other, Halle² claims to have demonstrated that under the action of an enzyme contained in the adrenal, tyrosin is converted into adrenin. As I suggested in the first edition of this work, it may, therefore, be surmised, in the first place, that the adrenal cortex takes up tyrosin and other chromogenic compounds from the circulating blood, and that these are converted into the adrenin of the medulla; and, in the second place, that the pigmentation of Addison's disease is due to a deficient conversion of these chromogenic substances, owing to the destruction of the adrenal, with its enzyme. These are the indications; the full proof is still wanting.

Another matter not yet fully worked out is the relationship and mutual antagonism of the thyroid, the adrenal, and the pituitary-as, indeed, of all the organs affording internal secretions. We have shown that internal secretions of the latter two here mentioned favor growth and proliferation of particular tissues. The thyroid secretion, on the other hand, favors increased oxidation and dissociation. As is generally known, properly applied, it reduces obesity. Nevertheless, we have the unexplained paradox, to which several workers have borne testimony, that ablation of the thyroid is followed by enlargement of the pituitary and increase of colloid in its vesicles. The only observation encountered by us that possibly throws light upon this apparent contradiction is that of Schäfer, when he found evidences of the existence in pituitary extract of opposing pressor and depressor substances, the one causing rise, the other fall, of the blood pressure. Possibly, according to circumstances, the gland may be activated to produce an excess now of the one. now of the other. In this association it may be noted that several observers (Hart, Hedinger, Wiesel, Crowell³) have called attention to the co-existence of hyperplasia of the thymus and the status lymphaticus

¹ Proc. Roy. Soc., B., 76: 1905: 491.

² Hofmeister's Beitr., 8: 1906: 276.

³ Proc. New York Path. Soc., N. S., 9: 1909: 75.

INTERNAL SECRETORY: OVARIES AND TESTES

with Addison's disease. Falta and others see a relationship between pancreatic disease and diabetes and hyperactivity of the chromaffin system. A full explanation of these relationships cannot yet be given.

Lastly, there deserves notice a striking fact, to which Langley and Starling call attention, namely, that adrenin does not merely act upon the smooth muscle fibres of the vessels, but would seem necessary for the normal functioning of the whole sympathetic system. The effects of adrenin upon a part are identical with the results of stimulating the sympathetic fibers distributed to that part.

These studies seem to be leading us toward a recognition of the chemical, and not merely a physical, basis of nerve excitation. However, the fact that adrenin directly excites muscle must make us proceed very cautionsly in reaching any such conclusion.

Ovaries and Testes.—It is a matter of familiar knowledge that capons and other castrated animals, in which the active organs of generation have been removed prior to puberty, differ materially in build and character from "entire" animals of the same sex and species; the external and secondary sexual features are imperfectly developed, the vocal organs do not reach adult development, the general bodily build approximates more to that of the other sex. Is this due to lack of nerve stimuli proceeding from the ovaries or testes to the higher centres of the brain, to lack of removal from the blood of substances necessary in the elaboration of the ova and spermatozoa; or, lastly, to absence of an internal secretion? It cannot be said that we know the complete answer; it is, indeed, difficult to see how it can be obtained by experimental means.

The full demonstration that these changes are due to lack of internal secretion must be obtained by removing ovaries and testes and then showing that ovarian or testicular extracts lead to full development of one or other sexual type. We deal here with growth. Now, normally, if this be due to the influence of internal secretions, these secretions must be continuously elaborated in small quantities over long periods of time. Experimentally, we cannot reproduce the process; we can only make periodical injections of uncertain quantities. At most, using these faulty methods, there is a possibility of obtaining approximate results, and this has been done in part.

Thus, Ancel and Bouin' have proved clearly that the secondary sexual characters in the male are due to the internal secretion of the interstitial cells of the testes—certain faiely large cells lying in the stroma, between the tubules. Removg the testes entirely, and the secondary characters do not develop. Ligature the vas deferens, and so bring about atrophy of the seminal tubules, the interstitial cells still persist and the secondary characters make their appearance. It is not, then, the secretory tubules of the organ that are responsible, but these interstitial cells, which, it may be added, are embryologically of like origin, although, as a result of position, their function becomes widely different. Shattock has confirmed and extended these observations.

¹ Compt. rend. Soc. de Biol., 55: 1903; 1397; and 56: 1904: 81.

THE ENDOGENOUS INTOXICATIONS

In the ovary there are homologous interstitial cells, derived, as shown by Miss Lane-Claypon,¹ from the common germinal epithelium, which gives rise indifferently to follicular epithelium, ova, and the intermediate cells. The indications are that these cells have equally important influence upon sexual life and growth, although, confessedly, it is difficult to distinguish between their effects and those of the follicular epithelium. Menstruation and "heat" depend upon the presence of the ovaries. It has been shown by Fränckel² and by Marshall and Jolly³ that, in castrated animals, heat can be produced by the injection of ovarian extracts.



Cells in ovary of young rabbit, derived from the germinal epithelium (a), which give rise to (b)primordial ovum; c. multinucleated interstitial cell. and d, interstitial cell becoming isolated; f, connective tissue; g, modified germinal cells. (Lane-Claypon.)

The corpus luteum has functions of another order. This develops rapidly after discharge of the ovum. These observers have shown that if the ovaries of an animal be removed within six days after coitus the ova do not become adherent to the uterus. Fränckel obtained the same results by destruction of the corpus luteum alone. To the internal secretion of the follicular cells of this little organ must be ascribed the stimulation of the uterine muccos so that it responds to the presence of the ovum and develops the maternal sections of the placenta and decidua. It is the interstitial cells lying external to the Graafian follicles that form, it would seem, the main cell layer of the corpus luteum and that supply the hormone.

The Fœtus, and its Bearing on the Growth of the Mammary Gland.—That the development of the mammary gland during pregnancy is not due to nervous reflex has been shown by more than one observer. The whole lumbosacral cord may be extirpated in a pregnant bitch, but pregnancy will continue, the gland enlarge, and the bitch suckle

¹ Proc. Roy. Soc., B., 77: 1905; 32.

² Arch. f. Gynäk., 68: 1903; Pt. 2,

³ Phil. Trans., B., 198:1906:99.

INTERNAL SECRETORY: THE FETUS

its pups. Ribbert transplanted a mammary gland in the guinea-pig to the region of the ear. With pregnancy, the gland underwent hypertrophy, and when the young were born milk could be expressed from it. Lane-Claypon and Starling' have shown that the stimulus to the hypertrophy of the gland comes not from ovaries, or uterus, or placenta, but from the developing foetus. Injections of watery extracts from rabbit foctuses into a virgin rabbit every one to three days, over a period of three weeks, led to the glands, which were at first almost invisible, becoming markedly hypertrophied, with enlarging ducts and epithelium, and discharge of a thin fluid; in multiparous rabbits, similar injections led to the discharge of true milk. They draw the conclusion that the amount of the substances leading to the glandular hypertrophy is greatest during the latter part of pregnancy, and lactation would seem due, we would add, primarily to the removal of these substances, the cells which had under its influence manifested anabolism and growth, in its absence proceeding to break down, and so form milk. These observations would appear to explain the cessation of lactation with the onset of a renewed pregnancy.

Eclampsia.—We are still wholly at a loss where to place eclampsia, that most dangerous combination of disturbances which may result from pregnancy. The kidneys are profoundly involved; indeed, the danger signal is found in the appearance of albuminuria during the last weeks of pregnancy; there is a relatively large proportion of globulin with low urea content, suggesting a relationship of the condition to uremia, which is strengthened by the convulsions and coma, leading to death. But anatomical studies show disturbances elsewhere, and notably in the liver, which are wholly foreign to the uremic state. The liver changes are remarkable; in typical cases there occur abundant necroses in this organ; where slight, these recall the focal necroses seen in many infectious and toxic conditions, but they may be so extensive as to give the appearance of that wide destruction of the hepatic parenchyma seen in acute yellow atrophy. Schmorl and others have ascribed these to liberated placental cells becoming blocked in the portal vein and hepatic arteries, but, while placental cell emboli do occur, they are incapable of explaining the widespread eclamptic disturbances. No positive or consonant results have been obtained from bacteriological studies, and the tendency nowadays is to regard the condition as an intoxication set up by abnormal or excessive substances diffusing into the maternal blood from either the placenta or the foetus. Inasmuch as the conditions may first become acute, it may be some days after the fortus has been removed, the tendency on the part of many is to regard the placenta as originating the irritant, and this view gains some support from Hitschmann's² case, in which eclampsia showed itself in a woman, the bearer of a placental mole (a mole is an aberrant growth of the chorionic villi placenta, the embryo having died or aborted).

¹ Proc. Roy. Soc., B., 77: 1906: 505.

² Centralbl. f. Gyn., 28; 1904; 1089, Quoted by Wells.

Opposed to this view is the fact that extract of placental tissue produces singularly little disturbance when injected, although Liepmann states that eclamptic placentas are definitely toxic. This awaits fuller confirmation. The most that can at present be said is that it is along these lines of investigation that attention is at present being directed, and that the tendency is to recognize two orders of cases, in one of which the main histological lesions are hepatic, in the other, renal.

The Digestive System.-A great step forward in the comprehension of the internal secretions and their mode of action was made by Bayliss and Starling in their already classical observations upon the secretion of pancreatic juice. Everyone should be familiar with Pawlow's1 notable advances in the study of digestive secretions. His studies established the existence of nervous reflexes as setting in action the flow of, more particularly, the gastric juice, just, as years previously, Ludwig had shown the nervous paths whereby salivary excretion is brought about. Bayliss and Starling went farther: studying the increased flow of pancreatic juice which follows the introduction of acid into the duodenum (it must be recalled that the chyme when it enters the small intestine is acid), they found that this occurred even when pancreas and duodenum were separated, and even when the acid was introduced into a loop of the upper jejunum that had been deprived of all nervous connections. The reaction could only be chemical, and further study demonstrated that, by scraping off the mucous membrane of the duodenum and upper part of the small intestine, pounding this with sand, and adding 0.4 per cent. HCl, they could, by boiling and neutralizing the fluid, extracting with alcohol, and evaporating the latter, obtain a substance, secretin, which, dissolved in water and injected into the veins of a mammal, leads to an abundant excretion of clear pancreatic juice. That acid in the small intestine leads to discharge of secretin from the mucous membrane into the blood, by which it is carried to the pancreas, stimulating that organ, has been further demonstrated by Wertheimer; establishing a cross-circulation between two animals, he found that acid introduced into the duodenum of the one led to increased secretion of pancreatic juice in the other. The indications are that, while there exists a nervous mechanism for the secretion of the digestive juices, as shown by the mouth watering at the sight, smell, or thought of food, and by Pawlow's demonstration of the pouring out of "appetite juice" in the stomach, there exists also a series of secretins developed by the mucous membrane of one segment of the alimentary canal after the other, which, passing into the blood, stimulate the specific glands of a neighboring segment. Thus, Edkins has proved the existence of a gastric secretin elaborated by the pyloric glands, leading to the discharge of hydrochloric acid from the cardiac end of the stomach. Hemmeter describes removal of the salivary glands as leading to dyspepsia and arrest of the cardiac secre-

¹ The Work of the Digestive Glands. Translated by W. H. Thompson, London, Griffin, 1902.

INTERNAL SECRETORY: DIABETES

tion.¹ Bayliss and Starling found that the duodenal secretion also influences the flow of bile. In other words, the various forms of indigestion, dyspepsia, absence of pancreatic juice, of biliary excretion, and of digestion in the lower portion of the small intestine, may, in the absence of gross obstruction and disturbance, be due to imperfect development of one or other secretin, or hormone. We thus open a new chapter of the pathology of digestion.

It may be added that, like adrenin and iodothyrin, and unlike toxins and enzymes, the secretins are not modified by heat, nor again by alcohol, in which they are readily soluble.

The Pancreas, the Liver, and Diabetes Mellitus.—As shown by von Mering and Minkowski (1889), complete removal of the pancreas in the dog is followed within twenty-four hours by glycosuria, by the appearance of abundant sugar in the urine, and this not temporarily, but of the diabetic type, *i. e.*, it persists. There may, in a few days, be as much as 8 to 10 per cent, and this although the food contains no carbohydrates; and, as in diabetes mellitus, in many cases, the animal exhibits increased appetite and excessive thirst, with abundant discharge of urine; progressive asthenia and emaciation show themselves, and the urine comes to contain acetone, diacetic and oxybutyric acids. The phenomenon is not confined to dogs, but results in all vertebrates, even down to eels (Capparelli), when the greater part of the pancreas—seven-eighths and more—is removed.

That the ordinary external secretion of the organ plays no part in the sugar regulation is evidenced by the facts: (1) That glycosuria does not supervene when the pancreas is transplanted into the abdominal wall (Lépine), and (2) that no diabetes occurs when a pancreatic fistula is made, draining away all the secretion as soon as it is elaborated.

There are thus the alternatives: (1) That the organ furnishes an internal secretion, affording something of the nature of a glycolytic ferment, in the absence of which the sugar is not transformed, but accumulates in the blood, and (2) that under normal metabolism there is produced a substance which hinders the above transformation, this substance being taken up and destroyed by the pancreas. When this organ is removed or diseased, the substance in question accumulates in the system and glycosuria results.


Beyond this, for long years, despite abundant experiments, no advance was made, no extract could be obtained from the pancreas having an action upon the sugar in the blood. It is true that Lépine, Bail, and others demonstrated the existence of glycolysis in normal blood, and that the former found this absent in cases of diabetes, but these observations could not be confirmed. Only in the last few years has Otto Cohnheim² afforded what appears to be the correct solution. Muscle and liver, it will be remembered, normally contain glycogen. Cohnheim expressed the juice from pancreas and muscle; adding each sepa-

¹ This is still a matter of debate.

² Zeitsch, f. Physiol. Chem., 39: 1903; 338; and 42: 1904; 401.

rately to a solution of glucose, there was no reaction; but when to the muscle juice he added a small quantity of the pancreatic extract, there followed a rapid conversion of the glucose into alcohol and CO_2 . Rahel-Hirsch independently announced the same results, as also that the other tissue, the seat of active carbohydrate metabolism, viz., the liver, gave corresponding results, the difference being that liver juice normally is able to act to some extent upon glucose; if, however, a little pancreatic extract be added, the decomposition is greatly accelerated.

Now, this substance produced by the pancreas withstands boiling, and is soluble in alcohol without loss of acivity; it has properties allied to those of adrenin and the hormones in general. We thus arrive at the conclusions: (1) That the normal pancreas affords a body of the



Section of island of Langerhans from pancreas of human adult. \times 200, (Dewitt.)

nature of a secretin; (2) that the liver and muscle are the main seats for the deposit and utilization of carbohydrates; (3) that these two tissues can convert soluble carbohydrates into insoluble glycogen, and *vice versa*; (4) that unaided they cannot to any extent break down these carbohydrates; (5) that for glycolysis to ensue, these tissues need to be activated by a hormone elaborated in the pancreas. Just as the pancreatic juice is unable to act upon proteins alone, but requires the addition of enterokinase before the trypsin is complete, so this hormone of pancreatic origin is necessary before the full glycolytic ferment is evolved in liver or muscle.

Were this all, our comprehension of the causation of diabetes would be easy, but there are many riddles still remaining to be solved. Dia-

betes in man, in the first place, is not always associated with pancreatic disease. In most cases the organ or its specific cells are greatly reduced through congenital aplasia (rare), interstitial fibrosis with atrophy of the glandular elements (the commonest cause), cancerous growths, etc.; but in the majority of these the reduction does not approximate to that which experimentally is found necessary to produce glycosuria. Opie¹ and, later and independently, Ssobolew² have laid down that it is not the pancreatic tissue in general, but the remarkable little "islands of Langerhans" scattered through the organ that are at fault; that these act as ductless glands (elaborating, we would now say, the hormone), and that diabetes is due to degeneration of the same. And certainly, as Opie has shown, in many cases there exists a form of hyaline degeneration of these bodies. But Dale,³ working under Starling, and, more recently, Swale Vincent and Mrs. Thompson,4 have shown conclusively that the islands are not separate organs; that they vary in number according to the state of nutrition and activity of the gland, becoming converted into active acini, and vice versa. We have encountered appearances in man difficult to explain save along these lines. These observations do not, it is true, wholly negative the contention that the islands bear some intimate relationship to a certain order of cases of diabetes; they suggest, however, that degenerative changes seen in them are an indication of other changes occurring in the intimately connected pancreatic tissue proper.

Then, again, Naunyn, Pavy, and others call attention to an hepatic form of the disease, in which, more particularly, the liver is enlarged and hyperemic; again, experimentally, by the exhibition of phloridzin, without there being any hyperglycemia, or excess of sugar in the blood, the urine comes to contain large quantities of sugar, and Zuntz has shown that if phloridzin be injected into one renal artery, the urine from the corresponding kidney contains sugar, that of the other remaining wholly free therefrom. Evidently, also, there is an adrenal glycosuria (Blum, Herter, etc.) produced by subcutaneous, intravenous, or peritoneal injection of adrenin, or even (Herter) by painting adrenin over the pancreas. Herter would explain this by the pronounced reducing powers of adrenin, whereby the function of the pancreatic cells becomes arrested; other reducing agents, he found, have the same effects.

Lastly, alimentary and nervous glycosuria have to be taken into account. Excessive consumption of carbohydrates is apt to be followed by the appearance of sugar (glucose) in the urine, while injuries to the head and neck, organic lesions of the brain, cerebral hemorrhage.

¹ Bull. Johns Hopkins Hosp., 11: 1900; 205, and Jour. Exp. Med., 1901; 397 and 527.

² Virch. Arch., 168:1902:61.

³ Phil. Trans., B., 197; 1905; 25. This view was first propounded by Lewaschew, Arch. f. mikr. Anat., 26; 1886; 453.

⁴ Proc. Physiol. Soc., Jour. of Physiol., 34; 1906, ibid., 35; 1907, and Trans. Roy. Soc., Canada, 1907.

cerebral tumors, more particularly involving the pons, cerebellum, medulla and posterior fossa, recall Claude Bernard's well-known "piqure" experiment, in which he established a transient glycosuria by puncture of the floor of the fourth ventricle between the vagus and auditory centres.

0

k

a

The causes and forms of glycosuria are, therefore, numerous, and no single theory, so far, suffices to explain all cases.

Conclusions.—Nevertheless, we think it possible to harmonize and bring into a common scheme the greater number of these diverse facts, and what is here said applies not merely to diabetes, but to all the group of conditions caused by disturbances in the discharge of the internal secretions.

1. As will be laid down more fully later,⁴ it must be recognized that if, in general, the function of the nervous system is to control and coordinate the various bodily functions, there are conditions under which it can stimulate one or other function in excess. Your drunken coachman may whip up the horses to a reckless and wholly unnecessary gallop. We must be prepared, for example, to find that glycosuria may present itself in the absence of any primary disturbance in the organs concerned in carbohydrate metabolism.

2. The elaboration and discharge of internal secretions has its limits. It is, therefore, possible, to have an intake or production of the substance acted on by those internal secretions, over and above the capacity of the internal secretions to convert or neutralize them, and this, again, without primary disturbance of the organs producing those internal secretions. An alimentary glycosuria is thus to be expected in case of excessive intake of carbohydrates.

3. The morbid states here discussed are not the outcome of one, but of the interaction of at least two factors; they represent a want of balance between amount of internal secretion and amount of the substance upon which that acts. These two must always be kept in mind. Myxœdema, we find, is the series of signs and symptoms associated with insufficient discharge of thyroid secretion; but if animals with removed thyroids be kept on a vegetable diet, and so the substance or substances be not elaborated which, under ordinary conditions, are neutralized by the thyroid secretion, then, notwithstanding the absence of the thyroid, no symptoms of myxœdema show themselves. The same symptoms may be brought about:

(a) By diminution of the internal secretion in the presence of normal production of the substratum or substance upon which it acts; and

(b) By no diminution in the amount of internal secretion elaborated and discharged, but by excess of the substratum upon which it acts.

In the first of these cases we may expect to find morbid changes in the organ affording the secretion; in the latter, none; in the first, the tissues, if any are known, affording the substratum may be perfectly normal; in the second, they may be diseased. In other words, we may

¹ In the discussion on Fever and Nervous Pyrexia.

INTERNAL SECRETORY

expect to find an identical syndrome in two cases, which, from the point of view of morbid anatomy or histology, differ widely.¹ That this must be so appears to be too often overlooked.

4. Where, as would seem to be the case in connection with glycolysis, we have the additional element of a hormone, the case becomes yet more complicated. The same syndrome it would seem, may be set up (a) by excessive development or intake of the substratum; (b) by lesion of the organ or organs in which that substratum undergoes disintegration preventing that disintegration; and (c) by lesion in the organ affording the hormone, without which the disintegration cannot be effected.²

¹ Adami, The Internal Secretions, Trans. Congr. of Am. Phys. and Surg., 4:1897: 103.

^a For fuller study of the pancreas and of diabetes mellitus, the reader is recommended to Opie, *The Pancreas*; Williamson on *Diabetes*, and Futcher's article on *Diabetes* in Osler and McCrac's *Modern Medicine*, vol. i. Starling's Croonian Lectures, Lancet, London, 1905; i and ii, give an admirable grasp of the trend of recent investigations on the internal secretions, as does the later address by Howell, Scientific American Suppl., 1797; June 11, 1910; 375.

CHAPTER XII.

fu

81

ge

m

te

11

ns

de

se

an

de

up

mi

sh

ret

ble pa

THE ENDOGENOUS INTOXICATIONS—(CONTINUED).

DISINTEGRATIVE INTOXICATIONS.

Autolysis.—There will be frequent occasion to refer to autolysis in discussing the various morbid processes and the part played in them by the self-digestion of the tissues; there is less to say regarding this action as a cause of morbid states. Nevertheless, to make ourselves clear, it is necessary to explain here in a little detail what we understand by this term, and what we know regarding the extent of the process.¹

It has long been known that proteolytic ferments are liberated in certain forms of cell disintegration. Thus, Leber, in the "eighties," showed that aseptic pus produced around copper filings had the power of dissolving white of egg and fibrin. Filehne, in 1877, Stolnikow, Fr. Müller, and others long ago obtained proleolytic ferments from gangrenous and pneumonic sputa. But the wide extent to which the tissues in general are capable of self-digestion (or autolysis) only became recognized after the publication of the works by Salkowski (1890), and more especially of Jacobi (1900). Hauser had previously noted that liver tissue, kept in a perfectly aseptic condition, underwent softening, the cells breaking up and becoming granular. These observers studied the phenomenon from its chemical aspect, and demonstrated that the process was of the nature of a proteolytic change. They showed that it still progressed when chloroform or toluol was added to the broken-up tissue, whereby bacterial activity was with certainty prevented and when all that could happen was the result of enzyme action; as, also, that all the soft tissues of the body undergo this autolysis, although at varying rates, the softening and disintegration of normal liver substance being most active, that of renal cortex the next; skin and brain substance (contrary to expectation) being among the slowest to be affected.

Very many different enzyme actions evidently take place in this process; that which is the most evident is the conversion of protein from an insoluble to a soluble state. In fresh liver tissue, 90.4 per cent. of the nitrogen in a given example, which had been boiled and kept for several days, was found to be in an insoluble form, 9.6 soluble. An emulsion from the same liver after digesting itself for twenty-two days, and then being boiled, afforded only 39.4 per cent. of insoluble

¹ The fullest and clearest résumé in English upon this subject is by Wells (Chemical Pathology, p. 86 et seq.), who himself has made material contributions to the study.

DISINTEGRATIVE INTOXICATIONS

nitrogen; 60.6 per cent. was soluble and contained in the filtrate.¹ But the nucleoproteins are broken down by nucleases; the purin bases are liberated, and, in their turn, acted upon by guanase and adenase; the fats are split up and fatty acids liberated, presumably by the action of lipase; the glycogen gives rise to glucose, and this, in turn, undergoes further splitting, being absent from the products of long-continued autolysis; the phosphatides are split up, and jecorin and allied bodies make their appearance; while, further, there is a marked increase in cholin and in cholesterin. What is interesting is that there may be, as shown by Hildesheim and Leathes, a marked increase in the fatty acids, suggesting, at first, that these are developed from proteins; the probability is that a large proportion of the fat in organs like the liver, kidneys, and muscle exists in a masked condition—whether in a state of loose combination with the proteins, or how, has not surely been determined.

This process proceeds most rapidly at, or slightly above, the body temperature, and in a slightly acid medium. Wiener has shown that the process does not begin until the normal alkalinity of the tissues has been neutralized by the production of organic acids (lactic, butyric, etc.). which takes place in all dying tissues. We deal, that is, not with ordinary tryptic digestion (which proceeds in an alkaline medium), although the products of the proteid cleavage are of the same type; nor do we deal with peptic digestion, for they are far more advanced than those effected by peptic activity. For us, what is of immediate importance, is that, as demonstrated by Jacobi and others, this process of autolysis can take place in the living organism; cut off the blood supply to a section of the liver and the central part softens and affords leucin, tyrosin, and the other cleavage products found in autolysis outside the body. The outer part does not show change to the same extent, autolysis being evidently arrested by the diffusion of the alkaline lymph. Thus, it is only in the more central portions of relatively large areas of necrosis or cell death that autolysis manifests itself-save where there is invasion by leukocytes. As shown by Opie and others, the leukocytes possess enzymes, proteases, which differ markedly from those of the tissues; those act mainly or almost entirely upon the cells in which they are developed, but, once liberated, the leukocytic enzymes act indifferently upon various tissues. These properties explain the softening of septic infarcts and of the outer zone of simple infarcts, as the result of the migration of leukocytes into them; they explain, also, as Fr. Müller has shown, the softening and absorption of the consolidated exudate in the pneumonic lung, that exudate being, in the stage of gray hepatization. little more than a dense mass of leukocytes.

There is, however, another factor, as determined by Opie,² which regulates the autolytic action of leukocytes, viz., the presence in normal blood serum of an antibody, neutralizing the leukocytic protease. In pathological exudates the amount of this antibody exhibits variation;

¹ Quoted from Wells, loc. cit.

² Journal of Experimental Medicine, 7:1905:316, and 8:1906:410,

thus, an exudate which, although containing abundant leukocytes, is at first fibrinous, may eventually show disappearance of that fibrin through diminution of the antibody and unrestrained digestive activity of the enzyme liberated from the leukocytes. Jacobi would distinguish this solution of dead matter by the agency of wandering in leukocytes as *heterolysis*.

a

81

W

W

st

St

a1

ki

There are, however, conditions under which autolysis-not heterolysis-may obviously manifest itself in individual cells or groups of cells. The conditions under which the process manifests itself are not wholly understood. The liver, again, affords the clearest example, and that in a curiously assorted group of cases-acute yellow atrophy, phosphorus and arsenic poisoning, chloroform necrosis, and, to a slighter degree, in the pernicious vomiting of pregnancy.¹ In all these cases we obtain histological evidence of death of cells throughout the organ, others in the immediate neighborhood still retaining their vitality. The death is not immediate, but is preceded by indications of profound nuclear and cytoplasmic disturbance; it leads to a breaking down and disappearance of the affected cells. Coincidently, as pointed out by Salkowski, the liver tissue and the urine come to contain the products of autolysis-leucin, tyrosin, etc. In the livers of the cases of phosphorus poisoning, Waldvogel and Tintemann² obtained increased amounts of cholesterin, jecorin, neutral fats, protagon, and fatty acids, with diminished lecithin, just as in the earlier stages of experimental autolysis. With Wells, we must conclude that in all these cases we have a death of the individual cells without inhibition or destruction of the intracellular enzymes; that just as chloroform, for example, kills bacteria, but leaves their ferments unaffected, so it acts also when exhibited in excess to the cells of the organism.

A partial explanation of this intravital autolysis is probably to be found in the coincident condition of acidosis. As already noted, the autolytic enzymes act best in a slightly acid medium; an alkalinity equal to 0.04 per cent. NaHO arrests their action.³ Now, as will be pointed out (p. 381), it is in just this group of conditions that the alkalinity of the blood is diminished and acetonuria is apt to manifest itself. It is not that the mere acidity of the blood kills the cells; in diabetic coma, for example, we do not have this acute destruction of the hepatic parenchyma; some toxic agent must be present having a specific action on the liver cells, and the coincident acidosis favors the autolysis.

Turning now to a more immediate discussion of autolysis as in itself a cause of morbid states, it will be recognized that three possible orders of disturbances may exist: (1) Disturbances due to liberation of the enzymes and diffusion of the same; (2) those due to possible toxic action of the diffused products of autolysis; and (3) alterations in the excretions due to discharge of these products. This last is but a sub-

¹See Williams, Johns Hopkins Hosp. Bull., 17:1906:71.

² Centralbl. f. Path., 15: 1904: 97.

³ Wiener, Centralbl. f. Physiol., 19: 1905: 349.

group of the second; it is not so much a morbid condition in itself as an indication and outcome of the existence of such, and we may dismiss it first.

Albumosuria.—Wherever there is extensive suppuration, with its attendant heterolysis, we gain indications in the urine by the development of what used to be termed *peptonuria*, but which now we know to be more correctly an *albumosuria*. This is most marked in the resolution of pneumonia, and in cases of empyema. A similar albumosuria, due to autolysis, has been noted in the case of large tumors undergoing softening and necrosis. In the case of diffuse autolysis of the liver, what is more marked is the appearance of amino-acids in the urine, with reduction of the urea.

T

۱,

e

)e

ie

of

10

16

Fever.—With reference to the liberated enzymes, it was demonstrated years ago by Hildebrandt that ferments of all orders injected into the blood set up marked *fever*, and the hyperpyrexia which is noted as following the development of an infarct, which follows internal hemorrhages and burns, and even that accompanying suppuration, might be ascribed to liberated intracellular enzymes. The fact, though, that there exist antibodies, or anti-enzymes, in the normal blood serum, which must tend to neutralize such enzymes, and that, save in the case of the leukocytes, we obtain no other evidence of action of these enzymes outside the organs which give origin to them, is, on the whole, against this supposition. Rather, we must regard the fever, which may show itself even in experimental aseptic suppuration, as due to the intermediate products of metabolism poured into the blood. We know that peptones and albumoses injected into the system set up high fever. To bedies of this order the pyrexia is best attributed.

Some of these bodies are distinctly hemolytic. The anemia and cachexia accompanying malignant growths are by some attributed to the extensive breaking down and autolysis of the new-formed tissue that accompanies these states. A similar anemia accompanies all oldstanding cases of suppuration.

Of the production and diffusion of acutely toxic substances we have little evidence save that of liberation of cholin in the autolysis of nerve substances. Cholin itself is but slightly toxic, but would seem to be easily convertible into the highly toxic neurin. Mott and Halliburton have found cholin in the cerebrospinal fluid in cases accompanied by nerve degeneration and softening, and have suggested that it is responsible for the convulsions and other toxic symptoms seen in these cases.

Burns.—The careful histological researches of Bardeen¹ and of my colleague, J. McCrae,² upon cases of extensive burns have shown that, even where death occurs before there has been time for adequate infection of the burned surfaces, the internal organs exhibit indications of an intense intoxication. Degenerative changes are seen in the liver, kidneys, and heart muscle, while the lymph glands present endothelial

¹ Johns Hopkins Hosp. Repts., 7:1899:137.

² Trans. Assoc. Am. Phys., 16:1901:153.

swelling and proliferation (McCrae) identical with that seen in typhoid, diphtheria, and other acute infections. The cell destruction has thus led to the formation, or liberation, from the burnt cells of bodies having an action resembling that of bacterial toxins. The rapidity with which these disturbances manifest themselves indicates that it is certain products of thermal disintegration of the cells that are directly responsible rather than the secondary products of autolysis. Whether these are the liberated enzymes, or modified proteins, or yet other substances, is not surely known.

Coagulation and Thrombosis.—Closely associated in origin with heterolysis is the coagulation of blood, either outside the vessels (*coagulation* proper), or within them (*thrombosis*), and this because everything points to these processes being due to the production of an enzyme or active discharge of the same from the leukocytes contained in the blood, and its action upon the fibrinogen of blood plasma. We shall have more to say regarding these processes elsewhere.

It is still a matter of debate whether, in breaking down, the leukocytes liberate a kinase (thrombokinase), which activates a prozymogen present in the blood plasma, or *vice versa*, or what part the blood platelets play in the process. What is pertinent in this connection is that an essential factor in the formation of fibrin is a body liberated from the disintegrating leukocytes.

It is generally accepted that in the sudden death of a mass of tissue cells within the organism there is a like discharge of a body leading to *congulation necrosis*, or congulation of the whole of the dead area. (See Section III, Chapter XXXII.)

IMPAIRED METABOLISM AS A CAUSE OF DISEASE.

It is well to bear in mind that the instances brought forward in the last chapter in connection with the internal secretions are not those of primary causes of disease. It is to the relative excess or deficiency of one or other internal secretion that the particular signs and symptoms are to be attributed, but behind this excess or deficiency are the causes which bring about the same, and these may be very various-inherited, or acquired. So, also, some of the signs and symptoms may not primarily be due to the direct toxic effect of relative excess of either internal secretion or of the bodies neutralized by the same, but may be secondary, due to imperfect metabolism, brought about by want of due amount of the internal secretion. There is, however, a group of cases of disease in which, so far as we at present see, wholly apart from internal secretions and their excess or deficiency, the cells of certain organs (once again from antecedent causes of varying nature) do not carry out the metabolic processes to their normal termination; as a consequence, there are discharged from these cells substances possessing a more or less toxic action, or, through deficient oxidation, there accumulate in the system bodies not themselves toxic, but obstructive to the proper activity of

1MPAIRED METABOLISM AS A CAUSE OF DISEASE 375

the tissues. These cases must be distinguished from another group in which it would seem that the intracellular metabolism proceeds aright, but the metabolites fail to be excreted, and, accumulating, give rise to disease. We make this distinction provisionally, or, more accurately, our knowledge is not sufficient to make it with absolute precision; nevertheless, it is well to attempt the separation. In the same rapid manner we must glance at the cases coming under these two categories.

Morbid Conditions Due to Various Products of Proteid Metabolism.—(1) Gout.¹—This is a condition characterized symptomatically by attacks of acute arthritis and other constitutional symptoms; clinically, further, by the presence of excess of uric acid in the blood; and anatomically, by the deposit of sodium biurate in the joint cartilages and elsewhere.

Here we shall not consider the underlying causes predisposing and indirect (sex, heredity, alcohol, high living, lead poisoning), but consider the metabolic disturbances which everything indicates as underlying the symptoms of this disease, these disturbances being associated with the nitrogenous metabolism. The presence of urates in the joints (Wollaston, 1797), and of an excess of loosely combined uric acid in the blood (Garrod, 1848), led naturally to this conclusion, that either excess in production of uric acid or deficient elimination of the same, is the cause of the disease. We are now convinced that this is not so. As first shown by Galvani, in 1766 (before Scheele discovered uric acid), and confirmed by Kionka, ligature of the ureters in birds leads to extensive gouty deposits in the joints, without, however, other gouty symptoms. There is excess of uric acid in the blood in leukemia and during the resolution of pneumonia, without a sign of the gouty syndrome manifesting itself. Further, large amounts of uric acid and the urates may either be given by the mouth or injected without setting up recognizable disturbance, save, possibly, slight necrotic change at the site of inoculation. They are, in fact, curiously inert bodies. At most, the *urates are an indicator*, or, in other words, while they do not themselves cause disease, the faulty metabolism which leads to their accumulation produces simultaneously other bodies having toxic effects. What these other bodies are has not been determined with absolute precision, but the study of the conditions under which uric acid is formed is that most likely to lead to their detection. As a matter of fact, this study has been pursued with great vigor of late years, and the researches of Emil Fischer and Kossel in one direction, and Jones, of Baltimore, and his colleagues, of Schittenhelm, and Burian in another, have very greatly increased our knowledge.

In the first place, it has been shown that uric acid is one of a group of substances termed *purin bodics* by Fischer, inasmuch as all are derived from, or have as nucleus, the compound $C_sH_sN_t$, or purin, the members

 $^1\,{\rm For}$ an admirable study of the modern theories regarding gout, the student is directed to the article on this subject by Futcher, in Osler and McCrae's Modern Medicine, vol. ii.

ast ry er ibng d. to or to he in sain blic er cem of

m

ıg

or

re

e-

ue

ee

of the group being derived from this by the replacement of the H atoms by hydroxyl, amide, or alkyl groups.¹

These bodies are uric acid, C₅H₄N₄O₃; xanthin, C₅H₄N₄O₂; hypoxanthin, C5H4N4O; guanin, C5H5N5O; adenin, C5H5N5; heteroxanthin, C₆H₆N₄O₂; paraxanthin, C₇H₈N₄O₂; episarkin, C₄H₆N₃O; carnin, C₇H₈-N₄O₈, and epiguanin, C₆H₂N₅O. Uric acid is thus trioxypurin; xanthin, dioxypurin; and hypoxanthin, oxypurin. Adenin is amino-oxypurin. Kossel terms them the *alloxuric bodies*, on the ground that each is made up of an alloxan and a urea nucleus, and (with the exception of uric acid) they are also referred to as the xanthin, purin, or alloxuric bases. As already noted in the first part of this work, members of this group are obtainable from nuclein. This in itself is a point of considerable importance. It indicates that in the body it is not from the ordinary cell proteins that they are derived, but from nuclear disintegration, the mother purin substance being present in the nucleic acid, combined, it is held, with phosphorus. Food, like milk, containing no purin bodies, gives rise to a minimal excretion of uric acid and purin bodies; on the contrary, feeding animals with nuclein, or substances like the thymus and pancreas (which are rich in nuclei and nucleins) leads to a great increase in uric acid excretion. There are, indeed, the two sources for the urinary purin bodies, the exogenous from the food undergoing assimilation, and the endogenous from the tissues. The excretion of the latter is relatively constant in amount for the individual, being derived from the normal disintegration of nuclear matter, and, it would appear, of muscle substance. Hypoxanthin is a constant product of muscle metabolism, and with increased exercise there is increased output of uric acid (Burian). The greatest discharge of this endogenous uric acid is encountered in leukemia, in which there is excessive production and breaking down of the leukocytes and their nuclei, as also in the resolution of pneumonia, through absorption of the autolyzed leukocytes.

We have already implied (in the statement regarding muscle hypoxanthin and urinary uric acid) that the conversion of one purin body into the other takes place within the organism. In witro, this conversion can be readily produced. Thus, if finely divided sterile pancreas be allowed to act upon guanin for some hours at 40° C., the guanin is converted into xanthin. As pointed out by Jones and Partridge, there is, obviously, a ferment present, by which the transformation is accomplished. This they term *guanase*. In like manner, Jones and Winternitz have found that through the action of *adenase* present in thymus, adrenals, pancreas, and liver, adenin is converted into hypoxanthin. Continuing these researches, they have discovered that the different glands contain different ferments or groups of ferments. The thymus, subjected to autodigestion, yields abundant xanthin, a little hypoxanthin and uracyl, but no guanin or adenin; the spleen, similarly treated, affords abundant hypoxanthin and guanin, but no adenin or

¹ I here follow Futcher's account of the relationship between these bodies, that being quite the clearest yet given.

xanthin. In both cases ammonia is split off; the ferments are typical de-amidizing agents (Chittenden).

$$\begin{array}{rll} C_5H_5N_5O &+ H_2O &= C_5H_4N_4O_2 + NH_3\\ & & \text{Guanin} & & \text{Xanthin} \\ C_5H_5N_5 &+ H_2O &= C_5H_4N_4O + NH_3\\ & \text{Adlenin} & & \text{Hypoxanthin} \end{array}$$

It will be noted that in these reactions uric acid does not make its appearance. Its development, as shown by Schittenhelm¹ and Burian,² is due to another ferment, an *oxidase*, present in the liver, lungs, muscles, and spleen, whereby the alloxur bases are oxidized to uric acid. The former observer has found another oxidase in the kidneys, liver, and muscle, which is capable of oxidizing uric acid into urea. Lastly, we have evidence of the existence of an intracellular *nuclease* (Sachs, Ivanoff), by the action of which nucleoproteids are disintegrated with the liberation of the purin or alloxur bases; and, even further, enzymes liberating the nucleoproteids from other proteins.

We thus have the following steps:

1. NUCLEOPROTEIDS, exogenous of nuclei of foodstuffs taken up by the cells; or endogenous and part of the cell structure, acted on by *nuclease*, yield

2. PURIN BASES, which, acted on by deamidizing enzyme (guanase, adenase), yield

3. XANTHIN and HYPOXANTHIN, which, acted on by an oxidase, yield

4. URIC ACID, which, acted on by an oxidase, may yield

5. UREA.

-

e

e

e

e

ľ

r

e

e

-

e

e

It need scarcely be said that this is neither the only, nor more than an insignificant, source of the urea normally excreted. We mention it here, as it throws light upon the variations in the amount of uric acid that, under different conditions, may be derived from the same diet. Nay, more, it offers a suggestion as to the cause of the accumulation of urates in the blood and tissues in the gouty state.

What is requisite now is a fuller study of the toxic effects of the purin bases. The evidence is that these are distinctly more toxic than the urates; they cause fever independent of the presence of any infective agent (Mendel),³ and the administration of adenin to dogs and rabbits leads to degenerative changes in the kidneys, with deposits of spheroliths of uric acid (or quadriurate) in the tubules, and of ammonium urate in the kidney substance. This recalls the condition of chronic nephritis, which so commonly accompanies gout. We can, at most, say that the trend of modern work is to the conclusion that the presence of the deposits of sodium urate in the joints and elsewhere, and the presence of loosely combined uric acid in the blood, indicates disturbed purin metabolism. An interesting observation in this direction is that of Beebee, working with Chittenden. Alcohol, taken with a purin-free

¹ Zeitschr. f. physiol. Chemie, 43: 1905: 354.

³ Amer. Jour. Physiol., 10: 1904: 452.

² Ibid., p. 497.

diet leads to no increased output of uric acid. Given with a diet containing a definite quantity of purin, there is an immediate increase in the excretion of uric acid. We know that alcohol is one of the factors leading to the gouty state; we know also that it reduces oxidation.

W

33

1

It would seem indicated that gout is the outcome of insufficient oxidation, whereby, in the first place, the precursors of uric acid are not wholly oxidized, and so, accumulating, set up morbid changes; while what uric acid is formed is in its turn imperfectly oxidized, and so tends to accumulate in the blood; and that this diminished oxidation is due to a constitutional deficiency of oxidases, whether inherited or acquired.

What leads to the formation of *tophi* and urate deposits in the joint cartilages is still an open question, although, contrary to the view of Ebstein, that originally maintained by the English workers is gaining ground, viz., that the deposit is primary, the necrosis secondary. It is still an open question, also, regarding the form in which the uric acid is present in the blood. The acid itself is most insoluble; it is a dibasic acid. The burate, or acid sodium urate NaHU, $(U = C_sH_4N_4O_3)$, is deposited in the joints and tophi; the neutral urates, Na₂U, have never been found in the body. Sir William Roberts, following certain work of Bence-Jones, held that the form present in the blood is the soluble, but easily decomposed, *quadriurate*, NaHU.H₂U, and made a series of observations strongly supporting the view, but it has not gained complete acceptance.

These, however, are matters not wholly bearing upon causation.

(2) **Cystin and Cystinuria**.—There is a rather remarkable condition, frequently of an hereditary nature, being observed in members of the same family, and, it may be, for successive generations, in which *cystin* is excreted in the urine. It is relatively harmless, save for the disturbance set up by calculus formation. This cystin is a sulphur-containing amino-acid—

$$S \rightarrow CH_2 \rightarrow CHNH_2 \rightarrow COOH$$

|
 $S \rightarrow CH_2 \rightarrow CHNH_2 \rightarrow COOH$


It occurs in the urine in hexagonal crystalline plates, which, as above indicated, may accumulate into concretions. There has been considerable debate as to its origin, whether exogenous and due to abnormal disintegration of proteins in the alimentary tract, or endogenous and due to some aberrant intracellular metabolism. The fact that it may be present in the absence of products of intestinal putrefaction, such as *cadaverin*; that it shows itself in the absence of proteid diet and is excreted over long periods independently of variation in the amount and nature of the food taken, together with the hereditary nature of the change, all indicate that we deal with an abnormal disintegration or conversion of the sulphur-containing portion of the protein molecule. It is possible that, normally, through the action of some special enzyme, this is in part into the neutral sulphure of the urine.

CYSTINURIA, ALKAPTONURIA, OBESITY

(3) Alkaptonuria.—A condition in many respects parallel to the last is alkaptonuria, in which, over long periods—in fact, through life—and with little apparent effect upon the health of the individual, the urine turns dark upon exposure to the air. The phenomenon has been found due to the presence of two aromatic substances, homogentisic and uroleucic acids. These evidently are due to incomplete burning up of the aromatic constituents of the protein molecule—*tyrosin* and *phenylalanin*. When those exhibiting this condition are fed with these two substances, there is a marked increase in the amount of the two acids excreted; whereas, in normal individuals, similarly fed, not a trace of either acid is to be found. The same is true in feeding with homogentisic acid. The alkaptonuric individual is incapable of carrying out the final stage of oxidation. By analogy it would seem that here also, as in gout, we have to deal with the deficiency or absence of a specific intracellular oxidase.

The normal steps would seem to be: (1) Separation of the tyrosin and phenylalanin from the protein molecule (alanin side chain); (2) oxidation of the tyrosin and phenylalanin into homogentisic and uroleucic acids; (3) further oxidation, with splitting up of the benzene (aromatic) ring. It is this last stage which is not carried out in the alkaptonuric.

There are, it may be noted, yet other causes of darkening of the urine, *i. e.*, the presence of a chromogen (melanogen) in cases of melanotic newgrowths, and of the absorption and excretion of phenol, etc.



Crystals of cystin spontaneously voided with urine. (Roberts.)

Morbid Conditions Due to Impaired Metabolism of Other Orders. —Liposis, or Obesity.—Here, in passing, we may note another condition which, like alkaptonuria, is morbid, though not of a toxic type. We refer to the accumulation of fat in the tissues. At most, where extreme, this hinders activity, and, doing this, brings about diminished oxidation, thus setting up a vicious circle. For, primarily, this accumulation of fat must be regarded as brought about by inadequate oxidation of the foodstuffs. The result may be impaired locomotion, dyspnœa, and palpitation. How much fat may be stored up is shown by Meyer and Falta's observation that the body of an individual 111 kilograms in weight afforded 51 kilograms of fat, or 38 per cent, of the total weight.

To what are we to ascribe this storing up of fat? The normal fate of fat is to be burnt up, CO_2 and water being the ultimate products. From this it follows that in obesity we have to deal either with: (1) Excessive absorption of food, either fats themselves, or substances which,

50

a

like carbohydrates, afford fat in the process of katabolism; or (2) inadequate combustion of the fats so acquired; in other words, diminished oxidation. There is a certain amount of clinical evidence that liposis may be due to both of these causes; we have the plethoric type of corpulency, occurring in those with blood rich in corpuscles and hemoglobin, and the anemic type, in pale individuals, with reduction in the amount of hemoglobin and corpuscles. In the latter, at least, the indications are those of lowered oxidative powers. Of the former, we can distinguish two groups, one largely yielding an hereditary history in which a normal diet is associated with progressive obesity. In these, two explanations afford themselves: (1) That these individuals have such excellent absorptive powers, that the digestive availability of the food consumed is in them rendered much above the normal, the excess absorbed becoming stored up in the form of fat. In a second group there is excessive feeding. As regards the first, the fact that the obese heredity is, in a very large percentage of cases, associated with the gouty diathesis, suggests that we deal with more than mere excessive consumption. (2) If gout be an indication of imperfect oxidation of one group of metabolites, so, probably, in these cases also a lack of oxidase is connected with the lack of fat consumption.

Acetonuria.—While thus, normally, we have no indication of disturbances other than mechanical, set up by inadequate metabolism of fats, there are morbid conditions best explained on the supposition that the toxic substances accumulating in the system are products of intermediate fatty katabolism. In diabetes, the peculiar odor of the breath —like that of an apple cupboard—is due to *acctone*, and this is found in the urine, in severe cases, in considerable quantities. Acetone itself is productive of little disturbance.

But associated with this in diabetes, the acetone bodies appear in the urine-*B*-oxybutyric acid and the oxidation product of the same, diacetic acid. These acetone bodies might be derived from any one of the three main groups of foodstuffs and cell constituents-from the amino-acids of the protein molecule, from the fatty acids and from the carbohydrates. Their excretion in cases of diabetes mellitus naturally suggests abnormal carbohydrate metabolism. But they may also show themselves in a group of other conditions—in high fever, wasting diseases. in cancer and in starvation, in which the store carbohydrates of the system -the glycogen-have become used up; the fact that the administration of sugar lowers their discharge is against the theory of impaired carbohydrate metabolism. We know so little regarding the normal stages of fat disintegration that it is not possible to formulate how these bodies are derivable from the fatty acids; we can only suggest that just as β -oxybutyric acid by oxidation becomes converted into diacetic acid, and then into acetone, so the higher fatty acids may similarly become changed into these acetone bodies by progressive oxidation. Eppinger's¹

¹ Wien, klin, Woch., 1906: 111.

ACETONURIA, ACIDOSIS

observation that the administration of amino-acids to rabbits arrests the appearance of acidosis would suggest that normally the ammonia evolved in ordinary protein metabolism combines with these lower fatty acids and prevents their appearance in the urine. It may be that their appearance in starvation is due to the fact that the fats of the organism are disintegrated at a greater rate than are the proteins. Nevertheless, it has to be admitted that if we accept this progressive disintegration of the fatty acids, we must also recognize the possibility that the aminated fatty acids—the amino-acids of the protein molecule—are capable of undergoing a like series of disintegrations. In other words, the matter of the origin of the acetone bodies is still very largely an open question.

The symptoms, it may be added, are those common to acidosis—they are those of a grave intoxication, with air hunger, and nervous symptoms passing into coma and death.

Acidosis.—That the accumulation of β -oxybutyrie and diacetic acids in the blood is the cause of the main symptoms of diabetic coma is evident from the fact that almost identical conditions follow the treatment of animals, especially herbivora, by repeated doses of inorganic acids. They become stuporous, their gait is unsteady, the breathing extremely rapid, the blood is bright red, containing much less CO₂ than normal, and with this there is a marked diminution in the alkali of the blood. Administer alkalies, and these symptoms pass away. The explanations would seem to be that, normally, the alkalies of the blood take up the CO₂ as it is formed in the tissues, convey it to the lungs, where, aided by the oxidase present, the CO₂ is split off and the salt, once more rendered basic, is prepared to join with another molecule of CO₂ in the tissues. Where there is an excess of other acid in the blood, it is this that combines with the alkaline salts, and as a result the CO₂ accunulates in the tissues, symptoms of asphyxia ensuing.

In carnivorous animals the combination affects not only the alkalies proper, Na, K, etc., but also ammonia. There is in these cases a diminution of urea in the urine, with great increase in ammonia compounds, indicating that in the disintegration of proteins there is an ammonia antecedent of urea, and that this in the blood combines with and neutralizes the excess of acid present.

As already noted, there are other conditions besides diabetes which afford these symptoms of acidosis with discharge of acetone bodies starvation, fevers, cachectic conditions, and, we should add, the pernicious vomiting of pregnancy, chloroform anesthesia, retained placenta and fœtus. These last conditions have this in common, that they may exhibit profound hepatic disturbances. Whether here the glycogenic function is gravely disturbed, or the lipolytic, must be left an open question.

Another acid which has been detected in the urine in increased quantities is *lactic acid*—in rheumatism, osteomalacia, and rickets. This is never in quantities sufficient to set up the extreme changes produced by the acetone group, and, as regards the effects on individual tissues,

there is so much debate and contradiction as to display our almost complete ignorance of its action as a cause of morbid states.¹

SOI

m

be

ac

me

of

of

 \mathbf{p}_i

Dyspncea and Asphyxia.—These conditions may now strictly be included among the acidoses. For long there was debate as to whether the main symptoms were due to lack of oxygen or excess of CO_2 . The researches of Haldane and Priestley appear to have finally answered the question.

It is well established that muscular exercise most materially leads to a using up of the oxygen in the blood and to an active discharge into the same of corresponding increased amounts of CO_2 . In the circulating blood the above observers have shown that the amount and tension of the oxygen may be altered from 20 to 8 per cent, without any increase in the depth or frequency of the respiratory movements, but that relatively slight increase in the above, increasing its activity, whereas diminution of the tension of the CO_2 in the blood, finds the respiratory centre extraordinarily sensitive, increasing its activity, whereas diminution of the tension of the gas depresses the activity of the centre and causes slowing of the respiration, or even apnea—total arrest of respiratory movements. A rise of only 0.5 per cent, in the tension of the carbon dioxide in the air in the alveoli, and so in the blood circulating around those alveoli and supplying the brain, was found to increase the volume of air respired 100 per cent.

This may appear contradictory to what has just been noted regarding the air-hunger of diabetic coma, in which the amount of CO_2 in the blood has been found greatly decreased (in one case Minkowski found it reduced from normal 36 to 3.3 per cent.). It must be remembered that, as Haldane and Priestley show, the phenomena depend not upon the total amount of CO_2 which can be extracted from the blood, but upon the tension of the (free) CO_2 . In diabetic coma other acids combine with the alkalies present in the blood plasma. Very much less CO_2 , in consequence, is taken up; but it still diffuses into the blood, and is present in a free state.

In this connection might well be discussed the subject of *impaired* carbohydrate metabolism. It has, however, seemed more serviceable to take up the main example of this—diabetes mellitus—in the previous chapter in connection with the internal secretions.

THE INTERMEDIATE INTOXICATIONS.

Gastro-intestinal Intoxications.—Constipation.—It is a familiar experience to those regular in their habits, or, in other words, to all healthy adults, that failure to pass the morning evacuation is apt to be

¹ To this lactic acid Zweifel attributes the acidosis of pregnancy, finding this acid both in the urine and in the blood of eclamptic cases. For a criticism of modern views on this form of acidosis see Leathes, Proc. Roy. Soc. of Medicine, 1;1908; No. 5.

THE INTERMEDIATE INTOXICATIONS

followed by a certain amount of heaviness and feeling of being "out of sorts," with mental dulness, accompanied, it may be, later in the day, by slight headache and malaise. It is true that a habit of constipation may be developed, and may continue for years without grave lesions being manifested. The late Sir George Humphrey, in his lectures, was accustomed to cite the case of two elderly maiden ladies, patients of his, who, living abstemious lives, indulged with regularity in but a monthly evacuation. We do not imagine that those two old ladies were cheerful; the common experience is that habitual constipation is associated with irritability and depression of spirits, other symptoms of a low toxic state showing themselves in lack of energy, muddy skin, and tendency to "spottiness."

It is more particularly in those cases in which there is a sudden or complete obstruction of the upper bowel that pronounced indications of intoxication show themselves. The subnormal temperature, vomiting, muscular weakness, and actual collapse encountered in these cases of ileus can only be attributed to a resorption of intestinal contents, and this is proved by the appearance in the nrine of considerable amounts of the products of imperfect or abnormal metabolism of the foodstuffs, of indoi, and other sulphuric acid nitrogenous compounds.

The graver and more immediate results ensuing upon obstruction of the upper intestine, as compared with the lower, indicate one of two things-either that normally in the digestive process toxic bodies are elaborated in the stomach, which undergo modification into harmless compounds lower down the digestive tract, or that the result of obstruction is to lead to abnormal fermentations in the upper portion of the canal, to the abnormal growth of bacteria there, and to the production of toxic products of their activity. There is a certain amount of evidence in favor of each of these suppositions. It has been shown by Magnus-Alsleben¹ that the contents of the stomach and duodenum of the normal dog are distinctly more toxic than are those of the lower part of the bowel. On the other hand, we know that in the healthy individual, while abundant bacteria may be taken into the stomach with the food, they undergo in the main a fairly rapid destruction, and that in the duodenum of the healthy animal that has been starved for a short time absolutely no bacteria may be present. The case is very different in the rest of the small intestines, whose alkaline contents favor the rapid proliferation of bacteria, so that in the neighborhood of the lower end of the ileum they reach their maximum abundance and activity; that, passing into the colon, with the concentration of the feces, their number now undergoes rapid diminution, so that from the stools of a healthy individual it may be that relatively few colonies are obtainable, although other observations demonstrate that quite a considerable proportion of the matter constituting the healthy stool is in the form of bacterial remains. We know, further, by experiment, that obstruction to any part of the alimentary canal leads to active proliferation of bacteria

¹ Hofmeister's Beiträge, 6: 1905: 503,

fo

above the point of obstruction and to the assumption of definitely increased virulence by those bacteria. Thus, it is more than probable that the toxic effects of obstruction are associated with this increased bacterial activity. Granting all this, it has to be admitted (1) that the substance or substances setting up the particular train of symptoms seen in ileus has not yet been isolated, and (2) that intestinal bacteria grown outside the body have not, so far, been shown to produce substances setting up these particular symptoms. We are, in short, only at the beginning of a knowledge of the intestinal intoxications.

Indol.—Of the known products of the putrefactive or bacterial decomposition of proteins, but one, or one group, stands out with any prominence, and this more as an indicator of intestinal putrefaction than, it may be, as an active toxic agent. This is indol, and with it other substances derived from the aromatic radicals of the protein molecule. We know that what is the commonest of the bacterial inhabitants of the intestine-the B. coli-is capable of forming this from peptones outside the body. We detect it in the urine in those cases of obstruction. Indol, as shown by Herter,¹ has distinct and suggestive toxic properties. It is capable of setting up irritability, mental dulness, and headache-symptoms, it will be seen, closely resembling those of constipation. The amount found in the urine of well-marked cases of obstruction is, however, so small, compared with that necessary to induce these symptoms when administered by the mouth, that it is gravely debated whether this can be regarded as the agent setting up these particular symptoms in obstruction. Our inclination is toward accepting Herter's view, and this from the fact that when indol is administered per os in quantities sufficient to produce definite symptoms, only a very small proportion can be regained from the urine; it would seem, therefore, quite possible that in obstruction there may be absorption of amounts of indol adequate to induce symptoms, and yet there be little excretion in the urine. But something more would seem necessary to account for the graver symptoms of ileus. Other decomposition products of the aromatic radicals are skatol, phenol, paracresol, indolacetic acid, paraoxyphenylacetic acid, and paraoxyphenylpropionic acid. All these may be detected in the urine.

Ptomaines.—For a time great expectations were based upon these bodies, not only as affording an explanation of the intestinal intoxications, but also as explaining the specific toxic effects of pathogenic bacteria. As regards the second of these, we now know that the ptomaines are not specific—*i. e.*, particular species of bacteria do not lead to the formation of ptomaines peculiar to those species; wherefore, to the ptomaines cannot be attributed the specific toxic symptoms of the different diseases; and secondly, that the amount of ptomaines present in bacterial culture fluids is inadequate to explain the toxic effects of the fluids. As regards intestinal intoxication, it may be, as we shall point out, that one group of ptomaines is responsible for certain disturbances. If animal matter of various orders be allowed to undergo putrefaction

¹ New York Medical Journal, 68: 1898: 89 and 116.

THE INTERMEDIATE INTOXICATIONS

for several days, the product was found by Gautier to contain small but definite quantities of nitrogenous bases, allied chemically to the vegetable alkaloids. To these Selmi, in 1881, gave the name *phomaines* $(\pi \tau \omega \mu \alpha, a \text{ corpse})$. We owe the fullest study of the substances and their properties to Brieger. The active and abundant studies of the "eighties" demonstrated the existence of a large group—some two score—of these bodies; among them may be noted:

Methylamin	$CH_2 - NH_2$
Dimethylamin	$CH_3 \longrightarrow NH \longrightarrow CH_3$
Trimethylamin	$CH_3 - N - CH_3$, or $N(CH_3)_3$
	CH ₃
Cholin	$CH_2OH - CH_2 - N(CH_3)_3 - OH$
Neurin	$CH_2 - CH - N(CH_3)_3 - OH$
Muscarin	$CH(OH)_2 - CH_2 - N(CH_3)_3 - OH$

Certain basic substances, such as the tyrotoxicon separated by Vaughan from putrefying cheese and milk products, while intensely toxic, have not vet been obtained in sufficient amounts to permit full analysis. While the majority of these bodies are non-toxic, a few are intensely toxic, having properties and constitution very similar to those of muscarin and other vegetable alkaloids; muscarin itself has, indeed, been isolated by Brieger from the cod. The central radicals of these bodies are amino groups; this fact and the mode of origin indicate that they are penultimate¹ products of the disintegration of proteins. Nevertheless, it may be that these are not in all cases directly derived from the pro-Among the most widespread components of the cell are the compound fatty bodies, the phosphatides, compounds of fatty acid molecules with phosphoric acid and cholin. The disintegration of these bodies will afford cholin, and this, indeed, has been isolated from brain matter (affording abundant phosphatides) that has undergone putrefaction. Cholin itself has relatively slight toxic powers; in large doses, however, Mott and Halliburton were the first to it has muscarin-like effects. call attention to the presence of free cholin in the cerebrospinal fluid in degenerative conditions of the brain matter. Donath² and Rosenheim⁴ have confirmed their work. The former finds that, introduced directly under the dura, cholin has a powerful action upon the nerve substance, setting up severe convulsions, tonic and clonic. He suggests thus that cholin may be responsible for the development of epileptic attacks.

On the other hand, introduced into the alimentary tract, cholin is found relatively harmless; the case is very different with neurin and muscarin, of which the former can be gained from cholin. As shown by the formula already given, it differs from cholin by the loss of a molecule of H_sO ; it is intensely toxic. That neurin is formed in the organism

25

a

21 2.

le

le

re

se

ic

10

¹ Penultimate, because if putrefation be continued beyond a certain limit they diminish in amount, giving place to nitrates and ammonium compounds.

² Zeitschr. f. physiol. Chemie, 39; 1903; 526.

³ Journal of Physiology, 35: 1907: 465.

is indicated by Kutscher's detection of it in human urine. It is thus well within the bounds of possibility that some of the depressant effects of constipation are due to the development and absorption of bodies of the neurin and muscarin type from fermenting fecal matter. But if such be present, it must be purely as the result of bacterial activity, and the same is true of the aromatic derivatives, indol, skatol, etc. Investigators have been unable to gain either category of bodies by the action of the digestive ferments upon foodstuffs in the absence of bacterial growth.

Here a word of caution should be introduced against the acceptance of not a few of the results announced by workers, more particularly of the Bouchard school. Toxic effects have been ascribed to not a few products of protein disintegration recognized as present in fecal matter, the urine, and other excretions, which now we are assured have been due to accompanying potassium salts.

It would appear to be bacterial fermentation more than the action of the digestive ferments that leads to the production of irritative and noxious products of carbohydrate disintegration. Foremost of these are the series of organic acids—lactic, butyric, formic, etc. Little is known regarding the deleterious effects of these in individual cases, but, on the other hand, it has been shown by Boix and others that the administration of them in repeated doses leads to very definite disturbances in the different tissues, of a chronic type; such as, for example, cirrhosis of the liver (Boix).

Besides cirrhosis, many of the more chronic or constitutional disorders have, indeed, been ascribed to absorption of the products of abnormal gastric and intestinal fermentations: chlorosis, pernicious anemia, rheumatoid conditions, Graves' disease, myxedema, migraine, and epilepsy—and the series could be greatly extended. It is, indeed, well within the bounds of possibility that the ascription is in many of these cases correct, but indubitable evidence is wanting. It is also possible, as I have pointed out elsewhere,¹ that in some of these, at least, we deal not with intestinal absorption: excessive growth of one or other form of intestinal bacteria and the resulting irritative conditions of the mucosa being followed by increased carriage into the portal circulation of bacteria of low virulence, the destruction of the same, and liberation of their endotoxins within the vessels and tissues, causing hemolysis and hepatic irritation.

It will be seen that these considerations all tend to point in the one direction—that the gastro-intestinal intoxications are, strictly speaking, exogenous, and in no sense auto-intoxications.

OBSTRUCTED ELIMINATION OF THE PRODUCTS OF KATABOLISM AS A CAUSE OF DISEASE.

Two closely related conditions may be considered under this heading: (1) The failure of excretory organs to eliminate, in consequence of

¹ Jour, Am. Med. Assoc., 33; 1899; 1503 and 1572,

disease; and (2) the retention of the eliminative powers by these organs but subsequent resorption of constituents of the excreta, owing to obstruction in the ducts.

of if

of

h.

ts

е,

se

it,

e,

ne

he

he

ne

M

The results of the two processes are not necessarily identical; there is, for example, a marked difference, as pointed out by Rose Bradford, between the results of removal of the two kidneys and ligature of the two ureters, a difference showing clearly that certain, at least, of the final metabolic processes are conducted by the gland cells of the excretory organ. In other conditions, as in obstructive jaundice, there are indications that the metabolites, the results of the cell activities of the excretory organ, may, as the result of the obstruction, not necessarily be excreted and then reabsorbed, but may pass direct from the excretory cells into the blood and lymph. No sharp line is to be drawn between the two conditions.

The Resorption of Excretions.-Excretions, or certain constituents of the same, undergo resorption under normal conditions to a far greater extent than is generally realized, and this as a physiological process. The solidification of the feces in the colon is in itself evidence that the watery constituents of the digestive juices become reabsorbed; the view nowadays most generally accepted regarding the mode of secretion of urine is that the urine becomes concentrated in its passage down the urinary tubules by resorption; the remarkable increase in the solid constituents of gall-bladder bile, as compared with that collected direct from the bile ducts, indicates a similar absorption. Not only this, but the complete disappearance of certain constituents from the intestinal contents is most simply explained in the supposition that they undergo resorption there. Nay, more, as shown experimentally by Aschoff, if the full gall-bladder be occluded, within a few days it may be found empty, all the constituents being taken up again. As I point out elsewhere, we are compelled to recognize a *reversibility* not merely of enzyme action, but also of cell activity. According to the influences acting upon either side of a secretory cell, so will it in certain cases either secrete or absorb. So long as the resorption is within limits, little or no disturbance is set up. Serious results are apt to occur if the process be long continued, and this with greater ease if the excretory products be taken up by cells other than those which have

The most striking example of such resorption occurs in *obstructive jaundice*. There has been, and continues to be, debate as to the exact process which occurs here. Both Harley and Ziegler conclude that the liver cells may, in consequence of the obstruction, discharge the substances elaborated by them, not into the bile duct, but into the system; they differ as to the details of this discharge. The former, by coincident ligature of the common bile duct and the thoracic duct (which receives the lymphatics from the liver), shows clearly that the onset of jaundice is delayed for several days over cases in which the bile duct alone is ligated, and concludes, therefore, that the bile constituents pass from the liver cells into the lymphatic system of the organ. The

be

W

at

pi

11

a

latter demonstrated equally surely by histological methods that the bile can be seen making a direct entry into the hepatic blood capillaries. The probability is that both are right and both wrong; that neither process is exclusive; and it may be that one or other predominates, according to the activity of the liver cells. But over and above this, where the obstruction is not in the bile capillaries, but lower down, the intrahepatic ducts are found widely distended, with signs of irritation around them, and, as Ford' more particularly has shown, working in our laboratory, this irritation may lead to the development of a characteristic type of duct cirrhosis. It is evident that in these cases there is resorption from the bile ducts.

A fuller description of jaundice is given in the discussion of the various forms of pigmentation in Section III, Chapter XXXI. Here it need only be said that jaundice is more than a mere pigmentation; accompanying this are cerebral symptoms, slowed pulse, itching of the skin, lowered coagulating power of the blood, with tendency to hemorrhages. The more important of these symptoms, cerebral dulness, slowed pulse, and even hemolysis, can be reproduced by the experimental injection of the bile salts. It is those that are in the main responsible for the symptoms other than the jaundice itself.

The indications are that these bile salts, under physiological conditions, undergo resorption in the intestines, setting up there no disturbance, and that because, as Sir Lauder Brunton has pointed out, taken up thence into the portal system and conveyed once more to the liver, they are there again taken up and discharged by the liver cells, and so, as it were, undergo a lesser circulation. There is a like intestinal resorption of the bile pigments, but this after they have in the bowel been converted into hydrobilirubin. Neither icterus nor cholemia would ever seem to be set up by absorption of these bodies from the alimentary canal.

The Pancreas.—Opie has called attention to the deleterious local effects of resorption of the pancreatic juice.

If the pancreatic ducts of a cat be ligated and the animal killed at the end of two or three weeks, the gland is found to be the seat of chronic interstitial inflammation. Schulze and Ssobolew have performed similar experiments with similar results. The inflammation specially shows itself around the ducts.

As Opie,² Halsted, and others have shown, obstruction of the ampulla of Vater by a biliary calculus may, under certain conditions, lead to the obstructed bile making its way into the pancreatic duct, and then its absorption leads to the more acute condition of hemorrhagic pancreatitis.

The Kidneys and Uremia.—Under the heading Uremia we include all the symptoms associated with retention in the system of the urinary constituents. Such retention may be of more than one order. It may

¹ American Journal of the Medical Sciences, 121: 1901: 60.

² Am. Jour. of Med. Sci., 121: 1901: 27, and Diseases of the Pancreas, 1903: 71.

OBSTRUCTED ELIMINATION

be due (1) to disease of the kidneys and renal incompetence, so that these constituents fail to be abstracted from the blood; or (2) to ureteral obstruction, so that, primarily at least, the kidneys perform their function, and, as in the case of the obstructed bile duct, in part we deal with resorption of the urine, in part with return from the renal epithelium of the products of their metabolism into the blood or lymph; or (3) it may be due to resorption of the urine from the bladder, where there is a prostatic or urethral obstruction. In this last case more particularly there is apt to be infective and fermentative change in the urine, and it is the modified constituents that are resorbed. German authorities have spoken of this condition as ammoniemia, the earlier idea being that the symptoms were more particularly due to the absorption into the system of the ammonia salts, the result of decomposition of the urea. It is true that ammonium carbonate has certain toxic properties, and that its absorption in considerable quantities may possibly set up disturbances; it has not been shown that this produces the symptom complex seen in vesical obstruction; it would seem more probable that intermediate products of protein metabolism are here to blame.

The symptoms are most marked and most characteristic in the cases of the first order, which are the most frequent; it is in these that we are most apt to have pronounced headache, nausea, and vomiting, convulsions of an epileptic type, passing on to coma, dyspnœa, with asthmatic attacks and Cheyne-Stokes breathing, and evidence of gastric and intestinal catarrh, going on to ulceration. It is remarkable and suggestive as to the nature of uremia, that where the kidneys have not been primarily at fault, then, in case of urethral or vesical obstruction, there may be complete anuria for several days without uremia showing itself; when in these cases it does supervene, it suggests itself that the back pressure has eventually led to renal disturbance; in other words, the most satisfactory view is that uremia is due to the heaping up in the blood of substances not acted upon duly by the renal epithelium. This view is supported by the experiments of Rose Bradford upon the different effects of complete removal of the kidneys in animals, as compared with the fact that nephrectomized animals may be kept alive for several days and uremia be warded off by injecting extracts of healthy kidney substance. Such kidney extract cannot excrete and discharge urea and its precursors; they must still tend to accumulate in the blood; but it may convert certain of these substances from a toxic into a relatively harmless state.

ia

10

What the substance or substances may be that are responsible for the nervous and other symptoms has been the source of abundant debate, experiment, and theory. One theory after another has been propounded, only to be shown inadequate; the present standpoint is one of a healthy agnosticism. As above stated, ammonium carbonate does not suffice, neither does ammonium carbamate (a possible precursor of urea). The potassium salts which should be discharged are toxic, but if they accumulate slowly in the organism they set up little

disturbance. Bouchard has described toxic substances present in normal urine, possibly ptomaines (diamines), differing in the overnight and daily urine. Stadthagen has wholly denied his findings. There is, as you Jaksch showed, a distinct accumulation of urea in the blood of most (but not all) uremic cases, but urea is curiously inactive, save upon the kidney itself. The same observer finds also an increase of uric acid, but the yet greater amount of this in gouty blood sets up no uremia. Kreatin and kreatinin have been invoked because the latter applied to the cerebral cortex causes convulsions; but these are not increased in the uremic blood. The family likeness of the conditions to diabetic coma has suggested that acidosis is the cause. Von Jaksch calls attention to the diminished alkalescence of the blood; A. E. Taylor denies that there is any such.

> cai end the foc

arr (car and act

eve all

thi firs or of (ane blo res mu acc ind org tie exc mo the

CHAPTER XIII.

BODILY STATES AS DIRECT AND PREDISPOSING CAUSES OF DISEASE,

NUTRITIONAL DISTURBANCES.

WHILE, as we have pointed out, the number of chemical bodies which can be dissolved in the fluids of the body and taken up by the cells is enormous and varying in effects, it is of importance to consider apart those substances which essentially subserve nutrition, which are used as foodstuffs for the daily replacement of cell matters discharged in the carrying out of cell activities, and for the building up of additional cell matter in growth. We must thus rapidly pass in review the effects of arrested or inadequate supply, and of excessive exhibition of the same.

Such foodstuffs belong to five great groups—oxygen, water, proteins, carbohydrates, and fats; to those must be added certain mineral matters and salts not merely essential for the due performance of metabolic activities, but, like the iron of hemoglobin, the phosphoric acid of the nucleus and the calcium and magnesium salts of bone, actually built in the structure of growing tissue. We may, in short, include as foodstuffs every substance the continued deprivation of which—as is the case with all the orders of matter above indicated—leads to the death of the individual. It is usual, we know, to regard the first two here mentioned, oxygen and water, more as intermediaries than as actual foodstuffs, but this is incorrect; they are built into the growing cell. We would, in the first place, indicate the disturbances set up by deficiency or excess of one or other order of substances, and then note the effects of lack and excess of food in general.

Oxygen.—The disturbances resulting from deficiency in the oxygen inhaled are treated more fully in the section upon respiratory disturbances. We would but recall that the sudden cutting off of the oxygen supply, accompanied as it is by progressive increase in the CO_2 of the blood and tissues, induces *aspheria*; the accompanying changes in the respiratory act—the altered rate and employment of the accessory muscles of respiration—constituting the condition of *dyapnava*; the accompanying congestion of superficial vessels with dark venous blood inducing *cyanosis*. Post-mortem, besides the congestion of various organs, and the capillary hemorrhages (ecchymoses) of, more particularly, the pericardium and pleura, it is noticeable that through excess of CO_2 the blood fails to cagulate, while if the condition of moderate asphyxia and O deficiency has been prolonged for several days, the tissues show fatty degeneration. The effects of excess of oxygen

inspired are relatively slight, and this because with ordinary respiration the arterial blood becomes so nearly saturated that the exhibition of more oxygen has little effect.

ec

ar

W

ne

ec

ec

in

 \mathbf{p}_{1}

80

SC

W

tł

SC

of

pa rł

m

as

ir

a

Water.—Deprivation of water leads to death more swiftly than does deprivation of solid food. There is a steady drain of water from the body through respiration, through the skin and the excretions, and if this be not counterbalanced, it would seem that the accumulation of waste products in the cells brings about their disorganization. The first indication of this lack of water is *thirst*, the next a general shrinkage and falling in of the tissues as the interstitial and cell fluids are given up to maintain the circulation; with this there is complete loss of appetite for solid foods. Mental excitement and delirium supervene, followed by coma and complete inanition. The dryness and the temperature of the air make a profound difference in the period during which the individual can sustain total deprivation of water. Death may supervene on the second or third day in a tropical desert, may not present itself till the second week in north temperate regions (in those immersed in damp caves, mines, etc.).

Excess consumption of fluid, through ready absorption of the water through the intestinal walls, may also lead to disturbances. One form of torture in the Middle Ages was to induce death by forcible continued pouring of water down the throat. The consequent lowering of the specific gravity of the blood tends toward grave changes in the erythrocytes, swelling and liberation of their hemoglobin, with an edematous condition of the body cells, more particularly those of the excretory organs. Pure water, it may be noted, applied to the internal tissues, as distinct from isotonic saline solutions, is capable of bringing about cell death. Excessive consumption of a lesser grade, and repeated, as in sundry German beer drinkers, leads more particularly to cardiac dilatation and hypertrophy (Munich "beer heart"), in consequence of the increased volume of blood and work thrown upon that organ.

Proteins, Fats, and Carbohydrates.—The abundant studies of the physiologists have shown that none of these can be cut out of the dietary without eventual grave disturbances and death. Deficiency in any one of them leads to intense desire for articles of food containing them.⁴ To a certain extent, but not permanently, earbohydrates and fats can replace one another. This question of dietetics is norwadays so fully discussed in the leading text-books of physiology, that it is unnecessary to enter into it here. The studies of Atwater Bubner, and others, on the one hand, upon the caloric values of different foodstuffs, and of Chittenden and his pupils, upon the minimum make requisite to sustain active life, are matters of active and general interest. Suffice it to say, that in the temperate zone, for the adult of merage size, a daily consumption of these foodstuffs in proper processor, at least advisable. The views already expressed upon the unstance in practically all the tissues of

¹ See Shackleton's Heart of the Antartic, 2: 1909: 4.

NUTRITIONAL DISTURBANCES

reserve material and reserve force, strongly suggest that it is not an economic principle of the organism in general to limit itself narrowly to an exact preparation to supply what is demanded under ordinary conditions, and no more. The indications are, thus, that the system is well prepared to consume more food than is necessary for the daily needs, and that such luxus consumption of a moderate grade is, if not economic, at least physiological. This, we would urge, is far from counselling persistent overfeeding, and is wholly compatible with the belief that most of us consume too much, more particularly of animal proteins, nor is it incompatible with the conviction that an occasional fast, as carried out by many peoples, is of distinct advantage, by insuring physiological rest and permitting the elimination of intermediate products which may tend to accumulate harmfully.

The most striking disturbances set up by an improper dietary are scurvy (scorbutus) in adults, and in infants, Barlow's disease, or infantile scurvy. The former may show itself in those consuming the proper proportion of proteins, fats, and carbohydrates; or, again, in those who, living mainly on potatoes and tea, have a deficient intake of proteins, When abundant meat is taken, and the disease shows itself, it is noteworthy that the meat has been salted or canned. So also in Barlow's disease, the infants may have been given abundant milk foods, but these either boiled or Pasteurized, or artificially prepared by heating. In these conditions disturbances of a like order manifest themselves; in scurvy-anemia, swelling, and softening of the gums, with loosening of the teeth, and a tendency to hemorrhages here and elsewhere, with pains in the bones and joints; in Barlow's disease-anemia, hemorrhages from the gums, extensive periosteal hemorrhages, and evidently intense pain in the bones and joints, so that the child fears the slightest movement. With this, actual changes in the bones are indicated by the increased liability to fracture. In both conditions the administration of fresh fruit or of acid fruit juices brings about rapid amelioration.

The indications are that the disturbing influence in both cases is not so much the altered proportion of the different forms of food administered as what has been termed the devitalized condition of essential elements of the food. More accurately, it would seem, that in the preparation the calcium, the phosphates, and, it may be, other salts, become associated in such a way that they cannot be absorbed and utilized by the organism. It is becoming increasingly recognized that particular salts in loose association are necessary for the due activity of the digestive and other enzymes.

It is difficult to write positive and briefly regarding the effects of the excessive exhibition of the difficult orders of foodstuffs. As already indicated, an excessive diet of piceuins, contrary to the general opinion, does not necessarily produce good shough it may be a factor in the production (p. 375); so also, excessive consumption of fats and carbo-hydrates, while one cause of *obesity*, is in the only cause (p. 379). What is becoming more and more recognized is that the continued consumption of excessive food, by maintaining a condition of overwork and over-

BODILY STATES AS CAUSES OF DISEASE

wh

des

de

up

un

de

sto

the

lea

m

by

alt

lar

of

no

tw

sei

of

va

tai

exi

me

wi

en

an

of

du

po th

T

of

or

stimulation of the digestive apparatus, eventually leads to exhaustion of the same, which may show itself, according to the individual, by degenerative atrophic and chronic interstitial disturbances, now in one organ, now in another. There is, for example, a certain amount of evidence that the lactic, butyric, and other acids freed in the dissociation of carbohydrates given in excess lead to cirrhotic changes in the liver, and that the indol compounds, as, again, certain of the purin bases and their compounds, dissociated in the breaking down of proteid and nucleoproteid foodstuffs when in excess have deleterious effects upon the liver, kidneys, etc. (p. 381).

Iron, Calcium, Etc.—Indications have already been afforded of the value to the economy of minute amounts of various mineral matters. Experimentally, it has been shown that diets specially prepared so as to be deficient in iron, are followed by grave disturbances of the hematopoietic system, those deficient in calcium by inadequate development and anomalies in the growth of bone, together with grave disturbances in general metabolism; those deficient in phosphoric acid, whether inorganic or organic and combined as in the phosphatides, by delayed growth and marantic conditions. The necessity for the various salts is indicated by a study of their rate of excretion in the urine; thus, as much as 16 grams of sodium chloride, and from 2.5 to 3.5 grams of phosphoric acid are excreted daily in the urine.

Just as we shall proceed to show that a distinction is to be drawn between starvation, or withdrawal of food, and marasmus, in which food is exhibited but cannot be utilized, so it would seem that two orders of events obtain in connection with certain of the more important of these salts. Apart from conditions due to insufficient administration, there are those in which, either while present, they are administered in such combination as the tissues are unable to utilize them, or from constitutional defects these tissues may be unable to assimilate or employ them when presented in the normal form. There is, thus, for example, a series of conditions of imperfect development of bone, some members of which may even be inherited, others congenital and showing themselves during intra-uterine life, others, again, acquired, in which there is no adequate evidence that the calcium and phosphoric acid are not exhibited, but in which certainly the bone corpuscles are incapable of building them to form normal bone at the proper period. The best known of these is rickets (rachitis), in which, with excessive preparation for the laying down of bone, in the form of cartilaginous overgrowth of the epiphyses, etc., the actual deposit of bone salts is delayed, and, as a consequence, the bones yielding to the various strains to which they are subjected, become deformed. It is interesting to note that after this preliminary period there is apt to be a reactive excessive deposit of bony salts, the bones becoming denser, and in some cases larger and even longer than normal. Into this same category come also, in all probability, the conditions of scurvy and Barlow's disease already mentioned. While in all of them we are still ignorant as to the exact causation, everything points to the fact that we deal with disorders of assimilation.

Starvation.—The opposite condition of withholding of foodstuffs, whether complete or partial (as through disease leading to stenosis and destruction of the upper part of the digestive canal), leads to a very definite train of symptoms, the most obvious of which is progressive emaciation. It would seem that the circulating proteins are first drawn upon, for in complete deprivation the nitrogenous excretion continues unchanged for forty-eight hours or thereabouts; next, the glycogen deposits in the liver and muscles become used up, and with this the fatty stores of the body begin to disappear. Following upon this it is noted that the muscles undergo pronounced shrinkage and diminution, those least used suffering most. Those actively exercised, like the heart muscle, remain in good condition for relatively long periods. Excretion by all the organs becomes rapidly diminished, and may sink to nil, although, if water be taken, the urinary excretion continues and now contains members of the acetone group of bodies. The leukocytes of the blood undergo reduction in number, the erythrocyte count remains largely unaffected, although, taking into consideration that the fluids of the body suffer general diminution, it is probable that this maintenance of the number of red corpuscles is only apparent (Stengel). Death may not occur until the weight has been reduced 50 per cent.

With these changes there is the sensation of hunger, which can be lessened by taking water, and in general is most marked during the first two or three days, being followed by a period of listlessness and lowered sensation; the mental and bodily powers become sluggish, and effort is followed by rapid exhaustion. With this there is progressive depression of the body temperature, and the development of the state of complete *inamition*.

The period during which the organism can endure complete deprivation of food depends upon several factors. Of these the most important are: (1) The bodily state at the beginning of the period. The existence of abundant reserve stores of fat, etc., prolongs the period. (2) The consumption of water. Without water, whether taken by the mouth or absorbed through the skin, life persists for at most a week; with water, as in professional "fasters," it may be prolonged for fifty days and more. (3) Rest. Active exercise uses up reserve material and energy. (4) Practice. The individual can train himself to stand longer and longer periods of fasting. The surrounding temperature, the state of health prior to the deprivation of food, the incidence of infection during the period, are other modifying factors.

Marasmus, or progressive wasting, differs from starvation in this important particular, that in it abundant food may be exhibited, but through imperfect assimilation an inadequate amount of food can be used. The condition may be either congenital, showing itself from the moment of birth in syphilitic infants, for example, or acquired either in infancy or later life.

BODILY STATES AS CAUSES OF DISEASE

OVERSTRAIN.

di ch

mi

pa

alı

no

asi

the

be

wi

ge

re

ca

aft

di

an

of

in

hu

ca

ca

n

ac

SO

m

di

gr

at

wl

as

th

m an

A condition that of late years has come in for not a little study, either as directly causing morbid states or as rendering the organism more susceptible to disease, is that of overwork and fatigue. It is necessary that we should call attention to the more important data bearing on the subject.¹

It may, at the outset, be noted that there is some little latitude in the employment of terms—some would limit fatigue to the physiological result of work, and would speak of exhaustion, surmenage, or overstrain, as a severer and pathological state, resulting from overwork. Others, on the contrary, would speak generally of fatigue as resulting from overwork. For ourselves, the meaning implied by "overstrain" is so obviously that of a pathological state that we are prepared to employ this term in a pathological sense, and "fatigue" to indicate more physiological states.

It does not need the evidence of exact studies upon the action of isolated muscles of cold-blooded animals to inform us that work within natural limits is followed by fatigue, so that what at the beginning was done with ease, with repetition of the act demands increasing effort for its accomplishment. Whereas such fatigue passes off if followed by adequate rest, and what is more, given such adequate rest, the individual is benefited by the work, and finds himself as the result of successive periods of work and rest, able to perform a particular act with greater ease and over longer periods without experiencing the sensation of fatigue; if adequate rest be not taken between successive work periods; or if, again, a given action is continued over too long a period, so that the sensation of effort and of fatigue becomes excessive; or, lastly, if a sudden violent effort be made and continued, then the result is overstrain; and, if return to the normal be gained-which is not always the case-it is after a period of rest wholly out of proportion to that needed after mere fatigue. What is more, at the end of this period the organ that had been overstrained, instead of being found stronger from the exercise, is definitely weaker—less capable of responding to a given demand.

The results of such overstrain are various, according to the organ or tissues involved, and according to the grade of work or intensity of the effort that has led up to the state. As already noted, there may be either direct production of morbid states or the development of a state of susceptibility to disease. It is in connection with the most widespread tissue of the body—the muscular—and with overwork of this tissue, that, both clinically and experimentally, the fullest observations have been made. This, then, may be considered in the first place and in more detail, other tissues, of necessity, receiving briefer consideration.

¹Two valuable articles may be especially commended for fuller study, that on "Surmenage," by Marfan, in the first volume of Bouchard's *Pathologie Generale*, p. 445; and that on "Fatigue," by Professor F. S. Lee, Journal of the American Medical Association, May 19, 1906.

OVERSTRAIN

Direct Effects of Physical Overstrain.—With Marfan, we may divide these into (1) superacute; (2) acute and subacute; and (3) chronic.

1. Into the first of these categories enter the cases of sudden excessive muscular action. Of these, we observe various degrees, from the painful dyspncea of the man who makes a spurt to catch his morning train, who suffers from violent heart action and a breathlessness that is almost suffocating—through a severer stage of extreme dyspncea, cyanosis, temporary cardiac dilatation, and irregular pulse—up to fata asphyxia, with death in a few minutes. Such may occur during or at the culmination of bursts of speed or violent effort, the classic example being that of the soldier who dropped dead when he reached Athens with the news of Marathon. Cases have not been unknown in recent times among "sprinters" and other athletes.

It may well be that in this series we deal with two categories; it suggests itself, that is, that the symptoms in the slighter cases are largely referable to cardiac inadequacy, the heart being unable to pass on the blood as rapidly as is demanded by the muscles, so that preëxisting cardiac weaknesses or disease may be regarded as the efficient cause. But these conditions may show themselves in those who, before and after, afford no indications of cardiac disease, the only noticeable condition being that they have been unaccustomed to and untrained to "sprint;" while, again, identical conditions are exhibited in the lower animals. It has been suggested that where death occurs as the result of prolonged intense effort, we have to deal with more than mere cardiac inadequacy, and with a state of intoxication. It is striking that animals hunted to death enter almost immediately into a state of cadaveric rigidity. Authentic cases are on record in which similar immediate rigidity has shown itself in man. During severe engagements, headless cavalrymen-their heads shot off-have retained their seat and been carried over the field by their horses, rigidity developing so immediately that the lower limbs continue to grip the saddle. This rigidity passes off rapidly, and gives place to very early putrefaction, indicating that the antibodies of the organism have been neutralized. Similarly, as noted by Hunter, the blood, dark and venous, fails to coagulate, and, according to Arloing, the capillaries are widely dilated, as though by some vasodilator drug. These facts all point to the presence in the muscles and discharge into the blood of products of muscle activity and dissociation. What those products are we will discuss later. But, granting all this, it must be admitted that cardiac inadequacy, with its attendant asphyxia, dominates the scene.

2. Muscular overwork of a less violent but more prolonged type, while leading to no noticeable cardiac irregularity or symptoms of asphyxia, may set up disturbances of another type. Such cases, for example, we meet with in those going straight from their city life into the country, and indulging, without due training, in the ascent of a mountain or a brisk twenty-mile walk. The symptoms then are extreme and prolonged lassitude, with pains in the muscles that have been

BODILY STATES AS CAUSES OF DISEASE

m

OF;

mi

is

wi

sit

It

is

ac

sti

the

in

ex

aff

thi

qu

wh

wit

un

the

in

wh

one

mo

by

aff

alr

OVE

bec

chi

effe

cha

cat

Fo

net

398

most used, sleeplessness at uight, and, it may be, next day anorexia and a definite low febrile state. Cases are on record in which the fever has been of the typhoidal type, lasting some five or six days and then suddenly disappearing, though usually it terminates in twenty-four or forty-eight hours, and with its termination the urine, previously diminished in amount, containing a large amount of urates, phosphates, and chlorides, becomes abundant and loaded with urea.

3. In addition, there is a certain class of cases in which no one act or series of acts may have seemed excessive, in which, nevertheless individuals performing muscular exercise above the normal eventually experience symptoms which can only be referable to overwork. Such, in those having to walk about and keep on their feet for a large part of the day, are: pains in bones and joints, with slight periarticular swelling, pains in the tendons, and, as seen in adolescence, among boys and young adults, in active exercise without adequate rest, as also in soldiers, what is known as the "irritable heart," a condition of cardiac hypertrophy, with palpitation and more or less marked irregularity of pulse, with signs pointing to mitral incompetency.

In this category, it would seem, are also to be placed the various occupational paralyses which may follow the excessive employment of particular groups of muscles—writer's and pianist's cramp, to mention the most familiar forms, labioglossal paralysis of players upon the flute and other wind instruments. The myopia which is apt to follow excessive use of the eyes is essentially due to exhaustion of the muscles of compensation. The earlier view, that these conditions are primarily nervous, due to exhaustion of particular nerve centres, has given place to the opinion that these states are essentially the outcome of muscular overstrain.

Overstrain as a Predisposing Cause of Disease.—It is a familiar experience clinically that overwork favors infection, that those engaged in hard labor, with late hours and inadequate periods of rest and recuperation, are apt to succumb to tuberculosis, pneumonia, influenza, etc. The difficulty in determining the importance of overwork as a factor in the development of such cases lies in the fact that most often there are associated conditions which also tend to be predisposing factors—inadequate nourishment, foul air, etc. Experimentally, however, as demonstrated more particularly by the studies of Charrin and Roger, it can be shown:

I. That animals subjected to forced labor over long periods (turning a wheel, etc.) are apt to die with naturally developed infections, either through secondary infection of abrasions, or from intestinal infection, it being presumed that pathogenic microbes of low virulence, which, in the healthy animal, live on the skin and mucous membranes without gaining entrance, now in the lowered state of the system manage to gain a foothold. Thus, Charrin and Roger, taking four guinea-pigs, placed them in a cage so constructed with a rotary cylinder that, to keep their balance, they were forced to keep moving; of these four so treated for one or two days, three died in from two to nine days after the experi-

OVERSTRAIN

ment. Smears made from the livers and spleens and cultures from these organs and from the blood gave positive results.

2. That animals subjected to forced labor succumb more rapidly to the effects of the injection of pathogenic microbes than do resting animals, or are killed by doses of the same, which resting animals resist. It is suggestive in this connection to note that sundry organisms of little virulence, which, injected into normal animals, undergo destruction, will gain a foothold, grow, and produce their specific effects, if there be simultaneously injected along with them a small quantity of lactic acid. It may well be that this greater liability of exhausted animals to infection is associated with the increased acid production accompanying muscular activity.

Predisposition of another order is well exemplified in these effects of overstrain. It may be laid down as a broad principle that such overstrain is apt to tell especially upon the parts which bear the brunt of the strain. The most familiar and striking example of this principle in action is seen in connection with the heart. During fortal life the burden of the circulation is borne by the right heart; in postnatal existence, by the left. We find, accordingly, that foetal heart disease affects the valves of the right heart, postnatal heart disease those of the left. With the greater intracardiac pressure, greater strain is thrown upon the valves of the one or other side, and these, in consequence, are more liable to become damaged, and, as a result, lesions, whether of an infective or of a purely mechanical origin, are apt to develop. The greater number of the conditions already noted in connection with the direct chronic disturbances set up by overstrain strictly come under this category. Those lesions, in one sense, are directly set up by the action of some strain upon one or other tissue especially involved; in another sense, it is the strain that has predisposed to the lesions, which, it may be noted, as affecting any particular tissue, may be of one order, i. e., diverse noxæ, acting upon a predisposed tissue, lead to more than different manifestations.

In the second volume we discuss at some length the part played by strain in the production of arteriosclerosis, that commonest of all affections of a physical, as distinct from an infectious, origin. This brings to us a second principle deserving notice, namely, that tissues already weakened by other agencies are particularly susceptible to overstrain; or, in other words, what is a simple strain for normal tissues becomes overstrain for those that are damaged. Here, again, the circulatory system affords well-marked examples. It is in the subjects of chronic intoxications, by syphilis, alcohol, tobacco, etc., that muscular effort, accompanied by increased intravascular pressure, is peculiarly liable to cause the production of aneurysms and extensive arteriosclerotic changes. Here, of course, it is the overstrain that acts as the immediate cause of the disturbance, the intoxication as the predisposing.

The Physiological Basis of Muscular Fatigue and Overstrain.— For long years it was held that muscular fatigue was the criterion of nervous exhaustion, and that the grave conditions of writer's cramp

BODILY STATES AS CAUSES OF DISEASE

at t

into

and

trac

lism

with

stim

proc

cult

-is

activ

has

fuse

of fa

com

syste

and

that

proc

gene

in n

from

Al

ing

prod

the f

Co

1.

2.

lesse

of a

may.

resis

muse

the l Ca

pass

that

4.

3.

H

and other occupational palsies were similarly of central origin. More recent studies by Woodworth,¹ Joteyko,² and others have profoundly modified our ideas. It has been shown, in the first place, that the peripheral nerve fibres are practically inexhaustible, and that the extent of the fatigue is identical in a pair of muscles, one of which is stimulated directly, the other through its nerve. Joteyko's experiments indicate also that the reflex centres in the cord are not capable of exhaustion. Accepting these views, there are those who hold that the nervous system must be wholly left out of account, that fatigue shows itself in the muscle fibres themselves. Nevertheless, Sherrington has shown that this cannot wholly be accepted. Selecting a motor centre in the spinal cord influencing a particular muscle, a centre acted upon by several afferent tracts, he has shown that, setting up reflex stimulation of the muscle along one path, he can bring about exhaustion so that the muscle no longer responds, and when this happens, by stimulating along another path to the same centre, the muscle responds as actively as at first. From the earlier studies, we know that the nerve fibers are not exhausted; we see that the muscle fibres are not exhausted. What is "exhausted," says Sherrington,³ is the "synapse," or membrane of junction between the first afferent tract and the motor neuron. It may be recalled that according to the neuron theory the individual cells, or neurons, are independent units; there is no true junction between them; that thus, when a stimulus passes from one to the other, it must be, at most, by contact action between the processes of one neuron and the body or processes from another. It is, suggests Sherrington, at this membrane of contact that repeated stimuli lead to physical and chemical changes, whereby the conducting power is modified and the nerve current encounters increasing difficulty in its transference from the one cell to the other.

Sherrington is so sound an observer that his experiments must be accepted, and from them it is difficult to arrive at any other conclusion than the above; nay, more, his conception of the cause of the difficulty in passage of the nerve current is in harmony with what we know regarding the hindrance to the passage of the electric current through an arc formed of different elements. While in his experiments there resulted no direct muscular exhaustion, we know, from abundant experiments, that it is possible to fatigue muscle fibres by direct stimulation. The only satisfactory conclusion, therefore, is that there are two orders of fatigue: (1) The immediate or direct muscular fatigue, brought about by the using up of muscle substance in the course of its activity, or, more exactly, due to the inhibiting action of the products of contraction; and (2) what we may term "conductive" fatigue, the neurons, as such, not being worn out, but increased obstruction being established

¹ New York University Bulletin of the Medical Sciences, 1:1901:133.

² Art. Fatigue, Richet's Dict. de Physiol., Paris, 1904.

³ Schaefer's Text-book of Physiology, 2:1900:831; and Journal of Physiology, 34:1906:12.

OVERSTRAIN

at the synapses, or, more broadly, at the sites at which the neurons come into closest communication.

Here, however, as regards the first of these, we must clearly distinguish between two allied but distinct conditions: muscular exhaustion and the sense of fatigue. Through overwork, undoubtedly, the contraction of the muscles becomes hindered by the products of metabolism. This can be demonstrated by repeated direct stimulation of a muscle until it fails to respond. If such a muscle be now washed out with blood, or even with salt solution, it very rapidly responds to further stimulation. In such cases the muscle is put out of action by the products of its own activity. On the other hand, the increasing difficulty in voluntarily repeating a given muscular act-the sense of fatigue -is of central origin, and due to the action of the products of muscular activity, whether directly or reflexly on the nervous system. As Mosso¹ has shown, if a dog be fatigued by a long run and his blood be transfused into another dog, that second animal exhibits all the phenomena of fatigue-dyspnœa, rapid heart action, etc. It is clear that the blood comes to contain substances having a deleterious effect on the nervous system and the tissues in general. Experiments by Zuntz, F. S. Lee, and others show that these products are largely of an acid naturethat sarcolactic acid, potassium monophosphate, and carbonic acid produce similar effects upon the isolated muscle and the organism in general; in other words, that the sense of fatigue is brought about by a mild form of acid intoxication. More particularly, it would seem that in muscular activity it is the glycogen of the fibres that is used up, and from this the sarcolactic acid and the carbonic acid would in the main appear to be derived.

Absence of glycogen, as in the diabetes produced by phloridzin poisoning and inhibition of further glycogen metabolism by the presence of products of muscle activity, leads to a like muscular weakness and exhaustion.

Conclusions.—Thus far, then, it would seem that we must accept the following conclusions:

1. The nerve fibres as such are incapable of fatigue.

 By direct repeated stimuli muscles can be made fatigued, their lessened response being due largely to the accumulation of the products of active function.

3. The progressive difficulty, in response to successive reflex stimuli, may, under certain conditions, not be due to exhaustion, but to increased resistance to the passage of stimuli from one neuron to another.

 The sense of fatigue is due to the accumulation of the products of muscle activity in the circulating blood and the action of the same on the higher centres.

Can we accept unreservedly Joteyko's observations that stimuli may pass through a nerve cell without leading to its exhaustion, to indicate that there is no such thing as nervous fatigue?

¹ Verhandl. d. Internat. med. Congr., Berlin, 1890; 2; Pt. 2; 13, 26

BODILY STATES AS CAUSES OF DISEASE

Personal experience tells us that the mental activities are capable of being overworked; not merely does attention become fagged (which might be ascribed to fatigue of the accessory muscles of eye, ear, and other sense organs-and not necessarily, therefore, to fatigue of the nerve centres themselves), but even in the domain of pure reason the philosopher also is apt to exhaust himself. Nor would this appear to be wholly a matter of synaptic resistance. The one definite series of observations we possess bearing on this nervous fatigue is that initiated by Hodge,¹ and expanded and confirmed by Vas,² Gustav Mann,³ Lugaro,⁴ and others. Histologically, that is, it can be shown that the nerve cells controlling the wing muscles of the bee present recognizable differences between their state in early morning, after the night's rest, and at night, after hours of active flight. Like distinctions are to be made out between resting motor cells of higher animals and those that have been repeatedly stimulated to induce muscular activity. (See Fig. 11, p. 48.)

If, therefore, recognizable differences can be made out in the size and appearance of the cell body, the Nissl granules, and even the nucleus, it is difficult to believe that the nerve cell itself is incapable of fatigue, even if the nerve fibres are; there must be exhaustion of the cell and nuclear matter, which, beyond a certain point, makes itself felt.

As regards glandular and other organs, so little has been determined along these lines that, at most, we can apply by analogy like conclusions.

CELL DISUSE AND LACK OF ACTIVITY AS A CAUSE OF DISEASE.

In an earlier chapter, discussing the states of cell activity, it was pointed out that cell disuse, equally with cell overwork, led to alterations in cell constitution. Here, it may be added that we have indications that this disuse predisposes to disease, just as it may be a direct cause of morbid conditions. We shall but call attention to these matters as briefly as possible.

Cell Disuse as a Direct Cause.—We have in the chapter just referred to pointed out that continued disuse of functional cells tends eventually to complete atrophy and death of the same. Such atrophy and death, if widespread, is apt to destroy the equilibrium between the different tissues of which the organism is composed. Atrophy of a part, in short, has the same effects as removal of that part, and, in the case of the organs supplying an internal secretion, induces identical metabolic disturbances.

The state of the cells in disuse atrophy approaches closely to that

seen it is in the of f nut three of t larly we not

-ar D

For wan gene the nerv satis whe and lacr kept orbi invo asso nutri from to be T effec

whice those name and

¹ American Journal of Psychiatry, 1:1888:479, and 2:1889:376, and Journal of Morphology, 7:1892:95.

² Archiv f. mikroscop. Anatomie, 40: 1892: 375.

³ Journal of Anatomy and Physiology, 29: 1894: 100.

⁴ Lo Sperimentale, Sez. biol., 49: 1895; 159.

DISUSE AS A PREDISPOSING CAUSE OF DISEASE -

seen in simple atrophy resulting from reduced blood supply; indeed, it is difficult to say whether, in the atrophies of inanition, the reduction in the size and the number of the cells of different tissues is to be attributed primarily to impoverished nutrition, or, on the contrary, to lack of functional activity; for, while adequate blood supply favors adequate nutrition, so also functional activity leads to improved circulation through a part, as also, within normal limits, it leads to a healthy state of the nourishing medium—the blood. It is, however, more particularly in these cases in which the nerve supply of a part is cut off, that we in the main encounter disuse atrophy. An organ, such as a muscle not stimulated by nervous influences, affords the most striking example —and commonest—of this type of lesion.

Disuse as a Predisposing Cause of Disease .- The last word has still to be said regarding the means whereby disuse predisposes to disease. For a considerable period this was ascribed in the main to the action of want of action of trophic nerves, which were supposed to govern the general nutrition of the tissues. To this, for example, was attributed the inflammation of the cornea following section or paralysis of the fifth nerve. But no incontrovertible demonstration has been afforded of the existence of such trophic nerves; the above noted keratitis may more satisfactorily be ascribed to the resulting insensibility of the cornea. whereby irritant dust, etc., settling on its surface, is allowed to remain and is not reflexly swept off by the eyelids or by increased flow of the lacrymal fluid. Experimentally, it is found that where the evelids are kept closed, or the surface of the eyeball is protected by covering over the orbit with a watchglass, no keratitis results. In herpes zoster, which involves the area of distribution of particular cutaneous nerves, it is found that there is a lesion of particular posterior root ganglia. This association does not necessarily demand that we deal here with either irritation or paralysis of trophic nerves. Lack of coordination between nutrition, vascular supply, and cell activity under the influence of direct stimuli, together with the lowered condition of cell vitality, resulting from this want of coordination of the cells of a tissue cut off from central control, would appear sufficient to explain the liability for such tissues to become more easily subjected to inflammations and infections.

This lowered vitality from disuse, it must be laid down, appears effective in increasing the susceptibility to infection in parts also in which the nerve supply is intact. It may be pointed out that it is in those regions of the lungs which, from their position, are least in action, namely, the apices, that the tubercle bacillus most easily gains lodgement and growth.

CHAPTER XIV.

PREDISPOSITION AND SUSCEPTIBILITY.

ALL living matter depends upon its environment for its continued existence; upon the stimuli which act upon it from without, whether these be of a chemical or a physical nature, and from its constitution it is so adjusted to that environment that life is only possible within a comparatively narrow range of intensity of stimuli. If this be increased beyond a certain point, it becomes irritation, causing injury; beyond this, again, it renders life impossible, and death ensues. But, in the course of their development under varying environments the different forms of life have come to respond to different agencies in varying degrees—a temperature, for example, which is a stimulant for one form, favoring increased metabolism and increased growth, may be fatal for another; and when we come to the members of the same species, we note at times similar differences; in fact, it may be laid down that no two individuals respond identically to influences acting upon them from without, from the grosser chemical influences of foodstuffs absorbed to the intangible influences of psychical impressions. And even in the individual himself the different tissues present different grades of reaction to stimuli or irritants of one and the same order. Such heightened sensitiveness to stimuli or irritants above what is normal for the species, the tissue, etc., we speak of as susceptibility, or, more narrowly, as predisposition, by this last term indicating that, constitutionally, there is a liability to be more affected by particular influences than is usual; and, pathologically speaking, whenever either of these terms is employed, it indicates an abnormal liability to be so influenced that the development of morbid conditions is favored.

Such predisposition may be either (1) inherited, or (2) acquired. In our discussion of inheritance we have already referred to this (pp. 160 and 210), and here need but briefly note that inherited predisposition may be: (a) Specific or ex-specie (e. g., cattle are peculiarly susceptible to contagious pleuropneumonia; dogs to distemper; genorrhea, typhoid, and the main exanthemata affect man alone). (b) Racial (e. g., those of European origin are susceptible to yellow fever, the Hebrew race, predisposed to diabetes, etc.). (c) Familial, as to scarlet fever, measles, tuberculosis, to particular neuroses, and weaknesses of individual tissues (Friedreich's disease, pseudohypertrophic paralysis), etc., to metabolic disturbances, gout, etc.

We have so fully discussed the subject of heredity that here it is unnecessary to enter again into the principles involved. But we would in passing note that, as regards all these forms of inherited predis-

hav the app for the the race ever of a whe afte As in f bac ato of t mic imr with reg sive the gra que I gest are tive of t diet diet inci inh and to s the

INDIVIDUAL PREDISPOSITION

position, more particularly the susceptibility to infectious diseases, we have to weigh with some caution the evidence that is presented to us: the specific susceptibility may not be so marked as on the surface it appears to be. That there is such specific susceptibility, we do not for a moment pretend to question. Certain influences tell more upon the cells of one species and of one family than on those of another, and the "survival of the fittest" is fitted, to some extent, to explain how races long accustomed to liability to infection by one disease may eventually show but a small percentage of cases of infection, and those of a milder type. What we would more especially point out is that, where a disease is endemic in a region, it is probable that a certain proportion of the inhabitants do not exhibit inherited, but acquired, immunity; that these inhabitants have suffered from transient, unrecognizable, or unrecognized, attacks of the specific disease, which have thereafter protected them. They may not even have had definite "attacks." As Meltzer has pointed out, if the wandering cells are constantly passing in from mucous surfaces, bearing with them individual bacteria, which bacteria tend to be destroyed, even if virulent, provided the number at one focus or place be not too great; then immunity may be brought about, not by the supervention of a mild attack, but by accustomance of the tissues to repeated minimal doses of the toxins of the specific microbes thus introduced. The negro, for example, may not owe his immunity to yellow fever entirely to heredity. Indeed, this has of late years been conclusively shown by Koch and others in connection' with malaria, that the apparent immunity of the natives of malarial regions is explicable by the fact that the young children become extensively affected; the malarial organism may abound in the blood without there being pronounced indications of an acute infection. To such a gradual process of immunization, without definite attack, Sir James Paget explained the freedom of the hardened frequenter of the postmortem room from the blood poisoning which may overtake the infrequent performer of autopsies. To it, also, we may add, is to be ascribed largely the immunity of the practitioner to epidemic disorders.

In this connection, certain observations of Hankin are, at least, suggestive. Rats as a species are refractory to anthrax; even young rats are little influenced. Taking a brood of newly born rats and feeding one moiety with the ordinary mixed food of these animals, with relatively large amounts of meat, the other moiety with bread and milk, he found the former moiety relatively insusceptible, while all the members of the latter succumbed to the disease. Here we have the influence of diet upon the bactericidal properties of the tissues. It may well be that diet and state of nutrition are factors helping to explain the relative incidence of disease among various races.

(d) Individual. While many forms of individual predisposition are inherited (p. 210), it is not at all times easy to distinguish between these and acquired conditions. Here it will be more serviceable, not so much to seek to attempt to distinguish between these two orders, as to classify the different forms of individual predisposition.

PREDISPOSITION AND SUSCEPTIBILITY

We note thus predisposition according to:

 Sex.—It need but be remarked that the sexual life of the female exposes her to the liability to contract a special series of disorders in connection with menstruation, childbirth, and the menopause.

 Life Period.—The liability to the incidence of particular disorders at different life periods may be briefly expressed in the following table, the brackets indicating the year or period at which the condition in question is most fatal:

Infancy.—Disorders of maldevelopment and inanition (to the end of first year); athrepsia, various forms of enteritis with diarrhoea; meningitis.

Childhood.—Rickets, measles, scarlatina, diphtheria.

- Puberty and Adolescence.—Chlorosis (in female); acute rheumatism and rheumatic heart disease (ten to fifteen); typhoid; tuberculosis.
- **Adult.**—Typhoid (twenty to twenty-five); tuberculosis (twenty to thirty).
- Middle Age.—Gout, lithiasis, and chronic Bright's disease (thirty-five onward); arteriosclerosis, aneurysms (thirty to fifty); cancer (forty to sixty).

Old Age.—The same continued along with atrophic conditions; cerebral apoplexy (fifty-five to seventy-five); low infections.

The immaturity of the cells and tissues, the fact that they have not become immunized, coupled with the fact also that particular tissues are undergoing either great strain or very rapid growth, would seem to explain to a very considerable extent the susceptibility of infants to digestive disturbances and meningitis. The weight of the brain is doubled in the first year of life (from 400 to 800 grams).¹ Like considerations would seem to explain the susceptibility of infants to digestive disturbance and of children to acute infections, the tissues being more vulnerable when first exposed to the toxic influence of fermented food, and to the influences of sundry specific microbes. There are, however, certain features in connection with the age incidence of infectious disease which must be pointed out. Both in animals and in man we observe that the newly born and the young are little affected by-in fact, are immune to-certain diseases which cause a high mortality in those of older years. Babies under three months old scarcely ever suffer from diphtheria, and this not because they are not exposed. The same is true with regard to young children and typhoid. While this is somewhat more common among the young than is usually supposed, it is almost unknown among infants, and in such young children as it attacks it causes in general a mild disease, compared with what we find in young adults. Many similar examples may be called to mind.

A priori, the more immature, the more unprepared the cells, the more vulnerable we should expect the tissues to be, but clearly this is not

¹ F. W. Beneke, *Die Altersdisposition*, Marburg, 1879,

alwa by t whi app part that stin it n with sub in t pro ent the not ord pre " c(loc the cor of lial exe clo the wh fui gri in the mi ore su tu

> ty] wa

always so. That certain tissues, liable especially to primary infection by the diseases in question, are more active and reactive in early life, while later they become exhausted and more susceptible, would not appear to afford a complete explanation, though that this may be a partial factor cannot be neglected. There is yet another possibility indicated by the essential nature of toxicity and infection. For a substance to be toxic and injurious to a cell, it is necessary that, entering that cell, it sets up such a molecular disturbance as either to arrest or to stimulate excessively the metabolic processes of that cell, or otherwise it must enter into chemical relationship with the biophores. It is well within the bounds of possibility that, as suggested by Abbott, a diffusible substance which sets up excessive molecular and destructive disturbances in the fully developed cell may have but little effect upon the more inert protoplasm of the immature cell, and that if certain bacteria gain entrance into the tissues, the cell may digest and otherwise destroy them, their toxins not combining with the biophores, and, as a consequence, not arresting the cell functions.

3. Habit of Life at Different Life Periods.—During infancy and early childhood the digestive organs are relatively most active, and, in order to bring about the absorption of food necessary for rapid growth, they are peculiarly liable to strain. The lack of power of locomotion prevents much mingling with others and the extensive exposure to "contagion" and the zymotic diseases which supervene with active locomotion and mingling with other children. With adolescence and the forsaking of an outdoor and active, for a more sedentary and confined existence in workshops and other places, where large bodies of men are collected and ventilation often defective, tuberculosis is liable to supervene. With increasing corpulence and inability to take exercise in middle age, constipation, gallstone formation, etc., tend to be favored. Other environmental influences, such as those of climate, clothing, and social influences, come under the same category.

4. **Previous Infection.**—While many diseases, more particularly the acute exanthemata, are followed by immunity, there are others in which this immunity is but short-lived; others, again, like eryspelas, furunculosis, acute rheumatism, and, we may add, influenza or la grippe, in which one attack actually predisposes to a second. Whether in these cases, the germs of the disease are not all destroyed, but some linger in the system and exhibit themselves actively if anything lowers the vitality, or whether a new infection occurs, is not precisely determined. It may be that either occurs.

What is even more noticeable is that an attack of one infectious disorder is frequently followed by an infection of a different nature. The tissues are weakened by the one disease, and in this condition are more susceptible. Thus, the acute exanthemata may follow one another and tuberculosis supervene upon any of them, or upon typhoid.

 Malnutrition.—'The terrible mortality from infectious diseases typhus, relapsing fever, typhoid, septicemia—which has followed in the wake of famine in Russia, India, Ireland, during the last century, is an

of :

bac

doi

fac

ent

gre

me

live

is

thi

oth

str

res

(1)

or

rea

de

ex

SIL

SPI

po

up

adequate example of the effects of malnutrition in predisposing to disease. Here may be included local malnutrition, such as that brought about by lessened functional activity, due to impaired nerve supply. Paresis and paralysis, with imperfect function, imperfect metabolism, and weakening of the tissues predisposes to local infections and suppurative disturbances. Impaired blood supply has similar effects. To these we shall refer more fully in a subsequent paragraph. The effects of overstrain in predisposing to disease have been discussed in the previous chapter.

Tissue Susceptibility.—Just as the general susceptibility of the organism as a whole is noted to be increased by the means just indicated, so we can increase the local susceptibility of the different tissues and favor the growth of microbes within them, or the development of functional disorders, by injury, by malnutrition (impaired blood supply or nerve supply), and by lessened functional activity. To these, indeed, we have already referred in passing. But apart from this, which we may term acquired tissue susceptibility, we have also to recognize an inherent susceptibility to disease on the part of various tissues.

It is a matter the significance of which is too little recognized, that very many pathogenic organisms show a predisposition to grow in special tissues; or, more correctly, that certain tissues exhibit a particular predisposition to permit these to grow within them. As regards the primary focus of infection, we see that the channel of entrance in general affords a partial explanation why this should become the seat of growth, that inhaled germs should especially affect the respiratory system, ingested germs the digestive tract. But even here it has to be noted that the diplococcus pneumonia, for example, has little effect upon the pharyngeal mucous membrane, growing there as a harmless saprophyte, whereas the diplotheria bacillus causes intense disturbance.

When we pass beyond, to the secondary foci, this tissue predisposition is still more marked. The tubercle bacillus flourishes in the lung, upon serous surfaces, in the different glandular organs, but rarely in muscle; infrequently in the brain substance, as compared with the pia-arachnoid; infrequently in the stomach wall, as compared with the small intestines; in the epiphyscal ends of bones, and in the joints of the young, but not commonly in the same regions in those of mature years. If the colon bacillus be injected into the blood stream, it sets up more especially a condition of acute enteritis. In like manner, passing in review each separate infectious disease, this specific tissue susceptibility can more or less definitely be pointed out.

This specific tissue susceptibility, then, is well marked; different pathogenic agents find in different tissues circumstances specially favoring their growth; the cells of these tissues react less perfectly against these specific bacteria. In certain individual cases explanations may be suggested for this predisposition. In tuberculosis, for instance, the acid production of the muscle, stomach walls, and brain substance has been suggested as explaining why the bacillus in general does not thrive in these organs—although Ficker's observations, that of all media the acid brain matter furnishes that upon which the tubercle bacillus flourishes most rapidly and abundantly, may well make us doubt this. In other cases, the nature of the circulation through the tissues implicated has been invoked. But that does not satisfy the fact that particular microbes may multiply in these particular tissues. In short, it is not possible to find one common basis of explanation for this selective action beyond this, that in each tissue there is a varying environment, and certain environments are specially favorable for the growth of special bacteria.

It is the corollary to this condition of tissue susceptibility that deserves more recognition. If the typhoid bacillus is found growing in the spleen, liver, skin, and kidneys, and is difficult to isolate from other organs, it is obvious that to reach these particular organs it must have travelled through the blood stream, and have been equally liable to enter the others. So, also, if in young children the ends of the bones become tuberculous, obviously the tubercle bacilli must have entered the blood stream and have been carried through the system generally before reaching these remote regions. We are thus bound to conclude either: (1) that infectious microbes circulate passively through the various organs, in which they show no growth, causing no reaction until they reach a susceptible tissue, or (2) that, circulating thus, they tend to be destroyed in every other tissue of the body save those that are susceptible. The first alternative is not only eminently unlikely, but is negatived by experiments, which show that the vascular endothelium of organs like the liver, which later may exhibit no special foci of disease (i. e., are not susceptible), actively takes up and destroys pathogenic bacteria. The second, thus, is the only adequate deduction from the facts at our disposal; and this leads us, further, to a very important conclusion, that in infection the body is never involved as a whole. Coincidently with the growth of the specific germs in individual organs, there tends to be a reaction to, and destruction of, the same in other parts. The bearing of this upon the recovery from infection we shall point out later.

We may now sum up the various forms of predisposition; they are: 1. *Inherited*:

- (1) Specific, characterizing the species.
- (2) Racial.
- (3) Familial.
- (4) Individual, as regards
 - (a) sexual incidence;
 - (b) age incidence;
 - (c) tissue incidence.

2. Acquired, as a consequence of

(1) Social and environmental conditions.

- (2) Injury.
- (3) Malnutrition.
- (4) Previous attacks of
 - (a) the same disease:
 - (b) other infectious disease
- (5) Exhaustion.

PREDISPOSITION AND SUSCEPTIBILITY

Idiosyncrasy.-This term is applied to the exhibition of extreme susceptibility on the part of the individual to the influence of substances which not only have no disturbing action upon the normal individual. but often are to him the source of distinct pleasure or benefit. It is an extreme form of susceptibility, and that manifested in unusual directions. Thus, some individuals are unable to eat sundry not unusual articles of diet (strawberries, porridge, certain shellfish, mackerel, or other fish) without a train of symptoms showing themselves, which seem to indicate a distinct grade of intoxication, manifested by urticaria and abnormal skin eruptions, headache, running at the eyes, abdominal disturbance, etc. One well-known London physician of our acquaintance, recently deceased, dared not take the pudding or cream at dinner away from home, for fear it be flavored with ginger, the least trace of which gave him acute misery for the better part of the next twenty-four hours. Similarly, there are drug idiosyncrasies, often accompanied by skin eruptions-from quinine, potassium iodide, opium, iodoform, and so on. Some of the most remarkable are in relation to animals, most commonly with cats; the presence of a cat in the room, even if unknown to and unsuspected by anyone, and hidden from sight, being sufficient to cause intense discomfort, and even a state of nervous terror, that is painful to the individual and all around. This, as Weir Mitchell¹ has pointed out, can only be due to the action of some emanation from the animal upon the sensitive olfactory mechanism, even though, in most cases, the affected persons state that they perceive no special odor. It thus becomes allied to that other more common idiosyncrasy, hay fever, in which, again, no odor is necessarily perceived, but in which it has been fully proved that the fine, floating pollen of flowering plants and grasses is the irritative agent, the intense coryza and discomfort making its appearance at the period when plants are in flower, and disappearing if the individual take a sea voyage or by any other means escapes to where little pollen is likely to be. The tendency nowadays is to regard some at least of these idiosyncrasies as anaphylactic phenomena. (See Section III, Chapter VIII.)

I

pas

we

iec

rec

det

pro

we

inc

an

the

ca: we

the

the

oth is sp co se or ar vi to of

¹ Trans, Assoc. Amer. Phys., 20; 1905; 1.

SECTION III.

THE MORBID AND REACTIVE PROCESSES.

INTRODUCTORY.

HAVING in the previous section discussed the causes of disease, we pass now to discuss how these causes act, and, doing so, observe that we can, from the point of view of general pathology, approach our subject from two directions; we can, that is, studying disease generally, recognize, underlying its various manifestations, certain common series of events, or morbid processes-processes which, it is true, vary in their details in individual cases; nevertheless, the broad features of groups of cases are alike, and once we establish that morbid conditions constituting a particular group are allied and have a common basis, we can proceed to inquire what it is in any particular case that leads to what we may term divergence from type; or, on the other hand, rather than inquiring into the course of the different processes, we can make the tissues and the changes they undergo the main object of our inquiry, and classify thus the morbid changes affecting these tissues, rather than the morbid processes proper. It may well be urged that in the latter case we are dealing with the results of disease. All depends upon how we approach the treatment of the subject. If we seek purely to describe the histological alterations in the cells brought about by disease, then these conditions should not be dealt with in this section. If, on the other hand, we study the succession of changes leading ultimately to the different morbid states seen in the cells and tissues, then our study is that of processes. Further, it may be propounded that, broadly speaking, the changes in an individual cell do not constitute disease (as generally accepted); that depends upon the cumulative effect of the combined cell disturbances, and more than that, upon the disturbances set up by the perverted activities of these cells upon other cells and organs not primarily involved. Seen in this light, morbid cell changes are factors in the production of particular results. It is, however, a minor matter under which heading we consider these conditions, provided their nature be recognized. In dealing with them we shall have to discuss the changes leading to the production of each individual form of cell disturbance, and at the same time to describe the resultant effect of these changes upon the individual cell; conformably with usage, we

THE MORBID AND REACTIVE PROCESSES

class them with the morbid processes. We take up the discussion of the same between that of the morbid processes proper and that of the results of disease upon the tissues and the organism as a whole.

In the heading to this section we refer to the morbid and reactive processes. It will be seen that, with the exception of conditions of arrest of cell activity and cell death, and the possible exception of neoplasia, or of some neoplastic conditions, all morbid processes are at the same time reactive; they are the expression of the reaction on the part of the tissues to noxe of various orders.

Thus, to repeat: In the first part of this section the morbid and reactive processes proper will be discussed; in the second, the morbid cell changes.

> I the tion the As of cor lea ess the doc has stu eas wa tor In tui nit læ eig it an ing tra the of sta

PART I.

THE MORBID AND REACTIVE PROCESSES PROPER.

CHAPTER I.

THE LOCAL REACTION TO INJURY; INFLAMMATION,

For the development of the sound pathologist, a full knowledge of the factors concerned in the inflammatory process and a right appreciation of the doctrine of inflammation is as essential as to the orthodox theologian is a right attitude in respect to the doctrine of the Trinity. As regards the one, there have been bitter fights and wide divergences of opinion; so with regard to the other—and these divergences in both continue to exist, and with them a wide tolerance. Nevertheless, the leaders of the pathological world, though they may differ in nonessentials, are at the present time in substantial agreement regarding the subject of inflammation. There is an orthodox doctrine, and that doctrine we shall proceed to expound.

Definition .- The condition of local "flaming"-inflammationhas of necessity been recognized from the very beginning of medical studies; but so long as little was known concerning the causes of disease and less regarding the processes, all that could be accomplished was to regard this as a state characterized by certain particular symptoms, and the first attempt at a definition, that of Celsus, so regarded it. Inflammation, he laid down, was a condition characterized by "rubor, tumor, calor, dolor"-redness, swelling, heat, and pain-to which definition later writers added a fifth cardinal symptom, that of "functio læsa"-disturbed function. The great English pathologist of the eighteenth century, John Hunter, said that it was something more, that it was a process rather than a condition-a process set up by injury, and tending toward counteraction of the same. But he was before his time. Three-quarters of a century elapsed before Cohnheim, employing a microscope to study the changes ensuing in an injured area-the transparent web of the frog's foot-saw and first described with accuracy the changes that occur in the vessels of the injured part-the dilatation of those vessels-the eventual slowing of the current, and, it might be, stasis, or complete arrest-the exudation of fluid and migration from

cyt

me

ter

sav

DO

inf

to

we

an

cel

tu

re

an

tic

at

pa

re

of

se

th

er

in

al

W

al

te

of

p

al

01

M

them of leukocytes; saw in these the full explanation of all the cardinal symptoms, and concluded that the vascular changes were the essential feature of the inflammatory process. And, undoubtedly, in tissues well supplied with vessels, these are the most striking feature, the feature dominating the whole field following upon injury, whether mechanical or chemical (by caustics or bacterial poisoning) above a certain grade. But with a lower grade of injury, by it may be, identical noxæ, these vascular changes are little noticeable, and are replaced by connectivetissue overgrowth; there is still injury, but the results differ, or rather correspond to those seen in the later stages of the process that follows injury of a severer grade; and secondly, a like order of end results is attained in tissues that are unprovided with vessels. It was these circumstances that led Burdon Sanderson, in the "seventies," to advance a broader definition, namely, that "inflammation is the succession of changes occurring in a part, as a result of injury, provided that that injury be not so excessive as to destroy the vitality of the part," or, briefly, and better, that it is "the (local) reaction to injury" (for some of the changes that occur in an injured part, hemorrhages, etc., are not reactive, and play no direct part in the inflammatory process). In Germany, Marchand in the early "eighties" propounded similar views.

From this date onward there has been an increasing recognition of the fact that the various processes which together make up the picture of inflammation are not merely the passive and deleterious outcome of local injury, but, on the contrary, are factors which tend to counteract the local effects of that injury; an increasing conviction that the distinction so much insisted upon by the older writers between inflammation and repair is non-existent, and several writers of this generation-Leber,¹ Neumann,² Councilman,³ and others-have advanced definitions containing a clause to the effect that the tendency of all those processes is in the direction of repair of the injury. We ourselves have elsewhere stated⁴ that inflammation "is the series of changes constituting the local manifestation of the attempt at repair of actual or referred injury to a part;" or, briefly, is "the local attempt at repair of actual or referred injury." Grawitz expresses the like idea in his definition of inflammation as the reaction of irritated and damaged tissues which still retain vitality.

To Metchnikoff,⁵ through his most remarkable and long-continued studies upon the functions of the leukocytes in disease, and upon phago--

¹ Ueber die Entstehung der Entzündung, Leipzig, 1891.

² Ziegler's Beitr, 5: 1889; 345.

³ Article Inflammation, Dennis' System of Surgery. See also, in a similar sense, H. Buchner, Fortschritte d. Med., 10:1892;363; Marchand, Wundheilung, Leipzig, 1900; Chantemesse and Podwyssotsky, Les processus généraux, 1905; Bier, Die Stauma als Heilmittel, 1904.

⁴ Article Inflammation, Allbutt's System of Medicine, 1:1896.

⁸ La pathologie comparée de l'Inflammetion, Paris, 1892. A work of the highest value, to which we shall frequently refer. It has been translated into English by Starling, 1893.

DEFINITION

cytosis, both by leukocytes and the fixed cells of the organism, we owe, more than to any other single individual, the recognition of these counteracting forces. Just as Cohnheim, studying the vascular changes, saw in them the essential factors in the inflammatory process, so Metchnikoff, studying the leukocytes, would elevate them to an exclusive position in his definition, and would lay down that phagocytosis is inflammation. This involves too wholesale a neglect of other factors to be acceptable. "In studying the reactions of the organism to injury. we must be impressed by the multifariousness of natural processes; the end may be attained, not in one way only, but in many. It is not by cells of one order alone-by phagocytes-or by leukocytes in general, and only leukocytes, or merely by the reaction on the part of the fixed cells of the tissues, or by vascular changes alone, or by altered temperature, or solely by the chemical and mechanical action of the exudate, that repair is effected. All means are employed to antagonize the irritant . . . now the one, now the other more parand to effect healing . . . but none exclusively." ticularly

Such processes must not, we would insist, be regarded as primarily purposeful; they are adaptive. What we understand by adaptation we have already laid down on p. 117.² And considering them as adaptations, we arrive at a further simplification of our definition, namely, that inflammation is the succession of changes which constitute the local effort at adaptation to the changes initiated by actual or referred injury to a part; or, in short, the local adaptive changes resulting from actual or referred injury.

More recently there has been a tendency to widen still further the scope of the inflammatory concept. It is seen that with local injury there are set up not merely local changes, but that tissues at a distance are also involved, and contribute toward counteracting the irritant and its effects; that, for example, there may be fever, the lymph glands may become enlarged, the bone-marrow stimulated to activity, with resulting increase in the number of circulating leukocytes and in the amount of antitoxic and protective substances in the blood serum. Thus, Ribbert³ lays down that inflammation is "a vital manifestation on the part of the whole body, if primarily of the local tissues that have undergone injury, and is the sum of all the exalted vital processes," and Aschoff, we find, teaches similarly. While we admit, and that most fully, that as a result of injury there may be this participation by the rest of the organism, we feel strongly that to introduce these further processes into our conception of inflammation brings about an immediate confusion between the" phenomena of inflammation and those of the general reaction to injury, and infection. As will be seen later, general reaction on the part of the organism as a whole, with or without marked primary local disturbances,

¹ Adami, Montreal Med. Jour., May, 1895.

² For the distinction between adaptation and purpose, see Adami, Inflammation, Macmillan, 1909: 232.

¹ Die Bedeutung der Entzündung, Bonn, 1905: 11 and 45.

416

is the distinguishing feature of "infection." We would thus emphasize that, to prevent confusion, it is advisable to restrict, as in the past, our conception of the inflammatory processes to those which are of local origin and occurrence.

THE COMPARATIVE PATHOLOGY OF INFLAMMATION.

We cannot better gain a grasp of the main features of the reaction to injury than by studying first the simplest cases, and gradually following up the processes seen in more and more advanced conditions. In other words, it is best to begin with the simple individual cell, with the unicellular organism, and trace the reactions that occur as we advance from the unicellular to the simpler metazoan, or multicellular forms; from these to the higher metazoan forms, and man himself. This we shall do rapidly, not taking up division after division, but noting merely those forms of life in which progressive evolution has led to the introduction of new factors in the reactive process.

Taking first the protozoa, we note that destruction of a portion of the cytoplasm-provided this does not include the nucleus-is followed by the closing together of the uninjured parts. In other words, provided the nucleus be uninjured, the cell recovers; provided also, we may note, that not too large a portion of the cytoplasm be destroyed or removed (p. 43). We note also other properties closely bearing upon the properties of the cells of higher forms of life, including man, namely: (1) The property of chemiotaxis-of being attracted toward certain substances, of being repelled from others; (2) the property of adaptation, whereby a negative may be changed into a positive chemiotaxis. It is found that by the gradual and careful addition to the fluid in which it lives a protozoan can be gradually accustomed to substances from which at first it was repelled, until eventually it will move actively toward stronger solutions of these substances;¹ and (3) the property of phagocytosis, that, namely, of ingesting solid particlesof eating them (&ayzev, to eat). This, at least, is the primary meaning of the term. It is inevitable, although unfortunate, that, with the development of Metchnikoff's studies and the evolution of his "phagocytic theory" into its present form, several other protective properties beyond that of ingestion and digestion have become attached to the conception of phagocytosis. Here we employ the term in its primary sense, and observe that one of three events may follow the ingestion of foreign particles by such a form as the amœba: (1) The particle may prove assimilable, in which case, as shown by Miss Greenwood and La Dantec, it becomes surrounded by a vacuole containing digestive fluid, and undergoes solution, portions proving indigestible being subsequently cast out from some region of the surface of the organism; (2) it may prove unassimilable, in which case, after a short sojourn within

¹ Stahl, Flora, 76: 1892; 247.

the in s hos its may exh libe see, cell I a d pro Bet

> fort end der the a p

> > ra

see

is ca

th

Wa Au


fu

m

THE COMPARATIVE PATHOLOGY OF INFLAMMATION 417

the cytoplasm, it is similarly discharged; or (3) if a living organism, in some cases a symbiosis, or living together, is set up; the protozoan host is not irritated by the parasite, and does not discharge it, and, on its part, the parasite, living upon the material assimilated by the host, may multiply until its progeny so fill the cytoplasm of its host, that they exhaust the foodstuffs, and lead to death of the host, when they become liberated into the surrounding medium. All these events, as we shall see, may likewise occur in the leukocytes and other cells of the multicellular organism.


Passing next to the lowest metazoan forms, we immediately observe a division of labor among the cells, the outer layer being more especially protective, the invaginated inner layer more particularly digestive. Between the epiderm and endoderm there develops a third set of cells,



Larva of one of the simplest metazoan forms (Astropecten) to show ed., ectoderm; end, endoderm; mea., wandering mesodermal cells which at pl, have attached themselves to a foreign body and formed a plasmodium around it. The plasmodium of fused mesodermal cells seen in the previous figure, higher magnification; *nucl.*, nuclei of individual cells, (After Metchnikoff.)

the mesoderm, derived from the other two, which, as we advance, rapidly assumes great importance, forming the connective-tissue layer of all the tissues, the internal skeleton, the muscles, the heart, and, subsequently, the closed vascular system. Here, in the lowliest stages, it is represented by relatively few cells, with long, fine processes, which can be retracted so that the cells are capable of floating about freely in the body cavity. These are, in fact, the earliest representatives of the wandering cells, white corpuscles, or leukocytes of the higher animals. And already in these lowest forms we find them possessing protective functions, though at the same time it must be noted that they are not the only class of cells that react. If any of the living cells be destroyed, those in the immediate neighborhood spread out by anneboid movement to fill the gap temporarily, and subsequently multiply, thus bringing about *regeneration* (and the regenerative process of these

lowest metazoan forms is nothing short of marvellous; a few cells may regenerate the whole animal). But if, as Metchnikoff has pointed out, foreign particles gain entry into the body cavity, it is the wandering cells that deal with them. If small, they may ingest them; if large, attaching themselves to and surrounding the objects, they fuse into a *plasmodium*, or multinucleated mass, thereby fixing it in position; if it be capable of digestion, forming a common stomach, as it were, and bringing about its dissolution; if inert and not digestible, then cutting it off from the body cavity. The power of forming a plasmodium around foreign bodies is possessed by leukocytes, from the lowest multicellular form up to man; one class of giant cells, those, for example, we see in tuberculosis, and those that form around foreign bodies introduced into the tissues are plasmodia of this nature (Fig. 140).



Foreign-body giant cells from man; at a the leukocytes form multinucleated plasmodia around foreign fibres (silk), at b the giant cells have broken away containing debris of the fibres. (Ribbert.)

Throughout the invertebrata we observe these two processes of regeneration and leukocytic activity, with little material addition; we note, indeed, that there gradually appears—in the crustacea, for example—more than one order of wandering cells, with different properties; some are phagocytic; others, as, for instance, those that pass out on to the surfaces of certain invertebrates, are explosive, and apparently with their disintegration liberate digestive and (it may be) bactericidal substances, whereby foreign organisms adhering to the carapace become destroyed and the surface cleansed (Hardy and Alcock). But though there is a heart, with rare exception there is no closed system of bloodvessels. The large vessels open into the body cavity, and the heart but serves to keep the body fluid circulating. Thus not until we reach the vertebrata, does a fully developed vascular system enter as a prominent factor into the reaction to injury. At most, as in the arthropoda, the body fluid has extensive powers of clotting.¹

¹ See Leo Loeb, Jour. of Med. Research, N. S., 2: 1902: 145.

THE COMPARATIVE PATHOLOGY OF INFLAMMATION 419

so that when injury is such as to cause communication between the body cavity and the external medium, the wound may gain a temporary closure through the agency of the coagulated fluid.

Nevertheless, the different classes of invertebrata afford some most remarkable examples both of regeneration' and of leukocytic activity. Of the latter, quite one of the most striking is that afforded by a parasitic disease of the *Daphnia*, or water flea, a minute crustacean, translucent, and just visible to the naked eye, and common in some fresh-water ponds and ditches. Metchnikoff, observing that the daphnias in his aquarium were becoming opaque and evidently dying off, found that the disease was produced by a blastomycete, or yeast, growing in the affected aquarium. This was an elongated form, termed by him *Monospora*, and characterized by producing long, sharp-pointed spores. These spores, taken in by the daphnia, when they passed along the intestine were apt to penetrate the delicate wall, and so gain entrance into the surrounding body cavity. If those so entering were few in number, the leukocytes attached themselves to them, often several together, forming a plasmodium, and eroded and destroved them.

If, on the other hand, too many penetrated into the body cavity, the number of leukocytes was inadequate to deal with all, and some not attacked germinated. Whereas the wandering cells freely attached themselves to the spores, they were seen to leave the germinated torulæ severely alone, with the result that the body became filled with the yeast, became opaque in consequence, and death ensued.

With the vertebrate we find a fully developed and closed vascular system, distinct from the body cavity (lymphatic) system, though the latter receives its fluid from the blood, and yields its fluid back to it, so that there still exists a body cavity circulation, but that secondary to the main vascular system. And with this we find (1) that the vessels of an injured part become directly concerned in the inflammatory process; (2) that here, in consequence, we have "flaming" as the result of injury, the vessels of the part undergoing dilatation, with its attendant results; (3) that the reaction shows itself more rapidly to the naked eye, and, indeed, has a more rapid course; and (4) that instead of the tardy and haphazard accumulation of leukocytes from the surrounding tissues and from the slowly circulating body cavity fluid. with the more rapid flow of blood through the injured area, leukocytes are constantly being carried into the area, and once there, provided the irritant has not too virulent properties, their chemiotactic activities lead them to make their way through the walls of the vessels toward the foreign body.

Undoubtedly, then, in the higher animals the vessels are a most important factor in the reaction to injury; they add greatly to the power of the individual in responding fully and promptly to local injury. It has, however, to be noted that the difference they introduce is one of degree rather than of kind; we find the same underlying local processes

¹ For examples, see the chapter on Regeneration.

at work in the invertebrate as in the vertebrate; we find them proceeding in non-vascular areas in the vertebrates just as in the vascular. There is, thus, no essential difference introduced by the participation of the vessels in the reaction to injury; and that being so, to draw a sharp line of demarcation between the two sets of cases, to say that in the one set of cases we have inflammation, in another we have not, is to make a distinction where no fundamental difference exists. While all the vertebrates possess this closed vascular system, it is deserving of note that in the lowest forms the development is not nearly so perfect as in the higher-the capillaries are not so abundant, nor is the vasomotor apparatus nearly so complete, whether nervous, controlled from the central nervous system, or idiomuscular, depending upon direct stimulation for the dilatation and contraction of individual vessels. Among the urodela, or tailed amphibians, it is very evident that the response to injury is slower, the diapedesis less, the part taken by the wandering cells already in the lymph spaces more noticeable, than it is in the mammalia.

THE CAUSES OF INFLAMMATION.

It follows from what we have said that anything which causes local injury to the tissues is a cause of inflammation, be it a mechanical trauma, a physical insult, as by heat, cold, or electricity, a disturbance brought about by altered metabolism and abnormal internal secretions, or bacterial or microbic invasion and growth. This last is the commonest cause of acute reaction, and differs from the physical and mechanical causes (though not from metabolic disturbances) in that, as a cause, it is not of momentary duration, but of continued. It is not the mere physical entry of microbes into the tissues that induces inflammation, but the liberation by them of their products in growth or disintegration. And so long as those products are being liberated, for so long is the cause in action. It differs from the metabolic causes in that the latter induce tissue irritation of a milder grade, and so do not induce acute, but rather chronic, reactions. But to say, as some surgeonpathologists insist, that because bacteria cause the acute inflammation most commonly encountered and that which most frequently demands treatment, therefore inflammation is always the result of bacterial activity, and anything not caused by bacteria is not inflammation, convicts one, to put it in moderate language, of a certain lack of breadth of vision.1

We distinguish two grades—not forms—of inflammation, the *acute* and the *chronic*. Of each, it is true, we recognize subgrades, and the one order of cases passes imperceptibly into the other, but for practical purposes the division is useful. In the former we have a frank

⁴ I have discussed the heresy in the Medical Record, March, 1896; Middleton-Goldsmith lectures, and the article *Inflammation*, in Keen's System of Surgery,

ACUTE INFLAMMATION EXPERIMENTALLY PRODUCED 421

reaction of rapid development and course; in the latter, a reaction characterized by little evidence of the cardinal symptoms, and slow, progressive course. In the one, the condition unfolds itself in the course of hours or days; in the other, of weeks and months; and thus the salient features in typical cases show marked differences, even though, as we say, the study of intermediate stages convinces us that essentially the two are grades of one and the same process. We shall take up the subject of chronic inflammation later. Here we shall, for the time, confine ourselves to the consideration of the acute type; and first, in order to have a mental picture of the sequence of events, it will be well to describe with some little detail what can be seen when we produce experimentally such an acute inflammation, and follow the successive steps under the microscope.

THE STAGES OF ACUTE INFLAMMATION EXPERIMENTALLY PRODUCED.

As affording the more typical picture, it is best to consider first what happens in a part well supplied with vessels. And two methods are



Inflamed mesentery of frog: a, margination of leukocytes in the dilated capillaries; b, migration of leukocytes; c, escape of red corpuscles; d, accumulation of leukocytes outside the capillaries. (After Ribbert.)

open to us—either to choose some region that is transparent, and injuring one spot in that region, to follow under the microscope the unfolding of the resulting disturbances. This was first done system-

atically by Cohnheim, and there are several such regions which may be selected—the delicate web of the frog's foot, the frog's tongue, which is attached in front only, and normally is directed backward; it can easily be pulled forward and out; or, selecting animals higher in the scale, the mesentery of the cat or one of the animals of the laboratory is admirably delicate and so sensitive to injury that exposure to the air alone suffices to set up pronounced disturbance. Or, on the other hand, we can cause injury, bacterial or otherwise, to some internal vascular tissue, and, removing and studying the injured area from a series of animals at successive periods, can in this way trace the successive steps. The advantages and disadvantages of the methods are obvious.



1, adhesion of leukocytes to the wells of a capillary in an inflamed area; 2, mode of migration of a polynuclear leukocyte. (Lawdowsky.)

The first is particularly useful for following the earliest stages, the changes in the caliber of the vessels and the rate of blood flow; even the relative position and movement of certain cells can be well observed; but the intervening skin or endothelium, along with the absence of nuclear staining, renders a study of the finer cell changes impossible. The animal has to be immobilized by some drug, or by "pithing," which may well lower the excitability of the tissues, and prolonged examination is virtually impossible. The second gives permanent preparations, which can be studied at leisure, and permits a very thorough study of individual cell forms, but often leaves us in doubt as to whence the cells come; to be carried out thoroughly requires the sacrifice of a large number of animals, and even then leaves us in doubt whether some important stage may not have been

passed over; while, owing to the mode of preparation, the vascular relationships and the blood contents of the part fail to be retained exactly as in life. The two combined afford us a very full picture.

Combining the results gained by these two methods, we gain the following picture:

The first result of local injury is temporary *contraction of the arteries* of the part, leading to a brief period of lessened blood supply.

This is succeeded slowly by progressive *arterial dilatation*, so slow that it cannot be of reflex nervous origin; indeed, it shows itself where the nerves to the part have been severed, and, as a result, within an hour or so (in the frog's web, injured by the application of a caustic)

ACUTE INFLAMMATION EXPERIMENTALLY PRODUCED 423

there is a greatly increased blood flow through the part, the capillaries become distended, and many, previously empty and unrecognizable, are now visible with the blood corpuscles hutrying through them.

Gradually—in the course of the next hour (in the frog)—the blood stream becomes slower. Whereas, at first, the individual corpuscles in a capillary were unrecognizable, all that was visible being a darker central band, or current, with clearer fluid (blood plasma) on either side, now the band becomes broader, the side currents narrower and nar.ower, and with this, as the slowing continues, *individual blood* corpuscles can be distinguished. And, what is more, the two forms can be made out (this especially in the frog, in which the red corpuscles are relatively much larger). And it is seen that the white corpuscles more particularly are carried along the outer part of the stream.

At this stage there is already manifest increased swelling of the part, and, as determined by placing a cannula in the main lymphatic of the leg of a dog whose foot has been inflamed by placing it in hot water, there is, at what corresponds to this period, a greatly increased *outflow* of lymph. The swelling, that is, is due to increased *exudation from the dilated vessels*.

If the white corpuscles be studied, it is seen that they lag along, now adhering momentarily and becoming elongated and pear-shaped, now becoming free. What is more, while the red corpuscles progress onward, the white corpuscles lag behind, in consequence of these temporary adhesions, until it is noted that, proportionally, there can be a remarkable number of white blood cells in the inflamed area. As the current becomes greatly slowed, there may be *oscillation* of *the current*, and the leukocytes move to and fro with it, until, finally, they become firmly adherent. This is the stage of *margination* of the leukocytes.

Eventually—but by no means necessarily in every case—the slowing of the current in some of the vessels may give place to complete standstill, or *stasis*.

Next, it is to be observed that certain of the leukocytes are appearing at the outer side of the capillary wall, and, if in a favorable specimen the process be followed carefully, it is seen that it is brought about by a process—pseudopodium—being extended or insinuated, which passes through the delicate wall, gradually gets larger on the outer side, and there often shows finer processes streaming away from the wall into the tissue or tissue spaces, until more of the cell is in the outer side than in the inner, and eventually the whole corpusele comes to be external. To use an old illustration, the process closely resembles the trick of passing a bladder three-quarters full of water through a keyhole. Thus is accomplished the *diagedesis*,¹ or *migration of the leukocytes*.

¹ Hamilton and some other authorities object to this use of the term diapedesis. They would confine it to the passive discharge of red corpuscles which may occur in very acute inflammation, or in greatly dilated capillaries that lack support, as in the lungs (hemorrhage "per diapedesin"). Nevertheless, the term diapedesis means primarily a "stepping through," not a "pushing through," and implies an acute process. Etymologically, the modern usage of the term is correct.

As regards the particular orders of leukocytes that pass through, this is best studied by examination of specimens fixed rapidly, cut and stained, when these cells can be seen fixed in the process of passage. In an acute inflammation it is the *polymorphonuclears* that *especially migrate*, though it must be remembered that in special conditions the other forms—the eosinophiles and the lymphocytes—are capable of

Fig. 143

Migration of lymphocytes through wall of capillary x; y y, nuclei of vessel wall cells; Lmc, lymphocytes; x, irregular shape of nucleus of lymphocyte in process of passage. (Maximow.)

migration, the latter more particularly in subacute and chronic inflammation (Fig. 143). This has recently been amply proved by Schridde,¹ using his acetone method and differential stain.

This much for the earlier stages—the stages of development. Studying them we gain an explanation of the redness, the heat (increased determination of the blood to the part), and the swelling. Regarding the pain, we shall speak later.

¹ Münch, med. Woch., 52: 1905: 1862.

HEALING BY FIRST INTENTION

These earlier stages are essentially the same in all cases of acute inflammation. The later stages can only be studied fully by the method of section, and what happens in an individual case depends upon the nature of the injury. Thus, it will be most satisfactory to consider now the different grades and varieties of injury, beginning with the simplest and advancing to the more severe, in each case discussing the process of events.

INFLAMMATION RESULTING IN HEALING BY "FIRST INTENTION."

The simplest case is that in which, without bacteria gaining entry into a wound, there is section of the tissues, with destruction of a certain number of cells and injury to others not of necessity so serious as to destroy them, and in which either the wound is so small that the edges come together naturally, or, being larger, the edges are brought into apposition by one or other surgical means. Here, as a result of the first stages, there is an oozing or exudation of "serum" into the wound, along with some leukocytes and with an inconsiderable amount of blood from divided capillaries. Even if there be no blood, the interaction of the leukocytes, as sundry of them break down and liberate their fibrin ferment, of the destroyed cells similarly undergoing disintegration, and of the serum, lead to coagulation of the latter, and on the surface and between the apposed edges fibrin is laid down, and this acts as a provisional cement substance. Within the first twenty-four hours, and without mitosis, the tissue cells bordering upon the wound enlarge and send out processes and undergo division; epithelial cells of the deeper layers may even, according to Leo Loeb, exhibit a certain grade of amceboid movement, with alteration in relative position, coming thus into immediate apposition with those on the opposite side. In this way there is a bridging over and interlacing of the cells of the two sides of the wound. Next, as we are now convinced, largely through the agency of the leukocytes, the debris of dead cells is carried awayeaten and removed-absorbed. The same is the case with the fibrin; this, too, undergoes absorption. In three days or so sections show occasional mitoses in the cells on either side of the wound; the capillaries in the two sides are seen to throw out bud-like processes, which unite across this gap, and repair is practically complete. For a few days longer there may continue some redness and slight tenderness along the wound, but this disappears, and the parts are practically in statu quo.

It is, or used to be, taught that every wound must leave its scar. That this is not the case everyone knows who shaves. We have performed an autopsy on a case in which laparotomy had been performed within three months, in which close examination externally failed to detect the operation wound, while internally it was only indicated by omental adhesions and the absence of important pelvic viscera.

Essentially, a similar course follows *contusions*, whether subcutaneous or affecting deeper viscera. There are the same preliminary stages, with redness, swelling, heat, and pain, though the condition is complicated generally by more extensive rupture of capillaries and greater escape of blood into the part. The fluid of such blood dra'ns away by the lymphatics, as may, also, some of the corpuscles. The mass of the corpuscles, being out of place, degenerate; their hemoglobin dissolves out and undergoes a series of reactive processes, characterized by the color changes familiar in the "black eye." Eventually the leukocytes carry away the debris, and the parts, provided the tissue destruction has not been severe (as in laceration), return absolutely to the normal.

INFLAMMATION LEADING DIRECTLY TO THE FORMATION OF GRANULATION TISSUE.

If there has been loss of tissue, whether in communication with the exterior or through some cause leading to local necrosis or tissue death in an internal organ, provided bacteria do not grow in the wound, we have the same preliminary stages, only here, the injury being more severe and extensive, there is a greater reaction. More fluid is apt to be exuded, and, attracted by the dead cells and the products of their disintegration, more leukocytes migrate into the area. As a result, in an open wound the exudation, mixed with leukocytes and escaped blood, coagulates and forms a scab, which, drying on the surface, is in itself protective against bacterial invasion; and, if the wound has been effected in a clean manner, and due precaution be taken, the wound may remain sterile. Bacteria, it is true, may gain entrance from the wounded skin-for this is never sterile, and cannot be rendered germfree-but under favorable conditions the phagocytic activity of the exuded leukocytes and the bactericidal properties of the exudate may destroy them before they have time to multiply and gain the upper hand. In an internal organ or part the dead tissues cannot be immediately removed (e. g., an infarct, or limited area of blocked blood supply in an organ, or the various structures in dry gangrene); the exudate infiltrates it, its cells in dying coagulate, and the infiltrate between those cells also coagulates (coagulation necrosis).

Following upon this stage, we find that the dead cell material becomes either eaten up (phagocytosis) or gradually dissolved; it disappears. As shown by Leber, aseptic collections of leukocytes and the fluid from the same have distinct digestive powers, but, in addition, we have come to recognize of late years the autolytic properties of the cells (p. 370). Whichever process plays the greater part, already, in three days or so after such tissue destruction, more particularly at the edge between the dead and living cells, there is observable a marked relative diminution in the amount of debris, and with this the appearance of a clearer zone between the intact tissue and the scab or area of coagulation necrosis, with its more closely packed migrated leukocytes.






Granulation Tissue. The Upper is toward the Outer Surface. (Maximow.)

end., capillary endothelium: new cap., endothelium of newly forming capillary; fN., fibroblasts; l., polynuclear leukocytes; x., the polyblasts of Maximow (including plasma calls; deb., debris of leukocytes; deg., a degenerating "polyblast."



FORMATION OF GRANULATION TISSUE

The next stage observable is that, into this clearer zone new capillary bands project from the surrounding capillaries. This is the first stage, or almost the first stage, of *organization*, and deserves close study. We say almost the first, because, even before this shows itself, changes have been noticeable in the tissue cells bordering on the wound. They are found markedly swollen, showing unusual processes directed toward the area of injury, may, as in the previous case, undergo direct division, and



Formation of new vessels in granulation tissue: 1, from a Ziggler's chamber (formed of two coverslips) left in the peritoneal cavity of a rabbit for forty-eight days; portion of field bounded by two fully formed new capillaries; between them can be seen the solid buds and processes of developing new capillaries; 2, from a similar preparation to show formative cells, or fibroblasts, in direct connection with the endothelial processes. (Ziegler.)

within twenty-four hours, according to some authorities, although usually the period is later, the more orderly form of indirect division may be exhibited.

There still remains doubt as to the structure of the finest capillaries whether they possess anything of the nature of the limiting membrane or whether they consist merely of a tube, or potential tube of endothelial cells in immediate communication with the surrounding tissue

no

wi

of

m

The changes they undergo in this process of organization would cells. seem to indicate that the latter is the case. It would seem natural that where a tube under pressure passes from a region where it is surrounded and supported by tissues to a region where it is superficial and unsupported, it should give way on the side toward that surface, and that so, new capillaries should arise, as local "givings way," or projections on the free aspect of preëxisting capillaries, brought about by the internal blood pressure. But this is not seen to be the case. At the most, in an exposed wound the preëxisting capillaries tend to present general dilatation with curvature toward the wound surface. But, where a new capillary is to be formed, instead of a thinning of the wall, there is a thickening. The endothelial cell or cells at that point form a swelling, from which projects a primarily solid process. As this grows it becomes hollowed out at its base and blood gains entrance into and distends it. Either before this occurs, or later, it is to be seen (in sections) that the thin filamentous end of the process has either come into contact with the wall of some other capillary, or has become connected with the delicate process of some fibroblast (growing connective-tissue cell), or with a similar process from some other capillary. In the latter case more particularly we encounter specimens showing that the process from either end becomes hollowed out, the "mother" endothelial cells of the buds multiply, and in this way is gained a new capillary identical in structure with the old.

It is in this way that granulation tissue is developed, by the formation of successive loops of new vessels. In this way the free surface of a wound becomes covered with a granular or velvety surface of reddened, minute projections, individual sets of these new loops being surrounded by young fibroblasts, the products in the main of the multiplication of the preexisting connective-tissue cells bordering on the wound. Of late the view has once more gained support that some, at least, of these fibroblasts are derived from one order of leukocytes, while very possibly the proliferating vascular endothelium may contribute some. The fact that new tissues never arise in isolated foci, but always in direct connection with preexisting tissue, is strong evidence against cells that have wandered in from elsewhere playing a primary part in this process of new connective-tissue formation.

The identical process is to be met with in internal organs and in the organization of congulated blood—thrombi—within the vessels, though here, there being no free surface, the orderly formation of loop on loop is not so obvious; there is first removal of the dead matter by the agency of leuko-cytes, autolysis, and absorption. Into the clear zone so formed the vascular loops project forward in an identical manner; these become clothed with fibroblasts, while farther forward there is simultaneously occurring the clearance of debris and the projection of new vascular processes.

Thus, in an orderly manner the dead tissue is removed and replaced and the free space filled in.


As the fibroblasts become mature, from being at first rounded or oval and fairly stout, with relatively large nucleus, they become elongated

JASSIL NOLLVIANVUD JO NOLLVINUOJ

and spindle-shaped (spindle cells), or of elongate, stellate form; and not only do fine fibrillar processes project from their ends or sides, but with appropriate reagents they are to be made out traversing the body of the cell (Fig. 145), and these increase in number, replacing the cell andstance proper, until the cell is represented by a meagre, attenuated nucleus, with but a trace of cytoplasm, lying surrounded by a bundle of fibrils—while connective tissue.

Whether the white connective-tissue florils are formed wholly out of the cell substance has been a matter of some debate; the weight of evidence indicates that the interstitial substance also contributes. Certainly the bundles of fibrils, and, indeed, the individual fibrils, extend through many cell areas. In this connection Leo Loeb's observations, made in many cell areas.

Ftc, 145



Portion of a fibroplast, undergoing milosis, greatly enlarged, to show the interrellular and intracellular fibrils. (Lubarsell.)

our laboratory, are at least suggestive. He has pointed out hat, if a little blood plasma containing loukocyctes be expressed gendy between two slides, and then the one slide be pulled over the other, fibrin formation sets in, and upon examination all the fibres are seen to lie in the line of traction, and, what is more, to traverse the bodies of the leukocytes. While the fibroblasts in granulation tissue are hid down with no trace of regularity, the bundles of white connective tissue are remarkably regular, in this resembling the bone lamellae, in which, clearly, we have to deal with intercellular deposits.

Such newly formed white connective tissue, as it grows older, undergres contraction and condentation. The area of the new tissue becomes markedly smaller, the abundant capillaries become in the main compressed and oblicitated; from being very vascular, the new "scar" perconnes distinctly areacular; the sides of the wound, infract, or thrombus approximate, and we have the formation of the characteristic cientrix,

According to the tissue involved, so may we find some variation in the process. Thus, the brain contains a relatively small amount of ordinary connective tissue, and its vessels, it would seem, do not easily form new capillaries. Hence, if there be a hemorrhage or other local tissue destruction, while there is congestion around the area and abundant passage in of leukocytes, which load themselves up with debris (Gluge's corpuscles),¹ the formation of granulation tissue is very meagre; a cyst wall only may be formed, and, with the carriage away and absorption of the dead matter, as the brain, from its relationships, cannot fall in, it becomes replaced by serous fluid, so that a cyst develops. (See chapter on Cysts.)

ACUTE FIBRINOUS INFLAMMATION OF SEROUS SURFACES.

What is but a variant of the above process is seen in the fibrinous inflammation of serous surfaces. Simple acute serous inflammation is a variant of the previous grade, characterized by abundant exudate from the greatly ingested vessels of the serous surface and relatively slight migration of leukocytes. With its arrest, the parts undergo restitutio ad integrum. In the fibrinous form, and the same is true of the serofibrinous, there is abundant migration of the leukocytes out of the surface vessels. The irritant-bacterial in this case-leads to widespread necrosis and casting off of the outer endothelial layer, and, just as thrombosis in a vessel is induced by destruction of the vascular endothelium, the blood coagulating or conglutinating over the site of injury, so here the leukocytic exudation, breaking down, coagulates upon the injured surface, with the result that a deposit of fibrin occurs-the homologue of the scab over an external wound. If the deposit be but slight, it is absorbed by subsequent leukocytes, the areas of serous endothelium that have not been destroyed exhibit proliferation, and new cells cover the surface. If it be more extensive, complete return to the normal does not occur; we encounter replacement of the dead fibrinous matter by granulation tissue. New vessels pass in from the serous surface, organization occurs around these, and, where the fibrinous exudate completely fills a cavity, or is adherent to both aspects, then the new vessels coming from opposite sides anastomose, and we have complete obliteration of the serous cavity (concretio, or synechia), or the formation of bands and veil-like membranes of connective tissue passing from one side to the other (organized adhesions).

SUPPURATIVE INFLAMMATION.

The essential feature of this grade of inflammation is the profound attraction of leukocytes to the area of injury, followed by the death of a large proportion of the same, and this unaccompanied by the rapid

¹ Fritz Marchand points out that these are largely glial cells, not leukocytes Ziegler's Beitr., 45:1909:161).

SUPPURATIVE INFLAMMATION

disintegration and dissolution of the dead cells, which characterized the preceding grades. There is thus found a thick, more or less creamy and opaque fluid at the site of inflammation, and, on microscopic examination, this is found to be a suspension of leukocytes—the polymorphonuclear form being in overwhelming majority in all acute cases. Some of them exhibit well-staining nuclei, some have intracellular foreign matter—bacteria in various stages of degeneration—fatty and other matters, the result of phagocytic activity. These are cells presumably still active at the time of the removal of the pus. Others, again, show badly staining, broken-up nuclei. These are, clearly, degenerated and dead cells. Occasionally, notably in peritoneal inflammations, cells of other type may be observed, along with abundant free bacteria. If the fluid in which the cells are suspended be filtered off, it is found to be thick and singularly rich in protein, the results of tissue destruction and leukocytic disintegration.

Such is *pus*, and, experimentally, it can be produced by many means —by the local injection of oil of turpentine, the local action of certain metals, such as mercury and copper and their salts; by the products of growth of certain bacteria, or by the bacteria themselves. Experimentally, that is, as Leber, Councilman, and others showed many years ago, it is possible to produce aseptic pus. This, however, is not the case in ordinary clinical practice; in man, suppurative inflammation, in nine hundred and ninety-nine cases out of a thousand, means the presence of bacteria. The abscesses that may follow the mercurial intramuscular injections, which constitute one method of treatment of syphifis, may be aseptic, as also, it may be, that the suppuration following deep burns may not in all cases be wholly ascribable to microbic agencies; the destroyed cell matter, that is, tends to cause an active migration of leukocytes. But in these cases it is impossible to exclude the presence of surface bacteria, and so decision is difficult.

The course of such suppurative inflammation can easily be followed in the lower animals by means of subcutaneous injection of pusproducing bacteria, thereby causing abscess formation away from the surface, and excluding the possible action of other agencies. The variation from the processes hitherto described consists in (1) the progressive growth of the pathogenic organsims at the site of inflammation; (2) the progressive destruction of the tissues; (3) the more pronounced migration of leukocytes to the area; and (4) the indication of interaction between those leukocytes and the bacteria, and of the existence of yet other reactive processes, whereby the growth of the bacteria becomes limited and arrested, and their destruction brought about. Experimentally, that is, just as in more natural conditions, there is a tendency for the abscess to come to a head, and following the successive stages, the most striking feature is that the bacteria grow freely and abundantly, and then the time comes when the indications of abundant growth are succeeded by those of arrest and disappearance.

There is, in short, a period of incubation, during which the bacteria multiply locally without setting up a very marked reaction. Soon there

is such reaction; the cells of the part become swollen, the capillaries in the immediate neighborhood full of blood, but only after this does the migration of leukocytes become noticeable. In other words, some little time is necessary before the bacteria, growing, are able to discharge sufficient toxins, and for those toxins to diffuse in sufficient concentration to attract the leukocytes out of the surrounding capillaries. From this time on the migration is the prominent featureis so extensive that the leukocytes at the focus of bacterial growth completely obliterate the tissue cells and by their abundance and the increased exudate compress the surrounding tissues; they take up the bacteria actively, remain in the centre of attraction, and-whether by the toxins of the ingested microbes, or by the toxins of the still abundant bacteria that have remained uningested-undergo destruction and dissolution. For some days, therefore, the organism is unable to restrain the bacterial growth. Despite the abundant leukocytes, the area involved increases; the tissues of this area are killed and disintegrated, their place being taken by the abundant leukocytes, which continue to pour in from an increasing ring of surrounding dilated capillaries. Bacteria are to be detected outside the limits of the area, having grown there along the tissue spaces or been carried by the currents of the exudate.

And then, at last—after three or four days—the organism gets the upper hand, and we see our fully developed *abscess*. There is what Ribbert describes as a dense wall of leukocytes (Wallbildung) around the area; immediately outside and merging into this is a zone of dilated capillaries, the *pyogenic zone*, or "*membrane*," there are no longer any bacteria outside, and, while the outer zone of leukocytes stains well, the central mass of leukocytes stains poorly and shows signs of nuclear disintegration and of death. Bacteria are still present in the central pus, and still living, as may be determined by making a culture, but their number is diminishing.

In the chapters upon general reaction and immunity (see particularly Chapter IX) we shall discuss more closely what has occurred to make the tissues more powerful.

Now, follow the stages of resolution of the abscess: The micrococci or other microbes become fewer and fewer, and as the production of toxins diminishes, and the attractive force is lessened, the undestroyed leukocytes pass away from the area into the lymph and bloodvessels—there is absorption. If the abscess be of small extent, the area closes in; if larger, the destroyed tissue is replaced by granulation tissue, and a cicatrix results.

The process here described as occurring in abscess formation holds also for inflammatory *ulceration*, with the difference that the latter process originates in close connection with an epidermal or mucous surface, and results in destruction of these surface layers, so that there is produced a loss of continuity of the tissues and exposure of the deeper layers—covered by pus—to the external medium. With resolution, there is healing by a like process of formation of granulation tissue. The like process obtains also in wounds, whether open or closed, that

SUPPURATIVE INFLAMMATION

have become infected. Intermediate between the abscess and the ulcer is the *fistulous* inflammation, in which what was originally a deep-seated abscess, through the pressure of its exudate, "points" or undergoes extension in the direction of least resistance of the surrounding tissues. As a result the suppurative process extends along a tract which sooner or later makes its way to a surface, from which the pus becomes discharged, the communicating passage being known as a *fistula* or *sinus*.

In suppurative inflammations of serous surfaces, another factor may be introduced. We have already seen that irritation of such highly vascular regions is liable to result in a fibrinous exudate. Where pyogenic bacteria gain entrance locally to a serous cavity-in appendicitis, for example-their toxins, before they become too concentrated, lead, first, to a fibrinous exudate, whereby the viscera in the immediate neighborhood becomes cemented together; and, while at the region of entrance, the bacteria may multiply and induce pus formation, the surrounding fibrinous adhesions prevent the escape of these bacteria into the serous cavity in general, and, in virtue of the fibrinous adhesion, dense, and permitting very slow diffusion of toxins, with arrest of bacterial passage, a localized abscess results in place of widespread suppuration-an abscess whose circumference, in part, is formed of inert matter, and not of living tissue. The products of bacterial growth, it is true, can dissolve this fibrin; thus, such an abscess tends to increase in size, but as it grows, so in favorable cases the outer zone of irritation, through the diffused toxins, leads to more fibrin being laid down, and if with active reaction on the part of the tissues and quiescence of the viscera, such fibrin formation can be adequately procured, the inflammation remains local. Indeed, with the increase in size, the visceral layers forming the walls of such an abscess may become eroded and the abscess discharge into the lumen of the intestines (for example) before the fibrinous adhesions give way; we thus may have *exogenous ulceration* of viscera.¹ Where, on the other hand, the fibrin formation is inadequate from one or other cause, from lack of reactive powers or from amount and intensity of action of the entering bacteria, there the bacteria become spread through the serous cavity, and general or diffuse suppurative inflammation occurs.

It is, perhaps, serviceable to banish the term general, as applied to peritonitis, as being too vague, and to distinguish between universal and diffuse. When, for example, there is perforation or rupture of the bowel, there may be immediate infection of a considerable neighborhood, and a purulent condition set up too diffused and indefinite in its boundaries to be spoken of as a local abscess; nevertheless, all around the outer limits fibrin formation is seen to have occurred. In this way the whole pelvis or one segment of the abdominal cavity may be involved, the rest being free. On the other hand, there may be no fibrin formation, and suppuration everywhere throughout the cavity universal peritonitis.

¹ See Adami, Montreal Med. Jour., 32: 1903: 401.

A similar spreading suppurative condition may show itself in the solid tissues, where the "wall building" by the leukocytes is incomplete, and the bacteria, as a consequence, proliferate and spread in the tissue spaces and along the lymph channels. In this way we have set up a *cellulocutaneous inflammation*, with diffuse suppuration of a limb or subcutaneous tissues. Where the toxic properties of the bacteria are still more pronounced, the extensive tissue destruction which follows their spread becoming the most marked feature, we have a *phlegmonous* or *gangeneous* inflammation.

In all these latter cases in which the bacteria are not successfully retained in the locality of their primary manifestation, gaining entrance into the lymph and blood stream they may be carried to a distance, and there, being deposited and finding conditions favorable for growth, may set up similar reactions, causing *metastatic* or secondary abscesses.



Fibrinous exudate in lung alveoli; case of acute lobar pneumonia; to show fibrinous network, (After Ribbert.)

More particularly in acute bacterial inflammations, stasis is apt to occur in surrounding vessels. By this means they become plugged, and, as an abscess extends, it is noteworthy that despite erosion of the tissues hemorrhage is not liable to occur. Such plugs of coagulation may extend up a vein for some distance (thrombophlebitis), and as an abscess extends, causative bacteria gaining entrance into them may grow along them, and so into the vessels. Growing, they soften the clot, and from it infect the vessel walls (pulephlebitis); portions of the disintegrated clot may become loosened, dislodged into the main vein beyond the point of arrested circulation, and carried by the blood to the lungs, liver, or other organ, and there become arrested in some smaller vessels, thus causing an infective embolus, or plug, and setting up an embolic abscess.


We have not as yet exhausted the varieties of acute inflammatory manifestations. In this progressive description of the different orders

MEMBRANOUS INFLAMMATION

we have purposely left out some which were not of the direct line. These we may now note.

Hemorrhagic Inflammation.—The exudate may contain not merely leukocytes, but also red blood corpuscles, under two conditions: (1) Where the causative agent is very toxic, setting up a greater dilatation and thinning of the capillary walls, along with, it may well be, a direct degeneration of the capillary endothelium; and (2) where the abundant capillaries are unsupported, so that their dilatation is not limited by the pressure of surrounding tissues. In both these cases erythrocytes may be forced through spaces in the capillary walls and escape with the exudate. In some cases of intense inflammation the number of leukocytes in the exudate may, from negative chemiotaxis, be very few, and, *per contra*, in inflammation of moderate grade, a stray erythrocyte may be found outside the vessels which, possibly, has passed out through the channel made by the actively migrating white cell (Fig. 141). In those other cases it has to be recognized that lack of continuity of the capillary wall is not necessarily produced by this means.





Schematic representation of a diphtheritic inflammation of a mucous surface. Complete loss of epithelial layer, the necrosis extending into the subepithelial layer; f, fibrinous layer; em, submucosa.

Such hemorrhagic inflammation often shows itself in connection with serous surfaces. The commonest site, however, is the lung substance, where the hemorrhagic exudate into the delicate walled alveoli is the distinguishing feature of acute pneumonia. Here once more we note that, so soon as the blood (for the exudate in these cases is blood to all intents and purposes) escapes from the vessels, it tends to coagulate. In this way is produced *acute fibrinous inflammation* (Fig. 146).

Membranous Inflammation.—Similar acute fibrinous inflammation may also affect mucous membranes, as of the throat and upper respiratory channels, but a distinction must be made between it and the *membranous* or *diphtheritic* inflammation. In this latter form an irritant acting from without or from the surface brings about necrosis of the surface layers—there is profound engorgement of the superficial vessels with exudation—and the same process happens as we noted in connection with infarcts, namely, the exudate and the dead cells undergo a common coagulation, and surface exudate and surface cell layers

together form a (false) membrane, which, in the earlier stages, is so intimately connected with the underlying living tissue that it cannot be detached, herein differing from the slighter fibrinous exudate. Such diphtheritic false membrane may occur not only in the throat, but in the intestines, bladder, and other mucous surfaces, and even on exposed wounds.

Later, the accumulation of leukocytes between the false membrane and the sounder tissues below leads to a digestion of the fibrinous connections and loosening of the membrane.

It is worthy of note in this connection that where there exists a wellformed basement membrane the diffusion inward of toxins is arrested to a considerable degree, only the surface epithelial layer undergoing necrosis and being cast off. This, at least, would seem to be the anatomical explanation of the firmly adherent false membrane in the pharynx in diphtheria, the looser, detachable membrane in the trachea and bronchi.

The terminology here is misleading and unfortunate; it dates from a time when the real nature of diphtheria was unknown and two conditions were recognized—croup and diphtheria, the former affording a detachable fibrinous membrane, the latter a more adherent membrane. Needless to say that nowadays we know no "fibrinous croup," but, as regards false membranes, we observe that a diphtheritic membrane is by no means always due to the diphtheria bacillus. Our own custom is to employ the term *diphtheritic* for all false membranes; the term *diphtherial*, when we wish to indicate that this is associated with the specific disease, diphtheria.



Schematic representation of a catarrhal inflammation: cpi. 1, columnar epithelium still in situ but with increased number of goblet cells discharging mucus; cpi. 2, epithelial cells liberated into exudate; cpi. 3, remaining vegetative epithelial cells,

Gatarrhal Inflammation.—A slighter grade of change, affecting mucous membranes, results in the *catarrhal inflammation*. Here, although we note that many of the epithelial cells are cast off, and can be recognized in the abundant exudate, the dominant feature is irritation rather than necrosis. With the surface irritation there is pronounced congestion of the underlying vessels, abundant exudation of serum, and some leukocytes, but most marked of all, the columnar, epithelial cells are stimulated to excrete abundant mucus, and the

INFLAMMATION IN NON-VASCULAR AREAS

copious mucus forms a layer over the inflamed surface, containing, also, leukocytes and cast-off epithelial cells. This discharge is regarded as, to a considerable extent, protective; not only are bacteria detained in it, but it is claimed to have a certain amount of bactericidal activity, and acids and other irritants diffuse through it with difficulty.

Inflammation in Non-vascular Areas.—There are several such non-vascular areas in the body—cartilage, the outer third (at least) of the cusps of the heart valves, the lens, and cornea. This last, more particularly, affords a favorable field for the study of the effects of local injury.



Mild grade of inflammation of cornes in man (keratitis e lagophthalmo), characterized by enlargement and direct division of the nuclei of the corneal corpuscles c_i with but slight invasion of polynuclear leukocytes p_i and lymphocytes I. (Tooke.)

Here, as in vascular tissues, we may have various grades. The very slightest destruction of surface cells is followed merely by swelling and overgrowth of the surrounding cells and replacement by these new cells. A little more severe injury is followed by attraction of leukocytes from the surrounding tissue spaces of the cornea and from the lacrymal fluid bathing the surface of the eye (this always contains a few leukocytes). These first fill the wound, then, as the surrounding corneal corpuscles multiply, they pass away and with them the debris of the dead cells is found to have gone also, and the part returns to the normal.

More severe injury is inflicted by injecting suppurative microbes, such as the *Pyococcus aureus*, into the corneal tissue. As in the case of the experimental abscess, we recognize a period of growth of the bacteria, with swelling, degeneration, and destruction of the cells at the focus of growth; some extension of the bacteria along the surrounding tissue spaces, the more peripheral cornuscles becoming

swollen, in their turn, and their processes more prominent. Only after many hours—it may be next day—do we see evidence that the toxins have diffused through the corneal lymph spaces and affected the circular marginal vein. This now becomes dilated and prominent, and microscopic examination at this stage shows margination of leukocytes and migration of the same *toward* the site of injury. Leber performed an admirable experiment to demonstrate that this migration is active, and that the mass of leukocytes is derived from the bloodvessels. He injected cocci into the cornea at three different points around the centre, and found that there was no leukocytic accumulation toward the central aspect of these foci, but on the peripheral aspect each showed a more or less wedge-shaped radial accumulation.

From now on the process closely resembles that which we have described in connection with the abscess-or, more exactly, what happens in the ulcer: progressive tissue destruction, with-in cases that end in healing-even more marked accumulation of leukocytes. And, to complete the similarity, the vessels become definitely involved. We mentioned in connection with granulation tissue that the formation of the buds or processes which become new vessels was difficult to explain, save as due to chemiotactic influences. It is impossible to resist a like conclusion here. From the marginal vein processes are given off at several points, directed toward the ulcer. If the ulcer be not central, it is particularly on the side closest to the ulcer that these show themselves. They are not developed on the side of the vein away from the cornea. And as these processes enlarge, they become canalized, become new vessels in the previously non-vascular cornea, passing to the region of ulceration. Once formed, they may be observed, in man, for years afterward, the indication of old corneal inflammation. Similar new vessels are formed in the inflamed hear, valves, and make their way into inflamed cartilage.

Whether, now, in the stages of healing, definite granulation tissue is formed at the site of the ulcer, or merely absorption takes place, with proliferation of the corneal corpuscles and filling in of the wound, depends upon the extent of the ulceration and the duration of the process. In all cases, however, the new tissue is imperfect, the cells more irregularly disposed than normal corneal tissue, the cicatrix fibrous, white, and opaque. Where the process is not duly arrested, the whole thickness of the cornea becomes eroded, the aqueous humor escapes, and the inflammation may become general, affecting the whole eve.

In the *heart valves* the like process is complicated by the fact that the ulcerated surface is exposed to the blood stream, and on it fibrin becomes deposited, forming "*vegetations*." The causative organism may grow into these vegetations, and lead them to soften and become detached into the blood stream; if small, they may be completely absorbed by the agency of leukocytes; if not absorbed, the vascularization of the ulcerated cusp extends into them, their fibrin becomes replaced by granulation tissue; they become organized and fibroid.

CHAPTER II.

THE LOCAL REACTION TO INJURY (CONTINUED).

CHRONIC INFLAMMATION.

THERE is another group of cases in which (a) there is a long-continued process (b), accompanied by a few or none of the cardinal symptoms (c), in which the proliferative changes and formation of cicatricial fibrous tissue are more prominent than the vascular disturbances (d), more prominent also than are the evidences of leukocytic migration. And yet, studying these chronic conditions, we are convinced that they represent but another grade of the one process of inflammation. In the first place, they follow upon definite insult or injury to the tissue affected. Here, too, we have a series of intermediate cases passing imperceptibly from the most acute to the most chronic conditions, with no sharp line of distinction at any point; and, further, we recognize cases in which the beginning is of the acute type; others in which there are recurrent outbreaks of more acute disturbance, the changes progressing slowly but steadily in between these.

The word chronic is apt to be used very loosely. It is customary, for example, to speak of indications of previous acute inflammation as chronic—of old pleural, pericardial, or peritoneal adhesions as *chronic* pleurisy, pericarditis, or peritonitis. This is both wrong and misleading, and only permissible when—as often happens in chronic endocarditis—the altered condition of the parts in itself leads to further progressive changes. We should make the distinction between *old* pleurisy, or pleuritic adhesion, and *chronic* pleurisy, using the latter term only where we recognize slowly progressive changes, due to the continuance of irritation.


Like the acute conditions, the chronic changes may be set up either by bacterial or by non-bacterial irritation; in the latter case we may have to deal either with the effects of repeated slight mechanical injuries, or, more often, with the effects of perverted metabolism and continued slight intoxications affecting particularly certain tissues.

The Infectious Granuloma.—It is more instructive to begin with the study of cases which may verge on the subacute; for this purpose, the so-called *infectious granulomas* afford the best examples.

The term "infectious granuloma" is applied to the localized effects of the growth within the organism of a class of microbes which, while in particular weakly or very susceptible individuals they may induce a rapidly fatal disease, in general set up chronic disturbances, which induce no intense immediate reaction, but at the same time are not

easily antagonized once they gain a foothold in the tissues. The process of the arrest of their activities is a slow one, and often incomplete, the microbes being cut off from the surrounding tissues, but not of necessity killed. Often this process of arrest fails, and there is progressive extension of the condition. Each localized growth of the microbes in question sets up a localized reaction, resulting not in the formation of an abscess, but of a nodular tissue overgrowth and cell destruction a tubercle. Such microbes are those of tuberculosis, leprosy, syphilis, and actinomycosis, to mention the more important. Relatively inert foreign particles and the ova of certain larger parasites set up similar changes. The tubercle proper, caused by the tubercle bacillus, affords us the best example; its mode of formation has been studied more especially by Baumgarten and Borrel.

When tubercle bacilli, injected into the circulation, come to rest within a capillary, they are apt to be taken up by the endothelial cells



Schematic representation of a tubercle: a, giant cell, with necrotic centre and multiple nuclei more peripherally arranged; b, epithelioid cells; c, lymphocytes. of that capillary within a few minutes; or, if too many to be taken up, it is found that they become surrounded by large cells, which now we recognize are swollen and proliferated endothelial cells. The earliest reaction, thus, is on the part of the tissues, and is not a cell destruction, but cell growth. Along with these large so-called *epitheloid* cells a certain but relatively not large number of leukocytes also collect. These are first polynuclear, later lymphocytes predominate to the exclusion of the polynuclears.

Through this endothelial swelling and proliferation the capillary first involved becomes closed up, and later, the outer capillaries in


the neighborhood become similarly closed, so that the tubercle becomes strictly an *extravascular* formation. During the first stages, then, cell growth around the bacilli continues to be the most marked feature. The large endothelial cells immediately surrounding the bacilli may fuse together, forming a *giant cell*, having the mass of bacilli in the centre. This is not always seen, but, when present, is very characteristic. Nor would it seem that the numerous nuclei of the giant cell are purely due to cell fusion; as shown in the accompanying figures from Dr. Duval's⁴ studies upon chronic glanders, the multiplication is in part by amitosis. As the bacilli multiply²—and also, it may be, as some of them

¹ Duval and White, Jour. of Exp. Medicine, 9: 1907: 352.

² Prudden and Hodenpyl have shown that the injection of dead tuberele bacilli will lead to these earlier stages, in which case the gradual diffusion out of the intracellular toxins can be the only explanation of the succession of changes induced.

CHRONIC INFLAMMATION

are destroyed and their intracellular toxins liberated—from acting in the less concentrated form as stimulants to cell proliferation, in the more concentrated form they act as intoxicants, and lead to the death of the cells in the immediate neighborhood. Thus, the typical giant cell comes to exhibit a central, badly staining, necrosed area, around which—often more on one side than the other—other endothelial cells have fused, whereby it comes to pass that the typical tuberculous giant cell exhibits on section a peripheral circle, or crescent, of nuclei surrounding a more or less hyaline necrosed mass, the tubercle bacilli being found more particularly at the edges of the necrosed area, although indications of broken-down bacilli may, with care, be detected throughout the central area. It must here be emphasized that the giant cell is not an essential component of the recent tubercle; cases of miliary tuberculosis of the lung may be encountered in which the abundant isolated tubercles are wholly devoid of this order of cell.





The complete early tubercle shows, thus, either a more centrally disposed cluster of larger, more hyaline, epitheloid cells or a giant cell surrounded by cells of this order, and a certain number of interspersed lymphocytes—the typical "small round cells" of subacute and chronic inflammation—or, less frequently, of polynuclear leukocytes. Outside this there may be some dilatation of the surrounding capillaries, but it is not very marked.

As the bacilli continue to propagate, the area of cells destroyed increases, and we gain a larger and larger central area of necrosis of that type known as *caseation*. The affected cells completely lose their power of staining, the cell bodies undergo fatty granular disintegration and lose their outlines—an opaque granular cheesy mass fills the centre. Now, around this there may be found several newly formed giant cells, and as the central area increases so does the peripheral ring of active cells increase in circumference until the mass may be clearly visible to the naked eye. After this stage it may be noted that the bacilli are

not merely within the giant cells, but interstitial also, lying between the cells in the central area.

Some, indeed, escape-grow, or are carried by leukocytes-outside the bounds of the tubercle, and, coming to rest in the lymph spaces, there may form foci for the development of new young tubercles, whereby, in place of one, there may be a conglomeration of several small tubercles around a central necrotic area. These give origin to yet other tubercles, so that eventually a gross conglomerate mass of tubercles, with large central area of caseation, becomes developed. Such a mass, in its continued growth, may come to involve a surface, or the wall of a large vessel, or a bronchus, and, causing the destruction of the superficial layers, may rupture and discharge the cheesy matter with its bacillary contents, causing thus a *tuberculous ulcer*, and favoring the spread of the bacilli to other regions. In such case there develops a progressive tuberculosis. On the other hand, as the tissues become accustomed and adapted to the toxins (we shall discuss the means later), and as the development of the tubercle proceeds at a less rapid rate, time is given for the fixed connective-tissue cells of the periphery of the tubercle to develop from fibroblasts into definite connective-tissue cells. And so we may obtain the formation of a well-marked connective-tissue capsule surrounding the growth. With this there is arrested activity on the part of the bacilli-indeed, if the tubercles are small, they may undergo complete absorption; if, through fusion, larger caseous masses have been formed, there may be consolidation of the same, with calcification, and hence calcareous nodules, surrounded by a dense fibrous capsule (often seen at the apex of the lungs), come to represent the old tubercle. In such caseous and caseocalcareous nodules the bacilli may, it would seem, persist for years, for the matter is found capable of infecting guinea-pigs. If the health of the individual be greatly lessened, the indications at postmortem are that they may form foci for renewed growth of the bacilli, and, it may be, rapidly progressive extension of the disease.

There is now no doubt that tubercles may undergo complete absorption. This has been demonstrated in dogs having experimental peritoneal tuberculosis, and there are surgical records to the same effect. We have ourselves seen in the postmortem-room only a few larger caseous masses left in the abdominal cavity of a woman on whom operation had been attempted some months before, and given up as hopeless on account of the universal peritoneal tuberculosis. The patient, after the attempted operation, had been treated with Röntgen rays for some months, as a last resort, and died, not of tuberculosis, but of obstruction of the bowel.

With variation, rather in detail than in principle, this description applies to the other infective granulomas. In syphilis the giant cells are not so frequent, and the necrosed centre matter becomes "gummy" rather than caseous; in actinomycosis the tendency is for the necrosis to be of a more suppurative type—leukocytes making their way into

the dead area and causing dissolution. Still more marked liquefaction is seen in chronic glanders.

Around foreign particles, while there may be extensive giant-cell formation, there is little necrosis—purely fibroblastic accumulation passing on to fibrosis, with subsequent hyaline degeneration, which must be regarded as a necrobiosis rather than a necrosis. The same is true regarding the eggs and larvæ of parasitic worms and the inflammation around them.

Chronic Diffuse Inflammation.—But tuberculosis and syphilis may also set up not these localized granulomas, but diffuse fibroid changes in organs. Such may be the after-results of multiple miliary granulomas. We have seen the transition from the one to the other well marked in the heart of a child with congenital syphilis. More often it would seem to be caused by the action of the toxins apart from the bacteria. Such fibrosis—cirrhosis—is common in the syphilitic liver, both of the congenital and the acquired disease. French writers have called attention to its existence in the liver in tuberculosis. The fibroid changes of recurrent rheumatism, seen more particularly in the heart valves, are also, it would seem, primarily of bacterial origin.

There exists, further, a widespread group of conditions that we class as chronic-"itis" (the term "itis" attached to the name of an organ should always denote a state of inflammation)-chronic nephritis, hepatitis, thyroiditis, and so on, in which, although in some cases the irritant may be of bacterial origin, faulty metabolism and imperfect nutrition appear to be the main causes. In these, so far as we can see, excessive or disturbed secretory activity leads to degeneration and atrophy of the specific secreting cells of the organism, and either with the reduction in the nobler elements, the less noble connective tissue undergoes proliferation, to replace the lost tissue—a process similar to what we see takes place in the formation of a cicatrix-or substances which act as irritants to more delicate and highly organized cells act as stimulants to the more lowly connective-tissue elements, and so, destroying the former, lead to the proliferation of the latter. It may be that both factors are at work. In all such cases we are apt to see proliferation accompanied by little vascular dilatation, and if leukocytes be present in any number, they are of the lymphocytic origin, and not polynuclear, and are most abundant around the larger vessels; nay, more, they have some tendency to assume the plasma-cell type, being present as relatively large polygonal and irregularly shaped cells with eccentrically situated nuclei.

Other examples of injury leading almost entirely to proliferation of the connective elements we have already noted, namely, the fibroid capsular formation around relatively inert foreign bodies.

How far injury may lead to proliferation of the higher tissues we shall discuss in connection with neoplasia.

THE MAIN DATA REGARDING INFLAMMATION.

In another place⁴ we have analyzed in some detail the various factors concerned in the process of inflammation. Apart from unwillingness to repeat ourselves unduly, we cannot here devote the same space to such analysis. It is, however, necessary that we should call attention to the conclusions that are to be deduced from the study of the different grades of the process, for certain facts stand out very prominently, and must be made the basis for our judgment regarding the essential nature of inflammation, and in so doing we can call attention to matters which could not easily be introduced in a straightforward description of the stages and different varieties of inflammatory manifestation.

 The Cause of Inflammation.—First, then, all the grades we have described have followed upon injury of one or other nature, and even where the injury has been continued, in all we have evidence of reaction.

2. The Two Prominent Factors in the Process of Inflammation.— This reaction shows itself in many ways. Whether vessels be present or not, we observe two predominant factors coming into play: (a) Sooner or later, proliferation of the cells immediately around the injured area; (b) attraction of the wandering cells to the area of injury.

3. Secondary Role of Vessels.—The presence of vessels adds no new features to the process; it does but reinforce processes already in action where vessels were absent, rendering the reaction more rapid and more complete. More particularly is this the case in respect to the migration of leukocytes and the conveyance of increased fluid (exudation) to the part.

4. **Tissue Proliferation.**—In the slightest grades, and again in microbic inflammation, where the toxins are not too concentrated, cell proliferation may show itself as the very beginning of the reaction. In severer grades it is only with resolution that it becomes marked. In intermediate grades cell destruction may be proceeding at the centre, while at the periphery, where the irritant is less intense, proliferation may be in evidence. It has been customary to draw a sharp line of distinction between inflammation and repair; this is irrational.

5. Capacity for Proliferation of Different Tissues.—As regards the individual tissues, we note that it is the lowest, the most undifferentiated of all, namely, connective tissue, that is most apt to undergo proliferation. And here we may note, although we shall have to consider the matter further when considering regeneration, that the more highly differentiated a tissue, the less is its capacity to proliferate and exhibit reparative processes. As, also, that where a tissue under favorable conditions can manifest proliferation and new-growth, that same tissue under conditions of acute or chronic inflammation may exhibit

¹ Inflammation, an Introduction to the Study of Pathology, Maemillan, 4th edition, 1909.



0

t

ehe ed f

ď

·I'

0 11

d nll n n e, n of

;0 1re id rie it n, Film Made from Peritoneal Fluid in Case of Peritonitis set up by Inoculating B. Coli Twenty-four Hours Previously into the Abdominal Cavity of a Rabbit. (Beattie.)

polym, polynuclear leukocytes, many containing bacilli; *l.m.*, large hyaline mononuclear cells, many acting as phagocytes for polynuclear cells, red corpuscles, etc.



LEUKOCYTIC MIGRATION

little tendency to reproduce itself. Where two tissues are proliferating together, the one actively and rapidly, the other at a slower rate, the former tends to hinder and arrest the growth of the other. Thus, in general in inflammation it is the connective tissue that tends to replace or supplant all other tissues, and on surfaces the simplest forms of epithelium or mucous membrane, and not the more elaborate glands, hair follicles, etc.

 Leukocytes in Inflammation.—Turning now to the leukocytes, a word must first be said regarding the varieties of the same. We now recognize three groups:

(a) That which we may term the "bone-marrow group," because, although there are indications that these may also be produced elsewhere, it is in the bone-marrow more particularly that in the adult we encounter the mother cells from which these develop. The members of this group are more especially the polynuclear (finely granular oxyphile), the commonest form present in the blood, and the cosinophiles.

(b) The lymph-follicle group consists of the lymphocytes and the plasma cells; Schridde' has recently confirmed the conclusions reached by Ribbert, Saxer, Marschalko, and Marchand, by demonstrating by a special method that both these cells possess identical granules.

(c) The endothelial and tissue-cell group, of which the large "hyaline" cells (Metchnikoff's macrophages) are the representatives (see Plate XVI). Each of these varieties may be encountered in one or other grade of the inflammatory process. In acute inflammation, it is the polynuclears that dominate the scene. It is these that are seen actively migrating and actively phagoextic for bacteria.

In acute inflammation, also, we observe that the cosinophiles are involved. Those that happen to be in the tissue are among the first to be found at the site of injury; in the very earliest stages several observers have found them relatively abundant. They are not, or scarcely at all, phagocytic. The part they play is in debate. In inflammation of the peritoneum, W. G. MacCallum has found them massed in the capillaries of the mesentery and omentum.

In the infectious granulomas the hyaline and endothelial cells play a prominent part, as again in inflammation of moderate intensity of the peritoneum and serous surfaces. These large, clear cells, with relatively pale-staining nuclei, are actively phagocytic for other cells and cell debris—as are all the tissue-cell group; as also for bacteria of medium grade of virulence (of the type of the tubercle and glanders bacilli), not so actively phagocytic for the microbes of suppuration.

It is in chronic inflammation also that we encounter the lymphocytes and plasma cells as prominent features, both more particularly clustered around the vessels. They are not markedly phagocytic. Several workers regard these as actively concerned in the production of specific antibodies.

7. Leukocytic Migration.—There is a definite object in this accumulation of leukocytes at the site of injury and irritation. We have

¹ Die Kornelungen der Plasmazellen, Wiesbaden, Bergman, 1905.

noted that they are attracted by dead-cell matter, and that they remove the debris. (Experimentally, if a fine glass tube containing tissue extract be inserted in one of the large veins, the leukocytes from the circulating blood are found to accumulate in it.) We have seen, also, that they actively ingest pathogenic microbes, if these be not too virulent, and, when ingested, may digest and destroy them. And it is found that even where either the ingested bacteria are too virulent, and lead to the death of the leukocytes, or where the cells are destroyed by the bacterial toxins diffused in the medium, they still may be of service, for in their disintegration and dissolution they liberate antibacterial" and antitoxic substances. It has been found, for instance, that if an acute aseptic inflammation be set up in the pleural cavity, by powdered glass, aleurone, etc., an exudate is given off containing abundant leukocytes. If these leukocytes be filtered off, the filtrate is much more bactericidal than is the lymph or blood serum of the same animal, and this is traced to the breaking down and dissolution of many of the migrated leukocytes. How far, also, it is due to an active excretion from the living leukocytes is still a matter of debate. Hardy and Kanthack described very definitely the active discharge of the granules from eosinophilous leukocytes in the frog, comparing them with the discharge of secreting cells; nor is it difficult to repeat their experiments. Undoubtedly, the exudate comes to contain more bactericidal and antitoxic substances than the circulating blood plasma.

8. Fate of Leukocytes.—As to the fate of the leukocytes, it is various. Many, we have noted, undergo dissolution *in situ*; this, indeed, seems to be the fate of the majority; some become ingested by other leukocytes and by the proliferating tissue cells of the part, some pass back out of the area of inflammation into the lymph stream, or actually back into the capillaries.

The members of the bone-marrow and lymph-follicle groups never form tissue. It had been thought by Maximow and others that the plasma cells have this power. Schridde, by his method of granule staining, has shown that, though plasma cells may come to be retained in connective tissue, and even to take on a spindle-cell shape, lying between the fibres, they retain their particular granulation; in other words, never become typical fibroblasts. Of the tissue-cell group, some, at least, have the power of tissue formation—as might, indeed, be expected.

9. Fluid Exudate.—Beyond favoring materially the carriage and migration of leukocytes into the inflamed area, the vessels, through their dilatation, are the direct cause of the increased exudation of fluid into the injured area. Possibly, this is not the correct way of stating the relationship. We know that there is a close interaction between the tissues and the blood; that when, for example, blood is drained from the body, the tissues rapidly give up fluid into the vessels. It may, thus, well be—nay, probably is—that physical changes in the tissues, due to cell irritation in the first place, act on the vessels, and call for increased flow of fluid from the vessels into the tissue spaces; that the

PARTICIPATION OF THE ORGANISM IN INFLAMMATION 447

dilatation of the vessels is the direct result of the tissue disturbances, and not primarily brought about by nervous influences. That this is so we have already indicated. But once the vessels have dilated, as indicated by the increased lymph flow from the parts, the exudation is in excess of the immediate needs of the tissues. We have developed, in short, a flushing process, serviceable so far as it dilutes the irritant, harmful so far as it carries the irritant out of the primary focus; serviceable in bacterial inflammation in that it brings a constant supply of antibacterial serum to the part, harmful in the same in that, also, it brings more foodstuffs for those bacteria; serviceable, lastly, as pointed out by Hilton,' by acting as a fluid splint, securing immobility, and harmful if too long continued, because it favors the formation of adhesions, rendering the part rigid.

10. **Repair**.—Indeed, while we cannot but see throughout the whole study of inflammation that the tendency is toward an arrest and repair of the injury, at the same time it is constantly being impressed upon us that the reaction is not directly proportioned to the injury inflicted; it may, in some respects, be inadequate; in others excessive.

Of this, instances will occur to the reader. The leukocytes may be repelled instead of attracted; may take up bacteria that they cannot destroy; may carry them away to other parts, and so originate new foci of inflammation. The tissue proliferation may be wanting, or may be superabundant, leading to exuberant granulations, keloid, etc. The fibrinous deposit on serous surfaces may be greatly lacking (as often happens in typhoidal perforations), so that there is no limitation of the inflammatory process; or excessive, so that full absorption is impossible; the organization of the same may lead to constriction and kinking of the gut, or to the formation of bands, causing strangulation.

Inflammation is not repair; it is, as we have elsewhere expressed it, the attempt thereat. It is an adaptation on the part of the organism to unusual conditions, and as such, being outside the normal range of tissue reactions, it is almost inevitably less perfect than is the normal reaction to normal stimuli. As we have pointed out (p. 125), adaptation is not an immediate process; while the cells can accomplish more than is normally demanded of them, it takes time for them to respond appropriately to stimuli that are beyond the normal, and in inflammation, before the cells have, as it were, learnt their lesson, such changes may have been produced that return to the normal is impossible—may, more, before it has educated itself the organism may be overcome.

11. **Participation of the Organism in Inflammation**.—This leads us to consider the further point: How is it that in microbic inflammations that are recovered from, the bacteria at first thrive, and then later are arrested in their growth? Is it purely due to local accustomance and adaptation of the leukocytes and tissue cells to the changed conditions, so that, becoming used at first (those that are not immediately destroyed)

¹ Rest and Pain, 1863; 89 (a work that should be a "companion book" of every medical student).

to small doses of the toxins, they eventually become able to neutralize much larger doses; or is there assistance from the rest of the organism? We must assume that both are in action (p. 415). Here attention must again be called to the fact that, whereas inflammation is, we hold, essentially a local process, it may be, and generally is, accompanied by general disturbances-by a certain amount of fever and malaise, even by general leukocytosis. Clearly, the products of cell destruction in any extensive injury, and, again, the toxins of bacteria, diffuse out, or are flushed out, of the local area into the lymph and blood, and so may tell upon the organism in general. It is not, let us repeat, only bacteria that cause fever and malaise; the development of an infarct or an internal hemorrhage is followed by a distinct rise of temperature and general discomfort. As Hildebrandt and others have shown years ago, the injection of enzymes into the circulation, or tissue extracts, leads to the febrile state-and when tissues break down such enzymes and dissociation products become liberated. Where there is leukocytosisincreased number of leukocytes in the blood-it is a clear indication that the inflammatory products circulating in the blood and carried to the bone-marrow, have there attracted the leukocytes from the tissue spaces into the vessels, if they have not directly stimulated the mother cells to increased proliferative activity. Such leukocytosis is clearly a means whereby the rest of the body may aid in arresting the inflammatory

12. The Nervous System in Inflammation.—There is another and important means of relationship between the injured area and the organism in general, which now we must take into consideration, having thus far wholly neglected it, its mode of action not being observable by histological observations. We refer to the nervous system. Pain, in the first place, is one of the cardinal symptoms of the condition. It has been usual until within the last few years to explain this as due in the main to pressure of the exudate upon the delicate nerve-endings, to point out that where a tissue is loose and can accommodate itself to greater increase in size, there is no pain accompanying inflammation; where it is dense, and the exudate cannot escape, there pain is severe. We now see that this explanation is invalid when, as by Schleich's and other methods, the injection of fluids into a part is found to induce local anesthesia. We can, thus, only conclude that pain is the expression of irritation of the nerve-endings by the diffused toxins (using this term in its widest sense), and recognize that the slighter pain in loose tissues is due to the greater affusion of fluid and dilution of those toxins. In more severe cases the actual exposure of the nerves and destruction of their endings affords an adequate explanation. While, ordinarily, the early stages of the inflammatory process are the result of local influences and independent of stimuli or direction from the central nervous system. undoubtedly the higher centres can, and do, play a part at a later period -nay, they may actually initiate a process indistinguishable from true inflammation. As proving the capacity of the central nervous system to intervene, it will be best to consider these cases first. There is ample

THE NERVOUS SYSTEM IN INFLAMMATION

evidence that imagined injury to a part may be followed, and that rapidly, by all the essential symptoms of inflammation, save, it may be, the migration of leukocytes. As we can have grave inflammatory changes unaccompanied by this migration, the exception is not sufficient to warrant us in declaring that, therefore, these conditions must not be regarded as inflammation. Such nervous mimicry doubtless explains the spontaneous and apparently causeless inflammatory manifestations which may show themselves in hysterical subjects; explains also to a large extent localized, or sometimes widespread cutaneous and other acute congestions and inflammatory disturbances seen in cases of idiosyncrasy (p. 410). Its existence is best demonstrated in certain hypnotized subjects, in whom the suggestion of burning or other local injury may lead to the sensation of acute pain in the part, followed within a very short time by local vascular congestion, heat, swelling, and pronounced exudation. As an indication of the central origin, it has frequently been noted that these manifestations tend to be bilateral and symmetrical, although the supposed injury has been unilateral. The condition of herpes zoster, with its pronounced inflammatory manifestations and serous exudation along the course of distribution of certain cutaneous nerves, appears to come in this category, the observations, more particularly of Head and Campbell,1 demonstrating that the primary lesion in these cases is in connection with the posterior ganglia of spinal nerves. It is from these considerations that we have been forced to include in our definition the clause to the effect that the process may follow referred injury.

Now, clearly this referred injury and its results are frequent factors in inflammatory manifestations. It is by this that we have to explain the swollen, and reddened check and the earache, and, it may be, discharge from the ear in cases of abscesses of the root of a tooth, the extensive involvement and swelling of the surrounding tissues in cases of joint injury. The areas of the face, above noted, have a common innervation; the same is true of the individual joints and the cutaneous and other tissues surrounding them; or, more exactly, these parts are supplied from the same level of the spinal cord. Irritation of the nerveendings in the injured area induces excessive stimulation of the nerve cells of the posterior horns, with irradiation of the simuli not alone along the reflex area particularly involved, but to other associated cells in the immediate neighborhood, with, as a result, not only referred pain, but reflex vasomotor changes initiated in the areas controlled by these nerve cells.

That the vasomotor nerves play a part in modifying the inflammatory process is well demonstrated in regions such as the rabbit's ear, in which the course of the vasoconstrictors has been differentiated from that of the vasodilators. Where the former nerves are divided and the latter alone in action, the vascular dilatation and exuation are more pronounced, and the process of inflammation has a more rapid

¹ Brain, 23: 1900: 353.

29

of

1e

od

ue

ple

course, than in the reverse condition of uncontrolled action of vasoconstriction.

13. **Temperature Changes.**—Here, too, a word must be said regarding the *increased heat of inflamed areas*. In plants, by the use of delicate electrothermometric methods, it has been demonstrated that injury is followed by distinct, if small, local rise of temperature. There is here no circulation, and such rise can only be the expression of the increased metabolic activity of the surrounding cells. If such a rise occurs in animals, it is too small to be appreciated, and is wholly masked by the vascular changes, to which—to the increased pouring in of arterial blood —must be ascribed the increased local heat. Just as it is a general law that within certain limits increased temperature leads to increased metabolism, so have we evidence that the increased local temperature leads to a more rapid and complete evolution of the inflammatory process.

14. Adequacy of the Inflammatory Reaction.—For long years it has been the custom to regard inflammation as a harmful process. The above considerations show, as the great old surgeon, John Hunter, recognized long ago, that nothing could be farther from the truth. What is true is that, where it shows itself, it indicates that harm has been, or is being, done to the organism, and that the system is reacting and tending to counteract that harm. It is, thus, a danger signal, and if the surgeon can assist matters by removing the cause of the harm, such assistance is clearly indicated.

It is further evident that the reaction on the part of the tissues may be (a) inadequate, (b) adequate, or (c) excessive. The natural tendency, observing the grave local disturbances, is to regard the third of these as most usual, and to seek to mitigate the symptoms. This, again, is seen to be an error; most often the reaction is inadequate, and modern surgeons, like the late von Mikulicz, are increasing the leukocytosis prior to operation; or, like Bier, are favoring and increasing the hyperemia and exudation, and this with excellent results, when the process is carried to the proper limits.⁴

FIBROSIS AND ITS RELATIONSHIP TO INFLAMMATION.

In considering chronic inflammation, we have called attention to the constant, or almost constant, development of increased fibrous tissue of fibrosis—as a consequence of the process. Here it becomes necessary to inquire whether fibrosis, which is so widespread a condition, is always of inflammatory origin.

The answer to this question must be in the negative; there are conditions of fibrosis which, by no process of reasoning, we can include among

⁴ We have discussed more fully the rationale of these methods in the article Inflammation in Keen's System of Surgery. Webb and Williams have recently shown that induced passive hypermin promotes lymphocytosis, as also (Trans. 5th Ann. Meeting Nat. Assoc. Study and Prev. of Tubereulosis) that high altitude increases the number of "mononuclears" in the blood. To this they aseribe the beneficial results of altitude upon tubereulosis.

FIBROSIS AND ITS RELATIONSHIP TO INFLAMMATION 451

the results of inflammation-conditions in which we can recognize no preceding local injury as the primary cause. We have to acknowledge, that is, that stimulation within physiological limits, as well as grades of pathological irritation, may lead to the overgrowth of fibrous connective, as of other tissues. We have to recognize physiological strain as a cause of growth. We see this in the case of muscle in the increased growth dependent upon exercise, in bone in the greater size of the processes and ridges of insertion of muscles in those of active muscular, compared with those of poor muscular, development. We must admit the same for the connective tissues. What we regard as the best illustration is one afforded by Carrel and since repeated and confirmed by other workers, namely, if a length of artery, such as the common carotid of the cat, be removed and in its place there be transplanted a length of vein from the same animal of approximately the same dimensions, that vein at first, under the increased blood pressure to which it is subjected, shows some dilatation; but if after several weeks the animal be killed and the transplanted vein be examined, the only signs of inflammatory change to be recognized are in the immediate neighborhood of the points of junction. Away from these all the coats show an extraordinary connective-tissue overgrowth and thickening, more particularly the media and adventitia, and the dilatation is now little marked. This overgrowth is not of inflammatory type, and can only be ascribed to the increased strain to which the vein has been subjected. coupled with adequate, not to say improved, nutrition.

In our second volume (Chapter VIII) we demonstrate that the marked connective overgrowth of the intima in the arteriosclerotic artery is of the same order, due to the increased strain thrown on the intima through giving way of the inflamed or degenerated media.⁴

We observe a similar, though less marked, change in the walls of the veins in cases of prolonged, though not extreme, passive congestion. (In extreme cases the venous condition of the blood, and consequent lack of adequate nourishment, leads to dilatation pure and simple.) We ourselves and several other observers have called attention to its existence in some cases of passive congestion of the liver (nutmeg liver.) And in lymphatic territories the same may occur. Mere obstruction to the main lymphatic trunks from a part does not lead to complete stagnation of the lymph; on the contrary, there is a continual interchange between it and the blood in the capillaries. Nevertheless, there may be set up continuous and prolonged distension of the parts, and this, similarly, is followed by fibrosis—diffuse in this case. Such appears to be the explanation of the commonest form of *elephantiasis*, the fibrosis of macroglossia, and other cases of lymphatic obstruction, whether congenital or acquired.

Another condition of fibrosis, the development of multiple fibroid growths, where a true *fibromatosis* and not, as will be pointed out later, a state of neurinomatosis, would seem, from the specimens we

¹ See also Adami, The Nature of the Arteriosclerotic Process, Amer. Jour. Med. Sci., October, 1909.

have examined, to originate in a congenital fault of the lymph vessels of the affected parts. The condition differs from fibroma formation proper in that the overgrowths are not encapsulated, but pass imperceptibly into the related normal connective tissue; early specimens show a well-marked lymphangiectasis, or obstructive dilatation of the lymph vessels and channels.

The *fibromas* proper, or neoplasms, formed of fibrous tissue, must also be separated from the inflammatory fibrosis. As to their causation, we are still uncertain.

We thus are able to classify the fibroses as follows:

I. Of inflammatory origin.

1. Replacement fibroses, in which the fibrous tissue takes the place of other tissue that has been destroyed (cicatricial fibrous tissue, including that of infarcts). The "seleroses" of the central nervous system come under this heading, although in them neuroglia in the main, and not ordinary fibrous tissue, is usually concerned. Here, also, is to be included, in part, the fibrosis of chronic interstitial nephritis and hepatitis.



Section of the norta from a case of nodose arteriosclerosis, to show the bulging and thinning of the media, prepared by Dr. Mathewson. ~ 8 diameters. The section shows also the bryaline degeneration of the deeper hyers of the overgrown intimu, and the persistence of a fine layer of less altered intima tissue immediately beneath the media. The media in this case showed evidences of calcarcous degeneration in patches with some hyaline change.

2. Proliferative Fibroses. Here more particularly we have (a) the capsular fibroses of the infective granulomas, around inert bodies, etc., and (b) postinflammatory fibroses, in which the connective tissue continues to grow, as in *keloid*, after the irritant has ceased to act.

3. Postfibrinous fibroses, if we may so term them. The new connective-tissue formations which replace (a) thrombosed blood within the vessels, and (b) fibrinous exudates on serous surfaces, adhesions, etc., occupy an intermediate position between the two; they are replacement-fibroses to the extent that they replace the fibrinous coagulation, proliferative in that they are tissue where previously no tissue proper existed.

II. Of non-inflammatory origin.

1. Due to strain: (a) arterial, (b) venous, and (c) lymphatic fibroses, as above indicated.

2. Neoplastic.

CHAPTER III.

THE SYSTEMIC REACTION TO MICROBIC INJURY, THE PROCESS OF INFECTION.

FROM this consideration of the local reaction to injury we must now pass on to that of general systemic reaction, and, continuing in due sequence, before considering any one particular order of reaction, should first analyze the various noxe causing general bodily disturbance, whether physical or chemical, endeavoring to recognize sundry broad groups, each of which sets up disturbance of a particular order. Attempting this, we could distinguish certain conditions set up in which blood changes are primary or predominant; others, nervous disturbances; others, in which certain glands are picked out to bear the brunt of the reaction, and should have to consider, in turn, the effects of disorders involving one or other system, upon the rest of the organism. This systematic survey of processes affecting the several systems and their results we shall take up at a later date. To enter into it now would lead us to consider in series the progressive and regressive changes which may affect individual systems and organs before gaining an insight into progressive and regressive disturbances in general, and would thus lead to extensive repetition. It will be more serviceable to select for consideration the more common general processes, and that in the order of their frequency rather than of their relationship to one or other system. And, doing this, undoubtedly the first general process to be considered is the systemic reaction to microbic injury, or *infection*.

Definition.-Attention must be called to the double meaning of this term, as employed by the hygienist and the pathologist. For the hygienist, water, air, and other media may be infected, *i. e.*, infection consists in the mere presence of potentially harinful microbes, and the mere act of their coming into contact with the animal organism. The hygienist distinguishes between (1) sporadic infections, isolated cases; (2) endemic, where a notable proportion of cases of a given microbic disease is met with year after year affecting the inhabitants of a given region; and (3) epidemic, where a disease, of a sudden, affects a large number of inhabitants, the number of cases rapidly increasing and later decreasing. Diseases of animals may, similarly, be sporadic, enzootic, or epizootic ($\partial \chi \mu o \zeta$, the people; $\xi \omega o \nu$, animal). For the pathologist, infection is a process; for him the mere presence of pathogenic bacteria in the mouth or the skin, in the digestive tract, does not constitute infection: that is brought about by the growth of those bacteria, the diffusion of their products, and the reaction they induce in the organism is the essence of the process. For the pathologist, therefore, infection is the

INFECTION

succession of changes induced in the organism generally by the growth within it of microbes; or, in other words, it is the interaction between the organism and the microorganism. This interaction may, in the main, be local, and then the processes, occurring locally, constitute infective inflammation; the general disturbances which follow such local growth, or which are brought about by widespread proliferation of the microbes, constitute "general infection," or, briefly, "infection."

Causation.—We have already considered the mode of entrance of pathogenic organisms into the organism, as also, to some extent, the circumstances favoring their growth, and have considered also the subject of *susceptibility*, of imperfect reactive powers, so that there is inadequate destruction of microbes coming in contact with, or gaining entrance by any means into, the tissue. There is, however, another aspect of the subject. Bacteria may grow in the tissues, not because the tissues are weaker than normal, but because those bacteria possess a virulence over and above the power of the cells to counteract. We thus, to recapitulate, may make the following table of circumstances leading to the growth of bacteria with the organism:

I. Imperfect reactive powers (susceptibility).

(A) Individual susceptibility.

- 1. Inherited, whether as specific, racial, familial, individual.
- 2. Acquired.
 - As the result of previous attacks of disease set up by (a) the same, or (b) another species of microbe; as the result of injury, as the result of malnutrition, or as the result of exhaustion.

(B) Tissue susceptibility.

- 1. Inherent, the special susceptibility of certain tissues to become the seat of growth of certain microorganisms.
- Acquired through injury, local malnutrition, impairment of nerve supply, local exhaustion, or local disease.
- Pathogenicity of microbes (virulence) due to simultaneous entrance of either a small number of highly virulent microbes, or a large number of lowly virulent microbes.

It is the interaction of I and II which determines the development of infection; microbes of low virulence are capable of infecting susceptible individuals, and are without effect on those of normal resisting powers when introduced in equal numbers in the two cases; microbes of high virulence, if they induce disease in those relatively refractory, do not induce so grave a disease as in those relatively susceptible. The influence of the amount of infecting material or number of infecting organisms has been well demonstrated in the recent studies upon acute anterior poliomyelitis, in which it has been clearly shown that the incubation period is materially lengthened when, by filtration, through a Berkefeld filter, the amount of infective material is reduced.

THE COURSE OF INFECTION

THE COURSE OF INFECTION.

We have now to consider the results of bacterial growth, and how these results are brought about—the *process of infection*.

It will be well, in the first place, to sketch the course and features of some typical uncomplicated case of infectious disease; for such a purpose a case of typhoid fever affords a good example.

The patient has upon a given date taken, it may be, some milk coming from a farm where the hygienic arrangements have been imperfect, and where recently there has occurred one or more cases of the disease. For some days no ill effects are experienced, but then symptoms of malaise show themselves—slight but persistent headache, lassitude, some abdominal discomfort, with constipation, or, it may be, diarrhæa, pain in the back, and so on. These disturbances at first are so slight as to be regarded as transient, and do not prevent the patient continuing his daily duties; but they continue, and grow steadily worse, until, eight days or so after the contaminated milk had been drunk, the patient feels so weak and feverish that work is impossible, and he has to take to bed and call in a medical man.

We note, that is, a stage of *incubation*, during the latter part of which *prodromal* or *premonitory* symptoms show themselves, this incubative stage continuing until the onset of a definite *febrile state*. As a rule, for clinical purposes we date the illness from the first day of recognized fever.


Not to dwell upon unessentials, the medical man, when called in, finds the following condition: heightened bodily temperature, general muscular weakness, alterations in (a) the nervous system, manifested either by irritability and excitement or by lassitude and dulness; (b) in the circulatory system, shown by a rapid, full pulse, with evidences of vasomotor disturbance (flushing, dilatation of superficial vessels, etc.); (c) respiratory system—increased rapidity of respiration; (d) digestive system—dryness of the mouth (diminished salivary secretion), distaste for food (anorexia), obscure abdominal disconfort or pain, with constipation, giving place to looseness of the bowels and foul motions (or these may be present from the first).

Day by day, for a week or so, the temperature rises, until it may attain to 103° to 104° F., and, with this, all the other symptoms mental, nervous, muscular, circulatory, respiratory, and abdominal become more pronounced; and in addition (e) disturbances of the urinary system show themselves, in the shape of diminution of the salts of the urine, notably of the chlorides, and of increase in other constituents, notably the urates; while (f) a characteristic cutaneous eruption begins to show itself toward the end of the first week, in the shape of scattered "rose spots." Examination of the blood shows that this, also, is modified, for, while typhoid differs from the majority of infections, in there being little or no increase in the number of circulating leukocytes, in common with most, there is a definite diminution in the number of

INFECTION

red corpuscles per cubic millimeter, and, as shown by the Grünbaum-Widal test, the serum gains new properties, being able to cause the typhoid bacilli to become motionless and adhere together, or agglutinate, even when it is diluted fifty times.

This is the stage of *fervescence*, or pyretogenic stage. Following upon it, for a fortnight or so, is the *fastigium*, or stage of high fever, a period in which the febrile temperature is maintained at first at a constant high level; later, toward the end of the period, while toward the night it may reach the previous high level, it tends during the daytime to descend two or more degrees. Accompanying this febrile temperature, the other disturbances continue, and, as a consequence, the patient becomes increasingly weak and emaciated. In an uncomplicated case, however, toward the end of the third week an improvement manifests



Course of typhoid fever. (Modified from Musser.)

itself. The temperature shows a distinct tendency to fall, and each twenty-four hours the maximum may be a degree or more lower than the maximum of the previous day; there is refreshing sleep, the mind becomes clearer, appetite and craving for food manifest themselves, pulse and respirations become more normal, the motions less foul. We reach, in short, the stage of *defervescence*, and in four to six days the normal temperature is again attained. Now follows the stage of convalescence. The great emaciation and exhaustion of the tissues leaves the patient very weak, and attempts to move actively easily tell upon the circulation, etc., while any but the most easily digestible food tells upon the digestive system, and may easily favor a *relapse*. Gradually the patient returns to the status quo ante. I need not here dwell upon sequela (morbid conditions which follow a disease, and are due to the action of the same original cause, e. g., in typhoid, the formation of

abscesses in various parts, due to the B. typhosus; inflammation of the gall-bladder, set up by the same, etc.); or complications (morbid conditions of other causation, which may accompany the primary disease, e, g, in typhoid, the development of acute pneumonia, due to the Diphococcus pneumoniae, and inflammatory disturbances, brought about by B. coli infection)—all that we wish here is to give a fair picture of a typical infection resulting in recovery, so as to have a basis for the presentation of our subject.

The Period of Incubation.-Once bacteria or other microbes begin to grow in the tissues, from that moment we have the beginning of the infective process. But, although it begins thus, we are not able to recognize it. If the entrance of the germs be local, there is, inevitably, a preliminary period of local growth, with absence of general disturbance. Strictly, this, and only this, should be regarded as the period of incubation; in practice we cannot carry out this idea. We have to determine upon some one easily recognizable symptom, from the onset of which we can date the onset of active disease, and the supervention of fever affords us this useful starting point. Whence it follows that the period of incubation, in the clinical acceptation of the term, is made up of two stages—that of purely local growth of the microorganisms and local disturbance, and that of prodromal symptoms, in which the bacterial products, or even, in some cases, the bacteria themselves, have become to some extent generalized, and have originated disturbances in the system at large, but have not as yet caused a febrile reaction with other pronounced systemic disturbances. This period of incubation varies greatly in the different infections and in different individual cases. Certain bacteria are so virulent toward certain of the smaller animals of the laboratory that they cause death within four to six hours. In such cases there is not much time for the manifestation of an incubation period, but, even here, in cases in which to produce the most rapidly fatal results the infective germs are injected directly into the vessels, it has been shown by Lemaire¹ that if we take that blood and make cultures from it at intervals of half an hour there is to be noted a preliminary period in which the number of circulating bacteria is greatly reduced-a period of reaction-of destruction of the bacteria and removal of them from the circulation; then, apparently, the cells of the body become exhausted, and there follows a second period of rapid proliferation and increase in the number of the circulatory bacteria. At the other extreme, the incubation period may last for weeks or months. The longest period has been noted in rabies. Ordinarily in this disease the period of incubation varies from a fortnight to a month; but there are certain well-authenticated cases, though these are exceptional, in which as much as six months have intervened between the entrance of the virus into the system and the development of symptoms.

The relationship between purely local growth and the diffusion of

¹ Buxton has recently confirmed and amplified these observations.

INFECTION

the bacterial products during the prodromal period is most variable, so that only in a certain class of cases is it possible to recognize this period of incubation with any definiteness; add to this that in certain diseases (*e. g.*, cholera) actual fever may be largely wanting, and thus we have to accept other symptoms as indicating the commencement of the active stage of infection.

The infective microbes also vary greatly in the extent to which they proliferate before they, through their products, induce general disturbances. Thus, to give a few examples of the variations met with:

1. There may be a minute boil or furuncle on the face, due to the local growth of a streptococcus. Although this growth is so local, there nevertheless is rapidly induced a slight febrile condition, with general malaise. The germ remains local, and the infection is singularly local; nevertheless, the general disturbances set up by diffusion of the bacterial products is relatively profound—out of all proportion to the extent of local disturbances. The development of the febrile state here is not coincident with any diffusion of the bacteria themselves. It even precedes the period of maturation of the boil.

2. In tetanus there is a similar strictly local growth of the bacteria, and the same is the case in diphtheria. In both these cases there is a well-marked period of incubation, in both the supervention of the febrile state indicates only that the diffusion of the bacterial products has reached a point at which the amount of concentration of those products is sufficient to induce severe disturbances in certain tissues away from the period of local growth.

3. On the other hand, in smallpox the development of the febrile state coincides with the earliest appearance of the cutaneous papules. The presence of these papules undoubtedly indicates that the virus has entered the blood from the focus of primary local infection, and has been carried through the system, and so to the vessels of the skin, during the period of incubation. Here, also, it may be noted, the site of primary entrance and primary local growth remains still to be discovered.

4. Lastly, in disease like tuberculosis there is primary local growth, but that localized growth develops so gradually, the toxins are so gradually diffused, that it is difficult, if not impossible, to recognize any one period at which incubation develops into general infection. We can, at most, speak of a pretuberculous stage, during which the tuberculin reaction affords an indication of the presence of the tubercle bacilli within the tissues.

From these instances it is clear that what determines the development of symptoms of general infection is not the presence of the specific bacteria circulating throughout the body, or even the extent of the local inflammatory disturbances set up by them, but is the toxicity of their products and the relative amount of the same. And the length of the incubation period in any given case is determined by several factors:

1. The toxicity of the products of a specific microbe.

2. The amount of toxic substances developed in a given time (this

depending upon the number of germs gaining entrance, and the virulence of those germs).

3. The neutralizing powers of the organism (whereby, in the same disease, the incubation period in different individuals is found to vary, often to the extent of several days).

4. A fourth factor is indicated by the researches of Roux and Yersin, and Sidney Martin, already noted (p. 315), namely, the time taken by the enzymes primarily excreted by the pathogenic organisms to act upon certain constituents of the fluids or tissues of the body and convert them into toxic albumoses, the direct agents in setting up the symptoms of generalized disturbance. This, however, is still a matter of debate.

Thus, to sum up: The period of incubation, as generally understood, consists of at least two stages, the first, of varying length, that of purely local growth and local disturbance, the second in which there are prodromal symptoms—general disturbances of a relatively mild type, due, not to the bacteria themselves, but essentially to the diffusion of their products of growth into the general circulation. These products of growth may not, in all cases, be themselves toxic; it is possible that they may by enzyme action convert certain of the body proteins into toxic bodies (albumose).

Grades and Types of Infection.-As noted upon page 458, we can classify to some extent the pathogenic microorganisms according as to whether, in the main, their growth in the system is local or whether it is diffuse, setting up a *bacteriemia*. This distinction, useful or otherwise, is for the present purposes largely useless. In the first place, there are many intermediate grades, the typhoid bacillus, for instance, has mainly local growth, but to be present in and produce the rose spots, must circulate in the blood; the sequelæ of pneumonia convince us more and more that the pneumococcus, while it has its seat of election in the lungs, passes readily into the blood. Secondly, one and the same organism, like the Pyococcus aureus or the streptococcus, may in one individual have a purely local growth; in another, set up extensive septicemia; and in the two cases the symptoms vary in degree rather than in kind. While thirdly, in both orders the symptoms are brought about by one common cause, namely, the diffusible toxins. It is the nature and toxicity of those products that are the prime factors in the development of infections; they cause the characteristic disturbances of metabolism. According as to how these products tell upon the organism, and how the organism is capable of neutralizing or destroying them, so can we recognize the following types of infection: (1) Fulminating or malignant. (2) Acute. (3) Chronic (persisting subacute). (4) Subinfection.

1. Fulminating or Malignant Infection.—Occasionally we find in man, or in the lower animals, that an infection is from the first so rapid in its development that we are hopeless of obtaining a favorable issue; the organism appears to be incapable of adequate reaction. Some reaction there is in each case, but it is wholly ineffectual, and becomes less and less marked. The symptoms are those of a rapid intoxication,

INFECTION

with depression of the functions. The heart beat and pulse become more and more rapid and feeble, with great lowering of the blood pressure; the respirations very rapid and shallow; there is pronounced "typhoid state," with coma and depression of the higher mental centres, and no excitement; the temperature may from the first rapidly sink, until it is far below the normal; if the disease be one characterized under ordinary conditions by well-marked leukocytosis, there is found pronounced and progressive leukopenia (or lack of leukocytes), and a condition of collapse and coma quickly gives place to death. All this may take place within a few hours. Experimentally, by "passage," we can so exalt the virulence of certain bacteria that we can kill the animals of the laboratory within six hours, and that with the minimal injections. But not with all bacteria; no amount of passage will reduce the period of experimental tuberculosis to less than ten days. Difference has to be acknowledged. In man, both exaltation of virulence and entrance of excessive numbers of bacteria at a time would seem to play a part. Such fulminant cases are more particularly seen in tropical countries-in cases of cholera, yellow fever, the plague. Here both the general bodily health is lowered by the surroundings, and pathogenic organisms at the higher external temperature can proliferate and multiply outside the body and at the same time retain their virulence. But such cases are far from being unknown in temperate climates. We have ourselves obtained autopsies and gained pure cultures from a case of Streptococcus peritonitis fatal within twelve hours from the onset of the first symptoms, and of epidemic cerebrospinal meningitis, fatal within six hours; malignant hemorrhagic cases of smallpox, scarlet fever, typhoid, fatal within twelve hours, are far from being unknown. These hemorrhagic conditions are indications that the circulatory toxins are sufficiently concentrated to act upon and cause degeneration of the epithelium of the capillaries, with weakening of the wall and subsequent rupture. With this often there is rapid destruction of the corpuscles and diffusion of their hemoglobin. We can, to some extent, reproduce this hemorrhagic state by injection of the sterile fluids of growth of virulent organisms into the vessels of lower animals.

In all these cases the slight, or absent, febrile reaction is especially noticeable. Indeed, it may be laid down as a general rule that if, in a case of infection, the temperature becomes rapidly subnormal, and the condition of the patient, with the lowering of the temperature, instead of improving, becomes worse, death may be expected in the course of a few hours. The cells of the body, instead of being stimulated, are paralyzed by the toxic substances; metabolism, and in consequence heat production, is arrested, and the fall of the temperature is an indication of intense intoxication.

2. Acute Infection.—This is the ordinary "type" infection already described, that characterized by a definite incubation period and a febrile stage or fastigium, which, according to the intensity of the process and the resisting powers of the system, either terminates in death or in defervescence and convalescence. The process of recovery may be interrupted by the recurrence of one or more *relapses* or *remissions*

GRADES AND TYPES OF INFECTIONS

(repetitions of the symptoms and disturbances characteristic of the primary disease), or by *sequelæ* or by *complications*. In most cases defervescence is gradual, a matter of some days; we speak then of recovery by *lysis*. In other cases (as often in acute lobar pneumonia) the temperature may fall to the neighborhood of the normal within twenty-four hours; we speak then of recovery by *erisis*.

In accordance with custom, and, we must admit, there is a certain convenience in so doing, we shall devote a separate chapter to the febrile state and fever, *i. e.*, to a consideration of the processes occurring in actual infection; and in that connection, also, it will be best to discuss the process of resolution and the cause of relapse, these being all allied subjects.

3. Chronic Infections.—Another well-marked class of infectious is characterized by insidious development, long continuance, with termination either in death after the disease had lasted for months, or it may be years, or gradual recovery. The whole process is prolonged, and may lack any sharp definition into successive stages. Examples of this class are to be seen in tuberculosis, syphilis, glanders, and actinomycosis.

In syphilis, a division into successive stages is more a matter of clinical convenience than of absolute fact—at least, as between the secondary and tertiary stages, for so-called tertiary lesions may exist along with secondary, and, as between primary and secondary, the transition is variable, and may be very insidious.

In all of these it would appear that there exists a focus of primary infection; in some, as in syphilis, it is very obvious. For weeks, as in syphilis, or months, as in actinomycosis, or permanently, as in mycetoma pedis (a disease closely allied to actinomycosis, but in general only affecting a lower extremity), we have but to deal with the local growth of the microorganism originating the local disturbances, with more or less pronounced general disturbances of a febrile nature. Of the nature of this local growth we have already spoken (p. 440).

A second subgroup is the *chronic remittent*, best represented by the rheumatic group of disorders, in the causation of which the indications at this present time are that more than one form of organism is concerned. Here we deal with conditions which may begin insidiously, but often acutely. In either case the progressive nature of the disorders set up indicates that there is not with defervescence complete recovery or total destruction of the pathogenic organism, and, while the progress continues slowly to act upon the joints or the heart valves, from time to time the condition lights up again into an acute form.¹

¹ The more recent work upon this etiology of acute rheumatism, in connection with Poynton and Paine's organism (Lancet, November 11, 1905; 860 and 932) and the Streptococcus pyogenes, is given by Beattie. Journal of Medical Research, 14: 1906; 399. We fail to see that any etiological distinction can be drawn between acute and chronic remittent type; indeed, in our laboratory at the Royal Victoria Hospital, from the discorganized hip-joint of a man who has suffered from such remittent rheumatism for twenty years, and was wholly crippled thereby, we gained abundant diplococci, which, cultivated by Dr. G. A. Charlton, exhibited all the characters of Poynton and Paine's organism.

INFECTION

In all these cases it would seem that the microbes, while developing their toxins at a relatively slow rate, are themselves distinctly resistant to the action of the tissues and bodily humors. On the other hand, growing and producing their toxins slowly, there is neither the same extent of intoxication nor the same well-marked reaction on the part of the system which we see in acute infections. There is, nevertheless, fever, though this tends to be of a more remittent type than is seen in the majority of the acute infections, and there is progressive emaciation and weakness as the disease advances.

Subinfection.-The appreciation of the fact that from the alimentary and respiratory tracts bacteria are constantly being taken into the system (p. 319), leads us to recognize the existence of yet another condition-that in which those bacteria, pathogenic and non-pathogenic instead of proliferating, are destroyed in the various tissues and organs to which they may be carried by the lymph and blood streams, a condition for which we have suggested the name subinfection.1 Normally, as we have indicated, the taking in of bacteria is relatively slight, and the exercise of what are strictly the physiological functions of the cells in bringing about the destruction of the same leads, we may be assured, to no disturbance, either local or general. There are, however, conditions of congestion and chronic slight inflammation of the intestinal mucosa in which there is an accompanying great increase in the passage of leukocytes into the submucosa, and thence between the epithelial cells into the lumen of the bowel; and as a consequence both of the increased passage out of the leukocytes and of the increased proliferation of intestinal bacteria, which accompanies, if it does not cause, the inflammation in question, there is increased taking in of these bacteria.

We obtain evidence of this increase by examination of the mesenteric glands and liver cells. In cases of chronic intestinal irritation we find in them abundant minute granules, which, at first, one is liable to regard as pigment granules. Indeed, we acknowledge that it takes long study before one can rid himself of the conviction that this is not the case. But thorough study and careful focussing of the sections under a very high power-a one-eighteenth inch immersion, for example -has convinced us that these granules are some of the final stages of bacterial destruction. They may be single; most often they are in pairs, resembling diplococci; rarely in sets of three, or rows of four, and often a distinct halo, as of a digestive vacuole, can be seen around them. Some hours after injection of B, coli intravenously into the lower animals, identical appearances are to be seen in the liver cells, and we have noted that the first stage in the taking up of the bacilli by the endothelial cells of the hepatic capillaries is the conversion of those bacilli into similar though somewhat larger diplococcoid bodies and sets of three or four granules. Then these bodies disappear from the endothelium, and, we hold, are represented by those more reduced bodies in the liver cells and, it may be added, in the bile.

¹ Journal of the American Medical Association, 33: 1899: 1506 and 1572.

SUBINFECTION

The effects of such continued passage of considerable numbers of bacteria into the system must be equivalent to the growth of the same within the tissues, and their destruction, if long continued and excessive, should bring on cell exhaustion. The frequency with which, in cases of cirrhosis of the liver, we have been able to gain cultures of intestinal bacteria from the liver, as also from the ascitic fluid during life, has led us to suggest that these are at least one factor in the production of that condition, and Weaver and Hektoen have isolated a form allied to the B. coli, with which they have set up cirrhosis in guinea-pigs.¹

We do not pretend that these are a constant factor in all cases of cirrhosis, even of Laennec's type, or that in any one case they are the only factor. There must first be intestinal irritation, whether by alcohol,

by acid fermentation, or otherwise. But we believe that they play an important role in the etiology of a large proportion of cases.

The almost constant evidence of old gastritis, the known hemolytic powers of members of the coli group, the existence in abundance of similar diplococcoid bodies in the liver cells, leads us to believe that in pernicious anemia some member of this group, or some other species of bacteria possessing strong hemolytic powers, is likewise involved. In support of this view, Charlton, ² working in our laboratory, has been able to produce not, it is true, a typical pernicious anemia, but a singularly grave anemia, accompanied by poikilocytosis and the presence of normoblasts, by successive inoculations of



Swollen endothelial cell of capillary of rabbit's liver containing Bacillus col in various stages of degeneration, within thirty minutes of injection of the bacilli into the blood stream. Part only of the nucleus is shown in the section,

a B. coli of low virulence into rabbits, the original strain having been obtained from the intestines of a healthy normal rabbit.

Nicholls,³ also, has demonstrated the existence of similar minute diplococcoid bodies in the cells of the tubuli contorti and elsewhere in the kidneys, and concludes that they are a factor in some cases of chronic interstitial nephritis.

It is but right to warn the reader that these observations and views have not as yet gained general acceptance. We mention them because

¹ As this work has been passing through the press, Opic (Trans. Assoe. Amer. Physicians, 1910, about to be published) has shown that while agents like chloroform, which cause necrosis of the liver cells, given alone do not lead to cirrhosis in animals of the laboratory, and while inoculations of B. coli alone may give negative results, the combined action induces a well-marked typical cirrhosis,

² Journal of Medical Research, N. S., 3: 1902: 344.

³ Montreal Medical Journal, 28: 1899: 161.

we are firmly convinced of their correctness, and because we believe that they throw light upon certain very obscure forms of disease.

Exogenous Bacterial Intoxication.-From such conditions we pass next to what, in our classification of intoxications, we speak of as exogenous intoxications of saprophytic origin. There is, it will be seen, but a slight step from the introduction of bacteria into the tissues, destruction of them forthwith, and liberation of their toxins, to the absorption of the toxins liberated by bacterial growth in the intestinal canal, the bacteria themselves not being to any extent taken into the system. Such conditions do not come within our definition of the process of infection. It is well, however, to call attention to the same in this place, the more so because Hunter' and others have ascribed pernicious anemia and allied conditions to such absorption of toxins. In diphtheria and cholera we have two pronounced infections which only just come within the terms of our definition, for the bacteria here only grow upon the surface of the mucous membranes and scarce enter the tissues. It is true that they destroy the surface layers, and this destruction doubtless aids very materially the absorption of the toxins. But even when the surface layers are not destroyed, it may well be that extensive absorption occurs of bacterial toxins. Where fermented feces are retained we know that malaise and an actual febrile state may ensue. But in such cases it is impossible to analyze and distinguish between the absorption of bacterial toxins and that of the products of fecal disintegration, and, we may add, the entrance of bacteria into the tissues. At most, we can, it seems to us, admit the existence of this order of condition.

The various orders of infection proper pass the one into the other (save that a malignant case does not become less acute), and there may be various intermediate stages. An acute infection may suddenly assume the malignant type, or may from the first tend toward malignancy. We have noted how an acute or subacute may recur and be associated with chronic disturbance. A microorganism which ordinarily induces chronic infection (e, q), the B. tuberculosis) may set up a disturbance that is distinctly acute, either by gaining entrance into the blood stream and consequent wide distribution and multiple foci of growth, or by attacking an individual of greatly lowered vitality, and, lastly, a study of "terminal infections" would indicate that eventually subinfection is liable to give place to acute infection-germs of low virulence, ordinarily destroyed by the cells coming to grow actively in the tissues when the resisting powers become lowered beyond a certain minimum. The study of such terminal infections supports the dictum of Osler, that "persons rarely die of the disease with which they suffer."

¹ Pernicious Anemia, London, Griffin, 1901.

CHAPTER IV.

THE SYSTEMIC REACTION-(Continued).

THE FEBRILE STATE.

This study of the different types of infection prepares us to consider the nature of the reaction between the organism as a whole and the microorganism, of that chain of disturbances which, collectively, we speak of as *fever*; or, if it be objected that this term should be only employed to indicate heightened temperature, as *the febrile state*.

Here, once again, we confront a difficulty, due to the prevailing laxity in the employment of terms. The oldest definition, that of Galen, describes fever as *calor prater naturam*, and, if influenced by tradition alone, as such and such only should we consider the condition. But two very distinct orders of events, at least, bring about this calor proter naturam, namely, the diffusive microbic products and certain other substances, and traumatic and other disturbances of the nervous system, regarding both of which, it is needless to say, Galen was in supreme ignorance. To retain the old definition demands that we persist in grouping together phenomena of unlike origin, in opposition to all right principles of classification. The more modern acceptation of the term is thus to be commended-that which regards fever as a particular train of symptoms and changes in the organism, associated with heightened bodily temperature, constituting the infective reaction. We would even say that the phrase "associated with heightened temperature" is inserted in accordance with usage and clinical demands, and this because the same train of disturbances may present itself without materially heightened temperature. We must consider all these cases together, and thus, for convenience, lay down that:

1. The reaction to infection may be with or without rise of temperature (may be "febrile" or "afebrile").

2. That throughout, mere heightened bodily temperature, however produced, will be referred to as *pyrexia*.

This course is not absolutely logical, but it is the only one practical under the circumstances. The causation of pyrexia and the subject of thermogenesis in general will be discussed in a separate chapter.

In the febrile state, then, we observe increased bodily temperature, along with that series of disturbances of the circulatory, nervous, respiratory, and digestive systems and the excretory organs which we have noted in our description of acute infection. Passing, now, to describe and analyze more fully these various changes in the different systems and organs, we shall be in a position to recognize more surely the nature of the fever.


The Febrile Temperature Changes.-The Stages of Fever.-As already noted, we can, in most acute infections, distinguish three stages:

1. The Pyretogenic.—This may be short, with rapid rise in the course of an hour or two, as in ague, or, as in typhoid, may extend over several days.

2. The Fastigial.—This, again, according to the nature of the process, may be brief, or may extend over several days.

3. The Defervescent.—This may be brief, the temperature diminishing rapidly by crisis, or gradual and prolonged-lysis.

In some fevers, as in typhoid, toward the conclusion of the fastigium, but before defervescence occurs, the temperature, from having previously shown but slight daily oscillation, may present a daily variation of several degrees. It is then termed amphibolous. It must be kept in mind that during the fastigium the daily temperature variations are in general of the same order as those seen in health, with rise toward evening and fall in the early morning.





0

The Varieties of Fever.-According to the temperature changes, we can distinguish:

1. Continued Fever.-In these, during the fastigium, the daily temperature changes, though occurring at a higher level, are little if at all greater than those seen in health (typhoid, pneumonia, etc.).

2. Remittent Fever.- Here the daily temperature changes may extend over several degrees and the temperature curve, while continuing above the normal, is very variable (pyemia, suppurative changes, tuberculosis accompanied by secondary infection).

3. Intermittent and Recurrent Fevers .- In intermittent fever there is a succession of febrile attacks, each characterized by pyretogenic, fastigial, and defervescent stages, and separated by intervals of twenty hours or more, the temperature during the interval being normal (malarial fever). A variety of the form, in which the interval is of

THE ASSOCIATED FEBRILE DISTURBANCES

several days' duration, is sometimes spoken of as *recurrent* (Relapsing fever proper, Malta or Mediterranean fever).

Stages of Pyrexia.—Here it will be useful to note the terms employed in describing and classifying the different grades of rise of body temperature occurring in febrile states. For in fevers we have every grade of rise of bodily temperature, and in the same infection different individuals react differently. In children, for example, a very slight disturbance is liable to cause a profound rise; in the aged, on the contrary, severe infection may be associated with relatively little increase. Wunderlich's divisions are usually followed.

Subfebrile, or high normal, fro	m .	37.5° to 38° C., or below 100° F.
Low febrile		38.0° to 38.5° C., or 100° to 101° F.
Moderately febrile		38.5° to 39.5° C., or 101° to 103° F.
High febrile		39.5° to 40.5° C., or 103° to 105° F.
" " morning, above		39.5° C.
" " evening, about		40.5 ⁵ C.
Hyperpyrexial .		$41.0^{\circ}\mathrm{C},$ and over, or $105^{\circ}\mathrm{F},$ and over.

Reversing the usual procedure, we shall not now discuss the causation of this increased temperature, but will discuss the disturbances other than pyrexial that characterize the febrile state, believing that by this means it is possible to gain a more thorough grasp of the subject and to come more fully prepared to the discussion of pyretic phenomena.

THE ASSOCIATED FEBRILE DISTURBANCES.

Nervous Disturbances. Chills. During the pyretogenic stage in very many fevers a marked feature is the supervention of one or a succession of chills. The patient feels cold, the teeth chatter, the sensations experienced are identical with those which follow exposure to cold with rapid cooling of the surface of the body. But now the hand, and the thermometer, often indicate that the surface is distinctly hotter than normal, while, at the same time, the thermometer in the rectum shows there a vet greater rise of temperature. We are dealing clearly with a nervous phenomenon, and one that is not the direct effect of cooling upon the cutaneous nerve-endings. It is true that the surface phenomena, save for this frequent increase in heat, are closely allied to what is seen in actual cooling; the extremities and the face may be This view that we are dealing with a nervous phenomenon, incited from the central nervous system, is supported by the fact that in those of unstable nervous constitution identical chills may occur without exposure to cold and without infection. What the nervous change is, that is, at the root of these chills, it is difficult to say. Marey' has

La circulation du sang, Paris, 1881.

THE SYSTEMIC REACTION

suggested that relative increase in the temperature of the central organs may produce the same results as relative decrease in surface temperature, and that relative temperature possibly plays a part is suggested by Recklinghausen's⁴ observation that when the chills have passed and high fever has developed, they may be brought on again by exposing an extremity. With relapse, also, in the course of a fever they may show themselves. Mere local anemia of the surface vessels, which is common both in cooling and in chills, would not seem to be a satisfactory explanation, for the same may occur under other conditions without chills being produced.

But, underlying these chills, we must see that the local anemia indicates that the blood is attracted to other organs, that there is a corresponding congestion elsewhere; as, also, that the relative cutaneous anemia indicates during this period a relative storage of heat in the system. Nor is this all; calometric observations demonstrate that more rapid and marked increased production of heat occurs during the chills than at any other period of the febrile state. Why this is will be explained on the next page.

With the onset of the fastigium the surface vessels become congested, and now there is a sensation of general surface and body heat—of feverishness.

Other Febrile Nervous Disturbances.—These are but one of a series of nervous disturbances, which we may divide into two categories, namely, states of nervous irritation and of nervous depression. Among the former we must class headaches and mental irritability, photophobia, sleeplessness, hallucinations, and the graver conditions of active delirium, with confusion of intellect. This may pass on into the next state, that of exhaustion and depression. Other conditions of the second category are apathy, arrest of mental activity, prostration and involuntary passage of excreta, quiet muttering delirium, and complete coma and collapse.

While the different conditions are, to some extent, an index of the severity of the fever, mental and nervous exhaustion being matters of graver import than are irritability and active delirium, we have to recognize that in different forms of infection there is a wide variance in the extent of the nervous disturbance. While in some cases we have to deal with active meningitis and presence of the specific organisms directly affecting the surface of the brain, and in others, it may be, with alterations in the circulation of the brain rather than with direct stimulation, in the main these nervous disturbances have to be ascribed to the action of the diffusible toxins circulating in the blood. They correspond with the nervous disturbances see up by other toxic agents, and, while they vary so greatly, we can reproduce one or other group by the intravenous injection of the sterile culture fluids or products of growth of specific pathogenic microbes. The most striking demonstration of this direct action of toxins upon nerve tissues has

¹ Hdbch. d. allgem. Pathol. des Kreislaufs, Stuttgart, 1883: 451.

MUSCULAR DISTURBANCES

been afforded by Meyer and Ransom,¹ in their studies of the course pursued by the tetanus toxin, in which they showed with absolute precision that this toxin has a direct affinity to, and selective action upon, the nervous tissues, and passes up the peripheral nerves. Di Vestea and Zagari² had, some years previously, demonstrated the same remarkable passage in connection with rabies, but there it is still undetermined whether we deal with the passage of the toxin or of the infective agent. Recently, a like predilection has been demonstrated by Flexner and Lewis,³ and by Landsteiner,⁴ in connection with the virus of anterior poliomvelitis.

In diphtheria, which likewise is characterized by definite nervous symptoms, Rainy⁵ and others have shown that the toxins produce direct histological changes upon the motor cells of the cord in cases in which paresis was present during life. Sidney Martin had previously noted the destructive effects of those toxins upon the nerve fibres. Yet earlier, Charrin,⁶ by injection of culture fluids of the B. pyocyaneus, had brought about in the rabbit the same spastic state of the hind limbs, with paralysis of the sphincters, etc., which are features of the blue pus disease in that animal, and, as Williams and Cameron,⁷ of Montreal, were among the first to point out, are features of the disease in the human infant. The evidence, therefore, that bacterial toxins are capable of acting directly upon the nervous system is very definite, and such action affords the simplest and most direct explanation of those nervous disturbances in infections generally which are not obviously the result of exhaustion.

Muscular Disturbances.—Rigors.—Associated with chills are rigors —fine fibrillary contractions of the muscles of the face, trunk, and extremities, involuntary in nature. Like chills, they are common to exposure to cold and to the incipient stage of many fevers. They represent a reflex stimulation of the muscles whereby, through the frequent and rapid contraction of the individual fibres, no definite movements are induced; in fact, opposing muscles are synchronously affected. The most that is produced is what corresponds to an increased stiffening up of the muscles generally, well indicated by the term "rigor." Every whit as much as the flapping of the arms of the chilled coachman, these rapid individual contractions mean work, and muscular work means the giving off of heat. Rigors, in short, are a mechanism whereby there is produced reflexly increased heat production, and, occurring simultaneously with the febrile chill, suggest strongly that they are the effective means of inducing increased heat produced is the second strong in the second strong in the second strong is the second strong in the second strong is not a second strong in the second strong is not a second strong in the second strong is not a second strong in the second strong is not a second strong in the second strong is the second strong is not a second strong in the second strong is not a second s

¹ Schmiedeberg's Arch., 49: 1903. An excellent résumé of this important article is given by Archibald, Montreal Medical Journal, 34: 1905; 874.

² Archiv p. l. Scienze mediche, 9:1887.

³ Jour. Amer. Med. Assoc., 53: 1909: 1913, and 2095.

⁴ Landsteiner and Popper, Ztschr. f. Immunitätsforsch., Orig., 2; 1909; 377.

⁵ Journal of Pathology, 6: 1900: 444 (with good bibliography).

⁶ La maladie pyocyanique, Paris.

⁷ Journal of Pathology, 3: 1895: 344.

production. Similarly, it has been shown that the muscular activity associated with the more rapid respirations of the febrile state is accom- **#** panied by increased heat production.

Other Muscular Disturbances.—Atrophy.—It is apt to be forgotten that the normal muscle is not only working when it is undergoing active contraction, but works and produces heat in the apparently resting condition. The condition of tonus is a state of partial contraction, and it can be demonstrated by recording the finger or hand movements after active exercise, or, in cases of paralysis agitans, that the fine twitchings have a definite rate per second, the irregularity of the curves tending to be regular, and brought about by interference between the rates of stimuli passing to opposing groups of muscles. Increased contraction and tonus clearly play a part in the earliest stages of the febrile state; later, they give way to muscular relaxation and exhaustion, and the muscles of the body in general are noticed to diminish in size at a greater rate than is to be explained by the combined lack of exercise and diminished assimilation. This rapid "burning up" of the muscles is another indication of increased heat production.

Circulatory Disturbances.—In certain cases, the so-called *sthenic* fevers, and, in certain stages of acute infections in general, we find the pulse full and bounding; in others—the *asthenic* fevers—it is weak and easily compressible. In all cases it and the heart beat are markedly increased in rate. While it may be laid down as a valuable rule for prognosis that strengthening of the pulse and lowering of the rate in any individual case of fever is a favorable sign, we have to confess that we know sadly little concerning the meaning and the causation of the febrile pulse; the factors possibly concerned in any given case are too many to permit a sure analysis. What these factors are we will briefly indicate.

1. Whenever the bed of the blood stream is widened in any considerable area without corresponding contraction of the bed in other areas, there is lessened resistance to the inflow of the blood, the pressure sinks, and the heart rate increases. The general lack of bodily tone and the actual vascular dilatation observed during the fastigium indicate that this factor is at work.

2. Increased temperature of the blood and organism generally has an identical action. If the frog's—or the cat's—heart be removed from the body and kept beating by supplying it with blood, warming that blood leads to more rapid and less powerful heart action; as, again, warming the rabbit's ear leads to obvious vascular dilatation and increase in stream bed.

3. The direct action of bacterial toxins upon (a) the peripheral vessels has also to be considered. And this action varies in the different infections. With diphtheria, Sharp¹ found that the toxins applied direct to the heart muscles cause, first, a more powerful heat, followed by weakening, with less complete systole and more prolonged diastole.

¹ Journal of Anatomy and Physiology, 31: 1897; 199.

CIRCULATORY DISTURBANCES

until complete arrest ensued in the diastolic state. Woodhead found that the same toxin leads directly to a condition of fatty degeneration. But, at the same time, we know that it has a specific action on the peripheral nerves, and disintegration of the vagus has been determined in some cases of sudden death, which vagus degeneration would seem the most satisfactory explanation of this terrible sequel of diphtheria. A similar action upon the vagus has been determined for influenza toxins (la grippe). Arloing and Courmont have found that the products of the Pyococcus aureus lead to capillary dilatation; those of the B. pyoeyaneus, according to Charrin and Gley, lead to contraction of the arterioles, but these at the same time (Mora and Doyon) act directly on the vagus, arresting its inhibitory action and causing increased rapidity of beat. These examples, alforded one and all by capable observers, will serve to indicate the intricacy of the subject and the need to study the cascular changes in each infection separately.






FIG. 159

Trainee taken from successive starge of a persystem of ague (tertian malaria). Fig. 15

Tracings taken from successive stages of a paroxysm of ague (tertian malarin). Fig. 150 from the period before an attack. Fig. 157, beginning of the attack (stage of shivering). Fig. 158, in height of attack. Fig. 150, beginning of the stage of defervescence (sweating stage). (Mannaberg.)

4. Yet another factor in affections such as cholera, which are accompanied by profuse diarrheea, is actual diminution in the amount of the blood; this, again, leads to rapidity of heart action and weak pulse. The results, so far as the heart is concerned, are the same as diminution of resistance to onward passage.

It follows, naturally, that the different stages of a fever afford different types of pulse and pulse tracings; that in the stage of contraction of the peripheral vessels the pulse is small and firm; in the fastigium it is of a fuller, softer type. It further is particularly liable to exhibit dicrotism, an indication, as now generally accepted, of loss of tone of the walls of the larger arteries. On these, also, it would seem, toxins have a direct action. The accompanying tracings of the pulse in the apyretic period and the period of chills (pyretogenic), heat (fastigium), and perspiration (defervescence) of two malarial paroxysms on successive days exhibits well the changes in type and rate of the pulse in these different conditions.

Alterations in the Blood.—We shall not attempt to describe minutely the blood changes in the different infections; they are very various, and are detailed fully in the many excellent works on hematology. We can but note the main features.

Red Corpuscles.—It may be laid down that, even in those cases in which there is a severe drain of fluid from the organism, as in cholera, when, in consequence of the concentration of the formed elements, they appear to undergo a great increase, infection, even in mild grades, if continued for but a few days, leads to diminution in the number of the erythrocytes. Their destruction in the febrile state is greater than their reproduction. Certain microbes, streptococci, and many members of the B. coli group have, through their toxins, a distinct hemolytic action upon the corpuscles in the test-tube, and if a smear be made from the swollen spleen, in typhoid and other fevers, it is often possible to note that the large endothelial cells from the splenic sinuses contain more or less degenerated red corpuscles.

White Corpuscles.-Here a very considerable variation exists in the different infections both as regards the total number of leukocytes found in the blood and the relative proportion of the different forms. In most cases there is a distinct increase, in some, as in pyemic conditions, streptococcic infections, pneumonia, the general increase is very marked; in others, notably in typhoid, not only is there no leukocytosis, but the number may be below the normal. Where, in a disease characterized by absence of leukocytosis, a sharp rise in the number of leukocytes is observed, we have indications of a second infection. In typhoid, for example, such leukocytosis should suggest the possible development of perforation, but it must be kept in mind that a rapidly developing septic condition, particularly where the patient is already in a weak state, may be accompanied by no leukocytosis. Where, in infection characterized by leukocytosis, the number of leukocytes steadily, or even rapidly, increases during the fastigium, this need occasion no great concern; it is, if anything, an indication of good reactive powers; while, when the fever is at its height, and when the number shows a pronounced fall, or, throughout, the leukocytosis is deficient, we have a condition of serious, not to say grave, import-an indication of deficient reaction and not improbable fatal ending. Where a moderate fall occurs in a case that has proceeded in a natural manner for some period, we may have indications of approaching resolution; indeed, Kanthack by this sign was able in a large number of cases to predict the oncoming crisis in pneumonia twenty-four hours before the temperature began to drop, or any other symptom of the event had declared itself.

Regarding the different forms of leukocytes noticeable by their frequency in the different infections, it may be said that, in acute infections, the polynuclears are the predominating form; in more chronic diseases, such as tuberculo is the lymphocytes are relatively frequent, without the

RESPIRATORY DISTURBANCES

other forms showing great increase in number. In the febrile conditions accompanying the presence of intestinal worms, as already noted (p. 319), there is a liability for a striking increase in the number of cosinophiles, with or without pyrexia.

Respiratory Disturbances.—The increased rate of respiration is a cardinal symptom in the course of infection. Such increased rate is common to all conditions in which either the body is exposed to heightened external temperature, or the bodily temperature itself is raised. It is one of the means of discharging and reducing the heat of the body, for, not merely is heat lost in the process of warming the inspired air, and more heat lost by inhaling and exhaling a large quantity of air in a given time, but also, by internal evaporation, the moistening of the air in the lungs, further loss of heat occurs.

But even when, as in a vapor bath, the external air is already both saturated and heated, the same increased rate of breathing is to be noted, rendering it doubtful whether loss of bodily heat is the main function of the increased rate. We must pass beyond the mere act of increase in rate to its cause, and lay down that anything which leads to increased temperature of the blood bathing the respiratory centre is accompanied by increased rate of respiration; this is very largely an automatic process, primarily resulting, it is true, in lessening the bodily temperature, and so that of the blood passing to the cord, but still occurring when the end cannot be obtained, or when the increased temperature is beneficial to the rest of the organism.

There is another cause of increased rate, acting also, we believe through the respiratory centre in the cord, which appears to have a more immediate bearing on the febrile state.1 We would refer to this as oxygen-hunger, even though it acts through its results, namely, accumulation of carbonic acid gas in the blood. We know from abundant observations that increased temperature favors increased metabolism in the individual tissues, and heightened metabolism means increased using up of oxygen, and results in an increased discharge of carbonic acid. Under ordinary physiological conditions, if the body be exposed to heat, we note a tendency to combat this heightened metabolism; the individual indulges in a minimum of exercise, and little food is taken. In fevers we have the striking phenomenon that despite the anorexia, the bedridden condition, and complete lack of muscular exercise, the amount of oxygen absorbed in a given time continues to be greatly increased, as also does the amount of carbonic acid given off. And the higher the body temperature, the greater the absorption of the one and the discharge of the other. As Haldane has shown (p. 382), it is the increased CO, tension in the circulating blood that is immediately concerned in acceleration of respiration. In one case of febrile chill,

¹ The conflicting views of Hering, Breuer, and Head (whose studies on the vagus led them to conclude that respiration is largely a reflex act, determined by the condition in the alveoli of the lungs) are weighed and criticised by Pembrey in Recent Advances in Physiology and Biochemistry, 1906; 563. Liebermeister found that two and a half times the usual amount of CO_2 was being given off. Leyden, from his observations, laid down that in general the increase is one and a half times the normal. It is this actual increase in CO_2 production and discharge rather than the respiratory quotient that must be taken into account. That quotient may, indeed, be abnormally low.⁴ Herein is the primary cause of increased respiratory rate, and here, also, we have a striking demonstration of what must be regarded as the prominent underlying feature of fever, namely, greatly increased metabolism, that metabolism leading to the increased temperature, to increased discharge of CO_2 into the blood, and to the increased rate of respiration. As with defervescence the temperature falls, the discharge of CO_3 returns to the normal.

Urinary Disturbances.—Here, in general, we find that (1) the amount of urine passed *per diem* is diminished; (2) what is passed is concentrated and high-colored; (3) relatively more urea and nitrogenous constituents (kreatinin and the ammonium salts of organic acids) are passed *per diem*; (4) more potash salts; (5) the chlorides are noticeably deficient, as, again, are the phosphates. The data regarding uric acid are conflicting.

That the amount, *i. e.*, the water, is diminished, is explicable by (1) the lowered blood pressure (and rate of flow); (2) the increased discharge of water by the lungs, the skin, and, in some cases, the feces, The increased pigmentation is another evidence of that destruction of the red corpuscles already referred to; and this explains, also, to a large extent, the increase in potassium salts (for the red corpuscles are relatively rich in potassium). Most characteristic is the increase in nitrogenous constituents. The urea in a continued fever is increased from 70 to 100 per cent., sometimes threefold; the increase in uric acid and kreatinin is more irregular. While the increase is not in all cases nearly parallel to the temperature variations, it is, nevertheless, a very constant phenomenon; and while, again, the amount of nitrogen discharged per diem by a healthy person may, under certain circumstances, exceed that shown by the febrile patient, the amount of urea passed by the latter is much greater than would be passed by a healthy individual on the same diet.

Thus the study of the urine affords one more convincing demonstration that the febrile state is characterized by increased metabolism, increased breaking down of proteins. And these proteins cannot be food and reserve materials; they must be derived from the tissues. It is deserving of note that the evidences of increased metabolic processes and the rise of temperature do not go absolutely hand-in-hand. There may be indications of increased proteid destruction before the temperature begins to rise, and, *per contra*, often after defervescence, especially when critical, the augmented elimination of nitrogenous bodies is found to continue several days after the temperature has returned to normal ("epicritical excretion of urea"). In pneumonia, and possibly in the

¹ Riethus, Arch. f. exp. Pathol., 44:254.

URINARY DISTURBANCES

other conditions in which this is encountered, a possible explanation is the absorption and discharge of inflammatory products.

Regarding the diminution of the chlorides excreted, it cannot be said that as yet we possess a satisfactory explanation. That the diminished consumption of food is responsible for a large proportion of the decrease must be regarded as well established, but this is not everything.1 Large quantities of sodium chloride may be given by the mouth and yet the excretion be deficient. Against this, it might be urged that absorption from the alimentary canal is greatly lessened, but such argument will not explain why, in fevers, the amount of the chlorides in the circulatory blood may not be found diminished. We are thus driven to conclude either that in fever there is an altered selective secretory activity on the part of the glomerular epithelium, or absorptive capacity on the part of the tubular epithelium (if the view be accepted that constituents of fluid filtered through the glomeruli undergo reabsorption), or lastly, with Forster and Sollman,² that excretion or non-excretion of the chlorides is dependent upon the relative amounts free in the plasma or combined there with colloid materials. But, accepting this last view, there is no explanation why, in fever, there should be this increased combination with colloids. Leyden³ has urged that in the febrile state there is a retention of water in the organism; that often the main loss of weight is observed to take place, not during the acute stage, but during convalescence. Such retention of water might be held to explain the retention of sodium chloride. There has been not a little controversy regarding this view. On the whole, the evidence indicates that his observations are correct for febrile states of some duration, such as typhoid, but are not substantiated for fevers of shorter length. As we shall point out in discussing the cell degenerations, active dissociative changes in the cell demand increased intracellular water to hold the increased number of molecules of simpler type in solution. Herein would seem to be the explanation of the retention.

Yet another feature of febrile urine is worthy of note, namely, its toxicity. Although most observers in this field have failed to take into account the toxic actions of the increased potash contents, there can no longer be reasonable doubt that different toxic substances are discharged by the urine in different infections. The urine, in short, affords one means of removal of the toxins from the organism.

Lastly, in every febrile condition we are liable to find albumin in the urine, and, associated with this, we find cloudy swelling and a more or less well-marked grade of parenchymatous nephritis set up, it would seem, in the main by the elimination of toxins and the deleterious products of tissue disintegration. The congestion of the kidneys set up by altered blood pressure would not, that is, appear to afford an

¹ Hatcher and Sollman, Amer. Jour. Physiol., 8: 1903: 117.

² Sollman has contributed a valuable series of papers on this subject, Amer. Jour. Physiol., 8: 1903: 155, 9: 1903; 425 and 454, 13: 1905: 241, 291.

³ Arch. f. klin. Med., 5:366.

THE SYSTEMIC REACTION

adequate explanation for the pronounced parenchymatous nephritis frequently noted.

Other Metabolic Disturbances.—Of these, the most noticeable is the rapid burning up of glycogen. This is found greatly diminished in the liver and muscles of the febrile animal. Although a factor, this consumption of glycogen is not the essential cause of the increased heat production, since that may show itself in starved and glycogen-free animals.

Digestive Disturbances.—Anorexia.—Distaste for food and loss of appetite is characteristic of all fevers in animals, as well as man. With this, as indicated by the mouth (for there is a marked sympathy between all parts of the intestinal canal), there is a marked diminution in the digestive secretion; as indicated also by the stomach by the frequent presence of masses of coagulated case in those on milk diet, who have died from typhoid and other high fevers. In the less severe, continued febrile states, as indicated by tuberculosis, the anorexia and lessened absorptive powers do not necessarily go together; there forced overfeeding leads certainly to increased absorption and positive benefit.

To attempt to explain the anorexia and arrest of intestinal activities is at present to indulge in little beyond pure theory; *i. e.*, to conclude from the frequent discharge of mucus all along the stomach and intestines that the cells are engaged mainly in excretive processes, tending to discharge the toxins, and cannot, therefore, be equally active in absorptive processes; or, again, that it is a means whereby the cells of the tissues in general devote their energies to the elaboration of antitoxins instead of to commonplace assimilation of foodstuffs. Briefly, we know nothing adequate to explain it, and can only note that it is one of the striking features of fever. At most, there are certain indications that bodies of the nature of complements exist in greater amounts in the blood of moderately starved than in that of full-fed animals; but these observations require to be materially extended before they can safely be built upon. The more recent observations by Rankin and Martin' in our laboratory upon opsonins in starvation point in the opposite direction.

Into the changes occurring in the liver and accessory digestive glands we would not here enter at length, beyond stating that these tend to exhibit the same cloudy swelling and parenchymatous inflammation noted as occurring in the kidneys. Like changes may be produced in experimental hyperpyrexia, or by subjecting small animals to greatly increased external warmth, but that the mere increase of temperature is not the essential cause is shown by the fact that pronounced disturbances of like nature may occur in diphtheria and in severe infection characterized by lack of pyrexia. Increased excretion of toxins and products of cell disintegration and physical change in the cells brought about by this excretion must be regarded as the cause.

⁴ Proc. Soc. Exp. Biol. and Med., 4; 1907; 81.

EMACIATION

Cutaneous Disturbances .- Here, as in connection with the loss of bodi'y heat through the respiration, we note a want of evidence of accurate relationship between heat production and heat discharge. The rule of the organism under physiological conditions is, that increased bodily heat is accompanied by dilatation of the cutaneous vessels, favoring more rapid discharge of heat, and, in addition, by increased perspiration, which yet more materially, by the evaporation of the sweat, brings about loss of heat. As regards surface dilatation, while this is well marked in the fastigium, it is noticeably absent in the period of pyretogenesis. As regards perspiration, it is, to say the least, irregular. There is no constant relationship between it and the height of the fever. It is generally pronounced, not when the fever is rising, but when it is going down, in the stage of defervescence; it is most marked accompanying the rapid fall of temperature to the normal, and below (cold sweats), heralding the fatal event in malignant fevers; may occur locally, as in tuberculosis, or may be a feature throughout the disease, as in acute rheumatism. In short, like the respiratory changes, it is not primarily related to the temperature needs of the body; more extended observations than have as yet been made require to be undertaken to determine its relationship to the excretion

Of the exanthematous manifestations, some, like the pocks in smallpox and syphilis, and the roseole of typhoid, are directly infective, due to the presence and proliferation of the specific microbes in and around the cutaneous capillaries; some, particularly those of hemorrhagic type, are merely toxic, caused by circulating toxins. It is possible experimentally by the injection of sundry toxins to induce cutaneous petechie and hemorrhages, indication that those toxins directly affect the walls of the vessels.

Emaciation.—This further cardinal symptom of the continued febrile state may be far more marked than can be explained by lessened intake of food. Notably, there is a reduction of the fatty tissues—a burning up of the fat—and, with this, also of the muscles. Of fat, it must be noted that in its combustion it is capable of giving off more calories of heat than any other constituent of the body. There is in the febrile state, to repeat, no mere retention of heat, but a most evident increased production, and this whether the loss of heat is not proportionately increased, so that there is a rise in the body temperature, or whether it is proportionately increased so that there is no fever in the narrow sense of the term.

ritis the

he

CHAPTER V.

THE SYSTEMIC REACTION-(Continued).

THERMOGENESIS AND PYREXIA.

PASSING in review the data afforded in the last chapter, these conclusions stand out prominent: (1) The febrile state is characterized by an increased metabolism altogether out of proportion to the amount of food and energy producing material taken in, the breaking down processes being greater than the building up; and (2) that with the increased heat production there is not in general a corresponding discharge of heat, so that the body temperature tends to rise. To this a third may be added, that there is evident lack of coordination between heat production and heat discharge. More correctly, this may exist—as is indicated by the regularity of the temperature curve during the fastigium, but at a higher level than the normal, or, in other words, the bodily heat becomes regulated to maintain a higher temperature.

From the respect mingled with awe with which from childhood upward he has seen the thermometer treated, and from the prominence given to the temperature charts in the wards, the student very naturally concludes that the temperature changes are the all-important factor in the febrile state. Let us here state emphatically that this is not the case. The temperature changes are but the expression and the outcome of other underlying conditions, and these it is that, so far as the organism is concerned, are all important. The temperature chart is important to the physician as indicating how these other conditions are telling upon the general state of the organism. While thus we would not dwell too long upon those temperature changes, there are other states besides infection in which there is rise of bodily temperature, and, in order to understand these and their relationship, it is necessary to recall what we know concerning thermogenesis and heat regulation. This knowledge should be familiar; we shall, therefore, state the main data as succinctly as possible.

Heat Production.—Heat is liberated in the organism under these conditions:

1. From the food, *i. e.*, from the recombination of dissociated food-stuffs (p. 101).

2. From tissue katabolism, i. e., from the oxygenation of tissue products. All work performed by the cells leads to dissociation in the cell substances, and it is ultimately the union of these products of dissociation with oxygen that produces heat.

If, therefore, in febrile conditions there be increased production of heat, despite lessened intake of foodstuffs, and despite a loss of heat

HEAT DISCHARGE

that is not less than normal, that increase can only be due to tissue disintegration and oxidation. The same is true of other conditions in which there is increased heat production irrespective of food taken. If, also, through nervous influences increased heat production be brought about, it is not the nervous centres themselves that develop heat; it is the dissociative changes in the cells of sundry tissues set up by nervous stimulation that is the cause.

Heat Discharge.—There may be loss of heat from the body through: 1. Surface radiation and conduction.

2. Surface convection or evaporation-of sweat and in the lungs.

3. The passage from the body of excreta.

The discharge may be *increased* and the temperature of the body lowered:

1. By dilatation of the surface vessels.

2. By increased pouring out of sweat.

at.

a

3. By increased respiration, whereby more air is warmed in passage over the respiratory surfaces and greater evaporation takes place in the lungs. (It is by this increased respiration (panting), in the main, that the dog, unable to perspire, cools himself down.)

4. By increased excretion (a very minor factor).

It may be *diminished* by the opposite conditions—contraction of surface vessels and arrest of perspiration, slower or shallower respiration.

According to Vierordt if the total income of available energy is 2,500,000 calories (a calorie is the amount of heat necessary to raise 1 gram of water 1° C, at the normal atmospheric pressure), then:

1.8 per cent, is lost in the urine and feces	27,500 ca	lories.
3.5 per cent. is lost in the expired air	84,500	44
7.2 per cent. is lost in the evaporation of water from		
the lungs	182,120	
14.5 per cent. is lost in the evaporation of water from		
the skin	364,120	66
73.0 per cent. is lost in the radiation and conduction		
from the skin	1,791,820	4.8

The lower the temperature of the external medium in immediate contact below that of the body, the greater, other things being equal, the loss of heat; the more nearly these approach equality, the less the loss of heat; and if the external temperature exceed that of the body, and at the same time, by saturation with moisture, evaporation is prevented, there is an actual gain of heat. Thus, for example, the body immersed in a warm bath (over 104° F.) shows a very evident rise of temperature.

Despite the fact that at different periods, through muscular exercise and the taking of food, the heat production undergoes great changes, and through alteration in the external medium the heat discharge is similarly liable to vary greatly, the temperature of the warm-blooded keeps remarkably constant, herein differing from the cold-blooded animal, in which the bodily temperature rises and falls with the external temperature. The infant is intermediate, and its temperature is modified

THE SYSTEMIC REACTION

considerably by that of its surroundings.⁴ But in the adult man, whether within the Arctic circle or in the Tropics, the mean temperature (rectal) is maintained in the near vicinity of 37.2° C. (98.96° F.).³ Conditions of great heat, as already indicated, will raise it above this point; of great cold and exposure, if forced exercise and food be not taken, will lower it. In cases of sunstroke the rectal temperature has been found as high as 42.9° C. (109.2° F.), but such temperatures are fatal, and it may be laid down that 42° C. (107.5° F.) is the upper limit of temperature compatible with continued existence.³ (See also p. 285.)

Cases are on record in which, with a rectal temperature of 24° C., those who have been so exposed have recovered, but when the temperature below falls 20° C. death is inevitable. Men thus are not "frozen to death," they die before reaching the frozen state.

 In the conditions of health there is a daily variation of temperature, as might be expected, the minimum being during sleep when the muscles are relaxed and the respiratory change is lowest.⁴

Heat Regulating Mechanism.—The existence of a heat regulating mechanism is thus evident; of a mechanism which within very wide limits is marvellously exact and precise in action. As a matter of fact, we possess indications pointing definitely to the existence and site of portions of such a mechanism, although there is still debate as to the number and mode of action of the same. The very existence of two sets of sensory nerves, one for the sensation of heat, the other for that of cold, in itself indicates the existence of a controlling mechanism. Both by clinical observation from the time of Brodie, in the beginning of the last century, onward, and by direct experiment it has been noted that injury or stimulation of certain areas of the brain or medulla have been followed by marked rise of bodily temperature, of others by lowering of the same.⁴

It is not difficult to understand that laceration or section of the spinal cord, high up, by paralyzing the muscles, leads to lowering of the temperature, as also by paralysis of the vasomotor mechanism. It is this latter that mainly is effective by leading to dilatation of the entaneous vessels, etc., for if in those cases the body be properly swathed the temperature rises to the normal. Cutting between the medulla and pons was found, by Horatio Wood,⁶ to lead occasionally to a considerable rise, which was also noted at times by Heidenhain.

¹ The same, also, is true of warm-blooded animals in the state of hibernation.

² Or in the mouth, 36.87° C. (98.36° F.). These are Pembrey's figures.

³ Much higher temperature than this has been recorded of axillary, anal, and rectal temperatures—even of 150° F, and over—and in those not seriously ill; but in so many of these careful detective work has shown the existence of some trick placing the thermometer in the tea or hot water, pressing, etc.—that imposture must always be diagnosticated in these cases (see Professor Welch, discussion of Dr. A. Jacobi's paper on Hyperthermy, Trans. Assoc. Amer. Phys., 10; 1895; 189).

⁴Vide Pembrey, Jour. of Physiol., 15: 1894: 401.

^b For clinical cases see more particularly Hale White, Guy's Hosp. Rep., 27: 1883–84:48, and Jour. of Physiol., 12:1891:233.

⁶ Fever, Smithsonian Contributions to Knowledge, Washington, No. 357, 1880.

HEAT REGULATING MECHANISM

These results have been opposed by others, but the reputation of both observers for careful, conscientious work is so great that the fact must be accepted. There is more abundant confirmation of the observation of Richet and of Aronsohn and Sachs, that puncture and electrical stimulation of an area in the corpus striatum or puncturing the midbrain with a long needle leads to marked elevation of temperature, beginning several hours after the puncture and lasting for several days, and due not to lessened loss but increased production of heat.

It may be urged that we have no right to speak of these as heatproducing centres;1 that there exist in the brain centres controlling tissues, such as the muscles and the liver, which, in activity, produce heat all must admit, and it is is such visceral centres that are stimulated. To a certain extent we agree with this objection. Heat can only be produced under these conditions by tissue metabolism. Indeed, it has been shown by Hirsch and Rolly² that, following midbrain puncture, the liver is the warmest organ in the body, and other observations3 indicate that, unlike the ordinary febrile state, the increased metabolism in these cases is non-nitrogenous rather than nitrogenous. This, however, does not prevent us from regarding centres which thus may lead to a rapid rise of heat as heat-producing centres. Similarly centres governing the sweat glands and cutaneous vasodilators become heat-discharging centres; and looking at the mechanism in this light we must, with Mac-Alister, predicate the existence of some central heat-controlling centre regulating the (various) heat-producing and heat-discharging apparatus -a centre which stimulates the former and inhibits the latter in order to raise the body temperature, and does the reverse in order to lower it. Without such it is difficult to see how regulation can occur. And we must regard this as being called into action (1) by reflex means; (2) by the temperature of the circulating blood telling directly upon its activity; and (3) by substances diffused in the circulating blood acting upon its constituent nerve cells. Here we trespass into a region of physiology that awaits fuller explanation. All that is sure is that within the brain and spinal cord are nerve cells which on stimulation lead, some of them, to increased production of heat by the tissues, others to increased loss of heat from the body surfaces. The wonderful regulation of the bodily temperature under ordinary conditions is a strong indication that controlling the production and the loss is one pair or an intimately connected

In fever this heat regulating mechanism is gravely disturbed, and the facts we have brought forward are in themselves adequate to prove that the disturbance is in the direction of increased heat production rather than of lessened discharge.

hether rectal) ons of great ver it. igh as ay be ature

4° C., perarozen

sture, iscles

wide fact, e of the two that ism. g of that peen ring inal ratter els, ure nd,

ire

¹This is Mosso's view, Arch. ital. de Biol., 13:1890;459. Reichert, attaches more importance to centres in the spinal cord (Univ. Med. Mag., Phila., 5:1893;406 and 6:1894;303).

² Deutsch. Arch. f. klin. Med., 75; 1905; 307.

³ Rolly, ibid., 78: 1905: 289, and Martin, Arch. f. exp. Pathol., 40: 1898: 453.

THE SYSTEMIC REACTION

Calorimetric Observations.-It must be emphasized that the thermometer can only afford information regarding the balance or resultant at a given moment between the heat production and income on the one hand, and the heat loss and expenditure on the other, of a particular part; it gives no information regarding the actual *amount* of heat that is being developed by the body, or the extent of heat loss. To determine these, observations of a totally different order have to be undertaken, namely, methods of indirect and direct calorimetry. We have, that is, to compare the heat-producing capacity of the ingested food, and of the oxygen absorbed in respiration, during a given period (after deducting the number of unused calories represented in the excreted and discharged matter from the organism during that period) with the contemporaneous amount of heat actually lost by the organism (from the skin, the lungs, and in the excreta), in order to assure ourselves either that there is a relative increase or a decrease of heat production and of heat loss. The methods for determining these data are very elaborate, and many sources of error have to be guarded against. But to such a perfection have the two methods been brought that, in the normal animal, practically identical results are now gained by the two methods. Naturally, it is more difficult to make calorimetric observations upon the patient. It is not surprising, therefore, that there has been not a little divergence on the results obtained. Long and exact studies have been made by capable observers in many countries, by Liebermeister, Rosenthal, and Rubner in Germany, by Lavoisier and D'Arsonval in France, by Ott, Horatio Wood, Reichert, and Atwater in America. The pioneer observer was Crawford, of Edinburgh (1778). Some of the fullest and most exact calorimetric studies upon the febrile states are those of Wood.¹ It may in general be said of acute fevers that (1) during the initial period of fever there is increased production with some diminution of loss; (2) during the fastigium both heat production and heat loss are increased above the normal: (3) a daily temperature variation occurs parallel to that seen in health, but differing in being at a higher level; (4) heat inhibition is not paralyzed, but the individual is not so responsive to those stimuli which in the normal individual stimulate heat loss when heat production rises above the normal level; (5) in defervescence there is lowered heat production with definite increase in loss of heat.² Certain annotations must be affixed to these broad statements: (1) It is found that this increased heat production on the average amounts to from 20 to 30 per cent., and is greatest at

¹ Fever, a Study in Morbid and Normal Pathology, Philadelphia, 1880, and Smithsonian Contributions, *loc. cit.*

² For a general review of the subject of calorimetry and calorimetric methods the reader may be referred to Reichert's article in the American Text-book of Physiology, 1: 1903; 467, and that by Kemp in Buck's Reference Handbook; the latter gives the more important bibliographical references. An excellent treatise, although dealing in the main with the calorimetric values of foodstuffs, is Atwater's Methods and Results of Investigations on the Chemistry and Economy of Food, United States Department of Agriculture, Bulletin 21, 1896.

DEVELOPMENT OF THE FEBRILE TEMPERATURE

ler.

me

xd.

he

the onset of chills and rigors. (2) The increase is not marked in long continued fevers. In some few cases of high fever the rise of temperature would seem to be due wholly to reduced discharge of heat. (3) In certain fevers, notably of the septic type, the rise and fall of temperature is wholly unrelated to the normal curve. Observations upon tuberculous patients strongly suggest that varying rates of diffusion of the bacterial toxins from the foci of bacterial growth play an important part in these irregular accessions of temperature. (4) It is not a matter of partial paralysis of the heat-discharging centre; on the contrary, the cutaneous vessels in fever are seen to be more responsive to temperature changes ihan normal, while cold baths or sponging reduce the temperature more rapidly than in the normal individual. Rather there is stimulation of the (chemical) heat production, which now is not in the normal manner followed by an equivalent heat discharge. There is disturbed equilibrium. What is the essential nature of this disturbance we do not know.

Lastly, before attempting to sum up, it will be well to classify the conditions in which we obtain the febrile state, *i. e.*, a condition of increased bodily temperature with allied increased metabolism. Up to this point we have purely considered the matter in relation to infection; evidently, from what has just been brought forward, the subject is a much wider one.

Causes of the Development of the Febrile Temperature.—1. First and foremost, *injection*, the proliferation within the organism of pathogenic microbes, both bacterial and animal forms. Closely allied to these, and strictly belonging to the infective group of causes, must be included sundry disturbances set up by gross parasites, of which the trichina spiralis affords the best example.


2. The microbes act through their toxins; it is the diffusion of these toxins that produce the symptoms, as can be demonstrated experimentally by the injection of toxins apart from the microbes, when a like train of symptoms is set up. Absorption of toxins from the alimentary canal is, judging from the analogy of cholera and diphteria, a probable cause of the subfebrile state which may accompany constitution.

The researches upon the bodies of the nature of toxins in their relationship to fever extend back over a space of fifty years. Billroth and O. Weber independently observed that the inoculation of putrefactive material, whether of animal or vegetable origin, led to the production of a febrile state, reaching its maximum in from two to twenty-eight hours with preliminary pyretogenic stage and a fastigium, simulating typical infection, but their material obviously was impure, containing infective agents. Panum, from putrefying solutions, extracted a body insoluble in alcohol, but soluble in water, of which 0.012 grains would kill a small dog, producing febrile disturbances and the symptoms of acute infection.

Hermann,¹ whose most thorough work and recognition of the importance of what we now term toxins is generally forgotten, concluded that

⁴ Experimentelle Studien über die Wirkung faulender Staffe auf den thierischen Organismus, 1866, the body he gained, having like properties to that described by Panum was of proteid origin, that it worked as a ferment, and that there were specific differences in the essential toxins of the different infections.

In more recent years it has been noted that the products of growth of bacteria in various media induce febrile states simulating in their peculiar features the fevers produced by the individual pathogenic bacteria. But further extracts of the actual body substances, proteins, as Buchner termed them, or, as is more usual at the present moment, endotxins, are found to cause pyrexia and the febrile state. Roussy first showed this in connection with yeast, Buchner proved it in the case of bacteria. Koch's tuberculin R.—the juice expressed from the bodies of tubercle bacilli under high pressure—causes, when inoculated, a very definite fever, with an incubation period, *i. e.*, several hours elapse before any change is noted, then a sharp rise to a maximum, some twenty-four hours after inoculation is attained, followed by a sharp fall.



Pyrexia following thrombosis in the course of typhoid fever. (T. McCrae.)

3. As Hildebrandt pointed out, many years ago, all enzymes and ferment-like bodies, when inoculated, lead, after a short preliminary period of definite lowering of the temperature, to an even more marked rise of the same. It is to be noted that a like temporary depression is observable after injecting most toxins. With pyocyaneus toxins we have found it well marked, lasting for an hour or two, and followed by a rapid rise. Hildebrandt's observations have been denied by some who urge that where pure ferments are employed no rise occurs. But "pure" ferments in our experience are very largely inactive ferments.

4. Hemorrhagic Products (hemolysis). A large internal hemorrhage, after a like period of depression, is always followed by pyrexia, which may last for some days, and this in the absence of any signs of sepsis. The subcutaneous injection or intravenous inoculation of the blood of another species when not immediately fatal has the same results, as has like inoculation of the blood of another individual of the same species. Something clearly is liberated in the process of coagulation and breaking

CAUSES OF PYREXIA

down which leads to the rise. It has been shown that fibrin ferment will cause these results, from which it would seem that in all the group of cases the common cause is liberation of fibrin ferment. If this be so, then here we have but a special case of the preceding group of causes. This, however, is still under debate.

5. The last case suggests strongly that fibrin ferment alone is not the cause of the pyrexia in these cases, the hemolysis mainly affecting the red corpuscles. As a matter of fact, the sterile extracts or juices of tissues in general produce rise of temperature. This rise is well seen in the administration of thyroid extract. Such diffusion out of tissue juices, or of the contained enzymes of the same of one or other nature, affords the most satisfactory explanation of the sharp rise in temperature in cases of infarct formation and of thrombosis, a rise so characteristic that we are justified in attributing sharp rises occurring in the course of acute endocarditis to the supervention of infarcts and necrosis, in the spleen, kidney, or other organ. Comparing the temperature chart in such cases with the lesions found, we have, on several occasions, been able to convince ourselves of this relationship. Allied to this is the pyrexia which may follow simple fracture of the large bones. In von Volkmann's clinic, of fourteen cases of simple uncomplicated fracture of the femur without sign of septic infection, no less than eleven manifested a pronounced and persistent pyrexia; in five the temperature remained for several days between 39° and 40° C. The local hemorrhage may be the main cause of this rise, or, again, the local tissue laceration and destruction. In short, it may be laid down as the result of experiment, that a large number of proteids, both simple and complicated, when injected induce fever. Krehl and Matthes and others have discussed whether the dissociated albumoses, so often found in the febrile urine, may not be an important factor in the temperature rise, but admit that as they are not constantly present they cannot be a constant factor.

A definite febrile state may follow burns and scalds if at all extensive. The more recent observations of Bardeen, J. McCrae, and others attribute this not to the actual exposure to heat, for it may become noticeable only some hours after the injury, nor, again, to nervous shock, but, judging from the accompanying evidence of selective toxic disturbances, more particularly in the lymph glands and the kidneys, to the absorbed products of cell destruction. This form, therefore, should be included under our fifth heading.

6. Water. Hort⁴ has recently directed renewed attention to the fact that the introduction of sterilized water into the circulation, in relatively small quantities, induces pyrexia. We know that water thus introduced "lakes" the erythrocytes, causing discharge of hemoglobin into the plasma. It may well affect the endothelial and other cells. Presumably, therefore, we have here another example of discharged cell products causing rise of temperature.

¹ Proc. Roy. Soc., June 24, 1910.

vere

ces,

ite.

in

om

cu-

Irs

me

ill.

THE SYSTEMIC REACTION

7. Certain drugs also induce pyrexia. O. Weber first pointed this out in connection with sulphuretted hydrogen. Strychnine working on the muscles through the nerves may definitely raise the temperature. Hale White has shown that β -tetrahydronapthalamin has very pronounced pyretogenic properties.

8. Subjection to greatly increased external temperature. This produces not merely pyrexia, but if that pyrexia be considerable or continued, it is associated with tissue degeneration, cloudy swelling, and parenchymatous degeneration, disturbances closely simulating those which accompany infection; the increased internal temperature leads, that is, to increased metabolism and, it may be, perverted. Here, as throwing light upon-or complicating?-the problem before us, must be noted the fact that rise in the general body temperature in itself stimulates increased heat production, so that fever in itself, beyond a certain grade, is not merely a source of higher fever, but stimulates excessive cell disintegration and degeneration. Years ago Pflüger laid down that each rise of 1° C. in the temperature of the rabbit was followed by no less than 6 per cent. of increase in heat production, and his conclusions have been substantially confirmed by Linser and Schmidt¹ in the human being. In infections, therefore, while bacterial toxins, etc., initiate the increased heat production, the resultant increased bodily temperature is in itself a potent factor in the continuance of the process. Herein is the danger of hyperpyrexia. As we have already emphasized, the heat-regulating centre is not out of action in the ordinary fever, but if through intense primary irritative action of the toxins or deficient discharge the temperature be permitted to rise above a certain point, there is the danger of the establishment of a vicious circle leading to a constantly augmenting temperature up to a point incompatible with life. In heat stroke, which we have discussed elsewhere (p. 289), it would seem that an external physical agency alone may bring about this vicious circle.

9. Nervous Pyrexia.—Lastly, we have those cases of pyrexia produced by nervous influences purely and solely. It has already been noted that this can be produced experimentally (p. 481), and while in any given case it is necessary to be cautious in arriving at a conclusion, while, for example, the run of temperature following a cerebral hemorrhage may be due to diffusion outward of products of the escaped blood, and while the extraordinary hyperthermy of some hysterical patients already noted may be due to some simple trickery, and the rise following upon conculsions may be due to muscular contractions, nevertheless there are undoubted cases of marked pyrexia accompanying brain tumors and other gross lesions of the central nervous system which can, by exclusion, be explained in no other way than by disturbances of the heat-regulating mechanism.

The occasional cases of high temperature supervening upon severe injury to the spinal cord, more especially in the cervical region, would seem to be due more to lack of regulation of the heat loss than to

⁴ Deutsches Arch, f. klin, Med., 79; 1904; 514,

THE SITES OF HEAT-PRODUCTION

increased heat-production. Men, and animals in these conditions, are apt, like cold-blooded animals, to be poikilothermic; raise the external temperature, and so reduce the heat loss, and the body temperature becomes raised above the normal; reduce the external temperature, and the body heat becomes lowered.

re.

Pd.

n-

ht

he

n-

·h

T

Are we justified in classing all these different forms of pyrexia together? Certainly there are differences. In infection alone do we have the absolutely typical example of continued fever, incubation period, pyretogenic stage, fastigium, and defervescence, characterized by what we may express as universal metabolic disturbances. But with the products of bacterial growth-toxins-with enzymes and tissue extracts, we approach very near to the typical condition, the only difference being the absence of that portion of the incubation period dependent upon the proliferation of the microbes until such time as they produce sufficient toxic matter to affect the organism generally, and those differences produced in the continuance of the fever by the continued development of toxins. Yet in cases of gradual diffusion of tissue products and enzymes (internal hemorrhages, burns, etc.), there may be similar continuance, and in all of them we note the existence of a distinct period between the introduction or formation of the disturbing causes, and the development of pyrexia, a period the significance of which is indicated by the observations of Sidney Martin on diphtherial toxins (p. 315) and of Preston Kyes1 on snake-venom lecithin, to which we shall refer shortly (pp. 498 and 540). Toxins, as such, it would seem, gain full toxic powers only after forming certain combinations within the organism, and for this combination time is requisite. This period is wanting in the pyrexia induced by drugs, and in these the pyretogenic period is greatly shortened. In these, also, the evidence of widespread metabolic disturbances are, it may be, of a somewhat different order, even if histologically we can determine no points of essential difference in the processes observed.

Between the nervous pyrexia and the infective and toxic one striking difference has been observed, namely, as shown by Rolly² in the rabbit, whose midbrain has been punctured, the augmented heat production is almost wholly due to the burning up of glycogen and non-nitrogenous material, whereas in ordinary fever there is in addition a disintegration of the cell proteids. Where, as in the starving animal, there is no store of glycogen in the system, then puncture of the midbrain is followed by no pyrexia.

On the Site of Heat-production in Febrile States.—We recognize as a general principle that heat production has its main sources in the muscles and the liver. In Rolly's studies, above noted, where the other indications were that the increased metabolism of nervous pyrexia is non-nitrogenous, the liver was found to be the warmest organ in the body. In ordinary fever, experiments give somewhat divided results. Thus,

¹ Berl. klin, Woch., 1903; 957 and 982.

² Deutsches Arch. f. klin, Med., 78; 1904; 289.

THE SYSTEMIC REACTION

Heidenhain and Körner, in a dog that had been given an injection of pus, found the blood coming from the leg warmer than that in the right ventricle. The general consensus of more recent experiments is that the blood from the liver and even that from the kidney is warmer than that from the extremities. It must, however, be remembered that the blood returning from a limb is mixed, *i. e.*, includes some that has been cooled by passage through the skin. It is thus safe to conclude that in ordinary fever both the muscles and the major glands contribute to the increased heat production.

THE ESSENTIAL SIGNIFICANCE OF THE FEBRILE STATE.

Clearly the infective, enzyme, and tissue extract reactions form a welldefined group; they represent the development of a common process. Now it is noteworthy, as we shall point out in the succeeding chapters upon immunity, that in all of these the reaction results in the production of antibodies, of substances which neutralize the toxic causes. And recognizing this, we appear coincidently to gain insight into the essential nature of fever. Just as we saw that inflammation is the process of adaptation of the tissues to local injury, so is fever the process of adaptation to such toxic agencies as can be neutralized by the development of antibodies.

In further support of this view it is to be noted that if an excessive dose of a toxin be injected—one that is fatal within a few hours instead of a rise of temperature there is a progressive fall, or, at most a transient rise is followed by a rapid fall, which continues until a point far below the normal is reached, when death ensues. The tissues, in short, are poisoned, and there is no adequate general reaction. And the same is to be noted in malignant fever, and in the fatal termination of acute fevers; the fatal event is heralded by arrest of the febrile reaction and falling temperature. To this it is true, there are exceptions. We note cases, both experimental and clinical, in which, on the contrary, death occurs in hyperpyrexia. But studying these, we observe that they are instances in which the toxins have a markedly selective actioneither on the nervous system, as in tetanus (there is a similar selective action on the nervous centres in sunstroke), or the lungs, as in pneumonia, or the heart, as in acute rheumatism-so that evidently in these cases death is due to some one of the vital trinity becoming inactive while the general reaction is proceeding vigorously.

It may be in some cases the very intensity of the reaction defeats itself, and that death is directly caused by the hyperpyrexia. We are, however, inclined to believe that such cases are primarily examples of selective action on the nerve centres; that, as indicated, in fever in general the heat regulating centres are still in action, although working at a higher level, and that in these particular cases of death in hyperpyrexia it is through the complete inhibition or intense stimulation—of the centres that the hyperpyrexia is brought about.

RELATIONSHIP OF THE NERVOUS SYSTEM TO PYREXIA 489

Regarding the drug pyrexias, it will be interesting to determine—for such has not yet been done—whether as a body chemical substances which induce pyrexia lead, upon inoculation or absorption, to the development of a similar class of antibodies. In one member of the group, namely, strychnine, as Professor Meltzer has pointed out to me, the fact that there is no acquired tolerance for the drug is against such assumption.

tht

he

10

THE RELATIONSHIP OF THE NERVOUS SYSTEM TO PYREXIA AND OTHER PROCESSES.

And finally, with nervous pyrexia and hyperpyrexia, here again we have the close parallelism with what we determined in connection with inflammation. Just as there we noted that without actual local injury the nerve centres could independently originate the succession of processes which ordinarily require to be initiated by actual injury, so here we have to acknowledge that, without the presence of what we may broadly refer to as "toxins," the higher nervous centres can set going the series of general changes throughout the organism, which, with the nervous system in normal state, are only called into being by the presence of such toxins. Just as in neurotic inflammation we saw that the complete succession of manifestations was wanting, so here, while there is induced by the nervous centres increased metabolism and heat production, with its associated pyrexia, there are of necessity wanting the tissue changes leading to antitoxin formation (although what these are we do not surely know) and the actual production of antibodies.

Dangerous, as from the essential unlikeness of distinct phenomena all similes must be, occasionally we encounter one that is of more use than many paragraphs, and that because it impresses forcibly some important point, or basal relationship. There has been a long and sometimes angry debate regarding the relative importance and activity of the nervous system and the tissue cells, respectively, in the performance of function, whether under physiological or pathological conditions. Some would see every little act of every individual cell under normal conditions initiated and governed by impulses proceeding from the nervous centres. For them nervous matter is the fountain and origin of all bodily activi-Others see the cells as largely independent, stimulated in the main ties. by alterations in their environment. For these the difficulty is to explain and harmonize cell activities undoubtedly set up by nervous stimulation; they would deny that processes of direct and nervous origin can in their sequence and results be identical. The parable of the coach, the coachman, and the horses places the interrelationship in a right light. Those horses have been foaled, have learned to eat and to run and to perform the natural functions without any necessary supervision by the coachman, but when grown up he has had to train them to run together in harness and pull the coach. And, well trained, so accustomed have they become to the daily round that, without direction, they will fall into their appointed

THE SYSTEMIC REACTION

places, will go cautiously down the decline, or work harder, pulling the coach up hill. Undoubtedly, the constant tension of the reins upon the bit and the variations in the same, guide and direct them. So also the whip will suddenly stimulate them to increased exertion, and that without any necessary knowledge on their part of why such call has been made upon them. The coachman may be excited and flog them to a gallop downhill, where they have been trained to go cautiously, and they gallop; or he may be drunk, and lead them off the road, and bound they are to go where he guides. But such is their training, that if, being on the road with the coach running behind, he becomes incapacitated or falls off. they will keep to the road, will even manage the curves, will strain up hill, and proceed with caution down; nay, even will draw up at the accustomed halting place. Only their pace may be uncertain, and if some unaccustomed obstacle presents itself before them, then, lacking guidance, they may come to a standstill; or, seized with panic and each acting for itself, may break the pole and tangle the harness; or, dragging the coach hither and thither off the road, may bring it and themselves to destruction.1

The application of this parable to these states of nervous hyperpyrexia and their relationship to fever proper is obvious. Kept in mind, it time and again is of help in harmonizing apparently antagonistic data bearing upon the functional activity of the different tissues in relationship to the nervous system.

⁴I find that I have been forestalled in this simile by Wenkebach (*Die Arrhythmie*, 2:1903;3) and very possibly by others. Wenkebach compares the regularly beating heart to a horse trotting.

CHAPTER VI.

he he hde

IMMUNIZATION AND IMMUNITY.

WE have laid it down in the preceding chapter that in the course of infectious fevers and those set up by certain organic substances there are developed within the system certain "antibodies," by means of which toxins become neutralized, so that the system becomes protected, and, with this the febrile disturbance comes to an end. It is now in place to inquire more closely into these processes of immunization, and into the steps whereby these antibodies are formed and immunity is acquired. Already we have regarded the subject from one aspect. We have seen that continued existence means continued adaptation, that an environment suitable for individuals of one species may be fatal for those of another, and that, consequently, if all species are primarily of common origin, then during evolution the ancestors of the different existing species, subjected to different environments, have undergone adaptations in different directions, have become modified, and, indeed, immune to influences which, without such modification, would have brought about cessation of activity and death. Speaking broadly, the individual must be regarded as having gained through inheritance a relative immunity, within certain limits, in respect to the action of all the agencies, physical, chemical, and organized, which constitute its normal environment. This statement is, perhaps, too extreme; there are agencies with which living matter shows no interaction, in respect to which it is absolutely inert; it makes no combination with certain chemical substances, and is uninfluenced, for example, by alternating electric currents beyond a certain range of frequency; is not stimulated, likewise, by certain sound waves, or by light waves of more than a certain amplitude. As regards such agencies, we must recognize (1) an absolute immunity. For all the other agencies, however-and their number is practically infinite-which are capable of affecting the molecular arrangement of living matter we must recognize (2) a relative immunity-that living matter has gained the capacity to withstand the action of such up to a certain limit without being destroyed. We must recognize, further, as above implied, (3) that this relationship is *quantitative*, that there are limits beyond which the action of an agency becomes detrimental; and next, that as regards any particular agent, not only the different species, but the different individuals of the same species, whether through inheritance or acquirement, exhibit different grades of reaction. We must recognize, that is, (4) the existence of ex-specie and individual suscep*tibility* or predisposition.

IMMUNIZATION AND IMMUNITY

With reference to the mechanism whereby inherited immunity is brought about, little can here be said; nothing, in fact, beyond what was stated in our discussion of inheritance in general (p. 162), and, with regard to acquired immunity, many acquirements are so gradual that our study becomes limited. Experimentally, that is, we can but inquire into those cases in which the development of this relative immunity is rapid, occurring within a few days or weeks; or, to be correct, it is these cases alone that have so far been studied, and that form the basis of our knowledge. Thus, to all intents and purposes, we are confined to the process of immunization with which is associated the development of fever, to a study of immunity produced through infection and toward toxins, enzymes, and tissue extracts.

Such has been the outburst of work during the last twenty years upon this subject, so great the accumulation of data of a wholly new order, so diverse the views of individual workers, and so appalling the array of new terms introduced, that, when in addition the solution is still unsettled, the unessential matter not having yet undergone precipitation, there is no branch of pathology more difficult to teach or that offers greater difficulties to the student to master. We shall have to touch upon many problems yet unsolved, and to note data which are apparently contradictory, along with others that must be regarded as established and of the very highest importance.

In these circumstances, the only satisfactory introduction to the subject is the historical, whereby, first and foremost, a proper perspective is gained of the successive steps in advance in our knowledge of the subject.

That one attack of many of the infectious diseases protects the individual against subsequent attacks, or renders those subsequent attacks mild and harmless, has been known through the ages; nay, more, for centuries in India and the East advantage has been taken of this fact, and, to protect them against the severe disease, individuals-chiefly children-have been purposely inoculated with matter taken from those suffering from a particular form of infection, or (and this not only in the East) have been made to sleep in the same bed, etc., it having been found that disease so communicated to those enjoying good health is apt to assume a mild form. In other words, infectious disease, under ordinary conditions, "selects" those of weakly constitution, or temporarily in poor condition, and gains a stronger hold than it does when conveyed to those previously sound in health. Such a method of inoculating smallpox, then endemic throughout Europe, and terribly rife, was introduced into England from Constantinople by Lady Mary Wortley Montagu, and became extensively practised there and elsewhere in Europe and North America in the middle of the eighteenth century, until the danger of the process became so evident that it had to be given up. The very mildness of the inoculated disease led to carelessness on the part of the patient and those around him, and, while the patient became immunized, his *entourage* became infected; the disease became more and more common as a consequence.

HISTORICAL INTRODUCTION

The next step, and this definitely in advance, must always be associated with the name of Edward Jenner; not that he was the first to note that cowpox conveyed to man protected against smallpox, or, indeed, to inoculate with cowpox to this end. One or two others had practised the method sporadically before him, but without fully testing the results or making them generally known. Jenner went into the matter as thoroughly as it was possible for him to do at that period, tested with smallpox virus those whom he had "vaccinated" (vacca, a cow), and found them resistant; and by the publication of his methods and results before the Royal Society, in 1796, brought the matter to public knowledge and public use, and thereby converted what had been the greatest endemic scourge into a relatively rare disease, capable of control. Thanks to the process of immunization properly carried out, smallpox is today non-existent in the German Empire, save for cases introduced over the borders, and affects other countries in inverse proportion to the rigor of their vaccination laws and the stringency with which these are enforced. The essence of Jenner's advance was that, conveying a mild disease, harmless, noninfective, save by direct contact, he protected against an often fatal and always disfiguring disease. But how these results were brought about he could not explain; he held that he was dealing with a modified form of smallpox, and in this we now know that he was right. In modern terms, he employed a germ attenuated by passage through another species of animal to set up a mild attack. One hundred and fourteen years have elapsed, and even today we are not positive what is the organism of smallpox.

With the next and greater epoch the name of Pasteur¹ must ever be associated. He it was who, first isolating the causative agent of an infection, found the means of attenuating it (1880), of making it so weak that, when inoculated, it set up a transient illness, after which the animal was found protected against the natural infection. The chickens around Paris were being decimated by a virulent diarrhœa (chicken cholera); Pasteur isolated the organism, a minute bacillus, and noted that old cultures did not kill the chickens inoculated; utilizing these chickens for subsequent inoculations with powerful fresh virus, he found that they did not succumb; and he possessed the genius to recognize the full significance of the observations; not that something had gone wrong with his experiments, but that by keeping, the culture had become attenuated, and that here he had at hand a means of conferring immunity by inoculation the chickens with attenuated virus.

That a plague of diarrhœa in a poultry yard, studied by a professor of chemistry, should be the seed from which has grown the vast development of latter years is a strange fact, but fact, nevertheless. Armed with the knowledge gained from the study of chicken cholera, Pasteur and his lieutenants, Roux and Chamberland, attacked next the subject of anthrax, a disease enzootic in certain parts of France, and causing a

¹ Compt. rend. de l'Acad. de Science, 90: 1880: 239.

IMMUNIZATION AND IMMUNITY

great annual loss to the farmers. The simple means of attenuation employed in the first case was here of no use; old cultures were as virulent as were new, and this, it was found, because the bacilli were spore-producing, and the spores formed during the early days of growth, and remaining dormant in the culture fluid, when they developed possessed all the properties of the bacilli in which they had been formed. Some method had to be found to lower their virulence. Toussaint, in 1880, had already published a method of attenuating the bacilli, namely, by heating the blood of an infected animal to 55° C. for a few minutes. The bacilli in such blood, we now know, contain no spores. Chauveau, later, gained similar results by taking fresh cultures of the bacilli and heating them momentarily to 80° C. Both these methods undoubtedly attenuated the bacilli; either continued for a little time would cause the death of the organism. A variation of a few seconds in the treatment would thus make a very material difference in the grade of attenuation; in fact, by neither could exactitude be obtained. To obtain practical results, exactitude in dosage was essential, and his training as a chemist led Pasteur to seek after exact reactions. Thus, after many trials, a way was found, namely: it was discovered that spores were not produced when the bacilli were cultivated close to the maximum temperature limit of growth. Grown at between 42° C. and 43° C., the bacilli became slowly attenuated, until, in thirty days, growth ceased altogether, through the weakening and death of the bacilli; between the eighth and the thirtieth day the loss of virulence was progressive, and, what is more, subcultures made from the original flasks of growth in the high temperature incubator, when kept at 37° C., did not regain their original virulence, but maintained for generations the grade of attenuation impressed upon the original culture by growth for a certain number of days at the high temperature.

Here, then, he possessed a method of accurate graduation, and now he was able to demonstrate that progressive inoculation of sheep and cattle, employing in each case first a weakened culture, then, some eight days later, when the reaction had subsided, inoculating with a more powerful one, gave immunity against large injections of the most powerful virus. His method was so precise that it was found of immediate benefit, and with its general adoption the reduction of the anthrax mortality in France was very remarkable. As a practical indication of its value (for at the time Koch vigorously criticised Pasteur's method, and even today many German text-books do not, in our opinion, appraise at its full value this great achievement), it may be stated that for some years French insurance companies refused to insure farms in the infected districts unless the sheep and cattle had been "vaccinated," and even supplied their own veterinarians, to make sure that the treatment was properly carried out.

Still more remarkable was the genius displayed in dealing with rables—a condition in which, for the first time, Pasteur came to deal with disease in man. Even at the present moment we are doubtful as

to the causative agent, and Pasteur was literally working in the dark; notwithstanding, he succeeded in gaining virus of definite grades of intensity with which to inoculate. First, it was determined that the virus was constantly present in the brain and cord of dogs affected with the disease; that an emulsion of this nerve matter inoculated into the animals of the laboratory would set up the disease in them, and that the most rapid and sure method of inoculating the disease was by the subdural method, by trephining the animal and introducing the matter beneath the dura mater. But with matter obtained thus from different rabid animals, the period of inoculation in the inoculated animals was very variable, and often very long; it was necessary to make it constant and, if possible, to shorten it. Now, Pasteur discovered that he could intensify the virus. He had, two years before, made certain remarkable observations upon "rouget du porc" (swine erysipelas), finding that by passage of a germ through a series of pigeons he increased its virulence for pigeons, through rabbits for rabbits, but that, whereas the virus rendered more virulent for pigeons was also more virulent for swine, with the rabbit virus the results were the contrary. Just as smallpox from man passed through a series of calves or monkeys1 is rendered less virulent for man, so with the rabbit "rouget" for swine. This was the first case in which this was definitely proved. Since then several like instances of exaltation or attenuation of virulence by passage have been reported. He found that he could intensify his rabies virus by passage through rabbits until, after some two hundred passages, he had brought down the incubation period with remarkable regularity to six days; further passages reduced it farther, but slightly and very slowly. He now possessed a "virus fixe," or sufficiently constant for purposes of obtaining attenuated material of definite grades of lessened virulence. We have said that the causative agent is unknown. Pasteur took the rabbit's cord, under strict aseptic precautions, and, falling back upon the chicken cholera experience, tested the effects of exposing it to the air for several days and drying it over caustic potash. As a matter of fact, he found that by this means it eventually lost all its virulence, and-not to enter into details-he was able to elaborate the method of developing immunity in man and protection against the disease by daily inoculations of emulsions of such dried cord, beginning with that dried twelve days, and devoid of all virulence, and gradually ascending until the most virulent material was injected. The remarkable success of this method in a disease which in man has so long an incubation period that preventive inoculations can be practised during that period, is known to all.

We have described these observations in a little detail, not to remind the reader of the greatness of Pasteur's genius, though that is worth the doing, nor, again, to impress the moral of the need in studies upon immunity to gain material of known and constant strength, though that is most important, but more particularly to impress the fact that

¹Vide Copeman, Jour. of Path., 2:1894:407.

IMMUNIZATION AND IMMUNITY

each specific disease is a special entity, having characters of its own that have to be taken into consideration and dealt with along special lines.

The next great step forward was the determination by Salmon and Theobald Smith, in Washington, in 1886, that immunity against hog cholera is to be gained by the inoculation of products of growth of specific organisms-observations which during the next few years were abundantly confirmed in connection with a large number of pathogenic organisms (tetanus by Brieger and Kitasato; diphtheria, by Roux and Yersin; Bacillus pyocyaneus, by Chantemesse and Charrin; and the list might be extended). It was thus definitely established that (1) the symptoms of infectious disease are caused by the diffusible products of bacterial activity, and (2) that immunity is more particularly the development of the capacity to neutralize those products. But it soon became obvious that here, again, the different pathogenic bacteria did not all possess like properties. With one group the filtered culture fluids were eminently active; with another group they were, if not wholly inactive in producing immunity, so weak as to be practically of no value, although immunity in this latter group could be in duced by inoculating minute doses of the living culture. We owe more especially to Pfeiffer,¹ in 1891, and to his studies upon the cholera spirillum, the recognition wherein the difference lies. In the latter cases immunity can also be induced by inoculating the killed bacilli. Whatever processes happen within the organism (and we are still uncertain regarding these), in cultures outside the body the toxins are present but do not diffuse out of the bacteria into the culture. e now speak of these as endotoxins (p. 497).

As to the chemical constitution of these toxins the observations so far have led us, as already noted, to no sure conclusions. For this reason the recognition by Ehrlich,² that there exist vegetable toxins (*phytotoxins*) that can be isolated in a state of relative purity has been of distinct importance. Abrin (from *Abrus precutorius*, the jequirity bean), ricin (from the eastor oil plant), robin, and crotin have all been found to possess toxic properties resembling those of toxins proper; and, what is more, as Ehrlich showed, it is possible to immunize against them. Another class of albuminoid bodies, the snake venoms, were shown by Calmette, in the same vear, 1894, to belong to this category.

Coincidently, while one series of observers were investigating the properties of the offensive toxins, another set was giving their attention more to the defensive mechanisms of the animal body. Traube, in 1874, and Lister,³ in 1881, noted that blood to which putrefying matter was added, had, within certain limits, the power of remaining sweet, and deduced that it had definite bactericidal properties. Metchnikoff, from 1884 onward, demonstrated with wonderful ingenuity the powers of the leukocytes to take up and destroy bacteria, and, what is more, to

¹ Zeitsch, f. Hygiene, 11: 1892; 393.

² Deutsch. med. Woch., 17:1891:976.

³ Trans. Inter. Med. Cong., London, 1:1881.

HISTORICAL INTRODUCTION

become adapted to and destroy bacteria from which, at first, they were repelled. Nuttall,⁴ in Flugge's laboratory, was the first to demonstrate under the microscope and by cultural methods the destruction of bacilli by the fluids of the organism. The solution of bacteria, whether within the cells or in the fluids of the body, must clearly be a chemical process, and if the bacteria produce specific chemical substances, it would seem certain that the organism provides particular chemical substances to counteract them. To Hankin, of Cambridge, whose brilliant work has been arrested by routine departmental work in India, belongs the credit of first isolating (1888) defensive bodies from the tissues, and showing that these neutralize the toxins. Independently, Buchner, a little later, made like observations, terming the bodies he isolated *alexines*. These two laid the foundation of our knowledge of the *antibodics*.

More immediately, however, in the direct line of advance during the next few years were the observations of Richet and Héricourt,2 and of Babes and Lepp,³ that the blood serum of animals immunized against pyococci and rabies, respectively, conferred immunity on other animals. These observations led up to the great work of Behring and Kitasato⁴ (1890) upon tetanus and diphtheria, work which not merely showed that immunity could thus be conferred by the serum of immunized animals, but that cure could be attained by the inoculation of such serum into those already affected-in tetanus, if the inoculations be made during the incubation period; in diphtheria, after the disease has definitely showed itself. With the development of the practical methods of employing diphtheria antitoxin Roux's name must always be asso-These observations showed the existence of another form of immunity, termed by Ehrlich passive immunity, not brought about by the reacti n of the infected animal, but due to introduced antitoxic substances, in contradistinction to the *active immunity* induced by such reaction.

Thus, in 1890-91, these men established the fundamental data of immunity. Since then there have been great advances in detail, in confirmation of the results here indicated, in further analysis and determination of the nature and mode of action of the bodies which we have grouped under the general terms of toxin and antitoxin. Of these, the greatest was first indicated by Pfeiffer's work upon the cholera spirillum, but its meaning was only comprehended through Bordet's able work upon the process of hemolysis⁶ and Ehrlich's⁶ expansion of his "sidechain theory" to cover "cytolytic" phenomena in general. We refer to the recognition of immunity being dependent, in most cases, not upon

¹ Nuttall, Zeit. f. Hyg., 4: 1888; 253.

² Compt. rend. de l'Acad. de Sci., 107:1888.

² Ann. de l'Inst. Pasteur, 3: 1889.

⁴ Deutsch. med. Woch., 1890:49.

⁵ Ann. de l'Inst. Pasteur, 12: 1898.

⁶ Wertbestimmung des Diphtherieheilserums, Jena, 1897.

IMMUNIZATION AND IMMUNITY

one, but upon two bodies, the one, "complement," present in the normal organism; the other, the "specific immune body," developed in reaction to the presence of the toxins. Of, it may be, equal importance, is the discovery of Preston Kyes,¹ that a body of approximately known character and composition, lecithin, can take the place of the complement (for thereby we come nearer to an exact chemical knowledge of the process of immunity), and, for the same reason, the earlier demonstration by Martin and Cherry,² confirmed by Brodie, that toxin and antitoxin combine to form a chemical compound, stands out as one of the landmarks of more recent advance.

At a relatively early period Hildebrandt showed that immunity could be obtained against the action of various enzymes by progressive inoculation of the same. Ehrlich discovered the production of antitoxins against the plant poisons (phytotoxins); Phisalix and Bertrand, Calmette, and Fraser, independently, antitoxins against snake venom; Ehrlich and Bordet, the formation of cytolysins, *i. e.*, the development in the organism of substances which protected against and destroyed the cells of other species or of other individuals of the same species, these being developed as the result of progressive inoculation of the particular order of cells.

The widespread interest taken in these investigations, opening up, as they have done, a new world, and the eager participation in the researches in all parts of the civilized world, is sufficiently indicated by the names we have mentioned in this rapid sketch. French and German, and Russian, English and American, Japanese and Australian, each and all are to be credited with one or other notable advance. And as the discovery of the new world, while it rounded geographical science, doubled the data with which the student of geography had to become familiar, so these discoveries of pathogenic bacteria, the nature of their action, and of the reaction to them, while they have wonderfully rounded our whole conception of the processes of disease in general, have undoubtedly doubled the data the medical student of today must master, as compared with the student of a quarter of a century ago.

¹ Berl, klin, Woch., 1903: loc. cit.

² Proc. Roy. Soc., 63: 1898; 423, and Brit. Med. Jour., 1898; ii; 1120.

CHAPTER VII.

IMMUNIZATION AND IMMUNITY-(CONTINUED).

THE VARIOUS ORDERS OF IMMUNITY.

HAVING afforded this short account of the development of the study of immunity, it is unnecessary to continue in the strict chronological order; to discuss first what we know concerning toxins and antitoxins, and the development of immunity to infectious disorders. On the contrary, it will, we think, be found more helpful to take into consideration, first, the data bearing upon what may be termed non-specific immunity; next, the simpler cases of immunity to bodies of known constitution; next, those of specific immunity, in which, if still we are not fully acquainted with the chemical composition of the agent against which immunity is obtained, we, nevertheless, are able to isolate that agent, and can be assured that we are dealing with substances of constant value. Having done this, we can with greater security, consider the data associated with the more complicated cases, in which we deal with the reaction to bodies which so far we have been unable to isolate in a state of purity.

Non-specific Immunity.—It is inevitable that the researches upon immunity have been almost wholly confined to the study of the specific antibodies, their development, and properties; in other words, selecting one or other agent, or antigen, the inevitable investigation has centred itself upon the particular antibody or antibodies developed in the organism whereby that particular antigen is counteracted and immunity is acquired. As a result the existence of what may be termed non-specific immunity has been very largely neglected. Nevertheless it is essential that this be continually kept in mind. Unless this be done it is impossible to comprehend the phenomena of the infective process and the foundation is wanting upon which to build up our conception of the nature of specific immunity.

Reference has already been made to some of the more important observations bearing upon this non-specific immunity in the discussion upon the normal defences of the organism, (pp. 319 et seq.). It has been shown that even definitely pathogenic bacteria are indifferently destroyed by the tissues if the number gaining entrance at one spot is below a certain minimum: that if pathogenic bacteria be introduced into the blood stream, although millions, not to say billions, be so introduced within an hour the blood may afford no cultures: that these bacteria are taken up by the endothelial cells in the liver, spleen, etc., and that examination of these organs shows that the bacteria undergo active phagocytic destruction; although, if the microbes be highly pathogenic or their

IMMUNIZATION AND IMMUNITY

numbers be such as to exhaust the bactericidal mechanism, some of the bacteria instead of being destroyed, themselves lead to the death of the cells that have ingested them, undergo proliferation and again enter the blood, so setting up a secondary bacteriemia. We recall these data because their relationship to the problem of immunity may not have been realized.

Or let the matter be regarded from another aspect. Take for example a case of malignant endocarditis. That condition in itself, we are now convinced, is the outcome of a bacteriemia. The streptococci or other causative agents have in the first place to be present in the blood in order to be arrested by the endothelial cells covering the heart valves, to proliferate within and destroy those cells and set up a focus of localized inflammation. Once that focus is set up there is repeated discharge of the cocci or other microbes from the inflamed surface, and these microbes are carried by the blood stream to all parts of the body. They must, for example, find their way in abundance into the minute capillaries of the muscles and the brain, and if in this disease spleen and kidneys are the seats of election for infarcts, if, that is, gross vegetations from the affected valves are swept into these organs, a fortiori individual bacteria and groups of bacteria must find their way into the capillaries of these organs. And yet how rarely in a case of malignant streptococcus endocarditis do we encounter metastatic foci and miliary abscesses in the muscles, the brain, the spleen, or kidney!

Our only conclusion can be that in an infection which passes on to a bacteriemia while there are foci in which the specific pathogenic bacteria find favorable conditions for growth, the site of these foci varying with the different pathogenic bacteria (see p. 408), coincidently the main mass of the tissues of the organism—or perhaps more correctly the vessels supplying the main mass of the tissues—being possessed of nonspecific bactericidal properties are capable of destroying the bacteria. This is an indifferent or non-specific immunity. We can demonstrate its existence before time has been given for the production of specific bodies: we can demonstrate it in connection with microbes like the exotic cholera spirillum against which the animal of the laboratory cannot possibly have acquired a previous immunity.

At this juncture it must suffice to state (1) that in this indifferent immunity we have the first general reaction on the part of the organism to infection: (2) that herein we have the first participation on the part of the tissues away from the site of infection in the reactive process, such participation being all important in the arrest of infection (p. 321), and (3), that the existence of this indifferent immunity is the starting point from which specific immunity becomes developed.

Possibly as throwing light upon this non-specific immunity and as the first steps in the development of specific immunity there should be noted certain recent and striking observations of Abderhalden and his pupils.¹ They show that if there be injected subcutaneously into dogs

¹ Numerous papers in Zeitschr. f. Physiol. Chemie 63: 1909 and 65: 1910.

IMMUNITY AGAINST SUBSTANCES OF KNOWN CONSTITUTION 501

solutions of various proteins, within twenty-four hours the blood contains enzymes of tryptic type which peptonise and split up not merely the particular protein injected but various other proteins. The same occurs when carbohydrates are subcutaneously injected. Only now there appears in the blood an indifferent glycolytic ferment acting both on sugars and starches. The reaction is specific to the extent that it is a group reaction, non-specific in that it is not narrowly confined to the particular protein or carbohydrate employed for injection.

Participation of Leukocytes in Non-specific Immunity.- It is interesting to note that contrary to what happens in the removed blood serum, within the vessels when bacteria are introduced there is relatively little phagocytosis. The case is different when bacteria and leukocytes are extravascular and here we have abundant evidence of the capacity of the leukocytes to take up indifferently pathogenic bacteria. Issaeff¹ was the first to call active attention to this phenomenon. A peritoneal leukocytosis may be induced by various non-toxic agencies; Issaeff employed intra-abdominal injections of saline solutions, serums, etc., and then twenty-four hours later when the leukocytosis was at its height gave what in the untreated animal would be lethal doses of sundry pathogenic spirilla. As a result either no ill results ensued or the course of the disease was materially prolonged. Durham² confirmed, showing that the leukocytosis set up lasted some four or five days, and that during this period there was increased resistance not only to spirilla but also to B. typhosus and other microbes. It is clearly non-specific, disappearing within a fortnight. These observations have been given practical application. Utilizing this "Issaeff resistance period" von Mikulicz and other surgeons have made intraperitoneal inoculations of nucleins, saline solutions, serum, etc., the day before undertaking any serious abdominal operation, and thereby they have reduced the damages of subsequent infection. The leucocytes present in abundance in the abdominal cavity take up and destroy any bacteria before they have time to multiply.

Immunity against Substances of Known Constitution.—Beyond the fact that there exists such an immunity, the data so far gained from the study of immunization lead us, unfortunately, a very short distance. It is a matter of familiar knowledge that there exist arsenic eaters in Styria who can accustom themselves in the course of a few years to the consumption of four times the ordinary fatal dose with no effect beyond a sensation of general well-being. If this be the case in man, and Hausmann^a has called it in doubt, no such definite immunity can be conferred upon the animals of the laboratory. Most observers have obtained nothing beyond merely negative results; have found rabbits and dogs to become increasingly susceptible. Even Besredka,⁴ though

- ³ Arch. f. Pharmakodynamie, 2: 1903, and Deutsch. med. Woch., 1903.
- ⁴ Ann. de l'Inst. Pasteur, 13: 1899; 49, 209, and 465.

¹ Zeitschr. f. Hygiene, 16:1894:287.

² Journ. of Path. 4:1897:370.

he discovered the cause of this ill success, can only be said to have induced a triffing immunity. Nevertheless, his observations are of great interest.

The normal peritoneal fluid contains farly abundant leukocytes, in the proportion of 66 per cent. of mononuclears to 33 per cent. of polymorphonuclears (in the rabbit). Inject an arsenical salt into the peritoneal cavity, whether in solution or in suspension, and if the dose be fatal, there is an almost immediate *hypoleukocytosis*, the diminution especially affecting the polynuclears. Employing Metchnikoff's terminology, there is a negative chemiotaxis. If the dose kills the rabbit in less than twenty-four hours, the reduction continues and becomes more and more marked. If the dose be below the fatal, there is, at most, a transitory hypoleukocytosis, followed by a hyperleukocytosis and increase, particularly in the number of polynuclears, so that the peritoneal fluid may become quite milky. This leukocytosis, first seen in the peritoneal fluid, subsequently affects the blood also.

If a sparingly soluble salt be exhibited, while in fatal cases it is untouched and found lying free on the peritoneal surface, in other cases it is taken up by the "polymorphs" and hyaline mononuclear leukocytes, and can be seen within them in the form of fine colored granules. If soluble salts be employed, the same process evidently occurs, for, if the blood be drawn with due precautions against clotting and be rapidly centrifugalized, the serum and red corpuscles afford no signs of the presence of arsenic; the layer of white corpuscles afford no signs of the presence.⁴ The nearer the dose is to the fatal limit the larger and more pronounced the stage of hyperleukocytosis, and where a sparingly soluble suspension had been given, Besredka found that in the course of days the foreign granules in the leukocytes became finer, until at last none was to be recognized.

That there are also fixed phagocytes was further obvious; the liver in all cases contained arsenic in excess of other organs. Employing sparingly soluble salts, it would seem that the leukocytes break down or otherwise discharge their contents, which are taken up by the liver cells by absorption, from which they are slowly discharged into the bile. The poison is also slowly passed out from the kidneys. It appears to produce its fatal effects by action upon the nervous tissue, for one one-hundredth of the dose which produces death when inoculated under the skin leads thereto when injected directly into the brain, and, injected thus, sets up all the symptoms of acute arsenical poisoning.

These are the basal facts. Now, as to the production of immunity. Besredka took advantage of the stage of hyperleukocytosis. With a solution of arsenical salts of such a strength that 10 c.e. surely kills a rabbit of given weight in forty-eight hours, he found that, injecting

¹ This is an observation of some importance; it has been freely taken for granted that leukocytes absorb soluble substances, like all other cells, and possess the power of selective absorption. This, to our knowledge, is the first positive demonstration of that selective absorption.

IMMUNITY AGAINST SUBSTANCES OF KNOWN CONSTITUTION 503

2 c.c. of this under the skin at night, and then giving the fatal dose (10 c.c.) subcutaneously the next morning, no ill results ensued. Arsenic is very slowly eliminated; the animal had in its system more than the fatal dose; 12 c.c. usually caused death in twenty-four hours or less, and yet animals so treated recovered. Examining into the matter, it was found that the preliminary dose had caused a pronounced polymorphonuclear hyperleukocytosis. The consequence is that, absorbing the poison, these leukocytes remove it from the blood and body fluids, and so prevent it from reaching the nervous system in sufficient concentration to set up fatal effect.

This, so far, is not immunity of the classic order; it is in line with the phenomena of Issaeff's "resistance period" (p. 501). At most, a grade of positive protection is given.

But there is more to be said. If immunity of this order be produced in a rabbit, and the animal be bled six or eight days later and the serum taken, that blood serum is found to have acquired new properties. Whereas the serum of an ordinary rabbit is absolutely without effect when injected into an animal, which has received a fatal dose, 8 c.c. of the serum of a treated rabbit injected into a fresh animal either at the same time as, or antecedent to, a minimal fatal dose of arsenic, acts as a preventive, and the animal recovers from the poisoning.

²This latter is what we term *passive immunity*; a substance not elaborated by the animal under experiment, when introduced into its organism, acts as an antidote—aids the system in neutralizing or destroying the poison.

It is obvious that the serum of a rabbit that is actively immunized against arsenic comes to contain something not present in recognizable amount in the serum of the normal rabbit.

What is this something? That we do not know. We only know that it is not arsenic-containing, is not a combination of the arsenic given to the first animal with its cell substance; for if the serum be obtained with all due care, despite the fact that the methods at our disposal indicate the presence of extraordinarily minute traces of arsenic, the serum may give absolutely negative reaction; and yet that same serum exhibits these protective properties when injected into a second animal.

We incline to the belief that these are examples of non-specific immunity, but mention these observations because Besredka is a worker of distinction, and, if true, we have here indications that as the organism becomes accustomed to the presence of toxic substances of known composition its cells elaborate something which discharged into the bodyfluids, is capable of neutralizing that toxic substance. But Ehrlich and his pupils have entirely failed to confirm Besredka's observations, Ehrlich himself laying down that such antibodies can only be produced by the reaction to substances which are capable of direct assimilation, and enter into synthetic union, with the cell substance. He draws the distinction between alkaloids, glucosides, and other drugs, that these, while entering the cell substance and acting upon it, do not become

fixed by the cytoplasm, but can be dissolved out, whereas, the whole group of toxins, to be presently noted, do become assimilated and fixed (p. 476). It is interesting to note that Ehrlich himself has supplied a striking example of acquirement of inherited insusceptibility to arsenic salts, in his observations upon trypanosomes subjected to the action of trypanroth, etc. How this is brought about is still a matter of debate. Ford, 'however, has recently obtained a definite experimentally induced immunity against a glucoside isolated by him from the poisonous mushroom, *Amanita phalloides*.

To these observations, also, Ehrlich has objected that Ford was not dealing with pure substances. Since then, employing pure material furnished by Professor Abel, Ford has repeated and confirmed his results.² It is at least suggestive that, as regards arsenic (and antimony), we have evidence from other directions of fixation of these elements by certain orders of cells (liver cells, etc.), and, as regards the glucosides, that there is a glucoside-like compound present in nucleic acid, rendering it not impossible that the cells may thus show an affinity toward bodies of this nature. In other words, it is possible that Ehrlich draws too sharp a distinction; there may be intermediary cases.

With regard to morphine, Faust's' observations are of importance. When injected subcutaneously into the ordinary dog, the greater part of the morphine is discharged through the bowels. By gradually increasing the amount injected, a point was reached at which at last the excretion through the intestine ceased. Analysis of the tissues in such animals showed that it was not retained within them. The only conclusion is that the organism gradually, under increasing doses, gains the power to dissociate the morphine. Where this happens, and what the process is, demands further investigation. There are indications that similar dissociations, or destructions, are developed in the process of immunization to alcohol, strychnine (?), and cocaine. In none of these cases, however, has so far any sure evidence been obtained of antioxin production. The cases that have been recorded 'ail to stand criticism.

It may be that in all this group there are developed processes of another nature. Ransom, for example, has shown that cholesterin, which is a common constituent or by-product of the tissues, neutralizes saponin, and several observers have noted that the loosely combined sulphur of the organism can neutralize highly poisonous nitrites up

¹ Jour. of Infec. Diseases, 3:1906:191; Jour. of Exp. Med., 8: 1906:437; and Jour. of Biol. Chemistry, 2:1907:273.

² Jour, of Infec. Dis., 5; 1908; 116. He finds that the poison of this mushroom contains two principles, the one toxic, the other hemolytic. It is the latter that is of glucoside nature, and employing it in proteid-free condition, he was able to gain an active antihemolytic serum, operative in a dilution of 1 to 5000. For analysis of this glucoside, see Abel and Ford, Arch. f. Exp. Path. u. Pharm., Suppl. Bd. (Schmideberg Festakr), 1908; 8.

² Arch. f. Exp. Path. u. Pharm., 44:1900;217. This has been confirmed by Cloetta, ibid., 50:1903:453.

PHYTOTOXINS

to a certain point by converting them into more harmless rhodan compounds. If we regard the antitoxins as more particularly albuminoid compounds, split off from the cell molecule, it may be that in these other cases there is induced an increased production of non-albuminous bodies, capable of neutralizing one or other poison. But it may also be that more exact research and the employment of other methods show eventually that the processes are of the same nature as those obtained with the bodies we shall next study, and that Ehrlich's distinction, useful as it has been, is not absolute.

Immunization against Albuminoid Vegetable Poisons: Phytotoxins. There are, as already indicated, certain vegetable poisons of a proteid nature, as distinct from the more usual alkaloids and glucosides, and these are extremely toxic. Abrin, ricin, robin, and crotin are the best-known members of the group. The seeds of the praver bead, or jequirity, and of the castor oil plant both yield when suitably treated bodies of this proteid nature. They are, it may be, conjugated proteins. Thus, the former, abrin, has been resolved into an albumose and a globulin, destroyed at different temperatures, both producing similar symptoms. It is still a matter of debate whether the toxic properties are due to the actual g obulin or albumose, or whether each has associated with it a third substance, which is the actual toxin. Nay, more, it has been suggested that the toxin is not so much an actual substance as a property; that the albumose and globulin have a force inseparable from them, comparable to the conferment of magnetism upon previously non-magnetized iron-a suggestion in line with that we have noted in connection with ferments (pp. 71 et seq.).

As the result of a very full study of ricin, Cushny' came to the conclusion that the poison is itself a protein, a globulin, or is so bound to a protein that the usual methods will not bring about separation. Heating to 51° C. leads after a considerable period to coagulation, with loss of its toxic properties. It is so powerful that 0.04 mmg, per kilo. kills a rabbit; or, according to Ehrlich, 1 gram is adequate to kill a million and a half guinea-pigs, with symptoms of acute cedema, inflammation, and necrosis of the tissues in the region of the injection. It is to be noted that there is a *period of incubation* before the symptoms make their appearance, in this resembling what we are familiar with in connection with the bacterial poisons. After seeming to be quite well for four or five days following upon inoculation, the rabbits suddenly show symptoms, and die in the course of a very short period. This, it is true, strongly suggests the action of a ferment and the accumulation in the system

¹ Arch. f. Exp. Path., 41; 1898; 439. Jacoby (ibid., 46; 1901; 28), by digesting the seeds, claims to have isolated a strongly toxic substance with no proteid properties; but Osborne and Mendel (Amer, Jour, of Physiol., 10; 1904; xxxvi) confirm Cushing, gaining an intensely toxic body, which they found inseparably associated with an albuminous body. We shall refer later (p. 540) to Liebermann's observations that certain non-proteid dissociation products of ricin possess hemolytic properties.

of some second substance able to directly combine with or act upon the cell substance. But Cushny could obtain no signs of accumulation of poison in the blood, nor of the presence of any new toxic substance possessing properties differing from those of the ricin injected.

Ehrlich was the first to call attention to the fact that, if mice or other animals be fed upon slowly increasing doses of ricin, they gain immunity such that they can take one hundred times the fatal dose with impunity. To immunize by subcutaneous inoculation, very much smaller doses have to be employed, and care has to be taken that the animal recovers from any signs of local or general disturbances before a second dose be given. But the immunity can be carried to such an extent that these animals will now stand *five thousand times the lethal dose*. In other words, it is possible to obtain an extreme grade of active immunity.

If the serum of animals thus highly immunized be mixed with ricin outside the body, it is found that the poison is rendered inert, and the mixture inoculated into another animal in large amounts is without any serious effects. The indications point to the fact that there is thus brought about a union between the poison and a substance formed by the body and present in the blood, and this union appears to resemble the formation of a double salt (p. 511).

Striving to analyze what happens in poisoning with ricin, and in the production of immunity to the same, we are met with a series of very curious facts. In the first place, the ricin is seen to have two properties which are separable. In toxic doses it causes agglutination of the blood corpuseles, and this agglutination can be well brought about outside the body, a mixture of a sufficiency of the poison with defibrinated blood causing the corpuscles to clump together and to be precipitated, the supernatant fluid becoming quite clear and limpid. And this agglutinating power marches hand-in-hand with the toxic properties of the substance, so that the extent of agglutination has been used by various observers as an index of toxicity. But, as Flexner has shown, the agglutination and the consequent thrombosis and blocking of the vessels is not the esential cause of the lethal action of ricin. The essential poisonous properties are shown in the extensive focal necroses in the liver and other organs. Indeed, as Müller has pointed out, under the action of pepsin and hydrochloric acid, the agglutinative action is destroyed without the toxicity being diminished.

Are we dealing here with a mixture of two substances or with one very complex molecule, which is capable of undergoing sligh chemical change without all its specific functions being destroyed? In this connection there are observations of very considerable interest. If ricin be heated to 100° C, for two hours, it undergoes coagulation and is rendered wholly inert, so that large doses may be injected without any toxic effects showing themselves, and yet animals treated with this modified toxin are found to become immunized.¹ The modified

⁴ Such modified toxins Ehrlich terms *toxoids*. We shall meet with them again in discussing bacterial poisons,

PHYTOTOXINS

ricin is no longer poisonous, but is still able to set up those changes in the organism which result in the production of an antibody.

In this connection we may refer to robin, the active principle of the seeds of *Robinia pseudacacia*. This is a proteid body of the same order as ricin, but much weaker in its effects; yet the blood of animals immunized against it becomes strongly antitoxic to ricin. As Ehrlich suggests, robin may be a toxoid of ricin. In this relationship there is no question regarding duality of the active substance. We have to regard the molecule of the poison as becoming so modified by heat that it cannot enter into *harmful* combinations with the cell substance of the animal inoculated, but not sufficiently modified to prevent it reacting, to a certain extent, with that cell substance, and thereby stimulating the production of antitox is.

Yet another remarkable fact has been noted. If sufficient ricin be added to the serum of immunized animals, with its contained antiricin, so as to make a neutral and inert mixture, which, upon inoculation, produces no toxic effects, that mixture *is still capable of producing active immunity* when injected into the species of animals which yielded the antiricin. It is clear, therefore, that in this combination between toxin and antitoxin the toxin is not destroyed; it continues to exist as such in the combination, and is able to stimulate to production of antiricin by the organism, although it is unable to cause toxic effects.

Little is known regarding the process whereby the antiricin is developed —regarding the cells which give rise to it. During the incubation period there has been noted an increase in the leukocytes, but the general opinion is that the leukocytes play here but little part of the process of immunization. That the stroma of the red blood corpuscles with which the agglutinin becomes connected plays any part is extremely doubtful. We shall refer later to Römer's suggestive observations on the site of production of antiabrin (p. 516).

Obviously, the presence of the ricin in the system leads to the active production by the tissues and passage into the blood of a substance which is able to neutralize large amounts of ricin. Is this directly derived from ricin? Is it a substance normally present in the tissues which is developed and poured out in increased quantities under the stimulus of the ricin? Or, on the other hand, is it a product new to the organism?

Once an animal is immunized against ricin it may be bled and re-bled, and still its blood continues to be antitoxic; its tissues have acquired the habit of discharging antitozin, a fact which scarce favors the view that this antiricin is directly derived from the ricin injected. According to Bashford, the most careful studies fail to reveal that the blood serum of the normal rabbit or guinea-pig has the faintest sign of the presence of an antiricin present under normal conditions. We are here dealing, it would seem, with the development of a substance wholly new to the organism—a fact of considerable importance in connection with the theory of adaptation. Something wholly beyond any principle of "survival of the fittest" is requisite to explain this new development.

Immunization against Substances of Unknown Constitution.-Enzymes and Anti-enzymes.-The toxins in their properties so closely resemble ferments that it might be expected that, at least as regards unusual ferments or ferments out of place, a series of facts would be ascertained closely parallel to what has been determined in regard to the former set of substances. And this is so, though with certain differences, which, in our opinion, form the only serious data-as distinguished from hypotheses of mode of action-upon which he found a distinction between the two orders of bodies. Hildebrandt⁴ showed, in 1893, that repeated injections of emulsin lead to a tolerance of the same, due to the development of an antiferment. The development of such antiferments was confirmed by von Dungern² as regards the proteolytic enzymes of bacteria, and more especially by Morgenroth³ as regards the rennet ferment. Morgenroth demonstrated that, as the result of the repeated injections of goats with rennet, their serum became able to neutralize that rennet; and, further, that this reaction is quantitative, a given amount of serum rendering inactive a fixed amount of rennet solution of known strength. But just as an animal immunized against a specific microorganism is not rendered immune to microbes which, morphologically and in cultural characters, are evidently closely allied, so such immunization of an animal with animal rennet will not protect against rennet ferments obtained from plants (e. g., cynarase, gained from the *Cynara cardunculus*); or, more exactly, the blood serum will not prevent the latter from coagulating milk.4 In like manner. Sachs⁶ has demonstrated the development of an antipepsin: Gessard,⁶ of an antityrosinase; Moll,⁷ of an antiurease (urea-splitting enzyme); Schütze,⁸ of an antisteapsin (fat-splitting enzyme), and of an antilactase (enzymes which split milk sugar), etc.

An important distinction between the immunization against these enzymes, or, at least, against enzymes which normally are in relationship to the organism, and that against bacteria and their toxins, is that the development of anti-enzymes is limited, and does not become extreme.⁹ This, it may be, has its explanation in a surplus of such anti-enzymes, disturbing normal metabolism and absorption of foodstuffs, and so stimulating sundry cells to produce anti-anti-enzymes, whereby a regulating mechanism is set up preventing the anti-enzymes from accumulating in the organism beyond a certain point. Another expla-

⁵ Fortschr. d. Med., 20: 1902: 425.

⁹ So far, to our knowledge, there are no adequate studies upon the upper limit of development of antibodies in relationship to enzymes which, in their activities, are foreign to the animal body (as are the toxins).

¹ Virchow's Arch., 131: 1893: 5.

² Münch, med. Woch., 1898, and Centralbl. f. Bakt., Abt. 1:24:1898,

³ Centralbl. f. Bakt., Abt. 1:26; 99, Nos. 11 and 12, and 27, 1900.

⁴ Briot, Thèse de Paris, 1900.

⁶ Ann, de l'Inst. Pasteur, 15: 1901.

⁷ Hofmeister's Beitr., 2: 1902.

⁸ Zeitschr, f. Hygiene, 48: 1904; 457, and Deutsch. med. Woch., 30: 1904, 308.

nation is afforded by certain remarkable observations of Beitzke and Nauberg.¹

According to these observers, the serum of animals immunized against emulsin possesses the property of synthesizing glucose and galactose into lactose; it combines, that is, the products of the ferment action. The process, it is true, is a slow one, and may be, after all, but a special case of reversibility of enzyme action, but, if confirmed, it would suggest that between enzymes and anti-enzymes a self-limiting cycle is developed.

It is worthy of note, also, that bodies of the nature of antiferments to the normal enzymes of the organism are normally present in the organism; an antirennin, or antirennins, have been isolated from normal blood, and of particular interest is the discovery of Weinland² that the cells of the gastric mucosa contain antipepsin, amplified by Pollack's³ determination that pancreatic extract similarly contains an antiferment against the digestive action of the pancreas. Here we have gained at last the needed explanation why the digestive organs do not digest themselves.

So, also, it has to be noted that there are certain common enzymes of the organism against which, so far, observers, by injection of these ferments, have been unable to obtain antiferments. If fibrin ferment, for example, be inoculated into an animal, no antifibrin ferment is produced—only precipitins; although Bordet has brought forward evidence that the inoculation of *fibrin-containing* blood into another species of animal stimulates the production of an antibody. No antipepsin or antidiastase, so far, has been developed. It may be that here some principle is in action, as already suggested, inhibiting the development of antibodies to ferments which are widely distributed throughout the organism; for peptic and diastatic ferments, it must be remembered, are not developed by the gastric and salivary glands alone, but, as shown by the results of autolysis, are producible by the cells of all organs, and the same is true of fibrin ferment, as indicated by the process of coagulation necrosis; that just as in normal assimilation the cells are not stimulated to produce antipeptones and anticarbohydrates, so the normal cell enzymes, when absorbed and presented to the cell, set up no antagonistic reaction. The general principle would seem to be that molecules which afford particular side-chains are not stimulated to produce these in excess in the presence of free side-chains of the same order; still less does that presence stimulate them to produce antibodies to their normal side-chains. We thus are not inclined to lay great weight upon the distinction between normal ferments and toxins, more particularly when we recall that, as in tuberculosis, a similar lack of reaction, or imperfect reaction, may present itself toward bacterial products.

¹ Verhandl. d. deutsch. pathol. Gesellsch., 1905; 160.

² Zeitsch. f. Biol., 44: 1902.

³ Hofmeister's Beitr., 6: 1904.

To this matter of the parallelism or identity between enzymes and toxins we shall return later, when we have noted the data bearing upon the production of antitoxic bodies proper. We shall leave until then the discussion of the action of enterokinases, etc.

Toxins and Antitoxins.—That certain bacteria produce diffusible, poisonous substances, or toxins, was noted very early in the study of bacteriology; indeed, even before specific forms of bacteria were distinguished, it was observed that putrefying material afforded substances like Panum's sepsin, which were intensely toxic; and when Koch introduced the method of growing pathogenic bacteria upon solid media, and the solution of gelatin was found by Fermi and others to be due to proteolytic enzymes, it was natural that observers should conclude that the toxic action of bacteria in the body was due to a similar excretion or diffusion out from the bodies of bacteria of ferment-like substances.

We know that, *in vitro* at least, and probably, also, in the body, this diffusion out of toxins by living pathogenic bacteria is the exception and not the rule. Nevertheless, a very important step forward was made when Roux and Yersin studied the diffusible toxins (ectotoxins) of the diphtheria bacillus, and Brieger (though less thoroughly) those of the tetanus bacillus. The toxins of both of these organisms were discovered to be of extraordinary virulence, or to have intense toxicity. By no means could they be gained in a pure state. Nevertheless, the impure precipitate gained, for example, by salting out, or precipitation by phosphates, and present in only minute amounts, was found so powerful that with the diphtheria toxin 0.0002 mg, would kill a guineapig of 250 grams within three days.

At a relatively early date (1887) it was shown (by Salmon and Smith) that such diffusible toxins inoculated into animals would produce immunity, but the most notable advance was made when, in 1890, Behring, with his colleagues, Kitasato and Wernicke, demonstrated that when such immunity was produced, the blood serum of the immunized animals contained substances which would neutralize the diffusible toxins, and so discoverd the antitoxins, proving, further, that such immune serum injected into animals suffering from either tetanus or diphtheria would in them neutralize the toxins and induce a condition of passive immunity. The more intimate studies upon the development of protective substances by the organism date from this discovery. It became necessary, in the first place, to determine with as great an accuracy as possible what was the advisable dose of antitoxin to be administered in order to neutralize the toxin; to do this a standard toxin had to be developed. What should constitute the minimal lethal dose had to be agreed upon-the smallest amount which would surely kill a given animal (guinea-pig) of normal size (250 grams) within a given time (four days). We owe to Ehrlich the fullest studies upon this subject and the most important deductions from the data obtained in these studies.

Behring had noted that outside the body in the test-tube the antitoxin acted upon and neutralized the toxin. (Like observations had

BACTERIAL TOXINS AND ANTITOXINS

also been made by Kanthack,1 for snake venom and its antitoxin, and by Denvs and Van der Velde2 for leukocidin-the leukocytedestroying substance obtained from cultures of pyococci-and by Ehrlich for ricin and antiricin.) Now, Ehrlich established clearly that this action in vitro follows the laws regulating the formation of double salts: (1) If an amount, x, of a particular antitoxin solution neutralizes an amount, y, of a particular toxin in a given time, then 3x is requisite to exactly neutralize 3y; (2) the higher the temperature (below the limits beyond which the antitoxin became inert) the more rapid the reaction; and (3) the reaction proceeds more rapidly in concentrated than in dilute solution. Obviously, while the formation of antitoxins is a vital process, the reaction between these and the toxins is purely chemical, occurring outside as well as within the organism. These conclusions were confirmed when Martin and Cherry and Brodie demonstrated, by their filtration experiments, that, whereas toxin alone will pass, as the latter showed, through porcelain filters impregnated with gelatin under high pressure, there is no passage when they have been acted upon by antitoxins, *i.e.*, they have undergone combinations with the same. This doctrine of Ehrlich, namely, that toxin and antitoxin, or, stated broadly, antigen³ and antibody, in their combination conform with the law of multiple proportions must, we think, be accepted for simple cases. There are some curious apparent divergences from the law when, instead of equivalent portions being brought together, less of the antibody is added than is necessary to effect neutralization. These and other considerations have led Arrhenius and Madsen to conclude, in opposition to Ehrlich, that instead of antigen and antibody acting like strong acids and strong bases and conforming with the law of multiple proportions, they act like weak acids and weak bases. and follow more nearly the law of "mass action" of Guldberg and Waage, doing which they are apt to exhibit reversible action. From this it would follow that the union is unstable and easily dissociated. To enter into a full account of this controversy would carry us too far afield.⁴ Here it can only be said that the matter is not yet settled one way or the other, nor is it likely that a sure decision can be given until further advances have been established in our knowledge of the chemistry of colloid bodies. Biltz's' endeavor to employ the data at our disposal regarding that chemistry to a solution of the problem is suggestive but not absolutely convincing. At present our decision must be the wise one of old Sir Roger de Coverley-"that much may be said on both sides." (See also p. 522.)

⁴ Jour. of Physiol., 13: 1892: 272.

² La Cellule, 11: 1896: 359.

³ The term antigen $(\delta n \tau_i)$ against, $\gamma \gamma n m$, I produce) has of late come into general use to signify any body against which, when injected into the organism, an antibody is developed.

⁴An impartial account is afforded by Emery in his *Immunity and Specific Ther*apy (London, Lewis, 1909).

⁸ Zeitschr. f. Physiol. Chemie, 18: 1904: 615.

Toxins.—What, then, are toxins? Unfortunately, no precise definition would appear to be possible. To describe them, as is often done, as poisons, against which it is possible to gain immunity by means of antibodies, is to include under this term a very large number of different substances—bacterial products, venoms of various animals, animal cell substances, sundry vegetable origin. It seems difficult, however, to approach any clearer definition, and, for practical purposes, this is adequate. If we attempt to characterize them more closely, we note: (1) that they are, one and all, the products of cell metabolism; (2) that they act in most minute doses; (3) that they diffuse with difficulty; (4) that hitherto not one of them has surely been obtained in a state of purity.

We are accustomed to regard all these bodies, if not as proteins proper, nevertheless as closely allied thereto. It must be remembered that the evidence here is presumptive and far from being positive. Like proteins, they are of colloid nature, as indicated by their low diffusibility. But as a body they are not absolutely non-diffusible. Nothing like a splitting-off of amido-acids can be determined, as with proteins proper. Nor, again, when obtained in a state of relative purity, do they necessarily afford those two characteristic reactions of proteids, namely, the binret reaction and Millon's test. But, on the other hand, they act in such extraordinarily small amounts, that to gain from them recognizable amounts of amido-acids would be beyond the scope of the chemist, and the same is true of the two reactions above noted. For these tests a certain minimum of material must be present; with the toxins, it may well be that we are below that minimum.

On the whole, although we have to admit that the evidence is far from complete, it is most satisfactory to regard the toxins, and particularly the bacterial ectotoxins, as cleavage products of protein metabolism, and as approximating in their nature to primary, non-polymerized, protein molecules. In favor of this view are the observations of Arrhenius and Madsen upon the rate of diffusion of toxins compared with salts and (definitely proteid) antitoxins, in which it was found that the former occupy a position between the latter two; their molecules are thus evidently larger than those of crystallizable salts, but smaller than those of the ordinary proteins.

But Arrhenius and Madsen¹ were dealing only with *cctotoxins* diffusible toxins; their observations do not apply to the less diffusible *cudotoxins*. And here we encounter identically the same problem regarding chemical constitution that we noted with the diffusible extracellular and the non-diffusible intracellular ferments, so that, taking merely bacterial toxins as a body, it is extremely doubtful if we are justified in regarding them all as cleavage products. It appears to be a sounder course not to speak of *toxins*, as a whole, as a specialized group

¹ Festschr. z. Eröffnung d. Scrum Instituts, Kopenhagen, 1902.

TOXINS: THEIR NATURE

of chemical substances, but of *toxin action*, and the capacity to elicit the production of antitoxins as a property of the protein molecule and of sundry of its dissociation products. Toxin action, that is, may be regarded as a form of enzyme action which may manifest itself either in connection with large and complicated protein molecules or with molecules dissociated from the cell, so few in number and so small (relatively to the complete protein molecule) as to render chemical detection difficult. Nay, it may well be that the almost constant association of "toxins" with albuminous matter is an indication that the toxin action is a property of this albuminous matter which may still be retained by certain molecules, the products of dissociation of the same, when an attempt is made to separate toxin from albumin.

It is our duty to point out that this is not the usually accepted view; it is almost universal to regard toxins as a particular class of chemical substances and to neglect the consideration of toxin action as a physical property of matter, having a particular molecular arrangement. The view here enunciated leads to the consideration of the phenomena of immunity as of an order intimately allied to those of enzyme action. This we admit, while admitting at the same time that "toxins" and "enzymes" are not wholly identical. They have many points of identity: Thus, to epitomize from Emery: (1) Both are produced by living animal or vegetable cells and belong to the two orders-either extracellular (free enzymes, ectotoxins) or intercellular and attached (intracellular enzymes, endotoxins). (2) Both begin their action by forming a combination with the substratum, $e, q_{..}$ fibrin placed in gastric juice at 0° C., taken out and repeatedly washed in the cold from all traces of enzyme, will undergo digestion when raised to the body temperature. The fluid of growth of the tetanus bacillus contains a toxin (tetanolysin) which dissolves the red corpuscles. If red corpuscles be placed in such fluid at 0° C, and then be centifugalized and thoroughly washed in the cold, when raised to a temperature of 37° C, they undergo hemolysis) (see also later). (3) As shown by the above examples, both enzymes and toxins demand suitable temperature for the development of their activity. (5) Similarly in both, combination and the exercise of specific activity are separate functions.

The one important distinction between the two is that toxins do not demonstrate to the same extent as do the enzymes the capacity for a minimal quantity eventually to dissociate a maximal amount of the substrate. It is difficult to prove or disprove this with toxins proper in the living body; indeed, it may with some justice be urged that toxins acting in a living system induce the formation of antitoxins which eventually arrest their activities; whereas, the characteristic enzyme action in vitro proceeds without any such anti-enzyme development. But with the closely allied hemolysin s it can be surely demonstrated that there is an exact quantitative relationship between the amount of hemolysin present and the volume or number of red corpuscles dissolved.

Not to confuse the reader, we will continue to speak of toxins and describe their properties in the accepted way. In the succeeding

ALINAKKI GNV NOLLYZINAKKI

'ram matter' this same binding of the neurotoxic component of snake venom with pig is employed. Flexner and Noguchi² have shown the occurrence of the toxin, which is not the case when kidney or other organ of the guineamade, be centrifugalized, the supernatant, clear fluid is wholly free from animals. And, as Michner has shown, if such a mixture, properly portions, the emulsion being without effects when injected into other -ord annihola in a babba neuron summa sailaruna yasala os ot bruot firmed, can be demonstrated outside the body, when brain substance is combination, as shown by the last two observers, and abundantly conit enters into intimate combination, and cannot be separated. The (more particularly the gray matter), and in the rabbit's liver and spleen, easily be separated; it is, at most, "adsorbed." In the nervous tissue the other organs the absorbed toxin is in loose combination, and can has been taken up by these three organs, but the very reverse. In all the like properties. This does not mean that none of the tetanotoxins horse, and the guinea-pig, in the rabbit the liver and spleen manifest Wassermann and Takaki? found that, while this was true of man, the are found toxic. That one exception is the brain and nerve matter, blood loses its toxicity, extracts from all the organs, with one exception, ample, as Ransom' pointed out, tetanotoxin be employed, while the Doing this, the organs become separated into two classes. If, for exso can be determined by making extracts from the various organs. tion by the cells of various tissues, and by the leukocytes. That this is process of neutralization occurring in the circulation, but to an absorpinnocuous. This, in the non-immunized animal, is not due to any in some cases, three or four minutes after inoculation the blood is found blood of an untreated animal, these disappear with relative rapidity; toxins-that we have mainly under consideration. Inoculated into the paragraphs it will be the toxins in the narrower sense-the bacterial ecto-

We call attention to these joints because everything indicates that it is those cells and issues which anchor the toxins that eventually develop the unitoxius. The mere loose adsorption of the toxin is not sufficient for this purpose.

Here Metchnikoff has recorded a very important observation. The toroise is uninfluenced by tetanus toxin, and after inoculation it is found that none is present in its brain or other organs, and no amount of injections of the toxin will in it produce antitoxins. The alligator,

¹ Neitschr. f. physiol. Chem., 31: 1901; 282.

³ Berl, klin, Woch, 35, 1898; 5.

1225 : 7061 : 9 ' Pak dx3 10 .mor ;

⁴ This same barring with specific cells has been abundantly proved as between the red blood corpuscies and "toxins," which lead to hemolysis—tetanolysis, staphytolysis, ricin, sunke venom, etc. Suchs found that spider venom (archeolysis), which has no arcino on genine-pige "crythrocytics, remained in solution on centrifugationind, but, noised to rats' crythrocytics (which it destroys), the supermutant fluid, tion, but, added to rats' crythrocytics (which it destroys).

upon centrifugalizing, was free from it.

PRODUCTION OF ANTITOXINS

while similarly showing no nervous effects upon inoculation, is found to have the toxin bound in certain organs, *but not in the nerve matter*, and in it successive inoculations lead to abundant antitoxins appearing in the blood. Ford¹ has, by other methods, demonstrated that antitoxin production only occurs in individuals whose cells have the specific power of binding the toxin.

For specific antitoxins to be produced, there must first have been a direct union with, and action upon, the cell substance on the part of the toxin. As Ehrlich points out, alkaloids only enter into the looser combination, and to this is to be attributed the non-formation of anti-alkaloids. With Wassermann, we can in this respect compare alkaloids and toxins to saccharin and sugar. Both act on sundry cells and give the sensation of sweetness; the latter only undergoes assimilation.

The observations of Metchnikoff demonstrated clearly that to stimulate the production of antitoxins it is not necessary that the toxins so act upon the cells as to set up the symptoms of disease. Ehrlich has, indeed, shown that by subjection to heat and other methods the diphtheria toxin can be so modified that its toxic action is greatly lowered or entirely lost; it may be rendered quite harmless, and yet, inoculated, such modified toxin leads to the development of antitoxins. The existence of these modified toxins, or *toxoids*, and their capacity to induce immunity afford proof that the toxin molecule consists of at least two subordinate groups—one that anchors the toxin on to the cell substance, the *haptophoric* group of Ehrlich's terminology; the other, the *toxophore*, which, when present, so influences the cell activities as to set up toxic disturbances.

A further proof of the existence of these two constituents has been afforded by Morgenroth.² He took three frogs and injected tetanus toxin into them without obvious effect, for at ordinary temperature these cold-blooded animals are unsusceptible to tetanus. These he kept for three, four, and five days to some weeks in the cold. Had the toxin been still circulating in the blood the subsequent injection of tetanus antitoxin would have neutralized it; had it been excreted, again no symptom would have developed. But now, upon warming these frogs, symptoms of tetanus appeared. The only possible explanation is that the toxins had become anchored to the nerve cells, etc., and, being so combined, could not unite with the circulating antitoxins. But, if so, then the haptophoric constituents must unite with the cell substance in the cold, and the toxophoric constituents only become active when the temperature is raised.

Ehrlich has brought forward evidence of the existence, more particularly in old toxin solutions, of a series of modifications of the toxins: protoxoids, syntoxoids, epitoxoids, and bodies which have little affinity to antitoxins. These last he has termed *toxones*. The active toxins themselves may also exhibit variations in their avidity to combine with


¹ Zeitsch, f. Hygiene, 40; 1902; 363.

n-

² Arch. Internat. de Pharmako-dynamie, 1900.

antitoxins, and in this way he has distinguished between proto-, deutero-, and tritotoxins.

The Antitoxins Proper.—Mode of Development.—It is evident from the above that the toxins enter into combination with the cell substance, and as we find that when this combination occurs the toxin becomes neutralized, it becomes most probable that the particular cells in which the toxins become anchored are those which eventually discharge into the blood the substances—antitoxins—which are capable of neutralizing them. The conclusive proof of this local development of antitoxins has been afforded by Römer, not, it is true, with bacterial toxins, but with abrin. This, it has long been known, has a peculiarly powerful effect upon the conjunctiva. By the exhibition of increasing strengths of abrin



in the right conjunctiva of a rabbit, Römer¹ gained a local immunity, so that the conjunctiva was no longer sensitive; then, at the beginning of the third week, before this should affect the organism in general, he killed the rabbit, took the conjunctive, and triturated each separately with a fatal dose of abrin. The injection of the emulsion from the right (immunized) conjunctiva was without effect; the like injection from the left was fatal. Thus. clearly, the cells which had absorbed the abrin had developed and contained anti-abrin in sufficient amounts to neutralize the poison. And, while there is evidence that certain leukocytes-Metchnikoff's

macrophages—play an active part in this development, it is obvious that they are far from being the only cells involved.

How, then, is the antitoxin developed? It used to be thought that there was a direct conversion of the toxin into antitoxin. This certainly is not the case, for the amount of antitoxin is altogether out of proportion to the amount of toxin injected. We have exactly the same continued production of the antitoxins as noted in the case of anti-abrin (p. 506). Knorr² has shown that a toxin unit in a horse (immunized against tetanus) leads to the production of about 100,000 antitoxin units, and McFarland³ has made similar observations. While the toxins stimulate the cells in the first place, it is the cells which assimilate the necessary constituents and build up and discharge the antitoxins.

The Side-chain Theory.-How are we best to picture the manner in which this is brought about? Here Ehrlich's conception of the process

¹ Arch. f. Ophthalmologie, 1901. ² Münch. med. Woch., 1898; 321 and 362.

³ Text-book of Bacteriology, third edition.

THE SIDE-CHAIN THEORY

is the only one which seems to be satisfactory. We have already noted that the cell substance enters into relatively firm combination with the toxin, just as it combines with constituents of the foodstuffs in order to assimilate them. We have seen, also, that the toxin possesses a haptophorous constituent which unites with the cell substance, and this whether the toxin is actively toxic or not; or, in Ehrlich's terminology, whether it possesses an active toxophorous constituent or not. Now, if we predicate these constituents for the minute toxin molecules, the specific matter of the relatively large cell molecule must be very much more complex. If the toxin molecule has its haptophore, whereby it anchors itself onto the cell substance, so we must imagine that the cell molecule has not one, but several orders of "anchors," whereby it attracts and combines with itself all the various orders of foodstuffs. The cell molecule, that is, also possesses haptophores, which, in this

case, Ehrlich terms *receptors*; and it is to some special order of anchor that the haptophore of the toxin molecule becomes attached. These receptors, it will be seen from what we have said regarding the constitution of the proteidogenous or biophoric molecule, are *side-chains* (p. 67); and certain orders of unsatisfied side-chains must be regarded as having affinity for, attracting and combining themselves with the toxin molecules. Nay, more, it is with one particular portion of the toxin molecules (the haptophorous) that the combination is



side-chains must be regarded as having affinity for, attracting and combining themselves with the itoxin molecules. Nay, more, it is with one particular portion of the toxin molecules (the hapto-

effected; these may or may not have a toxophorous moiety. That is not concerned in the act of combination, but, when present, after the combination has taken place, it may so effect the total constitution of the molecule as to lead to grave cell disturbances.

This is the groundwork of Ehrlich's now celebrated "side-chain theory," and up to this point we think our readers will agree with us that the theory is well grounded upon experimental data, and is strictly in harmony with our general conception of cell activity. But this simply carries us to the point at which the toxin has become associated with the cell substance. We have simply reached the point at which the toxin, unless too strong for the cell, or unless too many toxins have become joined on to the receptors of a given cell, is neutralized. We may speak of such neutralizing receptors as intracellular antitoxins. Now we have to explain how there is set up a discharge of free antitoxins into the blood, and here it is that, while forced to accept it, we cannot but feel that the theory is deductive, and on less secure ground.

The very fact, as pointed out by Cobbett,' that the blood serum of human beings who have never suffered from diphtheria contains a recognizable amount of diphtheria antitoxin would suggest that antitoxins are not primarily specific; in other words, that the cells of the body normally discharge substances capable of combining with and neutralizing diphtheria toxins. To this it may be objected that human beings may be subject to what we have termed "subiffection" with diphtheria. But the same is true also of horses' blood serum (Wassermann³); and when, as von Dungern⁵ has shown, rabbits' blood serum contains an fantitoxin against the poison of starfish eggs, our point becomes established. There is, that is, a normal discharge into the blood plasma of a large number of potential antitoxins, of bodies having affinities for one or other toxin, and it is only when these toxins gain entrance into the system that the particular antitoxins become developed in relatively enormous quantities.

If, with Ehrlich, we regard these as discharged cell receptors or sidechains of particular orders, then we have to assume that the very act of combination of the toxin molecules with the receptors stimulates the cell substance to reproduce more of these particular receptors than are necessary, and that the overproduction is discharged out of the cell. This assumption, it is true, appears to accord strictly with the facts of the case; but we find it difficult to picture the process whereby this overproduction is brought about. Ehrlich bases himself upon Weigert's law of inertia—the law, namely, that once a cell is stimulated to perform a certain act, it continues to perform that act for some time after the stimulus has ceased to act. But this production of antitoxins continues for a longer period than was contemplated by Weigert, and that law does not explain why the anchoring of the toxin by a particular side-chain acts as a stimulus to the production of new side-chains of the same order.

While offering these criticisms, it must not be thought that we are opposed to Ehrlich's theory; on the contrary, we accept it almost in its entirety; only at this one point it seems to us that the chain of events depicted has some weak links. Ehrlich throughout purposefully speaks in most general terms of the constitution of the cell, and draws no distinction between cytoplasm, paraplasm, and nucleoplasm. He, in fact, leaves it an open matter which portion of the cell is the seat of the changes demanded by his theory. The conception we have given of the cell structure (p. 156) leads us to consider that a poison can only interfere seriously with cell activities when either mediately or immediately it tells upon the nuclear biophores. We would regard a toxin molecule diffused into or absorbed by the cytoplasm not, with Ehrlich, as becoming fixed to the biophore, but rather as detracting and dissociating a side-chain from the biophoric molecule. The next stage would depend

¹ Centralbl. f. Bakt., 26; 1899.

¹ Zeitsch, f. Hvg., 19:1895:408.


³ Zeitsch, f. Allg. Physiol., 1:1901.

THE SIDE-CHAIN THEORY

upon the number of toxin molecules gaining entrance to the cell, and, it may be, upon their ferment-like activity. An excessive number would lead to such a dissociation of the biophoric molecules as to entail biophoric dissolution and cell death. The known facts regarding the extraordinary minute, unmeasurable quantities of toxin capable of causing death of relatively large animals leads us to believe that ferment action must play a part; namely, that the toxin molecule, having dissociated one side-chain, multi, dissociating the biophore more rapidly than that can build itself up, it causes dissolution of the same, and cell death. Along these lines the difference between toxins and alkaloids, and other cell poisons which can readily be dissolved out of the cell, is that the latter, if they set up cell disturbances, do this by a single act of union and dissociation, the latter by the repeated enzyme-like attraction and dissociation of side-chains from the cell molecules. So far as they

go, the histological appearances of the nuclei—of the brain cells, for example, in cases of tetanus—are in favor of such nuclear dissolution.

Where, on the contrary, the action is not so intense, the dissociation of a sidechain by a toxin molecule is followed by the building up or assimilation by the biophore of another to replace it. Here, also, to explain subsequent events, the toxin molecule must be regarded as



Ehrlich's conception of the regeneration of the cell haptophores, or receptors, to compensate for the loss of those neutralized and removed by the toxin molecules.

endowed with enzyme-like properties. The loss of a single side-chain cannot, no more than any other momentary stimulus, set up the habit of manufacturing side-chains in excess of the needs of the individual cell; but if for some little period that molecule is dissociating sidechains of a particular order, but these are reformed more actively and in excess of the rate of destruction, it appears to us possible to realize the gradual development of a habit of production and casting loose of these side-chains which shall continue long after the primary stimulus has ceased to act. That toxins in a potentially active state may remain in the cells for weeks is demonstrated by Morgenroth's observations upon the tetanotoxins in frogs, to which we have already referred!

¹ Along the lines here indicated Morgenroth's observations are capable of explanation, not so much as evidence of the existence of haptophores and toxophores (though they afford this also), as of persistence of the active poison within the cells, which poison is only able to manifest its results when, by abnormal temperature conditions, the cell metabolism and side-chain production is so lowered that now the toxin gets the upper hand

(p 515). The diagram opposite gives more fully our conception of the way in which the toxin acts in this respect.


It is, we would emphasize, the continued existence of the toxins for some little time within the cell, but not as a part and parcel of the biophoric molecules, that best satisfies the conditions of the case, i. e., the development of the habit of side-chain, receptor, or antitoxin production. And these, developed in excess of the needs of the cell, become discharged into the circulating medium.

On the Nature of the Union of Toxin with Antitoxin .- We have already referred to Martin and Cherry's evidence that toxin and antitoxin unite to form a distinct compound. The union, we now know, is not immediate. This explains the contrary evidence brought forward by Calmette, Wassermann, and others; they did not wait long enough to permit the complete combination. The antivenin to snake venom is rendered inert by subjection to a temperature of 68° C., whereas the venom itself is uninfluenced by a temperature of 80° C.1 Calmette found that the venom-antivenin mixture becomes toxic again if subjected to a heat of 70° C. Wassermann came to similar conclusions with regard to the pyocyaneus toxin-antitoxin mixture. It has since been found that by allowing the mixture to stand for a few hours these results are not obtained. Ehrlich further has shown that the mixture follows the law of true chemical combination in that the union is accelerated by heat and takes place more quickly in concentrated than in dilute solutions. Admitting this, there are certain data which are not immediately recon-Thus, for example, it is found that a combination of toxin cilable. and antitoxin which is absolutely neutral for individuals of one species is fatal for those of another.

This has been explained by Weigert as due to the presence in the blood of the second animal of substances having a stronger affinity for the antitoxin moiety, in consequence of which the toxin becomes set free. Aschoff affords the alternative explanation, that in the first animal (the mouse, for example, treated with a tetanotoxin-antitoxin mixture) tissues other than the nerve cells can combine with the toxin, as, indeed, has been shown to be the case, and that so, with subcutaneous inoculation of what for the mouse is found to be a neutral mixture, there is, nevertheless, an excess of free toxin. If, now, this same mixture be inoculated into a guinea-pig, whose other tissues have not this affinity for the toxin, such free toxin is capable of exercising its full effects on the nerve cells.

Of late years several obvious departures from the simple law of the formation of double salts have been noted in the reaction between toxin and antitoxin—exceptions which it is difficult to explain adequately. We shall not here mention all, but would refer to Emery's conscientious discussion of the subject. Of these, the most remarkable is *Danysez's*

¹ But, as a general rule, it must be noted that toxins are more sensitive to heat than are antitoxins; heating such more sensitive toxins, they are apparently converted to toxoids, which still remain attached to the antitoxins.



'n

a d

it

it

)ľ

n,

10

18

The author's conception of side-chain and antitoxin production. 1. The biophoric molecule situated within the cell, and possessing side-chains of various orders, A, B, C, D. 2. Mode of formation of side-chains. Free molecules (X, Y) diffuse into the cell (or are produced within the cell by dissociation of more complex molecules, also absorbed): these are attracted by an unsatisfied affinity of the biophore, and are built up by it to form the side-chain A. Such side-chain, when formed, may become satisfied by attracting to it other (foodstuff) molecules such as E, having the right order of haptophorous grouping. It is conceivable (but not shown in the diagram) that molecules of the E order may not merely satisfy the side-chain, but detach it so that the compound A-E becomes free in the cytoplasm, or discharged from the cell (active katabolism). 3. A toxin molecule F, diffusing into the cytoplasm has a stronger affinity for the side-chain A than has the biophore, combines with it and detaches it. But when detached and free in the cytoplasm other molecules (G) present in the cytoplasm have now a stronger affinity for the A moiety of the compound A-F and combine with it, liberating the toxin moiety F, which again becomes free in the cytoplasm and capable of dissociating another side-chain A. Or the compound G-A may become discharged from the cell (circulating antitoxin), and then in the altered surroundings the extracellular toxin molecules F may exert the greater affinity and joining with the A moiety become neutralized. It is the G-A compound, and not the side-chain A alone, that constitutes the extracellular antitoxin. 4. In the presence of abundant X and Ymolecules the side-chains A become built up in series, and this whether attached to the biophore or free in the cytoplasm; the more there are freed by the action of the toxin, the greater under these conditions will be the production of antitoxins. Thus, the presence of the toxin molecule F stimulates the cell to the production of increased numbers of the molecules of the particular side-chain order upon which it exerts specific action,

phenomenon, first noted with ricin, and later found to obtain with a great number of toxins by von Dungern¹ and Sachs.² If the amount of toxin has been accurately determined which, added to a given amount of antitoxin serum, completely neutralizes it, and if, now, half the amount of the toxin only be added, then if, at a later period, the other half be added, the result is not a neutral, but a poisonous, mixture. More antitoxin has now to be added to bring about neutralization.

A possible explanation of this phenomenon is to be found in Ehrlich's recognition of multiple toxins and toxoids. It is not unlikely that in a toxin solution, besides the active toxin proper, having the greatest avidity for antitoxins, there are present toxoids with less affini y, which, when the amount sufficient for complete neutralization is added, do not combat with the antitoxin; when only a fraction of that amount is added, are taken up by the unsatisfied antitoxin molecules. By this means, when further toxin solution is added, the molecules cannot find adequate antitoxin molecules with which to unite. But Bordet's argument that the union of toxin and antitoxin partakes more of the nature of the absorption of a dve by a colorable substance cannot be neglected. That absorption is what is now terred an "adsorption" phenomenon. The stainable substance can take up 1, 2, 3 volumes, etc., of the dve, and fix these, without obviously forming thereby a succession of distinct chemical substances. Now colloids (and such presumably are toxins and antitoxins) as a group manifest adsorption phenomena. Hence these anomalies in the combining volumes of toxin and antitoxin may possibly find an explanation without calling into existence Ehrlich's long series of toxoid bodies.

An enzyme-like action on the part of the toxins would also explain the phenomenon. We freely admit that, observing in general the remarkable obedience of toxin-antitoxin mixtures, to the law of multiple proportions, it would at first sight appear that toxins do not act like ordinary ferments. Nevertheless, as we have already shown, from Morgenroth's observations, in relationship to antibodies, ferments do obey this law; a given amount of antirennet serum renders inactive a fixed amount of rennet solution. More suggestive of enzyme action is Behring's^a observations that if a mixture of tetanus toxin and antitoxin be taken in which the toxin is in excess, there is a marked increase in toxicity when the mixture is diluted with water. This, as Jacobi indicates, suggests hydrolytic dissociation.

In general the law of multiple proportions is strongly in evidence, *i. e.*, if 10 volumes of antitoxin neutralize 100 of a toxin solution, 100 volumes of antitoxin are found to neutralize 1000 of the same toxin. The combination, however, is not immediate; while it is relatively rapid with diphtheria, in tetanus preparations it is slow, requiring often

¹ Deutsch. med. Woch., 1904; 275; and 310,

² Centralbl. f. Bakt., 37: 1904.

³ Beitr. z. Exper. Therapie, 1904; 7.

some hours; the age of the antitoxin preparations also introduces variations in rate.

On the Action of Antitoxin upon Toxin within the Organism.— We cannot leave this subject without discussing the mode in which antitoxins introduced into the organism bring about the cure of infection. Experimentally, we find that the smallest amount of antitoxin is required, and the least evident disturbance of the organism occurs, when toxin and antitoxin are introduced simultaneously; and here, again, least of all when the two have been in contact for some period.¹

When symptoms are already present, much larger quantities are required, and when the disease has been active for some days, no amount of antitoxin will arrest the fatal result. But, as just stated, the antitoxin can, under certain conditions, arrest the disease, even when symptoms are present, and when clearly the toxins have already gained entrance into the cells. These, indeed, are the conditions under which diphtheria antitoxin is most often employed in practice, with brilliant results. It is evident, therefore, that the antitoxins can act upon toxins which already have become bound in the cell, but that, just as with the lapse of time the toxin in vitro becomes more and more firmly bound to the antitoxins, so in the cell in vivo there is a similar increasingly firm anchoring of the toxin to the receptors, until eventually the free antitoxins introduced from without are unable to loosen the union. That this is so is confirmed by several observers of the phenomena of so-called "cure in vitro," as observed in connection with hemolytic agents. For a certain period it is found that the addition of an antihemolytic secum to blood corpuscles which have absorbed hemolysins will remove and neutralize these "toxins," but with longer interaction the "antitoxin" is without effect.

This very fact that foreign antitoxins entering the cell can in the carly stage arrest the infective process appears to us to support our contention that the toxins, at a time when they are already setting up active cell disturbances, are not in direct combination with the biophores, but are acting upon them from without—in the cytoplasm. We would conceive antitoxins introduced into the system as: (1) neutralizing any free toxin molecules in the circulating fluids of the body, and so preventing their action upon the cells, and (2) as gaining entrance into the cells, and there not so much acting directly upon the toxin molecules present (for those are already combined) as affording to the biophores that excess of side-chains necessary to build them up again and to neutralize the toxin molecules should [they become temporarily free—as, in short, affording to the cell physiological rest, together with the particular assimilable matter which has been used up by the activity of the toxin.

¹ Thus, as Wassermann has demonstrated, if a fresh neutral tetanus toxin-antitoxin mixture be introduced subcutaneously, symptoms of the disease will, nevertheless, manifest themselves if the animal has, beforehand, been treated with adrenal extract. That, by contracting the arteries, prevents the absorption of the antitoxin by means of the circulation, and the toxin is still able to make its way to the higher centres along the nerves. If the two have been in contact for two hours previously, no symptoms show themselves.

CHAPTER VIII.

IMMUNIZATION AND IMMUNITY-(Continued).

IF diffusible enzymes and the diffusible products of bacterial growth cause reaction and, under certain conditions, the development of antibodies which neutralize their action, it becomes probable that other diffusible cell products, and more particularly those of an essentially organic nature and proteid type, will be found to have the same proporties. And this, as a matter of fact, has been found to be the case.

PRECIPITINS.

The first discovery of these bodies was, it is true, not the simplest example. If, as pointed out by Kraus,1 in 1897, one inoculates an animal with fluid cultures of typhoid, cholera, or plague bacilli, and then, after some days, gains some of the blood serum of that animal and adds a little to the germ-free culture fluid, a specific precipitate appears in the mixture. Slowly the fluid, which has been clear, becomes hazy, then turbid, then the suspended particles aggregate into fine flocculi, or, to employ a term employed in a parallel process, undergo agglutination; and next the flocculi fall to the bottom as a precipitate. The same results appear when the fluid of growth minus the microorganisms is used for inoculation. We know that it is the proteid substances present in the broth which cause the reaction, whether these become modified in particular directions, by the growth therein of the particular bacteria, and in part, perhaps, have been excreted by those organisms, or whether they are unaltered, for simple broth will bring about a like result. Through the inoculation of these constituents of the broth antibodies are developed and appear in the blood, which, combining with the bodies present in the culture fluid, cause a precipitate. Such antibodies, first known as *coagulins*, are now grouped together as *precipitins*, and of them a large number have been developed by treating (injecting) animals with protein-containing fluids of various orders, and these not merely of animal, but also of vegetable origin-milk, horse serum, eel serum (Bordet² and Tschistovitsch), egg albumin (Meyer), globulin from sheep and ox blood, peptones (Wassermann), human blood serum (Uhlenhuth), albuminous urine, pleural exudate, etc. (Mertens, Leclainche and

¹ Wiener klin, Woch., 1897: 736.

² Ann. de l'Inst. Pasteur, 13: 1899. A full study of the literature is afforded by Nuttall, Journal of Hygiene, 1: 1901: 367, and 60 the whole subject of blood precipitation, in his Blood Immunity, and Blood Relationship, Cambridge, 1904.

PRECIPITINS

Vallé, and others), muscle albumin (Schütze), serum albumin, pseudoglobulin (Ide), casein, and milk albumin (Hamburger) vegetable albumins (Schutz), wheat, rye, and barley albumoses (Kowarski).

And these are to a large extent specific; thus, to give simple examples, the blood serum of an animal treated with wheat albumose will cause a precipitate in a solution of wheat albumose, but none in solutions of rye or barley albumoses, and, as Bordet showed, the serum of rabbits treated with cow's milk will cause a precipitate of cow's, but not in goat's milk,¹ and several others have shown that the serum of rabbits injected with human blood will cause a heavy precipitate in human blood serum, but none in goat's or dog's blood serum. Here, indeed, we possess an excellent medicolegal method for detecting the presence of human blood. Certain precautions, however, as pointed out by Nuttall, have to be taken into account. The specificity, as, indeed, might be expected, is not complete; the more nearly allied two animals are the greater the probability that their proteins of one or other order will be of closely similar constitution and lead to the production of allied antibodies, and this is found to be the case. Grünbaum has shown that the precipitins developed by injecting into other species of animals the blood of the gorilla, chimpanzee, and ourang-outang will each of them cause precipitates in the blood serum of all three species of ape and in that of man, although they are without effect upon the blood sera of lower animals. Conversely, as indicated by Nuttall, if a given precipitin leads to precipitation in a series of bloods from different animals, it is an evidence that they are "of the same blood" and genetically closely allied, and these data of Grünbaum are to be accepted as new and convincing evidence of the cousinship between man and the higher apes. A like common reaction is found in the blood sera of different species of birds, amphibia, reptiles, etc.

1

y

T

8

18

p

While this is the case, it has to be emphasized that the *precipitation* is most marked with the homologous serum, i. e., with the serum of the species against which the proof animal was inoculated, and if this precipitin-containing serum be diluted, a point will be reached at which it will still react in a given time with the homologous serum, but not with others. And, indeed, as showing that there are even differences in the constitution of the proteins of individuals of the same species it has been repeatedly noted that the reaction is most complete with the blood or other protein-containing fluid of the *individual* animal that has afforded the material for inoculating the proof animal.

We see, also, another order of allied phenomena, indicating that the different proteins of the one individual have certain constituents in common, so that the precipitin developed by inoculating the one protein is capable of associating itself with, or acting upon, other proteins, although, again, not to the same extent as it acts upon its specific protein. Thus Besredka has found that the serum gained by injecting

¹ As pointed out by Moro, Wiener klin. Woch., 1901: 1074, the specificity here is not complete.

animals with *cell-free blood serum* will, added to blood, cause *hemolysis*, *i. e.*, will act on erythrocytes, liberating their hemoglobin. So, also, the serum of rabbits, injected with human muscle albumin, is intensely hemolytic. And, we may add, this action upon the red corpuscles is extremely common on the part of the sera of animals injected with cells of various orders.

The Nature of the Precipitate.-When we come to inquire what are these precipitins, and what the substances precipitated by them, we meet with difficulties. We know so little regarding the inner constitution of protein bodies. Thus, to take the latter question first, it is generally held that the *precipitable substance* is the main proteid constituent of the fluid used for purposes of injection; the resulting precipitate is so large-the amount of casein precipitated, for instance, when milk has been used to develop the precipitin, is very considerable; and when we use what we regard as a simple protein, egg albumin, or globulin from the blood, as antigen, the precipitin acts specifically upon that substance. Thus, what we may term the commonsense view is that these particular proteins are specifically acted upon. The critical, or hypercritical, view is that we know next to nothing about the constitution of proteins, and that there may be an intermediary tertium quid produced which, in its fall, brings down with it the bulk of the protein.

Jacobi instances as a parallel the case of ricin, which, added to blood, precipitates the corpuscles, so that the fluid becomes quite clear, the ricin having no action upon the hemoglobin, but upon the stroma of the red blood corpuscles. The illustration appears to us far-fetched; we fail to see how, under the circumstances, the ricin, if it acts upon the stroma of these corpuscular elements, could do anything else. Welsh and Chapman's' argument is more weighty: that it is not the test protein that supplies the main bulk of the precipitate, but the antiserum. Quantities of the test protein many times too minute to yield an appreciable precipitum with ordinary proteid precipitants may yield distinct precipitates with the specific antiserum.

Let us admit that in the fall of the main protein other substances, including other proteid bodies, are apt also to be brought down;² whether linked to the precipitable substance, or directly acted upon by the precipitin, or as a purely mechanical matter, we cannot, in most instances, state with sureness. The sensible view, undoubtedly, is that the precipitin acts directly upon the main constituent of the precipitation. This view demands, it is true, that we admit the existence of an enormous number of proteid bodies; that, for example, the globulins in the blood of different species are distinct bodies, for otherwise the specific action of the precipitin on one particular serum becomes

⁴ Proc. Roy. Soc., Biol., 78: 1906; 297.

² Thus, Dehne and Hamburger have shown that a precipitin developed against horse serum will, if added to the serum of a horse immunized against toxins, bring down the antitoxins along with the serum proteids,

inexplicable. But this we are already prepared to admit; the widely different percentage composition in C,H,N,O, etc., gained by the various observers who have analyzed hemoglobin, has already shown us that this must be the case. What is true of hemoglobin is likely to be true of other proteins.

As regards the precipitins, these resemble, in properties, the other antibodies producible in blood serum; they can be precipitated by various reagents; can be isolated from the bulk of proteins present in that serum by fractional precipitation, *i. e.*, by adding successive increments of ammonium sulphate and filtering off the successive precipitates that show themselves, the precipitins being brought down along with certain globulins within certain narrow limits; can be redissolved along with these globulins, and so on. Here a like question presents itself: Are these particular euglobulins, or pseudoglobulins, the precipitin; in other words, is the precipitating property a function of the modified globulin (for the two cannot be separated), or are we to regard the precipitin as a distinct chemical entity, brought down at the same time as the globulin? The fact above noted, that the two cannot be separated, again renders the former the more direct and practical view, the one that should, provisionally, be accepted. But here, also, the indications are that we must admit the existence of a multiplicity of these modified globulins, and that even in the one blood serum more than one protein may be developed possessing these powers of precipitating.¹ These globulins appearing in the blood serum of the inoculated animals, it must always be kept in mind, appear there as the result of cellular reaction and activity, and, if different orders of cells coincidently take up the inoculated material, there is nothing improbable in their reactions varying and in their discharging into the blood bodies-precipitins-showing some variation in constitution.

t -

1-

k

r,

n

et

e

se

ıg

Lastly, we may note the existence or possibility of development of *precipitoids* corresponding in properties to the *toxoids*, and of *anti-precipitins*.

AGGLUTININS AND AGGLUTINATION.

Gruber and Durham² were the first to make a special study of the phenomenon of agglutination, which has previously been noted cursorily by Metchnikoff,³ Charrin and Roger,⁴ Pfeiffer,⁵ and several others who had made growths of bacteria in the serum of immunized animals. In June, 1896, Widal published his paper, in which he showed that, during the course of an attack of typhoid, the blood serum of the

² Münch, med. Woch., 43: 1896; 285.
 ³ Ann. de l'Inst. Pasteur, 5: 1891; 473,
 ⁴ Compt. rend. Soc. de Biol., 1889; 667.

⁵ Pfeiffer and Issaeff, Zeitsch. f. Hyg., 17: 1894: 355,

³ This has been conclusively demonstrated by Piek, Beitr, zur. Chem. Physiol. u. Pathol., 1:1901:351 and 445, and coincides with Durham's observations upon agglutinins, to be presently noted.

patient acquires the power of clumping typhoid bacilli to which it is added. Already, in March of the same year, Grünbaum, working under Gruber, had observed this phenomenon, and but for an unfortunate delay in the publication of his paper, for which he was not to blame, what is now known as the Widal reaction would have been known by his name. As it was Gruber had already, some monihs prior to Widal's publication, announced the general principles of the method. The test should, therefore, be known as the Gruber-Widal or the Grünbaum-Widal.

Agglutination consists in the aggregation into clumps of free bacteria suspended in physiological salt solution, or broth, when there is added to the same some of the homologous serum, *i. e.*, serum of an animal which has been inoculated with bacteria of the same species. With this clumping, the bacteria, if previously motile, become motionless. The reaction is obtained with almost all species of bacteria, although the extent of the reaction varies with different species. It obtains with non-motile as well as motile, with dead as well as with living bacteria. It obtains also where cells are used as antigens-washed¹ erythrocytes, leukocytes, and body cells of various orders. It is specific to this extent, that, with relatively high dilutions of the homologous serum, the specific antigen bodies alone exhibit elumping; non-specific, to the extent that, with more concentrated serum, bodies of allied species may also exhibit agglutination. Such specific agglutinins may be present in the blood serum, not merely during an infection, but for months and even years, after, though the statement is made that, unlike antitoxins, as a result of removal by bleeding they are not readily reproduced. The reaction may be followed under the microscope, but it is more easy to follow by the naked eye, by observation of the gradual flocculus formation and sedimentation, whether in pipettes, long tubes, or watchglasses. This flocculus formation is of the same order as that seen under the action of precipitins. Only with concentrated solution is it of rapid development; toward the upper limit of dilution it shows itself slowly. Most English and Continental observers recommend the use of the unaltered homologous serum, as permitting more exact dilution. For purposes of diagnosis at a distance, the dry method elaborated by our late colleague, Wyatt Johnston,2 after Widal had shown that the agglutinins are unaltered by drying, has, throughout North America, been found to give equally accurate results.

The Properties of Agglutinins.—Here it is not our purpose to enter into the fine details of the process of agglutination; these will be found carefully recorded by Ewing, Cabot, and other writers upon the blood and its properties. Our object is to note the main facts bearing on the agglutinins in their relationship to immunity, and to the

¹ Washed, in order to remove any accompanying serum and to prevent the development of a complicating and similar precipitin formation.

² Centralbl. f. Bakt., 21: 1897: 523; Brit. Med. Jour., 1896; ii: 1629, and Johnston and McTaggart, Montreal Medical Journal, 25: 1897; 709.

AGGLUTININS AND AGGLUTINATION

other bodies now under discussion. They are relatively highly resistant bodies; withstand drying for many months, even when freely exposed to the air; are little affected by light or putrefaction; the majority, in a moist condition, can be heated to 62° C. without loss of properties, although some (e. g., tuberculosis agglutinins) are rendered inactive at 56° C. Like the antitoxins, they may be present in normal serum. Thus, normal human serum has been found, when undiluted, to cause agglutination of many bacteria-B. coli, B. typhi, B. pyocyaneus, Spp. choleræ, and danubicus, and pyococci. Even when diluted thirty times, it may clump the typhoid bacilli; or, one hundred times, the Pyococcus aureus. And the same is true of animal sera. In general, foetal blood, and, indeed, that of healthy children, under seven years, has very feeble agglutinating powers, a fact which would suggest that processes of subinfection play some part in the development of these normal agglutinating powers. As shown by Stäubli,¹ the sixteenth part of an agar culture of the typhoid bacilli suffices, when injected, to cause the appearance of relatively abundant agglutinins, and J. McCrae,² in our laboratory, was able to obtain their production even when the bacilli were enclosed in celloidin capsules. Other observers have gained agglutining by feeding animals with the homologous bacteria, without setting up obvious infection. By inoculation of cultures, the agglutinating power of the serum can be so increased that, in dilutions of 1 in 1,000,000, Van der Veldt³ gained clumping of B. typhosus; and with 1 in 2,000,000, Durham⁴ was able to "clump" the B. coli.

In general, agglutinins show evidence of development in from three to six days after the first injection; after this they increase with considerable rapidity. McCrae found that after removal of the celloidin capsules they rapidly disappeared from the blood, and it may well be that the cases which have been noted, particularly with typhoid, in which they have been present a year or more after injection, are cases of latency of the bacilli within the organism, for the existence of "typhoid carriers" is now well established. Their mode of disappearance is not wholly understood. They have repeatedly been determined in the milk of nursing mothers, and have been detected in the tears. This would point to their excretion; there is a certain amount of evidence that they undergo destruction in the liver and spleen, but the observations upon their presence in the urine are very discordant.

As to where the agglutinins are formed observations are curiously conflicting. According to some authorities, of whom the latest is Meek, extracts of the leukocytes of animals injected with one or other microbe afford no agglutinins. Van Anden found in certain cases only, that the

³ Bull, de l'Acad. Roy, de Méd., Brussels, April, 1897.

⁴Lancet, (Lond.), 1898; i:15.

34

:0

10

16

pl-

¹ Centralbl. f. Bakt., Abt. 1:36: Nr. 2.

² Jour. of Exp. Med., 5; 1901; 635. The same has been noted by d'Espine and Mallet, Revue Méd. de la Suisse Romande, 1907.

spleen, bone-marrow, and lymph glands contained more agglutinins than did the blood, while others have determined that removal of the spleen has no effect in the production of these bodies, and several observers have throughout found less agglutinins in the organs than in the blood, while Figari and others have brought forward evidence to show that in the intravascular blood (plasma) agglutinins are wholly absent, only appearing with coagulation and in the serum, and yet others have suggested that in the circulating blood there occur free agglutinis, which gain full activity under the influence of the oxygen of the air. From this it will be seen that despite very numerous observations, we are still in complete ignorance.

Group Reactions.—Of late years, a very extensive literature has made its appearance upon the specificity of the agglutinins. With regard to Pfeiffer's phenomenon (p. 543), the Berlin school had demonstrated that this was remarkably specific, and at first it appeared that this was true with regard to agglutinins and the Gruber-Widal reaction. Very soon, however, it was found that a serum that would agglutinate the typhoid bacillus would agglutinate also the B. coli—to a less extent, it is true, but, nevertheless, definitely. At first this was ascribed to the existence in the serum employed of normal agglutinins, and the complication, it was thought, would be overcome by employing high dilutions. To this day there are those like Paltauf, who, save in connection with the B. coli group, would ascribe all aberrant results to the presence of these normal agglutinins. They base themselves upon the strong specificity of the reaction with the different allied species of spirilla and the clear results gained with the plague bacillus.

But when one has personally dealt with sera which will agglutinate two distinct species when diluted in the one case to 1 in 1000, in the other to 1 in 3000,⁴ dilutions far beyond the limits of normal agglutinins, it is impossible to accept this explanation. The conclusion is inevitable that both the typhoid and the colon bacilli and their allies exhibit what are termed group reactions.

The most satisfactory explanation of this "group agglutination" has been afforded by Durham,² in his study of the B, enteritidis. According to him, a given bacillus introduced into the system may cause the development not of one, but of several different agglutinins, and some of these may, likewise, be developed when bacteria of other species are introduced. We may regard the agglutinins as antibodies to the proteins of which the cell, bacterial or otherwise, is composed, and as these proteins are multiple, and some of them only specific to the particular cell, so some of the antibodies will be specific, others common to several orders of cells, or bacteria. If bacillus X possesses the agglutinogens a, b, c, d, e, stimulating the production of agglutinins A, B, C, D, E, and bacillus Y has similarly agglutinogens d, e, f, g, h, causing the production of agglutinins D, E, F, G, H, then, if bacillus X be introduced into

¹ Adami and Chopin, Jour. of Med. Research, N. S., 6: 1904: 469.

³ Brit, Med. Jour., 1900, and Jour. of Exp. Med., 5; 1901; 353.

its homologous serum, there will be the strongest reaction, but bacillus Y will also show a reaction with the same serum, although not so perfect, for although agglutinins A, B, C are unable to influence it, agglutinins D and E can.

We arrive, thus, at the conception of "idio-agglutinins," or specific agglutinins proper, developed by each species, and common, or group agglutinins which are able to act upon more than one species. In short, agglutinins come into line with what has been noted regarding precipitins.

Relationship of the Agglutinins.—As will be pointed out shortly, the injection of bacterial or cells into the body of an animal leads to the production of bacterialysins or cytolysins—of antibodies which cause the dissolution of the particular antigen. Are agglutination and lytic action stages of one and the same process, or are they distinct-processes? They are distinct bodies. Both, it is true, are thermostable and both become attached to the bacterial substance, both occur in the blood serum, and are to be found there for some time after infection, but as shown by Pfeiffer and Kolle,¹ bacteriolysis may occur without a sign of agglutination, and after inducing immunity, the blood serum, months later, may still be strongly bactericidal when it has completely lost the power of agglutinating.

There is, further, as well shown by Evans,² no relationship between the agglutinative and protective powers of a given blood; the blood of a typhoid convalescent agglutinated at 1 to 500 had a bactericidal value of 5 units, whereas another, which only agglutinated at 1 to 20, had a bactericidal value of 500,000 units. And lastly, as pointed out by Ehrlich and Morgenroth, bacteriolytic action is arrested by subjecting the serum to a heat of 56° C, whereas agglutinins are unaffected by this temperature. It will be shown later that for bacteriolysis and evolysis the interaction of two bodies is necessary amboceptor and complement—and that a heat of 56° C, destroys the latter. Nor, although they have many features in common, are the agglutinition can occur without any sign of precipitation.

The Nature of the Agglutination Process. —Of the many views regarding the nature of the agglutination process, that of Bordel is the most acceptable. As the result of the action of the agglutinins, he postulates an alteration in the molecular attraction or tension between the bacteria and the fluid medium. In the first phase of the process there is a junction of the agglutinis with the constituents of the bacterial cell; the second is purely physical, the salts in the medium playing a part in the process. It is, in short, a process of the same nature as the gathering of red corpuscles into rouleaux; in fact, a series of observations upon the agglutination of red corpuscles and other cells has been made which in many respects parallel those observed in connection with bacteria. As Sir Lauder Brunton³ has shown, if matches (to represent bacilli) or

¹ Centralbl. f. Bakt., 20: 1896; 129,
 ³ Ibid., 7; 1900; 53,

1-

0-

11.

b.

² Jour. of Path., 9: 1903: 42.

disks of cork (to represent erythrocytes) be covered with hard soap and thrown into water, they float about free and isolated until that water is slightly acidulated, when they immediately draw into clumps. Render the water faintly alkaline, and the clumps, if broken up, will not form again. More recently, Albrecht' has demonstrated that red corpuscles, as a matter of fact, possess such a fatty (lecithin) containing surface layer as is demanded in Brunton's experiment, and, in ignorance of the latter, he comes to a like conclusion regarding the mechanism of agglutination. Through alteration in the physical condition of the environment he concludes that the surface of corpuscles (and bacteria) becomes so modified as to lead to the physical attraction and adhesion of the bodies. Whatever the process, it would seem to be of the same order as that occurring under the action of precipitins, namely, a change in the colloidal state favoring floccular formation.

IMMUNIZATION AGAINST CELLS. CYTOLYSIS AND THE CYTOLYSINS.

It has long been known that the inoculation of a foreign blood is likely to set up grave if not fatal disturbances, and even the inoculation of the blood of another animal of the same species has been found so dangerous that, recommended in man for a time in cases of grave wasting disease, experience has led to its being almost wholly given up. Experimentally we find that it may lead to dissolution of the red corpuscles of the host and to intravascular coagulation. We owe to Bordet and his brilliant observations a fuller knowledge of this process of hemolysis or solution of the corpuscles, observations that can easily be confirmed and that have led to abundant studies upon the destruction of cells of other nature by the organism; so that now there is a rich literature upon the wider subject of cytolysis, and incidentally not a little light has been thrown upon the subject of cell destruction in general within the organism. Bordet showed that if an animal, A, be inoculated with successive small amounts of the blood corpuscles of an animal of another species, B, within a few days (not immediately), upon bleeding A and obtaining some of its blood serum, that serum added to the blood or blood corpuscles of B, outside the body, will cause the dissolution of B's red corpuscles, so that the fluid becomes "laked," or, if inoculated into the vessels of any animal of B's species, will cause an extensive intravascular destruction of the erythrocytes. The inoculation of the corpuscles leads to the appearance in the serum of a cytotoxin, or cytolysin.

In rapid succession, by a long series of observers, similar cytolysins or cytotoxins were demonstrated as being developed when cells of various organs were injected into an animal of another species. From their free condition spermatozoa soon suggested themselves for such experiments, and Metchnikoff and Landsteiner independently demonstrated

¹ Verhandl. d. Deutsch. pathol. Gesellsch., 5: 1903: 7.

IMMUNIZATION AGAINST CELLS

the existence of *spermatoxins* (the spermatozoa of bulls being injected into rabbits). Leukocytes were similarly suitable objects, and Metchnikoff, by inoculating polynuclear and mononuclear leukocytes respectively, gained cytotoxins acting specifically on one or other form. These *leukotoxins* are also known as *leukocidins*. Ciliated epithelium was shown to have its cytotoxin (*trichotoxin*), as have kidney cells (*nephrotoxin*), liver cells (*hepatotoxin*), pancreatic, adrenal, and, in fact, every form of animal cell that has been tested. Not only do the sera so gained act on the particular form of cell in the test-tube, but inoculated into the vascular system of animals of the species affording the original cells, the subjects of cytolysis, the cytotoxins act preëminently on the organs containing these particular cells, setting up grave degenerations of the same. Charrin has gone so far as to demonstrate that a hepatotoxin acting specifically upon the liver cells of the goat will pass through the placenta and lead to degeneration and atrophy of the liver of the feetal kid.

The exhibition of such cytotoxins is most effective when an animal is inoculated with the cells of a widely different species, but, as shown by Ehrlich and Morgenroth, if the red corpuscles of one goat be injected into another, its serum is capable of laking the blood of other goats, a fact which might perhaps be expected from the clinical observation that the serum of one patient (even without inoculation) may lake the blood of another. There are thus *isolysins* as well as *heterolysins* (lysins active in another species). But it is impossible to develop experimentally autolysins, that is, to bring about cell destruction by any method of reintroducing into the organism the separated cells of the one individual, a fact which is parallel perhaps to what was noted regarding the impossibility of gaining anti-enzymes to the common enzymes of the organisms. There are, however, indications that in disease, as, for example, in cases of cancerous and other new-growths, a form of autolysis manifetss itself, reaction being set up not by the products of normal cell activity, but by the abnormal disintegration products of the tumor cells.

01

Dr-

he

·11-

Also, it has to be observed that while these cytotoxins are specific to the extent that they act most powerfully upon the one particular cell form through which they were derived, they are liable to have some action on other forms of cells. This is not wholly surprising when we remember that all the cells of the organism have a common origin, and are likely thus to have certain constituents in common. Particularly the liability of cytotoxins in general to induce hemolysis has been often noted.

Here, again, by the cautious inoculation of an animal of the susceptible species with progressive small doses of cytolysins, *anticytolysins* can be developed and serum obtained which will neutralize the action of the cytolytic serum.

Bacteriolysins.—We have until now been silent regarding what is quite the most important of this series of cytolysins. Just as inoculation of *animal* cells leads to the production of bodies causing the destruction of those cells, so has the inoculation of *vegetable* cells a like result;

bacteriolysis and the *bactericidal* activity of the blood serum is of the same order as the cytolysis here described.

The existence of this bacteriolysis was known years before the other cases; but although Bordet's observations upon the cholera spirillum demonstrated the nature of the process, it is through his later observations upon hemolysis and through the researches of Ehrlich and his pupils on cytolytic phenomena in general that we have gained our grasp —such as it is—of the phenomena of bacteriolysis in particular. We believe that the student will gain a clearer understanding, if we detail first the main facts regarding the cytolysis of animal cells and then pass on to bacteriolysis.

The Mechanism of Gytolysis.—'To repeat Bordet's fundamental experiment: The serum of guinea-pigs' blood has normally very little effect upon rabbits' blood and blood corpuscles; but if a guinea-pig has injected into it rabbits' blood corpuscles its serum becomes in a few days extremely active (such serum is termed *immune*). If now a suspension of rabbits' red corpuscles be taken and have a little of this serum added to it, there results extensive dissolution of the corpuscles, with escape of the hemoglobin and "laking." So as to obviate any complicating appearance of precipitins, we now inoculate the centrifugalized and *washed* red corpuscles and use a suspension of such washed corpuscles for the test.

But if we warm the G. P. serum to 55° to 60° C., the hemolytic action is wholly arrested. It is usual to speak of such warmed serum as *inactivated serum*.

Now to this mixture of washed corpuscles and inactivated serum add a little blood serum from a normal, untreated guinea-pig or rabbit, and immediately the hemolysis takes place.

Or, briefly:

Normal non-immuneG , P, serum + washed R, erythroeytes.		No hemolysis
Active (unheated) immune G. P. serum + washed R. erythrocytes		Hemolysis
Inactivated (heated) immune G. P. serum + washed erythrocytes		No hemolysis
Inactivated immune G. P. serum)		
Normal G. P. or R. serum and	-	Hemolysis
Washed R. erythrocytes		

It is obvious that heating the treated guinea-pigs' serum has destroyed something which is restored by adding normal blood serum, or otherwise that there is something present both in unheated immunized guineapigs' serum, and in normal guinea-pigs' serum which is a necessary factor in the process of hemolysis.

But this is not the only factor, for non-immune G. P. serum is without effect in the corpuseles. There must, therefore, be a second body present and developed in the serum of the immunized guinea-pig which is equally essential; and it is the combined action of these two which leads to the hemolysis.

The existence and combined action of these two factors can similarly be demonstrated in every case of cytolysis.

We speak of the body developed in the serum of the immunized animal

CYTOLYSIS AND THE CYTOLYSINS

as the *immune body*. We shall also refer to it, for reasons presently to be given, as the *intermediate body* or *amboceptor*. The body present both in the normal and the immune serum, which is indeed a normal constituent of all healthy sera, is, most commonly, referred to as the *complement*. The term is not wholly fortunate, for while admittedly this body is necessary for the completion of the process, the term tends to suggest that it is accessory rather than, as we now regard it, the essential agent in the cell destruction. Many—altogether too many—alternative names have been given to these two bodies, to recite which at this point would only cause confusion; we will tabulate them at a later period.

We can demonstrate that both are present in the cytolytic serum in the following manner. Take the cytolytic serum in two parts:

1. Heat the one part to 55° to 60° C. This, as already noted, destroys the complement and leaves the immune body unaltered, that being only destroyed by a heat of 70° C. or over.

2. Cool the cytolytic serum to 0° C.; add to it the washed cells, upon which it has a specific action, these having also been carefully

cooled down to 0° C. At this temperature cytolytic action is arrested. But, while this is the case, as first demonstrated by Ehrlich and Morgenroth, the immune body attaches itself to the washed cells, and now, by carefully filtering it at zero, a complement-containing serum can be gained free from the immune body. And if now the cells in the filter be washed so as to remove all traces of the serum with contained complement, the cells may be suspended in physiological salt solution and brought to room temperature without showing the least trace of cytolysis. But, if the filtrate of (2) be now added, cytolysis rapidly ensues; or, if again, this filtrate be cautiously added to (1) and to the mixture the particular cells be added, then the reaction occurs, although, in this case, it is not Through the action of heat, as always perfect. we now know, the complement may be converted into complementoids, which combine with the im-

a-

lly

he

rly



Combination of cell a, amboceptor b, and complement c. The amboceptor may unite with the cell, but cannot affect it alone. The complement cannot unite with the cell except through the amboceptor, having no adaptation to the cell directly.

mune body, and while unable to disintegrate the cells, nevertheless, to a greater or less extent, prevent the active complement added later from combining and completing the reaction. In general, however, as shown by Sachs, the complement has a greater affinity to the immune body than have these complementoids.

These observations—and they have been repeatedly confirmed indicate very clearly that for cytolysis to occur there must first be combination between the immune body and the cell, and that then the additional combination of the complement brings about the cell disintegration, the immune body alone having no disintegrating action. They do not, it is true, exclude the possibility of the immune body and complement being in a state of loose combination in the serum, but even if

so, it must be through the immune body that such a combination attaches itself to or acts upon the cell. It is for this reason that the immune body is also referred to as the *intermediate* body, or *amboceptor*,¹ it being regarded as capable of a double attachment, seizing on to both cell and complement. That such combination between complement and immune body actually takes place has been shown by Preston Kyes in his observations upon cobra poison, to which we shall refer later, in which he showed clearly that lecithin acts as a complement, the cobra poison acting as the immune body in the destruction of the red corpuscles, and poison and lecithin becoming combined to form a most active and rapid hemolytic agent. The existence of such compounds strongly supports Ehrlich's view of the intermediary nature of the immune body, as against Bordet's that the immune body first acts directly on the cell and then, also directly, the complement.

The Immune or Intermediate Body (Amboceptor).—There have been very numerous observations made upon these two constituents, immune body and complement. The more important results must here be indicated, for largely through methods introduced by Ehrlich, the reactions obtainable approach the procedures of the chemist in their exactitude. We are dealing with bodies having well-defined properties; a precise amount of the complement-containing normal serum must be added to a particular quantity of the inactivated serum (containing immune body) to act completely on a given mass of cells; anything more or less leads to an imperfect reaction.

Multiplicity of Amboceptors.—Regarding the properties of immune bodies, it must in the first place be recognized that these are multiple. It is found, for instance, that goats' blood serum will dissolve both guinea-pigs' and rabbits' red corpuscles. If the proper amount be taken to hemolyze G. P. corpuscles, the serum still contains amboceptors capable of attaching themselves to rabbits' corpuscles and (with the complement) causing their dissolution, or otherwise, the goats' serum contains one set of immune bodies having immunity for the guinea-pigs' corpuscles, another for the rabbits' (Ehrlich). In like manner, Neisser has shown that when a serum is at the same time hemolytic and bacterriolytic for certain bacteria, the bacteriolytic power may be removed by letting it act upon the bacteria, and it still retains its hemolytic powers. We shall show that the same is true for the immune bodies or amboceptors which are developed to act against specific bacteria. We have to acknowledge then a pronounced multiplicity of immune bodies.

Multiplicity of Receptors.—But, as a corollary, it must be equally admitted that the cells have multiple affinities, or, as Ehrlich terms them, receptors. The very fact that red corpuscles are capable of being acted upon by so large a group of diverse substances as the phytotoxins (ricin, etc.), snake venom, spider and scorpion venom, bacterial products, and hemolysins proper can only be explained on the supposition that the corpuscles have manifold affinities. The alternative that there is some

Ambo, both; capio, I seize.

MULTIPLICITY OF COMPLEMENTS

common atom group in all these different lysins having an affinity for a special atom group in the red corpuscles is shown to be wrong, or at least inadequate to explain the whole series, because, as regards hemolysis alone, the same hemolytic serum will not act on the red corpuscles of all members of one species. Inoculate a goat with the corpuscles of another goat, and the serum developed will not hemolyze the corpuscles of all goats indifferently; it will act on some specimens, but not on others. As Aschoff expresses it, employing Durham's illustration regarding agglutination, if we regard the goats' corpuscles as capable of possessing a possible full series of receptors a, b, c, d, e, f, then if we treat a goat with corpuscles possessing only the receptors a, b, c, its serum will come to contain amboceptors for a, b, c, and not for d, e, f. Such serum coming into action with the goats' corpuscles possessing receptors a, b, c, will actively destroy them; possessing only a and c, will destroy them, but not so actively; possessing receptors d, e, f, will have no action.

While this is the case, it is also evident that certain amboceptors appearing in the serum of different species, if not throughout identical, may, nevertheless, be so closely allied structurally that, as regards any particular reaction, they may replace each other, their molecular constitution, in certain respects, causing them to have like affinities.

e

e

e

e

le

Amboceptoids.—Whether amboceptors can undergo modification; whether, for example, bodies can be developed which will combine with the complement but not with the cell, is still a matter of some debate. Weehsberg,' it may be, encountered such amboceptoids in dogs' blood, in which he gained "Complementablenkung" (diversion of complement), but found that there was no action on the cell. In Ehrlich's terminology the body or bodies in question possessed a complementophile group, but no evtophile.

Anti-amboceptors.—Regarding these, *i. e.*, the production of antibodies by inoculating a third animal with the immune bodies developed in the second, it has to be noted that so far they have not been surely determined in connection with hemolysins. Pfeiffer and Friedberger,² Bordet,³ and Ehrlich and Sachs have, however, produced these against other cytolytic agents. The first of these produced them by inoculating cholera immune serum into animals, and made the interesting observation that the anti-immune body so developed hindered the action of typhoid immune serum also. On the other hand, the anti-immune bodies developed from the amboceptors produced in immunity against the different snake venons are strictly specific in their action.

Seat of Development of Immune Bodies.—This we shall discuss when dealing later with the origin of antibodies in general (see also p. 514).

Complements.—There has been and continues to be discussion as to whether in a given blood one or more complements exist. It will

¹ Wiener klin. Woch., 15: 1902: 337.

² Centralbl. f. Bakt., 34: 1903 and 37; 1904.

³ Ann. de l'Inst. Pasteur, 13: 1904.

be seen that the evidence is in favor of there being a *multiplicity of complements*. Bordet more particularly has championed the unitarian theory. A given complement-containing serum, he showed, will, when added to inactivated immune hemolytic serum, activate it and cause hemolysis. Similarly, added to inactivated immune bacteriolytic serum, it causes bacteriolysis. But if now such complement-containing serum, after acting upon the blood corpuscles, be tested with inactivated bacteriolytic serum it has no effect, and vice versa. This, he concluded, indicated the existence of one complement active in both processes, and used up in the first. And Kyes' observations upon the hemolytic action of snake venom appear at first and up to a certain point to favor this contention. Kyes found that if a watery solution of cobra poison be



Schema of neutralizing netion of A, anti-ambegencor, and B, anti-complements, respectively. In A, the amboceptor cannot combine with the redureceptor b because of the junction of the anti-amboceptor aa. In B, the amboceptor aa muite with the cell receptor, but cannot be netivated because of the junction of the anticomplement ae with the complement. (Mitre Levalit). shaken up with a solution of lecithin in ether, the neurotoxic element in the venom remained in the watery solution; the hemolytic combined with the lecithin and could now be gained as a definite compound of lecithin-a lecithide insoluble in ether and so distinct from lecithin proper and possessing intense hemolytic powers. He showed, further, that in all respects such lecithin acts as a complement, and that not merely for cobra venom, but for other snake poisons; that where cobra poison injected alone causes hemolysis, the action is to be explained by the preëxistence of lecithin as a constituent of the corpuscles; that the mere existence of lecithin is not sufficient; all ervthrocytes contain lecithin, but not all are laked by cobra poison; therefore the lecithin, to act, must be in a free or disposable state.

To this extent lecithin, a body capable of crystallization, and therefore a definite chemical compound, is a complement common for a large number, at least, of hemo-

lytic snake venoms. But, if this be so, everything indicates that lecithin is able to form a large number of compounds, or, more exactly, to become attached to and an intimate portion of various cell substances. In the organism, at least, the lecithin may be very variously combined; and we are justified in supposing that under intravascular conditions the complement action of lecithin is not necessarily exerted by it as a free substance; but that, combined, it is capable of uniting the cobra poison, *provided* that in the combination the particular affinity for the poison is unsatisfied.

A like chain of reasoning would seem to harmonize Bordet's obser-

MULTIPLICITY OF COMPLEMENTS

vations with those we are about to note.1 If, as shown by Ehrlich and Morgenroth, goats' or horses' blood serum be filtered (through Pukall's filter), two complements are obtained, the one passing through with difficulty, the other under certain conditions coming through the filter alone: the former acting upon the immune bodies from rabbits' blood. the latter on those of guinea-pigs' blood. Neisser and Dohring were able to make a similar separation in human blood serum. Further, it has been noted that in some cases the complement is thermostable, in others From these and other observations, Ehrlich and thermolabile. Morgenroth concluded that (1) in every normal serum there exists a series of complements, and again that (2) in different animals there exist a certain number of identical complements either absolutely identical or identical so far as regards their haptophore groups (i, e_i) having identical affinities toward the cell, but not being throughout of the same composition). As above suggested, one common type of substance may be the basis of complements of all orders, but the modifications and accretions this gains in different species and different individuals may determine whether there be association with particular immune bodies or not. As Aschoff points out, this variation in the properties of the complements has a practical bearing upon the relative benefit gained by different individuals from the injection of protective sera.

Variation in Amount of Complement.—And, we may add, the amount of complement present is also a factor. There are numerous observations indicating that, in the course of disease and by experimental methods (Abbott and others), the amount may become greatly reduced, and that through their relative absence protective sera may fail to antagonize the bacteria.² On the other hand, it may be increased by the injection of indifferent substances, blood plasma, broth, etc., and such increase may, in part, explain the favorable results of Issaeff's method.

Complementoids and Anticomplements.—If a serum which has been heated up to the point at which it is inactivated and the complement as such destroyed, the existence in it of complementids is demonstrated from the fact that anticomplements become developed in the serum of the inoculated animal, just as toxoids (p. 506) will induce antitoxin formation. Such anticomplements when added to an active serum arrest its activity.

Structure of the Complement.—The existence of these complementoids, together with the considerations previously detailed, would indicate that the complement is formed of two essential parts, the haptophoric portion, whereby it attaches itself to the immune body; the toxophoric or symo-

¹Jacobi, loc. cit., p. 68, brings together other convincing proof in favor of the plurivalent hypothesis, notably Ehrlich and Marshall's observations upon an anticomplement found present by chance in a specimen of human ascites fluid and its capacity to inhibit the lytic complement in some cases and not in others.

² Vide Ehrlieh and Morgenroth, Berl, klin, Woch., 31: 1900 (phosphorus poisoning); Metchnikoff (chronic suppurations), Ann. de l'Inst. Pasteur, 14: 1900; 577; Bentivegua and Corini (hunger), Lo Sperimentale, 5: 1900; 400.

phoric or cytotoxic, which is the essential agent in bringing about cell destruction. This latter may be destroyed or modified (complementoid), whereby, although the altered complement combines with the immune body, no cytolytic results ensue, and active complement is prevented from uniting. A like absence of the cytotoxic moiety must be predicated for the anticomplement. The former also may be modified so that the complement is unable to join on to one or other amboceptor.

On the Nature of Amboeptor and Complement.—While it is useful to form an hypothesis regarding the nature of the process of cytolysis, and by means of Ehrlich's diagrams to visualize what we imagine to be the steps of that process, we must keep in mind that essentially we are dealing with chemical reactions, and that until we have established the nature of those reactions we are upon insecure ground. Have we, it may be asked, any definite chemical data which will help us to understand these reactions? We deal, in the first place, with colloid bodies, and admittedly are but at the beginning of a knowledge of the chemistry of colloids. Nevertheless, these are data which, to say the least, are most suggestive.

The first step along the road was afforded by Preston Kyes when he demonstrated that in the hemolytic action of cobra poison a constituent of the venom acts as amboceptor, while lecithin acts as complement. There have, it is true, been discussions regarding Kyes' cobra lecithide. What we know as lecithin is a notably unstable body; it is practically impossible to obtain it in a pure state, there being always fatty acids (oleic) and soaps present, the result of dissociation. We shall revert to this instability later, here merely stating that the very bulk of the "lecithide" obtainable by Kyes' method strongly favors the view that it is a phosphatide compound, while the fact that it is hemolytic after treatment with ether, in which it is insoluble, indicates that the action is not due to any free fatty acid, for that would be removed by the ether.

The next important observations were those of Noguchi.¹ Making alcoholic extracts of blood and various tissues, and then treating these with ether, he obtained a precipitate soluble in water, having, in short, the properties of soaps. Dissolved in normal salt solution these soaps are strongly hemolytic. He next turned his attention to the salts of the fatty acids, and found that all the soluble soaps, but notably those of oleic acid, have the same properties. He found, further, that, like his extracted soap, these pure soluble oleates mixed with serum are (1) rendered inactive by heating to 56° C, for half an hour; (2) they are inactive also at 0° C.; (3) acids and alkalies render them inactive: (4) the addition of yeast, kidney, and other cells also render them inactive. These serum-soap compounds, in short, are, as Noguchi describes them, "artificial complements." Independently, von Liebermann,² studying the nature of the hemolysis induced by ricin, came to practically identical conclusions. His observations led him to conclude that the complements of the blood are to be sought for in the soaps of the serum. that these soaps are inactivated through the serum albumin present, and

¹ Proc. Soc. Exp. Biol. and Med., 4: 1907: 107, and Biochem. Ztschr., 6: 1907.

² Biochem, Ztschr., 4: 1907, and (with Fenyvessy) ibid., 5: 1907.

THE CHEMICAL NATURE OF COMPLEMENTS

that the action of the immune bodies is, in the first place, to free the soaps from their combination with serum albumin, when they become the active hemolytic agents. Ricin, for example, he found to possess an acid constituent having (as he held) this capacity, and, proceeding farther, he found that oleic acid might act as an immune body, or amboceptor. Thus, a little oleic acid added to an inactive soap-protein combination, rendered it actively hemolytic. A mixture of sodium oleate (0.5 gram) and serum albumin (1.2 grams) is inactive as a hemolytic agent. Add oleic acid (0.5 gram), and it is rendered actively hemolytic. He concludes that in such a mixture the oleic acid plays the part of amboceptor, the albumin and soap compound that of complement. He and Fenyvessy have shown that such a mixture of soap, serum, and oleic acid exhibits the Ehrlich-Morgenroth phenomenon. Brought into contact with washed erythrocytes at 0° C., no hemolysis occurs, but the oleic acid becomes fixed to the corpuscles and removed from the mixture. And now both the corpuscles and the separated fluid are inactive, although when brought together at body temperature active hemolysis occurs.

The resemblance between these reactions, between the soap-serum combinations and complements, and oleic acid and amboceptor is certainly very striking, and most suggestive. They suggest, for example, that the amboceptor-complement action is much more than a mere linkage, that the amboceptor, on the one hand, has an affinity to certain constituents of the cell or bacterial body, on the other, acts upon the complement separating its constituents, separation enabling one of these constituents to become active in dissociating constituents of the cell body; that thus it is not the mere act of linkage of the complement that causes hemolysis, evtolysis, etc., but the dissociation of the complement and liberation from it of substances which actively combine with and dissociate the cell substance. At last thus we obtain a reasonable conception of how linkage of amboceptor and complement to the cell molecule may bring about cell disintegration. To harmonize with Kyes' data we must hold that in the action of cobra lecithide, the venom moiety of the compound acting as immune body has affinities which allow the compound to be taken into the erythrocytes; that so soon as this becomes fixed in the cell the lecithin moiety (like the albumin soap) undergoes dissociation, giving rise similarly to the oleic acid soap¹ which is the active agent in the hemolysis.

We must, however, freely admit that several objections have been brought against these observations of Noguchi and von Liebermann. While they fit in well with the admitted paucity of complemental bodies, it is difficult to harmonize them with the known abundance of specific immune bodies; those immune bodies as a class cannot be simple lipoids like oleic acid; they must in nature be much more complex. It

¹ Possibly a choline oleate, which we have found may be formed when choline and oleie acid (decomposition products of lecithin) are brought together (Adami and Aschoff, Proc. Roy. Soc., June, 1906).

has been noted that, like bile salts, these artificial hemolysins completely dissolve the corpuscles, whereas ordinary hemolysins simply dissolve out the hemoglobin, leaving the stroma undissolved. Nevertheless, the points of resemblance between the properties of these artificial complements and the natural are so many and so striking that we look for very material advance during the next few years in our knowledge of the chemistry of immunity along these or parallel lines.¹

BACTERIOLYSIS AND BACTERIOLYSINS.

While the essential data bearing upon the destruction of bacteria and the production of bacteriolysins are identical with those of cytolysis in general, it has seemed better to treat the subject separately, so as to present it to the reader in a clearer light when he is more fully prepared to grasp the main details.

Already, thirty years ago, Traube² concluded that the blood was able to destroy bacteria. In 1881 Lister noted that extravascular blood kept sweet despite the addition of small amounts of putrefying material, i. e., that within certain limits it arrests the activities of putrefactive microbes; in 1884 Grohmann³ published confirmatory results; v. Fodor⁴ followed the course of the destruction in intravascular blood. noting a preliminary destruction followed by increased proliferation: facts which were confirmed by Flügge and carried vet farther by Nuttall⁵ working in his laboratory, who found that the destruction could occur in fluids containing few leukocytes such as the aqueous humor and pericardial fluid, and that so the destruction was not, as Metchnikoff had laid down, essentially intracellular. He made the further important observation that a heat of 56° C, destroyed the bactericidal activity of sera and other body fluids. From these observations we pass to the more definite studies of Hankins and of Buchner and his school upon the nature of the essential bactericidal substance-Buchner's alexine"-

⁴ Although this does not directly bear upon complements and amboecptors, the chemical observations made with regard to the inhibiting action of cholesterin may here be noted.

Ransom has observed that this body inhibits hemolysis by suponin; Sachs and Kyes, that it inhibits the activation of cobra venom by lecithin. Preti has noted that while lecithin increases the hemolytic activity of extracts from the anchylostomum, cholesterin depresses it, and allied to these observations Reicher, who, with Morgenroth, had found that by feeding rabbits with cholesterin it was possible to arrest the anemia due to treatment with cobra lecithide, reports later that he has obtained favorable results from treating anemic patients with cholesterin (Berl. klin, Woch., 1907, No. 43). And lastly, Kurt Meyer has observed that the resistance of the red corpuscles to saponin hemolysis corresponds to the amount of cholesterin they contain compared with the lecithin.

² Jahr. d. Schles. Ges., 1874.

³ Inaug. Diss., 1884.

4 Deut. med. Woch., 1886; 617, and 1887; 745.

⁵ Zeitsch, f. Hygiene, 4:1888:253.

⁶There is an unfortunate and confusing tendency nowadays to employ this term as synonymous with complement.

while here must be noted the important observation of Vaughan, Novy, and McClintock,' confirmed by A. Kossel,' that the nuclei of leukocytes contain nucleic acid, which in itself is a definitely bactericidal substance. Here also must be mentioned, what we must discuss in more detail later, Metchnikoff's observations upon phagocytosis, the determination more particularly by Buchner and his pupils that leukocytes afforded the main source of the alexines found in the body fluids and the objections that have been raised to these conclusions.

The next great step forward was undoubtedly the observation by Pfeiffer that the cholera spirillum (and the same was quickly shown to be the case with the majority of pathogenic bacteria) contains what we now term *endotoxins* in contradistinction to the diffusible toxins of the diphtheria bacillus, and that immunity against these bacteria is, therefore, produced by means other than the neutralization of diffusible toxins, which had already been shown to occur by von Behring. These observations led to the demonstration by Pfeiffer and Bordet of the complex nature of the bactericidal process; in short, to the demonstration of the existence of complements in normal sera and the development of immune bodies or amboceptors in the immunized animal.

ia

le

od

te-

*0-

n:

115

DD

10.

he

Further, it was noted that forms which, like the diphtheria bacillus and the B. pyocyaneus, affords diffusible toxins contain at the same time endotoxins, and that the process leading to the destruction of such bacteria by the organism (as distinct from the neutralization of their toxins) is identical with that occurring in cytolysis.

Pfeiffer's Reaction.—The basal methods for studying bacteriolysis have been afforded by Pfeiffer and Bordet. Pfeiffer has described two methods of gaining his reaction which in principle are, however, identical.

 Take a guinea-pig that by successive inoculations has been rendered highly immune to virulent cholera spirilla and introduce into its peritoneal cavity five to ten times the ordinary fatal dose of an agar culture of the cholera spirillum.

 Or, inject into the normal guinea-pig a like dose of the spirillum mixed with an excess of cholera immune serum from another guinea-pig.

In either case, by removing with a pipette some of the peritoneal fluid from time to time, it is seen that the injected bacteria undergo destruction, and this apart from any phagocytosis and in a remarkable manner. They become motionless, swell, become rounded and like micrococci; therewith (more particularly in the peritoneum, not so clearly *in vitro*) they become progressively smaller, their substance undergoing solution, as Pfeiffer described it, like sugar in water. Radziewski³ has carefully followed the reaction with Sp. cholera, B. pyocyaneus, B. typhi, B. pneumonie, Streptococcus pyogenes, B. anthracis.

Metchnikoff and Bordet showed that the identical process occurred

¹ Medical News, 62:1893:536.

² Arch. f. Physiol., 1893: 164.

^a Zeit. f. Hygiene, 37:1901:1.

in vitro, and that the bacteriolysis could be brought about by taking definite proportions of bacilli, inactivated (heated) immune serum, and normal serum, the former containing the amboceptors, the latter the complement (although they use different terms for the two, which to avoid confusion we do not mention). Not to repeat ourselves unduly, we may sum up in brief sentences the main facts that have been ascertained regarding bacteriolysis and *mutatis mutandis* for all the different forms of evtolysis.

1. Bacteriolysis is brought about by the interaction of amboceptors and complements upon the bacterial body.

2. Antibodies, including both amboceptors and complements, may be found present in the blood of normal animals, and this not only in the serum, as Gengon has urged, but in the plasma (von Dungern and others). These are not, that is, entirely derived from the dissolution of leukocytes at the time of the removal of the blood. The amount of amboceptors is small, however, compared to what may be developed by specific inoculation.

 The amboceptors are multiple; an animal immunized against both cholera and typhoid provides a serum which after destroying the cholera spirilla will, added to typhoid bacilli suspended in normal serum, destroy these also.

 By immunization of animals against a specific microbe specific immune bodies are developed acting (specifically) upon the species of microorganisms employed for inoculation.

5. This specificity, while, as Pfeiffer has shown with the cholera spirilla and strains of the same, it may be very strongly marked and practically absolute, may in other cases be more diffuse; thus Loeffler and Abel⁴ found that typhoid immune scrum had a slight bacteriolytic action upon some strains of B. coli and Dünschmann² that quarter evil (Rauschbrand), serum could act on the bacillus of malignant cedema. Such "group reaction" on the part of amboceptors is but slight compared with what is seen in the case of agglutinins (see p. 530).

6. It is possible to develop temporarily non-specific protective powers on the part of the organism. Such increased resistance against pathogenic bacteria in general may be developed at the height of inflammation, or, as Issaeff has shown, by preliminary inoculation of various fluids (sterile broth, urine, physiological salt solution, etc.). Such increased protection is temporary—Issaeff's resistance period—and has no effect on the blood serum; it disappears in ten to fourteen days. His observations indicate that it is associated with increased leukocytic activity. We must recognize, that is, that the organism employs more than one means to protect itself (see p. 499).

7. The antibacterial amboceptors as a class are unaffected by heating for several hours to 60°, but are destroyed at 70°.

8. They are not immediately produced upon inoculating animals

¹ Centralbl, f. Bakt., 19:1896:51.

² Ann. de l'Inst. Pasteur, 8; 1894: 403.

with bacteria; usually three or more days clapse before they are recognizable in the blood.

 Once developed by the organism, they are to be recognized in the blood serum for a considerable period, varying with the different species, but extending in some cases to a year and more.

 If they disappear from the blood serum, a relatively very slight inoculation of the specific microorganism will result in their reappearance in abundance.

11. They may be developed either by progressive inoculations of the living microbes, by larger doses of the killed microbes, or by a combination. Such inoculations (Haffkine, Wright, etc.) may be employed to produce immunity in man against cholera, plague, typhoid, streptococcus infections, etc.

12. The complements of different animals are not necessarily identical, and for this reason the immune serum developed by one species will not necessarily protect another; thus, anthrax immune serum will protect one species and not another (Sobernheim). Vibrio Metchnikovi immune serum gained from the rabbit will protect rabbits, but not pigeons; *i. e.*, while the amboceptors produced to combine with the bacteria will so combine whether they encounter the bacteria in rabbits' or in pigeons' blood, in the second case the complements afforded by the pigeons' blood, being different, will not combine with the amboceptor, and the bacteria are not destroved.

13. So also there may be more than one order of complement; in the same blood there may be those of more than one order. Normal rabbits' serum heated to 56⁵ loses its power of activating cholera and typhoid immune serum, but can still act upon anthras bacilli.⁴

14. Virulent bacteria possibly possess more specific receptors than non-virulent, a larger amount of the amboceptor-containing immune serum being required to neutralize and destroy them. An alternative view is that virulent bacteria produces more antibodies of the nature of aggressins (p. 557) which antagonize the action of the amboceptors.²

15. Anti-ambaceptors which can readily be obtained in hemolytic studies are not so easily gained against bacteriolytic agencies. The haptophoric constituents of the ambaceptors tally with the corresponding bacterial receptors, and it is unlikely that these ambaceptors incoulated into normal animals (to develop anti-ambaceptors in their blood) will find in the organism of those animals receptors which correspond with those of the bacteria; unless such receptors be present the anti-ambaceptors cannot be developed (Friedberger).

Nevertheless, occasionally such have been found, as Pfeiffer and Friedberger^a chanced to gain anti-immune sera which neutralized the protective action of goat typhoid and cholera immune sera in guinea-pigs, by inoculating rabbits with a goat typhoid and goat cholera immune sera respectively (*i. e.*, the sera of immunized goats).

¹ Bail, Centralbl. f. Bakt., 27; 1900; 10 and 517.

² Weehsberg, Zeit. f. Hygiene, 39: 1902; 171. ³ Berl. klin. Woeh., 1902; 204. 35

16. Diversion of Complement.—Lastly, a word must be said regarding a peculiar phenomenon first noted by Neisser and Wechsherg.¹ If a suspension of bacteria be taken in a normal serum containing sufficient complement to cause bacteriolysis when a known amount of inactivated serum is added (*i. e.*, containing *x* amboceptors), then if, say, 10 *x* amboceptors be added, instead of the solution of the bacteria being hastened, the opposite occurs; it may be wholly arrested. There is evidently a diversion of the complements, the excess unattached amboceptors have a greater avidity or attraction for the complement molecules than have those that have become partially satisfied by attachment to the bacteria, or, conversely, it may be that the avidity of the bacterial receptors is greater for amboceptors pure and simple than for the combined amboceptors plus complement.

The phenomenon has been variously explained. Thus, Metchnikoff suggests agglutination of the bacteria by the excess of inactivated serum,



Diversion of complement: when there is excess of amboseptors 1, those which have not combined with the complement molecules C are supposed to enter more readily into combination with the bacterial receptors B than do the compound amboseptorcomplement molecules. or the presence in the excess serum of sufficient anticomplements to take the place of the complements. Gruber also has suggested anti-complements already present and made active by the addition of large doses of inactivated serum. To us the simplest explanation appears to be afforded by the diagram (Fig. 167), namely: an amboceptor which is within the sphere of influence of a complement, and is being attracted by it, is less likely to succumb to the attraction of the cell receptor than is an amboceptor not subjected to such influence, and so the receptors become satisfied by unlinked amboceptors.² In favor of this view are certain observa-

tions of our colleague, Dr. Meakins.⁹ Employing erythrocytes that had been repeatedly washed to remore every trace of serum, he found that no hemolysis occurred if he added to these a relatively large amount of heated hemolytic serum and a normal amount of normal serum. Centrifugalizing the corpuseles, he found that the amboceptors had become attached, for now the addition of fresh normal serum was followed by hemolysis. In other words, the corpuseles take up the unattached amboceptors in preference to those which had already in the serum become attracted to the complement.

This diversion of complement is a matter of practical importance in immunization, namely, the amount of immune serum injected, in order to be effective, should be within certain limits. Loeffler and Abel' report,

³ Johns Hopkins Hosp. Bull., 1907: 259. Centralbl. f. Bakt., Orig., 19: p. 51.

¹ Verhandl, d. Internat, Cong. d. Hygiene, Brussels, 1901; 697, and Münch, med. Woch., 1901, No. 18,

² Buxton, Jour. of Med. Research, N. S., 8: 1905: 431.

DIVERSION AND FIXATION OF COMPLEMENT

for example, that obtaining a serum which protected against B. coli, and giving animals a lethal dose of the organism, injections of more than 0.25 cm. of the serum did not prevent the fatal result. The protective dose lay between 0.02 and 0.25 cm. of the serum,

я

a

a

a,

е

17. From the practical point of view, that of establishing passive immunity by inoculating immune serum, it is of importance to recall what has already been said regarding the variation in the amount of complements in the individual and the reduction these may undergo in the course of the disease, as also the fact that the complements in the blood of one animal do not necessarily correspond with those of another, and are not so active in the blood of that other animal as are the complements proper thereto. Thus, passive immunity and the destruction of bacteria are not always complete. Mixed immune sera are thus sometimes found more satisfactory than the immune serum of a single animal or species; or it might be suggested that, as a final result, normal human serum is most likely to afford the right order of complements for human patients, a relatively small amount of serum containing it is found, sufficient complement to satisfy a large bulk of amboceptors.

Fixation of Complement.—The Bordet-Gengou Phenomenon.—Certain allied phenomena noted independently by Bordet and Gengou have led to the development of what has already become a valuable diagnostic method, even if opinions are still largely at variance as to the exact nature of those phenomena. To Bordet we owe the observation that if sensitized red corpuscles (i. e., corpuscles which placed in immune serum have taken up amboceptors) be placed in normal unheated serum they take up all the complement (as he held), or all the complements (according to Ehrlich), present in that serum, so that now this serum becomes wholly inactive for bacteriolytic or other cytolytic purposes. Similarly, if bacteria be sensitized, they absorb or fix all the complemental substance present in normal serum subsequently added. Accepting Ehrlich's view that there exists a (limited) multiplicity of complements, it is evident that the amboceptor-laden cells absorb or render inactive much more than the amount of complement necessary for the cytolytic process, and absorb indifferently all orders of complement.

Gengou showed that a like "fixation of complement" takes place under conditions in which complement plays no part in the main process. Thus, if to an immune precipitin serum a trace of the protein employed as antigen be added, even although the precipitate caused be so minute as to be invisible, the complemental bodies present in the serum become fixed and the serum subsequently cannot be employed to activate sensitized erythrocytes, etc. It is found, for example, that when toxin joins with antitoxin there is a like coincident fixation of complement. In short, any serum in which antigen and antibody undergo union is, as regards the function of its complements, rendered inactive. As already stated, what is the explanation of these processes is still debated. Along the line of certain most interesting observations by Gay,¹ it may be sug-

¹Centralbl. f. Bakt. Orig., 39: 1905; 172 and 603, and 40: 1906; 695. See also Moreschi, Berl. klin. Woch., 42: 1905; 1181.

gested that in all these cases in the production of the immune serum there is the coincident production of precipitins, and that as a consequence, when the antigen is added to the mixture, with it some serum is also added, leading to the formation of a precipitate, which, separating out, absorbs and carries down the complement. That certain proteins of the serum are responsible for the fixation would seem indicated by Noguchi's observations, to be presently noted.

The Wassermann Reaction.—⁷These observations upon fixation of complement render it possible to determine the presence of either antigen or antibody in a given fluid. Put briefly, if either be present, then the addition of the other in the presence of complement-containing serum leads to the fixation of that complement. If the fluid under examination contains antigen or antibody, respectively, then by this fixation the serum is rendered inactive, and will not, for example, activate sensitized erythrocytes and bring about hemolysis. If it is devoid of the specific antigen (or antibody), the complement does not undergo fixation, and as a result hemolysis occurs.

This is the rationale of the now well-known Wassermann reaction. Wassermann proceeded on the assumption that if he could obtain a syphiltic antigen he could along these lines determine the presence or absence of a syphiltic antibody in the serum of those suspected of syphilis, and could thus diagnosticate the existence of the disease. As antigen he selected the liver of a syphilitic feetus (or other feetal tissue rich in spirochetes) and made an extract of the same. To a given amount of this antigen extract (1) there is added:

2. The serum to be tested, diluted with normal saline solution and heated to destroy its complement.

3. Normal unheated guinea-pig serum (containing complement). The mixture or mixtures, containing varying proportions of these three ingredients, are placed in the incubator for one hour. If the serum to be tested contains the syphilitic antibody, then in its union with the antigen the complement will undergo fixation. Coincidently there has been prepared a mixture of

4. Washed erythrocytes of man, sheep, or other animal.

5. Heated immune serum from a rabbit which had been injected with the crythrocytes in question, or (5a) instead, the sensitized crythrocytes may be employed (*i. e.*, crythrocytes already treated with heated immune serum).

If, now, either the mixture of 4 and 5, or 5a, be added to the previous mixture, if the serum to be tested is negative, the complement will not have been fixed, but will be free to activate the sensitized erythrocytes, hemolysis being the result. A positive diagnosis is afforded when no hemolysis ensues. Needless to say, as control the mixture of 3, 4 and 5, should afford hemolysis.

The results of this method of diagnosis have, on the whole (with certain reservations), been satisfactory.

The majority of known cases of syphilis afford the reaction (90 per cent. of untreated primary and secondary cases, and the majority of

THE WASSERMANN REACTION

tertiary cases). It has been of distinct service in demonstrating that almost every case of general paralysis (well over 90 per cent.) affords the reaction.¹ In locomotor ataxia (*tabes dorsalis*) a smaller proportion react. But certain authorities have found diseases due to animal microparasites afford frequently the reaction, and that scarlet fever cases more particularly may yield positive results.

But interesting, not to say valuable, as are the proved results of the method, we now know that the reaction is of an order distinct from the Bordet-Gengou phenomenon proper. Equally valuable, and, in fact (by the use of Noguchi's method), more delicate, reactions can be obtained if Wassermann's syphilis antigen be wholly dispensed with and in its place: (1) an alcoholic extract of a normal organ,² (2) soluble soaps,³ (3) lecithin,⁴ or (4) bile salts⁵ be employed. This enumeration will vividly recall the description given (p. 540) of the bodies concerned in artificial hemolysis. It is evident that we deal with another lipoid reaction.

In conformity with these other observations, we are led to conclude that the blood serum of syphilities differs from normal blood serum in the relationship existing between its proteins and the lipoid bodies, and this to a certain extent has been proved. As shown by Klausner⁸ and Noguchi,⁷ syphilitic blood serum differs from the normal in the relative abundance of globulins it contains; indeed, Klausner⁸ has recommended the globulin reaction as a test for syphilis. In conformity with Noguchi and von Liebermann's observations (p. 540). I would suggest that lipoids added to syphilitic serum combine with these abundant globulins to form complement-like bodies, which, now joining with the immune bodies present in this process, coincidently bind the natural complements of the guinea-pig's blood present in the mixture. We have to deal with a nonspecific or generalized Bordet-Gengou phenomenon.

Saying this, attention must be called to the pitiable confusion that has resulted from this group of lipoids concerned in the modified Wassermann reaction being popularly referred to as "syphilitic antigens." They are nothing of the kind. Not one of them inoculated into an animal will lead to the production of specific antibodies against syphilis. All the true antigens known thus far are nitrogen-containing bodies. I do not say that it is not possible that there exist carbohydrates and fatty bodies which stimulate the development of antibodies to specific carbohydrates

¹ This, coupled with the fact that such general paralytics are found immune to reinfection with syphilis, confirms the deductions already made by clinicians that this disease is the outcome of, in general, mild and incompletely cured syphilis.

² Landsteiner, Müller, and Plötze, Wien, klin, Woch., 20; 1907; 1421 and 1565.

^a Sachs and Altmann, Berl. klin, Woch., 45; 1908; 349.

⁴ Porges and Meier, Wiener klin, Woch., 21: 1908:246.

^b Sodium glycochocolate and sodium taurochocolate, I evaditi and Yamanouchi, Comptes rendus Soc. de Biol., 62: 1907; 740, and 64: 1908; 349. They found cholin and protagon to have a similar action.

⁶Wien, klin, Woch., 21:1908:214 and 363.

⁷ Jour. Exp. Med., 11: 1909: 84.

8 Loc. eit.

and fats; but these carbohydrates and fats are not syphilis. Rather, it would seem that these lipoids, if not themselves complements, areif so barbarous a term be permitted-complementogens, and that the active agent in the reaction is, in the first place, the globulin or globulins of the syphilitic blood serum. It is through the agency of these globulins that the lipoids are brought into relationship with the amboceptors of the syphilitic serum.

ANIMAL VENOMS.

We have already more than once called attention to the toxic action of animal venoms-the venom of snakes, scorpions, certain spiders, etc.--and should in strict order have considered them after the toxins, but, as we shall point out, in some respects they are more nearly allied in their mode of action to the cytolysins and occupy thus an aberrant position. Here it may be briefly noted that, by cautious repeated injections, whether of minute quantities, or, better, of the modified toxin, into the lower animals, antitoxins-antivenins-can be obtained to all the animal venoms. Such antivenins were first obtained against cobra poison by Phisalix and Bertrand,1 Calmette,2 and Fraser3 by different methods of procedure. Calmette's antivenin from the horse has been produced extensively, and has been found of service in cases of actual snake bite in human beings. It was with such antivenom that Kanthack⁴ first demonstrated the neutralization of toxin and antitoxin in vitro. Since then there have been produced experimentally the following: antiscorpion venom (Calmette), anti-arachnolysin (against the poison of the spider, Epeira diadema, Sachs),⁵ antiphrynolysin (against toad poison [Bombinator bufo] Pröscher),⁶ antisalamander poison (Phisalix), anti-eel poison, antifish poison, etc.

Needless to say, it is snake venom that has attracted the greatest attention and that has materially contributed to our knowledge of toxic actions, and here Kyes,7 Flexner and Noguchi,8 and others have demonstrated that the raw poison contains several separate toxins-a hemolysin, a neurotoxin, a nephrotoxin, an endotheliotoxin, etc. Nor have all of these the same mode of action. The majority act directly, but others, like the hemolysin, require the intermediation of complement. We seem, indeed, with these venom hemolysins to have a series of bodies intermediate between or at least indicating the relationship between the toxins which act directly on the cell and those which for their action require the intermediation of a complement. Both spider and toad

¹ Compt. rend. Soc. de Biol., 1894.

³ Brit, Med. Jour., 1895; ii: 416.

⁵ Hofmeister's Beitr., 2: 1902.

6 Ibid., 1:1901. ⁷ Berl, klin, Woch., 1902; Nos. 38 and 39. Also in Ehrlich's Ges. Abhandl., 1904

² Ibid., 1894.

4 Jour. of Physiol., 3: 1893.

⁸ Jour. of Exp. Med., 6: 1901-05: 278.

VENOMS AND OPSONINS

poison act directly, indeed rapidly, upon the red corpuscles and give rise to bodies having all the characters of antitoxins. Cobra venom acts immediately on the red corpuscles of some animals (e. g., the guinea-pig), has no action on others (ox, sheep, goat), but can be activated in the latter cases by the addition of the blood serum of another animal (e. g., guinea-pig). That added blood serum must contain the complement; and in the case of the susceptible animals we can only conclude that the erythrocytes already possess an endocomplement. As a matter of fact, as already noted (p. 538), Kyes demonstrated that if lecithin does not constitute this endocomplement, at least it can act as a complement and so bring about the hemolysis.

OPSONINS.

For the recognition of another, and important factor in the interaction between the body cells and bacteria, we are indebted to Sir A. E. Wright¹-not that he was the first to call attention to the phenomenon. That the presence of serum aids phagocytosis by the leukocytes had been observed by certain workers² before he, with Douglas, published the first of his studies upon this subject (1903). But to Wright is due the credit of devising a precise method of determining quantitatively the influence of the serum upon the leukocytes and of recognizing the significance of these observations upon the arrest of active disease, and of the establishment of "vaccine therapy" upon a firm basis. Wright and Douglas employing Leishman's method of observing phagocytosis by leukocytes outside the body and under the microscope,³ have made out that, under ordinary conditions, the polymorphonuclear leukocytes do not freely take up and digest bacteria unless certain substances be present in the serum, which, in consequence of their auxiliary action, they term opsonins,4 and to the presence or relative absence of these substances they attribute to a large extent the eventual destruction of bacteria by pagocytic activity or their continued multiplication within the organism. They find, for example, that the leukocytes from a person suffering from chronic furunculosis or other pyococcic or streptococcic diseases are able to take up abundant cocci if removed from the patient's blood serum by centrifugation and placed in the serum of a normal healthy individual. In fifteen minutes, kept at 37° C., such leukocytes may each have taken fifteen to thirty cocci added to their suspension, whereas under the same conditions the leukocytes in the patient's own serum may have been almost inert. And contrariwise, the leukocytes from a healthy individual placed in the blood serum of the patient may also show scarcely any activity. Phagocytic activity is, therefore, not merely a matter of leukocytic activity, but is favored and stimulated by the action of some

¹See his Studies in Immunization, London, Constable, 1909.

² Denys and Leclef (La Cellule, 11: 1895; 175), and Markl (1903).

³ Brit, Med. Jour., 1902; i; 73. From "opsono," I cater for.

substance in the blood serum. How this acts is still a matter of some debate, $^{\rm i}$

Independently, but later, Neufeld and Rimpau² described the same bodies as *eptotropic* substances. They found that the serum of animals immunized against streptococci, pneumococci, and erythrocytes contains substances which act upon the bacterial cells or erythrocytes in such a way as to favor their ingestion by leukocytes. Like Wright, Neufeld further noted that they are thermostable, that they can be heated to 59° C, for half an hour without being destroyed, and that they become fixed by the microbes, but not by the leukocytes.³

To obtain Wright's phenomenon certain precautions are necessary: (1) The bacteria employed must be in an emulsion, so made that the individual microbes are separate and not massed into clusters. (2) The emulsion must not be too thick, *i. e.*, too great an abundance of bacteria by overlying the leukocytes in the preparation gives false ideas regarding the extent of the phagocytosis. (3) The observer must have considerable training so as to reduce or render constant the personal factor in the bacterial counts. It is better that the counter should not know beforehand the history of the preparation he is engaged upon. (4) The same pipettes should be employed for the same stages of the process, so as to insure accurate mensuration and mixture. (5) The greater the number of leukocytes counted the less the possibility of error. These are but some of the more obvious precautions; there are abundant minute details of technique as developed by Wright, all making for accuracy, but with the greatest perfection of technique, it has to be admitted, that the limit of experimental error remains high.⁴ Nevertheless, as the accompanying examples show,⁵ in the hands of a careful worker closely accordant results are obtainable.

Experiment I.—Rabbits' serum mixed with emulsion of staphylococci and human leukocytes (from seven persons) in the proportion of 3: 1: 3. Phagocytic count obtained by counting the number of cocci in 35 polymorph leukocytes and then calculating the number per leukocyte.

											lei	occi p tkocy	
1.	Rabbits'	serum	+	cocci	+	corpuscles of	W. B.	(normal	male)				9.8
2.						11	F. T.					{	0.3
3.		11	\pm		+	-11	O. G.					- 1	9.7
4.			÷	4.6	\pm	11	R. D.	11					9.6
5.			+		+	4.6	C. H.	44	. 12			- 1	0.0
6.			÷	++	÷	**	H. M.	(an anen	nic fem	ale)		- 1	9.9
7.		14	÷			**	S. M. ((male, fa	cial act	ne)		= {	9.0

⁴ Wright and Douglas, Proc. Roy. Soc., 72: 1903: 357; 73: 1904: 125, and 74: 1904: 159. See also Bulloch, Practitioner, November, 1905.

² Deut. med. Woch., 40: 1904: 1458.

^a Centralbl. f. Bakt., 38: 1905.

⁴ Vide Fitzgerald, Whiteman, and Strangeways, Bull. of Comm. for Study of Special Diseases, 1: 1907; 115; and Hort, Brit, Med, Jour., 1909; i; Feb. 13, ⁵ Bull, and Advis, 15; and Hort, Brit, Med, Jour., 1909; i; Feb. 13,

⁵ Bulloch and Atkin, Proc. Roy. Soc. Lond., 74: 1905; 381.

OPSONINS

Here it will be seen that, using the same serum, but different leukocytes, the counts are practically identical; differences between 9.0 and 9.9 do not exceed the limits of error of observation. In Experiment II the sera are different, but the leukocytes of one individual are employed throughout.

Experiment II.—Various human sera + cocci + one kind of leukocyte (from a normal male individual):

											Cocci per eukocyte,
1.	Serum of	W. B.	+	cocci	+	corpuscles of	W. B.				21.3
2.		F. T.		11	÷						20.3
3.		O, G.	\pm								21.1
4.		R. D.			$-\pm$						20.0
5.		C. H.									19.8
6.		H. M.									15.5
7.		S. M.									14.0

Here it will be seen that there are more differences; not much between the sera of the healthy males, but that of the anemic H. M. stimulates to a distinctly lessened phagocytosis, while the slightest of all is induced by S. M., who suffered from a disease, acne, due to growth of a particular microbe, the pyococcus.

These observations of Sir A. E. Wright have thus far been abundantly confirmed in numerous laboratories. It has been demonstrated that:

1. In a large number of infections protective substances (opsonins) exist in the blood serum.

2. There is a multiplicity of these bodies, which may be divided into two main groups: The *natural* opsonins present in normal serum are in the main thermolabile; the serum containing them is inactivated, if heated to 56° C. for a few minutes. Others appearing in the serum during the course of tuberculosis (Wright), and after strong immunization to such bacteria as those of typhoid and dysentery, are thermostable, unaffected by heating to 60° C. for ten minutes.¹

3. A certain grade of specificity can be recognized. A given serum may be active in promoting the phagocytosis of one species of bacteria (e. g., pyococci), inactive in connection with another species (e. g., tubercle bacilli). It is suggested (Hektoen and others) that there is a common opsonin (thermolabile) in normal serum, and that after vaccination specific opsonins are developed, some of which, only, are thermolabile.

4. The opsonins act upon the bacteria, so that the latter can subsequently be ingested by the leukocytes.

⁵5. When different bloods are compared, the variable factor is the serum, and not the leukocytes. This is not the same as stating that leukocytes of different individuals do not vary in their phagocytic activity. In certain diseases of the hemopoietic system, as shown by

¹ See Rosenow, Jour. Inf. Dis., 3: 1906; 683; Hektoen, ibid., 5: 1908; 249; Muir and Martin, Brit. Med. Jour., 1906; ii: 178; Dean, ibid., 1907; ii (with bibliography).

Ledingham,¹ apart from changes in the opsonic content of the serum, these may vary greatly in their phagocytic power, and this variation may explain occasional aberrant results. But, as a rule, the variations in the cells are so slight compared with the-range of variation of opsonic power of sera that it may be neglected. So also the matter is not affected by the fact, brought out in Metchnikoff's laboratory, that some slight ingestion of bacteria is seen to take place by leukocytes suspended in normal saline solution and in the absence of serum and opsonin. Such observations do not indicate that opsonins do not exist. In a normal serum at body temperature within fifteen minutes normal polymorphs take up abundant bacteria, not an occasional coccus, but from thirty to fifty may be counted in a single leukocyte. This difference between the effects of a normal serum and of an inactivated serum or physiological salt solution is very striking.

6. The specific opsonin is used up when bacteria are added in sufficient quantity to a serum, so that on removing the bacteria the serum used with a second portion of the same emulsion is inactive. In general, also, the serum is rendered inactive toward bacteria of other species.

7. The opsonins become combined or at least absorbed, by the bacteria, so that these bacteria removed after treatment and placed in an inactivated serum are freely taken up by leukocytes mixed with the same.

8. That there is a definite combination is suggested by the fact that, whereas the opsonin in serum is destroyed by a heat of 60° C., the mixture of serum and bacteria that has undergone opsonization may be heated to 60° C. for long periods without abolition of the opsonic effect (Bulloch and Atkin).²

9. By careful vaccination with measured small quantities of dead cultures of various pathogenic microbes (Pyococcus aureus, gonococcus, B. coli, B. tuberculosis, etc.), it is possible to increase markedly the opsonizing power of the serum of the individual. Such vaccination is followed by what Wright terms the "negative phase," during which the specific opsonin becomes reduced in amount; if the vaccinations succeed each other too rapidly there may be a summation of the negative phases and great lowering of opsonic content of the serum; secondly, there is a positive phase of increased opsonic power of the blood serum.

10. Regarding phagocytosis as the main process by which bacteria are destroyed within the organism, and the opsonins as the means whereby the bacteria are prepared for ingestion, Wright has concluded that the relative amount of opsonins in a given serum gives an indication of the defensive powers of the individual; for this purpose he has established an "opsonic index." This is the ratio between the average number of bacteria found within 20 to 40 polymorphonuclear leukocytes of an emulsion made with the patient's serum and the number found in the same number of like leukocytes in an emulsion made with normal

¹Lancet, London, June 16, 1906. See also Shattock and Dudgeon, Proc. Roy. Soc. of Medicine, 1:1908: Medical sect., 169.

² Proc. Roy. Soc., Biol., 74: 1905; 379.

OPSONINS

serum, the latter being taken as 1.0. For greater sureness a "pooled normal serum" may be employed, *i. e.*, a combination of the sera of five or more apparently normal individuals.

In most infections the index is found to be below 1.0. With carefully measured subcutaneous injections of dead specific bacteria there results a rise of the opsonic index, and this rise corresponds to an obvious improvement in the general condition of the patient and the local manifestations of the disease. By carefully watching the index it is possible by successive vaccinations to bring up the index in successive steps, until it reaches and exceeds the normal, and coincidently in these diseases a very material improvement is to be recognized, if not complete arrest of the morbid process. This is particularly the case with conditions due to the Pyococcus aureus. Good results are also obtainable in certain cases of gonorrhea, B. coli infections, tuberculosis, etc., although in the latter disease it cannot be said that the opsonic index affords clear indications.

There are admittedly difficulties in connection with Wright's mode of determining the opsonic index. One of these we have already noted, namely, the factor of error; another is the smallness of the scale and slight difference in the end reaction; a difference of a few tenths of a degree, say from 0.8 to 1.0, may be due to the personal equation of the observer, or may mean a very material alteration in the opsonic contents of the serum. Simon has recommended that using a moderately thin emulsion of bacteria, a large number of leukocytes be counted and the percentage of those which have taken up bacteria be estimated and compared with a control. This method gives results which tally closely with those obtained by Wright. A more serious objection is that neither of these methods give an estimation of the opsonic contents of the same order as those gained in the determination of the agglutinating and bactericidal powers. We determine the agglutinating powers, for example, by the process of dilution, by finding the limits of dilution beyond which the bacteria no longer undergo clumping. There is no reason why a similar method should not be employed for the opsonins, and, as pointed out by Neufeld and Hüne,1 Klien,2 and Meakins,3 such a method gives us a greatly expanded scale, reduces the personal error and demonstrates features in the reaction which are not revealed by the older method. As unit is taken the average number of bacteria per leukocyte in a control made with salt solution. When the serum has been diluted to such an extent that the leukocytes now only present the like average number of bacteria, it is seen that the point has been reached at which the opsonins cease to be active. It is found that frequently this point is not reached until (as happens with some agglutinating sera) a dilution of 1000 and more is reached. As with agglutinins, the count of opsonic activity differs with different species, dilutions of 300 to 1500 being

¹ Arb. a. d. k. Gesundheitsamte, 25: 1907; 164.

² Johns Hopkins Hosp. Bull., 18: 1907: 245.

^a Jour. of Exp. Med., 11: 1909: 100.

effective with animals immunized against tuberculosis, of 3000 after inoculation with Streptococcus pyogenes. The method demonstrates, as does Wright's, the existence of a negative phase; demonstrates, further, that bactericidal agglutination and opsonic powers, while often showing a coincident rise or fall, may nevertheless vary independently. Meakins has noted the curious fact that occasionally in the undiluted state a normal serum causes greater phagocytosis than an immune serum, yet the latter may show a much greater power when both are diluted a hundred times or more.

The want of certainty in the readings has, indeed, rendered many very skeptical regrading the full carrying out of Sir A. E. Wright's technique; it has to be admitted that all do not react similarly to successive vaccinations, and that even in his own practice Wright encounters not a few obstinate cases which do not react satisfactorily to his vaccinations.

On the Nature of Opsonins.—Are we here dealing with a series of bodies *sui generis*, or with a particular function of antibodies already known to us? The general employment of Wright's method for their estimation has obscured matters. A comparison of that and Neufeld's method shows that the extent of phagocytosis is not an index of the amount of opsonins developed, and the latter method shows clearly that with immunization the amount of opsonins present rises very much in the same manner as do the agglutinins and bacteriolytic substances. But, as already noted, there is not an exact parallelism between opsonins and either the one or the other order of these substances.

Undoubtedly, the way in which the opsonins unite with the bacterial or cell bodies suggests that they are of the nature of amboceptors. But here comes the difficulty. The natural opsonins are thermolabile, destroyed by heating for half an hour to 56° C. Amboceptors are characteristically thermostable. Rather, therefore, these natural opsonins are of the character of complements; like them, they are present in the normal blood; and, as Muir has pointed out, substances which remove or fix the complements remove also the thermolabile opsonins. But if we are to regard them as complements, then we have to assume that complements undergo absorption by bacteria, and so sensitize them. This we admit is possible, although contrary to our usual conception of their action.

The induced opsonins, on the contrary, are many of them thermostable; they make their appearance in increasing amounts during the course of immunity; they unite with the antigen; they are largely specific; they agree in type with the amboceptors.⁴

We are thus confronted with the dilemma that both bodies of the type of complements, and others of the type of amboceptors, function as opsonins. Our previous studies have scarce prepared us to find complements and amboceptors performing similar or interchangeable functions.

⁴ Dean, Proc. Roy. Soc., Biol., 76: 1907; 399; see also Chapin and Cowie, Jour, Med. Research, 17: 1908; 213,

AGGRESSINS

Here in this open state the problem lies at present. As possibly pointing to a solution, it may be recalled that Liebermann found oleic acid acting as a hemolytic amboceptor, and serum-soaps as complement; found that it is a common main constituent (oleic acid) acting in both orders of bodies; as, again, that several observers have detected thermostable opsonins present, if in small quantities, in normal serum, and thermolabile opsonins in immune sera; as also that heated immune serum plus normal serum, leads to more extensive phagocytosis than either serum produces separately. Can it be that after all we deal with one phase or aspect of complement-amboceptor activity?

AGGRESSINS.

As pointed out some years ago by Ainley Walker,¹ as also by Welch, in his Huxley lecture,² if the cells of multicellular organisms coming into relationship with bacteria and their products are stimulated to produce antibodies, we may premise that bacteria, as living cells, encountering the cells of the organism and their products, are, under favorable circumstances, stimulated to produce reciprocal antibodies, and to produce them in increasing amounts. It is in this way that we best explain those phenomena which we group together under the term "exaltation of virulence" by passage of bacteria through a succession of animals of one species. Now, the virulence of an organism is not merely dependent upon the production of toxins in the strict sense. This is immediately evident when we consider the case of the cholera spirillum, the anthrax bacillus, and other microbes which produce endotoxins almost exclusively. We have not a particle of evidence that, when these become more virulent, the production of ectotoxins undergoes increase; the filtered culture fluid from a twenty-four hour culture of the most virulent strain produces as few symptoms as does that from the most attenuated strain. Nevertheless, inject the attenuated bacilli into the organism, and phagocytosis is immediate; inject the virulent, and there is no phagocytosis. In other words, the indications are that the living virulent microbes excrete or discharge substances which are not toxins proper, but which, nevertheless, have an inhibitive or "anti" action upon the cells of the organism, substances which are not necessarily taken up by the body cells leading to their destruction, but which either neutralize the action of the opsonins or directly repel the body cells, the repulsion being greater than the attraction exerted by the other bacterial

Of late, certain interesting observations have been made by Bail^a and others which demonstrate the existence of bodies of this order. Inoculating cholera and typhoid bacilli into the pleural and peritoneal

¹ Jour. of Pathol., 8: 1902: 34.

² Brit, Med. Jour., 1902; ii; 1105,

³ Arch. f. Hygiene, 52; 1905; Heft 3 und 4. See also Wassermann and Citron, Deutsch. med. Woch., 1905; 1101.

cavities, Bail set up local infection. Taking the inflammatory fluid containing the bacteria, he removed the latter by centrifugalization and killed the few remaining organisms in the decanted supernatant fluid by antiseptics, or by heat at 44° C. This clear fluid has no toxic properties; it may be inoculated with impunity into animals of the same species. When, however, it is inoculated into an animal along with a sublethal dose of the particular (homologous) microbe, an acute lethal result follows. His associates, Kikuchi, Weil, and Hoke, report like results with dysentery, chicken cholera, and pneumonia organisms. Instead of the bacteria of a sublethal dose undergoing destruction, they multiply. There is something in the inflammatory exudate that has paralyzed the protective agencies of the body. The production of these aggressins is the more active the greater the resistance to the bacteria. They are produced in greater quantities during the strife between the bacteria and the body cells, while little is produced in the test-tube. Some, however, are so produced; they are, that is to say, normal products of bacterial activity. Thus, Kolle found that when bacteria are grown in pleural fluid or blood serum in the test-tube, or even in distilled water, then develops a substance which, when the sterile culture fluid is inoculated along with a sublethal dose of the bacteria, leads to fatal results.

An immunity may be developed against the sterile aggressin-containing fluids, and this immunity may be transferred from one animal to the other by inoculation of its immune serum.

Bail regards these aggressins as new undescribed substances; others regard them as free bacterial receptors, holding that these discharged receptors combine with the amboceptors, producing, as it were, a diversion of the amboceptor, so that the bacteria themselves are not attacked, and thus continue to proliferate. But even granting this, it is obvious that these receptors are not of the nature of endotoxins or of ectotoxins, for the fluid containing them is devoid of toxic effects. At the most, if of the nature of receptors, they are haptophorous and devoid of a toxophorous moiety.

The existence of these aggressins very probably explains certain observations of Wright,¹ Douglas and Reid,² which have been confirmed by Opie,⁸ namely, that exudates produced by the local growth of a given pathogenic microbe contain no opsonins. More correctly it may be, that there is not an absence of opsonins under these conditions, but a neutralization of the same by the bacterial aggressins. It may, indeed, be suggested that the aggressins are to the bacterial organism what the opsonins are to the animal.

¹ Proc. Roy. Soc. Lond., 74; 1904; 147.

² Ibid., 77: 1906: 194.

³ Jour. of Exp. Med., 9; 1907; 515,

ANAPHYLAX18

ANAPHYLAXIS.

Yet another order of phenomena deserves notice in this connection. From the early days of the employment of antidiphtheritic serum, occasional cases have been reported of sudden death following upon the inoculation of the serum. In 1905 von Pirquet¹ published the first full study on this subject; in 1906 Rosenau and Anderson² were able to collect nineteen such cases out of the literature. The symptoms may come on within five minutes of the treatment, with collapse, unconsciousness and convulsions; milder cases, of urticarial rashes with some nausea, are comparatively common, and it has been clearly proved that they are induced not by the toxins or antitoxins, but by the serum, horse serum producing identical effects. Along with these cases of serum sickness, attention may be called to the fatal effects which have followed the transfusion of the blood of sheep and other animals into man, in cases of grave anemia. Such transfusion led to after-effects so severe-high fever, hemorrhages, and intravascular clotting-and was so often fatal, that it was rapidly given up. Some cases, not so severe, showed merely urticaria with fever.

Experimental observations upon these phenomena have led to some very remarkable results. If a moderately large dose of a foreign serum be injected into an animal, either subcutaneously or into the peritoneum, no immediate effects are produced, and the animal in a few days becomes immunized to that serum. But, if instead of a dose of 5 c.c. of foreign serum, a guinea-pig be given a little, as 0.0025 c.c., and now in twelve days a second injection of 5 c.c. be given, the guinea-pig is apt to die, it may be, within a few minutes, or at most a few hours (Theobald Smith's phenomenon). Instead of being rendered immune, the very opposite result has been brought about; the animal has been "sensitized," rendered much more susceptible to the foreign serum. This process of sensitization has received the name of anaphylaxis.3 This hypersensitiveness may be induced, athough more slowly, by the injection of repeated large doses (in themselves non-toxic) of an otherwise harmless serum. It then shows itself by the development of ædematous and necrotic changes in the region of the later inoculations, followed by progressive cachexia and death after many weeks (Arthus' phenomenon).

It has been found that in herbivorous animals the same results may be gained by feeding with the foreign serum. Gay and Southard[†] and others have shown that the subjects of anaphylaxis exhibit hemorrhages in the stomach, cecum, hungs, spleen, heart, and adrenals; these appear to be associated with a definite fatty degeneration of the capillary

¹ Pirquet and Schick, *Die Serumkrankheit*, Leipzig and Vienna, F. Deuticke, 1905.

² U. S. Hygienic Laboratory Bulletin, No. 29, Washington, 1906.

³ By Richet, as opposed to *prophylaxis*; von Pirquet has given to it the name *allergia*, by which it is still referred to by some German writers.

⁴ Jour, of Med. Research, 16: 1907: 143.

endothelium. Further, the blood of the sensitized animals comes to contain a substance which, when the blood is injected into other guineapigs, sensitizes them. In man and omnivorous animals a single dose has sometimes the effect that the two doses possess in rabbits and guineapigs.

Such sensitization is so wholly opposed at first sight to all our experience in experimental immunity, that a totally different explanation would seem necessary. Nevertheless, for long years we have been familiar with what would evidently seem to be a phase of the same process. We refer to the tuberculin reaction, the intense local congestion of the tissues around a visible tuberculous focus that follows the inoculation of a minute quantity of tuberculin at a distance. Possibly the accelerated appearance of the vaccine erythema and eruption in those already vaccinated is a process of the same order. Various theories have been adduced, none of which is wholly satisfactory. In our first edition we pointed out that Vaughan's' remarkable studies upon the bacterial and other proteins seemed to furnish a clue. In a long series of papers, from 1901 until the present time, he has shown that the bacterial proteins may be split up into two portions-the one poisonous, the other nonpoisonous-and now, taking what is apparently the most innocuous of proteins, namely, egg white, he has determined that the same is true of this also. One may inject the white of three hens' eggs into the peritoneal cavity of a rabbit, with no untoward effects. Nevertheless, by extraction of the purified egg albumin with 2 per cent. sodium hydroxide in absolute alcohol, two bodies are obtained-one poisonous, soluble in absolute alcohol, the other non-poisonous, and insoluble. This poisonous moiety, apparently of proteid nature, kills just as promptly as that obtained from the proteins of the colon or typhoid bacillus; the minimum fatal dose for the guinea-pig ranges from 8 or 10 up to 100 mg., according to the grade of purification.

Vaughan and Wheeler find that animals may be sensitized to egg albumin either with unaltered egg white, or with the non-poisonous moiety, but not with the poisonous moiety. What is more, the non-poisonous moiety does not sensitize to itself, but only to the unbroken egg white.

These facts can only be satisfactorily explained on the supposition that, under the conditions of these experiments, when a small dose of a foreign protein is introduced into the organism, for it to be assimilated the cell substance has affinity for the non-poisonous moiety. The same results ensue, it is seen, whether the whole egg white or only its non-poisonous residue be exhibited. The cells become habituated to attract themselves to the non-poisonous moiety alone, and to form and discharge a series of receptors which combine with this. When, therefore, after this habituation or immunity has become established (in ten or twelve days), the unbroken egg white is again exhibited, the cells and these receptors—actively attract this non-poisonous moiety, liber-

¹ Vaughan and Wheeler, Jour. of Inf. Dis., 4:1907:476, give a full bibliography of the publications from the University of Michigan bearing upon the subject.

ANAPHYLAXIS

ating the poisonous moiety, which now, free in the body fluids, enters the blood, circulates to the brain, and there sets up those disturbances, more particularly in the respiratory centre, which lead to death. Vaughan and Wheeler show very clearly that the second dose must contain enough egg white to furnish a fatal dose when split up in the animal body. Wells,¹ however, is of the opinion that the amount of serum necessary to sensitize the cells is so extraordinarily small that this dissociation process cannot be invoked. But certain observations of Besredka² and of Auer and Lewis³ indicate that the fatal action of anaphylactic substances is peculiarly localized. The former showed (confirmed by Vaughan) that the toxic action is cerebral in type, and that under the influence of anesthetic doses of ether anaphylaxis does not show itself; the latter have demonstrated that the fatal result in guinea-pigs is due to a spasm of the bronchial muscles, with inhibition of respiration—in short, to what may be regarded as acute asthma, with striking insufflation of the lungs. Incidentally, as pointed out by Meltzer, their experiments throw a strong light upon the anaphylactic nature of at least an important group of cases of asthma, as also, it may be suggested, of hay fever. The reaction, however, is not the same in all species of animals: in the dog the chief demonstrable disturbance is a sharp fall in blood pressure⁴ resembling that of shock, brought about, as shown by Pearce and Eisenbrey, by loss of tone, or paralysis, of the smooth muscle of the splanchnic veins. As they point out, disturbance of the functions of non-striated muscle is the feature common to both orders of cases.

⁴ Proc. Soc. Exp. Biol. and Med., 6: 1909.

² Ann. de l'Inst. Pasteur, 21: 1907: 117 and 384.

³ Jour, Amer. Med. Assoc., 53: 1909: 458.

⁴ Biedl and Kraus, Wiener klin, Woch., 22: 1909; 383, and Pearce and Eisenbrey, Proc. Soc. Exp. Biol. and Med., 7: 1910; 30.

CHAPTER IX.

IMMUNITY-(Continued).

THEORIES OF IMMUNITY.

To master and to keep level with the vast number of individual observations that are now being poured out upon this one subject of immunity is in itself a life's work. Appalling as must be the mass of data contained in the last two chapters, to the reader who approaches the subject for the first time, these represent but a selection of the more important and generally accepted observations, and such conclusions as we have already drawn represent a sifting of opinions often very widely at variance. To have indicated and discussed the divergent views would have expanded those two chapters into two volumes. As it is, we have detailed but one theory-that of Ehrlich-and have not carried that to its conclusion. Now, it is for us to sum up, so far as is possible, the reasonable deductions that, we think, may be drawn from the data afforded, treating immunity and the process of immunization not as a subject apart, but as a branch of pathology, and indeed of general biology, so that our conclusions harmonize with those that are to be gained from the study of other vital reactions.

We note, in the first place, that all the antigens, namely, all those bodies which, gaining entrance into the system, lead to the formation of "antibodies," are themselves either cell substances, or the products of cell activity, and that the antibodies are likewise the products of cell activity. The two groups, in fact, are seen to be curiously similar in very many properties—reflections one of the other. A little consideration shows that this is not surprising; were we bacteria, we would regard the animal antibodies as toxins and our own toxins as protective antibodies, or, at least, as preparatory digestive ferments. It may well be that the bacterial toxins, being developed by organisms extremely low down in the scale of living beings, are of simpler constitution than the antitoxins developed by warm-blooded animals, but when we ascend to the venoms of vertebrate animals, we find that the cobra discharges a hemolytic *toxin*, which in its properties and mode of action is identical with the hemolysin formed as an antitoxin in the serum of warm-blooded animals consequent upon inoculation with erythrocytes. Here clearly, whether we regard the hemolysin as a toxin or an antitoxin, depends wholly upon the point of view. So far does this parallelism or reflection proceed, that as Welch suggested in his wellknown Huxley lecture, where two living organisms, the animal and the microbe, are pitted against each other, the increase in virulence which

THE PHAGOCYTOSIS THEORY

may be acquired by the latter may be the expression of the development by it of antibodies (which form the point of view of the microbe are simple antitoxins) corresponding to the development of antitoxins by the warm-blooded organism, and tending to neutralize the same.

We are dealing, it will be seen, throughout this whole study, with the methods in which the living matter, whether animal or vegetable, reacts toward other living matter, whether animal or vegetable, and the products of the same which come into contact with it, and although the statement may at first encounter appear to be both novel and extreme, further thought will confirm it; the problems of immunity narrow themselves down to special problems bearing upon the assimilation and digestion of unusual proteid matter, or, at least, of the primary products of cell metabolism.

Metchnikoff, indeed, would regard immunity as a matter of digestion and adaptation to the same by one order of cells, the phagocytes, free and fixed, and it will be well before proceeding farther to note his main observations and arguments, for they have had an extraordinary influence in stimulating work upon this subject, and, as we shall see, his conclusions do not, in their essentials, oppose the more generally accepted theory of Ehrlich; they are the expression of the same views regarded from another aspect, in some respects wider, in others narrower;1 wider in that throughout it keeps prominently in the foreground that immunity, like all other vital processes, is a matter of cellular activity, whereas the study of sera and their properties which constitutes the main method of the Ehrlich school is apt to cause neglect of this fact; narrower in that it is essentially morphological, and thus largely overlooks the fact that chemical processes underlie morphological phenomena, and again, that it would refer every important reaction to one order of cells, the leukocytes, and other potential phagocytes, and somewhat obstinately is unwilling to credit other cells and tissues with any part in the process

The Phagocytosis Theory.—We have already, on several occasions, called attention to the basal facts regarding phagocytosis, and have shown that the unicellular organism and sundry cells in the multicellular organism have the power of actively taking up foreign particles, organized and living, organic and inorganic, and that this process of phagocytosis is primarily nutritional—a means whereby the individual cell gains food material; that particles so taken up, if unfitted for assimilation, are discharged; if capable of affording food material, stimulate the formation of a digestive vacuole around them, in which vacuoles we have indications of the presence of digestive ferments, and lying thus in the fluid the foreign matter is seen to undergo solution,

¹ M. Metchnikoff has detailed his theory in his work "*L'immunité dans les maladies infectieuses*," Paris, Masson, 1901, of which an English translation has appeared, We would, however, recommend the student for practice in French to read it in the original, for it is of sustained interest and great value. A digest of his views is given by him in German in the fourth volume of Kolle and Wassermann's *Bacteriologie*.

until all that remains are a few granules of unassimilable debris which become eventually cast out.

Among such foreign bodies these phagocytic cells are able to take up the various microbes, animals and vegetable. They can, indeed, as abundantly proved by Metchnikoff, take them up in a living condition, and, observing what happens, whether in the unicellular organism, like the amoeba, or in the free leukocytes, cells of higher animals, we observe that there is the same formation of digestive vacuoles and destruction of the microbes by digestive processes. This under favorable conditions. And, as a matter of fact, the more carefully we study under the microscope the processes occurring within the organism during the course of infection, the more we become convinced that phagocytosis, whether by leukocytes or by endothelial cells, or by the newly developed "embryonic" fixed tissue cells, is extraordinarily common, and is obvi-



Mode of destruction of a bacillus (B, anthracis) by a cell. A digestive vacuole has formed around one portion of the bacillus, and that portion has lost its power of staining. (After Metchnikoff.)

ously a most important factor in the destruction of pathogenic organisms and in the cure of infectious disease. There is, indeed, not a single disease due to known animal or vegetable microbes, where there is definite reaction on the part of the organism. in which at one period or other of its course phagocytosis, and that often verv extensive, has not been observed. One has but to inject into the peritoneal cavity of one of the animals of the laboratory a little of the fluid culture of some mildly pathogenic microbe and examine a drop of the peritoneal fluid, or make a smear from the surface of the omentum in the course of a few minutes to a few hours (the time varying according to the virulence of the

microbe), to find abundant leukocytes containing the bacteria in different stages of digestion. Or, more instructive still, take, after Leishman's method, a few drops of human blood from the finger, dilute, centrifugalize, pipette off the layer of white corpuscles and suspend these in the serum, add to the suspension a small platinum loopful of a suspension of some pathogenic microbe—the Pyococcus aureus, for example—and place the mixture for a quarter of an hour in the incubator at 37° C. Upon examining a drop of the mixture under the microscope, the number of bacteria seen within the leukocytes and that have been taken up in this short time is very remarkable.

There can then be no question regarding the importance of phagocytosis as a factor in the destruction of microbes that gain entrance into the system. There may be a question, however, whether it is the one supreme method of destruction of microorganisms and what is its rela-

THE PHAGOCYTOSIS THEORY

tionship to the development of continued immunity. For the wandering cells of the organism, we know, have but a short life period, and if they gain the power of destroying bacteria they do not convey it to their descendants, for such descendants normally, we believe, do not exist. Before answering these questions there are other properties of the phagoevtes which we must recall.

In the first place, with very virulent microbes no phagocytosis may show itself; there is a *negative chemiotaxis*, or, perhaps more accurately, a lack of positive chemiotaxis; the leukocytes are not attracted, and those in the neighborhood, acted upon by the strong toxins, undergo dissolution. In the second we observe that in a typical localized infection, such as the abscess we described (p. 431) or in the pneumonic lung, for a considerable period the phagocytosis, if in evidence, is insufficient to destroy the microbes; the mere act of taking up living bacteria does not necessitate their destruction, and the number taken up may not correspond to the rate of proliferation. Only at a later period, if the infection results in healing, does the phagocytosis become adequate. To explain these phenomena, Metchnikoff invokes the property of adaptation. By accustomance, by dealing at first with small amounts of less concentrated toxins the leukocytes gain the power of withstanding and neutralizing larger amounts until a negative chemiotaxis or a weak positive becomes converted into an active positive chemiotaxis; and whereas at first the digestion was feeble, now it becomes powerful and rapid in action. These conclusions we cannot but accept; we have in Metchnikoff's remarkable studies the most convincing examples of the development of the individual cell adaptation.

All these considerations, however, throw no light (1) upon the observations made, from 1888 onward, by Nuttall and others upon the destruction of bacteria within the tissues without the intervention of leukocytes; (2) upon the destruction of bacteria in witro, by serum and other body fluids devoid of cell contents; (3) upon immunization against soluble bacterial ectotxins and the development of antitoxins, and, in fact, upon the development of antibodies against ferments, phytotoxins, and soluble "toxins" of all orders. How are all these phenomena brought within the terms of the theory?

As to the first, that such destruction takes place cannot be controverted; it is most obvious in Pfeiffer's original method of gaining his reaction in the peritoneal cavity of the guinea-pig immunized against cholera or typhoid; the peritoneal fluid is found to contain relatively few leukocytes and abundant swollen and globular bacteria undergoing destruction. Metchnikoff ascribes this extracellular destruction to a preliminary plasmolysis or leukolysis, to a destruction of leukocytes, whereby their digestive ferments become liberated into the plasma and able to act upon the bacteria in an extracellular manner. This view, that it is the leukocytes that in the main afford the bacteriolytic substance, is supported by the observations of Denys and his pupils and of Buchner and his pupils, that if a "sterile" suppurative inflammation be induced, as by the introduction into the pleural cavity of finely powdered glass or

IMMUNITY

aleurone (a vegetable protein), the exudate, rich in leukocytes, possesses distinctly more pronounced bactericidal properties than does the blood plasma and other body fluids, or even the blood serum of the same animal. Metchnikoff admits fully that the amboceptors (his *fixateurs*) pass out and become free in the ordinary plasma. These he regards as discharged by the leukocytes, but he is unwilling to accept or suggest excretion by the living leukocytes as an explanation of the presence of the complement (his *cutase*),¹ and he adduces certain important observations made by his pupil, Gengou,2 in support of his contentions and as a proof that the plasma, even of an immunized animal, contains no complements. Gengou's statements are certainly most positive. If, employing Freund's method, the blood of an animal be received direct into paraffined test-tubes, there is no destruction of leukocytes and no clotting. Such blood centrifugalized vields according to Gengou a plasma wholly devoid of complements. Therefore-concludes Metchnikoff-the complement (cytase) present in ordinary serum has been derived from the dissolution of leukocytes-they act normally as intracellular enzymes-and the theory of phagocytosis becomes expanded to this extent, that destruction of bacteria is recognized as being brought about either intracellularly by the digestive action of the leukocytes, or extracellularly by the enzyme-like action of the cytase, or complement, working through the intermediation of the fixateur, or amboceptor. But within the infected organism such liberation is infrequent, and throughout, it is cells which are potentially phagocytic that give origin to the

Further, to complete the outline of Metchnikoff's theory, it has to be noted that he recognizes two broad groups of phagocytes, each having the power of acting more particularly upon one set of substances; the *microphages* (polymorphonuclear leukocytes, eosinophiles, etc.), and the *macrophages* (hyaline leukocytes, endothelial cells, and fixed phagocytes); the former he finds more particularly active in opposing the bacteria of acute disease, the latter those of chronic disease, as also acting upon cells. Instead of a multiplicity of complements, as demanded by

⁺Here it may be well to afford a table of the various terms employed by different observers to indicate these two bodies, the amboceptor or immune body and the complement respectively:

SINON	1 M.S.
	Complement.
	Addiment (Ehrlich's first name).
ciffer).	Cytase (Metchnikoff and Bordet).
ŕ).	Alexine (Buchner).
ce (Bordet).	
Not now emplo	oyed.
	eiffer), f). ce (Bordet).

² Ann. Pasteur, 15; 1901; 68 and 232, See also Levaditi, ibid., 15; 1901; 894, and 16; 1902.

THE PHAGOCYTOSIS THEORY

Ehrlich, he recognizes only two *macrocytase* developed by the macrophages and *microcytase* by the microphages. The immune bodies he regards as derived also from the leukocytes.

His grounds for this are the following. In immunizing an animal against crythrocytes, spermatozoa, and other cells it is seen that the foreign cells are taken up by one order of cells, the macrophages. It is these cells alone that are directly involved, and thus, in the process of digesting, the cells must develop the antibodies. Of these, the ambocceptors (fixateurs) are excreted, the complement (macrocytase) remains within the cell.

If, finally, these are the deductions to be drawn from cases in which we are able to follow the fate of discrete bodies, such as bacteria and various cells, where we can see that *within the organism* it is the phagocytic cells that act upon them, and where we can determine that these phagocytes afford the antibodies, then it is justifiable to assume a like activity on the part of the potentially phagocytic cells in the case of soluble toxins.

These latter conclusions are far from having been accepted in their entirety, and we must briefly note some of the more important contradictory observations. As regards *plasmolysis* affording an adequate explanation for the liberation of the complement in Pfeiffer's reaction, Durham and Gruber have pointed out that the destruction of leukocytes in the peritoneum is only apparent. Following upon the introduction of the bacterial culture, the leukocytes are not destroyed; they become clumped or balled upon the surface of the omentum and mesentery (these observations, however, do not exclude wholly the dissolution of a certain number of the leukocytes). Pfeiffer has pointed out that the presence of increased complements in connection with Buchner's aleurone experiment does not necessarily indicate that these are derived from broken-down leukocytes, and shows that if leukocytes so gained be centrifugalized and carefully washed they afford not a trace of complement. (To this Metchnikoff objects that the repeated washing has artificially removed the cytase-but if they pass out with such ease under these conditions, may they not also diffuse out freely in the living organism?) Several observers also have given the "lie direct" to Gengou's observations. The majority, however, have not employed his method, using other means (oxalate, leech extract, etc.) to arrest the coagulation of the blood—methods which possibly lead to alteration in the leukocytes and liberation of the complements. But Lambotte,1 employing the same paraffined test-tubes, detected the definite presence of complements in the centrifugated plasma (not, so far as we can see, in the same amounts as when clotting has occurred and certain leukocytes have undergone dissolution).

Nor is it possible, in the light of Ehrlich's observations, to accept Metchnikoff's dictum that there are only two cytases (complements); and lastly it is, in our opinion, impossible to harmonize Metchnikoff's

¹ Centralbl. f. Bakt., Abt. 1, 34: 1903; 453,

IMMUNITY

theory with the observations of Wassermann, Ransom, and others that tetanus and other toxins are absorbed by the nervous tissues and there neutralized. In connection with hemolysis Metchnikoff lays down very precisely that it is the cells which absorb the "toxin" that furnish the antibody. The same process of reasoning would lead us to conclude that the nerve cells which neutralize the tetanotoxin are capable of furnishing the antitoxin. Römer's striking observations, already noticed (p. 516), upon the local production of anti-abrin in the conjunctiva of the rabbit are absolutely opposed to this narrower view.

On the other hand, as regards hemolysis in general, we would accept Metchnikoff's view rather than that of Ehrlich regarding the production of the hemolysins. These, by the supporters of the side-chain theory, are regarded as being developed by the red corpuscles, according to the general law above noted. But when foreign erythrocytes are injected into the body they are taken up by the macrophages; it is these cells in the spleen and elsewhere that more probably provide both hemolysins and antihemolysins. If, that is, a hemolysin be developed by inoculation of erythrocytes of another species, that hemolysin, it is true, inoculated into animals of that other species acts directly upon the red blood corpuscles and leads to the liberation of their hemoglobin; but in the organism the stromata of the destroyed blood cells are taken up by endothelial and other cells, and these living persistent cells it must be that produce the antibodies. The same must be true in connection with the development of the hemolysin in the first place. The brief existence of the ervthrocyte, its incapacity to reproduce itself and convey acquired properties to its descendants, are also against the supposition that these cells actively produce antibodies. We have, indeed, the observations of Calmette¹ and Jacobi² that the red corpuscles of animals rendered highly immune to such hemolytic agents as cobra venom and ricin still remain highly susceptible to the hemolytic action of these poisons, and Ehrlich³ himself has regarded an apparent exception, recorded by Kossel in the case of the rabbit immunized against eels' blood, as due not to the acquirement of antibodies by the corpuscles, but to the loss or exhaustion of the susceptible receptors. It is, in short, contrary to experience that a non-nucleated cell should exhibit the highest forms of biophoric activity. and as such must be regarded the power of multiplying its specific receptors and to this extent of undergoing growth. This does not prevent us from accepting Ehrlich's view that the red corpuscles are provided with relatively abundant receptors of weak combining powers, so that they readily give up, to the more avid receptors of cells proper, substances which have entered into combination with them, and act as the great common carriers of the economy.

The observations of Wright, also, upon opsonins, abundantly confirmed as they have been by numerous observers, and slighted but not

¹ Compt. rend. Acad. des Sciences, 134; 1902; No. 24.

² Hofmeister's Beitr., 2:1902.

³ Loc. cit., 569.

THE PHAGOCYTOSIS THEORY

contradicted by Levaditi,¹ indicate that the phagocytes while playing an all-important part in the destruction of bacteria possess an adjuvant in the surrounding fluid. There is, indeed, one striking demonstration that they are not the main source, as again, that the leukocytes alone are not to be invoked as the essential factor in immunity. While, as Metchnikoff has pointed out, the higher the grade of immunity induced the more extensive the phagocytosis, this is not a proof that the leukocytes have acquired increased phagocytic properties. My colleague, Dr. Meakins, points out to me that if two animals be taken, the one highly immunized the other as control, and the leukocytes of the two be taken after Leishman's method, no difference is to be recognized in their phagocytic activities when they are placed with pyococci in normal serum. The difference lies not in the cells, but in the properties of the serums. Where this adjuvant is developed we do not as yet know; the fact cited by Levaditi that in the absence of the serum, leukocytes suspended in an inert fluid can, very slowly, take up bacteria, suggest that the opsonins also may be developed by the leukocytes, but certainly does not prove that these form the only or the main source.

Conclusions.—A careful balancing of all the facts appears, therefore, to lead to the following conclusions:

1. Phagocytosis proper (*i. e.*, the ingestion and digestion of bacteria) is a great factor in the destruction of microbes entering the system.

2. With accustomance and adaptation to the products of bacterial growth and other toxins, leukocytes, and certain fixed cells, exhibit increased phagocytic activity. This, however, is not primarily due to increased digestive capacity on their part, but to increase in bodies of the nature of opsonins in the body fluids, preparing the bacteria for ingestion.

3. The cells which Metchnikoff regards as phagocytes and potential phagocytes, while they are the most commonly invoked to neutralize bacteria and bacterial and other toxins, are not the only cells of the organism possessing these powers.

4. With the exception of the red corpuscles (which are not cells proper), the rule would appear to be that those cells which take up microbes and microbic and other toxins are the cells which provide the antibodies.

5. Antibodies, whether present in the normal organism or developed in response to the introduction of particulate or dissolved toxins, are the products of cell activity, and their presence in the blood is a secondary process, either a true secretion (in this resembling glandular secretion proper) or to some extent, where there is cell destruction, the result of that cytolysis and of the freeing of substances previously bound in the cells.

6. Produced within the cells these antibodies can act within the cell and then bring about a condition undistinguishable from ordinary intracellular digestion, though their strikingly specific powers suggest

¹ Ann. de l'I. Pasteur, 1905.

IMMUNITY

thus that intracellular digestion, instead of being a simple single process, may be made to vary to an almost infinite extent, according to the nature of the substance entering the cell.

7. They can act also outside the cell, and in this case clearly neutralize the toxin by entering into combination with it.

8. Whether, therefore, we regard and study the processes associated with the development of immunity as occurring within the cell or apart from it, eventually we arrive at a common underlying chemical and physical groundwork for all the phenomena, and as we approach this point, although differences exist in respect to details, fundamentally the phagocytic and the side-chain theories are not contradictory; they merely view the one set of phenomena from different aspects.

The Side-chain Theory of Immunity.—Antibodies and the Sidechain Theory.—Rather than summing up in epitome the data acquired regarding the antibodies and their mode of action, it will be well to gather together and discuss the various data recorded in the previous chapters, and this first along the lines of Ehrlich's side-chain theory.

We have indicated the groundwork of that theory so far as it refers to simple toxins (p. 516); since then, discussing the cytolysins, we have become acquainted with a new order of phenomena, with the whole group of cases in which there is not a simple union between the protoplasmic molecule of the cell, whether of the organism or the microorganism and the toxin—or the complement—but combination brought about by the intervention of an intermediate body.

Here, before going farther, it will be useful to draw up, in a tabular form, the various forms of toxins and antibodies which we have had under consideration:

Enzymes Phytotoxins Bacterial ectotoxins	leading to the	production of	Antienzymes Anti (phyto) toxins Antitoxins	
Proteins animal vegetable			Precipitins	
Bacterial proteins (?)			Agglutinins	
Bacterial aggressins (?).			Opsonins	
Animal venoms (simple)			Antivenins	Acting singly.
Animal venoms (com- plex, requiring inter- mediation of comple- ment for action)			Antihemolysins, etc.	
Foreign complements .			Anticomplements	
Foreign amboceptors .			Anti-amboceptors	
Vegetable cells (bacteria) Animal cells of various)			Bacteriolysins	Requiring in- teraction of
*orders }			Cytolysins Hemolysins Leukotoxins Hepatolysins, etc,	1 amboceptor (specific), 2 complement (non-specific),

Studying this table, it will be noted that we advance from (presumably) simpler to recognizably very complex substances. To quote Ehrlich and Morgenroth:¹ "If relatively simple bodies have to be assimilated

¹Zur Theorie der Lysinwirkung, Ehrlich, Gesammt, Abhandl., 1904; 15,

THE SIDE-CHAIN THEORY OF IMMUNITY

by the side-chains, the presence of a single binding group is adequate, and, obviously, it is side-chains of such simple structure that attach the toxins. The conditions are, however, quite different in dealing with giant molecules (protein molecules). In this case, with fixation of the molecule for purposes of cell nourishment, only the preliminary stage is a matter of concern. Giant molecules are, as such, useless for the cell, and can only be rendered serviceable when they become dissociated and broken up into smaller parts by fermentative processes. Such dissociation would fittingly be gained if the 'seizing arm' (Fangarm) of the protoplasm, carrying simultaneously a fermentative group, brought this latter into immediate relationship with the booty that needed digestion and assimilation. We find an identical tendency for the seizing apparatus to be, at the same time, provided with digestive action in the whole series of the insectivorous plants, and this in the most varied The tentacles of the Drosera, for instance, are secreting and digestive 'fang arms' in the broadest sense; they cover the object seized with a juice having powerful digestive properties.


"When we note that lytic action occurs not in association with toxins, but where cell contents, whether of bacteria or blood cells, have to be absorbed, the simplest explanation is that here we are dealing with proteins of high molecular composition which are of much more complicated structure than cell secretions, for such we must regard ectotoxins. We must conclude that to combine with these and other highly complicated bodies the cell molecule possesses side-chains of a special order, which, in addition to the seizing complex, possess another complex capable of fixing the particular ferment, and so bringing about the digestion of the seized molecule. If, through immunization, there is developed an excessive production of side-chains, then the whole of this side-chain with its two complexes will be generated and discharged into the blood as an immune body. The remarkable process whereby, as a result of introducing bacteria into the system, a matter is generated which leads to the dissolution of the bacteria receives in this way a simple and natural explanation. We deal once more with the reproduction of one of the processes of normal cell activity."

Ehrlich's Three Orders of Receptors.—Continuing along these lines, Ehrlich distinguishes three orders of receptors in the cell, whether for the assimilation of food or for seizing of toxin molecules. (1) The relatively simple toxins and ferments are anchored by a receptor of the first order (Fig. 169), I, a), a side-chain possessing simply a haptophorous complex e, to which the toxin b becomes anchored by its haptophore c. (2) For (compound) protein molecules he holds that more complex receptors are requisite. In the first place, it is evident that the cell in the process of assimilating ordinary protein food molecules by its fermentative activities dissociates them, and the same would appear to be true regarding agglutinins and precipitins. The side-chains which anchor molecules of this order must possess both a haptophore and a zymophore (corresponding to the toxophorous moiety of the simple toxin molecule). This form of receptor is indicated in Fig. 169, II, in which

IMMUNITY

e represents the haptophore, d the zymophore. It will be seen that this is the converse of the first case: the haptine or free molecule or sidechain b of Fig. 169, I, is identical in properties with the attached receptor of Fig. 169, II. (3) For the cell to act upon the yet more complicated substance of bacterial and animal cells it has to anchor not only the cell molecule, but also the complement. This type of receptor is indicated in Fig. 169, III. There the complement k is represented as possessing a haptophore, h, and a zymophorous or, more accurately, zymotoxic moiety, z, while f represents the cell molecule that is acted upon.

When these receptors are produced in excess and become discharged, they are termed by Ehrlich *haptines*. They possess the same properties of attachment as they do when existing as fixed side-chains of the



The three orders of side-chains, according to Ehrlich.

protoplasmic molecule. Thus, there are recognized haptines of three orders—those possessing a single haptophore group, those like the toxin molecule above noted, with a haptophore and zymophore group (these two are both regarded as uniceptors), and those with two haptophoric affinities, the amboceptors or immune bodies proper. Of these free receptors or haptines, Ehrlich, as a result of a consideration of their properties, lavs down that:

Haptines of the first order include antitoxin and anti-enzymes.

Haptines of the second order include agglutinins and precipitins.

Haptines of the third order include cytolysins and bacteriolysins (amboceptors).

It will be observed that in this scheme no note is taken of, or, at least, no stress is laid upon, the group of original attachment of the haptine to the protoplasmic molecule. Nevertheless, there must here be an unsatisfied affinity.

We cordially accept Ehrlich's scheme in its general bearings, and cordially admit, as must everyone who has followed the development of this department since first the theory was enunciated, that from the results obtained by working along the lines of the theory it has more than justified itself. As by Mendeljeff's theory the chemists and physicists have been able to predict and discover new elements, so, by this theory, Ehrlich and his fellow-investigators have been able to predict with confidence the existence and properties of a series of antibodies. When such a statement can be made regarding any theory it is obvious that, if not complete, it approximates to the truth. Nevertheless, as indicated in our discussion of the toxins, in certain of its aspects-in matters which are of some importance although not fundamental-we are not quite satisfied that the theory perfectly represents the relationships between the toxins and the cell molecule. Just as we doubt whether primarily, at least, the bacterial toxins become directly anchored on to the cell molecules or biophores, so in cytolysis we doubt whether, under any circumstances, there is direct union between the biophores of the animal cell and the bacterial or animal cell introduced into the system, even when that foreign cell is ingested by a phagocyte. We doubt, that is, the correctness of Ehrlich's diagram which we have reproduced. That, it is true, is merely a diagram—a graphic simile—and, like similes in general, must not be tested too severely. It expresses, however, a relationship which we are convinced does not exist. As indicated by what occurs in ordinary gastric and tryptic digestion, as indicated also by the formation of digestive vacuoles around particles of foodstuff ingested by the phagocytic cell, the dissolution of foreign protein molecules is mediate and not immediate; it is always by free side-chains, or haptines, and it is the loss of these side-chains, the condition of partial unsatisfaction thereby produced, and not the direct stimulus of direct contact and combination of the foreign matter into the protoplasmic molecule, that affords the stimulus for the production of new side-chains and so the active development of immunity.

Constantly in striving to comprehend the processes which are concerned in the destruction of bacteria and their products, the neutralization of cell products, and development of immunity, we find ourselves brought back to the fundamental fact that these constitute but special cases of the dissociation of foodstuffs, of assimilation and digestion. The processes which occur when a cell destroys an ingested microbe must be identical with those which occur when it digests any other foreign matter, and bacteriolysis in the body fluids must be brought about by procedures of the same order as occur when fibrin is digested in the gastric juice. If we admit that enzyme action is throughout operative in the one case, we must accept its operation in the other. That there may be—nay, that there are—different grades and orders of enzyme

IMMUNITY

action must be freely acknowledged; the mode of action of ptyalin upon starches, which appears to be direct, is different from that of enterokinase upon proteins, which requires the intermediation of trypsin to render it complete.¹ It follows, therefore, that we must regard all these processes as examples of one or other stage of enzyme action, that we must conclude that toxins and cytolysins are enzymes, and that the data we have acquired regarding these and their mode of action reciprocally gain their explanation from what we know concerning the laws of enzyme action, and advance our knowledge of these laws.

This, it may be said, is proceeding farther than many are prepared to advance at the present time. There is, indeed, a remarkable nervousness exhibited toward the proposition that toxins are ferments, which is to be explained, in the first place, as due to the fact that we do not know the composition and structure of either the one or the other, and in the second, to the extreme specialization of modern science, so that we have at the present time three distinct groups, distinct in methods and distinct in aims, working at this very subject, each, with the best of good-will, recognizing with difficulty what for the other two groups are the points of fundamental importance. The chemists are interested in bringing enzyme action into line with the catalysis of inorganic matter, and at the present time the trend of their observations upon the latter is to indicate that it is of the nature of a physical contact action, and not of a true chemical combination, however temporary; they deny also that enzyme action is chemical, and, so, logically, when it is demonstrated that toxin combines with antitoxin, they are bound to deny that the toxin is an enzyme. The physiologists are torn asunder between their attempts to render physiological chemistry an exact science and to follow the guidance of the Lure inorganic chemists on the one hand, and their affinities with the pathologists and bacteriologists on the other; the pathologists and bacteriologists by methods widely different from those employed by the chemists, although at the same time exact and of a subtlety unapproached by chemical tests, have accumulated a vast mass of important data on enzymes, toxins, cytolysins and so on, and their powers of combination, and as these hang together and do not harmonize with the deductions drawn by the chemists as to enzyme action from the study of inorganic compounds, they are content to group their data as a class apart and to continue gathering more facts and applying the same.2

¹ It is usual to express this reaction in the reverse manner. The observations of Bayliss and Starling show learly, as opposed to Delezenne, that the activating substance is the kinase, which has properties corresponding to those of the bodies we recognize as ferments.

² The present confused state is well exemplified by Benjamin Moore's always suggestive—but it seems to us not always wholly logical—recent article (*Recent Advances in Physiology*, edited by Leonard Hill, 1905). Moore concludes that "it is most probable that the influence of the enzyme as an energy transformer is one of a physical character; at any rate, the formation of chemical compounds must be taken as unproved." At the same time, he accepts the fact that in all cases the same time, he accepts the fact that in all cases the same time.

TOXINS AND ENZYMES

We, on the other hand, cannot but see, in all the evidence that has been brought forward regarding enzyme action, the aptness of E. Fischer's conclusions in his studies upon fermentative glycolysis, namely, that the enzyme must be regarded as fitting into the molecule group it dissociates as does a key into a lock, and see here not the evidence of contact action, but of chemical combination, and see this same striking specific action in connection with toxins and antitoxins. We admit that the specificity is, as regards the substratum as a whole, not absolute. The same enzyme may act upon two different bodies, or two different enzymes can dissociate one substance, as, for example, both invertase from yeast and amygdalase from the bitter almond will act on the glucoside, amygdalin. But what is specific in all these cases is that for common action there must be present identical molecule goups to be acted upon. We need not adduce instances in which this is demonstrated to be the case in connection with toxins and antitoxins.

In short, the remarkable parallelism between the toxins and the enzymes, the proved existence of toxoids and zymoids, of antitoxins and anti-enzymes, of natural and experimentally acquired anti-enzymes (Bayliss), and other natural and acquired antitoxins, of complements and kinases, the evidence that a minimal amount of enzyme or toxin will, under favorable conditions, in the one case convert a maximum amount of the substrate, in the other case so induce a dissociation of the substance of certain cells that death ensues, whereas under unfavorable conditions both enzyme action and toxic action can be arrested; the arrest of action of both by the products of dissociation; the exact quantitative neutralization of enzyme by anti-enzyme, toxin by antitoxin —all these facts indicate that we are dealing with one common group of

products of reaction of an enzyme exercise a protective action against rise in temperature, *i. e.*, as Vernon has pointed out (Jour. of Physiol., 27; 1901; 288), whereas an active trypsin in nearly pure state is destroyed by a temperature of 38° C.; if protein on which it can act be present, a much higher temperature of 38° C.; if protein on which it can act be present, a much higher temperature of phenomenon save the formation of a compound between the enzyme and either the protein or its dissociation products. And elsewhere, discussing Adrian Brown's observations, which indicate that the enzyme forms a compound which persists for an appreciable time, all that he can suggest is that the 'time interval of combination be regarded as constant in all cases," *i. e.*, he admits here that combination does occur. The combination of toxin with antitoxin he does not take into consideration.

It is interesting to note that Moore, in his very full study of the properties of catalysers, lays down that there is a clear distinction between inorganic eatalysts and organic enzymes. He lays down that, while similar in not being altered by the reaction they set up, the enzymes differ in that they require external energy in order to do their work; instead of causing energy to be given out by the ehemical system, they cause the system to take up energy, and, lastly, they cause a movement away and not toward the equilibrium point. He admits that all are not prepared to accept all these conclusions.

IMMUNITY

substances and with a group that act not by physical contact, but by chemical combination. ${}^{\rm t}$

Not knowing the structure of either class of enzymes or toxins, we must be content for the time to employ symbols to indicate the different stages of the process, after the manner introduced by Ehrlich. The molecule endowed with enzyme or toxic properties we must indicate as possessing surface (or cell complex) accurately adjustable to a corresponding surface on the fermentescible substance—Ehrlich's haptophore. We must admit with Ehrlich, also, the existence of a zymophorous or toxophorous moiety, and our study of enzyme action in general suggests how this acts. A chemical study of the stubstrate or fermentescible substance and the products of action demonstrates to us that enzyme action—as a general rule—proceeding in the one direction acts by hydrolysis, proceeding in the other results in the union of two molecules with the liberation of a molecule of water. A familiar example of this first process is the following:

$\begin{array}{l} C_{12}H_{22}O_{11}\,+\,H_2O\,=\,C_6H_{12}O_6\,\,\pm\,C_6H_{12}O_6,\\ 1 \mbox{ molecule of dextrose } +\,H_4O\,=\,2 \mbox{ molecules of glucose}, \end{array}$

Obviously the molecule of maltose cannot be split into two equal portions; either that molecule is really a multiple formed of repetitions or polymerization of $C_{12}H_{22}O_{11}$ to some power of 2, or, if single, it becomes divided into 2 unequal portions. The simplest case of such unequal division under the action of an enzyme is:

$$\begin{array}{rcl} C_{12}H_{22}O_{11} \,=\, C_{6}H_{11}O_{5} \,+\, OH \,=\, C_{6}H_{12}O_{6} \\ C_{6}H_{11}O_{6} \,+\, H \,\,=\, C_{6}H_{12}O_{6} \end{array}$$

namely, that the enzyme splits the dextrose molecule into two moieties, one of which has positive, the other negative affinities, which when separated attract, the one a basic hydroxyl ion, the other an acid hydrogen ion. Our conception of the enzyme molecule must, therefore, be that it acts, whether as a base or as an acid, attracting and detaching one moiety of the fermentescible molecule. Take, for example, that it acts as an acid, then it detaches the complex $C_{\rm e}H_{\rm in}O_{\rm e}$; but so soon as it accomplishes this, an H ion free in the solution, which could not act on the complete molecule, exhibits a greater affinity for the moiety than does the ferment, replaces this, and so the enzyme is free to act upon a second molecule of the substrate. Following Ehrlich's method, we can express reactions of this order as in Fig. 170.


Or otherwise even in the simplest enzyme action we must recognize the cooperation of three factors: (1) the enzyme, (2) the fermentescible substance, and (3) the recipient. The existence of zymoids shows that the body or zymophorous portion of the enzyme may be so altered

³ But it may be said the action of toxin on antitoxin is quite different from that of enzyme on its substrate. Certainly it is. It is not here that the parallelism comes in. What is comparable is the relationship *between enzyme and anti-enzyme action*.

ANTITOXINS AND ANTI-ENZYMES

that whereas the haptophoric portion still is able to be attracted to and attach itself to the fermentescible molecule, the zymoid molecule as a whole is unable to split the latter, which, being already satisfied, cannot now be acted upon by other and active enzyme molecules.

In the more complicated enzyme action, such as has been demonstrated to occur in tryptic (proteolytic) digestion, following Bayliss and Starling we must regard the kinase, or, perhaps, more strictly the kinase plus the trypsin, as playing the part of the enzyme; the trypsin alone cannot spilt up the protein molecule and detach a peptone group; it has to be reinforced by the kinase, which, in its turn, cannot directly associate itself with the protein, its haptophoric group not corresponding to any of the haptophoric groups of the proteid.



Simple enzyme action: F_i the enzyme molecule, has affinity for and detaches A_i a side-chain of a protein molecule, forming a temporary combination with it. When $A \cdot F$ is free the recipient Ghas a greater affinity for the side-chain moiety $A \cdot F$ and combines with it, the enzyme molecule Fbecoming detached and ready to dissociate a second similar side-chain.

The difficulty experienced-and this we would emphasize-in applying ideas gained from the study of enzymes to toxins and antitoxins is that we are continually apt to confuse the latter with the fermentescible substances. The two are distinct, as will easily be recognized when we recall that the rennet ferment acts upon the caseinogen of milk, converting it into casein, but is completely arrested in its action by antirennin ("antilab"). In the one case the enzyme molecule acts as a carrier, as indicated in our diagram, in the other it becomes fixed and exercises no dissociative effects. Applying Ehrlich's terminology, the fermentescible and the anti-enzyme molecules possess identical haptophorous groups; the enzymes can become attached to both, but in the first place the affinities between enzyme and anti-enzyme are the greater, so that when both anti-enzyme and fermentescible molecules are present in a solution the enzyme is attracted to the former and not to the latter, and in the second place, when so attracted it is fixed and, what is more, is unable to dissociate the anti-enzyme molecule. The process of junction between fer-


IMMUNITY

ment and antiferment and between toxin and antitoxin is associative and self-limiting; that between ferment and fermentescible substances dissociative and recurrent, only arrested by the accumulation of the products of the reaction.

It is this distinction which is not clearly drawn by Professor Ehrlich,¹ and to us it appears to be of the very highest importance. If toxins are bodies of the same order as enzymes (and we have indicated that we cannot conclude otherwise), it follows that the antitoxic side-chains developed in reaction to the presence of toxins are not identical with those dissociated by the toxins, whose dissociation leads to the symptoms of disease. Or, otherwise, something in addition to the mere proliferation and overreplacement of the side-chains attacked by the toxin molecules is necessary to explain immunity.

18

F10, 171



GA (the side-chain A combined with the recipient G), when discharged from the cell into the surrounding fluid as an antitoxin molecule, is dissociated by the enzyme or toxin molecule F, which thus joining with A becomes neutralized.

We would tentatively suggest that but a slight although important modification of Professor Ehrlich's conception will meet the case. As we have noted in passing (p. 573), Professor Ehrlich in his conception of the junction of toxins and antibodies takes no note either in toxin or antibody of the group of junction with the original protoplasmic molecule. We would suggest that when dissociated, there must be in this position of the side-chains complex an unsatisfied or unsatisfiable affinity. If, now, we regard the toxin molecule when it gains entrance into the cell not as becoming attached to the biophoric or protoplasmic molecule by means

¹ It is but right to state that throughout Professor Ehrlich is extremely reserved in his judgment regarding the relationship of toxins to enzymes. Wherever he approaches the subject he is most careful to leave the question an open one. As we have indicated in several places, M. Metchnikoff is very definite in his conclusions that the complements (cytases) are enzymes and that the whole process is one allied to the digestive processes.

ANTITOXINS AND THE SIDE-CHAIN THEORY

of one of the side-chains of same, but, on the contrary, as detaching that side-chain, we can solve the difficulty. Namely, we can regard it as in this relationship, within the cell, acting as an enzyme, delivering over the detached side-chain to a recipient which has greater affinities for it, and itself becoming free to act upon another side-chain.

It is this side-chain plus recipient which now becomes the antitoxin. In the cell itself it cannot act, the toxin having greater affinities for the still adherent similar side-chains of the protoplasmic molecules, unless the point is reached when equilibrium is established by the accumulation of products of the enzyme action, and by the overproduction and discharge of side-chains of the particular order into the cytoplasm or paraplasm. But when the excess of such side-chains plus recipients is discharged into the blood stream, then any circulating toxins, not having the greater attraction of the adherent side-chains of the cells they specifically influence, join with these antitoxins and become neutralized and are not attracted to and taken up by the cells.

Along these lines, and by considerations of this order, we most simply indicate the relationship between enzyme and enzyme-antienzyme action, toxic and toxin-antitoxin action. We regard the same orders of cell groups as being involved in both cases, we call in no external factor save the "recipient" which is demanded by any chemical theory of enzyme action, which recipient must be some simple but active ion present in all solutions in which the enzyme or toxin is able to act.

We might very considerably expand the considerations here brought forward, but to do so would render our treatment of this subject out of all proportion to the rest of the work. It is only necessary to point out that the same considerations can be applied to the case of cytolysis, save that we are not as yet fully prepared to grasp the relationship of the lipoid bodies to the processes of hemolysis, fixation of complement, etc. The indications are that in all other respects the data which have been brought forward in support of Professor Ehrlich's theory can, so far as we can see, be applied to this modification of the same.¹

¹ Works of Reference.--Among the more important and authoritative works upon the subject of immunity are the various articles by the leading German workers on the subject, and by Metchnikoff, in the fourth volume of Kolle and Wassermann's "Handbuch der pathogenen Mikroorganismen" (1904); Metchnikoff's "L'Immunité dans les Maladies Infectieuses," Paris, Masson, 1901, translated into English by Binnie, Cambridge University Press, 1905; Aschoff, "Die Seitenkettentheorie," Jena, 1901; von Dungern, "Die Antikörper," Jena, 1901; Jacobi, "Immunität," Heidelberg, 1906. In English, Bolduan has published a clearly written "Immune Sera," New York, Wiley, 1907, and has translated most serviceably Ehrlich's "Collected Studies upon Immunity," New York, Wiley, 1907. Arrhenius' lectures upon "Immuno-Chemistry," delivered in San Francisco (Macmillan, 1908), have been painfully translated, and while full of important matter, are scarcely to be recommended to the beginner. Another excellent account of the main phenomena of immunity is afforded (by the late Dr. Ricketts) in a series of articles contributed to the Journal of the American Medical Association from January to July, 1905, and republished in separate form at a very moderate price. Emery's "Immunity and Specific Therapy," Lewis, London, 1909, is the fullest and most recent English work on the subject.

CHAPTER X.

SYNCOPE, SHOCK, COLLAPSE.

SYNCOPE.

WITH the reactions to injury must be considered a series of conditions of another order, conditions which are most often brought about by injury, which, nevertheless, just as we saw happened in the case of inflammation, may be wholly initiated by the higher nervous centres without any local tissue disturbance being at fault. These conditions are syncope, shock, and collapse. Some writers include the last two as synonymous; the symptoms are almost identical, but, following Cobbett,¹ we shall make some distinction between them. Admittedly, however, we have not a full grasp of the etiology of these states; much has still to be determined, even if, of late, material advance has been made.

Syncope, or fainting, is the slightest of the three conditions. The face suddenly becomes blanched, the pulse small, rapid, and at times almost imperceptible; a brief period of giddiness is followed by complete unconsciousness, the individual falling in a lax heap, often without time to hold on to a support or make any preparation. The condition is of relatively short duration.

In one series of cases we note nothing beyond the mechanical filling of the abdominal vessels, as when fainting supervenes upon emptying the distended bladder. Evidently closely allied to this is the syncope which at times follows the sudden change from to the supine to the erect position In another, injury to or strong stimulation of sensory nerves is the exciting cause—pain of various degrees. Numerous cases are of purely emotional origin, and that not only in the weaker sex or in those of weak health. I recall vividly a football match years ago, and the scattered, dropping, like pole-axed steers, of close upon half a score of the undergraduate onlookers, consequent upon the sharp loud snap of a leg bone of one of the players and the sight of his fall helpless to the ground.

SHOCK AND COLLAPSE.

In shock and collapse we have severer states. In extreme cases death ensues, and this in the case of shock with absolute suddenness. Sir Lauder Brunton cites the case of a mock trial conducted by certain Aberdeen students upon an obnoxious janitor, who, having been led to

¹ Allbutt's System of Medicine, 3: 1898: 320.

SHOCK AND COLLAPSE

the block and there struck on the neck with a wet towel, was taken up dead. In less extreme cases there is recovery, but this after hours and days, instead of minutes. There is in both conditions blanching, with rapid, feeble, and, it may be, imperceptible pulse; the eyes become sunken, the cheeks also, the cheek bones prominent (*facies hippocratica*). There is complete muscular laxity; the breathing is irregular and oppressed, the breath cold; the external temperature of the body is very noticeably lowered. The pupils are dilated; retching and vomiting are frequently prominent. The patient lies limp and regardless of his surroundings. Unlike syncope, neither in shock nor collapse is there complete unconsciousness. Upon rousing the patient, the answers, if show and obtained with difficulty, are quite rational; but all volition is abolished. In short, there is a marked general depression of function.

Etiology.—If we attempt to classify the conditions under which this syndrome presents itself, we find the following:

1. **Operations**, or wounds, associated with injuries to nerves. These may be: (a) Peripheral, affecting the nerve terminations, as in the shock that follows extensive burns, sharp blows upon the testicle, operative exposure and irritation of the peritoneum, irritation of the periosteum, as in amputation of the thigh (it has been found that the bone substance, as such, is comparatively insensitive, the periosteum very sensitive). (b) In continuity, as after severance of a large nerve, such as the sciatic. (c) Central, as after operations upon the brain and removal of cerebral substance. According to Howell,⁴ the last of these (in the dog) is the most effective in producing this train of symptoms. With regard to all these it may be noted that pain is not essential. Shock may follow operation conducted under anesthesia sufficient to abolish all sensation.²

2. **Pain**—Relatively intense irritation, without gross injury to peripheral nerves, may induce not merely syncope, but the more profound state here described. Closely allied is:

3. Emotional Disturbance.—To this we have already referred.

4. Severe hemorrhage, whether (a) external, or (b) internal into the cavities of the body, induces a similar train of symptoms, as does

5. Loss of fluid from the blood through (a) persistent vomiting (as in hyperemesis gravidarum), or (b) excessive diarrheea (as in cholera).

It is these last two conditions that we distinguish as *collapse*; all the former we include under the term *shock*. The only gross clinical difference between the two is that collapse in general is of relatively gradual development, whereas shock is of rapid onset. We shall point out shortly another partial distinction.

Studying these three states, we find one feature common to all, namely, a combination of cardiovascular disturbance, with grave arrest of cerebral


² For full studies upon the relative liability to shock following operations upon different regions and viscera, the student is referred to Crile's "Surgical Shock," Lippincott, 1899, and "Blood Pressure in Surgery," ibid., 1903.

¹ Contributions to Medical Research, dedicated to Victor C. Vaughan.

SYNCOPE, SHOCK, COLLAPSE

activity. Evidently the cardiovascular disturbance may be primary, as where fainting supervenes upon emptying the bladder, or the sudden assumption of the erect posture, or, in cases of collapse, where there is great loss of blood or of its fluid constituents. In the majority of cases, however, the indications are those of primary irritation or disturbance of the higher centres, whether direct or reflex.

The primary vascular disturbance is easily understood. We know that the vessels of the splanchnic area are capable of holding all, and more than all, the blood of the organism. The vessels of the liver alone are so distensible that they can contain all the circulating blood. That they do not is due (1) to the tone of the abdominal walls, whereby the viscera are definitely compressed, and (2) to the vascular tone, whereby under normal conditions, the arteries more particularly, but also, as Goltz has shown, the veins, are in a state of partial contraction. Sudden removal of fluid from the abdominal cavity may so lessen the pressure upon the visceral veins that these may undergo rapid dilatation, the blood pouring into them to fill the void, and this to such an extent that, more especially when the individual is in an erect posture, the blood is



Schema of cardiac conditions in the ''Klopfversuch'' experiment on the frog: a, normal state of filling of inferior vena cava and heart; b, dilatation of splanchnic veins (erect position), heart in consequence empty; c, heart filled and circulation restored when the mimal is placed on its back. (Sir Lauder Brunton.)

drained out of the vessels of the upper half of the body, so that little or none enters the right side of the heart. The result is not merely greatly lowered blood pressure in the radial and other arteries, but the brain is profoundly affected; the intracranial pressure is lowered, not merely by drainage away of the venous blood, but by lack of arterial blood supply, and this lack of intracranial pressure, coupled with the anemia and lack of blood supply, is adequate to induce the unconsciousness seen in fainting. One can, indeed, produce temporary insensibility by compressing both carotids.

This is the simplest case; in other cases of syncope, brought on by nervous influences, it is evident that a similar disturbance of the circulation is brought about by nervous stimulation. We can, indeed, as in Goltz's well-known "Klopfversuch," induce reflexly this syncopal condition. It is but necessary, as Goltz showed, to tap a frog over the intestines to induce an arrest of the circulation, and, as he showed, the arrest is accompanied by an accumulation of the blood of the body in the abdominal veins. Hold the animal erect, and the exposed heart is seen to be still beating, but devoid of blood; place it on its back, or

THE "KLOPFVERSUCH"

with the head lowered, and the heart and the arteries become filled and circulation is resumed. Here I may note that the late Professor Roy and I⁴ observed time and again how rapidly and extensively the arterial blood pressure could be raised by temporary compression the abdominal area, whereby the blood becomes driven out of the abdominal veins into the right heart. It is a matter of familiar knowledge that recovery from a faint is best brought about by placing the individual in a recumbent position. It is not usual to reinforce this by steady pressure, applied to the region of the waist, although our experiments indicated that this procedure is likely to be even more effective.

As Sir Lauder Bruinton² points out, a fuller study of the Klopfversuch indicates that, reflexly, by a tap on the abdomen, one, or both, of two effects may be produced—namely, inhibition of the splanchnic vasoconstrictor centres (whereby the abdominal veins become dilated), or stimulation of the cardiac inhibitory centre (vagus), whereby the beat of the heart is arrested. There are indications that in man either of these events may lead to syncope—and to sudden death. In deaths during operation under chloroform, which appear to come under this category,³ it would seem that either or both of these events may be responsible; at times death is sudden, through arrest of heart action; at others, the heart continues to beat feebly for some time after the pulse has become imperceptible (vasodilatation).

What, then, is the difference between temporary syncope and more prolonged shock? In both we have indications of local splanchnic vascular dilatation and of cerebral disturbance, but the one condition is temporary, the other of long duration; the one is accompanied by complete unconsciousness, the other by depression merely of cerebral functions. It may be that we are not yet prepared to give the full answer. There are, it appears to me, sundry suggestive observations pointing to the solution of this question. We have already referred to the striking depression of all functions seen in shock. There is no perverted metabolism; the blood, it has been noted, is not toxic. But it is found that easily soluble drugs, such as alcohol, ether, strychnine (Roger), produce little effect; nay, more, they are not absorbed by the cells, and, with recovery from the state of shock, such drugs given in excess have then produced their physiological symptoms, and have even caused death. As pointed out by Wesley Mills4 in connection with surgical shock, food may remain the whole day in the stomach

¹ On Waist Belts and Stays, National Review, 1889.

² Practitioner, 11:1873:246. Collected Papers on Circulation and Respiration, 1906:402.

³ Such deaths, it has been noted by several observers, are apt to occur where in trivial operations the patient is not brought fully under the influence of the drug—when the vagus centre is still capable of reflex irritation, but anesthesia has proceeded sufficiently far to arrest the splanchnic vasomotor constriction which normally follows severe stimulation of a sensory nerve; normally, these antagonize each other.

⁴ Brit. Med. Journ., 1910; ii.

undigested. The mere act of dilatation of the vessels must favor rather than arrest the diffusion of soluble drugs; that they do not act would seem to indicate an actual inhibition of the tissue cells and arrest of cellular activities. Even in syncope, as first noted by John Hunter, the venous blood may become arterial in hue, but, as Sir Lauder Brunton explains, this is seen also in other cases, in which there is dilatation of the_arterioles; it is not necessarily a sign of arrested cell activities in the capillary areas.

Yandell Henderson¹ ascribes the sequence of changes in the blood pressure, the filling of the splanchnic veins, the tachycardia and weakened heart beat not to primary vasomotor influences, but to acapnia, or the effects of diminished CO, in the blood. Exposure of the viscera, he shows, is accompanied by exhalation of CO₂, and this and mere aëration of the abdominal cavity afford the anatomical and clinical picture of shock. Bathing the exposed viscera with salt solution saturated with CO, prevents the same. We must, however, hold that shock is more than acapnia, even if this explains the main symptoms. The dominant feature is arrest of bodily function from profound nervous disturbance. Mere splanchnic vasodilatation and cerebral anemia may easily be recovered from; such, and such only, occur in syncope. In shock, if in many cases the splanchnic dilatation is secondary to lessened CO₂, and in other cases (as in the klopfversuch) is reflex, the underlying change is something more; it is a general inhibition or arrest of the activity of the higher nerve centres.

It would seem that we have evidence of this nervous inhibition in the "spinal shock" of the lower warm-blooded animals. Divide the spinal cord in the dog in the lower thoracic or lumbar region and the reflexes of this area below the region of section disappear. The same is true in man, but, unlike what happens in man, gradually these reflexes in the dog again make their appearance—although the cord still remains divided. For a time the centres do not react to stimuli. We have, that is, a profound inhibition of centres other than the vasomotor, which is, in turn, recovered from. In man the spinal centres are so largely under control of those in the brain that, with central stimuli cut off, there is no assumption of automatic independent action.

Admitting this, it must also be admitted that the circulatory changes, if secondary, are the most prominent features in shock. And here it is not merely the diversion of blood into the splanchnic area, and the weak heart action that accompanies it, but, as shown by Roy and Colbett,² this diversion leads to changes in the blood itself and in the tissues. By the most ingenious device of simultaneous testing of the specific gravity of the blood and tissues these observers demonstrated that in the course of shock the specific gravity of the blood becomes reduced, while that of the tissues increases. This can only be interpreted to mean that during the course of shock the volume of the blood is increased,

¹ Amer. Journ. of Physiol., 21:1907:126, 23:1908-09:345 and 24:1909:66.

² Allbutt's System of Medicine, loc. cit.

and that at the expense of the tissue fluids. This is in harmony with the observations made by Sherrington and others upon the specific gravity of the blood following on hemorrhage—loss of blood is followed by a rapid fall in the specific gravity of that still remaining in circulation, and a similar rise in the specific gravity of the tissues. In other words, actual or rélative loss of blood in a great part leads to a protective passage of body fluid into the bloodvessels, tending to preserve the circulation. This loss of fluid on the part of the tissues explains the development of the facies Hippocratica, the hollow orbits, the sinking in of the cheeks.

In collapse due to hemorrhage there is a similar reduction in the specific gravity of the remaining blood; in that due to profuse diarrheea or vomiting, while the specific gravity of the tissues becomes raised, that of the blood becomes raised also. 'The loss of fluid from the tissues into the vessels is not sufficient to counterbalance the loss of fluid constituents from the blood into the intestinal tract.

Collapse thus differs from shock in mode of causation, rate of onset. and (in some cases only) in the specific gravity of the blood. If the main symptoms are identical, and if we ascribe shock to widespread cerebral depression, then a like cerebral depression must be present in collapse. This we firmly believe to be the case; only, while in shock we regard the nervous disturbance as initiating the depression, in collapse we see that the depression of the higher centres is secondary to the continued cerebral anemia. We confess, however, that we are not inclined to draw too sharp a line of demarcation between these two states, while, further, we cannot but recognize that there are states in which shock and the blood changes characteristic of collapse are capable of being combined. Thus, as Crile has pointed out, one of the most sensitive tissues in the organism is the parietal peritoneum; severe handling of this in laparotomies is apt to bring on rapidly the syndrome already described. Nevertheless, Cobbett points out that prolonged exposure of the abdominal cavity and of the peritoneum is followed by rise in the specific gravity of the blood; there is evidently developed not merely a congestion, but a discharge of fluid from the vessels of the affected area.

Lastly, a word must be said regarding the condition of "shock with excitement," rather than depression; this, accepted as an allied form by the writers of last century, is apt to be dismissed by more recent writers as a bastard due to coincident stimulation of the cerebral centres by bacterial products. This we doubt. We have known it develop rapidly after a moderately extensive burn of the second degree affecting the front of the neck and upper part of the chest, the patient dying after six days, with symptoms of cardiac failure. A condition of almost maniacal, uncontrollable excitement supervened within an hour. The patient in this case presented a history of long-continued mental instability. We would only point out the familiar fact that sensory stimuli, which in one series of cases lead to inhibition of sundry centres, in another series of cases may produce, on the contrary, irritation of those same centres.



PART II.

THE TISSUE CHANGES.

CHAPTER XI.

THE PROGRESSIVE TISSUE CHANGES: HYPERTROPHY.

THUS far we have regarded morbid processes essentially from the aspect of their causation, discussing the changes brought about in the tissues by one or other order of disturbances. But these processes may be regarded from another point of view—one which, for an ordinary grasp of the subject, is most important—that, namely, of distinguishing and classifying these processes according to the alterations they produce in the tissues themselves. Or, briefly, while up to the present we have regarded these processes from the point of view of the irritant, from now on we shall regard these from the point of view of the tissues

For any tissue, and for the cells of that tissue, there is a certain normal condition; within the limits of that condition we have the healthy state; outside of these limits, disease. Studying what are the factors determining this state of health, we recognize that these are two in number: (1) the nutrition of the cell, and (2) the functioning or activity of the cell. These are closely dependent the one upon the other. The more we study the more we realize this interdependence.

It is obvious, in the first place, that for the cells to remain healthy and active they must have nourishment; otherwise the destructive processes associated with life will exceed the constructive, and atrophy will ensue, and eventual death. Further, this nourishment is not a mere passive process. *Absorption is active*; the food taken in before it can be used up in the manifestation of the various forms of energy and of construction, must become part of the protoplasm of the cell.

The mere presence of a food molecule within the cell neither yields up energy nor increases the amount of living matter. It has to undergo dissociation, and some portions of the molecule, whether temporarily or more permanently, must become directly bound up with the living matter of the cell. And here, while discussing nourishment, it has to be remembered that *quality* as well as *quantity* of the absorbed material has to be taken into account. Molecules of one order may supply energy to the cell complex, whether active or latent (bound up in the assimilated material); of another order, may so act upon the cell

THE PROGRESSIVE TISSUE CHANGES

substance as to withdraw energy or inhibit molecular activities, acting as toxic agents.

As regards function we recognize that up to a certain point the more active the cell the more extensive are the chemical changes proceeding in the protoplasm of the cell, and, with this, the more active the absorption of new material. Within certain limits, that is, increased activity becomes associated with increased assimilation. Contrariwise, diminished cellular activity demands diminished nourishment. Thus, the condition of the cell, healthy or otherwise, depends directly upon the functioning of that cell—a law applicable not only to the unit cell, but to the organism or individual as a whole.

As we have pointed out (p. 101), cell and organismal growth is intimately connected with the interaction of these two factors. Having already so fully discussed this subject of growth, we shall not enter into the matter in detail. We would, however, emphasize here the fact that a grasp of the factors influencing growth is essential for a proper understanding of the conditions we are about to discuss. That we possess a full knowledge of the factors underlying cell growth and cell shrinkage in all their aspects we do not in the least pretend to suggest; there is much that still remains to be determined. We do not know what it is that permits one cell, like the bird's ovum, to accumulate an enormous store of food material, and another, though bathed in nutritious fluid, like the bird's leukocyte, to remain small. We do not know what is the normal inhibitory force preventing a cell from developing, it may be for months or years, or the force which suddenly stimulates that cell to undertake active growth and proliferation. At most, obscurely, we see a progressive unfolding, as it were, of environmental conditions, which modify the forces acting upon that cell from without. We are still debating, without having arrived at any sure conclusions, what are the forces which permit the regeneration of a part to proceed until the new development may reach, but does not exceed, the size of the lost organ. / There are, however, certain general principles which we can recognize as being in action: that inadequate nutrition or lack of exercise of function, either of them, may lead to inanition and shrinkage of the cell unit to such a point that function is wholly arrested, and death may ensue; that excessive activity may lead to such using up of the cell substance that the absorption cannot keep pace with the disintegration, from which cause, also, cell death may ensue; that there develops an equilibrium between assimilative and functional disintegration such that the normal fully developed cell may remain for long in statu quo; that stimulation and functional activity somewhat above the normal, accompanied by adequate nutrition, may lead to growth. until again an equilibrium is reached (p. 109); that, also, an equilibrium tends to become established between cell mass and cell surface, nuclear mass and nuclear surface of such an order that the accumulation of living matter within cell and nucleus is beyond a ceratin point selfinhibitory (p. 36), or is apt to be followed by cellular and nuclear proliferation (p. 111); that storage of energy, which is implied by growth

WEIGERT'S HYPOTHESIS

and accumulation of cell material, and dissipation of energy, which inevitably accompanies functional activity, are opposed processes, which can only occur simultaneously within narrow limits (p. 103); that specialization of the cell through and for the performance of function in itself limits growth and proliferative capacity (p. 140); so that we find that the highly differentiated cell, as such, does not proliferate, and that the actively growing vegetable cells of the organism are either those which have never undergone specific differentiation to any degree. or, having been differentiated, have reverted to the undifferentiated, vegetative type. These are general conclusions which it is well to keep in mind. Here, as bearing directly upon the causation of pathological overgrowth, it is necessary to call attention to a controversy which dates back for its origin close upon twenty years, and which is still smouldering, upon the primary cause of that growth. We confess that we do so with some impatience. It seems to us that it is an outcome of overspecialization; that greater breadth of view, in the first place, and comparison with physiological growth in the lower and particularly the unicellular organisms would have indicated that the position on the one side was untenable. Nevertheless, the debate has had the good result of leading to many valuable observations.

In 1889 Weigert¹ laid down as the result of general consideration of the relationship of functional and vegetative activity, that pathological tissue growth "only occurs when, from any cause, there is disturbance of the reciprocal normal equilibrium of the tissue and tissue elements, and when the physiological restraint is removed which one tissue element exercises upon another." There cannot, that is, be a direct stimulus of growth outside the cell; the tendency to grow is within the cell, and this is restrained by environmental conditions; these environmental conditions must be removed, the resistance diminished, and then proliferative changes show themselves. So great was Weigert's ability and authority, that his dictum was for a considerable time generally accepted, and conditions which, prima facie, appeared to be examples of growth due to stimulation were by hook or crook explained according to his hypothesis, even to the extent of explaining the giant cell as primarily due to defect in the cytoplasm, whereby the restraint was removed from the nucleus, so that now this proceeded to undergo direct division.

Ribbert² has expanded this hypothesis; he acknowledges that many factors bring about restraint of growth, not alone cell pressure, but relationship to the vessels and nerves, and differentiation for functional purposes; in short new-growth on the part of the cell is regarded as being initiated by one and all the disturbances in function or cell relationship which disturb the equilibrium between the cells forming a tissue. He, too, it will be seen, denies that direct stimuli from without can initiate growth.

¹ Fortseh, d. Med., 1889; No. 16. This, it may be added, is in opposition to Virehow.

² Virch. Arch., 150; 1897; 391.

THE PROGRESSIVE TISSUE CHANGES

100

But what is the cause of growth in unicellular organisms? Is it not obvious that when an amœba ingests a food particle it does so in consequence of a stimulus from without, that such stimulus precedes assimilation, and it is the assimilation that is the first step in growth and proliferation? Studying also these lower forms, animal and vegetable, we observe that physical and chemical agencies, acting from without, are capable of stimulating growth; that increased temperature renders it more rapid; that, as Jacques Loeb has shown, alteration of the surrounding medium will initiate the nuclear and proliferative changes in the ovum of lower forms of life, even in the absence of fertilization. And if thus clearly direct external stimuli can initiate growth in the lower unicellular organisms, what reasons are to be adduced against their doing so in the multicellular organisms? As a matter of fact, there are numerous instances of increased proliferative activity in the warmblooded animal that can only be explained as brought about by direct stimulus. These have been well summed up by Levin⁴ and by Marchand.²

How, as Marchand points out, are we to explain the regeneration of the red corpuscles and the appearance of nucleated erythroblasts in the red marrow, following upon extensive hemorrhage, by the Weigert hypothesis? There has been no alteration of the surrounding cells. Or how explain the hypertrophy and hyperplasia, the overgrowth in cases of exercise and increased work-work hypertrophy? As we have already noted, and as we shall note again in discussing work hypertrophy, we are compelled to recognize that, within certain limits, increased function and the stimulus thereto is followed by growth: that, though beyond those limits cell work leads to a greater dissociation than assimilation, within those limits it favors growth. Thus, increased strain leads to growth. There may be, similarly, what Marchand terms a *tactile stimulus*, instancing the thickening of the epithelium upon exposed surfaces, and, as well shown by Bizzozero and Penzo's³ experiments, increased temperature favors increased growth.

They showed that if one ear of a young rabbit be kept at 12° to 15° . the other at 37° to 39° C. for a fortnight, the latter might become 1 cm. longer than the former, and skin, hair bulbs, and glands showed pronounced proliferation.⁴ So, also, that if they caused symmetrical fractures of a metacarpal bone, warmed one extremity and cooled the other, in forty-eight hours in the region of the one fracture the periosteum showed abundant mitoses, while around the other there was none.

There are, also, the instances of chemical stimulation. How, for example, is the endothelial swelling and proliferation in the vessels in cases of inflammation to be accounted for by the Weigert hypothesis, or the marked swelling and proliferation of the endothelium of the lymph glands throughout the body, so abundantly demonstrated by

² Die Wundheilung, Leipzic, 1901: 89.

³ Gaz. Med. di Torino, 42: 1891: 242. ⁴Sacerdotti, by similar means, obtained increased growth in length of one lower extremity as compared with the other.

¹ Journ. Med. Research, 6: 1901:145.

FUNCTION AND NUTRITION

104

Mallory in cases of typhoid and by J. McCrae in cases of burns? This we can only ascribe to the circulating toxins. We can induce similar proliferation by the inoculation of toxins. How explain the growth of the mamma in pregnancy? There is here no removal of cell restraint (p. 363). How, lastly, by this hypothesis, are we to account for the fact that free cells remain small—that the different forms of leukocytes, for example, circulating in the blood do not actually proliferate, and that when we excise a part of the brain, for example, although restraint is removed, the nerve cells show no sign of proliferation?

We are forced to admit that there can come into action an external stimulus to active cell growth. This, however, by no means necessarily means that tissue tension is not a factor in arresting growth, or that the interrelationship of the different components of a tissue is not of influence; only, for example, so long as the capillaries develop or dilate at the same rate as the specific cells of a gland can those cells continue to proliferate. With Ribbert, we must acknowledge that these are factors; only, over and above these restraining influences, there may be a direct stimulus which at times is sufficiently powerful to neutralize that restraint.

It follows from the above considerations that there is a relatively large number of combinations of conditions which may lead either to cell overgrowth or to cell shrinkage and degeneration. Leaving out of account those conditions in which there is equilibrium between functional activity and nutritional supply, we may have:

1. Normal functional activity of the cells, with increased nutrition,

2. Increased functional activity of the cells, with increased nutrition and assimilation.

3. Reduction in the external forces inhibiting cell growth; diminished tissue tension,

4. Normal functional activity, with reduced nutrition.

5. Normal functional activity, with perverted nutrition.

6. Increased stimulation and functional activity of the cells, with relatively insufficient nutrition (including here overstimulation of the cells).

7. Arrest of function of the cells.

8. Increase in the external forces arresting cell growth.

These conditions, it will be seen, fall into two groups, which we may entitle the *progressive* and the *regressive* cell and tissue changes. A third group is to be noted in which we have to deal, not so obviously with changes in the living cell matter of the cell as with alteration in the paraplasmic matters stored within the cell. Such alteration, either of excess or defect, we find to be either due to, or to lead to, regressive changes in the cell substance proper; it is thus usual to include them among the regressive changes.


Lastly, there is the important series of cases in which we observe excessive cell overgrowth without our being able as yet to state with precision what is the primary cause. These we include, naturally, among the progressive changes, but with this admission.

Of these progressive changes we meet with several forms; these we will consider in order,

THE PROGRESSIVE TISSUE CHANGES

OVERGROWTH.

The overgrowth of a tissue in which the individual elements retain their physiological relationships and functions may be manifested either by an increase in the size of the individual elements-simple hypertrophy—or by an increase in the number of those elements—hyperplasia (or as some would term it numerical hypertrophy)-or in the two combined. Yet another condition of increase in the size of organspseudohypertrophy-must here be distinguished. In this we have no overgrowth in the specific elements; but, on the contrary, an



Longitudinal section through muscle of calf of leg in pseudohypertrophic paralysis. The muscle fibres exhibit atrophy, the increase in bulk is due to the excessive development of fat cells. (Orth.)

atrophy of the same, with replacement in excess by another tissue. Thus, in pseudohypertrophic paralvsis, the great size of the muscles is seen to be brought about by an excessive interstitial deposit of fat cells, with accompanying degeneration, and, in the main, diminution in the number and in the size of most of the muscle fibres. "Hypertrophic cirrhosis" is, in almost every respect, an unfortunate and misleading term; the condition most often indicated by this term is truly a pseudohypertrophy of the liver-or a hyperplastic cirrhosis or fibrosis of the organ.

In other words, in speaking of the hypertrophy (or hyperplasia) of an organ, it is, for the sake of clearness, necessary to regard the specific elements of that organ, and refer to them only. In the liver, for example, while connective tissue is a normal constituent of this or every other organ, it is but the framework -the liver cells are the important specific constituents.

Once again we have to deal with imperfection of our pathological terms. "Hypertrophy" is employed loosely and commonly to designate all forms of overgrowth in which the elements retain their physiological relationships and functions-no matter whether there is increase in size or number of the same. Even if we overlook this, the term is in itself indefensible; its etymological meaning is "overnutrition;" thus, in itself, it is false, for overnutrition is, we now see, not the essential cause of the condition it connotes. The term is, however, so generally employed that we cannot cast it off, but must continue to employ it, regardless of its primary significance.

Such overgrowth may be either inherited or acquired. We have

OVERGROWTH

already considered the first group, the general or local giantisms, as also the inherited overgrowth of individual tissues, when discussing anomalies (p. 227). The considerations then brought forward throw some light upon certain aspects of the acquired condition, which, first, we will pass in rapid review, and later discuss as regards the causes which have been in action leading to this development.

Acquired Overgrowths.—Of these the majority appear, as we shall point out, to come under the heading of functional or *work* hypertrophies, increased demands upon the tissue and increased activity, coupled with adequate nutrition, being the feature noticeable in their production; and in these we pass from examples purely physiological to those wholly pathological. Thus, at one end of the series must be placed the pregnant uterus.

1. Physiological Hypertrophy.—'The overgrowth, both hypertrophic and hyperplastic, of the uterine musculature during pregnancy is most remarkable. The plain muscle fibres, according to Kölliker, become seven to eleven times as long and four times as broad as in the resting normal uterus. Associated with it we note: (1) Increasing distension of the cavity of the womb by the growing embryo, with pressure upon the walls; (2) greatly increased blood supply; (3) initiation of muscular contraction (from a very early period in pregnancy palpation shows that the muscle undergoes slow periodic contraction).

Other causes of distension of the uterus—fibroids, retained menses which are unaccompanied by any pronounced increase in vascularity, result also in hypertrophy. The increased nutrition, therefore, cannot be regarded as the primary cause of overgrowth.

Next, we have a series of cases in the borderland between the physiological and the pathological: the blacksmith's arm and the excessive development of muscle by exercise—a development which may approach the abnormal.

Here there can be no question concerning increased functional activity, coupled, we may add, with corresponding increase in nutrition, for, as is well known, coincident with more active contraction of the muscles there is increased circulation through them.


Nor is muscle the only tissue that undergoes growth as a result of increased work. The strain brought to bear on the bones leads to increased growth of the tissue, showing itself more especially along the ridges and tuberosities of muscular attachment and muscular "pull." And, as we noted in discussing fibrosis (p. 451), the remarkable overgrowth of connective tissue following upon lymphatic obstruction, or in the walls of the bloodvessels following upon increased blood pressure and increased tension (when not extreme) comes in the same category.

2. Adaptive Hypertrophies.—Allied to the above conditions, and clear examples of functional hypertrophies, are what we would term the adaptive hypertrophies. When there is obstruction to outflow, the nuscular walls of hollow viscera are apt to undergo great hypertrophy.

Enlargement of the prostate, with obstruction to the passage of urine, or, again, stricture of the urethra, leads to great hypertrophy

THE PROGRESSIVE TISSUE CHANGES

of the bladder, the individual muscle fibres being found twice as broad as normal; this, again, when the obstruction is not extreme; where it is, hypertrophy gives place to dilatation of the viscus and thinning of the walls. In like manner, stenosis of the cosophagus induces hypertrophy of the upper cosophageal muscular coat; of the pylorus leads to hypertrophy of the stomach; narrowing of the rectum or of the bowel at any point to hypertrophy above that point. Quite the most oommon and characteristic example of this class is afforded by the hypertrophy of the heart which follows upon narrowing of the valvular outlets or obstruction to the arterial circulation. As Bollinger showed in the case of the Munich "beer heart," a like hypertrophy follows an increased load, *i. e.*, a larger amount of fluid to be propelled through the vessels.



A, normal heart of rabbit; B, hypertrophied heart of rabbit due to repeated inoculations with small does of adrenalin extending over several weeks. The adrenalin causes contraction of the arterioles, beightened blood pressure, and increased heart work. (From specimens by Prof. Klotz in the McGill Medical Museum. Natural size.)


The increased work leads to a very rapid overgrowth, both of the size and the number of the muscular elements. The organ, which, inormally, in man weighs from 250 to 300 grams, may come to weigh as much as 1980 grams. (Stokes' case). Here it must be noted that the hypertrophy is relatively much greater when brought about in the young individual, in whom the thickness of the right ventricle may come to equal four times the normal.

Closely allied is the overgrowth of the middle coat of the muscular arteries (as in the kidneys) in cases of increased blood pressure. In every one of these cases the increased work is of the nature of increased strain or tension acting on the individual cells of the tissue.

3. Compensatory Overgrowth.—Such muscular overgrowth is frequently referred to as compensatory, as, indeed, to a certain extent, it is. We prefer, however, to distinguish it as *adaptive*, and to employ the term compensatory for another series of cases, in which there is

OVERGROWTH

overgrowth of tissue to make up for *loss* of tissue of the same order. Of such, many examples may be cited. If parts of several organs be removed, the cells of the remaining parts become enlarged (simple hypertrophy) and undergo proliferation (hyperplasia), and, as in the thyroid, collections of cells which had been in a latent or persistent embryonic condition develop into fully formed active constituents. Thus, as Ponfick showed, if three-quarters of a rabbit's liver be removed, the remainder may hypertrophy in this way until it attains the size of the original organ. Or, again, if one of a pair of organs be (a) congenitally lacking in development, (b) destroyed by disease, or (c) removed



Compensatory hypertrophy: R. Kidney, congenital hypoplasia; L. Kidney, compensatory hypertrophy (length, 14.5 cm.); Norm., a normal adult kidney (length, 11.0 cm.). (Outlines drawn to scale from specimens in the McGill Medical Museum.)

experimentally or surgically, the other in a very short time shows enlargement, until it approximates toward the size and the weight of the original pair combined. And even, as has been demonstrated in the case of the thyroid and the kidney, if, after such hypertrophy of the remaining single organ, half of that be removed, the remaining quarter will still show great overgrowth. When, for example, one lung, or the kidney, through some intra-uterine disturbance, fails to undergo development, the other half is found greatly enlarged; the same is true where one kidney or one testicle is the seat of destructive disease or injury to its blood supply, or where the kidney or testicle is removed by operation.

It may, indeed, be laid down that in all the paired organs of the body

THE PROGRESSIVE TISSUE CHANGES

one member of the pair is capable of undertaking the work of both, and in doing this undergoes hypertrophy. Indeed, in the brain there are indications that, as regards several centres, one is active and the other largely latent, so much so that through disuse it may later, if called upon to work by the destruction of the active member of the pair, be unable to respond wholly. This, however, is not the case in intra-uterine or early life, when, in addition to the hypertrophy, it would seem that hyperplasia can also occur in the nerve cells, though later this latter becomes impossible.

It may be laid down that in all the paired organs of the body one member of the pair is capable of undertaking the work of both, and, doing increased work, undergoes hypertrophy.

Certain limitations have to be made to this statement. One has already been indicated in the case of the brain centres. A more correct statement is that *primarily* one member of a pair has this capacity. In general, it may be haid down that, while hypertrophy occurs thus, *complete compensation* only shows itself in intra-interine life. The extent of the compensation is modified by the age of the tissues, as, again, is the rate of compensation. The younger a tissue the greater its capacity for growth; the older, the less. It is in cases of congenital and youthful valvular disease of the heart that we encounter the greatest cardial hypertrophy and hyperplasia; in cases of congenital absence of one kidney, that the other shows the greatest overgrowth. Only in these cases do we find one kidney attaining the volume and weight normal for the pair of organs. With removal or destruction late in life the compensation is much less complete, and, being slowly established, death may be brought about by disturbance of function.

Hypertrophy versus Regeneration.—A distinction must in these cases be drawn between hypertrophy and regeneration. It is not, it is true, perfect, because wherever hyperplasia occurs, there we have a regenerative process, and, in addition, a most typical regeneration often to some extent accompanies the hypertrophy where there has been loss of a portion of an organ. The feature, however, of hypertrophy is that, in an organ made up of cell complexes, the number of these complexes is not increased. There is no increase, for instance, in the *number* of liver lobules in the hypertrophying liver, or of glomeruli in the kidney undergoing compensatory hypertrophy (save in very early life); the individual complexes enlarge and become more cellular-larger liver lobules, larger glomeruli; increase, not in number, but in length, of the renal tubules. In regeneration, on the other hand, new cell complexes become budded off from the old. As might be expected, the more complicated the structure of an organ, and the greater the number of tissues entering into its composition, the less is there of orderly regeneration, the more pronounced is the hypertrophy. But even in the liver, part of which has been destroyed by removal or disease, as again in the kidney, we gain evidence of an imperfect regeneration.

4. Vicarious Overgrowth. In yet another series of cases the compensation is more indirect; where one organ fails, there is overgrowth

OVERGROWTH AND REGENERATION

affecting organs of another order, though apparently of allied function, these organs vicariously undertaking the work of the diseased, destroyed, or overworked tissues. One of the clearest examples of these cases has been afforded by Rogowicz and Boyce and Beadles,1 who have pointed out that where the thyroid gland is atrophied or removed, the pituitary gland undergoes definite enlargement. There is a certain amount of evidence of similar relationship between the thyroid and the thymus, while, according to some observers, atrophy of the pancreas is accompanied by enlargement of Brunner's glands in the duodenum. Where the spleen has been removed, the bone marrow and certain lymph nodes appear to take on some, at least, of its functions, becoming enlarged, more particularly the hemolymph glands;² while the enlargement of the spleen occurring in certain cases of anemia and the changes then recognizable in the same are regarded by Leube and others as vicarious to this extent, that normally the spleen is a blood-building organ during feetal life only; now, when the bone-marrow, the main source of red corpuscles during adult life, is inadequate to supply these in sufficient quantity, the spleen comes into activity and shows hypertrophy.

5. Irritative Overgrowth.—I have already dwelt upon the fact that substances which, in larger amounts and greater concentration, are toxic, leading to degeneration of the tissues or arrest of function, often, in smaller amounts, act as direct stimuli to the cells—and one of the effects of this stimulation of the cells to increased activity may be increased growth—this not, of necessity, secondary to tissue destruction, as in many inflammatory lesions, but primarily, as has been proved more especially by Wegner's observations on the increased growth of bone when minute doses of phosphorus are given,³ and those of Ziegler and Obolonsky upon the influence of arsenic and phosphorus on the liver and kidneys.⁴ It will be remembered that a study of the development of tubercles (see p. 440) proves that, under the action of bacterial toxins, there may be a similar local tissue overgrowth induced, and that we have evidence that some, at least, of the *productive* fibrosis met with in various organs is probably of a like origin.

Mechanical stimuli, or irritation, of moderate grade may result in similar overgrowth. Years ago the late Sir James Paget pointed out that, whereas constant pressure, by cutting off the capillary blood supply of parts, leads to atrophy, recurrent intermittent pressure has the reverse effect (as in "corns").

 Nutritional Hypertrophies.—It may, however, well be asked whether in these cases of toxic irritation leading to cell growth we are not dealing—as Virchow held—with increased nutrition, due to hyperemia;

¹See also (and for bibliography) Herring, Q. Journ. Exp. Physiol., 1: 1908: 281.

² For these, see more particularly Swale Vincent, Proc. Physiol. Soc., 1898; xl, and Warthin, Contrib. to Med. Research (Vaughan Festschr.), Ann Arbor, 1903; 216. Meek, however (Q. Journ. of Med., July, 1910), brings evidence to show that the hemolymph nodes are not distinct organs, but lymph nodes with, it may be, only temporary alteration of circulation.

³ Virehow's Arch., 55; 1872; 11.

⁴ Ziegler's Beiträge, 2:1888:291.

whether the presence of the toxin in the cell does not lead to increased absorption and assimilation on its part? This may well be, but the entrance of the toxin is the primary event, and acts as the stimulant. If, however, we regard toxins, like other assimilated bodies, as potential foodstuffs, then we, perhaps, solve a long-standing difficulty. We refer to the following:

Mere hyperemia, alone, due to increased arterial blood passing to a part does not lead to hypertrophy. Or, otherwise, the mere abundant bathing of the cell with its ordinary nutritive fluid, without coincident call upon that cell to increased activity, does not cause it to grow. This fact is difficult to explain, nor do we know that we can offer an adequate explanation, there being one apparent exception that is classical. We refer to John Hunter's experiment of grafting the cock's spur on to the cock's comb, when, within a few weeks, the spur grows to a relatively enormous size. And the experiment has been confirmed by more than one modern observer. Here the point that immediately strikes one is, that the spur comes from a relatively non-vascular region and is implanted in most vascular erectile tissue. The increased hyperemia appears to be an essential factor in the overgrowth. There is this difficulty, however, that a few weeks later the spur begins to shrink and atrophy, and ultimately drops off. What relationship has this to the hyperemia? In speaking of implantation we shall meet with similar examples of preliminary overgrowth followed by atrophy. Not, by any means, all tissues exhibit the phenomenon when transplanted, nor even all embryonic actively vegetative tissues. It is not the mere hyperemia of the region of implantation, therefore, that induces the overgrowth. Some additional factor has to be invoked, and the only factor that can be suggested is that the blood, even the arterial blood, as it passes through different regions, becomes modified—here it has more oxygen, there less; here there diffuse into it, even into the smaller arteries, certain substances, there others have been absorbed from it-that, in short, the transplanted tissue receives a different pabulum from the normal, and that in some cases this in itself acts as a stimulant to active assimilation and growth. The mere diffusion of fluid into and through a cell. if the matter contained in that fluid is not taken up by the cytoplasm, does not favor growth. Assimilation is an active, we might say chemiotactic, process, depending upon the state of the cell at the time and the nature of the substance presented to the cell. Where a cell is not, by nervous or other mechanisms, stimulated to activity from without, and is in a state of equilibrium, it requires some abnormal constituent of the absorbed fluid to disturb that equilibrium and favor either increased building up, or increased breaking down of the cell substance.

If these views be correct, then there may be a nutritional hypertrophy, just as there may be a toxic degeneration of the cells, and John Hunter's experiment and the cell growth set up by dilute toxins fall into a common group.

In this group there are to be included also the tissue growths occurring in myxodema, acromegaly, and osteo-arthropathy. In the first

NUTRITION AND OVERGROWTH

two of these we observe that there is an intimate relationship between the symptoms in general and disturbance of the internal secretions; the third in its general characters exhibits so strong a family likeness to the other two that we must associate it with them.

Myxadema is brought about by atrophic disease, arrested function, or removal of the thyroid gland, and its main symptoms appear to be brought about by absence of the internal secretion of that organ and accumulation of products in the fluids of the body which are normally neutralized by that internal secretion, for if thyroid extract be exhibited the symptoms disappear. It has been noted that in the earlier stages of the disease there is increased interstitial mucin in the subcutaneous connective tissues; later, this gives place to an hypertrophic fibrosis.

Similarly, acromegaly, in which there is a remarkable overgrowth of the bones of the head and extremities, has frequently been found to be associated with extensive disease of the pituitary body. That this condition is of the same order as, and allied to, myscedema was strongly suggested by a case in the wards of my colleague, Professor James Stewart, in which tumor of the pituitary body was diagnosticated, but in which a myscedematous condition of the hands replaced any bony enlargement. At autopsy we discovered a large endotheliomatous tumor of the pituitary.

The main feature of *ostco-arthropathy* is bony overgrowth of the extremities developing in adult life, secondary to chronic disease elsewhere. Marie, who first described the condition, regarded it as essentially secondary to chronic lung disease, and gave it the top-heavy name of "osteo-arthropathie hypertrophiante pneumonique." Cases have since been described in which the lungs have not been found involved.

Yet another class of cases is possibly to be included here. We refer to the so-called *sympathetic* overgrowths. These are largely physiological; the most marked example is the development of the breasts during pregnancy. We have no evidence that the nervous system is the exciting cause in these cases. The enerves, it is true, as in the case of muscles, may, by stimulating to increased function and dissociative activity, secondarily induce overgrowth, but in the cases under consideration the growing cells are not, so far as we can see, functionally active, and the tendency is to regard the effective stimulus as of an internal secretory nature, the increased assimilation and cell growth and proliferation being induced by products circulating in the blood. This view has recently received striking confirmation from certain experiments by Lane-Claypon and Starling,¹ in which it was shown that, by inoculating nonpregnant animals with an extract of foctal tissues, enlargement of the manume is brought about (see p. 363).

If we regard this subject of nutritional hypertrophy in its broadest aspects, we see, as a matter of fact, that abundant nourishment has not led to the development of individuals of fullest growth. As with apples and strawberries, so with races of men, the best developed are found

¹ Proc. Roy. Soc. Biol., 77: 1905; 505.

THE PROGRESSIVE TISSUE CHANGES

near the limits of tolerable existence. Not in the tropics, but in the upper temperate zone, do we find the races of men of the highest stature and best bodily development. The intestinal parasites bathed in food already partly digested are degenerates compared with the non-parasitic members of the same orders. Where food is easily obtainable, that very ease reduces the need of active exercise, and want of use leads to diminished development. Activity and function in the cell, as in the individual, mean more active metabolism, better excretion of waste products, and better conditions of growth.

Passing beyond the limit in which active exercise assures good food to within the arctic circle—we come again on small races of men. There is a certain mean between nutrition and activity of the tissues that is favorable. Abundant nutrition, with active exercise, would seem to afford the optimum conditions. We have indicated that in some cases it may be that increased nutrition, when it is of such an order as to lead to increased anabolism, may thus be the stimulus to increased activity on the part of the cell, but in the majority of instances the reverse obtains; increased functional activity precedes and leads to overgrowth. We would again recall what we have already emphasized, namely, that excessive functional activity has the very opposite result. There may be dissociation of the cell substance and liberation of energy beyond the compensatory assimilative capacity of the cell.

Simulated Hypertrophy.—Finally, we must note some cases, usually included among the hypertrophies, which externally appear to present some of the most marked examples, which, however, are not hypertrophies at all. We refer to cases in which parts persist owing to lack of attrition. The most marked examples occur in connection with the teeth of certain animals. Normally, in these animals, teeth that are opposed are undergoing constant and fairly rapid attrition, and to compensate for this there is constant growth. If, now, the opposed tooth be destroyed, the growth of that which is left proceeds without attrition, and the organ may attain enormous dimensions. Rats' teeth, for example, have been known in their growth upward from the lower jaw to curve inward into the orbit or pierce the roof of the skull.

CHAPTER XII.

REGENERATION.

Loss of substance, however produced, provided it be not excessive, and that it does not affect certain "vital areas," so causing immediate death, tends, in all living organisms, to be remedied by either regeneration of the lost tissues, or, as we have already indicated, by compensatory overgrowth of other parts and increase in their functions.

The extent to which regeneration occurs varies greatly in the different forms of life. Broadly, it may be laid down, with many exceptions, that the simpler and lower the form, the greater and more complete the capacity for regeneration.


The mere fact, however, that a form is low in the scale of living beings does not necessarily have associated with it the capacity for regeneration. We find some curious exceptions, in which members of allied species vary greatly in their regenerative capacity. Thus, a more accurate statement is, that among simpler forms of life we find the greater capacity for regeneration; the lower in the scale, the greater is its extent.

The illustrations of this principle are trite and familiar, the study passing back to the observations of Abbé Trembley, in 1740, upon the Hydra viridis. His were the first demonstrations that the power of parts to grow into the whole organism was a property of animal forms as well as of plants. He cut his hydras longitudinally into two and four parts, when each became a whole animal, some of which he preserved as long as two years; split them along one side when they grew together again, bisected the head and gained thus two-headed hydras; bisected these again, and gained many-headed hydras. Another abbé, Spallanzani, in 1768, made further advance, obtaining the regeneration of the head and tail segments of earth-worms when these had been cut off. In one experiment he obtained a new head five times in the one animal. He showed that in the salamander the tail was regenerated, even lost vertebræ being replaced, and that regeneration, as of the tadpole's tail, or the limbs of the salamander, is more complete the younger the animal.

The abundant observations of late years have amply confirmed these earlier observations. Regeneration has been studied by Gruber and others in the protozoa, in which the sections containing the nucleus develop into the complete individual, while non-nucleated portions fail to regenerate,' in sponges, coelenterates, echinoderms, the various

¹ The observations of Gruber (Biol, Centralbl., 1904;717, and 1905;137) and of M. Nussbaum (Arch. f. mikr. Anat., 26:1886;485) are of special importance in establishing this fundamental importance of the nucleus in cell regeneration. orders of worms, crustaceans, arthropods, vertebrates—in fact, through practically the whole animal kingdom.

It is found that the extent of growth in the normal individual is largely a function of relative position and relationship to neighboring cells. Remove the obstacle to continued growth by removing these neighboring cells, and in the hydra, for example, orally, mouth organs and tentacles develop; aborally, toward the foot, the newly proliferated cells take on the arrangement of the cells forming the foot.



A. Hydra split in two, hanging vertically downward; later, the halves separated completely; B. two posterior ends united by oral surfaces; B', same, it regenerated two heads, each composed of parts of both pieces; B', absorption of one piece leading to a later separation of halves; C, two posterior eads united by oblique surfaces; later, one piece partially cut off, as indicated by line; C', later still, two heads developed; D, similar experiment in which only one head developed; E, five pieces united as shown by arrows; four heads regenerated, one being composed of parts of two pieces. (King.)

But the mere inter-relationship of the cells is not the only factor; external influences, also, determine the nature of the growth. Thus, as shown by Jacques Loeb,¹ geotropism may be a factor. Take a tubularian (a form allied to the hydra), cut off the head and foot ends, and invert it; place it upside down in the sand of the aquarium, and now a new head develops from what was the foot end, a new foot from the original head end. Indeed, so sensitive are these organisms to position that, if the position of the whole animal in the water be reversed, the foot end, even though uninjured, becomes converted


¹ Amer. Jour. of Physiol., 4: 1900: 60.

HETEROPLASIA

into a head. We dwell on these facts because certain observers, studying regeneration in higher animals only, deny this influence of external agencies to stimulate growth. The ultimate function of a cell and its capacity to proliferate are functions (1) of its position relative to other cells, and (2) of the action upon it of external physical and chemical agencies, as well as (3) of its primary constitution.

A very full consideration of regeneration throughout the animal kingdom is afforded by Morgan's work upon *Regeneration*⁴ which well deserves perusal. Some of the most interesting and valuable recent studies have been upon the vermes, notably upon the flat-worms (*Trematodes*), by Morgan, Flexner,² and others. Thus, as indicating the

F1G. 177



Heteromorphosis: regeneration of an antenna to replace an eye in the crustacean *Palinurus*, (Herbst.)

influence of one tissue upon the other, it has been noted that the planarian head only undergoes perfect regeneration when the ventral nerve ganglion has not been destroyed. Similar observations have been made in the earth-worm. Allied to this is the remarkable phenomenon of *heteromorphosis.*² In certain crustaceans, for example, if one eye be removed, there develops in its place, not a new eye, but an antenna-like organ, unless the ganglion cells connected with the eye have been left behind, in which case an eye is redeveloped. From these facts it would appear that, while the nerve cells do not initiate the regenerative process, they may influence the ultimate cell relationships and functions.

⁴ New York, Macmillan, 1905; see also his studies upon regeneration of Planarians, Arch. f. Entwickelungsmeeh., 7; 1898; 364.

² Jour. of Morphol., 14:1898:337.

³ This is also described as *heteroplasia*, but this latter term is also used to indicate a somewhat different phenomenon (see p. 641).

REGENERATION

Obviously, it is not the active functioning of the cells that initiates the regenerative process, for, in the earliest stages of a regenerating eve or limb of an arthropod, the new parts are, obviously, incapable of functioning.

In the higher vertebrates and in man this capacity to reproduce lost *parts* and lost organs is wholly wanting. At most, we recognize the capacity to reproduce lost *tissues*, but this only within certain limits.

1. If an organ be completely removed or destroyed it cannot regenerate. Where portions have been destroyed or removed, the remaining parts may proliferate and bring about regeneration. With development, that is, the cells forming the anlagen of the different organs become so specialized that other cells cannot take their place. Remove a whole bone, and it is not replaced; leave what we have termed the cambium layer—the periosteum—and this may reproduce the lost substance.

2. The higher and more specialized the tissue the less its capacity for regeneration. Portions of nerve cells, nerve fibres, may grow again after destruction, but not the whole neuron. Muscular reproduction is very imperfect, as is that of most glands. When we consider that the regeneration of an organ means that not one but several tissues must undergo coordinate development, the higher specific cells of the organ, the connective-tissue framework, the vessels, and the nerves all assuming particular relationships, it becomes marvellous that so elaborate a part as the limb of a crustacean can be reproduced, and not difficult to realize that, in man, organs like the liver, while they may hypertrophy, do not regenerate. It is only simple glands, like the salivary glands and, to some extent, the thyroid, that actively regenerate, and that by a process of budding from the preëxisting ducts, and even then the regeneration is imperfect, and as the new-formed connective tissue contracts, the new acini are apt to atrophy.

Morgan, in his Harvey lecture, New York, 1906, comes to the same conclusion, as an explanation of the lack of regeneration of parts in the higher animals. The tendency on the part of the newly formed connective tissue to contract, and so induce atrophy of other cells, is, in our opinion, the main hindrance to adequate regeneration.

In the liver, after the removal of a part, the surrounding bile ducts may give off cell processes, but, again, the new connective tissue arrests development into acini proper. The only cases in the liver in which true regeneration occurs are those of acute atrophy, with complete destruction of the specific cells. In these the framework is retained, and here, from both cells and bile ducts, regeneration may occur regeneration, that is, of the old lobules (Ribbert). In the kidney, as pointed out by Galeotti and Santa,⁴ true regeneration occurs in very young animals, the organ showing an increased number of glomeruli (and, therefore, of (tubules); in older animals, there is purely hypertrophy. We here have the working of Spallanzani's law, that regeneration is more complete in the vounger individual. In the skin, the hairs,

⁴ Ziegler's Beitr., 31: 1902: 121.

REGENERATION OF THE VARIOUS TISSUES IN MAN 605

sweat and sudoriparous glands do not regenerate, though in mucous membranes there is more or less imperfect reproduction of the follicles. Even with the lowest forms of tissue—connective tisue—as we have already noted (p. 429), the regenerated areas are apt to be of imperfect development.

the

or

he

it.

n

17

11

3. So, also, in a given tissue, where the cells of one order exhibit different degrees of specialization and differentiation, it is the less differentiated cells which most easily regenerate; in glands, for example, the neck and duct cells more than the actively secreting cells.

4. Lastly, to express what has already been indicated in a somewhat different form, where an organ contains two or more tissues of different profiferative capacities, in the regenerative process the more actively pro-liferative arrests the growth of the less active tissue, and leads to imperfect reproduction of the lost part. Of all tissues, the ordinary connective tissue has the greatest proliferative capacity, and, in the processes of healing, we observe that this overwhelms the other regenerating tissues.

In the study of regeneration we frequently observe that the more specialized tissues are not without the power of regeneration. Muscle cells, for example, at the cut end of a muscle show what are clearly the first stages of regeneration; but the process is a slow one, and too often before it is nearly complete the developing cells atrophy, as a consequence, it would seem, of the disturbance of nutrition and pressure exerted by the already developed fibroblasts and fully formed connective tissue.

REGENERATION OF THE VARIOUS TISSUES IN MAN.

Connective Tissue .- White Fibrous Connective Tissue .- It is unnecessary here to repeat in detail the successive stages of observations and opinions based thereon whereby we have attained to our present state of-it must be confessed-some uncertainty regarding the development of new connective tissue, interesting though that history is. This may be admitted, that connective tissue falls into line with other tissues, and is, in the main, if not entirely, produced by proliferation from preëxisting connective tissue. Ziegler, who was, for a time, the great upholder of the view that leukocytes are the main source of new connective tissue, was also, with his pupils, the main influence in controverting that view. Nevertheless, during the last few years the pendulum has shown a tendency to swing back. Metchnikoff has pointed out that in lower vertebrates wandering cells can be seen to assume the characters of fibroblasts within the tissues; and Maximow's' very full studies upon what he terms "polyblasts" would indicate that one of the fates of these wandering cells is to come to rest in the newly forming tissue and assume all the characters of a connective tissue. These polyblasts he regards as lymphocytes, modified by sojourn in

¹ Ziegler's Beitr., Supplemental Heft 5, 1902.

REGENERATION

the tissues. The recent observations of Schridde' throw doubt upon these conclusions. By the employment of a stain which differentiates the cell granulations in sections, Schridde points out that, while it is true that in form these cells are identical with the fibroblasts, their granulation remains distinct, and, as a consequence, they do not become



Formative cells, or fibroblasts in direct connection with the endothelial processes. (Ziegler.) identical, while it is justifiable to assume that their properties remain different.

This much may be stated with definiteness, that the polymorphonuclear leukocytes, the commonest of the infiltrating leukocytes, never become converted into connective tissue; also, that the main mass of fibroblasts are derivatives from the preëxisting connective-tissue cells. As regards cells of the lymphocytic and endothelial type, opinions remain divided. We ourselves are inclined to see an extremely close relationship between the vascular and lymphatic endothelium and the connective tissue, and cannot recognize any distinguishing marks between the cells and their processes, given off as buds from the new capillary loops, and the fibroblasts and their processes, with which they enter into connection. We would, indeed, include such endothelial derivatives along with the connectivetissue derivatives. But this view is not generally accepted.

The undoubled authority of His has led most pathologists to accept his conclusions that the vascular endothelium is a special derivative of the mesenelyma, derived from an order of cells that is distinct from the connective tissues. But, granting that this is so, they are but later developments

from the common mesoderm, from which connective tissue, and mesonchyma, are derived. They are closely related. And we have the observations of practically all recent students of the organization of thrombus (Waldeyer, Cornil and Ranvier, Thiersch, Baumgarten, Raab, Riedel, etc.) that there the new connective tissue is almost, if not entirely, derived from the endothelial cells. We do not see how this testimony can be refuted, and, if the process occurs in one region, we fail to see why it should be denied in another. Nevertheless, Thoma² and Heuking are the only ones who have boldly laid down that endothelial proliferation

¹ Munchener med. Woch., 1906, No. 4.

⁷ Ziegler's Beitr., 10: 1891; 434,

OF CONNECTIVE TISSUES

precedes fibroblastic development, and is the main origin of the fibroblasts, though not a few others tend toward this conclusion.

To the view of Leo Loeb' and others, that epithelial cells may assume a fibroblastic type, we shall revert in discussing metaplasia (p. 648). It is unnecessary to repeat here the various stages whereby the vegetative young connective-tissue cell, oval and stout, becomes successively stellate, then spindle-shaped, then surrounded by fine fibrils, and eventually but the skeleton, as it were, of a cell, with long, thin, greatly compressed nucleus and scarce any body substance (see p. 426); or to discuss the relationship of the primary mucinous matrix to the eventual connective fibrils (p. 39). We would only emphasize that throughout the body it is this tissue which is most active in regenerating, replacing in the process tissues of higher development.

Elastic Connective Tissue.—This is not only able to exhibit overgrowth, but may be definitely regenerated, appearing in areas of new tissue. Thus it may appear in abundance in areas of new connective tissue; in the intima of the arteries, for example, and in interstitial new tissue, or in some cases of cirrhosis of the liver. No distinction has as yet been made out between the cells connected with its development and the fibroblasts and connective-tissue cells proper; indeed, the nature of its development is still unsettled.

According to Loisel² (1897) the elastic fibres of the ligamentum nuclea originate within the cell, either as a peripheral sheath, or as a fine process at either end. In regeneration Enderlen³ and Jores⁴ note that they develop from preëxisting fibres, according to the latter as fine granules lying in series in the spaces between connective-tissue fibrils and cells, or in other cases as fibres which at first do not take on the specific elastic-tissue stain (Weigert's orcein). Development is slow, and for long a cicatrix may show no elastic tissue save at its edges.

Fatty Tissue.—It is doubtful whether we can truly speak of the regeneration of fatty tissue, for it is still an open question whether fatty tissue exists as a distinct entity, or whether it is to be regarded as a local modification of connective tissue. We are inclined to regard it as the latter. There are, however, certain regions in which, in the adult, fat is constantly present under normal conditions, e, g., the subcutaneous tissue, the appendices epiploice, and the auriculoventricular grooves.

There is still some uncertainty regarding the normal process of development of the fat cell. The generally accepted view is that certain somewhat large endothelioid cells in the immediate neighborhood of capillaries undergo multiplication, and in their relatively abundant cytoplasm fat gradually becomes stored in the form of larger and larger droplets, until, by fusion of the same, the cell becomes distended by a homogeneous spherule of fat, the protoplasm coming to form a mere peripheral ring, being collected in rather greater amount face.

⁴ Johns Hopkins Hosp. Bull., 9: 1898: 157.

² Quoted by M. Heidenhain, Plasma und Zelle, Pt. 1: 1907; 37.

⁴ Zeitsch, f. Chir., 45; 1897; 453, ⁴ Ziegler's Beitr., 27; 1900; 381,

REGENERATION

nucleus, the cell assuming thus a signet-ring shape. The view of Grawitz' and his pupils, that the fat cell arises from the fusion of several cells, whose nuclei, with one exception, fade into those of "slumber cells," is generally discredited. For ourselves, we cannot accept the slumber-cell theory. We are impressed, however, with the fact that the development by fusion of several cells has not been refuted, and, recalling how constantly degeneration is a reflection of development, the fact that the degenerating fat cells afford multiple cells is in our opinion very significant. In the degenerating muscle fibre, which is generally accepted as being developed from a cell-complex, we get a very similar appearance. We have repeatedly noted the appearance of characteristic polygonal cells, with transparent bodies and fine vacu-

> oles, in cases of inflammation and in degenerating areas of lipomasoccupying the space of a previous fat cell, which remains clearly indicated by its persistent membranecells which are wholly unlike any surrounding leukocytes. Nevertheless, not a few observers have regarded them as of leukocytic origin; most modern observers (Craiewicz, Flemming, Cornil and Ranvier, Marchand) agree that these are derived from the original fat cell. Sometimes, in a like position, a multinucleated mass is found, while Marchand figures definite mitotic figures in the cells. Eventually the membrane is absorbed, or gives way and the cells may appear irregularly distributed.

In cases of inflammation passing on to healing in a wound (as of the

abdominal wall) in which cicatrization is already well developed, cell masses like those above described may still be seen lying within a more or less perfect membrane of the old fat cell. We are apt to regard these as still in the process of degeneration, but it appears to be as permissible to suppose that here the reverse process is gradually developing with fusion and loss of nuclei until a single large vacuolated cell is left.

It is most difficult to arrive at a conclusion regarding the appearances seen, for, as noted, some of the cells are free, and these may show indications of increased size. In incluing in part to the Grawitz view, we wish to indicate that we cannot consider the matter definitely settled, nor that we are satisfied that it is correct.

In the appendices epiploicæ and in the atrophying thymus there

A Degeneration of a fat cell in an area of inflammation. At a can be seen the main nucleus of the fat cell; at band c other nuclei with surrounding cytoplasm (regarded by Maximow as degenerated polyblasts). The homogeneous fat globule has disappeared from the cell, but fine fatty droplets are present in the cytoplasm. (Maximow.)

F10. 179

⁴ See more particularly Schmidt, H., Virch, Arch., 128: 1892; 58.

OF CARTILAGE

of

eral

ber

the

nd,

ent, our

t is

t a

nce

C11-

JUS

di-

INV

he-

re-

er,

tre

ell.

a

ile

he

av

rly

ng

he

tre

SP

de

li-

W.

d.

re

appears to be an inverse relationship between the development and increase in the fatty tissue and the stroma of lymphoid tissue (each appendix epiploica contains a minute lymph nodule). This has been noted by more than one observer, but the nature of the relationship has not been surely determined. There is also a marked tendency for atrophied acini of the pancreas to be replaced by fat cells, which are wholly absent from normal pancreatic tissue.

Cartilage.—As noted by all observers, the regeneration of cartilage is a very slow process, so slow in some cases that certain observers have denied its occurrence, but it is well established that two distinct processes are possible: (1) perichondrial regeneration, and (2) regeneration from the cartilage proper.

1. In the former the perichondrium is detached, swollen, and inflamed, the space between it and the injured cartilage filled with fibrin. The inner aspect of the perichondrium becomes intensely cellular, and gradually these cells replace the fibrin, projecting into the area of injury and loss of substance. The cells, like those of the inner aspect of the perichondrium, at first closely resemble ordinary connective-tissue corpuscles and spindle cells. Those that are oldest and farthest from the perichondrium become rounded or polygonal, and are seen to lie in a transparent matrix; they have the characters of young cartilage cells. More commonly, however, the matrix is not transparent, but fibrillar. If, for example, a cut has been made into the costal cartilage of a young animal, resulting in a wedge-shaped wound, this becomes filled with a wedge-shaped, fibrillar mass, and, on making sections, the fibrils are found to originate from the injured aspect of the cartilage, or to be attached there in bundles which cross the wound. Here and there between the bundles are cartilage cells, at first without definite capsule, later enclosed and multiple, lying in a clear space; later again, judging from the statements of several observers, the fibrils disappear or become transparent, and a hyaline, completely formed cartilage

2. The process of regeneration from the cartilage proper is, in general, imperfect. To a slight extent it may be found accompanying the process just noted. In the immediate neighborhood of the wound the matrix undergoes softening, the "capsules" surrounding individual cartilage cells become larger, the cells divide, so that clusters of daughter cells replace single cells. There is, thus, proliferation—and, after more or less proliferation of the specific cells, each would seem to govern the formation of a new surrounding matrix, either hyaline or more or less fibrillar. According to Tizzoni,⁴ the fibrillar modification may extend into the uninjured surroundings, and previously hyaline, cartilage. It was clearly stages in this process that Redfern³ described in that remarkable series of experiments which constitutes the first satisfactory microscopic study of the process of inflammation and repair.

¹ Arch. per le sei, med., 2:1878:27.

² Monthly Journal of Medical Science, March, 1850; September, 1851, 39

REGENERATION

Bone.-We have noted in the preceding section that the cells in the earliest stages of perichondrial proliferation are undistinguishable from fibroblasts. Cartilage must, indeed, be regarded as but a modified form of connective tissue, and the same applies to bone. The character of periosteum very closely resembles those of perichondrium. Cartilage may, we need scarce remark, become converted into bone, and, what is more, under certain conditions periosteum may give origin to fibrous tissue-to a fibrous, instead of an osseous, union between the fractured ends of a long bone. Therefore, although we distinguish two forms of bone, according to the mode of normal development, the chondriform and the membraniform, the one passing through a cartilaginous fore-stage, the other not, it must be realized that in both we are dealing with modified connective tissue, and be prepared to find that in regeneration no distinction is to be drawn between the two, as, also, that in its essence, regeneration from the medulla is of the same order as from the periosteum.

Periosteal Regeneration.—It used to be held that when once periosteum becomes stripped from the bone that inevitably undergoes necrosis. We now realize that this is by no means necessary. Periosteum can regenerate, and Marchand and others have published cases in which an area, deprived of its periosteum has become eventually covered with a new layer. Only when there is suppuration and infiltration of the outer table of the bone by the suppurative microorganisms, and destruction of the bone cells, does necrosis become inevitable. According to Marchand's description, the new periosteum is firmly attached to the bone and separated from the overlying connective tissue; the old periosteum at its edge shows thickening and its fibres, instead of being arranged longitudinally in the direction of the long axis of the bone, exhibit a radial development, suggesting definitely that growth, in the main, has been inward from the periphery. There may be a definite osteoblastic layer on its inner surface.

Osseous Regeneration.—Bone is constantly undergoing regeneration. One has but to study a longitudinal section of a long bone, such as the femur, to be convinced that the lamellae are laid down along the lines of strain. The adaptation, indeed—the economical use of a minimum of material to support a maximum load as indicated, in the first place, by the tubular nature of the bone; in the second, by the arrangement of the lamellae—is most wonderful. But at different periods of life the load to be borne and the direction of the strain varies, and the old lamellae become absorbed and replaced by new; or, more correctly, can be seen to be undergoing absorption along one aspect while new material is deposited upon the other, so that the position of an individual lamella becomes shifted. Howship's lacunae, in normal bone, are areas in which the osteoclasts are thus absorbing the old bone, and a microscopic section of such normal bone exhibits layers of bone, evidently of different ages.

Such regeneration is wholly apart from the periosteum, though at the same time, in the process of growth there is steady addition to the

OF BONE

s in

able

odi-

The

um.

one.

igin

'een uish

the

carwe

> wo, the

um isis.

can

the

de-

ord-

the

the

rth.

e a

on.

the

nes

the

old

tly,

WB

ual

are .nd ne,

the

amount of bone, with increase in diameter, due to periosteal activity. As thus, normally, there is both internal and medullary, as well a periosteal regeneration, so after injury and fracture, both play a part in the regenerative process.

Regeneration of Membraniform Bone.—In the following description we follow Marchand,[†] and the statements apply both to what is seen in the medullary spaces of an implanted portion of one of the bones of the skull and in the medullary spaces of a long bone after injury. The first change to be observed is the development of a mass of young spindle cells, identical with ordinary fibroblasts. These, which become the eventual osteoblasts, give origin, first, to fibrille in increasing amount. In the meshes of the fibrillary network the cells are seen as irregular, angular, and rounded or clongated bodies, now free in the spaces of the network, now lying on the walls. At this period they have no processes. The nuclei are large, rounded, and with large nucleolus. The cells are not distinguishable from the laver of osteoblasts beneath the periosteum.

Next, the fibrillary substance becomes more homogeneous, previous to its forming the new bone substance, and the cells above described become included, and shrinking and becoming stellate, form the bone corpuscles; whereas the free osteoblasts and these cells in their earlier stages exhibit mitosis, as bone corpuscles this is wholly wanting. The successive stages from the osteoblast to the bone corpuscle are difficult to follow, but the successive processes closely resemble those observed in the development of hyaline cartilage from perichondrium; there is the same preliminary fibrillation, but while in that this eventually becomes homogeneous, here there is not so much as impregnation as an intimate organic binding of the calcareous salts into the matrix.

Regeneration of Ohondriform Bone.—Where there is a preliminary laying down of cartilage either of two processes may be noted: (1) The cartilage is at first relatively non-vascular. Now, vascular loops make their way into the cartilage, which becomes absorbed before them, and upon the cartilaginous remains, again, through the agency of osteoblasts, layers of young bone are deposited; or (2) the cartilage cells become converted into bone cells, and the surrounding matrix becomes converted into osseous matter. The two processes may occur simultaneously. The latter is often well marked in the conversion of cartilaginous into bony callus.

Regeneration of the Medulla.—There have been many recent studies upon the regeneration of the bone-marrow, of which the more notable are those of Enderlen² and Haasler.³ The appearances vary largely, according to the age and condition of the individual, whether we deal with the red marrow of youth and certain anemic conditions, white fatty marrow, or the later gelatinous serous atrophy. These marrow cells

¹ Die Wundheilung, Leipzig, 1901. The fullest recent work upon the regeneration of tissues.

² Deutsch, Zeitsch, f. Chirurg., 52: 1899; 293,

^a Areh. f. klin, Chirurg., 50; 1895; 75.

form the specific constituents of the marrow, but with them, it is unnecessary to say, are abundant vessels, connective tissue, and cells of osteoblastic type. According to the different observers, these can readily be distinguished. According to Enderlen, the degenerative changes in the marrow cells, which show themselves during the first twenty-four hours after injury and hemorrhage, give place within forty-eight hours to active mitosis and proliferation, while similar proliferation occurs in the connective tissues around the capillaries at the margin of the injured areas, and fibroblasts make their way into the hemorrhagic area. In seventy-two hours there is definite formation of new capillaries; on the fourth day a network of new fibrillar tissue has invaded the area of injury, and in the network are new marrow



Diagram of early stage of regeneration of bone— i, e_i , repair of fracture of a long bone; a_i external callus; b_i medullary callus; e_i region of fracture; d_i medulla; e_i shaft; f_i periosteum. (Perls.) cells, with, in addition, foreign-body giant cells around splinters of the injured bone; and gradually the young marrow cells, which at first had been rare, become abundant, more particularly at the periphery. Here, on the sixth day, hematoblasts can be distinguished, and, more deeply, developing fat cells. Certain cells with giant and often lobulated nuclei (distinct from the multinucleated osteoblasts) are evidently modified myelocytes (marrow cells).

Beyond the proliferation of the vessels it is deserving of note that in these cases of injury of the marrow there is little evidence of reactive inflammation; there is, for example, no migration of polynuclear leukocytes into the injured area.

The Healing of Fractures.—This subject is so fully treated in works upon surgery that it is unnecessary to recall here but the outlines of the process.

1. The variation in the process of healing depends primarily upon two factors: (a) the apposition of the fragments; (b) the nutrition of both fragments. The more perfect the apposition, and the more perfectly apposition is maintained, the less is the amount of exudate and subsequent callus, and the more rapid the knitting together; the greater the amount of riding of one fragment on the other, the greater is the irritation, exudation, and callus; while, further, if one fragment be without due blood supply, the reaction on its part is imperfect, and union is delayed or completely arrested.

 Both periosteum and medulla take part in the process. To this we have already referred.

3. The formation of the callus exhibits the following stages:

(a) First, hemorrhage and some exudation from the surrounding vessels, with coagulation.

OF LYMPHADENOID TISSUE

(b) Invasion of the coagulum by cells – polymorphonuclear from the surrounding tissues, fibroblastic from the periosteum and marrow.

(c) Organization of the clot from periosteum and marrow, with absorption of the fibrin and replacement by tissue,

(d) Conversion of the cells derived from the periosteum into cartilage cells and formation of a cartilaginous callus. This is not a necessary stage, but it is seen where the callus is extensive.

(e) Conversion of the cartilaginous into osteoid callus, *i. e.*, modification of the cartilage cells into bone corpuscles. Here two stages are described by some authors: (1) A vascularization of the cartilage from the periosteum and marrow, with conversion of the cartilage into bone cells, without, at first, any deposit or combination of calcareous salts in the matrix. This soft tissue is regarded as *osteoid* tissue proper. When, later, the salts become deposited, but the tissue has still not the perfect features of bone, some still speak of this as osteoid tissue, others as *osseous or bony tissue*, as distinct from perfect bone. The distinction is unnecessary. (2) Absorption of the imperfect bone originally deposited, whether through fibrillar or cartilaginous development, and replacement by lamellar bone, the lamellae being laid down along the lines of strain.

This absorption is a very gradual process, extending over years where there is extensive callus or grave displacement. The medullary cavity is, in general, completely closed by new bony growth; this in time becomes absorbed, the ends of overriding fragments become rounded off; and, eventually, whereas at first there had been an excess of imperfect bone, there remains only sufficient properly formed bone to secure perfect solidity of the part.

Lymphadenoid Tissue.—It is, perhaps, difficult to speak of regeneration affecting a tissue, which, as regards its specific element, the lymphocytes, is always, normally, regenerating. There are indications, however, that these are not of the same origin as the framework, and that the lymphocytes and the framework, along with the large endothelial cells (macrophages) of the lymph spaces are of different origin.¹ The observations of Gulland show that the lymph cells

¹ I have to confess that my conceptions of the relationships of the different orders of leukocytes (see "Inflammation" 4th edition Maemillan, 1909, pages 65–84) have been not a little disturbed by the observations of Professor Downey, of Minneapolis (Folia harmatologica, 8: 1909; 41), upon the leukocytes of that very early form of fish: Polyodon spathula (the "sponobill" of the Mississippi). These ganoid fishes have a cartilaginous skeleton and therefore no bone and no bone marrow. The main seat of leukocyte formation is in the kidneys, and here, as Professor Downey has demonstrated to me, all types of lymphocytic and granule cells are developed. They are so intimately intermixed that it is difficult to believe that all have not a common origin. It may, however, be reasonably urged that conditions found in the lowly ganoids do not obtain in the higher mammals, that with evolution there has become a progressive differentiation, whereby the mother cells of the lymphocytes have become distinct from those of the "polymorphs" and granule cells. This best fits in with what we observed in the higher minals.

is unells of se can erative ie first within ries at iv into mation tissue arrow of the v the it first ndant, phery. more Cer-· muldently cells). of the e that · marreacor exuclear 28. works of the upon

> pere and reater is the at be , and

on of

> this

ading

wander into certain areas, and these, in the connective-tissue framework, form germ centres, and it would seem established that in the embryo the system of lymph spaces is developed before any leukocytes show themselves in the blood.

Beard, confirming an older observation of Kölliker, finds that the earliest leukocytes originate in the thymus by a remarkable conversion of the *epiblastic* cells of the follicles; and his observations have been corroborated by others. What form of leukocyte originates thus is left undetermined—a matter of some importance when we regard the lymphocytes and the polymorphonuclears as wholly distinct types. More recent observers find leukocytes in the blood before there is any sign of change in the thymus.

Under certain conditions new lymph nodes develop in various situations, in the subperitoneal tissue, the liver, etc., where, normally, these are unrecognizable. It is possible that lymphadenoid tissue is "latent" in these positions. Such latent or potential tissue is present in the sheaths of veins, though here, again, we note that the lymphocytes are capable of migrating from the lumen into the perivascular lymph spaces. The generally accepted view is that this latter process is the usual course, namely, that lymphocytes come to rest in these positions, and then, proliferating, lead to a modification of the inclosing tissue.

When a lymph node is injured, according to Ribbert, proliferative changes, with new-growth, are to be observed, affecting all the elements of the follicle—reticulum, endothelium of lymph spaces, bloodvessels, and germ centres—in the immediate neighborhood of the injury.

Leukocytes.—We have noted that the lymphocytes undergo active development in the lymph nodules, which contain accumulations of "mother cells," the germ centres. As regards the leukocytes proper (of some authorities), namely, the polymorphonuclear leukocytes (including the cosinophiles), these, in postnatal life, and under normal conditions, are developed (along with the erythrocytes) more particularly in the bone marrow, where there can be no question that they originate directly from the myelocytes (Ehrlich). The opinion enunciated by Gulland and Uskoff, that they are an older mature form of the lymphocyte, would not seem to be tenable. In pathological conditions the spleen, the liver, and other organs may exhibit myelocytes and be, therefore, a seat of formation. Mitotic forms have been observed in the normal blood, but very rarely; more abundantly in pathological conditions (e. q., leukemin).

It must be noted that the sudden exhibition of an increased number of leukocytes in the blood is not a necessary indication of regeneration, but may be merely evidence of attraction of the cells out of the bone marrow, lymph glands, etc. To such wandering out is to be ascribed the physiological leukocytosis that follows a meal. At the same time it must be remembered that there is a constant normal destruction of leukocytes and constant new development.

Blood-vascular Tissue. During the stage of growth three methods of vascular formation have been distinguished: (1) A remarkable pro-

OF BLOOD-VASCULAR TISSUE

cess of cell *cavitation*, certain cells (of the vascular area in the chick, for example) becoming hollowed out, and during the process giving rise to blood corpuscles in their interior, the cavities of apposed cells fusing in series, so as to form tubes, which eventually become connected with vessels containing circulating blood. (2) A process of canalization, the blood making its way between rows of cells, which cells become converted into the endothelial lining of the capillary channels thus formed. It is now regarded as doubtful whether this method of formation occurs in the developing organism. (3) A process of budding. Certain cells of the endothelium of capillaries already formed give off buds or long protoplasmic processes, at first non-nucleated; the process from one capillary fuses with that from another, and the solid strand thus formed becomes hollowed out, thus giving passage to the blood from one capillary to the other, while eventually, by mitosis, nuclei pass into the walls of the tube, which becomes converted into a capillary with endothelial walls.

'Thoma, who has made peculiarly full and painstaking observations upon vessel formation, wholly denies the existence of the first process. He holds that all new vascular formation is truly *inter-* and not *intra*cellular. In the vascular area of the chick he describes the mesoblastic cells as becoming arranged in strands; rounded spaces appear between the cells, these spaces become filled with clear substance, probably fluid, open into one another, and thus form the first capillaries, the surrounding cells, primarily polygonal, becoming gradually transformed into pavement endothelium. Following upon this stage, further new capillaries are formed by the third or budding process. It will be seen that he regards the canalization as primary and independent of any blood pressure.⁴

In regeneration, the first of the processes above described has never been observed. The nearest approach to the second is seen in the somewhat rare cases of complete dissecting aneurysm.

In this condition, owing to disease of the inner wall of the aorta, sudden exertion and sudden rise of blood pressure lead to rupture of the intima, and now the blood forces its way between the fibres of the middle coat. If nothing further happens, then the blood thus leaving the vessel coagulates, forming a solid clot; at times, however, dissecting a passage for itself, it reënters the aorta or one of its branches at some point lower down. As a result, the blood expelled from the heart finds its way down the dissected, as well as down the natural passage, and the aorta appears to be doubled. Where the current is free but little coagulation occurs along the walls of the artificial passage, and, if death occurs a week or more after the rupture, what coagulum is formed is found to be covered by a distinct endothelial coat.

In granulation tissue, whether superficial or internal, the third process is that encountered. This we have already described (p. 428).

Red Corpuscles.-After great loss of blood, and in profound anemias,

¹Thoma, Pathology, English edition. Translated by A. Bruce, 1:1896:474.

methe vtes

the

sion

ieen

s is

the

pes.

any

ua-

lese

nt"

the

rtes

aph

the

ms,

els,

Det

11-

1ev

he

ms

be.

ы,

ne

ne

ds

the red marrow of the bones becomes markedly increased in amount, and the spleen is frequently found enlarged. In the red marrow, as Neumann first pointed out, there are normally present nucleated cells having hemoglobin in their cell substance. In anemia, and after loss of blood, coincident with the increase in red marrow, nucleated red corpuscles are to be detected in the circulating blood. Further, increased mitosis of these *hematoblasts* is to be observed in the bone marrow. This is an indication that there is an increased production and discharge of hemoglobin-containing cells from the bone marrow. As Howell has more especially pointed out (in this confirming Bizzozero and Salvioli), similar hematoblasts are recognizable in the spleen under similar conditions.

We therefore conclude that the red bone marrow and the spleen are preëminently the seats of regeneration of red corpuscles. The hemolymph nodes of the abdominal area, organs intermediate in histological structure between the lymph glands and the spleen, which have of late been studied by Swale Vincent, Warthin, and others, would seem also to be concerned in this production. Regarding these nodes, the recent studies of Meek demonstrate that they are not structures *sui* generis, but lymph nodules which undergo modification in function in particular conditions of the circulation. Wherefore, we conclude that lymph nodules, like the spleen, may be the seat of red corpuscle formation.

It is when we come to determine the origin of these hematoblasts, and, again, the method whereby the nucleated hematoblasts give rise to the non-nucleated red corpuscles, that doubts arise. We are not certain how far hematoblasts arise from preexisting hematoblasts, or from less differentiated "mother cells." Löwit, indeed, has described certain small, colorless, nucleated cells in the bone marrow—"erythroblasts"—which he regards as the precursors of the hematoblasts, and upon *a priori* grounds, arguing from what occurs in the embryo, where colorless mesoblastic cells give rise to the nucleated hematoblasts, this may well be the case; while, again, it is not impossible that the endothelial cells of certain areas may, by division, give off, as in the embryo, such hematoblasts. According to Bizzozero, the hematoblasts are not in the lymph spaces, but actually within the capillaries of the bone marrow.

We must accept that the non-nucleated red corpuscle arises from the hematoblast, not by a process of budding (the hemoglobin containing moiety of the cell substance dividing off from the perfect cell, which then is capable of elaborating more hemoglobin-containing cell substance, and giving rise to a second, or a series of red corpuscles), but, as noted by Macallum, by a process of discharge of material from the nucleus which undergoes eventual disintegration.

Epithelia.—'There is no known exception to the rule that, in regeneration, epithelial elements develop from preëxisting epithelium. The apparent exceptions, *i. e.*, where an island of new epithelium begins to develop in the middle of an area of gramulating tissue, are, judging from

OF EPITHELIA

transplantation experiments, explicable by accidental transplantation of living epithelial cells on to the granulating surface, or, in some cases, by the persistence of epithelial elements upon the eroded surface – the deeper parts of hair follicles, or skin glands, which, proliferating, take on simpler epithelial characters.

A rather remarkable example of this persistence has been met with in connection with the lens, which, it need scarce be said, is of epithelial origin. Occasionally it has been noted that, after complete extirpation of this organ in cataract operation, a new lens has developed. Experiments upon the rabbit' show that, where this occurs in mammals, portions of the posterior ligament or capsule have been left behind, in close connection with which is the "cambium" layer, from which throughout life new lens cells have been developed. It is from this,

after removal of the main mass of the organ, that regeneration takes place. In certain of the lower animals the new lens has a different origin; to this reference will be made in discussing metaplasia (p. 643).

An apparent exception to this statement has been noted by Saxer² and others in certain gliomas. Here there is a tendency to the formation of cysts containing serous fluid, and in some cases these cysts, which are clearly secondary developments, are found to possess a more or less regular lining of fairly columnar cells; they gain, that is, an epithelium which evidently is derived from the glial cell of the body of the tumor. While this is clearly the case, we gravely doubt if this can be spoken of as a true epithelium. It must be remembered that the neu-



"Pseudo-epithelium," or secondary epithelium without basement membrane living a cyst in a glioma, formed by modification of the superficial layer of glioma cells. (Saxer,)

roglia is itself of epiblastic origin, so that, were a true epithelium found, it would not be an example of conversion of cells of one order into those of a wholly different type. But specimens which we have seen, and Saxer's figures, show that no basement membrane is formed; the lining cells pass imperceptibly without demarcation into the underlying cellular tissue.

Along with the vascular endothelium and endothelia in general, the epithelia of the body stand preëminent in their capacity for *complete* regeneration, and, what is more, offer particularly favorable conditions

¹Vide R. Randolph, Welch Festschrift (Johns Hopkins Hosp. Repts.), 10: 1900; 237, who gives full reference to earlier literature.

² Ziegler's Beitr., 38: 1904.

for a study of the process, which thus has been investigated by a large number of observers.¹

Here we may rapidly note the data which may be regarded as well established. Within two hours of the removal or destruction of epidermis, whether of the outer skin, the tongue, or of mucous membrane, in warm-biooded animals, as also in amphibians, the cells of the deeper layers, and even columnar epithelium, exhibit translation. Keratinized cells are degenerated and inert, but prickle cells and those of the lower layers are seen to alter their shape, to become pyriform, and gliding one over the other, while still retaining connection, there is thus early exhibited a tendency for the uninjured cells to close over the defect. Within twenty-four hours the epithelium surrounding a wound is distinctly thinned, composed of fewer layers, while, at its edge, a single layer of flattened cells covers over the wound. Direct division of the cells is frequently noted, so that many cells contain two nuclei.

Cellular masses may be seen extending between the layers of the fibrinous scab which by now covers the wound, but more especially they extend along the surface of the injured tissue and into depressions on that surface. Occasionally, it would seem, small collections of these cells may, by movement, become detached, and form islands of growth on a granulating surface. Within forty-eight hours mitoses may be observed in the Malpighian layer of the surrounding less altered epidermis and prickle-cell layer, as also in those cells which have spread over the surface. Briefly, by this combination of translation and cell multiplication the denuded surface tends to be covered, at first, by a single or irregular layer of flattened cells encroaching from the edge of the wound.

At first these are relatively loosely attached to the underlying surface and granulation tissue, but at a comparatively early date the formation of a basement membrane is observable, apparently by the fusion of fibrils passing from the underlying connective tissue, upon which the overlying cells become sessile. The exact relationship to this basement membrane is still undetermined. At first there is free passage out of leukocytes between the cells and their communicating bridges, and a fairly extensive phagocytosis has also been observed, the young epithelial cells evidently utilizing the leukocytes as one source of nutrition. Later, both these processes become restricted; the proliferation leads to the formation of several cell layers, of which the outer layer, at first, shows irregular keratinization (Unna), without an underlying granular layer.

¹ See, more especially, Mayzel (who first established that epithelium arises from preëxisting epithelium), Sitzber, der Warschauer ärztl, Gesell. (April, 1874); Ref. Virch.-Hirsch Jahresber, 1; 1878; 3, and Arbeiten a. d. Lab. A. Med, Fae, Warschaw, pt. 4; W. Flemming (on Mitoses), Arch. f. mikr, Anat., 19; 1880; 347; and ibid., 24; 1885; S. Garten (On the Arrangement of the Intercellular Bridges in Regeneration), Arch. f. Anat. u. Physiol, Physiol. Abth., 1895; 401; Leo Loeb (on the migration movements of epithelial cells), Bull. Johns Hopkins Hosp., 9; 1898; 157; Ranvier (Regeneration of Conjunctiva), Compt. rend. Acad. de Sei, 123; 1806; 1228; Spuder (Regeneration of Hairs), Verhaull, Anat., Geschl. 1899; 17.

Eventually, a granular layer shows itself, and a complete simple epidermis becomes developed, with all the normal layers, from the Malpighian upward.

r-

r

r

e

n

Hairs, sweat and sebaceous glands are not reproduced, unless the original injury had not been deep enough to destroy the deeper lying portions of their structure, in which case they may develop anew. In this case it is interesting to note, as regards the glands, that there is a definite down-growth of solid processes of the overlying epithelium to meet them, which is difficult to explain save on the theory of reciprocal attraction or chemiotaxis between cells of like nature.

In cases of chronic ulceration, tuberculous and syphilitic, we observe at the edges of the ulcer a marked tendency for the epithelium to grow downward into the underlying tissue in the form of processes. These may become snared off, and develop into epithelial pearls; or, if thin, may become infiltrated by, or themselves infiltrate, the surrounding tissue, in which case isolated epithelial cells, or small clusters of such, with no exact demarcation, are encountered in the dermis. Where this is the case it is ofttimes practically impossible to determine whether we are dealing with mere chronic inflammation or with the earlier stage of epithelioma. As a matter of fact, chronic inflammation may be succeeded by definite epithelioma, and this not merely of surface tissues; but where fistulous tracts occur leading down to bone, for example, owing to this spreading property of epithelium, the epidermal cells may spread down the fistula into the bony cavity, and there be a cause of apparently primary cancer of bone. Of hairs, it may be repeated that they only regenerate when the root beds have not been destroyed. Where, as in certain parasitic inflammations and anemic conditions of the scalp, there is destruction or death of the root-bed, permanent baldness is the result.

The same is, in general, true of the *nails*. The nail-bed passes farther back than is generally imagined, and this would seem to explain how it is that after removal of a terminal phalanx an imperfect nail occasionally shows itself at the end of the finger. In forming the flap, a portion of the nail-bed has been retained. There are, however, several cases on record in which this explanation is not adequate—in which, after removal of two phalanges, an apology for a nail ultimately develops at the end of the remaining part. We have to accept, it appears to us—however unwillingly—that environmental conditions may stimulate a metaplasia of the ordinary skin into nail-producing matrix.

Regeneration of Mucous Membranes.—The same general process described for the epidermis is found to apply in connection with the mucous membranes. At the edge of a wound even the fully formed columnar cells lose their cilia, become more cubical, and ultimately rounded and flattened, and undergo a translation over the exposed surface, with relatively considerable rapidity. Later, proliferating, they form again a columnar epithelium identical in its character with the normal. Unlike what occurs upon the skin surface, the simple gland follicles become reproduced, the reproduction being hastened

if the lower parts of the primary follicles have not been entirely destroyed.

Such regeneration of mucous membrane takes place in the uterus, to some extent, after every menstrual period, and very extensively at the placental attachment after pregnancy and delivery.

There has been some discussion as to the cells from which this puerperal regeneration originates. Certain giant cells are to be met with in the upper layers of the muscularis which disappear later, and some observers have regarded them as latent remains of the mucous glands. Aschoff doubts that this is their origin, and points out that throughout pregnancy, below the placental site, recognizable remains of the mucous glands are to be made out. It is more natural to accept these as the site of origin of the new mucosa.

Endothelia.—In the regeneration of endothelium there is observed the same tendency toward translation and proliferation by both direct and indirect nuclear division, so as to cover a denuded surface, as is seen in the case of epithelia, and the surface may be covered with extreme rapidity. Such endothelium may form a covering not only over the tissue proper of the part, but over fibrin, as, again, over newgrowths which, as is sometimes seen in the peritoneum, have clearly originated by surface transplantation. The generally accepted view is that such endothelium arises from preëxisting superficial endothelial cells, and provisionally, until this is surely determined, this is the safer view to accept. But, as already noted, the relationship of endothelial cells to fibroblasts and connective-tissue cells has not been wholly settled, and if, as Baumgarten maintains, and claims he has demonstrated, this vascular endothelium can, by proliferation, give rise to underlying fibrous tissue, the reverse process must also be regarded as possible.

Regeneration of the Glandular Tissue.—It is impossible to read the more recent studies upon the results of wounds or excisions of glandular organs without being impressed by the singularly small amount of regeneration that is found in most cases. In saying this, as we shall point out in our remarks on hypertrophy, a distinction must be drawn between that condition and regeneration proper. But the development of new glandular tubules, or acini, is wholly wanting, or slight and of little or no functional value until we study the simplest glands, such as the Lieberkühnian follicles of the intestines, the uterine glands, or the salivary glands. These may undergo extensive regeneration, though, in the case of salivary glands, the newly budded-off outgrowths from the ducts, which develop into acini proper, are apt to be surrounded by a new connective-tissue growth which is not normal, and subsequently, with contraction of the same, to undergo more or less atrophy.

Liver.—According to Podwyssozky,¹ the changes here vary according to the animal employed. In guinea-pigs and rabbits there may be a certain amount of regeneration following upon excision of the

¹ Ziegler's Beitr., 1:1886.

OF VARIOUS TISSUES

part. Already, two days after the injury, the epithelium of the bile ducts in the neighborhood exhibits mitoses, and processes are formed of new bile ducts, forming a network in the newly developed cicatricial connective tissue, while some of them at their termination develop into liver cells, the result is very incomplete, and the regeneration bears no proportion in the amount excised. In cats and rats, the liver cells show active proliferation, but in many this stops short of cellular multiplication, and cells with double nuclei result. The process here is one of hypertrophy and enlargement of the preëxisting lobules, without regeneration, in the true sense of the term. In his studies upon experimental cirrhosis Opie' has demonstrated clearly the co-existence of the two processes of budding from the bile ducts and proliferations of the liver cells with formation of pseudo-bile ducts.

·ly

-r-

ne

ls.

ut

ite

ect

is

ith

ilv

11-

rly

is

ing

the

WD

op-

rht

ds.

ds.

on.

ths

ur-

ord-

nay

the

Kidney.—In this organ the fairly numerous studies that have now been made in man, after injury or operative excision, afford no evidence of new formation of tubules, although in certain of the lower animals, more particularly from the medullary collecting tubules, new tubules may make their way into the region of the wound; but, in the adult animal at least, these have no glomerulus formed at their upper end, are small, imperfect, and cannot function.

There is evidence, however, that in quite young animals, not so much in an injured kidney, as accompanying the hypertrophy of the opposite kidney, new glomeruli and tubules are capable of arising, for the number of these is found greater than in the normal kidney, and certain cell accumulations have been detected in the growing organ, more particularly in the outermost portion of the cortex, which are now accepted as latent glomerular anlagen.

A study of cases of acute parenchymatous nephritis brings to light abundant evidence that there exists regeneration of the epithelium to replace that cast off in the course of the inflammatory process. Neighboring cells become multinucleated and flattened, spreading over the area of denudation, so that a single multinucleated cell may at times encircle the tubule. Later, cell division takes place with enlargement of the individual cells to the normal size.²

Thyroid.—In the thyroid, also, Wölfler³ drew attention some years ago to similar cell masses, which he likewise regarded as latent anlagen, and, after partial destruction of the organ, he recognized active growth in these with development into the typical follicles. We have recognized Wölfler's clusters in several cases of thyroid disease. Whether they are persistent anlagen, or reversions through atrophy, it is difficult to say, but transitions may be recognized from these cell masses to others having a small lumen, and so to typical follicles. Where, as by Halsted and others, portions of the thyroid have been removed, the

³ Die Entwickelung und der Bau der Kropfes, Berlin, 1883.

¹ Trans. Assoc. Am. Phys., 25; 1910.

² For study and bibliography, see Oertel, Publ. Russell Sage Inst. of Pathol., No. 1, 1909.

regeneration, which is here not inconsiderable, is by a process of budding and separation of new follicles from the old.

Pancreas.—The observations here are practically unanimous that no regeneration takes place.

Spleen.—Here, also, the balance of evidence is to the effect that, while there may be hypertrophy, and compensatory (or vicarious) hypertrophy of the hemolymph glands,[†] which may take on the characters of splenic tissue, at the edge of a wound in the spleen no true regeneration occurs.

Testicle.—Griffini² found that there might in the frog be regeneration of the tubules by budding and growths from the ducts, though Maximow,³ who, in part, confirms the observation, denies that this is a perfect regeneration. In higher animals, although there may be a marked overgrowth of the characteristic interstitial cells, the tubules do not regenerate.

Ovary.—It is generally accepted that this organ also is incapable of regeneration. Pugnet's' observation, that after removal of onehalf of the rabbit's ovary the wound becomes covered by germinal epithelium, which then gives origin to abundant ova, has not been confirmed.

Muscle.—Plain Muscle Fibres.—After injury in their neighborhood, as shown by observations upon the stomach wall, muscularis mucose, and uterus, these may, in from two to five days, exhibit abundant mitoses, and in the newt (Stilling and Pfitzner⁵) there may be formation of new fibres as a result; but in the rabbit (Ritchie⁶) it is followed by no proper new formation of fibres; the cicatrix in the uterus and elsewhere is formed entirely of connective tissue.

Striated Muscle.—The regeneration of striated muscle after injury is a slow and most often an incomplete process, for where there has been any extensive laceration and separation of the fibres, these, in the first place, contract apart, and, in the second, we find the rule in operation already noted, namely, that the more rapidly regenerating connective-tissue elements usurp the place and hinder the development of the higher tissue. Thus most often a fibrous cicatrix unites the two ends of a wounded muscle. But this is not always the case. Where a wound or cut has been, in the main, longitudinal, or where the cut edges can be kept apposed, there may, in a few months, be found very little indication of fibrous cicatrix. Or where, as happens in connection with Zenker's degeneration of muscle in typhoid, and may occur in contusions, the substance of the fibre is gravely injured without the sarcolemma sheath being destroyed, there the regeneration may be complete. The series of changes in the two orders of cases exhibits

¹ See Dock and Warthin, An., Jour, Med. Sci., 1904, and for full literature, Weidenreich, Arch. f. Mikros, Anat., 1905.

² Arch, per le sci. méd., 5:1887:11.

⁴ Compt. rend. Soc. de Biol., 1900.


6 Vireh. Arch., 109: 1887: 507.

³ Ziegler's Beitr., 26: 1899: 2.

⁵ Arch. f. mikr. Anat., 28: 1886,

OF STRIATED MUSCLE

certain differences. Needless to say, the two may be combined to a greater or less extent. (1) The latter series of cases offers the simpler picture, *i. e.*, where the sarcolemma remains intact. Here, first the separated fragments of the muscle substance proper contract into broad, swollen masses. All the muscle nuclei are not destroyed, and it is from them and the undifferentiated zone of protoplasm around them that the regeneration proceeds. They multiply with relative rapidity, and become abundant, each surrounded by an increasing



Successive stages in the regeneration of voluntary muscle; A, formation of bud of cytophasm with loss of strainton and multiplication of muscle mulcip? R, the nuclei acquire cytophasmic territories and cells, uninucleate and multinucleate, separate from the bud (sarcoblasts); a, unaltered end of muscle fibre; b, sarcoblasts; c, multinuclear sarcoblasts, one nucleus at d showing mitosis; C, carly stage of new nuscle fibre, multinucleate and exhibiting longitudinal strainton, becoming fused with the original fibre; D, regeneration complete but irregular, the original fibre being continued into three processes. (After Volkmann.)

zone of cytoplasm, and forming masses or clumps from which separate individual mononucleated cells, or multinuclear. As they grow in size they cause erosion in the now homogeneous glassy masses of the original striated matter, or make their way between these and the sarcolemma. It is obvious that they absorb this old material, employing it as a foodstuff, until little of it is left. In the meantime some of these new cells become elongated and obscurely spindle-shaped, irregular in size, others still remaining small and polygonal. These larger forms exhibit, first, a longitudinal fibrillation; later, the nucleus

ling

ha

hat, bus) hartrue

genugh is is be a ules

able oneinal been

ood, osæ, dant ition d by else-

jury has the operconnt of two cre a cut very nnecoccur it the y be

eiden-

: 2. : 1886,

or nuclei become lateral, and away from it, or them, the first signs of transverse striation show themselves. With this the old sarcolemma sheath becomes absorbed, and the new muscle elements, of very irregular size, lie free. Some (the smaller cells) migrate or disappear, just as in the tadpole's tail Metchnikoff showed that like elements from the degenerating muscle could become wandering cells; the others gain a sarcolemma from the surrounding connective tissue, though the stages of this process are not clearly understood. The nuclei are now prominently lateral, and the breadth of the fibres markedly increased, although still smaller than the normal. It will be seen that in this process, when uncomplicated, small-celled infiltration and fibroblastic overgrowth play no part.

2. When the muscle fibres are ruptured, as by a cut or laceration, the abundant capillary network is also ruptured, and hemorrhages, fibrin formation, inflammation, and fibroblastic regeneration complicate the picture. As regards the ruptured fibres themselves, again the separated portions contract into clumps, which largely lose their striation. Within twenty-four hours the muscle nuclei show a remarkably active direct division, giving rise to chains, sometimes of thirty to forty members, and these collect more particularly in the homogeneous unstriated clumped end of the fibre. Sometimes collections of the nuclei in a homogeneous protoplasm form lateral buds at the side of the injured fibre. More often they are terminal, and the fibre may divide into two or more parts, each terminating in one of these nucleated clumps or buds. The process, it will be seen, is a modification of that described above, and here, also, at times, individual nuclei, with surrounding cytoplasm, or multinucleated masses, separate themselves off, though, owing to the accompanying leukocytosis, the nature of the cells seen cannot always surely be made out.

The buds elongate, extending between the fibrils of new connective tissue derived from the growing intermuscular tissue, and in favorable cases, and in the course of weeks, the number of nuclei becomes reduced, longitudinal fibration and transverse striation show themselves, and the new-formed extension of the fibre becomes indistinguishable from the old, save that its direction may be irregular. More often the wouldbe fibre is strangled by the cicatricial tissue, and after development up to a certain point, atrophy and absorption occur.¹

Nerves.—In discussing the regeneration of nervous tissue, three component parts have to be considered—the neuroglia, the neurons, or nerve cells, and the nerve fibres, or peripheral portions of the neurons. In addition, it has to be kept in mind that there is yet another element, both in the central nervous system and the peripheral nerves, namely,

¹ The more important papers on this subject are by Zenker, Regeneration des quergestreiftes Muskelgenebes, Leipzig, 1864; Waldeyer, Vireh, Arch., 34:1865:473; E. Neumann, Arch. f. mikr, Anat., 4:1868; 323: R. Volkmann, Ziegler's Beitr., 12:1892:233; Stendel, Diss. Tübingen, 1887; Askanazy, Vireh, Arch., 125:1891:520; Nauwerek, Ueber Muskelregeneration, Jena, 1890.

OF NERVES

ordinary connective tissue, not only of the pia mater and of the endoneurium and perineurium, but also within the substance of the brain and cord accompanying the vessels, and this plays an important, and, as usual, a disturbing part in arresting regeneration of the specific elements proper.

Neuroglia.-The glial cells originate, like the nerve cells proper, from the lining of the medullary groove, and form, it must be remembered, a connective tissue of epiblastic origin. In the early stages it is impossible to differentiate between the glial and the eventual nerve cells, and it would appear possible that, after atrophy of certain cells of the cord in the growing human foctus, either latent neuroblasts or glial cells can undergo development and replace the atrophied cells.⁴ In the amphibia, where the terminal portion of the spinal cord has been removed, there is a partial regeneration of the nervous elements, proceeding from the epithelial cells lining the central canal, and giving rise to glial cells, along with other cells provided with fibres, and so of the neuron type, though imperfect.² In the adult there is no sign of this transformation. But there is abundant evidence that these glial cells in the human adult retain their proliferative capacity. Not only can they form tumors (see later), but in wounds of the brain and cord they exhibit abundant mitoses and subsequent proliferation. Their proliferative capacity is, however, hindered by the greater activity of the fibroblasts, and eventually, in a wound, they form a relatively narrow, dense zone, within or underlying the connective-tissue cicatrix.

Nerve Cells.—Where the whole neuron or its cell body undergoes destruction, there is no regeneration in man or the higher animals. This must be regarded as definitely settled. At most, mitoses have been observed in the neurons following injury, but, as Sanarelli⁹ points out, these are imperfect, nor is there any indication that they lead to subsequent cell division and proliferation. Certain observers have described new-growths—true neuromas—containing nerve cells, and from this have concluded that presence of the latter indicated abnormal proliferation of neurons. (See Chapter XIX.) This conclusion is regarded as most doubful; such neurons are either inclusions in the tumors (Solokoff), or are of the nature of cell rests, portions of nerve tissue isolated during the course of development and incapable of coordinated function.

There is no regeneration of gray matter of the brain or of the cord. The same is true, also, of the sympathetic or spinal ganglia.⁴

Peripheral Nerves: Nerve Fibres.—The essential portion of a peripheral nerve fibril, that establishing the communication between the nerve body and the peripheral organ or other neuron, is the axis cylinder. The older, well-established view is that this is a direct outgrowth and portion of the neuron. For a time the view gained credence that it is formed by relavs—that the cells of the sheath of Schwann practically

s of ima regjust the n a gess miigh nen lay on, es, obliin eir rk-

to

us

he

he

de

r-

ff,

1-

р

e

r

¹Adami, Jacobi Festschrift, 1900. ² See Caporaso, Ziegler's Beitr., 5: 1889: 67.

³ Acad. dei Lincei Ser., 4:7:1890 (Ref. Centralbl. f. Path., 2:1891:429).

⁴ Monte and Fieschi, Arch. ital. de Biol., 24: 1895: 401; Tirelli, ibid., 23: 1895: 301, 40

govern the development of successive sections, and many authorities saw in the facts of regeneration a considerable amount of support for this view.¹ Fortunately, this may be laid down with precision, that any new development of an axis cylinder always originates from a preëxisting axis eylinder, and, what is more, from such axis cylinder in organic connection with a neuron. Solution of continuity of a nerve fibre or of its axis cylinder leads to degeneration distalward of the distal portions separated from the cell body, and of the proximal and still connected portions upward as far as the next node of Ranvier, or, in some cases, a node or two higher.

There are not a few debatable points regarding this Wallerian degeneration and its extent that are of great importance; they are, however, not immediately germane to our present inquiry, and we must forbear to dwell upon them. Such, for example, are the questions of arrest of degeneration in the peripheral section of a cut nerve, by maintaining the tone of the muscle to which it runs, the question of the existence of peripheral nerve cells, and peripheral regeneration.²

Regeneration occurs, provided (1) that the shock (as in tearing out whole nerves) has not been too severe and the destruction has not led to complete arrest of function of the nerve cell; (2) that the organ or part innervated has not been destroyed or does not become atrophied and degenerated in consequence of the severance of its nerve supply; and (3) that the path of the regenerating fibres does not become blocked by cicatricial tissue.

In regard to the second of these conditions it is to be noted (and the same applies to the third) that regeneration of an imperfect order may show itself in these cases; it may commence, but is unable to effect any satisfactory result. This is well seen in the amputation neuroma, so called, in which the axis-cylinder processes grow out from the end of the severed nerve, but become wound and twisted in all directions its the connective-tissue overgrowth of the perineurium and endorium, a nodular mass resulting. We call attention to the degeneration of the organ innervated in consequence of certain important observations of our colleague, Dr. Shirres,3 that if, by massage and electrical stimulation, the healthy conditions of the muscles which are supplied by the severed nerves be preserved, even in the spinal cord there are indications that the axis-cylinder processes are capable of some regeneration, a result which previous observers had failed to attainapparently because this preservation of the innervated parts has not been sought after-and so had denied its possibility. Forsmann⁴ would seem to have approached this point in his recognition, as the results of many experiments, of a form of "chemiotropism," or positive neuro-

⁴ Galeotti, and Levi. Ziegler's Beitr., 17:1895: Kennedy, Phil. Trans. Roy. Soc. Biol., 1877;188; Wieting (under Marchand), Ziegler's Beitr., 23:1899;42.

² Upheld by Bethe, Allgem. Anat. u. Physiol. d. Nerrensystems, Leipzig, 1903, and by Stewart and Ballance, The Healing of Nerves, London, Maemillan, 1901.

³ Montreal Med. Jour., 34: 1905: 239, ⁴ Ziegler's Beitr., 24: 1898; 56,

OF NERVES

tropism, attracting the newly formed axis cylinders downward to join the distal portion of the nerve. Apparently, that is, the condition of the muscle has some influence upon the distal, severed portion of the nerve in connection with it.

With reference to the third condition, as we noted in connection with muscle, so here, regeneration is most complete when there is not complete section, but only contusion and destruction of the axis cylinder by any means, without rupture of the sheaths of Schwann.

There is, indeed, an interesting parallelism between the muscle and the medullated nerve fibre in this, that both are compound structures provided with a sheath (the sarcolemma and neurilemma, respectively) closely applied to which in normal development are the specific muscle nuclei in the one case, the nuclei controlling the myelin sheath in the other.

Considering this third, simpler case first, the stages in connection with regeneration are the following:

The first result of injury in the course of a fibre is traumatic degeneration. This is apt to show itself first in the medulla, the myelin dividing up into irregular masses and globules; it is followed by fibrillation, imperfect staining, and disintegration of the axis cylinder. The fragments of the myelin sheath become smaller and more numerous, and by the second day these conditions are very pronounced, the disintegrated axis cylinder becoming wholly unrecognizable.

But by the second or third day there is already noticeable a distinct proliferation of the nuclei of the sheath of Schwann. They multiply by direct division, no longer lie immediately beneath the neurilemma, but pass between the myelin globules, and, like the future sarcoblastic nuclei under similar conditions, it is to be observed that they gain a surrounding cytoplasm, increasing in amount. Mitoses, with further multiplication, may become evident on the third and fourth day, and, as with hyaline, degenerated muscle masses, so here, the myelin droplets diminish as these cells grow and enlarge. Of these remarkable cells,1 while some degenerate and disappear, others become elongated and spindle-shaped. As to their further development, there is still some debate. It is generally admitted that they give origin to the new sheath of Schwann and to myelin. It is in regard to their connection with the new axis cylinder that there has been difference of opinion. We see that the new axis cylinder originates from the central end of the damaged nerve, not necessarily from the last node of Ranvier; some of the fine young processes may be traced to an origin higher up the nerve. We see, also, that there has been merely crushing, without destruction of the sheath, the axis cylinders, often multiple, or, more correctly, with fine aberrant fibrillæ, proceed down the old sheath surrounded by the myelin cells, and, doing this, pass eventually beyond the region of injury into the distal undamaged part of the nerve.


¹ Now recognized as being like the glial cells, of epiblastic origin (Verocay, Ziegler's Beitr., 48; 1910).

Recent studies, more particularly those of Marinesco¹ and Ross Harrison, throw light upon the positive neurotropism of Forsmann above mentioned. It is seen that the end of regenerating axis cylinder is not filiform, but formed of a nodular mass of protoplasm possessing a most remarkable power of apparently pseudopodial motion. Following the development of the axones in removed portions of the primitive nerve tube of the larval frog, Ross Harrison, of Johns Hopkins, has shown that this nodular mass creeps forward, in this way lengthening the axone behind it, and this at the rate of about 1 micromillimeter in two minutes. The regenerating axone creeps thus across the area of destruction and along the lines of the old sheaths of Schwann until it enters the nerve bundle beyond. Not all necessarily reach this; some become diverted and arrested in the damaged area, but eventually a certain number of the swollen ends can, by appropriate staining methods, be detected in the distal nerve bundle. Presumably these fuse with the still intact axones of this distal portion: the exact mode of fusion has not been followed (Figs. 183 and 184).

Regeneration after Section or Rupture.—Unlike what occurs in the divided muscle fibre, there is, so far as we can see, no marked difference in the behavior of the myelin nuclei (or nuclei of the sheath of Schwann) when the nerve fibres have been divided, from what occurs when they remain intact, save this, that they wander out of the sheath, and, wandering in various directions, the axis-cylinder processes also are apt to curve and be distributed very irregularly, some wandering directly backward instead of forward. Whether, under these conditions, complete *functional* regeneration occurs, depends upon whether any considerable proportion of the new fibres find their way to the tract of the distal portion of the nerve, and is governed largely by two conditions, namely, (1) the distance apart of the two ends of the severed nerve, and (2) the extent of the cicatricial formation between these two ends.

For complete restoration of function many months may be necessary, and experiments show that the wider apart are the two ends of the nerve the longer is the time necessary (*ccteris paribus*). Beyond a certain distance no results are obtainable. Vaulair, in the dog, had negative results when a length of 4 cm. of the vagus was excised, though where a tubular-bone suture, or canal, was placed between the two ends, Huber obtained regeneration after excision of 6 cm. of the ulnar nerve of a dog (see also Tiedemann's results below). Regeneration, indeed, is materially aided by affording a path along which the new fibres are directed, whether a hollow bone (Vaulair), a small artery from some other animal (Bünger), a bundle of catgut threads (Assaky, Gluck, etc.), a lappet or slip from the nerve bundle itself turned over to join the severed ends (Létiévant), a length of nerve from another animal, etc. Where suppurative inflammation and leukocytic inflitration

¹ Marinesco and Minea, Revista Stüntelor Med., Bucharest, 1905, No. 5. For a fuller description of recent studies see Halliburton, Science Progress, 2: 1908; 413, and Mott, Trans. Rov. Soc. of Medicine, Path. Sect., 1909–10.



Regenerating nerve fibres. These are of varying thicknesses (A, B, C), and each is provided with a terminal swelling. The figure also shows the marked tendency on the part of a certain proportion of the regenerating fibres to take a spiral course (M). (Marinesco.) Longitudinal section of nerve from dog twenty-one days after division: A, central end. B, cieatrix of union in which the nerves take a very irregular course. Many regenerating fibres with club-like ends; "", "", ""," have already penetrated into C, the peripheral end of the divided nerve. Some, like m", have become so diverted in the cieatrix as to be turning directly backward. Others, " and "', growing from above, have not yet reached the cieatrix. (Marinesco.)

m

m"

-m4

occur, it will readily be understood that there is subsequent dense cicatrization, forming a barrier preventing union. There are, however, some exceptional cases on record in which, eventually, regeneration has taken place. Of these, one of the most remarkable is that of Tiedemann, in which, after excision of from 10 to 12 cm. from the brachial plexus of a dog, almost suddenly, at the end of two years, there was complete restoration of function in the limb. In fact, many of the negative observations upon this subject of regeneration by one or other means of favoring the passage of the fibrils downward appear to have allowed too short an interval before making the record.¹

⁴ Other important papers on this subject of the regeneration of peripheral nerves are: Howell and Huber, Jour. of Phys., 13 and 14: 1892, 1893; Huber, Jour. of Morphol., 11: 1895; 629 (a classical study of the subject); Stroebe, Ziegler's Beitr, 13: 160, and Centralbi, f. Pathol., 6; 1898 (a useful review of this literature); Vaulair, Areh. de Biol., 3: 1882; 379, and Areh. de Physiol., 8: 1886; Willard (nerve suturing), Internat. Med. Mag., April, 1894; Assaky, Areh. gén. de méd., 1886.

CHAPTER XIII.

GRAFTING OR TRANSPLANTATION.

HERE, as there is a tendency to use terms somewhat loosely, it may be well to lay down that under *implantation* we include all processes of inserting solid matter into the tissues of a living animal, whether living or dead tissue, or inert material of animal origin, whereas *transplantation*, or grafting, refers only to the one order of cases of insertion of living tissue. Such transplantation is spoken of as *autoplastic* when it is sought to graft tissue from the same individual; *heteroplastic*, where the tissues of another animal are employed. *Replantation*, the replacement of an organ or tissue after removal (c. g., tooth, end of nose, etc.), explains itself.

Such grafting is a common operation in gardening and arboriculture, and there it has been known for centuries that not only is it possible to mate and gain perfect organic autoplastic or heteroplastic union, but that successful grafts can be made upon a stock of wholly different species. It generally is found that the more nearly allied the species the greater the measure of success attained. Nevertheless, the growth of the parasitic mistletoe upon the apple and other trees shows how widely apart may be stock and graft.

There is now no doubt, more especially from the observations of Joest¹ upon earth-worms and of Born² upon amphibian larvae, that in lower animals transplantation is possible to an extraordinary extent. Thus, the latter was able to join together larvae, or parts of larvae, of different species, and, taking individuals of one species and cutting out portions of their bodies, he could replace these with like portions of the bodies of other individuals, and found that, if parts were accurately applied, corresponding parts of such organs, as the heart, the intestine, and nerves would fuse neatly together, and that growth continued with functional unity.

When, however, we come to study warm-blooded animals, we find that the capacity no longer obtains to anything like the same extent. Ribbert,³ Lubarsch,⁴ and many others have conducted full and most careful studies upon the effects of transplantation of very many tissues with eventually little more than negative results. Transplantation of the tissues of animals of another species is practically wholly ineffective.

¹ Arch. f. Entwickelungsmechanik, 5: 1897: 419.


2 Ibid., 4: 1896.

³ Ibid., 6: 1898: 131, etc.

⁴Zur Lehre d. Geschwälste, Wiesbaden, 1899. The observations of Paul Bert, De la Greffe animale, Paris, 1863, call for mention; they were performed on the rat, and were very extensive and more successful than those of later observers.

GRAFTING OR TRANSPLANTATION

In the heteroplastic transplantation various portions of various glands of the animals of the laboratory into regions well supplied with blood —into the liver (Lubarsch) or the abdominal lymph glands (Ribbert) —it is found that for a time there is complete healing in, and complete union, eventually absorption and disappearance of the transplanted tissue, takes place, a fibrous cicatrix alone remaining.



Heterophastic transplantation in the earth-worm: A, of tail end of another individual of the same species (Alloloophora terr.); B, intercalation of mid-body region of another individual; C, lateral grafting of anterior half of another individual. (Joest.)

There has, it is true, been evidence of primary growth; thus, the ducts of glands, especially, have shown mitosis and active proliferation, and the appearance of the tissue has shown what we elsewhere speak of as reversion, or reversionary degeneration, the cells assuming a simpler,

FIG. 186

Transplantation in an oblique plane between two distinct species of earth-worm, Lumbricus rubellus and A. terrestris. (Joest.)

more embryonic type, and with this the parts which, developmentally, are found to exhibit most active growth now show the same properties. In bone transplanted into a lymph gland, also, the periosteum may remain active, and there may even be new development of either cartilage

IN THE LOWER ANIMALS

or bone. But within a few weeks or months all the specific cells of the transplanted tissue atrophy and disappear. This has been the experience with such tissues as liver, kidney, and testis.

Embryonic Tissues.—What at first appear to be satisfactory results are gained by grafting embryonic tissues into the fully grown animal. Knowing the active vegetative capacities of the embryonic cells, this is readily understandable. Such observations have been carried out more particularly by Ribbert,1 Birch-Hirschfeld,2 and Feré,3 The latter observer, planting portions of forty-eight-hour chick embryos under the skin of young chickens, gained, rarely, progressive growth, the growth persisting and being recognizable in one case for five months, in another for thirty-three months. In one case, by the end of two months, small black feathers appeared on the graft, in another, simple epithelium. As a rule, the "tumors" were composed of mesodermal elements, vessels, plain muscular tissues, etc., with, in one case, cartilage. But in the majority of the cases, by the end of two months, the grafts began to diminish and undergo absorption. The same was true in Ribbert and Birch-Hirschfeld's cases, carried out more particularly with pieces of rabbit embryos. In these there was found a greater persistence of cartilage than of other tissues, and this at times formed masses of considerable size, which, however, eventually underwent regressive changes and atrophy. This growth of perichondrium and cartilage has been noted by several observers.

Hunter's famous experiment' upon transplanting the cock's spur into the cock's comb shows, in general, the same phenomena. There is a primary, most active growth, and the spur with its bony centre may attain the length of some inches. Such spurs have been reported as still being present at the end of two years. More often, after a few weeks or months, they undergo atrophy and fall off.

Thyroid.—More satisfactory results follow transplantation of the thyroid. Von Eiselsberg⁶ first transplanted one-half of the cat's thyroid into the animal's abdominal wall, and when this was healed and appeared to have united, he transplanted the other half into the abdominal wall or cavity. The animals so treated bore the operation well and showed no symptoms, but so soon as the transplanted portions were removed they rapidly died with the symptoms of tetany which follow extirpation of this gland in the cat. Munk,⁶ Enderlen,⁷ and Sultan⁸ have confirmed these observations, and have proved that both in the cat and the dog the central parts of the body undergo necrosis, the peripheral follicles remaining

¹ Loc. cit.

² Ziegler's Beitr., 26: 1899: 132.

³ Compt. rend. de la Soc. de Biol., 1895 and 1897 (several papers); also Arch. d'Anat. microscop., 1: 1897; 193 and 417.

⁴ This experiment, however, it must be noted, was the repetition of an experiment by Aldrovandi in the sixteenth century, confirmed by Worm in 1685 and by Duhamel in 1746.

⁵ Wiener klin. Woch., 1892; 5.

⁷ Mitth. a. d. Grenzgeb., etc., 3: 1898.

⁶ Virch. Arch., 150: 1897: 271.
⁸ Centralbl. f. Path., 9: 1898.

GRAFTING OR TRANSPLANTATION

unaffected, while as granulation tissue, derived from sucrounding parts. passes in between these into the central necrotic area, there is an actual growth of the thyroid-gland tissue, in the shape of solid cell processes or buds given off from the persisting follicular epithelium. These become separated off, and eventually, with secretion of colloid, may gain a lumen and become quite typical. Lubarsch¹ transplanted thyroid tissue into the kidney, found the new-growth imperfect, and eventually, within six months, becoming atrophied and absorbed. Transplanting into the abdominal cavity, other observers have gained more successful results. Thus Enderlen, grafting the dog's thyroid into its abdominal cavity, found that, at the end of five and a half months, the tissue there was of the normal type, and Christiani,² making autoplastic grafts of the cat's thyroid into the abdomen, found the graft throughout glandular, very vascular, and the follicles full of colloid at the end of two years; and was justified thus in concluding that the organ can function in its new position during the whole course of the animal's existence.

Mammary Glands.—Ribbert's observations indicate that the mammary glands of young individuals, if grafted subcutaneously, are capable of growing permanently in this new position. Taking a guinea-pig a few days old, he grafted the mammary glands below the ears; the skin did not heal over them completely, and when, five months later, the animal became pregnant, the glands underwent enlargement and secreted milk, while at the same time new mammary glands showed themselves in the normal position, apparently from regeneration of portions left behind.

Ovaries.—'There remain to be mentioned those more important instances in which, in the higher animals, transplantation would seem to be attended by continued growth and vitality of the transplanted tissue, namely, the transplantation of the ovaries and of the skin and of the periosteum.

As pointed out by Knauer³ and by Grigorieff,⁴ if *both* ovaries of a rabbit be transplanted to other regions of the peritoneum, it is found that while the central portions of these organs necrose, the other portions remain normal, exhibiting normal Graafian follicles, and successive corpora lutea—and the follicles still produce ova. Three of Grigorieff's animals became pregnant after the operation, and Knauer notes the birth of young sixteen months after the transplantation. The transplantation would, therefore, seem to be perfectly successful, although further experiments are necessary to determine the exact length of time during which the ovaries continue to perform these functions. Like observations were conducted, some years earlier, in the human female by R. T. Morris,⁵ in one case, in a girl, aged twenty years, who had never menstruated. Menstruation followed the grafting of part of the erany of another woman into the uterine wall; in another, after the removal of both ovaries and tubes, he grafted a portion of one of the removal.

¹ Loe, eit., p. 251.

³ Centralbl, f. Gynäk., 22; 1898; 201,

⁵ New York Med. Jour., 62: 1895; 436,

² Arch. de Physiol., 7: 1895: 65.
 ⁴ Ibid., 21: 1897: 663.

SKIN GRAFTING

ovaries upon the stump of the right tube. Pregnancy followed later, ending in abortion at the end of the third month. Knowledge of the data gained regarding cytolytic phenomena will prepare the reader to realize that the surest success is gained with autoplastic transplantation; that just as the blood serum of one animal of a species may be hemolytic for the corpuscles of another member of that same species, so there may be antagonism between the tissues of the host and the graft leading to cytolysis and absorption of the cells of the latter. It is demonstrated clearly from these and allied experiments that the continued growth of the graft is intimately dependent upon functional activity. Remove only one ovary and the likelihood is that the grafted ovary undergoes early absorption; remove both, and even a heteroplastic ovary not merely gains perfect union, but continues to perform function.

Skin Grafting,—The facts in connection with transplantation of the skin are so well known that here we need but summarize the results obtained:

(a) The deepest layer of the epidermis—the Malpighian layer—most surely undergoes proliferation when skin is transplanted; thus, for successful operation, the grafts must pass down well into the papillary layer of the cutis. It is true that, as McLeod has pointed out, these deeper cells can be obtained by blistering and employing the blister serum, but the surest results are obtained by Thiersch's or by Krause's methods, in which the papillary layer or even a large portion of the cutis is removed with the graft.

(b) In all cases the greater part of the graft dies, but as the vessels of the underlying granulation tissue make their way through the cutis and reach the under surface of the grafted epidermis, the cells of the Malpighian layer exhibit mitosis (generally about the third day) and now undergo active multiplication, spreading out in centrifugal manner. As pointed out by Loeb,¹ these new cells have amceboid properties.

(c) In this way the denuded area becomes gradually covered with a new epithelium, which, however, is unprovided with hair follicles or sweat glands; the more differentiated portions of the skin are not reproduced.

In some cases, at least, this new transplanted epithelium is quite permanent, although Loeb's interesting results, obtained by transplanting pigmented skin (in guinea-pigs) into unpigmented areas, and vice versa, would show that unpigmented skin so transplanted gradually becomes pigmented; that there is a definite migration of pigment cells into the surrounding epidermis; that in albinos pigmented grafts become, eventually, colorless; and they raise a doubt as to whether, after all, there may not be a gradual replacement of the graft, piecemeal, by cells derived from the epithelium of the "host," and whether the continued vitality of the graft is not more apparent than real.

Certain remarkable results have been obtained by Allen² and others, by the employment of the skin of frogs and other animals for the purposes

¹ Medicine, Chicago, March, 1898.

² Lancet, London, 1884; ii; 875.

GRAFTING OR TRANSPLANTATION

of grafting. For a time much attention was called to experiments and the surgical employment of grafts of this nature. We may sum up the results of a fuller study by stating that in no case is the skin of another species found to be successfully grafted in man; in all cases the cells undergo necrosis. At the same time, there are indications that the existence of epithelial cells on the healthy surface of a wound, even when not those of the same species, have a stimulating effect upon the epithelium of the host, causing it to spread more rapidly over the denuded surface. It is difficult to explain the observations to this effect, save on the basis of the existence of a homotropism, an attraction of cells to others of like order, to the possible existence of which we have more than once referred (e. q_{e} , pp. 615, 619, and 627).

Mucous Membranes.—Mucous membranes show a like capacity for transplantation. Thus, several observers have, with greater or less cosmetic success, grafted the mucous membranes of the lips and mouth upon the conjunctiva and evelids.

Teeth and Bone.—The grafting of teeth and bone are frequently cited as examples of successful transplantation. In reality, they belong to a different order of phenomena, being examples of implantation.

It appears to have been known for some centuries in India, that after removal of a relatively sound tooth, a similar tooth removed from another man and placed in its socket becomes perfectly united, and healed in, and in the time of the Roman Empire there are indications that something was known of the implantation of artificial teeth formed of bone, while, following upon Ambroise Paré, the employment of freshly drawn teeth from one individual to replace those of another seems to have been somewhat frequently practised in France and England, until the conveyance of syphilis, in several instances, threw the procedure into discredit.

Here it need only be noted that equally good results are gained, whether an entirely fresh and healthy, newly drawn tooth is employed, a tooth that has had its pulp removed, one that has been thoroughly sterilized so as to kill off any living cells, or one that has been out of the body for years. There is not, that is to say, organic union in the strict sense. Vessels, and even nerves, may penetrate into the pulp cavity, and osteoblasts also passing in, it may eventually become filled with bone and so become firmly fixed; but in all cases—even those in which the recently drawn tooth retains active alveolar periosteum, which can become grafted on to the alveolar periosteum of the jaw, and so cause firm fixation of the cement substances—the tooth proper is an inert substance whose scanty cells do not persist. And, while in some cases the implantation is successful, in a considerable proportion the roots become absorbed, and the tooth loosens and eventually falls out.

We observe very largely the same order of events in connection with bone. The implantation of dead, sterilized bone, or particles of such, whether of man or the rabbit, or even, following Senn's method, of decalcified bones, gives every bit as good results—in some respects better results—than does the living or recently removed bone of the individual

CARREL'S EXPERIMENTS

grafted upon. Employing this latter, the bone corpuscles examined a few days later are found dead and non-staining. Such bone, in fact, forms, like any other porous material, a framework into which may penetrate the cells and vessels from the surrounding periosteum and living tissues. It acts, in short, very much as does the fibrin and bloodclot found in a wound under natural conditions, with the additional advantage that it is rigid and is so more likely to preserve the natural contour of the part.

Periosteum.-With periosteum, as again with perichondrium, the case is very different. The autoplastic transplantation of both is most successful and most often the heteroplastic (i, e_i) from one individual to another of the same species), although here the younger the animal that affords the graft, the greater the measure of success. Saltykoff's¹ observations indicate that in the days following the operation the greater number of the periosteal cells undergo necrosis; a certain proportion of them, in the inner osteoblastic or cambium aver, remains alive, and by the third day shows mitoses; in five days the proliferation of these cells is abundant and they form into rows of osteoblasts, which give rise to new bone. It is the inner layer of the periosteum that is active in this bone formation. There may be a preliminary formation of hyaline cartilage followed by the development of the bone. In general, it is to be noted that transplantation into the bloodvessels (Cohnheim and Maas²) or into the soft tissues does not lead to such perfect results as when the transplantation occurs over old bone or in the area of previous bone. In the first of these cases there is eventual absorption, in the last the formation of apparently normal bone, even to the development of a medullary cavity.

With regard to bone marrow the results of different observers have been contradictory, but by the autoplastic grafting of red marrow under the skin Bruns³ gained the formation of cartilage, osteoid tissue, and, after twenty-two to twenty-four days, of true bone, results which were confirmed by Kölliker⁴ as regards transplantation into the anterior chamber of the eve and the abdominal cavity.

Carrel's Experiments.—The above was the state of our knowledge on this subject until within the last few years, when certain remarkable observations of Carrel⁵ have materially altered our point of view. With great surgical ingenuity, Carrel has developed the method of vascular anastomosis and union, and of the accurate adaptation of nerves, ducts (like the ureter), etc.; and by this method he has proved convincingly that, *provided the circulation be restored fully*, not merely can arteries and other vessels from one animal be transferred into another, remaining functional and apparently healthy for long weeks, but that this may even happen when the graft is taken from an animal of a different species,

¹ Arch. f. Entwickelmech., 9:1900.

² Virch. Arch., 70:1877:161.

³ Arch. f. klin. Chir., 26,

⁴Centralbl. f. Chirurg., 1881: 577.

⁵ Proc. Soc. Exp. Med., 4: 1907; Carrel and Guthrie, Science, 22: 1905; 473, and 23: 1906; 394.

GRAFTING OR TRANSPLANTATION

and even when the organ to be grafted has been preserved for a day or more in the ice chest; further, that organs of very considerable size, such as the kidney, may be transplanted from one animal to another of the same species and exhibit evidences of normal function.⁴ He has recorded cases in which he has transplanted the whole limb from one animal into another, with indications of at least temporary success. It is but right to sound the warning that these successes are extraordinary in a double sense. It is no disparagement of Carrel's wonderful skill to point out that results of this order are the exception rather than the rule where heteroplastic transplantation of organs is attempted. It is to his credit that he has demonstrated that this is possible. Nor should the surgeon who has not thoroughly familiarized himself with the technique of vascular union even dream of performing similar operations upon man.

Conclusions.—We reach, therefore, the following conclusions:

 In lower animals (as in plants), transplantation and grafting is possible to a remarkable extent, but in the higher warm-blooded animals it is possible to but a very limited extent, unless there be gained an accurate adaptation and anastomosis of the nutrient vessels.

2. In the latter, without such anastomosis, there may be temporary success, the grafted part gaining complete union and showing cell multiplication; but, with relatively few exceptions (epidermis, mucous membranes, periosteum, perichondrium, thyroid, and ovaries), the cells of the graft sooner or later undergo atrophy and absorption.

3. Autoplastic grafting is more successful than heteroplastic, and this, again, than grafting with tissues of a different species. In vertebrates the latter is only possible where immediate vascular anastomosis is brought about between the vessels of the host and of the graft.

4. The more vascular the site of transplantation, the greater the likelihood of obtaining (temporary) union.

5. The younger and more actively proliferating the tissue composing the graft, the greater the likelihood.

6. The more the graft is in position to satisfy the needs of the organism and to actively function, the longer, in general, would its cells appear to retain their vitality, *c.g.*, portions of liver grafted into other organs when the liver as a whole is still functioning rapidly degenerate; the thyroid or *both* ovaries removed from their natural site and transplanted elsewhere retain their functions. The skin transplanted on to a superficies forms a perfect graft.²

¹ Jour, of Exp. Med., 10: 1908; 98,

² Quite the most thorough study of this subject of transplantation and implantation will be found in the work to which we have already more than once referred, Marchand's *Die Wundheilung*. The literature is there given very fully.

CHAPTER XIV.

METAPLASIA AND HETEROPLASIA.

METAPLASIA.

IF an eve that through traumatism has been rendered functionless be removed some years later, it is, in general, found that from the choroid coat there has developed a layer or deposit of true bone. In one case, studied in our laboratory at the Royal Victoria Hospital, by Dr. Mathewson, not only the choroid, but also the lens was definitely implicated in this bony formation. To account for this remarkable development in a region where, normally, bone is wholly wanting, three theories may be adduced: (1) That in the process of formation of the eve, a few cells destined to form bone become accidentally carried into the eye, along with the invaginating membranes, and remain latent and inactive so long as the organ performs its functions, but when by accident it becomes functionless, then the altered conditions are such as to favor the active growth of the cells and the eventual formation of true bone; (2) that the bone formation is due to modified function and nutrition of certain choroidal (connective tissue) cells. Normally, in the uniniured eye, the choroidal cells have definite duties, and their activity appears to bear a direct relationship to the light-receiving function of the eye. When either the anterior or posterior chamber of the eye is injured, the functional activity of these cells is arrested; the vascularity of the choroid undergoes modification, and now certain of the choroidal cells become modified in their action and form bone; (3) the third theory, that of Ribbert, of conveyance of bone-forming cells to the part by the blood, we will discuss later (p. 647).


The first of these theories must, I think, be discredited. If we came across this choroidal bone formation only exceptionally, it might well be urged, but the eventual formation of this bone in the useless eye is the rule, not the exception. Holding to this theory, we should have to hold that the inclusion of aberrant tissue or "mother cells" in the developing eye—and in other_developing organs—is a matter of constant occurrence. As we shall proceed to show later, while we must admit the occurrence of these "cell-rests," and admit that they are not uncommon, we cannot believe that cell-rests of osteoblasts are practically constant in the coats of the eye. We have absolutely no ground for such an assumption. In studying normal tissues under the microscope, it is only very exceptionally that we encounter appearances which we can ascribe to the persistence of cell inclusions. In other words, metaplasia, or the postnatal production of specialized tissues from cells, which normally produce

METAPLASIA AND HETEROPLASIA

tissues of other orders, affords a more satisfactory explanation of the appearances above described than does the theory of *cell-rests*.

We are apt to forget that metaplasia is a constant physiological process. The conversion of ordinary connective-tissue corpuscles into fat cells is a typical example of metaplasia, and in early natal life, as again as the result of regeneration, it is now clearly established that cartilage cells and cartilaginous tissue may undergo direct transformation into bone cells and bony tissue.

Before proceeding to enumerate what are to be regarded as examples of true metaplasia, it is necessary, with Schridde and Orth,¹ to note the conditions which cannot be included under this term.



Island of squamous epithelium in cervix of uteri of newborn infant (? developmental metaplasia or inclusion). (R. Meyer.)

Heterotopia, congenital or acquired, is the abnormal snaring of cells of an organ or tissue with subsequent growth out of place. To the congenital form belong the various "cell-rests" of Cohnheim, aberrant adrenal nodules, accessory spleens, etc.; to the acquired, traumatic subcutaneous epidermal cysts ("see Cysts"), periosteal and bony growths from displaced periosteum, etc. In none of these cases is there tissue transformation, but continued growth and cell differentiation along the ordained line. A variety that may be mistaken for metaplasia is brought about by the invasion of one tissue into the territory of another as the result of trauma or inflammation. Thus, after tracheotomy, the epidermis may not only cover the wound but grow some little distance down the trachea, supplanting its columnar celled epithelium, or in inflammation of the cervix uteri the squamous epithelium of the os may spread upward toward the uterine cavity.

¹ Sixteenth International Congress of Medicine, Lisbon, 4 B, Section,

Heteroplasia.-We owe the clear recognition of this condition to Schridde,1 Careful examination shows that in practically every œsophagus examined, about the middle there are to be detected one or more small islands of columnar epithelium resembling that of the gastric mucosa. Now the hypoblast gives origin in the main to a columnar celled epithelium (respiratory passages, alimentary canal and its appendages from the stomach downward), but in the cosophagus the lining layer becomes converted into a squamous epithelium with cell bridges and fibrils wholly like those of the (epiblastic) skin. There is in these islands no conversion of the one type of epithelium into the other, but a persistence of developmental tendencies. A similar explanation has been afforded for the occasional islands of squamous epithelium encountered in the respiratory passages, stomach, gall-bladder, and cervix uteri in cases in which there is no sign of inflammatory or other disturbance which might account for the change. It cannot be neglected, however, that under the influence of chronic irritation, somewhat similar areas of squamous epithelium in the above regions may be examples of true metaplasia.

Anaplasia or Reversionary Atrophy (*Entdifferenzierung*).—The mere loss of differential characters by cells subjected to abnormal conditions is not metaplasia, *e. g.*, the simplification of the cells lining the convoluted tubules in cases of granular contracted kidney, or the reversion of the cells of the pulmonary alveoli to a large, more or less cubical form, in cases of chronic interstitial pneumonia with fibrosis and contraction of the interstitial tissue.

In this category must be placed the development of mucoid tissue in adult life. For mucoid tissue developmentally is always found as an intermediate stage in the growth of some other mesoblastic tissue. How such retrogressive change may be associated with active growth will be pointed out in discussing neoplasia, when also the subject of anaplasia will be seen to have an important bearing. Closely allied to this is what Orth terms allomorphism, simple morphological change of cells due to mechanical action, e. g., the flattening of a cubical or cylindrical epithelium lining a cyst, due to the distension and pressure of the contents of the same; as again is Schridde's prosoplasia, the further differentiation under modified environment of a tissue already performing a particular function, as, for example, the cornification of the epithelium of the prolapsed vagina.² Metaplasia comprises both morphological and functional change.

Physiologically and pathologically, we are forced to recognize that there are certain narrow limits bounding metaplasia, at least after birth. *Epithelial (epiblastic and hypoblastic) tissues can only be converted into other forms of epithelial tissue, one form of mesoblastic into another form of mesoblastic*. Epithelium and gland cells, for example, never

¹ Virch. Arch., 191:1908:178.

² While of squamous type, the normal epithelium of the vagina, as of the mouth and œsophagus, exhibits no keratinous change.

METAPLASIA AND HETEROPLASIA

become converted into bone or cartilage, or vice versa, while, again, it may be laid down that among epiblastic and hypoblastic tissues, on the one hand, and mesoblastic tissues, on the other, there is no new development or metaplasia of the most highly specialized tissues from less specialized tissues; a simple epithelium cannot in the vertebrate give rise to the more complex glandular tissue, or to nerve cells; in regeneration of epithelium there is no new formation of hair roots or cutaneous glands. The cells of white fibrous connective tissue have not been seen to form striated or even non-striated muscle. Within these relatively marrow limits numerous examples of pathological metaplasia present themselves.


FIG. 188

Metaplasia from a case of ectopia of the bladder: the ordinary squamous epithelium becomes replaced by a columnar epithelium. (After Enderlen.)

Epithelial.—The mucous membrane of the uterus is a columnar epithelium; if the organ be everted so that it projects from the vagina, forming a pear-shaped mass, exposed to the air, the mucosa covering it becomes eventually smooth and dry, and now, upon examination, in place of the columnar there is found a stratified squamous epithelium, of which the outer layers may show very definite keratinous or horny change, as in the true skin. Similarly, as the result of chronic irritation the columnar, ciliated epithelium which covers the greater part of the larvnx may here and there show thickening, and, on microscopic examination, these thickenings are found to be formed of a squamous epithelium of many layers of flattened cells Dr. Long, working in our laboratory, has recently encountered two cases of multiple osteoid growths in the submucosa of the trachea over which the conversion of the columnar cells into a squamous epithelium with true prickle cells was admirably demonstrated. In the gall-bladder also similar transformation of the columnar mucosa has been observed following chronic catarrh.

METAPLASIA OF CONNECTIVE TISSUES

In the bladder there may be changes of two orders. This organ is lined by a distinct, but loose, pavement epithelium in several layers. As a result of chronic inflammation, either there may develop areas of wholly typical squamous epiderm, with prickle cells (this over the enlarged prostate), or overgrowth of a papillomatous type may be initiated, when in place of the squamous epithelium there is developed an epithelium of the columnar type. This is often noted in cases of ectopia vesice, and here even simple glandular crypts may show themselves (Fig. 188).





Stages in the metaplastic regeneration or formation of a new lens from the iris, in the larval newt: 1, edge of iris becoming swellen; 2, 3, 4, progressive overgrowth of the edge; 5, separation of the hypertrophied mass of cells to form the lens. (Fischel.)

The Lens.—One of the most remarkable examples of metaplastic regeneration has been studied by G. Wolff¹ and Fischel.² The lens in vertebrates is formed by an invagination of the *surface* epithelium of the head into the primary optic vesicle, which in its turn is formed by a similar invagination from the forebrain. The inner wall of this primary vesicle forms eventually the retina and iris. If, now, in the larval newt or salamander the lens be extirpated, in a short time a new lens is developed *from the iris*. Further observations have shown that the retina itself can produce

¹ Biol. Centralbl., 14: 1894.

² Abhandl, d. Deutsch, pathol. Gesell., 1902. For a discussion of the cases see Schwalbe, Morphol. der Missbildungen, Jena, 1906, Pt. 1: 88.

lens-like bodies, and that if the old lens be not removed but simply pushed to one side, the iris will form a new one. There can be no more remarkable example of cells of one tissue taking on the functions and properties of another. But it must be kept clearly in mind that the normal lens and the iris, even if of widely different origin, are both epiblastic. The case comes clearly within the limits noted by us, that epiblast can only produce epiblast; mesoblast, mesoblast. The same is true in Saxer's case of the development of an epithelium lining the cysts in gliomas (p. 565). While, as we have noted, we do not regard this as a proper epithelium, the cells in question, like the glia cells, are of epiblastic origin. In regenerating planarian worms, as Flexner' has pointed out, epithelium can even give rise to distinct nervous elements. Braun² has shown that the same occurs in the larval frog.

Mesoblastic.—The most marked examples are afforded by the metaplastic formation of *bone*. This may be:

1. From cartilage, as in the ossification of the laryngeal and tracheal cartilages in advancing age, in which it is to be noted that, as in all these true metaplasias, there is merely conversion of one tissue into another—replacement and not new-growth. There may, however, be some admixture of replacement and growth. Occasionally the smaller bronchi are found converted into rigid bony tubes. It is a question here whether we have to deal with metaplasia of the small cartilaginous plates in the bronchial walls alone, with subsequent slight growth and fusion, or with metaplasia affecting both the cartilage and the intervening connective tissue. The replacement of cartilage by bone in callus after fracture is an example of the same process.

2. Osseous metaplasia of connective tissue. In the lungs we may encounter small irregular spiculated masses of true bone, best explained as originating in this manner, while relatively large plates of true bone are to be met with in old pleural and pericardial adhesions, where the inflammation has been prolonged and the formation of new tissue excessive. Both bone and cartilage are occasionally met with in the arterial wall in arteriosclerosis, as also in the fibroid valve of chronic endocarditis. Professor J. J. MacKenzie³ and W. Harvey,⁴ of Toronto, and others have noted the liability for bone to form in the walls of arteries of rabbits that have been experimentally injured.

3. Osseous metaplasia of tendons. Occasionally the tendons of origin or insertion of a muscle are found replaced by true bone, forming large bony prominences. This may be the first stage of a remarkable and rare condition, to which the mistaken designation of *myositis* ossificans has been given.

There is no certain evidence that we have to deal with an inflammatory

² Jahresber, d. Anat. u. Entwick., 1903 and 1904.

⁴ Jour, of Med. Res., N. S., 12; 1907; 25. For cartilage similarly produced see Trachtenberg, Centralbl, f. Path., 17; 1906; 614.

¹ Jour. of Morphology, 14: 1898: 337.

³ Brit. Med. Assoc., Toronto, 1906,

OF MESOBLASTIC TISSUES

process, nor is it certain that the muscle fibres are primarily involved. In this condition, in the course of years, the muscle tendons and the bodies of one set of muscles after the other become replaced by bone, until at last the patient is unable to move his limbs, or to rotate the head or bend the back bone. Further and fuller studies are needed into the essential nature of this very remarkable condition.

dy

re

3.5

ni-


m

11-

SP

de

Cartilage.—The most marked example of cartilaginous metaplasia (except forms that are observed in connective tissue tumors) is seen in the development of the provisional callus of long bones after fracture (see p. 611). In connection with new-growths derived from the connective tissues (osteomas, fibromas, sarcomas, lipomas, myomas), islands of cartilage are frequently observable. In mixed tumors of cer-



Osseous metaplasia in the wall of a bronchus (so-called ''osteoma''): a, mucosa; b, submucosa; c, mucous glands; d, cartilage; ϵ , connective tissue; f f, masses of bone in submucosa; g, fat cells. (Demnig.)

tain organs, such as the parotid and testis, it is usual to ascribe this cartilaginous formation to the existence of cell-rests. A consideration of cartilage development in general leads us to consider that we have no adequate ground for separating these from other cases of metaplasia. The development of cartilaginous tumors, not uncommon in the mammary gland of the bitch (and very rare in that of the human being), must equally, we think, be regarded as originating from a primary metaplasia of the connective tissue of this gland. This, however, is not the prevalent view.

Of the metaplasia of fibrous tissue into cartilage and of cartilage into bone, the most striking instance that we have encountered is in the cases of osteoid development in the trachea already referred to. The cartilaginous rings of the trachea are developed from the inner aspect

METAPLASIA AND HETEROPLASIA

of the fibrous perichondrium. The specimens demonstrate with absolute precision that occasional cartilage cells may develop within the layers of the perichondrium, and that here and there, without any obvious covering layer of fibrous tissue, plates of true fibrocartilage form from the outer aspect of the fibrous perichondrial layer. In these in several places the cartilage cells can be observed becoming separated, more or less stellate, and surrounded by an osteoid matrix. In this way nodular projections become developed of laminated osteoid material possessing large marrow canals, with vessels and relatively abundant fat cells.⁴

Fibrous Tissue .- As might be expected, we meet with frequent examples of the conversion of more specialized mesoblastic tissues into the simpler fibrous connective-tissue type. As Thoma well points out, one of the clearest examples is met with as the result of immobilization of a joint by surrounding adhesions, etc., when the cartilages covering the opposed surfaces disappear, being replaced by connective and mucoid tissue. There is, it is true, a primary neoplastic development of vessels in the part, but the appearance of the fibrous and mucous tissue must be regarded not as a degenerative change (for inactivity of the joint would lead to atrophy), but as directly due to change of function. Immobility of the joint does away with the condition which necessitated joint and cartilage, and slowly a new-formed tissue develops. Whether the more highly differentiated mesoblastic tissues can undergo metaplasiawhether for example, the plain muscle cells in a uterine myoma undergo conversion into fibroblasts, or whether they undergo atrophy and replacement-is still a moot point.

Such metaplasia is in all cases to be regarded as an adaptation on the part of the cells to altered environment, not of necessity and primarily to altered function. One can, for example, recognize no change of function in the laryngeal and other cartilages with advancing life, but can recognize alterations in nutrition leading to absorption here and there. entrance of bloodvessels and conversion into bone, of the same order as, to employ an instance afforded by Lubarsch,² is the conversion of spherical unicellular organisms, like the Micrococcus prodigiosus, into bacillary or even spirillar forms under the influence of acid added to the medium of growth; and in the lower multicellular forms of animal life the modification thus brought about may affect not merely the cells. and groups of cells, but entire organs, when instead of metaplasia we speak of *heteromorphosis*. To this we have already referred (p. 603) and quoted the example of the regeneration of an antenna in place of an eve noted by Herbst³ in *Palinurus* and other crabs (Fig. 173). Another remarkable example, noted first by Spallanzani in the eighteenth century, and confirmed by Morgan, is that occasionally the earth-worm,

¹ Aschoff points out that this is not the universal form of tracheal "osteopathy;" in some cases the bony nodules develop in the submucosa unrelated to the periehondrium. Abstr. Centralbi. f. allg. Path., 21:1910;442.

³ Arch. f. Entwickelungsmech., 9:1899:260 and 13:1902:436.

² Allgemeine Pathologie, Wiesbaden, 1905: 53.

te

or

ng

it,

1g

nt

10

he

е,

er

of

re

n

whose head end has been removed, regenerates not a head, but a tail. Very similar in character are the cases to which Loeb first applied the term heteromorphosis. The hydra, the tubularia, and the anemone *Cerianthus* all possess a simple digestive *cul-de-sac*, the mouth opening into a digestive pouch. If in any of these an opening be made through the body wall into the digestive sac, so that fluid so soon as it flows through the mouth flows out through this opening, it is found that the opening, in a short time, becomes provided with a ring of tentacles and comes to resemble, and function as, a true mouth.

The cells here through primary alteration in environment are capable of taking on altered function—and so it is in the more restricted condition of metaplasia. To deny or to minimize metaplasia almost to the vanishing point, as is the tendency of certain pathologists at the present day, appears unwarranted; on the other hand, we are not prepared to proceed to the other extreme and recognize that in the developed individual the cells of hypoblastic and epiblastic origin can take on mesoblastic functions.

One of the strongest opponents of the frequent occurrence of metaplasia is Ribbert.¹ He admits the existence of physiological metaplasia, but criticises, with scarce an exception, the examples here brought forward. Metaplasia, he holds, is an "extraordinarily much rarer occurrence" than is usually held. Where in phthisis bulbi, or in the vessel walls or lungs, or in sites of old calcification there occurs development of bone, what happens, according to him, is the appearance of a richly cellular tissue which either is directly converted into bone owing to the development between the cells of a homogeneous ground substance, which undergoes further modification (as may occur in callus), or rows of osteoblasts are to be seen as in normal ossification. But, says Ribbert, this is no metaplasia. These cells and the marrow cells, which subsequently form, *may* have been derived from the blood and indirectly thus from the marrow of bones elsewhere.

This possibility has to be admitted, but is it the greater probability? We think not. How does bone arise first in the fœtus? From mesoblastic cells; and even in the periosteum of the adult the future bone corpuscles pass through a stage in which they are indistinguishable from fibroblasts, and in delayed union and in osteogenesis imperfecta² they may actually be converted into connective-tissue corpuscles.

It is the environment and relationship to the vessels and other influences acting upon them that lead certain mesenchyme cells in embryonic life to become osteoblasts and marrow cells. No satisfactory reason exists for denying that this may be in action later, just as all through life osteoblasts, which have never functioned as such hitherto, become converted into bone corpuscles. Similarly, he would explain the replacement of columnar by squamous epithelium as most often due to the existence of included islands of squamous epithelium, as due to heteroplasia, in Schridde's sense, followed by an invasive heterotopia. This,

⁴ Geschwülstlehre, Bonn, 1904: 5. ² Vide Klotz, Journ. of Path., 13: 1909: 467.

METAPLASIA AND HETEROPLASIA

however, does not explain the reverse condition seen in ectopic bladders, where the columnar epithelium makes its appearance. He admits that the development of a many-layered epithelium of squamous type may replace a single columnar duct epithelium, under conditions in which this explanation cannot hold, as in the extensive alteratian which he has himself figured as occurring in the submaxillary duct of the rabbit, after ligature of the same. 'This he explains as an "innate tendency" on the part of this epithelium, a derivative of the squamous epithelium of the mouth, to form squamous epithelium, a tendency which under ordinary conditions has not the opportunity to show itself and admits the same for like changes in the urinary passages under altered states. So he concludes that only tissues that, while externally different, possess, nevertheless, the same histogenetic capacities can undergo metaplasia one into the other. But this is what all pathologists will admit. The histogenetic capacities of a cell are brought out by its surroundings; all we would urge is that they are not so narrow as Ribbert would make them out to be.

The cases we are not prepared at present to accept are such as those placed on record by Leo Loeb, and confirmed, we may add, by other observers, in which in the course of epithelial regeneration certain epithelial cells have been noted to pass into the underlying tissues and assume the appearance of fibroblasts. While we admit that the process occurs, we would urge that the fate of these cells has still to be determined. As shown by Schridde, the mere fact that a cell of the plasma-cell type comes to simulate in size and shape the connective-tissue corpuscle of connective tissue does not make it into a connective-tissue corpuscle. It has still to be shown that the fibroblast-like cells of epithelial origin function as fibroblasts and become fully formed connective-tissue cells. While it is true that in the past too great a stress has been laid upon the tendency of epiblast and hypoblast to form what we have termed lining membrane or lepidic tissues, and too little attention to the fact that each of the primary cell layers can form tissues of both orders (matters to which we shall revert in discussing neoplasia1), epiblastic connective tissue (e. q., the neuroglia) exhibits constant differences from the mesoblastic connective tissue, and for the present we must continue to lay down that while there may be conversion of one epiblastic or hypoblastic tissue into another epiblastic or hypoblastic tissue, and of one mesoblastic form of cell into another mesoblastic form, this conversion is of a limited extent; metaplasia of mesoblastic tissue into epiblastic or hypoblastic and vice versa does not occur.

Lastly, we recognize the difficulty that presents itself in the conception of fully formed cells of one order becoming directly converted into cells of a different type. But this, I believe, never happens. As I pointed out some years ago,² "metaplasia is never direct, but is only

² On Growth and Overgrowth and on the Relationship between Cell Differentiation and Proliferative Capacity, Jacobi Festschrift, 1900, and Med. Chron., Manchester, June, 1900.

¹ See page 700.

THE MODE OF CELL CONVERSION

brought about by preliminary reversion to a more embryonic type, or where mother cells are present, by the modified development of cells derived from the mother cells, the influence of environment altering the character of those cells during the period of growth."

The nearest approach to such direct conversion is to be seen in the conversion of cartilage into bone cells. The former cells are histologically so simple in type that it is difficult to recognize any change in them. But functional change there must be, since the matrix which they govern exhibits a preliminary retrogressive modification.

More recently Schridde has emphasized the same general principle. Thus, in the corporeal, as in the spiritual world, to undergo "conversion" it is seen to be necessary to "become like little children," if not to be "born again," or, as Orth expresses it, "In the majority of cases at least, it is not mature cells that change themselves, but it is youthful, not yet completely differentiated cells that take on the new character."¹

¹He similarly quotes bony metaplasia as presenting difficulties in formulating an absolute rule.

CHAPTER XV.

THE NEOPLASMS: TERATOMAS AND TERATOBLASTOMAS.

WHATEVER branch of biological science we make the subject of our study, the more deeply we enter into it, the more do we realize that classification, which is the goal and outcome of our knowledge-so far as it concerns the knowledge in itself—is not the sorting of data into sharply defined departments; such departments do not exist. Rather, it is the arrangement of our data in progressive order in such a way that we most satisfactorily gain a comprehension of their relationships and the place they occupy in one harmonious scheme. The classes of living objects and of vital phenomena are not distinct; classes as such have no absolute existence; they pass imperceptibly one into the other by many transitional forms. While this is so, classification is nevertheless necessary. To measure the grade we have to set up posts at convenient intervals; to grasp the progression of forms we have to select types here and there at suitable points, group the forms most nearly allied around these, and so constitute classes. Nowhere do these considerations gain a better illustration than in this study of the different forms of neoplasms.

Terminology.-We speak familiarly, and rightly, of any unusual swelling recognizable in any part of the organism as a tumor, for "swelling" is the root-meaning of that term. Under this heading we may include (1) examples of dislocation of parts, (2) abnormal collections of fluid or gas, whether sharply encapsulated, as in cysts, or, though localized, more diffuse, as in inflammatory and hemorrhagic conditions, and along with these, (3) actual tissue growths, whether (a) physiological (as in the case of the pregnant uterus), or (b) hypertrophic, or (c) due to localized abnormal growths of part of a tissue or organ, or within an organ. Save for gross descriptive purposes, unless preceded by a qualifying adjective, the term tumor has no value. If, therefore, we wish to classify and distinguish from other forms of tumor a series of solid overgrowths which are included under some of the conditions hitherto studied, namely, the class which Thoma has, we think, appropriately termed the autonomous tumors (i. e., those which are, or which possess, a law unto themselves), we have to select some more definite term, and for some years it has been usual to speak of neoplasms (literally, newgrowths) and of *neoplasia*, the process of new-growth. But, as Klebs justly points out, these terms are applicable also to conditions of regeneration, hyperplasia, and all forms of new or renewed growth of tissues. Conventionally, however, when we speak of neoplasms we only take into consideration the autonomous tumors we are now about to discuss. Where accuracy of description is required we distinguish two distinct

orders of these neoplasms proper, the *teratomas* and the *blastomas*, and, as 1^t have pointed out, it is serviceable to recognize a third intermediate order, the *teratoblastomas*. What we understand by these terms will be made clear in the following pages.

The termination *oma* following the Greek root for tissue of one or other nature, or even of some descriptive adjective, conventionally indicates an overgrowth of the types about to be considered.

Here, again, owing to the evolution of our science, exceptions are to be noted. Before it became possible to make clear distinctions between the different forms of neoplasia, the termination oma was employed indifferently to indicate swellings of any order. We thus still refer to subcutaneous collections of blood as *hematomas*,² and use it when referring to specific inflammatory overgrowths—*luberculoma*, *spphiloma*, *condyloma*, etc. These conditions are, however, not autonomous, and we no longer include them under the neoplasms proper.

It is this autonomy, this growth independent of function and of either present or future needs of the organism in which they occur and from which they gain their nourishment, independent also of obvious stimulation from without, that distinguishes the neoplasms proper from all other forms of tissue growth. And it is this also that renders it difficult to define them in terms applicable to other vital processes. In seeking for such a definition the natural course to follow would be to consider processes apparently most nearly allied, and carefully to analyze the points of likeness and of difference. Now such processes exist; there are overgrowths the result of inflammation, and others of the nature of congenital hypertrophies, in which it is almost if not wholly impossible to state where the division comes between inflammatory disturbances or hypertrophy on the one hand, and blastomatosis on the other. But these we shall consider later, when discussing the blastomas, for it is in connection with that order of tumors that these difficulties arise. In order to present as clear a picture as possible, we shall at first consider autonomous neoplasia in general, define that, and then take into consideration the different orders and their relationships.

Definition.—Too often have theories as to the causation of these autonomous neoplasms entered into the definitions. Thus, Cohnheim defined them as "circumscribed atypical productions of tissue from a matrix of superabundant or erratic deposit of embryonic elements." Here we have introduced the untenable theory that all autonomous neoplasms arise from embryonic tissue which has remained latent. We are still uncertain as to the causation of these growths, and so etiology must not enter into our definition. Thus, Ziegler's definition is more

² When words so formed were new and foreign, it was correct to employ the Greek form of plural, and to speak of sarcomata, osteomata, etc. But these words have now become so familiar a part of every-day language that they may be regarded as naturalized and given the ordinary English plural. We shall use the two forms of plural indifferently.

¹ Adami, Montreal Med. Jour., July, 1908.

satisfactory: "A tumor is a new formation of tissue possessing an atypical structure, not exercising any function of service to the body, and presenting no typical limit of growth." 'The use and limitations of the term "atypical structure" require here a little explanation, add to which, the pure teratomas to be presently described do present a limit of growth; and so we prefer C. P. White's statement that "a tumor proper is a mass of cells, tissues, or organs resembling those normally present, but arranged atypically. It grows at the expense of the organism without at the same time subserving any useful function." Von Rindfleisch characterizes them as a "localized degenerative excess of growth;" i. e., the very excess of growth is regarded as in itself a degeneration; Birch-Hirschfeld, as originating spontaneously, becoming separate from the physiological tissues in their physiological and functional relationships, as developing from the cells of the body, and possessing progressive growth; Ribbert, as "selfconfined, dependent upon the organism for their nourishment, but otherwise largely, if not quite independent, corresponding more or less but never absolutely with the tissues of the natural body, and presenting no definite limit to their growth." Lubarsch's definition is closely allied: "Under tumor proper we have to understand those growths of apparently independent origin which histologically correspond in structure more or less completely with the matrix from which they originate, but in form are atypical; which further, in spite of their organic connection with that matrix, and in subjection apparently to laws of their own, pursue an independent existence which is not, or only exceptionally, of advantage to the organism as a whole."

How next can we classify the growths possessing these characters? As indicated by the definition we have selected, these neoplasms are composed of cells, tissues, or organs resembling those normally present in the body; in fact, we cannot but conclude that they have a like origin. It would seem to follow, therefore, that classification is possible according to the type of cellular tissue present, just as we are able to classify the cells and tissues of the normal organism. But before proceeding to do this, it is well to take into account the variation noted in the definition (White's), namely, that some of the tumors are composed of cells of one particular type, others show a tendency toward arrangement of those cells in definite order with intervening stroma, such as we can see in normal tissues; a third group shows cells derived evidently from more than one type of tissue-the mixed tumors; a fourth shows even greater variation in the type of cells with tendency to the development and presence not merely of irregular cell collections, but of such fully formed organs as brain, teeth, masses of bone, skin, sebaceous and other glands.

We will consider these last first.

TERATOMAS.

All monstrosities are *terata*, and such terata we have discussed in an earlier period of this work, pointing out the successive grades, from the dichorial and monochorial twins, through the symmetrical double

TERATOMAS

monster to the asymmetrical parasitic monster, such as we see in the foetal inclusion. The study of that series has demonstrated that we there dealt with, in the simplest cases (the dichorial twins), the development of two ova side by side in the uterus, in the next with the formation of two separate (twin) individuals from a common ovum, until, reaching the feetal inclusion, we find that, of two individuals so developing from a common ovum, one, the feebler, became during early embryonic development infolded into the other and that it gains its blood supply from that other stronger focus, becoming ingrafted into it. Now, such a foetal inclusion, an imperfect grafted individual, we may regard as our type of teratoma. It is not an independent individual; it is incomplete; it is nourished from its host; but it has begun existence as a separate individual, its tissues have developed from an independent primitive streak. Even if, as in our earlier chapter we pointed out, both parasite and host originated primarily from a single ovum by a single act of fertilization, nevertheless, at an early period, that single ovum came to exhibit two independent centres of growth, and it is the autonomous growth of one of these that has given rise to the mass of tissues constituting the parasite.

If, then, we take this as our type, we may define the teratoma as an autonomous growth, the product of the continued development within one individual of another individual of the same species. We place an emphasis upon the "continued development" in order to exclude the normal fortus, which possesses only a temporary development of this order, and then through its placenta, which penetrates into the maternal tissues.

There are several different types of tumor which fulfil this definition. but before describing them it will be well for the purposes of orderly classification if we consider what cells in the organism in its different stages are capable of giving origin to all the orders of cells which constitute the individual. Our first inclination is to lay down that the fertilized ovum alone can do this; a little consideration shows that the potentiality is more extensive. Every *totipotent cell*, to employ the terminology of the enbryologist Barfurth, must be regarded as capable of giving origin to an individual, every cell, that is, possessing the power of giving origin to cells of every order.

Of such totipotent cells, in addition to the fertilized ovum, we recognize the following:

I. The primordial blastomeres. We know from abundant experiments that these, even among the vertebrates (the frog, Roux, Morgan), can be broken apart and each give origin to a complete dwarfed individual.

II. The primitive germinal area cells.

III. The "growing point" cells of the germinal area (p. 238), which give origin to the successive mother cells for the various fissues. The powers of these cells, it is true, is more restricted; they are not, like the blastomeres, yolk-containing, and can only give origin to the embryo so long as they are in connection with the body of the ovum.

1 in rom uble

he

·h,

th;

188

em

of

the

elf-

ely

in

ep-

200-

the

lore

IV. The germinal blastomeres. These from the very earliest period of segmentation of the ovum appear to be set apart, becoming eventually lodged in the generative glands, and there give origin to the eventual ova or spermatozoa. There is a succession of generations from the primordial germinal blastomere down to the mother cells of the ova and spermatoza, all retaining and conveying onward totipotential characters. Here we do not include the ova and spermatozoa as such, the results of a reduction process affecting the mother germ cells (see p. 150).

V. The fertilized ovum.

1. As already noted (p. 228), separation of the primordial blastomeres can only be regarded as giving rise to dichorial twins, not to teratomas.

2. Monochorial twins would seem to originate at a rather later date, and then not so much from a single primitive blastomere (although this cannot be wholly excluded) as from an early division or dichotomy of the cells set apart to form the germinal area, cells which, it is true, are totipotential. From this same order of cells we must regard the individuals as originating which become the eventual *fatal inclusions*.

3. Excess production of growing point cells affords the most satisfactory explanation, as already noted (p. 239), of those remarkable forms of teratoma, *Epignathus* (at the superior pole) and *Congenital Sacral Teratoma* (at the inferior). They may be regarded as examples of polar or serial deduplication.

4. The germinal blastomeres. (a) It has been noted by several embryologists who have followed the germinal blastomeres from the earliest stages of development, that these undergoing multiplication do not necessarily, in every individual, all find their way into the ovary or testis. Certain of them may come to be included in other organs and regions—in the cranium, the gill clefts, thoracic cavity, etc. To the later development of these misplaced germinal blastomeres have been ascribed the *embryomas* showing themselves in different regions. (b) But if this be so, a similar origin from germinal blastomeres, or more clearly from the mother cells capable of giving origin to ova or spermatozoa, most satisfactorily explains the much more common development in the ovaries and testes of complicated tumors containing tissues derived from all three cell layers (epiblast, mesoblast, and hypoblast), the *ovarian* and *testicular "dermoids*").

These, then, we regard as the teratomas.

Elsewhere¹ we have proposed that they should be classified into:

1. Twin teratomas (when host and parasite are of equal age).

2. Filial teratomas (in which the teratoma is the product of one of the germ cells of the host); and have subdivided these last into:

(a) Parthenogenetic, from germ cells multiplying without previous fertilization, and

(b) Gamogenetic, the product of growth of a fertilized germ cell.

Further consideration has made us a little doubtful as to the expedi-

¹ The Classification of Tumors, Jour. of Path., 4:1902:233.

CLASSIFICATION: FETAL INCLUSIONS

ency of this classification; while we believe that the conception of twin and filial teratomas is a useful one, we would replace the term germ cell by totipotent cell, to include growths of the epignathus type. So, also, we have noted that using the term "parthenogenetic," it has been assumed that what we meant was that teratomas of this order (ovarian and testicular embryomas) originate from the mature ova or spermatozoa. This is not our belief. Lastly, while formerly we were inclined to classify placental moles and that remarkable form of growth, the chorio-epithelioma malignum, both derived from cells of the fertilized ovum, among the teratomas proper—a position for which much may be said—we are now of the opinion that as such growths do not represent the individual, but only the aberrant growth of one set of cells belonging to the individual, it is better to discuss them as a class apart—of *Teratogenous blastomas*.

Here the classification of Wilms¹ deserves attention:

1. Developments of two anlagen on one germinal vesicle with partial fusion: *Double monsters*.

(a) With equal growth: Duplicia symmetros.

va

a,

rea

te.

re

ble

he

ter

32.

(b) With early arrest (lagging behind) of growth of one: Duplicia asymmetros,

(c) With parasitic inclusion: Fatal inclusion.

2. Developments from one anlage on the germinal vesicle producing excess blastomeres which become included in the growing individual.

(a) The included blastomere undergoes development at a very early age, the growth being relatively elaborate: *inclusions recognizable at birth*.

(b) The included blastomere lies latent within the organism for some period, and starts active growth only at a later period (often in the fully developed individual) as a result of some altered conditions (Gelegenheitsursache): most cases of abdominal inclusions, embryomas of the genital glands, and embryoid (mixed) tumors.²

Accepting for the time being the view that the tumors of class 2 are derived from aberrant blastomeres, it is difficult to suggest a term which will succinctly indicate them. Wilms first labelled them embryomas, in the belief that they were derived, parthenogenetically, from fully formed germ cells, and were, in short, of the same rank as the embryo. Similarly we have classed them as filial teratomas. If for "blastomeres" we employ, as suggested, the term "totipotent cells," then Wilms' classification and that here adopted become practically identical.

Fœtal Inclusions,—The inclusions may be incomplete or incomplete and projecting (Fig. 94). Our conception of such as a weaker and smaller embryo carried into the body of the more fully developed embryo, during the process of closure of the great anterior fissure,³ makes it

¹Wilms, Die Mischgeschwulste, Leipzig, Georgi, 1899, 250. This is the locus classicus for the forms here under consideration, and is a masterpiece of clarity. Si sie omnest.

 2 This, as Wilms states, does not pretend to be a full classification of the double nonsters.

³ The existence and early development of the amnion surrounding the embryo on its lateral and dorsai aspects prevents such inclusion anywhere save at this fissure.

essential that those embryomas only can be regarded as fœtal inclusions which are in relationship to that fissure, *i. e.*, whose situation is (1) median, (2) ventral, (3) thoracic or abdominal. We cannot, that is, conceive inclusion as occurring in any other region. Teratomas, for example, having a retroperitoneal and more dorsal abdominal position, posterior mediastinal, or cranial cannot be explained on this supposition.

Epignathus and Congenital Sacral Teratomas.—For a description of the mode of origin of these forms see p. 238.

Schwalbe distinguishes four groups:

1. There is inserted into the roof of the mouth of the one factus the umbilical cord of a second, which can be more or less well developed (very rare and cases poorly described).

In Baart de la Faille's case there was a large epignathus of the type of group 3, attached to which were the umbilical cords of two acephalic acardiae fœtuses—indications, that is, of three terata. This case in itself makes it impossible to regard the monstrosities of this type as of bigeminal origin with inclusion, and would seem to favor the theory of liberation of totipotential cells from the superior growing point. We are not, however, prepared to make definite pronouncement regarding the mode of origin of this very remarkable and rare order of monstrosity.

2. A mass of tissue presenting definitely formed organs (lower extremities, sexual organs, etc.) projects out of the mouth of the host.

3. A mass of tissue having its root at the base of the skull in the midpharyngeal region projects out of the mouth of the host, but this presents no definite organs, only an irregular mass of tissues—skin, connective tissue, cysts formed by epithelium both of epiblastic and hypoblastic type, cartilage, etc. This is the common form.

4. A larger or smaller tumor of the palate or oral cavity composed of a mixture of tissues of simpler type than the above (not from all three cell layers).

The indications are that not all of these are of median origin and attachment, and that they come under the teratoblastomas to be described later, rather than the epignathi proper. We must recall the difficulty in classification already referred to (p. 650). It is evident that in the same growing point region at a later period aberrant multipotential, rather than totipotential cells may be produced, giving rise to a less complicated form of tumor.

Sporadic Embryomas.—As above indicated, we occasionally encounter teratomatous growths of the same type as, but not conforming in position to, the preceding forms, bearing no relationship to the fissures or the poles of the body, nor again to the generative glands, and these we can only, *per exclusionem*, regard as due to the independent development of aberrant germinal blastomeres, blastomeres which, instead of finding their way into ovary or testis, have become displaced and arrested in one or other region of the developing organism. Such blastomeres may take on growth and the production of a mass of various tissues (epiblastic, mesoblastic, and hypoblastic) either during feetal life or, lying latent, only during postnatal existence. In this way we

OVARIAN TERATOMAS

best explain the organoid tumors of the deep thoracic and abdominal regions, of the cranium (independent of the sella turcica), neck (often in close association with the thyroid), anterior mediastinum (often associated with and apparently originating in the thymus), etc.

The careful study of these various teratomas made during the last few years has shown conclusively that by far the greater number contain tissues derived from all three germinal layers, although characteristically these do not present themselves in the proportions seen in the normal individual; one or other (notably nervous tissue) may be present in marked excess, and there is wanting in a very striking manner the orderly relationship of these tissues one to another: they are "jumbled" together. With this two orders of tumor may be distinguished, which, employing the terminology employed for the blastomas (p. 668), we may speak of as the typical and the atypical. Among the sporadic teratomas the former is the rarer class; in it the tissues are of the adult, fully formed type, and growth is characteristically limited. The indications are that development has been pari passu with that of the host. The latter is the more frequent class; while they may show themselves in early life, more often they are just noted at or after puberty, and they grow with relative rapidity, and, what is more, tend to afford metastatic newgrowths in distant organs, also composed of a mixture of cell elements. The cells are of imperfectly differentiated type, hence they are known as *embryonal* teratomas. We have here orders comparable with the benign and malignant blastomas to be presently studied. In other cases one or other of the component tissues of a typical teratoma may take on excessive growth, affording metastases of the one cell type, This formation of a "tumor in tumore" we shall take up later. (See

Ovarian Teratomas.-But quite the commonest seat of teratomas is in the ovary. Here similarly we encounter two forms: (1) The largecysted teratoma, or ovarian dermoid, and (2) the solid or small-cysted teratoma. The former is much the commoner. It presents itself as a cyst occupying the situation of an ovary, which may attain the size of an orange, a child's head, or larger; the contents are characteristically fatty debris with long hairs. It is lined with squamous epithelium provided with sebaceous and sudoriparous glands, and often bony masses are to be detected in the walls. Examination generally reveals what Rokitansky described as the "insular protuberance," a region which we now recognize as the representative of the head. From it arises the tuft of long hairs, and around a depression to one side there projects one or several teeth, frequently embedded in an amorphous bone which may be taken to represent a jaw. Serial sections demonstrate that into this depression opens a tube, which proximally has the characters of a trachea, distally takes on the structure of the digestive canal with muscular sheaths. Sections through the protuberance exhibit the presence of five layers: (1) The skin and subcutis; (2) central nervous system elements (glial cells, neurons, occasionally pigment cells) and the meninges; (3) hypoblastic tissues (tracheobronchial, gastro-intestinal);

ions (1) conple, erior

tion

the

pped type nalic tself s of y of We the '. ower nost, nidents tive ustic

d of hree ach-

ibed y in ame ther ated

ures hese loptead and such ious


ig in

tetal we

(4) sympathetic nerve cells and their processes; and (5) the remains of the ovarian cortex.

While recognized most commonly in adult life, operation or postmortem may reveal the presence of the form of tumor in the young child, and it is to be noted that the cells of the various tissues are as fully formed as are those of the host; we deal with a *typical* teratoma. Very rarely—only four cases are on record (Axel Key, Repin, Askanazy, and Shattock)—some proportion and relationship of parts is preserved, with development of extremities and genital parts, so that a definite fectus may be inferred.

The solid ovarian teratoma is distinctly uncommon. It has all the characters of the atypical sporadic teratoma already described.



Interior view of an ovarian teratoma ("dermoid cyst"), showing Rokitansky's island bearing c, hairs with d, teeth surrounding. (Schwalbe.)

Testicular Teratomas.—Here the relative frequency is reversed: the solid small-cysted form is the usual type, whereas examples of a large solitary epithelial cyst or dermoid are few in number. To this solid form would seem to belong the greater number of the malignant mixed to be regarded as chondrosarcoma or sarcoma carcinomatodes. Study of all parts of such a tumor in general reveals elements derived from all three germ layers, although in contradistinction to the ovarian growths the ectodermal elements are in a minority. This, with the one exception of the chorio-epithelium, to be referred to later (p. 666).

THEORIES OF ORIGIN

THEORIES OF ORIGIN OF OVARIAN AND TESTICULAR TERATOMAS.

Needless to say there have been very numerous theories regarding the origin of these tumors: (1) That they are due to inclusions in the ovary or testis of portions of all three layers during the course of growth. Against this the frequent bilateral nature of the growth was seen to militate. (2) That they are the product of fertilized polar bodies (Marchand). But no less than five separate embryomas have been noted in the one ovary (Wilms), and the ovum in preparation for fertilization only casts out three polar bodies. (3) That they are due to parthenogenesis, a mature germ cell, ovum, or spermatozoon, under certain unknown conditions, taking upon itself to start growing and segmenting without throwing off the polar bodies, and without fertilization. This very lack of normal stimulus, and the abnormal site in which growths occur, is held to explain the aberrant nature of that growth. Parthenogenesis, we know, is common among the lower forms of life, and even the virgin chick has been known to lay eggs which may exhibit an apology for a germinal area, and rarely, later stages of aberrant development. The grounds taken by Bonnet,¹ denving the possibility of such parthenogenetic development, do not appear to us absolutely convincing. What, however, is strongly against such a theory is that experimentally we know that for a cell to proliferate, the nucleus must be surrounded by a certain not inconsiderable quantity of cytoplasm. The spermatozoon, as such, exhibits much less than what we regard as that minimum, and nothing of the nature of yolk to support the growth during the early stages of segmentation. It is against all we know regarding the conditions favoring cell multiplication that the adult spermatozoon should give rise to an embryo, or an embryoma. And if the testicular embryoma cannot thus be a parthenogenetic development, the identical ovarian growth is unlikely to be of that nature. (4) That they are due to the aberrant development of cells of the theca interna of the Graafian follicle. For this suggestion absolutely no evidence of any weight has been advanced.

There remain, so far as we can see, only two other possibilities: (5) That they are derived not from ova or spermatozoa matured for fertilization, which have undergone the reduction process, but from oocytes or spermatocytes, or even the forerunners of the same; this we may term the modified parthenogenetic or germ cell theory; or (6) that they spring from dislocated blastomeres. This last is the theory at present in vogue. It supposes that in the early embryonic stage certain blastomeres capable of producing all three cell layers become dislocated with some arrest of their proliferative activity, and carried into the anlage of the future ovary or testis, when sooner or later they may take on growth.

Bonnet, who first propounded this theory, based it upon the facts to which we have already referred, that in many of the lower animals, the early segmenting ovum can have its cells shaken or otherwise broken

¹ Ergebnisse der Anatomie und Entwickelungsgesch., 9:1899.

ins of

postoung tre as toma. nazy, rved,

ll the

id 1: the

large solid nixed used Study m all owths xcep-

apart, when each is capable of developing into a complete if small individual, and that the segmentation and mitosis in mammalian eggs is known to be most irregular.

We confess that this theory does not appeal to us. The very fact which Wilms, who supports it, urges, in another connection, that he has found five separate embryomas in one ovary, would suggest altogether too extensive a dislocation and carriage of the not too numerous blastomeres into one particular locality to be within the bounds of probability. We, ourselves, must accept the germ cell theory, and this, for several considerations. We know that these germ cells from the first are dedicated to eventual reproduce all the tissues seen in the parent. We know, as pointed out by Beard and other zoologists, that there is an actual dislocation of many of these cells in the process of development. The total number of germinal blastomeres (already specialized and recognizable at a significantly early stage in the segmenting ovum) does not find its way into the ovum or testis. Here, in short, we have the cells which will fulfil all the needs of the case.

These germinal blastomeres, as distinct from Bonnet's primitive blastomeres of any order, we know may be carried to various parts of the developing organism. This carriage and deposition afford an adequate explanation for the development of three-layered teratomas in the abdominal and thoracic cavities and in other regions, teratomas of the same type as those in the ovary and testis. The sporadic teratomas outside the ovary and testis obtain their simplest explanation in an origin from such aberrant germinal blastomeres. Within these organs it does not seem to be necessary to postulate, however, that the embryomas arise from germinal blastomere that have lain intact *ab initio*. The function of the germinal blastomere is to give rise to cells having like properties, and thus we note that it is probable that any cell along the direct line between the primitive germinal blastomere and the mature and *reduced* ovum or spermatozoon (these last being excepted) may potentially give origin to an embryoma.

What is the cause of such cell taking on aberrant growth will be discussed when we come to deal with the theories of neoplasia (p. 835).¹

TERATOBLASTOMAS.


This is the proper place to consider another order of growths, growths which are not true embryomas, because all three germinal layers are not represented by the tissues found present. Under this heading we include the most striking examples of what are known as the "mixed tumors."

¹The most masterly survey of the present status of our knowledge regarding teratomas is afforded by Askanazy (Verhandl. Deutsch. Pathol. Gesellschaft, 1907, 11;1908;39). With his conclusions I find myself largely in agreement. The succeeding article by Borst gives valuable bibliographical references.

TERATOBLASTOMAS

Certain of these "Mischgeschwülste" are of another type (see p. 705), but the most characteristic examples come under this head.

1. Renal Teratoblastomas.—The type example of this form of growth is found in certain remarkable tumors of the kidney present, it may be, at birth, or developing during infancy or early childhood. We deal with relatively large localized growths of the body or the pelvis of this organ; soft and sarcoma-like, with great tendency to internal hemorrhages and necrosis. These on section show a more or less spindle-celled, sarcoma-like matrix, but in them we encounter epithelial elements of the nature of gland tubules recalling, but different from, the typical renal tubules, fat and cartilage cells, plain muscle fibres, striated muscle fibres, fibrous and elastic tissue, all in no apparent order. The tumors have been recorded under very different names—adenosarcoma, carcinoma sarcomatodes, rhabdomyoma, spindle-celled sarcoma, according to the tissue most in evidence.



Section of a "mixed tumor" of the kidney, showing gland tubules with surrounding sarcoma-like cells of the plain muscle type, fat cells, etc. (Ribbert.)

They are to be explained as follows (Wilms): The mesoderm of the kidney, or primitive kidney ("Urniere," or Wolffian body) region, gives rise, first, to the *myotome* (primitive segment), later gives off the *nephrotome*, or matrix for the future kidney tissues. From the myotome is given off also the *selerotome*, whence develop the mesenchymatous elements of this region of the body (striated muscle, vertebre, etc.). A tumor containing all these tissues—namely, kidney tissue proper, striated muscle, and different connective tissues—can only have originated from cells themselves capable of originating all these forms. Or otherwise to explain this particular form we are forced to the conclusion that in the course of development of this region certain of the primitive mesoderm cells, potentially capable of originating both sclerotome and nephrotome, are carried in a latent condition into the area of the future kidney, and growing later, give rise to all the tissues in question. Such is a onelayered embryoma, *i. e.*, all the contained tissues are of mesoblastic origin.

Muus, from Marchand's laboratory at Marburg, in one such tumor, from a child, aged eighteen months, found definite epidermal inclusions

LS

ggs is fact

e has gether lastobility. everal dedient of know, actual The recoges not e cells

blastoof the adenas in tomas teranation these act *ab* o cells ny cell nd the repted)

be dis-835).¹

rowths are not include mors."

ing teraft, 1907, The suc-

exhibiting strata mucosum et granulosum, and stratum corneum. For the explanation of such we must pass back to a still earlier date, to the inclusion of an epiblastic cell over the future kidney region, a cell capable of giving rise to both epiblastic and mesoblastic structures, and must regard this as a two-layered or diphyllic embryoma.⁴

2. Mixed Tumors of the Parotid.—The mixed tumors of the parotid are apt to be even more complicated, for here, not rarely but frequently, extensive epithelial overgrowth is present, both squamous epithelium (Hinsburg points out that many of the larger scattered cells regarded as endothelial, or even connective-tissue cells, are of epithelial nature and origin), cubical, and cylindrical epithelium. Adult goblet cells, elastic fibrils, cartilage, mucoid interstitial tissue, osteoid tissue and even true bone and spindle-celled, actively growing connective tissue, may all be present in these tumors.

3. **Submaxillary Gland.**—More rarely similar mixed growths are found in connection with the *submaxillary gland*.

The cells which originate such tumors we must ascribe to a developmental period when they were capable of giving rise to both squamous epithelium (of the mouth) and parotid glandular tissue, that is, to a period before the mouth proper had become differentiated. So also for the mesenchymatous elements, the cells of origin must have been capable of producing connective tissue, cartilage, and bone. There is greater power of transformation (metaplasia) between connective-tissue elements; nevertheless, proceeding on the same lines as we followed in connection with the kidney tumors, we must conclude that cells of the primitive epiblast have become displaced at a period when they had not as yet undergone differentiation and had thrown off the mesenchymatous elements, so that, passing into the parotid area, they there eventually give origin to all these forms of tissue.

4. Of the Vagina (in children).—Such exhibit round and spindlecelled sarcoma elements and striated muscle fibres; due, it would seem, to misplaced mesoderm of the inferior region of the body, misplaced during the growth of the Wolffian duct, which in the female, as Gärtner's duct, is present in early fortal life opening into the vaginal area.

5. Of the Cervix Uteri.—Similar sarcomatous elements with plain and striped muscle, and sometimes cartilage, may be found in mixed tumors of the cervix. These occur later in life. Wilms suggests for them also displacement along the course of Gärtner's duct, traces of which may also be found in the cervix.

6. **Mammary Glands.**—Cystofibrosarcomatous growths with, in parts, squamous epithelial tissues present in this region may be ascribed a like origin.

7. Lacrimal Glands, Cheeks, and Gums.—Here also are rarely found mixed tumors allied to the parotid and submaxillary tumors.

In conclusion, we would repeat the axiom laid down by Wilms, that

¹While most of the primitive mesoblast cells are derived from the hypoblast, some are given off from the epiblast.

TERATOGENOUS BLASTOMAS

"the mixed tumors always correspond wholly in structure with the normal processes of differentiation, occurring in the particular region of the body in which they originate." This, it may be repeated, is not the only, although it is the most characteristic, form of "mixed tumor." We shall have occasion to refer to other types.

TERATOGENOUS BLASTOMAS.

87.53

Placental Moles and Chorio-epithelioma Malignum.—The developing ovum consists of two parts, the fœtus and its membranes, including the fœtal placenta—an organ developed primarily from the chorionic villi, which, at first wholly epiblastic, come, with the development of the allantois, to gain a vascular and mesodermal core. As first demonstrated by Peters,¹ in his study of a singularly early human embryo, and since repeatedly confirmed, long before the development of the placenta the outer cell layers of the chorion show active erosive properties, for already his ovum was practically buried in the uterine mucosa. The outer layer of the villi has, in fact, intensely active phagocytic properties, and as the placenta forms, by their rapid growth and erosive powers, the villi penetrate through the mucous membrane into the underlying venous sinuses, where, by selective absorption, they gain nourishment for themselves and the foctal organs.

Under normal conditions, with the maturation of the foctus, these vill withdraw; their outer layer of fused cells, the syncytium, has long previously undergone extensive atrophy, so that only here and there small remnants are to be detected; the layer beneath, Langhans' layer, also undergoes degeneration, and so the attachment between foctus and mother is loosened in preparation for birth. But this does not always happen completely. More particularly, in cases of abortion (in which the foctus is discharged before these placental changes are complete), and in some cases of blighted ovum in which there is no fectus which by its metabolism must regulate the placental changes, the vill, or some of them, may persist in intimate contact with the maternal tissues and maternal blood, and may continue to grow after the normal period of gestation has been attained. According to the nature and the rate of this growth, so do we obtain two orders of tumor—the placental mole and the chorio-epithelioma malignum.

The Placental Mole.—We occasionally encounter cases in which the placenta and membranes are the sole product of conception, the fectus either being absent or dying and undergoing absorption at a very early age. In these cases the placenta, growing within the uterine cavity, tends to become converted into an irregular flashy mass, the fleshy mole, often infiltrated and surrounded by much recent and coagulated blood (hemorrhagic mole), while secondary to the hemorrhage and

¹ Die Einbettung des menschlichen Eies, Leipsic and Vienna (Deuticke), 1899.

arrest of blood supply, putrefaction may ensue (*putrefactive mole*), with or without eventual infection of the maternal organism.

• More particularly in cases of premature birth the portion or portions of the placenta remaining attached may exhibit a series of remarkable modifications of the chorionic villi. Continuing to be nourished by the maternal blood and to absorb fluid, they may hypertrophy and become distended by an cedematous mucoid infiltration so as to form a relatively huge mass of series of small, clear, grape-like vesicles of varying size, distending the uterus to even a greater extent than does a full-term fortus. Such is the hydatid mole. In the majority of cases there is no sign of an associated fectus; in some there is the history of abortion; in others a



A small portion of an bydatid mole; natural size.

dead foctus at some period of arrested growth has been found. In a very few cases the hydatid formation has affected one portion only of the placenta, and the foctus nourished by the remaining portion has been born alive. As the hydatid mole develops, there may be frequent hemorrhages until finally it is extruded,

Careful examination of an otherwise healthy placenta occasionally shows here and there a rare cyst in its substance. Such cysts of the foctal placenta are caused by a similar ocdema of portions of an individual villus in which circulation has been arrested.

Chorioepithelioma.—In the above cases the growth remains within its normal limits, but this is not always so. There are cases in which such a mole, continuing to grow, may also fill the maternal uterine sinuses with polypoid masses (destructive placental polypi), and thus we have transition to


a yet more remarkable condition, recognized only within the last few years, the fatal form of new-growth which after several changes of name—deciduoma, syncytioma—is now usually termed "chorioepithelioma malignum."¹¹ This occurs within the uterine wall, but may, indeed, first show itself quite outside the uterus, in the vaginal wall, etc. The outer surface of the villi, as we have said, consists of the layers of feetal epiderm, the most external, or *syncytium*, formed of cells which stain deeply, showing, as the name implies, a fusion of the cell bodies, so that they appear as large, multinucleated, protoplasmic masses covering the surface of the villi, the more internal Langhans' cells

⁴ For fuller description, see Marchand, Monatssehr. f. Geb. u. Gyn., 1:1895:419 and 513, and Ztschr. f. Geb. u. Gyn., 39:1898;173.

CHORIOEPITHELIOMA MALIGNUM

being of fair size, but individual, not fused, and not staining so deeply. It is the former that possess the intense erosive and phagocytic properties whereby the villi penetrate and come to lie within the maternal blood sinuses of the uterus. As Schmorl pointed out in his study of eclampsia,' certain of those cells are apt to be swept away by the blood current and deposited in the capillaries of the lungs and other organs. It may well be that this is a not infrequent occurrence in the course of pregnancy, although usually these cells become destroyed by the counteraction of the endothelium of the capillaries in which they become lodged. In cases of abortion, more particularly where the normal course of placental development and involution is interrupted, either (a) within the uterine sinuses, these cells, both the syncytial and those of Langhans, take on an

FIG. 194



Chorio-epithelioma growing within uterus: V, wall of uterine sinus; Syn., multinucleate cells of syncytial type; L.c., cells of Langhans' type. (Teacher.)

abnormal proliferation, or (b) this proliferation shows itself, not locally, but in some of the cells transplanted to another region, and that not necessarily immediately, but it may be after a considerable period.

There thus develops a tumor, that is:

1. Cellular, formed entirely of large cells of embryonic type.

2. Entirely within the vessels.

 Provided with no vessels of its own, and exhibiting no sign of structure save at times an obscure development into finger-like processes (although this would seem to be largely due to its mode of growth along the maternal vessels).

4. Unprovided with a capsule or any definite limitation; and so,

5. Extensively infiltrating the uterus or other organ involved.

 Peculiarly liable to form secondary growths by detachment of some of its cells, and proliferation of the same in vessels at a distance.


¹ Verhandl, Deutsch, Pathol, Gesellsch., 8:1902:39.

 Liable to induce hemorrhages by erosion of the walls of the vessel it distends.

8. Rapidly fatal.

Most often in sections of such tumors the characteristic deeply staining, multinucleated masses of syncytium are to be recognized in the growth. Occasionally cases are met with in which these are absent. It may be that the cells forming tumors in these cases date their origin from a relatively late period of pregnancy.

The angry debate waged for some years as to the nature of these tumors was based upon the denial by many leading pathologists of the feetal origin of these cells, and this notwithstanding that it had already been well established by Hubrecht and other embryologists that the



Cells of a chorio-epithelioma malignum, higher magnification: a, syncytial cell mass; b, cells of Langhans' type; c, broken-down erythrocytes. (Von Franque.)

syncytium was of foctal epiblastic development. It was held to be decidual, *i. e.*, of maternal origin. If we are not mistaken, Young,¹ of Manchester, was the first member of the medical profession to grasp the true nature of these tumors.

As bearing upon the latency of cells destined to give rise to malignant growths, it is instructive to note that several cases of chorio-epithelioma malignum are now on record in which the growth has developed several years after abortion, no subsequent pregnancy having occurred.

More recent studies have demonstrated that cell masses of this type may develop not as the outcome of the fertilized ovum and of uterine pregnancy, but also as the result of teratomatous or embryomatous growth. They have, for example,

been found in the male, in connection with testicular embryomas; have been described in connection with mediastinal and cranial teratomas. The origin in all cases must be regarded as the same. The included potential individual in its growth must develop or tend to develop a chorion, and primarily its nourishment must be gained through this chorion, with its cell masses invading the veins of the host. As in the case of the fertilized ovum, so here these chorionic cells may take an aberrant excessive growth.

It may be questioned how are we to classify this form of tumor. If we define the teratoma as the product of the aberrant growth of cells from one individual within the tissues of another, then the destructive placental mole and the chorio-epithelioma are teratomas. Histologically,

¹ Manchester Medical Chronicle, 1892.

however, they do not represent the embryo, but only the sac in which that is developed; they represent only one tissue; they are formed of cells which under no conditions are capable of giving rise to the individual, cells which are unipotential, not totipotential. It makes for clearness to regard the teratomas as essentially developed from totipotential cells, and as these tumors are derived from unipotential cells it is best to regard them as blastomas, as teratogenous or heterochthone blastomas. Here also must be grouped those cases in which there eventually develops a malignant sarcoma, epithelioma, or gland cancer from one element of a typical teratoma, ovarian, testicular, or sporadic. Regarding them thus, we gain material help in our comprehension of the common autochthone blastomas, the ordinary tumors composed of one type of cell, tumors which, as we shall point out, originate from the aberrant growth of unipotential cells of the individual.

Thus, to sum up, we may make the following broad divisions of the neoplasms proper:

- I. **Teratoma**.—Tumors derived from cells capable of giving rise to all the tissues of the individual (totipotential cells).
 - 1. Twin teratoma (geminal or heterochthonous).
 - Example. Fœtal inclusion.
 - Filial teratoma (or autochthonous), due to the segregation and subsequent growth of totipotential cells of the individual.
 - (1) From non-germinal blastomeres.
 - *Example*. Epignathus, congenital sacral teratoma.
 - (2) From germinal cells.
 - (a) From aberrant germinal blastomeres.
 - *Example*. Sporadic teratoma of cranium, etc.
 - (b) From unreduced ovarian and testicular germ cells.
 - *Example*. Ovarian and testicular teratomas.
 - Members of this second class may be:
 - i. Complete, exhibiting derivatives of all three germ layers, or
 - ii. Reduced, derivatives of one germ layer failing to develop.
 - They may also be (i) typical or (ii) atypical.
- II. **Teratoblastoma**.—Tumors (autochthonous) derived from pluripotential cells of the individual: mixed tumors.
 - 1. **Diphyllic**, containing derivatives of two germinal layers. *Example*. Certain parotid and renal mixed tumors.
 - 2. **Monophyllic**, containing derivatives from one germinal layer. *Example*. Most renal mixed tumors.

III. Blastoma.—Tumors derived from unipotential cells.

- 1. Heterochthonous or teratogenous, the cells being derived from another individual.
 - *Examples.* Destructive placental mole and chorio-epithelioma malignum; epithelioma derived from an ovarian dermoid.
- Autochthonous, from the independent growth of unipotential cells of the host individual.

Examples. All other tumors composed of cells of one order.

CHAPTER XVI.

THE AUTOCHTHONOUS BLASTOMAS (ORDINARY TUMORS).

FROM these tumors, which are obviously the products of the growth of the cells of one individual within the other, even if those cells be only unipotential, we pass on to the main mass of neoplasms formed by the overgrowth of a single type of tissue, and derived evidently from the aberrant, autonomous growth of tissue cells of the individual or host. That this is their origin must for the moment be taken upon trust; the data to be afforded in the study of the individual forms will amply confirm this view. We have now to deal with the *autochthonous blastomas*.

Before attempting to classify them it will be well to gain a general grasp of their characters. We here include, as has been stated, all tumors, not teratogenous, exhibiting an independent localized growth of tissue cells of one order. To this statement a qualification must be added; all normal tissues possess a stroma or framework of connective tissues in which run the capillaries and lymphatic channels, and so we find in these blastomas that save where we are dealing with tumors formed or developed from the simplest form of connective tissue, similar to that constituting the stroma of organs, a stroma is also present, and we shall see this may vary in amount and importance. With this qualification these neoplasms exhibit the localized growth of one order of tissue. We thus recognize a large number of different forms of tumors -fibromas, composed of fibrous tissue; chondromas, of cartilaginous; osteomas, of bony; odontomas, of tooth; myelomas, of bone-marrow cells; myomas, of muscle fibres; neuromas, of nerve cells, gliomas, of neuroglia cells; epitheliomas,¹ of squamous epithelium; adenomas, of glandular tissue; and the list might be very considerably extended. For there are many more forms of blastomas than there are forms of individual tissue; and this because the cells composing a tumor, while originating from a mother cell of a particular type, may not present the characters of the fully differentiated tissue. It may, indeed, be laid down that they never present complete differentiation; taking the very simplest form, the densest fibroma always contains more cells and larger than does fully formed connective tissue. Nevertheless, in one series of tumors the cell characters are strikingly like those of the normal adult tissue, and with this reservation we speak of them as *typical*. On the other hand, a great number of blastomas depart very far from type. Whether from

¹This term should be employed with caution, so many meanings having been given to it. Thus, French writers include all glandular tumors of a malignant type under it.

BENIGN TUMORS

certain properties or certain relationships of the component cells, or it may be only from the region of primary growth and comparison with other tumors of like properties which in some of their parts afford the necessary clue, we are most often (but not always) able to recognize the tissue from which they originated. The component cells are only partially differentiated or it may be purely of the mother cell or vegetative type. These we speak of as *atypical* neoplasms.

And, as a general rule, the more we study, what appear at first sight to be exceptions are more and more found not to be such, and it may be laid down that the properties of these two groups, the typical and the atypical, manifest well-marked differences.

Benign Tumors.—The *typical blastoma* is composed of cells which in their characters approximate to those of some adult tissue. It is *circum*scribed and slow growing. The slowness of its growth permits a reaction on the part of the surrounding tissue, so that it is, in general, encapsulated and sharply defined, capable of being shelled out in its entirety from the tissue in which it grows (in some tissues, as in the brain, and to some extent in the bone, in which the capacity for fibrous overgrowth is slight, the capsule formation may be deficient). Growth, indeed, may for a time be arrested and then slowly proceed again. Save when situated in some position in which it comes to press upon some vital part, or when it gradually attains a size so great that it compresses the other organs in its neighborhood and disturbs their proper functions, the growth is harmless so long as it retains the characters here indicated. The shape of the mass varies according to position; embodied within a tissue and subjected to like pressure on all sides, it tends to be globular; situated on or near a surface, so that the pressure on one aspect is less than on another, it may spread laterally or become lobate, or nodular, or, as in the case of some epithelial outgrowths, cauliflower-like or papillose.

Such a tumor we speak of as *benign*, *i*, *e*., harmless in itself. It may grow slowly during the course of long years; may never attain any great size, or if it does, as in the case of some abdominal lipomas (or fatty tumors), which have been recorded as attaining a weight of 60 pounds and more, even then it is not an immediate, but at most an indirect cause of death through mechanical disturbance of other functions.

Yet another feature of the typical blastoma is the character of its growth. This occurs not merely at the periphery, but throughout the mass, in the central parts as well as toward and at the periphery. Such a mode of growth in a mass that is already spherical tends merely to result in the production of a larger sphere. It is spoken of as the central or expansive type of growth. As distinct from the peripheral, we may refer to it as universal, although neither of these expressions is quite happy.

Not all typical blastomas present this form; a chondroma, for example, grows only at the surface of its lobules, and so exhibits what is strictly a peripheral growth. That growth, however, from the inner aspect of its perichondrium, is, if we may so express it, centripetal rather than centrifugal.

THE AUTOCHTHONOUS BLASTOMAS

Malignant Tumors.—The atypical blastomas, on the other hand, are formed of *imperfectly differentiated tissue*, and, as we have repeatedly had occasion to note (p. 103 and elsewhere), with lack of functional differentiation there is a corresponding manifestation of increased vegetative and proliferative capacity on the part of the cells. As a matter of fact, in these tumors the cells are characterized by *active proliferation*. There is relatively *rapid growth*, increase in the number of cells, and increase in size. The rapidity of this growth and expansion of the mass prevents an adequate reaction on the part of surrounding tissue; there is *little or no* sign of encapsulement; the tumor is thus not precisely circumscribed. It may appear so to the naked eye, but examination of sections of the removed material under the microscope shows proliferating cells extending between the fibres of the surrounding tissue, and by their active growth, if by no other means, causing the compression and atrophy of the specific cells of that tissue, leaving eventually a framework of connective tissue and vessels which becomes the stroma of the advancing growth; or otherwise, the atypical blastoma possesses the power of infiltration. What is more, in this expansive growth, either by extension along the lymph channels, or by erosion and rupture of a surface membrane or of the walls of a vessel, certain of these actively growing cells may become detached from their fellows and, becoming conveyed to a distance either by the lymph stream or by the lymph or other fluid bathing a surface, or by the blood stream, may become arrested in some locality where conditions favor, or do not prevent, their continued growth, and there multiplying, they develop new tumors of the type of the parent growth. Such new-growths are termed metastases.

Tumors manifesting these properties are known as malignant. Histologically, we recognize two main types, the sarcomas, or atypical cellular tumors of the connective-tissue type, and the *carcinomas*, or cancers of a more glandular type. We shall later inquire more fully into other conditions associated with malignancy. To note them here would perhaps raise the false impression that all atypical blastomas exhibited That is not the case. We must, indeed, emphasize that the them. properties above recorded are those of the "type" atypical blastoma. There are, indeed all transitions, from the typical to the atypical form. A tumor may for years have been slowly growing and then in one portion take an active growth; in such cases it may as a whole be surrounded by a capsule, but in one or more areas the examination of sections shows that that capsule is becoming infiltrated; tumors of perfectly benign type-well-formed and typical chondromas and even myomas-are on record as giving rise to metastases; on the other hand, definitely malignant growths rapidly growing and rapidly fatal may exhibit no metastases. Ehrlich,¹ for example, has recently noted that this is the case with his experimental and intensely malignant mouse cancer. After subcutaneous inoculation, there may be metastases in the lungs, but although the primary transplanted growth attains a huge size, these metastases

¹ Zeit. f. Aerztl. Fortbild., No. 7:1906.

MALIGNANCY

are so small as to be recognizable with difficulty and only by careful microscopic search. More than the mere escape of cells from the region of primary growth is necessary to cause metastatic growth.

We shall discuss these variations in properties later. In the meantime it is well to gain, at the outset, a general grasp of these relationships between tumor-cell differentiation and benignancy, on the one hand, and proliferation and vegetative type of cell and malignancy on the other. And that for practical purposes; because it is on these characters and their relationship that we base our diagnosis and prognosis; from them we determine whether a given tumor may be left in the organism, or must be removed, if not already too far developed to render removal in vain. We would only add the warning that for prognosis everything depends upon what is the adult type of the cell in the case of any par*ticular tumor*; the tumor cells may appear to be of a distinctly vegetative type, as in the so-called giant-celled sarcomas (myeloma), and yet the tumor not be malignant to any marked degree, this vegetative appearance being characteristic of the adult marrow cells from which the tumor has originated; on the contrary, the melanoma which appears to be composed of more highly differentiated spindle-like cells than is the giant-celled myeloma is intensely malignant; no tumor shows more abundant and widespread metastases. From these examples it is obvious that while the general rules that we have laid down are capable of application and are of distinct use for the grouping of phenomena, nevertheless, for the purposes of safe prognosis, it is essential to have an intimate acquaintance with the life history of each individual form of neoplasm. It must be realized, in short, that what has just been stated regarding the characteristics of benign and malignant tumors is of the nature of a "rule of thumb" rather than an exact generalization. There are cases to be noted which it is difficult to harmonize.

MALIGNANCY.

Here it is appropriate to consider what constitutes this state of malignancy which is so striking and important a feature of the atypical tumors. We have already indicated some of the conditions which are factors in its development. These are, briefly:

1. Vegetative (embryonic) character of the tumor cells.

2. Rapidity of growth.

3. Peripheral extension, lack of capsule and infiltration of the surrounding tissues.

4. Tendency to develop metastases.

These, it will be seen, are all related; th γ are the expression of an inherent augmented vegetative and proliferative activity of the cells constituting the tumor, over and beyond that possessed by the surrounding tissues. There are, however, other associated features which must be noted.

5. Tendency to central degenerative changes. The activity of the cell

THE AUTOCHTHONOUS BLASTOMAS

multiplication in a mass which cannot freely expand owing to the pressure of surrounding parts must, it will be seen upon consideration, lead to some compression of the vessels supplying the mass as the volume of the cells increases; there must, that is, with growth be increased internal pressure, and, as a matter of fact, we find in all malignant tumors that the peripheral cells exhibit evidences of active growth and mitosis; they are in the best position to gain nourishment from without; the central mass of the tumor shows tendencies toward degeneration and atrophic changes, and this to such an extent that in some large cancer masses we may find the cells in the centre completely absorbed (through autolysis), and a central cavity containing serous fluid. Similarly, where the tumor is superficial, the parts farthest from the blood supply, *i.e.*, the most superficial parts, are liable to undergo necrosis, ulceration being the result.

Certain authorities regard this liability of malignant tumor cells to degenerate as an indication of low vitality on their part. The very reverse would seem to be the case, as we shall point out later. It is the anarchical growth of the tumor cells that brings about the central degeneration.

6. Liability to Recurrence after Removal.—This is a property associated with the infiltrating character of these growths. Although not noticeable to the naked eye, the tumor may spread along the lymphatics for a long distance away from the main mass, and this not by detachment of individual cells and groups of cells, but, as indicated more particularly by studies of mammary cancer, by continuous growth of the cells in series along these channels. Such cells, extending far beyond the obvious border of the tumor, may not be excised, and may subsequently, by proliferation, develop into nodules of new-growth. It is this wide extension and danger of recurrence that is the basis of the modern radical and most extensive operations for the removal of cancers.

7. Cachexia.—By cachexia we imply a lowered impoverished state of the system, indicated especially by a wasting of the tissues coupled with an abnormal complexion. Several chronic wasting diseases induce a cachexia. The malignant cachexia is more particularly characterized by the extreme degree of wasting which may ensue (it is not always present), by the fact that while the tissues atrophy the tumor continues to grow, and by a peculiar sallowness of the skin. It is difficult, if not impossible, to describe this sallowness; the healthy color disappears and is replaced by an anemic yellowish-gray appearance, which once seen is easily recognized.

8. Anemia.—Anemia is a constant accompaniment of malignancy, and, indeed, this altered condition of the blood must be held to underlie the cachexia. In extreme cases this is indistinguishable upon blood examination from idiopathic pernicious anemia (which, indeed, is accompanied by an allied although distinct cachexia). These latter conditions suggest immediately that the actively growing tumor absorbs the nutritive elements of the circulating blood and thereby starves the rest of the system. This may—doubtfully—be a factor. We find, however, that the cachexia

672

x

MALIGNANCY

is not proportional either to the size or rate of growth of the tumor. It differs also according to the nature of the growth; there may be a marked cachexia associated with a small carcinoma or atypical glandular neoplasm, and little or none with a relatively large sarcoma or atypical connective-tissue growth. There are, it is true, in some cases of extreme cachexia and wasting, complications which are evidently in part responsible; thus certain tumors of the alimentary tract, notably of the œsophagus and stomach, may so narrow the lumen of the affected part as to arrest the passage of food and lead to starvation. Other superficial malignant growths undergo extensive ulceration, and by the absorption of the foul products from their surfaces and by low forms of infection the general bodily condition may be greatly lowered. But even when these cases are excluded we still encounter cachectic and anemic states associated with malignant growths.

If the tumor cells are not functional, they, nevertheless, in their growth -and it may be more particularly in their degeneration-discharge certain soluble substances into the lymph and blood. The more marked cachexia which accompanies malignant tumors of a glandular typetumors that are derived from cells which normally secrete bodies of the nature of enzymes-is at least, as Borst points out, suggestive. It has been noted by more than one observer that even the metastatic growths, outside the liver, of primary liver-cell tumors, secrete bile, and, as we pointed out some little time ago, the remarkable alternation in the mental and other conditions which may follow the removal of adenomatous or glandular tumors of the thyroid is difficult to explain, save in the assumption that the tumor supplies an internal secretion which has a direct influence upon the nervous and other systems.¹ More recently, Waring² and M. B. Schmidt³ have called attention to this active secretion of products by cancer, and Buxton⁴ and his associates have investigated the enzymes obtainable from malignant growths, while several observers have called attention to the hemolytic activities, more particularly of extracts and products of autolysis of cancers of glandular type. It is probable, then, that the malignant cachexia is primarily the outcome of deleterious products discharged or diffused from the malignant growth, both active modified secretions and the products of autolysis.

While the above conditions *in general* accompany malignancy, this is not, it must be remembered, equivalent to stating that all are *essential*

³ Triennial Congress of Amer. Surg. and Phys., 4:1897:103. It is, however, debatable whether the ordinary localized colloid goitres should be classed as blastomas proper.

² Jour, of Anat. and Physiol., 28:142.

³Virchow's Arch., 148:1897 (on secretory processes in secondary growths of a thyroid cancer in the liver).

⁴ Jour. of Med. Research, 9: 1903; 356, and Buxton and Shaffer, ibid., 13: 1905; 543; see also Macfadyen and Harden, Laneet, Lond., 1903; ii: 224. Proteolytic enzymes amylase, oxidase, lipase, and other enyzmes have been detected, but somewhat irregularly and independent of the character of malignancy of the growths, and similar to those found in other cellular tissues.

THE AUTOCHTHONOUS BLASTOMAS

accompaniments. Thus a highly differentiated tumor may afford metastases and, on the contrary, an infiltrating and destructive tumor, such as is characteristically the rodent ulcer, may form none; if imperfectly removed, a benign, well-encapsulated tumor may recur; if removed at an early period, and sufficiently thoroughly, a growth of malignant type will not.

Active vegetative and peripheral, as distinct from universal, growth with accompanying infiltration would seem to be the essential features of malignancy; the other features may or may not be added, although most often they are present. Or, with yon Hansemann, we can concisely express it, that the essential distinguishing marks of a benign growth are that it does not infiltrate and destroy and does not form metastases. To this we would add that the benign neoplasm shows not merely peripheral, but universally diffused growth, if slow, throughout its substance. Our position, in short, with regard to malignancy is very much that of the astronomers with regard to the solar path. Recognizing certain general laws, they can calculate the sun's position on a given date in relation to the stars in general with very fair accuracy. But always there is a certain error or correction to be made to the figures gained by working out the law. The value of that error or correction, it is true, has been determined by the astronomers; it represents some unknown influence or force acting upon the sun and deflecting it from its path in space; we have not arrived at this point. Here there would seem to be some other factor not as yet clearly determined acting in addition to those we have noted, and causing a tumor to behave otherwise than we would infer it should behave from its structure and relationship.

In his "Geschwülstlehre," Ribbert lavs stress upon the non-existence of malignant tumor cells per se. There are, he says, no malignant cells. Without discussing the reasons which led him to this conclusion, we will frankly say that this view is untenable, and this with all due deference to Ribbert's great authority, and recognition of the much we owe to him in advancing in many directions our knowledge of neoplasia. As well, it appears to us, might one deny the existence of virulent bacteria. Apart from mere general considerations, one fact alone, acquired, it is true, since his work was published, demonstrates the incorrectness of this dictum. We refer to Ehrlich's demonstration¹ that, by passage through mice, mouse cancer can be rendered more and more malignant, until it will surely "take" in close upon 100 per cent, of the animals, instead of, as in the first transplantation, from 5 to 35 per cent.; and grows so rapidly that in seven or eight days the minute portion of tissue transplanted has reached the size of an almond, which, compared with the size of a mouse, is something enormous.

It is difficult to state the exact figures of "takes" at first transplantation. The figures here given are excessive. They are gained from Ehrlich's statements regarding the forms which from the first were of the more virulent type, and were used for the purposes of passage.

¹ Zeitschr. f. Aerztl. Fortbild., loc. cit.

MALIGNANCY

With 21 tumors of the same adenocarcinomatous type, taken in succession, some more typical and adenomatous than others, inoculating 282 mice, he only gained 2 positive results. In such cases, in which the resisting powers of the tissue of the animals inoculated remain constant, and the vegetative powers of the inoculated cells undergo increase, to deny that we have evidence of increased malignancy on the part of the latter is to juggle with words. It is the grade of vegetative power of the cells which determines their malignancy, though, as we shall point out (pp. 682 and 686), the malignancy of a given tumor in a given tissue of a given animal is the expression of the interaction between the cell malignancy and the resisting powers of that tissue toward the growth of that particular type of cell.

18

1

a

g

1

The above are the characteristics of malignancy *proper*. There are, however, two other conditions, to which, unfortunately, too often the same term is employed without the use of distinguishing adjectives—conditions which are, it is true, malignant in the sense that they tend to a fatal termination, but in which the chain of events is of a different order. These are:

Malignancy in Virtue of Site, or Malignancy of the Second Order .--- A slowly growing tumor of any order which, developing elsewhere, would be perfectly harmless, may, by pressure upon vital organs, arrest their function, and so induce death. A relatively minute gliomatous or fibroid tumor of the brain, or its membranes, by pressing upon the medulla, for example, may cause death, and so may be termed malignant. A lipoma of the skin may grow to enormous dimensions, and cause little disturbance; a small lipoma of the intestinal mucosa, by blocking the lumen or causing torsion of the gut, may soon be fatal. Allied to these conditions, when tumors attain great size, they may eventually so press upon surrounding organs as to lead to their atrophy and to the obstruction of their ducts, and so attain to malignancy of this order; and it is noticeable that, attaining great size, they may also induce a cachectic condition. But in all these cases it will be seen that we are dealing with another order of affairs.

Malignancy of Recurrence.—Local Malignancy.—Certain tumors, again, slowly growing and of a typical rather than of atypical, cellular type, are malignant in so far that, after apparent extirpation, they tend to recur. They may be allowed to grow slowly for years without exhibiting any tendency to invade the surrounding tissues or to form metastases elsewhere, but, if extirpated, a second tumor of the same nature is peculiarly liable to develop in the same neighborhood; and what is more, the recurrent tumor tends to be more cellular, to grow more rapidly, to invade the surrounding tissues, and to be definitely malignant of the first order. The cause of this malignancy of recurrence is either (1) that the extirpation has not been complete, and certain of the tumor cells left behind are incited to active proliferation by the hyperemia and the lessened pressure in their neighborhood after removal of the main mass; or (2) that the tissues of the part have a predisposition toward tumor formation, and that after *complete* removal

THE AUTOCHTHONOUS BLASTOMAS

of the primary growth the two factors above mentioned favor the active proliferation of the neighboring cells, or foci of cells of the same order. For the present it must be left an open question whether we have to deal with one or both of these processes. Fibroid and mucoid tumors (e. g., nasal polyps) more especially present this liability to local recurrence and the eventual taking on of true malignant properties. It is usual, nowadays, to state that these tumors are from the first sarcomatous; my experience leads me to hold that this is not always so, that certain so-called recurrent fibroids may at first show absolutely no signs of sarcomatous nature, but belong to the blastomatoid group, to be presently noted (p. 714).

Before leaving this subject of malignancy, it is necessary to point out that a tumor which has for long been benign in its properties may eventually assume malignant characters (of the first order). The preceding pages indicate that true malignancy is a function of the rate and extent of cellular proliferation. From causes not as yet classified and fully studied, cells which at first undergo slow proliferation and complete differentiation may assume rapid growth, and, with this, all the characteristics of true malignancy may be developed.¹ From a prognostic point of view, this is a matter to be continually kept in mind.

METASTASES AND THEIR PROPERTIES.

We saw that in infective inflammation, in pyemia, or in tuberculosis, for example, the specific organisms might be carried from the primary lesion, and, becoming arrested in some more or less distant organ, might there set up new foci of inflammation, leading to, it might be, abscess formation, or new formation of infective granulomata.

These metastatic inflammations have, by many, been compared with the neoplastic metastases, and the similarity of the two processes has been made an argument by those who uphold the parasitic origin of tumors. It must, however, be kept in mind that these two processes of metastasis are absolutely distinct. In infection, the bacteria which are carried to distant parts set up a local reaction, and it is the cells of the part, together with migrating leukocytes, which are the factors in the local tissue disturbance, so that the so-called infective granuloma is composed of new tissue derived from the region affected; no matter what part or organ is the seat of the process, the results are the same, namely, the production of more or less well-developed fibrous tissue, inflictated with migrating leukocytes.

In the neoplastic metastases the local reaction is purely secondary; the new-growth is a development of cells which have been carried to the part from the primary tumor, and these cells give rise to a tumor tissue,

¹ Despite overwhelming elinical evidence, there are still those who, holding fast to the belief that the vegetative powers of the cell cannot undergo augmentation, strenuously deny this conversion.

METASTASES AND THEIR PROPERTIES

which varies in its character in strict accordance with the characters of the primary tumor. It is not the local cells which form the tumor metastasis, but derivatives from the original tumor. Here, indeed, the migrating cells are the parasites.

It is a general impression that the metastasis faithfully reproduces the parent growth. This, while most common, is not universally the case. It may but reproduce the *general* type of the parent growth. There is, indeed, a tendency often noted for the metastasis to be more actively growing, to be of a more vegetative type, the cells reverting

FIG, 196



Typical cells, drawn to scale from a tumor of the adrenal cortex and its metastases: 4, from the normal adrenal cortex; 2, 3, from the tumor in the adrenal cortex; 1, 5, 7, from a metastasis in the lung; 8, 9, from metastasis in the brain (5, a polynuclear leukocyte for comparison. (Woolley)

to a yet simpler condition. An extreme example of this variation was studied in our laboratory by Woolley, ¹ in a case of a tumor of the adrenal cortex, in which every transition was found, from the primary cancerlike mesotheliomatous growth to pure round-celled, sarcoma-like metastases. Jores² and Meakins⁸ have published similar cases. This variation has been treated fully by Beneke and von Hansemann.⁴

¹ Vireh, Areh., 172: 1903: 30, and Trans. Assoc. Amer. Phys., 17: 1902: 627.

² Deutsch. med. Wochenschr., 1894:208.


e d l

³ Proc. New York Path. Soc., N. S., 9:1909:19,

^eDie Spezifizität, etc., der Zellen, Berlin, 1893. See also Low and Lund, Jour. Med. Research, 7:1902.

THE AUTOCHTHONOUS BLASTOMAS

In such cases a microscopic study of all the metastases demonstrates clearly, from the various transitions found, that what is here stated is true, and that they are derivatives from the original growth. Occasionally, indeed, the original growth is comparatively insignificant, while the secondary growths obtain an enormous size and expansion. There may, for example, be a small, insignificant ulcer of the stomach, in the walls of which we may detect evidences of the existence of fibrous or scirrhous cancer or gland tumor, while in the liver the growth may be several inches in diameter, the organ being enormously enlarged. A study of the characters of the two growths points to the conclusion that the insignificant stomach ulcer is older and earlier, and is the point of origin of the metastasis. Or, again, in metastatic sarcoma (chromatophoroma) of cutaneous origin, the primary growth in a congenital pigmented mole may be no larger than a pea, and yet in



Extension of a small round-celled sarcoma (S) beneath the endothelium (E) of a vein; at I the sarcoma cells are infiltrating the middle coat (M) of the vessel. (Martin.)

numerous tissues we may find metastatic nodules the size of a cherry, and larger. These cases are the exception rather than the rule; in general, the primary growth is larger than any individual metastasis.


Mode of Origin.—Such metastases, as distinct from local infiltration, may originate in four ways:

1. The cells of the neoplasm, infiltrating between the tissue cells of the part, penetrate into the lymph spaces, and from these certain cells become carried into lymph vessels, and so the new-growths recur, in the first place, along the course of the *lymphatic system*. Often, in the more immediate neighborhood of the tumor, what appear to be separate metastases are found to be in direct continuity with the original tumor, solid strands of tumor cells, injecting the lymph vessels, passing from the primary tumor, and then at one or other point, where the conditions are favorable, more active growth of the cells has taken

METASTASES AND THEIR PROPERTIES

place, forming definite little tumor masses. As shown by Stiles, in mammary and other cancers such growths of the tumor cells along the lymph channels beyond the main mass may be demonstrated by treating the fresh excised surrounding tissue with nitric acid, the cells of epithelial type affording the xanthoproteic reaction, and standing out an opaque yellow.

2. The tumor in its growth may erode a vein, and certain of the cells may thus pass directly into the blood stream, and by that be carried to different portions of the body. Or, again, the tumor may originate in immediate connection with the vessel walls, and tumor cells, as in the case of some malignant atypical connective-tissue growths (sar-comata), may replace to a very large extent, or entirely, the normal endothelium lining the vessels both preexisting and newly formed in



Sarconntous transplantation. Abdominal lymphosarcoma; section through two of the close-set small growths: T T, covering the surface of the liver G. These have gained a covering layer of endothelium. (Martin.)

the tumor. More often they are found lying immediately beneath the endothelium. In such cases these tumor cells, by the result of slight injury to the tumor, or merely from the very character of the growth, may become free in the vessels. When either of these events happens, and the cells are carried along the veins, they may occasionally be arrested in the heart and multiply there. In general, however, it may be laid down that they are liable to be arrested in the first system of capillaries to which they are carried by the blood stream. Thus, sarcomas occurring along the branches of the portal system are peculiarly apt to form secondary growths in the liver; those occurring along the main venous system are peculiarly apt to form secondary growths in the lungs. Rarely, as pointed out by Zahn,' in cases of small-celled tumors in which growth is on the venous side of the heart, with no

¹ Virch. Arch., 117:1889:1.

secondary growth in the lungs, multiple metastases may be present in distant organs, and can only be explained by passage of individual cells or minute cell masses through the lung capillaries without arrest, the foramen ovale being found closed.

Such passage must not be regarded as impossible. Along with other observers, inoculating the relatively large cells scraped from a rabbit's liver into a vein in another rabbit, we have found these cells in the renal vessels and those of the renal capsule.

It has further to be remembered that where extension is mainly along the lymphatics, eventually the cancer cells from the lymphatic secondary growths may find their way into the thoracic duct, and so from this into the main circulation, and thus eventually, in these cases also, there may be extension and metastatic growth by means of the main circulatory system.

3. A third means is but a modification of the first. The various serous cavities of the body are essentially large lymph spaces. If a tumor penetrates and affects the lining membrane of one of the serous cavities, certain of its cells may become free, and thus pass to one or other region of the serous cavity. This is especially well seen in connection with the peritoneal cavity; the cells may become arrested here or there upon the surface, grow, and so there may be formed numerous metastases, scattered all over the serous surface of such a cavity. This method of extension is spoken of as *transplantation*.

Nor may this be wholly confined to the lymphatic system. There are cases in which cancer of the oral cavity has been found accompanied by secondary growths in the stomach, or of the stomach in the intestines. The first and most natural explanation in such cases is that of surface transplantation, and possibly in some cases this is the correct explanation. But careful study in others (more particularly as between stomach and intestines) has demonstrated marked lymphatic involvement, implicating the (retroperitoneal) glands, and thence extending to a distance. This possibility must always be carefully excluded before the decision is reached that a case is one of true transplantation.

4. Somewhat allied to this last form is transplantation by apposition. Cases are recorded in which (a) where one lip has been the seat of cancerous growth, the other has become, later, involved at a corresponding point, (b) where the skin of an arm, coming into contact with an ulcerated, cancerous breast, has become involved, or (c) where the parietes opposite to a viscus presenting superficial malignant growth, without showing adhesions, becomes the seat of growth. Here, clearly, cells become transplanted upon another surface, which, presumably by attrition or other cause, has lost its protective covering. Such transplantation does not, by any means, necessarily ensue.

Retrograde Metastasis.—It deserves note that, while the general rule is that tumor cells are carried in the direction of the normal blood or lymph current, and secondary growths occur in conformity therewith, we may encounter paradoxical cases in which the opposite is the case, or in which, at first sight, there appears to be no relationship

RETROGRADE METASTASES


between the site of the secondary growth and the blood and lymph flow from the primarily affected organ. Thus, in cancer of the breast, it is by no means infrequent to find the head of the humerus of the same side infiltrated with the growth. Such cancer extends in the main by the lymphatic system. Now, the lymphatics from the head of the humerus and shoulder region pass toward, and not away from, the axillary glands. Or a primary malignant growth of the kidney or of the heart region may show extensive secondary growths in the liver, and that with little evidence of growth in the lungs. The kidneys have no relationship with the portal veins. In such cases we have to fall back upon the retrograde passage of cells along vessels. In the first case

a

dv

ie

10



Schema of retrograde embolism to illustrate mode of retrograde metastasis: T, may be taken to represent a tumor mass in one of the branches of the inferior vena cava, part of which becomes iliberated at A and earried upward toward the heart. By temporary stasis and back pressure the mass may pass into the renal or hepatic vein and become lodged there. (After Lubarsch),

we have to suppose that the extension of cancer into the axillary glands blocks the normal channels, so that its lymph has to find a collateral or roundabout circulation whereby it and its contained cells may pass along certain channels in the reverse direction.

Another possibility in these cases which has to be borne in mind, and excluded, is that of direct, continuous growth along the lymphatics from this axillary gland. This, however, will not explain all the cases. Thus, Belin¹ has called attention to the relative frequency of left supraclavicular nodules in cases of visceral cancer, best explained by variation in pressure in the thoracic duct at its junction with the jugular vein, leading to reflux of cancer cells into the neighboring glands.

¹ Thèse de Paris, 1888.

THE AUTOCHTHONOUS BLASTOMAS

In the second case, we conclude that when there occurs a negative pressure in the inferior vena cava, cells may be actually drawn back from the right auricle, or fall back from the inferior vena cava into the hepatic veins, until they become arrested in some of the smaller vessels of the liver, and there, growing, cause the development of metastatic tumors.

Tissue of Predilection.—There is a feature characterizing metastatic tumors which has been little dwelt upon, which, nevertheless, is of high importance, as throwing light upon one important factor determining tumor growth in general. It is a feature parallel with what is seen in infective conditions. It is notable, for example, that in pyemia, where there may be a development of multiple abscesses, this development does not take place indifferently in all the tissues and organs of the body. They may be numerous in the lungs and kidneys, and yet in the same case scarce an abscess may show itself in the liver and spleen, and none at all in the muscles of the body. And, as was pointed out in discussing tissue predisposition (p. 408), to explain the distribution of these abscesses, we are forced to recognize that the resistance or reaction of the tissues plays a part in determining the fate of the bacteria.

Now, the same is true with regard to neoplastic metastasis. We find, for example, that melanotic tumors are especially liable to form new-growths in the liver; malignant glandular tumors of the thyroid are peculiarly liable to form secondary growths in bone; only rarely do we come across metastatic growths of any order developing in the muscles. Numerous other examples might be given, but this curious distribution renders it evident that the escaping cells of malignant tumors gaining entrance into the blood stream, and becoming carried to various organs, do not by any means necessarily proliferate; only under certain special local conditions is proliferation in general possible; or, otherwise, in many regions of the body the preexisting cells react against the invading cells, and lead to their destruction or the arrest of their growth. Only in this way is it possible to explain the remarkable distribution we occasionally meet with in connection with these metastatic growths.

Recklinghausen first called attention to the predilection of prostatic cancer to form metastases in bone, a predilection so marked that we have made the diagnosis of prostatic cancer from the existence of extensive involvement of the pubic bones. Leuzinger¹ noted that while osseous metastases occur to the extent of 2.3 to 3.5 per cent. in cases of uterine cancer, they are 14 per cent. in cancer of the breast, and 20 to 25 per cent. in thyroid cancer. Menetrier³ and Handford³ note a liability to muscular metastases in cases of primary lung cancer. Rolleston and others, to bone metastases in the part of adrenal cortical tumors.

¹ Inaug. Diss., Zurich, 1886.

² Art. Neoplasie, Bouchard's Pathologie Génér., vol. ii.

³ Trans. Path. Soc., London, 39:1888:48.

METASTASES: TISSUES OF ELECTION

Indeed, considering all the facts at our disposal with regard to metastasis, we are forced to recognize that it is not the mere escape of cells from a primary tumor which determines the development of these secondary growths, but that the fact of prime importance is the proliferative capacity of these cells as compared with the reactive powers of the tissue in which they find themselves arrested. Taking into consideration what we know regarding the attempts at transplantation of normal tissues, we are, I think, forced to conclude that the same conditions are in existence here in connection with the tumors. It is only when tumor cells have peculiarly active vegetative powers that, being carried into regions and surroundings widely different from their original habitat, they are capable of continued proliferation. Or, conversely, it is eminently probable that tumor cells are liable to escape, not merely from malignant, but also from benign growths; but, when they escape from benign growths, their vegetative activity is not in general sufficient for them to grow in altered surroundings, and so it comes to pass that no metastases are formed, and, even escaping from malignant growths, it is only in particular localities that they can grow. Nay, more, as with pyemia we have to recognize that the resisting power of the body and of the individual tissues may vary during the progress of the condition. The abundant studies of late years upon the transplantation of tumors in the lower animals have indeed demonstrated the existence of three periods in connection with such growths: (1) A preliminary, succeeding the primary transplantation, in which further transplants may be made with success; (2) a reactive. in which, as a result of the general reaction set up through the system by the growth of the tumor, further transplantation has negative results, the "graft" undergoing necrosis and absorption; and (3) a terminal, in which metastases show themselves, and further transplantation is successful; in which, evidently, the products of the growing primary tumor have exhausted or inhibited the antagonistic mechanisms of the system. Depression of the resisting powers on the part of the tissues may ensue whereby metastatic growths eventually can occur within them.

We met with a striking demonstration of this fact some few years ago in the case of an elderly woman, in which the following history was obtained: There had been noticed in the left breast, for some years, a small, dense, scirrhous cancer, which, as sometimes happens, had remained practically stationary. Some months before her death this patient had fallen on some steps and had hurt her back so severely that she was confined to her bed for several days, and, following upon this, it was her lumbar symptoms that most troubled her. At the autopsy a dense, and mainly fibrous scirrhous cancer was found in the breast (which had shown no increase during the eight months she was under observation), with involvement of the axillary and supra- and infraclavicular glands; but most marked was the extensive infiltration of the lumbar vertebrae, more cellular, it is true, than that of the breast, but clearly of the same type. It was impossible to resist the conclusion

that here trauma had lowered the tissue resistance, so that now cancer cells, brought to the bone, had found conditions favorable for growth.

This matter of tissue resistance, or relative insusceptibility, has, as we shall later show, a most important bearing upon the etiology of blastomas and upon the arrest and cure of the same.

The Production of Metastases by Tumors of a Benign Type.-Lastly, before summing up, attention has to be called to the fact that certain tumors of a benign type are liable to produce metastases. The chondromas, or tumors formed of cartilage, afford frequent examples of this, both in man and the lower animals. I have come across more than one example of localized and well-defined chondroma of the mammary glands in the bitch (a not infrequent condition) showing multiple small secondary cartilaginous nodules in the lungs. It cannot be imagined that fully formed cartilage cells invade the bloodyessels or lymphatics, become detached, and are then brought to rest in the pulmonary capillaries, and continue to grow there. A more rational explanation is afforded by the mode of growth of this particular form of tumor. Unlike the majority of benign neoplasms, the chondroma grows essentially by peripheral cell multiplication. Just as normal cartilage grows from the perichondrium, so at the periphery of a nodular chondroma there is a vascular zone containing actively proliferating cells-chondroblasts-small, actively vegetative cells, of "embryonic" types, and these it is which gain entrance into the circulation and, carried elsewhere, set up metastases. The same procedure, I would add, explains the development of osteomatous or osteosarcomatous metastases. An arm, removed by my colleague, Dr. James Bell, for osteosarcoma of the upper end of the humerus, and studied by Dr. Keenan, showed the axillary glands converted into nodules of solid bone. Here it was not the adult bone cells that had found their way into the lymph stream, but proliferating osteoblasts, which, arrested in the glands, and undergoing further growth, had fulfilled their normal function and had given rise to true bony tissue. Similarly there are cases on record of secondary osteosarcomatous growths in the lung. Krische,1 in 1889, from Orth's laboratory, described multiple metastases of a fibromyoma of the uterus, of a form of tumor, that is, which, while most common, is characterized, by its slow growth and benign properties. Cases of this nature are exceptional; nevertheless, since that date some half-dozen others have been placed on record.

A condition more difficult to name and classify is that seen in cases of so-called *malignant adenomas*. The adenoma is a tumor differing from the glandular cancer in that it is formed of an excessive growth of glandular elements, tubules, and acini, which still retain a typical glandular structure. It is in connection with the liver and with its ducts that we are apt to meet with tumors of this type. In connection with cirrhosis numerous cases are on record—on this continent by Finley and myself, Fussell and others—in which multiple tumors have

¹ Inaug. Diss., Göttingen, 1889.

developed formed of masses of proliferated liver cells. It is most difficult to draw the line between compensatory hypertrophy, occurring in the lobules of the liver secondary to cirrhosis, simple adenoma complicating cirrhosis, and adenocarcinoma; but, certainly, in the advanced cases metastases may form in other organs; and what is interesting is that both in the primary nodules and in these metastases there may be a formation of bile. The cells, that is, still retain certain of their functions. In connection with the bile ducts there may be a development of tumors formed of tubes repeating in their structure the normal bile ducts, and these, again, may give rise to metastases.

Equally, if not more remarkable tumors are met with in connection with the thyroid gland; indeed, the tumors of this organ-and they are common and very varied in character-present many aberrant features. In this connection I would point out that not only do wellmarked infiltrating cancers of this gland very frequently exhibit still the tendency toward the formation of colloid within the newly formed but irregular acini, but that we encounter, rarely, it is true, remarkable glandular masses growing within the bones of the skeleton which reproduce in structure, and in the presence of true colloid material within the alveoli, the young or growing thyroid tissues. These tumors are structurally, therefore, thyroid adenomas, and, what is most remarkable, is that in more than one case careful study of the thyroid gland proper has failed to reveal evidence of any primary tumor there. Structurally, these tumors are of benign type; clinically, they are found to grow extensively, replacing the bone, and manifesting definitely malignant properties.

In short, all these malignant adenomas exhibit the same want of correspondence between structure and properties. In some, at least, namely, those of the liver, a cause of this development of metastases has been determined; the normal liver cells are in very close relationship to the bloodvessels, lying immediately beneath the capillary endothelium, and it has been observed that in their growth the tumor masses project into, distend, and grow along the hepatic vessels; thus, portions of these finger-like masses are liable to be detached and carried to the pulmonary capillaries, etc., where, coming to rest, they produce metastases. The same course of events may take place in other instances. In the thyroid, in the production of these adenomata within the bones, another process would seem to be at work. In the first place, the bones affected may be at such a distance from the organ that the theory of fœtal inclusion cannot, in reason, be advanced, by which I mean it is unreasonable to suppose that we are dealing with cases in which, during development, portions of thyroid tissue have become detached and included in the growing bones, to lie dormant for years, and eventually take an active growth. Rather, it would seem, that here we have a state of affairs similar to that noted in connection with enchondromata. The thyroid, structurally, is peculiar in this, that even during adult life there can be detected in it small accumulations of indifferent cells, which, as Wolfler has pointed out, are truly "mother

cells," and, under certain conditions, are capable of active proliferation and the production of new acini. As we have pointed out (p. 680) in connection with the subject of cell emboli, it is probable that even cells of considerable size, like liver cells, not infrequently become liberated into the circulation, and so become carried to different parts of the organism. Under ordinary conditions these cells become destroyed. There appears, in fact, to be normally in the system a very definite intercellular antagonism, so that cells out of place are acted upon and destroyed by those with which they come into contact. As I have already noted in connection with transplantation, embryonic and actively proliferating cells are not so surely destroyed as are adult and functioning cells. The simplest explanation of these intra-osseous adenomas is that, during the course of the development of new acini in the thyroid (at a time, that is, when these areas of small vegetative mother cells become richly vascular), certain of these cells become detached into the lymph or blood stream, and, gaining entrance thus into the circulation, if they happen to be carried into a bone, there find conditions favorable to continued growth. No other explanation appears adequate to meet all the circumstances.¹

The facts and observations here recited are ample to prove that a classification based upon the existence or non-existence of malignant properties cannot be satisfactory. If malignancy itself has different meanings according to circumstances, if a tumor, benign in one region, is malignant in another, in virtue of its position; if tumors, structurally similar, like the adenomas or the chondromas, can indifferently assume either benign or malignant properties, it is hopeless to seek to arrange neoplasms according to this one feature, however important it be from a clinical standpoint. Some other basis for classification has, therefore, to be sought.

Latency in Relationship to Metastases.—It is a significant fact that tumor cells conveyed to other regions may not immediately begin to grow and develop into a metastatic neoplasm, but may remain latent and inactive for months, and, indeed, for years, and then only take an active growth. The proof of this statement is afforded by the fact that a tumor of malignant type may be removed surgically with apparently perfect success, and then months or years later a progressive enlargement is noted in some lymph gland or other organ which, on removal, is seen to present growth of the type of the original tumor. In this way our colleague, Dr. Shepherd, removed a cancerous cervical gland from a woman on whom, eight years previously, he had performed total excision of the breast for cancer. Beeckel and Verneuil have reported recurrences after twenty-nine and thirty years, respectively.² In con-

¹ Several cases are on record of "struma thyroidea ovarii," in part associated with teratomatous masses, in part solitary. Doubt still exists as to whether these last are truly thyroid tissue or follieular adenomas of the ovary. (See Borst, Verhandl. Deutsch. Pathol. Gesellsch., 11: 1008: 98; with literature.)

² For literature, see Bircher, Centralbl. f. Chirurg., 1907: No. 26.

STROMA

nection with melanotic sarcoma of the choroid of the eye, some few cases are on record in which, after total extirpation of the affected eye, growth of similar characteristically pigmented tumors have shown themselves after long intervals in the liver and elsewhere. The longest period of such latency after operation removal that we have found recorded is twenty-one years.¹ Allied to these observations is that of Ehrlich,² that, whereas in general a mouse chondroma transplanted into other mice grows almost immediately, and with great vigor, in two cases four months elapsed before any sign could be made out of (subcutaneous) development.

On the other hand, transplanting cancerous and adenomatous tumors, of 94 such tumors, of each of which portions were transplanted into 20 to 30 mice, only 11 showed signs of growth, and this usually in but 1 to 3 of the animals inoculated; rarely in 6 or 7. In the great majority of the cases the transplanted material underwent atrophy and absorption. As we have pointed out already, when normal tissues are transplanted, the same is most often the case. When, as in cases of mammary cancer, we find secondary growth only in a single distant organ, such as the liver, we can only conclude that this same destruction has overtaken cancer cells transplanted into other organs. And so it is evident that one of three things may happen to the transplanted cells of neoplasms: (1) Immediate growth in the area in which they become arrested; (2) latency for long periods, with or without eventual multiplication; and (3) degeneration and absorption.

The demonstration of this capacity for tumor cells to lie latent has an important bearing upon what is known as the "cell-rest" theory of tumor formation, to which we shall refer later. And from these same data another most important deduction may be drawn: If cells of like order, transported to various areas in the one organism gain growth in some, lie latent in some, and undergo absorption in others, and if, again, portions of the same tumor transplanted into the same tissues in different animals of the same species shows a like succession of results, it is obvious that, whether new-growth occurs or not, is not merely dependent upon the inherent vegetative powers of the transplanted cells, but is also governed by conditions obtaining in the tissue which receives these cells; that, in short, *tissue resistance and the extent* of the same is a factor in determining blastomatosis. To this we have already referred.

The Nature of the Stroma.—There are certain other data and considerations regarding the characters of the blastomas which must be noted: first and foremost, the nature of the organic relationship between these growths and the organism in which they develop. The blastomas gain their nutrition from the organism of the host, and possess both a blood and lymph supply. The capsule of the typical blastomas, as we have pointed out, is formed by the tissues of the host, and not

¹ Olshausen, Ztschr. f. Geb. u. Gyn., 48: 1903.

² Ztschr. f. Aerztl. Fortbildung, 1906: No. 7.

only that, but the stroma of all such tumors must also be regarded as afforded by the host. We must conceive, in short, the tumor as originating by overgrowth of a cell, or cluster of cells, which, as they proliferate, make their way between the connective tissue of the region. Even in what become benign, well-encapsulated tumors, showing expansive or diffuse growth, as suggested by Ribbert's observations upon what appear to be early stages of benign kidney tumors, there appears to be this primary infiltration. Such primary stroma may now, as the tumor expands, exhibit a growth *pari passu* with that of the specific tissue elements of the tumor; this in cases of benign neoplasms. In

Fig. 200

Transplanted carcinoma of mouse. First stages of formation of stroma by host. Site of transplantation after three days: a, growth with mitoses of cancer cells; b, degeneration of introduced stroma. Two zones in surrounding connective tissue: r, amitois in outer zone, d, mitosis of connective-tissue corpuseles next the tumor. $\times A_{+}^{a_{\pm}a_{\pm}}$. (Bashford.)

atypical infiltrative tumors the stroma is continually being added to as the tumor advances into the surrounding tissue, causing degeneration and absorption of the specific cells of the tissue, but leaving the connective-tissue stroma to be the framework of the growing tumor, as has been well demonstrated by Bashford in experimental mouse cancer. Frequently we encounter indications of the same order as those observed in connection with chronic inflammation and tissue regeneration. A tumor which is infiltrating destroys the more highly differentiated cells of a region, while coincidently it may stimulate the more lowly

connective-tissue cells to increased proliferative activity, and this at times, to such an extent that, as seen in certain cancers of a scirrhous type, the abundant stroma that is developed actually chokes or strangles the contained masses of tumor cells.

It is somewhat more difficult to understand the relationship in an atypical connective-tissue tumor (sarcoma). Here evidently a double process occurs; the stroma itself multiplies, and again, as the surrounding tissue is infiltrated, its stroma also becomes part of the stroma of the growth.

Blood and Lymph Vessels.—Such stroma of the host is equivalent to the tissue in which run the vessels and lymph channels. The bloodvessels and lymph channels of the host are retained by the growing tumor; and thus it is that the tumor gains nourishment and discharge of its products.


As the tumor grows there may even be a certain amount of vascular growth, this especially in neoplasms of sarcomatous type. Such vessels never pass beyond the capillary type; they may become distended to great size (and this is true of persisting capillaries), but there is never formation of muscular walls, of arteries and veins proper; nay, more, it is remarkable that what we must regard as arteries and veins enclosed in the growing tumor become simplified and lose their characters. Even at the outer part of an infiltrating growth it is noticeable how few arteries and veins proper are to be detected. So far as we can see, a blastoma has no power of regulating its blood supply.

Nerves. —The host supplies no nerves to the blastoma. Careful study may show a few filaments passing into the peripheral parts of an infitrating tumor,' but these are evidently the nerves of the persisting tissue of the part, and undergo degeneration, for the deeper parts of a blastoma are wholly nerveless. There is, thus, no nervous control, whether vasomotor, or trophic, or of any order on the part of the organism. The activities, vegetative and otherwise, of the neoplasm cannot be influenced by the organism, save through the composition of the blood and fluids supplied by it, and by alterations in the resisting powers of the surrounding tissues. The control, such as it is, is indirect.

Degenerative Changes.—This lack on the part of the tumor to control its own nutrition, and on the part of the organism to govern the tumor cells, renders it not surprising that blastomas are peculiarly apt to exhibit degenerative changes, and, as favoring these, another factor comes in, namely, the absence of any secreting or discharging passage proper over and above the imperfect change afforded by the blood and lymph. The products of the more outwardly placed cells may diffuse into the surrounding tissues; the internal cells are apt to "stew in their own juice," and to be subjected to a form of auto-intoxication. Practically, any of the forms of cell degeneration to be noted in a later section may present themselves, notably necrotic changes, cell death, and absorption. And, favoring these, and favored by them, in all those forms of growth in which degeneration is most liable to occur, we are apt to have hemorrhages, the thin-walled, dilated vessels giving way.

¹ Young, H. H., Jour, of Exp. Med., 2: 1897: 1.

Nuclear Changes.—Associated with these degenerations we meet with nuclear changes. Whereas, the cells of an actively growing tumor, and particularly those of the peripheral portions of actively infiltrating atypical tumors, are noticeable for their relatively large nuclei, with abundant




Irregular mitoses in cancer cells: 1, hypochromatic mitosis; 2, asymmetric mitosis, the upper daughter nucleus hyperchromatic; 3 to 7, various forms of multipolar mitoses. (Galeotti.)

chromatin, so that in the stained section there is a very pronounced difference between them and the nuclei of the surrounding tissue; the nuclei of the more central parts of such tumors are pale, and poor in chromatin (chromatolysis); often they are vesicular; at other times, shrunken and wrinkled looking; and, when degeneration is extensive,

the nuclei of the dead cells do not take the stain at all, while the degenerating cells bordering on such necrotic areas exhibit nuclear fragmentation (karyorrhexis).

These are not the only changes. The more aberrant the growth, the more do we encounter (in freshly removed sections suitably stained) irregular mitoses, and these of a very remarkable order. We owe to von Hansemann,⁴ more particularly, a study of the same in connection with tumors. We may encounter forms with *hypochromatic* mitoses (reduced number of chromosomes below the number normal for the species), asymmetric mitoses (one daughter cell receiving more chromosomes than the other), *hyperchromatic* mitoses (the number of chromosomes in excess, sometimes greatly, of that normal for the species), *multipolar* mitoses (there being more than two centrosomes, each attracting chromosomes)



 mitosis (homotype) in ordinary somatic cell, showing usual form of chromosome; 2, heterotype mitosis in germ cell; 3, irregular heterotype mitosis, with "ring" chromosomes in a cancer cell: at the lower pole aberrant chromosomes passing into the cytoplasm. (Moore.)

so that there may be developed three, four, six, or, according to Hansemann, as many as twelve to twenty daughter nuclei). Again, we may have scattered *chromosomes*, some becoming free in the cytoplasm, as though by rupture of some of the achromatic spindle elements. It must, however, be remembered that these mitotic irregularities are not peculiar to atypical malignant growths. They may be experimentally produced in various tissues in a variety of ways.⁷ They indicate, however, that the cells are subjected to abnormal influences.

Farmer, Moore, and Walker³ have called attention to the existence of⁶ heterotype mitosis" in cancer cells similar to those found, almost specifically, in the stage of maturation of germ cells, and have suggested that in the existence of this type of mitosis and cell is to be found the explanation of the vegetative properties of the blastomata. We believe

- ² Galeotti, Ziegl. Beitr., 20: 1896, and Lubarsch, Allgem. Path., 1905; 44 et seq.
- ³ Proc. Roy. Soc., 72: 1903: 499. See also Bashford and Murray, ibid., 73: 1904: 66.

¹ Die Mikrose, Diagnose d. bosartige Geschwülste, Berlin, 1897.

we are correct in saying that further study has demonstrated that these are found to be only one of a series of mitotic aberrations; that they bear no relationship to the malignancy of the tumor; are not, that is, necessarily present in highly malignant growths and that these observers are not now inclined to lay any stress either upon these appearances,' or on the other appearances recorded of nuclear migration and conjugation.

Retrogression and Healing.-These degenerative changes lead to the consideration of the absorption and disappearance of tumors, and from this to the data bearing upon the active healing of the same. Such absorption and disappearance is the rare exception. A blastoma, once it becomes recognizable, even of the most benign type, may remain stationary, but most often grows; rarely does it recede and undergo natural absorption. And yet every surgeon of large experience can recall one or more cases which he can only explain by such recession. Too often the cases are imperfect and unsatisfactory as evidence; there has been no histological examination of the growth when in its prime, to establish its exact nature. There are, however, cases on record about which there can be no doubt.² Thus, Nasse and Starck have each recorded cases of the spontaneous disappearance of multiple exostoses; Kaposi, of a lymphosarcoma of the upper jaw; Reichel, of a spindle-celled sarcoma of the temple; Nitze, of a papilloma of the bladder; Rotter, of a malignant adenoma of the rectum, involving the vaginal wall, which, after repeated removal and recrudescence, eventually disappeared spontaneously. Shepherd has recorded a case of cervical sarcoma with similar history, and even that most malignant form of growth, the chorio-epithelioma malignum, has been seen to recede by two observers (von Franqué and Fleischmann).

If, thus, there can be natural absorption of tumors even of the most malignant type, sooner or later we must penetrate the secret, and find how, by medical or surgical procedures, to bring about cure, *i. e.*, the degeneration, death, and subsequent absorption of the tumor cells. Undoubtedly this has been already secured in a number of instances, and that by very varied procedures—but the results so far have been very uncertain and most often incomplete. Either all the tumor cells have not been destroyed (and this by the recurrence of the original growth appears to be the most frequent event), or the tendency on the part of the organism to produce new-growths has not been arrested. Probably a more correct statement, which will include both these cases, is that the resisting powers of the organism have not sufficiently exalted to arrest the aberrant cell proliferation. To this matter of resisting power and vegetative power we shall again revert when dealing with the theory of blastomatosis.

Of those methods, apart from operative interference, which have been employed with more or less success, may be mentioned the exhibition of arsenic, Coley's method of inoculation of sterilized culture fluid of mixed streptococcus and B. prodigiosus growths (based on the old experience

¹ Cf. Bashford and Murray, Proc. Roy. Soc. Biol., 77: 1906: 226.

² We here in the main produce those selected by Ribbert, who gives the references.

UNICENTRIC, PLURICENTRIC, AND MULTIPLE TUMORS 693

that intercurrent erysipelas may lead to the absorption of malignant growths), Beatson's method of removal of the ovaries to bring about absorption of mammary cancer, the absorption of uterine myomas following upon electropuncture, and, more recently, the employment of ultraviolet light and the Röntgen rays to cause the disappearance of superficial growths.

Unicentric, Pluricentric, and Multiple Primary Growths.—Lastly, before taking up the subject of the different forms of blastoma, a word must be said regarding the foci of origin.

The majority of primary blastomas are single, and, what is more, appear to originate from a single focus, either a single cell or small collection of cells, separated from the rest of the tissue in which they find themselves. It will be understood that this point cannot be determined. We cannot recognize in the normal tissues a single cell which is destined to give rise to a tumor. By analogy with what occurs in the transplantation of tumors, it would seem that several cells coincidently manifest the aberrant growth. Tumors which appear to grow from a single focus are spoken of as unicentric. Some growths are clearly pluricentric, or multicentric. This would appear to be most often the case in mammary cancer, as has been beautifully demonstrated by Petersen.1 Whereas, in a single section of such a growth the masses of cancer cells in the alveoli appear to be all separate, this is really not the case. By studying serial sections it can be seen that the cell groups form a branching mass, all directly continuous, springing from common centres, the real foci of growth. By making wax models of successive layers, cutting away the parts representing the stroma, and building the successive layers together, Petersen was able to show that the alveoli, or, more correctly, the cell groups, originate from several separate centres.


In the adrenal timor studied by Woolley, to which we have already referred, it was possible to see that cells, clearly belonging to different strands or cell collections of the cortex, were undergoing the cancerous change.² The cells were larger, the nuclei richer in chromatin, so as to stand out in marked contrast to the unaffected cells next to them. There could be no question in this case regarding fœtal cell-rests; it was the cells of the developed tissue that were undergoing the change. We have recently had a second case of the same nature (Fig. 203). Van Heukelom, Horst Oertel, and others have noted a like series of transitions of liver cells into cancer cells in cases of carcinomatosis of the liver (p. 836).

Occasionally we meet not with single or pluricentric single tumors, but what we can only regard as multiple independent primary growths. The commonest example of this is seen in uterine myomas. We may find two, five, ten, to twenty or more isolated muscular tumors in the uterus. In the ovaries, also, cases are on record of as many as five separate dermoids (teratomas) in an ovary, and it is relatively common to find

² Jores had previously recorded a similar observation, Deutsch. med. Woch., 20: 1894: 208.

¹ Virch. Arch., 164: 1901: 570.

coincident dermoids, one in each ovary. With such tumors there can be no question of the one being a secondary, metastatic growth, derived from the other. But the same is true, also, of glandular (adenomatous and eystadenomatous) growths of the ovaries; these show a curious tendency to be bilateral. Here, the growths being of simpler type, it is not always possible to draw a definite conclusion. One might be metastatic from the other, due to tissue predilection. Were it the latter, then



From the edge of a small nodule of newgrowth in the adrenal cortex, showing every transition from a, cells undistinguishable from the surrounding cells of the cortex to small cells with deeply staining nuclei of sarcomatous type.

Similar conversion or modification of cortical cells of adrenal into tumor cells. (Woolley.)

occurring in the same tissue, it should be identical. As a matter of fact, we often find differences in the two growths, which suggest strongly that both are primary. Similarly, multiple fibromas, osteomas, and chondromas are not uncommon, and we approach thus close to a condition which we shall treat separately, that, namely, in which a particular tissue in all parts of the body shows a peculiar tendency toward overgrowth.

Passing a stage farther, there have now been a considerable number of cases reported of multiple primary growths in the one individual of different orders; uterine myomas with uterine or breast cancer; different forms of growth along the digestive tract, and this apart from the transitions which may be found from benign papilloma to carcinoma. The largest collection of these for this literature has been by Walter.¹ Woolley² has analyzed the cases in the literature up to 1903, and Nicholls³

¹ Arch. f. klin. Chir., 53: 1896: 1.

² Boston Med. and Surg. Jour., 148: 1903: 1.

³ Montreal Med. Jour., 32: 1903; 326. See also Wells, Jour. of Pathol., 7: 1901; 357, and Warthin, Jour. of the Amer. Med. Assoc., 32: 1899: 963.

MULTIPLE PRIMARY TUMORS

has collected quite a series from our autopsies in Montreal. In one of Walter's cases there was an angiosarcoma of the stomach, a sarcoma of the gall-bladder, an aberrant adrenal tumor, a lipoma of the kidney, and an enchondroma of the right pleura.

Such cases have been taken by the upholders of the cell-rest theory of blastomatosis to indicate a vice in development whereby several cells or masses of cells become segregated, and liable thus to form foci for subsequent overgrowth. But segregation alone does not explain blastomatosis. A more likely explanation is the lowering not merely of tissue, but of general bodily resistance, so that, simultaneously, cells in various parts find conditions possible for active and independent proliferation.⁴

¹ Of general works upon tumors the fullest and most recent is that by Borst, Die Lehre von den Geschwilsten, 2 vols., Wieshaden, 1902. Ribbert's Die Geschwästlehre, Bonn, 1904, is not so detailed but gives fully the author's much discussed theory. Lubarseh and others give important studies of the recent literature of different orders of tumors in the successive volumes of Lubarseh and Ostertag's invaluable Ergebnisse. It cannot be said that there is any authoritative book on the subject of tumors in our language. Senn's work, valuable from a surgeial point of view, is hurried and ill-digested in its pathology. More important is Bland-Sutton's Tumors, Innocent and Medigmant, which is individual and replete with matter difficult to encounter elsewhere, but its pathology is gross rather than minute.

CHAPTER XVII.

THE AUTOCHTHONOUS BLASTOMAS-(CONTINUED).

CLASSIFICATION OF THE AUTOCHTHONOUS BLASTOMAS.

In what order are we to treat the individual forms of tumors? How are we, that is, to classify them so as to bring together those which are most nearly related, and by its position in the scale gain a grasp of the properties of any particular form?

We have already discussed at length the one main division which for practical purposes is most important, that into typical and atypical blastomas, and the conclusion gained from the study can only be, that while most useful this is not wholly satisfactory because of the existence of (1) transitional forms between the two groups, and (2) apparent or real exceptions to the laws we have noted as, in the main governing either group.

Two courses are open to us: Either, studying these exceptional cases and noting the variation in properties of different forms, we may assume the agnostic position-may say that, despite the enormous amount of material collected, we still have not sufficient data to permit us to make a pronouncement, and, doing this, fall back upon a purely histological and admittedly provisional arrangement based almost entirely upon the characters of the cells constituting the tumors, with no regard to the properties of the individual forms save that which follows from a coincident separation of the typical from the atypical forms. Or, on the other hand, we can start from the basis that the properties of any given form of cell are an inheritance; have been impressed upon that cell by the successive forces to which its ancestry have been subjected; that these inherited properties, along with the forces acting upon the cell itself, determine its characters; so that if we can surely determine the derivation of the different forms of tumors, then an embryogenetic classification must be a natural classification.

The first of these courses is that which from Hamilton¹ (1889) onward has been increasingly adopted, and nowadays it is that employed by von Hansemann.² Lubarsch, Menetrier, Prudden, and the writers of the two most important recent treatises on the subject, Ribbert and Borst. Hansemann goes so far as to state that the only logical course is to take each organ in turn and describe separately the primary tumors which may originate from its component cells, or, in other words, to make as many classes as there are different tumors of different organs or tissues. But this is to construct a Chinese alphabet.

¹ Text-book of Pathology.

² Die bosartige Geschwülste, Berlin, 1897; 22.

CLASSIFICATION OF NEOPLASMS

Borst's classification is:

- Connective-tissue tumors of maturer tissue (so-called benign connective-tissue tumors).
 - (a) Connective-tissue tumors proper.
 - Fibroma, myxoma, lipoma, chondroma, osteoma, angioma.
 - (b) Tumors of the muscle and nervous systems,
 - Myoma, neuroma, glioma.
- 2. Endothelial tumors.
 - Lymphangio-endothelioma, hemangio-endothelioma, and perithelioma, cylindroma, psammoma, cholesteatoma.
- 3. Connective-tissue tumors of immature tissue (sarcoma).
 - (a) Sarcomas of simplest type.
 - (a) Round-celled, spindle-celled, giant-celled.
 - (b) The more highly developed sarcomas
 - Mixed sarcomas (fibroma sarcomatosum, osteoma sarcomatosum, etc.), melanosarcoma, chloroma, lymphoma sarcomatosum, myeloma multiplex, angioma sarcomatosum, myoma, neuroma, glioma sarcomatosum.
- 4. Epithelial tumors.

(a) Of mature type.


- Papilloma, adenoma, cystadenoma.
- (b) Of immature cell type (carcinoma).
 - Of skin, squamous epithelioma; of mucous membrane, cylindrical-celled cancer; of glands, carcinoma adenomatosum. Appendix. Adrenal tumors: Chorionic tumors.
- 4. Mixed tumors.
 - (a) Cystic mixed tumors.
 - Dermoid cysts of skin, testes, ovaries, branchiogenic cysts, ciliated epithelial cysts of brain, enterocysts.
 - (b) Mixed tumors in the narrower sense.
 - Of kidneys, vagina, bladder, testes, mamma, face.
 - (c) Teratoids and teratomata.
 - Of testes and ovaries; of anterior and posterior ends of body axis; bigerminal sacral teratoma, monogerminal sacral teratoid, teratoids and teratomas of the body cavities, teratoids and teratomas of neck, cranium, and ventricles.

If we analyze this we find that it is constructed on the principle of recognizing three groups of tissues, the connective, the endothelial, and the epithelial, of which the first two afford atypical tumors of like order (sarcoma); the last affords the carcinoma. It is interesting to see how close a carefully thought-out classification constructed purely on these histological principles brings us to the embryogenetic classification to be presently noted. There are it will be seen certain "jumble" departments; the myoma and the glioma have little in common; the adrenal tumors and the chorionic tumors have to be treated as an appendix, an admission of doubt as to their exact place in the scheme; simple epithelial cysts and the complicated ovarian and testicular teratomas come into the same section. But, on the whole, the teratomas and the teratoblastomas (mixed tumors) range themselves very much according to the classification we have already afforded from embryogenetic considerations.

In this connection may be mentioned a suggestive grouping of tumors proposed by Lubarsch, who would primarily divide them into three main groups:

 Those departing but slightly from the type of mother tissue and showing little or but temporary growth (teratoma, congenital nevus, congenital adenoma, myoma, lipoma, osteoma, chondroma). In all of these cases we have probably to deal with a local transposition of tissue.

2. Those which while showing autonomy still comply with the ordinary rules of life and respect physiological limits (the larger myomas, adenomas, angiomas, etc.; these may for long remain in a resting state, with periodical accessions of growth and absorption).



Cross-section of a human embryo of 1.54 mm.; ek, epiblast (ectoderm); en, hypoblast (endoderm); me, mesoblast (mesoderm); I_i , dorsal furrow, giving origin to nerve cord and brain; eh, region of hypoblast that gives rise to the notochord; g_i junction of the extra-embryonic sonatic mesoblast (eL) and splanchnic mesoblast (dL); p_i beginning of the embryonic excloss. (Graf von Spec.)

3. Tumors fully emancipated from physiological laws (malignant tumors proper, sarcoma, and carcinoma).

While this division is suggestive, and valuable as calling our attention to the properties of different orders of tumors, it is not a classification of the different forms of tumors in the proper sense. Growths of the same type occur in more than one class; an adenoma may belong to all three groups; a congenital mole may assume malignant properties and pass from the first to the third group; a tumor due to local transposition of tissue, for instance, the aberrant suprarenal growths in the kidney, may, while congenital, be fully emancipated from physiological laws and show malignancy even before birth.

Embryogentic Classification (Waldeyer).—With the development of the science of embryology it was noted that from the primitive germ layers

HISTOGENETIC CLASSIFICATION

different tissues were derived; that the connective tissues of the body, including bone, cartilage, and muscle, were of mesoblastic origin, while broadly (and, as I shall point out later, incorrectly) the specific cells of the epithelia and the acini of glands were seen to be derived from either epiblast or hypoblast; and as neoplasms originate from preëxisting tissues or their precursors, Waldeyer introduced the division of tumors into those of epiblastic or hypoblastic and those of mesoblastic origin, subdividing according to the nature of the tissue, and again according as to whether the arrangement of the component cells was typical or atypical. A further class had to be made for the mixed tumors, those, namely, containing overgrowths of both epi- (or hypo-) blastic and mesoblastic elements.

It is unnecessary that we here give the full classification according to this scheme. Such classification was popular during the last quarter of last century, but even those who used it recognized its defects. Of these, the greatest was that it ignored the fact that the mesoblast also gives rise to definite glandular organs; another, that it separated the gliomas (tumors derived from the neuroglia and so of epiblastic origin) from the sarcomas or atypical connective-tissue tumors, of mesoblastic origin, although histologically the growths are closely related, the glioma showing no close relationship to the carcinomas or atypical glandular and epithelial tumors. These were grave defects, and their recognition it was that led to the reaction indicated by the present attitude of most modern writers.

Histogenetic Classification.—Now, the principle underlying the above attempt at classification was a right one. Just as the form and structure of the individual of any species is the outcome of the phylogeny of that species, is the resultant of the special conditions to which that individual and its progenitors have been exposed in the course of countless generations, so the component tissues of the individual with their special characters are the resultant of both past and present forces.

A given cell of the embryo in a given relationship to the rest of the embryo has inherent tendencies to give origin to cells of a particular order. The weakness of Waldeyer's classification lay in this, that it was based upon an inadequate embryology. It does not follow that a fuller and more accurate knowledge of histogenesis will not afford us valuable aid. Each tissue has a definite origin and mode of development, and if neoplasms are derived from definite tissues, and their component cells represent stages in the development or degeneration of these tissues, then it is possible to establish a rational classification of tumors upon histogenetic lines.¹ We have to start from the very earliest stage of the developing ovum to gain a proper grasp. The earliest stage to be recognized in the development of the fertilized ovum, once it has proceeded to segment, is the *morula*, in which the blastomeres or cells form a cluster or group of cells of the same order with almost complete lack of differentiation.

⁴ I here follow very largely my article Upon the Classification of Tumors, Jour, of Pathology, 4:1902;243.

Rapidly this gives place to a second stage, in which these cells arrange themselves into two layers, into the primordial layers of *epiblast* and *hypoblast*. In this way at a singularly early stage the future epiderm and endoderm become recognizable. The next stage to be noted is that the hypoblast, or more internal of the two primitive layers, gives rise by proliferation of its cells to a group or mass of cells showing no definite arrangement among themselves and not forming a true layer. This is the *mesoblast* and *Anlage*² of the organs derived from that layer. The hypoblast, while it in the main gives origin to these cells, still remains as a distinct layer or membrane. The epiblast participates to a less extent. Waldeyer went so far as to recognize these three layers, but there he stopped.

The reader must dispel as erroneous the old deeply rooted idea that connective tissues, and connective tissues only, arise from mesoblast; epithelia and glandular tissues and nerves, and these alone, from epiblast and hypoblast.

From the epiblast, whose cells in general are from the earliest period arranged in regular order so as to form a definite layer, there is developed, along the dorsal groove, a marked proliferation of the cells, those away from the surface being no longer arranged in strata. Indeed, it is legitimate to compare this development of the *neuroblast* or anlage of the nervous system with the earlier development of the mesoblast. With the further infolding of the dorsal groove this portion of the original epiblast becomes cut off from the rest, the only portion recalling the original epiblast being the ependymal cell layer immediately around the central canal, the cells or descendants of cells which have originally been the outer layers of the dorsal epiblast. A very similar ingrowth of cells, irregularly arranged, occurs from the hypoblast to form the basis of the *notochord*.


The mesoblast in its turn undergoes changes: with the development of the primitive body cavity, or ceelom, those cells abutting in that cavity become arranged as one orderly layer, the mesothelium, the remaining portion of this "layer" not thus arranged constituting the mesochyme. From the mesothelium again, by a process of active growth and heaping up of cells, are developed localized masses of cells on either side, which we may compare with the neuroblast and notochord; these are the myotomes, the anlagen of the future striated muscles of the body, and later from the mesonchyme a final true layer is developed, the endothelium, lining the vascular cavities, both blood and lymph vascular.

We thus find that the embryo comes to exhibit cell collections of two orders, which may be termed "lining membranes" and (for lack of a more expressive word) "pulps," the lining membranes being the persistent epiblastic, hypoblastic, mesothelial, and endothelial layers, the "pulps" being the main mass of the neuroblast (of epiblastic origin), the notochord (of hypoblastic), and the mesonchyme (of mesoblastic).

¹ We have no word equivalent to the German Anlage—hence must employ it to indicate the primordial source, or developmental origin of any organ or tissue.

LEPIDIC AND HYLIC TISSUES

And now, following up the further development of these different cell collections, we observe that the adult tissues derived from these two series exhibit well-marked differences, so that we can divide adult tissues into two great groups, the *lepidic* (from $\lambda z \pi z$, $\lambda z \pi z \partial z$, a rind, skin, or membrane) and the *hylic* ($\lambda z z$, crude undifferentiated material).



Diagrammatic representation of section through vertebrate body to show ontogenetic relationship of the various order of tissues. A. Of lepidic type: 1. epiderm and its glands (epiblastic); 2, mucous membrane of digestive canal and its glands, liver, etc. (hypoblastic); 3, endothelium lining serous cavities (mesohlastic) and glands like renal cortex of mesothelial origi; 4, vascular endothelium of late mesohlastic origin. R. Of bylic type: 5, spinal cord, brain, and nerves (epiblastic); 6, notochord (hypoblastic); 7, connective tissues of the body (mesenchymatous); 8, myotomes, striated muscle of body (mesothelial). C. Cavities: 9, lumen of digestive tabe; 10, body eavity.

The characteristic of the lepidic tissues is that the specific cells which give them their main features are arranged either in layers or clusters in direct apposition; they are not separated by lymph spaces or by bloodvessels; they possess, nevertheless, a supporting framework or stroma of hylic tissue in which run the nutrient vessels. Of hylic tissues the features are the opposite: separating the cells there is a matrix of intercellular

substance either homogeneous or fibrillated, while *lymph spaces and blood* capillaries tend to separate and run between the individual cells. If in the lepidie tissues there is a stroma of hylic tissues, so here in the hylic there always enters lepidic tissue in the shape of the living endothelium of the blood and lymph vessels. In either case the elements of the other order occupy a subordinate position. While some pathologists, like O. Israel¹ and Buxton,² have already noticed this distinction, the histologists and embryologists have laid little stress upon it. The more we study tumors the more we realize this importance of the distinction.

On this basis we obtain the following classification of normal tissues:

1. LINING MEMBRANE OR LEPIDIC TISSUES,

in which the bloodvessels do not penetrate the groups of specific cells and in which there is an absence of definite stroma between the individual cells, although such stroma, of mesenchymatous origin, may be present between the groups of cells.

1. Epiblastic:

Epidermis. Epidermal appendages of hair, nails, enamel of teeth, etc. Epidermal glands. Epithelium of the mouth and salivary glands. Epithelium and glands of nasal tract and associated spaces. Epidermal (anterior) portion of hypophysis cerebri. Lens of eye. Epithelium of membranous labyrinth of ear, anus, male urethra (except prostatic portion).

2. Hypoblastic:

Epithelium of digestive tract and glands connected with it. Specific cells of liver, pancreas, tonsils, thymus, thyroid. Epithelium of trachea, lungs, bladder, female urethra, male urethra (prostatic portion).

3. Mesothelial:

Lining cells of pleure, pericardium, peritoneum. Specific cells of suprarenals, kidneys, testes, ovaries (Graafian follicles). Epithelium and glands of Fallopian tubes, uterus, vagina, vasa deferentia, vesiculæ seminales, etc.

4. Endothelial:

Lining endothelium of bloodvessels and lymphatics.

II. HYLIC OR PRIMITIVE PULP TISSUES.

Organs and tissues in which the special characteristic is that the specific cells lie in, and are separated by, a definite stroma, homogeneous or fibriltar, in which there may or may not be blood and lymph vessels.

1. Epiblastic:

Nerve cells, neuroglia.

2. Hypoblastic

Notochord.

¹ Berl, klin, Woch., 37; 1900; 609, 644, and 667.

² Jour, Cutan, and Genito-urin. Dis., New York, February and April, 1901.

3. Mesenchymatous:

Fibrous connective tissues, cartilage, bone, reticulum of lymph glands, bone marrow, fat cells, involuntary muscle tissue, spleen, bloodvessels, blood corpuscles.

4. Mesothelial:

Striated muscle, including cardiac muscle.

Following this scheme of classification of the normal tissues, we may now divide the tumors arising from the specific constituent cells of the various tissues into two main genera—the *lepidic* tumors, or *lepidomas*, originating from the above "lining membrane" tissues and the hylic tumors (*hylomas*), originating from tissues derived from the embryonic "pulp." We can further distinguish two broad groups of lepidic tumors, the *primary*, those whose cells are derived in direct descent from the original epiblast and hypoblast; and *secondary*, or *transitional*, whose cells are derived in indirect descent from the same, *i. e.*, have, in the course of development, passed through a mesoblastic or mesenchymatous stage before coming to form portions of a lining membrane. We shall explain the use of the term *transitional* later.

L LEPIDIC, OR RIND TUMORS.

(.1) Lepidomas of the First Order.

1. Of epiblastic origin.

- Tumors whose characteristic constituents are overgrowths of tissues derived directly from the epiblastic lining membranes, or epiderm.
- (a) Typical.—Papilloma, epidermal adenomas (of sweat, salivary, sebaceous, and mammary glands, etc.).
- (b) Atypical.—Squamous epithelioma, carcinoma of glands of epiblastic origin.

2. Of hypoblastic origin.

- (a) Typical. –Adenoma and papilloma of digestive and respiratory tracts, thyroid, pancreas, liver, bladder, etc.
- (b) Atypical.—Carcinoma developing in the same organs and regions.

(B) Lepidomas of the Second Order, or Transitional Lepidomas.

3. Of mesothelial origin.

- Tumors (mesotheliomas) whose characteristic constituents are cells derived in direct descent from the persistent mesothelium of the embryo.
- (a) Typical.—Adenoma of kidney, testicle, ovary, urogenital ducts; adenoma of uterus and prostate; adenomas originating from the serous membranes, "mesothelioma" of pleura, peritoneum, etc.
- (b) Atypical.—Cancer of the above-mentioned organs; squamous endothelioma, so called, of serous surfaces, epithelioma of vagina; adrenal mesotheliomas, "hypernephromas."

4. Endothelial Lepidomas.

Tumors originating from the endothelium of the blood and lymph vessels; lymphangio-endothelioma, hemangio-endothelioma, perithelioma, the commoner forms of cylindroma, psammoma, cholesteatoma of brain (but not of ear).

II. HYLIC, OR "PULP" TUMORS.

1. Of epiblastic origin.

- Tumors whose characteristic constituents are overgrowths of tissues derived from the embryonic pulp of epiblastic origin.
 - (a) Typical.—True neuroma, glioma.
 - (b) Atypical.—Gliosarcoma, sympathetic neurocytoma.

2. Of hypoblastic origin.

Tumors derived similarly from embryonic pulp of hypoblastic origin.

Chordoma.

3. Of mesenchymal origin.

- (.1) Mesenchymal Hylomas.—Derived from tissues originating from the persistent mesoblastic pulp, or mesenchyme.
 - (a) Typical.—Fibroma, lipoma, chondroma, osteoma, myxoma, leiomyoma, angioma, myeloma.
 - (b) Atypical.—Sarcoma (derived from mesenchymatous tissues), with its various subdivisions, fibrosarcoma, spindle-celled sarcoma, oat-shaped-celled sarcoma, chondrosarcoma, osteosarcoma, myxosarcoma, lymphosarcoma, chloroma, angiosarcoma; of origin still debated, melanosarcoma.
- (B) Mesothelial Hylomas.—Tumors which are overgrowths similarly of tissues derived from embryonic pulp of definitely mesothelial origin. Rhabdomyoma.

If this classification be studied, it will be seen that we have done away with that deficiency in the earlier embryological classifications, whereby tumors of unlike orders and histological appearances were grouped together, and those of like characters separated. Gliomas, for example, come to be placed close to the mesenchymatous tissues; the glandlike tumors of mesoblastic origin become grouped along with those of epiblastic and hypoblastic origin, while tumors of the same type, from whichever laver they may originate, are grouped together.

Have we, in accomplishing this, introduced any new difficulties? One objection will undoubtedly present itself, namely, that among the lepidomas of mesothelial origin we have grouped together tumors some of which are of a strongly epithelial or glandular type; for example, the cancers of the uterus, with others like the hypernephromas, tend to take a definitely sarcomatous character. I fully admit this difference in properties.

Two possibilities exist: either the neoplastic properties of the different portions of a given germ layer become differentiated, according to the ultimate function assumed by those portions—to admit which is, we

TRANSITIONAL LEPIDOMAS

confess, tantamount to acknowledge that little weight can be attached to embryogenetic considerations; or that the epithelium lining certain organs in which we find tumors not of the characteristic transitional type, is not mesothelial; that, for example, whereas primarily the vagina, uterus, and Fallopian tubes originated from Müller's duct, and so were of mesothelial origin, in the course of development the cloacal hypoblast has overgrown and replaced the original mesothelial lining of uterus and tubes, the epiderm from without has grown into and replaced the mesothelium of the vagina and cervix uteri. This has indeed been suggested by more than one embryologist. Certainly the characters of the vaginal epithelium are unlike those of any other mesothelial structure, and primary vaginal tumors are of an epiblastic, and not a mesothelial, character; while, similarly, the mucous membrane of uterus and tubes strongly recalls that of the alimentary tract, as do the tumors arising from the same.




Transition from adenomatous to sarcomatous type of growth in a renal mesothelioma, (Birch-Hirschfeld.)

With this admission, and, it may be, only apparent exception, the striking feature of these secondary lepidic tumors, as a class, *is their liability* to present transitional characters—and this in the lack of recognition of the underlying cause has created an appalling amount of confusion. A tumor of the adrenal, a "hypernephroma" of the kidney, a testicular neoplasm or ovarian growth, and the same is true of the whole class of endotheliomata, may, if of slow growth, present all the characters of a cancer—a glandular tumor—if actively vegetative be indistinguishable from a sarcoma; and frequently in this group we meet with intermediate types, in which one part of a growth shows the cancerous, lining membrane type of structure, and other parts have taken on the hylic, sarcomatous type. Such tumors form an important proportion

of the cases of so-called *carcinoma surcomatodes*. Nay, more, in such a tumor, as Woolley,¹ from our laboratory, has pointed out: using Mallory's connective-tissue stain, so as to follow accurately the ramifications of the stroma, the transition from the lepidic to the hylic type is found to be not apparent, but actual; certain cell clusters, as in cancer proper, lie wholly free from any intervening stroma; others, on the contrary, are separated and isolated by a matrix, which contains connective-tissue fibrils, a stroma proper, such as we find in surcoma.⁷

And, what appears to be an adequate reason for this difference in properties, suggests itself. As we have emphasized more than once, properties which are of oldest acquirement are those which are last to be lost; those of later acquirement are yielded up with greater case.



Section of carcinoma simplex of breast, treated with Mallory's connective-tissue stain, to demonstrate a complete absence of passage of intercellular fibrils between the individual members of the alveolar cell groups. (Woolley.)

Section of an endothelioma similarly treated. The alveolus below reacts almost wholly like an epithelial cancer, that above exhibits intercellular connective-tissue fibrils, like a streoma. (Woolley.)

The primary lepidic tumors are derived in direct descent from cells which, from the earliest embryonic period, have taken on lepidic, or lining-membrane characters; whereas, these transitional tumors one, and all, are derived from cells which, from being lepidic (in hypoblast or epiblast), have become hylic, and only at a later embryonic period have again taken on lepidic characters. Such cells in new-growth revert

¹ Johns Hopkins Hospital Bulletin, 14:1903:21.

² This has since been confirmed as between sareoma and earchnoma by Huruzo-Kuru (Verhandl, D., path. Gesell, 13; 1909; 386) who finds that the lattice ("Gitter") fibrils of the stroma revealed by Maresch's silver method do not invade the elusters of cancer cells.

more easily to the hylic, sarcomatous type, than do the cells of the primary lepidic tumors. Here, indeed, histogenetic considerations show themselves of singular value in clearing up one of the enigmas and great difficulties in the study of tumors.¹

In laying this down I do not coincidently imply that primary lepidic tumors, under these conditions, never manifest the same tendency to reversion or conversion to a hylic type. It is, I know, the general impression and the common teaching that epiblastic and hypoblastic "rind" tumors, squamous epitheliomas, glandular cancers of the mamma and digestive tract, for instance, are always typically cancerous. This is not so; one has but to study the advancing edge of a highly malignant, rapidly growing epithelioma to see that here and there individual cells, of epithelial type, become surrounded by (or probably directly make their way into) the connective tissue; while still farther out from the main mass of the growth it is impossible to say whether the largest cells seen are of epithelial or connective-tissue origin. And more recent studies of what Krompecher has termed "basal-celled" cancers, have established, it would seem beyond any doubt, that cells of epidermal, epiblastic origin can give origin to tumors indistinguishable from connective-tissue sarcomas in histological structure. What we would say is, that such reversion is so frequent as to be a distinguishing feature of the secondary lepidic tumors; it is the exception in the case of the primary.

I am strongly adverse to the coinage of new terms in our subject, but, at times, when a new idea or new relationship has to be expressed. such coinage becomes essential, and this was the case when I suggested lepidic and lepidoma, hylic and hyloma, respectively. They were necessary for the expression of my conception of tumor relationships. Whether others will find them so useful, not to say essential, as I have found them, time must tell. At present I regard them as a framework around which to group ideas, and do not suggest their employmentin fact-personally never employ them for daily clinical purposes. For such, the names of the different typical tumors and the terms carcinoma, sarcoma, and endothelioma are adequate. As will have been gathered, it is obvious that the terms carcinoma and sarcoma must be given apurely morphological significance. It is impossible nowadays to attach to them any histogenetic significance, once we recognize that tumors of identical type, hylic or lepidic, may originate from any of the germ layers. Here we find ourselves wholly in accord with Lubarsch,2 and strongly urge that his recommendation be put into general practice: "So I come back to this, that a combination of morphological and histogenetic mode of nomenclature is necessary. The chief word must be determined by the morphological structure; if we can with certainty give

¹ We do not claim credit, save in establishing this as a general principle explaining the feature of secondary lepidic tumors as a body. For O. Israel had already recognized fully this same dependence of the characters of the endotheliomas upon the embryogeny of the mother tissues.

² Ergebnisse, 6:1900:968,

the genesis, then indicate that by an adjective, as, for example, endothelial adenoma, epithelial adenoma, etc." To these examples we would add, as further indications of the method, osteosarcoma, cutaneous melanoma, choroidal sarcoma, mesothelial cancer. For practical purposes, the binomial and trinomial method is essential; there are marked differences in the malignancy of endothelial and epithelial growths; thus, to label both cancer is to afford no information, or to mislead the surgeon or clinician.

This, it may well be repeated, we note in all tumors, that the more rapid the growth, and the more the cells depart from their normal and mature environment, the more do we observe that those features of the tumor cells which are specific for one or other tissue tend to disappear. In the most rapidly growing and most aberrant tumors the individual cells, afford us little or no clue to the tissue of origin. It is the general arrangement of the cells that aids us in making our diagnosis,



"Pseudo-epithelium," or secondary epithelium without busement membrane lining a cyst in a glioma, formed by modification of the superficial layer of glioma cells. (Saxer.)

and even then the general arrangement is not so much that peculiar to the fully formed tissue as that common to connective tissue in general, or to glandular and lepidic tissues in general. We recognize a reversion to an earlier, simpler, or, as it is often expressed, a more embryonic type. The essential feature of the cell of the atypical tumor is the more or less complete replacement of functional by vegetative or proliferative activity, and the consequent loss of those features directly associated with the performance of function.

Lastly, as bearing upon the subject of classification, it may be asked, Can cells which, with neoplastic proliferation, have lost specific functional properties regain them? The answer to this must be that everything indicates that the power of reacquirement is minimal. A hylic tumor cannot take on lepidic characters. At most, modified relationships may bring about modification in properties, but this must be regarded as an adaptation, an assumption of new properties, not, it seems

ANAPLASIA

to us, an awakening into activity of properties which we would regard not as merely dormant, but actually lost. Here we may be mistaken, but it is thus we would explain Saxer's' case of the eventual clothing of degeneration cysts in gliomas with an imperfect layer of glial cells taking on epithelial characters; those cells do not form a true epithelium, and become cut off from their fellows; no basement membrane is formed, and we find every transition from the typical glioma cell to cells which, lying in apposition to the fluid of the cyst, take on a more epithelioid type; now there is a single layer of such cells, now two or three layers. As Marchand has pointed out, the difference between an embryonic cell proper and a vegetative tumor cell is that the former has the potentiality, given favorable environment, to undergo full differentiation in a particular direction: the latter in becoming a tumor cell has lost that potentiality: under the most favorable conditions it can only develop up to a certain stage. This, indeed, is the meaning of von Hansemann's term anaplasia, a reversion or loss of power of full development. At most, some of the powers of a tumor cell may lie latent. Thus, Ehrlich and Apolant have noted that an adenocarcinoma of the mouse which in the course of numerous transplantations had taken on a more and more atypical and cancerous appearance, in some later passages tended to revert to the less atypical and more adenomaous appearance. Nevertheless, it still possessed all the earmarks of a malignant tumor.

¹ Ziegler's Beitr., 38: 1905.

CHAPTER XVIII.

TYPICAL HYLIC TUMORS OF MESENCHYMATOUS ORIGIN.

BENIGN "CONNECTIVE-TISSUE" TUMORS.

IT would, perhaps, seem natural to discuss now the causation of neoplasia or blastomatosis. But, without a knowledge of the mode of recurrence and properties of the different forms of growth, it is difficult to treat this most difficult subject in a satisfactory manner, or to grasp the relative importance of the different arguments brought forward. To prevent undue digression and repetition, it is better first to pass in review the various forms, thereby forming a basis for our treatment of causation. In so doing it will be better, also, not to follow slavishly the order of the classification just given, but to consider first the simpler hylic or connective-tissue tumors, and later the more complicated lepidic and glandular forms. And here, following the example of descriptive biologists, it will be well to describe type forms first; as, also, to call attention to certain departures from type-certain impure blastomas, if we may so describe them-forms which do not conform in all respects with the definition of blastomas in general, and, indeed, possess different properties. It would be better to consider these as a class apart, and this we may be able to do in the future; at present it is so much the custom to include them under the same heading, that to divorce them absolutely would confuse the student consulting other works on the subiect. Thus, where necessary, we shall call attention to these examples of blastomatoid growth. Indeed, the frequent notes of the existence of these conditions may be of more service to calling attention to the difference than would a special section devoted to the subject.

FIBROMA.

As its name implies, the fibroma is a tumor composed of fibrous connective tissue, and as such connective tissue is peculiarly widely distributed, so tumors of this nature may be met with in all regions of the body, although, as will be pointed out, there are certain regions and tissues in which these tumors are especially apt to develop. And as ordinary connective tissue varies in its composition, being in some regions loose and areolar, with loose bundles of fibrils and relatively frequent cells, being in others dense and firm, with abundant fibrillar substance and relatively few cells, and those much compressed, so, largely according to the seat of origin, do we meet with fibromas of

FIBROMA

different density of formation. Thus, we are accustomed to distinguish soft and hard fibromas respectively, the latter more particularly developing from connective tissues of a dense type, as, for example, from tendons, fasciae, and periosteum; the former from looser, more areolar tissue, e. q_{\star} , subcutaneous connective tissue.

Wherever growing, the fibroma has for its essential and predominant constituent connective-tissue elements. As such, it is composed of connective-tissue cells, bands of white connective-tissue fibrils, bloodvessels, and, to a greater or less extent, elastic fibres. Nay more, as pointed out by Mallory,' just as in normal connective tissue two orders of fibrils are recognizable, so are these present in fibromas, forming thus a means of diagnosis. These are (a) the *fibroqlia* fibrils running along



Fibrosarcoma: cells viewed flat-wise; fibroglia fibrils black; wavy collagen fibrils barely visible, (Mallory,)

the surface of the cytoplasm in the direction of the long axis of the cell, passing from one cell to the next, and (b) the fine wavy collagen fibrils lying alongside of the cells but not attached. In addition, elastic fibrils may be present. The fibroglia fibrils are most abundant in relatively slowly growing spindle-cell sarcomas (fibrosarcomas), the collagen or ordinary wavy connective-tissue fibrils form the bulk of the tumor in fibromas proper. Lymph spaces and channels are also present, few and inconspicuous in the hard variety, frequent and large in the softer forms.

Typically, such growth forms a well-defined nodule, which, as it enlarges, leads to the atrophy, absorption, and replacement of the tissues immediately surrounding it. Growth is slow and expansive.

¹ Journ, of Med. Research, 10; 1903; 334 and 13; 1905; 113 (where are given the methods for staining the fibrils of different orders of tissue). See also Journ, of Exp. Med., 10; 1908; 575.

TYPICAL HYLIC TUMORS

Where rapid, there histological examination shows the existence of abundant cells, not of the typical, fully formed, connective-tissue type, but resembling fibroblasts, and, like them, flattened, possessing deeply staining nuclei of fair size and a relatively abundant protoplasm.

We have here the vegetative type of connective-tissue cell. The existence of great numbers of these fibroblasts, or spindle cells, indicates a transition to the sarcomatous condition and the assumption of more malignant characters. When the cellular character is prominent, we speak of a *fibrosarcoma*, herein, with Mallory, including *spindle-celled* sarcomas.

It must be borne in mind that all fibromas are more cellular than normal adult connective tissue. It is when this fibroblastic overgrowth is a striking feature, and particularly where it is marked in one or more areas of the tumor, that we are justified in speaking of fibrosarcoma. Growth in all cases is from such fibroblasts, and not from fully formed connective-tissue cells.

All typical fibromas are pale on section, and those of the firmer type are glistening, the light glinting from the cut surface somewhat as from watered silk. This is due to the fact that the fibrils run in bands, the various bands being cut in different directions. This structure is to be explained by the development of the connective tissue, which occurs in the main around the bloodvessels in the tumor; newly formed fibrils are laid down roughly parallel and concentric to these vessels, and as the vessels course in various directions, so do these bands of fibrils. The tumors are readily enucleated, and, in most cases, the border of the tumor is to the naked eye sharply circumscribed. It will, however, be easily understood that both the more concentrated tissue immediately around the tumors and the tumors themselves being composed of fibrous tissue, there is, under the microscope, no sharp outline to be distinguished between the two; the neoplastic and the surrounding non-neoplastic tissue appear, under the microscope, to pass into one another.

Degenerative Changes.—Fibromas of long standing are apt to exhibit degenerative changes; through arrest of the blood supply, by tension, or other cause, they may undergo necrosis, with abundant formation of cholesterin and fatty debris, or they may become so in-filtrated with calcareous salts as to be converted into calcareous nodules. Bony and cartilaginous masses have been noted in some cases of old standing; it is not always easy to decide whether we have to deal with original osteoid or cartilaginous inclusions, or with metaplasia induced by modified nutrition and cell relationships. Through obstructed lymph discharge, tumors may be found edematous, or lymphangiectatic, cystic or mucoid (*fibroma mucinosum*). Such conditions must be distinguished from the conversion of areas of a fibroma into definite myxomatous tissue (p. 719) when we deal with a *myzofibroma*.

Occasionally, in the kidney and elsewhere, we accidentally encounter what appears to be the earliest stage of fibromatous growth; small collections of proliferative fibroblasts, which are infiltrating the immediately surrounding tissue. At a later stage, it appears that infiltration

ceases, and the surrounding tissues are pushed aside by the expansive diffuse growth of the tumor, which thus gains a capsule and becomes sharply defined. In this way it would seem to be that occasionally we encounter in what is otherwise a pure fibroma rare glandular acini, and, it may be, other tissue elements. So, also, it is possible that in adult life what had been at first purely an interstitial inflammatory fibrosis in a gland, such as the mammary gland, takes on, in parts, active tumor growth; and, in like manner, there become developed isolated tumor masses which contain glandular or other elements. So long as these included tissues show no sign of independent proliferation, all such apparently mixed tumors should still be referred to as fibromas, as fibromas with inclusions of one or other order; only when there is coincident aberrant growth of the other element, along with evidence of like growth of the interstitial fibrous tissues, is it permissible to speak of fibro-adenoma, osteofibroma, etc. This rule, unfortunately, is more honored in the breach than in the observance; indeed, the majority of the fibro-adenomas, of which the commonest example is offered in the mammary gland, are not fibromas in the true sense. The fibroid overgrowth is not limited and sharply defined. It passes diffusely into the surrounding tissue; it is, at most, *fibromatoid*, and is to be considered along with the blastomatoid conditions, to be presently noted.

Fibromata proper do not form metastases, and, similarly, it may be laid down that they do not recur. If recurrence does happen, either the primary tumor, on examination, is found to exhibit fibrosarconatous changes, or we are dealing with a fibromatoid condition, *i. e.*, the original tumor was not a sharply defined, limited mass, but possessed a definite root or base, through which it passed imperceptibly into the surrounding connective tissue, recurrence being thus the manifestation of a tendency toward diffuse regional overgrowth on the part of the surrounding tissue, which, it may be, has been stimulated by the removal of the primary growth (see p. 714). This would appear to be the most satisfactory explanation of the recurrence of fibroid, fibromyxomatous, and other nasal polypi.

Hard Fibromas.—Hard fibromas, as isolated nodular growths, occur more especially in connection with tendons. While this is most often the case, it is not the absolute rule, for occasionally we meet with soft fibromas in fasciae, and those developing in "soft" tissues, such as the kidney, may be hard. Most often there is no history of previous injury or irritation. Sometimes, as in the mammary gland, we encounter the hard, well-defined variety, and here there may be a history of previous inflammation.

Another variety of hard fibroma develops in connection with the jaws, the characteristic *epulis*, a term properly applied only to these fibromatous growths, but often given to osteoid and osteosarcomatous growths. These develop from the periosteum, and, according to Bland-Sutton, originate always in connection with the root of a decayed tooth. In their growth they cause absorption and replacement of the bone. Fibromas of the uterus will be discussed along with the myomas of that organ. **Soft Fibromas.**—These may be single, but frequently are multiple. Upon analysis of the cases, it appears that the majority come under the fibromatoid growths, to be presently noted. They occur more




Hard fibroma. (Ribbert,)

especially in connection with the skin and submucosa of the pharynx and digestive tract. Those in connection with the nose and throat are peculiarly soft—mucoid polyps—and of the true fibromyxoma type. Here and in the nasal region there may be inclusions of mucous glands.

Fibromatoid Growths.—In the group of blastomatoid growths must be placed a series of conditions intermediate between simple hypertrophy and true tumor formation. Failure to recognize their peculiar properties have frequently introduced vagueness into the treatment of

simple hypertrophy on the one hand, true tumor formation on the other. These are conditions which Klebs and other German writers have classified as one form of Riesenwuchs (giant growth), which, also, C. P. White has recognized as "progressive hypertrophy." Such growths as a class (1) affect one particular tissue; (2) are multiple; (3) of congenital if not hereditary origin, frequently manifest in early life, and affecting several members of a family; (4) may be diffuse, or if

Fig. 213



Soft fibroma,

not diffuse, show no demarcation from the surrounding unaltered tissue, merging into this imperceptibly; (5) the apparent encapsulation which such growths may exhibit on one or more aspects is due to their strictly respecting the limits of the part in which they find themselves, and represents those limits; (6) they are of very slow growth, extending over years; (7) eventually, they may take on sarcomatous characters, but this is an epiphenomenon; it is but in accordance with the principle that tissue

FIBROMATOID

which has developed in excess of function is liable to take on aberrant growth.

As already noted, we regard the submucous connective tissue of the posterior nares and pharynx as a favorite seat for this fibromatoid development. In the previous edition, following von Recklinghausen, we included here the remarkable condition of multiple neurofibromators,

More recent studies, as will be pointed out in due season, necessitate that this be regarded as a particular form of blastomatoid affection of the



Fig. 214

Multiple fibromatoid overgrowths along the course of the cutaneous nerves. (Herczel.)

medullated peripheral nerves. It is true that, in not a few cases of the multiple tumors of this order, in some, or many, of the individual growths there is a preponderance of true fibrous connective-tissue elements, but this is comparable with the preponderance of fibrous tissue in scirrhous cancers, and if in the latter case this does not convert the tumor into a fibroma, the same principle must obtain here. Nevertheless a true fibromatosis does affect one particular nerve, and may possibly be found to affect another. While the optic and olfactory nerves' are never involved

¹ It may be recalled that, unlike all the other eranial and spinal nerves, the optic and olfactory are devoid of sheaths of Schwann and of the cells giving origin to the same. in cases of multiple "neurofibromas," the first of these may exhibit what would seem to be a true fibromatosis.

A very full study of this condition has been made by my colleague, Dr. Byers,¹ who has shown that all true intradural tumors of the optic nerve are of this nature. Of these, more than one hundred are on record. His studies indicate that there is some relationship between these growths and obstruction of the lymph channels of the affected parts. There was a dilatation of the lymph channels and development of appearances resembling those seen in elephantiasis. It is deserving of note that the lymphatics of nerves form a system distinct from that of the tissues they traverse. In nasal polyps, judging from the frequent cedematous and mucinous condition, we have a very similar state of affairs, and here, also, we encounter the *fibroma cavernosum*—forms with greatly dilated vessels, evidently brought about by a similar blood-vascular obstruction. In elephantiasis proper we encounter a like tendency to subcutaneous overgrowth and production of conditions which, if more diffuse, have, nevertheless, much in common with fibroma molluscum. Such disturbed nutrition, if, as we suggest, a factor in these cases, must in its turn be due to a vice of development, for all these conditions of fibromatosis characteristically make their appearance in early life, or are familial.

CHELOID.2

Closely related, though distinctive in etiology, and to some extent histologically, is the condition of cheloid. This consists in an excessive development of subcutaneous fibrous connective tissue, sometimes so excessive as to produce large overlapping masses, or lobes, of newgrowth, covered by stretched skin. Two factors would seem to be at work leading to the conditions, namely: (1) a congenital predisposition; (2) irritation or injury. Thus, cheloid is especially common, in negroes, male and female, and in those, both of colored and white races, who present the condition, slight cutaneous injuries, which, in ordinary individuals, lead to but temporary disturbance, are liable to be followed by excessive growth of connective tissue and formation of a tumor-like mass. In a case studied by Martin, working in my laboratory, the mere running the point of a pin along the forearm, with a force sufficient to cause reddening without bleeding, was followed by the development of little fibroid nodules along the track of the pin.

It has been the custom to divide the cases into the traumatic and the spontaneous, but the more fully cases are investigated, the more


³ Some authorities write this *keloid*, from $sq\lambda x_{2}$, a crab's claw (from the cancer-like way in which the processes spread into the surrounding corium); others, *cheloid*, from $\chi q \lambda q$, a sear (from the relationship of the growth to a cicatricial tissue). The mode of spelling is thus still in dispute. The latter is the less fanciful and the more appropriate.

¹ Studies from the Royal Victoria Hospital, 1:1900:1.

are we convinced that in all cases the growth follows irritation, though this irritation is often such as in ordinary individuals leads to no aftereffects; thus it is at times seen to follow vaccination. Cases are on record in which the pressure and rubbing of a shirt stud have been followed by one of these growths, and in one frequently quoted instance there was a massive development in the skin over the shoulder, following upon the carriage of a basket on the naked skin of that region.

Here, as in cases of fibromatosis proper, microscopic examination reveals the absence of a capsule; the process of fibrous connectivetissue overgrowth extends, by more or less radially situated processes, in the subcutaneous tissue, and there is an imperceptible transition from the overgrown, cicatricial-like, to the normal connective tissue. The fibrous tissue of the keloid itself is often, though not always, characterized by the presence of extremely thick homogeneous bundles or strands of almost hyaline character, between which lie well-developed fibroblast-like cells.

FIG. 215



Section from a growth in a case of cheloid to show the coarse, hyaline connective-tissue bundles. (After Ribbert.)

Another feature is the liability of these cheloid growths to spontaneous absorption. Recent observations indicate that steady pressure on the growths is followed by their atrophy and disappearance.

Thus the distinguishing features of cheloid growths are:

1. They are composed of fully formed connective tissue.

2. They develop in consequence of relatively slight irritation or trauma.

3. They develop in those showing congenital or racial predisposition.

 They have no capsule, but merge imperceptibly into the surrounding connective tissue.

5. They are liable to retrogression and absorption.

Just as in connection with the previous forms of fibromatosis we observed a transition between the blastoma proper and strain hypertrophy, so here it will be seen we observe a relationship or transition between blastomatosis and irritation overgrowth. It will be useful to express these relationships and differences in tabular form:

TYPICAL HYLIC TUMORS

Nature of growth,	Characters of tissues.	Capsule.	Hereditary predisposition,	Other pos- sible factors in etiology.	Recutrence,
Fibroma	Fully formed connective tis- sue, but of atypical ar- rangement, growing inde- pendent of surrounding tissue,	Definitely present and complete, save where vessels en- ter. Encap- sulation easy.	Sometimes well marked.	Rare history of previous irritation. Most often no cause suggested.	Very doubt- ful after complete removal.
Fibromatoid growth (Fibromatosis)		Incomplete; possesses a definite hi- lum or hila.	Characteristie.	None.	Very fre- quent.
Cheloid	Fully formed tissue merging gradually into surrounding tissue; exag- geration of cicatricial type of fibrous tissue.		Often marked,	In all cases with full history, a history of (minimal) trauma,	Very fre- quent,
Elephantiasis	Fully formed tissue; ar- rangement least atypical; hypertrophy of normal con- nective tissue of part.	Wanting.	Absent.	Lymphatic obstruction main cause,	Absent; where cause can be removed condition undergoes absorption,
Fibrosis	Fully formed connective tis- sue; dense; overgrowth more moder- ate, diffuse or localized,	whole over-	Absent.	Irritation, or replace- ment.	Absent, un- less cause continues in action.

MYXOMA.

The maxyomata are tumors composed, in the main, of a tissue resembling none found normally in the adult organism, namely, a tissue composed of well-formed isolated cells of a somewhat stellate or polyhedral appearance, giving off delicate processes, the individual cells being separated one from the other by a matrix containing varying amounts of mucin, which takes on a differential stain with thionin. In this matrix there run large but thin-walled vessels. Some leukocytes are also present. We say formed in the main of such tissue, for it is very rarely that we come across what may be termed pure myxoma; in general, areas of the tumor show more condensed fibrous tissue, or cartilaginous masses, or frequently lobules or collections of fat cells, while in other cases portions are of sarcomatous type and show close collections of spindle cells. Thus, many pathologists doubt whether we ought to regard the myxoma as a separate form of tumor, and urge that we should speak rather of myxomatous modification or degeneration of some one or other form of connective-tissue neoplasm-of lipoma, chondroma, or fibroma myxomatodes, rather than of myxolipoma, etc. As such the majority of so-called myxomas must be regarded. But Ribbert has described small pure myxomatous tumors of the endocardium. Further, cases have been recorded as congenital myxoma, the tumors being recognized at the time of birth.



Section from typical portion of a mucoid polyp, (Collection of Royal Victoria Hospital,)

These tumors are slowly growing, are soft and fluctuating, so as to give the impression, at times, of being cystic or fluid masses. They never form metastases, but, if imperfectly removed, are liable to recur, while, again, a certain number take on sarcomatous properties, and so may become malignant; in such cases the metastases are not myxomatous, but wholly sarcomatous.

A frequent seat of these myxomas is the nasopharynx, where they are either pure, or present a condition of fibromyxoma or myxosarcoma; they form multiple soft polyps, appearing in the upper portion of the nasopharynx, or, again, actually in the nasal passages. Others develop, at times in the interstitial tissue between the muscles, and then, as a rule, are solitary, and, growing slowly, may attain a large size. A favorite seat for such tumors is the buttocks, between the glutei. These are the largest uncomplicated myxomas that we have encountered. Other forms of great size occur beneath the peritoneum, but then are found to be associated with fatty tissue, forming lipoma myxomatodes. Apparently associated also with fatty tissue are occasional small

subcutaneous myxomata. Chondromas are peculiarly liable to show myxomatous areas, as, again, occasionally do large fibromas and fibromyomas. The mixed tumors of the testis and parotid very commonly, also, show more or less extensive myxomatous development.

The tissue which in appearance the tumor tissue most nearly approaches is the developing connective tissue, more especially the Wharton's jelly of the umbilical cord and the developing fatty tissue of the factus, as, for example, the developing subcutaneous fatty tissue. Of pathological conditions, it is to be noted that in the newly forming fibrous tissue around areas of inflammation we at times meet with fibroblasts lying in a more or less mucinous matrix; indeed, mucin is a constituent of all the connective tissues, even, as recently pointed out, of bone.

Resembling thus developing connective tissue of certain orders, it might seem that this form of tumor ought to be of a malignant type; this, as above stated, is not the case. It may be pointed out that in the myxoma the cells in general are fully formed, and that the appearance of delicate branches is due to the fact that here the individual connective-tissue cells are well dissected out by the surrounding transparent matrix. In short, the presence of processes is no indication of arrest of these cells at an early stage of development; on the contrary, when the myxoma does take on malignant characters and becomes sarcomatous, these processes become unrecognizable, and the mucinous matrix disappears; and it may be pointed out that the amount of matricial mucin is by no means an indication of vigorous vegetative growth, but of the reverse. Rather, indeed, it would seem that there is some relationship between the vascular supply of the tumor and the development of the mucid matrix.

Whether the œdematous condition of the matrix favors the nonremoval of the mucin, or whether, on the other hand, the existence of mucin leads to increased absorption and retention of the fluid which diffuses out from the vessels, is an open question. But, certainly, on observing a large series of connective-tissue tumors, we appear to have every transition from simple œdema of the neoplasm to extensive mucinous infiltration and true myxomatous condition. To distinguish between the two conditions, acetic acid is to be employed. If mucin be present, the interstitial substance of the section becomes granular, and shows a network.

With regard to the causation, as already remarked, some cases are apparently congenital, and must be ascribed to isolated rests of either imperfectly developed fatty or fibrous connective tissue.

Of such congenial myxomas, Borst reports a colossal growth upon the mesentery of a nine-months-old child. This exhibited extensive lymphangiectases, to which he ascribes the soft nature of the growth; there were correspondingly dilated bloodvessels. He ascribes the tumor to a persistence and continued growth of the embryonal mucoid tissue of the mesentery.

Other cases, notably the mucoid nasal polyps, which are apt to show

LIPOMA

themselves between the ages of twenty and fifty, appear to follow chronic catarrh and inflammatory conditions of the region of development. As already stated, it is only a relatively small proportion of such nasal polyps that are truly mucin-containing; the majority are simply codematous.

LIPOMA.

The lipomas are sharply defined tumors composed typically of pure fatty tissue, that is to say, of fat cells lying in a vascular connectivetissue matrix. These cells are so abundant that but little else is to be recognized. The fat tends to differ from normal fatty tissue in being paler and not so deep-colored, while in general the individual cells are larger than those of normal tissue. Thus, even where situated in a fat-containing tissue, the neoplasm is well defined from the surrounding parts. In shape, these growths appear as rounded masses or frequently the tumor, while forming a single mass, is separated into a number of finger-like processes radiating from the central portion. This is especially noticeable in connection with subcutaneous lipomas.

In number, these tumors are most often single, but they may be multiple; in size they vary from minute, almost microscopic growths, such as are not infrequently met with in the kidney, to masses more than 30 kilos in weight (63 pounds), as in the retroperitoneal lipoma recorded by Waldeyer. We have recorded a similar case weighing more than 41 pounds.¹ These larger lipomas are often composed of multiple rounded or lenticular lobules, and show no tendency to form finger-like processes.


These tumors are essentially benign and of slow growth, nor do they recur after complete extirpation. They are liable to exhibit a restricted series of modifications; thus, the connective-tissue matrix may predominate, and separate off relatively small lobules formed of fat cells, in which case the tumor is of firm consistency; not infrequently the fat cells appear to give place to a more mucoid tissue, and the amor then assumes a more jelly-like consistency; such cases are spoken of as lipoma myxomatodes. The pure lipoma, however, is also very soft and fluctuating, and the larger growths have very frequently been mistaken for cysts and localized collections of fluid. More rarely, portions of the tumor take on a sarcomatous development; at times the central portions of these tumors undergo necrosis, and thus oil-containing cysts may be formed within them. Several cases are on record in which nodules of cartilage, and, in at least one case (Dreschfeld), of true bone, have been found within the tumor mass. Some hold that here we are dealing with preëxistence of cartilaginous or bony "rests" within the primary tumor mass; we incline to the view that these are metaplastic, a modification of the tissue being brought about by altered nutrition and cell relationship.

Regions of Occurrence.—Lipomas have been recorded from several regions of the body; most frequently they are found as sub-

¹ Perirenal and Retroperitoneal Lipomata, Montreal Med. Jour., 25:1897:529 and 620.

cutaneous outgrowths of varying size; these are especially common in the region of the shoulder; occasionally they are multiple; more rarely they may be symmetrical. Another form occasionally met with develops in the submucosa of the intestine, and here it is liable to develop into solitary pendulous or pedunculate masses, which at times have led to intestinal obstruction. In the kidneys it is not uncommon to meet with minute nodules of fatty growth; rarely these may attain large dimensions (Warthin).¹ An organ in which they have rarely been met with is the

F1G, 217



Semidiagrammatic cross-section through a perirenal lipona at the level of the renal vessels, seen from above. The perirenal and retrorenal fascia unite to form the transversalis fascia. The whole intestinal tract lies in front of the perirenal fascia: a_i descending colon; b_i perirenal fascia; e_i perironemic, d_i retrorenal fascia; e_i small intestine; f_i superior mesenteric artery; g_i duodenuu; b_i ascending colon. (Reprodels and Wadsworth.)

brain and its membranes; and here, again, they are of small size. Perhaps rarest of all is the lipoma originating in the bone marrow.² Occasionally, also, there may be a lipomatous development in connection with the synovial membrane of the joints, in which cases rather flattened, much fringed processes project into the cavity. Tumors have also been

⁴ I cannot accept the guarded conclusions of Archibald and Keenan (Jour. of Med. Research, N. S., 11:1907;121) that these renal lipomas originate from aberrant and included adrenal cells. A simpler, and I think, adequate view is that, like the renal fibromas and hypernephromas, they owe their origin to cells, in this case of connective-tissue origin, which have become nipped in between the reneuli in the process of development.

² Wehrsig, Ctbl. f. Path., 21; 1910; 243,

XANTHOMA.

met with in connection with the peritoneum, occurring here either in the mesentery or the omentum, or as excessive developments of one or more of the appendices epiploicae, originating beneath the pelvic peritoneum. Larger forms have been recorded developing from the peritoneum and the colon in front of them, the kidney tending to be compressed at the back of the tumor. In these huge retroperitoneal tumors it is noticeable that the mass continues to grow and to enlarge at the expense of the rest of the body; the patient may become markedly emaciated, the fat disappearing from the subcutaneous tissues and elsewhere.

Lipomatoid Conditions.—Just as in connection with the fibromas we noticed that there might be a progressive overgrowth of fibrous tissue, or fibromatosis, so here, to repeat, in connection with fatty tissue we have to recognize the existence of a condition of lipomatosis tending to be regional.

Steatopygy (p. 210), crural lipomatosis (of the hips), and coxæ (of the thighs) are clearly racial or stock inheritances.¹ Other conditions of general lipomatosis, even that developing late in life, are often familial. As indicated by the effects of thyroid treatment, they may be an indication of congenital lack of equilibrium between the tissues. Cases of *adiposis dolorosa* suggest strongly some trophic disturbance, and, as a matter of fact, Dercum, who has especially called attention to this malady, has reported two cases in which he has found disturbances of the pituitary, growths involving the nervous portion of this organ.

XANTHOMA.

The xanthoma is a small, benign, fatty tumor, as to the exact nature of which there has been much debate. Single or symmetrical xanthomas occur not infrequently as flat, subentaneous growths of a yellow color affecting the skin in the immediate neighborhood of the eyelids near the inner canthus (X, palpebrarnm). They may be present as multiple small nodules beneath the skin, (X, multiplex), forming slight prominences on the palms of the hands, in the neighborhood of the flexures of the joints, etc.; these are occasionally found also in the internal organs. These tumors have a more or less abundant connective-tissue stroma, in which are larger cells, containing markedly yellow, fatty globules, which give the xanthomas their peculiar color and name. As noted by Störk² and confirmed by Pinkus and Pick³ these fatty globules are doubly refractive and thus of the nature of myelin. The pigment is of the nature of a lipochrome. The careful study by S. Pollitzer⁴ affords strong support to the view that X, palpebrarum is in no sense

⁴ For an admirable study of circumscribed lipomatoses, see Shattock, Trans. Path. Soc. Lond., in Proc. Roy. Soc. Med., 2:1909; Path. Sect. 225.

² Sitzungsber, d. Kais, Akad, Wien, Math.-nat, Klasse., 115:96; Abt. 3:31.

³ Deutsch. med. Woch., 1908, Nr. 33.

⁴ New York Med. Jour., 1889; July 15.

a tumor proper, but is a fatty¹ or lipochrome metamorphosis or degeneration of muscle fibres connected with the orbicularis muscle. *X. multiplex*, on the other hand, he regards as somewhat of the nature of cheloid, as minute fibroid nodules, undergoing a central fatty degeneration and, like cheloid, liable to disappear. While multiple they never form metastases or take on malignant characters. This type has not infrequently been found associated with diabetes (*X. diabeticorum*). In other cases there is observed a relationship between the development of these and a condition of cholelithiasis without definite jaundice, although the skin is often found distinctly sallow; others are obviously of congenital origin. Pinkus and Pick include these varieties of the *X.* multiplex under the common heading of *Xanthoma tuberosum*. Yet another form is described by French writers, the "xanthome en tumeurs," in which tumors usually solitary and of the nature of giant-cell sarcoma present numerous xanthoma cells.

CHONDROMA.

Chondromas are tumors formed of one or other variety of cartilage, hyaline, fibrous, or reticulated (hyalo-enchondroma, fibro-enchondroma, reticulated enchondroma). They may be single or multiple, possessing in general a well-marked fibrous capsule, and are globular, or, if of any size, distinctly lobulated. Two varieties are in general distinguished, the *ecchondroma* and the *enchrondroma*, or chondroma proper, the former occurring as overgrowths of regions where cartilage is normally present and persistent, e. g., they arise from the cartilages of the ribs, the larvnx, intervertebral disks, and perichondrium of the trachea. When these, as is most often the case, are in direct continuity with those cartilages, they should certainly not be classified as true tumors, for they have not an independent existence, but, rather, are local hypertrophies; I therefore include those ecchondromas under the heading of chondromatoid. Only where they have no connection with a parent matrix do we have true tumors. It will thus be seen that, clinically and histologically, it is at times difficult to draw a distinction between the ecchondroses, or such localized hypertrophy, and true and proper ecchondromatous development.

Therefore, in the chondroma proper, or enchondroma, we have to deal with well-marked independent nodules of cartilaginous growth. These enchondromas may occur in many regions of the body, notably in connection with the bones; they are also found in connection with the parotid and submaxillary glands, in the testes, mammary glands, the lungs, and, very rarely, in the corpus cavernosum, the ovaries, and other internal viscera. It is interesting to note that they never develop from the articular cartilages of joints, although they may form in the fringes of the synovial membranes, and thus give rise to single or multiple "loose cartilages" in joints.

¹ More accurately, lipoid or myelinic.

CHONDROMA

As in normal cartilage, few or no bloodyessels are to be found in the substance of these tumors. This would seem to be the explanation of the lobulated character of the larger growths. Bands of connective tissue containing vessels divide the mass of the tumor, and afford nourishment. Growth occurs at the periphery, along the zone of the perichondrium, although in soft hyaline or myxo-enchondromata we obtain frequent evidence that the cells of the partly formed cartilage multiply actively; in short, obtain appearances such as are seen in the growing portions of foetal cartilage. Where the growth is at all large there is a peculiar liability for the deeper cartilage to be replaced by true bony tissue (osteo-enchondroma); or, without the formation of true bone, there may be extensive calcareous infiltration (enchondroma petrificum). Again, in the large enchondromas, from the increasing lack of nourishment of the central parts, as the tumor develops at the periphery, there may be a central degeneration and necrosis, the cells showing fatty degeneration, the interstitial substance undergoing liquefaction. If the

tumor project beneath the skin, and, through ulceration, the necrotic centre comes to communicate with the exterior, a fistulous track is formed, which shows no signs of healing; then, if the whole tumor be not excised, the atonic character of the process is peculiarly apt to favor general sepsis. In the parotid and testes these tumors are in general mixed, a combined overgrowth of cartilage and of the glandular tissue, and these mixed tumors, more especially of the

e.

a-

er

X

m

a,

a. se

1-

s,

ie

le

Fig. 218

Enchondroma exhibiting calcareous infiltration (*E. petrificum*). (Ribbert.)

testes, are liable to take on sarcomatous characters, and to become extensively malignant; these are solid teratomas (p. 658).

In these and in other chondromas there is a tendency to transition into mysomatous conditions; in place of a definite cartilaginous matrix there is a mucinous intercellular substance, and the cells take on all the characters of mucoid tissue, with long, delicate processes traversing the matrix (*myxo-enchondroma*); or sometimes appearances are not so definitely myxomatous; then we speak of *enchondroma mucinosum*.

As already stated, although these tumors are slowly growing and of firm consistency, and are in general of a benign type, only being dangerous on account of local disturbance of function, it is not infrequent that they form metastases. Here, at least, three different conditions have to be distinguished as favoring metastatic development: (1) in some cases, as, for example, in connection with the mixed cartilaginous tumors of the testis, there is a marked tendency for the growths to become more cellular and more sarcomatous in character; there is a definite assumption of malignant properties; (2) in other cases, as pointed out by Virchow, the tumor in its growth may penetrate into a vessel,

and there may be extension along the vessel, portions of a young, rapidly growing process becoming detached and carried to the lungs and other organs; (3) in some cases, as in a slowly growing nodular chondroma of the mammary gland of the bitch, studied by Boutelle in our laboratory, neither of these conditions is discoverable, and the only explanation afforded of the numerous metastases in the lungs, etc., is that certain cells from the rather richly cellular perichondrium—chondroblasts—have gained entrance into the veins, and been brought to rest in the lung, and there have developed into fully formed cartilaginous tissue. This may be regarded as a variant to the first case; what we would again point out is that in such cases no portion of the tumor need show any evidence of multiplication of a surcomatous type.

Studying the etiology of these tumors, it is noticeable how, in the majority of cases, they develop in childhood and early life. This is noticeably the case in chondromas developing in connection with bones. where, again, the condition is frequently multiple. Virchow has afforded a well-studied explanation of this condition. It is especially in those showing signs of rickets that these multiple chondromas develop. Now, in rickets the prominent disturbance is an undue preparation for the formation of cartilaginous bone, coupled with an incomplete or tardy formation of that bone. Thus, in the region of the epiphyses we find a very extensive cartilaginous development, and processes of this new cartilage project into what is destined to be the shaft of the bone, and here may become isolated and cut off from the main mass, and, as Virchow has shown in examining rickety bones, we may frequently meet with these minute islands of persistent cartilage. It would seem evident that these islands may, under certain conditions, take on an independent and aberrant growth, and give origin to enchondromas, or osteo-enchondromas, or sometimes true osteomas. In a certain number of cases the development appears to be hereditary, appearing in several members in one or two successive generations of a family, and in other cases traumatism appears to be a predisposing factor. According to Otto Weber, a history of traumatism can be obtained in 50 per cent, of cases of this form of tumor.

In yet other regions of the body it is probable that embryonic or developmental rests afford the nidus for the development of these tumors; indeed, most modern authorities are inclined to lay down that in every case we have to deal with these as a prime factor.

Thus, it has been pointed out that parotid enchondromas originate from cartilaginous remains of the hyoid arch; chondromas occurring in the neck, from the remains of other branchial arches; chondromas of the testicle, from portions of the anlage of the vertebral column included in the developing testis when it lay against the vertebral column before its migration to the scrotum; and chondromas of the mammary gland, from included portions of the sternal cartilages.

For ourselves, while in certain of these instances we are inclined to believe that this is the explanation, we are far from prepared to accept it as the general rule. Why, for example, should cartilaginous tumors of

ap-

, 18

OII-

OUS

We

the

es,

380

W.,

he

as

)r.

or

1S

to

the testis be relatively so common, while those occurring in the homologous organ, the ovary, are so very rare, save as one of the constituents of an embryoma or dermoid? The cartilaginous constituents of a parotid tumor gain their most satisfactory explanation either along the line of Wilms' theory of the "Mischgeschwülste" (p. 661), or of the same coupled with metaplasia of the connective-tissue elements of such pluripotential developments. Why, again, should enchondromas be liable to occur in mammary glands and not in the skin and other structures situated over the sternum, or in the anterior mediastinum? The mammary gland develops as a downgrowth from the skin, and has no special close relationship to the costal cartilages. I cannot, therefore, but consider that in some cases the anlage of the chondroma has to be sought for in the metaplasia of the connective-tissue cells of a glandular organ, brought about by some modification in the nutrition and vascular supply of the part. The not infrequent occurrence of cartilaginous islands in other forms of connective-tissue tumors of old standing would seem to favor this view; as, again, does the obviously metaplastic origin of some overgrowths formed of the closely allied tissue-bone (see p. 730).

The loose cartilages in joints, when truly cartilaginous, and not merely fibrous, are found to occur more especially in those of a rheumatoid tendency, and to develop not in connection with preexisting cartilage, but in the synovial fringes, which either are originally finger-like, or, through the growth of the cartilage within them, become more pedunculated, the peduncle then becoming twisted or broken off.

Here, then, in connection with the chondromas, we again see the action of certain tendencies leading to the formation of the tumor: (1) heredity in some cases; (2) the production and subsequent growth of developmental "rests" in others; while, again, (3) traumatism or irritation in a certain proportion of cases seems to afford the stimulus for progressive growth.

OSTEOMA.

In the sense in which we speak of fibromas and lipomas, that is to say, as defined tumors having a growth of their own, independent of the tissue in which they find themselves, *true osteomas are distinctly rare*. Osteomatoid conditions, on the other hand, are very common; of these, in fact, there is a bewildering variety. A mere metaplastic formation of bony masses in another tissue ramifying between the cells of that tissue cannot be regarded as a tumor proper; it has not an independent growth; the growth is determined by the vessels and nutritional conditions in that tissue. Rarely, however, such metaplastic bone is found to take an independent growth. We will, however, distinguish the various forms of what are not osteomas after laying down what constitutes a true osteoma and describing such.

The true osteoma may occur in connection with preëxisting bone (homoplastic), or apart, and in other tissues (heteroplastic). In the former case it may be superficial and derived from the periosteum,

or may be within the bony substance (endosteal), and then originating either from (a) a misplaced area of epiphyseal cartilage, which takes an independent development, as other cell rests are capable of doing; or (b) from the medulla, in which case it exhibits no cartilaginous forestage. It may either be dense and ivory-like, or loose and spongy, but always, in the case of true osteoma, instead of there being gradual and imperceptible transition from the new-growth into the surrounding bone, there is central growth and expansion outward, so that there is produced a condensation of that surrounding bone, followed, as the tumor continues to expand, by progressive absorption, until, in the shaft of a long bone, one of these tumors may, even if itself of loose, spongy structure, eventually cause such thinning of the shaft that fracture ensues.

Tumors of this nature, endosteal, though rare, do occur. Another, more compact form, must here be included. Occasionally within the shaft of long bones, somewhat more frequently in the antrum of Highmore, or other nasal fossa, again at the base of the skull in the region of the cribriform plate, there develop dense, indurated, bony masses. In such cases there is frequently the history of inflammation—thus, in the antrum, the growth most commonly proceeds from around the root of a tooth that has been inflamed or displaced, and so, at first, the condition in these cases would appear to be one of inflammatory overgrowth. But later the growth appears to be independent, and of a wholly different type, either from rormal bone or inflammatory osteophytes, and, though such masses show more or less of a pedicle of attachment, the fact that cases are on record in which the nodules have been found free—and dead—in the antrum, would seem to show that, with continued growth, this pedicle of normal bone is liable to be absorbed.

Of the *heteroplastic osteomas*, the simplest type is the periosteal isolated tumor occurring free in the immediate neighborhood of a bone. Here, either congenitally, in the course of development, a portion of periosteum becomes displaced, and sooner or later takes on independent growth, producing a bony nodule, which it surrounds, or, as appeared to be the case in a youth, a patient of my colleague, Dr. Archibald, through injury a portion of periosteum became wholly displaced from the bone into the surrounding tissues, and eventually gave rise to an independently growing nodule. Such tumors of periosteal origin, as, also, some of the endosteal tumors, show no preliminary cartilaginous stage.

The other type is that of the ossifying chondroma, or true osteochondroma. We have shown that, whether by metastasis, or metaplasia, or as cell rests, chondromas may occur in the soft tissues. Here they may become the seat of calcareous infiltration (which is not true bone formation); or, on the other hand, by vascularization they may in their main bulk be converted into true bone. The growing surface remains cartilaginous, but within, true Haversian canals and a marrow, red or fatty, become developed. Such osteochondromas may be met with also in connection with bone, originating from misplaced epiphyseal cartilage.

OSTEOMATOID

ating

oing:

fore-

dual

ding

re is

the

the

lose.

her.

the

In

the

of a

th.

gh

th,

1e

'y

y

Here must be mentioned another true blastoma, the osteofibroma or fibro-osteoma. We have once encountered a small specimen of this, by chance, in the medullary cavity of the femur, lying sharply defined, and shelling out without difficulty. The main constituent is a spindlecelled fibrous tissue, which, however, has the tendency to give origin to bony lamellae and spicules. This form evidently originates from bonemarrow.

The sharply defined nature of our specimen, despite its cellular character, showed that it was not to be included among the sarcomas. Other cases closely resembling this, but infiltrating and more cellular, must be classed as *osteofibrosarcoma*. But of the bone sarcomas we will speak later.

Let us repeat, the true osteomas are rare compared with the large number of cases which familiarly receive this name. These we must attempt to classify.

Osteomatoid.—All cases, in the first place, of localized or general overgrowth of the bones in which the growth is not defined from the normal bone, is not independent, and is of unknown cause (save that we may, in some cases, recognize congenital or inherited influences), must be classed under this heading. Here are to be included *hyperostoses proper*; among these:

1. Idiopathic general periosteal and endosteal *hyperplasias* affecting one or several bones. The long bones here are specially found affected, and, through the periosteal overgrowth, may assume remarkable forms.

2. Enostoses, localized and fairly circumscribed growths within bones, not so circumscribed and defined as to be independent of the surrounding bone.

3. Exostoses, processes of various grades rising from the surface of a bone, more particularly along muscular and tendon attachments. They may, however, be apart from these, as on the bodies of the vertebræ (where they are nodular, and often symmetrical, and so, evidently, of congenital origin). A special form is the *ivory exostosis* of the skull, small, button-like elevations of the cranial vault, which, by their intense hardness, are sharply demarcated. The periosteum over these is continuous with the surrounding, and, while the outer layers are singularly dense and compact, internally the growth passes imperceptibly into the underlying bony tissue. These cannot, therefore, be regarded as true osteomas. At the ends of the long bones may be found one or several exostoses cartilaginew. These arise either from the epiphyseal region or are terminal, and, like the joint surface, are covered with cartilage. They may thus be regarded as *ossifuing ecchondroses*. Closely allied to these is the *exostosis bursata*, in which, in addition, a synovial covering, with synovial cavity and fluid, is present over the free end. Such appear to have developed originally from the cartilage at the edge of the joint.

Their synovial cavity may communicate with that of the joint, or be free. Sometimes an extraordinary number of minute secondary synovial cysts is in connection.

In the second place, there are some conditions of bony growths apart

from the skeletal elements which should, perhaps, be here included. Such, for example, are: (a) Bony masses occurring in tendons, and not connected with the bone insertion. Every transition occurs from the tendinous exostosis of varying length to these free masses. They grow with the organism, and then remain stationary, exhibiting no tendency to independent overgrowth. (b) Bony plates in the dura mater and falx cerebri. These are generally small and flat, and are best ascribed to a persistence of the primordial bone-producing property of the dura. More extensive bony development in the membranes of the cord are associated with chronic inflammation. (c) Penile bone formation, as seen in some animals and in certain African tribes, a congenital ossification of the connective tissue in the part (here, again, inflammatory processes may produce the condition when not congenitally present). (d) Myositis ossificans. Here we deal with an extraordinary and slowly progressive development of bone within one set of muscles after the other, replacing the muscle tissue proper, and bringing about complete rigidity and fixation of successive portions of the body. Despite several studies of recent years (the condition having shown itself not to be so rare as was for long thought to be the case), we are still far from understanding the succession of events, save that it is generally accepted that it is not a myositis, not an inflammatory process. After having replaced the muscle, the bone shows no tendency to progressive growth. Provisionally this appears to be the place in which to note this condition in immediate relationship to:

Metaplastic Ossification.-In this, as already noted in discussing metaplasia, we have in general an alteration of the structure of a part. most often brought about by inflammation, though it may also accompany senile changes in a tissue, and, following this, a development of true bone. We may here note in order the main examples: (1) The myositis ossificans above described, if due to general constitutional vice, is also a metaplasia; (2) ossifications in the adductors in cavalrymen, in the deltoids of infantry, etc.; in both cases the result of irritation-in the one case, from repeated contusions of the muscles in riding; in the other, likewise, from carrying and firing the gun; (3) in the choroid coat of the eye: (4) in the brain (rare), apparently following upon inflammation; (5) in the spinal pia mater, as already noted, from leptomeningitis; (6) in the heart valves and arterial walls; (7) in the respiratory channels and lungs. Here several conditions are to be noted. In the trachea, besides ossification of the cartilages, there may be definite nodular submucous bony outgrowths, which in some cases are extensive, constituting the osteopathia trachealis of Aschoff.¹ These originate as pedunculate cartilaginous outgrowths from the tracheal perichondrium, undergoing direct osteoid and osseous metaplasia (p. 645). The pedicle may undergo absorption and then a slow independent growth may continue. In the lungs there may be scattered, irregular masses, which

¹Verhandl, deutsch, pathol, Gesellsch., 1910. As already noted, Aschoff is of opinion that some of them have an independent origin in the submucosa.

ODONTOMA

ded.

rom

hey

no

me

m-

in,

lly

try

ite

19

develop from the bronchial cartilages, or a wholly different condition of ramifying, coral-like tubes, apparently the outcome of chronic periarteritis or periphlebitis, with laying down of bone in the overgrown and degenerating fibrous tissue; (8) in old pleural and pericardial exudates. Here we most often have calcification, but true bone may be later developed; (9) in various tumors, lipoma, fibroma, angioma, and even in malignant surromatous growths and cancer (carcinoma ossificans); (10) in the middle coat and deeper layers of the intima of the aorta. The variety of the growths in which this may occur and the variety of their sites make it impossible to regard the process in them as other than metaplastic. The same must be true in a specimen prepared by Dr. Klotz, in which the outer layer of a phlebolith from the prostatic plexus exhibited true bone formation.



Odontoma, (Garretson,)

ODONTOMA.

As with bone proper, so with the teeth; pure blastomas would seem to be rare, blastomatoid conditions the commoner, namely, overgrowth not independent of the cement or dentine or alveolar periosteum. Yet reue odontomas undoubtedly occur.

CHAPTER XIX.

TYPICAL HYLIC TUMORS-(CONTINUED).

BONE-MARROW TUMORS: MYELOMAS.

THERE is a remarkable-and debatable-class of tumors which, at first sight, morphologically resemble the atypical connective-tissue tumors; but although in them, as we so frequently note with other hylic tumors, there are transitions, and, from what is apparently a benign growth, we gain sarcomatous development, yet characteristically the members of this group are not malignant; they do not infiltrate other tissues; they do not form metastases; and, studying these neoplasms more carefully, we become convinced that we are dealing, not with anaplastic cell elements-with the growth of cells, that is, which have become undifferentiated-but with cells which, when at their fullest development, are of relatively simple type; with what are benign overgrowths of one or other of the constituent cells of the bone-marrow. On these grounds it is well to separate tumors so formed from the sarcomas proper; the evidence, we think, is sufficient to justify us in treating, and, what is more, the clinical characters render it eminently wise to treat, these tumors as a class apart, as typical, and not as atypical, growths.

The bone-marrow, it must be remembered, contains many different elements, and the genetic relationship of these elements is still a matter of active discussion. But these elements have, obviously, specific functions which are strongly impressed upon them, and persist through the life of the individual. On the one hand, we have the cells directly concerned in bone function and regulation, the osteoblasts and osteoclasts, or myeloplaxes; on the other hand, the remarkable series of ervthroblasts, megalocytes, myeloblasts, and also lymphoblasts-of mother cells of the red blood corpuscles and leukocytes. The functions of these two orders of cells are wholly different, and, with recognition of the existence of the different forms, evidence has rapidly accumulated of late years that not only may there be overgrowth of one or other order, but even overgrowth of one particular form, while the apparently divergent facts that have been gained come very largely into line if we apply the same principles to these as to other connective-tissue tumors, and recognize three orders of growths: (1) typical blastomas; (2) blastomatoid diffuse overgrowth; and (3) atypical blastomas, either (a) of primary origin, or (b) as secondary developments from typical blastomas, or (c) developing secondarily from blastomatoid overgrowth. These atypical developments come all under the heading of sarcoma, and will in the main be treated along with other atypical hylic tumors. The other two conditions we may here briefly pass in review.


It is after not a little debate that I have determined to classify this

BONE-MARROW TUMORS: GIANT-CELLED

order of tumor with the myelomas. I fully admit the force of Borst's contention' that a tumor exhibiting any immature stage in the development of one of the connective tissues is a sarcoma; that bone is the completed development of the osteogenic tissue whether of the marrow or of the periosteum: anything less mature than this is either (as the least mature) a cellular sarcoma or (more mature) an osteoblastic sarcoma. But I encounter this crux, that in the osteogenic apparatus we encounter the myeloplaxes which as such are mature cells not passing on to form bone, and in these giant-cell tumors we encounter as the main feature those identical myeloplaxes—cells, that is, which are mature and not anaplastic. While all other cellular derivatives of osteogenic tissue are sarcomas, these as regards their characteristic element do not come under this heading, and their non-malignant properties sustain this contention.

The Giant-celled Myeloma ("Giant-celled Sarcoma").-Giant cells-multinucleated cells-are common to, or may be met with in, almost all orders of tumors. Their characters differ. Von Hanse-

mann recognizes three distinct forms: (1) The *forcign body giant cell* found in connection with necrotic and degenerating areas. This is identical in character with that already described as occurring in the tubercle, having a peripheral or polar collection of nuclei and central hyaline or necrotic substance, although, as shown in Fig. 140, the distribution of the nuclei is not of this order when the foreign body is lateral to rather than surrounded by the cell plasmodium. (2) *Parenchymatous giant*



Giant-celled myeloma of bone (After Ribbert.)

cells. The former class is of endothelial and leukocytic origin; here we deal with modifications of the cells proper of the tumor, brought about by irregular mitosis and lack of cell division, following upon nuclear division. Where this is the case the constituent nuclei are characteristically variable in size; often some are joined by bridges, or fibrils; others lobulated. Giant cells of this order may be found in cancers; they are especially common in endotheliomas. They are most frequent in the neighborhood of degenerating areas of such tumors, and would seem to be indications of irregular growth of the specific tumor cells; they are in no sense constant or regularly distributed constituents of such tumors. (3) Myeloplaxes. These differ from the first class in that the nuclei are distributed evenly through the cell body, and in the absence of central degeneration of that body, and from the second, in that the nuclei are well formed and of uniform size. They are present normally in the red marrow of bone, as osteoclasts in Howship's lacunæ, and are the characteristic constituent of the form of growth now under

¹See also Buerger, Surgery, Gynec., and Obstetr., October, 1909: 431.

which. -tissue other ntly a ically ltrate lasms with have On and, reat, ifie etly eo--of

r.

le

1-

consideration; here they may be present in enormous numbers. Under pathological conditions they are developed both in the marrow and from the periosteum. What is their relationship to osteoblasts is still a moot point. Some regard them as derived from the adventitial cells of the bone vessels, and certainly, even in neoplastic growths, they appear to be intimately associated with fine capillaries; others as of endothelial origin. Ritter regards bone building through osteoblasts, and bone absorption through osteoclasts, as brought about by two states of one and the same order of cell. Kölliker held similar views; Pommer and Wegner have noted transition from osteoclasts into spindle cells and fibroblasts. (4) Yet another giant cell, the sarcoblastic type, will be described later (p. 752).

The giant-celled myeloma, then, has the following characteristics: It grows locally, most often in the shaft (marrow) of long bones, or of the jaw; it may also be of periosteal origin (as in the giant-celled epulis of the jaw); its growth is expansile, leading to absorption of the surrounding bone; it is abundantly vascularized; it does not form metastases, save in the infrequent cases in which it undergoes sarcomatous modification (see later); it does not recur on complete removal, and, as recently shown, to prevent recurrence it is only necessary to remove the portion immediately involved, with very small surrounding zone; histologically, it exhibits a body formed mainly of short spindle-celled elements of fibroblastic type, somewhat irregular in shape, varying from the typical spindle to polygonal cells, and among these are abundant giant cells of the type described above. As in the fibro-osteoma (which is apparently closely related), so here, there is a tendency to the formation of bony lamellæ and spicules throughout the tumor, although the more abundant the myeloplaxes in an area, the less the amount of bone formation. A smaller, more round-celled matrix is the indication, we hold, that the tumor has taken on sarcomatous characters, and that metastases and recurrence may be feared.¹

Myelomatosis (**Myeloma Multiplex**).—From this we pass to a condition that has recently attracted much attention. As indicating its general character, we cannot do better than quote Borst's² description:


¹ Recently, at the meeting of the Association of American Pathologists and Bacteriologists, Washington, 1910, Mallory has dissented from the view here taken. This order of tumor, he holds, is a fibrosarcoma in which the giant cells are a fortuitous or at least a non-essential feature. If I followed his argument aright, giant cells of similar character may show themselves in the bone marrow for weeks after boring fine passages into the shaft of long bones and then around the particles of disintegrated bone; he would, in short, regard the giant cells in this form of tumor as more nearly allied to foreign body giant cells. For myself, I cannot but hold that they are specific constituents of the tumors every whit as much as the osteoclasts of Howship's lacunæ are specific constituents of normal marrow. I would, indeed, lay down that the bone-forming and bone-destroying cells are the primary constituents of the bone-marrow, giving rise to myclomas of the second order, which, indeed, may originate elsewhere, wherever these hemal mother cells have a normal existence.

² Loc. cit., 1: 493.

BONE-MARROW TUMORS: MYELOMAS

"The condition always (according to Winkler) affects the red marrow (vertebre, ribs, cranium), and this is converted into a dark-red or reddishgray or grayish-yellow tumor mass. It is a *primary multiple process*, showing itself simultaneously in bones widely apart; a sharp limitation of the multiple growths is often impossible; these are at times soft and pulpy, at times more firm. The spongiosa and even the firm shafts of the bones become absorbed, so that the tumor masses may show themselves immediately under the periosteum, which may here and there be ruptured. Fractures are common, as are also giving-way and distortion of the vertebre. What is remarkable is that the affection remains confined to the boy system; in pure cases there is no involvement of the lymph glands, no swelling of the spleen, no metastases in the internal organs. At times the progress of the disease is accompanied by intermit-

FIG, 221



Section of myeloma of vertebra. × 600. (S. Saltykow.)

tent fever and pains in the bones and joints. The blood in general is gravely altered; severe forms of anemia, pernicious anemia (Grawitz), and albumosuria are, to a certain extent, characteristic of the affection (Seegelken, Bence Jones, Naunyn, etc.)."

The condition, indeed, has been observed for many years, but under different names and with most diverse ideas to its nature (malignant osteomyelitis, myelogenous pseudoleukemia, sarcomatous osteitis, lymphadenia ossium). This clear presentation of its features demonstrates that the condition comes under our class of blastomatoid states. To speak of it as myeloma is incorrect if we respect our definition.

When we come to study the histology of these cases we meet with great variations in individual cases. Taking, first, the pure cases, it is noticeable that there is no overgrowth of the myeloplaxes, nor any indication of implication of the osteogenic marrow elements. The

ler ot be in. ne ve et It he IIes, ifion ly, of cal he tly ny int A he nd en.

ht, eks eles of but the I prider

ond ells tumor cells are all such as are derivable by overgrowth from the cytoblastic elements of the marrow, are of the myeloblastic or lymphoblastic type; sometimes mixed, more often corresponding to one or other type, so that it is becoming recognized as possible to distinguish different forms of the affection.

It cannot be said that the exact relationship of these myeloid cells has been absolutely determined: some authorities refer to them as myeloblastic—and such may well be the case; others describe the form most commonly seen as composed of cells of the plasma-cell type with eccentric nucleus.¹ If we are correct in regarding cells of this last order as of lymphoblastic origin, then such cases must be classed as lymphomatoid rather than myelomatoid. It may indeed be that we have to recognize myelomatosis without leukemia and blastomatoid overgrowth of the lymphocytic elements of the bone marrow without leukemia.

There are also impure cases: (1) in which the condition develops into a definite surromatous state, with metastases and infiltration; and (2) in which the lymphoblastic tissues and myeloblastic tissues in other parts of the body are, from the first, or become eventually, affected, and take part in the overgrowth, when we have myelomatosis complicating, or being complicated by, pseudoleukemia; (3) in Hammer's case, and possibly, also, in Baumgarten's, there was, in addition, extensive osteosclerosis and thickening of the spongiosa in all the bones; the former of these was of the first type, with eventual round-celled sarcoma; the latter of the second, with coincident involvement of spleen and lymph glands; whether here was an accompanying osteogenic myelomatosis must be left an open question.

Myelogenous Leukemia.—In the above cases while there is diffuse overgrowth of specific orders of cells of the bone-marrow, these cells do not become discharged into the blood to any marked extent. There are other conditions in which with similar diffuse overgrowth of the bone-marrow there is abundant discharge of the cells into the circulation, so that we obtain a condition of myelocythemia, myelogenous leukemia, or, as it is often termed, mixed leukemia. In this what is characteristic is the presence in the blood of large numbers of an abnormal element namely, of large mononuclear cells with neutrophilic granulations, although there is at the same time a relative or actual increase in the number of eosinophiles; "mast cells" are npt to be increased in frequency and similarly normoblasts are often met with—*i. e.*, nucleated red cells. The ordinary red cells are reduced in number, the white cells enormously increased, until they may be more numerous than the red, and these myelocytes may form one-third or more of the total white corpuscles.

When the marrow of the bones is examined it is seen to exhibit a striking hyperplasia: often it is reddened. On examination the main elements present are the *myelocytes*—large cells with neutrophile granulation similar to those seen in the blood—so also there are nucleated red corpuscles (crythroblasts) and frequent cells, large and small, with

⁴ See Christian, H. A., Trans. Assoc. Am. Phys., 22:1907:145,

LEUKEMIA: LYMPHOMA

eosimophilous granules. An associated anatomical character is the great enlargement of the spleen, causing this to be often spoken of as "splenomyelogenous leukemia." With Ehrlich we regard this not as a primary but as a secondary condition, due to accumulation rather than formation of the blood cells. In the liver and in the kidneys there are in some cases leukemic tumors due it would seem to active growth of the myelocytes outside the capillaries.

We have, briefly, the picture of an excessive growth of "leukoblastic" elements of the bone marrow, at times confined to the bone marrow, and at times—and it would seem secondarily—in other tissues of the body.

Chloroma.—In close relationship, in fact, as one member of this group, must be included chloroma. This is a form of growth characteristically multiple found in association with the bones of the face (orbit) and skull, affecting also the vertebræ and, more rarely, the ribs and bone marrow, which, on first removal and examination, show many of them (not necessarily all) a striking greenish or greenish-yellow tint, which may tend to fade upon keeping the specimens.

The disposition of these tumors is strongly suggestive of periosteal or bone growths, and as such Chiari has regarded them; but, as Dock¹ pointed out, they show none of the elements found in ordinary periosteal tumors—no spindle or giant cells, and no tendency to bone formation. Their appearance is that of a lymphoid overgrowth, with a well-marked reticulum, in which lie medium-sized round cells. Still more suggestive is the fact, to which Dock particularly has drawn attention, that, associated with this condition, is a type of leukemia in which, as shown by Dock and Warthin,² the prevailing cells are of the large lymphocyte, or, more accurately, of the myeloblast type. We deal here, evidently, with an aberrant form of myelomatosis. In some cases a co-existent lymphglandular tuberculosis has been recorded, but this bears no direct relationship.

There is still debate as to the nature of the pigment, which some find present in the form of highly refractible minute droplets or granules, others regard as universally diffused. Some (Chiari, Huber) hold it to be a lipochrome; others (Dock) can find no evidence of the presence of fat.

LYMPHOMA AND LYMPHOMATOSIS.

Although, inevitably, such a course leads us to consider various conditions on the borderland between the atypical blastoma and blastomatoid conditions³ rather than states of typical blastoma, we here, naturally, pass from the bone marrow to the allied lymphoid tissues and their overgrowths, and, doing so, have to admit that the time is not ripe

² It will have been observed that similarly we recorded no typical leukoblastic myeloma as distinct from myelomatoid conditions.

cytolastic type, erent

s has yelomost ccenas of utoid mize the

lops and other sted, npliase, nsive rmer ; the mph tosis

s do are onei, so , or, ic is nent , alumand

fuse

hese it a nain

The

red

¹ Am. Jour. Med. Sci., 106; 1893; 152.

² Medical News, N. Y., 1904.

for a full classification of this most complicated group of conditions; as, again, that our treatment of the problems involved can be but summary.

Nevertheless the work of the last few years has very materially illuminated what had been a very obscure subject. We would note especially the studies of Kundrat,⁶ Löwit,⁷ Sternberg,⁹ Türck,⁴ Yamaska,⁶ Dorothy Reed,⁸ and MacCallum,⁷ and when to these we add a recognition of the existence of blastomatoid states, as here put forth, and apply this, we are inclined to believe that the different conditions fall into a natural grouping, even if we have to admit that there are transitional cases, intermediate between the various classes, along with other cases of relationships that are still uncertain.

The terminology of these lymphomatous states is, we admit, appalling - not so much the terms themselves as the diverse meanings given by different observers: pseudoleukemia, lymphadenoma, lymphosarcomatosis, to mention but a few, have very different meanings in the writings of different men. These terms we will not discuss. Rather we will lay down in the first place (1) that just as we recognize that the lymphocytes and what it is convenient to term the leukocytes (the polymorphonuclears and the eosinophile cells) have distinct origins, so it must be kept in mind that we have distinct if fairly parallel series of blastomatous and blastomatoid conditions, owing their origin to aberrant growth in connection with the tissues giving origin to these two orders of cells; (2) that just as upon analysis we determined that among the overgrowths of fibroid tissue a series of distinct conditions were to be made out from chronic inflammatory hyperplasia to the typical fibroma, and, eventually, to the fibro-sarcoma, so here we have an identical series. Accepting these two postulates, the various conditions fall into a comprehensible order. To discuss them it is well to clear the deck by first casting overboard the conditions due to overgrowth of the "leukocyte" producing tissues (as distant from the lymphocyte). This we have already attempted to do. The leukocytes originate from the myeloblasts. We have seen that these myeloblasts have their seat in the bone marrow: that this marrow contains specific cells of different orders, namely, osteoblasts (with osteoclasts) and myeloblasts, along with other cells which are not specific-lymphocytes (lymphoblasts) and connective-tissue cells (fibroblasts). Tumors derived from these specific cells we term myelomas, and of these we recognize the two main orders of the giant-celled myelomas and the various conditions due to overgrowth of the myeloblasts (and myelocytes), namely, myelomatoid or myelomatosis (myeloma multiplex), and what may be termed myelomatosis gravior, with escape

⁴ Berlin, klin, Woch., 1901.

⁴ Wiener klin, Woch., 1893, No. 12.

² Lubarsch and Ostertag, Ergebnisse, 8 (7th year):1902:36.

³ Ibid., 12 (9th year, Pt. 2): 1905: 360.

^b Zeitschr. f. Heilkunde, 25:1901:200 and 313,

^e Johns Hopkins Hospital Reports, 10, 1902;133.

⁷ Trans. Assoc. American Physicians, 22: 1907: 350.

HODGKIN'S DISEASE

of the proliferating cells into the blood, namely, myelogenous or mixed leukemia (in which the cells escaping into the blood possess granules and resemble myelocytes), and chloroma, in which the escaping cells are of the more primitive myeloblastic type without granules. We now can approach the lymphadenoid overgrowths proper. These are:

1. **Chronic Hyperplasia** (comparable with chronic inflammatory fibrosis). Some of the best examples of this are seen in connection with tuberculosis. In this connection there may be either (a) specific granulomatous change in the lymph glands with caseation, or (b) at times a diffuse enlargement of the glands without caseation, with marked fibrosis of the glands, prominence of large endothelial cells and diminution over large areas of the masses of lymphocytes. Along with other observers' we have encountered cases showing distinct evidences of tuberculosis elsewhere with general firm enlargement of the messence in them of recognizable tubercle bacilli.

Such cases may be described as *paratuberculous lymphadenitis*. Cattle, more particularly, show this tendency to profound hyperplasia, suggesting that in them (and in those human beings exhibiting a like change) there is a constitutional tendency to excessive overgrowth of this tissue following upon the stimulus of certain toxins.

2. Hodgkin's Disease (comparable with cheloid). In the above cases we deal with a known irritant as exciting cause of the lymphadenoid hyperplasia. There is a condition of widespread enlargement of the lymph glands of chronic inflammatory type in which so far the exciting cause is not known, although many cases (although very far from all) regarded clinically as such have been found eventually to afford evidences of tuberculosis, and to have possessed similar histological features. These cases with unknown cause constitute Hodgkin's disease.

In general, the cervical lymph glands are first affected, becoming greatly enlarged, and consolidated into dense masses; then, progressively, the axillary, inguinal, retroperitoneal, bronchial, mediastinal, and mesenteric glands may show enlargement, and enlargement of the spleen makes its appearance. Later, the liver undergoes enlargement, and at autopsy may show a diffuse lymphoid infiltration along the bloodvessels, notably the portal branches. The same is true in the kidney. In extreme cases the peritoneal and other serose, the submucosa of the intestines and other areas become also involved. The lungs remain free. The blood shows no pronounced change, no advanced leukocytosis. There is often an intermittent febrile or subfebrile state.

The histological characters are very different from those of leukemia. Taking first the enlarged groups of lymph glands, these show no signs of infiltration. In this we have a strong distinction from lymphosarcoma. The marked feature is the reticular and connective-tissue overgrowth similar to what is noticed in the tuberculous group. There is a relative, if not actual, reduction in the lymphocytes and cells of the germ centres, a marked prominence of the larger cells of endothelial type, with often a

¹ Notably Weichselbaum and the Viennese School and Harbitz, of Christiania.

tions; sum-

y illuespeaska,⁵ nition y this, atural cases,

alling en by oomaitings ill lay uphotrant ers of hastotrant ers of and, comfirst cormade and, corfirst rrow: stoich lasts, rrow: stoich cells yelolona


scape

01.

marked abundance of eosinophile cells.¹ The connective tissue, reticular and endothelial overgrowth is the marked feature. The same is true in the spleen. The organ is dense, fibrous. There is a remarkable increase in the connective-tissue elements of all parts, capsule, trabecule, vessel walls, reticulum of the sinuses, so that splenic pulp proper is greatly reduced, and the narrowed sinuses show endothelial overgrowth, or large size of the cells, with loosening of the same. The lymphoid tissue along the vessels is present, but is not a marked feature.

The appearances are different in the *liver*, *kidney*, and other regions where normally lymphoid tissue is little noticeable. Here it is the lymphoid cells that are the main feature. Some lymphocytes are always present in connection with the perivascular sheaths of the vessels. Now, around these vessels we find dense collections, but on examination these are found to be provided with a reticulum, and not to infil-





Section of enlarged lymph gland from a case of Hodgkin's disease, showing prominence of cells of large endothelial type, with occasional eosinophiles. (D. M. Reed.)

trate the tissues, though they may cause atrophy of the constituent cells. Here we deal with a *lymphoid hyperplasia*. The same characterizes the subserous growth, if present. *The condition is not one* of metastasis.

The marked contrast between the involved normal glands and spleen on the one hand, and these secondarily involved organs on the other, strongly suggest that in the one we are dealing with reduction of the lymphoid elements, in the other with a compensatory hyperplasia. It may well be that in the

early stage in the lymph glands there is similar lymphoid hyperplasia,² followed by exhaustion or atrophy, as the connective-tissue overgrowth progresses.

The picture, it will be seen, is wholly unlike that of malignant growth. To speak of Hodgkin's disease as a form of lymphosarcoma or as a lymphomatosis is absolutely unjustified. It approaches much more nearly to the results of chronic irritation; it may be compared with cheloid and regarded as an excessive overgrowth of the lymphoid stroma secondary to a minimal or unrecognizable irritation.

3. Closely allied to Hodgkin's disease is a condition which has been described by Schottelius and by Gross,³ and of which two cases have been studied at the Royal Victoria Hospital by my colleague, Dr. Keenan, in which the earliest lesion has been an enlarged and apparently inflamed subcutaneous lymph gland tending to ulcerate, unaffected by ordinary

¹ To this Miss D. M. Reed (loc. cit.) has called particular attention.

² This has been described by MacCallum and others,


³ Ziegler's Zeit., 39; 1907; 405,

LY MPHOMA

treatment and in which, despite excision, there has been extension locally along the lymphatics with involvement of the related lymph glands, which on section resemble those of Hodgkin's disease. In one of these Professor Ewing, of New York, by intensive staining reported the presence of tubercle bacilli. These have been described as granulomatous lymphomatosis: more accurately they are granulomatous hyperplastic lymphadenitis.

4. In **leukemia**, on the other hand, we have what is characteristically a lymphomatoid condition, comparable in every respect with fibromatosis. Here we find localized *overdevelopment* of *lypical lymphoid*

tissue, the different constituents, reticulum, sinuses, and cells, being developed in due proportion. Now this is more restricted to the spleen or certain groups of lymph glands, now more widely developed, involving both spleen and glands, and also the lymphoid tissues of the bone marrow. Eventually it may, also, as in Hodgkin's disease, secondarily affect the liver and other organs, but the growth is not to the same extent as may be found in the latter condition. Of indications of inflammatory overgrowth the indications are slight; through distension the capsules of indi-



Lymphosarcoma, pencilled to remove the cells in large part and show the characteristic stroma. (Le Count.)

vidual glands may become thickened, and they may become compressed into large consolidated masses, but the reticulum shows no corresponding overgrowth. So, also, in the spleen the trabeculæ may be somewhat thicker than normal, but in general this organ, despite its frequently enormous size, shows singularly little recognizable departure from the normal under the microscope, save that the lymphoid elements, the Malpighian bodies, are large and somewhat diffuse. As distinguished from malignant growth, the lymphoid hyperplasia respects boundaries and shows no signs of infiltration.

The main clinical and diagnostic distinction between this and the

ticular rue in rkable eculæ, greatly r large tissue

egions is the dways essels. minao infily may ituent a *lym*y charvth, if ot one

tween s and these on the in the action n the typerin the lasia,² rowth

owth. *lym*nearly d and ndary

been been an, in amed linary

Hodgkin's type is that with this overdevelopment of otherwise normal formphadenoid tissue there is the passage into the blood of excessive numbers of lymphocytes, whence the name of leukocythamia (Bennett) or leukemia (Virchow). As already indicated, we believe in the existence of a group of cases which essentially belong to this class (corresponding to that of myelomatosis without leukocytosis), namely, of lymphomatosis without lymphocytosis, or a pre-leukemic stage of leukemia. This may eventually develop into a true clinical leukemia, or, on the other hand, may develop up to a certain stage and there remain latent, with no further overgrowth; or, lastly, may slowly retrograde. Here, and not with Hodgkin's disease, must be placed some cases, at least, of idiopathic splenic tumor of children and adults.



Abdominal lymphosarcoma; section through two of the close-set small secondary growths: T T, covering the surface of the liver G. (Martin.)

5. **Typical Lymphoma** (comparable with typical fibroma). It is obvious that when lymphadenoid tissue is laid down so diffusely through the organism, it becomes a matter of peculiar difficulty to recognize a typical lymphoma and distinguish it from mere localized hyperplasia of a preëxisting lymph gland. The fact that atypical lymphoma occurs indicates that the typical blastoma must also exist. We know of one such case recorded, that by Le Count. Kundrat likewise recognizes the simple local regionary lymphoma.

6. Atypical Lymphoma.—Lymphosarcoma.—When we recall that the lymph follicle contains tissues of more than one order—vessels and endothelium—along with the specific lymphocytes, it becomes evident that there may be several forms of sarcoma originating therein—spindlecelled, simple round-celled, etc. These scarce come into the category of atypical lymphoma, save that it is almost impossible to state whether certain round-celled forms of sarcoma represent the most anaplastic and malignant type of lymphoma. Here we refer more particularly to two

LYMPHOMATOID CONDITIONS AFFECTING THE SPLEEN 743

orders of tumors, well distinguished by MacCallum, both coming under the class of Kundrat's lymphosarcomatosis. The first order is encountered in the thoracic region, apparently originating most often from the remains of the lymphoid tissue of the thymus. This is the commonest form of mediastinal tumor. It is a small round-celled growth, difficult to distinguish from the ordinary small round-celled sarcoma, but differing therefrom in the growth being *purely local and infiltrative*, spreading around the great cardiac vessels into the parietal pericardium, heart walls, and roots of lungs, involving the mediastinal and lymphatic glands, but not forming distant metastases—the spread, that is, is local and by the lymphatics not by the blood. The other originates as a primary lymphosarcoma of the intestinal submucosa infiltrating locally so as to enclose organs like the adrenal and pancreas and similarly spreading in the main by the lymphatics. The cells of this type are larger and there are interspersed phagocytes.

7. Lymphosarcomatosis.—Such tumors, however widely they infiltrate, are, I hold, wrongly termed "lymphosarcomatosis:" they are and they remain local. The termination—osis—is employed more correctly to indicate a generalized condition, and such multiple diffuse true lymphosarcomatosis is on record—what may be termed a malignant lymphomatoid state. Thus Goppert has noted the case of a boy of three years in which there was great enlargement of the thymus, enlarged torsils and cervical glands, some enlargement of the spleen, enlarged liver and kidneys, and everywhere the affected areas exhibited infiltrating masses of small round cells. These were present also in the pericardium, renal pelvis, and periosteum. There was a slight leukocytosis.

It will thus be seen that these various conditions fall into three main types: (1) productive lymphadenitis; (2) lymphomatoid (with and without leukenia); (3) lymphoma (typical, and atypical or sarcomatous).

The infinite relationship between the potential producers of the different cell forms of the blood, both in the bone-marrow and in the spleen, renders it obvious that there may be simultaneous involvement of the myeloblastic and lymphoblastic elements; indeed, Kundrat holds that myelogenous leukemia is always mixed celled. For practical purposes it is sufficient to recognize two types of leukemia: the lymphatic and the myeloblastic (myelogenous), respectively.

LYMPHOMATOID CONDITIONS AFFECTING THE SPLEEN.

Splenomegaly or Splenic Anemia.—The time is not yet ripe to make an assured statement regarding the nature and relationships of a group of conditions in which, for long years, splenic enlargement is the main symptom, unassociated with hyperplasia of the lymph glands. Osler distinguishes four types: (1) Simple splenomegaly, persisting for years without accompanying anemia. (2) Splenomegaly with marked anemia of a secondary type, associated with cutaneous pigmentation and liability to hemorrhages; these cases last from ten to twelve years, and then are

normal accessive bennett) distence bonding matosis nis may r hand, vith no nd not of idio-

growths:

It is rough nize a sia of occurs of one es the

that s and ident indleegory ether c and o two

apt to develop a secondary cirrhosis of the liver, with jaundice and ascites (Banti's disease). (3) The familial or infantile form (Frederich Taylor, Gilbert and Fournier, etc.) begins in infancy, affecting several members of the same family, and showing the same symptoms of Banti's disease, with associated stunted growth. (4) Gaucher's type of splenomegaly, with secondary anemia, in which the spleen presents what may be termed an endotheliomatosis, there being a very remarkable diffuse swelling and proliferation of the endothelium of the splenic sinuses, which dominates the whole picture. In Stengel's case this had advanced to the development of localized neoplastic growths in the organ. Here, also, the splenic enlargement and condition extend over many years. Intimately associated pathologically, though not clinically, is the condition of splenomegaly, not with anemia, but with cyanosis, and extraordinary increase in the number of erythrocytes, up to 9,000,000 to 13,000,000 per cm. (Polycythemia). Osler' and Weintraub² have brought together and analyzed the cases.³

Whatever the primary cause in the various conditions, histologically we appear to deal with a localized blastomatoid condition, now involving the splenic tissue in general, now the endothelium of the sinuses, and, in the last case, characterized, it would seem, by a continuance of, or reversion to, the foctal condition of this organ; for, in the foctus, the spleen is one of the active seats of proliferation of erythroblasts and production of erythrocytes. Recent observations show that this capacity to give origin to red corpuscles is latent in the organ throughout life; that it may be stimulated into activity by extensive loss of blood, etc.

MYOMA (LEIOMYOMA).

The uterine fibromyoma, generally, from its appearance and characters, spoken of as the uterine fibroid, is the commonest of all tumors, and in most respects a most typical example of the benign blastoma. Possibly for this very reason we know less regarding its exact nature than we do of many of the rarer forms. And yet it has been extensively studied.

Gross Characters.—The tumors are most frequently multiple; if one large growth be present (and this may reach and exceed the size of an infant's head), careful examination will generally reveal other minute growths in the uterine muscle; or the whole body of the uterus may appear replaced by a collection of hard nodules, closely packed, and varying from just recognizable, pinhead nodules upward, the organ gaining a most irregular, nodose appearance. These tumors affect the body rather than the cervical portion, and may be either (a) interstitial, (b) submucous, or (c) subserous. In the last two situations they may, through progressive growth, become pedunculate.

¹ Amer. Jour. Med. Sci., 126; 1903; 187, ^a See vol. ii, 1st edit.; 86, ² Naunyn's Festschrift, 1905.

LEIOMYOMA

Each individual tumor is to the naked eye sharply defined from the normal uterine tissue, though, while some shell out easily, others appear to be more intimately connected with the organ, and, microscopically, are seen to have a more diffuse "capsule," the multiple small vessels as they enter being surrounded by fibrous tissue, which thus connects organ and tumor. On section, the appearance is characteristic, but here two types may be noted, between which is every transition: (1) the *pure myoma*, which is the softer, though still firmer than normal uterine muscle, and has a reddish-gray appearance (in the uterus this is most often noted in connection with small and apparently recent tumors); and (2) the fibroid proper, pale, almost white in color, and extremely dense, so that it cuts with difficulty. Like the fibroma, the cut surface has a "watered-silk appearance," caused by the bundles passing in all directions, reflecting the light differently.



Section of portion of a pure myoma, showing the character of the nuclei and the appearance of the cells cut longitudinally and transversely. (Perls.)

These are the characteristic forms, but we may also meet with the following variations:

1. Telangiectatic, the bloodvessels being widely dilated (rare).

2. Lymphangiectatic, the lymph vessels greatly distended; and here we can again distinguish between (a) the ordematous fibroid, with diffuse infiltration of serous fluid, and a spurious myxomatous-like separation of the tumor cells; (b) lymphangiectatic proper, with widely dilated lymph channels; and (c) cystic fibroids (see p. 712). Hemorrhages are rare.

3. Calcified, either surface layers, or central part, or the whole tumor, becoming converted into a dense stony mass, which can only be cut through by the saw. Associated with this, the microscope may reveal a preceding stage of fatty degeneration, as, again, of extensive hyaline degeneration.

4. Necrotic. Occasionally, through arrest of the blood supply, or through infection, a tumor may soften and undergo a colliquative necrosis. A curious form of necrosis occasionally met with and affecting

l ascites Taylor, iembers ti's dissplenos what arkable splenic se this vths in extend gh not it with tes, up Wein-

tologii, now of the a confor, in ythroshow organ ye loss

charmors, toma. than sively

le; if size other terus ked, the mors ither tua-

)5,

either the whole fibroid or scattered areas of the same is what is commonly known as "red degeneration." This is frequently but not constantly associated with pregnancy.

Microscopically, we have every grade from a pure myoma to tumors in which no muscle tissue at all can be made out, but only bands of connective tissue—pure fibroma. To the beginner it is not always easy to recognize these two tissues; both run in bands, which are apt to spread and feather out. But, if the nuclei be studied, in bands which have been cut longitudinally, the distinction is soon made. Those of connective tissue and developed fibroblasts are spindle-shaped and relatively short; those of plain muscle fibres are larger, characteristically long, rod-like, and with blunt, rounded ends. So, also, cut transversely, the nucleus of connective tissue has a naked appearance; one

Coarse myoglia fibrils from an ordematous leiomyoma, (Mallory,)

observes round it no cytoplasm; the muscle fibre has a distinct body, and in well-preserved and well-stained specimens this appears roughly polygonal. As pointed out, more particularly by Benda and Mallory, examination of perfectly fresh material, appropriately stained, shows that the smooth muscle fibre contains in the outer layer of its cytoplasm numerous fine longitudinal striations (myoglia fibrils) which toward the poles coalesce characteristically into coarser fibrils.

This fact that the small myoma is preponderatingly muscular, whereas the large is preponderatingly fibroid, indicates forcibly that, in the course of growth and aging of the tumors, there is a progressive development of the connective-tissue elements, with, eventually, atrophy and replacement of the muscular bands. The relatively small vascular supply, which must become progressively and proportionately smaller as the tumor increases in size, well accounts for this gradual effacement

LEIOMYOMA

mmonly nstantly

tumors bands of always e apt to s which Those bed and istically transce; one



ighly llory, hows lasm l the

ereas the elopand ular aller nent of the more highly differentiated tissue. Whether there occurs what may be termed a degenerative metaplasia, the muscle fibres actually taking on the general characteristics of fibroid tissue, is still an open matter.

As a matter of experience, we would advise those desiring to demonstrate the structure of a typical myoma not to select portions of the largest and apparently finest examples, but to take a small growth —the smaller the better.

The tumors are essentially benign, showing purely local growth, and that extending, it may be, over long years. They may, however, set up, or be associated with, other uterine disturbances—menorrhagia and metrorrhagia, while a considerable number of cases are on record of secondary cancer of the body of the uterus which may invade a myomatous tumor; so, also, of sarcoma, similarly infiltrating the growth.

It can no longer be questioned that sarcomas may arise from plain muscle elements. This has been convincingly proved by the demonstration of myoglia fibrils in actively growing spindle-celled sarcomas (Mallory). Indeed, sarcomas originating in uterine fibroids have frequently been placed on record, and while some of these may originate from connective-tissue elements, the use of Mallory's stain shows that others are of muscular origin. I have, indeed, recently enjuntered a case in which the transition from the muscle bands to bands of short spindle cells and from this to a diffuse development of short, almost oat-shaped cells was most striking.⁴

This, however, does not detract from our statement that the tumors are characteristically benign, nor does the fact, already noted, that some dozen cases are on record in which metastatic myomatous growths have shown themselves outside the uterus. The paucity of such cases must be contrasted with the many thousands of cases in which neither of these events develops.

Etiology.—What is the mode of origin of these tumors? To this no very satisfactory answer can be given. We do not find them in the young; nor in the child have there, to our knowledge, been noted anything of the nature of cell rests which could account for their later growth. On the other hand, we note a distinct family tendency to their development. Two possibilities are, therefore, present: either that there exist cell rests so inconsiderable as to escape notice, or that some constitutional condition favors the segregation of the tumor cells in late life. We see no reason why both conditions may not obtain, but at present there are no adequate data for coming to any decision.

They may occur in both the single and those who have borne children. Pregnancy, therefore, cannot be regarded as an important factor. On the contrary, those who have borne children regularly and in number appear to be less liable than others, but here, again, no decision can be reached, for it may be that the presence of the tumors is the cause of the relative infertility of those with smaller families. At the same

¹ The condition was so diffuse that I am uncertain whether it should be regarded as a senile change or a sarcomatosis.

time it will be noted that the presence of myomas is not in itself incompatible with pregnancy.

Risger and Gottschalk, Lubarsch, Borst, and French pathologists in general regard the muscle fibres of these tumors as primarily originating from the muscular coats of the uterine arteries. They have reported cases in which, in small myomas, this relationship could definitely be made out. What is noticeable in all myomas is the relationship of the muscular and connective-tissue bands to the vessels, and a marked feature of the myoma, which renders it distinct from all other tumors that we can recall, is the frequent presence not merely of capillary vessels, but of arteries with well-developed muscular coat. Their presence must be regarded as supporting this view, which, if substantiated, renders it additionally likely that myomata may develop in later life from abnormal local overgrowth of what is not a cell rest. At the same time it must be kept in mind that other small myomas do not show



Gland acini included in a"myoma. (After Ribbert.)

this clear presence of arteries, and exhibit little more than a simple mass of aberrant plain muscle fibres. A priori, it is reasonable to suppose that in an organ formed essentially of plain muscle, these plain muscle tumors arise from the mass of the tissue rather than from the included vessels.

"Adenomyomas." — There is, however, an interesting group of myomas which, for a time, was held definitely to prove the cell-rest origin of myomas. These are the adenomyomas, wrongly so called,

for more accurately they are diffuse myomas containing scattered gland tubules, that show no signs of active growth. To these von Recklinghausen's writings⁴ have directed attention, and of late years they have been much studied. Such occur more particularly in myomas of the hinder wall of the uterus and below the angle of entrance of the tubes and in the wall of the tubes. But some cases are on record where these gland tubules have been seen in myomas of the body of the uterus. These tubules are lined by a columnar epithelium; sometimes they are distended into cysts, and often they have the characteristic appearance of a main channel, from which, on one side, several secondary tubules are given off more or less at right angles.

It was the last character which more particularly led von Recklinghausen to conclude that they represented included portions of the Wolffian body (paroophoron, or, more exactly, of the upper end of

¹ Die Adenomyome u. Cystadenome der Uterus u. Tubenwandung, Berlin, 1896.

ADENOMYOMAS

elf incom-

athologists arily orig-'hey have could dee relationssels, and n all other f capillary 'heir preslater life . At the o not show of arteries. ore than a rant plain riori, it is se that in entially of ese plain from the essels. '-There

interesting which, for finitely to origin of e the adeso called, scattered von Reckthey have nas of the the tubes here these he uterus. es they are earance of ubules are

Recklingons of the per end of

lin, 1896.

the Wolffian duct), the remains of which, in the broad ligament, frequently show this comb-like arrangement; to conclude, therefore, that the muscle elements of the tumor were overgrowths of the muscular coat of the included duct. This view cannot be accepted. The more the matter has been studied the more difficulties are presented by this view, though it still has its supporters. Most significant is the fact that, in the course of development, the Wolffian duct does normally enter into the uterus, but this not in the region where these "adenomyomas" are found, but in the cervical region, where myomas are strikingly rare. Others have pointed out, what impresses all who have studied any extensive series of uteruses, that processes of the uterine mucosa may extend and be found deep down in the muscle tissue. We have seen such more than half the distance between the inner and outer surface, and Ribbert has actually traced such sinuslike processes into an interstitial myoma; and seen that the tubules within the growth have all the characters of other uterine adenomyomas. In confirmation of this view, Cullen,¹ of Baltimore, has been able to trace this continuity in fifty-five out of fifty-six adenomyomas studied by him, and quotes cases in which during menstruation and pregnancy the included ducts have been found to exhibit changes of the same character as those undergone by the uterine mucosa. Such deep processes may also occur in connection with the intra-uterine portion of the Fallopian tubes. And lastly, Aschoff has called attention to the fact that in subserous myomas there may be peritoneal downgrowths which take on glandular characters. Our colleague, Dr. Goodall, has encountered an "adenomyoma" developing in the fundal portion of the uterine wall-a region in which, developmentally, the Wolffian duct could not enter. These facts are sufficient to show that these "adenomyomas" can be explained otherwise than by the theory of cell rests of portions of the Wolffian body, or Wolffian duct, or (as yet others suggest) of the Müllerian duct: they are diffuse myomas-if not "myomatoid"-with uterine glandular in-

Myomas of Other Regions.—We have thus far only discussed the one form of myoma, and that because the uterus is far and away the commonest site for these tumors, and there they gain their greatest development. More rarely we encounter them in other areas, notably: (1) Other portions of the *genito-urinary system*, the walls of the Fallopian tubes, the broad ligament (in both these cases they may be "adenonyomatous," in the latter they may occupy the site of the parovarium); the round ligament (ditto), testes, prostrate (rarely pure), kidney (often lipomyoma), ureter, mammary gland. (2) The digestive tract. Here, in connection with the non-striated muscle coats, they occur more frequently than in any other site outside the uterus. Usually small, they may in the stomach wall attain very large size; usually single, some cases of multiple intestinal myomas have been recorded. Most often they project inward, and then may lead to obstruction or invagination or

¹ Adenomyoma of the Uterus, Philadelphia and London, 1908; 194.

ulceration of the covering mucous membrane. Others, as in the case of a large gastric myoma described by our colleague, Dr. Nicholls, project into the abdominal cavity. The stomach and the intestines are more frequently involved than the œsophagus. One case is on record (Cohen) in which the tumor contained pancreatic lobules. (3) *Skin*. Cutaneous myomas are quite small, apt to be multiple. A rare case is on record of their appearance here in early childhood. Whether they arise from the muscle of the arteries or from the arrectores pili and muscular sheaths of the hair follicles is still an open question. The fact that when multiple on the skin, they are not found in other parts of the body, would rather favor the latter view. Indeed, considering the singularly widespread presence of plain muscle fibres throughout the arterial and venous trees, the rarity of localized myomas that can surely be recognized as originating around arteries and veins, is very remarkable and is in itself against the vascular origin of uterine myomas.

TYPICAL HYLIC TUMORS OF MESOTHELIAL ORIGIN.

RHABDOMYOMA.


With rare exceptions, new-growths of striated muscle fibres is found combined with growth of other tissues and sarcoma-like elements in tumors of the pluripotential, teratoblastomatous type, as, again, in teratomas proper. It is more particularly in the mixed tumors of the kidney, vagina, testes, etc., that we encounter more or less imperfectly formed striated fibres. Occasionally, however, we meet with pure rhabdomyomas, in general small and sharply encapsuled, and this in the kidney, genital tract, and other regions where mixed tumors may be found; but also in other regions, notably the heart muscle, and in areas where normally striated muscle is present—extremities, nates, orbit, etc.

In all these cases the fibres are of embryonic, imperfectly differentiated type; they may only show longitudinal fibrillation; or, if transverse striation is present, it affects only part of the fibres; in the other part, as in the developing muscle, are clusters or rows of nuclei, and, laterally or terminally, the fibre may be clubbed, without striation, and showing the nuclear clusters characteristic of sarcoblasts; or, lastly, large cells, with abundant cytoplasm and many nuclei of the perfect sarcoblastic type are present. These characters, coupled with the fact that, under normal conditions, the striated fibres show such imperfect regenerative powers, incline us to accept the prevalent view at the present time, that rhabdomyomas always arise from cell rests.

We encounter, in fact, cases in which these less differentiated stages are predominant; cases in which we have transitions toward the sarcomatous type. The difficulty in deciding whether we encounter true rhabdomyosarcomas is created by the fact that so often these tumors are of pluripotential type, and the vegetative cells seen may have origin-


RHABDOMYOMA

ated, not from sarcoblasts proper, but from a still earlier type, or from cells differentiated already toward the connective-tissue type. This, however, is to admit that sarcomas do originate from a forestage of the striated muscle fibre. There is, at least, one case on record, that of



Imperfectly formed striated muscle fibres from a rhabdomyoma of the asophagus, (Wolfensberger.)

Fig. 229



Rhabdomyvearcoma _miant-celled or sarcoblastic—from lateral muscle of trout. The spherical tumor was almost wholly composed of these giant cells, which here and there showed both longitudinal and transverse striation, more particularly in the relatively numerous elongated cells.

Störk,' of malignant rhabdomyoma (primary in the testicle, with secondary growths containing striated muscle elements in the retroperitoneal, mediastinal, and cervical glands). In others, while the primary growth has contained sarcoblasts and more highly differentiated elements, the metastases could only be described as pure sarcoma.

¹Ztschr. f. Heilk., 22

in the case of cholls, project ines are more ecord (Cohen) . Cutaneous s on record of arise from the dar sheaths of when multiple would rather ly widespread venous trees, as originating elf against the

, ORIGIN,

bres is found > elements in as, again, in umors of the s imperfectly et with pure l, and this in tumors may uscle, and in mities, nates,

rfectly differ-; or, if trans-; in the other ! nuclei, and, striation, and s; or, lastly, f the perfect with the fact uch imperfect at the present

ntiated stages vard the saracounter true these tumors i have origin-

We have described⁴ a pure rhabdomyoma, or rhabdomyosarcoma of this order, obtained from the red lake trout, in which the whole tumor was composed of cells of the sarcoblastic type, the majority multinucleated, some of the giant cells containing several score of nuclei; these giant cells had a tendency to be elongated and in parts showed definite cross-striation. The nearest approach to this pure type of "embryonic" rhabdomyoma in man is seen in a group of multiple myomatous tumors recorded in connection with the infantile heart muscle.² These are clearly congenital and often associated with diffuse cerebral scleroses,

TYPICAL HYLIC TUMORS OF EPIBLASTIC ORIGIN.

THE NEUROBLASTOMAS.

Only such tumors as contain nerve cells is it wise to class as *neuromas*; if, for example, we find a tumor containing abundant nerve fibres-axones -but nothing that can be construed as even an imperfectly developed nerve cell body, we know full well that each of these fibres is in connection with some cell body outside the tumor, and that thus, so far as regards its nervous elements, the growth is not autonomous and independent of the rest of the organism; and that, therefore, the term neuroma is inapplicable. But if these limitations reduce very materially the number of tumors that can be included under this heading, and if, as might be expected from the high degree of differentiation of the neuron, the gauglioneuroma, or true neuroma, is one of the rarest of tumors, on the other hand the more recent researches into the histology and embryogeny of the nervous system have proved clearly that other constituents of that system originate from the same order of cells as do the neurons. Just as in the ovary the "interstitial cells" and those of the Graafian follicles have a common origin, so in the nervous system four distinct types of cell can all be traced back to the primitive neuroblast. These are (1) the neurons; (2) the neuroglia cells; (3) the ependymal cells lining the canal of the spinal cord and its ventricular expansions in the brain; and (4) cells of the sheath of Schwann enclosing the medullated peripheral nerves. Hence, if for precision we employ the term neuroma in a restricted sense, it must be clearly borne in mind that at least four types of blastoma may originate from neuroblastic elements. We may name these (1) the neuroma proper, ganglioneuroma, or neurocytoma; (2) the *glioma*; (3) the *ependy*moma or adenoglioma, and (4) the neurinoma.

¹ Adami, Montreal Medical Journal, 37: 1908: 163.

² The fullest studies of these cases, some twelve in number, are by Seiffert, Ziegler's Beitr., 27: 1900; 145, and Wolbach, Jour. of Med. Research, 6:1907; 495. Carnegie Dickson has recently published a careful study of another case.

myosarcoma whole tumor ajority mulore of nuclei; parts showed pure type of nultiple myoeart muscle.² fuse cerebral

DRIGIN.


as neuromas; bres-axones ly developed in connection as regards its endent of the is inapplicaber of tumors be expected ganglioneuher hand the f the nervous tem originate the ovary the e a common all be traced cons; (2) the of the spinal of the sheath Hence, if for e, it must be nav originate the neuroma) the ependy-

iffert, Ziegler's 495. Carnegie

NEUROMA.

Tumors, in one case as large as a child's head, have been observed, more especially in the abdominal cavity, and apparently in close connection with the sympathetic system—the cardiac plexus—which, upon microscopic examination, have been found to contain abundant cells of the type of sympathetic ganglion cells, with numerous processes histologically identical with the axones and dendrites of the nerve cell proper. These, without exception, are recognizable early in life, and date back to the embryonic period. They would seem, thus, clearly to be explained as due to developmental anomaly—to the segregation or displacement of a portion of the developing "neuroblast," which now takes on independent growth, the typical neuroblast cells proliferating and giving rise

FIG. 230




Cells from a benign and a malignant neurocytoma (or true neuroma), respectively, the iormer from the sacral region, the latter from the retroperitoneal region at the level of the pancreas. (R. Beneke,

to nerve cells. Similar isolated tumors have been described in connection with the ependyma and the ventricles, evidently of like origin. We possess no instances of ganglion-celled neuromas of postnatal and irritative origin.

Of these tumors we distinguish two orders, the one form possessing as characteristic constituents clusters of large cells of the neuron type—typical ganglioneuromas—the other formed of aggregations of smaller round or pyriform cells, apt at first sight to be mistaken for a round-celled alveolar sarcoma. Closer study, as shown by Dr. J. H. Wright, who has been so good as to afford me the accompanying figures, reveals the existence of sheaves of fine fibrils, not staining by the ordinary connective-tissue stain and proceeding from the individual clusters. The 48

appearance, in short, is strikingly like that presented by the developing sympathetic ganglion, and, as in that, there may be smaller cell accumulations in the form of circles of the small cells with a central collection of fibrils cut transversely and presenting a punctate appearance. This is clearly a more embryonic type of growth approximating to the neurosarcoma, and as might be expected it has been found to afford metastases.¹

False Neuroma.—A condition of a different order is the so-called *am putation nuroma*. It happens occasionally that, following upon amputation—more rarely after mere section or rupture of a nerve—the proxi-



Atypical neurocytoma resembling the anlare of a sympathetic gauglion. From a child at the Massachusetts General Hospital. Multiple tumors of skull, mediastina, retroperitonead tissue, and liver. The masses of cells often contained cystic cavities. (Prof. J. H. Wright.) Still more "embryonic" neurocytoma from a child, sixteen months old. (Drs. Tileston and Wolbael.). The ball-like arrangement of the cells with central fibrils is characteristic of an early stage in the development of sympathetic gamption. (Prof. J. H. Wright.)

mal end of one or more nerves become swollen to two or three times the diameter of the nerve, forming a firm, painful, bulbous mass. Sometimes this can be freed without difficulty, but frequently cicatricial fibrous tissue unites it to the surrounding tissues, making dissection difficult. Upon microscopic examination the enlargement is seen to be composed of bundles of nerve fibrils, often curving upon themselves, and passing in different directions, embedded in a dense overgrowth of the fibrous endo-

⁴ Tumors of this nature have been reported by Marchand, Amberg, Kuster, Richards, Lapointe and Leeene, Tileston and Wolbach, many originating in the adrenal medulla.

and perineurium. Though the nerve bundles entering are, in general, in the main medullated, but little medullary substance can be recognized around the fibres involved in the growth.

The process of events leading to this condition would seem to be an outcome and variation of what normally happens after section of a nerve trunk (see p. 575), but here, owing to absence of a proper channel of least resistance down which to advance, owing, also, to the formation of surrounding cicatricial tissue, the growing fibrils turn upon themselves, and, together with the proliferated endoneurial tissues, form a definite enlargement, which ceases after attaining a certain size. What we have to deal with is, therefore, an *aberrant regenerative process*, and in no sense a true tumor formation. It nevertheless shows some relationship to the condition of a limb (and so of its nerves); *i.e.*, it only occurs in certain apparently predisposed individuals; and (2) where once in an individual such an "amputation neuroma" has been removed, there is a distinct liability for a second tumor to form at the freshly exposed end of the nerve stem.¹

GLIOMA.

A second element derived from the neuroblast is the glia, or neuroglia, formed of cells and singularly fine fibrils, which, together, give the stroma of the central nervous system its peculiar appearance. Compared with the neurons, these glial cells are small. They tend to be of oval shape, with a single nucleus and moderate amount of protoplasm, and, when teased out, show an extraordinary number of fine processes radiating from the body in all directions. These can be demonstrated in properly prepared 'material by Mallory's phosphotungstic hematoxylin or Weigert's neuroglia stain.

With regard to the nature and relationship of the fine fibrillae which form a felting all through the stroma of the central nervous system, opinion has been divided. There has been a discussion similar to that regarding the relationship of the fibrillae, yellow and white, of ordinary connective tissue. Weigert has laid down that these are not processes of the glial cells, but are independent derivations, formed and given off by the cells. Tavlor² and Pusey,⁴ employing Mallory's neuroglial stain.

⁴ Dr. Alex. Bruce, of Edinburgh, has shown me, and permits me to refer to, a unique case of multiple minute tumors of this order within the central nervous system. In this the axons can be traced passing down the vascular sheaths from the meningeal surface and ending in nodular masses formed of fibrils wound upon each other in every direction. He holds that a fortal meningeal lymphangitis had caused a diversion of the growing axons, so that instead of passing into the eranial and spinal nerves, a certain number of them became side-tracked into the meninges and thence made their way downward along the vessels into the brain and cord. He has demonstrated his specimens at the Bristol meeting of the British Pathological Society.

² Jour. Exper. Med., 2: 1897: 611. ³ Tr

10

³ Trans. Chic. Path. Soc., 4: 1899-01: 44.

TYPICAL HYLIC TUMORS

both found a definite connection between cells and fibrils. Bonome' has arrived at a similar conclusion, but holds that, in pathological conditions, this connection may be more intimate than normal, by which I understand that he takes the intermediate position, of regarding the fibrillae as being formed as processes of the cells which may later become separated, and so independent. By analogy with ordinary white connective tissue, this would appear to be the more likely relationship.

Tumors formed of these glial cells, with abundant or rare fibrils, are found only (a) in the brain, (b) along certain cerebral nerves (very rarely), (c) in connection with the retina, and (d) over the coccys (evidently from the remains of the neural canal). The increased growth of neuroglia elements in connection with syringomyelia is, nowadays, generally regarded as hyperplastic rather than of the nature of true tumor formation—as a gliomatosis, and not a glioma.

Glioma; numerous neuroglia fibrils surround the cells and run in all directions. (Mallory.)

Two varieties may be distinguished, the hard and the soft, the former found in connection with the ventricular walls, and projecting into the ventricles. These are well defined and easily enucleated. The latter are in the form of diffuse infiltrating growths, with no trace of a capsule, are very vascular, and peculiarly liable to be the seat of hemorrhages. In the brain, after death, their existence is indicated by the appearance of greater translucency than the surrounding brain tissue, and somewhat more bluish tinge, and by their more pulpy consistence; they often show signs of old and recent hemorrhages. The softness is due largely to the greater amount of glairy fluid in the matrix.

¹ Virch. Arch. 163; 1901; 469.

GLIOMA

These soft gliomas are found only in connection with the cerebral hemispheres and corpus callosum, and may attain a very considerable size, gravely compressing and replacing the normal brain tissue. Neither the hard nor the soft form of cranial growth shows any tendency to metastasis. These diffuse widespread growths suggest a condition of gliomatosis, rather than development from a single focus.

Retinal gliomata, on the other hand, exhibit more evident signs of malignancy, both in their capacity to invade the surrounding tissue of other orders and to give rise to metastatic growths. These tumors originate as soft, grayish, finely nodular tumors projecting from the retina into the vitreous. If not detected and removed at this stage, the growing tumor may invade the sclera, and so extend into the orbit; or, after completely filling the bulb, may erode the cornea and project externally as a fungating mass. It may also extend along the optic nerve into the cranium. Histologically, appearances vary.

Retinal gliomas are only exceptionally—and then only in part formed of typical glial tissue. Their structure, indeed, has led to considerable discussion as to their exact

considerable discussion as to their exact nature. In general, they are formed of small cells without processes, arranged characteristically in relationship to the vessels. Intercellular glial fibrils may be wholly wanting, or present only in parts of the tumor. The cells are grouped radially around the smaller capillaries, appearing to be in direct connection with the walls of the same, though, in the larger vessels, there is a layer of intervening connective tissue between the vessel wall and the surrounding cells. In the immediate neighborhood of the vessels the more or less radially disposed layers of cells stain



Section of retinal glioma, showing relationship of cells to vessels and formation of "rosettes." (Ribbert.)

well; farther away they tend to stain poorly, and to be broken-down and necroic. The appearance thus, to some extent, recalls that of the perithelioma (or periendothelioma). There are, in addition, to be determined certain characteristic cell groups, or "rosettes"—collections of cells arranged radially around an apparent lumen, recalling the tubules of an adenoma or glandular tumor, save that between the cells and the cavity proper there is a clear layer or membrane, from which occasional minute, conical processes project into the lumen proper; suggesting the rods and cones and external limiting membrane of the retina. These processes take the stain for glial fibrils; similar rosettes have been observed in cerebral gliomata. These retinal tumors are, thus, true gliomas, but formed of less differentiated cells, so that if, as I urge, we employ the term sarcoma purely in a histological sense, it is proper to speak of them as gliosarcomas.

ner nto l'he of ted ain onl'he ix,

10

re

th

ue

TYPICAL HYLIC TUMORS

Etiology.—In the newborn and in young children we occasionally recognize multiple small nodules, not well defined to the naked eye, but firmer than the surrounding tissue, which, under the microscope, are seen to be formed of glia, with included urve cells. That these are found in the white matter as well as the gray points clearly to the fact that here we have to deal with developmental abnormalities—with inclusions, or overproductions of nerve tissue. It is in young subjects that we are specially liable to encounter gliomas. Further, as pointed out more particularly by Bonome, the existence of cysts lined by epithelium, in not a few examples of this condition, can only be satisfactorily explained on the assumption of developmental anomaly, either by lateral branching of the central canal and its epithelium, or by the inclusion of growth, fulfils its destiny of producing epithelium of the nature of that lining the central canal.

There are, however, other secondary cysts which may appear in cerebral gliomas, the results evidently of cell necrosis, and these may become more or less imperfectly lined by a layer of more cubical cells, which, at first sight, may be mistaken for an epithelium. As we have pointed out elsewhere (p. 617), this is not a true epithelium.

EPENDYMOMA.

In the gray matter of the brain we occasionally encounter cysts lined by a definite epithelium which can only be ascribed to foetal inclusions of ependymal tissue. We have met with one such cyst in the frontal lobe lined by a ciliated epithelium. Further, the existence of cysts lined by a true epithelium in not a few gliomas can only be satisfactorily explained on the assumption of developmental anomaly, either by lateral branching of the central canal and its epithelium, or by the inclusion of undifferentiated neuroblast at a very early period, which in the process of growth fulfils its destiny of producing epithelium of the nature of that lining the central canal. The relationship of the neural epithelium to the other neuroblastic elements-the fact that neurous and glial cells are the direct derivatives from the epithelium-explains why we never encounter pure ependymomas; active growth of these vegetative ependymal cells leads coincidently to the development of cells of glial and even of neuron type, whereby the ependymal cysts appear as inclusions in a tumor of gliomatous type. It may be added that the simple cysts, above referred to, like other cell rests cannot be included in the category of tumors proper.

NEURINOMA AND NEURINOMATOSIS.

There can be no clearer demonstration of the dependence of pathology and more particularly the study of tumors upon embryology than is afforded by the controversy that has ranged for long years regarding the

NEURINOMA AND NEURINOMATOSIS

nature of a remarkable order of multiple tumors, which according to their position and to the prevailing histological teaching of the time have received very various names (fibroma, neuroma, "Rankenneurom," molluscum fibrosum, neurofibromas, fibromatosis nervorum, Recklinghausen's disease, etc.). The cases we refer to are those in which multiple tumors along the course of the main nerve stem and peripheral nerves are associated with multiple tumors of the subcutaneous tissue, now the one, now the other group preponderating. These subcutaneous nodules, from the size of a pinhead to that of an orange and larger, may be several hundred in number. In some few cases the growths have shown themselves also in the intestinal wall. It was in 1881 that von Recklinghausen called attention to the frequency of the association of "multiple neuromas and cutaneous fibromas." He termed these neurofibromas, but later (1898), holding that the essential feature was an overgrowth of the connective tissue envelopes of the nerve bundles and of the endoneurium, and that the term neuroma should be restricted to tumors formed of proliferative nerve cells, he concluded that these should be placed among the fibromas,¹ the nerve fibrils present not participating in the tumor

This view of von Recklinghausen gained general acceptance and the teaching of Klebs² (1889) that the primary growth originated from the medullated nerve fibres which became converted into bundles of fibrillæ, with later connective-tissue overgrowth and fibromatosis proper, was generally discredited. Nevertheless, of late years there have appeared careful studies of cases difficult to explain according to the prevalent views. Knauss^a has described a case of multiple subcutaneous growths of this order in a girl, aged eleven years, in which he discovered abundant ganglion cells; Askanazy⁴ a collection of cases of multiple tumors in the gastro-intestinal walls with or without associated tumors along the peripheral nerves and under the skin; and both workers call attention to the existence of ganglion cells along the fine sympathetic plexus around the cutaneous blood and lymph vessels and in the plexus mesentericus respectively, and see the origin of the growth in immediate connection with these. From another point of view Durante,5 regarding each interannular segment of the medullated nerve as a cell, or cell compound, governing the formation and nutrition of the contained axis cylinder, has postulated that, with aberrant growth, the differentiated products of these segmental cells disappear, or fail to be found to a greater or less extent, the vegetative cells (of the sheath of Schwann) enlarge and proliferate, forming nucleated bands or spindle cells. Thus are obtained various grades of more fibrous or more cellular (sarcomatous) tumors.

¹ As such or rather as a condition of blastomatoid fibromatosis, I considered them in the first edition of this work.

² Allgemeine Pathologie, Jena, 2; 1889. ⁴ Arbeit a. d. path.-anat. Inst. z. Tübingen, 2; 1894-99.

⁵ Cornil and Ranvier's Manuel d'Histol. Pathol., Paris, 3d edit., 3:1907:823; so also Francini, I Neuromi, Siena, 1909.

n is the

16

1g

чr,

abe

the

the


pe.

TYPICAL HYLIC TUMORS



Tumors of sciatic nerves and their branches. At a, large tumor connected with small intermuscular nerve. (Preble and Hektoen.)

Fig. 236



Section through a fibromatoid entaneous nodule showing the nerve fibers (n, f_{\cdot}) separated by fibroid overgrowth. (After Ribbert.)

As will be gathered from our account of the regeneration process (p. 625) we cannot accept this view of Durante, regarding the segmented nature of the peripheral medullated nerves. But as shown by Verocay,1 Durante's view contains this element of truth, namely, the recognition that the cells of the sheath of Schwann are of neuroblastic and not, as had been taught by Kölliker, of mesoblastic origin. That this is their nature has now been fully established by Kohn.2 Not a few observers (notably Genersich and Soyka) have noted the proliferation of the sheath of Schwann in these cases, but, regarding these as mesoblastic elements, have regarded the tumors as essentially fibromatous. Certain French observers, on the other hand, from Bard³ onward, while making the same observations, have regarded these cells as of nervous origin and the tumors derived from them as neuromas of the peripheral nerves. Verocay now shows convincingly that this is the view that must be

¹ Ziegler's Beiträge, 48:1910.

² Arch. f. Mikr. Anat., 70 ; 1907.

^a Arch, de physiol, et pathol, 1885; see also Grall, Thèse de Lyon, 1897, and Gautier, ibid., 1899.

CHORDOMA

accepted, or, more accurately, that the growths are formed from a "neurogenous" tissue either derived from the cells of the sheath of Schwann, or the less differentiated precursors of the same, from cells, that is, which give origin to characteristic nucleated bands and pale delicate bundles of fibrils. Just as the neuroblast may give origin to neurons, glia cells, and these fibre cells, so in such tumors may occasionally be encountered the simplest form of neurocyte-small cells resembling lymphocytes and various gradations from these up to cells of the imperfect neuron type. The tumor is neither a neuroma nor a glioma, but a neurinoma.¹ We would go farther and say that we deal not with neurinomas (with a blastomatous condition), but with a condition of neurinomatosis (blastomatoid). The multiple nature of the growths along the continuity of the nerves demonstrates this. What is of interest in this connection is that in a large number of cases, study of the central nervous system shows the coincident existence of gliomatous areas in the brain and spinal cord; we deal with a tendency toward excessive vegetative activity of the nervous tissue, and where the condition affects the sympathetic ganglia there may develop true neuromas.

Lastly, to complete the picture, with the overgrowth of the neuroblastic elements there may be coincident increase in the true connective tissue elements, but this is to be regarded not as a neurofibromatosis, but as parallel with what occurs in scirrhous cancer. There may also be serous infiltration and œdema passing on to the formation of cystic or lymphangiectatic cavities, mucinous infiltration, hyaline degeneration of the specific tissue, or, contrariwise, active vegetative growth of the cells, so that certain of the nodules take on a sarcomatous appearance.

CHORDOMA.

This is a remarkable form of tumor, first noted by Virchow, and regarded by him as cartilaginous; fully studied by Ribbert. Remains of the notochord are to be found in the intervertebral disks as small collections of large vesicular cells separated by a homogeneous interstitial substance. As in cartilage, there are no vessels. They rarely form here anything that may be regarded as a tumor, but there is one site where a small tumor, never attaining great size, and formed by overgrowth of these cells, is to be found, and that, on careful examination, not infrequently-according to Ribbert,² in 2 per cent. of all autopsies. This position is the Clivus Blumenbachii at the spheno-occipital synchondrosis, corresponding to the original upper end of the notochord, behind the pituitary body. Here the growth, originating in the bone in the middle line, is apt to penetrate the dura mater and project as a mass the size of half a pea, often intimately attached to the basilar artery; so that, on removal of the brain, the fine pedicle in the dura is apt to rupture, and the tumor be found hanging to the artery. This little tumor is

vevpov, a nerve; iç, ivóç, a sinew or fibre.
 ² Centralbl. f. Pathol., 1905.

see and

PSS

iew

seg-

DW.D

elv.

ight

hn.2 ner-

010-

tese

re-

ally

ob-

the

ded

and


1 as

ves.

ngly

be

composed of tissue distinct from cartilage, and showing all the characters of notochordal tissue. From the presence of interstitial substance between the cells, on full consideration, we cannot, with Minot,¹ regard



Section of a chardona. To the right the cells are of the being type, not unlike in arrangement to those of cartilage; to the left through active multiplication the cells are taking on a more screomatous type and the growth is becoming malignant. (Fischer.)

these cells as strictly epithelial, and must class the tumors thus formed as hypoblastic hylomata.

¹ Adami, Jour. of Pathol., 1902:216

CHAPTER XX.

ATYPICAL HYLIC TUMORS.

SARCOMAS.

I HAVE already of necessity referred on several occasions to the sarcomatous growths when discussing the typical form of hylic growth. Indeed, I heartily agree with Mallory that the more rational procedure is not to treat the atypical apart from the typical form of any tumor, but with each tissue in turn to discuss the whole series of growths that may originate from it. Nevertheless, in a work of this nature such a course would necessitate undue repetition: it makes for conciseness of handling to discuss the sarcomas as a class. Here, before writing of these extensively, it will be well once more to lay down what we understand by a sarcoma. Let me repeat that the term has nowadays first and foremost a histological significance. (1) First and foremost a sarcoma is a richly cellular tumor of the connective-tissue type, the cells being of the vegetative, imperfectly differentiated order, or "embryonic;" and the component cells develop and present characteristically interstitial substance. This may be minimal and little beyond granular matter, but careful examination of different parts of a tumor will show that cells of identic nature show between them here and there such granular passing into definitely fibrillar interstitial substance. We have, that is, the hylic arrangement. (2) Such arrangement is not confined to tumors derived only from the mesoblast (whether mesenchymatous or mesothelial); it is characteristic of certain typical and atypical tumors of epiblastic and hypoblastic origin. Therefore, certain atypical epiblastic tumors must also be regarded as sarcomas, and as we shall show later, the actively growing tumors of transitional lepidic characters have also from this standpoint to be included as sarcomatous. (3) Secondarily we have to give to tumors possessing these characters the clinical significance of infiltrative growth and the possession of malignant characters. But in doing this we must always keep in mind that malignancy depends upon more than the mere form of cell present; of two tumors composed of equally small round cells, one may exhibit rapid generalization, the other may be at most locally malignant. The tissue of origin, where it can be determined, should largely influence our diagnosis. At most, we can lay down (1) that the more embryonic the type of cell the greater the presumptive evidence of malignancy, and (2) that as between two tumors of the same origin the more vegetative the type of cell and the greater the departure from the adult cell standard, the greater is the malignancy.

ATYPICAL HYLIC TUMORS

All such sarcomas present certain features in common. They are not encapsulated, but exhibit a peripheral growth and invasion of the surrounding tissues. This invasion is along the tissue spaces and leads to progressive destruction of the preëxisting tissue, with general absorption of all that tissue save a supporting framework around the vessels and capillaries. Sometimes this is not so extreme, and so we obtain one form of so-called alveolar sarcoma, in which the tumor cells are arranged in groups separated by well-marked connective tissue. We recall a case of Professor Delépine's in which infiltration of the diaphragm and replacement of the muscle fibres by advance of the sarcoma within the sheaths gave this appearance with singular clearness. But even in such cases examination of the primary growth, or of the central area of the tumor mass, shows that this appearance is secondary, only the capillaries and vessels being eventually left, with a small amount of the preëxisting connective tissue.

The sarcoma cells, in short, grow in the immediate neighborhood of the capillaries. This is a marked feature of all sarcomas. We observe throughout the tumor that the vessels are composed of a single endothelial layer, immediately beneath which are the tumor cells. The capillaries may be widely dilated; in fact, another feature is the *abundant vascularity* of the growths.

While it is difficult to convince one's self over this point, it is generally accepted that there is a new formation of capillaries, and that the surroma cells grow along these, just as the fibroblasts appear to extend outward among the growing loops of granulation tissue; in fact, the close relationship between the sarcoma cells and the capillaries closely resembles that seen in granulation tissue. In certain small round-celled sarcomas we occasionally encounter channels, blood-vascular, that are bare of endothelium as through here the blood makes its way directly between the tumor cells.

From these relationships it will be readily understood that (1) hemorrhages into the tumors are very apt to occur, and (2) that sarcoma cells are liable to become free in the blood stream, and that metastases along the blood stream are characteristic of these growths. Such metastases, it must be remembered, are not confined to the blood-vascular system; they may occur along the lymphatics, so that malignant enlargement of superficial and other lymph glands is not absolutely diagnostic of cancer.

Borst ascribes this liability to lymphatic extension especially to small round-celled sarcomas of lymphosarcomatous type. Our own experience leads us rather to the conclusion that, while undoubtedly sarcomas of that type show this tendency, all sarcomatous growths of the abdominal area, whether small round-celled, small spindle-celled, or mixed-celled sarcomas derived from pluripotential mixed tumors of one or other abdominal viscus, may form such metastases. We recall also a case in which, in an arm amputated at the shoulder by our colleague Dr. James Bell, Dr. Keenan found extensive osteosarcoma with bone formation in the axillary glands.

But extension by the bloodvessels is undoubtedly the commonest pro-

PROPERTIES

cedure, and thus it is that secondary sarcomatous growth is peculiarly apt to show itself in the lungs. So also it must be noted that the growth may directly invade and grow along the bloodvessels.¹ Apart from such continuous growth, relatively large cell collections may become detached, and may even grow free within the heart cavity (as in a well-known case of malignant mixed tumor of the testis recorded years ago by Sir James Paget²), eventually gaining secondary attachment.

According to Ziegler, the sarcomas possess no lymph vessels proper, only occasional spaces and channels.

It will be readily understood that such rapidly growing tumors present abundant mitoses. Irregular mitoses, as also cell inclusions and socalled cancer or sarcoma parasites, which we must regard as a sign of degeneration, also occur, but not so frequently as in cancers. Indeed, the variety of degenerative changes is not so marked a feature as are the necrotic changes and death of the cells which affect portions of the growth, often associated with hemorrhages and pigmentation.

Forms of Sarcoma.—In accordance with what I have stated regarding cell differentiation and vegetative activity (pp. 141, 670), it will be recognized that lack of cell differentiation is to a very large extent accompanied by retention, or acquirement, of increased vegetative activity. We now may be prepared to find that the stages of undifferentiation, or *anaplasia*,² in the different forms of tissue are not wholly identical. A glial cell, for instance, in its development never passes through a spindle-celled stage; thus, vegetative glial cells never passes through a spindle-celled stage; thus, vegetative glial cells never produce a spindlecelled sarcoma; the mature lymphocyte is a smaller cell than the vegetative mother cell which produces it; thus, lymphosarcoma formed of vegetative lymphoid cells may be of larger cell type than the adult lymphocyte, and, it also, is not of the spindle-celled type. Only cells which in the course of their (normal) development pass through a spindlecelled stage can give origin to spindle-celled sarcoma—connective-tissue cells, plain muscle fibres, etc.

And here a common misconception must be noted, that it is inconceivable that a typical, fully developed tissue, or a typical blastoma, formed of well-differentiated cells, should become converted into unripe sarcoma tissue. Birch-Hirschfeld laid it down that such an anaplasia is neither probable nor proved, and Ribbert and Borst reëcho the sentiment. Such an argument shows a want of realization of vegetative

¹This seems to be a feature of the not very common pure sareomas of the kidney; both Wyler (Diss. Zurich, 1897) and Borst quote cases in which such growth has extended into the inferior vena cava, and so into the right auricle, and we have met with a similar case; the specimen is in the museum at McGill College. MacCallum (Johns Hopkins Hospital Reports, 9) has reported a case in which a malignant tumor of the left testiele extended without a break along the left spermatic vein, left renal vein, and also along the left lika veins, the growth passing into the inferior vena cava to above the diaphragm. According to our present knowledge this ought, however, to be classed, from his description and figures, as a chorio-epithelioma of the testis rather than with the sareomas proper.

² Also now regarded as a chorio-epithelioma of the testis. ³ See pp. 641 and 841.

id of erve othearies arity rally roma ward tionnbles omas re of ween emori cells along ses, it : they supersmall rience has of minal -celled

SHT-

s to

and

one

iged

II a

and

the

a of

the

f the

nation st pro-

other

ase in

James

ATYPICAL HYLIC TUMORS

Underlying it is the idea that the fully differentiated cell processes, gives origin to the fully differentiated cell. The whole study of regeneration shows that this never occurs. Either there are undifferentiated mother cells, or cambium cells, normally present, from which the differentiated cell is developed, or, as in muscle and many other tissues, to become vegetative the differentiated loses largely its specific features and passes back into the simple vegetative type. It is perfectly conceivable that in a highly developed tissue, or in a typical Hastoma, certain cells can lose their specific properties and revert to a simpler stage; that as the regenerating muscle fibre reverts toward the sarco lastic type, so the cells in a rhabdomyoma, being unable to function, are therefore all the more liable to lose their functional differentiation and assume a vegetative sarcoblastic type. In this there is nothing improbable. According to the hereditary characters impressed upon them, according also to the surroundings in which they find themselves, so will such cells attain to a certain stage of differentiation. And so a tumor may show any stage, from the very lowest vegetative round-celled type up to the (not quite perfectly) differentiated tissue cell. Being under abnormal conditions and unable to function normally, the tumor cell can never, and never does, acquire perfect differentiation.


These vegetative or "embryonic" types of cell are simple and their range is comparatively small, from the small round cell, to that with larger amount of cytoplasm and rounded nucleus, to the oval cell, likewise with relative abundant cytoplasm and oval nucleus, and the spindle cell, still larger, with oval or even spindle-shaped nucleus and relatively less cytoplasm; though here we note a difference: we may have a small spindlecelled or a larger spindle-celled type of cell. In this way we distinguish the several forms of sarcoma, (1) small round-celled, (2) round-celled, (3) large round-celled. (4) oat-shaped cell, (5) small spindle-celled, and (6) large spindle-celled. We classify according to the predominant type of cell. Where, as in one order of growth, we find considerable variation in type we speak of (7) the mixed-celled sarcoma.¹

All these we may speak of as *pure* sarcomas. In addition there exist the *intermediate* sarcomas, in which the undifferentiation has not proceeded so far, so that some of the constituent cells attain a considerable degree of differentiation and tend to reproduce the tissue characteristics, whereas others are of the actively vegetative type. These are mixedcelled sarcomas in another sense. In this way we recognize the fibrosarcoma, osteosarcoma, chondrosarcoma. The terms *fibroma sarcomatosum*, *osteoma sarcomatosum*, etc., would more accurately express the nature of these tumors. It is from the study of these transitional forms that we learn to recognize the origin and relationships of the different types of pure sarcoma, for in them we see the different anaplastic stages exhibited by cells of one order. It cannot be said that this field has as

⁴As noted (p. 753), I do not place the so-called "giant-celled sarcoma" here, but among the myelomas; the melanotic sarcomas are also considered separately (p. 825).

VARIETIES OF SARCOMA

yet been adequately worked; the finer details of the individual cell forms, and equally of the intercellular fibrils, in these intermediate sarcomas need a fuller study in order to afford adequate data for a sure recognition of the tissue of origin of the pure forms. We have thus far been satisfied to regard the sarcoma as an atypical connective-tissue tumor and leave it at that. The matter becomes one of practical and diagnostic importance once we admit, as we must, (1) that with a given tissue cell the more vegetative the type the greater the malignancy of the tumor, and (2) that the stages through which the cells of one tissue pass to attain full differentiation differ from those affecting the cells of another tissue, so that (3) tumors which superficially appear to be composed of cells of like size and arrangement, if derived from different tissues, may vary widely in malignancy.



Small round-celled sarcoma, infiltrating liver, advancing along a portal sheath: V P, portal vein; B D, bile duct; A H, hepatic artery; L, liver cells.

The Small Round-celled Sarcoma.—The small round-celled sarcoma is in general the most malignant of all, or, concisely, the most intensely malignant and infiltrative growths with which we become acquainted are members of this class. The closely packed cells have deep staining, round nuclei with little cytoplasm; the interstitial reticulum is at a minimum; the arrangement of the cells immediately beneath the vascular endothelium is very characteristic. They are extremely vascular and liable to exhibit hemorrhages here and there throughout their bulk. Metastases occur both through the blood stream and along the lymphatics.

They originate, it is generally held, from connective tissue in the most various parts of the organism, and the similarity they present in general to granulation tissue is striking. It is, however, possible that they represent the least differentiated, most actively vegetative stage in the development of all tissues. We may recall that the simplest, most vegetative type of neuroblast cell is of this order (p. 757).

' here, arately

era-

ren-

be-

able

cells

s the

the

| the

reta-

ordalso

cells show the rmal , and

arger with , still cytondleguish elled, , and

type .

exist

pro-

rable

istics.

nixed-

coma-

is the

forms

stages

ias as

Of the ordinary *round-celled sarcoma* nothing can be stated with definiteness, save that it also is actively malignant, but not to the extent that is the previous form. It also appears most often to originate from




Large round-celled sarcoma. (Ribbert.)

connective tissues. Its cells, while small, are not strikingly so; their cytoplasm is more obvious.

The Large Round-celled Sarcoma. —The large round-celled sarcoma shows, however, distinct difference in type, and we regard it as belonging to another class. The cells impress one as being distinctly large and of the more "epithelioid" type; *i.e.*, with abundant cytoplasm forming a cell body not perfectly round, but rather variable in shape, now suboval, now obscurely polygonal; the nuclei tend to be suboval, paler,

larger, and more vascular than those of the previous forms. The reticulum is more marked, and there is a slight connective-tissue stroma, nay, more, the peripheral portions of the growth are apt to take on an alveolar



Oat-shaped cell sarcoma. (Leo Loeb.)

type. There is not the same extensive destruction and absorption of the tissue infiltrated as we note in the previous forms. Such tumors are often found in connection with striated muscle. It is at least possible

VARIETIES OF SARCOMA

that one group represents the most vegetative form of rhabdomyosarcoma, though more evidence is required upon this point. Another group of forms originates in the testis, with characteristically large cells, and derived possibly, as suggested by Hansemann, from the large interstitial cells of this organ (they thus are of mesothelial origin). The least differentiated and most actively vegetative form of lymphoma presents this order of cell.

The Oat-shaped Cell Sarcoma.—The oat-shaped cell sarcoma is not very usual, but when encountered is a very characteristic form, because of the regularity of the blunt cells with long oval nuclei which form the main mass of growth. We have not been able to satisfy ourselves that this form is characteristic of sarcomas originating from any particular tissue, save that in one case, as already noted (p. 747), we have possibly traced a plain muscle origin for this type.

The Small Spindle-celled Sarcoma.—The small spindle-celled sarcoma is, as the name implies, formed of relatively small spindle cells, varying in

length from 10 to 20 μ (whereas in the large spindle-celled growths the cells are often from 50 to 80 μ long). The cell nuclei are oval or, like the cells, spindle-shaped, the cells collected in bundles with surrounding stroma, and these bundles appear to conform to or surround the capillaries of the growth. Here and there one notes often that some cells have produced definite fibrillae.

The appearance so closely resembles that of organizing cicatricial tissue that we have no hesitation in regarding this form as of connective-tissue origin, and, as a matter of fact, these forms are found in connection with connective-tissue areas, the corium, fascia,



Large spindle-celled sarcoma. (Ribbert.)

etc., while, further, both, fibroglia and collagen fibrils characteristic of white connective tissue can be detected in them. Cells with similar fusiform nuclei, but with less definite cell bodies, are encountered in the neurinomas, here associated with fine nerve fibrils. Compared with the round-celled forms, these are more benign. Metastases may occur, but are rare.

Large Spindle-celled Sarcoma.—The differences between this and the small spindle-celled form are very largely parallel to those between the large and small round-celled groups. The nuclei are larger and clearer, often vesicular, and a variability is noted in the size and shape of the cells, as also in the terminal processes, which may be simple or forked. The tumors are found in connection with periosteum⁴ (the large spindle cells of the giant-celled myeloma may here be recalled), fascia, and either the connective tissue of muscles, or it may be, from striated muscles themselves.

¹ In this type of large spindle-celled sarcoma the cells, while large, are in general "stocky" and not of great length.

49

defictent from vhile their

oma.

oma ence ongimarge ype; prmund, now mal; aler, ticunay, solar

> the are aible

ATYPICAL HYLIC TUMORS

We refer here more particularly to a group of large spindle-celled sarcomas, in which, besides spindle cells of great length somewhat irregularly disposed, and possessing often two or it may be three nuclei, we encounter also large oval or irregularly shaped cells recalling the sarcoblasts.

Intermediate Types.—Fibrosarcoma.—It is singularly difficult to draw the line between the fibroma proper and what we would term the fibroma sarcomatosum, and this because the ordinary fibroma in general is more cellular than ordinary fibroid tissue. It thus become a matter of individual experience to decide when this cellularity is sufficient to label the tumor as fibrosarcoma, and so attribute to its malignant characters. A well-marked form may still be well encapsulated, but show abundant "naked" spindle cells along with cells, still spindle-shaped, that have attained to the stage of forming interstitial bundles of fibrille. Such tumors are often singularly rich in large dilated vessels, and are of soft consistency.

Myxosarcoma.—In such, while the main mass of the tumor may show a rather richly cellular myxomatous appearance (the individual cells presenting the characteristic processes), here and there are islands and masses of more closely collected round cells of fair size unprovided with processes: less differentiated cells, evidently more rapidly growing. Such tumors are apt to increase in size rapidly and to form metastases.

Liposarcoma.—A lipoma, after growing slowly for years, may take on more rapid growth, and with this on removal may show one or more areas of sarcomatous change, in which the fat cells become replaced by a richly cellular tissue. In one such tumor examined by us the cells and their nuclei were of an oval type. Rindfleisch would restrict the term to a class of cases in which a round-celled tumor of sarcomatous type shows throughout cells having the tendency to become infiltrated with fat in the form of larger or smaller globules.

As will have been gleaned from the general treatment of the subject, we wholly fail to see the propriety of this ruling, or that it accords with fact. The doctrine upon which it is based, of absolute fixity of properties on the part of tumor cells, is untenable. Once it is admitted that a metastasis, while retaining the same basal characters, may be of a simpler, more vegetative type than the parent tumor, it becomes illogical to hold that in that parent tumor the same process cannot occur and even be progressive until certain cells acquire the most undifferentiated characters.

Chondrosarcoma.—The various stages of undifferentiation are, indeed, frequently exemplified in a rapidly growing chondroma. We see there in the centre of an area unmistakable cartilage, though more cellular than is the normal tissue. At its edge we note the cells still more abundant, and here the chondriform interstitial tissue becomes replaced by a more mucoid matrix, and the cells become stellate. There is no sharp boundary, no island of cells possessing other properties, but a gradual transition from the more highly differentiated—

INTERMEDIATE FORMS

a reversal, we may describe it, of what occurs in the normal condition of cartilage. Passing farther out we have every transition to larger cells, still more closely packed, without processes, or at most of blunt spindle shape; a true, rather large-celled sarcoma, the cells exhibiting abundant indications of active growth.

The contrary view, that the tumor is from the first essentially chondrosarcomatous is incompatible with the fact that such tumors present, along with the development and vascularization of this sarcomatous tissue, evidence of progressive removal of the cartilage. As the vascular sarcoma tissue becomes formed, here and there it can be seen advancing into, absorbing, and replacing the previous cartilaginous substance. The first stage has been that of cartilage formation, the latter that of sarcomatous modification.

the

to

101

llæ.

· of

WOI

nds

mo-

· on

hly

OWS

t in

per-

at a

pler.

and

eed,

here

by a

ed-

Osteoid Sarcoma, Osteochondrosarcoma, and Osteosarcoma.—By these terms we distinguish three different types of intermediate sarcoma, exhibiting different grades of the ossification process.

The osteoid sarcoma is fairly common, and is, as regards malignancy, a true sarcoma, growing rapidly and forming metastases. In it we find areas which it is best to describe as intermediate between cartilage and bone. There is a homogeneous, cartilage-like matrix, but the cells in this, where single, resemble more bone corpuscles than cartilage cells; often there are several in one space, and where this is so they show all transitions to the sarcoma cells, surrounding thickly the osteoid lamella or mass. These cells are polymorphous, and away from the lamellae giant cells occur. One cannot study such a tumor without being convinced that the osteoid tissue is an integral portion of the tumor mass, that it is the tumor cells which have produced and governed the deposit.

From this we pass to cases a stage less undifferentiated, in which there is deposit of calcareous salts in the lamellae in certain areas; we are a stage nearer to true osteosarcoma. Other cases exhibit areas both of true cartilage and true bone; for these the name osteochondrosarcoma should be reserved, although the last-mentioned form is evidently very closely allied.

The ostcosarcoma proper shows lamellae and masses having the chemical composition of true bone. The histological picture may be, nay, generally is, imperfect—imperfect lamellation—and, while showing corpuscles, these have not the typical branching character. But, nevertheless, it must be regarded as true bone to the same extent as the fibres in a myoma are true plain muscles fibres. It is what von Hansemann describes as rudimentary bone formation. This may be present in irregular isolated spicules, or as a thin irregular spongy mass, well shown when the tumor is macerated; or, as in periosteal osteosarcomas, as a series of radiating spikes, osteophytes, adherent to and apparently growing from the shaft.

It is interesting to note how the adherents of the fixity of tumor-cell properties theory dispose of this bone formation in these surcomas. The presence of true bone as an actual intimate constituent of the tumor proper cannot be allowed. It is, according to Borst, following Rind-

ATYPICAL HYLIC TUMORS

fleisch, (1) calcified intercellular substance, and it is to be distinguished from (2) the true bone, the stroma-like remains of the bony tissue invaded by the sarcoma, or (3) a reactive inflammatory proliferation of the bone involved by the sarcoma. The benign ossifying forms, such as we see developing from the periosteum, cause superficial erosion of the shaft of the bone and a reactionary osteophytic inflammatory reaction. It is not explained (because it cannot be) how this inflammatory growth of the osteophytes is brought about; that, for example, each "osteophyte" is surrounded by a normal vascular tissue, with normal osteoblasts which deposit the new bone corpuscle and bone layers on the surface of the same, and so bring about increase in length. As a matter of fact, the osteophytes are directly surrounded by the tumor cells, and the tumor cells, and they only, can give rise to the new-growth. As in normal bone there is a layer of cells around the vessels which do not themselves undergo ossification, whereas away from the vessels these cells or their descendants become osteoblasts, so in this order of tumors the cells have not become so completely undifferentiated that they cannot still, in certain surroundings, manifest a certain amount of functional activity. Borst has to admit that in osteoid sarcomas the cartilage-like ground substance is a constituent portion of the tumor, and does not attempt to explain how the metastases in the lungs and elsewhere contain bony elements.

This group of osteoid sarcomas exhibit most often mixed sarcoma elements: spindle cells large and stumpy, polygonal cells varying in size, giant cells, the latter not so frequent, as a rule, in periosteal, superficial sarcomas as in central growths. Such central growths are apt to be expansive, causing absorption of the shaft and spontaneous fracture. With the growth there is still some periosteal bone formation on the surface, as indicated by the "egg-shell crackle" over them. In general they do not form metastases until the periosteum becomes ruptured and infiltration occurs of the surrounding tissues. In general, also, it may be laid down that the greater the development of osseous matter the less is the malignancy of the tumor. Thus the very "osteophytic" periosteal sarcomas of the long bones and face are only mildly malignant. The most malignant cases in our experience are those which show least bone and most cells of the small type.¹

Lymphosarcoma.—This form we have already discussed. We will here only recall (1) that it is distinguished from the round-celled sarcoma proper by its more marked reticulum, well brought out by pencilling, or by washing the section in running water, and (2) that the larger-celled form (equivalent in size to the ordinary round-celled sarcoma) is of the more vegetative type. Such growths have a tendency to be local and infiltrative and to form metastases in the nearest lymph glands.


Leiomyosarcoma.—These again we have already noted (p. 747). It is probable that a group of spindle-celled sarcomas of the uterus and

¹ For a careful study of bone sarcoma, see Buerger, Surgery, Gynecology, and Obstetries, 1909; 431.

INTERMEDIATE FORMS

alimentary tract—possibly also of the genito-urinary tract—come under this category, tumors exhibiting moderately large spindle cells, variable in length, down to the blunt out-shaped form.

Rhabdomyosarcoma.—The pluripotential tumors of the kidney and other regions are apt to take on this type, to exhibit, that is, large and very long, imperfectly formed muscle fibres, which may in part show transverse striation, others of long spindle shape showing only longitudinal fibrillation, large polymorphous and often multinucleated cells of the sarcoblastic type, and with these cells with one or two nuclei of the epithelioid type. Whether all the latter are of sarcoblastic origin remains an open question. We are inclined to consider that some of the cases of "mischgeschwülste" of the simpler type reported as occurring in the vagina and genital tract are tumors composed of sarcoblasts of different



A. From the more typical portion of a glioma. B. Another region from the same growth of more malignant type, a true gliosarcoma. (Thomas and Hamilton.)

grades of development. The existence of this group renders it possible that a group of large spindle-celled tumors of muscle showing also great irregularity and some polymorphism may be sarcomata derived from muscle elements. (For the giant-celled form see page 751.)

Gliosarcoma.—We have noted this form in our discussion of the gliomas, and pointed out that (1) we regard the term as of perfectly correct usage (though we prefer the alternative Glioma sarcomatosum); (2) it is of most common occurrence in connection with the retina, and here is malignant, with infiltration, and liable to form metastases. The features of such a form are, briefly, that it is composed of small round-celled elements, in the main indistinguishable from those of ordinary small round-celled sarcoma. Careful investigation shows that some of these

ATYPICAL HYLIC TUMORS

retain, though imperfectly, the glial processes. The cells are especially well stained around the vessels of the tumor, *i. c.*, those at a distance are apt to degenerate. Here and there the peculiar rosette-like arrangement of the cells, characteristic of the glioma proper, is to be detected.

Greeff has even made out in these retinal tumors the existence of imperfect, undeveloped nerve cells, and ascribes the origin of these growths, we think justly, to cell rests of embryonic nerve tissue within the retina.¹

³ For reviews of the more recent literature upon individual forms of tumors the reader is referred more particularly to Lubarsch and Ostertag's Ergebnisse der allg. Pathologie, in which every other year or so there are afforded admirable studies upon neoplasia.

CHAPTER XXI.

PRIMARY LINING MEMBRANE, OR LEPIDIC TUMORS (LEPIDOMAS).

It has become the fashion of late years to speak of these as fibroepithelial tumors. We doubt the utility of the term. It is true that there is always more or less of a fibrous connective-tissue stroma, but this is secondary, even though, through its variation in amount it affords us to some extent a classification of certain forms, and although its presence is characteristic. But it exhibits no independent blastomatous growth of its own, save in the singularly rare cases which we shall have to note later. The essential part of all these tumors is the epithelial or glandular, or what we have termed the lining membrane element; it is this that takes on independent growth. For that growth the stroma, containing vessels, is essential, and, what is more, the very presence and activity of the "lining membrane" elements influences or sets up proliferative changes in the stroma, but these, at most (with the rarest exceptions), of an irritative, non-blastomatous type. These tumors, then, are best understood when we regard them as essentially of epithelial or lining membrane type. We can, as usual, divide them into the typical and the atypical.

TYPICAL LEPIDIC GROWTHS.

PAPILLOMA.

In these we have to deal with outgrowths from surfaces presenting a covering layer of epithelium, whether squamous or columnar, and having a more or less pronounced connective-tissue core to each individual process.


1. Of J'ritative Origin (Non-blastomatous).—Here, again, we must rer we a group of cases which, as their growth is obviously due to arritation, cannot be regarded as true blastomas, though, as these are papillary tumors, the name papilloma still adheres to certain of them.

(a) Warts.—In these we deal with overgrowths of a collection of papille of the corium, covered by a common, thickened, and somewhat hypertrophied epiderm. They would seem to arise from irritation, are commonest in childhood, and have a marked tendency to disappear eventually. Some ascribe to them a definite infective origin, and clear evidence has been brought forward of transmissibility of the condition.¹

¹Jadassohn, Verhandl. d. deutsch. dermatolog. Kongr., 1898; Lanz, Deutsch. med. Wochenschr., 1899; Nr. 20.

This may, however, indicate mere transplantation, although, as Lanz was able to produce them upon the tip of his finger by rubbing it over the wart of a patient, an infective origin appears the more likely.

(b) Molluscum Contagiosum.—This is a definitely contagious skin disease affecting the face and head, the hands, and other parts of the body, which presents itself first in the form of small red elevations, which grow into warty elevations, continually breaking down in the centre and discharging whitish, cheesy matter. The growth is not so much superficial, according to Kaposi, as deeper, affecting the epithelium of the sebaceous glands, or (O. Israel) the hair follieles. The



The various grades of warts and cutaneous papillomas. (Perls.)

surface layer, indeed, may show little hypertrophy; only, therefore, from its warty appearance does the condition come to be considered here. Characteristic bodies are found in the affected epithelial cells, regarding which there has been much debate as to whether they are parasitic protozoan forms or merely cell degenerations—or both.

In a somewhat similar condition affecting the head and comb of the fowl, it would seem evident that there are parasitic organisms present and setting up the disturbance. The consensus of opinion at the present time tends rather toward the degenerative view regarding the molluscum corpuscles; they are classed with the cancer bodies (see p. 793).

The Pointed Condyloma.—This projecting epithelial overgrowth occurs more especially as a multiple development upon the external genitalia-vulva, vagina, penis-or, again, in the anal region, or, more rarely, in the mouth.

It presents itself as a warty, nodular, sometimes mulberry-like, or even cauliflower-like growth, the outer wall of which is formed of thickened, overgrown, squamous epithelium, lying upon a stock of hypertrophied connective tissue, vascular, or very often showing small-celled infiltration. There is always a history of irritative discharges bathing the part, and in almost all cases in man one of chronic venereal disease to be gained. It is deserving of note that, while themselves of a benign type, they may eventually become the seat of epitheliomatous developments. On the other hand, if in the early stage the source of irritation

be removed, their growth is apt to be arrested. If left, the growth becomes progressive and independent, and may attain an extraordinary extent, with abundantly branching processes, as upon the penis, where it may completely encircle the glans. We here deal with a borderline condition.

Cutaneous Horns.-We occasionally encounter these, more often in the old than in the young-very slowly developing processes of true horny matter, often of bizarre shape, projecting from one or other region (most often of the scalp and face), and movable, their bases being soft.

They represent a condition of hyperkeratosis, or excessive development of the keratinous matter, an overdevelopment of the horny laver of the skin, coupled with a failure of the scales to peel or be rubbed off, so that they accumulate and form these Condylomata of the vulva. (Orth.) masses. But as the horn grows in length



it is to be noted that the underlying skin papilla, or papilla, become elongated, to form a vascular core, passing, it may be, almost to the end of the growth.' An overgrowth of the epithelium appears here to be the primary event, but as the process in the older areas appears to be self-limiting, the cells throughout the whole thickening of the epidermis becoming keratinized, and the stratum Malpighii almost completely disappearing, it is difficult to regard this as a true blastoma.

In reference to the relationship between inflammation or irritation and papillomatous growths, two interesting conditions have to be noted-coccidiosis and the disturbances caused by the ova of the worm Bilharzia.

Coccidiosis.—In the ordinary rabbit bred in captivity it is very common to find in the liver a variable number of rounded, whitish nodules, some hardly visible to the naked eye, others reaching the size of a pea, and even larger, and distributed along the branches of the bile ducts. When

large, these are soft and not unlike small abscesses, and when the soft material is removed from the centres of the nodules, a cystic cavity is left, with a papillomatous wall.

Upon examining these under the microscope, it is seen that we have a very remarkable condition of dilatation and proliferation of the bile ducts. The surrounding fibrous tissue is considerably increased, and the overgrowth of the epithelium is such as to form numerous fine papillary projections; in fact, the appearance is very much that of a cystadenoma.

If, now, we examine still more carefully, many of the individual columnar cells of this epithelium are seen to contain inclusions, and these inclusions cause the cells to be, many of them, greatly enlarged.



Section of portion of the wall of a coecidial cyst in the liver of a rabbit: a, fibrous capsule; b, proliferated epithelium of bile duct, with papillomatous outgrowths; c, coecidia free in lumen.

If we follow the series of stages, at first little can be made out beyond that there is an indefinite body in the cell substance; but, as this grows, it gains a definite wall of double contour, and takes on a somewhat oval shape. Outside the body, kept in a moist condition, sporulation takes place, and four spores are produced, in each of which there develop two somewhat crescentic germs. These germs become anneboid, and in this motile state are evidently capable of entering the epithelial cells of a second host, there to repeat the life history. According to Delépine, they may be found frequently affecting the cells of the duodenum of the rabbit, and it is supposed that here the motile forms are elaborated, pass up the bile duct, and into the liver, and again, either directly or through a second generation, enter the epithelium of the bile ducts.

A remarkable part in this disease is that the presence of these small animal forms within the cells leads to a very marked proliferation resembling what we find to occur in tumors proper of the adenomatous

INFECTIVE PAPILLARY OVERGROWTHS: COCCIDIOSIS 779

type. What is further of interest is that there is no spread of the growth apart from the presence of these coccidia, and that there is apparently no general disturbance set up in the vast majority of cases. Again, the presence of these parasites does not necessarily lead to necrosis and death, but rather to proliferation of the cells. As Delépine remarks, "the parasite appears to be almost entirely devoid of any marked irritating properties; its presence leads to a setting up of irritation which only slightly exceeds physiological stimulation, and a result of this slight increase of death among the cellular elements."

Tyzzer¹ also has made a study of coccidiosis in the rabbit. He points out that the parasites attack only the epithelial cells, and that at the termination of the process of growth of the parasite the cell is reduced to a sac containing the parasite, having on one side a darkly stained crescent, representing the degenerated nucleus. Ruptured cells are found from which the parasites have been set free. Degeneration and destruction of epithelial cells thus follow their invasion by parasites. But, adds Tyzzer, "numerous mitoses are seen in the epithelium, and, where the infection is not overwhelming, proliferation is in evidence. The epithelium is markedly thickened and its cells are crowded." His opinion is that the formation of the papilliform projections is to be explained by hyperplasia of the connective tissue, which pushes through defects in the epithelial layer; that both epithelium and connective tissues are stimulated to increase, and that the epithelium proliferates in an attempt to repair the defect in its continuity. For myself, I am more than doubtful whether the epithelial overgrowth is so particularly in relation to previous destruction, or to the papilliform projections of the connective tissue. It is often generalized all around a dilated bile duct, and irregularly of several layers. We have here a case, that is, in which irritation of low intensity, acting in a more concentrated form, leads to cell destruction; where less concentrated, it leads to cell proliferation; where overgrowth both of the epithelial cells and of the neighboring connective tissue is initiated by the coccidial products, the interesting and remarkable feature being that the irritation here leads to localized overgrowth of epithelial elements. In this we have a condition unusual in ordinary inflammation, and more like what we see in epithelial tumors or adenomas. There are, however, two features which would seem to distinguish these coccidial growths from tumors proper: first, the continuance and further growth is directly dependent upon the continuance of the coccidia, so that we frequently come across evidence of old cicatrized areas, showing no coccidia, or containing them in the encysted and resistant stage, the epithelium having undergone complete degeneration; and, secondly, unlike true adenomas, showing cystadenomatous change and atypical epithelial proliferation, we never meet with evidence of metastasis.

Therefore, while coccidiosis is interesting and of importance as

¹ Journal of Medical Research, 2:1902.

indicating the existence of irritants which lead to epithelial overgrowth, it cannot be quoted as affording us examples of true tumors or blastomata of parasitic origin. At most, it can be adduced as one of the intermediate stages between inflammatory and blastomatous conditions, and as an illustration that irritation of low intensity, insufficient to cause cell degeneration, may lead to proliferation of specific cells.

Bilharziasis.—There is, however, another condition of new-growth due to parasites which appears, from all descriptions, to be definitely blas-



Ova of Bilharzia (schistosoma) hæmatobium, to show *a*, terminal; *b*, lateral spike, (Perls.)

tomatous. It is possible that further and more minute studies will demonstrate recognizable differences, but for the present 1 do not see how we are to distinguish tumors of this order from tumors proper, save in that here the direct inciting cause is known. We refer to the rectal and vesical growths initiated by the ova of the Bilharzia.

The parasite is extremely common in Egypt and Abyssinia; the adult female, when mature, shows a predilection for the portal veins, more especially for those of the pelvic area. Here the eggs are discharged, and, passing into the smaller veins, mechanically, through the agency of their terminal spikes, they penetrate into the surrounding tissues. More particularly



Bilbarziasis of the rectum, to show papillomatous overgrowth of the mucosa: H, cavities filled with blood, (Looss.)

are they found in great numbers in the mucous coats of the large intestine and rectum and in the walls of the bladder, a considerable number making their way out, passively, into the cavity of the bladder and into the gut—inducing thereby conditions of hematuria and melena. This passage out of the ova induces chronic overgrowth of the rectal mucosa and the vesical epithelium, so that the mucosa of lower portions of the rectum becomes greatly swollen, in fact, papillomatous, and the same

is true of the inner coat of the bladder. What is more, numerous cases are on record in which this condition of chronic proliferative inflammation has given place to definitely cancerous growths.

ise

110

11

10

ie

We spoke of the assumption of malignant properties in cases of fibromatosis, as in epiphenomenon. It might be said that the same is the case here. The difference in the two is that here we recognize the presence and influence of a continuously acting irritant, which eventually, in a certain proportion of cases, if in action for a sufficiently long period—possibly, also, in those having a special predisposition leads to something beyond mere inflammatory changes—leads to an aberrant progressive and excessive tissue growth, with the proliferation of atypical tissue cells invading the surrounding tissues and, indeed, capable of forming metastases.

A point of not a little interest is that, while the ova are abundant in areas of chronic proliferative inflammation, T. Harris found them absent in the definitely cancerous areas. In other words, while they appear to start the cancerous process, once it is started they do not appear necessary for its continuance. This absence, I am assured by Professor Symmers, who has extensively studied the condition, is not by any means constant.¹ Goebel, from a study of the bladder tumors thus induced, lays down that changes in the connective tissue precede and lead up to the malignant epithelial growth.² This is eminently probable as the eggs are present in the submucosa and not in the epithelium.

2. Blastomatous Papillomas.—(a) Soft Papillomas.—These grow from nuccus membranes, and in general afford the most satisfactory examples of the form of tumor which develops in direct continuity with, and clearly from, a normal epithelial membrane. What is the direct cause of the cells in one particular locality taking on excessive growth is not easy to say. Even if, as is often the case with intestinal papillomata, we gain a history of previous inflammation or of dysentery in cases of so-called colitis polyposa, of gastric ulceration in gastritis polyposa, why in some individuals these conditions set up the overgrowth we can, at the most, suggest. But these matters we will discuss later. Of these papillomas we may have every form, from a simple nodular protuberance of the nuccus membrane, either sessile or subpedunculate, such as we often meet with in the intestine, up to a brushlike mass of delicate long processes, such as may be present in the bladder.

Such growths show themselves in the nasal passage, the stomach, and intestines, gall-bladder, urinary bladder, ureters, pelvis of kidney, and uterus. They show a framework or stock of connective tissue, which follows faithfully the branching of the growth, and is distinctly vascular. A transverse section of one of the finger-like processes of the many branching forms exhibits usually a central artery, with vein,

² Zeitschr. f. Krebsforschung, 3 : No. 3.

⁴ See also Symmers, *Festschrift* by the pupils of Professor D. J. Hamilton, Aberdeen, 1907.

or veins, surrounding a soft connective tissue somewhat infiltrated, and outside this the epithelial layer. That epithelium is apt to show





One of the fine processes of a papilloma of the bladder more highly magnified to show the central fibrous core or stock with vessels.

abundant mitoses. At times it is highly differentiated and very typical; in the intestines it may exhibit abundant goblet cells. But often,



Intracystic papilloma of breast. (Orth.)

especially in the more exposed parts of the growth, it is modified. A papilloma of the bladder, for instance, may exhibit an undifferentiated

Papilloma of bladder to show the long, fingerlike papillomatous outgrowths. (Ribbert.)

round-celled epithelium, resembling a proliferation of the cells of the lower layers.

(b) Intracystic Papillomas.—Another group of papillomas, the intracystic, is found developing in cystic adenomas, filling up the cysts with branching and complicated masses of epithelial processes. Such we find, notably, in connection with ovarian cystic growths and mammary adenomas. More rarely we encounter these in cystic growths of the kidneys and bile passages. As in the other form, these intracystic growths possess a connective-tissue stock-though it is interesting to observe, in ovarian adenomas, that the first stage of papillary overgrowth presents itself as a folding outward of the layer of columnar cells, suggesting that by this means the increased number of cells accommodate themselves; in this first stage nothing beyond basement membrane is present in the fold. The connective tissue and vascular growth into the process is secondary. These grow into the space offered by the outward projection of the epithelium. This must be regarded as the mode of development of all papillomas, even of the most complicated and many-branching forms, such as we encounter in the bladder. Thus, Kürsteiner¹ points out that when the cells of an ovarian cystadenoma "seed" themselves upon the peritoneal surface in their growth they stimulate proliferation of the peritoneal connective tissue which now forms the stock of the developing metastatic papilloma, and Steiner² has similarly called attention to the fact that when the cells of a neighboring papillomatous growth spread over a granulating or other surface devoid of epithelium, under their influence the underlying connective tissue develops papillary processes.

Not infrequently these soft papillomas become the seat of cancerous growth, with increasing proliferation and accompanying changes in the character of the cells. In certain areas the cells grow inward, instead of outward, and proceed to infiltrate. A study of papillomas exhibiting the early stages of the change is most instructive (Hauser). To this we shall refer later.

ADENOMA.

By adenoma we understand a growth composed of relatively typical glandular epithelium, arranged, that is, according to the manner in which it is found in the mother tissue; if that tissue be composed of glandular acini, with definite lumina, there the adenoma is likewise in the form of acini, with lumina; if, as in the liver, the acini are formed of solid cell masses, then the adenoma is of solid cell masses without lumina; if developing from duct epithelium, then the adenoma partakes of the character of duct epithelium. The form exhibited by the different varieties of adenoma is thus capable of very considerable variation; each has to be considered in relationship to the tissue of origin. Nevertheless, certain features are common to all.

¹ Vireh. Arch., 130: 1893: 463.

n,

ited

2 Ibid., 149: 1897: 307.

Exhibiting relatively slight anaplasia, such growths of secretory cells are apt to retain some power of secretion. Adenomas of the digestive tract still exhibit goblet cells and discharge mucus; of the thyroid, many still form colloid; of the liver, still produce bile; and, as such growths are independent, where contained within the tissues and encapsulated they are incapable of discharging their secretion, which is apt to accumulate, and, distending the constituent tubules, to form cysts—*cyst-adenomas*.

Here we immediately encounter the great crux in the study of this order of growths; although histologically of the same grade of development, of two tumors of the same organ one may become cystic, the other not; some tumors are sharply encapsuled from the mother tissue, and, in fact, may lie heterotopically, far removed; others, while



Adenoma of bile ducts, formed of acini resembling those of normal bile ducts. (Watzold.)

apparently equally encapsuled, but within the mother tissue, clearly (as in the so-called adenomatous hypertrophy of the prostate) retain through their ducts a communication with the body of the gland from which they have developed, and are still able to discharge their secretions. This is notably the case in the polypoid adenomas of the digestive tract, the nasal cavities, etc. From the frequent communication and direct connection with the surrounding normal tissue it is impossible by any means at present at our command to make a separation between adenomas of this order and conditions of localized glandular hyperplasia, in which, likewise, the communication with the exterior is not arrested. Given two sections, for example,—one from a case of diffuse chronic fibroid induration of the mammary gland of known irritative origin, another taken from the centre of a localized encapsulated fibro-adenoma

ADENOMA

of the manuma of this type, and it is frequently impossible to determine which is which. The same is notoriously true in the liver. We meet with a succession of cases from simple diffuse regenerative hypertrophy, as after acute yellow atrophy, through others in which, as in cases of cirrhosis, the regeneration, while regular, is more localized, confined to separated islands—or peninsulas—of liver lobules (regenerative hyperplasia); to others, in which nodules of liver tissue exhibit an expansive growth, and, growing, cause atrophy of the surrounding liver cells, a definite boundary of capsular nature distinguishing these growths from the rest of the liver tissue. Histological examination of such tumors show the cells of typical character, arranged in typical masses, but these

le

et.


ia.

٠d.

nic

in.

na



Adenoma of liver, formed of columns of cells (c), causing compression of surrounding normal liver tissue (d). (Wātzold.)

masses are not arranged into lobules. The same is true even lower down in the scale, in the regenerative hypertrophy of cirrhosis, but here it becomes even more pronounced, and from these cases we proceed, again by almost imperceptible transition, to others in which the cell growth is more and more irregular and atypical and locally malignant, columns and processes of the tumor cells infiltrating and spreading into the surrounding liver tissue, causing atrophy of the normal cells, the two forms being easily distinguishable by the deeper staining of the nuclei of the invading cells. Cases giving like histories will present one or other of these stages. It is true that by no means all cases of portal cirrhosis exhibit regenerative changes, and that of those which do present 50

them, a very small percentage exhibit the adenomatous type, still fewer the cancerous type of change. There are, obviously, individual differences in reactive and regenerative powers, and these differences—in other words, the tendency or absence of tendency to excessive cell growth—is an all-important factor in determining whether a given insult to the tissue leads merely to an orderly regeneration or to tumor growth. But it is equally clear that simple irritative and regenerative hyperplasia, adenomatous growth, and carcinoma, are stages which can be manifested in succession by the same tissue; that the differences are those of degree and not of kind.

We have, in short, conditions which are largely parallel to those to which we called attention in connection with hylic growths, and here we may make the like distinction between conditions of adenoma proper (independent encapsuled growths), adenomatosis (not properly encapsuled), and irritative and congenital glandular hyperplasia.

It is interesting to observe how the upholders of the unmitigated cell-rest theory, those who hold that all tumors arise from cells congenitally displaced, dispose of these cases. The facts have to be admitted, and are: but the straightforward explanation cannot be accepted. Judgment is suspended, and we are advised that it is necessary to be very cautious; it is suggested that the adenomas of the liver, for example, are not truly adenomas, the cancer not true cancer, and this, although they conform to all the usual postulates; that the irritative adenomas of the digestive tract are not true adenomas, although in the next paragraph, it may be, their liability to become the seat of malignant growths is acknowledged. It is pointed out that there are adenomas which being heterotopic, can only arise from displaced cell masses; that occasionally, in the liver, for instance, we encounter homotopic masses sharply encapsulated, and it is urged that, therefore, invisible and insignificant congenital displacement of cells is at the bottom of all these cases; or, driven farther back, that, although duly placed in the tissues, certain particular cells have from the first had a congenital weakness toward overgrowth. This, it will be seen, is coming very near to our point of view. The cell rest is cast overboard. No sign of it can be seen in simple regeneration of the liver, and equally none in adenomatous hyperplasia of the same.

Is it not more rational to take the view that, while cells (in cell rests) which have never attained full differentiation may, with relatively slight stimulation, take on independent and blastomatous characters, nevertheless fully differentiated tissues have not wholly lost the same power? We see that in inflammation these cells can revert to the undifferentiated vegetative stage and atypical arrangement. Why not accept that under these conditions the same cause that sets up independent growth in cell rests, may set up independent growth in cells produced from differentiated tissues?

We shall have to call attention to a like order of phenomena when we discuss the cancers.

Regarding the structure of the adenomas, it is necessary to say a

ADENOMA AND ALLIED CONDITIONS

-in

ISIA,

gree

· We

oper cap-

conadoted. o be uple, ough omas oarawths hich occaisses and f all o the

t can

ests)

ively

ters,

same

5 the

r not

inde-


cells

n we

ay a

few words. As with the papillomas, so here, the stroma is an essential part. The most typical adenomas show a well-marked basement membrane between the cell layer and the underlying stroma; where growth is rapid and becoming atypical this may be absent. With their growth, also, we must recognize that, while gland cells and stroma are essential to one another, the former are the dominant agents; the growth of the stroma follows that of the epithelium. The appearances seen in ovarian adenomas (see p. 783) can only be explained along these lines. So, also, it is evident, from the results of transplantation of adenocarcinomas in mice, it is the transplanted gland cells that form the new tumors, the stroma being furnished by the new host. The process by which the one tissue follows the other, so as to form an essential whole, is very remarkable. The growth of the stroma, with its accompanying vessels, must be regarded as reactive, as of a chemiotactic nature.

Fig. 253



The so-called fibro-adenoma of the mammary gland. The glandular acial and duets are prominent and show some irregular overgrowth of the epithelium, but the main feature is the develop ment of connective tissue, both peri-acinous and interstitial, the latter not sharply defined. (Ribber,)

It may well be that this reaction on the part of the normal tissue is a factor in the arrest of adenomatous and cancerous growths. Two orders of conditions have to be recognized. If, on the one hand, a given cell, entering a tissue, induces no reaction, its proliferation becomes arrested after attaining a certain point, because no vessels and stroma enter into the cell mass. If, on the other hand, an excessive reaction is produced, then the connective-tissue overgrowth cuts off the nutrition of the developing new-growth. In other words, as we have already pointed out, the development of a blastoma is the resultant of two factors, the proliferative capacity of the invading cells and the reactive properties of the organism.

We notice that the amount of stroma in the different forms of adenoma varies very greatly; it may be of the very slightest, little more than a network of vessels, with their supporting connective tissue; it may be so dense as to be the main feature in a growth. Then we speak of a *fibro-adenoma*—although, from the above considerations, *fibroid adenoma* (adenoma fibrosum) is seen to be the more correct term.

Let us now attempt to classify these allied conditions, calling attention briefly to the more important forms.

Congenital Glandular Hypertrophy.—This may affect any glandular organ, though what is perhaps the most remarkable example occasionally is met with in connection with the mammary glands. These, with the development of puberty, may take on enormous growth, and be a source of so much disfiguration as to demand excision. If not removed, it is found that they function normally and excrete milk. If removed, as in a case of my colleague, Dr. Bell, which came under my examination, they are found to be composed merely of an excess of normal mammary gland tissue.



Multiple adenomatous polyps (adentiis polyposa) of stomach: D, duodenum; P, pylorie ring, (Orth.)

Irritative Hyperplasia.—This in general is marked by an enlargement of the gland, in the main due to increased fibrosis—as in chronic interstitial mammitis. With this, however, there may be some glandular overgrowth of the same order as that which we note affecting the epithelium in chronic ulceration when also the glandular elements, the sebaceous glands more particularly, may undergo actual hyperplasia. Indeed, the growth at times may be so extreme as to stimulate tumor growth. Similar irritative hyperplasia leads to marked overgrowth of the mucous membrane of the digestive tract at the edge of an old ulcer, and of this, again, the same is true. It is being increasingly recognized that *prostatic hypertrophy* comes into the same category, that the prostatic adenoma, so called, is not a blastoma proper.

Here, also, though forming a different class, is *regenerative hyperplasia*, such as occurs in the liver after acute yellow atrophy and cirrhosis (p. 785).

ADENOMA AND ALLIED CONDITIONS

Adenomatosis. - We would confine this term to the conditions often, but not necessarily, multiple, in which, while maintaining organic connection with the surrounding tissue, portions of a glandular tissue or surface become the seat of exuberant irregular adenomatous overgrowth, with evident disturbance of function. No sharp line of demarcation can be drawn between this and the preceding class. Under this heading come the multiple adenomatous polyps of the alimentary canal, some of the adenomas (fibro-adenomas) of the mammary gland, the multiple adenomas of the liver, advanced cases of prostatic hypertrophy, and uterine adenomas. From their general properties we are inclined to regard the multiple adenomas of the thyroid gland and ovarian adenomas and cyst-adenomas as belonging to this group, although regarding the last we shall have more to say when discussing the transitional lepidomata. It must be emphasized that the transitional lepidomas also afford adenomatous growths; we think it better to discuss them as a separate class.

Adenoma Proper.-Here, finally, we include all the sharply demarcated and completely encapsulated benign glandular growths. Their number, compared with the examples of adenomatosis, is relatively small, and the main members will be noted when discussing the transitional lepidomas. So far as we can see, these must, one and all, be regarded as originating from cell rests. The only case regarding which we have doubt is a sharply defined and extremely typical adenoma of the sudiparous glands of the wrist which came into our hands, in which, however, the absence of any cystic enlargement suggested that it must have possessed a communication with the exterior. When the cell rest is formed of glandular tissue which normally communicates with the exterior, the complete encapsulement, coupled with but slight anaplasia, must (we are inclined to think) inevitably result in cyst formation; with progressive growth of the epithelium lining these cysts, papillary projections occur into the cyst cavity (intracystic papilloma), and may almost completely fill them. Absence of secretion, as in some bile-duct adenomas, indicates either origin from non-secreting duct cells or a further grade of anaplasia.

Here we would include the well-encapsuled cystic adenomas of the mammary gland, whether situated within or separate from that organ, certain isolated adenomas of the liver and pancreas, and detached adenomas of the thyroid. The large and important group of adenomas of Wolffian-duct origin together with renal and adrenal adenomas come under the heading of transitional lepidomata.

THE ATYPICAL LEPIDIC GROWTHS.

CANCER.

We group together all the atypical growths from epithelium and gland tissues, as cancers. The properties of the two orders are the same; indeed, the most atypical members of the two groups can with difficulty be distinguished.

ring.

ma

net-

e so

oma

tion

adu-

icca-

tese.

and

not

r my

ss of

ronic idular e epis, the plasia. tumor vth of ulcer, gnized pros-

hyperid cir-

In all we meet with a greater grade of undifferentiation, or anaplasia of the constituent lepidic cells, than is noticeable in the adenomas and papillomas, though, at the same time, we cannot but recognize that forms showing *histologically* a relatively slight degree of anaplasia may, nevertheless, be as malignant and as liable to form metastases as are more anaplastic forms—or even more so, so that here we possess some notorious examples in apparent—may, actual—contradiction to the general rule that the extent of anaplasia is the index of malignancy.

The most pronounced example of the contradiction is seen in the condition of rodent ulcer proper, which Krompecher-we hold unfortunately-has rechristened basal-celled carcinoma.1 Ordinarily epithelioma is prone to form metastases in the lymph glands. Here we deal with an epithelioma of the most aberrant and anaplastic type, which, nevertheless, for long months, and, it may be, years, continues to grow and locally infiltrate and destroy the surrounding tissues, which, nevertheless, characteristically does not form metastases-which possesses local and not general malignancy. As we note later, this assignment of these growths is not accepted by all. Under this category comes also the large group of adenocarcinomas, which some would term malignant adenomas. Carefully analyzed, it seems to us that here the exception is more apparent than real. While these growths are of the adenomatous type, showing well-formed gland tubules, with lumina, etc., compared with benign adenomata of the same order, the arrangement is seen to be less typical. More particularly a basement membrane is found very largely absent; here and there, instead of a single layer of cells, the acini show several layers, and some are simple solid cell groups without lumina. In other words, of the adenomas of the same origin, one benign and the other malignant, the latter is the more anaplastic. Yet it has to be admitted that of these glandular tumors certain forms, showing relatively little evident anaplasia, have powerful infiltrative tendencies, with capacity to form metastases.

These examples, as before noted, we cannot explain. It has been suggested that gland cells which, under normal conditions, actively produce proteolytic and other enzymes may, when they take on blastomatous growth, then elaborate and discharge enzymes whereby they easily overcome the resistance of the neighboring tissues. This, however, does not appear to explain all cases.


Thus, then, we regard as cancer all cases in which there is infiltrative, and apparently independent, growth of epithelial or gland cells into the surrounding tissues, and this whether of only slightly atypical or markedly atypical cells.

Relations of Specific Tumor Cells and Stroma.—As with the adenomas, so here, the primary tumor element is the gland cell; it is this that makes its way into the tissues, and, doing this, sets up a reaction on the part of that tissue. Such reaction is often very well marked at the growing edge of these tumors, best, perhaps, in the epitheliomas.

¹ Krompecher, Der Basalzellenkrebs, Jena, Fischer, 1903,

ATYPICAL CARCINOMA

We observe that there is set up a reaction of a distinctly inflammatory type, with marked small-celled infiltration. More study is needed of the form of cells exhibited in this process, but some clearly are leukocytes, and such leukocytes may be seen to penetrate into the masses of cancer cells, to accumulate especially in areas where such cells have undergone necrosis, and, what is more, either actively to penetrate or to be taken up by the tumor cells, forming a definite order of *cell inclusions*. Probably both events occur, for while, on the one hand, we may find well staining masses of leukocytes occupying the site of previous cells, in others, more especially at the growing edge, the included leukocytes stain badly, and are evidently undergoing disorganization.



16

be

elv

iey

the

dly

the

t is

eac-

nas.

Early adenocarcinoma of the rectum, to show the marked difference in staining powers between the cancerous (a and b) and the unaltered epithelium (d). (Petersen.)

There are thus indications that the actively proliferating cancer cells feed upon the tissues of the organism; and it would seem that by phagocytosis, as by extracellular ferments and preparatory solution, the cancer cells replace the preexisting tissues, using them as foodstuffs.

Such process, however, has its limits. Often we can note that, the growing cells making their way into lymph spaces, those spaces still present their endothelial lining, and the vessels with surrounding connective tissue of the infiltrated area are retained to form the stroma of the advancing growth. Such stroma may, indeed, itself proliferate under the stimulus of the tumor cells, just as we noted in the case of adenomas, and there may be new-growth, not only of new simple connective-tissue elements, but of even higher forms. Thus, in secondary

LEPIDIC TUMORS

cancer of bone, such as frequently follows prostatic cancer, the stroma may exhibit not only remnants of the old bone lamellæ of the area invaded, but what is obviously new-formed bone. What is more, we occasionally encounter mixed growths that can only be explained upon the assumption that in what had been an original adenoma or cancer the stroma has secondarily assumed malignant properties. Growths of this nature have been encountered in the thyroid more often than in any other organ of man. That this acquisition of malignant properties by the stroma may occur has been demonstrated by Ehrlich and Apolant, and confirmed by other observers, in the course of serial transplantation of adeno-carcinomatous mouse tumors. Occasionally, that is, in one animal of the series the stroma (derived from that animal) becomes strikingly cellular and infiltrative, and in subsequent transplants, the sarcoma tissue may overgrow and replace the original tumor.

According to the extent of this reaction, so do we distinguish four forms of cancer: (1) medullary, when the cell growth is abundant and predominant, the stroma inconsiderable; (2) scirrhous, in which the development of the stroma and its overgrowth is the most marked feature, the cancer cells in such cases being small and compressed; and (3) carcinoma simplex, in which neither element can be spoken of as taking the upper hand. (4) Only very rarely, indeed, does the stroma overgrowth pass beyond the irritative stage and assume an independent blastomatous type, leading to the production of the true carcinoma sarcomatodes.

At the growing edge the cancer cells are characteristically hyperchromatic, their nuclei stain deeply, they are of the intensive vegetative type. Cells of the same order may be found also in the centre of a growth, with mitotic and other indications that the growth of these tumors may be expansive (from within) as well as peripheral. But frequently we note that, more particularly in the deeper portions of a growth, the cells show evidences of extensive degeneration, and the degeneration varies, often according to the mother tissue from which a cancer is derived. Thus, fatty changes are common in mammary gland tumors (recalling the active part taken by the normal cells of the mammary gland in passing on absorbed fats into the milk), mucoid degeneration common in tumors derived from the stomach and intestinal epithelium (in evident relationship to the normal function of the goblet cells of the mucous membrane).

Cancer Bodies.—There exists, in fact, a very remarkable series of localized degenerative changes in cancer cells that have been the cause of active controversy for now close upon twenty years; nor can it be said that the controversy is as yet at an end, although the main body of pathologists of all countries is now of the opinion that these appearances are degenerative, and not parasitic. For some years, however, the parasitic theory of cancer had active and enthusiastic supporters. We shall have to notice the various arguments in support of the parasitic theory when discussing the etiology of tumors.

Here we need only refer to these particular changes. They are of

two orders—intercellular and intracellular. The first we may dispose of rapidly.

These are the so-called "Russel's bodies," small, hyaline, spherical bodies of varying size, the mean size being that of a red corpusele. They stain an intense red with fuchsin, and thus are easily recognized. From the fact that they often lie in little groups attached one to the other, like vegetative yeast cells, Russel, of Edinburgh, who first called attention to their frequent presence in malignant growths, was led to regard them as blastomycetes. Their presence, however, is not confined to malignant conditions; they may be encountered in a great variety of inflammatory states, and are now accepted generally as examples of hyaline degeneration; from the fact that examination shows that some are intracellular, we are inclined to regard their origin as from

F10, 256

se ut a

be


itic

Russel's bodies, stained with fuchsin, highly magnified, from epithelioma of the hp. It will be seen that the majority are extracellular. (Klien.)

modified or degenerated chromidial bodies developing frequently in cells of the plasma-cell type (Fig. 257), which become extracellular by the disintegration of the cells.

The intracellular bodies assume a variety of forms. There may be a single rounded homogeneous mass within the cell pushing the nucleus to one side; or, like bodies may have a metachromatic central part; or they may be surrounded by a clear space in the cytoplasm; or show a peripheral ring or case, with different staining powers; or this peripheral ring may exhibit an obscure radiation, or present processes connecting it with the cytoplasm; or, again, a large, rounded, central body may be surrounded by a ring of smaller globules; or a ring or sphere of these globules may surround a poorly staining space; or, throughout the cytoplasm there may be scattered abundant small bodies of the first type lying in apparent vacuoles. Lastly, Sjöbring has described a large amœboid, gregarine-like form, at times within the cells, at times free and intracellular.

As will be seen from this rapid review of the main forms described, they are very various, nor have any two observers quite corroborated each other and the rest regarding what are truly parasitic, what are to be regarded as cell degenerations. As here set forth, it will be seen that the series of forms reads something like the description of the successive stages of maturation of an intracellular protozoon—of the malarial organism, for example—with progressive enlargement and continual production of peripheral spores, which may when set free, continue to grow within the parent host; or it may be that several small forms invade a cell at once. And, as such sporozoon forms, they were regarded by Ruffer and Metchnikoff, Sjöbring (first papers), Plimmer, and many other observers. Celli and the Italian school, from the result of experimental inoculations, were led to regard them as blasto mycetes—yeast-like bodies.



Cell inclusions in cancer cells—the supposed parasites. It will be seen that the bodies are to the inner side of the cell toward the lumen;

Intracellular bodies of the type of Russel's fuchsin bodies from a case of cancerous leukoplakin in cells of the plasma-cell type. (Krompecher.) The position, that is, of modified secretory products, (Greenough.)

Fabre Domergue was the first to call attention to the fact that these bodies have the same reactions as degenerative products within the cells, and Pianese, as the result of a most thorough and elaborate investigation, in which he first studied very thoroughly the microchemical staining reactions of mucin, hyalin, amyloid, keratin, and other matters, the products of cell degeneration, carried these observations much farther, and showed that all the forms described corresponded in their staining reactions with one or other of these forms of degeneration; as also that while by selection it was possible to recognize sharply defined and globular bodies in some of the cells of a given cancer, other cells in the same sections show matter giving like reactions, but diffused or so irregular in its disposition that there could be no question regarding its paraplasmic and non-parasitic nature.

These observations of Pianese have never been refuted; nay, more, Borrell and Farmer, Moore and Walker,⁴ Greenough,² Borrel,³ and

⁴ Proc. Roy. Soc. Biol., 76; 1905, ⁴ Jour. of Med. Research, N. S., 8; 1905; 137, ⁵ Bull. de l'Inst. Pasteur, 5; 1907; 497.

ATYPICAL PROPERTIES

several recent workers, have recognized the close similarity between these presumed cancer bodies and the products of nucleolar discharge into the cytoplasm, comparing them more particularly to the changes undergone by the "archoplasm" of developing sperm cells of mammalia. It is, of course, possible that there may exist one cycle of intracellular bodies which cannot be explained in this way, but so far that cycle has not been determined, and other considerations, to be noted later, show that no one specific agent can be ascribed as cause of malignant growths in general, or of glandular or epithelial cancers in particular.

Mode of Extension.—While sarcomas exhibit a predilection to extend by means of the circulatory system, cancers show a predilection for lymphatic extension. This does not mean that they also cannot form metastases along the vascular channels; a growth may penetrate the wall of a large vessel and its cells be "seeded" in the next capillary area; thus, we have seen pancreatic cancer invade the splenic vein and cause an extraordinarily diffuse secondary growth in the smaller portal veins in the liver. But in general the cells infiltrating the lymph spaces of a tissue find their way thus into the lymph channels, and then, either by continuous growth along those channels or by becoming detached, are found in the group of lymph glands draining the region affected. In this way, after proliferating here, some cells may be carried to the next group of glands in communication, so to the thoracie duct, and so eventually into the iugular vein and vascular system.

Site of Origin.—Often we can come to no conclusion as to the first stage of a malignant growth; it has become too extensive before death ensues. But even in these cases, in the majority of cases it is evident that the condition originated in a single limited area of an organ. If the growth has an external situation, we see that it develops at a single point. Most often (though not always) this is the case with cancer of the breast, and the appearances in other organs point generally in the same direction. This does not, however, mean necessarily that a cancer represents the progeny of a single cell. This matter has been studied by Hauser, and more fully by Petersen.

SP

ie

al

er,

ng

HC

Petersen employed the ingenious means originated, by the embryologist Born, of taking serial sections, plotting out each elaborately by means of the camera lucida upon a wax slab, cutting out the parts representing the stroma, and building the slabs one upon the other.

It has thus been demonstrated very clearly that, though in sections the alveoli of cancer cells appear separate, all are connected in series. The growth is like a bush of branching processes. But while in some cases there is a single root, or centre, very often it is pluricentric—a fact which indicates that at the same time several cells in the same region may take on aberrant growth.

Here, as with the adenomas, we have to conclude that these malignant growths may originate either from cell rests or from hitherto functioning cells. The best illustration of the former condition we have met with is, it is true, in a transitional lepidoma, in a case of Professor Aschoff's, in the museum at Marburg, in which the one adrenal shows a large

LEPIDIC TUMORS

infiltrating adenocarcinomatous growth, the size of a child's head, the other, a well encapsuled benign mass, the size of a cherry, a segregated mass of adrenal tissue, which has developed up to a certain point and then remained stationary. The clearest examples of carcinomatosis are afforded by intestinal papillomas. We have already called attention to the irritative origin of several of these growths. They frequently pass on to a malignant stage. Hauser' has shown, we think, convincingly that if a series of such papillomas be examined some manifest what must be regarded as the earliest carcinomatous modification.

The cells of certain of the follicles, which otherwise are quite typical, are, some of them, seen to have a lessened mucin production; others show no signs of such production, are smaller, with more deeply staining nuclei, and exhibiting a tendency to form not one, but two and more cell layers. With this the follicles become larger, irregular, form lateral projections, and bridges with neighboring follicles while others form solid processes projecting evidently through the basement membrane into the underlying tissue. We have ourselves noted a like series of changes in the same condition.

Occasionally several primary growths are encountered. Sometimes, as may happen in the breasts, one being the larger and noted before the other, it is a question where the smaller is not secondary, due to metastasis from the first. In other cases, as in primary carcinomas of the liver, this cannot reasonably be advanced. These belong to the same category as the multiple epitheliomatous growths of chimneysweepers, workers in paraffin, and following upon the use of arsenic (J. Hutchinson). The most striking example of the condition is found in adenomatous and adenocarcinomatous growths in the ovary. Here, practically always, both organs are involved. There may be no secondary growths found elsewhere at operation, and only the one organ may appear to be involved, but, if the other be left, it may require subsequent removal, and then presents the same type of disturbance.

So, also, more frequently than is usually recognized, the one individual may exhibit two or more distinct forms of primary growth in different parts of the body. Rarely are these both, or all, malignant; most often we have one malignant and one or more benign tumors, uterine fibromas, or thyroid adenomas, along with epithelium or cancer of one or other organ. But occasionally we encounter two different types of cancer in organs remote from one another. To these cases we have already referred (p. 693).

While one group of these cases (multiple malignant adenomas of the alimentary canal, hepatic "carcinosis," and the occupation epitheliomas) comes under the heading of carcinomatosis, the other group as evidently suggests a general tendency to aberrant growth on the part of the tissues, and is best explained as the outcome of vices of development and multiple development of cell rests.

We have here a condition of affairs curiously parallel with what we

⁴ Ziegler's Beitr., 33 ; 1903 ; J.

EPITHELIOMA

e

y

e

ic

e,

·S.

observe in infection. There, also, most frequently we recognize a single focus of origin, though in other cases there may be more than one, but these in general simultaneous, or almost simultaneous. And if in these cases we conclude¹ that the growth of the bacteria in the one focus sets up a general reaction of the organism, so that its resisting powers are raised, and the same species of microbe, gaining entrance elsewhere, is successfully resisted, so it may be the general rule in cancer, that once the cancerous process has developed at one focus, the diffusion of the products of the new-growth causes a general reaction, which, not sufficiently strong to arrest the process once actually started, is, nevertheless, sufficiently powerful to prevent the manifestation of like tendencies on the part of other tissues. This, indeed, has been shown to be the case by the experiments of Sticker, Gaylord and Clowes, and Ehrlich.2 Sticker showed that a mouse with inoculated cancer was immune to secondary inoculation, but the immunity disappeared soon after the primary tumor had been removed (see also p. 683).

The subject of cancer is so large and so important that much more might here be said. We will, however, proceed now to call attention to some of the more important forms. In describing them, we shall have occasion to call attention to matters of importance as bearing either upon etiology or properties of these growths in general.

SQUAMOUS-CELLED CANCER: EPITHELIOMA.

The epitheliomas, or epidermoid carcinomas, originate always from a squamous epithelium, and as such may be of epiblastic (skin, mouth, tongue), hypoblastic (e, g_{i}, α sophagus), and even of mesothelial origin (cervix uteri, vagina), no broad distinction can be made between epitheliomas and gland cancers along the lines of regarding the one as of epiblastic origin and the other of hypoblastic. According to its relationships and its functions, so does a given lining membrane develop either into the squamous-cell or the columnar or cylindrical type; and according to the type of the mother tissue so is the type of the malignant growth developed from it. At the same time it is necessary to keep in mind the existence of metaplasia, whereby through continued irritation a columnar may be converted into a squamous epithelium, which in its turn may take on blastomatous characters, so that epitheliomas may develop in regions where squamous epithelium is not normally present. This explains to a very large extent the aberrant epitheliomas of the larynx, bronchi, gallbladder, stomach, uterus, etc. While it is possible that some of these cases are not of metaplastic but of cell-rest origin, it is difficult to explain several on this assumption. For example, we cannot explain the appearance of squamous epithelium in the gall-bladder on the cell-rest theory (see Fig. 263). As an instance of this metaplastic epithelioma, Tyzzer³

¹ See Adami, Brit. Med. Jour., 1905, i : 1133.


7 See p. 848.

³ Fifth Report Cancer Com., Harvard Univ., 1909: 153.

LEPIDIC TUMORS

has recently called attention to the frequency of a form of chronic bronchitis in the mouse's lung, with proliferation of the bronchial epithelium into a many-celled layer, with loss of ciliated columnar cell character. This he regards as due to the irritation of large intrabronchial crystals, and as throwing light upon the frequency of epidermoid carcinoma in the lungs of those animals.

Characteristically, the squamous-celled epithelioma presents solid columns of cells passing in various directions and cut in sections, now longitudinally, now transversely, lying in a relatively abundant and moderately vascular stroma. As already noted, this stroma is apt to show considerable small-celled infiltration. In the most typical members of this group the epidermal characters of the growth are very marked. There is an outer layer of closely set cells with deeply staining nuclei, representing the Malpighian layer. Within this there may be several rows of prickle cells, which, as the centre of the mass is approached, become indefinite and flattened, and eventually keratinized. Thus, the centre of a process may be formed of concentrically arranged.



Epithelioma: earliest stage of cancerous metamorphosis and proliferation of cells. (Petersen.)

flattened, and keratinized cells, taking on the cosin stain strongly when hematoxylin and cosin is the stain employed. Such concentric bodies constitute the *epithelial pearls*.

The formation of these pearls is understood if we imagine, instead of a solid downgrowth of epithelium from the surface, a folliele-like downgrowth of skin. Such would exhibit keratinization of the oldest cells farthest away from the Malpighian layer. When the process is solid, these oldest central cells show still the same tendency toward keratinization.

In less typical growths the outer Malpighian layer is not so distinct, and, as von Hansemann points out, the mitoses, instead of being in the main confined to it, occur more irregularly through the whole mass; the prickle-celled elements, also, are not so distinct. In the true rodent ulcer occurring in the upper part of the check, below or at the angle of the eye, and sometimes a little farther out over the zygomatic area, the cell masses are still more atypical. Epithelial pearls are wanting; the constituent cells are rounded, polygonal, or even spindle-celled in

EPITHELIOMA

appearance (Fig. 267). Nor are the cell masses so sharply defined from the stroma, so that some observers (Braun) have classified these growths as alveolar sarcomas, or endotheliomas, and Borst still holds that the characters and region of growth show this to be the correct view. Careful study seems to show that the processes originate definitely in connection with the overlying skin (though Borst urges that the junction is secondary). The region of development of this particular form of tumor is curiously limited, and, as noted (see p. 790), while of a very anaplastic type, and exhibiting considerable, though slow, local malignancy, these tumors rarely form metastases.

111

11-

be psd.

of

ells

act,

igle

rea,

ng;



Early epithelioma of tongue, to show (a) region of origin by downgrowth from preexisting epithelium; b b, epithelial pearls; c, small-celled infiltration in surrounding tissue. (Petersen.)

Krompecher, recognizing that all conditions classified as rodent ulcer do not belong to the type, has labelled this form *basal-celled carcinoma*, on the mistaken ground that, as it shows no prickle cells or keratinization, it is derived wholly from the basal undifferentiated cells of the rete Malpighii. But this is so, also, for all epitheliomas. In the more highly developed forms the prickle cells present do not arise from preëxisting prickle cells, but also from the basal mother cells. It is the stage of undifferentiation, or anaplasia—the capacity or incapacity to develop beyond a certain point—that determines the form of the cells.

There are certain differences to be made out in the appearance of

LEPIDIC TUMORS

the tumors, according to the region of origin. Skin and tongue epitheliomas are apt to give the most well-marked pearls. Œsophageal growths in general show them poorly developed, just as normally in the



Portion of edge of a rodent ulcer.





Fig. 262

Part of the same at a more highly magnified, to show assumption by the epithelial cells of a spindle-shaped type, (Krompecher.)

cesophagus the keratin development is not marked. In rodent ulcers they are wholly absent,

The more rapid the growth, the deeper the infiltration, the more atypical is apt to be the growth; we have encountered metastasis of a



Aberrant squamous epithelioma of gall-bladder. (Von Hansemann.)

very malignant epithelioma of the tongue which it has been impossible to distinguish from those of a medullary round-celled cancer.

Regarding the metaplastic epidermoid carcinomas, it deserves note that these very often are not typical epitheliomata; that they show no

GLANDULAR CARCINOMA

801

epithelial pearls, and nothing corresponding to a Malpighian layer. This we hold is the case: they are not typical epitheliomata, but, also, they are not tumors of the type ordinarily arising in these organs—not adenocarcinomas, but solid cell-growths. Were we dealing with cell rests, they should afford tumors true to type.

We would scarce expect acquired characters to be so tenaciously preserved as those of primary endowment, but clearly, in these cases, the metaplasia has modified the nature of the resulting tumor. To the importance of this conclusion I shall revert (p. 843).

Lastly, an abnormal form of epithelioma is to be noted, in which, instead of epithelial pearls, degeneration occurs in the centre of the large epithelial cell masses, whereby false lumina are produced, giving the growth a transitional appearance. We have come across this form in man, and once in the horse, in connection with the antrum, the



Epithelioma of the antrum of Highmore, with degeneration and liquefaction of the centre of the cell masses, producing large pseudolumina. (Krompecher.)

pharynx, and the esophagus. Krompecher includes this as one of the types of his *carcinoma basecellulare*. Similar appearances are sometimes to be encountered in secondary growths from gland cancers, *e. g.*, from the stomach.

GLANDULAR CARCINOMA.

According to the structure of the mother tissue, and the stage of anaplasia, so do we encounter a series of forms of uncomplicated carcinoma. From mucous membranes which, normally, are provided with simple follicles lined with a cylindrical epithelium, as again from ducts, like the bile duct, and from tubular glands, we are apt to obta'n cancers which, in some parts at least, exhibit a tubular arrangement. From acinous glands we may gain cancers whose alveoli have a more solid 51

ssible

epi-

geal

the


nified.

more of a

> note w no

LEPIDIC TUMORS


type and grape-like arrangement; from the liver, alveoli formed of solid irregular strands. On the other hand, with greater anaplasia all glandular organs may give rise to growths formed of solid masses of cells. Once again the cancer has to be judged according to its region of origin; what in the one case is a relatively typical form of growth, in another may be most atypical. In general, however, we may distinguish certain main types.



Carcinoma simplex, (Ribbert.)

 The adenocarcinoma, including forms of highly glandular type, and differing but slightly, as we have noted, from the simple adenoma (the so-called malignant adenoma) (see Fig. 255), and others in which only here and there the alveoli possess lumina, with cells arranged around

FIG, 266



Medullary cancer. (Ribbert.)


them with some approach to the arrangement seen in the parent gland, the rest of the alveoli showing solid cell masses with, it may be, several imperfect small lumina scattered through, or only *pseudolumina* (produced by cell degeneration), or altogether devoid even of this distant imitation of the normal glandular arrangement.

GLANDULAR CARCINOMA

2. The cell-mass carcinoma, departing still farther from type, in which the alveoli are formed of an aggregation of cells without sign of orderly arrangement—cells either large, full, and rounded, or irregular in form, polygonal and compressed. More particularly in the latter type we may make the further classification (see p. 792) into the (1) medullary forms, formed of large cell masses, with little stroma, and that very vascular; (2) scirrhous, with abundant stroma, compressing the cells, and showing alveoli formed of few individual cells, and those compressed and small; and (3) carcinoma simplex, the intermediate form.

A yet further classification of the adenocarcinomas may be made according to the tissue of origin into:

(a) Cylindrical-celled carcinomas arising from mucous membranes, c. g., the less atypical cancers of the alimentary tract, corpus uteri, and, in part, of the cervix uteri (for here there may also be squamous epithe-



Scirrhus of breast. The cells are compressed and degenerated and the stroma relatively abundant. \times 250.

lioma, Fallopian tubes, gall-bladder, and portions of the respiratory passages (nose, trachea, bronchi). In all these there is a columnar-celled epithelium of one or more layers, in all a distinct tendency, not merely to infiltrate, but to grow outwardly into fungating polypoid masses; in all, again, so long as the adenomatous type of growth is preserved, some liability to formation of goblet cells and continued formation of mucin.

(b) Duct carcinoma, arising from ducts provided with a more or less cubical epithelium, in which, again, the less atypical forms present characteristically alveoli, recalling the characters of these ducts. Such we may get in the liver, from overgrowth of the smaller bile ducts. This is the most characteristic form of primary cancer of the liver, and may, indeed, represent not so much a growth of these ducts as an

d of a all es of egion th, in guish

> type, enoma which round

> > gland, several a (prodistant

LEPIDIC TUMORS

anaplasia or reversion of the liver cells primarily affected to the earlier stage of their development. In the mammary gland occasionally we meet with cancers in which the ducted appearance is predominant.

(c) Gland cancer, reproducing imperfectly the structure of the component acini of the glandular tissue of origin. Such, according to the tissue of origin, may be composed of more tubular alveoli (C. tubulare), or of grape-like acini (C. acinosum), as in adenocarcinoma of the pancreas and of the prostate, or of follicles tending to be separate (C. folliculare), as in cancer of the thyroid and of the ovary.

We may meet with a combination of these forms; thus, in the mammary gland we meet with all possible combinations of duct, tubular, and acinous cancer, with, in addition, another form, which has here to be noted, cvstadenocarcinoma.



Colloid cancer, showing the large alveoli, within which is contained the gelatinous colloid insterial. \times 300. (Rindfleisch.)

The organs and parts which give rise to these various forms of adenocarcinoma may, it must be remembered, give rise also to the more atypical and undifferentiated cell-mass cancers.

Degeneration.—What we have said regarding the degeneration affecting the adenomas applies here also. We may note particularly:

1. The tendency that superficial cancers have to undergo extensive ulceration. This applies especially to the cancers of mucous membranes (as, indeed, also, to the necessarily superficial epitheliomas). The new tissue is of a lower order not under the governance of the nervous system, cannot control its blood supply, and so is capable of offering little resistance to insults and infective agents.

2. The extensive mucoid and "colloid" degeneration that this same order of cancers may undergo, especially those of the digestive tract, leading to the development of what is termed *colloid cancer*. The cells of an adenocarcinoma, while still retaining the power of forming mucin, may, nevertheless, be unable to excrete it properly, with the result that it becomes heaped up in the cells to such an extent that they become

GLANDULAR CARCINOMA

greatly distended and eventually die. Whole alveoli may be found composed of the more or less inspissated and fused cell bodies—and the growth, filled with this modified mucin ("colloid"), presents a remarkable, massive, translucent appearance. So extreme may be the condition that only here and there, upon careful search, may alveoli be found showing relatively healthy cells, and affording a clue to the nature of the change.

Metaplastic Glandular Cancer.—Just as we noted that from a columnar-celled surface occasionally a squamous epithelium is found to arise, so, but much more rarely, do we get indications of the opposite process. The majority of the cases so far recorded must be regarded as the results of tissue disturbances and cell-rests. But Schridde has pointed out recently the normal presence of islands of columnar epithelium in the œsophagus, having a common origin with the surrounding squamous epithelium (p. 641), and this would seem well to explain the occasional occurrence of a columnar-celled cancer in this organ. And Enderlen has noted the frequent conversion of the many layered vesical epithelium into a definite columnar-celled type in cases of ectopia vesica, and in one case has seen this metaplastic tissue give rise to a definite adenocarcinomatous condition (see pp. 642 and 800).

CHAPTER XXIL

THE TRANSITIONAL LEPIDOMATA (MESOTHELIOMAS AND ENDOTHELIOMAS).

It will be recalled that we classed together as a main group of lepidic tissues all those lining-membrane tissues of mesothelial and mesenchymatous origin, derived secondarily, that is, from the mesoblast, and suggested that tumors developed from these be placed in a separate class, as the secondary or transitional lepidomas. Of these, upon consideration, it will be seen that we can make four groups: (1) the tumors arising from the developments and vestiges of the Wolffian and Müllerian ducts; (2) those arising from organs which, while they come into intimate relationship with these, nevertheless, as regards their essential constituents, are of separate mesothelial or mesoblastic origin (ovaries, testes, kidneys). With this group may be included the adrenals; (3) other mesothelial tumors derived from the serous surfaces, and (4) the endothelial tumors.

1. UROGENITAL DUCT TUMORS.

We make these separate classes because the urogenital duct tumors occupy a position by themselves. Whether, as we have hinted, these ducts gain a secondary lining of hypoblast or epiblast, or whether, from their singularly early differentiation, the properties of their mucous membrane are more fixed, certain it is that the tumors derived from them are most often of pure lepidic type—true adenomas and true carcinomas with little tendency, so far as we can see, to take on secondarily hylomatous (sarcomatous) development. Thus, in the uterus, we encounter in general pure adenomatoid or cancerous growths, indistinguishable in this respect from the adenomas and cancers of the digestive tract. The same is true in the prostate.

We have encountered a tumor of the prostate, certain portions of which, submitted to other well-known pathologists, were diagnosticated by them as possible alveolar sarcoma. Examination of all parts of the growth showed that in the body of the organ itself the appearance was that of a typical adenocarcinoma. As the growth infiltrated the bladder wall, and grew freely into the vesical cavity, the cell growth became most abundant, the stroma correspondingly diminished, until there developed a collection of large alveoli filled with moderately large, round cells, the alveoli being surrounded by thinnest of vascular stroma. The growth thus was still a cancer, though of the extreme medullary type.

TUMORS OF THE OVARY, TESTIS, ADRENAL, AND KIDNEY S07

The same holds for tumors of the Fallopian tubes, and, if we mistake not, for here our experience is very limited, for those of the ureters, vas deferens, and vesiculæ seminales. It is thus, also, for the interesting series of tumors which arise in the various developmental remnants of, more particularly, the Wolffian ducts in the female.

So far as we can see, few or no observations exist upon tumors originating from the corresponding remains of the Müllerian duct in the male (from the prostatic sinus or uterus masculinus, and the sessile hydatid of the epididymis). When these remains take an aberrant growth, their tubes being blind, they inevitably produce cysts-cystadenomas and cystocarcinomas. It is still a matter of debate among the embryologists to what extent the tubules of the primitive kidney (pronephros or Urniere) are contributed by the Wolffian tube itself; to what extent they are formed by the nephrotome, or mesoblastic blastoma; and by the pathologists how far ovarian and testicular growths are derived from components of the primitive kidney which take part in the development of those two organs. It is impossible, therefore, at the present time to make a positive embryogenetic differentiation of the tumors affecting these two organs. While tumors of the broad ligament are regarded as largely due to Wolffian duct remains, there is a conflict of opinion as to the part played by the Wolffian duct remnants in the development of ovarian cystadenomas.

2. TUMORS OF THE OVARY, TESTIS, ADRENAL, AND KIDNEY.


In the primitive kidney (Urniere), while the Wolffian duct provides the distal, collecting portion of the tubules, the glomerular epithelium and the main portion of the tubules are of mesoblastic origin; the same, following Balfour and Sedgwick,¹ is nowadays more and more accepted for the kidneys—that here the glomeruli and convoluted tubules are of mesoblastic (mesothelial) development. In the development of ovary and testis, not the Wolffian duct, but the primitive kidney, intimately connected as it is with the duct, is similarly involved, along with the germinal mesothelium.

This much may be said with regard to all these organs, that while in all four we may encounter pure adenomas showing no tendency to reversion, in all we encounter a remarkable series of transitional tumors, tumors in certain areas definitely of adenomatous type, in others, formed of solid cell masses which are not truly adenomatous, because, on employing Mallory's stain, we find that here and there connective-tissue fibrils are present beneath the cells. These portions are of the nature of alveolar sarcoma, and, on careful study, we can make out the transition from the truly adenomatous to the alveolar sarconatous areas. And from these latter areas we may pass to regions of purely sarcomatous type, round, or even blunt spindle-celled. The

¹ See p. 856 for a fuller study of the embryogeny of the kidney.

THE TRANSITIONAL LEPIDOMATA

picture is an extraordinary one, wholly at variance with the older views of the "sanctity" of sarcomatous and carcinomatous properties. Here, absolutely without any manner of doubt, a tumor shows transition from carcinomatous to sarcomatous characteristics. The condition has been regarded as inexplicable, has been labelled carcinoma sarcomatodes, or sarcoma carcinomatodes, has been treated as a ne er-do-weel member of the family, and too often left out of account in general discussions upon the family relationships of neoplasms. Some have thought to dismiss these cases by the ruling that mesoblast cannot form true gland tissue and true adenomas or carcinomas; that wherever, as in the kidney, we obtain typical gland tubules, these must be of epiblastic or



Hypernephroma of kidney, section from same tumor shown in Fig. 270. In this portion the growth retains the columnar arrangement characteristic of adrenal cortex, the columns of cells (sho) being separated by a capillary network, Bdg.

nbr. f

Transition from adenomatous to sareomatous type of growth: *ubnz*¹, adenomatous overgrowth of solid columns or masses of cells of adrenal type; *ubn*¹¹, transition to sareomatous arrangement; K, a kidney tubule involved in the growth, (Debernardi.)

hypoblastic origin; others have denied the transition. But the fact is there that such transition occurs, and is to be found in tumors of just these organs, as, again, in the endotheliomas, to be presently mentioned.¹

¹ Here it will be well to distinguish the four different orders of "mixed tumors." They are: (1) The teratoblastomatous, or "Mischgeschwüste" of Wilms, products of multipotential cells (p. 661). (2) The carcinoma surromatodes proper, in which the stroma of a cancer takes on independent malignant growth, such as has been noted by Ehrlich and Apolant (p. 792). (3) The mesotheliomatous, or transitional lepidomatous, as in the cases here described. (4) The invasive, in which, as has been noted in the thyroid and uterus, a surroma originating outside an area of adenomatous or cancerous change infiltrates into it, replacing the fibrous stroma.

ADRENAL TUMORS

Adrenal Tumors.—More particularly in the ovary and testis, as above noted, and to a less extent in the kidney, we at the same time may encounter tumors of fixed type, showing no signs of transition. These (possibly, as we suggest, originating from the more stable Wolffian epithelium) introduce an element of confusion. There is an organ free from constituents of this nature, and in it we find the most unequivocal examples of the transitional tumors in question. We refer to the adrenal bodies.

These are formed of two constituents, which in the lower vertebrates (e. g., selachian fishes) remain permanently separate, but in the higher vertebrates become joined. The medulla originates in connection with the sympathetic system; the cortex from mesothelial elements closely related to those which originate the cortex of the primitive and the permanent kidnevs.

From the medulla we obtain grayish tumors, never attaining any good size. These have been studied more especially by Marchand, who has demonstrated that they are anyelinic neurocytomas—true neuromas; that they contain rudimentary ganglion cells and nonmedullated fibres, and, in short, must be regarded as developed from cell-rests of sympathetic constituents (see p. 753). At times tumors are noted containing characteristic pigmented cells, the so-called chromaffin cells, which are present in the normal adrenal medulla, and also are in relationship to the sympathetic system, for tumors composed of those elements have been described in connection with the solar plexus. The cortical tumors are of wholly different order.

It is of frequent occurrence to find accessory suprarenals. These are nodules which are composed only of cortical tissue, either lying separate in the adrenal capsule, or outside it, or even as isolated, capsulated little masses within the adrenal tissue. Or, again, these have become adherent to and enclosed within the liver during development, or, more often, to and in the kidney. As Marchand has pointed out, the ovary, also, and the testis may carry down with them portions of adrenal tissue in their descent. In the rat, for example, such accessory adrenals are constantly found adjoining the ovary and testis.

Such accessory adrenals contain typical cortical tissue, columns of cells lying in a meshwork of capillaries, the cells containing abundant droplets of what appears to be fat, along with others of the nature of myelin. It is not infrequent to find that one or more of these, whether homotopic or heterotopic (under the kidney capsule, for example), have undergone hypertrophy, so as to be as large as a cherry. It is difficult, perhaps, to know whether to speak of this as hyperplasia or as an adenomatous condition; the fact that they are isolated from the normal tissue, and have grown, despite want of normal relationships, must, we think, place them among the benign adenomas. Others evade the difficulty by terming the condition *struma suprarenalis*, just as like nodules in connection with the thyroid are also labelled struma a convenient word, which means simply "nodular swelling." But, in addition to these, we may have much larger growths, often as large as a large as large as a large as a large as a solarge as large as a slowed by the superplaced by the superplasion of the set of the superplasion of the set of the superplasion of the set of the set

THE TRANSITIONAL LEPIDOMATA

child's head, developing in connection with the adrenal.¹ Studying them, we find the remarkable series of transitions noted. In some the growth throughout retains the character of the normal cortex, and is of adenomatous type. We have solid columns of cells, lying in a meshwork of capillaries, the cells much larger than normal, densely filled with fat and fat-like globules and glycogen (the last a constant constituent of the growing adrenal). Here and there may be giant nuclei, or cells with three or four nuclei, but the type of structure is well preserved. In other tumors of this type we encounter an occasional definite tubule, a column of cells possessing a lumen. Such tumors we must regard as adenomatous.

The existence of lumina is common in the cortex of the bird's adrenal, and is occasionally met with in the otherwise normal adrenal of mammals. It is a further support for the view now held that the adrenal cortex is of like origin to the renal cortex, derived from the same order of cells, and for the contention that these tumors are adenomas.

But other tumors exhibiting in the main these characters show in various areas a development of more irregular cell masses. The cells in those masses become smaller, less fatty, the nuclei more deeply stained, and in one and the same section we may have every transition, from the adenomatous, through the alveolar sarcomatous, to the diffusely sarcomatous appearance, with the appearance not merely of round, but also of irregular spindle cells. What is more, as well shown by the early cases studied in our laboratory by Dr. Woolley, a tumor of the adenomatous type in the adrenal may furnish metastases of purely sarcomatous type. The same has been noted by Jores and Askanazy, the latter noting also that after removal of the adenomatous tumor the recurrence was sarcomatous. More recently Meakins² has described another case.

It is but necessary to glance over modern literature to see what confusion exists regarding these tumors. Some speak of them as alveolar sarcomas (Beneke), others as angiosarcomas, or among the peritheliomas; others, again, as carcinomas (Ribbert); others, not to commit themselves, as hypernephromas (*hypernephros*, the suprarenal), hypernephroid tumors (Lubarsch), suprarenal epitheliomas (Marchand).

⁷The same doubt as to their exact nature extends to the tumors developing from heterotopic adrenal tissue, notably in the kidney (hyperuephroma or struma suprarenalis aberrata). Grawitz was the first to recognize the relationship between a group of large and eventually malignant tumors which affect the kidney, most often after the fortieth year, and the frequent adrenal rests in this organ. His views are now very generally accepted, though there is still debate as to whether all the tumors included by him and others under this term are truly of adrenal origin, and whether the cortex of the kidney itself, being so closely allied to its origin, may not give origin to tumors of like order.

⁴ As also, rarely, in the liver. See Pepers, Arch. de Méd. exp. et d'Anat. pathol., and White, C. P., and Mair, Jour. of Pathol., 12:1907:107.

² Proc. New York Path. Soc., N. S., 9:1909:19.

ADRENAL TUMORS

We are of opinion that this latter view must be accepted, that whereas, aberrant adrenal tissue is, from its heterotopic nature, more prone to become blastomatous, and whereas, it may well be that a large number of the kidney tumors of this type are hypernephromas, others are "nephromas." We certainly encounter typical tubular or cystic adenomas of renal origin, and know now that what we once regarded as the conclusive demonstration of adrenal nature, namely, the presence of glycogen (Lubarsch), is of little diagnostic value. A very large number of embryonic tissues, as, again, of freely growing tumors, contain glycogen in their cells. When a tumor of the kidney shows a special liability to form tubules rather than solid cell masses, we would suspect a renal rather than an adrenal origin. Not to enter exhaustively into the



Section of portion of a hypernephroma of the kidney. A characteristic area showing columns of clear polygonal cells: a, lying in immediate apposition to the endothelium (d) of the capillary sinuses (c). At b, areas of infiltration and degeneration. Section from another portion of the same tumor, more highly magnified, showing tubular arrangement: a, swollen translucent tumor cells surrounding a definite lumen b, capillary c, fat droplets in tumor cells. (Buday.)

subject, we would say that the two organs are embryogenetically so closely related that *tumors arising from homologous tissues must possess closely related characters*. Where, as is most frequent, but by no means constant, one of these growths arises from the upper pole of the kidney, there the probabilities are that it is of adrenal origin. It may, indeed, be, as Wilson holds,¹ that such rests as occur in the kidney are Wolffian body rests—again, it may be added, an homologous organ—and that because in development this intervenes between the kidney and the adrenal, and is the more likely to have certain of its cells included in the growing kidney.

¹ Trans, Assoc. Amer. Physic., 1910.

THE TRANSITIONAL LEPIDOMATA

Certain other properties of these more malignant tumors as a group remain to be noted. They are apt to be extremely vascular, the cells lie in close contact with the capillaries, with, as a rule, little connectivetissue stroma; the vessels are apt to be greatly dilated. Thus, hemorrhages are frequent, and necrotic areas, and infiltrations of the cell masses, which may gain thus pseudolumina, the cells in immediate connection with the vessels retaining their vitality, the central cells of a column or mass undergoing necrosis and becoming replaced by blood, so that an endotheliomatous appearance may be produced. So, also, not only are metastases mainly, as in sarcomas, by the blood stream, but both in the adrenal and in the kidney the tumors are curiously apt to grow in continuity along the veins into the inferior vena cava.

If a convenient term is required for all this order of tumors, the transitional adenocarcinomas of adrenal, kidney, ovary, and testis, we have, from embryogenetic considerations, suggested the term *mesothelioma* Briefly, a mesothelioma is:

1. A tumor arising from such tissues or portions of organs as, being of mesothelial origin, possess in the adult state lepidic characters.

2. When typical, and growing slowly, it is of pure adenomatous type.

3. When atypical or more anaplastic, and growing rapidly, it reverts first to an alveolar sarcomatous type, and later to a structure, or want of structure which renders it indistinguishable from a round or even a short spindle-celled sarcoma.

4. The tumor, when it takes on this undifferentiated type, affords metastases of sarcomatous order. The primary growth, in general, if studied, exhibits indications of the successive stages through which it has passed, from the adenomatous to the sarcomatous form of growth.

I would recall that in the chapter upon Classification (p. 705 et seq.) there is given what, to me, appears to be a reasonable explanation why this order of tumors has these particular properties.

3. MESOTHELIOMAS OF SEROUS SURFACES.

Occasionally on the pleural surface, more rarely in connection with the peritoneum, and still more rarely in connection with the pericardium, there is encountered a form of tumor which, from all the attendant circumstances, is of primary origin at these sites. They are flattened, nodular tumors, spreading locally over the serous surface, causing what at first sight appears to be extreme inflammatory thickening, of the nature of a localized hyaloserositis; but in general, on microscopic examination, they are found of a distinctly cancerous type. A relatively abundant fibrous stroma contains elongated acini, lined with irregular, swollen cells, large and sometimes almost cubical, resembling the curiously epithelioid type of cells we encounter in some endotheliomas. In parts these are flattened, lining long narrow spaces, and then they recall endothelioma proper. In one case that came under our observation it was possible to follow the cells covering the peritoneal surface directly.

first into more solid cellular downgrowths into the omental tissue, and thence, apparently, into the lymph spaces of the part. Sometimes, in place of a single layer of tumor cells, we encounter more solid masses, and in one case (in the pleura) we have noted a transition into the alveolar sarcomatous type.

Taking into consideration all these characters, it is difficult to regard these tumors as other than mesotheliomas, originating from the endothelium (or epithelium, whichever term be preferred) covering the serous surfaces, that cell layer being of mesothelial origin.

Here, again, there has been a great debate as to what these should be called, some regarding them as endotheliomatous, others (Ribbert) as true carcinomas. Histologically, they most often present a strikingly cancerous appearance. In certain early vertebrate ancestors, as pointed out by His, the body cavity originates as an invagination of the hypoblast; there is no sign of such invagination in the higher vertebrates, in whom the serous cavities seem clearly to originate by a splitting of the mesoblast and formation of a lining mesothelium.⁴ The fact that in certain of our ancestors the body cavity was lined by a true epithelium does not convert the lining of later generations into an epithelium when in them the cavity originates in another manner.

4. ENDOTHELIOMA.

Of those tumors in which overgrowth of the lining cells of vessels is the most prominent feature, we distinguish naturally two groups: the hemangio-endotheliomas, originating from the lining membrane of bloodvessels; the lymphangio-endotheliomas, from that of the lymph vessels. The two groups present many features in common, nor is it always easy in tumors of large size and long establishment to determine with which form we have to deal; but even in such cases, in one or other part of a tumor wells, which, if they contain blood corpuseles, are clearly of blood capillary nature, if free throughout from formed contents, must, per exclusionem, be regarded as lymph spaces.

Ribbert objects that no proof is afforded of the nature of these tumors by finding vessel spaces lined by cells approximating in type to those of the tumor. We cannot but feel that the objection is hypercritical. While, with him, we agree that it is contrary to experience to find that the normal cells in the neighborhood of a tumor take on progressively, blastomatous features, the other possibility has to be admitted, that proliferating endothelial cells replace the normal cells in their outward extension. How, indeed, otherwise does he, or are we to, explain the continued enlargement of these tumors? For grow they do, and such is their structure that the growth cannot be central. We see such a

⁴ See Miller and Wynn, Jour. of Pathol., 12:1908:267, for bibliography of these conditions,

THE TRANSITIONAL LEPIDOMATA

process occurring in the uterus in pregnancy, when the fortal syncytial cells absorb and replace the endothelium of the maternal blood sinuses, and it is in full harmony with all we know concerning cell properties that such tumor cells, when in relationships which more nearly approach the normal, should themselves exhibit characters more nearly approaching the normal.

Typical Endotheliomas.—We find some little difficulty in treating these growths, and that because, under this heading, it is customary to consider only growths of an atypical nature. Following our custom, we think it necessary to call attention first to typical growths. These, along with a collection of what are not blastomas at all, have usually been discussed as a class apart, and this by modern writers, who, nevertheless, admit that they are of the nature of benign endotheliomas. We refer to the *angiomas*. These, then, we will first take into consideration.

5. ANGIOMA.

We cannot but conclude that the majority of so-called angiomas, or tumors having vessels as their main constituent, are spurious blastomas, whether formed of bloodvessels ("hemangiomas") or of lymph vessels ("lymphangiomas"), and this because these exhibit no power of independent growth. Mere dilatation and filling of vessel spaces with fluid is not growth, even if preceded by aplasia and followed by atrophy of the tissue proper to the part. And in the majority of cases the evident increase in length of the vessels (such as must occur in *cirsoid aneu*rusms) or thickening of the walls of the individual dilated loops (such as we see in *cavernomas*) is apparently not in excess of the physiological requirements. We find, that is, no evidence of proliferative capacity, at most a widening of preëxisting vessels, either of congenital origin, and ascribable to a primary want of coördination in the growth of the vessels of a part and of the tissue or cells they should nourish, or of postnatal origin, due to alteration in blood pressure of the nature of local venous obstruction, as, for instance, in the multiple capillary telangiectases, which can be produced in the liver by partial obstruction and stenosis of the hepatic vein-hemorrhoids are of this nature-or due to local atrophy of the cells of a restricted area in an organ, the capillaries undergoing what we may speak of as compensatory dilatation. We do not mean to imply that true angiomatous blastomas are non-existent; as we shall point out, they exist, only, compared with the spurious form, they are less frequent.

Here once more we must dwell on the meaning of words. A cellrest or inclusion within a tissue is not a blastoma so long as it lies latent and is not growing; similarly, a mass of tissue which, owing to developmental defects, is aberrant in structure is not a blastoma so long as it strictly respects physiological laws and at most grows coincidently with the rest of the organism. Independent growth is the test of what constitutes a tumor of this order. Does this mean that we are to remove

SPURIOUS HEMANGIOMAS

most of the conditions now included under the heading of angiomas, and place them under the class of, say, telangiectases? Frankly, this would be the better course; it would make for precision. We must be governed here by our conception of the meaning of the word *angioma*. If that is to be taken as meaning simply a swelling composed of bloodvessels, then all these remain as angiomas. If we restrict it to mean a tumor, due to the independent growth of vessels, they must be cast out of this class. To provide a class for them Albrecht has suggested the term *Hamartoma*.⁴

BLOOD-VASCULAR TUMORS ("HEMANGIOMAS") WHICH ARE NOT BLASTOMAS.

1. Obstructive Telangiectases.—The type example of such is the *hemorrhoid*, or pile.

The hemorrhoidal veins of the anal region communicate, it will be recalled, with both the main and the portal venous systems. Situated, as are these veins, immediately beneath the surface, and so, poorly supported, obstruction to the onward passage of the blood in either system is liable to lead to a dilatation of the capillary loops and smaller veins.

Similar capillary telangiectases are not infrequent in the liver; notably they occur in connection with the nutmeg liver (chronic passive congestion), as, again, upon the nose and cheeks of elderly individuals, apparently secondary to localized fibrotic changes and venous obstruction; while venous telangiectases (*varices*) are also common, and due, if not to obstruction, at least to giving way of the walls against the weight of the column of blood (pampiniform plexus, superficial veins of the lower extremities—*varicose* veins, etc.). In like manner, weakening of the arterial wall, so that it is unable to withstand the internal blood pressure leads to the production of:

2. Aneurysms.—Of these, one variety has been the cause of debate whether it should be regarded as angioma, namely, the *cirsoid* aneurysm. This is evidently congenital, may show itself at birth, but sometimes only "grows" rapidly in adult life. The favorite seat is the scalp, where the worm-like pulsating arteries beneath the skin give a very character-istic sensation. The dilatation of the vessels may lead to erosion of the skull. The condition, we hold, can only be ascribed to a congenital weakening of the arterial wall, relative to the blood supply and blood discharge.

We had occasion to examine one such some weeks after it had been reduced by ligaturing the carotid on the same side. The case was that of a young adult, in whom, from being inconsiderable, the aneurysm had rapidly increased. We found practically nothing. There was no evident increase, in our sections, in the number of arteries of the temporal

¹ Apparently from *aµapτia*—error, *i. e.*, due to developmental defects.

region, which had been involved; at most a condition of fibrosis, and that not of recent type.


It has been argued that in these conditions there is a condition of angiomatosis, because on ligature or removal of such cirsoid aneurysms they are liable to recur. In face of this finding we can only conclude as above, namely, that there is a constitutional weakness of the arterial



Cavernoma of liver. Gross appearance. (After Ribbert.)

walls, coupled with inadequate discharge, and so with relative and local increased pressure an artery and its branches become widely distended. A like cirsoidal aneurysmal condition occasionally follows trauma.

3. Congenital Telangiectases.—(a) Telangiectactic Nevi: Some nevi (pigmented moles) are purely cutaneous overgrowths, with collection of subcutaneous melanin-containing cells; others are yellow, and of the nature of xanthoma; the majority contain, in addition, dilated capil-



Section of small cavernoma of liver, showing the cavernous and communicating vascular spaces, from which the blood has been removed, (Ribbert.) laries, or may, indeed, be areas of simple telangiectases, such as the ordinary "birthmarks." An extreme grade of the same condition is the blue nevus, which may be extraordinarily extensive, affecting the whole side of the face, or even larger areas.

At Manchester, as a student, I assisted at the autopsy of an adult, a patient, if I remember aright, of the late Dr. Dreschfeld, in which the whole head and neck were involved. The skin was purple, the face a collection of coarse nodules. At Markurg, an infant, upon which Professor Aschoff performed the autopsy, showed the whole lower half of the body involved.

The frequent association of the telangiectasis with congenital pigmental disturbances in the same area indicates strongly that here we are dealing with a vice of development. We find two grades: in the simple birthmark (N. flammans) it is obvious that we are dealing with a capillary dilatation; such may also be found in bone, muscle, and (rarely) brain. In the *blue nevus* the vascular spaces are apt to be of greater size, sometimes septate, showing where, by pressure atrophy.

TYPICAL ENDOTHELIOMA: HEMANGIOMAS

neighboring spaces have fused into one, and there is a more cavernous appearance. Some authorities regard these as of venous origin, and speak of venous telangiectases. We are inclined to regard them as of capillary origin, the thickening of the walls and the surrounding fibrosis, which give the walls the venous appearance, being regarded as of the nature of overgrowth through pressure—strain hypertrophy.

(b) Cavernoma.—From these cases we pass imperceptibly to the cavernoma. These cavernomas are one of the commonest abnormalities to be met with in the liver, where they most often are small—from the size of a pea to that of a hazehunt—but, rarely, may be of great size—as large as an orange, and larger. They are to be found in the livers of young as well as of elderly individuals, and evidently a considerable proportion are of congenital origin, though probably some are acquired through localized atrophy of a group of liver cells and compensatory dilatation of the capillaries of the part.

Examined microscopically, they are found to be formed of large, irregular, distended, and communicating blood spaces, lined with endothelium; the walls between these are relatively thick and fibroid. They may contain groups of pigment particles, suggesting, when present, possible previous liver cells which have undergone atrophy, but of persistent liver cells there are none in the affected area. There is a liability to thrombosis in these cavities, indicated by the presence of recent blood clot, or organized blood clot, and fibrous bands; or, again, of calcification and formation of phleboliths.

Here, also, we deal apparently with what are capillary ectases. As we have noted, some are obviously of congenital origin, and it is suggested that they are due to a vascular branch not becoming clothed with, or not entering into connection with, liver cells; while Ribbert and others have called attention to the fact that the spaces do not communicate with the surrounding capillaries, and cannot be injected through the hepatic vein. We have noted these 20 times in 1400 complete autopsies, more frequently, in proportion, in adults and elderly people than in the young, and so are inclined to regard some, at least, as due to localized atrophy of liver cells. The majority show no sign of independent growth.

TRUE TYPICAL BLASTOMATOUS HEMANGIOMAS.

Angioma Simplex.—There are, however, true hemangiomas in which we encounter what can only be regarded as a progressive new development of capillaries. The slightest grade is seen in what are otherwise simple capillary angiomas, characterized by no pronounced ectasis. Some there is. But what is the marked feature is that the endothelium is very prominent; not only are the cells relatively large, with much cytoplasm, but they are two, and it may be more, layers thick. The appearance is not that which would be given by contraction of dilated capillaries in the process of preparation; there has been a definite over-

growth. Such cases have been recorded from the skin, the chorion, and muscle. A definite tumor is thus formed, composed throughout of these proliferating endothelial tubules. This we may speak of as the true benign hemiangioma. A more advanced condition is seen, in what Ziegler termed, we think unfortunately, angioma hypertrophicum, for there is more than hypertrophy-there is a true blastoma formation. In this, while some of the capillaries have the appearance noted above, in others the endothelial overgrowth has become so extensive that solid columns of cells are formed, and, what is more, these appear to be budding or projecting into the surrounding tissue. Save that the capillary tubes are the prominent feature, the condition is scarcely removed from what is characteristic of the hemangio-endotheliomas. A full study of a case of this nature, while of benign type, afforded metastases, has recently been published by Borrmann.¹ In fact, it will be seen from this description that we regard the angiomas proper as typical endotheliomas; that we regard the capillaries alone as giving rise to the condition, and see, in the capillary endothelium, the one primary factor in new vessel growth. All other conditions come under the heading of telangiectases.

Next may be included, possibly, a remarkable and rare form, whose nature, we believe, was first recognized by Ziegler. This has its seat in the skullcap, and presents itself as a sharply defined nodular growth, composed of large blood-filled spaces, with little interstitial tissue, each lined with a relatively large and very regular cubical epithelium. This form is most nearly related to the hemangio-endothelioma of bone and kidney, but differs in the remarkable regularity of its gland-like endothelium, and not infiltrating, sharply defined character.

We encountered this form once many years ago, before we knew much regarding endothelial possibilities, and shall not easily forget how it mystified us. The epithelium had an obviously glandular appearance, and the cavities were as obviously filled with blood.

Ribbert denies that vascular endothelium ever takes on such epithelial type as is seen in tumors of this nature, and so excludes them from the class of endotheliomata. In so doing he forgets the swollen epithelioid character that the endothelium may take on (along with increased proliferation) in various forms of inflammation, notably in endarteritis affecting arterioles. There the cells become strikingly epithelioid.


LYMPHANGIOMA.

 Here we have an exactly parallel series of cases to those observed in connection with the bloodvessels. The majority of cases termed lymphangiomas are strictly lymphangiectases. These may be wholly apart from any other form of growth, or, like capillary ectases, may accompany various forms of tumor.

¹ Ziegler's Beiträge, 40:1907.


HEMANGIO-ENDOTHELIOMA





Section from a hemangio-endotheliona of bone: a, large vascular spaces filled with erythroeytes and surrounded by large, elear, eubical endothelial cells, which in parts, as at e, form solid masses; b, stroma; d, larger and e, smaller bloodvessels. (Driessen,)

F1G, 276



Section from a case of hemangioma simplex, exhibiting progressive enlargement and extension. (Borrmann.)

In the preceding section no reference was made to what we have called attention to elsewhere, namely, the frequent occurrence of dilated capillary spaces in every form of overgrowth, which, more particularly in the connective-tissue group, may afford indication of actual new formation of vessels. When such occurs, it is strictly subordinate to the growth of the tumor, and in no sense independent, save in the hemangio-endotheliomas, so that to speak of an angiofibroma, etc., is incorrect. "Telangiectatic fibroma" expresses more accurately the condition. The same is true in respect to the frequent occurrence of dilated lymphatics in tumors; the condition here is subordinate, and, indeed, may occur in company with the former. When independent of other new-growth, these lymphangiectases may be either inherited or acquired. We thus distinguish three grades, although between them there is every transition.

1. Simple Lymphangiectasis ("Lymphangioma Simplex").—(a) Congenital.—One or several lymphangiectatic areas may be present in the skin, slightly protuberant, sometimes breaking through, and then "weeping" persistently (lymphorrhœa). Like "mothers' marks," they are most frequent on the face and neck, the frequency in both cases suggesting some slight vice of development in connection with the closure of the fissures present in these areas during development. The affection may be confined to the papillary layer of the corium, or may extend more deeply. On section, the tissue presents abundant moderately dilated and cylindrical lymph channels, lined with endothelium, lying in a fibrous, somewhat cellular stroma.


(b) Acquired.—Of allied type is the condition seen in filarial (tropical) elephantiasis. Here we have definitely to deal with lymphatic obstruction as a cause of the development of the condition, and it is worthy of note that, as we have pointed out elsewhere, the obstruction leads to surrounding connective-tissue overgrowth. We do not think it necessary, with Ribbert, to assume that there is an essential connection between the lymph channels and the surrounding connective tissue to account for the fibrous overgrowth seen in so many cases of lymphangiectusis. It is a secondary result of the expansion and "lymphedema."

2. Cavernous Lymphangiectasis ("Lymphangioma Cavernosum'). —This corresponds to the cavernomas, only, in place of blood, the wide, irregular chambers contain lymph. Under this heading we have some remarkable congenital conditions: macroglossia, in which children are born with relatively large tongue, which may continue to enlarge after birth; macrocheilia, similar enlargement of the lip, congenital elephantiasis (E. lymphangiectatica). When occurring in the mesentery, these cavities have milky, chylous contents. Here, also, we find extensive overgrowth, with fibrosis of the parts between the dilated lymph spaces. The conditions are all congenital, and we must conclude that there is obstruction to onward flow of the contained fluid, due to some abnormal relationship of the different vessels.

3. Cystic Lymphangiectasis ("Lymphangioma Cysticum").—The most extreme and remarkable examples of this condition are encountered

SPURIOUS LYMPHANGIOMAS

in the neck, causing the condition known as *cystic hygroma*—multiple large clear cysts, either below the ear or, more commonly, in the submaxillary region; or, again, below the level of the larynx, and extending to the supraclavicular region; usually unilateral, and forming a large, tense, fluid swelling, which may extend outward to the shoulder, or deeply beneath the sternum. The tumors are formed of a collection of large cysts, lined with endothelium, containing clear lymph, and having fibroid walls. Many of the large cysts appear to be absolutely closed off, not communicating with their neighbors. We must suppose that, with increasing distension, there has been a valve-like closing of the channel of which they are a dilatation, that the endothelium has grown *pari passu* with the dilatation, and that this endothelium has secretory powers. The mere force of the lymph flow cannot explain such extreme



Lymphangio-epithelioma of the lung. (Adler.)

development; we have to assume active excretion, which, indeed, is indicated by many other considerations and actual experiments (Heidenhain) (see also p. 861).

This condition must be distinguished from *cervical hydrocele* brought about by secretion into one of the persistent cervical ducts or fissures. Such consist of one cyst, and are lined, not by endothelium, but by squamous or columnar and sometimes ciliated epithelium.

An allied form is the *sacral hygroma*, one variety of the congenital sacral tumor (p. 239), in which, again, from imperfect lymph discharge, there develop congeries of relatively large, lymph-containing cysts. In uterine myomata occasionally we meet with a like condition.

4. Lymphangioma Proper.—From these telangiectatic cases we pass to what must be regarded as true lymphangiomas. Here let us repeat that the mere existence of growth of lymph channels, *pari passu* with other changes in tumors, but subordinate to the main tissue overgrowth.

THE TRANSITIONAL LEPIDOMATA

must not be considered as lymphangiomatous, even if actual budding of new channels be observed, as noted by three or four observers.

Borst describes and figures a localized lymph-vascular nodule observed by him in a lipoma, composed of a close collection of apparently new lymph channels. We have noted a somewhat similar condition innormal, but very fatty, appendices epiploice, and have regarded it as a latent lymphoid nodule. Whereas, in the normal state, such can be detected only with difficulty, in acute peritonitis each appendix is found to contain an easily recognizable and typical lymph follicle.

The cases that we would regard as true lymphangioma are characterized by a notable proliferation of the lymphatic endothelium. In some cases of congenital simple lymphangicetasis in the young, this is to be observed here and there; in older individuals the same is not observable. In some cases, for example, of cystic hygroma of the neck it is difficult not to believe that there is an actual growth and extension of the tumor. Schwalbe has described a case in which, in addition to the marked proliferation, he found solid endothelial cords being produced, and even solid processes, as of advancing new formation. Here we seem to have the transition to a lymphangio-endothelioma.

Possibly the lymphangioma tuberosum multiplex of Kaposi comes under this heading; multiple small nodular cutaneous appearances in the adult, which, on section, are found to be due to dilatation of a group of lymph spaces in the cutis, each of which shows marked endothelial proliferation and is filled with a jelly-like matter, due, according to Beneke, to a hyaline degeneration of the proliferated endothelium, and, according to the same observer, there is here a pronounced new budding of lymphatics.

ATYPICAL ENDOTHELIOMAS.

Hemangio-endothelioma.—Of this form of growth quite the most frequent and characteristic example is that developing on the inner side of the dura, over the skullcap, where it forms smaller or larger sessile or rounded nodules; or from the membranes at the base of the brain, where it may either form a large nodule or take on a spreading growth. More rarely, it may originate within the sheath of the optic and other nerves.

The appearance of a favorable section of such a growth is very striking. The whole field may be found composed of a collection of whorls of concentrically disposed cells. Between these whorls, and not sharply marked off from them, is a somewhat cellulur stroma. The cells composing these whorls are flattened, the centre of each whorl is not close packed, but rather loose, and with careful examination, or good fortune, here and there may be seen a whorl that has a lumen containing red blood corpuscles. Elsewhere, where the section has not been transverse through the component whorls, they have an oval appearance, or may be elongated and curved. In some cases there is a fair amount of cellular

HEMANGIO-ENDOTHELIOMA

fibrous tissue separating the individual whorls; in others the whorls exhibit hyaline changes, and their central cells become homogeneous, fused and translucent; while frequently we encounter the further change of deposit of calcareous salts in these degenerated areas, so that the specimen has scattered through it small calcareous nodules. Tumors in which this calcareous change is marked have been given the name of "psammoma." These psaumomas, and they are not uncommon, are, so far as we can determine, always of endothelial origin. There may be multiple small tumors of this nature scattered under the dura, or in the choroid plexus. In other cases, and these more particularly where there is a more rapidly spreading growth, the picture is not so clear; only in some parts of the section can these whorls be made out, and then somewhat indefinitely; elsewhere, and not sharply defined from the previously

mentioned areas, we have the appearance of a moderately small-celled sarcoma—a diffuse oval-celled growth.

The appearance is most satisfactorily explained as due to localized overgrowth of the capillary endothelium, at first in the main concentrically, so that the capillary becomes enlarged by the deposit of layer within layer of these endothelial cells. But, as we have had repeatedly occasion to note, the capillary is formed merely of endothelium, and growth may thus be outward as well as inward. Ribbert figures an early endothelium of the optic-nerve sheath, showing this outward growth. We have encountered the like appearance in the same region. It is in this way that, with more active proliferation, a more diffuse sarcomatous growth is developed.



Portion of an endothelioma of the dura mater, showing the characteristic whorled arrangement of the tumor cells and at a a concentrically arranged calcareous deposit or psammoma body. (P. Ernst.)

Ribbert, with whose views on this group of tumors we find ourselves largely in disagreement, regards these tumors as developed from the endothelium covering the dura mater and pia arachnoid space. He does not explain why a growth from such a primarily covering endothelium should take the remarkable form of cells concentrically disposed as though along a series of vessels, as, indeed, serial sections show is their arrangement. We are prepared to find that tumors exist of the order he describes, corresponding with the pleural and peritoneal mesotheliomas, but this is not the common type.

As a rule, it would seem that this form is relatively slow growing, in the main causing death by pressure upon the brain substance. We have, however, found it extensively spreading over the surface of the brain, and in a case where the optic nerve was also affected, have found recurrent nodules of alveolar sarcomatous nature developing in the orbit after extirpation, whereas the cranial growths showed still the endotheliomatous type.

• THE TRANSITIONAL LEPIDOMATA

Atypical Lymphangio-endothelioma.—More often, in other regions of the body, similar whorled cell masses of endothelioid type show no relationship to blood capillaries, and must be regarded as originating from the endothelium of lymph channels. These, equally with those just described, when exhibiting more active growth, show areas of round-celled sarcomatous type; indeed, the greater part of the tumor may appear sarcomatous.

Perithelioma.—A striking form of tumor is occasionally encountered exhibiting capillary channels cut in various directions and lined by recognizable endothelium, around each of which capillaries is a collection of cells, many layers deep, arranged radially. The individual cells are not specially elongated, but the arrangement in rows at right angles to the capillary axis is most characteristic. The general opinion



Section of a perithelioma of Luschka's or the coccygeal gland, (Von Hleb-Koszanka,)

is that these cells gain their origin from the lymphoid endothelium of the perivascular space—that thus these tumors form one variety of lymphangio-endothelioma.

Apparently those cells farthest removed from the central bloodvessel are the oldest. Whether from this cause, or from their more remote position, they are liable to exhibit degenerative change, and more particularly to undergo hyaline change. To such modification of a perithelioma it would seem that we owe the most typical form of *cylindroma* —tumors formed of a collection of hyaline tubes, or cylinders, cut in various directions, having a central dilated capillary vessel, surrounded by a zone of oval cells, somewhat radially disposed. Saying this, however, it is necessary to interject a word of caution. The cylindroma is by no means always a perithelioma; in fact, it is rarely such. Very many orders of tumors, even including carcinomas, when poorly nourished,

MELANOMA

show a zone of persistent tumor cells around the vessels, with hyaline necrobiosis of cells further removed, and there is the tendency to confuse these with peritheliomas proper. A frequent seat for this order of nonperitheliomatous eviladroma is in partotid tumors.

MELANOMA.

We have purposely left to the last the consideration of a series of tumors regarding which there is still hot debate and violent conflict of opinion. While, as will be seen, we take a definite position on one side rather than the other, believing that the facts brought forward up to the present turn the scale in that direction, and that it is wiser to present positive opinions, we confess that our inclination is to treat the matter as still open.



Benign papillary nevus, or papillary mole, to show the accumulation of "nevus cells" (Ribbert's chromatophores). (From photograph by Dr. J. A. Fordyce,)

Pigmented moles are, as everyone knows, a very common minor malformation. Most individuals, if we mistake not, are possessed of one or more. In the adult these exhibit no very clear histological picture; in the child their structure is more definite. They consist of a fibrous stroma immediately beneath the epidermis, in which are situated clusters of cells of fair size, irregularly polygonal, and containing brown pigment (Fig. 280). In the slighter cases these cells are noted as encircling closely the vessels. The condition is notoriously congenital. The mole represents an area in which there has been some vice of development. The specific cells are of a peculiar order, peculiar, not to these moles, but to the skin, and (to a less extent) the mucous membranes, as, also,

THE TRANSITIONAL LEPIDOMATA

the choroid coat of the eye, where they are most abundant. These pigment-bearing cells are known as *chromatophores*.

In the ordinary skin of the white man they cannot be clearly made out, save in the anal region and the pigmented areola of the nipple. In animals with deeply pigmented skin they are most abundant, and are to be seen not merely in the corium, but between the cells of the deeper layers of the epidermis. They are characterized by possessing two or more long, rather coarse processes tending to be branched, and relatively abundant cytoplasm, in which are pigmented granules of melanin, a pigment differing from hemoglobin in being iron-free and relatively much richer in sulphur (see p. 970). That the cells of the pigmented mole possess these properties can be demonstrated, according to Ribbert, by examining a teased-out preparation. In sections they appear merely polygonal.

Whether these moles should be termed definite benign tumors melanomas, or, with some, melanofibromas—is at least debatable. Though to the naked eye they appear sharply differentiated, under the microscope their connective-tissue stroma passes imperceptibly into the surroundings. Nor, although clearly due to some vice in development, can we with absolute precision speak of them as cell-rests. The appearances indicate more the excessive development of what is a constituent of the normal skin rather than a dislocation—a constituent which, for some reason (possibly increased vascularity, for these moles are most often nevoid), has taken on the active heaping up of pigment. Here we should explain that everything indicates that the chromatophore is a cell which has the capacity to manufacture melanin, but which, however, is not always melanin-containing. But the cell relationships are here disturbed.

In the choroid coat of the eye, and from the skin, frequently originating from such moles, we gain the development of highly malignant melanotic tumors. It is interesting to note that in the eye similar aberrant cell clusters have been noted, either in the iris (in areas showing coincident tumor growth) from the choroid, or even in and upon the sclerotic, as though in this latter case, in the course of development, a portion of the ultimate choroidal tissue had been pinched off. Virchow has described a primary growth from the brain membranes, which often show some pigment cells; Störk, from the pia and spinal cord. And primary growths have been recorded from other regions, more particularly the liver and gall-bladder. We confess to having had considerable doubt concerning the primary nature of these growths until recently our colleague, Dr. Duval, showed us his material from what was clearly a small primary growth originating beneath the mucosa of the common bile duct.¹ So, also, what appear to be authentic cases of primary growth have been reported from the ovary (Rösole), thyroid (Fränkel), adrenal (Orth), lymph nodes (Marchand, Birch-Hirschfeld, Martini, and others) and prostatic urethra.

¹ Montreal Med. Jour., 37 : 1908 : 270.


MELANOMA.

These tumors are very striking. According to the amount of pigment they contain, they may be coal black, or various shades of brown, or, on section, show pigmented areas, while the rest of the growth is colorless; or the primary growth or some of the metastases may be colorless, while other growths are heavily pigmented. As a rule, they grow rapidly, and are nearly always fatal within three years; often the period is but a



Pigment containing cells from a spindle-celled melanoma. (Ribbert.)

few months; there are, however, exceptions, of cutaneous melanomas of slow development over many years. The original tumor does not often attain any great size, but metastases are extraordinarily abundant. No other form of tumor affords so many obvious metastases, and these both by means of the bloodvessels and the lymphatics, so that the nearest lymph glands are apt to be involved, along with the liver (a specially favorable seat for abundant and relatively large secondary nodules), the



Section from an alveolar melanoma or chromatophoroma of the great toe. The cells in general are here seen to be free from melanin granules, but these are present in occasional cells both of the tumor (a) and of the stroma (b). At c, some of the melanin-containing cells are drawn separately.

lungs, practically all the viscera, including the brain, heart, bone marrow, the coats of the intestine, and the serous membranes. As usual, the muscles show little involvement.

When we come to examine various cases histologically, we find a marked divergence in the characters of the various tumors, and this whether a series be examined originating from the eye or from the skin.

THE TRANSITIONAL LEPIDOMATA

It is regarding the translation of these appearances that there is such active difference of opinion. We meet with two main types: the first, more common within the eye, composed of relatively small spindle cells; the second, more common in cutaneous growths, formed of large epithelioid cells, with or without obvious large spindle forms, and these tending to be arranged in alveolar masses surrounded by a fairly abundant fibrous stroma. The first, in general, shows not the slightest sign of alveolar structure.

In both forms the pigment, contained as small brown granules within the cells, varies considerably in amount; in some cases it is so densely packed that nothing can be seen of nucleus or cell structure; in these, and in cells containing less amounts, it is present also in the long cell processes. As a rule, in the tumor cells themselves the individual pigment granules are slightly rod-shaped. In the spindle-celled type we encounter densely pigmented globular cells without processes, in which the granules are more rounded and conglomerated. These Ribbert regards as dead cells which have undergone contraction. So, also, in the stroma we see cells containing irregular rounded pigment granules. In both forms we may encounter what are the cells proper of the tumor, wholly devoid of pigment; indeed, Ribbert, as the result of his studies, goes so far as to lay down that all sarcomas of the uvea and interior of the eye (excluding the gliosarcomas) are of the one origin, whether pigmented or unpigmented. In both, but more particularly in the alveolar form, there may be more pigment in the stroma than in the tumor cells proper.

Ribbert lays down that when this is intracellular in the stroma it is still within the same order of cells. This we are inclined to doubt. Certainly there is not the same sharp distinction between tumor cells and stroma as we meet with in ordinary cancers, but when we see that, in advanced cases, leukocytes and the endothelial cells lining the vessels take up the pigment, we cannot deny the same properties to the connective tissue and wandering cells of the stroma. And when, as frequently is to be noted, the tumor cells degenerate and form areas of softening, so that cavities full of a black fluid appear in the growths, then leukocytes pass into these areas, take up the pigment, and, judging from appearances, deposit it in the stroma.

So extensive may be these degenerative processes that free pigment passes into the blood (*melanemia*), and may be discharged into the urine (*melanuria*), besides tinging the tissues in general.

As to the meaning of these two forms of melanotic growth, two opinions are possible: (1) that they are distinct, and that there is a connective-tissue type of growth producing the spindle-celled type, a cancerous form which is the alveolar type; and (2) that they are both produced by one order of cell under different conditions, or at different stages of vegetative activity. There are still those who hold to the former view, but the existence of transitional and combined forms, and the fact, as pointed out by Ribbert, that teased-out specimens of the spindle-celled form afford cells with long and branching processes

MELANOMA

of the same type as those afforded by the other form, renders this view untenable. We must conclude, therefore, that cells of the same type give origin to both, and these cells are the chromatophores, that particular order of cells which, in the normal skin, may be pigment-containing.

But what is the nature of these cells, and what their origin, that they can give origin to tumors of varying type? It is around this question that the controversy ranges itself. Some years ago Unna, in his extensive studies upon the pathology of the skin, first brought forward the view that the cells which give rise to cutaneous melanomas are of epithelial origin, and this view has from many quarters gained adherents.

It must be recalled that the cells of the rete Malpighii contain pigment -melanin-and this in colored races in easily recognizable quantities.



Section taken through epidemis parallel to surface, or somewhat obliquely over a small entancous melanoma, showing typical prokle cells, as at b; others oval (c), containing a few granules of melanin, and others apparently of the same order as at d, densely filled with melann granules.

Prior to 1889 there was no doubt regarding the epithelial origin of melanin; in that year Aeby called attention to certain pigmented cells, now known as chromatophores, lying in the corium and between the cells of the Malpighian layer. These are stellate cells of connectivetissue type, and he, not unnaturally, concluded that they act as carriers, absorbing certain substances from the blood, elaborating them into melanin, and passing them on to the epithelial cells. Even up to the present moment this view has its upholders, and is supported by the sarcomatous, *i.e.*, connective-tissue type of melanotic tumors. Unna, in his extensive studies upon the skin, first brought forward the view that these cells are of epithelial origin, derived from the Malpighian

THE TRANSITIONAL LEPIDOMATA

layer, and that, so, tumors derived from them are more allied to the epitheliomas or cancers than to the sarcomas proper. It has been pointed out, although never, it seems to us, with absolute conviction, that frequently in nevi and in early cutaneous melanotic growths collections of cells of epithelioid, chromatophoric type are present in the epidermis, and constitute downgrowths, passing down into the masses of tumor cells, in which they are with difficulty distinguishable from the tumor elements. Were these cells to become the parent cells of the tumor, the pigmentation would be no new assumption. We have met with one case in which it was difficult to conclude that the epithelium had not taken on a melanotic metamorphosis, but could not determine whether this was directly associated with the underlying growth (Fig. 283).

Now, by analogy, just as we admit that in the transitional lepidomas a retrogression is possible from the lepidic to the hylic cell type, so, I hold, must we be ready to recognize that in epiblastic and hypoblastic tissues extreme anaplasia may be accompanied by a similar order of events; we must be prepared to find that, without conversion into true fibroblasts, cells derived from the epiderm may take on hylic arrangement in the corium and underlying tissues. Indeed, we cannot shut our eves to the fact that this conversion happens at the advancing edge of not a few actively malignant cancers and notably of epitheliomas. In a condition, xeroderma pigmentosum, which is on the borderline, and frequently passes into epithelioma, Ziegler,¹ by subjecting an area of the skin to the action of ultraviolet light for twenty minutes, set up necrosis of the papillary bodies, followed after several weeks by the development of a pigmented scar. In this he found pigment in both epiderm and cutis, with groups of pale cells of epithelial character passing into and present within the scar tissue, of the same order as those seen in the basal epidermal layer. He calls attention to their close resemblance to nevus cells.² As a matter of fact, many pathologists of the widest experience -such as Marchand and Lubarsch-hold the view that chromatophores are of epithelial origin, and melanomas, therefore, epiblastic in nature. But against this view are the following considerations:

 Pigmented tumors of pure epitheliomatous type are curiously rare, and, what is more, even in melanomas of the most characteristic alveolar carcinomatous type, employing Mallory's stain we never encounter an alveolus which throughout is devoid of interstitial substance.

2. If the comparison be made—as it has been—between the melanomas and the rodent ulcer, it is worthy of note that this latter, of all

¹ Quoted by Fordyce, Jour. Amer. Med. Assoc., 54: 1910; 91.

² So, too, Meriowsky (Monatssch. f. prakt. Dermatol., 42, 43, and 44) has followed the development of pigment in the cells, more particularly of the rete Malpighii, after subjecting small areas of skin for a short time to the Finsen light. He has seen the pigment collected more particularly on the side near the source of light, and later has observed those pigmented cells send out processes between the other epithelial cells and into the cutis—assume, that is, the characteristic chromatophore type.

epitheliomas, forms the fewest metastases; of all tumors, the melanoma forms the most.

3. What is noticeable in early nevi, in xeroderma pigmentosum and several other slighter states of cutaneous pigmentation is that the pigment cells in the cutis have a relationship not to the overlying epidermis, but to the vessels. As pointed out by both Ribbert and Borst, the normal habitat of the cuticular chromatophore-of the ordinary chromatophore, that is-is in the lymph spaces immediately around the vessels. Among the latest workers, Staffel,1 studying these perivascular collections, finds every transition from collections of lymphocytes through others composed of lymphocytes, plasma cells, and the allied mast cells showing transition into pigment-bearing cells, either branching and with processes, or of the spindle-celled type. Studying the development of the two orders of cells he notes that the pigment granules are coarser in the cuticular than in the epidermal chromatophores. We are inclined to favor this view of the duality of origin of the cuticular and epidermal chromatophores. We know from the observations of Schridde and others that the plasma cells, like their congeners, the lymphocytes, have the habit of wandering; that they may assume a spindle shape in the tissues; that they may also, according to Maximow, give origin to cells with processes of the clasmatocyte order. Cells of this nature undergoing an orderly proliferation in the lymph spaces would give origin to growths of the alveolar type; growing more actively, would infiltrate and exhibit a more purely sarcomatous structure.

In their habit of growth it will be seen that these tumors approximate to our class of transitional lepidomas; nevertheless, neither in the earliest stages, nor in their mode of spread do they wholly fall into it. There is perhaps a closer alliance to the large-celled lymphosarcoma of the intestine, in which more than once we have found it difficult to determine whether we dealt with a sarcoma or a loosely growing and abundantly infiltrating carcinoma. In other words, the slowly growing lymphosarcoma has a tendency to respect lymph spaces. In this uncertain state, with a leaning toward mesoblastic rather than epithelial origin, we must be content to leave the problem. The existence of transitional forms between the alveolar and spindle-celled types renders it difficult to accept Fordyce's dual hypothesis.

Before closing, an attempt must be made to answer a question which will naturally have arisen, namely, How are we to explain the malignancy of cells apparently so highly specialized? To this the answer must be another question, namely, Is the deposit of melanin granules an indication of specialized function, or, on the contrary, is it one of imperfect metabolism. Nor is this question easy to answer; it demands a knowledge of the nature and origin of melanin, which more appropriately is taken up when we discuss pigments in general (p. 969). We shall not, therefore, discuss here the rival views regarding the functions of the chromatophores—whether they supply melanin to the epidermis

¹ Verhandl, deutsch, pathol, Gesell., 11: 1908: 136.

(Kölliker, Delépine,¹ M. B. Schmidt²), or, on the contrary (as urged by Jarisch, Port, and others), procure their melanin from the epidermal cells. Nor shall we discuss the relationship of melanin to hemoglobin, as emphasized by Ehrmann.³ The modern view leaves these in abeyance, and regards the melanin, as developed within the cell, as a derivative from the nucleolar matter of the nuclei of the melanin-bearing cells (Rössle,⁴ Meirowsky,⁵ Staffel⁶), associated with distinct signs of nuclear exhaustion, not to say degeneration. This, however, does not in our opinion explain everything. These identical nuclear changes have been described in connection with so many cell deposits that we can only conclude that each individual deposit is not a direct development from the plasmasomes or chromidia, but is due to the interaction between the discharged nuclear matter and certain cytoplasmic or paraplasmic substances. Yet other recent indications are that melanin is of the nature of an oxidized product of the aromatic radicals gained from disintegration of the protein molecule; that its presence in the cell represents either an excessive production and activity of an oxidase or a deficiency of the enzyme or other body, which carries the process farther and converts the melanin into its colorless chromogen (melanogen). It may well be that the extraordinary deposit of melanin in melanotic tumors, far from being a progressive acquirement, indicates a deficiency in the disintegrative mechanisms of the cell, whereby the normal final stage of colorless chromogen formation, or of protein disintegration, is not

OTHER TUMORS OF DOUBTFUL RELATIONSHIP.

There remain, it must be emphasized, other tumors whose relationship are not wholly determined; of them may be noted the following:

Cholesteatoma.—This form is found more particularly in association with the membranes of the brain, and is characterized by the presence of little pearly nodules. These are formed of layers of cells of epithelioid or endothelioid type, and in the centre of the masses may be a cluster of cholesterin crystals. In appearance and structure they most closely resemble the endotheliomas, also encountered in these regions, and as such the majority of observers regard them. But Ziegler⁷ in some cases has encountered hairs and hair follieles, and regards them thus as of epithelial origin, secondarily, it may be, to foetal inclusions. They are wholly benign. This form must not be confounded with another condition affecting the antrum of the middle

¹ Delépine, Trans. Path. Soc. Lond., 1891. ² Schmidt, Virch. Arch., 115.


³ Arch. f. Dermat. u. Syph., 64. ⁴ Zeitschr. f. Krebsforschung, 2:1904:291.

⁵ Monats, f. prakt. Dermat., 44 ; 1907 ; 166.

⁶ Verhand, deutsch, pathol, Gesellsch., 11: 1907.


⁷ Vide Boström, Centralbl. f. Pathologie, 8:1897:1.

OTHER TUMORS OF DOUBTFUL RELATIONSHIP



Tumor of the enrotid gland: G, vessels; Bl, hemorrhage into a column of cells; at d the cells of the growth are taking on a more connective-tissue type; at c, hyaline degeneration.

F16, 285



Portion of the same tumor more highly magnified, to show peritheliomatous arrangement of the tumor_cells in relationship to the vascular endothelium, (Paltauf.) 52

THE TRANSITIONAL LEPIDOMATA

ear and external auditory meatus to which the same name is applied. This latter is not a true blastoma, but merely an abnormal heaping up of an onion-like series of layers of epithelial cells undergoing keratinization, degeneration, and necrosis, with frequent deposits of crystals of cholesterin. With progressive accumulation a dense mass is produced, which may entirely fill the antrum and cause erosion of its bony case.

Tumors of the Carotid Gland.—Seated upon the carotid artery close to its bifurcation may be found, upon careful search, a minute cell cluster embedded in a capillary meshwork. It cannot be said that the exact relationship of this carotid body has been determined. The cells are chromaffine—*i. e.*, take up and are strongly stained by chrome salts, and as the cells of the adrenal medulla have the same properties, and they, as now accepted, are of nervous (sympathetic) origin, it is held by some that these are of like origin. In this relationship it may be recalled (p. 757) that retinal gliomas, also of neuroblastic origin, present the liability to exhibit cylindromatous and so peritheliomatous appearance. The occasional tumors originating from the carotid bodies are of a type recalling the peritheliomas, and verge into the sarcomas proper-

Other tumors of irregular type and uncertain relationships have been recorded in connection with another small organ of undetermined relationship, namely, the coccygeal gland (Luschka's gland). These also, as indicated in Fig. 279, p. 824, tend toward the peritheliomatous type. We have, further, encountered two tumors of the palate, apparently primary, curiously resembling the hypernephromas in structure. What is their origin we do not know.

CHAPTER XXIII.

THEORIES OF NEOPLASIA.

THE examples afforded in the preceding chapters indicate:

1. That some of these growths owe their origin to cell rests. In addition to the whole groups of the teratomas and the teratoblastomas and allied forms, which we have considered separately, we have adduced as examples of this order the tumors originating from persistent rudiments of embryonic structures (gill clefts, branchial arch cartilages, etc.), from cells displaced during the course of development (aberrant hypernephromas, columnar-celled cancer originating in areas of squamous epithelium, etc.).

Cohnheim's Theory.—According to the oft-quoted theory of Cohnheim, it is these cells —cells which in the course of development have been displaced from their normal relationship, or have failed to undergo a normal atrophy—that are the essential nidus from which neoplasms originate. The theory, it must be noted, went very little farther; it did not lay down anything beyond the mere postulate of displacement favoring retention of embryonic properties and eventual aberrant growth. It did not explain why it is that not all of the abundant cell rests, which we may encounter in the human body take on aberrant growth, but only an occasional one, and that in exceptional circumstances; nor did it attempt to explain why after remaining latent for years, these begin active proliferation.

2. We are forced to admit that there are cases in which tumors, and more particularly those of a malignant type, originate from cells which must be regarded as having undergone *not congenital*, *but postnatal displacement;* cases, for example, of squamous epithelioma originating in the scar of an old ulcer, of columnar-celled cancer of the stomach or intestines similarly originating in the edges of an ulcer, and it may be of surcoma originating at the site of a contusion. That this is so was emphasized by Roger Williams,¹ but the expansion of Cohnheim's theory to include postnatal displacements is more widely known in connection with Ribbert and his theory.

3. We accept, therefore, that the autochthonous blastomas can originate from cells that have been displaced, and that very many examples—it may be the majority—of tumors come into this category. But does this include all cases? To this question we cannot but return a negative answer. Cell displacement is not the essential. There are undoubted cases in which cells exhibit the cancerous change, and exhibit the cancerous type without any indication of preexisting displacement. Our

¹ Principles of Cancer and Tumor Formation, London, 1888; in many respects a remarkable work, based on broad biological considerations,

THEORIES OF NEOPLASIA

own attention was called to this fact some years ago in a study of an early multiple tumor of the adrenal cortex, in which occasional cells in the immediate neighborhood of the small mass of new-growth, while still retaining their relationship to the columns of the zona radiata, by their enlarged and deep-staining nuclei stood out as of the cancerous type. The number of these and their relationship, wholly unconnected with the main growth, appeared to preclude any possibility of their being outgrowths from the latter; they were part and parcel of the gland substance proper.¹ It is admittedly a matter of chance, and must be rare, to encounter tumors at this early stage, but I have recently come across another tumor of the adrenal showing the same peculiarity, and Jores and others have recorded similar observations. Foremost among these must be mentioned Hauser² and his observations upon the malignant metamorphosis of cells of the intestinal mucosa while still in situ, and Van Heukelom, L. P. Daniels, Tolot, and Horst Oertel upon the direct cancerous transformation of the liver cells in case of multiple carcinomas of that organ. As Oertel states regarding his case: "It was plainly revealed that the origin of the cancer was a transformation of multiple growths, of multiple groups of liver cells, sometimes only involving few within one lobule. These microscopic areas, best observed in those parts of the liver which showed as yet no gross cancer formation, demonstrated a direct change of atrophic degenerated, wasting liver cells into cancer cells while they were still in perfect continuity with each other and still entered into the formation of the lobule." Oertel describes three stages through which those cells pass: a first with extensive loss and granular degeneration of the protoplasm and disappearance of nuclear chromatin; a second of marked nuclear enlargement, the cell body still remaining granular; and a third, in which there is abundant smooth protoplasm around a very large well-formed nucleus, this last being identical with that of the cells of the fully formed cancerous areas. Not until cells of this order had proliferated was their connection with each other lost, and that independent growth observed characteristic of the ordinary carcinoma. This is a process which has for years been actively denied by many of the leading writers upon the subject. For my part I regard it as wholly demonstrated that it does occur; and it must be taken into account in the development of any adequate theory of neoplasia.

4. It is through examples of this order that we link the blastomats with the blastomatoid growths referred to in detail in the preceding chapters, growths originating as multiple diffuse localized hypertrophies of particular tissues, developing obviously not from a single cell, but by a generalized proliferation of the specific elements of a region. These blastomatoid tumors may show every gradation from simple idiopathic hypertrophy to pronounced malignancy.

To recapitulate: From the point of view of the relationship of the cells giving origin to neoplasms, we recognize the following classes:

¹ See Woolley, Trans. Assoc. Am. Phys., 17: 1902: 627.

² Das Cylinder epithelium des Magens und des Darms, Jena, 1890.

1. Teratomas from totipotential cells.

Twin teratomas (feetal inclusions).

Filial teratomas (ovarian and testicular, etc.).

2. Teratoblastomas from multipotential cells.

"Mixed" tumors.

3. Blastomas from unipotential cells.

Originating from congenitally displaced cells.

Originating from cells of postnatal displacement.

Originating from cells that assume neoplastic characters without displacement, and rapidly assume malignancy.

4. Blastomatoid growths.

Originating as a diffuse though local hypertrophy of the specific elements of a tissue, which may or may not pass from the hypertrophic to the malignant type.

The adequate theory of neoplasia must cover all these forms. But a distinction needs to be drawn between the teratomas pure and simple, and all the other orders; once again we find that classification is gradation, with transitional types linking the one class to the other. As was pointed out when discussing these, the typical teratomas exhibit in their cells the orderly progression from embryonic to differentiated tissue, and associated with this we observe that they have a restricted power of growth comparable with the like restricted power of growth of the normal individual. Their cells, it is true, do not form perfect, but incomplete organs, and with this certain of them may sooner or later take a blastomatous growth. When they do so this development of a "tumor in tumore" is an epiphenomenon. There is no primary anaplasia of the cells of origin of these typical teratomas, although when from them there develop secondary blastomatous growths there must be secondary anaplasia of certain of the component cells. And as regards the atypical teratomas (p. 657), it has to be noted that the growths develop, not from cells which have suffered a reversionary anaplasia, but from those which have never passed beyond the vegetative stage or attained the stage of full differentiation. In the teratoblastomas as a body we have indications of transition: some cells (in the mixed tumors of the kidney, for example) exhibit a capacity to develop into recognizable striated muscle fibres-into one of the highest and most specialized of the tissue cellsbut along with these are cells which do not pass beyond the vegetative or "embryonic" type.

It is obvious that to cover all these forms, even of what we classify as blastomas, Cohnheim's well-known theory is inadequate. The same is true of the more elaborate theory of Ribbert.

Ribbert's Theory.—This theory is so frequently referred to at the present time that it is necessary to state its main contentions. According to it, cell displacement is the first essential, and these displaced cells take on active growth, not from any active exaltation of proliferation activity on the part of the cells themselves. With Weigert Ribbert holds that the vegetative powers of the cells cannot be stimulated from without. If, therefore, a cell-rest exhibits active growth, it is because of a diminished external

THEORIES OF NEOPLASIA

resistance, because of a reduction of the antagonistic forces. Changes in the surrounding tissues precede the specific tumor cell overgrowth. He holds that the cell-rest giving rise to a tumor cannot have its cells arranged in the normal order because in such case there would be the normal growth-restraining tension. That a retrograde change in the cells themselves is favorable to growth is admitted, but is regarded as secondary and not indispensable.

Now, isolation of cell groups, irregular disposal of cells, and lack of restraining tissue tension are to be encountered in the healing of wounds in connection notably with the epithelium which actively pushes over and into the underlying granulation tissue. Nevertheless, in such cases neoplastic development is the rare exception, not the rule. What is more, as we have pointed out elsewhere, Weigert's hypothesis is wholly untenable. In this very matter of tumor growth Ehrlich's observations upon transplanted adenocarcinoma of the mouse demonstrate the presence and increase of vegetative power on the part of the tumor cells, for with successive transplantation into fresh animals they may steadily manifest greater malignancy. And were further disproof needed, we have it in Bernard Fischer's interesting studies upon the chemiotactic and proliferative stimulation of the squamous epithelium of the rabbit's ear by Scharlach R dissolved in oil.

Fischer,¹ from one side of the ear, injected into the other side through the cartilage, a solution of Scharlach R in olive oil, and found that this produced a great proliferation of the Malpighian layer of the skin of the other side, so that this formed finger-like processes passing into the subcutaneous tissue down to the droplets of oil, and even, in some cases, into the track of the needle through the cartilage. With this the accumulation of injected oil diminished, leaving the Scharlach R in a solid form in the tissues. With the absorption of the oil the proliferation ceased, and the epithelial processes underwent atrophy. There was no development of autonomous new-growth, but the convincing demonstration of a stirring-up of the vegetative activities of the epithelial cells. These observations have been abundantly confirmed, among others by Klotz in our laboratory at the Royal Victoria Hospital.

Lastly, it must be noted that the theory takes no account of the assumption of malignant properties by cells *in situ*. Valuable as it has been in its time, the theory has shown itself inefficient.

Parasitic Theories.—There are certain data which have accumulated during the last fifteen years regarding the incidence of atypical malignant blastomas, and more particularly of cancer proper, which we freely admit are, upon their face, difficult to reconcile with any theory save one requiring the increasing spread of some microbic causative agent. These are: (1) The rapid increase in the mortality from cancer in most civilized countries; (2) the greater incidence of the disease, more particularly in certain low-lying localities, estuaries, and the borders of sluggish streams;

⁴ Verhandl, d. deutsch, pathol, Gesell., 10: 1907; 20, and Münchener med, Woch., 53: 1906; 2041.

PARASITIC THEORIES

(3) house incidence, certain houses affording a mortality from cancer over a series of years in striking excess over the average.

Upon the last we lay no weight; by the law of chance, just as one individual in a thousand may be of gigantic proportions, so one house in a thousand may show a great excess of cases of cancer-or of twin birthsover the ordinary run of houses. But Behla's full study of the incidence of cancer in the different sections of a little German town is certainly most suggestive; the cases occurring in the low-lying houses near the sluggish stream were found far in excess of those in other situations.¹ In this he confirmed the earlier work of Haviland in England, and while as regards the general increase in cancer some weight must be laid upon the fact that improved hygiene and care of the infant favors the survival of the weakly, and even brings them to middle age, so that (a) the average length of life has been increased, or, in other words, more individuals are protected from death by other diseases that they may reach the cancer age (after thirty-five years) and succumb to malignant growths, and (b) more individuals with constitutional weakness survive than was the case in former years; nevertheless, the increase would seem to be proceeding at a more rapid rate than can reasonably be explained along these lines. Newsholme, it is true, a leading English authority upon statistics of morbidity, still doubts whether the increase is not merely apparent and explicable along these lines. But in San Francisco the relative number of deaths from cancer increased seven times in thirty-two years, from 16.5 per 100,000 in 1866, to 103.6 in 1898. In Boston the rate trebled between 1863 and 1887. In New York State, according to Roswell Park, there were 2363 deaths from cancer in 1887; eleven years later there were 4456. According to Tatham,2 in the period 1861 to 1870 the annual rate of cancer mortality per million living in England and Wales was, in males, 242; in 1891 to 1900 this had increased to 597, an increase of 150 per cent. For females the increase was 74 per cent., from 519 to 903.

Wützdorff's³ statistics for Germany are equally remarkable. Taking deaths in hospitals, in 1879, 6330 were attributed to cancer; in 1898, the number had risen to 24,000. Even when the correction is made for increase in the number of hospital patients, the increase in the cancer death rate is 266 per cent. In the ordinary returns for deaths: in 1892, 2.6 per cent. of all deaths were returned as from new-growths; in 1808, (in but six years) the number had reached 3.5, an increase of 18.5 per cent.; it is attacking also at an earlier age than before, and attacks more men.

Nor is it that nowadays more cases are correctly diagnosticated than formerly; the postmortem statistics of certain old-established hospitals reveal an increasing ratio of cases found to be cancerous at autopsy, and this in different parts of the world. One of the most careful recent papers on this subject is by Barlow and 'Taylor upon the statistics of St. George's and Middlesex Hospitals.⁴

¹ Poppelmann has recently confirmed these results in the case of another small town. Zeitschr. f. Krebsforschung, 4: 1906; 39.

² Lancet, London, 1902, i, 745. ³ Brit. Med. Journ., 1902; i; 805.

⁴ Med. Exam. and Prac., New York, 1905: 719.

But, as already noted, if malignant growth be due to microparasites, there is no general consensus as to the nature of the causative organism; on the contrary, among those in favor of the parasitic theory there has been a most extraordinary diversity of findings and of opinions. Bacilli (Schuller), micrococci (Doyen), blastomycetes (Russel, Sanfelice, Leopold), ameebæ and rhizopod forms (Sjöbring, Schmidt), sporozoa (Sjöbring, Malassez-1889); "coccidia"-(Metchnikoff, Soudakewitsch, Ruffer, Plimmer), a gregarine "Rhopocephalus" (Korotneff)—have all had their advocates, and, what is more, not infrequently the same observer has not hesitated to describe now a causative agent of one order, later one of a wholly different order (Sjöbring, Gaylord). The latest form found associated with malignant growths is almost naturally the spirochete. Observed first by Borrel and some European workers, Gaylord has made the fullest study of the same, and has found a remarkable form of spirochete associated with a large proportion of cases of inoculated and natural adenocarcinoma of the mouse. There is no doubt that this form is present at least in some of these, it may be in all, and that it is of the nature of a spirochete. What is more, Forbes Robertson¹ has encountered identical forms in a small number of cases of human carcinoma. Warned by previous experience, Gaylord exhibits a wise caution in drawing any deductions; nay, more, has emphasized that the distribution of the form makes it impossible to regard it as the essential causative agent.

Emphasis has been laid upon the relationship between the ova of Bilharzia and the development of vesical and rectal carcinoma (p. 777), but these would appear to be at most mechanical irritants. So, also, attention has been drawn to the relatively huge tumors caused in the cabbage by the *Plasmidiophora brassica*, a minute intracellular vegetable parasite, and other local overgrowths in plants due either to allied vegetable parasites or, as in the case of the abundant galls of different orders, to irritation induced by insects. There can be no doubt that in plants at least there exist irritants, and even intracellular parasites, setting up just that order of stimulation which induces cell proliferation rather than cell degeneration, the growth continuing long after the primary irritant has ceased to act.

Attention may also be called to the enzootic goitre and thyroid tumors affecting the fresh-water fish in certain waters in Bavaria and in the United States, a condition so widespread and leading to so great a loss of fish in certain hatcheries as to cause great concern to the United States Government. This has been studied more particularly by Gaylord and the New York State Cancer Commission. The fact that fish from healthy districts develop these thyroid growths when placed in the incriminated waters strongly suggests infection. It is noticed that if young fry be placed in a series of tanks fed the one into the other, those in the upper tank, supplied with perfectly fresh water, may wholly escape; whereas the greatest incidence of the condition occurs in the lowest tank

¹ Lancet, London, 1907 : ii.

ANAPLASIA

of the series. With Marine we admit that in the majority of the fish the condition is one of goitrons hyperplasia of the diffuse non-encapsulated thyroid tissue of these animals, but with Gaylord must recognize that in a certain proportion of cases infiltration of the vessel walls and other tissues is definitely present, *i. e.*, a state of malignancy. But contrary to our conception either of ordinary infection or of ordinary neoplasia is the fact (parallel to what has been noted with regard to early goitres in man and the domestic animals) that removal of the affected fish to unaffected waters may lead to disappearance of the tumors. This strongly suggests that for the progressive hyperplasia with eventual blastomatosis we must invoke the action of a progressive or recurrent irritant which, if of infective nature, can only be of the class of subinfections (p. 462). For the present we can only safely ascribe the condition to overcrowding of the fry, with malnutrition, in certain predisposing waters.

Very much more work, and widespread confirmation of results, is necessary before we can be prepared to lay down that any form of microparasite is the specific causative agent of any form of malignant growth in man. Our present stand must be one not of absolute denial, but of agnosticism. But even granting that it is ultimately found that certain microbes set up certain orders of growth, it must be recognized that the microbic theory obviously cannot be applied to neoplasms in general. It can only have a limited application, and cannot be the foundation of the general theory of neoplasia. And this because it is equally obvious that there are certain orders of tumors which are to be ascribed to a totally different mode of causation-to the inherent, if aberrant, vegetative power of the constituent cells, and to that alone. We need no parasite to explain the aberrant growth of the syncytium which produces the chorio-epithelioma. There we deal with cells possessing naturally invasive and erosive powers, cells of another individual.¹ The same is true of all the teratomas and teratoblastomas and tumors originating during fœtal existence. Nor can those congenital cases of what we have termed blastomatoid be explained thus. In other words, the adequate theory of neoplasia must be one which will explain not cancer alone, but all types of tumor formation. No parasitic theory suffices to do this.

It is evident from the above that we are driven back to a *change in* the biological properties of the cells giving origin to tumors, and to look for some explanation of what it is that initiates the change. Here we have to choose between a long series of hypotheses.

Anaplasia.—It is to von Hansemann that we owe the first thorough study of the histological characters of cells of malignant growths. Cornil, in 1886, and Klebs, had previously called attention to the existence in these tumors of irregular and atypical mitoses. Von Hansemann regarded these as evidence of cell change, of the production of generations of cells which through altered distribution of nuclear matter do not so much undergo degeneration proper, but become incapable of attaining perfect structure and function. This modification he has termed

¹ Vide Adami, Syncytioma Malignum, Clinical Journal, Lond., June 18, 1902.

anaplasia, cell races being formed, possessing abnormal properties, one of which is that of increased vegetative activity.

But (1) among neoplasms this unequal distribution of chromatin is characteristic only of malignant tumors; (2) it has been observed in other conditions besides malignancy (in forms of inflammation in the lower animals, which by experience we know do not lead to neoplasia); and (3) no explanation is afforded of the cause of the irregular mitoses.

Farmer, Moore and Walker,¹ Bashford, and others have more recently enunciated somewhat parallel views, describing ring form of chromosomes, such as are seen in the process of nuclear reduction of the oocyte and the spermatocyte, but upon further study have withdrawn their contentions.

Others, again, of whom, if we mistake not, Creighton some twenty years ago was the first, have described an adulterous connection, with nuclear fusion between the cells in malignant growths, and to this "rejuvenation" of the cells have ascribed their increased vegetative activity. The latest of these is Moore.² Certain of the stages of amitotic nuclear division are curiously like the figures afforded by this last observer. We see no adequate support for this theory, which again has no bearing upon the large mass of non-malignant tumors.

Bashford's Theory.—Another English worker, Bashford, from a study of the age incidence of malignancy in man and various species of animals, concludes that the lighting up of aberrant proliferative activity is a function of cell senescence: that as different orders of cells have different life periods, so may they give origin to tumors at different periods during the life of the individual. This hypothesis again covers only a small part of the ground (*e. g.*, does not include the teratoblastomas), and is inadequate.

From these we pass to theories of a wider scope.

Hauser's Theory.—One which, in point of time, we should have mentioned among the first is that of Hauser. Just as among the members of a species of animal or plant there may be variation, so here he postulated that among the descendants of a single cell, the ovum, cells might make their appearance exhibiting active vegetation coupled with modified properties, the descendants of which constitute a neoplasm. He regards this variation as favored by change in nutrition with excess of the same. As pointed out elsewhere (p. 597), it does not appear to be surely established that mere excess nutrition is in general a cause of hypertrophy and proliferation.

The Habit of Growth.—We ourselves^a have laid stress upon the consideration that the cell that is differentiated for the performance of function in the performance of that function uses up energy and cannot simultaneously store up energy to any extent for purposes of proliferation; it is the cell that either has not yet undergone differentiation or the one that has passed from a fully formed to a less differentiated state

^a Adami, Brit. Med. Jour., 1901 : i : 621.

² Ibid., 79: 1907.

¹ Proc. Royal Soc. London, 72: 1903.

THE HABIT OF GROWTH

that is capable of active proliferation, not merely, as Bashford would hold, cells of the exhausted and senescent type, but premature or immature cells also; that the mere existence of cells of this order in the body does not in itself initiate blastomatosis, nor even a stimulus of the ordinary type acting on such cells; that just as cells modified by inflammation or irritation undergo metaplasia, and from these modified cells we find originated tumors departing from the type (for example, in the inflamed gall-bladder, the everted uterus, and other mucous membranes in which a flattened epithelium has replaced one of more columnar type, we meet with carcinomas approximating toward the epitheliomatous type), so in like manner cells which for a sufficient length of time have been so placed or so acted upon that they have been unable to perform function, while they have continued to gain nourishment, assuming the less differentiated, actively vegetative stage, gain the habit of growth, or, in other words, lose the habit of function; and it is the assumption of the habit of growth that is the point of origin of the neoplasm. This hypothesis, it is true, covers both cells, which as in the teratomas and teratoblastomas have been so situated that they have never been able to assume full function, and the blastomas developing whether from cell rests or from cells in situ. It demands that the grade of lack of assumption of complete adult type of the tumor cells is the expression of lack of development, or extent of reversion, of the parent cells of the tumor at the time when the habit of growth manifested itself.

Beneke¹ has given utterance to somewhat similar views. He regards blastomatosis as due to increase in growth energy on the part of the cells with contemporary lowering of the functional activity.

Other observers, however, have sought for a more tangible explanation of the process of *Entdifferenzierung*, anaplasia, or kataplasia, that is so obvious a characteristic of tumor cells in general.

Marchand² emphasizes the fact that the difference between the normal embryonic cell and the so-called embryonic cell of tumor lies in this, that the former has the potentiality with continued growth to undergo eventual differentiation, the latter has not merely passed into a latent state as regards functional and structural differentiation, but has actually and permanently lost the potentiality to undergo such differentiation. He urges that there must first be a degenerative change in the cell leading to faulty metabolism, and that the products of the perverted cell have a toxic action upon the other cells in their neighborhood, weakening them and in this way leading to unrestrained growth. This appears to apply specially to the malignant blastomas, and is not so easily applied to benign growths or to teratomas and teratoblastomas. Undoubtedly the trend of recent work is to show that malignant tumors excrete or afford substances, some of them of the nature of enzymes, which are of toxic nature, and it is a reasonable view that these tell especially upon the immediately surrounding tissues, but the theory leaves it open to determine what is the cause of the primary degeneration.

¹ Berl. klin. Woch., 1905 : No. 22.

² Deutsch. med. Woeh., 1902 ; Nos. 39 and 40,

Oertel's Theory.-On the basis of his studies already noted, Horst Oertel seeks to give an explanation why the tumor cells have so completely lost the power of undergoing subsequent full differentiation; why, to employ my terminology, they have gained the habit of growth, and lost that of function. The researches of the embryologist have shown that nuclear chromatin is not homogeneous; the existence and passage on from cell to cell of various orders of chromatin loops with curious persistency would seem to indicate that the different orders of loops convey different properties, while among the protozoa we find repeated examples of the existence of double nuclei, the one evidently associated with propagation, the other associated with the general or functional activities of the cell. Oertel thus suggests that in man and the metazoa in general the single nucleus contains chromatin of two orders-the one governing the functional, the other the proliferative or vegetative activities. What constitutes the primary cancer (or, we would add, other tumor) cell is a cell which has undergone loss of certain chromatin constituents of the former order; where these are lost the cell cannot reproduce them. The cell which has lost the chromatin controlling certain differentiating attributes. can only give rise to daughter cells minus these, but endowed still with vegetative attributes. In this way races of cells are developed in which vegetative attributes are predominant, but functional attributes to a greater or less extent have become lost. It is a matter for future research whether these two orders of chromatin actually exist in the mammalian cell, but the conception is admirable and has some justification, while it tumors in general approximate to the vegetative rather than the highly differentiated type, as also for Marchand's postulate.

Conclusions.—We have, it would seem, arrived thus far, that we recognize definitely among the blastomas some change in the biological properties of particular cells as an essential for neoplasia proper. We recognize, that is, an inherent and permanent alteration in the properties of the cells that constitute, or are to constitute, the neoplasm. It is not something from without that determines the continued growth, not an external stimulus, nor again a diminished external resistance. An external stimulus, it may be, starts the cells on that path which leads eventually to their assuming neoplastic properties; diminished external resistance may well favor active neoplastic growth; nay, more, it may well be that cells of a malignant type afford secretions inhibiting the growth, by depressing the vitality, of surrounding tissue cells. But all these are subsidiary. What is of primary importance is that the cells giving origin to an autochthonous new-growth are so modified that the energy acquired by the assimilation of food is not in the main discharged in the performance of function, as in the healthy cell in normal relationship, but is characteristically retained and accumulated for purposes of cell growth and cell multiplication.

What is the nature of this change we do not know with certainty; it is here that we have to turn to hypothesis. To lay down along the line of Hauser's theory that certain cells of a tissue in the course of their multi-

CONCLUSIONS

plication give rise to mutations, while satisfying the conditions that it is only one or a very few cells in a given tissue that give origin to a newgrowth, just as in a given species it is only one or a very few individuals that undergo mutation and become sports, does not carry us very far; on the contrary, it is apt, unless employed with proper appreciation, to check further advance; whereby we mean that too often the term sport, or mutation, is regarded as synonymous with chance variation. So employed it would in this instance indicate that neoplasia is without definite cause and that it is hopeless to seek farther. Biologists, however, are coming to see that mutations in animals and plants are not chance occurrences. It is being recognized that alterations in environment favor their more frequent appearance. Regarded thus, it is reasonable to regard the cancer cell as a mutation of the normal cell, for such a view is quite in harmony with the clinical observation that certain conditions-life period, trauma, chronic inflammation, etc.-favor the appearance of the blastomatous change. Like considerations apply also to the conception of anaplasia and to Beneke's view of modification in the cell properties, and to Oertel's most recent theory. There is, indeed, a close relationship between von Hansemann's theory of irregular distribution of chromosomes and Oertel's of using up or destruction of specific constituents of the nuclear chromatin of the adult cell. All recognize a change in cell properties, but none lays down what is the underlying influence or stimulus which brings this change about. Here it seems to us that our theory goes somewhat farther in postulating more fully than the others the nature of this change, namely, that there is no one stimulus, microbic or physical, that is responsible for the change in cell properties. As shown by a study of the life histories of different types of tumors, now it would seem to be merely long continuance in a state in which, through cell displacement, vegetative activity is possible but functional activity is inhibited, now a grade of chronic inflammation with accompanying arrest of function without arrest of nutrition, at one time of mechanical, at another of microbic origin; now, it may be a senescent exhaustion of the functional capacities of the cell that favors the assumption by that cell of vegetative to the exclusion of functional activities, an assumption accompanied by alteration in the histological characters of the cell. The more we study the life histories of the different orders of tumors the more we investigate the circumstances associated with the development of different examples of one particular order of tumor (such as the carcinomas), the more it is impressed upon us that there is no one specific causative agent, that a multiplicity of agents induce a particular grade of cell reaction; all these causes may lead to a modification in the cell properties, a modification that is not transient but permanent and conveyed to subsequent cell generations.

In a criticism of the first edition of this work Dr. C. P. White' inquires how, in accordance with the views here expressed, the origin of the squamous-celled epithelioma is to be explained. Can the cells of the

¹ Med. Chron. Manchester, June, 1909.

epidermis be said to cease to perform function before they give rise to a carcinoma? The admirable studies of my colleague, Dr. Wolbach,⁴ upon *x*-ray dermatitis and carcinoma (which, as has rightly been stated, is the first carcinoma produced experimentally) give an answer to this question wholly in harmony with the views here expressed, and what is more, to a large extent they harmonize our views with those of Ribbert, which are that, in cancer, connective-tissue changes precede those occurring in the epithelial elements.

There have now been numerous cases of epithelioma following x-ray burns. Of the tissues of the skin the epidermis is the least susceptible to direct injury by the Röntgen rays, but marked atrophic changes of a progressive type take place in the underlying corium, with dilatation, thrombosis, and eventual obliteration of the superficial vessels and lymphatics, resulting in a dense, relatively avascular layer immediately beneath the epidermis, with here and there areas of necrosis manifesting themselves months after the original injury. The malnutrition of the epidermis thus induced leads in some areas to distinct atrophy of the same, in others to a reactive proliferation² with incomplete or absent differentiation into normal epithelium, a clear evidence of change of function. "The factors responsible for the acquisition of great power to proliferate, and eventually the properties of malignancy, are those furnished by the changes in blood supply and in the connective tissue." In this way a slow augmentation of growth power is achieved, attended with loss of differentiation. The downgrowth of the epithelial processes even into thrombi in the underlying vessels and into the dense collagenous material, gains its simplest explanation along the lines of Fischer's Sudan III experiment in the rabbit's ear, namely, that with their normal nutrition reduced, the epidermal cells make their way, chemiotactically, through the dense corium toward regions of adequate blood supply. This acquisition of malignant powers in these cases of x-ray dermatitis is completed during years, it may be, of active proliferation accompanied by progressive impairment of nutrition. Of this we have microscopic evidence, and the hypothesis that the former is a direct sequence of the latter seems wholly justifiable In short, it is seen that the cells gradually acquire greater capacity for securing nutrition, with coincident manifest increased vegetative capacity and loss of specific functional activity, and finally become capable of living at the expense of other cells. This explanation would seem valid for the forms of epithelioma following upon lupus, syphilitic ulcers, burns, and other conditions inducing super-

¹ Jour. of Med. Research, 21: 1909;415. Dr. Wolbach's conclusion regarding the histological changes in these cases are in substantial accord with those of Unna (Fortschr, a. d. Gebiete d. Röntgenstrahlungen, 1904-05; No. 8) and Wyss (Beitr. z. klin. Chirurg., 44: 1906, and D. Arch, f. Chirurg., 93; No. 6).

² This would appear to be a principle of wide application that lowered but not wholly arrested nutrition leads to a reactive proliferation; we see it occurring in the amoba when the water in which it floats is about to dry up, and at the other end of the scale in the paradoxically large family of the poor curate. It is one of the outcomes of the struggle for existence.

CONCLUSIONS

ficial cicatrization, as also for the occupational epitheliomas (of chimneysweeps, workers in tar, paraffin, anilin, etc.) and for the Khangri cancer of India, induced by the application of charcoal braziers over the abdomen for purposes of warmth and comfort. It would seem also to satisfy the conditions which favor the development of senile cancers. Nevertheless, it must be remembered that it is not a universal generalization. To mention an example, it does not explain the development of papillomas. In them, instead of there being an atrophy with cicatrization of the underlying connective tissue, the proliferation of the epithelium, however brought about, leads to a reactive growth outward of that connective tissue. It is not pretended, that is, that Wolbach's observations are of general application; but they afford valuable information regarding the genesis of one important group of malignant growths. The broader generalization is that the specific tumor cells gaining increased vegetative activities, according to circumstances either stimulate the underlying stroma to afford increased nutrition, or make their way toward areas of increased nutrition: or, according to the circumstances of the case, either the Mountain comes to Mahomet, or Mahomet goes to the Mountain.

Finally, it must be asked what bearing has such a conclusion upon the quest for means of arrest and cure of malignant and other growths?

It is obvious, in the first place, that we hold that little is to be gained from the search for any parasitic cause. Even if found, we believe that once the cancer has taken on active growth, the mere destruction of the parasite would not modify the properties already impressed upon the cell.

It this be so, our attention should be directed in the immediate future not to the search for the cause of malignant and other growths, but to a careful investigation of the properties of tumor, as distinct from normal, cells, and more particularly to the inquiry into the factors which influence the growing powers of those cells.

Two possibilities suggest themselves, one along the lines of active bacterial immunity, the other along the lines of passive immunity.

1. As shown by Gaylord and Clowes,⁴ and later by Ehrlich, by Bashford, and others, in a definite proportion of cases of successful inoculation of certain mouse tumors there is to be observed a later shrinkage and disappearance of the growths, and if now it is attempted to reinoculate these mice with the same or allied tumor material, the results are negative. As also it has been found by Sticker² that where one of these experimentally induced tumors is growing actively in one region, a reinoculation in another place is apit to be without results.⁹ Sticker noted that

¹ Jour, Amer. Med. Assoc., 47 ; 1905 ; 206. ² Münch. med. Woch., 1904 ; 39.

³ While this has been passing through the press Bashford points out that this statement is true regarding reinoculation with the same type of tumor from another animal, but not as regards inoculation with portions of the animal's own tumor into another region (Eighth Annual Report, Imperial Cancer Research Fund, Brit, Med. Jour., 1910; ii ; 206.

THEORIES OF NEOPLASIA

while two implantations of mouse cancer made simultaneously into a mouse would both grow, after successful implantation in *one* region, inoculation in another was negative.

The phenomenon appears to be identical with what we encounter in syphilis, for example, and has been shown by Koch and others to occur in tuberculosis. And as in the latter case we are convinced that it is due to the development of antibodies by the organism, so in these cases we must conclude that the growth of the tumor cells at a focus leads to a like production of antibodies, and that here, as well as there, it is only when the growth of the parent tumor is so active, and the discharge of its products so great, that, as suggested by Ribbert, the antibodies are neutralized, the tissues becoming exhausted and the production of antibodies inadequate, that secondary inoculation or spontaneous metastasis formation becomes possible. If by inoculation of tuberculin into one in not too advanced a stage of tuberculosis it is possible to so exalt the general resistance of the organism, and so to increase the production of antibodies that the focal development of the tubercle bacilli is arrested and their death eventually brought about, so it would seem possible that extracts from the removed primary growth of a tumor might be employed to exalt the specific antineoplastic substances of the body at large, and so to prevent recurrence and bring about the atrophy and disappearance of metastases.

This has been proposed by Ribbert,¹ and at the present time Coca and other workers are engaged along these lines with results that are promising, the indications being against Ehrlich's opposing theory of "atreptic immunity." Employing successive subcutaneous injections of extracts of malignant growths sterilized with 0.5 per cent, carbolic acid Coca and Gilman³ report a series of cases of absorption of cancers with absence of recurrence for periods up to seven months, and this with extracts made from the main mass of the patient's own tumor and from tumors in others of like type.

Too much must not be expected of this method. The general experience is that antibodies to the ferments and other substances developed by the functioning body cells are produced to a very slight extent. The observations of Flexner would seem 'to indicate that not all inocutable tumors induce sufficient general reaction to lead to the inhibition of secondary growth. Further, it is difficult to see how the wise dosage is to be determined, for each individual and each individual case. By analogy it might easily be possible, as with tuberculin, to give doses of the extract which would have the contrary effect.

2. The second mode of destruction of new-growths that has to be

⁴ Philippine Jour. of Science, December, 1909.

¹ Deutsch, med. Woch., 32 ; 1906 ; 1693.

² Ehrlich could find no evidence that malignant growths induce the production of antibodies in the system, and thus postulated that failure of tumors to grow in a host, or retrogression of the same, depended upon the absence of substances necessary for the growth of the tumor cells.

CONCLUSIONS

tested is along the lines of passive immunity. It would seem quite within the bounds of possibility that substances may be discovered, whether drugs or animal products, or agents like the Röntgen rays and radium, to which the vegetative cells of the different orders of neoplasms will be more sensitive than are the fully differentiated cells of the organism. The indications are that we are approaching the successful application of bodies of this order. However inaccurate the reasoning that led Beard to suggest the use of pancreatic ferments, there is evidence that in certain cases these ferments act upon the cells of malignant growths, leading to their destruction and absorption, whereas they do not influence similarly the healthy cells. Von Leyden has extracted from the normal liver of animals a preparation of ferments which, applied to malignant tumors, is stated to cause their destruction with extraordinary rapidity, and Bier states that he obtains extremely favorable results of like order by hypodermic injections of pig's serum.¹ It cannot be said as yet that any of these enzyme methods have proved satisfactory in practice. The most striking results have been obtained in a few cases by Hodenpyl² through the employment of what we may term pure passive immunity, by treatment of cancer patients with the peritoneal fluid of a woman in whom a mammary cancer with extensive abdominal metastases underwent spontaneous retrogression and absorption. There was a persistent ascites, apparently from cicatricial obstruction of abdominal vessels. According to the more recent reports, although some patients have shown a remarkable arrest in and reduction of their cancers, others are either unaffected by inoculations of the fluid or actually harmed by the same. This might be expected. Nevertheless the outlook for the development of a rational antineoplastic therapy and for ultimate triumph over one of man's most terrible scourges is far from hopeless.

¹ Articles bearing upon this subject are Blumenthal, Ergebnisse d. exp. Path. u. Therapie, I: 1901; pt. 165. (It may be noted that this observer was working on the effects of trypsin upon cancer before Beard's mode of treatment was indicated.) Beard, Lancet, 1902; i: 17; 1904; ii: 1200; 1905; i: 281; and New York Med. Rec., (1907; 24. Martin, New York Med. Rec., (09): 1906; 893. Neuberg and Ascher, Arb. a. d. Pathol. Instit., Berlin, 1906. Bergell, Ztschr. f. Krebsforschung, 5; 1906; 204. Pinkuss and Pinkus, Med. Kluik, 1907; No2 Sa and 29. Von Leyden, Deutsch. med. Woch., 1907; 913. Bier, Deutsch, med. Woch., July 18, 1907.

² Med. Record, N. Y., February 26, 1910.

CHAPTER XXIV.

CYSTS.

By the term "cyst" we understand a sharply limited and abnormal accumulation of fluid in any area unprovided with a channel of outflow. The mere localized infiltration of a tissue with fluid, such as occurs in acute inflammation, in local cedema or elephantiasis, does not constitute a cyst; there must be a well-defined boundary wall or sac. An abscess, likewise, does not come under this term; its boundary is not defined with sufficient precision. The existence of a channel of outflow removes a well-defined collection of fluid from the category of cysts; an aneurysm, for example, communicating as it does with the arterial lumen, is not a cyst. We speak of all such localized dilatations of channels containing fluid as examples of *ectasia*; in addition to sacculated aneurysms, the varices or phlebectases (localized dilatations of veins) come under this heading, as do lymph varices and the sacculations or diverticula that may occur along the course of the digestive tube.

That the accumulation is spherical or oval is also involved in the conception, as, again, that the fluid fills the sac; thus, we never speak of the accumulation of fluids occupying the larger serous cavities as cysts, even though etiologically these be identical in origin with accumulations in smaller serous cavities which we include among the cysts. It is a matter of illogical convention that a hydropericardium is not regarded as a cyst, whereas a hydrocele of the tunica vaginalis is so regarded.

Understood thus, the true cysts form a very heterogeneous collection, so various in origin and character that save as a study in etiology little benefit is gained from bringing them together under a common head. It is usual to discuss them in association with tumors, and they may as well be grouped together here as anywhere, although it must be understood that, save for the fact that they constitute local swellings, as a body they have nothing in common with the neoplasms. At most, this is to be recognized, that as a cyst grows in volume there occurs *pari passu* a growth in the tissue constituting its wall, but that growth does not constitute the cyst; it is not primary, but is governed by the pressure of the fuid accumulation. The formation of multiple cysts is a feature in one group of neoplasms, but these *cystadenomas* form a relatively small proportion of the cysts that occur in the organism. The term "cystoma"

Classification.—Broadly we may classify the cysts into four groups of wholly distinct origin: (1) Those due to abnormal dilatation of preexisting cavities of the organism as a result of secretion into those cavities at a greater rate than absorption proceeds from the same. (2) Hemorrhagic cysts, due to the escape of blood out of the vessels into the tissue

SECRETORY CYSTS

and subsequent encapsulation of the same. (3) Necrotic cysts, due to local death of tissue and liquefaction with encapsulation. (4) Parasitic cysts, due to the development (in itself normal) of metazoan parasites within the organism, such parasites possessing a cystic or saccular stage of development.

I. SECRETORY CYSTS.

Of these, the first constitutes the most important and largest group. We may subdivide this according to two methods, either according to the nature of the lining cells, which afford the secretion, or according as to whether the cysts be of congenital origin, due to developmental defects, or postnatal, due to acquired conditions. According to the first of these classifications we can further form classes in which the secreting cells lining the cysts are originally formed of:

1. Cubical or columnar "glandular" epithelium.

2. Endothelium.

3. Ependyma.

4. Squamous epithelium.

5. Composite.

Inasmuch as the congenital cysts with a few exceptions come into the first class, this would seen the more convenient classification to adopt, and this more particularly because there is a certain number of cases in which a given form of cyst of sundry organs may be either antenatal or postnatal in origin.

Regarding all of these, it is to be noted that when cellular activity leads to secretion of fluid into a cavity unprovided with a duct or passage of outlet, or when the passage is obstructed, that fluid is secreted against a certain pressure. The pressure in general is low, only a little above that of the mean blood pressure in the capillaries, but as the secretion is continuous and the absorption through the surrounding vessels and lymphatics is less rapid then the discharge into the cavity, there results a gradual distension of the cavity. It is under these conditions of moderate as distinct from excessive strain that, as pointed out elsewhere (pp. 103 and 593), cell multiplication is favored, and, as a matter of fact, we find that a cyst developed from a narrow tube or duct, as it grows in diameter, continues for long to be lined by epithelium or cells of normal type; there is actual growth and increase of the lining membrane, as of the underlying fibrous stroma, and this adapts the chamber to the increased contents. In this way a cyst may attain very great size and continue to be lined by a typical epithelium of one or other order. Eventually the lining cells exhibit a tendency toward atrophy and flattening, whether through overstimulation and subsequent exhaustion of the growth energy of the cells, through increased internal pressure, through malnutrition of the lining cells as a consequence of the progressive dilatation of the cyst, whereby the nutritive capillaries of the outer wall become flattened and the circulation is continued with increasing

CYSTS

difficulty, or, lastly, through the deleterious effects of the cyst contents. The watery contents of such a cyst are constantly undergoing absorption; the result is that the less diffusible products of secretion come to be more and more concentrated, until in the kidney and other organs we may encounter cysts filled not with water, but with inspissated, thick even jelly-like or firm colloid contents. Which of these features plays the predominant part we do not know, but in old cysts it may be impossible to recognize any specific lining epithelium.

We shall in our review of the different forms of cysts call attention to the various factors leading to the formation of these cysts: obstruction of ducts, whether congenital or postnatal, persistence of portions of glandular and tubular organs which should have undergone complete retrogression during feetal life, inflammation, distension of ductless vesicles present in the adult organism, etc.

"Glandular" Cysts.—I. Of Antenatal Origin.—Congenital Cysts Due to Persistence of Parts of Embryonic and Fetal Duets.—A very large group of cysts comes under this category. In the complicated metamorphoses of the embryo and foctus it may happen that, with the closure of certain passages and atrophy of the same, portions of these do not undergo complete absorption. Such portions tend to form isolated tubular structures embedded in other tissues, and either immediately or it may be only after many years their cells may take on active secretory functions, cyst formation being the result.


Of such may be mentioned: (a) Thyrolingual custs in the median line of the neck, through distension of isolated remains of the primitive thyrolingual duct leading downward from the foramen cecum of the tongue. It is the distal branches of this that form the thyroid gland. (b)Branchial custs, found in the remains of the second and lower branchial clefts; these occur upon the side of the neck from behind the angle of the jaw (second branchial cleft) to just above the sternoclavicular articulation.¹ Cysts of this order may either contain mucous fluid or sebaceous material, according as to whether they originate from the inner end of the branchial cleft (lined with mucous membrane) or from the outer (lined with squamous epithelium). (c) Vitello-intestinal, projecting internally or externally in the neighborhood of the navel. These are lined with mucous membrane and arise from a remnant of the vitello-intestinal or omphalo-mesenteric duct, the old communicating channel between the small intestine and the yolk sac. Similar cysts have been described in the neighborhood of the small intestines, communicating with this by a fibrous cord. (d) Urachal cysts, situated in the lower abdominal region, due to persistence of portions of the urachus.

(e) Cysts of the Primordial Genito-urinary Passages in the Female.— The Wolffian body and the Wolffian and Müllerian ducts undergo a

¹See Bland Sutton, *Tumors, Innocent and Malignant*, for a careful study of these cysts and the associated fistulas (London, Cassel & Co., 1894, p. 324). In this work Bland Sutton gives a mass of data regarding the different forms of cysts which it is difficult to meet with elsewhere. His classification, however, strikes me as capable of improvement.

"GLANDULAR" CYSTS

most complicated series of metamorphoses in the female, and as a result cysts are liable to form in unabsorbed remnants of the same, *i. e.*, in the residue of the Wolffian body, situated in the substance of the ovary, the head of the Wolffian duct and its side tubes (the organ of Rosenmüller), the continuation of the Wolffian duct through the broad ligament and alongside, or in the outer wall of the vagina where it opens near to the urethral orifice. This duct, when present, as not infrequently happens in the sow and cow, is known as Gärtner's duct. In connection with all of these remnants cysts may arise.





Cysts of the Wolffian body are apt to be multilocular (although they may be unilocular), to grow downward in the broad ligament, to have papillomatous vegetations, and to develop after maturity is reached.

Cysts of the free outer tubes of the *organ of Rosenmüller* (or paroophoron), which do not communicate with the Wolffian body and ovary (and are known as Kobelt's tubes), are small, rarely larger than a pea.

Those from the vertical *connecting tubes of the paroophoron* are apt to become of great size; more often they are solitary. The smaller cysts are transparent and thin walled, with lining of ciliated columnar epithe-

lium; with increase in size the walls become thicker and opaque, the epithelium flattened, if not stratified; in the largest the epithelium has disappeared, and the contents are turbid, containing cholesterin and other products of cell degeneration. These cysts are apt to show themselves between puberty and the thirtieth year.

Cysts of Gärtner's duct are to be recognized as occurring in two positions: (1) in the broad ligament to the inner side of the ovary, giving rise to certain isolated cysts of the broad ligament, in general of no great size; and (2) in the vaginal wall, there also of no great size (compared with the huge cysts of the paroophoron), and either solitary or, rarely, in series.

There has been and continues to be discussion regarding certain small cysts that may occur in the uterine muscle, lined with columnar epithelium and often associated with new-growths of the muscle in the form of "adenomyoma." Many German authorities regard these cystic tubules as remains of Gärtner's duct; other investigators, more correctly, ascribe the tubules to deep ingrowth and sequestration of the uterine mucosa. We have considered these when dealing with the uterine "adenomyomas" (p. 748).

Hydatid of Morgagni.—The term hydatid of Morgagni is by many used laxly to indicate small cysts in the outer part of the broad ligament. This is a mistake; there is but one hydatid of Morgagni, the term, used correctly, being employed for the cystic dilatation of the longest of the finbriae of the Fallopian tube. Fig. 286 indicates the relationship of the various ovarian and tubal cysts.

(f) Gysts of the Primordial Genito-urinary Ducts in the Male.—In the male the mesonephros, or Wolffian body, is represented by the paradidymis, the Wolffian tubules by the vasa. Of the latter, we recognize three groups; the main group constitutes the epididymis; the most distal give rise to the *stalked hydatid* or *hydatids* at the upper pole of the testis or globus major of the epididymis.

These are sometimes mistakenly termed hydatids of Morgagni; it will be seen that they are not the homologues of the cysts of this name in the female.

The proximal remnants in connection with the lower end of the testis form a series of blind tubular remnants, forming the *paradidymis*. Along with these and resembling in its relationships the Kobelt's tubes. (*i. e.*, being blind distally, but communicating proximally with the efferent tubes) is the vas aberrans. All these vestiges may be, rarely, the seat of cyst formations; to the blind tubules of the paradidymis are ascribed certain multiple cystic growths of the testis. Just as in connection with the Wolffian tubules in the female there may arise parovarian cysts, so apparently from the vasa efferentia, the homologous organs, there may arise solitary cysts, constituting what English authors describe as *encysted hydrocele* of the testis. Such "encysted hydrocele" has a totally different origin from the hydrocele of the tunica vaginalis, to be presently noted.

Of the Müllerian duct in the male only two remnants have so far been determined, namely, the two ends, and of these, one only gives rise to cysts. This is the distal or anterior end in association with the globus

"GLANDULAR" CYSTS

major of the epididymis, which may be present as a small cyst, the sessile hydatid. The lower end fusing with that of the other side forms the sinus prostaticus, lying between the openings of the seminal ducts, the homologue of the female uterus and vagina.

(g) Congenital Cysts Due to Imperfect or Arrested Development of Glandular Organs.—The kidney and the liver supply the best examples of this order. Occasionally birth is rendered difficult by the huge size of the feetal kidneys, or following birth, death ensues quickly from obstruction to the action of the diaphragm by the greatly enlarged kidneys. At other times the condition is less extreme and is partial, and the individual may reach maturity. In such cases the kidneys may be found to be converted into a mass of cysts and dilated channels, varying in size



Congenital cysts of the kidney from a newborn child dying shortly after birth. Both organs, greatly enlarged, consisted of an aggiomeration of elongated cysts as shown here. (From the collection of the Royal Victoria Hospital, Zeiss objective DD without occular.)

from those just visible to the naked eye to others the size of a marble, or in the adult the size of a plum or larger. In these cases the renal tubules or the larger proportion of them are found not to communicate with the pelvis. A layer of dense fibrous tissue occupies the region of the calices. Above this the tubules are dilated and cystic.

There has been a long-continued debate as to the cause of such congenital cystic kidneys. Virchow regarded them as due to foctal inflammation affecting the medulla and leading to constriction and atrophy of the terminal portions of the collecting tubules. Others have regarded the condition as adenomatous. The consensus of opinion at the present moment is that we deal with a condition of arrested development. As to the exact nature of that arrest observers continue to be

CYSTS

widely at variance, although the more recent exact studies of Schreiner and Huber, made with the aid of reconstructions of serial sections of the fœtal kidney, by Born's wax-plate method as again of Herring, would seem definitely to have settled the matter.

The original view of Remak and Kölliker, accepted by a long list of later observers, was that the whole of the renal tubules arise as evaginations and branches from the Wolffian duct. According to this view congenital cysts of the kidney can only be brought about by obstruction of the tubules after their formation. Virchow, as already noted, taking this view ascribed the obstruction to inflammation of the medulla.

So long ago as 1865 Kupfer recognized a separate origin for Bowman's capsules and the convoluted (secretory) tubules, and his researches have been confirmed by a succession of embryologists of repute (Balfour, Adam Sedgwick, Wiedersheim, etc.). The above-noted researches of Schreiner' and Huber' prove:

 That the first indication of the fortal kidney is an evagination from the Wolffian duct near the cloaca. This clongates, and becomes bulbous at its end. The stalk is the future ureter, the ampulla expands longitudinally and is the anlage of the future pelvis and collecting tubules.

 Running alongside the Wolffian duct is a cell mass of mesenchymatous origin, the nephrogenic tissue. At the upper end this gives independent origin to the Wolffian tubes. Below, the renal ampulla penetrates into the mass.

3. The elongated ampulla (pelvis) gives rise to pairs of evaginations or branches, which proceed to dichotomise, developing the future collecting tubules.

4. The swollen or ampullary end of each branch becomes surrounded by a segregated mass of the nephrogenic tissue, and *in the inner zone of this tissue* there now form cell collections developing into vesicles—*renat vesicles*.

5. These vesieles elongate and fuse with the collecting tubules, so that their lumen opens into that of the ampullary ends of the collecting tubules. Elongating still farther and becoming S-shaped, the distal end by invagination gives rise to the Bowman's capsule with its contained glomerulus; the other portions of the S give origin to the first convoluted tubule, the loop of Henle and the second convoluted tubule; in fact, to all the secretory portions of the renal tubule.

Solution of the medulla in the later months of factal life might bring about obstruction, the simpler explanation of the congenital cystic kidney is that of a want of relationship or fusion between the renal vesicles and the outgrowths from the Wolffian duct. If the nephrogenic tissue proceeded to form renal vesicles without the renal ampulla proceeding to dichotomise and form the collecting tubes we should expect the case of universal cystic degeneration of the kidney. Now we encounter kidneys of this order—kidneys with no pelvis, and

¹ Zeitschr. f. Wiss. Zool., 71:1902.

² Amer. Jour. of Anat., 4: 1905, Supplement 1.

again, kidneys with a pelvis, but no signs of a medulla or of collecting tubules. We could expect also cases exhibiting in general a well-formed medulla and collecting tubules, but with these scattered cysts, due to local independent development of the mesenchymal renal vesicles without fusion with the collecting tubules, and, again, as a matter of fact we encounter these cases.

In the kidney in the newborn it is not uncommon to encounter what are evidently isolated imperfectly developed Malpighian bodies situated immediately beneath the cortex. These must be regarded as renal vesicles of latest development which have not as yet gained connection with a collecting tubule; in congenital syphilis causing, as it does, arrested development, these are apt to be relatively abundant. I cannot, however, accept the view propounded by Ribbert⁴ that the cysts in the congenital cystic kidney are developed purely from the glomerular bodies. The renal vesicles are not merely the glomerular anlagen, but give rise to the whole secretory tubules, and the cysts in certain cases are clearly of tubular type both as regards their form and their epithelial lining.

Aschoff would proceed farther and give the like origin to all the isolated "retention cysts" encountered in later life. Here we cannot entirely follow him. It is true that such cysts form most often immediately beneath the capsule, in the region, that is, where we find latent and rudimentary renal vesicles. We do not deny that some, perhaps a considerable proportion of isolated cysts, are of this nature, but the renal tubules are certainly capable of dilatation as a result of obstruction. We may, indeed, have acute cystic dilatation of tubules in acute parenchymatous nephritis as a result of blockage of the narrow descending loop of Henle by the shed cells of the first convoluted tubules. It is rational to believe that interstitial fibrosis, by contraction and obliteration of lumen of tubules, may lead to similar obstructive cyst formation. It is striking how frequently these scattered cysts are associated with chronic interstitial nephritis.

In the liver we more rarely encounter similar congenital cysts. In those cases we have studied there has always been a marked degree of fibrosis around the bile ducts in the immediate neighborhood of such cysts. A diffuse enlargement of the bile ducts, and even of the bile capillaries throughout the organ, may be associated with congenital stenosis or obstruction of the common bile duct. We regard them all, then, as hepatic bile cysts of obstructive origin.

Congenital Cystic Diathesis.—Still more rarely we encounter cases of what has been termed the *cystic diathesis* with multiple cyst formation in both the kidney and liver, or as in a case studied by us in kidney, liver, and pancreas.

In our case the kidneys were of the typical congenital cystic type, as seen in the adult, *i. e.*, there was a pelvis and in certain areas wellformed collecting tubules, although large portions of both kidneys were converted into dense congeneries of cysts with no sign of collecting

¹ Verhand, d. deutsch, path. Gesellsch., 2:1900.

CYSTS

tubules or pyramids. The bile cysts were relatively few and small, and there was marked fibrosis around the bile ducts in their neighborhood; they were obstructive cysts. The distal half of the pancreas was alone affected, but this to an extreme degree all the tissue being replaced by large cysts. The rest of the organ was normal and the cystic transformation began immediately beyond a distinct groove on the surface of the organ formed by the tense band of the mesenteric vessels. These clearly had pressed upon the organ and compressed, and to some extent obstructed itsduct. We had clearly to deal with obstructive retention cysts.

Here clearly the occurrence of multiple cystic disturbances was a coincidence, or, at most, the diathesis was of the nature that obstructions, which in the normal individual would be overcome, in this individual were followed by dilatation and giving way and atrophy of the surrounding tissues, so that cyst formation resulted more easily than in normal individuals. A study of the recorded cases leads me to regard them as of the same nature.

11. Glandular Cysts of Postnatal Origin.-1. Originating in Tubular Glands through Obstruction of their Ducts.-Retention cysts of this order are widely distributed. The ducts may become plugged and obstructed by mucus, calculi, etc.; may be stenosed through the pressure of surrounding fibrous tissue, the result of a chronic productive inflammation, or compressed by the pressure of new-growths from without, etc. Cysts of this order lined by characteristic columnar or cubical glandular epithelium may develop in any tubular gland. The examples are very numerous: Ranula of the floor of the mouth through obstruction of the duct of the sublingual glands, or of the glandula incisiva situated farther forward in the floor of the mouth; salivary cysts through blocking of a salivary duct; mucous cysts, small and multiple, of the intestinal mucosa through blocking of the crypts; pancreatic cysts (Ranula pancreatica) of like origin, cysts of the mucous glands of the epiglottis, trachea, Cowper's glands, glands of Bartholin; bile cysts of the liver; the small, solitary, or sparse retention cysts of the kidney (see above), of the glands of the uterine cervix (Ovula Nabothi), of the lacrimal gland (*Dacryops*), of the ducts of the mammary gland (*Galactocele*), etc.; wens, atheromatous or sebaceous cysts (arising from the sebaceous glands in the hair follicles). In this group must also be included conditions of hollow organs lined with a glandular epithelium, as, for example, hydrops cystidis fellew or cystic dilatation of the gall-bladder due to obstruction of the cystic duct, and appendicular cyst due to stenosis of the appendix vermiformis with dilatation above the region of obliteration. Some of the most remarkable examples of cyst formation are encountered in connection with the Fallopian tubes, when, through inflammation, they become closed at either end (hydrosalpinx). When through laceration in parturition and subsequent cicatrization the cervix uteri becomes occluded, there develops the condition of distension of the uterus with retained discharges known as hydrometra. These latter cases may, with propriety, be relegated to the group of composite cysts to be presently noted.

CYSTADENOMAS

2. Originating in Ductless Glands.—The thyroid and the pituitary body are both formed, as regards their main constituent, of closed vesicles. These under normal conditions become, many of them, excessively distended with secretion, and thus may present cysts visible to the naked eye. In the thyroid, which is more frequently affected, such enlargement constitutes one form of cystic goitre. To other forms of thyroid cysts we shall refer later.

Among the cysts of the ductless glands must be included *cysts of the corpora lulea* and *Graafian follicular cysts* of the ovary. Both are somewhat aberrant in their mode of development. A further note regarding the former is given on p. 864. The Graafian follicles are of mesothelial origin, originating from gland-like crypts of the germinal epithelium. The normal follicle has a fluid centre surrounded by

several cell layers. The function of these specific cells would appear to be that of nourishing the developing oocyte situated within the discus proligerus. At times they actively secrete fluid, and small follicular cysts have been found still containing an ovum. More commonly the ovum has undergone disintegration, and cysts, the size of a pea, or larger, persist in the ovary; these are frequently multiple.

111. Glandular Cysts of Neoplastic Origin: Cystadenomas.—

All adenomas of tubular glands or reproducing vesicular cell groups may, through continued production of (abnormal) secretion, give



Section from a case of multilocular ovarian cyst, showing early papillomatous ingrowths into certain of the cysts. (Ribbert.)

rise to multiple cystic growths. There are adenomas of certain organs which have a special tendency toward this cystic modification. First and foremost must be mentioned those of the ovary, or, as some would hold, of the Wolffian body. This *cystadenoma papilliferum* of the ovary, or multilocular ovarian cyst, may attain a great size and be composed of innumerable loculi or individual cysts, some as large as a child's head, others searcely visible—all of them are lined by a columnar epithelium. As indicated by the first title given, one group of these cysts from the ovarian region has, in addition, a pronounced tendency toward the development of intracystic papillomatous growths.

The breast is another region in which adenomata are prone to become cystic, although here it has to be pointed out that many of the conditions termed cystadenoma are not true neoplasms, but are primarily, at least, retention cysts of the milk ducts caused by chronic inflammation, or are involution cysts, from stenosis of the ducts and subsequent cyst or mation accompanying retrogressive changes and atrophy of the gland

CYSTS

substance. It is from the walls of such cysts that there is a tendency for papillomatous and even carcinomatous intracystic masses to develop.

Cysts of other nature may develop in tumors—lymph cysts (p. 861) and necrotic cysts (p. 864). Relatively small cysts of glandular type may occur in one form of tumor not of the adenomatous or lepidic type, namely, in gliomas. To these we have already referred (p. 757).

IV. Endothelial Cysts.—The cases of endothelial or serous cysts of congenital origin are relatively few in number; they may fittingly be considered along with those of postnatal acquirement. The characteristic examples are serous cysts—sacs distended with serous fluid or lymph. Such sacs are lined with endothelium, either of some cut-off portion of a serous cavity, more particularly of the peritoneum, or of the lymphatic system. Examples of the former are scrotal hydrocele, and cysts of the canal of Nuck; of the latter, bursal cysts, "ganglia" (cysts formed by the cutting off of hernial protrusions of the synovial lining of tendon sheaths), many mesenteric and subpleural cysts, lymph cysts in association with macroglossia, and other conditions of congenitally obstructed lymph channels.

Of these serous cysts of congenital origin, some of the most remarkable are those known as *hygromas*. These lymph cysts of congenital origin may occur in various situations, but the largest and most remarkable examples occur in the neck—*hygroma colli*.

Here huge and progressive collections of lymph cysts may develop, either a few large, or a sponge-like congeries of small cysts, and these ramify, not merely in the fascice between the neck muscles, but in the muscles themselves, the parotid and salivary and other tissues of this region, extending both upward and downward (into the thoracic cavity).

It is doubtful whether these should be regarded as simple serous cysts; some are of this nature, but many exhibit a very cellular stroma, and, as pointed out by Boyce,¹ there is evidence not merely of connective tissue, but of endothelial proliferation: ramifying columns of endothelial cell masses are to be made out, which, judging from transitional stages observed, ultimately become cystic. In other words we deal with definite hymphangio-endotheliomas (Borrmann²).

Other lymph cysts deserving of note are those which occur in the choroid of the eye, the chylangiomas of the intestinal wall (dilated lymphatics filled with inspissated milky chyle), and the serous cysts which may complicate tumors, notably uterine fibroids.

It is obvious, that since the tunica vaginalis is normally present as a closed sac, without accumulation of fluid, the accumulation of fluid in the same to form a hydrocele can only be brought about by modified activities of the lining membrane; there must, that is, be either increased secretion into the sac, or obstruction to the normal diffusion of fluid out of the sac, or both. It is probable that we have to deal with both. We cannot explain the first stage in the formation of a hydrocele (before

¹ Pathological Histology, p. 192.

² Ziegler's Beiträge, 40: 1906: 372.

EPENDYMAL CYSTS, NEURAL CYSTS

distension of the sac shows itself) without assuming increased secretion and this secretion must be governed by the state of the endothelial lining; in the developed hydrocele there is definite thickening of the tunica forming the sac, and this must hinder diffusion outward.

The same considerations govern bursal and lymph cysts in general; we have to recognize both obstruction to outflow and discharge, and active secretion. To form a cyst the pressure within the sac must be greater than that of the lymph in the immediately surrounding tissues, and this demands the latter condition. Such bursal cysts may be of intra-uterine or postnatal acquirement. The conditions leading to their development are increased pressure on a part, of an intermittent type, with movement of the loose connective tissue over some more rigid prominence, with rupture or separation of the strands of the connective tissue, resulting in a low form of inflammation, with the accumulation of serous fluid. The cavity thus formed, at first irregular, through proliferation of the endothelium of the lymph spaces involved becomes lined with endothelium, becomes also smooth-walled and cystic.

V. **Ependymal Cysts, Neural Cysts.**—The ependyma lining the hemispheres and the spinal canal is embryologically as an epiblastic tissue more nearly related to the "glandular" lining membranes, although functionally it is more of the nature of an endothelium, such as that lining the serous cavities; the cysts developed in connection with it may thus be treated as a special class.

Through stenosis, whether from imperfect development or intrauterine inflammation, there may be brought about localized closure of some portion of the spinal canal, or, again, of the canals of communication between that canal and the external lymphatics. Either condition results in accumulation of the cerebrospinal fluid and cystic dilatation of the cerebral ventricles or of the spinal canal. In this way result the conditions of hydrocephalus internus, hydrocele of the fourth ventricle, cysts of the spinal canal (syringomyelocele) (see p. 265).

These conditions must be distinguished from the various forms of meningoccle, serous or endothelial cystic states, partial or complete, involving the meninges. The ordinary hydrocephalus involves the lateral and third ventricles, the obstruction occurring in the iter. Hydrocele of the fourth ventricle, a condition first described by Virchow,¹ is a much rarer condition, brought about by obstruction of the lateral recesses of the fourth ventricle. These passages contain the choroid plexus of the fourth ventricle, and form the means of communication between that cavity and the subarachnoid space at the base of the flocculus; obliteration of one of these recesses, or of both, lead to unilateral or bilateral bulging and cyst formation at the base of the brain, with pressure upon and paralysis of the facial, auditory, and, it may be, of the glossopharyngeal and vagus nerves, all of whose root filaments lie in the neighborhood of the cyst. The condition, as in a case of my colleague, Dr. Bell, may be associated with a moderate grade of hydrocephaly.

¹ Die Krankhaften Geschwülste, 1:183.

CYSTS

VI. Epithelial Cysts.—These form a large group. The congenital "sequestration dermoids," and acquired implantation cysts of the skin, are more properly considered among the composite cysts, inasmuch as the fluid filling the cyst is in general the product of sweat and sebaceous glands, and not derived directly from the squamous epithelium; and there they will be discussed. Here it may be noted that the simplest form of implantation cyst, such as may be found beneath the skin of the fingers or palm in sewing-women and those performing rough manual labor, may, under the microscope, exhibit a simple squamous epithelium without glands, and nevertheless possess pultaceous or atheromatous fluid contents. Such cannot be regarded as true secretion; the fluid is the product of the disintegration and autolysis of the more centrally situated cells. We refer to cysts of this order in this place to call attention to the fact that their walls are in the main composed of squamous epithelium. Occasionally, as in the orbital tissue, simple cysts are found lined by stratified epithelium without glands which, nevertheless, have clear fluid contents.

VII. **Composite Cysts.**—Under this category are to be included those cysts whose walls are formed of more than one form of epithelium, and in which the secretion of the cystic fluid is from glands discharging into the main cavity. The sequestration cysts above mentioned are of this character, and closely allied are the teratomatous cysts, such as the ovarian dermoids; hydronephrosis of the kidney and its pelvis is a typical example.

Sequestration cysts are due to the inclusion and detachment of portions of the true skin in the region of the fissural lines of the body below the line of fusion of the two sides of the dorsal groove, of the thoracic abdominal cleft, the facial clefts, etc. In all these regions cysts may be encountered lined with squamous epithelium, and either containing glairy fluid (from sudoriparous glands) or pultaceous fatty fluid (from sebaceous glands); or, in addition to these, hairs both loose and attached.

Such cysts may not be merely superficial, but may be situated relatively deep in the tissues. It must be remembered, as pointed out by Bland Sutton, that the bony vault of the cranium is of relatively late development; originally the skin over the head was in contact with the dura mater; thus, through abnormal infolding and sequestration a skin cyst may eventually form attached to the dura mater and beneath the cranial vault. The same is true of the anterior cleft of the body and the sternum; sequestration dermoids may develop in the mediastinum beneath the sternum.

Postnatal *implantation cysts* may be of similar structure; through trauma, a portion, large or small, of the surface epithelium may become lodged in the underlying tissues; may become grafted there and proceed to grow. The active growth of a squamous epithelium, it will be recalled, occurs in the lower Malpighian layer; the inevitable tendency, therefore, is for the growing edge of this outer surface to curve inward until a globular mass is formed (Fig. 289).

HEMORRHAGIC CYSTS

If the original graft have sebaceous or sudoriparous glands associated with it, the secretion from these distends the cavity and a cyst results.

We have dealt with teratomatous cysts in discussing the teratomata (p. 659).

The subject of hydronephrosis will be dealt with in more detail in the second volume. Here we would only call to mind that hydronephrosis is brought about by obstruction, congenital, or more often acquired, of the urinary passages, at any point between the prepuce (phimosis) and the pelvic origin of the ureter. Such obstruction leads to distension of the passage above the point of closure, and that distension tells especially upon the pelvis of the kidney. There are cases of hydronephrosis in which the pelvis alone is affected, the kidney-substance not recognizably involved. Most often intrapelvic pressure leads to atrophy of the pyramids, and this atrophy is progressive until the kidney may be represented by a large sac, a foot or more long, and of proportional girth, the kidney substance proper being represented by a thin layer of tissue a millimeter, or little more, in diameter.



Diagram to illustrate mode of formation of an implantation cyst of skin: I shows a fragment of skin depressed into the underlying tissues: the actively growing cells of that fragment are upon its under aspect, at a (the padisode layer of the rete Malpighii): the stratified flattened cells of the epidermis (at b) have lost the power of growth; 2 and 3 show the continued growth of the cells of the rete Malpighii, which from the want of growth of the cells at b must come to surround those cells, and form as at 4 a solid, and later, as at δ_a hollow sphere.

In this connection may be noted the cysts of the jaw, both simple and multilocular, due to aberrant development of the teeth sacs and enamel germ remains. These may contain clear or atheromatous fluid, may be lined with squamous or merely a single layered cubical epithelium and may or may not contain, projecting into them, one or more teeth; when teeth are present we have the true *dentigerous cust*.

II. HEMORRHAGIC CYSTS.

Extensive hemorrhages into the substance of sundry organs may result, not in the ultimate absorption of the exuded fluid, but in cyst formation. The hemorrhage leads to the destruction of the tissue of the infiltrated area; eventually a capsule is formed around the exuded blood, but while this is proceeding, through the combined agencies of leukocytes and autolysis, the bodies of the corpuscles and the cell debris

undergo removal, as does also the diffused and altered hemoglobin; so that, after a few weeks, the cyst is found to contain a thin blood-stained fluid, and, eventually, all the pigment being removed, the contents come to be a clear serous fluid. The last indication of the hemorrhagic origin of such a cyst is the presence of modified blood pigment in and around the fibrous capsule. The organs in which such hemorrhagic cysts are specially liable to be found are the brain (substance of the hemispheres, base, internal capsule, and pons), ovary (corpora lutea), the goitrous thyroid, scalp of the newborn and children (*Cephalhematoma*), pinne of ears (in football players and lunatics—*Othematoma*).

Bradley,¹ following Rokitansky, has pointed out that the large cysts with serous contents found in the thyroid, from their structure, and the various stages found, can only be regarded as the results of hemorrhage into a nodular goitre.


The cephalhematomas are interesting in that there is a tendency toward the development of a ring or margin of true bone at their base, proceeding from the pericranium where it is raised from the bone, the hemorrhage, in these cases at least, being situated beneath the pericranium. A crateriform, osseous growth upon the skull is the result. Bland Sutton figures an extreme example of this condition found in a monkey.

The corpus luteum of pregnancy occupies an intermediate position; in its earliest stage it is definitely a blood cyst, but soon, without further hemorrhage, the volume of its contents increases, the contents with this becoming paler. The recent researches of Prenant, Born, Jolly and Marshall² denote that the corpus luteum affords an internal secretion, and indicate that the lutein cells, far from being degenerative, play a very active part in connection with the secretion of diffusible substances which influence the nuccus membrane of the uterus and the economy in general (see p. 362).

III. NECROTIC CYSTS.

These, in mode of origin, are closely allied to the hemorrhagic cysts. When there has been an extensive necrosis of tissue of a non-infective type, as in large infarcts, complete absorption of the dead matter and replacement by cicatricial tissue does not occur, but in its place the central area of such a mass undergoes autolytic liquefaction, the soluble products diffuse out and lymph diffuses in; around the space a cicatricial capsule becomes formed, and thus a cyst results. A not uncommon seat of such necrotic cysts is the centre of large cancerous nodules. Through the centrifugal growth of the tumors the central cells become cut off from due nourishment, and it is held that those cells of glandular type afford relatively abundant autolytic enzymes; the nature of the

¹ Journal of Experimental Medicine, 1:1896:401.
 ² Philosophical Trans. Royal Soc. Biol., 198:1906.



Eosmophilia (Trichiniasis).

Copy of an actual next containing three normal polynacian cosinophiles, one broken cosinophile, one polynacicar contractific, and one lymphocyta



PARASITIC CYSTS

growth hinders due cicatrization and replacement of the dead matter by granulation tissue. Thus, where under these conditions the growth does not fall in and become umbilicated, central cyst formation results.

IV. PARASITIC CYSTS.

Throughout the animal kingdom various metazoan parasites of the order of vermes, and of these preëminently the *Tamiada*, pass one cycle of their existence in an encysted stage within the tissues of their host, and the cysts formed by them may assume relatively large dimensions. In man the one parasite that forms cysts of large size is the Taenia echinococcus. Smaller cysts, just visible to the naked eye, are formed by the Trichina spiralis. These most commonly are situated within muscles, and are of an elongated oval shape, having the larval trichina coiled within them. The echinococcus cysts, on the other hand, may be as large as an orange, and yet larger, and may be variously distributed, being commonly known as—

Hydatid Cysts.—The liver is the commonest seat, although they may occur in very many organs and tissues—in the brain, spinal canal, eye, kidney, heart wall or cavity, lungs, mammary gland, bone, omentum, peritoneum, etc. They may be either single or multiple. Microscopically they are characterized by possessing an external capsule of laminated hyaline tissue, within which is a cellular and granular layer, from which at first there projects a single club-like head, with water-vascular system, and a circle of characteristic hooklets. As the cyst grows secondary heads with hooklets and surrounding cysts bud off from the wall internally, and these cysts are apt to become free in the hydatid fluid. In this way the single original hydatid may become filled with very numerous daughter cysts. Examination of the fluid tapped or expressed from such a tumor reveals the presence of hooklets in the sediment. Occasionally, certain of the cysts are sterile, *i. e.*, contain no heads and no hooklets.

More rarely in the liver, and the same is usually in bone, the daughter cysts, instead of being developed from the inner wall of the parent cyst, project outward, in which case there is developed a multilocular cystic formation, which, in the old days, was mistaken for a cystadenoma. The character of the hyaline wall, apart from the hooklets in the contents, suffices to determine the true nature of this more diffuse cystic state. It should be added that, outside the cyst wall of the hydatid proper, there is developed an adventitial fibrous wall, at times of considerable thickness, through the irritation and productive inflammation set up in the tissues of the host.

CHAPTER XXV

THE REGRESSIVE TISSUE CHANGES.

NORMAL HISTOLYSIS AND CYTOLYSIS.

JUST as, for a proper study of the progressive tissue changes, it was found advisable to consider first the phenomena of normal growth, so, as an introduction to the consideration of the regressive tissue changes, it is well to take first into account the natural decay of tissue elements. Although we give scant attention to the fact, it is almost needless to say that, from the embryonic period of life onward, individual cells and indeed individual tissues undergo a process of natural displacement and decay. Tissues and organs in the embryo, representatives of ancestral structures, make their appearance and then disappear; the thymus, functional in early life, attains its maximum development during the first two years of extra-uterine existence, then slowly shrinks and undergoes absorption; the lymph glands are largest during the early years, after which they exhibit a progressive diminution in size; as the milk teeth grow and emerge, they cause the atrophy of the tissue immediately surrounding them; after a few years their roots, in turn, undergo absorption before the growing permanent set of teeth; after each pregnancy the uterine tissue undergoes involution; that is to say, its cells which had undergone hypertrophy or had become hypertrophied, now, to a large extent, atrophy; the ovaries, again, atrophy at the menopause, and, in general, with advancing years, one and all of the tissues of the body give evidence of shrinkage and loss of substance; that is, loss of component cells. And while any one tissue-with the possible exception of the nervous system-may, for long years, appear to be unchanged, we know that this is far from being the case. Red corpuscles and leukocytes have a life period of, at most, a few weeks, they disintegrate or are eaten up by other cells and younger corpuscles take their place. The same is true of the cells forming the epidermis and hair, although there the life period is longer. The same again would appear to be true of gland cells, and, even so stable and solid a structure as bone, when carefully studied, is seen to be undergoing constant change, the cells of the older trabeculæ dying, the bone substance becoming dissolved, and new cells and new bone substance being laid down in their place.

We know relatively little concerning these processes of normal cytolysis and histolysis, although our knowledge has, of late, undergone a material increase. But, leaving out of account for the moment the condition of simple atrophy, so called—conditions, that is, in which

NORMAL HISTOLYSIS AND CYTOLYSIS

we can recognize more or less clearly some departure from the normal in the direction of either reduced nutrition or modified function as the cause of the regressive change-and, considering only what could be regarded as strictly normal cell and tissue destruction, even here we become impressed with the fact that not one but several processes are at work. These processes group themselves into two orders, namely: (1) changes occurring in the cells themselves, and (2) processes acting from without upon the moribund cells whereby they are destroyed and removed. Taking first the changes occurring in the tissues and the cells themselves, we have to admit that under certain conditions, nothing very obvious is to be made out in the cells that are about to die or be destroyed. We cannot with certainty state that a given leukocyte, seen in the tissue or in the blood stream, was at the height of its activity when fixed and stained or was about to disintegrate. We are only too apt to forget that our histological studies are made upon dead material and that we must be most cautious in our interpretation of what we observe in the cells which have been fixed-and killed-by ourselves as compared with cells which have died without our intervention. So, also, while it is true that, examining a given smear of the blood we may observe certain red corpuscles which are paler and more cedematous looking, we perhaps rightly judge that these cells are approaching the limit of their existence, yet, when we examine the cells of the spleen pulp, which, acting as phagocytes, have removed what are presumably damaged or moribund erythrocytes from the circulation, we cannot state confidently that the cells so removed present any marked difference when first ingested from those that are left behind.

Often, however, we can make out that the cells composing a given tissue are becoming older. In senile atrophy, for example, which, as already pointed out, is truly a physiological process, we can see that the individual cell elements undergo a definite amount of shrinkage. They become smaller, and the finer details of cell structure tend to disappear. The transverse striation of muscle fibres, for example, is not so sharply marked. This is the true normal, simple atrophy.

Nevertheless, although the process be a normal one, we are forced to recognize that, as a cell fails and shrinks, so does it become less and less able to carry out its metabolic processes. Substances absorted by it may thus not be properly broken down and converted, and the products of cellular activity will not be properly excreted, while, again, the products of the gradual breaking down of the cell substance, if they are not of a soluble nature, are apt to become heaped up within the cell. We should therefore expect to find—and, as a matter of fact, we do find that, under what are wholly normal conditions, cells undergoing atrophy and about to be disintegrated may exhibit much more than mere simple shrinkage. And, thus, under normal conditions, we may find the prototypes of disturbances which are apt to affect the cells in pathological states. Many of the so-called degenerative processes are liable to show themselves in this physiological or normal histolysis. The most familiar example of this is to be seen in senile atrophy of the heart-muscle fibres,

There is, perhaps, no more common histological finding in the tissues of old people than the so-called brown atrophy of the heart muscle. In this condition, hand in hand with the shrinkage in size of the fibre, there is to be recognized in the cell, at either pole of its nucleus, a deposit of fine reddish-brown granules. This deposit we suppose is the product of breaking down of the myohemoglobin which normally is diffused through the muscle fibre. (See p. 962 for discussion of this subject.) As the cell substance breaks down, this is supposed to undergo conversion in such a way that an insoluble residue is left behind.

Perhaps the best illustration of such changes is afforded in connection with the involution of the uterus after parturition. During pregnancy the muscle cells here have undergone an extreme hypertrophy. Now, during the week following delivery, the shrinkage is equally remarkable. From being on an average (according to Sänger¹) 208.7 µ long, they become reduced to an average length of 24 µ. The shrinkage is, by no Accompanying it, the cells have a more *cloudy* means, everything, appearance and some, but not all, show along their length fine, refractile globules which, treated with osmic acid, turn black. There is thus a condition of what is termed *fatty degeneration* (p. 909), which in some cells may be extreme. The nuclei also become diminished in size. Nor are the muscle cells the only portions of the tissue to show change. In between the more internal bands of fibres the connective-tissue cells become the seat of a deposit of fat, in fact, become converted temporarily into fat cells, losing this fat again by the fortieth day. The vessels also become markedly altered; many of the smaller ones become completely obliterated and disappear, the larger vessels show a characteristic swelling and overgrowth of the intima leading to considerable diminution of the lumen, with coincident hvaline or vitreous change in the internal elastic lamina of the arteries (see p. 896), atrophy of the media and eventual building up of a smaller new vessel within the dilated old one.

New conceptions with regard to forces at work leading to the physiological atrophy of cells as again to pathological atrophy, have been afforded by the studies originated by Bordet upon the cytolysis of cells derived from a different species of animal (p. 532). We can go still farther. We have observations showing that when individuals of one species only are employed, we can, by successive inoculations, cause the blood serum of one animal to destroy certain cells of another animal of the same species. If this be so, may it not be that, throughout life, a due development of the various tissues and their keeping within bounds, so that one tissue does not invade the other beyond a certain point, may be due to the development of *heterolysius* (p. 372), which developed by cells of one order destroy those of another order? So that, for example, the healthy cell prevents the excessive growth of cells of another nature in its immediate neighborhood by the discharge or excretion of these heterolysins. If this be so it is conceivable that when an

¹ Festschr, f. E. Wagner, Leipsic, 1888,

DISUSE ATROPHY

individual cell through functional activity or through other causes becomes worn out, it is no longer able to sufficiently protect itself by its heterolytic substances, and, as a result, it is killed and undergoes dissolution under the action of heterolysins secreted by other orders of cells. The French school, and more particularly Metchnikoff, have propounded these views. However attractive they may be, it has to be confessed that much more work has yet to be done before they can be regarded as established.

Leaving out of account, for the time being, inherited or acquired cell weakness and the conditions which lead to abiotrophy (p. 876), we are forced to recognize that the two prime factors which tell upon the health of the individual cells are (as we have already pointed out in the chapters upon Overgrowth and Progressive Tissue Changes) nutrition and the performance of function. If either of these be seriously disturbed, we are liable to have cell failure and the development of atrophic states.

Theoretically, we can imagine a series of disturbances which will lead to simple atrophy of the cells. The nutrition of a part not being primarily affected, if there be loss or arrest of function, the cells will undergo atrophy from disuse. Or, again, if the work of the cell be excessive, exhaustion of that cell will set in, the cell substance will be broken down more rapidly than it is formed, and the cell as a whole will shrink. Again, if primarily there be no disturbance of function but inadequate nutrition, then the cell substance cannot be built up at the same rate as it is being broken down and shrinkage inevitably results. Or, lastly, it is possible that, if there be excessive nutrition, the work of the cell may be hampered by overabundant material, so that, again, there is a relative disuse of the cell and, with that, a tendency, if not primarily to undergo atrophy, eventually to do so. But in general it is clear that when one of this group of disturbances shows itself, then, as a secondary result, the other comes into action and favors the production of cell shrinkage and general atrophy. For example, when, through section of its nerve, a given muscle is rendered incapable of function, or, again, when, without section of the nerve, the muscle is given enforced rest, as, for example, when a healthy limb is cased in plaster bandages, then, accompanying this arrest of function, the vessels contract and there is diminished nutrition, and this, in its turn, must favor the rapid development of the atrophic state. While, where the nutrition is diminished, the functional using up of the cell substance (we use this term in its broadest sense) and the inadequate supply of material to replenish the loss, render the cell less and less capable of function, so bringing it to rest.

Now, as a matter of fact, it would seem that all these cases present themselves in practice as leading to atrophic conditions of the tissues. We thus recognize: (1) Disuse atrophy. (2) Atrophy due to excessive function. (3) Atrophy due to lack of nourishment. (4) Atrophy due to excess of nourishment.

1. **Disuse Atrophy.**—Cases in which atrophy follows primary arrest of function frequently manifest themselves. We have already mentioned

one, namely, enforced rest of muscular tissues. In such enforced rest, in a very few days, there is a noticeable shrinkage in the size of a muscle and the girth, for example, of a limb. The shrinkage, in fact, is remarkable. There is not, in the earlier stage, any diminution in the number of muscle fibres; the individual fibres diminish in size; they show the phenomena, that is, of simple atrophy. The same is the case when a nerve going to a muscle becomes divided or destroyed. As a general rule, it may be laid down, not only for muscular tissues, but also for glandular organs, and even for structures so solid as bone, that lack of exercise is shortly followed by atrophy.

Some of the most interesting examples of atrophy from disuse have been observed in connection with the nervous system. It used to be taught that when a nerve was divided there resulted purely what is known as Wallerian degeneration: that only that portion of the nerve away from its trophic centre became degenerated. But, for some years past, with the development of more careful observation and of more delicate technique, this has been proved not-to be the case. So that now it may be laid down very definitely that, when by destruction of the main axon a neuron can no longer perform its function of conveying impulses of a definite order, that loss of function is followed by a very definite, if in some cases a slowly developed, atrophy. So long ago as 1874 von Gudden' pointed out that section of the optic nerve led to shrinkage and atrophy of the external geniculate body. The importance of this observation for long failed to be realized. Only in 1894 did Nissl, by his special method of staining, show that very definite alterations could be recognized in nerve cells soon after solution of continuity of their axons. Within twenty-four hours of excising part of the facial nerve in the rabbit, he obtained recognizable changes in the Nissl bodies of the cells of the seventh nucleus with rarefaction of these bodies, and a faintly granular change. In from a week to a fortnight the change was very much more advanced. Frey, in 1887, next pointed out the cellulipetal degeneration of the proximal end of a divided nerve. This is characterized by diminution of the caliber of the nerve fibres and shrinkage in size of the nerve cells. He showed further that if the rabbit's eve be extirpated, there is a considerable diminution in the size of the lateral geniculate body through disappearance, not of its nerve cells, but of the gelatinous substance between those cells which represents the terminal ramifications of the optic fibres. If, on the other hand, the visual area of the cerebral cortex be destroyed, the lateral geniculate body degenerates, not by disappearance of the gelatinous matter, but by atrophy of the nerve cells. He concluded, therefore, that two groups of neurons make up the optic path, one passing from the retina toward the optic centre at the base, the other from the optic centre at the base to the cortex. The true disuse atrophy is that seen in the second series of experiments. The more recent papers of R. A. Fleming² and of

> ¹ Gesamm, u. hinterlass, Abhandl., Wiesbaden, ² Lancet, London, 1896; ii : 508.

ATROPHY FROM OVERWORK

Warrington' still further prove this atrophy of disuse. In section of efferent nerves the axis cylinders become thinner, but do not disappear, there being great reduction in the myelin. The ganglion cells of posterior roots show degeneration much earlier than do the multipolar cells of the anterior horn. By the seventh day it may be noted that the nuclei have diminished in size, often becoming excentric, that the Nissi's bodies have become grouped more centrally and have diminished in number; occasionally they become broken up into a dust-like cloud; while, as a result of the shrinkage of the whole cell, there is an enlarge

ment of the pericellular lymph spaces. Only after four weeks are similar changes to be observed in the cells of the anterior horn. Fleming concludes, it seems to me very reasonably, that as in health the cells of the posterior horn receive their stimuli from without, they should suffer more and suffer at an earlier date than do the motor cells whose axiscylinder processes are directed toward the region of injury. The anterior motor cells can well be stimulated and so metabolism be initiated by the stimuli proceeding down the cord; whereas, in the posterior horn, through paralysis of the muscles, the cells are completely or almost completely cut off from their usual stimuli



Disase atrophy. Section from bulk at level of the middle of the hypoghosus nucleus, from a case of multiteral cortical atrophy (infinitile polyemerphalitis), to demonstrate disuse atrophy of right pyramid (normal); p^{\prime} , left pyramid (degenerated); st, stratum interolivare, greatly diminished on left side; pc, night opyramid (normal); p^{\prime} , left pyramid (degenerated); st, stratum interolivare, greatly diminished on right side; pcp, area of posterior column which on the right side shows fibres annihy resorbed; XII, reduction and diminution in size of cells of hypoglossus nucleus. (Mingazain.)

and so are brought to a condition of much more complete disuse. In short, as Monakow² pointed out now some years ago, neurons cannot be regarded as independent and automatic units. Their maintenance in a state of health and vigor demands that they receive stimuli from without, and unless they receive these stimuli they are incapable of action. In the absence of such they undergo a disuse atrophy.

 Atrophy from Overwork.—A similar atrophy may result from excessive function. Here, again, the best examples occur in connection with the neuromuscular system, special groups of muscles and their nerve centres which are overworked in the performance of certain

¹ Journ. of Physiol., 1898.

² Arch. f. Psychiatrie, 27 : pt. 1.

movements are apt to exhibit, following upon hypertrophy which has resulted from increased work, the onset, more or less rapid, of atrophy. Neurons, or it may be the muscles, become exhausted and worn out, and, as a consequence, we have paralysis with atrophy of the involved muscles (p. 397). We have examples of this in the various professional paralyses; atrophy of the muscles of the upper arm in blacksmiths, etc.; of the muscles of the forearm controlling the finger movements in piano players, etc.

3. Atrophy from Malnutrition.—Of simple atrophy also secondary to diminished nutrition, abundant examples can be afforded. Here we leave out of account, for the moment, those cases in which nutrition is wholly cut off, resulting in the rapid death of the part supplied by a given artery. The cases we have to deal with are those in which with general diminished nutrition, without the quality of the blood being altered, there is diminished quantity passing to a part. Thus, where tumors of various natures press upon a large arterial trunk, we have shrinkage and atrophy of the part supplied by it, or where there is pressure not so much upon the arteries as upon the small capillary vessels, the same is the case; notably, continued pressure upon any viscus leads to the simple atrophy of the parts pressed upon. Even so dense a tissue as bone may thus be gradually destroyed by a more or less elastic fluid mass pressing upon it. In this way we find that, at the edge of a growing tumor, the cells of the surrounding tissue become shrunken and diminished in size, their nourishment has been interfered with, and they atrophy. We frequently come across cases in which more particularly thoracic aneurysms lead to atrophy and absorption, either of the sternum or the subcutaneous tissue (the aneurysm coming to point, sometimes even to rupture, through the skin of the chest); or, on the other hand, of the vertebral column, the bodies of the vertebræ becoming eroded.

Lack of nourishment of the body as a whole—as in prolonged fasting and hunger—leads also, it need scarce be said, to a very definite atrophy of the tissues. This has already been discussed (p. 395) but here we may note the cell disturbances. Beyond simple shrinkage there are no marked changes until the animal has lost 10 per cent. of its weight, then cloudy and granular alterations are to be seen in the cells of the larger glands—liver and kidney—and in the muscle fibres. In the liver cells, according to Statkewitsch, the glycogen disappears at a comparatively early stage, and there is a cloudy swelling, giving place later to a more coarsely granular swelling; later, again, in the outer cells of the lobules, there is extensive fatty infiltration; large fat globules distending the cells. In the mucous cells of the salivary glands there are the appearances of fatty degeneration. Nerve cells come to exhibit vacuolation.

4. Atrophy from Overnutrition.—With reference to atrophy from excessive nutrition the case is not quite so clear. Notwithstanding, the sterility of the overfed and obese is an indication of what may be termed atrophic changes, occurring in the germinal cells, while the cardiac weakness of the very obese and of those suffering from fatty

SENILE ATROPHY

infiltration of the heart and the general lack of active performance of function on the part of the sundry glands, is not wholly explained by the fatty infiltration and by the disturbance of function brought about, by the heaping up of fat either in between or actually within the individual specific cells of those tissues.

SENILE ATROPHY.

Very intimately allied to this normal histolysis is the condition of senile atrophy, which also is, strictly speaking, a physiological process a process of natural wearing out of the elements of various tissues, although in it we see—perhaps not very clearly—that something else comes into play, namely, in some individuals, it makes its appearance at an earlier period and is much more marked than it is in others. So that we have to admit a certain constitutional state either favoring or delaying the process, as, again, we have to see that the incidence of the atrophic process is apt to vary in individual tissues. For example, while, as a general rule, the brain is one of the organs which is latest to show the signs of atrophy, we come across cases in which shrinkage of the brain and the coincident mental enfeeblement are perhaps the most marked feature.

While there are these exceptions, there is a certain order in the appearance of senile atrophy of the various tissues. The first to atrophy are organs which become functionless during the course of natural life. These we have already discussed. Closely allied to them may be placed the lymphadenoid tissues. Judging from their relative development during childhood and youth, these are most active at that period. They all-lymph glands, Malpighian bodies of the spleen, and, we may add, the red bone marrow-undergo great diminution with advancing years. Next are to be placed tissues not in themselves active, but containing reserve material, e. g., fatty tissue. Nervous tissue is, as a rule, the last to show marked atrophy, while a large group composed of the connective tissues, muscles and glands, occupy an intermediate position. Of these last three groups, referring rapidly to the changes there made out, beginning with the fat cells, it may be noted that here two orders of events show themselves: either the gradual shrinkage and disappearance of the fatty contents, the cells reverting to a connectivetissue type, or as the fat in the individual cells disappears its place is taken by a serous fluid and the tissue assumes a semitransparent appearance (see p. 904). Whether this latter, so-called serous atrophy, should be regarded as a pure senile change is doubted by some. It is in those exhibiting other senile changes that we most frequently encounter it, noticeably in connection with the epicardial fat; but, when present, it is usually associated with some marantic condition; yet it is so common in the purely senile atrophy of the marrow of bones that it is, I think rightly, mentioned here. In both conditions there is a distinct tendency toward proliferation of the nucleus of the fat cell, as Flemming was the first to point out.

Passing now to the active cellular tissues, muscles, and glands, the first result of this physiological senile atrophy is diminution in the size of the individual elements of the tissue. This is very characteristic in connection with muscle and is to be noted also in glandular organs like the kidney. Only when, at a later stage, there is an actual diminution in number of the component cells of the tissue by complete shrinkage and death do we have the condition of hypoplasia. But with this we are apt to encounter other changes, namely, the presence of deposits within the cells. These deposits are well marked in those cells which normally contain pigment. The muscle cells, for example, owe their color to myohemoglobin and, as they shrink, we observe in them a deposit of insoluble reddish-brown pigment granules which appear to be derivatives from the myohemoglobin of the lost cell substance. This is particularly well marked in the muscle cells of the heart. In fact, we know no more common change in the bodies of those past middle age than the exhibition of this so-called brown atrophy of the heart muscle. Here the pigment lies heaped up in the undifferentiated cytoplasm at either pole of the nucleus. In senile atrophy of the liver, similarly, a considerable deposit of brownish pigment is met with throughout the cytoplasm of the shrunken hepatic cells.

In bones the most marked feature is a process of rarefaction whereby the individual bones become markedly lighter, and, as a result of the loss of solid osseous substance, the liability to fracture is distinctly increased. There is no great reduction in the size of the individual bones; in other words, the loss of substance is largely central, the outer periosteal layers showing little absorption; it is the internal trabeculæ and lamellæ that are, in the main, absorbed. The medullary cavity becomes greatly increased in size, the Haversian canals enlarged, the individual trabeculæ become eroded and thinned and reduced in number by the complete absorption of some, and, with this, the medulla shows pronounced change. The red cellular medulla disappears and its place is taken by fat cells, which, in the later stages, show the serous atrophy just described. Here, again, we have an example of the replacement of a lymphoid tissue by fat cells.

The senile skin, with its characteristic wrinkling, presents a series of changes affecting, not so much the epidermis as the dermis and subcutaneous tissue; loss of subcutaneous fat, diminution of vascularity by thickening of the arterial coats and obliteration of some of the smaller vessels accompanied by lessened lymph in the interstices of the tissues, all of these together with the shrinkage of the underlying and muscular tissues, favor the wrinkling. But, in addition, as has been recently pointed out, there is a distinct alteration in the elastic fibrils of the dermis.

These alterations in the elastic-tissue fibrils (accompanied by loss of elasticity and of resilience) are important factors in the senile changes occurring in the two very important tissues, the arteries and the lungs. As pointed out by Aschoff and by Foster, working under Klotz, there is a natural alteration in the elastic laminae of the aorta and vessels. As

SENILE ATROPHY

the elastic sheaths of the large arteries lose their resilience-and this most frequently occurs first at the region of greatest strain, although in some cases the loss may be general-the arterial tube, dilating under the blood pressure, is unable to return to its normal caliber; it remains permanently expanded. One of two orders of events may now take place, either (1) the expansion is permanent, and where it is generalized we find diffuse dilatation of the artery, which may be so extreme as to develop a *fusiform aneurysm*, or, where it is localized, there is formed a saccular aneurusm. In either case microscopic examination shows that all the coats of the artery in the affected region are thinned and atrophied, this atrophy going on to complete disappearance of the inner coats in some cases of the latter condition. The increased caliber of the artery leads to slowing of the flow of the blood and all the consequences of the same in the region and the tissues supplied by that artery. Such dilatation and aneurysm formation is, however, rarely the result of pure senile atrophy. As a matter of fact, in such physiological atrophy we see a correlation between the loss of elasticity of the arterial wall and the force of the heart beat. Associated senile changes in the heart lead to lessened blood pressure. It is when the loss of elasticity of the arterial wall is premature from excessive effort, syphilis, gout, and the like, and the blood pressure above the normal, that these changes are most often encountered. Or (2) we find evidences of adaptive changes. As the arterial wall gives way, there is a connective-tissue overgrowth in the intima. There is developed, in short, a condition of arteriosclerosis, a condition so common in those advancing in years as to be physiological. But the adaptation is not complete; the new tissue growth, while it eventually comes to contain elastic fibrils, does not replace the lost elasticity. On the contrary, the artery is now rendered more rigid, and this increased rigidity of the tube leads to the pulse waves being conveyed with greater force into the smaller arteries and arterioles, which, in their turn, have to undergo compensatory changes, to the detriment of the tissues supplied by them.

A like senile loss of elasticity and degeneration of elastic tissue in the lungs underlies the *emphysema* so common in old people. Here, again, the loss of resilience may be premature as the result of increased strain thrown upon this tissue, as by forced expiration, etc., and in short by anything leading to overdistension of the air sacs. But, as a normal condition, the elastic fibrils of the walls of the air sacs lose their resilience with advancing years.

As a result, if by any cause the air sacs become overdistended, they have not the inherent power of contracting to the normal size, but tend to remain distended, and, when so distended, instead of the encircling capillaries being circular on section, they become flattened out, that is to say, oppose a greatly increased resistance to the onflow of blood. With this the alveolar walls atrophy, become thinned, and, in parts, wholly absorbed, so that neighboring alveoli run together to form large, thin-walled, and largely useless air sacs. This emphysematous atrophy, together with increased laying down of connective tissue around the

arteries and the bronchi, are the most characteristic changes occurring in the senile lung.

Throughout the organs of the body, with the exception of the central nervous system, there is this same tendency in senile atrophy for loss of the specific constituents to be accompanied by an increased fibrosis, This is, in part, *relative*; as the cells of the higher order diminish in size and in number, and the cells of the lower order-those of connective tissue-remain unaffected, the necessary result is that, in the shrunken organ, the fibrous tissue appears more prominent, and its proportion increases. But it is, in part, also, an actual increase—a replacement fibrosis: as the one element of a tissue diminishes the other proliferates and takes its place. The senile liver and spleen exhibit well these compensatory changes, the increased size of the trabeculæ of the spleen being very marked. Over and above both this relative and replacement fibrosis, there is a third form as already indicated, namely, the *perivascular* and, more particularly, the peri-arterial. How far this is primary, leading to the atrophy of the specific constituents of the tissue by reduced nutrition, and due to the alteration in the conditions of the circulation already noted, how far secondary to the atrophic changes already occurring in the organ, cannot be said with precision, but its intimate relationship to the vessels would seem to indicate that it is very largely primary. In any case, peri-arterial fibrosis is an important feature in senile changes.

We have, in the preceding paragraph, excepted the central nervous system from these fibrotic changes. A certain but relative slight amount of fibrosis is apt to occur in this, and then in connection with the vessels. Anything like the productive fibrosis in the brain substance proper is, under all circumstances, curiously rare; the glial cells have little tendency toward such. Remembering the liability shown by newly formed connective tissue to contract and compress included structures, it is fortunate for proper cerebration that this is so. When the brain in its rigid case atrophies, not solid tissue but fluid replaces the shrunken substance; increased cerebral fluid is poured out between the membranes and a condition is developed of *hydrops ce vacuo*.

ABIOTROPHY.

Somewhat closely allied to senile atrophy is a condition which has recently been brought into prominence, and has, by Gowers, been termed *abiotrophy.* This is the condition of premature death of the tissues or portions of tissues, not as the result of any immediate irritant. Possibly this should be regarded merely as a conception and explanation of premature cell decay. Nevertheless, if but a conception, it is a valuable one, inasmuch as it appears to explain in a manner more satisfactory than any other the development of certain otherwise obscure conditions. There is a series of morbid disturbances of the nervous system in which certain cells and systems of cells and the associated

ABIOTROPHY

tracts present degeneration, and, eventually, complete disorganization, the rest of the nervous system, apparently, showing no change. The development of these conditions is progressive. Many cases are hereditary. Nor can we find any one factor or set of factors to explain them unless we suppose that these cells have a shorter life than have the other neurons, that they exhibit a premature senility leading to precocious death. Such would seem to be a most satisfactory explanation of conditions like Thomsen's disease and other familial paraplegias. In these diseases, for a time, the mental and nervous conditions develop in a normal manner. In a few years one particular set of muscles undergoes atrophy with corresponding paralysis, and the motor centres governing these muscles show localized atrophy of their cells. Somewhat similar, it would seem, to these hereditary conditions, are the nervous disorders which may follow long years after an attack of syphilis, notably the condition of tabes or locomotor ataxia. One method of regarding these conditions is to imagine that syphilis is a disease that is never wholly cured; that, once in the system, the germs continue to grow and to produce their toxins, and that these toxins have, as it were, a cumulative affect until, at last, owing to their continued irritation, they bring about the death of certain groups of nerve cells which are more susceptible to their influence than are others. The difficulty in accepting this view is that, in such cases, we have no other sign of the continued existence of the germs of syphilis. We do not find, for example, indications of active gummata or other syphilomata. The individual is incapable of infecting others with his disease, and, judging by macroscopic and microscopic appearances, the disease is, and has been for years, wholly arrested. It is more satisfactory to suppose that there has been a general intoxication of all the nerve elements to such an extent that, although the intoxication has been recovered from temporarily, the cells have, notwithstanding, been weakened so that now, under the normal strain, these cells, being called upon to perform no more than the normal amount of work, become easily exhausted and undergo premature dissolution. This is Edinger's "Ersatz theorie" (or offset theory) of the varying symptomatology of tabes:1 "Function determines the symptom complex." While most often in tabetic patients the symptoms of incoordination show themselves in the more used lower limbs, in coachmen, tailors, and cobblers, the weakness in coordination and lightning pains first manifest themselves in the arms; whereas where occupation demands eye strain, it is disturbances of vision and optic atrophy that are the prominent early symptoms. It is the nerve centres controlling the groups most commonly in use that first undergo atrophy.

A most suggestive example of what may be termed general abiotrophy, throwing light upon these more specialized abiotrophies, has recently been adduced in Bardeen's studies upon the effects of x-rays upon

¹ Edinger, Deutsch. med. Woeh., 1904 : Nos. 45, 49, 52, and 1905 : Nos. 1 and 4; Dublin Journ. of Med. Sci., 1901, and Russel, Montreal Med. Journ., 39 ; 1910 : 162.

frogs' spermatozoa. By subjecting the sperm to the rays for a few minutes it is found that they are still capable of fertilizing the ova; the individual life begins, but the larvæ, growing, all die prematurely, none survive beyond the second week. There is, that is, cell exhaustion after a certain early period.

REVERSIONARY METAMORPHOSIS; KATAPLASIA.

In studying the various conditions of atrophy, we cannot fail to be impressed by the fact that, in addition to mere shrinkage of the individual specific cells of a tissue, another order of changes is frequently met with which I have elsewhere spoken of as reversionary degeneration,1 but for which, perhaps, a better term is that of kataplasia, introduced by Beneke. We find frequently, that is, that in the process of atrophy accompanying the gradual disappearance of their specific features, certain highly organized cells present appearances which simulate very closely and very curiously those presented by the developing cell. We may without hesitation say that there is a harking back or reversion to an earlier and more embryonic condition. The most striking example of this is to be seen, as Fujinami and others have pointed out, in striated muscle fibres. Such striated muscle fibres, if their development be studied, originate from cells rich in cytoplasm, known as sarcoblasts. In these, at first, no signs of striation are visible, but gradually, along one side of the cell, the striæ are seen to make their appearance, and thus, at one stage, we find cells frequently multinuclear, with undifferentiated protoplasm in the immediate neighborhood of the nuclei, and differentiated protoplasm, with striation showing itself in the cell substance away from this. As the fibres become more developed, the number of nuclei diminishes; the amount of undifferentiated protoplasm becomes less and less until at last in the complete fibre we have relatively rare nuclei situated in the outer aspect of the fully formed fibre.

If we examine the muscle fibres at the edge of an invading newgrowth, in those that are undergoing compression and diminution in nutrition, we find that the very reverse processes are occurring. The first disorganization of the fibrillar structure takes place around the nuclei more particularly, so that the nuclei become surrounded by a definite area of undifferentiated cytoplasm. They multiply and, as this process continues, we find, in place of fully formed striated fibrils, large multinucleated protoplasmic masses, while, as a last stage, we have evidence that these masses may divide up, the cytoplasm accumulating around individual nuclei, so that separate cells pass off from the multinucleated masses closely resembling the individual sarcoblasts of the embryonic period. The stages here closely reproduce in reverse order those seen in the process of development. And this is far from being

¹ Jacobi Festschrift, 1900; 422,

REVERSIONARY METAMORPHOSIS

the only example of this process. The bile ducts and the liver cells have a common origin. The liver, for example, if we follow its development, first shows itself as a series of separate cell columns, and only as the organ becomes larger and more important does this tubular arrangement of the hepatic cells become unrecognizable, the vascular relationship of the individual liver cells becoming more prominent than the relationship toward each other, although, throughout, the bile capillaries represent the lumina of the primitive hepatic tubules. As already stated, in the development of the human liver there is a stage in which there is an absence of differentiation between the epithelium lining the bile ducts and the hepatic cells proper. When, now, as happens frequently in cirrhosis, there is progressive atrophy of the hepatic parenchyma, it is frequently possible to notice that the transition from hepatic cells to bile ducts becomes gradual. At the periphery of the lobules clusters of liver cells are to be seen separated off from each other by intervening fibrous tissue, and these cells are of an intermediate type. They are small; the nuclei also are smaller than those of the ordinary liver cells, but larger than those of the bile-duct epithelium; the amount of cytoplasm is greatly diminished; there is a want of a perfect, welldefined lumen. This well-defined lumen, we may add, is also wanting in the developing bile duct. We have, in short, a reversion to the period in which bile-duct epithelium and liver cells were undifferentiated. It is these imperfectly formed strings of cells which compose the socalled proliferated bile ducts observed in many cases of cirrhosis of the liver. Let us repeat, these are not true bile ducts, for the arrangement of the cells is not perfect.

A certain amount of confusion has originated with regard to these from a want of recognition that two separate orders of events may show themselves in the cirrhotic liver; namely, that liver may, in the main, show evidences of progressive atrophy, or on the other hand, may exhibit compensatory hypertrophy. Undoubtedly, in the latter case, we have the reverse process occurring, namely, where the liver cells have been quite destroyed there may be a development of new liver cells from the still persistent bile ducts. Acknowledging this, the process of formation of apparent bile ducts while there is progressive atrophy is also to be recognized.¹

And other instances may be afforded, notably the cubical appearance taken on by the lining cells of the pulmonary alveoli in cases of chronic compression of the lung, or of interstitial fibrosis, reverting thus to the type of epithelium seen in the lung before birth; in cases of subacute inflammation of the kidneys, not only the epithelium of the tubules, but also that of the glomeruli may assume the embryonic cuboidal type.

Herein, it must be admitted, is the difficulty in arriving at a conclusion regarding individual cases; it is not always easy to determine

¹ A full study and discussion of the debated subject of regeneration *versus* degeneration of the liver cells in different conditions is by Muir. Journ. of Pathol , 1908; 287, where is afforded a good bibliography,

whether we deal with cells in the process of reparative development or with these kataplastic changes. In the first case here cited there can be no question; the processes seen at the periphery of a growing tumor, although similar to those described as occurring in repair (p. 622), are degenerative, not regenerative. As regards the liver, Opie' has recently shown, in his study of the effects of combined effects of intoxication and bacterial infection in the production of cirrhosis, that a distinct interval may intervene between the old bile ducts and these "pseudobile ducts" within the lobule; such are clearly of parenchymatous origin, whether we regard them as degenerative or regenerative. In fact, from the point of view of function, these processes, like the closely allied anaplasia of tumor cells, are degenerative; from the point of view of vegetative activities, they are regenerative.

Such kataplasia comes very close to the anaplasia which von Hansemann demands as the starting point for new-growths, the distinction being that the anaplastic cell in the process of simplification loses the power to rereturn to the normal differentiation, whereas in this kataplasia that power would seem to be preserved. Possibly in senile conditions this tendency to reversion is of the anaplastic rather than the kataplastic type—a matter of some importance in connection with the theory of neoplasia.

¹ Trans. Assoc. Amer. Phys., 1910 (about to be published).

CHAPTER XXVI.

THE REGRESSIVE TISSUE CHANGES-(CONTINUED).

THE DEGENERATIONS AND INFILTRATIONS.

WHILE we are forced, for various reasons, to recognize the condition which we term simple atrophy, we have already said enough to show that, even in the simplest cases, we in general have to deal with more than merely a progressive reduction in the volume of the cell constituents. The very heaping up of what we may term by-products must, in itself, tell upon the cell and its activities. So that, both from a histological and physiological standpoint, the cell undergoing simple atrophy eventually becomes degenerated. The attempt has been made in the past to distinguish between the two orders of events, either of which might lead to, or might accompany, regressive disturbances in the cell. On the one hand, it was thought that there could be recognized processes of pure regressive metamorphosis, the abnormal products that appear within the cell being due to the breaking down of the cytoplasm; on the other hand, that there was a process of laving up of preformed material gained from the lymph or blood. Conditions exhibiting the former process have been spoken of as degenerations proper; those showing the latter, as infiltrations, Undoubtedly we do encounter examples of what are true infiltrations. The leukocytes in the coal-miner's lung, containing inert particles of coal, certainly contain substances which have been obtained from outside, and which have not been acted upon by the living protoplasm of the cell. But the more we study the various regressive metamorphoses, the more is it brought home to us that uncomplicated infiltration is comparatively rare. We are apt, for example, to speak of fatty and glycogenous infiltration-in the liver cell, for example-but if we study the physiology of these processes, we are rapidly forced to the conclusion that we have to deal with something much more complicated than mere absorption of fat or of glycogen from the blood or lymph. The cells, it is true, become infiltrated with or contain the substances in question, but the process is not that of direct absorption of the fat or glycogen in a preformed condition from body fluids. Everything points to a series of synthetic processes, the activities on the part of the cytoplasm leading to these deposits. We do not, for example, under normal conditions, detect fat as such in the blood. On the contrary, we have evidence that, to a very large extent, it is saponified before it is absorbed by the intestinal mucosa. We may, as Heidenhain pointed out, find a few leukocytes containing fatty globules in the terminal portion of the villi after a meal containing fat, but, as we pass

down the villus, it seems very clear that before the villus is left, these cells dissolve up and break down, and in their place no fat is seen; it has been converted into some soluble compound. The liver cell, however, absorbs that soluble compound from the blood, and reconverts this by the activity of its ferments into fat. The same would seem to be true in connection with glycogen. On the other hand, as we shall proceed to point out more fully, in discussing the subject of fatty degeneration, when in the diseased cell there appear minute globules of fat, and the cell shows evidences of breaking down, the old idea that in these special cases we had the actual process of breaking down of the proteid framework of the cell substance, with liberation of the fatty molecules, has also to be given up, at least to a considerable extent. Recent observations point to the fact that degeneration of this nature is not the prevailing type. The evidence would seem to prove that the minute globules of fat have, in the main, been absorbed from without. We can rarely, therefore, make a clear distinction between the degenerations and the infiltrations, although for convenience we retain these terms for particular conditions. Here I shall treat these conditions together, and, in order to pass them in review in due order. I shall consider the various disturbances of metabolism and their outcome in recognizably altered conditions of the tissues in the following order:

1 Disturbances of proteid metabolism (excluding pigmentation).

2. Disturbances of intracellular water (cedema and vacuolization of cells).

3. Disturbances of fat metabolism.

4. Disturbances of carbohydrate metabolism.

5. Calcareous deposits.

6. Deposition of other products of cell metabolism.

7. Pigmentation and pigmental deposits.

DISTURBANCES OF PROTEID METABOLISM.

OF SIMPLE TYPE: CLOUDY SWELLING.

Under various conditions—in fact, this is the most common morbid change we encounter in certain tissues at autopsy—muscular tissues and certain glandular organs exhibit a condition which is now most frequently known as cloudy swelling, or albuminous degeneration. Upon making a section of the affected organs, they have a duller appearance. Instead of the healthy look of the heart muscle, for example, it appears as though, to use an oft-quoted description, the heart had been momentarily dipped in boiling water. With this there is a certain amount of swelling, best seen in the kidney, where the cortex, which is particularly affected, is on section found raised_slightly above the level of the medulla, the diameter of the cortex being at the same time somewhat increased. And, now, upon examining sections of the tissue.

CLOUDY SWELLING

whether freshly cut, or after treatment with the ordinary hardening reagents, the individual cells are no longer so transparent as normal. They have a cloudy, ground-glass appearance, while, in well-developed cases, the nuclei look as though obscured by the deposit of a finely granular material in the surrounding cytoplasm, and stain more feebly than normal. These are the main features of cloudy swelling. The main chemical reactions are that, by the agency of weak acid, or weak

alkali, this cloudiness can be cleared up; something is dissolved out of the cells which now become transparent. That something would seem to be, from its reaction, a body of a proteid or albuminous nature, unacted upon by alcohol or chloroform, but stained brown by iodine, and giving the xanthoproteic reaction.

Conditions Leading to Cloudy Swelling. —As already stated the conditions under which we find these particular changes are very many. Most commonly they are met with in cases of acute infection and high fever. They may show themselves, however, under the action of certain poisons, as, for example, in the early stages of phosphorus



Cloudy swelling of cells of convoluted tubules of kidney. \times 400, (Ribbert.)

poisoning: following upon extensive burns, and here as early as six hours after the infliction of the burn; in cases of subjection of the individual to a high external temperature; in cases where there is no sign of febrile disturbances, or even, as already indicated a few pages back, in conditions of prolonged hunger, in which the irritant setting up the disturbance, if any, must be a direct produce of metabolism, or when, on the other hand, we have to deal with the first stage in the disorganization of the cytoplasm.



The Altmann granules in normal cells of the in convoluted tubules of the kidney. (Lubarsch.) ((Fro. 293

Enlargement and irregularity of the Altmann granules in renal epithelium, with cloudy swelling (experimental inflammation of the kidney), (Lubarsch.)

Examination of the cells of the convoluted tubules of the normal kidney that has been suitably hardened after death, reveals the fact that their cytoplasm is not homogeneous, but exhibits closely set rows of minute globules or granules running across them from the basement membrane to the lumen. These minute globules are so closely packed that they almost simulate rodlets; in Henle's tubules, indeed, employing

884

Müller-formalin, they may be quite indistinguishable from closely packed rodlets traversing the cell, although by other modes of hardening their composite nature is revealed. These stain intensely with iron hematoxylin, and are dissolved out by weak acid. In cloudy swelling they are replaced by an irregular distribution of what are apparently similar globules, but on the whole larger, varying considerably in size. These globules are evidently of the same order as those seen in the healthy cell, but now they are swollen and disordered. With this, nuclear changes show themselves. Contrary to the usual teaching, the nuclei of the cells are not always obscured. One has but to study a series of kidneys exhibiting the naked-eye appearance of cloudy swelling to be convinced that this is not so; there may be well-marked swelling, with opaque, finely granular appearance of the cytoplasm, and with the nuclei more deeply stained and larger than usual. There appear to be three stages: the first, of increase in the chromatin of the nuclei; the second, of accumulation of the chromatin in clumps at the periphery of the nucleus, the achromatic substance being accumulated in the centre (chromatolysis), the final stage is such extensive loss of chromatin that the nucleus is almost unrecognizable, if it does not undergo karvorrhexis. The indications are those of stimulation, giving place to exhaustion of the nuclear material, with loss of chromatin.

The exact relationship of these changes to those occurring in the cytoplasm have not been determined. Lukjanow' is of the opinion that the development of the albuminous granules is associated with the actual giving off of "plasmosomes," or minute globular extrusions from the periphery of the nucleus, and that these undergo alteration and conversion into the cell granules; but this view still lacks confirmation, save to this extent, that fatty degeneration and cloudy swelling are very intimately connected, that in all cases of definite cloudy swelling minute fatty globules are present also in the cytoplasm (Bennario⁵); and several observers have noted the plasmasome formation in wellmarked conditions of fatty degeneration.

The preliminary increase in size and staining power of the nucleus, together with the increase in the bulk of the cytoplasm, would suggest that in cloudy swelling we deal with increased absorption on the part of the cell, and that the albuminous globules are the indication of matter assimilated and not utilized. Virchow, indeed, regarded the cloudy cell as supporting his view that inflammation was, at base, a stimulus to increased autrition. On this view cloudy swelling is an indication of increased absorption of foodstuffs with imperfect conversion and utilization of the same. It is possible that both opinions are correct up to a certain point. Cloudy swelling manifests itself in the active tissues of the body, the muscles and the main excretory glands; in the latter case, in evident connection with the removal from the blood of toxic matters. It may, indeed, be produced by overwork. Thus,

Grundzüje einer allg. Pathologie der Zelle, Leipzic, Veit, 1891.
 Die Lehre von der trüben Schwellung, Würzburg, 1891.

CLOUDY SWELLING

Schilling¹ has demonstrated that in the rabbit, if one renal vein be ligatured, and its kidney, therefore, rendered functionless, in fortyeight hours there is developed well-marked cloudy swelling in the convoluted tubules of the second order in the other kidney. Evidently, in this case, the cells are stimulated to increased work and increased absorption by the excess of normal urinary constituents, and the cloudy swelling is a precursor of subsequent hypertrophy. The nuclear changes can only be regarded as exemplification of the fact that the nucleus takes an active, if not a controlling, part in cell function; and, further, of the principles already laid down, that increased activity within certain limits leads to increased growth, beyond those limits to increased disintegration of living matter—in this case of the nuclear chromatin.

That the cytoplasm in the ordinary cases of cloudy swelling undergoes actual growth is at least debatable. The cells obviously increase in size, but this increase is in part due to the increase in paraplasmic deposits (the "cloudy" granules or globules), in part due to a hydropic condition, and increase in watery constituents. Cloudy swelling may, indeed, pass on imperceptibly to a vacuolar or vesterilar degeneration of the cell (p. 903), as may be demonstrated in intexication with progressive amounts of cantharidin.

The albuminous globules, we would repeat, appear to be of the same order as the smaller paraplasmic globules seen in the normal cell. Regarding the processes of assimilation and disintegration of living matter as largely reversible, such paraplasmic matter may be indifferently either matter absorbed and in part built up, or be matter dissociated from the cell substance proper and in part disintegrated.


This condition of cloudy swelling must not be confused with another condition, that of granular disintegration to which Durante² more especially has called attention, or "tropfische Entmischung," as it has been termed by Albrecht.³ The latter is a disintegrative condition of the cytoplasm itself, an indication of cell death. If, as pointed out by Landsteiner,4 the kidney cells be taken and placed in water, they become filled with small, cloudy, packed vacuoles. The condition appears to be allied to the granular degeneration noted by Verworn⁵ in injured infusorians. In many of these, if the unicellular organism be cut in two, from the surface of the wound inward, the previously homogeneous cytoplasm now, when exposed to the water, becomes progressively converted into an agglomeration of minute droplets. Durante and others have noted this granular disintegration in muscle cells in severe febrile conditions; it may follow upon cloudy swelling, but where, as in the latter condition, weak acetic acid, dissolving out the droplets, brings back the striated condition of the fibres, in the former the striæ are wholly lost; we deal with a liquefactive necrosis. Similarly, Landsteiner notes that in the kidney cells above described,

⁵ General Physiology. Translated by Lee, p. 236.

¹ Virchow's Arch., 135; 470. ² Bull. de la Soc. Anatomique, Février, 1900,

³ Lubarsch's Ergeb., 2: 1895; 151. ⁴ Ziegler's Beitr., 33: 1903 (237.)

staining with iron hematoxylin demonstrates the presence of the finer albuminous granules between the larger droplets of disintegrated cytoplasm. It is not improbable that imbibition of increased fluid is, as Albrecht suggests, a factor in the production of the larger albuminous



Hyalopus (Gromia) dujardinii, granular dieintegration: I, whole individual; numerous pseudopodia are extended from the egg-ahaped membranous shell; at the left they are being drawn in; II and III, pseudopodia eu off; granular dieintegration is developing; the globules and droptets of protoplasm are held together simply by a loose viscous ground substance; between them lie scattered large hyaline protoplasmic droptets and viscous globules. (Verworn.)

droplets of cloudy swelling, as compared with the extremely fine albuminous granules of the normal, but these remain clearly distinct from the vacuoles of granular disintegration.

To epitomize, cloudy swelling is the expression of overstimulation of

GRANULAR, WAXY, AND FIBRINOUS DEGENERATION 887

the cell by absorbed substances leading to disordered metabolism and the heaping up of paraplasmic matter of albuminous nature. It is not in itself a necessary cause of cell death. Judging by the constant presence of the condition in deaths from febrile disorders, it is a constant accompaniment of bacterial intoxication, and as such must often be followed by return to the normal state.

Like the granular disintegration just noted, "*waxy degeneration*" of the muscle is an expression, not of reaction on the part of the living cell, but of cell death. The condition, is therefore, more appropriately considered along with the necroses (see p. 979), and this notwithstanding the fact that the albuminous constituents of the cytoplasm are in the main involved.

Fibrinous degeneration, in the strict sense, *i. e.*, the formation and deposit of fibrin in the living cell, is, if it ever occurs, very rare. Mallory¹ gives pictures of intracellular fibrin in vacuoles within the liver cells, but these cells, as he points out, are undergoing necrosis, and in general the fibrinous or fibrinoid coagulation within cells is an evidence of cell death, and so must be considered along with the necroses.

¹ Journal of Medical Research, 1: 1901: 264,

CHAPTER XXVII.

THE DEGENERATIONS AND INFILTRATIONS.

DEGENERATIONS ASSOCIATED WITH THE DEPOSIT OF SCLERO-PROTEINS AND CONJUGATED PROTEINS.

THERE is a complicated series of degenerative conditions in which there is laid down in the tissues or spaces of the body material which, in the unstained condition, has a translucent or glassy appearance. When colorless and firm, we speak of the deposits as hyaline; when colorless and fluid, or semifluid, as mucoid; when semisolid, or solid, and of brownish, glue-like appearance, as colloid. There was a time when these appearances and terms were regarded as indicating the presence of specific substances, and, as a consequence, the terms hyaline, mucoid, and colloid degeneration are still employed, with, as a result, very considerable confusion. We now know that these different appearances may be brought about by a multiplicity of substances, but from the wide use of the terms it is still necessary to bring together the various conditions under the old names, distinguishing under each title the various orders of substances which may give rise to the different orders of deposit. One exception may be made, that, namely, of amyloid infiltration. This, in an unstained condition, is preëminently hyaline in appearance, but its reactions are so characteristic that for long it has been admitted as a separate entity.

MUCOID DEPOSITS.

Mucoid Degeneration and Mucinous Deposits.—Physiologically, mucin—or, more correctly, the mucins (for there is considerable variation in composition of mucinous material gained from different regions) —is laid down in the organism in two conditions: (1) as of definitely *intracellular* origin, and (2) as *intercellular* matter, without obvious secretion from cells. The type examples of the former are afforded by the mucous salivary glands, and the goblet cells of the intestinal mucosa, of the latter by the mucin of "Wharton's jelly" in the unbilical cord, and the mucinous intercellular matrix of embryonic tissues in general.

In either case we deal with material which has definite physical and chemical characteristics; it is viscid, swells up with water, is soluble in weak alkalies, is precipitated by acetic acid, not being dissolved in excess, as also by alcohol. It stains with basic dyes, more particularly with thionin.

Its composition, as above noted, varies,¹ but the true mucins have this in common, that they are composed of, and on decomposition yield, a

¹ See Cutter and Gies, Amer. Jour. of Physiol., 6: 1901: 155.

MUCOID DEPOSITS

protein and a carbohydrate, which reduces Fehling's solution—glucosamin, or, according to Levene,' chondroitin-sulphuric acid. They thus show some relationship to cartilage and the amyloid material, to be presently noted.

Closely allied in physical properties are the *pseudomucins*, bodies also yielding a reducing substance, and being of the nature of glycoproteins, the reducing substance, according to Leathes, being a reduced *chondrosin* (the carbohydrate constituent of chondroitin-sulphuric acid). Of these pseudomucins, more than one has been distinguished. They differ from the mucins proper in not being precipitated by acetic acid. We may dismiss them by saying that they are found in considerable quantities in ovarian cysts, which never contain true mucin, and that in this situation they are clearly products of excretion from the lining columnar cells.

Intracellular Mucinous Production.—'The main condition in which we observe a condition of excessive production of mucin is in catarrhal conditions of mucous membranes. In these not only is there a marked increase in the number of goblet cells, and excessive discharge from these, so that the surface becomes covered by a layer of mucus, but the individual cells may degenerate, their whole substance, the nucleus included, appearing to become used up, so that we can truly speak of a mucoid or, more correctly, mucinogenous degeneration of the cells.

For mucus, as such, would not seem to be present in the cells, but a precursor, *mucinogen*, and this in the form of small globules. The histological studies upon the development of goblet cells indicate that a succession of events occurs of the same order as that described by Nissen for the mammary gland cells, namely, direct division of the nucleus, passage of one daughter nucleus into the outer part of the cell, when it undergoes chromatolysis and gives off plasmosomes, which, as they pass farther from the nucleus, swell up and take on the characters of mucinogen globules. When the goblet cell ruptures and discharges these globules they swell up and fuse into a homogeneous mass of mucin. The other nucleus left behind becomes surrounded by an increasing mass of cytoplasm, and so the cell becomes restored.

An even more active, not to say excessive, development of mucous cells and production of mucin occurs in that form of carcinoma originating from mucous membranes, more particularly of the intestines, known, unfortunately, as "colloid" cancer. In this the production may be so extreme that, through pressure, if not through the actual mucoid degeneration of the cells already noted, the cells of the cancer alveoli undergo destruction, and the alveoli become represented by masses of dense, inspissated mucin. The semisolid translucent material found in these cancers is not true colloid; it gives all the reaction for mucin.

Interstitial Mucinous Infiltration.—This may be found pathologically in: 1. Senile atrophic tissues, as in the cartilage and bones (medulla) of old people.

¹ Medical Record, 1900; i: 188

DEGENERATIONS AND INFILTRATIONS

2. In the connective tissues in experimentally induced myxœdema, and in the same areas in the early stages of the disease in man.

This mucoid nature of the swollen subcutaneous tissues gave the name to the disease, and the earlier observers regarded increased interstitial mucin as the characteristic change in cases of atrophic disease of the thyroid. Halliburton has shown that in cases of longer duration there is no increase in mucin beyond what is found in normal connective tissue. These findings suggest that in the first stage of the disease the connective tissues take on active growth, and that, as the proliferated tissue cells mature, the mucin, as in the developing fectus, undergoes diminution.

 In actively developing tumors of the connective-tissue type, sarcomas, fibromas, etc., and in the interstitial tissue of carcinomas.

The frequency of mucoid changes in tumors is more apparent than treal (see p. 720), and true myxomas are rare.

4. In inflammatory new-growths, as in developing granulation tissue.



Section of thyroid gland, showing vesicles with contained colloid: a, colloid; b, secretory cells with granules. (After Bozzi.)

In all these cases it will be observed that we have to deal with either active tissue growth, with immature tissue, or, on the other hand, with tissues undergoing reversion. The remarks made (p. 878) upon reversionary metamorphosis will have prepared the reader to comprehend why it is that these apparently opposed conditions present the same change.

Our knowledge of the intermediate metabolism in cells is so slight that we know nothing of the stages leading up to the formation of glycoproteins. The observations above recorded upon goblet cells indicate that the nucleus in the one series of cases controls their formation; there is no evidence that it does this in connection with interstitial mucin, save that Hektoen records that.

in advanced cases of senile atrophy of bone and cartilage, globules or masses of mucin (mucinogen) are to be recognized within the cells of the affected areas.

COLLOID DEPOSITS.

I am inclined to restrict the use of this term to a single deposit, namely, the material accumulating in the vesicles of the normal, and, to an excessive extent, in those of the ordinary goitrous thyroid. It is more accurate to describe the so-called colloid cancer as a mucoid cancer, for the material forming in such cases is of the nature of inspissated mucin. Through inspissation, brownish, solid, colloidal matter may

AMYLOID INFILTRATION

accumulate and fill ovarian cysts, as also isolated and multiple cysts of the kidney. In each case the composition differs from that of the type colloid of the thyroid. The main constituent of the latter is a compound protein, a compound of globulin with an iodine-containing body, iodothyrin, associated with which is a nucleoproteid.¹

Allied to this, although not identical, is the colloid material of the vesicles of the anterior portion of the pituitary body, to which the researches of Saint-Remy and Benda, and more recently of Schäfer and Herring, have drawn attention.

AMYLOID (CHONDROID) INFILTRATION.

There is an infiltration which has suffered from a succession of infelicitous names still in use—waxy (when the deposit has no relationship to the waxes), lardaceous (and it is devoid of fats), amyloid (and unrelated to the starches). A satisfactory appellation has at last been suggested by Wells, namely, "chondroid," but we fear that it might be regarded as pedantic to foist this upon the student, and out of respect to Virchow will continue to call it amyloid.

We noted that mucin is to be regarded as a glycoprotein, a compound between protein and a nitrogenous carbohydrate. Another substance of the same order is amyloid—a hyaline deposit not found in normal tissues, though allied to the matricial matter of cartilage. Into what knowledge we possess regarding its composition and mode of formation we will enter after having detailed the well-ascertained facts regarding its microchemical detection and distribution in the organism.

Amyloid material laid down in the tissues produces characteristic changes in the appearance of those tissues, and gives most characteristic reactions. The deposit may be either (a) generalized, affecting several organs, and this is the most common condition, or (b) localized, then affecting relatively small areas of tissue and inflammatory and other new-growths. The statements which follow refer in the main to the generalized form, the localized being discussed later.

Generalized Amyloid Infiltration (Amyloidosis).—This, when advanced, affects a large number of organs, but is most noticeable in the spleen, liver, and kidneys. The tissues which, so far, have not been found affected are the epidermis and cutis, bone, lung tissue, and nervous tissue proper. If the spleen and liver be unaffected, there is little use in studying the other organs for this change.

Naked-eye Appearances.—The spleen is found distinctly enlarged, and with rounded edges, pale, and usually much firmer and denser than normal. Upon section it has a semitranslucent, waxy appearance (hence the terms *waxy*, *bacony*, and *lardaccous* degeneration), and either this appearance is evenly diffused (*bacony spleen*), or rounded bodies project from the cut surface, somewhat of the size and appearance

¹ See more particularly Oswald, Virchow's Arch., 169: 1902: 444.

DEGENERATIONS AND INFILTRATIONS

892

of boiled grains of sago, embedded in the same (*sago spleen*). These little elevations are the affected Malpighian bodies. The liver shows a similar or more marked enlargement, is firm, with obtuse edges, and is pale and waxy upon section; the kidney also shows distinct enlargement and pallor; whether it is firm or flaccid depending upon the existence or non-existence of extensive parenchymatous disturbances.

Reaction of Amyloid Material.—In these and other organs the presence of amyloid change is most rapidly and most surely determined by the iodine reaction. On to the cut surface is poured official tineture of iodine diluted until it is the color of port wine, or Lugol's solution, or, best of all, according to Kyber,¹ iodine, gr. 10; pot. iod., 1 dram; aq., 10 ounces. This should be left on until the surface assumes a pure yellow tint, when any amyloid material present will take on a red or brownish color. Care has to be taken to wash off previously any blood which may have exuded on to the surface. If, now, a 5 to 10 per cent. solution of sulphuric acid be poured on, the amyloid areas assume a darkviolet to black color, the non-amyloid parts remaining unaffected (Virchow). It should be noted that the sulphuric acid reaction is not absolutely constant. In general the iodine reaction is reaction as decisive.

The iodine reaction is also of use in sections, though here, by transmitted light, it is the affected parts that have a semitransparent, vellowish appearance, the rest of the section being granular and more brownish. Such sections are best mounted in glycerin, or Farrant's solution, to which some iodine has been added. With a little experience, the existence of anything beyond the slightest grades of amyloid infiltration can be detected in sections of tissue stained in the ordinary way with hematoxylin or hematoxylin and eosin. The position of the affected parts in the spleen, liver, or kidneys, and the peculiar translucency and lack of stain are most characteristic. The condition is, however, most clearly demonstrated by the differential stain afforded by watery solutions of many of the basic aniline dyes, notably gentian violet, methyl violet, and methyl green. A long series, including safranin, might be given. To give good results, material, if not fresh, should have been kept in alcohol; or, if in Müller's fluid, this must be well washed out and the tissue kept for some time in alcohol before applying the test. To clear the groundwork, and to fix the stain more thoroughly in the amyloid areas, it is advisable to place the section for a short time in very dilute tannic, or even hydrochloric, acid, after washing out the excess of stain in water. Methyl violet sections, for example, so prepared, show the amyloid material standing out sharply as a rich rose pink against a paler, often somewhat slaty colored background.


When we come to study such preparations carefully, the seat of the amyloid change becomes evident. Most often it involves the capillaries; this is especially noticeable in the liver. Here it is the intermediate zone of the individual lobules that is at first affected. It is in connection with the capillaries of this region that the deposit occurs, and becomes so

⁴ Virchow's Arch., 81: 1880: 278 and 420.

AMYLOID INFILTRATION

pronounced that, apparently through the pressure as well as through disturbance of nutrition, the cells lying between the thickened capillaries become atrophied.

Careful examination of suitable specimens shows that the endothelium of the capillaries is not the seat of the charge—that the endothelium still remains, although its cells may undergo fatty degeneration. The amyloid material is laid down *outside* the endothelium, and laid down irregularly, so that one side of a capillary may have a much thicker deposit than the other. As a result of this infiltration two things happen, namely, that the capillary itself is compressed and its lumen diminished, and that the liver cells, both by pressure and by disturbance of nutrition, show evidences of fatty degeneration and atrophy, until in advanced cases, in this intermediate zone, scarcely any liver cells may be seen, and there appears to be little more than a belt of translucent amyloid material. As the process advances, the amyloid deposit trespasses more and more upon the periphery of the lobule, as also, to a slighter extent, toward the centre, until very little healthy liver tissue is left.



Amyloid degeneration affecting the liver; slighter grade: the cells are still present with but moderate atrophy; the irregular deposit of amyloid around the capillaries is well marked. (After Ribbert.)

Amyloid degeneration of liver advanced: a, atrophied liver cells; b, transverse section of a capillary surrounded by a broad ring of amyloid material; c, a capillary cut longitudinally. (Ribbert.)

At the same time the process affects the branches of the hepatic artery. Here it is the middle coat that is primarily affected, and in that, apparently, not the muscle cells, but the connective tissue. From here the change extends more particularly into the deeper layers of the intima. In very advanced cases the connective tissue of the walls of the veins may also show anyloid change. One note of caution is here to be given, namely, that employing the iodine reaction upon the liver, this also acts upon and causes brownish discoloration of the glycogen within the cells, so that at first sight it may be thought that there are intracellular deposits of the amyloid material. Treatment with the aniline stains, however, has no effect upon the glycogen, and demonstrates that we are dealing with a different substance.

DEGENERATIONS AND INFILTRATIONS

While thus the deposit occurs most often immediately outside the endothelium of the finer capillaries, it may also affect the middle coats of the smaller arteries. Only in advanced cases is it seen affecting the walls of the veins. More frequently, as in the spleen, it may affect the reticulum of lymphoid tissue and the connective tissue. So, too, in advanced amyloid change in the kidney, the basement membrane of the collecting tubules is seen clearly to become the seat of these deposits, though care has to be taken to distinguish between the appearances thus produced (particularly around the small Henle's tubes) and the very similar appearances brought about in the longitudinally arranged capillaries of the medulla. We have never been able to satisfy ourselves regarding the intracellular development of amyloid material in the form of spherules, though some observers have described such deposit within the cells. Nor can we accept Maximow's conclusion that the Altmann's granules in the liver cells play a part in the development of the amyloid material.



Amyloid degeneration of the media of a small artery of the kidney: the amyloid deposit is around the muscle fibres leading to their atrophy. (Ribbert.)

for the amyloid deposits around the hepatic capillaries clearly continue to grow after the liver cells have undergone total atrophy.

Conditions under which Amyloidosis Shows Itself.—Amyloid deposits show themselves most characteristically in conditions characterized by prolonged suppuration and discharge from the system of proteins in one or other form. The most frequent precursor is tuberculosis of bones in the form of Pott's disease, with cold abscess formation, or of osteomyelitis of the extremities, though it frequently follows, also, intestinal and abdominal tuberculosis; in un-

complicated pulmonary tuberculosis it is relatively rare. Chronic ulcerative syphilis is at times responsible, as is subacute or chronic supprative osteomyelitis with sinus formation. More rarely it has been found associated with leukemia and malarial cachexia; still more rarely with chronic Bright's disease (albuminuria) and prolonged and excessive lactation.

Localized Amyloid.—Apart from the generalized amyloidosis, there is encountered occasionally a restricted local amyloid deposit, with no signs of the change in the usual sites, the liver, spleen, etc. Such may be found in localized granulomatous masses, of tuberculous or syphilitic origin; it has been noted in granulation tissue of the conjunctiva, in connection with the cartilages of the larynx and upper part of the respiratory tract, and somewhat characteristically in tumors—fibromas and sarcomas—of the upper air passages. It is noticeable that in these conditions, as pointed out by Ribbert, the smaller vessels are relatively unaffected; the amyloid change affects the interstitial tissue, forming a

AMYLOID INFILTRATION

network which, Eden suggests, follows the lymph channels. It has been noted also in the lymph glands nearest to areas of local suppurative inflammation.

The Nature of the Amyloid Matter.-The blue color gained by treating the amyloid material with iodine and sulphuric acid led Virchow to suspect some relationship between it and the vegetable products, starch and cellulose, hence the name by which it is now generally known (amylum, starch). Needless to say, this was a mistaken deduction, and soon its protein nature was demonstrated (Friederich, Kekule). But for long the nature of this protein baffled analysis. It appeared to be related to hyaline material, and cases were reported in which there was apparently a combination of, or a transition between, hyaline and amyloid material -cases of only partial reaction with iodine and the aniline dyes. But the composition of hvaline matter was equally difficult to determine. Others regarded it as modified fibrin. Without entering into the details of the various theories regarding its nature, it may be said that the first sure advance was made by Krakow,1 who demonstrated clearly that it is a compound protein, a combination of a proteid (histon) with chondroitin-sulphuric acid (C18H27NSO17).

Chondroitin-sulphuric acid in its turn yields chondroitin ($C_{18}H_{27}NO_{11}$), and from this can be gained chondrosin, a reducing substance, of the nature of a nitrogen-containing carbohydrate. Pure amyloid separated from nucleoproteid is an almost white powder, and, like the nucleoproteins, is resistant to digestion with pepsin, though Neuberg found it to be acted upon by trypsin. It is this resistance to peptic digestion that enables us to isolate it from the main mass of proteins.

Amyloid thus comes into the category of the glycoproteins, and, by containing chondroitin-sulphuric acid, is found to be allied in characteristic constituents to cartilage and yellow elastic tissue, both of which yield the same acid. Indeed, from the normal aorta, presumably from its elastic tissue, there can be gained a substance closely allied to amyloid. What we have said regarding the mucins will indicate that they are of an allied group. Like the one group of mucins, and the specific substance of cartilage and yellow elastic tissue, it is an extracellular deposit. How it is formed, how it comes to occupy the position in which it is found, is still a matter of debate. In the first place, it has never been found within the vessels; but the way it is deposited outside the vessels suggests a discharge from the blood. It is most reasonable to assume that one of the eventual constituents, upon diffusing out from the blood, meets with the other outside the capillary walls, and, combining, amyloid is produced. Eden suggests that in local amyloid the lymph vessels and channels play a corresponding part, the conveyance of the one constituent being by the lymph.

Yet another advance in our knowledge of amyloid during the last decade has been the determination that it can be produced experimentally in various animals. The experiments are not always successful, but

¹ Arch. f. exp. Path. u. Pharm., 40 ; 1897 ; 196.

DEGENERATIONS AND INFILTRATIONS

more particularly in hens and rabbits repeated inoculations of sublethal doses of attenuated pyogenic organisms, or, again, of the sterile fluids of growth, or toxins of pyococci, diphtheria bacilli, etc., will, in a certain proportion, eventually produce amyloid deposits. Pease and Pearce¹ have noted its not uncommon presence in the organs of "antitoxin horses." In the hen the deposits have been found developing as early as ten days after inoculation. It is in the spleen that these experimental deposits are first noticeable. Nor is it only bacteria and their products that initiate the infiltration; it is developed after inoculations of turpentine. Turpentine, it may be noted, is capable of giving rise to aseptic suppuration, and this association of pus-producing organisms and turpentine might suggest that leukocytic disturbances are factors in the process. For a time this was held by certain investigators, who called attention to the existence of globules reacting with iodine in the leukocytes in cases of suppuration, suggesting that these were the intermediate stage between glycogen and amyloid. Our fuller knowledge of the chemical nature of the latter has demonstrated that there can be no such relationship, nor has investigation shown that the leukocytes play any part in the conveyance and deposit of amyloid matter.

To sum up, the indications are (1) that amyloid material is allied to, but not identical with, certain compounds of chondroitin-sulpluric acid and protein found normally in cartilage and yellow elastic tissue; (2) that the protein constituent differs from that found in the abovementioned tissues, but, also, judging from analysis, exhibits not a little variation in amyloid obtained from different tissues and cases; (3) that presumably amyloid, as such, is not conveyed by the blood or lymph, but is the result of local interaction between a chondroitin-sulphuric acid moiety (brought by the blood or lymph?) and a modified local protein moiety.

ELASTOID, OR VITREOUS DEGENERATION OF THE ELASTIC TISSUES.

From the main mass of hyaline conditions we have thus separated three distinct changes, the mucoid, the amyloid or chondroid, and the colloid. There is still left a residuum which it is the custom to speak of as the hyaline degenerations proper. This residuum is characterized by affording no specific staining reactions recognized up to the present moment; save a not very characteristic coloration with acid aniline dyes; nevertheless, it embraces conditions of widely different origin, and the time has come to attempt a further separation.


The terms elastoid, or vitreous degeneration of elastic tissue, are employed by my colleague, Dr. Goodall,² to distinguish one of the most striking members of the group of hyaline changes. Sections through

⁴ Journ, of Inf. Dis., 3:1906:619. See also Lewis, Journ. of Med. Research., N. S., 10:1906:449.

² Studies from the Royal Victoria Hospital, Montreal, ii: No. 5: 1910,

ELASTOID DEGENERATION

more particularly the inner third of the wall and beneath a previous placental site in the uterus of a multiparous woman are apt to present extraordinary convoluted clumps and masses of hyaline material. They are in intimate relationship to the vessels, and, as demonstrated by Pankow and Ssasz-Schwarz, are involution products, associated with the reduction in caliber of the uterine vessels, which during pregnancy undergo a huge dilatation. The process of reduction is very remarkable. So great is the distension of the vessels and coincidently the increase in the substance of their walls that evidently contraction down to the previous diameter is an impossibility. The arteries adapt themselves to the lessened demand for blood by building what may be an entire new artery or part of the same within the lumen of the old. In this way, in the woman who



Elastoid degeneration: Section of small artery from beneath placental site of uterus exhibiting subinvolution. The patient suffered from renal and hepatic disturbances, and died seven months after delivery. Section stained by Weigert's elastica and Van Gieson's stains, (Dr. Goodall.)

Vitreous hypertrophy of elastica interna at a_i vitreous degeneration of the same at b_i at c_i transition from stained hypertrophic to unstained degenerated elastica; d_i lumen of the new vessel surrounded by irregular new nuevalar and intimal tissui; at c_i remains of old media with hypertrophy of its elastic fibres; outside the degenerated elastica interna at f the atrophy of the media is more extreme.

has borne many children, there may be traces of even as many as five vessels developed, one within the other. Studying a succession of cases at different periods after pregnancy, Goodall was able to trace the steps in the process. These, in the main, involve the internal elastic lamina. Shortly after parturition it is possible to encounter arteries in which in part the wavy internal elastic lamina is fairly normal, but following it around the circumference it may exhibit a swelling to ten times or more of its normal thickness. There is thus interposed between the media and

57

the swollen cedematous intima a broad hyaline band which at first takes on the characteristic reaction for elastic tissue by Weigert's stain. but later is liable to have ill-staining properties and, becoming much folded upon itself, resembles not a little the hvaline corpora candicantia of the Graafian follicles. In the earlier stages it can be made out that muscle and connective-tissue cells make their way through spaces in the internal elastic lamina, and, at first irregularly distributed through the greatly swollen intima, eventually with proliferation develop a new media and adventitia within the hyaline elastic lamina. The extent of their development depends upon the size of the older artery; the old media and adventitia undergo degeneration and absorption, so that the hvaline masses appear eventually to lie external to the arteries. The picture is further complicated by the fact that in some vessels the process only involves one portion of the internal elastic lamina, in which case the old wall is preserved on the one side and a new series of layers develops upon the other. So, also, in smaller vessels it may happen that there is not room for the full development of all the layers within the hyaline surrounding mass, in which case there may be no adventitia, and only a partial media reproduced.

Time will show to what extent this vitreous degeneration of elastic tissue is responsible for hyaloid changes elsewhere. Evidently it explains the hyaline deposits around the arteries of the ovaries after menstruation, for Sohma, from Aschoff's laboratory, has shown that the same process of redevelopment of the arterial walls occurs in them as is seen in the uterus. So, also, I have encountered it present, to a moderate degree but very definitely, in the arteries of the spleen. Hyaloid degeneration of these arteries has been noted previously by Klein and Greenfield,' who, however, placed the development between the internal elastic lamina and the endothelium. In this connection it may be pointed out that elastin chemically is closely allied to keratin and collagen bodies, which, while classified as scleroproteids. It may also be recalled that from elastic tissues, besides elastin, there has been gained a body of glycoproteid nature (p. 895).

OTHER FORMS OF HYALINE CHANGE.


Hematogenous Hyaloid (Hematohyaloid). — The type example of this form is seen in the hyaline thrombi, to be more fully studied in the second volume of this work, due to intravascular and intravital conglutination (a) of the entire erythrocytes as under the action of agglutinins, or (b) of the same after a preliminary disintegration into smaller globular masses, or (c) of the blood platelets. As a result there is developed a translucent soft solid mass without a trace of structure, obstructing or occluding the vessel in which it becomes formed.

¹ Quoted by Woodhead, Practical Pathology, 4th edit., Oxford Med. Publ., 1910; 332, who gives a figure of this condition.

HYALIN

A closely allied but probably distinct condition is occasionally seen when, instead of forming the characteristic fine fibrils of fibrin, a fibrinogen-containing serous exudate undergoes coagulation in the form of homogeneous masses or small clumps. This has been noted, for example, in connection with inflamed mucous and serous surfaces.

Hyaline Casts.—The clear translucent casts seen in the urine, and within the tubules of the kidney, are, to be honest, of unknown constitution. By one set of writers they are regarded as of the same order as the hyaline fibrinoid deposits just noted, and are supposed to result




Hyaline thrombus in dilated venule of hemorrhoid. This was perfectly homogeneous, Reichert, obj. 7a, ocular 4. Camera lucida, reduced one-third,

from the coagulation of constituents of the blood escaping into the tubules. In favor of this view is the fact that occasionally in parts they take Weigert's fibrin stain. But others attribute them to the fusion and inspissation of discharges from the epithelium of the tubules. A study of well-preserved kidney tissue from mild cases of parenchymatous nephritis (in connection with which hyaline as distinct from granular casts are apt to show themselves in the urine) frequently affords the appearance of loose accumulations of delicate spherules with a fine limiting membrane, so transparent as to be just visible, filling the slightly dilated lumina of

the convoluted tubules. These appear to be given off or to break off from the free surface of the renal epithelium, and it is held that by fusion and concentration they form the delicate hyaline casts. Where the irritation is more intense and the cell disintegration of a coarser type not passing on to fusion, the *granular* cast is produced; where still more intense with desquamation, the *cellular*. When the hyaline or granular cast is detained within the tubule, and undergoes both fusion of the globules and progressive inspissation, then the *waxy* or *colloid* cast is produced and this last may be metachromatic, taking on a stain resembling amyloid. The advantage of the latter view is that it brings into relationship the whole series of casts; its disadvantage that it does not explain the variations in the staining properties of the different members of the series.

Other Orders of Hyaline.—Of the considerable remnant of other hyaloid conditions still recognizable, the most important form is that



Hyaline degeneration of a glomerulus, from a kidney showing chronic interstitial nephritis.

presenting itself in connection with connective tissue. although possibly one and the same process involves cells of different orders. Thus, a common seat of such hyaline change is in the heart muscle in cases of so-called chronic myocarditis, in which scattered irregular areas of hvaline matter, transparent and with rare shrunken nuclei, are interposed in the mass of still unchanged muscle tissue. It is usual to regard this as the product of a replacement fibrosis, necrobiosis of the muscle fibres supplied by a particular

branch of the coronary artery being followed by absorption, with growth of new connective tissue; and as malnutrition in the first place led to the necrobiosis, so now the same malnutrition is held responsible for the swollen translucent condition of the fibrous tissue. Studying relatively early cases of *myomalacia* (softening of areas of the ventricle through arrest of the circulation in a branch or branches of the coronary artery). I have repeatedly been impressed by the fact that these may present regions in which the faintly and finely granular shadows of dead muscle fibres, whose nuclei no longer stain, pass almost imperceptibly into the completely hyaline areas. It is difficult not to conclude that by a process of the order of coagulation necrosis (p. 984) the nuscle cells contribute to the subsequent hyaline formation. We may hazard the opinion that the corpus candicans stage of the Granafan follicle repre-

HYALIN

sents a transformation of the same order. Nor is it uncommon to find a hyaline conversion of the tubules of the old or diseased testis, while a similar hyaline necrobiosis of the cells of several orders of tumors away from the nutrient vessels of the same, brings about the development of the *culindroma*.

Nevertheless, it is the connective tissue framework of tissues that is peculiarly apt to be affected—of the thyroid, and the kidney. Scar tissue is liable to be involved, as also the organized fibroid deposits on serous surfaces, which may attain a notable thickness (half an inch and more) and a porcelain-like appearance (*hyaloscrositis*). Similarly the new connective tissue growths of tuberculous and syphilitic granulomas may exhibit hyaline change.

Another well-marked group of hyaline conditions is met with in connection with the capillaries. Either their walls become converted into thickened hyaline tubes, or the whole capillary is changed into a solid, impervious hyaline mass. The former is occasionally seen in the lymph

follicles, in the brain and in the thyroid, the latter in old areas of interstitial nephritis, here involving the glomeruli, these becoming converted into small homogeneous glassy nodules (Fig. 302). Such changes recall amyloid disease, and as a matter of fact cases have been recorded in which the deposits have in part given the amyloid reaction, in part acted like hyaline. In advanced amyloid we may find the basement membrane of the renal tubules presenting hyaline meta-



Hyaline degeneration of the membrana propria of two renal tubules with loosening of the epithelium, (Ribbert.)

morphosis, while in the experimental production of amyloid more than one observer has noted that material giving the amyloid reaction is preceded by deposits of hyaline type, not affording the same.

Intracellular Hyaline.—More particularly in cancers we are apt to encounter small accumulations within the cells, either globular or of irregular shape, having the appearance and reactions of hyalin. These have been studied more especially by Pianese and Fabre-Domergue. From these studies it is evident that they differ to some extent among themselves in chemical composition, for they do not uniformly take on the different stains; some take fuchsin with considerable intensity (Russel's fuchsin bodies). These may occur in chronic inflammations and in enlarged lymph glands, as well as in cancers; and, through degeneration of the cells, may come to be extracellular. Other deposits of a somewhat similar appearance in epithelial cells may be of keratinous nature. While calling attention to these, it has to be admitted that they occur so sparsely that anything of the nature of an exact chemical study of their constitution is out of the question.

PATHOLOGICAL KERATINIZATION.

We occasionally encounter either excessive development of the horny layer of the skin (as in the cutaneous horns which may develop from various parts of the body and conditions of hyperkeratosis and ichthyosis), or the presence of horny layers in regions which normally exhibit no horny layer. The mouth, cesophagus, and vagina, for example, have what may be termed squamous mucous membranes, exhibiting no keratinous change of the outer cells; but, under conditions of irritation, the tongue may present keratinized processes of the epithelium (hairy tonque), the vagina show horny ridges (pachydermia), the resophagus exhibit longitudinal bands of leukoplakia (notably in alcoholics). Similarly the mucous membrane of the middle ear may give origin to a dense accumulation of keratinized cells of pearly appearance (cholesteatoma). The same process may show itself in the pelvis of the kidney and the urinary bladder. The pia mater may also be the seat of the tumor-like cholesteatoma, the development in this last region being due to epidermal inclusions during development. Keratinization may occur in tumors proper, leading to the epithelial pearls of epitheliomas, present even when normally the epithelium is devoid of keratin. In regions where through metaplasia squamous epithelium may replace columnar cells, there both the metaplastic epithelium and malignant growths developing from the same may show keratinization, although rarely so clearly defined and pronounced as where the growths originate from a naturally squamous epithelium.

Keratinization is under normal conditions a form of physiological degeneration associated with necrobiotic changes in the cell. As the cells originating from the mother-cell palisade layer of the epidermis are pushed outward and become further removed from the vessels of the corium by the development beneath them of new generations of daughter cells, they are at first free from cytoplasmic granules, but when they reach the level of the stratum granulosum they exhibit fine granules of keratohyalin. These stain blue with hematoxylin. Passing farther out into the stratum lucidum they become translucent and evenly diffused throughout the cells (eleidin). Farther outward they exhibit a second development of granules, presenting fine granules of keratin (taking a blue color with Gram's stain). The cells become increasingly flattened and shrunken, their nuclei no longer stain, until, finally, flattened scales of compact keratin, staining vellow with Van Gieson are all that represent the original cells. The fully keratinized cell is dead, and the cholesteatomas, cutaneous horns, and other massive accumulations are not therefore hypertrophies, but abnormal collections of matter not properly cast off. At most the underlying rete Malpighii may show hypertrophy and more active cell development.

The opposite condition of *Parakeratosis* or deficient development of keratin is encountered in atrophic conditions of the skin from inflammation of the corium, in psoriasis, etc.

It may be recalled that keratin is, like elastin, and collagen one of the scleroproteins.

¹ In this rapid review of Keratinization I largely follow Gierke.

CHAPTER XXVIII.

THE DEGENERATIONS AND INFILTRATIONS-(CONTINUED).

Disturbances of the Aqueous, Fatty, and Carbohydrate Contents of the Cell.

HYDROPIC DEGENERATION.

It will be remembered that attention was called to the fact that in cloudy swelling there is a definite increase in the watery contents of the cell. What would seem to be a further stage in the same condition, and one associated with yet graver disturbances in the cell, is the appearance of definite vacuoles in the cytoplasm, containing a watery fluid, which vacuoles may attain so great a size that the cell undergoing disorganiza-

tion bursts, and, with its neighbors, becomes represented by a vesicle visible to the naked eve.

The most striking example of this hydropic degeneration is met with in the lower layers of the epidermis in cases of smallpox; the vesi-ular stage of the pock is essentially brought about by the acute hydropic swelling and disintegration of neighboring cells. Experimentally, a similar condition can be induced in the cells of the convoluted tubules of the kidney by the exhibition of cantharidin.


This rapid imbibition and accumulation e^{*r*} a fluid in a cell can, upon physical grounds, have only one explanation. The constitution of cyto-



Hydropic degeneration: epithelium from a smallpox papule. The epidermal cells greatly swollen, distended by large vacuoles. X 300. (Ribbert.)

plasmic matter, as also of the nucleus, is colloidal, and colloidal membranes (for such we can regard the surface layers of cells) have characteristic properties. They hinder the diffusion of crystalloid molecules to a considerable extent. Although animal cells possess no well formed outer membrane (as do plant cells), we are led to believe that in animal cells a fine layer of similar nature acts physiologically as such a membrane. We therefore conclude that the essential cause of hydropic degeneration is some dissociation of the complex colloid material of the cytoplasm, whereby, either by cleavage or ionization, crystalloid bodies make their appearance in the cytoplasm. As an illustration of conversions

of this order, it may be noted that the peptones, leucin, tyrosin, etc., which are the products of the breaking down of (colloidal) proteins, are of distinctly crystalloid nature. So long as such products are present within the cell body in greater concentration than they exist in the surrounding medium, there will be a tendency to osmotic diffusion inward of watery fluid until such time as the osmotic pressure on the two sides of the membrane becomes equal. In other words, the cell swells up and becomes hydropic.



Serous atrophy of fatty tissues of neck of pig after prolonged feeding with fat-free diet; A, fat cells which have undergone serous atrophy (contents not stained by osmic acid); B, capillaries; C, bundles of elastic fibres. (After Herter.)

Vacuolar Degeneration.—A condition which may possibly be closely akin to the hydropic degeneration is vacuolar degeneration, in which isolated vacuoles of medium size make their appearance in sundry cells, and, it may be, actually within the nuclei. The condition has been noted more particularly in connection with the voluntary and cardiac muscle fibres, and the ganglion cells of the central nervous system. Thus, in typhoid fever, oval and spindle-like vacuoles have been observed within well-striated muscle fibres; vacuolation of the heart muscle cells has been observed in numerous conditions, and Nerlic and others have called attention to similar vacuoles in the ganglion cells of nerve centres in tetanus and acute infections. Hodge has shown that strong electrical stimulation brings about diminution in the size of the ganglion cells, with

FATTY INFILTRATION

the development of numerous vacuoles. It is suggested that in all these cases there is scrong stimulation of the nerve cells with increased dissociation of the cytoplasm, and that here, as in the case of cloudy swelling, the products of dissociation lead to osmotic absorption of increased fluid.

Serous Atrophy.—This is seen in wasting diseases affecting more particularly the epicardial and perirenal fat deposits; in place of the normal fatty tissue there appears a translucent gelatinous tissue. It occurs most extensively in the senile fatty marrow of bones, affecting the same tissues also in wasting diseases, here from the surroundings partaking clearly of the nature of a hydrops ex vacuo. It is essentially, as pointed out by Flemming, a fat atrophy, with disappearance of the fat out of the cells, so that they no longer react with osmic acid and other stains for fat. The cells, however, do not wholly shrink, the place of the fat being taken by a serous fluid, which further infiltrates the extracellular tissue.

Herter' has reproduced the condition in the pig by prolonged fat starvation, and found that he could arrest its appearance by giving a marked excess of carbohydrates.

DISTURBANCES OF FAT METABOLISM.

Of the fatty accumulations, two conditions are to be recognized, which, for want of a better terminology, it is usual to describe, following Virchew's original distinction, as fatty infiltration and fatty degeneration. Well-marked conditions of the two are strongly contrasted; there are, however, intermediate states which it is difficult to distinguish surely; the cause of this difficulty will be evident when we come to consider the nature of the processes.

FATTY INFILTRATION.

Neutral fat is a constituent of most of the tissues of the body, but this in a state in which it is not recognizable within the cells, either by simple microscopic or by microchemical means. The kidney tissue, for example, may, by all the usual microscopic methods, by osmic acid, or by Sudan III, show not a trace of fat; nevertheless by appropriate chemical means, as much as 23 per cent, of the total solids, nay be demonstrated to consist of fats.² In one tissue, however—fatty tissue—the amount present is extreme, so that the cells are distended with fat in the form of large globules; so distended that the nucleus is pushed to one side, and the cell assumes a signet-ring appearance. There are certain regions of the body in which this fatty tissue is normally present, notably in the subcutaneous connective tissue, in the

¹ Jour. of Exp. Med., 3: 1898: 293.

² Rosenfeld, Arch. f. exp. Path., 55; 1909; 179. See also Orgler, Virch. Arch., 167; 1902; 310.

omentum and appendices epiploicæ, around the kidneys (suet), in the cardiac grooves, etc. We must regard the cells accumulating the fat in these regions as normal. It is when connective-tissue cells elsewhere, more particularly in the interstices of tissues, become the seat of identical accumulation of, and distension with, fat, and assume the identical appearance, that we speak of the fatty infiltration of a tissue. Such may occur between the cardiac muscle cells, between the fibres of skeletal muscles (as in pseudohypertrophic paralysis, p. 592, Fig. 178), or, again, in the pancreas. There is yet another order of cells that becomes physiologically (as during pregnancy) the seat of notable accumulations of fat, namely, the liver cell, and this also, through the accumulation, is apt to assume the signet-ring appearance. Where fat is heaped up in the liver in this manner we also speak of fatty infiltration. And the accumulation may be extraordinary: in the fatty liver of alcoholism, Perls determined that four-fifths of the total solids, and close upon 41 per cent. of the total cell substance (including water) might be fat. It must be clearly borne in mind that in all other parts of the body fatty infiltration involves the connective-tissue cells; in the *liver*, the connective-tissue cells are not affected, but the liver cells proper. Such accumulation of visible, neutral fats in the otherwise normal



Liver cells in various stages of fatty accumulation, \times 300, (Rindbleisch.)

cells of connective tissues and the liver occurs in a variety of conditions; these may be classified:

1. *Physiological.*—As already noted there is heaping up of fat in the liver (particularly in the more central cells of the lobules) during the latter months of pregnancy and during lactation, apparently as a preparation for the latter.

2. Fatty Growths "Ex Vacuo."—Occasionally the atrophy of a part, or tissue, is accompanied by a compensatory appearance of fat cells. The most frequent

example of this nature is seen in the pelvis of the kidney in senile or other atrophy of that organ; another striking example is in pseudohypertrophic paralysis, where series of fat cells replace the atrophied muscle fibres. The replacement of the red marrow by fat cells in normal bones is of the same order.

3. Overnutrition.—More fatty matters being taken in or elaborated than can be burnt up in the performance of function. The "aldermanic" type of individual and the overfed Strassburg goose, with its "foie gras," are the familiar examples of this form of fatty infiltration.

4. Substitution (?).—Fatty infiltration, and not, as vulgarly supposed, cirrhosis, is the commonest affection of the liver to be met with in those addicted to alcohol—and this notwithstanding the fact that the confirmed alcoholic is a small eater. Two explanations have been afforded for this fact: (1) that alcohol, acting on the nerve centres, or directly on the cells of the body, lowers functional activity and oxidation, and

FATTY INFILTRATION

so the fat absorbed is not burnt up; (2) that alcohol is in itself a foodstuff capable of easy oxidation, and that it replaces more particularly the fats, so that these, not being oxidized, remain and accumulate in the liver cells. The more recent studies upon metabolism in animals treated with alcohol favor the latter view. Probably both factors must be regarded as operative.

5. Diminished Oxidation: (a) Congenital.—There are those so constituted that, despite small appetite and consumption of food, if anything, below normal, they appear inevitably to become fat; others, on the contrary, who, with ravenous appetites, remain as lean as Pharaoh's kine.

Waldvogel' has thrown light upon these phenomena. Injecting a solution of β -oxybutyric acid into the chest wall of healthily lean persons, there was a slight rise of temperature, but no oxybutyric or acetic acid appeared in the urine, nor was there increased excretion of acetone; it was completely consumed. The same dose administered to obese individuals was followed by no rise of temperature, this suggesting that the oxidation is slower, while the excretion of acetone was increased the next day, and acetone could be recognized in the breath. He concluded, therefore, that the obese are unable to oxidize the fat acids reaching the intermediate metabolism with the same intensity as do healthy persons, and that the defective transformation of the fatty acids leads to accumulation of neutral fat in the cells. It is suggestive in this connection that the administration of thyroid extract, which materially accelerates the oxidative processes of the organism, materially reduces obesity.

(b) Through Discase.—It is noteworthy that in tuberculosis there is a tendency in some to become obese (this is noted more often in cattle); in others, and this is more frequent, despite general emaciation, the liver is found at autopsy in a state of well-marked fatty infiltration. The most satisfactory explanation is that of diminished oxidation lowered vitality with lowered functional activity of the tissues, and, as a consequence, lessened burning up of the fats taken as food. Along with this we have to recognize a certain amount of transposition of fat from the normal stores in the subcutaneous and other tissues to the liver.

Here may be noted a rare condition in infants, due apparently to a congenital incapacity to transform fat, rather than to any excess of fat in the diet, in which the liver attains a great size. Not a single liver cell may be seen which is not distended by a large globule of fat. Save for the regularity of the vessels the sections almost resemble those of a lipoma.

FATTY DEGENERATION.

In this condition we deal with what is *primarily a cell degeneration* —the deposit of fat accompanying and being the result of depressed

¹ Epstein Festschr., Deut. Arch. f. klin, Med., 89; 1906; 342.

and damaged cell activities. The cell nuclei exhibit marked indications of chromatolysis and degenerative changes; the cytoplasm becomes filled with minute, dust-like fatty globules, so that, stained with Sudan III, or Scharlach R, the whole cell body takes on diffusely the characteristic orange-reddish color, the high power demonstrating that this is due to abundant minute fatty dots.

The long-established test for the presence of fats, osmic acid, is now recognized to be imperfect. This blackens the globules of oleic acid and its compounds, but has no effect upon palmitin and stearin compounds. Among the oleic acid compounds it affects "myelin," not merely turning the myelin globules gray, as Kaiserling and Ogler point out, but eventually a deep black (apparently through the gradual dissociation of the oleic acid constituent). This is true, also, we find with lecithin (Reidel's lecithol). Sudan III, on the other hand, is taken up by and so stains all fats present in the globular form within the tissues, and exhibits a differential stain for soaps (Fischler, confirmed by Klotz). Scharlach R has similar properties. Both are employed in weak (60 per cent.) alcoholic solutions, and the staining is due to the fact that fats dissolve these dyes more readily than does the alcoholic solvent.¹

The tissues especially liable to be affected are (1) those liable to exhibit cloudy swelling (gland cells, more particularly those of the liver and kidney and muscle fibres, particularly those of the heart); (2) endothelial cells of bloodvessels; (3) certain cells undergoing normal retrogressive changes (cells of the sebaceous glands).

There is histological evidence that the substance composing these finer granules or globules is not, in some cases at least, of the same nature as that present in fatty infiltration. Thus, in a study of fatty degeneration of the liver, shortly to be published, McCrae and Klotz call attention to the fact that we deal frequently with polygonal or granular deposits of a subcrystalline type rather than with spherical globules.

Intermediate Conditions.—The accumulation of fat in the form of small globules, without pronounced nuclear degeneration, occurs normally in several orders of cells, namely, in the cortical cells of the adrenal, and in the hypertrophied plain muscle cells of the uterus during the process of involution following upon parturition, as also in the decidual cells of the fourth month (Holsti⁷). Even in what had hitherto been recognized as ordinary fatty tissue Shattock³ points out that certain areas (the "fat-masses" of the necks of hedgehogs, cretins, and young children and the subpleural fat in infants) present fat cells of the glandular type with multiple discrete globules. Contrariwise, conditions may be noted in which, with well-marked evidence of nuclear degeneration, there accumulate in the cell large fatty globules along with small. This

¹ For the chemistry of the staining of fats and lipoids see Lorrain Smith, Journ. of Pathol., 12: 1907: 134, and Dietrich in Lubarsch and Ostertag's Ergebnisse, 13: 1909; pt. 2: 295.

² Anat., Heft 37; 1908; 179. ³ Proc. Roy. Soc. Med., 2; 1909; Path. Sect. 252,

FATTY DEGENERATION

in phosphorus poisoning affecting the liver. There are thus, histologically, intermediate states between well-marked fat infiltration and fatty degeneration.

Herxheimer's observations upon the thymus indicate yet another intermediate stage, which may, however, be closely allied to that seen in the uterine muscle fibres.¹

According to Gegenbaur, the thymus continues to grow in size until the end of the second year. Herxheimer found that even before the end of the first year this organ contains fat in abundance. This is in the form of fine granules tending to coalesce, in part in the lymphocytes, which form the main constituents of this organ, and here more

F1G. 306	F16, 307

Fatty degeneration of liver cells: b_i fresh cells, eloudy and granular, nuclei not clear; a_i the fine, fatty globules seen more clearly after treatment with acetic acid. (Ribbert.)

Fatty degeneration of heart-muscle fibres, showing different grades of involvement of the individual fibres; fresh specimen, (Ribbert.)

particularly at the periphery of the follicle; in part, in the connectivetissue cells of the stroma. Here it was in greatest abundance in the connective-tissue cells around the capillaries and smaller vessels. This clearly is not a degeneration proper; it attacks the cells at a period of active growth. Later the fine granules run together, and then are found the large fat cells characteristic of the thymus in its later retrogressive stage. It may be here noted that the levator palpebræ superioris, one of the most active muscles of the body, contains normally abundant fat, in the form of fine globules; here, again, we cannot deal with a degeneration. Lastly, Shattock and Dudgeon² have noted that in chlorotic and secondary toxemic conditions the polymorphonuclear leukocytes of the circulating blood exhibit fine fatty globules, staining

¹ Verhandl, d. deutsch, pathol. Gesell., Jena, 1904; 258.
 ² Proc. Roy. Soc. Biol., 79; 1907; 427.

with Scharlach R, without there being any sign of nuclear disturbance (although such may be present along with fatty change in pus cells). Cesaris-Demel has recorded similar observations.

Gausation.—These intermediate cases we will discuss later; leaving them aside for the moment, and considering only the typical cases, as regards the condition leading up to the degeneration, two groups of cases may possibly be distinguished: (1) Those in which the fatty degeneration follows upon cloudy swelling, and so may be regarded as the second stage in parenchymatous inflammation of organs, and (2) simple uncomplicated fatty degeneration—(a) physiological, and (b) pathological.

Fatty degeneration of the first order is apt to accompany all severe fevers-pyemia, septicemia, the acute exanthemata and typhoid; the cause is clearly identical, namely, bacterial intoxication. Subjection to high temperature equally produces cloudy swelling followed by fatty degeneration. As regards the second order, the physiological fatty degeneration seen in the cells of sebaceous glands, as, again, in the cells of the mammary glands during lactation, is a somewhat remarkable process. In the former case there is a constant multiplication of the gland cells proper; of the two cells, the products of an act of multiplication, the outer, which we may term the daughter cell, becomes filled with fatty globules of fair size (larger than those seen in parenchymatous inflammation), but much smaller than those of fatty infiltration, and uniformly scattered through the cytoplasm. With this the nucleus becomes paler and shows evidences of chromatolysis. Eventually the cells of this order become liberated and break down, and, as a result, a fatty emulsion fills the lumen of the gland.

In the mammary gland a parallel condition has been observed.

The nucleus of the gland cell multiplies by direct division until two or three nuclei are present in the cell. Next, the nucleus nearest to the lumen undergoes chromatolysis, and it can be observed that its chromatin passes into the surrounding protoplasm. With this there is not true cell division, but in the outer portions of the cell fine fat droplets collect, and this outer portion, containing the degenerated nucleus, is discharged into the lumen, giving rise there to the fatty globules of the milk. Nissen¹ calls attention to this breaking down of the nucleus and disintegration of the phosphorus-containing nucleoproteid, and the characteristic presence in milk of a phosphorus-containing protein, namely, casein.

In man the chromatolysis of the nucleus in these glands is very obvious; according to Altmann² the process is somewhat different in the sebaceous glands of the inguinal folds of the rabbit (where they are very abundant). The process here is more of the nature of a secretion. The cells are filled with granules, which, in the centre around the nucleus, do not react with osmic acid, but show transition toward the periphery into the fatty granules, which, discharged, fuse into definite globules.

¹ Arch, f. mikr, Anat., 26: 1886: 337,

² Die elementäre Organismen, 1890.

.910

One of the most recent studies upon this subject is by Professor Arnold.1 He admits that the first signs of fat in the cell show themselves in the immediate neighborhood of the nucleus, and regards this as an indication that the formation of milk fats is a synthetic process, in which the nucleus takes a part; also, that amitotic nuclear changes may occur during the process, but lays down very definitely that the process of fat accumulation and discharge may proceed without nuclear disintegration; nay, more, may be of the nature of a selective absorption and secretion from the lymph. He has pointed out² that sundry tissues placed in a soap solution present in their cells a granular deposit of fat globules and in this connection may be noted Jacobsthal's observation³ that the Scarlet R dueing the fats given as food may be detected in the milk of those animals.⁴ Arnold holds that these absorbed fats become associated with the cell plasmosomes.

There are thus divergent views regarding the nature of milk secretion, but, evidently, the cells of the mammary gland do not absorb and excrete the droplets of fat as such; the process is much more complicated; the fat is absorbed in a soluble form, the process of converting it into neutral fat is accomplished by intracellular enzymes, and the production and activity of these enzymes is accompanied by using up and eventual disorganization of the nuclear and cytoplasmic material.

Simple *pathological* fatty degeneration of the second order occurs (1) in certain cases of acute non-bacterial intoxications-by arsenic, antimony, bismuth, carbon-monoxide poisoning, mineral acids, pyrogallic acid, chloroform, phloridzin, etc., and (2) in conditions of malnutrition, notably in certain anemias (pernicious anemia, advanced chlorosis, and cachexias, and the anemia following severe hemorrhages), as, again, in the later stages of starvation. With these, although, as above noted, the appearances are intermediate, must be included the pronounced fatty degeneration of phosphorus poisoning.

We do not pretend that all these intoxications produce fatty degeneration not preceded or complicated by cloudy swelling. This point has not been sufficiently studied. We are inclined to believe that fuller study will demonstrate a preliminary or accompanying cloudy swelling in many of these cases, as in starvation and in phosphorus poisoning; but this is not marked, and in certain conditions, as in chloroform poisoning, there is no evidence of cloudy change, although it is true that in the neighborhood of the nucleus non-fatty granules, the socalled plasmosomes, make an appearance.

1 Ziegler's Beitr., 38: 1905: 421.

² Zentralbl. f. Pathol., 14: 1903; 785. ³ Verhandl, deutsch, path. Gesell., 13:1909:

⁴ I would suggest that the two sets of observation are not necessarily contradictory. Thus, in the mammalian pancreas there is no suggestion of nuclear multiplication in connection with the formation of the zymogen granules; nevertheless, studying recently along with Professor Downey, of Minneapolis, certain sections of the pancreas of the ganoid fish, Polyodon spathula (the spoonbill), which he had prepared, we found what appeared to be various stages of double nucleus with the disintegration of one of the two in intimate association with the appearance of zymogen globules.

Etiology.—Fatty Infiltration.—What are the underlying causes of these two conditions of fatty infiltration and fatty degeneration? Regarding the former there can be no question as to the origin of the fat; it is storage fat, accumulated in the cells, either as the result of an intake of food material affording neutral fats over and above the capacity of the tissues to oxidize, or of oxidative capacities of the tissues below the normal, so that foodstuffs reach the stage of neutral fat, but do not forthwith pass beyond that.

But even in this case, we would repeat, the process is not that of simple taking up of already formed neutral fats from the blood and lymph. Neutral fats-glycerides of the fatty acids-do not exist as such in the fluids of the body under normal conditions; only in pathological conditions, as in diabetes, advanced alcoholism, and some cases of arteriosclerosis, do we encounter lipemia or an emulsion of fine fatty droplets in the blood. In what form they exist there is a matter of debate. The more recent studies upon immunity point definitely to the formation of loose compounds between fats and the proteins of the plasma (more particularly the globulins), and pointing in the same direction are the observations of Wilson and Williams¹ that in diabetes we deal with a lipoidemia rather than a lipenia. The laving down of neutral fats in the cells necessitates, therefore, a dissociation and a subsequent combination of fatty acids and glycerin; and this, it has been demonstrated, is accomplished by the agency of intracellular enzymeslipases (see p. 83). Nay, more, that the nucleus of the fat cell is concerned in the process, is indicated by the remarkable presence of a vacuole within it. We do not see vacuoles in any other normal cells of the human organism, and that the vacuole is related to the deposition of neutral fat is indicated by Shattock's² observation that it reacts with Sudan III, i. e., is of a fatty nature. Similar fatty vacuoles have been recorded in the nuclei of the other potential fat cells, namely, the liver cells in cases of cirrhosis³ and other conditions.

Fatty Degeneration.—The long-accepted view was that fatty degeneration is, as the name implies, the result of a breaking down of the cell substance, with liberation of the nitrogen-containing element of its proteins, and retention of its carbon-containing moiety, and conversion of the same into fat.

Many arguments were adduced in favor of this view: the cells were seen clearly to be undergoing disorganization; in conditions favoring fatty degeneration, the N. output was found increased, the CO₂ output

¹ Biochem. Journ., 2: 1907; 20. Fugoni and Marchetti, however, in an extreme case found that the main bulk of the ether extract consisted of neutral fats (21.98 per cent. of total blood), although fatty acids and soaps (3.45 per cent.), cholesterin (1.06 per cent.), and lecithin (0.5 per cent.) were also present (Berl, klin, Woch., 1908; No. 44). Klemperer and Umber confirm Wilson in finding in a long series of diabeties that the main increase is in cholesterin esters and phosphatides (Zeitschr. f. klin, Med., 61: 1907; 145; see also Adler, Berl, klin, Woch, 1909; No. 31).

² Trans. Path. Soc. Lond., 54: 1903: 215.

³ Brandts, Ziegler's Beiträge, 45 : 1909 : 457.

FATTY CHANGES: ETIOLOGY

diminished;1 there is an actual increase in the fat in the fatty degenerated liver, even in starving animals;² and this fat has been regarded as formed at the expense of the carbohydrate constituents of the cell, for, in phosphorus poisoning, with increase in the fat, there is notable absence of glycogen (Stolnikow and others). There is evidence that fats are capable of development from proteins. F. Hofmann³ demonstrated that the larvæ of the fly, Musca vomitoria, grown from eggs placed on ox blood containing a known quantity of fat, contained considerably more fat than was present in the control eggs and the ox blood combined. Similarly, Burdach⁴ found that in the development of the eggs of the snail Limnaus stagnalis, the fat increases, it may be three- or fourfold. Pettenkofer and Voit,⁵ feeding dogs on meat free from fat, determined on analysis that the C, was retained in the organism in the form of fat. Hoppe-Seyler determined that, upon keeping, the fat of milk increases, the casein diminishes. The formation of adipocere in the corpse was explained by Virchow⁶ along these lines, namely, of conversion of the proteins of the corpse into fats.

Much of this evidence has been discredited or put on one side as not bearing upon the case in point. It has been found, for example, by numerous observers, that, while the fat in the liver may be increased, the total fat of the body is not increased, but may be definitely diminished in cases of fatty degeneration.

This was well demonstrated by A. E. Taylor,⁷ who, taking two series of frogs, one as control, the others, in which he had induced fatty degeneration, killing and desiccating them, and then extracting the total fats, found that there was an actual loss, and not a gain, of fats.

Kobert has noted that, while in the living animal phosphorus easily sets up fatty degeneration of the cardiac muscle, if the removed ("überlebendes") heart be taken and transfused with fluid containing relatively enormous doses of phosphorus, not the slightest trace of fatty degeneration is to be made out. More recent and exact studies have shown that Pettenkofer and Voit's observations are valueless, from the fact that meat, which, to the naked eye, is free from fat, contains, nevertheless, a very considerable proportion; their dogs were fed with fat. The most convincing series of experiments are those of Rosenfeld,⁸ Rosenfeld demonstrated, in the first place, that if a starving animal be poisoned with phosphorus, the accumulation of fat in the (fatty degen-

[†] Fränkel and Gebbert, Centralbl. f. med. Wissensch., 21:1883;583 (abstr.); see also Bauer, Zeitschr. f. Biol., 7:1871;63.

⁴ Diss. Regensburg, 1853.

² Stolnikow, Du Bois-Raymond's Arch., 1887, Suppl. Bd. 1.

³ Zeitschr. f. Biol., 8: 1872; 153.

⁵ Liebig's Ann., 1862, Suppl. Bd. 2, 52, and 361.

⁶ Würzburger Verhandl, 3:1852.

⁷Journ. of Exp. Med., 4: 1899: 399 This has been confirmed by Kraus and Sommer with mice poisoned with phosphorus. The total body fat might be reduced to one-half the normal, and of this, one-third to one-half might be present in the liver (Hofmeister's Beitr., 2: 1902: 86).

⁸ Verhandl. d. deutsch. path. Gesellsch., 6:1904:71.


erated) liver is accompanied by a corresponding diminution of the fat elsewhere in the organism, in the skeletal muscles, for example; and, secondly, that if a dog be poisoned with phosphorus or phloridzin, and coincidently fed with a foreign fat, such as tallow (mutton fat), in which the relative proportion of palmitic, stearic, and oleic acids are widely different from those present in dog fat, the composition of the fat obtained from its "fatty degenerated" liver approximates to that of the foreign fat. The same is true, according to Schwalbe, when it is fed with the patent fatty preparation known as iodipin, although Wells¹ could not confirm.

It is obvious, from these experiments, that the bulk of the fat making its appearance in the liver cells in these experiments is absorbed, and



Double contoured myelin bodies of irregular rounded shape with processes. (Perls.)

is not the product of the breaking down of the cell cytoplasm, and that in fatty degeneration what we have to deal with in the main is a translation of fat in the organism from the fat cells and customary fat deposits of the organism to the liver, and, as Leick and Winckler have



Juice expressed from adrenal cortex, seen under crossed Nicol's prisms, showing isotropous, fatty globules and anistropous myclin globules (with black cross).

shown, to the myocardium (for in the heart-muscle fibres there may be found the same accumulation of foreign fat), and Löhlein and Landsteiner and Mucha (contrary to Rosenfeld) show an increase in fat in the kidney also. The other organs of the body, with the exception of the pancreas, lose their fat. The experiments appear to be so convincing that there has been a movement to replace the terms "fatty infiltration" and "degeneration" by "physiological" and "pathological fat infiltration," respectively. Nay, more, Ribbert² would hold that a distinction can scarce be made between the infiltration of normal cells and the degeneration of morbid cells; that the size of the globules is no guide.

¹ Zeitschr. f. physiol. Chemie, 45: 1905: 412.

² Sitzungsber, d. Gesellsch, zur Beforder, d. ges. Naturwiss., Marburg, 1902; No. 4,

MYELINIC OR LIPOID DEGENERATION

Lipoid Degeneration.—Already, however, there is a reactive movement; the matter is seen not to be so simple. We have to take into account the existence of what, to distinguish it from pathological fatty infiltration, I would term lipoid degeneration,¹ or what Kaiserling and Ogler have termed *myelinic metamorphosis*.

Attention has been called to this more especially through the study of autolysis. If liver, kidney, or muscle tissue be removed from the organism, and placed for twenty-four hours in the incubator at 37° C., under strict aseptic precautions, it is found that the cells now contain abundant irregular doubly contoured globules or granules, which swell up with water, undergoing change of shape, which may be doubly refractive, and are soluble in ether and alcohol. They possess, in short, the properties of the substance or substances to which, in 1854, Virchow directed attention, and, from their resemblance to the brain marrow and its properties, termed myelin. These, then, are myelin bodies.²

As to the chemical nature of myelin, there has been abundant debate. Virchow was not sure that he dealt with a single substance; the elder Beneke regarded it as of the nature of cholesterin compounds; Liebreich held that the protagon which he had isolated from brain substance must be present to afford the myelin reactions. Quincke pointed out that these reactions are afforded by many substances, among them the simple soaps. Recent of servers have largely favored the hypothesis that they are of the nature of lecithin, or, more accurately, of phosphatides, compounds of the nitrogenous base cholin, with glycerophosphoric acid, and two atoms of fatty acid, one of which, according to Thudichum, must be oleic acid. There are those who, like F. Müller, still favor the protagon hypothesis; and others, like Aschoff, who favor the cholesteryl compounds. The study made by Professor Aschoff and myself³ of the physical properties of the myelins leads us to the conclusion that the only known substances which have the property of forming doubly refractive "fluid crystals" at room temperature (as have the myelins) are compounds of oleic acid. The indications are that several compounds are present in pathological conditions-and some physiologicalwhich can form doubly refractive globules, and, with water, swell up into bizarre shapes-cholesteryl oleate, cholin oleate, and phosphatides: and that, therefore, in brief, the myelin bodies of the organism are one and all lipoid bodies-allied to the soaps-in which the fatty acid essentially concerned is oleic acid.

¹ If the term *myelin* is to be retained—and we have doubts as to its utility—then the property of double refraction must be regarded as an essential attribute of bodies to which the name is applied. Now this property is not always demonstrable in the tissues; wherefore, until the chemistry of these bodies is worked out more fully, it is better to employ this more non-committal term in place of the *myelinic degeneration* of the previous edition.

 $^2\,{\rm For}$ a full review of the literature of the myelins, see the article by Schultze in Lubarsch and Ostertag's Ergebnisse, 13:1909: pt. 2:253 to 281.

³ Proc. Roy. Soc. Lond., B., 78: 1906; 359. See also Aschoff, Verhandl. deutsch. pathol. Gesellsch., 10: 1907: 166, and Adami, Harvey lectures, 2d series; 1908: 117,

As to the myelin developed in autolytic processes, indications all point to its being of the nature of a phosphatide. Analysis shows that in the early stages of autolysis there is a pronounced increase in the lecithin from the liver and other organs. This may, in the liver, be as high as 15 per cent. of the solids of the liver at the end of twentyfour hours' autolysis, according to Waldvogel' and Dietrich. With prolonged autolysis, the lecithin undergoes marked diminution, with corresponding increase in the fatty acids, neutral fats, and cholesterin. Clearly there is dissociation of the lecithin with appearance of simpler fatty bodies and cholesterin. These observations of Waldvogel are not accepted by all, but if for "lecithin" we substitute "phosphatides," and among these include *jecorin* (p. 92) then the statement is accurate.

Whence are derived the phosphatides in these cells? A considerable proportion is present in the normal cell. It is suggestive, in the first place, that a characteristic constituent of phosphatide is glyccrophosphoric acid; another, the nitrogenous base, cholin. Now, the one prominent protein or nitrogenous compound, or group of compounds, in the cell which contain glyccrophosphoric acid is nuclein; or, more accurately, the nucleus—and these, it has been demonstrated, are associated with the nuclear chromatin or stainable material.

It is more than suggestive that, as pointed out by Albrecht and Dietrich, coincident with the appearance of the myelin granules in the cell undergoing autolysis, there is solution and disappearance of the nuclear chromatin. In other words, a glycerophosphoric acid compound appears in the cytoplasm coincidently with the disappearance of such compound from the nucleus. It would seem, therefore, that lecithin-like bodies are primarily derived from the nucleus. Whether this occurs from simple splitting off has not yet been determinedwhether, that is, the nuclear chromatin has combined in its molecule certain fatty acid groups, or whether, in dissociation, the nucleinic cholin and glycerophosphoric acid unite with fatty acids present in the cytoplasm, or dissociate the fatty acid from its previous state of combination in the cytoplasm. The very abundance of phosphatides in the normal liver cell would suggest that the latter is the more likely.² Indeed, the whole trend of modern studies upon the physiological chemistry of the fats is opposed to the view that any fatty bodies are formed directly from the proteins-a very different matter from the existence of compounds between fats and proteins or the disintegration products of the same. This, at least, is certain, that in the cell as a whole, as demonstrated by Hildesheim and Leathes,3 after autolysis for three

¹ Münch, med. Woch., 53 ; 1906 ; 402.

² Hefter (Arch. f. Exp. Pathol. u. Pharm., 28: 1891; 97) found that half of the fatty bodies in the normal liver are in the form of lecithin; Dunham (Berl, klin Woch., 1904; 750) found similarly that from 30 to 70 per cent. 5f the kidney extract is lecithin.

^a Jour. of Physiol., 31: 1904; Proc., p. 1. See also Leathes, Arch. f. exp. Pathol. u, Pharm., Supplement-Band (Schmiedeberg Festschr.), 1908; 327.

days, there may be from 10 to 40 per cent, more recoverable fat than from the fresh organ. Whether here we deal with a synthesis of fatty acid from glycogen (which undergoes diminution) or a dissociation of fatty acid from relatively firm combination with proteid matter, or both, is not determined. In favor of the latter view the observations of Gies and others indicate the existence of compounds between proteins and fatty acids, as, again, do the observations of Dormeyer,⁴ that the "fixed" fat of parenchymatous organs—the fat, that is, unrecognizable by microchemical means—can be extracted by ether after the tissue has been digested with pepsin—a mode of liberation which suggests strongly that the protein moiety of the compound becomes dissociated.

Autolysis and cell degeneration are two very different conditions, the one occurring in the dead, the other in the still living cell. But both are disintegrative, and we have dwelt at such length upon these autolytic phenomena because, in our opinion, they are to some extent paralleled by observations upon the degenerating cell, and throw light upon the process of degeneration.

In the first place, as shown many years ago by Stolnikow,² and confirmed by Ziegler and Obolonski,² Albrecht and Schmorl, and others, in the acute fatty degeneration, such as is produced by phosphorus, study of the cells of the liver and kidney shows that, coincident with the appearance of fatty granules in the cell, there is a remarkable process of chromatolysis with discharge of plasmosomes, or minute masses of chromatin from the surface of the nucleus into the surrounding cytoplasm. According to Stolnikow, more than half of the fat present in the phosphorus liver is in the form of lecithin. If the neutral fats are increased (by transportation), so also are the phosphatides.

As to the cause of the disintegration of the nucleus and discharge of the chromosomes, a suggestion of Wells' deserves consideration, namely, that phosphorus and the other poisons already mentioned act by inhibiting or destroying the higher cell activities, and notably the production of oxidases, while not influencing the lipases or enzymes associated with the elaboration of fats from fatty acids, etc.

It is interesting to recall in this connection the cases of the co-existence of fat and myelin in the conditions of normal "fatty degeneration" in the cortex of the adrenal (where the cells are filled with small, fatty globules; and, as first shown by Kaiserling and Orgler,⁵ doubly refracting myelin globules are also present—(see Fig. 309 on p. 914), and in the thymus undergoing retrogressive change; as, also, the association already referred to between the fat and casein (a phosphorus-containing protein) derived from the cells of the mammary gland.

¹ Pflüger's Arch., 65:1897:90; see also A. E. Taylor, Jour. of Med. Research, 2:1903:57.

² Du Bois-Raymond's Arch. f. Physiol., 87; Suppl. Bd. 1.

³ Ziegler's Beiträge, 2: 1887; 291.

⁴ Chemical Pathology, p. 341.

r

e

2

d

⁵ Virehow's Arch., 167: 1902: 296.

Lastly, in a very exact series of studies, Löhlein¹ has demonstrated that in the human kidney, as had previously been urged by yon Hansemann, two distinct conditions are recognizable-a fatty "infiltration," in which fatty globules alone are to be detected (in the cells of the convoluted tubules and the ascending limb of Henle's tubes), and a "fatty degeneration" in which abundant doubly refractive globules, the smallest only recognizable with the immersion lens, are present, along with definitely fatty, simply refractive globules, both in the cells of the convoluted tubules and in the endothelial and other cells of the interstitial tissue. This condition was always associated with indications of cell and nuclear degeneration, and was found by him in conditions of acute and chronic inflammation (Bright's disease), and very well marked in the different stages of amyloid kidney. These observations have been further developed by Klotz,² who in certain cases of the "large white kidney" has shown that, as already noted, the cells contain abundant doubly refractive myelin globules, but unlike Störk,3 who regarded these as of the nature of "protagon," his analyses indicate that these are relatively simple soaps of Na and K. The largefatty kidney is truly the large soapy kidney.

Löhlein has introduced another method of recognizing the myelin: sections of tissues left sufficiently long in Müller-formalin, and cut frozen, when washed in physiological salt solution and mounted in glycerin, exhibit minute, needle-like crystals, with stumpy ends and in clusters, in place of the myelin globules. Upon heating gently the crystals became reconverted into doubly refractive globules.⁴

All these data point to the existence of a lipoid degeneration leading to the appearance of fatty globules in the cells. Possibly this presence of fatty compounds is the explanation of the histological differences between the fine globules of the "degenerated" cell and the coarser globules of the "infiltrated," already noted.

There is not a little to be said in favor of the presence of cholesteryl oleate and other cholesterin fatty compounds as the cause of the myelin droplets in at least certain cases of degeneration. These are favored by Aschoff,⁵ and according to Craven Moore⁶ the action of formalin upon cholesteryl oleate affords the crystals noted by Löhlein. The two views are not absolutely contradictory: if lecithin be treated with

¹ Virehow's Arch., 180: 1905: 1.

⁷ Proc. Soc. Ex. Biol. and Med., 1908. Aschoff suggests that in most of these cases some cholesterin is present, and quotes analyses by Windaus in favor of this view (Ziegler's Beirr, 47: 1909: 1).

³ Sitz-Ber, d. Kais, Akad. d. Wiss, in Wien, Math.-Naturw, Kl., 115:1906, Abth. 3; 1).

⁴ It should be noted that Schmidt had previously recorded the same phenomena after treatment of the myelin bodies of the sputum with 1 to 2 per cent. KOH or NaOH. (Berl. klin. Woek., 1898; 73). For differentiation of intracellular crystals, see White, Jour. of Pathol., 13; 1908; 11.

⁵ Loc. cit.

⁶ Med. Chronicle, Manchester, 47: 1907: 204.

FAT PHANAROSIS

sterile liver juice it affords cholesterin, fats, and fatty acid, from which cholesteryl compounds may well be developed.¹

Wells,² in his valuable work on *Chemical Pathology*, would ascribe the fat in the kidney, spleen, nervous tissue, lung, in cases of fatty degeneration, to a rendering visible of the previously fixed fat already referred to-to what Klemperer3 has christened "fat phanarosis."4 He emphasizes Rosenfeld's observation that the rendering visible of the fat in the kidney is often accompanied by an actual decrease in the amount of fat recoverable from this organ; a kidney containing 16 per cent. of fat (i. e., below the normal quantity) may exhibit marked fatty degeneration, whereas another yielding 23 per cent. may show none. On the other hand, he holds that fatty degeneration in the liver and heart muscles is not due to such liberation of combined fat, but to accumulation from the blood, the difference in appearance from that seen in normal infiltration being due to cytoplasmic disintegration.

We cannot believe that the process is quite so simple. As we have pointed out, we can have similar accumulation of fat in fine droplets in the adrenal—in which the cells are not undergoing disintegration—and in liver and kidney we encounter identical nuclear changes and discharge of plasmosomes. We are inclined to question, also, the probability of the fats as such, and not rather fatty acids, being in combination with cytoplasmic matter. We can only conclude that there is a basal difference in the mode in which the fat is laid down in the cell in infiltration and degeneration, respectively, a difference which suggests strongly that we deal with the exhibition of different fatty compounds.

Lastly, we must note certain observations, so recent that their exact bearing upon the problem before us is difficult to estimate. Continuing certain observations of Leathes, Leathes and Hartlev⁵ show that, while the connective-tissue fats are almost entirely formed of the neutral fats of oleic, stearic, and palmitic acids, from the liver, kidney, and heart muscle of man and the higher animals, are to be obtained in fair amount by a process of saponification, members of the higher fatty acid series, fatty acids soluble in ether but insoluble in petroleum ether-acids of the linoleic and linolenic series, etc. (CnH2n-4O2, CnH2n-6O2, and possibly CnH2n-8O2). Dunham,6 as regards the kidney, has recently demonstrated the presence of another of the higher fatty acids-carnaubic acid, ${\rm C}_2 {}_4 {\rm H}_4 {\rm O}_2$. Conclusion.—To sum up and endeavor to harmonize these contra-

dicting views, it would seem that:

1. There is a physiological process of absorption from the fluids

- ¹ Waldvogel and Mette, Münch, med. Woch., loc. cit.
- ² Chemical Pathology, 1907: 334 et seq.
- ³ Deutsch. med. Woch., 1909: 89.
- 4 øaivø, to bring to light or make evident.
- ⁵ Jour, of Physiol., 36: 1907: 17; see also Leathes, ibid., 31: 1904: 1.
- ⁶ Proc. Soc. for Exp. Biol. and Med., 5: 1908: 58.

of the body of the precursors of the neutral fats, which precursors, absorbed into the cell in a soluble state, are by the action of lipases converted into neutral fats. By reversible action of the same enzymes (p. 82) these neutral fats so formed may be redissolved and discharged from the cells. When not so discharged, the neutral fats may accumulate in large globules, and the accumulations may be so excessive as to assume pathological proportions, and be known as fatty infiltration.

2. In this process there is no indication of the intermediate formation of myelin-like bodies. That the nucleus takes part in or controls the process is suggested by the presence of fat-containing vacuoles in the nuclei of normal fat cells.

3. In many tissues fats, or fatty acids, are present in a form unrecognizable under the microscope or by microchemical tests; this would indicate that a certain proportion of fatty matter must be present in the cell in a combined state.

4. Observations upon autolysis and phosphorus poisoning indicate a process in which the appearance of fatty globules within the cell is preceded by the increased formation of phosphatides and bodies of myelinic nature.

5. This lipoid degeneration is apparently a process distinct from ordinary fatty infiltration. In well-marked cases there is obvious disintegration of the nucleus, with discharge of the nuclear chromatin into the cytoplasm.

6. The fact that the nuclear chromatin contains nucleins as a main constituent, that both nucleins and phosphatides contain characteristically glycerophosphoric acid, and afford cholin as a disintegration product, indicate that the phosphatides, as regards these two constituents, are derived from nuclear matter.

7. The observations of Rosenfeld and others upon the translation of fats in phosphorus poisoning from the ordinary fat deposits of the body to the liver and heart muscle, are best harmonized with the above observations by regarding the glycerophosphorie acid and cholin of the lecithins as derived from nuclear material; the fatty acid constituents in these organs as, in the main, derived from fatty acid compounds brought to the cells from the other tissues in a soluble state, and there disintegrated; the fatty acid molecules combining with the glycerophosphoric acid and cholin to form phosphatides. That phosphatides and other lipoids (e. g., cholesterin compounds) are capable of undergoing further disintegration, and that thus the cells may exhibit both lipoid and pure fatty contents, must also be kept in mind. On the other hand, in those organs in which, during degeneration, the fat is increased, the formation of lecithin-like bodies may be brought about by combination with, and dissociation of, the fixed fats of the cytoplasm.

8. The disappearance of glycogen from the cells undergoing fatty degeneration suggests that it may be one of the sources of ultimate fat; as, indeed, it may be concerned in physiological fat formation.

GLYCOGENOUS INFILTRATION

DISTURBANCES OF CARBOHYDRATE METABOLISM.

GLYCOGENOUS INFILTRATION.

THE evidence that we have concerning pathological alterations in the glycogen contents of the tissues is at most meagre.

We recognize that glycogen plays an important part in normal metabolism; that, through ferment action, the starches of the food are converted into sugar; that, despite a large meal of carbohydrates, starches, or sugars, there is no marked increase in the sugar of the circulating blood; that absorbed sugars are rapidly taken up from the circulating more particularly by the liver cells and the muscles of the body; that in these organs they are stored as a less soluble modification, glycogen, or, as it has been termed, "animal starch;" that the liver may be regarded as the main storehouse and controller of the carbohydrate equilibrium of the system, the muscles as the main consumers of glycogen, muscular activity being dependent largely upon the dissociation of glycogen into carbonic acid, lactic acid, etc., carbohydrates affording

FIG. 310



Glycogen globules in cells of Henle's loops of kidney from ease of diabetes mellitus: each dark intracellular globule represents a red-staining globule by Best's carmine method. (After Gierke.)

by their dissociation the most easily and rapidly utilizable energy. In the liver we have evidence that points to the presence of reversible glycolytic enzymes which, under the one order of conditions, convert the soluble sugars into the less soluble glycogen; in the other, convert the glycogen into easily diffusible sugars, which pass into the blood, and so to the muscles and other tissues of the body (p. 83).

The physiology of glycogen is thus fairly well understood up to a certain point, and, histologically, if care is taken to deal with fresh material, or material which, when fresh, has been hardened in absolute alcohol, it is not difficult to differentiate and recognize the glycogen within the cells in the form of discrete vacuoles. As pointed out long ago by Claude Bernard, the discoverer of glycogen, if the tissues be kept, then through enzymic action the glycogen becomes converted into sugar, dextrine, and then it is unrecognizable microscopically; or,

more accurately, the process of conversion gradually slows, but what is left is no index of the amount originally present.

Glycogen is only relatively insoluble, forming a colloidal solution, and the fresh tissues must, therefore, not be brought into contact with water; they may be hardened in alcohol saturated with iodine and cut in iodine mucilage (Ehrlich) and mounted in iodine glycerin; or, after hardening in alcohol and passing through the ordinary procedure for cutting, may be treated with a mixture of tincture of iodine, 1 part to alcohol absolute 4 parts, and cleared with origanum oil. The glycogen by these methods is stained brownish red to claret color; unlike amyloid, it gives no reaction with sulphuric acid.

By using a special carmine stain with material prepared in celloidin, Best' has recently made a great advance in the ease with which glycogen can be detected in tissues.

It is in the liver that we most easily recognize glycogen histologically, and that we find the greatest variations. By chemical analysis it can be gained from both liver and muscles, as also from embryonic tissues. In these growing tissues it may be present in large amounts.

In moderate conditions of diabetes mellitus it has been found in considerable abundance in the liver cells, in severe cases it wholly disappears; but in these it has been noted in the heart muscle, and, characteristically, in the cells of the ascending loops of Henle in the kidney. What is its significance in this position has not been surely determined. In starvation and wasting diseases it disappears largely from the skeletal muscles and the liver; its absence from the former has been regarded as the explanation of the muscular weakness that accompanies these conditions; there is no store of readily convertible "fuel."

Just as glycogen is abundant in embryonic tissues, so has it been found abundant in new-growths of "embryonic" type; it may be detected in many neoplasms of a distinctly cellular and actively growing type, but more particularly in chorio-epitheliomas, myomas, endotheliomas, testicular and adrenal tumors.

Lubarsch² found osteomas, fibromas, hemangiomas, gliomas, and colloid cancers to contain no glycogen, and that it was rarely present in adenomas, lipomas, and lymphangiomas. The correspondence which Brault³ thought to exist between the embryonic type of a tumor and its glycogen content is not, therefore, complete, though, on the other hand, Gierke's⁴ contention that deficient oxidation is the cause of its appearance, cannot be supported.

The presence of glycogen in large quantities in renal hypernephromas was regarded a few years ago, by Lubarsch, as one of the arguments in favor of the origin of these growths from adrenal tissue, for this also is apt to contain considerable glycogen; it was later determined that richness in glycogen characterizes very many cellular tumors.

- ¹Ziegler's Beitr., 33:1902:585; see also Gierke, ibid., 37:1905:564.
- ² Virchow's Arch., 183 : 1906 : 188.

³ Jour. de Physiol. et de Path. gén., 6: 1904 ; 295 and 720.

⁴ Ziegler's Beitr., loc. cit., gives full bibliography.

GLYCOGENOUS INFILTRATION

Lastly, glycogen has been detected in pus cells, though here there has been some debate by Czerny and others regarding the existence of other iodine-staining globules possessing intermediate properties between glycogen and amyloid material. 'This view, however, has not received acceptance.

Save for a doubtful observation by Frerichs, no relationship had until recently been determined between glycogen and the cell nucleus; now Huebschmann' draws attention to the fact (and Rössle confirms) that in advanced cases of diabetes, while there may be no glycogen in the liver cells, certain nuclei are to be seen, sometimes in abundance, which are distended with one or several globules of glycogen. More rarely he encountered these glycogen-holding nuclei in other conditions (nutmeg liver, etc.). Glycogenous infiltration and fatty degeneration do not coexist; although glycogen deposits, contrary to the usual teaching, have occasionally been observed in liver cells, the seat of coincident fatty infiltration.

¹ Verhandl, d. deutsch, path. Gesellsch., 11:1908:35.

CHAPTER XXIX.

CALCIFICATION AND CALCAREOUS DEPOSITS.

THERE are few tissues which may not become the seat of interstitial deposits of calcareous salts. The deposits, it is true, most frequently occur in one or other of the connective tissues; in cartilage, in the connective tissue of the vessels, in the stroma of the organs; they are relatively infrequent in the parenchyma of glands, in muscle, and in nervous tissue, but even this last may be affected, the deposits occurring within the bodies of the cells.

In general, these deposits are large enough to be visible to the naked eye as opaque, whitish masses within the affected tissues. Their density varies from a crumbling, cheesy consistency, such as we encounter in caseous tuberculous foci of some little standing, when fine, gritty particles can be detected between the fingers, to a hardness greater than that of bone, as in old calcified fibroids of the uterus.

As in bone, these deposits are composed mainly of calcium salts, but there are wide differences between calcification and ossification. There is, in the former, a want of organization; cells of the nature of bone corpuscles and osteoblasts are wholly wanting. So, too, there is wanting anything resembling an orderly disposition in relationship to the vessels and matrix of the affected area. In bone there is a ratio, constant within relatively narrow limits, between the calcium and magnesium salts and the phosphoric and carbonic acids. The analyses of calcified tissue show no such constant ratio. In calcareous plaques from advanced arteriosclerosis von Kossa could detect no magnesium salts; in experimentally produced calcification of the kidney, no carbonates, though these were clearly present in experimental calcification of the liver, and are met with in normal bone. Kockel also cites examples of calcification in the lungs in which strong acid led to no evolution of carbonic acid. These statements are refuted by Wells,¹ who, on the contrary, calls attention to the pronounced similarity in composition between examples of calcification, studied by him, and normal bone, as regards calcareous salts. The conditions studied by him were calcified tuberculous masses in man and the ox, a calcified nodule from the thyroid, and a thrombus, the seat of calcification; these are compared with analysis by Zalesky and Carnot of human and ox bone.

	$Mg_3(\mathrm{PO}_4)_2$	$CaCo_3$	$\mathrm{Ca}_{5}(\mathrm{PO}_{4})_{2}$
Calcified matter	0.84 to 1.5	7.6 to 13.4	85,4 to 90.6
Normal ossification	1.02 to 1.75	9.2 to 12.8	83.8 to 87.8

¹ Jour. of Med. Research, 7: 1906: 491.

CALCIFICATION AND CALCAREOUS DEPOSITS

The only point of any importance, according to these figures, is that in calcification in general there is apt to be a wider range of variation in the percentage amount of the different salts than in ossification. It must be admitted that Wells' material was of a restricted character, and did not involve the grosser conditions of calcified tumors and serous plates, nor the commonest of all, calcified areas in the aortic wall.

Gierke¹ has called attention to the existence of iron in minute quantities in connection with normal ossification and its frequent but not constant presence in calcified areas. While he found it in psammomas in a calcified thyroid and a kidney, the seat of experimental calcification, it was absent in the common conditions of arterial and tuberculous calcification. Rarely, traces of calcium oxalate are encountered.

Chemical Reactions.—Treatment with acetic or mineral acid leads to the dissolution of the calcareous salts more rapidly than is the case with bone, more slowly than with pure phosphates and carbonates of calcium. When so dissolved, the extract, treated with ammonium oxalate, gives a heavy precipitate of crystals of calcium oxalate; treated with molybdic acid, a heavy deposit of phosphate. Apparently in all natural, as distinct from experimental calcification, when of any extent, treatment with mineral acids causes an evolution of bubbles of gas—carbonic acid gas—indicating the presence of calcium carbonate. Sulphuric acid causes solution, followed by the appearance of fine crystals of calcium sulphate (gypsum). Such solution by acids leaves behind an organic matrix.

Microchemical Appearances and Reactions.—Sections examined under the microscope show the salts differing according to the grade of calcification. The earliest appearance is that of a fine dust, scattered through the affected areas; more frequently there are rather coarse, obscurely crystalline or angular particles, highly refractive. They may run together to form solid plaques and masses, or occasionally, as in brain sand, there are evidences of growth by accretion into globular or polyhedral, somewhat crystalline masses, so as to form small concrements within the tissue. Whatever the size, the granules are insoluble in ether and in dilute caustic potash. They are slowly dissolved by formalin. With rare exceptions—to be mentioned later—upon removal of the salts, the matrix is found to be composed of dead tissue, in which the nuclei no longer stain, and cell boundaries are not to be detected.

In unstained sections, undeprived of their salts, the affected portions have a curiously opaque appearance. The simplest microchemical test is with hematoxylin, with which calcareous masses assume a deep-blue color. Upon treating sections for five minutes with pyrogallic acid (2 parts in 80) to which one part of caustic soda has been added, there is, after washing with distilled water, a decolorization of normal tissues, while the calcified parts stand out as a deep brown, becoming brownish black after the course of a few days. Yet more characteristic, according to von Kossa,² is the action for five minutes of a 5 per cent, solution of silver nitrate. This forms a vellow phosphate of silver, and, upon

¹ Virchow's Arch., 167 ; 1902 : 318.

² Ziegler's Beitr., 29:1901:63.

CALCIFICATION AND CALCAREOUS DEPOSITS

standing and exposure to the air, the salt is reduced and metallic silver precipitated, so that the smallest granules of calcareous salt within the tissues stand out prominently as coal-black dots.

Conditions under which Calcification is Found.—It is the custom to divide cases of pathological calcification into two distinct classes of (1) calcareous metastasis, and of (2) necrotic calcification; calcification being held to take place within the living tissue in the former case, in dead tissue in the latter.

Calcareous Metastasis.—In 1855 Virchow called attention to the fact that in certain cases of extreme resorption of bone from extensive caries, from malignant growths within the bone, and (doubtfully) osteomalacia, there may be widespread deposits of calcareous salts in cartilage, in the lungs, in the mucous membrane of the stomach, in the walls of



926

Section of human aorta of elderly individual, treated by von Kossa's method, to demonstrate calcification of media, and more particularly of the muscular bands. (Klotz.)

the arteries and capillaries, etc./ the presumption being that the excess of calcareous salts liberated from the destroyed bone becomes metastatically deposited in these other tissues. Such diffuse deposit is very rare. Little more than a dozen cases have been recorded in half a century.

For ourselves, we doubt the existence of this metastatic calcification; or, more correctly, would hold that its existence or non-existence depends upon what we regard as living tissue. We admit that calcareous deposits occasionally occur in tissues which still contain living cells, but those deposits occur not in the living cells themselves, but in the intert interstilial matter between

the cells; they occur, that is, in non-living material (p. 39), and if the calcareous matter be dissolved out by acid, it is seen that it had been contained in a swollen homogeneous matrix. Living functional cells do not take up and become the seat of deposit of calcareous salts.

There are certain possible exceptions to this statement which are still *sub judice*:

1. Schläpfer, Virchow, Grohe, and Roth have described cases of calcareous deposit in the outer layers of the mucous membrane of the stomach, rectum, and colon. It is suggested that when there is excess in the blood there is an actual excretion of calcareous salts through the mucous membrane of the intestinal tract, and that excessive excretion is accompanied by the presence of calcareous salts within the cells. We know of no recent studies in which the modern, more exact, micro-chemical methods have been employed, whereby it has been determined that the deposits occur in cells retaining nuclear stain.

2. Similarly, calcareous deposits occur in the kidney; while these must frequently show themselves in the contents of small cortical cysts, in old fibroid and hyaline glomeruli, and in the substance of retained casts, occasionally cells of the tubules can be seen containing dust-like, calcareous particles (Beer' and others). Most often these cells are loosened and free in the lumen, recognizable rather by their shape than by their nuclear stain; in the rare cases in which the cells remaining *in situ* show the deposits, it is probable that they are already necrosed, or at least necrobiotic. Beer, in his study of one hundred kidneys, from various conditions, demonstrates that macroscopic and microscopic calcification in the kidney is of the same type in cases of extensive disease of the bone and in cases in which no resorption of bone has been present indeed, is apt to be very pronounced in the latter; that such calcification is absent before the twenty-fourth year, constant after the thirty-fifth.

3. The same would seem true of the cases described by Kockel,² in which the cells of the capillary endothelium in the lungs have shown the deposit. Kockel shows that the so-called calcareous metastases in the lungs are of infarctous nature, due, when present, to emboli of cancer cells (in cases of malignant bone disease), etc., and identical in histo-logical character with the deposits found in "non-osseous" cases in long-continued passive congestion. In such infarctous areas, as in lung infarcts in general, there is no complete necrosis of all the tissues, but there is lowered vitality and necrobiosis. The lime salts are not deposited in the alveolar epithelium, but in the interalveolar fibrous tissue, and more particularly in the elastic tissue of the arteries and in the capillary wills. It is evident from this description that the cells involved are either dead or dving.

Necrotic Calcification.—There is, then, no satisfactory evidence that the process occurring in metastatic calcification is distinct in nature from that which occurs under other pathological conditions. At most, this is determined, that there may be calcareous infiltration, not merely of necrobiotic and necrotic tissue *en masse*, but also of certain interstitial substances. Of these, more particularly the matrix of cartilage and yellow elastic tissue appear to take up calcareous salts with some readiness. Allied to these, areas of pathological hyaline transformation possess the same tendency. As already noted, such hyaline areas are largely devoid of cells; they, too, are inert. Such deposits are often associated with senile changes.

In those advanced in years there may be very extensive calcareous deposits, and, combined with this, a progressive resorption of bone, so that the individual bones are thin and very light, and their salts greatly diminished in amount. The earliest deposits occur in the cartilages of the ribs, the larynx, and the respiratory system in general, in tissues, that is, which still retain their cells and normal structure, although the deposits are not in, but between, the cells. But with this there are usually deposits in the arterial walls, deposits occurring in definitely

¹ Jour. of Pathol., 9:1903:225.

² Arch, f. klin, Med., 64: 1899: 332.

CALCIFICATION AND CALCAREOUS DEPOSITS

degenerating tissue that has undergone necrobiosis. And, in addition, other deposits occur, which may be one or other order, in the interstitial tissues of the thyroid, more rarely in the testes or ovaries and other glandular organs, in the membranes of the brain, the parenchyma of the lungs, the subcutaneous tissues of the shins, etc., and in fibroid areas—scar tissue—the outcome of old inflammatory disturbances.

✓ Many of these are clearly examples of necrotic calcification. Upon histological examination it is obvious that there has been an antecedent necrobiotic change. Examples of such, apart from senile processes, are abundant. The commonest, as already suggested, is in connection with the arteries. In arteriosclerosis (see p. 451) the condition of calcareous atheroma follows hvaline and fatty degeneration and necrobiosis of the media and hypertrophied intima. But also, as Klotz more especially has pointed out, after middle age there may be calcification of the media, not necessarily accompanied by intimal overgrowth, not visible to the naked eve, but very marked when the specific tests are employed for calcium salts. Frequently also, there is calcification of old tuberculous foci in the lungs, lymph glands, spleen, and other organs. Here, again, calcification follows caseation, and caseation is the outcome of necrobiosis. Other inflammatory processes of the chronic type result often in calcification, notably chronic inflammation of the serose. More particularly where there has been chronic suppurative disturbance, with imperfect absorption and resolution, and dense fibroid adhesions are present, the central areas of such adhesions, cut off from adequate blood supply, undergo necrosis. Thus, large calcareous plaques are to be found in the pleura after old empyema. Similar plaques occur in the pericardium, associated with extensive adhesions. The process is not so common in connection with the peritoneum, though here localized areas of calcification may occur in the serosa of individual organs whose capsules have been the seat of chronic inflammation, notably in the capsule of the spleen, less frequently in that of the gall-bladder. In the chronic inflammatory group may be placed the calcification of capsules around foreign bodies, of cysts, etc.

Tumors whose blood supply has been wholly or in part cut off are liable to undergo petrifaction. That which is the commonest tumor of all—the uterine fibromyoma—frequently exhibits it. And we meet with general or localized calcification in other slow-growing and benign tumors, in fibromas and lipomas; it has even been recorded as occurring in slow-growing scirrhous cancers. One form of tumor, the slowgrowing endothelioma of the brain membranes, is so apt to exhibit areas of calcification that it has come to be regarded as a special variety, the *psanmoma*.

Organs, or portions of organs, whose blood supply has been cut off may exhibit the process. Thus, we at times encounter calcified infarcts. Individual cells undergoing necrobiosis may exhibit it, as, for example, nerve cells after traumatism. According to Durante, calcification of muscle fibres is almost physiological in the herbivora. In man the same may be encountered in the neighborhood of sutures and abscesses

THE EXPERIMENTAL PRODUCTION OF CALCIFICATION 929

and as the result of traumatism; here, again, the calcification, according to Schnjeninoff,¹ is preceded by necrosis. In this connection we may include the impregnation with lime salts of the dead factus, the result of extra-uterine gestation retained within the abdominal cavity (*lithopedion*). Similarly, the cysts of parasites within the tissues, hydatids, trichine, etc., are liable to before impregnated.

The Experimental Production of Calcification.—Before discussing the factors at work in the production of calcification, and as an aid to that discussion, it will be well to note the facts gained from experimental observations. These experiments have, it is true, been mainly upon one organ, the kidney, and the process is not, therefore, in all respects parallel with the commoner examples encountered in man, but, notwithstanding, they establish certain points very definitely.

Litten,² in 1881, made the first full study in this direction, and pointed out that if, in the rabbit or dog, the renal artery be ligatured for one and a half to two hours, and then the ligature be removed, at first little disturbance is to be detected; the epithelium of the tubules appears uninfluenced. But in twenty-four hours the cortex is found swollen and hyaline; the nuclei in the cells of the convoluted and some of the straight tubules no longer stain, while here and there the cells have fused into cylinders, completely filling the enclosing tube of basement membrane. Some of these cells already show irregular, highly refractile granules, soluble in acid. By the second day the necrosis, the formation of hyaline cylinders, and the deposit of granules within the cells and cylinders is most pronounced. The whole tubule becomes filled with dense granular matter; the deposit becomes more and more intense, until, by the tenth day, the organ is so hard that the razor is notehed in attempting to cut it.

Here, obviously, the result of the ligature has been to induce a necrobiosis of the tubules, and this precedes the deposit of the salts. Litten laid great weight upon the fact that not all forms of necrosis lead to calcification. With the necrosis there must be continuance of an adequate arterial supply. But herein Litten was mistaken; this is not absolutely essential. Complete ligature of all the vessels at the hilus of the kidney may be followed—slowly—by calcification of the organ. But then the process is somewhat different. It occurs at the periphery, slowly progressing toward the deeper parts. The process, in fact, is of the same type as that which we encounter in the lithopedion, in calcifying caseous tubercles and uterine fibroids. But in both cases, obviously, the results are to be explained by infiltration; where the arterial supply is preserved this takes place from the arteries, and the deposits occur throughout the organ; where it is cut off the infiltration is from the lymph at the periphery of the organ.

A calcification identical in its stages with that produced by Litten had already been observed in sublimate poisoning, by Salkovsky, in 1866; Kaufmann and other observers have since noted it in subacute cases of

² Virehow's Arch., 83: 1881: 508.

¹ Zeitschr. f. Heilkunde, 18:1897.

CALCIFICATION AND CALCAREOUS DEPOSITS

corrosive poisoning in man, and have produced similar changes in the kidney by the employment of neutral potassium chromate, and now a long series may be given of drugs leading to renal necrobiosis and calcification—aloin (Gottschalk, 1882), glycerin (Afanassiew, 1884), bismuth subnitrate (Langhans, 1885), cyanide of mercury (Virchow, 1888), phosphorus (Paltauf, 1888), acetate of lead (Prevost and Binet), copper sulphate, iodine, and iodoform (von Kossa, 1901), formalin (Putti, 1904), copper acetate (Klotz, 1905).

Kaufmann, in his cases, noted an intense contraction of the renal artery and arterioles, with great venous engorgement; he attributes the necrobiosis to this rather than to the direct action of the sublimate upon the cells. In passing, we may note that these conditions of extensive cell death are conditions which favor autolysis. There would seem to be some relationship between the changes which occur in the cell undergoing such autolysis and the subsequent development within it of calcareous deposits.

Similar appearances are produced in the liver cells of the rabbit by the ingestion of iodoform, calcification being preceded by extensive fatty degeneration.

For all these experiments the rabbit has been found more serviceable than the dog, because its epithelium more easily undergoes necrosis; and, secondly, its blood contains a much larger proportion of calcium salts. Dried rabbit's blood contains from 1 to 2 per cent. of calcium; dried dog's blood only 0.05 to 0.07 per cent.

We may say in passing that, experimentally, poisoning with oxalic acid (Kobert and Keusner and Neuberger) leads to abundant deposits within the kidney; but these do not stain with hematoxylin; they are deposits of calcium oxalate.

The Causation of Calcareous Deposits.—We are now in a better position to discuss the causation of these deposits.

In the first place, remembering that line salts are constituents of practically all the tissues and fluids of the body, have we to deal simply with a local change in these salts from the soluble into the insoluble precipitated form, after the manner of Lot's wife? Certainly not. The amount of calcium phosphate and carbonate in the calcified uterine fibroid is very far in excess of the amount present in an ordinary fibroid. As von Kossa has shown, the kidney of a rabbit in which calcification has been brought about by aloin may contain three hundred times as much calcium as does the normal kidney.

Or, in the second place, have we to deal with an alteration of the blood and lymph, so that either (a) the actual amount of soluble calcium salts has been increased until saturation occurs, or (b), on the other hand, the state of the fluid is so modified that it is unable to hold the normal proportion of salts in solution? There is not a particle of evidence in support of either of these views. We never find the blood so full of calcium salts that these become deposited within the vessels or precipitated after the removal of the blood from the organism. In all cases the amount of calcium salts in the blood is far below the point of satura-

tion. Cases of calcareous metastasis cannot be explained upon any theory of saturation. At most, this can be said, that, as indicated by von Kossa's analyses, increase in the amount of calcium salts in the body fluids favor the precipitation in certain tissues. There is, indeed, one tissue of election, namely, the matrix of cartilage. What stands out very prominently is that in the majority of cases a deposit of these salts occurs in dead tissue, and where, as in cartilage, the tissue is not dead, deposits do not occur within the bodies of the living cells, but in the interstitial substance, in matter which is extracellular and is of low vitality, if, indeed, it can rightly be regarded as living.

We arrive, therefore, at this, that calcification occurs only in dead tissue or in the inanimate, intercellular parts of living tissue, and that it is, not a precipitation of salts normally present in the affected areas.

Thus, it follows that, for calcification to occur, it is necessary that lime salts be brought to the parts. This can only be through the agency of the blood, or, more exactly, of the lymph. We know that both of these fluids contain calcium salts in solution. As the lymph diffuses into the parts, chemical processes ensue, such that the contained lime salts become converted into insoluble salts, and are precipitated in situ. This is very evident from a study of experimental calcification.

But what is the nature of the chemical change leading to the deposit? Here we enter upon debatable ground. While we know that calcium is present in the blood and lymph, we do not know with certainty how it is there combined. The amount is so small that it may be present in the form of ions, or, as Brailsford Robertson' has recently shown, it may be combined with protein ions to form an ion-protein compound with some of the proteins of the lymph, resembling the compounds between calcium and casein. That is to say, it is debatable whether calcium exists in normal lymph as definite phosphate or carbonate. Analysis indicates that it cannot all be thus combined.

There are, it will be seen, two possible methods whereby the calcium and magnesium salts become deposited, which may be termed the physical and the chemical, respectively. By the former, the salts are to be regarded as deposited without direct chemical interaction with the matrix, the conditions in that matrix being such as to favor the interaction between the components of the salts, so that they join into insoluble forms. Several theories have been advanced along these lines, but none has proved satisfactory.

10

H

d,

3-

Thus, Askanazy, to explain calcification in the stomach mucosa and the kidneys, has urged that the deposits occur in parts which secrete an acid fluid, and so are rendered increasingly alkaline; but there are other areas, like the rectum and colon, secreting an alkaline fluid, and, nevertheless, calcification has been recorded in them. Chabrié has suggested that in necrotic areas the CO_2 becomes fixed, and, with its removal from solution, the calcium phosphate and carbonate become precipitated. Irvine and Woodhead, on the contrary, suggest that nascent CO_2 plays

¹ Jour. of Biol. Chem., 2: 1907: 317.

CALCIFICATION AND CALCAREOUS DEPOSITS

the main role, pointing out that in the presence of free CO_2 a solution of carbonate of lime and phosphate of soda yields a precipitate of phosphate of lime. Others, again, have called attention to the special affinity between colloids and crystalline substances, and more particularly the liability for the latter to undergo abnormal crystallization in the former. Certainly, as pointed out by Wells, dead cartilage placed within the tissues becomes rapidly impregnated with calcium salts. This, however, does not explain why functional cartilage remains for long years in the organism continually percolated with lymph without a sign of calcification. Some chemical changes must take place in senile cartilage favoring this precipitation. The evidence in favor of such physical deposit of the salts is thus singularly unsatisfactory.

Is there any evidence in favor of deposition through chemical activities in the affected areas, of junction between products of tissue degeneration and the calcium and magnesium brought by the lymph? Here various possibilities may be suggested: (1) That the disintegration of the necrotic tissue supplies the phosphoric and carbonic acid which combines with the calcium and magnesium brought by the blood; (2) that the simpler products of disintegration of the protein substances present combine with the calcium and magnesium, there being formed calciumprotein compounds in the first place, the proteid moiety acting as a weak acid, being subsequently replaced by the stronger phosphoric or carbonic acid; or (3) that the fatty acids which make their appearance in degenerating areas play a similar part.

It is along these lines of testing those various possibilities that the most active work is being engaged in at-the present time. The first may be dismissed; the amount of phosphoric acid accumulating in, for example, a densely calcified uterine fibroid is far in excess of what could be supplied by the decomposition of the nucleoproteids previously present. The phosphoric and carbonic acids must in the main be conveyed to the part by the lymph.

As regards the second possibility, this cannot be neglected. The affinity of dead cartilage for calcium salts may well indicate an active process of association between the two; the fact, also, that the dissolution of calcareous deposits by acid always reveals a hyaline matrix is at least suggestive, although it does not necessarily imply that the salts have been in direct chemical union with the matrix; they may merely have been adsorbed. Until we know more concerning the compounds between calcium and protein, or the disintegration products of protein, and have isolated such compounds from areas of calcification, this must remain but an hypothesis.

Theory of Calcium Soap Formation.—The third possibility, suggested sporadically by occasional workers during the last fifty years (Weber, Wagner, Diakonow), had received little attention or acceptance untilthe investigations of Dr. Klotz in our laboratory at McGill brought it prominently to the fore; at the present time it may be said to afford the most promising theory so far advanced.

Many years ago Virchow demonstrated the presence of calcium soaps

THEORIES OF CALCIFICATION

in a lipoma; more recently, Jaeckle¹ has analyzed a calcifying lipoma in which 29.5 per cent. of the calcium was present in the form of soap. Klotz² showed that if celloidin capsules, filled with fat or fatty acid, be inserted into the peritoneal cavity of a rabbit, in the course of a few days these are found to contain amounts of calcium far in excess of that present in the body fluids of the animal. In other words, the calcium, percolating into the sac, becomes fixed. As already noted, calcification occurs, in the main, in areas of necrotic disintegration. By the improved microchemical methods for the detection of fats and calcium salts in the tissues, Klotz was enabled to show that in the atheromatous aorta, the most frequent seat of calcification, as also in calcifying tumors and calcareous plaques of the pleura, and in the experimentally induced calcification of the rabbit's kidney, the process of calcification was in all cases observed to be associated with fatty degeneration. The oldest and densest parts of the deposit did not show this; it was present at the peripheral zone, where the process of deposition was more recent, and here cell bodies could be detected which gave both the fat and the calcium salt reaction. What is more, in this confirming Fischler, he called attention to the fact that fats and soaps take on a differential stain with Sudan III, globules of the latter assuming a yellower tint. In this way he demonstrated that soaps are present in recognizable amounts in areas of progressing calcification. By Fischler's methods fatty acids and their salts are also to be demonstrated in the areas of calcification. He concluded that the stages in pathological classification are:

1. Fatty degeneration, with liberation of fatty acids.

2. Combination of these with calcium to form compound calcium soaps.

3. Interaction between the soaps and the phosphates and carbonates brought by the lymph, resulting in the replacement of the weaker fatty acid by phosphoric and carbonic acids and deposit of insoluble calcium phosphate and carbonate in the dead tissues.

His observations led him to regard the soaps thus formed not as simple soaps, but as compounds of fatty acid, calcium, and some protein or product of protein disintegration. The weak point in the investigation is that Klotz did not determine quantitatively the amount of calcium soap obtainable in his cases; qualitatively such were clearly present, although, obviously, in small amounts. Both Wells and Baldauf have called attention to the fact that the quantity present is very small; according to Baldauf³ it is in general non-existent; but the latter's methods are open to criticism, and Wells⁴ would appear only to have studied advanced conditions—not advancing. If the fatty acid play the part of an intermediary body, bringing together successive molecules of calcium and phosphate, a very minute quantity of calcium soap might be present at any moment So, also, was Klotz unable

¹ Zeitschr. f. Physiol. Chem., 36: 1902: 53. ² Jour of Exp. Med., 7: 1905: 661

³ Jour. of Med. Research, N. S., 10: 1906: 355.

⁴ Ibid., 9: 1906: 491, and 12: 1907: 14.

CALCIFICATION AND CALCAREOUS DEPOSITS

to indicate more than vaguely the nature of his presumed compound soap. From our own studies on the myelius, and their relationship to autolysis and tissue degeneration, we are prepared to find that these play an important part in the process. Although the possibility cannot be overlooked that the fatty degeneration and the calcareous deposit may be independent processes, their relationship, as indicated in Dr. Klotz's sections, is so striking that it is difficult to regard it as merely a coincidence, while his collodion sac experiments have proved that fats and fatty acids are capable of forming calcium soaps when free within the organism. Here, in short, is the simplest and the most satisfactory explanations of the mode of development of calcareous deposits.

CONCREMENTS AND CALCULI.

In addition to this deposit of calcareous salts within the tissues, there may be a deposit of the same in the ducts and passages of the body, leading to the formation of sharply defined solid masses, either round or oval, or assuming the shape of the duct in which they are found. They constitute one order of the solid bodies occurring in these regions and growing by concretion, to which the term *concrement* or *calculus*' is applied. The terms, in short, are applied vaguely, and some would regard them as interchangeable, though even these are accustomed to use the term calculus for the agglomerated deposits occurring in the large excretory passages, hepatic, urinary, and pancreatic, and concrement for those occurring in less usual sites.

We may divide these concrements into three main groups: (1) Those composed characteristically of calcium salts; (2) those formed of the precipitated constituents of one or other excretion or disintegration product; and (3) those formed of accumulations of foreign matter. It will be readily understood that these groups are not absolutely defined: where there is retention of the products of excretion in a duct or deposit in one or other channel of foreign matter, a certain amount of calcareous impregnation may occur; nor can there be deposit of calcareous salts without at the same time inclusion of excretory matter, but for general purposes the distinction is well marked and of utility.

CALCAREOUS CONCREMENTS.

The calcareous concrement, therefore, we regard as a firm and solid mass forming in one of the passages of the body, which, on analysis, furnishes a notable amount of calcareous salts, in addition to varying

¹ In our first edition we urged that the term concrement be confined to calcarcous deposits of this order, calcult to deposits of the specific excretions of various glands. Experience shows that this employment is apt to lead to confusion. It is better to retain the older view that any isolated deposit growing by concretion is a concrement, and to employ calculus as an alternative term commonly applied to certain orders of concrements.

CONCREMENTS

amounts of other constituents, these calcareous salts being greatly in excess of the quantities usually present in the fluids discharged along the affected passages. On dissolving out the salts, there is constantly left a matrix of proteid or mucinous type, and in general there is an admixture of fatty acids, soaps, cholesterin, and products of proteolysis.

This composition and the clinical history of these cases, in which the development of the concrements can be followed, affords a clue to the mode of their development. We deal, that is, in general, with the results of a catarrhal process—an inflammation—whereby, in the first place, there is exuded into the passage a mucinous discharge, together with exfoliated cells. The disintegration of the latter affords the products of proteolysis and the fatty matters, and in such a matrix, just as in necrotic areas within the tissues, there next occurs a deposit of calcareous salts, through diffusion of serum of the inflammatory exudate into the mass, as, again, of the secretion normal to the passage. Of concretions of this order, the following may be noted:

Rhinoliths are concretions occurring in the nasal passages, most often as the result of chronic ozena and obstruction. Berlioz,¹ as the result of an analysis of four specimens, obtained on the average 17.2 per cent. of organic matter; calcium phosphate, 57.9; calcium carbonate, 13.6; magnesium sulphate, 5.5 per cent.

Tonsillar concretions forming in the crypts of the tonsil as the result of chronic obstructive inflammation have like characters.

Salivary Concrements.—Formed in one or other of the salivary ducts. The analysis of various specimens has given very varying results from as high as 83 per cent. of calcium phosphate to as low as 15 the amount of organic matter being correspondingly varied.

Other concrements of the same order are *lacrimal* concrements: *cutaneous* concrements (the latter often multiple, and formed, it would seem, in obstructed and dilated sebaceous glands); *preputial* concrements (found in conditions of phimosis with the accumulated smegma as base); in these there is an admixture of relative large amounts of calcium phosphate with urinary salts, notably the ammonia-magnesium phosphate precipitated by ammoniacal fermentation of the urine; and *appendicular* concrements. The commonest accumulation in the appendix is of semisolid fecal matter with abundant organic and relatively small proportions of calcareous salts. Rarely, large solid concrements are encountered in an obstructed appendix, having abundant calcareous salts.

Intestinal Sand.—With these may be mentioned the true "sable intestinale," small, brownish concretions (from staining with fecal pigment) showing abundant organic matter, but also abundant calcium phosphate, and in some cases abundant magnesium phosphate.² These deposits are comparatively rare and small in man, but in cattle huge concretions may form in the lower bowel around retained vegetable

¹ Jour. de Phar. et Chem., 23: 1891: 497.

² Vide Duckworth and Garrod, Lancet, 1: 1902: 653.

CALCIFICATION AND CALCAREOUS DEPOSITS

foodstuffs; around such a nucleus there may be deposited a thick, solid layer of calcareous salts.

Pancreatic Concrements and Calculi.—Occasionally, concretions form in the pancreatic duct, leading to cyst formation (ranula). The commonest form is the true concrement with matrix of organic matter and mixed cholesterin, and preponderance of phosphates and carbonate of calcium. But here, again, the percentage of the salts exhibits great variations. We would suggest that here, as in the appendix, the age of the concretion is a factor determining variation. A recent catarrhal deposit may show abundant products of cell disintegration, proteins, fatty acids, soaps, cholesterin, etc.; one of long duration, disappearance of fatty acids and soaps, and with preponderance of calcium and magnesium salts.

Phleboliths.—In this category are to be included phleboliths, small oval stones formed occasionally in veins. Their commonest site is in the prostatic plexus in man; the uterine plexus in woman. Every transition may be found, from a recent thrombus in one of the anastomosing branches of such a plexus to extremely dense, pearl-like bodies. We deal, obviously, with the gradual deposition of lime salts in isolated thrombi which have not undergone organization, so that the masses lie free in the lumen of a vein.

We have not seen discussed the conditions under which a thrombus undergoes this characteristic metamorphosis. The necessary factors would seem to be: (1) Small size of thrombus; (2) bland, non-infected nature; (3) development in a communicating vein, the thrombosis affecting but one of the series of alternative channels, and leading thus to no secondary disturbance in the nutrition of an area. Under these conditions it would seem that the thrombus undergoes a certain amount of contraction within its bed; some absorption takes place at either extremity, but before this process can be complete three has been an impregnation of its substance by the blood plasma and precipitation within it of calcareous salts, arresting further dissolution. Processes of heterolysis (see p. 372), due to the disintegration of the contained and infiltrating leukocytes, may also be involved to liberate in the thrombus those bodies which fix the calcareous salts.

Calcareous Incrustations,—Deposits upon surfaces may undergo similar impregnation with calcareous salts derived from the body fluids and secretions. The commonest example is *tartar* of the teeth, in which mucinous epithelial debris and foodstuffs form the matrix. In catarrh of the bladder and ammoniacal cystitis a deposit of mixed phosphates not infrequently forms an incrustation upon the vesical mucosa. Incrustation may also form on the surface of foreign bodies gaining entrance into the tissues and cavities of the organism.

CHAPTER XXX.

DEPOSITION OF OTHER PRODUCTS OF CELL METABOLISM.

CONCREMENTS FORMED FROM PRECIPITATED CONSTITUENTS OF THE EXCRETIONS.


PROSTATIC CALCULI.

SECTION through the prostate glands of most men beyond middle age reveals the presence of little dark-colored granules here and there throughout the substance of the organ. Examination of sections show that these are situated within the lumina of the glands. They are sufficiently soft to be cut with the knife (in paraffin sections), are hyaline in appearance, rounded or polygonal, and exhibit concentric structure. Rarely do they exceed the size of millet seeds. They exhibit imperfectly the reactions of amyloid matter—they may or may not give the iodine reaction, and may or may not take on a metachromatic stain with methyl violet; like amyloid, they are thoroughly resistant to acids and alkalies. They resemble in many respects the corpora amylacea about

F1G, 312



Corpora amylacea from brain, to show laminated character. × 250,



Section through a "corpus amylaceum" from a sternal tumor, yet more highly magnified to show the subcrystalline deposit of successive layers of closely packed needles of amyloid material. At athe needles radiate from a small focus. (Ophdia)

to be noted, but should perhaps be classed apart as occurring in a normal secretion. Rarely, larger concrements are encountered in the prostate, in which successive layers of calcareous salts have become deposited around a nucleus formed of one of these "amyloid" bodies.

CORPORA AMYLACEA.

Minute concretions, often showing several concentric layers, may be found in the brain tissue and spinal cord of elderly individuals, in the

CALCULI AND CONCREMENTS

alveoli of the lungs (most often, it has seemed to us, in the lower lobes in cases of chronic congestion) and in certain tumors. They have the general appearance of hyaline material, and may, or may not, exhibit a certain amount of metachromatism with gentian violet and other aniline dyes, suggesting thus some relationship to amyloid. The observations of Ophüls' show that the growth of some, at least, of these is by the successive deposit of layers of obscurely crystalline needles of matter of a protein nature.

URINARY CALCULI; LITHIASIS.

Normally, the urine contains several salts in solution; abnormally, it may contain yet others, and under certain conditions members of either group may be precipitated in the course of the urinary tract and be deposited, layer upon layer, so as to form urinary calculi. We have to consider not merely what are the different forms of urinary calculi, but what are the conditions favoring their formation. It cannot be said that we have a full knowledge concerning these, but something has been gleaned, more particularly as bearing upon the three more common classes, the uratic (including uric acid and the urates), the calcium oxalate, and the phosphatic. These we will discuss in a little detail, treating the less common forms more briefly.


To prevent repetition, certain features common to all calculi may here be mentioned; and first, it must be pointed out that the calculus is of relatively slow formation, and that it grows by accretion upon the surface. There is a nucleus of mucus, cell debris, or foreign matter, such as blood clot, or some body actively introduced into the urinary tract, and in and upon this the salts become deposited. The presence and nature of this nucleus can often be determined by careful observation. As in ordinary crystallization, so here, salts present in solution must have some solid or semisolid as a base upon which to separate out from solution. The rate of separation and deposition must depend primarily upon the relative amount of salts present in solution, and as the urine varies both as regards amount of contained water and portion of various salts in solution, so the rate of deposition necessarily varies. Periods of relatively abundant deposit alternate with periods of arrest. As a consequence, most vesical calculi are, on examination, seen to be formed of concentric lamina. Not only is this the case, but in a fluid of such varying composition as is the urine, now one, now another salt may be present in relative excess, so that at one period conditions may favor the deposit of one particular salt or combination of salts; at another, the deposit of other constituents of the urine. As a matter of fact, it is the exception and not the rule to have vesical calculi composed throughout of one constituent. There may be alternate layers of uric acid and calcium oxalate, or often of urates and phosphates. It is very common

¹ Jour. of Exp. Med., 5: 1900: 111.

URINARY CALCULI; LITHIASIS

to have an outer coat of phosphates, this latter deposit being due to the irritation caused by the growing stone having eventually favored the development of infection in the urinary tract, with bacterial fermentation and production of alkaline urine.

But the mere presence of excess of one particular salt is not the only factor. If, by suitable reagents, the salts of one or other order in the urinary calculus be dissolved out, there is left behind a *matrix* of organic matter, of homogeneous, gelatinous, or colloid type, more or less pigmented or yellowish. What is more, if, after the methods of the geolo-



F16, 314,-Calculus formed of uric acid followed by oxalate appearance on section,

FIG. 315.—Oxalate of lime calculus ("mulberry" calculus), exterior view.

FIG. 316.—Ammonium urate followed by oxalate and eventually by mixed phosphates. (These three figures are from the catalogue of the Royal College of Surgeons.)

gist, calculi be rubbed down until fine sections are obtained, it is seen that, as a rule, they are not formed of well-made typical crystals of one or other order. The nearest approach to such perfect crystalline form would seem to be in cystin calculi; calcium oxalate and uric acid are present most frequently in spheres and dumb-bell shapes. It is worthy of remark that in bone, egg-shell, and other normal deposits of salts, examination shows that these also, in general, are not deposited in a crystalline form. Attention was called to this fact by George Rainey, in 1857, and he formulated a theory of molecular coalescence to account

for the peculiar method of deposit of salts in organic material. In 1873 Vandyke Carter' applied Rainey's observations to calculus formation. He explained the development of urinary calculi as due to the production in certain states, either in the renal pelvis, the ureter, bladder, or urethra, of a mucous secretion; a colloidal matrix was thus afforded in which the salts were precipitated in an imperfectly crystalline, rounded, or amorphous form. In this way was produced, according to him, the most common type of nucleus, and most often this contained oxalate of lime and uric acid or urates. Once the nucleus is formed, other layers of mucoid material are successively deposited, and in these there occurs a like process of decomposition, the stone in this way gradually growing.

Ord,² in 1879, confirmed and advanced these observations by his studies upon change in the form of uric acid and various salts precipitated from saturated solution in albuminous and sugary media. To Ebstein³ is usually but wrongly given the credit for these observations.

Now, mucin, or, more correctly, a mucinous body, apparently derived from nucleoproteids, is thrown off in abundance in conditions of inflammation of the urinary passages, and the greater the amount of this in suspension in the urine, the greater the liability of salts which would otherwise have remained in solution, to become precipitated in the colloid menstruum. Whether loose combination occurs between the salt and the colloid matter is still an open question. Thus, for the formation of urinary calculi, we recognize two factors: (1) The presence of one or other crystallizable body in relative abundance in the urine, and (2) irritation of some portion of the urinary tract leading to increased discharge of mucinous material.

Moritz⁴ has pointed out that an organic base is present in every urinary crystal, whatever its nature, and, as the urine always contains mucinous and other organic material which can serve as a base, Krehl,⁵ following him, doubts whether it is necessary to invoke a preliminary local irritation or inflammation. In taking this stand, however, the physical influence of concentration of the colloid material is left out of account.

Irritation, it would seem, there must be, but not necessarily of bacterial nature. Chemical or mechanical irritation alone, as, for instance, in alkaline urine the presence of ammoniacal salts, can set up increased mucinous discharge. But infection of one or other part of the urinary tract is also capable of favoring the process, and, once it has begun, the local irritation set up by the presence of the foreign body leads to deposit of a surface layer of mucinous matter, and so the stone grows. The indications are that in the urinary passages, as in the gall-bladder, low forms of infection, notably by the B. coli, initiate the process in a large number of cases, but not necessarily in all. Without infection the urates precipi

¹ On the Microscopic Structure and Formation of Urinary Calculi, London, 1873.

² On the Influence of Colloids upon Crystalline Form and Cohesion, London, 1879.

³ Die Natur und Behandlung der Harnsteine, Wiesbaden, 1884.

⁴ Cong. f. inn. Med., 1896; 532. ⁵ Path. Physiologie, Leipsic, 1898.

URATIC CALCULI

tating in the collecting tubules of infants can set up a very definite irritation.

Uric Acid and Uratic Calculi.-Uric acid may be a constituent of a stone, either as the acid or in the form of amorphous urates. The first of these is the more common. Calculi formed mostly of urates are soft and of gravish-vellow color, and are found chiefly in children. Combined uric acid and urates are fairly common at all ages, the uric acid predominating. In fact, uric acid is the commonest constituent of calculi, whether as the main constituent or forming the central portion of the stone. Frequently there is some admixture with oxalate of lime. Uricacid calculi may be found either in the pelvis of the kidney or in the bladder. The typical vesical uric-acid calculus is of flattened, rounded, or oval shape; its surface is smooth or finely mammillated; the color varies from pale fawn to a brick red, according to the amount of associated pigment (*uroerythrin*) which is brought down in the urine when uric acid separates out. Upon section the individual laminæ are well recognized. Frequently the calculus is solitary, but they may be multiple, in which case the individual stones may exhibit smooth, faceted surfaces.

These calculi apparently originate in the pelvis of the kidney, and, instead of being arrested in the bladder, they may, while still small, be voided in the form of gravel. These little calculi have a smooth surface of reddish tinge. On the other hand, the calculus may never pass beyond the pelvis of the kidney, and here it may grow until it forms a large "stag-horn" mass, forming a mould of the dilated pelvis. These calculi may be found at any period of life, but their frequency in early years is noticeable.

Uratic Inspissation of Infancy.—The liability for the development of uric acid and uratic calculi in the young appears to be intimately associated with another condition, that, namely, of the formation of so-called *uric-acid infarcts*—a term which, while old established and etymologically correct as regards usage of the term infarction, is unfortunate, since our ordinary employment of that term is for a totally different process. Nor do these infarcts contain crystals of uric acid. *Uratic inspissation* would be a much more satisfactory term.

Autopsies upon infants a few weeks old not infrequently afford kidneys in which the calices and a considerable part of the medulla have an opaque, dirty whitish appearance, and, upon examination, this is seen to be due to the obstruction of the collecting tubules with abundant small spherical masses lying in a mucinous matrix. These are doubly refracting under the Nicol's prisms. Chemically these masses have all the properties of amorphous urates—quadriurates. The urine of the newborn contains normally twice as much uric acid as does that of the adult. Why this should be so is a matter of debate. Woods Hutchinson explains it as due to the breaking down of the nucleated red cells, which, present in the foctal blood, rapidly disappear upon birth. Fry's observations indicate that these deposits may only occur some days, or even weeks, after birth. The acute disturbance of metabolism, the tissue destruction following upon birth, and, more particularly, the leukocytosis of the newborn

CALCULI AND CONCREMENTS

appear to afford a more satisfactory explanation. So grave is the disturbance which this obstruction of the tubules may set up that, as Fry¹ and Martin point out, there may be developed not merely a coincident albuminuria, but a definite condition of infantile uremia. This excessive discharge would appear to be the origin of the uratic calculi in the young. Occasionally in young children, in the pelvis of the kidney, more rarely in the bladder, we encounter small, soft agglomerations of the same type having a mucoid matrix containing these spherical grains of quadriurates, and examination of fully formed uric-acid calculi shows, most often, a nucleus of this nature.

Both in the young and in adults who are the subjects of gravel and calculus, the urine passed is characterized by its high acidity and by the deposit, on standing, of a brick-dust precipitation of urates, from which uric acid gradually crystallizes out. While, as already pointed out, freshly passed urine rarely contains free uric acid, these stones are formed mainly of this substance. We are led to conclude, therefore, that, in the process of decomposition in the mucoid matrix, the quadriurates are broken down, uric acid is precipitated, and the alkaline urates are set free.

The usual teaching is that lithiasis—the condition in which there is this passage of urine containing precipitated urates—exhibits increase discharge of urates. Such increased discharge does occur, but it is, we think, more than debatable whether this is essential for calculus formation, or even for many cases of lithiasis, for, in a large proportion of cases, brickdust deposits in urine are due not to increased discharge of urates, but to a concentrated and more acid state of the urine than is normal. There is frequently to be noted a relationship between gout and lithiasis such that those afflicted with gout have been liable to show abundant urate deposits, or to pass gravel or, again, some members of the family are gouty, the others the subjects of lithiasis. But it is not the family that forms the main factor in lithiasis, and the relationship to gout suggests that, in some cases at least, the actual amount of uric acid formed may be below rather than above the normal.

Calculi of this nature may remain in the system for years, setting up relatively little disturbance; or, on the other hand, may, by their passage, cause obstruction in the ureters, the orifice of the bladder, or the urethra (renal colic). There has been much debate as to whether they ever become spontaneously dissolved, for uric acid is, to some extent, soluble in alakaline solutions. As a matter of fact, the evidence seems to be conclusive that, through keeping the urine alkaline through long periods by giving sodium bicarbonate, alkaline soaps, etc., not merely is gravel arrested, but the stones within the bladder, after such treatment, show clear evidence of erosin. In examining any large collection or uric acid calculi obtained in the postmortem room, etc., certain specimens have a rugged, wormenten appearance, and, upon section, are found lose in

¹ Trans. Amer. Pediatric Assoc., 1903:150.

CALCIUM OXALATE CALCULI

texture, with evidence of lamination very indistinct. Upon analysis such calculi are apt to yield relatively considerable amounts of quadriurates. These calculi, is seems to us, are clearly undergoing slow solution, and the quadriurates are an evidence of the action of the alkaline urine upon the uric acid. Sometimes, also, relatively large calculi undergo spontaneous fracture. We had two well-marked examples of this in the collection at McGill University before it was destroyed, and i thas been noted ever since the time of Hunter. With such breaking up, the individual smaller pieces may be voided in the urine.

CALCIUM OXALATE CALCULI.

Calculi formed in the main of calcium oxalate are of three types. The commonest and the most characteristic is the so-called "mulberry calculus." In general solitary and occurring in the bladder, this form is relatively the heaviest of all calculi, densest and most resistant to fracture. The main feature is the heavily bossed or mulberry-like appearance. The color tends to be brown, or even brownish black. Often there is an incrustation of phosphates secondary to the great mechanical irritation induced by the presence of the heavy, irregularly nodular stone in the bladder. Upon section the appearance is also characteristic. The outlines of the concentric laminæ recall the plan of a fortress, with bastions and reëntering angles.

More rarely we encounter white, colorless oxalate calculi. These, upon fracture, have a more crystalline appearance, are relatively pure, and their surface shows small crystals of the salt. There may also be a condition of oxalate gravel. The minute concretions then are in the form of small, smooth, round bodies, blackish or dark gray in color. Occasionally multiple small oxalate calculi are found in the pelvis of the kidney, or occupying sacculations of the same, possessing spherical bodies with spinous processes recalling those compromising iron balls chained to a staff which princes of the church militant took into battle during the middle ages.

A greater or less proportion of uric acid is usually present in these oxalate calculi. The nucleus also is frequently of uric acid or urates. This combination can be easily understood when it is remembered that, like the urates, calcium oxalate is deposited from acid urine.

Regarding the conditions under which these are formed, it may be recalled that oxalic acid is normally present in the urine to the extent of some 50 milligrams daily. In certain conditions this amount is greatly increased, and that during long periods, so that a definite condition of *oxaluria* and the oxalic-acid diathesis is to be recognized. What is the essential cause of this is little understood. Under normal conditions it is found that vegetable food containing oxalic acid (rhubarb, etc.) causes an increased discharge, but this does not explain persistent oxaluria This oxaluria has been found in certain conditions of dyspepsia of a somewhat neurotic type, in confirmed obesity, and in association with

CALCULI AND CONCREMENTS

diabetes. Excess of cane sugar has also been noted to cause it.¹ According to E. G. Smith,² there is often a condition of hyperchlorhydria. It may, therefore, be that some perverted metabolism of the carbohydrates plays a part in the development of this state. Once formed, it would appear that oxaletes are not easily dissolved.

PHOSPHATIC CALCULI.

Two, or probably three, forms of phosphatic calculi are to be recognized, namely: (1) Those in which the phosphate is wholly in the form of calcium phosphate; (2) the mixed phosphatic or "fusible calculus" in which ammonium magnesium phosphate is the main ingredient, and in which calcium phosphate is also present; and (3) the pure ammonium phosphate calculus. The second of these is by far the commonest. All these are thrown down from alkaline urine, but the form of salt precipitated depends upon the cause of the alkalinity. It is where the urine is secreted in an alkaline state, the alkalinity depending more upon sodium salts, that calcium phosphate is precipitated. It is where the urine has become secondarily alkaline through fermentation and the bacterial breaking down of the urea into ammonia and carbonic acid gas that the triple phosphate is thrown down.

As a matter of fact, while calcium phosphate is freely precipitated in ordinary alkaline urine, it then shows relatively little liability to accumulate in the form of stones, and there may be, for months or years, a passage of such precipitated phosphates without any symptoms pointing to calculus formation. It may be pointed out that, under these conditions, the urine is not irritating, so there is no stimulus to the production of any excess of mucoid material to form the matrix; hence pure calcium phosphate stones are definitely rare. When found, they are white, rather firm-much firmer than the form about to be mentioned-and they rarely attain to large size. The mixed phosphates, on the other hand, are relatively loose and friable, the different layers splitting apart with fair ease, and they may attain to an enormous size. It is almost unknown to find such mixed phosphates as a nucleus of a calculus; the salts are usually deposited around other stones, or, again, around foreign bodies introduced into the bladder. It is rare also to find that other salts form the outer layer of phosphatic calculi, the reason being that, once cystitis is set up, where such stones are present, it continues, so that urine very rarely becomes once more acid. It is generally stated in text-books that these stones are not absorbed, and this because of the difficulty in rendering the urine acid by therapeutic means. With our knowledge that cystitis and the alkaline fermentation of the urine

¹ Thus, Dr. Helen Baldwin has found that by feeding dogs with large amounts of sugar a condition of mucous gastritis is set up with appearance of oxalie acid both in the stomach contents and in the urine. See Herter, Harvey Lectures, 1906-07: 84, and *Chemical Pathology*, 144.

² Ref. Handbook Med. Sci., article Calculi, 1903.

OTHER URINARY CALCULI

are of bacterial causation, and the recent production of drugs like urotropin, which are capable of sterilizing the urine, the possibility is certainly open that these calculi may become dissolved in the organism.

In such cases of cystitis, not only are we apt to find stones within the bladder or pelvis of the kidney, but on the inflamed mucous membrane there may be deposits of the phosphates.

OTHER URINARY CALCULI.

Outside these three main groups above mentioned, other forms of urinary calculi are distinctly rare. Of these, one of the most interesting is the *cystin* calculus. These are of a honey-yellow color, becoming green upon exposure to light; they have a radiate structure, and, upon fracture, a peculiarly transparent brilliancy, like that of beeswax. They are soft, so that they can be indented with the nail, and are usually pure. Sometimes they are large and single, but a condition of cystin gravel also occurs, in which the patient, from time to time, over long periods, voids minute calculi of this nature. The collection at McGill contained a set of eighteen small calculi passed by one patient in this way.

Cystin is an abnormal constituent of urine. It contains a relatively large amount of sulphur (as much as 26 per cent.), and dissolves in liquor ammoniæ. From this solution, as the ammonia evaporates, beautiful six-sided tablets separate out along with some square prisms, the substance being dimorphous. (See Fig. 137, p. 379. Its formula is $C_{\rm s}H_{\rm s}NSO_{\rm s}$, and thus it is not far removed from that of taurin.

What leads to its appearance in the urine is wholly unknown, save this, that it tends to be formed by more than one member of a family, and that the condition of cystinuria may show itself for long years without any impairment in health or obvious change in the other constituents of the urine, the only serious danger being a tendency to form calculi (see p. 379).


Still rarer are *xanthin* calculi, first noted by Marcet, of London, in 1817. Altogether, little more than half a dozen cases are on record of the formation of these calculi, which are of pale-yellow color, and exhibit a waxy lustre when rubbed. Their main constituent is xanthin, and nothing is known with regard to the conditions leading to extensive discharge of this alloxur body. In cattle, somewhat parallel calculi of *guanin* have been encountered. Yet rarer, again, are the concretions termed by Heller *urostealiths*. These are formed of fatty and soapy matter, lime soaps being apparently their main constituent. While some, obtained from the bladder, have been explained as due to the passing in of soapy medicaments, others have been found actually within the kidney, and so must have been derived from fatty matter discharged from this organ.

Lastly, there are at times to be met with small, firm, *fibrinous* concretions, more or less blood-stained, which clearly are secondary to hemorrhage in the urinary tract.

CALCULI AND CONCREMENTS

BILIARY CALCULI; CHOLELITHIASIS.

The composition of the calculi which may form in the gall-bladder and bile passages differs widely from that of the urinary calculi, and, as a body, they are characterized by being less dense and much lighter in weight. The main constituents are the modified pigments of the bile and cholesterin; a third constituent present not infrequently as an admixture, but rarely as a predominant constituent, is calcium carbonate. We thus distinguish the following varieties of gallstone.



Cholesterin calculus, cut and polished to show radiate crystalline structure. (Naunyn.)





Section of common mixed bilirubin calcium gallstone, (Naunyn.)



Pure bilirubin calcium calculi: bile gravel. (Naunyn.) Section of "amorphous" cholesterin gallstone exhibiting central cavitation. (Naunyn.)

1. Pure (or Almost Pure) Cholesterin Calculi — These are characteristic stones, most often single and oval, of white or yellowish color, waxy exterior, with finely nodulated surface. But two or, very rarely, three stones are encountered: our collection at McGill contained one set of five, partially faceted. If broken, these have a distinctly radial, crystalline surface; if sawed through and polished, an obscure crystalline, rather nacreous surface, with little or no sign or stratification. There may be a small nucleus of pigment. It is, in short, rare to obtain a calculus of pure cholesterin, but from 95 to 98 per cent. of the total contents may be of this substance. There is a minimal amount of associated calcium.

CHOLELITHIASIS

2. The Laminated Cholesterin Gallstone.—This is also a solitary stone, often attaining a large size, oval, whitish or yellowish, but not so transparent as the first form. On section it is laminated, although the centre may be of the radial type. Examined chemically it differs from the previous form in containing a fair amount of calcium carbonate, and, according to Bacmeister, it is the coincident laying down of chalk and cholesterin that alters the type of crystallization. The lamine are formed of closely set fine needles, radially disposed. German authorities unfortunately give to the next group the name that should rightly belong to this form, namely *Cholesterin-kalksteine*, when more truly that is the Cholesterin-bilirubin-calcium group. To avoid confusion we speak of these as laminated cholesterin. The amount of cholesterin varies from 75 to 95 per cent. Some of the layers may be deeply pigmented, although the total amount of bile pigment is small.

3. The Common Gallstones.—These may be (1) single and large, when they are often barrel-shaped (as though other stones had previously been present in series, and had compressed and faceted the two ends); or there may be two to four relatively large stones faceted on their apposed aspects, and forming together a cast of the gall-bladder; or (2)multiple small faceted stones, from a score or so to several hundred in number, relatively equal in size. It is more uncommon to meet with an admixture of large and small stones. The faceted surfaces are smooth; the larger stones, where the surfaces do not appose, are usually finely nodular. The external color varies from a deep brown that is almost black, through reddish brown (bilirubin), to green (biliverdin). Or, again, it often is pale yellow, through surface layers of cholesterin, or, rarely, is white, from a surface layer of calcium carbonate. On section, such calculi exhibit a characteristically concentric structure of successive layers of varying depth of color, according to the extent of admixture of cholesterin with the calcium salt of bilirubin or of biliverdin. The nucleus may be either of cholesterin or of bile pigment, or, as Naunyn points out, may be represented by a cavity¹ (Fig. 320).

Such calculi, it wilf be seen, are mixed, the main constituents are cholesterin and bilirubin calcium, although biliverdin, another higher oxidation form of bile pigment, may take the place of bilirubin; with these may be minute quantities of Cu and Fe, apparently in combination with the bilirubin calcium. Calcium carbonate is a not uncommon constituent, laid down in minute nodules throughout the mass of a calculus; more rarely, inconsiderable quantities of calcium sulphate and phosphate have been detected.

4. **Pure Bilirubin Calcium Calculi.**—These most often are present as what may be termed bile gravel—multiple blackish granules, averaging about 1 mm. in diameter, lying in a distinctly mucoid bile. When fresh they are very soft, easily crumbling between the fingers. They are difficult to dry and preserve, in consequence of shrinkage and

¹ Although most often this cavity, according to Bacmeister, is due to the drying out of the softer nucleus,

CALCULI AND CONCREMENTS

fracture into small fragments. Frequently, through the mucus in which they are embedded, they agglomerate into small mulberry-like masses. Naunyn describes another firmer form, with metallic lustre, which we have encountered but once, and that doubtfully.

Calculi of this class are composed almost entirely of the calcium salts of the bile pigments, with but a trace of cholesterin. In addition to the bilirubin salt, there may be large proportions of bilihumin calcium (bilihumin being the most oxidized of the bile pigments).

5. Calcium Carbonate Stones.—Stones formed almost entirely of calcium carbonate are rare and small in man; among the herbivora they are more common, and may attain a considerable size. Naunyn differentiates two forms in man—the one brown and spicular, the other pale and smooth, both of relatively great hardness. As already noted, nodules of calcium carbonate may occur more especially in connection with biliverdin calcium.

Another rare form is what may be termed cholesterin gravel—fine deposits or accumulations of what Naunyn regarded as amorphous cholesterin. This is the first stage of stone formation; later, the cholesterin becomes crystalline.

Etiology.—There have been abundant theories regarding the causation of gallstones; these have been largely forgotten since Naunyn's classical work gave a foundation of more thorough knowledge. The observations of Brockbank, Kramer, Aschoff, and Bacmeister have most materially advanced our knowledge.

It had for long been recognized that gallstones are more common (1) in middle and advanced life; (2) in females, rather than in males; (3) in those of sedentary, as distinct from active, habits. Frerichs more particularly laid down that stagnation of the bile is the main factor leading to their formation. The more recent observations have demonstrated that this is an essential although not, as Frerichs thought, the primary factor.

The main data upon which our knowledge of the etiology of gallstones is based are:

I. Bacteriological studies have demonstrated the frequent presence of bacteria in the ordinary (mixed) calculi. In 1802 Naunyn and Gilbert and Girode independently called attention to the relationship between the B. coli and cholelithiasis. Dufort, in 1893, found a history of a previous attack of typhoid in 19 cases of cholelithiasis, in 12 of which a first attack of biliary colic came on within six months after the fever. That same year Chiari called attention to the presence of typhoid bacilli in the gall-bladder, obtaining these in 19 out of 22 cases of cholelithiasis. This, according to later observers, is an excessive proportion. Very frequently the B. coli is found, at times the B, proteus. The presence of streptococci is not associated with the development of gallstones.

II. Experimentally, cultures of one or other form introduced into the unobstructed gall-bladder lead to no result. If by ligation obstruction be produced, cholangitis is set up and precipitation occurs in the bile. In

CHOLELITHIASIS

this way Cushing⁴ has produced gallstones by the injection of **B**, typhosus into the gall-bladder of the dog. On the other hand, the simple ligation of the uninfected gall bladder leads to absorption of the bile with no cholelithiasis.

III. Outside the body Kramer² has shown that if to filtered normal bile dilated with peptone broth there be added cultures of B. coli or B. typhosus there slowly appears (in the course of months) a deposit of all the constituents of gallstones—of bile pigment, cholesterin and calcium salts. These observations have been confirmed by Bacmeister,³ who has found that paratyphoid forms, B. pyocyaneus and yet other bacteria added to sterilized bile have the same effect. Our own experience is that whenever the bile removed by a pipette at postmortem is turbid with fine granular flocculi, that bile affords cultures of bacteria—of members of the B. coli group.

IV. As pointed out by Brockbank,⁴ bile removed postmortem, in which there is much desquamated epithelium, when first removed shows abundant cells; but rare crystals of cholesterin, if kept for a few days, may eventually show no cells but abundant deposit of cholesterin. Bacmeister shows that this same effect is brought about when gall-bladder bile to which scraped-off epithelium has been added is sterilized by heating to 59° C, on successive days. Sterile gall-bladder bile alone will show eventual precipitation of cholesterin, but it does this very slowly: the more the epithelial cells the more rapid and abundant the precipitation.

VI. Pure (as distinct from laminated) cholesterin stones are always obtained from bladders which at the most show signs of distension, but never chronic inflammation (Aschoff), and frequently, in our experience, the contents of the bladder in these cases are sterile. All other forms occur in gall-bladders exhibiting clear indications of chronic cholecystitis.

VII. If the deposited salts of gallstones be dissolved out there is left a matrix of albuminous or inspissated mucoid matter. The amount of this is minimal in the case of pure cholesterin stones in marked contrast to the other forms.

Weighing these data, we conclude:

1. That the separation out of the constituents of the bile in such a form and to such an extent as to favor calculus formation is a relatively slow process, and that therefore stagnation of the bile is an essential factor for cholelithiasis.

2. That the development of pure cholesterin calculi occurs in the absence of infection; the natural desquamation of the gall bladder epithelium, such as occurs with all epithelia, occurring where there is obstruction and retention of the bile favors the crystallization out of the cholesterin.

3. The amount of calcium salts in normal bile is inadequate to

⁴ On Gallstones or Cholelithiasis, London, Churchill, 1896 ; p. 21.

¹ Johns Hopkins Hospital Bull., 10: 1899: 166 (with bibliography).

² Jour. of Exp. Med., 9:1907:319.

³ Münch. med. Woch., 1908: Nos. 5, 6, and 7; Ziegler's Beitr., 44: 1908: 528.

explain that found in all the other forms of biliary calculi. Chronic inflammation and discharge of mucus is known to be accompanied by discharge of calculum salts. All these other forms of calculi are therefore the results of conditions of chronic or subacute cholecystitis.

4. As regards the common mixed gallstones, this chronic inflammation is of infective nature, due to the presence in the gall-bladder of, more particularly, members of the typhocolon group.

As regards the laminated cholesterin stones, we cannot as yet arrive at such conclusions, the relative absence of precipitated bile pigments is against these being of infective production; the presence of calcium carbonate in fair amounts is an indication of inflammatory changes in the gall-bladder. It is quite possible to conceive a chronic irritative inflammation of the gall-bladder of a non-infective type. It may be urged with some force that although differing in structure all calculi save the pure cholesterin are members of one series, differing not in their components, but in the proportions of the same, so that we may presume that they are all of infective origin, the different proportions of these constituents, being the outcome of an interaction between the grades of irritation set up by bacteria of varying virulence and the constitution of the bile in different individuals. This it will be seen is a matter of theory. The matter must be left open.

On the ground that experiments in vitro lead to precipitation out of the pigments and cholesterin only after an extraordinarily long interval, Bacmeister regards bacteria as a problematical factor in cholelithiasis, pointing out further that colon bacilli may be found even in pure cholesterin calculi. Even in the common calculi the bacilli may well have wandered in secondarily. Thus, we admit that it is not the actual presence of bacilli in the gallstones that has led to the formation of those stones, rather it is their presence in the bile, their action in modifying the chemical composition of the same, and more especially their action upon the mucous membrane of the gall-bladder and the cystitis they induce that is the essential cause. To us it is abundantly evident that the typhoid bacillus, for example, growing in the gall-bladder and there setting up a cholecystitis is apt to set up gallstone formation in young individuals twenty years and more earlier than the usual period of cholelithiasis. The bacillus is clearly the prime cause of the gallstone formation, and if in vitro in diluted normal bile, with little mucin and less calcium salts, these different bacteria nevertheless bring down the constituents of gallstones. much more are they likely to do so where these are adequately provided; in fact, as already noted, the precipitation is striking in postmortem bile which is infected; and this, we may add, where there is no evidence of obstruction and, save for the somewhat mucoid character of the bile, no signs of cholecystitis. In general we may lay down that it is a low and not an acute form of cholecystitis that favors gallstone formation.

Origin of Biliary Cholesterin.—Despite this material advance in our understanding of cholelithiasis, there remain several moot points regarding the process of stone building. Most of these revolve around the cholesterin component, whence it originates, whether it is laid down primarily or secondarily, its state of suspension or solution in normal and

CHOLELITHIASIS

pathological bile. Naunyn held that the greatly increased amount of this substance in gall-bladder bile as compared with that obtained by fistula from the hepatic bile ducts was due to active excretion from the gall-bladder epithelium. His pupils, Jankau and Thomas, showed that the amount of cholesterin in fistula bile compared with the other constituents remains remarkably constant and is not increased by feeding animals with cholesterin. Recently, however, Goodman¹ has shown that it is subject to considerable variation, and that a rich proteid diet, for example, leads to a pronounced increase in this constituent of the bile. Bacmeister in his experiments has found no evidence of discharge of cholesterin from the gall-bladder mucous membrane, and Aschoff has demonstrated that, on the contrary, when abundant cholesterin dissolved in oil is introduced into the ligated gall-bladder of the dog, it undergoes complete absorption within 11 days, while similarly if the gallbladder be emptied and then ligated the only cholesterin to be found in it at the end of a month is in the inspissated mass representing apparently the bile which had not been completely removed at the time of operation. The same able investigator has shown conclusively that fatty matters as well as cholesterin are absorbed by the gall-bladder epithelium and removed thus out of the bile. He thus joins with those who hold that the cholesterin of gallstones is furnished by the biliary excretion alone.

There are, however, data equally convincing on the other side. Thus, Bristowe² found crystalline masses of cholesterin embedded in crypts in the wall of a thickened and obstructed gall bladder. Brockbank also has found small accumulations of cholesterin lying in retention cysts in the mucous membrane. We ourselves in a case of hydrops cystidis fellew found the obstructed gall-bladder distended with thin colorless fluid, without a trace of bile pigment, and clear save for abundant shimmering crystals of cholesterin. The cause of the obstruction was a pear-shaped gallstone of fair size, the larger end, projecting into the bladder, being formed of crystalline cholesterin, the narrower end, imprisoned in the ducts, was of mixed formation and pigmented. The contracted shape of the neck suggested strongly that the stone at the time of obstruction had been of small size, and that the larger free end had been formed since that Nevertheless, it was impossible that the bile imprisoned in the gallbladder had provided all this mass of cholesterin in addition to the abundant free crystals floating in the gall-bladder contents. It was difficult to conclude other than that this cholesterin had in the main been discharged from the mucous membrane of the gall-bladder; just as it is difficult to explain the two previous observations, save on this assumption. Positive evidence of such discharge has been afforded by Herter and Workman,3 who have shown that the injection of minute amounts of corrosive sublimate or of phenol into the gall-bladders of dogs leads to an increase in the amount of cholesterin in the contained bile. Lastly, if lipoids and cholesterin are absorbed freely through the cholecystic mucosa it is difficult to explain why fresh bile from the gall-bladder is

¹ Beitr. z. Chem. Physiol., 9:1907. ² Lancet, London, 1887; 1: February, 19. ³ Trans. Triennial Congress Amer. Phys. and Surg., 6:1903:158. constantly from three to six times as rich in cholesterin as is that recoverable from a bilary fistula. This might be explained as due to resorption of fluid and concentration of the bile during its stay in the bladder, but the variation in the relative amounts of the other contents makes it very difficult to accept this assumption; the increase in fats and soaps is even more marked (about 13 times). The only means of harmonizing these apparently wholly opposed observations is to accept both, and conclude that just as enzymes are reversible, so-cell activities may be reversible: that just as the endothelial cells and walls of the venous capillaries are capable both of permitting a discharge of fluid from the blood and of absorbing fluid from the surrounding lymph, so the mucous membrane of the gall-bladder in one order of conditions takes up cholesterin and fatty matter from the bile, passing it over into the lymph, in another order of conditions takes up these bodies from the surrounding bloodvessels and lymph and excretes them.¹

I am inclined, therefore, to conclude that the cholesterin of gallstones is derivable in part from the liver cells, in certain conditions largely from the gall-bladder or bile duct epithelium.

How and why the cholesterin present in the bile passes from the soluble or apparently soluble into the crystalline state is still a matter of debate. According to Lichtwitz,² it is not dissolved in normal bile, but is in a state of colloidal suspension. Now, if two or more colloids in a solution possess the same electric charge, they keep each other in suspension; precipitation is induced when either another colloid possessing the opposite charge enters into the solution, or when a colloid already present by any means loses its charge. According to this view the action of epithelial cells in the bile in promoting the appearance of crystals of cholesterin is the introduction of another colloid of opposite charge as a result of the cell autolysis. Bacteria by causing various dissociations of colloids are supposed to have a like effect.

I am not prepared to say that this is the adequate explanation. The older view regarded the fats and soaps of the bile as playing an important part in the process, and I cannot so lightheartedly as Bacmeister, in his second paper, put on one side the evidence that there is an intimate relationship between the fats and cholesterin of the bile, disturbance of which leads to precipitation of the latter in crystalline form. These observations may be briefly noted. With olein, cholesterin forms compounds having the characters of myelin, developing in favorable conditions bodies having a double contour and doubly refractive (p. 915). Now. more particularly in the gall-bladder mucosa of cases of cholelithiasis, the cells are apt to exhibit abundant fatty globules together with other anisotropic globules; occasionally the cells are completely filled with the myelin bodies. Naunyn³ observed that through breaking down of the discharged cells these myelin bodies are liberated into the bile, where, swelling, they undergo fusion into small clumps. In these free glassy,

¹ Adami, Western Canada Med. Jour., Winnipeg: 1909 ; January,

² Deut, Arch. f. klin. Med., 92: 1907; see also Porgess and Neubauer, Biochem. Ztschr., 7: 1907.

³ Klinik der Cholelithiasis, Leipzig, 1892:25.

URATIC DEPOSITS IN THE TISSUES

highly refractile masses of what he termed "amorphous cholesterin" he saw the origin of cholesterin calculi, for in them he could see the development of the crystals, as he thought, by direct conversion. By the addition of a little weak acid he could see under the microscope the production of cholesterin crystals in these glassy clumps. Our fuller knowledge of the constitution of myelin bodies convinces us that the "amorphous cholesterin" is, if not cholesteryl oleate, at least a chemical compound between the two constituents.¹

As Brockbank has shown, we can reverse the process: place crystals of cholesterin in a solution of ordinary soap and a most striking picture is rapidly produced. Elongating villus-like doubly refractive processes form all over the surface of the crystals, and become detached into the surrounding fluid, so that the solid cholesterin is eventually converted into "myelin," which Brockbank recognized as identical with Naunyn's amorphous cholesterin. Thus, as Aschoff pointed out,2 the myelin cholesteryl compound present in the bile under the action of the alkalies of the alkaline bile undergoes dissociation; the fatty acid is liberated to form soaps (which are capable of absorption), and the cholesterin is precipitated in a crystalline form. Unlike Aschoff, I am inclined to hold that one and the same process is largely responsible for the mucin in the bile, the myelin "cholesteryl oleate," and the increased calcium content, namely, irritation or other disturbance of the mucous membrane. It is deserving of note that Lichtwitz's observations were conducted not on natural bile, but upon made-up mixtures containing cholesterin. In normal bile, we would lay down, the colloid present is not pure cholesterin but is this lipoid compound of cholesterin and oleic acid. This substance it is which is in a state of colloidal suspension.

URATIC DEPOSITS IN THE TISSUES.

Gout and the so-called uratic infarcts of the newborn are discussed at some length elsewhere (pp. 375 and 941). It is but necessary to recall here in due order the fact that there exist deposits of urates of sodium within the tissues, and these of two orders. In gouty states these are in the form of closely packed, fine, needle-like crystals of sodium biurate. The seats of election are primarily the cartilages of the joints and particularly of the more distal joints, so that where goutiness

¹ Craven Moore (Med. Chron., Manchester 47:1908:204) brings forward evidence to show that in the body cholesterin does not exist as a true ester of the fatty acids; that where the two are combined the fatty acid is present as "acid of crystallization." It must be said in opposition that while he supports his views with certain strong arguments, they are opposed to the conclusions of a considerable number of careful observers. What is of most importance is that cholesterin alone does not exhibit the "fluid crystalline" state; the combination of oleic acid and cholesterin does, as do also the myelin bodies of the bile. These, therefore, clearly are independent chemical bodies possessing different physical properties from cholesterin.

² Verhandl. d. deutsch, path. Gesellsch. (1906), 10:1907:166. Without knowing of Aschoff's work, I enumerated similar views before the Buffalo Academy of Medicine early in 1907.

is suspected a routine examination should be made *postmortem* of the metatarsophalangeal articulation of the great toe. In the more advanced conditions there may be accumulations or nodules of the deposit (tophi) outside the joints, either in cartilage elsewhere—e. g., in the pinna of the ear, or around synovial sheaths and tendons. We may possibly compromise over the long-standing debate as to whether tissue necrosis precedes (Ebstein) or follows the deposit (Garrod and others) by laying down that the infiltration is never primary in the living cells, but occurs in the inert, intercellular matrix, and that as indicated by the experiments of His¹ once laid down the crystals act as irritants or slow poisons, setting up eventual death of the surrounding cells.

The deposits in the medulla and collecting tubules of the newly born are totally different in appearance. They occur in the form of spherules of minute, doubly refractive spherocrystals² or spheroliths, present both in the collecting tubules and in their investing cells. What is the exact composition of these is still a matter of debate. We incline to the view that they are identical with the similar so-called amorphous urates of the urine, which have the same globular character, and according to Sir William Roberts are quadriurates of sodium (p. 378). Somewhat similar deposits have been produced in the dog³ and in the rabbit by feeding⁴ with adenin, but according to Nicolaier⁵ these are neither uric acid nor urates, but amino-dioxypurin, a body not present in human urine.

OTHER DEPOSITS.

Intestinal Concretions.—Besides those already described, showing deposits of calcareous salts, cases are on record of resinous gastric concrements. In certain of the herbivora these are derived from the food eaten—and such form the true oriental *bezoars*, to which, in the East, marvellous healing and preventive properties are still attributed. These bezoars are said to be obtained from the wild goat. In man there are rare examples of the formation of similar concretions in painters and others, who have taken to drinking spirit varnish for the alcohol therein contained.

In animals, also, notably cattle, sheep, and pigs, hair balls (ægagropiles) are not uncommon in the stomach and intestines, derived from licking their own coats and those of their fellows. These form felted masses, which may assume a large size. More rarely, the habit of hair eating is acquired by the human female, and the hair balls growing during the course of months and years may form a cast completely filling the stomach. Our collection at McGill contains two examples. Keenan has collected some thirty cases from the literature. Scotch museums contain examples of oat-hair balls—smaller, finely felted

¹ Deutsch. Arch. f. klin. Med., 63: 1899: 266.

² Adami and Aschoff, Proc. Roy. Soc. Biol., 78: 1906: 367.


³ Minkowski, Arch. f. exp. Pathol. u. Pharm., 41: 1898: 428.

⁴Schittenhelm, ibid., 47: 1902; 432. ⁵Zeitschr. f. klin. Med., 45: 1902; 359.

FATTY CONCRETIONS

masses, composed in the main of the minute hair-like processes derived from the outer scales of the oat in improperly prepared oatmeal. These are largely absent from modern properly milled oatmeal; but, recalling Johnson's dictum concerning that article of diet, it has to be noted that these oat-hair balls are still to be met with in horses, more particularly those fed on sweepings of flour-mills.

Fatty Concretions.—Besides those mentioned as rarely encountered in the urinary tract (urostealiths), similar soapy masses are occasionally passed in the stools of those consuming large quantities of fat or oil, where they may, at first sight, be mistaken for gallstones. Of allied nature is *ambergris*, that rare and powerfully scented wax-like body



Hair ball of the stomach. The hair forms a complete cast of the stomach and duodenum, (Case of Dr. James Bell, Royal Victoria Hospital, Montreal.)

found in masses from 1 to 182 pounds in weight, either floating in the sea, or, as has been known for generations to the New England whalers, in the last seven feet or so of the large intestine of emaciated and diseased sperm whales.¹ That this is of intestinal origin is further shown by the presence in the masses of numerous "beaks" of the giant octopus, which as readers of *The Cruise of the Cachalot* will recall, forms the food of the sperm whale. Another familiar example is the accumulation of cerumen which may form in the outer auditory meatus.

¹ See Schwediawer, Phil. Trans. Roy. Soc., 73: 1783, and Fawkener, ibid., 81: 1791. Mr. J. Y. Buchanan, to whom I am indebted for these references, tells me that the Prince of Monaco has confirmed these observations, seeing the lumps of ambergris voided in the death struggle of a sperm whale, which at the same time vomited the tentacles and body of a giant octopus.

CHAPTER XXXI.

PIGMENTATION AND PIGMENTARY CHANGES.

THE property possessed by various chemical compounds of being colored affords no adequate ground for bringing them together into one common class. The colored state is an accident to this extent, that, so far as we can see, it connotes no common underlying physiological feature. Bodies as wide apart as elements like iodine, and coal-tar products, like the aniline dyes, are equally colored. Thus, at first sight, it would seem unscientific to bring together into one common group the various pigmentary changes occurring in the organism. There is, however, a certain usefulness in so doing, for the more numerous and the more pigmented bodies present in the system, under both normal and abnormal conditions, are closely allied, while the remainder are so varied that it is not easy to group them according to any other scheme.

ENDOGENOUS PIGMENTS.

With this understanding we would proceed to redivide the colored substances of the body appearing under pathological conditions into two broad subgroups: (1) *Endogenous pigments*, the direct products of cell metabolism or disintegration, and (2) the *exogenous* colored matters foreign to the organism and absorbed from without. The endogenous we may further divide into (a) hemoglobin and its derivatives; (b) other metabolic pigments. Each main subgroup contains two orders of bodies, namely: (1) Soluble; (2) insoluble, or precipitated pigments.

It is usual in works dating from the days when morbid histology was considered as pathology to exclude very largely the first of these, and in this connection to discuss more particularly the pigmentary deposits. Such a distinction only leads to confusion.

Abnormal Pigmentation Due to Hemoglobin and its Derivatives.— Hemoglobin and its derivatives are so fully studied in works upon physiology and physiological chemistry that it is unnecessary here to do more than recall its presence in a soluble condition, not only in the blood corpuscles,but also in the muscles; its remarkable chemical properties; its constant liberation from dying red corpuscles, more particularly in the portal system (including the spleen); its disintegration, more particularly through the agency of the liver cells, with discharge into the bile of the iron-free portion of the pigment as bilirubin and other bile pigments. It is probable, though this is not wholly determined, that the urinary pigment, or *urochrome*, is likewise a derivative of the normal

ABNORMAL PIGMENTATION

disintegration of hemoglobin. Certainly, under pathological conditions, the liver and the kidneys are the two organs through which hemoglobin and its compounds are discharged from the blood.

It is further deserving of note, as contributing to a natural classification of metabolic disturbances, that hemoglobin is a conjugated protein after the type of the nucleoproteins and the glycoproteins recently discussed. It is a compound of a nitrogenous coloring matter—*hematin* $(C_nH_nN_FeO_i)$, with the basic protein *globin*.

Experimentally, by various means—injections of large amounts of water, dilute glycerin, potassium chlorate, arseniuretted hydrogen, toluylenediamin, certain acids, and by the transfusion of the blood or blood serum of another species of animal—it is possible to cause the red corpuscles to break up and liberate their hemoglobin, which then becomes free in the blood plasma, and may diffuse out of this into the various tissues, and, indeed, undergo absorption by various orders of cells. Similar destruction of the corpuscles may be brought about by severe thermal changes.

In disease we have numerous examples of a similar liberation, and may either encounter this hemoglobin in the unaltered state or find it modified. A somewhat frequent example of the former condition is seen in *hemoglobin-*, or, as it is often termed, *postmortem imbibition*. Here we find the heart valves and the intima of the aorta and larger arteries assuming a bright rose-pink color. This occurs more particularly in cases of general sepsis. Two conditions, it seems to us, have to be distinguished. In hot weather, with the rapid onset of decomposition, a staining of the intima may show itself, which is distinctly a postmortem change. In acute sepsis the same process is seen, even when the autopsy is performed within an hour after death. Here it is not merely postmortem, but, through the extreme toxic state, there has been destruction of the erythrocytes during the last hours of life, and it is a combination of antemortem and postmortem diffusion of the hemoglobin which leads to this characteristic appearance.

Certain poisons-some of them already mentioned-are apt to cause, in man, a similar rapid liberation of the hemoglobin, notably potassium chlorate, certain poisonous mushrooms, and snake venoms. The abundant observations of late years upon hemolysis have revealed the existence of a long series of agents causing this liberation of the blood pigment out of the corpuscles, among them an important group of organic substances -bile salts, soaps, and the specific antibodies developed by the inoculation of foreign erythrocytes and other cells. Where the destruction of the red cells occurs in the systemic circulation, the liberated hemoglobin may be discharged, unaltered, through the kidneys (hemoglobinuria); slightly altered, as after potassium chlorate (methemoglobinuria). Methemoglobin, it may be added, is apparently of the same composition as oxyhemoglobin, but the oxygen is more firmly combined, and the reaction is acid. The modification occurs often in the bladder rather than in the system before excretion. Where the destruction of the red cells has occurred, not in the blood stream, but in the tissues and cavities of the

PIGMENTATION AND PIGMENTARY CHANGES

body, the pigment before discharge may undergo still further change into hematoidin or urobilin (*urobilinuria*).

Care must be taken to distinguish between these different states and hematuria, in which we have to deal, not with the mere excretion of blood pigment, but with escape of blood into any portion of the urinary apparatus and the consequent presence of all the constituents of blood in the blood-stained urine.

Hemoglobin, as Miss Adams first demonstrated, may be discharged through the glomeruli of the kidneys, or also, as Afanassiew was the first to show, may be taken up by the cells of the convoluted tubules. In the latter case it is clearly modified, being present in these cells, not merely in a diffused form, but also in the form of fine brownish granules. These granules are iron-containing, with the iron in looser combination than occurs in hemoglobin proper.

Paroxysmal Hemoglobinuria.—A very remarkable condition, associated with liberation of hemoglobin, deserves more than passing notice. This is the condition first described by Dessler (1854) and by Dr. George Harley (1865), to which Pavy has given the name of *paroxysmal hemoglobinaria*.

In this condition there is a sudden appearance of urine, tinged, or, it may be, deeply colored, by the presence of hemoglobin. For an hour or two what urine is passed is thus colored, and the next passage may be perfectly clear and limpid. This little understood condition is not in itself fatal, even though two or three attacks per diem may occur over a considerable period. Patients have been known to be affected from time to time for as many as eleven years. When the condition first manifests itself it is noted that the paroxysms come on after slight exposure to cold, as, for example, the chilling of the body upon rising in the morning, and first and other attacks are most frequent during the winter months. In inveterate cases they occur during the summer also, and frequently without obvious cause. Yet it has been observed that, in these severe cases, living in a warm climate prevents the paroxysms. There is, indeed, a certain relationship between this condition and cyclical albuminuria. As Copeman points out, such cyclical albuminuria also follows chills, and is truly a globinuria, and not an albuminuria; while, in the condition we are now discussing, a condition of globinuria, associated with a rapid reduction in the number of red corpuscles in the circulating blood, may precede the appearance of hemoglobin in the urine. In both conditions the red corpuscles appear to be abnormally sensitive to temperature changes, and a feeling of coldness and shivering in the extremities frequently precedes the paroxysms.

That the condition is due to a liberation of hemoglobin from the corpuscles was first fully demonstrated by Küssner, in 1879, by withdrawing blood from the peripheral circulation during a paroxysm, and finding that, after coagulation, the serum was distinctly colored by hemoglobin. Yet clearer demonstration was afforded by Ehrlich, who showed that, after ligaturing the finger of a patient subject to these attacks, and diping it in cold, and then in hot, water, not only had the serum become

PAROXYSMAL HEMOGLOBINURIA

tinged, but, among the corpuscles, numerous shadows of red cells, which had lost their hemoglobin, could be distinguished. We owe to Donath and Landsteiner¹ most material advance in our knowledge of this morbid process. Briefly, they have shown that if the blood serum be centrifugalized from the blood of a hemoglobinuric patient and the erythrocytes be added to this at the body heat, no hemolysis ensues; on the other hand, if the serum be cooled, if only to room temperature, again no hemolysis ensues, but upon bringing the suspension to body temperature, the mixture becomes laked. From this and other studies they conclude that the serum contains amboceptors, which attach themselves to the corpuscles at a low temperature, but not at a high; but once attached, on rewarming, the corpuscles are acted on by complementary substances present in the serum, so that hemolysis is brought about. What is of interest is that the hemoglobinuric serum has the like action upon the erythrocytes of normal individuals-nay, more, these are even more susceptible to the action-a proof of the existence and development of a definite if slight grade of auto-immunity in the bodies of affected individuals. These observations have been confirmed by several observers.2 What leads to the development of these auto-hemolytic bodies is still undetermined. Donath and Landsteiner found that the blood of apparently normal rabbits occasionally affords the phenomenon, and suggest that they are of the nature of sports. Rössle calls attention to the fact that a large number of sufferers from paroxysmal hemoglobinuria give the history of syphilis, and when to this we add the data afforded by the studies upon Wassermann's reaction, namely, the increase in globulins in the serum of syphilitics, and the knowledge that these with lipoids are concerned in hemolysis (p. 549), this suggestion that syphilis plays a part in at least some cases is plausible.

A similar but more severe and more pronounced paroxysmal hemoglobinuria is seen in horses, and here also exposure to cold has been noted to play a part. This condition is sometimes spoken of as *azoturia*.

Intective Hemoglobinuria.—But not all the hemoglobinurias in cattle and in man are of this same obscure origin. Some are undoubtedly due to the effects of blood parasites, sporozoa. The organisms of ague or malaria, for example, would appear to be the essential cause of "blackwater fever" in man, though, according to Koch, an idiosyncrasy toward quinine, taken to ward off the ague, has also to be invoked. In Texas fever in cattle the piroplasma similarly leads to a dissolution of the erythrocytes and the discharge of hemoglobin.

Modified Hemoglobin.—Free hemoglobin undergoes extensive modifications in cases in which there is prolonged destruction of the red corpuscles; and, again, where there is localized hemorrhage in the tissues. As indicated by the succession of tints assumed by subcutaneous hemor-

¹ Zeitschr, f. klin, Med., 58; 1906; 173; Centralbl, f. Bakt., Orig., 45; 1907; and Wiener klin, Woch., 1908; No. 45.

² Widal and Rostaine (quoted by Rössle, Lubarsch's Ergebnisse, 13: 1909: Abth. 2, Langstein, ibid., see also Eason, Edin. Med. Journ., N.S., 1: 1908; 121.

PIGMENTATION AND PIGMENTARY CHANGES

rhage—in a "black eye," for example—there is a succession of modifications of its pigment constituent, the *hematin*, leading to eventual deposit in the tissues of two, or, more correctly, three substances—*hematoidin*, *hemosiderin*, and *hemofuscin*. Of these, hematoidin is a pigmented, rather ruby-red, iron-free substance, which may be present either in the form of definite rhombic crystals, or in a more granular state, with occasional faint indications of crystalline structure. This, it has been noted, occurs more frequently in relatively large hemorrhages, or, more correctly, in the more central portion of hemorrhagic areas; it may be found in cerebral hemorrhages, or in the hemorrhagic contents of corpora lutea.



Rhombic plates and needles of hematoidin. × 500. (Ziegler.)

It is soluble to a slight extent, and is supposed by some to be the cause of the yellowish pigmentation of the skin that accompanies severe paroxysmal hemoglobinuria and



From a section of the liver from a case of periicious anemia, treated by Perl's test to demonstrate the iron-containing pigment lying in the liver cells close to the bile capillaries a, and away from the blood capillaries b. (After Ribbert.)

other conditions like pernicious anemia, in which there is extensive destruction of the red corpuscles. Chemically it is identical with bilirubin and closely allied to urobilin.

Hemosiderin, on the other hand, is always amorphous, in the form of fine granules, and is iron-containing, so that it can always be recognized by certain microchemical tests. By treating sections with ammonium sulphide, the granules appear brownish black, through the formation of sulphide of iron (Quincke's test). The more beautiful test is the so-called Perl's test, which consists of treatment of sections with moderately dilute solutions of potassium ferrocyanide, followed by treatment with weak hydrochloric acid. By this test the iron granules become converted into cyanide of iron (Prussian blue), and each granule has then a characteristic rich blue appearance. Deposits of this hemosiderin are regularly to be encountered, more particularly in the outer zone of hemorrhagic areas, both free and in leukocytes or other cells. Cellular activity appears to be necessary for its development. It occurs also in various organs of the body after repeated hemorrhages, or where, from one cause or another, there has been extensive destruction of red

HEMOCHROMATOSIS

corpuscles-in the spleen, liver, pancreas, lymph glands, walls of the intestines, etc. More particularly under such conditions do the liver cells come to contain relatively great quantities of hemosiderin. This is particularly noticeable in pernicious anemia, and here, as Quincke, Hunter, and others have pointed out, the liver may contain as much as ten times the normal amount of iron. In addition to pernicious anemia there may be large deposits of hemosiderin in the liver and other tissues in the condition known as hemochromatosis.

What is the cause of this condition is still a matter of debate. Von Recklinghausen was the first to call attention to it and point out that, in the lighter stages, it is to be noted in the walls of the small intestines. which, as a consequence, assume a yellowish or light-brown tint. In more advanced cases it is associated with cirrhosis of the liver, fibrosis of the pancreas, and, not infrequently, with diabetes, in the condition known by the French as diabete bronzé. As Kretz has pointed out, and as Dr. Maude Abbott has confirmed,¹ a considerable proportion of cases of cirrhosis of the liver show an excessive deposit of iron-containing pigment in that organ. Dr. Abbott found this in not less than nine out of sixteen cases of portal cirrhosis studied in Montreal. But what is the cause remains doubtful. The condition occurs more commonly in men; indeed, Dr. Abbott's case is the only one so far recorded in the female. Opie² is of the opinion that hemochromatosis is a specific disease. This we are somewhat inclined to doubt, for, as already indicated, a large proportion of cases of ordinary cirrhosis show slight grades of the condition. This, however, I think must be admitted, that the deposits of the pigment indicate, as in pernicious anemia, a destruction of the red corpuscles extending over a long period, and the heaping-up of the pigment, more particularly in the liver, as an indication that the cells of that organ are incapable of dealing adequately with the iron-containing portion of the hemoglobin, which thus remains in a fixed state in the liver cells and other cells throughout the organism. That the destruction is of the nature of a subinfection is supported by the recognition by Blackader, of Montreal, Gibson, of Edinburgh, Stokvis and others of an acute hemochromatosis, or bacterial cvanosis, due to B, coli bacteriemia.

The indications are that this hemosiderin is an albuminate of iron in which the iron is relatively loosely combined. In livers containing a considerable amount of the hemosiderin turning blue by Perl's test. other granules can be seen of the same size and general appearance, which, however, are unaffected, remaining of a yellowish brown. These granules are known as hemofuscin, and, as pointed out by Dr. Abbott. employing the Perl test, with heat and somewhat stronger acids a certain proportion of these now take on a blue color. The probability, therefore, is that hemofuscin represents an albuminate of iron in which, as in hemoglobin itself (which also does not react to Perl's test), the iron is in a more stable and firmly fixed combination, though other observers regard it as hemosiderin which has lost its iron. Such hemofuscin is

¹ Journ. of Pathol., 7: 1901: 55. ² Journ. of Exp. Med., 4: 1899: 279.

⁶¹

PIGMENTATION AND PIGMENTARY CHANGES

962

found also in other conditions—in extravasations of blood at a late stage; similar brownish-yellow granules have been noted in certain gland cells of the stomach and intestines, of mucous and sweat glands. Whether these granules are all of the same order is debatable. We are inclined to regard them, with von Recklinghausen, as, in the majority of cases, clearly derived from hemoglobin.

The nature of the pigment seen within the muscle fibres in brown atrophy is still a matter of debate; from histological considerations, regarding the atrophy of the fibres with which the pigmentation is associated, it is most natural to regard the pigment as a product of dissociation of myohemoglobin. Lubarsch' and others have regarded it as a lipochrome (see p. 972), and undoubtedly from the heart with brown atrophy there can be dissolved out a certain amount of such fatty pigment. But Taranonkhine² cannot recognize that it is either a derivative of hemoglobin or a lipochrome, although he regards it as a proteid derivative. Yet it is deserving of note that Rosenfeld,³ studying the pigment of the muscles of the small intestines (the pigmentation which von Recklinghausen has regarded as the first stage of hemochromatosis), found in this so large a proportion of sulphur that he regarded as more essentially allied to the melanins.

As against this view we would point out the impossibility of adequately isolating this pigment; and secondly, the ease with which sulphur, in the form of sulphuretted hydrogen, diffuses from the bowel, and so is liable to be taken up and absorbed by the surrounding tissues and their constituents.

Pseudomelanosis.—In this connection we may note that a marked grade of pseudomelanosis is occasionally to be met with. Where there has been breaking down of red corpuscles, with liberation of hemoglobin and formation of hemosiderin, and, further, the presence of sulphuretted hydrogen, an iron sulphide is formed and the tissues become black. Such liberation of sulphuretted hydrogen occurs most often in the stomach during digestion and fermentation of the food. Thus, at autopsy, it is most frequently the stomach walls and the organs in the immediate neighborhood—the liver and the spleen—that exhibit the change. But sulphuretted hydrogen may also be liberated by bacterial activity in suppurating wounds, gangrenous extremities, etc., and a similar pseudomelanosis be so produced. Mere postmortem decomposition rarely seems adequate to induce the change; more often there has been during life local or general hemolysis, with setting free of hemorylobin.

Hematoporphyrin and Hematoporphyrinuria.—Like hematoidin, hematoporphyrin is an iron-free derivative of hemoglobin, or, more truly, of hematin. It is not, to our knowledge, found within the tissues, but minute quantities have been recognized in normal urine, and in certain conditions of disease the amount in the urine may be notably increased;

¹ Centralbl. f. Path., 13: 1902: 881.

² Roussky Arch. patol., 10: 1900: 441; ref. in Lubarsch-Ostertag's Ergebnisse, 1901.

³ Arch. f. exp. Path., 48:1900.

JAUNDICE

acute rheumatism and sundry hepatic conditions are liable to be accompanied by a slight grade of hematoporphyrinuria. The greatest discharge is liable to occur after the injection of certain drugs, such as sulphonal, whose acid constituent is supposed to be liberated within the system, and then to act upon free hemoglobin.

JAUNDICE; ICTERUS.

That there is a constant, if somewhat intermittent, excretion of bile is easily realized. It is more difficult to realize that the abundant pigment of that bile is derived from the never-ceasing destruction of red corpuscles. This is surely the case. The individual erythrocytes have but a short life period, apparently not exceeding four weeks; either they die in the blood stream-fade away, their hemoglobin dissolving out. and the colorless shadows later breaking up and dissolving in the blood plasma—or the corpuscles are taken up in a moribund condition by the phagocytic cells of the spleen and liver. Other endothelial cells, notably those of the lymph glands, possess the same property, although they exercise it to a less extent. The hemoglobin from these corpuscles, whether in an unaltered state or modified, is absorbed by the endothelium of the hepatic capillaries, is passed on to the liver cells, and is, by them, broken up. The exact stages of this breaking-up process, first demonstrated by Kühne in 1858, have not been followed; the result is that the iron-free portion of the pigment is discharged into the bile capillaries in the form of bilirubin, the pigment of the bile. What becomes of the iron-containing moiety under normal conditions is a matter of conjecture rather than of precise knowledge. It may be returned direct to the blood: it may pass into the lymph channels. Compared with the amount of bile pigment discharged, the amount of iron present in the washed-out liver is relatively small. There is no great storage of iron in a combined state in this organ; the amount in the bile is so small as not to be worth consideration. Evidently the iron of hemoglobin is a valuable asset, and is not lightly parted with by the organism. If discharged into the bile it is capable of being absorbed again in the small intestines, where, as Macallum, of Toronto, has demonstrated, albuminates of iron are taken up by leukocytes which have wandered on to the free surface and are, by them, conveyed into the blood and lymph stream.¹

Any disturbance of the regular discharge of altered blood pigment leads to a more or less definite grade of jaundice, by which we understand a condition in which the bile pigment, failing to be discharged duly from the liver, accumulates in the same until it regurgitates into the lymph and bloodvessels, and so, being carried, in a soluble state, to the other tissues of the body, it is absorbed by those tissues, causing them to assume a bile-stained appearance. It is this staining that is the essential feature of jaundice (French, *jaunisse; jaune*, yellow); the accompanying symptoms, itching of the skin, slowing of the heart, mental

⁴ For a discussion upon the fate of the iron liberated from hemoglobin, see also Morishima, Arch, f. Pharm. u. Path., 41 ; 1898 ; 291,

PIGMENTATION AND PIGMENTARY CHANGES

depression and melancholy (literally; black bile), etc., while due to hepatic incompetency and regurgitation of other constituents of the bile, must be regarded as subsidiary. They may not show themselves in the milder stages of the condition, and will be discussed later when treating of the general pathology of hepatic disturbances. That this staining is due to bile pigment is easily proved. The blood, the urine, and the tissue give, upon analysis, the chemical reactions for bilirubin, or, in some cases, for its more oxydized modification, biliverdin. That in all cases of true jaundice the liver is the organ in which this pigment is formed, if not wholly, at least in preponderating quantities, is shown by the fact that this organ is the first to be involved, and may exhibit a greenish color when other parts of the body are but slightly involved. Microscopically, also, it can be made out that the pigment is present in excess within the liver cells. There is clearly regurgitation or passage of this soluble pigment from the liver into the circulation, and when it passes into the blood it is taken up by and stains the various tissues.

Distribution.—The tissues are affected variously. At a very early stage the bilirubin may be detected in the blood serum, and at an early period also the kidney takes on a distinctly vellow color; for it is by the kidney that the excess pigment of the blood tends especially to be discharged, and it is the convoluted tubules, or certain portions of the same, which take an active part in this process. The cells of these loops are not only diffusely pigmented, but contain vellowish-brown granules, which are either granules of inspissated bilirubin, or, by some observers, are thought to be the small granules of the cell deeply pigmented. But, clinically, the condition is first noticeable in the sclerotics of the eye, and next in the mucous membrane of the mouth, being more particularly capable of detection over the hard palate. Then the skin becomes involved, taking on a pale vellow tinge. From this period onward, upon postmortem examination, it is found that the ordinary connective tissues throughout the body are especially affected. The spleen takes on a pronounced stain, as do also the vessel walls. Certain tissues are relatively unaffected, noticeably cartilage, the cornea, the brain, and nervous tissues (save in the very young). In the brain the perivascular lymph may be relatively deeply stained, but the nervous tissue proper remains colorless. The lungs also are not greatly involved, save where there is intercurrent pneumonia, when the pneumonic areas and the expectoration become jaundiced. The bile pigment passes into several of the secretions; first and foremost is the urine; next in importance is the sweat, whereby the underclothing of the patient becomes stained yellow. The saliva is not unfrequently affected, the milk more rarely; the tears are said to be free from pigment even in the most advanced cases, and, as evidenced by the pale color of the feces in those cases in which there is complete obstruction of the common bile duct, there is very little, if any, discharged from the glands of the stomach and intestines.

The tint of the skin varies according to the intensity and duration of the condition, from pale yellow, through an obtrusive sulphur yellow,

JAUNDICE

to olive green, or, in advanced cases, to a dirty greenish black. In those cases in which the cause of the jaundice passes away, the pigment becomes again discharged from the tissues; but this not immediately; it may be present, slowly fading, for weeks afterward.

Etiology.—What, now, is the exact nature of this process? Are all the conditions we include under the term jaundice due to staining with bile pigment; or, more exactly, is it essential that, when bodies of the nature of bile pigment cause a staining of the tissues, they have been formed in the liver itself? Or, on the other hand, can bodies of this nature be formed by the breaking-down of the hemoglobin in the circulation, and thus there be a hematogenous as well as a hepatogenous jaundice? As regards hepatogenous jaundice, how does the bile pigment gain entry from the liver cells into the circulation? Is it discharged or diffused backward from the liver cells, through the vascular endothelium into the portal capillaries; is there a process of distension of the bile capillaries and bile ducts, ending in the rupture of the same into the portal vessels; or does the pigment enter the blood outside the liver by way of the hepatic lymph vessels and the thoracic duct?

We will endeavor to answer these questions, if not in the order here given, at least in such a way as to give a clear indication of their relationship and importance. There can, in the first place, be no question that pronounced cases of jaundice are due to the production of bile pigment in the liver and passage of the same from the liver into the blood, for in these cases not only is the liver the organ which, to the naked eye, shows the earliest and most advanced pigmentation, but the microscopic appearances also fully bear out this conclusion. There is here abundant evidence of the obstruction of the outflow of the bile. All true jaundice, in fact, is of an obstructive nature, with regurgitation of pigment and other bile constituents into the circulation. We can distinguish the following:

1. Obstructive jaundice pure and simple (hepatogenous jaundice), with no primary blood disturbance. Cases of this order may be due to any one of the many causes which lead to complete or partial obstruction of the bile channels in any part of their course, from the hepatic lobule down to the duodenal papilla; congenital absence or narrowing of the main bile ducts; inflammatory swelling of their walls, with narrowing of the lumen (catarrhal jaundice); growths within the passage; presence of foreign bodies within the lumen, such as gallstones or parasites, enlarged lymph glands, new-growths, or inflammatory cicatrices compressing the bile ducts from without; spasmodic stricture of the ducts. This is, as it were, but the skeleton of the many causes which have been found operative in leading to obstructive jaundice.

In these cases not only do we have indications of distension of the bile channels and the filling of the same with bile, within the liver, but the liver cells show a diffuse pigmentation, together with the presence of deeply stained masses of pigment. And careful examination in advanced cases shows that this pigment deposit within the liver cells has a definite and characteristic arrangement. As first shown by

PIGMENTATION AND PIGMENTARY CHANGES

Nauwerck,¹ of Königsberg, and independently, a few months later, by Fütterer,² of Chicago, this pigment lies in and injects a system of intracellular channels which are in direct connection with the bile capillaries. This network encloses the nucleus of the cell, but never enters it.

How, then, does this pigment enter the circulation? The more usual method was first demonstrated by Saunders a century ago.³ He ligatured the common bile duct, and then was able to trace the lymphatics of the liver, distended with bile, up to their junction with the thoracic duct. It has also been clearly demonstrated by Vaughan Harley.⁴ If two dogs be taken, and in both dogs the common bile duct be ligated, and in one, in addition, the thoracic duct be also closed, in the one bile pigment appears in the other, with the thoracic duct closed, it may be eight or even fourteen days before there is any such discharge.

Clearly, the normal path by which the bile reaches the circulation is by way of the lymphatics. There is here a certain analogy between what occurs in the liver and what has been observed in the pancreas after obstruction of the pancreatic duct. When the pancreatic duct becomes overdistended, the pancreatic juice makes its way into the lymph spaces round about the ducts, and there leads to very definite disturbances. There would seem to be a similar passage out of the distended bile capillaries and ducts into the lymphatics. But, on attempting to repeat Vaughan Harley's experiment, it does not succeed in every case. At times, although the operation has been performed most carefully, bile may appear in the blood and in the urine within a day or two, and Ziegler demonstrated very clearly that, in advanced grades of obstructive jaundice, there may be rupture of the distended bile capillaries into the neighboring bloodyessels. This is probably what occurs in the exceptional cases above referred to. And lastly, there is the possibility to consider, that even within the liver cells there may be a diffusion or reverse discharge of the bile pigment into the blood capillaries of the lobules, and this would seem to be favored by a second series of extremely fine intracellular channels communicating with the blood capillaries. The existence of such was indicated by Nauwerck and, even more definitely, by Browicz; their actual existence and connection with the lobules has been shown by Professor Schäfer, of Edinburgh, in the rabbit's liver. It may well be that reverse currents might thus be set up in the liver cells themselves, with discharge through these fine channels into the blood. But of such a *parapedesis* we have no positive evidence. The prolonged period which may supervene before bile appears in the blood when the thoracic duct has been ligated would seem to contra-indicate any such reversal of secretion, at least in cases of simple obstruction.

¹ Deutsch, med. Woch., 1895, and Münch, med. Woch., 1897.

² Medicine, Chicago, June and July, 1898.

³ The Structure, Economy and Disorders of the Liver, London, 1809.

⁴ Arch. f. Anat. und Physiologie, Physiolog. Abt., 1893.

FORMS OF JAUNDICE

2. A second form of jaundice is secondary to extensive breakingdown of the red corpuscles in the circulation, with, as a result, overloading of the liver cells and excretion by them of a concentrated or inspissated bile, whereby the *fine bile channels* become blocked, and the bile pigment now makes its way into the general circulation. This we may term, with Afanassiew, *hemohepatogenous jaundice*, or, as it is frequently now termed, *toxemic*. It may be manifested in all those conditions which lead to excessive destruction of the red corpuscles—in acute septic disturbances, for example.

In these cases the jaundice is not so severe as in the preceding group; there is not complete obstruction, the feces remain colored, and the gall-bladder may contain a thick, intensely black bile. Experimentally, the condition can be produced, as Hunter and others have pointed out, by the employment of toluylenediamin and other drugs which set up great destruction of the erythrocytes. Recent studies by Widal, Riesman,¹ and others have called increasing attention to the state of undue fragility of the erythrocytes, either present from earliest years and familial, or acquired in later life, characterized by persistent discoloration of the skin, with recurrent conditions of more severe icteroid states, if not actual jaundice. The general symptomatology is that of the type here described. There is still debate as to whether this is a true jaundice—*i.e.*, whether obstruction to the discharge of bilirubin is an integral part of the condition, or whether this is more correctly to be placed among the conditions of hematogenous "urobilin" pigmentation, to be noted on the next page.

3. A similar grade of jaundice is set up in another group of conditions, namely, in acute yellow atrophy of the liver, phosphorus poisoning, and acute infective jaundice, or Weil's disease. It is questionable where to place this group. There is here, clearly, an acute toxic condition chieffy affecting the liver cells and not affecting the blood. Probably it should be included in the purely hepatogenous group, for the disturbance or obstruction is in the liver cells themselves, or in the finest capillaries.

4. Lastly, there are cases on record of the rapid supervention of jaundice after severe shock and nervous disturbance. The condition develops so rapidly that it is difficult to suppose that it is due to any spastic contraction of the ducts, and there must either be rapid concentration of secretion, with blocking of the finer channels, or reversed circulation, as already indicated when discussing the intracellular passage in the liver cells.

Hematogenous (Urobilin) Pigmentation.—Can bilirubin or any allied derivative of hemoglobin be produced in the circulating blood outside the liver, and this to such an extent that the tissues become pigmented? We know that hematoidin, which, as Neumann has shown, is identical with bilirubin, can be formed in tissues (see p. 960). We know that it has been detected by Naunyn and Minkowski, Löwitz, and others actually within the circulating leukocytes. We recognize, also, that in cases of known destruction of the red corpuscles within the vessels—

¹ Trans. Assoc, Amer. Physic., 1910.

PIGMENTATION AND PIGMENTARY CHANGES

in paroxysmal hemoglobinuria, in pernicious anemia, in hemochromatosis, in hepatic cirrhosis, in sepsis, etc.—the skin is apt to assume not, it is true, a frank jaundiced hue, but at least a distinct tinge, varying from pale lemon yellow to ashen or bluish gray (as in advanced hemochromatosis). Whether we are dealing with one and the same substance in all these cases is, perhaps, doubtful; but it is evident to every clinical observer that in this group of cases we have pigmental changes which appear to form a distinct group by themselves. And in this group of cases, when we come to make postmortem examination, the liver in general is not jaundiced, and, when examined microscopically, its cells exhibit no excess of bile pigment, nor is there any evidence of inspissation in the bile capillaries. We say "in general," because there is a certain proportion of such cases of more acute type in which the livers exhibit the characters associated with obstructive jaundice. The ordinary septic liver, for example, is not jaundiced, even though the skin be of a lemon tinge; but in some cases of sepsis we have all the indications of an acute hepatitis, with disturbed secretion and accumulation of bile pigment within the cells.

Now, in this group of cases, judging from the blood counts, there is abnormal and continued destruction of the red corpuscles, and the urine tends to be high colored and to contain urobilin, which analysis shows is a modified form of bilirubin, a reduction product of the same, closely allied to, if not identical with, hydrobilirubin (bilirubin = $C_{32}H_{36}N_4O_6$; hydrobilirubin = $C_{32}H_{36}N_4O_6$; hydrobilirubin = $C_{32}H_{36}N_4O_6$; hydrobilirubin e dation the intestines,¹ and in others directly produced in the liver, it is also admitted that it may be due to production from blood pigment in the organism independent of the agency of the liver.

But, these facts notwithstanding, the accepted teaching of the present day is that all these conditions of pigmentation are also hepatogenous and that the pigment is derived from the liver. We believe that this teaching is wrong. It is based upon certain apparently most decisive observations of Naunyn and Minkowski (1886) upon the goose, of Stern upon the pigeon (1885), and on other confirmatory observations upon the frog. In these animals not all the portal blood passes through the liver; there is a collateral vessel carrying a portion directly into the inferior vena cava. It is thus possible to close off or extirpate the liver without completely arresting the portal circulation. An animal so treated may, under favorable conditions, continue to live for a length of time sufficient to make observations about its metabolism. Naunyn and Minkowski found that if, for instance, they allowed geese so treated to inhale arseniuretted hydrogen (which ordinarily brings about great destruction of red corpuscles), no jaundice was set up. And, further, no bilirubin or urobilin was excreted from the kidneys. Hemoglobin alone was discharged. In other words, for the formation of these pigments the liver is indispensable.

The more recent observations by Croftan² appear to explain this

¹ The main fecal pigment, stereobilin, is now recognized as identical with urobilin.
² Phila. Med. Jour., 1902: 75 and 142.

non-appearance. According to him, the breaking-up of hemoglobin to bilirubin within the system is brought about by tryptic ferments in the presence of a carbohydrate (glycogen or dextrose). It is in the liver, he points out, that this combination of ferments with carbohydrate and hemoglobin is most liable to occur. If the liver be removed, any excess of glycogen or carbohydrate within the blood or tissue is immediately utilized. Hence the above experiments do not demonstrate that all bilirubin is formed within the liver. The liver, it is true, is a factor, and, when it is present, bilirubin may be formed within its cells. But it may be formed in other parts of the organism—wherever, in fact, free hemoglobin is present along with dextrose and trypsin. And trypsin has been detected within every organ.

This explanation harmonizes the data in our possession. There may be a true hematogenous pigmentation of the tissues with bilirubin derived from hemolysis within the vessels, without any participation by the liver, save that this organ supplies to the blood the carbohydrate necessary for the conversion. This is not jaundice proper; for want of a better term, we may speak of it as *hematogenous pigmentation*.⁴ If the hemolysis be more intense, then the overstimulation of the liver cells leads to a disturbance of their function, obstruction of the finer bile channels, and regurgitation into the circulation of the bilirubin produced within the liver cells (toxemic hematogenous jaundice). And lastly, and this is the most pronounced form, there is the purely hepatogenous or obstructive jaundice developed without any primary blood disturbances.²

OTHER ENDOGENOUS PIGMENTATIONS.

Melanotic Pigmentation.—Of the autochthonous pigments not derived from hemoglobin, the most important is melanin. Melanin —or, perhaps more correctly, the melanins, for the divergent analyses suggest that we have to deal with not a single body, but with a group is characterized by absence, or minimal quantity, of iron and, with rare exception, by relatively considerable sulphur content.

The variation in the sulphur present is striking; as high as 10 per cent, has been recorded in the melanin from some cases of melanotic sarcoma. Abel and Davis³ found from 2 to 4 per cent, in that from skin and hair, but that from the choroid of the eye has been found free from sulphur. As regards the iron, it would appear that the more the melanin is purified the less is the amount of iron detectable.

¹ We use this term in preference to *urobilin pigmentation* because, while urobilin appears in the urine in this order of cases, evidence is wanting that this is the pigmentcirculating in the blood and deposited in the tissues. Saying this we would not imply that the pigment is not urobilin, which as a reduction product of bilirubin might well be present in the tissues; merely that in the absence of sure knowledge it is more accurate not to employ this label.

² For a fuller study of jaundice, Hunter's article in Allbutt's System, vol. iv, may well be recommended.

³ Jour. of Exp. Med., 1:1896:361.

970 PIGMENTATION AND PIGMENTARY CHANGES

According to Abel and Davis, to whom we are indebted for the most thorough study of the pigment of the negro's skin and hair, melanin granules are insoluble in dilute alkali, dilute hydrochloric acid, alcohol, or other solvents, in the order here named, although after treatment for some days with dilute hydrochloric acid, dilute alkali now causes them to give up their pigment, leaving behind fine shadows of an organic substratum. What is more, these observers detected a definite amount of silicates in the granules, from both the skin and the hair. It exists normally in the choroid coat of the eve, in the deeper cells of the Malpighian layer of the skin, as also in certain cells-chromatophoresof the upper layers of the corium, and is found also in the membranes of the brain, more particularly in the neighborhood of the choroid plexus. Its coloring power is intense. Abel and Davis calculate that the entire skin and hair of the negro do not contain more than 1 gram of the substance. In melanotic growths, however, it is present in great quantities; from the affected liver alone in a case of melanotic sarcoma as much as 300 grams have been gained. It may be recalled that it is not only in the colored races of mankind that it is present in the skin, but in all human beings, with the exception of *Albinos*, as those are termed who exhibit an inherited lack of melanin formation.

What we regard as the normal production of melanin in members of the human family varies within wide limits, the fair-haired Saxon and the swarthy negro representing the extremes. What we regard as abnormal is, with the one striking exception, not very extreme. A physiological increase in the pigmentation is observed in pregnant women in the increased color of the areolæ around the nipples; this pigmentation, or chloasma uterinum, affects also other areas already pigmented, and is most marked in those having already dark skinsbrunettes. A somewhat similar pigmentation, but of irregular distribution, is observed in many cases of exophthalmic goitre and certain neurotic states (melasma). What is generally regarded as a pigmentation of the same order, and still more marked, is encountered in Addison's disease. But the most extreme abnormal development of melanin is associated with the development of new-growths and of melanotic tumors (see p. 825). Where these are extensive and rapidly developing, there is an escape of pigment into the blood (melanemia) and discharge through the kidney (*melanuria*). This excretion may be either of the fully formed melanin or of its chromogen: the urine, at first relatively colorless, taking on a dark-brown color on standing or after treatment with certain reagents. Among the domestic animals, notably the horse, a condition has been described not found in man, namely, a diffuse melanosis, in which pigment-containing cells are found throughout the tissues. Such melanosis and the presence of melanotic tumors have been found to affect white and not dark-colored horses.

Von Fürth,¹ to whom we are indebted for the fullest recent study upon

¹ Centralbl. f. Path., 15: 1904; 617; see also von Fürth and Schneider, Hofmeister's Beiträge, 1: 1901; 229.

MELANIN AND OCHRONOSIS

the nature of melanin, has brought forward an ingenious and plausible theory regarding the nature and origin of melanin. French observers, more particularly, have called attention to the existence of oxidases within the living tissues, through whose action certain proteins are darkened. It has, for example, been shown that the browning of the cut surface of an apple is due to this process, while, similarly, the conversion of the relatively colorless juice of certain Japanese plants into black lacquer by exposure to air' has been found to be due to the presence of oxidases which act upon the tyrosin and other aromatic products of protein decomposition. Experimentally, also, it can be shown that the action of strong acids upon proteins produces a dark-brown substance-"artificial melanin"-which is regarded as produced from the trytophane, tyrosin, and other aromatic bodies resulting from proteid decomposition by the addition of oxygen. Von Fürth would regard melanin and the melanoid bodies as developed by the action of intracellular oxidases ("tyrosinase") upon the aromatic or chromogen groups of the protein molecule. In favor of this view is the fact that a tyrosinase has been shown to be present in the ink sacs of cuttlefish, the pigment developed in these sacs, sepia, being allied in composition to melanin. And he would regard both the sulphur and iron as combined secondarily. It is somewhat against this view that tyrosin is not one of the products gained from the decomposition of melanin, although indol and skatol are obtained. Indol, therefore, and the allied bodies, rather than tyrosin, would appear to be involved in the process.

Independently what would seem to be a striking support to this view has been adduced by W. L. Halle.² He has demonstrated that, under the influence of an enzyme contained in the adrenal, tyrosin is converted into adrenin. We would point out that it is when the adrenal or its secretion is deficient that the characteristic pigmentation—bronzing—of Addison's disease shows itself. If the above view be correct, that pigments of the melanin group are of the nature of members of the aromatic series of derivatives of the protein molecule, then the bronzing gains its explanation: it is due to the want of conversion of tyrosin and allied bodies in the relative absence of the adrenal and to their consequent accumulation in the tissues, and we would add the greater darkening of the superficial parts most exposed to light and air gains its explanation from the more active oxidation of these "aromatic" bodies in these regions.

In the section upon neoplasms the nature of the cells containing the melanin has already been discussed (p. 829).

Melanin, or the melanins, are also the cause of the color of the hair.

¹ For the fullest data upon the oxidases with review of the literature, see Kastle, *The Oxidases*, Publications of the Department of Hygiene, Washington, 1910.

² Hofmeister's Beitr., 3 Chem. Physiol. u. Pathol., 8: 1906; 276. I owe this reference to Professor Schäfer's lectures upon the adrenal, Brit. Med. Jour., 1908; 1; 1281. These and Dr. Rolleston's address, Montreal Med. Jour., 36: 1907; 671, give an admirable summary of the present status of knowledge regarding the adrenals.

PIGMENTATION AND PIGMENTARY CHANGES 972

We do not encounter pathological-as distinguished from artificialexcess of hair pigment, although one of my students has called my attention to the fact that after erysipelas his hair, which had been of a light brown color, became jet black and curly; only gradually in the course of six years is it returning to a brown color, being still darker than it was originally. A few similar cases are recorded in the literature. The opposite condition of loss of pigment-turning gray, or canitiesis common. According to Metchnikoff, this process is essentially brought about by the increased phagocytic activity of the epidermal cells of the medullary layer of the hair, cells which Metchnikoff terms *piqmentoph*ages. We cannot but regard this as at most a partial explanation. It may well explain certain cases of loss of color, but some cases of white hair, like leukoderma of the skin, are surely due to failure on the part of the cells of the hair bulb to assimilate or elaborate the melanin. Following von Fürth's theory, it may be suggested that there is in these conditions a lack of that intracellular oxidase or tyrosinase, whereby the chromogen or melanogen is converted into melanin. In this connection it is suggestive that Spiegler⁴ has isolated from white hair and wool a body closely related to melanin, which he regards as a white chromogen or melanogen.

Ochronosis.-Possibly allied to the melanins-although its nature is still a matter of debate—is the pigment which, in very rare cases, causes a striking blackish discoloration of cartilage. In the sixties Virchow described the first case noted; since then scarce half a dozen cases have been recorded. The tendons, tendon sheaths, and synovial membranes may also be involved. The pigment is iron-free.

Hansemann's² case had associated with it the conditions of melanuria; Hecker and Wolff and Pick have added other cases, if not of melanuria proper, at least of darkening of the urine upon standing; and the latter, in a very thorough study of the condition, comes to a conclusion closely allied to that of von Fürth regarding melanins, namely, that the pigment in ochronosis is derived from aromatic compounds through the action of tyrosinase.

Lipochromes.—'There is a little understood series of colored fatty bodies occurring in normal tissues. In the human body these more particularly give the color to the fat of the organism, and one of them lutein-is present in considerable abundance in the cells of the corpora lutea. The pigment that accumulates in the nerve cells in advancing life, and under certain pathological conditions, would seem to belong to this order of bodies; as also, according to some observers, that of brown atrophy of muscle cells, though, possibly, in this last case we have to deal with combinations between fat and derivatives of hemoglobin. What is apparently a true lipochrome is the light-yellow, fatty body present in the cells of *xanthomas* (p. 723).

There is yet another form of neoplastic growth-the chloroma-in

¹ Hofmeister's Beiträge, 4:1903:40.

³ Berl. klin. Woch., 1892; Pick, ibid., 1906: 478.

PNEUMONOKONIOSIS

which the characteristic (green) pigmentation is evidently of fatty nature. We have discussed this elsewhere (p. 737), and here would only note that, according to Dock and Huber, the pigment is dissolved by ether and alcohol. On exposure to the air it loses its color.

We have encountered a pale greenish discoloration of a fatty tumor, of another order, not to be mistaken for chloroma proper, namely, in the neighborhood of a trocar puncture into a large abdominal lipoma. The discoloration here was evidently secondary to hemorphage.

EXOGENOUS PIGMENTATIONS.

Of the exogenous pigmentations, three main groups are to be distinguished:

1. Colored substances gaining entrance into the organism in a solid state and becoming deposited here and there as collections of colored particles.

2. Colored substances entering in a state of solution, becoming absorbed by the cells of one or other tissue, and so staining them.

3. The colored derivatives or decomposition products of substances themselves not colored, absorbed by the organism.

1. The most familiar example of the first of these groups is exhibited in the widespread practice of *tattooing*, in which insoluble particles of colored matter—charcoal, vermilion, etc., are rubbed into fine punctures of the skin in such a way as to produce designs or patterns of varying degress of crudeness or elaboration. In this way there are deposited in the outer layer of the corium collections of isolated particles. Though the tattoo marks may last a lifetime, they gradually become paler, there being a slow transposition of the particles along the lymph spaces and channels by the agency of the leukocytes. The pigment is always to be found in the nearest lymphatic glands. It is further possible to cause almost complete disappearance of the marks by inducing a cutaneous inflammation of some duration. That inflammation causes an active determination of leukocytes to the part and accelerates the removal of the pigment.

Of more serious import is the group of inhalation pigmentary deposits, the so-called *pneumonokonioses*, deposits in the lung tissue or elsewhere of colored particles inhaled in the form of dust (zozz, dust). The commonest of these is *anthracosis*, the deposit of coaldust or **qf** soot, present to a moderate degree in the lungs of every adult town dweller, and present to an extreme grade in the lungs of miners working in the dusty soft-coal mines. These deposits, it is needless to say, are coal black. Of a more gravish color are the deposits of siliceous particles in the lungs of quarrymen and workers in granite and other hard stones (*chalicosis*-zuzz, a pebble—or *silicosis*). The lungs of knifegrinders, glasspolishers, and others subjected to iron or iron oxide dust, take on a rusty red (*pulmonary siderosis*); workers in the potteries, inhaling kaolin or clavdust, obtain similarly dirty white deposits in the lungs

974 PIGMENTATION AND PIGMENTARY CHANGES

(aluminosis); workers in tobacco obtain rusty brown lungs from the tobacco dust (tabacosis), etc.

By causing animals to inhale air laden with one or other of these dusts, and by studying their lungs at successive periods, as, again, by an examination of sections of the lungs of human beings affected by these disturbances, the process by which the deposits are formed can be well followed. Where the air is full of dust, not all the particles are arrested by the moist lining of the nasal passages and the pharynx. As a result some particles are conveyed into the pulmonary alveoli. Unlike the bronchi and bronchioles, in that it is not ciliated, the epithelium lining the air sacs is unable to expel these solid particles, which would remain within the sacs were it not for the phagocytic activity of the epithelium and, more particularly, of scavenging leukocytes, which make their way from the vessels into the air sacs. Free cells can be seen in the alveoli laden with these foreign particles. While some of these wandering cells make their way into the bronchioles, and so are discharged with the sputum, others wander back into the lymph spaces of the alveolar wall and from thence to the lymph channels. In either of these positions the leukocytes may break down and the contained pigment be taken up by the endothelial connective-tissue cells of the region; or the breakingdown process, with liberation of the particles, may not occur until the lymph glands are reached at the root of the lungs. It is along the course of the lymph channels that the interstitial deposits mostly occur, namely, in the interlobular lymphatics and in those around the bronchi and the pulmonary vessels. There is also a peculiar liability for the pigment to be deposited where the interlobular lymphatics approach the surface of the lung to join the subpleural network of the lymph channels. As already stated, it can be seen that the endothelium of the air sacs also takes up these foreign particles. What happens to these has not been so clearly followed.

In support of the contention now urged by not a few workers (we think excessively) that pulmonary tuberculosis is most often secondary to the taking up of tubercle bacilli from the intestines, Calmette has recently published observations to the effect that the pneumonokonioses are due not to the direct inhalation of particles into the lungs, but to a swallowing of the same and selective collection of the same in the lung tissue through the agency of the leukocytes. A large number of observers have been thus stimulated to researches on the subject, with the result that Calmette's conclusions cannot be accepted, and the mechanism here laid down has become more surely established.⁴

The foreign particles act as mild irritants. Certain cells of the connective-tissue type seem particularly to take them up, and become, as a consequence, enlarged. Eventually there is a development of new

¹ Vide Calmette, Ann. de l'I. Pasteur, 19: 1905; 601, 20: 1906; 353 and 609; as also Whitla (and Symmers), Brit. Med. Jour., 1908; ii: 61; Bulloch, Allbutt and Rolleston's System of Medicine, 5: 1909; 299, and Cobbett, Jour. of Pathol., 14: 1910; 563.

PLATE XVIII


FIG 1



Two Sections from the Same Kidney of a Rabbit Treated with Injections of Corrosive Sublimate. (Klotz.)

Fig. 1.—Section stained with Sudan III to demonstrate fatty degeneration of certain tubules.

Fig. 2.—Section stained with silver nitrate to demonstrate calcareous deposits in the same groups of tubules. (By combined staining it could be shown that the identical tubules took on both the fatty and the calcareous reactions.)



EXOGENOUS PIGMENTATION

connective tissue in their neighborhood, with fibrosis or interstitial pneumonia. This may be both diffuse and nodular, so that masses of new tissue resembling tubercles may be formed around larger accumulations of the particles. It is noteworthy, in some districts, at least, that these changes are frequently followed by tuberculosis proper, so that a combination of anthracosis, or of stonemason's phthisis, with true tuberculosis, is often encountered. Nor are these deposits confined to the lungs. At times, even in the absence of adhesions, they may be observed in the parietal pleura, the pigment having evidently been conveyed by the leukocytes across the pleural spaces. And, in advanced cases, they are to be detected in other organs. Thus, we have encountered clusters of silicious particles with an obscure development of fibrous tubercles around them in the liver of a stonemason. Here it may be observed that not all dusts necessarily set up these conditions. W. P. Kaufmann¹ has shown that starch packers, although exposed to an atmosphere laden with floating starch granules, are remarkably free from pulmonary disturbances after the first few days, and from experiments on guinea-pigs, concludes that inhaled starch is dissolved by an augmented secretion of the amylolytic enzymes present in the blood. Millers, on the other hand, are affected, the gluten of wheat not being readily dissolved.

2. The second group, that, namely, of absorption of colored matter in solution, with staining of the different tissues, is of only experimental interest. As first shown by Daddi, certain of the aniline colors, such as Sudan III, used commercially to color waxes and fats in candle-making, when given by the mouth are absorbed, and, gaining entrance to the circulation, they color fat cells *intra vitam*. Recently, in his studies upon trypanosomiasis, Ehrlich has found that trypanroth, given to rats, while destroying the trypanosomes in their blood, at the same time colors the skin and other tissues of the animal a very distinct red. Such diffusible coloring matters may be discharged in the urine, the milk (p. 911), and other excretions.

3. In the third group of pigmented decomposition products the commonest example is a blue line on the gums in cases of chronic lead poisoning. Lead may enter the system either through the *digestice tract*—as in drinking soft water which has been conveyed in lead pipes; through the *respiratory system*, as in a series of cases observed recently in the Royal Victoria Hospital, in which several members of a family became the victims of acute lead poisoning as the result of using old white lead barrels for fuel in a defective stove; or, it would seem, by alsorption through the *skin* of workers in lead and lead paints. The blue line, when examined after death, is found to be due to a deposit of fine, brownish-black granules in clusters in the subepithelial connective tissue of the gums. The clusters apparently indicate endothelial and other cells. In those with clean mouths and well-brushed teeth the blue line is often wanting. It is more pronounced at the bases of decaying teeth, or, where the teeth are badly kept, near the accumulations of

¹ Jour. of Hygiene, 1909 : 220.

PIGMENTATION AND PIGMENTARY CHANGES

debris of food at their bases. What happens in these cases is that the sulphuretted hydrogen liberated from the food material diffuses into the tissues of the gums and acts upon the soluble salts of lead which have diffused out of the blood into the lymph spaces of the gums. The granules are a precipitate of insoluble sulphide of lead. Rarely a dirty greenish line has been noted on the gums of workers in copper.

Another metallic deposit is seen in argyria. Thirty years or so ago a treatment for epilepsy came into vogue, consisting of rather small doses of silver nitrate. This mode of treatment ceased when it was found that the unfortunate patients assumed an earthy-or unearthly-bluishgray color, and this of a most unfortunate permanency; for those who have survived their epilepsy, and the treatment, are today as blue, or almost as blue, as they were at the expiration of a few months. Experiments upon the lower animals to determine the cause of the phenomenon have demonstrated that soluble silver salts given by the digestive tract and absorbed into the circulation pass into the lymph. In the ground substance of the tissues the salts are reduced with the deposit of excessively fine granules of metallic silver. The process, in fact, is the same as that which occurs when we employ silver nitrate to make silver preparations of the tissues for histological purposes. The epithelial and glandular tissues are unaffected; the brain also remains free, but connective tissues are the seat of the deposits, notably the connective-tissue framework of the medulla of the kidney, the papillæ of the skin, the intima of the larger arteries, and serous membranes.

CHAPTER XXXII.

NECROSIS.

UNDER the term necrosis are included all those conditions of local death of cells, of tissues, and even of parts of the organism composed of many tissues, the organism as a whole continuing to live.

Gausation.—All those classes of noxe, mechanical, physical, chemical (including the bacterial), which may set up disease and cell degeneration, may induce necrosis when they act more intensely upon local cell areas. It will thus be recognized that there are all grades of cell disturbance, from the slighter degenerative conditions, through graver degenerations leading to eventual cell disintegration, to sudden death of the cells and tissues. It is usual to make a distinction between this intermediate form of gradual death, and necrosis in the narrower sense; it is spoken of as *necrobiosis*, and in the discussion of many of the degenerations, if sufficiently severe, induces necrobiotic changes; among these even the physiological atrophies, such as the constant wearing out and death of the outer layers of the epidermis, and the physiological accompany the formation of milk and sebum.

We have discussed in some detail the various mechanical, physical, and chemical causes of disease in the second part of this work, and there indicated how these may induce cell death. It is unnecessary now to do more than refer to what is there written. It is necessary, however, to refer in somewhat fuller detail to those conditions of necrosis set up by circulatory and nervous disturbances.

Girculatory Disturbances.—Two different orders of disturbance tend to produce cell and tissue death: (1) Arrest of blood supply; (2) deficient or perverted quality of the blood, with, as a result, deficient nutrition.

Many orders of local disturbance may cause the arrest of the blood supply to a part—ligation of the nutrient artery; ligation of the efferent veins; pressure upon the vessels by tumors, cysts, etc.; thrombosis, or coagulation of the blood within artery or vein; embolism, or obstruction of certain types of artery by foreign bodies, which, from their size, become blocked in the course of the vessel; the direct constricting and obliterating action of a poison, such as ergot; lowered action of the heart, so that the pressure within the vessels is unable to propel the blood onward; lastly, actual disease of the arterial wall, with proliferation of the intima, leading to occlusion.

Of these, widely different as are the effects upon the vessels of the part, occlusion of the afferent arteries and occlusion of the efferent 62

NECROSIS

vein both lead to the same death of the tissues of the part; the result is the same whether the blood be cut off from the region, or whether it can pass into the region but cannot leave it. In both cases there is developed a lack of oxidation of the tissues. The subjects of thrombosis and embolism will be found treated in detail in the second volume of this work. Here it is necessary to remind the reader that obstruction of an artery or of a vein only leads to necrosis in those cases in which there is an inadequate collateral circulation; provided that where an artery is blocked nutrition can be gained from blood provided by other arteries, and that where a vein is blocked the blood can drain from a region through collateral veins, a sufficient circulation may be maintained to preserve the vitality of the cells of a tissue or part. It is only where vessels are what is termed absolutely or relatively *terminal* that necrosis ensues; it is only under these conditions that we have developed the state of *infarct*, using this term in its broadest sense, and such infarct may be either anemic or hemorrhagic. It deserves emphasis that the infarctous state may be brought about either by arterial obstruction (the more common) or by venous. According as to whether there is sudden obliteration or gradual, so do we have either necrosis or necrobiosis.

Inadequate Nutrition.—Under this heading we include more particularly cases of general malnutrition and cachesia. Associated with these there is weakened heart action and inadequate blood supply. In all these cases the result is not so much a sudden necrosis as a progressive condition of necrobiosis; and in this the different orders of cells react differently; the more highly differentiated cells, such as those of glandular epithelium, are more easily influenced by nutritional disturbances than are the more lowly cells of connective-tissue type. In this way not all the cells of an affected area are necessarily involved.

Nervous Disturbances.-There has been, and there continues to be, much debate as to whether central stimuli proceeding from the higher nervous centres can in themselves induce necrosis, as also whether the removal of nervous influences is a direct factor. There are undoubted cases of impoverished nutrition and local anemia, more particularly of the extremities, which can only be referred to functional or hysterical conditions. In general, it is becoming more and more accepted that, while vasomotor influences, by constricting the vessels of a part, may induce necrosis, loss of nerve supply, while it may lead to cell inanition, does not of itself set up necrotic conditions. To produce these some other factor is regarded as necessary; thus, to cite a familiar example, it used to be held that section of the fifth nerve led to necrosis and ulceration of the cornea, and that there existed a definite condition of neurotrophic keratitis. It is now well established that, after such section of the nerve, provided that the surface of the eye be protected from light and dust and coarser injury no inflammation and no necrosis show themselves. The section of the nerves supplying a part affords an important predisposing but not a direct inciting cause of cell death. A like explanation is to be given for the so-called perforating ulcers of the sole

and other regions associated with Charcot's joint disease and advanced cases of locomotor ataxia.

Here may be recalled what has been stated (p. 869) regarding inanition atrophy and the gradual shrinkage and the gradual death of cells which have been cut off from receiving the normal nervous stimuli. The necrobiosis in these cases is so gradual as to be almost imperceptible. We would not, however, have it believed from the above paragraphs that loss of function does not eventually lead to the death of, more particularly, more highly differentiated cells.

Forms of Necrosis. --It is difficult to make a wholly rational and satisfactory distinction between the forms of necrosis and the resultant

changes that take place in a necrosed area. It is, however, possible in the first place to distinguish between (1) the necroses affecting individual cells; (2) those affecting small groups of cells—focal necroses; (3) those affecting circumscribed areas of one tissue —as result of vascular of struction infarcts; (4) necroses involving parts rather than tissues—mortification.

1. Necrosis of Individual Cells.— Apart from the conditions of local atrophy and fatty degeneration and necrobiosis which have already been referred to, certain rather characteristic necrotic changes in individual cells deserve mention. Of these, the aoost characteristic is that known as Zenker's degeneration, or waxy degeneration of muscle. In this condition individual muscle fibres are found which have lost all their striation and have become converted into masses of a waxy, almost glassy, appearance, lying within a still intact screolemma. The condition is found



Wax-like degeneration of muscle fibres (a, b) seventeen hours after temporary ligation of the same. In b there is already some accumulation of leukocytes. (Oberndörffer.)

most frequently in the muscles in typhoid fever, and more particularly, as first pointed out by Zenker, in the recti muscles of the abdomen. This, is however, by no means the only condition; it may be induced in individual muscle fibres by sharp blows, or by trauma; it has been noted in the skeletal muscle fibres in the neighborhood of tumors, as also in the heart muscle in cases of diphtheria (Ribbert). Opinion is divided as to the exact nature of the change; apparently it is of the nature of a coagulation of the muscle substance, a coagulation associated with the death of the same, for the waxy matter may undergo disintegration or absorption, and is not involved in the new regenerative process. Wells and Mathews suggest that the relatively abundant acid present in muscles leads to a

NECROSIS

swelling of the coagulated muscle substance similar to that of fibrin under the action of acids.

2. Focal Necroses. —More particularly in certain cases of severe infections there are encountered in different tissues minute areas of necrosis scarcely visible to the naked eye. Such, for example, are present in the lymph follicles in diphtheria and typhoid, as, again, after severe burns (Bardeen, J. McCrae). The most common example is, however, seen in the liver in typhoid fever, though similar conditions have been recorded in cases of sepsis, of scarlet fever (Pearce), and even in the more chronic states of tuberculosis and glanders; the most extreme are in the liver in the liver in cases of purperal eclampsia.



Multiple focal necroses in the liver of a rabbit subjected to experimental glanders. (Duval.)

Experimentally, in addition to injections of sundry bacteria and their toxins, focal necroses of many organs may be produced by abrin and ricin, by the toxic substance present in dog's blood serum,⁴ and by hemolytic agents in general (Pearce²).

Studying these cases, it is noted that small capillary areas are involved, and that here the cells in the first stage lose their nuclear stain undergoing karyorrhexis and chromatolysis. They thus, in stained specimens, contrast strongly with those of the surrounding tissue. McCree and Klotz¹ have made the interesting observation that while the surrounding liver tissue may show abundant fat, the recent focal necroses in

³ Journ. of Pathol., 12:1908:79.

³ Flexner, Johns Hopkins Hosp. Rep., 6: 1897 : 259.

³ Journ. of Med. Research, N. S., 7 ; 1904 ; 329.

FOCAL NECROSES

typhoid fever wholly fail to react with Sudan III. Later, there is attraction of leukocytes to the part, with disintegration and eventual absorption of the dead cells, the appearance suggesting that of early abscess formation. Nevertheless, careful staining for microörganisms shows that in the majority of cases these are characteristically absent. Only, to our knowledge, in the focal necroses of tuberculous marasmus has Le Count' detected the presence of bacilli, and attributed the condition to the local action of toxins diffused from these. In the allied necroses seen in chronic glanders Duval² has found no such relationship.

There has been much debate regarding the mode of causation of these focal necroses—nor can the matter be regarded as definitely settled. The probability is that there is more than one mode of formation. The following solutions have been advanced:

1. Local diffusion of toxins by bacteria *present in the tissues*. This as a possible cause is indicated by Le Count's observation, but is clearly the exception, and not the rule.

2. That soluble toxins *circulating* in the blood are directly responsible (Flexner and Opie). It is suggested that stasis of the blood in restricted capillary areas permits these toxins to affect the capillary endothelium, and, diffusing into the tissue cells of this area, produce upon them more severe and fatal effects.

3. That the causation is *embolic*. Schmorl more particularly called attention to the productions of capillary emboli by placental cells in cases of puerperal eclampsia, and Mallory has demonstrated experimentally that if the spleen of the guinea-pig be compressed so as to drive some of the splenic cells (endothelial) out of the sinuses into the splenic vein (or if active contraction of the spleen be induced by passing an electric current through the upper abdomen), within a few minutes capillary emboli of splenic corpuscles may be obtained in the liver.

4. Thrombotic causation. There is so extensive a collateral circulation in the hepatic capillaries that it is difficult to realize that capillary emboli alone are able to induce focal necroses. And Mallory³ has suggested, in connection with the typhoidal focal necroses, that the enlarged and proliferated endothelial cells seen in this disease, reaching the liver as emboli, there undergo degeneration and disintegration, lead to the local formation of thrombi extending along the capillaries, arresting the nutrition of the surrounding cells, and so leading to their necrosis.

It is possible that Flexner's and Mallory's theories may be harmonized by the determination that in one series of cases the toxins act directly on the capillary vascular endothelium, and, by destroying it, lead to the development of capillary thrombi; in the other, the thrombus is induced by the disintegration of cells within the capillary lumina.

In a careful examination of some 40 cases at the Royal Victoria Hospital, by J. McCrae and Klotz,⁴ they were unable to convince themselves that cell emboli played any active part in association with the

⁴ Journ. of Pathol., 12:1908:279.

¹ Jour, of Exp. Med., 2:1897:657.

¹ Trans. Assoc. Amer. Phys., 22; 1907; 398.

³ Jour, of Exp. Med., 3 ; 1898,

focal necroses; on the other hand, they found frequent evidences of hyaline thrombi. We were led to the same conclusion from a study of Dr. Duval's specimens from the liver in experimental glanders.

5. That the thrombi are due to a primary hemolysis. As an expla nation of these hyaline masses within the liver capillaries, the more recent observations of Pearce and his associates appear to afford valuable indications. These observations show that the toxins associated with the formation of these necroses exhibit, one and all, a very definite action upon the red corpuscles in vitro. Many of them are markedly agglutinative, but, what is more important, the more marked the production of focal necroses the greater the hemolytic activity of the toxin. Thus, Pearce suggests that hemolysis takes place throughout the system, and that small masses of broken-down erythrocytes become arrested in the liver capillaries, there giving rise to hvaline thrombi. As a confirmation of this view, Benno Schmidt, of Zurich, has recently shown that disintegrated masses of red blood cells can be recognized in the capillaries of the spleen and other organs, and this more particularly in the course of typhoid fever. We are thus inclined to hold that, whereas at times the endothelial and other cells may block the hepatic capillaries, more frequently the focal necroses are due to hemolytic action. It is quite possible that this hemolytic action may be (1) general, from the action of some agent in the circulating blood; or (2) local, the hemolysin being generated or discharged from disintegrating endothelial or other cells.

Fat Necrosis.—The condition of fat necrosis was first described by Balser, in 1882, but to Fitz, of Boston, we owe the first recognition of its intimate relationship to pancreatic disease or disorder, and other observers in the United States (Opie, Flexner, Williams, Wells) have been foremost in establishing our knowledge of the condition and its causes, although to Langerhans is due the credit of first establishing experimentally the relationship.


Fat necrosis reveals itself by the striking appearance of opaque whitish yellow areas or islands of small size —a few millimeters in diameter in the fatty tissues of the organism, standing out prominently against the surrounding more translucent yellow fat or zone of hemorrhagic tissue. Most frequently it is the fat upon and in the immediate neighborhood of the pancreas that is involved. In more advanced cases the omentum, mesenteric fat, appendices epiploice, and subperitoneal fat have these little areas scattered through them extensively. In the most advanced cases the mediastinal and pericardial fat are recorded as having been involved.

Microscopically, examining sections which have been frozen and not treated with alcohol or clearing fluid, collections of fat cells are seen having a greatly altered appearance. Their outline may still be determined, but instead of clear contents, they present clusters of "margarin crystals" (a combination of palmitic and stearic acids), while others are filled with a granular debris which, according to Langerhans, is composed largely of calcium soaps detectable by microchemical means.

FAT NECROSIS

It is noticeable that frequently the affected fat cells take on little or no stain with osmic acid, although the surrounding normal fat cells assume the usual dense black. As osmic acid stains only oleic acid compounds, this would appear to indicate that the olein, in dissociation, passes rapidly into a diffusible modification, leaving the more insoluble palmitic and stearic acid compounds behind.

If careful examination be made of the peripancreatic fat at autopsies, it is not unusual to distinguish an isolated area or two of fat necrosis and this in cases affording no history indicative of pancreatic disturbance and showing no obvious lesion of the organ. To these slight cases reference will be made later. But any extensive manifestation of the condition confirms Fitz's observation that there is associated pancreatic lesion. In the majority of cases hemorrhagic pancreatiis is present; not infrequently there is actual gangrene of the organ, which may lie



Fat crystals (margarin). × 250. (Perls.)

almost dissected from the surrounding tissues. It is deserving of note that suppurative disorders of the organ are rarely accompanied by fat necrosis.

The numerous experiments that have been made upon the subject, from Langerbans onward, demonstrate that the condition is due to the escape into the tissues of the fat-splitting ferment, normally present in the pancreatic juice. Thus, fat necroses have been observed to follow:

1. The injection into the fat tissue of rabbits and dogs of an aseptic infusion of rabbit's pancreas (Langerhans).

 The introduction into the peritoneal cavity of one animal of pieces of fresh pancreas taken from another (Jung).

 The ligature of the tail of the pancreas with ligation of its veins (Hildebrand, Flexner, Williams), or by multiple ligatures (Katz and Winkler).

NECROSIS

4. The temporary obstruction (20 min.) of the circulation of part of the organ (Blume).

5. Ligation of the pancreatic ducts (Opie).

6. The escape of the pancreatic juice from the divided duct into the surrounding, or into the subcutaneous, fat (Milisch, Opie).

 Severe injury to the pancreatic tissue, as by injecting into the duct, or the tissue direct, turpentine, artificial gastric juice, etc. (Hlava, Körte, Oser, Flexner).

8. Injection of steapsin (the fat-splitting ferment) into fatty tissue (Flexner).

9. Injection of steapsin plus trypsin (Wells).

The evidence is thus abundant that the pancreatic juice and its constituent steaps induce fat necrosis. Nevertheless, it has been objected (1) that a remarkably large proportion of experiments fail to produce the disturbance, and (2) that the experimental necroses are not nearly so extensive as those occasionally met with in man. The explanation, according to Opie, is that (*a*) suppurative complications prevent the development, and (*b*) that time is not usually afforded for extensive diffusion of the pancreatic juice. In some of Opie's experiments, in which the animals died or were killed at the end of two or three weeks, necroses as extensive as those in man were obtained. Wells, in addition, doubts whether, under ordinary conditions of experiment, steaps alone will induce necrosis; there must be coincident action of tryps in.


Chiari has afforded what appears to be an adequate explanation for the cases of slight fat necrosis unaccompanied by obvious pancreatic disease. Such constantly exhibit postmortem self-digestion of the organ, and must be due to diffusion outward of the juice from the cells undergoing this form of autolysis into the surrounding fat. He is inclined to the opinion that such self-digestion may at times be agonal. The observations of Opie, Flexner, and Halsted suggest the mechanism whereby hemorrhagic pancreatitis would seem most often to be brought about, namely, by injury to the walls of the duct, or pressure within the duct sufficient to permit a transfusion of the excreted juice into the surrounding tissues, with digestion of the same. Opic more particularly has pointed out that obstruction of the ampulla of Vater by a gallstone, or of the termination of the duct by cancer of the head of the pancreas, reproduces the conditions developed in his experiment. Similar obstruction of larger and smaller ducts may be induced by chronic interstitial pancreatitis.1

Infarcts and Coagulation Necrosis.—The mode of formation of infarcts, their relationship to embolism and thrombosis, and the varieties, will be discussed in the second volume. Here only must be taken into account a characteristic form of necrosis which accompanies some, but not all, cases of infarct formation, a form to which Weigert gave the

⁴ For a fuller discussion of this subject with literature, see Opie's Diseases of the Panceas; Wells' Chemical Pathology, Lippincott, 1907, gives the more recent literature.

COAGULATION NECROSIS

name of *coagulation necrosis*. The condition is best seen in anemic infarcts of the kidney and spleen; the affected areas become firm, pale, and relatively swollen, in the early state projecting distinctly above the surrounding surface. Under the microscope the nuclei have wholly lost their chromatin, and the cells have a hyaline appearance, with loss of sharp outline, the whole area appearing to be converted into a solid, somewhat homogeneous mass, as though coagulated uniformly. In some cases, by the use of Weigert's fibrin stain, actual fibrin can be recognized, laid down here and there in filamentous form. The comparison may be instituted between this condition and thrombus formation, in which the



Anemic infarct of cortex of kidney to show congulation necrosis, with surrounding zone of congestion: a, artery. (Orth.)

coagulation of the blood, while often leading to the production of a fibrinous network, may also be of the hyaline type, with coagulation *en masse*. Weigert regarded the process as essentially of the nature of a fibrinous coagulation, the dying cells liberating a fibrin ferment, the lymph and fluids of the area, together with the cell bodies, affording substances of the nature of fibrinogen. That the whole mass of cells forming the infarct becomes a mass of fibrin may well be doubted. Rather we must admit that blood fibrin is but one of a group of coagulated proteins, and compare the process with what obtains in muscle in Zenker's degeneration, where the myosinogen becomes converted into coagulated

myosin (p. 979). The process, it may be noted, does not attack all tissues, as might reasonably be expected were it the result of reaction between the protein-containing fluids diffusing into those tissues and enzymes liberated in the death of the cells. Anemic infarcts of the brain do not exhibit it, but instead, a form of colliquative necrosis.

Colliquative Necrosis.-Two distinct processes are often confused under this term, namely, colliquative necrosis proper, due to the liquefaction of the dead tissue as a process of self-digestion unassociated with any bacterial decomposition, and *putrefactive necrosis*, due to a not dissimilar liquefaction brought about by the proteolytic activities of bacteria. The latter we shall refer to later. Colliquative necrosis proper can only be regarded as autolytic in nature; the dead area softens, the cells undergoing a granular disintegration, with production of myelin, fat, cholesterin, etc. It is well seen, as above noted, in connection with infarcts (both anemic and hemorrhagic) of the brain. A fluid granular debris is the result, containing a certain number of migrated leukocytes (or, according to Fritz Marchand, of glial cells), which become engorged with fat in the form of small fatty globules, forming large characteristic granule cells, or Gluge's corpuscles. Some of these migrate into the tissue immediately surrounding. According as there is more or less blood present and involved in the necrotic area, so do the fluid contents assume different colors (see p. 960).

Similar colliquative necrosis occurs in other tissues and conditions apart from obvious circulatory disturbances, though in all cases, strictly speaking, we deal with a primary cutting off of the blood supply to the affected areas. The atheromatous softening of the deeper layers of the intima in one stage of the arteriosclerotic process is of colliquative type. A similar colliquative necrosis may involve the central parts of tumors and lead to eventual falling in and umbilication, or to cyst formation (see p. 864). So, also, old standing thrombi, more particularly the sessile thrombi of the heart cavities, undergo a central colliquation, and may be represented by a shell of fibrin enclosing a turbid fluid.

Gaseation.—In this connection another type of necrosis may be noted, in which the necrosed area exhibits neither congulation nor colliquative change, but, undergoing a slower necrobiotic change, the cells exhibit a change akin to fatty degeneration, become granular, and break up into fatty granular debris, in which no sign of the earlier cellular structure can be recognized. The area thus becomes converted into a mass of the appearance and consistence of a rather dry cream cheese; hence the term caseation. The change characteristically occurs in connection with tuberculous new-growths. Tubercles are, from their mode of development, extravascular; but the cutting off of the blood supply, while tending to produce necrobiosis and necrosis, would not produce this special type of change. That must be attributed to the action of the tuberculous toxins and their effect upon the cells. Through inspissation of pus, produced by other bacterial agents, we occasionally encounter similar caseous accumulations.

Gummata.-That toxins are factors is indicated by the fact that gum-

MORTIFICATION AND GANGRENE

mata, or syphilitic tubercles, which, histologically, are of closely allied formation, although due to the presence of microbes of very different type, do not exhibit caseation proper. Their necrotic centres, while showing, similarly, no trace of cell structure, are "gummy" rather than caseous. There is not the same abundant fat present. Beyond this little is known of the exact constitution of the gummatous necrotic matter.

Mortification and Gangrene.—The death of large areas of tissue and of parts composed of many tissues may be brought about by very many causes: by vascular obstruction and arrest of the blood supply to a part, or of the outflow from a part; by enfeebled circulation; temporary stoppage of the circulation of a part or organ, as in Litten's experiment on the kidney (p. 929); acute infection (as, for example, phlegmonous cellulitis, hospital gangrene, and emphysematous gangrene, due to the growth of B. Welchii); by burns, as an after result of intense cold (frostbite), action of chemical agents (caustics and acids), and of physical (electricity, z-rays, radium).

The Results of Necrosis.—This description of the different forms of necrosis prepares us to find that the results vary widely. (1) Where the necrosed area is small and there is no infection, *absorption* occurs. By autolysis the cells undergo disintegration and more or less solution. Leukocytes attracted to the area, by their phagocytic activity aid the process, and not only may there be absorption, but this may be followed by regeneration. Even where the necrotic process has been very extensive such regeneration may tend to show itself, as, for example, in those cases of very extensive necrosis of the liver cells which are met with in acute yellow atrophy and chloroform poisoning, in which the specific cells of the liver are in the main involved. In such cases where death as not been brought about within a few days, proliferating liver cells can be recognized advancing into the spaces left by the autolysis and dissolution of the preexisting liver cells.


(2) More often, where the area of dead tissue is a fair size, and where, again, there is no infection, we encounter a *cicatrization* as the result of organization. This is especially well seen in the latter stages of the non-infected infarct. The death of the cells and diffusion outward of their products of disintegration lead to the production of a surrounding zone of inflammation, with well-marked congestion of capillaries and migration of leukocytes into the dead area. As in the previous case, these leukocytes reinforce the autolytic processes and aid in the removal of the dead matter, but with this the surrounding capillaries send new vascular loops into the region, and what is truly a granulation tissue is developed, which gives place eventually to well-contracted cicatricial fibrous tissue. A similar process of organization is the end result in many cases of intravascular congulation of the blood and thrombosis.

(3) Where, as in the brain, the tendency is toward colliquative necrosis, there the end result of the autolytic process is *eyst formation*, rather than organization and cicatrization, though small necrotic foci in the cerebellum may give place to a complete organization.

NECROSIS

(4) Where, as in bone, the tissue is so dense that disintegration of the dead matter is a long-drawn-out process, there, through leukocytic action more particularly, the surface portions of the dead area may be disintegrated and loosened from the surrounding more healthy tissue, and in such cases the still unabsorbed necrosed mass remains as a *sequestrum*, lying in a more or less well-defined cavity or tract, and bathed in a fluid of purulent nature.

(5) Another sequel to colliquative change is *inspissation*, the fluid portion of the dead material draining away, leaving thus a more or less cheesy accumulation. Indeed, in caseation proper, as seen in tuberculosis, there is a certain grade of inspissation present. Such cheesy, inspissated matter is especially prone to become the site of calcification (p. 928).



Senile gangrene of the great toe, from a case of arterial thrombosis. The toe is shrunken and its epidemis as being exfoliated. At the line of demarcation the skin has retracted (a) and the deeper parts are separating (b).

Gangrene.—In the condition of gangrene, according to the extent of blood entering the dead part from the vessels, and the rate of evaporation of fluid from the surface, so do we have developed either the condition of (6) moist gangrene or sphacelus, or of (7) mummification or dry gangrene. It is in the extremities and the ears that the latter condition alone can show itself. The necrotic portion becomes shrunken, wrinkled, and assumes the dark brownish-black color, the appearance, in short, of mummy flesh. As in an infarct formation, so here, at the zone of junction of the dead and living tissue, there develops an infarct formation, leukocytes passing beyond this aid in the solution of the dead matter, whereby the mummified and living matter become separated and the former becomes eventually detached.

Where, as in the lung and the intestine, evaporation cannot take place, and in the extremities of those cases in which blood is still able to enter the

GANGRENE

dead area, there, on the contrary, the dead matter becomes waterlogged, and inevitably putrefaction sets in, through the entrance from the surface of various microbes. The appearance in these cases is striking; the affected tissue becomes greatly swollen and livid; on the skin large blebs may form, filled with ordematous fluid; the discharge from such blebs becomes foul and stinking through bacterial growth; through the same putrefactive agents the blood corpuseles become broken down, and their pigment becomes diffused through the tissues. So, also, the various soft tissues become decomposed and deliquesce, a foul fluid resulting, filled with fatty globules, pigment, and various products of proteolysis.

Both in dry and in moist gangrene any bony portions involved are the last to undergo decomposition.

CHAPTER XXXIII.

DEATH.

In this life of ours, with all its uncertainties, there stands out one certainty—that we shall die. Sooner or later death comes to all men. It is as inevitable as that on this rotating world of ours night follows after day; nor can any care on our part, any precise regulation of the course of our days—can science or prescience—ward it off for more than a few years. There is no *elixir vita*, no philosopher's stone.

This supreme fact has profoundly affected all human thought, and has been the pivotal point of all philosophies; nay, more, the various religions of the world may be regarded as the evidence of man's determination to rise superior to the dissolution of his body. Mere philosophy, however, cannot tell us why death is inevitable; we can only find an explanation by the study of living matter and its attributes. Making such a study, it is seen that death is not inherent in living matter as such; that it is the price paid for advance and increased power over Nature. For death is not inherent in the constitution of the simplest unicellular organisms. With such relative constancy of environment as Nature provides-within the natural limits of heat and cold, of dryness and moisture-the schizomycete microbe assimilates and grows and divides, and if, over long periods of time, the environment undergoes slow change, the property of adaptation, to which we referred at length in our opening chapters, permits the organism to adjust itself surely to the new conditions. At each division each half carries on the flame of life. There is in such a process no inherent death; at most, there may be accidental death by temporary lack of nutrition, by desiccation, by physical and chemical bactericidal agents in general. In like manner, with unicellular organisms higher in the scale, death is only apparent; when, for example, the nuclear matter of the hematozoon of malaria undergoes division and becomes distributed into numerous spores, and those spores break away, leaving behind a collection of fine pigment granules and debris to represent what had been the previous single individual, there is to the eve, it is true, a cessation of individual existence. but in reality the living matter, far from being dead, is increased in amount, and is given the occasion to increase itself still farther. There has been no destruction of the nucleoplasm, the essential living matter, but a multiplication of the same; each spore carries on the life.

It is with the appearance of the metazoon, of the multicellular organisms, that natural death enters into the world. Multicellularity connotes division of labor among the component cells, specialization of function and increased capacity of the individual as a whole. Certain cells, the germ cells, become set apart to carry on the living matter which shall develop into new individuals. In these germ cells, then, death is not inherent. It becomes inherent in the somatic cells, and this through the functional differentiation that they have undergone. An ideal cell republic might be imagined, in which the division of labor and function among the constituent cells was so allotted that each nourished and contributed to the exact needs of the other, all developing their powers simultaneously. In such a case there would be no need for somatic death. As a matter of fact, this ideal state has not been attained, nor, under the conditions of development, is it even possible. The individual, that is, undergoes growth and development, and this process of growth demands that different orders of cells are required to be mature and active at different life periods. Herein, it seems to us, is the essential explanation of somatic death.

Take, for example, the case of man himself: Even in the embryo, organs are developed-such as the yolk sac-which are of merely temporary use; they perform temporary service until other parts are developed which are of greater value; and when they become useless, their cells tend to atrophy and disappear; a new equilibrium has to be established. And so, throughout foetal life there is constant change in the relationship and interaction of parts. Of all embryonic and fœtal organs of active function which have this temporary character, the placenta stands out preëminent. Postnatal existence affords abundant examples of the same order. The individual tissues have periods of development to full activity and maturity which are not synchronous. Some, like the heart and kidneys, are fully functional before birth; subsequent increase in size and activity are of the nature of adaptations to the increased work thrown upon them by the growth and increase of the body in general. Others, like the thymus and the lymphoid tissues of the organism, are at their maximum in the early years of life, and already show diminution and atrophy before the adult state of the organism as a whole has been attained. Others, like the brain, attain their full anatomical development in childhood. Yet other organs and tissues lie latent, showing little signs of development for years. Such are the genitalia and accessory organs of generation, the mammary glands, etc., and these, again, like the ovaries, may, from purely physiological causes, have a period of active life shorter than that of the organism as a whole.

If these various organs in the performance of special function not only extract from the blood the materials necessary for their growth and nutrition, but afford internal secretions to the same which are of definite service to other tissues and to the organism as a whole, it will be seen that the atrophy and disappearance of the same induces not only loss of special function, but leaves the blood and remaining active tissues impoverished in one or other direction. Up to a certain point there may be an internal adaptation; it would seem there is always a tendency thereto, other tissues taking on certain of the functions of those that have activity throws additional strain upon them and brings the still active cells nearer to the margin of their reserve force, nearer to the point at which these in turn become exhausted and undergo atrophy.

By such processes of continually modified equilibrium and of increased strain thrown upon the surviving cells and tissues the period is reached at which disintegrative changes in the organism exceed the assimilative, at which, also, the reserve force of the surviving cells becomes diminished. and in this way inevitably the time is reached when sundry cells, unable through loss of this reserve force to respond to stimuli, bring about the condition of somatic death. As Bichat long ago pointed out, most of the tissues and even parts of the organism may be destroyed and yet life still persist. There is, however, a triumvirate of organs-the circulatory, the respiratory, and the nervous-each one of which is indispensable, and any one of which, if injured in particular regions, alone may bring about the state of somatic death. The reseaches of the last half century, and more especially of the last few years, have impressed upon us that other organs play an almost equal part. While those which for the race are of the foremost importance-the ovaries and testes-may be removed without in any way influencing the span of life of the individual whose "by-play," to quote Sir Michæl Foster, may still continue, certain small and hitherto little regarded organs cannot be removed without death being the inevitable result. Such are the adrenals, the minute parathyroids, and, as Paulesco, Reford and Harvey Cushing, and others have shown, the yet smaller pituitary body. Insignificant in size, as it is, remove this last and death (at least in the laboratory animals so far tested) supervenes within forty-eight hours.¹ The results are not so immediate, it is true; but in physiological death, such as that here suggested, it must be that certain cells in one or other of these fail to react, and so bring about arrest of function and cessation of life.

Rarely do we encounter this natural process—the passing of the quiet sleep of exhausted old age imperceptibly into death. Usually in those who have attained great age what happens is that the resisting and protective powers of the organism, more particularly, shows signs of exhaustion, with the result that sundry pathogenic organisms normally present upon the surface of the body, but normally prevented from gaining entrance into the tissues through the agency of the protective cells, eventually, despite slight virulence, gain entrance into the tissues, multiply, and set up a *terminal infection*. Such terminal infections are the immediate cause of death not merely in old age, but in all the states of progressive disease with progressive sapping of the reserve force of the individual. It is the prevalence of the terminal infections that gives force to Osler's dictum, that the individual rarely dies of the disease from which he suffers, save, it may be added, when that disease is in itself of the nature of an acute infection.

¹ As Schäfer and Herring point out, the active portion of the pituitary is not the posterior neuroglial lobe, but the anterior with its contained glandular vesicles, and it may be also the glandular *pars intermedia*. (Herter Lectures, Baltimore, April, 1908.)

THE SIGNS OF DEATH

We have thus to recegnize two orders of death—the *physiological* and the *pathological*; of the latter the terminal infection is the most common cause, though there are abundant others that will lead to the same end. Just as it has been pointed out that all the causes of disease may lead to local cell death when acting with a certain intensity, so, without exception, acting with greater intensity or acting directly upon one of the three vital organs, mechanical, physical, chemical, and bacterial agents may induce somatic death.

This somatic death may briefly be described as the cessation of function of the three vital organs, followed by the signs of disintegration and decomposition of the tissues in general. It is difficult to describe the state otherwise, inasmuch as it is not necessarily accompanied by the immediate death of all the component cells of the body.

Examples confirmatory of this statement are abundant and familiar. The head may be cut off a snake and the body for long continue to wriggle actively. We have ourselves seen the heart of the tortoise removed and a strip of the cardiac muscle still exhibiting spontaneous contractions eight days after such removal.¹

The so-called "überlebendes," cat's heart, or mammalian kidney wholly removed from connection with the body, will similarly, under favorable conditions, continue to function for hours. I have already afforded several examples (p. 638, etc.) of the persistence of life in isolated tissues, notably the continued existence and proliferative capacity of epidermal cells, for several weeks, and of tumor cells from the lower animals for several months, if kept under appropriate conditions.

The Signs of Death.—These are discussed fully in text-books of medical jurisprudence; here it is but necessary to refer to them briefly. They are many in number, and from a medicolegal point of view are of different value as affording indication of the period that has elapsed since death occurred. Among the more important may be mentioned;

1. **Cessation of respirations**, so that a cold mirror held in front of the nostrils does not become moistened and dulled.

2. **Stoppage of Heart Beat.**—Neither this nor the preceding are absolute signs, as it has been shown experimentally that after poisoning a dog with chloroform until both heart and respiration have stopped, transfusion of saline fluid or defibrinated blood under pressure may be followed by resumption of the heart beat and gradual recovery. They are, however, the first signs for which one looks. They are corroborated by—

3. Loss of Transparency of the Cornea.—The pupils usually dilate at the moment of death; the eyes stare directly forward; the cornea in a short time becomes cloudy.

4. The development of **rigor mortis**. Of those parts of the body which can be observed and studied without section, the eyelids are the first to pass into a state of rigidity with contracture, leaving the eyes to remain open. According to Fuchs, the heart ventricles are the first muscles to become rigid, and certainly in those cases in which a post-

¹ This in Dr. Gaskell's laboratory at Cambridge, in 1884.

DEATH

mortem is performed within an hour or two after death, the ventricles are constantly found firmly contracted, so as to suggest to those ignorant of this rapid rigidity the presence of the so-called "concentric" hypertrophy. There is great variation, however, in the period of onset of this change in the muscles of the body. In those engaged in active and violent exercise, it may be practically coincident with death. Strychnine poisoning and tetanus also exhibit rapid rigidity. On the other hand, prolonged wasting conditions, with muscular atrophy, may exhibit a rigidity only showing itself after many hours; the same is true in cases of death from asphysia and hemorrhage. The duration of the rigidity also varies very greatly.

The nature of the rigidity would appear to be that of a coagulation of the *myosinogen* of the normal muscle, *myosin* being the term given to the cocgulated product. This coagulation would appear to be brought about chiefly by the lactic acid of the muscle. The passing off of the rigor mortis would appear to be due to development of autolytic changes, favored, we may recall, by the acid state of the myosin.

5. **Cadaveric** Lividity.—Through gravitation of the blood to the dependent capillaries, the under or lower parts of the dead body show within a few hours a livid reddening, or, where the blood is more venous, a bluish-purple color. Where there has been cyanosis with great distension of the superficial capillaries before death, as not infrequently happens in the vessels of the face and neck where death has occurred from asphyxial disturbances, a similar, and even more intense, lividity may be present over surfaces that are not dependent. Where there is pressure, as upon the nates and over the shoulder-blades, the mere weight of the body prevents the filling of the capillaries, and such regions of pressure remain pale.

6. Decomposition and Putrefaction.-Decomposition of the organs shows itself most frequently first over the abdomen, as a greenish discoloration. The onset is very variable, it being delayed or completely arrested by great cold and materially hastened by warmth. Those organs and parts which normally are moist and contain abundant bacteria exhibit the putrefactive changes earliest; thus it is that the intestinal canal is most markedly affected. There are, however, other factors: Cases of acute infection and of bacteremia are especially apt to early decomposition, not merely, it would seem, through the action of the specific pathogenic organisms, but because in the course of the infection the protective substances of the organism have been exhausted, and there is no inhibition to the growth of putrefactive bacteria. A similar rapid decomposition has been noted after snake poisoning, in which also, there is a rapid destruction of antibodies. Arsenical and certain other intoxications may very materially delay the onset.

Other slighter signs of death are:

7. Relaxation of sphincters; and

 Loss of tissue elasticity.—The latter would seem to be largely due to solidification of the subcutaneous fat, so that the position assumed by the surface tissues at the moment of solidification tends to become fixed.

APPENDIX.

THE ULTRAMICROSCOPIC MICROBES IN RELATIONSHIP TO DISEASE.

THE existence of pathogenic microbes so minute as to be undistinguishable under the highest powers of the ordinary microscope save as fine points too small to be resolved, was first established by Roux and Nocard¹ in 1898. These observers showed that the clear exudate from the lungs in cases of contagious pleuropneumonia in cattle, while presenting no microorganisms visible under the microscope, nevertheless conveys infection when inoculated into other cattle; that it gained increased powers when placed in celloidin capsules within the tissues, and that when a drop of the exudate was placed in an appropriate serum broth and incubated at body temperature, there gradually developed in the upper part of the tube a cloudy or milky layer. Examination of this laver, as of the contents of the celloidin capsules, demonstrated abundant minute points with Brownian movements. Growths could be made from tube to tube, retaining their virulence for long periods. What is more, the virus is so minute that it is filterable through a Berkefeld or Chamberland (F) filter (Dujardin-Beaumetz).


With this definitely determined, it is natural that the existence of similar ultramicroscopic organisms has been predicated in connection with all those infectious diseases in which, despite all the resources of modern technique, neither bacteria nor animal microparasites have thus far been detected. And inevitably, as every other form of microorganism has been tried and found wanting, there are those who ascribe malignant growths to these ultramicroscopic organisms. The long-delayed discovery of the spirochete in syphilis indicates that such conclusions are not wholly secure. Nevertheless, with regard to not a few diseases, there are indications favoring this view-where, for example, as in cases of vellow fever, as shown by Thomas, and of measles, as shown by Hektoen, the blood of a patient conveys infection although it affords no cultures and microscopically is devoid of any signs of microorganisms; where, again, the virus is filterable, as in yellow fever, as shown by the American Yellow Fever Commission in foot-and-mouth disease (Löffler and Frösch), South African horse sickness (MacFadyean), dengue (Ashburn and Craig), avian pest or cyanophilia; in smallpox and vaccinia,

⁴ Annales de l'Inst. Pasteur, 12; 1898; see also Dujardin-Beaumetz, Le Microbe de la péripneumonie, Paris, Doin, 1900.

MICROBES IN RELATIONSHIP TO DISEASE

as noted by Casagrandi,¹ Licheri,² and several observers, and as in acute epidemic poliomyelitis (Flexner and Lewis), the presumption is distinctly in favor of ultramicroscopic microbes as the causative agents. So also both Ballah,³ in our laboratory in connection with vaccinia, and Flexner and Lewis,⁴ with epidemic poliomyelitis, have obtained clouding of the upper layers of fluid media with appearance of minute granules. It would seem not unlikely, therefore, that yellow fever, acute epidemic poliomyelitis, smallpox and vaccinia, and measles are due to microbes of the nature of those found in contagious pleuropneumonia.

What are these microbes? Are they more nearly related to the bacteria or the animal microparasites? By analogy the high grade of immunity developed after an attack of any of the above-mentioned diseases indicates that they are of bacterial nature; immunity of this order is not



From a culture of the microbe of contagious pleuropneumonia of cattle, For description see text, (After Dujardin-Beaumetz.)

characteristic of diseases due to animal microparasites. And while this volume has been passing through the press there have appeared two articles, by Bordet,⁵ and Dujardin-Beaumetz⁶ respectively, bearing out this contention. The former points out that grown upon media containing a large proportion of rabbit's blood, in three days or so cultures of

¹ Boll, d. Soc, tra i colt, d. Scienz, in Cagliari, 1908; Boll, 1-4; 11 and Boll, 5; 32.

² Ann. d'Ig. sperim., 19: 1909: 291.

³ Brit. Med Journ., 1906; ii: 1779.


4 Journ. Amer. Med. Assoc., 54: 1910: No. 1.

⁵ Bulletin de la Soc. roy. des Sci. Méd. et Nat., Brussels, November, 1909, and Annales de l'Inst. Pasteur, 24:1910:161.

⁶ Ibid., 24:1910:168,

APPENDIX

the organism of boyine pleuropneumonia exhibit irregular curved and twisted filaments resembling, he holds, spirilla rather than spirochetes. Dujardin-Beaumetz and his collaborators, making a fuller study and employing the ultramicroscope and the highest powers of the ordinary microscope, with magnification of 5000 diameters, describe a very remarkable series of forms according to the age of the cultures, Bordet's spirals being only one stage. At first coccoid and diplococcoid forms are to be made out, followed by the appearance of chains and ring-like collections of coccoid bodies; later irregular filaments of Bordet's type, with remarkable star-like bodies (Fig. 329). By overstaining they find evidence that these are enclosed in a mucoid envelope. The organism is extraordinarily polymorphic. Dujardin-Beaumetz suggests the name of *Asterococcus mycoides* and points out that in not a few features these organisms recall the curious bacterial organisms of the root nodules of leguminose.



INDEX.

A

ABDERHALDEN'S experiments, 500 Abnormalities, 226 inheritance of, 209 from imperfection of germ cells, 208 from infectious diseases, 219 a hemolytic, 306 immunization against, 505 in descendants, 194 Römer's experiment, 516 Abscess, embolic, 434 formation, 431, 432 resolution of, 432 stitch, 324 Acardiac monsters, 231 Accessory chromosome, 152, 153 digits, inheritance of, 163 mammæ, 181 organs, 258 Acetonuria, 372, 380 Acetylene, action of, 306 Achondroplasia, 224 Acid intoxication and fatigue, 401 Acidosis, 372, 381 Acquired characters, inheritance of, 192 disease, antenatal, from infection, of intra-uterine acquirement, 215 parturient, 225 Acrania, 263 Acromegaly, 228, 357 hypertrophy in, 598 Actinomycosis granuloma, 442 melanin granules in, 48 Acute infection, 260 yellow atrophy, autolysis in, 372 Adaptation, 116 bisphores and, 157 immunity and, 565 inflammation and, 415, 447 to parasitic life, 344

Adaptation to physical alterations in physical basis of, 122 in protozoa, 416 Addiment, 566 Addison's disease, 358 in autolysis, 371 Adenin, 64, 376 Adenocarcinoma, 802 Adenoma, 783, 789 malignant, 784 of prostate, 789 of thyroid, 789 Adenomyoma, 748 Adipocere, 913 Adiposis dolorosa, 723 Adrenal, accessory, 359, 640 cortex of, enzymes in, 360 tumors of, 677, 836 mode of origin of, 693 glycosuria, 368 internal secretion, 358 neuroma of, 809 tumors of, 807, 809 metastases of, 682 Adrenalin, 306 action of, 308 Adrenin, action of sympathetic nerves aromatic protein derivatives and, source of, 358, 360 enzyme action and, 79 Ægagropile, 954 Affections, diseases and, 21 Age, a factor in hypertrophy, 594, 516 Agenesia, 210 Agglutination, nature of process of, 531 by ricin, 506 Agglutinins, 527 group reactions of, 530 properties of, 528 relationship, 531 Aggressins, 557 Agnathia, 272 Alanin, 59 Albinism, 163 a recessive condition, 174 due to mutation, 182

1000

Albuminoids, 56 Albumins, 56 Albuminuria, cyclical, 958 Albumosuria, 373 Alcohol, action of, on heart, 307 effects of, on nervous system, 304 immunity, 504 Alcoholism, effects of, upon offspring, 196, 220 Aleppo button, 334 Alexines, 497, 543, 566 Alkaptonuria, 379 Allemorphs, 190 Allergia. See Anaphylaxis. Allomorphism, 641 Alloxuric bodies, 376 Aloes, action of, 309 Altmann's elementary organisms, 331 granules, cloudy swelling and, 883 Aluminosis, 974 Amanita, 306 phalloides, 504 Ambergris, 955 Amboceptoids, 537 Amboceptor, 536 group reactions of, 544 multiplicity of, 536 synonyms of, 566 (note) Amino-acids, 57, 59 Amino-dioxypurin, deposits of, 954 Amitosis, 113 Ammonium, carbamate, 389 Amniotic adhesions due to pressure, 217 Amaba, digestion in, 416 of dysentery, 333 excretion of slime in, 47 Amphiaster, 115 Amphimixis, adaptation and, 118 as a factor in variation, 188 its influence, 159 Amphioxus, experiments on eggs of, 138 Amputation, intra-uterine, 217 Amygdalin, 82, 575 Amyl nitrite, action of, 308 Amyloid, 40, 57 infiltration, 891 Amyloidosis, 891 Amylopsin, 91 Anadidymus, 240 (note) Anaphase, mitotic, 115 Anaphylaxis, 559 idiosyncrasy and, 410 Anaplasia, 641, 709, 765, **841** Andalusian fowl, hybrids of, 171 (note) Anemia, due to autolysis, 373

pernicious. See Pernicious anemia.

Anemia, splenic, 743 Anencephaly, 263 from pressure in utero, 217 Aneurysm, 875 atrophy and, 872 dissecting, development of, 615 sacculated, 850 Angioma, 814, 817 Aniline dyes, intravital staining of tissues by, 975 Animal venoms. See Venoms Ankylostoma duodenale, 346, 348, 350, Ankylostomiasis, 350 Anopheles, 337 Anorexia, 476 Antedon, 139 bifida, nucleoli in, 31 ova, chromidia and yolk granules of, 49 Antenatal acquirement of disease, 215 disease, acquired, from infection, 221 from malnutrition, 218 Anthracosis, 973 Anthrax, Pasteur's observations on, 493 Antiamboceptors, 537, 545 Antiarin, action of, 307 Antibodies, in blood of normal animals, of tapeworms, 348 Anti-enzyme, distinct from fermentescible negative catalysts and, 78 (note) Antimony, salts of, action of, 307 tartrate, mode of action of, 309 Antipepsin, 508 Antiphrynolysin, 550 Antitoxin, action of, on toxin within organism, 523 diphtheria, amount of serum produced, 125 production as compared with of horses, amyloid infiltration and, 896 Antityrosinase, 508 Antiurease, 508 Antivenin, 550 Antler's, inheritance of, 180 Anus, atresia of, 275 Aorta, atheroma of, soaps in, 94 Apicopolar fusion, 248

INDEX

Autorennin, 508

Azoturia, 959

Azoospermia, 294

Apomorphine, 304 mode of action of, 308, 309 Appendicitis, 433 Aprosopia, 271 Arachnida, 343 Arginin, 58 Argyria, 976 Arrhincephalus, 260 Arsenic as cause of overgrowth, 597 immunity toward, 501 salts of, action of, 307 treatment of malignancy with, 692 Arsenious acid, elimination of, 310 Artemia, 148 Arteries, media of, hypertrophy of, 594 uterine, involution of, 896 Arteriosclerosis, senile, 875 Arthrospores, 314 Arthus' phenomenon, 559 Articular laxity, 182 Ascariasis, 350 Ascaris, chromosomes in, 148 germ cells of, 145 lumbricoides, 347, 348, 350 megalocephala, 348 bivalens, spermatogenesis and oögenesis in, 149 Aseptic supportation, 431 Asparagin, 59 Aspartic acid, 59 acidosis and, 382 Asterococcus mycoides, 997 Asthma, anaphylactic nature of, 561 Atavism, 175 definition of, 164 Atheroma, 986 soaps in, 94 Athyrea, 355 Atreptic immunity, 848 Atresia ani, 275 Atrophy, brown, 874 disuse, 402, 869 non-inheritance of, 196 from excessive work, 111 from overnutrition, 872 from overwork, 871 from subnormal cell activity, 109 reversionary, 641 senile, 873 serous, 873, 904 Atropine, 305, 306 action of, on skin, 312 on vessels, 308 Autointoxication, 500 Autolysis, 370 myelin bodies in, 97

B

ACILLUS ACIDI LATICI, tolerance
of, to acids, 318
avrogenes capsulatus. See Bacillus
welchii.
anthracis, immune serum of, 545 mode of action of, 327
choleræ gallinarum, aggressins of, 557 coli, cholelithiasis and, 948
exaltation of virulence of, in
body, 324 fermentation of sugars by, 119
phagocytosis of, by endothelium, 322
tissue predisposition to, 408 tolerance of, to acids, 318
toxins of, effects of, on muscle, 306
urinary calculus formation and, 940
vaccination against, 555
diphtheria, antitoxin production of, 125
ectotoxins of, 329
endotoxins of, 543
tissue predisposition to, 408
toxins of, 315
effects upon nervous sys-
tem, 304
dysenteria, 119
aggressins of, 557
enteritidis, group reactions, 530
megatherium, 120
mesentericus vulgatus, 120
paratyphoid forms, fermentation by, 119
perturbans, 119
pestis, transmission of, by insects, 326
pyocyaneus, ectotoxins of, 329
endotoxins of, 543
infection through skin and, 327
septicus putidus, toxins, action of, on heart, 308
tetani, ectotoxins of, 329
tuberculosis, endotoxins of, 329
growth of, in particular tissues, 408
not inherited in man, 206
passage of, from alimentary canal into tissues, 320
presence in hens' eggs, 206
typhosus, aggressins of, 557
cholelithiasis and, 948
endotoxins of, 329
indol production by, 119
toxins, action of, on heart, 308
transmission of, by insects, 326
welchii, 987
acony degeneration. See Amyloid.
acteria, adaptation in, 118
in alimentary canal, 383

Bacteria as causes of disease, 313 channels of entry of, 326 composition of, Vaughan, 316 exaltation of virulence of. See Virulence. excretion of, 322 grouping of, as regards toxicity, 314 influence of traces of metals upon growth of, 89 metaplasia of, 646 in normal organs, 321 presence within tissues, 319 nuclear matter in, 29 pathogenic effects of passage of, 328 influence of number present on, 325, 327, 330 normal presence of, on body surfaces, 317 specificity, 328 photogenic, 69 properties of, 313 rate of multiplication of, 121 (note) Bacteriemia, 459 Bacteriolysins, 533 Bacteriolysis, 542 by body fluids, 565 Balantidium coli, 342 Barium chloride, 306 action of, 308 salts of, action of, 307 Barlow's disease, 393 Basedow's disease. See Exophthalmic goitre Bashford's theory of malignant growths, 842 Basset hound, inheritance in, 174 "Baustein" theory, 38 Bedsores, 284 Bee, nerve cells of, fatigue and, 402 Begonia, regeneration in, 140 Benign tumors, 669 metastasis of, 684 Benzoic acid, 81 Beri-beri, racial susceptibility to, 162 Bezoar, 954 Bile, bactericidal action of, 318 resorption of, 388 salts, effects of, on leukocytes, 307 hemolytic action of, 306 Wassermann reaction and, 549 secretins and, 365 Bilharzia, 350 cancer and, 840 Bilharziasis, 350, 780 Biliary calculi, 946 Bioblasts, 33 Biogen, 75 (note) Biont, 75 (note) Biophores, chromatin and, 146, 152 definition of, 75 differentiation of, 124, 130 growth of, 98 inheritance and, 187 interaction of, parental, 188 Weismann's conception of, 134

Biophoric molecules, 54 theory, résumé of, 156 Bioplastic activities of cell, 101 Biotic energy, 59 Birth palsies, 225 Bladder, hypertrophy of, 594 metaplasia of, 643 papilloma of, 781 Blastoma, autochthonous, 668 benign, 669 degenerative changes in, 689 heterochthonous, 667 malignant, 670 stages of, 683 multiple, 695 nerves of, 689 nuclear changes in, 690 retrogression of, 692 teratogenous, 663 unicentric and pluricentric, 693 vessels of, 689 Blastomeres, displaced, cause of teratomas, 659 Blastomycetes, cancer and, 794 Blended inheritance, 167 Blood coagulation. See Coagulation. glycolytic properties of, 365 myelocytes of, 736 poisons acting on, 306 specific gravity of, alterations of, in shock and collapse, 584 Bloodvessels, regeneration of, 614 Bombinator bufo, 550 Bone, atrophy of, senile, 874 chondriform, 610, 611 fracture of, pyrexia following, 485 grafting, 636 hypertrophy of, 593 lipoma of, 722 marrow, vicarious hypertrophy of, 597 membraniform, 610, 611 metaplasia of, 730 regeneration of, 610 Bordet-Gengou phenomenon, 547 Bordet's experiment, 534 Bothriocephalus latus. See Dibothrio-Bradydactylism, due to mutation, 182 inheritance of, 209 Brain cells, compensatory hypertrophy of, 596 Branchial, 852 Breast. See Mammary gland. carcinoma of, metastasis in, 682 Bromides, 305 action of, on skin, 312 Brown-Séquard's experiments on guineapigs, 199 Brunner's glands, vicarious hypertrophy of, 597 Burns, 290, 373 cloudy swelling in, 883 pyrexia following, 485

Bursal cysts, 860 Butterflies. See Lepidoptera. Butyric acid, 93, 386

С

CACHEXIA, 672 strumipriva, 354 Caffeine, 305 Caisson-disease, 285 Calcareous concrements, 934 incrustation, 936 Calcification, 924 causation of, 930 chemical reactions in, 925 experimental production of, 929 metastatic, 926 microchemical reactions in, 925 of myoma, 745 necrotic, 927 theories of, 932 Calcium, bone formation and, 394 carbonate calculi, 948 oxalate calculi, 943 deposits of, in kidney, 930 salts, in parathyroidectomy, 355 soaps, 94 calcification and, 932 in fat necrosis, 982 Calculus. See Concrements. Calories, 479 Calorimetry, 482 Cambridge mathematical tripos, 178 Cancer, 789. See Carcinoma. bodies, 341, 792 incidence of, according to locality, 839 parasites, 838, 840 therapy, 847 Canities, 972 Cantharidin, 311, 885 apillaries, growth of, in granulation tissue, 427 Capillaries. Caproie acid, 93 Carbohydrates, excessive consumption of, 393 Carbon dioxide, 304 effects of excess of, in blood, 382 monoxide, 306 Carbonic acid, fatigue and, 401 Carbonization, 290 Carcinoma, 670, **789** basal-celled, 790, 799 of breast, metastases in, 682 colloid, 804 nature of colloid matter in, 889 definition of, 707 duct of, 803 experimental production of, by x-rays, 846 glandular, 801 of intima, 682 latency of, 686 of lung, 682 metaplastic, 800, 805 modes of extension of, 795 origin of, from inflammation, 619

Carcinoma, parasitic theory of, 792 of prostate, 682 sarcomatodes, 705, 808 squamous-celled, 797 treatment of, principles of, 847 Carnaubic acid, 919 Carnin, 376 Carotid gland, tumors of, 834 Carrel's transplantation experiments, 637 Cartilage, inflammation of, 437 metaplasia of, 644 into bone, 640 regeneration of, 609 transplantation of, 633 Caseation, 441, 986 Casein, 57 Casts, urinary, 899 Cat, idiosynerasy toward, 410 Catalysis, 73 Catalysts, negative, 78 Causes of disease, 201 abnormal bodily states, 311 cell disuse, 402 nutritional, 391 overstrain, 396 starvation, 395 chemical, 298 endogenous, 352 disintegrative, 370 intermediate, 382 internal secretory, 352 metabolic, 375 exogenous, 303 non-parasitic, 303 parasitie, 313 bacterial, 313 metazoan, 343 protozoan, 331 mechanical, 281 physical, 285 Cavernoma, 814, 817 Cell activities, reversibility of, 387 bioplastic, katabiotic activities and, See Cytoplasm. body. chromidia. See Chromidia, carbohydrates of, 90 chemistry of, 54 non-proteid constituents, 85 connections of, 35 differentiation of, 129 adaptation and, 130 in terms of biophoric theory, disuse as cause of disease, 402 division. See also Mitosis, economics of, 38 epithelioid, 440 fatty acid crystals in, 94 germ, origin of, 145 giant. See Giant cells. granules, 33 histology of, 29 inclusions, 791 malignancy, 674

Cell membrane, part played by phos-phatides in, 96 "mother cells," 141 Chloroma, 737, 972 multiplication, 112 indirect, 115 nucleus of. See Nucleus. physiology of, **42** reduction of chromatin in, 146, 147 regeneration, limits of, 141 relationship of surface area and mass. "rests." 695, 835 metaplasia and, 639, 640 as origin of cancer, 795 simple atrophy of, 867 salts of, 88 size of, in relationship to shape and function, 37 states of cell activity, 109 functional, 111 hyperactivity, 111 subnormal, 109 vegetative, 110 theory, intercellular substance in relationship to, 38 totipotent, 653, 655 vegetative versus functional, 142 water of, 85 Cellulitis, phlegmonous, 987 Cellulose, nuclear activity in secretion of, Centrosome, 34 in amitosis, 114 in mitosis, 115 Cephalhematoma, 225, 864 Cephalothoracopagus, 247 Cercomonas, 336 (note) Cerebratulus, eggs of, 34 Cerebrosides, 93, 97 Cerianthus, heteromorphosis of, 647 Cerumen, 955 Cestodes, 343, 350 Chalicosis, 973 Cheek, tumors of, mixed, 662 Cheiloschisis, 271 Cheloid, 716 of skin, 716 Chemiotaxis of capillary bulbs, 615 epidermal, 838 homotropism, 636 negative, 565 neurotropism, 627, 628 in protozoa, 416 of tissue cells, 619 Chemiotropism, 626 Chicken cholera, 493 Chigger, 351 Chilblains, 287 Chills, 467 Chloasma uterinum, 970 Chloral hydrate, action of, 308 action of, on heart, 307

death under, 583

Chlorophyll, metabolism, 90 Chlorosis, abnormal gastro-intestinal fermentations and, 386 Cholelithiasis, 946 Cholesteatoma, 832, 902 Cholesterins, 92, 94 biliary, origin of, 950 calculi, 946 inhibitory action of, on hemolysis, 542 (note) neutralization of nitrites, 504 oleate, myelins and, 918 Cholin, 385 in autolysis of nervous tissues, 373 Chondrin, 39 Chondroalbuminoid, 57 Chondroid infiltration. See Amyloid infiltration. Chondroitin-sulphuric acid, 889, 895 Chondroma, 724 of mammary gland, 227 metastases of, 684 of mouse, latency of, 687 ossifying, 728 of testis, 726 Chondromucoid, 57 Chondrosarcoma, 770 Chondrosin, 889 Chordoma, **761** Chorio-angiopagus. See Fortus acardiacus. Chorio-epithelioma, 664 Chromaffin system, 358 Chromatin, biophoric matter and, 152 reduction of, in germ cells, 147 in somatic cells, 140 relationship to biophoric matter, 146 Chromatolysis, 52, 690 Chromatophoroma. See Melanoma. Chromatophore, 826, 829 Chromidea, 32 cancer bodies and, 341 in cloudy swelling, 884 melanin granules and, 48 mode of development of, 48 Chromoproteins, 57 Chromosomes, aberrant, in malignant growths, 841 as anatomical basis of heredity, 153 in mitosis, 115 number of, in different species, 147 reduction process in germ cells, its effects, 190 in spermatogenesis, 148 Chronic, definition of term, 421, 439 infection, 461 Chylangioma, 860 Ciliata, 341 Circulation, organs of, poisons acting on,

Circulatory system, effect of toxins on, Complement, fixation of, **547** 470 junction of, with amboce

Circumcision, non-inheritance of results, 195 Cirrhosis, hepatic, as a subinfection, 463

hypertrophic, 592 of liver, abnormal gastro-intestinal fermentations and, 381 from lower fatty acids, 394 obstructive, 388 pigmentation in, 961 Cirsoid aneurysm, 814 Cloaca, defects of, 274 persistent, 268, 276 Cloacal membrane, defects of, 268 Cloudy swelling, 882 in starvation, 872 Clover, mutation in, 181 Clupeine, 56 Coagulation of blood, absence of, in asphyxia, 397 in death from overstrain, arrested by ankylostoma extract, 348 necrosis, 374, 426, 984 hyaline degeneration and, 900 Coagulins, 524 lecithide, 538, 540 of duodenum, 778 Coccygeal gland, tumors of, 834 Cock's spur experiment, 598 grafting of, 633 Cohnheim's theory, 835 Cold as a cause of disease, 286 as a predisposing agent, 288 Coley's fluid, 692 Collagen, 39, 56 fibrils, 711 Colliquative necrosis, 985 Colloid menstruum, crystallization in, 940 properties of, 87 Colloidal membranes, 45, 88 suspension, 952 Color-blindness, 163 sex-limited, 180 Coma, diabetic, 381 Complement, 535 action of heat upon, 535 artificial, 540 diversion of, 546

junction of, with amboceptor, 536 multiplicity of, 538, 567 soaps acting as, 540 source of, 566 structure of, 539 synonyms of, 566 variation in amount of, 539 Complications (of infections), 456 intra-uterine, 217 Concrements, 934 biliary, 946 bilirubin, 947 calcium carbonate, 948 cholesterin, laminated, 947 pure, 946 common, mixed, 947 etiology of, 948 calcareous, 934 corpora amylacea, 937 intestinal, 953, 954 phleboliths, 936 prostatic, 937 rhinoliths, 935 salivary, 935 tonsillar, 935 urinary, 938 calcium oxalate, 943 fibrinous, 946 guanin, 945 uratic, 941 Concussion, 282 Conditions that cannot be inherited, 205 Condyloma, 777 Congenital cysts of neck, 273 meaning of term, 201 sacral teratoma, 239, 654, 656 Conglutination as a property of colloids, See Elastic connective tissue. fatty. See Fatty tissue metaplasia of, 644 new formation of, in inflammation, 428 tumors, 710 white fibrous, regeneration of, 605 Consanguines, marriage of, 163, 213 Constipation, 382 Contagious disease, 326 Contrecoup, 282 Convalescence, 456 Convallamarin, action of, 307

Coolie itch, 346

1006

Copper, action of, on liver, 311 aseptic suppuration by, 431 pigmentation, 976 poisoning, 976 salts of, action of, 307 Coprosterin, 95 Copula, 566 Cornea, inflammation of, 437 Corpora amvlacea, 937 Corpus luteum, 864 cysts of, 857 internal secretion, 362 Craniopagus, 247, 248 Cranioschisis, 263 Creatin, 305 Cretinism, 353 Cretins, 259 Crinoids, experiments upon development Daltonism. See Color-blindness. of, 139 Crisis, resolution by, 466 Croton oil, action of, 309 Croup, fibrinous, 436 Crow, hematozoa of, 337 Crustacea, reaction of, to injury in, 418 regeneration in, 603 Cryptogenic infection, 324 Cumulative inheritance, 180 Curare, 304 Curves of variation, 19 Cyanogen, action of, on blood, 306 Cyanophycex, 30 Cyanophilia, 995 Cyanosis, 311 Cyclops, 260 Cyclotia, 273 Cylindroma, 824 hyaline degeneration and, 901 Cynarase, 508 Cystadenoma, 783, 784, 850, 859 papilliferum ovarii, 859 Cysticercus cellulosa, 345 Cystin, 378 calculi, 945 Cystinuria, 378 Cytotropic substances, 552. See also Opsonins. Cysts, 850 congenital, of neck, 273 hemorrhagic, 863 intracystic papilloma, 783 lymphangiomatous, 820 necrotic, 864, 987 ovarian, contents of, 889 parasitic, 865 secretory, 851 composite, 862 endothelial, 860 ependymal, 861 epithelial, 862 glandular, of antenatal origin, of neoplastic origin, 859 of postnatal origin, 858 Cytase, 566. See also Complement. Cytolysis, 532 enzyme action and, 574

Cytolysis, experimental, 582 maternal, effects upon fœtus, 219 mechanism of, 534 normal, 866 Cytoplasm, formation of starch by, 43 granules in, 33 influence of, upon nucleus, 139 motility and, 44 nucleus and, 157 relationship to nucleus, 38 respiration and, 42 structure of, 32 Cytosin, 64

D

DACRYOPS, 858 Danyscz's phenomenon, 520 Daphnia, 419 Darwin's experiment on reversionary inheritance, 176 Death. 990 from adrenal ablation, 359 antenatal, 215 from overstrain, 397 from pituitary ablation, 357 pathological, 993 physiological, 992 signs of, 993 Deciduoma. See Chorio-epithelioma. Decomposition, 994 Decubitus, 284 Deduplication, 234 fusional, 244 Defences of organism, 316 Defensive proteins, 497 Defervescence, 456 Degeneration, amyloid, 891 calcareous, 924 chondroid. See Amyloid, colloid, 890 elastoid, 896 familial, 177 fatty, 908 of uterine muscle, 868 fibrinous, 887 glycogenous, 921 granular, 885 hematohyaloid, 898 hyaline, 896, 900 hydropic, 903 infiltration and, 881 keratinoid, 902 lipoid, 915 mucoid, 888 of nucleus, stages and forms of, 53 pigmentary, 956 red, 746 vacuolar, 904 vitreous. See Elastoid degeneration. Wallerian, 626, 870 waxy, 887. See also Zenker's degeneration. Delhi boil, 334 Demarcation, line of, 988

Dengue, 995 Dentigerous cysts, 863 Dentition, disorder of, racial susceptibility to, 163 Dentoplasm. See Paraplasm. Dermoid, ovarian, 256, 654 sequestration, 862 Desmon, 566 Determinants, 133, 134 argument against, 163 Development, influence of cytoplasm upon, 139 "Diabéte bronzé," 961 Diabetes, diathesis for, 210 gouty diathesis and, 212 mellitus, 365 glycogen in, 922 hepatic type, 367 phloridzin, 91, 367 racial susceptibility to, 163 Diacetic acid, 380 Diapedesis of leukocytes, 423 Diaphoresis, 311 Diaphragm, abnormalities of, 268 Diarrhœa, 309 collapse and, 581 Diathesis, 163, 204, 210 anatomical conditions and, 201 congenital cystic, 857 nervous, 211 Diathetic reversion, 177 Dibothriocephalus latus, 347, 348, 349 Dicephalus, 243 Differentiation of cells and tissues, 129 Digestive system, action of poisons upon, 308 in fibrile state, 476 Digitalin, action of, 307 Digits, accessory, inheritance of, 163 Dimethylamin, 385 Dioctophyme renale, 345 Dipeplids, 60 Diphtheria antitoxin, 510 Diplococcus pneumonia. See Pneumo- Electricity as a cause of disease, 295 Disease, conditions of, classification, 203 contagious, 326 conveyance of, by food, 327 definitions of, 21, 22, 128 as distinguished from affection, 21 of foetus, 224 local and general, 20 postnatal acquirement, 280 variability and, 17 Disintegration, granular, 885 Disintegrative intoxications, 370 Distension, 284 Distoma hepaticum, 348 Disuse atrophy. See Atrophy. as predisposing cause of disease, 403 Diversion of complement, 546 Dominant characters, 168

Dominant properties, 164 Dorsal groove, defective closure of, 263 Dorsipolar fusion, 248 Drosera, 47 insectivorous tentacles of, 571 Dum-dum fever. See Kala-azar. Duodenum, coccidiosis of, 778 potential sterility of, 383 sterile state of, 317 Duplicia symmetros, 655 Dwarfism, 259 from division of ovum, 152 Dyspepsia, racial susceptibility to, 163 Dyspncea, 381, 391

E

EARTH-WORM grafting, 631 heteromorphosis, 646 regeneration in, 601 Eechymoses in asphyxia, 391 from drugs, 312 Echinococcus, 345 Echinus, chromosomes in, 148 connections between blastomeres, 36 eggs of, effects of shaking of, 217 experiments on eggs of, 138 Eclampsia, 363 Ectopia cordis abdominalis, 270 vesica, 268 Ectotoxins, 510 Eczema, gouty diathesis and, 212 Edestin, 56 Eel, glycosuria in, 365 poison, 550 Egg albumin, 56 formula of, 55 (note) Ehrlich side-chain theory of immunity, 516, 570 Elastic connective tissue, atrophy of, senile, 874 regeneration of, 607 Elastin, 39, 56 Elastoid degeneration, 896 constant and alternating currents, 296 fulguration, 295 Electrolytes, 86 Electropuncture, 693 Eleidin, 902 Elephantiasis, 228, 718 congenital, 224 Emaciation in febrile state, 477 Embolism, 977 capillary, 312 in malaria, 339 cell, 680 placental-cell, eclampsia and, 363 septic, 434 Embolus, placental-cell, 665 Embryo, amitosis in, 113 definition of, 202

1008

Epithelium, connections between cells Embryoma, ovarian, 654 sporadic, 656 testicular, 654 chorio-epithelioma and, 666 Embryonic tissues, transplantation of, 633 Emphycema, pulmonary, 875 Emulsin, 508, 509 Endocarditis, acute non-specific, immunity in, 500 Endogenous intoxications, 352 Endothelioma, 813 Endotheliotoxin of snake venom, 550 Endothelium, connection between cells of, 35 regeneration of, 620 vascular, bactericidal activity of, Ether, 304 relationship of connective-tissue cells, 606 Endotoxins, 315, 496, 512, 543 ectotoxins and, 329 Enostoses, 729 Entambed dysenteria, 333 "Entdifferenzierung," 141, 641 Enteric fever. See Typhoid fever. Enzyme, 70 action, 70 growth and, 75 immunity and, 573 intracellular and reversible, 83 reversibility, of, 79 surface action and, 78 in adrenal, 360 bacterial, 314 comparison with nucleoproteins, 66 immunization against, 508 intracellular and extracellular, 72 ionic action and, 576 of leukocytes, 371 mode of action of, 73 production of, adaptive, 118 properties of, 70 Eosinophilia, 740 produced by metazoan parasites, 348 Ependyma, cysts of, 861 Ependymoma, 758 Epidermis, regeneration of, 618 Epigenesis, **131** Epignathus, 239, 656 Epiguanin, 376 Epilepsy, 386 diathesis and transmission, 211 diathetic reversion and, 179 from parental intoxications, 220 inheritance of, 211, 214 traumatic, inheritance of, 199 Episarkin, 376 Epispádias, 268 Epithelioid cells, 440 Epithelioma, 797

x-ray, 846

Epithelium, amorehoid movement of, 425

of, 35 metaplasia of, 648 regeneration of, 616 Epitoxoids, 515 Epoöphoron, 274 Epulis, 713 Ergot, 305 action of, 308 "Ersatz" theory, 877 Ervthema, 312 Erythrocytes, changes of, in febrile state, 472 fragility of, 967 life period of, 42 origin of hemoglobin, 50 regeneration of, 615 Ethyl alcohol, a constituent of normal tissues, 54 butyrate, 82 Etiology, 201 definition of, 25 Eustrongylus gigas, 345 Evening primrose. See Œnothera. Eventration, 267 principles not an explanation of, 187 in terms of biophores, 157 Exencephaly, 264 Exogenous intoxication, bacterial, 464 non-parasitic, 303 parasitic causes of, 313 Exophthalmic goitre, 354, 355 abnormal gastro-intestinal fermentation and, 386 Exophthalmos, 356 Exudate, inflammatory, 423, 446 Eye, choroid of, metaplasia of, 639 iris of, pigmentation of, Mendel's principles affecting, 173 particulate inheritance of, lens of. See Lens. retina of, glioma of, 757

FACIAL elefts, defective closure of, 270 Facies Hippocratica, 585 Factors of safety, 106 Fallopian tube, an open channel, 316 Familial degeneration, 177 inheritance, 163 Fasciola hepatica, 348 Fastigium, 456 Fasting, 393 Fat cells, lymphoid tissue and, 60 origin of, 607 senile changes in, 873 vacuoles in nuclei of, 50 concretions, 955 9

Fat, excessive consumption of, 393 metabolism of, 93 necrosis, 982 soaps in, 94 neutral, 92 Fat phanarosis, 919 Fatigue, 396 Sherrington's synapse theory of, 400 Fatty acids, 92 in autolysis, 371, 372 crystals of, in cells, 94 degeneration, 907 infiltration, 905 tissue, regeneration of, 607 Febrile state, 465 associated disturbances, blood, Fissura buccalis, 271 472 circulatory, 470 cutaneous, 477 digestive, 476 emaciation, 477 muscular, 469 nervous, 467 respiratory, 473 urinary, 474 pyrexia in, grades of, 467 site of heat production, 487 thermogenesis and, 479 significance of, 488 temperature in, causes of development of, 485 changes of, 466 varieties of fever, 466 Ferment. See also Enzyme. action. See Enzyme action. organic and inorganic, 70 Fertilization, 144 self-, 169 Fervescence, 456 Fever. See Febrile state. from overstrain, 398 Fibrinogen, 56 Fibrinous adhesions, 430, 433 Fibroblast, development of, 429 Fibroglia fibrils, 711 Fibroin, 56, 60 Fibroma, 452, 710 Fibromatoid, 451, 714 Fibromatosis of olfactory nerve, 715 of optic nerve, 716 Fibromyoma. See Leiomyoma. metastases of, 684 Fibrosarcoma, 712 growths, 718 inflammatory, 452 postfibrinous, 452 non-inflammatory, neoplastic, 452

Fibrosis, non-inflammatory, strain, 452 peri-arterial, 876 of placenta, 222 senile, 876 strain-, 451 Filaria bancrofti, 351 medinensis, 346, 350 nocturna, 344, 346 Filariasis, 350 First intention, healing by, 423 Fischer's experiment, 838 Fish eggs, separation of blastomeres in, malignant goitre in, 840 poison, 550 sterni, 267 vesicogenitalis, 267 Fistula, 433 congenital, of neck, 273 Fixateur, 566. See also Amboceptor. Fixation of complement, 547 Flowering plants, accessory somes in, 154 Flukes, 343 Fluorescent substances as sensitizers to Fortus acardiacus, 218, 231 amorphous, 231 diseases peculiar to, 224 distinction from embryo, 202 infection of, 206 internal secretion and lactation, 363 Foraminifera, calcification in, 46 Formic acid, 386 Fowls, eggs of, conveyance of tuberculosis through, 206 Mendel's principles affecting, 173 Fracture of bones, pyrexia following, 485 intrauterine, 257 of parturient origin, 225 Freezing, effects of, on tissues, 286 Frog, blastomeres of, 137 development of, 137 skin, employed for grafting, 635 species and varieties of, 172 tadpole, absorption of tail, 624 adaptation in, 128 regeneration in, 601 tongue of, in inflammation, 422 web of foot of, in inflammation, 422 Fulguration, 295 Fundulus, effects of crossing with men-

G

GALACTOCELE, 858 Gallstones, 946 Galton's law, 174 "Ganglion," 860 cells. See Neurons,

Ganglioneuroma. See Neuroma. Gangrene, 987, 988 dry, 988 emphysematous, 987 frost, 287 hospital, 987 moist, 988 Gartner's duct, 853 development of, 275 Gastric juice, a protective mechanism, secretions and, 365 Gastro-intestinal intoxications, 382 Gelatin, 56 reaction in, 549 Genital passages, infection through, 327 Geotropism, 602 Germ cells, arrested development of offspring due to imperfections of. 208 influence of environment on, 187 of parental intoxication on, Gout. 375 196 origin of, 145 Germinal period, 202 Giant cells, 32 foreign body in, 417 tuberculosis, 440 types of, 733 Giantism, 227 itism, 227 local, from intra-uterine obstruction, Gregarina, 339 Gromia, 37, 886 pituitary and, 357 Glanders, giant cells in, 439, 440 granuloma, 443 Gliadins, 56 Glial cells. See Neuroglia. Glioma, **755** metaplasia of, 644 of retina, 757 Gliosarcoma, 757, 773 Globin, 56 Globinuria, 958 Globulins, 56 as a factor in immunity, 550 Glossina morsitans, 336 palpalis, 336 figure of, 335 Glucosamin, 889 Glucose, 80 Glucosides, action of, on heart, 307 immunity toward, 504 Glutamic acid, 59 Glutelins, 56 Glutenin, 56 Glycin, 60 Glycocoll, 59 form hippuric acid, 81 dehydration products, 60 Glycogen, absence of, fatigue and, 401 in febrile state, 476

Glycogen metabolism, 91 mode of accumulation of, in liver cells, 83 in starvation, 872 Glycogenous infiltration, 921 Glycosuria, 365 adrenal, 368 alimentary, 368 nervous, 368 phloridzin, 367 renal, 367 Glycyl, 60 Glycyl-proline anhydride, 60 General paralysis of insane, Wassermann Goblet cells, development of mucin in, 50 Godlewski's experiment, 139 Goitre, colloid, 354, 890 congenital, 219, 224 exophthalmic, 354, **355** in fishes, 840 Gonococcus, phagocytosis of, 325 Gonorrheea, vaccination against, 555 diathesis toward, 210, 212 sex limited inheritance of, 180 Graafian follicles, cysts of, 859 Grafting. 631 Granulation tissue, 426 syncytium and plasmodium, 31 Granuloma, infectious, course of, 439 set up around foreign bodies, 443 Gravel, urinary, 941 Graves' disease. See Exophthalmic goitre. Ground itch, 346 Group reactions, 530 of agglutinins, 530 of amboceptors, 544 of precipitins, 525 Growth, as distinct from multiplication, influence of minute traces of metals upon, 89 its essential nature, 75, 98 nucleoproteins and, 76 nucleus and, 42 nutrition and function, 599 problems of, 588 to other cell activities, 101 Guanase, 376 in autolysis, 371 Guanin, 64 calculi, 945 Guanylic acid, 64 (note), 65 (note) Guinea-pig, anaphylactic phenomena of, Brown-Séquard's experiment on, 199 Mendel's principle affecting, 173 combination with benzoic acid to Guinea-worm. See Filaria medinensis. Gumma, 442 Gummous change, 986 Gums, tumors of, mixed, 662

H HABIT of growth theory, 842 origin of, 105 Hair balls, 954 dominance of dark pigmentation in, regeneration of, 619 Halteridium, 206, 335 Hamartoma, 815 Haptine, 572 Haptophore, 515, 517 Harelip, 270 Hauser's theory of neoplasia, 842 Hay fever, 410 anaphylaxis and, 561 Heart, disease of, effects upon placenta, 218 hypertrophy of, 594 concentric, 994 and dilatation of, from excess of fluid, 392 irritable, 398 poisons acting on, 307 reserve force of, 107 valves of, inflammation of, 437, 438 Heat of body, calories, 479 calorimetric observations of, 482 discharge, modes of, 479 production of, 478 regulating mechanism in, 480 where produced, 487 as a cause of disease, 286 Heatstroke, 289 Helminthiasis, 343 Hemamæba. See Hematozoon. Hemangio-endothelioma, 813, 822 Hemangioma, 817 Hematin, 57 Hematoblasts, 616 Hematobyaloid degeneration, 898 Hematoidin, 960 Hematoporphyrinuria, 962 Hematosporidia, 337 Hematozoon malaria, 337 Hematuria, 958 Hemicephaly, 263 Hemochromatosis, 961 Hemoclastics, 306 Hemocyanin, 57 Hemofuscin, 960, 961 Hemoglobin, 57 formula of, 55 imbibition, 957 origin of, 50 Hemoglobinemia, 306 due to hypisotonic blood plasma, 90 Hemoglobinuria, 306, 957 infective, 959 paroxysmal, 958 Hemolymph glands, 597, 616 Hemolysin, artificial, 540 of snake venom, 550 source of, 568

Hemolysis, 532

inheritance of coat color of, 175

Hemolysis, agents causing, 306 diversion of complement in, 546 drugs causing, 957 focal necrosis and, 982 influence of concentration of medium upon, 90 pyrexia and, 484 solvents of phosphatides as agents in, 96 Hemophilia, 210 due to mutation, 182 familial inheritance of, 163 sex limited inheritance, 180 Hemorrhage as a cause of collapse, 581 effects of, upon specific gravity of blood, 584 pyrexia and, 484 Hemorrhagic cysts, 863 Hemorrhoids, 814, 815 Hemosiderin, 960 Hen, embryos of, transplantation of, 633 Hepatic incompetence, 311 Hepatolysins, Charrin's experiment on gravid goats, 219 Hepatotoxin, 533 Heredity, 159 anatomical basis of, 153 biophores and, 157 forms of, 160 Hermaphroditism, 278 Hernia, diaphragmatic, 268 funiculi abdominis, 267 Herpes zoster, 449 Heterolysin, 533, 868 Heterolysis, 372 Heteromorphic inheritance, 179 Heteromorphosis, 603, 646 Heteroplasia, 641 Heterotopia, 640 Hexone bases, 58 synthesis of, in kidney, 81 Histidin, 58 Hippuric acid, 81 Histolysis, normal, 866 Histones, 56, 63 Histozyme, 81 Hodgkin's disease, 739 Hofmeister's theory of the protein molecule, 61 Hog cholera, immunization against, 496 Homeomorphic inheritance, 179 Homogentisic acid, 379 Homotropism, 636 Hook worm. See Ankylostoma. Hordein, 56 Hormones, 353 pancreatic, 366 properties of, 365 Horns, 777 of sheep, inheritance in, 180 Horse, albino breed, mutation and descent, 213 callosity of foreleg, 182

1012

Hunter's grafting experiment, 598, 633 Hyaline degeneration, 896 matter, 40 Hyalopus dujardinii, 886 Hybridization, principles of, 168 Hydatid cysts, 345, 351, 865 mole, 664 of Morgagni, 274, 854 sessile, 855 Hydra, regeneration in, 140, 601 Hydramnios, 224 Hydrobilirubin, 968 Hydrocele, encysted, of testis, 854 of fourth ventricle, 861 of scrotum, 860 Hydrocephalus internus, 861 of parturient origin, 225 Hydroeyanic acid, 306 Hydrometra, 858 Hydronephrosis, 863 Hydropic degeneration, 903 osmosis and, 90 Hydrops cystidis fellea, 858 ex vacuo, 876, 905 Hydroquinone, 306 Hygroma, 860 Hyloma, 703 Hyocyanine, 305 Hypamnios, 224 Hyperchromatosis, 53 nuclear, 691 Hyperemesis gravidarum, collapse and, Hyperemia, induced, Bier's method, 450 Hypernephroma, 810 Hyperplasia, 592 irritative glandular, 788 lymphadenoid, 740 Hyperpyrexia, 467, 480 (note) Hyperthermia. See Pyrexia. Hyperthyrea, 355 Hyperthyroidism, 353 Hypertrophy, 587 adaptive, 593 compensatory, 594 concentric, 994 congenital glandular, 788 as distinct from regeneration, 596 from hyperactivity, 111 rritative, 597 glandular, 788 of liver, 786 neoplasia and, 785 nutritional, 597 physiological, 593 progressive, 714 simulated, 600 sympathetic, 599 vicarious, 596 Hypnosis, inflammation and, 449 Hypnotic drugs, 304 Hypochromatosis, nuclear, 691

Hypogenesis, polar, 260

Hypoleukocytosis, 502

Hypophysis cerebri. Sce Pituitary. Hypophasia, 260 senile, 874 Hypospadias, 278 inheritance of, 209 Hypothermia, 288 Hypoxanthin, 64, 376 action of, on muscles, 305 Hysteria, diathesis and transmission, 211 Hystero-epilepsy, racial susceptibility to,

I

due to mutation, 182 Icterus. See Jaundice. Idants, 134 Idiocy from parental intoxications, 220 a result of familial degeneration, 177 Idiosyncrasy, 312, 410 Ids, 134 Ileus, 383 Ilyanassa, 140 dwarf larvæ from isolated blastomeres, 216 Imbecility from parental intoxications, Immune body, 535. See also Amboceptor, orders of, 572 serum, 510 Immunity, 491 absolute and relative, 491 acquired, inheritance of, 194 active and passive, 497 adaptation and, 121 anaphylaxis and, 559 atreptic, 848 comparison of, with enzyme action, historical review of, 492 inheritance of, 179 non-specific, 409 of offspring from parental intoxica-tion or infection, 209 orders of, 499 against phytotoxins, 505 abrin, 505 ricin, 505 robin, 507 substances of unknown constitution, 508 cells. bacteriolysins, cytolysins, 532 enzymes, 508 proteins, 524 agglutinins, 527 aggressins, 557 opsonins, 551 venoms, 550 non-specific, 491 leukocytes and, 501 toward substances of known constitution, 501

NDEX

Immunity, orders of, toward substances | Inflammation, acute catarrhal, 436 of known constitution, arsenic, 502 glucosides, 504 morphine, 504 passive, 503 theories of, 562 Immunizations, 491 Implantation, 631 cysts, 862 of teeth and bone, 636 Inbreeding, dwarfism and, 260 Inclusions, abdominal, 655 foetal, 254, 655 Incompetence, hepatic, 311 Incrustation, calcareous, 936 Incubation period, 455, 457 Indol, action of, on liver, 311 as a cause of gastro-intestinal intoxications, 384 production of, by bacteria, 119 Indolacetic acid, 384 Inertia, physiological, 105, 109 Infancy, insusceptibility during, 406 Infantile scurvy. See Barlow's disease. Infarct, pyrexia produced by, 373 Infection, 453 bacterial, 314 causation of, 454 cause of stillbirth and abortion, 219 conveyance of, through foetal memcourse of, 455 cryptogenic, 324 definition of, 453 distinguished from inflammation, 415 epidemic, 453 epizootic, 453 favored by overwork, 399 febrile state and, 465 significance of, 488 incapable of inheritance, 206 incubation period, 457 malnutrition as predisposing cause, 407 modes of, 323 predisposition toward, 405 previous, influence of, 406 pyrexia and thermogenesis, 478 sporadic, 453 stages of pyrexia, 467 terminal, 992 transmitted to offspring, 221 types of, 459 acute, 460 chronic, 461 fulminant, 459 subinfection, 462 varieties of fever, 466 Infiltration, amyloid, 891 degeneration and, 881 fatty, 905 glycogenous, 921 Inflammation, 412

cellulocutaneous, 434 fibrinous, 430, 435 healing of, by first intention, 425 by granulation, 426 hemorrhagic, 435 leukocytes in, 445 membranous, 435 or diphtheritic, 435 in non-vascular areas, 437 phlegmonous, 434 stages of, 421 suppurative, 430 adaptation in, 447 adequacy of, 450 causes of, 420 chronic, 439 diffuse, 443 infectious granulomas, 439 leukocytes in, 445 comparative pathology of, 416 definition of, 412, 415 distinguished from infection, 415 exudate in, 446 fibrosis and its relationship to, 450 general reaction accompanying, 448 hypnosis and, 449 leukocytes in, 445 main data regarding, 444 metastasis in, 434 nervous system in, 448 part played by vessels in, 444 referred, 449 repair and, 447 temperature changes in, 450 tissue proliferation in, 444 ulcerative, 432 Infusoria, 341 adaptation in, 128 Inheritance, 159. See also Heredity. of acquired characters, 192 acquired immunity, 194 disuse atrophy, 196 intra-uterine arrests of development, 195 maternal impressions, 193 mutilations, 195 retrogressive changes due to intoxications, 196 use acquirements, 194 of anatomical abnormalities, 209 atavistic, 175 collateral, 175 cumulative, 180 diathetic reversion in, 177 of epilepsy, 211, 214 false, 201 familial, 163 degeneration, 177 forms of, classification of, 186 homeomorphic and heteromorphic, homeomorphous and heteromorphous, 211 individual, 167

Inheritance, mosaic, 175 mutation or spontaneous variation, parental, 167 particulate, 167 of polydactylism, 209, 214 racial, 162 reversionary, 176 sex limited, 164, 179 theory of, 185 true and false, 201 Inherited morbid conditions, 205 Iniencephaly, 265 Injury, local reaction to, **412** Inosinic acid, 64 (note) Intensification of virus, 495 Insecta, accessory chromosome in, 154 germ cells of, 145 Insects, amitosis in, 113 parasitic, 343 transmitters of bacterial infection, 326 KALA-AZAR, Leishman-Donovan bodies lation, 290 in, 332, 334 Insolation, 290 Intercellular substances, 38 Intermediate body. See Amboceptor. intoxications, 382 Intermittent fever, 466 Internal secretions, as causes of disease, 352 general principles of, 368 in hypertrophy, 598 Intestinal concretions, 954 sand, 935 Intestines, hypertrophy of, 594 obstruction of, results of, 383 Intoxications, acid, fatigue and, 401 effects of, upon offspring, 196 endogenous, **352** exogenous bacterial, 464 and endogenous, 300 gastro-intestinal, 382 "intermediate," 382 paternal, influence of, upon offspring, 194, 197, 198, 220 Invertase, 575 reversible action of, 82 Involution of uterine muscle fibers, 868 Iodides, action of, on skin, 312 Iodipin, 914 Iodoform, action of, on skin, 312 Iodothyrin, 353, 891 Ion-proteids, 86 Ions, 85 evolution of energy in combination of, 102 growth and, 100 Ipecacuanha, action of, 509 Iron in calcareous deposits, 925 carriage of, by leukocytes, 320 need for, in organism, 394 as a nuclear constituent, 63 Ischiopagus, 247, 253 Islands of Langerhans, 367 Isocholesterin, 95

Issaeff's resistance period, 501

JANICEPS, 247 Janus, 247 Jaundice, 963 hemohepatogenous, 967 infective, 967 neurotic, 967 obstructive, 387, 965 toxemic, 967 Jecorin, 92, 371, 392, 916 Jelly fish. See Medusa. Jennerian vaccination, 493 Joints, articular laxity, due to mutation, loose cartilages in 727

к

Kallima paralecta, 134 Karyorrhexis, 53 Katabiotic activities of cell, 101 Katadidymus, 240 (note) Katalysis. See Catalysis. Kataplasia, 878 anaplasia and, 880 Keloid, 716 (note) Kephalin, 92 Kephir lactase, 82 Kerasin, 97 Keratin, 56 Keratinization, pathological, 902 Keratohyalin, 902 Kidneys, action of poisons in, 311 cloudy swelling in, 838 compensatory hyperplasia of, 595 congenital cystic, 855 disease of, effects of, upon offspring, 219 hypertrophy of adrenals and, 360 embryogeny, 856 glycosuria, 367 infiltration of, fatty, 918 lipoma of, 722 parenchymatous nephritis of, soaps in, 94 regeneration of, 604, 621 reserve force of, 108 resorption of urine, 388 retention cysts, 857 teratoblastoma of, 661 tumors of, 807 "Klopfversuch," 582 Kobelt's tubes, 853 Kyes' experiment, 538 Kyrins, 62

L

LACERATION, 284 Lacrimal gland, tumors of, mixed, 662 Lacrymaria olor, 44

NDEX

Lactalbumin, 56 Lactation, pregnancy and, 363 Lactic acid. 386 bacterial growth in tissues and, disturbances set up by, 381 Lamblia, 336 Lardaceous degeneration. See Amyloid degeneration. Latency of life, 286 (note) Latent period, 105 Law of chance, 18 of multiple proportions, 511, 520, 522 Lead, action of, on liver, 311 poisoning, 304 blue line on gums in, 975 effects of, on offspring, 198, 219 Lecithalbumin, 92 Lecithins. See Phosphatides acting as complement, 538 Lecithoproteins, 57 Leiomyoma, 744 of digestive tract, 749 of skin, 750 Leiomyosarcoma, 772 Leishman-Donovan bodies, 332 Leishman's method, 551, 564 Lens, metaplasia of, 643 regeneration of, 617 Lepidoma, 703, 775 transitional, 703, 705, 806 evolution and preformation in, 133 compounds with glycocoll, 60 produced in autolysis, 371, 372 Leukemia, myelogenous, 736 Leukocidin, 511, 533 Leukocytes, classification of, 445 eosinophile, diapedesis in, 424 origin of, 445 in peritonitis, 307 explosive of crustacea, 418 in febrile state, 472 histogenous, 624 origin of, 445 hvaline, 445 in inflammation, fate of, 446 leukolysis of, 307 lymphocyte, diapedesis in, 424 origin of, 445 margination of, 423 migration of, 423, 445 in non-specific immunity, 501 origin of, 418 passage of, on to mucous surfaces, phagocytosis of, 319 polymorph, connective-tissue formation and, 606

variations in phagocytic power of, 553

Leukocytes, polynuclear, diapedesis in, 424 origin of, 445 regeneration of, 614 Leukocythemia. See Leukemia. Leukocythosis, induced, resistance period and, 450 Leukolysis, 306, 565 Leukopenia, 306 Leukoplakia, 902 Life, definition of, 67 enzyme action and, 77 growth and, 77 latent, 286 (note) period of, predisposition to disease, 406, 407 potential or latent, 68 Light as a cause of disease, 292 Finsen, 830 (note) Röntgen ray dermatitis, 846 ultraviolet, action of, on skin, 830 Lightning, effects of, 295 Limnaus stagnalis, fat formation and, 903 Line of demarcation, 988 Linoleic acid, 93, 919 Lipase, 93 reversible action of, 81, 82 Lipemia, 912 Lipochromes, 93, 97, 972 Lipoids as agents in osmosis, 46 classification of, 92 degeneration of, 915 hypnotics and, 304 Wassermann reaction and, 549 Lipoma, **721** of bone, 722 myxomatodes, 719, 721 of peritoneum, 721 Lipoproteins, 57, 93, 97 Liposarcoma, 770 Liposis. See Obesity Lithiasis, 938 gouty diathesis and, 212 Liver, adenoma of, 684 bacteria in normal, 321 carbohydrate metabolism in, 366 corrhosis of, pigmentation in, 961 coccidiosis of, 778 compensatory hyperplasia of, 95 congenital cysts of, 857 disease of, effects of, upon offspring, fatty infiltration of, 906 glycogenic activity of, 352 metabolism in, 83 hypertrophy and adenoma of, 786 incompetence of, 311 kataplasia of, 879 multiple carcinoma of, 836 origin of myelocytes in, 614 poisons acting on, 310 regeneration of, 604, 620

Liver, reserve force of, 107 Lividity, cadaveric, 994 Living matter, orders of, 77 Locomotor ataxia, 877 Wassermann reaction in, 549 Longinymph state, 211 Luminosity of photogenic bacteria at low temperatures, 69 Lungs, carcinoma of, metastases in, 682 compensatory hyperplasia of, 595 kataplasia of, 879 reserve force of, 108 Luschka's gland, 834 Lymph cysts, 860 nodes, intestinal, bacteria in, 319 peribronchial, bacteria in, 319 tonsillar, bacteria in, 319 Lymphadenoma. See Lymphoma. Lymphadenoid tissue, action of x-rays upon, 294 development of erythrocytes in, Melanin, 828 endothelial cells of, in burns, 373 hyperplasia of, 739 regeneration of, 613 vicarious hypertrophy of, 597 Lymphangiectasis of myoma, 745 Lymphangio-endothelioma, 813, 824 Lymphangioma, 818, 820, 821 Lymphatic vessels, extension of tumors in, 681 Lymphoid tissue, fat cells and, 609 Lymphoma, 737, 742 Lymphomatosis, granulomatous, 741 Lymphosarcoma, 742, 772 Lymphosarcomatosis, 743 Lysin, 58, 59 Lysis, resolution by, 466 M MACROCYTASE, 567 Macrodactyly, 228 Macroglossia, 228 Macronucleus, 34 Macrophages, 566 in tonsils, 319 Macrostomia, 271 Magnesium sulphate, mode of action of, Malaria, 337

in children, 405

Malformations due to intra-uterine disturbances, doubtful inheritance of, 195 Malignancy, 671

Malignancy, 671 Malignant infection, 459

tumors, 670

Malnutrition as predisposing cause of disease, 407

Maltase, 79

Maltose, 80

Mammary gland, accessory, 181 carcinoma of, pluricentric origin of, 693

cause of enlargement of, in pregnancy, 363

Mammary gland, chondroma of, 727 fat in cells of, 910 hypertrophy of, during pregnancy, 599 polymastia, 258 transplantation of, 634 tumors of, mixed, 662 Man, chromosomes in, 148 Mandelonitrile glucoside, 82 Marasmus, 394, 395 Margarin crystals, 982 Margination of leukocytes, 423 Mastigophora, 334 Maternal impressions, 193 Measles, organisms of, 995 Medusa, experiments on eggs of, 138 Megastoma, 336 Meiosis, 150 Melanemia, 828, 970 Melanotic pigmentation, 969 granules, chromidial nature in protozoæ, 48 properties of, 969 Melanoma, 678, 825 latency of, 687 Melanuria, 828, 970 Melasma, 970 Membranes, foetal, infection through, 207 Mendel's principles, 168 governing multiple pairs of features, 172 limitations of, 170, 187 Menidia, 148 Meningocele, 266 Meningomyelocele, 266 Menstruation, internal secretion of ovaries and, 362 Mental capacity, 178 Mercurial poisoning, effects of, upon offspring, 198, 219 Mercury, action of, on liver, 311 on skin, 312 aseptic suppuration by, 431 salts of, elimination of, 310 salivation of, 309 Mesodidymus, 240, 244 Mesothelioma, 812 adrenal, 677 mode of origin, 693 Mesothelium, 700 Metabolism, carbohydrate, 90 glycogenesis and ycogenesis and glycolysis through reversible enzyme action, 83 pancreas and muscle in, 366 chemical composition of protein molecule and, 66 of chlorophyll, 90 fatigue and, 401 of fats and lipoids, 92 fatty, nuclear activities in, 51 growth and, 75 impaired, as cause of disease, 374 nuclear activities in relationship to,

44

Metabolism, nucleoli and their relation- "Mother cells," metaplasia and, 639 ship to, **47** part played by water in, 85 proteid, impairment of, 375 of starch, 90 Metachondria. See Chromidia. Metals, effect of, upon growth, 89 Metaphase, mitotic, 115 Metaplasia, 639 of bone, 730 of cartilage into bone cells, 611 of nails, 619 Metaproteins, 57 Metastasis, 670, 676 calcareous, 926 in inflammation, 434 Metazoa as causes of disease, 343 reaction of, to injury, 418 Methemoglobinemia, 306 Methemoglobinuria, 957 Methylamin, 385 Miasmatic disease, 326 Micrococcus epidermidis albus, Welch's observations, 324 gonorrhaa. See Gonococcus. lanceolatus. See Pneumococcus. prodigiosus, metaplasia of, 646 Microcytase, 567 Micrognathia, 273 Micromelia, 224 Micronucleus, 34 Microphages, 566 Microphthalmos, 261 Migraine, 386 diathesis and transmission of, 211 Milk, bactericidal action of, 318 Mirabilis jalapa, hybrids of, 171 Mistletoe, 631 Mitosis in bacteria, 29 heterotype, 691 irregular, of cancer cells, 690 stages of, **115** Mixed tumors, 667 forms of, 808 (note) Modification versus variation, 159 Mole, hydatid, 664 placental, forms of, 663 Molecules, size of, 136 Molluseum contagiosum, 776 Momentum, physiological, 105 Monera, nuclear matter in, 29 Monospora, 419 Monsters, acardiac, 231 double, 234 parasitic, 254 Monstrosities, **226** causation of, effects of parental infection on, 197 familial degeneration and, 177 Morphine, 306 immunity toward, 504 Mortification, 987 Mosaic inheritance, 175 theory, 137 "Mother cells." 141

Mouse, adenocarcinoma of, 709 carcinoma of, 674 stroma of, 688 transplantation of, 687 Mendel's principles affecting, 173 Weismann's experiment of docking tail, 195 " Moving equilibrium" of living matter, 67 Mucigen, 889 Muein, 57 chromidial, development of, 50 Mucinogen, 889 Mucoid, 57 degeneration, 888 tissue, 641 Mucous membranes, grafting of, 636 regeneration of, 619 Mucus, a protective mechanism, 317 Mulatto, blended inheritance, 167 Müller's duct, cysts of, 854 Multiple primary tumors, 694 Multiplication of cell, 112 Mummification, 988 Musca vomitoria, fat formation and, 913 Muscarin, 385 action of, on heart, 307 Muscle, adductor, ossification of, 30 carbohydrate metabolism in. 91 connection between cells of, 35 deltoid, ossification of, 30 non-striated, involution of, 868 plain, regeneration of, 622 uterine, hypertrophy of, 593 poisons acting on, 305 reserve force of, 107 striated, atrophy of, brown, 874, 962 in fever, 470 a factor in glycolysis, 366 hypertrophy of, 593 kataplasia of, 878 regeneration of, 622 vacuolation of, 904 Mutation, 180 causation of, 183 due to interaction of parental biophores, 189 inheritance of, 209 neoplasia and, 842, 845 Mutilations, non-inheritance of, 105 Myelin, 915 bodies, in autolysis, their nature, 97 droplets, 95 Myelinic degeneration. See Lipoid degeneration. Myelocele, 266 Myelocystocele, 266 Myeloma, 732 giant-celled, 733 multiplex, 734 Myelomatoid, 735 Myeloplaxes, 733 Myiasis, 351 Myoglia fibrils, 746

1018

Myoma. See also Leiomyoma. calcification of, 745 necrosis of, 745 Myomalacia, 900 Myosarcoma, 747 Myosin, 56 Myositis ossificans, 730 Mysinogen and myosin, 994 Myxcedema, 353 hypertrophy in, 598 mucoid deposits in, 890 Myxo-enchondroma, 726 Myxofibroma, 712 Myxoma, 718 of peritoneum, 721 recurrent, 676 Myxosarcoma, 770 Myxosporidia, 339

N

INDEX

NAILS, regeneration of, 619 "Nebenkern," 50 Necrobiosis, 977 Necrosis, 977 calcification and, 927 caseation, 986 coagulation, 984, 986 fat, 982 fatty acid crystals in cells, 94 focal, 980 of eclampsia, 363 from arrested circulation, 977 from burns, 290 from inadequate nutrition, 978 gangrene, 988 gummatous, 986 of individual cells, 979 "Zenker's degeneration," 979 mortification, 987 of myomas, 745 necrobiosis and, 977 neuropathic, 978 results of, 987 Necturus, ova of, nucleoli of, 49 phagocytes of, 320 Negative catalysts, 78 phase, 554 Nemathelminthes, 343 Nematodes, 343 Neoplasia, 650 Neoplasm, 650 classification of, 667 Borst's, 697 histogenetic, 699 Waldeyer's (embryogenetic), 698 Neosporidia, 337 Nephrotoxin, 533 of snake venom, 550 Nerve cells, connections between, 35 fatigue and, 402

Nerves, injuries to. shock and, 581 olfactory, fibromatosis of, 715 optic, fibromatosis of, 716 peripheral, regeneration of, 625 regeneration of, 624 trophic, 403 Wallerian degeneration of, 870 Nervous affections, racial susceptibility to, 163 system, heat regulating centres in, 480 in inflammation, 448 poisons affecting, 304 relationship of, to heat pro duction, 488 Neural cysts, 861 groove. See Dorsal groove. Neurin, 385 Neurinoma, 758 Neuroblastoma, 752 Neurocytoma, 753 Neurofibroma, 716 Neuroglia fibrils, 756 regeneration of, 625 Neuroma, 753 adrenal, 809 amputation, 754 false, 754 Neurons, disue atrophy of, 110, 870 gemmules of, 305 influence of, upon development, 603 motor, nucleus and metabolism, 47 regeneration of, 604, 625 Neurotoxin of snake venom, 550 Nevus, pigmented, 826 telangiectatic, 816 Newl, See Triton. N'gana, 336 Nicotine, 304, 305, 306 Nissl bodies of nuclear origin, 50 Nitrites, poisonous, neutralization by cholesterin, 504 Nose, polyp of, 719 Nosema bombycis, 205 Nuclear membrane, 45 Nuclease, 377 in autolysis, 371

Nuclei of blastomatous growths, 690 Nucleic acid, 63 bactericidal properties of, 543 classification of, 64 (note) Nuclein, 63 Nucleoalbumins, 57 Nucleoalbumins, 57

melanin production and, 832

metabolism and, 47 Nucleoproteins, 57, 63

growth and, 76

Nucleus, chemistry of, 62 chromatin of. See Chromatin. in cloudy swelling, 884 constituents of, 30 cytoplasm and, 157 differentiation of, 100 degenerative changes in, 53

Nucleus, division of, arrangement of chro- Osteoma, 727 matin in bacteria, 29 dominance of, 36, 42 fat vacuoles in, 912 in fatty degeneration, 917 glycogen in, 922 incidence of, in living forms, 29 metabolism, and, 44 milk formation and, 910 oxidative powers of, 65 pancreatic zymogen and, 911 its relationship to cytoplasm, 38 size of, in relationship to function, 37 vacuoles in, 30 variation in shape of, 30 Nutrition, growth and, 590 Nutritional disturbances, 391

OAT-HAIR balls, 955 Obseity, 379 diathetic, 210 effects of thyroid secretion, 360 gouty diathesis and, 212 premature, 358 Obstructed elimination, as a cause of disease, 386 Occupational palsies, 872 Ochronosis, 972 Odontoma, 731 (Edema of chorionic villi, 222 determined by saline contents of body fluid, 90 (Enothera, mutation in, 183, 184 Oertel's theory of neoplasia, 844 (Esophagus, hypertrophy of, 594 Oleic acid, action of, as amboceptor, 541 hemolysis and, 306 Olein, formula of, 93 Omphalomesenteric duct, persistent, 267 Ontogeny, relationship to phylogeny, 133, Occytes, primary and secondary, 150 Oogenesis, 150 Oogonia, 150 Ophthalmia neonatorum, 225 Opsonic index, 554 Opsonins, 551 relation of, to complements and am-Organ of Rosenmüller, 274 Organism, defences of, 316 Oriental sore, 334 Ornithin, 59 Osmosis, 45 cellular, 89 hyperisotonic and hypisotonic solutions, 90 Ossification, distinction of, from calcifi- Pancreatin, effects of, on leukocytes, 307 cation, 924 Osteo-arthropathy, hypertrophy in, 598 Pandorina, 144 Osteoclasts, 610 Papilloma, **775**

Osteogensis imperfecta, 218, 647

Osteomatoid, 729 Osteomyelitis, cryptogenic, 325 Osteopathia trachealis, 730 Osteopsathyrosis, 224 Osteosarcoma, 771 metastases of, 684 Othematoma, 864 Otocephaly, 273 Ovary, adenoma of, mode of origin of, 694 action of x-rays upon, 294 degeneration in, elastoid, 898 extract, 352 internal secretion, 361 regeneration of, 622 removal of, in mammary cancer, 693 reserve force of, 108 transplantation of, 634 tumors of, 807 rgrowth. See Hypertrophy. Overgrowth. Overstrain, 396 Ovoglobulin, 56 Ovula Nabothi, 858 Ovum, blighted, 215 infection as a cause of, 219 conveyance of infection through, 205 maturation of, 150 polarity of, 138 separation of, blastomeres in, 217 Oxalate calculi, 943 Oxalic acid, poisoning by, 930 Oxidase, 377 melanin and, 971 Oxybutyric acid, 380 metabolism of, 907 Oxygen, deficiency and excess, as causes of disease, 391 Oxytricha, 43 Oxyuris vermicularis, 346, 351

p

PACHYDERMA, 902 Pain, causation of, 448 as a cause of shock, 581 of syncope, 580 Palinurus, heteromorphosis in, 603 Palmitin, formula of, 93 Pancreas, calculi of, 936 cells, origin of, secretory granules, 50 congenital cysts, 857 effects of extirpation of, 365 fat necrosis and, 982 soaps in, 94 fatty infiltration of, 609 regeneration of, absence of, 622 reserve force of, 107 resorption of juice, 3 self-digestion of, 984 reversible action of, 82 of bladder, 781

1020

Papilloma, non-blastomatous, 775 true, 781 Paradidymis, cysts of, 854 Paradiphtherial lesions, 208 Paralyses, occupational, 398, 400 Paramacium, parthenogenetic multiplica-tion of, 145 Paranuclear body, 50 Paranuclein, 82 Paraoöphoron, 274 Para-oxyphenylacetic acid, 384 Para-oxyphenylpropionic acid, 384 Parapedesis, 966 Paraplasm, 33 Parasites in cancer, 838 classification of, 280 Parasitic monsters, 254 Parasyphilis, 197 Parasyphilitic lesions, 205, 207 Paratuberculous lesions, 208 Paraxanthin, 376 Paresis, traumatic, inheritance of, 199 Paroophoron, cysts of, 853 Parotid, tumors of, mixed, 662 Paroxysmal hemoglobinuria, 958 Parry's disease, 355 Parthenogenesis, accessory chromosome and, 154 in infusoria, 145 natural and experimental, 129 Particulate inheritance, 167 Parturition, disease acquired during, " Passage," 495 of bacteria, 328 Passive immunity, 503, 510 Pathology, cellular, 22 definition of, 17, 25, internal and external, 25 special, 25 systemic, 26 Peas, cumulative inheritance in, 180 hybridization of, 168 Pearls, epithelial, 798 Pebrine, 205 Pellagra, 293 Penis, development of, 275 double, 244 Pentose in nucleic acids, 64 Peptones, 57 Perichondrium, transplantation of, 637 Periosteum, regeneration of, 610 transplantation of, 637 Peripheral nerves. See Nerves, peripheral. Perithelioma, 824 Peritoneum, lipoma of, 721 myxoma of, 720 Peritonitis by infection through Fallopian tube, 316 obstructive, 324 perforative, 317 universal and diffuse, 433

Pernicious anemia, abnormal gastro-intestinal fermentations and, 386 due to dibothriocephalus, 348 as a subinfection, 463 Peyer's patches, normal pressure of bacteria in, 319 Pfeiffer's reaction, 543 Phagocytosis, 565 on mucous surfaces, 318 part played by opsonins, 551 in protozoa, 416 Phenol, 384 Phenylalanin, 59, 379 Phlebolith, 936 ossification of, 731 Phloridzin, glycosuria, 367 Phosphate calculi, 944 Phosphatides, 57, 58, 92, 96 in autolysis, 371 classification of, 92 distribution of, 96 lipoid degeneration and, 916 relationship of, to cholin, 385 Wassermann reaction and, 549 Phosphoproteins, 57 Phosphorus, action of, on liver, 311 as cause of overgrowth, 597 fatty degeneration caused by, 913, 917 as a nuclear constituent, 63 poisoning, autolysis in, 372 cloudy swelling in, 883 Phrenosin, 93, 96 Phylloxera, 154 Phytonucleic acids, 65 (note) Phytosterins, 92, 95 Phytotoxins, 496 immunization against, 505 Picrotoxin, 304 Pigeon, cross-breeding of, 176 Pigmentation, abnormal, of urine, 379 endogenous, 956 of hemoglobin and its derivatives, 956 hematogenous (urobilin), 967 hematoporphyrinuria, 962 hemochromatosis, 962 hemoglobinuria, 957 infective, 959 paroxysmal, 958 icterus or jaundice, 963 modifications of, 960 postmortem imbibition, 957 pseudomelanosis, 962 lipochromes, 972 melanotic, 969 ochronosis, 972 exogenous, 973 and pigmentary dangers, 956 Pigmentophages, 972 Pilocarpine, action of, 312 Pin worm. See Oxyuris. "Piqure" experiment, 368

Piroplasma, 335 Piscicola, nucleoli in, 31 ova of, chromidial discharge, 49 Pituitary body, colloid matter of, 891 internal secretion of, 357 vicarious activity of, 108 hypertrophy of, 597 Placenta, disease of, effects upon foctus, hemorrhages of, causation of, 222 infection through, 207, 327 influence of disease of, upon fortus, 221 moles of, 663 transmission of metallic and other poison through, 220 Placental mole, 663 Plague, racial susceptibility to, 162 Planaria, regeneration in, 603 Plasmidiophora brassica, 840 Plasmodium, 31, 417, 418 Plasmolysis, 565, 567 Plasmosomes. See Chromidia. Pleuropneumonia, contagious, of cattle, Plumbism. See Lead poisoning. Pneumococcus, tissue predisposition to, Pneumonokonioses, 973 Poisons, 298 double monsters and, 234 dichotomy, 240 hypogenesis, 260 Polarity of ovum, 138 organisms of, 996 Pollen as cause of hay fever, 410 Polyblasts, 605 Polycyanic acids, 64 (note) Polydactylism, 64 a dominant condition, 174 inheritance of, 209, 214 Polydactyly, 258 Polymastia, 258 Polydon spathula, pancreatic cells of, 911 (note) Polypeptids, 57, 59 Potash, salts of, 307 Potassium soaps, 94 monophosphate, fatigue and, 401 Precipitins, **524** extent of specificity of, 525 nature of precipitate, 526 Precipitoids, 527 Precocious development, 358 Predisposition, 404

Predisposition, forms of, 404 of individual tissues, 408 to infectious disease, inheritance of, racial, 162 Preformation, 131 Pregnancy, influence of, upon mammary glands, 363 Preparator, 566 Pressure, atmospheric, 285 effects of, in developing ovum and embryo, 217 Prezymogens, 50 Prodromal, 455, 457 Professional palsies. See Occupational palsies. Progressive tissue changes, 587 Prolamins, 56 Prophase, mitotic, 115 Prostate, adenomatosis of, 789 carcinoma of, metastases in, 682 Protagon, 96 Protamins, 56 Proteans, 57 Proteases, 371 Proteidogenous matter, 54 essential to life, 77 Proteins, alcohol soluble, 56 classification of, 55 conjugated, 56 constitution of, 55 defective intake of, 393 derived, 57 general survey of, 54 molecular weight of, 55 non-specific immunity in, 501 simple, 56 synthesis through reversible enzyme action, 82 Protozoa, asexual multiplication of, 339 as causes of disease, 331 reaction to injury in, 416 regeneration in, 601 Prozymogens. See Prezymogens. Psammoma, 823 Pseudohermaphroditismus, 279 Pseudohypertrophy, 592 Pseudoleukemia. See Lymphoma. Pseudomelanosis, 962 Pseudomucin, 889 Ptomaines, 384 Ptyalin, 70, 91 Puncture, 283 Purgatives, mode of action of, 308, 309 Purin bases, classification of, 64 toxic effects of, 377 Pus, 431 Pygopagus, 247, 253.

Pyknosis, 53 Pylephlebitis, 434 Pyococcus, infection through skin and, 327 aureus, inflammation set up by, 437 opsonins and treatment, 555 Pyogenic membrane, 432 Pyrexia, 465, **478** causation of, 483 from heatstroke, 289 from increased external temperature, 486 neurogenic, 486 difference from infective, 487 Pyretogenic stage, 456 Pyrimidin bases, 64 Pyrocatechin, 306 Pyrogallic acid, 306

Q

QUADRIURATES, 378 Quinine, 304, 306 action of, on vessels, 308

R

RABBIT, bacteria in lymphadenoid tissue of normal, 319 coecidiosis of, 777 embryos of, transplantation of, 633 Mendel's principles affecting, 172 Rabies, incubation period of, 458 transmission of, acquired immunity against, 194 Rachischisis, 263, 265 Rachitis. See Rickets Racial inheritance, 162 Radium rays, 293 "Rankenneurom," 759 Ranula, 858 pancreatica, 858, 936 Rat, grafting, 631 (note) reaction of, to anthrax, 405 Receptor, 517 multiplicity of, 536 orders of, 571 Recessive characters, 168 Recklinghausen's disease, 759 Recurrent fever, 467 Red corpuscles. See Erythrocyte "Red degeneration," 746 Reduction of chromosomes, 146, 151 influence of, on zygote, 190 Referred inflammation, 449 Regeneration, 418, 601 capacity of different orders of cells, as distinct from hypertrophy, 596 Relapse, 456 Relapsing fever, 467 Remittent fever, 466 Rennet, 508 Rennin, 71, 577 Repair, 447

Replantation, 631 Reserve force, 106 Resistance period, 501 Resolution of abcess, 432 Resorption of excretions, 387 Respiratory system in febrile state, 473 Retrograde metastasis, 680 Reversibility of enzyme action, 79 Reversion, 164 diathetic, 177 general and systemic, 211 Reversionary atrophy, 641 degeneration, 632 inheritance, 176 metamorphosis, 878. See also Kataplasia Rhabdomyona, 750 Rhabdomyosarcoma, 752, 773 Rheumatism, acute, etiology of, 461 (note) gouty diathesis and, 212 Rheumatoid conditions, 386 Rhinolith, 935 Rhubarb, elimination of, by liver, 310 mode of action of, 309 Ribbert's theory of neoplasia, 837 Ribose, 65 (note) Ribs, accessory, 257 due to mutation, 182 deficiency of, due to mutation, 182 Ricin, 306 adaptation to, 122 immunization against, 505 Rickets, 394 fcetal, 224 giantism and, 228 "Riesenwuchs," 714 Rigor, 469 mortis, 993 immediate, 397 Robin, 507 Rodent ulcer, 790, 798 Römer's experiment, 516 Röntgen rays, 293 effects of, on frog's sperm, 877 Round-worm. See Ascaris. Roux's experiment, 137

S

SALAMAN DER poison, 550
 regeneration in, 601
 Salleylie acid, action of, on skin, 312
 Salivar, a protective mechanism, 317
 Salivary concrements, 935
 glands, cells of, metabolic activities
 of nuclei of, 47
 regeneration of, 620
 secretions and, 365
 serous, formation of secretory
 granules, 50
 Salivation, 309, 312
 Salponin, hemolysis and, 306

NDEX

Saprophytes, 314 Sarcolactic acid, fatigue and, 401, Sarcoma, 670, 763 definition of, 707 giant-celled. See Myeloma, giantintermediate forms, 770 melanotic. See Melanoma. pure forms of, 766 Sarcopsylla penetrans, 351 Sarcosporidia, 339 Saturnine poisoning. See Lead poisoning. Scarlet fever, Mallory's bodies in, 332, 340 Wassermann reaction in, 549 "Schlummerzellen," 110, 608 Scleroproteins, 56 Scombrone, 56 Scorbutus. See Scurvy Scorpion, antivenin, 550 Scrotum, hydrocele of, 860 Scurvy, 393 See Echinus. Seat-worm. See Oxyuris Secretin, 364. See also Hormone Semipermeable membranes, 89 of cells, part played by phosphatides, 96 Senna, action of, 309 Sequelæ, 456 Sequestrum, 988 Serosæ, fibrinous inflammation of, 430 hemorrhagic, inflammation of, 435 mesothelioma, 812 suppurative, inflammation of, 433 Serous atrophy, 904 globulin, 56 soap compounds as complements, 540 Sex, accessory chromosomes and, 153 differentiation of, 278 as example of particulate inheritance, influence of parental nutrition upon, 199 (note) mosaic inheritance and, 175 predisposition according to, 406 Sex-limited inheritance, 179 Shamrock, mutation in, 181 Sheep, Ancona breed, mutation and descent, 213 Shock, 580 after burns, 291 distinction of, from collapse, 585 Side-chain theory, 516 of immunity, **516**, **570** Siderosis, pulmonary, 973 Silicosis, 973 Silk-worms, Mendel's principles affecting, Siphonocladus, 42 Siren, 261

Situs inversus, 277 Skatol, 384 action of, on liver, 311 Skin, action of poison on, 311 atrophy of, senile, 874 bronzing of, 358 cheloid of, 716 in febrile state, 477 grafting, 635 heterotopia of, 640 horns of, 777 ichthyosis of, 182 leiomyoma of, 750 regeneration of, 618 Sleeping sickness, 336 Smallpox, Guarnieri's bodies in, 332, 340 incubation period of, 458 inoculation against, 492 ultramicroscopic organisms of, 995 vaccination in, 493 vesiculation in, 903 Snail, Mendel's principles affecting, 173 Snake venom, 511 neurotoxin, combination of, with nerve matter, 514 Soaps, 92, 94 action of, as complement, 540 Wassermann reaction and, 549 Sodium soaps, 94 sulphindigotate, elimination of, 310 Solution, crystalloid and colloid, 87 hyperisotonic and hypisotonic, 90 solute and solvent, 87 water-solid and solid-water phases, 88 Sow, mamma of, 181 Spallangani's law. 694 Species and varieties, what should determine, 172 Spermatocytes, primary and secondary, Spermatogenesis, 148 Spermatogonia, 148 Spermatoxin, 533 Spermatozoa, effects of x-rays on, 878 evolution of, 145 human, incapable of conveying infection in fertilization, 206 maturation of, 148 Spheroliths, 954 Sphingomyelin, 93, 96 Spider antivenin, 550 parasitic, 351 Spina bifida, 265 causation of, 217 ¹¹ Spinal shock,¹¹ 584 Spirillum cholera, aggressins of, 557 endotoxins of, 543 Pfeiffer's reaction in, 543 Metchnikori, immune serum of, 45 "Nasik," 300 Spirochata pallida, 207 Spirochete associated with malignant growth, 840 Spirogyra, amitosis and mitosis in, 114

Spleen, accessory, 640 degeneration in, elastoid, 898 lymphomatoid conditions of, 743 origin of myelocytes in, 614 regeneration of, 622 sago, 892 Splenic anemia, 743 Splenomegaly, 743 Spontaneous variation, 180. See Mutation. Spores of bacteria, 313 Sporozoa, 336 cancer and, 341, 794 Squamous-celled cancer, 797 Staphylococcus pyogenes aureus. Pvococcus aureus. Staphyloschisis, 271 Starch, absorption of, from lungs, 975 metabolism in vegetable cell, 90 Star-fish (Asterias), connections between blastomeres, 36 between larval mesoderm cells, 36 Starvation, 395, 872 acidosis in, 381 Status lymphaticus, 360 Steapsin, 71. See also Lipase. fat necrosis and, 984 Stearin, formula of, 93 Steatopygia, 210, 723 Stentor, nucleus of, 30 and growth of, 43 Stercobilin, 968 (note) Sterility from modification of germ cells, Sternopagus, 257 Stillbirths from alcohol, 221 from cardiac disease, 219 from imperfection of germ cells, 208 a result of familial degeneration, 177 Stomach hypertrophy of, 594 Streptococcus, cryptogenic infection by, TABACOSIS, 974 infection through skin and, 327 normal presence of, in pharynx, 317 pyogenes, opsonins and, 556 Stroma of neoplasms, source of, 687 of tumors, origin of, 787, 790 Strongylocentrotus, 139 Strongyloides intestinalis, 346, 347, 348 stercoralis. See Strongyloides intes-Struma suprarenalis, 809 thyroidea ovarii, 686 (note) Strychnine, 304 immunity, 504 Sturine, 56, 58 Subinfection, 462 Submaxillary gland, tumor of, mixed, 662 Teratoblastoma, 660, 750 Substance sensibilitrice. See Ambocep-

Suctoria, 342

Sugars, fermentation of, by bacteria, 119 Sulphocyanates, action of, 306 Sulphuretted hydrogen, 306 Sunstroke, 290 Suppuration, 430 aseptic, 431 Surface action, 78 membrane of colloids, 88 "Surmenage," 396 Susceptibility. See Predisposition. increased in descendants by abrin poisoning, 194 in offspring by parental infection, 197 to infection, 454 Symmelia, 261 Sympathetic nervous system, abdominal. and Addison's disease, 358 Sympus, 261 Synapse theory of fatigue, 400 Synapsis, 141 its significance, 153 Syncephalus, 248 Syncope, 580 Syncytioma. See Che Syncytium, 31 (note) See Chorio-epithelioma. Synechia, 430 Synotia, 260, 272, 273 Syntoxoids, 515 Syphilis, congenital, not inherited, 207 effects of, upon foctus, 218 upon offspring, 197 upon placenta, 218, 223 Wassermann reaction in, 548 "Syphilitic antigens," 549 Syringomyelocele, 266

т

Tabes dorsalis, 877 Twnia echinococcus, 345, 348, 351 cystic stage of, 865 Talipes equinovarus from pressure in Tape-worms, 343 Tattooing, 973 Taurin, 378 Teeth grafting, 636 Telophase, mitotic, 115 Temperature, adaption to changes in, amphibolous, 466 as a cause of disease, 285 of body, centres for regulation of, 480 of inflamed areas, 450 influence of, upon Lepidoptera, 135 vital manifestations and, 69 Teratogenic termination period, 235 Teratogenous blastoma, 663

Teratoma, 255, 650, 652 congenital sacral, 239, 654, 656 difference in properties from blastoma, 837 filial, 257 ovarian, 256, 657 testicular, 658 twin, 257 Testes, action of x-rays upon, 294 chondroma of, 726 compensatory hyperplasia of, 595 extract, 352 internal secretion of, 361 interstitial cells of, 362 regeneration of, 622 tumors of, 807 Tetanotoxin, 523 junction of, with nerve matter, 514 Tetanus antitoxin, 510 incubation period of, 458 toxin, properties of, 514, 515 Tetany, 356 Tetrahydronapthalamin, 486 Texas fever, 335 conveyance of, through ovum, 206 Thalassicola pelagica, 46 Thebaine, 304 Theobald Smith's phenomenon, 559 Thermogenesis, 478. See also Pyrexia. Thomsen's disease, 877 Thoracopagus, 246 Thread-worm. See Trichocephalus. Thrombokinase, 374 Thrombophlebitis, 434 Thrombosis, 977 Thrombus, organization of, 428 Thymin, 64 Thymonucleic acids, 65 (note) Thymus, fat cells of, 608 involution of, fatty degeneration and, 909 Thyroid, adenomatosis of, 789 colloid matter of, 890 compensatory hyperplasia of, 595 effects of removal of, upon offspring, internal secretion, 352 regeneration of, 621 transplantation of, 633 tumors of, 685 Thyrolingual cysts, 852 Ticks, conveyance of Texas fever through, 206 Tissue differentiation, 129 extracts, pyrexia and, 485 hylic, 701 of predilection, 682, proliferation of, in inflammation, 444 resistance of to freezing, 286 susceptibility of. 408 Toad venom, action of, 307 poison, 550 Toluylenediamin, 306, 967 Tongue, "hairy," 902

Tonsillar concretions, 935 Tonsils, protective action of, 319 Tophi, 378 Totipotent cells, 653, 655 Toxins, 315 action of, on liver, 311 through nervous system, 468 on vessels,308 adaptation to, 121 antitoxins and, 510, 520 nature of union of, 511 production of, 125 their reciprocal nature, 562 bacterial, 314 comparison of, with enzymes, 576 distinction of, from ordinary poisons, intracellular, 315 mode of action of, 519 nature of, 512 of union of, with antitoxin, 520 pyrexia and, 483 relationship of, to enzymes, 513, 519, Toxoids, 515 Toxones, 515 Toxophone, 515, 517 Trachea metaplastic bone in, 645 ossification of, 730 Transplantation, 631 of tumor cells, 674, 680 Transposition of viscera, 277 Trauma as cause of infection, 324 Trematodes, 343 regeneration in, 603 Tricephalus, 243 Trichina, 346 spiralis, 865 Trichinosis, 350, 351 Trichocephalus, 346, 347 Trichomonas, 336 Trichotoxin, 533 Trigonocephaly, 260 Trimethylamin, 385 Triplets, 235 Triton, experiments on embryos of, 140 "Tropfische Entmischung," 885 Trophic nerves, 403 Trypanosoma, 334 brucei, 334 evansii, 334 gambiense, 334 figure of 331 ingeus, 334 noctua, 206 theileri, 334 Trypanosomes, adaptation of, 120 transmitted immunity to arsenic, 195 Trypanroth, 335 Trypsin, 71 Tryptophane, 59 Tsetse fly, 336 Tubercle, absorption of, 442 development of, 440 Tuberculin, action of, on vessels, 308 reaction, anaphylactic nature of, 560

Tuberculosis of cervical glands, 325 effects of, upon offspring, 197 giant cell in, 444 mesenteric glands in, 325 of placenta, 223 predisposition toward, inheritance of, pretuberculous stage of, 458 racial susceptibility to, 162 ulcerative, 442 vaccination and opsonins in, 555 Tubularia, geotropism in, 602 Tulip, mutation in, 180 Tumors, action of x-rays upon, 294 benign, 669 cells, latency of, 686 connective tissue, 710 due to mutation, 182 glycogen in, 922 malignant, 670 autolysis in, 373 mixed, 667 forms of, 808 (note) multiple, 796 primary, 694 ordinary. See Blastoma. stroma of, origin of, 787, 790 in tumore, 657, 837 Turpentine, aseptic suppuration by, 431 VACCINATION, Jennerian, 493 Twins, 228 Typhoid fever, course of, 455 rose spots in, 312 Tylosis, 224 Tyrosin, 59, 379 produced in autolysis, 371, 372 Tyrotoxicon, 385

U

ULCERATION, exogenous, 433 inflammatory, 432 tuberculous, 442 Ultramicroscopic organisms, 995 Ultraviolet light, treatment of malignant growths by, 693 rays, 292 Umbilical cord, disturbances of, 224, 225 effects of compression by, 217 Uncinaria. See Ankylostoma. Uniceptor, 572 Urachal cysts, 852 Uracil, 64 Uracyl, 376 Uramia, 388 Uranoschisis, 271 Urates, amorphous, 954 in gout, 375 Uratic deposits in tissues, 953 inspissation of infancy, 941 Urea from alloxuric bases, 377 non-toxic properties of, 390 Ureters, ligature of, effects of, 387 Urethra, development of, 275 Uric acid, 64, 376 calculi, 941

INDEX

Uric acid, causes leading to increased production of, 377 infarcts, 941 Urinary calculi, 938 Urine, bactericidal action of, 318 in fever, 474 resorption of, 388 Urobilin pigmentation, 967 Urobilinuria, 958 Urogenital ducts, tumors of, 806 Uroleucic acid, 379 Urostealith, 945 Urtica, hybrids of, 169 Urticaria, 312 Use acquirements, non-inheritance of, 194 Uterus, arteries of, degeneration of, elastoid, 896 carcinoma of, mestastases in, 682 cervix of, tumors of, mixed, 662 duplex, 278 mucosa of, regeneration of, 20 muscle of, involution of, 868, 909 pregnant, hypertrophy of, 593

τ

Pasteurian, 494 Wright's, against various pathogenic organisms, 554 Vaccinia, Guarineri's bodies and, 340 ultramicroscopic organisms of, 995 Vacuolar degeneration, 904 Vacuoles in nuclei of fat cells, 51 Vagina, tumors of, mixed, 662 Variability, disease and, 17 law of chance and, 18 Variation, adaptation and, 116, 122 biophores and, 157 correlated, 182 as distinct from modification, 159 due to interaction of parental biophores, 188 spontaneous, 180 Varieties, distinction from species, 172 Variola. See Smallpox. Varix, 815, 850 Vegetations, 438 Venoms, animal, 550 Ventricle, fourth, hydrocele of, 861 Veratrine, 304, 306 Veronica, hybrids of, 171 mutation in, 183 Vertebræ, accessory, 257 supernumerary, due to mutation, 182 Vesication from burns, 290 Vessels, blood, regeneration of, 614 of invertebrates, inflammation and, 418 part played by, in inflammation, 415 poisons acting on, 308 secondary role of, in inflammation, 444 transplantation of, 637

Vessels of vertebrates, inflammation and, 419

Vibratory motion, 298

Vibrio. See Spirillum.

Vicarious activity, 108 Virility, premature, 358

Virulence, 327

of bacteria producing endotoxins, 557 exaltation of, 495 bacterial antibodies and, 557 by passage, 328

through obstruction of passages, 323

by transference in vitro, 328 Vitellin, 57

Vitello-intestinal cysts, 852

Vitreous degeneration. See Elastoid degeneration. X-RAYS, 293 dermatitis

Vomiting, 309 collapse and, 581 pernicious, of pregnancy, 381

W

 "WALLBILDUNG," 432
 Walterian degeneration, 626, 870
 Warts, 775
 Wassermann reaction, 548
 Water, as a constituent of cell, 85 deficiency and excess of, as causes of disease, 392
 hemolysis and, 306
 inoculation and, 85
 Water scorption, nucleus of, 30

Waxy degeneration. See also Amyloid degeneration.

Weigert's hypothesis, 589 law of inertia, 518 Weil's disease, 967 Weismann's theory, 134 Wharton's jelly, 720 Whart, serviceable hybrids of, 173 Whip-torm, See Trichoeephaus. Widal reaction, 528 Wolfian body, 274 eysts of, 853 duct, 274 eysts of, 854 Worms, intestinal, 343 Wright's phenomenon, 552

INDEX

х

X-RAYS, 293 dermatitis, 846 treatment of malignant growths by, 693 Xanthelasma. See Xanthoma. Xanthin, 64, 376 calculi, 945 Xanthoma, **723**, 972 Xanthoma, **723**, 972 Xaroberna pigmentosum, 293, 830 Xiphopagus, 246 Xiphopagus, 246 Xiphopagus, 246

Y

YEAST, Buchner's experiment, 315 Yellow fever, organism of, 995 racial succeptibility to, 162 Yolk granules, origin of, 49 nucleus, 50

Z

ZEIN, 56 Zenker's degeneration, 979 Zymogen granules, 50