Technical and Bibliographic Notes / Notes techniques et bibliographiques


Canadiana.org has attempted to obtain the best copy available for scanning. Features of this copy which may be bibliographically unique, which may alter any of the images in the reproduction, or which may significantly change the usual method of scanning are checked below.

Canadiana.org a numérisé le meilleur exemplaire qu'il lui a été possible de se procurer. Les détails de cet exemplaire qui sont peut-être uniques du point de vue bibliographique, qui peuvent modifier une image reproduite, ou qui peuvent exiger une modification dans la méthode normale de numérisation sont indiqués ci-dessous.

Coloured covers / Couverture de couleur		Coloured pages / Pages de couleur
Covers damaged / Couverture endommagée		Pages damaged / Pages endommagées
Covers restored and/or laminated / Couverture restaurée et/ou pelliculée		Pages restored and/or laminated / Pages restaurées et/ou pelliculées
Cover title missing / Le titre de couverture manque	\square	Pages discoloured, stained or foxed/ Pages décolorées, tachetées ou piquées
Coloured maps / Cartes géographiques en couleur		Pages detached / Pages détachées
	\checkmark	Showthrough / Transparence
Coloured ink (i.e. other than blue or black) / Encre de couleur (i.e. autre que bleue ou noire)	\checkmark	Quality of print varies / Qualité inégale de l'impression
Coloured plates and/or illustrations / Planches et/ou illustrations en couleur		Includes supplementary materials /
Bound with other material / Relié avec d'autres documents		Comprend du matériel supplémentaire
Only edition available / Seule édition disponible		Blank leaves added during restorations may appear within the text. Whenever possible, these have been omitted from scanning / II se peut que
Tight binding may cause shadows or distortion along interior margin / La reliure serrée peut causer de l'ombre ou de la distorsion le long de la marge intérieure.		certaines pages blanches ajoutées lors d'une restauration apparaissent dans le texte, mais, lorsque cela était possible, ces pages n'ont pas été numérisées.

Additional comments / Commentaires supplémentaires:

.



. · .

INDEX.

I

Å.

Acmo Tomat	page	
Acme Tomato, Advice to farmers on fault mounting	184	
	82	
Agawam Grape,	183	
	20	
A Plea for Flowers, A Plea for our Small Fruits	37	
Apples in Minnesota,	190	
Apples, how to Pack,	14	
	185	
Ben Davis,		
	178	
Bienheim Orange, 17 Boston Russet,		
Baldwin,	$179 \\ 78, 180$	
Cayuga Red Streak, 17	8, 180	
Uranberry Pipuin.	178	
Unenango Strawberry 15	8, 180	ł
Colvert,	178	
Canada Baldwin,	50	1
Decarie,	51	i
Duchess of Oldenburg	14, 164	i
Dutch Mignonne	180	i
Eigin Beauty.	14	ì
Early Harvest	140	1
Lsopus Spitzenberg	178	
rameuse, 14 H	79, 180	
raneuse Sucree.	49	1
rall Pippin, 1	78, 180	
rall Janetting,	180	ì
Gravenstein, [52.]	78, 180	
Green Newtown Pippin	166	Ì
Golden Russet,	-178	
Gloria Mundi,	178	
Hastings Apple,	33	
Keswie Coullin,	180	1
King of Tompkins, 186, 1		
McIntosh Red,	41	
Mountain Beet,	50	
Maiden's Blush,	180	
Mielon, Northurn Sam	178	
Northern Spy,65, 1 Newtown Spitzenburg	79, 180	
- Pomme Grise, 45, 1	179	
* VALUE de For		
" CON 8 Pleasant	5(
	17:	
	-5(1-1	
	14: 170 IG	۴. ۱
Rambo, Rostury Pass	1787, 181 171	י ר
TO A DULLY IN ISEAF	179 179	
LUNSCOLL LUNIN	179, 180	
Serioner ispitzenner	1	ò
St. Lawrence,14,	$170 \ 1.0$	o n
······································	10, 10	U

	page
APPLES-Snow Apple,14, 179), 180 -
SwayziePommeGrise45,17	
Strawberry of Montreal	51
Seek-no-Further,	180
Swaar,	179
Sherwoods Favorite.	178
Tetofsky,	14
Talman Sweet,	179
Twenty Ounce,	178
Utter's Red,	14
Vandevere,	179
Wagner, 179	9, 180
Wealthy,	14
White Astracan,	14
YellowNewtownPippin17	79,180
Yellow Bellflower, 17	9, 180
Apple Trees, new hardy	<u> </u>
Apple Market, glut in	172
Apple Tree Pruner.	-170
Apple Trees, diseases of	22
in Dundas Co.,	41
" Grafting, bearing	71
Asparagus,	31
Conover's Colossal	40
" Special Manures for	66
" How to raise Colossal	71
Attacus Cecropia,	8, 28
Autumn Meeting of F. G. A.	152

B.

180		
180	B .	
166	Baldwin Apple,	117
178		183
178	Bartlett Pear, 181,	182
33 :	Ben Davis Apple,	178
180	Begonias,	167
180	Belle Incrative Pear,	181
41	Beurre d'Anjou Pear, 181.	182
50	Beurre Gris de Hiver Pear,	181
180	Beurre Clairgeau Pear,	182
178	Bearre Diel Pear.	182
180	Beurre Bose Pear,	69
179	Belle Strawberry,	20
180	Beets,	26
50	" Blood Turnip,	$\tilde{26}$
179	" Early Bassano,	26
50	" Egyptian,	$\tilde{26}$
140	Beans, Snap,	63
, 180	" E Rachel,	63
179	" E. Mohawk,	63
179		64
. 180	"Wax or Butter,	64
13	• Windsor,	64
, 150	" White Marrowfat,	- 64

INDEX TO VOL. I.

Bearing year, changing.	$\mathbf{p}\mathbf{a}_i$
Bearing Apple Trees, grafting.	(
Blackberry,	- 7
Blackberry,	
Ularkson's Early	Ċ
VOI, Wilder	č
UTYStal White	
" Cumberland,	
" Dolomona	Į
" Delaware,	ę
" Dorchester,	ç
Dr. Warder	0
LOOSAC Thornless	ò
Holcomb.	9
Kittatinny,	
" New Rochelle,	9
" Sipolois	0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0
" Sinclair,	9
onvuer.	c c
	a a
	170 10
Bliss' American Wonder Pea,	178, 18
	11
Bradshaw Plum, Black Current mill	17
Black Cumula	18
	3
Burnet Grape,	J

C.

Calchis Maria	Dorchester Plash	78
Calchis Maria, Cabbage Butterfly, 29 Canadian Phoenbath, 36	Dorchester Blackberry,	90
Canadian Phosphates,	Doctor Warder Blackberry,	90
Canada Baldwin 43	Downing Gooseberry,	165
Canada Baldwin, 43 Cayuga Red Streak Apple, 178, 180 Chinese Primroses, 178, 180	Duchess of Oldenburg Apple, 14	4, 164
Chinese Primer Apple, 178, 180	Dunuas County, Annie Troop in	4[
Chinese Primroses,	i Uutuu Mignonne Annie	*
Changing the Bearing Plum, 54, 94	Duchess (LAngenheime Pean 10)	1 1 9 9
Changing the Bearing Year, Chenango Strawberry 4 - 64	Duane's Purple Plum, 18	182
Chenango Strawberry Apple, 64 Cherry, the Old Kentish		102
Cherry, the Old Kentish	Ε.	
Clapp's Favorite Pear, 187 Clarkson's Faely Picture 17		
	Early Bassano Beet,	26
ansectos, one or		52
Attacus Cecropia e no		
Archinnus Data - d		63
		63
		140
		162
		150
" Grape-vine flor headle 22 and 75		182
Grape-vine flen beetle 62,92,111 Plum Curculio, 72		182
" Rose Share 72		26
" Rose Slug, 99		14
" Rose-Leaf Hopper, 99		178
" Rose Bug,	Exorista militaris,	84
Conover's Coloscol Astronomical 18	Experiments in Hybridizing,	29
Conover's Colossal Asparagus, 18 Corvius Americana 40	Experice in winter Guilding,	126
	Exper'ce in wintering Geraniums,	183
Lisparagus, Conover's	_	-+0
	Famouac A J F.	
Conotrachelus Nenuphar, 71 Colonel Wilder Plant, 72	Fameuse Apple, Fameuse Sucree	14
	Fameuse Sucree, Farmers, Advice on fruit grant	49
Colvert Apple,	Farmers, Advice on fruit growing,	82
Columbia Plum,	Fall and rise of Sap,	
102	Fall Pippin,	190
	110	, 180

•ge		
64	Coe's Golden Drop Plum,	page
71	Concord Grape, Crystal White Blackberry, Cranberry Pinnin America	182
90	Crystal White Blackhorny	183
91		
91	Creveling Grape, Criterion Tomato	178
91		
91	Crescent Seedling Strawberry,	184
91		
91		
91		
91		129
91		
91	Cultivation of Small Fruits,	121
91	or of anal Fruits,	109
91	D.	
91	-	
91	Danais Archippus.	1 4 1
80		141
19	Delaware Blackberry,	49
79		90
82	" crenata fl. pl.	94
30	De Tongres Pear,	94
2	· I/Claware (Franc	180
	Diseases of Apple Trees,	183
	Double Flowering Hawthorns,	22
	Dorchester Blackberry,	78
2 9 ;	Doctor Warder Blackberry,	90
36		90
43	Duchess of Oldenburg Apple,	165
50	Dundas County, Apple Trees in	14, 164
80	Dutch Mignonne Apple,	4[
23	Duchess d'Angonluma Desa	180
94	Duchess d'Angouleme Pear,	181, 182
64	Duane's Purple Plum,	182
30	Ε.	
87		
17	Early Bassano Beet,	
90		26
28		52
28	Early Mohawk Bean, Early Harvest Appl	63
41	Early Harvest Apple, Early Canada Peach	63
70		140
36	Early Conqueror Tomato,	162
37	Easter Beurre Pear,	150
75		182
11	Egyptian Beet, Elgin Beauty Apple	182
12		26
99	Esopus Spitzenberg Apple, Evergreeps Transplant	14
99	Evergreens, Transplanting of	178
99	Excrista militaris, Experiments in Hybridizing, Experice in wintering Com-	84
8	Experiments in Hybridizing	29
ю	Exper'ce in wintering Germanium	126

iv.

	page
Fall Janneting Apple,	130
rerunzers for Fruit Trees	6
Flowering Shrubs, how propagate,	94
" Chinese d'bl fl'g plum	54
" Deutzia Gracilis,	
" crenata fl. pl.	94
" Early Flowening	52
" Early Flowering,	
" Japan Quince,	53
" Plum-leaf Spirea,	53
oyringa,	94
"Weigela Rosea,	94
riemish Beauty Pear. 1	81, 182
r towering mawinorns	94
rorest lent Caternillar	75
roster reach.	21
rolest nose Strawberry	125
Fruits of Recent Introduction,	20
Fruit Prospects,	67
Fruit Shipped from Meaford,	
" Ome- Sum)	139
Fruit Gromer And	139
Fruit Growers Ass'n, of Quebec,	151
Fruits in the Ottawa Valley,	168
Fruit at the Provincial Exhibition,	177
Fruit Growing near Sarnia,	155
Francis H. Hora, in memoriam	79

G.

Consolla D. D.	. 1
Gansel's Bergamot Pear,	181
Garden, Poor Man's,	-74
Gen. Sherman Strawberry,	20
Geraniums, How to Winter, 160,	183
General Grant Tomato.	150
unout morceau Pear,	182
Gloria Mundi Apple,	178
Gladiolus,	174
" Varieties,	176
Glut in the Apple Market,	172
Glass Seedling Plum,	152
Goodale Pear,	181
Golden Russet Apple,	178
Gooseberry, The	131
Downing,	165
Grape, Agawam,	183
	183
"Barry,	183
" Concord,	-
" Creveling,	183
Delaware,	183
Iona,	183
martha.	183
moore's Early 09	188
I CIKIUS,	183
nebecca,	183
Kogers No. 44	183
Senasqua	183
Wilder.	109
whape vine Flea Beetle 69 09	111
Grane Vine Summer D	, 114

page 27, 71 71 ge]

 grafting over Old Orchards,

 50
 Grafting Bearing Apple Trees, ...

 94
 Gravenstein Apple,

 94
 Green Peas,

 94
 Green Newtown Pippin,

152119 166 Green Gage Plum, 182 94 $\overline{52}$

H.

94	Hastings Apple,	33
82	Haltica Chalybea,	92
94	Hathaway Tomato,	184
75	Hawthorns,	78
21	Hardy Apples, New	49
$\overline{25}$	Hardy Apples, New Herry,	20
20	Hervey Davis Strawberry,	158
67	Henrietta Raspberry, History of Wilson Strawberry,	57
139	Hora, F. H., In memoriam	79
139	Horticulture, Woman's work in.	84
151	Horticultural Gossip,	
168	Holcomb Blackberry,	90
177	Hoosac Thornless "	
155	Howell Pear	181
79	How to propogate some fl'g shrubs	94
	Horticultural Eccentricities,	127
	Hubbard's Imp Curl Leaf Tomato	150
	· ·	
101	Hybridization, Experiments in ••	126

I.

flag 1 (1), 1104 00 ((11001, **** 100,	183	
General Grant Tomato,	150	Imperial Gage Plum,
Glout Morceau Pear,	182	
Gloria Mundi Apple,	178	" Jared P. Kirkland, 25
Gladiolus,	174	
" Varieties,	176	Insects, one of our common, 8, 28
Glut in the Apple Market,	172	"Attacus Cecropia, 8, 28
Glass Seedling Plum,	182	
Goodala Boar		Charles Maria,
Goodale Pear,	181	Construction Menaphan, 72
Golden Russet Apple,	178	Danais Aremppus, 141
Gooseberry, The	131	" Exorista militaris, 29
Downing.	165	" Forest Tent Caterpillar, 75
Grape, Agawam,	183	" Haltica Chalybea,
Barry,	183	" Macrodactylus subspinosa 99
" Concord,	183	(f Norr found fails).
" Creveling	183	66 Nonnahar
"Creveling,		ff Orbign measures
Delaware,	183	29
rona,	183	" Parasitical, 29, 37
martha.	183	
moore's Early	188	
"Perkins, 22,	183	
"Rebecca,		
" Rogers No. 44,	183	100 T
4 Sonnooue	183	istenocerus putator,
" Senasqua,	-183	" Tent Caterpillar, 18
Wilder.	183	" Tettigonia Rosæ, 99
where wine Flea Beetle co oo	. 111	Iona Grape, 183
Grape Vine, Summer Pruning,	114	Iris, Japanese,

J,

Jared P. Kirkland, In memoriam,
Japan Quince,
Japanese Iris,
Jefferson Plum,
John Freed in memorias
Josephine de Malines Pear,

ĸ.

Kentish Cherry, The old	18
Keswic Codlin,	10
King of Tompkins Apple, 178, 180,	13
aring of Tompkins Apple, 178, 180	15
ALL MANUE VALUE I. HE MOMONIANS	
Kittatinny Blackberry,	-2
· · · · · · · · · · · · · · · · · · ·	- 9

L,

Lawrence Pear,		100
Late Spring Frosts	10-	130
Lettuce,	100,	
Lombard Pinn		157
Louise Bonne Pear,		-81
	181.	152

M.

Martha Grape,	
Maiden's Blush Apple,	
Manure for Asparagus,	
Macrodactylus subspinosa,	•
Market Club in handlings,	
Market, Glut in Apple	
McIntosh Red Apple,	
Meeting of F ()	
Meeting of F. G. Ass'n, Winter.	
" Summer.	
Mildow on the D	
Moore's Early Grape, Mountain Beet Apple,	22,
Montreal II	
Montreal Horticultural Society,	
Mount Vernon Pear,	
······································	

N.

New	found Fr	ienda		
Nen	/ found Fr latus venta	ionais,	• • • • • • •	
New	atus venta Rochelle	ricosus,		
V	Rochelle Raspberr	Blackb	erry	
11 G W	'Raspberr Varieties	¥		•••
.Ne₩	Varieties town Spit	of Con		• • •
New	town Snit	7051	senerry,	
	town Spit	zenourg	Apple.	

0.

Old Orchards, Grafting over

page	Ophion macrurum, Ottawa Valler, Em.	page
25	Ottawa Valley, Fruits in the	29
52, 94		168
-130	and ampped from	139
81	5	
47	P.	
181	Packing Apples	
101	Packing Apples,	185
	Parasitical Insects, Peach Growers, Warmin	29, 37
		15
187		15, 172
	- cach, the roster	21
180	Lariy Canada	
80, 186	"Crates,	162
25	"Ladder,	55
90		55
	Peach Plum	111
	Peach Plum, Pears, Bartlett,	182
	Pears, Bartlett,	1, 182
-180	" Beurre Bose	80, 181
5, 114	i vourre chalreean	30, 182
157	Deurre Gris de Hivan	181
81	Deurre d'Anion 10	1, 182
1, 152	. Denrie Diel.	182
-, 102	1 1010 111091100	181
	Dustum,	182
	Ciabb S Favorito	
183	De l'oncres	17
180		180
66	"Easter Beurre,	
99	" Plenush Roomter	182
172	" Gansel's Bergamot, 18	1, 182
	tt Goodalo	181
182	Goodale,	181
182	Glout Morceau,	182
41	"Howell,	181
3	voscumie de Mahnos	181
105	THE ALENGE	180
153	House Donne to	
139	- Vullt) ernon	182
178		
30	NOCE E CI. + 4 4 5	
43	" Tyson,	· . ·
14		116
2, 188 ;		· · · ·
- 50 İ	"Winter Nelis,	
151		181
182		153
		119
		119
		179
18	Petaloid Pollen	183
67	Petaloid Pollen, Phosphates, Canadian, Plea for Flowers	126
90	Plea for Flowers	43
113	Plea for our Small E	37
131	Plea for our Small Fruits, Plum-Leaved Source	
1791	Plum-Leaved Spirea,	190
110	Plum Curculio,	i3, 95 70
		72
	"Columbia, Coe's Golden David	182
7 71	Coe's Goldon D	182
7, 71	" Duane's Purple,	182
	urpie,	182

27, 71

τ7

State of the second sec

INDEX TO VOL. I.

Pluma Red Ol	$\mathbf{pa}_{\mathbf{g}}$
Plums—Early Orleans,	- 18
lass Seedling	- 18
oreen Gage	- 18
imperial Gage.	8
Jenerson.	5
Lontoar(έ
ALCLANGHIN	18
plarten s	18
"Peach Plum,	
Prince Englebert,	18
"Prince of Wales,	18
" Pond's Soodling	18
" Pond's Seedling,	18
" Quackenboss,	18
" Red Gage,	18
Reine Claude de Bavay,	18
" Saint Catharine,	18
* ICLOFIA	18
r chow r.gg	18
rianus and Insects	1
LUMME Grise Ande	
roug s Seedling Plan	1
roume de rer	_
r oor man's Garden.	,
rithrose, Uninese	7, 5
I Fince Englebert Plum.	1
rince of Wales "]
Prenus Triloha	
Propagation of Flowering Shrubs	54,
Pride of the Hudson Raspberry,	
Provincial Exhibition Raspberry,	1
Provincial Exhibition,	1
Pruner, The Apple Tree]
I A SHARE SUBDEPT OF FIRENCE	1
r ver omanus puparnim	
Purple-Leaved Filbert,	54,

) ;

Q.

Quackenhoss Plum,	18
Oneboo E	14
	14
the state of the s	52, 9
Cultivation of	15

0. 1

÷.,

R.

Rambo A.	le, Ne w
P	10,
maspberry.	New Notes
"	Notes
••	Notes,
41	High the second se
44	rightand Handr
	Highland Hardy, Pride of the Herd
Record II	Pride of the Hudson, ly Apples,
heccent Har	V Annles
Refugee Ro.	ny Apples, un
RALES DO	tween plants & insects
relations be	Ween plant
Red Com 1	lum,
D vage 1	lum
neine Clau	la da D
Roda	le de Bavay Plum,
Astrac	an Apple
	an Apple,

page		page
	Rebecca Grape,	
182		
182		147
81	Rhode Island Greening Apple,	166
81	RoLers' No. 44 Grape,	183
81	Roseau Apple,	50
182	Roses,	97, 105
$182 \\ 182$	" Tree and Weeping,	112
		99
18 2	Rose Slug,	
182	Rose-Leaf Hopper,	99
182	Rose Bug,	99
182	Rose, The, a Type of Infinity,	132
182	Rose, A good	160
182	Roxbury Russet Apple,	179
182		
182	S	
182	N.	
182	Salutatory,	1
-00	Си т	

s.

182	Salutatory,	1
-88	St. Lawrence Apple,	14
45	Sable Queen Blackberry,	90
182	Saint Catharine Plum,	182
49	Sarnia. Fruit culture near	154
74 23	Sap, Rise and Fall of	147
182	Scribner Spitzenburg Apple,	13
	Scarlet Flowering Thorn,	78
182	Senasqua Grape,	183
, 95 04	Seek no Further,	180
$\frac{94}{135}$		181, 182
135	Shelter,	5, 11
170	Sherwood's Favorite Apple,	178
114	Shrubs, Chinese Doub, Fl'g Plum	52
	Deutzia gracilis,	94
37	" " crenata fl. pl.,	94
l, 95	" Early Flowering,	52
	" Japan Quince,	52
	" Plum-leaved Spirea,	52
100	" Purple leaved Filbert,	52
182	" Syringa,	94
151	"Weigela Rosca, Sheldon Pear,	191 100
2, 94	Sinclair Blackberry,	
121	Small Fruits, A plea for our,	90 190
	Snow Apple,	130
	Snap Beans,	63
	Snyder Blackberry,	90
179	Souvenir du Congres Pear	21
113	Sphinx Quinque-maculatus	184
123	Special Manure for Asparague	66
109	Stenocerus putator.	169
158	Strawberry Notes	120
109	"Belle,	20
135	" Col. Cheney,	108
49	" Crescent Scedling,	108, 125
-63	** Cumberland Triumph	108
- 88	" Duchesse	108
182	" Forest Rose	12
182	" Gen. Sherman,	21
140	Great American,	108

page 21

108

108

 $\mathbf{48}$

105 114

179

45

94

Strawberry,	Hervy Davis,
"	Long John,
44	Monarch of the West,
* *	New Dominion,
	Wilson,
* *	New Varieties,
Strawberry	Apple of Montreal
Success in 1	ransplanting.
Summer Me	eting of the F. G. A.
Summer Pro	ning of Grape,
Swaar Appl	6,
Swayzie Por	mme Grise Apple,
Syringa,	

717

oyrmga,	- 94)	wax or butter bean,	63
		Wealthy Apple,	14
154		Weeping Roses,	112
A •		Wojgala Passa	94
		Weigela Rosea,	
Talman Sweet,	179	White Astracan Apple,	14
	3, 75	White Marowfat Bean,	63
Tetotsky Apple,		White Doyenne Pear, 181,	182
Tettigania Pose	14	Wilder Grape,	183
Tettigonia Rosa,	100	Wilson's Early Blackberry,	91
Three Popular Plums,	- 81	Willie en A	
Tomatoes, 150,	184	Wilson Strawberry,	57
4 Juno		Winter Meeting,	3
Acme,	184	Windsor Bean,	63
" Criterion,	184		183
Parly Conqueror.	150	Wintering Geraniums, 160,	
General Grant,	150	Winter Nelis,	181
"Hathaway,	184	Woman's Work in Horticulture, .	84
¹⁴ Hubbord's imp. and last			
"Hubbard's imp. curl leaf,	150	· I.	
Trubny	-150)	
rausplanting Evergreens.	34	Yellows in the Peach, 15,	172
Trees	48	V. W. M.	180
Tree Roses,	110	Yellow Newtown Pippin, 179,	190
Twenty Ounce Areal	112	1 ellow Bellflower Apple	179
Twenty Ounce Apple,	178	Yellow Egg Plum,	182*
Tyson Pear,	316	Yucca Filamentosa,	161
	.10	- wood 1 Hamontosa,	101

INDEX TO CONTRIBUTORS.

 $page_{\perp}$

BONDWOORTIG UT CI	1 · · · · · · · · · · · · · · · · · · ·
Bonnycastle, Henry, Campbellford, 62, 111	McAinsh, J. M., St. Mary's, 71, 160
Bond, Hiram, Meaford,	
Ramot D. D. T.	Did Digger, Hamilton,
Burnet, Rev. R., London, 11, 45, 166	Peacock, George, Mount Salem, 64
DUCKE, P. E., Ottawa 43 168	Roy, Wm., Owen Sound 183
	Saunders Wm., London, 8, 28, 72, 77, 92
Crement, nev. v., Peterborough, 184	Qual 6 1 1 11, London, 0, 20, 12, 11, 92
Cron, John, Aultsville 41	Smale, S. B., Wroxeter,
Darling, Rev. W. S., Toronto, 30	Smith, A, M., Drummondville, 15, 120
Darling, Rev. W. S., Toronto, 30	Weakerstern T
LIAIIIgiou, John, Omnah cc cc	(wonverton, Linus, 55 65 116 169 185
Fellow Worker, Paris,	The Editor, 1, 2, 13, 14, 17, 18, 20, 22, 23,
renow worker, Paris, 187	05 00 01 2, 19, 14, 17, 18, 20, 22, 23,
	40, 20, 30, 31, 33, 36, 38, 49, 52, 57, 63,
Gott, B., Arkona, 67	69, 78, 79, 81, 82, 90, 94, 97, 105, 112,
Grenny, F. J., Brantford, 165	112 114, 02, 90, 94, 97, 100, 112,
Ives H Batania N V	1 10, 114, 110, 110, 117, 119, 121, 125,
Ives, H., Batavia, N. Y., 20, 27	126, 129, 130, 131, 135, 137, 140, 141,
Mills, W. H., Hamilton 47	147 160, 100, 101, 100, 107, 140, 141,
Morrow All, T. Y. OL V. A	1 = 100, 101, 102, 100, 100, 101, 102, 100, 101, 102, 100, 101, 102, 100, 101, 102, 100, 101, 102, 100, 100
Moyer, Allen, Jordan Station, 123	164, 167, 174, 177, 188, 190.

Utter's Red Apple, Varieties of Gladiolus,

w,

Wagner Apple, 179, 180

Victoria Plum,....

Warning to Peach Growers,

Wax or Butter Bean,

14 176

182

15

63

nage

U--**V**.

The Canadian Borticulturist.

VOL. I.]

ST. CATHARINES.

[NO. 1.

THE CANADIAN HORTICULTURIST.

The Directors of the Fruit Growers' Association have long felt the importance of having a monthly publication as a medium of communication between the members, and a means of imparting information on subjects of interest, more frequently and promptly than can be done by the Annual Report. And now, after careful deliberation, they have decided to make the experiment, and commence to-day the issue of the HORTICULTURIST, in the hope that it will find favor with the It will be devoted chiefly to the publication of such informembers. mation as is sought after by those who are interested in fruit culture, yet not neglecting those kindred subjects which are closely connected with that pursuit. The lover of fruits is also usually a lover of flowers, and delights to surround the house with a well kept lawn. It will therefore contain accasional articles intended to guide and help those who seek to cultivate flowering plants and shrubs, and to make their grounds bright with summer flowers. And if the less showy, but not less important vegetable garden should have a place now and then in these pages, there are those among the readers, it is believed, who will welcome any timely information in this department also.

But while the Directors will spare no pains to make the HORTICUL-TURIST acceptable and profitable, it will nevertheless be, in a very large degree, what the members shall make it. If they shall use it as the medium through which they tell each other of success and of failure with particular fruits, flowers, trees, &c., and in which they ask for information upon doubtful points, then will it become what the Directors hope, a mirror, in which is reflected continually the Horticultural progress and skill of Ontario. They ask therefore that the members will regard it as their publication, put forth in their interests, to help them in whatever way it can, and to be used by them for the promotion of Horticulture in this Canada of ours.

THE BURNET GRAPE.

As long ago as in the Autumn of 1873, Mr. P. C. Dempsey, Albury, Prince Edward County, exhibited at the Fruit Growers' meeting a few bunches of a grape that on account of the beauty of its appearance, its earliness of ripening, and delicacy of flavor, attracted much attention and called forth universal praise. In due time a committee was appointed to visit Mr. Dempsey's grounds and examine the vine and fruit; and such was the character of their report that the Directors requested Mr. Dempsey to propagate it largely, so as to be able to supply the Association with vines sufficient to give one to each member. Since this arrangement was made, the members have become familiar with its general appearance through the colored lithograph which was presented to them in the Report for 1876: Mr. Dempsey has given to this excellent grape the name of our honored President, and henceforth it will be known in the Pomological world as the "Burnet" grape.

This grape was raised by fertilizing the Hartford Prolific with pollen from the Black Hamburgh. The vine seems to possess much resemblance to the Hartford Prolific, is a vigorous grower, of robust and healthy constitution, very productive and hardy. The fruit is very like that of the Black Hamburgh, the bunch is large, slightly shoulder ed; berries large, sweet, and delicately flavored, having nothing of the foxiness of the Hartford Prolific. The flesh is tender, almost melting, with none of the tough pulpiness of the most of our hardy grapes. also ripens early, somewhat earlier than the Hartford Prolific, and considerably before the Concord. Our members are to be congratulated on the reception of so valuable a grape-one that gives promise of being held in lasting estimation as a variety of unusual excellence, and adapted to general cultivation in nearly all parts of our Province. It will be sent to all who are members this year as early in the Spring as the season will permit.

WINTER MEETING.

The regular Winter meeting was held in the City of Hamilton, on Wednesday, the sixth of February. The President, Rev. R. Burnet,

took the chair; and after the reading of the Minutes by the Secretary, introduced to the members Mr. Craig, Secretary of the Agricultural and Arts Association, of Ontario, and Mr. J. B. Jones, delegate from the Horticultural Society of Western New York. The gentlemen were most enthusiastically welcomed by the members, and addressed the meeting in a few well-timed words of hearty interest in the object of our Association.

Mr. Chas. Arnold—our accredited delegate to the Winter meeting of the W. N. Y. Horticultural Society-read his Report of what he heard and saw on that occasion. He stated that there was an average attendance during the two days of the meeting, 23rd and 24th of January, of about one hundred and forty intelligent fruit growers from all parts of the State of New York, and adjoining States. The evening session of the 23rd was largely taken up with a discussion upon the best means of destroying the Codlin Moth. One gentleman spoke for nearly two hours, advocating the merits of his patent invention for catching the larve of this Moth. (Our cousins are highly gifted in the talking line, and are an exceedingly inventive people.) This invention consisted of a piece of water-proof paper or pasteboard, lined with cotton batting. This was to be placed, in the form of a band of about three inches in width around the trunk of each tree, with the cotton batting next to the tree, and occasionally taken off and the larvæ found therein destroyed. Another man had applied for a patent for substantially the same thing, only in this case the cardboard was punched full of holes, and the cotton batting pressed into the holes. It was fully admitted by all who took part in the discussion, that the Codlin Moth was a very serious pest, and that every owner of a pear or apple tree should wage a war of extermination against it. The larvæ will take refuge under anything that gives them shelter and security, hence any contrivance that offers them a hiding place will be sought by them, and can be used as a trap for eatching and killing them.

A very convenient trap has been made by fastening a strip of old carpeting or of cotton fiannel around the trunk of the tree, and removing it every week or ten days and passing it through an old clotheswringer, so as to crush the larve that have taken refuge in it, and then putting it back around the tree. Those who desire to inform themselves more fully on the subject of the Codlin Moth, will find much valuable information in the entomological part of the Report for 1870, page 91; for 1872, page 5; for 1874, page 43; and those who have only the Report for 1877, will find the insect figured in all its stages of existence at page 46 of the entomological part.

t

Mr: Arnold further reported that the morning session was taken up by the reading of essays, some of them containing much valuable information. The essays were upon our public roads; gathering, marketing, and preserving apples; small fruits; spring flowering shrubs; the kitchen garden; horticultural botany; roses, and weeping or drooping trees. But few fruits were exhibited. A plate of the Columbia pear was the finest plate of Winter pears he ever saw, judging from the appearance merely, as no opportunity was given him to test their flavor-

Reports from different parts of the State shewed that an immense revenue is derived from the sale of apples. Niagara County alone reported sales amounting to three hundred thousand dollars. Other counties reported as high as five hundred thousand dollars worth of apples, besides large sums for pears and other fruits.

Mr. Arnold close l his Report by expressing the hope that the day was not far distant when reports similar to those made to the Western New York Society, will come from many counties in Ontario, where both soil and climate are certainly equal to any portion of the State of New York; and ventured the prediction that in view of our already great and yearly increasing facilities for shipping, the growing of first class fruit in Ontario must be profitable for many years to come.

The subject of fruit statistics, brought before the meeting by Mr. Arnold's closing remarks, was briefly discussed, and Messrs. Burnet, Beadle and Bucke were appointed a Committee to interview the Government, and devise means for obtaining reliable statistics of the quantity and value of the fruits raised and exported from Ontario. A resolution was also passed requesting the railways to incorporate in their annual report on the crops, the condition and extent of the apple crop.

The discussion now turned upon the Canker Worm—an insect pest that is doing considerable damage to apple orchards in some sections. A full description of the Canker Worm, and engravings shewing the insect in all its stages, from the egg to the moth, will be found in the entomological part of the Report for 1870, at page 86; also a very rull article on the Canker Worms in the same part of the Report for 1875, page 25. Mr. Bowman, of Hamilton, said that for the past two years

they had stripped the leaves completely off from some two or three hundred of his apple trees-they did their work early in Spring, and disappeared about the 15th of June. He had read that syringing the trees with a mixture of Paris Green and water was complete destruction to the worms. Mr. Woolverton, of Grimsby, had suffered severely from the Canker Worms, and had tried several means of preventing their ravages. He had tied bandages around the trunks of the trees and smeared them with pitch tar, and found this a very easy and successful method of destroying the female moths. The tar must be renewed as often as it becomes hard, or the moths will crawl over it. Last year he had applied Paris Green in water with a garden engine, and found that also very beneficial. This must be done very early in the season, as soon as the buds burst, to be effectual. He had also tried fall ploughing of his orchard in the end of October, and thought this also had been beneficial, by lessening their numbers. Mr Smith, of Glanford, suggested that a mixture of castor oil and resin, ---such as is used in making the sticky fly-paper---might be found useful, though in cold winter weather it would become too hard. Molasses mixed with tar was also suggested, but rains will wash the molasses out and leave only the tar. D. W. Beadle, of St. Catharines. remarked that the use of some sticky substance, over which the wingless female moths could not crawl, would be found to be the most certain and convenient method of preventing their ravages.

P. E. Bucke, of Ottawa, read an able paper on irrigation, which was heard with marked interest and attention. This paper has been handed to the Secretary, and will appear in full in the Annual Report.

A. M. Smith, of Drummondville, called attention to the Yellows in peach trees, a disease which has been very destructive to the trees in many places, and was making its appearance in this Province. His views are given more fully in an article on this subject which will be found in this number.

The meeting proceeded to the consideration of the benefits of shelter to peach orchards, and the trees which are the best to plant for this purpose. C. M. Honsberger, of Jordan Station, had planted his peach trees between the rows of apple trees, and let them take their chances, but now, however, had been induced to plant some evergreens on the south-west side for a wind-break, and had set out a row of Norway spruce. W. Haskins, Hamilton, spoke of fifty acres of peach orchard

at Navy Island in which he was interested, and said that the best trees and the best fruit were to be found in that part of the orchard that was sheltered. He was also convinced that good cultivation of the soil was just as necessary for the production of fine peaches as for anything else. A. M. Smith would protect peach orchards on the south, south-west W. Holton, Hamilton, remarked that the peach orchards about Brantford seemed to thrive best on a poor soil where they were sheltered, and that in the rich hollows they did not succeed. thought that our native arbor-vita, or as it is often called, white cedar, and the native white pine, and black spruce were excellent trees to plant for shelter, and easily procured. Chief Johnson, of Tuscarora, thought the sugar maple an excellent tree to plant for shelter. Dempsey, Albury, advocated planting the basswood, because it grew rapidly, afforded as good shelter as any deciduous tree, and from its blossoms the bees gather the best honey, fully equal to, if not better, than white clover honey. W. McKenzie Ross, Chatham, spoke favorably of the Scotch pine, because it was a hardy tree and rapid grower. Aultsville, thought that the Norway spruce was the most valuable tree for shelter belts, it being even a more rapid grower than the Scotch pine, very dense in its habit and symmetrical in form. D. W. Beadle, St. Catharines, concurred fully in this opinion; he had seen this tree planted around a large field devoted principally to a pear orchard; in a very few years it had attained to a height of ten or twelve feet, and was quite He believed also that at present it was the cheapest tree that could be planted, cheaper than gathering up the white pines and spruces of our forests, for the reason that the Norway spruce having been several times transplanted, was very sure to grow, and could be bought, of small sizes, about as cheap as the cost of digging up the native trees. W. Roy, Owen Sound, spoke favorably of the Norway spruce, Austrian pine, and Scotch pine as shelter trees. J. B. Jones, Rochester, N. Y., spoke highly of the Norway spruce, saying that it was a hardy tree, easily transplanted, easily kept within any desired limits, and com-The European larch was also a graceful tree, of rapid growth, and very cheap.

On the subject of fertilizers for fruit trees, Mr. Robertson, of Oakville, said that in sandy soils he had found that the application of clay around the trees proved to be very beneficial and lasting in its effects. L. Woolverton, Grimsby, had also used clay around trees growing in sandy soil with marked benefit. P. E. Bucke, Ottawa, suggested the use of mineral phosphates, and spoke of the large beds which had been found near Ottawa, whence considerable quantities were being shipped to Europe. J. McGill, Oshawa, thought wood ashes to be one of the very best fertilizers for fruit bearing trees. C. Arnold, Paris, preferred barnyard manure, this he considered preferable to all other fertilizers, believing it contained all that was needed both for the tree and the fruit. J. B. Jones, Rochester, N. Y., would apply lime and ashes liberally to orchards growing in heavy soils, occasionally plow under some green crop, and apply barn-yard manure. He remarked that the practice of composting barn-yard manure, and allowing it to stand some time in large heaps, where it would ferment and decay, was now believed to be erroneous, and that the best results were obtained by applying it to the land as quickly as possible, without allowing any opportunity for fermentation.

The Report of the Committee on fruits was read. This occasioned a short discussion on the value of the Ben Davis apple. W. Holton, Hamilton, remarked that he feared many planters of this variety would be disappointed in the quality of the fruit, it not being equal in this respect to many of our older sorts. The tree was hardy, and it might on that account be a valuable sort to plant where the higher flavored kinds could not be grown. P. C. Dempsey, Albury, remarked that one of his neighbors had found it a very profitable orchard variety.

The Summer meeting will be held in the city of St. Catharines, on Wednesday, July 10th, at ten o'clock A. M.

CHINESE PRIMROSES.

We commend these beautiful plants to our readers for the reason that we have found, them among the most desirable and satisfactory for window cultivation of all the various things we have grown in the sitting room. They are very abundant bloomers, and keep up a succession of flowers for many months, so that from December to May they are continually bright and beautiful; they are easily grown by the merest novice in plant culture, requiring only to be kept from the frost, and regularly supplied with water. They can be had of several shades, red, pure white, and striped red and white, and both single and double.

ONE OF OUR COMMON INSECTS.

BY W. SAUNDERS, LONDON, ONT.

Most of our readers will recognize in the accompanying cut, FIG. 1, an object with which they are more or less familiar, although they may know little of its origin or the nature of its contents.

During the Winter months, when our trees and shrubs are leafless these curious silky structures are readily seen, and are found on many different trees and shrubs, but perhaps oftener on the twigs of apple trees and currant bushes than anywhere else. They are the cocoons of a very large and beautiful moth, called the Cecropia moth, (Attacus Cecropia,) which thus spends the winter in a quiet and torpid condition.

If you cut a twig on which one of these cocoons has been hung, and shake it, you will feel that it contains a heavy body which is to some extent moveable, and

you can feel a slight dull thud as it falls from side to side. This winter home of the insect is about three inches long. shaped something like a pod, tapering toward**s** each end, and invariably fastened lengthwise to the twig. It is of a dirty brown colour; the exterior is very close and apery like, although nuch wrinkled, and is juite impervious to wet. Let us look inside of it; underneath the close exterior we find a mass of loosely woven threads of strong yellow silk which surround the dark brown chrysalis and fill the intervening space, the upper end of the cocoon where the moth is eventually to make its escape, being

FIG. 1.

much looser in texture than the other portions. The crysalis itself, the object of all this care, is smooth, of a dull brown colour, and about one and a half inches long, and $\frac{5}{8}$ of an inch broad in the widest portion.

Early in June—or if the cocoon is kept in a warm room, many weeks before this-a marvellously beautiful moth issues from this snug enclosure. When the time has come for its escape, the shelly structure of its prison-house is rent, split open along the back, and at once restless movements begin within; the struggling creature as it tries to free itself, making a scratching noise as it tears away the silken bars which stand between it and the outer world, and this noise can be distinctly keard at some distance from the object. At this juncture a fluid is secreted from the mouth of the insect which so softens the silk as to make the escape of the moth a comparatively easy matter, while without this wise provision it might remain in its cell and exhaust itself in fruitless efforts to get out. Presently the fore legs appear, thrust out of the upper end of the cocoon, then the head crowned with its beautiful feather-like antennæ; and very soon a heavy looking object with a large plump body and soft clumsy little wings is drawn slowly out of the orifice, and stands before you in the free air.

The first care of the moth is to place itself in such a position that its wings may hang downwards, the only favourable posture for their proper development, then a rapid process of growth or expansion begins, resulting in full maturity in about half an hour, during which time the wings enlarge from the size of an ordinary humble bee until they measure from five to six inches across.

This magnificent creature is nicely represented in Fig. 2. Both front and hind wings are of a rich brown; the anterior pair greyish, shaded with red, while the posterior are more uniformly brown; about the middle of each of the wings there is a nearly kidney-shaped white spot, shaded more or less with red, and margined with black. A wavy dull red band crosses each of the wings, bordered within on the front wings, more or less faintly with white, while on the hind pair the band is widely and clearly margined with the same colour. The outer edges of the wings are of a pale silky brown, in which, on the anterior pair, runs an irregular black line which on the hind wings is replaced by a narrow, double broken band of the same hue. The front wings next to the shoulders are dull red with a curved white and black band, varying in distinctness in different specimens, and near their tips there is an eye-like black spot with a bluish-white crescent. The upper side of the body and the legs are dull red, with a wide band behind the head,

y

and the hinder edges of the rings of the abdomen white; the under side of the body is also irregularly marked with white. Below, the wings are very much like the upper surface, but paler.




Fig. 2.

These gay creatures are nocturnal in their habits, flying like bats in the dim twilight and dusky night. After pairing, the female deposits her eggs, numbering 200 or more, a process which occupies some time, as the eggs are not laid in patches, but fastened singly with a glutinous material, usually on the under side of a leaf. The eggs are about onetenth of an inch long, nearly round, of a dull creamy white colour, with a reddish spot near the centre. They usually hatch in about a week or ten days.

The subsequent history of this curious and beautiful insect will be given hereafter.

SHELTER

BY REV. R. BURNET, LONDON, ONT.

Few situations are there but require shelter, either from the bleak and stormy winds of Winter, or from the bare and neglected look of a place without trees. A very little thought and planning would accomplish all that is wanted. Having matured a plan for ornamentation and shelter, the best way to carry it out is to do it piecemeal; continuous planting is much more satisfactory to the man of taste than making a rush at the work, and then leaving it very much to take care of itself. This continuous planting, too, commends itself to the pecuniary means of most people.

To a large extent the farmer and amateur gardener should be his own nurseryman. As a general rule, the trees that are most recommended and used for shelter are easily grown from seed; the requisite is to know the *how* of growing them. At Arkona, Mr. B. Gott, the orchardist, raises an erection about three feet from the ground and covers it with brush; the seeds are sown, spring up, and make a good growth even in the course of one season. Let them be transplanted in due time in some suitable spot, and all trees necessary for shelter are at hand.

Wind-breaks, either for shelter or ornamentation, are of great moment to the fruit grower, in fact almost essential. The one tree that stands out pre-eminently is the Norway spruce, (*abies excelsa;*) its hardiness, rapidity of growth, and ease of cultivation strongly recommend it; for a windy country, or where ice and snow are apt to injure the trees, the elasticity and strength of the branches render them safe from damage. Planted in rows for screens, eight feet apart is not too close.

Perhaps the Scotch and Austrian pine follow next, as the most valuable trees for shelter. Planted alternately they make a fine appearance; the former very dark in the foliage, and the latter a whitish green. They are both hardy, and afford a large amount of protection.

In Europe we have seen the Himalayan pine interspersed among other varieties with good effect.

For a hedge, nothing can equal the arbor vike, or white cedar; it forms a perfect wind-break and stands our Winters perfectly. The planting must be closely done, and if slightly pruned in the tips, the spray becomes intensely thick and impervious.

At our Winter meeting in Hamilton, Mr. Holton urged the planting of our common white pine—what might be called the pine of the country. It is one of the best trees for shelter, however you look at it; perfectly handy, moderately quick in growth, acclimatized, and valuable for its timber. The same gentleman also mentioned the hardy black spruce.

In Essex, when recently there, and also in the county of Elgin, we saw the European larch planted as a wind-break; its growth is something wonderful, and the shelter perfect; although deciduous, the spray is so small and close that it cuts the wind, and makes a complete calm on the sheltered side. We might add perhaps, that this variety should be planted in the Fall.

At Tyrconnell the walnut is used for shelter, and a noble screen it makes. It is planted on Mr. Comis' grounds in triple rows. Few finer sights can be seen when in the full season of flowering and fruiting. We greatly wish that its popularity were on the increase, the more especially so, as our soil in many districts is admirably adapted for the walnut. The nuts should be planted as soon as they fall, four feet apart, and three inches deep. The first year they will make a growth of fourteen or fifteen inches, the next, three feet, and in the third year they will be of sufficient size to render cultivation unnecessary.

Many advocate the planting of the sugar maple, and the soft maple; both have their advantages, being deciduous however, they cannot compare to the many varieties of the conifere.

In answer to the question, when, and how to transplant evergreens, much has been, and will be said. Some say, just when the buds first begin to swell; others affirm as strongly and persistently in the latter end of May and during the month of June. One or two requirements are absolutely necessary to insure its growth; first, see that the earth is firmly in contact with the roots. Mr. Meehan, of Philadelphia, uses beaters in planting; we are persuaded that the plan is a good one. Mulching is an essential; with proper mulching a single tree need not be lost.

Regard is also to be had to the particular district where the shelter is needed. In some of our western counties, in many parts of Essex and Kent, the white ash (*Fraxinus Americana*,) might be planted with beneficial results; the arboriculturist looking to future recompense, as well as to present benefit; its strength, elasticity, and durability making it invaluable for the manufacture of agricultural implements. The walnut, butternut, and larch are well adapted for the extreme points of the western peninsula. Norway spruce flourishes everywhere, and the same may be said of the Scotch and Austrian pine.

A cheap and effective wind-break can be formed by protecting the trees that spring up around uncultivated fences. In the neighborhood of Hamilton, several farmers have derived benefit from this means, both for their crops and fruits.

This question, however, is not unlikely shortly to assume larger proportions. The stripping of the country of its forests; the long bleak tracts, inviting the violence and injurious influence of winds, will speedily demand a remedy. Arboriculture should be allied to the other efforts pat forth by the F. G. A. of Ontario; not till then will the subject meet with that attention which it so justly deserves. That all efforts to accomplish this laudable result, may meet with an able advocate in our HORTICULTURIST, and that this paper may lend a helping hand in the good work, is the aim and object of your essayist.

SCRIBNER SPITZENBERG APPLE.

A correspondent of the *Gardeners' Monthly* residing at Plattsburg, N.Y., claims that he has in this variety an apple of the peculiar flavor of the Esopus Spitzenburg, while the tree is more hardy and vigorous, and an abundant bearer. In 1859 it was exhibited at the annual meeting of the New York State Agricultural Society, and received a silver medal. The fruit is very like the Esopus Spitzenburg in form, but somewhat more angular, and in color a lighter red. This may prove

to be a valuable fruit for general cultivation in Ontario, for Plattsburg lies in about the same latitude with Peterboro' and Barrie, and if hardy and productive at Plattsburg, it should do well in many of our colder sections.

APPLES IN MINNESOTA.

Seeing that the climate of the State of Minnesota is very severe upon fruit trees, our readers who live in the more trying sections of the Province will be interested to know what kinds of apples are found to answer there. At St. Paul, Minnesota, the mean temperature for the three Summer months is about the same as that of London, Ontario; while the mean temperature for the three Winter months is about that of Pembroke, in the County of Renfrew, or Three Rivers, in the Province of Quebec. Thus it will be seen that fruit trees in Minnesota are exposed to very severe cold in Winter, while the Summer heat is also quite considerable.

On looking at the transactions of the Minnesota State Horticultural Society for the last year, we notice that only two apples are recommended for general cultivation, namely, the Duchess of Oldenburg, and Wealthy. This indicates that the climate of that State must be very unfavorable to the apple tree. The Duchess of Oldenburg has been cultivated in this Province for a number of years, and has borne the severity of our Winters unharmed. The Wealthy is not as well known, indeed it is doubtful if it has been planted in many of our orchards. The Tetofsky is recommended for planting in limited quantities. The reason for this limitation does not seem to be brought out in the discussion, unless it be because one gentleman lost a number of trees of this variety in the Winter of 1873. The St. Lawrence, Utter's Red, and Snow apple or Fameuse were recommended for favorable localities; and the White Astracan and Elgin Beauty for general trial throughout the State. So far as the White Astracan has been cultivated here, it has been considered a fruit of poor quality, not to be grown where better sorts will thrive. The Wealthy is a very good, medium sized fruit, in use from December to February, which originated near St. Paul, in Minnesota, and seems to have maintained a character for extraordinary hardiness; it is worthy the attention of those who require a tree capable

of enduring a very low temperature. The Elgin Beauty originated in the township of Elgin, in Wabasha county, Minnesota; it is a medium sized fruit, streaked with red on a yellow ground, moderately juicy, sub-acid, in season from November to March. The writer is not aware that it has been planted in Ontario.

A WORD OF WARNING TO PEACH GROWERS OF ONTARIO.

BY A. M. SMITH, DRUMMONDVILLE, ONTARIO.

Perhaps it is not generally known, but it is nevertheless a fact, that the disease so destructive to peach orchards called the yellows, has made its appearance in our midst. Quite a number of orchards along the frontier, particularly in the vicinity of Drummondville and Stamford, have had affected trees in them the last season, and some in the great peach growing section of Grimsby. The symptoms of this disease are, 1st, an enfeebled vitality, the foliage looks sickly; and, 2nd, the fruit ripens prematurely, sometimes two or three weeks before its usual season for maturing, it is usually high coloured, red and flecked or spotted, and is red around the stone. This occurring in young trees newly planted, has led many to think they had some new variety which was very early; but the flavor is universally insipid and watery, and the fruit nearly Hundreds of bushels of them were sold in Western New York last season, their color recommended them, but no one would care to buy them a second time. This disease, according to Downing, showed itself about the year 1800, in the vicinity of Philadelphia, where many orchards showed decay and death without any apparent cause, and it has since spread into nearly all the peach growing sections of America. That the disease is contagious there is no doubt, and it is also hereditary. Seeds from diseased trees will produce diseased nursery stock, and buds taken from them will produce disease where inserted, and the pollen from the flowers of diseased trees is also believed to carry the disease to trees that are contiguous. the disease has been spread over the country. It is therefore of great In these ways importance to those planting trees to procure them from localities that are free from this disease, and from parties who would use the utmost care in getting seeds and buds not affected with it.

Is there a remedy for the disease? It is said that when the disease shows itself in an orchard it is difficult to eradicate it, unless the trees showing the first symptoms are taken out, root and branch. In Michi gan they have a law compelling people to dig them out; and I under stand that there is such importance attached to this matter that vigilance committees are appointed in some localities, who visit orchards, ordering out every tree that shows any symptoms of this disease. But notwithstanding all their vigilance many of their orchards have been destroyed If such extreme measures are necessary there, I should think by it. that self-interest at least would dictate to every peach grower of Ontario to be on the alert, and remove all symptons of it as soon as it appears I know there are some people who langh at the idea of the yellows being here, and attribute the sickly condition of their trees to the cold Winter of three or four years ago, and I do not doubt that the Winter referred to injured the trees in some sections, and by enfeebling them, made them much more susceptible to disease. But I saw this same disease on several trees in an orchard in Niagara Co., N. Y. the Fall before the severe Winter spoken of, (and I think some of it in Canada,) and now that orchard is totally destroyed, and several others in its immediate vicinity are badly affected with it.

Professor Beal, microscopist of the Michigan Agricultural College, has been making observations to learn if possible the cause of the disease He has detected several forms of fungoid growth attached to the roots of the trees, and this matter has so adapted itself as to enter into the circulation of the sap of the tree. Professor Redgie, of the same college, thinks the disease may be traced to this cause, and that an enfeebled condition of the tree caused by excessive bearing while young, or other enfeebling causes, may greatly increase the liability of the attack. In the analysis of the ashes of healthy, and diseased peach trees, it has been found that diseased trees lack two important elements, potash and phosphoric acid. Now it is an established fact that these are of the first importance among inorganic elements of tree growth, and this deficiency suggests a remedy. ("An ounce of prevention is worth a pound of cure.") Keep your trees supplied with plenty of potash in the soil, give them good cultivation, thin out the fruit and not let them exhaust themselves by over-bearing, particularly when young, and they will be less liable to an attack of this disease; and be vigilant in taking out all diseased trees when first attacked, and you may save your peach orchards.