





**IMAGE EVALUATION
TEST TARGET (MT-3)**



**Photographic
Sciences
Corporation**

23 WEST MAIN STREET
WEBSTER, N.Y. 14580
(716) 872-4903

**CIHM/ICMH
Microfiche
Series.**

**CIHM/ICMH
Collection de
microfiches.**



Canadian Institute for Historical Microreproductions / Institut canadien de microreproductions historiques

© 1983

Technical and Bibliographic Notes/Notes techniques et bibliographiques

The Institute has attempted to obtain the best original copy available for filming. Features of this copy which may be bibliographically unique, which may alter any of the images in the reproduction, or which may significantly change the usual method of filming, are checked below.

L'Institut a microfilmé le meilleur exemplaire qu'il lui a été possible de se procurer. Les détails de cet exemplaire qui sont peut-être uniques du point de vue bibliographique, qui peuvent modifier une image reproduite, ou qui peuvent exiger une modification dans la méthode normale de filmage sont indiqués ci-dessous.

- Coloured covers/
Couverture de couleur
- Covers damaged/
Couverture endommagée
- Covers restored and/or laminated/
Couverture restaurée et/ou pelliculée
- Cover title missing/
Le titre de couverture manque
- Coloured maps/
Cartes géographiques en couleur
- Coloured ink (i.e. other than blue or black)/
Encre de couleur (i.e. autre que bleue ou noire)
- Coloured plates and/or illustrations/
Planches et/ou illustrations en couleur
- Bound with other material/
Relié avec d'autres documents
- Tight binding may cause shadows or distortion along interior margin/
La reliure serrée peut causer de l'ombre ou de la distortion le long de la marge intérieure
- Blank leaves added during restoration may appear within the text. Whenever possible, these have been omitted from filming/
Il se peut que certaines pages blanches ajoutées lors d'une restauration apparaissent dans le texte, mais, lorsque cela était possible, ces pages n'ont pas été filmées.
- Additional comments:
Commentaires supplémentaires:

- Coloured pages/
Pages de couleur
- Pages damaged/
Pages endommagées
- Pages restored and/or laminated/
Pages restaurées et/ou pelliculées
- Pages discoloured, stained or foxed/
Pages décolorées, tachetées ou piquées
- Pages detached/
Pages détachées
- Showthrough/
Transparence
- Quality of print varies/
Qualité inégale de l'impression
- Includes supplementary material/
Comprend du matériel supplémentaire
- Only edition available/
Seule édition disponible
- Pages wholly or partially obscured by errata slips, tissues, etc., have been refilmed to ensure the best possible image/
Les pages totalement ou partiellement obscurcies par un feuillet d'errata, une pelure, etc., ont été filmées à nouveau de façon à obtenir la meilleure image possible.

This item is filmed at the reduction ratio checked below/
Ce document est filmé au taux de réduction indiqué ci-dessous.

10X	12X	14X	16X	18X	20X	22X	24X	26X	28X	30X	32X
							X				

The copy filmed here has been reproduced thanks to the generosity of:




La Bibliothèque de la Ville de Montréal

The images appearing here are the best quality possible considering the condition and legibility of the original copy and in keeping with the filming contract specifications.

Original copies in printed paper covers are filmed beginning with the front cover and ending on the last page with a printed or illustrated impression, or the back cover when appropriate. All other original copies are filmed beginning on the first page with a printed or illustrated impression, and ending on the last page with a printed or illustrated impression.

The last recorded frame on each microfiche shall contain the symbol \rightarrow (meaning "CONTINUED"), or the symbol ∇ (meaning "END"), whichever applies.

Maps, plates, charts, etc., may be filmed at different reduction ratios. Those too large to be entirely included in one exposure are filmed beginning in the upper left hand corner, left to right and top to bottom, as many frames as required. The following diagrams illustrate the method:



L'exemplaire filmé fut reproduit grâce à la générosité de:

La Bibliothèque de la Ville de Montréal

Les images suivantes ont été reproduites avec le plus grand soin, compte tenu de la condition et de la netteté de l'exemplaire filmé, et en conformité avec les conditions du contrat de filmage.

Les exemplaires originaux dont la couverture en papier est imprimée sont filmés en commençant par le premier plat et en terminant soit par la dernière page qui comporte une empreinte d'impression ou d'illustration, soit par le second plat, selon le cas. Tous les autres exemplaires originaux sont filmés en commençant par la première page qui comporte une empreinte d'impression ou d'illustration et en terminant par la dernière page qui comporte une telle empreinte.

Un des symboles suivants apparaîtra sur la dernière image de chaque microfiche, selon le cas: le symbole \rightarrow signifie "A SUIVRE", le symbole ∇ signifie "FIN".

Les cartes, planches, tableaux, etc., peuvent être filmés à des taux de réduction différents. Lorsque le document est trop grand pour être reproduit en un seul cliché, il est filmé à partir de l'angle supérieur gauche, de gauche à droite, et de haut en bas, en prenant le nombre d'images nécessaire. Les diagrammes suivants illustrent la méthode.


aire
détails
ues du
modifier
ger une
filmage

ées

re

y errata
d to

nt
ne pelure,
çon à



THE

North American Sylva;

OR, A DESCRIPTION OF THE

FOREST TREES

OF THE

UNITED STATES, CANADA, AND NOVA SCOTIA.

CONSIDERED PARTICULARLY

WITH RESPECT TO THEIR USE IN THE ARTS AND THEIR
INTRODUCTION INTO COMMERCE.

TO WHICH IS ADDED

A DESCRIPTION OF THE MOST USEFUL OF THE

EUROPEAN FOREST TREES.

ILLUSTRATED BY 166 COLORED ENGRAVINGS.

TRANSLATED FROM THE FRENCH OF

F. ANDREW MICHAUX,

MEMBER OF THE PHILOSOPHICAL SOCIETY OF PHILADELPHIA, ETC. ETC.

WITH NOTES BY J. JAY SMITH,

EDITOR OF THE HORTICULTURIST, MEMBER OF THE ACADEMY OF NATURAL SCIENCES, ETC.

IN THREE VOLUMES.

VOL. I

PHILADELPHIA:
W. M. RUTTER & CO.,
SEVENTH & CHERRY STREETS.

G

183, 1860

1871

1868

Entered according to Act of Congress, In the year 1865, by
 RICE, RUTTER & CO.,
 In the Clerk's Office of the District Court of the United States for the Eastern District of
 Pennsylvania

STEREOTYPED BY L. JOHNSON AND CO.
 PHILADELPHIA.

1871



CAXTON PRESS OF
 SHERMAN & CO., PHILADELPHIA.

TO

SAMUEL GEORGE MORTON, M.D.,

PRESIDENT OF THE ACADEMY OF NATURAL SCIENCES, PHILADELPHIA.

This Edition

OF


MICHAUX'S NORTH AMERICAN SYLVA

IS AFFECTIONATELY INSCRIBED

BY HIS FRIEND AND BROTHER,

J. JAY SMITH.

February, 1850.



PREFACE TO THE EDITION OF 1857.

THE Philadelphia editions of this important work have had a wide circulation in the several States of the Union, proving how extensively the taste of the public is turned to the study of Arboriculture. The whole of the sheets of the last imprint were destroyed by a fire at the bindery, whither they had been sent for collation; but fortunately the original French copper-plates were in another building. This has enabled the new publishers to issue the work in a much improved style, and has allowed opportunity for additional notes and remarks. These might have been more extended; but it was thought best not to swell the work beyond a reasonable charge.


Issued as it is in connection with Nuttall's continuation, the whole forms a work of reference of unrivalled interest and beauty.

The period which has elapsed between the editions has taken from us the individuals to whom the public is indebted for the lucid descriptions here reprinted. F. André Michaux died in Paris in November, 1855. Dr. Samuel George Morton also paid the debt of nature on the 15th of May, 1851, aged fifty-two years, beloved and lamented by a large circle of friends. Thus pass away the admirers of nature's works: even

"Art, Glory, Freedom, fall; but
Nature still is fair."—BYRON.

J. J. S.

[NOTE.—When Michaux wrote, *Louisiana* included all the territory west of the Mississippi, excepting Texas and New Mexico and the territory west of the Rocky Mountains. This is so well known that it has not been thought necessary to alter the text in any instance where that State is mentioned.]



P R E F A C E.

THE foundation of the North American Sylva was laid by the laborious researches of the elder Michaux; who, under the auspices of the French Government, devoted ten years, from 1785 to 1796, to a thorough exploration of the country, from the sunny sub-tropical groves of Florida to the cold and inhospitable shores of Hudson's Bay; repeatedly visiting all the higher peaks and deepest recesses of the Alleghany Mountains, and extending his toilsome journeys westward to the prairies of Illinois and the banks of the Mississippi. He proposed to Mr. Jefferson, then Secretary of State, to extend his researches to Oregon, but was prevented from doing so by untoward circumstances.

Soon after his return to France, and the year before he fell a victim to scientific zeal upon the coast of Madagascar, the elder Michaux published his history of North American Oaks, which may be deemed the nucleus of this more comprehensive work, subsequently issued by his son, who accompanied his father in the earlier portions of his travels. Revisiting this country in 1801, and again in 1807, the son made the extended and toilsome researches of which these volumes are the result: they were first published in Paris, in 1810-13.

They were translated into English by Hillhouse, and printed in Paris with French types, in 1819. This edition has been long since exhausted; the second English edition was produced at New Harmony, Indiana, but was carelessly executed on very inferior paper, though, like the present, the engravings were printed from the original copper-plates partly engraved by the celebrated Redouté, which had been brought from Paris by the liberal friend of education and science, the late William McClure, with a view of making the work more generally known among the American people. His brother and executor, Alexander McClure, Esq., of New Harmony, still keeping in view the future utility to the community of these expensive engravings, presented them to the late Dr. Samuel George Morton, successor of William McClure in the Presidency of the Academy of Natural Sciences of Philadelphia.

In passing this edition through the press, I have not thought it advisable to make extensive alterations in the text, but have left it, with some corrections in the translation, as it was written by its distinguished author, adding a few observations on soil, propagation, &c. &c. These additions may always be distinguished by their being enclosed in brackets. For corrections of Hillhouse's translation, and in other particulars, I cannot but acknowledge my great indebtedness to my friend Thomas Forrest Betton, M.D., of Germantown, Pennsylvania.

An improvement in the work would have consisted in rearranging the plates according to the demands of modern Science and nomenclature; but this would have required the renumbering of them, and thus all the numerous references to these in other books would have been erroneous and confused.

It was a singular circumstance, and a happy one it has proved for advancing science, that Mr. Nuttall arrived in this country the very year that the younger Michaux left it. From that time he devoted his talents to Botany, and after visiting a large portion of the United States, with an aptitude for observation, a quickness of eye, tact in discrimination, and tenacity of memory, rarely possessed by one man, he published his extended and most happily-executed botanical work, the "Genera of North American Plants." In 1834 he crossed the Rocky Mountains and explored the territory of Oregon and Upper California. With his peculiar qualifications, he prepared the supplement to Michaux's Sylva, in three handsome volumes, corresponding in size with the present, the publication of which, after many delays, was completed in 1849, by my son, in Philadelphia. The two works are now one and homogeneous, the former most highly valued by all lovers of trees, and the latter destined to be equally so, when the fine products of our newly-acquired Western regions make their way to our gardens and plantations.

The frequent references I have made to Mr. Nuttall's volumes will show the reader that his additions to our Sylva are both extensive and important; inspection will convince him that both authors stand on the highest pedestal of merit.

J. JAY SMITH.

CONTENTS OF VOLUME FIRST.

	PAGE
WHITE OAK. <i>Quercus alba</i>	22
Common European Oak	30
European White Oak	32
Mossy-Cup Oak	33
Over-Cup White Oak.....	35
Post Oak.....	36
Over-Cup Oak.....	39
Swamp White Oak.....	41
Chestnut White Oak.....	44
Rock Chestnut Oak	46
Yellow Oak	49
Small Chestnut Oak.....	50
Live Oak.....	52
Cork Oak.....	55
Willow Oak	58
Laurel Oak.....	60
Upland Willow Oak.....	61
Running Oak	63
Bertram Oak	64
Water Oak.....	65
Black Jack Oak.....	67
Bear Oak	69
Barrens Scrub Oak	71
Spanish Oak.....	73
Black Oak	76
Scarlet Oak	79
Gray Oak	81

	PAGE
Pin Oak.....	<i>Quercus palustris</i> 83
Red Oak.....	<i>Quercus rubra</i> 84
Common European Walnut	<i>Juglans regia</i> 97
Black Walnut	<i>Juglans nigra</i>104
Butternut.....	<i>Juglans cathartica</i>109
Pecanutt Hickory	<i>Juglans olivæformis</i>114
Bitternut Hickory	<i>Juglans amara</i>116
Water Bitternut Hickory	<i>Juglans aquatica</i>119
Mockernut Hickory	<i>Juglans tomentosa</i>120
Shellbark Hickory	<i>Juglans squamosa</i>123
Thick Shellbark Hickory	<i>Juglans laciniosa</i>128
Pignut Hickory	<i>Juglans porcina</i>132
Nutmeg Hickory.....	<i>Juglans myristicæformis</i>135
White Maple.....	<i>Acer eriocarpum</i>146
Red-Flowering Maple.....	<i>Acer rubrum</i>149
Sugar Maple.....	<i>Acer saccharinum</i>153
Black Sugar Tree.....	<i>Acer nigrum</i>163
Norway Maple	<i>Acer platanoïdes</i>165
Sycamore.....	<i>Acer pseudo-platanus</i>167
Moose Wood	<i>Acer striatum</i>169
Box Elder.....	<i>Acer negundo</i>172
Mountain Maple.....	<i>Acer montanum</i>175
Dogwood	<i>Cornus florida</i>176
Georgia Bark.....	<i>Pinckneya pubens</i>180
Coffee Tree.....	<i>Gymnocladus Canadensis</i>182

.....	PAGE
.....	83
.....	84
.....	97
.....	104
.....	109
.....	114
.....	116
.....	119
.....	120
.....	123
.....	128
.....	132
.....	135
.....	146
.....	149
.....	153
.....	163
.....	165
.....	167
.....	169
.....	172
.....	175
.....	176
.....	180
.....	182

THE
NORTH AMERICAN
SYLVA.

OAKS.

In the greater part of North America, as well as in Europe, there is no tree so generally useful as the Oak. It is everywhere the most highly esteemed in the construction of houses and of vessels, and is commonly selected for implements of husbandry. It seems, also, to have been multiplied by nature in proportion to its utility. Without insisting upon the diversity of climates to which it is indigenous, we may observe that the number of its known species is already considerable and is daily increasing, particularly on the Western Continent, and that its varieties are infinite. These considerations determined my father in 1801, after his return from the United States, to publish a treatise containing drawings and descriptions of the Oaks of that country, which was favorably received by the lovers of botany and agriculture.

The following extract from his work exhibits a just outline of this tree:—"The genus of the Oaks (Introduct. p. 4) comprises many unknown species; most of those which grow in America

exhibit such various forms while young that they can be ascertained with certainty only when arrived at maturer years. Often an intermediate variety so nearly resembles two species that it is difficult to determine, from the foliage, to which of them it belongs. Some species are so variable that it is impossible, by the leaves, to recognise their identity in youth and at a more advanced age. Others are so similar that specific characters must be derived from the fructification, which is itself liable to variations and exceptions. It is only by a comparison of stocks of different ages that analogous species can be distinguished and varieties correctly referred to their species.

“I have endeavored to arrange the American Oaks in a natural series, the characters of which I first sought in the fructification: but this afforded only unimportant distinctions, such as the position of the barren flowers, whether pedunculated or nearly sessile, and the size and period of the fruit. Neither was I able to find my distinction on the structure of the cup. I was obliged therefore to have recourse to the foliage, which has been made the basis of a division into two sections, the first containing the species with beardless leaves, and the second, those in which the summit or lobes are terminated by a bristle.

“The interval between the appearance of the flower and the maturity of the fruit is different in different species; and this distinction I have admitted as a secondary character.

“All the Oaks are proved to be monoecious. We know, too, that on the European White Oak and other species the female flowers are situated above the male upon the shoots of the same season; that both are axillary; and that, immediately after the fecundation, the male flowers fade and fall, while the female blossom continues advancing through the natural stages till, in the course of the year, it ripens into perfect fruit. But there are some species whose fertile flowers remain stationary a whole year and begin to develop their germ the second spring, probably because they are not fecundated the first season; so that

eighteen months elapse between the appearance of the flower and the maturity of the fruit. Hence I have formed a subdivision into species of annual and species of biennial fructification. The female flower which is axillary the first season ceases to be so, of course, at the falling of the leaf. Several species are found upon the Old Continent whose fructuation is biennial, such as the Cork Oak, *Quercus suber*, etc."

I have derived great assistance from my father's work, and have adopted his arrangement, which perfectly accords with my own observations. But I have inserted several new species, and have suppressed two that were not well ascertained: the existence of one of them is doubtful, and the other is evidently a duplicate.

The chief distinction between my work and his consists in the more extended practical observations, which are the fruit of my own researches. My constant aim was to appreciate the utility of each species in the mechanical arts, and to point out those which are the most deserving of attention in Europe and America. If in this respect mine has some advantage, my father's work will always preserve its title to the attention of botanists and amateurs of foreign plants, by other details not consistent with my plan. They will find, for example, quotations from all the authors who had previously taken notice of the species he describes, and, in the plates, leaves of the young plant as well as of the full-grown tree.

I have described twenty-six American species, which I have divided into two sections, according to the term of fructification: the first comprising ten species that bear fruit every year; and the second, sixteen, of which the fructification is biennial. I have learned by multiplied observations that, with the exception of the Live Oak, the wood of the first section is of a finer texture, more compact, and consequently more durable.

Linnaeus, in the third edition of his *Species Plantarum*, published in 1774, described fourteen species of Oak, of which five

only are natives of the New World. Since that period such additions have been made to the list that the new edition of Willdenow's *Species Plantarum*, published in 1805, contains forty-four American species; of which sixteen were recognised by Messrs. Humboldt and Bonpland in Old Mexico, and twenty-six by my father and myself in the United States and the adjacent countries. Probably the American series will be still further augmented by discoveries in the western part of Louisiana, and in the interior provinces of New Spain,—a country 1200 miles in extent, lying between the United States and Old Mexico, which no naturalist has explored.

In America, as we have just observed, are found forty-four species, which are all comprised between the 20th and 48th degrees of north latitude; on the Old Continent are enumerated only thirty, which are scattered on both sides of the equator, beginning at the 60th degree north.

This sketch is not without utility, and appears naturally in this place; such parallels might perhaps contribute more than is generally thought to the progress of botany and agriculture, and they deserve particular attention from naturalists travelling in foreign countries. It would be interesting to possess comparative tables of those plants which are found in the higher latitudes of both continents, and of the trees and shrubs of the temperate climates of America with the analogous species found in nearly the same latitude in Asia. I have long entertained a wish, which will doubtless be shared by all who interest themselves in the science, that botanists would go more deeply into the geography of plants. The rapid progress of the young Americans who are beginning to devote themselves with ardor to the study of Natural History will soon afford the requisite information concerning their own portion of the globe.

[For a continuation of the subject, and for further interesting particulars respecting the Oaks, see Nuttall's Supplement to this

work, vol. i. p. 1, *et seq.* Six new species are there figured, with additional information regarding several treated of by Michaux.

Soil, Situation, and Climate. The Oaks, to attain their full size, require a deep, loamy soil, a situation low rather than elevated, and a climate not liable to late spring frosts, which injure both the blossoms and leaves. In elevated situations, or in the extreme North, those species which, under favorable circumstances, form the most magnificent trees, become, as in the case of other trees, mere shrubs. The Oaks which flourish on the worst soils are the low-growing kinds belonging to the section *Ilex*, and especially those belonging to the group *Phellos*; and those which require the best soil are the *Quercus sessiliflora*, the *Q. cerris*, and most of the sorts composing the American group *Rubræ*.

Propagation, &c. The Oak is propagated with difficulty by every other mode except from seed; and, generally, time will be gained when the acorns are sown where the plants are intended finally to remain. It is only, therefore, where peculiar varieties are to be continued that the process of grafting is resorted to; and the mode by approach is almost the only one that is certain to be attended with success. The best stock for grafting on is *Q. cerris*, on which some sorts may be successfully budded.

The acorns need not be gathered from the tree, but may be collected from the ground immediately after they have dropped, and may either be sown then or kept till the following spring. If they are to be kept, they should be made perfectly dry in the sun or in an airy shed, mixed with dry sand, in the proportion of three bushels of sand to one of acorns, or with dry moss, and then excluded from the air and vermin, by being put into barrels or boxes, or laid up in a cellar, or buried in heaps and covered with a sufficient thickness of earth to exclude the weather. Very few of any species will germinate after having

been kept a year. When acorns are to be sown in a nursery, the soil ought to be thoroughly prepared and rendered fine; and after the earth is drawn off the beds, or the drills opened, the acorns may either be scattered over the beds, or along the drills, so that the nuts may be about two inches apart. The acorns, before covering, must be patted down with the back of a spade in the beds, and with the back of a wooden-headed rake in the drills. The covering, of well-broken soil, should vary in depth according to the size of the acorn, $1\frac{1}{2}$ inches being enough for those of the largest size and half an inch for those of the smallest size. No mode of depositing acorns in the soil can be worse than that of dropping them in holes made by too small a dibble. The acorn drops into the hole, and becomes wedged by its sides before it gets to the bottom; and, if the upper extremity should be downward instead of upward, it can hardly be expected to grow. Sown late in March, the period between the depositing the acorn and its becoming a plant is lessened and the danger from destruction by vermin somewhat diminished. When it is necessary to remove the plant, the tap-root should be shortened a year at least in advance; side-pruning is soon necessary when the object is a straight clean trunk.

The American Oaks vary so exceedingly in their leaves at different seasons of the year, in different stages of their growth, and in different localities, that some experience is necessary in deciding on them.

Like most other trees, the Oak seldom bears an abundant crop of fruit for two years in succession, and it increases in productiveness with age. All the species push up shoots from the collar when cut down, but only one or two species from the root. After Oaks have stood in good soil and a suitable climate for five or six years, they grow with rapidity till they have attained the age of thirty or forty years; and the life of some species is known to extend to upward of one thousand years.

There are some Oaks in Britain which are believed to have

been old trees in the time of William the Conqueror; and Pliny mentions a *Quercus ilex* which was an old tree when Rome was founded, and which was still living in his time. The Merton Oak measures at the surface of the ground sixty-three feet two inches.

The Cowthorp Oak, in Yorkshire, measures seventy-eight feet in circumference near the ground, and its age is estimated as nearly coeval with the Christian era. An Oak in Lower Charante, in France, is declared, on good authority, to measure from eighty-five to ninety-four feet.

Particular attention should be given to the remarks of the author on the subject of planting the Oak for future use. The General or State Governments should never grant a charter for a railroad or canal, without a clause requiring the planting of useful trees, such as the White Oak, for instance, at the North, and the Live Oak wherever the climate will admit, along both sides of the route. A store of ship-timber would thus be accumulating for national or mercantile service, whence it could easily be transported to the seaboard in emergencies,—a plan which would shade the road and be advantageous to the banks of a canal. The French Government has shown a wise foresight in this particular; her turnpikes are often thus planted, and the product is at the call of the authorities.]

METHODICAL DISPOSITION
OF THE
OAKS OF NORTH AMERICA,
INCLUDING
THREE EUROPEAN SPECIES.

Monecia polyandria. LINN. *Amentaceæ*. JUSS.

FIRST DIVISION.

Fructification annual; leaves beardless.

FIRST SECTION.—*Leaves lobed.*

1. White Oak *Quercus alba.*
2. Common European Oak *Quercus robur.*
2. European White Oak *Quercus robur pedunculata.*
3. Mossy-cup Oak *Quercus olivæformis.*
4. Over-cup White Oak *Quercus macrocarpa.*
5. Post Oak *Quercus obtusiloba.*
6. Over-cup Oak *Quercus lyrata.*

SECOND SECTION.—*Leaves toothed.*

7. Swamp White Oak *Quercus prinus discolor.*
8. Chestnut White Oak *Quercus prinus palustris.*
9. Rock Chestnut Oak *Quercus prinus monticola.*
10. Yellow Oak *Quercus prinus acuminata.*
11. Small Chestnut Oak *Quercus prinus chincaquin.*

SECOND DIVISION.

Fructification biennial; leaves mucronated, (except in the 13th species.)

FIRST SECTION.—*Leaves obtuse or entire.*

- 12. Live Oak *Quercus virens.*
- 13. Cork Oak *Quercus suber.*
- 14. Willow Oak *Quercus phellos.*
- 15. Laurel Oak *Quercus imbricaria.*
- 16. Upland Willow Oak *Quercus cinerea.*
- 17. Running Oak *Quercus pumila.*

SECOND SECTION.—*Leaves lobed.*

- 18. Bartram Oak *Quercus heterophylla.*
- 19. Water Oak *Quercus aquatica.*
- 20. Black Jack Oak *Quercus ferruginea.*
- 21. Bear Oak *Quercus banisteri.*

THIRD SECTION.—*Leaves multifid or many-clefted.*

- 22. Barrens Scrub Oak *Quercus catesbæi.*
- 23. Spanish Oak *Quercus fulcata.*
- 24. Black Oak *Quercus tinctoria.*
- 25. Scarlet Oak *Quercus coccinea.*
- 26. Gray Oak *Quercus ambigua.*
- 27. Pin Oak *Quercus palustris.*
- 28. Red Oak *Quercus rubra.*

CA,

culata.

lor.

stris.

icola.

inata.

capin.

WHITE OAK.

QUERCUS ALBA. *Q. foliis subæqualiter pinnatifidis; laciniis oblongis, obtusis, plerumquæ integerrimis; fructu majusculo; cupulâ crateratâ; tuberculoso-scabratâ; glande ovatâ.*


THROUGHOUT the United States, and in Canada, this tree is known by the name of *White Oak*. The environs of the small town of *Trois Rivières*, in Canada, latitude $46^{\circ} 20'$, and the lower part of the river Kennebeck, in the district of Maine, are the most northern points at which it was observed by my father and myself. Thence we traced it along the sea-shore to a distance beyond Cape Canaveral, latitude 28° , and westward from the ocean to the country of the Illinois,—an extent of more than 1200 miles from northeast to southwest, and nearly as much from east to west. It is, however, by no means equally diffused over this vast tract; in the district of Maine, Vermont, and Lower Canada, it is little multiplied, and its vegetation is repressed by the severity of the winter. In the lower part of the Southern States, in the Floridas and Lower Louisiana, it is found only on the borders of the swamps with a few other trees which likewise shun a dry and barren soil. This region is generally so sandy that it is covered with a continued growth of Pines, as will be more particularly mentioned in the description of the Long-leaved Pine. The White Oak is observed also to be uncommon on lands of extraordinary fertility, like those of Tennessee, Kentucky, and Genesee; and, of all the spacious valleys watered by the Western rivers, I have travelled whole days in those States without seeing a single stock, though the few that exist, both there and in the Southern States, exhibit the most luxuriant vegetation.

The White Oak abounds chiefly in the Middle States and in Virginia, particularly in that part of Pennsylvania and Virginia

blongis, ob-
atâ; tuber-

s tree is
the small
the lower
, are the
y father
to a dis-
ard from
of more
early as
equally
ermont,
tation is
part of
ana, it is
er trees
egion is
growth
descrip-
ved also
ke those
spacious
d whole
ugh the
exhibit

and in
Virginia





Ed. H. B. K.

White Oak.
Quercus alba.



which lies between the Alleghanies and the Ohio, a distance of about 150 miles, beginning at Brownsville on the Monongahela. Near Greensburg, Macconellsville, Unionville, and Washington Court-house, I have seen large forests, nine-tenths of which consisted of White Oaks, whose healthful appearance evinced the favorable nature of the soil, though in general they were not more than fifteen inches in diameter. East of the mountains this tree is found in every exposure, and in every soil which is not extremely dry or subject to long inundations; but the largest stocks grow in humid places. In the Western districts, where it composes entire forests, the face of the country is undulated, and the yellow soil, consisting partly of clay with a mixture of calcareous stones, yields abundant crops of wheat.

By the foregoing observations, it appears that the severity of the climate, the fertility of the soil, its dryness or humidity, are the causes which render the White Oak so rare over three-quarters of the United States that it is inadequate to supply the local demand, though the country does not contain a fourth of the population which it is capable of supporting.

Among the American Oaks this species bears the greatest analogy to the European Oak, especially to the variety called European White Oak, (*Quercus pedunculata*.) which it resembles in foliage and in the qualities of its wood. The American White Oak is seventy or eighty feet high, and six or seven feet in diameter; but its proportions vary with the soil and climate. The leaves are regularly and obliquely divided into oblong, rounded lobes, destitute of points; the sections appeared to be the deepest in the most humid soils. Soon after their unfolding, they are reddish above and white and downy beneath; when fully grown, they are smooth and of a light green on the upper surface and glaucous underneath. In the fall they change to a bright violet color, and form an agreeable contrast with the surrounding foliage which has not yet suffered by the frost.

This is the only Oak on which a few of the dried leaves persist till the circulation is renewed in the spring. By this peculiarity and by the whiteness of the bark, from which it derives its name, it is easily distinguished in the winter. The acorns are of an oval form, large, very sweet, contained in rough, shallow, grayish cups, and borne singly or in pairs, by peduncles eight or ten lines in length, attached, as in all the species with annual fructification, to the shoots of the season.

The fruit of the White Oak is rarely abundant, and frequently for several years in succession a few handfuls of acorns could hardly be collected in a large forest where the tree is multiplied. Some stocks produce acorns of a deep blue color; but I have found only two indications of this variety, one a flourishing tree in the garden of Mr. W. Hamilton, [now the Woodlands Cemetery,] near Philadelphia, and the other in Virginia.

The trunk is clad in a white bark, variegated frequently with large black spots. On stocks less than sixteen inches in diameter the epidermis is divided into squares; on old trees, growing in moist grounds, it is in the form of plates laterally attached. The wood is reddish, and very similar to that of the European Oak, though lighter and less compact, as may be proved by splitting billets of each of the same size; in the American species the vessels which occupy the intervals of the concentric circles are visibly less replete. But, of all the American Oaks which I shall describe, this is best and most generally used, being strong, durable, and of large dimensions. It is less employed than formerly in building only because it is more scarce and costly.

At Philadelphia, Baltimore, and in the smaller towns of the Middle States, the frame of all well-built houses, whether of brick or wood, is of White Oak. West of the Alleghanies, where pine boards are not easily procured, the White Oak is substituted for the floors and for the exterior covering of the frame, notwithstanding its liability to warp and split.

It is much used in the construction of mills and dams, particularly for such parts as are exposed to be alternately wet and dry.

The wooden bridge, nearly 3000 feet long, that unites Boston and Cambridge, is supported by posts of White Oak, from fourteen to fifty feet in length, which replaced those of White Pine on which it originally stood.

The excellent properties of this wood cause it to be preferred for a great variety of uses, among which are many articles manufactured by the wheelwright. This trade is carried to the greatest perfection at Philadelphia, and its wares are highly esteemed for solidity both at home and abroad. White Oak perfectly seasoned is employed for the frame of coaches, wagons, and sledges, for the mould-board of ploughs, the teeth of wooden harrows, the felloes and spokes of wheels, particularly the spokes of coach-wheels. In the Northern, Middle, and Western States, the naves are also made of Oak, in the country; but it splits too easily to be proper for this object. Except in the district of Maine, it is always chosen for the bow or circular back of windsor-chairs. The wood of the young stocks is very elastic, and is susceptible of minute division; hence it is preferred for the large baskets used in harvesting, for the hoop of sieves, the bottom of riddles, and the handles of coach-whips, which are braided and covered with leather; at Boston it is chosen for pail-handles, and in Maine for axe-helves.

In many parts of the Middle States the White Oak is selected for the posts of rural fences, and beyond the Laurel-Hill Mountain, in Pennsylvania, where it is common, it forms the entire enclosure.

The bark is considered by many tanners as the best for preparing leather for saddles and other similar objects: it is little used, however, because in the United States the bark of the trunk and large limbs only is employed; and on these the cellular tissue is much thinner in the White than in the Red Oak, which is, besides, more abundant.

I have been told that the bark yields a purple dye. Though I have not witnessed the fact, I am disposed to believe in its existence, as I received the information from persons residing several hundred miles from each other. But, if the color was not defective in permanence or intensity, it would have found its way into commerce, like the *Quercitron* of the Black Oak.

Of all the species that grow east of the Mississippi, the White Oak alone furnishes staves fitted for containing wine and spirituous liquors. The domestic consumption for this purpose is immense, and vast quantities are exported to the West Indies, Great Britain, and the islands of Madeira and Teneriffe. The Post Oak might, indeed, be applied to the same use; but even in Maryland and Virginia, where it is most common, it is not sufficiently multiplied to supply the local demand.

The Rock Chestnut Oak and the Swamp White Oak in the Northern and Middle States, the Chestnut White Oak and the Over-cup Oak in the South, are compact enough to prevent the escape of spirits and fine oils, yet porous enough to absorb them. If they united every requisite quality, and were employed for this purpose, they would be consumed in less than ten years.

It is well understood at Bordeaux that the wood of the European White Oak is closer-grained than that of the American species, and the preference is given to our domestic growth, or to that imported from Dantzic. The American Oak is exclusively employed in Maderia and the West Indies, only because it is cheaper and more easily procured.

White Oak staves are exported from all the parts of the Northern and Middle States, and from New Orleans. Those which come from Baltimore, Norfolk, and New Orleans, are far superior to those of the Northern States; the difference results naturally from that of the soil and climate.

The quantity of Oak staves exported to England and the West Indies appears, by two official documents that I have examined, to be considerable. In 1808, the value received by

England amounted to more than \$146,000, and the number of staves sent to the West Indies exceeded 53,000,000. I am unable to fix the proportion of the two species of White and Red Oak; probably more of the first are sent to England, and of the second to the Colonies. The price of both has varied surprisingly within a hundred years. In 1720, staves for barrels were sold at Philadelphia at \$3 a thousand; in 1798, at \$18; and in 1818, at \$30. In August, 1807, before the American Embargo, they were advertised at \$55, and in April, 1808, after that municipal regulation became known, at \$100.

The young White Oak, on account of its elasticity, is very proper for hoops, but it has less strength and less durability than the Hickory.

Among the uses of this wood, the most important is in ship-building. In all the dock-yards of the Northern and Middle States, except in the district of Maine, it is almost exclusively employed for the keel, and always for the lower part of the frame and the sides; it is preferred for the knees when sticks of a proper form can be found. In the smaller ports south of New York, the upper part of the frame is also of White Oak; but such vessels are less esteemed than those built of more durable wood.

At Boston, the tree-nails, or the pins by which the side-planks are attached to the ribs, are of this species.

To obtain correct notions on the comparative value of the American White Oak and the European Oak, I consulted French, English, and American shipwrights, in almost all the ports of the United States. They generally agreed that the European Oak was tougher and more durable from the superior closeness of its grain, but that the American species was more elastic and required a shorter time, with only half the weight, to bend it. This advantage, though important in ship-building, does not compensate for the openness of its pores. Experience, however, every day shows that by growing in places long inhabited its

quality is improved; and, if the American vessels are less durable than those built in Europe, it is because the timber is not thoroughly seasoned.

The greater part of the immense quantity of White Oak exported from the United States is sent to England. It is shipped from the Northern and Middle States in the form of boards and of square timber: what goes to England from Quebec is brought from the shores of Lake Champlain, for Canada probably furnishes hardly enough for its own consumption.

By an extract from the custom-house books of St. John, which I have already quoted, 143,000 cubic feet of Oak would appear to have entered by this port during the first six months of 1807. Oddy, in his *Treatise on the Commerce of Europe*, says that in the English dock-yards the White Oak from British America is esteemed excellent timber. The opinion, simply considered, is correct; but that which comes from Baltimore and Philadelphia must still be superior.

Before I conclude this article, I must be allowed to hazard a conjecture on the consequences of the neglect of all means of preserving and multiplying this tree in the United States,—consequences which neither the Federal Government nor the States have taken any measures to prevent. From the increase of population, and from the impoverishment of the soil produced by a gradual change in the climate, the White Oak will probably, in less than fifty years, be the most rare in the Middle States, where it is now the most abundant, and in Tennessee, Kentucky, Genesee, and farther north, where it is the least multiplied, it will be the most common, and will replace the species which now compose the forests, but which the soil will then be too feeble to sustain. Thus, near the river Kennebeck, in the midst of the primitive forests, composed of the Beeches, the Canoe Birch, the Sugar Maple, and the Hemlock Spruce, I have observed small tracts, formerly cleared and since abandoned, which are naturally repeopled with the White and Gray

Oaks; and, in the lower part of Virginia, poor Red Oaks, Yellow Pines, and Loblolly Pines, are extensively replacing trees of a better quality. East of the mountains, the valleys that lie along the rivers are, with a few exceptions, the only places where the Oak could be advantageously reared; but these fertile lands are more profitably devoted to husbandry.

The American White Oak cannot, in my opinion, be regarded as a useful acquisition to the forests of Europe. Its elasticity, which renders the young stocks proper for hoops, is doubtless a valuable property; but the Chestnut of France is superior for this purpose, because it is more durable.

The White Oak is used in the royal dock-yards of England, probably because it has been found impossible to procure supplies of European Oak. Perhaps it is employed only for the lower part of the frame, while the European Oak is reserved for the upper timbers.

If the advantage of this comparison be allowed to be on the side of the European species, the Americans should lose no time in introducing it into their forests. To corporations particularly, whose property is less frequently alienated, I take the liberty of addressing this advice, which, if followed, would be productive of great advantage to themselves and to the public. The analogy of the climates leaves no doubt of the perfect success of this tree in the United States, an example of which is found in the garden of Messrs. J. & W. Bartram, three miles from Philadelphia, where there is a large stock which has yielded seed for several years, and which continues to expand with vigor.

[In ornamental planting, the White Oak should have abundant space around it for expanding; under such circumstances it will throw out long limbs and lateral branches of the most picturesque beauty.

This tree extends much farther to the west than is stated by

our author. Mr. Douglas considers Lake Winnipeg its northern limit, where it is rarely over ten or twenty feet high. The White Oak everywhere shrinks from the sea-breeze.

A tree of this species, near Boston, Massachusetts, measured in 1840 twenty-five feet and nine inches in circumference.]

PLATE I.

A branch with leaves and acorns of the natural size.

[See Nuttall's Supplement, vol. i. pp. 24, 33.]

COMMON EUROPEAN OAK.

QUERCUS ROBUR. *Q. foliis petiolatis, oblongis, glabris, sinuatis, lobis rotundatis; fructibus oblongis, scssilibus.*

To the particular attention bestowed upon this interesting tree in modern times is owing its division into two species,—the Common European Oak, *Quercus robur*, and the European White Oak, *Quercus pedunculata*.

These two species, which are much alike and are usually considered as the same, grow in the same countries, and frequently together. They constitute the greater part of the European forests, from the 60th to the 35th degree of north latitude, overspreading a great part of the north of Asia and the northern extremity of Africa. They are most abundantly multiplied on the shores of the Black Sea, in Germany, England, France, and some parts of Italy, where the climate is particularly favorable to their growth.

The Common European Oak is from sixty to eighty feet in

•

northern
gh. The
measured
ce.]


, *lobis ro-*


teresting
species,—
European

usually
and fre-
t of the
of north
and the
ntly mul-
England,
particu-

y feet in

•





Michx.


European White Oak.

Quercus pedunculata

Common European Oak.

Quercus robur.

Common



height, numerously ramified, and crowned with an ample and majestic summit. The bark upon the trunk is thick, and, upon old stocks, deeply furrowed. The leaves are petiolated, smooth, and of a uniform color on both sides, enlarged toward the summit, and very coarsely toothed. The acorns are oval and *sessile*, which is the principal difference between the two species.

This tree prefers high places and the declivities of hills, with a barren gravelly soil; hence it grows more slowly, and its wood is more compact, tougher, and heavier, than that of the European White Oak. It is less used for household stuff and other kinds of joinery, because it is less easily wrought; but is more esteemed for building and for works that require great strength and durability.

The Common European Oak is subdivided into many varieties, the most valuable of which are the European Black Oak, *Quercus robur lanuginosa*, and the *Quercus robur glomerata*. The first is only thirty or forty feet high, with small thick leaves, very downy underneath; its timber is compact and excellent for fuel. The second never rises to a great height; the leaves are small, but smooth on both sides; the acorns are of an inferior size and collected in clusters upon a short common peduncle.*

PLATE II.

A branch of the Common European Oak, with leaves and acorns of the natural size.

* [For a highly-interesting account of this tree and the ensuing one, *Q. pedunculata*, see Loudon's Arboretum Britannicum, vol. iii. p. 1740.]

EUROPEAN WHITE OAK.

QUERCUS PEDUNCULATA. *Q. foliis subsessilibus, glabris, sinuatis; fructibus oblongis, pedunculatis.*

THE European White Oak grows of choice in rich bottoms, where the soil is deep and moderately humid. It reaches the height of ninety or one hundred feet, and has a large, well-proportioned trunk, which is often undivided for a considerable distance, and which spreads into a large, commanding summit. The bark upon the body is very thick, and, on old trees, deeply furrowed; upon the limbs and the young stocks it is grayish, smooth, and glossy. The leaves are of a light green on the upper surface, whitish beneath, widened toward the summit, deeply sinuated with blunted points, and supported by short petioles like those of the American White Oak. They are more or less divided according to the age of the tree and to the moisture of the soil. A part of the dry discolored foliage persists through the winter and falls the ensuing spring.

Besides the difference of the foliage, this species is constantly distinguished from the preceding by its fruit, which is supported singly or in pairs by slender peduncles, two, three, or even four inches long. The acorns are of an oval shape, from nine to eighteen lines in length, according to the age and vigor of the tree, and contained in shallow cups; they fall about a fortnight before those of the Common Oak.

The wood of the European White Oak is of the same color with that of the American species, the sap being white and the heart reddish; but the texture is closer and the pores fuller, which is probably the reason of its being less elastic, but stronger and more durable. It is generally preferred to the Common Oak, as it furnishes larger timbers, splits more regu-

atis; fruc-

bottoms,
reaches the
well-pro-
siderable
summit.
es, deeply
s grayish,
n on the
summit,
by short
are more
nd to the
ed foliage
g.

constantly
h is sup-
three, or
ape, from
and vigor
l about a

same color
e and the
res fuller,
astie, but
ed to the
ore regu-



P. H. R. H. H. H.

Mossy Cup Oak
Quercus chrysolepis





larly, and is more easily wrought; hence it is highly esteemed for the construction of houses and ships, and extensively employed by the joiner, the wheelwright, and the cooper.

Throughout Europe, except in the north of Russia, the bark of the Common Oak and the White Oak is almost exclusively used in tanning. That which is taken from the branches and from small stocks is preferred, because the epidermis is thinner, and the cellular tissue, which contains the tannin, more abundant.

Oak wood is more generally used in Europe than in the United States, where the different species of Ash, Birch, etc. in some measure supply its place. The European White Oak would be a valuable addition to the American forests, and I have sent out acorns to begin the formation of nurseries.

PLATE II.

A branch of the European White Oak, with leaves and acorns of the natural size.

MOSSY-CUP OAK.

QUERCUS OLIVÆFORMIS. *Q. foliis oblongis, glabris, subtus glaucis, profunde inæqualiterque sinuato-lobatis; fructu ovato; cupulâ profundius crateratâ, superne crinitâ; glande olivæformi.*

I HAVE observed this species of Oak only in the State of New York, on the banks of the Hudson above Albany and in Genesee, where it is so rare that it has hitherto received no specific name.

Its leaves are of a light green above and whitish beneath: they resemble those of the White Oak in color, but differ from

them in form, being larger, and very deeply and irregularly lacinated, with rounded lobes so various in shape that it is impossible to find two leaves that are alike. The acorns are of an elongated oval form, and are enclosed in cups of nearly the same configuration, of which the scales are prominent and recurved, except near the edge, where they terminate in slender, flexible filaments. From this peculiarity I have derived the name of *Mossy-cup Oak*.

This tree is sixty or seventy feet in height, with a spacious summit and an imposing aspect. The bark is white and laminated; but the tree is chiefly remarkable for the form and disposition of its secondary branches, which are slender, flexible, and always inclined toward the earth. This peculiarity alone would render it a valuable acquisition for parks and gardens.

As I have met with this species only in uninhabited places, I have had little opportunity of examining its wood: as far as I can judge, it is not better than that of the White Oak, though far superior to that of the Red Oak.

[Pursh found this Oak on iron-ore hills in Pennsylvania and Virginia, and adds that in general appearance it resembles *Q macrocarpa*.]

PLATE III.

Leaves of the natural size. Fig. 1. An acorn with the cup. Fig. 2. An acorn without the cup.

[See Nuttall's Supplement, vol. i. p. 24.]

irregularly
that it is
corns are of
nearly the
ent and re-
in slender,
derived the

a spacious
e and lami-
rm and dis-
der, flexible,
liarity alone
gardens.

ted places, I
: as far as I
Oak, though

sylvania and
it resembles

e cup. Fig. 2.

]



P. J. Roberts


Roberts

Over Cup White Oak.
Quercus macrocarpa.

Pl. 4.



Rem. 10.



OVER-CUP WHITE OAK.

QUERCUS MACROCARPA. *Q. foliis subtomentosis profunde lyratimque sinuato-lobatis, obtusis; fructu maximo; cupulâ profundius crateratâ, superne erinitâ; glande turgide-ovalâ.*

THIS interesting species is most multiplied beyond the Alleghanies, in the fertile districts of Kentucky and West Tennessee, and in Upper Louisiana near the Missouri.* It is called by the Americans Bur Oak and Over-cup White Oak, and, by the French of Illinois, *Chêne à g. ou gland.*

It is a beautiful tree, more than sixty feet in height, laden with dark, tufted foliage. The leaves are larger than those of any other Oak in the United States, being frequently fifteen inches long and eight broad: they are notched near the summit, and deeply lacinated below. The acorns, which are also larger than those of any other American species, are oval, and enclosed for two-thirds of their length in a thick, rugged cup, bordered with fine flexible filaments. Sometimes, however, in compact forests or in very temperate seasons, the filaments do not appear, and the edge of the cup is smooth and bent inward.

The fructification of this tree is not abundant, and, as its wood is inferior to that of the White Oak, it is little esteemed in the United States.

I have observed, as well as my father, who first made the remark, that the young branches are frequently covered with a yellowish fungous substance, like those of the Elm and Sweet Gum.

[A specimen of this tree standing about three miles from Troy, Ohio, has been measured; its dimensions are as follows:—the

* [According to Pursh, on dry slate or limestone hills.]

diameter at one foot above the ground, seventeen feet; at six feet above the ground, fourteen feet nine inches. The trunk rises about fifty feet without limbs, and with scarcely a perceptible diminution in size. The top branches rise one hundred feet above the earth.

As a tree for ornamental planting, the *Over-cup Oak* is attracting much attention; its introduction is the beginning of a taste for the finest trees of our own land. The *macrocarpa* is surely one of these, both from its fine growth, large leaves and fruit, and its magnificence. This Oak is remarkable when young for the *corky* appearance of its bark, in which it differs from other Oaks and resembles the Cork-bark Elm.]

PLATE IV.

A leaf of half the natural size. Fig. 1. An Acorn in the cup, of the natural size.

POST OAK.

QUERCUS OBTUSILOBA. *Q. foliis sinuatis, subtus pubescens, lobis obtusis, superioribus dilatatis, bilobis; fructu mediocri; glande breviorati.*

Quercus stellata, WILLD, Sp. Pl

In New Jersey, near the sea, and in the vicinity of Philadelphia, this species is thinly disseminated in the forests, and has hitherto been considered as a variety of the White Oak. In Maryland and a great part of Virginia, where it abounds, and where its properties are better understood, it is called Box White Oak, and sometimes Iron Oak and Post Oak. The last denomination only is used in the Carolinas, Georgia, and East Tennessee.

et; at six
The trunk
r a percep-
e hundred

is attract-
of a taste
a is surely
and fruit,
young for
from other

ne cup, of the

ibus, lobis ob-
glande brevi-

HILLD, Sp. Pl

of Philadel-
ests, and has
te Oak. In
abounds, and
called Box
k. The last
gia, and East




Richard



Richardson

Post Oak
Quercus obtusiloba.

Richardson



The steep banks of the Hudson, nearly opposite to the city of New York, are the most northern point at which I have observed it. Even here its existence seems to be secured only by the influence of the sea-air, which tempers to a certain degree the severity of the winter. A little farther inland it is not found in the forests. In the vicinity of South Amboy, thirty miles nearer the sea, where the soil is dry and sandy, it is more multiplied; and it becomes still more vigorous and more common in advancing toward the South. Westward, in Pennsylvania, I saw the last individual of this species a little beyond Carlisle, on the road to Pittsburg, 150 miles from Philadelphia. Near Baltimore, at the distance of 210 miles from New York, it abounds in the woods and attains its utmost expansion. In Kentucky and Tennessee it is rare, except on the edges of the swamps enclosed in the forests, about which it is multiplied, though not fully developed. It probably exists in Lower Louisiana; for we met with it in East Florida, of which the climate is the same.

But it is nowhere more abundant than in Maryland and in Virginia, between the Alleghamies and the sea. Wherever the soil is dry, gravelly, and unsubstantial, it forms a considerable proportion of the forests, which are composed principally of the Black, Scarlet, Spanish, and Black Jack Oaks, the Dogwood and the Yellow Pine. These woods exhibit a squalid appearance, occasioned not only by the sterility of the soil, but by the injury they are constantly sustaining from the cattle which range through them at all seasons, and which in winter are compelled, by the want of herbage, to subsist upon the young sprouts and the shoots of the preceding year. The upper part of the two Carolinas and Georgia, particularly where the Pine and Oak forests unite, is analogous in soil to that portion of Virginia of which we have been speaking, and abounds in the Post Oak; but nearer the sea the barren wastes are covered with the Long-leaved Pine, and the Oak is seen only in the lowest parts of the

swamps, about the plantations, and on tracts that have been exhausted by cultivation and abandoned.

The leaves are borne by short petioles, and are of a dusky green above and grayish beneath. They are four or five inches in length, thick, and even coriaceous toward the end of summer, deeply and regularly sinuated, and are divided into four or five rounded lobes, of which the two nearest the summit are the broadest. Toward the fall the ribs are of a rosy tint, instead of a purplish red, like those of the Scarlet Oak. The fructification seldom fails. The acorns are small, oval, and covered for a third of their length with a slightly-rugged grayish cup. They are very sweet, and form a delicious food for squirrels and wild turkeys; hence the tree is sometimes called Turkey Oak.

The height of this species rarely exceeds forty or fifty feet, with a diameter of fifteen inches. Its summit, even when compressed in the forests, is disproportionately large, owing probably to the early division of the trunk into several limbs, with which the secondary branches form more open angles than is common on other trees. The branches also are bent into elbows at certain distances, which gives so peculiar an appearance to the tree that it is easily distinguished when the leaves are fallen. The bark upon the trunk is thin and of a grayish white. The wood is yellowish, with no tinge of red. Growing upon a less humid soil, it is less elastic, but finer-grained, stronger, and more durable, than the White Oak; hence it is preferred for posts, and is used with advantage by wheelwrights and coopers.

In ship-building it is used principally for the knees, and is admitted into the lower part of the frame. It rarely furnishes side-planks or timber of considerable length: for this reason it is less esteemed than the White Oak, and it is, besides, less common, except in Maryland and certain parts of Virginia.

The preference given in the West Indies to the staves from Baltimore and Norfolk is due in a great measure to their being made of the Post Oak.

have been

f a dusky
five inches
f summer,
our or five
it are the
nt, instead
e fructifica-
vered for a
up. They
s and wild
Oak.

e fifty feet,
when com-
ng probably
with which
is common
ows at cer-
to the tree
allen. The

The wood
less humid
more dura-
posts, and is

nees, and is
ly furnishes
is reason it
besides, less
Virginia.


staves from
their being



Ed. B. Kent. 1850.

Over Cup Oak.
Quercus lyrata.





This tree, though only of secondary size, should be propagated in America and introduced into the forests of Europe.

[Emerson mentions this tree as growing on Martha's Vineyard, much beyond the northern limit assigned to it by Michaux. Elliot says, "Its timber is supposed, in strength and durability, to surpass that of any other species of the Oak; and therefore it is highly prized when it can be obtained sufficiently large to be used in the construction of vessels." Staves of it are greatly esteemed: its timber is sometimes confounded with the White Oak, which it greatly resembles.]

PLATE V.

A branch with leaves and fruit of the natural size.

[See Nuttall's Supplement, vol. i. p. 23.]

 OVER-CUP OAK.

QUERCUS LYRATA. *Q. foliis subsessilibus, glabris, lyrato-sinuosis, summitate dilatata, divaricato-tribata, lobis acutangulis, terminali tricuspide; cupula depresso-globosa, muricato-scabrata; glauca subtecta.*

In the United States I have met with this interesting species only in the lower part of the Carolinas and of Georgia. It probably exists on the banks of the Mississippi in Lower Louisiana; and I have observed it on the St. John, in East Florida, in situations analogous to those in which it flourishes a little farther north. In Georgia and Carolina it is not extensively multiplied, and has been distinguished only by the inhabitants of the places where it grows. It is called Swamp Post Oak,

Over-cup Oak, and Water White Oak. The first of these denominations indicates an analogy between its foliage and that of the Post Oak, and the second a remarkable peculiarity of its fruit, of which the acorn is covered by the cup. The name of Over-cup Oak is the most common in South Carolina, and that of Swamp Post Oak on the Savannah in Georgia.

The Over-cup Oak grows in more humid situations than any other species of this genus in the United States. It is never seen in the long narrow marshes which intersect the pine-barrens, but is found exclusively in the great swamps on the borders of the rivers, which are often overflowed at the rising of the waters and are inaccessible during three-quarters of the year. In these gloomy forests it is united with the Large Tupelo, White Elm, Wahoo, Planer Tree, Carolinian Poplar, Water Bitternut Hickory, and Water Locust.

It expands to a majestic size, and the influence of a deep and constantly-humid soil is shown in the luxuriance of its vegetation. On the banks of the Savannah I have seen stocks which were more than eighty feet high and from eight to twelve feet in circumference. The leaves are six or eight inches long, smooth, narrow, lyre-shaped, deeply sinuated, and borne by short petioles. The lobes, particularly the two upper ones, are truncated, and from their resemblance in this respect to those of the Post Oak is derived the name of Swamp Post Oak. The foliage is thick and of a light, agreeable tint. The acorns, unlike those of the Oaks in general, which are of an elongated oval shape, are broad, round, and depressed at the summit: they are sometimes from twelve to eighteen lines in diameter from side to side, and from six to ten lines from the base to the summit. The cup, which is nearly closed, is thin, and its scales are terminated by short, firm points.

The bark upon the trunk is white, and the wood, though inferior to that of the White Oak and the Post Oak, is more compact than would be supposed from the soil in which it grows;

f these de-
and that of
arity of its
e name of
a, and that

s than any
It is never
the pine-bar-
the borders
ising of the
of the year.
rge Tupelo,
plar, Water

f a deep and
of its vegeta-
stocks which
twelve feet
inches long,
d borne by
per ones, are
et to those of
t Oak. The
e acorns, un-
an elongated
the summit:
s in diameter
e base to the
and its scales

od, though in-
is more com
ich it grows;



Baron del

Chalms. et

Swamp White Oak
Quercus Pr^{nc} discolor.





the pores are observable only between the concentrical circles, and are more regularly disposed than in other trees.

This species is the largest and the most highly esteemed among the Oaks that grow in wet grounds. Its propagation should be attempted in the forests of Europe, where no doubt can be entertained of its success. The acorns which I sent to France several years since, though sown upon uplands, have produced flourishing plants, which bear the winter of Paris without injury.

PLATE VI.

A branch with leaves and fruit of the natural size.

SWAMP WHITE OAK.

QUERCUS PRINUS DISCOLOR. *Q. foliis oblongo-obovatis serratis albo-tomentosis, grosse dentatis, basi integerrimis, dentibus inequalibus dilatatis; fructibus longè pedunculatis.*

Quercus bicolor. WILLD.

THIS species is known in the United States only by the name of Swamp White Oak, which indicates at once the soil which it prefers and its analogy to the White Oak.

I first observed it near Portsmouth in New Hampshire; but it is less multiplied in this latitude than in the Middle and Western States. It particularly attracted my attention in New Jersey near the city of New York, on the Delaware in Pennsylvania, on the Susquehanna in Virginia, and beyond the mountains on the Ohio in Kentucky, and on the Holston, near Knoxville, in East Tennessee; I have also seen it on the shores of Lake Champlain and Lake Ontario. Except the district of

Maine and the maritime parts of the southern section, it is diffused throughout the United States: in comparison, however, with several other species, it is not common, being found only on the edges of swamps and in wet places exposed to inundations, and not in the forests at large, like the White Oak, the Black Oak, &c. In New Jersey it is associated with the Pin Oak, the Red-flowering Maple, the White Ash, the Tupelo, and the Shellbark Hickory. On the shores of Lake Champlain, which occasionally offer similar situations, particularly at a little distance from Skeensborough, it is mingled with the White Maples, which occupy the next line to the Willows in retiring from the shore.

The Swamp White Oak is a beautiful tree, more than seventy feet in height, of which the vegetation is vigorous and the foliage luxuriant. The leaves are six or eight inches long and four inches broad, smooth and of a slightly-dark green above, downy and light-colored beneath; they are entire toward the base, which is cuneiform, but are widened and coarsely-toothed for two-thirds of their length toward the summit. The tree is distinguished when young by the form of its base and by the down upon its leaves, which is more sensible to the touch than on any analogous species. At a riper age the lower side of the leaf is of a silvery white, which is strikingly contrasted with the bright green of the upper surface; hence, the specific name of *discolor* was given it by Dr. Muhlenberg.

The acorns are sweet, but seldom abundant; they are rather large, of a brown complexion, and contained in a spreading cup edged with short, slender filaments, more downy within than those of any other Oak, and supported by peduncles one or two inches in length.

The trunk is clad in a scaly grayish-white bark. The wood is strong, elastic, and heavier than that of the White Oak. In stocks more than a foot in diameter the grain is fine and close and the pores are nearly obliterated. It splits easily and in a

straight line, and is esteemed next in quality to the White Oak, though from its rareness it is but accidentally employed in the arts.

If, as I incline to believe, the Swamp White Oak is found by more accurate experiments to be superior to the White Oak, it must be considered as a very valuable tree, and its increase should be favored at the expense of the Red-flowering Maple, the Bitternut Hickory, the Hornbeam, and other species which grow in the same exposures. It seems also to deserve a place in the forests of Europe, where, in moist grounds, it might be blended or alternated with the Ashes, the Alders, and the Poplars.

[This tree occasionally attains a large size. One is mentioned growing in a wet clayey soil, measuring twelve feet and one inch in circumference at four feet from the ground. The wood is of a brownish color, heavy, compact, and fine-grained, and is preferred by some for boat-building to the White Oak.]

PLATE VII.

A branch with leaves and fruit of the natural size.

[See Nuttall's Supplement, vol. i. p. 23.]

CHESTNUT WHITE OAK.

QUERCUS PRINUS PALUSTRIS. *Q. foliis oblongo-ovalibus, acuminatis acutis, subuniformiter dentatis; cupulâ crateratâ, subsquamosâ; glande ovalâ.*

Quercus prinus. WILLD.

THE Chestnut White Oak is first seen within ten miles of Philadelphia; but it is less multiplied and less amply developed than farther south. It is most abundant in the maritime parts of the Carolinas, Georgia, and East Florida, and is probably found on the banks of the Mississippi, which are analogous to those of many rivers of the Southern States.

In Pennsylvania this species is confounded with the Rock Chestnut Oak, which it strikingly resembles; farther south, where the Rock Chestnut Oak is unknown, it is called Chestnut White Oak, Swamp Chestnut Oak, and generally, on the Savannah, White Oak.

The Chestnut White Oak is adorned with beautiful foliage; the leaves are eight or nine inches long, four or five inches broad, obovate, deeply toothed, of a light shining green above and whitish beneath.

The acorns are brown, oval, larger than those of any other species except the Over-cup White Oak, and contained in shallow scaly cups. Being sweet-flavored, and sometimes abundant, they are sought with avidity by wild and domestic animals, such as deer, cows, horses, and swine.

The Chestnut White Oak, like the Over-cup Oak, grows only in the large swamps that border the rivers or are enclosed in the forests; but it always chooses spots that are rarely inundated, where the soil is loose, deep, constantly cool, and luxuriantly fertile.

In the Carolinas and Georgia it is usually accompanied by

minatis acu-
osâ; glande

WILLD.

a miles of
developed
itime parts
s probably
analogous to


the Rock
ther south,
ed Chestnut
on the Sa-

iful foliage;
five inches
green above

of any other
ned in shal-
es abundant,
tic animals,

. grows only
e enclosed in
rarely inun-
ol, and luxu-

ompanied by






Desse del.

Chesnut del.

Chesnut White Oak.
Quercus P.^{mo} palustris.



the White Elm, the Wahoo, the Big Laurel, the Umbrella Tree, the Sweet Leaves, the Beach, the Poplar, the Bitternut Hickory, and the Devil Wood. In this latitude it attains its utmost development, which is eighty or ninety feet in stature, with a proportional diameter. Its straight trunk, undivided and of a uniform size to the height of fifty feet, and its expansive tufted summit, form one of the most beautiful and majestic trees of the North American forests.

Its wood, which is affected by the richness of the soil, is inferior to that of the Post Oak, the White Oak, and even the Over-cup Oak; and its pores, though nearly obliterated, are more open. But it is superior to many other species, and is employed for wheelwrights' work and other objects which require strength and durability. As it splits in a straight line and may be divided into fine shreds, it is chosen by the negroes for baskets and brooms. Its pores are too open to contain wine or spirituous liquors. In the form of rails it lasts twelve or fifteen years, or a third longer than the Willow Oak. At Augusta in Georgia it is considered as the best fuel, and is sold at two or three dollars a cord.

The Chestnut White Oak endures the winter of Paris, but its vegetation would be quicker in the more southern departments. It is to be regretted that a tree which seems formed to be one of the finest ornaments of our forests should have nothing to recommend it but its beauty. Other properties it possesses only in a secondary degree, and in Europe it will probably be confined to the pleasure-grounds of amateurs.

PLATE VIII.

A branch with leaves and fruit of the natural size.

ROCK CHESTNUT OAK.

QUERCUS PRINUS MONTICOLA. *Q. foliis obovatis acutis grossè dentatis, dentibus subæqualibus; fructu majusculo, cupulâ turbinatâ, scabrosâ; glande oblongâ.*

Quercus montana. WILLD.

THIS Oak is among the species which are not scattered promiscuously in the forests, but which grow only in particular situations and easily escape observation; hence, it is difficult to assign its limits with precision. It probably does not extend northward far beyond Vermont nor eastward beyond New Hampshire. I have never seen it in the district of Maine nor in Nova Scotia, and it is not mentioned in my father's botanical notes upon Lower Canada; it is likewise a stranger to the maritime parts of the Southern States. It is most frequently met with in the middle and in some parts of the northern sections, but is rarely mingled with other trees in the forests, and is found only on high grounds thickly strewn with stones or covered with rocks. Thus, it is often seen on the steep and rocky banks of the Hudson and on the shores of Lake Champlain, and still more frequently on the Alleghanies in Pennsylvania and Virginia. It forms nine-tenths of the growth on some parts of these mountains, but the soil is so meagre that it is thinly disseminated and does not exceed twenty or twenty-five feet in height and eight or ten inches in diameter. I made this observation particularly on the Dry Ridges, fifteen miles from Bedford.

In that part of Pennsylvania, as well as in Maryland and Virginia, it is known by the name of Chestnut Oak, and by that of Rock Oak on the banks of the Hudson and the shores of Lake Champlain to the distance of 400 miles from New York. Both are significant: the first, of a remarkable resemblance of

rossè dentatis,
atâ, scabrosâ;

na. WILLD.

attered pro-
n particular
is difficult to
not extend
eyond New
of Maine nor
er's botanical
to the mari-
requently met
ern sections,
rests, and is
th stones or
he steep and
Lake Cham-
es in Pennsylv-
e growth on
neagre that it
ty or twenty-
eter. I made
fifteen miles

Maryland and
k, and by that
the shores of
m New York.
resemblance of






Burns del.

L'Esperance sc.

Rock Chestnut Oak.
Quercus P.^{us} monticola.



the bark to that of the Chestnut; and the second, of the situations in which the tree is exclusively found. For this reason, and to avoid confounding it with the preceding and following species, which also grow in Virginia, I have blended the two denominations.

The beautiful appearance of this tree when growing in a fertile soil is owing equally to the symmetry of its form and to the luxuriance of its foliage. The leaves are five or six inches long, three or four broad, oval and uniformly denticulated, with the teeth more regular but less acute than those of the Chestnut White Oak. When beginning to open in the spring they are covered with a thick down; but when fully expanded they are perfectly smooth, whitish beneath, and of a delicate texture. The petiole is of a yellow color, which becomes brighter toward the fall.

The acorns are brown, of an oblong-oval shape, and sometimes an inch in length, a third part of which is contained in a spreading cup covered with loose scales; they are sweet-tasted, and are a favorite nourishment of wild and domestic animals.

The Rock Chestnut Oak is sometimes three feet in diameter and more than sixty feet high; but, as its growth is usually repressed by the poverty of the soil, it rarely attains these dimensions. In open, elevated situations it spreads widely and forms a head like that of the apple-tree. When the trunk exceeds a foot in diameter it is covered with a thick, hard, deeply-furrowed bark. At New York, and near the Alleghanies in Pennsylvania, this species of bark is esteemed the best for tanning. Only that of the secondary branches and of stocks less than six inches thick is employed. It is sold at New York for ten or twelve dollars a cord. The epidermis is strongly impregnated with the tanning principle, which in other species resides only in the cellular tissue.

The wood is reddish, like that of the White Oak; but its pores are more open, though its specific gravity is greater: pieces

of both species being thrown into water, the White Oak remains on the surface and the other at the bottom. Its staves are not used to contain spirituous liquors. At New York, and on the banks of the Hudson, it holds the next place to the White Oak in the construction of vessels. It is employed for the lower part of the frame, and oftener for the knees and the ribs. Pieces of White Oak suited to these objects are procured with difficulty; but the Rock Chestnut Oak, growing up in a continual controversy with the winds, is more frequently bent into the proper shape. For fuel, it is next in price to the Hickory. I have been told in several forges, especially those at the foot of the *North Mountain*, 200 miles from Philadelphia, that it is superior in this respect to every other species of its genus except the Live Oak

A tree like this, which grows in stony soils, in abrupt uninhabitable exposures, and whose bark and timber are so valuable, deserves the particular attention of American and European foresters. They should sow the acorns in the crevices of the rocks and wherever the soil is incapable of cultivation. Thousands of young plants already exist in the vicinity of Paris.

PLATE IX.

A branch with leaves and fruit of the natural size.

[See Nuttall's Supplement, vol. i. pp. 28, 33.]

Oak remains
aves are not
, and on the
e White Oak
or the lower
nd the ribs.
procured with
up in a con-
tly bent into
the Hickory.
at the foot of
ia, that it is
s genus except

abrupt unin-
re so valuable.
nd European
erevices of the
vation. Thou-
y of Paris.

size.

3, 33.]



P. H. Ravn, J. P.

1860

Yellow Oak

Quercus P^m acuminata.

Pl. 10.



100



YELLOW OAK.

QUERCUS PRINUS ACUMINATA. *Q. foliis longè petiolatis, acuminatis, subæqualiter dentatis; fructu mediocri; cupulâ subhæmisphæricâ.*

Quercus castanea. WILLD.

THE banks of the Delaware may be assumed as the northern limit of the Yellow Oak. It scarcely exists in the maritime parts of the Southern States, where I have seen only a few stocks near Two Sisters' Ferry on the Savannah in Georgia, and a single one on the Cape Fear, a mile from Fayetteville, in North Carolina. In the Middle and Western States, though more common, it is still rare in comparison with many other trees, and is sometimes lost sight of by the traveller for several days in succession. I have most particularly observed it on the small river Conestoga near Lancaster in Pennsylvania, on the Monongahela a little above Pittsburg, and in several small tracts near the Holston and Nolachuky in East Tennessee. In the Monography of American Oaks, my father takes notice of its existence in the country of the Illinois.

Near Lancaster this tree is called Yellow Oak, from the complexion of its wood; but in other parts of the United States it is confounded with the Chestnut White Oak and Rock Chestnut Oak, to which it bears some resemblance in its foliage.

The leaves are lanceolate, acuminate, regularly toothed, of a light green above and whitish beneath. The small acorns are contained in slightly-sealy cups, and are sweeter than those of any other species in the United States.

The Yellow Oak is a fine tree, seventy or eighty feet high and two feet in diameter, with branches tending rather to close round the trunk than to diffuse themselves horizontally. I invariably found it in valleys where the soil was loose, deep, and fertile. The bark upon the trunk is whitish, very slightly fur-

rowed, and sometimes divided into plates, like that of the Swamp White Oak. The wood is yellowish, though the tint is not bright enough to fit it for peculiar uses. Its pores are partly obliterated, irregularly disposed, and more numerous than those of any other American Oak: this organization must impair its strength and render it less durable than the Chestnut White Oak and the Rock Chestnut Oak.

As this tree is so thinly disseminated, it will not appear surprising that I should not have witnessed the application of its wood in the arts, or have found occasions of accurately appreciating its qualities. Its agreeable form and beautiful foliage render it proper for the embellishment of picturesque gardens.

PLATE X.

A branch with leaves and fruit of the natural size.

SMALL CHESTNUT OAK.

QUERCUS PRINUS CHINECAPIN. *Q. foliis obovatis, grosse dentatis, subtus glaucis; cupulâ hemispharica; glande ovatâ.*

Quercus prinoides. WILLD.

IN the Northern and Middle States this pretty little species is called Small or Dwarf Chestnut Oak, from the resemblance of its leaves to those of the Rock Chestnut Oak; as there is also a likeness between its foliage and that of the Chincapin, it is known in East Tennessee and in the upper part of the Carolinas by the name of the Chincapin Oak.

The Small Chestnut Oak is not generally diffused, but is rare in many places adapted to its constitution, and is usually found

that of the
the tint is
s are partly
than those
t impair its
tnut White

appear sur-
cation of its
ately appre-
tiful foliage
que gardens.

ze.

K.


e dentatis, subtilis

l.

oides. WILLD.

little species is
resemblance of
as there is also
Chincapin, it is
of the Carolinas

used, but is rare
is usually found



Name





Brown del.




W. C. C. 1/2

Small Chesnut Oak
Quercus P^{no} chincaipi.





**IMAGE EVALUATION
TEST TARGET (MT-3)**




**Photographic
Sciences
Corporation**

23 WEST MAIN STREET
WEBSTER, N.Y. 14580
(716) 872-4503

1.5 2.8
1.6 3.2
1.8 3.6
2.0 4.0
2.2 4.5
2.5 5.0

10
11
12
13
14
15



in particular districts, where, alone or mingled with the Bear Oak, it sometimes covers tracts of more than 100 acres. The presence of these species is a certain proof of the barrenness of the soil. I have particularly observed the Small Chestnut Oak in the vicinity of Providence in Rhode Island, of Albany in New York, of Knoxville in Tennessee, and on the Alleghany Mountains in Virginia. It grows spontaneously in the park of Mr. W. Hamilton, near Philadelphia.

This species, and another which is found in the Pine forests of the Southern States, rarely exceed thirty inches in height: they are the most diminutive of the American Oaks, and are mentioned only to complete the series.

The leaves of the Small Chestnut Oak are oval-acuminate, regularly but not deeply denticulated, of a light green above and whitish beneath. The acorns are enclosed for one-third of their length in scaly sessile cups; they are of middle size, somewhat elongated, similarly rounded at both ends, and very sweet.

Nature seems to have sought a compensation for the diminutive size of this shrub in the abundance of its fruit: the stem, which is sometimes no bigger than a quill, is stretched at full length upon the ground by the weight of the thickly-clustering acorns. United with the Bear Oak, which is of the same size and equally prolific, perhaps it might be cultivated with advantage for its fruit.

PLATE XI.

A branch with leaves and fruit of the natural size.

[See Nuttall's Supplement, vol. i. p. 33.]

LIVE OAK.

QUERCUS VIRENS. *Q. foliis perennantibus, coriaceis, ovato-oblongis, junioribus dentatis, vetustioribus integris; cupulâ turbinatâ, squamulis abbreviatis; glande oblongâ.*

THIS species, which is confined to the maritime parts of the Southern States, the Floridas, and Louisiana, is known only by the name of Live Oak. The climate becomes mild enough for its growth near Norfolk in Virginia, though it is less multiplied and less vigorous than in a more southern latitude. From Norfolk it spreads along the coast for a distance of 1500 or 1800 miles, extending beyond the mouth of the Mississippi. The sea-air seems essential to its existence, for it is rarely found in the forests upon the mainland, and never more than fifteen or twenty miles from the shore.

It is the most abundant, the most fully-developed, and of the best quality, about the bays and creeks and on the fertile islands which in great numbers lie scattered for several hundred miles along the coast. I particularly observed it on the islands of St. Simon, Cumberland, Sapelo, etc., between the St. John and the St. Mary, in an excursion of 400 or 500 miles in a canoe from Cape Canaveral in East Florida to Savannah in Georgia. I frequently saw it upon the beach, or half buried in the movable sands upon the downs, where it had preserved its freshness and vigor, though exposed during a long lapse of time to the fury of the wintry tempest and to the ardor of the summer's sun.

The Live Oak is commonly forty or forty-five feet in height and from one to two feet in diameter; but it is sometimes much larger. Mr. S., President of the Agricultural Society of Charleston, assured me that he had felled a trunk, hollowed by age, which was twenty-four feet in circumference. Like most other trees, it has, when insulated, a wide and tufted summit. Its

*ablongis, junio-
amulis abbre-*

parts of the
own only by
l enough for
ss multiplied
ude. From
1500 or 1800
issippi. The
rely found in
an fifteen or

ed, and of the
fertile islands
undred miles
islands of St.
John and the
a canoe from
n Georgia. I
n the movable
freshness and
to the fury of
er's sun.

feet in height
ometimes much
iety of Charles-
ollowed by age,
like most other
d summit. Its





QUERCUS virens.
Live Oak.



trunk is sometimes undivided for eighteen or twenty feet, but often ramifies at half this height, and at a distance has the appearance of an old Apple-tree or Pear-tree. The leaves are oval, coriaceous, of a dark green above and whitish beneath; they persist during several years, and are partially renewed every spring. On trees reared upon plantations, or growing in cool soils, they are one-half larger, and are often denticulated; upon stocks of two or three years they are commonly very distinctly toothed.

The acorns are of an elongated oval form, nearly black, and contained in shallow, grayish, pedunculated cups. The Indians are said to have expressed an oil from them to mingle with their food; perhaps, also, they ate the kernel, which, though not agreeable to the taste, is less rough and bitter than that of many other species. The fruit is sometimes very abundant, and it germinates with such ease that, if the weather is rainy at the season of its maturity, many acorns are found upon the trees with the radicle unfolded.

The bark upon the trunk is blackish, hard, and thick. The wood is heavy, compact, fine-grained, and of a yellowish color, which deepens as the tree advances in age. The number and closeness of the concentric circles evince the slowness of its growth. As it is very strong, and incomparably more durable than the best White Oak, it is highly esteemed in ship-building, and is consumed not only in the country which produces it but still more extensively in the Northern States. From its great durability when perfectly seasoned it is almost exclusively employed for the upper part of the frame. To compensate its excessive weight it is joined with the Red Cedar, which is extremely light and equally lasting.

The Live Oak does not afford large timber; but its wide and branching summit makes amends for this disadvantage by furnishing a great number of knees, of which there is never a sufficient quantity in the dock-yards.

The vessels built at New York and Philadelphia, with the upper frame of Red Cedar and Live Oak and the lower timbers of White Oak, are as durable as those constructed of the best materials in Europe. Brekel, whom I have already quoted, says that the best tree-nails are of Live Oak; but at present it is replaced, in the Southern States, by the Locust and the heart of the long-leaved Pine.

In the South, particularly at Charleston and Savannah, this species is used for the naves and felloes of heavy wheels, for which it is far superior to the White Oak; it is more proper, also, for screws and for the cogs of mill-wheels.

The bark is excellent for tanning, but is only accidentally employed.

Besides the Live Oak timber exported to England, great quantities are used in ship-building in the United States, particularly at Boston, New York, Philadelphia, and Baltimore. The consumption has trebled within twenty years, in consequence of the immense development of American commerce. Hence the price has doubled, and the species is rapidly diminishing. The clearing of the islands for the culture of cotton, which they yield of a superior quality, has contributed greatly to its destruction. It is already difficult to procure sticks of considerable size in the Southern States, and they are sought on the western coast of East Florida between the St. Mary and the St. John. From St. Augustine to the Cape the species is rarer; but we are informed that it abounds on the shores of West Florida, whither the English of the Bahama Islands resort for supplies.

As the Live Oak, from the peculiarities of its construction, is multiplied with difficulty, I cannot but consider its disappearance throughout the United States within fifty years as nearly certain. It will then be found only in the form of a shrub, like the *Quercus ilex*, which formerly skirted the southern coast of France and Italy.

ia, with the
lower timbers
of the best
ready quoted,
at present it
and the heart

avannah, this
heavy wheels, for
more proper,

y accidentally

England, great
ed States, par-
and Baltimore.
ears, in conse-
can commerce.
apidly diminish-
of cotton, which
d greatly to its
cks of consider-
e sought on the
Mary and the St.
species is rarer;
shores of West
islands resort for

its construction,
consider its dis-
thin fifty years as
in the form of a
irted the southern



Bacca del


Gibb.

Cork Oak.
Quercus suber.

Pl. 15.



Cyrtus



[Every individual that has access to Government authorities in successive administrations should urge the planting of this most valuable tree. Since our author penned the above remarks, great havoc has been committed on our Southern coast for want of Government protection. The tree is produced in TEXAS; but from there also its disappearance may be deemed almost certain.]

PLATE XII.

A branch with leaves and fruit of the natural size.

[See Nuttall's Supplement, vol. i. pp. 28, 29.]

 CORK OAK.

QUERCUS SUBER. *Q. foliis ovato-oblongis, indivisis, serratis, subtus glaucis; cortice rimoso, fungoso.*

THE Cork Oak grows naturally in the southern parts of France, in Spain, Portugal, Italy, and the States of Barbary, which are comprised between the 44th and 35th degrees of latitude. It rarely exceeds forty feet in height and three feet in diameter. Its leaves are evergreen, but the greater part of them fall and are renewed in the spring; they are ovate, thick, slightly toothed, of a light green on the upper surface and glaucous beneath. The acorns are rather large, oval, and half enclosed in a conical cup, and, being of a sweetish taste, are eagerly devoured by swine.

The wood is hard, compact, and heavy, but less durable than that of the Common European Oak, particularly when exposed to humidity. The worth of the tree resides in its bark, which

begins to be taken off at the age of twenty-five years. The first growth is of little value; in ten years it is renewed; but the second product, though less cracked than the first, is not thick enough for corks, and is used only by fishermen to buoy up their nets. It is not till the tree is forty-five or fifty years old that the bark possesses all the qualities requisite for good corks, and from that period it is collected once in eight or ten years. Its thickness is owing to the extraordinary swelling of the cellular tissue. It is better fitted than any other substance for the use to which it is appropriated, as its elasticity exactly adapts it to the neck of the bottle and its impenetrable structure refuses exit to the fluid.

July and August are the seasons for gathering it. For this purpose two opposite longitudinal incisions are made through the whole length of the trunk, and two others, transverse to the first, at the extremities; the bark is then detached by inserting a hatchet-handle shaped like a wedge. Great care must be taken not to wound the alburnum, as the bark is never renewed upon the injured parts. After being scraped, the bark is heated on its convex side and laden with stones, to flatten it and render it easier of transportation. In Catalonia it is cut into pieces and boiled to improve its quality. Its excellence consists in being compact, supple, and elastic, and it should be from fifteen to twenty lines thick.

The cork produced in France may be reckoned at 17,000 or 18,000 quintals, and when the sheets are smooth and even each quintal affords 7000 or 7500 corks eighteen lines long. The common price is a dollar and seventy cents a thousand, of which fifty cents must be allowed for the expense of making. It is computed that 110 or 115 millions of corks are annually consumed in France.

This tree would be an important acquisition to the United States, and would grow wherever the Live Oak subsists. The soil of the pine-barrens is in general too meagre to sustain its

vegetation; the bed of vegetable mould is in many places too thin, and the sand beneath so homogeneous that the roots of the Pines, instead of shooting downward, fold themselves back, as if repelled by a solid rock.

Both public and private interest require the inhabitants of the Southern coast, and especially the neighboring islands, to rear the Cork Oak about their plantations and in places that are unfit for the cultivation of cotton. It should also be introduced into West Tennessee, and with the more reason as the Vine may be cultivated there with success.

As the young stocks are injured by transplanting, they should be permanently fixed the second or third year. To favor their growth, the earth should be loosed about the roots two or three times a year; and, to render them tall and well shaped, the lower branches should be cut even with the trunk. Their vegetation is in this manner strengthened and the bark improved; without further attention they will continue to afford a valuable product during two or three centuries.

This tree has great advantages over several others which would likewise flourish in the same parts of the United States, such as the Olive and the White Mulberry. To fit their products for consumption, particularly that of the Mulberry, requires complicated processes, which can be performed with advantage only in populous countries. Hence the attempts made seventy or eighty years ago in Georgia to introduce the rearing of silkworms proved abortive; and the old White Mulberry-trees that still remain are monuments of that ill-calculated speculation. The bark of the Cork Oak, on the contrary, might be transported to the Northern States, or made into corks upon the spot by a simple operation performed by a single person with implements of which the price does not exceed two or three dollars.

[It gives us pleasure to record here that the acorn of the Cork Oak has been introduced into the Southern States by importa-

tions made at the Patent-Office. This first effort should be followed by others, till we are independent of foreign countries for an article of prime importance.]

PLATE XIII.

A branch with leaves and fruit of the natural size.

WILLOW OAK.

QUERCUS PHELLOS. *Q. foliis linear-lanceolatis, integerrimis, glabris, apice setaceo-acuminatis, junioribus dentatis lobatisve; cupulâ scutellatâ; glande subrotundâ, minimâ.*

THIS species, which is remarkable for its foliage, makes its first appearance in the environs of Philadelphia; but it is more common and of a larger size in Virginia, the Carolinas, and Georgia, where the milder temperature of the winter is evidently favorable to its growth. It is seen, however, only in the maritime parts of those States, and is a stranger to the inland districts, where the surface is mountainous and the climate more severe. From the analogy of soil and climate, it is probably found in Lower Louisiana; but I have never observed it beyond the Alleghanies in Kentucky and Tennessee.

The Willow Oak commonly grows in cool moist places, and, with the Tupelo, the Small Magnolia, the Red-flowering Maple, the Red Bay, and the Water Oak, it borders the swamps in the lower part of the Southern States. In these situations it attains its greatest expansion, which is fifty or sixty feet in height and from twenty to twenty-four inches in diameter. The trunk, even at an advanced age, is covered with a smooth bark, re-

ould be fol-
n countries

e.

is, glabris, apice
cutellatâ; glande

age, makes its
but it is more
Carolinâs, and
winter is evi-
er, only in the
to the inland
and the climate
climate, it is
never observed
nessec.

ist places, and,
owering Maple,
e swamps in the
ations it attains
et in height and
er. The trunk,
smooth bark. re-






H. J. Redouté del.

Remond'is

Willow Oak.
Quercus pubescens



markable for the thickness of its cellular tissue. The leaves are two or three inches long, of a light green, smooth, narrow, entire, and similar to those of the Willow, whence is derived the name of Willow Oak, which is used in every part of North America where the tree is known.

Though the Willow Oak, as I have just observed, is almost always seen in moist grounds, by an exception for which it is difficult to account it is sometimes found among the Live Oaks, near the sea, in the dryest and most sandy soils. At a distance it resembles the Live Oak in its shape and in its foliage, which persists during several years; but on a closer examination it is easily distinguished by the form of its leaves, which are shorter and much narrower, and by the porous texture of its wood.

The fruit of this species is rarely abundant; the acorns are of a dark brown color, small, round, very bitter, and contained in shallow cups lightly coated with scales; kept in a cool place they preserve the faculty of germination for several months.

The wood is reddish and coarse-grained. It is too porous to contain wine or spirituous liquor, and its staves are classed with those of Red Oak. The quantity, however, is small, as the tree is so little multiplied that alone it would not supply the consumption for two years. In some of the lower parts of Virginia, particularly in the county of York, it is found to possess great strength and tenacity and to split less easily than the White Oak; hence, after being thoroughly seasoned, it is employed for the felloes of wheels. These are the only uses to which it seems adapted, and for these it is less proper than the Post Oak and White Ash. On several plantations near Augusta, in Georgia, the fences are made partly of Willow Oak, which lasts only eight or nine years. As fuel, it is sold at the lowest price

PLATE XIV.

A branch with leaves and fruit of the natural size.

[See Nuttall's Supplement, vol. i. p. 26.]

LAUREL OAK.

QUERCUS IMBRICATA. *Q. foliis subsessilibus, ovali-oblongis, acutis, integerrimis, nitidis; glauca subhemisphaericâ.*

EAST of the Alleghanies this species is rare, and has received no specific name; west of the mountains, where it is more multiplied and has attracted more attention, it is called Jack Oak, Black Jack Oak, and sometimes, from the form of its leaves, Laurel Oak. The last denomination I have preserved as the most appropriate, though perhaps it is less common than the first.

I observed this tree for the first time in Pennsylvania, near Bedford, on the Juniata, upon the road from Philadelphia to Pittsburg; and it does not exist in the more northern States. I found it abundant only beyond the mountains, and particularly near Washington Court-house, and in some parts of Kentucky and Tennessee. From my father's observations, it appears to be more multiplied in the country of the Illinois than in the places I have just mentioned, and it is called by the French *Chêne à lattes*,—Lath Oak.

In the western parts of Pennsylvania and Virginia, small lawns, covered only with tall grass, are frequently seen in the forests, around which the Laurel Oak forms entire groves; insulated stocks are also found in cool, humid situations. It is probable, from its flourishing in open exposures, that it is most abundant in the country of the Illinois, which consists of immeasurable savannas stretching in every direction, to which the forests bear no sensible proportion.

The Laurel Oak is forty or fifty feet high and twelve or fifteen inches in diameter. Its trunk, even when old, is clad in a smooth bark, and for three-fourths of its height is laden with branches. It has an uncouth form when bared in the winter,

is, acutis, inte-

l has received
t is more mul-
led Jack Oak,
of its leaves,
served as the
mon than the

sylvania, near
Philadelphia to
thern States. I
nd particularly
ts of Kentucky
it appears to be
an in the places
French *Chêne à*

Virginia, small
tly seen in the
re groves; insu-
ions. It is pro-
that it is most
n consists of im-
on, to which the

twelve or fifteen
old, is clad in a
ght is laden with
ed in the winter,





Laurel Oak
Quercus imbricaria.

W. H. Sargent del.

Boquet 36




P. J. Rebouté del.

Reynolds co.

Upland Willow Oak
Quercus emerau.





but is beautiful in the summer when clad in its thick, tufted foliage. The leaves are long, lanceolate, entire, and of a light, shining green.

The wood is hard and heavy, though its pores are open. As the trunk is branchy and often crooked, it is considered, wherever I have observed it, as fit only for fuel; but my father, who first described it, says that the French of Illinois use it for shingles. Probably in that region it attains much greater dimensions; but in my opinion the want of better species only can account for its use. Its wood is inferior to that of the Willow Oak, which it nearly resembles.

This tree has no merit but its singular foliage; and it deserves the attention only of amateurs desirous of adorning their rural retreats with a variety of exotic trees.

PLATE XV.

A branch with leaves and fruit of the natural size.

 UPLAND WILLOW OAK.

QUERCUS CINEREA. *Q. foliis petiolatis, lanceolato-oblongis, acutis, integerrimis, subtus cinereo-pubescentibus; cupulâ scutellatâ; glande subhemisphærica.*

THE Upland Willow Oak is confined to the maritime parts of the Southern States. It is little multiplied in comparison with many other species, and is dispersed in small groups in the forests of White Pine. It is found also upon the sea-shore, and upon the islands, where it covers tracts of several acres still more barren than the main. But the stocks which grow in

these different situations are so different in appearance that they might easily be mistaken for distinct species: in the *pine-barrens* they are eighteen or twenty feet high and four or five inches in diameter, with the leaves entire, two and a half inches long, and whitish underneath; on the islands and on the shore of the continent, where the soil is extremely dry, as near Wilmington, N.C., they are only three or four feet in height, and the leaves are denticulated, are an inch in length, and persist for two years. I have ascertained that both varieties belong to the same species, by examining the sprouts of the larger stocks in the *pine-barrens*, of which the foliage is perfectly similar to that of the smaller ones on the shore.

The Upland Willow Oak is one of the abject trees that succeed the Pines on lands which have been cleared for cultivation and abandoned on account of their sterility. In these places, as in the *pine-barrens*, it is twenty feet in stature, and its trunk, crooked and covered with a thick bark, begins at a third of this height to divide by numerous ramifications. In the spring it is distinguished at a distance by the reddish color of its leaves and male aments. The acorns, which are contained in shallow cups, are round and blackish, with the base of a bright rose-color when freshly exposed. It is rare to meet with a tree which yields a quart of fruit.

The bark of this species, like that of the Black Oak, affords a beautiful yellow dye; but the tree is so small and so little multiplied that it is of no utility in this respect, nor even for fuel.

The *Quercus nana* of Willdenow is certainly the smaller variety of this species.

PLATE XVI.

A branch with leaves and fruit of the natural size. Fig. 1. A leaf of the smaller variety of the natural size.

t they
arrens
hes in
long,
of the
ngton,
leaves
years.
pecies,
arrens,
smaller

at suc-
ivation
places,
s trunk,
of this
ing it is
ves and
ow cups,
ose-color
e which

k, affords
so little
even for

smaller

leaf of the



H. J. Foster, Jr.

1900

Running Oak.
Quercus pumila.

Pl. 17



R. 17

C

h
d
W
G
sp
va
gr
la

sp
de
an
an
fe
gr
fo
th
de

di
pa
O
O
po
siz

RUNNING OAK.

QUERCUS PUMILA. *Q. foliis deciduis, lanceolatis, integerrimis, basi attenuatis, apice dilatatis; cupulâ scutellatâ; glande subhemisphericâ.*

Quercus sericea. WILLD.

THIS species, which is rarely more than twenty inches in height and two lines in diameter, is the smallest Oak hitherto discovered in the Old or the New World. Like the Upland Willow Oak, it is confined to the maritime parts of the Carolinas, Georgia, and the Floridas, where it is called Running Oak. It springs with that species in the *pine-barrens*, amid the numerous varieties of Whortleberry and other plants which overspread the ground wherever there is a little moisture in the soil and the layer of vegetable mould is a few inches thick.

The leaves of this dwarfish shrub are of a reddish tint in the spring and turn green as the season advances. When fully developed they are entire, smooth, of an elongated oval shape, and about two inches in length. The acorns are small, round, and similar to those of the Willow and Water Oaks; they are few in number, because the stem is burned to the surface of the ground, almost every spring, by the fire which is kindled in the forests to consume the dead grass. As this species belongs to the division whose fructification is biennial, the acorns are destroyed before they arrive at maturity.

My own observations, and those of Messrs. Bosc and Delille, distinguished botanists who resided several years in the southern part of the United States, have led me to consider the Running Oak as a distinct species, and not as a variety of the Willow Oak, as my father has treated it in his *Monography* of this important genus. It is hardly necessary to remark that from its size it can be interesting only to botanists.

[In situations where it is not liable to being burned, as described above, this plant is very prolific of acorns; and it has been suggested that it might be cultivated for its fruit. It sometimes covers tracts of more than 100 acres in extent, along with the Bear Oak. It is well worthy of cultivation in small villa gardens or miniature arboretums.]

PLATE XVII.

A branch with leaves and fruit of the natural size.

 BARTRAM OAK.


QUERCUS HETEROPHYLLA. *Q. foliis longè petiolatis, orato-lanceolatis, integris vel inæqualiter dentatis; glande subglobosâ.*

EVERY botanist who has visited different regions of the globe must have remarked certain species of vegetables which are so little multiplied that they seem likely at no distant period to disappear from the earth. To this class belongs the Bartram Oak. Several English and American naturalists who, like my father and myself, have spent years in exploring the United States, and who have obligingly communicated to us the result of their observations, have, like us, found no traces of this species except a single stock in a field belonging to Mr. Bartram, on the banks of the Schuylkill, four miles from Philadelphia. This is a flourishing tree, thirty feet in height and eight inches in diameter, and seems formed to attain a much greater development. Its leaves are of an elongated oval form, coarsely and irregularly toothed, smooth above, and beneath of a dark green.

s de-
has
It
long
small


ccolatis,

e globe
are so
riod to
Bartram
like my
United
e result
of this
Bartram,
delphia.
t inches
develop-
sely and
k green.





Bartram's Oak
Quercus heterophylla



Water Oak
Quercus aquatica

Pl. 12



THE
cup

La
lea
exi
S
hin
pre

[S
that
cord
tram

QUE
sub
sub

T
Rich
south
Geor
is so
alwa
V

The acorns are round, of a middle size, and contained in shallow cups lightly covered with scales.

I was at first disposed to consider this tree as a variety of the Laurel Oak, to which it bears the greatest affinity; but the leaves of that species are never indented, and not a stock of it exists within a hundred miles of Philadelphia.

Several young plants, which I received from Mr. Bartram himself, have been placed in our public gardens to insure the preservation of the species.

PLATE XVIII.

A branch with leaves and fruit of the natural size.

[See Nuttall's Supplement, vol. i. p. 24, by which it appears that this tree has been discovered near Cincinnati, Ohio. According to Meehan's Hand-Book of Trees, the specimen at Bartram's Garden is seventy feet high and six feet in circumference.]

WATER OAK.

QUERCUS AQUATICA. *Q. foliis obovati-cuneatis, basi acutis, summitate subintegris, varivae trilobis, glabris; cupulâ modicæ crateratâ; glande subglobosâ.*

THIS species first attracted my attention in the forests near Richmond in Virginia; it becomes more common in proceeding southward, and abounds in the lower parts of the Carolinas and Georgia and in East Florida. Under the name of Water Oak it is sometimes confounded with the Willow Oak, by which it is always accompanied in the ponds and narrow swamps enclosed

in the *pine-barrens*. It is inferior in size to the Willow Oak, and rarely exceeds forty or forty-five feet in height and twelve or eighteen inches in diameter. On full-grown trees the leaves are smooth, shining, and pyriform,—or broad and rounded at the summit and terminated in an acute angle at the base. In the severe climate of Virginia they fall with the first frost; but on the sea-shore of the Carolinas, Georgia, and Florida, they persist during two or three years. There is no Oak in the United States of which the foliage is so variable and so different from that of the tree on the young stocks, and on the sprouts from an old trunk or from the base of a limb that has been lopped: the leaves are commonly oval and deeply and irregularly toothed.

The acorns, which are contained in shallow, slightly-sealy cups, are brown, small, and extremely bitter: the largest tree rarely yields more than five or six quarts. Like those of the Willow Oak, when kept cool they preserve their fecundity for several months.

The bark upon the oldest trunks is smooth and very slightly furrowed; it is little used in tanning, either because it is inferior to that of the Spanish Oak or because the tree is less abundant.

The wood is very tough, but less durable and less esteemed by carpenters and wheelwrights than that of the White Oak and Chestnut White Oak.

As this species is destitute of interest, it will probably become extinct, like many others which are rapidly diminishing. In France it would flourish only in the southern departments.

PLATE XIX

A branch with leaves and fruit of the natural size.

[See Nuttall's Supplement, vol. i. p. 33.]

ak,
lve
ves
l at
In
but
they
the
erent
routs
been
regu-

scaly
t tree
of the
ty for

ightly
nferior
ndant.
ned by
k and

become
g. In
s.



Black Jack Oak
Quercus ferruginea



C
A
s
t
i
n
o
t
a
a
t
v
d
-
s
v
c
s
l
S
c
t
l
g
s

BLACK JACK OAK.

QUERCUS FERRUGINEA. *Q. foliis coriaceis, summitate dilatatis, retuso-subtrilobis, basi retusis, subtus rubiginoso-pulverulentis; cupulâ turbinatâ, squamis obtusis, scariosis; glande brevi ovatâ.*

Quercus nigra. WILLD.

I OBSERVED this species for the first time in the forests near Allentown and Cranberry, small towns of New Jersey, about sixty miles east of Philadelphia; but it is smaller and less multiplied than farther south. In New Jersey and Philadelphia it is called *Barren Oak*, and Black Jack Oak in Maryland and the more Southern States. I have adopted the last of these names only because it is the most generally used, and have changed the specific epithet *nigra*, because the name of the Black Oak is appropriated in the United States to the *Quercus tinctoria*.

This species is commonly found upon soils composed of red argillaceous sand mingled with gravel, and so meagre as to be totally exhausted by five or six crops when they are thought worthy of cultivation. Unhappily, from Baltimore to the borders of North Carolina—an extent of four or five hundred miles—the greater part of Maryland and Virginia consists of this soil. The whole of this interval, with the exception of the valleys and the swamps with their surrounding acclivities, is covered with forests impoverished by fire and the cattle that subsist in them during a great part of the year. They are composed principally of Yellow Pine, Post Oak, Black Oak, and Scarlet Oak. In the Carolinas and Georgia, where the soil gradually improves in retiring from the shore toward the mountains, the same trees form a band fifteen or twenty miles wide, between the *pine-barrens* and the forests of a more generous growth. In Kentucky and Tennessee the Black Jack Oak is seen only in the savannas, where it is widely diffused, and

where, preserved by the thickness of its bark and its insulated position, it survives the conflagrations that almost every year consume the grass: the fire, driven forward by the wind, has only time to devour its foliage. In the *pine-barrens* it grows chiefly on the edges of the *branch-swamps*, where the soil is a little stronger than is necessary for the Pines. With the Upland Willow Oak and the Scrub Oak, it possesses itself of the pine-lands that have been cleared for cultivation and abandoned on account of their sterility; and in these situations it is larger than in the forests.

The Black Jack Oak is sometimes thirty feet high and eight or ten inches in diameter, but commonly does not exceed half these dimensions. Its trunk is generally crooked, and is covered with a very hard, thick, and deeply-furrowed bark, of which the epidermis is nearly black and the cellular tissue of a dull red. The summit is spacious even in the midst of the woods. The leaves are yellowish, and somewhat downy at their unfolding in the spring; when fully expanded they are of a dark green above, rusty beneath, thick, coriaceous, and dilated toward the summit like a pear. In the autumn they turn reddish and fall with the earliest frost.

The oldest trees bear only a few handfuls of acorns, which are large and half covered with very scaly cups.

When the stock is more than eight inches in diameter the wood is heavy and compact, but coarse-grained and porous before it has reached this size. As it speedily decays when exposed to the weather, it is not used in the arts. It forms excellent fuel, and is sold at Philadelphia only one dollar a cord less than Hickory, while other kinds of wood are a third cheaper.

The species deserves the attention of amateurs in Europe for the singularity of its foliage.

PLATE XX.

A branch with leaves and fruit of the natural size.

lated
year
, has
grows
l is a
n the
elf of
aban-
s it is

eight
l half
covered
ch the
ll red.

The
ling in
above,
unmit
ith the

which

ter the
porous
s when
t forms
e a cord
heaper.
rope for



H. J. Rehnke del.


Fig.

Bear's Oak
Quercus banisteri.

11



12



BEAR OAK.

QUERCUS BANISTERI. *Q. foliis longe petiolatis, acutangulo-quinque-lobis, margine integris, subtus cinereis; cupulâ subtruncatâ; glande subglobosâ.*

THIS diminutive species is known in the Northern and Middle States by the names of Bear Oak, Black Scrub Oak, and Dwarf Red Oak, of which the first is the most common in New Jersey, where the shrub abounds. The Latin specific name was given it in honor of Banister, an English writer, by whom it was first introduced to notice.

I do not remember to have seen the Bear Oak in the lower part of the Southern States; it is common in those of the North, and still more so, I believe, in New York, New Jersey, and Pennsylvania. I have more particularly observed it at Fishkill, Catskill, and Albany in New York, near Paramus in New Jersey, and on that part of the Alleghanies in Pennsylvania which is crossed by the road to Pittsburg. It is never found insulated nor mingled with other shrubs in the forests, but always in tracts of several hundred acres, which it covers almost exclusively: a few stocks of the Dwarf Chestnut Oak are often united with it.

The ordinary height of the Bear Oak is three or four feet; but, when accidentally insulated and nourished by a vein of more fertile soil, it sometimes equals eight or ten feet. It usually grows in compact masses, which are traversed with difficulty, though no higher than the waist. As the individuals which compose them are of a uniform height, they form so even a surface that at a distance the ground appears to be covered with grass instead of shrubs.

The trunk, which is numerously ramified, is covered, like the branches, with a polished bark. It has more strength than

would be supposed from its size, which is rarely more than an inch in diameter. The leaves are of a dark green on the upper surface, whitish beneath, and regularly divided into three or five lobes. The acorns are small, blackish, and longitudinally marked with a few reddish lines: they are so abundant as sometimes to cover the branches. The lowly stature of the shrub renders it easy for bears, deer, and swine to reach them by lifting their heads or rising on their hindfeet.

The presence of this Oak is considered as an infallible index of a barren soil; and it is usually found on dry, sandy land mingled with gravel. It is too small to be adapted to any use; but near Goshen, on the road to New York, I observed an attempt to turn it to advantage by planting it about the fields for the purpose of strengthening the fences. Though this experiment seemed to have failed, I believe the Bear Oak might be usefully adopted in the Northern States for hedges, which might be formed from twenty to twenty-four inches thick, by sowing the acorns in three parallel furrows. They would be perfected in a short time, would be agreeable to the eye, and probably would be sufficient to prevent the passage of horses and cows. Hedges of the European thorn would doubtless be preferable; but they require a good soil and more labor than can at present be afforded in America: those that exist in the neighborhood of Philadelphia are left in a condition which would give a very unfavorable opinion of the farmer on whose lands they were seen in the North of France.

As the Bear Oak grows on the most sterile soils and resists the most intense cold and impetuous winds, perhaps it might serve to shelter the infancy of other more valuable trees in such exposures. The want of some protection is the greatest obstacle to the success of plantations on the downs, as I was told, near the Hague upon the coast of Holland.

Proprietors of large estates, who are addicted to the chase, might find this species and the Dwarf Chestnut Oak convenient

an
per
or
ally
me-
rub
lift-

dex
land
use;
d an
fields
s ex-
might
which
t, by
d be
, and
orses
ess be
than
n the
which
whose

resists
might
n such
st ob-
s told,

chase,
renient




W. A. S. 187

Barrens Scrub Oak.

Quercus coccinea





Q
th
a
in
vi
wh
nu
sec
un
mi
ter
rel
sm
dec
pe

for copses: they would afford nourishment to the game during several months in the year, and would allow the sportsman a fair aim at the birds as they rose upon the wing.

PLATE XXI.

A branch with leaves and fruit of the natural size.

BARRENS SCRUB OAK.

QUERCUS CATESBÆI. *Q. foliis brevissimè petiolatis, basi angustatis, acutis, subpalmato lobatis, lobis interdùm subfalcatis; cupulâ majusculâ; squamis marginalibus introflexis; glande brevi ovatâ.*

ACCORDING to my own observations, this species is confined to the lower part of the Carolinas and Georgia. I first saw it a few miles south of Raleigh, N.C., latitude 35° 40'. It grows in soils too meagre to sustain any other vegetation, such as the vicinity of Wilmington, N.C., where the light movable sand is wholly destitute of vegetable mould. It is the only species multiplied in the *pine-barrens*, and from this circumstance it seems to have derived its name.

In traversing these forests, I nowhere saw the Scrub Oak more uniformly disseminated than between Fayetteville and Wilmington, an interval of sixty miles, where it forms nearly one-tenth of the woods: the Pines themselves, throughout the barrens, are scattered at the distance of fifteen or twenty feet.

The foliage of this tree is open, and its leaves are large, smooth, thick, and coriaceous toward the close of summer, deeply and irregularly lacinated, and supported by short petioles. With the first frost they change to a dull red, and fall

the ensuing month. The acorns are pretty large, of a blackish color, and partly covered with a fine gray dust, which is easily rubbed off between the fingers; they are contained in thick cups swollen toward the edge, and distinguished from all other species by having the upper scales bent inward. The oldest trees alone are productive, and their fruit never exceeds a few handfuls.

In the winter it is difficult to distinguish the Scrub Oak from the Black Jack Oak, which it nearly resembles. Like that, it is crooked, ramified at the height of two or three feet, and covered with a thick, blackish, deeply-furrowed bark: it is, besides, perfectly similar in the color, texture, and weight of its wood. At Wilmington the Scrub Oak is the best fuel, and is sold separately; but, notwithstanding its abundance in this district, it is insufficient for the supply of the inhabitants. Its size alone would exclude it from use in the arts.

The general character of this tree forbids the hopes of advantage that might be conceived from its flourishing upon the most sterile soils.


PLATE XXII.

A branch with leaves and fruit of the natural size.

ackish
easily
k cups
er spe-
t trees
hand-

k from
that, it
et, and
: it is,
t of its
, and is
in this
ts. Its

s of ad-
upon the



Quercus filicata

Quercus filicata

Spanish Oak.
Quercus filicata.





SPANISH OAK.

QUERCUS FALCATA. *Q. foliis longè petiolatis, subpalmato-lobatis, subtùs eximie tomentosis, lobis falcatis; cupulâ crateriformi; glande subglobosâ.*

Quercus elongata. WILLD.

THIS species, like the Black Jack Oak, begins to show itself in New Jersey, near Allentown, about sixty miles from Philadelphia. But even at this distance it is smaller than in the immediate vicinity of the city, where it acquires its perfect development and where its leaves exhibit their appropriate form. Farther south it is constantly found among the most common trees in the forests. I have observed that it is less multiplied near the mountains and in the country beyond them. In Delaware, Maryland, and Virginia, it is known only by the name of Spanish Oak, and in the Carolinas and Georgia by that of Red Oak. In an old English work which I found in the library of Charleston it is said to have been called Spanish Oak by the first settlers, from the resemblance of its leaves to those of the *Quercus velani* which grows in Spain. Whether this etymology is just or not I am unable to say; but it is unknown to the inhabitants who have adopted the name. The denomination of Red Oak, which is used only in the more Southern States, was probably given it on account of the great analogy between its wood and that of the species thus called in the Northern and Middle States, where the Spanish Oak is much less common than in the South.

This tree is more than eighty feet in height and four or five feet in diameter. Its leaves are very different on different individuals: thus, in New Jersey, where the tree is only thirty feet high and four or five inches thick, they are three-lobed, except a few on the summit, and not falcated as on the large stocks in

the Southern States. On young plants, and on the lower branches of the most vigorous stocks growing in moist and shaded situations, they are also trilobed; and on the upper limbs they are more acutely laciniated, with the sections more arching, than those represented in the figure. This remarkable difference led my father to describe as a distinct species, under the name of *Quercus triloba*, the individuals whose foliage had not acquired its perfect form. Sometimes on the sprouts of trees that have been felled the leaves are deeply denticulated at right angles to the main rib. One of their constant characters is a thick down upon the lower side of the leaf and upon the young shoots to which they are attached.

The acorns are small, round, of a brown color, and contained in slightly-scaly cups supported by peduncles one or two lines in length. They resemble those of the Bear Oak, and, like them, preserve for a long time the faculty of germination.

The bark upon the trunk is blackish and deeply furrowed, with a cellular tissue of middling thickness. The wood is reddish and coarse-grained, with empty pores and all the characteristic properties of the species known in commerce by the general name of Red Oak: hence its staves are fit only to contain molasses and provisions and dry-goods. I have been told that in the West Indies the Red Oak staves from the Southern States, where this species abounds, are the most esteemed; from which it seems probable that its wood is better than that of the Red, Scarlet, and Black Oaks that furnish almost all the Red Oak staves from the Northern and Middle States: this superiority, however, is not sufficiently marked to occasion a difference in the price.

From its want of durability, the Spanish Oak is less esteemed than the White Oak, the Port Oak, and other species of annual fructification. It is rarely employed in building, and is used by wheelwrights only at Baltimore, where it is preferred to the White Oak for the felloes of large wheels.

The principal merit of the Spanish Oak, which gives it a superiority over most other species in the United States, resides in its bark. This is preferred for tanning coarse leather, which it renders whiter and more supple; it is consequently sold at Philadelphia and Wilmington a fourth dearer than that of the other Oaks: the leather is said to be improved by the addition of a small quantity of the bark of the Hemlock Spruce.

The Spanish Oak is adapted to the climate of the centre of France, if we may judge from its multiplication in the nurseries and in the gardens of amateurs. The stocks that have sprung from the acorns which I sent home during my residence in America bear as yet only three-lobed leaves; but they will become falcated at a maturer age.

From the inferiority of its wood, this species would not, in my opinion, deserve a place in our forests, though its bark should prove equal to that of the European Oak. But in the Southern States, when some species of trees are to be multiplied in preference to others, the Spanish Oak alone should be spared among the Red Oaks, as, besides its superiority in other respects, it has the advantage of flourishing on lands of a middling quality, such as compose a large part of that section of the United States.

PLATE XXIII.

A branch with leaves and fruit of the natural size.

BLACK OAK.

QUERCUS TINCTORIA. *Q. foliis profundè sinuosis, subtus pulverulentis; cupulâ turbinatâ, squamosâ; glande brevi oratâ.*

EXCEPT the district of Maine, the northern part of New Hampshire, Vermont, and Tennessee, this species is found throughout the United States on both sides of the Alleghanics, and it is everywhere called Black Oak. It is more abundant in the Middle States and in the upper part of the Carolinas and Georgia than on the Southern coast.

The Black Oak flourishes in a poorer soil than the White Oak. In Maryland and certain districts of Virginia, where the soil is lean, gravelly, and uneven, it is constantly united in the forests with the Scarlet, Spanish, and Post Oaks, and the Mockernut Hickory, with which the Yellow Pine is also frequently mingled.

This Oak is one of the loftiest trees of North America, being eighty or ninety feet high and four or five feet in diameter. The leaves are large, deeply laciniated, and divided into four or five lobes: they resemble those of the Scarlet Oak, but have less deep and open sinuses, are less shining, of a duller green, and in the spring and during a part of the summer have their surface roughened with small glands, which are sensible to the eye and to the touch. The same appearance is observed on the young shoots. I have remarked that the leaves of the young stocks change in the autumn to a dull red, and those of the old trees to yellow, beginning with the petiole.

The trunk is covered with a deeply-furrowed bark of middling thickness, and always of a black or very deep brown color, whence probably is derived the name of the tree. Northeast of Pennsylvania the complexion of the bark is the only cha-


ntis;

New
ound
nies,
nt in
and

White
e the
n the
ecker-
ently


being
meter.
our or
have
green,
e their
to the
on the
young
the old

iddling
color,
rtheast
ly cha-





Chêne quercitron.
Quercus tinctoria



racter by which it can be distinguished from the Red, Scarlet, and Gray Oaks, when the leaves are fallen. Farther south this character is not sufficient to distinguish it from the Spanish Oak, the bark of which is of the same color; and recourse must be had to the buds, which, on the Black Oak, are longer, more acuminate, and more scaly. All doubt may be removed by chewing a bit of the cellular tissue of each; that of the Black Oak is very bitter and gives a yellow tinge to the saliva, which is not the case with the other.

The wood is reddish and coarse-grained, with empty pores: it is, however, more esteemed for strength and durability than that of any other species of biennial fructification. At Philadelphia it is employed, for want of White Oak, in building; and the farmers of the Northern States, with false economy, substitute it in the place of the White Oak for fences.

As this species is abundant in the Northern and Middle States, it furnishes a large proportion of the *Red Oak staves* exported to the colonies or employed at home to contain flour, salted provisions, and molasses.

The bark is extensively used in tanning, as it is easily procured and is rich in tannin. The only inconvenience which attends it is imparting a yellow color to the leather, which must be discharged by a particular process, to prevent its staining the stockings: it is a great error to assert that this color augments its value.

From the cellular tissue of the Black Oak is obtained the *quercitron*, of which great use is made in dyeing wool, silk, and paper-hangings. According to several authors who have written on this subject, and, among others, Dr. Bancroft, to whom we are indebted for this discovery, one part of quercitron yields as much coloring-matter as eight or ten parts of woad. The decoction is of a brownish yellow, which is rendered deeper by alkali and lighter by acids. A solution of alum causes a small portion of the coloring-matter to fall in a deep yellow precipitate; solu-

tions of tin afford a more abundant precipitate of the same color but of a much brighter hue.

To dye wood, it is sufficient to boil the quercitron with an equal weight of alum: in dipping the stuff, the deepest shade is given first, and afterward the straw-color; to animate the tint, the stuff may be passed, in coming out of the dye, through water whitened with a little washed chalk. A brighter color is obtained by means of a solution of tin. Quercitron may be substituted for woad in giving all the shades of yellow to silk; the proportion is one part by weight to twelve parts of silk. In the advertisements of Philadelphia for February, 1808, this substance is rated at forty dollars a ton, and from that city chiefly it is exported to Europe.

Though the wood of the Black Oak is of a better quality than that of the Scarlet, Spanish, Red, Pin, Gray, Willow, and Water Oaks, which are all comprehended under the name of *Red Oak*, it is much inferior to that of the European Oak. But its stature, the rapidity of its growth in the coldest climates and on the most indifferent soils, and, above all, the value of its bark in dyeing, recommend it powerfully to the notice of European foresters.

[This Oak produces a small acorn, sometimes striped with bars of yellow and brown, in a very deep cup. The leaves figured are such as can always be found on the young, lower, vigorous sprouts; the leaves on old trees are deeply lobed,—almost as much so as those of the Scarlet Oak. Upon these, as well as on those of the Red and Scarlet, are found smooth, round, light excrescences, called oak-apples, from one to two inches in diameter, formed by the extension of the cuticle of the leaf. They are produced by an insect, *Cynis confluentus*, which punctures the healthy leaf and deposits therein an egg, about which the apple forms. “A single grub,” says Harris, “lives in the kernel, becomes a chrysalis in the autumn, when

e color

with an
made is
the tint,
a water
obtained
stituted
he pro-
In the
is sub-
chiefly

quality
ow, and
name of
k. But
climates
value of
ctice of

ped with
e leaves
g, lower,
lobed,—
these, as
smooth,
e to two
uticle of
influentus,
n an egg,
s Harris,
nn, when




Bosa del

Gibbert Sculp

Scarlet Oak.
Quercus coccinea.





C
is
a
a
c
a
if
V
S
S
s
b
d
th
b

the oak-apple falls from the tree, changes to a fly in the spring, and makes its escape out of a small round hole which it gnaws through the kernel and shell.]

PLATE XXIV.

A leaf of the natural size.

NOTE.—*The small branch with the acorns belongs to the Scarlet Oak.*

 SCARLET OAK.

QUERCUS COCCINEA. *Q. foliis longè petiolatis, oblongis, profundè sinuatis, glabris; lobis dentatis, acutis; cupulâ insigniter squamosâ; glande brevi ovalâ.*

THE Scarlet Oak is first seen in the vicinity of Boston; but it is less multiplied than in New Jersey, Pennsylvania, Virginia, and the upper part of the Carolinas and Georgia, where it forms a part of the forests that are still standing: it is much less common in the lower parts of these States, which, as I have already observed, produce nothing but Pines. I have not seen it in the district of Maine, the States of New Hampshire and Vermont, nor beyond Utica in Genesee. In the Northern States it is confounded with the Red Oak, and in those of the South with the Spanish Oak. The name of Scarlet Oak was given it by my father, and, though not in use by the inhabitants, it will probably be adopted, as the tree is evidently a distinct species.

This is a vegetable of more than eighty feet in height and of three or four feet in diameter. The leaves, which are supported by long petioles, are of a beautiful green, smooth, shining on

both sides, and lacinated in a remarkable manner, having usually four deep sinuses very broad at the bottom. They begin to change with the first cold, and, after several successive frosts, turn to a bright red, instead of a dull hue like those of the Red Oak. At this season the singular color of the foliage forms a striking contrast with that of the surrounding trees, and is alone a sufficient inducement to cultivate the tree for ornament.

The acorns are large, somewhat elongated, similarly rounded at both ends, and half covered with scaly cups. As this fruit varies in size with the quality of the soil, it is difficult to distinguish it from that of the Black Oak; the only constant difference is in the kernel, which is yellowish in the Black Oak and white in the Scarlet.

The wood of this Oak is reddish and coarse-grained, with open pores. As it decays much more rapidly than the White Oak, it is employed by the builder and wheelwright only from necessity or economy. It is poor fuel, and is used principally for staves. In the Middle States a large part of the *Red Oak* staves are furnished by this species.

The bark, though very thick and generally employed in tanning, is in no respect preferable to that of the Gray and Red Oaks.

That this tree will flourish in France is shown by an example at Rambouillet, where it makes part of a beautiful plantation forty-five feet in height, formed in 1786, of species sent home by my father soon after his arrival in the United States. It is to be regretted that so fine a tree, which is so well adapted to our soil, should afford such indifferent wood that we cannot recommend its introduction into the forests of Europe nor its preservation in those of the United States.

[The leaves of the Scarlet Oak so much resemble those of the Black that a novice will easily confound them. In the Black

having
They
cessive
hose of
foliage
g trees,
tree for

rounded
his fruit
t to dis-
tant dif-
ack Oak

ed, with
e White
nly from
incipally
Red Oak

loyed in
and Red

example
lantation
home by

It is to
ed to our
ot recom-
ts preser-

ose of the
the Black




h. s. d.

wood. s. p.

Gray Oak.
Quercus ambigua





Q
f
f
Q
a
n
t
n
f
o

the leaves are broader and fuller toward the ends, larger, more nearly entire, and usually darker and thicker. In the Scarlet Oak the leaves are fuller toward the middle, smaller, thinner, more deeply cut, and of a lighter and livelier color; on small plants especially more deeply cut, but sometimes running down along the footstalk: the footstalk is more slender and longer, and both surfaces and the axils of the veins are always less downy. In the fall the rich and beautiful deep scarlet color, red dotted with crimson, or orange-scarlet, of the foliage of this Oak, make it unmistakable from any other species.]

PLATE XXV.

A leaf of the natural size.

NOTE.—The acorns in this plate belong to the *Black Oak*.

 GRAY OAK.

QUERCUS BOREALIS. *Q. foliis serratis, glabris, sinibus subacutis; cupulâ subscutellatâ; glanâ turpide ornatâ.*

THE Gray Oak appears, by my father's notes, to be found farther north than any other species in America: in returning from Hudson's Bay he saw it on the St. Lawrence between Quebec and Malabie, in latitude 47° 50'. Under that parallel, and near Halifax in Nova Scotia, where I first observed it, it is not more than forty feet high; and, though the bloom is annual, the winter is so rigorous and long that the fruit is said to be matured only once in three or four years. Three degrees farther south, in Maine and New Hampshire, and on the shores of Lake Champlain in Vermont, it is more multiplied, and is

fifty or sixty feet in height and eighteen inches in diameter. It is called by the inhabitants Gray Oak; but it has been confounded by botanists with the Red Oak, to which it bears a close analogy in its foliage, as it does to the Scarlet Oak in its fruit: on this resemblance I have founded the Latin specific name *ambigua*.

The leaves are large, smooth, and deeply sinuated at right angles to the main rib. The acorns are of the middle size, rounded at the end, and contained in scaly cups.

The wood is similar to that of the other species included under the common name of Red Oak. Its coarse and open texture renders it unfit for any use except to contain dry wares; but, in districts where Oak wood is rare, recourse is had for other purposes to several species of inferior quality, which are still superior to the Birch, the Beech, and the Pine. Thus, the Gray Oak is employed for the knees of vessels and for wheelwrights' work: it is even preferred to the Red Oak, as being stronger and more durable.

This tree is without interest, as the regions in which it grows possess other species in every respect more valuable, such as the White Oak, the Swamp White Oak, and the Rock Chestnut Oak.

PLATE XXVI.

A branch with leaves and fruit of the natural size.

diameter.
been con-
s a close
ts fruit:
e name

at right
dle size,


included
nd open
tain dry
course is
quality,
the Pine.
s and for
Oak, as

it grows
ch as the
nut Oak.



Pin Oak
Quercus palustris.





PIN OAK.



QUERCUS PALUSTRIS. *Q. foliis profundè sinuatis, glabris, sinibus latis; fructu parvo; cupulâ scutellatâ, levi; glande subglobosâ.*

THIS species, like the preceding, grows in Massachusetts, but is less common than in the vicinity of New York, in New Jersey, Pennsylvania, and Maryland. I saw it abundant beyond the mountains near Pittsburg, in Ohio, and in East Tennessee; and my father found it multiplied in the country of Illinois: I feel assured that it does not exist in Maine, Vermont, and the Southern States. It is called Pin Oak in the lower part of New York and in New Jersey, and Swamp Spanish Oak in Pennsylvania, Delaware, and Maryland. The last of these denominations is sufficiently appropriate; but I have preferred the second, because it is less liable to mistake, and is indicative of a characteristic arrangement of the branches.



The Pin Oak is a tall tree, which grows constantly in moist places and prefers the swamps enclosed in the forests. In these situations it is frequently more than eighty feet high and three or four feet in diameter. Its secondary branches are more slender and numerous than is common on so large a tree, and are intermingled so as to give it at a distance the appearance of being stuffed. This singular disposition renders it distinguishable at first sight in the winter, and is perhaps the cause of its being called Pin Oak. The leaves are smooth, of a pleasing green, supported by long petioles, deeply lacinated, and very similar to those of the Scarlet Oak, from which they differ principally in their proportions. The acorns are small, round, and contained in flat shallow cups, of which the scales are closely applied one upon another.

The bark upon the oldest trunk is scarcely cracked, and consists almost wholly of a very thick cellular tissue. The wood is





**IMAGE EVALUATION
TEST TARGET (MT-3)**



**Photographic
Sciences
Corporation**

23 WEST MAIN STREET
WEBSTER, N.Y. 14590
(716) 873-4503

15 128
18 132
22 136
25 140
20 144
18 148

10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50

coarse-grained, with the pores open and larger than those of the Scarlet and Red Oaks. Though stronger and more tenacious than those species, it is little esteemed for durability. It is used for the shafts of mill-wheels when White Oak of sufficient dimensions cannot be procured; it is also sometimes, though rarely, made into staves, as the species is little multiplied compared with the Scarlet, Red, and Black Oaks.

The Pin Oak in its youth assumes an agreeable pyramidal shape, and its light elegant foliage contributes greatly to its beauty. It deserves a conspicuous place in parks and gardens. It should never be deprived of its interior branches. The most beautiful stock of this species with which I am acquainted in Europe is in a garden near Antwerp; it was about twenty feet high in 1804, and its brilliant and vigorous vegetation proved how well the soil and climate were adapted to it.

PLATE XXVII.

A branch with leaves and fruit of the natural size.

RED OAK.

QUERCUS RUBRA. *Q. foliis longe petiolatis, glabris, obtuse sinuatis; cupulâ scutellatâ, sublævi; glande subovatâ.*

NEXT to the Gray Oak, this species is found in the highest latitude of all the American Oaks, and is one of the most common species in the Northern States and in Canada. Farther south, particularly in the lower part of New York, in New Jersey, the upper districts of Pennsylvania, and along the whole range of the Alleghanies, it is nearly as abundant as the Scarlet

of the
acious
s used
imen-
arely,
pared

amidal
to its
ardens.
e most
ted in
ty feet
proved

; *cupulā*

highest
ost com-
Farther
in New
e whole
Scarlet





Red Oak.
Quercus rubra



and Black Oaks; but it is much less common in Maryland, Lower Virginia, and the maritime parts of the Carolinas and Georgia. This remark confirms an observation which I have often made,—that its perfect development requires a cool climate and a fertile soil. It is universally known by the name of Red Oak, except near Lancaster in Pennsylvania, where it is sometimes confounded with the Spanish Oak.

The Red Oak is a tall, wide-spreading tree, frequently more than eighty feet high and three or four feet in diameter. Its leaves are smooth and shining on both sides, large, deeply lacinated, and rounded at the base. They are larger and have deeper and narrower sections on the young stocks than on the middle or the summit of the full-grown tree: these last resemble the leaves of the Spanish Oak, which, however, are always downy beneath, while those of the Red Oak are perfectly smooth. In the autumn they change to a dull red, and turn yellow before they fall.

The acorns are very large and abundant, rounded at the summit, compressed at the base, and contained in flat cups covered with narrow, compact scales. They are voraciously devoured by wild animals, and by the cows, horses, and swine which are allowed to range in the woods after the herbage has perished.

The wood is reddish and coarse-grained, and the pores are often large enough for the passage of a hair; it is strong but not durable, and is the last among the Oaks to be employed in building. Its principal use is for staves, which at home are used to contain salted provisions, flour, and such dry wares as are exported to the islands, and, in the colonies, to receive molasses and sugar.

The bark consists of a very thin epidermis and a very thick cellular tissue. It is extensively used in tanning, but is less esteemed than that of the Spanish, Black, and Rock Chestnut Oaks.

The Red Oak was one of the earliest American trees intro-

duced into Europe. Large stocks are found on the estate of Duhamel, which yield seed abundantly and even multiply naturally; but the quality of its wood is so inferior that I cannot recommend its propagation in our forests.

[The Red is the most northern of the Oaks, Dr. Richardson reporting it as far north as Saskatchewan and the rocks of Lake Namakeen. Though its usefulness is not great, its beauty is unsurpassed, as are also its dimensions, which give an idea of nobility and strength. It grows rapidly from stoles, sometimes six feet in a season.]

PLATE XXVIII.

A branch with leaves and fruit of the natural size.

ADDITIONS TO THE OAKS.

IN the *Flora Americae Septentrionalis* of Pursh, published in England in 1814, the following species of Oak are added to those which I have described:—

QUERCUS MARITIMA. *Q. foliis perennantibus, coriaceis, integerrimis, glabris, basi attenuatis, apice mucronatis; cupulâ scutellatâ; glauca subrotundâ.*

A shrub from three to eight feet high, found on the sea-coast in Virginia and Carolina. I consider it as a variety of the Willow Oak, *Quercus phellos*.

QUERCUS MYRTIFOLIA. *Q. foliis perennantibus, coriaceis, oblongis, integerrimis, glabris, utrinque acutis, supra nitidis, margine revolutis.*

This species, of which Mr. Pursh appears to have seen neither the blossoms nor the acorns, escaped my researches: perhaps it

is the variety of the Water Oak which I found among the Live Oaks, and which preserves its leaves for three or four years.

QUERCUS HEMISPHERICA. *Q. foliis perennantibus, oblongo-lanceolatis, trilobis sinuatisque, lobis mucronatis, utrinque glabris.* Willd.

Mr. Pursh has inserted this species from Willdenow, and believes it to be a variety of the Water Oak, *Quercus aquatica*.

QUERCUS NANA. *Q. foliis cuneiformis, glabris, apice trilobis, basi subsinuatis, lobis divaricatis, mucronatis, intermedio majore; cupulâ scutellatâ.*

According to Mr. Pursh, this species is a low-growing shrub, distinct from the Water Oak, *Quercus aquatica*.

QUERCUS DISCOLOR. *Q. foliis oblongis, pinnatifido-sinuatis, subtus pubescentibus, lobis oblongis, dentatis, setaceo-mucronatis; cupulâ turbinatâ.*

This species of Mr. Pursh I consider as a variety of the *Quercus tinctoria*.

OAKS found in New Spain by Messrs. Humboldt and Bonpland, and described in their "Nova Genera et Species Plantarum." PARIS, 1816.

QUERCUS CONFERTIFOLIA. *Q. ramulis abbreviatis; foliis brevissime petiolatis, confertis, lanceolatis, acuminatis, mucronato-aristatis, integerrimis, coriaceis; margine subreflexis, subtus pubescentibus; fructibus subgeminis, sessilibus.*

This tree is ten or twelve feet in height: it is evergreen, grows in the temperate and mountainous regions of New Spain, between Guanajuato and Santa Rosa, and fructifies in September.

QUERCUS CRASSIPES. *Q. ramulis tuberculosis; foliis breviter petiolatis, lanceolato-oblongis, mucronatis, basi rotundatis, integerrimis, coriaceis, sultus cinereo-tomentosis; fructibus pedunculatis subgeminis; pedunculis incrassatis; cupulis subturbinatis.*

This tree is about twenty feet high: it is found on the low mountains of New Spain, near Santa Rosa, and fructifies in September and October.

QUERCUS MEXICANA. *Q. ramulis foliisque, subtus stellatim pubescentibus, supra nitidis, linearis-oblongis, acutis, submucronatis, subcordatis, undulato-subsinuatis, subcoriaceis; fructibus solitariis, breviter pedunculatis; cupulis cyathiformibus.*

This species rises from fifteen to twenty feet: it is very

abundant between Acapulco and the city of Mexico, near Mojonera, Quajiniquilapa, and Chilpalcingo, and is also found near Moran, Regla, Guanajuato, and Santa Rosa: it fructifies in September.

QUERCUS LANCEOLATA. *Q. ramulis tuberculatis; foliis oblongo-lanceolatis, utriusque acutis, undulato-repandis, coriaceis, supra nitidis, subtus stellatim pubescentibus; fructibus subternis, brevissime pedunculatis; cupulis cyathiformibus.*

This tree equals and sometimes exceeds twenty feet in height: it abounds in the temperate regions of Mexico, between Moran and Santa Rosa, where it forms immense forests: it fructifies in September.

QUERCUS TRIDENS. *Q. ramis lævibus; foliis oblongis, basi rotundatis, apice cuspidato-tridentatis, membranaceis, supra pubescentibus, subtus tenuiter cinereo-tomentosis; fructibus ternis aut quinis, breviter pedunculatis.*

This tree rises from ten to twenty feet: it grows in the mountains near Moran in Mexico, and fructifies in May.

QUERCUS LAURINA. *Q. ramulis glabris, foliis oblongis, acuminatis, basi subrotundatis; apicem versus subdentatis, coriaceis, glabris, nitidis; fructibus solitariis aut ternis, sessilibus; cupulis cyathiformibus.*

This is a large tree, which resembles the Laurel, and attains the height of forty feet: it is found in the temperate parts of New Spain, in the environs of Pachuca, Totonilco, and Grande: it flowers in May.

QUERCUS REPANDA. *Q. fruticosa, procumbens; ramulis foliisque subtus albidotomentosis, subsessilibus, oblongis, obtusiusculis, basi inæqualibus, sinuato-repandis coriaceis; fructibus subsolitariis, sessilibus.*

This is a shrub about two feet in height: it grows in moist, shady situations, between Real del Monte and Moran, and flowers in May.

QUERCUS DEPRESSA. *Q. fruticosa procumbens; ramulis pubescentibus; foliis sempervirentibus, oblongis, acutis, basi rotundatis, argute et remote dentatis, rigidus, glabris, nitidis; fructibus geminis aut ternis, breviter, pedunculatis.*

This species is an evergreen shrub, numerously ramified, and only one or two feet in height: it abounds in the same situations with the preceding, and flowers in May and June.

QUERCUS CHRYSOPHYLLA. *Q. ramulis sulcatis, pubescentibus; foliis oblongis, basi rotundatis, apicem versus cuspidato-dentatis, membranaceis, supra nitidis, subtus tenuissime ureo-tomentosis; fructibus ternis aut quinis, pedunculatis.*

This tree, which has a thin foliage, rises to the height of thirty or forty feet, and is from eighteen to twenty-four inches in diameter: it grows in the temperate and stony parts of New Spain between Moran, Pachuca, and Regla, and flowers in May.

QUERCUS JALAPENSIS. *Q. ramis tuberculatis; foliis longe petiolatis, ovato-oblongis, acuminatis, remote cuspidato-dentatis, subcoriaceis, glabris; fructibus solitariis aut geminis, breviter pedunculatis; cupulis cyathiformibus.*

This is a very lofty tree, about two feet in diameter: it is very common in the forests near Jalapa, on the eastern side of the mountains: it fructifies in January.

QUERCUS ACUTIFOLIA. *Q. foliis ovato-lanceolatis, acuminatis, inequaliter subcordatis, subtus puberulento-tomentosis, ferrugineis, sinuato-dentatis; dentibus elongatis, cuspidatis; fructibus geminis aut quaternis, brevissime pedunculatis.*

A very lofty tree, about two feet in diameter, which grows on the west side of the mountains between Venta de Acaguisolta and La Majonera: it fructifies in May.

QUERCUS STIPULARIS. *Q. ramulis ferrugineo-tomentosis; foliis obovato-oblongis, subcordatis, argute et grosse dentatis; crasse coriaceis, supra nitidis, subtus flavido-tomentosis; stipulis persistentibus; fructibus solitariis, aut geminis, sessilibus.*

This tree rises to the height of about fifty feet, and has a thick foliage: it is found on the mountains of Mexico near Actopan, and fructifies in May.

QUERCUS SYDEROXyla. *Q. ramulis pubescentibus; foliis obovato-oblongis, basi rotundatis, apicem versus argute serratis, cordatis, remote et obtuse dentatis, membranaceis, subtus tomentosis; fructibus longe pedunculatis, subspicatis; cupulis hemisphericis.*

A very lofty tree, from one to two feet in diameter: it grows in the temperate regions of New Spain near Villalpando and Santa Rosa, and fructifies in September.

QUERCUS PULCHELLA. *Q. ramis tuberculatis; foliis obovato-oblongis, subcordatis, argute dentatis, coriaceis, supra nitidis, subtus tenuissime ineano-tomentosis; fructibus geminis, breviter pedunculatis; cupulis depresso-sphericis.*

This tree is from fifteen to twenty feet in height, and grows

in the mountainous regions of New Spain between Guanajuato and Santa Rosa : it fructifies in September.

QUERCUS RETICULATA. *Q. ramulis subpubescentibus; foliis subsessilibus, obovatis, cordatis, remote dentatis, coriaceis, rugosis, subtus tenuissime tomentosis; fructibus geminis, longe pedunculatis; cupulis cyathiformibus.*

A tall tree, which grows in the mountainous parts of New Spain between Santa Rosa and Guanajuato, and fructifies in September.

QUERCUS CRASSIFOLIA. *Q. ramulis sulcatis, foliisque subtus, flavescenti-tomentosis, breviter petiolatis, obovatis cordatis, remote dentatis, crasse coriaceis; fructibus subternis, pedunculatis; cupulis subsphericis.*

This tree is from thirty to forty feet high, and has a very thick foliage: it is found in the stony and mountainous regions of New Spain near Chilpancingo, and fructifies in April.

QUERCUS SPICATA. *Q. ramulis tomentosis; foliis ellipticis, sinuato-dentatis, coriaceis, supra nitidis, subtus tenuissime canescenti-tomentosis; fructibus subgeminis, brevissime pedunculatis; cupulis cyathiformibus.*

This species is from thirty to forty feet in height, and is found in shady situations near El Oyamel, El Jacal, and Cerro de las Nabajas : it fructifies in May.

QUERCUS PANDURATA. *Q. ramulis levibus, hirtis, foliis obovato-oblongis, subpanduratis, subcordatis, sinuato-dentatis, coriaceis, subtus pubescentibus; fructibus subquinis, pedunculatis; cupulis cyathiformibus.*

This species is about fifteen feet high: it is found on the sides of the mountains in the kingdom of Mechoacan, between Ario and Patzeuaro : it fructifies in September.

QUERCUS AMBIGUA. *Q. ramulis glabris, obovato-oblongis, obtusis, basi rotundatis, subrepandis, membranaceis, subtus tenuissime pubescentibus; fructibus quinis aut septenis pedunculatis; pedunculis geminis, elongatis.*

This tree is about twenty feet high, and is found in the temperate regions of Mexico near Moran, Cerro Ventoso, and Omilton : it flowers in May.

QUERCUS GLAUDESCENS. *Q. ramulis angulatis; foliis brevissime petiolatis, obovato-oblongis, obtusiusculis, basi cuneatis, dentato-sinuatis, membranaceis, glaucescentibus; fructibus quinis aut septenis, pedunculatis.*

A tall tree, very common in the warm parts of New Spain between La Venta de la Majorena and Acaguisotla: it blooms in April.

QUERCUS OBTUSATA. *Q. ramulis tuberculatis, tenuiter pubescentibus, foliis oblongis, obtusis basi inæqualibus, repandis, coriaceis, subtus pulverulento-pubescentibus; fructibus subquinis, pedunculatis; cupulis campanulata-globosis.*

This species is very tall, with a remarkably straight trunk: it is found in the elevated and dry parts of New Spain near Ario, and flowers in September.

WALNUTS.

Of the various trees which compose the vast forests of North America east of the Mississippi, the Walnut ranks, after the Oak, among the genera the species of which are most multiplied. In this particular the soil of the United States is more favored than that of Europe, in no part of which is any species of this tree indigenous. I have distinguished in the United States ten species of Walnut, and others will probably be discovered in Louisiana. Travellers who visit these regions to explore their natural history should direct their attention to this class of vegetables, so interesting from the useful applications of their wood in the arts. There is room to hope, also, that species may be discovered, susceptible, like the Pecanutt Hickory, of rapid improvement by the aid of grafting and attentive cultivation. Some weight is given this consideration by an observation which I have often heard repeated by my father, that the fruit of the Common European Walnut, in its natural state, is harder than that of the American species just mentioned, and inferior to it in size and quality. To the members of agricultural societies in the United States it belongs to extend their observations and experiments on this subject, after the example of our ancestors, to whom we are indebted for a rich variety of fruits, equally salutary and beautiful.

The Walnuts of North America appear to present characters so distinct as to require their division into two sections. These characters consist principally in the form of the barren aments or catkins, and in the greater or less rapidity of vegetation in the trees. The first section is composed of Wal-

nuts with single aments, (Pl. 29 and 30,) and includes two species,—the Black Walnut and the Butternut; to which is added the European Walnut. The second section consists of such as have compound aments, (Pl. 36,) and comprises eight species,—the Pecannut Hickory, Bitternut Hickory, Water Bitternut Hickory, Mockernut Hickory, Shellbark Hickory, Thick Shellbark Hickory, Pignut Hickory, and Nutmeg Hickory. The first three species of the second section bear some relation to those of the first in their buds, which are not covered with scales. For this reason I have placed them immediately next, beginning with the Pecannut Hickory, which, by its numerous leaflets, most nearly resembles the Black Walnut and the Butternut, whose buds are also uncovered.

Throughout the United States the common name of Hickory is given to the species of the second section. This universal appellation is due to certain properties of their wood, which, however modified, are possessed by them all in a greater degree than by any other tree of Europe or America. These species exhibit also a striking analogy in their form and in their leaves, though they differ in the number and size of their leaflets. To these sources of confusion must be added another in the fruit, which is often so various in its appearance that it is easy to mistake the species to which it belongs. It is not, then, on the most remarkable differences alone that our distinctions must be founded; recourse must also be had to an examination of the shoots of the preceding year, of the buds, and of the aments. It was only by constant observation in the forests of the country, pursued through the course of a summer, that I became able readily to distinguish between mere varieties and species. M. Dehille, of the Institute of Egypt, who was at that time in the United States, took an active part in my researches, and resorted with me daily to the woods. Our investigations, I flatter myself, have had the result which may always be hoped for from unwearied perseverance.

From the considerations alleged, and principally from the striking resemblance of their wood, I have thought proper, in describing the species of Hickory, to speak but summarily of their respective properties, and to treat of this part of the subject collectively and comparatively more at large, in a separate article which will complete their history.

[For additional information on the Walnuts, see Nuttall's Supplement, vol. i. p. 39.

The genus *Carya* has been separated from *Juglans* by Nuttall chiefly on account of a technical distinction in the fruit.

Propagation, &c. The species is propagated by the nut; which, when the tree is to be grown chiefly for its timber, is best sown where it is finally to remain, on account of the taproot, which will thus have its full influence on the vigor and prosperity of the tree. In soils on moist or otherwise unfavorable subsoils, if sown where it is not finally to remain, a tile, slate, or flat stone should be placed under the nut at the depth of three or four inches, to give the taproot a horizontal direction; or, if this precaution is neglected, the taproot may be cut through with a spade six or eight inches below the nut. In a dry or rocky subsoil, or among rocks, no precaution of this kind is necessary. The varieties may be propagated by budding, grafting, in-arching, or layering, and, possibly, by cuttings of the root.

The nuts may be sown as soon as gathered if there is no danger from vermin; but, if there is, defer sowing till February. The most convenient mode is to deposit the seed in drills two feet apart from each other, placing the seeds at from three to six inches apart in the drills. If germinated in a heap before sowing, the points of the taproot may be pinched off before planting. Whether sown in drills or broadcast, almost the only attention required in their culture while in the nursery is to shorten once a year the tap or main root, in order to induce

them to throw out fibres, for the purpose of facilitating their transplantation, which, if performed in the autumn, should be followed in the spring, before the sap begins to rise, by cutting the head of the tree entirely off, leaving only a main stem terminating in the stumps of the principal branches. The wounds of these stumps are carefully covered with plaster composed of loam and cowdung, or grafting-clay, secured from the weather by straw and cords. Trees thus treated push out shoots of great vigor the first year, and, these being thinned out or rubbed off, the remainder soon form a head.

Soil and situation. The Walnuts attain the largest size in a deep loamy soil, dry rather than moist; but the fruit has the best flavor, and produces most oil, when the tree is grown in calcareous soils or among calcareous rocks: in a wet-bottomed soil it will not thrive. The Walnut is not a social tree, and neither produces good timber nor fruit when planted in masses. The Walnut is generally considered injurious by its shade both to man and plants.

Hickories planted in masses should be thinned when the plants have attained the height of from five to eight feet, the larger trees being left for timber, for ornament, or for fruit. Managed in this way, and gradually exposed to the action of the sun and air, they will have their peculiar beauties developed in the fullest manner. The wood which has grown most rapidly is the most valuable, having least of the heart-wood. The ashes of the Hickories abound in alkali, and are considered better for the purpose of making soap than any other of the native woods, being next to those of the Apple-tree. The Shell-bark grows best on the border of cultivated land, or on the edge of a forest. Some of our gardeners have paid attention to procuring the best nuts for cultivation, and the "true thin-shelled" may now be purchased from the nurseries; the nuts, however, will differ in different soils and situations, and even on individual trees growing in immediate proximity.]

METHODICAL DISPOSITION

OF THE

WALNUTS

OF

NORTH AMERICA.

Monocia polyandria. LINN. *Tercbinthaceæ*. JUSS.

FIRST SECTION.

Simple aments. (Pl. 29 and 30.)

VEGETATION RAPID.

1. Common European Walnut *Juglans regia*.
2. Black Walnut *Juglans nigra*.
3. Butternut *Juglans cathartica*.

SECOND SECTION.

HICKORIES.

Compound aments, each peduncle bearing three. (Pl. 36, fig. 3.)

VEGETATION SLOW.

4. Pecannut Hickory . . . *Juglans (Carya) olivæformis*.
5. Bitternut Hickory . . . *Juglans (Carya) amara*.
6. Water Bitternut Hickory *Juglans (Carya) aquatica*.
7. Mockernut Hickory . . . *Juglans (Carya) tomentosa*.
8. Shellbark Hickory . . . *Juglans (Carya) squamosa*.
9. Thick Shellbark Hickory *Juglans (Carya) lacineosa*.
10. Pignut Hickory *Juglans (Carya) porcina*.
11. Nutmeg Hickory *Juglans (Carya) myristicæformis*.

COMMON EUROPEAN WALNUT.

JUGLANS REGIA. *J. foliolis subseptenis, ovalibus, glabris, subserratis, subaequalibus: fructibus subovalibus.*

THE Walnut which for several centuries has been cultivated in Europe is a native of Asia. According to an ancient but uncertain tradition, its fruit was brought from Persia with the Peach and the Apricot. My father, who, in the years 1782, '83, and '84, visited that part of the East to examine its natural productions, first ascertained with exactness the origin of this tree: he found it, in the natural state, in the province of Ghilan, which lies on the Caspian Sea, between the 35th and 40th degrees of latitude.

The period of its introduction into Europe—a point on which ancient authors leave us in obscurity—is proved to be remote by several rites in use among the Romans; such, for instance, as the distribution of nuts in the *Cerealia*. In the village festival of the *Rosière*, instituted by St. Medard, at Salency, Department of the Oise, twelve hundred years ago, it is directed that an offering composed of nuts and other fruits of the country shall be presented to the young maid who is crowned; which proves the tree to have been already naturalized in that part of France.

The Walnut is common throughout the centre of Europe; but it flourishes most in the western and southern departments of France, in Spain and in Italy, which approach nearest to the latitude in which it grows naturally. In France it is only in the west and south, where the vegetation of the Walnut is perfectly secure from frost, that its wood is of a superior quality, and that its fruit is regularly yielded in sufficient abundance to become an article of commerce.

The European Walnut is one of the tallest and most beauti-

ful among fruit-trees, and one of the most remarkable for the amplitude of its summit and the thickness of its shade. On the trunk of old trees, which frequently are several feet in diameter, the bark is thick and deeply furrowed; on the upper branches it is gray and smooth, a good deal resembling that of the Butternut. The leaves are borne by long petioles, and are composed of two, three, and sometimes four pair of leaflets, surmounted by an odd one. The leaflets are oval and smooth; when bruised, they exhale a strong aromatic odor. In the extreme heat of summer, the emanations from the Walnut are so powerful as to produce unpleasant effects upon some persons if they slumber in its shade.

The flowers of the Common European Walnut, like those of the Black Walnut and Butternut, appear before the unfolding of the leaves; the barren ones in single, pendulous, imbricated aments, the fertile ones on separate branches, at the end of the young shoots, and commonly in pairs. The fruit is green and oval, and in the natural state contains a small hard nut. In the most esteemed cultivated species, the fruit is oval and strongly odoriferous, about an inch and two-thirds long and from an inch and a quarter to an inch and a half in diameter. The nut occupies two-thirds of its volume.

Toward autumn the husk softens, and, decaying from about the nut, allows it to fall. The shell is slightly channelled, and so thin as to be easily crushed by the fingers. The kernel is of a very agreeable taste: it is large, covered with a fine pellicle, and separated by a thin partition, which may readily be detached both from the shell and from the kernel.

The nuts are better-tasted and easier of digestion soon after their maturity than later in the season, when the oily principle becomes perfectly formed; they are then oppressive, if immoderately eaten.

A dessert of an excellent relish is made by extracting the kernels a fortnight before they are ripe, and seasoning them

with the juice of green grapes and salt. They should be thrown into water as soon as they are taken from the shell, and allowed to remain till the moment when they are seasoned to be set upon the table. They are sold in Paris by the name of *Cer-
neaux*; and a greater quantity of walnuts is consumed in this way by people in easy circumstances than after they are perfectly ripe: the use of them is then almost exclusively confined to the lower classes.

The Common Walnut is more multiplied in the departments of France which lie between the 45th and the 48th degrees of latitude than in any other part of Europe. In these departments it is planted in the midst of cultivated fields, like the Apple-tree for cider in those of the north and the centre: the fruit, the oil, and the wood, may be considered as forming one of their principal branches of commerce.

In extracting the oil of Walnuts, certain delicate attentions are necessary to insure its fineness. When the fruit is gathered and the nuts are separated from the husks, they should be kept dry and occasionally moved till they are used. The proper time for the operation is at the close of winter, as in this interval the change by which the mucilage of the fruit is converted into oil has become completely effected, and by longer delay the kernel grows rancid and the oil is of a vitiated quality. The nut is cracked by striking it on the end with a small mallet, and pains are taken not to bruise the kernel. The slight ligneous partition is detached, and such kernels as are partially spoiled are selected and thrown aside. The sound kernels, thus cleared from every particle of the shell, should be sent immediately to the mill, as they soon become rancid by exposure to the air. They are crushed by a vertical stone, which turns in a circular trough, and is moved by a horse or by a current of water. The paste is next enclosed in bags of strong linen and submitted to the press. The oil which flows under this first pressure without the application of heat is of the best quality. It is very clear,

and is proper for food; but it sensibly retains the taste of the nut, which in general is not agreeable to persons unaccustomed to it, so that the consumption is limited to the departments where it is made. To be kept sweet for the table, it should be drawn off several times during the first months, carefully corked, and stored in the cellar, as it is more easily affected than any other oil by the action of air and heat.

After the first expression, the paste is emptied from the sacks, moistened with warm water, and moderately heated in coppers. It is then replaced in the sacks and returned to the press. The oil of the second discharge is highly colored, and very speedily becomes rancid; it is therefore employed only in the preparation of colors. The cakes which remain after the expression is finished are used for fattening fowls.

Although nut-oil, as an article of diet, is in general use in the departments where the tree abounds, it serves a still more important purpose in the preparation of fine colors. It is preferred on account of the complete and rapid manner in which it dries, and of the facility of obtaining it perfectly limpid, which is done by diffusing it upon water in large shallow vases.

In copper-plate printing, walnut-oil is considered, in Paris, indispensably necessary for a fine impression in black or in colors. But there are peculiar modes of preparing it for the several colors with which it is to be mixed. Thus, for white, blue, light green, and the intermediate shades, it is reduced by boiling to two-thirds of its bulk; but, for dark green and black, to one-fifth, which leaves it a thick, semifluid substance. To facilitate the process, one-tenth part of linseed-oil is added to it; it is then placed, in an iron or copper vessel, over an active, clear fire. When it begins to boil rapidly, the vessel is uncovered, and the oil takes fire by contact with the flame and burns till it is reduced to the proper consistency: sometimes it is not allowed to kindle, but, when the ebullition commences, crusts of bread are thrown into it, which remain till the necessary

evaporation is effected, and are then taken out, charged with mucilaginous particles. The principal advantage of this oil, in the preparation of white lead for painting the interior of houses, as well as of the colors employed in copper-plate printing, is the longer and more perfect preservation of the tints. The backs of prints done with it do not turn yellow like others.

A fine stomachic liquor is made with the fruit of the Walnut gathered a month before its maturity. Twelve green nuts in the husk are bruised and thrown into a pint of good brandy; after they have steeped three weeks, the brandy is filtered through brown paper, and a quarter of a pound of loaf-sugar is added. This cordial improves by age.

Dyers obtain, by boiling the husks when they begin to decay, and the bark of the roots, a substantial dark brown with which they dye woollens. Cabinet-makers also make use of it in staining other pieces of wood in imitation of Walnut.

Among the American Walnuts which are found east of the Mississippi, the Black Walnut bears the greatest resemblance to the European Walnut in its general appearance, in its flowers and fruit, and in the qualities of its wood: in foliage they are strikingly different. The wood of the European Walnut is inferior in strength and weight, and, I believe, far more liable to injury from worms.

Twenty or thirty years ago, before Mahogany was imported in such abundance into Europe, Walnut wood was employed almost exclusively in cabinet-making. In the country it is still in general use, and the furniture made of it is far from being inelegant, especially pieces obtained from such old trees as bear small and thick-shelled nuts. It is preferred for the stocks of muskets; and in Paris and Brussels no other wood is used for the panels of carriages. The old trees furnish excellent screws for large presses. Great quantities of wooden shoes are manufactured of Walnut, which are more highly esteemed than others.

The wood of the European Walnut is largely exported from

the south of France to the north and to Holland and Germany: formerly it was carried to England.

Like other fruit-trees, whose perfection is among the "noblest conquests of industrious man," the Walnut has been greatly improved by long and careful cultivation. There are seven or eight cultivated varieties, whose superiority is principally apparent in the augmented size of the fruit and in the diminished thickness of the shell. Of these the most esteemed, after that which I have described, are the *St. Jean* and the *Jauge* Walnuts. The St. John Walnut is a variety obtained within a few years. It yields fruit as large and as abundant as the common Walnut, and for that part of Europe which lies beyond the 45th degree of latitude it possesses an advantage in opening its vegetation three weeks later and in being thus secure from the injuries of frost. The *Jauge* Walnut is chiefly remarkable for the size of its fruit, which is twice as large as the variety represented in the plate. It is unproductive, and the kernel does not fill the shell. The *Jauge* nut is made into cases by jewellers, and furnished with trinkets, for the amusement of children.

The wood of the Black Walnut is already superior to that of the European Walnut; and it will acquire a still finer grain when it is raised on lands that have been long under cultivation. It is solely for the excellency of its fruit and the decided superiority of its oil in the preparation of colors, that the European Walnut should be warmly recommended to the attention of Americans. It would thrive better than elsewhere in places where the Black Walnut naturally abounds.

In some parts of Pennsylvania and Maryland the Black Walnuts have been preserved in clearing the lands. Great advantage would be found in grafting them with the European Walnut. The limbs should be cut fifteen inches from the trunk, and from the stumps will spring vigorous shoots, which, the second year, may be grafted by inoculation. Fifty or sixty buds should be set upon each tree, as is practised near Lyons, where

it is found that, by inserting the Walnut of St. John on the common Walnut, the fruit is rendered finer and the crop more certain. Black Walnuts thus grafted begin to bear the fifth year. On estates where no Black Walnut exist, the deficiency may be supplied by planting the nuts and grafting the young stocks when they come to the height of eight or ten feet.

It should be observed, that in the Walnut, more than any other tree, it is necessary, on account of the loose texture of the wood and the large volume of the pith, to protect the amputated limbs from the weather. A covering of clay should be so nicely adapted to the exposed surface as entirely to exclude the rain; otherwise, decay will commence and spread itself into the body of the tree.

In those parts of France, Belgium, and Germany, where the Walnut is not cultivated for commerce, the trees have generally sprung from the seed, which is the cause of the inferiority of their fruit. For it is observed that, with a few accidental exceptions, the finest fruits and flowers degenerate in reproduction. This inconvenience would be experienced in the United States; and, as there do not perhaps exist in that country, south of the Hudson River, ten European Walnut-trees,* I should recommend the importation from Bordeaux of young grafted trees, which will soon furnish the means to such proprietors as wish to enrich their estates with this useful and magnificent tree.

PLATE XXIX.

Fig. 1. A leaf of half the natural size. Fig. 2. Barren flowers. Fig. 3. Fertile flowers. Fig. 4. A nut in its husk of the natural size. Fig. 5. A nut without its husk. Fig. 6. A nut deprived of half the shell, to show the kernel.

* [Since this was written, the European Walnut has been extensively introduced in America, but as far north as Philadelphia it does not produce fruit abundantly except in sheltered situations or when surrounded by hard surface-ground.]

BLACK WALNUT.

JUGLANS NIGRA. J. foliolis quinqueis, subcordatis, superne angustatis, serratis; fructu globoso, punctato, scabriusculo; nucē corrugatā.

THIS tree is known in all parts of the United States where it grows, and to the French of Upper and Lower Louisiana, by no other name than Black Walnut. East of the Alleghany Mountains, the most northern point at which it appears is about Goshen in the State of New Jersey, in the latitude of 40° 50'.* West of the mountains it exists abundantly two degrees further north, in that portion of Genesee which is comprised between the 77th and 79th degrees of longitude. This observation, as I shall have occasion to remark, is applicable to several other vegetables the northern limit of whose appearance varies with the climate; and this becomes milder in advancing toward the west. The Black Walnut is multiplied in the forests about Philadelphia; and, with the exception of the lower parts of the Southern States, where the soil is too sandy, or too wet, as in the swamps, it is met with to the banks of the Mississippi, throughout an extent of two thousand miles. East of the Alleghany Mountains in Virginia, and in the upper part of the Carolinas and of Georgia, it is chiefly confined to the valleys where the soil is deep and fertile, and which are watered by creeks and rivers: in the western country, in Genesee, and in the States of Ohio and Kentucky, where the soil in general is very rich, it grows in the forests, with the Coffee-tree, Honey Locust, Red Mulberry, Locust Shellbark Hickory, Black Sugar Maple, Hack Berry, and Red Elm:—all which trees prove the goodness of the soil in which they are found.

It is in these countries that the Black Walnut displays its

* [It is found in Massachusetts. EMERSON.]

angustatis,
ugald.


s where it
anna, by no
ny Moun-
s is about
of 40° 50'.*
ees farther
ed between
vation, as I
veral other
varies with
toward the
rests about
parts of the
o wet, as in
Mississippi,
of the Alle-
part of the
the valleys
watered by
essee, and in
n general is
tree, Honey
Black Sugar
s prove the

displays its





Black Walnut
Juglans nigra.



full proportions. On the banks of the Ohio, and on the islands of that beautiful river, I have often seen trees of three or four feet in diameter and sixty or seventy feet in height. It is not rare to find them of the thickness of six or seven feet. Its powerful vegetation clearly points out this as one of the largest trees of America. When it stands insulated, its branches, extending themselves horizontally to a great distance, spread into a spacious head, which gives it a very majestic appearance.

The leaves of the Black Walnut when bruised emit a strong aromatic odor. They are about eighteen inches in length, pinnate, and composed in general of six, seven, or eight pair of leaflets surmounted by an odd one. The leaflets are opposite and fixed on short petioles; they are acuminate, serrate, and somewhat downy. The barren flowers are disposed in pendulous and cylindrical aments, of which the peduncles are simple, unlike those of the Hickories, (Pl. 30, fig. 1.) The fruit is round, odoriferous, of rather an uneven surface, and always appears at the extremity of the branches: on young and vigorous trees it is sometimes seven or eight inches in circumference. The husk is thick, and is not, as in the Hickories, divided into sections, but when ripe it softens and gradually decays. The nut is hard, somewhat compressed at the sides, and sulcated. The kernel, which is divided by firm ligneous partitions, is of a sweet and agreeable taste, though inferior to that of the European Walnut. These nuts are sold in the markets of New York, Philadelphia, and Baltimore, and served upon the tables. The size of the fruit varies considerably, and depends upon the vigor of the tree and upon the nature of the soil and of the climate. On the banks of the Ohio, and in Kentucky, the fruit with the husk is seven or eight inches in compass, with the nut proportionally large: in Genesee, on the contrary, where the cold is intense, and in fields exhausted by cultivation, where these trees have been preserved since the first clearing of the land, it is not of more than half this volume. Some variations are observed in

the form of the fruit and in the moulding of the shell; but these I consider as merely accidental differences. Indeed, there is no genus of trees in America in which the fruit of a given species exhibits such various forms as in the Walnut; and doubtless this circumstance has misled observers, who, being acquainted only with the small number of trees existing in European gardens, have described them as distinct species.

The bark of the Black Walnut is thick, blackish, and, on old trees, deeply furrowed. When the timber is freshly cut, the sap is white and the heart of a violet-color, which after a short exposure to the air assumes an intenser shade and becomes nearly black; hence probably is derived the name of Black Walnut. There are several qualities for which its wood is principally esteemed. It remains sound during a long time, even when exposed to the influences of heat and moisture: but this observation is applicable only to the heart; the sap speedily decays. It is very strong and very tenacious; when thoroughly seasoned, it is not liable to warp and split; and its grain is sufficiently fine and compact to admit of a beautiful polish. It possesses, in addition to these advantages, that of being secure from worms. On account of these excellencies, it is preferred and successfully employed in many kinds of work. East of the Alleghanies, its timber is not extensively used in building houses; but in some parts of Kentucky and Ohio it is split into shingles eighteen inches long and from four to six inches wide, which serve to cover them: sometimes also this timber enters into the composition of the frame. But it is chiefly in cabinet-making that the Black Walnut is employed wherever it abounds. By selecting pieces from the upper part of the trunk, immediately below the first ramification, furniture is sometimes made which from the accidental curlings of the grain is highly beautiful; but, as its color soon changes to a dusky hue, the Wild Cherry wood is frequently preferred for this use. The Black Walnut is also employed for musket-stocks; it is stronger and

tougher than the Red-flowering Maple, which, from its superior lightness and elegance, is chosen for fowling-pieces. In Virginia posts are very commonly made of Black Walnut; and, as it lasts undecayed in the ground from twenty to twenty-five years, it appears every way fit for this purpose. I have been assured that it makes excellent naves for wheels, which further proves its strength and durability. At Philadelphia coffins are very frequently made of it.

The timber of this tree is also excellently adapted to certain uses in naval architecture. It should never be wrought till it is perfectly seasoned, after which it is asserted to be more durable, though more brittle, than the White Oak. Breckel, in his "History of North Carolina," affirms that it is not liable, like the Oak, to be attacked by sea-worms in warm latitudes. This advantage, if it is real, is highly important, and deserves to be ascertained by further observation. In the marine lumber-yards of Philadelphia I have often seen it used for knees and floor-timber; but in the vessels built at Wheeling and Marietta, towns on the Ohio, it constitutes a principal part of the frame. On the river Wabash canoes are made of it which are greatly esteemed for strength and durability. Some of them, fashioned from the trunk of a single tree, are more than forty feet long and two or three feet wide.

The Black Walnut is exported in small quantities to England in planks of two inches in thickness, which are sold at Philadelphia at four cents a foot.*

The husk of the fruit yields a color similar to that which is

* [The demand for Walnut wood in the Atlantic cities, and the want of attention to its cultivation, have since made it necessary for the cabinet-makers, &c. to import from the West the greater portion of their supplies. This resource must fail in time, and the wood may not improbably become nearly as costly as Mahogany, which it resembles in many of its properties.]

obtained from the European Walnut. It is used in the country for dyeing woollen stuffs.

This tree has long since been introduced, in England and France, into the gardens of the lovers of foreign culture. It succeeds perfectly, and yields fruit abundantly. Though differing widely from the European species, it bears a nearer resemblance to it than any other American Walnut. By comparing the two species as to their utility in the arts and in commerce, it will appear that the wood of the Black Walnut is more compact, heavier, and much stronger; that it is susceptible of a finer polish, and that it is not injured by worms,—qualities which, as has been seen, render it fit not only for the same uses with ours, but also for the larger works of architecture. These considerations sufficiently evince that it is a valuable tree, and that it is with great reason that many proprietors in America have spared it in clearing their new lands. On highroads, I am of opinion that it might be chosen to succeed the Elm; for experience has proved that, to insure success in the continued cultivation of trees or herbaceous plants on the same soil, the practice must be varied with species of different genera.

Nuts of the European Walnut and of the Black Walnut have been planted at the same time in the same soil: those of the Black Walnut are observed to shoot more vigorously, and to grow in a given time to a greater height. By grafting the European upon the American species at the height of eight or ten feet, their advantages, with respect to the quality both of wood and of fruit, might be united.

PLATE XXX.

A leaf of half its natural size. Fig. 1. A nut with its husk. Fig. 2. A nut without its husk. Fig. 3. A barren nut.

the country

England and
culture. It
though differ-
a nearer re-
ut. By com-
s and in com-
ck Walnut is
is susceptible
ms,—qualities
the same uses
cture. These
able tree, and
rs in America
n highroads, I
ceed the Elm;
n the continued
same soil, the
genera.

ck Walnut have
l: those of the
sly, and to grow
g the European
ght or ten feet,
oth of wood and

sk. Fig. 2. A nut
t.



P. J. Redouté del.


Reuss sculp.

Butter Nut.
Juglans cathartica.

Pl. m.



Rinari



BUTTERNUT

JUGLANS CATHARTICA. *J. foliolis subquindenis, lanceolatis, basi rotundato-obtusis subtus tomentosis, leviter serratis; fructu oblongo, ovato, apice mammoso, viscido, longè pedunculato, nuce oblongâ, acuminatâ, insigniter insculptè-scabrosâ.*

THIS species of Walnut is known in North America under different denominations. In Massachusetts, New Hampshire, and Vermont, it bears the name of Oilnut; in Pennsylvania and Maryland, and on the banks of the Ohio, it is generally known by that of White Walnut; in Connecticut, New York, New Jersey, Virginia, and in the mountainous districts of the upper parts of the Carolinas, it is called Butternut. The last of these names I have retained, because it is not wholly unknown in those parts of the United States where the others are in general use, and because the wood is employed in the neighborhood of New York for a greater variety of uses than elsewhere. I think also that the Latin specific name *Cathartica*, which was long since given it by Doctor Cutler of Massachusetts, should be definitively substituted for that of *Cinerea*, by which it has hitherto been distinguished among botanists. This last appellation, derived from the color of the secondary branches, whose bark is smooth and grayish, suggests only an unimportant characteristic, while the first expresses one of the most interesting properties of the tree.

The Butternut is found in Upper and Lower Canada, in the district of Maine, on the shores of Lake Erie, in the States of Kentucky and Tennessee, and on the banks of the Missouri; but I have never met with it in the lower part of the Carolinas, of Georgia, and of East Florida, where the nature of the soil and the intemperate heat of the summer are unfavorable to its vegetation. In cold regions, on the contrary, its growth is luxu-

riant; for in the State of Vermont,* where the winter is so rigorous that sledges are used during four months in the year, this tree attains a circumference of eight or ten feet. I have nowhere seen it more abundant than in the bottoms which border the Ohio between Wheeling and Marietta; but the thickness of these forests, which are hardly penetrated by the sun, appears to prevent its utmost expansion. I have seen here no trees as large as some in New Jersey, on the steep and elevated banks of the Hudson, nearly opposite to the city of New York. The woods in this place are thin, and the soil cold, unproductive, and interspersed with large rocks, in the interstices of which the biggest Butternuts have their root. I have measured some of them, which, at five feet from the ground, were ten or twelve feet in circumference, and which were fifty feet in height, with roots extending even with the surface of the ground, in a serpentine direction and with little variation in size, to the distance of forty feet. The trunk ramifies at a small height, and the branches, seeking a direction more horizontal than those of other trees, and spreading widely, form a large and tufted head, which gives the tree a remarkable appearance.

The buds of the Butternut, like those of the Black Walnut, are uncovered. In the spring its vegetation is forward, and its leaves unfold a fortnight earlier than those of the Hickories. Each leaf is composed of seven or eight pair of sessile leaflets, and terminated by a petiolated odd one. The leaflets are from two to three inches in length, lanceolate, serrate, and slightly downy. The barren flowers stand on large cylindrical aments, which are single, four or five inches long, and attached to the shoots of the preceding year; the fertile flowers, on the contrary, come out on the shoots of the same spring, and are situated at their extremity. The ovary is crowned by two rose-colored stigmata. The fruit is commonly single, and suspended by a

* [It occurs in all the New England States. EMERSON.]

thin, pliable peduncle, about three inches in length; its form is oblong-oval, without any appearance of seam. It is often two and a half inches in length and five inches in circumference, and is covered with a viscid adhesive substance, composed of small transparent vesicles, which are easily discerned with the aid of a glass. The nuts are hard, oblong, rounded at the base, and terminated at the summit in an acute point; the surface is very rough, and deeply and irregularly furrowed. They are ripe in the neighborhood of New York about the 15th of September,—a fortnight earlier than the other species of Walnut. Some years they are so abundant that one person may gather several bushels of them in a day. The kernel is thick and oily, and soon becomes rancid; hence, doubtless, are derived the names of Butternut and Oilnut. These nuts are rarely seen in the markets of New York and Philadelphia. The Indians who inhabited these regions pounded and boiled them, and, separating the oily substance which swam upon the surface, mixed it with their food. When the fruit has attained about half its growth, it is sometimes used for making pickles, being first plunged into boiling water, and thoroughly wiped, to clean it of its down, and afterward preserved in vinegar.

The Black Walnut and Butternut, when young, resemble each other in their foliage and in the rapidity of their growth; but when arrived at maturity, their forms are so different as to be distinguishable at first sight. Remarkable peculiarities are also found on examining their wood, especially when seasoned: the Black Walnut is heavy, strong, and of a dark brown color, while the Butternut is light, of little strength, and of a reddish hue; but they possess in common the great advantage of lasting long and of being secure from the annoyance of worms. From its want of solidity, and from the difficulty of procuring pieces of considerable length, Butternut timber is never used in the cities in the construction of houses, though it is sometimes employed for this purpose in the country. In some districts of New

Jersey it is often taken for the sleepers which are placed immediately on the ground, in the framing of houses and barns. As it long resists the effects of heat and moisture, it is esteemed for the posts and rails of rural fence and for troughs for the use of cattle. For corn-shovels and wooden dishes, it is preferred to the Red-flowering Maple, because it is lighter and less liable to split; consequently, articles made of it are sold at a higher price. Near New York I have observed it to be made use of for canoes formed of one or two logs, and for the futtocks designed to give them solidity; but in boats of considerable size some stronger wood is selected for this purpose. At Pittsburg, on the Ohio, the Butternut is sometimes sawn into planks, for the construction of small skiffs, which, on account of their lightness, are in request for descending the river. At Windsor, in Vermont, it is used for the panels of coaches and chaises; the workmen find it excellently adapted to this object, not only from its lightness, but because it is not liable to split and receives paint in a superior manner. Indeed, I have remarked that its pores are more open than those of the Poplar and Basswood.

The medicinal properties of Butternut bark have long since been proved by several eminent physicians of the United States, and, among others, by Doctor Cutler.* An extract in water, or even a decoction sweetened with honey, is acknowledged to be one of the best cathartics afforded by the materia medica; its purgative operation is always sure, and unattended, in the most delicate constitutions, with pain or irritation. Experience has shown that it produces the best effects in many cases of dysentery. It is commonly given in the form of pills, and, to adults, in doses from half a scruple to a scruple. It is not, however, in general use, except in the country, where many of the farmers'

* [Previously, by Dr. Rush, in the war of the Revolution: the official extract is the only one now used by practitioners.]

wives provide a small store of it in the spring for the wants of their families and of their neighbors. They obtain it by boiling the bark entire in water till the liquid is reduced by evaporation to a thick, viscid substance, which is almost black. This is a faulty process: the exterior bark, or the dead part which covers the cellular integument, should first be taken off; for, by continued boiling, it becomes charged with four-fifths of the liquid, already enriched with extractive matter. I have also seen this bark successfully employed as a revulsive in inflammatory ophthalmias and in the toothache: a piece of it soaked in warm water is applied in these cases to the back of the neck. In the country it is sometimes employed for dyeing wool of a dark brown color; but the bark of the Black Walnut is preferable for this purpose.

On a live tree the cellular tissue, when first exposed, is of a pure white; in a moment it changes to a beautiful lemon-color, and soon after to a deep brown.

If the trunk of the Butternut is pierced in the month which precedes the unfolding of the leaves, a pretty copious discharge ensues of a slightly-sugary sap, from which, by evaporation, sugar is obtained of a quality inferior to that of the Sugar Maple.

Although the Butternut, as has been seen, possesses useful properties, I do not think it sufficiently valuable, either in the arts or for fuel, to recommend its introduction into the forests of the Old Continent: it should find place only in our pleasure-grounds.

PLATE XXXI.

A leaf of half its natural size. Fig. 1. A nut with its husk. Fig. 2. A nut without its husk.

PECANNUT HICKORY

JUGLANS OLIVÆFORMIS. *J. foliolis plurimis, subpetiolatis, falcatis, serratis; fructu oblongo, prominulo-quadrangulo; nucce olivæformi, levi.*

Carya olivæformis. Nutt.

THIS species, which is found in Upper Louisiana, is called, by the French of Illinois and New Orleans, *Pecanier*, and its fruit *Pecanes*. This name has been adopted by the inhabitants of the United States, who call it Pecannut. On the borders of the rivers Missouri, Illinois, St. Francis, and Arkansas, it is most abundantly multiplied: it is also common on the river Wabash; on the Ohio it is found for 200 miles from its junction with the Mississippi; higher than this it becomes more rare, and is not seen beyond Louisville. My father, in traversing this country, learned from the French inhabitants, who ascend the Mississippi in quest of furs, that it is not found on that river beyond the mouth of the Great Mackakity, which discharges itself in the latitude of 42° 51'.

This tree grows most naturally in cold and wet grounds. There is a swamp of 800 acres, situated on the right bank of the Ohio, opposite to the river Cumberland, which is said to be entirely covered with it, and which is called, by the French, *La Pecanière*.

The Pecannut is a beautiful tree, with a straight and well-shaped trunk; in the forests it reaches the height of sixty or seventy feet. Its wood is coarse-grained, and, like the other Hickories, heavy and compact: it possesses also great strength and durability; but in these respects it is inferior to some spe-

* [It bears fruit in gardens near Philadelphia, but can scarcely be said to perfect it.]

s, *serra*
levi.

NUTT.

called,
and its
abitants
orders of
as, it is
the river
unction
re rare,
aversing
o ascend
on that
which dis-

grounds.
bank of
aid to be
French,

nd well-
sixty or
the other
strength
some spe-

said to per-





Pacament Hickory.
Juglans oliviformis.

E. W. Brown



cies hereafter to be described. Its buds, like those of the Black Walnut and Butternut, are uncovered. The leaves are from twelve to eighteen inches in length, and are supported by petioles somewhat angular, and slightly downy in the spring. Each leaf is composed of six or seven pair of sessile leaflets and terminated by a petiolated odd one, which is commonly smaller than the pair immediately preceding. The leaflets on flourishing trees are from two to three inches long, ovate, serrate, and remarkable for the circular form of the upper edge, while the lower one is less rounded. It is also to be noticed that the main rib is placed a little below the middle of the leaflet.

The nuts, which are usually abundant, are contained in a husk from one to two lines thick, and have four slightly-prominent angles corresponding to their internal divisions. They vary in length from an inch to an inch and a half, are pointed at the extremities, of a cylindrical form, and of a yellowish color, marked at the period of perfect maturity with blackish or purple lines. The shell is smooth and thin, though too hard to be broken by the fingers: the kernel is full, and, not being divided by ligneous partitions, is easily extracted. These nuts, which are of a very agreeable taste, form an object of petty commerce between Upper and Lower Louisiana. From New Orleans they are exported to the West Indies and to the ports of the United States. They are not only better than any other species of North American Walnuts, but they appear to me to be more delicately flavored than those of Europe. And, besides, wild varieties of the Pecannut are found, the fruit of which is much larger than that of the European Walnut unimproved by culture. I am of opinion, then, that this tree merits the attention both of Americans and Europeans, and that by assiduous cultivation it may be brought to a high degree of perfection. These advantages, it is true, are balanced in part by the slowness of its growth; there are trees in France which

have been planted more than thirty years, and which do not yield fruit.

If the practice should be successfully adopted of grafting the Pecanutt on the Black Walnut, or on the Common Walnut, its vegetation would be incomparably more rapid, and no motive should discourage its propagation in Europe.

PLATE XXXII.

A leaf of half its natural size. Fig. 1. A nut with its husk. Fig. 2. A nut without its husk.

BITTERNUT HICKORY.

JUGLANS AMARA. *J. arbor maxima, foliolis 7-9^{nis}, glabris, conspicuè serratis, impari breviter petiolato: fructu subrotundo-ovoideo, supernè suturis prominulis; nucè levi, subgloboſâ, mucronatâ: putamine fragili, nucleo amaro.*

Cárya amara. NUTT.

THIS species is generally known in New Jersey by the name of Bitternut Hickory; in Pennsylvania, and particularly in the county of Lancaster, it is called White Hickory, and sometimes Swamp Hickory; farther south, it is confounded with the Pig-nut Hickory. The French of Illinois, like the inhabitants of New Jersey, give it the name of Bitternut, which, as it indicates one of the peculiar properties of the fruit, I have chosen to retain.

The Bitternut Hickory, I believe, is nowhere found much beyond the boundaries of Vermont, in the 45th degree of latitude. It is not seen in the province of Maine, where the borders of the rivers offer situations analogous to those in which it

do not

ng the
ut, its
motive


ig. 2. A

conspicua
superna
e fragili,

NUTT.

ne name
y in the
ometimes
the Pig-
tants of
it indi-
e chosen

nd much
e of lati-
e borders
which it





W. L. A. 11

Bitter Nut Hickory
Amelanchier amara.

1133

shell is white, smooth, and thin enough to be broken by the fingers. The kernel is remarkable for the deep inequalities produced on every side by its foldings. It is so harsh and bitter that squirrels and other animals will not feed on it while any other nut is to be found.

In some parts of Pennsylvania where this tree is multiplied an oil is extracted from the nuts, which is used for burning in lamps and other inferior purposes. But from these experiments, in which individuals have succeeded, it is not to be concluded that a sufficient product of this sort can be obtained to form a branch of industry; neither this nor any other species of Walnut is abundant enough in the United States.

In the texture of its bark, and in the color of its heart and sap, the Bitternut Hickory resembles the other Hickories, and its wood possesses, though in an inferior degree, the weight, strength, tenacity, and elasticity which so plainly distinguish them. At Lancaster it is used for fuel; but it is not considered superior to the White Oak nor sold at a higher price.

The Bitternut Hickory exists and bears fruit in several gardens in France; but it is of no value for its nuts, and flourishes only in very fertile soils. As its wood, also, is proved in America to be inferior to that of the following species, I think it should not be propagated in the forests of Europe.

PLATE XXXIII.

A leaf of the natural size. Fig. 1. A nut with its husk. Fig. 2. A nut without its husk.

by the
qualities
and bitter
while any

Multiplied
forming in
experiments,
included
to form a
of Wal-

heart and
series, and
weight,
distinguish
considered

gardens
shows only
America to
should not

2. *A nat*




Fig. 1

Fig. 2

Water bitter Nut Hickory.
Juglans aquatica





WATER BITTERNUT HICKORY.

JUGLANS AQUATICA. *J. foliis 9-11^{nis}, lanceolato-acuminatis, subserratis, sessilibus, impari breviter petiolato: fructibus pedunculatis, nuce subdepressâ, parvâ, rubiginosâ, tenerâ.*

Carya aquatica. NUTT

No specific name has hitherto been given this species, which is confined to the Southern States; it is confounded with the Pignut Hickory, though differing from it in many respects. The name which I propose appears sufficiently appropriate; for I have always found this tree in swamps, and ditches which surround the ricefields, where it is accompanied by the Red-flowering Maple, Tupelo, Cypress, and Carolina Poplar. The Water Bitternut Hickory grows to the height of forty or fifty feet, and in its general appearance resembles the other Hickories. Its leaves are eight or nine inches long, and of a beautiful green. They are composed of four or five pair of sessile leaflets surmounted by a petiolated odd one. The leaflets are serrate, four or five inches long, eight or nine lines broad, and very similar to the leaves of the Peach-tree.

The husk is thin, and the nuts are small, angular, a little depressed at the sides, somewhat rough, of a reddish color, and very tender. The kernel is very bitter, formed in folds like that of the Bitternut Hickory, and, as may be supposed, is not eatable. The wood of this species, though partaking of the common properties of the Hickories, is in every respect inferior to the others, from the nature of the grounds on which it grows.

The Water Bitternut Hickory, which I have introduced into France, flourishes unchecked by the rigor of our winters; but I do not think it deserves to find a place in the forests of Europe, nor to be spared in clearing the new lands of America. The southern parts of the United States possess many sorts of timber

more useful in building, to which purpose this, like the other Hickories, is poorly adapted.

PLATE XXXIV.

A branch with leaves of the natural size. Fig. 1. Nuts with their husks.

Fig. 2. Nuts without their husks.

MOCKERNUT HICKORY.

JUGLANS TOMENTOSA. *J. foliolis 7-9^{uis}, leviter serratis, conspicuè villosis, impari subpetiolato; amentis compositis, longissimis, filiformibus, eximie tomentosis: fructu globoso vel oblongo; nucè quadrangulâ, crassâ, durissimâque.*

Carya tomentosa. NUTT.

In the parts of New Jersey which lie on the river Hudson, and in the city of New York and its vicinity, this species is known by the name of Mockernut Hickory, and, less commonly, of White-heart Hickory; at Philadelphia and Baltimore, and in Virginia, that of Common Hickory is the only one in use. The French of Illinois call it *Noyer dur*, or Hard Walnut. The first of these denominations, which is descriptive of the fruit, I have for that reason adopted.

This species is not, as the name which it bears in that country would indicate, more multiplied in Pennsylvania, and farther south, than the other Hickories. I have not seen it north of Portsmouth in New Hampshire; though 100 miles south, in the neighborhood of Boston and Providence, it is common. It is most abundant in the forests that still remain on the coast of the Middle States and in those which cover the upper parts of the Carolinas and of Georgia; but in the last-mentioned States

e other


r husks.

villosis,
us, eximiè
â, duris-

NUTT.


Hudson,
pecies is
mmonly,
e, and in
e. The
The first
e, I have

country
farther
north of
h, in the
n. It is
coast of
parts of
d States





Mocker Nut Hickory
Juglans tomentosa



it becomes more rare in approaching the sea, as the sterility of the soil, in general dry and sandy, is unpropitious to its growth. I have noticed, however, that this is the only Hickory which springs in the pine-barrens: the sprouts are burnt every year, and never rise higher than three or four feet. I have made the same observation in traversing the Big Barrens of Kentucky and Tennessee, where the Mockernut Hickory and Black Jack Oak alone are seen. They survive the conflagrations which almost every spring envelop the prairies; but their vegetation is checked by the fire, and they do not exceed the height of eight or ten feet.

Like most of the Walnuts, the Mockernut Hickory flourishes in rich soils, and chiefly on the gentle acclivities which surround the swamps, where it grows mingled with the Sweet Gum, Poplar, Sugar Maple, Bitternut Hickory, and Black Walnut. In these situations it reaches its greatest size, which is commonly about sixty feet in height and eighteen or twenty inches in diameter. I remember to have seen larger Mockernut Hickories near Lexington in Kentucky; but this extraordinary growth in several species of trees is rarely seen on this side of the Alleghany, and is attributable to the extreme fertility of the soil in the Western country. Of all the Hickories, however, the Mockernut succeeds best on lands of a middling quality; for it forms a part of the waste and impoverished forests which cover the meager, sandy soil of Lower Virginia, though under these disadvantages it exhibits but a mean and stunted appearance.


The buds of this species are large, short, of a grayish white, and very hard: in the winter, after the falling of the leaf, they afford the only characteristic by which the tree can be distinguished when it exceeds eight or ten feet in height. In the beginning of May the buds swell, the external scales fall off, and the inner ones soon after burst and display the young leaf. The leaves grow so rapidly that I have seen them gain twenty inches in eighteen days. They are composed of four pair of

sessile leaflets and terminated by an odd one. The leaflets are large, oval-acuminate, slightly serrate, odorous, pretty thick, and hairy underneath, as is also the common petiole to which they are attached. With the first frost the leaves change to a beautiful yellow, and fall soon after. The male flowers appear on pendulous, downy, axillary aments, six or eight inches long; the female flowers, which are not very conspicuous, are of a pale rose-color and are situated at the extremity of the young shoots.

The fruit is ripe about the 15th of November. It is odorous, sessile or rarely pedunculated, and commonly united in pairs. In form and size it exhibits remarkable varieties: on some trees it is round, with depressed seams, on others oblong, with angular or prominent seams; it is sometimes two inches long and twelve or fifteen lines in diameter, and sometimes of less than half this size. It differs also in weight as well as in configuration and volume, varying from one drachm to four. The largest nuts might be confounded with those of the Thick Shellbark Hickory, and the smallest with those of the Pignut Hickory: I have selected for the drawing a nut of the most common size. The shell is very thick, somewhat channelled, and extremely hard. The kernel is sweet, but minute, and difficult to extract, on account of the strong partitions which divide it: hence, probably, is derived the name of Mockernut, and hence, also, this fruit is rarely seen in the markets.

The trunk of the old Mockernut Hickory is covered with a thick, hard, and rugged bark. Its wood is of the same color and texture with the other Hickories, and characterized by the qualities which render this class of trees so remarkable. It is particularly esteemed for fuel, for which use trees of six or eight inches in diameter are preferred. At this stage of its growth, while the heart, the proper color of which is reddish, is not yet developed, it frequently goes by the name of White-heart Hickory. In the country a greenish color is sometimes extracted from the bark; but it is not extensively in use.

fillets are
thick, and
which they
to a beau-
appear on
ches long;
e of a pale
ng shoots.
s odorous,
in pairs.
some trees
with angu-
long and
less than
configura-
four. The
the Thick
the Pignut
of the most
channelled,
minute, and
ions which
Mockernut,
ets.
ered with a
same color
rized by the
kable. It is
es of six or
stage of its
is reddish, is
ne of White-
is sometimes
in use.



Woods del.

Walters sc.

Shell bark Hickory
Anglans squamosa.



Tabac. 36.



Of all the Hickories this species is of the slowest growth,—a fact which I have proved by planting nuts of the several species and by comparing the length of their annual shoots. I have also been led to believe that it is the most liable to be attacked by worms, and especially by the *Callidium flexuosum*, whose larva eats within the body of the tree. These considerations appear sufficiently weighty to induce cultivators, in forming large plantations, to prefer some of the species which are described in the sequel.

PLATE XXXV.

*A leaf of the third of its natural size. Fig. 1. A nut with its husk. Fig. 2.
A nut without its husk. Fig. 3. Callidium flexuosum.*

SHELLBARK HICKORY.

JUGLANS SQUAMOSA. J. foliis quinis, majoribus, longè petiolatis, ovato-acuminatis, serratis, subtus villosis, impari sessili; amentis masculis, compositis, glabris, filiformibus: fructu globoso, depresso, majore; nuce compressâ albâ.

Carya alba. NUTT.

THE singular disposition of the bark in this species has given rise to the descriptive names of Shellbark, Shagbark, and Scalybark Hickory, the first of which, as being most generally in use in the Middle and Southern States, I have adopted. Many descendants of the Dutch settlers, who inhabit the parts of New Jersey near the city of New York, call it *Kisky Thomas nut*, and the French of Illinois know it by the name of *Noyer tendre*, or Soft Walnut.

Beyond Portsmouth, in New Hampshire, I have not observed

the Shellbark Hickory; and even there, its vegetation being impeded by the rigor of the climate, its stature is low and its fruit small. I have not found it in the forests of the district of Maine,* nor in those of Vermont, situated a little higher toward the north. It abounds on the shores of Lake Erie, about Geneva in Genesee, along the river Mohawk, in the neighborhood of Goshen in New Jersey, and on the banks of the rivers Susquehanna and Schuylkill in Pennsylvania. In Maryland, in the lower parts of Virginia, and in the other Southern States, it is less common. In South Carolina I have not noticed it nearer Charleston than the parish of Goose Creek, about twenty-four miles distant. It is met with in the Western States, but not as frequently as the following species,—the Thick Shellbark Hickory, to which it bears a striking analogy, and with which it is confounded by the inhabitants. East of the Alleghanies the Shellbark Hickory grows almost exclusively about swamps and wet grounds, which are exposed to be inundated for several weeks together: in these situations it is found in company with the Swamp White Oak, Red-flowering Maple, Sweet Gum, Buttonwood, and Tupelo. Of all the Hickories this species grows to the greatest height with proportionally the smallest diameter, for it is sometimes seen eighty or ninety feet high and less than two feet thick. The trunk is destitute of branches, regularly shaped, and of an almost uniform size for three-quarters of its length, thus forming a very fine tree. The greatest peculiarity in its appearance, and that by which it is most easily distinguished, is the surface of the trunk. The exterior bark is divided into a great number of long, narrow plates, which bend outward at the ends and adhere only in the middle. Bristling in this manner with projecting points, the Shellbark Hickory attracts the attention of the most careless observer. This remarkable exfoliation of the epidermis takes place only

* [The Shellbark Hickory is found in the county of York, in Maine. EMERSON.]

in trees which exceed ten inches in diameter, though it is much earlier indicated by seams. This characteristic, by which the tree may be recognised in winter when stripped of its leaves, does not exist during the seven or eight first years of its growth; and during this period it may easily be confounded with the Mockernut Hickory and Pignut Hickory, if recourse is not had to the buds. In these two species, and generally in all trees, the buds are formed of scales closely applied one upon another; in the species which we are considering, the two external scales adhere for only half the length of the bud and leave the upper part uncovered. It is my opinion that in this disposition of the scales, which is peculiar to this and the following species, should be sought the origin of the exfoliation of the bark. When the sap begins to ascend in the spring, the outer scales fall, and the inner ones swell and become covered with a yellowish silky down: after a fortnight, the buds, which are already two inches long, open and give birth to the young leaves. The growth of the leaves are so rapid, that in a month they attain their full length, which on young and vigorous trees is sometimes twenty inches. They consist of two pair of leaflets with a sessile odd one. The leaflets are very large, oval-acuminate, serrate, and slightly downy underneath. The male flowers, which in the State of New York appear from the 15th to the 20th of May, are disposed, as in the preceding species, on long, glabrous, filiform, pendulous aments, of which three are united on a common peduncle, attached at the base of the young shoots; the female flowers, of a greenish hue, and scarcely apparent, are situated at the extremity. The fruit of the Shellbark Hickory is ripe about the beginning of October. Some years it is so abundant that several bushels may be gathered from a single tree. It varies in size, according to the soil and the exposure in which it is produced; but five and a half inches may be assumed as the average of its circumference. The shape is uniformly round, with four depressed seams, in which the husk opens at

the season of perfect maturity, dividing itself completely into equal sections. The entire separation of the husk, and its thickness disproportioned to the size of the nut, form a character peculiar to the Shellbark Hickories. The nuts of this species are small, white, compressed at the sides, and marked by four distinct angles, which correspond to the divisions of the husk.

The Shellbark nut contains a fuller and sweeter kernel than any American Walnut except the Pecanutt. The shell, though thin, must be cracked before being brought upon the table, as it is too hard to be crushed in the fingers like the European Walnut, which is certainly a superior fruit. These nuts are in such request that they form a small article of commerce, registered on the list of exports of the products of the United States. This exportation, which does not exceed four or five hundred bushels annually, takes place from New York, and from the small ports of Connecticut, to the Southern States, to the West India Islands, and even to Liverpool, where the fruit is known by the name of Hickory nuts. In the market of New York they are sold at two dollars a bushel. They are gathered in the forests, and from insulated trees which in some places have been spared in clearing the lands,—a precaution which I have particularly noticed to have been used near Goshen in New Jersey, and on several estates about thirty miles beyond Albany.

The Indians who inhabit the shores of Lakes Erie and Michigan lay up a store of these nuts for the winter, a part of which they pound in wooden mortars, and, boiling the paste in water, collect the oily matter which swims upon the surface, to season their food.

Before speaking of the properties of the wood, I cannot forbear mentioning a fine variety of Shellbark nuts produced upon a farm at Seacocus, near Snakehill in New Jersey. They are nearly twice as large as any that I have seen elsewhere, and have a white shell with rounded prominences instead of angles.

A century of cultivation, perhaps, would not advance the species generally to an equal degree of perfection, and probably this variety might still be improved by grafting.

The wood of the Shellbark Hickory possesses all the characteristic properties of the Hickories, being strong, elastic, and tenacious. It has also their common defects of soon decaying and of being eaten by worms. As this tree stretches up to a great height with nearly a uniform diameter, it is sometimes employed at New York and Philadelphia for the keels of vessels; but it is now seldom used for this purpose, most of the large trees near the seaports being already consumed. Its wood is found to split most easily and to be the most elastic; for this reason it is used for making baskets, and also for whip-handles, which are esteemed for their suppleness, and of which several cases are annually exported to England. For the same excellence, and for the superior fineness of its grain, it is selected in the neighborhood of New York and Philadelphia for the back-bows of Windsor chairs, which are wholly of wood. I have frequently observed, that among the Hickory wood brought to New York for fuel, this species predominated.

Such are the uses to which the Shellbark Hickory appears peculiarly adapted. It has before been seen to be a tree of lofty stature and majestic appearance: I should therefore recommend its introduction into the European forests, where it should be consigned to cool and humid places, congenial with those in which it flourishes in America. In the north of Europe it could not fail of succeeding, as it securely braves the most intense cold.

PLATE XXXVI.

Fig. 1. A nut with its husk. Fig. 2. A section of the husk. Fig. 3. A nut without its husk. Fig. 4. A barren ament divided into three parts.

THICK SHELLBARK HICKORY.

JUGLANS LACINIOSA. *J. foliis majoribus, foliolis 7-9^{nis}, orato-acuminatis, serratis, subtomentosis, impari, petiolato: fructu majore, orato; nucce oblongâ, crassâ, medioeriter compressâ.*

Carya sulcata. NUTT.

THIS species bears a striking analogy to the preceding, and is frequently confounded with it by the inhabitants of the Western country: some of them distinguish it by the name of Thick Shellbark Hickory, which should be preserved as its appropriate denomination. East of the Alleghanies, this tree is rare and is found only in a few places; it grows on the Schuylkill River thirty or forty miles from its junction with the Delaware, and in the vicinity of Springfield, fifteen or twenty miles from Philadelphia, where its fruit is called Springfield nut. It is also found in Gloucester county, in Virginia, under the name of Gloucester Walnut. These different denominations confirm my observation that this species is little multiplied on the eastern side of the Alleghany Mountains,—a fact of which I became assured in travelling through the country. It abounds, on the other hand, in the bottoms which skirt the Ohio and the rivers which empty into it, where it unites with the Honey Locust, Black Maple, Hackberry, Black Walnut, Wild Cherry, White and Red Elm, Box Elder, White Maple, and Buttonwood, to form the thick and gloomy forests which cover these valleys. Like the Shellbark Hickory, it grows to the height of eighty feet, and its ample head is supported by a straight trunk, in diameter proportioned to its elevation. The bark exhibits the same singular arrangement with that of the Shellbark Hickory: it is divided into strips from one to three feet long, which are warped outward at the end and attached only at the middle. They fall, and are succeeded by others similarly disposed. It


Y.


acuminatis,
erato; nuce

. Nutt.

ng, and is
e Western
of Thick
ppropriate
are and is
kill River
aware, and
from Phi-

It is also
e name of
onfirm my
the eastern
. I became
nds, on the
d the rivers
ney Locust,
erry, White
onwood, to
ese valleys.
t of eighty
t trunk, in
exhibits the
k Hickory:
; which are
the middle.
isposed. It





Brown del.

1850 del.

Thick Shell bark Hickory.
Juglans laciniata.

is only observable that in this species the plates are narrower, more numerous, and of a lighter color; from which differences I have thought proper to give it the specific name of *laciniosa*. The outer scales of the buds do not adhere entirely to the inner ones, but retire as in the Shellbark Hickory. The leaves also, which vary in length from eight to twenty inches, observe the same process in unfolding, and are similar in size, configuration, and texture; but they differ in being composed of seven leaflets and sometimes of nine, instead of five, the invariable number of the Shellbark Hickory. The male aments are disposed in the same form, though they are, perhaps, a little longer than in the other species. The female flowers appear, not very conspicuously, at the extremity of the shoots of the same spring. They are succeeded by a large oval fruit, more than two inches long and four or five inches in circumference. Like that of the Shellbark Hickory, it has four depressed seams, which, at its complete maturity, open through their whole length for the escape of the nut. The nut of this species is widely different from the other; it is nearly twice as big, longer than it is broad, and terminated at each end in a firm point. The shell is also thicker and of a yellowish hue, while that of the Shellbark nut is white.

From the color of its nut, the Shellbark Hickory received the specific name of *alba*, which I have changed, as it indicates a character possessed by it in common with another species found in the Royal Gardens of the *Petit Trianon*. This species, originally from North America, belongs to the Scalybark Hickories. The nuts are white, and the entire fruit, though a little inferior in size, resembles that of the proper Shellbark Hickory. By its foliage it is related to the Thick Shellbark Hickory, each leaf being composed of four pair of leaflets with an odd one. The specific name of *ambigua* might with propriety be given to it.

The nuts of the Thick Shellbark Hickory are brought every

autumn to the market of Philadelphia, but the quantity does not exceed a few bushels, and they are generally sold mixed with those of the Mockernut Hickory, which resemble some varieties of this species. The Gloucester Hickory I consider only as a variety of the Thick Shellbark Hickory, to which it bears the strongest resemblance in its young shoots, in the number of its leaflets, and in its barren aments. The only essential difference is in the nuts; those of the Gloucester Walnut are a third larger, with the shell one-half thicker, and so hard that it requires pretty heavy blows of a hammer to crack them. In color they resemble the nuts of the Mockernut Hickory, with the finest varieties of which they might, from this circumstance, be confounded.

The Thick Shellbark Hickory, as has been said, is nearly related to the Shellbark Hickory; and its wood, which is of the same color and texture, unites the peculiar qualities of that species with such as are common to the Hickories. Its fruit, though larger, is inferior in taste; and this consideration should induce proprietors in the Western country, in clearing their new lands, to spare the true Shellbark Hickory in preference when both species are found upon the same soil. For the same reason, and for its favorable growth in less fertile grounds, and even in elevated situations,—a fact which I have observed near Brownsville on the Monongahela River,—the same preference should, I think, be given to it in the forests of Europe.

In the description of the Scalybark Hickories, it has been seen that they exhibit many striking traits of resemblance, which may warrant the grouping of them into a secondary section. Besides their generic and specific characters, they possess others peculiar to themselves, by which they are so nearly related that, were it not for some remarkable differences, they might be treated as a single species. The general characters of the Hickories are, three-clefted, pliable, and pendulous male aments, and certain common properties of the wood. To these are

added, in the Scaly Hickories, a very thick husk, covering the nut completely, and divided into four parts when ripe; a shaggy bark on the trunk, indicated, in my opinion, by the external scales of the buds not adhering to those beneath; and leaves composed of very large leaflets of a uniform shape and texture. In comparing the three species with each other, essential differences are observed. The Shellbark Hickory, for instance, and the *Juglans ambigua*, are constantly distinguished by the number of leaflets, which is always five in the first species and nine in the last. The nuts and the entire fruit, on the other hand, are so much alike, that they might be mistaken for the product of the same tree: the fruit of both is round, with depressed seams, and the nuts are similarly moulded and equally white. If, on a more attentive examination, the Gloucester Hickory is determined to be a distinct species from the Thick Shellbark Hickory, it will be observed that they resemble each other in their leaves, composed of seven and sometimes of nine leaflets, and in the luxuriant force of their vegetation, but that they differ in their fruit, which in the Thick Shellbark Hickory is oblong, with a compressed nut, like that of the Shellbark Hickory, of twice the size, and of a yellowish color, and in the Gloucester Hickory spherical and very large, with a large grayish-white nut, nearly round, whose shell is two lines thick and extremely hard. In fine, it is to be remarked that the species and the variety of the Scalybark Hickory which have been described grow, or at least are most abundantly multiplied, in regions far remote from each other.

PLATE XXXVII.

A leaf of one-third of its natural size. Fig. 1. A section of the husk.

Fig. 2. Nuts.

PIGNUT HICKORY.

JUGLANS PORCINA. *J. foliolis 5-7^{mis}, ovato-acuminatis, serratis, glabris; amantibus masculis compositis, filiformibus, glabris; fructu pyriformi vel globoso; nucis minimâ, levi, durissimâ.*


Carya porcina. NUTT.

THIS species is generally known in the United States by the name of Pignut and Hognut Hickory; sometimes also by that of Broom Hickory. The first of these names is most commonly in use; the others are known only in some districts of Pennsylvania, and particularly in the county of Lancaster. Portsmouth in New Hampshire may be considered as limiting, toward the north, the climate of this tree. A little farther south it is abundant, and in the Atlantic parts of the Middle States it helps, with the Mockernut Hickory, White Oak, Swamp White Oak, Sweet Gum, and Dogwood, to form the mass of the forests. In the Southern States, especially near the coast, it is less common in the woods, being found only on the borders of the swamps and in places which are wet without being absolutely marshy or exposed to be long inundated. This tree is met with in the Western country, but less frequently, I believe, than the Thick Shellbark and Mockernut Hickories. I have observed that the last-mentioned species grows wherever the Pignut is found, but that the Pignut does not always accompany the Mockernut, which is satisfied with a less substantial soil. This remark I have made more particularly in the lower parts of Virginia, of the two Carolinas, and of Georgia. It appears, then, that, with the exception of the States of Vermont and New Hampshire, of the district of Maine, of the Genesee country, and of the cold and mountainous tracts along the whole range of the Allegheny Mountains, this tree is more or less abundant in the forests throughout the United States.

glabris;
formi vel


NUTT.

s by the
by that
commonly
of Penn-
. Ports-
g, toward
outh it is
it helps,
hite Oak,
cests. In
s common
e swamps
marshy or
th in the
the Thick
l that the
found, but
Lockemat,
remark I
irginia, of
that. with
mpshire, of
of the cold
Allegheny
the forests





Pignut Hickory
Juglans pycnantha



The Pignut Hickory is one of the largest trees of the United States. It grows to the height of seventy or eighty feet, with a diameter of three or four feet. In the winter, when stripped of its leaves, it is easily known by the shoots of the preceding summer, which are brown, less than half as large as those of the Mockernut and Shellbark Hickories, and terminated by small oval buds. At this season it is easy also to distinguish the Bitternut Hickory by its naked and yellow buds. The buds of this species, as in the other Hickories with scaly buds, are more than an inch in length a few days before their unfolding. The inner scales are the largest, and of a reddish color. They do not fall till the leaves are five or six inches long. The leaves are compound, and vary in size and in the number of leaflets, according to the moisture and fertility of the soil. In rich grounds, they are eighteen inches long; and the complete number of leaflets is three pairs with an odd one. The leaflets are four or five inches long, acuminate, serrate, nearly sessile, and glabrous or smooth on both sides. On vigorous trees, which grow in shady exposures, the petiole is of a violet color.

The male aments are smooth, filiform, flexible, and pendulous: they are two inches long, and in their arrangement resemble those of the other Hickories. The female flowers are greenish, not very conspicuous, and situated at the extremity of the shoot: the fruit succeeds them in pairs as often as singly. The husk is thin and of a beautiful green: when ripe, it opens through half its length for the passage of the nut. The nut is small, smooth, and very hard on account of the thickness of the shell. Its kernel is sweet but meagre, and difficult to extract, from the firmness of the partitions. These nuts are never carried to market, but serve for food to swine, raccoons, and the numerous species of squirrels which people the forests.

In the Pignut Hickory, the form and size of the nuts vary more than in the other species. Some are oval, and when covered with their husks resemble young figs; others are

broader than they are long, and others are perfectly round. Among these various forms, some nuts are as large as the thumb and others not bigger than the little finger. Although the same tree yields fruit of the same form every year, I cannot, after an attentive examination of the young shoots and of the aments, consider these differences in any other light than as varieties. The two most remarkable of them are described in the new edition of the *Species Plantarum*, by Willdenow, as distinct species. That with oblong fruit is called *Juglans glabra*, and that with round fruit and a husk somewhat rough, *Juglans obcordata*. Dr. Muhlenberg admits this distinction; but, with all the deference which I owe to his botanical knowledge, I cannot adopt his opinion.

The wood of the Pignut Hickory resembles that of the other species in the color of its sap and of its heart: it possesses also their excellencies and their defects. I have conversed with wheelwrights in the country, who affirmed that it is the strongest and the most tenacious of the Hickories, and who, for that reason, preferred it to any other for axle-trees and axe-helves. These considerations lead me to recommend its introduction into the forests of Europe, where its success would be certain.

PLATE XXXVIII.

A branch with its leaves of one-third of the natural size. Fig. 1. A nut with its husk, (oblong variety.) Fig. 2. A nut without its husk. Fig. 3. A nut with its husk, (round variety.) Fig. 4. A nut without its husk.

y round.
e thumb
ough the
I cannot,
nd of the
e than as
cribed in
w, as dis-
glabra,
a, *Juglans*
but, with
nowledge, I

the other
esses also
rsed with
he strong-
o, for that
axe-helves.
roduction
e certain.

Fig. 1. *A nut*
usk. Fig. 3.
t its husk.




Nutmeg Hickory Nut .
Juglans myristiciformis .

Boiss. & Hausskn.

Pl. 30.



Boeam. Sculp.



NUTMEG HICKORY.

JUGLANS MYRISTICÆFORMIS. *J. foliis quinis, foliolis ovato-acuminatis, serratis, glabris: fructu ovato, scabriusculo; nucce minimâ, durissimâ.*

Carya myristicaeformis. NUTT.

No specific denomination has hitherto been given to this species by the inhabitants of the Middle States, to which it is peculiar: that of Nutmeg Hickory, which I have formed, appears sufficiently appropriate, from the resemblance of its nuts to a nutmeg.



I have not myself found this tree in the forests, and hence I conclude that it is not common. It is true I had not, at the period of my residence in that part of the United States, conceived the design of the present work, and did not devote myself entirely to the researches which have since given birth to it. I am acquainted with the Nutmeg Hickory only by a branch and a handful of nuts given me at Charleston, in the fall of 1802, by the gardener of Mr. H. Izard, which he had gathered in a swamp on his master's plantation of the Elms, in the parish of Goose Creek. From this specimen alone I have included the tree among the Hickories.

The leaves, which are composed of four or six leaflets with an odd one, are symmetrically arranged. I remarked also that the shoots of the preceding year were flexible and coriaceous.



The nuts are very small, smooth, and of a brown color, marked with lines of white: the husk is thin, and somewhat rough on the surface. The shell is so thick that it constitutes two-thirds of the volume of the nut, which, consequently, is extremely hard and has a minute kernel. The fruit is inferior even to the Pignut.

I suspect that the Nutmeg Hickory is more common in Lower





**IMAGE EVALUATION
TEST TARGET (MT-3)**



**Photographic
Sciences
Corporation**

23 WEST MAIN STREET
WEBSTER, N.Y. 14590
(716) 672-4502

1.5
1.6
1.8
1.9
2.0
2.2
2.5

1.5
1.6
1.8
1.9
2.0
2.2
2.5

Louisiana:* it belongs to inquirers who engage in researches analogous to those which I have pursued in the Atlantic and Western States, to study this tree more fully than I have been able to do, and to complete the imperfect description which I have given of it.

PLATE XXXIX.

A branch and nuts with their husks. Fig. 1. A nut without its husk.

RECAPITULATION
OF THE
PROPERTIES AND USES
OF
HICKORY WOOD.

In the summary introduction to the History of the Walnuts • of North America, it was remarked, that those of the second section, or the Hickories, exhibit great variations in the size and shape of their fruit, in the number of leaflets which compose their leaves, and in their general appearance, from the effect of soils of different degrees of moisture. Hence result, in many cases, mutual resemblance so striking, that a person not familiar with this class of trees might easily confound distinct species, or

* In the interesting work of Mr. W. Darby on Louisiana, published at Philadelphia in 1817, the Nutmeg Hickory is said to abound on the waters of Red River in the Mississippi Territory.

describe as different species what are mere varieties. On taking off the epidermis, or dead part, the same organization is observed in the bark of all the Hickories. In other trees the fibrous and the cellular tissue are confounded; here, on the contrary, they are separate, and the fibrous is regularly disposed in the form of lozenges, which are smaller in young trees than in such as are more fully grown. An arrangement so peculiar and remarkable has a beautiful effect, and great advantage might be taken of it in cabinet-making, if this bark was not, like other species, liable to warp. It affords, nevertheless, an interesting object in vegetable physiology. So close an analogy exists in the wood of these trees, that, when stripped of the bark, no difference is discernible in the grain, which is coarse and open in all, nor in the color of the heart, which is uniformly reddish. To these conspicuous properties are added others worthy of remark, which, as has been observed, though modified in the several species, are possessed by them all in a higher degree than by any other tree of the same latitude in Europe or America. These are—great weight, strength, and tenacity, a speedy decay when exposed to heat and moisture, and peculiar liability to injury from worms. According to these prominent excellencies and defects, the uses of their wood are pretty well determined, and to these uses they are indiscriminately applied.

Hickory timber is employed in no part of the United States in the building of houses, because, as has been before observed, it is too heavy, and soon becomes worm-eaten. But if its defects forbid its employment in architecture, its good qualities, on the other hand, adapt it to many secondary uses, which could not be as well subserved by any other wood. Throughout the Middle States it is selected for the axle-trees of carriages, for the handles of axes and other carpenters' tools, and for large screws, particularly those of bookbinders' presses. The cogs of mill-wheels are made of Hickory heart thoroughly seasoned; but it is proper only for such wheels as are not exposed to moisture;

researches
antic and
ave been
which I

its husk.

ES

the Walnuts
the second
the size and
ch compose
the effect of
lt, in many
not familiar
t species, or

ished at Phila-
waters of Red
F. A. M.

and for this reason some other wood is, by many millwrights, preferred. The rods which form the back of Windsor chairs, coach-whip handles, ramrods, rake-teeth, flails for thrashing grain, and the bows of ox-yokes,—all these are objects ordinarily made of Hickory. At Baltimore it is used for the hoops of sieves, and is more esteemed than the White Oak, which is equally elastic, but more apt to peel off in small shreds into the substance sifted. In the country near Augusta in Georgia I have remarked that the common chairs are of Hickory wood. In New Jersey it is employed for shoeing of sledges,—that is, for covering the runners or parts which slide upon the snow; but to be fit for this use, it must have been cut long enough to have become perfectly dry.

Of the numerous trees of North America east of the Alleghany Mountains, none except the Hickory is perfectly adapted to the making of hoops for casks and boxes. For this purpose vast quantities of it are consumed at home, and exported to the West India Islands. The hoops are made of young Hickories from six to twelve feet high, without choice as to the species. The largest hoop-poles sold at Philadelphia and New York in February, 1808, at three dollars a hundred. Each pole is split into two parts, and the hoop is crossed and fastened by notches, instead of being bound at the end with twigs, like those made of Chestnut. From the solidity of the wood, this method is sufficiently secure.

When it is considered how large a part of the productions of the United States is packed for exportation in barrels, an estimate may be formed of the necessary consumption of hoops. In consequence of it, young trees proper for this object have become scarce in all parts of the country which have long been settled. The evil is greater as they do not sprout a second time from the same root, and as their growth is slow. The cooper cannot lay up a store of them for future use; for, unless employed within a year, and often six months, after being cut, they

are attacked by two species of insect; one of these, which eats within the wood, and commits the greatest ravages, is represented on the plate of the Mockernut Hickory, the wood of which species I have observed to be peculiarly liable to its attacks.

The defects which unfit the Hickory for use in the building of houses equally exclude it from the construction of vessels. At New York and Philadelphia, the Shellbark and Pignut Hickories have been taken for keels, and are found to last as long as those of other wood, owing to their being always in the water. Of the two species, the Pignut would be preferable as being less liable to split; but it is rarely found of as large dimensions as the other.

In sloops and schooners, the rings by which the sails are hoisted and confined to the mast are always of Hickory. I have also been assured, that for attaching the cordage it makes excellent pegs, which are stronger than those of Oak; but they should set loosely in the holes, as otherwise, for want of speedily seasoning, they soon decay. For handspikes the Hickory is particularly esteemed on account of its strength: it is accordingly employed in most American vessels, and is exported for the same purpose to England, where it sells from 50 to 100 per cent. higher than Ash, which is brought also from the north of the United States. The Hickories are cut without distinction for this use; but the Pignut, I believe, is the best.

All the Hickories are very heavy, and in a given volume contain a great quantity of combustible matter. They produce an ardent heat, and leave a heavy, compact, and long-lived coal. In this respect, no wood of the same latitude, in Europe or America, can be compared to them: such, at least, is the opinion of all Europeans who have resided in the United States. At New York, Philadelphia, and Baltimore, people in easy circumstances burn no other wood; and, though it is sold 50 per cent. higher than Oak, it is found profitable in use. It sold at

New York, the 20th of October, 1807, at fifteen dollars a cord, and Oak wood at ten dollars. From its superior quality, the Hickory is always sold separately. I have noticed that, at New York, the Shellbark predominated in the fuel, and at Philadelphia and Baltimore, the Mockernut. At Baltimore, the Shellbark, easily recognised by its scaly bark, is never seen.

The quantity of the respective species of Hickory consumed in the cities is regulated by a soil and climate more favorable to one than another, and not by an opinion entertained of their comparative excellence; though experience shows the Mockernut to be the best, and the Bitternut the poorest. This difference, however, is too slight to be generally regarded.

Of the uses to which the Hickory is devoted in the United States, two will principally contribute, together with the slowness of its growth, to its entire extermination: these are the cutting of the saplings for hoops and of the trees for fuel. These considerations, independently of many accessory causes, which hasten the destruction of the forests in this part of the New World, lead me to believe that in less than fifty years they will not furnish a tenth part of the hoops demanded in commerce. Hence arise motives sufficiently powerful to engage proprietors, who seek to preserve their forests and to augment their value, to multiply in them the most useful trees, and especially the Hickories. The object might be fully attained by planting the nuts, previously made to germinate in boxes filled with earth, and kept moist in the cellar. The success of this simple method is certain. It would be advantageous, also, to plant a greater number than the soil can sustain, that, when the poles are an inch in diameter, a part of them may be cut for hoops,* while the rest are left to grow for fuel, or for other uses to which the Hickory is appropriate.

* [Or for walking-sticks, for which the consumption is considerable, and the demand constantly increasing. EMERSON: "Trees and Shrubs of Massachusetts."]

It has been seen, by what precedes, that, though the Hickory wood has essential defects, they are compensated by good properties which render it valuable in the arts, and which entitle it to the attention of Europeans,—above all, as a combustile. Though its growth is slow during its early years, it should form a part of our forests. But I doubt whether this can be effected except by planting nuts in the woods; for the trees, even when very young, with difficulty survive transplantation. Before they are three lines in diameter and eighteen inches tall, they have a tap-root three feet long and destitute of fibres. Hence, it has happened that, of more than a hundred thousand young plants produced by nuts which I have at different times sent to France, very few are found alive. They have perished in the removal from the nursery, or in the second transplantation to the place of their ultimate destination. The Black Walnut and Bitternut, on the contrary, whose roots do not descend deeply and are plentifully garnished with fibres, easily recover after transplantation, even when six or eight feet high at the time of their removal.

In concluding this article, I recommend particularly for propagation in European forests the Shellbark Hickory and the Pignut Hickory, whose wood unites in the highest degree the valuable properties of the group. I think, also, that the Pecan nut merits attention from promoters of useful culture, not so much for its wood as for its fruit, which is excellent, and more delicate than that of the European Walnut. It might probably be doubled in size if the practice was successfully adopted of grafting this species upon the Black Walnut or upon the Common European Walnut.

M A P L E S.

OF the species which compose this genus, the number known is already considerable, and will probably be augmented by the future researches of Botanists, especially on the continent of North America.

The Maples, in general, are lofty and beautiful trees. One of their principal characters consists in opposite leaves divided into several very distinct lobes. Capable of enduring an intense degree of cold, they form, in the north of the Old and of the New Continent, extensive forests, which, with those of the Beech, appear to succeed the Spruce, the Larch, and the Pine, and to precede the Chestnut and the Oak. Such, at least, seems to be in America, between the 43d and 46th degrees of latitude, the place assigned by nature to the true Sugar Maple.

The species of Maples hitherto described amount to fourteen, of which seven belong to Europe and seven to North America. Among these last I have not included the Dwarf Red Maple, *Acer coccineum*, which is a diminutive species, and concerning which I do not possess adequate materials for a description. It abounds in Nova Scotia, and has always been confounded with the proper Red Maple; I have also observed it in the upper parts of New Hampshire. It scarcely exceeds twelve or eighteen feet in height, and its flowers and seeds are of a more vivid red than those of the Red Maple. The Black Sugar Maple grows to about the same height with the Sugar Maple, but it is plainly a distinct species. In the collection of dried plants made by Captains Lewis and Clarke during their journey to the South

Sea, I saw specimens of a beautiful Maple from the banks of Columbia River.

From this brief summary it results that the North American species are more numerous than those of Europe. The wood of the Maples differs so widely in quality in different species, that it becomes difficult to characterize it by general observations: it may be remarked, that it speedily ferments and decays when exposed to the weather, that it is liable to be injured by worms, and that hence it is unfit for building. It possesses properties, however, which compensate in part for these defects, and which render it useful in the arts and in domestic economy. For more particular information I must refer the reader to the descriptions of the respective species.

[*Propagation and culture.* The *Aceráceæ* prefer a free, deep, loamy soil, rich rather than sterile, and neither wet nor very dry. The situation that suits them best is one that is sheltered, and shady rather than exposed. They are seldom found on the north sides of lofty mountains, or on mountains at all, except among other trees; but in the plains they are found by themselves. Though the species only attain perfection in favorable soils and situations, they will spring up and live in any situation whatever.

They are chiefly propagated from seeds; but some sorts are increased by layers, cuttings of the shoots or roots, or by budding or grafting. The seeds of most of the species ripen in October, and may be gathered by hand, or shaken from the tree, when the keys begin to turn brown. The maturity of the seed may be proved by opening the key and observing if the cotyledons are green, succulent, and fresh; if the green color is wanting, the seeds are good for nothing. The seeds may be sown either in autumn or in spring; the latter is preferable where moles abound, as they are very fond of the seeds. Sown in spring, they come up in five or six weeks, with the exception of *Acer*

campestre, which never grow till the second or third year. The seeds should not be covered with more than from a quarter to half an inch of soil. The surface of the ground in which they are sown may be advantageously shaded with leaves, fronds of firs, or straw.

The *Acer argenteum*, or Silvery-leaved, and *Acer rubrum*, or Scarlet Maple, perfect their seeds in May; and these should be sown immediately after having been collected: they will vegetate directly, and produce fine plants the first season, if kept free from weeds. The seeds of the former do not keep well till spring.]*

* [For a large additional list of Maples, see Nuttall's Supplement, vol. ii. p. 24 *et seq.* Many of these are deserving of the attention of our planters; especially the large-leaved Maple, sometimes ninety feet high, with leaves nearly a foot in diameter, affording an impervious and complete shade. See also Emerson's "Trees and Shrubs of Massachusetts," p. 481.]

ar. The
quarter to
which they
fronds of

brum, or
should be
will vege-
n, if kept
o well till

vol. ii. p. 24
; especially
rly a foot in
son's "Trees

METHODICAL DISPOSITION
OF THE
MAPLES OF NORTH AMERICA,
INCLUDING
TWO EUROPEAN SPECIES.

Polyandria dioecia, LINN. *Acera*, JUSS.

FIRST SECTION.

Sessile flowers. (Fructification vernal.)

1. White Maple *Acer eriocarpum.*
2. Red-flowering Maple *Acer rubrum.*

SECOND SECTION.

Pedunculated flowers. (Fructification autumnal.)

3. Sugar Maple *Acer saccharinum.*
4. Black Sugar Maple *Acer nigrum.*
5. Norway Maple *Acer platanoides.*
6. Sycamore *Acer pseudo-platanus.*
7. Moose Wood *Acer striatum.*
8. Box Elder *Acer negundo.*
9. Mountain Maple *Acer montanum.*

WHITE MAPLE.

ACER ERIOCARPUM. *A. foliis oppositis, quinquelobis, profundè sinuatis, inæqualiter dentatis, subtis candidissimis: floribus pentandris, apetalis.*

Acer dasycarpum. EHRENBURG.

IN the Atlantic parts of the United States, this species is often confounded with the Red Maple, which it nearly resembles; west of the mountains, they are constantly distinguished, and the *Acer eriocarpum* is known by no other name than White Maple.


The banks of Sandy River in the district of Maine, and those of the Connecticut near Windsor in Vermont, are the most northern points at which I have seen the White Maple. But, like many other vegetables, it is pinched by the rigorous winters of this latitude, and never reaches the size which it attains a few degrees farther south. It is found on the banks of all the rivers which flow from the mountains to the ocean, though it is less common along the streams which water the southern parts of the Carolinas and of Georgia. In no part of the United States is it more multiplied than in the Western country, and nowhere is its vegetation more luxuriant than on the banks of the Ohio and of the great rivers which empty into it. There, sometimes alone, and sometimes mingled with the Willow, which is found along all these waters, it contributes singularly by its magnificent foliage to the embellishment of the scene. The brilliant white of the leaves beneath forms a striking contrast with the bright green above; and the alternate reflection of the two surfaces in the water heightens the beauty of this wonderful moving mirror, and aids in forming an enchanting picture, which, during my long excursions in a canoe in these

andè sinuatis,
, apctalis.

SHRENBURG.

species is
nearly resem-
bling, distinguished,
than White

, and those
the most
ample. But,
vigorous win-
ch it attains
ks of all the
though it is
thern parts
the United
country, and
the banks of
it. There,
the Willow,
es singularly
f the scene.
striking con-
ate reflection
eauty of this
an enchanting
noe in these





Boiss del

Gabriel sculp

White Maple.
Acer spicatum.



regions of solitude and silence, I contemplated with unwearied admiration. Beginning at Pittsburg, and even some miles above the junction of the rivers Alleghany and Monongahela, White Maples twelve or fifteen feet in circumference are continually met with at short distances.

The trunk of this tree is low, and divides into a great number of limbs, so divergent that they form a head more spacious than that of any other tree with which I am acquainted. It is worthy of remark, that the White Maple is found on the banks of such rivers only as have limpid waters and a gravelly bed, and never in swamps and other wet grounds enclosed in forests, where the soil is black and miry. These situations, on the contrary, are so well adapted to the Red Maple that they are frequently occupied by it exclusively. Hence the last-mentioned species is common in the lower parts of the Carolinas and of Georgia, where the White Maple is no longer seen; for as soon as the rivers, in descending from the mountains toward the ocean, reach the low country, they begin to be bordered by miry swamps covered with the Cypress, Blackgum, Large Tupelo, etc.

The White Maple blooms early in the spring: its flowers are small and sessile, with a downy *ovarium*. The fruit is larger than that of any other species which grows east of the Mississippi. It consists of two capsules joined at the base, each of which encloses one roundish seed, and is terminated by a large membranaceous, falciform wing. In Pennsylvania, it is ripe about the 1st of May, and a month earlier on the Savannah River in Georgia. At this period, the leaves which have attained half their size are very downy underneath; a month later, when fully grown, they are perfectly smooth. They are opposite and supported by long petioles; they are divided by deep sinuses into four lobes, are toothed on the edges, of a bright green on the upper surface, and of a beautiful white

beneath. The foliage, however, is scattered, and leaves an open passage to the sunbeams.

The wood of this Maple is very white, and of a fine grain; but it is softer and lighter than that of the other species in the United States, and, from its want of strength and durability, it is little used. Wooden bowls are sometimes made of it when Poplar cannot be procured. At Pittsburg, and in the neighboring towns, it serves in cabinet-making, instead of Holly, for inlaying furniture of Mahogany, Cherry-tree, and Walnut: though, as it soon changes color, it is less fitted for this purpose. The hatters of Pittsburg prefer the charcoal of this wood to every other for heating their boiler, as it affords a heat more uniform and of longer continuance. Some of the inhabitants on the Ohio make sugar of its sap, by the same process which is employed with the Sugar Maple. Like the Red Maple, it yields but half the product from a given measure of sap; but the unrefined sugar is whiter and more agreeable to the taste than that of the Sugar Maple. The sap is in motion earlier in this species than in the Sugar Maple, beginning to ascend about the 15th of January; so that the work of extracting the sugar is sooner completed. The cellular tissue rapidly produces a black precipitate with sulphate of iron.

In all parts of the United States where this tree abounds, many others are found of superior value. Its secondary consequence is evinced by the unimportant uses to which it is devoted.

In Europe, the White Maple is multiplied in nurseries and gardens. Its rapid growth affords hopes of cultivating it with profit in this quarter of the world, which is less rich in the diversity of its species. In forming plantations, more care than has hitherto been taken should be paid to the choice of the ground, which should be constantly moist, or exposed to annual inundations: in such situations its vegetation would be surprisingly beautiful and rapid.

es an open

fine grain ;
pecies in the
urability, it
of it when
e neighbor-
Holly, for
d Walnut :
his purpose.
his wood to
a heat more
inhabitants
rocess which
ed Maple, it
of sap ; but
to the taste
on earlier in
ascend about
ng the sugar
y produces a

tree abounds,
secondary con-
o which it is

nurseries and
vating it with
ss rich in the
more care than
choice of the
osed to annual
would be sur



Banks del.

Walpole sculp.

Red flowering Maple.
Acer rubrum.

1141



cauliculus




PLATE XL.

A branch with leaves of the natural size. Fig. 1. Barren flowers. Fig. 2. Fertile flowers. Fig. 3. A seed of the natural size.

[For measurements of several White Maples, see EMERSON'S "Trees and Shrubs of Massachusetts," p. 489.]

RED-FLOWERING MAPLE.

ACER RUBRUM. *A. foliis oppositis, trilobis, inaequaliter dentatis, subtus glaucis; floribus rubris, aggregatis; germine glaberrimo; umbellis sessilibus; capsulis rubris, pedunculatis.*

DIFFERENT names are given to this tree in different parts of the United States: east of the Alleghany Mountains, it is called Red-flowering Maple, Swamp Maple, and Soft Maple; in the Western country, simply Maple. The first denomination, which is most generally in use, is also most appropriate, as the young shoots, the flowers, and the fruit are red.

Toward the North, the Red-flowering Maple appears first about Malabaic in Canada, in latitude 48°; but it soon becomes more common in proceeding southward, and is found abundant to the extremities of Florida and Lower Louisiana. Of all the trees which flourish in wet grounds occasionally overflowed, this species is most multiplied in the Middle and Southern States. It occupies, in great part, the borders of the creeks, and abounds in all the swamps which are often inundated and always miry. In these situations it is accompanied by the Blackgum, Sweetgum, Shellbark Hickory, Swamp White Oak, Black Ash, and White Ash. To these are added, in the Carolinas and Georgia, the small Magnolia or Swamp Bay, the

Water Oak, Loblolly Bay, Tupelo, and Red Bay. It is a remarkable fact, that west of the mountains, between Brownsville and Pittsburg, the Red-flowering Maple is seen growing on elevated ground with the Oaks and the Walnuts. I have nowhere observed it of as ample dimensions as in Pennsylvania and New Jersey: in these States exist extensive marshes, called Maple swamps, exclusively covered with it, where it is found seventy feet high and three or four feet in diameter.

The Red-flowering Maple is the earliest tree whose bloom announces the return of spring; it is in flower near New York from the 10th to the 15th of April. The blossoms, of a beautifully-deep red, unfold more than a fortnight before the leaves. They are sessile, aggregate, and situated at the extremity of the branches. The fruit is suspended by long flexible peduncles, and is of the same hue with the flowers, though it varies in size and in the intensity of its coloring, according to the exposure and dampness of the soil. The leaves are smaller than those of the preceding species, but in some respects they resemble them. They are glaucous or whitish underneath, and are palmated or divided into three or four acuminate lobes, irregularly toothed. The extremities of this tree, which are formed by numerous twigs united at the base, have a remarkable appearance when garnished with flowers and seeds of a deep red, before vegetation has begun generally to revive.

Before the Red-flowering Maple exceeds twenty-five or thirty feet in height and seven or eight inches in diameter, its bark is perfectly smooth, and marked with white blotches, by which it is easily distinguishable. Afterward, the trunk, like that of the White Oak and Sweet Gum, becomes brown and chapped. In this tree, as in others which grow in wet places, the sap bears a large proportion to the heart, if indeed the name of heart can properly be given to the irregular star which occupies the centre of large trunks, with points from one to three inches in length projecting into the sap.

The wood of the Red-flowering Maple is applicable to interesting uses. It is harder than that of the White Maple, and of a finer and closer grain: hence, it is easily wrought in the lathe, and acquires by polishing a glossy and silken surface. It is sufficiently solid, and for many purposes it is preferred by workmen to other kinds of wood. It is principally employed for the lower part of Windsor chairs: the pieces are turned in the country, and so considerable is the demand, that boats laden with them arrive at New York and Philadelphia, where an extensive manufacture is carried on for the consumption of the neighboring towns, and for exportation to the Southern States and to the West India Islands. The whole frame of japanned chairs is of this wood, except the back, for which Hickory is chosen on account of its superior strength and elasticity. The frame, the nave, and the spokes of spinning-wheels are made of it: at Philadelphia it is exclusively employed for saddle trees, and in the country it is preferred for yokes, and also for shovels and wooden dishes, which are brought to market and purchased by the dealers in wooden ware.

It sometimes happens that, in very old trees, the grain, instead of following a perpendicular direction, is undulated, and this variety bears the name of Curled Maple. This singular arrangement, of which I am able to assign no cause, is never witnessed in young trees nor in the branches of such as exhibit it in the trunk; it is also less conspicuous at the centre than near the circumference. Trees offering this disposition are rare, and do not exist in the proportion of one to a hundred. The serpentine direction of the fibre, which renders them difficult to split and to work, produces, in the hands of a skilful mechanic, the most beautiful effects of light and shade. These effects are rendered more striking if, after smoothing the surface of the wood with a double-ironed plane, it is rubbed with a little sulphuric acid and afterward anointed with linseed-oil. On examining it attentively, the varying shades are found to be owing entirely to the

inflection of the rays of light, which is more sensibly perceived in viewing it in different directions by candle-light.

Before Mahogany became generally fashionable in the United States, the most beautiful furniture was of Red-flowering Maple, and bedsteads are still made of it, which, in richness of lustre, exceed the finest Mahogany. At Boston some cabinet-makers saw it into thin plates for inlaying Mahogany. But the most constant use of the Curled Maple is for the stocks of fowling-pieces and rifles, which to elegance and lightness unite a solidity resulting from the accidental direction of the fibre.

The cellular tissue of the Red-flowering Maple is of a dusky red. By boiling, it yields a purplish color, which, on the addition of sulphate of iron, becomes dark blue, approaching to black. It is used in the country, with a certain portion of alum in solution, for dyeing black.

The wood of the Red-flowering Maple does not burn well, and is so little esteemed for fuel that it is rarely brought into the cities.

The French Canadians make sugar from the sap of this Maple, which they call *Plaine*; but, as in the preceding species, the product of a given measure is only half as great as is obtained from the Sugar Maple.

It should be observed that the Red-flowering Maple never attains its full dimensions except in swamps where the bottom is composed of fertile soil. When the population of the country becomes denser, these tracts will be cleared and improved by some mode of culture more profitable than the growth of woods, and especially of this species, which is fit neither for the uses of the wheelwright nor for any other solid work; for it possesses little strength, is liable to injury from worms, and ferments and speedily decays when exposed to the alternations of dryness and moisture. Though at present it is extensively used, its importance in the arts is not such as to entitle it to preservation, and it will doubtless one day become rare. When the period arrives

y perceived

the United
ring Maple,
s of lustre,
inet-makers
t the most
of fowling-
te a solidity

of a dusky
n the addi-
roaching to
cion of alum

rn well, and
ght into the

sap of this
ling species,
great as is

Maple never
the bottom is
the country
mproved by
th of woods,
for the uses
r it possesses
erments and
dryness and
l, its import-
ervation, and
eriod arrives



H. J. Redouté del.

G. B. C. sculp.

Sugar Maple.
Acer saccharum.

Pl. 42



Asplenium nidus



that it is necessary in the United States, as in Europe, to renew the forests, or to preserve those which have escaped destruction, the American forester will find among the Oaks, the Walnuts, and the Ashes, many species more deserving of his care. The Sugar Maple also will be preferred, which grows on uplands, and possesses in a superior degree all the good properties of the other. From these considerations, the Red-flowering Maple appears to have no pretensions to a place in European forests.*

PLATE XLI.

A branch with leaves of the natural size. Fig. 1. Barren flowers. Fig. 2. Fertile flowers. Fig. 3. Seeds of the natural size.

SUGAR MAPLE.

ACER SACCHARINUM. *A. foliis quinque partito-palmatis, glabris, margine integris, subtus glaucis; floribus pedunculatis, pendentibus.*

THIS species, the most interesting of the American Maples, is called Rock Maple, Hard Maple, and Sugar Maple. The first of these names is most generally in use; but I have preserved the last, because it indicates one of the most valuable properties of the tree.

According to my father's researches into the topography of American vegetables, the Sugar Maple begins a little north of Lake St. John in Canada, near the 48th degree of latitude, which, in the rigor of its winter, corresponds to the 68th degree

* [See Emerson's "Trees and Shrubs of Massachusetts" for some additional particulars, and for remarks on the autumnal color of leaves, in which it is asserted that frost has very little influence on them.]

in Europe. It is nowhere more abundant than between the 46th and 43d degrees, which comprise Canada, New Brunswick, Nova Scotia, the States of Vermont and New Hampshire, and the district of Maine; in these regions it enters largely into the composition of the forests with which they are still covered. Farther south, it is common only in Genesee in the State of New York, and in the upper parts of Pennsylvania. It is estimated by Dr. Rush, that, in the northern parts of these two States, there are ten million of acres which produce these trees in the proportion of thirty to an acre. Indeed, I have noticed, in traversing these districts, large masses of woods formed of them almost exclusively. In Genesee, however, a great part of the Maples belong to a species which I shall describe, which has hitherto been confounded by botanists with the Sugar Maple.

In the lower parts of Virginia, of the Carolinas, and of Georgia, and likewise in the Mississippi Territory, this tree is unknown or very rare. It is rapidly disappearing from the forests about New York and Philadelphia, where it is no longer drained for sugar, but is felled for fuel and other purposes.

Between the parallels mentioned as bounding the tracts where this tree is most abundant, the forests do not resemble those of a more southern latitude: they are composed of two different descriptions of trees divided into two great classes, which alternately occupy the soil, and which exist in nearly equal proportions. The first class comprises the resinous trees, such as Pines and Spruces, and covers the low grounds and the bottoms of the valleys; these forests are called *Black-wood lands*. The second class consists of leafy trees, such as the Sugar Maple, the White and the Red Beech, the Birch and the Ashes, of which the Sugar Maple is most multiplied. They grow on level grounds or on gentle declivities, and form what are denominated *Hard-wood lands*. In proceeding from the 46th degree of latitude northward, the trees of the second class are observed to become more rare, and the resinous trees in the same proportion more

abundant: below the 43d degree, on the other hand, the resinous trees are found less common, and the others lose their predominance in the forests, as they become mingled with the numerous species of Oaks and Walnuts.

The Sugar Maple covers a greater extent of the American soil than any other species of this genus. It flourishes most in mountainous places, where the soil, though fertile, is cold and humid. Besides the parts which I have particularly mentioned, where the face of the country is generally of this nature, it is found along the whole chain of the Alleghanies to their termination in Georgia, and on the steep and shady banks of the rivers which rise in these mountains.

The Sugar Maple reaches the height of seventy or eighty feet, with a proportional diameter; but it does not commonly exceed fifty or sixty feet, with a diameter of twelve or eighteen inches. Well-grown, thriving trees are beautiful in their appearance, and easily distinguishable by the whiteness of their bark. The leaves are about five inches broad; but they vary in length according to the age and vigor of the tree. They are opposite, attached by long petioles, palmated and unequally divided into five lobes, entire at the edges, of a bright green above and glaucous or whitish underneath. In autumn, they turn reddish with the first frosts. Except in the color of the lower surface, they nearly resemble the Norway Maple. The flowers are small, yellowish, and suspended by slender, drooping peduncles. The seed is contained in two capsules united at the base and terminated by a membranous wing. It is ripe near New York in the beginning of October, though the capsules attain their full size six weeks earlier. Externally, they appear equally perfect, but I have constantly found one of them empty. The fruit is matured only once in two or three years.

The wood, when cut, is white; but, after being wrought and exposed for some time to the light, it takes a rosy tinge. Its grain is fine and close, and when polished it has a silken lustre.

It is very strong and sufficiently heavy, but wants the property of durability, for which the Chestnut and the Oak are so highly esteemed. When exposed to moisture it soon decays, and for this reason it is neglected in civil and naval architecture. In Vermont, New Hampshire, the district of Maine, and farther north, where the Oak is not plentiful, this timber is substituted for it in preference to the Beech, the Birch, and the Elm. When perfectly seasoned, (which requires two or three years,) it is used by wheelwrights for axle-trees and spokes, and for lining the runners of common sleds. It is also employed, as well as the Red-flowering Maple, in the manufacture of Windsor chairs. In the country, where the houses are wholly of wood, Sugar Maple timber is admitted into the frame; and in the district of Maine it is preferred to the Beech for the keels of vessels, as it furnishes longer pieces: with the Beech and the Yellow Pine, it forms also the lower frame, which is always in the water.

This wood exhibits two accidental forms in the arrangement of the fibre, of which cabinet-makers take advantage for obtaining beautiful articles of furniture. The first consists in undulations like those of the Curled Maple; the second, which takes place in old trees which are still sound, and which appears to arise from an inflexion of the fibre from the circumference toward the centre, produces spots of half a line in diameter, sometimes contiguous, and sometimes several lines apart. The more numerous the spots, the more beautiful and the more esteemed is the wood: this variety is called Bird's-eye Maple. Like the Curled Maple, it is used for inlaying Mahogany. Bedsteads are made of it, and portable writing-desks, which are elegant and highly prized. To obtain the finest effect, the log should be sawn in a direction as nearly as possible parallel to the concentric circles.

When cut at the proper season, the Sugar Maple forms excellent fuel. It is exported from the district of Maine for the consumption of Boston, and is equally esteemed with the

Hickory. The opinion entertained of it in this respect, in the north of America, accords with the interesting experiments of Mr. Hartig on the comparative heat afforded by different species of European wood, from which it results that the Sycamore (*Acer pseudo-platanus*) is superior to every other.

The ashes of the Sugar Maple are rich in the alkaline principle; and it may be confidently asserted that they furnish four-fifths of the potash exported to Europe from Boston and New York.

In the forges of Vermont and the district of Maine, the charcoal of this wood is preferred to any other, and it is said to be one-fifth heavier than the coal made from the same species in the Middle and Southern States,—a fact which sufficiently evinces that this Maple acquires its characteristic properties in perfection only in northern climates.

The wood of the Sugar Maple is easily distinguished from that of the Red-flowering Maple, which it resembles in appearance, by its weight and hardness. There is, besides, a very simple and certain test: a few drops of sulphate of iron being poured on samples of the different species, the Sugar Maple turns greenish, and the White Maple and Red-flowering Maple change to a deep blue.

The extraction of Sugar from the Maple is a valuable resource in a country where all classes of society make daily use of tea and coffee.

The process by which it is obtained is very simple, and is everywhere nearly the same. Though not essentially defective, it might be improved and made more profitable by adopting hints which have been thrown out in American publications.

The work is commonly taken in hand in the month of February, or in the beginning of March, while the cold continues

intense and the ground is still covered with snow. The sap begins to be in motion at this season, two months before the general revival of vegetation. In a central situation, lying convenient to the trees from which the sap is drawn, a shed is constructed, called a sugar-camp, which is destined to shelter the boilers, and the persons who tend them, from the weather. An auger three-quarters of an inch in diameter, small troughs to receive the sap, tubes of Elder or Sumac, eight or ten inches long, corresponding in size to the auger and laid open for a part of their length, buckets for emptying the troughs and conveying the sap to the camp, boilers of fifteen or eighteen gallons' capacity, moulds to receive the syrup when reduced to a proper consistency for being formed into cakes, and, lastly, axes to cut and split the fuel, are the principal utensils employed in the operation.

The trees are perforated in an obliquely-ascending direction, eighteen or twenty inches from the ground, with two holes four or five inches apart. Care should be taken that the augers do not enter more than half an inch within the wood, as experience has shown the most abundant flow of sap to take place at this depth. It is also recommended to insert the tubes on the south side of the tree; but this useful hint is not always attended to.

The troughs, which contain two or three gallons, are made, in the Northern States, of White Pine, of White or Black Oak, or of Maple; on the Ohio, the Mulberry, which is very abundant, is preferred. The Chestnut, the Black Walnut, and the Butternut should be rejected, as they impart to the liquid the coloring-matter and bitter principle with which they are impregnated.

A trough is placed on the ground at the foot of each tree, and the sap is every day collected and temporarily poured into casks, from which it is drawn out to fill the boilers. The evaporation is kept up by a brisk fire, and the scum is carefully taken off during this part of the process. Fresh sap is added

from time to time, and the heat is maintained till the liquid is reduced to a syrup, after which it is left to cool, and then strained through a blanket or other woollen stuff, to separate the remaining impurities.

Some persons recommend leaving the syrup twelve hours before boiling it for the last time; others proceed with it immediately. In either case, the boilers are only half filled, and, by an active, steady heat, the liquor is rapidly reduced to the proper consistency for being poured into the moulds. The evaporation is known to have proceeded far enough when, upon rubbing a drop of the syrup between the fingers, it is perceived to be granular. If it is in danger of boiling over, a bit of lard or of butter is thrown into it, which instantly calms the ebullition. The molasses being drained off from the moulds, the sugar is no longer deliquescent, like the raw sugar of the West Indies.

Maple Sugar manufactured in this way is light-colored in proportion to the care with which it is made and the judgment with which the evaporation is conducted. It is superior to the brown sugar of the Colonies,—at least, to such as is generally used in the United States; its taste is as pleasant, and it is as good for culinary purposes. When refined, it equals in beauty the finest sugar consumed in Europe. It is made use of, however, only in the districts where it is made, and there only in the country: from prejudice or taste, imported sugar is used in all the small towns and in the inns.

The sap continues to flow for six weeks, after which it becomes less abundant, less rich in saccharine matter, and sometimes even incapable of crystallization. In this case, it is consumed in the state of molasses, which is superior to that of the Islands. After three or four days' exposure to the sun, Maple sap is converted into vinegar by the acetous fermentation.

In a periodical work published at Philadelphia several years

since, the following receipt is given for making Sugar Maple beer:—Upon four gallons of boiling water pour one quart of Maple molasses; add a little yeast or leaven to excite the fermentation, and a spoonful of the essence of spruce; a very pleasant and salutary drink is thus obtained.

The process which I have described for extracting the sugar is the most common one, and it is the same from whatever species of Maple the sugar is made.

The amount of sugar manufactured in a year varies from different causes. A cold and dry winter renders the trees more productive than a changeable and humid season. It is observed that, when a frosty night is followed by a bright and brilliant day, the sap flows abundantly; and two or three gallons are sometimes yielded by a single tree in twenty-four hours. Three persons are found sufficient to tend 250 trees, which give 1000 pounds of sugar, or four pounds from each tree. But this product is not uniform; for many farmers on the Ohio have assured me that they did not commonly obtain more than two pounds from a tree.

Trees which grow in low and moist places afford a greater quantity of sap than those which occupy rising grounds, but it is less rich in the saccharine principle. That of insulated trees, left standing in the middle of fields or by the side of fences, is the best. It is also remarked that in districts which have been cleared of other trees, and even of the less vigorous Sugar Maple, the product of the remainder is, proportionally, most considerable.

While I resided in Pittsburg, the following curious particulars appeared in the Greensburg Gazette:—“Having introduced,” says the writer, “twenty tubes into a Sugar Maple, I drew from it the same day twenty-three gallons and three quarts of sap, which gave seven and a quarter pounds of sugar: thirty-three pounds have been made this season from the same tree; which supposes one hundred gallons of sap.” It appears here that

only a little more than three gallons was required for a pound, though four are commonly allowed.

In the foregoing experiments, five quarts were drawn in one day from each tube, which is about equal to the quantity discharged when two pipes are employed. Might it not hence be concluded that the sap escapes only from the orifices of the vessels which have been divided by the auger, without being diverted to this issue from the neighboring parts? I am the more inclined to this opinion, as, in rambling one day in the profound solitude of the forests on the banks of the Ohio, the idea suggested itself to me of cutting into a Maple which had been bored the preceding year. I found, amid the white mass of its wood, a green column, equal in width and in depth to the hole beneath. The organization appeared not to be affected; but this is not sufficient to warrant the conclusion that these vessels would be in condition to give passage to the sap the succeeding year. It may be objected that trees have been drained for thirty years, without diminution of their produce. But a tree of two or three feet in diameter presents an extensive surface, and the tubes are every year shifted; besides, the successive layers of thirty or forty years would restore it nearly to the state of one that never had been perforated.

In the United States, Maple sugar is made in the greatest quantities in the upper part of New Hampshire, in Vermont, in the State of New York, particularly in Genesee, and in the counties of Pennsylvania which lie on the eastern and western branches of the Susquehanna; west of the mountains, in the country bordering on the rivers Alleghany, Monongahela, and Ohio. The farmers, after reserving a sufficient store for their own consumption, sell the residue to the shopkeepers in the small towns of the neighborhood at eight cents a pound, by whom it is retailed at eleven cents. A great deal of sugar is also made in Upper Canada, on the Wabash, and near Michilimackinac. The Indians dispose of it to the commissioners of

the Northwestern Company established at Montreal, for the use of the numerous agents who go out in their employ, in quest of furs, beyond Lake Superior. In Nova Scotia and the district of Maine, and on the highest mountains of Virginia and the Carolinas, where these trees are sufficiently common, the manufacture is less considerable, and probably six-sevenths of the inhabitants consume imported sugar.

It has been stated, and doubtless correctly, that the northern parts of New York and Pennsylvania contain Maples enough to supply the whole consumption of the United States. But the annual produce by no means answers to this patriotic calculation. The trees grow upon excellent lands, which, by the influx of emigrants from the older settlements, and by the surprising increase of the population already established, are rapidly clearing; so that in less, perhaps, than half a century, the Maples will be confined to exposures too steep for cultivation, and will afford no resource, except to the proprietor on whose domain they grow. At this period, also, the wood will probably produce a greater and more ready profit than the sugar. It has been proposed to plant Sugar Maples in orchards or about the fields; but would it not be more certainly advantageous to multiply the Apple-tree, which grows in soils too dry to sustain the vegetation of the Maple? All that has been said on this subject must be considered as speculative merely, since, in the Eastern States, where information is generally diffused, no enterprises of this nature have been undertaken by which the importation of sugar might be diminished.


Wild and domestic animals are inordinately fond of Maple juice, and break into enclosures to sate themselves with it.

The details into which I have entered concerning the Sugar Maple furnish the means of estimating its importance with reference both to its sap and to its wood. I have indicated the regions where it grows and the soil in which it thrives; and I feel authorized in seriously recommending it for propagation in

ontreal, for the
employ, in quest
ia and the dis-
of Virginia and
ly common, the
six-sevenths of

hat the northern
Maples enough to
States. But the
patriotic calcula-
ch, by the influx
by the surprising
are rapidly clear-
tury, the Maples
ivation, and will
on whose domain
will probably pro-
e sugar. It has
ards or about the
advantageous to
too dry to sustain
been said on this
erely, since, in the
ly diffused, no en-
ken by which the

ely fond of Maple
selves with it.
cerning the Sugar
importance with
have indicated the
h it thrives; and I
for propagation in



H. J. Rehnke junr

Black Sugar Maple
Acer nigrum.

July 1894

Pl. 43.



July 1887



the north of Europe. Its sap and its wood are superior to those of the Norway Maple and of the Sycamore, and, in the same countries where these two species abound in the forests, its success would be most complete, and its cultivation more profitable.

PLATE XLII.

A branch with leaves and seeds of the natural size. Fig. 1. A small twig with flowers.

[For later and distinct remarks on the product of this tree, see Emerson's "Trees and Shrubs of Massachusetts," p. 489. Count Wingersky is said to have planted a great many of these trees on his estates in Moravia, and to have made very good sugar from their sap when they had attained the age of twenty-five years. But, in consequence of drawing sap every year, the trees became sickly, and soon died.]

 BLACK SUGAR-TREE.

ACER NIGRUM. *A. foliis quinque-partito-palmatis, sinibus apertis, margine integra, subtilis pubescentibus, atroviridibus; floribus corymbosis; capsulis turgidè subglobosis.*

In the Western States, and in the parts of Pennsylvania and Virginia which lie between the mountains and the Ohio, this species of Maple is designated by the name of Sugar-tree, and frequently by the more characteristic denomination of Black Sugar-tree: probably on account of the dark color of its leaves in comparison with those of the true Sugar Maple, which sometimes grows with it. In the extensive country of Genesee, both

species are indiscriminately called Rock Maple and Sugar Maple. This confusion seems to have arisen from the country's being settled principally by emigrants from the Eastern States, who, finding the Black Sugar-tree applicable to the same uses with the other, and equally productive of Sugar, have given it the same name. The two species have also been confounded by botanists, in describing the vegetable productions of America.

Toward the north, I first observed the Black Sugar-tree near Windsor in Vermont, on the Connecticut River. But, from its inferior size and its scarceness, it may be inferred that it belongs to a more southern climate. Accordingly, a few degrees lower, it forms a large part of the forests of Genesec, and covers the immense valleys through which flow the great rivers of the West. In these bottoms it is one of the most common and one of the loftiest trees.

The leaves are four or five inches long, and exhibit, in every respect, nearly the same conformation as those of the true Sugar Maple. They differ from them, principally, in being of a deeper green and a thicker texture, and in having more open sinuses: they are also slightly downy, which is most sensibly perceptible on the main rib.

The flowers, like those of the Sugar Maple, are suspended by long, flexible peduncles: the seeds, also, are similar, and are ripe about the same time; that is, about the 1st of October.

The wood is much like that of the other species, but it is coarser-grained and less brilliant when polished. It is little used, because, wherever it abounds, other trees are found, such as the Oak, the Walnut, the Cherry-tree, and the Mulberry, which are more esteemed for building and for cabinet-making. It is, however, preferred for the frames of Windsor chairs, and is considered, after the Hickories, as the best fuel. Its most important use is for making sugar, of which it annually yields a vast amount in the neighborhood of Pittsburg.

When the Black Sugar-tree stands alone, it naturally assumes

and Sugar
the country's
Eastern States,
the same uses
have given it
confounded by
of America.

ugar-tree near
But, from its
that it belongs
degrees lower,
and covers the
rivers of the
common and one

hibit, in every
the true Sugar
ing of a deeper
open sinuses:
ly perceptible

suspended by
imilar, and are
of October.

pecies, but it is
d. It is little
are found, such
the Mulberry,
abinet-making.
sor chairs, and
fuel. Its most
annually yields

naturally assumes



Sycamore,
Acer pseudoplatanus.

Norway Maple,
Acer platanoides.

T. Brown del.

T. Brown sculp.

Pl. 44



a regular and agreeable shape. Its foliage, of a darker tint and more tufted than that of the other Maples, renders it proper for forming avenues and for adorning parks and gardens; in short, for every situation where thick shade is desired, as a shelter from the sun.

PLATE XLIII.

A branch with a leaf and seeds of the natural size.

NORWAY MAPLE.

ACER PLATANOIDES. *A. foliis quinque-lobis, acuminatis, utrinque glabris, lobis dentatis; corymbis erectis, pedunculis glabris.*

THIS species of Maple is found in the same parts of Europe with the Sycamore, but it is most multiplied in Sweden and Norway, whence it has received the name of Norway Maple.

Like the Sycamore, it attains a lofty height and a diameter of several feet, and ranks among the largest trees of the north of Europe. Its leaves are broad, of a fine texture, and of a light green color; in shape they resemble those of the Black Sugar-tree and the Sugar Maple. They are not whitish underneath like those of the Sugar Maple, and when the petiole is broken a milky fluid distils from it, which does not take place in the American species.

The flowers of the Norway Maple are small, yellowish, and suspended by pretty long peduncles. The seeds grow in two capsules, which are united at the base, compressed, and garnished with large, divergent, membranous wings. They are ripe in the month of September.

In the winter, when the Norway Maple and the Sycamore

are stripped of their leaves, they may still be distinguished by their buds. On the Sycamore, the last year's shoots are larger than on the Norway Maple, and the buds are of a yellowish color, while those of the other species are of a reddish complexion, and are united in groups of three. On the two species of American Sugar Maple, the shoots are still more tapering and slender, and the buds are nearly black.

The wood of the Norway Maple is very white and very fine-grained: it is easily wrought, and is employed for nearly the same purposes with that of the Sycamore. Among cabinet-makers in Germany, such trees are in request as present agreeable accidental variations in the direction of the fibre, similar to the Curled Maple and the Bird's-eye Maple.

The rapid and beautiful vegetation of the Norway Maple in soils inferior to such as are required by the Sycamore causes it to be extensively planted in Europe for the embellishment of gardens; for which purpose trees are preferred that develop their foliage early and shed it late, and that afford through the intemperate season a refreshing shade; all which advantages are united in the Norway Maple.

PLATE XLIV.

Fig. 1. A leaf of half the natural size. A seed of the natural size.

[This is unquestionably one of the best ornamental trees; its beautifully-formed, round head, and the great density of its shade, should give it a preference over most of the Maples for the landscape-gardener. Scarcely too much can be said in its praise.]

SYCAMORE TREE.

ACER PSEUDO-PLATANUS. *A. foliis quinque lobis, inæqualiter dentatis subtilis glaucescentibus; floribus subspicatis, pendulis.*

THIS beautiful tree is diffused over all the centre of Europe, and abounds especially in Bohemia, Hungary, and Poland. It thrives most luxuriantly in moist and fertile soils, and when expanded to its full dimensions it is sixty or seventy feet in height and two or three feet in diameter. Its head is spacious, and its foliage thick. On old trees, the bark of the trunk is deeply furrowed; on such as are less than six inches in diameter, it is perfectly smooth. The leaves of the Sycamore are opposite, with long petioles, large, and distinctly divided into five unequal lobes; they are of a dark green above and whitish underneath. In the heat of midsummer they are covered with a very sweet viscid substance, which is gathered with avidity by bees. The flowers appear toward the end of April: they are small, greenish, and grouped into pendulous clusters from three to four inches in length. The seed is in capsules about an inch in length, united at the base and terminated by a membranous wing.

When the Sycamore is fully grown, its wood is fine-grained and susceptible of a brilliant polish. In those parts of Europe where it is most common, it is in demand with turners for making wooden ware. It is used for making violins, and, when its grain is undulated, for ornamenting forte-pianos. By the interesting experiments of Mr. Hartig, Grand Master of the forests of Prussia, on the comparative value of different species of wood as combustibles, the Sycamore was found to afford more heat than any other wood of the north of Europe.

Sugar has been made from the Sycamore in Bohemia and Hungary. Though the attempt has completely succeeded, it

appears certain that the sugar is yielded in a smaller proportion than by the Sugar Maple.

In France and England, the Sycamore is a rare tree in the forests, but it is multiplied in pleasure-grounds, on account of its rapid growth, the early development of its foliage in the spring, and the fine shade which it affords through the summer.

It has been observed in England that the foliage of this tree is less injured than that of others by the saline vapors wafted from the sea; hence it is chosen for situations exposed to these winds. The justness of the observation I have never had an opportunity of examining.

The Sycamore appears to me to possess no one superior property which entitles it to preference in the United States over the Sugar Maple and the Black Sugar-tree.

PLATE XLIV.

Fig. 2. A leaf of half the natural size. A seed of the natural size.

[Its growth is very rapid, particularly when it is in a deep, free, rich soil, and in a mild climate. It arrives at full growth in fifty or sixty years. In marshy soil, or dry sand, the tree never attains a great size.

“There is a very interesting tree of this species standing at the entrance of the village of Trons, in the Grisons, the cradle of liberty in the Rhoetian Alps. Under the once spreading branches of this now hollow and cloven trunk, the Gray League was solemnly ratified in 1424. Upon the supposition that it was only a century old when the meeting, to which its celebrity is owing, took place, and a younger tree would hardly have been selected for the purpose, it has now attained the age of five hundred and twenty years, and may be much older.” *Dr. Gray, in N. American Rev., July, 1844.*]

r proportion


tree in the
account of
age in the
he summer.
of this tree
spors wafted
sed to these
ever had an

superior pro-
States over

natural size.

is in a deep,
t full growth
and, the tree

s standing at
ns, the cradle
nce spreading
Gray League
sition that it
h its celebrity
dly have been
e age of five
r." *Dr. Gray,*




H. J. Rehnke del.

J. G. Smith sculp.

Moose Wood.
Acer striatum.

Pl. 45



July 1845

MOOSE WOOD.

ACEP STRIATUM. *A. foliis infernè rotundatis, supernè acuminato-tricuspidibus, argute serratis; racemis simplicibus, pendentibus.*

A. demissylvanicum. LINN.

IN the Province of Nova Scotia and New Brunswick, in the district of Maine, and in the States of Vermont and New Hampshire, this Maple is known only by the name of Moose Wood: in New Jersey and Pennsylvania it is called Striped Maple. This last denomination, which is preferable as being descriptive, I have thought proper to reject, because it is in use only in a part of the United States where the tree is rare, and is wholly unknown in those parts in which it abounds. The name of Moose Wood was given it by the first settlers from observing that the moose—an animal now rare in this region—subsisted during the latter part of winter and the beginning of spring upon its young twigs.

This tree makes its first appearance near Lake St. John in the 47th degree of latitude, that is to say, a little farther north than the preceding species. In Nova Scotia and the district of Maine, where I have most particularly observed it, it fills the forests. Toward the Hudson it becomes more rare, and beyond this boundary is confined to the mountainous tracts of the Alleghanies, on which it is found in cold and shaded exposures, along the whole range to its termination in Georgia.

In the district of Maine I have always found the Moose Wood most vigorous in mixed forests, or what are called *Mixture lands*; where the woods are composed of the Sugar Maple, the Beech, the White Birch, the Yellow Birch, and the Hemlock Spruce. In these forests it constitutes a great part of the undergrowth;

for its ordinary height is less than ten feet, though I have found individual trees of more than twice this stature.

The trunk and branches of the Moose Wood are clad in a smooth green bark, longitudinally marked with black stripes, by which it is easily distinguishable at all seasons of the year.

This is one of the earliest trees of North America whose vegetation announces the approach of the genial season. Its buds and leaves, when beginning to unfold, are rose-colored, and have a pleasing effect; but this hue soon changes to green. On luxuriant trees, the leaves are of a pretty thick texture, and finely serrate. They are four or five inches broad, rounded at the base, and divided into three deep acute lobes. The flowers are of a greenish cast, and grouped on long, pendulous peduncles. The fruit, which in the main resembles that of the other Maples, is remarkable for a small cavity on one side of the capsules: it is produced in abundance, and is ripe about the end of September.

The inferior size of the Moose Wood forbids its use in any kind of carpentry; but, as it is white and fine-grained, the cabinet-makers of Halifax employ it instead of the Holly, which does not grow in so northern a climate, for forming the white lines with which they inlay Mahogany. Its principal advantage consists in furnishing the inhabitants at the close of winter, when their forage is exhausted, a resource for sustaining their cattle till the advancing season has renewed the herbage. As soon as the buds begin to swell, the famished horses and neat cattle are turned loose into the woods, to browse on the young shoots, which they crop with avidity. Poor as this resource may appear, it is not wholly inadequate, since the twigs are tender and full of saccharine juice. A similar practice prevails also in the new settlements of the West.

This species of Maple has been long cultivated in Europe, in parks and extensive gardens. It is in request as one of the earliest trees to feel the approach of spring, but more particu-

larly on account of the pleasing effect of the white veins which variegates its trunk. In the primitive forests, where it grows beneath a canopy of impervious shade, these veins are black: the change of color seems owing to its being planted in drier soils more open to the sun. Most of the trees of this species which now grow in Europe have been grafted on the lofty Sycamore, whose vigor is felt by the Moose Wood, and expands it to four times its natural dimensions.

PLATE XLV.

A branch with fruit of the natural size. Fig. 1. Bark of a tree in the forests of North America. Fig. 2. Bark of a tree cultivated on dry and open ground.

[From the great beauty of its bark, this tree deserves a place in every collection. The delicate rose-color of the buds and leaves on opening, and the elegance of the ample foliage afterward, the graceful, pendulous racemes of flowers succeeded by large, showy keys, not unlike a cluster of insects, will also recommend it.]

Europe, in
one of the
more particu-

BOX ELDER,
OR
ASH-LEAVED MAPLE.

ACER NEGUNDO. *A. foliis pinnatis ternatisve, inæqualiter serratis; floribus dioicis.*

In the country west of the Alleghanies, where this tree is common, it is called Box Elder; east of the mountains it is more rare, and, having been less attentively observed, it has received no specific name. Some persons, however, distinguish it by that of Ash-leaved Maple, which is a perfectly appropriate denomination. I have chosen the other, though absolutely insignificant of any characteristic property of the tree, because it is sanctioned by general use. The French of Illinois call it *Erable à Giguères*.

The leaves of the Box Elder are opposite, and are from six to fifteen inches long, according to the vigor of the tree and the moisture of the soil in which it grows. Each leaf is composed of two pair of leaflets with an odd one. The leaves are petioled and oval-acuminate, and sharply toothed; toward fall, the common petiole is of a deep red. The male and female flowers are borne on different trees, and are supported by slender, pendulous peduncles, six or seven inches in length.

Of all the Maples of the United States, this species ventures least into northern latitudes, for in the Atlantic States it is first seen on the banks of the Delaware, in the neighborhood of Philadelphia, and even there it is rare. In the maritime parts of the Southern States, also, it is far from being a common tree; which is less attributable to the heat of the summer than the marshy nature of the soil on the borders of the rivers. West of the mountains, on the contrary, it is extremely multiplied,

; *floribus*

s tree is
ins it is
l, it has
stinguish
ropriate
ely insig-
ause it is
it *Erable*

from six
e and the
composed
are petio-
d fall, the
le flowers
y slender,

s ventures
s it is first
oorhood of
time parts
nmon tree;
r than the
ers. West
multiplied,



Wm. A. D.

Gabriel S. S.

Box Elder.
Acer negundo.

and, instead of being confined, as in the upper parts of Virginia and of the Carolinas, to the river-sides, it grows in the woods, with the Locust, Wild Cherry, and Coffee-tree. But in the bottoms which skirt the rivers, where the soil is deep, fertile, constantly moist, and often inundated, this tree is most abundant and most fully expanded. Even here it can be considered only as a tree of secondary size: the largest Box Elders that I have seen were not more than fifty feet in height and twenty inches in diameter, and trees of these dimensions are found only in Tennessee and in the back parts of Georgia, which lie far to the south. In Kentucky, they are of only half this height. Though growing in thick forests, the Box Elder expands into a head like that of the Apple-tree. I have remarked this form particularly on the banks of the Ohio, where I have also observed that the trunk bulges into knots at unequal distances, and is often decayed at the heart. A fine row of Box Elders in the botanical garden of Paris, along the *Rue de Buffon*, affords a sufficiently just idea of their appearance in the forests on the rivers Monongahela and Ohio. It may be concluded, from what has been said, that, to obtain its full proportions, this tree requires a climate three or four degrees milder than that of Philadelphia, Pittsburg, and Paris.

The Box Elder branches at a small height. The bark of its trunk is brown, and I have remarked a disagreeable odor in the cellular tissue. The proportion of the sap to the heart is large, except in very old trees: in these the heart is variegated with rose-colored and bluish veins. Some cabinet-makers in the Western country employ it to ornament furniture made of Mahogany or Wild Cherry-tree. The wood is of a fine and close grain, and is said to split with difficulty; but, as it soon decays when exposed to the air, it is little used. It has been erroneously asserted that sugar is made from the sap of this species.

More than fifty years since, the Box Elder was introduced

into France by Admiral La Gallissonière. Subsequently it has spread into Germany and England, where it is in great request for adorning pleasure-grounds, on account of the rapidity of its growth and the beauty of its foliage, whose bright green forms an agreeable contrast with the surrounding trees. Its young branches, of a lively green, contribute to the favor in which it is held, and serve to distinguish it in the winter, when its leaves are fallen.

The utility of its wood, I believe, has of late been exaggerated; both Europe and America possess many trees superior in strength as well as in size. It appears certain, however, that, growing in copses, and cut every three or four years, it would afford a profitable product in its sprouts, which are very numerous, and which, during the first years, shoot with astonishing rapidity. The success of this experiment will be more certain if it is made on grounds constantly moist and cool: though the Box Elder may seem, for a few years, to prosper in dry and meagre soils, it sooner or later pines and perishes. Of this I became convinced in conversing with several proprietors in the environs of Paris, who, after some recent publications on this tree, had made an unsuccessful attempt to derive profit from their poor lands by planting them with the Box Elder.

PLATE XLVI.

A branch with leaves and seed of the natural size.

E.

it has
request
y of its
n forms
s young
which it
ts leaves

exagge-
perior in
ver, that,
it would
ry nume-
tonishing
e certain
ough the
n dry and
Of this I
ors in the
s on this
rofit from
c.



A. montanum

Mountain Maple,
Acer montanum

Robert Gray

17 +



Salvia

MOUNTAIN MAPLE.

ACER MONTANUM. *A. foliis tri-subquinculobis, acuminatis, dentatis, rugosis; racemis spiciformibus, suberectis, petalis lucaribus.*

THIS species is more abundant in Canada, Nova Scotia, and along the whole range of the Alleghany Mountains, than in any other part of North America. It is called Mountain Maple and Low Maple. Though the last of these names indicates the stature of the tree, I have retained the first, which is more generally in use, and which is likewise appropriate, as this Maple grows of preference on the declivities of mountains exposed to the north, and in cool, moist, and shady situations, on the abrupt and rocky banks of torrents and rivers. On the Mohawk, for instance, near the Little Falls, it abounds among the enormous rocks which lie scattered along its sides.

The Mountain Maple is six or eight feet in height, and it blooms even at a smaller elevation. It most frequently grows in the form of a shrub, with a single and straight stock. The leaves are large, opposite, and divided into three acute and indented lobes: they are slightly hairy at their unfolding, and, when fully grown, they are uneven and of a dark green upon the upper surface. The blossoms are small, of a greenish color, and produced in semi-erect spikes from two to four inches in length. The seeds, which are smaller than those of any other American Maple, are fixed upon slender, pendulous footstalks: they are reddish at maturity, and each of them is surmounted by a membranous wing, and has a small cavity upon one side.

The Mountain Maple is too small to be profitably cultivated for its wood; and, as its flowers, its roots, and its bark are destitute of any very sensible odor, it promises no resources to medicine. It is found in the gardens of the curious, rather to

complete the series of species than for any remarkable property of its foliage or of its flowers.

This species is commonly grafted upon the Sycamore, and, like the Moose Wood, it is thus augmented to twice its natural dimensions. This surprising development evinces how great are the advantages which may be derived from this process and from continued cultivation in improving inferior vegetables.

PLATE XLVII.

A branch with leaves and flowers of their natural size. Fig. 1. A bunch of seeds of the natural size.

[The Mountain Maple sometimes attains the height of twenty-four feet; the leaves assume, toward autumn, various rich shades of red, with the seeds yellowish, and in this state it has considerable beauty.]

DOGWOOD.

CORNUS FLORIDA. *C. foliis ovalibus, acuminatis subtis albicantibus; floribus sessiliter capitatis; involuero maximo, foliolis apice deformi quasi obcordatis; fructibus oratis, rubris.*

Tetrandria monogynia. LINN. Caprifolia. JUSS.

AMONG the eight species of Dogwood which have been observed in North America, this alone is entitled by its size to be classed with the forest-trees. It is the most interesting, too, for the value of its wood, the properties of its bark, and the beauty of its flowers. In the United States at large, it is known by the name of Dogwood, and in Connecticut it is also called Boxwood.

ble property

amore, and,
its natural
how great
process and
vegetables.

1. *A bunch of*

ht of twenty-
as rich shades
e it has con-

*bicantibis; flori-
e deformi quasi*

trifolia. Juss.

have been ob-
y its size to be
interesting, too,
bark, and the
ge, it is known
is also called



Dogwood.
Cornus florida

The Dogwood is first seen in Massachusetts, between the 42d and 43d degrees of latitude; and in proceeding southward, it is met with uninterruptedly throughout the Eastern and Western States, and the two Floridas, to the banks of the Mississippi. Over this vast extent of country it is one of the most common trees, and it abounds particularly in New Jersey, Pennsylvania, Maryland, and Virginia, wherever the soil is moist, gravelly, and somewhat uneven; farther south, in the Carolinas, Georgia, and the Floridas, it is found only on the borders of swamps, and never in the pine-barrens, where the soil is too dry and sandy to sustain its vegetation. In the most fertile districts of Kentucky and West Tennessee, it does not appear in the forests except where the soil is gravelly and of a middling quality.

The Dogwood sometimes reaches thirty or thirty-five feet in height and nine or ten inches in diameter; but it does not generally exceed the height of eighteen or twenty feet and the diameter of four or five inches. The trunk is strong, and is covered with a blackish bark, chapped into small portions, which are often in the shape of squares more or less exact. The branches are proportionally less numerous than on other trees, and are regularly disposed nearly in the form of crosses. The young twigs are observed to incline upward in a semicircular direction.

The leaves are opposite, about three inches in length, oval, of a dark green above and whitish beneath; the upper surface is very distinctly sulcated. Toward the close of summer they are often marked with black spots, and at the approach of winter they change to a dull red.

In New York and New Jersey, the flowers are fully blown about the 10th or 15th of May, while the leaves are only beginning to unfold themselves. The flowers are small, yellowish, and collected in bunches, which are surrounded with a very large involucre composed of four white floral leaves, sometimes inclining to violet. This fine involucre constitutes all the beauty

of the flowers, which are very numerous, and which in their season robe the tree in white like a full-blown Apple-tree, and render it one of the fairest ornaments of the American forests.

The seeds, of a vivid glossy red and of an oval shape, are always united. They remain upon the trees till the first frosts, when, notwithstanding their bitterness, they are devoured by the Robin, *Turdus migratorius*, which about this period arrives from the northern regions.

The wood is hard, compact, heavy, and fine-grained, and is susceptible of a brilliant polish. The sap is perfectly white, and the heart is of a chocolate-color. This tree is not large enough for works which require pieces of considerable volume: it is used for the handles of light tools, such as mallets, small vices, etc. In the country some farmers select it for harrow-teeth, for the hames of horses' collars, and also for lining the runners of sledges; but, to whatever purpose it is applied, being liable to split, it should never be wrought till it is perfectly seasoned. The shoots, when three or four years old, are found proper for the light hoops of small, portable casks; but the consumption in this way is inconsiderable. In the Middle States, the cogs of mill-wheels are made of Dogwood, and its divergent branches are taken for the yokes which are put upon the necks of swine to prevent their breaking into cultivated enclosures. Such are the profitable uses of this tree: it affords also excellent fuel, but it is too small to be brought into the markets of the cities.

The *liber* or interior bark of the Dogwood is extremely bitter, and proves an excellent remedy in intermittent fevers. It has been known and successfully used by the country-people, as a specific in these maladies, for more than fifty years. Its medicinal properties were made the subject of a thesis defended in the College of Physicians at Philadelphia, in 1803, in which was presented an analysis of the bark of the Dogwood and of the Blue-berried Dogwood, compared with the Peruvian bark.

By the experiments made on this occasion, the Dogwood bark was shown to have a close analogy to the Peruvian bark, and to be capable, in many cases, of supplying its place with success. The author of this excellent piece cites a physician of Pennsylvania, who during twenty years had constantly employed it, and who estimated thirty-five grains of it to be equivalent to thirty grains of the Peruvian bark. The only inconvenience accompanying its use was that, if taken within a year after being stripped from the tree, it sometimes occasioned acute pains of the bowels; but this evil was remedied by adding to it five grains of Virginia Snake-root, (*Aristolochia serpentaria*.)

The same author gives a receipt for making an excellent ink, in which this bark is substituted for gall-nuts. Put half an ounce of Dogwood bark, two scruples of sulphate of iron, and two scruples of gum-arabic, into sixteen ounces of rain-water: during the infusion shake it repeatedly.

The Dogwood merits the attention of Europeans for the value of its wood, and especially for the brilliancy of its flowers, by which it is better adapted than almost any other North American tree to the embellishment of forests, parks, and extensive gardens.

PLATE XLVIII.

A branch with leaves and flowers of the natural size. Fig. 1. A branch with fruit of the natural size.

[From the bark of the smaller roots the Indians obtained a good scarlet color. The smaller branches, stripped of their bark and used as a brush, are said to render the teeth extremely white. EMERSON.]

GEORGIA BARK.

PINCKNEYA PUBENS. *P. foliis oppositis, ovalibus, utrinque acutis; subtomentosis.*

Pentandria monogynia. LINN. Rubiaceæ. JUSS.

Obs. Floribus majusculis, pallentibus et purpureo-lineatis, fasciculato-paniculatis. Capsulis subrotundis, modicè compressis: seminibus numerosis alatis.

THIS tree, still more interesting by the properties of its bark than by the elegance of its flowers and of its foliage, is indigenous to the most southern parts of the United States: probably it grows also in the two Floridas and in Lower Louisiana. My father found it for the first time, in 1791, on the banks of the St. Mary. He carried seeds and young plants to Charleston, and planted them in a garden which he possessed near that city. Though intrusted to an ungrateful soil, they succeeded so well that in 1807 I found several of them twenty-five feet high and seven or eight inches in diameter; which proves that the vegetation of this tree does not require a very warm climate nor a very substantial soil.

With a great affinity to the *Cinchona*, which yields the Peruvian bark, my father discerned in the Georgia Bark sufficient differences to distinguish it as a new genus. In testimony of his gratitude and respect, he consecrated it to Charles Cotesworth Pinckney, an enlightened patron of the arts and sciences, from whom my father and myself, during our residence in South Carolina, received multiplied proofs of benevolence and esteem.

The Georgia Bark is a low tree, dividing itself into numerous branches, and rarely exceeding the height of twenty-five feet and the diameter of five or six inches at the base. A cool and shady exposure appears the most favorable to its growth. Its leaves are opposite, four or five inches long, of a light green

s; sub-

Juss.

miculatis.
tis.

ts bark
indige-
probably
na. My
s of the
arleston,
hat city.
l so well
high and
the vege-
ate nor a

the Peru-
sufficient
timony of
Cotesworth
nces, from
in South
nd esteem.
o numerous
ty-five feet
A cool and
rowth. Its
light green

Pr

Or

th
no
it
fa
S
a
T
tl
so
ti
v

v
d
l
l
v
C

l
r
r



W. Bot. del.

W. Bot. sculp.

Georgia Bark
Puckneya pubens

color, and downy underneath, as are also the shoots to which they are attached. The flowers, which are white, with longitudinal rose-colored stripes, are pretty large, and are collected in beautiful panicles at the extremity of the branches. Each flower is accompanied by a floral leaf, bordered with rose-color near the upper edge. The capsules are round, compressed in the middle, and stored with a great number of small, winged seeds.

The wood of the Georgia Bark is soft, and unfit for use in the arts; but its inner bark is extremely bitter, and appears to partake of the febrifuge virtues of the *Cinchona*, for the inhabitants of the southern parts of Georgia employ it successfully in the intermittent fevers which, during the latter part of summer and the autumn, prevail in the Southern States. A handful of the bark is boiled in a quart of water till the liquid is reduced one-half, and the infusion is administered to the sick. From the properties of its bark, the *Pinckneya* has taken the name of *Georgia Bark*. It is to be wished that some intelligent physician would examine these properties with care, and indicate with accuracy the manner of employing this indigenous remedy, and the effects to be expected from it. The tree which produces it so nearly resembles the Peruvian vegetable, that some botanists have included them in the same genus.

PLATE XLIX.

*A branch with leaves and flowers of the natural size. Fig. 1. A seed-vessel.
Fig. 2. A seed.*

COFFEE TREE.

GYMNOCLADUS CANADENSIS. *G. jolii bipinnatis, amplissimis, deciduis; foliis ovalibus, acuminatis; floribus racemosis; leguminosis polyspermis.*

Dicecia decandria. LINN. Leguminosæ. JUSS.

UPPER CANADA beyond Montreal, and that part of Genesee which borders on Lake Ontario and Lake Erie, are the most northern countries which produce the Coffee-tree: but it is much less abundant in these climates than in the States of Kentucky and Tennessee, and in the tract which is bounded by the rivers Ohio and Illinois, between the 35th and 40th degrees of latitude. The large dimensions which it exhibits in these regions is attributable to the milder temperature of the seasons and to the extreme fertility of the soil.

The French of Canada call this tree *Chicot*; those of Illinois, *Gros Favier*; and the inhabitants of the Western States, Coffee-tree.

The presence of the Coffee-tree is an index of the richest lands; on which it habitually grows in company with the Black Walnut, the Red Elm, the Poplar, the Blue Ash, the Honey Locust, and the Hackberry. These trees it equals in height, but not in bulk; for a Coffee-tree fifty or sixty feet high does not generally exceed twelve or fifteen inches in diameter.

In summer, this tree, when fully grown, has a fine appearance: its straight trunk is often destitute of branches for forty feet, and supports a summit not very widely spread, but of a regular shape and of tufted foliage: such at least is its form in primitive forests, where it is confined by the trees which grow around it. In the winter, when its leaves are fallen, the paucity of its branches and the size of the terminal ones, which are very

imilis, deciduis;
uminosis poly-

minosæ. Juss.

of Genesee
are the most
e: but it is
the States of
bounded by
40th degrees
bits in these
of the seasons

se of Illinois,
ates, Coffee-

the richest
th the Black
, the Honey
ls in height,
et high does
meter.

fine appear-
ces for thirty
ad, but of a
is its form in
which grow
n, the paucity
hich are very



P. J. Boiss. del.

Bonard sculp.

Coffee Tree.
Cynocladus canadensis.



large in comparison with those of other trees, give it a peculiar appearance, somewhat resembling a dead tree. This is probably the reason of its being called *Chicot*—Stump-tree—by the French Canadians. To this peculiar character is added another of the epidermis, which is extremely rough, and which detaches itself in small, hard, transverse strips, rolled backward at the ends, and projecting sufficiently to render the tree distinguishable at first sight. I have also remarked that the live bark is very bitter, so that a morsel no bigger than a grain of maize, chewed for some time, produces a violent irritation of the throat.

The leaves are three feet long and twenty inches wide on young and thriving trees: on old ones they are not more than half as large. These leaves are doubly compound, with oval-acuminate leaflets from one to two inches long. The leaflets are of a dull green, and in the fall the petiole is of a violet color.

The Coffee-tree belongs to the class *Dioecia* of Linnæus, which includes all vegetables whose male and female flowers are borne by different plants; in which case those only that bear the female flowers produce fruit: to effect the fecundation, it is necessary that there should be male plants growing near them. The flowers and the fruit are large, bowed pods, of a reddish-brown color, and of a pulpy consistency within. They contain several large, gray seeds, which are extremely hard. The French of Upper Louisiana call them *Gourygués*.

The name of Coffee-tree was given to this vegetable by the early emigrants to Kentucky and Tennessee, who hoped to find in its seeds a substitute for coffee; but the smaller number of persons who made the experiment abandoned it as soon as it became easy to obtain from the seaports the coffee of the West Indies.

The wood of the Coffee-tree is very compact and of a rosy hue. The fineness and closeness of its grain fit it for cabinet-making, and its strength renders it proper for building. Like

the Locust, it has the valuable property of rapidly converting its sap into perfect wood, so that a trunk six inches in diameter has only six lines of sap, and may be employed almost entire. These qualities recommend it for propagation in the forests of the north and of the centre of Europe.

The Coffee-tree was sent to France more than fifty years since. It thrives in the environs of Paris, where there are trees that exceed forty feet in height; but it does not yield fruit, and is multiplied only by shoots obtained by digging trenches round the old trees. The divided roots produce shoots three or four feet long the first year. The young trees are sought, on account of their beautiful foliage, for the embellishment of parks and picturesque gardens.

PLATE I.

A branch with flowers of the natural size. Fig. 1. A pod of the natural size. Fig. 2. A seed of the natural size.

[The Coffee-tree thrives as far north as Massachusetts. It requires a rich, deep, free soil, and, when isolated, spreads over a large space, and is extremely beautiful. It is readily propagated from the seeds.]

END OF VOL. I.



verting
iameter
entire.
ests of

s since.
es that
and is
round
or four
account
cks and




e natural

etts. It
nds over
y propa-





**IMAGE EVALUATION
TEST TARGET (MT-3)**



**Photographic
Sciences
Corporation**

23 WEST MAIN STREET
WEBSTER, N.Y. 14580
(716) 872-4903

18
20
22
25

18
20
22
25

FICHE 5 NOT REQUIRED



