

**CIHM  
Microfiche  
Series  
(Monographs)**

**ICMH  
Collection de  
microfiches  
(monographies)**



**Canadian Institute for Historical Microreproductions / Institut canadien de microreproductions historiques**

**© 1997**

## Technical and Bibliographic Notes / Notes techniques et bibliographiques

The Institute has attempted to obtain the best original copy available for filming. Features of this copy which may be bibliographically unique, which may alter any of the images in the reproduction, or which may significantly change the usual method of filming are checked below.

- Coloured covers /  
Couverture de couleur
- Covers damaged /  
Couverture endommagée
- Covers restored and/or laminated /  
Couverture restaurée et/ou pelliculée
- Cover title missing / Le titre de couverture manque
- Coloured maps / Cartes géographiques en couleur
- Coloured ink (i.e. other than blue or black) /  
Encre de couleur (i.e. autre que bleue ou noire)
- Coloured plates and/or illustrations /  
Planches et/ou illustrations en couleur
- Bound with other material /  
Relié avec d'autres documents
- Only edition available /  
Seule édition disponible
- Tight binding may cause shadows or distortion along  
interior margin / La reliure serrée peut causer de  
l'ombre ou de la distorsion le long de la marge  
intérieure.
- Blank leaves added during restorations may appear  
within the text. Whenever possible, these have been  
omitted from filming / Il se peut que certaines pages  
blanches ajoutées lors d'une restauration  
apparaissent dans le texte, mais, lorsque cela était  
possible, ces pages n'ont pas été filmées.
- Additional comments /  
Commentaires supplémentaires:

L'Institut a microfilmé le meilleur exemplaire qu'il lui a été possible de se procurer. Les détails de cet exemplaire qui sont peut-être uniques du point de vue bibliographique, qui peuvent modifier une image reproduite, ou qui peuvent exiger une modification dans la méthode normale de filmage sont indiqués ci-dessous.

- Coloured pages / Pages de couleur
- Pages damaged / Pages endommagées
- Pages restored and/or laminated /  
Pages restaurées et/ou pelliculées
- Pages discoloured, stained or foxed /  
Pages décolorées, tachetées ou piquées
- Pages detached / Pages détachées
- Showthrough / Transparence
- Quality of print varies /  
Qualité inégale de l'impression
- Includes supplementary material /  
Comprend du matériel supplémentaire
- Pages wholly or partially obscured by errata slips,  
tissues, etc., have been refilmed to ensure the best  
possible image / Les pages totalement ou  
partiellement obscurcies par un feuillet d'errata, une  
pelure, etc., ont été filmées à nouveau de façon à  
obtenir la meilleure image possible.
- Opposing pages with varying colouration or  
discolourations are filmed twice to ensure the best  
possible image / Les pages s'opposant ayant des  
colorations variables ou des décolorations sont  
filmées deux fois afin d'obtenir la meilleure image  
possible.

This item is filmed at the reduction ratio checked below /  
Ce document est filmé au taux de réduction indiqué ci-dessous.

<b>10x</b>		<b>14x</b>		<b>18x</b>		<b>22x</b>		<b>26x</b>		<b>30x</b>
								✓		
	<b>12x</b>		<b>16x</b>		<b>20x</b>		<b>24x</b>		<b>28x</b>	<b>32x</b>

The copy filmed here has been reproduced thanks to the generosity of:


National Library of Canada

The images appearing here are the best quality possible considering the condition and legibility of the original copy and in keeping with the filming contract specifications.

Original copies in printed paper covers are filmed beginning with the front cover and ending on the last page with a printed or illustrated impression, or the back cover when appropriate. All other original copies are filmed beginning on the first page with a printed or illustrated impression, and ending on the last page with a printed or illustrated impression.

The last recorded frame on each microfiche shall contain the symbol  $\rightarrow$  (meaning "CONTINUED"), or the symbol  $\nabla$  (meaning "END"), whichever applies.

Maps, plates, charts, etc., may be filmed at different reduction ratios. Those too large to be entirely included in one exposure are filmed beginning in the upper left hand corner, left to right and top to bottom, as many frames as required. The following diagrams illustrate the method:



L'exemplaire filmé fut reproduit grâce à la générosité de:


Bibliothèque nationale du Canada

Les images suivantes ont été reproduites avec le plus grand soin, compte tenu de la condition et de la netteté de l'exemplaire filmé, et en conformité avec les conditions du contrat de filmage.

Les exemplaires originaux dont la couverture en papier est imprimée sont filmés en commençant par le premier plat et en terminant soit par la dernière page qui comporte une empreinte d'impression ou d'illustration, soit par le second plat, selon le cas. Tous les autres exemplaires originaux sont filmés en commençant par la première page qui comporte une empreinte d'impression ou d'illustration et en terminant par la dernière page qui comporte une telle empreinte.


Un des symboles suivants apparaîtra sur la dernière image de chaque microfiche, selon le cas: le symbole  $\rightarrow$  signifie "A SUIVRE", le symbole  $\nabla$  signifie "FIN".

Les cartes, planches, tableaux, etc., peuvent être filmés à des taux de réduction différents. Lorsque le document est trop grand pour être reproduit en un seul cliché, il est filmé à partir de l'angle supérieur gauche, de gauche à droite, et de haut en bas, en prenant le nombre d'images nécessaire. Les diagrammes suivants illustrent la méthode.



MICROCOPY RESOLUTION TEST CHART

(ANSI and ISO TEST CHART No. 2)



APPLIED IMAGE Inc

1653 East Main Street  
Rochester, New York 14609 USA  
(716) 482-0300 - Phone  
(716) 288-5989 - Fax

**Canada**  
**Department of Mines**

Hon. LOUIS CODERRE, Minister;  
R. W. BROCK, Deputy Minister.

**Geological Survey**

**Museum Bulletin No. 5**

GEOLOGICAL SERIES, No. 21

NOVEMBER 23, 1914


A BEATRICEA-LIKE ORGANISM FROM THE  
MIDDLE ORDOVICIAN

by

Percy E. Raymond

OTTAWA  
GOVERNMENT PRINTING BUREAU  
1914

No. 1467



November 23, 1914.

**Canada**  
**Geological Survey**  
**Museum Bulletin No. 5.**  
**GEOLOGICAL SERIES, No. 21.**

---

*A Beatricea-like Organism from the Middle Ordovician.*

By PERCY E. RAYMOND.

Species of *Beatricea* have long been known from the Richmond formations of Anticosti, the Manitoulin islands, Manitoba, and Kentucky, but until very recently it was not suspected that this type of organism occurred in any older strata. In a paper by George W. Stose on the "Cambro-Ordovician limestones of the Appalachian Valley in Southern Pennsylvania,"<sup>1</sup> Ulrich has listed *Beatricea* n. sp. as being very common in the Lowville. In later papers the name *Beatricea gracilis* was given to this species by Ulrich, but so far as I know, no description has yet been published. Through the kindness of Dr. Bassler, of the United States National Museum, I have been able to see specimens purporting to be *Beatricea gracilis*, and from them it would appear that the form so named is much more slender than the species about to be described. The material which I have seen showed no internal structure, and was not suitable for sectioning, and it is, therefore, impossible to make any real comparison with *Beatricea gracilis* at the present time.

The specimens about to be described were found just below the range of *Tetradium cellulosum* and *Bathyrurus extans*, and, therefore, just below the base of the Lowville. The writer's first acquaintance with these fossils was in the autumn of 1910, when he found a bed containing great numbers of them on the hill north of Aylmer, Que., in the highest layer of the formation

<sup>1</sup>Journal of Geology, Vol. 16, 1908, p. 714.

which underlies the Lowville. Although abundant, all the specimens were so badly preserved that they were at first thought to be "sea-weeds" or other indeterminate objects, until finally a single weathered specimen containing septa (Plate III, figure 2) was obtained. Being then unaware of Ulrich's finds in Pennsylvania, the writer believed he had been the first to extend the geological range of these *Beatricea*-like forms, but soon came upon Ulrich's list in Stose's paper. Further investigation revealed the presence of these fossils both in and below the Lowville at Mechanicsville, on the Ontario side of the river near Ottawa. These occurrences, listed as "*Beatricea* sp." were noted in a paper in the Ottawa Naturalist for February, 1911.<sup>1</sup>

In June, 1911, the writer had an opportunity of visiting northern New York, and there, at the top of the Upper Pamela and just below the Lowville, found the layer containing the same organisms at a number of places south of Clayton. About the same time, Mr. W. A. Johnston, of the Geological Survey, to whom I had shown my specimens, sent in from the field specimens much better than any I had found. They were obtained, as usual, from strata just below those containing *Tetradium celulosum* and *Bathyurus extans*, and the localities are near a wood road along the northern fence of lot 25, concession VI, of the town of Carden, east of Lake Simcoe, Ontario, about 150 miles west of the localities near Ottawa. This locality was visited again later in the year by Mr. Johnston and the writer, in 1912 by Mr. E. J. Whittaker, and a second time by the writer in 1913, so that a considerable amount of material has been accumulated. A list of the fossils from this locality so far determined was published by Mr. Johnston in the Summary Report of the Director of the Geological Survey of Canada for 1911 (1912), page 255. The most striking fossils are a remarkable *Tetradium* (*T. halysitoides* Raymond), *Onchometopus simplex* Raymond and Narraway, and *Bathyurus johnstoni* Raymond. On the basis of these species, which have been found by the writer both at Ottawa and in northern New York at this same horizon just

<sup>1</sup>Preliminary Notes on the "Chazy" Formation in the vicinity of Ottawa. Ottawa Naturalist, vol. 24, p. 193, 1911.

00936568

below the Lowville, these beds are correlated with the upper part of the Pamela formation of New York.

At the locality in Carden, the *Beatricea-like* fossils are found on the weathered surfaces of the flat-lying strata, and have the form of long, narrow chambered tubes. They have the general appearance of cephalopods, but are easily distinguished from them because they do not taper appreciably, though some specimens are 18 inches long, the chambers are of irregular depth, there is no siphuncle, and while usually nearly straight, many of the specimens are irregularly or abruptly curved. Most of the specimens lie parallel to the bedding planes, and are usually incomplete; but some are found which were buried in a vertical position and on cutting and polishing these, some are found to preserve a good deal of their original structure.

It will be remembered that *Beatricea* is one of those unfortunate genera whose systematic position has not been satisfactorily determined. Originally described by Billings,<sup>1</sup> these fossils have been called plants, foraminifera, rugose corals and cephalopods, and at last have found a resting place as a family among the Stromatoporoids, this position having been fixed for them by Nicholson.<sup>2</sup>

#### DESCRIPTION OF THE SPECIMENS.

This fossil has a long flexuose, cylindrical form of unknown length, only the upper termination having been seen. Fragments from 12 to 18 inches long are not uncommon. The diameter appears to be approximately uniform throughout the length of the fragments observed. Most of the specimens are from 10 to 20 mm. in diameter, but as will be hereafter shown, this does not show the true thickness of the fossil, as in nearly all specimens a part has failed of preservation. No well preserved outer surface has been seen, but from the evidence of sections of embedded specimens, it would appear that the surface is covered with small papillae. Sections perpendicular to the long axis show that the fossil is made up of concentric zones of

<sup>1</sup>Geol. Surv. of Canada, Rept. of Prog. for 1856, 1857, p. 343.

<sup>2</sup>Mono. Graph. British Stromatoporoids, Palaeontographical Society, London, 1886, p. 86-89, pl. 8, figs. 1-8.

very different structure. The inner zone seems to consist of a tube with a definite wall, this tube being divided by deeply concavo-convex sac-like transverse partitions which have the appearance of the tabulae of corals. These tabulae seem to be convex upward, or at least they occupy that position in specimens found embedded in an upright position in the strata. In sections, the centre of this tube is open and filled with transparent calcite, or often empty, while around this space, close to the wall of the tube, is an area with numerous incomplete tabulae or cystose diaphragms (Plate II, figures 1, 2; Plate IV, figures 2, 4). This central tube, therefore, corresponds to the whole section of *B. nodulosa* Billings, as shown by Nicholson in figures 2 and 3 of Plate VIII, of the publications cited above, but in *B. nodulosa* there is evidently much more of the cystose tissue than in the present species. This axial tube is nearly central in position in most of our specimens and is of variable diameter in proportion to the total diameter of the fossil. In one good section it is 3 mm. in diameter, and the total diameter is 14 mm. In another it is 5 mm. of a total diameter of 24 mm. In other forms it seems to occupy a much larger proportion, as in Plate I, figure 2.

Outside the axial tube there is a zone which in the best specimens is filled with clear calcite or is hollow, while in others it is filled with a very fine-grained brownish lime-mud containing more or less clear calcite. None of the sections show any trace of structure in this zone, and there is nothing to indicate that it ever contained any continuous skeletal structures, though it is very possible that at intervals it was crossed by some sort of supporting processes (See Plate II, figures 1 and 2; Plate IV, figures 2, et al).

The outer zone consists of a number of concentric sheaths traversed by radial canals. This structure is well shown in the best preserved of the specimens (Plate II, figures 1, 2).

The radial structure is well shown in several figures, particularly in Plates II and III. The appearance produced is that of the radial septa of corals. That these are not septa, but radial tubes with intervening pillars is, however, shown both by sections and by weathered specimens. For instance, figures

1, Plate III, and 1, Plate IV, in which the sections are not cut exactly perpendicular to the axis show wavy instead of straight lines, and a tangential section clearly shows the true nature of the structure (Plate III, figures 4, 5). In the weathered specimens the outer shell is frequently preserved, and the filling of the tubes weathered away, so as to reproduce almost exactly the original structure.

In only a single case have I found a specimen showing an end, and this specimen shows only one end (Plate I, figure 2). The specimen as now preserved is about 210 mm. long, and assuming that the natural position of the diaphragms is really convex upward, it shows the upper end of the specimen. The specimen enlarges into a sort of bulb at the top, both the inner tube and the outer sheath being enlarged. The specimen was originally weathered away about one-half at the lower end and apparently somewhat more than half at the bulbous end. Figure 2, Plate I, shows the specimen as it was found. This specimen, like the others, shows the inner tube to be entirely distinct from the outer sheaths, and where the specimen is weathered, a zone 2 mm. wide has weathered away, so that there is a narrow and deep trench on each side of the inner tube. The whole diameter of the specimen at the smaller end is 23 mm. and the tube is 13 mm. in diameter at this point. At the widest part of the bulb the whole specimen is 35 mm. across, and the bulb of the tube itself is 21 mm. wide. The specimen seems, at the bulb, to be cut down below the median plane so that these dimensions probably do not express the full size of the bulbous end. The outer sheath seems to consist of 4 or 5 thin layers and is most decidedly Stromatopora-like in appearance, even suggesting that it is an incrustation upon the inner tube. The sheath seems to have entirely enveloped the upper end, and sends out a curved portion of the sheath which extends some 20 mm. beyond the apex of the bulb. The presence of this sheath over the upper end of the organism effectually disposes of the idea that the animal could have been a coral.

The condition of preservation of the inner tube of this specimen is of some interest in connexion with the question of the possible nature of the organism. In the untapering part

of the cylinder, the tabulae or diaphragms are mostly well preserved and apparently in their natural positions. In the enlarged, bulbous part, all of the diaphragms are broken and none extends across the full width of the bulb. This portion of the tube is filled with the same limestone in which the fossil is embedded, whereas, in the cylindrical part lower down, where the diaphragms are unbroken, the chambers are, generally, filled with crystalline calcite. Beginning at the bulbous end the inner tube shows, first, 4 mm. with cystose tissue, such as is seen around the inner part of the tube in some sections. Then 39 mm. in which all diaphragms are broken and the filling is like the matrix. Next, 20 mm. of clear calcite filling. Next, three unusually deep chambers (18 mm. in all) filled with limestone matrix. Then follow 33 mm. in which the diaphragms are rather close together (twelve chambers in the interval), and the filling material is all calcite. Next a single deep chamber (6 mm.), filled with matrix, then three (6 mm. in all) with calcite, one partially empty. Next two (9 mm.), empty except for a lining of calcite crystals. Then three (13 mm.), with clear calcite. Then a deep one partially filled with clear calcite and partially with matrix. The filling with limy mud seems to have taken place after the death of the organism and after it had fallen from its erect to a horizontal position. The chambers with unfractured walls and diaphragms were evidently cut off from the supply of mud, and the filling consists of such material as could be filtered in in solutions. The mud-filled parts must, on the other hand, have had openings by which rather coarse material could enter, and that these openings were accidental and not natural is indicated by the fact that the diaphragms or side walls are seen to be fractured in most cases. The supply of mud does not, however, seem to have entered directly from the outside to the chambers which it now fills, but has been conveyed along the "empty zone" surrounding the tube. This zone is filled with clay, and served to feed mud into the inner tube through any fractures which might be present. In most cases this zone is filled with clear calcite (as in figure 1, Plate III, and figures 1 and 2, Plate IV, for instance), but in such cases there is often another and outer zone mud-filled (figure 2, Plate IV). The explanation of these "empty" zones is

not obvious. In no section so far cut does this zone show anything more than the vaguest trace of structure, and the only explanation which suggests itself to the writer is that the inner tube may have been entirely surrounded by a sheath of organic matter which depended for support on the calcareous walls on either side of it. The presence of this zone seems to militate against the idea of the outer sheaths being a separate organism parasitic upon the interior tube-like portion, for if the tube were entirely coated with organic matter, it would not be apt to attract a parasite which secreted a calcareous skeleton.

Judging from the nature of the upper end of the specimen just described, it would appear that the inner tube of this structure was not the habitation of an animal, otherwise the top would not have been enclosed. The one conception which appeals to the writer is to think of this as a colonial organism built up by numerous small polyps which secreted a skeleton at the base, the skeleton being, therefore, internal. The inner tube was a structure deposited as an axial support giving rigidity to the colony, and the zooids were on the exterior of the mass, and secreted the successive sheaths as they grew outward. The radial pores may have been the lodging places of the separate zooids, all of the pores being of the same size, and, therefore, no differentiation into gastropores and dactylopores as in *Millepora*.

This way of thinking of the organism also serves to explain the empty rings which are otherwise so puzzling. Among the Hydromedusae we have the order Tubulariae, in which a chitinous periderm is secreted. It is quite possible that the zooids of this early Palaeozoic organism secreted a basal layer of calcareous matter and then surrounded themselves at the sides with chitinous material. The next generation founded their calcareous basement upon the chitinous layer of their progenitors, thus producing alternating bands of calcareous and chitinous material. After death, the calcareous material being much more resistant than the chitinous is preserved, while the more easily decomposed material is replaced by an infiltration of mud or crystalline calcite.

Owing to the slight development of the cystose tissue and

the great development of the sheath zone in the form under consideration, it does not seem proper to refer it to *Beatricea* and the following new generic name is, therefore, proposed.

Genus CRYPTOPHRAGMUS *nov.*

(*Kryptos, concealed phragmos, partition.*)

Colonial organisms, presumably Hydromedusae, of elongate, upright, unbranched form, small diameter, and slight taper. Skeleton consisting of internal camerate tube with distinct wall, the tube crossed by irregularly spaced partitions which are convex upward, and further strengthened by a deposit of incomplete partitions (cystose material) on the inner side of the wall of the tube. The inner tube enveloped by concentric, often slightly separated sheaths of calcareous material, the sheaths traversed by numerous circular openings at right angles to the long axis of the colony. The sheaths extend over the upper end of the inner tube, which may be enlarged and bulb-like. Type, *Cryptophragmus antiquatus* sp. nov.

CRYPTOPHRAGMUS ANTIQUATUS *sp. nov.*

Plates I-IV.

Since I have not differentiated the rather numerous varieties that are found in the locality at Carden, the generic description will serve also for the species. Should Ulrich's *Beatricea gracilis* prove to belong to this genus, it may be necessary to enter into the question of specific characters. At present, *Beatricea gracilis* has the status of *nomen nudum*. It may be possible to differentiate species on the basis of the relation of the diameter of the inner tube or support to the total diameter of the colony, in which case the specimens figured on Plate I, figure 2, would not belong to the same species as those shown in figures 3 and 4, Plate I, and 1 and 2, Plate II. In view of this contingency, I have designated the specimens with the large tubes as the types of the species *C. antiquatus*, and, in particular, would designate the most complete specimen Plate I, figure 1, as the holotype of the species.

Some of the specimens have the axial tube 15 mm. in diameter

and one such has the clear band 2.5 mm. wide. The presence of this structureless "clear zone" explains the state of preservation of most of the specimens found on the weathered surfaces of the rocks. It being either hollow or filled with a soft substance, the outer sheath zone and the axial tube were easily separated, and in the specimens usually found such separation had occurred before the specimens were buried in the rocks.

Summary. Transverse sections of *Cryptophragmus antiquatus* show that it is composed of three distinctly separated and well marked concentric layers. The inner axial tube has a large cavity, divided into chambers by bulging partitions, and around this cavity is a narrow band of cystose tissue, without radial elements of any kind. The axial tube has a well-defined outer wall, and is surrounded by the second zone which seems to be structureless and is usually marked by a band of clear calcite. The outer zone is composed of concentric sheaths traversed by numerous radial canals.

Comparison with *Beatricea nodulosa* and *B. undualata*.

The skeleton of this species has a much smaller habit than either of Billings's species and the surface is evidently much more nearly smooth. As to the internal structure, as revealed by thin transverse sections, there seem to be rather striking differences. A distinct differentiation into three zones exactly comparable to those described above has not been noted by students of the above species. As noted above, the sections given by Nicholson in figures 2 and 3 of Plate VIII of his article do not appear to represent a section of a complete skeleton, as Nicholson supposed, but are comparable to the axial tube of *Cryptophragmus antiquatus*. If this latter comparison is correct, then the central tabulate zone of *B. nodulosa* is very small as compared with that in the axial tube of *B. antiquata*, and the surrounding cystose zone very thick. In *C. antiquatus* there are no indications of radial elements in the cystose zone, nor are there anywhere the granular deposits on the walls, such as both Nicholson and Parks have described. As seen in this section the walls are perfectly sharp, and though each wall has a cloudy border, this border, as seen under a high power, seems to be due to alter-

ation products of the wall distributed along minute cracks in the calcite which fills the cells.

While descriptions of *Beatricea nodulosa* and *B. undulata* do not specifically describe the three zones, yet an inspection of the macroscopic character of specimens from Anticosti shows that both the outer or "sheath" zone and the axial tube are present in well preserved specimens, and certain of them indicate the presence of a "clear" zone as well. Parks, in his "Ordovician Stromatoproids,"<sup>1</sup> thus describes a section of *B. undulata* which, unfortunately, he does not figure: "This specimen is 3 mm. thick and presents in cross section a series of concentric layers of very different aspect. The inner tube has a radius of 3 mm. This is surrounded by a ring, 20 mm. thick, of ordinary vesicular tissue with the granular element well developed, but with scarcely a trace of radial pillars. Surrounding this ring is an outer zone, 15 mm. thick, which is fairly well demarcated by a sharp line of separation. This outer layer is strikingly different from the middle annulus, being composed of continuous laminae and well marked radial pillars."

Parks does not state that his inner tube, only 3 mm. in radius, has a distinct wall, and as he copies Nicholson's figures where such a wall is absent, we are inclined to think that both his inner tube and the 20 mm. thick band of vesicular tissue around it are to be correlated with the "inner tube" of *C. antiquatus*. The outer band is probably the same as our "sheath" zone, and there is apparently no "clear" zone present.

Some of Nicholson's sections seem to have been cut from the "sheath" zone of *B. nodulosa*, for he states in a footnote that in one section he noticed perpendicular calcareous septa crossing the vesicles. The relation of the present form to *Beatricea* is now being studied by the writer, and will probably be set forth in a later paper.

The photographs which illustrate this article were, with the exception of figure 2, Plate I, made at the Geological Survey of Canada and all the types are in the collections of this Survey. Except where otherwise noted, the specimens are from the locality in Carden.

<sup>1</sup>Univ. of Toronto Studies, Geological Series, No. 7, p. 44, 1910.


te cracks in


*undulata* do  
ection of the  
shows that  
are present  
indicate the  
Ordovician  
*undulata*,  
specimen is 75  
concentric  
diameter of only  
of ordinary  
developed, but  
this ring  
demarked  
strikingly  
continuous

3 mm. in  
s figures,  
that both  
lar tissue  
of *C. anti-*  
"sheath"

cut from  
footnote  
is septa  
form to  
probably

e, with  
urvey of  
Survey.  
om the






## EXPLANATION OF PLATE I.


*Cryptophragmus antiquatus* Raymond.

- Figure 1. Fragment of a specimen from which only a part of the outer sheath have been removed by weathering. In places where the specimen is broken the transverse partitions may be seen. The figure is in a reversed position, up side down. About natural size.
- " 2. A photograph of the only specimen so far found preserving a termination. The bulb-like enlargement of the upper end is shown, also the curved extension of the outer sheath above. (Indicated by arrow). Notice the calcite-filled chambers (A), the empty ones (B), and those filled with lime-mud (C). Six-tenths natural size.
- " 3. Photograph of polished longitudinal section showing inner tube with convex partitions, the "clear zone" near the tube, and the sheaths outside. Natural size.
- " 4. The upper, naturally weathered end of the same specimen, enlarged. Notice the differential weathering, and the faintly indicated radial tubes of the sheaths.

outer sheaths  
specimen  
figure is  
size.  
termination.  
n, also the  
by arrow).  
(B), and  
e.  
tube with  
the sheaths  
enlarged.  
indicated









## EXPLANATION OF PLATE II.

*Cryptophragmus antiquatus* Raymond.

Figures 1 and 2. Transverse and longitudinal thin sections of a specimen, enlarged 4 diameters. Note the inner tube with its convex partition and wall of vesicular tissue, the alteration of wall preservation and lime-filled zones, and the radial tubes. For other sections of this same specimen, see Plate III, figures 4 and 5.



specimen, ca-  
x partitions  
t preserved  
er sections



The first part of the document discusses the importance of maintaining accurate records of all transactions. It emphasizes that every entry should be supported by a valid receipt or invoice. This ensures transparency and allows for easy verification of the data.

In the second section, the author outlines the various methods used to collect and analyze the data. This includes both primary and secondary data collection techniques. The analysis focuses on identifying trends and patterns over time, which is crucial for making informed decisions.



The third part of the document provides a detailed breakdown of the results. It shows that there has been a significant increase in sales volume, particularly in the middle and lower income brackets. This suggests that the current marketing strategy is effective in reaching a wider audience.

Finally, the document concludes with several key recommendations. It suggests that the company should continue to invest in research and development to stay ahead of the competition. Additionally, it recommends a more targeted marketing approach to maximize the return on investment.

## EXPLANATION OF PLATE III.

*Cryptophragmus antiquatus* Raymond.

- Figure 1. A somewhat diagonal transverse thin section, showing the central tube with its vesicular wall, the cross-section of the partition and the radial tubes of the outer sheath zone. Between the tube wall and the sheath zone is an area of calcite in which the crystals are coated with a brown substance, thus producing, in the photograph, an appearance of structure in the "clear zone." Enlarged 3 diameters.
- " 2. A specimen from near Ottawa, showing the somewhat abrupt curvature of some specimens. From the top of the Pamela, 2 miles north of Aylmer, Que. Natural size.
- " 3. Transverse section just at the top of one of the convex partitions, the dark circle at the centre being the top. Just what the meaning of the structures seen in this section may be has not been determined. X 3.
- " 4. Tangential section of the specimen shown in Plate II, showing the exceedingly numerous tubes in the sheath. X 3.
- " 5. Section similar to figure 4, and from the same specimen, but not strictly tangential. At the centre it cuts into one of the structureless zones. X 3.



g the central  
ee partitions,  
een the tube  
a the crystals  
n the photo-  
" Enlarged

abrupt curv-  
elia, 2 miles

t partitions,  
he meaning  
been deter-

showing the


n, but not  
structure-

2

3

4

5





## EXPLANATION OF PLATE IV.

*Cryptophragmus antiquatus* Raymond.

- Figure 1. A transverse section, not at right angles to the axis. X 3.  
 " 2. A transverse section at right angles to the axis, shows the inner wall with definite wall, a "clear zone," then a pore-pierced sheath zone, a lime-mud filled structureless zone, and finally an outer sheath zone. Notice the papillate nature of the outer surface where the specimen is still in contact with the matrix, and note that the elevations are at the apertures of the tubes, and not at the tube walls. Also that the tubes are filled with calcite and not with mud. X 3.  
 " 3. A rather badly preserved transverse section. X 3.  
 " 4. A transverse section in which the vesicular wall of the inner tube is especially well preserved. X 3.  
 " 5. A specimen in which the inner tube only is preserved, the sheath being entirely destroyed. Such specimens are very frequently found  $\frac{1}{2}$  natural size.



X 3.  
the inner tube  
and sheath zone,  
an outer sheath  
surface where the  
note that the  
not at the tube  
and not with

the inner tube  
and, the sheaths  
very frequently

5





The first number of the Museum Bulletin was entitled, *Victoria Memorial Museum Bulletin Number 1*.

The following articles of the Geological Series of Museum Bulletins have been issued.

*Geological Series*

1. The Trenton crinoid, Ottawacrinus, W. R. Billings, by F. A. Bell.
2. Note on Mercopinus, Walcott, by F. V. Bathur.
3. The occurrence of Helicot teeth at Roche Mouton and vicinity, Alberta, by F. M. Lande.
4. Notes on Cyclocoptoides, by P. L. Raymond.
5. Notes on some new and old Tribolites in the Victoria Memorial Museum, by P. L. Raymond.
6. Description of some new Asaphids, by P. L. Raymond.
7. Two new species of Tetradium, by P. L. Raymond.
8. Revision of the species which have been referred to the genus Bathynurus (preliminary paper), by P. L. Raymond.
9. A new Brachiopod from the base of the Trias, by A. F. Wilson.
10. A new genus of dicotyledonous plant from the Tertiary of Kettle river, British Columbia, by W. F. Wilson.
11. A new species of Lepidostrobus, by W. F. Wilson.
12. Prehnite from Adams sound, Admiralty Inlet, Bathin Island, Franklin, by R. A. A. Johnston.
13. The origin of granite-micropegmatites in the Purcell hills, by S. J. Schofield.
14. Columnar structure in limestone, by F. M. Kindle.
15. Supposed evidences of subsidence of the coast of New Brunswick within modern time, by J. W. Goldthwait.
16. The Pre-Cambrian (Beltian) rocks of southeastern British Columbia and their correlation, by S. J. Schofield.
17. Early Cambrian stratigraphy in the North American Cordillera, with discussion of the Albertella and related faunas, by E. D. Burling.
18. A preliminary study of the variations of the plications of Pterostrophia hemiplicata, Hall, by A. E. Wilson.
19. The Anticosti Island faunas, by W. H. Fyviebedel.
20. The Crownsnest Volcanics, by J. D. Mackenzie.

