

748, m

Flora Americæ Septentrionalis;
OR, A
SYSTEMATIC ARRANGEMENT
AND
DESCRIPTION
OF
The Plants
OF
NORTH AMERICA.
CONTAINING,
BESIDES WHAT HAVE BEEN DESCRIBED BY PRECEDING AUTHORS,
MANY NEW AND RARE SPECIES,
COLLECTED DURING TWELVE YEARS' TRAVELS AND RESIDENCE
IN THAT COUNTRY.

BY FREDERICK PURSH.

IN TWO VOLUMES.
WITH TWENTY-FOUR ENGRAVINGS.

VOL. II.

SECOND EDITION.

London:

PRINTED FOR JAMES BLACK AND SON,
TAVISTOCK-STREET, COVENT-GARDEN.

MDCCXVI.

QK110.P9
1816
V.2

XII.

423. BEJARIA
7-locula
424. TILIA.
sub-5-lo
5-sperm
425. HELIAN
qualibus
locularis,
426. HUDSON
Stam. 10
427. TALINUM
cularis, 3
428. PORTUL
cularis, c
429. CHELIDONI
Caps. 1-
430. GLAUCI
Siliqua 2
tata,
431. PAPAVEI
1-locular
432. SANGUINI
ovata, 1-
433. ARGEMONE
Caps. ser
434. PODOPHY
1-loculari
435. ACTÆA.
1-loculari
436. SARRACENIA
Pet. 5.
peatus.
VOL. II.

XII. POLYANDRIA.

I. MONOGYNIA.

423. BEJARIA. *Cal.* 7-fidus. *Pet.* 7. *Stam.* 14. *Caps.* 7-locularis, polysperma.
424. TILIA. *Cal.* 5-partitus, deciduus. *Pet.* 5. *Stigma* sub-5-lobum. *Caps.* immatura 5-locularis, 5-valvis, 5-sperma: matura submonosperma, basi dehiscens.
425. HELIANTHEMUM. *Cal* laciniis 5. sæpius inæqualibus: 2. extimis minoribus. *Pet.* 5. *Caps.* 1-locularis, 3-valvis; valvis medio septiferis.
426. HUDSONIA. *Cal.* 3-partitus, tubulosus. *Pet.* 5. *Stam.* 10—15. *Caps.* 1-locularis, 3-valvis, 3-sperma.
427. TALINUM. *Cal.* 5-phyllo. *Pet.* 5. *Caps.* 1-locularis, 3 valvis, polysperma.
428. PORTULACCA. *Cal.* 2-fidus. *Pet.* 5. *Caps.* 1-locularis, circumscissa.
429. CHELIDONIUM. *Cal.* 2-phyllo, deciduus. *Pet.* 4. *Caps.* 1-locularis, linearis. *Sem.* plurima, cristata.
430. GLAUCIUM. *Cal.* 2-phyllo, deciduus. *Pet.* 4. *Siliqua* 2-locularis, linearis. *Sem.* plurima, punctata.
431. PAPAVER. *Cal.* 2-phyllo, deciduus. *Pet.* 4. *Caps.* 1-locularis, sub stigmate persistente poris dehiscens.
432. SANGUINARIA. *Cal.* 2-phyllo. *Pet.* 8. *Caps.* ovata, 1-locularis.
433. ARGEMONE. *Cal.* 3-phyllo, deciduus. *Pet.* 6. *Caps.* semivalvis.
434. PODOPHYLLUM. *Cal.* 3-phyllo. *Pet.* 9. *Bacca* 1-locularis, coronata stigmate.
435. ACTÆA. *Cal.* 4-phyllo, deciduus. *Pet.* 4. *Bacca* 1-locularis. *Sem.* semiobtusata.
436. SARRACENIA. *Cal.* duplex, 3-phyllo et 5-phyllo. *Pet.* 5. *Caps.* 5-locularis. *Stylus* stigmate clypeatus.

437. LEWISIA. *Cal. 7—9-phillus, scariosus. Pet. 14—18. Stylus 3-fidus. Caps. 3-locularis, polysperma. Semina nitida.*
438. NYMPHÆA. *Cal. 4—5-phillus. Pet. plurima, germini sub staminibus inserta. Stigma radiatum, sessile, medio nectariferum. Baccæ multilocularis, polysperma.*
439. NUPHAR. *Cal. 5—6-phillus. Pet. plurima, receptaculo cum staminibus inserta, dorso nectarifera. Stigma radiato-sulcatum, sessile. Baccæ multilocularis, polysperma.*
440. CERATOPHYLLUM. *Cal. multipartitus : femineus 6-phillus, imbricatus. Cor. 0. Anth. tricuspidatae. Nux 1-sperma. Monoicum.*

II. DI-PENTAGYNYIA.

441. CALLIGONUM. *Cal. 5-partitus. Cor. 0. Styli 4. Nux 1-locularis, 4-alata. Dioicum.*
442. DATISCA. *Cal. 5-phillus : femineus 2-dentatus. Cor. 0. Anth. sessiles, oblongæ, 15. Styli 3. Caps. 3-angularis, 3-cornis, 1-locularis. Dioica.*
443. MENISPERMUM. *Cal. 0. Pet. 4. exteriora ; 8. interiora. Stami. 6. 8. 12. 16. Pist. 2—3. Baccæ 1—2, monospermæ. Dioicum.*
444. DELPHINIUM. *Cal. 0. Pet. 5. Nectar. 2-fidum, postice cornutum. Caps. 1—3. seu 5.*
445. ACONITUM. *Cal. 0. Pet. 5. supremo fornicato. Nectar. 2. pedunculata, recurva. Caps. 3. seu 5.*
446. AQUILEGIA. *Cal. 0. Pet. 5. Nectar. corniculata, inter petala. Caps. 5. distinctæ.*
447. CIMICIFUGA. *Cal. 4—5-phillus. Pet. 4—8. cartilaginea. Caps. 1—5 seu plures, oblongæ, sutura laterali dehiscentes, polyspermæ.*
448. ASCYRUM. *Cal. 4-phillus : 2. interioribus majoribus. Pet. 4. Filam. in 4. phalanges digesta. Caps. oblonga, 1-locularis, 2-valvis, calyce inclusa.*
449. HYPERICUM. *Cal. 5-partitus, laciniis subæqualibus. Pet. 5. Filam. vix basi connata. Caps. subrotunda : loculis numero stylorum : 1—2—3—5.*
450. ELODEA. *Cal. 5-partitus, æqualis. Pet. 5. unguibus nectariferis. Filam. 10—15. in 3. phalanges connata. Glandulæ inter phalanges. Styli 3. divergentes. Caps. 3-locularis.*

451. ILLICIUM. in orbe
452. MAGNOIA. 2-valves, pendula.
453. LIRIODEMA. maræ im non dehis.
454. PORCELIANA. jora. St abortione spermæ.
455. ATRAGEA. 4. spathu dilatata su
456. CLEMATIS. cauda plu
457. ANEMONE.
458. THALICTRUM. gissima.
459. HYDRASTIS. acinis mo
460. BRASENIA. riora long
461. CALTHA. polyspermæ
462. COPTIS. cucullata. polyspermæ
463. TROLLIUM. 5—8. lir spermæ.
464. HEPATICUM. nuda.
465. RANUNCULUS. gues poro
466. SAGITTARIUS. gregatæ,
467. CYAMUS. turbinatus. 1-spermis.

Pet. 14—
polysperma.

irima, ger-
atum, ses-
cularis, po-

ma, recep-
iectarifera.
multilocu-

femineus
cuspitatae.

1. *Styli* 4.
-dentatus.
3. *Caps.*
eriora; 8.
Bacce
2-fidum,
fornicato.
.. seu 5.
orniculata,
4—8. car-
æ, sutura
as majori-
ta. *Caps.*
sa.
ubæquali-
Caps. sub-
—5.
5. ungu-
phalanges
yli 3. di-

III. POLYGYNIA.

* *Flores vagi.*

451. ILLICIUM. *Cal.* 6-phyllus. *Pet.* 27. *Caps.* plures, in orbem digestæ, 2-valves, 1-spermæ.
 452. MAGNOLIA. *Cal.* 3-phyllus. *Pet.* 6—9. *Caps.* 2-valves, 1-spermæ, in strobilum imbricatæ. *Sem.* pendula.
 453. LIRIODENDRON. *Cal.* 3-phyllus. *Pet.* 6. *Samarœ* imbricatae in strobilum. *Caps.* 1—2-spermæ, non dehiscentes.
 454. PORCELIA. *Cal.* 3-phyllus. *Pet.* 6. interiora majora. *Stigmata* sessilia, obtusa. *Baccæ* plures aut abortione subsolitariae, ovoideo-subreniformes, polyspermæ. *Sem.* sutura interna adnexa, arillata.
 455. ATRAGENE. *Cal.* 0. *Cor.* 4—6-petala. *Nectaria* 4. spathulata, petalis alternantia. *Filam.* exteriora dilatata subantherifera. *Sem.* aristata, aristis pilosis.
 456. CLEMATIS. *Cal.* 0. *Pet.* 4—6. *Sem.* caudata, cauda plumosa.
 457. ANEMONE. *Cal.* 0. *Pet.* 6—9. *Sem.* ecaudata.
 458. THALICTRUM. *Cal.* 0. *Pet.* 4—5. *Stam.* longissima. *Sem.* ecaudata, striata, teretia.
 459. HYDRASTIS. *Cal.* 0. *Pet.* 3. *Bacca* composita, acinus monospermis.
 460. BRASENIA. *Cal.* 0. *Pet.* 6. persistentia: 3. interiora longiora. *Caps.* 6—10. oblongæ, dispermæ.
 461. CALTHA. *Cal.* 0. *Pet.* 5. decidua. *Caps.* plurimæ, polyspermæ.
 462. COPTIS. *Cal.* 0. *Pet.* 5—6. caduca. *Nectar.* 5—6. cucullata. *Caps.* 5—8. stipitatæ, stellatæ, rostratæ, polyspermæ.
 463. TROLLIUS. *Cal.* 0. *Pet.* 5—8. decidua. *Nectar.* 5—8. linearia. *Caps.* plurimæ, sessiles, polyspermæ.
 464. HEPATICA. *Cal.* 3-phyllus. *Pet.* 6—9. *Sem.* nuda.
 465. RANUNCULUS. *Cal.* 5-phyllus. *Pet.* 5. intra unguis poro mellifero. *Sem.* nuda.
 466. SAGITTARIA. *Cal.* 3-phyllus. *Pet.* 3. *Caps.* aggregatae, 1-spermæ. *Polygama.*
 467. CYAMUS. *Cal.* 4—5-phyllus. *Pet.* plurima. *Fructus* turbinatus, in disco truncato foveis plurimis excavatus 1-spermis. *Nuces* ovatae, stylo persistente coronatae,

** *Flores spathacei. Spadix simplex.*

468. POTHOS. *Spatha cucullata. Spadix floribus tectus.*
Cal. O. Pet. 4. Baccæ 2-spermæ.
 469. CALLA. *Spatha planiuscula. Spadix floribus tectus.*
Cal. et Cor. O. Baccæ polyspermæ.
 470. ARUM. *Spatha cucullata. Spadix supra nudus,*
inferne feminineus, medio stamineus. Cal. et Cor. O.
Baccæ utplurimum 1-spermæ.
-

I. MONOGYNIA.

423. BEJARIA. *Gen. pl. 811. Vent. hort. cels p. 51.*

racemosa.

1. *B. foliis ovato-lanceolatis glabris, floribus racemoso-paniculatis terminalibus.* — *Vent. l. c.*
Befaria paniculata. Mich. fl. amer. 1. p. 280.
Icon. Mich. l. c. t. 26. Vent. l. c. t. 51.
α. B. foliis utrinque acutis, caule hispido.
β. B. foliis obovatis obtusiusculis, caule glabro.
 In sandy woods, on the banks of swamps and ponds :
 Georgia and Florida : particularly plentiful on the
 Island of St. Mary's. $\frac{1}{2}$. June. v. v. A beautiful
 shrub, from three to four feet high ; flowers white,
 tinged with red, of an agreeable scent.

424. TILIA. *Gen. pl. 894.*

glabra.

1. *T. foliis suborbiculato-cordatis abrupte acuminatis argute serratis subcoriaceis glabris, petalis apice truncatis crenatis, stylo petalis subæquali, nuce ovata subcostata.* — *Vent. in mem. de l'instit. sc. phys. 4. p. 9. t. 2.*
T. americana. Willd. sp. pl. 2. p. 1162.
T. caroliniana. Wangh. amer. 56.
T. canadensis. Mich. fl. amer. 1. p. 306.
Icon. Vent. l. c. t. 2. Mich. arb. t.
 In the woods of Canada and the northern United States,
 and on the mountains, as far south as Carolina. $\frac{1}{2}$.
 May, June. v. v. This tree is known by the name
 of Lime- or Line-tree; Basswood; Spoonwood; and is
 both useful and ornamental.

2. *T. foliis coriaceis natis, stylo amer. 1.*
Icon. Mich.

Near the sea.
 June. v. generally

3. *T. foliis bas serratissimis su petalis en bosa laevi.*
t. 3. Wi

T. caroliniana
T. americana

β. T. foliis laxe l. c. p. 11.
 In close copses
 Georgia.
 can scarce flowers and
 have not seen

4. *T. foliis ovati datisque arg globosa sul*
 On the banks
 v. s. A ver

425. HELI

1. *H. exstipulata lanceolatis libus paucifl capsulis caly sub Cisto.*

In dry fields at
 June. v. v.
 yellow.

2. *H. exstipulata foliis alternis tomentosis, ris, calycis globosis lor p. 307.*

2. *T. foliis cordatis sensim acuminatis rarer dentatis membranaceis glabris, paniculis laxifloris, petalis emarginatis, stylo petalis longiore, nuce globosa.* — *Mich. fl. amer. 1. p. 306.*

Icon. Mich. arb. t.

Near the sea-coast: Maryland to Georgia. $\frac{1}{2}$. May, June. *v. v. s. fl.* A very distinct species, though generally confounded with the foregoing one.

3. *T. foliis basi truncatis obliquis acuminatis denticulato-serratis subtus pubescentibus, paniculis confertifloris, petalis emarginatis, stylo petalis subæquali, nuce globosa lœvi.* — *Vent. in mem. de l'instit. sc. phys. 4. p. 10. t. 3. Willd. sp. pl. 2 p. 1162.*

T. caroliniana. Mill. dict. no. 4.

T. americana. Walt. fl. car. 153. Wangh. amer. 55.

- β . *T. foliis laxe serratis tenuissimis subpapyraceis. Vent. leptophylla. l. c. p. 11.*

In close copses and on the banks of rivers: Virginia to Georgia. $\frac{1}{2}$. May—July. *v. v.* The variety β . can scarcely be considered a distinct species, unless the flowers and fruit should prove it so, both of which I have not seen.

4. *T. foliis ovatis basi oblique aut æqualiter truncatis cor-datisque argute serratis subtus niveo-tomentosis, nuce globosa subcostata.* — *Vent. l. c. p. 16 t 5.*

On the banks of the Ohio and Mississippi. $\frac{1}{2}$. June. *v. s.* A very handsome and desirable ornamental tree.

425. HELIANTHEMUM. Juss. gen. pl. 326.

Cisti spec. Gen. pl. 913.

1. *H. exstipulatum, erectum; foliis alternis erectis linearilanceolatis planis subtus tomentosis, racemis terminibus paucifloris, calycis laciniis lato-ovatis acuminatis, capsulis calyce brevioribus.* — *Willd. sp. pl. 2. p. 1199. sub Cisto.*

In dry fields and pine woods: Canada to Carolina. $\frac{1}{2}$. June. *v. v.* Flowers, as in all the following species, yellow.

2. *H. exstipulatum, erectum, pulvriulento-tomentosum; ramuliflorum.* foliis alternis oblongis acutis margine revolutis subtus tomentosis, racemis brevissimis, summitate subtrifloris, calycis laciniis orbiculatis pulverulentis, capsulis globosis longitudine calycis.

— *Mich. fl. amer. 1. p. 307.*

laxiflora.

pubescens.

canadense.

POLYANDRIA MONOGYNIA. Helianthemum.

Menandra ramis alternis. *Clayt. fl. virg.* 20.

In sandy fields: New Jersey to Carolina. 4. July.
v. v. Flowers small.

427.

1. *T. foliis cy dunculat*
On sunny
v. v. F

428.

1. *P. foliis cu pl. 2. p.*
Icon. *Schk*
Purslane is

429.

1. *C. foliis alt dunculat*
Icon. *Fl. a*
Near old
and Pen
June. i
breaking
in curing

2. *C. catifibus*
silibus lo
capsula c
In shady w
Michaux
very mu
both gen

430.

1. *G. caule gl peduncu*
Smith fl
Chelidoni
Icon. *Fl. a*
On the se
July. v

4.

1. *P. capsulis subpinna*
—*Willa*

corymbosum.

3. *H. exstipulatum, erectum, ramosum, minutim pubescens; foliis alternis lanceolatis subtus cano-tomentosis, corymbis fastigiatis confertim multifloris, calycis lacinias ovatis acutis, capsulis calyce vix longioribus.* —
Mich. fl. amer. 1. p. 307.

In dry barren sand-fields, particularly along the sea-coast: New Jersey to Georgia. 4. June—Aug.
v. v. Flowers very small, sometimes without petals; from which circumstance this and the foregoing species have been mistaken for *Lechea major*.

*rosmarinifo-
lium.*

4. *H. exstipulatum, stricte-erectum, simplex, canescens; foliis linearibus margine revolutis subtus cano-tomentosis, racemulis axillaribus interrupte confertifloris, calycis lacinias ovatis acutis, petalis calyce triplo longioribus.*

In pine-barrens: Georgia. *Enslen.* 4. July. v. s.
in *Herb. Enslen.* Flowers bright yellow.

carolinianum.

5. *H. exstipulatum, hirsutum, erectum; foliis alternis oblongo-ovalibus subdenticulatis, imis obovalibus utrinque hirsutis, pedunculis terminalibus paucis calycibusque villosissimis, calycis lacinias oblongis acutis petalis brevioribus.* —
Mich. fl. amer. 1. p. 307.

Cistus carolinianus. *Walt. fl. car.* 152.
Icon. *Vent. cels* 74.

In pine-barrens of Carolina and Georgia. 4. June—
Aug. v. v. Flowers and leaves larger than in any other species.

426. HUDSONIA. *Gen. pl.* 822. *Willd. hort. berol.* 15.

ericoides.

1. *Hudsonia.* *Willd. sp. pl.* 2. p. 858.

Icon. *Willd. hort. berol.* 15. *Berg. act. holm.* 1778. t. 2.
In pine-barrens of New Jersey and Virginia. 4. May,
June. v. v. Flowers yellow. This delicate little
shrub, not above six inches high, covers large tracts
of sandy pine-woods in a similar manner as common
heath.

427. TALINUM. *Juss. gen. pl.* 346.

1. *T. foliis cylindricis carnosis, corymbis terminalibus pendunculatis.* — *teretifolium.*

On sunny rocks: Delaware and Virginia. 4. July.
v. v. Flowers purple.

428. PORTULACCA. *Gen. pl.* 824.

1. *P. foliis cuneiformibus, floribus sessilibus.* — *Willd. sp.* *oleracea.*
pl. 2. p. 859.

Icon. *Schkuhr handb.* 130.

Purslane is a common weed in gardens.

429. CHELIDONIUM. *Gen. pl.* 880.

1. *C. foliis alternis pinnatis lobatis, umbellis axillaribus pendunculatis.* — *Willd. sp. pl. 2. p. 1141.*

majus.

Icon. *Fl. dan.* 542. *Engl. bot.* 1581.
 Near old buildings and among rocks: New England
 and Pennsylvania; probably introduced. 4. May,
 June. *v. v.* The yellow fluid which it affords by
 breaking the stems, is considered an infallible remedy
 in curing warts and corns.

2. *C. catilobis in summitate geminatim diphylloides, foliis sessilibus lobato-pinnatifidis, pedunculo solitario unifloro,*
capsula oblongo-ovata. — *Mich. fl. amer.* 1. p. 309.

diphyllum.

In shady woods, near rivulets: Kentucky and Tennessee.
Michaux. 4. May. *v. s.* Resembles *Sanguinaria*
 very much, and seems to be the connecting link to
 both genera.

430. GLAUCIUM. *Juss. gen. pl.* 261.

1. *G. caule glabro, foliis caulinis amplexicaulibus repandis,*
pedunculis unifloris, siliqua tuberculato-scabriuscula.
Smith fl. brit. 2. p. 563.

luteum.

Chelidonium Glaucium. — *Willd. sp. pl. 2. p. 1142.*

Icon. *Fl. dan.* 585. *Engl. bot.* 8.

On the sea-coast of Virginia and Carolina. ①. June,
 July. *v. v.* Flowers large, yellow.

431. PAPAVER. *Gen. pl.* 881.

1. *P. capsulis hispidis, scapo unifloro nudo hispido, foliis subpinnatis, foliolis lanceolatis; inferioribus subincisis.*
— Willd. sp. pl. 2. p. 1145.

nudicaule.

POLYANDRIA MONOGYNIA. Papaver.

Icon. *Fl. dan.* 41. *Dill. elth. t.* 224. *f.* 291.
In Labrador. *Colmaster.* ♂. *v. s. in Herb. Dickson.*
Flowers yellow.

432. SANGUINARIA. Gen. pl. 878.

canadensis.

1. *S. folio subreniformi sinuato-lobato, scapo unifloro.*—
Willd. sp. pl. 2. p. 1140.

Icon. *Bot. mag.* 162. *Dill. elth. t.* 252.

stenopetala.

- β. *S. petalis linearibus.*
In dry woods, generally in fertile soil : Canada to Florida. ♀. March, April. v. v. Flowers white ; the number of petals is variable ; and with little care and attention a fine double variety might be produced. It is known by the name of *Blood-wort.*

433. ARGEMONE. Gen. pl. 882.

mexicana.

1. *A. capsulis 5-valvibus, foliis pinnatifidis incisis spinosis,*
floribus axillaribus.—*Willd. sp. pl. 2. p. 1148.*

Icon. *Bot. mag.* 243.

On banks of rivers : Carolina to Florida. ♂. July, Aug.
v. v. Flowers yellow, large. In Georgia is a variety
with white flowers, which has a beautiful appearance.

434. PODOPHYLLUM. Gen. pl. 879.

peltatum.

1. *P. caule unifolio unifloro, folio peltato palmato : lobis*
cuneatis incisis.—*Willd. sp. pl. 2. p. 1141.*

Icon. *Trew ehret. t.* 29. *Catesb. car. 1. t.* 24.

In shady woods, in large patches, generally on moist ground : New England to Carolina. ♀. March, April v. v. Flower white ; the fruit is the size of a common plum, green, eatable, and known by the name of *May-apple* : the root is sometimes used as Ipecacuanha.

435. ACTÆA. Gen. pl. 877.

americana.

1. *A. foliis decompositis : foliolis oblongis sensim acutis*
inciso-serratis, racemo ovato, petalis staminibus bre-
vioribus.

Α. *spicata.* *Mich. fl. amer. 1. p. 308.*

Α. *rubra.* *Willd. enum. 500.*

α. *A. baccis niveis.*

Icon. *Corn. canad. t. 77.* *Moris. hist. 2. s. 1. t. 2. f. 7.*

β. *A. baccis*
In shady
to Vir
May.
knowl
consid
spicata
the let

43

1. *S. foliis*
tato, a
subreni
p. 310
Icon. *Bot*
t. 70.
In cedar
Carolin
large,
the foli
of the a
pots fill

2. *S. foliis el*
pendicei
ala vet
p. 310.
S. minor.
S. adunca.
In open sw
rida. ♀
transpare
this spec
it has oft

3. *S. foliis sti*
patula, a
coarctata
bulato.
p. 1150.
Icon. *Bot.*
t. 69. P
In open swa
v v. Flo

rubra.

5. A. baccis rubris.

In shady rocky woods, in rich vegetable mould: Canada to Virginia; principally on the mountains. ♀. April, May. v. v. Flowers white; berries white or red; known by the name of *Red and White Cohosh*, and considered by the natives as a valuable medicine. *A. spicata* differs not only in its foliage, but the petals are the length of the stamens.

436. SARRACENIA. *Gen. pl. 885.*

1. *S. foliis brevibus, tubo ventricoso-gibbo, fauce coarcata, ala ventrali arcuatim elatiore, appendice erecta subreniforme-cordata mutica sessili.* *Mich. fl. amer.* 1. p. 310. *Willd. sp. pl.* 2. p. 1150. *Icon. Bot. mag.* 849. *Mill. ic.* 241. *Catesb. car.* 2. t. 70. *Pluk. amalh.* t. 376. f. 6. *purpurea.*
In cedar swamps and sphagnum marshes: Canada to Carolina. ♀. June, July. v. v. Flowers purple, large. This singular and handsome plant is, with all the following, a very desirable object in the collection of the admirers of nature; they bear cultivation, in pots filled half way with sphagnum, extremely well.
2. *S. foliis elongatis, tubo superne dorso maculato, in appendicem fornicate incircumflexum, ala ventrali linear-lanceolata.* *Mich. fl. amer.* 1. p. 310. *S. minor.* *Walt. fl. car.* 153. *S. adunca.* *Smith exot. b.* t. 1. t. 53. *variolaris.*
In open swamps, on sandy soil: North Carolina to Florida. ♀. June, July. v. v. Flowers yellow. The transverse spots on the back of the leaf distinguish this species very easily from the following, with which it has often been confounded.
3. *S. foliis strictis longissime infundibuliformibus, fauce patula, ala ventrali subnulla, appendice erecta basi coarctata ima parte lateribus retroflexis, mucrone subulato.* *Mich. fl. amer.* 1. p. 310. *Willd. sp. pl.* 2. p. 1150. *Icon. Bot. mag.* 780. *Bot. rep.* 381. *Catesb. car.* 2. t. 69. *Pluk. amalh.* t. 376. f. 5. *flava.*
In open swamps: Virginia to Florida. ♀. June, July. v. v. Flowers yellow. This is the tallest growing sort; the leaves are often two feet long.

psittacina.

4. *S. foliis brevibus superne coloratis venoso-reticulatis, ala ventrali sursum subcuneatim latescente, tubo sensim in appendicem recurvatum rotundatim fornicatam mucronatam desinente.* *Mich. fl. amer.* 1, p. 311.
S. rubra. *Walt. fl. car.* 152.
Icon. Pluk. amalh. t. 152, f. 3?
 In swamps of Georgia and Florida. ♀. June, July.
 v. v. Flowers on very long peduncles, purple; leaves small, very handsomely marked with purple veins.

rediviva.

437. LEWISIA. *Pursh in linn. trans.* v. 11.
 1. *Lewisia. l. c.*
 On the banks of Clark's river. ♀. July. v. s. in *Herb. Lewis.* Petals white; calyx elegantly red-veined, of a consistency like paper.
Radix fusiformis, ramosa, sanguinea. Folia radicalia linearia, subcarnosa, obtusiuscula. Scapus uniflorus? an biflorus. *Pedicellus* basi geniculatus. *Calyx coloratus, scariosus, 7—9-phylus, patens: foliolis ovatis, acutis, concavis, nervoso-venosis, interioribus angustioribus. Petala 14—18. alba, lanceolata, patentia, calyce vix duplo longiora. Filamenta 14—18. receptaculo petalis opposite inserta, filiformia, calyce breviora. Antheræ oblongæ, erectæ. Germen superum, ovatum, glabrum. Stylus filiformis, staminibus paulo longior, superne trifidus. Stigmata 3. bifida. Capsula oblonga, 3-locularis: loculis bispermis. Semina lenticularia, nitida, nigra.*

This elegant plant would be a very desirable addition to the ornamental perennials, since, if once introduced, it would be easily kept and propagated, as the following circumstance will clearly prove. The specimen with roots taken out of the Herbarium of M. Lewis, Esq. was planted by Mr. M'Mahon of Philadelphia, and vegetated for more than one year: but some accident happening to it, I had not the pleasure of seeing it in flower.

438. NYMPHÆA. *Gen. pl.* 886.*Castalia. Salisb. in annals of bot.* 2, p. 71.*odorata.*

1. *N. foliis orbiculato-cordatis integerrimis subemarginatis, lobis divaricatis, acumine obtuso, petalis calyce 4-*

4-p
 W
 N. alt
 Castal
 Icon.
 β. N. fol
 acu
 caly
 Icon.
 In pon
 ♀.
 riety
 ther

439. 1

Nympha

1. N. foli
 5-ph
 umb
 Nympl
 Nymph
 Icon. 1
 In lakes
 v. v.

2. N. folii
 calyc
 perica
 Nymph
 Nymph
 Icon. B
 In all p
 to Ca
 foreg
 its up

3. N. foliis
 stigma
 p. 29²
 Nympha
 Nympha
 Icon. B
 In ponds
 v. v.

nia.

reticulatis, ala
tubo sensim
nigritatam mu-
b. 311.

June, July.
purple; leaves
with veins.

11.

v. s. in
red-veined,

a radicaria
uniflorus?
Calyx co-
oliolis ova-
ioribus an-
, patentia,
18. recep-
tacle bre-
superum,
bus paulo
Capsula
mina len-

dition to
duced, it
following
en with
is, Esq.
bia, and
accident
ing it in

ginatis,
yce 4-

POLYANDRIA MONOGYNIA. *Nymphaea*.

369

4-phyllo æqualibus, stigmate radiis 16—20. erectis.—
Willd. sp. pl. 2. p. 1153.

N. alba. Mich. fl. amer. 1. p. 311. Walt. fl. car. 155.
Castalia pudica. Salisb. l. c. p. 72.
Icon. Bot. mag. 819. Bot. rep. 297.

β . N. foliis cordatis integrerrimis obtusis, lobis divaricatis
acutis, petiolis pedunculis foliisque subtus purpureis,
calycibus petalisque extus roseis.

Icon. Willd. hort. berol. 39?

In ponds and slow-flowing waters: Canada to Carolina.
 γ . July. v. v. Flowers white, fragrant. The va-
riety β . is probably a distinct species, which only fur-
ther observation can decide. It resembles *N. pygmæa*.

439. NUPHAR. *Smith prod. fl. græc. 1. p. 361.*

Nymphaea. Gen. pl. 886. Salisb. in annals of bot. 2.
p. 71.

1. N. foliis cordatis integrerrimis, lobis approximatis, calyce
5-phyllo, stigmate repando 14—20-radiato profunde
umbilicato.—*Ait. kew. ed. 2. v. 3. p. 295.*

Nymphaea lutea. Willd. sp. pl. 2. p. 1151.

Nymphaea umbilicalis. Salisb. l. c.

Icon. Engl. bot. 159.

In lakes on the Allegany mountains. γ . June, July.
v. v. Flowers yellow, as are all the following species.

2. N. foliis erectis cordatis integrerrimis, lobis divaricatis,
calyce 6-phyllo, stigmate leviter umbilicato 13-radiato,
pericarpio sulcato.—*Ait. kew. ed. 2. v. 3. p. 295.*

Nymphaea advena. Willd. sp. pl. 2. p. 1152.

Nymphaea arifolia. Salisb. l. c.

Icon. Bot. mag. 684. Willd. hort. berol. 38.

In all ponds and ditches, even in salt water: Canada
to Carolina. γ . July, Aug. v. v. Resembles the
foregoing species, but is distinguished at first sight by
its upright leaves.

3. N. foliis cordatis, lobis subapproximatis, calyce 5-phyllo,
stigmate inciso 8—12-radiato. *Ait. kew. ed. 2. v. 2.*
p. 295.

Nymphaea lutea β . Kalmiana. Mich. fl. amer. 1. p. 311.
Nymphaea lutea. Walt. fl. car. 154.

Icon. Bot. mag. 1243.

In ponds: Canada to Carolina; rare. γ . July, Aug.
v. v. Flowers very small in comparison with others.

rosea.

lutea.

advena.

Kalmiana.

- sagittifolia.* 4. N. foliis elongatis sagittato-cordatis obtusis, calyce 6-phyllo, petalis nullis, antheris subsessilibus.—*Salisb. l. c.*
Nymphaea sagittifolia. *Walt. fl. car.* 155.
Nymphaea longifolia. *Mich. fl. amer.* 1. p. 312.
 In slow-flowing waters: Carolina and Georgia. ♀. July, Aug. v. s. Flowers small.

440. CERATOPHYLLUM. *Gen. pl.* 493. *Willd. act. ac. berol.* 1798. p. 84.

- demersum.* 1. C. foliis verticillatis octonis dichotomo-bigeminis dorso dentatis spinulosis, floribus axillaribus solitariis sessilibus, fructibus 3-spinosis.—*Willd. sp. pl.* 4. p. 405.
Icon. Engl. bot. 947.
 In ditches and stagnant waters, floating: New York to Virginia. ♀. July, Aug. v. v. Flowers exceeding small.

II. DI-PENTAGYNIA.

441. CALLIGONUM. *Gen. pl.* 834.

- canescens.* 1. C. dioicum, pulverulento-tomentosum; foliis lanceolatis, floribus axillaribus glomeratis in apice ramulorum subspicatis, fructibus alatis, alis venosis cristato-dentatis.
 In the plains of the Missouri, near the Big-bend. h. July, Aug. v. s. in *Herb. Lewis.* Flowers exceeding small. Goats delight to feed upon this shrub.

442. DATISCA. *Gen. pl.* 1543.

- kirta.* 1. D. caule hirsuto, foliis pinnatis, foliolis basi confluentibus.—*Willd. sp. pl.* 4. p. 823.
 In Pennsylvania. *Kalm.* ♀. +. Flowers yellow, small, in terminal panicles. I never had an opportunity of seeing this plant myself.

443. MENISPERMUM. *Gen. pl.* 1544.

- canadense.* 1. M. foliis peltatis cordatis subrotundo-angulatis, racemis compositis.—*Willd. sp. pl.* 4. p. 824.

POL
 1. M. fo
 M. vi
 elth
 On the
 ame
 gree
 M.
 to A
 2. M. fol
 ract
 In Ken
 fl.;
 seede
 leave

1. D. pun
 lacini
 flor,
 pressi
 p. 31
 On the
 rivers
 about
 April,
 flower
 2. D. caule
 florib
 petalis
 p. 31
 D. caroli
 On the b
 Georg
 sippi.
 blue, 1
 some sp

3. D. caule
 ciniis la
 longitud
 p. 1230
 D. tridacty
 D. urceola
 Icon. *Mil*

6. *M. foliis lobatis.*
M. virginicum. Willd. sp. pl. 4. p. 824. excl. syn. Dill. elth. lobatum.
 On the banks of rivers and sides of fertile hills, climbing among other shrubs. ♂. July. v. v. Flowers greenish yellow; berries black, resembling grapes. *M. folio hederaceo*, *Dill. elth. t. 178. f. 219.* belongs to *M. carolinianum* or *Wendlandia*.
2. *M. foliis cordatis palmato-lobatis longissime pedicellatis, racemis simplicibus, floribus hexapetalis-dodecandris.* In Kentucky and Tennessee. ♀. June, July. v. v. s. fl.; v. s. in *Herb. Lyon.* Berries large, black, one-seeded. Stem climbing to the height of twenty feet; leaves large.

444. DELPHINIUM. Gen. pl. 924.-

1. *D. pumilum*; caule simplici, foliis palmato-multifidis, lacinii sublanceolatis obtusiusculis, fasciculo paucifloro, cornu recto corolla breviore, capsulis 3. compressis arcuatissimis subreflexo-patulis.—*Mich. fl. amer. 1. p. 314.*
- On the side of shady and fertile hills, on the banks of rivers: Virginia to Carolina; particularly plentiful about Harper's Ferry on the Potowmac, Virginia. ♀. April, May. v. v. Not above eight inches high; flowers beautiful blue.
2. *D. caule stricto simplici, foliis linearis-multipartitis, spica floribus breviter pedicellatis, cornu sursum arcuato, petalis villosissimis, capsulis ternis. Mich. fl. amer. 1. p. 314.*
- D. carolinianum. Walt. fl. car. 155.*
 On the borders of woods, in sandy soil: Carolina and Georgia. On the banks of the Missouri and Mississippi. ♀. May, June. v. s. Flowers beautiful sky-blue, large. W. Bartram mentions this very handsome species in his Travels.
3. *D. caule glabro subramoso, foliis tripartito-palmatis, lacinii lanceolatis subtrifidis, spicis strictis, cornu recto longitudine corollæ, capsulis ternis.—Willd. sp. pl. 2. p. 1230.*
- D. tridactylum. Mich. fl. amer. 1. p. 314.*
D. urceolatum. Jacq. collect. 1. p. 153.
 Icon. *Mill. ic. t. 250. f. 2. Jacq. ic. rar. 1. t. 101.*

POLYANDRIA DI-PENTAGYNIA. Delphinium.

- In rocky shady places, in the moutains of Virginia and Carolina. $\frac{3}{4}$. July. v. v. From two to four feet high.
- Consolida.* 4. D. pubescens; caule erecto ramoso, foliis linearis-multipartitis, cornu ascendentem corollam subæquante, nec-tario monophyllo, capsula solitaria.—*Willd. sp. pl. 2. p. 1226.*
Icon. *Fl. dan. 683.*
In fields and woods: Pennsylvania and Virginia. $\frac{3}{4}$. June, July. v. v. Common *Lark-spur*, probably introduced with the seeds of grain from Europe.

445. ACONITUM. *Gen. pl. 928.*

- uncinatum.* 1. A. floribus subpentagynis, foliis 3—5-lobis inciso-dentatis, corollarum galea longius extensa.—*Willd. sp. pl. 2. p. 1238.*
In swamps and on the side of rivulets, on the high mountains of Virginia and Carolina. $\frac{3}{4}$. June, July. v. v. Flowers large, of a fine blue and singular structure. On the foot of the Peaks of Otter and about the Sweet-springs another species occurs, with smaller flowers, and a climbing stem which sometimes attains the height of nine feet; but unfortunately I have no materials at present to give a correct description thereof.

446. AQUILEGIA. *Gen. pl. 934.*

- canadensis.* 1. A. cornibus rectis, staminibus exertis.—*Willd. sp. pl. 2. p. 1247.*
Icon. *Bot. mag. 246.* *Rob. ic. 296.*
In the crevices of rocks: Canada to Carolina. $\frac{3}{4}$. April, May. v. v. Flowers scarlet, mixed with yellow.

447. CIMICIFUGA. *Gen. pl. 993.*

- Serpentaria.* 1. C. foliis decompositis, foliolis ovato-oblongis incisis dentatis: dentibus mucronatis divaricatis, racemis virgatim paniculatis elongatis, floribus submonogynis, capsulis ovatis.
Actaea racemosa. Willd. sp. pl. 2. p. 1139.
Actaea monogyna. Walt. fl. car. 151.
Icon. *Dill. elth. t. 67. f. 78.* *Schkuhr handb. 139.* *Pluk. amalh. t. 383. f. 3.*

PO
In sha
Aug
gene
2. C. folii
tibus
5-gy
p. 12
Actæa
Icon. 1
On the
3. C. folii
serrati
C. ame
In shad
July.
foreg
4. C. foliis
culati
subgl
Actæa ra
In the l
lina.
some
great a

4

1. A. pusilli
longis
p. 77.
A. numm
In pine-l
Not at
as are a
2. A. multi
obtusis,
gynis,
A. multici
In sandy
 $\frac{1}{2}$. Jul
pale yel
3. A. erectu

In shady, stony woods: Canada to Florida. ♀. July, Aug. v. v. A tall stately plant; spikes long, white; generally known under the name of *Black Snake-root*.

2. *C. foliis decompositis, foliolis ovatis incisis dentatis, dentibus acutis, racemis paniculatis nutantibus, floribus 5-gynis, capsulis ovatis uncinatis*.—*Willd. sp. pl. 2. p. 1244.*

Actaea Cimicifuga. Sp. pl. 722.

Icon. *Lam. illustr. 487. Amoen. acad. 7. t. 6. f. 2.*
On the north-west coast. ♀. July, Aug. v. v.

3. *C. foliis decomposito-binternatis, foliolis cordatis lobatis serrato-dentatis, racemis virgatim-paniculatis elongatis, floribus 2—5-gynis, germinibus glabris*.

C. americana. Mich. fl. amer. 1. p. 316.

In shady woods, on high mountains of Carolina. ♀. July. v. v.; v. s. in *Herb. Lambert*. Resembles the foregoing very much in general habit.

4. *C. foliis simplicibus palmatis, floribus dichotome-paniculatis subcorymbosis polygynis, capsulis brevissimis subgloboso-capitatis*.—*Mich. fl. amer. 1. p. 316.*

Actaea racemosa. Herb. Walt.

In the beds of mountain rivulets in Virginia and Carolina. ♀. July, Aug. v. v. A very tall and handsome plant; the leaves very large, and the flowers in great abundance.

448. ASCYRUM. Gen. pl. 1225.

1. *A. pusillum, simplex; foliis ovalibus obtusis, pedicellis longis reflexis, floribus 1—2-gynis*.—*Mich. fl. amer. 2. p. 77.*

pumilum.

A. nummularifolium. Herb. Banks. mss.

In pine-barrens of Georgia. ♀. June—Aug. v. s. Not above three inches high; flowers small, yellow, as are all the following species.

2. *A. multicaule, diffusum; foliis sublanceolato-oblongis obtusis, corymbo terminali, floribus subsessilibus 2-gynis, caule subtereti*.—*Willd. sp. pl. 3. p. 1472.*

Crux Andreæ.

A. multicaule. Mich. fl. amer. 2. p. 77.

In sandy fields and woods: New Jersey to Carolina. ♀. July. v. v. Not above a foot high; flowers pale yellow; petals narrow.

3. *A. erectum, ramosum; ramis ancipitibus, foliis oblongo-*

hypericoides.

gis basi biglandulosis, floribus terminalibus solitariis breviter pedicellatis 3-gynis.—*Willd. sp. pl. 3. p. 1473.*
A. stans. Mich. fl. amer. 2. p. 77.
 In low overflowed open places, generally in sandy soil : New Jersey to Carolina. $\frac{1}{2}$. July, Aug. v. v. Flowers larger than the former species : it is from one to two feet high.

*amplexicaule. 4. A. erectum ; caule superne dichotome-paniculato, ramis
 ancipitibus, foliis ovato-oblongis amplexicaulibus, foliis calycinis exterioribus cordatis, floribus 3-4-gynis.—Mich. fl. amer. 2. p. 77.*
A. stans. Willd. sp. pl. 3. p. 1475.
Hypericum tetrapetalum. Lam. encycl. 4. p. 153.
 In low grounds and woods : Virginia to Florida. $\frac{1}{2}$. July, Aug. v. v. The flowers and leaves larger than any other sort.

449. HYPERICUM. *Gen. pl. 1224.** *Pentagyna.*

Kalmianum. 1. H. fruticosum, ramosissimum ; ramulis 4-gonis, foliis linear-lanceolatis, corymbis terminalibus 3—7-floris, laciniis calycinis lanceolatis obtusiusculis, capsulis longitudine calycis.—Willd. sp. pl. 3. p. 1438.
H. Bartramiaenum. Mill. dict. n. 10.

In Pennsylvania and Virginia. $\frac{1}{2}$. June, July. v. s. in *Herb. Lambert.* v. v. in *Hortis.* From three to five feet high : the flowers of the whole genus are yellow.

pyramida-tum. 2. H. herbaceum, erectum ; caule subtetragono ramoso, foliis sessilibus oblongis acutis glabris, floribus terminalibus, stylis staminibus brevioribus, calycinis foliis ovatis acutis.—Willd. sp. pl. 3. p. 1444.

H. amplexicaule. Lam. encycl. 4. p. 147.
Icon. Vent. malm. 118.

In Canada ? $\frac{1}{2}$. July, Aug. v. v. in *Hortis.* Flowers the size of *H. Ascyrum.*

ascyroides. 3. H. herbaceum, glabrum ; caule simplici 4-gono, foliis sessilibus oblongis acutis glabris, floribus terminalibus, stylis longitudine staminum, calycinis foliis ovato-lanceolatis.—Willd. sp. pl. 3. p. 1443.

H. macrocarpum. Mich. fl. amer. 2. p. 82.
 In Canada and the western parts of New York and Pennsylvania. $\frac{1}{2}$. July, Aug. v. v. Flowers the largest of all the species here enumerated.

4. *H. ramis
 natis b
 calycin
 æquanti
 Mich. j*

*H. elatun
 In shady
 July, A
 the leav
 than the*

5. *H. ramis
 revoluti
 nalibus
 acamina
 lis coad
 In South C
 v. v. t
 about tw
 one to th
 the petal
 of stami
 beautiful
 flower a*

6. *H. ramis a
 siusculis,
 floris, flo
 lisque lon
 latis.—W
 H. Kalmian
 In Virginia.
 The most
 gardens.*

*nada, as in
 southern F*

7. *H. ramis ala
 siusculis s
 nudiuscula
 lacinis la
 coadunatis
 p. 78.
 H. lævигатум
 H. ligustrinum*

VOL. II.

** *Trigynæ. Fruticosæ.*

4. *H. ramis ancipiibus, foliis ovali-oblongis obtusis mucronatis basi angustatis, floribus terminalibus subsolitariis, calycinis foliolis oblongis acutis staminibusque petalis æquantibus, stylis coadunatis longitudine staminum.*
Mich. fl. amer. 2. p. 81.
H. elatum. Ait. kew. 3. p. 104?
In shady rocky places in Kentucky and Tennessee. $\frac{1}{2}$. July, Aug. v. v. Resembles *H. Ascyrum* very much; the leaves of the calyx are broad, and sometimes longer than the petals.
5. *H. ramis ancipiibus, foliis ellipticis margine crispato-revolutis supra opacis subtus glaucis, floribus terminalibus subsolitariis sessilibus, calycinis foliolis ovatis acuminatis, petalis deflexis staminibus longioribus, stylis coadunatis longitudine staminum.*
In South Carolina and Georgia. *Lyon.* $\frac{1}{2}$. July, Aug. v. v. This elegant species grows to the height of about two feet or more. Every branchlet has from one to three large flowers, of a bright golden yellow; the petals turn downwards, and leave the large crown of stamens in an upright situation, which, with the beautiful red ovate germen in the centre, gives the flower a particularly pleasing appearance.
6. *H. ramis ancipiibus, foliis lanceolato-linearibus obtusiusculis, corymbis axillaribus et terminalibus paucifloris, floribus primordialibus sessilibus staminibus stylisque longitudine petalorum, calycinis foliolis lanceolatis.*—*Willd. sp. pl. 3. p. 1453.*
H. Kalmianum. Du Roi harbk 1. p. 310.
In Virginia. *Gronov.* $\frac{1}{2}$. June—Aug. v. v. in *Hortis.* The most common and most hardy species in the gardens. I rather suspect its native country to be Canada, as most of the shrubby kinds coming from the southern parts are extremely tender.
7. *H. ramis alato-4-gonis, foliis oblongo-lanceolatis obtusiusculis sessilibus, panicula pedunculata dichotoma nudiuscula, floribus solitariis subsessilibus, calycinis lacinias lanceolatis petalis duplo brevioribus, stylis coadunatis staminibus longioribus.*—*Mich. fl. amer. 2. p. 78.*
H. laevigatum. Ait. kew. 3. p. 106?
H. ligustrinum. Herb. Banks. MSS.

In Carolina. *Michaux.* $\text{h}.$ July, Aug. *v. v.* Flowers small. Willdenow in the Species Plantarum considers this to be an herbaceous plant, whereas it certainly is a shrub.

glaucum.

8. *H. ramis teretibus, foliis oblongis cordatis amplexicaulis obtusiusculis lœvigatis glaucis coriaceis, panicula divaricato-dichotoma foliosa, calycinis foliolis ovatis acutis petalis longioribus, staminibus petalis subæquantibus, stylis coadunatis.* — *Mich. fl. Amer.* 2. p. 78.

H. myrtifolium. *Lam. encycl.* 4. p. 180.

In Florida. *Michaux.* $\text{h}.$ July, Aug. *v. v.* Flowers large.

densiflorum.

9. *H. ramosissimum; ramulis subteretibus, foliis linearilanceolatis obtusiusculis basi attenuatis, paniculis terminalibus composite-dichotomis subfoliosis densifloris, floribus solitariis pedicellatis, calycinis laciniis brevissimis ovatis deciduis, staminibus stylisque coadunatis petalis brevioribus.*

H. rosmarinifolium. *Lam. encycl.* 4. p. 159?

On the dry ridges and savannahs of the Virginia mountains. $\text{h}.$ June, July. *v. v.* About two feet high; flowers middle-sized, in very abundant and close panicles.

galoides.

10. *H. ramulis tetragonis, foliis linearibus sessilibus margine revolutis, axillis fasciculato-foliolosis, paniculis terminalibus divaricato-dichotomis, petalis staminibus æquantibus calyce linearis vix longioribus, stylis coadunatis subexertis.* — *Lam. encycl.* 4. p. 161.

In sandy moist places, near rivulets: New Jersey to Carolina. $\text{h}.$ July—Sept. *v. v.* About two feet high; flowers the size of No. 9.

aspalathoides.

11. *H. ramis subteretibus dichotomis, foliis fasciculatis linearibus acutis striatis, floribus solitariis axillaris subsessilibus, staminibus petalis brevioribus, stylis coadunatis.* — *Willd. sp. pl.* 3. p. 1451.

H. fasciculatum. *Lam. encycl.* 4. p. 160.

In Carolina. *Fraser.* h. +. Flowers of middle size.

fasciculatum.

12. *H. ramulis teretibus, foliis quasi verticillatim-fasciculatis linearilanceolatis acutis basi attenuatis petiolatis margine revolutis, pedunculis in apice ramorum axillaris trifloris, flore medio sessili, calycibus foliiformibus, stylis coadunatis.* — *Willd. sp. pl.* 3. p. 1452. excl. *syn. Michaux.*

H. axill.

H. Cori

In pine

Flowe

13. *H. ram*

fascic

bus, |

floris,

H. fascic

In Georg

bles th

14. *H. erect*

obtusis

chiato-

longior

H. virgin

Icon. Fl.

In old fiel

June—

probabl

one of

very inj

blindne

quence.

15. *H. erectu*

ramoso,

tusis, c

lycinis |

p. 1457.

H. macula

H. puncta

In shady w

in the ra

Flowers

16. *H. erectu*

gono, fo

sis sessili

rymbosis

Willd. sp

H. quinque

H. mutilum

In overflow

Sept. v.

ericum.

v. v. Flowers
rum considers
it certainly is

amplexicauli-
ceis, panicula
foliolis ovatis
lis subæqua-
2. p. 78.

v. v. Flowers

foliis linearis-
paniculis ter-
ris densifloris,
ciniis brevis-
e coadunatis

ginia moun-
'o feet high ;
nd close pa-

bus margine
iculis termi-
minibus æ-
tylis coadu-

ersey to Ca-
o feet high ;

culatis line-
laribus sub-
tylis coadu-

niddle size.
-fasciculatis
iolatis mar-
mum axillari-
foliiformi-
1452. excl.

POLYANDRIA DI-PENTAGYNIA. Hypericum.

377

H. axillare. Lam. encycl. 4. p. 161.

H. Coris. Walt. fl. car. 190.

In pine woods of Georgia and Florida. 1 $\frac{1}{2}$. July. v. v.
Flowers the size of *H. perforatum*.

13. H. ramulis angulosis, foliis confertis quasi verticillatim fasciculatis filiformi-linearibus convexis obtusis sessilibus, pedunculis in apice ramulorum axillaribus 1—3-floris, calycibus foliiformibus, stylis coadunatis. *tenuifolium*.

H. fasciculatum. Mich. fl. amer. 1. p. 80.

In Georgia. Enslen. v. s. in Herb. Enslen. Resembles the foregoing species in many respects.

*** Trigynæ. Herbaceæ.

14. H. erectum, ramosum; caule ancipi, foliis oblongis obtusis pellucido-punctatis, panicula terminali brachiatofolia, petalis calyce acuto lanceolato duplo longioribus.—Willd. sp. pl. 3. p. 1460.

H. virginicum. Walt. fl. car. 189.

Icon. Fl. dan. 1043. Engl. bot. 295.

In old fields, meadows, and on dry hills; common. 1 $\frac{1}{2}$. June—Aug. v. v. Common St. John's Wort has probably been introduced from Europe, but become one of the most pernicious weeds, and is considered very injurious to horses; for, when they feed upon it, blindness and other diseases are said to be the consequence.

15. H. erectum, glabrum, nigro-punctatum; caule tereti ramoso, foliis amplexicaulibus oblongo-ovalibus obtusis, corymbis terminalibus brachiatis densifloris, cacycinis laciiniis lanceolatis acutis.—Willd. sp. pl. 3. p. 1457.

H. maculatum. Walt. fl. car. 189.

H. punctatum. Lam. encycl. 4. p. 164.

In shady woods: New England to Carolina; particularly in the range of the mountains. 1 $\frac{1}{2}$. June—Aug. v. v.
Flowers pale yellow, smaller than the foregoing one.

16. H. erectum, ramosissimum, glabrum; caule subtetragono, foliis ovato-oblongis subcordatis obtusis nervosis sessilibus, paniculis terminalibus dichotomo-corymbosis, petalis calyce lanceolato brevioribus.—Willd. sp. pl. 3. p. 1456.

H. quinquevium. Walt. fl. car. 190.

H. muticum. Willd. sp. pl. 3. p. 1471.

In overflowed places: Canada to Carolina. 1 $\frac{1}{2}$. June—Sept. v. v. Flowers small; size very variable.

- triplinerve.* 17. *H. erectum, subramosum ; caule subancipi, foliis linearibus patentissimis triplinerviis, floribus racemoso-paniculatis, corolla inaequali, calycibus acutis petalisque serrato-glandulosis.* — *Vent. hort. cels 58.*
On the banks of the Ohio. *Michaux.* 4. July, Aug.
+. Flowers the size of *H. perforatum*, pale yellow.
- dolabriiforme.* 18. *H. erectum, ramosum ; caule ancipi, foliis linear-lanceolatis patentibus obtusis, corymbo terminali dichotomo, calycibus inaequalibus lanceolatis, petalis dolabriiformibus.* — *Vent. hort. cels 45.*
On dry hills of Kentucky. *Michaux.* 4. June, July.
+. Flowers golden yellow.
- angulosum.* 19. *H. erectum ; caule tetragono, foliis oblongis acutis arce sessilibus, panicula terminali dichotoma, ramis divaricatis distanter alternifloris, calycibus subcampanulatis, lacinias lanceolatis acutis inferne carinatis longitudo fere corollæ, petalis dente unico laterali.* — *Willd. sp. pl. 3. p. 1454?* *Mich. fl. amer. 2. p. 78.*
H. denticulatum. *Walt. fl. car. 190.*
In bogs and cedar-swamps: New Jersey and Carolina.
4. June, July. v. v. About a foot high; flowers of a beautiful copper colour, of considerable size.
- sphaerocarpum.* 20. *H. erectum, glaberrimum ; foliis oblongis, panicula nuda dichotoma, dichotomis omnibus unifloris, stylis coadunatis, capsula globosa.* — *Mich. fl. amer. 2. p. 78.*
In Kentucky. *Michaux.* 4. +.
- canadense.* 21. *H. erectum, pusillum, parviflorum ; caule tetragono superne dichotomo, foliis sessilibus linearibus basi attenuatis, paniculæ ramis primariis oppositis, secundariis dichotomis, capsulis longis conoideis coloratis.* — *Willd. sp. pl. 8. p. 1455.*
In low gravelly soil; Canada to Carolina. 5. June—Aug. v. v. Flowers very small; capsules red.
- Sarothra.* 22. *H. pusillum, erectum, setaceo-ramosissimum ; quasi aphyllum ; foliis minutis appressis, floribus in ramulis solitariis alternis 5—10-andris, capsulis oblongis tilocularibus.* — *Mich. fl. amer. 2. p. 79.*
H. nudicaule. *Walt. fl. car. 190.*
Sarothra gentianoides. *Willd. sp. pl. 1. p. 1515.*
Icon. *Plak. mant. t. 342. f. 2.*
In dry sunny situations: New England to Carolina. 5. June—Aug. v. v. A singular little species; flowers very small; the number of stamens is very variable, according to soil and situation.

- POLYANDRIA DI-PENTAGYNIA. Hypericum.
23. *H. erectum*
foliis
— *M.*
H. pilosum
H. ferrugineum
Ascyrum
Icon. P
Ih. Lowe
24. *H. procumbens*
mitate
lycinis
corolla
2. p. 8
On dry sand
calyx i
1. *E. foliis* *tusissimis*
et term
9—12.
Hypericum
Hypericum
Icon. Bot
- β. *E. foliis* *coloratis*
bus elon
Hypericum
In bogs at
July—S
ers yellow
2. *E. foliis* *se*
quam ad
Hypericum
In Carolina
of the I
belonging
in my H
species al
and acute
leaves, sc
and acute
3. *E. foliis* *pe*

ericum.

ipiti, foliis linearibus racemoso-acutis petalis 58.

♀. July, Aug.
pale yellow.

foliis lineariterminali dilatatis, petalis

. June, July.

jis acutis arcte
ramis divaricata
abcampanulatis longitu-
rali.—Willd.
p. 78.

nd Carolina.
igh ; flowers
ible size.

is, panicula
ifloris, stylis
ter. 2. p. 78.

e tetragono-
bus basi at-
is, secunda-
coloratis.—

○. June—
s red.
am ; quasi
; in ramulis
longis 1-lo-

5.

rolina. ○.
s ; flowers
y variable,

POLYANDRIA DI-PENTAGYNIA. Hypericum.

379

23. H. erectum, simplicissimum, lanulosum ; caule tereti, foliis oblongis adpressis, floribus paucis terminalibus.—*Mich. fl. amer.* 2. p. 80.
H. pilosum, *Walt. fl. car.* 190.
H. ferrugineum, *Herb. Banks.* mss.
Ascyrum villosum. Willd. sp. pl. 3. p. 1474.
Icon. *Pluk. alm.* t. 245, f. 6.
In Lower Carolina and Georgia. ○. July, Aug. v. s.
24. H. procumbens, glaberrimum ; ramis ancipitibus summitate paucifloris, foliis linearibus enerviis patulis, calycinis foliolis oblongo-lanceolatis acutis, staminibus corolla brevioribus, stylis coadunatis.—*Mich. fl. amer.* 2. p. 81. *Willd. sp. pl.* 3. p. 1450?
On dry sunny hills of Kentucky. *Michaux.* +. The calyx is as large as the corolla.

450. ELODEA. *Adanson.*

Hyperici sp. *Linn.*

1. E. foliis sessilibus amplexicaulibus cordato-oblongis obtusissimis, fasciculis pedunculatis panicifloris axillaribus et terminalibus, corollis subcampanulatis, staminibus 9—12. levissime basi coalitis.
Hypericum virginicum. Willd. sp. pl. 3. p. 1455.
Hypericum campanulatum. Walt. fl. car. 191.
Icon. *Bot. rep.* 552.
β. E. foliis cordato-ovatis emarginatis, pedunculis axillari- bus elongatis. emarginata.
Hypericum emarginatum. Lam. encycl. 4. p. 154.
In bogs and wet meadows : Canada to Carolina. ♀.
July—Sept. v. v. β. v. s. in *Herb. Lambert.* Flowers yellowish-red, very handsome.
2. E. foliis sessilibus, corollis tubulosis, staminibus plus quam ad medium connatis.
Hypericum tubulosum. Walt. fl. car. 191.
In Carolina. *Walter.* +. I observed in the vicinity of the Dismal-swamp in North Carolina, a species belonging to this genus, of which I have no specimen in my Herbarium, and which probably may be the species alluded to by Walter. The leaves are elliptic and acute ; the flowers nearly sessile in the axils of the leaves, solitary or by two or three ; the petals are long and acute, pale red ; capsules long, prismatic.
3. E. foliis petiolatis oblongo-ovalibus rotundato-obtusis, petiolata.

floribus oppositis axillaribus subsessilibus subtornis,
staminibus ad medium usque connatis, capsulis oblongis.

Hypericum petiolatum. *Walt. fl. car.* 191.

Hypericum axillare. *Mich. fl. amer.* 2. p. 81.

On banks of lakes and rotten wood: Virginia and Carolina. $\frac{1}{2}$. July, Aug. v. v.

III. POLYGYNIA.

451. ILLICIUM. Gen. pl. 940.

floridanum.

1. *I. foliis acuminatis, petalis numerosis oblongis linearibusque*.—*Willd. sp. pl.* 2. p. 1254.

Icon. Bot. mag. 439.

On the banks of the Mississippi, in West Florida. $\frac{1}{2}$. May, June. v. v. Flowers dark purple, large; leaves of a very aromatic anise-scent, when bruised.

parviflorum.

2. *I. foliis obtusis, petalis sex calyceque rotundatis concavis*.—*Mich. fl. amer.* 1. p. 326.

I. anisatum. *Batr. trav.* 24.

Icon. Vent. hort. cels 22.

In East Florida, on the banks of the river St. John. *Michaux*. $\frac{1}{2}$. May, June. v. v. in *Hortis*. Flowers small, yellow.

452. MAGNOLIA. Gen. pl. 942.

grandiflora.

1. *M. foliis persistentibus ovalibus crassis coriaceis, petalis dilatato-ovobatis abrupte in unguem angustatis*.—*Willd. sp. pl.* 2. p. 1255.

α. *M. foliis oblongo-ellipticis subtus ferrugineis, floribus subcontractis*. *Ait. kew.* 2. p. 251.

Icon. Bot. rep. 518. *Trew ehret.* 33.

β. *M. foliis obovato-oblongis, floribus expansis*. *Ait. l. c.*

γ. *M. foliis oblongo-lanceolatis apice flexis concoloribus, floribus subcontractis*. *Ait. l. c.*

Icon. Mill. ic. 172. *Catesb. car.* 61. *Mich. arb.* t. 1.

In the forests of South Carolina and Georgia, reaching west to the Mississippi. $\frac{1}{2}$. May—Aug. v. v. This stately tree, with its elegant foliage and large white flowers, is the prince of the forests of North America. The varieties here mentioned might easily be considered as good species, if there were not so many intermediate ones, which connect them too closely to-

gether
Laure

2. *M. foliis attenuata*

α. *M. foliis kew.* 2

Icon. D

Trew e

β. *M. foliis l. c.*

In swamp

β. Geo

small tr

β. is a

leaves

consider

by the

Swamp

3. *M. ramis*

longe su

subtus

{ *amer.* 1.

Icon. Mic

In the deep

s. fl.

leaves ar

red, whi

the most

4. *M. foliis a*

juniorib

exteriorib

M. umbrell

Icon. Cates

In the moun

July. v.

of *M. gr*

known by

tains they

5. *M. foliis ova*

tal obova

Icon. Catesb

In high moun

Carolina.

mixed wit

gether. In Carolina it is known by the name of *Big Laurel*.

2. M. foliis ovalibus subtus glaucis, petalis obovatis basi attenuatis.—*Willd. sp. pl. 2. p. 1256.*
- α. M. foliis deciduis ovali-oblongis obtusiusculis.—*Ait. kew. 2. p. 251.*
Icon. *Dill. elth. t. 168. f. 205. Catesb. car. 1. t. 39. Trew ehret. q. Mich. arb. t. 2.*
- β. M. foliis perennantibus ellipticis utrinque acutis.—*Ait. l. c.*
In swamps covered with wood: New Jersey to Carolina.
β. Georgia and Florida. $\frac{1}{2}$. May—July. v. v. A small tree; flowers white, very fragrant. The variety β. is a taller tree, the branches more upright, and the leaves narrow and long; it might with propriety be considered a distinct species. This species is known by the names of *Swamp Sassafras*, *Sweet Bay*, *Swamp Laurel*, and *Beaver-wood*.
3. M. ramis medullosois fragilibus, foliis amplissimis ob-
longe subcuneato-ovalibus basi sinuata subauriculatis
subtus glaucis, petalis 6. ovatis obtusis.—*Mich. fl. amer. 1. p. 327.*
Icon. *Mich. arb. t. 7.*
In the deep forests of Tennessee. $\frac{1}{2}$. May—July. v. v.
s. fl. This stately small tree, with its extremely large leaves and white flowers, tinged at the bottom with red, which are larger than *M. grandiflora*, is one of the most ornamental trees America produces.
4. M. foliis amplis oblonge subcuneato-ovalibus acutis,
junioribus holosericeis, petalis 9. ellipticis acutis,
exterioribus reflexis.—*Willd. sp. pl. 2. p. 1258.*
M. umbrella. *Lam. encycl. 3. p. 644.*
Icon. *Catesb. car. 2. t. 80. Mich. arb. t. 5.*
In the mountains of Virginia and Carolina. $\frac{1}{2}$. June,
July. v. v. Leaves large; flowers white, the size of *M. grandiflora*, and very fragrant. It is generally known by the name of *Umbrella-tree*; in the mountains they call it *Elk-wood*.
5. M. foliis ovalibus acuminatis subtus pubescentibus, pe-
talibus obovatis obtusiusculis.—*Willd. sp. pl. 2. p. 1257.*
Icon. *Catesb. car. 3. t. 15. Mich. arb. t. 3.*
In high mountain-valleys, in fertile soil: Pennsylvania to Carolina. $\frac{1}{2}$. June, July. v. v. Flowers a dim yellow mixed with a faint blue, not much larger than those

*glaucia.**latifolia.**longifolia.**tripetala.**acuminata.*

POLYANDRIA POLYGYNIA. Magnolia.

of *M. glauca*. It is generally known by the name of *Cucumber-tree*, and its fruits are used as a wholesome bitter among the inhabitants.

cordata.

6. *M. foliis cordatis subtus subtomentosis, petalis lanceo-lato-oblongis acutis.* — *Mich. fl. amer.* 1. p. 328.

Icon. *Mich. arb.* t. 4.

On dry ridges of mountains in Upper Carolina and Georgia. $\frac{1}{2}$. v. v. s. fl. Flowers yellow.

auriculata.

7. *M. foliis amplis obovato-lanceolatis acutis subtus glancis basi cordatis auriculatis, lobis approximatis, petalis ovatis acutis sub-unguiculatis.* — *Willd. sp. pl.* 2. p. 1258.

M. auricularis. *Parad. lond.* 43.

M. Fraseri. *Walt. fl. car.* 159.

Icon. *Bot. mag.* 1206. *Bot. rep.* 573. *Mich. arb.* t. 6.

In the Allegany mountains, from the head-waters of the Susquehanna to Carolina. $\frac{1}{2}$. April, May. v. v. A beautiful tree; flowers yellowish white, large. The bark of this and some of the foregoing species is esteemed a valuable medicine, particularly in intermitting fevers; from which circumstance it is known in some places by the name of *Indian Physic*.

pyramidata.

8. *M. foliis rhomboideo-obovalibus abrupte acutis concoloribus basi subcordatis auriculatis, lobis divaricatis, petalis lanceolatis sensim acutis.* — *Bartr. trav.* p.

M. auriculata. *Mich. fl. amer.* 1. p. 328.

In the western parts of Carolina and Georgia. $\frac{1}{2}$. April, May. v. v. This species has generally been confounded with the foregoing one, from which it not only differs in the distinctions given above, but in general habit: the tree is of a more upright pyramidal growth, and the leaves not one fourth the size of *M. auriculata*.

453. LIRIODENDRON Gen. pl. 941.

tulipifera.

1. *L. foliis absciso-truncatis 4-lobatis, calyce triphylo.* — *Willd. sp. pl.* 2. p. 1254.

acutiloba.

- a. *L. lobis acutis acuminatisque.* *Mich. fl. amer.* 1. p. 326. Icon. *Bot. mag.* 275. *Schmidt arb.* 48. *Catesb. car.* 1. t. 48. *Mich. arb.* t.

obtusiloba.

- β . *L. lobis rotundato-obtusissimis.* *Mich. l. c.* In fertile ground: Canada to Florida. β . in Pennsylvania. $\frac{1}{2}$. June, July. v. v. A highly ornamental and use-

ful in
green
or W

454

Ore

1. *P. foliis exterie*
Pers. i

Anona tr

Oridoc

Icon. *Ca*

arb. 1.

On the o

rida. 1

ers dar

2. *P. foliis e*
pubesca
tis subc

Orchidoc

In shady v

gia. $\frac{1}{2}$

times :

flowers

3. *P. foliis el*
exterior
longiori

Orchidoc

Anona pyg

Icon. *Bart*

In sandy fi

the size

above a 1

4. *P. foliis cur*
pubescen
terior
Orchidocarp

Anona obov

Icon. *Bart*

In sandy w

$\frac{1}{2}$. May,

in propor

ful timber tree; flowers yellow and red, mixed with green. Generally known by the name of *Tulip-tree*, or *White and Yellow Poplar*.

454. PORCELIA. *Prodr. fl. peruv. p. 84.*

Orchidocarpum. Mich. fl. amer. 1. p. 329.

1. *P. foliis glabriusculis oblonge cuneato-obovatis, petalis exterioribus orbiculatis, fructibus magnis carnosis.*—
Pers. syn. 2. p. 95.

Anona triloba. Willd. sp. pl. 2. p. 1267.

Orchidocarpum arietinum. Mich. fl. amer. 1. p. 329.

Icon. Catesb. car. 2. t. 83. Trew ehret. 5. Duham. arb. 1. t. 19.

On the overflowed banks of rivers: Pennsylvania to Florida. $\frac{1}{2}$. March, April. v. v. A small tree; flowers dark brown; fruits large, eatable.

2. *P. foliis cuneato-obovatis basi acutis ramulisque rufopubescentibus, petalis omnibus ovalibus, fructibus ova-tis subcarnosis.*—*Pers. syn. 2. p. 95.*

Orchidocarpum parviflorum. Mich. fl. amer. 1. p. 329.

In shady woods, near rivers and lakes: Virginia to Georgia. $\frac{1}{2}$. April, May. v. v. A low shrub, sometimes not above two feet high, when in full fruit; flowers small, dark purple.

3. *P. foliis elongato-lanceolatis cuneatisve lavigatis, petalis exterioribus majoribus obovato-oblongis, interioribus longioribus.*—*Pers. syn. 2. p. 95.*

Orchidocarpum pygmæum. Mich. fl. amer. 1. p. 330.

Anona pygmæa. Willd. sp. pl. 2. p. 1268.

Icon. Bartr. trav. t. 1.

In sandy fields: Georgia and Florida. $\frac{1}{2}$. v. v. Flowers the size of *Anona squamosa*; the whole shrub not above a foot high.

4. *P. foliis cuneato-ovovalibus basi obtusis ramulisque rufopubescentibus, petalis exterioribus majoribus obovatis, interioribus oblongis.*—*Pers. syn. 2. p. 95.*

Orchidocarpum grandiflorum. Mich. fl. amer. 1. p. 330.

Anona obovata. Willd. sp. pl. 2. p. 1269.

Icon. Bartr. trav. t. 2.

In sandy woods, in shady places: Georgia and Florida. $\frac{1}{2}$. May. v. s. A small shrub: flowers very large in proportion, white.

triloba.

parviflora.

pygmæa.

grandiflora.

gnolia.

by the name of
as a wholesome

, petalis lanceo-
1. p. 328.

r Carolina and
yellow.

is subtus glancis
oximatis, petalis
llid. sp. pl. 2.

Mich. arb. t. 6.
d-waters of the
, May. v. v.
ite, large. The
; species is es-
in intermitting
rown in some

acuteis concavo-
is divaricatis,
. trav. p.

jia. $\frac{1}{2}$. April,
ly been con-
which it not
e, but in ge-
ight pyramidal
the size of

941.

triphylo.—

r. 1. p. 326.
atesb. car. 1.

Pensylvania.
ital and use-

455. ATRAGENE. *Gen. pl. 949. Sims in bot. mag. 887.**americana.*

1. A. foliis quaternis ternatis, foliolis cordatis subintegerimis, nectaris acutis. *Sims l. c. cum icono.*
In shady places, on the side of rivulets, climbing and creeping among loose rocks : New York and Pennsylvania, near the foot of the Blue-mountains. ♂. May, June. v. v. Flowers large, purple.

456. CLEMATIS. *Gen. pl. 960.**virginica.*

1. C. scandens ; foliis ternatis, foliolis ovatis subcordatis inciso-dentatis lobatisque, corymbis dichotomis paucifloris, petalis staminibus longioribus, floribus dioicis. —*Willd. sp. pl. 2. p. 1290.*
In hedge-rows and among small shrubberies on the side of woods : Canada to Florida. ♂. June—Aug. v. v. Flowers small, white, and fragrant.

cordata.

2. C. scandens ; foliis pinnatis bijugis, foliolis cordatis lobatis acuminatis inciso-dentatis, dentibus mucronatis, racemis compositis elongatis multifloris, staminibus petalis subaequantibus, floribus dioicis.
On high mountains : Virginia. ♂. July. v. v. Flowers resembling the former species. In general appearance it approaches near to *C. Vitalba*.

holosericea.

3. C. scandens, tota holosericeo-pubescentes ; foliis ternatis oblongo-lanceolatis integris utrinque pubescentibus, corymbis trichotomis paucifloris, petalis linearibus staminibus longioribus, floribus dioicis.
In Carolina. ♂. v. s. in *Herb. Walter*. The flowers small, white ; tails of the seeds very long, feathered.

Walteri.

4. C. scandens ; foliis pinnatis cirrhosis trijugis, foliolis divaricatis petiolatis linear-lanceolatis acutis integerimis subtus glaucis, floribus solitariis, petalis ellipticis staminibus duplo longioribus.

In Carolina. ♂. v. s. in *Herb. Walter*. Flowers white.

erispa.

5. C. scandens ; foliis pinnatis ternatisque, foliolis divaricatis ovato-lanceolatis acutis trilobise integerimis, floribus solitariis, corollis campanulatis acuminatis revolutis margine undulatis, aristis seminum subulatis nudis.—*Willd. sp. pl. 2. p. 1289.*

Icon. Dill. elth. t. 73. f. 84.

In hedges and among bushes on the side of rivers : Vir-

ginia at
large,]

6. C. scandens
tis glab
nalibus
subcori
plumos
C. Viorna
In Virgin
Flowers

7. C. scandens
que obt
ceis utri
tal is sub
car. 156

C. rosea.
In Georgia
Walter.
pale pur

8. C. scandens
que, foli
integeri
crassis ac
Icon. Bot.
In hedges a
—Aug.

9. C. erecta, s
cisis, lac
multifidi
campanu
apice refl
On the pla
Herb. Le
close coat
it very m
all the di
belong to
Anemone.

10. C. erecta, s
libus integ
cano-seric
nuo.—*Mi*
C. ochroleuc
Icon. Bot. n

- in bot. mag.* 887.
atis subintegerri-
cione.
t, climbing and
ork and Pensyl-
tains. h. May,
30.
tis subcordatis
hotomis pauci-
loribus dioicis.
- ries on the side*
—Aug. v. v.
- is cordatis lo-*
is mucronatis,
s, staminibus
- v. v. Flowers*
al appearance
- foliis ternatis*
ibescentibus,
nearibus sta-
- The flowers
feathered.
- gis, foliolis
is integeri-
lis ellipticis
- wers white.
lis divarica-
rimis, flo-
natis revo-
bulatis nu-
- vers: Vir-
- ginia and Carolina. h. July, Aug. v. v. Flowers large, purple.
6. C. scandens; foliis pinnatis, foliis ovatis utrinque acutis glabris simplicibus pedicellatis, pedunculis terminalibus solitariis, corollis cernuis cylindricis, petalis subcoriaceis undulatis conniventibus, aristis seminum plumosis.—*Sims in bot. mag.* 1160.
- C. Viorna. *Bot. rep.* 71.
- In Virginia and Carolina. h. June—Aug. v. v. Flowers large, of a fine pale purplish blue.
7. C. scandens; foliis pinnatis 4-jugis, foliolis ovatis utrinque obtusis omnibus integris petiolatisque membranaceis utrinque reticulato-venosis, floribus solitariis, petalis subcoriaceis, aristis seminum plumosis.—*Walt. fl. car.* 156. *Mich. fl. amer.* 1, p. 318.
- C. rosea. *Abbot insect. t.*
- In Georgia and Carolina. h. June, July. v. s. in *Herb. Walter.* Flowers resembling the following species, pale purplish-red.
8. C. scandens; foliis divaricato-compositis decompositaque, foliolis ovali-lanceolatis utrinque acutis trifidisque integerrimis, floribus solitariis campanulatis, petalis crassis acuminatis.—*Willd. sp. pl.* 2, p. 1288.
- Icon. *Bot. rep.* 71. *Dill. elth. t.* 118, *f.* 144.
- In hedges and copses: Virginia and Carolina. h. June—Aug. v. v. Flowers dark blue, large.
9. C. erecta, simplex; foliis hirsutissimis bipinnatifidis incisis, laciniis linearibus acutis, caulinis composite multifidis, pedunculo terminali solitario, flore erecto campanulato, petalis 4. erectis ovatis obtusiusculis apice reflexis.
- On the plains of Columbia river. h. May. v. s. in *Herb. Lewis.* The whole plant is covered with a close coat of long hairs like *Anemone Pulsatilla*, which it very much resembles in several respects. I consider all the division of *Anemones* with caudated seeds to belong to this genus, or at least to one separate from *Anemone*.
10. C. erecta, simplex, pubescens; foliis simplicibus ovalibus integerrimis, junioribus calycibusque extus incano-sericeis, pedunculo terminali solitario, flore cernuo.—*Mich. fl. amer.* 1, p. 319.
- C. ochroleuca. *Willd. sp. pl.* 2, p. 1294.
- Icon. *Bot. mag.* 1175.
- cylindrica.*
- reticulata.*
- Viorna.*
- hirsutissima.*
- sericea.*

On the banks of rivers : Pennsylvania and Virginia ; rare.

\mathcal{U} . May—July. *v. v.* About a foot high ; flowers inside yellow.

457. ANEMONE. *Gen. pl. 948.*

nemorosa.

1. A. caule unifloro, foliis caulinis ternis ternatis, foliolis cuneatis inciso-lobatis dentatis acutis, corollis 5-6-petalis, seminibus ovatis stylo brevi uncinatis.—*Willd. sp. pl. 2. p. 1281.*

Icon. *Fl. dan. 549. Engl. bot. 355.*

- quinquefolia.* β . A. foliis laterilibus profunde bipartitis. *Willd. sp. pl. 2. p. 1281.*

Icon. *Pluk. alm. t. 106. f. 3.*

In shady woods, round the roots of trees : Canada to Carolina. \mathcal{U} . March, April. *v. v.* Flowers white, with a tinge of red on the under side ; sometimes nearly double. Var. β . can never be considered as a species, as there are intermediate varieties between it and the original species.

lancifolia.

2. A. caule unifloro, foliis caulinis ternis ternatis, foliolis lanceolatis crenato-dentatis, corollis 5-petalis, seminibus ovatis stylo brevi uncinatis.

A. trifolia. Willd. sp. pl. 2. p. 1281 ?

On high mountains, in boggy soil : Pennsylvania and Virginia. \mathcal{U} . May—July. *v. v.* Resembles the foregoing species ; flowers larger, and clear white.

cuneifolia.

3. A. caule unifloro involucrato, foliis ternatis, foliolis sessilibus cuneatis apice truncato-crenatis, involucro 3-phyllo, corollis 5-petalis, seminibus capitato-globosis lanatis acuminatis.—*Juss. in annal. du mus. 3. p. 248. t. 21. f. 1.*

*A. parviflora. Mich. fl. amer. 1. p. 319.
A. tenella. Herb. Banks.*

On the banks of rivulets, near Hudson's Bay and Labrador. \mathcal{U} . March—May. *v. s. in Herb. Dickson.* Flowers white, the size of No. 1.

tenella.

4. A. caule unifloro involucrato, foliis subbiternatis, foliolis 3-fidis seu 3-dentatis acutis involucro 3-partito, lacinias incisis, petalis 12—15-linearibus, seminibus lanatis mucronatis, radice tuberosa.

A. caroliniana. Walt. fl. car. 157.

On the banks of the Missouri. *M. Lewis.* In Carolina. *Walter.* \mathcal{U} . May. *v. s. in Herb. Lewis.* Flowers

PC

purple,
lilate ;

5. A. caule
corolla
Thalictrum
In Carolin

any inf
Herbari
genus ti

6. A. florib
ternatis,
phyllo,
flora, s
Willd.
Thalictrum
Icon. *Bo*
 $\mathfrak{C}.$ b.

β . A. involuc
In woods,
—May.
variety i
of figure
elegant g

7. A. floribus
tifido-pa
sessilibus
ero longi
In Canada
v. s. Fl

8. A. caule di
sessilibus
lobis acu
tariis unit
lato-roste
A. aconitifo
In meadows
Pensylvan
white ; ai

9. A. caule di
plexicauli
inciso-seri
capitulo g
sp. pl. 2. f
A. irregulari

purple, small; the whole plant very slender and delicate; roots tuberous.

5. A. caule unifloro nudo, foliis palmatis longius petiolatis, corolla 5-petala, radice tuberosa.
Thalictrum carolinianum: *Walt. fl. car.* 157.

In Carolina. *Walter*. 2. +. I was not able to find any information respecting this plant in the Walterian Herbarium, but consider it to belong rather to this genus than to *Thalictrum*.

6. A. floribus umbellatis involucratis, foliis radicalibus bitemnatis, foliolis subcordatis 3-dentatis, involucre 6-phyllo, foliolis petiolatis conformibus, umbella pauciflora, seminibus nudis striatis, radice tuberosa.—*Willd. sp. pl.* 2. p. 1284.

Thalictrum anemonoides. *Mich. fl. amer.* 1. p. 322.
Icon. *Bot. mag.* 866. *Annales du mus.* 3. t. 21. f. 2. a.
& b. *Pluk. aln.* t. 106. f. 4. *Willd. hort. berol.* 44.

- β. A. involucro unifloro. *Gronov. virg.*
In woods, frequent: Canada to Carolina. 2. March—May. v. v. Flowers white. A beautiful double variety is cultivated in the gardens, which in delicacy of figure and colour exceeds any other species of this elegant genus.

7. A. floribus umbellatis involucratis, foliis radicalibus multifido-palmatis incisis, lacinias linearibus, involucris sessilibus incisis, umbella pauciflora, pedicellis involucro longioribus.—*Willd. sp. pl.* 2. p. 1283.

In Canada and on the north-west coast. 2. April, May.
v. s. Flowers white.

8. A. caule dichotomo, foliis radicalibus palmatis, caulinis pensylvanica.
sessilibus amplexicaulibus 3-fidis: infimis ternatis: lobis acute lanceolatis inciso-serratis, pedunculis solitariis unifloris, seminibus capitulo globoso nudis subulato-rostellatis.—*Willd. sp. pl.* 2. p. 1280.

A. aconitifolia. *Mich. fl. amer.* 1. p. 320.
In meadows and on the borders of woods: Canada to Pennsylvania. 2. June, July. v. v. Flowers large, white; anthers yellow.

9. A. caule dichotomo, foliis caulinis omnibus oppositis amplexicaulibus trifidis, lobis cuneato-lanceolatis superne inciso-serratis, pedunculis solitariis unifloris, seminibus capitulo globoso nudis recurvato-mucronatis. *Willd. sp. pl.* 2. p. 1280.

A. irregularis, *Lam. encycl.* 1. p. 167.

Walteri.

uniflora.

dichotoma.

- Icon. *Linn. fil. dec. 2. t. 15.*
In wet woods and natural meadows: Canada and the western parts of New York. $\frac{1}{4}$. May, June. *v. v.* Flowers white, with a tinge of red on the under side, smaller than the foregoing species.
- virginiana.* 10. A. caule dichotomo, foliis ternis ternatis, superioribus oppositis, foliolis inciso-lobatis serratisque acutis, pedunculis solitariis unifloris elongatis, seminibus capitulo oblongo lanatis mucronatis.—*Willd. sp. pl. 2. p. 1279.*
- Icon. *Herm. parad. t. 18.*
In woods, on the side of dry sandy hills: Canada to Carolina. $\frac{1}{4}$. May, June. *v. v.* Flowers small, greenish-yellow.
- Cornuti.* 1. T. foliis supradecompositis, foliolis ovatis trifidis, paniculis terminalibus.—*Willd. sp. pl. 2. p. 1296.*
Icon. *Corn. canad. 186.*
On the banks of rivers and in wet meadows: Canada to New England. $\frac{1}{4}$. June, July. *v. v.* From two to three feet high; flowers small and greenish-yellow.
- dioicum.* 2. T. laevigatum; foliis supradecompositis, foliolis subrotundis cordatis obtuse-lobatis subtus glaucis laevis, paniculis axillaribus filiformibus folio brevioribus, floribus parvis in umbellis paucifloris.—*Willd. sp. pl. 2. p. 1296.*
- T. laevigatum. *Mich. fl. amer. 1. p. 322.*
In shady woods and on the banks of rivers: Canada to Virginia. $\frac{1}{4}$. May—July. *v. v.* Not above a foot high; flowers white.
- rugosum.* 3. T. caule striato, foliis supradecompositis, foliolis ovatis lanceolatisque rugosis venosis obtuse-lobatis, paniculis amplis terminalibus.—*Willd. sp. pl. 2. p. 1298.*
On the banks of rivers and in swamps: Pennsylvania to Carolina. $\frac{1}{4}$. June—Aug. *v. v.* Very tall, sometimes above five feet high; flowers white.
- pubescens.* 4. T. tenui tomento pubescens; foliis supradecompositis, foliolis ovatis subcordatis cuneatisque apice 3-lobis, supra subrugosis, subtus subtomentosis, paniculis terminalibus, pedicellis subumbellatis divaricatis, floribus polygamis.
- T. polygamum. *Muhl. mss.*

PO.

On the b
Virgini
flowers5. T. caule
subrotu
ribus c
p. 1295
On dry s
May, J
purple.6. T. foliis s
—Will
In Carolin
those of

45

1. H. caule t
emargin
nali soli
Icon. *Mil*
In shady w
da to Ca
 $\frac{1}{4}$. Apr
berry re

46

Hyd

1. B. foliis ce
solitariis
Hydropelti
Icon. *Bot.*
In lakes an
rolina.
ter; the
dark pur
surface
gelatinou

4

1. C. caule er
dato-ovat
petalis ell

- On the banks of ditches and rivulets: Pennsylvania and Virginia. ♀. June—Aug. v. v. A tall species; flowers white.
5. T. caule foliis duplo altiore, foliis compositis, foliolis *purpurascens*.
subrotundis 3-fidis incisis, paniculis nudiusculis, floribus cernuis, staminibus coloratis.—*Willd. sp. pl. 2.*
p. 1299.
- On dry sunny hills: Pennsylvania and Virginia. ♀.
May, June. v. v. A small plant; stem and filaments purple.
6. T. foliis simplicibus 5-lobis serratis, floribus corymbosis.
—*Willd. enum. 585.*
- In Carolina, *Willdenow.* ♀. +. Flowers resembling those of *T. speciosum.*

459. HYDRASTIS. *Gen. pl. 958.*

1. H. caule superne opposite diphylo, foliis petiolatis basi emarginatis palmatis serratis incisis, pedunculo terminali solitario unifloro.—*Willd. sp. pl. 2. p. 1340.*
Icon. *Mill. ic. t. 285.*
- In shady woods, on fertile soil and among rocks: Canada to Carolina, principally in the Allegany mountains. ♀. April, May. v. v. Flowers pale rose-coloured; berry red. The roots are yellow, and afford a fine dye.

460. BRASENIA. *Gen. pl. 938.*Hydropeltis. *Mich. fl. amer. 1. p. 323.*

1. B. foliis centro peltatis ovalibus integerrimis, pedunculis solitariis unifloris.
Hydropeltis purpurea. *Mich. fl. amer. 1. p. 324. t. 29.*
Icon. *Bot. mag. 1147.* *Pluk. alm. t. 349. f. 3.*
- In lakes and slow-flowing waters: New Jersey to Carolina. ♀. June—Aug. v. v. Floating on the water; the whole plant has a purple colour; the flowers dark purple, the size of *Caltha*; the petioles and under surface of the leaves are covered with a transparent gelatinous substance.

461. CALTHA. *Gen. pl. 957.*

1. C. caule erecto unifloro unifolio, foliis radicalibus cordato-ovatis obtusissimis paucidentatis multinervibus, petalis ellipticis.

*ranunculinum.**canadensis.**peltata.**canada**R. J. Davis**ficioides.*

Ranunculus Ficaria. *Walt. fl. car.* 159.

In shady cedar-swamps of New Jersey and Carolina.
2. June, July. v. v. Flowers deep yellow, the size
of *Ranunculus Ficaria*.

integerrima. 2. *C. caule erecto corymboso, foliis integerrimis sinu clauso*
orbiculato-cordatis, floralibus sessilibus reniformibus
basi obsolete crenatis, petalis obovalibus.

In wet meadows and small rivulets: New England to
Virginia. 2. May—July. v. v. Flowers yellow,
smaller than the following species.

palustris. 3. *C. caule erecto corymboso, foliis cordato-reniformibus:*
lobis patulis, circumcirca acute-crenatis, floralibus sub-
sessilibus, petalis ovatis.—*Willd. sp. pl. 2. p. 1338.*

Icon. *Fl. dan.* 668. *Engl. bot.* 506.
In low boggy meadows: Canada to Pennsylvania. 2.
April—June. v. v. Flowers golden yellow, large.

flabellifolia. 4. *C. caule procumbente, foliis dilatato-reniformibus: lobis*
patentissimis, circumcirca argute acuteque dentatis,
pedunculis axillaribus solitariis unifloris, petalis obova-
tis, capsulis rostro adunco.

In a large sand-spring on Pokono-mountain, Pennsylvania.
2. July, Aug. v. v. Flowers yellow, the size of
Ranunculus arvensis. This plant has some affinity
with *C. natans* *Willd. sp. pl. 2. p. 1339.* of which I
have seen perfect specimens in the Herbarium of A.
B. Lambert, Esq., collected by Pallas in the most
eastern parts of Siberia, under the name of *C. pusilla*,
which I distinguish by the following diagnosis: *Caltha*
natans, caule procumbente natante, foliis cordato-re-
niformibus: lobis subapproximatis, postice obsolete
crenatis, antice dentatis, petalis ovalibus, capsulis ros-
tro recto. The flowers of that species are white, with
a tinge of red, of the same size as the present species.

462. COPTIS. *Salisb. in Linn. trans. 8. p. 305.*

trifolia.

1. *C. foliis ternatis, foliolis rotundato-obovalibus acumina-*
to-crenatis, scapo unifloro.—*Salisb. l. c.*

Helleborus trifolius. *Willd. sp. pl. 2. p. 1338.*
Icon. *Fl. dan.* 566.

In cedar-swamps and mountain bogs: Canada to Vir-
ginia. 2. May—July. v. v. A delicate small plant;
flowers white.



Caltha palustris.

2. C. foliis b
—*Salisb.*
On the no
Banks.

40

1. T. petalis
ribus. S
T. america
T. pentape
In shady v
and Per
yellow,

464.

1. H. foliis ti
enum.
Anemone H
Icon. Bot. 1
α. H. lobis rot
β. H. lobis ova
In woods, o
to Carolin
purple; t

465.

1. R. glaber, c
tis integer
pedunculi
subreflexis
Icon. Fl. da
In inundated
Virginia.
low, small

2. R. adpresso-
acuminatis
Willd. sp.
Icon. Engl. 1
On the bank
plants : Pe

VOL. II.

2. *C. foliis bibernatis, foliolis subpinnatifidis, scapo bifloro. asplenifolia.*
—*Salisb. l. c. p. 306.*

On the north-west coast. *Menzies.* ♀. v. s. in Herb.
Banks. Flowers larger than the foregoing one.

463. TROLLIUS. *Gen. pl. 954.*

1. *T. petalis 5. obtusis patulis, nectariis staminibus brevioribus. Salisb. in linn. trans. 8. p. 303.*

T. americanus. Donn catal.

T. pentapetalus. Herb. Banks.

In shady wet places, on the mountains of New York
and Pennsylvania. ♀. May—July. v. v. Flowers
yellow, the size of *Ranunculus acris*.

laxus.

464. HEPATICA. *Willd. enum.....*

1. *H. foliis trilobis integerimis, scapis unifloris.—Willd. enum.*

Anemone Hepatica. Willd. sp. pl. 2. p. 1273.
Icon. *Bot. mag. 10. Fl. dan. 610.*

α. *H. lobis rotundatis obtusis.*

β. *H. lobis ovalibus acutis.*

In woods, on the side of fertile and rocky hills : Canada
to Carolina. ♀. March—May. v. v. Flowers blue or
purple ; generally smaller than in the European plant.

triloba.

obtusa.
acuta,

465. RANUNCULUS. *Gen. pl. 953.*

* *Foliis simplicibus, integris.*

1. *R. glaber, caule declinato, foliis angusto-lanceolatis acutis integerrimis denticulatisque, inferioribus petiolatis, pedunculis terminalibus axillaribus unifloris, calycibus subreflexis.—Willd. sp. pl. 2. p. 1307.*

Icon. *Fl. dan. 575. Engl. bot. 387.*

In inundated places and small rivulets : Pennsylvania and
Virginia. ♀. July—Sept. v. v. Flowers deep yellow,
small.

Flammula.

2. *R. alpresso-pilosus; foliis lanceolatis subdenticulatis acuminatis subsessilibus, caule erecto multifloro.—Willd. sp. pl. 2. p. 1308.*

Icon. *Engl. bot. 100. Fl. dan. 755.*

On the banks of rivers, among scirpus and other river
plants : Pennsylvania and Virginia. ♀. July, Aug.

Lingua.

- v. v.* Resembles the preceding one, but is upright, the flowers larger and in greater abundance.
- pusillus.* 3. *R. glaber*; foliis petiolatis, inferioribus ovatis dentatis, superioribus lineari-lanceolatis apice dentatis, supremis linearibus bracteiformibus, pedunculis alternis solitariis unifloris.
R. flammula. *Walt. fl. car.* 159. *Mich. fl. amer.* 1. p. 321.
q. June—Aug. *v. s. in Herb. Enslen.* A small species, with exceeding small flowers, petals pale yellow.
- filiformis.* 4. *R. glaber*, *pusillus*; caulinis filiformibus repentibus geniculatis, geniculis unifloris, foliis lineari-subulatis obtusis.—*Mich. fl. amer.* 1. p. 320.
 In inundated places, on the river St. Lawrence, Hudson's Bay, and Labrador. *q.* June, July. *v. s. in Herb. Banks.* Resembles *R. reptans* very much.
- Cymbalaria.* 5. *R. glaber*, *minimus*, *filiformis*, *repens*, ad genicula radicans; foliis cordato-reniformibus obtuse 5-dentatis, pedunculis radicalibus solitariis plerunque bifloris, petalis linearibus, fructibus oblongis.
 In saline marshes near the salt-works of Onondago, New York. *q.* June, July. *v. v.* Flowers the size of the foregoing one, pale yellow, sometimes white. This little plant has some resemblance to *R. salsuginosus* of Pallas, whose specimens I had an opportunity of seeing in the collection of A. B. Lambert, Esq. and which may be distinguished by the following diagnosis: *R. glaber*, *filiformis*, *repens*, ad genicula radicans; foliis ovatis rainer cordatis apice 3-dentatis, pedunculis radicalibus solitariis plerunque unifloris, petalis obovatis, fructibus subglobosis. The flowers are more than ten times larger than in *R. Cymbalaria*.
- ** *Foliis dissectis et divisis.*
- abortivus.* 6. *R. glaber*; caulinis striatis inferne nudis, foliis radicalibus cordato-reniformibus obtuse-crenatis, caulinis petiolatis ternatis angulatis, superioribus sessilibus, ranunculis subinfloris — *Willd. sp. pl.* 2. p. 1314.
 In wet places on the side of ponds and ditches: New York to Carolina. *q.* July, Aug. *v. v.* Flowers small, yellow.
- nitidus.* 7. *R. glaberrimus*; caulinis fistulosis, foliis radicalibus rotundato-subreniformibus obtuse-crenatis, caulinis sessili-

POLY

- silibus dig
bus subgl
encycl. 6.
 In inundate
July, Aug.
 and the f.
 and prob
 though di
 8. *R. glaber*; i
 silibus dig
 p. 1315.
 Icon. *Fl. da*
 In ponds and
 Aug. *v. v.*
 9. *R. pubescens*
 inciso-cren
 caule mult
 pl. 2. p. 1.
 Icon. *Fl. da*
 In old mead
 June. *v. v.*
 10. *R. pusillus*, g
 formibus u
 laciniis line
 lis oblongis
 In Labrador.
Herb. Dick.
 very near to
 11. *R. pilosus*, ere
 subtus pilosi
 petalis caly
 p. 1323.
R. canadensis.
 In low meado
 July, Aug.
 12. *R. hirsutissimu*
 sis dentatisq
 catis, calycib
 pl. 2. p. 132
 Icon. *Fl. dan.*
 In meadows an
 v. v.
 13. *R. hirsutus*; fo

silibus digitatis, foliolis incisis, laciniis obtusis, seminibus subglobosis glaberrimis.—*Walt. fl. car.* 159. *Lam. encycl.* 6. p. 126.

In inundated grounds: New York to Carolina. ♀. July, Aug. v. v. Flowers small; petals white. This and the foregoing one are nearly related to each other, and probably only varieties of the same species, though distinguished by Walter and Lamarek.

8. R. glaber; foliis inferioribus palmatis, superioribus sessilibus digitatis, fructibus oblongis.—*Willd. sp. pl.* 2. p. 1315.

Icon. *Fl. dan.* 571. *Engl. bot.* 681.

In ponds and ditches: Canada to Virginia. ♀. June—Aug. v. v. Flowers yellow, small.

9. R. pubescens; foliis radicalibus reniformibus tripartitis inciso-crenatis, caulinis sessilibus digitatis linearibus, caule multifloro, calyce colorato patente.—*Willd. sp. pl.* 2. p. 1314.

Icon. *Fl. dan.* 665. *Engl. bot.* 624.

In old meadows and woods: Pennsylvania. ♀. May, June. v. v. Flowers yellow, middle-sized.

10. R. pusillus, glaber; foliis radicalibus subcordato-reniformibus inciso-dentatis, caulinis sessilibus digitatis, laciniis linearibus integerrimis, caule paucifloro, petalis oblongis calyce subæquantibus.

In Labrador. *Colmster.* ♀. May, June. v. s. in *Herb. Dickson.* Flowers small, yellow; it approaches very near to *R. nivalis*.

11. R. pilosus, erectus, ramosus; foliis ternatis trifidis incisis subtus pilosis, pedunculis teretibus, calycibus reflexis, petalis calyce subæquantibus.—*Willd. sp. pl.* 2. p. 1323.

R. canadensis. Jacq. ic. rar. 1. t. 105.

In low meadows: Canada to Pennsylvania; rare. ♀. July, Aug. v. v. Flowers the size of *R. acris*.

12. R. hirsutissimus; foliis compositis: ternatis trifidis incisis dentatisque, caule erecto multifloro, pedunculis sultatis, calycibus retroflexis, radice bulbosa.—*Willd. sp. pl.* 2. p. 1324.

Icon. *Fl. dan.* 551. *Engl. bot.* 515.

In meadows and grass plots, common. ♀. May—Aug. v. v.

13. R. hirsutus; foliis ternatis inciso-lobatis, caule erecto

septiferatus.

auricomus.

pygmæus.

penylvani-
cus.

bulbosus.

Philonotis.

- multifloro, pedunculis sulcatis, calycibus retroflexis acuminatis, seminibus tuberculatis, radice fibrosa.—*Willd. sp. pl. 2. p. 1324.*
- R. hirsutus. Curt. lond. 2. t. 40. Engl. bot. 1504.*
In old meadows and low wet fields: New England to Pennsylvania. ♀. June—Oct. v. v. Resembles the foregoing one.
- repens.* 14. *R. hirsutus; foliis compositis: ternatis trifidis incisis, sarmenatis repentibus, pedunculis sulcatis, calycibus patulis. Willd. sp. pl. 2. p. 1325.*
Icon. *Fl. dan. 795. Engl. bot. 510.*
In shady wet woods, particularly in the mountains: Pennsylvania to Virginia. ♀. July, Aug. v. v. Flowers deep yellow.
- acris.* 15. *R. adpresso-pilosus; foliis tripartito-multifidis, summis linearibus, pedunculis teretibus, calycibus patulis.—Willd. sp. pl. 2. p. 1326.*
Icon. *Engl. bot. 652.*
In wet meadows, in the western parts of New York and Canada. ♀. May—Aug. v. v.
- lanuginosus.* 16. *R. hirsutus; foliis trifidis lobatis dentatis holosericeis, pedunculis elongatis teretibus, calycibus patulis.—Willd. sp. pl. 2. p. 1327.*
Icon. *Fl. dan. 397.*
In old fields and meadows: Pennsylvania to Carolina. ♀. June—Oct. v. v.
- tomentosus.* 17. *R. villosissimus, humilis; caule repente 1—2-floro, foliis tomentosis trilobatis, calycibus hispidis subreflexis.—Lam. encycl. 6. p. 127.*
In Carolina. *Bosc.* ♀. +. Flowers yellowish-white.
- marylandicus.* 18. *R. pubescens; caule simplici subnudo, foliis radicalibus ternatis, foliolis trilobatis, lobis acutis incisis, calycibus reflexis. Lam. encycl. 6. p. 126.*
In shady woods: Pennsylvania to Virginia. ♀. May—July. v. v. Flowers pale yellow.
- recurvatus.* 19. *R. pubescens; foliis trilobis, lobis basi cuneiformibus, apice incisis acutis, caule multifloro, corolla calycibusque recurvis, petalis linearibus.—Lam. encycl. 6. p. 125.*
In shady woods: New York to Carolina. ♀. June—Aug. v. v. Flowers small; petals almost white.
- septentrionalis.* 20. *R. glabriusculus, foliis membranaceis glabris ternatis, foliolis subtrilobatis incisis acutis, caule petiolisque*

basi hirs

Lam. en

In North yellow.

21. *R. hirsutis*

lobatis,

cifloris,

In wet field

Carolina,

pale yell

22. *R. foliis su**sp. pl. 2.**R. heterop*Icon. *Engl*

In ponds a

York.

23. *R. foliis on**Willd. sp*Icon. *Fl. da*

In tranquil

—Aug.

24. *R. glabellus*

lobatis,

ca

*pl. 2. p. 1*Icon. *Lam.*

In old fields

v. v. Flo

25. *R. glabellus,*

lobatis,

pe

cels 73.

Near Charle

ers yellow,

one.

1. *S. foliis lance*

tis rectis,

calycibusqu

p. 408.

Icon. *Fl. dan.*

In small pon

bibus retroflexis, pedunculis subbifloris, *calycibus reflexis*.
Lam. encycl. 6. p. 125.

In North America. *Lamarck.* \mathcal{U} . +. Flowers pale yellow.

21. *R. hirsutissimus*, erectus; foliis ternatis, foliolis acute lobatis, caulibus infra primum pedunculum nudis paucifloris, calycé appresso. *Mich. fl. amer. 1. p. 321.*
 In wet fields and on the banks of ditches: Virginia and Carolina. \mathcal{U} . June—Aug. *v. v.* Flowers small, pale yellow.

hispidus.

22. *R. foliis submersis capillaceis, emersis peltatis.* *Willd. sp. pl. 2. p. 1332.*
R. heterophyllum. Roth fl. germ. 1. p. 240.
Icon. Engl. bot. 101. Fl. dan. 370.
 In ponds and slow-flowing rivulets: Canada to New York. \mathcal{U} . June, July. *v. v.* Flowers white.
23. *R. foliis omnibus dichotomo-capillaceis, caule natante.* *Willd. sp. pl. 2. p. 1333.*
Icon. Fl. dan. 376.
 In tranquil rivers: Pennsylvania to Carolina. \mathcal{U} . June—Aug. *v. v.* Flowers white.

aquatilis.

fluvialis.

24. *R. glabellus, diffusus; foliis simplicibus subrotundis trilobatis, calycibus longitudine corollarum.—Willd. sp. pl. 2. p. 1329.*
Icon. Lam. illustr. t. 498. f. 2. Alp. exot. 262.
 In old fields: Virginia and Carolina. \odot . June, July. *v. v.* Flowers yellow.
25. *R. glabellus, simplex; foliis simplicibus subrotundis trilobatis, petalis calycé duplo longioribus.—Vent. hort. cels 73.*
 Near Charlestown, South Carolina. *Bosc.* +. Flowers yellow, more than twice the size of the preceding one.

muricatus.

echinatus.

466. SAGITTARIA. *Gen. pl. 1441.*
1. *S. foliis lanceolatis acutis sagittatis, lobis lanceolatis acutis rectis, scapo simplici, floribus monoicis, bracteis calycibusque lanceolatis acutis.—Willd. sp. pl. 4. p. 408.*
Icon. Fl. dan. 172.
 In small ponds and ditches: Pennsylvania to Carolina.

sagittifolia.
 $\beta.$ minor.

- latifolia.* POL
2. *S. foliis ovatis acutiusculis sagittatis, lobis ovatis tenuissime acuminatis rectis, scapo simplici, floribus monoicis, bracteis subrotundis obtusis.* — *Willd. sp. pl. 4. p. 400.*
S. sagittifolia. Mich. fl. amer. 2. p. 189.
3. *S. foliis amplis abrupte acutis, scapo subramoso, floribus dioicis.*
 In ponds and ditches: Canada to Carolina. $\frac{3}{4}$. June—Aug. *v. v.* Leaves large and broad.
- obtusa.*
3. *S. foliis dilatato-ovatis rotundato-obtusis mucronatis sagittatis, lobis approximatis oblongis oblique-acuminatis rectis, scapo simplici, floribus dioicis, bracteis ovatis acutis.* — *Willd. sp. pl. 4. p. 409.*
 In ditches and shallow ponds: Pennsylvania to Virginia.
 $\frac{3}{4}$. June—Sept. *v. v.* Leaves the size of *Calla palustris.*
- hastata.*
4. *S. foliis oblongo-lanceolatis sensim-acutis sagittatis, lobis patentibus lanceolatis longissime acuminatis, scapo simplici, floribus dioicis, bracteis calycibusque subrotundis obtusis.*
 In old ditches: Pennsylvania. $\frac{3}{4}$. July, Aug. *v. v.* Leaves the size of *S. sagittifolia*; the lobes very spreading.
- gracilis.*
5. *S. foliis linearibus obtusiusculis 3-nervibus sagittatis, lobis patentibus linearibus elongatis sensim tenuissime acuminatis, scapo simplici paucifloro, floribus dioicis, bracteis brevibus suborbiculatis.*
 In bogs and ditches: Pennsylvania to Virginia, particularly on the mountains. $\frac{3}{4}$. July, Aug. *v. v.* Leaves very slender, about three inches long, measured from the beginning of the petiole.
- heterophylla.*
6. *S. foliis simplicibus linearibus lanceolatisque utrinque acutis, seu ellipticis acutis basi obtusis sagittatis, lobis divaricatis linearibus, scapo simplici paucifloro, floribus monoicis, femaleis subsessilibus, bracteis latovatis acuminatis.*
 In small ponds and ditches: Pennsylvania and New Jersey. $\frac{3}{4}$. June—Aug. *v. v.* Leaves about two inches and a half long; this species is easily distinguished by the female flowers being almost sessile.
7. *S. foliis latmoso, se S. lancifolia In ponds: J. yon.*
8. *S. foliis angue acutis In still and falls, Ne cies grow the petio numerous*
9. *S. foliis line simplici n que rotun In small po Leaves ab flowers fr*
10. *S. foliis line plici pauc obtusis. S. graminea tillorum p In Carolina, Aug. v. :*
11. *S. foliis sub simplici patis acumin In small riv July, Aug with a dil*
12. *S. foliis natu basi atten paucifloro, fl. amer. 2 In rivulets of in Herb. I and a half*

POLYANDRIA POLYGYNIA. Sagittaria.

397

7. *S. foliis lato-lanceolatis inferne angustatis, scapo subramoso, seminibus planis falcatis rostratis.*

S. lancifolia. Mich. fl. amer. 2. p. 189.

In ponds: Carolina. Ʌ. July, Aug. v. s. in *Herb. Lyon.*

8. *S. foliis angusto-lanceolatis inferne carinatis rigidis utrinque acutissimis, scapo ramoso, floribus monoicis.*

In still and deep waters of Oswego river, near the great falls, New York. Ʌ. July, Aug. v. v. This species grows in a depth of more than seven feet water; the petioles are strong and stiff; the flowers very numerous.

9. *S. foliis linear-lanceolatis acutis inferne angustatis, scapo simplici multifloro, floribus dioicis, bracteis calycibusque rotundatis obtusis.*

In small ponds: New Jersey. Ʌ. July, Aug. v. v. Leaves about six inches long and half an inch wide; flowers from twelve to eighteen in a scape.

10. *S. foliis linearibus longissimis 3-nervibus, scapo simplici paucifloro, floribus monoicis, bracteis oblongis obtusis.*

S. graminea; foliis lanceolato-linearibus, capitulis pistillorum pusillis. Mich. fl. amer. 2. p. 190?

In Carolina. Lyon. Canada. Michaux. Ʌ. July, Aug. v. s. in *Herb. Lyon.*

11. *S. foliis subulatis basi vaginantibus dorso convexis, scapo simplici paucifloro, floribus monoicis, bracteis dilatatis acuminatis.*

In small rivulets and ditches, near Philadelphia. Ʌ. July, Aug. v. v. A small species; leaves sometimes with a dilated point, representing a leaf.

12. *S. foliis natantibus elliptico-lanceolatis obtusis 3-nervibus basi attenuatis, infimis subcordatis, scapo simplici paucifloro, pedunculis inferioribus elongatis.—Mich. fl. amer. 2. p. 190.*

In rivulets of Lower Carolina. Ʌ. July, Aug. v. s. in *Herb. Lyon.* Flowers small; leaves about an inch and a half long. ●

falcata.

rigida.

simplex.

graminea.

acutifolia.

natans.

of a single
whole p.
of *Skunk*
truncated

4

1. C. foliis sulcata.—*Icon. Fl. d*
- Icon. *Fl. d*
- In sphagnolets : *Calla*
- Flowers v

4

1. A. acaule ; spatha elongata culifloro.—*Calla virginiana*
- In overflow Scirpi an long, inv
2. A. acaule ; spatha curvad apicem
- Calla sagittifolia*
- Caladium sag.
- On the banks Carolina.
3. A. acaule ; dice cylind taliter pate
- Icon. Pluk.*
- In swamps of v. v. Spat

4. A. subcaulescens ; spadicea dunculata d
- sp. pl. 4. p.*
- α. A. spatha virens
- Icon. Pluk. a*
- β. A. spatha atrata
- Icon. Pluk. al*
- In shady wet

467. CYAMUS. *Salisb. in annals of bot. 2. p. 75.*Nelumbium. *Willd. sp. pl. 2. p.**flavicomus.*

1. C. foliis peltatis orbiculatis integerrimis, corolla polypetala, antheris superne linearibus.—*Salisb. l. c.*

Nelumbium luteum. *Willd. sp. pl. 2. p. 1259.**Nymphaea Nelumbo β. Sp. pl. 730.**Icon. Bartr. ic. ined. in mus. Banks.*

In ponds and lakes of Carolina and Virginia; also near Philadelphia, in ditches and ponds of Brobston's meadows. ♀. July. v. v. Flowers large, yellow, resembling a double tulip. The seeds are very agreeable to eat, and eagerly sought for by children and Indians. By the latter it is supposed they were introduced to those ponds near Philadelphia, as there is no other instance known to have been found so far north. Walter mentions his *Nelumbo* to have white flowers; this variety I have never seen.

pentapetalus.

2. C. foliis peltatis orbiculatis integerrimis, calyx pentaphyllo, corolla pentapetalum.

Nelumbium pentapetalum. *Willd. sp. pl. 2. p. 1259.**Nymphaea pentapetalum. Walt. fl. car. 155.*

In North and South Carolina. ♀. July. v. s. Flowers large, white. A specimen seen in the collection of a gentleman in Carolina ascertains the existence of this formerly doubtful plant; but unfortunately I took no notes at that time, being in expectation of seeing the living plant.

reniformis.

3. C. foliis reniformibus, corolla polypetala.

Nelumbium reniforme. *Willd. sp. pl. 2. p. 1260.**Nymphaea reniformis. Walt. fl. car. 155.*

In Carolina. *Walter.* ♀. +. This is a doubtful species, and probably a plant belonging to a different genus.

468. POTHEOS. *Gen. pl. 210.**fœtida.*

1. P. acaulis ; foliis ovatis cordatis, spadicea subgloboso.—

*Mich. fl. amer. 2. p. 186.*Dracontium fœtidum. *Willd. sp. pl. 2. p. 288.**Icon. Bot. mag. 836. Catesb. car. 2. t. 71.*

In low grounds, on the side of shady rivulets and springs :

Canada to Virginia. ♀. Feb.—April. v. v. Leaves large; flowers several, almost sessile on the ground,

Symplocarpus
fœtidus
Muhl.

2. p. 75.

...
rolla polype-
t. l. c.
259.

a; also near
boston's mea-
yellow, re-
very agreea-
dren and In-
were intro-
s there is no
so far north.
ite flowers;

ilyce penta-
p. 1259.

s. Flowers
lection of a
ence of this
I took no
seeing the

260.

ful species,
t genus.

globoso.—

d springs:
. Leaves
ground,

of a singular shape, spotted yellow and brown; the whole plant very foetid, from which it has the name of *Skunk-weed* or *Skunk-cabbage*. It has four distinct, truncated, fleshy petals, and four stamens to each germ.

469. CALLA. *Gen. pl.* 1388.

1. C. foliis subrotundo-cordatis acutis, spatha ovata cuspidata.—*Willd. sp. pl.* 2. p. 290. *palustris.*

Icon. *Fl. dan.* 422.

In sphagnum swamps, on the mountains and near rivulets: Canada to Pennsylvania. 2. July, Aug. v. v. Flowers white.

470. ARUM. *Gen. pl.* 1387.

1. A. acaule; foliis hastato-cordatis acutis, lobis obtusis, spatha elongata incurva, spadice superne longius masculifloro.—*Willd. sp. pl.* 4. p. 484. *virginicum.*

Calla virginica. *Mich. fl. amer.* 2. p. 187.

In overflowed places on the banks of rivers, among Scirpi and Junci. 2. June, July. v. v. Spatha long, involute, green.

2. A. acaule; foliis subhastato-sagittatis, lobis oblongis, *sagittifolium.* spatha cucullata superne ovali, spadice longo a medio ad apicem masculifloro.

Calla sagittifolia. *Mich. fl. amer.* 2. p. 187.*Caladium sagittifolium. Willd. sp. pl.* 4. p. 489?

On the banks of rivers and old swamps: Pennsylvania to Carolina. 2. July, Aug. v. v.

3. A. acaule; foliis ternatis, foliolis ovatis acuminatis, spadice cylindrico spatha sessili ovata acuminata horizontaliter patenti altiore.—*Willd. sp. pl.* 4. p. 481. *atrorubens.*

Icon. *Pluk. alm.* t. 148. f. 6. *bona.*

In swamps of Virginia and Maryland. 2. May, June. v. v. Spatha dark brown, of a disagreeable smell.

4. A. subcaulescens; foliis ternatis, foliolis ovatis acuminatis, spadice clavato spatha ovata acuminata plana pendulata dimidio breviore, floribus monoicis.—*Willd. sp. pl.* 4. p. 480. *triphyllum.*

α. A. spatha virescente.

virens.

Icon. *Pluk. alnath.* t. 376. f. 3.

β. A. spatha atropurpurea.

atropurpu-

Icon. *Pluk. alm.* t. 77. f. 5.

reum.

In shady wet woods: Canada to Carolina. 2. June,

July. *v. v.* Flowers handsomely green and yellow, or green and purple striped. Roots known as a medicine among the natives, particularly in colics: they are generally called *Indian Turnips*.

Dracontium. 5. *A. acaule; foliis pedatis, foliolis lanceolato-oblongis integerrimis, spadice subulato spatha oblonga convoluta longiore.* — *Willd. sp. pl. 4. p. 478.*

Icon. *Pluk. alm. t. 271. f. 2.*

In shady woods: Virginia to Florida. *v.* June, July.

v. v.

XI.

- 471. AJUGA
Stami.
- 472. TEUCHE
sura ej.
- 473. MENTI
emarg.
- 474. ISANT
*tubo re
subaequ.*
- 475. CUNIL
*sus. C.
emarginata.*
- 476. HYSSO
nula in
- 477. NEPET
*culo:
faux m*
- 478. LAMIU
*tum: 1
gine de*
- 479. GALEO
fornicata
- 480. STACH
*bium i
dia ma
tera ref*
- 481. LEONU
*bium si
tripartit*
- 482. GLECH
par in f

and yellow,
vn as a me-
colics : they

oblongis in-
ga convoluta

June, July.

XIII. DIDYNAMIA.

I. GYMNOSPERMIA.

* *Calyces subquinquedidi.*

471. AJUGA. *Cor.* labium superius minimum bidentatum.
Stamina labio superiore longiora.
472. TEUCRIUM. *Cor.* labium superius nullum, sed fis-
sura ejus loco in qua *Stamina* jacent.
473. MENTHA. *Cor.* subæqualis, 4-fida : lacinia latiore
emarginata. *Stam.* erecta, distantia.
474. ISANTHUS. *Cal.* campanulatus. *Cor.* 5-partita :
tubo recto, angusto : lacinias ovatis æqualibus. *Stam.*
subæqualia. *Stigm.* linearia, recurva.
475. CUNILA. *Cal.* cylindricus, 5-dentatus, fauce villo-
sus. *Cor.* ringens : labio superiore erecto, plano,
emarginato. *Stam.* 2. sterilia.
476. HYSSOPUS. *Cor.* labium inferius tripartitum, laci-
nula intermedia crenata. *Stam.* recta, distantia.
477. NEPETA. *Cal.* aridus, striatus *Cor.* tubo longius-
culo : labio inferioris lacinula intermedia crenata ;
faux margine reflexo. *Stam.* approximata.
478. LAMIUM. *Cor.* labium superius integrum, fornica-
tum : labium inferius bilobum : faux utrinque mar-
gine dentata.
479. GALEOPSIS. *Cor.* labium superius subcrenatum,
fornicatum : labium inferius supra 2-dentatum.
480. STACHYS. *Cor.* labium superius fornicatum : la-
bium inferius lateralibus reflexum : lacinia interme-
dia maiore emarginata. *Stam.* deflorata versus la-
tera reflexa.
481. LEONURUS. *Cal.* 5-gonus, 5-dentatus. *Cor.* la-
bium superius villosum, planum, integrum : inferius
tripartitum, lacinia media indivisa.
482. GLECHOMA. *Cal.* 5-fidus. *Antherarum* singulum
par in formam crucis connivens.

483. MARRUBIUM. *Cal.* hypocrateiformis, rigidus, 10-striatus. *Cor.* labium superius bifidum, lineare, rectum.
484. HYPTIS. *Cal.* 5-dentatus, crescens. *Cor.* ringens : labium superius bifidum, inferius tripartitum, lacinia media sacculiformi. *Stam.* tubi ventri inserta, declinata.
485. PYCNANTHEMUM. *Involucrum* multibracteatum, capitulis subjectum. *Cal.* tubulatus, striatus. *Cor.* labium superius subintegrum, inferius trifidum. *Stam.* subæqualia.
- ** *Calyces bilabiati.*
486. CLINOPODIUM. *Involucrum* multisetum, verticillo subjectum. *Cor.* labium superius planum, obcordatum, rectum.
487. ORIGANUM. *Strobilus* tetragonus, spicatus, calyces colligens. *Cor.* labium superius erectum, planum : inferius tripartitum : laciniis æqualibus.
488. DRACOCEPHALUM. *Cor.* faux inflata, labium superius concavum.
489. PRUNELLA. *Cor.* labium superius dilatatum. *Filam.* bifurca : altero apice antherifera. *Stigm.* bifidum.
490. SCUTELLARIA. *Cal.* ore integro, post florescentiam clauso, operculato. *Cor.* tubus elongatus.
491. THYMUS. *Cal.* campanulatus, faux villis clausa. *Cor.* labium superius planum, emarginatum.
492. CALAMINTHA. *Cal.* defloratus villis clausus. *Cor.* fauce inflata, labio superiore emarginato, inferiore tripartito : lacinia intermedia integra, subemarginata aut crenulata.
493. HEDEOMA. *Cal.* basi gibbus. *Cor.* ringens. *Stam.* 2. sterilia.
494. TRICHOSTEMA. *Cor.* labium superius falcatum. *Stam.* longissima.

II. ANGIOSPERMIA.

* *Calyces quinquefidi.*

495. LINNÆA. *Cal.* superus, duplex : fructus 2-phyllos ; floris 5-partitus. *Cor.* campanulata. *Bacca* sicca, 3-locularis.
496. PHRYMA. *Cal.* cylindricus, supra longior, trifidus, infra bidentatus. *Cor.* labium superius emarginatum, inferius majus. *Sem.* unicum.

497. VERBEI tubo in 5-fida.
498. ZAPAN obliqui nectens
499. CAPRAI 5-fida, sperma.
500. HERPE *Cor.* tu 2-valvis
501. LINDER gens, 1 basi bicla bilamell mento v
502. SCROPH resupina
503. BIGNON campan locularis
504. RUELLI limbo 5-tenuata,
505. BUCHNI æqualis :
506. AN TIRR ringens : locularis
507. GERARD feriore 3 tito. *Ca*
508. PEDICUL 2-locular
509. MIMULU ringens : crassum.
510. CHELON ringens, brevius, margine
511. PENTSTE ventricos superne 2-valvis.

nis, rigidus,
lum, lineare,

Cor. ringens :
tum, lacinia
inserta, de-

ibracteatum,
riatus. *Cor.*
fidum. *Stam.*

m, verticillo
m, obcorda-

catus, caly-
m, planum :

ita, labium

dilatatum.
Stigm. bi-

t florescen-
igatus.

llis clausa.
m.

ausus. *Cor.*
o, inferiore

emarginata
ens. *Stam.*

; falcatum.

3-phyllo ;
cca sicca,

, trifidus,
rginatum,

497. VERBENA. *Cal.* 5-fidus. *Cor.* infundibuliformis, tubo incurvo, limbo inaequali, 5-fido. *Sem.* 2—4.
498. ZAPANIA. *Flor.* capitati. *Cal.* 5-dentatus. *Cor.* 5-fida. *Stam.* 4. fertilia. *Stigm.* peltato-capitatum, obliquum. *Fructus* tectus utricularis, evanescens, nectens semina 2.
499. CAPRARIA. *Cal.* 5-partitus. *Cor.* campanulata, 5-fida, acuta. *Caps.* 2-valvis, 2-locularis, polysperma.
500. HERPESTIS. *Cal.* inaequalis, basi 2-bracteatus. *Cor.* tubulosa, subbilabiata. *Stam.* inclusa. *Caps.* 2-valvis, 2-locularis, dissepimento valvis parallelo.
501. LINDERNIA. *Cal.* 5-partitus, aequalis. *Cor.* ringens, labio superiore brevissimo, inferiore 3-fido, basi bicarinata. *Anther.* per paria cohaerentes. *Stigm.* bilamellatum. *Caps.* 2-locularis, 2-valvis, dissepi-mento valvis parallelo.
502. SCROPHULARIA. *Cal.* 5-fidus. *Cor.* subglobosa, resupinata. *Caps.* 2-locularis.
503. BIGNONIA. *Cal.* 5-fidus, cyathiformis. *Cor.* fauce campanulata, 5-fida, subtus ventricosa. *Siliqua* 2-locularis. *Sem.* membranaceo-alata.
504. RUELLIA. *Cal.* 5-partitus. *Cor.* subcampanulata, limbo 5-fido. *Stam.* conjugata. *Caps.* utrinque attenuata, dentibus elastice dehiscens. *Sem.* paucula.
505. BUCHNERA. *Cal.* 5-dentatus. *Cor.* limbus 5-fidus, aequalis : lobis cordatis. *Caps.* 2-locularis.
506. ANGIRRHINUM. *Cal.* 5-phyllo. *Cor.* calcarea, ringens : rictu clauso palato prominente. *Caps.* 2-locularis, 2-valvis.
507. GERARDIA. *Cal.* 5-fidus. *Cor.* 2-labiata : labio inferiore 3-partito : lobis emarginatis : medio bipartito. *Caps.* 2-locularis, dehiscens.
508. PEDICULARIS. *Cal.* 5-fidus. *Cor.* ringens. *Caps.* 2-locularis, mucronata, obliqua. *Sem.* tunicata.
509. MIMULUS. *Cal.* prismaticus, 5-dentatus. *Cor.* ringens : labio superiore lateribus replicato. *Stigm.* crassum. *Caps.* 2-locularis, polysperma.
510. CHELONE. *Cal.* 5-partitus, 3-bracteatus. *Cor.* ringens, ventricosa. *Filam.* quintum sterile, cæteris brevius. *Caps.* 2-locularis, 2-valvis. *Sem.* plurima, margine membranacea.
511. PENTSTEMON. *Cal.* 5-phyllo. *Cor.* 2-labiata, ventricosa. *Filam.* quintum sterile, cæteris longius, superne barbatum. *Caps.* compressa, 2-locularis, 2-valvis. *Sem.* numerosa, subglobosa.

512. MARTYNIA. *Cal.* 5-fidus. *Cor.* ringens. *Caps.* lignosa, corticata, rostro hamato, 4-locularis, 2-valvis.

** *Calyces quadrifidi.*

513. SCHWALBEA. *Cal.* ventricoso-tubulosus, 4-fidus: lacinia superiore minima, intima maxima, emarginata. *Cor.* ringens. *Caps.* 2-locularis, 2-valvis: dissepinamento duplicato. *Sem.* paleacea.

514. RHINANTHUS. *Cal.* ventricosus, 4-fidus. *Cor.* ringens: labio superiore compresso. *Caps.* 2-locularis, compressa, obtusa.

515. BARTSIA. *Cal.* 2-lobus, emarginatus, coloratus. *Cor.* minus ipso calyce: labio superiore longiore. *Caps.* 2-locularis. *Sem.* angulata.

516. EUPHRASIA. *Cal.* cylindricus, 4-fidus. *Cor.* bilabiata: labium superius 2-fidum, inferius 3-lobum: lacinias bifidis. *Anth.* inferiores lobis spinosis.

517. MELAMPYRUM. *Cal.* 4-fidus. *Cor.* labium superius compressum, margine replicato. *Caps.* 2-locularis, obliqua, hinc dehiscens. *Sem.* 2. gibba.

*** *Calyces bifidi.*

518. OBOLARIA. *Cor.* campanulata, 4-fida. *Stam.* ex divisuris corollæ. *Caps.* 1-locularis, 2-valvis, polysperma.

519. OKOBANCHE. *Cor.* ringens. *Caps.* 1-locularis, 2-valvis, polysperma. *Glandula* sub basi germinis.

*Chamaepi-
thys.*

I. GYMNO S P E R M I A.

471. AJUGA. *Gen. pl.* 959.

1. A. foliis trifidis, floribus axillaribus solitariis folio brevioribus, caule diffuso. *Willd. sp. pl.* 3. p. 10.
Teucrium Chamaepithys. Sp. pl. 787.
Icon. *Fl. dan.* 733. *Rivin. t.* 24.
In sandy fields of Virginia. *Gronov.* ○. June, July.
v. s. in *Herb. Gronov.* Flowers pale yellow.

47

1. *T. cano-his-*
omnibus
confertis
sp. pl. 3.
In low gro-
nada to 1
purple.

2. *T. pubescens*
subsessili-
tis, bract
p. 22.
Icon. *Schle-*
In low grou-
June—A
very muc

47.

1. *M. floribus*
tis pilosis,
sp. pl. 3. 1
In Canada.

2. *M. assurgentis*
latis utrinque
nibus exer-
On the bank
sylvania.

3. *M. erecta, g*
silibus, sp
staminibus
M. viridis. *h*
In wet places
June—Au-

474. ISA

1. *I. viscidopu-*
acutis 3-ni-
amer. 2. p
Icon. *Mich.* J
On dry hills,
○. July, t

472. TEUCRIUM. Gen. pl. 960.

1. T. cano-hirsutissimum; foliis ovato-lanceolatis serratis, omnibus petiolatis, caule erecto, spicis verticillatis confertis, bracteis calyce duplo longioribus.—Willd. sp. pl. 3. p. 22.

In low grounds, on the borders of ponds and lakes: Canada to New York. ♀. July, Aug. v. v. Flowers purple.

2. T. pubescens; foliis ovato-oblongis serratis, superioribus subsessilibus, caule erecto, spicis verticillatis confertis, bracteis longitudine calycis.—Willd. sp. pl. 3. p. 22.

Icon. Schkuhr handb. 160.
In low grounds and bogs: New York to Carolina. ♀. June—Aug. v. v. Resembles the foregoing species very much.

473. MENTHA. Gen. pl. 967.

1. M. floribus verticillatis, foliis lanceolatis serratis petiolatis pilosis, staminibus corollam æquantibus. Willd. sp. pl. 3. p. 82.

In Canada. Kalm. ♀. +.

2. M. assurgens, pubescens; foliis petiolatis ovali-lanceolatis utrinque acutissimis, floribus verticillatis, staminibus exsertis.—Mich. fl. amer. 2. p. 2.

On the banks of rivers and near springs: Canada to Pennsylvania. ♀. July, Aug. v. v. Flowers pale purple.

3. M. erecta, glabriuscula; foliis lanceolato-ovatis subsessilibus, spica gracili, verticillis minimis interrupta, staminibus non exsertis.—Mich. fl. amer. 2. p. 2.

M. viridis. Walt. fl. car. 161.

In wet places, near springs: Pennsylvania to Georgia. ♀. June—Aug. v. v. Flowers white.

*canadense.**virginicum.**canadensis.**borealis.**tenius.**cærulea.*

474. ISANTHUS. Mich. fl. amer. 2. p. 3.

1. I. viscidopubescent; foliis ovali-lanceolatis utrinque acutis 3-nervibus, pedunculis 1-2-floris. Mich. fl. amer. 2. p. 4.

Icon. Mich. fl. amer. 2. t. 30.

On dry hills, in lime-stone soil: Virginia and Kentucky. ○. July, Aug. v. v. Flowers pale blue.

gens. Caps.
locularis, 2-

as, 4-fidus:
na, emarginata,
2-valvis:

fidus. Cor.
ups. 2-locu-

coloratus.
e longiore.

. Cor. bi-
s 3-lobum :
osis.
rium super-
ps. 2-locu-
bba.

Stam. ex
ilvis, poly-
-locularis,
germinis.

folio bre-
10.

ane, July.

In cultivate
○. May

2. L. caule his
bescentib
In shady wo
large, wh

479

1. G. internodi
mis subec
lyce parue
p. 92.

Icon. Engl.
On road side
July, Aug

480

1. S. glabriusc
lanceolatis
ris.—Mich
S. palustris.
In Carolina.

2. S. caulibus ei
tiolatis lanc
lis subsexfl
Mich. fl. a
S. tenuifolia.
S. arvensis. *H*
In low fields a
Virginia,
lustris; flo

3. S. caule foliis
oblongis ob
cibus glabri
Galeopsis florii
In old fields at
July. *v. v.*
has been pl
widely diffi
seum suffici

4. S. erecta, pub
pra glabris
subsexfloris.

VOL. II.

mariana.

1. C. foliis ovatis serratis sessilibus, corymbis terminalibus dichotomis.—*Willd. sp. pl. 1. p. 122.*

Icon. *Pluk. mant. t. 344. f. 1.*

On dry shady hills: New York to Carolina. *U.* June, July. *v. v.* Flowers red. The whole herb has an aromatic scent, and is used as tea in severe colds and other complaints, under the name of *Dittany*.

476. HYSSOPUS. *Gen. pl. 963.**nepetoides.*

1. H. spicis verticillatis cylindricis, stylis corolla brevioribus, foliis subcordatis ovatis acuminatis acute dentatis.—*Willd. sp. pl. 3. p. 48.*

Icon. *Jacq. hort. 1. t. 69. Pluk. alm. t. 150. f. 3.*

In fertile shady woods, along hedges: Pennsylvania to Virginia. *U.* July, Aug. *v. v.* From three to five feet high; flowers in long close spikes, yellowish.

scrophulari-folius.

2. H. spicis verticillatis cylindricis, stylis corolla longioribus, foliis cordatis ovatis acuminatis obtuse dentatis.—*Willd. sp. pl. 3. p. 48.*

Agastache. *Gronov. fl. virg. 88.*

Icon. *Herm. parad. t. 106.*

In wet copses and on the banks of rivers: Pennsylvania to Virginia; rare. *U.* July, Aug. *v. v.* Seldom above two feet high; flowers purple.

477. NEPETA. *Gen. pl. 964.**Cataria.*

1. N. floribus spicatis, verticillis subpedicellatis, foliis petiolatis cordatis dentato-serratis. *Willd. sp. pl. 3. p. 49.*

Icon. *Fl. dan. 580. Schkuhr handb. 157.*

On road sides and in cultivated grounds: frequent. *U.* July, Aug. *v. v.* Flowers white. Common *Cat-nep*, so well known as a domestic medicine, has probably been introduced from Europe.

478. LAMIUM. *Gen. pl. 971.**amplexicaule.* 1. L. foliis floralibus sessilibus amplexicaulibus obtusis.

Willd. sp. pl. 3. p. 90.

Pollichia amplexicaulis. Roth germ. 1. p. 254.

Icon. *Fl. dan. 752. Schkuhr handb. 159.*

In cultivated grounds, common ; probably introduced.

⊖. May—Aug. v. v. Flowers purple.

2. L. caule hispido, foliis longe petiolatis lato-cordatis pu-

bescientibus, axillis unifloris.—*Mich. fl. amer. 2. p. 4.*

In shady woods of Tennessee. *Michaux.* +. Flowers large, white.

479. GALEOPSIS. *Gen. pl. 972.*

1. G. internodiis caulinis superne incrassatis, verticillis summis subcontiguis, calycibus pungentibus, corollis calyce parum longioribus, caule hirto. *Willd. sp. pl. 3. p. 92.*

Icon. *Engl. bot. 207.*

On road sides, in the western parts of Pennsylvania. ⊖.

July, Aug. v. v. Flowers white, with purple spots.

Tetrahit.

480. STACHYS. *Gen. pl. 974.*

1. S. glabriuscula, gracilis, erecta ; foliis sessilibus linearilanceolatis rarer subdentatis, verticillis subquadrifloris.—*Mich. fl. amer. 2. p. 4.*

S. palustris. *Walt. fl. car. 162.*

In Carolina. *Michaux.* +.

aspera.

2. S. caulis erectis, retrorsum hispidissimis, foliis subpetiolatis lanceolatis acute serratis glaberrimis, verticillis subsexfloris, calycibus divaricato-spinescentibus.—*Mich. fl. amer. 2. p. 5.*

S. tenuifolia. *Willd. sp. pl. 3. p. 100.*

S. arvensis. *Walt. fl. car. 162.*

In low fields and on the banks of rivers : New York to Virginia. ⊖. June—Aug. v. v. Resembles *S. palustris*; flowers purple.

hispida.

3. S. caule foliisque hispido-pilosus; foliis petiolatis ovato-oblongis obtuse-serratis, verticillis subsexfloris, calycibus glabriusculis.

Galeopsis floris galea rubente, &c. *Gron. fl. virg. n. 271.*
In old fields and low pastures : Virginia to Carolina. ⊖.

July. v. v. Flowers purple. Gronovius's synonym has been placed under *S. annua*, from which it is widely different, as his specimen in the Banksian Museum sufficiently demonstrates.

4. S. erecta, pubescens; foliis cordato-ovatis dentatis, supra glabris subtus albicanibus tomentosis, verticillis subsexfloris.

Foeniculum.

On the banks of the Missouri. ♂. July. v. s. Flowers blue. The whole of this plant has a scent extremely like that of fennel.

481. LEONURUS. *Gen. pl.* 977.*Cardiaca.*

1. L. foliis cuneiformi-ovatis trilobis dentatis, corollis calyce pungente majoribus, lacinia media labii inferioris acuta.—*Willd. sp. pl.* 3. p. 114.
Icon. *Fl. dan.* 727. *Engl. bot.* 286.
In old fields and hedges: Pennsylvania to Virginia; probably introduced from Europe. ♀. July, Aug. v. v. Flowers white, mixed with red.

482. GLECHOMA. *Gen. pl.* 970.*hederacea.*

1. G. foliis reniformibus crenatis. *Willd. sp. pl.* 3. p. 85.
Icon. *Fl. dan.* 789. *Curt. lond.* 2. t. 44.
In orchards and on grassy hills: Pennsylvania to Virginia. ♀. May, June. v. v. Flowers handsome, blue.

483. MARRUBIUM. *Gen. pl.* 976.*vulgare.*

1. M. foliis subrotundo-ovatis dentatis rugoso-venosis, calycibus dentibus setaceis uncinatis. *Willd. sp. pl.* 3. p. 111.
Icon. *Fl. dan.* 1036. *Engl. bot.* 410.
On road sides; frequent. ♀. July, Aug. v. v. Flowers small, white.

484. HYPTIS. *Gen. pl.* 969.*capitata.*

1. H. capitulis oppositis, pedunculo internodiis longitudine, bracteis lanceolatis calyce frugifero brevioribus, foliis oblongis utrinque attenuatis inaequaliter serratis.—*Willd. sp. pl.* 3. p. 84. *Poiteau in annal. du mus.* 7. p. 464.

Clinopodium capitatum. *Swartz. prodr.* 88.

Icon. *Jacq. ic. rar.* t. 114. *Annal. du mus.* 7. t. 27. f. 1.
In Lower Carolina. *Michaux.* ♀. July, Aug. v. v. Flowers white.

radiata.

2. H. capitulis oppositis, pedunculo internodiis longitudine, bracteis lanceolatis calyce frugifero longioribus, foliis oblongis serratis.—*Willd. sp. pl.* 3. p. 84. *Poiteau in annal. du mus.* 7. p. 466.

Clinopodium
Icon. *Dill.*
f. 2
In Lower C

485. PYCN

* Bract

1. P. foliis obli-
sis, capitu-
teis setaceis
Clinopodium
Icon. *Dill.*
In low field

July—Oc
white, tir

2. P. foliis lan-
subcordic-
—*Mich. fl.*
Nepeta virgin-

Icon. *Mich.*
In dry woods

♀. July,

3. P. foliis oval-
sessili, caly-
Mich. fl. a
On high mo-
Stem and l

4. P. hirsutum ;
bus serrati-
liatis, calyc-
p. 8.
Origanum inc

Icon. *Mich. J.*

On the moun

—Aug. . v

5. P. caule strict-
3-nervibus
culato-cory-
Brachystemum
Brachystemum
Thymus virgin

phys.

v. s. Flowers
cent extremely

'7.

corollis calyce
nferioris acuta.Virginia ; pro-
illy, Aug. v. v.

0.

pl. 3. p. 85.

nia to Virginia.
ome, blue.

76.

o-venosis, ca-
illd. sp. pl. 3.

v. v. Flow-

is longitudine,
oribus, foliis
er serratis.—
al. du mus. 7.7. t. 27. f. 1.
, Aug. v. v.s longitudine,
oribus, foliis
Poiteau in

Clinopodium rugosum. *Sp. pl.* 822.
 Icon. *Dill. elth. t. 75. f. 86.* *Annal. du mus. 7. t. 27.*
f. 2
 In Lower Carolina. 4. +.

485. PYCNANTHEMUM. *Mich. fl. amer. 2. p. 7.*Brachystemum. *Mich. fl. amer. 2. p. 5.** *Staminibus exsertis.*

1. P. foliis oblongo-ovatis acutis subserratis cano-tomento-
sis, capitulis compositis, lateralibus pedunculatis, brac-
teis setaceis.—*Mich. fl. amer. 2. p. 7.*

Clinopodium incanum. *Willd. sp. pl. 3. p. 132.*
 Icon. *Dill. elth. t. 74. f. 85.* *Pluk. mant. t. 344. f. 7.*
 In low fields and copses : New York to Carolina. 4.
 July—Oct. v. v. About three feet high ; flowers
white, tinged with red.

2. P. foliis lanceolato-ovatis subserratis brevissime petiolatis
subcandicantibus, capitulis sessilibus, bracteis aristatis.
—Mich. fl. amer. 2. p. 8.

Nepeta virginica. *Willd. sp. pl. 3. p. 56.*
 Icon. *Mich. fl. amer. 2. t. 33.* *Pluk. alm. t. 85. f. 2.*
 In dry woods, in lime-stone soil : Maryland to Carolina.
 4. July, Aug. v. v. Flowers white, very small.

3. P. foliis ovali-lanceolatis serratis subsessilibus, capitulo
sessili, calycibus approximatis erectis breviter dentatis.
Mich. fl. amer. 2. p. 8.

On high mountains of Carolina. *Michaux.* 4. +.
 Stem and leaves tinged with purple.

4. P. hirsutum ; foliis abrupte petiolatis subcordato-ovali-
bus serratis, bracteis magnis coloratis, bracteolis ci-
liatis, calycibus summitate barbatis.—*Mich. fl. amer. 2.*
p. 8.

Origanum incanum. *Walt. fl. car. 165.*
 Icon. *Mich. fl. amer. 2. t. 34.*
 On the mountains of Virginia and Carolina. 4. June
 —Aug. v. v. Flowers small, pale red.

5. P. caule stricto ramosissimo asperiusculo, foliis linearibus
3-nervibus integerrimis, capitulis terminalibus fasci-
culato-corymbosis.

Brachystemum virginicum. *Mich. fl. amer. 2. p. 6.*
 Brachystemum linifolium. *Willd. enum. 623.*
 Thymus virginicus. *Willd. sp. pl. 3. p. 145.*

*incanum.**aristatum.**montanum.**Monardella.**linifolium.*

Satureja virginiana. *Sp. pl. 2. p. 793.*

Origanum flexuosum. *Walt. fl. car. 165.*

Icon. *Herm. parad. t. 218. Pluk. alm. t. 54. f. 2.*

In dry swamps and mountain meadows: New England to Carolina. ♀. July, Aug. v. v. Flowers very small, entirely white. This and the following species approach so near to the genus *Westringia*, that they almost ought to be added to that genus.

** *Staminibus inclusis.*

- lanceolatum.* 6. *P. caule stricto ramosissimo asperiusculo, foliis linearilanceolatis venosis integerrimis, capitulis terminalibus fasciculato-corymbosis.*
Brachystemum lanceolatum. *Willd. enum. 623.*

In dry swamps and on the banks of rivers: Pennsylvania to Virginia. ♀. July, Aug. v. v. Resembles the foregoing species very much; flowers white; the lower lip of the corolla has purple dots.

- muticum.* 7. *P. foliis ovato-lanceolatis subdentatis glabriusculis, capitulis terminalibus, bracteis lanceolatis acutiusculis.—Pers. syn. 2. p. 128.*
Brachystemum muticum. *Mich. fl. amer. 2. p. 6.*

Icon. *Mich. fl. amer. 2. t. 32.*

In Upper Carolina. *Michaux.* ♀. +.

- verticillatum.* 8. *P. foliis ovato-lanceolatis integerrimis, verticillis sessilibus compactis, bracteis acuminatis.—Pers. syn. 2. p. 128.*
Brachystemum verticillatum. *Mich. fl. amer. 2. p. 6.*

Origanum clinopodioides. *Walt. fl. car. 165.*

Icon. *Mich. fl. amer. 2. t. 31.*

In the mountains: Pennsylvania to Carolina. ♀. July, Aug. v. v.

486. CLINOPodium. *Gen. pl. 980.*

- vulgare.* 1. *C. capitulis verticillatis, bracteis setaceis hispidis, foliis superne pilosis remote dentatis, caule subsimplici.*
Willd. sp. pl. 3. p. 131.

Icon. *Fl. dan. 930. Engl. bot. 1401.*

In dry stony fields and woods: Canada to Virginia. ♀. July, Aug. Flowers pale purple or rose-coloured.

48

1. *O. spicis* *calyce le*
Icon. *Fl. a*
In dry stor
♀. July

488. L

1. *D. spicis b*
cis dentil
oblongis
cum icor
Prasium inc
In Lower
blueish-r

2. *D. spicis el*
lycис der
lanceolati
Icon. *Bot. 1*
In mountai
July—Se
a taller va

3. *D. spicis e*
lycис dent
superne d
Prasium pur
Icon. *Bot. 1*
In the mour
Sept. v.
always sm

489.

1. *P. foliis peti*
labii inae
adscendent
Icon. *Fl. dan*
In fields and
July, Aug

2. *P. foliis peti*
labii æqu
adscendent

487. ORIGANUM. Gen. pl. 981.

1. O. spicis subrotundis paniculatis conglomeratis, bracteis calyce longioribus ovatis. Willd. sp. pl. 3. p. 135.
Icon. Fl. dan. 638.
In dry stony fields and woods: Pennsylvania to Virginia.
♀. July, Aug. v. v. Flowers pale red.

488. DRACOCEPHALUM. Gen. pl. 984.

1. D. spicis brevibus tetragonis, bracteis ovatis acutis, calycis dentibus inæqualibus, foliis arcte sessilibus ovali-oblongis superne denticulatis. Vent. hort. cels t. 44. *variegatum*.
Prasium incarnatum. Walt. fl. car. 165.
In Lower Carolina. ♀. Aug. Sept. v. v. Flowers blueish-red, with white stripes, very ornamental.
2. D. spicis elongatis confertifloris, bracteis subulatis, calycis dentibus brevibus subæqualibus, foliis linear-lanceolatis serratis.—Willd. sp. pl. 3. p. 149. *virginianum*.
Icon. Bot. mag. 467. Rob. ic. 207.
In mountain meadows: Virginia and Carolina. ♀.
July—Sept. v. v. Flowers reddish-purple. There is a taller variety with dark purple flowers.
3. D. spicis elongatis remotifloris, bracteis subulatis, calycis dentibus subæqualibus, foliis obovato-lanceolatis superne denticulatis.—Willd. sp. pl. 3. p. 150. *denticulatum*.
Prasium purpureum. Walt. fl. car. 166.
Icon. Bot. mag. 214.
In the mountains of Virginia and Carolina. ♀. July—
Sept. v. v. Resembles the former very much; it is always smaller in size, and weakly in stem.

489. PRUNELLA. Gen. pl. 990.

1. P. foliis petiolatis oblongo-ovatis basi dentatis, calycis labii inæqualibus, superiore truncato aristato, caule adscendente.—Willd. sp. pl. 3. p. 176.
Icon. Fl. dan. 910. Engl. bot. 961.
In fields and meadows: New England to Carolina. ♀.
July, Aug. v. v. Flowers violet purple.
2. P. foliis petiolatis ovato-lanceolatis basi dentatis, calycis *pensylvanica*. labii æqualibus, superiore truncato triaristato, caule adscendente. Willd. hort. berol. p. 9. t. 9. *vulgaris*.

P. laciniata. Walt. fl. car. 163.

P. sylvestris autumnalis, &c. Gron. fl. virg. 91.

In woods and meadows: Pennsylvania to Carolina. ♂.

July, Aug. v. v. Resembles the foregoing species; the spikes are longer, and the flowers pale blue.

490. SCUTELLARIA. Gen. pl. 989.

lateriflora.

1. *S. ramosissima*, glabriuscula; foliis longissime petiolatis ovatis dentatis, caulinis subcordatis, racemis lateraliibus foliosis.—*Willd. sp. pl. 3. p. 172.*

On the side of ditches and ponds: Canada to Carolina. ♀. July—Sept. v. v. Flowers blue, small.

nervosa.

2. *S. simpliciuscula*, glabra; foliis sessilibus ovatis dentatis nervosis, racemo terminali laxo folioso.

On the banks of rivulets: Virginia. ♀. July, Aug. v. v. Flowers, as the most of this genus, blue, larger than the foregoing species.

galericulata.

3. *S. simpliciuscula*, glabra; foliis subsessilibus ovato-lanceolatis basi subcordatis crenatis, floribus axillaribus solitariis.—*Willd. sp. pl. 3. p. 173.*

Icon. *Fl. dan. 637. Engl. bot. 523.*

In Canada and in the salt-marshes of Onondago, New York. ♀. July, Aug. v. v. Flowers large, pale blue.

angustifolia.

4. *S. simplex*, tenuissime pubescens; foliis linearibus, floribus axillaribus oppositis, genitalibus subexertis.

On the river Kooskoosky. *M. Lewis.* ♀. June. v. s. in *Herb. Lewis.*

parvula.

5. *S. simplex*, pusilla, dense pubescens; foliis sessilibus ovatis integerrimis, omnibus conformibus, floribus axillaribus solitariis.—*Mich. fl. amer. 2. p. 11.*

In Canada and the Illinois country. *Michaux.* On banks of rivers in Virginia. *Ph. ♂.* June, July. v. v. Not above two inches high; flowers small, pale blue.

caroliniana.

6. *S. ramosa*, glaberrima; foliis petiolatis linear-lanceolatis acutis integerrimis, racemis laxis foliosis, calycibus obtusis.—*Lam. encycl. 7. p. 706.*

Icon. *Lam. illustr. t. 515. f. 3.*

In Carolina. *Fraser.* ♀. +.

integrifolia.

7. *S. simpliciuscula*, dense pubescens; foliis subsessilibus oblongis linearibus obtusis integerrimis basi attenuatis, racemis laxiusculis foliosis.—*Willd. sp. pl. 3. p. 173.*

Icon. *Pluk.*

♂. *S. foliis or-*

On dry hill

July—Se-

species i

simple ai

♀. *S. ramosa,*

ratis brev-

plerumq;

Bot. rep.

S. integriflo-

In fields an-

—Sept.

handsom

9. *S. simpliciu-*

ovatis ob-

lis brevib-

mosis, br-

p. 11.

S. ovalifolia

S. carolinian-

Icon. Pluk.

In shady wo-

v. v. Re-

plant, wi

1. *T. floribus*

obtusis ba-

Icon. Engl.

Common T

state, par-

♀. h.

1. *C. pubescen-*

ratis, ver-

folio longi-

Melissa Nep-

Thymus Ne-

Icon. Engl.

Icon. *Pluk. alm.* 441. f. 6.

8. *S. foliis omnibus linearibus*.—*Willd. sp. pl.* 3. p. 174. *hyssopifolia*.

On dry hills, in rich soil : New York to Carolina. ♀.

July—Sept. v. v. Flowers large, handsome blue. This species is very variable, in size and figure of leaves, simple and branched stems, &c.

8. *S. ramosa*, elata, pubescens ; foliis ovatis acuminatis serratis breviter petiolatis, racemis terminalibus laxiusculis plerumque paniculatis, bracteis lanceolatis brevibus.—*Bot. rep.* 494.

S. integrifolia. *Auctorum*.

In fields and meadows : Virginia and Carolina. ♀. July—Sept. v. v. The tallest species ; flowers large and handsome.

9. *S. simpliciuscula*, pubescens ; foliis remotis rhomboideo-ovatis obtusis rotundato-crenatis basi attenuatis, petiolis brevibus, racemis terminalibus laxis plerumque ramosis, bracteis lanceolatis integris.—*Mich. fl. amer.* 2. p. 11.

S. ovalifolia. *Pers. syn.* 2. p. 136?

S. caroliniana. *Walt. fl. car.* 163.

Icon. *Pluk. alm.* t. 313. f. 4.

In shady woods : Virginia and Carolina. ♀. July, Aug. v. v. Resembles the foregoing one ; it is a smaller plant, with only a few leaves.

serrata.

pilosa.

491. THYMUS. *Gen. pl.* 982.

1. *T. floribus capitatis*, caulibus repentibus, foliis planis obtusis basi ciliatis. *Willd. sp. pl.* 3. p. 138.

Icon. *Engl. bot.* 1514. *Schkahr handb.* 144.

Common *Thyme* occurs frequently in an apparently wild state, particularly on the commons near Philadelphia. ♀. ♂. July, Aug. v. v. Flowers purple.

Serpyllum.

1. *C. pubescens*, ramosissima ; foliis ovatis obtusis subseratis, verticillis pedunculatis dichotomo-corymbosis folio longioribus, villis calycinis prominentibus.

Melissa Nepetæ. *Willd. sp. pl.* 3. p. 147.

Thymus Nepeta. *Smith fl. brit.* 2. p. 642.

Icon. *Engl. bot.* 1414. *Curt. lond.* 6. t. 40.

Nepeta.

492. CALAMINTHA. *Tourn. inst.* 193.

1. *C. pubescens*, ramosissima ; foliis ovatis obtusis subseratis, verticillis pedunculatis dichotomo-corymbosis folio longioribus, villis calycinis prominentibus.

Melissa Nepetæ. *Willd. sp. pl.* 3. p. 147.

Thymus Nepeta. *Smith fl. brit.* 2. p. 642.

Icon. *Engl. bot.* 1414. *Curt. lond.* 6. t. 40.

In lime-stone soil, on dry barren hills: Virginia. 2.
June, July. Flowers white, tinged with red.

- grandiflora.* 2. C. erecto-cespitosa; foliis rhomboideo-ovalibus superne obsolete dentatis glabriusculis, verticillis subpedunculatis subdecemfloris folio brevioribus, villis calycinis inclusis.

Thymus grandiflorus. *Bot. mag.* 997.

Thymus carolinianus. *Mich. fl. amer.* 2. p. 9.

Thymbra caroliniana. *Walt. fl. car.* 162.

On the banks of the river Savannah, in Georgia and Carolina. 2. h. July, Aug. v. v. Flowers large, pale purple.

493. HEDEOMA. *Pers. syn.* 2. p. 131.

- glabra.* 1. H. erecta, glabra; foliis inferioribus oblongis, superioribus lanceolatis remote serratis, pedunculis terminalibus ternis.—*Pers. syn.* 2. p. 131.

Cunila glabella. *Mich. fl. amer.* 1. p. 13.

On rocks on the river Tennessee, near Nashville. 2. +.

- pulegioides.* 2. H. pubescens; foliis oblongis serratis, pedunculis axillaribus verticillatis, calycis labio inferiore biseto, setis ciliatis.—*Pers. syn.* 2. p. 131.

Cunila pulegioides. *Willd. sp. pl.* 1. p. 122.

In fields and on dry hills: Canada to Carolina. 2. June, July. v. v. A small plant; flowers small, pale blue.

- hispida.* 3. H. ramosa, undique hispido-pilosa; foliis linearibus nervosis margine revoluto integrerrimis, verticillis subquadrifloris, calycibus hispidissimis lacinii subulatis. On the banks of the Missouri. 2. June, July. v. s. Not above two inches high; flowers very small.

494. TRICHOSTEMA. *Gen. pl.* 988.

- dichotoma.* 1. T. foliis rhomboideo-lanceolatis, ramulis floriferis bifurcatis, staminibus longissimis.—*Willd. sp. pl.* 3. p. 170.

- linearis.* 2. T. foliis linearibus. *Walt. fl. car.* 164.
In fields and on sunny hills: Pennsylvania to Carolina. 2. June—Aug. v. v. Flowers large, fine blue; the whole plant has an agreeable resinous scent.

2. T. foliis ovato-niculatis,
pl. 3. p. 17
Icon. *Dill. el.*
In Virginia.
ers blue.

II. A

495.

1. L. caule prost
subrotundis
Icon. *Fl. dan.*
In old fir-wooc
the mountai
tifully in th
and near th
Jersey. 2.
tinged with
little plant v
nist, should

496.

1. P. foliis ovatis
bus gracilis
pl. 179.
Icon. *Pluk. an.*
In shady, rock
—Aug. v.
Walter in
which he ca
oblongis se
plant I was
Herbarium,
urticifolia by

497.

1. V. assurgens;
corollarum la
serratis petio

2. T. foliis ovato-lanceolatis, ramulis floriferis oppositis paniculatis, staminibus brevibus inclusis.—*Willd. sp. pl. 3. p. 170.*
 Icon. *Dill. edit. t. 285. f. 369.*
 In Virginia. *Dillen. O. v. s. in Herb. Banks.* Flowers blue.

II. ANGIOSPERMIA.

495. LINNÆA. *Gen. pl. 1037.*

1. L. caule prostrato, ramulis floriferis erectis bifloris, foliis subrotundis antice crenatis.—*Willd. sp. pl. 3. p. 340.*
 Icon. *Fl. dan. 3. Engl. bot. 433. Fl. lapp. t. 12. f. 4.*
 In old fir-woods of Canada and New England, and on the mountains of New York and Pennsylvania. Plentifully in the woods near Wiscasset, Province of Main, and near the White Lake, on the mountains of New Jersey. $\frac{1}{2}$. May, June. v. v. Flowers white, tinged with red. This interesting and handsome little plant will for ever be the admiration of the botanist, should it even be only for the sake of its name.

*borealis.*496. PHRYMA. *Gen. pl. 994.*

1. P. foliis ovatis grosse serratis petiolatis, spicis terminalibus gracilibus, floribus oppositis.—*Willd. sp. pl. 3. p. 179.*
 Icon. *Pluk. amalh. t. 380. f. 5. Lam. illustr. t. 516.**
 In shady, rocky woods: Canada to Carolina. $\frac{1}{2}$. June—Aug. v. v. Flowers small, white with purple.—Walter in his Flora Caroliniana has another species which he calls *P. carolinensis*, foliis sessilibus obovato-oblongis serratis scabris, spica terminali. Of this plant I was not able to find any information in his Herbarium, and strongly suspect he means *Salvia urticifolia* by it.

497. VERBENA. *Gen. pl. 43.** *Foliis laciniatis.*

1. V. assurgens; spicis solitariis pedunculatis imbricatis, corollarum laciniis emarginatis, foliis ovalibus inciso-serratis petiolatis.—*Willd. sp. pl. 1. p. 119.*

Aubletia.

- V.** longiflora. *Lam. illustr.* 1. p. 57.
Buchnera canadensis. *Linn. mant.* 88.
Glandularia carolinensis. *Gmel. syst. nat.* 2. p. 920.
Anonymos carolinensis. *Walt. fl. car.* 164.
Icon. Jacq. hort. 2. t. 176. *Roxier journ. d'hist. nat.* 1. t. 2.
 In Carolina and Georgia. ♂. June, July. v. v. Flowers large, of a beautiful purple.
- bracteosa.** 2. *V. pumila, decumbens, hirsutissima; foliis laciniatis, spica floribus subimbricata, bracteis linearibus longissimis et undique patentibus squarrosa.* *Mich. fl. amer.* 2. p. 13.
Zapania bracteosa. *Lam. encycl.* 8. p. 843.
 In Illinois and Kentucky. ♀. July. v. s. in *Herb. Lyon.* Flowers light purple.
- spuria.** 3. *V. caule decumbente ramosissimo divaricato, foliis multifido-laciniatis, spicis filiformibus, bracteis calyces superantibus.* — *Willd. sp. pl.* 1. p. 119.
 On slate-hills and lime-stone rocks: Illinois, Virginia and Kentucky. ♂. July, Aug. v. v. Flowers very small, blue.
- hastata.** 4. *V. erecta, elatior; foliis lanceolatis acuminatis inciso-serratis nonnullis inciso-hastatis, spicis linearibus paniculatis subimbricatis.* — *Willd. sp. pl.* 1. p. 118.
Icon. Herm. parad. 242.
- pinnatifida.** β. *V. foliis inciso-pinnatifidis grosse serratis.* *Lam. illustr.* 1. p. 57.
 On road sides, in wet soil, and on the banks of rivers: Canada to Carolina. ♀. July, Aug. v. v. From two to five feet high; flowers purple, sometimes white.

** *Foliis integris.*

- paniculata.** 5. *V. erecta, scabriuscula; foliis lanceolatis grosse serratis indivisis, spicis filiformibus imbricatis corymboso-paniculatis.* — *Lam. encycl.* 8. p. 548.
 On the natural meadows of the high mountains of Virginia and Carolina. ♀. July, Aug. v. v. From four to six feet high; flowers very numerous, purple.
- urticifolia.** 6. *V. erecta, subpubescens; foliis ovatis acutis serratis petiolatis, spicis filiformibus distinctifloris axillaribus terminalibusque.* — *Willd. sp. pl.* 1. p. 119.
Icon. Riv. monop. 57. *Rob. ic.* 26.
 On road sides and cultivated grounds: New England to

Carolina
small, w

- 7.
- V. erecta,*
-
- subpubes-
-
- valde dif-
-
- In North
-
- very sma-
-
- near rese

- 8.
- V. humilis,*
-
- nuatis ra-
-
- bus solita
-
- 633.

V. angustif-
In dry soil,
Pennsylva-
Not abov

- 9.
- V. erecta,*
-
- æqualiter
-
- longissim
-
- p. 119.
-
- Icon. Dill.*

In dry sand
July. v.

- 10.
- V. hirsuta,*
-
- libus obo-
-
- spicis stri-
-
- 633.
- Ve*
-
- V. rigens.*
- A
-
- In Carolina
-
- v. s. in H

498. Z

- 1.
- Z. foliis ova-*
-
- pitato-con-
-
- herbaceo i
-
- Verbena no-*
-
- Lippia nodif*
-
- Blairia nodif*
-
- Icon. Lam.*
-
- On the coast
-
- of the Oh-
-
- small.

- 2.
- Z. foliis line-*

bena.

. 2. p. 920.

4.

i. d'hist. nat. 1.

y. v. v. Flow-

foliis laciniatis,
earibus longissi-
mich. fl. amer. 2.

3.
v. s. in Herb.

ato, foliis mul-
racteis calyces

inois, Virginia
Flowers very

minatis inciso-
nearibus pani-
p. 118.

Lam. illustr. 1.

nks of rivers :
v. From two
imes white.

grosse serratis
corymboso-pa-

ntains of Vir-
v. From four
, purple.

is serratis pe-
xillaribus ter-

w England to

DIDYNAMIA ANGIOSPERMIA. Verbena.

417

Carolina. 4. June—Sept. v. v. Flowers very small, white.

7. V. erecta, ramosissima ; foliis ovato-lanceolatis serratis subpubescentibus, spicis longissimis laxis paniculatis valde diffusis.—Lam. encycl. 8. p. 550.

In North America. Lamarck. 4. h. +. Flowers very small, somewhat purple. This species has a very near resemblance to the foregoing.

8. V. humilis, erecta ; foliis lineari-lanceolatis basi attenuatis rarer serratis lineato-venosis, spicis filiformibus solitariis terminalibus axillaribus.—Willd. enum. 633.

V. angustifolia. Mich. fl. amer. 2. p. 14.
In dry soil, on road sides, particularly in lime-stone soil : Pennsylvania to Tennessee. 4. June—Aug. v. v.
Not above a foot high ; flowers fine blue.

9. V. erecta, scabra ; foliis oblongo-ovoalibus obtusis inaequaliter serratis basi attenuatis subsessilibus, spicis longissimis filiformibus distinctifloris.—Willd. sp. 1. p. 119.

Icon. Dill. elth. t. 301. f. 388.
In dry sandy fields : Carolina to Georgia. 4. June, July. v. s. Flowers pale red.

10. V. hirsuta, albicans ; caulis rigide erectis, foliis sessilibus obovalibus serratis subomentoso-hirsutissimis, spicis strictis imbricatis subfasciculatis.—Willd. enum. 633. Vent. hort. cel. p. 53. cum iconē.

V. rigens. Mich. fl. amer. 2. p. 14.
In Carolina and the Illinois country. 4. July, Aug.
v. s. in Herb. Lambert.

diffusa.

rugosa.

caroliniana.

stricta.

498. ZAPANIA. Lam. illustr. 1. p. 58.

1. Z. foliis ovato-cuneiformibus superne serratis, spicis capitato-conicis solitariis elongato-pedunculatis, caule herbaceo repente.—Lam. illustr. 1. p. 59.

Verbena nodiflora. Willd. sp. pl. 1. p. 117.

Lippia nodiflora. Mich. fl. amer. 2. p. 15.

Blairia nodiflora. Gert. de fruct. et sem. 1. p. 266. t. 56.

Icon. Lam. illustr. 1. t. 17. f. 3. Rob. ic. 30.

On the coast of Virginia and Carolina, and on the banks of the Ohio. 4. July, Aug. v. v. Flowers white, small.

nodiflora.

2. Z. foliis linear-lanceolatis argute serratis, spicis capitato-

lanceolata.

conicis solitariis elongato-pedunculatis, caule herbaceo repente. — *Pers. syn. 2. p. 140.*

Lippia lanceolata. Mich. fl. amer. 2. p. 15.

In Carolina and Georgia, on the banks of rivers. ♀. July, Aug. v. v. Flowers white.

multifida.

1. *C. erecta, viscidio-pubescent; foliis superioribus terno-vorticillatis multifidis, pedicellis solitariis.—Mich. fl. amer. 2. p. 22. t. 36.*

On the banks of rivers, in the western parts of the settlements, particularly on the Missouri and Mississippi. ♂. July, Aug. v. s. Flowers small, greenish-white.

500. HERPESTIS. *Gært. carp. 3. p. 187. t. 214.*

Monnieria. Mich. fl. amer. 2. p. 22.

rotundifolia.

1. *H. minutum pubescens; foliis subovali-orbiculatis multinvibus, pedunculis passim oppositis folia subæquantibus.*

Monnieria rotundifolia. Mich. fl. amer. 2. p. 22.

On overflowed banks of rivers, in the Illinois and other western parts. ♀. July, Aug. v. s. Flowers pale blue, small.

amplexicaulis.

2. *H. caulibus lanatis, foliis breviter cordato-ovalibus amplexicaulibus integris obtusis, pedunculis folio breviribibus.*

Monnieria amplexicaulis. Mich. fl. amer. 2. p. 22.

In ponds and ditches: Carolina and Georgia. ♀. June — Aug. v. v. Flowers blue, larger than the other species.

cuneifolia.

3. *H. glaberrima; foliis cuneato-obovalibus superne obsolete subcrenatis, pedunculis folia subæquantibus.*

Monnieria cuneifolia. Mich. fl. amer. 2. p. 22.

On the overflowed banks of rivers, near the sea-shore: Pennsylvania to Carolina. ♀. Aug. v. v. Leaves thick; flowers very small, pale blue.

micrantha.

4. *H. glabra, succulenta; foliis arcte sessilibus ovalibus obtusis integerrimis striato-nervosis, pedunculis folio brevioribus, calyce 5-phyllo, stylo bifido.*

Gratiola repens. Willd. sp. pl. 1. p. 103?

Icon. Swartz. ic. t. 33.

DIDYN

On the bank
Pennsylvan
ers very si
dernia.

501.

1. *L. glaberrin
sessilibus,
lyce duplo
Willd. sp.*

*Gratiola inæc
Gratiola anag
Gratiola floril
fl. virg. 3.*

2. *L. foliis repa
On the bank
York to V
blue. The
Krocker, &
present one*

502. SC.

1. *S. foliis cord
inferne cili:
Willd. sp. 1*

*S. nodosa β. a
In low groun
Aug. v. i
often more*

2. *S. foliis lance
acutis, peti
In wet mead
Sept. v. v*

503.

1. *B. foliis conju
ovato-corda
muricato.
Icon. Plum. i
In shady wood
v. v. Flow*

2. *B. foliis conju
cordatis, in*

On the banks of rivers, to the edge of low water mark :
Pennsylvania to Virginia. ♂. July, Aug. v. v. Flowers very small, white. It approaches near to *Lindernia*.

501. LINDERNIA. *Gen. pl.* 1031.

1. *L. glaberrima*; foliis oblongo-ovatis obsolete dentatis sessilibus, pedunculis axillaribus unifloris, corollis calyce duplo longioribus, capsulis calyce brevioribus.—*Willd. sp. pl.* 3. p. 325?

Gratiola inaequalis, *Walt. fl. car.* 61. ^{ref}

Gratiola anagalloidea. *Mich. fl. amer.* 1. p. 6.

Gratiola floribus pedunculatis, foliis ovatis crenatis. *Gron. fl. virg.* 3.

2. *L. foliis repando-dentatis, pedunculis brevibus*.

On the banks of ponds and rivers, in gravelly soil : New York to Virginia. ♂. July, Aug. v. v. Flowers blue. The European plant, figured by Schkuhr and Krockier, seems to be a very different species from the present one, as the corolla is shorter than the calyx.

pyridaria.

major.

502. SCROPHULARIA. *Gen. pl.* 1014.

1. *S. foliis cordatis serratis acutis basi rotundatis, petiolis inferne ciliatis, paniculæ fasciculis laxè paucifloris*.—*Willd. sp. pl.* 3. p. 269.

S. nodosa β, *americana*. *Mich. fl. amer.* 2. p. 21.

In low grounds : Pennsylvania to Carolina. ♀. June—Aug. v. v. Flowers greenish-brown. The plant often more than four feet high.

2. *S. foliis lanceolatis inaequaliter serratis acuminatis basi acutis, petiolis nudis, paniculæ fasciculis corymbosis*.

In wet meadows and woods : Pennsylvania. ♀. Aug. Sept. v. v. Flowers greenish-yellow.

lanceolata.

503. BIGNONIA. *Gen. pl.* 1018.

1. *B. foliis conjugatis cirrhosis, inferioribus ternatis, foliolis ovato-cordatis acuminatis, racemis axillaribus, caule muricato*.—*Willd. sp. pl.* 3. p. 296.

Icon. Plum. ic. t. 58.

In shady woods : Carolina to Florida. ♂. June, July. v. v. Flowers yellowish-scarlet.

crucigera.

2. *B. foliis conjugatis cirrhosis, foliolis oblongis basi subcordatis, imis simplicibus*.—*Willd. sp. pl.* 3. p. 297.

capreolata.

Icon. *Bot. mag.* 864. *Jacq. hort. schoenbr.* 363.
In shady woods, on the banks of rivers and lakes: Virginia to Florida. $\frac{1}{2}$. July, Aug. v. v. Flowers scarlet.

radicans. 3. B. foliis pinnatis, foliolis ovatis dentatis acuminatis, corymbo terminali, tubo corollæ calyce triplo longiore, caule radicante.—*Willd. sp. pl.* 3. p. 301.

Tecoma radicans. Juss.
flammea. a. B. floribus luteo-coccineis.
Icon. *Schmidt arb.* t. 42.
coccinea. b. B. floribus coccineis.
Icon. *Schmidt arb.* t. 43 & 44. *Bot. mag.* 485. *Catesb. car.* 1. t. 65.

On the banks of rivers: Pennsylvania to Florida. $\frac{1}{2}$. July, Aug. v. v. This highly ornamental creeper ascends the tallest trees and highest rocks, and with its large scarlet flowers produces one of the most pleasing effects in ornamental gardening.

504. RUELLIA. *Gen. pl.* 1050.

strepsis. 1. R. erecta, hirsuta; foliis petiolatis lanceolato-ovatis integrerimis, pedunculis 1—3-floris, calycis laciniis lanceolatis hispidis tubo corollæ duplo brevioribus.—*Willd. sp. pl.* 3. p. 363.

Icon. *Schkuhr handb.* 177. *Dill. eth.* t. 249, f. 321.
On dry hills, in shady woods: Virginia to Carolina. $\frac{1}{2}$. June, July. v. v. Flowers large, pale blue, as are all the following.

oblongifolia. 2. R. assurgens, dense pubescens; foliis subsessilibus oblongato-oblongis, floribus solitariis, bracteis longitudine calycis, calycis laciniis filiformibus longitudine tubi corollæ.—*Mich. fl. amer.* 2. p. 23.

R. biflora. *Willd. sp. pl.* 3. p. 368?
In sandy pine-woods: Georgia. $\frac{1}{2}$. June, July. v. s.
Flowers yellowish-blue.

hybrida. 3. R. erecta, ramosissima, pilis albidis hirsuta; foliis subsessilibus oblongis utrinque subacutis dense hirsutis, bracteis calycé brevioribus, calycis laciniis linearibus tubo corollæ vix brevioribus.

In sandy fields near Savannah, Georgia. $\frac{1}{2}$. July. v. s.
in *Herb. Enslen.*

ciliosa. 4. R. erecta, ramosa; foliis subsessilibus ovato-oblongis margine nervis venisque pilis albis longe ciliatis, brac-

DIDYN

teis lanceolæ
corollæ qua
Near Savann
Enslen.

5. R. glabriusculi
giuscule ang
libus, capsu
In Georgia and

505. B

1. B. caule simpl
trinervibus,
p. 334.
On dry sandy l
July. v. v.

506. AN

1. A. procumbens
rimis, pedu
pl. 3. p. 234.
Icon. *Fl. dan.*
In sandy field
Aug. v. v.

2. A. erectum, gl
bus conferti
bus glabris
p. 253.

Icon. *Fl. dan.* g
In fields and dr
—Aug. v. v.
looking plant
some weeds in

3. A. assurgens, |
erectis angus
cemosis, stolo
p. 255.

Icon. *Vent. hort*
In low grounds :
v. v. Flowe

4. A. pusillum, sin
acutis, florib
bus campanul

teis lanceolatis brevibus, calycis laciniis subulatis tubo
corollæ quadruplo brevioribus.
Near Savannah, Georgia. 4. July. v. s. in *Herb.*
Enslen.

5. *R. glabriuscula*, diffusa, radicans; foliis in petiolum longo-
guscule angustatis ovalibus obtusis, floribus subsessi-
bus, capsulis linearibus. *Mich. fl. amer.* 2. p. 23.
In Georgia and Florida. *Michaux.* +.

505. BUCHNERA. *Gen. pl.* 1035.

1. *B. caule simplici*, foliis lanceolatis subdentatis asperis
trinervibus, spicis remotifloris.—*Willd. sp. pl.* 3.
p. 334.
On dry sandy hills: Pennsylvania to Carolina; rare. 4.
July. v. v. Flowers deep blue.

506. ANTIRRHINUM. *Gen. pl.* 1007.

1. *A. procumbens*, pilosum; foliis alternis hastatis integer-
rimis, pedunculis solitariis longissimis.—*Willd. sp.*
pl. 3. p. 234.
Icon. *Fl. dan.* 426. *Engl. bot.* 692.
In sandy fields: Pennsylvania to Virginia. ①. July,
Aug. v. v. Flowers yellow mixed with purple.
2. *A. erectum*, glabrum; foliis sparsis lanceolato-linearibus
confertis, spicis terminalibus densifloris, calyci-
bus glabris calcare brevioribus.—*Willd. sp. pl.* 3.
p. 253.
Icon. *Fl. dan.* 982. *Engl. bot.* 658.
In fields and dry hills, near houses; frequent. 4. June—
Aug. v. v. Flowers large, yellow. This fine-
looking plant is one of the worst and most trouble-
some weeds in several parts of Pennsylvania and Virginia.
3. *A. assurgens*, glabrum, simplicissimum; foliis sparsis
erectis angusto-linearibus obtusis remotis, floribus ra-
cemosis, stolonibus procumbentibus.—*Willd. sp. pl.* 3.
p. 255.
Icon. *Vent. hort. cels.* 49.
In low grounds: Canada to Carolina. ①. June—Aug.
v. v. Flowers small, purple.
4. *A. pusillum*, simplex, glabrum; foliis oppositis linearibus
acutis, floribus axillaribus brevi-pedunculatis, calyci-
bus campanulatis.

*americana.**Elatine.**Linaria.**canadense.**tenellum.*

On the banks of the Missouri. *M. Lewis.* ♂. July. v. v.; v. s. in *Herb. Lewis.* Flowers bright blue, large in proportion to the size of the plant.

Crontium.

5. *A. erectum*, ramosum, pubescens; foliis alternis lanceolatis, floribus subspicatis, calycibus digitatis colla longioribus, corollis ecaudatis.—*Willd. sp. pl. 3. p. 258.*

Icon. *Curt. fl. lond. 4. t. 45.* Riv. t. 82.

In fields: Virginia; rare. ♂. July, Aug. v. v. Flowers pale purple.

507. GERARDIA. *Gen. pl. 1004.** *Flores purpurei.**purpurea.*

1. *G. caule* opposite ramosissimo, foliis linearibus, floribus axillaribus oppositis subsessilibus.—*Willd. sp. pl. 3. p. 221.*

Icon. *Pluk. mant. t. 388. f. 1.*

crassifolia.

- β. *G. foliis* carnosis brevioribus.
In fields and woods: New England to Carolina. β. In salt-marshes, near New York, &c. ♂. July, Aug. v. v.

tenuifolia.

2. *G. paniculato-ramosa*; foliis linearibus, pedunculis axillaribus oppositis flore longioribus.—*Willd. sp. pl. 3. p. 222.*

G. erecta. *Walt. fl. car. 170.* *Mich. fl. amer. 2. p. 20.*
Icon. *Pluk. alm. t. 12. f. 4.*

In fields and woods: Pennsylvania to Carolina. ♂. July, Aug. v. v.

setacea.

3. *G. caule* ramosissimo, foliis setaceis, ramulis axillaribus folio longioribus plerumque unifloris.—*Walt. fl. car. 170.*

In sandy fields: Pennsylvania to Virginia. ♂. Aug. v. v.
Resembles the foregoing one; but the flowers are not axillary but terminal, very seldom two on a branch.

auriculata.

4. *G. subsimplex*, tota aspera; foliis ovato-lanceolatis basi 2-auriculatis integerrimis, floribus axillaribus oppositis folisque arcte sessilibus.—*Mich. fl. amer. 2. p. 20.*

In dry stony fields: Virginia and Illinois. ♂. July, Aug. v. v. Not above eight inches high, without branches.

cuneifolia.

5. *G. paniculato-ramosa*; ramis erectis, foliis cuneato-lanceolatis inaequilateri serratis superioribus alternis,

num.

s. ♂. July,
bright blue,
it.

alternis lan-
digitatis co-
lld. sp. pl. 3.

v. Flowers

bus, floribus
t. sp. pl. 3.

olina. β. In
July, Aug.

dunculis ax-
ld. sp. pl. 3.

r. 2. p. 20.

a. ♂. July,

is axillaribus
Valt. fl. car.

. Aug. v. v.
wers are not
in a branch.
ceolatis basi-
ribus oppo-
!. amer. 2.

, July, Aug.
at branches.

is cuneato-
bus alternis,



Gerardia fruticosa.



Gerardia quercifolia.

Drawn & Engraved

DIDYNAMI

pedunculis axi-
partitis.

In Georgia. Ba-

6. *G. ramosissima* ;
subpetiolatis,
bracteatis.

In great abundan-
mountains. A
Lewis.

Fruter elegans 3-
dense foliosi, 1
cialia, opposit-
cum mucrone,
petiolum atten-
cemi terminale
oppositi, pedicel-
purplei. Brad-
integerrimæ,
funde 5-partitu-
latis, acutis.

Corollæ: Tubu
tricosus. Lim-
ciniis rotundati
corollæ inserta.
lus longitudine

7. *G. pubescens*; cau-
lanceolatis integ-
pinnatifido-incis-
sessilibus.—*Wi*

Icon. *Pluk. mant*
In dry shady and
da. 4. July—
all the following
two feet high.

8. *G. glabra*; caule
laciiniis lanceola-
dentatis, summu-
bus oppositis pec-

Rhinanthus virginicus
Icon. *Pluk. alm.*

On the banks of ri-
to Carolina. 4
five feet high.

VOL. II.

pedunculis axillaribus folio longioribus, calycibus 5-partitis.

In Georgia. *Bartram.* v. s. in *Herb. Banks.*

6. G. ramosissima; foliis lanceolatis serratis basi attenuatis subpetiolatis, racemis terminalibus, floribus oppositis bracteatis.

In great abundance in the pine-forests of the Rocky-mountain. *M. Lewis.* 2. June. v. s. in *Herb. Lewis.*

Fruter elegans 3-4-pedalis ramosissimus. *Rami* teretes, dense foliosi, tenuissime pubescentes. *Folia* semuncialia, opposita, lanceolata, acuta, interdum obtusa cum mucrone, superne acute serrata, inferne quasi in petiolum attenuata, integerrima, utrinque glabra. *Racemi* terminales, simplices, erecti, bracteati. *Flores* oppositi, pedicellati, magnitudine et colore *Digitalis purpurei*. *Bracteæ* oppositæ, sessiles, oblongæ, acutæ, integerrimæ, pedicellis vix breviores. *Calyx* profunde 5-partitus, basi unibracteolatus: lacinia lanceolatis, acutis. *Bracteolæ* lineares, calyce breviores. *Corolla*: *Tubus* calyce quadruplo longior, superne ventricosus. *Limbus* patens, subæqualis, 5-fidus: *Lacinia* rotundatis. *Filamenta* tubo duplo breviora, imo corollæ inserta. *Antheræ* oblongæ, hirsutissimæ. *Stylus* longitudine tubi. *Capsula*.....

** *Flores flavi.*

7. G. pubescens; caulis subsimplicibus, foliis subsessilibus lanceolatis integerrimis vel dentatis, inferioribus subpinnatifido-incisis, floribus axillaris oppositis subsessilibus.—*Willd. sp. pl. 3. p. 223.*

Icon. *Pluk. mant. t. 389. f. 3.*

In dry shady and rocky woods: New England to Florida. 2. July—Sept. v. v. Flowers large, and, as all the following, yellow. It is seldom more than two feet high.

8. G. glabra; caule paniculato, foliis petiolatis pinnatifidis: lacinia lanceolatis acutis integris vel postice incisodentatis, summis lanceolatis integris, floribus axillariis oppositis pedicellatis.

Rhinanthus virginicus. *Willd. sp. pl. 3. p. 191.*

Icon. *Pluk. aln. t. 389. f. 1.*

On the banks of rivers, in rich shady places: Pennsylvania to Carolina. 2. July, Aug. v. v. From three to five feet high. This fine species has been generally

fruticosa.

flava.

quercifolia.

confounded with the foregoing one, but differs widely from it. The stem is smooth, purple, with a blue dew over it, and very much branched. The flowers are nearly the size, shape, and colour of the preceding species. The original specimen of Gronovius of *Rhinanthus virginicus*, in the Banksian Museum, is evidently our present plant.

- Pedicularia.* 9. *G. pubescens*, brachiatim-paniculata; foliis oblongis duplicato-inciso-serratis pinnatifidisque, floribus axillaribus oppositis pedicellatis, calycis lacinias foliaceis inciso-dentatis.—*Willd. sp. pl. 3. p. 223.*

Icon. *Lam. illustr. t. 529. f. 2.*

In dry shady woods: New York to Carolina. ♂. June—Aug. v. v. A very handsome species, of which there are several very striking varieties.

- cassiodes.* 10. *G. virgatim-paniculata*; foliis setaceo-pinnatifidis: lacinias linearibus.

G. Afzelia. Mich. fl. amer. 2. p. 20.

Afzelia cassiodes. Gmel. syst. nat. 927.

Anonymos cassiodes. Walt. fl. car. 171.

In dry sandy woods of Carolina and Georgia. ♂. July, Aug. v. s. Flowers small, yellow.

508. PEDICULARIS. Gen. pl. 1003.

- euphrasioides.* 1. *P. caule ramoso pubescente*, foliis pinnatifidis dentatis, calycibus tubulosis bipartitis truncatis, corollæ galea bidentata. *Willd. sp. pl. 3. p. 294.*

Icon. *Houttuyn. linn. syst. 8. t. 57.*

In Labrador. *Colmaster. Ʌ. v. s. in Herb. Dickson.* Flowers yellow; from six to eight inches high.

- pallida.* 2. *P. caule ramoso elato glabro*, foliis suboppositis lanceolatis crenatis dentatis, spica aphylla glabra, corollæ galea obtusa, calyce bifido rotundato.—*Herb. Banks.*

P. virginica. Lam. encycl. 5. p. 126.

In swamps, on lime-stone soil: Virginia. *Lamarck.* In a swamp near King's-bridge, New York. *Ph. Ʌ.* July, Aug. v. v.; v. s. in *Herb. Banks.* Flowers yellow; grows upwards of three feet high. I considered this species to be the *Rhinanthus virginicus*, which Linnaeus had adopted from Gronovius's Flora Virginica; but on examining the Gronovian Herbarium in the Banksian Museum, I found his *Rhinanthus* to be the *Gerardia quercijolia*.

- lanceolata.* 3. *P. caule simplici*, foliis lanceolatis inciso-dentatis, spica

DIDYNA

aphylla, ca
fl. amer. 2

In the Illinois

4. *P. caule simplici*, spica folios
nem gladiat
p. 18.

In rich soil, i
ginia. Ʌ.
with purple

5. *P. caule simplici*, pilulo basi t
tata, calycit
p. 211.

In woods and
—July. v. v
white, with

6. *P. caule simplici*, natis, calyci
Willd. sp. p

Icon. *Gmel. fl.*
In Canada. Ʌ
ers deep pur

7. *P. caule simplici*, colatis pinn
calycibus gla
sissima. *W*

Icon. *Jacq. fl.*
On the north-
throughout C
Flowers purp

8. *P. caule simplici*, neari-lanceolat
bus glabris 5.
Willd. sp. pl

In low plains, o
Ʌ. July. v
flowers purpl

9. *P. caule simplici*, lycibus bifidis
ta. *Willd. s.*

Icon. *Fl. dan. 2*
In Labrador. *Dickson. Fl.*

- aphylla, calycibus glabris, capsulis brevibus. *Mich.*
fl. amer. 2. p. 18.
- In the Illinois country. *Michaux.* ♀.
4. *P. caule simplici, foliis lanceolatis pinnatifidis dentatis, spica foliosa alterniflora hirsuta, capsulis in mucronem gladiatum exsertum protractis.*—*Mich. fl. amer. 3. p. 18.*
- In rich soil, in grassy moist places: Pennsylvania to Virginia. ♀. May, June. v. v. Flowers yellow, tinged with purple; sometimes a foot high.
5. *P. caule simplici, foliis pinnatifidis inciso-dentatis, capitulo basi folioso hirsuto, corollis galea setaceo-bidentata, calycibus deorsum truncatis.*—*Willd. sp. pl. 3. p. 211.*
- In woods and meadows: Canada to Carolina. ♀. May—July. v. v. About a span high; flowers yellowish-white, with a tinge of purple.
6. *P. caule simplici glabro, foliis lanceolatis dentatis crenatis, calycibus bifidis truncatis, corollæ galea acuta.*—*Willd. sp. pl. 3. p. 206.*
- Icon. *Gmel. fl. sib. 3. t. 44.*
- In Canada. ♀. July. v. s. About a foot high; flowers deep purple.
7. *P. caule simplici, foliis profunde pinnatifidis, pinnis lanceolatis pinnatifidis dentatis, spica compacta foliosa, calycibus glabris quinquedentatis, corollæ galea obtusissima.*—*Willd. sp. pl. 3. p. 210.*
- Icon. *Jacq. fl. austr. 258.*
- On the north-west coast. A. *Menzies*; and probably throughout Canada. ♀. July. v. s. in *Herb. Banks.* Flowers purple.
8. *P. caule simplici, foliis profunde pinnatifidis, pinnis linear-lanceolatis crenatis, spica laxa subfoliosa, calycibus glabris 5-dentatis, corollæ galea obtusa truncata.*—*Willd. sp. pl. 3. p. 210.*
- In low plains, on the waters of Clark's river. M. *Lewis.* ♀. July. v. s. in *Herb. Lewis.* Near two feet high; flowers purple.
9. *P. caule simplici, foliis pinnatifidis, pinnis serratis, calycibus bifidis oblongis, corollæ galea uncinata truncata.*—*Willd. sp. pl. 3. p. 207.*
- Icon. *Fl. dan. 2. Fl. lapp. t. 4. f. 1.*
- In Labrador. Colmaster. ♀. July. v. s. in *Herb. Dickson.* Flowers yellow.

grænlandica. 10. P. caule simplici, foliis profunde pinnatifidis, pinnis lanceolatis acuminatis serratis, calycibus glabris 5-dentatis, corollis clausis, galea uncinato-subulata. *Willd. sp. pl. 3. p. 212.*

Icon. *Fl. dan. 1166. Retz. obs. 4. t. 1.*

In low plains of the Columbia. *M. Lewis.* 4. July.
v. s. in Herb. Lewis. nec non Lambert. Flowers purple.

verticillata. 11. P. caule simplici, foliis caulinis profunde pinnatifidis quaternis, pinnis linearibus acute dentatis, spica capitata involucrata, calycibus villosis 5-fidis, corollæ galea obtusissima.

P. verticillata β. *Willd. sp. pl. 3. p. 214.*

P. lanata. Pallas in Herb. Lambert.

On the north-west coast. *Nelson.* 4. v. s. in *Herb. Banks. nec non Lambert.* Not above two inches high; flowers purple.

flammea. 12. P. caule simplici, foliis pinnatis, pinnis imbricatis ovatis obtusis duplicito-dentatis, calycibus 5-dentatis, corollæ galea obtusa. *Willd. sp. pl. 3. p. 215.*

Icon. *Fl. dan. 30. Fl. lapp. t. 4. f. 2. Hall. fl. helv. t. 8. f. 3.*

In Labrador. *Colmaster.* 4. v. s. in *Herb. Dickson.* Not above three inches high; flowers yellow, with a scarlet helmet.

509. MIMULUS. *Gen. pl. 1049.*

ringens.

1. *M. erectus, glaber; foliis sessilibus lanceolatis acuminatis serratis, pedunculis axillaribus oppositis flore longioribus, dentibus calycis oblongis acuminatis.* — *Willd. sp. pl. 3. p. 360.*

Icon. *Bot. mag. 283. Pluk. amalthe. t. 393. f. 3.*

In low grounds, on clay soil: Pennsylvania to Carolina. 4. July, Aug. v. v. Flowers pale blue, large.

alatus.

2. *M. erectus, glaber; foliis petiolatis ovatis acuminatis serratis, pedunculis axillaribus oppositis flore brevioribus, dentibus calycis rotundatis mucronatis, caule tetragono alato.* — *Willd. sp. pl. 3. p. 361.*

In ditches and on the borders of ponds: Pennsylvania to Carolina. 4. July, Aug. v. v. Flowers the size and colour of the foregoing.

luteus.

3. *M. erectus, glaber, stoloniferus; foliis subrotundo-ovatis nervosis dentatis, inferioribus obtusis petiolatis, superioribus acutis arcte sessilibus, racemo terminali,*

5.

pinnis lan-
5-dentatis,
Willd., sp.

24. July.
Flowers

innatifidis
pica capi-
ollæ galea

b. Banks.
; flowers

atis ovatis
s, corollæ

fl. helv.


son. Not
h a scar-

icumina-
dore lon-
—*Willd.*

Carolina.
ge.
uminatis
bréviori-
caule te-

vania to
the size

do-ova-
itis, su-
rminali,



Mimulus leucostoma

Scoparia 1819

DIDYNAN

pedunculis a
quali, dentit
Icon. Feuill. p
On the banks
north-west c
Herb. Lewis.
low. It has
and will be
plants.

4. *M. erectus*, pu
lanceolatis ac
ribus paucis te
tibus calycis a
On the head spr
age hill. *M*
Lewis. Not
three, larger t
iful pale purp

510. Cl

1. *C. foliis oppositi*
spicis termin
p. 225.

2. *C. foliis subsessil*
 β . *C. foliis brevi-pe*
C. obliqua. *Will*
C. purpurea. *Mi*
Icon. Schkuhr hu
t. 348. f. 3.

On the banks of
gland to Caroli
tains of Virgin
Flowers large,
in var. β . I h
stinctions betw
consider them

511. PENT

1. *P. caule glabro*, f
plexicaulibus te
tegerrimis, flori
perne barbato.—
Chelone Penstemo
Icon. Lam. illustr.
s. 5. t. 8. f. 6.



Lupinus villosus.

DIDYNAMIA ANGIOSPERMIA. *Mimulus.*

427

pedunculis axillaribus oppositis elongatis, calyce inaequali, dentibus acutiusculis.—*Willd. sp. pl. 3. p. 361.*

Icon. Feuill. peruv. 2. t. 34.

On the banks of Clark's river. *M. Lewis.* On the north-west coast. *Pallas.* $\frac{1}{4}$. July, Aug. v. s. in *Herb. Lewis. nec non Lambert.* Flowers golden-yellow. It has lately been introduced into the gardens, and will be a fine addition to our hardy perennial plants.

4. *M. erectus*, pusillus, pubescens; foliis sessilibus oblongo-lanceolatis acutis nervosis mucronato-denticulatis, floribus paucis terminalibus longissime pedunculatis, dentibus calycis acuminatis.

On the head springs of the Missouri, at the foot of Portage hill. *M. Lewis.* $\frac{1}{4}$. Aug. v. s. in *Herb. Lewis.* Not above eight inches high; flowers two or three, larger than any other known species, of a beautiful pale purple.

510. CHELONE. *Gen. pl. 1005.*

1. *C. foliis oppositis lanceolato-oblongis acuminatis serratis, spicis terminalibus densifloris.*—*Willd. sp. pl. 3. p. 225.*

glabra.

α . *C. foliis subsessilibus, floribus albis.*

alba.

β . *C. foliis brevi-petiolatis, floribus purpureis.*

purpurea.

C. obliqua. Willd. sp. pl. 3. p. 225.

C. purpurea. Mill. dict. n. 2.

Icon. Schkuhr handb. 172. Mill. ic. 93. Pluk. mant. t. 348. f. 3.

On the banks of rivers, ponds, and swamps: New England to Carolina. β In rivulets on the high mountains of Virginia and Carolina. $\frac{1}{4}$. July, Aug. v. v. Flowers large, in close spikes, in var. α . white; purple in var. β . I have not been able to find sufficient distinctions between *C. glabra* and *C. obliqua*, so as to consider them distinct species.

511. PENTSTEMON. *Gen. pl. 1758.*

1. *P. caule glabro, foliis lœvigatis ovato-oblongis basi amplexicaulibus tenuissime denticulatis, interioribus integerrimis, floribus paniculatis, filamento sterili superne barbato.*—*Willd. sp. pl. 3. p. 228.*

lœvigata.

Chelone Pentstemon. J. F. Miller. ic. 4.

Icon. Lam. illustr. 528. Arduin. spec. t. 5. Moris. hist. 2. s. 5. t. 8. f. 6.

In stony low grounds: Pennsylvania to Carolina. ♀. July—Sept. v. v. From one to two feet high; flowers pale purple.

pubescens.

2. *P. caule pubescente, foliis serrulatis lanceolato-oblongis sessilibus, floribus paniculatis, filamento sterili ab apice infra medietatem barbato.* — *Willd. sp. pl. 3. p. 227.*

Chelone Pentstemon. Mant. 415.

latifolia.

- α. *P. foliis laevigatis serratis latioribus.*

Icon. Moris. hist. 3. s. 11. t. 21. f. 2.

angustifolia.

- β. *P. foliis pubescensibus obscure denticulatis angustis.*

P. hirsuta. Willd. sp. pl. 3. p. 227.

Chelone hirsuta. Mill. dict. n. 3.

Icon. Moris. hist. 3. s. 11. t. 21. f. 3.

On the side of hills and in dry meadows: New York to Virginia. ♀. July—Sept. v. v. About a foot high; flowers pale purple, longer than the foregoing species.

frutescens.

3. *P. caule fruticoso, ramis angulatis superne pubescensibus, foliis lanceolatis obsolete denticulatis sessilibus glabriusculis, racemis terminalibus subcorymbosis, filamento sterili longitudinaliter barbato.* — *Lambert in Linn. trans. 10. p. 259.*

Icon. Linn. trans. 10. t. 6.

On the north-west coast. *M. Lewis.* Unalashka. *Pal-las.* ½. v. s. in *Herb. Lambert.* nec non *Lewis.* A small shrub, about a foot or more high; flowers purple.

512. MARTYNIA. Gen. pl. 1010.

Proboscidea.

1. *M. caule ramoso, foliis alternis cordatis integerrimis villosis.* — *Willd. sp. pl. 3. p. 264.*

- 2. *M. alternifolia. Lam. encycl. 2. p. 112.*

M. australis. Sp. pl. 862.

Icon. Bot. mag. 1056. Mill. ic. 286.

On the banks of the Mississippi. ♂. July. v. v. Flowers large, white, speckled with purple and yellow. The singular shape of its fruit, connected with the beauty of its flowers, makes this annual a desirable plant in all collections.

513. SCHWALBEA. Gen. pl. 1001.

americana.

1. *S. simplex, pubescens; foliis lanceolatis, racemo terminali, floribus alternis.* — *Willd. sp. pl. 3. p. 201.*

Icon. Lam. illustr. 520. Pluk. mant. Y. 348. f. 2.

In pine-barrens: New Jersey to Carolina. ♀. ♂. June, July. v. v. Flowers yellowish-brown.

1. *R. corollarum*

cinia medi-

p. 188.

Icon. Schkuh.

In fields and

v. v. Flow-

& Pedicula-

1. *B. foliis alter-*

latis, florali-

nibus 3-ner-

pl. 3. p. 180

Icon. Gmel. f.

In Canada and

Hampshire.

Lambert. Fl.

2. *B. foliis alter-*

longissime a-

floribus brac-

On the north

Herb. Lamb.

3. *B. hirsutissima*

dis: laciniis

longis obtusis

longioribus, c-

latis.

On the banks o-

v. s. in Herb

deep yellow;

4. *B. foliis alternis*

nearibus, bra-

longioribus,

Willd. sp. pl.

Icon. Pluk. alm.

f. 28.

- α. *B. bracteis cocci-*

- β. *B. bracteis lutesc-*

In wet meadows

July. v. v.

leaves, which

pearance.

514. RHINANTHUS. *Gen. pl. 997.*

1. *R. corollarum* labio superiore emarginato bidentato, lacinia media labii inferioris brevissima. *Willd. sp. pl. 3.*
p. 188.

Icon. *Schkuhr handb. 169. Fl. dan. 981. Engl. bot. 657.*
 In fields and meadows: Lower Canada. ♂. June, July.
v. v. Flowers yellow. *R. virginicus.* Vid. *Gerardia*
 & *Pedicularis pallida.*

515. BARTSIA. *Gen. pl. 996.*

1. *B. foliis alternis linearibus indivisis, superioribus lanceolatis, floralibus subovalibus summitate subdentatis, omnibus 3-nervibus, dentibus calycis acutis.* — *Willd. sp. pl. 3. p. 186. Mich. fl. amer. 2. p. 17.*

Icon. *Gmel. fl. sil. 3. t. 42.*
 In Canada and Labrador. On the White-hills of New Hampshire. *Prof. Peck. ♀. July. v. s. in Herb. Lambert.* Flowers pale yellow; the floral leaves purple.

2. *B. foliis alternis praelongo-linearibus, floralibus ovatis longissime acuminateis 3-nervibus, omnibus indivisis, floribus bracteis brevioribus, dentibus calycis acutis.*

On the north-west coast: Unalashka. ♀. *v. s. in Herb. Lambert.*

3. *B. hirsutissima; foliis alternis linearibus inciso-pinnatifidis: laciniis filiformibus, bracteis membranaceis oblongis obtusis basi utrinque hastato-unidentatis floribus longioribus, calycibus brevibus pilosis, dentibus subulatis.*

On the banks of Clark's river. *M. Lewis. ♀. July. v. s. in Herb. Lewis.* About a foot high; flowers deep yellow; floral leaves tinged with purple.

4. *B. foliis alternis linearibus inciso-pinnatifidis, laciniis linearibus, bracteis dilatatis plerumque trifidis floribus longioribus, dentibus calycis rotundato-obtusis.* — *Willd. sp. pl. 3. p. 185.*

Icon. *Pluk. alm. t. 102. f. 5. Moris. hist. 3. s. 11. t. 13. f. 28.*

α. *B. bracteis coccineis.*

β. *B. bracteis lutescentibus. Rarior.*

In wet meadows: Canada to Florida. ♂. ♀. June, July. *v. v.* Flowers yellow, involved in scarlet floral leaves, which give this plant a very elegant appearance.

pallida.

acuminata.

tenuifolia.

coccinea.

coccinea.
pallens.

alpina.

5. B. foliis oppositis ovatis subcordatis obtuse-serratis rugosus, antheris hirsutis.—*Willd. sp. pl. 3. p. 187.*
 Icon. *Fl. dan. 43. Engl. bot. 361. Schkuhr handb. 168.*
 In Labrador. *Colmaster.* ♀. July. v. s. in *Herb. Dickson.* Not above six inches high; floral leaves similar to the stem leaves, purple; flowers long, of a fine deep purple.

Gymnandra.

6. B. foliis suboppositis ovatis crenatis, binis radicalibus petiolatis, spica oblonga, verticillis bracteatis coarctatis.—*Willd. sp. pl. 3. p. 186.*
Gymnandra borealis. Fall. itin. 3. t. A. f. 1.
Lagotis glauca. Gært. act. petrop. 14. t. 18.
 On the mouth of the river Columbia. *M. Lewis.* In Unalashka. *Herb. Pallas.* ♀. v. s. in *Herb. Lewis.* nec non *Lambert.* About a span high; flowers pale blue. This singular plant grows on the most unfertile rocks, exposed to the most rigorous cold winds, in places where vegetation seems entirely to cease.

516. EUPHRASIA. *Gen. pl. 998.**officinalis.*

1. E. foliis ovatis obtuse-dentatis, laevis labii corollæ inferioris emarginatis. *Willd. sp. pl. 3. p. 193.*
 Icon. *Fl. dan. 1037. Engl. bot. 1416. Sabb. hort. 3. t. 9.*
 In Canada. *Michaux.* ♂. July—Sept. v. s. Flowers white, with purple veins.

latifolia.

2. E. foliis ovatis dentato-palmatis, floribus spicatis, corollis tubulosis, laevis labii inferioris obtusis. *Willd. sp. pl. 3. p. 192.*
 Icon. *Sabb. hort. 3. t. 7. Moris. hist. 3. s. 11. t. 24. f. 8.*
 In Labrador. *Colmaster.* ♂. July. v. s. in *Herb. Dickson.* Flowers smaller, pale purple.

517. MELAMPYRUM. *Gen. pl. 999.**lineare.*

1. M. gracile; foliis inferioribus linearibus integris, floralibus lanceolatis postice dentatis, floribus axillaribus distinctis.—*Lam. encycl. 4. p. 23.*
M. americanum. Mich. fl. amer. 2. p. 16.
 In shady woods, particularly on the mountains: Canada to Carolina. ♂. July, Aug. v.v. Flowers yellow.

1. O. caule sin. purpureis
Willd. sp.
 Icon. *Pluk. f. 23.*

In shady ro.
 Not above spike, sup.

1. O. caule ran-
 ciduis 4-di-
 Icon. *Moris.*
 Parasitic on t.
 ♀. July—
 plant is k
 tbrought to
 ulcers.

2. O. caule simp-
 lis obtecto
 staminibus
 Parasitic on tl
 ♀. July.
 yellow colo

3. O. scapis nudi-
 curvata.—l
 Icon. *Pluk. n.*
 Parasitic on th
 nada to Vir
 two or three
 pale purple.

518. OBOLARIA. *Gen. pl.* 1044.

1. *O. caule simplici, foliis oblongis truncatis carnosis subtus purpureis, floribus axillaribus solitariis sessilibus.*—*Willd. sp. pl. 3. p. 346.*

Icon. Pluk. alm. t. 209. f. 6. Moris. hist. 3. s. 12. t. 16. f. 23.

In shady rocky places: Pennsylvania to Virginia; rare.
Not above three inches high; flowers pale red, in a spike, supported by purple bracteas.

*virginica.*519. OROBANCHE. *Gen. pl.* 1045.

1. *O. caule ramoso, floribus alternis distantibus, corollis deciduis 4-dentatis.*—*Willd. sp. pl. 3. p. 351.*

Icon. Moris. hist. 3. s. 12. t. 16. f. 9.

Parasitic on the roots of beech-trees: Canada to Georgia.

4. July—Sept. v. v. Flowers small, purple. This plant is known by the name of *Cancer-root*, being thought to be a never-failing remedy in cancerous ulcers.

virginiana.

2. *O. caule simplicissimo squamis ovali-lanceolatis imbricatis obtecto, spica terminali glabra, corollis recurvatis, staminibus exsertis.*—*Willd. sp. pl. 3. p. 351.*

Parasitic on the roots of trees: Pennsylvania to Carolina.

4. July. v. v. About a span high, of a brownish-yellow colour.

americana.

3. *O. scapis nudis unifloris, calyce ebracteato, corolla recurvata.*—*Willd. sp. pl. 3. p. 352.*

Icon. Pluk. mant. t. 348. f. 3.

Parasitic on the roots of different plants and shrubs: Canada to Virginia. 4. June, July. v. v. Not above two or three inches high, of a tan colour; flowers pale purple.

uniflora.

feræ 4. s
reflexæ.

531. TURRITI

Cal. conr

532. BRASSIC*tum* pron

mina bre

533. DENTAR

volutis, 1

ginatum.

** *Caly*:

534. CARDAM*revolutis.*

535. SISYMBR*valvulis re*

536. CLEOME,
sinum caly
adscenden
ris, 2-valv

XIV. TETRADYDYNAMIA.

I. SILICULOSA.

* *Silicula integra.*

520. DRABA. *Silicula* integra, ovali-oblonga, valvis planiusculis dissepimento parallelis.

521. MYAGRUM. *Silicula* suberosa, evalvis, loculis fertilibus atque spuriis, sibi mutuo superimpositis. *Stylus* persistens.

522. CAKILE. *Silicula* lanceolata, subtetragona, medio utrinque dente instructa, biarticulata, ad articulos sedens: articulis monospermis, evalvibus.

** *Silicula apice emarginata.*

523. ALYSSUM. *Filamenta* quædam introrsum denticulo notata. *Silicula* emarginata, plerumque pilosa.

524. CORONOPUS. *Silicula* reniformis, compressa, corrugata, loculis evalvibus monospermis.

525. LEPIDIUM. *Silicula* emarginata, cordata, polysperma: valvulis carinatis dissepimento contrariis.

526. THLASPI. *Silicula* emarginata, obcordata, polysperma: valvulis navicularibus, marginato-carinatis.

I.

520.

1. D. scapo nudo
petalis bifid
p. 426.

Icon. *Fl. dan.*
In cultivated g
O. March—
nus are small

2. D. scapo nudo,
lis longo-line
amer. 2. p. 2
D. caroliniana,

In sandy fields :
of the Missou

3. D. scapo subuni
gerrimis subte
— *Willd. sp.* 1

II. SILIQUOSA.

* *Calyx clausus, foliolis longitudinaliter conniventibus.*

527. ERYSUMUM. *Siliqua* columnaris, tetraëdra.

528. CHEIRANTHUS. *Silicula* compressa. *Stigma* bilobum. *Germen* utrinque denticulo glandulato.

529. HESPERIS. *Petala* oblique flexa. *Siliqua* stricta. *Stigma* basi bifurca apice connivente. *Glandula* intra stamina breviora.

530. ARABIS. *Siliqua* compressæ. *Glandulae* nectari-

- seræ 4. singula intra calycis foliola, squamæ instar reflexæ.
531. TURRITIS. *Siliqua* longissima, angulata, striata. *Cal.* connivens, erectus. *Petala* erecta.
532. BRASSICA. *Cal.* erectus, connivens. *Dissepimentum* prominens. *Sem.* globosa. *Glandula* inter stamina breviora et calycem. *Siliqua* teres, rostrata.
533. DENTARIA. *Siliqua* elastice dissiliens, valvulis revolutis. *Dissepimentum* subfungosum. *Stigma* emarginatum. *Cal.* connivens.

** *Calyx hians*, *foliolis superne distantibus*.

534. CARDAMINE. *Siliqua* elastice dissiliens, valvulis revolutis. *Stigma* integrum.
535. SISYMBRIUM. *Siliqua* rostro brevi tereti dehiscens, valvulis rectiusculis. *Corolla* patens.
536. CLEOME. *Glandulæ* nectariferæ 3. ad singulum sinus calycis singulæ, excepto infimo. *Petala* omnia adscendentia. *Germen* stipitatum. *Siliqua* 1-locularis, 2-valvis.

I. SILICULOSA.

520. DRABA. Gen. pl. 1076.

1. D. scapo nudo, foliis oblongis acutis subserratis hirtis, petalis bitidis, stigmate sessili.—*Willd.* sp. pl. 3. p. 426.

Icon. *Fl. dan.* 983. *Engl. bot.* 586.

In cultivated grounds and old fields: Canada to Virginia.

○. March—May. v. v. All the species of this genus are small plants with white flowers.

2. D. scapo nudo, foliis ovalibus acutis hirsutissimis, siliulis longo-linearibus glabris approximatis.—*Mich. fl. amer.* 2. p. 28.

D. caroliniana. *Walt. fl. car.* 174.

In sandy fields: Virginia and Carolina, and on the banks of the Missouri. ○. April—June. v. v.

3. D. scapo subunifolio adscendente, foliis lanceolatis integrifinis subtomentosis: pilis stellatis, petalis retusis. —*Willd.* sp. pl. 3. p. 427.

verna.

hispidula.

nivalis.

TETRADYNAZIA SILICULOSA. Draba.

glabella.

- Icon. *Fl. dan.* 142. *Liljeblad nov. act. ups.* 6. t. 2. f. 2.
In Labrador. *Colmaster.* ♂. May. v. s. in *Herb. Dickson.*
4. D. scapo subbifolio erecto, foliis spathulato-lanceolatis
glabriusculis, racemo densifloro, siliculis ovato-lanceo-
latis acutis.

incana.

- In Hudson's Bay. ♂. v. s. in *Herb. Banks.*
5. D. foliis caulinis numerosis incanis, pilis implexis stella-
tis, siliculis oblongis obliquis stylo brevissimo mucro-
natis.—*Willd. sp. pl.* 3. p. 430.

arabisans.

- Icon. *Fl. dan.* 130. *Engl. bot.* 388. *Pluk. alm.* t. 42.
f. 1.
In Labrador. ♂. May, June. v. s. in *Herb. Lambert.*
6. D. caule folioso subramoso, foliis radicalibus cuneato-
lanceolatis, caulinis lanceolatis, omnibus acutis divi-
ricato-dentatis, siliculis stylo acuminatis.—*Mich. fl.*
amer. 2. p. 28.

On rocks, near the banks of rivers: Canada to Virginia.
♂. May, June. v. v. I collected specimens on the
rocks near Harper's Ferry, Virginia.

argenteum.

1. M. undique cano-tomentosum; siliculis oblongis pedun-
culatis tomentosis, foliis linearis-spathulatis sensim at-
tenuatis integrerrimis, petalis obovatis.
On the banks of the Missouri. ♂. v. s. Flowers yel-
low; the whole of the plant has some resemblance to
Alyssum saxatile.

maritima.

522. CAKILE. *Gært. sem.* 2. p. 287.
1. C. foliis pinnatis, pinnis linearibus subdentatis. *Willd.*
sp. pl. 3. p. 416.
Bunias Cakile. Sp. pl. 936.
Icon. *Engl. bot.* 234.
On the sea-coast: Canada to Virginia. ♂. July, Aug.
v. v. Flowers purple.

hyperboreum.

523. ALYSSUM. *Gen. pl.* 1081.
1. A. caulis herbaceis, foliis incanis dentatis, staminibus
4. bifurcatis. *Willd. sp. pl.* 3. p. 463.
Icon. *Kraschen. act. petr.* 1747. t. 15. f. 1.
On the north-west coast. +.

T

524. CC

1. C. siliculis int
rymbis pauc
Cochlearia Co
Icon. *Fl. dan.*
On road-sides
v. v. Flow
2. C. siliculis err
obsoleto, cor
Cochlearia hun
Lepidium didyi
Lepidium angl
Biscutella apeta
In cultivated gr
—Aug. v. v
both have pi

525. L

1. L. foliis radicalib
subinciso-seti
lentiformibus.
Icon. *Sloan. ja*
t. 21. f. 2.
In dry fields and
June—Aug.
conspicuous.

526. TI

1. T. siliculis orbicul
dentatis glabris
Icon. *Fl. dan.* 79
In stony fields; 1
June, July. +
2. T. hirsutum, silic
libus pinnatifidi
Icon. *Engl. bot.* 1
A common weed,
grounds. ♂.

524. CORONOPUS. *Gært. sem. t. 142.*

1. *C. siliculis integris cristato-muricatis, stylo porrecto, corymbis paucifloris.* *Smith fl. brit. 2. p. 690.*
Cochlearia Coronopus. *Willd. sp. pl. 3. p. 450.*
Icon. Fl. dan. 202. Engl. bot. 1660.
 On road-sides : Virginia and Carolina. ○. June—Aug.
 $v. v.$ Flowers very small, white.
2. *C. siliculis emarginatis didymis reticulato-rugosis, stylo obsoletō, corymbis multifloris.* *Smith fl. brit. 2. p. 691.*
Cochlearia humifusa. *Mich. fl. amer. 2. p. 27.*
Lepidium didymum. *Willd. sp. pl. 3. p. 439.*
Lepidium anglicum. *Huds. angl. 280.*
Biscutella apetala. *Walt. fl. car. 174.*
 In cultivated grounds of Virginia and Carolina. ○. June
 — Aug. $v. v.$ Smaller than the foregoing species ;
 both have pinnated leaves.

*Ruellii.**didyma.*525. LEPIDIUM. *Gen. pl. 1077.*

1. *L. foliis radicalibus pinnatifidis, caulinis linear-lanceolatis subinciso-serratis, floribus 4-petalis diandris, siliculis lentiformibus.* *Willd. sp. pl. 3. p. 440.*
Icon. Sloan. jam. 1. t. 123. f. 3. Moris. hist. 2. s. 3. t. 21. f. 2.
 In dry fields and orchards : New York to Carolina. ♂.
 June—Aug. $v. v.$ About a foot high ; flowers not conspicuous.

*virginicum.*526. THLASPI. *Gen. pl. 1078.*

1. *T. siliculis orbiculatis compressis lăvibus, foliis oblongis dentatis glabris.* *Willd. sp. pl. 3. p. 442.*
Icon. Fl. dan. 793 Engl. bot. 1659.
 In stony fields ; New York and Pennsylvania ; rare. ○.
 June, July. $v. v.$
2. *T. hirsutum, siliculis deltoideo-obcordatis, foliis radicibus pinnatifidis.* *Willd. sp. pl. 3. p. 447.*
Icon. Engl. bot. 1485. Schkuhr handb. 180.
 A common weed, every where to be found in cultivated grounds. ○. April—Sept. $v. v.$

*arvense.**ris.*

II. SILIQ UOSA.

527. ERYSIMUM. Gen. pl. 1090.

officinale.

1. E. siliquæ spicæ adpressis, foliis uncinatis. *Willd. sp. pl. 3. p. 509.*

Icon. *Fl. dan. 560. Schkuhr handb. 183.*

On road sides, frequent : Canada to Carolina. ♂. June, July. v. v. Flowers small, yellow.

cheiranthoides.

2. E. siliquæ erectis, pedicellis fructiferis patentibus, foliis lanceolatis subintegerrimis.—*Willd. sp. pl. 3. p. 511.*

Icon. *Fl. dan. 731 & 923. Engl. bot. 942. Jacq. fl. austr. 23.*

In old fields and on road sides : Virginia ; rare. ♂. June, July. v. v. Flowers yellow.

lanceolatum.

3. E. foliis lanceolatis inferioribus dentatis, unguibus calyce longioribus, laminis orbiculato-obovatis, stigmate sessili. *Ait. kew. ed. 2. t. 4. p. 116.*

Cheiranthus erysimoides. *Willd. sp. pl. 3. p. 514.*Icon. *Jacq. fl. austr. 74.*

On the banks of the Missouri. ♂. June. v. s.

528. CHEIRANTHUS. Gen. pl. 1091.

Pallasii.

1. C. caule simplici tereti glabriusculo, foliis glabris lanceo-lato-linearibus attenuatis repando-dentatis.

On the north-west coast. *Pallas. ♂. July. v. s. in Herb. Lambert.* Flowers dark red, the size of *C. Cheiri.*

529. HESPERIS. Gen. pl. 1093.

pinnatifida.

1. H. foliis argute serratis: superioribus lanceolatis, inferioribus inferne pinnatifidis, siliquis brevi-pedunculatis breviusculis. *Mich. fl. amer. 2. p. 31.*

In shady woods : Kentucky and Tennessee. ♂. June. v. v.

530. ARABIS. Gen. pl. 1049.

alpina.

1. A. foliis oblongo-lanceolatis amplexicaulibus acute dentatis. *Willd. sp. pl. 3. p. 534.*

Icon. *Bot. mog. 226. Fl. dan. 62.*In Labrador. *Colmaster. ♀. May, June. v. s. in Herb. Dickson.* Flowers, as all the following, white.

2. A. foliis radic latis sessilibi duplo longio
Icon. *Schkuhr*
In dry sandy f Virginia. C

3. A. foliis subrot tantibus. B
Icon. *Pluk. al.*
In sandy fields : v. v.

4. A. foliis glabris Willd. sp. pl.
In fields and on March—May

5. A. foliis dentati caulis hisp p. 539.
A. hispida. Ait.
Icon. *Engl. bot.*
In Labrador. D
Dickson.

6. A. foliis lanceol hastato-sessilib tis.—*Mich. fl.*

- ♦ A. canadensis. B
Turritis foliis lar siliquis compre Icon. *Pluk. alm.*
In rocky shady May, June. v. to five feet high

7. A. foliis glabris i infimis longe pe p. 204

- A. bulbosa. Muhl.*
A. amara. Herb.
A. foliis ovatis den On the borders of May. v. v. The taste resembles to have it purpose, as it wi

2. A. foliis radicalibus oblongis petiolatis, caulinis lanceolatis sessilibus, caule erecto basi hirto, petalis calyce duplo longioribus. *Willd. sp. pl. 3. p. 535.*
Icon. *Schkuhr handb. 185. Curt. fl. lond. 2. t. 49.*
In dry sandy fields and on old walls: New England to Virginia. ○. April—July. v. v.

3. A. foliis subrotundis integerrimis hirsutis, surculis rep- tantibus. *Willd. sp. pl. 3. p. 536.*

Icon. *Pluk. alm. t. 51. f. 5.*
In sandy fields: Pennsylvania to Virginia. ♀. June, July.
v. v.

4. A. foliis glabris, radicalibus lyratis, caulinis linearibus. *Willd. sp. pl. 3. p. 538.*

In fields and on dry hills: Canada to Pennsylvania. ○.
March—May. v. v.

5. A. foliis dentatis obtusis hispidis, radicalibus sublyratis, caulinis hispidis, petalis erectis.—*Willd. sp. pl. 3. p. 539.*

A. hispida. *Ait. kew. 2. p. 400.*
Icon. *Engl. bot. 614.*
In Labrador. *Colmaster. ♀. May. v. s. in Herb. Dickson.*

6. A. foliis lanceolatis utrinque angustatis remote dentatis hastato-sessilibus, siliquis pendulis ancipitibus falcatis.—*Mich. fl. amer. 2. p. 31.*

- * A. canadensis. *Willd. sp. pl. 3. p. 540.*
Turritis foliis lanceolatis dentatis, radicalibus maximis, siliquis compressis falcatis. *Gron. fl. virg. 1. p. 77.*
Icon. *Pluk. alm. t. 86 f. 8.*
In rocky shady situations: Canada to Virginia. ♀.
May, June. v. v.; v. s. in *Herb. Gronov.* From two to five feet high.

7. A. foliis glabris rhomboideis repando-obsolete-dentatis, infimis longe petiolatis, radice tuberosa.—*Pers. syn. 2. p. 204.*

A. bulbosa. *Muhl. in litt.*

A. amara. *Herb. Banks.*

A. foliis ovatis denticulatis glabris. *Gronov. virg. 99.*
On the borders of rivulets and springs. ♀. March—
May. v. v. This is one of the early spring flowers;
the taste resembling spring cresses. It would be de-
sirable to have it introduced into the gardens for that
purpose, as it will grow in any soil.

thaliana.

replana.

lyrata.

stricta.

falcata.

rhomboidea.

531. TURRITIS. *Gen. pl.* 1095.*lævigata.*

1. *T. foliis glabris, radicalibus obovatis serratis, caulinis lanceolato-linearibus integerrimis amplexicaulibus.* Willd. *sp. pl.* 3. p. 543.

On rocks: Pennsylvania to Virginia; rare. ♂. May, June. v. v. Very smooth and glaucous, about a foot high.

lvata.

2. *T. pubescens; foliis radicalibus petiolatis ovatis dentatis obtusis, caulinis amplexicaulibus oblongis serratis acutis.*

On rocks: Pennsylvania to Virginia. ♂. May, June. Resembles *T. hirsuta*.

532. BRASSICA. *Gen. pl.* 1096.*Napus.*

1. *B. radice caulescente fusiformi, foliis lævibus, superioribus cordato-lanceolatis amplexicaulibus, inferioribus lyratis dentatis.* Willd. *sp. pl.* 3. p. 547.

Icon. *Engl. bot.* 2146.

Rape frequently is found wild, but is of European origin. ♂. May, June. v. v.

Rapa.

2. *B. radice caulescente orbiculari depressa carnosa, foliis radicalibus lyratis scabris, caulinis integerrimis lævibus.* Willd. *sp. pl.* 3. p. 548.

Icon. *Engl. bot.* 2176.

Turnips are likewise apparently indigenous.

533. DENTARIA. *Gen. pl.* 1067.*laciniata.*

1. *D. foliis ternis ternatis, foliolis tripartitis linearis-oblongis incisis dentatis, radicis tuberibus moniliformibus.* — Willd. *sp. pl.* 3. p. 479.

D. concatenata. Mich. *fl. amer.* 2. p. 30.

In shady mountainous situations: New England to Carolina. ♀. May, June. v. v. Flowers large, pale purple.

diphylla.

2. *D. caulibus approximate diphylloides, foliolis ternis ovato-oblongis inaequilateris inciso-dentatis, radice dentata.* — Mich. *fl. amer.* 2. p. 30.

Icon. *Bot. mag.* 1465.

In shady beech-woods, on high mountains: Pennsylvania to Carolina. ♀. May, June. v. v. Flowers pale red, with a tinge of yellow. The roots of this plant

are of a
the native
generally
figure in
individual
which is

3. *D. caule distichio-*

liolis linea-

On the bank

Herb. Lea-

purple.

1. *C. foliis simplici-*

bus simplici-

amer. 2. p.

On rocks, in

ginia and

flowers of a

2. *C. caule nudum,*

tatis glabris

On the north-

Banks. A

3. *C. parvula;*

spathulatis

tegris dentat-

amer. 2. p.

On high moun-

4. *C. pusilla,*

acutis-

dunculis radi-

On rocks near

5. *C. pusilla,*

ere-

pinnatifidis,

retibus,

bre-

p. 29.

In overflowed

4. June, Ju-

6. *C. glabra,*

erecta-

auriculatis,

sp.

pl. 3. p. 488.

Icon. Pluk. aln-

In low meadow-

v. v.

are of a pungent mustard-like taste, and are used by the natives instead of mustard; in the mountains it is generally known by the name of *Pepper-root*. The figure in the Botanical Magazine is taken from an individual plant, in which the radical leaf is bilobated, which is not always the case.

3. *D. caule distanter diphylo, foliis sessilibus ternatis, foliolis linear-ellipticis integris, petalis cuneatis.*

On the banks of the Columbia. $\frac{1}{2}$. April. v. s. in *Herb. Lewis*. Very slender; leaves small; flowers purple.

*tenella.*534. CARDAMINE. *Gen. pl.* 1088.

1. *C. foliis simplicibus suborbiculatis integrisculis, caulis simpliciusculis debiliter procumbentibus.* *Mich. fl. amer. 2. p. 30.* *rotundifolia.*

On rocks, in the rivulets of the high mountains of Virginia and Carolina. $\frac{1}{2}$. May, June. v. v. The flowers of all the following species are white and small.

2. *C. caule nudo; foliis radicalibus lanceolatis inciso-dentatis glabris, siliquis turgidis subarticulatis.*

On the north-west coast. *Nelson*. $\frac{1}{2}$. v. s. in *Herb. Banks*. A doubtful species.

articulata.

3. *C. parvula; caulis decumbentibus, foliis radicalibus spathulatis pubescensibus, caulinis linearis cuneatis integris dentatisque, siliquis divaricato-laxis.* — *Mich. fl. amer. 2. p. 29.*

On high mountains of Carolina. *Michaux*. +.

spathulata.

4. *C. pusilla, acaulis; foliis obtuse lyrato-pinnatifidis, pedunculis radicalibus unifloris.* *Mich. fl. amer. 2. p. 29.* *uniflora.*

On rocks near Knoxville, Kentucky. *Michaux*. +.

teres.

5. *C. pusilla, erecta, ramosa; foliis omnibus sublyrato-pinnatifidis, siliquis brevibus linearibus acuminatis teretibus, brevissime pedicellatis.* *Mich. fl. amer. 2. p. 29.*

In overflowed places: New England to New Jersey. $\frac{1}{2}$. June, July. v. v.

virginica.

6. *C. glabra, erecta; foliis pinnatis, foliolis lanceolatis subauriculatis, siliquis longis stricte erectis.* — *Willd. sp. pl. 3. p. 488.*

Icon. *Pluk. alm. t. 101. f. 4.* *Petiv. gaz. t. 105. f. 18.* In low meadows: Canada to Carolina. $\frac{1}{2}$. May, June. v. v.

pensylvanica. 7. *C. glabra*, ramosa; foliis pinnatis, foliolis subrotundoblongis obtusis angulato-dentatis, siliquis angustis erectis.—*Willd. sp. pl. 3. p. 486.*

In low meadows: New York to Pennsylvania. ♀. June.
v. v.

pratensis. 8. *C. glabra*, erecta, simplex; foliis pinnatis, foliolis radiocalibus subrotundis dentatis, caulinis lanceolatis sub-integris, racemis subcorymbosis.—*Willd. sp. pl. 3. p. 487.*

Icon. *Fl. dan. 1039. Engl. bot. 776.*

In meadows: Hudson's Bay. ♀. May. v. s. in *Herb. Banks.* Flowers large, reddish purple.

multifida. 9. *C. pubescens*, ramosa; foliis interrupte pinnatis, pinnis bipinnatifidis, laciniis rotundatis incisia, siliquis brevibus.—*Herb. Banks.*

In Florida. *Bartram.* ♀. v. s. in *Herb. Banks.*

535. SISYMBRIUM. Gen. pl. 1089.

Nasturtium. 1. *S. siliquias declinatis brevibus*, foliis pinnatis, foliolis subrotundis repando-pauci-dentatis.—*Willd. sp. pl. 3. p. 489.*

Icon. *Fl. dan. 690. Schkuhr handb. 187.*

In rivulets and springs: New England to Virginia. ♀. June, July. v. v. *Water-Cress* is found in most countries; I observed it in abundance in the rivulets on the high mountains of the island of Dominica.

palustre. 2. *S. siliquias declinatis oblongo-ovatis*, foliis pinnatifidis serratis, petalis calyce brevioribus. *Willd. sp. pl. 3. p. 490.*

S. terrestre. *Smith fl. brit. 2. p. 701.*

Icon. *Fl. dan. 409. Schkuhr handb. 187.*

In inundated and low grounds: Canada to Carolina. ♂. July, Aug. v. v. Flowers yellow.

amphibium. 3. *S. siliquias declinatis oblongo-ovatis*, foliis oblongo-lanceolatis pinnatifidisve serratis, petalis calyce longioribus. *Willd. sp. pl. 3. p. 491.*

Icon. *Fl. dan. 984.*

In ponds and ditches: Pennsylvania to Virginia. ♀. June, Aug. v. v. Flowers yellow.

Sophia. 4. *S. foliis pinnato-decompositis subpilosis*, petalis calyce minoribus. *Willd. sp. pl. 3. p. 500.*

Icon. *Fl. dan. 528. Engl. bot. 963.*

On road sides: Virginia; rare. ♂. July, Aug. v. v. Flowers very small, pale yellow.

- 5
1. *C. floribui* *Willd. s.*
Icon. *Jacq*
In cultivate
○. July
structure
 2. *C. glabra*;
lanceolati
bracteis li
On the ban
Lewis. 1
 3. *C. glabrum*;
floribus a
Icon. *Sloan*
On the bank
ers purple

536. CLEOME. *Gen. pl.* 1099.

1. *C. floribus gynandris, foliis quinatis, caule inermi. pentaphylla.*
Willd. sp. pl. 3. p. 564.
Icon. *Jacq. hort. vind. 24. Rheed. mal. 9. t. 24.*
In cultivated grounds: Pennsylvania and Virginia; rare.
◎. July. v. v. Flowers pale red, of a singular structure.
2. *C. glabra; floribus hexandris, foliis ternatis, foliolis lanceolatis tenuissime serrulatis, racemo elongato, bracteis linearibus.*
On the banks of the Missouri. ◎. Aug. v. s. in *Herb. Lewis.* Flowers pale purple or white.
serrulata.
3. *C. glabrum; floribus dodecandris, foliis ternatis ellipticis, floribus axillaribus solitariis.—Willd. sp. pl. 3. p. 566.*
Icon. *Sloan. hist. 1. t. 124. f. 1.*
On the banks of the Mississippi. ◎. July. v. v. Flowers purple.

546. GERAN
Stigm. 3

XV. MONADELPHIA.

I. TRIANDRIA.

537. PHYLLANTHUS. *Cal. 6-partitus, campanulatus. Cor. 0. Styli 3. bifidi. Caps. 3-locularis. Sem. solitaria. Monoicus.*
538. MELOTHRIA. *Cal. 5-dentatus. Cor. campanulata. Stigm. 3. Bacc. 3-locularis, polysperma.*
539. MOMORDICA. *Cal. 5-fidus. Cor. 5-partita. Styl. 3-fidus. Pepo sicca elastice dissiliens. Sem. compressa. Monoica.*
540. SICYOS. *Cal. 5-dentatus. Cor. 5-partita. Styl. 3-fidus. Pepo monosperma. Monoica.*

II. PENTANDRIA.

541. PASSIFLORA. *Cal. 5-partitus. Pet. 5. calyci inserta. Nectar. corona filamentosa. Styli 3. Pepo pedicellata.*
542. IRESINE. *Cal. 2-phyllus. Pet. 5. Nectar. 5-6-7. Stigm. 2. sessilia. Caps. seminibus tomentosis. Dioica.*
543. ILLECEBRUM. *Cal. 5-phyllus, cartilagineus. Cor. 0. Squamulae 5. basi in tubum connatae. Stigma 2-fidum. Caps. 1-sperma, calyce tecta.*
544. GALAX. *Cal. 5-partitus. Pet. 5. imo staminum affixa. Urceolus 10-fidus, lacinias 5. alternis brevioribus, antheriferis. Stigm. 3-lobum. Caps. 3-locularis, 3-valvis, polysperma.*
545. LOBELIA. *Cal. 5-fidus. Cor. 1-petala, irregularis, saepius fissa. Caps. infera, 2-3-locularis.*

I.

537. PHY
1. *P. foliis obovatis, pedicellatis, pl. 4. p. 574.*

547. HALESI
Stam. 8-

548. STYRA
buliformi 2-sperma

549. HOPEA.
Stam. muc

550. GORDON
Styl. 5-g Recept, ce

551. STEWAR
connati a. Caps. 5-lo loculis 1-2

552. SIDA. *C
tus. Caps.*

553. CRISTARI
*culata. S
pellicula te*

554. MALVA.
Caps. pluri

555. MALOPE.
absque ordi

556. HIBISCUS.
Pet. 5. Ce

III. DECANDRIA.

546. GERANIUM. *Cal. 5-phylloides. Pet. 5. Stam. 10.
Stigm. 5. Arilli 5. monospermia, aristata.*

IV. POLYANDRIA.

547. HALESIA. *Cal. 4-dentatus, superus. Cor. 4-fida.
Stam. 8—12. Nux 4-angularis, 2-sperma.*
548. STYRAX. *Cal. subinteger, inferus. Cor. infundibuliformis, 5—7-partita. Stam. 8—16. Drupa 1—2-sperma, coriacea.*
549. HOPEA. *Cal. 5-fidus, superus. Cor. 5-petala.
Stam. multa, connata in 5. corpora. Styl. 1. Drupa nuce 3-loculari.*
550. GORDONIA. *Cal. 5-phylloides. Pet. 5. basi connata.
Styl. 5-gonus, stigmate 5-fido. Caps. 5-locularis.
Recept. centrale, columnare. Sem. bina, ala foliacea.*
551. STEWARTIA. *Cal. 5-partitus. Pet. 5. Styl. 5.
connati s. liberi. Stigm. capitatum, sub-5-lobum.
Caps. 5-locularis, 5-valvis, valvulis medio septiferis,
loculis 1—2-spermis. Sem. 1—2. ossea.*
552. SIDA. *Cal. simplex, angulatus. Styl. multipartitus.
Caps. plures, 1-spermæ.*
553. CRISTARIA. *Cal. simplex, 5-fidus. Pet. 5. unguiculata.
Styl. plures. Fructus orbiculato-depressus,
pellicula tectus, dehiscens in Arillos bialatos.*
554. MALVA. *Cal. duplex : exterior 3-phylloides. Pet. 5.
Caps. plurimæ, evales, 1-spermæ.*
555. MALOPE. *Cal. duplex : exterior 3-phylloides. Caps.
absque ordine glomeratæ, monospermæ.*
556. HIBISCUS. *Cal. duplex : exterior polyphylloides.
Pet. 5. Caps. 5-locularis, polysperma.*

I. TRIANDRIA.

557. PHYLLANTHUS. *Gen. pl. 1412.*

1. P. foliis obovatis obtusiusculis, floribus geminis axillaribus
pedicellatis, caule erecto ramoso tereti. *Willd. sp.
pl. 4. p. 574.*

obovatus.

P. carolinensis. Walt. fl. car. 228. Mich. fl. amer. 2. p. 209.

In gravelly soil, on the banks of rivers: Pennsylvania to Georgia. ☽. June, July. v. v. About a span or more high; the branches grow regularly in two rows.

538. MELOTHRIA. *Gen. pl. 68.**pendula.*

1. *M. foliis subreniformibus lobato-angulosus: lobo impari cæteris æquali, bacca cylindraceo-ovata. Mi h. fl. amer. 2. p. 217. Willd. sp. pl. 1. p. 189.*

Icon. *Pluk. alm. t. 85. f. 5.*

In sandy, wet places, particularly on the islands of large rivers: Virginia to Georgia. ☽. June. v. v. Flowers small, yellow. Fruit the size of a pea, black.

539. MOMORDICA. *Gen. pl. 1477.**echinata.*

1. *M. pomis tetraspermis subrotundis setoso-echinatis, foliis cordatis 5-lobo-angulatis acuminatis integerrimis glabris. Willd. sp. pl. 4. p. 605.*

Sicyos lobata. *Mich. fl. amer. 2. p. 217.*

In the western parts of the United States: On the banks of the Ohio and Mississippi. ☽. July. v. v. Fruit the size of a gooseberry.

540. SICYOS. *Gen. pl. 1481.**angulata.*

1. *S. foliis angulo obtuso cordatis 5-angularibus denticulatis scabris, fructibus capitatis, hispidis.—Willd. sp. pl. 4. p. 625.*

Icon. *Dill. elth. t. 51. f. 59. Herm. parad. t. 133.*

On the banks of rivers: Canada to Carolina. ☽. June, July. v. v.

II. PENTANDRIA.

541. PASSIFLORA. *Gen. pl. 509.**lutea.*

1. *P. foliis cordatis trilobis obtusis glabris, petiolis eglandulosis, pedunculis axillaribus geminis, petalis calyce duplo angustioribus. Willd. sp. pl. 3. p. 615.*

Icon. *Cavan. diss. 10. t. 267. Jacq. ic. 3. t. 607.*

MONA

On the banana to small, gr

2. *P. foliis lanceolatis culis solit pl. 3. p. 6*

Icon. *Cava*
In Virginia
the size.

3. *P. foliis tril glandulos glandulos Willd. sp.*

Icon. *Cava*
ic. 193.

On the bank
June—Au
riegated.

541

1. *I. foliis punctis perioribus caule sulca*
In overflowed
to Florida.
ing small.

543. I

1. *I. caulibus retiolatis, capi sp. pl. 1. p*

Gomphrena ¶
I. Achyrantha

Icon. *Solan. i*
On the sea-

about Chai
Aug. v. v.
ers, in spin-

2. *I. caulibus retiolatis, capi pl. 1. p. 120*

Achyranthes fi

- On the banks of rivers, in shady rocky situations : Virginia to Florida. ♀. June—Aug. v. v. Flowers small, greenish-yellow.
2. *P. foliis peltatis profunde trilobis glabris, lobis linearilanceolatis divaricatis, petiolis biglandulosis, pedunculis solitariis axillaribus, floribus apetalis.* *Willd. sp. pl. 3. p. 617.*
Icon. *Cavan. diss. 10. t. 274: Pluk. alm. t. 210. f. 4.*
In Virginia and Georgia. ♂. v. s. Flowers of a middle size.
3. *P. foliis trilobis serratis, lobis oblongis acutis, petiolis biglandulosis, involucro triphylo, foliolis lanceolatis glanduloso-dentatis, filis coronæ corolla longioribus.* *Willd. sp. pl. 3. p. 621.*
Icon. *Cavan. diss. 10. 293. Hern. mex. t. 888. Rob. ic. 193.*
On the banks of rivers of Virginia and Carolina. ♀. June—Aug. v. v. Flowers beautiful, blue and variegated.

*peltata.**incarnata.*542. IRESINE. *Gen. pl. 1519.*

1. *I. foliis punctato-scabris, inferioribus lato-ovalibus, superioribus ovato-lanceolatis, panicula ramosa conferta, caule sulcato.* —*Willd. sp. pl. 4. p. 765.*
In overflowed grounds, on the banks of rivers : Virginia to Florida. ♂. July, Aug. v. v. Flowers exceeding small.

*celosioides.*543. ILLECEBRUM. *Gen. pl. 407.*

1. *I. caulibus repentibus hirsutis, foliis lato-lanceolatis petiolatis, capitulis orbiculatis nudis sessilibus.* —*Willd. sp. pl. 1. p. 1208.*
Gomphrena polygonoides. Sp. pl. 225.
I. Achyrantha. Walt. fl. car. 103.
Icon. *Sloan. hist. 1. f. 86. f. 2.*
On the sea-coast : Virginia to Carolina ; plentifully about Charlestown, South Carolina. ♀. June—Aug. v. v. A low creeping plant, with small flowers, in spiny clusters.
2. *I. caulibus repentibus glabris, foliis lato-lanceolatis petiolatis, capitulis orbiculatis pubescentibus.* *Willd. sp. pl. 1. p. 1208.*
Achyranthes ficoidea. Pers. syn. 1. p. 259.

ficoideum.

MONADELPHIA PENTANDRIA. Illecebrum.

MONAD

Icon. *Jacq. amer.* t. 60. f. 4.
On the sea-coast: Carolina to Florida. ♀. July, Aug.

v. v. Resembles the former one.

vermicula-
tum.

3. I. glabrum; caulibus repentibus, foliis subteretibus carnosis, capitulis solitariis terminalibus oblongis.—*Sp. pl.* 300.

Gomphrena vermicularis. *Willd. sp. pl.* 1. p. 1322.

Icon. *Herm. parad.* t. 15. *Pluk. alm.* t. 75. f. 9.

On the sea-coast: Carolina to Florida. ♀. June—Aug.
v. s.

dichotomum.

4. I. caulibus suffruticosis diffusis, foliis linearibus planis acutis margine scabris, cyma dichotoma.—*Willd. sp. pl.* 1. p. 1196.

In Virginia. ♂. +. I never saw this plant either living or in a dried state.

544. GALAX. *Gen. pl.* 382.

Erythrorhiza. *Mich. fl. amer.* 2. p. 34.

rotundifolia.

1. G. glaberrima; foliis orbiculato-reniformibus dentatis, spica longissima.

Galax aphylla. *Willd. sp. pl.* 1. p. 1146.

Blandfordia cordata. *Bot. rep.* 343.

Solanandra cordifolia. *Vent. malm.* 69.

Erythrorhiza rotundifolia. *Mich. fl. amer.* 2. p. 36.

Icon. *Mich. fl. amer.* 2. t. 36. *Vent. l. c.* *Bot. rep.* 343.

On the side of shady hills, near rivulets, in the high mountains of Virginia and Carolina. ♀. May, June.
v. v. Flowers white, in long and close spikes. In the Virginia mountains this plant is known by the name of *Carpenter's-leaf*, being used in healing all kinds of wounds and cuts.

545. LOBELIA. *Gen. pl.* 1363.

Dortmanna.

1. L. foliis radicalibus linearibus recurvis fistulosis bilocularibus integerrimis, scapo simplici subnudo racemoso, floribus remotis pendulis.—*Willd. sp. pl.* 1. p. 938.

Icon. *Fl. dan.* 39. *Engl. bot.* 140.

On Hudson's Bay. *Michaux.* ♀. July, Aug. v. s.
Flowers light blue.

Kalmii.

2. L. tenuis, erecta, simpliciuscula; foliis radicalibus spatulatis, caulinis linearibus tenuissime denticulatis, floribus racemosis alternis remotis pedicellatis.—*Willd. sp. pl.* 1. p. 939.

In low sandy
♂. ○. J.

3. L. erecta, si-
siusculis,
denticulatis
Mich. fl. a-

Rapantium fo-
L. spicata. *L.*
L. goodenioi-
In shady wood-
lina. ♀. J.
high; flow-

4. L. erecta, g-
spica multi-
corollæ lac-
amer. 2. p.
On the moun-
Aug. v. v
two to three

5. L. erecta, sub-
ceolatis gla-
cemosis br-
dentatis.—
L. crassiuscula

In pine-swam-
v. v. From
dark blue.

6. L. erecta, sim-
libus repand-
sessilibus,
Mich. fl. a-

In the range
♀. July, A.
flowers mid-

7. L. erecta, sim-
serratis, rad-
reflexis.—
Icon. Rob. ic.

In low ground
New York
large, blue,
virtues; or
into Europe

In low sandy fields and meadows: Canada to Carolina.
♂. ♂. July, Aug. v. v. Flowers small, blue.

3. *L. erecta, simplex, subpubescens; foliis oblongis obtusiusculis, radicalibus integerrimis, caulinis superne denticulatis, racemo virgato, bracteis subulatis.*—
Mich. fl. amer. 2. p. 153.

Rapuntium foliis villosis mollibus, &c. *Gron. virg. 135.*

L. spicata. *Lam. encycl. 3. p. 587.*

L. goodenioides. *Willd. hort. berol. 30. cum icon.*

In shady woods and on grassy hills: New York to Carolina. ♀. July—Sept. v. v. About eighteen inches high; flowers pale blue, small.

4. *L. erecta, glaberrima; foliis lato-lanceolatis serratis, spica multiflora secunda, calycis laciniis integerrimis, corollæ laciniis inferioribus ovalibus acutis.*—*Mich. fl. amer. 2. p. 153.*

On the mountains of Virginia and Carolina. ♀. July, Aug. v. v. Flowers of a beautiful sky-blue. From two to three feet high.

5. *L. erecta, subramosa, subpubescens, lucida; foliis lanceolatis glanduloso-serrulatis subcarnosis, floribus rachemosis brevi-pedunculatis, calycis laciniis revolutis dentatis.*—*Walt. fl. car. 218.*

L. crassiuscula. *Mich. fl. amer. 2. p. 152.*

In pine-swamps: Virginia to Florida. ♀. July, Aug. v. v. From eight inches to a foot high; flowers dark blue.

6. *L. erecta, simplicissima, pubescens: foliis oblongo-ovalibus repando-serrulatis, floribus spicatis alternis sessilibus, germinibus hispidis, calycibus ciliatis.*—*Mich. fl. amer. 2. p. 152.*

In the range of mountains, from Virginia to Carolina. ♀. July, Aug. v. v. From one to two feet high; flowers middle size, sky blue.

7. *L. erecta, simplex, hirtella; foliis ovato-lanceolatis subserratis, racemo folioso, calycibus hirsutis, sinus reflexis.*—*Willd. sp. pl. 1. p. 945.*

Icon. Rob. ic. 136. Jacq. ic. 3. t. 597.

In low grounds, on the banks of rivers and near springs: New York to Carolina. ♀. July, Aug. v. v. Flowers large, blue. This plant is possessed of great medicinal virtues; on account of which it was introduced into Europe nearly two hundred years ago.

cardinalis.

8. L. erecta, simplex, pubescens; foliis ovato-lanceolatis acuminate eroso-denticulatis, racemo subsecundo multifloro, genitalibus corolla longioribus.—*Willd. sp. pl. 1. p. 944.*

Icon. *Bot. mag. 320. Rob. ic. 137. Knorr delic. 2. t. L. 2.*

In swamps and meadows near springs: Canada to Carolina. ♀. July—Sept. v. v. This beautiful plant, with its bright scarlet flowers, rises to the height of from two to four feet. I have seen a white variety of it.

fulgens.

9. L. erecta, simplex, subpubescens; foliis elongato-lanceolatis attenuatis subintegerrimis, racemo multifloro, genitalibus longitudine corollæ.—*Donn. catal.*

Icon. *Bot. rep. 659.*

On the Mississippi. ♀. June, July. v. v. in *Hortis.* This species exceeds in splendour of colour and in size the preceding one.

inflata.

10. L. erecta, ramosa, hirsutissima; foliis ovatis serratis, racemis foliosis, capsulis inflatis.—*Willd. sp. pl. 1. p. 946.*

In fields and woods, common: Canada to Carolina. ♂. June—Aug. v. v. Flowers blue, small.

Cliffortiana.

11. L. glabra, superne ramosa; foliis petiolatis ovatis crenato-dentatis, infimis suborbiculatis, racemis laxis, pedunculis elongatis.—*Willd. sp. pl. 1. p. 946.*

Icon. *Plum. ic. 235. f. 2.*

In old fields: Virginia. ♂. June, July. v. v. Flowers very small, purple.

II. DECANDRIA.

546. GERANIUM. Gen. pl. 1118.

maculatum.

1. G. erectum, retrorsum pubescens; caule dichotomo, foliis oppositis 3—5-partitis incisis, summis sessilibus, pedunculis elongatis bifloris, petalis obovatis.—*Willd. sp. pl. 3. p. 705.*

Icon. *Cavan. diss. 4. t. 86. f. 2. Dill. elth. t. 132. f. 159.*
In shady woods and meadows: Canada to Carolina. ♀. May—July. v. v. Flowers purple, large. This

species is
the name
taste of its
in curing
very preva

2. G. diffusum
trifido-inci-
tal is emar-
losis.—*W.*

G. lanuginosum.

Icon. *Cava-
elth. t. 13.*

In fields: V.
Flowers s

3. G. diffusum
3-fid is in-
petalis em-
pilosis, pi-

Icon. Cava-

In fields: N.
rose-colou

4. G. diffusum
tisque tri-
integrifoli-
culatis.—*W.*

Icon. Fl. da-

In shady m-
ginia; ran-
red. It is
that heavy
though its

547. H. foliis ova-
candris, fi-

Willd. sp.

Icon. *Bot.*
car. 1. t.

species is known in some parts of the mountains by the name of *Alum-root*, on account of the astringent taste of its roots, which are very successfully employed in curing the flux among children, which is a disease very prevalent in those countries.

2. *G. diffusum*, pubescens; foliis oppositis 5-lobis, lobis *carolinianum*. trifido-incisis, pedunculis bifloris subfasciculatis, petalis emarginatis longitudine calycis aristati, arillus villosus.—*Willd. sp. pl. 3. p. 711.*

G. lanuginosum. *Jacq. hort. schœnb. 2. t. 140.*
Icon. *Cavan. diss. 4. t. 84. f. 1.* ♂ *t. 124. f. 2.* *Dill. elth. t. 135. f. 162.*

In fields: Virginia to Georgia. ♂. June, July. v. v.
Flowers small, white, with red veins.

3. *G. diffusum*, pubescens; foliis oppositis 5-partitis, lobis 3-fidis incisis linearibus, pedunculis bifloris elongatis, petalis emarginatis longitudine calycis aristati, arillus pilosis, pilis glandulosis.—*Willd. sp. pl. 3. p. 712.*

Icon. *Cavan. diss. 4. t. 78. f. 2.*

In fields: New York. ♂. June, July. v. v. Flowers rose-coloured, larger than the foregoing one.

4. *G. diffusum*, ~~bifidum~~; foliis oppositis ternatis quinatisque trifido-pinnatifidis, pedunculis bifloris, petalis integris calyce aristato duplo longioribus, arillus reticulatis.—*Willd. sp. pl. 3. p. 714.*

Icon. *Fl. dan. 694.* *Cavan. diss. 4. t. 86. f. 1.*

In shady moist places, on rocks: New York to Virginia; rare. ♂. June—Aug. v. v. Flowers pale red. It is singular, that the American plant has not that heavy scent the European is so well known by, though its other characters agree exactly.

III. POLYANDRIA.

547. HALESIA. *Gen. pl. 814.*

1. *H. foliis ovatis acuminatis argute serratis, floribus dodecandris, fructu subæqualiter rhomboideo-tetraptero.* *tetraptera.*
Willd. sp. pl. 2. p. 849.
Icon. *Bot. mag. 910.* *Cavan. diss. 6. t. 186.* *Catesb. car. 1. t. 64.*

- In shady woods, on the banks of rivers: Carolina. $\frac{1}{2}$. April, May. *v. v.* A handsome small tree; flowers white, resembling those of Snow-drops, from which it has been called *Snow-drop Tree* or *Silver-bell Tree*.
- diptera.* 2. *H. foliis oblongo-ovatis obtuse-acuminatis membranaceis, floribus octandris, fructu subelliptico, alis alternis majoribus.* —*Willd. sp. pl. 2. p. 849.*
Icon. *Cavan. diss. 6. t. 187.*
In shady forests of Georgia. $\frac{1}{2}$. April, May. *v. v. s.* fl.; *v. s. in Herb. Lyon.* Flowers larger than the preceding species.
- parviflora.* 3. *H. fructu inaequaliter subtetraptero clavato floribusque parvulis.* —*Mich. fl. amer. 2. p. 40.*
In Florida near Matanza. *Michaux.* $\frac{1}{2}.$ +.

548. STYRAX. *Gen. pl. 753.*

- grandifolium.* 1. *S. foliis lato-ovobatis acuminatis subtus tomentosis, racemis multifloris elongatis, pedunculis inferioribus axillaribus solitariis unifloris.* —*Willd. sp. pl. 2. p. 622.*
S. officinale. Walt. fl. car. 140.
S. grandiflorum. Mich. fl. amer. 2. p. 41.
In woods, on the banks of rivers: Virginia to Georgia. $\frac{1}{2}$. June—Aug. *v. v.* A fine ornamental shrub, as the two following likewise are; flowers white.
- pulverulentum.* 2. *S. foliis subsessilibus ovalibus s. obovatis obtusis subtus pulverulento-tomentosis, floribus axillaribus et subterno-terminalibus brevissime pedicellatis.* —*Mich. fl. amer. 2. p. 41.*
S. laevigatum. Bot. mag. 921.
In woods of Virginia and Carolina. $\frac{1}{2}$. June—Aug. *v. v.* Resembles the foregoing one. The figure referred to represents the present species, *S. laevigatum* or the following species has to my knowledge not yet been introduced into Europe.
- glabrum.* 3. *S. foliis ovali-lanceolatis utrinque acutis glabris, pedunculis axillaribus unifloris solitariis binisve.* —*Cavan. diss. 6. t. 188. f. 1.*
S. laevigatum. Willd. sp. pl. 2. p. 624.
S. americanum. Lam. encycl. 1. p. 82.
S. laeve. Walt. fl. car. 140.
In the swamps of Virginia and Carolina. $\frac{1}{2}$. July, Aug. *v. v.* Not above three or four feet high; the number of stamens is from six to ten.

1. *H. foliis oblique aggregatis Symplocos ti.*
Icon. *Mich.* In low woods *v. v.* A which the taste; from inhabitants

1. *G. foliis lance floribus longatis.* —*Willd. Hypericum L.*
Icon. *Bot. mag. t. 161. C. f. 3.*
In cedar-swan rida. $\frac{1}{2}$. A green tree;
2. *G. foliis obov membranaceis.* —*Willd. G. Franklinia. Franklinia Alt. Lacathea florid. Icon. Cavan. On the banks of Georgia, whose large a most agreeable southern cli*

1. *S. foliis ovatis cibus ovatis S. Malachoder*

549. HOPEA. *Gen. pl.* 1222.

1. *H. foliis oblongo-lanceolatis glaucis, floribus axillaribus aggregatis brevissime pedicellatis.* — *Linn. mant.* 105. *Symplocos tinctoria.* *Willd. sp. pl.* 3. p. 1436. *Icon. Mich. arb. t. . . . Catesb. car. 1. p. 54.*
In low woods of Virginia and Carolina. ♀. April, May.
v. v. A small tree or shrub, variable in size, of which there is a variety with leaves pubescent underneath; flowers yellow, small, very sweet scented. The leaves dye a very fine yellow, and have a sweet taste; from which circumstance it is known by the inhabitants under the name of *Sweet-leaf*.

550. GORDONIA. *Gen. pl.* 1144.

1. *G. foliis lanceolato-oblongis glaberrimis nitidis coriaceis, floribus longe pedunculatis, capsulis conoideis acuminate.* — *Willd. sp. pl.* 3. p. 840. *Hypericum Lasianthus.* *Sp. pl.* 1101. *Icon. Bot. mag.* 668. *Mich. arb. t. . . . Cavan. diss. 6. t. 161. Catesb. car. 1. t. 44. Pluk. amaln. t. 352. f. 3.*
In cedar-swamps, near the sea-coast: Virginia to Florida. ♀. Aug—Sept. *v. v.* A beautiful small evergreen tree; flowers white, the size of a rose.
2. *G. foliis obovato-lanceolatis subtus pubescensibus submembranaceis, floribus subsessilibus, capsulis sphæricis.* — *Willd. sp. pl.* 3. p. 841. *G. Franklinia.* *Willd. sp. pl.* 3. p. 841. *Franklinia Alatamaha.* *Marsh. arb.* 48. *Lacathea florida.* *Parad. lond.* 56. cum iconē. *Icon. Cavan. diss. t. 162. Mich. arb. t. . . .*
On the banks of the Alatamaha, near Fort Barrington, Georgia. ♀. Aug. Sept. *v. v.* This elegant tree, whose large white flowers with yellow anthers have a most agreeable appearance, though a native of a very southern latitude, is able to stand a considerable northern climate.

551. STUARTIA. *Gen. pl.* 1142 & 1143.

1. *S. foliis ovatis acutis, floribus axillaribus subbinis, calycibus ovatis obtusis, petalis integris, stylis coalitis.* *virginica.*
S. Malachodendron. *Willd. sp. pl.* 3. p. 840.

MONADELPHIA POLYANDRJA. Stuartia.

Stewartia virginica. *Cavan. diss.* 5. t. 159. f. 2.

Stewartia marilandica. *Bot. rep.* 397.

Icon. *L'Herit. stirp.* 1. t. 73. *Catesb. car.* 3. t. 13.
Duham. arb. 2. t. 78.

In swamps in the lower counties of Virginia and Carolina. $\frac{1}{2}$. July—Sept. v. v. From five to seven feet high, very ornamental; flowers large, white. There are a number of varieties, with more or less serrated leaves, more or less pubescent, &c.

pentagyna.

2. *S. foliis ovatis acuminatis, floribus axillaribus solitariis, calycibus lanceolatis calyculatis, petalis undulato-in-cisis, stylis distinctis.* — *Willd. sp. pl.* 3. p. 840.

Malachodendron ovatum. *Cavan. diss.* 5. t. 158. f. 2.

Icon. *Exot. bot.* 101. *L'Herit. stirp.* 1. t. 74. *J. Miller. ic.* 3.

In the mountains of Carolina and Georgia. $\frac{1}{2}$. Aug. v. v. This beautiful shrub, whose large flowers are of a cream colour, deserves a place in the collection of every admirer of ornamental shrubs. It has generally been considered as a distinct genus; but I follow Dr. Smith's observations in Exotic Botany, considering it always unnecessary to split a small genus on slight differences, though the same differences might be sufficient reasons for establishing a distinction where the genus is numerous.

552. SIDA. *Gen. pl.* 1129.

spinosa.

1. *S. caule patulo, axillis subspinosis, foliis longe petiolatis ovato-lanceolatis obsolete cordatis dentatis, pedunculis solitariis axillaribus, stipulis setaceis pedunculo longioribus, capsulis birostratis.* — *Willd. sp. pl.* 3. p. 736.

Icon. *Cavan. diss.* 1. t. 1. f. 9. *Pluk. alm.* t. 9. f. 6.

Among rubbish and on road-sides: Pennsylvania to Carolina. \odot . July, Aug. v. v. Flowers yellow, small.

hispida.

2. *S. hispido-pilosa; foliis lanceolatis serratis, pedunculis solitariis axillaribus longitudine petolorum, calyce exteriore filiformi.*

In sandy plains of Georgia. *Lyon.* $\frac{1}{2}$. v. s. in *Herb. Lyon.* Flowers yellow.

rhombofolia.

3. *S. foliis oblongo-lanceolatis dentatis basi cuneiformibus integerrimis, pedunculis petiolis multo longioribus, capsulis bicornibus.* *Willd. sp. pl.* 3. p. 740.

Icon. *Cavan. diss.* 1. t. 3. f. 12. *Dill. elth.* t. 172. f. 212.

MO

In cultivat
Florida,
low, sma

4. *S. foliis obl
silibus, l
feris defl
Willd. s*

Icon. *Cava*
On the sea
Flowers

5. *S. foliis sub
tosis, pec
biaristatis*

Icon. *Schki*
In cultivate
July, Au

6. *S. foliis cor
tis denta
acuminat*

Napaea laevi
Napaea harr

Icon. *Cava*
In shady ro
 $\frac{1}{2}$. July-
flowers w

7. *S. foliis 7-lo
dentatis,
sp. pl.* 3. 1

Napaea scabi
Napaea dioic

Icon. *Cava*
In stony gro
July, Aug

8. *S. foliis infe
bus palma
hispidis.*

In stony field
Aug. v. v.

553. C

1. *C. undique c
3—5-fidis,
caule diffu*

In cultivated grounds and among rubbish : Virginia to Florida. ♂. June—Aug. v. v. Flowers yellow, small.

4. S. foliis oblongo-cordatis acuminatis crenatis summis sessilibus, pedunculis solitariis petiolo longioribus fructiferis deflexis, capsulis inflatis muticis undulato-crispis. *Willd. sp. pl. 3. p. 747.*

Icon. *Cavan. diss. 1. t. 7. f. 1. Dill. elth. t. 5. f. 5.*

On the sea-coast of Carolina. ♂. July—Sept. v. v. Flowers white, small.

5. S. foliis subrotundo-cordatis acuminatis dentatis tomentosis, pedunculis solitariis petiolo brevioribus, capsulis bariostatis truncatis. *Willd. sp. pl. 3. p. 750.*

Icon. *Schkuhr handb. 190.*

In cultivated grounds : Pennsylvania to Carolina. ♂. July, Aug. v. v. A tall plant; flowers dark yellow.

6. S. foliis cordatis 5-lobis glabris, lobis oblongis acuminatis dentatis, pedunculis multifloris, capsulis muticis acuminatis. *Willd. sp. pl. 3. p. 766.*

Napaea laevis. Syst. veg. 750.

Napaea hermaphrodita. Sp. pl. 965.

Icon. *Cavan. diss. 5. t. 132. f. 1. Herm. lugdb. t. 23.*

In shady rocky places : Pennsylvania to Virginia ; rare. ♀. July—Sept. v. v. From two to four feet high ; flowers white.

7. S. foliis 7-lobo-palmatis scabris, lobis lanceolatis inciso-dentatis, floribus dioicis corymbosis bracteatis. *Willd. sp. pl. 3. p. 766.*

Napaea scabra. Syst. veg. 750.

Napaea dioica. Sp. pl. 965.

Icon. *Cavan. diss. 5. t. 132. f. 2. Ehret. pict. 7 & 8.*

In stony ground, in the upper parts of Virginia. ♀. July, Aug. v. v. Flowers white.

8. S. foliis inferioribus triangulo-cordatis incisis, superioribus palmato-multifidis, corymbo terminali, calycibus hispidis. *Mich. fl. amer. 2. p. 44.*

In stony fields of Kentucky, Tennessee, &c. ♀. July, Aug. v. s. Resembles in habit *Malva Alcea*.

crispa.

Abutilon.

Napaea.

dioica.

alcooides.

coccinea.

553. CRISTARIA. *Cavan. ic. 5. p. 10.*

1. C. undique cano-tomentosa et pilis stellatis obsita ; foliis 3—5-fidis, lacinias incisis acutis, racemis terminalibus, caule diffuso.

Malva coccinea, *Fraser, catal.*

On the dry prairies and extensive plains of the Missouri.

♀. Aug. Sept. v. v.; v. s. in *Herb. Lewis*. Flowers scarlet.

Caulis teres, subpedalis, ramosus, ramis diffusis, tomentosus, pilis stellatis distinctis obsitis, uti tota planta. *Folia* alterna, petiolata, palmato-3—5-partita: lacinias sub-pinnatifido-incisis, lobis sinibusque acutis. *Petoli* teretes, longitudine folii. *Stipulæ* lineares, erectæ, marcescentes. *Flores* terminales, racemosi, coccinei, diametro pollicari. *Calyx* 5-fidus, lacinias lanceolatis acutis. *Petala* 5, oblique cuneato-obcordata, subunguiculata, ad ungues fasciculato-villosa, calyce duplo longiora. *Filamenta* monadelpha, numerosa, in columnam brevem connata, superne libera, patentia. *Stylæ* circiter 10. *Fructus* orbiculatus.

The singularly stellated fascicles of hair, placed on the close tomentum, and its bright scarlet flowers, make this plant particularly interesting.

554. MALVA. Gen. pl. 1134.

abutiloides.

1. M. foliis 5-angulari-lobatis tomentosis, pedunculis sub-4-floris bifidis axillaribus, capsulis polyspermis. *Willd. sp. pl. 3. p. 780.*

Icon. *Jacq. hort. schœnbr. 3. t. 293.* *Cavan. diss. 2. t. 16. f. 2.*

On the sea-coast of Carolina. ♀. June—Aug. v. v. Flowers large, purple.

caroliniana.

2. M. foliis 5-lobis palmatisve inciso-dentatis, pedunculis petiolo longioribus, petalis integris, fructu villoso, caule prostrato. *Willd. sp. pl. 3. p. 784.*

Icon. *Cavan. diss. 2. t. 15. f. 1.* *Dill. elth. t. 4. f. 4.* *Schkuhr handb. 192.*

In fields: Virginia and Carolina. ♂. July—Sept. v. v. Flowers dark red.

rotundifolia.

3. M. caule prostrato, foliis cordato-orbiculatis obsolete 5-lobis, pedunculis fructiferis declinatis. *Willd. sp. pl. 3. p. 786.*

Icon. *Fl. dan. 721.* *Cavan. aiss. 2. t. 26. f. 3.* *Schkuhr handb. 1.*

In cultivated grounds: Pennsylvania to Virginia. ♀. June—Sept., v. v. Flowers white, small.

1. M. foliis oblongo-ovalibus pedunculis sub-4-floris bifidis axillaribus, capsulis polyspermis. *Willd. sp. pl. 3. p. 799.*
Icon. *Cavan. Rob. ic. 18.*
In Carolina, plant having drawing the
Walter me

555

1. H. foliis ovatis vibus subtus calycibus tenuibus. *Willd. sp. pl. 3. p. 800.*
H. palustris. *Cavan. H. palustri.*
Icon. *Cavan. H. palustri.*
In swamps and
plentifully in
dago, New
large, white,
pale purple.

2. H. foliis lati-ovalibus subtus tomentosis gioribus.—
Icon. *Cavan. aiss. 2. t. 26. f. 3.*
In swamps and
—Oct. v. v.
been genera-
very distinct.

3. H. foliis ampliatis utrinque ex-
tomentoso-hirsutis. *Willd. sp. pl. 3. p. 46.*
In salt-marshes
of the Mississ.
very large, 1
fructification.

4. H. foliis ovatis non-tomentosis sub-
tomentosis sub-

555. MALOPE. *Gen. pl.* 1136.

1. *M. foliis oblongis obtusis integris crenatis supra glabris, pedunculis solitariis axillaribus.* —*Willd. sp. pl.* 3. *p. 799.*

*Icon. Cavan. diss. 2. t. 27. f. 1. Sall. hort. 1. t. 50.
Rob. ic. 189.*

In Carolina. *Walter.* ♂. +. Though I doubt of this plant having been found in America, yet it is worth drawing the attention of the botanist to find out what Walter means by it.

556. HIBISCUS. *Gen. pl.* 1139.

1. *H. foliis ovatis acuminatis serratis subtrilobis sub-5-nervibus subtus incano-tomentosis, pedunculis floriferis, calycibus tomentosis, capsulis glabris.* —*Willd. sp. pl.* 3. *p. 806.*

H. palustris. Walt. fl. car. 176.

Icon. Cavan. diss. 3. t. 65. f. 1. Bot. mag. 882. sub H. palustri.

In swamps and salt-marshes : New York to Carolina ; plentifully in the marshes round the Salt-lake, Onondago, New York. ♀. Aug.—Oct. v. v. Flowers large, white, with a purple bottom, or sometimes pale purple.

2. *H. foliis lato-ovatis obtuse-serratis subtrilobis 3-nervibus subtus tomentosis, pedunculis axillaribus petiolo longioribus.* —*Willd. sp. pl.* 3. *p. 808.*

Icon. Cavan. diss. 3. t. 65. f. 2.

In swamps and marshes : Canada to Virginia. ♀. Aug.—Oct. v. v. Flowers large, purple. This plant has been generally confounded with the former, but is very distinct.

3. *H. foliis amplis coriaceis triangulato-cordatis trilobis utrinque eximie tomentosis subtus incanis, capsulis tomentoso-hirsutissimis subtruncatis.* —*Mich. fl. amer.* 2. *p. 46.*

In salt-marshes of Georgia and Florida, and on the banks of the Mississippi. ♀. Aug.—Oct. v. v. Flowers very large, flesh-coloured, with a red bottom ; the fructifications yellow.

4. *H. foliis ovatis acuminatis obtuse serratis utrinque incano-tomentosis, pedunculis axillaribus, calycibus tomentosis subæqualibus.* —*Willd. sp. pl.* 3. *p. 807.*

palustris.

incanus.

Icon. *Wendl. hort. herrenh.* 4. t. 24. *Bartr. ic. ined. in Mus. Banks.*

In Carolina. ♀. Sept. +. Flowers very large, sulphur-coloured.

esculentus.

5. *H. foliis cordatis 5-lobis obtusiusculis dentatis, petiolis flore longioribus, calycibus exterioribus subdecaphyllis deciduis, interioribus longitudinaliter rumpentibus.* *Willd. sp. pl. 3. p. 827.*

Icon. *Cavan. diss. 3. t. 61. f. 2.* *Comm. hort. 1. t. 19.*
In cultivated grounds: Virginia and Carolina. ♂. June —Sept. v. v. Flowers yellow. It is generally cultivated by the name of *Okra* for the sake of its unripe capsules, which give a fine dish for the table.

virginicus.

6. *H. undique tomentosus, asper; foliis acuminatis inaequilater dentatis, inferioribus cordatis indivisis, superioribus cordatis oblongis trilobis, pedunculis axillaribus et racemoso-terminalibus, floribus cernuis, pistillis nubantibus.* *Willd. sp. pl. 3. p. 831.*

H. clypeatus. *Walt. fl. car. 177.*

Icon. *Jacq. ic. rar. t. 142.* *Pluk. phyt. t. 6. f. 4.*

In salt-marshes, on the sea-coast: New Jersey to Carolina. ♀. Aug. Sept. v. v. Flowers small, rose-coloured.

militaris.

7. *H. glaberrimus; foliis 3-lobo-hastatis acuminatis serratis, corolla tubulato-campanulata, capsulis ovatis acuminatis glabris, seminibus holosericeis.* —*Willd. sp. pl. 3. p. 808.*

H. laevis. *Scop. del. insub. 3. t. 17.*

H. virginicus. *Walt. fl. car. 177.*

H. hastatus. *Mich. fl. amer. 2. p. 45.*

H. riparius. *Pers. syn. 2. p. 254.*

Icon. *Scop. del. insub. 3. t. 17.* *Cavan. diss. 6. t. 198. f. 2.*

On the banks of rivers in Louisiana and the western parts of Pennsylvania and Carolina. ♀. Aug. Sept. v. v. Flowers purple, large.

speciosus.

8. *H. glaberrimum; foliis palmato-5-partitis, lacinias lancolatis acuminatis apice remote serratis, calycibus exterioribus decaphyllis, corolla patula, capsula glabra 5-gono-ovata.* —*Willd. sp. pl. 3. p. 822.*

H. coccineus. *Walt. fl. car. 177.*

Icon. *Bot. mag. 360.* *Wendl. hort. herrenh. t. 11.*

On the banks of rivers of South Carolina and Florida. ♀. Aug. Sept. v. v. A most elegant peren-

nial. W
to more t
grow to a

9. *H. caule pe
7-partitis,
exteriore
tissima py*

Icon. *Cavan.*

On the bank

Flowers su
large whe

10. *H. caule sca
tione subr
dilatatis cr
dissimis ex*

H. aculeatus.

Near the sea-

July—Sep
bright yell

nial. When in high perfection, the stem will rise to more than eight feet, and the deep scarlet flowers grow to a larger size than any other Hibiscus I know.

9. H. caule petiolisque inermibus, foliis palmato-digitatis 7-partitis, lacinis linearibus, pedunculis hispidis, calyce exteriore 1-phyllo, interiore hinc lacero, capsula hirsutissima pyramidata.—*Willd. sp. pl. 3. p. 825.*

Icon. *Cavan. diss. 3. t. 63. f. 2. Dill. elth. t. 156. f. 189.*
On the banks of the Mississippi. $\frac{3}{4}$. July—Sept. v. v.
Flowers sulphur-yellow, with a purple bottom, very large when in rich soil.

10. H. caule scabro, foliis asperis basi truncatis circumscriptione subrotundis, superioribus palmatis, lobis superne dilatatis crenatis, floribus subsessilibus, calycibus hispidissimis exteriore furcellato. *Mich. fl. amer. 2. p. 45.*
H. aculeatus. *Walt. fl. car. 177.*
Near the sea-coast, in marshes: Carolina to Florida. $\frac{3}{4}$.
July—Sept. v. v. Flowers very handsome and large, bright yellow with a dark purple bottom.

Manihot.

scaber.

in
ul-

olis
llis.
us.

9.
une
lti-
ripe

ju-
ori-
bus
nu-

aro-
ose-

atis,
ina-
l. 3.

198.

parts
v. v.

lan-
ex-
abra

lori-
ren-

XVI. DIADEPHIA.

I. PENTANDRIA.

557. PETALOSTEMUM. *Pet.* 4. staminibus interjecta : utraque in tubum fissum connata. *Vexillum* nullum, ejus loco quintum petalum. *Legum.* calyce tectum, 1-spermum.

II. HEXANDRIA.

558. CORYDALIS. *Cal.* diphyllo. *Cor.* ringens. *Filam.* 2. membranacea, singula *Antheris* 3. *Caps.* siliquosa, polysperma.
559. FUMARIA. *Cal.* diphyllo. *Cor.* irregularis, basi calcarata. *Filam.* 2. singula *Antheris* 3. *Caps.* 1-locularis, evalus, 1-sperma.

III. OCTANDRIA.

560. POLYGALA. *Cal.* 5. phyllus : foliolis duobus alæ formibus, coloratis. *Caps.* obcordata, 2-locularis, 2-valvis.

IV. DECANDRIA.

* *Stamina omnia connexa.*

561. AMORPHA. *Cal.* campanulatus, 5-fidus. *Cor.* vexillum ovatum, concavum. *Alæ* *Carinaque* nullæ. *Legum.* 2-spermum, falcatum.
562. ERYTHRINA. *Cal.* 2-lobatus. *Cor.* vexillum longissimum, lanceolatum. *Legumen* torulosum.
563. LUPINUS. *Cal.* 2-labiatus. *Anth.* 5. oblongæ, 5. subrotundæ. *Legum.* coriaceum.
564. CROTALARIA. *Cor.* vexillum cordatum, magnum, carina acuminata. *Filam.* connata cum fissura dorsali. *Legum.* pedicellatum, turgidum.

** *Stigm*

565. PHASEOL
raliter torta
566. DOLICHO
alas subta

567. PISUM. C
plicas 2. p
supra vilic

568. LATHYRU
Stylus pla

569. VICIA. C
ferne dent
tum. *Sti*

*** *Legun*

570. ASTRAGA
aut subbile

571. OXYTROP
nens. *Leg*
superiore i

572. APIOS. C
illum refle
denticulat
men coriac
interceptui

**** *Legum*

573. DALEA. C
et *Carina* C
Stam. omni
brevius.

574. PORALE
diadelpha.

575. MELILOTI
simplex, a
gius, rugos

576. TRIFOI IU
4-spermum

577. LUPINAST
dentibus se
tum. *Leg*

578. STYLOSA
rollifer. C
culatum, h

** *Stigma pulescens.* (Nec priorum notæ.)

565. PHASEOLUS. *Carina* cum *Staminibus Styloque* spiraliter torta.
 566. DOLICHOS. *Vexilli basis callis* 2. parallelis oblongis,
alas subitus comprimentibus.
 567. PISUM. *Cal.* laciniae foliaceæ æquales. *Vexillum*
plicas 2. protrudens. *Stylus* compressus, carinatus,
supra villosus. *Legum.* ad suturas nudum.
 568. LATHYRUS. *Cal.* laciniae superiores 2. breviores.
Stylus planus, supra villosus, superne latior.
 569. VICIA. *Cal.* superne emarginatus, 2-dentatus, in-
ferne dentibus 3. rectis longis. *Vexillum* emarginatum.
Stigma latere inferiore transverse barbatum.

*** *Legumen subbiloculare.* (Nec priorum.)

570. ASTRAGALUS. *Carina* obtusa. *Legum.* biloculare
 aut subbiloculare, sutura inferiore introflexa.
 571. OXYTROPIS. *Carina* in mucronem superne desi-
 nens. *Legum.* biloculare aut subbiloculare, sutura
 superiore introflexa.
 572. APIOS. *Cal.* campanulatus, bilabiatus. *Carina vex-*
illum reflectens. *Appendix*: tubulus ex centro disci,
 denticulatus, vaginans basin stipitis germinali. *Legu-*
men coriaceum, biloculare, dissepimento seminibus
 interceptum.

**** *Legumina submonosperma.* (Nec priorum.)

573. DALEA. *Cal.* glandulosus, subsemi-5-fidus. *Alæ*
 et *Carina* columnæ staminum adnatæ. *Vexill.* breve.
Stam. omnia connata. *Legum.* 1-spermum, calyce
 brevius.
 574. PSORALEA. *Cal.* longitudine leguminis. *Stam.*
 diadelpha. *Legum.* 1-spermum, subrostratum, evalve.
 575. MELILOTUS. *Cal.* tubulosus, 5-dentatus. *Carina*
 simplex, alis et vexillo brevior. *Legum.* calyce lon-
 gius, rugosum. *Flores* racemosi.
 576. TRIFOLIUM. *Legum.* calyce tectum, evalve, 2—
 4-spermum. *Flores* subcapitati.
 577. LUPINASTER. *Cal.* campanulatus, 5-dentatus,
 dentibus setaceis, uno sub carina. *Stigma* uncina-
 tum. *Legum.* enode, teres, polyspermum.
 578. STYLOSANTHES. *Cal.* tubulosus, longissimus, co-
 rollifer. *Germen* sub corolla. *Lomentum* 1—2-arti-
 culatum, hamatum.

579. GLYCYRRHIZA. *Cal.* 2-labiatus, basi gibbus. *Cor.* vexillum alas et carinam includens. *Legum.* ovatum, compressum, hispidum, 2—4-spermum.
 580. LESPEDEZA. *Cal.* 5-partitus, lacinis subæqualibus. *Cor.* carina transverse obtusa. *Legum.* lenticulare, inerme, 1-spermum.

***** *Legumen in articulos secedens.* (Nec priorum.)

581. HEDYSARUM. *Cal.* 5-fidus. *Cor.* carina transverse obtusa. *Lomentum* pluri-articulatum : articulis 1-spermis, compressis, utplurimum hispidis.
 582. ZORNIA. *Cal.* campanulatus, 2-labiatus. *Vexillum* cordatum, revolutum. *Antheræ* alternae oblongæ, alternae globosæ. *Lomentum* articulatum, hispidum.
 583. ÆSCHYNOMENE. *Cal.* bilabiatus. *Lomentum* compressum, sutura altera recta, altera lobata, articulis truncatis, 1-spermis.
 584. SESBANIA. *Cal.* dentibus subæqualibus. *Legum.* elongatum, subcylindricum, 2-valve.

***** *Legumen uniloculare polyspermum.* (Nec priorum.)

585. GLYCINE. *Cal.* bilabiatus : dente labii inferioris longiore. *Cor.* carina apice vexillum reflectens. *Stylus* incurvus. *Legum.* oblongum, compressum.
 586. GALACTIA. *Cal.* 4-dentatus, 2-bracteatus. *Petalæ* omnia oblonga, vexillo latiore incumbente. *Stigm.* obtusum. *Legum.* teres. *Sem.* subrotunda.
 587. CLITORIA. *Cal.* tubuloso-campanulatus. *Cor.* resupinata : vexillo maximo, patente, alas obumbrante. *Legum.* lineare, longissimum, acuminatum.
 588. ROBINIA. *Cal.* 4-fidus, lacinia superiore bipartita. *Vexillum* reflexo-patens, subrotundum. *Legum.* gibbum, elongatum.
 589. INDIGOFERA. *Cal.* patens. *Cor.* carina utrinque calcari subulato patulo. *Legum.* lineare, parvulum, teres, subquadrangulare.
 590. TEPHROSIA. *Cal.* dentibus subulatis, subæqualibus. *Stam.* monadelpha. *Legum.* compressum, subcoriaceum.
 591. LOTUS. *Cal.* tubulosus. *Alæ* sursum longitudinale conniventes. *Filam.* cuneiformia. *Legum.* cylindricum, strictum.
 592. MEDICAGO. *Carina* corollæ a vexillo deflectens. *Legum.* compressum, cochleatum.

I.

557. PETALO

1. *P.* spica cylindrica calycis glomerata amer. 2. *p.* Dalea candida Icon. Mich. In Tennessee July, white.
 2. *P.* spica cylindrica tubine calycis glaberrima p. 49. In pine-barre v. s. in H...
 3. *P.* spica cylindrica æquantibus sericeis, fo p. 50. Dalea violacea Icon. Mich. In Tennessee July— spikes are
 4. *P.* spicis cylindrica calyce brevis jugis elliptica Dalea alopecu In Georgia at Enslen. F...
 5. *P.* capitulis secundis corymbosis paniculatis.—*Mic* Dalea Kuhnii Kuhnistera ca Kuhniae affini In pine-barre

I. PENTANDRIA.

557. PETALOSTEMUM. *Mich. fl. amer. 2. p. 48.*Dalea. *Juss. gen. pl. 392.*

1. *P. spica cylindrica pedunculata, bracteis flore longioribus, calycibus glabris, foliis trijugis lanceolatis.* — *Mich. fl. amer. 2. p. 49.*

*candidum.*Dalea candida. *Willd. sp. pl. 3. p. 1337.*Icon. *Mich. l. c. t. 37. f. 1.*

In Tennessee, Illinois, and on the banks of the Missouri.
U. July, Aug. v. s. in Herb. Lewis. &c. Flowers white.

2. *P. spica cylindrica pedunculata, bracteis subulatis longitudine calycis, bracteolis setaceis persistentibus, calycibus glabris, foliolis lanceolatis.* — *Mich. fl. amer. 2. p. 49.*

carneum.

In pine-barrens of Georgia and Florida. *U. July, Aug. v. s. in Herb. Enslen. &c.*

3. *P. spica cylindrica pedunculata, bracteis calycem subaequantibus, bracteolis spatulatis deciduis, calycibus sericeis, foliis bijugis linearibus.* — *Mich. fl. amer. 2. p. 50.*

*violaceum.*Dalea violacea. *Willd. sp. pl. 3. p. 1337.*Icon. *Mich. l. c. t. 37. f. 2.*

In Tennessee, Illinois, and on the banks of the Missouri.
U. July—Sept. v. v. A beautiful plant; the close spikes are of a delicate rose-purple.

4. *P. spicis cylindricis pedunculatis terminalibus, bracteis calyce brevioribus, calycibus villosis, foliis sub-10-jugis ellipticis retusis mucronatis.*

*alopecuroides.*Dalea alopecuroides. *Willd. sp. pl. 3. p. 1336.*

In Georgia and Florida. *U. July, Aug. v. s. in Herb. Enslen.* Flowers pale blue.

5. *P. capitulis squamoso-involutatis, pedunculis paniculato-corymbosis, calycibus plumositis, foliolis linearibus multicoris.* — *Mich. fl. amer. 3. p. 50.*

*corymbosum.*Dalea Kuhnistera. *Willd. sp. pl. 3. p. 1337.*Kuhnistera carolinensis. *Lam. encycl. 3. p. 371.*Kuhniae affinis pinnata. *Walt. fl. car. 103.*

In pine-barrens of Carolina and Georgia. *U. June—*

Aug. v. v. Flowers white. This singular plant appears, at first sight, to belong to the class Syngenesia, and certainly is the connecting link between that class and the present.

II. HEXANDRIA.

558. CORYDALIS. *Vent. choix de pl.* 19.

Fumaria. Gen. pl. 1154.

* *Corollis bicalcaratis.* Perizomanthi.

Cucullaria.

1. C. scapo nudo, racemo simplici secundo, nectariis divaricatis longitudine corollæ, stylo inclusu, radice bulbosa.

—*Pers. syn.* 2. p. 269.

Fumaria Cucullaria. Willd. sp. pl. 3. p. 857.

Icon. Bot. mag. 1127. *Rob. ic.* 222. *Pluk. alm. t.* 90. f. 3.

On the side of shady hills, in rich vegetable mould, among rocks: Canada to Virginia. 4. May. v. v. Flowers white, with a yellow mouth. This singularly constructed flower is known among the inhabitants by the name of *Breeches-flower* or *Yellow-breeches*; and as I consider that all those species with two spurs or nectaries might form, with propriety, a good genus, I have given it the name of *Perizomanthus*.

formosa.

2. C. scapo nudo, racemo subcomposito nutante multifloro, nectariis brevissimis incurvis, stylo porrecto, laciniis foliorum oblongis inciso-pinnatifidis, radice tuberosa.

Icon. Bot. rep. 393. *Bot. mag.* 1335. sub *Fumaria*.

In Canada, and on the highest peaks of the Virginia and Carolina mountains, in the cliffs of shady rocks. 4. May—July. v. v. Flowers a lively bright red. This elegant perennial deserves a place in every collection. The Canada plant is somewhat different in aspect from the Virginian, but not sufficiently distinct to make it a species.

tenuifolia.

3. C. scapo nudo, racemo simplici nutante paucifloro, nectariis brevissimis incurvis, stylo longe exerto, foliis tenuissime laciniatis, radice tuberosa.

On the north-west coast. 4. June. v. s. in *Herb.*

DIAI

Lambert.
same sha

4. C. caule sc
corymbo
—*Pers.* :

Fumaria fu

Fumaria re

In moist an

da and P

licate twi

abundanc

5. C. caule ra
racemis
nearibus
p. 269.

Fumaria se

Icon. Bot.

In the cliffs
mountain
handsom

6. C. caule rat
linear-lan
bracteis l.
bus turg
enun. 74

On shady ro

July. v.

the size c
ing specie

559

1. F. caule ran
cuneiform
p. 867. \

Icon. Fl. de

In cultivate

○. May

with gree

Lambert. specimen Pallasianum. Flowers of nearly the same shape and colour as the foregoing species.

4. C. caule scandente, foliis cirrhosis, racemis axillaribus corymbosis cernuis, corollis monopetalis basi bigibbis.
—*Pers. syn. 2. p. 269.*

fungosa.

Fumaria fungosa. *Willd. sp. pl. 3. p. 857.*

Fumaria recta. *Mich. fl. amer. 2. p. 51.*

In moist and shady places, in the beech-woods of Canada and Pennsylvania. ♂. June—Sept. v. v. A delicate twining plant; flowers pale red and in great abundance.

** *Corollis unicalcaratis.*

5. C. caule ramoso erecto, foliis glaucis, caulinis bibernatis, racemis subcorymbosis, bracteis minutis, siliquis linearibus pedunculo triplo longioribus.—*Pers. syn. 2. p. 269.*

glaucia.

Fumaria sempervirens. *Willd. sp. pl. 3. p. 863.*

Icon. Bot. mag. 179. Corn. canad. 57.

In the cliffs of rocks, in Canada, and on the Allegany mountains. ♂. June—Aug. v. v. Flowers very handsomely variegated with yellow, red, and green.

6. C. caule ramoso diffuso, foliis bipinnatis, foliolis partitis linear-lanceolatis utrinque acutis, racemis secundis, bracteis lato-lanceolatis subdenticulatis, siliquis teretibus turgidis pedunculo duplo longioribus.—*Willd. enum. 740.*

aurea.

On shady rocks: Pennsylvania to Virginia. ♂. May—July. v. v. Flowers bright yellow; the leaves and the size of the flowers resemble those of the preceding species.

559. FUMARIA. *Gen. pl. 1154.*

1. F. caule ramoso diffuso, foliis supradecompositis, foliolis cuneiformi-lanceolatis incisis.—*Willd. sp. pl. 3. p. 867.*

officinalis.

Icon. Fl. dan. 940. Engl. bot. 589.

In cultivated grounds: Pennsylvania to Virginia; rare. ♂. May—Aug. v. v. Flowers small, red, mixed with green.

III, OCTANDRIA.

560. POLYGALA. Gen. pl. 1154.

* *Cristatae*. Floribus appendice penicilliformi.

incarnata.

1. P. caule simpliciusculo erecto, foliis sparsis subulatis, spicis ovali-oblongis, corollis tubo gracili elongato.—Willd. sp. pl. 3. p. 871.

In low sandy fields and on the edge of rivulets : Jersey to Carolina. ☺. June, July. v. v. A delicate little plant ; flowers of a bright flesh-colour.

rubella.

2. P. pubescens ; caule simplici erecto, foliis angusto-lanceolatis acutis, racemo terminali, bracteis subulatis solitariis caducis, alis calycinis nervoso-venosis truncato-rotundatis corolla longioribus.

P. rubella. Willd. sp. pl. 3. p. 875 ?

In woods and on dry shady hills : Pennsylvania to Georgia. ♀. June, July. v. v. Flowers pale red. This species approaches near to *P. vulgaris*. Whether Willdenow's plant is the same or not, I am not able to determine ; but I strongly suspect he had but an imperfect specimen, in which the very minute bracteæ had disappeared. If his specimen has come from Dr. Muhlenberg, it certainly is the same plant.

paucifolia.

3. P. pumila, grandiflora ; caulibus simplicissimis erectis inferne nudis, foliis ovatis acutis glabris, floribus terminalibus subternis.—Willd. sp. pl. 3. p. 880.

In sphagnum swamps and bogs : Pennsylvania to Virginia ; principally on the mountains. ♀. May—Aug. v. v. Not above three or four inches high ; flowers the largest of those mentioned here, about the size of *P. myrtifolia*, of a fine pale red colour.

** *Imberbes*.

uniflora.

4. P. pumila ; caulibus simplicissimis erectis inferne nudisculis, foliis paucis lato-ovatis in petiolum angustatis, floribus solitariis sparsis pedicellatis nutantibus.—Mich. fl. amer. 2. p. 53.

On the banks of lake Ontario. Michaux. ♀. +.

Senega.

5. P. caulibus erectis simplicissimis foliosis, foliis alternis lanceolatis, spica terminali filiformi, floribus alternis. Willd. sp. pl. 3. p. 894.

8

Icon. Bot.

t. 2.

α. P. foliis l.

floribus :

β. P. glabella

laxiuscula

amer. 2.

On the side

throughout

and Geoc

famous i

the bite

remedies

Liatris.

6. P. caule sin-

spatulat-

pitata c

p. 894.

Icon. Pluk

α. P. caule sul-

β. P. subacaul

In wet wo

low mead

A fine sp

β. does r

7. P. caule e

obtusiusc

Willd. s

In dry wo

v. v. Fl

general h

8. P. caule se

moso, fo

dense spi

In Carolina

Lyon.

9. P. caule er-

ternis lit-

rosis.—B

Icon. Pluk

In woods o

lina. ☺

10. P. caule a

attenuatis

- Icon. *Bot. mag.* 1051. *Mill. dict.* 3. t. 5. *Amoen. acad.* 2. t. 2.
- α . P. foliis lanceolatis, vel ovalibus, spica confertiuscula, floribus albidis subsessilibus.—*Mich. fl. amer.* 2. p. 53.
- β . P. glabella vel pubescens; foliis linear-lanceolatis, spica laxiuscula alterniflora, floribus roseis.—*Mich. fl. amer.* 2. p. 53.
- On the side of hills and in dry woods: α . Canada and throughout the Allegany mountains. β . in Carolina and Georgia. ♂. June—Aug. v. v. This is the famous *Seneca Snake-root*, formerly so celebrated for the bite of rattle-snakes; but other more efficacious remedies have supplanted it. Vid. *Prenanthes* and *Liatris*.
6. P. caule simplici vel ramoso; foliis radicalibus infimisque spathulatis, cæteris lanceolatis, spica cylindraceo-capitata confertiflora pedunculata.—*Willd. sp. pl.* 3. p. 894.
- Icon. *Pluk. amalh.* t. 438. f. 6.
- α . P. caule subvirgatim elatiore.
- β . P. subacaulis, foliis omnibus spathulatis, capitulo majore. In wet woods and bogs: New Jersey to Florida. β . in low meadows of Carolina. ♂. July, Aug. v. v. A fine species; flowers of a golden yellow; the var. β . does not grow above three or four inches high.
7. P. caule erecto simplicissimo, foliis linear-lanceolatis obtusiusculis, capitulo terminali globoso-capitato.—*Willd. sp. pl.* 3. p. 895.
- In dry woods: Pennsylvania to Carolina. ♂. July, Aug. v. v. Flowers greenish-white, with some red. In general habit it resembles *P. incarnata*.
8. P. caule setaceo subaphyllo simplici, summitate subramoso, foliis parvis setaceis sparsis, floribus minutis dense spicatis.—*Mich. fl. amer.* 2. p. 52.
- In Carolina and Georgia. ♂. July, Aug. v. s. in Herb. Lyon.
9. P. caule erecto summitate corymboso-ramoso, foliis alternis linearibus, spicis capitatis, pedunculis squarrosis.—*Willd. sp. pl.* 3. p. 896.
- Icon. *Pluk. mant.* t. 438. f. 5.
- In woods on the side of hills: New England to Carolina. ♂. July, Aug. v. v. Flowers rose-coloured.
10. P. caule a basi ramoso, foliis alternis ellipticis obtusis basi attenuatis, spicis terminalibus oblongis, floribus alter-

*albida.**rosea.**lutea.**elatior.*
*nana.**viridescens.**setacea.**sanguinea.**polygama.*

nis pedunculatis, spicis inferioribus depressis apetalis.
—Walt. fl. car. 179.

In the pine-barrens of Carolina. ♂. July, Aug. v. s.
Calyx greenish-white, with pale yellow petals.

- verticillata.* 11. P. caule erecto ramoso, foliis verticillatis linearibus, spicis setaceis pedunculatis, floribus distincte alternis approximatis.—Willd. sp. pl. 3. p. 897.

Icon. Pluk. mant. t. 438. f. 4.

On the sides of dry hills and in pine-woods : New York to Carolina. ♂. June—Aug. v. v. Flowers very small, white, or sometimes pale red.

- cruciata.* 12. P. caule erecto ramoso alato-angulosso, foliis quaternis linear-lanceolatis, capitulis spicatis sessilibus.—Willd. sp. pl. 3. p. 897.

In wet places, on the edge of bogs and rivulets ; Canada to Carolina. ♂. July—Sept. v. v. Flowers red mixed with green.

A. nana. i
On the b
in Herb
purple a

3. A. humilis
mucronat
latis elc
dentibus
p. 970.

A. herbace
A. pumila
In sandy
July. fine blu

4. A. undiqu
matis s
spicis pa
dentibus
On the ba
Aug. v.
flowers

IV. DECANDRIA.

561. AMORPHA. Gen. pl. 1170.

fruticosa.

1. A. glabra, subarborescens ; foliis petiolatis, spicis aggregatis elongatis, calycibus nudiusculis pedicellatis, dentibus 4. obtusis, unico acuminato, leguminibus oligospermis.—Willd. sp. pl. 3. p. 970.

Icon. Schmidt arb. 30. Schkuhr handb. 197. Hort. cliff. t. 19. Mill. ic. 27. Duham. arb. t. 46.

α. A. foliis mucronatis, calycibus glabris.

β. A. foliis emarginatis, calycibus canis.

γ. A. foliis linear-ellipticis basi subacutis.

On the banks of rivers : Carolina and Florida. ♂.
June, July. v. v. Var. γ. v. s. in Herb. Lewis. A shrub from nine to twelve feet high. Flowers of a very fine blue, in close and long spikes. It is generally known by the name of *Wild Indigo*. The leaves, as in all the following species, are beset with glands, which are transparent.

microphylla.

2. A. glabriuscula, pumila ; foliis brevissime petiolatis utrinque obtusis, spicis solitariis abbreviatis, calycibus nudiusculis pedicellatis, dentibus omnibus acuminatis, leguminibus monospermis.

1. E. pumila
gissimis
p. 912.
Icon. Bot.
Dill. elt
In sandy
June, J
scarlet.

5

1. L. perenn
digitatis
cibus al
ginato ;
Icon. Bot.
On dry hil
May—J
times w

2. L. perenn
foliolis (

A. nana. Fraser. catal. 1813.

On the banks of the Missouri. $\frac{1}{2}$. July, Aug. v. s. in *Herb. Lewis*. From one to two feet high; flowers purple and fragrant. A very elegant little shrub.

3. *A. humilis*, frutescens; foliis brevissime petiolatis sine mucrone utrinque obtusis glabriusculis, spicis paniculatis elongatis pubescentibus, calycibus subsessilibus, dentibus omnibus acuminatis.—*Willd. sp. pl. 3. p. 970.*

*pulescens.**A. herbacea. Walt. fl. car. 179.**A. pumila. Mich. fl. amer. 2. p. 64.*

In sandy dry fields: Carolina and Georgia. $\frac{1}{2}$. June, July. v. v. From two to three feet high; flowers a fine blue, smaller than No. 1.

4. *A. undique cano-tomentosa*, humilis; foliolis approximatis subsessilibus ovato-ellipticis acutis mucronatis, spicis paniculatis cano-tomentosis, calycibus sessilibus, dentibus omnibus acuminatis.

canescens.

On the banks of the Missouri and Mississippi. $\frac{1}{2}$. July, Aug. v. v. Resembles the foregoing one very much; flowers of a more beautiful blue.

562. ERYTHRINA. *Gen. pl. 1163.*

1. *E. pumila*; foliis ternatis rhomboidis glabris, spicis longissimis, caule herbaceo inermi.—*Willd. sp. pl. 3. p. 912.*

herbacea.

Icon. Bot. mag. 877. Trew. ehret. t. 58. Catesb. car. 49. Dill. elth. t. 90. f. 106.

In sandy woods and fields: Carolina and Florida. 2 $\frac{1}{2}$. June, July. v. v. A beautiful perennial; flowers scarlet.

563. LUPINUS. *Gen. pl. 1176.*

1. *L. perennis*, repens; caule foliisque glabriusculis, foliis digitatis: foliolis (8-9) lanceolatis obtusiusculis, calycibus alternis inappendiculatis: labio superiore emarginato; inferiore integro.—*Willd. sp. pl. 3. p. 1022.*

perennis.

Icon. Bot. mag. 202. Mill. dict. t. 170. f. 1.
On dry hills, in gravelly soil: Canada to Florida. 4. May—July. v. v. Flowers blue, purple, and sometimes white.

2. *L. perennis*; caule foliisque hirsutis, foliis digitatis: foliolis (7-8) lanceolatis obtusis, calycibus verticillatis

inappendiculatis : labio superiore emarginato, inferiore integro.—*Donn. catal. hort. cantab.*

Icon. Bot. mag. 1311.

On the north-west coast. *Menzies.* ♀. June, July. *v. s. in Herb. Lambert.* This stately perennial approaches near to the foregoing one in its general character, but the flowers are larger and beautifully variegated. In the Herbarium of A. B. Lambert, Esq. is a specimen from Professor Rudolph, of St. Petersburg, collected in Unalashka, under the name of *L. regius*, decidedly the same with the present species, the name of which sufficiently shows the grandeur of this plant in its native state.

sericeus.

3. L. perennis ; caule foliisque sericeo-tomentosis, foliis digitatis : foliolis (7-8) lanceolatis acutis utrinque sericeis, calycibus subverticillatis inappendiculatis : labio superiore inciso ; inferiore integro.

On the banks of the Kooskoosky. *M. Lewis.* ♀. July. *v. s. in Herb. Lewis.* Flowers pale purple, or rose-coloured.

argenteus.

4. L. perennis ; foliis digitatis : foliolis (5-7) linear-lanceolatis acutis supra glbris subtus argenteo-sericeis, calycibus alternis inappendiculatis : labio superiore obtuso ; inferiore integro.

On the banks of the Kooskoosky. *M. Lewis.* ♀. June, July. *v. s. in Herb. Lambert.* Flowers small, cream-coloured.

pusillus.

5. L. biennis, pusillus, undique pilis longis albidis hirsutissimus ; foliis digitatis : foliolis (7) linear-ellipticis supra glbris, calycibus alternis inappendiculatis : labio superiore bipartito ; inferiore integro, leguminibus subdispermis.

On the banks of the Missouri. ♂. June. *v. s. in Herb. Lewis.* Flowers small, a fine blue mixed with some red.

villosus.

6. L. perennis, villosissimus ; foliis simplicibus oblongis, spicis elongatis, calycibus alternis inappendiculatis : labio superiore bifido ; inferiore integro elongato.—*Mich. fl. amer.* 2. p. 56.

L. pilosus. Walt. fl. car. 180.

In dry sandy fields : Carolina and Florida. ♀. June—Aug. *v. v.* A beautiful perennial, but very shy of cultivation ; flowers very variable in colour : white, rose red, and purple.

1. C. hirsuta,
lanceolat
rentibus,
lyce min
C. sagittalis
In pine-bar
Flowers,

2. C. hirsuta,
ceolatis,
bidentati
noribus.—
C. sagittalis
In sandy fie
v. v.

3. C. hirsuta,
subrotund
bus brevi
lis calyce
C. sagittalis
Anonymos
In pine-bari
v. s. in H

4. C. glabra,
stipulis la
oppositiflo
C. sagittalis
alm. 122.
In pine-woo

1. P. volubilis,
subcordat
dunculis
pendulis.—
182.

P. paniculata
Dolichos pol
On the side
among ro
v. v. ; v. s
abundance
appearanc

564. CROTALARIA. *Gen. pl.* 1172.

1. *C. hirsuta*, erecta, ramosa ; foliis simplicibus oblongo-lanceolatis, stipulis lanceolatis acuminate decurrentibus, racemis oppositifoliis subtrifloris, corollis calyce minoribus.—*Willd. sp. pl.* 3. p. 972.
C. sagittalis β . *oblonga*. *Mich. fl. amer.* 2. p. 55.
In pine-barrens : Virginia to Georgia. ☽. July. v. v.
Flowers, as all the following, yellow.
2. *C. hirsuta*, erecta, ramosa ; foliis simplicibus linearis-lanceolatis, stipulis superioribus decurrentibus brevissime bidentatis, racemis oppositifoliis, corollis calyce minoribus.—*Willd. sp. pl.* 3. p. 973.
C. sagittalis α . *linearis*. *Mich. fl. amer.* 2. p. 55.
In sandy fields : New York to Carolina. ☽. June, July.
v. v.
3. *C. hirsuta*, diffusa, ramosa ; foliis simplicibus petiolatis subrotundo-ovalibus, stipulis summis vix decurrentibus brevissimis, racemis oppositifoliis elongatis, corollis calycem æquantibus.
C. sagittalis γ . *ovalis*. *Mich. fl. amer.* 2. p. 55.
Anonymous *rotundifolia*. *Walt. fl. car.* 181.
In pine-barrens of Carolina and Georgia. ☽. June, July.
v. s. in *Herb. Enslen*.
4. *C. glabra*, erecta, simplex ; foliis lanceolato-oblongis, stipulis lanceolatis acuminate decurrentibus, racemis oppositifoliis subtrifloris.
C. sagittalis *glabra*, longioribus foliis, *americana*. *Pluk. alm.* 122. t. 277. f. 2.
In pine-woods of Virginia and Carolina. ☽. July. v. v.

565. PHASEOLUS. *Gen. pl.* 1180.

1. *P. volubilis*, totus pubescens ; foliolis lato-ovatis impari subcordato, racemis geminatis folio longioribus, pendulis geminatis, bracteis oboletis, leguminibus pendulis.—*Willd. sp. pl.* 3. p. 1031. *Walt. fl. car.* 182.
P. paniculata. *Mich. fl. amer.* 2. p. 60.
Dolichos polystachyos. *Willd. sp. pl.* 3. p. 1049.
On the side of fertile hills, near the banks of rivers, among rocks : Pennsylvania to Carolina. 4. July, Aug.
v. v.; v. s. in *Herb. Gronov.* Flowers purple, in great abundance, which gives this plant a very ornamental appearance.



vexillatus.

2. *P. volubilis*, pubescens; pedunculis petiolo crassioribus capitatis, alis subfalcatis difformibus, leguminibus linearibus strictis. *Willd. sp. pl. 3. p. 1032.*
Icon. *Jacq. hort. t. 102.* *Dill. elth. t. 234. f. 302.*
On the sea-coast of Carolina and Georgia. ☽. July, Aug. v. v. Flowers large, purple.

helvolus.

3. *P. volubilis*, perennis, hirtellus; foliolis oblongo-ovalibus subdeltoideis obtusiusculis, capitulis longe pedunculatis, vexillis brevibus, alis expansis maximis, leguminibus tereti-linearibus.—*Willd. sp. pl. 3. p. 1032.*
Glycine umbellata. Willd. sp. pl. 3. p. 1058.
Icon. *Dill. elth. t. 233. f. 300.*

In dry gravelly soil, in old fields and pastures: New York to Carolina. v. v. Flowers of a rose-colour, carina purple.

trilobus.

4. *P. volubilis*, pubescens; foliolis inferioribus rhomboideo-ovalibus, superioribus trilobatis, capitulis longe pedunculatis, leguminibus linearibus. *Mich. fl. amer. 2. p. 60.*
P. vexillatus. Walt. fl. car. 182.
Glycine angulosa. Willd. sp. pl. 3. p. 1056.
On banks of rivers in gravelly soil: Pennsylvania to Carolina. ☽. June, July. v. v. Flowers resembling the foregoing species.

566. DOLICHOS. *Gen. pl. 1181.**luteolus.*

1. *D. volubilis*, pubescens; foliolis ovatis acuminatis, pedunculis folio longioribus, spicis brevibus subcapitatis, vexillo lato reflexo, alis rhomboideis.—*Willd. sp. pl. 3. p. 1038.*
Icon. *Jacq. hort. t. 90.*
In rice fields of Georgia. *Enslen.* ☽. June, July. v. v.
Flowers bright yellow.

567. PISUM. *Gen. pl. 1184.**maritimum.*

1. *P. petiolis* supra planiusculis, caule angulato, stipulis sagittatis, pedunculis multifloris. *Willd. sp. pl. 3. p. 1071.*
Icon. *Fl. dan. 338. Engl. bot. 1046.*
In rocky and sandy situations, near the sea-coast: Canada to New York. ♀. July, Aug. v. v. Flowers large, handsome, purple and blue.

1. *L. caule ala sex linearis. Willd. sp. Icon. Fl. de In low grou v. s. Flc*
2. *L. caule nudum acutis mucronatis longioribus. In salt-mars larly plen v.v. Reser purple and*
3. *L. caule nudum minatis, mucronatis 10-floris. In low mead Flowers pi*
4. *L. caule tetragrammum 5-jugis: fulculis 3-4-fid. On the ban purple, lar*

1. *V. pedunculis misagittatis ceohatis obtusis.—Willd. In low grassy July, Aug. with a ting*
2. *V. floribus b notatis, foli guminibus sp. pl. 3. p. In Lower Ca purplish-bl*
3. *V. pedunculis VOL. II.*

568. LATHYRUS. *Gen. pl.* 1186.

1. *L. caule alato, stipulis semisagittatis lanceolatis, foliolis sex linear-lanceolatis acutis, pedunculis sub-3-floris.* — *Willd. sp. pl.* 3. p. 1090.
Icon. *Fl. dan.* 399. *Engl. bot.* 169.
In low grounds of Canada. *Michaux.* ♀. July, Aug.
v. s. Flowers pale purple, of middle size.
2. *L. caule nudo tetragono, stipulis semisagittatis lanceolatis acuminatis, foliolis quatuor oblongo-lanceolatis acutis mucronatis venoso-reticulatis, pedunculis folio longioribus sub-3-floris.* — *Willd. sp. pl.* 3. p. 1091.
In salt-marshes: Pennsylvania and New York; particularly plenty about Lake Onondago. ♀. July, Aug.
v. v. Resembles the foregoing species; flowers smaller, purple and rose-coloured.
3. *L. caule nudo tetragono, stipulis semisagittatis ovatis acuminatis, foliolis numerosis subalternis ovatis obtusis mucronatis venosis, pedunculis folio brevioribus 5—10-floris.* — *Willd. sp. pl.* 3. p. 1092.
In low meadows: Pennsylvania. ♀. July, Aug. *v. s.*
Flowers purple.
4. *L. caule tetragono, stipulis semisagittatis linearibus, foliis 5-jugis: foliolis oblongo-ellipticis mucronatis, pedunculis 3—4-floris.* — *Willd. sp. pl.* 3. p. 1093.
On the banks of the Missouri. ♀. *v. s.* Flowers purple, large; pods large.

569. VICIA. *Gen. pl.* 1187.

1. *V. pedunculis solitarii unifloris capillaribus, stipulis semisagittatis integerrimis, foliolis subsenis linear-lanceolatis obtusiusculis, leguminibus parvis oblongis glabris.* — *Willd. sp. pl.* 3. p. 1106.
In low grassy grounds: Pennsylvania to Virginia. ♂. July, Aug. *v. v.* Flowers exceeding small, white, with a tinge of red.
2. *V. floribus binis subsessilibus, stipulis dentatis macula notatis, foliolis oblongo-ovatis retusis mucronatis, leguminibus erectis subtereti-linearibus glabris.* — *Willd. sp. pl.* 3. p. 1104? *Walt. fl. car.* 183.
In Lower Carolina. ♂. July, Aug. *v. s.* Flowers purplish-blue.
3. *V. pedunculis submultifloris folio brevioribus, stipulis* *pusilla.*
v. v. s.

semisagittatis dentatis, foliolis 8—12. elliptico-lanceolatis obtusis glabris.—*Willd. sp. pl. 3. p. 1096.*

In Pennsylvania. *Muhlenberg.* ♀. +. Flowers the size of *V. sylvatica*, to which it bears a great resemblance.

Cracca.

4. *V. spicis multifloris retrorsum imbricatis folio longioribus, stipulis semisagittatis linearis-subulatis integerimis, foliolis numerosis lanceolato-oblongis pubescens-tibus.*—*Willd. sp. pl. 3. p. 1098.*

Icon. *Fl. dan.* 804. *Mart. fl. rust.* 117.

In woods and meadows: Canada to Virginia. ♀. July, Aug. v. v. Flowers pale blue.

caroliniana.

5. *V. pedunculis multifloris, floribus distantibus, stipulis ovato-lanceolatis integerrimis, foliolis 8—10. elliptico-lanceolatis glabriusculis, caule glabro.*—*Willd. sp. pl. 3. p. 1094.* *Walt. fl. car.* 182.

V. parviflora. *Mich. fl. amer.* 2. p. 69.

In the mountains from Pennsylvania to Carolina. ♀. July, Aug. v. v. Resembles the foregoing species; but the flowers are white, with a vexillum black on the tip, and a great deal smaller.

570. ASTRAGALUS. *Gen. pl.* 1208.

carolinianus.

1. *A. caulescens, erectus; foliolis (41) oblongis subtus pubescentibus, spicis pedunculatis, bracteis lanceolatis pedunculi longitudine, leguminibus ovatis tumidis rotatis.*—*Willd. sp. pl. 3. p. 1273.*

Icon. *Dill. elth.* t. 39. f. 45.

In the mountains of Virginia and Carolina. ♀. June, July. v. v. Flowers pale yellow.

canadensis.

2. *A. caulescens, diffusus; foliolis (21) utrinque glabris, leguminibus subcylindricis mucronatis.*—*Willd. sp. pl. 3. p. 1274.*

Icon. *Dodart. mem.* t. 65. *Rob. ic.* 314.

In the mountains: Canada to Carolina. ♀. July, Aug. v. v. Flowers yellow.

glaber.

3. *A. caulescens, glaber; foliolis lanceolato-oblongis subciliatis, spicis longe pedunculatis, floribus laxiusculis, leguminibus distantibus incurvis teretibus glabris.*—*Mich. fl. amer.* 3. p. 66.

In the sands of Georgia. *Michaux.* ♀. +. Flowers white.

alpinus.

4. *A. caulescens, procumbens; floribus pendulis racemosis,*

legu
p. 12
Icon.
In Labi
Dick
5. A. caul
spicis
que a
In the
maste
6. A. caule
pedur
minib
On the
v. s. is
7. A. subac
bris, l
iris, le
longis.
In sandy
in Her

571.

1. *O. acancli
scapis
ceolati
p. 331.*
Astragalu
Icon. *Pa*
On the ba
v. s. in

1. *A. volubil
natis: t
tatis, sp
Glycine A
Icon. Bot.
In hedges
Carolina
sweet-sc
an enorn*

- leguminibus utrinque acutis pilosis. *Willd. sp. pl. 3.*
p. 1297.
Icon. *Fl. dan. 51. Pall. astrag. t. 32.*
In Labrador. *Colmaster. 4. July, Aug. v. s. in Herb.*
Dickson. Flowers pale purple.
5. A. caulescens, procumbens; foliolis ovatis pubescentibus,
spicis pedunculatis, leguminibus secundis rectis utrinque
acuminatis pendulis.—*Mich. fl. amer. 2. p. 67.*
In the north of Canada. *Michaux. Labrador. Col-*
master. 4. July, Aug. v. s. Flowers dark purple.
6. A. caulescens, glaber; foliolis linearibus alternis, spicis
pedunculatis folio longioribus, bracteis minutis, legu-
minibus subteretibus rostratis.
On the banks of the Missouri. *M. Lewis. 4. Aug.*
v. s. in Herb. Lewis. Flowers small, yellowish-white.
7. A. subacutis, pilosissimus; foliolis ovalibus supra gla-
bris, pedunculis folia subæquantibus, spicis pauciflo-
ris, leguminibus incano-villosissimis assurgentibus ob-
longis.—*Mich. fl. amer. 2. p. 67.*
In sandy pine-woods of Georgia. *4. July, Aug. v. s.*
in Herb. Enslen.

*secundus.**tenellus.**vilosus.**argentata.*

571. OXYTROPIS. *Decand. astrag. p. 24.*
1. O. acanthis; foliolis obovato-lanceolatis sericeo-argenteis,
scapis folia æquantibus, floribus capitatis, bracteis lan-
ceolatis longitudine calycis sericei.—*Pers. syn. 2.*
p. 331.
Astragalus argentatus. Willd. sp. pl. 3. p. 1310.
Icon. *Pall. astrag. t. 48.*
On the banks of Clark's river. *M. Lewis. 4. July.*
v. s. in Herb. Lewis.

572. APIOS.

Glycine. *Gen. pl. 1182.*

1. A. volubilis, glabra; radice tuberosa, foliis impari-pinnatis: foliolis 5—7. ovato-lanceolatis superne angustatis, spicis confertis folio *brevioribus.*
Glycine Apios. Willd. sp. pl. 3. p. 1067.
Icon. *Bot. mag. 1198. Schkuhr handb. 198. Rob. ic. 317.*
In hedges and on mountain meadows: Pennsylvania to
Carolina. *4. July. v. v.* Flowers dark brown,
sweet-scented; roots eatable, growing sometimes to
an enormously large size.

tulerosa.

frutescens. 2. A. volubilis ; foliis impari-pinnatis novenis ovatis pubescentibus, racemis terminalibus bracteatis, leguminibus coriaceis.

Glycine frutescens. Willd. sp. pl. 3. p. 1067.

Anonymos frutescens. Walt. fl. car. 186.

In swamps of Virginia, Carolina, and Illinois. ♀. June, July. v. v. Flowers purplish-blue. It is a fine ornamental plant.

573. DALEA. Hort. cliff. p. 363. Mich. fl. amer. 3. p. 56.

Cliffortiana. 1. D. spicis oblongis confertis pedunculatis terminalibus sericeis, bracteis calycis longitudine, foliis subsexjugis linear-i-cuneatis retusis apice subdentatis.—Willd. sp. pl. 3. p. 1336.

Psoralea Dalea. Sp. pl. 1076.

Icon. Hort. cliffort. t. 22.

In Georgia and Florida. ♂. v. v. in Hortis. Flowers blue.

alopécuroïdes. 2. D. spicis cylindricis pedunculatis terminalibus sericeis confertis, bracteis calyce brevioribus, foliis 10—12-jugis ellipticis retusis mucronatis. Willd. sp. pl. 3. p. 1336.

D. Linnæi. Mich. fl. amer. 2. p. 57. t. 38.

On the banks of the Mississippi. ♂. July, Aug. v. v. Flowers pale blue.

pedunculata. 3. D. glabriuscula ; spicis oblongis axillaribus subconfertis brevibus, pedunculis folio triplo longioribus, bracteis minutis longitudine pedicellorum, dentibus calycis subulatis, foliis subsexjugis linearibus acutis mucronatis. On the banks of the Mississippi. Enslen. ♂. July, Aug. v. s. in Herb. Enslen. Flowers rose-coloured.

parviflora. 4. D. cano-pubescent; spicis filiformibus elongatis laxifloris axillaribus, pedunculis folio duplo longioribus, bracteis minutis, calycis tomentosi dentibus acutis brevibus, foliis sub-5-jugis linearibus obtusis.

On the banks of the Missouri. ♂. July, Aug. v. s. Flowers small, white.

vatis pube-
leguminini-

ly June,
a fine or-

. amer. 3.

nalibus se-
subsexjugis
-Willd. sp.

Flowers

us sericeis
is 10—12-
sp. pl. 3.

Aug. v. v.

abconfertis
s, bracteis
calycis su-
nucronatis.
July, Aug.
ed.

tis laxiflo-
longioribus,
acutis bre-

Aug. v. s.



Borago esculenta.

1. P. tota ex spicis la sissimis. In barren v. s. Re
2. P. parce p spicis ob minatis, —Mich Melilotus Trifolium In pine-bar v. s. in F
3. P. pubescen tpticis u bus folio On the bar v. s. in F
4. P. pubescen tis, petio bus dens vix longi On the ba Flowers
5. P. undique lanceolat bracteis silibus. On the ba purple : ticularly
6. P. undique ceolatis in bus dens breviorib fusiformi On the ba July. v. Tota planta crassa, emosus, te

574. PSORALEA. Gen. pl. 1210.

1. P. tota canescens; foliis breviter petiolatis trifoliatis, spicis laxifloris, floribus pedicellatis, calycibus villosissimis. *Mich. fl. amer. 2. p. 57.*
In barren sandy fields of Carolina and Georgia. July.
v. s. Resembles in appearance *Cytisus Cajan.* *canescens.*
2. P. parce pubescens; foliis ternatis: foliolis lanceolatis, spicis oblongis, bracteis lato-cordatis longissime acuminatis, leguminibus rotundatis nervoso-rugosissimis.
—*Mich. fl. amer. 2. p. 58.*
Melilotus flore violaceo, odore remisso. *Clayt. n. 103.*
Trifolium psoraloides. Walt. fl. car. 184.
In pine-barrens: Carolina to Florida. $\frac{1}{2}$. June—Aug.
v. s. in Herb. Leconte. Flowers blue. *melilotoides.*
3. P. pubescens, ramosissima; foliis ternatis: foliolis ellipticis utrinque rugoso-punctatis, pedunculis axillaris folio longioribus sub-3-floris.
On the banks of the Missouri. *M. Lewis.* $\frac{1}{2}$. Sept.
v. s. in Herb. Lewis. Flowers very small, pale blue. *tenuiflora.*
4. P. pubescens; foliis ternatis: foliolis elongato-lanceolatis, petiolis crassis, spicis axillaribus folio vix longioribus densifloris, floribus pedicellatis, bracteis pedicello vix longioribus, dentibus calycis coloratis.
On the banks of the Missouri. $\frac{1}{2}$. July, Aug. *v. s.*
Flowers bright blue, small. *lanceolata.*
5. P. undique argenteo-tomentosa; foliis ternatis: foliolis lanceolato-oblongis, spicis terminalibus interruptis, bracteis ovatis acuminatis, floribus suboppositis sessilibus.
On the banks of the Missouri. $\frac{1}{2}$. *v. s.* Flowers purple: the silvery tomentum gives this plant a particularly handsome appearance. *argophylla.*
6. P. undique villosa; foliis digitato-quinatis: foliolis lanceolatis inaequalibus planis integerrimis, spicis axillaribus densifloris, lacinias calycis lanceolatis corolla paulo brevioribus, leguminibus ensiformi-rostratis, radice fusiformi.
On the banks of the Missouri. *M. Lewis.* $\frac{1}{2}$. June, July. *v. s. in Herb. Lewis.* Flowers pale blue. *esculenta.*
Tota planta pilis longis albidis tecta. *Radix fusiformis, crassa, exsiccatione farinosa. Caulis erectus, subramosus, teres, sulcatus, pedalis et ultra. Folia alterna,*

petiolata, digitato-quinata. *Foliola* subpetiolata, lanceolata, utrinque acuta, integerrima, plana, supra glabra, punctis vix conspicuis, confertissime adspersa, subitus pilosa, uncialia. *Petioles* communes teretes, folio longiores. *Stipulae* persistentes, ad basin petiolorum oppositae, oblongo-lanceolatae, acutae. *Spicae* axillares, pedunculatae, densiflorae, bracteatae, foliis longiores. *Bracteae* lato-ovatae, acuminatae, calyx breviores. *Flores* sessiles, caerulei, magnitudine et figura *Viciae* sutive. *Calyx* hirsutissimus, 5-fidus s. profunde 5-dentatus: dentibus lineari-lanceolatis: 2. lateralibus subfalcatis: inferius latior. *Corolla*: *Vexillum* obovatum, inferne utrinque glandula instructum, calyx paulo longiore. *Alae* semi-ovatae, vexillo breviores. *Curva* oblonga. *Legumen* calyx tectum, monospermum, hirsutum, rostro ensiformi, glabro, calyx longiore instructum. *Semen* reniforme, nigrum.

It approaches very near to *P. pentaphylla* Willd. sp. pl. 3. p. 1352. which is fully described and figured by B. Jussieu in *Act. par.* 1744. p. 381. t. 17. But not only the botanical difference, but the medical use made of the Mexican plant, proves their distinctness. The present plant produces the famous *Bread-root* of the American Western Indians, on which they partly subsist in winter. They collect them in large quantities, and if for present use, they roast them in the ashes, when they give a food similar to yams: if intended for winter use, they are carefully dried, and preserved in a dry place in their huts. When wanted for use, they are mashed between two stones, mixed with some water, and baked in cakes over the coals. It is a wholesome and nourishing food, and, according to Mr. Lewis's observation, agreeable to most constitutions; which, he observed, was not the case with the rest of the roots collected by those Indians for food. This root has been frequently found by travellers in the canoes of the Indians, but the plant which produces it has not been known until lately.

Lupinellus.

7. *P. foliis quino-digitatis: foliolis angustissime linearibus, spica pauciflora, legumiibus ovoideis, uncinato-mucronatis, nervoso-rugosis.* — *Mich. fl. amer.* 2. p. 58.

In barren fields of Carolina. $\frac{1}{4}$. v. s. in *Herb. Lyon.* Flowers very small. The leaves of the whole genus are covered with resinous dots.

575.

1. *M. cat*
bus
Trifoli
Icon.
On th
nia.

1. *T. pr*
obliqu
cella
sperz
On dry
and
Flow

2. *T. repe*
tis, c
quali
p. 13
Icon. F
In field
 $\frac{1}{4}$. M
by th
where
mote
forwa
dred
was f
have
cum
whose

3. *T. erect*
bris n
lis um
rollis
fl. am
T. arven
T. como
In sandy
June—
encycl

575. MELILOTUS. *Tournef. inst. p. 407. t. 229.*

1. *M. caule erecto, foliolis obovatis serratis, spicis axillariibus paniculatis, leguminibus dispermis rugosis acutis.*
Trifolium officinale. Willd. sp. pl. 3. p. 1355.
Icon. Fl. dan. 934. Mart. fl. rust. t. 72.
 On the gravelly shores of rivers: Pennsylvania to Virginia. ♂. ○. June, July. v. v. Flowers white.

576. TRIFOLIUM. *Gen. pl. 1211.*

1. *T. procumbens, pubescens; foliolis obovatis, stipulis oblique cordatis, capitulis multifloris, floribus pedicellatis demum omnibus reflexis, leguminibus sub-3-spermis.*—*Willd. sp. pl. 3. p. 1357.*
 On dry hills and among rocks: Pennsylvania, Kentucky, and the Carolina mountains. ♀. June, July. v. v.
 Flowers of a fine purplish-red.
2. *T. repens; foliolis ovato-oblongis emarginatis serrulatis, capitulis umbellaribus, calycinis dentibus subæqualibus, leguminibus tetraspermis.*—*Willd. sp. pl. 3. p. 1359.*
Icon. Fl. dan. 990. Mart. fl. rust. 34.
 In fields, pastures, and cleared lands on the mountains. ♀. May—Sept. v. v. This species, generally known by the name of *White Clover*, makes its appearance wherever lands are cleared of timber, in the most remote parts of the country; I have seen it springing forward on new plantations, from sixty to one hundred miles distant from any place where it naturally was found. A number of plants of this description have struck my attention, among which are *Verbas-cum Thapsus* and *Senecio hieracifolius* in particular, whose dissemination is really problematical.
3. *T. erectum, pusillum; foliolis rotundato-obcordatis glabris marginè ciliatis crenulatis, stipulis bifidis, capitulis umbellaribus pedunculatis reflexis paucifloris, corollis vix exsertis, leguminibus subtrispermis.*—*Mich. fl. amer. 2. p. 58.*
T. arvense. Walt. fl. car. 183.
T. comosum. Willd. sp. pl. 3. p. 1860?
 In sandy fields, near Charlestown, South Carolina. ♀. June—Aug. v. s. Flowers small, white. In *Lam. encycl. 8. p. 2.* it has been suggested, that *T. como-*

sum of Linnæus and the present plant are the same, and I believe with very good reason, as I suppose the little difference between them may have been occasioned by culture.

microcephalum.

4. T. adscendens, pubescens; foliolis obovatis emarginatis denticulatis, stipulis ovatis acuminatis, capitulis pusillis pedunculatis paucifloris, floribus sessilibus, involucris ovatis aristatis, calycibus aristatis longitudine collæ.

On the banks of Clarek's river. *M. Lewis.* ♀. July. v. s. in *Herb. Lewis.* Flowers exceeding small, pale purple.

pratense.

5. T. adscendens, glabriuscum; foliolis ovalibus subintegerrimis, stipulis aristatis, spicis densis ovatis, calycis dente infimo tubo corollæ monopetalæ inæqualis breviore.—*Willd. sp. pl. 3. p. 1366.*

Icon. *Fl. dan. 989. Mart. fl. rust. 3.*

In fields and woods, frequent. ♀. May—Sept. v. v. Common *Red Clover* is apparently in a native state, but very probably has been introduced.

pennsylvanicum.

6. T. adscendens; caule ramosissimo flexuoso, foliolis ovato-ellipticis obtusis integerrimis, stipulis aristatis, spicis ovato-cylindraceis solitariis densis, calycis dente infimo corollæ monopetalæ breviore.—*Willd. enum. p. 793.*

In woods and fields: Pennsylvania to Virginia. ♀. June — Sept. v. v. Resembles *T. medium* very much; flowers a fine red. It is known by the name of *Buffalo Clover.*

arvense.

7. T. erectum, ramosissimum, villosum; foliolis linearilanceolatis apice serrulatis, stipulis connatis apice subulatis, spicis villosissimis subcylindraceis, dentibus calycinis setaceis corolla longioribus.—*Willd. sp. pl. 3. p. 1373.*

Icon. *Fl. dan. 724. Curt. fl. lond. 50.*

In dry and sandy fields: Canada to Virginia. ♂. July, Aug. v. v. Flowers small, white; the wings have a red spot.

agrarium.

8. T. erectum, subpubescens; foliolis lanceolato-cuneatis obtusis: intermedio sessili, stipulis lanceolatis acutis, spicis ovalibus imbricatis, vexillis deflexis persistentibus, calycinis dentibus subulatis glabris inæqualibus — *Willd. sp. pl. 3. p. 1382.*

T. aureum. *Pollich. fl. palat. n. 708.*



Tigemaster macrocephalus.

DIADELE

- T. strepens. Cr.
Icon. Fl. dan.
In meadows :
v. v. Flowe
9. T. procumbens
marginatis gl
spicis ovalib
bus sulcatis.
Icon. Fl. dan.
On dry gravelly
—Aug. v.
the foregoing

577. LUPINA

Pentap

1. L. foliolis nove
pulis cuneati
formibus plu
At the head-w
April, May.
white, with
showy : the
tremely large
Caulis plures,
suprema opp
lanceolata, i
glabra, subt
tioli inferiore
tiolo adnata
apice inciso-
et pubescent
diatum. *Flor*
campanulatu
dentibus su
quantibus. O
ceolatum, a
volvens. *A*
cinata, integ
diadelpha :
monadelpha
brum. *Sty*
Legumen 3—

T. strepens. Crantz. fl. austr. 411. n. 8.

Icon. *Fl. dan. 558.*

In meadows : Pennsylvania to Virginia. ♂. June—Aug.
v. v. Flowers small, yellow.

9. *T. procumbens*; caule piloso; foliolis obovatis sub-
marginatis glabris, stipulis brevibus lanceolatis acutis,
spicis ovalibus imbricatis, vexillis deflexis persistenti-
bus sulcatis.—*Willd. sp. pl. 3. p. 1383.*

Icon. *Fl. dan. 796.*

On dry gravelly soil : Pennsylvania to Virginia. ♂. June
—Aug. v. v. Flowers yellow, resembling those of
the foregoing species.

577. LUPINASTER. *Moench. meth. suppl. p. 50.*

Pentaphyllum. *Pers. syn. 2. p. 352.*

1. *L. foliolis novenis* ob lanceolatis, petiolis longissimis, sti-
pulis cuneatis inciso-3-dentatis, calycinis dentibus fili-
formibus plumosis. *macrocephalus.*

At the head-waters of the Missouri. *M. Lewis.* ♀.
April, May. v. s. in *Herb. Lewis.* Flowers yellowish-
white, with a pale purple carina, very handsome and
showy : the flower-heads sometimes grow to an ex-
tremely large size.

Caulis plures, simplices, angulosi, pilosi. *Folia* petiolata,
suprema opposita, novena. *Foliola* inaequalia, cuneato-
lanceolata, inaequaliter mucronato-denticulata, supra
glabra, subtus pilosa, lineato-venosa, uncilia. *Pe-*
tio inferiores longi, subteretes. *Stipulae* inferiores pe-
tiolo adnatae, superiores liberæ, cuneato-oblongæ,
apice inciso-3-dentatae, margine serrulatae, consistentia
et pubescencia foliorum. *Capitulum* speciosum, di-
diatum. *Flores* lutescentes, carina purpurascente. *Cal.*
campanulatus, subaequaliter 5-dentatus, villosissimus :
dentibus subulatis piloso-plumosis corollam subae-
quantibus. *Corolla*: *Vexillum* magnum, oblongo-lan-
ceolatum, acutum, cæteras partes fructificationis in-
volvens. *Alæ* carinæ adnatae, lineares. *Carina* un-
cinata, integra, vexillo vix breviore. *Filamenta* 10.
diadelpha : 1. liberum, reliqua superne libera, inferne
monadelpha, carina adnata. *Germen* oblongum, gla-
brum. *Style* filiformis. *Stigma* simplex, hirsutum.
Legumen 3-4-spermum, inflatum, calyce longius.

578. STYLOSANTHES. *Gen. pl.* 1203.

- hispidia.* 1. *S. caule uno latere pubescente, foliolis lanceolatis glabris, bracteis lanceolatis ciliatis, capitulis 2-3-floris.* — *Mich. fl. amer. 2. p. 75.*
Trifolium biflorum. Sp. pl. 1088.
Arachis aprica. Walt. fl. car. 182.
- erecta.* α. *S. caule simplici erecto superne breviter ramoso.*
Icon. Swartz. act. holm. 1789. t. 11. f. 2.
- procumbens.* β. *S. caule procumbente a basi ramoso.*
 In dry gravelly fields and woods; Pennsylvania to Carolina. ♀. July, Aug. v. v. Flowers yellow, small.

579. GLYCYRRHIZA. *Gen. pl.* 1197.

- lepidota.* 1. *G. foliolis oblongis acutis sericeo-villosis, leguminibus ramosis oblongis hispidis.* — *Fraser. catal.*
 On the banks of the Missouri. ♀. July, Aug. v. s. v. s. fl.

580. LESPEDEZA. *Mich. fl. amer. 2. p. 70.*

- sessiliflora.* 1. *L. erecta; foliolis oblongis, fasciculis florum sessilibus numerosis, lomentis calyce minuto suboudatis acutis.*
Mich. fl. amer. 2. p. 70.
Hedysarum junceum. Walt. fl. car. 184.
Hedysarum sessiliflorum. Lam. encycl. 6. p. 414.
Hedysarum reticulatum. Willd. sp. pl. 3. p. 1194.
Medicago virginica. Syst. veg. 3. p. 573.
 In old fields and on the borders of woods: New York to Carolina. ♀. July, Aug. v. v. Flowers purple.
- capitata.* 2. *L. erecta, simplex; foliis subsessilibus: foliolis ellipticis, spicis capitatis brevi-pedunculatis axillaribus et conglobato-terminalibus, calycibus villosis longitudine corollæ lomentis calyce multo minoribus.* *Mich. fl. amer. 3. p. 71.*
Hedysarum conglomeratum. Lam. encycl. 6. p. 416.
 β. *L. foliolis linearibus, pedunculis longioribus.*
Trifolium erectum haud ramosum, foliis longis, &c.
Clayt. n. 934.
 In sandy soil, in old fields and borders of woods: New York to Carolina. ♀. June, July. v. v. From one to three feet high; flowers pale purple.
- polystachya.* 3. *L. erecta, ramosa, villosissima; foliis subsessilibus: foliolis rotundato-ovalibus, spicis axillaribus longe-pe-*

dun
lyce
Hedys
Icon.
In cop
June
flow
4. L. diffi
ellip
vibu
beis
Hedysa
On gra
v. v.
5. L. diffi
press
minib
Hedysa
In Nort
very
only
6. L. proc
pedat
bus c
Hedysa
In sand
June,
low s
7. L. prost
subtu
lio loi
Hedysa
In Pens
dysari
58
1. H. erect
ceolat
culis
pl. 3.
Icon. M
In sandy
♀. Ju
hands

dunculatis, corollis calycet tubaequalibus, lomento calyce subæquante.—*Mich. fl. amer.* 3. p. 71.

Hedysarum hirtum. *Willd. sp. pl.* 3. p. 1193.

Icon. *Mich. fl. amer.* 2. t. 40.

In copses and old fields: Pennsylvania to Carolina. ♀.

June—Aug. v. v. From two to three feet high; flowers small, white, with a red spot on the vexillum.

4. L. diffusa, ramosissima; foliis longe-petiolatis: foliolis ellipticis obtusis subtus adpresso-pilosus, racemis brevibus umbellatis, floribus geminatis, lomentis rhomboides reticulatis glabris.

Hedysarum violaceum. *Willd. sp. pl.* 3. p. 1195.

On gravelly dry soil: Virginia to Carolina. ♀. July. v. v. Flowers small, purple or violet.

5. L. diffusa, ramosa; foliolis oblongis obtusis subtus adpresso-pilosus, racemis petiolo longioribus, floribus geminis, lomentis ovatis reticulatis glabris.

Hedysarum divergens. *Willd. sp. pl.* 3. p. 1196.

In North America. *Muhlenberg.* ♀. +. Approaches very near to the foregoing species, and is probably only a variety.

6. L. procumbens, gracilis, pubescens; foliolis ovalibus, pedaneulis longissimis, setaceis spicifloris, leguminibus calyce minuto nudatis.—*Mich. fl. amer.* 3. p. 70.

Hedysarum Lespedeza. *Lam. encycl.* 6. p. 415.

In sandy and gravelly soil: New York to Carolina. ♀. June, July. v. v. Flowers violet-purple, with yellow spots.

7. L. prostrata; foliis petiolatis: foliolis ellipticis obtusis subtus pilis rarissimis tecta, racemis axillaribus folio longioribus, lomentis ovatis pubescentibus.

Hedysarum prostratum. *Willd. sp. pl.* 3. p. 1200.

In Pennsylvania. *Muhlenberg.* ♀. +. Resembles *Hedysarum repens* greatly.

violacea.

divergens.

prostrata.

581. HEDYSARUM. *Gen. pl.* 1204.

1. H. erectum, glabriuscum; foliis ternatis oblongo-lanceolatis, stipulis filiformibus, floribus racemosis, articulis lomenti obtuse triangulis hispidis.—*Willd. sp. pl.* 3. p. 1187.

Icon. *Moris. hist.* 2. s. 2. t. 11. f. 9.

In sandy barren fields and woods: Canada to Carolina. ♀. July, Aug. v. v. Flowers middle-sized, of a handsome purplish-red.

canadense.

canescens.

2. *H. erectum, pilosum; caule angulato-ciliato hispido, foliis ternatis subrotundis subtus adpresso-pilosis, stipulis ovatis acuminatis, racemis paniculatis, bracteis cordatis, lomenti articulis triangularibus hispidis.*—*Willd. sp. pl. 3. p. 1188.*

In dry barren woods: Virginia and Carolina. $\frac{1}{2}$. June—Aug. *v. v.* Flowers middle-sized, pale purple.

marylandicum.

3. *H. erectum, ramosum, pilosum; foliis ternatis oblongis subtus villosis, stipulis subulatis, racemis paniculatis, lomentis triarticulatis, articulis rhomboides reticulatis pilosiusculis.*—*Willd. sp. pl. 3. p. 1189.*

Icon. Dill. elth. t. 174. f. 171.

In dry swamps and woods: New York to Carolina. $\frac{1}{2}$. June—Aug. *v. v.* Flowers large, a fine purple, with a green spot at the bottom of the vexillum, which mark occurs in most species.

obtusum.

4. *H. erectum, pubescens; foliis ternatis ovatis obtusis basi subcordatis, stipulis lanceolato-subulatis, panicula terminali, articulis lomenti semiobtusatis reticulatis hispidis.*—*Willd. sp. pl. 3. p. 1190.*

In dry swamps, old fields and woods: Pennsylvania to Virginia. $\frac{1}{2}$. July, Aug. *v. v.* Flowers violet.

viridiflorum.

5. *H. erectum, ramosum, pubescens; foliis ternatis ovato-oblongis subtus scabris, stipulis lanceolato-cuspidatis, racemis paniculatis bracteatis, lomentis asperis, articulis ovalibus.*—*Willd. sp. pl. 3. p. 1192.*

Icon. Pluk. alm. t. 308. f. 5.

In woods and old fields: New Jersey to Carolina. $\frac{1}{2}$. July—Sept. *v. v.* About three feet high; flowers reddish-purple, which turns green after flowering.

glabellum.

6. *H. erectum, glabrum; foliis ternatis ovatis obtusis subtus subglauca, stipulis parvis subulatis, panicula terminali, lomenti articulis triangulo-rhomboides.*—*Mich. fl. amer. 3. p. 73.*

In fields and woods: Virginia and Carolina. $\frac{1}{2}$. July, Aug. *v. v.* Flowers purple, small.

bracteosum.

7. *H. erectum, glabrum; foliis ternatis oblongo-ovalibus acuminatis, stipulis subulatis, racemo terminali sparsifloro, bracteis ovatis acuminatis striatis glabris, lomenti articulis subovalibus.*—*Mich. fl. amer. 2. p. 73.*

In the western parts of Pennsylvania and Virginia. $\frac{1}{2}$. July—Sept. *v. v.* Flowers large, purplish-red.

ciliare.

8. *H. erectum, glabriuscum; foliis ternatis ovatis subtus*

DI

pubes
panic
culatis
In North
violet.

9. *H. erect
vel ell
nali, l
Willd.
Icon. Pl
In dry w
June—*

10. *H. strict
ternatis
tus gl
pedunc
In pine-
Flower*

11. *H. erect
oblong
pulis c
loment
scentib
In shady
Aug.*

12. *H. erectu
foliis te
adpresso
to-subr
duncula
In shady
Aug.
It appro
connect*

13. *H. erectu
ovalibus
caule fo
angulari
In woods
June—
purple.*

14. *H. erectu
rotundo*

- pubescentibus margine ciliatis, stipulis filiformibus,
panicula terminali, lomenti articulis (2-3) semioblique
culatis hispidis.—*Willd. sp. pl. 3. p. 1196.*
In North America. *Muhlenberg.* ♀. +. Flowers small,
violet.
9. *H. erectum, glabrum; foliis ternatis oblongo-lanceolatis paniculatum.*
vel ellipticis glabris, stipulis subulatis panicula terminali, lomenti articulis (4) rhomboides pubescentibus.—
Willd. sp. pl. 3. p. 1196.
Icon. *Pluk. mant. t. 432. f. 6.*
In dry woods and fields: New York to Carolina. ♀.
June—Aug. v. v. Flowers purple.
10. *H. stricte-erectum, glabrum, simplex; foliis petiolatis ternatis linear-ellipticis glabris reticulato-venosis subtus glaucis, stipulis subulatis, panicula terminali pedunculata pauciflora.*
In pine-woods of New Jersey. ♀. July, Aug. v. v.
Flowers small, pale purple; seeds not known.
11. *H. erectum, glabrum; foliis ternatis petiolatis ovato-oblongis longe-acuminatis glabris margine scabris, stipulis ovato-lanceolatis, panicula terminali, articulis lomenti triangularibus reticulatis glabris margine pubescentibus.* —*Willd. sp. pl. 3. p. 1198.*
In shady woods: Pennsylvania and Virginia. ♀. July,
Aug. v. v. Flowers violet.
12. *H. erectum, simplex, pubescens, summitate frondosum; foliis ternatis ovatis longe acuminatis utrinque rarer adpresso-pilosis longissime petiolatis: impari rotundato-subrhomboideo, panicula terminali longissime pendulata.* —*Mich. fl. amer. 2. p. 72.*
In shady woods: New England to Carolina. ♀. July,
Aug. v. v. Not above a foot high; flowers purple.
It approaches near the foregoing species, and is the connecting link between that and the following.
13. *H. erectum, simplex, glabriuscum; foliis ternatis latovalibus acuminatis, scapo paniculato glabro radicali caule foliifero altiore, lomenti articulis subrotundo-triangularibus glabriusculis.* —*Willd. sp. pl. 3. p. 1198.*
In woods and old fields: New England to Carolina. ♀.
June—Sept. v. v. About a foot high; flowers purple.
14. *H. erectum, simplex; foliis longe petiolatis ternatis subrotundo-ovatis acuminatis, panicula scapiformi e caulis nudiflorum.*
rotundo-ovatis acuminatis, panicula scapiformi e caulis *glutinosum.*

basi, pedunculis piloso-viscosis, lomenti articulis oblongo-triangularibus glabriusculis.—*Willd.* sp. pl. 3. p. 1198.

In dry shady places, in the western parts of Pennsylvania and Virginia. ♀. July, Aug. v. v. Flowers purple.

lineatum.

15. H. caule repente viridi lineato, foliis subsessilibus ternatis suborbiculatis, racemis elongatis laxe parvifloris, lomenti articulis lenticularibus.—*Mich. fl. amer.* 2. p. 72.

In Carolina. *Michaux.* ♀. +.

*rotundifoli-
um.*

16. H. prostratum, hirsutum; foliis ternatis orbiculatis utrinque pilosis, stipulis rotundato-cordatis reflexis, racemis axillaribus et paniculato-terminalibus paucifloris, bracteis cordatis, lomenti articulis subrhomboideis reticulatis scabris.—*Mich. fl. amer.* 2. p. 72.

In rocky, dry situations: Pennsylvania to Carolina. ♀. July, Aug. v. v. A handsome species: flowers purple.

*alpinum.
β. americanum.*

17. H. erectum; foliis pinnatis: foliolis ovali-oblongis pilosis, stipulis subvaginantibus, leguminibus patulis glabris ex utraque sutura articulatis: articulis brevi-ovalibus. *Mich. fl. amer.* 2. p. 74.

In the north of Canada, and near the cataracts of the Allegany mountains. *Michaux.* ♀. +. Flowers purple.

582. ZORNIA. *Mich. fl. amer.* 2. p. 76. *Gmel. syst. veg.* 1096.

tetraphylla.

1. Z. foliis digitato-quadrifoliatis: foliolis lanceolatis glabris, spicis axillaribus pedunculatis, floribus alternis bibracteatis, bracteis suborbiculatis.—*Mich. fl. amer.* 2. p. 76.

Z. bracteata. *Gmel.*

Hedysarum tetraphyllum. Lam. encycl. 6. p. 405.

Anonymos bracteata. Walt. fl. car. 181.

Icon. Mich. fl. amer. 2. t. 41.

In sandy fields of Lower Carolina. ♀. July, Aug. v. s. in *Herb. Enslen.* About a foot high, very branching; flowers yellow.

583.

1. A. caule

7—9. c
bescen-
mer. 3.

On the o
island,

2. A. caule

multiju-
branace-
ris, lori-
Hedysarum
On the ba-
as the ti-
Aug. 1
finely ve-

584. SES.

1. S. glabra;

glabris n-
ris folio-
sis longe-
S. platycar-
Robinia ve-
Phaca florid-
Æschynom-
In overflow-
Flowers:

58

1. G. caule vc

mis filifo-
nibus obl-
G. monoica

In low grou-

○. July,
a separa-
show the
Glycine.

2. G. caule gr

glabris me-
corollatis

583. ÆSCHYNOMENE. *Gen. pl. 1202.*

1. A. caule prostrato gracili viscido-pubescente, foliolis 7—9. obovalibus, pedunculis subbifloris, lomento pubescente profunda incisura articulata.—*Mich. fl. amer. 3. p. 74.*

On the overflowed sands of Florida, and Cumberland island, Georgia. *Michaux.* ♂. +.

2. A. caule erecto petiolisque tuberculato-hispido, foliis multijugis: foliolis linearibus obtusis, stipulis membranaceis semisagittatis, racemis simplicibus paucifloris, lomentis hispidis.—*Willd. sp. pl. 3. p. 1163.*

Hedysarum virginicum. Willd. sp. pl. 3. p. 1212.
On the banks and islands of the principal rivers, as far as the tide flows: Pennsylvania to Carolina. ♂. July, Aug. v. v.; v. s. in *Herb. Gronov.* Flowers yellow, finely veined with red.

584. SESBANIA. *Poiret in lam. encycl. 7. p. 126.*

1. S. glabra; foliis sub-20-jugis: foliolis oblongis obtusis glabris margine ciliatis, racemis pedunculatis paucifloris folio brevioribus, lomentis lato-lanceolatis compressione longe stipitatis subdispermis.

S. platycarpa. Pers. syn. 2. p. 316.

Robinia vesicaria. Jacq. ic. rar. 1. t. 48.

Phaca floridana. Pers. syn. 2. p. 331.

Æschynomene platycarpa. Mich. fl. amer. 2. p. 75.

In overflowed places: Carolina and Florida. ♂. v. s.
Flowers yellow.

viscidula.

hispida.

disperma.

585. GLYCINE. *Gen. pl. 1182.*

1. G. caule volubili, foliis ternatis ovatis acutis glabris, racemis filiformibus sub-3-floris, floribus apetalis, leguminibus oblongis dispermis.—*Willd. sp. pl. 3. p. 1055.*

G. monoica. Schkuhr bot. annal. 12. p. 20. t. 2.

In low grounds, among bushes: Virginia and Carolina. ♂. July, Aug. v. v. It might with propriety form a separate genus, if the following species did not show the strong claim it has to be still considered a *Glycine.*

sarmentosa.

2. G. caule gracili retrosum piloso, foliis ternatis ovatis glabris membranaceis, racemis caulinis pendulis laxis corollatis plerumque sterilibus, bracteis suborbiculatis

monoica.

concavis, pedunculis radicalibus floribus apetalis fructiferis.—*Willd. sp. pl. 3. p. 1055.*

G. bracteata. Sp. pl. p. 754.

On the banks of rivulets in stony and shady places: New York to Carolina. ♂. July—Sept. v. v. The flowers on the stem resemble those of *Vicia Cracca*, they are pale blue, and seldom produce ripe fruit; the long radical tendrils have exceeding small flowers which produce a pod, generally with only one seed, which drops into the ground before maturity.

comosa.

3. *G. caule volubili, foliis ternatis ovato-lanceolatis acutis hirsutis, racemis lateralibus pendulis confertifloris, floribus omnibus fructiferis.—Willd. sp. pl. 3. p. 1058.*

In hedges, near rivers and ditches: New York to Virginia. ♀. July—Sept. v. v. Flowers blue, of the size and shape of the preceding species, to which it has a strong resemblance, but differs in being a perennial and robust plant, besides the other characters given.

tomentosa.

4. *G. caule volubili anguloso, foliis ternatis ovato-oblongis acutis pubescentibus, subtus rugoso-venosis tomentosis, racemis axillaribus petiolo brevioribus, leguminibus oblongis dispermis.—Willd. sp. pl. 3. p. 1061.*

Icon. *Dill. ethn. t. 26. f. 29.*

erecta.

- β. *G. tomentosior; caule erecto, foliis oblongo-ovalibus.*

Trifolium erectum. Walt. fl. car....

In sandy ground, among hedges: β. on road sides and old fields: Virginia to Georgia. ♀. June—Aug. v. v. Flowers yellow.

reniformis.

5. *G. erecta, humilis, pubescens; foliis omnibus simplicibus reniformi-rotundatis rugosis reticulatis, junioribus tomentosis, racemis subsessilibus paucifloris, leguminibus compresso-ovatis dispermis.*

G. tomentosa γ. monophylla. Mich. fl. amer. 2. p. 63.

Trifolium simplicifolium. Walt. fl. car. 184.

In barren pine-woods: Carolina and Georgia. ♀. July, Aug. v. s. Flowers yellow.

586. GALACTIA. Brown. *Mich. fl. amer. 2. p. 61.*

mollis.

1. *G. volubilis, molliter pubescens, subcanescens; foliis ternatis ovato-oblongis obtusis glabriusculis subtus glaucis, racemis axillaribus simplicibus elongatis paucifloris, leguminibus villosis.—Mich. fl. amer. 2. p. 61.*

Hedysa
In pine
Aug
the s

2. *G. prostrata, lenticularis, glabra, ciliata, fl. am...*

Dolichos
Ervum
In pine-
na. 7
and ex
large,
mixed

1. *C. foliis lanceolatis subens...*
Icon. *Di...*
In hedges
Flower
namen

2. *C. foliis triangulatis n...*
Icon. *Ciebus c...*
fl. amer...
In hedge...
lina.
of a pal...

1. *R. foliis pendulibus.—H...*
Icon. *Schr...*
arb. t. 7
On dry fer...
Carolina
sweet-sco...

Hedysarum volubile. Willd. sp. pl. 3. p. 1204.
In pine-barrens: New Jersey to Carolina. ♀. July,
Aug. v. v. Flowers purple, with yellow and white,
the size of *Hedysarum paniculatum*.

2. G. prostrata, subvolubilis, glabriuscula; foliis ternatis el-
liptico-oblongis obtusis utrinque emarginatis utrinque
glabris, racemis axillaribus simplicibus abbreviatis pau-
cifloris, calycibus glabris, leguminibus villosis.—Mich.
fl. amer. 2. p. 62.

Dolichos regularis, Willd. sp. pl. 3. p. 1049.
Ervum volubile. Walt. fl. car. 187.

In pine-woods and sandy fields: New Jersey to Caroli-
na. ♀. June—Sept. v. v. The roots are fusiform,
and extend to a great depth: the flowers, though not
large, are extremely pretty, purple, red and white
mixed.

587. CLITORIA. Gen. pl. 1183.

1. C. foliis ternatis ovatis calyce bracteis longiore 5-partito:
lacinias subulatis incurvis divergentibus, leguminibus
subensiformibus.—Willd. sp. pl. 3. p. 1069.

Icon. Dill. elth. t. 76. f. 87.

In hedges: Virginia and Carolina. ♀. July, Aug. v. v.
Flowers large, beautiful blue, very handsome and or-
namental.

2. C. foliis ternatis, calyce bracteis linear-lanceolatis mul-
totes majore, oblongo-tubuloso 5-fido, legumine fa-
ciebus convexis toruloso, seminibus glutinosis. Mich.
fl. amer. 2. p. 62. Willd. sp. pl. 3. p. 1070.

In hedges on the banks of rivulets: Virginia and Caro-
lina. ♀. July—Sept. v. v. Flowers smaller and
of a paler blue than the preceding.

glabella,

virginiana,

marianna.

588. ROBINIA. Gen. pl. 1195.

1. R. foliis impari-pinnatis, stipulis spinescentibus, racemis *Pseudacacia*
pendulis, calycis dentibus muticis, leguminibus lœvi-
bus.—Willd. sp. pl. 3. p. 1131.

Icon. Schmidt arb. 32. Duham. arb. ed. nov. 16. Wangh.
arb. t. 7. f. 19. Pluk. alm. t. 4.

On dry fertile ridges on the mountains, from Canada to
Carolina. ♂. May, June. v. v. Flowers white,
sweet-scented. This highly ornamental tree is at the
same time a most useful one: its wood is almost in-

DIADELPHIA DECANDRIA. Robinia.

corruptible, and particularly calculated for posts of gates and fences. It is universally known by the name of *False Acacia*, in America *Locust-tree*.

viscosa.

2. *R. foliis impari-pinnatis, racemis axillaribus erectis confertifloris, calycibus acuminatis, ramis petiolis pedunculis leguminibusque glanduloso-viscosis.*—*Willd. sp. pl. 3. p. 1131.*

R. glutinosa. Bot. mag. 560.

Icon. Vent. desc. pl. nov. t. 4. Duham. arb. ed. nov. 17.
On the banks of rivers in South Carolina, particularly on the Savannah. $\frac{1}{2}$. June, July. v. v. Flowers pale red, with white. This tree is highly ornamental; but its creeping roots are a great nuisance in small plantations.

hispida.

3. *R. foliis impari-pinnatis : foliolis rotundato-ovalibus mucronatis, racemis axillaribus suberectis, calycibus acuminatis, caule subinermi, ramis pedunculis calycibus leguminibusque hispidis.*—*Willd. sp. pl. 3. p. 1132.*

R. montana. Bartr. catal.

R. rosea. Duham. arb. ed. nov. 18.

Icon. Bot. mag. 311. Schmidt arb. 31. Catesb. car. 3. t. 20. Mill. ic. 244.

rosea.

- β . *R. foliolis plerumque alternis, ramulis glabriusculis.*
On the high mountains of Virginia and Carolina. $\frac{1}{2}$. May—Sept. v. v. Flowers large, a beautiful rose-colour, inodorous. This elegant little shrub is one of the highest ornaments of our gardens and plantations. The var. β . is less hispid, and grows to a considerable large upright shrub, whereas the original *hispida* is a low straggling plant.

589. INDIGOFERA. *Gen. pl. 1205.*

caroliniana.

1. *I. herbacea, erecta ; foliis pinnatis : foliolis obovalibus, spicis folio longioribus pedunculatis, floribus distans, leguminibus pendulis subdispermis reticulato-venosis.*—*Walt. fl. car. 187. Mich. fl. amer. 2. p. 68.*

In sandy fields: Carolina to Florida. $\frac{1}{2}$. v. s. Flowers blue.

5
 1. *T. ere*
oblc
legu
Per
Galega
In san
July
som
 2. *T. pro*
bus
bris,
tis s
Galega
In Ge
Ens
of t
a tri
The
any c
tioni
most
 3. *T. gra*
long
lis o
long
Galega
Galega
In pi
4.

1. *L. folii*
losis,
flore
mine
On the
yellow

590. TEPHROSIA. Pers. syn. 2. p. 329.

1. *T. erecta*, candicanti-villosa ; foliolis (17—21) ovali-oblongis acuminatis, racemo terminali brevi subsessili, leguminibus retrofalcatis villosis, calycibus lanatis.—
Pers. syn. 2. p. 329.
Galega virginiana. Pluk. alm. t. 23. f. 2.
In sandy and dry woods : Canada to Florida. ¼. June, July. v. v. About a foot high ; flowers very handsome, rose-coloured and yellowish white.
2. *T. prostrata*, pubescens ; foliis pinnatis quinis subsessiliibus : foliolis cuneato-obovatis obtusissimis supra glabris, subtus sericeis, pedunculis oppositifoliis elongatis sub-3-floris, leguminibus rectiusculis.
Galega villosa. Mich. fl. amer. 2. p. 67?
In Georgia. *Enslen.* ¼. July, Aug. v. s. in *Herb. Enslen.* Flowers purple. The singular circumstance of the nearly sessile leaves, gives it the appearance of a trifoliate plant, the lower pair representing stipulæ. Though Michaux's description answers very well in any other respect, I was led to doubt by his not mentioning the number of leaves, which is one of the most striking characters.
3. *T. gracilis*, humifusa, pubescens ; foliolis (13—15) oblongo-ellipticis truncato-obtusis mucronatis, pedunculis oppositifoliis elongatis 3—5-floris, leguminibus oblongis falcatis hispidis.
Galega hispidula. Mich. fl. amer. 2. p. 68.
Galega spicata. Walt. fl. car. 188.
In pine-woods and on slate-hills : Virginia to Georgia. ¼. July—Sept. v. v. Flowers pale red.

591. LOTUS. Gen. pl. 1212.

1. *L. foliis ternatis subsessilibus oblongis acutis sericeo-villosis*, pedunculis axillaribus unifloris folio longioribus, flore unibracteato, calycinis laciniis linearibus, legume glabro longissimo.
On the banks of the Missouri. ♂. v. s. Flowers yellow.

592. MEDICAGO. *Gen. pl.* 1214.*lupulina.*

1. *M. spicis ovalibus, leguminibus reniformibus monospermis stipulis integerrimis, foliolis obovatis, caulinibus procumbentibus.*—*Willd. sp. pl.* 3. p. 1406.

Icon. *Schkuhr handb.* t. 212.In gravelly soil, on road sides: Canada to Carolina. ⊖.
June—Sept. v. v. Flowers small, yellow.*intertexta.*

2. *M. pedunculis subbifloris, leguminibus cochleatis ovalibus, aculeis pubescensibus setaceis distichis adpressis, stipulis ciliato-dentatis, foliolis obovatis dentatis.*—*Willd. sp. pl.* 3. p. 1411.

M. polymorpha intertexta. *Sp. pl.* 1098.Icon. *Schkuhr handb.* 212. b.

In sandy fields: Carolina. July, Aug. v. v. Flowers yellow.

Note.—In addition to the observations under *Psoralea esculenta*, p. 475, it may be interesting to state, that A. B. Lambert, Esq. raised in the summer of 1812 a number of plants, the seeds of which were taken out of a specimen collected in the year 1805 by M. Lewis, Esq. Also, that this highly interesting plant is now growing at Messrs. Frasers, Sloane Square.

X

593. CICE
lyph594. APA
silis595. SCO
sus,
gine596. LEO
sus,597. CHO
stipi
rie.598. PREN
subs599. LACT
pitat
bran600. SONC
silis,601. HIER
plex602. KRIG
5-ph
plex.603. HYOS
terior
calyc604. TROX
pus
plex,

XVII. SYNGENESIA.

I. AEQUALIS.

* *Ligulæ.*

593. CICHORIUM. *Recept.* subpaleaceum. *Pappus* polyphyllus, paleaceus. *Cal.* calyculatus.
594. APARGIA. *Recept.* nudum. *Pappus* plomosus, sessilis. *Cal.* imbricatus.
595. SCORZONERA. *Recept.* nudum. *Pappus* plomosus, substipitatus. *Cal.* imbricatus, squamis margine scariosis.
596. LEONTODON. *Recept.* nudum. *Pappus* plomosus, stipitatus. *Cal.* imbricatus, squamis laxiusculis.
597. CHONDRILLA. *Recept.* nudum. *Pappus* pilosus, stipitatus. *Cal.* calyculatus. *Flosculi* multiplici serie. *Sem.* rugosa.
598. PRENANTHES. *Recept.* nudum. *Pappus* simplex, subsessilis. *Cal.* calyculatus. *Flosculi* simplici serie.
599. LACTUCA. *Recept.* nudum. *Pappus* simplex, stipitatus. *Cal.* imbricatus, cylindricus, margine membranaceo. *Sem.* lœvia.
600. SONCHUS. *Recept.* nudum. *Pappus* pilosus, sessilis. *Cal.* imbricatus, ventricosus.
601. HIERACIUM. *Recept.* nudiusculum. *Pappus* simplex, sessilis. *Cal.* imbricatus, ovatus.
602. KRIGIA. *Recept.* nudum. *Pappus* membranaceus, 5-phyllus, setis 5. iterstinctis. *Cal.* polyphyllus, simplex.
603. HYOSERIS. *Recept.* nudum. *Pappus* duplex: exterior capillaceus; interior paleaceo-aristatus. *Cal.* calyculatus.
604. TRÖXIMON. *Recept.* nudum, puncticulatum. *Pappus* pilosus, sessilis. *Cal.* oblongus, conicus, simplex, aut squamis inæqualibus imbricatus.

** *Flosculosæ.*

605. STOKESIA. *Recept.* nudum. *Pappus 4-setosus.* *Cal.* foliaceus, subimbricatus. *Cor.* radiata : corollulis radii infundibuliformibus, irregularibus.
606. ARCTIUM. *Recept.* paleaceum. *Pappus setoso-paleaceus.* *Cal.* globosus : squamis apice hamis inflexis.
607. CNICUS. *Recept.* villosum. *Pappus plumosus.* *Cal.* imbricatus, ventricosus.
608. CARDUUS. *Recept.* villosum. *Pappus capillaris.* *Cal.* imbricatus, ventricosus.
609. LIAIRIS. *Recept.* nudum. *Pappus plumosus,* colossum. *Cal.* oblongus, imbricatus.
610. VERNONIA. *Recept.* nudum. *Pappus duplex :* exterior paleaceus, brevis; interior capillaris. *Cal.* ovatus, imbricatus.
611. KUHNIA. *Recept.* nudum. *Pappus plumosus.* *Cal.* imbricatus, cylindraceus.
612. EUPATORIUM. *Recept.* nudum. *Pappus pilosus.* *Cal.* imbricatus, cylindricus. *Stylus semibifidus,* longus.
613. MIKANIA. *Recept.* nudum. *Pappus pilosus s. plumosus.* *Cal.* 4—6-phylloides, 4—6-florus. *Stylus semibifidus,* longus.
614. CHRYSOCOMA. *Recept.* nudum. *Pappus pilosus.* *Cal.* imbricatus. *Stylus* vix flosculis longior.
615. CACALIA. *Recept.* nudum. *Pappus pilosus.* *Cal.* cylindricus, basi calyculatus.
616. SPARGANOPHORUS. *Recept.* nudum. *Sem.* coronatum cupula subcartilaginea, integerrima, nitida. *Cal.* subglobosus, imbricatus : squamis apice recurvatis.
617. HYMENOPAPPUS. *Recept.* nudum. *Pappus polypillus,* paleaceus. *Cal.* polypillus, patens.
618. MELANANTHERA. *Recept.* paleaceum : paleis flosculos inferne amplexantibus. *Pappus aristis inaequalibus,* 4—5-inermibus. *Cal.* imbricatus : squamis ovatis, appressis, subæqualibus.
619. MARSHALLIA. *Recept.* paleaceum. *Pappus paleis 5.* membranaceis, acuminatis. *Cal.* imbricatus : squamis sublanceolatis incumbentibus.
620. SANTOLINA. *Recept.* paleaceum. *Pappus nullus.* *Cal.* imbricatus, hemisphæricus : squamis carinatis apice scariosis.

621. ARTI. *Cor.*
datiæ
622. TAN. *gina-*
tus,
623. BACCI. *Flas-*
bris
624. CON. *pilla-*
mis
625. GNA. *sus.*
datiæ
626. BELL. *mis-*
627. CHR. *null-*
inti-
628. PYRE. *tus.*
629. ARN. *sus.*
longi
absorbi-
630. CINN. *Cal-*
631. SEN. *cylind-*
632. TUSS. *Cor-*
squæ-
nac-
633. INU. *imb-*
634. ERIC. *radi-*

II. SUPERFLUA.

* *Discoidei.*

621. ARTEMISIA. *Recept.* subnudum. *Pappus nullus.*
Cor. radii nulla. *Cal.* imbricatus : squamis rotundatis, conniventibus.
622. TANACETUM. *Recept.* nudum. *Pappus submarginalis.* *Cor.* radii obsoletæ, 3-fidæ. *Cal.* imbricatus, hemisphæricus : squamis acuminatis.
623. BACCHARIS. *Recept.* nudum. *Pappus pilosus.*
Fls. feminei hermaphroditi immixti. *Cal.* imbricatus, cylindricus : squamis ovatis subcoriaceis.
624. CONYZA. *Recept.* nudum. *Pappus simplex,* capillaris. *Cor.* radii 3-fidæ. *Cal.* imbricatus : squamis linearibus aut ovatis.
625. GNAPHALIUM. *Recept.* nudum. *Pappus plumosus.* *Cal.* imbricatus : squamis marginalibus rotundatis, scariosis, coloratis,

** *Radiati.*

626. BELLIS. *Recept.* nudum. *Pappus nullus.* *Cal.* hemisphæricus : squamis æqualibus. *Sem.* obovata.
627. CHRYSANTHEMUM. *Recept.* nudum. *Pappus nullus.* *Cal.* hemisphæricus, imbricatus ; squamis intimis scariosis.
628. PYRETHRUM. *Recept.* nudum. *Pappus marginatus.* *Cal.* squamis acutiusculis margine scariosi.
629. ARNICA. *Recept.* nudum. *Pappus simplex,* pilosus. *Cal.* hemisphæricus ; foliolis subæqualibus disco longioribus. *Cor.* radii (luteæ) saepius filamentis 5. absque antheris.
630. CINERARIA. *Recept.* nudum. *Pappus simplex.* *Cal.* simplex, polyphyllus, æqualis.
631. SENEPIO. *Recept.* nudum. *Pappus simplex.* *Cal.* cylindricus, calyculatus : squamis apice sphacelatis.
632. TUSSILAGO. *Recept.* nudum. *Pappus simplex.* *Cor.* femineæ ligulatæ s. edentulæ. *Cal.* simplex : squamæ æquales, discum æquantes, submembranaceæ.
633. INULA. *Recept.* nudum. *Pappus simplex.* *Cal.* imbricatus. *Cor.* radii numerosæ (luteæ).
634. ERIGERON. *Recept.* nudum. *Pappus pilosus.* *Cor.* radii capillares (coloratae).

635. SOLIDAGO. *Recept.* nudum, punctis exaratum. *Pappus* simplex. *Cor.* radii circiter 5. (flavæ). *Cal.* squamæ imbricatae, clausæ.
636. ASTER. *Recept.* nudum, scrobiculatum. *Pappus* pilosus. *Cor.* radii plures, saepius 10. (aut pauciores, colorati). *Cal.* imbricati squamæ inferiores patulæ.
637. DONIA. *Recept.* nudum, favosum. *Pappus* 3-4-aristatus, deciduus. *Cor.* radii plures (30-35 flavi). *Cal.* hemisphaericus, imbricatus : squamis interioribus cartilagineis, coloratis.
638. BŒBERA. *Recept.* nudum. *Pappus* pilosus. *Cal.* duplex : exterior polyphyllus, interior octophyllus.
639. ACTINELLA. *Recept.* nudum. *Pappus* polyphyllus (4-6), paleaceo-aristatus. *Cal.* polyphyllus, æqualis.
640. HÉLENIUM. *Recept.* nudum : radii paleaceum. *Pappus* 5-aristatus. *Cor.* radiis 3-fidis. *Cal.* simplex, multipartitus.
641. BOLTONIA. *Recept.* favosum. *Pappus* dentato-aristatus : aristis duabus oppositis interdum elongatis. *Cor.* radii plurimæ. *Cal.* imbricatus.
642. SIEGE-BECKIA. *Recept.* paleaceum. *Pappus* nullus. *Radius* dimidiatus. *Cal.* exterior 5-phyllus, proprius patens. *Sem.* sub-4-gona.
643. PHÆTHUSA. *Recept.* paleaceum. *Pappus* nullus. *Flosculi* radii 1-3. *Cal.* imbricatus. *Sem.* hispida.
644. ECLIPTA. *Recept.* paleaceum. *Pappus* nullus. *Cor.* disci 4-fidæ. *Sem.* subcompresso-marginata, obsolete 2-3-dentata.
645. ANTHEMIS. *Recept.* paleaceum : paleis planis, apice acuminatis, rigidis. *Pappus* nullus s. margo membranaceus. *Flores* radii plures quam 5. *Cal.* hemisphaericus, subæqualis.
646. ACHILLEA. *Recept.* paleaceum. *Pappus* nullus. *Flosc.* radii circiter 5., dilatati. *Cal.* ovatus, imbricatus. (*Flores* corymbosi.)
647. TETRAGONOTHECA. *Recept.* paleaceum. *Pappus* nullus. *Cal.* monophyllus, 4-gonus, 4-partitus, latissimus.
648. HELIOPSIS. *Recept.* paleaceum, conicum. *Pappus* nullus. *Cor.* radii lineares, magnæ. *Cal.* imbricatus : squamis ovatis, lineatis. *Sem.* 4-gona.
649. BUPHANTHALMUM. *Recept.* paleaceum. *Pappus* mar-
go obsoletus s. 4-dentatus. *Cal.* foliaceus. *Sem.* la-
tera, præstimum radii, marginata.
650. AME Cor
651. VER tus.
652. ZINI erec
653. BIDI caly
2. s
diat
sub.
654. COR radi
pres
tosa
655. HEL pus
rosu
656. GAL Pat
Cal
657. RUD pus
ma
658. CEN Cor
659. CHA Flo
Flo
660. SILP bico
pres
661. IPOL Ca
lus
662. CHR

- im.
Cal.
pus
res,
læ.
-4-
vi).
bus

al.
.lus
æ-
ip-
ex,
to-
nis.
us
is,
is.
la.
or.
te
ce
n-
e-

is.
i-
b-
s,
ts
1-
-
-
-
-
650. AMELLUS. *Recept.* paleaceum. *Pappus* pilosus.
Cor. radii indivisæ. *Cal.* imbricatus.
651. VERBESINA. *Recept.* paleaceum. *Pappus* aristatus.
Flosc. radii circiter 5. *Cal.* dupli ordine.
652. ZINNIA. *Recept.* paleaceum. *Pappus* aristis 2.
erectis. *Radius* 5-florus, persistens. *Cal.* imbricatus, ovato-cylindricus.

III. FRUSTRANEA.

653. BIDENS. *Recept* paleaceum: paleis extimis quasi calycom communem constituentibus. *Pappus* aristis 2. s. 4. retrorsum. *Flores* flosculosi aut etiam radiati: radiis neutris aut imperfecte masculis. *Cal.* subæqualis, calyculatus.
654. COREOPSIS. *Recept.* et *Cal. comm.* Bidentis. *Flores* radiati: radiis neutris, prolixis (luteis). *Sem.* compressa aut emarginata, aut bidentata, aut etiam biseptosa: setis inermibus nec retrorsum aculeatis.
655. HELIANTHUS. *Recept.* paleaceum, planum. *Pappus* diphyllos, caducus. *Cal.* imbricatus, subsquarrosus, foliaceus.
656. GALARDIA. *Recept.* paleaceum, hemisphæricum. *Pappus* paleacens, polyphyllus. *Cor.* radii 3-partitæ. *Cal.* polyphyllus, planus.
657. RUDBECKIA. *Recept.* paleaceum, conicum. *Pappus* margine 4-dentato. *Cal.* dupli ordine squararum.
658. CENTAUREA. *Recept.* setosum. *Pappus* pilosus. *Cor.* radii infundibuliformes, irregulares. *Cal.* varius.

IV. NECESSARIA.

659. CHAPTALIA. *Recept.* nudum. *Pappus* capillaris. *Flosculi* radii in dupli serie difformes, femininei. *Flosculi disci* masculi bilabiati.
660. SILPHIUM. *Recept.* paleaceum. *Pappus* marginatobicornis. *Cal.* squarresus, foliaceus. *Sem.* compressa, obcordata, marginata.
- 661 POLYMNIA. *Recept.* paleaceum. *Pappus* nullus. *Cal.* duplex: exterior 4-5-phyllos; interior 10-phyllos. foliolis concavis.
662. CHRYSOGONUM. *Recept.* paleaceum. *Pappus* 1-

phyllus, 3-dentatus. *Cal.* 5-phyllus. *Sem.* calyculo 4-phylo involuta.

663. **BALTIMORA.** *Recept.* paleaceum. *Pappus* nullus. *Flosc.* radii 5. *Cal.* cylindricus, polyphyllus. *Sem.* 3-quatra.

664. **PARTHENIUM.** *Recept.* paleaceum, planum. *Pappus* nullus. *Cal.* 5-phyllus. *Sem.* obovata.

665. **IVA.** *Recept.* pilosum. *Pappus* nullus. *Cor.* radii 5. *Antheræ* approximatæ, non coalitæ. *Styli* 2. longi. *Cor.* radii 5. *femineis* minimis. *Cal.* 3-phyllus.

666. **AMBROSIA.** *Monoica.* *Masc.* *Recept.* nudum. *Antheræ* approximatæ, non coalitæ. *Cal.* 1-phyllus, hemisphæricus, multiflorus. *Fem.* *Cal.* 1-phyllus, subinteger aut 5-dentatus, 1-florus. *Cor.* nulla. *Styli* 2. *Nux e calyce indurato*, 1-sperma.

667. **XANTHIUM.** *Monoicum.* *Masc.* *Recept.* paleaceum. *Antheræ* approximatæ, non coalitæ. *Cal.* polyphyllus, imbricatus, hemisphæricus, multiflorus. *Fem.* *Cal.* involucrum 2-phyllum, 2-florum. *Cor.* O. *Stigm.* 2. *Drupa* sicca, muricata, 2-fida. *Nucleus* 2-locularis.

V. SEGREGATA.

668. **ELEPHANTEPUS.** *Recept.* nudum. *Pappus* setaceous. *Calyculus* 4-florus. *Corollulæ* ligulatæ, hermaphroditæ.

I. $\mathcal{A}E$ Q U A L I S.

593. CICHORIUM. *Gen. pl. 1251.*

Intybus.

1. *C.* floribus axillaribus geminis subsessilibus, foliis runcinatis. *Willd.* sp. pl. 3. p. 1624.

Icon. *Fl. dan.* 907. *Engl. bot.* 539.

In fields and on road sides: Pennsylvania, near Philadelphia. Most certainly introduced from Europe. \mathcal{U} . July, Aug. v. v. Flowers beautiful blue.

1. *A.* scap
runc
p. 1st
Leonto
Hedyp
Icon.
In past
Sept
is pr

1. *S.* caul
ceola
dent
Leonto
In was

1. *L.* caly
natis
pl. 3
Icon.
Comm
24.
rally
anot
con
to g

1. *C.* gla
tege
flori
In Le
brig
are
leav

594. APARGIA. *Gen. pl. 1232.*

1. *A. scapis ramosis declinatis squamosis, foliis lanceolatis runcinato-pinnatifidis glabriusculis. Willd. sp. pl. 3. p. 1550.*
Leontodon autumnale Sp. pl. 1123.
Hedypnois autumnalis. Smith fl. brit. 2. p. 826.
Icon. Fl. dan. 501. Engl. bot. 830.
 In pastures and on road sides : New England. 4. Aug.
 Sept. v. v. Flowers golden-yellow, outside red. It
 is probably an introduced plant.

595. SCORZONERA. *Gen. pl. 1230.*

1. *S. caule erecto 3-floro, pedunculis elongatis, foliis lan. ceolatis acutissimis acute-pinnatifidis interdum rariter dentatis. Mich. fl. amer. 2. p. 89.*
Leontodon carolinianum. Walt. fl. car. 192.
 In waste places : Carolina. *Michaux.* +.

596. LEONTODON. *Gen. pl. 1237.*

1. *L. calyce exteriore reflexo, scapo unifloro, foliis runcinatis glabris, lacinii lanceolatis dentatis. Willd. sp. pl. 3. p. 1544.*
Icon. Fl. dan. 574. Engl. bot. 510.
 Common in fields and grass plots ; probably introduced.
 4. April—July. v. v. Flowers yellow ; it is generally known by the name of *Dandelion*. There is another species, found in shady wet woods, which I consider indigenous, but I have no materials in hand to give its description.

*Taraxacum.*597. CHONDRILLA. *Gen. pl. 1235.*

1. *C. glaberimum, multicaule ; foliis linear-lanceolatis integrinis, pedunculis 2-3. longissimis subaphyllis unifloris.*
 In Lower Virginia. 3. Aug. Sept. v. v. Flowers bright yellow ; pappus turns a pale red ; the peduncles are from six to ten inches long, with a few linear, small leaves.

lævigata.

598. PRENANTHES. *Gen. pl.* 1236.

- junccea.* 1. P. caule virgatim-ramosissimo sulcato glabro, foliis caulinis remotis subulatis brevissimis, ramulis unifloris, calycibus 5-fidis 5-floris: foliolis margine membranaceis.
On the banks of the Missouri. 2f. v. s. Flowers purple; it has a striking resemblance to *Chondrilla junccea*; but the leaves are shorter, and the pappus entirely sessile; the radical leaves I have not seen.
- altissima.* 2. P. caule ramoso, foliis trilobis petiolatis angulatis denticulatis **margine** scabris, racemis axillaribus, floribus nutantibus, calycibus sub-5-floris.—*Willd. sp. pl.* 3. p. 1537.
Icon. *Pluk. alm. t.* 317. f. 2.
In shady woods: Canada to Virginia. 2f. July—Sept. v. v. Flowers yellow. Generally a very tall robust plant.
- cordata.* 3. P. caule superne paniculato, foliis petiolatis cordatis dentatis ciliatis, floralibus sessilibus oblongis integerrimis, panicula laxa racemiflora, floribus nutantibus, calycibus 6-fidis 6-floris.—*Willd. hort. berol.* 25.
In shady woods, about rocks; New York to Virginia. 2f. Aug.—Oct. v. v. Flowers pale yellow. It is an intermediate species between the preceding and *P. alba*.
- virgata.* 4. P. glabra; caule simplicissimo, foliis omnibus lyrato-sinuatis, racemulis subsecundis, floribus pendulis, calycibus glabris 8-fidis 10-floris.—*Mich. fl. amer.* 2. p. 83.
P. autumnalis. *Walt. fl. car.* 193.
In sandy fields, near ditches: New Jersey to Carolina. 2f. Aug. Sept. v. v. From three to six feet; flowers pale purple. This plant has a very fine effect in large plantations. Michaux quotes Gronovius; but his *P. autumnalis* belongs to *P. rubicunda*.
- simplex.* 5. P. caule simplicissimo, foliis superioribus linear-lanceolatis integerrimis, radicalibus lanceolatis sinuatis, racemo terminali simplicissimo, floribus nutantibus, calycibus sub-5-floris.
P. virgata. *Willd. sp. pl.* 3. p. 1533.
In Carolina. *Enslen.* 2f. July, Aug. v. s. in *Herb. Enslen.* About two feet high; flowers purple.

foliis cau-
; unifloris,
membra-

wers pur-
la juncea;
tirely ses-

atis denti-
, floribus
sp. pl. 3.

ly—Sept.
all robust

datis den-
gerrimis,
, calyci-

Virginia.
v. It is
ding and

is lyrato-
dulis, ca-
amer. 2.

Carolina.
et ; flow-
effect in
ius ; but

i-lanceo-
atis, ra-
ibus, ca-

in *Herb.*
le.



Scironanthus serpentarius.

- 5
6. P. foliis l.
angulat.
cifloris
sub-20
In Illinois
chauz.
7. P. foliis r.
caulinis
lanceol.
tantibu
pl. 3. p
P. suavis.
Icon. Bo
In shady
Sept.
flowers
8. P. foliis c.
rimis, i
summis
ribus n
P. alba β.
P. autumn
In shady
Oct. 1
eighteen
9. P. foliis
caulinis
lacinia
cemas t
calycib
On the m
—Oct.
purple.
the na
specific
my trav
an oppo
this rem
the Swe
snake, a
inflamm
immedia
boiled in
steeped in
he was e

SYNGENESIA EQUALIS. Prenanthes.

400

6. *P. foliis lato-lanceolatis in petiolum attenuatis inæqualiter angulato-dentatis, panicula fasciculis terminalibus paucifloris nutantibus, calycibus hirsutis 10—12-fidis sub-20-floris.* — *Mich. fl. amer. 2. p. 84.*

In Illinois and on the high mountains of Carolina. *Mitchaux. 4. +.*

7. *P. foliis radicalibus angulato-hastatis dentatis sublobatis, caulinis subrotundo-ovatis dentatis petiolatis, summis lanceolatis, panicula laxa, fasciculis terminalibus nutantibus, calycibus 8-fidis 9—10-floris.* — *Willd. sp. pl. 3. p. 1536.*

P. suavis. Parad. lond. 85.

Icon. Bot. mag. 1079.

In shady woods: New England to Carolina. *4. Aug.*
Sept. v. v. About two feet high; calyces purple,
flowers white.

8. *P. foliis ciliatis, radicalibus hastato-angulatis subintegerimis, inferioribus obovatis basi attenuatis subangulatis, summis lanceolatis integerimis, racemo simplici, floribus nutantibus.* — *Willd. sp. pl. 3. p. 1537.*

P. alba β. Sp. pl. 1122.

P. autumnalis. Gron. fl. virg. 1. p. 89.

In shady woods: Pennsylvania to Carolina. *4. Aug.—Oct. v. v.* Resembles the preceding; not above eighteen inches high.

9. *P. foliis dentatis asperis, radicalibus palmato-sinuatis: caulinis longe petiolatis sinuato-pinnatifidis subtrilobis: lacinia intermedia 3-partita, summis lanceolatis, racemis terminalibus subpaniculatis brevibus nutantibus, calycibus 8-fidis 12-floris.*

On the mountains of Virginia and Carolina. *4. Aug.—Oct. v. v.* About two feet high; flowers pale purple. This plant is known by the inhabitants under the name of *Lion's-foot*, and is in high esteem as a specific in curing the bite of the rattlesnake. During my travels through the mountains of Virginia, I had an opportunity of being a witness of the efficacy of this remedy. A man living in Cove-mountains, near the Sweet-springs, was bit in the foot by a Moccasin snake, a species considered the most dangerous. An inflammation and swelling of his whole leg took place immediately; but by taking the milky juice of this plant boiled in milk, inwardly, and applying to the wound the steeped leaves, which were very frequently changed, he was cured in a few days. As this plant deserves

crepidina.

alba.

rubicunda.

Serpentaria.

the attention of the physician, I have given a figure of it, it being frequently confounded with another species of this genus, which probably may not have quite so strong an effect, as the inhabitants are very careful to have the true *Lion's-foot*, in case of accidents happening, and usually call the other species of *Prenanthes* *False Lion's-foot*. Gronovius, in his Flora, pag. 113, mentions Dr. Witt's *Snake-root* under *P. autumnalis* or Willdenow's *rubicunda*, as a remedy for the bite of the rattlesnake; which shows that he had information of the use made of this plant, though he did not know the genuine species. In the Banksian Herbarium is a specimen of *P. rubicunda*, with the following note in the hand-writing of Clayton: "This is the rattlesnake-root that Dr. Witt supposes to be the best cure for the bite of the snake, a very odd plant, hardly two leaves alike upon a plant, as to shape or the indentings of the leaves."

racemosa.

10. *P. caule simplici, foliis omnibus indivisis laevibus, radicibus ovali-lanceolatis, caulinis semiamplexicaulibus, racemo oblongo hirsutissimo, fasciculis cernuis, calycibus 8-9-partitis 9-12-floris.* *Mich. fl. amer. 2. p. 84.*

In the northern parts of Canada. *Michaux.* +.

illinoensis.

11. *P. caule simplici, foliisque aspermis, foliis omnibus indivisis ovali-lanceolatis, racemo longo, fasciculis subsessilibus erectis hirsutis.* *Pers. syn. 2. p. 366.*

P. aspera. *Mich. fl. amer. 2. p. 83.*

In natural meadows in the Illinois country. *Michaux.* +.

599. LACTUCA. *Gen. pl. 1234.*

elongata.

1. *L. foliis subtus laevibus, inferioribus runcinatis integerimis amplexi-caulibus, infimis dentatis, summis lanceolatis, floribus corymboso-paniculatis.* *Willd. enum. 817.* *Willd. sp. pl. 3. p. 1525.*

L. longifolia. *Mich. fl. amer. 2. p. 85?*

In woods, on road sides, in fertile soil: Canada to Carolina. ♂. Aug. Sept. v. v. From three to six feet high. Flowers small, pale yellow.

graminifolia.

2. *L. caule erecto simplici, foliis inernibus plerisque indivisis basi simplici longissime linearibus, panicula aplylla laxa, ramis rarifloris, floribus omnibus pedunculatis.* *Mich. fl. amer. 2. p. 85.*

In Lower Carolina. *Michaux.* +.

1. *S. pedunculata*
Smith
Icon. E
Among
4. A

2. *S. pedunculata*
Icon. F
A comm.—Sept

3. *S. pedunculata*
Icon. G
S. canadensis
Chondrilla
In shady
rolina.
feet h
bus;

4. *S. pedunculata*
Icon. H
Willd.
S. montana
S. cærulea
S. canadensis
Hieracium
Icon. Fl.
In Canad.
someti

5. *S. pedunculata*
Icon. I
S. cinerea
sp. pl.
S. spicata
On the b.
to Vir.
white,

6. *S. pedunculata*
Icon. J
S. rato-rum
p. 152

600. SONCHUS. *Gen. pl.* 1233.

1. *S. pedunculis calycibusque hispidis subumbellatis, foliis runcinatis denticulatis basi cordatis, radice repente.*
Smith fl. brit. 2. p. 817. *Willd. sp. pl.* 3. p. 1512.
Icon. *Engl. bot.* 674. *Curt. fl. lond.* 53.
Among rubbish and in cultivated grounds: Pennsylvania.
♀. Aug. Sept. v. v. Flowers large, deep yellow.
arvensis.
2. *S. pedunculis subtomentosis umbellatis, calycibus glabris, foliis oblongo-lanceolatis amplexicaulibus denticulatis subsinuatis.* *Willd. sp. pl.* 3. p. 1514.
Icon. *Fl. dan.* 682. *Curt. fl. lond.* 58.
A common weed in most gardens and fields. ♂. July—Sept. v. v. Flowers yellow.
oleraceus.
3. *S. pedunculis hirsutis nudis, floribus paniculatis, foliis macrophyllus, lyratis basi cordatis subtus hirtis.* *Willd. sp. pl.* 3. p. 1519.
S. canadensis. *Fraelich. in usteri annal.* 1. p. 29.
Chondrilla sylvestris alta. *Gronov. virg.* 115.
In shady low grounds, near springs: Pennsylvania to Carolina. ♀. Aug. Sept. v. v. From four to seven feet high; flowers blue, the size of *Cichorium Intybus*; root tuberous.
alpinus.
4. *S. pedunculis hirsutis nudis, floribus racemosis bracteatis, foliis runcinatis basi sagittatis glabris subtus glaucis.* *Willd. sp. pl.* 3. p. 1519.
S. montanus. *Lam. encycl.* 3. p. 401.
S. cæruleus. *Smith fl. brit.* 2. p. 815.
S. canadensis. *Sp. pl.* 1115.
Hieracium cæruleum. *Scop. fl. carn.* n. 976.
Icon. *Fl. dan.* 182.
In Canada. ♀. Aug. Sept. v. s. Flowers blue, and sometimes white.
5. *S. pedunculis squamosis, floribus racemosis, foliis runcinatis acuminatis, caule paniculato-virgato.* *Willd. sp. pl.* 3. p. 1520.
S. spicatus. *Lam. encycl.* 3. p. 401.
On the borders of woods, on road sides: New England to Virginia. ♂. July—Sept. v. v. Flowers small, white, tinged with blue.
leucophæus.
6. *S. pedunculis subsquamosis, floribus paniculatis, foliis lyrato-runcinatis denticulatis petiolatis.* — *Willd. sp. pl.* 3. p. 1520.
floridanus.

On road sides, in shady woods: Virginia and Carolina.
 ♂. July—Sept. v. v. Flowers very small, blue.
 This plant has been used for curing the bite of the
 rattlesnake, in the same manner as *Prenanthes Ser-*
pentaria, and is known by the name of *Gall of the*
Earth.

- acuminatus.* 7. S. pedunculis subsquamosis, floribus paniculatis, foliis
 radicalibus subruncinatis, caulinis ovatis acuminatis
 petiolatis medio denticulatis.—*Willd. sp. pl. 3.*
p. 1521.

Lactuca villosa. Jacq. hort. schœnb. 3. t. 367?
 In low shady places: Pennsylvania to Virginia. ♂. Aug.
 Sept. v. v. Flowers small, blue.

- pallidus.* 8. S. racemo composito terminali, foliis lanceolato-ensi-
 formibus amplexicaulibus dentatis. *Willd. sp. pl. 3.*
p. 1521.

Lactuca canadensis. Sp. pl. 1119.
Icon. Rob. ic. 148 ♂ 151.

On road sides and in woods: Canada to New England.
 ♀. July—Sept. v. v. Flowers small, yellow.

- pulchellus.* 9. S. pedunculis squamosis, floribus corymboso-racemosis,
 foliis caulinis cordato-amplexicaulibus ovato-oblongis
 acutis integerrimis glabris.
 On the banks of the Missouri. ♀. Sept. v. s. Flowers
 large, beautiful-blue.

601. HIERACIUM. *Gen. pl. 1238.*

- pusillum.* 1. H. pusillum, villosissimum; caule simplici erecto uni-
 floro remote subbifoliato, foliis lanceolatis attenuatis
 acutis integerrimis, caulinis linearibus, calyce villosis-
 simo.—*Willd. sp. pl. 3. 1561.*

In Labrador. *Colmaster.* ♀. July. v. s. in *Herb.*
Dickson. A very small plant; flowers, as in all the
 following species, yellow, the size of a common daisy.
 It strongly resembles *H. alpinum*, and is probably only
 a diminutive variety thereof.

- venosum.* 2. H. scapo nudo corymboso-paniculato glabro, pedicellis
 filiformibus, foliis obovato-lanceolatis supra rariter pi-
 losis, subtus nudis margine ciliatis glanduloso-denticu-
 latis, venis coloratis, calycibus glabris.—*Willd. sp.*
pl. 3. p. 1570.

In shady fertile woods: Canada to Carolina. ♀. June
 —Aug. v. v. Flowers middle size, of a beautiful

yel
 wit
 Po
 der
 3. H. sc
 culi
 geri
 sub
 a. H. cat
 ame
 β. H. cat
 H. ma
 Icon.
 In woc
 July
 4. H. pilo
 ciflor
 pauc
 p. 15
 Icon. J
 In Lat
 Herb
 5. H. glab
 albo-
 nudis
 p. 15
 In shad
 the r
 very s
 6. H. caule
 ceolat
 culis i
Willd
 In Pensy
 leaves
 Ameri
 scriptio
 7. H. caule
 latis ac
 argute-
 divaric
 culisqu
 H. canad
 In the we

yellow; the light green leaves are elegantly marked with blood-red veins. It is known by the name of *Poor Robin's Plantain*, and thought to possess considerable medicinal powers.

3. *H. scapo folioso corymboso-paniculato, calycibus pedunculisque glanduloso-pilosis, foliis obovatis obtusis integrerrimis, supra rarer strigoso-villosis, nervo medio subtus villosissimo.*—*Willd. sp. pl. 3. p. 1570.*

4. *H. caule subunifoliato, panicula subfastigiata.* *Mich. fl. amer. 2. p. 87.*

5. *H. caule parce folioso, panicula oblonga.* *Mich. l. c.*
H. Marianum. *Willd. sp. pl. 3. p. 1572.*

Icon. Pluk. mant. t. 320. f. 2.

In woods and on dry hills: Canada to Carolina. ♀.
July, Aug. v. v. Flowers small.

6. *H. pilosum; caule erecto simplici folioso, corymbo paucifloro, foliis oblongo-lanceolatis attenuatis inferne paucidentatis, radicalibus petiolatis.*—*Willd. sp. pl. 3. p. 1577?*

Icon. Jacq. austr. t. 119.

In Labrador. *Colmaster.* ♀. July—Sept. v. s. in *Herb. Dickson.* Flowers middle size.

7. *H. glabriusculum; caule erecto folioso paniculato inferne albo-lanato, pedicellis capillaribus, foliis lanceolatis nudis dentatis membranaceis.*—*Willd. sp. pl. 3. p. 1572.*

In shady woods: Canada to Carolina; principally on the mountains. ♀. July—Sept. v. v. Flowers very small.

8. *H. caule erecto multifloro glabro, foliis subsessilibus lanceolatis acuminatis extrorsum argute dentatis, pedunculis in apice caulis alternis subunifloris tomentosis.*—*Willd. sp. pl. 3. p. 1588.*

In Pennsylvania. *Kalm.* +. Flowers small, upright, leaves small. It is singular, that in no part of North America could I find a plant answering to this description given by Linnæus.

9. *H. caule erecto simplici viloso, foliis sessilibus lanceolatis acutis glabris subtus pilosis margine divaricato-argute-dentatis versus apicem integerrimis: dentibus divaricatis, panicula subcorymbosa, calycibus pedunculisque tomentosis.*—*Lambert. herb. ms.*

H. canadense. *Mich. fl. amer. 2. p. 86?*

In the western part of New York and Canada. ♀.

Gronovii.

nudicaule.

foliosum.

molle.

Kalmii.

virgatum.

July—Sept. v. v.; v. s. in *Herb. Lambert.* Flowers middle size.

- fasciculatum.* 8. H. glabriuscum; caule erecto folioso simplici glabro; foliis sessilibus oblongis acutis argute-dentatis: dentibus elongatis, ramis paniculae divaricatis brevibus, pedicellis subfasciculatis pubescentibus.

In Canada. *Masson.* ♀. v. s. in *Herb. Lambert.* A tall robust plant; flowers middle size.

- scrubrum.* 9. H. hirsutissimum; caule erecto folioso hispido punctis fuscis scaberrimo, foliis oblongo-ovatis integris utrinque hirsutis, panicula simpliciuscula multiflora calycibusque glanduloso-hispidis, pedicellis brevibus divaricatis.—*Mich. fl. amer. 2. p. 86.*

In woods, in shady, rocky situations: Pennsylvania. ♀. Aug. Sept. v. v. From three to five feet high; flowers small.

- macrophyllum.* 10. H. altissimum; caule erecto folioso hispido sulcato, foliis cordato-semiamplexicaulis ovato-oblongis remote grosse-dentatis nudiusculis, nervis venisque subtus pubescentibus, panicula divaricato-corymbosa, pedunculus elongatis nudis calycibusque glabris.

In Canada. ♀. v. s. in *Herb. Lambert.* A very robust plant; leaves the largest of the genus; flowers large.

602. KRIGIA. Gen. pl. 1244.

- virginica.* 1. K. pusilla; foliis lyratis glaucis glabriusculis margine ciliatis, scapo 1-floro foliis duplo altiore, calyce sub-8-phyllo.—*Willd. sp. pl. 3. p. 1618.*

Hyoseris virginica. Sp. pl. 1138.

Icon. Lam. journ. hist. nat. 1. t. 12.

In pastures, grassy hills, fields, and road sides: Canada to Florida. ♂. May—July. v. v. Flowers deep yellow, small.

603. HYOSERIS. Gen. pl. 1242.

- montana.* 1. H. glaberrima, procumbens; foliis lanceolatis integerris mis, scapo 1-floro.—*Mich. fl. Amer. 2. p. 87.*

In the mountains of Carolina. *Michaux.* +.

- angustifolia.* 2. H. foliis linear-lanceolatis sensim acutissimis glabellis, dentibus rarioribus exertis, scapo 1-floro. *Mich. fl. amer. 3. p. 87.*

Hyoseris major. Walt. fl. car. 194.

In fi
Jt
3. H. fi
tis
In C

604. TR

1. T. sc
fol
On t
Ho

2. T. gl
foli
cau
p. 1

Hyose
Hyose
Hyose

Trago
In me
rolit
flow

60

1. S. foli
1-fl

Cartha
Icon.
In Sou

large

1. A. folii
pani
pl. 3

Icon. 1
On road
frequ

Com

In fields and pastures: Virginia and Carolina. ♂. July, Aug. v. v. Flowers large, bright yellow.

3. *H. foliis lyratis basi acutis superne dilatatis apice truncatis hirsutis, scapis 1-floris.* — *Walt. fl. car.* 194. *caroliniana.*

In Carolina. *Walter.* +.

604. TROXIMON. *Gært. carp.* 2. p. 360. *Pers. syn.* 2. p. 360.

1. *T. scapo unifloro, calycinis foliolis imbricatis cuspidatis, foliis linearibus integerrimis utrinque glaucis.* *glaucum.*

On the banks of the Missouri. ♂. v. s.; v. v. in *Hortis.* Flowers bright yellow.

2. *T. glabrum, glaucum; caule erecto 2-3-fido subnudo, foliis glabris, radicalibus sublyratis, caulinis amplexis caulinibus lanceolatis integerrimis glabris.* — *Pers. syn.* 2. p. 260. *virginicum.*

Hyoseris amplexicaulis. *Mich. fl. amer.* 2. p. 87.

Hyoseris prenanthoides. *Willd. sp. pl.* 3. p. 1615.

Hyoseris biflora. *Walt. fl. car.* 194.

Tragopogon virginicum. *Sp. pl.* p. 1111.

In meadows and moist shady woods: Pennsylvania to Carolina. ♀. July, Aug. v. v. About a foot high; flowers handsome, yellow.

605. STOKESIA. *L'Herit. sert. angl.* 27.

1. *S. foliis lanceolatis, caule folioso, pedunculis axillaribus 1-floris.* — *Willd. sp. pl.* 3. p. 1703. *cyanea.*

Carthamus laevis. *Hill. kew.* p. 57. t. 5.

Icon. *L'Herit. l. c.* 38.

In South Carolina. ♀. v. s. in *Herb. Banks.* Flowers large, purple, very handsome.

606. ARCTIUM. *Gen. pl.* 1253.

1. *A. foliis caulinis cordatis petiolatis denticulatis, floribus paniculatis globosis, calycibus laevibus.* — *Willd. sp. pl.* 3. p. 1631.

Icon. *Fl. dan.* 642. *Engl. bot.* 1228.

On road sides, among rubbish, and in cultivated grounds; frequent. ♂. July, Aug. v. v. Flowers purple. Common *Burdock.* Has probably been introduced.

Lappa.

607. CNICUS. *Gen. pl. 1255.**lanceolatus.*

1. C. foliis decurrentibus hispidis pinnatifidis, laciinis bilobis divaricatis spinosis, calycibus ovatis arachnoideo-pubescentibus, squamis lanceolatis spinosis patentibus.
Willd. sp. pl. 3. p. 1666.

Carduus lanceolatus. Sp. pl. 1149.

On road sides; common every where. ♂. June—
Sept. v. v. Flowers large, purple.

altissimus.

2. C. foliis sessilibus oblongo-lanceolatis scabris subtus tomentosis dentatis ciliatis, radicalibus pinnatifidis, calycibus bracteatis ovatis, squamis ovato-lanceolatis spinosis adpressis. *Willd. sp. pl. 3. p. 1671.*

*Carduus altissimus. Sp. pl. 1154.**Carduus virginianus. Walt. fl. car. 195.**Cirsium repandum. Mich. fl. amer. 2. p. 89.**Cirsium altissimum. Gron. fl. virg. 117.*

In old fields: Virginia and Carolina. ♀. July—Sept.
v. v. Flowers large, purple.

arvensis.

3. C. foliis sessilibus pinnatifidis glabriusculis ciliatis spinosis, caule paniculato, calycibus ovatis mucronatis, squamis lato-lanceolatis adpressis margine lanatis.

*Carduus arvensis. Smith. fl. brit. 2. p. 850.**Serratula arvensis. Willd. sp. pl. 3. p. 1646.*

Icon. Fl. dan. 644. Engl. bot. 975. Mart. fl. rust. 132.
In fields and on road sides: Canada and New England.

♀. July—Sept. v. v. Flowers small, purple. It
is one of the most troublesome weeds, and when once
introduced on a plantation it is almost impossible to
get rid of it.

muticus.

4. C. foliis omnibus pinnatifidis subtus lanuginosum: laciiniis spinulosum sublanceolatis acutis, ramis in summitate pluribus nudiusculis unifloris, calyce globoso, squamis muticis.

*Cirsium muticum. Mich. fl. amer. 2. p. 89.**Carduus carolinianus. Walt. fl. car. 195.*

In the mountains of Virginia and Carolina. ♂. July—
Sept. v. v. Flowers middle-sized, purple.

virginianus.

5. C. simpliciusculus; foliis sessilibus lanceolatis subtus cano-tomentosis remote dentatis, dentibus spinosis, floribus solitariis, calyce globoso, squamis mucronatis.

*Carduus virginianus. Willd. sp. pl. 3. p. 1659.**Cirsium virginianum. Mich. fl. amer. 2. p. 90.**Icon. Jacq. obs. bot. 4. t. 99.*In th
Sej

6. C. el

nos

fol

Serra

Cirsit

Cardi

On th

var

flow

1. C. inc

pec

ris

squ

In Pe

Are

1. L. ca

liati

sess

app

L. spic

Anony

Serrat

Icon.

In nat

—C

all t

spik

2. L. ca

pub

bus,

ame

Anony

Icon.

In mo

♀.

flow

In the mountains of Virginia and Carolina. ♀. July—
Sept. v. v. Flowers purple.

6. C. elatius; foliis sessilibus pinnatifidis acute incisis spinosissimis, involucro terminali unifloro polyphyllo: foliolis geminatum spinosissimum, calycibus inermibus. Serratula discolor. *Lam. encycl. 6. p. 565.*
Cirsium horridulum. Mich. fl. amer. 2. p. 90.
Carduus spinosissimus. Walt. fl. car. 194.
 On the banks of rivers and borders of woods: Pennsylvania to Carolina. ♀. July—Sept. v. v. Very tall; flowers large, pale yellow.

*horridulus.*608. CARDUUS. *Gen. pl. 1254.*

1. C. inermis; foliis decurrentibus lanceolatis pinnatifido-pectinatis, pedunculis terminalibus longissimis unifloris subaphyllis, floribus defloratis cernuis calycinis squamis linearibus patulis.—*Willd. sp. pl. 3. p. 1655.*
 In Pennsylvania. ♂. +. Flowers purple, the size of *Arctium Lappa*. It probably is a species of *Cnicus*.

*pectinatus.*609. LIATRIS. *Gen. pl. 1263.** *Spicatae seu racemosae; bulbosae.*

1. L. caule simplici, elato; foliis linearibus glabris basi ciliatis nervosis et punctatis, spica longissima, floribus sessilibus, squamis calycinis linear-oblongis obtusis appressis.—*Mich. fl. amer. 2. p. 91.*
L. spicata. Willd. sp. pl. 3. p. 1636.
Anonymos graminifolius. Walt. fl. car. 197.
Serratula spicata. Sp. pl. 1147.
Icon. Bot. rep. 401. Pluk. alm. t. 424. f. 6.
 In natural meadows: Pennsylvania to Carolina. ♀. Aug.—Oct. v. v. From three to six feet high; flowers, as all the following species, purple, in very long and close spikes.
2. L. caule simplici hirsuto, foliis strictis angusto-linearibus pubescentibus, spica longa, floribus confertim sessilibus, calycibus appressis superne squarrosis.—*Mich. fl. amer. 2. p. 91.*
Anonymos pilosa. Walt. fl. car. 197.
Icon. Dill. eth. t. 72. f. 83.
 In mountain meadows: Virginia, Illinois, and Carolina. ♀. Aug.—Oct. v. v. From two to four feet high; flowers smaller than No. 1.

*macrostachya.**pyrenostachya.*

graminifolia. 3. L. caule simplici glabro, foliis linearibus longissimis glabris nervosis margine scabriusculis costa media interne subpilosis, floribus spicatis remotiusculis subsessilibus, calycinis squamis oblongis obtusis mucronatis ciliatis appressis, interioribus coloratis.—*Willd. sp. pl. 3. p. 1636 exlus. syn. Walteri.*

Serratula compta. Herb. Banks. mss.

In South Carolina and Georgia. *Bartram. 4. Aug.—Oct. v. v.* About two or three feet high; flowers the size of No. 2; leaves very long and narrow.

heterophylla. 4. L. caule simplici glabro, foliis lanceolatis glabris laevibus: superioribus lineari-lanceolatis multoties minoribus, calycibus spicatis brevissime pedunculatis subsquarrosis: squamis lanceolatis acutis nudis.—*Willd. enum. 503. L. varia. Herb. Banks. mss.*

In South Carolina and Georgia. *Fraser. Bartram. 4. Aug.—Oct. v. s. in Herb. Banks. nec non Lambert.* Flowers the size of the preceding.

aspera. 5. L. caule subramoso scabro-pubescente, foliis linearilanceolatis asperrimis, calycibus brevibus spicatis distincte alternis solitariis sessilibus: squamis rotundato-obtusis conniventibus.—*Mich. fl. amer. 2. p. 92.*

Anonymos ramosus. Walt. fl. car. 198.

In South Carolina and Illinois. *Walter. Michaux. 4. Aug.—Oct. v. s.* Flowers larger than any of the foregoing species.

cylindracea. 6. L. gracilis, tota hirsuta; foliis gramineis, spica rariflora, calycibus subsessilibus cylindraceis paucifloris, squamis apice rotundatis abrupte mucronatis. *Mich. fl. amer. 2. p. 93.*

In woods and meadows of South Carolina and Illinois. *4. Aug. Sept. v. s. in Herb. Fraser.*

pilosa. 7. L. caule simplici pubescente, foliis linearibus pilosis ciliatis, calycibus racemosis laxiusculis: squamis linearioribz oblongis obtusiusculis, pedicellis bracteolatis.—*Willd. sp. pl. 3. p. 1636.*

Anonymos ciliatus. Walt. fl. car. 197.

In pine-barrens and sandy fields: New Jersey to Carolina. *4. Sept.—Nov. v. v.* A low species; flowers the size of No. 2.

gracilis. 8. L. caule simplici glabro, foliis linearibus nudis, calycibus racemosis subglobosis: squamis oblongis obtusiusculis appressis, pedicellis elongatis patentibus squamoso-bracteolatis.

In Ge
Flo

9. L. cat
bro
caly
sp.

Serrat
Stæhe
Eupat
In san
v. v
spik

10. L. cat
peti
neal
bosí
liati
On big
—O

11. L. cau
que
mos
spatl
sp. t
L. squa
Serrato
Icon.
In moi
—O

12. L. caul
nerv
lios
rigid
Serratu
Pteroni
Icon. i
In -san
Caro
some
inhal
snak
anim
it to

In Georgia. *Bartram.* Aug.—Oct. *v. s. in Herb. Banks.*

Flowers smaller than any preceding species.

9. L. caule simplici villoso, foliis linearibus subfalcatis sca-
bro-punctatis, spica subfoliosa, pedicellis brevibus,
calycinis squamis intimis ligulatis coloratis.—*Willd.*
sp. pl. 3. p. 1635.

elegans.

Serratula speciosa. *Ait. kew. 3. p. 138.*

Stæhelina elegans. *Walt. fl. car. 202.*

Eupatorium speciosum. *Vent. hort. cels. 79.*

In sandy fields: Carolina to Florida. 2. Sept.—Nov.
v. v. A beautiful perennial; flowers in long close
spikes, purplish-red.

10. L. caule simplici pubescente, foliis lævibus: inferioribus
petiolatis lato-lanceolatis, superioribus lanceolato-li-
nearibus, calycibus racemosis solitariis alternis subglo-
bosis: squamis ovatæ acutiusculis erectis, margine ci-
liatis.—*Mich. fl. amer. 2. p. 92.*

sphaeroidea.

On high mountains of Virginia and Carolina. 2. Aug.
—Oct. *v. v.* Flowers large and handsome.

scariosa.

11. L. caule simplici subpubescente, foliis lanceolatis utrin-
que attenuatis glabris margine scabris, calycibus race-
mosis alternis distantibus inferne squarrosis; squamis
spathulatis margine membranaceo coloratis.—*Willd.*
sp. pl. 3. p. 1635.

L. squarrulosa. *Mich. fl. amer. 2. p. 92.*

Serratula scariosa. *Sp. pl. 1147.*

Icon. *Pluk. alm. t. 177. f. 4.*

In mountain meadows: Virginia to Carolina. 2. Aug.
—Oct. *v. v.* Flowers the size of No. 10.

squarrosa.

12. L. caule simplici pubescente, foliis longissime linearibus
nervosis margine scabriusculis, racemis paucifloris fo-
liosis, calycinis squamis superne foliaceis lanceolatis
rigidis patentibus.—*Willd. sp. pl. 3. p. 1634.*

Serratula squarrosa. *Hort. cliff. 392.*

Pteronia caroliniana. *Walt. fl. car. 202.*

Icon. *Dill. elth. t. 71. f. 82.*

In sandy woods and fields: Virginia, Kentucky, and
Carolina. 2. Sept-Oct. *v. v.* Flowers large, very hand-
some. This and the preceding are known among the
inhabitants of those countries by the name of *Rattle-
snake's Master.* In case of being bit by this horrible
animal, they bruise the bulbs of this plant and apply
it to the wound, while, at the same time, they make

a decoction of it in milk, which is taken inwardly, in the same manner as I have mentioned under *Prenanthes serpentaria*.

** *Paniculae seu corymbosae; non bulbosae.*

pauciflora. 13. L. caule simplici glabro; foliis linearibus, panicula virgata foliosa, ramis brevibus paucifloris, calycibus subsessilibus secundis 3—5-floris: squamis erectis lanceolatis acutis glabris.

In Georgia. *Bartram.* 4. v. s. in *Herb. Banks.*
Flowers small, the size of No. 4.

paniculata. 14. L. caule simplici panicula calycibusque piloso-viscosis, foliis inferioribus lanceolatis attenuatis nervosis glabris: caulinis minoribus sessilibus, panicula coarctata fasciculata, calycibus sub-5-floribus: squamis lanceolatis acutiusculis.—*Willd. sp. pl. 3. p. 1037.*

Anonymos paniculatus. *Walt. fl. car.* 198.

In pine-barrens: Virginia to Florida. 4. Aug.—Oct.
v. s. in *Herb. Enslen.* Flowers the smallest of all the species here enumerated.

odoratissima. 15. L. glaberrima; caule simplici; foliis radicalibus oblongis, caulinis amplexicaulibus, panicula corymbosa axodivaricata, calycibus sub-8 floribus: squamis lanceolatis obtusiusculis.—*Willd. sp. pl. 3. p. 1037.*

Anonymos odoratissimus. *Walt. fl. car.* 198.

Icon. *Bot. rep.* 633.

In open swamps: Carolina to Florida. 4. Sept. Oct.
v. v. Flowers a little larger than the preceding. It is generally called *Carolina Vanilla-plant*. The leaves, when dry, give a very agreeable scent, resembling somewhat that of Vanilla, which they maintain for years if kept dry and inclosed.

tomentosa. 16. L. caule simplicissimo foliisque cuneato-lanceolatis hirsutis, corymbo paucifloro depresso divaricato, ramis 1—2-floris, calycibus tomentosis: squamis ovatis acutis.—*Mich. fl. amer. 2. p. 93.*

In open swamps: Virginia and North Carolina. 4.
Aug.—Oct. v. v. About eighteen inches high; flowers the size of No. 1.

bellidifolia. 17. L. pumila, glabella; foliis cuneato-ob lanceolatis, corymbo inaequali, floribus omnibus longiuscule pedicellatis, calycinis squamis oblongo-ovalibus obtusis.
Mich. fl. amer. 2. p. 93.

Anony
In dry
4.

1. V. cau
oblo
lato.

Chrys

2. V. utr

3. V. alte

In Sou

as al

2. V. cau

bus

squa

Chryso

In barr

Sept.

figur

3. V. folii
basis

squa

In natu

4.

V. cau

bris

cory

ticis.

Serratul

Chryso

Icon.

alm.

On road

Carol

lookin

5. V. altis

coryn

bus.

Serratul

Chrysoc

Icon. L

On road

Aug.

Anonymos uniflorus. Walt. fl. car. 198.
In dry gravelly fields of North Carolina. *Michaux.*
 \mathcal{U} . +.

610. VERNONIA. *Gen. pl. 1262.*

1. V. caule simplici nudiusculo, foliis serratis: radicalibus oblongo-ovatis, caulinis lanceolatis, corymbo paniculato.—*Mich. fl. amer. 2. p. 94.*
Chrysocoma acaulis. *Walt. fl. car. 196.*
 a. V. utroque flore pedunculato. verna.
 b. V. altero flore subsessili. autumnalis.
In South Carolina. \mathcal{U} . v. s. in *Herb. Walter.* Flowers, as all the following, purple.
2. V. caule simplici, foliis crebris longe angusteque linearibus subintegris, corymbo subumbellato, calycinis squamis rigide mucronatis. *Mich. fl. amer. 2. p. 94.*
Chrysocoma angustifolia. *Walt. fl. car. 196.*
In barren sandy woods: Virginia to Georgia. \mathcal{U} . Aug. Sept. v. s. in *Herb. Enslen.* Flowers the size and figure of *V. præalta.*
3. V. foliis longo-linearibus rarer serratis, floribus corymbosis erecto approximatis, calycibus ovoideis lœvibus, squamis muticis. *Mich. fl. amer. 2. p. 85.*
In natural meadows: Illinois *Michaux.* Virginia. *Ph. U. S. N. H. 1820. p. 10.* \mathcal{U} . Aug.—Oct. v. v. Flowers small.
fasciculata.
4. V. caule altissimo anguloso dense pubescente, foliis crebris lanceolatis acute serratis subtus pubescensibus, corymbo fastigiato, calycinis squamis ovatis acutis muticis.—*Willd. sp. pl. 3. p. 1633.*
Serratula præalta. *Sp. pl. 1146.*
Chrysocoma tomentosa. *Walt. fl. car. 196.*
Icon. *Dill. elth. t. 264. f. 343.* *Mill. ic. 234.* *Pluk. alm. t. 280. f. 6.*
On road sides and borders of woods: New England to Carolina. \mathcal{U} . Aug.—Oct. v. v. A tall rough-looking plant.
præalta.
5. V. altissima; foliis crebris lanceolatis serrulatis scabris, corymbo fastigiato, calycinis squamis apice filiformibus.—*Willd. sp. pl. 3. p. 1632.*
Serratula noveboracensis. *Hort. cliff. 392.*
Chrysocoma gigantea. *Walt. fl. car. 196.*
Icon. *Dill. elth. t. 263. f. 342.* *Pluk. alm. t. 109. f. 3.*
On road sides and old pastures: Canada to Carolina. \mathcal{U} . Aug.—Oct. v. v. About five feet high.
noveboracensis.

611. KUHNIA. *Gen. pl. 1272.*

*Critonia. Gært. carp. 2. p. 411.
Dalea. Brown. jam. 314.*

eupatorioides. 1. *K. glabra*; foliis petiolatis lato-lanceolatis serratis, corymbo terminali paucifloro coarctato.—*Willd. sp. pl. 3. p. 1772.* *

Eupatorium alternifolium. Ard. spec. 2. p. 40. t. 20.

Icon. Linn. fil. dec. t. 11. Pluk. alm. t. 87. f. 2.

In shady woods, about rocks: Pennsylvania to Virginia.

4. Aug. Sept. v. v. Flowers white; resembles an *Eupatorium* exceedingly.

Critonia.

2. *K. pubescens*; foliis angusto-lanceolatis inferne subdentatis petiolatis subtus punctatis: superioribus linearibus integerrimis sessilibus, panicula terminali divaricata.—*Willd. sp. pl. 3. p. 1773.*

Critonia Kuhnii. Mich. fl. amer. 2. p. 101.

Icon. Gært. carp. 2. t. 174. f. 7.

On high mountains of Pennsylvania and Virginia. 4. Aug. Sept. v. v. Flowers pale yellow.

612. EUPATORIUM. *Gen. pl. 1272.*

* *Calycibus 3—5-floris.*

fœniculace-
um.

1. *E. caule paniculato, foliis glabris inferioribus pinnatis, superioribus fasciculatis, omnibus filiformibus.—Willd. sp. pl. 3. p. 1750.*

E. fœniculoides. Walt. fl. car. 199.

Chrysocoma capillacea. Mich. fl. amer. 2. p. 101.

In fields and on road sides, near the sea-coast: Virginia to Florida. 4. Aug.—Oct. v. v. From two to five feet high, resembling an *Artemisia* very much; flowers in great abundance, small, pale yellow.

coronopifoli-
um.

2. *E. caule paniculato, foliis infimis pinnatifidis, reliquis indivisis fasciculatis linearibus integerrimis. Willd. sp. pl. 3. p. 1750.*

E. compositifolium. Walt. fl. car. 199.

Chrysocoma coronopifolia. Mich. fl. amer. 2. p. 102.

In dry woods, on road sides: Carolina. 4. Aug. Sept. v. v. Flowers white, double the size of the preceding.

hyssopifoli-
um.

3. *E. foliis oppositis subverticillatis linearibus integerrimis pubescentibus trinervibus punctatis, radicalibus subdentatis. Willd. sp. pl. 3. p. 1749.*

Icon. L
In dry p
lina.
leaves
unless

4. *E. foliis bus p
superi fl. car
In low s
— No dots.*

5. *E. foliis ceolat briuse About r the A About*

6. *E. foliis basi tr Willd
In shady J species*

7. *E. foliis serratis lati sc In woods Aug.—*

8. *E. foliis basi pr —Wil
In low cnia. ceding*

9. *E. foliis basi gr pl. 3. p. 1
E. pilosu E. verber Icon. W
In swam 4. At*

- Icon. Dill. eth. t. 115. f. 120. Pluk alm. t. 88. f. 2.
In dry pine-barrens and sandy fields: New Jersey to Carolina. ♀. Aug.—Oct. v. v. About a foot high; leaves small: flowers white, as all the following are, unless a different colour is mentioned.
4. E. foliis linear-lanceolatis denticulatis obsolete 3-nervi *linearifolium*.
bus pubescentibus, inferioribus oppositis verticillatis, superioribus alternis. Willd sp. pl. 3. p. 1750. Walt. fl. car. 199.
In low sandy fields: New Jersey to Carolina. ♀. Aug.
—Nov. v. v. The calyx is covered with resinous dots.
 5. E. foliis sessilibus amplexicaulibus distinctis ovato-lan- *sessilifolium*.
ceolatis basi rotundatis serratis glaberrimis, caule gla-
briusculo. Willd. sp. pl. 3. p. 1751.
About rocks, particularly lime-stone, and throughout
the Allegany mountains. ♀. Aug. Sept. v. v.
About two feet high; very smooth.
 6. E. foliis sessilibus amplexicaulibus distinctis lanceolatis *truncatum*.
basi truncatis serratis glabriusculis, caule pubescente.
Willd. sp. pl. 3. p. 1751.
In shady woods, about rocks: Pennsylvania to Virginia.
♀. July—Oct. v. v. Resembles the preceding species.
 7. E. foliis subsessilibus oblongo-lanceolatis scabriusculis
serratis, calycinis squamis interioribus elongatis lanceo-
latis scariosis coloratis. Willd. sp. pl. 3. p. 1752.
In woods and swamps: Pennsylvania and Virginia. ♀.
Aug.—Oct. v. v. About eighteen inches high.
 8. E. foliis sessilibus distinctis oblongo-lanceolatis scabris *lanceolatum*.
basi profunde serratis, calycinis squamis concoloribus.
—Willd. sp. pl. 3. p. 1752.
In low copes and dry swamps: Pennsylvania to Virgi-
nia. ♀. Aug.—Nov. v. v. Resembles the pre-
ceding species.
 9. E. foliis sessilibus distinctis ovatis scabris, superioribus *teucrifolium*
basi grosse serratis, summis integerrimis. Willd. sp.
pl. 3. p. 1753.
 - E. pilosum. Walt. fl. car. 199.
 - E. verbenæfolium. Mich. fl. amer. 2. p. 98.
Icon. Willd. hort berol. 32
In swamps and low woods: New England to Carolina.
♀. Aug.—Nov. v. v. About two feet high.



- cuneifolium*. 10. E. foliis petiolatis obovato-lanceolatis apice subserratis triplinervibus utrinque pubescentibus. *Willd. sp. pl. 3. p. 1753.*
E. Marrubium. *Walt. fl. car. 199.*
In low sandy fields: Virginia and Carolina. ♀. Aug.—Oct. v. v. Not above a foot high.
- mellissoides*. 11. E. foliis petiolatis ovatis obtusiusculis obtuse serratis venosis glabriusculis. *Willd. sp. pl. 3. p. 1754.*
In Pennsylvania. ♀. Aug.—Oct. +. Resembles No. 9; but the leaves are smaller, petiolated, and smooth.
- rotundifoli- um.* 12. E. foliis sessilibus distinctis subrotundo-cordatis obtuse serratis venosis, calycinis squamis acuminatis.—*Willd. sp. pl. 3. p. 1754.*
Icon. *Pluk. alm. t. 88. f. 4.*
In swamps and shady woods: Canada to Carolina. ♀. Aug.—Nov. v. v. About a foot or eighteen inches high.
- pubescens*. 13. E. foliis sessilibus distinctis ovatis scabris venosis, inferioribus duplicato-serratis, superioribus subserratis, caule paniculato pubescente, ramis fastigiatis. *Willd. sp. pl. 3. p. 1755.*
E. glandulosum. *Mich. fl. amer. 2. p. 98?*
In dry sandy woods: New Jersey to Carolina. ♀. Aug.—Oct. v. v. Not above two feet high.
- ceanothifoli- um.* 14. E. foliis petiolatis ovatis acuminatis dentatis triplinervibus glabris. *Willd. sp. pl. 3. p. 1755.*
In shady woods, about rocks: New York to Virginia. ♀. Aug.—Nov. v. v. Resembles *Ceanothus americanus* in its foliage exceedingly.
- altissimum*. 15. E. foliis subsessilibus lanceolatis 3-nervibus utrinque attenuatis pubescentibus, inferioribus medio serratis. *Willd. sp. pl. 3. p. 1754.*
Icon. *Jacq. hort. vind. 164.*
In low sandy woods: Pennsylvania and Virginia, and on the banks of the Mississippi and Missouri. ♀. Aug.—Oct. v. v. From three to seven feet high.
- amoenum*. 16. E. foliis brevi-petiolatis oppositis ternatisque lanceolato-oblongis utrinque acutis serratis glabriusculis subrugosis subtus reticulato-venosis, panicula corymboso-fasciculata confertiflora, calycinis squamis oblongis acutiusculis coloratis.
On the New Jersey mountains. ♀. Sept. Oct. v. v.
- About
dunel
collec
purple
17. E. foliis
nuat
In swa
Virgin
out br
18. E. glab
acumi
nis e:
p. 99.
On the t
♀. +
19. E. foliis
rat
Willd
Icon. *R*
s. 7. t.
In swa
Canad
tall ; 1
20. E. foliis
æquali
sulcate
Icon. *H*
f. 3.
In swa
rolina,
purple
ers pur
21. E. foliis
serratis
eu,
E. macul
In the Bl
♀. At
flowers
22. E. foliis
basi cu
solido

About two feet high; stem solid, smooth, purple; peduncles tomentose; flowers small, but very closely collected in a corymbose panicle, of a beautiful pale purple inclining to flesh colour.

17. *E. foliis petiolatis ternis quaternis ovatis utrinque attenuatis serratis scabriusculis.* *Willd. sp. pl. 3. p. 1750.*
In swamps and on the banks of rivers: New York to Virginia. ♀. Aug.—Oct. v. v. A tall plant without branches.
18. *E. glabellum*; *foliis quaternis ovali-lanceolatis utrinque acuminatis subfalcatis rariter serratis, squamis calycinis extimis brevibus ovalibus.* —*Mich. fl. amer. 2. p. 99.*
On the banks of the Ohio and Scioto rivers. *Michaux.*
♀. +. A tall plant, resembling *E. album*.

** *Calycibus plus quam 5-florisi.*

19. *E. foliis petiolatis quaternis quinise ovato-lanceolatis serratis rugoso-venosis scabriusculis, caule fistuloso.* *Willd. sp. pl. 3. p. 1759.*
Icon. *Rob. ic. 217.* *Corn. canad. t. 72.* *Moris. hist. 3. s. 7. t. 13. f. 4.* *Dod. mem. t. 217.*
In swamps, wet woods, and on the banks of rivulets: Canada to Virginia. ♀. Aug.—Oct. v. v. Very tall; flowers purple.
20. *E. foliis petiolatis quaternis quinise ovato-lanceolatis inaequaliter serratis subtus pubescentibus, caule solido sulcato.* *Willd. sp. pl. 3. p. 1760.*
Icon. *Herm. parad. t. 158.* *Moris. hist. 3. s. 7. t. 18. f. 3.*
In swamps and near ponds and rivulets: Canada to Carolina. ♀. Aug.—Oct. v. v. Stem marked with purple lines, not so tall as the preceding species; flowers purple.

21. *E. foliis petiolatis quaternis quinise ovatis acuminatis serratis utrinque scabris, caule solidi tereti.* *Willd. enum. 853.*
E. maculatum. Hortul.
In the Blue-mountains: New Jersey and Pennsylvania.
♀. Aug.—Oct. v. v. Not so tall as the preceding; flowers purple, very ornamental.

22. *E. foliis petiolatis ternis quaternis ovato-lanceolatis, verticillatum.*
basi cuneatis inaequaliter serratis glabriusculis, caule solidi laevi. *Willd. sp. pl. 3. p. 1760.*

trifoliatum.

salcatum.

purpureum.

maculatum.

punctatum.

On the banks of rivers and in swamps; frequent: New York to Carolina. *v. v.* From four to six feet high; flowers purple. It differs from No. 18 in its stem, broader and smoother leaves, and white calyces; from No. 19 in its stem and smooth leaves.

- perfoliatum.* 23. *E. foliis connato-perfoliatis oblongis sursum angustatis serratis rugosis subtus tomentosis, caule villoso.* — *Willd. sp. pl. 3. p. 1761.*

E. connatum. Mich. fl. amer. 2. p. 99.
Icon. Pluk. alm. t. 87. f. 6.

In low meadows and about springs: Canada to Florida. *q. Aug.—Oct. v. v.* From two to three feet high; flowers white. The whole plant is exceedingly bitter, and has been used for ages past by the natives and inhabitants in intermitting fevers. It is generally known by the name of *Thorough-wort* or *Bone-set*. I have stated a case of its efficacy in those diseases in a letter to William Royston, Esq. who inserted it in the Medical and Physical Journal. In which I stated the benefits derived from this plant, by myself and others during my stay in the neighbourhood of lake Ontario, when both the influenza and lake fever (similar to the yellow fever) were raging among the inhabitants. It is generally used as a decoction, or, which I consider the more effectual way, as an infusion in gin or rum.

- cœlestinum.* 24. *E. foliis petiolatis cordato-ovatis obtusiusculis triplineribus obtuse serratis, floribus corymbosis.* *Willd. sp. pl. 3. p. 1764.*

Icon. Dill. elth. t. 114. f. 159.

In mountain-woods of Virginia to Carolina. *q. Aug.—Oct. v. v.* Flowers large, beautiful blue.

- aromaticum.* 25. *E. foliis petiolatis ovatis acutis trinervibus obtuse-serratis glabris, caule superne paniculato, floribus corymbosis, calycibus simplicibus.* *Willd. sp. pl. 3. p. 1765.*

Icon. Pluk. alm. t. 88. f. 3.

In woods and on the banks of rivers: Pennsylvania to Florida. *q. Aug.—Oct. v. v.* Flowers large, clear white, sweet-scented.

- ageratoides.* 26. *E. foliis petiolatis ovatis acuminatis trinervibus inæqualiter grosse serratis glabris, corymbo multifloro divaricato, calycibus subsimplicibus.* *Willd. sp. pl. 3. p. 1765.*

E. urticatum. Mich. fl. amer. 2. p. 100.

Ageratum alissimum. Sp. pl. ed. 2. p. 1176.

Icon. Moris. hist. 3. s. 7. t. 18. f. 11.

The mo
and i
q. A
white

27. *E. foliis*
serrat
pulve
On the
Micha

613.

1. *M. cau*
tatis a
coryn
Eupator
Icon. J
On the
q. J
very r
is and
loured
hand

61

1. *C. foliis*
ribus
sito fa
fl. am
In low g
rolina
appea
coryn
eveh

2. *C. glabr*
florib
glabri
C. biflor
in He

- On high
q. O
high

3. *C. caly*

The most common species in the Allegany mountains, and near the sea-coast from Canada to Pennsylvania. ♀. Aug.—Oct. v. v. About two feet high; flowers white.

27. E. foliis longe-petiolatis subdeltoideo-lanceolatis rarer serratis, calycinis squamis tomentosis obtusis, caule pulverulento.—*Mich. fl. amer. 2. p. 100.*

On the sea-coast of Carolina, among Scirpus, &c. *Michaux.* ♀. Sept.—Nov. +. Flowers white.

serotinum.

613. MIKANIA. *Willd. sp. pl. 3. p. 1742.*

1. M. caule scandente glabro, foliis cordatis repando-dentatis acuminatis lobis divaricatis inaequalibus, floribus corymbosis. *Willd. sp. pl. 3. p. 1743.*

scandens.

Eupatorium scandens. Sp. pl. 1171.

Icon. Jacq. ic. rar. 1. t. 169. Pluk. alm. t. 163. f. 3.

On the banks of rivers and ditches: Canada to Carolina.

♀. July—Sept. v. v. A twining plant; flowers very numerous, white with a tinge of blue. There is another species, found in Carolina, with flesh-coloured fragrant flowers, but I have no materials at hand to give its description.

614. CHRYSOCOMA. *Gen. pl. 1019.*

1. C. foliis radicalibus spatulato-lanceolatis, caulinis linearibus rarer sparsis, caule subnudo, corymbo composto fastigato, calycibus oblongis 3—4-floris. *Mich. fl. amer. 2. p. 101.*

nudata.

In low grounds, in the forests of North and South Carolina. ♀. Aug.—Oct. v. v. A plant of singular appearance, the stem almost leafless; the large flat corymbo consists of small flowers, entirely yellow; even the calyx is deeply coloured.

2. C. glabra; foliis linearibus 3-nervibus punctato-scabris, floribus corymbosis congestis, calycibus laxis 5-floris glabris.—*Lam. encycl. 2. p. 192.*

dracunculoides.

C. biflora. *Sp. pl. 1178. secundum specimen Pallasianum in Herbario Lambertiano asservatum.*

On high cliffs on the banks of the Missouri. *M. Lewis.* ♀. Oct. v. s. in *Herb. Lewis.* From one to two feet high; flowers large, yellow.

3. C. calycibus pedunculis foliisque sublanato-pubescenti-

nauseosa.

bus ; foliis angustissime linearibus, corymbo laxiusculo, calycibus laxis 5-floris : squamis interioribus divaricatis glabris.—*Pallas. mss. in Herb. Lambert.*

On the banks of the Missouri. *M. Lewis.* ♀. Oct. *v. s. in Herb. Lewis.* Flowers yellow, somewhat smaller than the preceding.

615. CACALIA. *Gen. pl. 1275.**suaveolens.*

1. *C. caule herbaceo, foliis petiolatis hastato-sagittatis serratis glabris concoloribus, floribus corymbosis erectis, calycibus multifloris.*—*Willd. sp. pl. 3. p. 1734.*

On the banks of rivers : Pensylvania to Carolina. ♀. Aug.—Oct. *v. v.* From three to four feet high ; flowers white ; anthers yellow.

atriplicifolia.

2. *C. caule herbaceo, foliis petiolatis glabris subtus glaucis, radicalibus cordatis dentatis, caulinis rhombis utrinque subbidentatis, floribus corymbosis erectis, calycibus 5-floris.*—*Willd. sp. pl. 3. p. 1734.*

Icon. Pluk. alm. t. 101. f. 1. Moris. hist. 3. s. 7. t. 15. f. 7.

In low and overflowed places, near rivers : Canada to Carolina. ♀. Aug.—Oct. *v. v.* Flowers white, considerably smaller than the preceding.

reniformis.

3. *C. caule herbaceo, foliis petiolatis glabris subtus ad venas pilosis, radicalibus amplis cordatis reniformibus repando-dentatis, caulinis oblongis dentatis basi cuneatis integrerrimis, corymbis fastigiatis, calycibus multifloris.* *Willd. sp. pl. 3. p. 1735.*

On the banks of rivers : Pensylvania to Virginia. ♀. Aug.—Oct. *v. v.* From five to ten feet high ; flowers white.

616. SPARGANOPHORUS. *Gært. carp. 2. p. 395.**verticillatus.*

1. *S. foliis setaceo-linearibus verticillatis, caule subunifloro, papo campanulato 5-dentato.*—*Mich. fl. amer. 2. p. 95.*

Ethulia uniflora. *Walt. fl. car. 195. Willd. sp. pl. 3. p. 1742.*

Icon. Mich. fl. amer. 2. t. 42.

In overflowed swamps and rivulets : New Jersey, Carolina, and Florida. ♀. Aug.—Oct. *v. v.* Floating in the water ; flowers purple, the size of a large daisy.

617. H.

1. *H. ca-*
lac
so-
Rothi
In Ca
tris

618. M.

1. *M. fo-*
ter
tato
leis
men
Biden
Athani
α. *M. fo-*
Icon.
β. *M. fo-*
Icon.
On the
—C

Tran
Per

1. *M. cau-*
long
thula
Persoon
Athana
In the
Flow
all th
2. *M. cau-*
triner
tis, p
Persoon

VOL. II.

617. HYMENOPAPPUS. *L'Herit.* monog. *Mich. fl. amer.* 2. p. 103.

Rothia. *Lam. illustr. t. 667.*

1. *H. candicanti-lanuginosus*; foliis profunde pinnatifidis, laciniis linearis-oblongis subdentalis, floribus corymbo-so-paniculatis.—*Mich. fl. amer.* 2. p. 104.

Rothia carolinensis. *Journ. de hist. nat. no. 1. cum icon.*
In Carolina. ♂. +. Flowers white, the size of *Liatris scariosa*.

618. MELANANTHERA. *Mich. fl. amer.* 2. p. 106.

1. *M. foliis triplinerviis ovatis acuminatis scabris inaequaliter dentatis, inferioribus subcordatis, superioribus hastato-trilobis, pedunculis terminalibus corymbosis, paleis receptaculi lanceolatis acuminatis.* —*Mich. fl. amer.* 2. p. 107.

Bidens nivea. *Willd. sp. pl. 3. p. 1721.*
Athanasia hastata. *Walt. fl. car. 201.*

α. *M. foliis profunde trilobatis.*
Icon. Dill. elth. t. 47. f. 55.

β. *M. foliis leviter lobatis panduriformibus.*
Icon. Dill. elth. t. 46. f. 54.

On the banks of rivers: Carolina to Florida. ♀. Aug.
—Oct. v.v. Flowers large, white.

619. MARSHALLIA. *Gen. pl.*

Trattenickia. *Pers. syn. 2. p. 403.*

Persoonia. *Mich. fl. amer.* 2. p. 104.

1. *M. caule simplici inferne folioso superne nudato, foliis longo-lanceolatis, calycinis foliolis obtusis, paleis spatulatis.*

Persoonia lanceolata. *Mich. fl. amer.* 2. p. 105.

Athanasia obovata. *Walt. fl. car. 201.*

In the Carolina mountains. ♀. v.s. in *Herb. Walter.*
Flowers resembling a *Scabious*, pale purple, and like all the following very ornamental.

2. *M. caule simplici, foliis lanceolato-ovalibus acuminatis trinervibus, infinis vaginantibus, calycinis foliolis acutis, paleis angusto-linearibus.*

Persoonia latifolia. *Mich. fl. amer.* 2. p. 105. t. 43.

scabiosaeus.

hastata.

lobata.

pandurata.

lanceolata.

latifolia.

Athanasia trinervia. Walt. fl. car. 201.

In the Carolina mountains. ♀. v. s. in *Herb. Walter.*

angustifolia.

3. M. caule ramoso, foliis infimis angusto-lanceolatis, ramis angustissime linearibus, calycinis foliolis rigidis superne angustatis acutissimis, paleis setaceis.

Persoonia angustifolia. Mich. fl. amer. 2. p. 106.

Athanasia graminifolia. Walt. fl. car. 200.

In Tennessee. *Michaux.* In swamps, near Wilming-

ton, North Carolina. *Ph.* ♀. June, July. v. v.

The handsomest species of this genus.

620. SANTOLINA. *Gen. pl. 1278.*

suaveolens.

1. S. glabra; caule corymboso-ramoso, foliis subbipinnatifidis, lacinias acutis linearibus, pedunculis terminalibus unifloris.

On the banks of the Kooskoosky. *M. Lewis.* ⊖. June—Aug. v. v.; v. s. in *Herb. Lewis.* A small plant of an agreeable sweet scent; flowers yellow.

Planta glabra. Caulis erectus, ramosus: rami corymbo-

bosi uniflori. Folia sessilia tenuissime pinnatifida, la-

ciniis inciso-2-3-fidis, omnibus acutis angusto-lineari-

bus. Pedunculi terminales, solitarii, uniflori. Calyx

hemisphæricus, subimbricato-polyphyllus: foliolis ova-

ibus, obtusis, glabris, viridibus, margine albido-mem-

branaceis, laceris. Cor. discoidea. Discus globosus,

luteus. Corollulae germine oblique insertæ, tubulosæ,

4-fidæ? laciñis erectis. Genitalia inclusa. Recepta-

culum conicum, paleaceum. Paleis sparsis, oblongis,

obtusis. Pappus nullus. Semina obliquæ obovatae.

II. SUPERFLUA.

621. ARTEMISIA. *Gen. pl. 1281.*

* *Foliis simplicibus.*

integrifolia.

1. A. cano-tomentosa; caule erecto virgato; foliis linearilanceolatis acuminatis subtus tomentosis subdentatis, floribus ovatis subsessilibus erectis tomentosis.—*Willd.* *sp. pl. 3. p. 1846.*

Icon.
On t
La
fee

2. A. gl
flo
pl.

Icon.
On th
in

3. A. ca
vos
3-le
ribu

On th
in 1

4. A. ca

trilc
sis p
Icon.
t. 3

On the
Lan

5. A. foli
sis,
floris

Icon.
On the
Oct.

6. A. foli
natis,
moso

pl. 3.
Icon.
On the
Nov.

7. A. foli
acutis
dente

p. 18

Icon. *Gmel. fl. sib. 2. t. 48. f. 1* ♂ 2.

On the cliffs and dry savannahs of the Missouri. *M. Lewis.* ♀. Oct. *v. s. in Herb. Lewis.* About three feet high.

2. A. glabra; foliis linear-lanceolatis utrinque attenuatis, *Dracunculus.* floribus subrotundis pedunculatis erectis.—*Willd. sp. pl. 3. p. 1848.*

Icon. *Gmel. fl. sib. 2. t. 59* ♂ 60. f. 1.

On the Missouri. *M. Lewis.* ♀. Aug.—Oct. *v. s. in Herb. Lewis.*

3. A. canescens; foliis longe linear-lanceolatis acutis nervosis utrinque canis: inferioribus cuneiformibus acute-3-lobis, floribus ovalibus glomeratis sessilibus axillaribus.

On the Missouri. *M. Lewis.* ♀. Sept.—Nov. *v. s. in Herb. Lewis.*

4. A. canescens: foliis inferioribus cuneiformibus obtusis trilobis, superioribus linearibus obtusis, floribus globosis pedunculatis cernuis. *Willd. sp. pl. 3. p. 1848.*

Icon. *Gmel. fl. sib. 2. t. 61. f. 1* ♂ 2. *Pluk. amalthe. t. 353. f. 5.*

On the north-west coast. ♀. Sept. Oct. *v. s. in Herb. Lambert.* Flowers large.

cana.

chinensis.

** *Foliis compositis; caule paniculato.*

5. A. foliis caulinis pinnatis linearibus glabris, ramis indistinctis, spicis secundis reflexis, floribus subsessilibus 5-floris. *Willd. sp. pl. 3. p. 1826.*

Icon. *Gmel. fl. sib. 2. t. 51.* *Lob. ic. 756.*

On the plains of the Missouri. *M. Lewis.* ♀. Sept. Oct. *v. s. in Herb. Lewis.*

Santonica.

6. A. foliis caulinis setaceis pinnatis glabris, radicalibus pinnatis, lacinias 3-fidis incanis, caule procumbente ramoso virgato, floribus ovatis pedunculatis. *Willd. sp. pl. 3. p. 1827.*

Icon. *Engl. bot. 338.*

On the plains of the Missouri. *M. Lewis.* ♀. Sept.—Nov. *v. s. in Herb. Lewis.*

campestris.

7. A. foliis incanis pinnatis, pinnis tripartitis linearibus acutis, floralibus pinnatis 3-partitis, caule adscendente, floribus globosis nutantibus. *Willd. sp. pl. 3. p. 1838.*

frigida.

On the plains of the Missouri. *M. Lewis.* 4. Oct.
Nov. v. s. in *Herb. Lewis.*

liennis.

8. A. foliis glabris, radicalibus tripartito-pinnatis, caulinis inferioribus pinnatifidis, superioribus indivisis linearibus, caule stricto, floribus subrotundis subsessilibus erectis. *Willd. sp. pl. 3. p. 1842.*

A. hispanica. *Jacq. ic. rar. 1. t. 172.*

On the banks of the Missouri. ♂. v. s. in *Herb. Lambert.*

vulgaris.

9. A. foliis subtus tomentosis, caulinis pinnatifidis, lacinii lanceolatis subdentatis acutis, floralibus indivisis linearibus lanceolatis, floribus subsessilibus oblongis erectis, calycibus tomentosis. *Willd. sp. pl. 3. p. 1845.*

Icon. *Engl. bot. 978.*

On the banks of rivers: Canada to New England. 4. Sept.—Nov. v. v.

canadensis.

10. A. subdecumbens, parce pubescens; foliis planis linearipinnatifidis, ramulis spicifloris, floribus subhemisphaericis, calycibus scariosis. *Mich. fl. amer. 2. p. 129.*

In the moveable sands about Hudson's Bay. *Michaux.* +. Receptacle smooth.

caudata.

11. A. erecta, glabra; foliis subsetaceo-pinnatifidis, lacinii convexis, ramulis confertis, racemis terminalibus longissimis strictis, floribus pedicellatis globoso-ovatis.—*Mich. fl. amer. 2. p. 129.*

On the gravelly banks of the Missouri. *Michaux.* +.

*** *Foliis compositis; caule simplicissimo.*

spitkamica.

12. A. pumila, sericeo-pubescent; foliis inferioribus linearibus superne pinnatifidis, lacinii paucis linearibus acutis, floralibus simplicibus, floribus racemosis brevipedunculatis hemisphaericis, calycibus scariosis.

In Labrador. *Colmaster.* 4. Sept. Oct. v. s. in *Herb. Dickson.* Flower large, in proportion to the size of the plant; receptacle naked.

622. TANACETUM. *Gen. pl. 1280.*

vulgare.

1. T. foliis bipinnatis incisis serratis. *Willd. sp. pl. 3. p. 1814.*

Icon. *Oed. dan. 871.*

In cu
to
Ta

1. B. gl
line
ealy
p. 1

On th
the
v. s
low

2. B. fol
glo
mis
p.

In wo

Aug

3. B. foli
posi
p. 1

Icon.

t. 27

On th

Nov

whi

1. C. her
acut
tern
squar
Mic

Erigeron
Icon.

In ove
to C
high
a str

2. C. herb
latis

In cultivated grounds and on road sides : New England to Pennsylvania. ♀. July—Sept. v. v. Common *Tansy* is probably introduced from Europe.

623. BACCHARIS. *Gen. pl.* 1285.

1. B. glaberrima, paniculato-ramosissima ; foliis angusto-linearibus integerrimis, panicula composita multiflora, calycibus parvulis sub-20-floris.—*Mich. fl. amer.* 2. p. 125.
On the sea-coast : Carolina to Florida ; on the banks of the Mississippi. *M. Lewis.* ♂. July—Sept. v. v. ; v. s. in *Herb. Lewis.* Flowers white, as all the following are.
2. B. foliis laevibus cuneato-ovalibus superne dentatis, *glomerulifloris* glomerulis florum axillaribus sessilibus remotis, squamis calycinis superne rans.—*Mich. fl. amer.* 2. p. 125.
In woods, on the coast of Virginia and Carolina. ♂. Aug.—Oct. v. v. Resembles the following.
3. B. foliis obovatis superne inciso-dentatis, panicula composita foliosa, fasciculis pedunculatis.—*Willd. sp. pl.* 3. p. 1915.
Icon. *Schmidt arb.* 82. *Herm. parad.* 225. *Pluk. alm.* t. 27. f. 2.
On the sea-coast : Maryland to Florida. ♂. Sept.—Nov. v. v. The whole shrub is covered with a white powder.

624. CONYZA. *Gen. pl.* 1286.

1. C. herbacea, pubescens ; foliis sessilibus lato-lanceolatis acutis serratis : serraturis apice cartilagineis, corymbis terminalibus fastigiatis coarctatis subaphyllis, calycinis squamis subulato-mucronatis flosculis brevioribus.—*Mich. fl. amer.* 2. p. 126. *exclus. syn. Walteri.*
Erigeron camphoratum. Willd. sp. pl. 3. p. 1960.
Icon. *Dill. elth.* t. 88. f. 104.
In overflowed places, near rivers and ponds : New York to Carolina. ♂. Aug.—Oct. v. v. About a foot high ; flowers reddish-purple ; the whole plant emits a strong scent of camphor, only more disagreeable.
2. C. herbacea, subpubescens ; foliis petiolatis ovato-lanceolatis acutissimis subrepando-denticulatis : denticulis

glandulosis, corymbis terminalibus et axillaribus folio brevioribus, calycinis squamis acutis flosculos subæquantibus.

Baccharis foetida. *Willd. sp. pl. 3. p. 1918. Walt. fl. car. 202.*

Icon. *Dill. elth. t. 89. f. 105.*

On the banks of rivers and in large swamps: South Carolina and Georgia. ♀. Aug. Sept. v. v. Flowers purple, smaller than the preceding; stem about three feet high.

bifrons.

3. C. herbacea, subglutinosa; foliis amplexicaulibus spatulato-oblongis acutis serratis, paniculæ corymbulis capitato-globose.—*Willd. sp. pl. 3. p. 1920.*

C. amplexicaulis. *Mich. fl. amer. 2. p. 126.*

C. uliginosa. *Pers. syn. 2. p. 427.*

Baccharis viscosa. *Walt. fl. car. 202.*

Icon. *Pluk. alm. t. 87. f. 4.*

In low and overflowed fields: Carolina. ♀. July, Aug. v. s. in *Herb. Gronov.* Flowers pale yellow.

pycnostachya.

4. C. caule alato, foliis lanceolatis subtus tomentosis subintegerrimis, spica cylindrica densiflora.—*Mich. fl. amer. 2. p. 126.*

C. spicata. *Cav. ic. 1. p. 8. t. 12?*

Gnaphalium undulatum. *Walt. fl. car. 203.*

In dry fields and barren woods: Carolina to Florida. ♀. July—Sept. v. v. A singular-looking species, nearly allied to *C. virgata.*

625. GNAPHALIUM. *Gen. pl. 1282.*

margaritaceum.

1. G. herbaceum; foliis linear-lanceolatis sensim angustatis acutis, caule superne ramoso, corymbo fastigiato, floribus pedicellatis.—*Willd. sp. pl. 3. p. 1881.*

Icon. *Engl. bot. 2018.*

In Canada and on the mountains of New York and New Jersey. ♀. Aug.—Oct. v. v. About eighteen inches high; flowers large, white, with a yellow disk, very handsome.

polycephalum.

2. G. herbaceum, erectum; foliis linear-lanceolatis acutis, supra glabris, subtus pubescentibus, caule paniculato tomentoso, corymbis terminalibus coarctatis.—*Mich. fl. amer. 2. p. 127.*

G. obtusifolia. *Dill. elth. t. 89. f. 105.*
In old July flow
3. G. hermaphroditica. *Gloucest. p. 18.*
Icon. In sand —O calyx
4. G. sarcocaulis. *Mich. fl. amer. 2. p. 126.*
Icon. 1 In woody in th Not plain
5. G. caeruleum. *Gnaphal. naceum.*
Icon. 1 In Lat. Not
6. G. caulinatum. *Mich. fl. 3.*
Icon. 1 In stony v. v. color
7. G. caulinatum. *Mich. fl. 18.*

- G. obtusifolium. Willd. sp. pl. 3. p. 1880. excl. syn.
Dillenii.
In old fields and woods: New England to Carolina. ♂. July—Sept. v. v. About a foot high, spreading; flowers yellowish-white, not shining.
3. G. herbaceum; foliis linearispathulatis subtus tomentosis, caule erecto simplicissimo, floribus sessilibus glomeratis terminalibus et axillaribus.—Willd. sp. pl. 3. p. 1884.
Icon. Dill. eth. t. 109. f. 132.
In sandy barren soil: New York to Carolina. ♀. July—Oct. v. v. Not above a span high; flowers small, calyces purple.
4. G. sarmentis procumbentibus, caule simplicissimo, foliis radicalibus ovatis nervosis micronatis, corymbo coarctato, floribus dioicis, squamis calycinis interioribus elongatis acutiusculis coloratis.—Willd. sp. pl. 3. p. 1882.
Icon. Pluk. alm. t. 348. f. 9.
In woods and on sunny hills: Canada to Carolina, and in the Allegany mountains. ♀. May—July. v. v. Not above six inches high; leaves large, resembling plaintain leaves; flowers white.
5. G. caule non sarmentoso simplicissimo, foliis radicalibus lanceolatis, floribus terminalibus aggregatis sessilibus, calycinis squamis interioribus elongatis acutis membranaceis. Willd. sp. pl. 3. p. 1883.
Icon. Fl. dan. 332.
In Labrador. Colmaster. ♀. v. s. in Herb. Dickson. Not above four inches high; resembling *G. dioicum*.
6. G. caule herbaceo simplicissimo erecto, foliis angustolanceolatis utrinque attenuatis et lanatis, floribus terminalibus axillaribusque sessilibus spicatis. Willd. sp. pl. 3. p. 1884.
Icon. Fl. dan. 254.
In stony woods: New York and Canada. ♀. July, Aug. v. v. About a foot or less high; calyces shining, straw-coloured with brown.
7. G. caule herbaceo erecto ramoso, foliis obovato-spathulatis subtus pubescentibus, floribus axillaribus et terminalibus glomeratis spicatis. Willd. sp. pl. 3. p. 1887.

*alpinum.**sylvaticum.**americanum.*

G. spathulatum. *Lam. encycl. 2. p. 728.*

In dry, sunny, and rocky situations: Pennsylvania to Virginia. ♂. July—Sept. v. v. About a span high; flowers small, pale straw-coloured.

uliginosum.

8. G. caule herbaceo ramoso diffuso lanato, foliis linearilanceolatis utrinque angustatis tomentosis, floribus terminalibus congestis. *Willd. sp. pl. 3. p. 1891.*

Icon. *Fl. dan. 859.*

In exsiccated pools: Canada to Virginia. ♂. Aug. Sept. v. v. About a span high; flowers small, calyces brown.

germanicum.

9. G. herbaceum, caule erecto dichotomo, foliis linearilanceolatis acutis tomentosis, floribus globoso-capitatis alaribus terminalibusque. *Willd. sp. pl. 3. p. 1894.*

Filago germanica. Sp. pl. 1311.

Icon. *Fl. dan. 997. Engl. bot. 946.*

In dry fields and pastures: Pennsylvania to Virginia. ♂. July, Aug. v. v. About a span high: calyces straw-coloured.

626. BELLIS. *Gen. pl. 1300.**integerrolia.*

1. B. caulescens, divaricato-ramosa; foliis integerrimis, inferioribus obovatis, supremis lanceolatis, calycinis foliolis acutissimis pilo acuminate.—*Mich. fl. amer. 2. p. 131.*

On banks of rivers and on shady hills in Tennessee. *Michaux. +.*

627. CHRYSANTHEMUM. *Gen. pl. 1307.**Leucanthe-*
mum.

1. C. foliis amplexicaulibus lanceolatis serratis, basi inciso-dentatis, caule erecto ramoso. *Willd. sp. pl. 3. p. 2142.*

Icon. *Fl. dan. 994. Engl. bot. 601.*

In meadows and fields: common every where, and probably introduced from Europe. ♀. June, July. v. v. A very troublesome weed; flowers large, white, resembling a daisy.

ercticum.

2. C. foliis radicalibus tripartitis inciso-dentatis, caulinis cuneiformibus 3-partitis obtusis. *Willd. sp. pl. 3. p. 2146.*

Icon. *Gmel. fl. sib. 2. t. 84.*

On
A
o
62

1. P. f
bi
p
Chr
Icon
f.
In
be
L

1. A. f
bi
ot
p.
Icon
f.
In L
D
ye

2. A. g
ne
tic
bu
In L
D
pr

3. A. p
ba
re
On t
or
ral
pr
4. A. h
go
pe

On the north-west coast. ♀. v. s. in *Herb. Lambert.*

About a foot high; flowers purplish-white, the size of the preceding species.

628. PYRETHRUM. *Smith fl. brit.* 2. p. 900.

1. P. foliis lanceolatis, inferioribus apice serratis, superioribus integerrimis, ramis corymbosis. *Willd. sp. pl. 3. p. 251.*

serotinum.

Chrysanthemum serotinum. Sp. pl. 1251.

Icon. *Pluk. aln.* t. 17. f. 2. *Moris. hist.* 3. s. 6. t. 9. f. 11.

In North America. ♀. Oct. Nov. v. s. in *Herb. Lambert.* Flowers white, smaller than *Chrysanthemum Leucanthemum.*

629. ARNICA. *Gen. pl. 1296.*

1. A. foliis remote dentatis subtus lanato-hirsutis, radicalibus petiolatis oblongis basi angustatis, caulinis alternis oblongo-lanceolatis, caule unifloro.—*Willd. sp. pl. 3. p. 2108.*

Doronicum.

Icon. *Jacq. fl. austr.* 1. t. 92. *Allion. fl. ped.* t. 17. f. 1 & 2.

In Labrador. *Colmaster.* ♀. July, Aug. v. s. in *Herb. Dickson.* Not above a span high; flowers large, yellow,

2. A. glabra; foliis integerrimis utrinque glabris acutis trinervibus, radicalibus spatulato-lanceolatis basi in petiolum angustatis, caulinis oppositis lanceolatis sessilibus, caule unifloro.

plantaginea.

In Labrador. *Colmaster.* ♀. July, Aug. v. s. in *Herb. Dickson.* About a span high; flower the size of the preceding, yellow, the rays broad and 3-dentated.

3. A. pubescens; foliis radicalibus lanceolatis obtusiusculis basi attenuatis petiolatis trinervibus, caulinis oppositis remotis linearibus, caule unifloro.

fulgens.

On the banks of the Missouri. ♀. v. s. About a foot or more high, very slender; leaves on the stem generally two pairs; flowers somewhat smaller than the preceding, of a very deep and beautiful yellow.

4. A. hirsuta; foliis radicalibus decussatim oppositis oblongo-ovatis subdentatis, caule subaphyllo summitate in pedunculos 1-floros diviso.

Claytoni.

maritima.

- Doronicum nudicaule. *Mich. fl. amer.* 2. p. 121.
 Doronicum acaule. *Walt. fl. car.* 205.
 Doronicum foliis Plantaginis, &c. *Clayt. fl. virg.* no. 37.
 In shady woods: Virginia to Florida. ♀. July, Aug.
 v. s. in *Herb. Gronov. nec non Walter.* About two
 feet high; flowers large, of a fine yellow.
 5. A. foliis lanceolatis, inferioribus serratis, caule folioso
 multifloro. *Willd. sp. pl.* 3. p. 2110.
 On the north-west coast. ♀. v. s. in *Herb. Banks.*
 Flowers large, very handsome.

630. CINERARIA. *Gen. pl.* 1294.*integerrifolia.*
β. minor.

1. C. lanato-tomentosa; caule simplici, foliis inferioribus
 spatulatis subdentatis, superioribus angusto-lanceola-
 tis integerrimis erectis, floribus subumbellatis.—*Willd.*
sp. pl. 3. p. 2082.
 Senecio tomentosus. *Mich. fl. amer.* 2. p. 119.
 On the banks of the Missouri. *M. Lewis.* In Carolina
 near Flat-rock. *Michaux.* ♀. Aug. Sept. v. s.
 in *Herb. Lewis.* About a foot high; flowers the size
 of *Senecio Jacobaea*, yellow; the rays very short.

heterophylla.

2. C. lanato-tomentosa; foliis radicalibus longe petiolatis
 spatulato-obovatis, ovatis acutiusculis pinnatifidisque,
 caulinis 2-3-linearibus pinnatifidis, floribus corym-
 bosis.
 On dry sunny rocks, in the Blue-mountains: Pensylva-
 nia. ♀. May, June. v. v. About a span high;
 flowers deep yellow, the size of the preceding.

canadensis.

3. C. villosa; foliis pinnatifidis subvillosis, laciniis sinuatis,
 floribus paniculatis.—*Willd. sp. pl.* 3. p. 2085.
 In Canada. *Kalm.* ♀. +. Resembles *C. maritima*,
 but is not tomentose.

631. SENEPIO. *Gen. pl.* 1290.* *Floribus flosculosiosis: radiis nullis.**vulgaris.*

1. S. foliis amplexicaulibus pinnatifidis dentatis, floribus
 corymboso-coarctatis.—*Willd. sp. pl.* 3. p. 1979.
 Icon. *Fl. dan.* 513. *Engl. bot.* 747.
 In cultivated grounds as a weed; about Philadelphia and
 some other places in Pennsylvania; introduced from
 Europe. ♂. April—Oct. v. v. Flowers, as all the
 following, yellow.

2. S. ca-
 lo-
 ca-
 Icon.
 On r-
 Ca-
 fee-
 in-
 is-
 wc-
 it i-
 ha-
 cle-
 fot-
 rou-
 3. S. gla-
 lur-
 sin-
 On re-
 syl-
 but-
 4. S. gla-
 rot-
 nat-
 bel-
 S. tus-
 In L-
 v. s-
 5. S. cau-
 nib-
 tisq-
 pari-
 About
 flow-
 6. S. foli-
 corc-
 oblo-
 vissi-
 radi-
 S. stric-
 On the
 —A

2. S. caule virgato-paniculato, foliis amplexicaulibus ob-longis acutis inaequilatero acuteque profunde dentatis, calycibus laevibus.—Willd. sp. pl. 3. p. 1974.

Icon. Phuk. phyt. t. 112. f. 1. mala.

On road sides and in newly cleared grounds: Canada to Carolina. ♂. July, Aug. v. v. From two to five feet high. This is one of the plants which spring up in the most remote western countries, when the land is cleared of timber, particularly when the brushwood is burnt on the ground; from which circumstance it is generally known by the name of *Fire-weed*. I have frequently found it covering a square piece of cleared land, when there was not a single plant to be found in any other place for a considerable distance round it.

3. S. glaber; foliis radicalibus spathulatis serratis in petiolum attenuatis, caulinis pinnatifidis dentatis remotissimis, pedunculis elongatis umbellato-corymbosis.

On rocks, near the banks of rivers: about Easton, Pennsylvania. ♀. July, Aug. v. v. Resembling No. 8, but is destitute of a ray.

4. S. glaber; foliis radicalibus longe petiolatis ovato-subrotundis subcordatis dentatis, caulinis 2. remotis pinnatifidis dentatis, pedunculis brevibus subternis umbellatis.

S. tussilaginoides. Walt. fl. car. 208?

In Labrador. Colmaster. In Carolina. Walter. ♀. v. s. in Herb. Dickson. Not above a span high.

elongatus.

pauciflorus.

** *Floribus radiatis.*

5. S. caule simplicissimo rigide erecto subnudo, foliis omnibus lanceolatis, radicalibus subintegris inciso-dentatis, corymbo paucifloro, calycibus glabellis, radiis parvulis.—Mich. fl. amer. 2. p. 120.

About lakes: Canada. Michaux. +. A small plant; flowers the size of *S. Jacobaea*.

6. S. foliis radicalibus longissime petiolatis orbiculatis subcordatis crenatis, caulinis paucis remotissimis linearioribus basi dilatatis inciso-dentatis, pedunculis brevissimis hirsutis subumbellatis, calycibus glabellis, radiis paucis brevissimis.

S. strictus. Herb. Banks. mss.

On the rocky banks of rivers; Pennsylvania. ♀. May—Aug. v. v.; v. s. in Herb. Banks. About a foot

gracilis.

- high, very slender; flowers the smallest of all the species here enumerated.
- olovatus.* 7. *S. foliis radicalibus obovatis crenato-serratis petiolatis, caulinis pinnatifidis dentatis, floribus subumbellatis longe pedunculatis, caule glabriusculo.*—*Willd. sp. pl. 3. p. 1999.*
- S. obtusatus. Herb. Banks. mss.*
On the side of hills and rocks, near rivers: New York to Virginia. $\frac{4}{4}$. June, July. *v. v.; v. s. in Herb. Banks.* Flowers nearly the size of *S. Jacobæa*.
- Balsamitæ.* 8. *S. foliis radicalibus oblongis serratis petiolatis, caulinis inferioribus lyrato-pinnatifidis serratis, summis pinnatifidis dentatis, floribus subumbellatis, caule pedunculatis basi villosa.*—*Willd. sp. pl. 3. p. 1998.*
- S. lyratus. Mich. fl. amer. 2. p. 120.*
S. glabellus. Lam. encycl. 7. p. 102.
In rocky fertile soil: Pennsylvania to Virginia. $\frac{4}{4}$. June, July. *v. v.* Flowers the size of the preceding.
- aureus.* 9. *S. foliis radicalibus ovatis cordatis serratis petiolatis, caulinis pinnatifidis dentatis, lacinia terminali lanceolata, pedunculis subumbellatis incrassatis.*—*Willd. sp. pl. 3. p. 1998.*
In shady woods, about rocks: Canada to Virginia. $\frac{4}{4}$. June, July. *v. v.* Flowers the same size as the preceding.
- Cymularia.* 10. *S. foliis radicalibus petiolatis subrotundis basi truncatis dentatis, petiolis appendiculatis sublyratis, caulinis sessilibus linearibus inciso-dentatis, caule subunifloro.*
On the north-west coast. *D. Nelson. v. s. in Herb. Banks.*
- canadensis.* 11. *S. foliis bipinnatis linearibus glabris, summis pinnatis, corymbis compositis fastigiatis.*—*Willd. sp. pl. 3. p. 1996.*
In Canada. *Kalm. +.*
- ciliatus.* 12. *S. caule piloso, foliis lanceolato-linearibus ciliatis.*—*Walt. fl. car. 208.*
In Carolina. *Walter. +.* From six to eight feet high; rays of the flower white.

1. T. th	co	sp.	Icon
In C	an	Fl	
2. T. th	bu	H	On l
			Be
3. T. th	rot	tua	Icon
In I			He
1. I. fol	cal		Icon
In le			Pe
2. I. vi	nu	tio	du
			Pl
I. gla			Icon
In dr			4
3. I. his	lor	du	ers
			I. sul

632. TUSSILAGO. *Gen. pl. 1289.*

1. T. thyrso fastigiato, floribus radiatis, foliis subrotundo-cordatis inaequaliter dentatis subtus tomentosis. *Willd. sp. pl. 3. p. 1968.*
Icon. *Fl. dan. 61.* *Gmel. fl. sib. 2. t. 70.*
In Canada, and on the highest peaks of the Vermont and New Hampshire mountains. ♀. May. v. v.
Flowers of the ray white, disk pale purple.
2. T. thyrso ovato fastigiato, floribus radiatis, foliis radicalibus oblongis acutis sagittatis integerrimis, lobis obtusis. *Herb. Banks. mss.*
On Hudson's Bay. *Hutchinson.* ♀. v. s. in *Herb. Banks.*
3. T. thyrso fastigiato, floribus obsolete radiatis, foliis subrotundo-cordatis semi-septemlobis inciso-dentatis subtus tomentosis. *Willd. sp. pl. 3. p. 1972.*
Icon. *Ait. kew. 3. t. 11.*
In Labrador. *Colmaster.* ♀. April, May. v. s. in *Herb. Dickson.*

*frigida.**sagittata.**palmata.*633. INULA. *Gen. pl. 1295.*

1. I. foliis amplexicaulibus ovatis rugosis subtus tomentosis, calycum squamis ovatis. *Willd. sp. pl. 3. p. 2089.*
Icon. *Fl. dan. 728.* *Engl. bot. 1546.*
In low meadows and on road sides: New England to Pennsylvania; originally brought from Europe. ♀. July, Aug. v. v. A tall rough-looking plant; flowers large, and, as all the rest, yellow.
2. I. villosa; foliis sessilibus oblongo-lanceolatis basi attenuatis obtusis glanduloso-denticulatis, inferioribus petiolatis serratis, pedunculis axillaribus corymbosis glanduloso-pilosus.—*Willd. sp. pl. 3. p. 2099. excl. syn. Plukenetii.*
I. glandulosa. *Lam. encycl. 3. p. 259.*
Icon. *Mill. dict. ic. t. 57.*
In dry sandy woods and fields: New Jersey to Carolina. ♀. Aug.—Oct. v. v. A foot or more high; flowers the size of *I. Britannica.*
3. I. hispido-pilosa, scaberrima; foliis arcte sessilibus oblongis acutis superne subdentatis utrinque scabris, pedunculis axillaribus corymbosis hispidis.

*Helenium.**mariana.**scabra.**I. subaxillaris. Lam. encycl. 3. p. 259.*

- Icon. *Pluk. mant.* 340, *f. 1. no. 5.*
In the pine-barrens of South Carolina. $\frac{u}{u}$. Aug. Sept.
v. s. in. Herb. Enslen. About two feet high; flowers
smaller than the preceding.
- falcata.* 4. *I. lanato-villosa*; foliis sessilibus linearibus acutissimis
subfalcato-patentibus nervosis utrinque pilosis, pedun-
culis paucis axillaribus corymbosis calycibusque vil-
losis.
Aster gracilens. Herb. Banks. mss.
In sandy pine-woods: New Jersey. $\frac{u}{u}$. Sept. Oct.
v. v. Not above a span high; flowers the size and
shape of No. 2.
- gossypina.* 5. *I. lanuginoso-candicans*; foliis sessilibus oblongo-spathu-
latis obtusis integerrimis, corymbo subfastigiato.—
Mich. fl. amer. 2. *p. 122.*
Inula lanata. Herb. Banks. mss.
On the sea-coast of Carolina and Florida. $\frac{u}{u}$. Aug.—
Oct. *v. s.* About a foot high; flowers the size of
No. 3.
- graminifolia.* 6. *I. argenteo-sericea*; caule simplicissimo, foliis longissi-
me lanceolato-linearibus integerrimis erectis nervosis,
corymbo composito laxo, calycibus turbinatis, squa-
nis acutissimis carina serrato-glandulosi.—*Mich. fl.*
amer. 2. *p. 122.*
In sandy dry woods: Pennsylvania to Florida. $\frac{u}{u}$. Aug.—
Oct. *v. v.* About a foot high; flowers small.
- argentea.* 7. *I. sericea*; foliis lanceolatis trinerviis erectis flexuosis,
corymbo subcomposito stricto, squamis calycinis planis
pubescentibus.—*Pers. syn.* 2. *p. 452.*
In Pennsylvania. *Persoon.* $\frac{u}{u}$. +. Flower a great deal
larger than the preceding species, with which it has a
great affinity.
634. ERIGERON. *Gen. pl.* 1287.
- alpinum.* 1. *E. villosum*; caule plerumque unifloro, calycibus villo-
sis, radio patente, foliis lanceolatis obtusiusculis.—
Willd. sp. pl. 3. *p. 1959.* *Smith fl. brit.* 2. *p. 877.*
Icon. *Engl. bot.* 464. *Fl. dan.* 292.
In Labrador. *Colnaster.* $\frac{u}{u}$. June, July. *v. s. in*
Herb. Dickson. The Labrador specimen is not above
two inches high; flower large, rays purple.
- lepidifolium.* 2. *E. hirsutum, incanum*; foliis radicalibus obovatis sub-
serratis, caulinis paucis distantibus lanceolatis integer.

- rimis, caule sub-3-flore, radiis elongatis.—*Willd. sp. pl. 3. p. 1958.*
- E. serpentaria. *Herb. Banks. mss.*
- E. pulchellum. *Mich. fl. amer. 2. p. 224.*
- E. caule simplicissimo, &c. *Gron. fl. virg. 122.*
- In shady woods, principally throughout the mountains: Canada to Carolina. $\frac{1}{2}$. June—Aug. v. v. A foot or eighteen inches high; flowers large, rays blueish-white; sometimes it produces only one flower, and sometimes more than three. It is known by the name of *Poor Robin's Plantain*.
3. E. glabrum; foliis radicalibus ovali-lanceolatis acutis subdentatis, caule simplicissimo subaphylo elongato, corymbis terminalibus paucifloris, radiis longitudine calycis.—*Mich. fl. amer. 2. p. 224.*
- Aster vernus. Willd. sp. pl. 3. p. 2029. secundum specimen Clayton.*
- In low grounds of Carolina and Georgia. $\frac{1}{2}$. June—Aug. v. v.; v. s. in *Herb. Gronov.* A foot high; flowers small, rays blueish-white.
4. E. tenue pubescens; foliis lanceolatis acutis, inferioribus *quercifolium.* sublyratis grosse deutatis supremis integerrimis, caule subsimplici summitate 3-floro, radiis calyce hemisphaerico duplo longioribus.—*Lam. encycl. 8. p. 491.*
- Icon. *Lam. illustr. t. 681. f. 4.*
- In Carolina and Florida. $\frac{1}{2}$. July, Aug. v. s. in *Herb. Enslen.* Not above a span high; flowers pale blue or white.
5. E. pubescens; foliis cuneato-oblongis rarer inciso-dentatis, caulinis semiamplexicaulibus, caule debili simplici superne corymbose, pedunculis elongatis unifloris, radiis capillaceis calyce hemisphaerico duplo longioribus.—*Willd. sp. pl. 3. p. 1957. Mich. fl. amer. 2. p. 223.*
- In fields and dry woods: Canada to Carolina. $\frac{1}{2}$. Aug. Sept. v. v. A delicate and handsome plant, from one to two feet high; flowers the size of a common daisy, rays of a beautiful blueish-purple.
6. E. pubescens; foliis oblongis dentatis amplexicaulibus, superioribus integerrimis, pedunculis corymbosis in-crassatis, inferioribus elongatis, calycinis squamis carina pilosis, radiis calyce duplo longioribus.—*Willd. sp. pl. 3. p. 1958.*
- In pastures and on fertile hills, near the banks of rivers:
- nudicaule.*
- philadelphi-cum.*
- purpureum.*

SYNGENESIA SUPERFLUA. Erigeron.

Canada to Virginia. ♀. July, Aug. v. v. A foot or eighteen inches high; flowers large, purple, and very ornamental.

strigosum.

7. E. *strigoso-pilosum*; foliis lanceolatis utrinque attenuatis medio grosse paucidentatis seu integerrimis, floribus corymboso-paniculatis.—*Willd. sp. pl. 3. p. 1956.*
In fields and dry pastures; common: Canada to Virginia. ♂. July, Aug. v. v. A very troublesome weed; flowers white, resembling the common daisy.

nervosum.

8. E. *albido-pubescent*; foliis linear-lanceolatis integerimis sericeis nervosis, floribus paniculatis.—*Willd. sp. pl. 3. p. 1953.*
In old fields and woods; common: Canada to Virginia. ♀. July—Sept. v. v. Likewise a troublesome weed; flowers resembling the preceding.

heterophyl-
lum.

9. E. foliis radicalibus subrotundo-ovatis profunde dentatis petiolatis, caulinis lanceolatis acutis medio serratis, cymbo terminali. *Willd. sp. pl. 3. p. 1956.*
Aster annuus. *Willd. sp. pl. 3. p. 2041.*
Icon. *Fl. dan. 486.*

In old waste fields and on road sides; common. ♂.
June—Aug. v. v. Flowers white.

canadense.

10. E. caule hispido paniculatum ramosissimo, foliis lanceolato-linearibus ciliatis, calycibus cylindricis, radiis multiplice serie confertis brevissimis.—*Willd. sp. pl. 3. p. 1954.*
In dry fields and on road sides; common: Canada to Florida. ♂. July—Sept. v. v. A common weed; flowers small.

divaricatum.

11. E. *divaricato-ramosissimum*, subfastigiatopaniculatum; foliis subulatis, floribus brevissime radiatis, corollulis disci 4-fidis.—*Mich. fl. amer. 2. p. 123.*
In fields and meadows of Kentucky, and on the banks of the Missouri and Mississippi. ♂. July—Sept. v. s. Resembles the preceding.

longifolium.

12. E. *glaberrimum*; caule virgatim-paniculato, ramis strictis, foliis longissime-linearibus strictis, calycibus ovatis, radiis vix calyce longioribus.—*Lam. encycl. 8. p. 480.*
In Carolina. ♀. July—Sept. v. s. About two feet high; rays yellow.

hyssopifo-
lum.

13. E. caule ramis sterilibus paniculato: paucis in pedunculos nudos protractis, foliis linearibus glabris ciliatis,

rad
ame
E. car
Icon.
In lov
—S
low
14. E. pilo
trip.
linis
unil
On th
July
high
chat
pale

*
1. S. cau
scab
abbr
Icon.
In hed
sylva
Ban
feet
like
2. S. cau
vibu
erect
sp. p
In swa
ginia
Fron
3. S. cau
tis gl
nicul
sp. p
In wood
Sept.
4. S. caule
scabri

radiis calyce cylindraceo multo longioribus. — *Mich. fl. amer.* 2. p. 123.

E. carolinianum. *Willd. sp. pl.* 3. p. 1953.

Icon. Dill. eth. t. 306. f. 394.

In low grounds: Canada to Carolina; rare. $\frac{1}{2}$. July—Sept. *v. s.* Resembles the preceding; rays yellow.

14. *E. pilosum, subacaule; foliis radicalibus longe petiolatis triplicato-3-partitis, lacinias linearibus divaricatis, caulinis linearibus plerumque indivisis, caule superne nudo unifloro.*

On the banks of the Kooskoosky. *M. Lewis.* $\frac{1}{2}$. July, Aug. *v. s.; v. v. cultum.* Not above a span high; flowers resembling a daisy exceedingly; they change during their flowering, from white to a lively pale red.

635. SOLIDAGO. *Gen. pl.* 1292.

* *Racemis secundis; foliis triplinervibus.*

1. *S. caule villoso, foliis lanceolatis serratis triplinervibus scabris, racemis paniculatis secundis recurvis, ligulis abbreviatis.* *Willd. sp. pl.* 3. p. 2055.

canadensis.

Icon. Pluk. alm. t. 263. f. 1. In hedges, old fields, and along fences: Canada to Pennsylvania. $\frac{1}{2}$. July—Sept. *v. v.; v. s. in Herb. Banks. et Lambert.* From eighteen inches to five feet high; the flowers of all the following species are, like the present, yellow and small.

2. *S. caule villoso erecto, foliis lanceolatis serratis triplineribus scabris subtus villoso, racemis spiciformibus erectis, innuptis nutantibus, ligulis abbreviatis.* *Willd. sp. pl.* 3. p. 2055.

procera.

In swamps, hedges, and about fences: Canada to Virginia. $\frac{1}{2}$. July—Sept. *v. v.; v. s. in Herb. Banks.* From four to seven feet high.

serotina.

3. *S. caule erecto tereti laevi, foliis lineari-lanceolatis serratis glabris margine asperis triplinervibus, racemis paniculatis secundis, pedunculis pubescentibus.* — *Willd. sp. pl.* 3. p. 2056.

In woods and hedges: New England to Pennsylvania. $\frac{1}{2}$. Sept. Oct. *v. v.; v. s. in Herb. Banks.*

gigantea.

4. *S. caule erecto glabro, foliis lanceolatis serratis margine scabris obsolete triplinervibus, racemis paniculatis se-*

- cundis, pedunculis hirtis, ligulis abbreviatis. *Willd.*
sp. pl. 3. p. 2056.
- In open low places, on the side of fertile hills: New England to Virginia. ♀. Aug.—Oct. *v. v.; v. s. in Herb. Banks.* From four to seven feet high.
- ciliaris.* 5. *S. caule erecto glabro, foliis lanceolatis subtriplinervibus glabris margine scabris subserratis, racemis paniculatis secundis, pedunculis glabris, bracteis ciliatis, ligulis abbreviatis. Willd. sp. pl. 3. p. 2056.*
In Pennsylvania. *Muhlenberg.* ♀. +.
- reflexa.* 6. *S. caule erecto villosa, foliis lanceolatis subserratis triplinervibus scabris reflexis, ramis paniculatis subsecundis reflexis. Willd. sp. pl. 3. p. 2056.*
In pine-woods and old fields: New Jersey to Carolina. ♀. Aug. Sept. *v. v.; v. s. in Herb. Banks.*
- lateriflora.* 7. *S. caule erecto pilosiusculo, foliis lanceolatis subtriplinervibus glabris margine scabris, inferioribus subserratis, racemis paniculatis subrecurvis secundis. Willd.*
sp. pl. 3. p. 2057.
In old fields and dry woods: Canada to Carolina. ♀. Aug.—Oct. *v. v.; v. s. in Herb. Banks.* Not more than half the size of No. 1; flowers larger than any of the preceding.
- ** *Racemis secundis; foliis venosis.*
- aspera.* 8. *S. caule erecto tereti piloso, foliis ovatis subellipticis scaberrimis rugosis serratis enervibus, racemis paniculatis secundis. Willd. sp. pl. 3. p. 2057.*
Icon. *Dill. elth. t. 305. f. 392.*
In barren dry fields and woods: New York to Carolina. ♀. Sept.—Nov. *v. v. v. s. in Herb. Banks.*
- altissima.* 9. *S. caule erecto hirto, foliis lanceolatis inferioribus profunde serratis scaberrimis rugosis, paniculis secundis. Willd. sp. pl. 3. p. 2057.*
Icon. *Mart. cent. 14.*
- vulgaris.* α. *S. caule 5-pedali piloso, serraturis profundis inæqualibus, racemis divaricatis. Ait. kew. 3. p. 212.*
- recurvata.* β. *S. caule 3-pedali villosa, serraturis profundis subæqualibus. Ait. l. c.*
- virginiana.* γ. *S. caule 5-pedali villosissimo, serraturis magnis, racemis vix divergentibus. Ait. l. c. Mill. dict.*
In hedges and fields; common. ♀. Aug.—Oct. *v. v.; v. s. in Herb. Banks. et Lambert.* It is a very variable species, and scarcely two individuals look alike.

10. *S. caule infuscata, eu dicata. S. altissima. Icon. In di-Occ.*
11. *S. caule rati-
Ba-
S. altissima. S. pilosa. In field
v. s. incl.*
12. *S. caule nua-
me-
pl. In old
♀.*
13. *S. caule plex-
sole-
mis-
In pu-
v. s-
leav-
whi-*
14. *S. caule pidi-
rat-
p. 2-
In san-
lina. The
or sc-*
15. *S. caule di-
cunc-
sp. f. In shad-
♀.*

SYNGENESIA SUPERFLUA. Solidago.

537

rugosa.

10. *S. caule erecto birto, foliis lanceolatis scaberrimis rugosis inferioribus adpresso-serratis, racemis paniculatis secundis patentissimis.* Willd. sp. pl. 3. p. 2058. Mill. dict.

S. altissima, e. Ait. kew. 3. p. 213.

Icon. Dill. elth. t. 308. f. 396.

In dry barren soil: Canada to Virginia. ♀. Aug.—Oct. v. v.; v. s. in Herb. Banks. et Lambert.

11. *S. caule erecto villosa, foliis lanceolatis molliusculis serratis enervibus, racemis paniculatis secundis.* —Herb. Banks. mss.

*villosa.**S. altissima, β.* Ait. kew. 3. p. 213.*S. pilosa.* Mill. dict.

In fields and woods; frequent. ♀. Aug.—Oct. v. v.; v. s. in Herb. Banks. et Lambert. From eighteen inches to three feet high.

12. *S. caule erecto hirto sulcato, foliis oblongis utrinque attenuatis acuminatis, supra glabris, subtus rugosis scabris, medio adpresso-serratis, racemis secundis.* —Willd. sp. pl. 3. p. 2059.

scabra.

In old fields and about fences: Pennsylvania to Virginia. ♀. Aug.—Oct. v. v.; v. s. in Herb. Lambert.

13. *S. caule erecto tereti birto, foliis oblongis acutis subamplexicauli-sessilibus glabris margine scabris rarer obsolete dentatis, panicula nuda secunda pyramidata, ramis reflexis, pedunculis glabris.*

pyramidalis.

In pine-barrens of Georgia. Enslen. ♀. Aug. Sept. v. s. in Herb. Enslen. About two feet high; the leaves decrease toward the beginning of the panicle, which consists of lively yellow and very small flowers.

14. *S. caule erecto tomentoso, foliis caulinis lanceolatis hispidis integerrimis, radicalibus subcuneiformibus serratis, racemis paniculatis secundis.* Willd. sp. pl. 3. p. 2059.

nemoralis.

In sandy and barren fields; common: Canada to Carolina. ♀. Aug.—Oct. v. v.; v. s. in Herb. Banks. The whole plant has a gray aspect, and is about a foot, or sometimes two, high.

15. *S. caule erecto glabro, foliis ellipticis serratis glabris, radicalibus oblongo-spathulatis, racemis paniculatis secundis patentibus, pedunculis pubescentibus.* Willd. sp. pl. 3. p. 2059.

patula.

In shady woods, in fertile soil: New York to Virginia. ♀. Sept. Oct. v. v. Two feet high.

ulmifolia.

16. S. caule erecto glabro striato, foliis ellipticis profunde serratis acuminatis subtus villosis, radicalibus obovatis, racemis paniculatis secundis, pedunculis villosis, ligulis abbreviatis.—*Willd. sp. pl. 3. p. 2060.*

In swamps and shady woods: New York to Virginia.
4. Aug.—Oct. v. v.

arguta.

17. S. caule erecto glabro, foliis glabris argute inaequaliter serratis, caulinis ellipticis, radicalibus ovato-oblongis, racemis paniculatis secundis, ligulis elongatis. *Willd. sp. pl. 3. p. 2060.*

In woods and meadows: Canada to Virginia. 4. Sept.—Nov. v. v.; v. s. in *Herb. Banks.*

juncea.

18. S. caule erecto glabro, foliis lanceolatis glabris margine scabris, inferioribus serratis, racemis paniculatis secundis. *Willd. sp. pl. 3. p. 2060.*

In sandy fields and woods: New Jersey to Carolina.
4. Aug.—Oct. v. v.; v. s. in *Herb. Banks.*

elliptica.

19. S. caule erecto glabro, foliis ellipticis laevis serratis, racemis paniculatis secundis, ligulis mediocribus. *Willd. sp. pl. 3. p. 2060.*

S. latissimifolia. Mill. dict.

In shady woods, among rocks: New York to Virginia.
4. Aug.—Oct. v. v.; v. s. in *Herb. Banks.*

asperata.

20. S. caule paniculato-corymboso, racemis suberectis, floribus adscendentibus, foliis lanceolatis serratis scabris. *Herb. Banks. mss.*

In Canada. 4. v. s. in *Herb. Banks.*

recurvata.

21. S. caule erecto pubescente, foliis lanceolatis serratis margine scabris, racemis elongatis secundis recurvatis paniculatis. *Willd. enum. 889.*

In shady woods: Pennsylvania and Virginia. 4. Sept.—Nov. v. v.

semperfiriens.

22. S. caule erecto glabro, foliis linear-lanceolatis subcarnosis laevis integerrimis margine scabris, racemis paniculatis secundis, pedunculis pilosis. *Willd. sp. pl. 3. p. 2060.*

Icon. Pluk. alm. t. 235. f. 5. Moris. hist. 3. s. 7. t. 23. f. 15.

In swamps and on the banks of rivers: Canada to Pennsylvania. 4. Sept.—Nov. v. v.; v. s. in *Herb. Banks. et Lambert.*

virgata.

23. S. caule glabro simplicissimo, foliis subcuneato-lanceolatis obtusis integerrimis glaberrimis appressis, supe-

ri
ap
fl.
In s
4
tre

24. S. ca
ge
sec

Icon
In dr
to
mc
Ba
am
use
full
abl
an
bigl

25. S. ca
flex
mai
fl. a
In ope
Sep

26. S. glat
pani
cibu
ame
In the
—O

27. S. caul
rami
sis,
Icon. i
In woo
Aug.
distir
racen

28. S. caule
tiolat
pl. 3.

- profunde
obovatis,
osis, ligu-
- Virginia.
- æqualiter
oblongis,
Willd.
- ¶. Sept.
- margine
ulatis se-
- Carolina.
- serratis,
liocribus.
- Virginia.
- tis, flori-
scabris.
- atis mar-
vatis pa-
- l. Sept.
- ibcarno-
emis pa-
p. pl. 3.
7. t. 23.
- to Pen-
Herb.
- lanceo-
, supe-
- rioribus sensim minoribus, ramis paniculæ elongatis apice racemifloris, pedunculis glabris secundis.—*Mich. fl. amer. 2. p. 117.*
In shady wet woods of Lower Carolina and Georgia.
¶. Aug.—Oct. v. v. About two foot high; extremely smooth and slender.
24. S. caule erecto pubescente, foliis linear-lanceolatis integrerrimis glabris margine scabris, racemis paniculatis secundis. *Willd. sp. pl. 3. p. 2061.*
Icon. *Pluk. alm. 116. f. 6.*
In dry sunny situations, on fertile woodlands: Canada to Carolina; principally throughout the Allegany mountains. ¶. Aug. Oct.—v. v.; v. s. in *Herb. Banks*. This is most generally and principally known among the inhabitants by the name of *Golden-rod*, and used as a wholesome tea. The flowers, gathered when fully expanded, and carefully dried, give a most agreeable substitute for tea, which for some time has been an article of exportation to China, where it fetches a high price.
25. S. caule erecto tereti aspero, foliis arcte sessilibus reflexis linear-lanceolatis mucronatis subtus subcarinatis margine asperis, panicula racemis recurvatis.—*Mich. fl. amer. 2. p. 117.*
In open swamps of Virginia and Carolina. ¶. Aug. Sept. v. v.
- *** *Racemis erectis.*
26. S. glabra, suffruticosa; foliis lanceolatis obtusis enervibus, paucifloscu-
panicula composita multiflora, fasciculis erectis, caly-
cibus angusto-oblongis 5-floris, radio unico.—*Mich. fl.
amer. 2. p. 116.*
In the sand-barrens of Virginia and Carolina. ¶. Aug.—Oct. v. v.
27. S. caule foliisque ellipticis pilosis, inferioribus serratis, ramis foliolosis, racemis erectis, foliolis calycinis obtusis. *Willd. sp. pl. 3. p. 2061.*
Icon. *Pluk. alm. t. 114. f. 3.*
In woods and on dry hills: Canada to Carolina. ¶. Aug.—Oct. v. v.; v. s. in *Herb. Banks*. It is easily distinguished from the rest, by the short and close racemes and white rays.
28. S. caule erecto villoso, foliis ellipticis scabriusculis petiolatis, racemis erectis, ligulis elongatis. *Willd. sp.
pl. 3. p. 2062.*
- odora.*
- retrorsa.*
- bicolor.*
- petiolaris.*

In dry sandy soil : New Jersey to Carolina. ♀. Sept.
—Nov. v. v.; v. s. in *Herb. Banks.*

stricta.

29. S. caule erecto glabro, foliis caulinis lanceolatis integerimis glabris margine scabris, radicalibus serratis, racemis paniculatis erectis, pedunculis glabris. *Willd.* sp. pl. 3. p. 2062.

In sandy woods : New Jersey to Carolina. ♀. Aug.
—Nov. v. v.; v. s. in *Herb. Banks. et Lambert.*
About two feet high, very smooth.

lanceolata.

30. S. caule hirto angulato ramosissimo, foliis lanceolato-linearibus integerrimis erectiusculis 3—5-nervibus scabriusculis : nervis subtus pilosis, axillis nudis, corymbis terminalibus fastigiatis, ramulis capitatis, ligulis altitudine disci.—*Willd. sp. pl. 3. p. 2062.*

Chrysocoma graminifolia. Sp. pl. 1178.

In fields and meadows, on the side of woods and rivers:
Canada to Pennsylvania. ♀. Sept.—Nov. v. v.; v. s.
in *Herb. Banks. et Lambert.* From four to five feet
high ; flowers small.

tenuifolia.

31. S. caule scabro angulato corymboso-ramoso, foliis angustissime-linearibus patulis obsolete 3-nervibus scabris, axillis foliosis, corymbis terminalibus fastigiatis, ramulis capitatis, ligulis disco vix altioribus.

S. *lanceolata* β. *minor. Mich. fl. amer. 2. p. 116.*

In pine-barrens : New Jersey to Carolina. ♀. Sept.
Oct. v. v. Not above a foot high ; leaves very small
and narrow.

Sarothræ.

32. S. caule angulato scabro inferne nudo, superne corymboso, ramis summitate paucifloris, foliis linearibus stricte-appressis obsolete 3-nervibus utrinque scabris, axillis nudis, ligulis disco duplo longioribus.

On the plains of the Missouri. *M. Lewis.* ♀. Sept.
v. s in *Herb. Lewis.* About a span high ; resembling
Hypericum Sarothra in general habit very much. This
and the two preceding have a great affinity to one another ; but I consider them good and permanent species,
as there are no intermediate varieties, which might lead to the suspicion that all belong to one genuine species.

væsia.

33. S. caule lœvi erecto, foliis lanceolatis glabris, racemis erectis, ligulis mediocribus. *Willd. sp. pl. 3.* p. 2062.

Icon. *Dill. elth. t. 307. f. 395.*

In w.
Au

34. S. caule
bris
elat

In sa
Sep

35. S. caule
inten
lis

In gra
Oc

36. S. caule
bris
ere

In sa
No

37. S. caule
scal
ered

In sa
Au

38. S. caule
mis
dum
pl.

In salt
v. v
rese
bly

39. S. caule
tege
tis,
Wi

Icon.
On th

July
Tal

40. S. caule
mer
infir
Wi

S. inte
On the

♀. Sept.

is integer-
erratis, ra-
s. Willd.

♀. Aug.
Lambert.

lanceolato-
eribus sca-
lis, corym-
, ligulis al-

and rivers:
v. v.; v. s.
o five feet

, foliis an-
rvibus sca-
fastigiatis,

16.
♀. Sept.
very small

cne corym-
linearibus
ue scabris,
i.

♀. Sept.
resembling
much. This
to one an-
cient spe-
cies, which
to one ge-

is, racemis
sp. pl. 3.

In woods and low fields : New England to Carolina. ♀.
Aug.—Oct. v. v.

34. S. caule glabro paniculato, foliis lanceolatis serratis gla-
bris margine scabris, ramis apice racemosis, ligulis
elongatis. Willd. enum. 891.

In sandy fields and woods : New York to Virginia. ♀.
Sept. Oct. v. v.

35. S. caule erecto hispido scabro, foliis lanceolatis scabris
integerimis, radicalibus serratis, racemis erectis, ligu-
lis mediocribus. Willd. sp. pl. 3. p. 2063.

In grassy fields and woods : New Jersey to Virginia. ♀.
Oct. Nov. v. v. Resembles No. 33.

36. S. caule paniculato hirsuto, foliis lanceolatis utrinque sca-
bris, caulinis serratis, rameis integerimis, racemis
erectis, ligulis elongatis. Willd. enum. 891.

In North America. Willdenow. ♀. +.

37. S. caule ramoso pubescente, foliis lanceolatis utrinque
scabris attenuatis 3-nervibus integerimis, racemis
erectis, ligulis elongatis. Willd. enum. 891.

In sandy barren woods : New Jersey to Carolina. ♀.
Aug.—Oct. v. v.

38. S. caule erecto laevi, foliis lanceolatis carnosis integeri-
mis undique laevibus, racemis paniculatis erectis, pe-
dunculis squamosis villosis, ligulis elongatis. Willd. sp.
pl. 3. p. 2063.

In salt-marshes : Canada to Virginia. ♀. Sept.—Nov.
v. v.; v. s. in Herb. Banks. A tall robust species,
resembling the following one very much, and proba-
bly nothing more than a variety.

39. S. caule obliquo glabro, foliis lanceolatis subcarnosis in-
tegerimis undique laevibus, racemis paniculatis erec-
tis, pedunculis squamosis glabris, ligulis elongatis.
Willd. sp. pl. 3. p. 2063.

Icon. Dodart. act. 4. t. 219.

On the banks of rivers and ditches, near salt-water. ♀.
July—Sept. v. v.; v. s. in Herb. Banks. et Lambert.
Tall; flowers large.

40. S. caule erecto subpubescente, foliis linear-lanceolatis
membranaceis basi attenuatis glabris margine scabris,
infimis subserratis, racemis erectis, ligulis elongatis.
Willd. sp. pl. 3. p. 2064.

S. integerima. Mill. dict.

On the banks of rivers and ditches : Canada to Virginia.

livida.

hispida.

hirta.

lithospermi-
folia.

laevigata.

mexicana.

viminea.

- U.* Aug.—Oct. *v. v.*; *v. s.* in *Herb. Banks. et Lambert.* Resembles the preceding, but the leaves are membranaceous, &c.
- erecta.* 41. *S. caule subviloso, foliis lanceolatis venosis glabris integrimis subpetiolatis.* *Herb. Banks. mss.*
In North America. *U.* *v. s.* in *Herb. Banks.*
- macrophylla.* 42. *S. foliis inferioribus ovatis acuminatis attenuatis inaequilater argute serratis glabris, caulinis lanceolatis utrinque attenuatis subsessilibus serratis, racemis axillariibus pedunculatis foliosis, longitudine foliorum, calycibus oblongis turgidis multifloris, ligulis subelongatis.* — *Herb. Banks. mss.*
In Canada. *U.* *v. s.* in *Herb. Banks.* About three feet high; it is intermediate between this genus and *Aster.*
- glomerata.* 43. *S. caule humili simplicissimo, foliis glabris oblongo-lanceolatis serratis, racemo simplici e glomerulis axillariibus: superioribus capitato-congestis, calycibus turgidis multifloris.* — *Mich. fl. amer. 2. p. 117.*
In the mountains of Carolina. *Michaux.* *U. +.*
- flexicaulis.* 44. *S. caule flexuoso glabro angulato, foliis ovatis acuminatis serratis glabris, racemis erectis axillaribus, ligulis mediocribus.* — *Willd sp. pl. 3. p. 2064.*
Icon. *Pluk. alm. t. 235. f. 3.* *Rob. ic. 22.*
- latifolia.* $\beta.$ *S. Pluk. alm. t. 235. f. 4.*
In woods and fields; common: Canada to Carolina. *U.* Aug.—Oct. *v. v.*; *v. s.* in *Herb. Banks.* This is the most common species.
- axillaris.* 45. *S. caule glabro tereti stricto, foliis lanceolatis serratis glabris, racemis axillaribus subglobosis erectis, ligulis elongatis.* — *Herb. Banks. mss.*
In shady woods: Canada to Virginia. *U.* Aug.—Oct. *v. v.*; *v. s.* in *Herb. Banks.*
- Virgaurea.* 46. *S. caule erecto tereti superne ramoso et pubescente, foliis caulinis lanceolatis utrinque attenuatis serratis, inferioribus ellipticis pilosiusculis, racemis erectis, ligulis elongatis.* — *Willd sp. pl. 3. p. 2065.*
Icon. *Engl. bot. 301.*
In labor. *Colmaster.* *U.* Aug.—Oct. *v. s.* in *Herb. Dickson.*
- multiradiata.* 47. *S. caule villosiusculo, foliis sessilibus lanceolatis glabris ciliatis, inferioribus apice serratis, racemo terminali*

erec
p. 2
In Lal
in F
48. *S. cau
glab
Ban
In No
49. *S. cau
culi
mss.
In No
50. *S. cau
integ
latis
pl. 3
Icon.
In the
Aug
Flow
51. *S. foli
diuse
In sand
roug
1. *A. folii
marg
tis, i
brevi
In sand
Aug.
From
tene
disk
2. *A. folii
mis
giatis
lycib
obtus
p. 20
Conyza******

- erecto, ligulis elongatis numerosis. *Willd. sp. pl. 3. p. 2066.*
In Labrador and Hudson's Bay. \mathcal{U} . Aug.—Oct. *v. s. in Herb. Dickson. et Banks.*
48. S. caule simplici erecto glabro, foliis lanceolatis serratis glabris basi attenuatis elongatis, racemo erecto. *Herb. Banks mss.* *humilis.*
- In North America. \mathcal{U} . *v. s. in Herb. Banks.*
49. S. caule piloso tereti, foliis lanceolatis subtus pilosiusculis, racemis erectis, ligulis elongatis. *Herb. Banks. mss.* *elata.*
- In North America. \mathcal{U} . *v. s. in Herb. Banks.*
50. S. caule foliisque ovato-oblongis pilosis scabris, caulinis integerrimis, infimis serratis, ramis floriferis paniculatis, racemis compactis, ligulis elongatis. *Willd. sp. pl. 3. p. 2067.* *rigida.*
Icon. *H-rm. parad. 243.*
In the tract of mountains: New Jersey to Carolina. \mathcal{U} . Aug.—Oct. *v. v.; v. s. in Herb. Banks. et Lambert.*
Flowers large; leaves very rough.
51. S. foliis radicalibus ovato-oblongis petiolatis, caule nudusculo ramoso fastigiat. *Willd. sp. pl. 3. p. 2067. sis.*
In sandy fields and woods. \mathcal{U} . Sept. Oct. *v. v.* Leaves rough; flowers large.
636. ASTER. *Gen. pl. 1291.*
* *Foliis integerrimis.*
1. A. foliis lineari-lanceolatis 3-nervibus punctatis acutis *hyssopifolius.*
margini scabris, ramulis corymboso-fastigiatis coarctatis, radio sub-5-floro, calycibus imbricatis disco duplo brevioribus. *Willd. sp. pl. 3. p. 2022.*
In sandy fields and woods: New Jersey to Carolina. \mathcal{U} . Aug.—Oct. *v. v.; v. s. in Herb. Banks. et Lambert.*
From a foot to two high; flowers in a close and flattened corymb; ray white, with a tinge of purple; disk yellow.
2. A. foliis lineari-lanceolatis obsolete trinervibus integerrimis obtusis margini scabris, ramulis corymboso-fastigiatis, floribus sessilibus aggregatis, radio 5-floro, calycibus disco brevioribus imbricatis, squamis oblongis obtusis adpressis apice subreflexis.—*Willd. sp. pl. 3. p. 2024.* *solidaginoides.*
Conyza linifolia. Sp. pl. 1205.

Icon. *Pluk. alm. t. 79. f. 2.*

In dry swamps and on the borders of woods : Pennsylvania to Carolina. 24. Aug.—Oct. v. v.; v. s. in *Herb. Banks. et Lambert.* About two feet high; the scales of the calyx white, with green tips; rays long, white.

tordifolius.

3. A. foliis cuneato-obovatis acutis enervibus scabris tortuos-patulis, corymbo composito subfastigato, calycibus cylindricis imbricatis basi bibracteatis, squamis oblongis acutiusculis adpressis, radio 5-floro.—*Mich. fl. amer. 2. p. 109.*

Conyzza bifoliata. Walt. fl. car. 204.

In Lower Carolina and Georgia. 24. Aug.—Oct. v. s. in *Herb. Enslen.* About a foot high; flowers the size of the preceding, rays white.

ledifolius.

4. A. foliis linear-lanceolatis basi attenuatis enervibus scabriusculis margine revolutis, ramis corymbosis, ramulis filiformibus unifloris nudiusculis, calycibus laxis imbricatis disco duplo brevioribus, foliolis acutissimis, radio sub-20-floro.

A. nemoralis. *Ait. kew. 3. p. 198.*

β. A. caule simplicissimo unifloro.

A. uniflorus. *Mich. fl. amer. 2. p. 110?*

In sphagnum bogs and about mountain-lakes : Canada to New Jersey. 24. Sept. Oct. v. v.; v. s. in *Herb. Banks. et Lambert.* About a foot high; flowers middle-sized; disk yellowish white; rays lilac or reddish-purple, twice the length of the disk. Sometimes the leaves have a small indenture on each side.

rigidus.

5. A. foliis linearibus mucronatis subcarinatis rigidis margine scabro-ciliatis, caulinis reflexis, rameis patentibus subulatis, caule erecto superne subramoso, ramulis 1-floris corymbosis, calycibus imbricatis disco duplo brevioribus, squamis carinatis obtusiusculis, radiis sub 10-floris reflexis.—*Willd. sp. pl. 3. p. 2032. Mich. fl. amer. 2. p. 118.*

A. nemoralis. *Willd. sp. pl. 3. p. 2021. excl. syn. Aiton.*
A. linariifolius. *Herb. Banks. et Lambert.*

Icon. *Pluk. alm. t. 14. f. 7.*

In sandy fields and pine-barrens: New Jersey to Carolina. 24. Aug.—Oct. v. v.; v. s. in *Herb. Banks. et Lambert.* About a foot or less high; flowers nearly the size of the preceding, rays a beautiful pale violet-blue, disk yellow.

6. A. foliis tatis c
subde imbri
Mich

A. squar

In dry,

sylvat

in *H*

more

rough

crowd

frne

naked

7. A. foliis

erecti

lycibi

mss.

At Hud

Flow

8. A. foliis

tentib

bus i

Willia

In dry s

to Vi

teen

large

9. A. foliis

villoso

uniflo

lance

A. villos

In natu

—Oc

10. A. glab

multi

ceis,

In salt-

Nov.

11. A. foliis

caule

calyci

sis.

- Pensylv.
v. s. in
high; the
rays long,
rays long,
- que sca-
fastigiat-
tis, squa-
5-floro.—
- Oct. v. s.
's the size
- vibus sca-
is, ramu-
bus laxis
utissimis,
- : Canada
in Herb.
vers mid-
reddish-
times the
- dis mar-
pa'entii-
, ramu-
isco du-
is, radiis
2. Mich.
- n. Aiton.
- to Caro-
. Banks.
's nearly
e violet-
6. A. foliis crebris linearibus mucronatis enervibus impunc-
tatis carinatis scabris rigidis, rameis recurvatis, caule
subdecumbente, ramis fastigiatis 1-floris, calycibus
imbricatis longitudine disci.—Willd. sp. pl. 3. p. 2024.
Mich. fl. amer. 2. p. 110.
A. squarrosus. *Herb. Banks. mss.*
In dry, sandy and gravelly fields, and among rocks: Pen-
sylvania to Carolina. ♀. Aug.—Nov. v. v.; v. s.
in *Herb. Banks.* Resembles the preceding, but is
more branching and of lower growth; the stems are
rough, and generally of a purple colour; leaves more
crowded, smaller, and more reflex; flowers of the
same colour, but smaller, and not raised on almost
naked peduncles.
7. A. foliis angusto-linearibus enervibus impunctatis glabris *graminifolius.*
erectis, ramulis terminalibus nudiusculis 1-floris, ca-
lycibus laxis discum subæquantibus.—*Herb. Banks.*
mss.
At Hudson's Bay. ♀. Aug. Sept. v. s. in *Herb. Banks.*
Flowers pale-purple; not above six inches high.
8. A. foliis linearibus enervibus punctatis scabris reflexo-pa-
tentibus, ramis corymboso-fastigiatis foliosis, calyci-
bus imbricatis brevibus, radiis disco subæqualibus.—
Willd. sp. pl. 3. p. 2024
In dry shady woods and exsiccated swamps: Pennsylvania
to Virginia. ♀. Sept.—Nov. v. v. About eighteen
inches or two feet high; flowers the size of a
large daisy; rays white or faintly purple.
9. A. foliis linear-lanceolatis albo-villosis, caule ramoso
villoso, ramis recurvato-patulis, ramulis subsecundis
unifloris, calycibus oblongis laxis imbricatis, squamis
lanceolatis mucronatis —*Willd. sp. pl. 3. p. 2025.*
A. villosus. *Mich. fl. amer.* 2. p. 113.
In natural meadows of the Illinois country. ♀. Aug.
—Oct. +.
10. A. glaberrimus, parviflorus; caule paniculato, ramis
multifloris, foliis linear-subulatis, calycibus cylindra-
ceis, ligulis radii minutis. *Mich. fl. amer.* 2. p. 111.
In salt-marshes: Pennsylvania to Carolina. ♀. Aug.—
Nov. +.
11. A. foliis linear-lanceolatis utrique attenuatis acuminatis,
caule pubescente paniculato erecto, ramis paucifloris,
calycibus imbricatis, squamis linearibus acutis adpres-
sis. *Willd. sp. pl. 3. p. 2025.*

*linariifolius.**linifolius.**pilosus.**subulatus.**foliolosus.*

Icon. *Dill. elth. t. 35. f. 39?*

In dry barren fields and on road sides: Canada to Carolina. ♀. Aug.—Oct. *v. v.; v. s. in Herb. Lambert.* Flowers small, white; scarcely a distinct species from the following. There is no specimen of it in the Banksian Herbarium.

tenuifolius.

12. A. foliis linear-lanceolatis utrinque attenuatis margine hispidis, caule glabro ramoso erecto, ramulis unifloris, calycibus imbricatis, squamis oblongis acutis laxis. *Willd. sp. pl. 3. p. 2026.*

Icon. *Pluk. alm. 78. f. 5.*

On road sides and barren fields. Aug.—Nov. *v. v.; v. s. in Herb. Banks. et Lambert.* Flowers the same as the preceding, which I consider only a variety of the present species.

dumosus.

13. A. foliis linearibus glabris, ramulorum brevissimis, ramis paniculatis, calycibus cylindraceis arcte imbricatis. *Willd. sp. pl. 3. p. 2026.*

violaceus.

α. A. radio pallide violaceo, caule pubescente. *Ait. kew. 3. p. 202.*

albus.

β. A. radio albo, caule glabro. *Ait. l. c.*

Icon. *Herm. parad. 95. Pluk. alm. t. 78. f. 6.*

In dry barren fields and sand-barrens: New England to Carolina: ♀. Aug.—Nov. *v. v.; v. s. in Herb. Banks. et Lambert.* Flowers small, rays white, disk yellow.

ericoides.

14. A. foliis linearibus glaberrimis, ramulorum subulatis approximatis, caulinis elongatis, calycibus subsquarrosis, foliolis acutis, caule glabro. *Willd. sp. pl. 3. p. 2027.*

In barren grounds and on road sides; common: Canada to Carolina. ♀. Aug.—Nov. *v. v.; v. s. in Herb. Banks. et Lambert.* Flowers the size and colour of the preceding.

multiflorus.

15. A. foliis linearibus glabriusculis, caule ramosissimo difuso pubescente, ramulis secundis, calycibus imbricatis, squamis oblongis squarrosis acutis. *Willd. sp. pl. 3. p. 2027.*

Icon. *Dill. elth. t. 36. f. 40.*

In dry fields, and on road sides: Canada to Carolina. ♀. Aug.—Nov. *v. v.; v. s. in Herb. Banks. et Lambert.* Probably only a variety of the preceding.

ciliatus.

16. A. foliis ciliatis, caulinis linear-lanceolatis nervosis, ramis brevissimis lanceolatis 3-nervibus, caule ramoso

pube

squa

p. 24

In Nor
the I

17. A. cana
bosa
mis

On the
Flow

18. A. folii
berri
cibus
pl. 3.

In mars
—No
large,

19. A. glat
subre
patuli
ricatis
sis acu
In salt-r
Nov.
smalle
yellow

20. A. humi
bus, t
spatu
lycis s
amer.

In wood
Micha

21. A. foliis
hispidi
floris,
tis obt
mer. 2.

In dry ba
v. s. R
blue.

22. A. foliis
hispidis

ada to Caro-
rb. Lambert.
species from
of it in the

itis margin-
lis unifloris,
cutis laxis.

v. v. ;
is the same
a variety of

ssimilis, ra-
e imbrica-
fit. kew. 3.

6.
England to
in Herb.
hite, disk

subulatis
ibsquarro-
sp. pl. 3.

: Canada
in Herb.
colour of

simo dif-
imbrica-
Willd. sp.

Carolina.
anks. et
ceding.

rosis, ra-
: ramoso

pubescente, ramis paniculatis, calycibus imbricatis,
squamis spathulatis squarrosis. Willd. sp. pl. 3.
p. 2027.

In North America. Muhlenberg. ♀. +. Resembles
the preceding one.

17. A. cano-pubescent; foliis linearibus, panicula corym-
bosa ramosissima foliosa, calycibus imbricatis acutissi-
mis disco longioribus.

On the banks of the Missouri. ♀. Aug.—Nov. v. s.
Flowers the size of a daisy, rays pale purple.

18. A. foliis remotis linearibus amplexicaulis erectis gla-
berrimis margine scabris, pedunculis fere nudis, caly-
cibus squarrosis basi foliis 2. suffultis.—Willd. sp.
pl. 3. p. 2033.

In marshes and swamps of Lower Carolina. ♀. Aug.
—Nov. v. s. in Herb. Banks. et Lambert. Flowers
large, rays blue, disk yellow.

19. A. glaberrimus; foliis subulato-linearibus subcarnosis
subreflexis, caule tenui ramosissimo, ramis ramulisque
patulis setaceis 1-floris, squamis peduncularum divar-
icatis subulatis, calycibus imbricatis, squamis adpres-
sis acutis.—Mich. fl. amer. 2. p. 112.

In salt-marshes: New York to Carolina. ♀. Sept.—
Nov. v. v. Resembles the preceding one; flowers
smaller; rays pale purple or white, disk brownish-
yellow.

20. A. humilis; radice repente, caulibus debilibus simplici-
bus, foliis longe lanceolatis glabriusculis, radicalibus
spathulatis, corymbo terminali incompte rarifloro, ca-
lycis squamis lineari-oblongis obtusiusculis. Mich. fl.
amer. 2. p. 113.

In woods of North Carolina, in the county of Burke.
Michaux. +.

21. A. foliis creberrimis linearibus obtusis reflexis margine
hispidis, caule ramoso diffuso glabro, ramulis uni-
floris, calycibus imbricatis, squamis lineari-spathula-
tis obtusis.—Willd. sp. pl. 3. p. 2028. Mich. fl. a-
mer. 2. p. 112.

In dry barren soil in Lower Carolina. ♀. Aug.—Nov.
v. s. Resembles the following species; flowers bright
blue.

22. A. foliis creberrimis ovatis acuminatis reflexis margine
hispidis, caule ramoso hirto, ramulis unifloris, caly-

canescens.

paludosus.

sparsiflorus.

surculosus.

coridifolius.

squarrosus.

cibus imbricatis, squamis linearis-cuneatis acutis squarrosis. Willd. sp. pl. 3. p. 2028. Walt. fl. car. 209.

In pine-barrens of Lower Carolina. ♀. Aug.—Nov. v. v.; v. s. in Herb. Lambert. Has a striking resemblance to *A. reflexus* from the Cape of Good Hope, and is a very ornamental plant; flowers blue and large.

concolor.

23. *A. foliis oblongo-lanceolatis utrinque cano-pubescentibus, caule simplicissimo erecto pubescente, racemo terminali, calycibus imbricatis, squamis lanceolatis sericeis adpressis.* Willd. sp. pl. 3. p. 2029.

In sandy pine and oak woods near the sea-coast: New Jersey to Florida. ♀. Aug.—Nov. v. v.; v. s. in Herb. Banks. et Lambert. About a foot high; flowers, both ray and disk, of an agreeable reddish-purple. This is one of the handsomest of the American *Asters*, and highly deserves to be introduced as an ornamental plant.

sericeus.

24. *A. foliis oblongo-lanceolatis sessilibus sericeo-tomentosis, caule gracili decumbente laxe ramoso, ramis ramulis subunifloris, calycibus foliaceis subsquarroso-*

—Willd. sp. pl. 3. p. 2016.

A. argenteus. Mich. fl. amer. 2. p. 111.

Icon. Vent. hort. cels. t. 33.

On the banks of rivers, among rocks, in the Illinois country, and on the Mississippi and Missouri. ♀. Sept. Oct. v. s. in Herb. Lambert. A beautiful species; leaves resemble those of *Protea argentea*; flowers large, rays a fine purple, disk pale-yellow.

reticulatus.

25. *A. undique cano-tomentosus; foliis lanceolato-oblongis utrinque acutis sessilibus margine revolutis, subtus reticulato-venosis triplinervibus, caule superne ramoso, ramis apice corymboso-racemifloris, pedunculis subaphyllis, calycibus imbricatis laxiusculis, squamis acutissimis.*

In dry swamps of Carolina and Georgia. ♀. Aug.—Oct. v. v. About three feet high; flowers middle-sized, rays and florets white.

cornifolius.

26. *A. glaber; foliis oblongo-ovatis acuminatis brevi-petiolatis margine sebris, caule glabro, panicula pauciflora, ramis bifloris, calycibus subimbricatis.* —Willd. sp. pl. 3. p. 2039.

In Pennsylvania. Muhlenberg. ♀. +.

humilis.

27. *A. foliis subrhomboideis ovali-lanceolatis utrinque acu-*

mina
diver
laxis
p. 20

A. infir

cimer

Icon. I

In sha

Aug.

Not

in pr

28. *A. foliis
scabr
lycib
Lam.*

A. umb

On the

rolina

et La

29. *A. foliis
glabri
squar
p. 20*

A. pree

Icon. K

In low

Virgi

Bank

flowe

30. *A. foliis
marg
ramu
tis ae*

In dry

♀. J

ber

the p

31. *A. foliis
auric
subse
laxis
pl. 3.*

Icon. I

icutis squar.
car. 209.
Aug.—Nov.
ting resemi-
Good Hope,
e and large.
pubescenti-
te, racemo-
lanceolatis
).
oast : New
v. ; v. s. in
b; flowers,
lish-purple.
ican *Asters*,
ornamental

o-tomento-
, ramis ra-
bsquarrosis.

the Illinois
souri. Ʌ.
autiful spe-
ea; flowers

to-oblongis
subtus re-
ne ramoso,
culis suba-
uamis acu-

Ʌ. Aug.—
rs middle-

brevi-pe-
ula pauci-
s.—Willd.

inque acu-

minatis subpetiolatis glabris margine hispidis, corymbo
divergenti-dichotomo nudiusculo paucifloro, calycibus
laxis imbricatis, radiis 8-floris.—Willd. sp. pl. 3.
p. 2038.

A. infirmus. Mich. fl. amer. 2. p. 109.

A. divaricatus. Willd. sp. pl. 3. p. 2044. secundum spe-
cimen Gronov. et Banks.

Icon. Willd. hort. berol. 67. Pluk. alm. t. 79. f. 1.

In shady and rocky places : Canada to Carolina. Ʌ.

Aug.—Oct. v. v.; v. s. in Herb. Banks. et Gronov.
Not above a foot high ; flowers clear white, and large
in proportion to the plant.

28. A. foliis lanceolatis basi attenuatis acuminatis margine *amygdalinus*.
scabris, caule simplici apice corymboso-fastigiatu, ca-
lycibus laxis imbricatis, squamis lanceolatis obtusis.—
Lam. encycl. 1. p. 305.

A. umbellatus. Ait. kew. 3. p. 199.

On the banks of rivers and in swamps : Canada to Ca-
rolina. Ʌ. Sept.—Nov. v. v.; v. s. in Herb. Banks.
et Lambert. Rays large, white.

29. A. foliis linear-lanceolatis subintegerrimis glabris, caule
glabro apice paniculato, calycibus laxis imbricatis,
squamis acutis apice patulis.—Willd. sp. pl. 3.
p. 2030.

A. præaltus. *Lam. encycl. suppl.* 1. p. 493.

Icon. Rob. ic. 307.
In low grounds, on the side of ditches : New York to
Virginia. Ʌ. Aug.—Oct. v. v.; v. s. in Herb.
Banks. et Lambert. From four to six feet high ;
flowers middle-sized, reddish-blue.

30. A. foliis lanceolatis subamplexicaulibus apice attenuatis
margine scabris, caule a basi ramoso erecto hispido,
ramulis pilosis, calycinis squamis laxis linearibus acu-
tis æqualibus.—Willd. sp. pl. 3. p. 2030.

In dry swamps and copses : New York and Pennsylvania.
Ʌ. July—Sept. v. v.; v. s. in Herb. Banks. et Lam-
bert. Not above two feet high ; flowers resembling
the preceding, rays blue.

31. A. foliis linear-lanceolatis pilosis amplexicaulibus basi *noveæ angliæ*.
auriculatis, caule simpliciuscculo piloso stricto, floribus
subsessilibus terminalibus confertis, calycinis squamis
laxis coloratis lanceolatis disco longioribus.—Willd. sp.
pl. 3. p. 2032.

Icon. Herm. parad. 98.

salicifolia.

aestivus.

On the banks of rivers and borders of meadows: Canada to Virginia. ♀. Sept.—Nov. v. v.; v. s. in *Herb. Banks. et Lambert.* A highly ornamental species; in rich moist soil it will grow to the height of ten feet: flowers large, rays blue or purple, and almost every shade between red and blue.

cyaneus.

32. A. foliis linear-lanceolatis amplexicaulibus laevigatis, caule virgato-paniculato glaberrimo, ramis racemifloris, calycinis squamis laxis lanceolatis discum æquantiibus, interioribus apice coloratis.—*Hoffm. phytogr.* 1. p. 71. t. B. f. 1.

A. rubricaulis. *Lam. encycl.* 1. p. 305.

A. spurius. *Willd. sp. pl.* 3. p. 2032.

A. Novæ anglie. β. *Ait. kew.* 3. p. 201.

In old meadows and dry swamps: New England to Carolina. ♀. Sept.—Nov. v. v.; v. s. in *Herb. Lambert.* From three to four feet high; flowers large, in great abundance, of various tints of blue and purple; the handsomest of the genus.

carolinianus.

33. A. foliis oblongis utrinque attenuatis sessilibus, caule subscandente ramosissimo, ramulis pubescentibus summitate paucifloris, calycinis squamis lanceolatis squarrosis.—*Willd. sp. pl.* 3. p. 2017. *Walt. fl. car.* 208.

In shady moist hedges and swamps: Carolina and Georgia. ♀. Aug. Sept. v. v. From six to fifteen feet high; flowers large; rays purple or flesh-coloured, disk yellowish-purple.

grandiflorus.

34. A. foliis linearibus rigidis acutiusculis subamplexicaulibus, rameis reflexis, margine ciliato-hispidis, caule ramoso hirto, ramis unifloris, calycinis squamis squarrosis lineari-lanceolatis. *Willd. sp. pl.* 3. p. 2033.

Icon. Mill. ic. 282. *Dill. elth.* t. 36. f. 41. *Mart. cent.* 19. *Hoffm. phyt.* 1. p. 65. t. A. f. 1.

In dry sandy woods: Virginia to Carolina. ♀. Sept.—Nov. v. v.; v. s. in *Herb. Banks.* About two feet high; leaves small; flowers large, rays purple and blue, disk yellow.

phlogifolius.

35. A. foliis lanceolatis cordatis amplexicaulibus subtus pubescentibus, margine scabris, caule simplicissimo pubescente, panicula terminali laxa rariflora, calycinis squamis laxis imbricatis lanceolatis.—*Willd. sp. pl.* 3. p. 2034.

A. amplexicaulis. *Mich. fl. amer.* 2. p. 114.

In shady moist places: New Jersey to Virginia. ♀.

*Aug.
high
36. A. foliis
bus u
paten
squar
p. 20.
A. diver
In hedge
rocks
v. v.
foot t
or blu

37. A. foliis
pilosis
serrati
pido,
pl. 3.
Icon. H
In dry w
♀. Se
bert.
violet.

38. A. foliis
serratis
petiolata
imbricata
In North
blue, t

39. A. foliis
radicali
petiolis
losis, c
p. 2035
Icon. C
In meadow
nada to
Herb. A
high; f
changin
low to p
whole p

lows: Ca-
; v. s. in
ornamental
the height
purple, and

lævigatis,
racemiflo-
n æquanti-
phytogr. 1.

and to Ca-
Herb Lam-
wers large,
e and pur-

bus, caule
tibus sum-
atis squar-
car. 208.
and Geor-
to fifteen
flesh-co-

plexicauli-
lis, caule
mis squar-
2033.
Mart. cent.

4. Sept.
it two feet
purple and

ubitus pu-
ssimo pu-
calycinis
sp. pl. 3.

inia. 4.

Aug.—Nov. v. v. From eighteen inches to a foot high; flowers middle-sized, rays violet.

36. A. foliis oblongo-lanceolatis ciliatis cordatis amplexicaulis utrinque scabris pilosis, caule ramoso hirsuto, ramis patentibus elongatis paucifloris microphyllis, calycinis squamis imbricatis lanceolatis patulis. — Willd. sp. pl. 3. p. 2034.

A. diversitolius. Mich. fl. amer. 2. p. 113?

In hedges, old drying up swamps, and sometimes among rocks: New England to Carolina. 4. Sept.—Nov. v. v.; v. s. in Herb. Banks. et Lambert. From a foot to two high; flowers middle-sized, rays purple or blue, disk yellow.

** *Foliis cordatis ovatisque, serratis.*

37. A. foliis oblongis cordatis amplexicaulis integerrimis pilosis subundulatis, inferioribus ovatis cordatis sub-serratis petiolatis, petiolis alatis, caule paniculato hispidi, ramulis secundis foliosis unifloris. Willd. sp. pl. 3. p. 2035.

Icon. Hoffm. phyt. 1. t. C. f. 1. Herm. parad. 96.

In dry woods and old fields: New England to Carolina. 4. Sept.—Nov. v. v.; v. s. in Herb. Banks. et Lambert. Flowers smaller than the preceding; rays pale violet.

38. A. foliis oblongo-lanceolatis acuminatis sessilibus medio serratis, radicalibus oblongis cordato-sagittatis serratis petiolatis, caule ramoso glabro, squamis calycinis laxis imbricatis lanceolatis. — Willd. sp. pl. 3. p. 2035.

In North America. Wedemeyer. 4. +. Flowers blue, the size of the following species.

39. A. foliis ovato-lanceolatis subserratis petiolatis glabris, radicalibus ovato-cordatis serratis scabris petiolatis, petiolis nudis, caule ramosissimo glabro, ramulis pilosis, calycibus laxis subimbricatis. Willd. sp. pl. 3. p. 2035.

Icon. Corn. canad. 65. Moris. hist. 3. s. 7. t. 22. f. 34. In meadows, hedges, and on the borders of woods: Canada to Virginia. 4. Aug.—Nov. v. v.; v. s. in Herb. Banks. et Lambert. From two to four feet high; flowers rather small, but in great abundance, changing from white to blue in the ray, and from yellow to purple or brown in the disk, which gives the whole plant a beautiful variegated appearance.

patens.

undulatus.

sagittifolius.

paniculatus.

cordifolius. 40. A. foliis cordatis subtus pilosis argute serratis petiolatis, petiolis alatis, caule paniculato glabriuscōlo, panicula divaricata, calycibus laxis subimbricatis.—*Willd. sp. pl. 3. p. 2036.*

In shady woods, throughout the mountains : Canada to Florida. ♀. Sept.—Nov. v. v.; v. s. in *Herb. Banks.* Flowers small, white, or pale purple.

corymbosus. 41. A. foliis ovatis argute serratis acuminatis glabriusculis, inferioribus cordatis petiolatis, petiolis nudis, caule glabro superne corymboseo-fastigato, ramis pilosis, calycibus oblongis imbricatis, squamis obtusis arcte adpressis.—*Willd. sp. pl. 3. p. 2036.*

In shady woods : Canada to Florida. ♀. Aug.—Nov. v. v.; v. s. in *Herb. Banks. et Lambert.* A foot or two high ; flowers white, larger than the preceding.

macrophyllus. 42. A. foliis ovatis petiolatis serratis scabris, summis ovato-cordatis sessilibus, inferioribus cordatis petiolatis, petiolis submarginatis, caule ramoso diffuso, calycibus cylindricis arcte imbricatis, squamis oblongis acutis.—*Willd. sp. pl. 3. p. 2037.*

In shady woods about rocks ; Canada to Virginia. ♀. Sept.—Nov. v. v.; v. s. in *Herb. Banks. et Lambert.* A foot or two high ; flowers above the middle size, rays white, and sometimes blue.

heterophyllus. 43. A. foliis glabris, caulinis ovatis subcordatis acuminatis profunde serratis apice integerrimis, ramorum ovato-oblongis subserratis, ramulorum lanceolatis minutissime reflexo-patentibus, caule paniculato glabro, calycibus arcte imbricatis. *Willd. enum. 882.*

In North America. *Willdenow.* ♀. +.

*** *Foliis lanceolatis ovatisque, inferioribus serratis.*

amplexicaulis. 44. A. foliis ovato-oblongis acutis amplexicaulibus cordatis serratis glabris, caule paniculato glabro, ramulis 1-2-floris, calycinis squamis lanceolatis arcte imbricatis.—*Willd. sp. pl. 3. p. 2046.*

A. pensylvanicus. Lam. encycl. suppl. 1. p. 498.

On the borders of woods, in sandy soil : New Jersey to Virginia. ♀. Sept.—Nov. v. v. Flowers middle-sized, blue.

prenanthoides. 45. A. foliis amplexicaulibus spathulato-lanceolatis acuminatis medio serratis basi cordatis, ramulis pilosis, calycinis squamis lanceolatis squarroso.—*Willd. sp. pl. 3. p. 2046.*

In Pen

46. A. foli
ratis
bus
neati

In Non
disk

47. A. foli
lævib
ris, c
quant

In swan
Sept.
large,
purple

48. A. foli
glabri
glabre
ribus.

In old fi
to Ca
Lamb
great
to a de
flower
has a v

49. A. foli
acumini
angust
brevior
p. 204

Icon. Pl
In old fie
Aug.—
middle
ing to

50. A. foli
mis lu
uniflori
bus ac
Willd.

In woods
York t
blueish

atis petiolatis,
culo, panicula
—Willd. sp.

s : Canada to
s. in Herb.
purple.

glabriusculis,
nudis, caule
ramis pilosis,
obtusis arcte

Aug.—Nov.
rt. A foot or
he preceding.
ummis ovato-
petiolatis, pe-
uso, calycibus
longis acutis.

Virginia. Ʌ.
is. et Lambert.
e middle size,

tis acuminatis
norum ovato-
latis minutis-
o glabro, caly-
2.

us serratis.
ilibus cordatis
ramulis 1-2-
imbricatis.—

498.
New Jersey to
owers middle-

latis acuminata-
is pilosis, caly-
Willd. sp. pl. 3.

In Pennsylvania. Muhlenberg. Ʌ. +. Flowers blue.

46. A. foliis subamplexicaulibus lanceolatis, inferioribus subserratis glabris, ramulorum linearibus squarrosis, calycibus squarrosis disco brevioribus, squamis lineari-cuneatis.—Willd. enum. 884.

In North America. Willdenow. Ʌ. +. Rays violet; disk yellowish-brown.

47. A. foliis subamplexicaulibus lato-lanceolatis subserratis laevibus, caule ramosissimo glabro, ramulis multifloris, calycinis squamis lanceolatis laxis discum subæquantibus.—Willd. sp. pl. 3. p. 2046.

In swamps and wet woods: Pennsylvania to Virginia. Ʌ. Sept.—Nov. v. v.; v. s. in Herb. Lambert. Flowers large, pale rose- or flesh-coloured, with a tinge of purple.

48. A. foliis subamplexicaulibus lato-lanceolatis subserratis glabris, radicalibus medio serratis, caule ramosissimo glabro, calycinis squamis lanceolatis laxis disco brevioribus. Willd. sp. pl. 3. p. 2045.

In old fields and on the borders of woods: New Jersey to Carolina. Ʌ. Aug.—Oct v. v.; v. s. in Herb. Lambert. A beautiful species; flowers large and in great abundance, disk yellow, rays white, changing to a deep violet; so that at the same time there are flowers of different colours on the same plant; which has a very pleasing effect.

49. A. foliis subamplexicaulibus, superioribus lanceolatis acuminatis integerrimis, inferioribus lanceolatis basi angustatis serratis, ramulis virgatis, calycibus disco brevioribus laxis, caule glabro. Willd. sp. pl. 3. p. 2045.

Icon. Plak. alni. t. 326 f. 1. Herm. lugdb. 67.

In old fields and woods: Pennsylvania to Virginia. Ʌ. Aug.—Oct. v. v.; v. s. in Herb. Lambert. Flowers middle-sized; rays deep purple, disk yellow, changing to purple.

50. A. foliis subamplexicaulibus remotis oblongis integerimis lucidis, radicalibus subserratis, ramis simplicibus unifloris, calycibus imbricatis, foliolis subcuneiformibus acutis apice incrassatis, caule glabro angulato.—Willd. sp. pl. 3. p. 2044.

In woods and on the sides of ponds and ditches: New York to Virginia. Ʌ. Sept.—Nov. v. v. Rays blueish-purple.

adulterinus.

lævigatus.

versicolor.

mutabilis.

lævis.

- concinus.* 51. A. foliis lanceolatis subamplexicaulibus inferioribus subserratis glabris, caule simplici apice paniculato, calycibus arte imbricatis. *Willd. enum.* 884.
In fields and woods: New York and Pennsylvania. ♀. Sept.—Nov. v. v.; v. s. in *Herb. Lambert.* Leaves resembling those of *Phlox maculata*; rays blueish-purple.
- puniceus.* 52. A. foliis amplexicaulibus lanceolatis serratis scabriusculis, ramis paniculatis, calycibus laxis discum superantibus, foliolis linear-lanceolatis subæqualibus, caule hispido. *Willd. sp. pl. 3. p. 2040.*
Icon. *Herm. lugdb.* 651.
- purpureus.* α. A. septempedalis, caulis saturate purpureis. *Ait. kew. 3. p. 208.*
- β. A. amoenus. *Lam. encycl. 1. p. 306.*
A. octopetalis, caulis viridi-rufescens. *Ait. l. c.*
- rufescens.* A. hispidus. *Lam. encycl. 1. p. 307.*
On the banks of rivers and ditches, particularly near salt-water: Canada to Carolina, and in the range of mountains, about lakes. ♀. Sept.—Nov. v. v.; v. s. in *Herb. Banks. et Lambert.* From three to ten feet high; flowers large, purple or blue.
- floribundus.* 53. A. foliis subamplexicaulibus lanceolatis, inferioribus serratis, caule glabro, ramis corymbosis, ramulis multifloris pilosis, calycinis squamis laxis lanceolatis imbricatis. *Willd. sp. pl. 3. p. 2048.*
In North America. *Willdenow.* ♀. +. Resembles the following species.
- novi belgii.* 54. A. foliis subamplexicaulibus lanceolatis glabris margine scabris, inferioribus subserratis, ramis subdivisis, calycibus laxe-imbricatis, foliolis linear-lanceolatis, caule tereti glabro. *Willd. sp. pl. 3. p. 2048.*
Icon. *Herm. lugdb.* 69.
In hedges and old fields: New England to Virginia. ♀. Aug.—Oct. v. v.; v. s. in *Herb. Lambert.* Flowers middle-sized, pale purple.
- bellidiflorus.* 55. A. foliis amplexicaulibus angusto-lanceolatis supra scabris, inferioribus subserratis, caule ramosissimo, calycis squamis lanceolatis patentibus. *Willd. enum.* 686.
In North America. *Willdenow.* ♀. +. Ray pale red, disk yellow, changing to brown.
- spectabilis.* 56. A. foliis lanceolatis scabriusculis subamplexicaulibus, inferioribus medio serratis, ramis corymbosis, foliolis

calyci
squarr

In low n

Virgin

Lamb

57. A. foliis
margi
sis gla
ceolati
p. 204

In low g

to Virg

Lamb

58. A. foliis
basi at
mis di
linearit
p. 204

In low c

rare.

dle size

59. A. foliis
natis se
axillari
culis to
disco bi

In Canada

ers abo

60. A. foliis
perne ir
simplici
cato-dic
breviori

α. A. caule a

A. latifoli

β. A. caule

altiore.

In Canad

shady s

Virginia

foot or

61. A. foliis ol
rioribus
latis ser

erioribus sub-
iculato, caly-
4.
sylvania. Ʌ.
bert. Leaves
rays blueish-

is scabriuscum
superan-
alibus, caule

s. Ait. kew. 3.

us. Ait. l. c.
ticularly near
the range of
v. v. v.; v. s.
ee to ten feet

ferioribus ser-
ramulis multi-
olatis imbrica-

Resembles

abris margine
bdivis, caly-
ceolatis, caule

Virginia. Ʌ.
ert. Flowers

itis supra sca-
nosissimo, ca-
Willd. enum.

+. Ray pale

xicaulibus, in-
habosis, foliolis

calycinis laxis foliaceis subcuneiformibus acutiusculis
squarrosis. Willd. sp. pl. 3. p. 2048.

In low meadows, on the sides of woods: Pennsylvania to
Virginia. 2l. Aug.—Nov. v. v.; v. s. in Herb.
Lambert. About two feet high; flowers large, blue.

57. A. foliis oblongo-lanceolatis acuminatis sessilibus glabris
margini scabris, inferioribus serratis, ramis corymbo-
sis glabris, ramulis unifloris, calycinis squamis lan-
ceolatis acuminatis patentibus.—Willd. sp. pl. 3.
p. 2049.

In low grounds and on the banks of ditches: New York
to Virginia. Ʌ. Sept.—Nov. v. v.; v. s. in Herb.
Lambert. About three feet high; flowers large, blue.

58. A. foliis sessilibus serratis glabris spatulato-lanceolatis
basi attenuatis et margini utroque latere deflexis, ra-
mis divaricatis, calycibus laxis, foliolis lanceolato-
linearibus subæqualibus glabris.—Willd. sp. pl. 3.
p. 2049.

In low overflowed grounds: New York to Virginia;
rare. Ʌ. Oct. Nov. v. v. Flowers below the mid-
dle size; rays pale blue.

59. A. foliis subamplexicaulibus oblongo-lanceolatis acumi-
natis serratis glabris, caule pyramidato-ramoso, ramis
axillaris racemifloris folio vix longioribus, pedun-
culis tomentosis nudis, calycibus laxis subæqualibus
disco brevioribus.—Herb. Banks. mss.

In Canada. Ʌ. Oct. Nov. v. s. in Herb. Banks. Flow-
ers above the middle size; rays pale purple.

60. A. foliis lato-lanceolatis inferne attenuatis integris su-
perne inæqualiter serratis longissime acuminatis, caule
simplici flexuoso anguloso, panicula corymbosa divari-
cato-dichotoma, calycinis foliolis laxis linearibus disco
brevioribus.—Mich. fl. amer. 2. p. 109.

α. A. caule altiore, panicula multiflora foliosia.

A. latifolius. Herb. Banks. et Lambert.

β. A. caule pumilo, corymbo paucifloro nudo vix foliis
altiore.

In Canada and on the Allegany mountains, in rocky
shady situations. β. On the highest mountains of
Virginia and Carolina. Ʌ. Aug.—Oct. v. v. About a
foot or more high; flowers middle-sized; rays white.

61. A. foliis oblongis 3-nervibus basi angustatis acutis, supe-
rioribus sessilibus subintegerrimis, inferioribus petio-
latis serratis, caule simplici apice corymboso, calyci-

serotinus.

tardiflorus.

blandus.

acuminatus.

elatior.

pumilus.

conyzoides.

- bus cylindricis- squarrosis, radiis 5. brevissimis.—
Willd. sp. pl. 3. p. 2043.
- A. marylandicus.* *Mich. fl. amer.* 2. p. 108.
Conyza asteroides. Sp. pl. 1206.
 In fields and woods : Pennsylvania to Carolina. ♀. Aug.—Oct. v. v. About a foot high ; flowers small ; rays only five, white.
- peregrinus.* 62. *A. foliis sessilibus remotis oblongo-lanceolatis sensim acutis glabris, caule erecto subsimplici glabro superne hirsuto summitate 1—3-floro, calycinis squamis lanceo-lato-linearibus villosis acutissimis laxis longitudine disci, radiis numerosis* — *Herb. Banks. mss.*
 On the north-west coast : Unalashka. *D. Nelson.* ♀. v. s. in *Herb. Banks.* Flowers large, violet!
- Radula.* 63. *A. foliis lanceolatis serratis acuminatis rugosis scaberrimis, caule erecto angulato simplici, corymbo terminali, calycibus imbricatis, foliolis lanceolatis obtusiusculis subsquarrosis.* — *Willd. sp. pl. 3. p. 2044.*
 In Nova Scotia and on the high mountains of New York and Pennsylvania. ♀. Sept.—Nov. v. v. ; v. s. in *Herb. Banks. et Lambert.* Flowers middle size ; rays short, white.
- strictus.* 64. *A. foliis sessilibus angusto-lanceolatis serratis scabris, caule superne uno aut paucifloro, calycinis squamis imbricatis adpressis oblongis acutis discum subæquantibus.* — *Herb. Banks. mss.*
A. biflorus. *Mich. fl. amer.* 2. p. 114 ?
 In Labrador, at Hudson's Bay, and on high mountains in Pennsylvania. ♀. Sept. Oct. v. v. ; v. s. in *Herb. Banks.* From four inches to a span high ; flowers middle-sized ; rays pale violet, disk brownish-yellow.
- Tradescanti.* 65. *A. foliis lanceolatis serratis sessilibus glabris, ramis virgatis, calycibus imbricatis, caule tereti glabro.* *Willd. sp. pl. 3. p. 2047.*
A. vimineus. *Lam. encycl.* 1. p. 306.
 In hedges and dry swamps; frequent: Canada to Virginia. ♀. Aug.—Oct. v. v. ; v. s. in *Herb. Lambert.* Three or four feet high ; flowers very small ; rays white, and sometimes pale purple.
- recurvatus.* 66. *A. foliis sessilibus angusto-lanceolatis basi attenuatis, inferioribus medio serratis, caule ramoso glabro recurvato, calycinis squamis laxis imbricatis linearie lanceolatis subæqualibus.* — *Willd. sp. pl. 3. p. 2047.*
A. salicifolius. *Lam. encycl.* 1. p. 306.

In field
nia.
cedin

67. A. foliis
infim
ris, o
Willd.

In Nort
in H
pale I

68. A. foliis
infim
tissim
catis,
enum.

In low s
—No

69. A. foliis
apice
apice
linearis

In Nort
disk y

70. A. foliis
serrati
imbric

In North
disk y

71. A. foliis
subser
glabro
sp. pl.

In swam
Aug.—
ers fle

72. A. foliis
infim
glabrit
subpat

In North
Resem
changi

73. A. foliis

67

brevissimis.—

ina. ♀. Aug.
owers small;polatis sensim
labro superne
juamis lanceo-
s longitudine
nss.Nelson. ♀.
plet!is scaberrimis,
terminali, ca-
asiusculis sub-of New York
v. ; v. s. in
idle size; rayserratis scabris,
ycinis squamis
im subæquan-igh mountains
v. s. in Herb.
high; flowers
wnish-yellow.
ris, ramis vir-
glabro. Willd.da to Virginia.
mbert. Three
; rays white,asi attenuatis,
> glabro recur-
lineari-lanceo-
2047.

In fields and on road sides: New England to Pennsylvania. ♀. Sept.—Nov. v. v. Resembles the preceding; flowers pale blue.

67. A. foliis linear-lanceolatis acuminatis margine scabris, infimis subserratis, caule paniculato, ramulis unifloris, calycibus laxe imbricatis, foliolis lanceolatis. *Willd. enum. 886.*In North America. *Willdenow.* ♀. Sept.—Nov. v. s. in *Herb. Lambert.* Flowers of middle size; rays pale Mac, disk yellow, changing to red.68. A. foliis linear-lanceolatis acuminatis margine scabris, infimis subserratis, caulinis subreflexis, rameis patenssimis, caule laxe apice paniculato, calycibus imbricatis, foliolis lanceolatis acutis apice reflexis. *Willd. enum. 886.*

In low sandy fields: New Jersey to Virginia. ♀. Sept.—Nov. v. v. Rays white; disk yellow.

69. A. foliis lanceolatis acuminatis margine scabris, caulinis apice serratis, rameis integerrimis, caule glaberrimo apice paniculato, calycibus laxe imbricatis, foliolis lineari-subulatis. *Willd. enum. 887.*In North America. *Willdenow.* ♀. +. Rays white, disk yellow.70. A. foliis linearibus integerrimis, radicalibus oblongis sub-serratis, caule ramosissimo pubescente, calycibus laxe imbricatis. *Willd. enum. 888.*In North America. *Willdenow.* ♀. +. Rays white, disk yellow, changing to red or brown.71. A. foliis lanceolato-linearibus sessilibus glabris, infimis subserratis, ramulorum lanceolatis, caule paniculato glabro, ramis virgatis, calycibus imbricatis. *Willd. sp. pl. 3. p. 2050.*

In swamps and ditches: Pennsylvania to Virginia. ♀. Aug.—Oct. v. v. From four to six feet high; flowers flesh-coloured.

72. A. foliis linear-lanceolatis sessilibus integerrimis glabris, infimis lanceolatis subserratis, caule ramoso diffuso glabriusculo, calycibus imbricatis, foliolis lanceolatis subpatulis.—*Willd. sp. pl. 3. p. 2050.*In North America. *Willdenow.* ♀. Aug—Nov. +. Resembles the preceding; flowers smaller, rays white, changing to a pale violet.

73. A. foliis linearibus acuminatis integerimis, inferioribus dracunculoides.

eminens.

laxus.

simplex.

juncus.

lanceolatus.

SYNGENESIA SUPERFLUA. Aster.

linear-lanceolatis subserratis, ramis corymbosis, calycibus imbricatis, caule glabriusculo. Willd. sp. pl. 3. p. 2050.

A. artemisiflorus. Lam. encycl. suppl. 1. p. 500.

In low grounds and on sides of ditches: New Jersey to Carolina. ♀. Sept.—Nov. v. v.; v. s. in Herb. Lambert. About four feet or less high; flowers about half the size of No. 71, rays white.

fragilis.

74. A. foliis linearibus acuminatis integerimis, radicalibus oblongis serratis, ramis corymboso-paniculatis, calycibus imbricatis squamis adpressis, caule glabriusculo. Willd. sp. pl. 3. p. 2051.

In North America. Muhlenberg. ♀. v. s. in Herb. Lambert. About two feet high; flowers white, smaller than the preceding. This and the three preceding species are very nearly allied to one another, so much so that it is difficult to find sufficient characters to distinguish them, though their general appearance and the size of the flowers separate them at first sight.

miser.

75. A. foliis sessilibus lanceolatis serratis glabris, calycibus imbricatis, foliolis acutis, disco radii æquali, caule villosuscule. Willd. sp. pl. 3. p. 2051.

In old barren fields: Canada and New England. ♀. Aug.—Oct. v. v.; v. s. in Herb. Lambert.

divergens.

76. A. foliis elliptico-lanceolatis serratis glabris, caulinis linear-lanceolatis elongatis, ramis patentibus, calycibus imbricatis, caule pubescente. Willd. sp. pl. 3. p. 2052.

In old fields and barren woods: New York to Carolina. ♀. Sept.—Nov. v. v.; v. s. in Herb. Lambert. From three to five feet high; flowers small; rays white, disk reddish-brown.

diffusus.

77. A. foliis elliptico-lanceolatis serratis glabris proportionatis, ramis patentibus, calycibus imbricatis, caule pubescente. Willd. sp. pl. 3. p. 2052.

In fields and hedges; frequent: Canada to Carolina. ♀. Sept.—Nov. v. v.; v. s. in Herb. Lambert. Flowers white, small.

pendulus.

78. A. foliis elliptico-lanceolatis serratis glabris, ramulorum remotiusculis, ramis divaricatissimis pendulis, caule pubescente. Willd. sp. pl. 3. p. 2052.

In old fields; common. ♀. Oct. Nov. v. v.; v. s. in Herb. Lambert. Resembles the preceding; but the

stem
the d

637

1. D. herb
calyci
rosis.

In open
is.
cult.

The v
rous o
Caulis ei
alterm
basi d
sa, se
bus re
mular
piose i
tus: .
caput
lanceo
aristis
favulo

It appro
t. 168
t. 4. p
gnosis
obcunc
ribus e

It appro
t. 168
t. 4. p
gnosis
obcunc
ribus e

It appro
t. 168
t. 4. p
gnosis
obcunc
ribus e

It appro
t. 168
t. 4. p
gnosis
obcunc
ribus e

It appro
t. 168
t. 4. p
gnosis
obcunc
ribus e

It appro
t. 168
t. 4. p
gnosis
obcunc
ribus e

It appro
t. 168
t. 4. p
gnosis
obcunc
ribus e

1. B. caule
acutis.

Tagetes |
Tagetes |

Dyssodia |
On the
sippi.
yellow

stem forms a loose panicle with pendulous branches; the disk turns brown.

637. DONIA. *Ait. kew. ed. 2. t. 4. p. 82.*

1. D. herbacea; foliis oblongis amplexicaulibus serratis, calycinis squamis apice filiformibus revoluto-squarrosis.

squarrosa.

In open prairies, on the banks of the Missouri. *M. Lewis.* 4. Aug. Sept. v. s. in *Herb. Lewis.*; v. v. cult. Flowers yellow, resembling those of *Inula*. The whole plant is viscous, and has a strong resiniferous or balsamic scent.

Caulis erectus, teres, albidus, corymboso-ramosus. *Rami* alterni, angulati, apice uni- aut triflori. *Folia* alterna, basi dilatata, amplexicaulia, oblonga, acuta aut obtusa, serrata, glabra, utrinque squamulis punctiformibus resiniferis obsita. *Flores* 1—3. in summitate rameorum, corymbosi, flavi, ante explicationem copiose resiniferi. *Calyx* hemisphaericus, arcte imbricatus: *Squamis* apice filiformibus, revolutis et quasi caput medusae referentibus. *Flosculi* radii linearilanceolatae, acutae. *Semina* oblongo-ovata. *Pappus*: aristis 2—4. deciduis. *Receptaculum* planum, nudum, favulosum.

It approaches near to *Aster glutinosus* Cav. ic. 2. p. 53. t. 168. which is the *Donia glutinosa* Ait. kew. ed. 2. t. 4. p. 82. which I distinguish by the following diagnosis: *Donia glutinosa*: trutescens; foliis oblongo-obcuneatis sessilibus serratis, calycinis squamis linearibus erectis.

638. BCEBERA. *Willd. sp. pl. 3. p. 2125.*

1. B. caule paniculato, foliis oppositis subbipinnatis glabris acutis.—*Willd. sp. pl. 3. p. 2125.*

chrysanthemoides.

Tagetes papposa. *Mich. fl. amer.* 2. p. 132.

Tagetes puinila. *Hbrt. pisan.*

Dyssodia glandulosa. *Cav. demonstr. bot.* p. 202.

On the overflowed banks of the Missouri and Mississippi. ⊖. Aug. Sept. v. s. Flowers small, golden-yellow.

639. ACTINELLA. *Pers. syn. 2 p. 469.*Actinea. *Juss. in annal. mus. 2. p. 425.**lanata.*

1. A. undique lanato-tomentosa; foliis linearibus superne pinnatifidis, pedunculis elongatis unifloris, radius bidentatis, ~~semim~~ glabris 5-gonis.

On the high lands of the Kooskoosky. *M. Lewis.* ♀. June, July. *v. s. in Herb. Lewis.* Flowers orange-yellow. It resembles in habit Jussieu's *Actinea*, in *Annal. du mus. 2. t. 61. f. 2.*

Planta omnibus partibus tomento albo lanato tecta. *Qua-*
lis ramosus, teres. *Rami* alterni, superne subdivisi.
Ramuli uniflori. *Folia* caulinis alterna, linearia, superne dilatato-pinnatifida, dentata; ramea linearia, integra. *Pedunculi* terminalis, elongati, superne in-
crassati. *Flores* aurantiaci, magnitudine *Tagetidis* erectae. *Calyx* oblongus, simplex, polyphyllus: foliola 12—14. linear-lanceolata, acuta. *Corolla* radii 12—14. oblongæ, bidentatae, nervosæ: disci tubulosæ, concolores. *Pappus*: paleis 4—6. scariosis, albidis, acutis, interdum laceris. *Semina* prismatico-
oblonga, 5-gona.

640. HELENIUM. *Gen. pl. 1299.**autumnale.*

1. H. foliis lanceolatis serratis subdecurrentibus, caule superne corymboso, corollulis disci 5-fidis: radii planis reflexis.—*Willd. sp. pl. 3. p. 2120.*

Icon. *Pluk. amalthe. t. 372. f. 4.* *Corn. canad. 63.*

β. H. foliis pubescens. *Willd. sp. pl. 3. p. 2121.*

On the banks of rivers, ponds, and ditches: Canada to Carolina. ♀. Oct. Nov. *v. v.* About two feet high; flowers bright yellow.

canalicula-
tum.

2. H. foliis lanceolatis serratis subdecurrentibus, caulis simplicibus, corollulis disci 5-fidis: radii canaliculatis. *Lamarek in journ. hist. nat. 213. t. 35.*

3. H. foliis lato-decurrentibus, inferioribus subpinnatifidis, superioribus lanceolatis integerimis glabris, corollulis disci 4-dentatis.—*Willd. sp. pl. 3. p. 2121.*

Rudbeckia alata. Jacq. in rar. t. 503.

Icon. *Labillard. in act. soc. hist. nat. par. 1. t. 4.*

On the banks of the Mississippi. ♀. July, Aug. *v. v.* From three to four feet high; flowers smaller than the preceding.

quadridenta-
tum.

1. B. foli
nibus
p. 2
Matrix
Chrys
Icon.
On the
nois
disk

2. B. foli
latis
bus
Wit
Icon.
On the
gini
cedi

6
1. S. foli
tis tu
Icon.
On the
v. s.

1. P. foli
cory
p. 2
Icon.
In No
be th
chau

1. E. erec
tenu
calyc
p. 2
Verbes

641. BOLTONIA. *Gen. pl.* 1309.

1. *B. foliis integerrimis, floribus longe pedunculatis, seminibus ovalibus glabris submuticis.* — *Willd. sp. pl.* 3. *p. 2162.* *Matricaria asteroides.* *Linn. mant.* 116. *Chrysanthemum carolinianum.* *Walt. fl. car.* 204. *Icon. L'Herit. sert. angl. t. 25.* *ined.* On the banks of ponds and swamps: Carolina and Illinois. \mathcal{U} . Aug.—Oct. *v. s.* Rays flesh-coloured; disk yellow.
2. *B. foliis inferioribus serratis, floribus breviter pedunculatis, seminibus obcordatis conspicue alatis pubescentibus, aristis pappi duabus ipsorum longitudine.* — *Willd. sp. pl.* 3. *p. 2163.* *Icon. L'Herit. sert. angl. t. 26.* *ined.* On the banks of rivers and ponds. Pennsylvania to Virginia. \mathcal{U} . July, Aug. *v. v.* Resembles the preceding very much.

642. SIEGESBECKIA. *Gen. pl.* 1320.

1. *S. foliis sessilibus ovatis dentatis, floribus disci tridentatis triandris.* — *Willd. sp. pl.* 3. *p. 2220.* *flosculosa.* *Icon. L'Herit. stirp. t. 19.* On the north-west coast. *M. Lewis.* \odot . July, Aug. *v. s. in Herb. Lewis.* Flowers yellow.

643. PHÆTHUSA. *Gen. pl.* 1319.

1. *P. foliis oppositis ovatis acuminatis serratis triplinervibus, corymbo terminali brachiatu.* — *Willd. sp. pl.* 3. *p. 2221.* *americana.* *Icon. Pluk. mant. t. 342.* In North America. \mathcal{U} . +. I strongly suspect this to be the same with *Verbesina Siegesbeckia*, although Michaux declares it to be different.

644. ECLIPTA. *Gen. pl.* 1316.

1. *E. erecta, dichotoma, strigosa; foliis lanceolatis basi attenuatis rarer serratis, pedunculis geminis elongatis, calycinis foliolis ovatis acuminatis.* — *Willd. sp. pl.* 3. *p. 2217.* *erecta.* *Verbesina alba.* *Sp. pl.* 1272.

Cotula alba. *Syst. nat.* 2. p. 564.
Icon. *Dill. eth.* t. 113. f. 137. *Pluk. alm.* t. 109. f. 1.

Moris. hist. 3. s. 6. t. 13. f. 16.

In dry gravelly soil: Virginia to Florida. ♂. June, July.
v. v. Flowers small, white.

procumbens. 2. E. procumbente-assurgens; foliis longo-lanceolatis inferne angustatis rarer subserratis, pedunculis longiusculis, calycinis foliolis acute-lanceolatis, flosculis 4-fidis.—*Mich. fl. amer.* 2. p. 129.

In Lower Carolina. ♂. July, Aug. v. s. Flowers small, white.

brachypoda. 3. E. divaricato-prostrata; foliis lanceolatis subserratis, pedunculis solitariis geminisque brevibus, calycinis foliolis ovali-lanceolatis, flosculis 5-fidis.—*Mich. fl. amer.* 2. p. 130.

Amellus carolinianus. *Walt. fl. car.* 213.

In low sandy fields: Carolina. ♂. July, Aug. v. v.

In No
lou

3. A. fol
mu
Wi

Icon.
In fiel
v.
whi

4. A. fol
cini
sito
Icon.
On th
v. s.

647. T

1. T. bir
lato
ner
Polym
Icon.
In fert
Virg
foui

6

1. H. fo
syn.
Bupht
Heliar
Rudbe
Silphit
On the
sylv.
plan

64

1. B. foli
peti
p. 2
Icon.
Cate

645. ANTHEMIS. *Gen. pl.* 1312.

1. A. receptaculis conicis, paleis lanceolatis, seminibus coronato-marginatis, foliis bipinnatis lacinii lanceolato-linearibus. *Willd. sp. pl.* 3. p. 2180.

Icon. *Engl. bot.* 602.

In waste grounds and on road sides: Pennsylvania to Virginia. ♂. ♂. Aug. Sept. v. v. Rays white, disk yellow.

2. A. caule repente, foliis ovali-lanceolatis subintegris, pedunculis terminalibus unifloris longissimis, calycibus subæqualibus 10—12-phylloides.—*Walt. fl. car.* 211.

Spilanthes repens. *Mich. fl. amer.* 2. p. 181.

On overflowed banks of rivers: South Carolina. ♂. July—Sept. v. v. Flowers yellow.

646. ACHILLEA. *Gen. pl.* 1313.

1. A. foliis linearibus acuminatis æqualiter argute serratis glabris. *Willd. sp. pl.* 3. p. 2191.

Icon. *Fl. dan.* 643. *Engl. bot.* 757.

In open dry swamps: Canada and New York. ♀. Aug. Sept. v. v. Flowers white.

2. A. foliis radicalibus bipinnatifidis, caulinis incisis, lacinii ovatis serrato-mucronatis obtusis.—*Vent. hort. cels* 95.

109. f. 1.
une, July.
solatis in-
is longius-
osculis 4-
Flowers
ratis, pe-
ycinis fo-
Mich. fl.

g. v. v.

nibus co-
ncelato-

ia to Vir-
nite, disk
gris, pe-
calycibus
211.
na. ♂.

e serratis
♀. Aug.
is, laci-
nt. hort.
- In North America. *Bosc.* ♀. +. Flowers rose-coloured.
3. A. foliis bipinnatifidis pilosis laciniis linearibus dentatis *Millefolium*. mucronatis, caulinis sulcatis. *Smith. fl. brit.* 2. p. 908. *Willd. sp. pl.* 3. p. 2208. *Icon. Fl. dan.* 737. *Engl. bot.* 758. In fields and on road sides; common. ♀. June—Aug. v. v. Probably introduced from Europe. Flowers white or rose-coloured.
4. A. foliis bipinnatifidis villosis, pinnis confertissimis, laciniis linear-lanceolatis integerrimis, corymbo composto. *Willd. sp. pl.* 3. p. 2209. *Icon. Bot. mag.* 498. *Gmel. it.* 1. t. 25. f. 2. On the banks of the Koooskoosky. *M. Lewis.* ♀. June. v. s. in *Herb. Lewis.* Flowers yellow. *tomentosa.*
647. TETRAGONOTHECA. *L'Herit. stirp.* p. 177.
1. T. hirsutissima; foliis oppositis amplexicaulibus spathulato-ovatis, summis ovatis acuminatis dentatis tripli-nervibus.—*Willd. sp. pl.* 3. p. 2116. *Polymnia Tetragonotheca. Syst. veg.* 658. *Icon. Dill. elth.* t. 283. f. 305. In fertile soil, on the borders of woods and along hedges: Virginia to Florida. ♀. July—Sept. v. v. From four to six feet high; flowers large, yellow.
648. HELIOPSIS. *Pers. syn.* 2. p. 473.
1. H. foliis oppositis ovatis serratis triplinervibus. *Pers. syn.* 2. p. 473. *Buphtalmum helianthoides. Willd. sp. pl.* 3. p. 2236. *Helianthus lœvis. Sp. pl.* 1278. *Rudbeckia oppositifolia. Sp. pl.* 1280. *Silphium solidaginoides. Sp. pl.* 1302. On the banks of rivers, and in rocky shady places: Pennsylvania to Carolina. ♀. Aug.—Oct. v. v. A tall plant, resembling a species of *Helianthus*.
649. BUPHTHALMUM. *Gen. pl.* 1231.
1. B. foliis oppositis cuneato-lanceolatis carnosis incanis, petiolis bidentatis, caule fruticoso.—*Willd. sp. pl.* 3. p. 2229. *Icon. Dill. elth.* t. 28. f. 4. *Pluk alm.* t. 115. f. 4. *Catesb. fl. car.* 1. t. 93. *frutescens.*

On the sea-coast: Virginia to Florida. $\frac{1}{2}$. Aug. Sept.
v. v. A shrub about five feet high.

angustifoli-
um.

2. B. foliis alternis linearibus superne lato-lanceolatis integerimis glabris, calycinis foliolis acute-lanceolatis.—*Herb. Banks. mss.*

In Georgia and Florida. *Bartram.* $\frac{1}{2}$. *v. s. in Herb. Banks.*

sagittatum.

3. B. tomentosum; foliis radicalibus longissime petiolatis oblongis sagittatis integerimis subtrinervibus, caulinis oblongis in petiolum attenuatis, caule subtrifloro, calycinis foliolis exterioribus disco longioribus.

On dry barren hills, in the Rocky-mountains. *M. Lewis.*
 $\frac{1}{2}$. June, July. *v. s. in Herb. Lewis.* Flowers large, yellow. The natives eat the young stems as they spring up, raw.

650. AMELLUS. *Gen. pl. 1315.*

villosus.

1. A. villosissimus; foliis sessilibus oblongis acuminatis integerimis, floribus axillaribus brevi-petiolatis, radiis integris, paleis setaceis.

On the Missouri. $\frac{1}{2}$. *v. s.* Rays yellow.

spinulosus.

2. A. canescens; foliis bipinnatifidis inciso-dentatis, laciiniis linearibus rigido-mucronatis, floribus lateralibus et terminalibus congestis, radiis bidentatis, paleis setaceis.

In open prairies on the Missouri. *M. Lewis.* $\frac{1}{2}$. Aug. Sept. *v. s. in Herb. Lewis.* Rays yellow.

*Tota planta rigida, scabra, cano-viridescens. Caulis ramosissimus. Rami angulosi. Ramuli laterales breves, uniflori. Folia alterna, bipinnatifida, rigida, laciiniis linearibus, spinula terminatis, inferioribus inciso-dentatis. Flores laterales solitarii et terminales congesti, subcorymbosi, lutei. Calyx imbricatus: squamis exterioribus brevioribus, acutis, hirsutis; interioribus linearibus, scariosis. Corollæ radii linear-lanceolatae, bidentatae. Receptaculum: Paleis brevibus, subulatis. Both species are an intermediate link between *Aster* and *Inula*. They approach the nearest to *Amellus*; but the receptacle is not, properly speaking, paleaceous, but only setaceous.*

651. VERBESINA. *Gen. pl. 1317.*

virginica.

1. V. caule angusto alato, foliis alternis lato-lanceolatis sub-serratis, corymbo composito, calycibus oblongis pubescentibus.—*Willd. sp. pl. 3. p. 2222.*

In shad
rolin
four,
2. V. caul
acum
sumt
Siegesb
In shad
v. v.
rešen
3. V. caul
coryt
subgl
Coreop
a. V. radi
Icon. J
β. V. sem
Athana
In the r
Carol
to se
riety
and I
4. V. caul
ter de
scenti
amer.
In the v
nasse
like F
1. Z. florit
subpe
Icon. I
149.
On the b
Rays :

im.
Aug. Sept.
s integerri-
tis.—Herb.
s. in Herb.
e petiolatis
is, caulinis
rifloro, ca-

M. Lewis.
wers large,
ns as they

iminatis in-
latis, radiis
tis, laciinis
ibus et ter-
is setaceis.
♀. Aug.

Caulis rati-
les breves,
da, laciinis
nciso-dent-
i congesti,
quamis ex-
rioribus li-
anceolatæ,
subulatis.
Aster and
s; but the
ceous, but

olatis sub-
longis pu-

In shady woods on the mountains: Pennsylvania to Ca-
rolina. ♀. July—Sept. v. v. Rays from three to
four, white.

2. V. caule alato, foliis oppositis ovato-lanceolatis utrinque Siegesbeckia.
acuminatis acute serratis, panicula brachiata, ramulis summitate fasciculifloris.—Willd. sp. pl. 3. p. 2224.

Siegesbeckia occidentalis. Sp. pl. 1209.

In shady woods: Virginia to Carolina. ♀. July—Sept.
v. v. Rays from one to three, yellow. It has a strong
resemblance to *Phæthusa*.

3. V. caule alato, foliis alternis lanceolatis serratis, panicula Coreopsis.
corymbosa laxa foliosa, calycibus laxis patulis, disco subgloboso.—Mich. fl. amer. 2. p. 134.

Coreopsis alternifolia. Willd. sp. pl. 3. p. 2257.

- α. V. radiata aut flosculosa luteiflora. Mich. l. c.

Icon. Jacq. hort. vind. t. 110. Pluk. alm. t. 159. f. 3.

- β. V. semper flosculosa albiflora. Mich. l. c.

Athanasia paniculata. Wilt. fl. car. 201.

In the mountains of Virginia and Carolina. β. On the
Carolina coast. ♀. July—Sept. v. v. From three
to seven feet high; flowers yellow. The white va-
riety I have only seen in the Walterian Herbarium,
and I strongly suspect it to be a very distinct species.

4. V. caule alato, foliis alternis lato-lanceolatis acutis leviter dentatis subtus albido-villosis, supra aspero-pubescentibus, pedunculis unifloris aggregatis.—Mich. fl. amer. 2. p. 135.

In the western parts of the Allegany mountains: Ten-
nessee and Illinois. ♀. Aug. Sept. v. s. Flowers
like *Helianthus*.

652. ZINNIA. Gen. pl. 1304.

1. Z. floribus pedunculatis, foliis oppositis ovato-lanceolatis subpetiolatis. Willd. sp. pl. 3. p. 2139.

Icon. Linn. dec. t. 12. Jacq. obs. 2. t. 40. Bot. mag.
149.

- On the banks of the Mississippi. ♂. July, Aug. v. v.
Rays yellow, orange, and sometimes brick-red.

Coreopsis.

lutea.

alba.

multiflora.

III. FRUSTRANEA.

653. BIDENS. *Gen. pl. 1267.**cernua.*1. B. floribus subradiatis cernuis, calyce exteriore flore longiore, foliis lanceolatis subconnatis dentatis. *Willd. sp. pl. 3. p. 1716.*Icon. *Fl. dan. 841. Curtis fl. lond. 3. t. 55. Pet. hort. brit. t. 20. f. 6.**minima.*β. B. foliis lanceolatis sessilibus, floribus seminibusque erectis. *Sp. pl. 1165.*Icon. *Fl. dan. 312. Dill. in Raii syn. t. 7. f. 2.**Coreopsis.*γ. B. foliis lanceolatis serratis oppositis amplexicaulis. *Sp. pl. 1281.*Icon. *Barrel. ic. t. 1209. Lass. fl. pruss. t. 11.*

About ponds, ditches and springs : Canada to Pennsylvania. ☽. Aug. Sept. The flowers of this and all the following species are yellow, and the most of them are either without or with rays.

*chrysanthemoides.*2. B. floribus radiatis cernuis, radio calyci subæquali triplo longiore, foliis oblongis utrinque attenuatis dentatis basi connatis. *Willd. sp. pl. 3. p. 1717. Mich. fl. amer. 2. p. 136.*Coreopsis flammula. *Herb. Banks. mss.*Coreopsis perfoliata. *Walt. fl. car. 215.*

In overflowed places and about ponds : New England to Carolina. ☽. Aug.—Oct. v. v. Flowers large, bright yellow. The awns of the seeds are by two, three, or four; not always by two, as Michaux asserts.

*frondosa.*3. B. floribus discoideis, calyce exteriore flore sexduplo longiore, foliolis basi ciliatis, foliis inferioribus pinnatis, superioribus ternatis lanceolatis serratis. *Willd. sp. pl. 3. p. 1718.*Icon. *Moris hist. 3. s. 6. t. 5. f. 21.*

In shady fertile woods and fields : Pennsylvania to Carolina. ☽. July—Oct. v. v.

*connata.*4. B. floribus discoideis, calyce exteriore flore triplo longiore, foliis caulinis ternatis, foliolis lateralibus connatis, floralibus oblongo-lanceolatis. *Willd. sp. pl. 3. p. 1718.*

In fields and woods; frequent : Canada to Carolina. ☽. July—Oct. v. v.

pilosa.

5. B. floribus discoideis, calyce exteriore longitudine interioris, foliis inferioribus pinnatis, superioribus ter-

natis,
duplo
Icon. *Dil*
In old fiel
rolina.
6. B. florib
terioris
dis.
Icon. *He*
f. 23.
A comm

6;
1. C. foliis s
tis, pe
scabris
pl. 3. t

Icon. *Di*
β. C. caule
p. 137.
C. crassifi

In mount
Aug.—
with fo

2. C. glabra
minatis
nalibus

In Caroli
Flower
bright y

3. C. foliis c
decurrens
In Virgin

4. C. foliis ov
cronatis
cuneato
p. 2257.
On the hi
July—S
rather si

natis, foliolis oblongis, terminali lanceolato reliquis duplo longiore. *Willd. sp. pl. 3. p. 1720.*

Icon. *Dill. elth. t. 43. f. 51.*

In old fields and cultivated grounds: Pennsylvania to Carolina. ♂. July—Oct. v. v. A common weed.

6. B. floribus subradiatis, calyce exteriore longitudine interioris, foliis bipinnatis, foliolis lanceolatis pinnatifidis. *Willd. sp. pl. 3. p. 1721.*

Icon. *Herm. parad. t. 123. Moris. hist. 3. s. 6. t. 7. f. 23.*

A common weed in all old fields. ♂. July—Oct. v. v.

lipinnata.

654. COREOPSIS. Gen. pl. 1325.

* *Foliis oppositis, indivisis.*

1. C. foliis sessilibus lanceolato-linearibus integerrimis ciliatis, pedunculis elongatis nudis, seminibus orbiculatis scabris alatis apice bidentatis emarginatis.—*Willd. sp. pl. 3. p. 2256.*

Icon. *Dill. elth. t. 48. f. 56. Mart. cent. 26.*

β. C. caule foliisque canescenti-villosa. *Mich. fl. amer. 2. p. 137.*

lanceolata.

villosa.

C. crassifolia. *Willd. sp. pl. 3. p. 2256.*

In mountainous situations: Virginia and Carolina. ♀. Aug.—Oct. v. v. Flowers deep yellow; rays broad with four or five teeth.

2. C. glabra; foliis petiolatis lanceolato-ovatis sensim acuminatis argute serratis, pedunculis axillaribus terminalibusque dichotome-corymbosis.

In Carolina. *P. Miller. 4. v. s. in Herb. Banks.* Flowers middle size. This and all the rest are of a bright yellow colour.

arguta.

3. C. foliis ovatis acuminatis serratis petiolatis triplinervibus decurrentibus.

In Virginia. *Bartram. 3. v. s. in Herb. Banks.*

alata.

4. C. foliis ovatis acuminatis crenato-dentatis, dentibus mucronatis, petiolis brevibus, radiis integris, seminibus cuneato-oblongis apteris apice nudis.—*Willd. sp. pl. 3. p. 2257. Mich. fl. amer. 2. p. 137.*

latifolia.

On the high mountains of Virginia and Carolina. ♀. July—Sept. v. v. A tall growing species; flowers rather small.

** *Foliis oppositis, divisis.*

- aristata.* 5. C. pubescens ; foliis quinato-pinnatis serratis, radiis integris lato-ovalibus, seminibus cuneato-obovatis bizarriatis, aristis divaricatis.—*Willd. sp. pl. 3. p. 2253.*
C. aristosa. Mich. fl. amer. 2. p. 140.
 In South Carolina and Illinois. ♀. Aug. Sept. v. s.
 Flowers large ; rays very broad.
- trichosperma.* 6. C. glabella, dichotoma ; foliis subquinato-pinnatis lanceolatis serratis, calycis exterioris foliolis (8) spathulatis ciliato-serratis, radiis integris, seminibus cuneatis sub-4-dentatis.—*Willd. sp. pl. 3. p. 2252.* *Mich. fl. amer. 2. p. 139.*
 In cedar-swamps : New Jersey and Carolina. ♂. Aug.—Oct. v. v. Flowers small ; resembles *C. coronata.*
- auriculata.* 7. C. pubescens ; foliis subsessilibus ovali-lanceolatis integerrimis, inferioribus ternatis, calyce exteriore profunde partito, radiis 4-dentatis, seminibus subrotundobovatis apice emarginato-bidentatis.—*Willd. sp. pl. 3. p. 2256.*
Icon. Pluk. alm. t. 83. f. 5. ♂ t. 242. f. 4.
 On the high mountains of Virginia and Carolina. ♀. Aug.—Oct. v. v.
- tripteris.* 8. C. glabra ; foliis petiolatis lanceolatis integerrimis, radicalibus pinnatis, caulinis ternatis, radiis integris, seminibus obovatis apice nudis.—*Willd. sp. pl. 3. p. 2253.*
Icon. Moris. hist. 3. s. 6. t. 3. f. 44.
 On high mountains : Virginia to Carolina. ♀. Aug.—Oct. v. v.
- senifolia.* 9. C. subpubescens ; foliis sessilibus integerrimis ternatis, radiis integris, seminibus cuneato-oblongis.—*Willd. sp. pl. 3. p. 2254.*
C. major. Walt. fl. car. 214.
C. stellata. Herb. Banks. mss.
 On dry stony hills, in the mountains of Virginia and Carolina. ♀. Aug.—Oct. v. v.
- aurea.* 10. C. foliis serratis, radicalibus 3-partitis, caulinis trifidis integris lanceolato-linearibus. *Willd. sp. pl. 3. p. 2253.*
 In North America. ♀. +.
- trifida.* 11. C. glabra ; foliis trifidis subulatis fasciculatis, pedunculis terminalibus unifloris, radiis 3-fidis, caule subflexuoso.—*Lam. encycl. suppl. 2. p. 353.*

- Icon. I
 In Noi
C. ve
 12. C. folii
 aribu
sp. pi
C. verti
Icon. F
 On higl
 Aug.
 13. C. folii
 aribu
sp. pi
C. delph
Icon. E
 On higt
 —Oc
 14. C. glab
 sup
 culi
 In low
 souri.
 15. C. folii
 florib
 In Geor
 16. C. caule
 rumq
 gusta
 ginali
C. gladi
 In shad
 na.
 17. C. folii
 decur
 altern
 In Nort
 18. C. folii
 oblon
pl. 3.
 In Caro
Bank.

SYNGENESIA FRUSTRANEA. Coreopsis.

569

- Icon. *Lam. illustr. t. 704. f. 2.*
In North America. *Lamarc. 4. +.* Resembles
C. verticillata.
12. *C. foliis verticillatis ternis quinisve pinnatis, pinnis line-
aribus tripartitis indivisisque, disco concolore. Willd.
sp. pl. 3. p. 2252.*
C. verticillata. Lam. encycl. 2. p. 108.
Icon. *Pluk. mant. t. 344. f. 4.*
On high dry mountain lands: Virginia and Carolina. 4.
Aug.—Oct. v. v.
13. *C. foliis verticillatis ternis quinisve pinnatis, pinnis line-
aribus tripartitis indivisisque, disco discolore. Willd.
sp. pl. 3. p. 2151.*
C. delphinifolia. Lam. encycl. 2. p. 108.
Icon. *Bot. mag. 150. Ehret pict. t. 9. f. 1.*
On high mountains: Virginia and Carolina. 4. Aug.
—Oct. v. v.
14. *C. glabra; foliis petiolatis, inferioribus bipinnatifidis,
superioribus linearis-partitis, calycibus simplicius-
culis, seminibus nudis.—Willd. sp. pl. 3. p. 2253.*
In low grounds of Carolina. *Michaux.* On the Mis-
souri. *M. Lewis.* ♂. July, Aug. v. v.

*tenuifolia.**verticillata.**mitis.**acuta.**dichotoma.**procera.**angustifolia.*

aspera.

19. *C. foliis lanceolato-linearibus asperis, superioribus alternis, inferioribus oppositis, caule unifloro.*
In Maryland. *v. s. in Herb. Banks.*

*atrorubens.*655. HELIANTHUS. *Gen. pl. 1322.** *Foliis oppositis.**pubescens.*

1. *H. totus hispidus; caule superne nudiusculo laxe paniculato, foliis spathulatis ovatis crenatis triplinervibus scabris, squamis calycinis ovato-lanceolatis longitudine disci atropurpureo.* — *Willd. sp. pl. 3. p. 2245.*
Icon. *Dill. elth. t. 94. f. 110.* *Mart. cent. t. 20.*
In gravelly and slate soil, in the western parts: Pennsylvania to Carolina. ♀. ♂. Aug. Sept. *v. v.* Rays yellow; disk dark purple.
2. *H. cano-pubescent; caule villoso, foliis sessilibus cordato-ovatis amplexicaulibus triplinervibus adpresso-subserratis, calycinis squamis lanceolatis villosis.* — *Willd. sp. pl. 3. p. 2244.*
H. mollis. Lam. encycl. 3. p. 81.
H. canescens. Mich. fl. amer. 2. p. 140.
In wet meadows of Kentucky and Tennessee. ♂. Aug.—Oct. *v. s.* Both ray and disk are yellow, which is the case with all the following species.

varicatus.

3. *H. caule glabro ramosissimo, foliis suboppositis sessilibus lanceolato-ovatis trinervibus, panicula trichotoma gracili parviflora.* — *Willd. sp. pl. 3. p. 2244.*
Icon. *Moris. hist. 3. s. 6. t. 7. f. 66. mala.*
On the borders of woods and in hedges: Canada to Pennsylvania. ♀. Aug.—Nov. *v. v.* About five or six feet high; flowers smaller than any other species here enumerated.

frondosus.

4. *H. caule inferne glabro, foliis ovatis argute serratis, pedunculis scabris, calycibus squarrosis undulatis frondosis ciliatis, radiis 8-floris.* — *Willd. sp. pl. 3. p. 2241.*
In Canada. ♀. +. About four feet high; resembles *H. decapetalus* and *multiflorus* in several respects, but is distinct.

trachelifolius.

5. *H. foliis ovato-lanceolatis acuminatis serratis triplinervibus utrinque scaberrimis, calycinis squarnis linearilanceolatis ciliatis, exterioribus longioribus.* — *Willd. sp. pl. 3. p. 2241.*
H. Gigas. Mich. fl. amer. 2. p. 141.

SYN

In wood

Aug.—

three

6. *H. glab-*

pauci-

triplin-

cinis se-

varicata

In the v

Oct.

purple

7. *H. foliis*

pliuver-

tis, cal-

pl. 3. p.

In dry sw

Carolin

looking

Stems

8. *H. foliis*

nervibu

calycin

p. 2243

Icon. *Jac*

In copse

lina.

ceding

recepta

ceding.

9. *H. foliis*

scabris,

atis. *W*Icon. *Boo*

In Canad

10. *H. caule*

scabris

calycin

p. 2242

In North

11. *H. foliis*

scabris,

In woods and dry swamps: Canada to Carolina. ♀.
Aug.—Oct. v. v. Resembles *H. decapetalus*. About
three feet high.

6. *H. glaberrimus*; caule paniculato, ramis summitate
paucifloris, foliis subsessilibus longissime-lanceolatis
triplinervibus integerrimis, inferioribus serratis, caly-
cinis squamis ovatis acutis, exterioribus linearibus di-
varicatis.

In the western parts of Georgia. Lyon. ♀. Aug.—
Oct. v. v. From four to seven feet high; stem dark
purple and very smooth; flowers middle size.

** *Foliis superioribus alternis.*

7. *H. foliis alternis lanceolatis serratis scabris obsolete tri-
plinervibus utrinque attenuatis subsessilibus basi cilia-
tis, calycinis squamis lanceolatis ciliatis.* Willd. sp.
pl. 3. p. 2242.

In dry swamps and on mountain meadows: Canada to
Carolina. ♀. Aug.—Oct. v. v. A tall, rough-
looking plant; horses are fond of its young shoots.
Stems rough and green.

8. *H. foliis alternis ovato-lanceolatis serratis scabris tripli-
nervibus, apice attenuatis petiolatis, petiolis ciliatis,
calycinis squamis lanceolatis ciliatis.* Willd. sp. pl. 3.
p. 2243.

Icon. *Jacq. hort. vind. t. 160.*

In copses and mountain meadows: Pennsylvania to Caro-
lina. ♀. July—Sept. v. v. Resembles the pre-
ceding; stems smooth and purple; the chaff of the
receptacle is green in this species, but black in the pre-
ceding.

9. *H. foliis ovatis acuminatis serratis triplinervibus subtus
scabris, calycinis squamis linear-lanceolatis basi cili-
atis.* Willd. sp. pl. 3. p. 2242.

Icon. *Bocc. sic. t. 27. f. 4.*

In Canada and New England. ♀. Aug.—Oct. v. v.

10. *H. caule debili prostrato, foliis lanceolatis acuminatis
scabris serratis triplinervibus, superioribus integerrimis,
calycinis squamis lanceolatis ciliatis.* Willd. sp. pl. 3.
p. 2242.

In North America. Willdenow. ♀. +.

11. *H. foliis ovatis acuminatis remote serratis triplinervibus
scabris, calycinis squamis lanceolatis subæqualibus*

longifolius.

giganteus.

altissimus.

strumosus.

prostratus.

decapetalus

subciliatis, radiis denis duodenisve.—*Willd. sp. pl. 3. p. 2241.*

Icon. *Rob. ic. 235.*

In dry stony soil, in woods, and on road sides: Canada to Virginia. 4 Aug—Oct. v. v. About three or four feet high. Resembles the following.

- multiflorus.* 12. *H. foliis triplinervibus scabris, inferioribus cordatis, superioribus ovatis, radio multifloro, calycinis squamis lanceolatis.* *Willd. sp. pl. 3. p. 2239.*

Icon. *Bot. mag. 227. Pluk. phytogr. t. 159. f. 2.*

In dry mountain woods: Pennsylvania to Carolina. 4 July—Sept. v. v.

- mollis.* 13. *H. foliis ovatis acuminatis triplinervibus adpresso-serratis supra scabris, subtus cano-pubescentibus mollissimis, calycinis squamis lanceolatis adpressis.* *Willd. enum. 920. Willd. sp. pl. 3. p. 2240.*

H. tomentosus. *Mich. fl. amer. 2. p. 141.*

In exsiccated swamps: Pennsylvania and Virginia. 4 July—Oct. v. v. Leaves very rough on the upper surface, and soft underneath.

- macrophyllus.* 14. *H. foliis ovatis acuminatis triplinervibus serratis supra scabris, subtus cano-pubescentibus, calycinis squamis linearibus squarrosis.* *Willd. hort. berol. 70. cum icono.*

In Pennsylvania and Virginia, on the borders of woods. 4 Aug.—Oct. v. v.

- angustifolius.* 15. *H. caulibus gracilibus subunifloris, foliis linearibus margine revolutis asperis.* *Willd. sp. pl. 3. p. 2244.*

Rudbeckia angustifolia. *Willd. sp. pl. 3. p. 2250.*

Icon. *Mill. ic. t. 224. f. 2.*

In cedar-swamps: New Jersey to Florida. 4 Sept.—Nov. v. v. Rays deep yellow, disk brown.

656. GALARDIA. *Gen. pl. 1323.*

bicolor.

1. *G. caule ramoso, foliis amplexicaulibus lanceolatis inciso-dentatis obtusis, paleis pappi integerrimis aristatis.* —*Willd. sp. pl. 3. p. 2245.*

G. lanceolata. *Mich. fl. amer. 2. p. 142.*

Gaillarda pulchella. *Fougeroux act. par. 1785.*

Colonnea pulcherrima. *Buchoz. ic. t. 126.*

Virgilia helioides. *L'Herit. monog. Smith exot. bot. p. 71. t. 37.*

Leysera
Icon. *L.*
In dry sa
v. v.
showy
dark p
2. G. caule
tis, ca
sp. pl.
In open
+.
3. G. hirsut
lanceol.
ceolatis
On dry hi
v. s. in
Planta h
Rami u
lanceol.
imbrica
tissimis
Corollu
lacinis
tubulos
sutissim
centro s
tis, long

657
1. R. aspera;
remote
acumina
dentibus
Icon. *Bot.*
t. 21. f.
In the m
Oct. v
large, ra
2. R. glabra
caulibus
co.—*W*
R. amplex
p. 29. t.

Leysera caroliniana. *Walt. fl. car.* 211. *secundum Herb.*
Icon. Lam. illustr. t. 708.

In dry sandy soil: Carolina to Florida. ♂. July, Aug.
v. v. About two feet high; flowers large and very
 showy; rays orange with red towards the base, disk
 dark purple with orange.

2. *G. caule simplicissimo unifloro, foliis radicalibus obovatis, caulinis linearibus, paleis pappi laceris.* *Willd. sp. pl. 3. p. 2246.*

In open swamps of Carolina and Florida. *Michaux.*

+

3. *G. hirsutissima, scabra; caule ramoso, foliis linearilanceolatis sessilibus, paleis pappi (5—7) ovato-lanceolatis longissime aristatis.*

On dry hills on the Rocky-mountains. *M. Lewis.* ♂.
v. s. in Herb. Lewis. Flowers orange-coloured.

Planta hirsutissima. Caulis erectus, teres, ramosus.
Rami uniflori. Folia caulina alterna, longe-linearilanceolata, sessilia, scabra, hirsutissima. Calyx subimbricatus, polyphyllus: foliolis lineari-oblongis, acutissimis, disco duplo longioribus, margine coloratis.
Corollæ: Radii circiter 12. obcuneatæ, 3-fidæ:
lacinias oblongis, subæqualibus, obtusiusculis; Disci
tubulosæ, 5-fidæ: lacinias extus purpurascensibus hirsutissimis. Receptaculum in ambitu subpaleaceum,
centro setosum. Pappus: paleis 5—7. scariosis, ovatis, longissime-setaceo-aristatis.

fimbriata.

aristata.

657. RUDBECKIA. *Gen. pl.* 1324.

1. *R. aspera; foliis inferioribus lato-ovatis basi attenuatis remote dentatis, caulinis lanceolato-ovatis utrinque acuminatis subintegerrimis, radiis longissimis dependentibus bifidis.—Willd. sp. pl. 3. p. 2249.*

Icon. Bot. mag. 2. Catesb. car. 2. t. 59. Pluk. alm.
t. 21. f. 1.

In the mountains: Virginia to Florida. ♀. Aug.—Oct. *v. v.* A highly ornamental perennial; flowers large, rays purple, disk brown.

purpurea.

2. *R. glabra; foliis oblongo-lanceolatis cordatis amplexicaulis, inferioribus serratis, disco cylindraceo-conico.—Willd. sp. pl. 3. p. 2249.*

R. amplexicaulis. Vahl in act. soc. nat. scrut. haf. 2.
p. 29. t. 4.

amplexifolia.

- R. perfoliata*. Cav. ic. 3. p. 27. t. 252.
Icon. Jacq. ic. rar. 3. t. 592.
In Lower Louisiana, on the banks of the Mississippi.
♂. July, Aug. v. v. Flowers deep yellow.
- lævigata*.
3. R. undique glaberrima; caule lævigato paniculato, ramis corymbosis, pedunculis elongatis unifloris, foliis ovato-lanceolatis utrinque acuminate triplinervibus integerimis aut dente uno altero instructis lævigatis, calycinis foliolis lanceolatis longitudine radii.
In Georgia. Lyon. ♀. v. s. in Herb. Lambert. Rays pale yellow, short.
- discolor*.
4. R. ramis corymbosis unifloris, pedunculis nudis elongatis, foliis lanceolatis strigoso-pilosis subintegerrimis, calycinis foliolis ovatis acutis, petalis lanceolatis integerrimis discoloribus longitudine calycis.
In Florida. Bartram. ♀. v. s. in Herb. Banks. Flowers small, rays yellow, and deep orange or purple underneath.
- spathulata*.
4. R. gracilis, pubescens; caulis unifloris, foliis obovato-spathulatis integerrimis, calyce patulo imbricato, radiis tridentatis.—Willd. sp. pl. 3. p. 2249.
R. gracilis. Herb. Banks. mss.
- aristata*.
5. R. caule hispido, ramis elongatis corymbosis unifloris, foliis lanceolato-oblongis serratis hispidis, disco subbenispærico, paleis pappi subulatis aristatis.—Herb. Banks. mss.
In the mountains of Carolina. Michaux. In Florida. Bartram. ♂. July, Aug. v. s. in Herb. Banks.
- fulgida*.
6. R. caule hispido, ramis virgatim elongatis unifloris, foliis oblongo-lanceolatis denticulatis hispidis basi angustatis subcordatis, calyce foliaceo radium subæquante, disco hemispærico, paleis lanceolatis.—Willd. sp. pl. 3. p. 2248.
In mountain meadows and woods: Pennsylvania to Carolina. ♀. July—Oct. v. v. Rays dark orange-coloured, disk deep purple.
- hirta*.
7. R. hirsutissima; caulis virgatis subramosis unifloris, pedunculo nudo, foliis ovato-spathulatis triplinervibus serratis hirtis, calyce foliaceo radium subæquante, disco conico, paleis lanceolatis.—Willd. sp. pl. 3. p. 2248.
Icon. Dill. elth. t. 218. f. 285. Pluk. alm. t. 242. f. 2.

- In the Sept. disk
8. R. caul dunc tubet ovati. In Geo
9. R. hisp multitis se aribu p. 22 Icon. I In the Sept. small
10. R. brevis etis si incirc R. trilob In mou Sept. which only:
11. R. caulis ele bus, lindra R. colui On the recep other
12. R. foli ovatis journ Icon. A On the ginia. feet h
13. R. foliis riorib

Mississippi.
low.uniculato, ran-
nifloris, foliis
triplinervibus
ctis lævigatis,
idii.

imbert. Rays

nudis elonga-
tintegerrimis,
ceolatis inte-lanks. Flow-
or purple un-oliis obovato-
nbrocato, ra-
9.In Florida.
b. Banks.osis unifloris,
s, disco sub-
tatis.—Herb.

ks. Flowers

unifloris, fo-
dis basi an-
subæquante,
Willd. sp. pl.ania to Caro-
k orange-co-sis unifloris,
riplinervibus
quante, disco
3. p. 2248.
. t. 242. f. 2.

In the mountains: Virginia to Florida. ♂. July—
Sept. v. v. Resembles the preceding; rays yellow,
disk dark brown.

8. R. caule inferne hispido, superne glabro nudiusculo, pe-
dunculis longissimis unifloris, foliis ovatis attenuatis
tuberculatis hispidis, calycibus imbricatis, squamis
ovatis acuminatis ciliatis.

In Georgia. Bartram. ♂. v. s. in Herb. Banks.

9. R. hispido-pilosa; caule paniculato, ramis divaricatis
multifloris foliosis, foliis lanceolatis utrinque acumina-
tis serratis, inferioribus trilobis, calycinis foliolis line-
aribus deflexis longitudine radii.—Willd. sp. pl. 3.
p. 2247.

Icon. Pluk. alm. 22. f. 2.

In the Virginia and Carolina mountains. ♀. ♂. Aug.
Sept. v. v. About four or five feet high; flowers
small, pale yellow.

10. R. brevi pubescens subtomentosa; caule ramoso, ra- *subtomentosa*.
mis erectis multifloris, foliis oblongo-lanceolatis acu-
tis subserratis, inferioribus trilobis, calycinis foliolis
incumbentibus radio brevioribus.

R. triloba B. Mich. fl. amer. 2. p. 144.

In mountain meadows: Virginia and Illinois. ♀. Aug.
Sept. v. v. Not near so high as the preceding, from
which it is very distinct, though Michaux considers it
only a variety.

11. R. caule stricto simplici summitate paucifloro, peduncu-
lis elongatis, foliis pinnatifidis incisis, lacinias linearis-
bus, calyce simplici 5-phyllo, radius 5—8., disco cy-
lindraceo elongato.

R. columnifera. Fraser catal. 1813.

On the Missouri. v. s. The singular appearance of the
receptacle of this plant distinguishes it from all the
other known species.

12. R. foliis inferioribus pinnatis, pinnis 3-lobis, summis
ovatis, pappo crenato, caule glabro. Schrad. new.
journ. 2. p. 61. Willd. enum. 921.

Icon. Moris. hist. 3. s. 6. t. 6. f. 53. Corn. canad. t. 179.

On the edges of swamps and ditches: Canada to Vir-
ginia. ♀. Aug.—Oct. v. v. From five to ten
feet high.

13. R. foliis inferioribus pinnatis, pinnis pinnatifidis, supe-
rioribus simplicibus pinnatis, summis 3-fidis, pappo

Radula.

triloba.

columnaris.*laciñata*.*digitata*.

crenato, caule lœvi. *Schrad. new. journ.* 2. p. 61.
Willd. enum. 921.

Icon. Moris. hist. 3. s. 6. t. 6. f. 54.

In the mountains of Virginia and Carolina. ♀. Aug.—
 Oct. v. v. Resembles the preceding.

pinnata.

14. R. foliis omnibus pinnatis, piona una alterave inferiorum bipartita, reliquis indivisis, pappo integerrimo, caule sulcato hispido. *Schrad. new. journ.* 2. p. 61.
Willd. enum. 921. *Mich. fl. amer.* 2. p. 144.

R. digitata. *Willd. sp. pl.* 3. p. 2247. excl. syn. *Morrison. et Aiton.*

R. odorata. *Hortul.*

Icon. Smith exot. bot. 38. *Vent. jard. cels* 71.

In the western parts of Carolina and Georgia. ♀. July
 —Oct. v. v. Flowers very showy, of an anise scent; rays long, bright yellow, hanging downwards; disk ovate, purple.

658. CENTAUREA. *Gen. pl.* 1331.

Cyanus.

1. C. squamis calycinis serratis, foliis linearibus integerimis, inferioribus dentatis. *Smith fl. brit.* 2. p. 911.
Willd. sp. pl. 3. p. 2291.

Icon. Engl. bot. 277. *Fl. dan.* 993.

In rye and wheat fields; rare. Brought from Europe with the grain. ♂. July, Aug. v. v. Flowers blue.

solstitialis.

2. C. calycibus palmato-spinosis solitariis, spinis rectis, foliis canescentibus linear-lanceolatis decurrentibus integerrimis, radicalibus lyratis. *Willd. sp. pl.* 3. p. 2309.

Icon. Engl. bot. 243.

In sandy fields and woods: New Jersey: introduced from Europe. ♂. July, Aug. v. v. Flowers yellow.

3

1. C. foliis
mento
l. c. ci

Tussilag-

Perdicut-

In sandy

Florida-

a tinge

1. S. caule
pinnat-
subcor-

Icon. Li

On the t

cularly

From

all the

2. S. caule
libus t-
culatis

S. lacinia

S. sinuat-

In gravel

rolina.

high;

3. S. caule
alterni-

p. 233

Icon. Ja

In the w

Oct.

large.

4. S. caule
serrati-

β . S. squar-

longio-

In the

IV. NECESSARIA.

659. CHAPTLIA. *Vent. hort. cels* 61.

1. *C. foliis ovato-oblongis integerrimis subitus argenteo-tomentosis, scapo nudo unifloro, flore nutante.* — *Vent. l. c. cum icon.*

Tussilago integrifolia. *Mich. fl. amer.* 2. p. 121.

Perdicium semiflosculare. *Walt. fl. car.* 204.

In sandy fields and woods, in moist soil : Carolina to Florida. 4. Aug., June. v. v. Flowers white, with a tinge of pale purple.

660. SILPHIUM. *Gen. pl.* 1334.

1. *S. caule superne hispido, foliis radicalibus caulinisque pinnatifidis, lacinia dentato-sinuatis, calycinis foliolis subcordatis acuminatis.* — *Willd. sp. pl.* 3. p. 2330.

Icon. Linn. fil. fasc. 1. t. 3.

On the banks of rivers in the western territories, particularly on the Mississippi. 4. Aug.—Oct. v. v. From eight to twelve feet high ; flowers large, and, as all the rest of the genus, yellow.

2. *S. caule lævi, foliis caulinis sinuato-pinnatifidis, radicalibus ternatis simplici- multifidis, floribus parvis paniculatis.* — *Willd. sp. pl.* 3. p. 2331.

S. laciniatum. *Walt. fl. car.* 217.

S. sinuatum. *Herb. Banks, mss.*

In gravelly woods, near the sea-coast : Virginia and Carolina. 4. Aug.—Oct. v. v. Not above two feet high ; flowers small.

3. *S. caule lævi, foliis radicalibus amplis cordatis, caulinis *terebinthina-* alternis ovatis serratis scabris.* — *Willd. sp. pl.* 3. p. 2331.

Icon. Jacq. hort. 1. t. 43.

In the western mountains and Louisiana. 4. Aug.—Oct. v. v. About five feet high ; radical leaves very large.

4. *S. caule tetragono lævi, foliis oppositis connatis ovatis *perfoliatum,* serratis.* — *Willd. sp. pl.* 3. p. 2331.

- β. *S. squamis calycinis quatuor exterioribus calyce interiore conjunctum.* longiore. *Willd. enum.* 933.

In the Allegany mountains : Pennsylvania to Carolina.

- connatum.* 5. *S. caule tereti hispido, foliis oppositis connatis remote serratis scabris.*—*Willd. sp. pl. 3. p. 2332.*
In high mountains of Virginia and Carolina. $\frac{4}{4}$. Aug.
Sept. *v. v.* Resembles the preceding, but is a very distinct species.
- Asteriscus.* 6. *S. caule tereti hispido, foliis oppositis alternis oblongis acutis serratis scabris, calycibus ciliatis.*—*Willd. sp. pl. 3. p. 2332.*
Icon. Dill. elth. t. 37. f. 42.
In mountain meadows: Virginia and Carolina. $\frac{4}{4}$. July—Sept. *v. v.*
- pumilum.* 7. *S. caule tenuiter tomentoso, foliis ovalibus leviter serratis obtusis subtomentosis, calycinis foliolis obovatis obtusis, seminibus muticis.*—*Willd. sp. pl. 3. p. 2332.*
S. patulum. Herb. Banks.
In Florida. *Michaux. Bartram.* $\frac{4}{4}$. *v. s. in Herb. Banks.*
- integrifolium.* 8. *S. caule tetragono aspero, foliis oppositis sessilibus oblongis integerrimis scabris, floribus paucis breviter pedunculatis.*—*Willd. sp. pl. 3. p. 2333.*
On the most western of the Allegany mountains and Illinois. $\frac{4}{4}$. Aug.—Oct. *v. v.* About four feet high.
- lævigatum.* 9. *S. caule simplici tetragono sulcato glabro, foliis oppositis sessilibus ovatis acuminatis tenuissime serratis basi subcordatis utrinque glabris, calycinis squamis ovatis ciliatis.*
In Georgia. *Enslen. $\frac{4}{4}$. v. s. in Herb. Enslen.* About two feet high; flowers in a close corymb.
- trifoliatum.* 10. *S. caule sexangulo lœvi, foliis terno-verticillatis ovato-lanceolatis inæqualiter dentato-serratis scabris, superioribus sessilibus, panicula trichotoma.*—*Willd. sp. pl. 3. p. 2333.*
S. ternifolium. Mich. fl. amer. 2. p. 146.
Icon. Moris. hist. 3. s. 6 t. 3. f. 68.
On the high mountains of Virginia and Carolina. $\frac{4}{4}$. Sept. Oct. *v. v.* About six feet high.
- ternatum.* 11. *S. caule tereti lœvi, foliis terno-verticillatis petiolatis lanceolatis subdenticulatis scabriusculis basi ciliatis, superioribus sparsis sessilibus, panicula dichotoma, calycibus ciliatis.*—*Willd. sp. pl. 3. p. 2333.*

In North
bert.12. *S. caule*
ceolatis
tis, sup
sp. pl. 3.
In Carolin
Resembl
distingui13. *S. caule*
alternis
nibus n
In Georgi14. *S. foliis al*
cinis ob
In Carolin15. *S. foliis alt*
usculis v
In Florida.

661

1. *P. viscidob*
ribus pi
Willd. s
Icon. Ama
In shady i
July. *v*
the who2. *P. foliis o*
bus, lob
sp. pl. 3.
P. maculat
Icon. Plu
f. 55. 1
In the mou
Sept. *v*

662. C

1. *C. pumilu*
tatis oval
Icon. Pluk
On dry hill
v. v. N

- In North America. *Retz.* \mathcal{U} . *v. s. in Herb. Lambert.*
12. *S. caule tereti laevi, foliis subquaterno-verticillatis lanceolatis scabris subintegerrimis subsessilibus basi ciliatis, superioribus sparsis, panicula dichotoma.* — *Willd. sp. pl. 3. p. 2334.*
- In Carolina and Georgia. *Lyon.* \mathcal{U} . Aug. Sept. *v. v.*
Resembles the preceding; but its purple smooth stem distinguishes it at first sight.
13. *S. caule petiolisque tomentosis, ramis unifloris, foliis alternis cordatis ovatis serratis petiolatis villosis, seminibus muticis.* — *Herb. Banks. mss.*
- In Georgia. *Bartram.* \mathcal{U} . *v. s. in Herb. Banks.*
14. *S. foliis alternis petiolatis cordatis sinuatis, squamis calycinis obtusis.* — *Herb. Banks. mss.*
- In Carolina. \mathcal{U} . *v. s. in Herb. Banks.*
15. *S. foliis alternis ovato-lanceolatis cordatis serratis obtusiusculis villosiusculis.* — *Herb. Banks. mss.*
- In Florida. \mathcal{U} . *v. s. in Herb. Banks.*

*atropurpureum.*661. POLYMNIA. *Gen. pl. 1335.*

1. *P. viscidó-villosa; foliis denticulatis acuminatis, inferioribus pinnatifidis, superioribus trilobis integris.* — *Willd. sp. pl. 3. p. 2335.*
- Icon. *Amaen. acad. 3. t. 1. f. 5.* *
- In shady mountains: Canada to Virginia. \mathcal{U} . June, July. *v. v.* Two or three feet high; flowers yellow; the whole plant has a strong balsamic scent.
2. *P. foliis oppositis trilobis acutis in petiolum decurrentibus, lobis anguloso-sinuatis, radiis elongatis.* — *Willd. sp. pl. 3. p. 2335.*
- P. maculata.* *Cav. ic. 3. t. 227.*
- Icon. *Pluk. alm. t. 83. f. 3. Meris. hist. 3. s. 6. t. 7. f. 55. Trew pict. suppl. 112.*
- In the mountains of Virginia and Carolina. \mathcal{U} . July—Sept. *v. v.* From four to eight feet high.

Uvedalia.

662. CHRYSOGONUM. *Gen. pl. 1337.*
1. *C. pumilum, lanato-villosum; foliis in petiolum angustatis ovalibus dentatis.* — *Willd. sp. pl. 3. p. 2337.*
- Icon. *Pluk. alm. t. 83. f. 4. & t. 242. f. 3.*
- On dry hills: Virginia and Carolina. \mathcal{U} . May, June. *v. v.* Not above a span high; flowers yellow.

663. BALTIMORA. *Gen. pl. 1333.**recta.*

1. *B. scabra*; caule dichotomo, foliis oppositis ovatis acuminatis serratis petiolatis.—*Willd. sp. pl. 3. p. 2329.*
 Icon. *Pluk. mant. t. 342. f. 3.*
 In Maryland near Baltimore. *Sp. pl. ♂. July. v. v.*
cult. Flowers small, yellow. I have never seen this plant in any part of the United States, and suppose it to be only an inhabitant of Vera Cruz.

664. PARTHENIUM. *Gen. pl. 1428.*

- integerrifolium.* 1. *P. foliis oblongis inaequaliter dentatis asperis, superioribus amplexicaulibus.*—*Willd. sp. pl. 3. p. 2385.*
 Icon. *Willd. hort. berol. t. 4. Dill. elth. t. 225. f. 292.*
Pluk. alm. t. 53. f. 5. ♂ t. 219. f. 1. Lam. illustr. t. 766.
 In dry places on the mountains of Virginia and Carolina.
♀. Sept. Oct. v. v. About a foot high; flowers white.

665. IVA. *Gen. pl. 1429.**ciliata.*

1. *I. caule herbaceo, foliis lanceolato-ovatis sub serratis, spica confertiuscula, bracteis lanceolatis acuminatis petiolatis que longe ciliatis.*—*Willd. sp. pl. 3. p. 2386.*
 In natural meadows: Kentucky and Illinois. ♂. July, Aug. v. s. About two feet high.

imbricata.

2. *I. fruticosa, glabra; foliis linear-lanceolatis cuneatis integrerrimis glabris, calycibus imbricatis.*—*Willd. sp. pl. 3. p. 2387. Walt. fl. car. 232.*
I. integrifolia. Herb. Banks. mss.
 On the sea-coast of Carolina and Georgia. ½. v. s. in *Herb. Walter.*

frutescens.

3. *I. fruticosa; foliis lanceolatis punctato-scabris profunde serratis, glomerulis florum depresso-globosis.*—*Willd. sp. pl. 3. p. 2387.*
 Icon. *Lam. encycl. 766. Pluk. alm. t. 27. f. 1.*
 On the sea-coast: New England to Florida. ½. Aug. Sept. v. v. A shrub about four feet high.

666. AMBROSIA. *Gen. pl. 1427.**integerrifolia.*

1. *A. foliis ovatis sessilibus acuminatis serratis utrinque hispidis basi ciliatis, racemis terminalibus plerumque ternis.*—*Willd. sp. pl. 4. p. 375.*

On the
Virginia
this g

2. *A. hirsuta*
gis su
tu tet
amer.

In the
July—

3. *A. hirsuta*
ceolat
Willda

Icon. M
On the
rolina
feet h

4. *A. foliis*
tis, ra
Willda

In old fi

5. *A. foliis*
nativid
Willda

In fields
♂. A

6. *A. foliis*
mis te
pl. 4. p

Icon. P
In old fi
July—

7. *A. foliis*
sumin
racem
Willda

On the b
♂. Ju

667.
1. *X. caule*
scabris
bus set

13. *is ovatis acutis. p. 2329.*
July. *v. v.* ever seen this
and suppose it

28. *is, superiori-
t. 2385.*
l. 225. f. 292.
Lam. illustr.
and Carolina.
high; flowers

serratis, spica
inatis petiolatis.
386. *is. ⊖ July,*

is cuneatis in-
i.—Willd. sp.
h. v. s. in
abris profunde
osse.—Willd.
f. 1.
la. h. Aug.
1.

7. *utrinque his-
lerumque ter-*

On the banks of ponds and ditches: Pennsylvania and Virginia. ⊖ July—Sept. *v. v.* The flowers of this genus are very inconspicuous.

2. A. hirsutissima; foliis arte sessilibus simplicibus oblongis supra basin latescentem utrinque unidentatis, fructu tetragono infra apicem quadrispinoso. *Mich. fl. amer. 2. p. 182.*

In the western parts of the Allegany mountains ⊖ July—Sept. *v. v.*

3. A. hirsuta, aspera; foliis 3-lobis serratis, lobis ovali-lanceolatis acuminatis, fructu infra apicem 6-spinoso.—*Willd. sp. pl. 4. p. 375.*

Icon. *Moris. hist. 3. s. 6. t. 1. f. 4.*
On the banks of rivers and ditches: Pennsylvania to Carolina. ⊖ Aug. Sept. *v. v.* From five to eight feet high.

4. A. foliis bipinnatifidis glabriusculis, petiolis longe ciliatis, racemis terminalibus paniculatis, caule virgato. *Willd. sp. pl. 4. p. 376.*

In old fields: Canada to Carolina. ⊖ June—Aug. *v. v.*

5. A. foliis bipinnatifidis subtus canescensibus summis pin- natifidis, racemis tert. is terminalibus, ramis fastigiatis. *Willd. sp. pl. 4. p. 370.*

In fields and waste grounds: Pennsylvania to Carolina. ⊖ Aug. Sept. *v. v.*

6. A. foliis glabris bipinnatifidis summis pinnatifidis, race- mis terminalibus solitariis, ramis fastigiatis. *Willd. sp. pl. 4. p. 376.*

Icon. *Pluk. alm. t. 10. f. 5.*

In old fields and on road sides: Canada to Florida. ⊖ July—Sept. *v. v.*

7. A. foliis caulinis pinnatifidis subdentatis petiolatis, rameis *heterophylla.* suminis lanceolatis sessilibus, petiolis longe ciliatis, racemis terminalibus solitariis, caule paniculato. *Willd. sp. pl. 4. p. 378.*

On the banks of rivers and ditches: near Philadelphia. ⊖ July—Sept. *v. v.*

667. XANTHIUM. Gen. pl. 1426.

1. X. caule inermi ramoso, foliis cordatis lobatis serratis *Strumarium.* scabris basi trinervibus, fructibus ellipticis pubescenti- bus setis rigidis uncinatis.—*Willd. sp. pl. 4. p. 373.*

bidentata.

trifida.

elatior.

paniculata.

heterophylla.

Icon. *Fl. dan.* 970.

On road sides, among rubbish and cultivated grounds :
New England to Pennsylvania. ♂. Aug. Sept. v. v.
The fruit resembles the common *Burdock*.

XV

V. S E G R E G A T A.

668. ELEPHANTOPUS. *Gen. pl.* 1347.

- carolinianus.* 1. E. foliis radicalibus caulinisque oblongis basi angustatis utrinque pilosis, caule simplici erecto piloso.—*Willd. sp. pl.* 3. p. 2390.
E. scaber. *Mich. fl. amer.* 2. p. 148.
Icon. *Dill. elth. t.* 106. f. 126. *Sloan. hist.* 1. t. 156. f. 1.
In shady woods: Virginia to Florida. ♀. Aug.—Oct. v. v. Flowers red.
- tomentosus.* 2. E. foliis radicalibus caulinisque ovatis basi attenuatis utrinque tomentosis, caule simplici tomentoso.—*Willd. sp. pl.* 3. p. 2390.
On dry sunny hills: Lower Virginia and Carolina. ♀. Aug.—Oct. v. v. Flowers deep red.

* *Caly*

669. THALI
duo int
mento
anthera
1-sper

670. CANNA
Nectar.
rollæ &
sperma

** *Ecalycula*

671. ORCHIS
Labellu
catum.

672. NEOTTI
rioribus
nexis.

673. ARETHU
tibus.
persiste

674. EPIPACT
ratum,
opercula

675. MALAXI
cavo-patu

676. CYMBID
basi conc
opercula

VOL. II.

ed grounds :
Sept. v. v.

XVIII. GYNANDRIA.

I. MONANDRIA.

* *Calyculatæ; filamentum basi styli insertum.*

669. THALIA. *Cal. minimus, triphyllus. Cor. 5-petala: duo interiora minora. Anthera simplex, ovata, filamento proprio depresso innixa. Stylus simplex, ab anthera deflexus. Stigma perforatum ringens. Baccæ 1-sperma.*
670. CANNA. *Cal. 3-phyllus. Cor. 6-partita, erecta. Nectar. bipartitum revolutum. Styl. lanceolatus, corollæ adnatus. Caps. muricata, 3-locularis, polysperma.*
- ** *Ecalyculatæ; filamentum nullum; anthera apice styli inserta, biloba.*
671. QRCHIS. *Cor. ringens: petalo superiore fornicato. Labellum dilatatum, basi subtus calcaratum aut sacrum. Anthera terminalis, adnata.*
672. NEOTTIA. *Cor. ringens: petalis lateralibus exterioribus antice circa basin labelli ventricosum conexis. Anthera stylo parallela, postice inserta.*
673. ARETHUSA. *Cor. subringens: petalis subconniventibus. Labellum ecalcaratum. Anthera opercularis, persistens. Pollen pulvereo-granulatum.*
674. EPIPACTIS. *Cor. erecto-patens. Labellum ecalcaratum, planum, pendulum, apice bifidum. Anthera opercularis, persistens.*
675. MALAXIS. *Cor. patens, resupinata. Labellum concavo-patulum, adscendens. Anthera opercularis.*
676. CYMBIDIUM. *Cor. erecta vel patens. Labellum basi concavum, ecalcaratum, lamina patula. Anthera opercularis, decidua. Pollen globosum.*

677. CALYPSO. *Cor. 6-petala: 5. sursum expansa. Labellum sacciforme, dorso basis 2-calcaratum. Stylus late alatus. Stigma operculo 2-lobo absconditum. Pollen lamellis 4. solidis.*

II. DIANDRIA.

678. CYPRIPEDIUM. *Cor. 4-petala, patens. Labellum ventricoso-inflatum, obliquum. Stylus superne lobo petaloideo appendiculatus.*

III. HEXANDRIA.

679. ARISTOLOCHIA. *Cor. 1-petala, tubuloso-ligulata, basi ventricosa. Styl. 0. Caps. infera, 6-locularis.*

IV. DODECANDRIA.

680. ASARUM. *Cor. tubuloso-campanulata, 3-4-fida. Antheræ mediis filamentis adnatæ. Caps. infera, coronata, 6-locularis.*

1. C. foliis j
teriore
pl. 1. p
C. glauca.
In deep
Enslen.
ers scar
bably m
gustifoli
2. C. foliis s
teriore
trans. 8.
C. glauca
C. angustif
Icon. Salis
douté lili
In swamp
Flowers
671. ORC

1. O. labello c
plo longi
p. 8.
Icon. Bot.
In old mead
rolina.
golden-ye
2. O. labello la
cornu ger
Orchis testic
In low swan
June, July
the preced
3. O. labello 3.
obtusis, c
longitudin
In old fields
nia. 4.
4. O. labello ol
binis later
Willd. sp.

dealbata.

I. MONANDRIA.

669. THALIA. *Gen. pl. 10.*

1. T. panicula albido-pulverulenta, spathis bifloris, foliis ovatis apice revolutis.—*Roscoe in linn. trans. 8. p. 340.*
Icon. *In a single plate of J. Fraser.*
In impenetrable swamps of South Carolina. 4. Aug.
Sept. v. v. s. fl.; v. s. c. fl. in *Herb. Lambert.* A tall and handsome plant; flowers small, purple. J. Millington, Esq. of South Carolina, is, to my knowledge, the first discoverer of this elegant plant. It was introduced into the English gardens by the Messrs. Frasers.

- expansa.
um. *Sty-*
conditum.
1. *C. foliis petiolatis lanceolatis costatis, corollæ limbo in-* *angustifolia.*
teriori trifido : laciniis strictis emarginatis.—Willd. sp.
pl. 1. p. 3?

C. glauca. Walt. fl. car. 59.
In deep and almost impregnable swamps of Georgia.
Enslen. 4. July, Aug. v. s. in Herb. Enslen. Flowers scarlet. It approaches near to *C. indica*, and probably may be a new species, if not the true *C. angustifolia*.

2. *C. foliis sessilibus lanceolatis glaucis, corollæ limbo in-* *flaccida.*
teriori trifido : laciniis flaccidis.—Roscoe in linn.
trans. 8. p. 339.

C. glauca β. flaccida. Willd. sp. pl. 1. p. 4.
C. angustifolia. Walt. fl. car. 59.
Icon. Salisb. stirp. rar. 3. t. 2. Thoms. bot. t. 1. Re-
douté liliac. 107.

In swamps of South Carolina. 4. Aug. Sept. v. v.
Flowers large, yellow.

671. ORCHIS. *Swartz act. holm. 1800, p. 205.*

* *Radicibus testiculatis.*

1. *O. labello oblongo-lanceolato pinnatum ciliato petalis du-* *ciliaris.*
plo longiore, cornu germe longiore. Willd. sp. pl. 4.
p. 8.

Icon. Bot. rep. 42.
In old meadows and exsiccated swamps: Canada to Carolina. 4. June, July. v. v. Flowers a bright golden-yellow, very handsome.

2. *O. labello lanceolato ciliato longitudine petali supremi, blephariglot-* *tis.*
cornu germe longiore. Willd. sp. pl. 4. p. 9.

Orchis testiculata floribus niveis, &c. Clayt. no. 560.
In low swampy bottoms: New Jersey to Carolina. 4.
June, July. v. v. Flowers clear white; resembles the preceding very much.

3. *O. labello 3-partito, laciniis capillaceo-multifidis, petalis* *psycodes.*
obtusis, cornu filiformi-clavato adscendente germinis
longitudine. Willd. sp. pl. 4. p. 39.

In old fields, on the side of swamps: Canada to Virginia. 4. June—Aug. v. s. Flowers yellow.

4. *O. labello oblongo pinnatum-ciliato, petalis rotundatis,* *cristata.*
binis lateralibus dentatis, cornu germe breviore.
Willd. sp. pl. 4. p. 9. Mich. fl. amer. 2. p. 156.

Labellum
perne lobo

io-ligulata,
-locularis.

3-4-fida.
ips. infera,

floris, foliis
i. 8. p. 340.

4. Aug.
imbert. A
all, purple.
is, to my
gant plant.
ens by the

- In exsiccated swamps and mountain meadows: Pennsylvania to Carolina. ♀. June, July. v. v. In colour and shape resembling No. 1; but the flowers are smaller and closer, in a short spike.
- lacera.* 5. *O. labello 3-partito, laciniis subdigitato-filiformibus, cornu germen subæquante, floribus alternis.* —*Mich. fl. amer. 2. p. 156.*
- O. radice palmata, foliis lilii, &c.* *Clayt. n. 644.*
In low meadows: Pennsylvania to Virginia. ♀. July. v. v. Flowers greenish-white. Roots not known.
- clavellata.* 6. *O. labello ovato integerrimo, petalis conniventibus, cornu clavato longitudine germinis, caule unifoliato.* *Willd. sp. pl. 4. p. 10.* *Mich. fl. amer. 2. p. 155.*
In Carolina. *Michaux.* ♀. +. A loose spike, with few flowers; bractes very short.
- quinqueseta.* 7. *O. labello 5-partito: laciniis setaceis, cornu germine duplo longiore.* —*Mich. fl. amer. 2. p. 155.*
In sandy low fields, on the side of swamps: Virginia and Carolina. ♀. June, July. v. v. A loose spike, with alternate and distant flowers; bractes acuminate; leaves ovate, acute.
- discolor.* 8. *O. labello petalis longiore tripartito: laciniis lateralibus brevibus acutis, intermedia producta spathulata, cornu filiformi germine subduplo-longiore, folio unico radi cali ovato-cordato.*
Orchis floribus sparsis, nectario pedunculum superante, &c. *Gron. virg. 137.*
In pine-barrens: New Jersey to South Carolina. ♀. July. v. v.
Bullis testiculatis, ovatis, acutis. Folium unicum, ovatum, petiolatum, plicato-nervosum, glabrum, subtus purpurascens. Petiolus margine plicato-undulatus. Scapus pedalis, nudus, vagina una alterave instructus, teres, glaber. Flores laxe spicati, brevisime pedicellati, nutantes, tristi- seu exviridi-purpurentes. Petala oblonga, acutiuscula, flexuosa. Labellum petalis longius, margine subundatum. Cornu ascendentis, filiforme, flexuose, purpureum.
- ** *Radicibus palmatis.*
- flava.* 9. *O. labello trifido integerrimo: lacinia media majore, cornu filiformi longitudine germinis, spica elongata congesta, bracteis flore longioribus.* —*Willd. sp. pl. 4. p. 33.*
- In Virgin yellow.
10. *O. labello cornu ribus.* *Satyrion Icon. Fl.*
In dry grass and Car three inc
11. *O. labello cornu longiore*
On the high June, J small w
12. *O. labello petalis su cornu ob ribus pat*
Icon. Parc
O. foliis d
Gron. vi
In low sha Aug. v
13. *O. labello cornu ob scapo nu*
Serapias foli
Gron. vir
In low sha ♀. July,
14. *O. labello o subulato*
Icon. Gmel
On grassy h ers in lon very long
15. *O. labello o lateralibus viore, bra*
sp. pl. 4.

s : Pensyl-
In colour
lowers are
liformibus,
vis.—Mich.

44.
4. July.
known.

ntibus, cor-
unifoliato.
p. 155.
spike, with

au germe

s : Virginia
loose spike,
ctes acumi-

s lateralibus
culata, cornu
unico radi-

n superante,
rolina. 4.

m unicum,
glabrum,
dicato-undu-
ana alterave
icati, brevis-
viridi-purpu-
exuosa. La-
um. Cornu
um.

edia majore,
ica elongata
llid. sp. pl. 4.

- In Virginia. 4. July. v. s. in Herb. Gronov. Flowers yellow.
10. O. labello linearis apice tridentato, petalis conniventibus, cornu obtuso scrotiformi, bracteis flore sesquilongoribus. Willd. sp. pl. 4. p. 33. *Satyrium viride*. Sp. pl. 1357. Icon. Fl. dan. 77. Hall. helv. t. 26. In dry grassy places, on the high mountains of Virginia and Carolina. 4. June, July. v. v. Not above three inches high; flower small, greenish-white. *viridis.*
11. O. labello lanceolato apice tridentato, petalis obtusis, cornu filiformi apice clavato-descendente germine longiore. Willd. sp. pl. 4. p. 41. On the high mountains of Pennsylvania and Virginia. 4. June, July. v. v. Spike consisting of but a few small white flowers. *tridentata.*
12. O. labello linearis apice emarginato obsolete 3-dentato, petalis subconniventibus, lateralibus ovatis latioribus, cornu obtuso scrotiforme, bracteis flore duplo longioribus patentibus. Willd. sp. pl. 4. p. 34. Icon. Parad. lond. 110. O. foliis duobus inferioribus ovatis, superioribus, &c. Gron. virg. 136. In low shady places: Pennsylvania to Virginia. 4. July, Aug. v. v. About a span high; flowers green. *bracteata.*
13. O. labello oblongo lanceolato indiviso, petalis erectis, cornu obtuso scrotiformi, germinibus pedicellatis, scapo nudo. Willd. sp. pl. 4. p. 34. Serapias foliis ovatis radicalibus, scapo nudo multifloro. Gron. virg. 137. In low shady places: New England to Virginia; rare. 4. July, Aug. v. v. The bractes are very short. *olsoleta.*
- *** *Radicibus fasciculatis.*
14. O. labello ovato basi dentato, petalis patentibus, cornu subulato germinis longitudine. Willd. sp. pl. 4. p. 35. Icon. Gmel. fl. sib. 1. p. 20 t. 4. f. 2. On grassy hills: Pennsylvania. 4. July. v. v. Flowers in long spikes, dark or brownish yellow; bractes very long, acuminate, and longer than the flowers. *fuscescens.*
15. O. labello obovato indiviso crenato retuso, petalis rectis, lateralibus longioribus, cornu clavato germine breviore, bracteis flore longioribus, caule aphylo. Willd. sp. pl. 4. p. 36. *spectabilis.*

- O. humilis. Mich. fl. amer. 2. p. 155.*
In shady, rocky situations: New York to Carolina;
particularly on the mountains. \mathcal{U} . May, June. *v. v.*
About a span high; flowers large, purple and white,
very handsome.
- orbiculata.* 16. *O. labello linearis integerrimo obtusiusculo, petalis 3. su-*
perioribus conniventibus, 2. lateralibus patentibus
basi obliquis, cornu germine longiore, scapo basi di-
phylo, foliis planis orbiculatis.
In shady beech-woods: on the mountains of Pennsylvania
and Virginia. \mathcal{U} . July, Aug. *v. v.* Resembles
O. bifolia. Two leaves of a fleshy texture are spread
flat on the ground, between which rises the stalk about
a foot or eighteen inches high, which bears a loose
spike of greenish-white flowers. It is known in the
mountains by the name of *Heal-all*.
- dilatata.* 17. *O. labello linearis integerrimo obtusiusculo, basi subro-*
tundato-dilatato, cornu longitudine labelli: germine
breviore, bracteis longitudine florum, caule folioso.
O. acuta. Herb. Banks. mss.
In Labrador. *Colmaster.* \mathcal{U} . *v. s. in Herb. Dickson.*
- virescens.* 18. *O. labello lanceolato crenato, petalis conniventibus, cor-*
nu obtuso scrotiformi, bracteis flore longioribus. Willd.
sp. pl. 4. p. 37.
In boggy meadows: Pennsylvania. \mathcal{U} . July. *v. s.* From
a foot to eighteen inches high; flowers green.
- hyperborea.* 19. *O. labello lanceolato magnitudine petalorum patentium,*
cornu subulato germine breviore, spica ovata, bracteis
longitudine florum.—Willd. sp. pl. 4. p. 37.
Icon. Retz obs. bot. 4. t. 3.
In Hudson's Bay, near Albany. *Hutchinson.* \mathcal{U} . June,
July. *v. s. in Herb. Banks.* Flowers greenish-yellow.
- obtusata.* 20. *O. labello linearis integerrimo cornu longiore, cornu lon-*
gitudine germinis, folio unico radicali subcuneiformi-
obtuso.—Herb. Banks. mss.
On Hudson's Bay, near Fort Albany. *Hutchinson.* \mathcal{U} .
v. s. in Herb. Banks. A small species, with a few
flowers only.
- rotundifolia.* 21. *O. labello 3-fido; intermedio bifido, cornu germine*
breviore, foliis ovali-subrotundis.—Herb. Banks. mss.
On Hudson's Bay. \mathcal{U} . *v. s. in Herb. Banks.*
- fimbriata.* 22. *O. labello tripartito, lacinias cuneiformibus ciliato-fim-*
biatis, petalis lateralibus ovatis dentatis, cornu fili-
- formi
p. 39.
In low m
Pensyl
high;
23. *O. labell*
intern
dentat
tudine.
In low r
v. v.
paler ir
like the
exceedi
24. *O. labell*
termedi
dente g
In mount
July, A
dark pu
672. NEC
1. *N. foliis r*
spiralite
lato.—*L*
Satyrium s
Ophrys æs
Limodorus
In low spo
June, Ju
flowers
known t
2. *N. foliis l*
oblonga
oblongo
Ophrys cer
Limodorus
In grassy a
July, Au
spikes ar
3. *N. foliis r*
vaginato
labello pe
Satyrium re

Carolina;
une. v. v.
and white,

etalis 3. su-
patentibus
ipo basi di-

f Pensyla-
Resembles
e are spread
: stalk about
ars a loose
own in the

basí subro-
i: germiné
e folioso.

rb. Dickson.
ntibus, cor-
bus. Willd.

v. s. From
en.

patentium,
ata, bracteis
7.

i. 4. June,
ish-yellow.

cornu lon-
cuneiformi-

hinson. 4.
with a few

iu germiné
Banks. mss.
ks.

ciliato-fim-
cornu fili-

formi clavato germine longiore. Willd. sp. pl. 4.
p. 39.

In low meadows and on high mountain bogs: Canada to Pennsylvania. 4. July. v. v. From a foot to two high; flowers purple, very handsome.

23. O. labello tripartito, laciniis cuneiformibus inciso-dentatis, intermedia emarginata, petalis lateralibus obtusis sub-dentatis, cornu subulato adscendente germinis longitudine. Willd. sp. pl. 4. p. 40.

In low meadows: New York to Virginia. 4. July. v. v. Resembles the preceding; flowers smaller and paler in colour. In the mountain-meadows it grows, like the following, to a very great height, sometimes exceeding three and four feet.

24. O. labello tripartito, laciniis cuneiformibus dentatis, intermedio bilobo, cornu filiformi apice clavato adscendente germine longiore. Willd. sp. pl. 4. p. 40.

In mountain meadows: Pennsylvania to Virginia. 4. July, Aug. v. v. Resembles the preceding; flowers dark purple.

672. NEOTTIA. Swartz. act. holm. 1800. p. 224.

1. N. foliis radicalibus linearibus, scapo vaginato, floribus spiraliter-secundis, labio trifido; medio majori crenulato.—Willd. sp. pl. 4. p. 74.

tortilis.

Satyrium spirale. Swartz. prodr. 118.

Ophrys æstivalis. Mich. fl. amer. 2. p. 157.

Limodorum præcox. Walt. fl. car. 221.

In low spongy meadows: Pennsylvania to Carolina. 4. June, July. v. v. From ten inches to a foot high; flowers white. This and the following species are known by the name of *Ladies Traces*.

2. N. foliis lanceolatis trinervibus, caule vaginato, spica oblonga densiflora, floribus recurvato-cernuis, labello oblongo integerrimo acuto.—Willd. sp. pl. 4. p. 75.

cernua.

Ophrys cernua. Sp. pl. 1340.

Limodorum autumnale. Walt. fl. car. 221.

In grassy and springy places: Canada to Carolina. 4. July, Aug. v. v. Resembles the preceding, but the spikes are closer and the flowers double the size.

3. N. foliis radicalibus ovatis petiolatis reticulatis, scapo vaginato floribusque pubescentibus, floribus secundis, labello petalisque lanceolatis. Willd. sp. pl. 4. p. 76.

repens

Satyrium repens. Sp. pl. 1339.

Icon. *Jacq. austr.* 369. *Hall. helv.* t. 22. *Com. hort.* t. 35.

In rocky, shady pine-woods, on the high mountains of Virginia. ♀. July. v. v. Not above three inches high; stem creeping; leaves delicately veined; flowers white.

pubescens.

4. N. foliis radicalibus ovatis petiolatis reticulatis, scapo vaginato floribusque pubescentibus, labello ovato-acuminato, petalis ovatis. *Willd. sp. pl.* 4. p. 76.

Satyrium repens. Mich. fl. amer. 2. p. 137.

In shady and rocky situations: Canada to Florida. ♀. July. v. v. Resembles the preceding; is more robust, and the leaves more distinctly and handsomely variegated with white veins. This plant has lately made a great noise among the country people, as infallibly curing the bite of a mad dog.

673. ARETHUSA. *Swartz. act. holm.* 1800. p. 230.

bulbosa.

1. A. aphylla; radice globosa, scapo vaginato unifloro, calyce laciniis superioribus incurvatis, labello subcrenulato.—*Willd. sp. pl.* 4. p. 80.

Icon. *Lam. illustr.* t. 729. f. 1. *Pluk. mant.* t. 348. f. 7.

In sphagnum swamps: Canada to Carolina. ♀. June. v. v. Flowers large, purple, sweet-scented.

ophioglossoides.

2. A. radice fibrosa, scapo dissite bifoliato 1—2-floro, foliis ovali-lanceolatis, labello fimbriato.—*Willd. sp. pl.* 4. p. 80.

Icon. *Lam. illustr.* t. 729. f. 2.

In bogs and sphagnum swamps: Canada to Carolina. ♀. June, July. v. v. Flowers nearly the size of the preceding, purple, sometimes white.

divaricata.

3. A. radice subpalmata, scapo remote bifoliato unifloro, foliis oblongis obtusiusculis, petalis exterioribus adscendentibus longo-linearibus, labello eroso-subcrenulato.—*Willd. sp. pl.* 4. p. 81.

Icon. *Lam. illustr.* t. 729. f. 3. *Catesb. car.* 1. t. 58.

In swamps near the sea-coast: Virginia to Florida. ♀. July. v. v. Flowers larger than any of the preceding, purple.

pendula.

4. A. radice tuberosa, caule folioso summitate subtrifloro, foliis ovatis alternis, floribus longo-pedunculatis alternis, labello integro.—*Willd. sp. pl.* 4. p. 82.

A. trianthophoros. *Swartz. act. holm.* 1800. p. 230.

A. parviflora.
Icon. Pl.

In shady
York
three c.

5. A. foliis
floro,
interior
undula
Icon. Pl.

In shady
to Vir
handsome
A. divar
has so

6. A. foliis
flore s
interior
talis c
In shady
July.
ceding

674. EPI

1. E. foliis
cente
vix pe
radice
Ophrys p
Arethusa
Serapias f
Gron. J

In pine-l
v. s. :
greenish

2. E. caule
tis, spi
obtuse
Willd.
Ophrys c
In shady
to Ne
small p

Com. hort.
mountains of
three inches
high ; flow-
ers, scapo
ovato-acu-
tus.
76.
orida. 2.
s more ro-
andsomely
has lately
purple, as in-

1. p. 230.
niflora, ca-
llo subcre-
t. 348. f. 7.
2. 2. June.
l.
floro, foliis
l. sp. pl. 4.

Carolina.
the size of
o unifloro,
ioribus ad-
-subcrenu-
1. t. 58.
orida. 2.
the preced-
subtrifloro,
ulatis alter-
2.
p. 230.

- A. parviflora. *Mich. fl. amer.* 2. p. 160.
Icon. *Pluk. mant.* t. 348. f. 6.
In shady woods, round the roots of beech-trees : New
York to Kentucky. 2. July. v. v. Not above
three or four inches high ; flowers pale purple, small.
5. A. foliis quinis oblongo-lanceolatis verticillatis, caule uni-
floro, petalis tribus exterioribus longissimis linearibus,
interioribus lanceolatis, labello trilobo, lacinia media
undulata. *Willd. sp. pl.* 4. p. 81.
Icon. *Pluk. mant.* t. 348. f. 1. *tona*.
In shady woods, near the banks of rivulets : Pennsylvania
to Virginia. 2. July. v. v. A very scarce and
handsome species ; flowers, the size and shape of
A. divaricata, of a dull purple mixed with yellow. It
has sometimes two or three flowers.
6. A. foliis verticillatis oblongis acuminatis, caule unifloro,
flore subsessili, petalis tribus exterioribus linearibus,
interioribus brevioribus oblongis obtusis, labello pe-
talum consimile.
In shady woods on the Blue-mountains. *Vanvleck.* 2.
July. v. s. in *Herb. Vanvleck.* Resembles the pre-
ceding in general habit.
674. EPIPACTIS. *Swartz. act. holm.* 1800. p. 231.
1. E. foliis radicalibus ovatis acutis, scapo aphylo pubes-
cente laxifloro, floribus pedicellatis, labello bilobo
vix petalis conniventibus longiore, capsulis clavatis,
radice palmato.
Ophrys pubera. Mich. fl. amer. 2. p. 158.
Arethusa racemosa. Walt. fl. car. 222.
Serapias foliis ovatis radicalibus, scapo nudo multifloro.
Gron. fl. virg. 137.
In pine-barrens of Carolina and Georgia. 2. June.
v. s. in *Herb. Gronov. et Walter.* Flowers small,
greenish-white.
2. E. caule bifolio, foliis oppositis cordato-subrotundis acu-
tis, spica parviflora, labello oblongo apice dilatato
obtuse bilobo, germe subgloboso, radice fibrosa.—
Willd. sp. pl. 4. p. 88.
Ophrys cordata. Mich. fl. amer. 2. p. 158.
In shady cedar-swamps and sphagnous woods : Canada
to New Jersey ; rare. 2. May. v. v. A very
small plant ; flowers dark-brown, mixed with green.

pubescent.

*convallarioi-
des.*

675. MALAXIS. *Swartz. act. holm.* 1800. p. 233.*ophioglossoides.**liliifolia.*

1. M. folio solitario ovato amplexicauli, scapo pentagono, labello apice bifido. *Willd. sp. pl.* 4. p. 90.

M. unifolia. *Mich. fl. amer.* 2. p. 157.

In shady woods: Pennsylvania to Florida. $\frac{1}{4}$. May, June. v. v.

2. M. foliis binis ovato-lanceolatis, scapo triquetro, petalis interioribus filiformibus reflexis discoloribus, labello concavo obovato apice acuto.—*Willd. sp. pl.* 4. p. 92.

Ophrys trifolia. Walt. fl. car. 221.

Cyphrus scapo nudo, foliis radicalibus, &c. *Gron. virg.* p. 138.

Epidendrum caule erecto simplicissimo nudo, racemo simplici erecto. *Gron. virg.* p. 140.

Icon. *Bot. rep.* 65.

In shady wet woods, round the roots of trees: Canada to Virginia. $\frac{1}{4}$. June, July. v. v. The three outside petals are sharp-pointed, white; the two inside ones thread-shaped, yellowish, and reflex; the lower lip broad-obovate with an abrupt point of a pale olive colour.

676. CYMBIDIUM. *Swartz. nov. act. ups.* 6. p. 70.*pulchellum.*

1. C. foliis radicalibus ensiformibus nervosis, scapo paucifloro, labello erecto basi attenuato, lamina expansa, disco concavo barbato.—*Willd. sp. pl.* 4. p. 105.

Limodorum tuberosum. *Syst. veg.* 680. *Mich. fl. amer.* 2. p. 159.

Limodorum pulchellum. Salisb. prodr. 8.

Ophrys barbata. Walt. fl. car. 221.

Icon. *Bot. mag.* 116.

In sphagnum swamps: Canada to Florida. $\frac{1}{4}$. July, v. v. Flowers purple, very handsome.

vereendum.

2. C. foliis radicalibus lato-lanceolatis plicato-nervosis, scapo multifloro, petalis interioribus conniventibus, labello ventricoso, lamina emarginata crispa sulcata. *Willd. sp. pl.* 4. p. 105.

Limodorum vereendum. Salisb. prodr. 9.

Limodorum tuberosum. Jacq. collect. 4. p. 108.

Limodorum altum. Jacq. ic. rar. 3. t. 602.

Limodorum trifidum. Mich. fl. amer. 2. p. 159.

Icon. *Mart. cent. t.* 50. *Mill. ic.* 145.

GY

In Caroli
semble

3. C. foliis simplicibus 1
vato r
p. 107

Arethusa
On the si
Pensyl
ish-pur
the nat
a small
when i
ingly d

4. C. scapo lanceol
pl. 4. p
Ophrys c
Icon. *Pla*
In shady
Virgini
yellow.

677. C

1. C. folio ra
uniflor
Limodoru
Cypripedi
Cymbidi
Icon. *Pa*
sib. 1. t
In Nova S
M. Leu
Flower
resembl

p. 233.
pentagono,
).

4. May,
tro, petalis
us, labello
sp. pl. 4.

Gron. virg.
o, racemo

s: Canada
three out-
two inside
the lower
pale olive

5. p. 70.
ipo pauci-
a expansa,
105.
Tich. fl. a-

4. July.
osis, scapo
is, labello
. Willd.

3.
9.

In Carolina and Florida. 4. July, Aug. v. v. Resembles the preceding, but is larger.

3. C. foliis radicalibus geminis ovatis multinervibus, scapo simplici vaginato erecto, petalis erectis, tribus exterioribus lanceolatis, interioribus oblongis, labello obovato margine undulato-crenato.—Willd. sp., pl. 4. p. 107.

Arethusa spicata. Walt. fl. car. 222.

On the side of fertile hills, under the shade of trees: Pennsylvania to Carolina. 4. May. v. v. Flowers greenish-purple. It is known among the inhabitants by the name of *Adam and Eve*. The roots bruised, with a small addition of water, give a strong cement, which when applied to broken china and glass is exceedingly durable.

4. C. scapo vaginato aphylo, floribus pedicellatis, petalis lanceolatis æqualibus, labello ovato obtuso. Willd. sp. pl. 4. p. 110.

Ophrys corallorhiza. Mich. fl. amer. 2. p. 158.

Icon. Pluk. alm. t. 211. f. 1 ♂ 2.

In shady woods, about the roots of trees: Canada to Virginia. Flowers small, dark purple mixed with yellow.

677. CALYPSO. *Salisb. in parad. lond. t. 89.*

1. C. folio radicali unico subtrotundo-ovato nervoso, scapo unifloro vaginato.—*Salisb. l. c.*

Limodorum boreale. Willd. sp. pl. 123.

Cypripedium bulbosum. Sp. pl. 1347.

Cymbidium boreale. Swartz. nov. act. ups. 6. p. 76.

Icon. *Parad. lond. 89*. *Smith spicil. t. 11*. *Gmel. fl. sib. 1. t. 2. f. 5*.

In Nova Scotia. *Menzies*. On the Columbia river.

M. Lewis. 4. May, June. v. s. in *Herb. Lewis*. Flower large, purplish rose-coloured; the lower lip resembling *Cypripedium*.

hyemale.

Odontorhi-
zon.

borealis.

II. DIANDRIA.

678. CYPRIPEDIUM. *Swartz. act. holm.* 1800, p. 250.*candidum.*

1. C. caule folioso, foliis oblongo-lanceolatis, lobo styli lanceolato obtusiusculo, labello petalis lanceolatis breviore compresso.—*Willd. sp. pl. 4. p. 142.*
In Pennsylvania. *Muhlenberg.* 4. +. Resembles *C. Calceolus*, but the flowers are white, and not half the size: the form of the leaves and lobe of the style distinguish it sufficiently.

parviflorum.

2. C. caule folioso, lobo styli triangulari acuto, petalis exterioribus ovato-oblongis acuminatis, interioribus linearibus contortis, labello petalis breviore compresso. *Willd. sp. pl. 4. p. 143.*
C. Calceolus. Mich. fl. amer. 2. p. 161.
Icon. *Salisb. in linn. trans. 1. p. 77. t. 2. f. 2. Pluk. mant. t. 418. f. 2.*
In shady fertile woods: New Jersey to Carolina. 4. May, June. v. v. Petals greenish-brown; lower lip yellow, with brown spots.

pubescens.

3. C. caule folioso, lobo styli triangulari-oblongo obtuso, petalis exterioribus ovato-oblongis acuminatis, interioribus longissimis linearibus contortis, labello petalis breviore compresso. *Willd. sp. pl. 4. p. 143.*

*C. flavescens. Redout. pl. liliac.**C. Calceolus. Sp. pl. 1346.*Icon. *Willd. hort. berol. 1. t. 13. Moris. hist. 3. s. 12. t. 11. f. 15.**C. Calceolus Walt. fl. car. 222.*

On sunny fertile hills, among small trees and bushes, generally in rocky situations: Pennsylvania to Carolina. 4. May. v. v. Petals greenish-yellow with red spots; lower lip yellow. It approaches near to the preceding. Sometimes it is found with two or three flowers.

spectabile.

4. C. caule folioso, lobo styli elliptico-cordato obtuso, petalis exterioribus lato-ovalibus obtusis, labello petalis longiore antice fisso.—*Willd. sp. pl. 4. p. 143.*

*C. album. Ait. kew. 3. p. 303.**C. canadense. Mich. fl. amer. 2. p. 161.**C. reginæ. Walt. fl. car. 222.*

C. Calceolus

Icon. *Bot. hist. 3.*In low moun-
tains tra-
From e-
its foli-
are get-
lower I-
side.5. C. caule foliis 5. d.
lateraliter oblongo-
trigono ob-
p.In shady
M'Ken-
greenish
veins;
front vi-
ing the
high;
It disti-
by havi-
its lowe6. C. scapo longis c-
nato de-
fissio.C. acaule.
Icon. *Bot.*Catesb.
In shady
Canada
much a-
observe
Pensylv
among
Flower,

C. Calceolus γ. Sp. pl. 1346.
Icon. Bot. mag. 216. Pluk. mant. t. 418. f. 3. Moris.
hist. 3. s. 12. t. 11. f. 17.

In low meadows and bogs, particularly in the mountainous tracts : Canada to Carolina. ♀. May, June. v. v. From eighteen inches to three feet high. The plant in its foliage resembles *Helleborus viridis*; the flowers are generally by two, sometimes three, very large, the lower lip white with red veins, and crimson spots inside.

5. C. caule folioso, lobo styli orbiculato obtusiusculo, petalis 5. duobus inferioribus linear-lanceolatis deflexis, lateralibus linearibus horizontalibus, supremo ovato-oblongo acuto, labello longitudine petalorum antice obverse-conico saccato.—Ait. kew. ed. 2. tom. 5.
p.

In shady sphagnum woods, about Montreal, Canada. M'Kenzie. ♀. May. v. v. Flower small, petals greenish-brown; lower lip white, with crimson veins; of a very singular structure, resembling at a front view a sheep's head, the two lateral petals forming the horns. The stem is pubescent, not a span high; leaves smooth, in general alternately by four. It distinguishes itself very strongly from all the rest by having five petals, and by the singular structure of its lower lip.

6. C. scapo aphylo unifloro, foliis radicalibus geminis oblongis obtusis, lobo styli subrotundo-rhombeo acuminate deflexo, labello petalis lanceolatis longiore antice fiso. Willd. sp. pl. 4. p. 144.

C. acaule. Ait. kew. 3. p. 161.

Icon. Bot. mag. 192. Salisb. in linn. trans. 1. t. 3. f. 4.
Catesb. ear. app. t. 3. Pluk. mant. t. 418 f. 1.

In shady rocky situations and on dry mountain bogs : Canada to Carolina. ♀. May, June. v. v. Not much above a span high; flower purple, large. I have observed a white variety on the Broad-mountains, Pennsylvania. The whole of this genus is known among the inhabitants by the name of *Moccasin Flower*, or sometimes *Ladies Slippers*.

Arietinum.

humile.

III. HEXANDRIA.

679. ARISTOLOCHIA. Gen. pl. 1383.

Sipho.

1. A. foliis cordatis acutis, caule volubili, pedunculis unifloris bractea ovata instructis, corollis adscendentibus, limbo trifido æquali. *Willd. sp. pl. 4. p. 155.*
A. macrophylla. Lam. encycl. 1. p. 252.
Icon. Bot. mag. 534. L'Herit. stirp. t. 7.
 On the mountains: Pensylvania to Carolina. $\frac{1}{2}$. June.
 $v. v.$ A very high running vine; leaves large, and well calculated to form shady bowers; flowers very singular in structure, yellowish-brown; the leaves are more or less pubescent.

Serpentaria.

2. A. foliis cordatis oblongis acuminatis, caule flexuoso adscendente, pedunculis radicalibus, corollæ labio lanceolato. *Willd. sp. pl. 4. p. 159.*
 In shady woods, on fertile soil: New England to Carolina; principally throughout the mountains. $\frac{1}{2}$. May, June. $v. v.$ This is the famous *Virginia Snake-root*, so very highly and, I think, deservedly esteemed by the Indians and white inhabitants of America. I have seen a very narrow and long-leaved variety of it, which, if there were any difference in the flowers, might claim to be a distinct species.

IV. DODECANDRIA.

680. ASARUM. Gen. pl. 801.

canadense.

1. A. foliis lato-reniformibus geminatis, calyce lanato profunde tripartito, lacinia sublanceolatis reflexis.—
Mich. fl. amer. 1. p. 279. Willd. sp. pl. 2. p. 838.
A. carolinianum. Walt. fl. car. 143.
A. latifolium. Salisb. prodr. 344.
Icon. Moris. hist. 3. s. 13. t. 7. f. 4. Corn. canad. t. 25.
 In shady rocky situations: Canada to Carolina. $\frac{1}{2}$. April.
 $v. v.$ The root is highly aromatic, and known by the inhabitants under the name of *Wild Ginger*. It is said to be made use of by the Indian females to prevent impregnation.

2. A. foliis flore sulcato.—
 $p. 838.$
 In shady $v.$
 $v. v.$ L
3. A. foliis infra linamer. 1.
 A. virginicIn shady v
 Leaves s

2. A. foliis solitariis rotundato-cordatis glabris coriaceis, *virginicum*.
flore subsessili, calyce extus glabro breviter campanu-
lato.—*Mich. fl. amer.* 1. p. 279. *Willd. sp. pl.* 2.
p. 838.
In shady rocky woods: Virginia and Carolina. ♀. May.
v. v. Leaves white speckled, very smooth.
3. A. foliis subhastato-cordatis coriaceis, calyce tubuloso
infra limbum brevissimo trifidum coarctato. *Mich. fl.*
amer. 1. p. 279.
A. virginicum. *Walt. fl. car.* 143.
In shady woods of Lower Carolina. ♀. May. v. v.
Leaves speckled.

XIX. DICLINIA.

I. SEGREGATÆ.

* *Uniloculares.*

681. VALISNERIA. *Dioica. Masc. Spatha* 2-partita.
Spadix conicus tectus flosculis. *Cal.* 3-partitus.
Stam. 2. *Fem. Spatha* tubulosa, 2-fida, 1-flora.
Cal. superus, longissimus, limbo 6-partito, laciinis
alternis linearibus. *Stigmata* 3. semibifida. *Caps.*
cylindrica, 1-locularis, polysperma.
682. NĀJAS. *Monoica. Masc. Cal.* cylindricus, 2-fidus.
Stamen 1. filamento longo, anthera 4-valvi: valvis
patentibus. *Fem. Cal.* O. *Styl.* 1. *Stigm.* 2. *Nux*
4-sperma.
683. DIOTIS. *Monoica. Masc. Cal.* 4-phylus. *Stam.* 4.
Fem. Cal. 1-phylus, bicornis. *Stylus* 2-partitus.
Sem. 1. villosum, calyce bicorni tectum.
- ** *Tricoccæ; Capsula* 3-locularis, 3-sperma. *Stamina*
monadelpha. *Monoica.*
684. RICINUS. *Masc. Cal.* 5-partitus. *Stam.* numerosa.
Fem. Cal. 3-partitus. *Styli* 3. bifidi. *Caps.* echinata.
685. JATROPHA. *Masc. Cor.* 1-petala, infundibuliformis.
Stam. 10. alterna breviora. *Fem. Cor.* 5-pe-
tala, patens. *Styli* 3. bifidi.
686. CROTON. *Masc. Cal.* cylindricus, 5-dentatus. *Cor.*
5-petala, aut O. *Stam.* 10—15. *Fem. Cal.* poly-
phylus. *Cor.* O. *Styli* 3. bifidi.
687. ACALYPHA. *Cal.* 3—4-partitus, 1-bracteatus. *Masc.*
Bractea squamiformis. *Stam.* 8—16. *Fem. Brac-*
tea magna cordata involucriformis. Styli 3. bifidi.
688. TRAGIA. *Masc. Cal.* 3-partitus. *Stam.* 3. subses-
silia. *Fem. Cal.* 5-partitus. *Stylus* 3-fidus.

689. EUPHOC
limbo
que 5.
ticulati
foliolis
Flos f.
Styls 3.
690. STILLII
in masc
tubulosi
Stam. 2.
Stigm. 1.

691. SALIX.
floris.
Stam. 1.
fidus. C
papposa.
692. POPULU
squamis
integer.
fidum.
Sem. pap
693. MYRICA.
tis, triflo
rius 6-and

694. BETULA.
floris, 3-f
12. *Fem*
1. utrinqu
695. ALNUS.
catis 3-fl
Stam. 4.
Sem. 1. co
696. CARPINUS
floris. C
6—10.
lanceolata
tum. St

1.
A.

2-partita.
3-partitus.
a, 1-flora.
to, laciniis
ia. *Caps.*

is, 2-fidus.
lvi : valvis
n. 2. *Nux*

Stam. 4.
2-partitus.

Stamina

numerosa.
Caps. echin-
ndibulifor-
Cor. 5-pe-
atus. Cor.
Cal. poly-
tus. *Masc.*
Fem. Brac-
3. bifidi.
. 3. subses-
us.

689. EUPHORBIA. *Involucrum* commune calyciforme, limbo extorsum appendicibus petaloideis, plerumque 5.; introrsum lacinulis totidem alternis et denticulatis. *Masc.* tot quot lacinulae. *Cal.* polyphyllus: foliolis pinnatum laciniatis. *Stam.* 4-5. aut plura. *Flos fem.* centralis, nudus, solitarius, stipitatus. *Styli* 3. bifidi.

690. STILLINGIA. *Involucrum* coriaceum, urceolatum, in masculis multiflorum, 1-florum in feminineis. *Cal.* tubulosus, infundibuliformis, limbo ciliato. *Masc.* *Stam.* 2. exerta, basi coalita. *Fem.* *Stylus* filiformis. *Stigm.* 3.

II. AMENTACEÆ.

* *Flores dioici.*

691. SALIX. *Amenta* cylindracea, imbricata: squamis 1-floris. *Cal.* O. *Masc.* *Glandula* baseos nectarifera. *Stam.* 1-6, interdum monadelpha. *Fem.* *Styl.* bifidus. *Caps.* 1-locularis, 2-valvis, polysperma. *Sem.* papposa.

692. POPULUS. *Amenta* cylindracea, laxe imbricata: squamis margine laceris, 1-floris. *Cal.* urceolatus, integer. *Masc.* *Stam.* 8. exerta. *Fem.* *Stigma* 4-fidum. *Caps.* 2-locularis, 2-valvis, polysperma. *Sem.* papposa.

693. MYRICA. *Amenta* ovata, imbricata: squamis lunatis, trifloris. *Cal.* O. *Masc.* *Squama* 4-andra, rarius 6-andra. *Fem.* *Styli* 2. *Drupa* 1-sperma.

** *Flores monoici.*

694. BETULA. *Amenta* imbricata: squamis peltatis, 3-floris, 3-fidis. *Masc.* *Cal.* 4-partitus. *Stam.* 10-12. *Fem.* *Squama* 2-flora. *Cal.* O. *Stigm.* 2. *Sem.* 1. utrinque membrana alatum.

695. ALNUS. *Amentum* receptaculis cuneiformibus truncatis 3-floris compositum. *Masc.* *Cal.* 4-partitus. *Stam.* 4. *Fem.* *squama* biflora. *Cal.* O. *Stigm.* 2. *Sem.* 1. compressum, ovatum, apterum.

696. CARPINUS. *Amenta* laxe imbricata: squamis 1-floris. *Cal.* O. *Masc.* *Squama* ovata, ciliata. *Stam.* 6-10. *Antheræ* apice villosæ. *Fem.* *Squama* lanceolata, ciliata, biflora. *Germen* apice denticulatum. *Stigm.* 2. *Nux* ovata, sulcata.

697. OSTRYĀ. *Amenta imbricata. Masc. Cal. squama. Fil. ramosa. Fem. Amentum nudum. Caps. inflatæ, imbricatæ, basi monospermæ.*
698. FAGUS. *Masc. Amentum subrotundum. Cal. 5-fidus, campanulatus. Stam. 8—12. Fem. Cal. 4-dentatus, setosus. Germ. 2. Nuces 2. calyce echinato, coriaceo, 4-fido inclusæ.*
699. CASTANEA. *Masc. Amentum nudum, lineare. Cal. sub-6-phyllo. Stam. 5—20. Fem. Cal. 5-6-phyllo, muricatus. Germ. 3. Stigm. penicilliformia. Nuces 3. calyce echinato inclusæ.*
700. QUERCUS. *Masc. Amentum nudum, lineare. Cal. sub-5-fidus. Stam. 4—10. Fem. Cal. plerumque 6-dentatus, scaber. Styli 2—5. Nux coriacea, calyce persistente basi cincta.*
701. CORYLUS. *Masc. Amentum imbricatum : squamis 3-fidis unifloris. Cal. 0. Stam. 8. Fem. Cal. bipartitus, lacerus. Styli 2. Nux ovata, calyce persistente cincta.*
702. LIQUIDAMBAR. *Masc. Amentum conicum, involucro 4-phyllo cinctum. Filam. numerosa. Fem. Amentum globosum, involucro 4-phyllo cinctum. Cal. urceolatus, biflorus. Styli 2. Caps. 2. basi cinctæ, 2-valves, polyspermæ.*
703. COMPTONIA. *Masc. Amentum cylindricum, lake imbricatum ; squamis 1-floria. Cal. 2-phyllo. Filam. 3. bifurcata. Fem. Amentum ovatum. Cal. 6-phyllo. Styli 2. Nux ovata, 1-locularis.*
704. PLATANUS. *Amentum globosum. Masc. Cor. vix manifesta. Antheræ filamentum circumnatæ. Fem. Cor. polypetala. Styli stigmate recurvo. Sem. subrotunda, stylo mucronata, basi papposa.*
705. JUGLANS. *Masc. Amentum imbricatum. Cal. 6-partitus. Filam. 4—18. Fem. Cal. 4-fidus, superus. Cor. 4-partita. Styli 2. Drupa coriacea, nuce sulcata.*
706. MORUS. *Amentum subglobosum. Masc. Cal. 4-partitus. Filam. 4. Fem. Cal. 4-phyllo. Styli 2. Cal. baccatus. Sem. 1.*

707. PINUS.
lus. St
bili s. c
membran

708. CUPRESSUM : s
4-andris
squamis
singulis
Stigm.
sperma.

709. THUYA
ordine o
substipit
longis a
bus. G
cavum.

710. JUNIPER
peltatis.
pite 2-3
cruciati
bulosum.
locularis.

711. TAXUS.
Cal. 0.
Germ. 1.
læforme,

712. ZAMIA.
obovata.
dehiscent
tæ. Ge

Zal. squama.
Caps. inflatæ,
n. *Cal.* 5-
'em. *Cal.* 4-
calyce echin-

ineare. *Cal.*
d. 5-6-phyl-
aicilliformia.

ineare. *Cal.*
d. plerumque
ux coriacea,

m: squamis
'em. *Cal.* bi-
calyce per-

onicum, in-
erosa. *Fem.*
lo cinctum.
Caps. 2. basi

lricum, laxe
phyllus. *Fi-*
im. *Cal.* 6-
ris.
fasc. *Cor.* vix
inatæ. *Fem.*
. *Sem.* sub-

im. *Cal.* 6-
dus, superus.
iacea, nuce

fasc. *Cal.* 4-
lus. *Styli* 2.

III. CONIFERÆ.

* Flores monoici.

707. PINUS. *Masc.* *Amentum* imbricatum. *Cal.* 4-phyl-
lus. *Stam.* plurima. *Anth.* nudæ. *Fem.* *Cal.* stro-
bili s. coni: squama biflora. *Pistill.* 1.. *Nux* ala
membranacea excepta.
708. CUPRESSUS. *Masc.* *Amentum* ovatum, imbrica-
tum: squamis (20) oppositis, apice peltatis, basi
4-andris. *Anth.* sessiles. *Fem.* *Conus* subrotundus:
squamis (8—10) oppositis, peltatis. *Germina* sub
singulis plurima minima serie simplici. *Styl.* 0.
Stigm. cylindricum, fistulosum. *Nux* angulosa 1-
sperma.
709. THUYA. *Masc.* *Amentum* ovatum: squamis 6. trino
ordine oppositis, apice obtusis, basi 4-andris; *Anth.*
substipitatae. *Fem.* *Conus* subovatus: squamis ob-
longis apice incrassatis, longitudinaliter conniventibus.
Germina 2. *Styl.* brevissimus. *Stigm.* con-
cavum. *Caps.* 1-spermæ, margine alato.

** Flores dioici.

710. JUNIPERUS. *Amenta* ovata: squamis stipitato-
peltatis. *Masc.* *Squamæ* ternatim verticillatae, sti-
pite 2—3—4-antherifero. *Fem.* *Squamæ* pauciores,
cruciati oppositæ. *Germ.* 1. *Styl.* 0. *Stigm.* tu-
bulosum. *Fructus* bacciformis, carnosus. *Nux* 1-
locularis.
711. TAXUS. *Gemma* polyphylla, opposite-imbricata.
Cal. 0. *Masc.* *Antheræ* peltate, 8-fidæ. *Fem.*
Germ. 1. *Styl.* 0. *Stigm.* 1. *Receptaculum* cupu-
læforme, nuce ovala, nuda.
712. ZAMIA. *Amentum* strobiliforme. *Masc.* *Squama*
obovata. *Cal.* 0. *Antheræ* plures, globosæ, rima
dehiscentes, in squama sessiles. *Fem.* *Squamæ* pel-
tatae. *Germ.* 2. *Styl.* 0. *Baccae* 2. 7-spermæ.

I. SEGREGATÆ.

681. VALLISNERIA. *Gen. pl.* 1491.

- americana.* 1. *V. foliis linearibus, pedunculis masculis feminineisque rectis.* *Willd. sp. pl.* 4. p. 651. *Mich. fl. amer.* 2. p. 220.

At the bottom of muddy and slow-flowing rivers: Mississippi and St. John's, Florida. *Michaux.* In Oswego river, New York: Delaware, Pennsylvania, and James's and Greenbriar rivers, Virginia. *Ph.* 4. ♂? Aug. Oct. v. v. This most interesting and singular plant I observed in the Delaware, near Philadelphia, in 1800; which discovery I communicated, accompanied with a drawing and description, to Dr. B. S. Barton, of Philadelphia. The economy of its fecundation is highly interesting to the young student of the sexual system. *Michaux* and *Willdenow* consider it as specifically distinct from *V. spiralis*; but I have always considered it only a local variety, as the peduncles of the female flowers are in deep water really spiral.

682. NAJAS. *Willd. act. acad. berol.* 1798, p. 85.

- canadensis.* 1. *N. pusilla, filiformis, laevis; foliis angustissime linearibus.* *Mich. fl. amer.* 2. p. 220.
In lakes: Canada, on Lake St. John's. *Michaux.* +.

683. DIOTIS. *Gen. pl.* 1423.

- lanata.* 1. *D. omnibus partibus cano-tomentosa; caule flexuoso, glomerulis spicarum confertissimis.* On the banks of the Missouri, in open prairies. *M. Lewis.* ♀. Aug. Sept. v. s. in *Herb. Lewis.* Resembles *Diotis ceratooides*, but is distinct at first sight by its long woolly tomentum.

684. RICINUS. *Gen. pl.* 1464.

- communis.* 1. *R. foliis peltatis palmatis, lobis lanceolatis serratis, caule herbaceo pruinoso, stigmatibus tribus apice bifidis, capsulis echinatis.* *Willd. sp. pl.* 4. p. 564.
Icon. *Blackw. t.* 148.

685.

In cultivate
ginia and
duced by
tor-oil. Bu

686.

1. *J. herbacea,*
mato-loba
cymis bre
J. urens. *W.*
Icon. Pluk.
In fields and
Aug. v.
eatable, th
rally culti
Cassada.
in some pa
when they
the name

1. *C. foliis ellip
tus tomen
ris.* *Wil*
C. disjunctifl
β. C. spicis dich
Mich. fl. a
On the sea
Nashville.
this genus
sometimes
a real shru
are covered

2. *C. suffruticos*
integerrimi
terminalibu
C. punctatum
In sterile woo
Aug. v. s.

3. *C. hispidissim*
basi subinte
spicis alarib
sp. pl. 4. p.
C. scordiooides.

In cultivated grounds, frequent in old plantations in Virginia and Carolina. ♂. Aug. Sept. v. v. Introduced by the Negroes. Known by the name of *Castor-oil Bean*.

685. JATROPHA. *Gen. pl.* 1463.

1. *J. herbacea*, stimulis patulis horridissima; foliis subpalmato-lobatis, lobis obtusiusculis subsinuato-incisis, cymis brevi-pedunculatis, *Mich. fl. amer.* 2. p. 216.

J. urens. *Walt. fl. car.* 239.

Icon. Pluk. alm. t. 220 f. 3. *Marcg. brass.* 79. f. 2.

In fields and plantations: Virginia to Florida. ♀. July, Aug. v. v. Flowers white; roots tuberous and eatable, the same as *J. Manihot*, which is so generally cultivated in the West Indies under the name of *Cassada*. The present plant is a very injurious weed in some parts of Carolina, as it ruins the Negroes' feet when they tread upon it; from which it is known by the name of *Tread-softly*.

686. CROTON. *Gen. pl.* 1462.

1. *C. foliis ellipticis integerrimis obtusiusculis incanis subtus tomentosis petiolatis, spicis terminalibus paucifloris*.—*Willd. sp. pl.* 4. p. 532. *Walt. fl. car.* 239.

C. disjunctiflorum. *Mich. fl. amer.* 2. p. 214.

- β. *C. spicis dichotomalibus, flore semineo unico pedicellato*. *monanthogynum*. *Mich. fl. amer.* 2. p. 215.

On the sea-coast of Carolina; β. in Tennessee, near Nashville. ♀. ♀. June, July. v. s. The flowers of this genus are small and inconspicuous. The branches sometimes live over winter, but it cannot be considered a real shrub. Most of the species here mentioned are covered with stellated hairs.

2. *C. suffruticosum, stellato-tomentosum; foliis oblongis integerrimis subitus tomentosis petiolatis, pedunculis terminalibus subbifloris*.—*Willd. sp. pl.* 4. p. 535.

C. punctatum. *Jacq. ic. rar.* 3. t. 621.

In sterile woods of Georgia and Florida. ♀. ♀. July, Aug. v. s.

3. *C. hispidissimum*; foliis oblongis serratis subtus hirtis glandulosum. basi subintegerrimis biglandulosis, caule trichotomo, spicis alaribus, capsulis glomeratim sessilibus.—*Willd. sp. pl.* 4. p. 540.

C. scordioides. *Lam. encycl.* 2. p. 214.

- Icon. *Jacq. ic. rar.* 1. t. 194.
On the sea-coast of Carolina and Florida. ⊖ July,
Aug. v. v.
- capitatum.* 4. C. tomentoso-lanatum; foliis oblongo-ovalibus obtusis
basi rotundatis integris, utrinque tomentosis, floribus
femineis ad basin spicillæ marium capitatiæ congestis.
—*Mich. fl. amer.* 2. p. 214.
In the Illinois country. *Michaux.* ⊖ +. Resem-
bles *C. argenteum.*

687. ACALYPHA. *Gen. pl.* 1461.

- virginica.* 1. A. floribus femineis ad basin spicæ masculæ, involucris
ovatis acuminatis dentatis, foliis breviter petiolatis ob-
longo-lanceolatis remote obtuse-serratis.—*Willd. sp.*
pl. 4. p. 521.
Icon. *Pluk. phyt.* t. 99. f. 4.
In fields, cultivated grounds, and on road sides: Canada
to Virginia. ⊖ June—Aug. v. v.
- caroliniana.* 2. A. floribus femineis ad basin spicæ masculæ, involucris
cordatis dentatis, foliis longe petiolatis subrhomboidali-
ovatis serratis basi integerrimis.—*Willd. sp. pl. 4.*
p. 521. Walt. fl. car. 238.
In cultivated grounds: Virginia to Florida. ⊖ July,
Aug. v. v.

688. TRAGIA. *Gen. pl.* 1410.

- urens.* 1. T. erecta; foliis lanceolatis sessilibus obtusis apice sub-
dentatis, cauleque ramoso pubescentibus. *Willd. sp.*
pl. 4. p. 325.
- subovalis.* α. T. foliis oblongo-ovalibus nonnullis subcuneatis. *Mich.*
fl. amer. 2. p. 175.
Icon. *Pluk. alm.* t. 107. f. 5.
- lanceolata.* β. T. foliis lanceolatis subdentatis integrisque. *Mich. l. c.*
T. innocua. Walt. fl. car. 229.
- linearis.* γ. T. foliis linearibus fere omnibus integris. *Mich. l. c.*
In waste places, on road sides and cultivated grounds:
Virginia to Carolina. ⊖ June—Aug. v. v.
- urticifolia.* 2. T. erecta, caule hirsutissimo, foliis cordatis ovatis ser-
ratis.—*Willd. sp. pl. 4. p. 324.*
T. mercurialis. *Walt. fl. car.* 229.
In fields: Virginia to Georgia. ⊖ July. v. s. in Herb.
Lyon.

3. T. scandens, hi-
dentatis.—*H.*
In Kentucky.
Lyon.

689. E

* *Flor*

1. E. fruticescens
riformibus, s-
subumbellati
E. heterophylla
Icon. *Jacq. ic*
p. 81. t. 1.
On the banks
Florida. h
feet high; u
red.
2. E. pumila, hi-
flores ad su-
p. 211.
In shady rocky
July, Aug.
3. E. pusilla, erect
aribus integer
fl. amer. 2. p.
On the sea-coas
Resembles *E.*

** *Dichotom*

4. E. glabra, rame-
tis, foliis opp
corymbis terr
Icon. *Comm. pr*
jam. 1. t. 126.
On the banks of
Canada to Flo
5. E. erecto-patula
floribus axilla
coloratis.—*W.*
Icon. *Jacq. hort*
In cultivated gr
July—Sept.
The involucru

- July,
obtusis
floribus
agrestis.
Resem-
volucris
atis ob-
lld. sp.
Canada
volucris
ombeo-
pl. 4.
. July,
ce sub-
lld. sp.
Mich.
. l. c.
' c.
ounds :
itis ser-
n Herb.
3. *T. scandens*, *hispida*; *foliis profunde cordatis ovatis acute macrocarpa*.
dentatis.—*Willd. sp. pl. 4. p. 323.*

In Kentucky. *Michaux.* ⊙. July. v. s. in Herb.
Lyon.

689. EUPHORBIA. *Gen. pl. 823.*

* *Floribus fasciculato-terminalibus.*

1. *E. fruticescens*; *foliis petiolatis ovatis subdentatis pandu-* *cyathophora*.
riformibus, summis involucellisque coloratis, floribus
subumbellatis.—*Willd. sp. pl. 2. p. 891.*
E. heterophylla. *Jacq. collect. 1. p. 157.*
Icon. Jacq. ic. rar. 3. t. 480. Murr. comm. gæt. 7.
p. 81. t. 1.
On the banks of the Mississippi, and in Georgia and Florida. ⊙. June, July. v. v. About three feet high; upper leaves and involucres painted deep red.
2. *E. pumila*, *hirsuta*; *foliis oppositis ovalibus dentatis,* *dentata*.
flores ad summitates congestis. *Mich. fl. amer. 2.*
p. 211.
In shady rocky situations: Pennsylvania to Tennessee. ⊙.
July, Aug. v. v. The upper leaves spotted.
3. *E. pusilla*, *erecta*, *ramosa*, *pubescens*; *foliis sparsis line-* *graminifolia*.
aribus integerrimis, supremis basi discoloribus.—*Mich.*
fl. amer. 2. p. 210.
On the sea-coast of Georgia and Florida. *Michaux.* +.
Resembles *E. hyssopifolia*.

** *Dichotomæ*. (Umbella bifida aut nulla.)

4. *E. glabra*, *ramosissima*, *patulo-erecta*; *ramis divaricatis*, *foliis oppositis serratis ovali-oblongis subfalcatis*
corymbis terminalibus.—*Willd. sp. pl. 2. p. 895.*
Icon. Comm. præl. 60. t. 10. Rob. ic. 11. Sloan. hist.
jam. 1. t. 126.
On the banks of rivers and in cultivated sandy grounds:
Canada to Florida. ⊙. June—Sept. v. v.
5. *E. erecto-patula*; *foliis oppositis serratis oblongis pilosis*, *maculata*.
floribus axillaribus solitariis, appendicibus calycinis
coloratis.—*Willd. sp. pl. 2. p. 896.*
Icon. Jacq. hort. vind. t. 186. Pluk. alm. t. 65. f. 8.
In cultivated grounds: Pennsylvania to Carolina. ⊙.
July—Sept. v. v. Leaves marked with a brown spot.
The involucrum of the florets white.

thymifolia.

6. *E. humifusa*, gracilis, pubescens; foliis oppositis ovalioblongis obtusis superne subserratis, capitulis axillariibus glomeratis subsessilibus.—*Willd. sp. pl. 2. p. 898.*
Icon. *Pluk alm. t. 113. f. 2.* *Burm. zeyl. t. 105. f. 3.*
On the banks of the Ohio and Mississippi. ☽. June—
Aug. v. v.

polygonifolia.

7. *E. glaberrima*, diffusa; foliis oppositis integerrimis lineari-lanceolatis obtusis, floribus solitariis axillaribus.—*Willd. sp. pl. 2. p. 900.*

Icon. *Jacq. collect. suppl. t. 13. f. 3.*

In cultivated grounds: Canada to Virginia. ♀. June—
Sept. v. v.

Ipecacuanhæ.

8. *E. procumbens*, pumila, glabra; foliis oppositis obovalibus seu lanceolatis, pedunculis axillaribus unifloris elongatis.—*Willd. sp. pl. 2. p. 900.*

Icon. *Bot. mag. 1494.*

In pine-barrens: New Jersey to Carolina. ♀. June,
July. v. v. This humble species has perhaps the
deepest root of any perennial I know; I frequently
have dug to the depth of more than six feet, and was
by appearance as far off from its end as ever.

portulacoides.

9. *E. erecta*; foliis integerrimis ovalibus retusis, pedunculis axillaribus unifloris folia æquantibus.—*Willd. sp. pl. 2. p. 901.*

In sandy soil, and cultivated grounds: Pennsylvania. ♀.
June—Aug. v. v.

pulcritissima.

10. *E. erecta*; caule calycibusque pubescentibus, foliis oppositis sessilibus subcordato-ovalibus obtusis, pedunculis solitarii capillaris unifloris, appendicibus calycinis coloratis petaloideis. *Mich. fl. amer. 2. p. 212.*

In Carolina. *Michaux.* ♀. +

*** *Umbellatæ; involucratæ.*

Peplus.

11. *E. umbella* 3-fida: dichotoma, involucellis ovatis, foliis integerrimis obovatis petiolatis. *Willd. sp. pl. 2. p. 903.*

Icon. *Engl. bot. 959.*

In cultivated grounds: Pennsylvania and Virginia. ☽.
July, Aug. v. v.

obtusata.

12. *E. umbellata* 3-fida: bis dichotoma, involucellis ovatis obtusiusculis subcordatis, foliis alternis sessilibus spathulatis involucellisque serrulatis glabris, capsulis muricatis.

DICL

In cultivated
July, Aug.

13. *E. caulinis*;
oppositis to
dunculus te
amer. 2. p.
In shady rock
♀. July, /
alis annua.

14. *E. umbella* 3-
involutellis
coloratis, 1
plexicaulib
cibus calyc
losis.

On the Yellow
in Herb. 1
white margi
pendices ha
green leaves

15. *E. umbella* 4
latis integreri
Icon. *Blackw.*
Not native, b
cultivated g
nerally kno
plant, it bei
ground whei

16. *E. umbella* 5-1
lisque oblor
loideis obova
Icon. *Pluk. ma*
In dry fields:
It is very var
are quite line

17. *E. umbella* 5-
petalis integrati
latis. *Willd.*
Icon. *Gmel. sil*
In shady wet w
Virginia. ♀
feet high.

- In cultivated grounds: Virginia, near Staunton. ♀.
July, Aug. v. v. Resembles *E. spathulata*.
13. *E. caulinis* gracile-debilibus, simpliciter 3-fidis, foliis *mercurialina*, oppositis ternisve subsessilibus ovalibus integris, pedunculis terminalibus solitariis unifloris.—*Mich. fl. amer.* 2. p. 212.
- In shady rocky situations: Pennsylvania and Kentucky.
♀. July, Aug. v. v. Resembles in aspect *Mercularis annua*.
14. *E. umbella* 3-fida: bis dichotoma, involucro foliiforme, involucellis oblongis cordatis margine membranaceis coloratis, foliis lanceolato-oblongis subcordato-amplexicaulibus acutis glaucescentibus glabris, appendicibus calycinis petaloideis subrotundis, capsulis pilosis. *marginalata*.
On the Yellow-stone river. *M. Lewis*. ♂. July. v. s. in *Herb. Lewis*. A very handsome species; the white margin of the involucre and white petal-like appendices have a fine contrast with the elegant soft green leaves.
15. *E. umbella* 4-fida: dichotoma, foliis oppositis lanceolatis integerrimis. *Willd. sp. pl.* 2. p. 906.
Icon. *Blackw. t.* 123.
Not native, but frequently found near gardens and in cultivated ground. ♂. July, Aug. v. v. It is generally known in America by the name of *Mole-plant*, it being supposed that no moles disturb the ground where this plant grows.
16. *E. umbella* 5-fida: 3-fida: dichotoma; involucellis foliisque oblongis obtusis, appendicibus calycinis petaloideis obovatis.—*Willd. sp. pl.* 2. p. 916.
Icon. *Pluk. mant. t.* 446. f. 3.
In dry fields: Canada to Carolina. ♀. July, Aug. v. v.
It is very variable in its foliage: sometimes the leaves are quite linear.
17. *E. umbella* 5-fida: trifida: bifida, involucris ovatis, petalis integris, foliis lanceolatis subpilosis apice serrulatis. *Willd. sp. pl.* 2. p. 917.
Icon. *Gmel. sib.* 2. t. 93.
In shady wet woods, in the mountains of Maryland and Virginia. ♀. June, July. v. v. From two to four feet high.

690. STILLINGIA. *Gen. pl. 1470.**sylvatica.*

1. S. herbacea ; foliis sessilibus oblongis obtusis basi attenuatis serrulatis, flosculis masculis squamam floralem vix superantibus.—*Willd. sp. pl. 4. p. 588.*

In pine-barrens: Virginia to Florida. $\frac{1}{2}$. May, June.
v. v. Flowers yellow, in a spike, resembling a catkin.

ligustrina.

2. S. fruticosa ; foliis petiolatis ovali-lanceolatis utrinque attenuatis integerrimis, flosculis masculis brevissime pedicellatis.—*Willd. sp. pl. 4. p. 588.*

In shady woods : Carolina and Georgia. $\frac{1}{2}$. June, July.
v. v.

sebifera.

3. S. arborea ; foliis petiolatis rhomboides acuminatis integerrimis, infra basin glandula petiolari, floribus masculis pedicellatis. —*Willd. sp. pl. 4. p. 588.*

Croton sebiferum. Sp. pl. 1425.
Icon. *Pet. gaz. t. 54. f. 3.* *Pluk. amalth. t. 390. f. 2.*
On the sea-coast of South Carolina; originally a native of China. $\frac{1}{2}$. July, Aug. *v. v.* Known by the name of *Tallow Tree.*

II. AMENITACEÆ.

691. SALIX. *Gen. pl. 1493.*

* *Foliis integerrimis aut obsolete serratis; vernatio revoluta.*

viminalis.

1. S. foliis lanceolato-linearibus longissimis acuminatis integerrimis subundulatis subtus sericeo-argenteis, versus marginem sparse-glandulosis, stipulis exiguis, amentis præcocibus, squaniis subrotundis pilosissimis, germinalibus sessilibus ovatis, stylo filiformi, stigmatibus acutis indivisis.—*Willd. sp. pl. 4. p. 706.*

Icon. *Hoffm. salic. 1. t. 2. f. 1 & 2. t. 5. f. 2. t. 21. f. e. f. g.*

On the banks of rivers and about plantations; introduced from Europe. $\frac{1}{2}$. April, May. *v. v.* Filaments yellow, anthers orange.

candida.

2. S. foliis linear-lanceolatis longissimis apice obsolete denticulatis supra pubescens subtus niveo-tomentosis margine revolutis, stipulis lanceolatis petiolum subæ-

quantib
obovato
pl. 4. p.
In dry sha
April, N
is not n
collectio
among a
me on t
question
indebet
revoluta
sent arra

3. S. foliis 1
scenti-ci
stipulis c
dris, squ
to-lance
brevi, st
S. tristis.
S. alpina.
S. flava. S
In shady d
v. v. Fr
yellow,
changin
a red tip
pearance
been abl
lute; bu
strongly

4. S. foliis lir
marginé
venosis t
oblongis
In dry sa
March,
proaches

5. S. foliis ol
glandulo
stipulis n
apice ni
ovatis b
stigmatil

basi atte-
floralem
ay, June,
mbling a
utrinque
revissime
ne, July.
i integer-
masculis
D. f. 2.
a native
1 by the
revoluta.
inatis in-
is, versus
amentis
germi-
bus acu-
21. f. e.
roduced
filaments
ete den-
nentosis
n subæ-

quantibus, amentis præcocibus cylindraceis, squamis obovato-lanceolatis longissime villosis.—Willd. sp. pl. 4. p. 708.

In dry shady woods: New York to Pennsylvania. $\frac{1}{2}$. April, May. v. v. The native place of this willow is not mentioned by Willdenow; but a plant in the collection of G. Anderson, Esq. which was found among a promiscuous collection of willows made by me on the Catskill mountains, New York, decides the question. To this gentleman's acute observations I am indebted for the present division of *Salices "vernatio revoluta,"* as likewise for the greater part of the present arrangement of American willows.

3. S. foliis lanceolatis acutiusculis subintegerrimis pubescenti-canis, subtus rugoso-venosis, margine revolutis, stipulis deciduis lanceolatis, amentis præcocibus diandris, squamis oblongis margine villosis, germinibus ova-to-lanceolatis sericeo-villosos longe pedicellatis, stylo brevi, stigmatibus bifidis.—Willd. sp. pl. 4. p. 692.
- S. tristis. Muhl. in annals of bot. 2. p. 68. t. 5. f. 9.
- S. alpina. Walt. fl. car. 243.
- S. flava. Schæpf. mat. med. amer.

In shady dry woods: New York to Virginia. $\frac{1}{2}$. April. v. v. From three to five feet high; branches greenish-yellow, with black dots; the anthers are purple, changing to yellow when burst; scales white with a red tip, which gives the catkins a very pleasing appearance. In this and the following species I have not been able to ascertain whether the vernatio is revolute; but their natural affinity to the preceding ones strongly indicates that they are so.

4. S. foliis linear-lanceolatis utrinque acutis integerrimis margine revolutis supra glabriusculis subtus rugoso-venosis tomentosis, stipulis nullis, amentis præcocibus oblongis.—Willd. sp. pl. 4. p. 693.

In dry sandy woods: New Jersey to Carolina. $\frac{1}{2}$. March, April. v. v.; v. s. in Herb. Banks. Approaches near to the preceding.

5. S. foliis obovato-lanceolatis acutis integerrimis margine glandulosis glabris subtus glaucis, junioribus sericeis, stipulis nullis, amentis præcocibus recurvatis, squamis apice nigris, pilis longitudine germinis, germinibus ovatis brevi-pedicellatis sericeis, stylo brevissimo, stigmatibus bifidis.

Muhlenber-
giana.

tristis.

recurvata.

In shady woods, in the mountains of New Jersey and Pennsylvania. ♀. April. v. v. A low shrub; branches brown, smooth; buds yellow.

** *Foliis integerrimis aut obsolete serratis; vernatio equitans.*

† *Caule repente, depresso aut diffuso.*

repens.

6. *S. repens*; foliis elliptico-lanceolatis integerrimis acutis glabris subtus subsericeis, stipulis nullis, amentis præcocibus ovatis diandris, squamis obovatis obtusis pilosis apice fuscis, germinibus ovato-oblongis pedicellatis pubescentibus, stylo brevissimo, stigmatibus bilobis, capsulis glabris.—*Willd. sp. pl. 4 p. 693.*

S. depressa. Hoffm. salic. t. 15 ♂ 16.

S. polymorpha. Ehrh. frut. 49.

In Nova Scotia and Newfoundland. ♀. May. v. s. in *Herb. Banks.* A very small creeping species.

reticulata.

7. *S. repens*; foliis elliptico-orbiculatis obtusis integerrimis glabris reticulato-venosis subtus glaucis, stipulis nullis, amentis serotinis pedunculatis diandris, squamis obovatis obtusis pubescentibus, germinibus ovatis sessilibus villosis, stigmatibus subsessilibus bipartitis.—*Willd. sp. pl. 4 p. 685.*

Icon. *Fl. dan. 212. Hoffm. salic. t. 15. 16. 17. Fl. lapp. t. 8. f. L. ♂ t. 7. f. 1 ♂ 2.*

In Labrador, Newfoundland, and on the north-west coast. ♀. June. v. s. in *Herb. Banks.* A very low creeping species; leaves very handsomely marked with coloured veins.

vestita.

8. *S. repens*; foliis suborbiculatis integerrimis supra glabris reticulato-venosis subtus villis sericeis longissimis adpressis tectis, stipulis nullis, amentis serotinis pedunculatis linearibus sericeis, germinibus ovatis sessilibus villosis, stylo profunde bipartito, stigmatibus bifidis.

In Labrador. ♀. June. v. s. in *Herb. Lambert. et Banks.* A very elegant species, resembling the preceding.

Uva ursi.

9. *S. depressa*; foliis spatulato-obovatis obtusis integerimis, inferne margine sparse glandulosis, glabris, supra nitidis, stipulis nullis, amentis serotinis laxis, squamis oblongis ciliatis, germinibus ovatis pedicellatis glabris, stylo bipartito, stigmatibus bilobis.

In Labrador. ♀. April, May. v. v. in *Hort. Anderson.*

For this
Andersc
Uva ursi.

10. *S. depressa*
gerrimis
lidis nervis
In Labrador
neral hat

11. *S. diffusa*;
bris, su
subcoœta
vatis apic
In Labrador
♀. May
species is
clined to

12. *S. erectissima*
longo-lan
berrimis
In Labrador
gular spe
remarkab
rise from
proaches

13. *S. ramulis la*
gerrimis u
amentis c
oblongis
minibus o
stigmatibus
S. pensylvan
On the Cats
This elega
den of G.
me from A
nel, which

14. *S. foliis sub*
apice subse
præcocibus
unico, ant
to ellipticis
tis emargin

Jersey and
branches

rnatio

nis acutis
entis præ-
sis pilosis
edicellatis
s bilobis,

v. s. in

gerrimis
is nullis,
mis obo-
sessilibus
—Willd.

17. Fl.

rth-west
A very
markeda glabris
mis ad-
nis pe-
atis ses-
matibusbert. et
he pre-tegerri-
, supra
quamis
glabris,

derson.

For this beautiful little species I am indebted to G. Anderson, Esq.; it has all the appearance of *Arbutus Uva ursi* in habit, as well as in the form of its leaves.

10. S. depressa; foliis ovalibus subacutis basi cordatis integrerrimis reticulato-venosis supra glabris, subtus palidis nervo margineque pilosis, stipulis semicordatis.

In Labrador. ♀. v. v. s. fl in Hort. Anderson. In general habit it resembles *S. myrsinoides*.

11. S. diffusa; foliis obovatis obtusis integerrimis supra glabris, subtus sericeo-villosis, stipulis nullis, amentis subcoætaneis sessilibus oblongis 2-andris, squamis obovatis apice nigris pilosis.

In Labrador. ♀. Colmaster. North-west coast. Nelson. ♀. May. v. s. in Herb. Dickson. et Banks. This species is allied to *S. Arenaria* Linn. and is more inclined to be upright than the preceding.

12. S. erectincola, divaricata; ramulis lævigatis, foliis oblongo-lanceolatis utrinque acutis medio serrulatis glaberrimis patentibus planis discoloribus, stipulis nullis.

In Labrador. ♀. v. v. s. fl in Hort. Anderson. This singular species distinguishes itself at first sight by its remarkable plain and patent leaves; it is inclined to rise from the ground on a single low stem, and approaches to the following division.

†† *Caule erecto.*

13. S. ramulis lævigatis, foliis obovato-lanceolatis acutis integrerrimis utrinque glabris concoloribus, stipulis nullis, amentis coætaneis pedunculatis glaberrimis, squamis oblongis pedicello duplo brevioribus vix pilosis, germinibus ovato oblongis longissime pedicellatis glabris, stigmatibus sessilibus bifidis.

S. pensylvanica. Hortul.

On the Catskill mountains, New York. ♀. April. v. v. This elegant and singular species flowered in the garden of G. Anderson, Esq., from a plant brought by me from America. He has one through another channel, which appears to be the male to this species.

14. S. foliis suboppositis obovato-lanceolatis acutis glabris *Lambertiana*, apice subserratis discoloribus, stipulis nullis, amentis praecocibus, squamis orbiculatis nigris, filamento unico, anthera geminata, germinibus sessilibus ovato ellipticis sericeis, stylo brevissimo, stigmatibus ovatis emarginatis.—Willd. sp. pl. 4. p. 673.

*cordifolia.**olovata.**planifolia.**pedicellaris.*

Icon. *Engl. bot.* 1359.

On the banks of rivers and in willow grounds; introduced from Europe. $\frac{1}{2}$. March, April. v. v. It is cultivated on account of its exceeding tough branches for the use of making baskets.

rosmarinifolia.

15. *S. foliis strictis linear-lanceolatis utrinque acutis integrerimis margine subglandulosis supra deciduo-pubescentibus, subtus sericeis, stipulis exiguis lanceolatis erectis, amentis præcocibus ovatis recurvatis, squamis oblongis obtusis ciliatis, germinibus pedicellatis lanceolatis villosis, stigmatibus subsessilibus bifidis.* — *Willd. sp. pl. 4. p. 679.*

Icon. *Engl. bot.* 1365.

In wet meadows and mountain swamps: Pennsylvania to Carolina. $\frac{1}{2}$. March, April. v. v. Not above three feet high. This species, though mentioned as a British plant in the Flora Britannica, is decidedly of American origin; as both sexes have been introduced from that country by G. Anderson, Esq. It has a great resemblance in general character to *S. recurvata*, but the *vernatio equitans* removes them far from one another.

fuscata.

16. *S. foliis obovato-lanceolatis acutis glabris subserratis subtus glaucis, junioribus pubescentibus, stipulis exiguis, amentis præcocibus nutantibus, squamis obtusis intus vix pilosis, germinibus brevi-pedicellatis ovatis sericeis, stigmatibus sessilibus bilobis.*

In low overflowed grounds, on the banks of rivers: New York to Pennsylvania. $\frac{1}{2}$. March, April. v. v. Branches of the preceding year covered with a dark brown or black tomentum.

*** *Foliis remote obtuseque serratis.*

conifera.

17. *S. foliis oblongo-lanceolatis remote serratis acutis supra glabris subtus planis tomentosis, annotinis glabris, stipulis lunatis subdentatis, amentis præcocibus diandris, squamis lanceolatis obtusis villosis, germinibus pedicellatis lanceolatis sericeis, stylo bifido, stigmatibus bilobis.* — *Willd. sp. pl. 4. p. 705.*

S. longirostris. Mich. fl. amer. 2. p. 226.

Icon. *Wangh. amer. t. 31. f. 72.*

In shady woods on gravelly dry soil: New York to Carolina. $\frac{1}{2}$. April. v. v. The cone-like excrescence at the end of the branches, occasioned by an insect, is not unfrequently found on other species belonging

to the I
noides.

18. *S. foliis ob-*
tuse serr-
tis gland-
foliosis,
minibus
bifido, s

Icon. *Muh-*
In wet mea-
 $\frac{1}{2}$. Ap-
purple, s

19. *S. foliis ov-*
glabris su-
tatis, am-
cellatis o-
tibus bif
- On the bar-
March, /
color.

20. *S. foliis ob-*
apice inte-
ceolatis se-
tomentos
minibus
mediocri-
p. 665.

Icon. *Muhl-*
In low gro-
New Eng-
dark bro-
when but-
basket-m:

21. *S. foliis la-*
nuatis set-
semicorda
germinib
stigmatib
- In shady wo-
Pensylvan-
long; res

22. *S. foliis line-*
tissime de-
guis lance

to the present division, but particularly on *S. prionoides*.

18. *S. foliis oblongo-lanceolatis acutis basi biglandulosis obtuse serratis glabris subtus glaucis, stipulis ovatis acutis glanduloso-serratis, amentis coætaneis villosis basi foliosis, squamis lanceolatis obtusis villosis atris, germinibus longe pedicellatis lanceolatis glabris, stylo bifido, stigmatibus bifidis.* — *Willd. sp. pl. 4. p. 666.*

Icon. *Muhl. in annals of bot. 2. t. 5. f. 2.*

In wet meadows and woods : New England to Virginia. $\frac{1}{2}$. April. v. v. Branches green ; younger ones purple, smooth.

19. *S. foliis ovali-oblongis acutis remote undulato-serratis glabris subtus glaucis, stipulis semicordatis inciso-dentatis, amentis præcocibus villosis, germinibus pedicellatis ovatis acuminatis sericeis, stylo longo, stigmatibus bifidis.*

On the banks of rivers : Pennsylvania to Virginia. $\frac{1}{2}$. March, April. v. y. Middle-sized ; resembles *S. discolor*.

20. *S. foliis oblongis obtusiusculis glabris remote-serratis apice integerrimis subtus glaucis, stipulis deciduis lanceolatis serratis, amentis subcoætaneis diandris oblongis tomentosis, squamis oblongis acutis atris pilosis, germinibus subsessilibus lanceolatis tomentosis, stylo mediocri, stigmatibus bipartitis.* — *Willd. sp. pl. 4. p. 665.*

Icon. *Muhl. in annals of bot. 2. t. 5. f. 1.*

In low grounds and on the banks of rivers ; common : New England to Carolina. $\frac{1}{2}$. April. v. v. Branches dark brown ; filaments white ; anthers red, yellow when burst. This is the most common in use for basket-making.

21. *S. foliis lanceolatis acutis longissimis basi sensim attenuatis serrulatis glaberrimis subconcoloribus, stipulis semicordatis, amentis præcocibus erectis glabriusculis, germinibus pedicellatis ovatis glabris, stylo bifido, stigmatibus 2-lobis.*

In shady woods on the banks of rivers : New York and Pennsylvania. $\frac{1}{2}$. March, April. v. v. Leaves very long ; resembles *S. prionoides*.

22. *S. foliis linearibus utrinque acuminatis elongatis remotissime denticulatis glabris concoloribus, stipulis exiguis lanceolatis denticulatis, amentis serotinis pedun-*

myricoides.

prionoides.

discolor.

angustata.

longifolia.

culatis tomentosis diandris, squamis planis retusis, filamentis basi barbatis squama duplo longioribus.—Willd. sp. pl. 4. p. 670.

Icon. Muhl. in Annals of Bot. 2. t. 5. f. 6.

On the banks of the Susquehannah. Muhlenberg. ½. July. +. Not above two feet high; branches brown; branchlets white.

**** *Foliis dense acuteque serratis.*

+ *Triandæ.* (Filamentis 3 - 6.)

- talyonica.* 23. S. ramis pendulis, foliis lanceolatis acuminatis serratis glabris discoloribus, stipulis exiguis subrotundis, amentis coætaneis, germinibus sessilibus ovatis glabris.—Willd. sp. pl. 4. p. 671.

On road sides and about plantations; introduced from Europe. ½. May. v. v. The Weeping Willow is now almost naturalized, being every where planted for ornament.

- Houstoniana.* 24. S. foliis linear-lanceolatis acutis tenuissime serratis utrinque glabris nitidis concoloribus, stipulis nullis, amentis coætaneis cylindricis villosis, squamis ovatis acutis, filamentis 3 - 5. usque ad medium barbatis.

S. tristis. Hortul.

In Virginia and Carolina. ½. v. v. s. fl.; v. s. c. fl. in Herb. Banks. specimen Houstonianum. This species, so frequently found in the gardens under the name of *S. tristis*, is very far from being in any way related to it. The specimen in the Banksian Herbarium was collected by Houston, and, as it is said, in Vera Cruz. But I am confident it is a more northern plant, as I frequently have seen it in Virginia. The branches are extremely brittle at their base.

- falcata.* 25. S. foliis longissimis linear-lanceolatis superne sensim attenuatis subfalcatis basi acutis approximato-serratis utrinque glabris, junioribus sericeis, stipulis lunatis dentatis deflexis.

On the banks of rivers: Pennsylvania to Virginia. ½. v. v. s. fl. A very smooth species; branches very slender and brown.

- nigra.* 26. S. foliis lanceolatis utrinque acutis serrulatis concoloribus glabris petiolis et nervo medio supra tomentosis, stipulis exiguis dentatis, amentis coætaneis erectis cylindricis villosis, squamis oblongis villosissimis, filamentis 3 - 6. basi barbatis, germinibus pedicellatis ovatis gla-

bris, styl
pl. 4. p.

S. carolina

S. pentand

S. vulgaris.

Icon. Mich

t. 5. f. 5

On the ban

May. v.

smooth,

27. S. foliis ov
dati serr
lis oblong
3-andris,
serratis g
bris, styl
pl. 4. p. 6

Icon. Muhl

t.

In low grou
½. May.
cies; bran
to the siz
shrub. It

28. S. foliis oblo
rigidis glab
gata, petio
glandulosos
squamis lar
dicellatis la
tibus bipart

S. cordata. M

S. cordifolia.

Icon. Muhl.

In swamps an
April, May.
end, young
much in use

29. S. foliis obl
argute serra
plis ovato-s
coætaneis si
atris, germi

VOL. II.

etus, fila-
is.—Willd.

berg. h.
es brown;

tis serratis
dis, amen-
glabris.—

uced from
Willow is
e planted

atis utrin-
lis, amen-
itis acutis,

v. s. c. fl.
This spe-
under the
any way
in Herba-
s said, in
northern
nia. The

e sensim
to-serratis
is lunatis

nia. h.
hes very

coloribus
is, stipu-
cylindri-
culamentis
vatls gla-

bris, stylo brevissimo; stigmatibus bifidis.—Willd. sp.
pl. 4. p. 657. Marsh. arb. 293.

S. caroliniana. Mich. fl. amer. 2. p. 226.

S. pentandra. Walt. fl. car. 243.

S. vulgaris. Clayt. fl. virg.

Icon. Mich. arb. 3. t. Muhl. in annals of bot. 2.
t. 5. f. 5.

On the banks of rivers: Pennsylvania and Virginia. h.

May. v. v. A tree about twenty feet high; branches
smooth, very brittle at the base.

27. S. foliis ovato-oblongis cuspidato-acuminatis basi rotundatis serratis glandulosis utrinque glabris nitidis, stipulis oblongis glanduloso-serratis, amentis coætaneis sub-3-andris, squamis lanceolatis obtusis basi pilosis apice serratis glabris, germinibus lanceolato-subulatis glabris, stylo bifido, stigmatibus obtusis.—Willd. sp.
pl. 4. p. 667.

Icon. Muhl. in annals of bot. 2. t. 5. f. 7. Mich. arb. 3.
t.

In low grounds, about springs: New York to Virginia.
h. May. v. v. A very smooth and handsome species;
branches yellowish brown; sometimes it grows
to the size of a tree, but more generally it is only a
shrub. It approaches to *S. amygdalina* of Europe.

28. S. foliis oblongo-lanceolatis acuminatis basi subcordatis rigidis glabris argute serratis, serratura infima elongata, petiolis villosis, stipulis amplis cordatis obtusis glanduloso-serratis, amentis coætaneis sub-3-andris, squamis lanceolatis atris lanatis, germinibus longe pedicellatis lanceolatis glabris, stylo brevissimo, stigmatibus bipartitis.—Willd. sp. pl. 4. p. 667.

S. cordata. Mich. fl. amer. 2. p. 225.

S. cordifolia. Herb. Banks. niss.

Icon. Muhl. in annals of bot. 2. t. 5. f. 4.

In swamps and hedges: New England to Virginia. h.
April, May. v. v. Branches green, red towards the
end, younger ones pubescent. It is very tough, and
much in use by the basket-makers.

29. S. foliis oblongo-lanceolatis acuminatis basi cordatis argute serratis glabris subtus pallidioribus, stipulis amplis ovato-subrotundis cartilagineo-serratis, amentis coætaneis sub-3-andris, squamis lanceolatis lanatis atris, germinibus pedicellatis lanceolatis glabris, stylo

lucida.

rigida.

cordata.

brevissimo, stigmatibus bifidis.—*Willd. sp. pl. 4. p. 660.*

Icon. *Muhl. in annals of bot. 2. t. 5. f. 8.*

In low and swampy grounds, on the banks of rivers: New York to Virginia. ♂. April, May. v. v. A very elegant species, about six or eight feet high.

†† *Diandrae.*

grisea.

30. S. foliis lanceolatis acuminatis serrulatis supra glabris nervo medio tomentosis subtus sericeis nudis, stipulis linearibus deflexis deciduis, amentis præcocibus, squamis oblongis pilosis apice atris, germinibus oblongis pedicellatis sericeis, stigmatibus sessilibus obtusis.—*Willd. sp. pl. 4. p. 699.*

S. sericea. *Muhl. in annals of bot. 2. t. 6. f. 8. Marsh. arb.*

In low overflowed grounds: Pennsylvania to Virginia. ♂. April. v. v. About eight feet high; branches greenish-purple, very brittle at the base.

petiolaris.

31. S. foliis lanceolatis undique serratis glabris subtus glaucis sericeis basi plerumque inæqualibus, stipulis lunatis dentatis exiguis, amentis præcocibus laxis, squamis obovatis obtusis nigris pilosis, germinibus longe pedicellatis ovatis sericeis, stigmatibus sessilibus bilobis.—*Willd. sp. pl. 4. p. 665.*

S. pensylvanica. *Hortul.*

Icon. *Engl. bot. 1147.*

In swamps and on the banks of rivers; common. ♂. April. v. v.; v. s. in *Herb. Lambert. et Anderson.* Branches slender, smooth, dark brown. It has been by mistake adopted as a native of Great Britain.

ella.

32. S. foliis lanceolatis acuminatis serratis utrinque sericeis, serraturis infimis glandulosis, stipulis obsoletis, amentis coætaneis elongatis, squamis elliptico-lanceolatis condoloribus pubescentibus, germinibus subsessilibus ovato-oblongis denique glabratibus, stylo brevi, stigmatibus bipartitis crassis.—*Willd. sp. pl. 4. p. 710.*

Icon. *Hoffm. salic. 1. t. 7. & 8. & t. 24. f. 3.*

On road sides and river banks; introduced from Europe. ♂. April, May. v. v. A tall tree.

vitellina.

33. S. foliis lanceolatis acuminatis crebro-serratis supra glabris subtus discoloribus subsericeis, stipulis nullis, amentis subcoætaneis cylindricis, squamis ovato-lan-

ceolatis co
sessilibus o
libus bilob

Icon. *Hoffm.
bot. 1050.*

On road sides
duced fror
sized tree.

34. S. foliis lance
duloso-serr
majuscule :
glabris, flos
In low ground
v. v. Resen
in fructifica

35. S. foliis lance
mis concolor
rotundatis,
villous, nec
subsessilibus
bipartitis.—
On road sides
Europe. ♂.
very brittle.

36. S. foliis elliptic
bris nitidis co
mulis hirtis, :
squamis lanceol
silibus bifidis.
Icon. *Fl. lapp. t.
salic. t. 18.*

- In Labrador. ♂.
in *Herb. La*
above a foot h

37. S. foliis orbicula
tidis, stipulis
squamis obvo
ovatis subsessil
sessilibus.—
Icon. *Fl. dan.
t. 8. f. h. t. 7.*
On the north-we

pl. 4.

f rivers:
v. A
high.bris ner-
, stipulis
is, squa-
oblongis
btusis.—

Marsh.

Virginia.
branchesis glaucis
lunatis
squamis
ige pedi-
lobis.—on. h.
nderson.
has been
in.sericeis,
s, amen-
ncoelatis
sessilibus
stigma-
to.

Europe.

apra gla-
nullis,
ato-lan-*ceolatis concoloribus extus pubescentibus, germinibus sessilibus ovato-lanceolatis glabris, stigmatibus subsessilibus bilobis.* —Willd. sp. pl. 4. p. 668.

Icon. Hoffm. salic. 1. t. 11. & 12. & t. 24. f. 1. Engl. bot. 1050.

On road sides and about plantations; common. Introduced from Europe. h. May. v. v. A middle-sized tree.

34. *S. foliis lanceolatis acuminatis concoloribus glabris glanduloso-serratis, amentis coætaneis, nectario gemino majusculo: lobis lanceolatis difformibus apice dentatis glabris, flosculis terminalibus 3-andris.*

In low grounds: New York and New Jersey. h. April. v. v. Resembles in leaves and habit the preceding, and in fructification the following.

35. *S. foliis lanceolatis acuminatis undique serratis glaberri- mis concoloribus, petiolis dentato-glandulosis, stipulis rotundatis, amentis subcoætaneis, squamis obovatiss villosis, nectario masculorum gemino, germinibus subsessilibus lanceolatis glabris, stigmatibus sessilibus bipartitis.* —Hoffm. salic. 2. t. 31. Engl. bot. 1937.

On road sides and about plantations. Introduced from Europe. h. April, May. v. v. A tall tree; branches very brittle.

36. *S. foliis elliptico-ovatis undique serrulatis utrinque glabris nitidis concoloribus, stipulis lanceolatis serratis, ramulis hirtis, amentis subpræcocibus erectis cylindricis, squamis lanceolato-oblongis longe villosis, germinibus ovato-lanceolatis sessilibus sericeis, stigmatibus subsessilibus bifidis.* —Willd. sp. pl. 4. p. 678.

Icon. Fl. lapp. t. 8. f. f. t. 7. f. 6. Fl. dan. 1054. Hoffm. salic. t. 18.

In Labrador. h. May. v. v. in Hort. Anderson.; v. s. in Herb. Lambert. A small straggling bush, not above a foot high; branches purplish-yellow.

37. *S. foliis orbiculatis subretusis serratis utrinque glabris nitidis, stipulis nullis, amentis serotinis paucifloris, squamis obovatiss obtusis villosis, germinibus oblongo-ovatis subsessilibus glabris, stigmatibus brevissimis subsessilibus.* —Willd. sp. pl. 4. p. 682.

Icon. Fl. dan. 117. Hoffm. salic. 1. t. 20. Fl. lapp. t. 8. f. h. t. 7. f. 3. & 4.

On the north-west coast. D. Nelson. h. June, July.

s 2

*amligua.**decipiens.**myrsinites.**herbacea.*

v. s. in Herb. Banks. The smallest of all the known species; the stems not above an inch high.

692. POPULUS. *Gen. pl. 1531.*

- balsamifera.* 1. *P. foliis ovatis acuminatis adpresso-serratis subtus albidis reticulato-venosis, gemmis resinosis.* *Willd. sp. pl. 4. p. 805.*
Icon. Mich. arb. 3. p. 306. t. 13. f. 1. Duham. arb. ed. nov. 2. t. 50. Pall. fl. ross. 1. t. 41. Wangh. amer. t. 28. f. 59. Trew ehret. t. 46. bona.
 In Canada, particularly plenty about Hudson's Bay. $\frac{1}{2}$. March. *v. v.* From seventy to eighty feet high; the young buds are covered with a very odiferous balsam, from which it is called *Balsam Poplar*; the natives call it *Tacamahac*.
- candicans.* 2. *P. foliis cordatis ovatis acuminatis obtuse inæqualiter serratis subtus albidis subtriplinervibus reticulato-venosis, petiolis hirsutis, gemmis resinosis, ramis teretibus.—Willd. sp. pl. 4. p. 806.*
P. latifolia. Mœnch. method. p. 338.
P. canadensis. Mœnch. weissenst. 81.
Icon. Catesb. car. 1. t. 34.
 In New England. $\frac{1}{2}$. March. *v. v.* From forty to fifty feet high; the buds are strongly covered with a balsamic fluid.
- trepida.* 3. *P. foliis suborbiculatis abrupte acuminatis dentatis basi supra biglandulosis glabris, junioribus seficeis.—Willd. sp. pl. 4. p. 803.*
P. tremuloides. Mich. fl. amer. 2. p. 243.
Icon. Mich. arb. 3. p. 285. t. 8. f. 1. Duham. arb. ed. nov. 2. t. 53.
 In extensive swamps: Canada to Pennsylvania. $\frac{1}{2}$. April. *v. v.* From twenty to thirty feet high. Generally known by the name of *American Aspen Tree*.
- monilifera.* 4. *P. foliis subcordato-deltoidibus glabris basi glandulosis, serraturis cartilagineis hamatis pilosiusculis, nervis partulis, petiolis superne compressis, ramis adultis teretibus.—Willd. sp. pl. 4. p. 805.*
P. glandulosa. Mœnch. method. p. 339.
P. carolinensis. Mœnch. weissenst. 81.
Icon. Mich. arb. p. 295. t. 10. f. 2.
 In North America. $\frac{1}{2}$. April. *v. v. in Hortis.* From sixty to seventy feet high. Michaux says that it has

- not been growing w
5. *P. foliis rho dentatis gla*
P. hudsonica.
P. nigra. Mic
 On the Huds
 rio. $\frac{1}{2}$. N
 high; petio
6. *P. foliis subr*
grandi-dent
perne comp
 In Canada. $\frac{1}{2}$
 high.
7. *P. foliis subro*
datis inæqua
lis compress
pl. 4 p. 803
P. canadensis.
 In high rocky
 the Western
 to eighty fee
Tree.
8. *P. foliis ovato*
dentatis glab
alato-angulos
P. heterophylla
P. basamifera.
Icon. Mich. ar
 In morasses, on
 and on the
 eighty feet hi
 It is known t
9. *P. foliis subrotu*
latis obtusis t
Willd. sp. pl
Icon. Mich. arb
t. 51.
 In swamps: Ne
 rivers. $\frac{1}{2}$.
 feet high. Th
 ing, more auri

known

tus albi-
illd. sp.arb. ed.
h. amer.Bay. h.
t high;
oriferous
ar; theliter ser-
-venosis,
tibus.—forty to
1 with aatis basi-
-Willd.

arb. ed.

ia. h.
h. Ge-
en Tree.
ndulosis,
ervis pa-
is tereti-s: From
t it has

- not been observed in any part of North America growing wild.
5. *P. foliis rhomboideis promisse acuminatis circumcirca dentatis glabris, ramulis junioribus pilosis.* *betulifolia.*
P. hudsonica. *Mich. arb. 3. p. 293. t. 10. f. 1.*
P. nigra. *Mich. fl. amer. 2. p. 244.*
 On the Hudson- or North-river, and about Lake Ontario. h. March. v. v. From thirty to forty feet high; petioles and young branches yellow.
 6. *P. foliis subrotundo-ovatis acutis inæqualiter sinuato-grandi-dentatis glabris, junioribus villosis, petiolis superne compressis.* *Mich. fl. amer. 2. p. 243.* *grandidentata.*
 In Canada. h. April. v. v. From forty to fifty feet high.
 7. *P. foliis subrotundo-ovatis deltoideis acuminatis subcordatis inæqualiter serratis glabris basi glandulosis, petiolis compressis, ramis junioribus angulatis.* *Willd. sp. pl. 4 p. 803.* *lævigata.*
P. canadensis. *Mich. arb. 3. p. 298. t. 11.*
 In high rocky situations: Canada to Virginia, and about the Western lakes. h. March. v. v. From seventy to eighty feet high. In Virginia it is called *Cotton Tree.*
 8. *P. foliis ovato-deltoideis acuminatis obtuse uncinato-dentatis glabris, junioribus amplissimis cordatis, ramis alato-angulosi.* *Willd. sp. pl. 4. p. 805.* *angulata.*
P. heterophylla. *Du Roi hærbk. 2. p. 150.*
P. basamifera. *Mill. dict. n. 5.*
Icon. *Mich. arb. 3. p. 302. t. 12.*
 In morasses, on the banks of rivers: Virginia to Florida, and on the Mississippi. h. April. v. v. About eighty feet high; leaves very large; branches brittle. It is known by the name of *Mississippi Cotton Tree.*
 9. *P. foliis subrotundo-ovatis sinu parvo cordatis subauriculatis obtusis uncinato-dentatis, junioribus tomentosis.* *heterophylla.* *Willd. sp. pl. 4. p. 806.*
Icon. *Mich. arb. 3. p. 290. t. 9.* *Duham. arb. ed. nov. 2. t. 51.*
 In swamps: New York to Carolina, and on the western rivers. h. May. v. v. About seventy or eighty feet high. The leaves are on their base, properly speaking, more auriculated than cordated.

Gale.

693. MYRICA. Gen. pl. 1510.

1. M. foliis cuneato-lanceolatis apice serratis obtusis, amentis masculis imbricatis, squamis acuminatis ciliatis, fructibus squamoso-capitatis.—Willd. sp. pl. 4. p. 745.
Icon. Fl. dan. 327.

In boggy grounds in Canada, and about lakes on high mountains; plentifully on the Broad mountains, Pennsylvania. $\frac{1}{2}$. May. v. v. A low shrub, like all the following, of an agreeable sweet scent.

cerifera.

2. M. foliis cuneato-lanceolatis apice rarer serratis acutis, amentis masculis laxis, squamis acutis, fructibus globosis minoribus.—Willd. sp. pl. 4. p. 745.

Icon. Catesb. car. 1. t. 69. Pluk. alm. t. 48. f. 9.

pumila.

- β . M. foliis lineari-lanceolatis. Mich. fl. amer. 2. p. 228.
In shady dry woods: Virginia to Carolina. β . In Carolina and Florida. $\frac{1}{2}$. May, June. v. v. A shrub sometimes more than twelve feet high. The berries of this and the following species produce a white vegetable wax, which the inhabitants manufacture into candles equal to those made of bees-wax.

carolinensis.

3. M. foliis cuneato-oblongis grosse dentatis, amentis masculis laxis, squamis acutis, baccis globosis majoribus.—Willd. sp. pl. 4. p. 746.

M. cerifera β . Mich. fl. amer. 2. p. 223.

Icon. Catesb. car. 1. p. 13.

In wet places about rivers and swamps: New England to Florida. $\frac{1}{2}$. May. v. v. Not above three or four feet high.

pensylvanica.

4. M. foliis oblongis utrinque acutiusculis integerrimis aut apice rarer subserratis margine revolutis, amentis masculis laxis, squamis acutis, baccis globosis majusculis.—Lam. encycl. 2. p. 592.

Icon. Duham. arb. ed. nov. 2. t. 55.

In shady rocky situations: New Jersey and Pennsylvania. $\frac{1}{2}$. May. v. v. Not above three feet high. It generally is confounded with the preceding, but is a very distinct species.

694. BETULA. Gen. pl. 1419.

populifolia.

1. B. foliis deltoidibus longe acuminatis inaequaliter serratis glaberrimis, strobilorum squamis lobis lateralibus subtundis, petiolis glabris. Willd. sp. pl. 4. p. 463.

B. acuminata.
B. lenta. Du.

t. 29. f. 38.

Icon. Mich. a.

In barren rock

sylvania. $\frac{1}{2}$

feet high.

or Old Field

2. B. foliis ovatis dunculo brevibus rotunda

B. lutea. Mich.

Icon. Mich. l.

In low grounds

May, June.

generally call

3. B. foliis rhomb pubescentibus squamis villos

sp. pl. 4. p. 4

B. lanulosa. Mi

B. rubra. Mich.

Icon. Mich. l. c

On the banks c

May. v. v.

grown; the

Red Birch.

4. B. foliis ovatis hirsutis, petiolatis notantibus, orbiculatis.— $\frac{1}{2}$

B. papyrifera. M

Icon. Mich. arb.

In the forests of C

as far south as

v. v. A large

constructing the

circumstance it

5. B. foliis cordato- o
subtus petiolisq
lobis obtusis æ
pl. 4. 464.

B. carpinifolia. Eh

- B. acuminata. *Ehrh. beitr. 6. p. 98.*
- B. lenta. *Du Roi harbk. 1. p. 92. Wangh. amer. p. 45. t. 29. f. 38.*
Icon. *Mich. arb. 2. p. 139. t. 2. Willd. arb. t. 2. f. 5.*
In barren rocky woods and old fields: Canada to Pennsylvania. $\frac{1}{2}$. June, July. v. v. From thirty to forty feet high. It is known by the name of *White Birch* or *Old Field Birch*.
2. B. foliis ovatis acutis serratis, petiolis pubescentibus pedunculo brevioribus, strobilorum squamis lobis lateribus rotundatis. *Willd. sp. pl. 4. p. 463.*
- B. lutea. *Mich. arb. 2. p. 152.*
Icon. *Mich. l. c. t. 5. Willd. arb. t. 2. f. 2.*
In low grounds: province of Maine, Hudson's river. $\frac{1}{2}$. May, June. v. v. From seventy to eighty feet high; generally called *Yellow Birch*.
3. B. foliis rhomboe-ovatis duplicato-serratis acutis subtus pubescentibus, basi integris, amentis feminineis ovatis, squamis villosis, lacinii linearibus æqualibus.—*Willd. sp. pl. 4. p. 464.*
- B. lanulosa. *Mich. fl. amer. 2. p. 181.*
- B. rubra. *Mich. arb. 2. p. 142.*
Icon. *Mich. l. c. t. 3. Willd. arb. t. 2. f. 6.*
On the banks of rivers: New Jersey to Carolina. $\frac{1}{2}$. May. v. v. About seventy feet high when full grown; the wood very fine. It is generally called *Red Birch*.
4. B. foliis ovatis acuminatis duplicato-serratis, venis subtus hirsutis, petiolo glabro, amentis feminineis pedunculis nutantibus, squamis lobis lateralibus brevibus suborbiculatis.—*Willd. sp. pl. 4. p. 464.*
- B. papyrifera. *Mich. fl. amer. 2. p. 180.*
Icon. *Mich. arb. 2. p. 133. t. 1. Willd. arb. t. 2. f. 1.*
In the forests of Canada to a great northern latitude, and as far south as the Hudson's river. $\frac{1}{2}$. May, June. v. v. A large tree, highly useful to the natives for constructing their large portable canoes, from which circumstance it is known by the name of *Canoe Birch*.
5. B. foliis cordato-ovatis argute serratis acuminatis, nervis subtus petiolisque pilosis, strobilorum squamis glabris, lobis obtusis æqualibus elevato-venosis.—*Willd. sp. pl. 4. 464.*
- B. carpinifolia. *Ehrh. beitr. 6. p. 99.*

*excelsa.**nigra.**papyracea.**lenta.*

B. nigra. *Du Roi harbk.* 1. p. 93. *Wangh. amer.* t. 15. f. 34.

Icon. *Mich. arb.* 2. p. 145. t. 4.

In Canada and throughout the Allegany mountains, to their termination in Georgia. $\frac{1}{2}$. May, June. v. v. This elegant and large tree is the most interesting of this genus, on account of the excellence of its wood. It is known by the name of *Mountain Mahogany*, *Black Birch*, *Cherry Birch*, and *Sweet Birch*. This last appellation it has from the sweet scent the branchlets give when bruised.

pumila.

6. *B. ramis pubescentibus impunctatis, foliis orbiculato-obovatis petiolatis subtus dense pubescentibus, amentis feminis cylindricis.* — *Willd. sp. pl. 4. p. 467.*

B. nana. *Kalm itin.* 2. p. 263.

Icon. *Jacq. hort. vind.* t. 122. *Du Roi harbk.* 1. t. 3. *Wangh. amer.* t. 29. f. 61.

In bogs of Canada, and on high mountains of New York and Pennsylvania. $\frac{1}{2}$. May, June. v. v. A low shrub, not above two or three feet high.

glandulosa.

7. *B. ramis glanduloso-punctatis glabris, foliis obovatis serratis basi integerrimis glabris subsessilibus, amentis feminis oblongis, squamis semitrifidis, seminibus orbiculatis angusto-marginatis.* — *Willd. sp. pl. 4. p. 466.*

In Canada, about Hudson's Bay, and on the borders of lakes on the high mountains of New Jersey and Pennsylvania. $\frac{1}{2}$. May. v. v. A handsome little shrub, not above two feet high.

nana.

8. *B. humillima, glaberrima; foliis orbiculatis crenatis subtus reticulato-venosis, amenti squamis profunde 3-partitis, lacinias oblongis, seminibus orbiculatis subapteris.* — *Willd. sp. pl. 4. p. 465.*

Icon. *Amœn. acad.* 1. t. 1. *Fl. dan.* 91. *Pall. fl. ross.* 1. t. 40. *Amman. act.* 9. t. 14.

In sphagnum swamps about Hudson's Bay and other parts of Canada. $\frac{1}{2}$. April, May. v. s. in *Herb. Lambert.*

glutinosa.

695. ALNUS. *Willd. sp. pl. 4. p. 334.*

Betula. *Gen. pl. 1419.*

1. *A. foliis subrotundo-euneatis obtusis subretusis glutinosis, axillis venarum subtus villosis.* *Willd. sp. pl. 4. p. 334.*

Betula Alna
Icon. *Engl.*
On the bank
on the no
tree, know

2. *A. foliis obl
petiolis ve
stipulis ov:*

A. undulata.
Betula Alnus
Betula crispa.
In Canada and
of Pensylva
three or fo

3. *A. foliis obo
tus pilosis,*
p. 336.

Betula serrula
Betula rugosa
Icon. *Abbot.*
In swamps a
 $\frac{1}{2}$. March.
growing in

696. (

1. *C. foliis oblon
stroblorum*
liqua ovato-l
pl. 4. p. 468

C. virginiana.
In shady wood
May. v. v.
European ve
qualities: it

697. (

1. *O. foliis ovato-
qualiter serra
nis acutis.—*

Carpinus virgin
Icon. *Abbot. in*
f. 1.
In shady woods

- t. 15.
is, to
v. v.
ng of
wood.
gany,
This
inch-
ilato-
entis
t. 3.
Fork
low
vatis
entis
s or-
466.
s of
Pen-
rub.
natis
inde
sub-
s. 1.
ther
erb.
sis,
4.
- Betula Alnus. Sp. pl. 1394.
Icon. Engl. bot. 1508. Læs. pruss. t. 1.
On the banks of rivers, in the interior of Canada, and
on the north-west coast. h. April. v. v. A large
tree, known every where under the name of Alder.
2. A. foliis oblongis acutis basi rotundatis duplicato-serratis,
petiolis venisque subtus pilosis, axillis venarum nudis,
stipulis ovato-oblongis. *crispa*
A. undulata. Willd. sp. pl. 4. p. 336.
- Betula Alnus crispa. Mich. fl. amer. 2. p. 181.
Betula crispa. Ait. kew. 3. p. 339.
In Canada and on high mountains, in sphagnum swamps,
of Pennsylvania. h. April. v. v. A shrub not above
three or four feet high.
3. A. foliis obovatis acuminatis venis et axillis venarum sub-
tus pilosis, stipulis ellipticis obtusis. Willd. sp. pl. 4.
p. 336. *serrulata*.
Betula serrulata. Ait. kew. 3. p. 338.
Betula rugosa. Ehrh. beitr. 3. p. 21.
Icon. Abbot. insect. 2. t. 92. Wangh. amer.
In swamps and on river sides; common every where.
h. March. v. v. A shrub from six to ten feet high,
growing in close thickets.

696. CARPINUS. Gen. pl. 1449.

1. C. foliis oblongo-ovatis acuminatis inæqualiter serratis,
strobilorum squamis tripartitis, lacinia intermedia ob-
liqua ovato-lanceolata uno latere dentata.—Willd. sp.
pl. 4. p. 468. *americana*.
C. virginiana. Mich. arb. 3. t. 8.
In shady woods and hedges: Canada to Florida. h.
May. v. v. This species of Hornbeam resembles the
European very much, and its wood is of the same
qualities: it rises to about fifteen feet high.

697. OSTRYA. Mich. gen. 223.

1. O. foliis ovato-oblongis basi subcordatis acuminatis inæ-
qualiter serratis, strobilis oblongo-ovatis erectis, gemi-
nis acutis.—Willd. sp. pl. 4. p. 469. *virginica*.
Carpinus virginiana. Ait. kew. 3. p. 363.
Icon. Abbot. insect. 2. p. 151. t. 75. Pluk. alm. t. 156.
f. 1.
In shady woods: New England to Carolina. h. May.

v. v. A small tree, of exceeding hard and heavy wood, from which it is generally known under the name of *Iron-wood*; in some parts they call it *Lever-wood*. I observed a variety in *Pensylvania*, of which the younger branches were covered with glandulous hairs. It is rather singular to observe that the plate in *Mich. arb. 3. t. 7.* intended for the present plant, represents *Ostrya vulgaris* with hanging cones, which is never the case in the American species.

698. FAGUS. *Gen. pl. 1448.**sylvatica.*

1. *F. foliis ovatis acuminatis leviter dentatis margine ciliatis basi acutis, nucibus ovato-triquetris obtusis cum mucrone.* — *Willd. sp. pl. 4. p. 459. Mich. arb. 2. p. 170.*

Icon. *Mich. l. c. t. 8.*

In extensive forests: New Hampshire to Georgia. $\frac{1}{2}$. May. *v. v.* This species of beech forms those extensive and beautiful woods generally known by the name of *Beech-woods*, and always gives the indication of a fertile soil. It is distinguished by the name of *White Beech* from the following.

ferruginea.

2. *F. foliis ovato-oblongis acuminatis subtus pubescentibus grosse dentatis basi obtusis subcordato-inæqualibus, nucibus acute ovato-triquetris acutissimis.* — *Willd. sp. pl. 4. p. 460.*

Icon. *Mich. arb. 2. p. 174. t. 9. Abbot. insect. 2. t. 75.* In rich level lands, at the foot of mountains: Canada to New England. $\frac{1}{2}$. May, June. *v. v.* It is distinguished by the inhabitants by the name of *Red Beech*, the wood being of a darker colour than the preceding. Both species are highly useful timber trees.

699. CASTANEA. *Tournef. inst. t. 352.**vesca : ame-
ricana.*

1. *C. foliis lanceolatis acuminatis mucronato-serratis utrinque glabris.* — *Willd. sp. pl. 4. p. 460.*

*Fagus Castanea. Hort. cliff. 447.*Icon. *Mich. arb. 2. p. 156. t. 6.*

In the mountainous parts: New England to Carolina. $\frac{1}{2}$. May, June. *v. v.* The *American Chestnut* differs so little from the European, that no specific distinction can be drawn. It is one of the largest and

DICLI

most useful
tremely dur
to construct

2. *C. foliis oblon
tomentosis.* —

Fagus pumila.
Icon. *Mich. c
f. 44. Abb
Pluk. alm. t*

In dry barren
to Georgia.
or rather sl
and upwards
it seldom ex
very sweet ap
by the name

700. C

* *Fructificatio bien
muero.*

1. *Q. foliis decidu
integerrimis
glande subrot
Q. Phellos sylva*

*Icon. Mich. l. c
car. 1. t. 16.*

t. 5. f. 11. F

- β. *Q. foliis brevior
amer. t. 5. f. 1*

In low swampy
to Florida. $\frac{1}{2}$
to the height
leaves are dent
gling growth.

2. *Q. foliis perenna
glabris basi att
scutellata, gl
p. 424.*

Q. Phellos mariti

*On the sea-coast
June. v. v.*

eight feet high

most useful trees of the forests, the wood being extremely durable, and in high esteem for posts and rails to construct fences. The nuts are very delicious.

2. *C. foliis oblongis acutis mucronato-serratis subtus albo-tomentosis*.—*Willd. sp. pl. 4. p. 461.*

Fagus pumila. *Sp. pl. 1418.*

Icon. *Mich. arb. 2. p. 166. t. 7.* *Wangh. amer. t. 19. f. 44.* *Abbot. insect. t. 57.* *Catesb. car. 1. t. 9. Pluk. alm. t. 156. f. 2.*

In dry barren and sandy fields and woods: New Jersey to Georgia. $\frac{1}{2}$. May, June. v. v. This small tree, or rather shrub, grows to the height of thirty feet and upwards in the southern regions, but to the north it seldom exceeds seven or eight feet. The fruit is very sweet and agreeable to eat, and is generally known by the name of *Chinquapin*.

pumila.

700. QUERCUS. *Gen. pl. 1446.*

- * *Fructificatio biennis; fructibus subsessilibus; foliis setaceo-mucronatis.* (in specie 1. muticis.)

+ *Foliis integerrimis.*

1. *Q. foliis deciduis linear-lanceolatis utrinque attenuatis integerrimis glabris mucronatis, cupula soutellata, glande subrotunda*.—*Willd. sp. pl. 4. p. 423.*

Q. Phellos sylvatica. *Mich. querc. n. 7.*

Icon. *Mich. l. c. t. 12.* *Mich. arb. 2. t. 12.* *Catesb. car. 1. t. 16.* *Abbot. insect. 2. t. 91.* *Wangh. amer. t. 5. f. 11.* *Pluk. amaln. t. 441. f. 7.*

Phellos.

- β . *Q. foliis brevioribus.* *Catesb. car. 1. t. 22.* *Wangh. amer. t. 5. f. 12.*

In low swampy forests, near the sea-coast: New Jersey to Florida. $\frac{1}{2}$. May. v. v. The *Willow Oak* grows to the height of about fifty or sixty feet. The young leaves are dentated. The variety β . is of low straggling growth.

humilis.

2. *Q. foliis perennantibus coriaceis lanceolatis integerrimis glabris basi attenuatis, apice acutis unicoloratis, cupula scutellata, glande subrotunda*.—*Willd. sp. pl. 4. p. 424.*

Q. Phellos maritima. *Mich. querc. t. 13. f. 1.*

maritima.

On the sea-coast of Virginia and Carolina. $\frac{1}{2}$. May, June. v. v. A low shrubby species, from three to eight feet high.

sericea.

3. *Q. foliis deciduis lanceolato-oblongis integerrimis subundatis basi attenuatis obtusis, apice dilatatis acutis subtus sericeis, cupula scutellata, glande subglobosa.* — *Willd. sp. pl. 4. p. 424.*

*Q. Phellos pumila. Mich. querc. t. 13. f. 1. ♂ 2.**Q. Phellos sericea. Ait. kew. 3. p. 354. Abbot. insect. 2. t. 51.**Q. pumila. Mich. arb. 2. p. 84. t. 15. Walt. fl. car. 234.*
Near the sea-coast : Carolina to Florida. $\frac{1}{2}$. May. v. v.

The *Running Oak* is perhaps the smallest of all the species, scarcely ever exceeding two feet in height.

myrtifolia.

4. *Q. foliis perennantibus coriaceis oblongis integerrimis glabris utrinque acutis supra nitidis margine revolutis.* — *Willd. sp. pl. 4. p. 424.*

In Carolina. *Willdenow.* $\frac{1}{2}$. +. Leaves resemble those of *Myrtus communis* with large leaves ; the fruit is not known.

virens.

5. *Q. foliis perennantibus coriaceis oblongo-ellipticis integerrimis margine revolutis basi obtusis apice acutis muticis subtus stellatum pubescentibus, fructibus penduculatis, cupula turbinata, squamis abbreviatis, glande oblonga.* — *Willd. sp. pl. 4. p. 425.*

*Q. Phellos β. Sp. pl. 1412.**Q. sempervirens. Walt. fl. car. 234.**Icon. Mich. querc. t. 10. ♂ 11. Mich. arb. t. 11. Catesb. car. 1. t. 16.*

Near the sea-coast : Virginia to Florida and Mississippi. $\frac{1}{2}$. May. v. v. The *Live Oak* grows to the height of forty or fifty feet, spreading its branches, when in open places, extremely wide : it yields the finest and most durable ship-timber of any species known ; for which reason it is considered one of the most valuable trees in America.

cinerea.

6. *Q. foliis perennantibus coriaceis lanceolato-oblongis integerrimis margine revolutis basi attenuatis apice obtusiusculis mucronatis subtus stellatum tormentosis, fructibus sessilibus, cupula scutellata, glande subglobosa.* — *Willd. sp. pl. 4. p. 425.*

*Q. Phellos β. Ait. kew. 3. p. 354.**Q. humilis. Walt. fl. car. 234.**Icon. Mich. querc. n. 8. t. 14. Mich. arb. 2. p. 80. t. 14.*

In dry barren soil and pine-forests : Virginia to Georgia. $\frac{1}{2}$. May, June. v. v. The *Upland Willow Oak* is

DICL

very variat
at others :
chaux con-
variety of
doubt.

7. *Q. foliis deci-
tegerrimis
tellata : se-
Willd. sp. 1*

*Icon. Mich.**p. 78. t. 13**On the banks**and in the**v. v. The**feet. The y**species of 1*

8. *Q. foliis decid-
tis integerr-
squamis la-
pl. 4. p. 427*

Icon. Mich. q

- β. *Q. foliis apice*

*In South Carol**Laurel Oak,**Oak, is ab-**cording to 1**almost prefe**son the you**cies, I canno**of the preced*

†† Ful

9. *Q. foliis subro-
remote spino-
mis laxis, &
p. 431.*

*Icon. Pluk. ph**On the north-**h. +.*

10. *Q. foliis longe-
tegris vel in
sphaerica, gla-*

Icon. Mich. t. c

- On the banks of
v. v. Of thi*

very variable in size, sometimes not above four feet, at others nearly twenty feet high. The younger Michaux considers *Q. nana* of Willdenow to be only a variety of the present species; which I very much doubt.

7. *Q. foliis deciduis oblongis utrinque acutis mucronatis integrerrimis nitidis subtilis pubescentibus, cupula scutellata : squamis lato-ovatis, glande subglobosa.*—Willd. sp. pl. 4. p. 428.

Icon. *Mich. querc. n. 9. t. 15. & 16. Mich. arb. 2. p. 78. t. 13.*

On the banks of rivers, within the Allegany mountains, and in the western countries thereof. ♂. May, June. v. v. The *Shingle Oak* rises to about forty or fifty feet. The younger Michaux considers the wood of this species of little value.

8. *Q. foliis deciduis obovato-lanceolatis acutis basi attenuatis integerrimis utrinque glabris, cupula scutellata : squamis lanceolatis, glande subovata.*—Willd. sp. pl. 4. p. 427.

Icon. *Mich. querc. n. 10. t. 17.*

9. *Q. foliis apice obtusis.* *Mich. querc. t. 18.*
In South Carolina and Georgia. ♂. May. v. v. The *Laurel Oak*, or as it is sometimes called *Swamp Willow Oak*, is about fifty or sixty feet high: its wood, according to the elder Michaux, is very valuable, and almost preferable to that of *Q. virens*. For what reason the younger Michaux does not mention this species, I cannot tell, unless he considers it only a variety of the preceding; which most certainly is not the case.

† *Foliis dentatis aut breviter lobatis.*

9. *Q. foliis subrotundo-ovatis subcordatis utrinque glabris remote spinoso-dentatis, cupula hemisphaerica : squamis laxis, glande ovata acuta.*—Willd. sp. pl. 4. p. 431.

Icon. *Pluk. phytogr. t. 196. f. 3?*

On the north-west coast, about Nootka Sound. *Nee.* ♂. +.

10. *Q. foliis longe petiolatis ovato-lanceolatis oblongisve integris vel inaequaliter grandidentatis, cupula hemisphaerica, glande subglobosa.*—*Mich. arb. 2. p. 87.*

Icon. *Mich. l. c. t. 16.*

On the banks of the Delaware, Pennsylvania. ♂. May. v. v. Of this singular species there is but one individ-

imbricaria.

laurifolia.

obtusa.

agrifolia.

heterophylla.

DICLINIA AMENTACEÆ. Quercus.

DICL

- dual known, which grows on the plantation of the Messrs. Bartrams near Philadelphia. It probably is only a hybrid plant on that account, and cannot with propriety be considered a genuine species: but the younger Michaux having given a good figure of it, I insert it here on that account only.
- aquatica.* 11. *Q. foliis obovato-cuneiformibus glabris integerrimis apice obsolete trilobis, lobo intermedio majore, cupula hemisphaerica, glande subglobosa.* — *Willd. sp. pl. 4. p. 441.*
Q. nigra. *Sp. pl. 1413.*
Q. uliginosa. *Wangh. amer. t. 6. f. 18.*
Icon. Mich. querc. n. 11. t. 19. t. 20. f. 1. 3. 4. 5. & t. 21. *Mich. arb. 2. p. 89. t. 17.*
 In swamps: Maryland to Florida. $\text{h}.$ May. $v.$ $v.$ The Water Oak is about forty feet high when full grown; its wood is but little valued. Its leaves vary, according to soil and age, *ad infinitum*. There is scarcely one tree found having leaves like the other, and the same tree is almost as variable in its different branches.
- hemisphaerica.* 12. *Q. foliis perennantibus oblongo-lanceolatis indivisis trilobis sinuatisque, lobis mucronatis, utrinque glabris.* — *Willd. sp. pl. 4. p. 443.* *Bartr. itin. 320.*
Icon. Mich. querc. t. 20. f. 2.
 In Georgia and Florida. $\text{h}.$ May. $v.$ $v.$ $s.$ This is probably nothing more than a young plant of *Q. aquatica*, which is so very apt to vary in its foliage.
- nana.* 13. *Q. foliis cuneiformibus glabris apice trilobis basi subsinuatis, lobis divaricatis mucronatis intermedio majore, axillis venarum subtus pubescentibus, cupula scutellata, glande ovato-subglobosa.* — *Willd. sp. pl. 4. p. 443.*
Q. aquatica elongata. *Ait. kew. 3. p. 357.*
Icon. Abbot. insect. 2. t. 59.
 In pine-barrens of South Carolina. $\text{h}.$ May. $v.$ $v.$ $s.$ $fr.$ A low-growing species, always keeping distinct from *Q. aquatica*.
- triloba.* 14. *Q. foliis oblongis cuneiformibus basi acutis, apice subtrilobis, lobis æqualibus mucronatis, intermedio longiore, subtus tomentosis, cupula scutellata, glande depresso-globosa.* — *Willd. sp. pl. 4. p. 443.*
Q. cuneata. *Wangh. amer. p. 78. t. 5. f. 14.*
Icon. Mich. querc. n. 14. t. 26. *Abbot. insect. 1. t. 50?*
 In pine-barrens, near the sea-coast: New Jersey to Geor-
- gia. $\text{h}.$ from twenty of very rapid inclosing of
15. *Q. foliis coriaceis dilatatis reticulatis pra glabris, turbinata:* *ovata.* — *Wi*
Q. nigra $\beta.$ *S*
Q. ferruginea. *Icon. Mich. q*
Catesb. car.
insect. t. 58.
- In barren sand ridges. $\text{h}.$ l Jack of the more numerous and furnished but excellent
16. *Q. foliis obovato-sessentibus, tis setaceo-nigro-globosus.* *Q. tinctoria angustifolia.* *Q. discolor.* *Q. velutina.* *L* In all large woods New England Black Oak, or the American its timber valuable in tanning to
17. *Q. foliis oblongibus, lobis obtusatis, cupula turbinata.* *p. 444.*
Q. tinctoria sinuata. *Icon. Abbot. insect. 1. f. 5.* In large forests $v.$ $v.$ Resembling *cinea*. The yellow

- gia. $\frac{h}{2}$. May. *v. v.* The *Downy Black Oak* is from twenty to forty feet high, according to Michaux, of very rapid growth, and extremely well calculated for inclosing of lands.
15. *Q. foliis coriaceis cuneiformibus basi subcordatis, apice dilatatis retuso-subtrilobis, junioribus mucronatis, supra glabris, subtus rubiginoso-pulverulentis, cupula turbinata: squamis obtusis scariosis, glande breviovata.* — *Willd. sp. pl. 4. p. 442.*
Q. nigra β . *Sp. pl. 1413.*
Q. ferruginea. Mich. arb. 2. p. 92.
Icon. Mich. querc. n. 12. t. 22. 23. Mich. arb. t. 18.
Catesb. car. 1. t. 19. Wangh. amer. t. 5. f. 13. Abbot. insect. t. 58.
- In barren sandy or gravelly woods: New Jersey to Florida. $\frac{h}{2}$. May. *v. v.* The *Barren Oak* or *Black Jack* of the Virginians is of low growth, especially in the more northern states: it bears very abundantly and furnishes a fine mast for hogs: the wood is small, but excellent for fuel.
16. *Q. foliis obovato-oblongis levissime sinuatis subtus pubescentibus, lobis oblongis obtusis obsolete denticulatis setaceo-mucronatis, cupula scutellata, glande depresso-globosa.* — *Willd. sp. pl. 4. p. 444.*
Q. tinctoria angulosa. Mich. querc. n. 13. t. 24.
Q. discolor. Willd. nov. act. soc. nat. sci. ut. berol. 3. p. 399.
Q. velutina. Lam. encycl. 1. p. 173
- In all large woods, particularly in the mountainous parts: New England to Georgia. $\frac{h}{2}$. May. *v. v.* The *Black Oak*, or *Quercitron*, is one of the largest trees of the American forest, and highly valuable on account of its timber as well as its bark, which is so very superior in tanning to any other species of oak.
17. *Q. foliis oblongis pinnatifido-sinuatibus, lobis oblongis dentatis setaceo-mucronatis, cupula turbinata, glande ovata.* — *Willd. sp. pl. 4. p. 444.*
Q. tinctoria sinuosa. Mich. querc. n. 13. t. 25.
Icon. Abbot. insect. 2. p. 111. & 56. Plák. alm. t. 54. f. 5.
- In large forests: Pennsylvania to Carolina. $\frac{h}{2}$. May. *v. v.* Resembles the preceding, and likewise *Q. coccinea*. The young expanding leaves are covered with

*nigra.**tinctoria**discolor.*

a white down on both sides, which is not the case with either *Q. rubra* or *Q. coccinea*. Whether the figure of *Q. tinctoria* Mich. arb. 2. t. 22. belongs to this or the preceding species, I have not been able to decide.

+++ *Folii profunde sinuatis lobatisque.*

coccinea.

18. *Q. foliis longe petiolatis oblongis profunde sinuatis glabris, lobis divaricatis dentatis acutis setaceo-mucronatis, cupula turbinata insigniter squamosa, glande breviovata.* — *Willd. sp. pl. 4. p. 446.*

Icon. *Mich. querc. n. 18. t. 31. 32. Mich. arb. 2. p. 116. t. 23. Wangh. amer. t. 4. f. 9.*

In fertile woods: New England to Georgia. ♀. May. v. v. The Scarlet Oak is a very large tree, and changes its leaves in autumn to a bright red, which give the woods a very picturesque appearance.

ambigua.

19. *Q. foliis sinuatis glabris basi acutis, sinubus subacutis, cupula subscutellata, glande turgide ovata.* — *Mich. arb. 2. p. 120.*

Icon. *Mich. l. c. t. 24.*

On Hudson's Bay and in Nova Scotia. *Michaux.* ♀. May. +. This is an intermediate species, between the preceding and following one, and probably only a hybrid plant. Michaux calls it *Gray Oak*.

rubra.

20. *Q. foliis longe petiolatis oblongis glabris obtuse sinuatis, lobis acutiusculis dentatis setaceo-mucronatis, cupula scutellata sublaevi, glande subovata turgida.* — *Willd. sp. pl. 4. p. 445.*

Icon. *Mich. querc. n. 20. t. 35. 36. Mich. arb. 2. p. 126. t. 26. Wangh. amer. t. 3. f. 7. Pluk. alm. t. 54. f. 4. Du Roi, harbk. 2. t. 5. f. 2.*

In fertile forests: Canada to Pennsylvania, and in the Alleghany mountains. ♀. May. v. v. This large and fine timber tree is exclusively known by the name of *Red Oak*, which name is likewise applied to *Q. tinctoria*, *coccinea*, *falcata*, *palustris*, *phellos*, *ambigua*, and *aquatica*, in several parts of America.

Catesbeei.

21. *Q. foliis brevissime petiolatis basi cuneatis oblongis profunde sinuatis glabris, lobis tribus quiniseve divaricatis dentatis acutis setaceo-mucronatis, cupula turbinata ampla: squamis obtusis marginalibus introflexis, glande subglobosa.* — *Willd. sp. pl. 4. p. 446.*

Icon. *Mich. querc. n. 17. t. 29. 30. Mich. arb. 2. p. 101. t. 20. Catesb. car. 1. t. 23.*

DICL

In pine-barren
The Barren
fifteen feet
those count

22. *Q. foliis longe trilobis sinu terminali ek sa.* — *Mich. l. elongata.* *Icon. Mich. qu*
t. 21.

In sandy soil, ♀.
May. Spanish Oak
the name of

23. *Q. foliis longe bris, axillis v dentatis acut laevi, glandes*
Icon. Mich. qu
t. 25. Du Roi
f. 10.

In low swamp
and in Illinois
nish Oak or I
small and han
generally in g

24. *Q. foliis longe quelobisve ma tosis, lobis set glande subglob*
Q. ilicifolia. Wi
t. 6. f. 17.

Icon. *Mich. qu*
t. 19. Abbot.

In dry barren fie
to Virginia.
four or six feet
wherever it oc
by the name o
Dwarf Red Oai

In pine-barrens: Carolina and Georgia. $\frac{1}{2}$. May. v. v. The *Barren Scrub Oak* is of shrubby growth, not above fifteen feet high, and occurs in the most sterile soil of those countries in great abundance.

22. Q. foliis longe petiolatis basi obtusis subtus tomentosis trilobis sinuatis, lobis subfalcatis setaceo-mucronatis, terminali elongato, cupula crateriformi, glande globosa.—*Mich. fl. amer. 2. p. 199.*
Q. elongata. Willd. sp. pl. 4. p. 444.
 Icon. *Mich. querc. n. 16. t. 28. Mich. arb. 2. p. 104. t. 21.*

In sandy soil, near the sea-coast: New Jersey to Georgia. $\frac{1}{2}$. May. v. v. A very large tree, commonly called *Spanish Oak*. In the southern states it is known by the name of *Red Oak*.

23. Q. foliis longe petiolatis oblongis profunde sinuatis glabris, axillis venarum subtus villosis, lobis divaricatis dentatis acutis setaceo-mucronatis, cupula scutellata laevi, glande subglobosa.—*Willd. sp. pl. 4. p. 446.*
 Icon. *Mich. querc. n. 19. t. 33. 34. Mich. arb. 2. p. 123. t. 25. Du Roi harbk. 2. t. 5. f. 4. Wangh. amer. t. 5. f. 10.*

In low swampy woods: New England to Pennsylvania, and in Illinois. $\frac{1}{2}$. May. v. v. The *Swamp Spanish Oak* or *Pin Oak* is a large tree; leaves rather small and handsomely divided; the acorns small, and generally in great abundance.

24. Q. foliis longe petiolatis obovato-cuneiformibus tri-quinquelobisve margine integerrimis subtus cinereo-tomentosis, lobis setaceo-mucronatis, cupula subturbinate, glande subglobosa.—*Mich. fl. amer. 2. p. 199.*

Q. ilicifolia. Willd. sp. pl. 4. p. 447. Wangh. amer. 79. t. 6. f. 17.
 Icon. *Mich. querc. n. 15. t. 27. Mich. arb. 2. p. 96. t. 19. Abbot. insect. 2. t. 79?*

In dry barren fields and on the mountains: New Jersey to Virginia. $\frac{1}{2}$. May. v. v. This shrub, about four or six feet high, covers large tracts of ground wherever it occurs, called *Oak-barrens*: it is known by the name of *Bear Oak*, *Black Scrub Oak*, and *Dwarf Red Oak*.

falcata.

palustris.

Banisteri.

- DICT
- ** *Fructificatio annua; fructibus pedunculatis; foliis muticis.*
- † *Foliis lobatis.*
- obtusiloba.* 25. Q. foliis oblongis sinuatis basi cuneatis subtus pubescens-tibus, lobis obtusis superioribus dilatatis bilobis, cupula hemisphaerica, glande ovali.—*Mich. fl. amer.* 2. p. 194.
Q.stellata. *Willd. sp. pl.* 4. p. 452. *Wangh. amer.* 78. t. 6. f. 15.
Icon. *Mich. querc. n.* 1. t. 1. *Mich. arb.* 2. t. 4.
In most forests: Canada to Florida. ♂. May. v. v. The *Upland White Oak* or *Iron Oak* is a spreading tree about fifty or sixty feet high; its timber is of great value in ship-building.
- macrocarpa.* 26. Q. foliis subtus tomentosis profunde lyratim sinuato-lobatis, lobis obtusis repandis, superioribus dilatatis, cupula craterata; squamis superioribus setosis, glande turgide ovata.—*Willd. sp. pl.* 4. p. 453.
Icon. *Mich. querc. n.* 2. t. 2. ♂ 3. *Mich. arb.* 2. t. 3.
Within the mountains, on dry slate or lime-stone hills: Kentucky, Tennessee, Illinois, Mississippi, and Missouri. ♂. May. v. v. A large tree, called *Overcup White Oak* by the inhabitants; the fruit is the largest of the American species; the wood very excellent.
- olivæformis.* 27. Q. foliis oblongis glabris subtus glaucis profunde inæquilater sinuato-pinnatifidis, fructu elliptico-ovato, cupula profunde craterata superne crinita, glande elliptico-ovali.—*Mich. arb.* 2. p. 32.
Icon. *Mich. arb.* 2. t. 2.
On the banks of Hudson's river and the western parts of New York. *Michaux.* In Pennsylvania and Virginia, on iron-ore hills. Ph. ♂. May. v. v. The *Mossy-cup Oak* is a large tree; the foliage handsome, and somewhat resembling the preceding species.
- lyrata.* 28. Q. foliis brevissime petiolatis glabris lyrato-sinuatis summitate dilatatis, lobis oblongis acutis superioribus angulato-truncatis, cupula depresso-globosa tuberculoso-muricata glande globosa subtecta.—*Willd. sp. pl.* 4. p. 453. *Walt. fl. car.* 235.
Icon. *Mich. querc. n.* 3. t. 4. *Mich. arb.* 2. p. 42. t. 5.
In swamps: Carolina to Florida, and on the Mississippi. ♂. May. v. s. This oak is from eight to fifteen feet

- high, and *Swamp-oak* is almost en
29. Q. foliis oblongibus, lobis li attenuatis, i tuberculosa, *pl.* 4. p. 449
Icon. *Mich. querc. n.* 4. t. 1. *Du Roi harbk Catesb. car.*
β. Q. foliis levissimis t. 5. f. 2.
In fertile forest *sylvana* and *Oak* is one of genus in America immense size
- ††
30. Q. foliis longe patentibus grosse d apice callosis, ovata.—*Willd. sp. pl.* 4. p. 450
Q. *Prinus palustris* *Mich. que Roi harbk.* 2. *Catesb. car.* 1.
In low shady woods *vania* to Florida useful tree is *Swamp Oak*, *Swampy Oak*. The fruit
31. Q. foliis brevi-petiolatis tomentosis grossa inæqualibus dilatibus geminis lobi terminante, eu ovata.—*Willd. sp. pl.* 4. p. 451
Q. *Prinus tomentosa* *Mich. que Roi harbk.* 2. *Catesb. car.* 1.
In low wet woods v. v. The *Sweet Oak*; the acorn, sweet.

high, and known under the names *Over-cup Oak*, *Swamp-post Oak*, and *Water White Oak*. The fruit is almost entirely covered with the cupula or calyx.

29. Q. foliis oblongis pinnatifido-sinuatibus subtus pubescentibus, lobis linear-lanceolatis obtusis integerrimis basi attenuatis, fructibus pedunculatis, cupula subcraterata tuberculosa, basi plana, glande ovata.—*Willd. sp. pl. 4. p. 449.*

Icon. *Mich. querc. n. 4. t. 5.* *Mich. arb. 2. p. 13. t. 1.*
Du Roi harbk. 2. t. 5. f. 5. *Wangh. amer. t. 3. f. 6.*
Catesb. car. 1. t. 21. f. 2.

- β. Q. foliis levissime lobatis utrinque viridibus. *Mich. l. c. t. 5. f. 2.*

In fertile forests: New England to Carolina. β. Pennsylvania and Carolina. h. May. v. v. The *White Oak* is one of the most abundant and useful of this genus in America; it grows in the middle states to an immense size.

†† *Foliis integris, dentatis.*

30. Q. foliis longe petiolatis obovatis acutis subtus pubescens-tibus grosse dentatis, dentibus subæqualibus dilatatis apice callosis, cupula craterata basi attenuata, glande ovata.—*Willd. sp. pl. 4. p. 439.*

Q. *Prinus palustris.* *Mich. fl. amer. 2. p. 196.*
 Icon. *Mich. querc. n. 5. t. 6.* *Mich. arb. 2. t. 7.* *Du Roi harbk. 2. t. 6. f. 3.* *Wangh. amer. t. 4. f. 8.*
Catesb. car. 1. t. 18. *Pluk. alm. t. 54. f. 3.*

In low shady woods and on the banks of rivers: Pennsylvania to Florida. h. May. u. v. This large and useful tree is known by the name of *Chesnut White Oak*, *Swamp Chesnut Oak*, and, to the south, *White Oak*. The fruit is large, and of a sweet taste.

31. Q. foliis brevi-petiolatis oblongo-obovatis subtus albotomentosis grosse dentatis basi integerrimis, dentibus inæqualibus dilatatis acutiusculis apice callosis, fructibus geminis longe pedunculatis, pedunculo in setam terminante, cupula hemisphærica, glande oblongo-ovata.—*Willd. sp. pl. 4. p. 440.*

Q. *Prinus tomentosa.* *Mich. querc. n. 5. t. 9. f. 2.*
 Q. *Prinus discolor.* *Mich. arb. 2. p. 46. t. 6.*

In low wet woods: Pennsylvania to Carolina. h. May. v. v. The *Swamp White Oak* grows to a very large tree; the acorn, as of most of the chesnut oaks, is sweet.

alba.

repanda.

Prinus.

bicolor.

montana.

32. Q. foliis modice petiolatis lato-obovatis oblongis subtus albo-tomentosis supra nitidis grosse dentatis basi obtusis inæqualibus, dentibus subæqualibus obtusissimis brevibus, fructibus geminis brevi-pedunculatis, cupula hemisphærica, squamis tuberculatis rugosis, glande ovata.—*Willd. sp. pl. 4. p. 440.*

Q. *Prinus monticola. Mich. fl. amer. 2. p. 196.*
Icon. *Mich. querc. n. 5. t. 7. Mich. arb. 2. t. 8. Abbot. insect. 2. t. 82.*

In rocky situations on the mountains: New England to Carolina. $\frac{1}{2}$. May. v. v. The *Rock Chesnut Oak* rises to about sixty feet; its timber is in high value, and its bark tans extremely well.

Castanea.

33. Q. foliis longe petiolatis oblongo-lanceolatis basi obtusis acuminatis subtus tomentosis grosse dentatis, dentibus subæqualibus dilatatis acutis apice callosis, cupula hemisphærica, glande ovato-subglobosa.—*Willd. sp. pl. 4. p. 441.*

Q. *Prinus acuminata. Mich. fl. amer. 2. p. 196.*
Icon. *Mich. querc. n. 5. t. 8. Mich. arb. 2. p. 61. t. 9.*
In the Allegany mountains and on the banks of the Delaware. $\frac{1}{2}$. May. v. v. The *Yellow Oak* is a beautiful and large tree; the acorns are eatable.

Chinquapin.

34. Q. foliis brevi-petiolatis obovatis basi acutis grosse dentatis subtus glaucis, dentibus subæqualibus dilatatis apice callosis, cupula hemisphærica, glande ovata.

Q. *Prinus Chinquapin. Mich. arb. 2. p. 65.*
Q. *prinooides. Willd. sp. pl. 4. p. 440.*

Icon. *Mich. querc. n. 5. t. 9. f. 1. Mich. arb. 2. t. 10. bona.*

On dry mountain lands: Pennsylvania to Carolina. $\frac{1}{2}$. May. v. v. The *Chinquapin or Dwarf Chesnut Oak* is a low-growing shrub, not exceeding three or four feet in height; it is highly ornamental when in full bloom.

701. CORYLUS. *Gen. pl. 1450.**americana.*

1. C. foliis subrotundis cordatis acuminatis, calycibus fructus subrotundo-campanulatis nucæ subglobosa majoribus, limbo dilatato dentato-serrato.—*Willd. sp. pl. 4. p. 471.*

Icon. *Wangh. amer. t. 29. f. 63.*
In low shady woods: Canada to Florida. $\frac{1}{2}$. March, April. v. v. The *Hazel Nut or Wild Filbert* is a shrub from four to eight feet high; the nuts very fine.

DICT.

2. C. foliis oblongo-
ceolatis, ca-
majoribus b-
pl. 4 p. 47
Icon. *Willd. a-*
In the mounta-
This species

702. LIG.

1. L. foliis palmata-
seos venarum
Icon. *Mich. ar-*
In low woods,
and in all
The *Sweet-g*
mense ~~size~~,
wood is of a
and furniture
ance.

703. CO.

1. C. foliis longo-
~~L~~—*Willd. sp.*
Liquidambar pe-
Liquidambar asp-
Icon. *Pluk. alm-*
In sandy, stony,
nia. $\frac{1}{2}$. Ma-
Bush rises to a
of the plant w

704. PL.

1. P. foliis quinqua-
mulis albentib.
Icon. *Catesb. car-*
arb. 3. t. 3.
On the banks of
siana. $\frac{1}{2}$. M
name of *Butt*
Plane Tree; in
largest tree in
the Ohio and t
from ten to six

2. *C. foliis oblongo-ovatis acuminatis, stipulis linearis-lan-*
ceolatis, calycibus fructus campanulato-tubulosis nuce
majoribus bipartitis, lacinias inciso-dentatis. Willd. sp.
pl. 4 p. 471.

Icon. *Willd. arb. t. 1. f. 2.*

In the mountains: Canada to Carolina. $\frac{1}{2}$. April. v. v.

This species is not above four or five feet high.

702. LIQUIDAMBAR. *Gen. pl. 1452.*

1. *L. foliis palmatis, lobis acuminatis, serratis, sinibus ba-*
seos venarum villosis.—Willd. sp. pl. 4. p. 475.

Icon. *Mich. arb. 3. p. 194. t. 4.*

In low woods, on fertile soil: New England to Florida, and in all the western countries. $\frac{1}{2}$. May. v. v. The *Sweet-gum Tree* is sometimes found of an immense size, particularly in the southern states; its wood is of an exquisite hard texture and fine grain, and furniture made of it has a very handsome appearance.

703. COMPTONIA. *Gen. pl. 1764.*

1. *C. foliis longo-linearibus alternatim crenato-pinnatifidis. asplenifolia.*
Willd. sp. pl. 4 p. 320.

Liquidambar peregrinum. *Syst. veg. 860.*

Liquidambar asplenifolium. *Sp. pl. 1418.*

Icon. *Pluk. alm. t. 100. f. 6.*

In sandy, stony, or slaty woods: New England to Virginia. $\frac{1}{2}$. March, April. v. v. The *Sweet-fern Bush* rises to about three or four feet high; the whole of the plant when rubbed has a strong resinous scent.

704. PLATANUS. *Gen. pl. 1451.*

1. *P. foliis quinquangularibus obsolete lobatis dentatis, ra-*
mulis albentibus.—Willd. sp. pl. 4. p. 474.

Icon. *Catesb. car. 1. t. 56. Duham. arb. t. 35. Mich.*
arb. 3. t. 3.

On the banks of rivers: Canada to Florida, and in Louisiana. $\frac{1}{2}$. May. v. v. This tree is known by the name of *Button-wood*, *Water Beech*, *Sycamore* and *Plane Tree*; in Canada *Cotton Tree*. It is perhaps the largest tree in North America: on the fertile banks of the Ohio and the Mississippi there are trees measuring from ten to sixteen feet in diameter.

rostrata.

styraciflora.

705. JUGLANS. *Gen. pl.* 1446.

* *Amentis masculis simplicibus, polyandris*: Walnuts,

magra.

1. *J. foliolis numerosis ovato-lanceolatis serratis subcordatis superne angustatis, subitus petiolisque subpubescentibus, fructibus globosis scabro punctatis, nuce corrugata.* — *Willd. sp. pl. 4. p. 456.*

Icon. Mich. arb. 1. p. 157. t. 1. Jacq. ic. rar. 1. t. 191. Wangh. amer. t. 8. f. 20. Catesb. car. 1. t. 67.

In fertile woods: New England to Florida. $\frac{1}{2}$. April, May. v. v. The *Black Walnut* is, like all the following, a tree of large size, and its nuts are eaten by men and several species of animals. The wood is put to various mechanical and æconomical uses; for which information I refer to the different authors on forest trees, but particularly to the works of the younger Michaux.

pinerea.

2. *J. foliolis numerosis lanceolatis serratis basi rotundatis subitus pubescenti-mollibus, petiolis villosis, fructibus oblongo-ovatis apice mammosis viscidis longe penduculatis, nuce oblonga acuminata insigniter insculpta-sebrosa.* — *Willd. sp. pl. 4. p. 456.*

J. cathartica. Mich. arb. 1. p. 165.

J. oblonga. Mill. dict. n. 3. Retz. obs. 1. p. 10.

Icon. Mich. arb. 1. t. 2. Jacq. ic. rar. 1. t. 192. Wangh. amer. t. 9. f. 21.

Near the sea-coast: Canada to Virginia, and on the Allegany mountains. $\frac{1}{2}$. April, May. v. v. It is known under the name of *Butter Nut*, *Oil Nut*, and *White Walnut*. The younger Michaux has altered the adopted name, and substituted one indicating the medicinal use made of it, which certainly would be a very good name if the old one had not been so long established.

** *Amentis masculis compositis, tetrandris*. Hickory.

ellipticiformis.

3. *J. foliolis numerosis (13—15) lanceolatis subfalcatis serratis, impari petiolato, fructibus oblongis tetragonis, nuce olivæformi laevi.* — *Willd. sp. pl. 4. p. 457.*

J. angustifolia. Ait. kew. 3. p. 361.

J. Pecan. Muell. in nov. act. soc. nat. scrut. berol. 3. p. 392.

Icon. Mich. arb. 1. t. 3.

On the banks of the Ohio, Mississippi, and other rivers

in Upper L generally kn Nut; the delicious kei

4. *J. foliolis subn ratis subtus l nuato, fructi bosa leviter sp. pl. 4. p. 4*

J. mucronata.

J. laciniosa. M

Icon. Mich. ar

In fertile valley

May. v. v.

Springfield o

fine tasted;

loose fragme

5. *J. foliolis subse acuminatis ar sili, amentis globosis, nuce*

p. 193. Sp.

J. compressa. H

act. soc. nat. s

J. squamosa. M

Icon. Mich. arl

carp. 2. t. 89.

In shady fertile

throughout

May. v. v.

of *Shell-bark*

kory, on acco

fragments like

a thin shell an

6. *J. foliolis subno ter serratis sub petiolato, am fructibus subg nuce subsexan*

Mich. fl. amer

J. alba. Willd. s

Icon. Mich. arb.

In fertile forests

Allegany moun

in Upper Louisiana. $\frac{1}{2}$. April, May. v. v. It is generally known under the name of *Pecan* or *Illinois Nut*; the fruit is small, with a very thin shell, and a delicious kernel.

4. *J. foliolis subnovenis obovato-lanceolatis acuminatis serratis subtus pubescentibus, impari subsessili basi attenuato, fructibus subrotundis 4-carinatis, nuce subglobosa leviter compressa laevi longe mucronata.* —Willd. sp. pl. 4. p. 457.

suleata.

J. mucronata. Mich. fl. amer. 2. p. 192.

J. laciniosa. Mich. arb. 1. p. 199.

Icon. Mich. arb. 1. t. 8.

In fertile valleys of the Allegany mountains. $\frac{1}{2}$. April, May. v. v. It is called *Thick Shell-bark Hickory*, *Springfield* or *Glocester Nut*. The nuts are large and fine tasted; the bark is torn lengthways in long loose fragments, as in the following species.

5. *J. foliolis subseptenis longe petiolatis oblongo-lanceolatis acuminatis argute serratis, subtus villosis, impari sessili, amentis filiformibus glabris, fructibus depresso-globosis, nuce compressa obliqua.* —Mich. fl. amer. 2. p. 193. Sp. pl. 1415.

alba.

J. compressa. Willd. sp. pl. 4. p. 458. Muhl. in nov. act. soc. nat. scrut. berol. 3. p. 390.

J. squamosa. Mich. arb. 1. p. 190.

Icon. Mich. arb. 1. t. 7. Pluk. alm. t. 309. f. 2. Gært. carp. 2. t. 89. sub *J. compressa*. Catesb. car. 1. t. 38.

In shady fertile forests: New England to Carolina, and throughout the Allegany mountains. $\frac{1}{2}$. April, May. v. v. This useful tree is known by the name of *Shell-bark Hickory*, *Shag-bark* and *Scaly-bark Hickory*, on account of its bark, which is torn in loose fragments like the preceding: its nuts are white, with a thin shell and very sweet and large kernel.

6. *J. foliolis subnovenis oblongo-lanceolatis acuminatis leviter serratis subtus pubescentibus scabris, impari subpetiolato, amentis filiformibus longissimis tomentosis, fructibus subglobosis laevibus, pericarpio crassissimo, nuce subsexangulata putamine crassa durissima.* —Mich. fl. amer. 2. p. 192.

tomentosa.

J. alba. Willd. sp. pl. 4. p. 457.

Icon. Mich. arb. 1. p. 186. t. 6.

In fertile forests: New England to Virginia, and on the Allegany mountains. $\frac{1}{2}$. April, May. v. v. This

is known under the name of *Mocker Nut*, *White-heart Hickory* or *Common Hickory*. The wood is excellent for mechanical purposes, and particularly esteemed as fire-wood; but the nuts are very hard, with but little kernel in them.

amara.

7. J. foliolis subnovenis ovato-oblongis acuminatis argute serratis utrinque glabris, impari breviter petiolato, fructibus subglobosis, superne suturis prominulis, nuce lœvi subglobosa mucronata, putamine fragili.—*Mich. arb.* 1. p. 177.

In dry fertile woods: New England to Maryland, and on the mountains. $\frac{1}{2}$. May. v. v. This is known by the name of *Bitter Nut*, *White* or *Swamp Hickory*. The nuts are small, the kernel bitter.

porcina.

8. J. foliolis subseptenatis lanceolatis acuminatis serratis utrinque glabris, impari subsessili, fructibus pyriformibus vel globosis, nuce lœvi durissima.—*Mich. arb.* 1. p. 206.

- α. J. fructu globoso, nuce obcordata.—*Mich. arb.* 1. p. 206. t. 9. f. 3. ♂ 4.

J. obcordata. *Willd. sp. pl.* 4. p. 458.

- β. J. fructu turbinato, nuce oblonga.—*Mich. arb.* 1. p. 209. t. 9. f. 1. ♂ 2.

J. glabra. *Willd. sp. pl.* 4. p. 458.

In dry fertile forests: New England to Virginia. $\frac{1}{2}$. May. v. v. It is known by the name of *Pig* or *Hog Nut*, also *Broom Hickory*. The natives and inhabitants make brooms of it by slitting the very tough wood into narrow slips, which finally form a very good and durable broom. The nuts are very small and extremely hard.

aquatica.

9. J. foliolis subundenis angusto-lanceolatis acuminatis sub-serratis sessilibus, impari breviter pedicellato, fructibus pedunculatis ovatis suturis 4. prominulis, nuce subdepressa rubiginosa tenera.—*Mich. arb.* 1. p. 182.

Icon. *Mich. arb.* 1. t. 5.

In swamps and rice-fields: South Carolina and Georgia.

$\frac{1}{2}$. May. v. s. Michaux calls it *Water Bitter-nut Hickory*. The fruit is extremely bitter.

myristicæfor-
mis.

10. J. foliolis quinis ovato-lanceolatis acuminatis serratis glabris, impari subsessili, fructibus ovalibus rugoso-scabris, nuce ovali brevi-acuminata sulcato-lineata durissima.—*Mich. arb.* 1. p. 211.

Icon. *Mich. arb.* 1. t. 10.

DIC

In South Ca
Hickory is
and brown

706

1. M. foliis con-
serratis scal-
cylindricis.
Icon. *Mich.* t.
f. 35. *Plu-*
In fertile valle
v. v. The
the wood u

III

707

* *Foliis s*

1. P. foliis solit-
glauca subsp-
tibus, conis
obovatis lon-
pl. 4. p. 50-
Abies balsamit-
Icon. *Lambert-*
In Canada, Ne-
legany mou-
May. v. v.
of *Balsam* or
Silver Fir.
fifty feet.

2. P. foliis solita-
glauca subsp-
oblongis ere-
cuneatis em-
latis.

On high moun-
mountains,
in Herb. La-
inhabitants

In South Carolina. *Michaux.* h. +. The *Nutmeg Hickory* is but little known; the nuts are very hard, and brown with white lines.

706. MORUS. *Gen. pl. 1424.*

1. M. foliis cordatis ovatis acuminatis trilobisve, æqualiter serratis scabris, subtus pubescentibus, amentis femineis cylindricis.—*Willd. sp. pl. 4. p. 369.*

Icon. Mich. arb. 3. p. 232. t. 10. Wangh. amer. t. 15. f. 35. Pluk. alm. t. 246. f. 4.

In fertile valleys throughout the middle States. h. June, v. v. The *Red Mulberry* is a very handsome tree, and the wood useful and very durable.

rubra.

III. CONIFERÆ.

707. PINUS. *Gen. pl. 1451.*

* *Foliis solitariis, basi distinctis.* Abies.

1. P. foliis solitariis planis emarginatis integrisve subtus glaucis subpectinatis supra suberectis recurvato-patentibus, conis cylindraceis erectis, bracteolis abbreviatis obovatis longe mucronatis subserrulatis.—*Willd. sp. pl. 4. p. 504.*

Abies balsamifera. Mich. fl. amer. 2. p. 207.

Icon. Lambert. monogr. p. 48. t. 31. Mich. arb. 3. t. 14.

In Canada, Nova Scotia, New England, and on the Allegany mountains, in high and cold situations. h. May. v. v. This elegant tree is known by the name of *Balsam of Gilead Fir*, *Fir Balsam*, and *American Silver Fir*. It grows to the height of about forty or fifty feet.

Balsamea.

2. P. foliis solitariis planis brevioribus emarginatis subtus glaucis subsecundis supra crebris erectis, conis ovato-oblongis erectis, bracteolis elongatis reflexis oblongo-cuneatis emarginatis brevi-mucronatis inciso-denticulatis.

On high mountains of Carolina. *Fraser.* On the Broad-mountains, Pennsylvania. Ph. h. May. v. v.; v. s. in *Herb. Lambert.* This species, known among the inhabitants by the name of *Double-balsam Fir*, re-

Fraseri.

sembles the preceding in several respects, but differs at first sight in being a smaller tree, the leaves shorter and more erect, and the cones not one fourth the size. Messrs. Frasers introduced this tree into England a few years ago.

taxifolia.

3. *P. foliis solitariis planis subdistichis, conis oblongis, antheris didymis.*—*Lambert. monogr. p. 51. t. 33.*

On the banks of the Columbia. *M. Lewis.* On the north-west coast. *Menzies.* $\frac{1}{2}$. *v. s. in Herb. Lewis.* This elegant and tall tree has some resemblance to the following one, but the leaves are more than twice the length. I have among my specimens two varieties, or probably distinct species, which for want of fructification I cannot decide: one has acute leaves, green on both sides; the other emarginate leaves, glaucous underneath.

canadensis.

4. *P. foliis solitariis planis denticulatis subdistichis, strobilis ovatis terminalibus vix folio longioribus.* *Lambert. monogr. p. 50. t. 32.*

P. americana. Du Roi harlk. 2. p. 107.

Icon. Lambert. l. c. Mich. arb. 1. p. 137. t. 13. Wangh. amer. t. 15. f. 36.

In the most northern regions of Canada and on the highest mountains, as far south as Carolina. $\frac{1}{2}$. May. *v. v.* The *Hemlock Spruce* is a very elegant tree, and grows in some situations to an enormous size; its bark is a fine substitute for oak-bark in tanning.

nigra.

5. *P. foliis solitariis tetragonis undique sparsis erectis strictis strobilis ovatis, squamis ellipticis margine undulatis apice eroso-denticulatis.*—*Lambert. monogr. p. 41. t. 27.*

P. mariana. Du Roi harlk. 2. p. 107.

P. denticulata. Mich. fl. amer. 2. p. 206.

Icon. Lambert. l. c. Mich. arb. 1. t. 11. Mill. ic. 1. t. 1.

From the more northern parts of Canada to Carolina, throughout the tracts of high mountains. $\frac{1}{2}$. May. *v. v.* The *Black Spruce* or *Double Spruce* is, besides its great mechanical use, the tree of which that wholesome beverage called *Spruce Beer* is made.

rubra.

6. *P. foliis solitariis subulatis, strobilis oblongis obtusis, squamis rotundatis subbilobis margine integris.* *Lambert. monogr. p. 43. t. 28.*

Abies pectinata. Lamarck. encycl. 6. t. 523.

Icon. *Lambert.*
In Nova Scoti
This specie
ceding. It
dens.

7. *P. foliis solit
dricis laxis,
monogr. p. 1
P. laxa. Eh.
P. canadensis.
Icon. *Lambert*
mer. 5. t. 1.
In similar sit
The *White
Spruce*, its
unfit for ma*

** *Foliis*

8. *P. foliis brevib
nicis longitu
latis rectis.—
P. virginiana.
Icon. *Lambert*
In dry barren s
v. v. The
middle size,
branches are
might be use
was not so ap
in his magi
and useful ge
the younger l
taken up as
Mr. Lambert
pecting this
Michaux, wh
in the follow
Mr. Michaux
are so inaccur
notice before
vent Mr. Pur
believe he is a
therefore beg
on the differen
belong, if th*

fers
ter
ize.
d a

an-
the
rb.
m-
ore
ens
for
ute
ate

ilis
rt.

gh.
ay.
nd
its

tis
tis
fl.

1.

ia,
ly.
es
e-

is,
n-

Icon. *Lambert. l. c.*

In Nova Scotia and about Hudson's Bay. $\frac{1}{2}$. May. v. s.

This species of *Spruce Fir* is nearly related to the preceding. It is, as yet, a very scarce tree in the gardens.

7. *P. foliis solitariis tetragonis incurvis, strobilis subcylindricis laxis, squamis obovatis integerrimis.* *Lambert. monogr. p. 39. t. 26.*

P. laxa. *Ehrh. beitr. 3. p. 24.*

P. canadensis. *Du Roi harbk. 2. t. 124.*

Icon. *Lambert. l. c.* *Mich. arb. 1. t. 12.* *Wangh. amer. 5. t. 1. f. 2.*

In similar situations with *P. nigra*. $\frac{1}{2}$. May. v. v.

The *White Spruce* is of lower growth than the *Black Spruce*, its wood not so good as that, and its branches unfit for making spruce beer.

alba.

** *Foliis pluribus, basi vaginatis.* *Pinus.*

8. *P. foliis brevibus geminis, strobilis recurvis oblongo-conicis longitudine foliorum, aculeis squamarum subulatis rectis.* — *Lambert. monogr. p. 18. t. 13.*

P. virginiana. *Du Roi harbk. 2. p. 35.*

Icon. *Lambert. l. c.* *Mich. arb. 1. t. 4.*

In dry barren soil: New Jersey to Carolina. $\frac{1}{2}$. May.

v. v. The *Jersey Pine*, *Pitch* or *Scrub Pine*, is of middle size, straggling growth, and full of resin. Its branches are tougher than any other pine I know, and might be used for several useful purposes if its wood was not so apt to an early decay. A. B. Lambert, Esq. in his magnificent Monograph on this interesting and useful genus, made similar observations, which the younger Michaux in his *Arbres forestiers*, p. 60. has taken up as a point of criticism. On this subject Mr. Lambert favoured me with his observations respecting this and some other remarks made by Mr. Michaux, which I think proper here to insert; it is in the following words: "The criticisms made by Mr. Michaux on my 'Description of the genus Pinus' are so inaccurate, that I did not think them worth my notice before, nor should I now, were it not to prevent Mr. Pursh from being misled by them, though I believe he is already well aware of their fallacy: I shall therefore beg of him to insert a few observations on the different subjects to which they respectively belong, if they meet with his approbation and con-

inops.

currence. Whether Mr. Michaux has been successful in changing the specific names of the pines, I shall leave to future botanists to decide: but I must correct him in respect to the *specific name* he has added to my own, which certainly does not belong to it." The observation which Mr. Lambert makes in this place is, " Michaux's observations relative to the branches of *P. inops* being not fit for hoops appear to be mere matter of opinion, as well as mine, without experience."

resinosa.

9. *P. foliis elongatis geminatis, vaginis elongatis, strobilis ovato-conicis basi rotundatis subsolitariis folio dimidio brevioribus, squamis medio dilatatis inermibus.*—*Lambert. monogr. p. 20. t. 14.*

P. rubra. Mich. arb. p. 45. t. 1.

In Canada and the western parts of New York. $\frac{1}{2}$. v. v. This tree, called by Aiton *Pitch Pine*, is generally known in its native country by the name of *Norway Pine*; sometimes, particularly among the Canadian French, *Red Pine*. It grows in close forests, is very tall, and its bark is remarkably smooth and red; the timber is very heavy, for which reason it is rejected for masts, though its size and shape are apparently very recommendable for that purpose. Mr. Lambert in his note, quoted above, observes, " In confirmation of what I have said relative to the timber of *P. resinosa*, Mr. Michaux mentions the exportation of it being much diminished, as it has been found to contain too much sap; but he still thinks, from some planks he has seen, it is without foundation."

Banksiana.

10. *P. foliis brevibus geminatis rigidis divaricatis obliquis, strobilis recurvis tortis, squamis inermibus.*—*Lambert. monogr. p. 7. t. 3.*

P. ropestris. Mich. arb. 1. p. 49. t. 2.

P. hudsonia. Lam. encycl. 5. p. 339.

P. sylvestris ð. *divaricata. Ait. kew. 3. p. 366.*

In cold barren and rocky situations: Nova Scotia, Hudson's Bay, parts of Canada and Province of Maine. $\frac{1}{2}$. April, May. v. v. The *Scrub Pine*, or, as Michaux calls it, the *Gray Pine*, is a small straggling tree, which in some instances, when growing among the barren rocks, does not rise above five or eight feet high, though it will grow to a considerable size when, by accident or culture, it is brought on good soil. Mr. Lambert observes in his note, " that the trees of this

species now sin from the

11. *P. foliis elongatis, strobilis, aculeis incurvatis, squamis medio dilatatis inermibus.*—*Mitch. In most pine-trees May. v. v. building of note on this P. variabilis saw it grow doubtless his*

12. *P. foliis ternis erecto-incunabulis, gregatis, squamis medio dilatatis inermibus.*—*Lambert. monogr. p. 25. Icon. Lambert. On the plains v. v. The favourable situation promiscuous Mr. Lambert were taken f.*

13. *P. foliis elongatis, bidentatis, strigatis, squamis medio dilatatis inermibus.*—*P. Tæda alopec. Mich. arb. f. 5. Icon. Lambert. On the edges of Carolina. $\frac{1}{2}$ species ripen*

I strongly suspect had not been figure of Mi such.

14. *P. foliis geminatis, aculeis inferioribus inermibus.*—*Lambert. t. 5. On high mountains*

species now in England exude a great quantity of resin from their branches."

11. *P. foliis elongatis binis ternatisque tenuibus canaliculatis, strobilis ovato-conicis subsolitariis, squamarum aculeis incurvis.*—*Lambert. monogr. p. 22. t. 15.*

P. mitis. Mich. arb. 1. p. 52. t. 3.

In most pine-forests: New England to Georgia. $\frac{1}{2}$. May. v. v. The Yellow Pine is the most in use for building of houses as well as shipping. Mr. Lambert's note on this species says, "The height and size of *P. variabilis* I have taken from Wangenheim, who saw it growing in America, and see no reason for doubting his authority."

12. *P. foliis ternis, vaginis abbreviatis, amentis masculis erecto-incubentibus, strobilis ovatis sparsis vel aggregatis, squamarum spinis reflexis.*—*Lambert. monogr. p. 25. t. 18. 19.*

Icon. *Lambert. l. c. Mich. arb. 1. p. 89. t. 8.*

On the plains: New England to Virginia. $\frac{1}{2}$. May. v. v. The common Black or Pitch Pine grows in favourable situations to a very large tree; it is found promiscuously in dry soil and very wet low grounds. Mr. Lambert says that his observations on this species were taken from Wangenheim.

13. *P. foliis elongatis ternis, amentis masculis erecto-incubentibus, strobilis ovatis, squamarum aculeis rectis tenuissimis.*—*Mich. fl. amer. 2. p. 205.*

P. Tæda alopecuroides. Ait. kew. 3. t. 368.

Icon. *Mich. arb. 1. p. 86. t. 7. Lambert. monogr. t. 19. f. 5.*

On the edges of ponds and swamps: New Jersey to Carolina. $\frac{1}{2}$. May. v. v. This and the preceding species ripen their seeds only after the second year. I strongly suspect them to be only varieties; and if it had not been introduced as a species by a very good figure of Michaux's, I should have only given it as such.

14. *P. foliis geminis brevibus acutis, strobilis ovato-conicis, aculeis squamarum elongatis subulatis incurvis, inferioribus reflexis.*—*Lambert. monogr. p. 91.*

Icon. *Lambert. monogr. t. 16. f. c. Mich. arb. 1. p. 61. t. 5.*

On high mountains: on the Grandfather and Table

variabilis.

rigida.

serotina.

pungens.

Tæda.

mountains, Carolina. $\frac{1}{2}$. v. s. in *Herb. Lambert.* The cones are large and armed with strong spines. The cone figured t. 16. f. c. in *Lambert. monogr.* is genuine, but the branch belongs to a separate species.

15. *P. foliis elongatis ternis, vaginis elongatis, strobilis oblongo-conicis deflexis folio brevioribus, spinis inflexis.*
—*Lambert. monogr.* p. 23. t. 16. 17.

Icon. *Lambert. l. c.* *Mich. arb. 1.* p. 97. t. 9.

In barren and sandy situations: Virginia to Florida. $\frac{1}{2}$. April, May. v. v. The *Loblolly* or *Old-field Pine* is found in large tracts in the southern states: all the woods seem to be seeded with it; for when any piece of cleared land is neglected for any space of time it will be covered with those pines. It is difficult and in some cases almost impracticable to recover those lands run over with young pines, as the ground appears to have lost all fertile properties for any other vegetable than those trees. The account Mr. Michaux gives of this tree is very correct and instructive; as also is the plate of Mr. Lambert; but unfortunately the latter, considering *P. tæda Wangh. amer. 41.* a genuine synonym, although it evidently belongs to *P. rigida* or *serotina*, has inadvertently been led into some errors in his description.

palustris.

16. *P. foliis ternis longissimis, stipulis pinnatifidis ramentaceis persistentibus, strobilis subcylindraceis muricatis.*
—*Lambert. monogr.* p. 27. f. 20.

P. australis. Mich. arb. 1. p. 64. t. 6.

Icon. *Lambert. et Mich. l. c.* *Abbot. insect. 1.* t. 42.

In the forests, near the sea-coast: North Carolina to Florida. $\frac{1}{2}$. May. v. v. The *Long-leaved, Yellow, Pitch, or Broom Pine* is a beautiful as well as very useful tree. Mr. Lambert's remarks respecting the quality of the wood he has taken from Wangenheim, whom he considers sufficient authority; but on the observations of Michaux respecting the male flowers, he mentions in his note, "I can only suppose Mr. Michaux has never seen my plate with the male flowers, or he never would have made such an observation about them."

Strobus.

17. *P. foliis quinis gracilibus, vaginis brevissimis, strobilis pendulis cylindraceis folio longioribus, squamis laxis.*
—*Willd. sp. pl. 4.* p. 501. *Lambert. monogr.* p. 31. t. 22.

Icon. *Lambert. amer. 1. t. 1.*
In fertile soil, $\frac{1}{2}$. May.
the largest & mentioned;
enormous size
masts.

18. *P. foliis fasciculatum marginib. acuminata.*
Ait. kew. 3. p.

In low cedar April, May.
Hack-matack resembling the in the exceller

19. *P. foliis fasciculata floris, squamis minatis.* *La*
P. pendula. Wil
P. larcina. Du
Larix americana.
Icon. *Lambert. l*
t. 16. f. 37.

About Hudson's York and Pens Larch resemble considered as specifically and both growing in other. Mr. La always kept dis

708. CUI

1. *C. foliis distichis pilo-paniculatis, sp. 512.*

Icon. *Mich. arb. 3*
Comm. hort. 1.

In extensive swan from Indian-rive

Icon. *Lambert. l. c. Mich. arb. 1. p. 103. t. 10. Wangh. amer. 1. t. 1. f. 1.*

In fertile soil, on the side of hills: Canada to Virginia. $\frac{1}{2}$. May. v. v. The *White* or *Weymouth Pine* is the largest and most useful of all the species above mentioned; in the State of Vermont they grow to an enormous size; it is the best timber in America for masts.

*** *Foliis fasciculatis.* Larix.

18. *P. foliis fasciculatis deciduis, strobilis oblongis, squamorum marginibus inflexis, bracteolis panduræformibus acumine attenuato.* *Lambert. monogr. p. 55. t. 36. Ait. kew. 3. p. 369.*

In low cedar swamps: Canada to New Jersey. $\frac{1}{2}$. April, May. v. v. The *Black Larch*, *Tamarack*, or *Hack-matack* of the Americans, is a beautiful tree, resembling the European Larch in appearance, as well as in the excellent qualities of its wood and bark.

19. *P. foliis fasciculatis deciduis, strobilis subrotundis paucifloris, squamis reflexis, bracteolis ellipticis obtuse acuminatis.* *Lambert. monogr. p. 56. t. 37.*

P. pendula. *Willd. arb. 215.*

P. laricina. *Du Roi harbk. 2. p. 83.*

Larix americana. *Mich. fl. amer. 2. p. 203.*

Icon. *Lambert. l. c. Mich. arb. 3. t. 4. Wangh. amer. t. 16. f. 37.*

About Hudson's Bay and on high mountains of New York and Pennsylvania. $\frac{1}{2}$. May. v. v. The *Red Larch* resembles the preceding, and both have been considered as one species by Michaux; but they are specifically and constantly different; I never saw them both growing in the same place, or even near one another. Mr. Lambert likewise observes that they have always kept distinct, when raised from seed.

708. CUPRESSUS. *Gen. pl. 1458.*

1. *C. foliis distichis planis deciduis, floribus masculis aphyllo-paniculatis, strobillis subglobosis.—Willd. sp. pl. 4. p. 512.*

Icon. *Mich. arb. 3. p. 4. t. 1. Catesb. car. 1. t. 11. Comm. hort. 1. t. 59. Fluk. alm. t. 85. f. 8.*

In extensive swamps and on the banks of large rivers from Indian-river Delaware to Florida, and on the Mis-

pendula.

microcarpa.

disticha.

thyoides.

sissippi. h . May. *v. v.* The *Bald* or *Deciduous Cypress* is one of the largest trees of the New Continent. The extensive use made of its wood, and other particulars respecting this interesting tree, are fully elucidated in the excellent work of the younger Michaux, who paid particular attention to the use and properties of the different American forest trees.

2. C. ramulis compressis, foliis quadrifariam imbricatis ovatis basi tuberculatis, strobilis globulosis.—*Willd. sp. pl. 4. p. 512.*
Icon. *Mich. arb. 3. p. 20. t. 2.* *Wangh. amer. t. 2. f. 4.*
Pluk. mant. t. 345. f. 1.

In deep swamps: New England to Carolina. h . April, May. *v. v.* The *White Cedar*, so well treated upon by Michaux, is the principal tree, composing what is called *Cedar-swamps*, throughout the middle and southern States.

709. THUYA. *Gen. pl. 1457.**occidentalis.*

1. T. ramulis ancipitibus, foliis quadrifariam imbricatis ova-to-rhombeis adpressis nudis tuberculatis, strobilis obovatis, squamis interioribus truncatis, infra apicem gibbosus. *Willd. sp. pl. 4. p. 508.*
Icon. *Mich. arb. 3. p. 29. t. 3.* *Wangh. amer. t. 2. f. 3.*
From Canada to the mountains of Virginia and Carolina.
 h . May. *v. v.* The *American Arbor Vitæ* is rather scarce in the southern States, and only found on the steep banks of mountain torrents. The branches are extremely tough.

710. JUNIPERUS. *Gen. pl. 1552.**communis.*

1. J. foliis ternis patentibus mucronatis bacca longioribus. *Willd. sp. pl. 4. p. 853.*

α . J. ramis erectis.
 β . J. ramis depressis.

α . About rocks, near the falls of rivers, in Canada and the western part of New York. β in New York, and particularly in the province of Maine, in rocky or gravelly situations. h . May. *v. v.* The *Common Juniper* may probably have been originally brought from Europe; but the variety β ., or probably a distinct species, seems to be really an original native; it does not grow above a foot or two high, and one single

DICI

root will cov
twenty feet i

2. J. foliis ternis |

ribus patulis.

Icon. *Mich. ar*

In dry and roc

to Georgia.

useful and du

to Michaux's

abundance in

termination is

tants will be

want of it wh

3. J. foliis oppositi

imbricatis, te

- β
- .
- Willd. sp. pl. 4.*

In the clefts of

the Rocky-mo

Lewis. Note

4. J. foliis oppositis

drifariam imbr

arboreo. *Wil**J. Sabina varietas*

On the banks o

M. Lewis. h

elegant tree,

of A. B. Lam

those collected

5. J. foliis omnibus

tis, senioribus a

Icon. *Pluk. alm.*

On the coast of F

711. T

1. T. foliis linearibus lis masculis globosus. *T. baccata minor.*
In shady, rocky banks of the rivers. h . April. *v. v.* They cover a great portion of the surface of the ground, under the shade of large trees, two or three feet high.

- root will cover sometimes a space of from fifteen to twenty feet in diameter.
2. J. foliis ternis basi adnatis, junioribus imbricatis, senioribus patulis. *Willd. sp. pl. 4. p. 863.*
Icon. *Mich. arb. 3. p. 42. t. 5. Wangh. amer. t. 2. f. 51.*
In dry and rocky woods and fields: Province of Maine to Georgia. $\frac{1}{2}$. May. v. v. The *Red Cedar*, so useful and durable a wood, for whose history I refer to Michaux's work so often quoted, is as yet in great abundance in most parts of that country; but its extermination is going on so rapidly, that future inhabitants will be very much at a loss, and will feel the want of it when it is too late.
3. J. foliis oppositis obtusis medio glandulosis, quadrisariam imbricatis, tenellis acutis oppositis, caule fruticoso. *Sabina.*
 β . *Willd. sp. pl. 4. p. 852.*
In the clefts of rocks: Canada. *Michaux. β .* Within the Rocky-mountains. *M. Lewis. $\frac{1}{2}$. v. s. in Herb. Lewis.* Not above six inches high.
4. J. foliis oppositis obtusiusculis, medio glandulosis, quadrisariam imbricatis, tenellis acutis ternis patulis, caule arboreo. *Willd. sp. pl. 4. p. 852.*
J. Sabina varietas. Pall. ross. 2. p. 15.
On the banks of the waters of the Rocky-mountains. *M. Lewis. $\frac{1}{2}$. May. v. s. in Herb. Lewis.* A lofty elegant tree. Specimens of Pallas in the Herbarium of A. B. Lambert, Esq. seem to be the same with those collected by Mr. Lewis.
5. J. foliis omnibus quadrisariam imbricatis, junioribus ovatis, senioribus acutis. *Willd. sp. pl. 4. p. 851.*
Icon. *Pluk. alm. t. 197. f. 4.*
On the coast of Florida. *Michaux. $\frac{1}{2}$. v. v. in Hortis. barbadensis.*

711. TAXUS. *Gen. pl. 1553.*

1. T. foliis linearibus distichis margine revolutis, receptaculis masculis globosis. *Willd. sp. pl. 4. p. 856.*
T. baccata minor. Mich. fl. amer. 2. p. 245.
In shady, rocky places: Canada. *Michaux.* On the banks of the Antietum, Maryland. Ph. March, April. v. v. The plants I observed on the Antietum cover a great part of the rocky banks of that river. Under the shade of other trees it does not rise above two or three feet.

712. ZAMIA. *Gen. pl. 1700.**integrifolia.*

1. *Z. frondibus pinnatis, foliolis lanceolatis rotundato-obtusis basi attenuatis latere exteriore ad apicem serrulatis, stipite glabro subtetragono.* Willd. *sp. pl. 4. p. 847.*
Z. pumila. Sp. pl. 1659. excl. synon.

Icon. *Jacq. ic. 3. t. 635.*

In East Florida. $\frac{1}{2} \text{ ft. +}$. This species of palm is only found in Florida, as I have made all inquiries to find it in Georgia, but without success.

XX.

I

713. EQUISETU
Indusium co

II. S

714. LYCOPODI
2-valves, po
riformia.715. BERNHARD
loculis super716. OPHIOGLO
in spicam ar
transverse de717. BOTRYCHI
natae in racen
valvulis binia
tibus.

III. SC

718. HYDROGLO
biseriales, late
centes. *Indu*
obtegens.719. SCHIZÆA.
gatae. *Capsula*
poro oblongo
e margine infl720. OSMUNDA.
striatae, semib

XX. CRYPTOGAMIA.

I. GONOPTERIDES.

713. EQUISETUM. *Recept.* peltata, pentagona, spicata.
Indusium corniculatum. *Stam.* 4. *Styl.* 0. *Sem.* 1.

II. STACHYOPTERIDES.

714. LYCOPODIUM. *Capsulae* reniformes, 1-loculares, 2-valves, polyspermæ. *Sem.* minutissima, pulv- riformia.
715. BERNHARDIA. *Capsulae* 3-coccæ, 3-loculares : loculis superne dehiscentibus, semibivalvibus.
716. OPHIOGLOSSUM. *Capsulae* nudæ, uniloculares, in spicam articulatam disticham connatae, 2-valves, transverse dehiscentes.
717. BOTRYCHIUM. *Capsulae* subglobosæ, rachi adun- natae in racemum compositum-dispositæ, 1-loculares : valvulis binis postice connexis transverse dehis- centibus.

III. SCHISMATOPTERIDES.

718. HYDROGLOSSUM. *Spicæ unilaterales.* *Capsulae* biseriales, latere interiore a basi ad verticem dehis- centes. *Indusium squamiforme* quamlibet capsulam obtegens.
719. SCHIZÆA. *Spicæ unilaterales*, flabellatæ, aggre- gatae. *Capsula* vertice radiatim striatæ, subturbinatæ, poro oblongo latere hiantes. *Indusium continuum* e margine inflexo spicæ formatum.
720. OSMUNDA. *Capsulae* subglobosæ, pedicellatæ, striatæ, semibivalves, paniculatæ. *Indusium nullum.*

IV. FILICES.

721. ACROSTICHUM. *Capsulæ* sparsæ, discum totum inferiorem frondis vel ejus partem occupantes. *Indusium* nullum.
722. POLYPODIUM. *Sori* subrotundi, sparsi. *Indusia* nulla.
723. WOODSIA. *Sori* subrotundi, sparsi. *Indusium* calyciforme, apertum, margine crinitum, includens *Capsulas* pedicellatas.
724. ASPIDIUM. *Sori* subrotundi, sparsi. *Indusium* umbilicatum vel uno latere dehiscens.
725. ONOCLEA. *Capsulæ* partem frondis inferiorem dense tegentes. *Indusia* squamiformia, in formam baccae connata non dehiscentia.
726. STRUTHIOPTERIS. *Capsulæ* partem frondis inferiorem dense tegentes. *Indusia* squamiformia, marginalia, interius dehiscentia.
727. ASPLENIUM. *Sori* lineares, transversales, sparsi. *Indusia* e venis lateralibus orta, costam versus dehiscentia.
728. SCOLOPENDRIUM. *Sori* lineares, transversales, sparsi. *Indusium* duplex, ad utrumque latus sori situm.
729. PTERIS. *Sori* continui, lineares, marginales. *Indusium* e margine frondis, inflexum, interius dehiscens.
730. VITTARIA. *Sori* lineares, longitudinales, continui vel in margine vel in disco frondis. *Indusium* duplex, ad utrumque latus sori.
731. BLECHNUM. *Sori* lineares, longitudinales, continui, ad utrumque latus costæ frondis. *Indusium* superficiarium, interius dehiscens.
732. WOODWARDIA. *Sori* oblongi, distincti, recti, ad utrumque latus costæ frondis. *Indusia* superficiaria, fornicata, interius dehiscentia.
733. ADIANTUM. *Sori* oblongi vel subrotundi, *Indusia* membranaceis, e margine ortis, interius debiscentibus inserti.
734. CHEILANTHES. *Sori* punctiformes, discreti, marginales, tecti *Indusio* squamiformi marginali interius dehiciente.
735. DICKSONIA. *Sori* punctiformes, marginales, subrotundi, distincti. *Indusium* duplex, alternum superficiarium, exterius dehiscens, alterum marginale interius dehiscens.

736. HYMENO
culo cylind
includens.

- V.
737. ISOETES.
frondis basi
receptaculis
738. SALVINIA.
uniloculare
trali inserta.
739. AZOLLA.
polysperma.

I. GO

713. EQ

1. E. caulinis sterili
culis tetragoni
cylindraceis in
enum. 1065.
Icon. Schkuhr fil
In low and sandy
April, May.
2. E. caulinis sterili
sis, ramis scat
subtriquetris,
p. 3.
Icon. Schkuhr fil
In low shady wood
—June. v. v.
3. E. caulinis subrat
glabris, spica cy
p. 4.
In swamps : New
Resembles E. lin
4. E. caulinis simpli
pentagonis apice
Willd. sp. pl. 5.

736. HYMENOPHYLLUM. *Sorus marginalis*, receptaculo cylindraceo insertus. *Indusium* bivalve, sorum includens.

V. HYDROPTERIDES.

737. ISOETES. *Capsula membranacea*, non dehiscens, frondis basi immersa, 1-locularis. *Semina angulata*, receptaculis pluribus filiformibus inserta.
738. SALVINIA. *Indusia imbricata*, connata, capsulam unilocularem mentientia. *Semina receptaculo centrali inserta*.
739. AZOLLA. *Capsula unilocularis*, radicalis, globosa, polysperma.

I. GONOPTERIDES.

713. EQUISETUM. *Gen. pl. 1614.*

1. E. caulis sterilibus simpliciter ramosis, ramis scabriusculis tetragonis, fructificationibus simplicibus, vaginis cylindraceis inciso-dentatis, dentibus acutis. *Willd. enum. 1065.* *Willd. sp. pl. 5. p. 1.*
Icon. *Schkuhr filic. t. 167.*
In low and sandy fields: Pennsylvania and Virginia. $\frac{1}{2}$. April, May. v. v.
2. E. caulis sterilibus fructificantibusque duplicato-ramosis, ramis scabriusculis deflexis tetragonis, ramulis subtiguetris. *Willd. enum. 1065.* *Willd. sp. pl. 5. p. 3.*
Icon. *Schkuhr filic. t. 166.* *Fl. dan. 1182.*
In low shady woods: New York to Virginia. $\frac{1}{2}$. April — June. v. v.
3. E. caulis subramosis, ramis subquaternis tetragonis glabris, spica cylindracea terminali.—*Willd. sp. pl. 5. p. 4.*
In swamps: New York to Virginia. $\frac{1}{2}$. May. v. v.
Resembles *E. limosum*.
4. E. caulis simpliciter ramosis glabris sulcatis, ramis pentagonis apice spiciferis. *Willd. enum. 1065.* *Willd. sp. pl. 5. p. 5.*

CRYPTOGAMIA GONOPTERIDES. Equisetum.

Icon. Schkuhr filic. t. 168, 169. Fl. dan. 1183.

On the banks of rivers and in deep swamps: Virginia.

♀. May, June. v. v.

scirpoideæ.

5. E. caulis simplicibus adscendentibus glabris filiformibus apice spiciferis, vaginis 3-dentatis sphacelatis, dentibus aristatis apice caducis. Willd. sp. pl. 5. p. 7. Mich. fl. amer. 2. p. 281.

E. reptans. Swartz. in litt.

In shady woods, round the mossy feet of trees: Canada and on the Green mountains, Vermont. ♀. July. v. v. A small species.

hyemale.

6. E. caulis simplicibus erectis scaberrimis apice spiciferis, vaginis discoloribus basi apiceque sphacelatis, dentibus aristatis omnino caducis. Willd. sp. pl. 5. p. 8.

Icon. Schkuhr filic. t. 172.

On the banks of rivers and in swamps: Canada to Virginia. ♀. July, Aug. v. v.

II. STACHYOPTERIDES.

714. LYCOPODIUM. Gen. pl. 1615.

* *Spicis pedunculatis.*

- carolinianum.* 1. L. caule repente, foliis subdistichis patulis lanceolatis superficialibus adpressis, pedunculo erecto solitario elongato monostachyo, bracteis sublanceolatis.—Willd. sp. pl. 5. p. 14.

Icon. Dill. musc. t. 62. f. 5.

In low meadows and woods: Pennsylvania to Carolina. ♀. July. v. v.

clavatum.

2. L. caule repente, ramis adscendentibus, foliis sparsis enervibus incurvato-setigeris, spicis geminatis cylindricis pedunculatis, squamis ovatis acuminatis erosodentatibus. Willd. sp. pl. 5. p. 16.

Icon. Schkuhr filic. t. 162. Fl. dan. 126. Dill. musc. t. 58. f. 1. Pluk. alm. t. 47. f. 8.

In dry pine woods: Canada to Pennsylvania. ♀. July. v. v. It sometimes has only one spike.

complanatum.

3. L. caule erecto, ramis alterni dichotomis, foliis bifariis connatis apice patentibus, superficialibus solitariis adpressis, pedunculis quadrifidis tetrastachyis, spicis teretibus cylindricis. Willd. sp. pl. 5. p. 19.

CRYPTOGAMIA

Icon. Fl. dan.

In shady wood

July. v. v.

4. L. caulis ei pressis, foliis pedunculis bus, squamis

On high moun

♀. July. v

the precedin

5. L. caule erecto lati acutis q solitariis mo cordatis acun

L. alpinum. M

Icon. Dill. musc.

In Canada. M

** *Spicis sessiliis.*

6. L. caule erecto, tentibus, foliis, spicis soli pl. 5. p. 21.

L. obscurum. L

Icon. Schkuhr fl

In dry shady v

v. v. This ar

name of Gro

menting room

properly tied i

7. L. caule repente ti, ramulis s lanceolatis mu increpenta an terminalibus.

Icon. Schkuhr fl

f. 9.

In shady woods:

8. L. caule subram erectis apice matis integerrimis sp. pl. 5. p. 25.

Icon. Schkuhr fl

t. 61. f. 7.

Icon. Fl. dan. 78. Dill. musc. t. 59. f. 8.

In shady woods, on rocky soil: Canada to Virginia. ♀.
July. v. v.

4. L. caulis erectis, ramis alternis dichotomis subcom-
pressis, foliis lanceolatis acutis quadrifariis adpresso-
pedunculis solitariis elongatis tristachyis, spicis tereti-
bus, squamis subrotundis acuminatis.

On high mountains in Virginia, near the Sweet-springs.

♀. July. v. v. This species is intermediate between
the preceding and following one.

5. L. caule erecto, ramis alternis dichotomis, foliis lanceo-
latiis acutis quadrifariis adpresso convexis, pedunculis
solitariis monostachyis, spicis teretibus, squamis sub-
cordatis acuminatis.—Willd. sp. pl. 5. p. 20.

L. alpinum. Mich. fl. amer. 2. p. 282.

Icon. Dill. musc. t. 58. f. 2?

In Canada. Michaux. ♀. July. +.

** *Spicis sessilibus, foliis caulem circumobsidentibus.*

6. L. caule erecto, ramis alternis confertis dichotomis pa-
tentibus, foliis sparsis sexfarulis linear-lanceolatis patu-
lis, spicis solitariis terminalibus sessilibus. Willd. sp.
pl. 5. p. 21.

L. obscurum. Sp. pl. 1566.

Icon. Schkuhr filic. t. 164. Dill. musc. t. 64. f. 12.

In dry shady woods: Canada to Carolina. ♀. July.
v. v. This and some other species are known by the
name of *Ground Pine*, and made use of for orna-
menting rooms and chimney-pieces, it forming, when
properly tied up, beautiful festoons.

7. L. caule repente, ramis adscendentibus basi bis biparti-
tis, ramulis simplicibus, foliis quinquefarulis linear-
lanceolatis mucronatis apice serrulatis patentibus, ad
incrementa annua contractis, spicis solitariis sessilibus
terminalibus. Willd. sp. pl. 5. p. 23.

Icon. Schkuhr filic. 162. Fl. dan. 127. Dill. musc. t. 63.
f. 9.

In shady woods: Canada to Pennsylvania. ♀. July. v. v.

8. L. caule subramoso repente, ramis simplicibus solitariis
erectis apice monostachyis, foliis linearibus sparsis acu-
tis integerim supra curvis, spica sessili foliosa. Willd.
sp. pl. 5. p. 25.

Icon. Schkuhr filic. 160. Fl. dan. 336. Dill. musc.
t. 61. f. 7.

CRYPTOGAMIA STACHYOPTERIDES. *Lycopodium.*

In cedar swamps and overflowed woods: Canada to New York. 4. July. v. v.

- alopecuroides.* 9. L. caule repente subramoso, ramis subsimplicibus elongatis adscendentibus apice monostachyis, foliis linearisubulatis basi ciliato-dentatis patulis, spica sessili foliosa. *Willd. sp. pl. 5. p. 26.*

Icon. *Schkuhr filic. t. 160. Dill. musc. t. 62. f. 6.*

In sphagnous swamps: New York to Carolina. 4. July. v. v. It is known by the name of *Walking Fern.*

- selaginoides.* 10. L. caule repente, ramis adscendentibus simplicibus, foliis sparsis lanceolatis patulis ciliato-denticulatis, spicis terminalibus solitariis foliosis sessilibus. *Willd. sp. pl. 5. p. 28.*

Icon. *Schkuhr filic. t. 165. Fl. dan. 70. Dill. musc. t. 68. f. 1.*

In mossy woods: Canada and New Hampshire. 4. July. v. v. The American plant is smaller than the European.

- rupestre.* 11. L. caule repente ramoso, ramis subdivisis adscendentibus, foliis sparsis imbricatis linearis-lanceolatis ciliatis apice piliferis, spicis solitariis sessilibus terminalibus. *Willd. sp. pl. 5. p. 30.*

Icon. *Schkuhr filic. t. 165. Dill. musc. t. 63. f. 11.*

On rocks: Canada to Virginia, and on the Carolina mountains. 4. July. v. v. A small species.

*** *Spicis sessilibus; foliis distichis.*

- albidulum.* 12. L. foliis bifariis ovatis acutis denticulatis, superficialibus alternis distichis adpressis, spicis terminalibus elongatis sessilibus tetragonis. *Willd. sp. pl. 5. p. 37.*

In moist shady places: Pennsylvania. 4. Aug. v. v. A very small, moss-like species, very much resembling the following.

- apodum.* 13. L. foliis bifariis subrotundo-ovatis acutis planis denticulatis cum superficialibus acuminatis alternantibus, caule ramoso basi radicante, spicis terminalibus sessilibus subsolitariis. *Willd. sp. pl. 5. p. 38.*

Icon. *Dill. musc. t. 64. f. 3.*
In shady woods: Pennsylvania to Florida. 4. July. v. v.

**** *Capsulis axillaris.*

- lucidulum.* 14. L. foliis octofariis linearis-lanceolatis denticulatis acutis patentia-reflexis, caule adscendente bifido. *Willd. sp. pl. 5. p. 51.*

CRYPTOGAI

L. reflexum.
Icon. *Dill. 1.*
In low sph
rivulets :
sembles *L*
of capsule

715. BERNHAF

1. B. caule nud
sp. pl. 3. p
Hoffmannia *a*
p. 17.

Psilotum triqu
Psilotum flori
Lycopodium
Icon. *Schkuh*
On the cliffs
4. June.

716. OPI

1. O. spica caulin
Willd. sp. 1
Icon. *Schkuh*
In low grassy
hills : New

2. O. spica caulin
bulbosa. *W*
O. crotalophor
In low sandy
May. v. v.

717. BOTR

1. B. scapo nudo,
bipinnatis, f
Willd. sp. p.
B. lunarioides.
Botrypus lunari
Osmunda biter
Icon. *Schkuhr*
In pastures and
June. v. v.

2. B. scapo inferne
oblongo-lanc
cordatis, spic

View
lon-
eari-
i fo-

U.
king

fo-
dicis
sp.

usc.

U.
the

nti-
atis
ous.

ina

ous
ga-

v.
m-

u-
is-

v.
p.

tia
p.

L. reflexum. Schkuhr filic. t. 159.
Icon. Dill. musc. t. 56. f. 2.

In low sphagnum grounds and on the banks of shady rivulets: Canada to Carolina. U. July. v. v. Resembles *L. Selago*; it frequently bears bulbs instead of capsules.

715. BERNHARDIA. Willd. act. acad. erford. 1802. p. 11.

1. B. caule nudo dichotomo ramisque triquetris.—Willd. sp. pl. 3. p. 56. *dichotoma*.

Hoffmannia aphylla. Wild. in Roem. et Usteri bot. mag. 6. p. 17.

Psilotum triquetrum. Swartz. synop. filic. 187.

Psilotum floridanum. Mich. fl. amer. 2. p. 281.

Lycopodium nudum. Sp. pl. 1564.

Icon. Schkuhr filic. t. 165. b. Dill. musc. t. 64. f. 4.

On the cliffs of rocks, near the sea-coast, in Florida.
U. June. v. s.

716. OPHIOGLOSSUM. Gen. pl. 1621.

1. O. spica caulina, fronde ovata obtusa, arcta reticulata. Willd. sp. pl. 5. p. 58. *vulgatum*.

Icon. Schkuhr filic. t. 153. Fl. dan. 147.

In low grassy open places, in woods and on the sides of hills: New York to Pennsylvania. U. May. v. v.

2. O. spica caulina, fronde subcordato-ovata obtusa, radice bulbosa. Willd. sp. pl. 5. p. 60. *bulbosum*.

O. crotalophoroides. Walt. fl. car. 256.

In low sandy grounds: New Jersey to Carolina. U. May. v. v.

717. BOTRYCHIUM. Swartz. synop. filic. 8.

1. B. scapo nudo, frondibus glabris radicalibus tripartito-bipinnatis, foliolis lunatis crenatis, spicis pinnatis.—Willd. sp. pl. 5. p. 63. *fumarioides*.

B. lunarioides. Swartz. synop. filic. 172.

Botrypus lunarioides. Mich. fl. amer. 2. p. 274.

Osmunda bibernata. Lam. encycl. 4. p. 608.

Icon. Schkuhr filic. t. 157.

In pastures and open woods: New York to Carolina. U. June. v. v.

2. B. scapo inferne unifrondoso, fronde subbibernata, foliolis oblongo-lanceolatis serrulatis basi dilatatis inaequaliter cordatis, spicis bipinnatis.—Willd. sp. pl. 5. p. 63. *obliquum*.

CRYPTOGAMIA STACHYOPTERIDES. Botrychium.

In open woods: Pennsylvania and Virginia. ♀. June, July. v. v. Resembles the preceding very much, and is probably only a variety.

- dissectum.* 3. B. scapo inferne unifrondoso, fronde tripartito-bipinnatifida, lacinis linearibus bipartitis apice bidentatis.—*Willd. sp. pl. 5. p. 64.*

Icon. *Schkuhr filic. t. 158. Pluk. amalh. t. 427. f. 5.*
In pastures and open dry woods: New York to Florida. ♀. June. v. v.

- virginicum.* 4. B. hirsutum; scapo medio frondoso, fronde subterna tripartito-bipinnatifida, foliolis inciso-pinnatifidis, lacinis obtusis subtridentatis, spicis bipinnatis divaricatis.—*Willd. sp. pl. 5. p. 64.*

Botrypus virginicus. Mich. fl. amer. 2. p. 274.

Osmunda virginica. Sp. pl. 1579.

Osmunda multifida. Gmel. nov. com. petrop. 12. t. 11. f. 1.

Icon. *Schkuhr filic. t. 156.*

In shady woods, on rich vegetable soil: Canada to Carolina. ♀. June, July. v. v. This is the largest of the species here described; it is known by the name of *Rattle Snake Fern*, probably from the circumstance of growing near the places where those venomous animals generally are found.

- gracile.* 5. B. glabrum; scapo medio frondoso, fronde tripartito-bipinnatifida, lacinis inciso-subpinnatifidis acutis subdentatis, spicis gracilibus pinnatis erectis.

In shady fertile woods: Virginia. ♀. June. v. v. This species approaches near to the preceding in its first appearance, but is by far a smaller and more slender plant, besides the other distinctions.

III. SCHISMATOPTERIDES.

718. HYDROGLOSSUM. *Willd. sp. pl. 5. p. 77.*

Ophioglossum. Linn.

- palmatum.* 1. H. caule flexuoso-scandente, frondibus conjugatis cordatis quinquelobo-palmatis, lobis lanceolatis integerimis obtusis obsolete sinuatis, spicillis terminalibus composite paniculatis oblongo-linearibus.—*Willd. sp. pl. 5. p. 94.*

CRYPTOGAMIA

Lygodium palmatum. Cteisium paniculatum. Icon. Schkuhr t. 1. f. 2.

In low copses, v. v. This the North & plant when

719. SCHIZOPHYLLUM

1. S. fronde simpliciteratis inflata

In barren sand-limestone coun Aug. Sept., scarce, and s of the botan

720. C

1. O. frondibus simpliciteratis ovato-cordatis, fructificans pl. 5. p. 98.

Icon. Schkuhr f. 3.

In low open woodland near to Florida fern of the sandstone the fertile ste down.

2. O. frondibus simpliciteratis apiculatis pl. 5. p. 96.

In Virginia. (probably only a specimen in my herbarium) the fructiferous grow out into formation of

3. O. frondibus simpliciteratis oblongo-quot interme p. 96.

Lygodium palmatum. *Swartz. synop. filic.* 154.
Cteisium paniculatum. *Mich. fl. amer. 2. p. 275.*
Icon. Schkuhr filic. 140. *Willd. act. acad. erford.* 1802.
t. 1. f. 2.
In low copses, climbing among small shrubs. $\frac{3}{4}$. July.
v. v. This is the most beautiful and singular of all
the North American ferns, and is a highly ornamental
plant when cultivated in pots.

719. SCHIZÆA. *Smith act. taur.* 5. p. 419.

1. *S. fronde simplicissima linear-compressa, spicis conglobatis inflexis secundis.* *pusilla.*
In barren sandy but moist grounds : New Jersey, Burlington county, near a place called Quaker Bridge. $\frac{3}{4}$. Aug. Sept. *v. v.* This singular little fern is very scarce, and so small that it generally escapes the notice of the botanist, unless accident points it out.

720. OSMUNDA. *Gen. pl.* 1622.

1. *O. frondibus sterilibus pinnatis, pinnis pinnatifidis, laciniis ovato-oblongis obtusis integerrimis, stipite lanato, fructificantibus bipinnatis lanuginosis.* — *Willd. sp. pl. 5. p. 98.*
Icon. Schkuhr filic. t. 146. Moris. hist. 3. s. 14. t. 4. f. 3.
In low open woods and exsiccated swamps : New England to Florida. $\frac{3}{4}$. June—Aug. *v. v.* This is a fern of the size and habit of *Polypodium Filix mas*; the fertile stems are covered with a cinnamon-coloured down.
2. *O. frondibus pinnatis ferrugineo-tomentosis, pinnis pinnatifidis apice coarctato-fructificantibus.* — *Willd. sp. pl. 5. p. 96.*
In Virginia. *Clayton.* $\frac{3}{4}$. April. +. This is probably only a variety of the preceding. I have no specimen in my herbarium; but recollect very well that the fructiferous stems of that species very frequently grow out into leaves, which may have given rise to the formation of the present species.
3. *O. frondibus pinnatis glabris, pinnis oppositis pinnatifidis, laciniis oblongis acutiusculis integerrimis, pinnis aliquot intermediis fructificantibus.* — *Willd. sp. pl. 5. p. 96.* *interrupta.*

- O. basilaris. *Sprengel. aul.* 3. p. 160.
Icon. *Schkuhr filic.* t. 144.
In low and swampy grounds: Canada to Virginia, and in
the western territories. ♀. June, July. v. v. This
singular fern is about two feet high.
- spectabilis.* 4. O. frondibus bipinnatis, pinnulis oblongis obtusiusculis
argute serrulatis basi truncatis inaequalibus, omnibus
alternis, panicula bipinnata fructificante in apice frondis.—*Willd. sp. pl.* 5. p. 98.
O. regalis β. *Sp. pl.* 1521. *Mich. fl. amer.* 2. p. 273.
Icon. *Pluk. alm.* t. 184. f. 4.
In mountain swamps: Canada to Carolina. ♀. July.
v. v. A very elegant species.

CRYPT

On the true
Willdenow
Virginia, a
wise never
under the

3. P. frondibus |
ribus integ |
ribus, subt |
soris solitar |
P. ceteraccin |
Acrostichum |
Icon. *Schkuh* |
On rocks and |
Kentucky a |
ral bridge, |
Dismal-swa

4. P. frondibus |
infimis deflu |
narum infel |
mis, infimi |
tariis. *Wi* |
Icon. *Pluk.* |
t. 20. b. |
In shady mois |
rolina, ♀. |
triangle in it

5. P. frondibus b |
tiguis adnat |
soris minutis |
In shady wood |
Resembles t

6. P. frondibus t |
obtusiusculis |
fluentibus. |
P. Dryopteris. |
Nephrodium |
In shady beec |
July. v. v. |
I only foun |
shady beech

IV. FILICES.

721. ACROSTICHUM. *Gen. pl.* 1625.

- aureum.* 1. A. frondibus pinnatis, pinnis alternis oblongo-lanceolatis integerimis, basi cuneatis aequalibus, omnibus acuminatis, superioribus fructificantibus. *Willd. sp. pl.* 5. p. 116.
Icon. *Schkuhr filic.* t. 1. *Petiv. fil.* t. 8. f. 5. *Plum. amer.* t. 7. *Pluk. alm.* t. 288. f. 2.
In deep swamps near the sea-coast: Florida, ♀. July.
v. v. This is the largest of the North American ferns,
the leaves being sometimes above five feet long.

722. POLYPODIUM. *Swartz. synop. filic.* 3.

- vulgare.* 1. P. frondibus profunde pinnatifidis, lacinias linear-lanceolatis obtusis crenulatis approximatis, superioribus sensim minoribus, soris solitariis, caudice paleaceo. *Willd. sp. pl.* 5. p. 172.
Icon. *Schkuhr filic.* t. 11. *Plum. fil.* t. A. f. 2.
On rocks and old rotten trees: Canada to Carolina. ♀.
July. v. v.
- virginianum.* 2. P. frondibus profunde pinnatifidis, lacinias lanceolatis obtusis integerimis approximatis, superioribus sensim minoribus, soris solitariis, caudice nudo. *Willd. sp. pl.* 5. p. 174.
Icon. *Petiv. filic.* t. 7. f. 9. *Plum. amer.* t. 36. filic. t. 77.

On the trunks of trees: Virginia? \mathcal{U} . July. +. Willdenow doubts whether this is really a native of Virginia, and I believe with much propriety, as I likewise never have seen any other species than *P. vulgare* under the name of *P. virginicum*.

3. *P. frondibus profunde pinnatifidis, lacinis alternis linearibus integerrimis obtusis, superioribus sensim minoribus, subtus stipiteque squamulosis, apice fructiferis, soris solitariis.* *Willd. sp. pl. 5. p. 174.*
P. ceteraccinum. Mich. fl. amer. 2. p. 271.
Acrostichum polypodioides. Sp. pl. 1525.
Icon. Schkuhr filic. t. 11. b.
 On rocks and trunks of old trees: Virginia to Florida, Kentucky and Tennessee; on the rocks of the Natural bridge, Virginia, and on the stems of trees in the Dismal-swamp. *Ph.* July. *v. v.*
4. *P. frondibus bipinnatifidis glabriusculis, pinnis duabus infimis deflexis, lacinis lanceolatis obtusis ciliatis, pinnarum inferiorum inciso-crenatis superiorum integerimis, infimis adnato-decurrentibus, soris minutis solitariis.* *Willd. sp. pl. 5. p. 200.*
Icon. Pluk. alm. t. 284. f. 2. bona. Schkuhr filic. t. 20. b.
 In shady moist woods on fertile soil: New York to Carolina. \mathcal{U} . July. *v. v.* The whole leaf forms a triangle in its circumference; the stem is smooth.
5. *P. frondibus bipinnatifidis ciliatis, pinnis oppositis contiguis adnatis, pinulis subellipticis, stipite paleaceo, soris minutis.* *Willd. sp. pl. 5. p. 200.*
 In shady woods: Canada to New York. \mathcal{U} . July. *v. v.* Resembles the preceding.
6. *P. frondibus ternatis bipinnatis rectis rigidulis, lacinis obtusiusculis subintegerrimis, soris marginalibus confluentibus.* *Willd. sp. pl. 5. p. 210.*
P. Dryopteris. Bolton filic. 53. t. 1.
Nephrodium Dryopteris. Mich. fl. amer. 2. p. 270.
 In shady beech-woods: Canada to Pennsylvania. \mathcal{U} . July. *v. v.* This is a very rare fern in Pennsylvania. I only found it on the western mountains in deep shady beech forests.

*incanum.**connectile.**calcareum.*

723. WOODSIA. *Brown in linn. trans.* 11. p. 173.*hyperborea.*

1. *W. frondibus pinnatis, pinnis tripartitis vel pinnatifidocrenatis cuneiformibus obtusis dentatis subtus hirtis, soris solitariis demum confluentibus.* — *Brown l. c. cum icon.*

*Polypodium hyperboreum. Willd. sp. pl. 5. p. 197.**Polypodium arvonicum. Smith fl. brit. 3. p. 1115.**Acrostichum ilvense. Huds. angl. 451.**Acrostichum alpinum. Bolton filic. 76. t. 42.**Acrostichum hyperboreum. Liljeblad act. holm. 1793. p. 201. t. 8.**Icon. Schkuhr filic. t. 17. b. Pluk. phytogr. t. 89. f. 5. Moris. hist. 3. s. 14. t. 3. f. 23.*

In the clefts of rocks: Canada and on the high mountains of Pennsylvania and Virginia. $\frac{4}{4}$. July. v. v. A small fern growing in large tufts, and covered with long brown chaffy hair.

ilvensis.

2. *W. frondibus bipinnatifidis, laciniis oblongis obtusis inferioribus repandis, superioribus integerrimis subtus hirtis, soris submarginalibus demum confluentibus, stipite superne villoso.* — *Brown l. c.*

*Polypodium ilvense. Willd. sp. pl. 5. p. 198.**Polystichum Marantæ. Roth fl. germ. 3. p. 92.**Acrostichum ilvense. Sp. pl. 1528.**Icon. Schkuhr filic. t. 19. Fl. dan. 391.*

On rocks: Canada to Virginia. $\frac{4}{4}$. July. v. v. Resembles the preceding very much; but the leaves are smaller divided and covered with fine woolly hair, not chaffy as in the other.

724. ASPIDIUM. *Swartz. synop. filic.* 3.*Nephrodium. Mich. fl. amer. 2. p. 266.*** Frondibus ternatis.**cicutarium.*

1. *A. frondibus ternatis, foliolis pinnatifidis, laciniis acuminatis sinuato-pinnatifidis integerrimis, lateralium infima et exteriore lacinia elongata, soris serialibus.* Willd. sp. pl. 5. p. 215.

*Polypodium cicutarium. Sp. pl. 1549.**Polypodium appendiculatum. Swartz. fl. ind. occid. 3. p. 1677.**Icon. Pluk. alm. t. 289. f. 4. & t. 296. f. 2.*

On rocks in the mountains of Virginia. $\frac{4}{4}$. +.

2. *A. frondibus ciliato-serratis, inferiore ciliatis, ribus, stipi confluentibus.*
A. auriculatum.
Nephrodium
On rocks, in
lina. $\frac{4}{4}$.
latum.

**** Frondibus*

3. *A. frondibus fidis, laciniis ciliatis, soris pl. 5. p. 248.*
A. thelypterioi

*Nephrodium**Polypodium**Icon. Schkuhr**On river sides.**nada to Carc*

4. *A. frondibus dis glabris, marginalibus sp. pl. 5. p. 2*

*Polypodium**Polystichum**Acrostichum**T**Icon. Schkuhr**1018.**In low grassy fi* *$\frac{4}{4}$. July. v**fication; th**ceding.*

5. *A. frondibus subcordato-otusis dentato-*

*pl. 5. p. 252.**Polypodium**cris**Icon. Schkuhr*

** *Frondibus pinnatis.*

2. A. frondibus pinnatis, pinnis lanceolatis falcatis acutis *acrostichoides*.
ciliato-serratis, basi superiore mucronato-auriculatis,
inferiore cuneatis, superioribus fructificantibus mino-
ribus, stipite rachique paleaceis, soris solitariis demum
confluentibus. *Willd. sp. pl. 5. p. 225.*

A. auriculatum. *Schkuhr filic. 31. t. 30. excl. synon.*
Nephrodium acrostichoides. Mich. fl. amer. 2. p. 267.
On rocks, in shady low places : New England to Caro-
lina. ♀. June—Aug. v. v. Resembles *N. auricu-*
latum.

*** *Frondibus subbitpinnatis; indusiis dimidiatis renifor-*
mibus.

3. A. frondibus pinnatis, pinnis linear-lanceolatis pinnati-
fidis, lacinis oblongis obtusiusculis subintegerrimis
ciliatis, soris marginalibus, stipite laevi.—*Willd. sp.*
pl. 5. p. 248.

A. thelypteroides. *Swartz. synop. filic. 57.*
Nephrodium thelypteroides. Mich. fl. amer. 2. p. 267.
Polypodium noveboracense. Sp. pl. 1552.
Icon. *Schkuhr filic. t. 46.*

On river sides and the banks of ditches and ponds : Ca-
nada to Carolina. ♀. July. v. v.

4. A. frondibus pinnatis, pinnis linear-lanceolatis pinnatifi-
dis glabris, lacinis ovatis acutis integerrimis, soris
marginalibus contiguis demum confluentibus. *Willd.*
sp. pl. 5. p. 249.

Polypodium Thelypteris. Mant. 505.
Polystichum Thelypteris. Roth fl. germ. 3. p. 77.
Acrostichum Thelypteris. Sp. pl. 1528.

Icon. *Schkuhr filic. t. 52. Fl. dan. 760. Engl. bot.*
1018.

In low grassy fields and swamps : Canada to Virginia.
♀. July. v. v. It is very seldom found with fructi-
fications ; the leaves are smaller than in the pre-
ceding.

5. A. frondibus lanceolato-ovatis glabris pinnatis, pinnis
subcordato-oblongis pinnatifidis, lacinis oblongis ob-
tusis dentato-serratis, stipite paleaceo.—*Willd. sp.*
pl. 5. p. 252.

Polypodium cristatum. Sp. pl. 1551.
Icon. *Schkuhr filic. t. 37. Afzel. act. holm. 1787. t. 9.*

- Callipteris. β . A. majuscum; pinnis alternis, lacinis oblongis inaequaliter adpresso-serratis, fronde apice fructificante.
Polypodium Callipteris. Ehrh. beitr. 3. p. 77.
- lancastriense. γ . A. pinnis suboppositis, lacinis triangulari-ovatis acute serratis, stipite nudo.—Willd. sp. pl. 5. p. 261.
Schkuhr filic. t. 41.
In moist rocky situations: New York to Virginia. 4. July. v. v. Of middle size.
- ***** *Frondibus bipinnatis triplicato-pinnatis; indusii dimidiatis reniformibus.*
- obtusum.* 6. A. frondibus bipinnatis, pinnulis oblongis obtusis superioribus coadunatis dentato-crenatis, rachi superne pubescente. Willd. sp. pl. 5. p. 254.
Polypodium obtusum. Swartz. synop. filic. 39. Schkuhr filic. 18.
Icon. *Schkuhr filic.* t. 21. ♂ 43. b.
In clefts of rocks: Pennsylvania and Virginia. 4. July.
v. v. A small species, not above four or five inches high.
- aculeatum.* 7. A. frondibus bipinnatis, pinnulis ovatis subfalcatis acutis mucronato-serratis, basi superiore truncatis, inferiore abbreviato-cuneatis, superioribus fructificantibus, stipite rachique paleaceis. Willd. sp. pl. 5. p. 258.
Polypodium aculeatum. Sp. pl. 1552.
Polystichum aculeatum. Roth fl. germ. 3. p. 79.
Icon. *Schkuhr filic.* t. 39. ♂ 40. Bolt. filic. t. 56. Pluk. alm. t. 180. f. 1.
In shady rocky places, on the Green mountains, Vermont. 4. Aug. v. v. The size of No. 5, of a shining and strong texture.
- marginale.* 8. A. frondibus bipinnatis, pinnulis oblongis obtusis decurrentibus crenatis, crenis baseos profundioribus, soris marginalibus, stipite paleaceo. Willd. sp. pl. 5. p. 259.
Polypodium marginale. Sp. pl. 1522.
Nephrodium marginale. Mich. fl. amer. 2. p. 267.
Icon. *Schkuhr filic.* t. 45. b.
In rocky shady places: Canada to Carolina. 4. July.
v. v. Resembles the following.
- Filix mas.* 9. A. frondibus bipinnatis, pinnulis oblongis obtusis serratis, serraturis muticis, soris costæ mediae approximatis, stipite rachique paleaceis. Willd. sp. pl. 5. p. 259.
- CRYP
- Polypodium F
Polystichum F
Icon. *Schkuhr*
s. 14. t. 3. f.
 β . A. pinnulis et
t. 45.
In shady pine
Carolina. 7
high; the le
quently doub
10. A. frondibus b
incisis, lacini
leaceo. Wil
In Pennsylvania.
following in s
11. A. frondibus bip
confluentibus
tis, stipite pal
pl. 5. p. 262.
Polypodium spin
Polypodium crist
Polystichum spin
Icon. *Schkuhr* fi
icon. t. 2. f. 2.
In shady woods,
4. July. v.
12. A. frondibus bipi
natifido-incisis,
leaceo, indusis
Polypodium dilata
Polypodium arista
Polypodium crista
Polystichum multi
Nephrodium crista
Icon. *Schkuhr fil*
ic. t. 2. f. 4. I
In shady rocky w
July. v. v. Re
broader, and ma
- ***** *Frondibus bip*
13. A. frondibus bipin
positis oblongis c

- Polypodium Filix mas. *Sp. pl.* 1551.
 Polystichum Filix mas. *Roth. Fl. germ.* 3. p. 82.
 Icon. *Schkuhr filic.* t. 44. *Bolt. filic.* t. 24. *Moris. hist.* 3. s. 14. t. 3. f. 6.
- β . A. pinnulis erosio-serratis remotioribus.—*Schkuhr filic.* t. 45. *erosum.*
- In shady pine forests: New Jersey to Virginia. β . in Carolina. \mathcal{U} . July. v. v. From two to three feet high; the leaflets are more obtuse, and more frequently doubly serrated, than in the European plant.
10. A. frondibus bipinnatis, pinnulis linearibus pinnatifido-incisis, lacinia apice submucronato-serratis, stipite paleaceo. *Willd. sp. pl.* 5. p. 262. *intermedium.*
- In Pennsylvania. *Muhlenberg.* \mathcal{U} . +. Resembles the following in several respects very much.
11. A. frondibus bipinnatis, pinnulis oblongis decurrentibus confluentibus inciso-dentatis, lacinia mucronato-serratis, stipite paleaceo, indusii glandulosis. *Willd. sp. pl.* 5. p. 262. *spinulosum.*
- Polypodium spinulosum. *Retz. prodr. fl. scand.* n. 1295.
 Polypodium cristatum. *Hoffm. fl. germ.* 2. p. 8.
 Polystichum spinosum. *Roth. fl. germ.* 3. p. 91.
 Icon. *Schkuhr filic.* t. 48. *Fl. dan.* 707. *Fr. Mill. icon.* t. 2. f. 2.
- In shady woods, in fertile soil: Pennsylvania to Virginia. \mathcal{U} . July. v. v. A large species.
12. A. frondibus bipinnatis, pinnulis oblongis distinctis pinnatifido-incisis, lacinia mucronato-serratis, stipite paleaceo, indusii laevibus. *Willd. sp. pl.* 5. p. 203. *dilatatum.*
- Polypodium dilatatum. *Hoffm. fl. germ.* 2. p. 7.
 Polypodium aristatum. *Pillars fl. delph.* 3. p. 844.
 Polypodium cristatum. *Huds. angl.* 457.
 Polystichum multiflorum. *Roth. fl. germ.* 3. p. 87.
 Nephrodium cristatum. *Mich. fl. amer.* 2. p. 209.
 Icon. *Schkuhr filic.* t. 47. *Bolt. filic.* t. 23. *Fr. Mill. ic.* t. 2. f. 4. *Pluk. alm.* t. 181. f. 2.
- In shady rocky woods: Pennsylvania and Virginia. \mathcal{U} . July. v. v. Resembles *A. Filix femina*; it is larger, broader, and more compound in its leaves.
- ***** *Frondibus bipinnatis triplicato-pinnatis; indusiis lateralibus.*
13. A. frondibus bipinnatis oblongo-lanceolatis, pinnulis oppositis oblongis obtusis serratis, inferioribus pinnatifido-

dis, rachi bulbifera, soris subrotundis. *Willd. sp. pl. 5.* p. 275.

Polypodium bulbiferum. Sp. pl. 1553.

Nephrodium bulbiferum. Mich. fl. amer. 2. p. 268.

Icon. Schkuhr filic. t. 57. Corn canad. 5. Moris. hist. 3. s. 14. t. 3. f. 10.

In shady woods, about rocks : Canada to Pennsylvania.

4. July. v. v. The leaf is narrow, and produces generally some small bulbs on its midrib.

asplenoides.

14. A. frondibus bipinnatis, pinnulis lineari-lanceolatis inciso-serratis, serraturis bi- s. tridentatis terminalibus acutioribus, soris oblongis lunatis. *Willd. sp. pl. 5.* p. 276.

Nephrodium asplenoides. Mich. fl. amer. 2. p. 268.

Polypodium pensylvanicum. Muhl. in litt.

Asplenium Athyrium. Sprengel. anleit. 3. p. 113.

Icon. Schkuhr filic. t. 78.

In shady woods, on the banks of rivulets and ditches : New England to Carolina. 4. July. v. v. A tall species, very much resembling the following in many respects.

Filix femina.

15. A. frondibus bipinnatis, pinnulis oblongo-lanceolatis inciso-serratis, serraturis bi- s. tridentatis acutiusculis, soris oblongis rectis. *Willd. sp. pl. 5. p. 276.*

A. alpestre. Schkuhr filic. p. 58.

Polyodium Filix femina. Sp. pl. 1551.

Polyodium molle. Schreb. spicil. 70.

Polyodium dentatum, incisum et trifidum. Hoffm. fl. germ. 2. p. 7.

Athyrium molle, trifidum, ovatum et Filix femina.

Roth. fl. germ. 3. p. 61—65.

Icon. Schkuhr filic. t. 58. 59. ♂ 60. Pluk. phytoogr. t. 180. f. 4. Moris. hist. 3. s. 14. t. 3. f. 8.

In low shady grounds : Canada to Virginia. 4. July. v. v.

angustum.

16. A. frondibus bipinnatis, pinnulis lanceolatis inciso-serratis, serraturis subbidentatis, infima superiore elongata, / soris oblongis sublunatis. *Willd. sp. pl. 5. p. 277.*

Nephrodium Filix femina. Mich. fl. amer. 2. p. 208.

In shady low woods : Canada to Pennsylvania. 4. July. v. v. Resembles the preceding.

punctilobum.

17. A. frondibus bipinnatis, pinnulis decurrentibus ovato-oblongis pinnatifidis, lacinias subquadridentatis, soris

CRYPTO

solitario inst
bro. *Willde*

Nephrodium p
In shady wood
da to Virgin
feet high.

18. A. frondibus
oblongis pin
Willd. sp. pl.
In Pennsylvania
above a span

19. A. frondibus
dentatis, soris
sp. pl. 5. p. 1

Nephrodium te
In clefts of roc
A small speci

20. A. frondibus bi
oblongis obtu
fluentibus. *W*

Nephrodium ru
On rocks : Ca

A small speci
Woodsia hyp

725. ONOCLE

1. O. frondibus ste
inciso-dentatibus
bus bipinnatis
chi glabra.—I

Icon. Schkuhr fi
Breyn. cent. t.

In low pastures
4. July. v.
green.

2. O. frondibus ster
fidis, lobis rotu
superioribus c

piunulis incisis
mosa.—*Schku*

In low grounds
so large as the |

- pl. 5.
ist. 3.
ania.
duces
inci-
acu-
pl. 5.
3.
ches :
A tall
many
is in-
culis,
m. fl.
mina.
/togr.
July.
serra-
gata,
7.
8.
July.
wato-
goro
- solitario instructis, rachibus pubescentibus, stipite gla-
bro. *Willd. sp. pl. 5. p. 279.*
- Nephrodium punctilobulum. *Mich. fl. amer. 2. p. 268.*
In shady woods, on the banks of small rivulets : Canada
to Virginia. ♀. July. v. v. From two to three
feet high.
18. A. frondibus bipinnatis, pinnulis decurrentibus ovato-
oblongis pinnatifidis, laciniis dentatis, soris sparsis. *atomarium.*
Willd. sp. pl. 5. p. 279.
In Pennsylvania. *Muhlenberg.* ♀. +. Not much
above a span high ; fructification small.
19. A. frondibus bipinnatis, pinnulis ovatis decurrentibus
dentatis, soris solitariis juxta dentes pinnularum. *Willd.*
sp. pl. 5. p. 279. *tenue.*
Nephrodium tenue. *Mich. fl. amer. 2. p. 269.*
In clefts of rocks : Canada to Virginia. ♀. July. v. v.
A small species ; fructifications large.
20. A. frondibus bipinnatis, pinnulis subtus paleaceo-setosis
oblongis obtusis crenatis coadunatis, soris demum con-
fluentibus. *Willd. sp. pl. 5. p. 282.*
Nephrodium rufidulum. *Mich. fl. amer. 2. p. 269.*
On rocks : Canada to Pennsylvania. ♀. July. v. v.
A small species, growing in large tufts ; it resembles
Woodsia hyperborea very much.
725. ONOCLEA. *Willd. in mag. ges. naturf. fr. z.
berl. 1809. p. 160.*
1. O. frondibus sterilibus pinnatis, pinnis lanceolatis acutis
inciso-dentatis, superioribus coadunatis, fructificantibus
bipinnatis, pinnulis recurvato-globosis glabris, rachi glabra.—*Willd. sp. pl. 5. p. 287.*
Icon. *Schkuhr filic. t. 102. Pluk. mant. t. 404. f. 2.*
Breyn. cent. t. 56. f. B.
In low pastures and open woods : New York to Florida.
♀. July. v. v. About a foot high ; leaves pale
green.
2. O. frondibus sterilibus pinnatis, pinnis oppositis pinnati-
fidis, lobis rotundatis : inferioribus sensim minoribus,
superioribus coadunatis, fructificantibus bipinnatis,
pinnulis incisis recurvato-globosis villosis, rachi squa-
mosa.—*Schkuhr filic. p. 95. t. 103.*
In low grounds : Pennsylvania. ♀. July. v. v. Not
so large as the preceding.

726. STRUTHIOPTERIS. *Willd. in mag. ges. nat. fr. z. berl. 1809. p. 160.*

- pensylvanica.* 1. S. frondibus sterilibus bipinnatifidis, lacinias integerimis obtusis, infima elongata acuta. *Willd. sp. pl. 5. p. 289.*
Onoclea nodulosa. Schkuhr filic. p. 96. t. 104.
 In Pennsylvania. *Muhlenberg.* ♀. +. Resembles *S. germanica.*

727. ASPLENIUM. *Swartz. synop. filic. p. 4.*

- rhzophyllum.* 1. A. frondibus lanceolatis stipitatis subcrenatis, basi auriculato-cordatis, apice longissimo linearifiliforme radicante. *Willd. sp. pl. 5. p. 305.*
 Icon. *Pluk. alm. t. 105. f. 3.* *Petiv. gazoph. t. 105. f. 7.*
Moris. hist. 3. s. 14. t. 5. f. 14.
 On wet shady rocks: Canada to Carolina. ♀. July.
v. v. This fern is highly ornamental to rock-works, and very singular in its striking root at the elongated points of the leaves.

- angustifoli- um.* 2. A. frondibus pinnatis, pinnis alternis, superioribus suboppositis, lineariflanceolatis subrepandis, basi superiore truncatis, inferiore leviter rotundatis. *Willd. sp. pl. 5. p. 313.*
A. pycnocarpon. Sprengel. anleit. 3. p. 112.
 Icon. *Schkuhr filic. t. 67. ♂ 69.* *Moris. hist. 3. s. 14. t. 2. f. 25.*

In shady woods, on the banks of rivers: western parts of New York and Pennsylvania. ♀. July. *v. v.* A very elegant fern about a foot high.

- eleneum.* 3. A. frondibus pinnatis, pinnis sessilibus lanceolatis serrulatis basi cordatis, sursum auriculatis. *Willd. sp. pl. 5. p. 329.*

- A. polypodioides. Swartz. synop. filic. 79. ♂ 272.*
A. trichomanoides. Mich. fl. amer. 2. p. 265.
Acrostichum platyneuron. Sp. pl. 1527. excl. synon.
 Icon. *Schkuhr filic. t. 73.* *Pluk. alm. t. 287. f. 2. et t. 89. f. 8.* *Moris. hist. 3. s. 14. t. 2. f. 12.*
 In clefts of rocks: New England to Carolina. ♀. July.
v. v. A small species.

- melanoau- lon.* 4. A. frondibus pinnatis, pinnis subrotundis obtusis crenatis basi cuneatis, stipe discoloro. *Willd. enum. 1072.*
Willd. sp. pl. 5. p. 332.

- A. trichomanes
 On rocks: Can
 small: stem
 5. A. frondibus p
 minatis profu
 obtuse-serrati
—Willd. sp.
 A. acrostichoide
 Icon. *Schkuhr J*
 In shady woods
 Carolina. ♀.
Thelypteris ve
 two feet high.
 6. A. frondibus ba
 pinnulis rhom
 culatis. *Will*
 Icon. *Schkuhr J*
 In clefts of rock
v. v. The An
 exceeding an i
 7. A. frondibus glat
 natifidis, lacini
p. 342.
 A. Adiantum nig
 In clefts of rocks
 rolinia. ♀. Ju
A. nigrum, to

728. SCLOOPENI

1. S. frondibus lato-l
Willd. sp. pl. 5
Asplenium Sclop
 Icon. *Schkuhr fili*
 In shady woods, a
 New York, ne
 J. Geddis, Esq.
 seen in no other
 ther have I ha
 found in any oth

- A. trichomanes. *Mich. fl. amer. 2. p. 264.*
On rocks: Canada to Carolina. ♀. July. v. v. Very
small: stem polished, black.
5. A. frondibus pinnatis, pinnis sessilibus lanceolatis acu- *thelypterioi-*
minatis profunde pinnatifidis, laciiniis oblongis obtusis *des.*
obtuse-serratis, soris parallelis demum confluentibus.
—*Willd. sp. pl. 5. p. 336.*
- A. acrostichoides. *Swartz. synop. filic. 82. ♂ 275.*
Icon. *Schkuhr filic. t. 76. b.*
In shady woods on the sides of rivulets: Pennsylvania to
Carolina. ♀. July. v. v. Resembles *Aspidium*
Thelypteris very much; it is from eighteen inches to
two feet high.
6. A. frondibus basi bipinnatis, apice simpliciter pinnatis, *Ruta mura-*
pinnulis rhomboe-oblongis obtusis, apice obtuse denti-
culatis. *Willd. sp. pl. 5. p. 341.*
Icon. *Schkuhr filic. t. 80. b. Fl. dan. 190.*
In clefts of rocks: New York to Carolina. ♀. July.
v. v. The American plant is very small, scarcely ever
exceeding an inch or two in height.
7. A. frondibus glabris bipinnatis, pinnulis lanceolatis pin- *montanum.*
natifidis, laciiniis tri- vel bidentatis.—*Willd. sp. pl. 5.*
p. 342.
A. Adiantum nigrum. *Mich. fl. amer. 2. p. 265.*
In clefts of rocks, on the mountains of Virginia and Ca-
rolina. ♀. July. v. v. A much smaller plant than
A. nigrum, to which it has some resemblance.
728. SCOLOPENDRIUM. *Smith act. taur. 5. p. 410.*
t. 9. f. 2.
1. S. frondibus lato-lanceolatis basi cordatis, stipite paleaceo. *officinarum.*
Willd. sp. pl. 5. p. 348.
Asplenium Scolopendrium. *Sp. pl. 1537.*
Icon. *Schkuhr filic. t. 83. Pluk. phytogr. t. 248. f. 1.*
In shady woods, among loose rocks in the western parts
New York, near Onondago, on the plantations of
J. Geddis, Esq. ♀. July. v. v. This species I have
seen in no other place but that here mentioned, nei-
ther have I had any information of its having been
found in any other part of North America.

fr. z.

gerri-
pl. 5.

mblea

auri-
radi-

f. 7.

July.
works,
gatedsub-
eriore
pl. 5.

i. 14.

parts
. Aerru-
pl. 5.

et

uly.

epa-
372.

729. PTERIS. *Gen. pl.* 1626.*pedata.*

1. *P. frondibus profundissime quinquelobo-palmatis, lobis pinnatifidis, lacinias linearis-lanceolatis acutis, infima lobi intermedii utrinque pinnatifida, lobo intermedio basi longissime cuneato, sinibus acutis.* — *Willd. sp. pl. 5. p. 358.*

*Icon. Schkuhr filic. t. 100. Plum. amer. t. 34. fil. t. 152.
Petiv. filic. t. 8. f. 12. Pluk. alm. t. 286. f. 5.*

On rocks: Virginia. ♀. July. v. s. About six inches high.

atropurpurea.

2. *P. frondibus pinnatis, pinnis inferioribus ternatis pinnatisve lanceolatis obtusis, basi obtuse truncatis subcordatis.* — *Willd. sp. pl. 5. p. 375.*

venosa.

- α. *P. pinnulis subtus venosis, stipite angulato.*

Icon. Schkuhr filic. t. 101. Pluk. mant. t. 349. f. 1.

- β. *P. pinnulis subtus punctatis, stipite terete.*

P. spiculata. Schkuhr filic. t. 100.

In clefts of rocks: Pennsylvania to Carolina. ♀. July. v. v. About a span high; grayish-green with dark purple stems.

gracilis.

3. *P. frondibus pinnatis, pinnis lanceolatis obtusis alternis sessilibus, inferioribus pinnatifidis, fertilibus integerimis, sterilibus crenatis rotundato-obtusis.* — *Willd. sp. pl. 5. p. 376.*

On rocks in Canada. *Michaux.* ♀. July. v. s. Not above four inches high; stems yellowish-green, smooth and shining.

caudata.

4. *P. fronde pinnato-tripartita, ramis sterilibus bipinnatis, pinnulis linearibus elongatis obtusis integerrimis, inferioribus bipinnatifidis, fructificantibus pinnatis, pinnis remotiusculis inferioribus basi pinnatifido-dentatis.* — *Willd. sp. pl. 5. p. 401.*

Icon. Jacq. ic. rar. 3. t. 645. Plum. amer. t. 22. filic. t. 29.

In shady stony woods, on the side of rivulets: Pennsylvania to Carolina. ♀. Aug. v. v.

aquilina.

5. *P. fronde pinnato-tripartita, ramis sterilibus bipinnatis, pinnis linearis-lanceolatis obtusis pinnatifido-dentatis, fructificantibus pinnatis, pinnulis pinnatifidis, lacinias acutiusculis, omnibus ciliatis.* — *Willd. sp. pl. 5. p. 402.*

Icon. Schkuhr filic. t. 95. 96. b. Plum. filic. t. A. f. 1.
In dry barren woods: Canada to Florida. ♀. Aug.

v. v. It c
plant, but
species, at
species its
ent soils.

730. VITTAR

1. *V. frondibus tariis intra V. angustifro Pteris lineata*
Icon. Schkuhr t. 143. P
On the banks *chaux.* ♀ common gr

731.

1. *B. frondibus obtusiusculi nearibus ac*
B. Spicant. R
Onoclea Spica
Osmunda Spic
Icon. Schkuhr
Moris. hist.
On the north-
in Herb. Le

2. *B. frondibus nuatis basi*
pl. 5. p. 411
On the banks *chaux.* ♀.

732. WOOD'

1. *W. frondibus repandis ten*
nis linearibus
p. 416.
W. angustifolia
W. floridana.
Onoclea nodula
synop. filic.

v. v. It deviates in some respects from the European plant, but is not sufficiently distinct to form a separate species, as it is well known how much the European species itself assumes different appearances in different soils.

730. VITTARIA. *Smith act. taur.* 5. p. 413. t. 9. f. 5.

1. V. frondibus linearibus longissimis pendulis, soris solitariis intra marginalibus. *Willd. sp. pl.* 5. p. 404.
V. angustifrons. Mich. fl. amer. 2. p. 201.
Pteris lineata. Sp. pl. 1530.
Icon. Schkuhr filic. t. 101, b. *Plum. amer.* t. 41. *filic.*
t. 143. Petiv. filic. t. 14. f. 3.
 On the banks of the rivulet Aisa-hatcha, Florida. *Michaux.* ♀. +. The leaves resemble those of a common grass.



lineata.

731. BLECHNUM. *Gen. pl.* 1627.

1. B. frondibus sterilibus pinnatifidis, laciniis lanceolatis obtusiusculis parallelis, fructiferis pinnatis, pinnis linearibus acuminatis. *Willd. sp. pl.* 5. p. 408.
B. Spicant. Roth. fl. germ. 3. p. 44.
Onoclea Spicant. Hoffm. fl. germ. 2. p. 11.
Osmunda Spicant. Sp. pl. 1522.
Icon. Schkuhr filic. t. 110. *Fl. dan.* 99. *Bolt. filic.* t. 6.
Moris. hist. 3. s. 14. t. 2. f. 23.
 On the north-west coast. *M. Lewis.* ♀. Aug. v. s. in *Herb. Lewis.*
2. B. frondibus pinnatis, pinnis lanceolatis utrinque attenuatis basi decurrentibus argute serratis. *Willd. sp. pl.* 5. p. 411.
 On the banks of the rivulet Aisa-hatcha, Florida. *Michaux.* ♀. +.

boreale.

732. WOODWARDIA. *Smith act. taur.* 5. p. 411.
t. 9. f. 5.

1. W. frondibus sterilibus pinnatifidis, laciniis lanceolatis repandis tenuissime serrulatis, fructiferis pinnatis, pinnis linearibus integrerrimis acutis. *Willd. sp. pl.* 5. p. 416.
W. angustifolia. Smith act. taur. 5. p. 411.
W. floridana. Schkuhr filic. p. 103. t. 111.
Onoclea nodulosa. Mich. fl. amer. 2. p. 272. *Swartz.*
synop. filic. 111.

onocleoides




virginica.

thelypteroides.



pedatum.

vestita.



- Acrostichum areolatum. Sp. pl. 1526.*
Osmunda caroliniana. Walt. fl. car. 257.
Icon. Schkuhr t. c. Pluk. phytogr. t. 399. f. 1. Moris. hist. 3. s. 14. t. 2 f. 24.
 In cedar and cypress-swamps: New Jersey to Florida.
 Ʌ. Aug. v. v. About a foot high.
2. *W. frondibus glaberrimis pinnatis, pinnis sessilibus lanceolatis pinnatifidis, lacinias oblongis obtusis crenulatis, fructiferis elongatis, stipite glabro tereti.—Willd. sp. pl. 5. p. 418.*
W. Banisteriana. Mich. fl. amer. 2. p. 263.
Blechnum virginicum. Mant. 307.
Blechnum carolinianum. Walt. fl. car. 257.
Icon. Pluk. alm. t. 179. f. 2.
 In swamps and shady woods: Virginia and Carolina. Ʌ. July. v. v. Above a foot high.
3. *W. frondibus pinnatis, pinnis sessilibus basi villosis linear-lanceolatis pinnatifidis, lacinias sterulum oblongis obtusiusculis, fertilium abbreviato-triangularibus acutis, omnibus integerrimis, stipite pubescente angulato.*
 In sandy swamps: South Carolina, near Charlestown.
 Ʌ. July. v. v. Resembles the preceding, but is not half the size.

733. ADIANTUM. *Gen. pl. 1633.*

1. *A. fronde pedata, ramis pinnatis, pinnis dimidiatis oblongis lunatis margine superiore incisis, lacinias sterilibus dentatis, fertilibus integerrimis, soris linearibus, stipite glabro. Willd. sp. pl. 5. p. 438.*
Icon. Schkuhr filic. t. 115. Pluk. alm. t. 124. f. 2. Corn. canad. t. 6.
 In shady woods: Canada to Virginia. Ʌ. July. v. v.
 A very elegant fern.

734. CHEILANTHES. *Swartz. synop. filic. p. 5.*

1. *C. frondibus bipinnatis utrinque hirtis, pinnulis pinnatifidis, lacinias oblongis obtusis integerrimis, stipite rachibusque hirtis. Willd. sp. pl. 5. p. 458.*
Adiantum vestitum. Strengel. anleit. 3. p. 122.
Aspidium lanosum. Swartz synop. filic. p. 58.
Neprodium lanosum. Mich. fl. amer. 2. p. 270.
Icon. Schkuhr filic. t. 124.

CR1

- On rocks:
 Ʌ. with long
 2. *C. frondibus apice*
 On rocks,
 A very c
C. tenuifl

735. DI

1. *D. frondibus natividis, chibus pil*
D. pubescens
Polypodium
 In shady wo
 Ʌ. v. v.
dilatatum.

736. HYMEN

1. *H. frondibus rioribus set bus obtusis alatis ciliat*
Trichomanes
 In shady fore
 nia and Vir

V. H

737.

1. *I. frondibus su*
p. 534.
Icon. Schkuhr
t. 80. f. 1.
 On rocks, on
 Oswego riv
 gular plant,
 high.

On rocks: Pennsylvania to Virginia and on the Missouri.
 ♀. July. v. v. A handsome very small fern, covered
 with long brownish hair.

2. C. frondibus triplicato-pinnatis, pinnulis ovalibus distinc-
 tis apice crenatis aut emarginatis subtus dealbatis.

dealbata.

On rocks, on the banks of the Missouri. ♀. July. v. s.
 A very delicate small fern, very much resembling
C. tenuifolia.

735. DICKSONIA. *L'Herit. sert. angl.* 30.

1. D. frondibus bipinnatis, pinnulis oblongo-lanceolatis pin-
 natifidis, laciniis margine superiore inciso-dentatis, ra-
 chibus pilosiusculis. *Willd. sp. pl. 5. p. 484.*

pilosiuscula.

D. pubescens. *Schkuhr filic. p. 125. t. 131.*
Polypodium pilosiusculum. Muhl. in litt.

In shady woods, among rocks: Pennsylvania and Virginia.
 ♀. v. v. A foot or more high, resembling *Aspidium*
dilatatum.

736. HYMENOPHYLLUM. *Smith act. taur. 5. p. 418.*

1. H. frondibus pinnatis, pinnis infimis majoribus, supe-
 rioribus sensim minoribus pinnatifidis, laciniis linear-
 bus obtusis bifidis ciliatis, venis hirtis, rachi stipiteque
 alatis ciliatis. *Willd. sp. pl. 5. p. 519.*

ciliatum.

Trichomanes ciliatum. *Swartz. prodr. 136.*
 In shady forests, on the trunks of old trees: Pennsylvania
 and Virginia. ♀. v. v.

V. HYDROPTERIDES.

737. ISOETES. *Gen. pl. 1620.*

1. I. frondibus subulatis semiteretibus. *Willd. sp. pl. 5.*
 p. 534.

lacustris.

Icon. *Schkuhr filic. t. 173. Fl. dan. 191. Dill. musc.*
t. 80. f. 1.

On rocks, on the bottom of rivers: New York, in the
 Oswego river, near the Falls. ♀. v. v. This sin-
 gular plant, resembling a zoophyte, is about two inches
 high.

738. SALVINIA. *Gen. pl.* 1617.*natans.*

1. *S. foliis ellipticis obtusis supra fasciculato-setosis, petiolis pilosis, fructibus subsessilibus aggregatis.* *Willd. sp. pl.* 5. p. 536.

Marsilea natans. Sp. pl. 1562.Icon. *Schkuhr filic. t.* 173. *Lam. illust.* 863.

Floating, like *Lemna*, on the surface of stagnant waters: in several of the small lakes in the western parts of New York and Canada. ○. v. v.

AN

739. AZOLLA. *Lam. encycl. 1. p.* 340.*caroliniana.*

1. *A. foliis imbricatis ovato-oblongis obtusis patentibus subtus rubris.* — *Willd. sp. pl.* 5. p. 541.

Floating on the waters of Lake Ontario. ○. v. v.

Resembles a floating moss or *Jungermannia*.

MO

Kyllingia. 47.
Canna. 585.

M

Agrostis. 64. :

DI

Salicornia. 2.
Lepidium.
Salix. 608.

Podostemum.
Holcus 78.

TR

Galium. 102.
Juncus. 23;
dica. 444.

T

Salix. 608. :

T

Amaranthus. 2.
Polygonum.

FINIS.

TABLE

OF

ANOMALOUS FLOWERS.

MONANDRIA. MONOGYNIA.

Kyllingia. 47. Cyperus. 50. Scirpus. 54. Thalia. 584.
Canna. 585. *ORCHIDEÆ* omnes 585—594.

MONANDRIA. DIGYNIA.

Agrostis. 64. Saccharum. 73. Uniola. 84. Alchemilla. 112.

DIANDRIA. MONOGYNIA.

Salicornia. 2. Boerhavia. 31. Kyllingia. 46. Anychia. 176.
Lepidium. 435. Cypripedium. 594. Vallisneria. 602.
Salix. 608.

DIANDRIA. DIGYNIA.

Podostemum. 3. Muhlenbergia. 61. Anthoxanthum. 65.
Holcus. 78. Festuca. 83. Salix. 608.

TRIANDRIA. MONOGYNIA.

Galium. 102. Anychia. 176. Prinos. 220. Brodiaea. 223.
Juncus. 237. Melothria. 444. Sicyos. 444. Momordica. 444.

TRIANDRIA. DIGYNIA.

Salix. 608. Comptonia. 635.

TRIANDRIA. TRIGYNIA.

Amaranthus. 206. Triglochin. 247. Phyllanthus. 443.
Polygonum. 269. Tragia. 604. Juniperus. 646.

TETRANDRIA. MONOGYNIA.

Gratiola. 12. Elytraria. 13. Monarda. 16. Collinsonia. 20.
Anychia. 176. Prinos. 220. Smilacina. 233. Clarkia.
260. Pothos. 398. *Class. XIII. DIDYNAMIA.* 401
—431. Diotis. 602. Alnus. 622.

TETRANDRIA. DIGYNIA.

Gentiana. 185. Myrica. 620. Juglans. 636. Morus. 639
Cupressus. 645. Thuya. 646.

TETRANDRIA. TETRAGYNIA.

Proserpinaca. 92. Myriophyllum. 274.

PENTANDRIA. MONOGYNIA.

Catalpa. 10. Cornus. 107. Echites. 178. Gonolobium.
178. Apocynum. 179. Asclepias. 180. Ceropagia. 184.
Amsonia. 184. Gelsemium. 184. Gentiana. 185.
Prinos. 220. Polygonum. 269. Cassia. 307. Passiflora.
444. Illecebrum. 445. Galax. 446. Lobelia. 446.
Petalostemum. 461. *Class. XVII. SYNGENESIA.*
491—582.

PENTANDRIA. DIGYNIA.

Polygonum. 270. Scleranthus. 315. Iresine. 445. Salix.
608. Quercus. 625.

PENTANDRIA. TRIGYNIA.

Polygonum. 270. Arenaria. 319. Hypericum. 378. Pas-
siflora. 444.

PENTANDRIA. PENTAGYNIA.

Spergula. 320. Cerastium. 320.

HEXANDRIA. MONOGYNIA.

Commelina. 31. Trientalis. 252. Laurus. 274. Lythrum.
334. *Class. XIV. TETRADYNAMIA.* 432—441.
Corydalis. 462. Fumaria. 463. Aristolochia. 596.

HEXANDRIA. DIGYNIA.

Polygonum. 271.

HEXANDRIA. TRIGYNIA.

Lechea. 90.

T

HE

Myriophyllum

HEP

Sabbatia. 137.

HE

Polygonum. 27

HEP

Saururus. 252.

OCT

Æsculus. 253.

notropis. 303

lus. 618.

OC

Polygonum. 271

OCTA

Myriophyllum. 2

OCTA

Sedum. 323.

ENNE

Acer. 267.

ENN

Lechea. 90. Elo

DECA

Sabbatia. 137. A

264. Jeffersonia

Galax. 446. Ge

Class. XVI. D

Carpinus. 623.

DEC

Chrysosplenium. 2

DEC

Jatropha. 603. C

nea. 624.

HEXANDRIA. TETRAGYNIA.

Myriophyllum. 274.

HEPTANDRIA. MONOGYNIA.

Sabbatia. 137.

HEPTANDRIA. TRIGYNIA.

Polygonum. 271.

HEPTANDRIA. TETRAGYNIA.

Saururus. 252.

OCTANDRIA. MONOGYNIA.

Aesculus. 253. *Gaultheria*. 282. *Vaccinium*. 284. *Monetropia*. 303. *Halesia*. 449. *Polygala*. 464. *Populus*. 618.

OCTANDRIA. DIGYNIA.

Polygonum. 271. *Corylus* 634.

OCTANDRIA. TETRAGYNIA,

Myriophyllum. 274.

OCTANDRIA. PENTAGYNIA.

Sedum. 323.

ENNEANDRIA. MONOGYNIA.

Acer. 267.

ENNEANDRIA. TRIGYNIA.

Lechea. 90. *Elodea*. 379.

DECANDRIA. MONOGYNIA.

Sabbatia. 137. *Azalea*. 151. *Asclepias*. 180. *Menziesia*. 264. *Jeffersonia*. 268. *Lythrum*. 334. *Portulaca*. 365. *Galax*. 446. *Geranium* 448. *Halesia*. 449. *Styrax*. 450. *Class. XVI. DIADELPHIA*. 460—490. *Betula*. 620. *Carpinus*. 623. *Ostrya*. 623.

DECANDRIA. DIGYNIA.

Chrysosplenium. 269. *Fagus*. 624. *Quercus*. 625.

DECANDRIA. TRIGYNIA.

Jatropha. 603. *Croton*. 603. *Acalypha*. 604. *Casta-nea*. 624.

DECANDRIA. PENTAGYNIA.

Drosera. 210. Portulacca. 365. Geranium. 448.

DODECANDRIA. MONOGYNIA.

Sabbatia. 137. Decumaria. 328. Lythrum. 394. Cuphea. 335. Bejaria. 362. Hudsonia. 364. Talinum. 365. Portulacca. 365. Cleome. 441. Halesia. 449. Styrax. 450. Asarum. 596. Betula. 620.

DODECANDRIA. DIGYNIA.

Agrimonia. 335. Calligonum. 370. Fagus. 624.

DODECANDRIA. TRIGYNIA.

Datisca. 370. Menispermum. 371. Eloëda. 379. Croton. 603. Acalypha. 604. Euphorbia. 605.

POLYANDRIA. MONOGYNIA.

Jeffersonia. 268. *Class. XV. MONADELPHIA.* Polyandria. 449.—456. Asarum. 596. Taxus 647.

POLYANDRIA. DIGYNIA.

Fothergilla. 335.

POLYANDRIA. TRIGYNIA.

Ricinus 602. Juglans 636. Zamia. 648.

POLYANDRIA. TETRAGYNIA.

Myriophyllum. 274.

POLYANDRIA. POLYGYNIA.

Zostera. 2.

DIDYNAMIA: GYMNOSPERMIA.

Lycopus. 16. Monarda. 16. Salvia. 19. Collinsonia. 20.

DIDYNAMIA. ANGIOSPERMIA.

Catalpa. 10. Gratiola. 12. Elytraria. 13.

MONADELPHIA. TRIANDRIA.

Sisyrinchium. 31. Pistia. 268.

MONADELPHIA. PENTANDRIA.

Anagallis. 135. Lysimachia. 135. Linum. 210. Petalostemum. 461.

T.

MON

Cardiospermum

MON

Acacia. 305.

DIADELPHI

MONA

Pinus. 639.

DIA

Podalyria. 307.

POLY.

Ascyrum. 373.

451.

SYN

Solanum. 156.

444. Mom

GYN

Passiflora. 444.

GYN

Cleome. 441.

GYN

Zostera. 2. Po

MO

Zostera. 2. C

Callitriches. 3

Arum. 399.

MI

Podostemum. 3.

MC

Sparganium. 33

Zea 40. M

Sorghum. 78.

88. Eriocal

Phyllanthus.

eyos. 444.

MONADELPHIA. OCTANDRIA.

Cardiospermum. 273. *Polygala.* 464.

MONADELPHIA. DECANDRIA.

Acacia. 305. *Schrankia.* 305. *Oxalis.* 322. *Class. XVI.*
DIADELPHIA. Decandria. 466—490.

MONADELPHIA. POLYANDRIA.

Pinus. 639.

DIADELPHIA. DECANDRIA.

Podalyria. 307. *Cercis.* 308. *Virgilia.* 309.

POLYADELPHIA. POLYANDRIA.

Ascyrum. 373. *Hypericum.* 374. *Elodea.* 379. *Hopoe.*
451.

SYNGENESIA. MONOGAMIA.

Solanum. 156. *Impatiens.* 171. *Viola.* 171. *Melothria.*
444. *Momordica.* 444. *Sicyos.* 444. *Lobelia.* 446.

GYNANDRIA. PENTANDRIA.

Fassiflora. 444.

GYNANDRIA. HEXANDRIA.

Cleome. 441.

GYNANDRIA. POLYANDRIA.

Zostera. 2. *Pothos.* 398. *Calla.* 399. *Arum.* 399.

MONOECIA. MONANDRIA.

Zostera. 2. *Caulinia.* 2. *Salicornia.* 2. *Hippuris.* 3.
Callitrichie. 3. *Zannichellia.* 4. *Chara.* 4. *Calla.* 399.
Arum. 399. *Liquidambar.* 635. *Platanus.* 635.

MONOECIA. DIANDRIA.

Podostemum. 3. *Fraxinus.* 8. *Lemna.* 22.

MONOECIA. TRIANDRIA.

Sparganium. 33. *Typha.* 34. *Carex.* 34. *Scleria.* 45.
Zea. 40. *Milium.* 63. *Andropogon.* 74. *Holcus.* 78.
Sorghum. 78. *Chloris.* 87. *Rottboellia.* 88. *Tripsacum.*
88. *Eriocaulon.* 91. *Empetrum.* 93. *Amaranthus.* 206.
Phyllanthus. 443. *Melothria.* 444. *Momordica.* 444. *Si-*
cyos. 444. *Tragia.* 604. *Comptonia.* 635.

MONOECIA. TETRANDRIA.

Serpicula 33. Boehmeria. 112. Urtica. 112. Parietaria. 114. Planera. 115. Pachysandra. 117. Myriophyllum. 274. Alnus. 622. Morus. 639.

MONOECIA. PENTANDRIA.

Nyssa. 177. Panax. 191. Atriplex. 199. Celtis. 200. Rhus. 204. Crotonopsis 206. Amaranthus. 206. Zanthorrhiza. 212. Schisandra. 212. Iva. 580. Ambrosia. 580. Xanthium. 581. Fagus. 624. Quercus. 625.

MONOECIA. HEXANDRIA.

Zizania. 60. Eriocaulon. 91. Planera. 115. Celtis. 200. Melanthium. 240. Veratrum. 242. Myriophyllum. 274. Betula. 620. Quercus. 625.

MONOECIA. OCTANDRIA.

Acer. 265.

MONOECIA. POLYANDRIA.

Sparganium. 33. Acer. 267. Myriophyllum. 274. Laurus. 275. Acacia. 305. Schrankia. 305. Chrysobalanus. 329. Ceratophyllum. 370. Sagittaria. 395. Pothos. 398. Calla. 399. Arum. 399. Betula. 620. Carpinus. 623. Ostrya. 623. Fagus. 624. Castanea. 624. Quercus. 625. Corylus. 634. Juglans. 636. Liquidambar. 635. Platanus. 635.

MONOECIA. MONADELPHIA.

Podostemum. 3. Phyllanthus. 443. Melothria. 444. Momordica. 444. Sicyos. 444. Ricinus. 602. Jatropha. 603. Croton. 603. Acalypha. 604. Euphorbia. 605. Stillingia. 608. Platanus. 635. Pinus. 639. Cupressus. 645. Thuja. 646.

MONOECIA. SYNGENESIA.

Schisandra. 212. Melothria. 444. Momordica. 444. Sicyos. 444.

DIOECIA. MONANDRIA.

Najas. 602. Salix. 608.

Olea. 7. Frax
lisneria. 602

Carex. 34. E
220. Salix.

DI
Urtica. 112.
115. Ilex. 1

DI
Caprifolium. 16
178. Zizyp
Cannabis. 199
Acnida. 208.
sine. 445. Sa

DI
Prinos. 220. G
mex. 249. St

DI
Diosyru. 265.
618.

DIC
Gymnocladus. 30

DIOE
Calligonum. 370

DIO
Spiraea. 343. Ru

DIOE
Menispermum. 37
Sagittaria. 395.

DIOEC
Iresine. 445. Sida

TABLE OF ANOMALOUS FLOWERS.

679

DIOECIA. DIANDRIA.

Olea. 7. Fraxinus. 8. Ceratiola. 21. Borya. 22. Vallisneria. 602. Salix. 608.

DIOECIA. TRIANDRIA.

Carex. 34. Empetrum. 93. Zanthoxylum. 209. Prinos. 220. Salix. 608.

DIOECIA. TETRANDRIA.

Urtica. 112. Viðcum. 114. Elæagnus. 114. Hippophæ. 115. Ilex. 117. Prinos. 220. Myrica. 620.

DIOECIA. PENTANDRIA.

Caprifolium. 161. Vitis. 169. Nyssa. 177. Hamiltonia. 178. Zizyphus. 188. Panax. 191. Atriplex. 199. Cannabis. 199. Humulus. 199. Celtis. 200. Rhus. 204. Acnida. 208. Zanthoxylum. 209. Prinos. 220. Iresine. 445. Salix. 608.

DIOECIA. HEXANDRIA.

Prinos. 220. Gleditschia. 221. Chamærops. 239. Rhus. 249. Smilax. 249. Dioscorea. 251. Laurus. 275.

DIOECIA. OCTANDRIA.

Diosyrus. 265. Acer. 268. Polygonum. 272. Populus. 618.

DIOECIA. DECANDRIA.

Gymnocladus. 304.

DIOECIA. DODECANDRIA.

Calligonum. 370. Datisca. 370. Menispermum. 370.

DIOECIA. ICOSANDRIA.

Spiræa. 343. Rubus. 346.

DIOECIA. POLYANDRIA.

Menispermum. 370. Clematis. 384. Thalictrum. 388. Sagittaria. 395. Arum. 399. Populus. 618. Zamia. 648.

DIOECIA. MONADELPHIA.

Iresine. 445. Sida. 453. Juniperus. 646. Taxus. 646.

POLYGAMIA. MONOECIA.

Fraxinus. 8. *Andropogon*. 74. *Atheropogon*. 75. *Holcus*. 78. *Sorghum*. 78. *Chloris*. 87. *Parietaria*. 114. *Planera*. 115. *Ilex*. 117. *Atriplex*. 199. *Celtis*. 200. *Rhus*. 204. *Zanthorrhiza*. 212. *Melanthium*. 240. *Vernatum*. 242. *Aesculus*. 254. *Acer*. 265. *Acacia*. 305. *Schrankia*. 305.

POLYGAMIA. DIOECIA.

Fraxinus. 8. *Empetrum*. 93. *Nyssa*. 177. *Hamiltonia*. 178. *Panax*. 191. *Gleditschia*. 221. *Chamærops*. 239. *Diospyros*. 265.

ACER.

— *saccharinum*.
— *Negundo*.
— *striatum*.
Achillea.
— *Millefolium*.
— *Pterniifolia*.
Acnida.
Aconitum.
Acorus.
Acrostichum.
Actaea.

Adiantum.
Aesculus.

Agave.
Agrimonia.
Agrostemma.
Agrostis.
Aira.
Ajuga.
Alchemilla.
— *Apophyllum*.
Aletris.
Alisma.
Allium.
Alnus.
Alopecurus.

Hol-
114.
200.
Ve-
.305.

ltonia.
.239.

LATIN AND ENGLISH

INDEX.

A C E R.

— saccharinum.
— Negundo.
— striatum.
Achillea.
— Millefolium.
— Ptarmica.

Acnida.
Aconitum.
Acorus.
Acrostichum.
Actaea.

Adiantum.

Æsculus.

Agave.

Agrimonia.

Agrostemma.

Agrostis.

Aira.

Ajuga.

Alchemilla.

— Aphanes.

Aletris.

Alisma.

Allium.

Alnus.

Alopecurus.

<i>Maple-tree.</i>	265
<i>Sugar-tree.</i>	266
<i>Flowering Ash. Box Elder.</i>	268
<i>Moose-wood.</i>	267
<i>Milfoil.</i>	562
<i>Yarrow.</i>	563
<i>Sneeze-wort.</i>	562
<i>River-hemp.</i>	208
<i>Wolf's-bane. Monk's-hood.</i>	372
<i>Sweet-flag.</i>	235
<i>Full-fern.</i>	658
<i>Herb Christopher. Baneberry.</i>	
<i>Red and White Cohosh.</i>	366
<i>Maiden-hair.</i>	733
<i>Horse Chesnut. Buck's-eye</i>	
<i>Tree.</i>	254
<i>Sweet Virginian Aloe.</i>	226
<i>Agrimony.</i>	335
<i>Corn Cockle.</i>	321
<i>Bent-grass.</i>	63
<i>Hair-grass.</i>	76
<i>Bugle. Ground Pine.</i>	404
<i>Ladies' Mantle.</i>	112
<i>Parsley-piert.</i>	112
<i>Star-wort. Devil's-bit.</i>	225
<i>Water Plantain.</i>	352
<i>Gaſtic.</i>	222
<i>Alder.</i>	622
<i>Fox-tail grass.</i>	65

y 2

Alyssum.	<i>Mad-wort.</i>	434	Asclepias tu
Amaranthus.	<i>Amaranth.</i>	206	Ascyrum Cr
— Blitum.	<i>Blite.</i>	207	Aspidium.
— hypochondriacus.	<i>Prince's Feather.</i>	207	Asplenium.
Amaryllis.	<i>Atamasco Lily.</i>	222	— Rut
Ambrosia.	<i>Hog- or Iron-weed.</i>	580	Aster.
Ammi.	<i>Bishop's-weed.</i>	192	Astragalus.
Ammyrsine.	<i>Sand-myrtle.</i>	301	Atriplex.
Amorpha.	<i>Bastard Indigo.</i>	466	Avena.
Anagallis.	<i>Pimpernel.</i>	135	Azalea.
Andromeda.	<i>Bastard Huckle-berry.</i>	290	
— arborea.	<i>Sorrel-tree.</i>	205	
— polifolia.	<i>Wild Rosemary.</i>	291	Baccharis.
Andropogon.	<i>Man's Beard.</i>	74	Batschia.
Anemone.	<i>Anemone. Wind-flower.</i>	386	Bejaria.
Anthemis.	<i>Chamomile.</i>	562	Bellis.
— arvensis.	<i>May-weed.</i>	562	Berberis.
Anthoxanthus.	<i>Spring- or Sweet-grass.</i>	65	Betula.
Antirrhinum.	<i>Snap-dragon.</i>	421	— lenta.
— Elatine.	<i>Flueulin.</i>	421	Bidens.
— Linaria.	<i>Toad-flax.</i>	421	Bignonia.
— Oronthium.	<i>Calf's Snout.</i>	422	Blechnum.
Apocynum.	<i>Dog's Bane. Indian Hemp.</i>	179	Blitum.
Aquilegia.	<i>Columbine. Honey-suckle.</i>	372	Botrychium.
Arabis.	<i>Wall-cress.</i>	436	Brassica.
Aralia nudicaulis.	<i>Sarsaparilla.</i>	209	Briza.
— hispida.	<i>Poison Elder.</i>	209	Bromus.
— spinosa.	<i>Angelica-tree.</i>	209	Buphthalmum
— racemosa.	<i>Spikenard.</i>	209	
Arbutus.	<i>Strawberry-tree.</i>	282	Cactus.
— Uva ursi.	<i>Bear-berry. Saca-commis.</i>	283	Cakile.
Arctium.	<i>Burdock. Clot-bur.</i>	505	Callitrichie.
Arenaria.	<i>Sand-wort.</i>	317	Caltha.
Aristolochia.	<i>Birth-wort.</i>	396	Calycanthus.
— Serpentaria.	<i>Virginia Snake-root.</i>	396	
Artemisia.	<i>Wormwood.</i>	521	Campanula.
— Dracunculus.	<i>Tarragon.</i>	521	Canna.
— vulgaris.	<i>Mug-wort.</i>	522	Cannabis.
Arum.	<i>Wake-Robin.</i>	399	Caprifolium.
— Dracontium.	<i>Green Dragon.</i>	400	Cardamine.
Arundo.	<i>Reed-grass.</i>	86	Cardiospermum.
— arenaria.	<i>Sea-mat-grass.</i>	87	Carduus.
Asarum.	<i>Asarubacea.</i>	596	Carex.
— canadense.	<i>Wild Ginger. Colt's Foot.</i>	596	Carpinus.
Asclepias.	<i>Swallow-wort.</i>	180	

434		
206		
207		
207		
222		
580		
192		
301		
466		
135		
290		
295		
291		
74		
386		
562		
562		
65		
421		
421		
421		
422		
179		
372		
436		
209		
209		
209		
209		
282		
283		
505		
317		
596		
596		
521		
521		
522		
399		
400		
86		
87		
596		
596		
180		
	Aselepias tuberosa.	
	Ascyrum Crux.	
	Aspidium.	
	Asplenium.	
	— Ruta muraria.	
	Aster.	
	Astragalus.	
	Atriplex.	
	Avena.	
	Azalea.	
	Baccharis.	
	Batschia.	
	Bejaria.	
	Bellis.	
	Berberis.	
	Betula.	
	— lenta.	
	Bidens.	
	Bignonia.	
	Blechnum.	
	Blitum.	
	Botrychium.	
	Brassica.	
	Briza.	
	Bromus.	
	Buphthalmum.	
	Cactus.	
	Cakile.	
	Callitriches.	
	Caltha.	
	Calycanthus.	
	Campanula.	
	Canna.	
	Cannabis.	
	Caprifolium.	
	Cardamine.	
	Cardiospermum.	
	Carduus.	
	Carex.	
	Carpinus.	
	Ipecacuanha.	
	Butterfly-flower.	
	St. Andrew's Cross.	
	Shield Fern.	
	Spleen-wort.	
	Wall-Rue.	
	Star-wort.	
	Milk Vetch.	
	Orache.	
	Oat-grass. Oats.	
	Upright or American Honey-suckle.	
	Groundsel-tree.	
	Puccoon. Indian Paint.	
	Cat-h-fly-bush.	
	Daisy.	
	Berberry. Pipperidge.	
	Birch-tree.	
	Mountain-Mahogany.	
	Bur-marygold.	
	Trumpet-flower.	
	Rib Fern.	
	Strawberry-blite.	
	Rattle Snake Fern.	
	Cabbage.	
	Quaking-grass.	
	Brome grass.	
	Ox-eye.	
	Prickly-pear. Indian Fig.	
	Sea-rocket.	
	Water Star-wort.	
	Marsh-marygold.	
	Carolina Allspice. Sweet-scented shrub.	
	Bell-flower.	
	Indian Reed or Shot.	
	Hemp.	
	Honey-suckle. Woodbine.	
	Ladies' Smock.	
	Heart-seed.	
	Thistle.	
	Sedge.	
	Hornbeam-tree.	

Cassia nictitans.	Sensitive Pea.	305
Castanea.	Chesnut-tree.	624
— pumila.	Chinquapin.	625
Catalpa.	Catalpa-tree.	10
Caulophyllum.	Blue Cohosh.	218
Ceanothus.	Red-root. New Jersey Tea.	167
Celastrus.	Staff-tree.	167
Celtis.	Nettle-tree. Hag-berry. Hoop-ash.	200
Cenchrus.	Bur-grass.	60
Centaurea.	Centaury.	576
— Cyanus.	Blue-bottle.	576
— solstitialis.	St. Barnaby's Thistle.	576
Centunculus.	Chaff-weed.	97
Cephalanthus.	Button-wood.	114
Cerastium.	Mouse-ear Chick-weed.	320
Ceratophyllum.	Horn-wort.	370
Cercis.	Judas-tree.	308
Chara.	Feather-beds.	4
Chærophyllosum.	Cow-parsley. Wild Cher- vil.	195
Chamærops.	Palmetto-tree. Cabbage- tree.	239
Cheiranthus.	Stock. Wall-flower.	436
Chelidonium.	Celandine.	365
Chenopodium.	Goose-foot.	197
— viride.	Lamb's-quarter.	198
— Anthelminticum.	Worm-seed.	198
Chimaphila.	Winter-green. Sipsisewa.	300
Chiococca.	Snow-berry.	160
Chionanthus.	Fringe-tree. Snow-flock-tree. Snow-flower.	7
Chloris.	Wire-grass. Couch-grass.	87
Chondrilla.	Gum Succory.	497
Chrysanthemum.	Ox-eye Daisy.	526
Chrysocoma.	Goldilocks.	517
Chrysosplenium.	Golden Saxifrage.	269
Cichorium.	Succory.	496
Cicuta.	Water Cowbane.	195
Cimicifuga.	Bug-wort.	372
— Serpentaria.	Black Snake-root.	372
Circæa.	Enchanters' Nightshade.	21
Cissus.	Pepper-vine.	170
— hederacea.	Virginia Creeper.	170

Clematis.
 Clinopodium.
 Cnicus.
 Collinsonia.
 Comarum.
 Comptonia.
 Conium.
 Convallaria.
 Convolvulus.
 Conzya.
 Coreopsis.
 Corisperma.
 Cornus.
 — sangi.
 Coronopus.
 Corydalis.
 — Cucu.
 Corylus.
 — rostra.
 Crataegus.
 Crotalaria.
 Cucubalus.
 Cunila.
 Cupressus.
 — thysoid.
 Cuscuta.
 Cynoglossum.
 Cyperus.
 Cypripedium.
 Dactylis.
 Datura.
 Daucus.
 Delphinium.
 Dentaria.
 — diphyll.
 Dianthus.
 Dionaea.
 Dioscorea.
 Diospyros.

305	Clematis.	<i>Virgin's Bower. Traveller's-joy.</i>	384
624	<i>Clinopodium.</i>	<i>Wild Basil.</i>	410
625	<i>Cnicus.</i>	<i>Thistle.</i>	506
10	<i>Collinsonia.</i>	<i>Horse-weed.</i>	20
218	<i>Comarum.</i>	<i>Marsh Cinquefoil.</i>	356
167	<i>Comptonia.</i>	<i>Sweet Fern-bush.</i>	633
167	<i>Conium.</i>	<i>Hemlock.</i>	195
	<i>Convallaria.</i>	<i>Lily of the Valley. Sneeze-wort.</i>	232
200	<i>Convolvulus.</i>	<i>Bind-weed.</i>	143
60	<i>Conzya.</i>	<i>Flea-bane.</i>	523
576	<i>Coreopsis.</i>	<i>Tick-seed Sunflower.</i>	567
576	<i>Corispermum.</i>	<i>Tick-seed.</i>	4
97	<i>Cornus.</i>	<i>Dogwood.</i>	107
114	— <i>sanguinea.</i>	<i>Red Osier.</i>	109
320	<i>Coronopus.</i>	<i>Swine's-Cress.</i>	435
370	<i>Corydalis.</i>	<i>Fumitory.</i>	402
308	— <i>Cucullaria.</i>	<i>Breeches-flower.</i>	462
4	<i>Corylus.</i>	<i>Hazel-nut. Filbert.</i>	634
	— <i>rostrata.</i>	<i>Cuckold's Nut.</i>	635
195	<i>Crataegus.</i>	<i>Hawthorn.</i>	336
	<i>Crotalaria.</i>	<i>Rattle-pea.</i>	469
239	<i>Cucubalus.</i>	<i>Campion.</i>	315
436	<i>Cunila.</i>	<i>Dittany.</i>	406
365	<i>Cupressus.</i>	<i>Cypress.</i>	645
197	— <i>thyoides.</i>	<i>White Cedar.</i>	646
98	<i>Cuscuta.</i>	<i>Dodder.</i>	116
98	<i>Cynoglossum.</i>	<i>Hound's-tongue.</i>	133
100	<i>Cyperus.</i>	<i>Nut-grass.</i>	50
60	<i>Cypripedium.</i>	<i>Ladies' Slipper. Moccasin-flower.</i>	594
7	<i>Dactylis.</i>	<i>Cock's-foot-grass. Orchard-grass.</i>	84
87	<i>Datura.</i>	<i>Thorn-apple. Stramonium.</i>	141
97	<i>Daucus.</i>	<i>Carrot.</i>	191
26	<i>Delphinium.</i>	<i>Lark-spur.</i>	371
17	<i>Dentaria.</i>	<i>Tooth-wort.</i>	438
59	— <i>diphylla.</i>	<i>Pepper-root. Indian Mustard.</i>	438
16	<i>Dianthus.</i>	<i>Pink.</i>	314
35	<i>Dionaea.</i>	<i>Venus's Fly-trap.</i>	304
2	<i>Dioscorea.</i>	<i>Yam.</i>	251
2	<i>Diospyros.</i>	<i>Date-plum. Persimon.</i>	265
1			
0			
0			

LATIN AND ENGLISH INDEX.

Dipsacus.	<i>Teasel. Fullers-thistle.</i>	96	Gelsemin
Dirca.	<i>Leather-wood.</i>	268	Gentiana
Dodecatheon.	<i>American Cowslip. Mitre-flower.</i>	136	Geranium
Draba.	<i>Whitlow-grass.</i>	433	— n
Dracocephalum.	<i>Dragon's-head.</i>	411	— R
Drosera.	<i>Sun-dew.</i>	210	Geum.
Echium.	<i>Viper's Bugloss.</i>	130	Glaux.
Elaeagnus.	<i>Missouri Silver-tree.</i>	114	Glechoma
Elephantopus.	<i>Elephant's-foot.</i>	582	Gleditschi
Elymus.	<i>Lyme-grass. Wild Rye.</i>	89	Glycirrhiz
Empetrum.	<i>Crake- or Crow-berry.</i>	93	Gnaphaliu
Epigaea.	<i>Ground Ivy.</i>	297	Gordonia.
Epilobium.	<i>Willow-herb.</i>	259	Gratiola.
Equisetum.	<i>Horse-tail. Shave-grass.</i>	651	Gymnoclad
Erigeron	<i>Rag-weed.</i>	532	
— bellidifolium.	<i>Poor Robin's Plantain. Rattle-snake Plantain.</i>	532	Halesia.
Eriocaulon.	<i>Pipe-wort.</i>	91	Hamamelis
Eriophorum.	<i>Cotton-grass.</i>	57	Hamiltonia
Eryngium.	<i>Eryngio.</i>	189	Hedeoma
Erysimum.	<i>Hedge Mustard.</i>	436	Helianthen
— cheiranthoides.	<i>Worm-seed.</i>	436	Helianthus
Erythrina.	<i>Coral-plant.</i>	467	Heliotropiu
Erythronium.	<i>Dog's-tooth Violet.</i>	230	Helonias di
Eupatorium.	<i>Hemp Agrimony.</i>	512	Hepatica.
— perfoliatum.	<i>Thorough-wort. Bone-set.</i>	516	Heracleum.
Euphrasia.	<i>Eye-bright.</i>	430	Hesperis.
Evonymus.	<i>Spindle-tree.</i>	168	Hibiscus esc
— americanus.	<i>Burning-bush.</i>	168	Hieracium.
Fagus.	<i>Beech-tree.</i>	624	Hippophaë.
Festuca.	<i>Fescue-grass.</i>	83	Hippuris.
Fragaria.	<i>Strawberry.</i>	357	Holcus fragi
Fraxinus.	<i>Ash tree.</i>	8	Hopea.
Fumaria.	<i>Fumitory.</i>	463	Hordeum.
Galax.	<i>Carpenter's-leaf. Heal-all.</i>	446	Hottonia.
Galeopsis.	<i>Dead-nettle.</i>	407	Hudsonia.
Galium.	<i>Ladies' Bed-straw. Cleavers.</i>		Humulus.
Gaultheria.	<i>Goose-grass.</i>	102	Hydrastis.
	<i>Mountain-tea. Partridge-berry.</i>	283	Hydrocotyle.
			Hydrophyllu
			Hyoscyamus,
			Hyoseris.

LATIN AND ENGLISH INDEX.

687

96	Gelsemium.	Carolina Jasmine.	184
268	Gentiana.	Gentian.	185
	Geranium.	Crane's-bill.	448
136	— maculatum.	Alum-root. Flux-weed.	448
433	— Robertianum.	Herb-Robert.	449
411	Geum.	Avens. Herb Bennet.	351
210	Glaux.	Black Salt-wort.	176
	Glechoma.	Ground Ivy.	408
130	Gleditschia.	Honey Locust-tree.	221
114	Glycirrhiza.	Liquorice.	480
582	Gnaphalium.	Everlasting. Cudweed.	524
89	Gordonia.	Loblolly Bay.	451
93	Gratiola.	Hedge-hyssop.	12
297	— aurea.	Golden-pert.	12
259	Gymnocladus.	Kentucky Coffee-tree. Chicot.	304
651			
532	Halesia.	Snow-drop-tree. Silver-bell-tree.	449
91	Hamamelis.	Witch-hazel.	116
57	Hamiltonia.	Oil-nut.	178
189	Hedeoma.	Pennyroyal.	414
436	Helianthemum.	Rock rose.	363
436	Helianthus.	Sun-flower.	570
467	Heliotropium.	Turnsol.	130
230	Helonias dioica.	Star-wort. Colic-wort.	243
512	Hepatica.	Liver-wort.	391
516	Heracleum.	Cow Parsnep.	492
430	Hesperis.	Rocket.	436
168	Hibiscus esculentus.	Okra or Okro.	456
168	Hieracium.	Hauk-weed.	502
	Hippophaë.	Sea Buckthorn.	115
	Hippuris.	Mare's-tail.	3
624	Holcus fragrans.	Seneca-grass.	78
83	Hopea.	Sweet-leaf.	451
357	Hordeum.	Barley.	89
8	Hottonia.	Water-violet. Feather-foil.	138
163	Hudsonia.	Yellow American Heath.	364
146	Humulus.	Hop.	199
107	Hydrastis.	Yellow-root. Turmeric.	389
02	Hydrocotyle.	Penny-wort.	190
	Hydrophyllum.	Water-leaf.	134
63	Hyoscyamus.	Henbane.	141
	Hyoseris.	Swine's Succory.	504

<i>Hypericum.</i>	<i>St. John's-wort.</i>	374	<i>Lilium.</i>
— <i>densiflorum.</i>	<i>American Yellow-heath.</i>	376	— <i>car.</i>
<i>Jatropha.</i>	<i>Wild Cassava. Tread-</i>		<i>Linum.</i>
	<i>softly.</i>	603	<i>Liquidambar.</i>
<i>Ilex.</i>	<i>Holly. Dahoон.</i>	117	<i>Liriodendron.</i>
— <i>vomitoria.</i>	<i>South-sea Tea.</i>	118	
<i>Illecebrum.</i>	<i>Knot grass.</i>	445	<i>Lithospermum.</i>
<i>Illicium.</i>	<i>Aniseed-tree.</i>	380	<i>Lobelia Cardinalis.</i>
<i>Impatiens.</i>	<i>Balsam. Touch-me-not.</i>	171	— <i>infl.</i>
<i>Indigofera.</i>	<i>Indigo-plant.</i>	488	<i>Lotus.</i>
<i>Inula Helenium.</i>	<i>Elecampane.</i>	531	<i>Lupinus.</i>
<i>Ipomoea Quamoclit.</i>	<i>Cypress-vine.</i>	145	<i>Lycium.</i>
<i>Iris.</i>	<i>Flag. Flower-de-luce.</i>	29	<i>Lycopodium.</i>
<i>Juglans.</i>	<i>Walnut. Hickory.</i>	636	— <i>denc.</i>
— <i>cineraria.</i>	<i>Butter-nut. Oil-nut.</i>	636	<i>Lycopsis.</i>
— <i>alba.</i>	<i>Shell- or Shag-bark.</i>	637	<i>Lycopus.</i>
— <i>olivæformis.</i>	<i>Pecan-nut.</i>	636	<i>Lysimachia.</i>
— <i>porcina.</i>	<i>Hog- or Pig-nut.</i>	638	
<i>Juncus.</i>	<i>Rush.</i>	235	
<i>Juniperus.</i>	<i>Juniper.</i>	646	<i>Magnolia acuminata.</i>
— <i>virginiana.</i>	<i>Red Cedar.</i>	647	— <i>auriculata.</i>
<i>Iva.</i>	<i>Bastard Jesuit's-bark.</i>	580	— <i>glaucophylla.</i>
<i>Kalmia.</i>	<i>American Laurel.</i>	295	
— <i>angustifolia.</i>	<i>Sheep Laurel.</i>	296	— <i>grandiflora.</i>
— <i>latifolia.</i>	<i>Calico-flower.</i>	296	— <i>tripetala.</i>
<i>Lactuca.</i>	<i>Lettuce.</i>	500	<i>Malva.</i>
<i>Lamium.</i>	<i>Archangel.</i>	406	<i>Marrubium.</i>
— <i>amplexicaule.</i>	<i>Henbit.</i>	406	<i>Martynia.</i>
<i>Laurus.</i>	<i>Laurel. Sweet-bay.</i>	275	<i>Medicago.</i>
— <i>Benzoin.</i>	<i>Spice-wood. Benjamin-tree.</i>	276	<i>Medeola.</i>
— <i>carolinensis.</i>	<i>Red-bay.</i>	276	<i>Melampyrum.</i>
— <i>Sassafras.</i>	<i>Sassafras-tree.</i>	277	<i>Melica.</i>
<i>Ledum latifolium.</i>	<i>Labrador Tea.</i>	300	<i>Melilotus.</i>
<i>Lemna.</i>	<i>Duck's-meat.</i>	22	<i>Menispernum.</i>
<i>Leontodon.</i>	<i>Dandelion.</i>	497	— <i>canadense.</i>
<i>Lepidium.</i>	<i>Pepper-wort.</i>	435	<i>Mentha.</i>
<i>Liatris.</i>	<i>Button Snake-root.</i>	507	<i>Menyanthes.</i>
— <i>squarrosa.</i>	<i>Rattle-snake's-master.</i>	509	
<i>Ligusticum.</i>	<i>Lovage.</i>	193	<i>Miegia.</i>
<i>Ligustrum.</i>	<i>Privet. Prim.</i>	7	<i>Milium.</i>
			<i>Mimulus.</i>
			<i>Mitchella.</i>

374	Lilium.	Lily.	228
376	— carolinianum.	American Martagon.	229
	Linum.	Flax.	210
	Liquidambar.	Sweet-gum.	635
603	Liriodendron.	Tulip-tree. American Poplar.	382
117		Gromwell.	131
118	Lithospermum.	Cardinal's flower.	448
445	Lobelia Cardinalis.	Eye-bright.	448
380	— inflata.	American Mercury.	447
171	— siphilitica.	Bird's foot Trefoil	489
488	Lotus.	Lupine.	467
531	Lupinus.	Box-thorn.	97
145	Lycium.	Club-moss.	652
29	Lycopodium.	Ground Pine.	653
636	— dendroideum.	Walking-fern.	654
636	— alopecuroides.	Wild-Bugloss.	132
637	Lycopsis.	Water Horehound.	16
536	Lycopus.	Loose-strife.	135
538	Lysimachia.		
235			
546	Magnolia acuminata.	Cucumber-tree.	381
547	— auriculata.	Indian Physic.	382
580	— glauca.	Swamp Sassafras. Sweet Bay. Swamp Laurel. Beaver-wood.	381
595		Big Laurel.	380
596	— grandiflora.	Umbrella-tree. Elk-wood.	381
596	— tripetala.	Mallow.	454
00	Malva.	Horehound.	408
06	Marrubium.	Cuckold's Horns.	428
06	Martynia.	Medick. Nonsuch.	490
75	Medicago.	Indian Cucumber.	244
76	Medeola.	Cow-wheat.	430
76	Melampyrum.	Melic-grass.	77
76	Melica.	Melilot.	477
77	Melilotus.	Moon-seed.	370
80	Menispermum.	Sarsaparilla.	370
82	— canadense.	Mint.	405
7	Mentha.	Buck-bean. Marsh Tre- foil.	139
5	Menyanthes.	Canes.	59
7	Miegia.	Millet-grass.	62
9	Milium.	Monkey-flower.	426
3	Mimulus.	Eye-bright.	101
7	Mitchella.		

<i>Momordica.</i>	<i>Balsam apple.</i>	444	Passiflora.
<i>Monarda.</i>	<i>Mountain mint.</i>	16	Pastinaca.
— <i>Kalmiana.</i>	<i>Oswego Tea.</i>	17	Pediculari.
— <i>punctata.</i>	<i>Gravel wort.</i>	18	Peplis.
<i>Monotropa.</i>	<i>Yellow Bird's-nest.</i>	303	Phalaris.
<i>Morus.</i>	<i>Mulberry.</i>	639	Phaseolus.
<i>Myosotis.</i>	<i>Scorpion-grass.</i>	133	Phellandri.
<i>Myrica.</i>	<i>Candleberry Myrtle. Wax Myrtle.</i>	620	Philadelph.
— <i>Gale.</i>	<i>Sweet-gale.</i>	620	Phleum.
<i>Myriophyllum.</i>	<i>Water-milfoil.</i>	274	pr.
<i>Neottia.</i>	<i>Ladies' Traces.</i>	589	Physalis.
<i>Nepeta.</i>	<i>Cat-nep. Cat-mint.</i>	406	Phytolacca.
<i>Nicotiana.</i>	<i>Tobacco.</i>	141	Pinguicula.
<i>Nuphar.</i>	<i>Water Lily.</i>	368 369	Pinus.
<i>Nymphaea.</i>	<i>Tupelo-tree.</i>	177	Pisum.
— <i>candidans.</i>	<i>Ogochee Lime.</i>	177	Plantago.
— <i>villosa.</i>	<i>Sour Gum-tree.</i>	177	Platanus.
<i>Oenanthe.</i>	<i>Water drop-wort.</i>	194	Poa.
<i>Oenothera.</i>	<i>Tree Primrose.</i>	261	Podophyllum.
— <i>fruticosa.</i>	<i>Sun-drop.</i>	262	Polemoniace.
<i>Olea.</i>	<i>Olive tree. Devil's-wood.</i>	7	Polygala.
<i>Onoclea.</i>	<i>Sensitive-fern.</i>	665	sen.
<i>Ophioglossum.</i>	<i>Adder's-tongue.</i>	655	Polygonatum.
<i>Origanum.</i>	<i>Marjoram.</i>	411	Polygonum.
<i>Ornus.</i>	<i>Flowering-Ash.</i>	8	— avic.
<i>Orobanche.</i>	<i>Broom-rape.</i>	431	— bist.
— <i>virginiana.</i>	<i>Cancer-root.</i>	431	— Cor.
<i>Oryzopsis.</i>	<i>Mountain-rice.</i>	60	— Fag.
<i>Osmunda.</i>	<i>Flowering-fern.</i>	657	— Hyd.
<i>Ostrya.</i>	<i>Virginian Hornbeam. Iron-wood. Lever-wood.</i>	623	Populus.
<i>Oxalis.</i>	<i>Wood-sorrel.</i>	322	— bals.
<i>Oxycoccus.</i>	<i>Craneberry.</i>	263	— trep.
<i>Panax.</i>	<i>Ginseng.</i>	191	Porcelia.
<i>Panicum.</i>	<i>Panic-grass.</i>	66	Portulacca.
<i>Papaver.</i>	<i>Poppy.</i>	365	Potamogeton.
<i>Parietaria.</i>	<i>Pellitory.</i>	114	Potentilla.
<i>Parnassia.</i>	<i>Grass of Parnassus.</i>	208	Anemone.
			Pothos foetida.
			Prenanthes.
			Serp.

444	Passiflora.	Passion flower.	444
16	Pastinaca.	Parsnep.	196
17	Pedicularis.	Louse-wort.	424
18	Peplis.	Water-Purslane.	238
303	Phalaris.	Canary-grass.	65
639	Phaseolus.	Kidney-bean.	469
133	Phellandrium.	Water Hemlock.	195
620	Philadelphus.	Mock-Orange.	329
620	Phleum.	Cat's-tail-grass.	65
274	———— pratense.	Timothy-grass.	65
580	Physalis.	Ground Cherries.	157
406	Phytolacca.	Poke-root.	324
141	Pinguicula.	Butter-wort.	14
369	Pinus.	Pine. Fir tree. Hemlock.	
177	Pisum.	Spruce. Larch-tree.	639
177	Plantago.	Pea.	470
177	———— lanceolata.	Plantain.	98
194	Platanus.	Ribwort.	98
261	Poa.	Plane-tree. Sycamore.	
262	Podophyllum.	Button-wood.	635
7	Polemonium.	Meadow-grass.	78
265	Polygala.	Duck's-foot. May-apple.	366
55	———— seneca.	Greek Valerian. Jacob's	
11	Polygonatum.	Ladder.	151
8	Polygonum.	Milk-wort.	464
31	———— aviculare.	Seneca Snake-root.	464
31	———— bistortoides.	Solomon's Seal.	234
60	———— Convolvulus.	Persicaria.	269
57	———— Fagopyrum.	Knot-grass.	269
13	———— Hydropiper.	Snake-weed.	271
12	Populus.	Black Bind-weed.	273
13	———— balsamifera.	Buck-wheat.	273
12	———— trepida.	Water Pepper.	270
13	Porcelia.	Poplar.	618
11	Portulacca.	Tacamahac.	618
6	Potamogeton.	Asp or Aspen-tree.	618
5	Potentilla.	Papaw-tree.	383
4	———— Anserina.	Purslane.	365
8	Pothos foetida.	Pond-weed.	120
1	Prenanthes.	Cinquefoil.	358
6	———— Serpentaria.	Wild Tansey.	356
5		Scunk-weed. Scunk-cab-	
4		bage.	398
8		Mil-weed.	498
		Lion's-foot.	499

LATIN AND ENGLISH INDEX.

<i>Prinos.</i>	<i>Winter-berry.</i>	220	Sambuci
— <i>ambiguus.</i>	<i>Black Alder.</i>	220	Samolus.
— <i>glaber.</i>	<i>Ink-berry.</i>	220	
<i>Prunella.</i>	<i>Self-heal.</i>	411	Sanguina
<i>Prunus.</i>	<i>Plum-tree. Cherry-tree.</i>	329	Sanguisorba
<i>Pteris.</i>	<i>Brake.</i>	668	Santolina
<i>Pulmonaria.</i>	<i>Lung-wort.</i>	130	Sapindus
<i>Pycnanthemum.</i>	<i>Horse- or Mountain-mint.</i>	409	Saponaria
<i>Pyrola.</i>	<i>Winter-green.</i>	299	Sarracenia
<i>Pyrus.</i>	<i>Apple-tree,</i>	339	Saururus.
— <i>Botryapium.</i>	<i>May-bush.</i>	339	Saxifraga.
 <i>Quercus.</i>	 <i>Oak-tree.</i>	 625	n
 <i>Ranunculus.</i>	 <i>Crow-foot.</i>	 391	Schoenus.
<i>Rhamnus.</i>	<i>Buck-thorn.</i>	166	Schrankia
<i>Rhinanthus.</i>	<i>Yellow-rattle.</i>	429	Scirpus.
<i>Rhododendron.</i>	<i>Rose Bay. Mountain Laurel.</i>	297	Scleranthus
 <i>Rhus.</i>	 <i>Sumach.</i>	 204	Scolopendrium
— <i>Toxicodendron.</i>	<i>Poison Oak.</i>	205	Scorzonera
<i>Ribes.</i>	<i>Currant-tree. Gooseberry.</i>	163	Scutellaria
<i>Ricinus.</i>	<i>Palma-Christi. Castor-oil tree.</i>	602	Secale.
<i>Robinia.</i>	<i>False Acacia. Locust-tree.</i>	487	Sedum.
<i>Rosa.</i>	<i>Rose-Hep-tree.</i>	344	Selinum.
<i>Rottboellia.</i>	<i>Hard-grass.</i>	88	Senecio.
<i>Rubia.</i>	<i>Madder.</i>	102	
<i>Rubus.</i>	<i>Bramble. Raspberry.</i>	346	Seseli.
— <i>trivialis.</i>	<i>Dewberry.</i>	347	Silene.
— <i>Chamaemorus.</i>	<i>Cloud-berry.</i>	349	Sisou.
— <i>vilosus.</i>	<i>Black-berry.</i>	346	Sisymbrium
<i>Rumex.</i>	<i>Dock. Sorrel.</i>	247	am
<i>Ruppia.</i>	<i>Tassel Pond-weed.</i>	121	So
<i>Sabbatia.</i>	<i>Centaury.</i>	137	Sisyrinchium
<i>Saccharum.</i>	<i>Wild Sugar-cane.</i>	73	Sium.
<i>Sagina.</i>	<i>Pearl-wort.</i>	119	Smilax.
<i>Sagittaria.</i>	<i>Arrow-head. Arrow-root.</i>	395	Smyrnium.
<i>Salicornia.</i>	<i>Salt- or Glass-wort. Samphire.</i>	2	Solanum.
 <i>Salix.</i>	 <i>Willow. Osier.</i>	 608	Du
<i>Salsola.</i>	<i>Salt-wort.</i>	197	Solidago.
<i>Salvia.</i>	<i>Sage.</i>	19	Sorbus.
			Sorghum.
			Sparganium.
			Spergula.
			Spermacoce.
			Spigelia.

220	Sambucus.	Elder.	203
220	Samolus.	Brook-weed. Water Pim-	203
220		pernel.	158
411	Sanguinaria.	Blood-wort. Puccoon.	366
329	Sanguisorba.	Great Burnet.	116
668	Santolina.	Lavender Cotton.	520
130	Sapindus.	Soap-berry.	274
409	Saponaria.	Soap-wort.	314
299	Sarracenia.	Side-saddle flower.	367
339	Saururus.	Lizard's-tail.	252
339	Saxifraga.	Saxifrage.	310
	nivalis.	Sengreene.	310
625	Schoenus.	Bog Rush.	47
	Schrankia.	Sensitive Briar.	305
	Scirpus.	Club Rush.	54
	Scleranthus.	Knawel.	315
391	Scolopendrium.	Hart's-tongue.	667
166	Scorzonera.	Viper's-grass.	497
429	Scutellaria.	Scull-cap.	412
297	Secale.	Rye.	90
204	Sedum.	Stone-crop. Orpine.	323
105	Selinum.	Milk Parsley.	192
63	Senecio.	Groundsel. Ragwort.	528
	hieracifolius.	Butter-weed. Fire-weed.	529
02	Seseli.	Meadow Saxifrage.	197
87	Silene.	Catch-fly. Wild Pink.	315
44	Sison.	Hone wort.	194
88	Sisymbrium.	Water-cress.	440
02	amphibium.	Water-radish.	440
46	Sophia.	Flix-weed.	440
47	Sisyrinchium.	Blue-eyed Grass.	31
49	Sium.	Water Parsnep.	194
56	Smilax.	Green Briar. Poison Briar.	249
7	Smyrnium.	Alexanders.	196
1	Solanum.	Nightshade.	156
	Dulcamara.	Bittersweet.	156
7	Solidago.	Golden-rod.	535
3	Sorbus.	Service-tree. Mountain-	
9		Ash.	341
5	Sorghum.	Indian Millet. Broom grass.	78
2	Sparganium.	Bur-reed.	33
3	Spergula.	Spurrey.	320
7	Spermacoce.	Button-weed.	105
	Spigelia.	Worm-grass. Indian Pink-root.	139

LATIN AND ENGLISH INDEX.

Spiraea.	Meadow-sweet.	341
— opulifolia.	Virginian Guelder-rose.	342
— trifoliata.	Indian Physic. Ipecacuanha.	343
Stachys.	Hedge Nettle.	407
Staphylea.	Bladder-nut.	206
Statice.	Thrift. Sea Lavender.	212
Stellaria.	Stitch-wort.	317
— media.	Chick-weed.	317
Stillingia sebifera.	Tallow-tree.	608
Stipa.	Feather-grass.	72
Swertia.	Marsh Felwort.	100
Symporia.	St. Peter's-wort.	162
Tanacetum.	Tansey.	523
Taxus.	Yew-tree.	647
Tephrosia.	Goat's Rue.	489
Teucrium.	Germander.	405
Thalictrum.	Meadow Rue.	388
Thesium.	Bastard Toad-flax.	177
Thlaspi.	Bastard cress. Penny-cress.	
Thuya.	Shepherd's-purse.	435
Thymus.	American Arbor-vitæ.	646
Tilia.	Thyme.	413
 	Lime tree. Linden-tree.	
Tradescantia.	Bass wood. White-wood.	
Trientalis.	Spoon-wood.	362
Trifolium.	Spider wort.	218
Triglochin.	Winter green.	254
Triosteum.	Trefoil. Clover.	477
Triticum.	Arrow-grass.	247
Trollius.	Fever-wort. Gentian.	162
Turritis.	Wheat.	90
Tussilago.	Globe-flower.	391
Typha.	Tower-mustard.	438
Ulmus.	Colts'-foot.	531
Uniola.	Cat's-tail. Reed Mace.	34
Urtica.	Eelm.	199
Vaccinium.	Sea-side Oat.	82
— frondosum.	Nettle.	112
— stamineum.	Hawthorn-le- berry. Huckle- berry.	284
— Vitis Idæa.	Blue Tangles.	285
	Deer-berry.	284
	Bilberry.	289

341	Valeriana.	Valerian.	28
342	Veratrum.	Green Hellebore.	242
nha. 343	Verbascum.	Mullein. High-taper.	142
407	Verbena.	Vervain. Iron-weed.	415
206	Veronica.	Speedwell.	10
212	— serpyllifolia.	Paul's Betony.	11
317	— Beccabunga.	Brook-lime.	11
317	Viburnum dentatum.	Arrow-wood.	202
608	— lantanaoides.	Hobble-bush. Wayfaring-tree.	202
72	— laevigatum.	Cassicberry-bush.	203
100	— Oxycoccus.	Craneberry-tree. American Guelder-rose.	203
162	— prunifolium.	Sheep-berry.	202
523	Vicia.	Vetch.	471
647	Viola.	Violet. Heart's-ease.	171
489	Viscum.	Mistletoe.	114
405	Vitis.	Grape-vine.	169
388	Xanthium.	Small Burdock.	581
177	Yucca.	Adam's Needle. Cabbage-tree.	227
435	Zannichellia.	Horned Pond-weed.	4
646	Zanthorhiza.	Yellow-root.	212
413	Zanthoxylum.	Tooth-ach Tree.	209
362	Zea.	Maize. Indian Corn.	46
218	Zizania.	Canada or Indian Rice.	60
254	Zizyphus.	Supple-Jack.	188
477	Zostera.	Grass-wrack.	22
247			
162			
90			
391			
438			
531			
34			
199			
82			
112			
284			
285			
284			
289			

ENGLISH AND LATIN

INDEX.

A CACIA. False-	<i>Robinia.</i>	487
Adam's Needle.	<i>Yucca.</i>	227
Adder's Tongue.	<i>Ophioglossum.</i>	655
Agrimony.	<i>Agrimonia.</i>	335
—. Hemp-	<i>Eupatorium.</i>	511
Alder.	<i>Alnus.</i>	602
—. Black-	<i>Prinos.</i>	220
Alexanders.	<i>Smyrnium.</i>	196
Allspice. Carolina-	<i>Calycanthus.</i>	357
Aloe. Sweet Virginian-	<i>Agave.</i>	226
Alum-root.	<i>Geranium maculatum.</i>	448
Angelica-tree.	<i>Aralia spinosa.</i>	209
Aniseed-tree.	<i>Ilicium.</i>	380
Apple-tree.	<i>Pyrus.</i>	339
Arbor-vitæ.	<i>Thuya.</i>	646
Archangel.	<i>Lamium.</i>	406
Arrow-grass.	<i>Triglochin.</i>	247
— head.	{ <i>Sagittaria.</i>	395
— root.	<i>Viburnum dentatum.</i>	201
— wood.	<i>Asarum.</i>	596
Asarabæcca.	{ <i>Acer Negundo.</i>	265
Ash. Flowering-	<i>Ornus.</i>	8
—. Mountain-	<i>Sorbus.</i>	341
Ash-tree.	<i>Fraxinus.</i>	8
Asp. Aspen.	<i>Populus tremula.</i>	618
Avens.	<i>Geum.</i>	351
Balm of Gilead.	<i>Pinus Balsamea.</i>	639
Balsam.	<i>Impatiens.</i>	171
— Apple.	<i>Momordica.</i>	444

Banebe
 Barley.
 Bass-wc
 Bay-trec
 Bear-bir
 Bearberi
 Beaver-
 Beech-tr
 —. W
 Bell-flow
 Benjamini
 Bent-gra
 Berberry
 Bilberry.
 Bind-wet

—.
 Birch-tre
 Bird's-ne
 Birth-wo
 Bishop's-
 Bistort.
 Bitter-sw
 Blackberr
 Bladder-n
 Blite.
 —. St
 Blood-wo
 Bluebottle
 Blue-eyed
 Bluetangle
 Bog-rush.
 Boneset.
 Box Elder
 Box Thort
 Brake.
 Bramble.
 Breeches-fl
 Briar. Gre
 —. Poi
 —. Sen
 —. Swe
 —. Wil
 Brook-lime
 —. wee
 Brome-gras

487
227
655
335
511
602
220
196
357
226
448
209
380
339
646
406
247
395
201
590
265
8
341
8
618
351
639
171
444

Baneberry.	<i>Actaea.</i>	366
Barley.	<i>Hordeum.</i>	89
Bass-wood.	<i>Tilia.</i>	302
Bay-tree.	<i>Laurus.</i>	275
Bear-bind.	<i>Calystegia Sepium.</i>	142
Bearberry.	<i>Arbutus Uva ursi.</i>	283
Beaver-wood.	<i>Magnolia glauca.</i>	381
Beech-tree.	<i>Fagus.</i>	624
—. Water-	<i>Platanus.</i>	634
Bell-flower.	<i>Campanula.</i>	159
Benjamin-tree.	<i>Laurus Benzoin.</i>	276
Bent-grass.	<i>Agrostis.</i>	63
Berberry.	<i>Berberis.</i>	219
Bilberry.	<i>Vaccinium Vitis Idaea.</i>	289
Bind-weed.	<i>Convolvulus.</i>	143
—. Black-	<i>Polygonum Convolvulus.</i>	273
Birch-tree.	<i>Betula.</i>	620
Bird's-nest. Yellow-	<i>Monotropa.</i>	303
Birth-wort.	<i>Aristolochia.</i>	596
Bishop's-weed.	<i>Ammi.</i>	192
Bistort.	<i>Polygonum Bistorta.</i>	369
Bitter-sweet.	<i>Solanum Dulcamara.</i>	156
Blackberry.	<i>Rubus villosus.</i>	346
Bladder-nut.	<i>Staphylea.</i>	206
Blite.	<i>Amaranthus Blitum.</i>	207
—. Strawberry.	<i>Blitum.</i>	4
Blood-wort.	<i>Sanguinaria.</i>	366
Bluebottle.	<i>Centaurea Cyanus.</i>	576
Blue-eyed Grass.	<i>Sisyrinchium.</i>	31
Bluetangles.	<i>Vaccinium frondosum.</i>	285
Bog-rush.	<i>Schoenus.</i>	47
Boneset.	<i>Eupatorium perfoliatum.</i>	516
Box Elder.	<i>Acer Negundo.</i>	268
Box Thorn.	<i>Lycium.</i>	97
Brake.	<i>Pteris.</i>	668
Bramble.	<i>Rubus.</i>	346
Breeches-flower.	<i>Corydalis.</i>	462
Briar. Green-	{ <i>Smilax.</i>	249
—. Poison-	{ <i>Schrankia.</i>	305
—. Sensitive-	{ <i>Rosa.</i>	344
—. Sweet-	<i>Veronica Beccabunga.</i>	10
—. Wild-	<i>Samolus.</i>	158
Brook-lime.	{ <i>Bromus.</i>	85
—. weed.	{ <i>Sorghum.</i>	78
Brome-grass.		

Broom-rape.	<i>Orobanche.</i>	431	Cedar.
Buck-bean.	<i>Menyanthes.</i>	139	Celandin
Buck's eye-tree.	<i>Æculus.</i>	254	Centaury
— thorn.	<i>Rhamnus.</i>	166	
— Sea-	<i>Hippophaë.</i>	115	
— Wheat.	<i>Polygonum Fagopyrum.</i>	273	
Bugle.	<i>Ajuga.</i>	404	Chaff-we
Bugloss. Wild-	<i>Lycopsis.</i>	133	Chamom
Bug-wort.	<i>Cimicifuga.</i>	372	Cherry-tr
Bull-rush.	<i>Scirpus lacustris.</i>	54	Chervil.
Burdock.	<i>Arctium.</i>	505	Chesnut-
— Small-	<i>Xanthium.</i>	581	
Bur-grass.	<i>Cenchrus.</i>	60	Chickwee
— reed.	<i>Sparagnum.</i>	33	Chicot.
Eu-net. Great-	<i>Sanguisorba.</i>	116	Chinquap
Burning-bush.	<i>Erythronium americanum.</i>	168	Cinquefoi
Butterfly-flower.	<i>Asclepias tuberosa.</i>	183	
Butter-nut.	<i>Juglans cinerea.</i>	636	Cleavers.
— weed.	<i>Senecio hieracifolius.</i>	529	Clot-bur.
— wort.	<i>Pinguicula.</i>	14	Cloud-ber
Button-weed.	<i>Spermacoce.</i>	105	Clover.
— wood.	<i>Platanus.</i>	635	Club-moss
	<i>Cephalanthus.</i>	114	Club-rush
Cabbage.	<i>Brassica.</i>	438	Cockle.
— tree.	<i>Chamærops. Yucca.</i>	239	Cock's-foo
Calico-tree.	<i>Kalmia latifolia.</i>	296	Coffee tree
Calf's-snout.	<i>Antirrhinum Orontium.</i>	422	Cohosh.
Campion.	<i>Cucubalus.</i>	315	— R
Canary-grass.	<i>Phalaris.</i>	65	Colt's-foot.
Cancer-root.	<i>Orobanche virginica.</i>	431	Columbine
Candleberry Myrtle.	<i>Myrica.</i>	620	Coral-plant
Cane.	<i>Melia.</i>	59	Cotton-gra
Cardinal's-flower.	<i>Lobelia Cardinalis.</i>	448	— tre
Carpenter's-leaf.	<i>Galax.</i>	446	Couch-gras
Carrot.	<i>Daucus.</i>	191	Cowbane.
Cassava. Wild-	<i>Jatropha.</i>	603	Cowslip.
Cassioberry-bush.	<i>Viburnum laevigatum.</i>	201	Crakeberry.
Castor-oil tree.	<i>Ricinus.</i>	602	Craeberry.
Catalpa-tree.	<i>Catalpa.</i>	10	— t
Catchfly.	<i>Silene.</i>	315	Crane's-bill
— bush.	<i>Bejaria.</i>	362	Creeper.
Catmint.	<i>Nepeta.</i>	406	V
Cat's-tail.	<i>Typha.</i>	34	Cress.
— Grass.	<i>Phleum.</i>	65	Bast;
Cedar. Red-	<i>Juniperus virginiana.</i>	646	— Swir
			— Wal
			— Wat
			Crow-berry.

431	Cedar, White-	<i>Cupressus thyoides.</i>	646
139	Celandine.	<i>Chelidonium.</i>	365
254		{ <i>Centaurea.</i>	576
166	Centaury.	<i>Sabatia.</i>	137
115		<i>Centunculus.</i>	97
273	Chaff-weed.	<i>Anthemis.</i>	502
404	Chamomile.	<i>Prunus.</i>	329
133	Cherry-tree.	<i>Chærophyllum.</i>	195
372	Chervil.	<i>Castanea.</i>	624
54	Chesnut-tree..	<i>Æsculus.</i>	254
505		<i>Stellaria media.</i>	617
581	Horse-	<i>Gymnocladus.</i>	304
60	Chickweed.	<i>Castanea pumila.</i>	625
33	Chicot.	<i>Potentilla.</i>	353
116	Chinquapin.	<i>Comarum.</i>	356
168	Cinquefoil.	<i>Galium.</i>	102
183	— — —. Marsh-	<i>Arctium.</i>	505
636	Cleavers.	<i>Rubus Chamæmorus.</i>	345
529	Clot-bur.	<i>Trifolium.</i>	477
14	Cloud-berry.	<i>Lycopodium.</i>	652
105	Clover.	<i>Scirpus.</i>	54
635	Club-moss.	<i>Agrostemma.</i>	321
114	Club-rush.	<i>Dactylis.</i>	84
438	Cockle. Corn-	<i>Gymnocladus.</i>	304
239	Cock's-foot Grass.	<i>Caulophyllum.</i>	218
296	Coffee tree. Kentucky-	<i>Actaea.</i>	366
422	Cohosh. Blue-	<i>Tussilago.</i>	531
315	— — —. Red and White-	{i} <i>Asarum.</i>	596
65	Colt's-foot.	<i>Aquilegia.</i>	372
431	Columbine.	<i>Erythrina.</i>	467
620	Coral-plant.	<i>Eriophorum.</i>	57
59	Cotton-grass.	<i>Platanus.</i>	635
448	— — — tree.	{i} <i>Populus.</i>	613
446	Couch-grass.	<i>Chloris.</i>	87
191	Cowbane. Water-	<i>Cicuta.</i>	197
603	Cowslip. American-	<i>Dodecatheon.</i>	136
201	Crakeberry.	<i>Empetrum.</i>	94
602	Craneberry.	<i>Oxycoccus.</i>	263
10	— — — tree.	<i>Viburnum Oxyccos.</i>	203
315	Crane's-bill.	<i>Geranium.</i>	448
362	Creeper. Virginian-	<i>Cissus hederacea.</i>	170
400	Cress. Bastard-	<i>Thlaspi.</i>	435
34	— — —. Swine-	<i>Coronopus.</i>	435
65	— — —. Wall-	<i>Aralis.</i>	436
646	— — —. Water-	<i>Sisymbrium.</i>	440
	Crow-berry.	<i>Empetrum.</i>	23

Crow-foot.	<i>Ranunculus.</i>	391	Fescue.
Cuckold's-nut.	<i>Corylus rostrata.</i>	635	Fever-w.
— horn.	<i>Martynia.</i>	428	Filbert.
Cucumber-tree,	<i>Magnolia acuminata.</i>	381	Fire-we.
Cudweed.	<i>Gnaphalium.</i>	524	Fir-tree.
Currant-tree.	<i>Ribes.</i>	163	Flag.
Cypress.	<i>Cupressus.</i>	645	—.
— Vine.	<i>Ipomoea Quamoclit.</i>	145	Flax.
Dahoon.	<i>Ilex.</i>	117	Flea-ban.
Daisy.	<i>Bellis.</i>	526	Flix-w.
—. Ox-eye-	{ <i>Chrysanthemum Leucanth.</i>	526	Flower-
Dandelion.	{ <i>Bupthalmum.</i>	563	Flowerin.
Date-plum.	<i>Leontodon.</i>	497	Fluillin.
Deerberry.	<i>Diospyros.</i>	265	Flux-wet.
Devil's-bit.	<i>Vaccinium stamineum.</i>	284	Fly-catch.
— wood.	<i>Aletris.</i>	225	Fox-tail-.
Dewberry.	<i>Olea.</i>	7	Fringe-tr.
Dittany.	<i>Rubus trivialis.</i>	347	Full-fern.
Dock.	<i>Cunila.</i>	406	Fumatory.
Dodder.	<i>Rumex.</i>	247	Fuller's T.
Dog's-bane.	<i>Cuscuta.</i>	116	Garlic.
Dog's-foot Violet.	<i>Apocynum.</i>	179	Gentian.
Dog wood.	<i>Erythronium.</i>	239	Germand.
Dragon's head.	<i>Cornus.</i>	107	Ginger.
Drop-wort. Water-	<i>Dracoccephalum.</i>	411	Ginseng.
Duck's-foot.	<i>Oenanthe.</i>	194	Glass-wor.
— mead.	<i>Podophyllum.</i>	366	Globe-flov.
Elder.	<i>Lemna.</i>	22	Goat's Ru.
Elecampane.	<i>Sambucus.</i>	203	Golden-ro.
Elephant's-foot.	<i>Inula Helenium.</i>	531	— — pe.
Elk-wood.	<i>Elephantopus.</i>	582	Goldilocks.
Elm-tree.	<i>Magnolia tripetala.</i>	381	Gooseberry.
—. Witch-	<i>Ulmus.</i>	199	Goose-foot.
Eringo.	<i>Planera.</i>	115	— — gra.
Everlasting.	<i>Eryngium.</i>	189	Grape-vine.
Eye-bright.	<i>Gnaphalium.</i>	524	Grass of P.
Featherfoil.	{ <i>Euphrasia.</i>	430	— Wrac.
— beds.	{ <i>Mitchella.</i>	101	Gravel-wor.
— grass.	{ <i>Lokelia inflata.</i>	448	Green Dra.
Felwort. Marsh-	<i>Hottonia.</i>	138	Gromwel.
Fennel, Dog's-	<i>Chara.</i>	4	Ground Ivy.
	<i>Stipa.</i>	72	
	<i>Swertia.</i>	100	
	<i>Eupatorium fœniculaceum.</i>	512	Ground Pir.

391	Fescue-grass.	<i>Festuca.</i>	83
635	Fever-wort.	<i>Triosteum.</i>	162
428	Filbert. Wild-	<i>Corylus.</i>	634
381	Fire-weed.	<i>Senecio hieracifolius.</i>	521
524	Fir-tree.	<i>Pinus.</i>	669
163	Flag.	<i>Iris.</i>	29
645	—. Sweet-	<i>Acorus.</i>	235
145	Flax.	<i>Linum.</i>	210
117	Flea-bane.	<i>Conyza.</i>	523
526	Flix-weed.	<i>Sisymbrium Sophia.</i>	440
526	Flower-de-luce.	<i>Iris.</i>	29
503	Flowering-fern.	<i>Osmunda.</i>	657
497	Fluellin.	<i>Antirrhinum Elatine.</i>	421
265	Flux-weed.	<i>Geranium maculatum.</i>	440
284	Fly-catch-bush.	<i>Bejaria.</i>	362
225	Fox-tail-grass.	<i>Alopecurus.</i>	65
7	Fringe-tree.	<i>Chionanthus.</i>	7
347	Full-fern.	<i>Acrostichum.</i>	658
406	Fumatory.	<i>Funaria. Corydalis.</i>	463
247	Fuller's Thistle.	<i>Dipsacus.</i>	96
116	Garlic. Wild-	<i>Allium.</i>	222
179	Gentian.	{ <i>Gentiana.</i>	185
239		{ <i>Tristeum.</i>	162
107	Germanander.	<i>Teucrium.</i>	405
411	Ginger. Wild-	<i>Asarum.</i>	596
194	Ginseng.	<i>Panax.</i>	191
366	Glass-wort.	<i>Salicornia.</i>	2
22	Globe-flower.	<i>Trollius.</i>	391
203	Goat's Rue.	<i>Tephrosia.</i>	489
531	Golden-rod.	<i>Solidago.</i>	535
582	— pert.	<i>Gratiola aurea.</i>	92
381	Goldilocks.	{ <i>Chrysocoma.</i>	517
199	Gooseberry.	{ <i>Ranunculus Auricomus.</i>	391
115	Goose-foot.	<i>Riles.</i>	163
189	— grass.	<i>Chenopodium.</i>	197
524	Grape-vine.	<i>Galium.</i>	102
430	Grass of Parnassus.	<i>Vitis.</i>	169
101	— Wrack.	<i>Parnassia.</i>	208
448	Gravel-wort.	<i>Zostera.</i>	2
138	Green Dragon.	<i>Monarda punctata.</i>	18
4	Gromwel.	<i>Arum Dracunculus.</i>	400
72	Ground Ivy.	<i>Lithospermum.</i>	131
100	Ground Pine.	{ <i>Glechoma.</i>	408
512		{ <i>Epigaea.</i>	297
		{ <i>Lycopodium dendroideum.</i>	653
		{ <i>Ajuga Chamæpithys.</i>	404

Ground Cherry.	<i>Physalis.</i>	157	Hop.
Groundsel.	<i>Senecio.</i>	528	— He
— tree.	<i>Baccharis.</i>	523	Horehou
Guelder-rose, Ameri- can.	{ <i>Viburnum.</i> <i>Spiraea opulifolia.</i>	201 342	Hornbea
Gum-tree. Sour- — Sweet.	<i>Nyssa villosa.</i> <i>Liquidambar.</i>	177 635	Hornwor
Hag-berry.	<i>Celtis.</i>	200	Horse-ta
Hackmatack.	<i>Pinus pendula.</i>	639	Horse-w
Hair-grass.	<i>Aira.</i>	76	Hound's-
Hard-grass.	<i>Rottboellia.</i>	88	Hucklebe
Hart's-tongue.	<i>Scolopendrium.</i>	667	Hyssop.
Hawkweed.	<i>Hieracium.</i>	502	Jacob's L.
Hawthorn.	<i>Crataegus.</i>	376	Jalap.
Hazel-nut.	<i>Corylus.</i>	634	Jasmine.
Heal-all.	{ <i>Galax.</i> <i>Viola clandestina.</i>	446 171	Jesuit's B.
Heart-seed.	<i>Cardiospermum.</i>	273	Indian Ce
Heart's-ease.	<i>Viola.</i>	171	— Cu
Heath. Yellow-	{ <i>Hudsonia.</i> <i>Hypericum densiflorum.</i>	364 376	— Fig
Heliotrope.	<i>Heliotropium.</i>	130	— Mi
Hellebore. Green-	<i>Veratrum viride.</i>	242	— Ph
Hemlock.	<i>Conium.</i>	195	— Ric
— Spruce-	<i>Pinus canadensis.</i>	639	— Rec
— Water-	<i>Phellandrium.</i>	195	Indigo.
Hemp.	<i>Cannabis.</i>	199	Ink-berry.
— Indian-	<i>Apocynum.</i>	179	Ipecacuanl
— River-	<i>Acnida.</i>	208	Iron-weed.
Henbane.	<i>Hyoscyamus.</i>	141	— wood
Henbit.	<i>Lamium amplexicaule.</i>	406	Judas-tree.
Heptree.	<i>Rosa.</i>	344	Juniper.
Herb Pennet.	<i>Geum.</i>	351	Kidney Be
— Christopher.	<i>Actaea.</i>	366	Knawel.
— Robert.	<i>Geranium Robertianum.</i>	449	Knot-grass.
High-taper.	<i>Verbascum.</i>	142	Ladies' Bed
Hickory or Hickery.	<i>Juglans.</i>	636	— Mai
Hobble-bush.	<i>Viburnum lantanaoides.</i>	201	
Hog-nut.	<i>Juglans porcina.</i>	638	
— weed.	<i>Ambrosia.</i>	580	
Holly.	<i>Ilex.</i>	117	
Hone-wort.	<i>Sison.</i>	194	
Honey-suckle.	{ <i>Caprifolium.</i> <i>Aquilegia.</i>	160 372	
— Upright-	<i>Azalea.</i>	151	
Hoop-Ash.	<i>Celtis.</i>	200	

157	Hop.	<i>Humulus.</i>	199
528	— Hornbeam.	<i>Ostrya.</i>	623
523	Horehound.	<i>Marrubium.</i>	408
201	— Water-	<i>Lycopus.</i>	16
342	Hornbeam-tree.	<i>Carpinus.</i>	623
177	Hornwort.	<i>Ceratophyllum.</i>	370
635	Horse-tail.	<i>Equisetum.</i>	651
	Horse-weed.	<i>Collinsonia.</i>	20
200	Hound's-tongue.	<i>Cynoglossum.</i>	133
639	Huckleberry.	<i>Vaccinium.</i>	284
76	— Bastard-	<i>Andromeda.</i>	290
88	Hyssop.	<i>Hyssopus.</i>	405
667	— Hedge-	<i>Gratiola.</i>	12
502	Jacob's Ladder.	<i>Polemonium.</i>	151
376	Jalap.	{ <i>Ipomoea Jalapa.</i> <i>Convolvulus panduratus.</i>	147 143
634	Jasmine. Carolina-	<i>Gelsemium.</i>	184
446	Jesuit's Bark-tree.	Bastard- <i>Iva.</i>	580
171	Indian Corp.	<i>Zea.</i>	46
273	— Cucumber.	<i>Medeola.</i>	244
171	— Fig.	<i>Cactus.</i>	327
364	— Millet.	<i>Sorghum.</i>	78
376	— Physic.	{ <i>Magnolia auriculata.</i> <i>Spiraea trifoliata.</i>	382 341
130	— Rice.	<i>Zizania.</i>	60
242	— Reed or Shot.	<i>Canna.</i>	585
195	Indigo.	<i>Indigofera.</i>	448
639	— Bastard-	<i>Amorpha.</i>	466
195	Ink-berry.	<i>Prinos glaber.</i>	220
199	Ipecacuanha.	{ <i>Asclepias tuberosa.</i> <i>Euphorbia.</i>	183 605
179	Iron-weed.	{ <i>Spiraea trifoliata.</i> <i>Ambrosia.</i>	341 580
208	— wood.	<i>Verbena.</i>	415
141	Judas-tree.	<i>Ostrya.</i>	623
406	Juniper.	<i>Cercis.</i>	308
344	Kidney Bean.	{ <i>Juniperus.</i> <i>Cupressus.</i>	646 645
351	Knawel.	<i>Phaseolus.</i>	469
366	Knot-grass.	<i>Scleranthus.</i>	315
449	Ladies' Bedstraw.	{ <i>Illecebrum.</i> <i>Polygonum aviculare.</i>	445 359
142	— Mantle.	<i>Galium.</i>	102
636		<i>Alchemilla.</i>	112
201			
638			
580			
117			
194			
160			
172			
51			
300			

Ladies' Slipper.	<i>Cypripedium.</i>	594	Martago
— Smock.	<i>Cardamine.</i>	439	Marygol
— Traces.	<i>Nicotia.</i>	589	
Lamb's Quarters.	<i>Chenopodium viride.</i>	198	
Larch-tree.	<i>Pinus pendula.</i>	639	
Tarkspur.	<i>Delphinium.</i>	371	
Laurel.	<i>Laurus.</i>	275	
— Big-	<i>Magnolia grandiflora.</i>	380	
— Mountain-	<i>Rhododendron.</i>	297	
— Sheep-	<i>Kalmia angustifolia.</i>	296	
— Small-	— <i>latifolia.</i>		
— Swamp-	<i>Magnolia glauca.</i>	381	
Lavender Cotton.	<i>Santolina.</i>	520	
— Sea-	<i>Statice.</i>	212	
Leather-wood.	<i>Divca.</i>	268	
Lettuce.	<i>Lactuca.</i>	500	
Lever-wood.	<i>Ostrya.</i>	623	
Lily.	<i>Lilium.</i>	221	
— Water-	{ <i>Nymphaea.</i>	308	
— of the Valley.	{ <i>Nuphar.</i>	369	
— Atamasco-	<i>Convallaria.</i>	232	
Lime or Linden-tree.	<i>Amaryllis.</i>	222	
Lion's-foot.	<i>Tilia.</i>	362	
Liquorice.	<i>Prenanthes.</i>	449	
Liver-wort.	<i>Glycyrrhiza.</i>	480	
Lizard's-tail.	<i>Hepatica.</i>	391	
Loblolly-bay.	<i>Laururus.</i>	252	
Locust-tree.	<i>Gordonia.</i>	451	
— Honey-	<i>Robinia.</i>	487	
Loose-strife.	<i>Gleditschia.</i>	221	
Lovage.	<i>Lysimachia.</i>	135	
Louse-wort.	<i>Ligusticum.</i>	193	
Lung-wort.	<i>Pedicularis.</i>	424	
Lupine.	<i>Pulmonaria.</i>	130	
Lyme-grass.	<i>Lupinus.</i>	487	
	<i>Eigenus.</i>	89	
Madder.	<i>Rulia.</i>	102	
Madwort.	<i>Alyssum.</i>	434	
Mahogany. Mountain-	<i>Betula lenta.</i>	621	Myrtle.
Maiden-hair.	<i>Adiantum.</i>	670	
Mallow.	<i>Malva.</i>	454	
Man's-beard.	<i>Andropogon.</i>	74	
Maple-tree.	<i>Acer.</i>	265	Nettle.
Mare's-tail.	<i>Hippuris.</i>	3	
Majoram.	<i>Origanum.</i>	411	H
			Nettle-tree

594	Martagon.	<i>Lilium.</i>	229
439	Marygold. Bur-	<i>Bidens.</i>	566
589	_____. Marsh-	<i>Caltha.</i>	389
198	May-apple.	<i>Podophyllum.</i>	366
639	_____. bush.	<i>Pyrus Botryapium.</i>	339
371	_____. weed.	<i>Anthemis.</i>	562
275	Meadow-grass.	<i>Poa.</i>	72
380	_____. rue.	<i>Thalictrum.</i>	388
297	_____. sweet.	<i>Spiraea.</i>	388
296	Medick.	<i>Medicago.</i>	490
381	Melic-grass.	<i>Melica.</i>	77
520	Melilot.	<i>Melilotus.</i>	477
212	Mercury. American-	<i>Lobelia siphilitica.</i>	447
268	Milfoil.	<i>Achillea.</i>	562
500	_____. Hooded-	<i>Utricularia.</i>	15
623	_____. Water-	<i>Myriophyllum.</i>	274
221	Milk-weed.	<i>Prenanthes.</i>	498
308	_____. wort.	<i>Polygala.</i>	464
309	Millet-grass.	<i>Milium.</i>	62
232	Mint.	<i>Mentha.</i>	405
222	_____. Horse or Moun- tain-	{ <i>Monarda.</i> <i>Pycnanthemum.</i>	16 409
362	Mitre-flower.	<i>Dodecatheon.</i>	136
449	Mistle-toe.	<i>Viscum.</i>	114
480	Moccasin-flower.	<i>Cypripedium.</i>	594
391	Mocker-nut.	<i>Juglans.</i>	636
252	Mock Orange.	<i>Philadelphus.</i>	329
451	Monk's-hood.	<i>Aconitum.</i>	372
487	Monkey-flower.	<i>Mimulus.</i>	426
221	Moon-seed.	<i>Menispermum.</i>	370
135	Moose-wood.	<i>Acer striatum.</i>	267
193	Mouse-ear Chickweed.	<i>Cerastium.</i>	320
424	Mugwort.	<i>Artemisia vulgaris.</i>	322
130	Mulberry.	<i>Morus.</i>	639
187	Mullein.	<i>Verbascum.</i>	142
89	Mustard. Hedge-	<i>Erysimum.</i>	436
02	_____. Indian-	<i>Dentaria.</i>	438
34	_____. Tower-	<i>Turritis.</i>	438
21	Myrtle. Candleberry-	{ <i>Myrica.</i> <i>Ammyrsine.</i>	620 301
70	_____. Wax-		
54	_____. Sand- .		
74	Nettle.	<i>Urtica.</i>	112
65	_____. Dead-	<i>Galeopsis.</i>	407
3	_____. Hedge-	<i>Stachys.</i>	407
11	Nettle-tree.	<i>Celtis.</i>	200

Nightshade.	<i>Solanum.</i>	156	Pig-nut.
—. Enchanter's-	<i>Ciræa.</i>	21	Pimpern.
Nonsuch.	<i>Medicago Lupulina.</i>	490	
Nut-grass.	<i>Cyperus.</i>	50	
Oak.	<i>Quercus.</i>	625	Pine-tree
Oat. Sea-side-	<i>Uniola.</i>	83	Pink.
Oat-grass.	<i>Avena.</i>	85	Pink-roo
Ogeechee-lime.	<i>Nyssa candidans.</i>	177	—. W
Oil-nut.	{ <i>Hamiltonia.</i>	178	Pipe-wor
	<i>Juglans cinerea.</i>	636	Piperidge
Okro.	<i>Hibiscus esculentus.</i>	456	Plane-tre
Olive-tree.	<i>Olea.</i>	7	Plantain.
Orache.	<i>Atriplex.</i>	199	—.
Orchard-grass.	<i>Dactylis.</i>	84	Plum-tree
Orpine.	<i>Sedum.</i>	323	Poison El
Osier.	<i>Salix.</i>	608	—. Oa
—. Red-	<i>Cornus.</i>	109	Poke-root
Palmetto-tree.	<i>Chamærops.</i>	299	Pond-wee
Palma Christi.	<i>Ricinus.</i>	602	—.
Panic-grass.	<i>Panicum.</i>	60	Poplar-tre
Pansy.	<i>Viola tricolor.</i>	171	—. W
Papaw-tree.	<i>Porcelia.</i>	383	Poppy.
Parsley. Cow-	<i>Chœrophyllum.</i>	195	Prickly-pe
—. Milk-	<i>Selinum.</i>	192	Prim.
—. piert.	<i>Alchemilla arvensis.</i>	112	Prince's Fe
Parsnep.	<i>Pastinaca.</i>	196	Privet.
—. Cow-	<i>Heracleum.</i>	192	Puccoon.
—. Water-	<i>Sium.</i>	194	Purslane.
Partridge-berry.	<i>Gaultheria.</i>	283	—.
Passion-flower.	<i>Passiflora.</i>	444	Quaking G
Paul's Betony.	<i>Veronica serpyllifolia.</i>	10	Quamash.
Pea.	<i>Pisum.</i>	470	Rag-weed.
Pearl-wort.	<i>Sagina.</i>	119	— wort.
Pear.	<i>Pyrus.</i>	339	Raspberry.
Pecan-nut.	<i>Juglans.</i>	636	Rattle-pea.
Pellitory.	<i>Parietaria.</i>	114	Rattle-snak
Penny-cress.	<i>Thlaspi-arvense.</i>	435	
— royal.	<i>Hedera.</i>	414	
— wort.	<i>Hydrocotyle.</i>	190	
Pepper-wort.	<i>Lepidium.</i>	435	
— root.	<i>Dentaria.</i>	438	
— vine.	<i>Cissus bipinnata.</i>	170	
Persicaria.	<i>Polygonum.</i>	269	
Persimon.	<i>Diospyros.</i>	265	Red-root.

156	Pig-nut.	<i>Juglans porcina.</i>	638
21	Pimpernel.	<i>Anagallis.</i>	135
490	—. Water-	<i>Samolus.</i>	158
50	Pine-tree.	<i>Pinus.</i>	639
625	Pink.	<i>Dianthus.</i>	314
82	Pink-root.	<i>Spigelia.</i>	139
85	—. Wild-	<i>Silene.</i>	315
177	Pipe-wort.	<i>Eriocaulon.</i>	91
178	Piperidge-bush.	<i>Berberis.</i>	219
636	Plane-tree.	<i>Platanus.</i>	635
456	Plantain.	<i>Plantago.</i>	98
7	—. Rattle-snake-	<i>Erigeron bellidifolium.</i>	532
199	—. Water-	<i>Alisma.</i>	352
84	Plum-tree.	<i>Prunus.</i>	329
323	Poison Elder.	<i>Aralia hispida.</i>	209
608	—. Oak.	<i>Rhus Toxicodendron.</i>	205
109	Poke-root.	<i>Phytolacca.</i>	324
299	Pond-weed.	<i>Potamogeton.</i>	120
602	—. Horned-	<i>Zannichellia.</i>	4
60	—. Tassel-	<i>Ruppia.</i>	121
171	Poplar-tree.	<i>Populus.</i>	618
383	—. White-	<i>Liriodendron.</i>	382
195	Poppy.	<i>Papaver.</i>	305
192	Prickly pear.	<i>Cactus.</i>	327
112	Prim.	<i>Ligustrum.</i>	7
196	Prince's Feather.	<i>Amaranthus.</i>	207
192	Privet.	<i>Ligustrum.</i>	7
194	Puccoon.	{ <i>Sanguisorba.</i>	116
83	Purslane.	{ <i>Bartschia.</i>	132
44	—. Water-	{ <i>Portulaca.</i>	365
10	Quaking Grass.	<i>Peplis.</i>	238
70	Quamash.	<i>Briiza.</i>	82
19		<i>Phalangium.</i>	226
39	Rag-weed.	<i>Erigeron.</i>	532
36	— wort.	<i>Senecio.</i>	528
14	Raspberry.	<i>Rubus.</i>	340
35	Rattle-pea.	<i>Crotalaria.</i>	469
14	Rattle-snake's-master.	<i>Liatris squarrosa.</i>	509
30	— plantain.	<i>Erigeron bellidifolium.</i>	532
35	— root.	<i>Polygonata.</i>	464
38	— weed.	{ <i>Sonchus floridanus.</i>	501
70		{ <i>Prenanthes.</i>	408
19	Red-root.	{ <i>Ceanothus.</i>	167
15		{ <i>Leucisia.</i>	368

Reed-grass.	<i>Arundo.</i>	86	Seneca C
— mace.	<i>Typha.</i>	34	Sengreen.
Rib-fern.	<i>Blechnum.</i>	669	Service-tr
— wort.	<i>Plantago lanceolata.</i>	98	Shave-gra
Rice. Canada-	<i>Zizania.</i>	60	Shell- or
— Mountain-	<i>Oryzopsis.</i>	69	Sheep ber
Rock Rose.	<i>Helianthemum.</i>	303	Shepherd'
Rocket.	<i>Hesperis.</i>	436	Shield-fer
— Sea-	<i>Cakile.</i>	434	Side-saddl
Rose.	<i>Rosa.</i>	344	Silver-tree
Rose Bay.	<i>Rhododendron.</i>	297	— bel
Rosemary. Wild-	<i>Andromeda polifolia.</i>	291	Snake-roo
Rush.	<i>Juncus.</i>	235	—
Rye.	<i>Secale.</i>	90	—
— Wild-	<i>Elymus.</i>	89	—
Saca-commis.	<i>Arbutus Uva ursi.</i>	283	—
Sage.	<i>Salvia.</i>	19	weed
St. Andrew's Cross.	<i>Acyrum.</i>	373	Snap-drag
St. Barnaby's Thistle.	<i>Centaurea solstitialis.</i>	576	Sneeze-wc
St. John's-wort.	<i>Hypericum.</i>	374	Snow-berr
St. Peter's-wort.	<i>Symporia.</i>	162	Snow-drop
Salt-wort.	{ <i>Salicornia.</i>	2	— flock
— Black-	<i>Salsola.</i>	197	— flower
Samphire.	<i>Glaux.</i>	176	Soap-berry
Sand-wort.	<i>Salicornia.</i>	2	— wort
Sarsaparilla.	<i>Arenaria.</i>	317	Solomon's
Sassafras-tree.	{ <i>Aralia nudicaulis.</i>	209	Sorrel.
— Swamp-	<i>Smilax.</i>	249	— tree
Savin.	<i>Menispermum.</i>	370	Spear-wort
Saxifrage.	<i>Laurus Sassafras.</i>	277	Speedwell.
— Golden-	<i>Magnolia glauca.</i>	381	Spice-wood
— Meadow-	<i>Juniperus Salina.</i>	646	Spider-wor
Scorpion-grass.	<i>Saxifraga.</i>	310	Spikenard.
Scull-cap.	<i>Chrysosplenium.</i>	269	Spindle tre
Scunk-weed.	<i>Seseli.</i>	197	Spleen-wor
— cabbage.	<i>Myosotis.</i>	133	Spoon-woo
Sea-mat-grass.	<i>Scutellaria.</i>	412	Spring-grass
Self-heal.	{ Pothos.	398	Spruce-pine
Sensitive-fern.	<i>Arundo arenaria.</i>	87	Spurrey.
— pea.	<i>Prunella.</i>	411	Staff-tree.
— briar.	<i>Onoclea.</i>	665	Star-wort.
Sedge.	<i>Cassia nicticans.</i>	305	
	<i>Schrankia.</i>	365	
	{ <i>Carex.</i>	34	
	<i>Schœnus</i>	47	

86	Seneca Grass.	<i>Holcus fragrans.</i>	78
34	Sengreen.	<i>Saxifraga nivalis.</i>	310
669	Service-tree.	{ <i>Sorbus.</i>	341
98		<i>Pyrus.</i>	339
60	Shave-grass.	<i>Equisetum hyemale.</i>	651
69	Shell- or Shag-bark.	<i>Juglans alla.</i>	637
363	Sheep berry.	<i>Viburnum prunifolium.</i>	201
436	Shepherd's-purse.	<i>Thlaspi.</i>	435
434	Shield-fern.	<i>Aspidium.</i>	660
344	Side-saddle-flower.	<i>Sarracenia.</i>	367
297	Silver-tree.	<i>Eleagnus.</i>	114
291	— bell tree.	<i>Halesia.</i>	449
235	Snake-root.	<i>Aristolochia.</i>	596
90	— Black-.	<i>Cimicifuga Serpentaria.</i>	372
89	— Button-	<i>Liatris.</i>	507
283	— Plantain-	<i>Erigeron bellidifolium.</i>	532
19	— Fern.	<i>Botrychium.</i>	655
373	— weed.	<i>Polygonum bistortoides.</i>	271
576	Snap-dragon.	<i>Antirrhinum.</i>	421
374	Sneeze-wort.	{ <i>Achillea Ptarmica.</i>	562
162	Snow-berry.	<i>Convallaria.</i>	232
2	Snow-drop-tree.	<i>Chionocca.</i>	100
197	— flock-tree.	<i>Halesia.</i>	449
176	— flower.	{ <i>Chionanthus.</i>	7
2	Soap-berry.	<i>Sapindus.</i>	274
317	— wort.	<i>Saponaria.</i>	314
209	Solomon's Seal.	<i>Polygonatum.</i>	234
249	Sorrel.	<i>Rumex.</i>	247
370	— tree.	<i>Andromeda arborea.</i>	295
277	Spear-wort.	<i>Ranunculus Lingua.</i>	391
381	Speedwell.	<i>Veronica.</i>	10
646	Spice-wood.	<i>Laurus Benzoin.</i>	279
310	Spider-wort.	<i>Tradescantia.</i>	218
269	Spikenard.	<i>Aralia racemosa.</i>	209
197	Spindle tree.	<i>Erythronium.</i>	103
133	Spleen-wort.	<i>Asplenium.</i>	665
412	Spoon-wood.	<i>Tilia.</i>	302
398	Spring-grass.	<i>Anthoxanthum.</i>	65
87	Spruce-pine.	<i>Pinus.</i>	639
411	Spurrey.	<i>Spergula.</i>	320
565	Staff-tree.	<i>Celastrus.</i>	167
305	Star-wort.	{ <i>Aster.</i>	543
165	— Water-.	<i>Aletris.</i>	225
34		{ <i>Melanthium divaricatum.</i>	242
47		<i>Callitricha</i>	

Stitch-wort.	<i>Stellaria.</i>	317	Timothy-g
Stock.	<i>Cheiranthus.</i>	436	Toad-flax.
Stone-crop.	<i>Sedum.</i>	323	_____.
Stramonium.	<i>Datura.</i>	141	Tobacco.
Strawberry.	<i>Fragaria.</i>	357	Tooth-ach-
— Blite.	<i>Bütom.</i>	4	Tooth-wor-
— tree.	<i>Arlutus.</i>	282	Touch-me-
Succory.	<i>Cichorium Intybus.</i>	496	Traveller's-
— Gum-	<i>Chondrilla.</i>	497	Tread-sofily
Sugar-cane. Wild-	<i>Succharum.</i>	73	Tree Prim-
Sugar-tree.	<i>Acer saccharinum.</i>	266	Trefoil.
Sumach.	<i>Rhus.</i>	204	_____ Bi
Sun-dew.	<i>Drosera.</i>	210	_____ M
— drop.	<i>Oenothera fruticosa.</i>	262	Trumpet-fl
— flower.	<i>Helianthus.</i>	570	Tulip-tree.
— Tick-seed-	<i>Coreopsis.</i>	567	Tupelo.
Supple Jack.	<i>Zizyphus.</i>	188	Turnerick.
Swallow-wort.	<i>Asclepias.</i>	180	Turnsole.
Sweet-bay.	{ <i>Laurus.</i>	275	Umbrella-tr
— briar.	{ <i>Magnolia glauca.</i>	381	Valerian.
— fern-bush.	<i>Rosa.</i>	344	_____ Gi
— gale.	<i>Comptonia.</i>	635	Venus's-fly-t
— leaf.	<i>Myrica.</i>	620	Vervain.
— scented shrub.	<i>Hopea.</i>	451	Vetch.
Sycamore.	<i>Calycanthus.</i>	357	_____ Milk
Tacamahac.	<i>Platanus.</i>	635	Vine-grape.
Tallow-tree.	<i>Populus balsamifera.</i>	618	Violet.
Tamarack.	<i>Stillingia.</i>	608	_____ Water
Tansey.	<i>Pinus pendula.</i>	639	Viper's Bugle
— Wild-	<i>Tanacetum.</i>	523	_____ grass
Tarragon.	<i>Potentilla Anserina.</i>	356	Virgin's-bow
Tea. Labrador-	<i>Artemisia Dracunculus.</i>	521	_____.
— Mountain-	<i>Ledum latifolium.</i>	300	Wake-Robin.
— New Jersey-	<i>Gaultheria.</i>	283	Walking-fern.
— Oswego-	<i>Ceanothus.</i>	167	Wall-flower.
— South Sea-	<i>Monarda.</i>	17	— nut-tree.
Teasel.	<i>Ilex vomitoria.</i>	118	— Rue.
Thistle.	<i>Dipsacus.</i>	96	Water-leaf.
— Sow-	{ <i>Carduus.</i>	507	— Radish.
Thorn-apple.	{ <i>Cnicus.</i>		Wax Myrtle.
Thorough-wort.	<i>Sonchus.</i>	501	Way-faring-tr
Thrift.	<i>Datura.</i>	141	Wheat.
Thyme.	<i>Eupatorium perfoliatum.</i>	516	_____ Cow
Tick-seed.	<i>Statice.</i>	221	White-wood.
	<i>Thymus.</i>	413	
	<i>Corispermum.</i>	4	

317	Timothy-grass.	<i>Phleum pratense.</i>	65
436	Toad-flax.	<i>Antirrhinum Linaria.</i>	421
323	—. Bastard-	<i>Thesium.</i>	177
141	Tobacco.	<i>Nicotiana.</i>	144
357	Tooth-ach-tree.	<i>Zanthoxylum.</i>	209
4	Tooth-wort.	<i>Dentaria.</i>	438
282	Touch-me-not.	<i>Impatiens.</i>	171
496	Traveller's-joy.	<i>Clematis.</i>	384
497	Tread-softly.	<i>Jatropha.</i>	603
73	Tree Primrose.	<i>Oenothera.</i>	261
266	Trefoil.	<i>Trifolium.</i>	477
204	—. Bird's-foot-	<i>Lotus.</i>	489
210	—. Marsh-	<i>Menyanthes trifolia.</i>	139
262	Trumpet-flower.	<i>Bignonia.</i>	419
570	Tulip-tree.	<i>Liriodendron.</i>	382
567	Tupelo.	<i>Nyssa.</i>	177
188	Turnerick.	<i>Hydrastis.</i>	389
180	Turnsole.	<i>Heliotropium.</i>	130
275			
381	Umbrella-tree.	<i>Magnolia tripetala.</i>	361
344			
635	Valerian.	<i>Valeriana.</i>	26
620	—. Greek-	<i>Polemonium.</i>	151
451	Venus's-fly-trap.	<i>Dionaea.</i>	304
357	Vervain.	<i>Verbena.</i>	415
635	Vetch.	<i>Vicia.</i>	471
618	—. Milk-	<i>Astragalus.</i>	472
608	Vine-grape.	<i>Vitis.</i>	169
639	Violet.	<i>Viola.</i>	171
523	—. Water-	<i>Hottonia.</i>	138
356	Viper's Bugloss.	<i>Echium.</i>	130
521	— grass.	<i>Scorzonera.</i>	497
300	Virgin's-bower.	<i>Clematis.</i>	384
283			
167	Wake-Robin.	<i>Arum.</i>	399
17	Walking-fern.	<i>Lycopodium alopecuroides.</i>	654
118	Wall-flower.	<i>Cheiranthus.</i>	436
96	— nut-tree.	<i>Juglans.</i>	636
507	— Rue.	<i>Asplenium Ruta.</i>	666
501	Water-leaf.	<i>Hydrophyllum.</i>	134
141	— Radish.	<i>Sisymbrium amphibium.</i>	440
516	Wax Myrtle.	<i>Myrica.</i>	620
221	Way-faring-tree.	<i>Viturnum lantanoides.</i>	201
413	Wheat.	<i>Triticum.</i>	90
4	— Cow-	<i>Melampyrum.</i>	43
	White-wood.	<i>Tilia.</i>	362

Whitlow-grass.	<i>Drala.</i>	433
Whortle-berry.	<i>Vaccinium.</i>	284
Willow.	<i>Salix.</i>	608
— herb.	<i>Epilobium.</i>	259
Wind-flower.	<i>Anemone.</i>	386
Winter-berry.	<i>Prinos.</i>	220
— green.	<i>Pyrola. Chimaphila.</i>	299
— Chick weed.	<i>Trientalis.</i>	254
Wire-grass.	<i>Chloris.</i>	87
Witch Hazel.	<i>Hamamelis.</i>	116
Wolf's-bane.	<i>Aconitum.</i>	372
Wood-bine.	<i>Caprifolium.</i>	160
— sorrel.	<i>Oalis.</i>	322
Worm-grass.	<i>Spigelia.</i>	139
— seed.	<i>Chenopodium Anthelminticum.</i>	198
	<i>Erysimum cheiranthoideum.</i>	436
Yam.	<i>Dioscorea.</i>	251
Yarrow	<i>Achillea millefolium.</i>	503
Yellow Rattle.	<i>Rhinanthus.</i>	429
— Root.	<i>Zantorrhiza.</i>	212
Yew-tree.	<i>Hydrastis.</i>	389
	<i>Taxus.</i>	647

GEN

ABIES.

Acacia.

Acalypha.

Acer.

Achillea.

Achyranth

Acnida.

Aconitum.

Acorus.

Acrostichu

Acrostichu

Actaea.

Actaea.

Actinea.

Actinella.

Adelia.

Adiantum.

Ægilops.

Aeschynom

Æschynome

Æsculus.

Æzelia.

Agave.

Ageratum.

Agrimony.

Agrostemm

Agrostis.

Agrostis.

Aira.

433
284
608
259
386
220
299
254
87
116
372
160
322
139
n. 198
436
251
563
429
212
389
647

INDEX

GENERUM ET SYNONYMORUM.

<i>ABIES.</i>		<i>Aira.</i>	85
<i>Acacia.</i>	i. 373) 305	<i>Ajuga.</i>	ii. 471) 404
<i>Acalypha.</i>	ii. 687) 604	<i>Alchemilla.</i>	i. 139) 112
<i>Acer.</i>	i. 342) 265	<i>Aletris.</i>	i. 293) 225
<i>Achillea.</i>	ii. 646) 562	<i>Alisma.</i>	i. 331) 252
<i>Achyranthes.</i>	445	<i>Allionia.</i>	i. 118) 97
<i>Acnida.</i>	i. 268) 208	<i>Allium.</i>	i. 288) 222
<i>Aconitum.</i>	ii. 445) 372	<i>Alnus.</i>	ii. 695) 622
<i>Acorus.</i>	i. 309) 235	<i>Alopecurus.</i>	i. 77) 65
<i>Acrostichum.</i>	ii. 721) 658	<i>Alsine.</i>	317
<i>Acrostichum.</i>	659, 660, 661, 666, 670	<i>Alyssum.</i>	ii. 523) 434
<i>Actaea.</i>	ii. 435) 306	<i>Amaranthus.</i>	i. 267) 206
<i>Actaea.</i>	372, 373	<i>Amaryllis.</i>	i. 287) 222
<i>Actinea.</i>	500	<i>Ambrosia.</i>	ii. 666) 580
<i>Actinella.</i>	ii. 639) 560	<i>Amellus.</i>	ii. 650) 564
<i>Adelia.</i>	22	<i>Amellus.</i>	502
<i>Adiantum.</i>	ii. 733) 670	<i>Ammantia.</i>	i. 135) 107
<i>Ægilops.</i>	87, 88	<i>Ammi.</i>	i. 238) 192
<i>Aeschynomene.</i>	ii. 583) 485	<i>Ammyrsine.</i>	i. 360) 301
<i>Aeschynomene.</i>	485	<i>Amorpha.</i>	ii. 501) 466
<i>Æsculus.</i>	i. 333) 254	<i>Ampelopsis.</i>	170
<i>Afzelia.</i>	424	<i>Amsonia.</i>	i. 225) 184
<i>Agave.</i>	i. 294) 226	<i>Anagallis.</i>	i. 166) 135
<i>Ageratum.</i>	516	<i>Andromeda.</i>	i. 358) 290
<i>Agrimonie.</i>	i. 409) 335	<i>Andromeda.</i>	264, 286
<i>Agrostemma.</i>	i. 393) 321	<i>Androphylax.</i>	252
<i>Agrostis.</i>	i. 73) 63	<i>Andropogon.</i>	i. 85) 74
<i>Agrostis.</i>	61, 87	<i>Andropogon.</i>	73
<i>Aira.</i>	i. 89) 76	<i>Androsace.</i>	i. 170) 137
		<i>Anemone.</i>	ii. 457) 386

<i>Anemone.</i>	391	<i>Aspidium.</i>	ii. 724) 660	Boeb
<i>Angelica.</i>	i. 243) 193	<i>Aspidium.</i>	670	Boeh
<i>Anona.</i>	383	<i>Asplenium.</i>	ii. 727) 666	Boerl
<i>Anonymos</i> Walteri.	10, 13, 28, 31, 115, 132, 139, 185, 187, 231, 252, 334, 416, 424, 474, 484, 507, 508, 510, 511	<i>Asplenium.</i>	664, 667	Bolta
<i>Anthemis.</i>	ii. 645) 562	<i>Aster.</i>	ii. 636) 543	Borya
<i>Anthericum.</i>	242, 246	<i>Astragalus.</i>	532, 533, 534	Botry
<i>Anthoxanthus.</i>	i. 74) 65	<i>Astragalus.</i>	ii. 570) 472	Bract
<i>Anthoxanthum.</i>	73	<i>Astragalus.</i>	473	Brase
<i>Antirrhinum.</i>	ii. 506) 421	<i>Athanasia.</i>	519, 520	Brassi
<i>Anychia.</i>	i. 27) 176	<i>Atheropogon.</i>	i. 86) 75	Eriza
<i>Apargia.</i>	ii. 594) 497	<i>Athyrium.</i>	664	Brodi
<i>Aphanes.</i>	112	<i>Atragene.</i>	ii. 455) 384	Brom
<i>Apios.</i>	ii. 572) 473	<i>Atriplex.</i>	i. 255) 199	Brunn
<i>Apium.</i>	192	<i>Atropa.</i>	158	Buchi
<i>Apocynum.</i>	i. 223) 179	<i>Avena.</i>	i. 100) 85	Bume
<i>Aquilegia.</i>	ii. 440) 372	<i>Avena.</i>	76	Bunia
<i>Arabis.</i>	ii. 530) 436	<i>Azalea.</i>	i. 190) 151	Bupht
<i>Arachis.</i>	480	<i>Azalea.</i>	204	Bupht
<i>Aralia.</i>	i. 270) 209	<i>Azolla.</i>	ii. 739) 672	Burma
<i>Arbutus.</i>	i. 355) 282	<i>Baccharis.</i>	ii. 623) 523	Cabom
<i>Arbutus.</i>	283	<i>Baccharis.</i>	524	Cacalit
<i>Aretium.</i>	ii. 606) 505	<i>Balsamona.</i>	335	Cactus
<i>Arenaria.</i>	i. 389) 317	<i>Baltimora.</i>	ii. 663) 580	Cakile
<i>Arethusa.</i>	ii. 673) 590	<i>Bartonia.</i>	i. 400) 327	Caladi
<i>Arethusa.</i>	591, 593	<i>Bartonia.</i>	100	Calam
<i>Argemone.</i>	ii. 433) 360	<i>Bartsia.</i>	ii. 515) 429	Calla,
<i>Argolasia.</i>	224	<i>Batschia.</i>	i. 160) 132	Calla,
<i>Aristida.</i>	i. 82) 72	<i>Esfaria.</i>	362	Callicla
<i>Aristolochia.</i>	i. 679) 596	<i>Bejaria.</i>	ii. 423) 362	Callico
<i>Arnica.</i>	ii. 629) 527	<i>Bellis.</i>	ii. 626) 526	Calligo
<i>Artemesia.</i>	ii. 621) 520	<i>Berberis.</i>	i. 283) 219	Callitri
<i>Arum.</i>	ii. 470) 399	<i>Bernhardia.</i>	ii. 715) 655	Caloch
<i>Arundinaria.</i>	59	<i>Betula.</i>	ii. 694) 620	Caltha.
<i>Arundo.</i>	i. 101) 86	<i>Bidens.</i>	623	Calycan
<i>Arundo.</i>	59, 65	<i>Bignonia.</i>	ii. 653) 566	Calyde
<i>Asarum.</i>	ii. 680) 596	<i>Bignonia.</i>	519	Calypso
<i>Asclepias.</i>	i. 224) 180	<i>Biscutella.</i>	ii. 503) 419	Calyste
<i>Ascyrum.</i>	ii. 448) 373	<i>Blairia.</i>	10, 185	Campa
<i>Ascyrum.</i>	379	<i>Blandfordia.</i>	435	Canna
<i>Asparagus.</i>	i. 307) 235	<i>Blechnum.</i>	417	Cannab
		<i>Blechnum.</i>	ii. 731) 669	Cantua
		<i>Blitum.</i>	446	Caprari
			670	Capriffo
			4	

660			
670			
666	Boebera.	ii. 638) 559	Cardamine. ii. 534) 439
, 667	Boehmeria.	i. 140) 112	Cardiospermum. i. 349) 273
543	Boerhavia.	i. 39) 31	Carduus. ii. 608) 507
, 534	Boltonia.	ii. 641) 561	Carduus. 506
472	Borya.	i. 30) 22	Carex. i. 47) 34
473	Botrychium.	ii. 717) 655	Carex. 45
, 520	Botrypus.	655	Carpinus. ii. 696) 623
75	Brachystemum.	409, 410	Carpinus. 623
664	Brasenia.	ii. 460) 389	Carthamus. 505
384	Brassica.	ii. 532) 438	Cassia. i. 375) 305
199	Briza.	i. 94) 82	Cassine. 118
158	Brodiaea.	i. 289) 223	Castalia. 368
85	Bromus.	i. 98) 85	Castanea. ii. 699) 624
76	Brunnichia.	i. 348) 273	Catalpa. i. 16) 10
151	Buchnera.	ii. 505) 421	Caulinia. i. 2) 2
204	Buchnera.	416	Caulophyllum. i. 282) 218
672	Bumelia.	i. 191) 155	Ceanothus. i. 206) 167
523	Bunias.	434	Celastrus. i. 207) 167
524	Buphtalmum.	ii. 649) 563	Celtis. i. 259) 200
335	Buphtalmum.	563	Cenchrus. i. 66) 60
580	Burmannia.	i. 278) 217	Centaurea. ii. 658) 576
327	Cabomba.	239	Centaurella. i. 124) 99
100	Cacalia.	ii. 615) 518	Centunculus. i. 121) 97
429	Cactus.	i. 399) 327	Cephalanthus. i. 144) 114
132	Cakile.	ii. 522) 434	Cephalotis. 238
362	Caladium.	399	Cerastium. i. 392) 320
362	Calamintha.	ii. 492) 413	Ceratiola. i. 29) 21
526	Calla.	ii. 469) 399	Ceratophyllum. ii. 440) 370
219	Calla.	399	Cercis. i. 377) 308
655	Callicarpa.	i. 120) 97	Ceropegia. i. 224 b.) 184
620	Calligonum.	ii. 441) 370	Chærophillum. i. 249) 195
623	Callitricha.	i. 6) 3	Chamærops. i. 314) 239
566	Calochortus.	i. 316) 240	Chamærops. 239
519	Caltha.	ii. 461) 389	Chaptalia. ii. 650) 577
419	Calycanthus.	i. 422) 357	Chara. i. 10) 4
185	Calydermos.	158	Cheilanthes. ii. 734) 670
435	Calypso.	ii. 677) 593	Cheiranthus. ii. 528) 436
417	Calystegia.	i. 183) 142	Chelidonium. ii. 429) 365
446	Campanula.	i. 197) 159	Chelidonium. 365
569	Canna.	ii. 670) 585	Chelone. ii. 510) 427
570	Cannabis.	i. 256) 199	Chelone. 427, 428
4	Cantua.	i. 186) 147	Chenopodium. i. 254) 197
	Capraria.	ii. 490) 418	Chimaphila. i. 364) 300
	Caprifolium.	i. 199) 160	Chiococca. i. 198) 160

INDEX GENERUM ET SYNONYMORUM.

<i>Chionanthus.</i>	i. 13) 7	<i>Convolvulus.</i>	i. 184) 143	Dacty
<i>Chironia.</i>	i. 137	<i>Convolvulus.</i>	i. 142, 145, 146	Dacty
<i>Chlora.</i>	i. 137	<i>Conyzæ.</i>	ii. 624) 523	Dalea
<i>Chloris.</i>	i. 103) 87	<i>Conyza.</i>	543, 544, 556	Dalea
<i>Chondrilla.</i>	ii. 597) 497	<i>Coptis.</i>	ii. 462) 390	Daliba
<i>Chrysanthemum.</i>	ii. 627) 526	<i>Coreopsis.</i>	ii. 654) 567	Datisc
<i>Chrysanthemum.</i>	527, 561	<i>Coreopsis.</i>	565, 566	Datur
<i>Chrysobalanus.</i>	i. 403) 329	<i>Corispernum.</i>	i. 8) 4	Daucu
<i>Chrysocoma.</i>	ii. 614) 517	<i>Cornuçoipæ.</i>	61	Daucu
<i>Chrysocoma.</i>	511, 512, 540	<i>Cornuſ.</i>	i. 137) 107	Decod
<i>Chrysogonium.</i>	ii. 662) 579	<i>Coronopus.</i>	ii. 524) 435	Decun
<i>Chrysosplenium.</i>	i. 346) 269	<i>Corydalis.</i>	ii. 558) 462	Delph
<i>Cichorium.</i>	iii. 593) 496	<i>Corylus.</i>	ii. 701) 634	Demid
<i>Cicuta.</i>	i. 247) 195	<i>Corypha.</i>	239, 240	Denta
<i>Cicuta.</i>	192	<i>Cotula.</i>	562	Diant
<i>Cimicifuga.</i>	ii. 447) 372	<i>Cratægus.</i>	i. 410) 336	Dianth
<i>Cineraria.</i>	ii. 630) 528	<i>Cratægus.</i>	339, 340	Diapet
<i>Cinna.</i>	64, 75	<i>Cristaria.</i>	ii. 553) 453	Dicho
<i>Circæa.</i>	i. 28) 21	<i>Critonia.</i>	512	Dichro
<i>Cirsium.</i>	506	<i>Crotalaria.</i>	ii. 564) 469	Dicks
<i>Cissampelos.</i>	252	<i>Crotalaria.</i>	307, 308	Diervil
<i>Cissus.</i>	i. 210) 170	<i>Croton.</i>	ii. 686) 603	Digital
<i>Cistus.</i>	363	<i>Crotonopsis.</i>	i. 265) 206	Digital
<i>Clarckia.</i>	i. 337) 260	<i>Cteisium.</i>	657	Dilatris
<i>Claytonia.</i>	i. 215) 175	<i>Cucubalus.</i>	i. 386) 315	Dilepy
<i>Clematis.</i>	ii. 456) 384	<i>Cucubalus.</i>	316	Diodia
<i>Cleome.</i>	ii. 536) 441	<i>Cunila.</i>	ii. 475) 406	Dionæ
<i>Clethra.</i>	i. 367) 301	<i>Cunila.</i>	414	Diosco
<i>Cliftonia.</i>	302	<i>Cuphea.</i>	i. 407) 335	Diospy
<i>Clinopodium.</i>	ii. 486) 410	<i>Cupressus.</i>	ii. 708) 645	Diotis
<i>Clinopodium.</i>	408, 409	<i>Cuscuta.</i>	i. 149) 116	Diphyl
<i>Clitoria.</i>	ii. 587) 487	<i>Cyamus.</i>	ii. 467) 398	Dipsac
<i>Cnicus.</i>	ii. 607) 506	<i>Cymbidium.</i>	ii. 676) 592	Dirca.
<i>Cochlearia.</i>	435	<i>Cymbidium.</i>	593	Dodeca
<i>Collinsonia.</i>	i. 27) 20	<i>Cynanchum.</i>	179	Dolich
<i>Colonea.</i>	572	<i>Cynoctonum.</i>	139	Donia
<i>Comarum.</i>	i. 420) 356	<i>Cynodon.</i>	i. 80) 70	Doronic
<i>Commelinæ.</i>	i. 40) 31	<i>Cynoglossum.</i>	i. 163) 133	Draba.
<i>Commelinæ.</i>	32	<i>Cynosterus.</i>	87	Dracæn
<i>Comptonia.</i>	ii. 703) 635	<i>Cyperus.</i>	i. 56) 50	Dracoce
<i>Conium.</i>	i. 248) 195	<i>Cyperus.</i>	58	Dracon
<i>Conostylis.</i>	i. 292) 224	<i>Cypripedium.</i>	ii. 678) 594	Drosera
<i>Convallaria.</i>	i. 304) 232	<i>Cypripedium.</i>	593	Dryas.
<i>Convallaria.</i>	233, 234, 235	<i>Cyrilla.</i>	i. 211) 170	Dryas.

) 143	Dactylis.	i. 97) 84	Dulichium.	i. 57) 53
5, 146	<i>Dactylis.</i>	ii. 59)	<i>Dyssodid</i>	559
) 523	Dalea.	ii. 573) 474	Echites.	i. 221) 178
4, 556	<i>Dalea.</i>	461, 512	Echium.	i. 156) 130
) 390	Dalibarda.	i. 416) 350	Eclipta.	ii. 644) 561
) 507	Datisca.	ii. 442) 370	Elæagrus.	i. 145) 114
5, 566	Datura.	i. 179) 141	Elephantopus.	ii. 668) 582
) 4	Daucus.	i. 237) 191	Eleusine.	i. 102) 87
61	<i>Daucus.</i>	194	Ellisia.	i. 178) 141
) 107	<i>Decodon.</i>	334	Elodea.	ii. 450) 379
) 435	Decumaria.	i. 401) 828	<i>Elodea.</i>	33
) 462	Delphinium.	ii. 444) 371	Elymus.	i. 106) 89
) 634	<i>Demidofia.</i>	187	Elytraria.	i. 20) 13
, 240	Dentaria.	ii. 533) 438	Empetrum.	i. 116) 93
562	<i>Dianthera.</i>	13	Epigæa.	i. 360) 297
) 336	Dianthus.	i. 384) 314	Epilobium.	i. 335) 259
, 340	Diapensia.	i. 187) 147	Epipactis.	ii. 674) 591
453	Dichondra.	i. 229) 187	Equisetum.	ii. 713) 651
512	Dichromena.	i. 52) 47	<i>Erianthus.</i>	73, 74
469	Dicksonia.	ii. 735) 671	<i>Erica.</i>	264
, 308	Diervilla.	i. 202) 162	Erigeron.	ii. 634) 532
603	Digitaria.	i. 79) 69	<i>Erigeron.</i>	523
608	<i>Digitaria.</i>	70	Eriocaulon.	i. 113) 91
206	Dilatris.	i. 37) 30	Eriogonum.	i. 353) 277
657	<i>Dilepyrum.</i>	61	Eriophorum.	i. 60) 57
315	Diodia.	i. 133) 105	<i>Eriophorum.</i>	57
316	Dionæa.	i. 370) 304	<i>Eruvum.</i>	487
406	Dioscorea.	i. 328) 251	Eryngium.	i. 233) 189
414	Diospyros.	i. 341) 265	Erysimum.	ii. 527) 436
335	Diotis.	ii. 683) 602	Erythrina.	ii. 562) 467
645	Diphylla.	i. 281) 218	Erythronium.	i. 301) 230
116	Dipsacus.	i. 117) 96	<i>Erythrorhiza.</i>	446
398	Dirca.	i. 343) 268	<i>Ethulia.</i>	518
592	Dodecatheon.	i. 168) 136	Eupatorium.	ii. 612) 512
593	Dolichos.	ii. 566) 470	<i>Eupatorium.</i>	512, 517
179	<i>Doliches.</i>	469, 487	Euphorbia.	ii. 689) 605
139	Donia.	ii. 637) 559	Euphrasia.	ii. 516) 430
70	<i>Doronicum.</i>	528	Evolvulus.	i. 230) 187
133	Draba.	ii. 520) 433	Evonymus.	i. 208) 168
87	<i>Dracæna.</i>	233	Exacum.	i. 125) 100
50	Dracocephalum.	ii. 488) 411	Fagus.	ii. 698) 624
53	<i>Dracontium.</i>	398	<i>Fagus.</i>	624
594	Drosera.	i. 273) 210	Fedia.	i. 33) 28
593	<i>Dryas.</i>	i. 417) 350		
170	<i>Dryas.</i>	350, 352		

INDEX GENERUM ET SYNONYMORUM.

Ferula.	i. 240) 192	Gonolobium.	i. 222) 178	Hord
Festuca.	i. 96) 83	<i>Genolobus.</i>		Hott
<i>Filago.</i>	526	<i>Gordonia.</i>	ii. 550) 451	Hous
Fimbristylis.	i. 55) 49	<i>Gratiola.</i>	i. 19) 12	Huds
<i>Floerkea.</i>	239	<i>Gratiola.</i>	418, 419	Hum
<i>Forsythia.</i>	328	<i>Guilandina.</i>	304	Hydr
Fothergilla.	i. 408) 335	<i>Gymnandra.</i>	430	Hydr
Fragaria.	i. 421) 357	<i>Gymnocladus.</i>	i. 372) 304	Hydr
<i>Franklinia.</i>	451			Hydr
Frasera.	i. 127) 101	Halesia	ii. 547) 449	Hydr
Fraxinus.	i. 15) 8	<i>Hannamelis.</i>	i. 148) 116	Hydr
Fritillaria.	i. 300) 230	<i>Hamiltonia.</i>	i. 220) 178	Hydr
Fuirena.	i. 62) 58	<i>Hedema.</i>	ii. 493) 414	Hyos
<i>Fuirena.</i>	58	<i>Hedysotis.</i>	102, 107	Hyos
Fumaria.	ii. 559) 463	<i>Hedypnois.</i>	497	Hyos
<i>Fumaria.</i>	462, 463	<i>Hedysarum.</i>	ii. 581) 481	Hym
		<i>Hedysarum.</i>	480, 481,	Hym
Galactia.	ii. 586) 486		484, 485, 487	
Galardia.	ii. 656) 572	<i>Helenium.</i>	ii. 640) 560	Hype
<i>Gailarda.</i>	572	<i>Helianthemum.</i>	ii. 425) 363	Hype
Galax.	ii. 544) 446	<i>Helianthus.</i>	ii. 655) 570	
<i>Galega.</i>	489	<i>Helianthus.</i>	563	Hypo
Galeopsis.	ii. 479) 407	<i>Heliopsis.</i>	ii. 648) 563	Hypti
Galium.	i. 131) 102	<i>Heliotropium.</i>	i. 157) 130	Hyssc
Gaultheria.	i. 356) 283	<i>Helleborus.</i>	390	
Gaura.	i. 336) 260	<i>Helonias.</i>	i. 320) 242	Jatrop
Gelseminum.	i. 226) 184	<i>Helonias.</i>	241	Jeffer:
Gentiana.	i. 227) 185	<i>Hepatica.</i>	ii. 464) 391	Ilex.
Geranium.	ii. 546) 448	<i>Heracleum.</i>	i. 241) 192	Ilex.
Gerardia.	ii. 507) 422	<i>Heritiera.</i>	31	Illecel
<i>Geum.</i>	i. 418) 351	<i>Herpestis.</i>	ii. 500) 418	Illicio
<i>Gilia.</i>	147	<i>Hesperis.</i>	ii. 529) 436	Impat
<i>Gillenia.</i>	343	<i>Heteranthera.</i>	i. 42) 32	Indig
<i>Glandularia.</i>	416	<i>Heuchera.</i>	i. 231) 187	Inula.
Glaucium.	ii. 430) 365	<i>Hibiscus.</i>	ii. 556) 455	Ipome
Glaux.	i. 216) 176	<i>Hieracium.</i>	ii. 601) 502	Ipoma
Glechoma.	ii. 482) 408	<i>Hippophaë.</i>	i. 146) 115	Ipoma
Gleditschia.	i. 285) 221	<i>Hippuris.</i>	i. 4) 3	Iresin
<i>Globifera.</i>	10	<i>Hoffmannia.</i>	655	Iris.
Glycine.	ii. 585) 485	<i>Holcus.</i>	i. 91) 78	Isanth
<i>Glycine.</i>	470, 473, 474	<i>Holcus.</i>	82	Ischae
Glycirrhiza.	ii. 579) 480	<i>Holosteum.</i>	i. 110) 90	Isnara
Gnaphalium.	ii. 625) 524	<i>Hookera.</i>	223	Isoëte
<i>Gnaphalium.</i>	524	<i>Hopea.</i>	ii. 549) 451	Itea.
<i>Gomphrena.</i>	445, 446			Itea.

INDEX GENERUM ET SYNONYMORUM.

719

178		Hordeum.	i. 107) 89	Juglans.	ii. 705) 636
178		Hottonia.	i. 172) 138	Juncus.	i. 310) 235
451		Houstonia.	i. 134) 106	Juniperus.	ii. 710) 646
12		Hudsonia.	ii. 426) 364	Jussiaea.	i. 371) 304
, 419		Humulus.	i. 257) 199	Justicia.	i. 21) 13
304		Hydrangea.	i. 379) 309	Justicia.	13
430		Hydrastis.	ii. 459) 389	Iva.	ii. 665) 580
304		Hydrocotyle.	i. 234) 190	Ixia.	i. 35) 29
449		Hydroglossum.	ii. 718) 656	Kalmia.	i. 359) 295
116		Hydrolea.	i. 228) 187	Knoxia.	107
178		Hydropeltis.	389	Kochia.	i. 266) 206
414		Hydrophyllum.	i. 165) 134	Koeleria.	i. 99) 85
107		Hyoscyamus.	i. 180) 141	Krigia.	ii. 602) 504
497		Hyoseris.	ii. 603) 504	Kuhnbia.	ii. 611) 512
481		Hyoseris.	504, 505	Kuhnistera.	461
,		Hymenopappus.	ii. 617) 519	Kyllingia.	i. 50) 46
487		Hymenophyllum.	ii. 736) 671	Kyllingia.	58, 59
560		Hypericum.	ii. 449) 374	Lacathea.	451
363		Hypericum.	374, 379,	Lactuca.	ii. 599) 500
570			380, 451	Lactuca.	503
503		Hypoxis.	i. 291) 224	Lagotis.	430
563		Hyptis.	ii. 484) 408	Lamium.	ii. 478) 406
130		Hysopos.	ii. 476) 406	Larix.	645
390		Jatropha.	ii. 685) 603	Lathyrus.	ii. 568) 471
242		Jeffersonia.	i. 344) 268	Laurus.	i. 352) 275
391		Ilex.	i. 152) 117	Lechea.	i. 112) 90
192		Ilex.	221	Ledum.	i. 365) 300
31		Illecebrum.	ii. 543) 445	Ledum.	301
418		Illicium.	ii. 451) 380	Leersia.	i. 71) 62
136		Impatiens.	i. 213) 171	Lemma.	i. 31) 22
32		Indigofera.	ii. 589) 488	Leontice.	218
187		Inula.	ii. 633) 531	Leontodon.	ii. 596) 497
155		Ipomoea.	i. 185) 145	Leontodon.	497
502		Ipomoea.	147	Leonurus.	ii. 481) 408
601		Ipomopsis.	147	Lepidium.	ii. 525) 435
115		Iresine.	ii. 542) 445	Leptanthus.	32
3		Iris.	i. 36) 29	Lespedeza.	ii. 580) 480
155		Isanthus.	ii. 474) 405	Lewisia.	ii. 437) 308
78		Ischænum.	88	Leysera.	573
82		Isnardia.	111	Liatris.	ii. 609) 507
90		Isoëtes.	ii. 737) 671	Ligusticum.	i. 242) 193
23		Itea.	i. 212) 171	Ligustrum.	i. 11) 7
51		Itea.	171		

<i>Lilium.</i>	i. 299)	228	<i>Matuschkeae.</i>	352	Najas
<i>Lilium.</i>	230		<i>Mayaca.</i>	32	<i>Napa</i>
<i>Limnetis.</i>	i. 65)	59	<i>Medeola.</i>	i. 321)	<i>Nartt</i>
<i>Limodorum.</i>	589, 592, 593		<i>Medicago.</i>	ii. 592)	<i>Narti</i>
<i>Lindernia.</i>	ii. 501)	419	<i>Medicago.</i>	490	<i>Nectr</i>
<i>Linnæa.</i>	ii. 495)	415	<i>Melampyrum.</i>	ii. 517)	<i>Nelun</i>
<i>Linum.</i>	i. 272)	210	<i>Melanthera.</i>	ii. 618)	<i>Neott</i>
<i>Lippia.</i>		417, 418	<i>Melanthium.</i>	i. 317)	<i>Nepe</i>
<i>Liquidambar.</i>	ii. 702)	635	<i>Melanthium.</i>	242, 243,	<i>Nepet</i>
<i>Liquidambar.</i>		635	<i>Melica.</i>	246	<i>Neph</i>
<i>Liriodendron.</i>	ii. 453)	382	<i>Melilotus.</i>	i. 90)	Nican
<i>Lithospermum.</i>	i. 159)	131	<i>Melissa.</i>	77	Nicot
<i>Lithospermum.</i>		132	<i>Melothria.</i>	477	Nolin
<i>Lobelia.</i>	ii. 545)	446	<i>Menispernum.</i>	ii. 538)	Nuph
<i>Lonicera.</i>	160, 161, 162		<i>Menispernum.</i>	444	<i>Nymj</i>
<i>Lotus.</i>	ii. 591)	489	<i>Mentha.</i>	ii. 443)	<i>Nymp</i>
<i>Ludwigia.</i>	i. 138)	110	<i>Menyanthes.</i>	370	<i>Nyssa</i>
<i>Lupinaster.</i>	ii. 577)	479	<i>Menyanthes.</i>		
<i>Lupinus.</i>	ii. 563)	467	<i>Menzesia.</i>	i. 173)	
<i>Luzula.</i>		238	<i>Mespilus.</i>	139	Obola
<i>Lychnis.</i>	i. 394)	321		337, 338,	Oenan
<i>Lycium.</i>	i. 119)	97		339, 340	Oenot
<i>Lycopodium.</i>	ii. 714)	652	<i>Micranthemum.</i>	i. 17)	Olden
<i>Lycopodium.</i>		655	<i>Micropetalum.</i>	10	Olea.
<i>Lycopsis.</i>	i. 162)	132	<i>Mieglia.</i>	ii. 390)	Onocle
<i>Lycopus.</i>	i. 24)	16	<i>Mikania.</i>	319	Onosn
<i>Lygodium.</i>		657	<i>Milium.</i>	i. 613)	Ophio
<i>Lysimachia.</i>	i. 167)	135	<i>Mimosa.</i>	517	Ophio
<i>Lythrum.</i>	i. 406)	334	<i>Mimulus.</i>	ii. 613)	Ophry
<i>Lythrum.</i>		335	<i>Mitchella.</i>	62	Orchid
<i>Magnolia.</i>	ii. 452)	380	<i>Mitella.</i>	i. 128)	Orchis
<i>Majanthemum.</i>		233	<i>Mollugo.</i>	101	Origan
<i>Malachodendrum.</i>		452	<i>Momordica.</i>	i. 382)	Origan
<i>Malaxis.</i>	ii. 675)	592	<i>Monarda.</i>	92	Ornith
<i>Malope.</i>	ii. 555)	455	<i>Monnierae.</i>	ii. 539)	Ornith
<i>Malva.</i>	ii. 554)	454.	<i>Monotropa.</i>	444	Oryzop
<i>Malva.</i>		454	<i>Morus.</i>	i. 25)	Osmun
<i>Mapania.</i>	i. 51)	47	<i>Muhlenbergia.</i>	16	Osmux
<i>Mariscus.</i>	i. 63)	58	<i>Myagrum.</i>	i. 706)	Ostrya
<i>Marrubium.</i>	ii. 483)	408	<i>Mylocarium.</i>	639	Oxalis.
<i>Marshallia.</i>	ii. 619)	519	<i>Myosotis.</i>	i. 69)	Oxyco
<i>Marsilea.</i>		672	<i>Myrica.</i>	61	
<i>Martynia.</i>	ii. 512)	428	<i>Myriophyllum.</i>	ii. 521)	
<i>Matricaria.</i>		561	<i>Myrrhis.</i>	302	

352		Najas.	ii. 682)	602	Oxytropis.	ii. 571)	473
32		<i>Napaea.</i>		453	<i>Pachysandra.</i>	i. 151)	117
244		<i>Narthecium.</i>	i. 296)	227	<i>Panax.</i>	i. 235)	191
490		<i>Narthecium.</i>		246	<i>Pancratium.</i>	i. 286)	221
480		<i>Nectris.</i>	i. 312)	239	<i>Panicum.</i>	i. 78)	66
430		<i>Nelumbium.</i>		398	<i>Panicum.</i>	69, 70	
519		<i>Neottia.</i>	ii. 672)	589	<i>Papaver.</i>	ii. 431)	365
240		<i>Nepeta.</i>	ii. 477)	406	<i>Parietaria.</i>	i. 142)	114
246		<i>Nepeta.</i>		409	<i>Parnassia.</i>	i. 269)	208
77		<i>Nephrodium.</i>	659, 651,		<i>Parthenium.</i>	ii. 604)	580
477			663, 604, 605, 670		<i>Paspalum.</i>	i. 81)	70
413		<i>Nicandra.</i>	i. 194)	158	<i>Paspalum.</i>		69, 70
444		<i>Nicotiana.</i>	i. 181)	141	<i>Passiflora.</i>	ii. 541)	444
370		<i>Nolina.</i>	i. 315)	240	<i>Pastinaca.</i>	i. 250)	196
252		<i>Nuphar.</i>	ii. 439)	369	<i>Pedicularis.</i>	ii. 508)	424
105		<i>Nymphaea.</i>	ii. 438)	368	<i>Penthorum.</i>	i. 396)	323
39		<i>Nymphaea.</i>	369, 370, 398		<i>Pentstemon.</i>	ii. 511)	427
39		<i>Nyssa.</i>	i. 219)	177	<i>Peplis.</i>	i. 311)	238
64		<i>Obolaria.</i>	ii. 518)	431	<i>Perdicium.</i>		577
40		<i>Oenanthe.</i>	i. 245)	194	<i>Personoria.</i>		519, 520
10		<i>Oenothera.</i>	i. 338)	261	<i>Petalostemum.</i>	ii. 557)	461
19		<i>Oldenlandia.</i>	i. 129)	102	<i>Phaca.</i>		485
59		<i>Olea.</i>	i. 12)	7	<i>Phacelia.</i>	i. 177)	140
17		<i>Onoclea.</i>	ii. 725)	665	<i>Phæthusa.</i>	ii. 643)	561
62		<i>Onoclea.</i>		666, 669	<i>Phalangium.</i>	i. 295)	226
71		<i>Onosmodium.</i>	i. 161)	132	<i>Phalangium.</i>		240
25		<i>Ophioglossum.</i>	ii. 716)	655	<i>Phalaris.</i>	i. 75)	65
26		<i>Ophiorrhiza.</i>	i. 170)	139	<i>Phaseolus.</i>	ii. 565)	409
11		<i>Ophrys.</i>	589, 591, 592, 593		<i>Phellandrium.</i>	i. 246)	195
3		<i>Orchidocarpum.</i>		383	<i>Philadelphus.</i>	i. 402)	329
12		<i>Orchis.</i>	ii. 671)	585	<i>Phleum.</i>	i. 76)	65
14		<i>Origanum.</i>	ii. 487)	411	<i>Phlox.</i>	i. 188)	148
6		<i>Origanum.</i>		409, 410	<i>Phryma.</i>	ii. 496)	415
8		<i>Ornithogalum.</i>	ii. 297)	227	<i>Phyllanthus.</i>	ii. 537)	443
3		<i>Ornithogalum.</i>		222	<i>Phyllococe.</i>		264
9		<i>Ornus.</i>	i. 14)	8	<i>Physalis.</i>	i. 193)	157
1		<i>Orobanche.</i>	ii. 519)	431	<i>Phytolacca.</i>	i. 398)	324
4		<i>Orontium.</i>	i. 308)	235	<i>Pinguicula.</i>	i. 22)	14
2		<i>Oryzopsis.</i>	i. 67)	60	<i>Pinckneya.</i>	i. 190)	158
3		<i>Osmunda.</i>	ii. 720)	657	<i>Pinus.</i>	ii. 707)	639
1		<i>Osmurda.</i>	655, 669, 670		<i>Pistia.</i>	i. 345)	268
1		<i>Ostrya.</i>	ii. 697)	623	<i>Pisum.</i>	ii. 507)	470
1		<i>Oxalis.</i>	i. 395)	322	<i>Planera.</i>	i. 147)	115
1		<i>Oxycoetus.</i>	i. 339)	263	<i>Plantago.</i>	i. 122)	98

INDEX GENERUM ET SYNONYMORUM.

<i>Platanus.</i>	ii. 704)	635	<i>Pteris.</i>	669	Sabal.	
<i>Pleea.</i>	i. 354)	278	<i>Pteronia.</i>	509	Sabba	
<i>Pleuranthus.</i>		53	<i>Pulmonaria.</i>	i. 158)	130	Sacch.
<i>Poa.</i>	i. 93)	78	<i>Pycnanthemum.</i>		Sagina	
<i>Poa.</i>		85		ii. 485)	409	Sagin
<i>Podalyria.</i>	i. 376)	307	<i>Pyret'rum.</i>	ii. 628)	527	Sagitt
<i>Podophyllum.</i>	ii. 434)	366	<i>Pyrola.</i>	i. 363)	299	Salico
<i>Podophyllum.</i>		268	<i>Pyrola.</i>		Salix.	
<i>Podostemum.</i>	i. 5)	3	<i>Pyrularia.</i>		Salsol	
<i>Poiretia.</i>		106	<i>Pyrus.</i>	i. 411)	339	Salsol
<i>Polemonium.</i>	i. 189)	151	<i>Pyxidanthera.</i>		Salvia	
<i>Polemonium.</i>		140, 147			Salvin	
<i>Polygonum.</i>	i. 111)	90	<i>Quercus.</i>	ii. 700)	625	Sambu
<i>Polygonum.</i>	ii. 560)	464	<i>Queria.</i>		Samol	
<i>Polygonatum.</i>	i. 306)	234			Sangu	
<i>Polygonella.</i>		272	<i>Rafnia.</i>		Sangu	
<i>Polygonum.</i>	i. 347)	369	<i>Rojania.</i>		Sanicu	
<i>Polymnia.</i>	ii. 661)	579	<i>Ranunculus.</i>	ii. 465)	391	Santol
<i>Polymnia.</i>		503	<i>Ranunculus.</i>		Sapini	
<i>Polypodium.</i>	ii. 722)	658	<i>Raphis.</i>		Saponi	
<i>Polypodium.</i>	660, 661,		<i>Rhamnus.</i>	i. 205)	166	Sarrac
	663, 664, 671		<i>Rhamnus.</i>		Savoth	
<i>Polypodium.</i>	i. 123)	99	<i>Rhexia.</i>	i. 334)	257	Sature
<i>Polystichum.</i>	660, 661, 669		<i>Rhinanthus.</i>	ii. 514)	429	Satyri
<i>Pontederia.</i>	i. 290)	223	<i>Rhizanthus.</i>		Saurui	
<i>Pontederia.</i>		38	<i>Rhododendron.</i>	i. 361)	297	Saxifr
<i>Populus.</i>	ii. 692)	613	<i>Rhodora.</i>	i. 362)	208	Scandi
<i>Porcelia.</i>	ii. 454)	385	<i>Rhus.</i>	i. 262)	204	Scheuc
<i>Portulacca.</i>	ii. 428)	365	<i>Rhynchospora.</i>	i. 54)	48	Schisai
<i>Potamogeton.</i>	i. 154)	120	<i>Ribes.</i>	i. 204)	163	Schiza
<i>Potamogeton.</i>		274	<i>Ricinus.</i>	ii. 684)	602	Schoen
<i>Potentilla.</i>	i. 419)	353	<i>Robinia.</i>	ii. 588)	487	Schœn
<i>Pothos.</i>	ii. 468)	398	<i>Robinia.</i>		Schran	
<i>Prasium.</i>		411	<i>Rosa.</i>	i. 414)	344	
<i>Prenanthes.</i>	ii. 598)	498	<i>Rothia.</i>		Scirpu	
<i>Primula.</i>	i. 169)	137	<i>Rottboellia.</i>	i. 104)	88	Scirpu
<i>Prinos.</i>	i. 284)	220	<i>Rubia.</i>	i. 130)	102	
<i>Troserpinaca.</i>	i. 115)	92	<i>Rubus.</i>	i. 415)	346	Schwa
<i>Prunella.</i>	ii. 489)	411	<i>Rubus.</i>		Scleria	
<i>Prunus.</i>	i. 404)	329	<i>Rudbeckia.</i>	ii. 657)	573	Sclerai
<i>Psilotum.</i>		655	<i>Rudbeckia.</i>	500, 503,	572	Scolop
<i>Psoralea.</i>	ii. 574)	475	<i>Ruellia.</i>	ii. 504)	420	Scorz
<i>Psoralea.</i>		474	<i>Rumex.</i>	i. 326)	247	Scroph
<i>Ptelea.</i>	i. 136)	107	<i>Ruppia.</i>	i. 155)	121	Scutell
<i>Pteris.</i>	ii. 729)	668			Secale.	

669	Sabal.	i. 313)	239	Sedum.	i. 397)	323
509	Sabbatia.	i. 171)	137	Selinum.	i. 239)	192
130	Saccharum.	i. 84)	73	Senecio.	ii. 631)	528
409	Sagina.	i. 153)	119	Senecio.		528
527	Sagittaria.	ii. 466)	395	Serpicula.	i. 43)	33
209	Salicornia.	i. 3)	2	Serratula.	506, 507,	
300	Salix.	ii. 691)	608	Sesbania.	508, 509, 511	
178	Salsola.	i. 253)	197	Seseli.	i. 252)	197
339	Salsola.		206	Sibbaldia.	i. 274)	211
148	Salvia.	i. 26)	19	Sicyos.	ii. 540)	444
525	Salvinia.	ii. 728)	672	Sicyos.		444
176	Sambucus.	i. 261)	203	Sida.	ii. 552)	452
107	Samolus.	i. 195)	158	Sideroxylon.		155
173	Sanguinaria.	ii. 432)	360	Siegesbeckia.	ii. 642)	501
191	Sanguisorba.	i. 150)	116	Siegesbeckia.		505
190	Sancta.	i. 230)	191	Silene.	i. 387)	315
139	Santolina.	ii. 620)	520	Silene.		315
66	Sapindus.	i. 350)	274	Silphium.	ii. 660)	577
88	Saponaria.	i. 383)	314	Silphium.		503
57	Sarracenia.	ii. 430)	367	Sison.	i. 244 b.)	194
29	Sarothra.		378	Sison.		190, 195
23	Satureja.		410	Sisymbrium.	ii. 535)	440
97	Satyrium.		587, 589	Sisyrinchium.	i. 38)	31
98	Saururus.	i. 329)	252	Sium.	i. 244)	194
04	Saxifraga.	i. 380)	310	Smilacina.	i. 305)	232
48	Scandix.		195, 196	Smilax.	i. 327)	249
63	Scheuchzeria.	i. 325)	247	Smyrnium.	i. 251)	190
02	Schisandra.	i. 277)	212	Solanandra.		446
37	Schizæa.	ii. 719)	657	Solanum.	i. 192)	156
35	Schoenus.	i. 53)	47	Solidago.	ii. 635)	635
14	Schœnus.	47, 48, 49,	54, 57, 58	Sonchus.	ii. 600)	501
9	Schrankia.	i. 374)	305	Sophora.		307, 308
18	Scirpus.	i. 58)	54	Sorbus.	i. 412)	341
12	Scirpus.	47, 49, 51, 54,	57, 59	Sorghum.	i. 92)	78
6	Schwalbea.	ii. 513)	428	Sparganium.	i. 45)	33
0	Scleria.	i. 48)	45	Sparganophorus.	ii. 616)	518
3	Scleranthus.	i. 385)	315	Spergula.	i. 391)	320
2	Scolopendrium.	ii. 728)	667	Spermacoce.	i. 132)	105
0	Scorzonera.	ii. 595), 497		Spergulastrum.		319
7	Seriphularia.	ii. 502)	419	Spigelia.	i. 175)	139
1	Scutellaria.	ii. 490)	412	Spilanthes.		502
	Secale.	i. 108)	90	Spiræa.	i. 413)	341

INDEX GENERUM ET SYNONYMORUM.

<i>Staphylea.</i>	i. 263) 206	<i>Tillandsia.</i>	i. 279) 217	Valli
<i>Statice.</i>	i. 275) 212	<i>Tofieldia.</i>	i. 323) 246	Vera
<i>Stellaria.</i>	i. 388) 317	<i>Trachynotia.</i>	59	Vera
<i>Stellaria.</i>	317	<i>Tradescantia.</i>	i. 280) 218	Verb
<i>Stewartia.</i>	452	<i>Tragia.</i>	ii. 688) 604	Verb
<i>Stillingia.</i>	ii. 690) 608	<i>Tragopogon.</i>	505	Verbi
<i>Stipa.</i>	i. 83) 72	<i>Trattinickia.</i>	519	Verbi
<i>Stipulicida.</i>	90	<i>Trichodium.</i>	i. 70) 61	Vern
<i>Stokesia.</i>	ii. 605) 505	<i>Trichomanes.</i>	671	Vero
<i>Streptopus.</i>	i. 303) 232	<i>Trichophorum.</i>	i. 59) 57	Vibu
<i>Struthiopteris.</i>	ii. 726) 666	<i>Trichostema.</i>	ii. 494) 414	Vicia
<i>Stuartia.</i>	ii. 551) 451	<i>Trientalis.</i>	i. 332) 254	Villa
<i>Stylosanthes.</i>	ii. 578) 480	<i>Trifolium.</i>	ii. 576) 477	Vinc
<i>Styrax.</i>	ii. 548) 450	<i>Trifolium.</i>	475, 480, 486	Viola
<i>Sueda.</i>	197	<i>Triglochin.</i>	i. 324) 247	Virgi
<i>Swertia.</i>	i. 126) 100	<i>Trillium.</i>	i. 322) 244	Virg
<i>Swertia.</i>	138	<i>Triosteum.</i>	i. 203) 162	Visci
<i>Syena.</i>	i. 41) 32	<i>Tripsacum.</i>	i. 105) 88	Vitis
<i>Symphoria.</i>	i. 201) 162	<i>Tripterella.</i>	i. 34) 28	Vitis
<i>Symplocos.</i>	451	<i>Trisetum.</i>	i. 87) 76	Vitta
<i>Syntherisma.</i>	69, 70	<i>Triticum.</i>	i. 109) 90	Vogel
<i>Tabernemontana.</i>	184	<i>Trollius.</i>	ii. 463) 391	Wen
<i>Tagetes.</i>	559	<i>Troximum.</i>	ii. 604) 505	
<i>Talinum.</i>	ii. 427) 365	<i>Tubiflora.</i>	13	
<i>Tanacetum.</i>	ii. 622) 522	<i>Turritis.</i>	ii. 531) 438	
<i>Taxus.</i>	ii. 711) 647	<i>Turnera.</i>	i. 264) 206	
<i>Tecoma.</i>	420	<i>Tussilago.</i>	ii. 632) 531	
<i>Tephrosia.</i>	ii. 590) 489	<i>Tussilago.</i>	577	
<i>Tetragonotheca.</i>	ii. 647) 503	<i>Typha.</i>	i. 46) 34	
<i>Teucrium.</i>	ii. 472) 405	<i>Ulmus.</i>	i. 258) 199	
<i>Thalia.</i>	ii. 669) 584	<i>Uniola.</i>	i. 95) 82	
<i>Thalictrum.</i>	ii. 458) 388	<i>Uniola.</i>	84	
<i>Thalictrum.</i>	387	<i>Urtica.</i>	i. 141) 112	
<i>Thapsia.</i>	196	<i>Utricularia.</i>	i. 23) 15	
<i>Thesium.</i>	i. 218) 177	<i>Uvularia.</i>	i. 302) 231	
<i>Thlaspi.</i>	ii. 526) 435	<i>Uvularia.</i>	232	
<i>Thuya.</i>	ii. 709) 646	<i>Vaccinium.</i>	i. 357) 284	
<i>Thymbra.</i>	414	<i>Vaccinium.</i>	263, 283	
<i>Thymus.</i>	ii. 491) 413	<i>Vagnaria.</i>	i. 61) 58	
<i>Thymus.</i>	409, 413, 414	<i>Valantia.</i>	102	
<i>Tiarella.</i>	i. 381) 313	<i>Valeriana.</i>	i. 32) 28	
<i>Tigarea.</i>	i. 405) 333	<i>Valeriana.</i>	28	
<i>Tilia.</i>	ii. 424) 362			

217	Vallisneria.	ii. 681) 602	Woodsia.	ii. 723) 660
246	Veratrum.	i. 319) 242	Woodwardia.	ii. 732) 669
59	<i>Veratrum.</i>	243	<i>Wurmbea.</i>	225
218	Verbascum.	i. 182) 142	Xanthium.	ii. 667) 581
604	Verbena.	ii. 497) 415	Xylosteum.	i. 200) 161
505	<i>Verbena.</i>	417	Xyris.	i. 44) 33
519	Verbesina.	ii. 651) 564	Yucca.	i. 298) 227
61	<i>Verbesina.</i>	561	Zamia.	ii. 712) 648
671	Vernonia.	ii. 610) 511	Zannichellia.	i. 9) 4
57	Veronica.	i. 18) 10	Zanthorrhiza.	i. 276) 212
57	Viburnum.	i. 200) 201	Zanthoxylum.	i. 271) 209
414	Vicia.	ii. 509) 471	Zapania.	ii. 498) 417
254	Villarsia.	i. 174) 139	Zapania.	416
477	<i>Vincetoxicum</i>	179	Zea.	i. 49) 46
486	Viola.	i. 214) 171	Zigadenus.	i. 318) 241
247	Virgilia.	i. 378) 309	Zinnia.	ii. 652) 565
244	<i>Virgilia.</i>	572	Zizania.	i. 68) 60
162	Viscum.	i. 143) 114	Zizyphus.	i. 232) 188
88	Vitis.	i. 209) 169	Zornia.	ii. 582) 484
28	<i>Vitis.</i>	170	Zostera.	i. 1) 2
76	Vittaria.	ii. 730) 669		
90	<i>Vogelia.</i>	29		
91				
05	Wendlandia.	i. 330) 252		
13				
38				
06				
31				
77				
34				
99				
32				
34				
12				
5				
11				
2				
4				
3				
8				
2				
9				
8				

Lycopus
inte
Wi
Icon.
In low
July
to s
spec
in V
Sher
real

Monarda
subs
pilos
bii in
rugos
gulos
In Can
—Se
flowe

Valeriana
lato-
pinna
que o
Icon.
In swan
in Ha
colou

Fedia chen
ferne
menti

VOL. II.

SUPPLEMENTUM.

Lycopus exaltatus.—L. foliis basi pinnatifido-serratis, lacinias integerrimis subserratisque, calycibus spinescentibus.—

page 16.

Willd. sp. pl. 1. p. 121.

Icon. Fl. græc. t. 12.

In low shady woods : Virginia to Carolina ; rare. ♀. July, Aug. v. v. ; v. s. in Herb. Sherard. From three to six feet high. This species I omitted for want of a specimen in my Herbarium, though I found it myself in Virginia ; but seeing a specimen of Catesby's in the Sherardian Herbarium, I was convinced of its being a real American species.

p. 17.

Monarda rugosa.—M. glabriuscula ; capitulis mediocribus subsimplicibus, bracteis exterioribus ovatis undulatis pilosis, calycibus corollisque glabris, lacinia media labii inferioris bidentata, foliis ovatis subcordatis acutis rugosis, nervis subtus petiolisque pilosis, caule acutangulo glabriuscule.—Willd. sp. pl. 1. p. 125.

In Canada and on the mountains of Virginia. ♀. July —Sept. v. v. in Hort. Lambert. About four feet high ; flowers small, white.

p. 18.

Valeriana dioica.—V. glabra ; foliis radicalibus subspathulato-ovatis integris longissime petiolatis, caulinis paucis pinnatifidis, lacinias lanceolatis integerrimis subdentatisque obtusiusculis.—Willd. sp. pl. 1. p. 176.

Icon. Fl. dan. 687.

In swamps at Hudson's Bay. Tidden. ♀. June. v. s. in Herb. Sherard. About ~~a~~ span high ; flowers flesh-coloured.

Fedia chenopodifolia.—F. dichotoma ; foliis ovatis acutis inferne dentatis, cymis nudis divaricato-dichotomis, filamentis longis.

- Valeriana virginiana*, pedis anserinæ folio. *Pet. H. S. in Herb. Sherard.*
In Virginia. ♂. v. s. in *Herb. Sherard*. About a span high; flowers the size of *V. officinalis*.
- p. 72. *Aristida pallens*.—A. culmo erecto simplici, foliis filiformibus, radicalibus culmo brevioribus, panicula pauciflora, pedicellis alternis brevibus bifloris, aristis elongatis.—*Cav. ic. 5. t. 468. f. 2.*
In Upper Louisiana. *Bradbury*. ♀. v. s. in *Herb. Bradbury*. About a foot high; the awns more than three inches high.
- p. 73. *Stipa membranacea*.—S. foliis convoluto-filiformibus glabris, panicula laxa, pedicellis flexuosis, calycibus membranaceis 3-nervibus longe acuminatis, corollis calyce brevioribus sericeo-villosis, villis corollam superantibus, arista nuda recta calyce paulo longiore.
On the banks of the Missouri. *Bradbury*. v. s. in *Herb. Bradbury*. About eighteen inches high; flowers oval, covered with long silky hair.
- p. 85. *Bromus altissimus*.—B. panicula nutante, spiculis oblongis sub-6-floris, flosculis pubescentibus, valvula exteriore 3-nervata breviter aristata, foliis vaginisque glabris, culmo glabro altissimo.
On the banks of the Missouri. ♀. July, Aug. v. v. in *Hort. Lambert*. From five to eight feet high.
- p. 88. *Cynosurus secundus*.—C. glaber; foliis linearibus, spica elongata unilateralis, spiculis solitariis, defloratis reflexis, calycibus lineari-lanceolatis acutissimis.
In Upper Louisiana. *Bradbury*. v. s. in *Herb. Bradbury*. Resembles *C. retroflexus* very much.
- p. 97. *Allionia linearis*.—A. caule erecto tereti, foliis linearibus subcarnosis, floribus racemosis, pedicellis aggregatis, staminibus longissime exsertis.
Calymenia angustifolia. *Fraser Catal. 1813*.
In Upper Louisiana. *Bradbury*. v. s. in *Herb. Bradbury*.
- Allionia hirsuta*.—A. caule erecto angulato, foliis lanceolatis nervosis utrinque hirsutis, calycibus hirsutissimis sub-3-floris.
In Upper Louisiana. *Bradbury*. v. s. in *Herb. Bradbury*.

Planta

sc

bi

In

bi

Pulmo

bu

ot

In U

bu

cu

Pulmo

lit

Ancl

In C

Fl

vir

Cynoglo

ob

C. gl

In U

Br

Lysima

pet

qu

la

L. ca

In C

Flc

Calysteg

lon

bre

rib

In Ca

lar

Calysteg

lon

elo

aril

Conv

flo

He

Plantago elongata.—P. foliis linearibus integerrimis glabris, scapo spicis elongatis subimbricatis foliis longioribus, bracteis ovatis acutis margine membranaceis.

In Upper Louisiana. *Bradbury*. v. s. in *Herb. Bradbury*. Resembles *P. pauciflora*.

Pulmonaria lanceolata.—P. glabra, erecta; foliis radicalibus longissime pedicellatis lanceolatis, caulinis linearioribus, floribus subpaniculatis, calycibus abbreviatis.

In Upper Louisiana. *Bradbury*. 4. v. s. in *Herb. Bradbury*. Flowers small, resembling those of *P. paniculata*.

Pulmonaria sibirica.—P. calycibus abbreviatis, foliis radicalibus cordatis. *Willd. sp. pl. 1. p. 770.*

Anchusa. Gmel. sib. 4. p. 75. no. 15. t. 39.

In Canada. *Masson*. 4. July. v. s. in *Herb. Lambert*.

Flowers purple, in terminal corymbs; it resembles *P. virginica* very much.

Cynoglossum glomeratum.—C. hispidum; foliis spatulatis obtusis, floribus glomeratis.

C. glomeratum. Fraser. catal. 1813.

In Upper Louisiana. *Bradbury*. 4. v. s. in *Herb. Bradbury*.

Lysimachia lanceolata.—L. glaberrima; foliis quaternis subpetiolatis lanceolatis promissa acuminatis, pedunculis quaternis multifloris, floribus summis racemosis, corollae lacinias ovatis acutis, caule erecto simplici.

L. caroliniana quadrifolia, foliis acutis. *Herb. Sherard*.

In Carolina *Catesby*. 4. v. s. in *Herb. Sherard*. Flowers the size of *L. vulgaris*.

Calystegia Catesbeiana.—C. volubilis, tomentosa; foliis oblongis sagittatis acuminatis, pedunculis unifloris folio brevioribus, bracteis ovalibus obtusis calyce longioribus.

In Carolina. *Catesby*. v. s. in *Herb. Sherard*. Flowers large, purple.

Calystegia paradoxa.—C. humifusa, tomentosa; foliis oblongis cordato-sagittatis acutis, pedunculis unifloris elongatis folio longioribus, bracteis a flore remotis linearibus, calycibus nudis glabris acuminatis.

Convolvulus virginianus, laeviter hirsutis et oblongis foliis, flore maximo albicante. *Pluk. mant. 54. secundum Herb. Sherard*.

p. 99.

p. 131.

p. 133.

p. 135.

p. 143.

- In Virginia or Carolina. *v. s. in Herb. Sherard.* Flowers large, white.
- p. 143.* *Convolvulus Sherardi.*—C. humifusus, pubescens; foliis ellipticis utrinque retusis mucronatis, floribus solitariis subsessilibus.
C. forte monococco Banist. similis, sed vasculis sessilibus. *Herb. Sherard.*
In Carolina. *Catesby.* ♂. *v. s. in Herb. Sherard.* Resembles *C. trichosanthes* very much; I have only seen a specimen with fruit.
- p. 147.* *Cantua parviflora.*—C. pubescens, diffusa; foliis pinnatifidis, lacinias linearibus elongatis, pedunculis solitariis unifloris terminalibus et oppositifoliis, corollis calyce vix longioribus.
Ipomopsis inconspicua. *Smith exot bot. 1. t. 14.*
In North America. *Hoy.* ♂. June. *v. v. in Hortis.* Flowers small, pale blue. I insert this plant on the authority of Exotic Botany; but at the same time I doubt very much of its being a native of North America, and more strongly suspect it to come from Mexico.
- p. 148.* *Phlox acuminata.*—P. erecta, pubescens; caule angulato, foliis ovato-lanceolatis acuminatis inferne angustatis subpetiolatis subtus pubescentibus, floralibus subcordatis, corymbis paniculatis, corollæ lacinias rotundatis, tubo pubescente, calycibus aristatis.
Phlox decussata. *Lyon catal.* 1812.
On the mountains of Georgia and South Carolina. *Lyon.* ♀. Aug. Sept. *v. v. in Hort. Lambert.* From four to five feet high; flowers very handsomely variegated, with dark and light shades of purple.
- p. 149.* *Phlox nitida.*—P. erecta, glaberrima; caule tereti maculato lævigato, foliis ovato-oblongis subcarnosis supra nitidis atrovirentibus, subtus pallidis, corymbis fastigiatis, ramis inferioribus elongatis nudiusculis, corollæ lacinias lato-ovatis subretusis, dentibus calycinis lanceolatis mucronatis.
P. suffruticosa. *Hortul.*
In South Carolina. *Fraser.* ♀. Aug.—Oct. *v. v. in Hortis.* Near two feet high; flowers of a brilliant purple, which is deeper than in any other species. This species approaches near to *P. Carolina*; but besides its other characters it distinguishes itself very easily by flowering later and longer than that species.

Solanum
tis,
ob
for
Icon.
The J
rai
Mi
tio
Physalis
inte
cor
Icon.
In So
She

Campan
thu
lan
cor
C. am
In No
nec
size
kno
by
non
coni

Ribes la
den
long
gatu
hisp
On the
Ban
size

Ribes sp
foliu
sis
sub-
glan
ribu
On the
Ban

p. 156.

Solanum heterandrum.—S. annum; caule folisque aculeatis, foliis ovatis pinnatifidis utrinque hirsutis, laciinis obtusiusculis, antheris declinatis distantibus, ima corniformi productiore, bacca calyce echinato inclusa.

Icon. Tab. 7.

The present description has been taken from a living plant raised by A. B. Lambert, Esq. from seeds received of Mr. Nuttall. The native specimen the former description was made from was very imperfect.

Physalis somnifera.—P. perennis, tomentosa; foliis ovatis integrerimis, floribus confertis brevissime pedicellatis, corollis campanulatis. *Willd. sp. pl. 1. p. 1019.*

Icon. *Cav. ic. 2. t. 103.*

In South Carolina. *Catesby.* 4. July. v. s. in *Herb. Sherard.* Flowers small, brownish-yellow.

Campanula nitida.—C. glaberrima; foliis radicalibus spatulato-oblongis coriaceis cartilagineo-crenatis, caulinis lanceolato-oblongis subintegerrimis, floribus racemosis, corollis campanulato rotatis. *Willd. sp. pl. 1. p. 899.*

C. americana minor, flore cæruleo patulo. *Rob. ic.*

In North America. 4. July. v. s. in *Herb. Lambert. nec non Sherard.* About a span high; flowers middle size, fine blue. What part of America it is from is not known, but I suppose it to be Hudson's Bay. Misled by Michaux, this plant had been inserted as a synonym to *C. acuminata*, with which it has not the least connection.

Ribes laeviflorum.—R. inerme; foliis cordatis 5-lobis incisodentatis glabris, petiolis gracilibus, racemis laxis erectis longitudine foliorum, bracteis subulatis, pedicellis elongatis, calycibus campanulato-tubulatis, baccis globosis hispidis.

On the north-west coast. *Menzies.* ½. v. s. in *Herb. Banks.* The flowers appear to be yellow, about the size of *R. floridum*.

Ribes speciosum.—R. sub gemmis 3-plicato-aculeatum; foliis cuneato-subrotundis inciso-crenatis glabris nervosis subtus pallidioribus, petiolis brevibus, pedunculis sub-3-floris foliis longioribus, pedicellis germinibusque glanduloso-pilosis, calycibus tubulatis pedicellis longioribus, staminibus longissime exertis.

On the north-west coast. *Menzies.* ½. v. s. in *Herb. Banks.* The younger branches are hispid; the flowers

p. 157.

p. 159.

p. 163.

p. 165.

large, purple. The leaves are small, and have some resemblance to those of *Thalictrum*.

p. 165.

Ribes Menziesii.—R. sub gemmis 3-plicato-aculeatum; ramis hispidissimis, foliis basi truncatis sub-5-lobis inciso-dentatis, lobis lateralibus brevioribus, subtus tomentosis, pedunculis subbifloris foliis subaequantibus, calycibus tubulosis: limbo patente, staminibus longitudine calycis, stylo exerto, baccis globosis aculeatis.

On the north-west coast, near Fort Trinidad. *Menzies*.
h. v. s. in *Herb. Banks*. The flowers of the size and colour of *R. sanguineum*.

p. 173.

Viola asarifolia.—V. acaulis, pubescens; foliis dilatato-re-niformibus acutis circumcirca crenato-dentatis basi flabellatim in petiolum attenuatis, pedunculis foliis multo brevioribus.

Viola caroliniana maxima, foliis cucumerinis, floribus ad radicem pediculis brevibus natis. *Herb. Sherard*, In low rich woods: Virginia to Carolina. *Catesby*. 4. May. v. s. in *Herb. Sherard*. I have seen this species several times in Virginia, but generally without flowers; which has been the reason that no specimen was in my collection.

p. 190.

Hydrocotyle amligua.—H. folio radicali composito, caule subaphyllo 2-3-umbellifero, umbellulis sub-3-floris, petalis lanceolatis.

In Upper Louisiana. *Bradbury*. v. s. in *Herb. Bradbury*. A very small plant; flowers in proportion large, white, with purple anthers; resembling *H. composita*.

p. 192.

Selinum acaule.—S. glabrum, acaule; foliis bipinnatis, foliolis linearibus superioribus confluentibus, scapis radicibus foliis brevioribus, umbellulis radii masculis.

In Upper Louisiana. *Bradbury*. 4. v. s. in *Herb. Bradbury*. Not above two or three inches high; the seeds resembling those of *Pastinaca sativa*.

p. 197.

Seseli divaricatum.—S. caule ramosissimo divaricato, foliis bipinnatifidis, laciniis lanceolatis incisis, involucris nullis, involucellis linearibus.

In Upper Louisiana. *Bradbury*. v. s. in *Herb. Bradbury*. Resembles *S. tortuosum* very much.

p. 223.

Allium cernuum.—A. scapo nudo tetragono umbellifero, foliis linearibus planiusculis, umbella cernua, staminibus simplicibus, germe sex-dentato. *Roth in Annals of bot. 2. p. 27.*

Icon
On t
At
sc

Smilaci
tis

Poly
H

In H
As
it
di

Helonia
di
lis
ot
Be

On t
in
N

Rumex
ve

In U
Bi

Smilax
co
lat

S. flo
m

In Sc
ra
mo
sti

Gaura
ce

di
G. co

In U
Al
sc

as

I

Oenoth
vir

- Icon. *Cat. bot. fasc. 2. t. 2. Bot. mag. 1324.*
On the mountains of Virginia and Carolina. ♀. July,
Aug. v. v. Flowers large, rose-coloured, sweet-scented ; the leaves have scarcely any scent.
- Smilacina stellata*, β. uniflora.—S. foliis subfalcato-lanceolatis glabriusculis, caule unifloro.
- Polygonatum monanthemum*, flore parvo erecto. *Sherard. Herb.*
In Hudson's Bay. *Tilden.* ♀. v. s. in *Herb. Sherard.*
As I have seen but one individual specimen, I consider it only as a variety, though it is probable it may be a distinct species.
- Helonias graminea*.—H. scapo folioso paniculato, racemis divaricatis, bracteolis ovatis pedicellis longioribus, petalis oblongis acuminatis, staminibus corolla triplo brevioribus, foliis linearibus canaliculatis subtus glaucis.—*Bot. mag. 1599.*
- On the mountains of Georgia. *Lyon.* ♀. Aug. v. s. in *Herb. Lyon.* This species has a strong affinity with *Nolina Georgiana.*
- Rumex venosus*.—R. valvulis maximis integerrimis reticulato-venosis, foliis lanceolatis integerrimis parallelo-venosis.
In Upper Louisiana. *Bradbury.* ♀. v. s. in *Herb. Bradbury.* The seeds resemble those of *R. vesicarius.*
- Smilax circidifolia*.—S. aculeata ; foliis inermibus orbiculato-cordatis acuminatis 5-nervibus utrinque glabris reticulato-venosis breviter petiolatis.
- S. floridana ; viticulus spinosis judaicæ arboris folio maximo cordato. *Pluk. amalth. 194.*
- In South Carolina. *Catesby.* ♂. v. s. s. fl. in *Herb. Sherard.* Though the description was taken from a specimen without flowers, I am convinced of its being a distinct species from any one described.
- Gaura coccinea*.—G. sericeo-pubescentes ; foliis linear-lanceolatis subdenticulatis, spica conferta, petalis longitidine calycis, stigmate subintegro.
G. coccinea. *Fraser. catal. 1813.*
In Upper Louisiana *Bradbury.* v. s. in *Herb. Bradbury.* About a span high ; flowers smaller than *G. biennis*, scarlet. In the Catalogue quoted this plant is marked as a perennial, but by the appearance of the specimen I should take it to be only annual.
- Oenothera albicaulis*.—O. tenuissime pubescens ; caule nervisque foliorum albidis, foliis pinnatifidis, lacinias diva-

p. 243.

p. 248.

p. 249.

p. 260.

p. 262.

ricatis, floribus subspicatis, germinibus sessilibus cylindraceis, staminibus corolla brevioribus, petalis obcordatis.

O. albicaulis. Fraser. catal. 1813.

In Upper Louisiana. *Bradbury. ♂. v. s. in Herb. Bradbury.* Not above four inches high; flowers large, white. This species is intermediate between *O. tetrapeta* and *O. sinuata*.

Oenothera Fraseri.—*O. glabriuscula*; caule inferne simplici, superne ramuloso, foliis ovatis petiolatis glandulosodenticulatis, racemis foliosis, capsulis pedicellatis obovatis 4-gonis.

In South Carolina. *Fraser. ♀. June, July. v. v. in Hortis.* This species approaches so near to *O. fruticosa* that it is difficult to define the distinction, though their other habits are so very different. The present species flowers a short time, the stems decay, and form immediately a tuft of large radical leaves. *O. fruticosa* flowers till late in autumn, and never shows its radical leaves in that manner. The differentia specifica of *O. fruticosa* will be *O. fruticosa*, tenue pubescens; caule a basi ramosa, divaricata, foliis sessilibus lanceolatis subtendatis acutis, capsulis pedicellatis oblongo-clavatis angulatis.

Oenothera macrocarpa.—*O. glabriuscula*; caule ramoso, foliis lanceolatis, petiolatis, distanter glandulosodenticalatis margine nervisque sericeo-albidis, petalis cum acumine obcordatis, capsulis ellipticis 4-alatis brevisime pedicellatis.

O. missouriensis. Bot. mag. 1592.

On the banks of the Mississippi, near St. Louis. ♂. July—Oct. *v. v. in Hort. Lambert.* This species exceeds in size and beauty of its flowers any other one known; they are of a bright yellow, very large, and open about five o'clock in the evening. The calyx is covered with a very fine and white silky down, and is spotted with purple. The figure in the Botanical Magazine is apparently taken from a very weak and starved plant: the flowers of those in possession of A. B. Lambert, Esq., were nearly six inches in diameter, and the tube nearly seven inches long. There is an inequality on the edges of the petals, but by no means can it be called serrulate, which term has been introduced in the differentia specifica, and strongly but erroneously expressed in the figure. The specific name given there is inapplicable, as it never was found any where else,

bus cylindricus obcordatus

in Herb.
ers large,
O. tetra-

simplici-
nduloso-
latis obo-

Hortis,
that it is
ther ha-
lowers a
liately a
vers till
s in that
osa will-
ramosa,
acutis,

'amoso,
identi-
is cum
brevis-

♂.
ies ex-
r one
, and
lyx is
and is
1 Ma-
tarved
Lam-
id the
uality
it be
n the
y ex-
there
else,

but near St. Louis, where Mr. Nuttall gathered the ripe fruits of it, specimens of which I have seen.

Oenothera cespitosa.—*O. acaulis*; foliis lanceolatis inciso-dentatis, capsulis oblongis sessilibus, calycis tubo longissimo, petalis bilobis distantibus. *Sims in Bot. mag.* 1593.

O. scapigera. *Fl. amer. sept.* 1. p. 263.

On the banks of the Missouri. *Lewis*. ♀. June. July. *v. v. in Hort. Fraser*; *v. s. specimen imperfectum in Herb. Lewis*. This singular species resembles *O. acaule* of Cavanilles. The flowers open in the evening, are white, changing to red, and of an agreeable scent. The erroneous description given in the place above quoted was owing to a very imperfect specimen in the Lewisanian collection.

Eriogonum pauciflorum.—*E. caule simplici basi folioso, umbellis paucifloris, floribus subsessilibus, foliis linearilanceolatis utrinque cano-tomentosis*.

In Upper Louisiana. *Bradbury*. ♀. *v. s. in Herb. Bradbury*. This is the smallest species, generally from two to three inches high; the flowers appear to be white.

Cactus viviparus.—*C. subrotundus multiplex, tuberculis cylindraceis barbatis supra sulca exaratis proliferis*.

C. viviparus. *Fraser. catal.* 1813.

In Upper Louisiana. ♂. *v. v. s. fl. in Hort. Lambert*. Resembles *C. mammillaris*; the flowers are large and red; the fruit the size of a grape, green and eatable. In the same collection are also the two other species mentioned in Fraser's Catalogue; but only seeing the plants without flowers, I could not define their characters.

Crataegus turbinata.—*C. inermis, glabra; foliis cuneato-ovatis incisis serratis, corymbis paucifloris, pedicellis brevibus, fructibus turbinatis*.

In Carolina and Virginia. ♂. June. *v. v. ; v. s. in Herb. Sherard*. This species approaches near to *C. spathulata*, but its fruit distinguishes it from all the rest of the American species.

Rosa lutescens.—*R. germinibus globosis pedunculisque glabris, ramulis hispido-spinosis, foliolis (7) glabris ovalibus acuminato-serratis, petiolis inermibus, floribus so-*

p. 263.

p. 277.

p. 327.

p. 336.

p. 345.

litariis, laciniis calycis lanceolatis cuspidatis, petalis ovalibus obtusissimis.

In Carolina. $\frac{1}{2}$. May, June. *v. v. in Hort. Lee.* Flowers white, with a faint tint of yellow.

p. 352. *Geum triflorum*.—G. pilosum; caule simplici subtrifloro, foliis radicalibus interrupte pinnatis, foliolis cuneatis in ciso-dentatis, petalis oblongis longitudine calycis, aristis longissimis villosis.

In Upper Louisiana. *Bradbury*. $\frac{1}{2}$. *v. s. in Herb. Bradbury.* The aristas of the seeds are resembling those of *Clematis*.

p. 356. *Potentilla arguta*.—P. erecta, simplex, pubescens; foliis pin-natis, foliolis ovatis argute dentatis, exterioribus majoribus, stipulis rhomboideis incisis, floribus terminibus conferto-corymbosis.

In Upper Louisiana. *Bradbury*. $\frac{1}{2}$. *v. s. in Herb. Bradbury.* Flowers yellow; leaves large.

p. 384. *Clematis Catesbyana*.—C. scandens, pubescens; foliis bi-ternatis, foliolis subcordatis trilobis, paniculis divaricato-dichotomis, petalis staminibus longioribus, floribus subdioicis.

In South Carolina. *Catesby*. $\frac{1}{2}$. *v. s. in Herb. She-rard.* Resembles *C. virginica*. The filaments of the female flowers are membranaceous, linear, and without anthers.

p. 385. *Clematis ovata*.—C. erecta, simplex; foliis ovatis sensim acutis brevi-petiolatis glabris utrinque reticulato-veno-sis, inferioribus subcordatis pedunculo terminali solita-rio unifloro, aristis plumosis longissimis.

In South Carolina. *Catesby*. $\frac{1}{2}$. *v. s. in Herb. She-rard.* Resembles *C. integrifolia*. In the Sherardian Herbarium the figure of *Pluk. mant. t. 379. f 4.* is quoted; but I consider that figure to be an imperfect specimen of a *Clematis* with compound leaves.

p. 395. *Ranunculus multifidus*.—R. foliis tripartito-multifidis: laciniis incisis approximatis, caule natante, foliolis calycis subrotundis petalis paulo brevioribus.

In Upper Louisiana. *Bradbury*. $\frac{1}{2}$. *v. s. in Herb. Bradbury.* Flowers yellow; resembles *R. fluviatilis*.

p. 424. SEYMERIA. *Cal.* campanulatus, 5-partitus: laciniis li-nearibus, æqualibus. *Cor.* campanulata. *Tubus calyce*

Seyn

G.
In

Seyn

In

Chel

Ir

Pen

, petalis
Flowers
trifloro,
eatis in-
is, aris-
. Brad-
hose of
liis pin-
us ma-
ermina-
. Brad-
iiis bi-
varica-
toribus
. She-
of the
ithout
ensim
veno-
solita-
She-
urdian
4. is
erfect
laci-
lycis
Terb.
ilis.
s li-
lyce

vix longior. *Limbus* 5-fidus : laciinis subæqualibus, oblongis, patentibus. *Filamenta* 4. brevia, fauce inserita. *Antheræ* erectæ, oblongæ, nudæ, longitudine corollæ. *Stylus* declinatus, longitudine staminum. *Stigma* simplex. *Caps.* subrotunda, 2-locularis, 2-valvis, apice dehiscens.

Seymeria tenuifolia.—S. glabriuscula, virgatim paniculata ; foliis setaceo-pinnatifidis, laciinis divisis.

Gerardia Cassioides. Vol. 2. p. 424.

In dry sandy woods of Carolina and Georgia. ♂. July, Aug. v. s. in *Herb. Sherard*. Flowers small, dark yellow.

Seymeria pectinata.—S. viscidio-pubescent, brachiatim ramosissima ; foliis pectinato-pinnatifidis, laciinis indivisis linearibus acutis.

In South Carolina. *Catesby*. ♂ July. v. s. in *Herb. Sherard*. Flowers the size and colour of the preceding. While writing the genus *Gerardia*, I strongly suspected *G. cassioides* to be a distinct genus, which already had been considered so by Walter and Gmelin. Having no material then at hand, I followed the authority of Michaux. But finding in the Sherardian Herbarium not only that species, but also a new one described here, I was convinced of its generic distinction, the character of which is well expressed in *Walt. fl. car.* 171. I have named this genus in memory of the late Henry Seymer, Esq. of Hanford, Dorsetshire, a gentleman much celebrated for his extensive collections in natural history, and for whom Dr. Solander had so great a regard as to dedicate the plant which afterwards was called *Solandra grandiflora* to his memory, as I have seen it named *Seymera* in the Herbarium of A. B. Lambert, Esq., in Dr. Solander's own handwriting.

Chelone Lyonii.—C. glabra, ramosa ; foliis petiolatis cordato-ovatis serratis, spicis terminalibus densifloris.

In Upper Carolina and Georgia. *Lyon*. ♀. July—Sept. v. s. in *Herb. Lambert*. This is a very fine and large species ; flowers purple.

Pentstemon erianthera.—P. caule pumilo pubescente, foliis oblongis acutis obsolete denticulatis subhirsutis, floribus racemosis, pedicellis plerumque solitariis, filamento sterili exerto apice revoluto longitudinaliter barbato, calycis foliolis linearibus hirsutissimis.

p. 427.

p. 428.

P. erianthera. *Fraser. catal.* 1813.

In Upper Louisiana. *Bradbury*. *4. v. s. in Herb. Bradbury*. Not above a span high; flowers large in proportion, of a deep purple; the hairs of the infertile anthera are long and golden yellow. It approaches near to *P. frutescens*.

Pentstemon angustifolia.—*P. caule* vix pubescente, foliis glabris longe linearibus integerrimis, floribus racemoso-paniculatis, filamento sterili reliquis breviore superne sub-hirsuto, calycis foliolis lanceolatis glabris.

P. angustifolium. *Fraser. catal.* 1813.

In Upper Louisiana. *Bradbury*. *4. v. s. in Herb. Bradbury*. Flowers resembling those of *P. pubescens*.

Pentstemon glabra.—*P. caule* foliisque glabris, foliis sub-amplexicaulibus ovato-oblongis integerrimis, floribus racemoso-paniculatis, filamento sterili nudo clavato apice retuso, calycis foliolis subrotundis acuminatis, lacinias corollæ subæqualibus rotundatis, antheris hirsutis.

In Upper Louisiana. *Bradbury*. *4. v. s. in Herb. Bradbury*. A fine and large species; flowers dark purple.

Pentstemon Bradburii.—*P. glaberrima*; foliis subamplexicaulibus ovato-oblongis integerrimis, superioribus subrotundis, floribus verticillatis, filamento sterili apice brevi barbato, calycis foliolis oblongis acutis.

P. grandiflorum. *Fraser. catal.* 1813.

This is the finest and largest of all the species; the flowers very large, and variable in their colour.

p. 431.

Castilleja sessiliflora.—*C. foliis* apice palmato-incisis, floribus sessilibus.

In Upper Louisiana. *Bradbury*. *4. v. s. in Herb. Bradbury*.

Caulis ramosus, teres, dense pubescens. *Folia* alterna, sessilia, utrinque pubescentia, apice in lacinias 3—5. divisa: lacinias linearibus, elongatis, acutis. *Flores* in apice ramorum axillares, solitarii, sessiles, coccinei? *Calyx* tubulosus, basi ventricosus, 4-fidus. *Corolla* ringens. *Labium* superiorius acutum? *inferius* trifidum: lacinias linearibus, acutis, labio superiore vix brevioribus. *Stamina* fauci corollæ inserta, filiformia, longitude labii superioris, glabra. *Antheræ* lineares. *Stigma* exertum.

This
wi

Cleome
tis
su
lis

In U
Al
po

Polygao
mi
M
P. cy
In sw
Fl

Vicia st
bu
fol
sis

In U
Br
Cauli
lio
utr
bas
set:
Ra
Leg

Ervum
foli
4-sj
obt

Astrag
In Up
v. s

Caulis
seri
bric
nate
teæ
celle
bidi
briæ

1 Herb.
large in
infertile
soil near

leaves gla-
oso-pa-
ne sub-

Herb.
escens.

3 sub-
oribus
lavato
is, la-
s hir-

Herb.
dark

olexi-
sub-
apice

low-

lori-
erb.

rna,
di-
in
ei?
olla
m :
ori-
tu-
ig-

This highly interesting plant has some distant affinity
with *C. fissifolia*. *Smith ic. ined. t. 40.*

Cleome pinnata.—*C. subpubescens*, hexandra; foliis pinnatis, foliolis lanceolatis inferne attenuatis rarer dentatis, superioribus confluentibus, floribus racemosis, pedicellis aggregatis,

In Upper Louisiana. *Bradbury. v. s. in Herb. Bradbury.*
About two feet high; flowers of a bright yellow; the pods are on very long pedicels.

Polygala corymlosa.—*P. caule* inferne simplicissimo, summitate ramosa corymbose multifloro, foliis linearibus.—*Mich. fl. amer. 2. p. 54.*

P. cymosa. *Walt. fl. car. 179.*
In swamps: Carolina to Florida. 24. July, Aug. *v. s.*
Flowers yellow, when dried green.

Vicia stipulacea.—*V. racemis* subquadrifloris folio longioribus, stipulis foliformibus basi semisagittatis integris, foliolis 8—10. lineari-lanceolatis utrinque acutis nervosis glabris.

In Upper Louisiana. *Bradbury. 24. v. s. in Herb. Bradbury.*

Caulis subtetragonus, striatus, simplex, vix pedalis. *Mola* 8—10. opposita lineari-lanceolata acuta, glabra utrinque elevato-nervosa. *Stipulae* longitudine foliorum, basi semisagittatae, auriculis acutis. *Cirrus* obsoletus, setaceus. *Pedunculi* axillares folio duplo longiores. *Racemi* plerumque 4-flori. *Flores* magni, purpurei. *Legumen* glabrum.

Ervum multiflorum.—*E. racemis* pedunculatis multifloris folio longioribus, leguminibus lanceolatis glabris sub-4-spermis, stipulis ovatis acuminatis, foliolis linearibus obtusis, cirrhi nullis, caule paniculato.

Astragalus tenellus. *Vol. 2. p. 473.*
In Upper Louisiana. *M. Lewis and Bradbury. 24. Aug. v. s. in Herb. Lewis. nec non Bradbury.*

Caulis erectus, paniculato-ramosus, teres. *Ramuli* juniores sericei. *Foliola* plurima, alterna linearia, obtusa, glabriuscula. *Cirri* nulli. *Stipulae* ovatae longe acuminatae. *Racemi* axillares, elongati, pedunculati. *Bracteae* minutae, ovatae, acuminatae, vix longitudine pedicellorum. *Pedicelli* flore breviores. *Flores* parvi albidii, vexillo oblongo. *Calyx* sericeus, corolla triplobrevior, dentibus subulatis. *Stylus* filiformis, glaber,

p. 441.

p. 446.

p. 472.

SUPPLEMENTUM.

inflexus. Stigma acutum, glabrum. Legumina pendula, oblongo-lanceolata, glabra, 3-4-sperma.

Of this plant I had an imperfect specimen in the Lewisian Herbarium, together with some pods of an *Astragalus*, which led me into the error of placing it under that genus; but having since seen fine specimens in the collection of Mr. Bradbury, I was enabled to correct this error.

Astragalus racemosus.—A. caulescens, erectus, subpubescens; foliolis (21) lanceolatis superne sensim minoribus, stipulis ovatis acuminatis, spicis racemosis pedunculatis folio longioribus, densifloris, bracteis linearibus pedicellis brevioribus, calycibus glabriusculis, dentibus subulatis.

In Upper Louisiana. *Bradbury. ♀. v. s. in Herb. Bradbury.* Flower white. Resembles *A. carolinianus*.

Astragalus carnosus.—A. caulescens, diffusus, albido-sericeus, foliolis (21) ellipticis supra glabris, subtus sericeis, stipulis subulatis, spicis subsessilis, leguminibus globosis inflatis carnosis.

In Upper Louisiana. *Bradbury. v. s. in Herb. Bradbury.* The fruit resembles that of *A. psydodes*. Flowers appear to have been white. This is probably the *A. crassicarpus* *Fraser. catal. 1813*.

p. 473.

Astragalus triphyllus.—A. acaulis, argenteus, foliis ternatis, foliolis sessilis lanceolatis, scapo nullo, leguminibus sessilis oblongis.

In Upper Louisiana. *Bradbury. ♀. v. s. in Herb. Bradbury.* A most singular species, the leaves of a shining silver colour; the flowers were too much decayed when I examined the specimens.

Oxytropis Lambertii.—O. acaulis, sericeo-pilosa; foliolis (19) lanceolato-ellipticis utrinque acutis, scapis folia aequantibus, spicis capitatis, bracteis linearibus longitudine calycis.

On the Missouri. *Bradbury. ♀. Aug. Sept. v. v. in Hort. Lambert.* Flowers a beautiful purple. It approaches near to *O. argentata* and *setosa*.

p. 474.

Dalea aurea.—D. sericeo-villosa; spicis oblongis terminibus solitariis, bracteis oblongis acuminatis calycis longitudine ciliatis, calycibus villosissimis, foliis sub-3-jugis oblongis obovatisque obtusis.

D. at
In U
Bi
sp

Dalea l
mo
flo
ca
fol

D. ei
In U
Br
flo
sill
the

Psoralea
lis
rib
bre
In U
Br
pur

Psoralea
nat
dui
bra
In U
Br
Th
ral

Psoralea
liol
den
con
P. ma
In U
Br
escu

Cytisus
ere
long
obl

- D. aurea. *Fraser. catal.* 1813.
In Upper Louisiana. *Bradbury.* ♀. v. s. in *Herb.*
Bradbury. Flowers bright yellow, in oblong hairy spikes.
- Dalea laxiflora.*—D. glabriuscula; caule superne ramosissimo, ramis laxis elongatis, spicis paniculatis elongatis, floribus remotiusculis, bracteis subrotundis mucronatis carinatis, floribus enneandris, calycibus sericeo-villosis, foliis sub-4-jugis linearibus.
D. eneandra. *Fraser. catal.* 1813.
In Upper Louisiana. *Bradbury.* ♀. v. s. in *Herb.*
Bradbury. This is the tallest of the genus. The white flowers are in narrow spikes, and have remarkable silky calyces, which form a fine contrast with the rest of the plant, being entirely destitute of hair.
- Psoralea elliptica.*—P. subpubescens; foliis ternatis: foliolis ellipticis obtusis, pedunculis axillaribus folio longioribus, spicis conferto-paucifloris, dentibus calycis abbreviatis.
In Upper Louisiana. *Bradbury.* ♀. v. s. in *Herb.*
Bradbury. Flowers small, white with a tinge of purple.
- Psoralea longifolia.*—P. undique sericeo-villosa; foliis ternatis: foliolis linearibus elongatis, spicis axillaribus penduculatis laxifloris folio brevioribus, dentibus calycis bracteisque subulatis.
In Upper Louisiana. *Bradbury.* ♀. v. s. in *Herb.*
Bradbury. Flowers middle-sized, apparently white. The leaves have none of those glandular dots so general in this genus.
- Psoralea cuspidata.*—P. villosa; foliis digitato-quinatis, foliolis obovatis mucronatis integerrimis, spicis axillaribus densifloris, lacinias calycis bracteisque ovatis cuspidatis conspicue punctatis.
P. macrorhiza. *Fraser. catal.* 1813.
In Upper Louisiana. *Bradbury.* ♀. v. s. in *Herb.*
Bradbury. This species has some resemblance to *P. esculenta.*
- Cytisus rhombifolius.*—C. pubescens; racemis terminalibus erectis, leguminibus falcatis subarticulatis, foliolis oblongo-rhomboideis obtusis, stipulis rotundato-ovatis obliquis.

p. 475.

p. 476.

p. 489.

pli
cal
E. un
In U
Br
ray
Galardia
spa
In U
Bre
ply
Iva axil
pos
bus
liol
In U
Bra
Ambrosia
bipi
pan
In Sou
Abc
Orchis h
bell
den
Haber
p. 1
In'No
unk
fusc
Aristoloc
peti
riis
liter
On the
v. v
Alnus gle
subt
tis.-
In Ne
vani
is ki
sem
VOL. II.

C. rhombifolius. Fraser, catal. 1813.

In Upper Louisiana. *Bradbury. v. s. in Herb. Bradbury.* The stipulae are large and foliaceous; the specimen I have seen was in fruit, and Mr. Bradbury has seen only one plant in flower.

p. 505.

Troximum glaucum.—T. scapo unifloro glabriuscule, foliis lineari-lanceolatis planis integerrimis utrinque glaucis, calycinis foliolis imbricatis acutis pubescentibus.—Vol. 2. p. 505.

Troximum cuspidatum.—T. scapo unifloro superne subtomoso, foliis linearibus margine undato tomentosis, calycinis foliolis imbricatis cuspidatis glabris.

In Upper Louisiana. *Bradbury. Ʌ. v. s. in Herb. Bradbury.* Flowers large, yellow. Resembles the preceding.

p. 519.

Hymenopappus scabiosæus.—H. candicanti-lanuginosus; foliis profunde pinnatifidis, laciniis lineari-oblongis subdentalis, floribus simpliciter corymbosis, calyce sub-10-phyllo flosculis majore, seminibus pubescentibus.—Vol. 2. p. 519.

Hymenopappus tenuifolius.—H. candicanti-lanuginosus; caule corymboso-ramoso, foliis subbipinnatifidis, laciniis anguste linearibus integris, floribus composite corymbosis, pedicellis confertis, calyce sub-6-phyllo flosculis breviore, seminibus villosissimis.

In Upper Louisiana. *Bradbury. ♂. v. s. in Herb. Bradbury.* At first sight I considered this to be the same with the preceding species; but comparing the figure and description in the *Journal d'Histoire Naturelle*, vol. 1. p. 16. t. 1. with my specimen, I found them materially distinct. The flowers are white, and not half the size of the former.

p. 521.

Artemisia dracunculoides.—A. glabra; ramis nutantibus, foliis longo-linearibus subfalcatis, radicalibus trifidis, racemis foliosis, pedicellis brevibus, floribus erectis, calyculis subglobosis paucifloris, foliolis scariosis.

A. nutans. Fraser, catal. 1813.

In Upper Louisiana. *Bradbury. Ʌ. v. s. in Herb. Bradbury.* From three to four feet high; flowers very small.

p. 532.

Erigeron hirsutum.—E. undique hirsutissimum; caule sim-

- plici unifloro, foliis linearibus integerimis enervibus,
calycibus hirsutis
E. uniflorum. Willd. sp. pl. 3. p. 1960 ?
In Upper Louisiana. Bradbury. ♀. v. s. in Herb.
Bradbury. Not above five inches high; flowers large,
ray white and long.
- Galardia acaulis*.—G. pilosa; scapo unifloro, foliis linearis-
spathulatis, paleis pappi (5—7) ovatis brevi-aristatis.
In Upper Louisiana. Bradbury. ♀. v. s. in Herb.
Bradbury. Not above a span high; flower small, sim-
ply yellow.
- Iva axillaris*.—I. herbacea, subpilosa; foliis plerumque op-
positis linearis-lanceolatis integerimis, floribus axillari-
bus solitariis nutantibus, calycibus simpliciusculis, fo-
liolis ovatis.
In Upper Louisiana. Bradbury. ♂. v. s. in Herb.
Bradbury. About a span high; flowers white.
- Ambrosia hispida*.—A. undique canescens-hispida; foliis
bipinnatifidis, lacinias incisis, racemis terminalibus sub-
paniculatis.
In South Carolina. Catesby. ♂. v. s. in Herb. Sherard.
About a foot high; flowers larger than *A. trifida*.
- Orchis herbiola*.—O. cornu filiformi germine breviore, la-
bello oblongo obtuso basi utrinque dentato; palato uni-
dentato, bracteis flore longioribus.
Habenaria herbiola. Brown in Ait. kew. ed. 2. vol. 5.
p. 193.
In North America. Fothergill. ♀. +. This species,
unknown to me, appears to approach near to *Orchis*
fuscescens.
- Aristolochia tomentosa*.—A. caule volubili frutescente, foliis
petiolatis cordatis subtus tomentosis, pedunculus solita-
riis ebracteatis, corolla tubo retorto; limbo subaequaliter
trifido: ore hiante. Sims in bot. mag. 1369.
On the mountains of South Carolina. Lyon. ♀. June.
v. v. Resembles *A. Siphon*. Flowers yellow.
- Alnus glauca* —A. foliis oblongis acutis duplicato-serratis
subitus glaucis, axillis venarum nudis, stipulis lanceola-
tis.—Mich. arb. 3. p. 322. t. 4.
In New England and on the high mountains of Pensyl-
vania. ♀. March, April. v. v. This shrub, which
is known by the name of *Black Alder*, has a strong re-
semblance to *A. incana*.

pag.
26
30

ADDENDA ET CORRIGENDA.

VOL. I.

pag.		42
		43
		61
		62
		72
		95
		96
		98
		102
1	lin. 4 loco <i>Monoicum</i> lege <i>Monoica</i> . This error occurs in this and the next table of genera.	
15	adde <i>Monoicum</i> .	
22	adde <i>Monoica</i> .	
23	adde <i>Monoica</i> .	
3	7 <i>Hippuris vulgaris</i> . In the Sherardian Herbarium are specimens collected at Hudson's Bay which I compared with the European <i>Hippuris</i> , and found them entirely distinct. I rather suspect this plant ought to be placed among the <i>Filices</i> .	115
6	5 adde <i>Dioica</i> .	
7	12 adde <i>Dioica</i> .	
	13 adde <i>Dioica</i> .	
	15 adde <i>Monoica</i> .	
	7 ab infer, adde <i>Mich. arb.</i> 3. p. 50. t. 6.	128
8	17 ab infer, lege <i>Mich. arb.</i> 3. p. 122. t. 12.	138
	11 ab infer, lege <i>Mich. arb.</i> 3. p. 118. t. 11.	140
	3 et 5 ab infer, delendæ sunt.	
9	5 delenda est.	
	6 lege <i>Mich. arb.</i> 3. p. 106. t. 8.	158
	14 dele <i>Mich. arb.</i> ejus loco <i>Catesl. car.</i> 1. t. 80.	159
	19 dele <i>Catesb. car.</i> t. 80. et lege <i>Mich. arb.</i> 3 p. 128. t. 13.	163
	28 lege <i>Mich. arb.</i> 3. p. 112. t. 9.	164
	3 ab infer, lege <i>F. viridis. Mich. arb.</i> 3. p. 115. t. 10.	165
10	5 adde <i>Mich. arb.</i> 3. p. 217. t. 6.	166
10	Monarda Kalmiana, adde Icon. t. 1.	167
17	13 post Flowers adde pale.	169
21	Ceratiola ericoides, adde Icon. t. 13.	175
24	25 Rynchospora lege Rhynchospora.	177

- DA.
rror oce-
ra.

Herba-
n's Bay
Hippu-
I rather
among

80.
rb. 3

. 115.

169
- pag. lin.
 26 30 dele genus ANTHISTIRIA.
 30 Iris cuprea, *adde* I. fulva. *Bot. mag.* 1496
 Iris verna, *conf.* *Bot. mag.* 1393. *observ.*
 Iris prismatica, *adde* *Bot. mag.* 1504.
 42 16 *loco foliis lege foliisque.*
 43 18 *loco E lege C.*
 61 9 et 14 *loco Dilepycum lege Dilepyrum.*
 62 Milium amphicarpon *adde* Icon. *Tab. 2.*
 72 6 *ab infer. loco A lege S.*
 95 23 *dete 2. ante biloculari.*
 96 7 *ab infer. loco involucro lege involucris.*
 98 8 *ad marginem loco data lege cordata.*
 ultima, loco caule lege scapo.
 102 *Oldenlandia glomerata.* Willdenow's and La-
 marck's synonym seem to belong to a very
 ambiguous species of this genus, or, as I
 strongly suspect, to *Ludwigia nitida*, as I
 have seen this plant in the Sherardian Herba-
 rium under the name and synonyms of *Olden-
 landia uniflora*.
 115 Planera aquatica, *adde* P. ulmifolia. *Mich. arb.*
 3. p. 283.
 126 18 *loco Gonolobus lege Gonolobium.*
 20 *loco Appendix lege Appendices.*
 inter 224 et 225 insere CEROPEGIA. *Folliculi 2.*
 erecti. Sem. plumosa. Cor. urceolatæ limbus
 connivens.
 128 inter 244 et 245 insere SISON. *Fructus ovatus,*
 striatus. Invol. sub-4-phylla.
 138 8 *ab infer. post palustris adde* β. inflata.
 140 Ipomoea Jalapa, *adde* Convolvulus Jalapa. *Bot.*
 mag. 1572.
 158 Pinkneya pubescens, *adde* *Mich. arb.* 2. p. 276.
 t. . . .
 159 11 et 12 delendæ sunt.
 163 Ribes resinosum, *adde* Icon. *Bot. mag.* 1583.
 164 7 *ab infer. adde* *Mich. fl. amer.* 1. p. 109.
 165 7 *ab infer. loco lacustris lege lacustre.*
 166 1 *loco subgeminis lege sub gemmis.*
 167 6 *adde* *Bot. mag.* 1479.
 175 ultima, ante *utrinque adde foliis.*
 175 Claytonia lanceolata, *adde* Icon. *Tab. 3.*
 177 Nyssa villosa, *adde* N. sylvatica. *Mich. arb.* 2.
 p. 260. *t. 21.*
 Nyssa biflora, *adde* *Mich. arb.* 2. p. 265. *t. 22.*

2 c 2

pag.	lin.	pag.
177	<i>Nyssa candidans</i> , <i>adde Mich. arb.</i> 2. <i>p.</i> 257. <i>t. 20.</i>	
178	<i>Nyssa tormentosa</i> , <i>adde Mich. arb.</i> 2. <i>p.</i> 252. <i>t. .</i>	
181	1 loco 1187 lege 1181.	
186	1 <i>adde Bot. mag.</i> 1551.	
187	20 loco unifloribus lege unifloris.	
199	ultima, <i>adde Mich. arb.</i> 3. <i>p.</i> 269. <i>t. .</i>	
200	<i>Ulmus fulva</i> , <i>adde U. rubra. Mich. arb.</i> 3. <i>p.</i> 278. <i>t. .</i>	242
	<i>Ulmus alata</i> , <i>adde Mich. arb.</i> 3. <i>p.</i> 275. <i>t. .</i>	
	<i>Celtis occidentalis</i> , <i>adde Mich. arb.</i> 3. <i>p.</i> 226. <i>t. 8.</i>	
	<i>Celtis crassifolia</i> , <i>adde Mich. arb.</i> 3. <i>p.</i> 228. <i>t. 9.</i>	
206	206 17 <i>locu ramosissima lege ramosissimo.</i>	
219	<i>Berberis aquifolium</i> , <i>adde Icon. Tab.</i> 4.	
	<i>Berberis nervosa</i> , <i>adde Icon. Tab.</i> 5.	
221	<i>Gleditschia triacanthos</i> , <i>adde Mich. arb.</i> 3. <i>p.</i> 164. <i>t. .</i>	242
	<i>Gleditschia monosperma</i> , <i>adde Mich. arb.</i> 3. <i>p.</i> 169. <i>t. .</i>	243
223	223 <i>Allium angulosum</i> , <i>adde A. stellatum. Bot. mag.</i> 1576.	255
224	224 <i>Conostylis americana</i> , <i>adde Icon. Tab.</i> 6. <i>Lophiola aurea. Bot. mag.</i> 1596. In this figure the flowers are more than twice the size of those in its native state.	259
226	226 <i>Phalangium Quamash</i> , <i>adde Scilla esculenta.</i> <i>Bot. mag.</i> 1574. Having not seen the living plant when I made my description, I was misled by Mr. Lewis in describing one of the petals to be dependent and the rest ascendent, which is not the case.	264
228	228 <i>Lilium pudicum</i> , <i>adde Icon. Tab.</i> 8.	265.
232	232 <i>Streptopus roseus</i> , <i>adde Uvularia rosea. Bot.</i> <i>mag.</i> 1489.	266
	<i>Streptopus lanuginosus</i> , <i>adde Uvularia lanuginosa.</i> <i>Bot. mag.</i> 1490.	
240	240 <i>Chamærops Palmetto</i> , <i>adde Mich. arb.</i> 2. <i>p.</i> 186. <i>t. .</i>	267
	<i>Nolina georgiana</i> , <i>adde Asphodelus minor albus</i> ex Provincia mariana, <i>Pluk. mant.</i> 29. <i>t. 342.</i> <i>f. 1.</i> (pars paniculæ), secundum <i>Herb. Sherard.</i> Very fine specimens of this rare plant I found in the Sherardian Herbarium collected by M. Catesby on the Savannah river. It re- sembles <i>Helonias graminea</i> , <i>Bot. mag.</i> 1599.	268
		282

57. t. 20.
252. t. .
- arb. 3.
t....
p. 226.
28. t. 9.
4. 3. p.
arb. 3.
t. mag.
Lo-
; figure
size of
ulenta.
; living
I was
of the
ndent,

Bot.
ginosa.
; b. 2.
albus
. 342.
She-
plant
ected
It re-
1599.
- pag. lin.
- exceedingly, and I should have taken it to be the same if the pedicels were not solitary, and, as Mr. Kerr observes; the germ polyspermous, which in *Noline* is only 3-spermous.
- 242 Veratrum viride, *adde* *Helonias viridis*. *Bot. mag.* 1096.
Veratrum angustifolium.—Of this species I have seen a fine specimen in the Sherardian Herbarium, under the name *Phalangium canadense majus ramosum, Iridis folio*, by which it appeared to grow likewise in Canada. The panicle was monoic not dioic, as I observed it in Virginia, the seeds lanceolate, membranaceous.
- 242 14 *ab infer. loco legumine lege tegumine.*
 23 *loco thier lege their.*
- 255 *Æsculus flava, adde Mich. arb. 3. p. 238. t. .*
Æsculus glabra, adde Mich. arb. 3. p. 242.
- 259 4 *ab infer. adde nomen specificum, rosmarinifolium.*
- 263 dele *Oenothera scapigera. Ejus loco vid. Supple-*
mentum sub nomine Oenothera cæspitosa.
- 264 Menziesia globularis, *adde M. ferruginea* β .
Bot. mag. 1571. Though Dr. Sims considers this species only as a variety, I still think the distinctions given by A. Salisbury, Esq. are sufficient to keep it as a species.
- 265 *Diospyros virginiana, adde Mich. arb. 2. p. 195.*
t. 12.
- 266 *Acer rubrum, adde Mich. arb. 2. p. 210. t. 14.*
Acer dasycarpum, adde Mich. arb. 2. p. 205.
t. 13.
- 267 *Acer saccharinum, adde Mich. arb. 2. p. 218.*
t. 15.
- 268 *Acer nigrum, adde Mich. arb. 2. p. 238. t. 16.*
Acer striatum, adde Mich. arb. 2. p. 242. t. 17.
- 269 *Acer Negundo, adde Mich. arb. 2. p. 247. t. 18.*
inter AGROSTEMMA et OXALIS, adde 394.
- LYCHNIS. *Cal. 1-phyllo, oblongus, lœvis.*
Pet. 5. unguiculata, limbo subbifido. Caps.
1-5-locularis.
- 282 6 *loco 394 lege 395.*
 10 *loco 395 lege 396.*

ADDENDA ET CORRIGENDA.

pag.	lin.		pag.
	12	loco 396 lege 397.	
	15	loco 397 lege 398.	
283		<i>Gaultheria serpyllifolia</i> , <i>adde Tab. 13.</i>	
284		<i>Gaultheria Shallon</i> , <i>adde Tab. 12.</i>	
289		<i>Vaccinium nitidum</i> , <i>adde Bot. mag. 1550.</i>	
293		<i>Andromeda floribunda</i> , <i>adde Bot. mag. 1566.</i>	
		In the extensive collection of the Sherardian Herbarium I met with most of the species of <i>Vaccinium</i> and <i>Andromeda</i> described in the present work; but meeting with none that were new, convinced me of the idea that this extensive family is nearly fully described here.	333
294		<i>Andromeda mariana</i> , <i>adde var. ovalis. Bot. mag. 1579.</i>	339
295		<i>Andromeda arborea</i> , <i>adde Mich. arb. 3. p. 222. t. 7.</i>	340
297		<i>Kalmia latifolia</i> , <i>adde Mich. arb. 3. p. 147. t. 5.</i>	344
298		<i>Rhododendron maximum</i> , <i>adde Mich. arb. 3. p. 144. t. 4.</i>	
		<i>Rhododendron cataubense</i> . I have had an opportunity of seeing this species in flower at Messrs. Lee and Kennedy's; the plant was not above a foot high; the flowers rose-coloured, similar to those of <i>R. maximum</i> ; but having scarcely any marks or spots on the upper part of the corolla.	348
302		<i>Mylocarium ligustrinum</i> , <i>adde Icon. Tab. 14.</i>	350
		The original drawing of this plant was taken by me at Messrs. Bartrams' near Philadelphia.	
303		<i>Monotropa morisoniana</i> . I had the pleasure of seeing, in the Oxford Herbarium, the original specimen from which the figure in Morrison's History has been taken, and which so erroneously has been quoted under <i>Monotropa uniflora</i> .	362
304		<i>Gymnocladus canadensis</i> , <i>adde Mich. arb. 2. p. 272. t. 23.</i>	363
308		<i>Podalyria mollis</i> , <i>adde v. s. in Herb. Lambert.</i>	390
309		<i>Virgilia lutea</i> , <i>adde Mich. arb. 3. p. 266. t. 3.</i>	404
313		<i>Tiarella cordifolia</i> , <i>adde Bot. mag. 1589.</i>	
327		BARTONIA. Since writing the above article I had an opportunity of seeing both species there described in a living state; the first in the garden of A. B. Lambert, Esq. at Eoyton,	411
			423
			424
			427
			428

VOL

pag. lin.

where it flowered in very high perfection and formed seeds; the other at Messrs. Frasers', Sloane Square. From the latter I have taken the following description.

Bartonia nuda.—*Caulis ramosus, angulatus, glanduloso-pubescent. Folia oblonga, sessilia, basi attenuata, repando-dentata, utrinque scabra, viridia. Inflorescentia B. ornatae. Flores duplo minores, lutescentes, germine nudo, angusto. Petala 10. angusto-lanceolata. Filamenta exteriora longiora, plerumque castrata et petaloidea.*

- 333 *Tigarea tridentata, adde Icon. Tab. 15.*
 339 8 *ab infer. loco linearibus lege linearis*
 1 *ab infer. adde Mespilus arborea. Mich. arb. 3. p. 68. t. 11.*
 340 *Pyrus coronaria, adde Malus coronaria. Mich. arb. 3. p. 65. t. 10.*
 344 *Rosá blanda. Differentia specifica lege R. germinibus globosis pedunculisque hispidis, ramis aculeato-hispidae, foliolis &c.*
 348 *Rubus spectabilis, adde Icon. Tab. 16.*
 350 *Dalibarda fragaroides, adde Bot. mag. 1567.*

VOL. II.

- 362 *Tilia glabra, adde T. americana. Mich. arb. 3. p. 311. t. 1.*
 363 *Tilia pubescens, adde Mich. arb. 3. p. 317. t. 3.*
Tilia heterophylla, adde T. alba. Mich. arb. 3. p. 315. t. 2.
 365 *Talinum teretifolium, adde Pluk. alm. t. 223. f. 2. secundum Herb. Sherard.*
 390 *Caltha flabellifolia, adde Icon. Tab. 17.*
 404 *adde CASTILLEJA. Cal. tubulosus unilabiatus. Cor. bilabiata, labio inferiore trifido, glandulosa duabus inter lacinias. Caps. bilocularis.*
 411 *Prunella pensylvanica, adde Icon. Mart. cent. t. 5.*
 423 *Gerardia fruticosa, adde Icon. Tab. 18.*
Gerardia quercifolia, adde Icon. Tab. 19.
 424 *Gerardia cassiodes dele. Ejus loco vide Supplémentum sub nomine Seymeria.*
 427 *Mimulus Lewisii, adde Icon. Tab. 20.*
Pentstemon laevigata, adde Bot. mag. 1425.
Pentstemon pubescens, adde Bot. mag. 1424.

ADDENDA ET CORRIGENDA.

pag.	lin.		pag.
441	13	<i>loco glabrum lege glabra.</i>	
451		<i>Hopea tinctoria, adde Mich. arb. 3. p. 61. t. 9.</i>	
		<i>Gordonia Lasianthus, adde Mich. arb. 3. p. 131.</i>	
		<i>t. 1.</i>	
		<i>Gordonia pubescens, adde Mich. arb. 3. p. 135.</i>	
		<i>t. 2</i>	
460	inter 591 et 592,	<i>adde CYTISUS. Cal. bilabiatus :</i>	575
		<i>labio superiore bidentato. Carina erecta.</i>	
		<i>Legum. basi attenuatum, 1-loculare, compressum. Stam. monodelpha.</i>	
		<i>inter 587 et 588, adde ERVUM Cal. 5-dentatus, dentibus subulatis. Stigma simplex, incurvum. Legum. compressum, oligospermum. Stam. diadelpha.</i>	
465		<i>Polygala Seneca, adde var. γ tenuifolia—foliis linearibus, spicis confertifloris.</i>	560
		<i>On the Missouri. Lewis and Bradbury. v. s.</i>	
468		<i>Lupinus villosus, adde Icon. Tab. 21.</i>	585
471		<i>Vicia americana, adde v. s. in Herb. Bradbury. var. foliis linearibus.</i>	
475		<i>Psoralea esculenta, adde Icon. Tab. 22.</i>	587
479		<i>Lupinaster macrocephalus, adde Icon. Tab. 23.</i>	
480		<i>Glycyrrhiza lepidota, loco sericeo-villosis lege utrinque lepidotis.</i>	
487		<i>Robinia pseudacacia, adde Mich. arb. 3. p. 245. t. 1.</i>	588
488		<i>Robinia viscosa, adde Mich. arb. 3. p. 262. t. 2.</i>	
499		<i>Prenanthes Serpentaria, adde Icon. Tab. 24.</i>	
507		<i>Liatris macrostachya, adde Bot. mag. 1411.</i>	
529		<i>Senecio pauperculus, adde v. s. in Herb. Bradbury.</i>	589
559		3 <i>loco t. 4. lege vol. 5.</i>	
564		<i>Amellus villosus, adde Sideranthus integrifolius. Fraser. catal. 1813. v. s. in Herb. Bradbury.</i>	590
		<i>Amellus spinulosus, adde Sideranthus pinnatifidus. Fraser. catal. 1813. v. s. in Herb. Bradbury.</i>	592
571		7 <i>loco acutis lege acuminatis.</i>	
572		<i>Galardia bicolor, adde G. bicolor; caule ramoso, foliis caulinis lanceolatis subdentatis, radicalibus dentato-pinnatifidis, paleis pappi integerrimis aristatis. Sims in Bot. mag. 1002. Virginia grandiflora. Fraser. catal. 1813.</i>	594
		<i>This elegant plant, which has been proved to be a perennial, though it was always considered</i>	595

	pag.	lin.	
61. t. 9. p. 131.			an annual, is very variable in its size, colour, and number of florets in the ray. Native specimens collected by Mr. Bradbury are covered with rigid hairs, and have but one flower. Cultivated plants raised from seeds of the same species are more smooth, branch more or less; the leaves are deeper divided, and produce several flowers double the size.
p. 135.			
ibiatus : erecta. ompres-			
entatus, . incur- ermum.	575		Rudbeckia columnaris, <i>adde</i> R. columnaris, foliis pinnatifidis: foliolis lanceolatis, exterioribus ordinatim majoribus, receptaculo cylindraceo elongato. <i>Sims in Bot. mag.</i> 1601. This species is not above a foot high, and appears to be annual.
—foliis	580		Iva ciliata, <i>adde</i> Iva annua. <i>Mich. fl. amer.</i> 2. p. 184.
v. s.	585		Orchis ciliaris, <i>adde</i> Habenaria ciliaris. <i>Brown in Ait. kew. ed. 2. vol. 5.</i> p. 194.
adbury.			Orchis cristata, <i>adde</i> Habenaria cristata. <i>Brown in Ait. kew. ed. 2. vol. 5.</i> p. 194.
ab. 23. sis lege	587		Orchis viridis, <i>adde</i> Habenaria viridis. <i>Brown in Ait. kew. ed. 2. vol. 5.</i> p. 192.
p. 245.	588		Orchis bracteata, <i>adde</i> Habenaria bracteata. <i>Brown in Ait. kew. ed. 2. vol. 5.</i> p. 192.
2. t. 2. . 24. 1. . Brad-			Orchis hyperborea, <i>adde</i> Habenaria hyperborea. <i>Brown in Ait. kew. ed. 2. vol. 5.</i> p. 193.
ifolius. dbury. matifi- Herb.	589		Orchis fimbriata, <i>adde</i> Habenaria fimbriata. <i>Brown in Ait. kew. ed. 2. vol. 5.</i> p. 193.
ramo- is, ra- pappi 1002.	590		Neottia cernua, <i>adde</i> Bot. mag. 1568.
to be idered			Neottia repens, <i>adde</i> Goodyera repens. <i>Brown in Ait. kew. ed. 2. vol. 5.</i> p. 198.
	592		Neottia pubescens, <i>adde</i> Goodyera pubescens. <i>Brown in Ait. kew. ed. 2. vol. 5.</i> p. 198.
			Arethusa divaricata, <i>adde</i> Pogonia divaricata. <i>Brown in Ait. kew. ed. 2. vol. 5.</i> p. 203.
			Cymbidium pulchellum, <i>adde</i> Calopogon pulchellus. <i>Brown in Ait. kew. ed. 2. vol. 5.</i> p. 204.
			Cymbidium verecundum, <i>adde</i> Bletia verecunda. <i>Brown in Ait. kew. ed. 2. vol. 5.</i> p. 206. Bot. mag. 930.
	594		Cypripedium parviflorum, <i>adde</i> Bot. mag. 911.
	595		Cypripedium arietinum, <i>adde</i> Bot. mag. 1509

O
Fat
incl
Nat
F.R
16l.
frot
Ant
gus

T
BO
NA
NVA
Fell
post
whi

F
Par
Off

T
sups
to n
by I
to it
and

VALUABLE WORKS

IN

Natural History

PUBLISHED BY

WHITE, COCHRANE, AND Co.,

HORACE'S HEAD, FLEET STREET.

ORIENTAL MEMOIRS; selected and abridged from a Series of Familiar Letters written during Seventeen Years' Residence in India; including Observations on Parts of Africa and South America; and a Narrative of Occurrences in four India Voyages. By JAMES FORBES, F.R.S. &c. Elegantly printed by BENSLEY, in 4 vols. royal 4to. Price 16l. 16s. in boards. With a Portrait of the Author, and 93 Engravings from original Drawings, illustrative of the Natural History, Scenery, Antiquities, and Costume, and including Portraits of several distinguished Characters.

The Subjects of Natural History have been engraved and coloured by HOOKER in a style of great accuracy and beauty; and the Landscapes and other Subjects by C. HEATH, ANGUS, STOKE, GREIG, &c.

THE NATURAL HISTORY AND ANTIQUITIES OF SELBORNE, in the County of Southampton. To which are added: The NATURALIST'S CALENDAR; OBSERVATIONS ON VARIOUS PARTS OF NATURE; and POEMS. By the late Rev. GILBERT WHITE, formerly Fellow of Oriel College, Oxford. A new Edition, elegantly printed in post 4to. by BENSLEY, with Additions, and 12 Engravings, several of which are new. Price 2l. 12s. 6d. in boards.

Fifty Copies are printed on royal quarto paper, (with the copy of a curious old Painting, beautifully coloured,) and done up in 2 vols. Price 5l. 5s. in boards. Of these very few now remain.

To those who are acquainted with this Work, any recommendation of it is superfluous: to those who are not, the testimonies borne to its merits, both as to matter and style, by some of the most eminent Writers of the present day, by DR. WHITAKER, DR. AIKIN, MR. EDGWORTH, &c. may offer some inducement to its perusal. Without exaggeration, it may be said to stand at the head of, and to serve as a model for, all Works of a similar description.

Valuable Works lately published by

At the same time is published, handsomely printed in 2 vols. 8vo. with Four Engravings, price 16s. in boards, a new Edition of

THE NATURAL HISTORY OF SELBORNE: To which are added, THE NATURALIST'S CALENDAR; MISCELLANEOUS OBSERVATIONS; and POEMS.

Extract from the Advertisement.—“The NATURAL HISTORY OF SELBORNE has probably been supposed by many to be formed upon a more local and confined plan than it really is. In fact, the greater part of the Observations are applicable to all that portion of the island in which the Author resided, and were indeed made in various places. Almost the only matter absolutely local, is the account of the Antiquities of the Village of Selborne; and this seemed to stand so much apart, that, however well calculated to gratify the lovers of topographical studies, it was thought that its entire omission would be no loss to the Work, considered as a publication in Natural History.”—“The Editor flatters himself that, in its present form, it will prove an acceptable addition to the library of the Naturalist, and will in particular be useful in inspiring young persons, and those who pass their time in retirement, with a taste for the very pleasing branch of knowledge on which it treats.”

FAUNA ORCADENSIS; or, The Natural History of the Quadrupeds, Birds, Reptiles, and Fishes of Orkney and Shetland. By the Rev. GEORGE LOW, Minister of Birsay and Haray. From a Manuscript in the possession of WILLIAM ELFORD LEACH, M.D. F.L.S. &c. Elegantly printed in 1 vol. sm. 4to. Price 1l. 1s. in boards.

NATURAL HISTORY; General and Particular. By the COUNT DE BUFFON. Illustrated with above Six Hundred Copper-plates. The History of Man and Quadrupeds translated, with Notes and Observations, by WILLIAM SMELLIE, Member of the Antiquarian and Royal Societies of Edinburgh. A new Edition, carefully corrected, and considerably enlarged by many additional Articles, Notes, and Plates, and some Account of the Life of BUFFON, by WILLIAM WOOD, F.L.S. In 20 vols. 8vo. Price 12l. in boards.

AN ESSAY ON THE PHILOSOPHY, STUDY, AND USE OF NATURAL HISTORY. By CHARLES FOATHERGILL. Beautifully printed in foolscap 3vo. Price 8s. in boards.

LACHESIS LAPponica: or, A TOUR IN LAPLAND. Now first published from the original Manuscript Journal of the celebrated LINNÆUS. By JAMES EDWARD SMITH, M.D. F.R.S. &c. President of the Linnean Society. Handsomely printed in 2 vols. 8vo. with above 60 Wood Cuts from the extemporaneous Sketches of the illustrious Author. Price 1l. 1s. in boards.

“Although the preceding notices and extracts can convey to our readers a very inadequate idea of the quantity of minute details which has been pressed into this interesting diary, they may serve to furnish exemplifications of its nature and spirit. Whoever peruses the entire document with candour and attention, will be induced to regard it as the unaided and unpremeditated production of a student of medicine, in his 25th year, labouring under the daily privation of many comforts, and braving cold, tempest, and danger, in

8vo.

ded,
and

: has
split
were
l, is
med
s of
loss
itor
tion
ring
for

ru-
the
mu-
xe.

NT
he
ca-
val
in-
In

F
ly

st
c-
ie
o.
r.

a
d
s
i
l
e

White, Cochrane, and Co. Fleet Street.

the eager prosecution of natural science. In this Itinerary we perceive no traces of that vanity for which its Author was subsequently too much distinguished."—*Monthly Review, March 1812.*

"Before we close our report of this very extraordinary production, we deem it only an act of justice to the learned and laborious Editor, to mention, that he has bestowed much trouble in deciphering the original manuscript, and in procuring a faithful version of its miscellaneous contents. Even the *fac-similes* of the rough drawings, though executed in a very different style from the pretty plates of Mr. Ackermann's Repository, contribute, nevertheless, to the graphic and ghostly air of the whole performance."—*Edinburgh Review, Feb. 1812.*

PETRALOGY: A TREATISE ON ROCKS. By JOHN PINKERTON, Author of Modern Geography, &c. Handsomely printed in 2 large vols. 8vo, with 25 Vignette and other Engravings. Price 2l. 2s. in boards.

Extracts from the Introduction.—"I would propose, in the present advanced state of science, that the mineral kingdom be divided into three provinces: 1. PETRALOGY, or the knowledge of rocks, or stones which occur in large masses, 2. LITHOLOGY; the knowledge of gems and small stones, 3. METALLOGY; or the knowledge of metals. Each of these branches is even at present so important, and offers such numerous topics of disquisition and research, that in the course of no long period a Professor of each will appear in Universities; and each might occupy the sole pursuit of an Author who is zealous to make discoveries, or to compose complete and classical works."—"In the present Work it is to be hoped that the want of attention, care, or labour will not be accused. The Author has sedulously employed the intervals of ten years in this production, particularly three which he passed at Paris, where he had constant opportunities of seeing the most opulent cabinets, and of conversing with men eminent in the science; as PATRIN, GILLÉT-LAUMONT, DAUBUSSON, and others: not to mention interviews with Haüy and with Werner, during his short stay in that capital.—He hopes the Work may at least aspire to the humble praise of utility, the chief aim of his labours."

MINERALOGICAL TRAVELS THROUGH THE HEBRIDES, ORKNEY AND SHETLAND ISLANDS, AND MAINLAND OF SCOTLAND; with Dissertations upon PEAT and KELP. By ROBERT JAMESON, Regius Professor of Natural History, Keeper of the Museum, and Lecturer on Mineralogy in the University of Edinburgh, President of the Wernerian Society, &c. &c. In 2 vols. 4to, with Maps and Plates. Price 1l. 16s. in boards.

POMONA HEREFORDIENSIS: being a descriptive Account of the old CIDER and PERRY FRUITS of HEREFORDSHIRE, which have always been esteemed the best of their kind; with such new Fruits as have been found to possess superior Excellence. Accompanied by 30 highly-finished coloured Engravings, by WILLIAM HOOKER, from Drawings principally by Miss MATTHEWS. Published under the Patronage of the Agricultural Society of Herefordshire. The Descriptions by T. A. KNIGHT, Esq. In 1 vol. royal 4to. Price 4l. 10s. in boards.

POMONA LONDINENSIS. Containing COLOURED REPRESENTATIONS of the BEST FRUITS cultivated in the British Gardens; with De-

Valuable Works lately published by

Scriptions, in which the Author is assisted by the President and Members of the Horticultural Society. By WILLIAM HOOKER, F.H.S. Draughtsman and Engraver to the Society.

The FIRST and SECOND NUMBERS, handsomely printed on extra super-royal 4to, with seven Plates in each, Price 1l. 1s. per Number, are now ready for delivery.

A few Copies are taken off on atlas 4to, and coloured in a superior manner; Price 1l. 15s.

The probable extent of the Work will be about Eighteen Numbers. A Number will be published every Four Months.

POMONA BRITANNICA: or, A COLLECTION of the most esteemed FRUITS at present cultivated in this Country: together with the Blossoms and Leaves of such as are necessary to distinguish the various Sorts from each other. Selected from the Royal Gardens at Hampton Court, and the remainder from the most celebrated Gardens round London. Accurately drawn and coloured from Nature; with full Descriptions of their various Qualities, Seasons, &c. By GEORGE BROOKSHAW. In Atlas folio; the Letter-press printed by BENSLEY in the first style of elegance; with 90 Engravings, engraved and coloured in a novel and beautiful style. Price 45l. in boards.

This Work was originally published in Thirty Numbers, at 1l. 10s. each. Subscribers who have not yet completed their Copies, may do it by an early application to the Publishers.

HORTUS CANTABRIGIENSIS: or, A CATALOGUE of PLANTS, INDIGENOUS and EXOTIC. By JAMES DONN, Curator, Fellow of the Linnean and Horticultural Societies. The Seventh Edition, in small 3vo. Price 10s. in boards.

MAWE'S GARDENER'S CALENDAR, AND GENERAL DIRECTORY. By JOHN ABERCROMBIE. The Twentieth Edition, corrected and greatly enlarged, in 1 thick vol. 12mo. Price 7s. 6d. in boards.

AN ESSAY ON THE PROBABILITY OF SENSATION IN VEGETABLES. By J. P. TUPPER, F.L.S. Member of the Royal College of Surgeons. In 8vo. Price 6s. in boards.

FLORA GRÆCA: sive Plantarum rariorum Historia, quas in Provinciis aut Insulis Graecis legit, investigavit, et depingit curavit, JOHANNES STIBTHORP, M.D. S.S. Reg. et Linu. Lond. Socius, Bot. Professor Regius in Academia Oxoniensi. Hic ille etiam insertae sunt pauca species quas Vir idem clarissimus, Græcam versus navigans, in Ithere, presertim apud Italianam et Siciliam, invenerit. Characteres omnium, Descriptiones et Synonyma elaboravit JACOBUS EDWARDUS SMITH, M.D. &c. Societatis Linneanae Londinensis Praeses.

VOLUME the First, containing Two FASCICULI; and the First FASCICULUS of the Second Volume, are now published. This Work is beautifully printed in folio, and each Fasciculus contains 50 Plates engraved and coloured in the most finished manner by SOWERBY, from drawings by FERD. BAUER. Price Twelve Guineas each.

The whole Work will extend to Ten Volumes, or Twenty Fasciculi. From the nature of the Work, and the great expenses attending it, only Fifty Copies are prepared for sale.

WHITE, COCHRANE, and Co. Fleet Street.

FLORÆ GRECE PRODROMUS: sive Plantarum omnium Enumeratio, quas in Provinciis aut Insulis Græciæ inventis JOHANNES SIBTHORP, M.D. &c. Hic illie etiam insertæ sunt pauculae Species quas Vir idem clarissimus, Græciam versus navigans, in Itinere, præsertim apud Italiam et Siciliam, invenerit. Characteres omnium, Descriptiones et Synonyma, elaboravit JACOBUS EDWARDUS SMITH, M.D. &c. Societatis Linneanae Londinensis Praeses.

VOLUME the FIRST, containing Two Parts, and the First Part of the Second Volume, handsomely printed in royal 8vo. are published, Price 10s. 6d. each Part in boards.

The Fourth Part (which is in progress) will complete the Work.

FLORA BRITANNICA, Auctore JAC. EDV. SMITH, M.D. Societatis Linneanae Praes. &c. &c.

Volumes the First, Second, and Third, in 8vo. Price 4l. 6s. 6d. in boards.

The Fourth and Fifth Volumes are in preparation.

AN INTRODUCTION TO PHYSIOLOGICAL AND SYSTEMATIC BOTANY. By JAMES EDWARD SMITH, M.D. F.R.S. &c. &c. President of the Linnean Society. The Third Edition, in 8vo. with Engravings. Price 14s. in boards.

A few Copies with coloured Plates. Price 1l. 8s. in boards.

The Edinburgh Reviewers, at the conclusion of their review of "Dr. Smith's Introduction to Botany," observe that "Botany is daily becoming a more fashionable female study"; and this is an elementary book which may be put with confidence into the hands of women, without any risk of wounding the most delicate mind. We are happy, for the sake of those fair students, to observe, that "Dr. Smith promises a translation of his *Flora Britannica*; for this, we doubt not, will, when accomplished by Dr. Smith himself, form the best popular Herbal ever published. In the mean time he very candidly recommends Dr. Withering's Arrangement of British Plants: to which we would take the liberty to add Mr. Galpine's Compendium of British Botany, (which is indeed nearly a translation of Dr. Smith's *Compendium Floræ Britannicae*,) as a most useful and commodious pocket companion in botanical excursions."

A SYSTEMATIC ARRANGEMENT OF BRITISH PLANTS: with an easy Introduction to the Study of Botany. Illustrated by Copper-plates. By W. WITHERING, M.D. F.R.S. &c. The Fifth Edition, corrected and considerably enlarged, by WILLIAM WITHERING, Esq. In 4 vols. 8vo. Price 2l. 8s. in boards.

A SYNOPTICAL COMPEND OF BRITISH BOTANY; (from the Class Monandria to Polygamia inclusive;) arranged after the Linnaean System; and containing the essential Characters of the Genera, the specific Characters, English Names, Places of Growth, Soil, and Situation, Colour of the Flowers, Times of Flowering, Duration, and References to Figures. By JOHN GALPINE, A.L.S. In foolscap 8vo. with a Frontispiece. Price 10s. 6d. in boards.

LETTERS ON THE ELEMENTS OF BOTANY. Addressed to a Lady, by the celebrated J. J. ROUSSEAU. Translated into English;

Works lately published by WHITE and Co. Fleet Street.

with Notes, and Twenty-four additional Letters fully explaining the System of Linnaeus. By THOMAS MARTYN, B.D. F.R. & L.S.S. Regius Professor of Botany in the University of Cambridge. The Seventh Edition, with Corrections and Improvements, in 8vo. Price 8s. in boards.

THIRTY-EIGHT PLATES, with Explanations, intended to illustrate Linnaeus's System of Vegetables, and particularly adapted to the LETTERS ON THE ELEMENTS OF BOTANY. By the SAME AUTHOR. In 8vo. Price 9s. in boards, or coloured, Price 18s. in boards.

THE LANGUAGE OF BOTANY; being A DICTIONARY of the TERMS made use of in that Science, principally by Linnaeus; with familiar Explanations, and an Attempt to establish significant English Terms. By the SAME AUTHOR. The Third Edition, corrected and enlarged, in 8vo. Price 7s. in boards.

AN INTRODUCTION TO THE SCIENCE OF BOTANY, chiefly extracted from the Writings of Linnaeus. By the late JAMES LEE, Nurseryman and Florist. To which are added several new Tables and Notes, and a Life of the Author. Fourth Edition, corrected and enlarged by JAMES LEE, Son and Successor to the Author. In 8vo, with a Portrait and Plates, Price 14s. in boards; or with the Plates coloured, 21s. in boards.

AN INTRODUCTORY ON TO THE STUDY OF CRYPTOGAMOUS PLANTS. By KURT SPRENGET, M.D. Professor of Botany at Halle, &c. Translated from the German. Neatly printed in 8vo. with 10 4to Plates, Price 9s. in boards; or with the Plates coloured, 18s. in boards.

FERDINANDI BAUER ILLUSTRATIONES FLORÆ NOVÆ HOLLANDIE, sive ICONES GENERUM quæ in Prodromo Flora Novæ Hollandiae et Insulæ Van Diemen descriptis ROBERTUS BROWN.

Nos. I and II, handsomely printed in folio, with Five Plates in each, price 7s. per Number.

THEOPHRASTI ERESII DE HISTORIA PLANTARUM Libri X. Grecè, cum Syllabo Generum et Specierum, Glossario et Notis. Curante J. STACKHOUSE, Arm. Soc. Linn. S.

Pars I. continens Libros V. priores. Handsomely printed in crown 8vo. with a Portrait of the Author, Price 12s. in boards.

MR. PENNANT'S BRITISH ZOOLOGY; a new Edition, in 4 vols. 8vo. with 298 Engravings, many of which are new. Price 6l. in boards; or in royal 8vo. 8l. 8s. in boards.

✓ In this Edition the Additions by the Editor constitute an increase of nearly a third to the preceding Edition.