




**IMAGE EVALUATION  
TEST TARGET (MT-3)**



**Photographic  
Sciences  
Corporation**

23 WEST MAIN STREET  
WEBSTER, N.Y. 14580  
(716) 872-4503

2.8  
2.5  
2.2  
2.0  
1.8

**CIHM/ICMH  
Microfiche  
Series.**

**CIHM/ICMH  
Collection de  
microfiches.**



**Canadian Institute for Historical Microreproductions / Institut canadien de microreproductions historiques**

10  
01

**© 1982**

Technical and Bibliographic Notes/Notes techniques et bibliographiques

The Institute has attempted to obtain the best original copy available for filming. Features of this copy which may be bibliographically unique, which may alter any of the images in the reproduction, or which may significantly change the usual method of filming, are checked below.

L'Institut a microfilmé le meilleur exemplaire qu'il lui a été possible de se procurer. Les détails de cet exemplaire qui sont peut-être uniques du point de vue bibliographique, qui peuvent modifier une image reproduite, ou qui peuvent exiger une modification dans la méthode normale de filmage sont indiqués ci-dessous.

- |  |  |
|--|--|
| <input type="checkbox"/> Coloured covers/<br>Couverture de couleur   | <input type="checkbox"/> Coloured pages/<br>Pages de couleur   |
| <input type="checkbox"/> Covers damaged/<br>Couverture endommagée  | <input type="checkbox"/> Pages damaged/<br>Pages endommagées   |
| <input type="checkbox"/> Covers restored and/or laminated/<br>Couverture restaurée et/ou pelliculée  | <input type="checkbox"/> Pages restored and/or laminated/<br>Pages restaurées et/ou pelliculées  |
| <input type="checkbox"/> Cover title missing/<br>Le titre de couverture manque   | <input checked="" type="checkbox"/> Pages discoloured, stained or foxed/<br>Pages décolorées, tachetées ou piquées   |
| <input type="checkbox"/> Coloured maps/<br>Cartes géographiques en couleur   | <input type="checkbox"/> Pages detached/<br>Pages détachées  |
| <input type="checkbox"/> Coloured ink (i.e. other than blue or black)/<br>Encre de couleur (i.e. autre que bleue ou noire)   | <input checked="" type="checkbox"/> Showthrough/<br>Transparence   |
| <input type="checkbox"/> Coloured plates and/or illustrations/<br>Planches et/ou illustrations en couleur  | <input type="checkbox"/> Quality of print varies/<br>Qualité inégale de l'impression   |
| <input type="checkbox"/> Bound with other material/<br>Relié avec d'autres documents   | <input type="checkbox"/> Includes supplementary material/<br>Comprend du matériel supplémentaire   |
| <input type="checkbox"/> Tight binding may cause shadows or distortion<br>along interior margin/<br>La reliure serrée peut causer de l'ombre ou de la<br>distortion le long de la marge intérieure   | <input type="checkbox"/> Only edition available/<br>Seule édition disponible   |
| <input type="checkbox"/> Blank leaves added during restoration may<br>appear within the text. Whenever possible, these<br>have been omitted from filming/<br>Il se peut que certaines pages blanches ajoutées<br>lors d'une restauration apparaissent dans le texte,<br>mais, lorsque cela était possible, ces pages n'ont<br>pas été filmées. | <input type="checkbox"/> Pages wholly or partially obscured by errata<br>slips, tissues, etc., have been refilmed to<br>ensure the best possible image/<br>Les pages totalement ou partiellement<br>obscurcies par un feuillet d'errata, une pelure,<br>etc., ont été filmées à nouveau de façon à<br>obtenir la meilleure image possible. |
| <input checked="" type="checkbox"/> Additional comments:/<br>Commentaires supplémentaires:   | Pagination continued from Vol. II.   |

This item is filmed at the reduction ratio checked below/  
Ce document est filmé au taux de réduction indiqué ci-dessous.

10X	14X	18X	22X	26X	30X
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
12X	16X	20X	24X	28X	32X

The copy filmed here has been reproduced thanks to the generosity of:



Library.  
Department of Fisheries and Oceans

The images appearing here are the best quality possible considering the condition and legibility of the original copy and in keeping with the filming contract specifications.

Original copies in printed paper covers are filmed beginning with the front cover and ending on the last page with a printed or illustrated impression, or the back cover when appropriate. All other original copies are filmed beginning on the first page with a printed or illustrated impression, and ending on the last page with a printed or illustrated impression.

The last recorded frame on each microfiche shall contain the symbol  $\rightarrow$  (meaning "CONTINUED"), or the symbol  $\nabla$  (meaning "END"), whichever applies.

Maps, plates, charts, etc., may be filmed at different reduction ratios. Those too large to be entirely included in one exposure are filmed beginning in the upper left corner, left to right and top to bottom, as many frames as required. The following diagrams illustrate the method:



L'exemplaire filmé fut reproduit grâce à la générosité de:

Bibliothèque.  
Ministère des pêches et océans

Les images suivantes ont été reproduites avec le plus grand soin, compte tenu de la condition et de la netteté de l'exemplaire filmé, et en conformité avec les conditions du contrat de filmage.

Les exemplaires originaux dont la couverture en papier est imprimée sont filmés en commençant par le premier plat et en terminant soit par la dernière page qui comporte une empreinte d'impression ou d'illustration, soit par le second plat, selon le cas. Tous les autres exemplaires originaux sont filmés en commençant par la première page qui comporte une empreinte d'impression ou d'illustration et en terminant par la dernière page qui comporte une telle empreinte.

Un des symboles suivants apparaîtra sur la dernière image de chaque microfiche, selon le cas: le symbole  $\rightarrow$  signifie "A SUIVRE", le symbole  $\nabla$  signifie "FIN".

Les cartes, planches, tableaux, etc., peuvent être filmés à des taux de réduction différents. Lorsque le document est trop grand pour être reproduit en un seul cliché, il est filmé à partir de l'angle supérieur gauche, de gauche à droite, et de haut en bas, en prenant le nombre d'images nécessaire. Les diagrammes suivants illustrent la méthode.

UN

A DESC  
TI

17-2

71185

SMITHSONIAN INSTITUTION.

UNITED STATES NATIONAL MUSEUM. 59231  
11

---

# BULLETIN

OF THE

UNITED STATES NATIONAL MUSEUM.

No. 47.

---

## THE FISHES

OF

NORTH AND MIDDLE AMERICA:

A DESCRIPTIVE CATALOGUE OF THE SPECIES OF FISH-LIKE VERTEBRATES FOUND IN  
THE WATERS OF NORTH AMERICA, NORTH OF THE ISTHMUS OF PANAMA.

BY

DAVID STARR JORDAN, Ph. D.,

PRESIDENT OF THE LELAND STANFORD JUNIOR UNIVERSITY AND OF THE  
CALIFORNIA ACADEMY OF SCIENCES,

AND

BARTON WARREN EVERMANN, Ph. D.,

ICHTHOLOGIST OF THE UNITED STATES FISH COMMISSION.

16. 71

PART III.

WASHINGTON:

GOVERNMENT PRINTING OFFICE.

1898.

N

SMITHSONIAN INSTITUTION.

UNITED STATES NATIONAL MUSEUM.

THE FISHES  
OF  
NORTH AND MIDDLE AMERICA:

A DESCRIPTIVE CATALOGUE

OF THE

SPECIES OF FISH-LIKE VERTEBRATES FOUND IN THE  
WATERS OF NORTH AMERICA, NORTH OF  
THE ISTHMUS OF PANAMA.

BY

DAVID STARR JORDAN, PH. D.,

PRESIDENT OF THE LELAND STANFORD JUNIOR UNIVERSITY AND OF THE  
CALIFORNIA ACADEMY OF SCIENCES,

AND

BARTON WARREN EVERMANN, PH. D.,

ICHTHYOLOGIST OF THE UNITED STATES FISH COMMISSION.

PART III.

WASHINGTON:

GOVERNMENT PRINTING OFFICE.

1898.



CLA

918  
J82  
V 3

## TABLE OF CONTENTS, PART III.

### CLASS III. PISCES—Continued.

#### ORDER BB. ACANTHOPTERI—Continued.

<i>Family CLXXXIV. Triglidae—Continued.</i>	Page.
Group Gobioldea.....	2184
<i>Family CLXXXVII. Callionymidae</i> .....	2184
Genus 799. <i>Callionymus</i> , Linnaeus.....	2185
2511. <i>bairdi</i> , Jordan.....	2185
2512. <i>himantophorus</i> , Goode & Bean.....	2186
2513. <i>callurus</i> , Eigenmann & Eigenmann.....	2187
2514. <i>pauciradiatus</i> , Gill.....	2188
<i>Family CLXXXVIII. Gobiidae</i> .....	2188
Genus 800. <i>Ioglossus</i> , Bean.....	2192
2515. <i>callurus</i> , Beau.....	2193
Genus 801. <i>Philypnus</i> , Cuvier & Valenciennes.....	2194
2516. <i>dormitor</i> (Lacépède).....	2194
2517. <i>lateralis</i> , Gill.....	2195
Genus 802. <i>Dormitator</i> , Gill.....	2195
2518. <i>maculatus</i> (Bloch).....	2196
Genus 803. <i>Gnævina</i> , Bleeker.....	2198
2519. <i>gnævina</i> , Cuvier & Valenciennes.....	2198
Genus 804. <i>Eleotris</i> (Gronow) Bloch & Schneider.....	2199
2520. <i>amblyopsis</i> (Cope).....	2199
2521. <i>abaenrus</i> , Jordan & Gilbert.....	2200
2522. <i>pisonis</i> (Gmelin).....	2200
2523. <i>perniger</i> (Cope).....	2201
2524. <i>pietus</i> (Knor & Steindachner).....	2201
Genus 805. <i>Alexurus</i> , Jordan.....	2202
2525. <i>armiger</i> , Jordan & Richardson.....	2203
Genus 806. <i>Erotelis</i> , Poey.....	2203
2526. <i>amaragdus</i> (Cuvier & Valenciennes).....	2204
Genus 807. <i>Gymneleotris</i> , Bleeker.....	2204
2527. <i>seminudus</i> (Günther).....	2204
Genus 808. <i>Chriolepis</i> , Gilbert.....	2205
2528. <i>minutillus</i> , Gilbert.....	2205
Genus 809. <i>Sicydium</i> , Cuvier & Valenciennes.....	2205
2529. <i>plumieri</i> (Bloch).....	2206
2530. <i>antillarum</i> , Ogilvie-Grant.....	2206
2531. <i>vincente</i> , Jordan & Evermann.....	2207
2531 (a). <i>punctatum</i> , Perugia.....	2207
Genus 810. <i>Cotylopus</i> , Gulchenot.....	2207
Subgenus <i>Sleya</i> , Jordan & Evermann.....	2207
2532. <i>gymnogaster</i> (Ogilvie-Grant).....	2207
2533. <i>salvini</i> (Ogilvie-Grant).....	2208
Genus 811. <i>Evorthodus</i> , Gill.....	2208
2534. <i>breviceps</i> , Gill.....	2208
Genus 812. <i>Lophogobius</i> , Gill.....	2209
2535. <i>cyprinoides</i> (Pallas).....	2209

507  
818  
J82  
V. 3

## CLASS III. PISCES—Continued.

## ORDER BB. ACANTHOPTERI—Continued.

## Family CLXXXVIII. Gobiidæ—Continued.

	Page.
Genus 813. <i>Gobius</i> (Artedi) Linnaeus .....	2210
Subgenus <i>Gobius</i> .....	2216
2536. <i>soporator</i> , Cuvier & Valenciennes .....	2216
Subgenus <i>Ctenogobius</i> , Gill .....	2218
2537. <i>nicholsii</i> , Bean .....	2218
2538. <i>eigenmanni</i> , Garman .....	2218
2539. <i>glaucofrœnum</i> (Gill) .....	2219
2540. <i>manglicola</i> , Jordan & Starks .....	2220
2541. <i>stigmaturus</i> , Goode & Bean .....	2220
2542. <i>quadriporus</i> , Cuvier & Valenciennes .....	2221
2543. <i>stuefeldti</i> , Jordan & Eigenmann .....	2221
2544. <i>boleosoma</i> , Jordan & Gilbert .....	2221
2545. <i>fasciatus</i> (Gill) .....	2222
2546. <i>eucæopus</i> , Jordan & Gilbert .....	2223
2547. <i>stigmaticus</i> (Poey) .....	2224
2548. <i>lyricus</i> , Girard .....	2224
2549. <i>garmani</i> , Eigenmann & Eigenmann .....	2225
2550. <i>zebra</i> , Gilbert .....	2226; 2867
Subgenus <i>Euctenogobius</i> , Gill .....	2226
2551. <i>poeyi</i> , Steindachner .....	2226
2552. <i>badius</i> (Gill) .....	2227
Subgenus <i>Gobionellus</i> , Girard .....	2227
2553. <i>microdon</i> , Gilbert .....	2227
2554. <i>smaragdus</i> , Cuvier & Valenciennes .....	2227
2555. <i>strigatus</i> , O'Shaughnessy .....	2228
2556. <i>sagittula</i> (Günther) .....	2228
2557. <i>hastatus</i> , Girard .....	2229
2558. <i>oceanicus</i> , Pallas .....	2230
Subgenus <i>Lythrypnus</i> , Jordan & Evermann .....	2230
2559. <i>dalli</i> , Gilbert .....	2230
Genus 814. <i>Garmannia</i> , Jordan & Evermann .....	2231
Subgenus <i>Garmannia</i> .....	2232
2560. <i>paradoxa</i> (Günther) .....	2232
2561. <i>hemigymina</i> (Eigenmann & Eigenmann) .....	2233
Subgenus <i>Enypnius</i> , Jordan & Evermann .....	2233
2562. <i>seminuda</i> (Günther) .....	2233
Genus 815. <i>Awaous</i> , Steindachner .....	2234
2563. <i>flavus</i> (Cuvier & Valenciennes) .....	2235
2564. <i>nelsoni</i> , Evermann .....	2235
2565. <i>taiasica</i> (Lichtenstein) .....	2236
2566. <i>mexicanus</i> (Günther) .....	2237
Genus 816. <i>Bollmannia</i> , Jordan .....	2237
2567. <i>ocellata</i> , Gilbert .....	2238
2568. <i>chlamydes</i> , Jordan .....	2238
2569. <i>macropona</i> , Gilbert .....	2239
2570. <i>stigmatura</i> , Gilbert .....	2239
Genus 817. <i>Aboma</i> , Jordan & Starks .....	2240
2571. <i>etheostoma</i> , Jordan & Starks .....	2240
2572. <i>lucretie</i> (Eigenmann & Eigenmann) .....	2241
2573. <i>chiquita</i> (Jenkins & Evermann) .....	2241
Genus 818. <i>Microgobius</i> , Poey .....	2242
2574. <i>gulosus</i> (Girard) .....	2243
2575. <i>eulepis</i> , Eigenmann & Eigenmann .....	2244
2576. <i>thalassinus</i> , Jordan & Gilbert .....	2245
2577. <i>signatus</i> , Poey .....	2246
Genus 819. <i>Zalypnus</i> , Jordan & Evermann .....	2246

## CLASS III. PISCES—Continued.

## ORDER BB. ACANTHOPTERI—Continued.

Family CLXXXVIII. *Gobiidae*—Continued.

	Page.
2578. <i>cyclolepis</i> (Gilbert) .....	2246
2579. <i>emblematis</i> (Jordan & Gilbert) .....	2247
Genus 820. <i>Eucyclogobius</i> , Gill .....	2248
2580. <i>newberryi</i> (Girard) .....	2248
Genus 821. <i>Lepidogobius</i> , Gill .....	2249
2581. <i>lepidus</i> (Girard) .....	2249
Genus 822. <i>Gillichthys</i> , Cooper .....	2249
2582. <i>mirabilis</i> , Cooper .....	2250
2583. <i>detrusus</i> , Gilbert & Seefield .....	2251
Genus 823. <i>Quiletula</i> , Jordan & Evermann .....	2251
2584. <i>y-canda</i> (Jenkins & Evermann) .....	2251
Genus 824. <i>Hypnus</i> , Jordan & Evermann .....	2253
2585. <i>gillerti</i> (Eigenmann & Eigenmann) .....	2253
Genus 825. <i>Clovelandia</i> , Eigenmann & Eigenmann .....	2254
2586. <i>ios</i> (Jordan & Gilbert) .....	2254
2587. <i>rosa</i> , Jordan & Evermann .....	2255
Genus 826. <i>Evermannia</i> , Jordan .....	2256
2588. <i>longipinnis</i> (Steindachner) .....	2256
2589. <i>zosterura</i> (Jordan & Gilbert) .....	2256
Genus 827. <i>Gobiosoma</i> , Girard .....	2257
2590. <i>histrion</i> , Jordan .....	2258
2591. <i>molestum</i> , Girard .....	2258
2592. <i>bosei</i> (Lacépède) .....	2259
2593. <i>crescentale</i> , Gilbert .....	2259
2594. <i>multifasciatum</i> , Steindachner .....	2260
Genus 828. <i>Barbulifer</i> , Eigenmann & Eigenmann .....	2260
2595. <i>centhæus</i> (Jordan & Gilbert) .....	2260
Genus 829. <i>Typhlogobius</i> , Steindachner .....	2261
2596. <i>californiensis</i> , Steindachner .....	2262
Genus 830. <i>Tyntlastes</i> , Günther .....	2262
2597. <i>brevis</i> (Günther) .....	2262
2598. <i>sagitta</i> (Günther) .....	2263
Genus 831. <i>Gobioides</i> , Lacépède .....	2263; 2268
2599. <i>brunssonnetii</i> , Lacépède .....	2263
2600. <i>peruanus</i> (Steindachner) .....	2264
Genus 832. <i>Cayennia</i> , Sauvage .....	2265
2601. <i>guichenoti</i> , Sauvage .....	2265
SUBORDER DISCOCEPHALI .....	2265
Family CLXXXIX. <i>Echeneidae</i> .....	2265
Genus 833. <i>Phtheiroichthys</i> , Gill .....	2268
2602. <i>lineatus</i> (Menzies) .....	2268
Genus 834. <i>Echeneis</i> (Artedi) Linnaeus .....	2268
2603. <i>naukrates</i> , Linnaeus .....	2269
2604. <i>naukrateoides</i> , Zuiuw .....	2270
Genus 835. <i>Remilegia</i> , Gill .....	2270
2605. <i>australis</i> (Bennett) .....	2270
Genus 836. <i>Remora</i> , Gill .....	2271
Subgenus <i>Remora</i> .....	2271
2606. <i>remora</i> (Linnaeus) .....	2271
Subgenus <i>Remorina</i> , Jordan & Evermann .....	2272
2607. <i>albescens</i> (Temminck & Schlegel) .....	2272
Subgenus <i>Remoropsis</i> , Gill .....	2272
2608. <i>brachyptera</i> (Lowe) .....	2272
Genus 837. <i>Rhombelistrus</i> , Gill .....	2273
2609. <i>osteocheir</i> (Cuvier) .....	2273
Group <i>Trachinoidea</i> .....	2273

## CLASS III. PISCES—Continued.

## ORDER BB. ACANTHOPTERI—Continued.

	Page.
<i>Family CXO. Malacanthidae</i> .....	2274
Genus 838. <i>Malacanthus</i> , Cuvier.....	2275
2610. <i>plumieri</i> (Bloch).....	2275
Genus 839. <i>Caulolatilus</i> , Gill.....	2276
2611. <i>princeps</i> (Jenyns).....	2276
2612. <i>microps</i> , Goode & Bean.....	2277
2613. <i>cynnops</i> , Poey.....	2278
Genus 840. <i>Lopholatilus</i> , Goode & Bean.....	2278
2614. <i>chamaeleonticeps</i> , Goode & Bean.....	2278
<i>Family CXCI. Opisthognathidae</i> .....	2279
Genus 841. <i>Opisthognathus</i> Cuvier.....	2280
2615. <i>lonchurum</i> , Jordan & Gilbert.....	2281
2616. <i>punctatum</i> , Peters.....	2281
2617. <i>macrognathum</i> , Poey.....	2281
2618. <i>ornatum</i> , Jenkins & Evermann.....	2282
Genus 842. <i>Gnathypops</i> , Gill.....	2283
2619. <i>scops</i> , Jenkins & Evermann.....	2283
2620. <i>maxillosa</i> (Poey).....	2284
2621. <i>macrops</i> (Poey).....	2284
2622. <i>rhomblea</i> (Jordan & Gilbert).....	2285
2623. <i>snyderi</i> , Jordan & Evermann.....	2285
2624. <i>mystacina</i> , Jordan.....	2286
Genus 843. <i>Loucheopisthus</i> , Gill.....	2286
2625. <i>micrognathus</i> (Poey).....	2287
<i>Family CXCII. Bathymasteridae</i> .....	2287
Genus 844. <i>Bathymaster</i> .....	2288
2626. <i>signatus</i> , Cope.....	2288
Genus 845. <i>Rouquillus</i> , Jordan & Starks.....	2289
2627. <i>jordani</i> (Gilbert).....	2289
Genus 846. <i>Rathbunella</i> , Jordan & Evermann.....	2289
2628. <i>hypoplecta</i> (Gilbert).....	2290
<i>Family CXCIII. Chaenodontidae</i> .....	2291
Genus 847. <i>Chiasmodon</i> , Johnson.....	2291
2629. <i>niger</i> , Johnson.....	2291
Genus 848. <i>Pseudoscopelus</i> , Lütken.....	2292
2630. <i>scriptus</i> , Lütken.....	2292
<i>Family CXCIV. Chaenichthyidae</i> .....	2293
Genus 849. <i>Hypsicoetes</i> , Goode.....	2293
2631. <i>gobioides</i> , Goode.....	2294
<i>Family CXCV. Trichodontidae</i> .....	2295
Genus 850. <i>Trichodon</i> (Steller) Cuvier.....	2295
2632. <i>trichodon</i> (Tilesius).....	2295
Genus 851. <i>Arctoscopus</i> , Jordan & Evermann.....	2297
2633. <i>japonicus</i> (Steindachner).....	2297
<i>Family CXCVI. Dactyloscopidae</i> .....	2297
Genus 852. <i>Gillellus</i> , Gilbert.....	2298
2634. <i>semicinctus</i> , Gilbert.....	2298
2635. <i>arenicola</i> , Gilbert.....	2299
2636. <i>ornatus</i> , Gilbert.....	2299
Genus 853. <i>Dactyloscopus</i> , Gill.....	2300
Subgenus <i>Dactyloscopus</i> .....	2301
2637. <i>pectoralis</i> , Gill.....	2301
2638. <i>tridigitatus</i> , Gill.....	2301
2639. <i>poeyi</i> , Gill.....	2302
2640. <i>lunatiens</i> , Gilbert.....	2302
Subgenus <i>Esloscopus</i> , Jordan & Evermann.....	2303
2641. <i>zelotes</i> , Jordan & Evermann.....	2303

## CLASS III. PISCES—Continued.

## ORDER BB. ACANTHOPTERI—Continued.

	Page.
<i>Family CXCVI. Dactyloscopidae—Continued.</i>	
Genus 854. <i>Dactylagmus</i> , Gill	2304
2642. <i>mundus</i> , Gill	2304
Genus 855. <i>Myxodagmus</i> , Gill	2305
2643. <i>opercularis</i> , Gill	2305
<i>Family CXCVII. Uranoscopidae</i>	
Genus 856. <i>Astroscopus</i> , Brevoort	2306
2644. <i>y-græcum</i> (Cuvier & Valenciennes)	2307
2645. <i>zephyreus</i> , Gilbert & Starks	2309
2646. <i>guttatus</i> (Abbott)	2310
Genus 857. <i>Kathetostomus</i> , Günther	2311
2647. <i>averruncus</i> , Jordan & Bollman	2311
2648. <i>albiguttum</i> , Bean	2312
SUBORDER HAPLODOCI	
<i>Family CXCVIII. Batrachoididae</i>	
Genus 858. <i>Batrachoides</i> , Lacépède	2313; 2868
2649. <i>surinamensis</i> (Bloch & Schneider)	2314
2650. <i>pacifæ</i> (Günther)	2314
Genus 859. <i>Opsanus</i> , Rafinesque	2315
2651. <i>tan</i> (Linnaeus)	2315
2652. <i>pardus</i> (Goode & Bean)	2316
Genus 860. <i>Porlethys</i> , Girard	2317
2653. <i>porosiasinus</i> (Cuvier & Valenciennes)	2319
2654. <i>notatus</i> , Girard	2321
2655. <i>margaritatus</i> (Richardson)	2322
Genus 861. <i>Thalassophryne</i> , Günther	2323
2656. <i>maculosa</i> , Günther	2324
2657. <i>reticulata</i> , Günther	2325
Genus 862. <i>Dector</i> , Jordan & Evermann	2325
2658. <i>dowi</i> (Jordan & Gilbert)	2325
SUBORDER XENOPTERYGII	
<i>Family CXCIX. Gobioidae</i>	
Genus 863. <i>Cantharus</i> , Gill	2327
2659. <i>maandricus</i> (Girard)	2328
Genus 864. <i>Brysseteres</i> , Jordan & Evermann	2328
2660. <i>pinniger</i> (Gilbert)	2328
Genus 865. <i>Gobiesox</i> , Lacépède	2329
Subgenus <i>Bryssophilus</i> , Jordan & Evermann	2330
2661. <i>papillifer</i> , Gilbert	2330
Subgenus <i>Gobiesox</i>	2331
2662. <i>gyrinus</i> , Jordan & Evermann	2331
2663. <i>nigripinnis</i> (Peters)	2331
2664. <i>cephalus</i> , Lacépède	2332
2665. <i>indes</i> , Richardson	2333
2666. <i>strumosus</i> , Cope	2333
2667. <i>virgatulus</i> , Jordan & Gilbert	2333
2668. <i>adustus</i> , Jordan & Gilbert	2334
2669. <i>funbris</i> , Gilbert	2334
2370. <i>puccilophthalmus</i> , Jenyns	2335
2671. <i>rhodospilus</i> , Günther	2335
2672. <i>macrophthalmus</i> , Günther	2335
2673. <i>cerasinus</i> , Cope	2336
Subgenus <i>Sicyases</i> , Müller & Troschel	2336
2674. <i>erythroptus</i> , Jordan & Gilbert	2336
2675. <i>rubiginosus</i> (Poey)	2337
2676. <i>carneus</i> (Poey)	2337
2677. <i>hæres</i> , Jordan & Bollman	2337

## CLASS III. PISCES—Continued.

## ORDER BB. ACANTHOPTERI—Continued.

	Page.
<i>Family CXCIX. Gobiesocidae—Continued.</i>	
2678. punctulatus (Poey) .....	2338
2679. fasciatus (Peters) .....	2338
Genus 866. Rimicola, Jordan & Evermann .....	2338
2680. muscarum (Meek & Pierson) .....	2338
2681. rigemanni (Gilbert) .....	2339
Genus 867. Arhaciosa, Jordan & Evermann .....	2340
2682. rhessodon (Rosa Smith) .....	2340
2683. humeralis (Gilbert) .....	2341
2684. rupestris (Poey) .....	2341
2685. zebra (Jordan & Gilbert) .....	2341
2686. eos (Jordan & Gilbert) .....	2343
Group Blennioides .....	2343
<i>Family CV. Blenniidae</i> .....	2344
Genus 868. Enneanectes, Jordan & Evermann .....	2349
2687. carminalis (Jordan & Gilbert) .....	2350
Genus 868(a). Dialommus, Gilbert .....	2668
2687(a). fuscus, Gilbert .....	2668
Genus 869. Heterostichus, Girard .....	2350
2688. rostratus, Girard .....	2351
Genus 870. Gilbousia, Cooper .....	2351
2689. evides (Jordan & Gilbert) .....	2352; 2869
2690. elegans (Cooper) .....	2353
Genus 871. Neoclinus, Girard .....	2354
Subgenus Neoclinus .....	2354
2691. blanchardi, Girard .....	2354
Subgenus Pterognathus, Girard .....	2355
2692. satiriens, Girard .....	2355
Genus 872. Malacoctenus, Gill .....	2356
2693. ocellatus (Steindachner) .....	2356; 2869
2694. varius (Poey) .....	2357
2695. macropus (Poey) .....	2357
2696. lugubris (Poey) .....	2357
2697. gillii (Steindachner) .....	2358
2698. bimaculatus (Steindachner) .....	2358
2699. delalandi (Cuvier & Valenciennes) .....	2358
2700. versicolor (Poey) .....	2359
2701. biguttatus (Cope) .....	2360
Genus 873. Labrisomus, Swainson .....	2360
2702. herminier (Le Sueur) .....	2361
2703. nuchiplanis (Quoy & Gaimard) .....	2362
2704. xanti, Gill .....	2362
2705. bucciferus, Poey .....	2363
2706. microlepidotus, Poey .....	2363
Genus 874. Mniroperes, Jordan & Evermann .....	2364
2707. macrocephalus (Günther) .....	2364
Genus 875. Gobioclinus, Gill .....	2364
2708. gobio (Cuvier & Valenciennes) .....	2365
Genus 876. Starksia, Jordan & Evermann .....	2365
2709. crennobates (Gilbert) .....	2365
Genus 877. Cryptotrema, Gilbert .....	2366
2710. crystallinum, Gilbert .....	2366
Genus 878. Exerpes, Jordan & Evermann .....	2367
2711. asper (Jenkins & Evermann) .....	2367
Genus 879. Auchenopterus, Günther .....	2369
Subgenus Corallicola, Jordan & Evermann .....	2369
2712. nigriplanis (Steindachner) .....	2369

## CLASS III. PISCES—Continued.

## ORDER BB. ACANTHOPTERI—Continued.

## Family CC. Blenniidae—Continued.

Page.		Page.
2338	2713. <i>altiveis</i> (Lockington) .....	2370
2338	2714. <i>marmoratus</i> (Steindachner) .....	2371
2338	Subgenus <i>Auchenopterus</i> .....	2371
2339	2715. <i>affinis</i> (Steindachner) .....	2371
2340	2716. <i>monophthalmus</i> , Günther .....	2372
2341	2717. <i>integripinnis</i> (Rosa Smith) .....	2372
2341	2718. <i>fasciatus</i> (Steindachner) .....	2373
2341	2719. <i>nox</i> (Jordan & Gilbert) .....	2373
2341	Genus 880. <i>Paraclinus</i> , Mocquard .....	2374
2341	2720. <i>charteri</i> , Mocquard .....	2374
2343	Genus 881. <i>Emmion</i> , Jordan .....	2375
2343	2721. <i>bristole</i> , Jordan .....	2375
2344	Genus 882. <i>Atopoclinus</i> , Vaillant .....	2376
2349	2722. <i>ringens</i> , Vaillant .....	2376
2350	Genus 883. <i>Ranula</i> , Jordan & Bollman .....	2377
2868	2723. <i>azalea</i> , Jordan & Bollman .....	2377
2868	Genus 884. <i>Blennius</i> (Artedi) Linnaeus .....	2377
2350	Subgenus <i>Lipophrys</i> , Gill .....	2378
2351	2724. <i>carolinus</i> (Cuvier & Valenciennes) .....	2378
2869	2725. <i>ficorum</i> , Cuvier & Valenciennes .....	2379
2353	2726. <i>stearnsi</i> , Jordan & Gilbert .....	2379
2354	2727. <i>favosus</i> , Goode & Bean .....	2380
2354	2728. <i>pilicornis</i> , Cuvier & Valenciennes .....	2380
2355	2729. <i>marmoratus</i> , Poey .....	2381
2355	2730. <i>truncatus</i> (Poey) .....	2381
2356	2731. <i>vinctus</i> , Poey .....	2382
2869	2732. <i>cristatus</i> , Linnaeus .....	2382
2357	Genus 885. <i>Scartella</i> , Jordan .....	2384
2357	2733. <i>microstoma</i> (Poey) .....	2384
2358	Genus 886. <i>Hypleurochilus</i> , Gill .....	2385
2358	2734. <i>geminatus</i> (Wood) .....	2385
2359	Genus 887. <i>Hypsoblennius</i> , Gill .....	2386
2360	Subgenus <i>Hypsoblennius</i> .....	2386
2361	2735. <i>gilberti</i> (Jordan) .....	2386
2362	2736. <i>gentilis</i> (Girard) .....	2387
2362	2737. <i>striatus</i> (Steindachner) .....	2388
2363	2738. <i>lonthas</i> (Jordan & Gilbert) .....	2388
2364	2739. <i>hantz</i> (Le Sueur) .....	2390
2364	Subgenus <i>Blenniolus</i> , Jordan & Evermann .....	2390
2364	2740. <i>brevipinnis</i> (Günther) .....	2390
2365	Genus 888. <i>Chasmodes</i> , Cuvier & Valenciennes .....	2391
2365	2741. <i>jenkinsi</i> (Jordan & Evermann) .....	2391
2366	2742. <i>quadrifasciatus</i> (Wood) .....	2392
2366	2743. <i>suburne</i> , Jordan & Gilbert .....	2392
2367	2744. <i>novemlineatus</i> (Wood) .....	2393
2367	2745. <i>bosquianus</i> (Lacépède) .....	2394
2369	Genus 889. <i>Honesthes</i> , Gilbert .....	2394
2369	2746. <i>caulopus</i> , Gilbert .....	2394
2369	Genus 890. <i>Scartichthys</i> , Jordan & Evermann .....	2395
2369	2747. <i>ruipunctatus</i> (Cuvier & Valenciennes) .....	2396
2369	Genus 891. <i>Rupiscartes</i> , Swainson .....	2396
2369	2748. <i>atlanticus</i> (Cuvier & Valenciennes) .....	2397
2369	Genus 892. <i>Entomacrodus</i> , Gill .....	2397
2369	2749. <i>chiostictus</i> (Jordan & Gilbert) .....	2398
2369	2750. <i>margaritaceus</i> (Poey) .....	2398
2369	2751. <i>decoratus</i> , Poey .....	2399



## CLASS III. PISCES—Continued.

## ORDER BB. ACANTHOPTERI—Continued.

Family CC. *Blenniidae*—Continued.

	Page.
2752. <i>nigricans</i> , Gill.....	2399
Genus 893. <i>Salarichthys</i> , Guckenhof.....	2400
2753. <i>textilis</i> (Quoy & Gaimard).....	2400
Genus 894. <i>Ophioblennius</i> , Gill.....	2400
2754. <i>welshii</i> (Valenciennes).....	2401
2755. <i>steindachneri</i> , Jordan & Evermann.....	2401
Genus 895. <i>Emblemaria</i> , Jordan & Gilbert.....	2401
2756. <i>atlantica</i> , Jordan & Evermann.....	2402
2757. <i>nivipes</i> , Jordan & Gilbert.....	2402
2758. <i>oculocirris</i> , Jordan.....	2403
Genus 896. <i>Chenopsis</i> , Gill.....	2403
2759. <i>ocellatus</i> , Peey.....	2403
Genus 897. <i>Lucioblennius</i> , Gilbert.....	2404
2760. <i>alepidotus</i> , Gilbert.....	2404
Genus 898. <i>Pholidichthys</i> , Bleeker.....	2405
2761. <i>anguilliformis</i> , Lockington.....	2405
Genus 899. <i>Pseudoblennius</i> , Jenkins & Evermann.....	2406
2762. <i>hyacanthus</i> , Jenkins & Evermann.....	2406
Genus 900. <i>Stathmonotus</i> , Bean.....	2407
2763. <i>hemphillii</i> , Bean.....	2407
Genus 901. <i>Bryostenma</i> , Jordan & Starks.....	2408
2764. <i>polyactcephalum</i> (Pallas).....	2408
2765. <i>nugator</i> , Jordan & Williams.....	2410
Genus 902. <i>Apodichthys</i> , Girard.....	2411
2766. <i>flavidus</i> , Girard.....	2411
2767. <i>univittatus</i> , Lockington.....	2412
Genus 903. <i>Xerocpes</i> , Jordan & Gilbert.....	2413
2768. <i>fuorum</i> (Jordan & Gilbert).....	2413
Genus 904. <i>Ulvicola</i> , Gilbert.....	2413
2769. <i>saneta-rose</i> , Gilbert & Starks.....	2413
Genus 905. <i>Pholis</i> (Gronow) Scopoli.....	2414
Subgenus <i>Urocentrus</i> , Kner.....	2415
2770. <i>pictus</i> (Kner).....	2415
Subgenus <i>Rhodymenichthys</i> , Jordan & Evermann.....	2416
2771. <i>dolichogaster</i> (Pallas).....	2416
Subgenus <i>Pholis</i> .....	2417
2772. <i>fasciatus</i> (Bloch & Schneider).....	2417
2773. <i>gunnellus</i> (Linnaeus).....	2419
2774. <i>ornatus</i> (Girard).....	2419
Genus 906. <i>Gunnellops</i> , Bleeker.....	2420
2775. <i>rosens</i> (Pallas).....	2420
Genus 907. <i>Asternopteryx</i> , Rüppell.....	2420
2776. <i>gunnelliformis</i> , Rüppell.....	2420
Genus 908. <i>Anoplarchus</i> , Gill.....	2421
2777. <i>atropurpureus</i> (Kittlitz).....	2422; 2869
Genus 908(a). <i>Alectrias</i> , Jordan & Evermann.....	2869
2778. <i>alectrolophus</i> (Pallas).....	2421; 2869
Genus 909. <i>Xiphistes</i> , Jordan & Starks.....	2423
2779. <i>ulvae</i> , Jordan & Starks.....	2423
2780. <i>chirus</i> (Jordan & Gilbert).....	2424
Genus 910. <i>Xiphidion</i> , Girard.....	2424
2781. <i>mucosum</i> , Girard.....	2425
2782. <i>rupestre</i> (Jordan & Gilbert).....	2426
Genus 911. <i>Cebedichthys</i> , Ayres.....	2426
2783. <i>violaceus</i> (Ayres).....	2427
Genus 912. <i>Plaglogrammus</i> , Bean.....	2427

## CLASS III. PISCES—Continued.

## ORDER BB. ACANTHOPTERI—Continued.

*Family CC. Blenniidae*—Continued.

	Page.
2784. hopkinsi, Bean .....	2428
Genus 913. <i>Opisthocentrus</i> , Kner .....	2428
2785. ocellatus (Tilesius) .....	2429
Genus 914. <i>Pholidapus</i> , Bean & Bean .....	2430
2786. dyhowskil (Steindachner) .....	2430
Genus 915. <i>Plectobanchus</i> , Gilbert .....	2431
2787. evides, Gilbert .....	2432
Genus 916. <i>Leptoclinus</i> , Gill .....	2432
2788. maculatus (Fries) .....	2433
Genus 917. <i>Poroclinus</i> , Bean .....	2433
2789. rothrocki, Bean .....	2434
Genus 918. <i>Lumpenus</i> , Reinhardt .....	2435
Subgenus <i>Anisarchus</i> , Gill .....	2435
2790. medius (Reinhardt) .....	2435
Subgenus <i>Lumpenus</i> .....	2436
2791. anguillaris (Pallas) .....	2436
2792. mackayi (Gilbert) .....	2436
2793. fabricii (Cuvier & Valenciennes) .....	2437
2794. lampetreformis (Walbaum) .....	2438
Genus 919. <i>Stichæus</i> , Reinhardt .....	2439
2795. punctatus (Fabricius) .....	2439
Genus 920. <i>Ulvaria</i> , Jordau & Evermann .....	2440
2796. subbifurcata (Storer) .....	2440
Genus 921. <i>Eumesogrammus</i> , Gill .....	2441
2797. præcisus (Krøyer) .....	2441
<i>Family CCI. Cryptacanthoidæ</i> .....	2442
Genus 922. <i>Delolepis</i> , Bean .....	2442
2798. virgatus, Bean .....	2442
Genus 923. <i>Cryptacanthodes</i> , Storer .....	2443
2799. maculatus, Storer .....	2443
Genus 924. <i>Lyconectes</i> , Gilbert .....	2444
2800. aleutensis, Gilbert .....	2444
<i>Family CCII. Anarhichadidæ</i> .....	2445
Genus 925. <i>Anarhichas</i> (Artdi) Linnæus .....	2445
2801. latifrons, Steenstrup & Hallgrímsson .....	2446
2802. minor, Olafson .....	2446
2803. lupus, Linnæus .....	2446
2804. lepturus, Bean .....	2447
2805. orientalis, Pallas .....	2447
Genus 926. <i>Anarhichthys</i> , Ayres .....	2447
2806. ocellatus, Ayres .....	2448
<i>Family CCIII. Cerdalidæ</i> .....	2448
Genus 927. <i>Cerdale</i> , Jordan & Gilbert .....	2448
2807. lonthas, Jordan & Gilbert .....	2449
Genus 928. <i>Microdesmus</i> , Günther .....	2450
2808. dipus, Günther .....	2450
2809. retropinnis, Jordan & Gilbert .....	2450
<i>Family CCIV. Ptilichthyidæ</i> .....	2451
Genus 929. <i>Ptilichthys</i> , Bean .....	2452
2810. goodii, Bean .....	2452
Group <i>Ophidioidæ</i> .....	2453
<i>Family CCV. Scytalinidæ</i> .....	2453
Genus 930. <i>Scytalina</i> , Jordan & Gilbert .....	2454
2811. cerdale, Jordan & Gilbert .....	2454
<i>Family CCVI. Zoarcidæ</i> .....	2455
Genus 931. <i>Zoarces</i> , Gill .....	2456

## CLASS III. PISCES—Continued.

## ORDER BB. ACANTHOPTERI—Continued.

Family CCVI. *Zoaridae*—Continued.

	Page.
Subgenus <i>Macrozoarces</i> , Gill.....	2457
2812. <i>anguillaris</i> (Peck) .....	2457
Genus 932. <i>Embryx</i> , Jordan & Evermann .....	2458
2813. <i>crassilabris</i> (Gilbert) .....	2458
2814. <i>crotalinus</i> (Gilbert).....	2458
Genus 933. <i>Lycodopsis</i> , Collett.....	2460
2815. <i>pacificus</i> (Collett).....	2460
Genus 934. <i>Aprodon</i> , Gilbert.....	2460
2816. <i>cortezianus</i> , Gilbert.....	2461
Genus 935. <i>Lycodes</i> , Reinhardt .....	2461
Subgenus <i>Lycodes</i> .....	2463
2817. <i>esmarkii</i> , Collett.....	2463
2818. <i>vahlil</i> , Reinhardt .....	2463
2819. <i>concolor</i> , Gill & Townsend .....	2463
2820. <i>zoarchus</i> , Goode & Bean.....	2464
2821. <i>reticulatus</i> , Reinhardt .....	2465
2822. <i>perspicillum</i> , Kröyer .....	2465
2823. <i>frigidus</i> , Collett .....	2465
2824. <i>terra-novæ</i> , Collett .....	2466
2825. <i>digitatus</i> , Gill & Townsend.....	2466
2826. <i>palearis</i> , Gilbert.....	2466
2827. <i>brevipes</i> , Bean .....	2467
Subgenus <i>Lycias</i> , Jordan & Evermann.....	2468
2828. <i>nebulosus</i> , Kröyer.....	2468
2829. <i>seminudus</i> , Reinhardt.....	2468
Genus 936. <i>Lycodalepis</i> , Bleeker.....	2468
2830. <i>polaris</i> (Sabine) .....	2468
2831. <i>mucosus</i> (Richardson).....	2470
Genus 937. <i>Lycenchelys</i> , Gill .....	2470
2832. <i>verrillii</i> (Goode & Bean).....	2470
2833. <i>paxillus</i> (Goode & Bean).....	2471
2834. <i>porifer</i> (Gilbert).....	2471; 2869
Genus 938. <i>Furcella</i> , Jordan & Evermann.....	2472
2835. <i>diaptera</i> (Gilbert).....	2472
Genus 939. <i>Lycodonus</i> , Goode & Bean.....	2473
2836. <i>mirabilis</i> , Goode & Bean.....	2474
Genus 940. <i>Lycinema</i> , Gilbert .....	2474
2837. <i>barbatum</i> , Gilbert.....	2474
Genus 941. <i>Bothrocara</i> , Bean.....	2475
2838. <i>pusilla</i> (Bean).....	2476
2839. <i>mollis</i> , Bean.....	2476
Genus 942. <i>Gymnelis</i> , Reinhardt.....	2477
2840. <i>viridis</i> (Fabricius) .....	2477
2841. <i>stigma</i> (Lay & Bennett).....	2477
Genus 943. <i>Lycocara</i> , Gill.....	2478
2842. <i>parrli</i> (Ross).....	2478
Genus 944. <i>Melanostigma</i> , Günther.....	2478
2843. <i>gelatinosum</i> , Günther.....	2479
2844. <i>paumelas</i> , Gilbert.....	2479
Family CCVII. <i>Derepodichthyidae</i> .....	2480
Genus 945. <i>Derepodichthys</i> , Gilbert .....	2480
2845. <i>alepidotus</i> , Gilbert .....	2480
Family CCVIII. <i>Ophidiidae</i> .....	2481
Genus 946. <i>Lepophichthium</i> , Gill.....	2482
2846. <i>marmoratum</i> (Goode & Bean).....	2482
2847. <i>emmelas</i> (Gilbert).....	2483

## CLASS III. PISCES—Continued.

## ORDER BB. ACANTHOPTERI—Continued.

Family OCVIII. *Ophidiidae*—Continued.

	Page.	
2457	2848. <i>stigmatistium</i> (Gilbert) .....	2483
2457	2849. <i>profunderum</i> (Gill) .....	2484
2458	2850. <i>corvinum</i> (Goode & Bean) .....	2484
2458	2851. <i>prorates</i> (Jordan & Bollman) .....	2485
2460	2852. <i>brevibarbe</i> (Cuvier) .....	2485
2460	2853. <i>pardale</i> (Gilbert) .....	2486
2461	2854. <i>microlepis</i> (Gilbert) .....	2480
2463	Genus 947. <i>Ophidiou</i> (Artedi) Linnaeus .....	2487
2463	2855. <i>beani</i> , Jordan & Gilbert .....	2487
2463	2856. <i>holbrookii</i> (Putnaun) .....	2487
2463	2857. <i>graellsii</i> , Poey .....	2488
2463	Genus 948. <i>Chilara</i> , Jordan & Evermann .....	2488
2463	2858. <i>taylori</i> (Girard) .....	2489
2463	Genus 949. <i>Rissola</i> , Jordan & Evermann .....	2489
2463	2859. <i>marginata</i> (Do Kay) .....	2489
2465	Genus 950. <i>Otophidium</i> , Gill .....	2490
2465	2860. <i>omostigma</i> (Jordan & Gilbert) .....	2490
2465	2861. <i>indetatigabile</i> , Jordan & Bollman .....	2490
2466	2862. <i>galeoides</i> (Gilbert) .....	2491
2466	Family CCIX. <i>Lycodapodidae</i> .....	2491
2466	Genus 951. <i>Lycodapus</i> , Gilbert .....	2492
2466	2863. <i>dermatinus</i> , Gilbert .....	2492
2466	2864. <i>fierasfer</i> , Gilbert .....	2493
2466	2865. <i>parviceps</i> , Gilbert .....	2493
2466	2866. <i>extensus</i> (Gilbert) .....	2494
2468	Family COX. <i>Fierasferidae</i> .....	2494
2468	Genus 952. <i>Fierasfer</i> , Cuvier .....	2495
2470	2867. <i>affinis</i> (Günther) .....	2495
2470	2868. <i>arenicola</i> , Jordan & Gilbert .....	2496
2470	2869. <i>bermudensis</i> (Jones) .....	2497
2471	Family COXI. <i>Brotulidae</i> .....	2498
2471	Genus 953. <i>Brotula</i> , Cuvier .....	2500
2471	2870. <i>harbata</i> (Bloch & Schneider) .....	2500
2472	Genus 954. <i>Stygicola</i> , Gill .....	2500
2472	2871. <i>dentatus</i> (Poey) .....	2500
2473	Genus 955. <i>Lucifuga</i> , Poey .....	2501
2474	2872. <i>subterraneus</i> , Poey .....	2501
2474	Genus 956. <i>Brosomphycis</i> , Gill .....	2502
2474	2873. <i>marginatus</i> (Ayres) .....	2502
2475	Genus 957. <i>Ogilbia</i> , Jordan & Evermann .....	2502
2476	2874. <i>ventralis</i> (Gill) .....	2503
2476	2875. <i>cayorum</i> , Evermann & Kendall .....	2503
2477	Genus 958. <i>Bythites</i> , Reinhardt .....	2504
2477	2876. <i>fuscus</i> , Reinhardt .....	2504
2478	Genus 959. <i>Catetyx</i> , Günther .....	2504
2478	2877. <i>rubirostris</i> , Gilbert .....	2505
2478	Genus 960. <i>Dicromita</i> , Goode & Bean .....	2506
2479	2878. <i>agassizii</i> , Goode & Bean .....	2506
2479	Genus 961. <i>Bassozetus</i> , Gill .....	2507
2480	2879. <i>normalis</i> , Gill .....	2507
2480	2880. <i>compressus</i> (Günther) .....	2508
2480	2881. <i>catena</i> , Goode & Bean .....	2509
2481	2882. <i>tenia</i> (Günther) .....	2510
2482	Genus 962. <i>Muebia</i> , Goode & Bean .....	2510
2482	2883. <i>promelas</i> (Gilbert) .....	2511
2483	Genus 963. <i>Neobythites</i> , Goode & Bean .....	2512

## CLASS III. PISCES—Continued.

## ORDER BB. ACANTHOPTERI—Continued.

	Page.
<i>Family CCXI. Brotulidæ—Continued.</i>	
2884. gillii, Goode & Bean.....	2512
2885. marginatus, Goode & Bean.....	2513
Genus 964. Benthocometes, Goode & Bean.....	2513
2886. robustus, Goode & Bean.....	2514
Genus 965. Bassogigas, Gill.....	2515
2887. gillii, Goode & Bean.....	2515
2888. stelliferoides (Gilbert).....	2516
Genus 966. Barathrodemus, Goode & Bean.....	2517
2889. manatinus, Goode & Bean.....	2517
Genus 967. Nematonus, Günther.....	2518
2890. pectoralis, Goode & Bean.....	2518
Genus 968. Porogadus, Goode & Bean.....	2519
2891. miles, Goode & Bean.....	2520
Genus 969. Penopus, Goode & Bean.....	2520
2892. macdonaldi, Goode & Bean.....	2521
Genus 970. Dierolone, Goode & Bean.....	2522
2893. intronigra, Goode & Bean.....	2522
Genus 971. Mixonus, Günther.....	2523
2894. laticeps (Günther).....	2523
Genus 972. Barathronus, Goode & Bean.....	2524
2895. bicolor, Goode & Bean.....	2524
Genus 973. Aphyonus, Günther.....	2525
2896. mollis, Goode & Bean.....	2525
<i>Family CCXII. Bregmacerotidæ</i> .....	
Genus 974. Bregmaceros, Thompson.....	2526
2897. maclellandii, Thompson.....	2526
2898. atlanticus, Goode & Bean.....	2527
SUBORDER ANACANTHINI.....	
<i>Family CCXIII. Merlucciidæ</i> .....	
Genus 975. Merluccius, Rafinesque.....	2529
2899. merluccius (Linnaeus).....	2530
2900. bilinearis (Mitchill).....	2530
2901. productus (Ayres).....	2531
<i>Family CCXIV. Gadidæ</i> .....	
Genus 976. Boreogadus, Günther.....	2533
2902. saida (Lepechin).....	2533
Genus 977. Pollachius, Nilsson.....	2534
2903. virens (Linnaeus).....	2534
Genus 978. Theragra, Lucas.....	2535
2904. chalcogramma (Pallas).....	2535
2905. fucensis (Jordan & Gilbert).....	2536
Genus 979. Eleginus, Fischer.....	2537
2906. navaga (Kölreuter).....	2537
Genus 980. Microgadus, Gill.....	2538
2907. proximus (Girard).....	2539
2908. tomcod (Walbaum).....	2540
Genus 981. Gadus (Artemis) Linnaeus.....	2540
2909. callarias, Linnaeus.....	2541
2910. macrocephalus, Tilesius.....	2541
2911. ogac, Richardson.....	2542
Genus 982. Molanogrammus, Gill.....	2542
2912. æglefinus, Linnaeus.....	2542
Genus 983. Lepedion, Swainson.....	2543
2913. vercundum, Jordan & Craner.....	2543
Genus 984. Antimora, Günther.....	2544
2914. viola (Goode & Bean).....	2544

## CLASS III. PISCES—Continued.

## ORDER BB. ACANTHOPTERI—Continued.

Family CCXIV. *Gadidæ*—Continued.

Page.		Page.
2512	2915. <i>microlepis</i> , Bean.....	2545
2513	Genus 985. <i>Uraleptus</i> , Costa.....	2545
2513	2916. <i>maraldi</i> (Risso).....	2545
2514	Genus 986. <i>Lotella</i> , Kaup.....	2546
2515	2917. <i>maxillaris</i> , Bean.....	2546
2515	Genus 987. <i>Physiculus</i> , Kaup.....	2547
2516	2918. <i>fulvus</i> , Bean.....	2547
2517	2919. <i>nematopus</i> , Gilbert.....	2548
2517	2920. <i>kaupi</i> , Poey.....	2548
2518	2921. <i>rastrelliger</i> , Gilbert.....	2549
2518	Genus 988. <i>Lota</i> (Cuvier) Oken.....	2550
2519	2922. <i>maculosa</i> (Le Sueur).....	2550
2520	Genus 989. <i>Molva</i> , Fleming.....	2551
2520	2923. <i>molva</i> (Linnaeus).....	2551
2521	Genus 990. <i>Urophycis</i> , Gill.....	2552
2522	Subgenus <i>Urophycis</i> .....	2553
2522	2924. <i>regius</i> (Walbaum).....	2553
2523	2925. <i>cirratus</i> (Goode & Bean).....	2553
2523	2926. <i>floridanus</i> (Bean & Dresel).....	2554
2524	Subgenus <i>Emphycus</i> , Jordan & Evermann.....	2554
2524	2927. <i>carlli</i> (Bean).....	2554
2525	2928. <i>tenuis</i> (Mitchill).....	2555
2525	2929. <i>elmus</i> (Walbaum).....	2555
2525	2930. <i>chesteri</i> (Goode & Bean).....	2556
2526	Genus 991. <i>Læmonema</i> , Günther.....	2556
2526	2931. <i>barbatulum</i> , Goode & Bean.....	2556
2527	2932. <i>melanurum</i> , Goode & Bean.....	2557
2528	Genus 992. <i>Gaidropsarus</i> , Rafinesque.....	2557
2529	2933. <i>ensis</i> (Reinhardt).....	2558
2529	2934. <i>argentatus</i> (Reinhardt).....	2559
2530	2935. <i>septentrionalis</i> (Collett).....	2559
2531	Genus 993. <i>Euchelyopsis</i> , Blech & Schneider.....	2560
2531	2936. <i>cimbrius</i> (Linnaeus).....	2560
2533	Genus 994. <i>Brosme</i> (Cuvier) Oken.....	2561
2533	2937. <i>brosme</i> (Müller).....	2561
2534	Family CCXV. <i>Macrouridæ</i> .....	2561
2534	Genus 995. <i>Bathygadus</i> , Günther.....	2563
2535	2938. <i>arenatus</i> , Goode & Bean.....	2564
2535	2939. <i>favosus</i> , Goode & Bean.....	2565
2536	2940. <i>macrops</i> , Goode & Bean.....	2566
2537	2941. <i>longifilis</i> , Goode & Bean.....	2566
2537	Genus 996. <i>Steindachneria</i> , Goode & Bean.....	2567
2538	2942. <i>argentea</i> , Goode & Bean.....	2568
2539	Genus 997. <i>Trachyrinchus</i> , Giorna.....	2563
2540	2943. <i>helolepis</i> , Gilbert.....	2569
2541	Genus 998. <i>Malacocephalus</i> , Günther.....	2569
2541	2944. <i>occidentalis</i> , Goode & Bean.....	2570
2542	Genus 999. <i>Moseleya</i> , Goode & Bean.....	2570
2542	2945. <i>cyclolepis</i> (Gilbert).....	2570
2543	Genus 1000. <i>Nematonurus</i> , Günther.....	2571
2544	2946. <i>goodei</i> (Günther).....	2571
2544	2947. <i>suborbitalis</i> (Gill & Townsend).....	2572
2544	Genus 1001. <i>Albatrossia</i> , Jordan & Evermann.....	2573
2544	2948. <i>pectoralis</i> (Gilbert).....	2573
2544	Genus 1002. <i>Bogoslovius</i> , Jordan & Evermann.....	2574
2544	2949. <i>clarki</i> , Jordan & Gilbert.....	2575

## CLASS III. PISCES—Continued.

## ORDER BB. ACANTHOPTERI—Continued.

<i>Family CCXV. Macrouridae</i> —Continued.	Page.
2950. firmisquamis (Gill & Townsend) .....	2575
Genus 1003. Chalinura, Goode & Bean .....	2576
2951. serrula, Bean .....	2576
2952. filifera, Gilbert .....	2577
2953. sinula, Goode & Bean .....	2578
Genus 1004. Coryhanoides, Gunner .....	2578
2954. rupestris, Gunner .....	2579
2955. carapinus, Goode & Bean .....	2579
Genus 1005. Hymenoccephalus, Giglioli .....	2580
2956. cavernosus (Goode & Bean) .....	2580
Genus 1006. Macrourus, Bloch .....	2581
2957. berglax, Lacépède .....	2582
2958. holotrachys, Günther .....	2582
2959. bairdii, Goode & Bean .....	2583
2960. lepturus, Gill & Townsend .....	2584
2961. acrolepis, Bean .....	2585
2962. stolidolepis, Gilbert .....	2585
2963. cinereus, Gilbert .....	2586
Genus 1007. Cælorhynchus, Giorna .....	2587
2964. oeca (Goode & Bean) .....	2588
2965. carminatus (Goode) .....	2588
2966. caribbæus (Goode & Bean) .....	2589
2967. scaphopsis (Gilbert) .....	2590
Genus 1008. Trachonurus, Günther .....	2591
2968. sulcatus (Goode & Bean) .....	2591
Genus 1009. Lionurus, Günther .....	2592
2969. filicanda (Günther) .....	2592
2970. liolepis, Gilbert .....	2593
SUBORDER TÆNIOSOMI .....	2594
<i>Family CCXVI. Regalecidae</i> .....	2595
Genus 1010. Regalecus, Brünnich .....	2595
2971. glæne (Ascanius) .....	2596
✓ <i>Family CCXVII. Trachypterida</i> .....	2597
Genus 1011. Trachypterus, Gouan .....	2599
—2972. rex-salmonorum, Jordan & Gilbert .....	2599
—2973. trachyrus, Poey .....	2600
<i>Family CCXVIII. Stylephoridae</i> .....	2601
Genus 1012. Stylephorus, Shaw .....	2601
2974. chordatus, Shaw .....	2601
SUBORDER HETEROSOMATA .....	2602
<i>Family CCXIX. Pleuronectida</i> .....	2602
Genus 1013. Atheresthes, Jordan & Gilbert .....	2609
2975. stomias (Jordan & Gilbert) .....	2609
Genus 1014. Reinhardtius, Gill .....	2610
2976. hippoglossoides (Walbaum) .....	2611
Genus 1015. Hippoglossus, Cuvier .....	2611
2977. hippoglossus (Linnaeus) .....	2611
Genus 1016. Lyopsetta, Jordan & Goss .....	2612
2978. exilis (Jordan & Gilbert) .....	2612
Genus 1017. Eopsetta, Jordan & Goss .....	2613
2979. jordani (Lockington) .....	2613
Genus 1018. Hippoglossoides, Gottsche .....	2614
2980. platessoides (Fabricius) .....	2614
2981. elassodon, Jordan & Gilbert .....	2615
2982. robustus, Gill & Townsend .....	2616
2983. hamiltoni, Jordan & Gilbert .....	2616

## CLASS III. PISCES—Continued.

## ORDER BB. ACANTHOPTERI—Continued.

Family COXIX. *Pleuronectidae*—Continued.

Page.	Page.
575	Genus 1019. <i>Psettichthys</i> , Girard ..... 2617
576	2084. <i>melanostictus</i> , Girard ..... 2618
576	Genus 1020. <i>Verasper</i> , Jordan & Gilbert ..... 2618
577	2985. <i>moseri</i> , Jordan & Gilbert ..... 2619
578	Genus 1021. <i>Hippoglossina</i> , Steindachner ..... 2620
578	2986. <i>stomata</i> , Eigenmann & Eigenmann ..... 2620
579	2987. <i>macrops</i> , Steindachner ..... 2621
2579	2988. <i>bolimani</i> , Gilbert ..... 2621
2580	Genus 1022. <i>Lloglossina</i> , Gilbert ..... 2622
2580	2989. <i>tetropthalma</i> , Gilbert ..... 2622
2581	Genus 1023. <i>Xystreurus</i> , Jordan & Gilbert ..... 2623
2582	2990. <i>holopsis</i> , Jordan & Gilbert ..... 2623
2582	Genus 1024. <i>Paralichthys</i> , Girard ..... 2624
2583	2991. <i>californicus</i> (Ayres) ..... 2625
2584	2991(a). <i>Paralichthys magdalena</i> , Abbott ..... 2627
2585	2992. <i>restuarinus</i> , Gilbert & Scofield ..... 2626
2586	2993. <i>brasiliensis</i> (Ranzani) ..... 2626
2587	2994. <i>shaloe</i> , Jordan & Abbott ..... 2627, 2627
2588	2995. <i>woolmani</i> , Jordan & Williams ..... 2628
2588	2996. <i>dentatus</i> (Linnaeus) ..... 2629
2589	2997. <i>lethostigmus</i> , Jordan & Gilbert ..... 2630
2590	2998. <i>squamilentus</i> , Jordan & Gilbert ..... 2631
2591	2999. <i>albignatus</i> , Jordan & Gilbert ..... 2631
2591	3000. <i>oblongus</i> (Mitchill) ..... 2632
2592	Genus 1025. <i>Ramularia</i> , Jordan & Evermann ..... 2633
2592	3001. <i>dendritica</i> (Gilbert) ..... 2633
2593	Genus 1026. <i>Ancylopsetta</i> , Gill ..... 2634
2594	3002. <i>quadrocellata</i> , Gill ..... 2634
2595	Genus 1027. <i>Notosema</i> , Goode & Bean ..... 2635
2595	3003. <i>dilectum</i> , Goode & Bean ..... 2635
2596	Genus 1028. <i>Gastropsetta</i> , B. A. Bean ..... 2636
2597	3004. <i>frontalis</i> , B. A. Bean ..... 2636
2599	Genus 1029. <i>Pleuronichthys</i> , Girard ..... 2637
2599	3005. <i>decurrens</i> , Jordan & Gilbert ..... 2637
2600	3006. <i>verticalis</i> , Jordan & Gilbert ..... 2638
2601	3007. <i>cenosus</i> , Girard ..... 2638
2601	Genus 1030. <i>Hypsopsetta</i> , Gill ..... 2639
2601	3008. <i>guttulata</i> (Girard) ..... 2639
2602	Genus 1031. <i>Parophrys</i> , Girard ..... 2640
2602	3009. <i>vetulus</i> , Girard ..... 2640
2609	Genus 1032. <i>Inopsetta</i> , Jordan & Goss ..... 2641
2609	3010. <i>ischyra</i> (Jordan & Gilbert) ..... 2641
2610	Genus 1033. <i>Isopsetta</i> , Lockington ..... 2642
2610	3011. <i>isolepis</i> (Lockington) ..... 2642
2611	Genus 1034. <i>Lepidopsetta</i> , Gill ..... 2642
2611	3012. <i>bilineata</i> (Ayres) ..... 2643
2612	Genus 1035. <i>Limanda</i> , Gotsche ..... 2644
2612	3013. <i>ferruginea</i> (Storer) ..... 2644
2613	3014. <i>aspera</i> (Pallas) ..... 2645
2614	3015. <i>proboscidea</i> , Gilbert ..... 2645
2614	3016. <i>beauii</i> , Goode ..... 2646
2615	Genus 1036. <i>Pseudopleuronectes</i> , Bleeker ..... 2646
2616	3017. <i>americanus</i> (Walbaum) ..... 2647
2616	3018. <i>pinnifasciatus</i> (Kner) ..... 2647
2616	Genus 1037. <i>Pleuronectes</i> (Arctid) Linnaeus ..... 2648
2616	3019. <i>quadrituberculatus</i> , Pallas ..... 2648



## CLASS III. PISCES—Continued.

## ORDER BB. ACANTHOPTERI—Continued.

Family CXXIX. *Pleuronectidae*—Continued.

	Page.
Genus 1038. <i>Liopsetta</i> , Gill .....	2649
3020. <i>glucialis</i> (Pallas).....	2649
3021. <i>putnami</i> (Gill).....	2650
3022. <i>obscura</i> (Herzenstein).....	2651
Genus 1039. <i>Platichthys</i> , Girard .....	2651
3023. <i>stellatus</i> (Pallas) .....	2652
Genus 1040. <i>Microstomus</i> , Gottsche.....	2653
3024. <i>kitt</i> (Walbaum) .....	2654
3025. <i>pacificus</i> (Lockington) .....	2655
Genus 1041. <i>Eubassichthys</i> , Jordan & Evermann .....	2655
3026. <i>bathybius</i> (Gilbert) .....	2655
Genus 1042. <i>Glyptocephalus</i> , Gottsche.....	2656
3027. <i>cynoglossus</i> (Linnaeus).....	2657
3028. <i>zachirus</i> , Lockington.....	2658
Genus 1043. <i>Lophopsetta</i> , Gill.....	2659
3029. <i>maculata</i> (Mitchill).....	2660
Genus 1044. <i>Platophrys</i> , Swainson.....	2660
3030. <i>spinousus</i> (Poey) .....	2662
3031. <i>constellatus</i> , Jordan.....	2663
3032. <i>ocellatus</i> (Agassiz) .....	2663
3033. <i>maculifer</i> (Poey) .....	2664
3034. <i>ellipticus</i> (Poey).....	2665
3035. <i>lunatus</i> (Linnaeus) .....	2665
3036. <i>leopardinus</i> (Günther).....	2666
Genus 1045. <i>Perissias</i> , Jordan & Evermann.....	2667
3037. <i>tæniopterus</i> (Gilbert) .....	2667
Genus 1046. <i>Engyphrys</i> , Jordan & Bollman .....	2668
3038. <i>santi-laurentii</i> , Jordan & Bollman .....	2668
Genus 1047. <i>Trichopsetta</i> , Gill.....	2669
3039. <i>ventralis</i> (Goode & Bean).....	2669
Genus 1048. <i>Syacium</i> , Ranzani.....	2670
3040. <i>papillosum</i> (Linnaeus) .....	2671
3041. <i>micrurum</i> , Ranzani.....	2672
3042. <i>latifrons</i> (Jordan & Gilbert).....	2673
3043. <i>ovale</i> (Günther) .....	2674
Genus 1049. <i>Cyclosetta</i> , Gill.....	2675
3044. <i>querna</i> (Jordan & Bollman).....	2675
3045. <i>chittendeni</i> , B. A. Bean .....	2676
3046. <i>imbriata</i> (Goode & Bean).....	2676
Genus 1050. <i>Azevia</i> , Jordan .....	2677
3047. <i>panamensis</i> (Steindachner).....	2677
Genus 1051. <i>Citharichthys</i> , Bleeker.....	2678
Subgenus <i>Orthopsetta</i> , Gill.....	2679
3048. <i>sordidus</i> (Girard).....	2679
3049. <i>fragilis</i> , Gilbert .....	2680
3050. <i>xanthostigmus</i> , Gilbert .....	2680
3051. <i>stigmaeus</i> , Jordan & Gilbert.....	2681
Subgenus <i>Citharichthys</i> .....	2682
3052. <i>dioceros</i> , Goode & Bean .....	2682
3053. <i>platophrys</i> , Gilbert.....	2683
3054. <i>aretrifrons</i> , Goode.....	2683
3055. <i>unicornis</i> , Goode .....	2683
3056. <i>uhleri</i> , Jordan.....	2684
3057. <i>macrops</i> , Dressel.....	2684
3058. <i>spilopterus</i> , Günther .....	2685
3059. <i>gilberti</i> , Jenkins & Evermann .....	2686

## CLASS III. PISCES—Continued.

## ORDER BB. ACANTHOPTERI—Continued.

Family CCXXIX. *Plenrocetidae*—Continued.

	Page.
Genus 1052. <i>Etopus</i> , Jordan & Gilbert.....	2687
3060. <i>microstomus</i> (Gill).....	2687
3061. <i>rimosus</i> , Goode & Bean.....	2688
3062. <i>crossotus</i> , Jordan & Gilbert.....	2689
Genus 1053. <i>Monoleus</i> , Goode.....	2690
3063. <i>scasilicuda</i> , Goode.....	2691
3064. <i>atrimum</i> , Goode & Bean.....	2692

Family CCXX. *Soleidae*.....

Genus 1054. <i>Achirus</i> , Lacépède.....	2693
Subgenus <i>Balostoma</i> , Bean.....	2695
3065. <i>achirus</i> (Linnaeus).....	2695
3066. <i>inscriptus</i> , Gosse.....	2696
3067. <i>klunzingeri</i> (Steindachner).....	2697
3068. <i>lineatus</i> (Linnaeus).....	2697
3069. <i>mazatlanus</i> (Steindachner).....	2698
3070. <i>fonsecensis</i> (Günther).....	2699
3071. <i>fischeri</i> (Steindachner).....	2699
3072. <i>scutum</i> (Günther).....	2700
Subgenus <i>Achirus</i> .....	2700
3073. <i>fimbriatus</i> (Günther).....	2700
3074. <i>fasciatus</i> , Lacépède.....	2700
3075. <i>panamensis</i> (Steindachner).....	2702
Genus 1055. <i>Apionichthys</i> , Kaup.....	2702
3076. <i>unicolor</i> (Günther).....	2702
Genus 1056. <i>Gymnachirus</i> , Kaup.....	2703
3077. <i>fasciatus</i> , Günther.....	2703
Genus 1057. <i>Symphurus</i> , Rafinesque.....	2704
Subgenus <i>Symphurus</i> .....	2705
3078. <i>piger</i> (Goode & Bean).....	2705
3079. <i>marginatus</i> (Goode & Bean).....	2706
3080. <i>atramentatus</i> , Jordan & Bollman.....	2706
3081. <i>fasciolaris</i> , Gilbert.....	2707
3082. <i>elongatus</i> (Günther).....	2707
3083. <i>atricaudus</i> (Jordan & Gilbert).....	2707
3084. <i>leel</i> , Jordan & Bollman.....	2708
3085. <i>plagusia</i> (Bloch & Schneider).....	2709
3086. <i>plagusia</i> (Linnaeus).....	2710
3087. <i>puellus</i> (Goode & Bean).....	2710
3088. <i>diomedeanus</i> (Goode & Bean).....	2711
3089. <i>williamsi</i> , Jordan & Culver.....	2711
Subgenus <i>Acedia</i> , Jordan.....	2712
3090. <i>nebulosus</i> (Goode & Bean).....	2712

## ORDER CC. PEDICULATI.....

Family CCXXI. *Lophiidae*.....

Genus 1058. <i>Lophius</i> (Arted) Linnaeus.....	2713
3091. <i>piecatorius</i> , Linnaeus.....	2713
Genus 1059. <i>Lophiomus</i> , Gill.....	2714
3092. <i>setigerus</i> (Vahl).....	2714

Family CCXXII. *Antennariidae*.....

Genus 1060. <i>Pterophryne</i> , Gill.....	2715
3093. <i>histris</i> (Linnaeus).....	2716
3094. <i>gibba</i> (Mitchill).....	2717
Genus 1061. <i>Antennarius</i> , Lacépède.....	2717
3095. <i>inops</i> , Poey.....	2718
3096. <i>principis</i> (Cuvier & Valenciennes).....	2719
3097. <i>tenebrosus</i> (Poey).....	2719

2690  
2691  
2692  
2693  
2694  
2695  
2696  
2697  
2698  
2699  
2700  
2701  
2702  
2703  
2704  
2705  
2706  
2707  
2708  
2709  
2710  
2711  
2712  
2713  
2714  
2715  
2716  
2717  
2718  
2719  
2720  
2721  
2722  
2723  
2724  
2725  
2726  
2727  
2728  
2729  
2730  
2731  
2732  
2733  
2734  
2735  
2736  
2737  
2738  
2739  
2740  
2741  
2742  
2743  
2744  
2745  
2746  
2747  
2748  
2749  
2750  
2751  
2752  
2753  
2754  
2755  
2756  
2757  
2758  
2759  
2760  
2761  
2762  
2763  
2764  
2765  
2766  
2767  
2768  
2769  
2770  
2771  
2772  
2773  
2774  
2775  
2776  
2777  
2778  
2779  
2780  
2781  
2782  
2783  
2784  
2785  
2786

## CLASS III. PISCES—Continued.

## ORDER CC. PEDICULATI—Continued.

Family CCXXII. *Antennariidae*—Continued.

	Page.
3098. <i>reticularis</i> , Gilbert .....	2719
3099. <i>strigatus</i> , Gill .....	2720
3100. <i>sanguineus</i> , Gill .....	2721
3101. <i>ocellatus</i> (Bloch & Schneider) .....	2721
3102. <i>scaber</i> (Cuvier) .....	2722
3103. <i>tigris</i> , Poey .....	2723
3104. <i>nuttingii</i> , Garman .....	2723
3105. <i>multiocellatus</i> (Cuvier & Valenciennes) .....	2724
3106. <i>radiosus</i> , Garman .....	2725
Genus 1062. <i>Chaunax</i> , Lowe .....	2726
3107. <i>pietus</i> , Lowe .....	2726
3108. <i>nuttingii</i> , Garman .....	2727
Family CCXXIII. <i>Ceratiidae</i> .....	2727
Genus 1063. <i>Ceratinus</i> , Kröyer .....	2729
3109. <i>holbollahi</i> , Kröyer .....	2729
Genus 1064. <i>Mancallas</i> , Gill .....	2729
3110. <i>uranoscopus</i> (Murray) .....	2729
3111. <i>shufeldti</i> (Gill) .....	2730
Genus 1065. <i>Cryptopsaras</i> , Gill .....	2731
3112. <i>conceili</i> , Gill .....	2731
Genus 1066. <i>Oncirodes</i> , Lütken .....	2732
3113. <i>esrichtii</i> , Lütken .....	2732
Genus 1067. <i>Himantolophus</i> , Reinhardt .....	2732
3114. <i>grœnlandicus</i> , Reinhardt .....	2733
Genus 1068. <i>Corynolophus</i> , Gill .....	2733
3115. <i>reinhardti</i> (Lütken) .....	2733
Genus 1069. <i>Lloceus</i> , Günther .....	2733
3116. <i>murrayi</i> (Günther) .....	2733
Genus 1070. <i>Linophryne</i> , Collett .....	2734
3117. <i>lucifer</i> , Collett .....	2734
Genus 1071. <i>Caulophryne</i> , Goode & Bean .....	2734
3118. <i>jordani</i> , Goode & Bean .....	2735
Family CCXXIV. <i>Ogcocephalidae</i> .....	2735
Genus 1072. <i>Ogcocephalus</i> , Fischer .....	2736
3119. <i>vespertilio</i> (Linnaeus) .....	2737
3120. <i>nasutus</i> (Cuvier & Valenciennes) .....	2737
3121. <i>radiatus</i> (Mitchill) .....	2738
Genus 1073. <i>Zalieutes</i> , Jordan & Evermann .....	2738
3122. <i>elater</i> (Jordan & Gilbert) .....	2738
Genus 1074. <i>Haliutichthys</i> , Poey .....	2739
3123. <i>aculeatus</i> (Mitchill) .....	2739
3124. <i>caribbaeus</i> , Garman .....	2741
Genus 1075. <i>Haliutæa</i> , Cuvier & Valenciennes .....	2741
3125. <i>spongiosa</i> , Gilbert .....	2741
Genus 1076. <i>Haliutella</i> , Goode & Bean .....	2742
3126. <i>lappa</i> , Goode & Bean .....	2742
Genus 1077. <i>Dibranchus</i> , Peters .....	2743
3127. <i>atlanticus</i> , Peters .....	2743

1  
spe  
Sicy  
Eury  
Gna  
Dact  
Dac  
Brys  
Gobi  
Cora  
Blen  
Hom  
Hom  
Scar  
Ophi  
Embl  
Eud  
Eubr  
Lyci  
Emph  
Albat  
Bogos  
Hippo  
Veras  
Veras  
Rauu  
Periss  
Carch  
Carch  
Mylio  
Aspis  
Galeic  
Tachy  
Aztec  
Notroy  
Notroy  
Pisood  
Murre  
Stolep  
Stolep  
Stolep  
Stolep  
Ceterg  
Argyr  
Os.ner  
Bathy

Page.  
 .. 2710  
 .. 2720  
 .. 2721  
 .. 2721  
 .. 2722  
 .. 2723  
 .. 2723  
 .. 2724  
 .. 2725  
 .. 2726  
 .. 2726  
 .. 2727  
 .. 2727  
 .. 2729  
 .. 2729  
 .. 2729  
 .. 2729  
 .. 2730  
 .. 2731  
 .. 2731  
 .. 2732  
 .. 2732  
 .. 2732  
 .. 2733  
 .. 2733  
 .. 2733  
 .. 2733  
 .. 2734  
 .. 2734  
 .. 2735  
 .. 2735  
 .. 2736  
 .. 2737  
 .. 2737  
 .. 2738  
 .. 2738  
 .. 2738  
 .. 2739  
 .. 2739  
 .. 2741  
 .. 2741  
 .. 2741  
 .. 2742  
 .. 2742  
 .. 2743  
 .. 2743

## LIST OF NEW NAMES.

The following is a list of the new generic, subgeneric, specific, and sub-specific names which appear as new in Part III of the present work:

	Page.
Sleydium vincente, Jordan & Evermann.....	2207
Enypnius, Jordan & Evermann.....	2231
Gnathypops snyderi, Jordan & Evermann.....	2285
Daetyloscopus zelotes, Jordan & Gilbert.....	2303
Dector, Jordan & Evermann.....	2325
Bryssophilus, Jordan & Evermann.....	2329
Gobieex gyrinus, Jordan & Evermann.....	2331
Corallicolu, Jordan & Evermann.....	2369
Blenniulus, Jordan & Evermann.....	2386
Homesthes, Gilbert.....	2394
Homesthes caulopus, Gilbert.....	2394
Scartichthys, Jordan & Evermann.....	2395
Ophioblennius steindachneri, Jordan & Evermann.....	2401
Emblemaria atlantica, Jordan & Evermann.....	2402
Enedrius, Jordan & Gilbert.....	2414
Embryx, Jordan & Evermann.....	2458
Lycias, Jordan & Evermann.....	2461
Emphyens, Jordan & Evermann.....	2552
Albutrossia, Jordan & Evermann.....	2573
Bogosalovius, Jordan & Evermann.....	2574
Hippoglossoides hamiltoni, Jordan & Gilbert.....	2616
Verasper, Jordan & Gilbert.....	2618
Verasper moseri, Jordan & Gilbert.....	2619
Rumularia, Jordan & Evermann.....	2633
Perissias, Jordan & Evermann.....	2667
Carcharhinus cerdale, Gilbert.....	2746
Carcharhinus velox, Gilbert.....	2747
Myliobatls asperrimus, Gilbert.....	2754
Aspistor, Jordan & Evermann.....	2763
Galeichthys xenancheni, Gilbert.....	2777
Tachysurus emmelane, Gilbert.....	2785
Azteenia, Jordan & Evermann.....	2799
Notropis chamberlaini, Evermann.....	2800
Notropis louisiana, Evermann.....	2801
Pisodonopsis daspilotes, Gilbert.....	2803
Muraena clepsydra, Gilbert.....	2805
Stolephorus rastralis, Gilbert & Pierson.....	2811
Stolephorus mundeolus, Gilbert & Pierson.....	2812
Stolephorus naso, Gilbert & Pierson.....	2813
Stolephorus etarksi, Gilbert & Pierson.....	2813
Cetengraulis engyemen, Gilbert & Pierson.....	2815
Argyrosomus alascanus, Scofield.....	2817
Osmerus albutrossis, Jordan & Gilbert.....	2823
Bathylagus milleri, Jordan & Gilbert.....	2825

	Page.
Zaphotias, Goode & Bean.....	2826
Churaeodon garmani, Jordan & Evermann.....	2831
Siphostoma sinaloe, Jordan & Starks.....	2838
Rhynchelas, Gill.....	2841
Oligoplites mundus, Jordan & Starks.....	2844
Hemicaranx zelotes, Gilbert.....	2845
Ulocentra meadia, Jordan & Evermann.....	2852
Lobotes pacificus, Gilbert.....	2857
Porocottus bradfordi, Rutter.....	2862
Sigmistes Kutter.....	2863
Sigmistes emillas, Rutter.....	2863
Crystallichthys, Jordan & Gilbert.....	2864
Crystallichthys mirabilis, Jordan & Gilbert.....	2865
Allineetes, Jordan & Evermann.....	2866
Prognurus, Jordan & Evermann.....	2866
Prognurus cypselurus, Jordan & Gilbert.....	2866
Sicyosus, Jordan & Evermann.....	2867
Alectrias, Jordan & Evermann.....	2869
Furcinanus, Jordan & Evermann.....	2869
Salmo clarkii tahoensis, Jordan & Evermann.....	2870
Oligocottus snyderi, Greeley.....	2871
Flammeo, Jordan & Evermann.....	2871
Paralichthys magdalena, Abbott.....	2872
Petalichthys sinaloe, Jordan & Abbott.....	2872

fis  
 gro  
 So  
 the  
 sur  
 arc  
 T  
 gro  
 I  
 wa  
 inc  
 III  
 II,  
 Atl  
 rec  
 T  
 fish  
 que  
 glo  
 wo  
 A  
 mo  
 the  
 T  
 seni  
 him  
 as c  
 ich'  
 T  
 autl  
 with  
 Mus

ago.  
2826  
2831  
2838  
2841  
2844  
2845  
2852  
2857  
2862  
2863  
2863  
2864  
2865  
2866  
2866  
2866  
2867  
2869  
2869  
2870  
2871  
2871  
2872  
2872

# THE FISHES

OF

## NORTH AND MIDDLE AMERICA.

BY DAVID STARR JORDAN AND BARTON WARREN EVERMANN.

### PART III.

#### PREFATORY NOTE.

This volume is the third of a descriptive catalogue of the fishes and fish-like vertebrates of North and Middle America. For the sake of greater completeness the marine fishes of the Galapagos Islands and the South American coast north of the equator have been included, as all of these are sure, sooner or later, to be found within our limits. For the same reason the few species known from Kamchatka and the Kuril Islands are included as a part of the fauna of the Alaskan Sea.

The pagination and the numbering of the species, genera, and higher groups are continuous throughout the three parts.

Part I, *Branchiostomatidæ* to *Priacanthidæ* inclusive (pages 1 to 1240), was published October 3, 1896; Part II, *Lutianidæ* to *Cephalacanthidæ* inclusive (pages 1241 to 2182), was published October 3, 1898; and Part III, *Callionymidæ* to *Ogcocephalidæ* appears on November 26, 1898. Parts I, II, and III have each their own table of contents, while in Part IV (the Atlas) is given a table of contents complete for the entire work and corrected to include the Addenda.

The present part includes also an artificial key to the families of true fishes, an addendum containing species overlooked or described subsequently to the publication or casting of the part to which they belong, a glossary of scientific terms, and a general index complete for the entire work.

A fourth volume, or Atlas of plates, containing illustrations of one or more species of each of the more important genera, will follow within the year.

The preparation of the manuscript for this work was begun by the senior author in 1891. In 1893 the junior author became associated with him, and since then both have given to it such of their time and energy as could be spared from engrossing official duties to which systematic ichthyology bears no relation.

The insertion of the comma between generic and specific names and the authorities for them, as practiced in this publication, is in accordance with the views held by the authorities of the United States National Museum, and does not express the views of the authors of this work.

## Class PISCES—Concluded.

## Subclass TELEOSTOMI—Concluded.

## Order BB. ACANTHOPTERI—Concluded.

## Group GOBIOIDEA.

## (THE GOBIES.)

Body elongate, variously scaled or naked; head usually large, armed or not, the suborbital ring without a bony stay for the preopercle; gill openings reduced, the membranes attached to the isthmus. Gills 4, a slit behind the last; pseudobranchiae present. Ventral rays I, 4 or I, 5, inserted below pectoral, the fins close together or united or widely separated or otherwise peculiar; dorsal fins separate or united, the first of a few weak spines, sometimes wanting; anal rather long, usually with a single weak spine, similar to soft dorsal; caudal rounded. Usually no air bladder nor pyloric caeca. Vertebrae 24 to 35. Carnivorous bottom fishes, mostly of small size in warm regions, some marine, others of the fresh waters. Two families.

a. Ventral fins widely separated; preopercle strongly armed; lateral line present.

CALLIONYMIDÆ, CLXXXVII.

aa. Ventral fins close together, usually united; preopercle with a weak spine or none; no lateral line.

GOBIIDÆ, CLXXXVIII.

## Family CLXXXVII. CALLIONYMIDÆ.

## (THE DRAGONETS.)

Body elongate, naked; head usually broad and depressed; the mouth narrow, the upper jaw very protractile; teeth very small, in jaws only; preopercle armed with a strong spine, which is usually branched. Eyes moderate, usually directed upward. Lateral line present, often duplicated. Dorsal fins 2, the anterior with 3 or 4 flexible spines; soft dorsal and anal short, the latter without distinct spine; ventrals I, 5, widely separated from each other; pectoral fins large. Gill openings small, the membranes broadly attached to the isthmus; gills 4, a slit behind the fourth; pseudobranchiae present; no air bladder. Vertebrae usually  $8 + 13 = 21$ . Small fishes of the shores of warm seas, chiefly of the old world. Allied to the Gobies, but often resembling the *Cottidae* in form. Genera 4, species about 30. (*Gobiidae Callionymina*, Günther, Cat. Fishes, III, 138-152.)

a. Ventrals entire, the outer ray not detached; head depressed; gill opening reduced to a very small foramen on upper surface of head; lateral line single.

CALLIONYMUS, 799.

799. CALLIONYMUS, Linnaeus.

*Callionymus*, LINNAEUS, Syst. Nat., Ed. x, 249, 1758 (*lyra*).

This genus includes Dragonets with the ventral fins entire, without detached ray, the gill opening reduced to a small foramen opening upward, and the lateral line single; head triangular, depressed; eyes directed upward; preopercular spine very large; sexual differences strongly marked. Species numerous, living on sea bottoms at some depth. (*κάλλις*, beauty; *ὄνομα*, name.)

- a. Dorsal rays IV, 8 or 9; anal rays 8; some of the dorsal spines filamentous.
- b. Preopercular spine very long, armed with about 9 hooks or spinules; caudal not filamentous. BAIRDI, 2511.
- bb. Preopercular spine strong, bifurcate; caudal fin more or less produced or filamentous. HIMANTOPHORUS, 2512.
- aa. Dorsal rays III, 6 or IV, 6; anal rays 4.
- c. Preopercular spine with 2 barba, the anterior turned forward; body with white spots. CALLIURUS, 2513.
- cc. Preopercular spine with 3 teeth above, ending in an acute point. FAUCIRADIATUS, 2514.

2511. CALLIONYMUS BAIRDI, Jordan.

Head  $3\frac{1}{2}$ ; depth  $9\frac{1}{4}$ . D. IV, 9; A. 8. Body long and low, very slender, the head much depressed, the least depth of the caudal peduncle about equal to the diameter of the eye. Head triangular as viewed from above, its breadth  $\frac{2}{3}$  its length, exclusive of the preopercular spine. Snout bluntish as seen from above, sharp in profile, its outline straight and moderately steep until above the eyes; profile behind the eyes considerably depressed. Snout  $2\frac{1}{2}$  in head to gill opening; eye 4; mouth small, inferior, the maxillary reaching front of eye, as long as snout; lower lip conspicuous. Teeth slender, in villiform bands in both jaws, none on vomer. Interorbital area a simple narrow ridge. Bones of head behind eyes rugose; a low rough tubercle of bare bone above the temporal region on each side, somewhat behind each eye. Preopercular spine very long, as long as eye, its exterior ridge with a single antrorse spinule at its base, its posterior edge with 8 conspicuous hooks turned forward and inward, these growing progressively smaller from the second. Gill opening reduced to a pore at upper posterior angle of opercle, its width rather less than that of pupil. Dorsal spines strong, the first ending in a slender filament, the whole as long as head; second and third spines broken (probably each with a short filament in life, as a short filament is still present on the fourth spine); fourth spine well behind third (leaving room for another spine, although no trace of such spine is present); soft dorsal high, most of its rays slightly filamentous at tip, the longest about  $\frac{2}{3}$  head; caudal subtruncate, not filamentous, about as long as head to base of preopercular spine; anal fin rather high, the length of its base 3 in body; pectorals about as long as ventrals, each as long as head without preopercular spine. Lateral line single. Color light grayish, mottled or spotted with yellowish and dark brown; cheeks with steel-bluish spots; first dorsal with dusky reticulations around pale gray spots; second dorsal and caudal with nar-



row dusky cross streaks; anal with its posterior half chiefly black, the anterior pale; ventrals black; pectorals pale. Type, a specimen  $4\frac{1}{2}$  inches long, in good condition, from the "spewings" of a Snapper or a Grouper (*Neomenis aya* or *Epinephelus morio*), taken on the Snapper Banks, between Pensacola and Tampa; 1 other specimen known. ("I have named this species for Prof. Spencer F. Baird, to whom I have been indebted for aids of many kinds in connection with my studies of American fishes." Jordan.)

*Callionymus bairdi*, JORDAN, Proc. U. S. Nat. Mus. 1887, 501, Snapper Banks off Pensacola. (Type, No. 39300. Coll. Silas Stearns.)

2512. CALLIONYMUS HIMANTOPHORUS, Goode & Bean.

Head  $3\frac{1}{2}$ ; depth of head equal to length of its postorbital portion or to greatest depth of body. Greatest depth of body at the head and the anterior portion of the trunk. D. IV, 8; A. 8; P. 19; V. I, 5. Body slender, moderately elongate, fins all well developed, the tail tapering and with some of its rays produced into a filament. Caudal peduncle very slender, the least height of tail scarcely more than  $\frac{1}{2}$  greatest height of body. Profile descending very rapidly at snout. Mouth small and the intermaxillary very protractile, but may be almost entirely concealed under the pre-orbitals. Intermaxillary reaching to front of orbit. Maxillary a roundish, slender bone, extending backward to end of intermaxillary. Mandible about as long as eye, extending to vertical through front of pupil. Teeth, in villiform bands on intermaxillary and mandible. Interorbital space very narrow, less than  $\frac{1}{2}$  length of eye, which is  $1\frac{1}{2}$  times as long as snout and nearly  $\frac{1}{2}$  of total without caudal. A strong bifurcated spine at angle of the preoperculum extending backward slightly beyond the gill opening; length of this spine at its upper articulation  $\frac{2}{3}$  length of eye. Gill opening reduced to a small slit, placed at a distance behind eye about equaling length of eye and above median line of body. Skin naked. Lateral line abruptly arched over gill opening and connected across nape with its fellow of the opposite side. Spinous dorsal somewhat elevated in front, the first spine nearly twice as long as last, its length about  $\frac{1}{2}$  total length of caudal; sixth and seventh rays longest, their length nearly equaling that of base of fin; caudal consisting of 4 simple and 8 divided rays; of the divided rays the fifth and sixth are the longest, the lower portion of the fifth and the upper portion of the sixth being produced into a filament, making these rays as long as the distance from the tip of the intermaxillary to the fourth anal ray. It is worthy of remark that in another example of the same species and of about the same size as the type, the sixth of the divided rays alone contributes to form the filament; and in a young example, about  $\frac{1}{2}$  as large as the type, the first dorsal spine when laid back reaches to the end of soft dorsal. Some of the numerous examples of this species have none of the caudal rays much produced, even in large individuals. Anal fin beginning directly under third ray of soft dorsal, its rays increasing in length to the sixth, which is the longest and twice as long as the first, its length  $5\frac{1}{2}$

in total without caudal. All the rays simple except the last, which is divided. The pectoral beginning under middle of spinous dorsal and extending to below the fifth ray of the soft dorsal, its rays all simple. The ventral base overlapping lower extremity of pectoral base, its origin under the gill opening. The fourth and longest ray equaling  $\frac{1}{2}$  of total length without caudal. A small but distinct anal papilla. Color generally light brown, the back with numerous narrow streaks and blotches of slightly darker brown; a dark blotch on membrane between the third and fourth dorsal spines, in some cases occupying nearly all the membrane, in other cases more limited and nearly elliptical in shape; anal with a broad subvertical dark band, the tips of rays and a small area of the membrane behind each ray pale; the lower caudal lobe with a narrow submarginal dark band; ventral with 2 indistinct narrow dark bands on its outer half. From Blake Station XXX, off Barbados, in 209 fathoms; Station CLXXX, at 137 fathoms; Station XXXIII, off Santa Cruz, at 115 fathoms; Station 2CCXVI, at 119 fathoms; Station CCXXX, at 84 fathoms. (*ἴμας*, whip; *φορέω*, bear.)

*Callionymus himantophorus*,\* GOODE & BEAN, Ocean. Ichth., 296, pl. LXXVI, figs. 268, 268a, b, 1906, off Barbados.

2513. *CALLIONYMUS CALLIURUS*, Eigenmann & Eigenmann.

Head  $3\frac{1}{2}$  to tip of opercular spine (5 in total); depth 7 (9). D. IV, 6; A. 4. Body flat below, the ventral surface bordered on each side with a fold of skin which is wider than the pupil; a single lateral line; diameter of eye equaling length of snout,  $3\frac{1}{2}$  in head; maxillary not extending to eye; preopercular spine with 2 barbs above, the anterior one larger and turned forward; gill opening a minute foramen opening upward. The last dorsal ray equaling length of head, and the first dorsal spine reaching its tip when the fin is depressed; ventral fins connected by a broad membrane to the middle of the outer pectoral region; pectoral fins as long as the head. Cheeks, opercles, connecting membrane of ventral fins and antepectoral region with milk-white spots; lower jaw black near the rictus; a series of black dots on branchiostegal membranes, 1 or 2 similar dots in front of pectorals, 2 on the cheek forming a series with the second branchiostegal spot; 4 black spots on the marginal membrane of the belly, other black spots above it; lower half of body with numerous dirty white spots; pectorals transparent, ventrals dusky; membrane of anal sprinkled with minute black points aggregated into black spots in places, and with opaque white spots; caudal transparent, having minute points, its upper half with opaque milk-white bars running obliquely downward and backward from ray to ray; lower half with interrupted longitudinal lines of opaque white, alternating with black spots; dorsal transparent, with white and dark dots most conspicuous between last rays; body marbled with light and darker. Key West, Florida; 1 specimen dredged in 5 fathoms. (*κάλλος*, beauty; *ὄψα*, tail.)

*Callionymus calliurus*, EIGENMANN & EIGENMANN, Proc. Cal. Ac. Sci. 1888, 76, South Beach, Key West. (Type, No. 20265. M. C. Z.)

\* The species was listed by Eigenmann, Proc. Cal. Ac. Sci., 2d ser. 1, 78, as "*Callionymus agassizii*, Goode & Bean," a name only, accompanied by no description.

## 2514. CALLIONYMUS PAUCIRADIATUS, GILL.

"D. III, 6; A. 4. The preopercular spine is armed with three teeth above and terminates in an acute point." (Gill.) Matanzas, Cuba; an imperfectly described species, known only from the above note. (*pauci*, few; *radiatus*, rayed.)

*Callionymus pauciradiatus*, GILL, Ann. Lyc. Nat. Hist. N. Y., VIII, 1865, 143, Matanzas, Cuba.

## Family CLXXXVIII. GOBIIDÆ.

## (THE GOBIES.)

Body oblong or elongate, naked or covered with ctenoid or cycloid scales. Dentition various, the teeth generally small; premaxillaries protractile; suborbital without bony stay. Skin of head continuous with covering of eyes. Opercle unarmed; preopercle unarmed or with a short spine; pseudobranchiæ present. Gills 4, a slit behind the fourth; gill membranes united to the isthmus, the gill openings thus restricted to the sides. No lateral line. Dorsal fins separate or connected, the spinous dorsal least developed, of 2 to 8 flexible spines, rarely wanting; anal usually with a single weak spine, similar to the soft dorsal; ventral fins close together, separate or fully united, each composed of a short spine and 5 (rarely 4) soft rays, the inner rays longest; the ventral fins, when united, form a sucking disk, a cross fold of skin at their base completing the cup; caudal fin convex; anal papilla prominent. No pyloric cæca; usually no air bladder. Carnivorous fishes, mostly of small size, living on the bottoms near the shores in warm regions. Some inhabit fresh waters, and others live indiscriminately in either fresh or salt water. Many of them bury in the mud of estuaries. Few of them are large enough to be of much value as food. Genera about 80; species nearly 600. The species are for the most part easily recognized, but their arrangement in genera is a matter of extreme difficulty. Until the multitude of Asiatic forms are critically studied, any definition of the American genera must be tentative only. (*Gobiidæ*, part; groups *Gobiina*, *Amblyopina*, and *Trypauchenina*, Günther, Cat. Fishes, III, 1-138.)

## ANALYSIS OF GENERA OF NORTH AMERICAN GOBIIDÆ.

## a. Ventral fins separate; body scaly.

## OXYMETOPONTINÆ:

## b. Ventral rays I, 4.

- c. Forehead bluntly rounded, without sharp keel; tongue very slender, sharp; body elongate, compressed, covered with very small scales; head short, compressed, rather broad above; mouth oblique, the lower jaw projecting; teeth in few series, some of them canine-like; isthmus narrow. Dorsals separate, the first of 6 slender spines; soft dorsal and anal elongate; caudal lanceolate. LOGLOSSUS, 800.

## ELEOTRIDINÆ:

## bb. Ventral rays I, 5.

- d. Vomer with a broad patch of villiform teeth; gill openings extending forward to below posterior angle of mouth, the isthmus thus very narrow;

teeth villiform, the outer scarcely enlarged; vertebrae 12 + 13 (*dormitor*); skull above with conspicuous elevated ridges, one of these bounding the orbit above, the orbital ridges connected posteriorly above by a strong cross ridge; a sharp longitudinal ridge on each side of the occipital, the two nearly parallel, the post-temporals being attached to the posterior ends. Insertions of post-temporals widely separated, the distance between them greater than the rather narrow interorbital width; the post-temporal bones little divergent; top of head depressed, both before and behind the cross ridge between eyes; a flattish triangular area between this and the little elevated supraoccipital region; preopercle without spines; lower pharyngeals with slender, depressible teeth, and without lamelliform appendages; scales of moderate size, ctenoid.

PHILYPNUS, 801.

dd. Vomer without teeth; isthmus broad; gill openings scarcely extending forward below to posterior angle of preopercle; skull without crests.

e. Body scaly, both anteriorly and posteriorly.

f. Lower pharyngeal teeth stiff and blunt; the bones with an outer series of broad flexible lamelliform appendages, which are rudimentary gill filaments; body short and elevated; teeth slender, those in the outer row scarcely larger, and movable; top of head without raised crests, flattish, its surface uneven; post-temporal bones rather strongly diverging, the distance between their insertions about  $\frac{1}{2}$  the broad flattish interorbital space; no spine on preopercle or branchiostegals; scales large, ctenoid. Species herbivorous.

DORMITATOR, 802.

ff. Lower pharyngeals normal, subtriangular, the teeth stiff, villiform, no lamelliform appendages; scales of moderate or small size; body oblong or elongate.

g. Body moderately robust, the depth 4 to  $5\frac{1}{2}$  times in the length to base of caudal; scales ctenoid; cranium without distinct median keel; a small supraoccipital crest.

h. Post-temporal bones little divergent, not inserted close together, the distance between their insertions greater than the moderate interorbital space, or  $3\frac{1}{2}$  in length of head; top of skull little gibbous; lower pharyngeals narrower than in *Eleotris*; preopercle without spine; scales very small, about 110 in a longitudinal series. Vertebrae 11 + 13; teeth moderate, the outer series on lower jaw enlarged.

GUAVINA, 803.

hh. Post-temporal bones very strongly divergent, their insertions close together, the distance between them about  $\frac{1}{2}$  the narrow interorbital space, and less than  $\frac{1}{2}$  length of head; top of skull somewhat elevated and declivous; interorbital area somewhat convex transversely; lower pharyngeals rather broad, the teeth bluntish; preopercle with partly concealed spine directed downwards and forward at its angle; scales moderate, 45 to 60 in a longitudinal series; vertebrae (*pisoniae*) 11 + 15; teeth small.

ELEOTRIS, 804.

gg. Body very slender, elongate, the depth 8 to 9 times in length to base of caudal; scales very small, cycloid.

i. Preopercle with a partly concealed antorse hook at its angle; caudal with numerous accessory rays at base.

ALEXURUS, 805.

- ii. Proopercle without spine; caudal without many accessory rays at base; post-temporal bones short strongly divergent, the distance between their insertions about equal to the narrow interorbital space, or about  $\frac{1}{2}$  length of head; top of head with a strong median keel, which is highest on the occipital region; no supraoccipital crest; mouth very oblique; the teeth small. EROTELIS, 806.
- ee. Body naked on the anterior part; head naked; lower jaw with 4 larger recurved teeth. GYMNELEOTRIS, 807.
- eee. Body entirely naked. CHRIOLEPIS, 808.
- aa. Ventral fins united.
- j. Dorsal fins separate, free from caudal.
- SICYDIINÆ:
- k. Ventral disk short, adnate to belly; body subcylindrical, covered with ctenoid scales; lips very thick; upper teeth mostly small and movable, lower fixed; dorsal spines 6.
- l. Teeth simple; no canines in front of lower jaw. SICYDIUM, 809.
- ll. Teeth trifold (or bifid); no canines in front of lower jaw. COTYLOPUS, 810.
- GOBIINÆ:
- kk. Ventral disk free from the belly.
- m. Dorsal spines 4 to 8; eyes well developed.
- n. Teeth emarginate, uniserial, those of the lower jaw nearly horizontal; dorsal spines 6; scales large, ctenoid; gill openings moderate. EVORTHODUS, 811.
- nn. Teeth simple.
- o. Body scaly, more or less.
- p. Maxillary normal, not prolonged behind the rictus; skull of the usual gobioid form, comparatively short and abruptly broadened behind the orbits; occiput depressed; supraoccipital and temporal ridges continuous.
- q. Dorsal spines 6; scales evidently ctenoid; head naked (the nape scaly as usual.)
- r. Interorbital area anteriorly elevated, with a large foramen-like depression in front of eye; body short, compressed, formed much as in *Dormitator*; nape with a fleshy crest; scales large. Vertebrae 11 + 15. LOPHOGOBIUS, 812.
- rr. Interorbital area not elevated in front; body more elongate; no fleshy nuchal crest; isthmus broad.
- s. Inner edge of shoulder girdle without fleshy cirri or papillae; cranium anteriorly short; interorbital space narrower, grooved, with a low median ridge or none; median crest on cranium low.
- t. Body scaly anteriorly and posteriorly (sometimes a naked strip on back or belly). Vertebrae 12 + 16 to 10 + 15. GOBIUS, 813.
- tt. Body entirely naked anteriorly, the posterior half scaled; scales moderate or small. GARMANNIA, 814.

- ss. Inner edge of shoulder girdle with 2 or 3 conspicuous dermal flaps; preorbital region very long; premaxillary and maxillary strong; interorbital groove with a conspicuous median crest; scales rather small (45 to 70.) AWAOUS, 815.
- qq. Dorsal apines 7 or 8 (very rarely 6, especially in *Eucyclogobius*.)
- u. Scales large, ctenoid; shoulder girdle without dermal flaps.
- v. Sides of head scaled; soft dorsal and anal rather short, of 11 to 14 rays each; deep-water species.  
BOLLMANNIA, 816.
- vv. Sides of head naked; soft dorsal and anal short, of 10 to 12 rays each; shore species. ABOMA, 817.
- uu. Scales very small, cycloid or nearly so.
- w. Inner edge of shoulder girdle without fleshy processes; head naked; body more or less compressed; mouth very oblique; teeth strong; interorbital groove with or without a median ridge. Vertebrae 11 + 15 or 16; soft dorsal and anal long, of 15 to 17 rays each.
- x. Body chiefly scaly, anteriorly as well as posteriorly.  
MICROGOBIUS, 818.
- xx. Body naked anteriorly, scaled posteriorly. ZALYPNUS, 819.
- ww. Inner edge of shoulder girdle with 2 or 3 dermal flaps, or processes, as in *Awaous*.
- y. Head naked, the interorbital groove with the median ridge high, not extending forward to orbit; body rather robust; soft dorsal and anal short; fresh-water species.  
EUCYCLOGOBIUS, 820.
- yy. Head scaled like the body; the interorbital groove with the median ridge little developed; soft dorsal and anal long; body elongate; marine species.  
LEPIDOGOBIUS, 821.
- pp. Maxillary much produced backward, extending beyond the gill opening in the adult; skull comparatively long, gradually (not abruptly) broadened behind orbits; median crest of cranium well developed; scales small, cycloid; head naked, occipital region narrowed forward; supraorbital and temporal crests not continuous.

es-  
rt  
eir  
tal  
ith  
oci-  
ery  
306.  
ger  
307.  
308.

with  
ble,  
800.  
810.

hori-  
nings  
, 811.

ctus;  
tively  
rbits;  
aporal

head

with a  
ont of  
rmed  
with a  
tebrae  
s, 812.  
; body  
crest;

without  
anium  
space  
a low  
n crest

poste-  
d strip  
rtebrae  
us, 813.  
eriously,  
; scales

IA, 814.

- z. Occiput depressed, with a blunt median keel.  
*a'*. Shoulder girdle without dermal flaps; dorsal spines 6; soft dorsal and anal short; mouth very large; isthmus broad; vertebrae 14 + 16 (*mirabilis*).

GILLICHTHYS, 822.

- aa'*. Shoulder girdle with 1 to 3 small dermal flaps on the inner edge; dorsal spines 5; soft dorsal and anal long.

QUIETULA, 823.

- zz. Occiput transversely rounded without median keel.

- b'*. Shoulder girdle with 1 to 3 small dermal flaps on its inner edge; dorsal spines 5; soft dorsal and anal long. LYPNUS, 824.

- bb'*. Shoulder girdle without dermal flaps; dorsal spines 4 or 5; soft dorsal and anal long. CLEVELANDIA, 825.

- oo. Body and head entirely naked.

- c'*. Dorsal spines 4; body long and slender; mouth large, the lower jaw projecting; no barbels; soft dorsal and anal long; male with ornate colors.

EVERMANNIA, 826.

- cc'*. Dorsal spines 7 (rarely 6).

- d'*. Chin without barbels; mouth small, little oblique; body robust, soft dorsal and anal short. GOBIOSOMA, 827.

- dd'*. Chin with a fringe of short barbels; mouth terminal, oblique; soft dorsal and anal very short. BARBULIFER, 828.

## CRYSTALLOGOBINÆ:

- mm*. Dorsal spines 2 (or 1); body wholly naked.

- e'*. Eyes reduced to small rudiments; interorbital area forming a sharp median ridge; skull rather abruptly widened behind orbits; anterior portion of skull unusually long; no flaps on shoulder girdle; skull highest at nape, depressed above the eyes; soft dorsal and anal short.

TYPHLOGOBIUS, 829.

## GOBIODINÆ:

- jj*. Dorsal fin continuous, the soft part and the anal joined to base of caudal; eyes minute; body elongate; scales minute or wanting; mouth very oblique, the lower jaw projecting; gill openings moderate.

- f'*. Dorsal rays VI, 16 to 23; anal rays 17 to 23.

- g'*. Teeth small, in a single series; scales present. TYNTLASTES, 830.

- gg'*. Teeth in a band, those of the outer series being very strong; scales present.

- h'*. Body entirely scaled. GOBIOIDES, 831.

- hh'*. Anterior part of body naked. CAVENNA, 832.

## 800. IOGLOSSUS, Bean.

*Ioglossus*, BEAN, in Jordan & Gilbert, Proc. U. S. Nat. Mus. 1882, 297 (*ocillurus*).

Body elongate, strongly compressed, of equal depth throughout, covered with very small, mostly cycloid, scales. Head short, compressed, not keeled above; mouth large, oblique, the lower jaw projecting; teeth in narrow bands or single series, some of them canine; no teeth on vomer or palatines; tongue very slender, sharp; opercles unarmed. Gill openings

*Ioglossus*  
sac  
949,  
Nat

very wide, the membranes narrowly joined to isthmus on median line. No lateral line. Branchiostegals 5. Dorsals separate, the first of 6 very slender, flexible spines; the second elongate, similar to the anal; caudal long and pointed, free from dorsal and anal; ventrals close together, separate, each of 1 spine and 4 rays, their insertion below or behind pectorals; anal papilla present. A remarkable type, belonging to the *Oryzotopontinae*, differing widely from our other Gobioid fishes. Gulf of Mexico, in rather deep water. (ὄσ, arrow; γλῶσσα, tongue.)

2515. IOGLOSSUS CALLIURUS, Bean.

Head 5; depth 7 to 7½. D. VI-22 to 24; A. I, 21 to 23. Body very elongate, slender, much compressed, of equal depth throughout; head compressed, without osseous crest; mouth very oblique, the lower jaw strongly projecting; premaxillaries in front on the level with pupil; maxillary extending to opposite front of pupil, its length 2½ in head; upper jaw with a narrow band of about 2 series of conical cardiform teeth, those of the outer row much larger than the others, behind these 2 small conical curved canines; lower jaw with a single row of smaller teeth, behind which are about 4 short canines directed somewhat backward; the posterior pair strongly curved; no teeth on vomer or palatines. Tongue narrow, pointed. Eye large, nearly twice length of snout, 3½ in head, its diameter considerably more than depth of cheek, about ½ more than interorbital width; opercles unarmed. Pseudobranchiæ present. Gill openings wide, extending forward below, the membranes attached mesially to the very narrow isthmus, across which they do not form a fold. Gill rakers long and slender. Dorsal fins separated by a short interval, the first of very slender somewhat filamentous spines, the longest about as long as head; second dorsal little more than ½ as high as first, apparently nearly uniform, separated from the caudal by an interval nearly ½ length of head; caudal lanceolate, its middle rays filamentous, about ½ the length of rest of body; anal rather high, similar to soft dorsal; ventrals I, 4, inserted very slightly in advance of base of pectorals, the 2 fins very close together, but apparently quite separate and without basal fold of skin, the fin little longer than head, the inner rays filamentous; pectoral with broad base, about 1½ in head. Anal papilla very short, midway between tip of snout and base of caudal. Body with very small, nonimbricate, embedded scales, these a little larger and imbricate on the tail; cheeks with embedded cycloid scales; scales very weakly etenoid, most of them appearing cycloid; no lateral line. Color light olive, everywhere densely punctate; dorsals edged with black; middle of caudal reddish, with paler bluish edgings. Length 4½ inches. Here described from specimens from off Pensacola. Gulf of Mexico; known only from the Snapper Banks off Pensacola, in rather deep water. (κάλλος, beauty; οὐρά, tail.)

*Ioglossus calliurus* (BEAN MS.), in JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1882, 297, Pensacola, Florida; BEAN, Proc. U. S. Nat. Mus. 1882, 419; JORDAN & GILBERT, Synopsis, 949, 1883; JORDAN, Proc. U. S. Nat. Mus. 1884, 437; JORDAN & EIGENMANN, Proc. U. S. Nat. Mus. 1886, 481.



## 801. PHILYPNUS,\* Cuvier &amp; Valenciennes.

(GUAVINAS.)

*Gobiomorus*,† LACÉPÈDE, Hist. Nat. Poiss., II, 699, 1798, in part (*dormitor*, etc.); restricted to *dormitor* by JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1882, 571; restricted to *Gobiomorus taibe* †, Lacépède (Valenciennes' *strigata*), by GILL, Proc. U. S. Nat. Mus. 1888, 70, in accordance with the law of exclusion.

*Philypnus*, CUVIER & VALENCIENNES, Hist. Nat. Poiss., XII, 255, 1837 (*dormitor*).

*Lembus*, GÜNTHER, Cat. Fishes, I, 505, 1859 (*maculatus*).

Body elongate, terete anteriorly, compressed behind. Head elongate, depressed above. Mouth large; lower jaw considerably projecting; teeth in jaw rather small, slender, recurved, the outer scarcely enlarged; teeth on vomer villiform, in a broad, crescent-shaped patch; gill openings extending forward to below posterior angle of mouth, the isthmus very narrow. Scales moderate, ctenoid, covering most of the head, 55 to 66 in a longitudinal series. Dorsal with 6 spines and 9 or 10 rays; anal rays I, 9 or 10; ventrals separate. No preopercular spine; insertion of post-temporals almost midway between occipital crest and edge of skull; parietals with a crest running from insertion of post-temporal forward to just behind eye, where they are connected by a thin, high, transverse crest; supraocular with a short, high crest, extending from above front of eye back to posterior edge of orbit, thence extending outward parallel with the transverse crest, leaving a deep groove between them; bony projections before and behind eye prominent. Vertebrae 12 + 13 = 25; lower pharyngeals triangular, with slender teeth. Largest of the Gobies, some of the species reaching a length of 2 or 3 feet and valued as food. Tropical rivers. (*φιλυπνος*, slumber-loving; *φιλος*, loving; *ὑπνος*, sleep.)

a. Coloration rather obscure, the dark lateral band indistinct or wanting; scales 55 to 57. DORMITOR, 2516.

aa. Coloration bright, the black lateral band distinct; scales 52 to 55. LATERALS, 2517.

## 2516. PHILYPNUS DORMITOR (Lacépède).

(SLEEPER; GUAVINA.)

Head  $2\frac{2}{3}$  to  $2\frac{3}{4}$ ; depth 5 to  $5\frac{1}{2}$ . D. VI-10; A. I, 9; scales 55 to 57; eye  $6\frac{1}{4}$  to  $7\frac{1}{4}$  in head; snout  $3\frac{2}{3}$ ; maxillary  $2\frac{2}{3}$ . Body elongate, terete anteriorly, compressed behind. Head elongate, depressed above. Mouth large; maxillary reaching to middle of pupil. Lower jaw considerably projecting. Teeth on jaws slender, depressible. Interspace between dorsals slightly greater than interorbital width; dorsal spines slender, the second the longest,  $2\frac{1}{2}$  in head; length of base of anal about  $2\frac{1}{2}$  in head; ventrals reaching  $\frac{2}{3}$  of the distance to vent; tips of pectorals reaching ventral.

\* The *Electrinae* have been made the subject of a special paper (A Review of the American Eleotridine, in Proc. Ac. Nat. Sci. Phila. 1885, 66-80) by Eigenmann & Fordice. The *Gobiidae* of America have been discussed in detail by Jordan & Eigenmann (Proc. U. S. Nat. Mus. 1888, 477-518) and later by Eigenmann & Eigenmann (Proc. Cal. Ac. Sci., 2d ser., vol. 1, 1888, 51-78). In this paper are valuable notes on the specimens in the Museum of Comparative Zoology.

† For the reasons in favor of the use of the name *Gobiomorus* for *Valenciennes*, Bleeker, instead of using it for the present genus, see GILL, Proc. U. S. Nat. Mus. 1888, 69.

Dark brownish or olive, lighter below; an interrupted dark lateral band extending from base of pectoral to base of caudal (not always present); fins dusky, and with the exception of the anal and ventrals, all distinctly mottled; spinous dorsal margined with blackish; head often with dark spots. Streams of the West Indies and Atlantic shores of Central America, Mexico, and Surinam; everywhere common, reaching a length of 2 feet or more. Here described from Cuban specimens. (*dormitor*, sleeper.)

- Guavina*, PARRA, Descr. Dif. Piezas Hist. Nat. Cuba, tab. 39, fig. 1, 1787, Havana.  
*Gobiomorus dormitor*, LACÉPÈDE, Hist. Nat. Poiss., II, 599, 1798, Martinique; from a drawing by PLUMIER; EIGENMANN & EIGENMANN, Proc. Cal. Ac. Sci. 1888, 52.  
*Platycephalus dormitor*, BLOCH, Ichth., 1801, Martinique; after LACÉPÈDE.  
*Batrachus guavina*, BLOCH & SCHNEIDER, Syst. Ichth., 44, 1801; based on *Guavina* of PARRA.  
*Eleotris longiceps*, GÜNTHER, Proc. Zool. Soc. Lond. 1864, 151, Nicaragua; GÜNTHER, Fish. Contr. Amer., 440, 1869.  
*Electris dormitatrix*, CUVIER, Règne Animal, Ed. II, vol. 2, 246, 1829, Antilles; GÜNTHER, Cat. Fish., III, 119, 1861.  
*Gobiomorus dormitor*, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1882, 572.  
*Philypnus dormitor*, CUVIER & VALENCIENNES, Hist. Nat. Poiss., XII, 255, 1837; POEY, Mem. de Cuba, II, 381, 1860; GIRARD, U. S. and Mexican Boundary Survey, Zool., 27, pl. 12, fig. 13, 1859; JORDAN & GILBERT, Synopsis, 631, 1883.

2517. PHILYPNUS LATERALIS, GILL.

(ABOMA DE MAR.)

Head  $2\frac{1}{2}$ ; depth  $5\frac{1}{2}$ . D. VI-10; A. I, 10; scales 52 to 55; eye 6 to  $6\frac{1}{2}$  in head; snout  $3\frac{1}{2}$  to  $3\frac{3}{4}$ ; maxillary  $2\frac{1}{2}$  to  $2\frac{3}{4}$ . Brownish, lighter or white below; a distinct dark brown or blackish band extending from base of pectoral to base of caudal; dorsals, pectoral and caudal dusky; ventrals and anal lighter; dorsals, caudal, and in some specimens the anal, distinctly blotched. The only constant difference between this species and *Philypnus dormitor* seems to be the brighter coloration of *lateralis*. Streams of Pacific Coast of Mexico and Central America, from Sonora to Panama, entering the sea; common, reaching a much larger size than any other of our Gobies. Here described from specimens from Rio Presidio, Mazatlan. (*lateralis*, pertaining to the side.)

- Philypnus lateralis*, GILL, Proc. Ac. Nat. Sci. Phila. 1860, 123, Cape San Lucas (Coll. Xantus); JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1882, 377.  
*Eleotris lateralis*, GÜNTHER, Cat., III, 122, 1861.

802. DORMITATOR, Gill.

(PUÑECAS.)

- Prochilus*, CUVIER, Règne Animal, Ed. 1, vol. II, 294, 1817 (*macrolepidota* = *maculatus*); name preoccupied.  
*Dormitator*, GILL, Proc. Ac. Nat. Sci. Phila. 1862, 240 (*gundlachi*).

Body short, robust; head broad and flat above; mouth little oblique; maxillary reaching to anterior margin of orbit; lower jaw little projecting; no teeth on vomer; lower pharyngeal teeth stiff and blunt, the bones with an external series broad, flexible, lamelliform, these being rudimentary gill filaments; scales large, ctenoid, 30 to 33 in a longitudinal

series; skull much as in *Eleotris*; D. VII-I, 8; A. I, 9 or 10; no spine on preopercle; post-temporals inserted midway between occipital crest and edge of skull; supraoccipital crest low. (*dormitator*, one who sleeps.)

2518. *DORMITATOR MACULATUS* (Bloch).

(GUAVINA MAPO; PANACA.)

Head  $3\frac{1}{2}$ ; depth about 3 in adult. D. VII-I, 8 or 9; A. I, 9 or 10; lateral line 33. Body short, robust; head broad and flat above; eye small, less than snout; caudal a little shorter than head; mouth little oblique; maxillary reaching to anterior margin of orbit; lower jaw little projecting; no teeth on vomer; interspace between dorsals equaling orbit; highest anal ray  $1\frac{1}{2}$  in head; highest dorsal ray  $1\frac{1}{2}$  in head; skull much as in *Eleotris*, but everywhere broader; no spine on preopercle; post-temporal inserted midway between occipital crest and edge of skull; supraoccipital crest low; scales large, becoming much smaller on belly, 25 series on median line from base of ventrals to vent; 18 series across breast from pectoral to pectoral; 18 on a median line from posterior border of orbit to dorsal. Dark brown, with lighter bluish spots; a faint dark stripe along sides; a conspicuous large dark blue spot edged with black above base of pectorals; a dark streak from eye to angle of mouth; 2 dark streaks on side of head; branchiostegal membrane blackish; dorsals barred with spots; anal dusky, barred with bluish, and with white margin; a dark bar on base of pectoral. Length 1 to 2 feet. Both coasts of America, ranging from South Carolina through the West Indies to Pará, Cape San Lucas, and Panama, in fresh or brackish water; everywhere abundant and used as food. Dr. Eigenmann observes:

There seem to be 2 forms of the adult—one with the profile gibbous, the dorsal outline forming a regular curve; the other having the profile depressed over the eyes, the anterior portion being subhorizontal. The specimens from Gurupa and the Rio Grande have the profile depressed; all the other specimens have a gibbous profile. A comparison in detail of the two forms is appended. Only extreme differences are given.

West Indian specimens 5 to  $7\frac{1}{2}$  inches.  
 Profile regularly curved from first dorsal spine to snout.  
 Head  $3\frac{1}{2}$  to 4; depth 3 to  $3\frac{1}{2}$ ; depth always greater than length of head.  
 Highest anal ray  $1\frac{1}{2}$  to  $1\frac{3}{4}$  in head.  
 Distance from first dorsal spine to snout greater than distance from first dorsal spine to first anal ray.  
 Scales in median series 29 to 32.  
 Color usually dark brown, a black spot above base of pectoral, a short bar on base of pectoral.

Rio Grande specimens 5,  $6\frac{1}{2}$ , and  $7\frac{1}{2}$  inches.  
 Profile depressed over eye, becoming horizontal anteriorly.  
 Head 3; depth 3 to  $3\frac{1}{2}$ ; depth usually less than length of head.  
 Highest anal ray  $1\frac{1}{2}$  to 2 in head.  
 Distance from first dorsal spine to snout equals distance from first dorsal spine to base of last anal ray.  
 Scales in median series 30 to 34.  
 Color gray, a jet-black spot above base of pectoral; a black bar at base of pectoral; a black line from eye to mouth; longitudinal black lines on cheeks and opercles; dark spots on back; some silvery scales on sides.

Among our specimens from Mazatlan are 3 markedly different forms which seem like distinct species. In view of the great variations to which this species is subject we do not, however, regard them as such, especially as none of the three corresponds exactly to the account above given of the 2 Atlantic forms.

I. DEEP-BODIED SPECIMENS (*Dermitator latifrons*, Richardson).

Head 3; depth 3. D. VII-I, 8; A. I, 8 or 9; scales 30 to 33; eye  $4\frac{1}{2}$  to  $5\frac{1}{2}$  in head; snout  $3\frac{1}{2}$  to  $3\frac{3}{4}$  in head; interorbital width  $2\frac{1}{2}$  in head; ventrals reaching  $\frac{2}{3}$  the distance to vent,  $1\frac{1}{2}$  to  $1\frac{3}{4}$  in head; highest anal ray  $1\frac{1}{2}$  to 2 in head. Body short, robust, the back elevated; head broad and flat above, the anterior profile from first dorsal spine to tip of snout oblique, descending abruptly; mouth oblique, maxillary reaching anterior margin of orbit; lower jaw little projecting. Color greenish, lighter below; body with cross bars of dark brown; fins dusky, the dorsals distinctly blotched with darker; a dark cross bar at base of pectorals; a dark-blue humeral blotch, becoming blackish in spirits; 3 or 4 dark cross bands extending from eye and below eye to posterior margin of preopercle; a dark band extending from below eye to below tip of maxillary. Two specimens from Rio Presidio, Mazatlan.

II. COMMON FORM, AT MAZATLAN.

Head  $3\frac{1}{2}$ ; depth  $3\frac{3}{4}$  to  $3\frac{1}{2}$ . D. VII-I, 7; A. I, 8; scales 33 or 34; eye  $4\frac{1}{2}$  to  $4\frac{3}{4}$  in head; snout  $3\frac{1}{2}$  to 4 in head; interorbital width 3 to  $3\frac{1}{2}$  in head; ventrals reaching about  $\frac{2}{3}$  the distance to vent,  $1\frac{1}{2}$  in head; highest anal ray  $1\frac{1}{2}$  to 2 in head. Body short, compressed, the back little elevated; head rather broad and slightly convex above, the anterior profile from first dorsal spine to tip of snout slightly convex; mouth oblique, maxillary reaching anterior margin of orbit; lower jaw little projecting. Color olive brown, with cross bars of darker brown, lighter below; fins dusky, the dorsals with about 3 darker cross bars; pectorals with a darker cross bar at base; a distinct dark-brown humeral spot slightly larger than eye; 3 or 4 dark cross bands extending from eye and below eye to posterior margin of preopercle; a distinct dark-brown bar extending from below eye to below tip of maxillary; a dark lateral band extending from base of pectoral to base of caudal. Many specimens from Mazatlan.

III. LARGE-HEADED FORM.

Head  $3\frac{1}{2}$ ; depth  $3\frac{1}{2}$ . D. VI-I, 8; A. I, 9; scales 32 or 33; eye  $4\frac{1}{2}$  in head; snout  $3\frac{1}{2}$  in head; interorbital width  $2\frac{1}{2}$  in head; ventrals reaching  $\frac{2}{3}$  the distance to vent,  $1\frac{1}{2}$  in head; highest anal ray 2 in head. Body moderately compressed, the back little elevated; head very broad above, convex; the anterior profile from first dorsal spine to tip of snout oblique, gently descending; mouth oblique, maxillary reaching anterior margin of orbit; lower jaw little projecting. Color brownish, middle of back darker, lighter below; body with darker cross bands; ventrals yellowish; other fins dusky; dorsals with darker blotches; a dark crossbar at base of pectoral; a dark humeral spot; four cross bands extending from eye and below eye to posterior margin of preopercle; a dark band extending from below eye to below top of maxillary; a dark lateral band extending from base of pectoral to base of caudal. One specimen from near Mazatlan. (*maculatus*, spotted.)

- Sciæna maculata*, BLOCH, Ichth., pl. 299, fig. 2, 1790, West Indies.  
*Eleotris mugloides*, CUVIER & VALENCIENNES, Hist. Nat. Poiss., XII, 226, 1837, Martinique; Surinam.  
*Eleotris sima*,\* CUVIER & VALENCIENNES, Hist. Nat. Poiss., XII, 232, 1837, Vera Cruz.  
*Eleotris latifrons*, RICHARDSON, Voy. Sulphur, Fishes, 57, pl. 35, figs. 4 and 5, 1837, locality unknown, supposed to be from Pacific coast, Central America.  
? *Eleotris grandisquama*,† CUVIER & VALENCIENNES, Hist. Nat. Poiss., XII, 229, 1837, America; locality unknown.  
*Eleotris somnolentus*, GIRARD, Proc. Ac. Nat. Sci. Phila. 1858, 169, near mouth of Rio Grande.  
*Eleotris omocyanus*, POEY, Memorias, II, 269, 1860, Havana.  
*Dormitator micropthalmus*, GILL, Proc. Ac. Nat. Sci. Phila. 1863, 170, Panama. (Coll. Capt. John M. Dow.)  
*Dormitator gundlachi*, POEY, Synopsis, 396, 1868, Cuba.  
*Dormitator lineatus*, GILL, Proc. Ac. Nat. Sci. Phila. 1863, 271, Savannah.  
*Dormitator maculatus*, JORDAN & GILBERT, Synopsis, 632, 1883; JORDAN & EIGENMANN, l. c., 482; EIGENMANN & EIGENMANN, Proc. Cal. Ac. Sci., 2d series, vol. 1, 1888, 52.

## 803. GUAVINA, Bleeker.

*Guavina*, BLEEKER, Esquisse d'un Syst. Nat. Gobioid., 302, 1874 (*guavina*).

This genus is allied to *Eleotris*, differing in having the post-temporal bones little divergent, not inserted close together, the distance between their insertions greater than the moderate interorbital space, or  $3\frac{1}{2}$  in length of head; top of skull little gibbous; lower pharyngeals narrower than in *Eleotris*; preopercle without spine; scales very small, ctenoid, about 110 in a longitudinal series. Vertebrae 11 + 13; teeth moderate, the outer series on lower jaw enlarged. Fresh waters of the West Indies and Brazil. Two species known; *Guavina brasiliensis* (Sauvage) from Bahia, and the following. (*Guavina*, the Spanish name.)

## 2519. GUAVINA GUAVINA (Cuvier &amp; Valenciennes).

(GUARUHACO; GUAVINA.)

Head  $3\frac{1}{2}$ ; depth  $4\frac{1}{2}$  to  $5\frac{1}{2}$ . D. VI, or VII-I, 10; A. I, 9 or 10. Body stoutish, oblong; mouth oblique; maxillary reaching opposite middle of eye, its length  $2\frac{1}{2}$  to  $3\frac{1}{2}$  in head. Lower jaw little projecting; teeth in broad bands, the outer ones on lower jaw enlarged. Scales on head embedded; those on body very small, ctenoid on sides, cycloid on back and belly, 100 to 110 in a longitudinal series. Isthmus very broad. Pectorals reaching to middle of spinous dorsal. Highest anal ray  $1\frac{1}{2}$  in head. Post-temporals inserted twice as far from occipital crest as in *Eleotris pisonis*. Parietals ending

\* Types, 2 specimens in poor order, from Vera Cruz, 0.09 mm. long. Snout a little more steep and convex than usual in *Dormitator maculatus*. Head  $3\frac{1}{2}$  in length; depth  $3\frac{1}{2}$ . Eye  $4\frac{1}{2}$  in head. D. VII, 9; A. 11; scales 31-11. Soft dorsal very high, with round black spots. Caudal and anal plain. This seems to be inseparable from *Dormitator maculatus*.

† We have the following note on the type of *Eleotris grandisquama*: Type specimen in fair condition, 0.14 mm. long, from "Amérique Méridionale?" Head slenderer than in *D. maculatus*, and much depressed, its depth at the eyes less than its width, which is less than that of body. Anterior profile almost concave. Caudal fin large; other fins moderate. Dorsal VI, 9; anal I, 9; scales about 29-11. A few dusky spots on dorsal and anal. According to Dr. Eigenmann, specimens of *Dormitator maculatus* from the Rio Grande agree fairly with this type, and it is not likely that it is different.

in a sharp point behind. Preopercular spine none; a broad, thin extension on the lower limb of preopercle taking its place. Lower pharyngeals triangular, normal, rather narrow; the teeth small. Vomer without teeth. Length 1 foot. East coast of tropical America, Cuba to Rio Janeiro, in fresh and brackish waters; very common. (*guavina*, Spanish name.)

*Eleotris guavina*, CUVIER & VALENCIENNES, Hist. Nat. Poiss., XII, 223, 1837, Martinique.  
*Guavina guavina*, JORDAN & EIGENMANN, l. c., 483.

804. ELEOTRIS (Gronow) Bloch & Schneider.

*Eleotris*, GRONOW, Zooph., 83, 1763 (nonbinomial).  
*Eleotris*, BLOCH & SCHNEIDER, Syst. Ichth., 65, 1801 (*pisonis*).  
*Culius*, BLEEKER, Esquisse d'un Syst. Nat. des Gobioid., 303, 1874 (*fuscus*).

Body long and low, compressed behind. Head long, low, flattened above, without spines or crests, almost everywhere scaly. Mouth large, oblique, lower jaw projecting. Lower pharyngeals rather broad, the teeth small,\* bluntish. Preopercle with a small concealed spine below, its tip hooked forward. Branchiostegals unarmed. Eyes small, high, anterior; isthmus broad. Post-temporal bones very strongly divergent, their insertions close together, the distance between them about  $\frac{3}{4}$  the narrow interorbital space, and less than  $\frac{1}{2}$  length of head; top of skull somewhat elevated and declivous; interorbital area slightly convex transversely; dorsal fins well apart, the first of 6 or 7 flexible spines; ventrals separate. Scales moderate, ctenoid, 45 to 62 in a longitudinal series; vertebrae (*pisonis*) 11 + 15. Tropical seas, entering fresh waters. (*ἡλεός*, bewildered.)

- a. Teeth subequal, those of inner or outer series enlarged.
- b. Cheek entirely scaled.
- c. Teeth of inner series of each jaw enlarged.
  - d. Scales in a median series 40 to 51, in a cross series 12 to 20.
  - e. Eye large, 5 to 6 in head; scales 40 to 44—12 to 14.
- ee. Eye small, 8 in head; scales 51-20. AMBLYOPSIS, 2520.  
 ABACURUS, 2521.
- dd. Scales in a median series 57 to 66; in a cross series 18 to 24.
- bb. Lower half of cheek naked; scales 61. PISONIS, 2522.  
 PERNIGER, 2523.
- aa. Teeth all equal; scales 60. PICTUS, 2524.

2520. ELEOTRIS AMBLYOPSIS (Cope).

Head  $3\frac{1}{2}$ ; depth  $4\frac{1}{4}$ . D. VI-9; A. I, 8; scales 46 (40 to 44—12 to 14 according to Eigenmann); eye  $5\frac{1}{2}$  in head, 2 in interorbital width; preopercular spine strong, decurved; width of head  $\frac{3}{4}$  in its length; chin prominent; premaxillary spines forming a projection in profile. Brown, a black spot above at base of pectoral; first dorsal and anal dusky; second dorsal and caudal delicately cross-banded with blackish; 3 black lines from orbit behind and below. Surinam. Described from 3 specimens each 3 inches long. (Cope.) Dr. Eigenmann mentions 15 other examples,

\* The characters of the skeleton are taken from *Eleotris pisonis* and have not been verified on other species. The hooked preopercular spine supposed to characterize *Culius* is found on the typical species of *Eleotris*, as well as in *Ateurus*.

the longest  $2\frac{1}{2}$  inches long, from Surinam, in the Museum of Comparative Zoology. (*ἀμβλύς*, blunt; *ὄψις*, face.)

*Eleotris amblyopsis*, COPE, Trans. Amer. Philos. Soc. 1870, 473, Surinam (Coll. Dr. Charles Hering); JORDAN & EIGENMANN, l. c., 483, 1886; EIGENMANN & EIGENMANN, l. c., 55.

2521. *ELEOTRIS ABACURUS*, Jordan & Gilbert.

Head 3; depth  $4\frac{1}{2}$ . D. VI-9; A. I, 8; scales 51-20; eye 8 in head,  $2\frac{1}{2}$  in interorbital width; pectoral  $1\frac{1}{2}$ ; ventral  $1\frac{1}{2}$ ; highest dorsal ray 2; highest anal ray 2; caudal  $1\frac{1}{2}$ . Body slender, compressed, the head depressed, becoming very narrow anteriorly, its width  $\frac{3}{4}$  its length; a notable depression above orbits, the premaxillary processes protruding before it; lower jaw the longer; maxillary reaching vertical behind pupil,  $2\frac{3}{4}$  in head. Teeth in jaws in narrow villiform bands, becoming a single series on sides of lower jaw, those of the outer and inner series in each jaw somewhat enlarged, the largest being a single series in sides of lower jaw. Preopercular spine as usual in the genus. Scales smooth above and below, ctenoid on sides. Color in spirits, brown, lighter above and below; each scale on middle of sides with a dusky streak, these forming obscure lengthwise lines; back anteriorly with a few small black spots; under parts, including sides of head, very thickly punctulate with black; no dark stripes from orbit; lips black; a dark streak from snout through eye to upper angle of preopercle; 2 dusky streaks from eye downward and backward across cheek; a very conspicuous black blotch as large as eye in front of upper pectoral rays; pectorals and ventrals transparent, dusky; vertical fins all barred with light and dark in fine pattern. Coast of South Carolina. Known from a single specimen, 4 inches long, taken in the harbor of Charleston. This species agrees very well with Cope's account of *Culius amblyopsis*, but the scales are larger, the eye is smaller, and there is some difference in color, besides the remote habitat. (*ἄβακος*, checker; *ὄψα*, tail.)

*Culius amblyopsis*, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1882, 610; not of Cope.

*Eleotris abacurus*, JORDAN & GILBERT, Proc. Cal. Ac. Sci. 1896, 228, Charleston. (Coll. Dr. C. H. Gilbert. Type, No. 2009, I. S. Jr. Univ. Mus.)

2522. *ELEOTRIS PISONIS* (Gmelin).

(GUAVINA TÉTARD; SLEEPER.)

Head 3 to  $3\frac{1}{2}$  in body; depth  $4\frac{1}{2}$  to 5. D. VI-9; A. I, 8; scales 62; eye  $5\frac{1}{2}$  to 8 in head; maxillary  $2\frac{3}{4}$ ; pectoral  $1\frac{1}{2}$ ; ventral 2; caudal  $1\frac{1}{2}$ . Body not much compressed; head somewhat depressed; mouth rather large, the maxillary reaching to below posterior margin of pupil; lower jaw much projecting, a knob at symphysis; wide bands of villiform teeth in jaws, none on vomer or palatines; interorbital region nearly twice as wide as the horizontal diameter of eye; top of head, cheeks, and opercles covered with small scales; a stout, concealed spine projecting downward on edge of preopercle. Origin of dorsal about midway between tip of snout and end of last dorsal rays; tips of first dorsal spines not reaching front of

see  
of  
tra  
wi  
wa  
des  
dar  
mo  
ma  
Pis  
Pri  
Bra  
Am  
Eleo  
Gobi  
Gobi  
Eleot  
M  
Eleot  
(  
E  
Eleot  
  
He  
wid  
cultu  
file;  
diam  
dusk  
1 alon  
spot  
(Cope  
but th  
Culius  
Rij  
Eleotri  
  
Hea  
scales  
  
\* Eleo  
scales  
betwee  
extendi  
flattene  
ment; o  
brownis  
Eviden

second dorsal when fin is depressed; origin of anal a little behind that of soft dorsal; pectorals reaching to posterior spine of first dorsal; ventrals inserted very slightly behind base of pectorals; caudal peduncle as wide as length of maxillary. Color brownish; fins with dark spots and wavy lines; ventrals dusky; 2 dark stripes behind the orbit. Here described from specimens, 6 or 7 inches long, collected in the Rio Almendares, Cuba, by Dr. Jordan. Streams of the West Indies, generally common from southern Florida to Rio Janeiro. Dr. Eigenmann enumerates many specimens from various localities in Brazil. (Named for Dr. William Piso, of the University of Leyden, associate of George Maregraf and Prince Maurice of Nassau, in 1648, in the study of the natural history of Brazil.)

*Amore pixuma*, MARCGRAVE & PISO, Hist. Brasil, IV, 166, 1648, Brazil.

*Eleotris capite plagioplateo*, GRONOW, Mus. Ichth., II, 168, 1757; after MARCGRAVE.

*Gobius pisonis*, GMELIN, Syst. Nat., 1206, 1788; based on *Eleotris* of GRONOW.

*Gobius amorsæ*, WALBAUM, Artedii Piscium, III, 205, 1792; based on *Eleotris* of GRONOW.

*Eleotris gyrrinus*, CUVIER & VALENCIENNES, Hist. Nat. Poiss., XII, 220, pl. 356, 1837, Martinique; San Domingo; Surinam.

*Eleotris (Culius) belizianus*,\* SAUVAGE, Bull. Soc. Philom. Paris 1879 (1880), 55, Belize (Coll. Morelet), Cayenne (Coll. Méllon); EIGENMANN & FORDICE, Proc. Ac. Nat. Sci. Phila. 1885, 75; EIGENMANN & EIGENMANN, Proc. Cal. Ac. Sci. 1888, 55.

*Eleotris pisonis*, JORDAN & EIGENMANN, l. c., 483; EIGENMANN & EIGENMANN, l. c., 55.

2523. ELEOTRIS PERNIGER (Cope).

Head  $4\frac{1}{2}$ ; depth  $4\frac{1}{2}$ . D. VI-I, 9; A. I, 9; scales 61; eye 3 in interorbital width; no vomerine teeth. A strong spine at posterior angle of preoperculum, directed downward. Premaxillary spines not prominent in profile; scaling of vertex extending to their extremities. Longitudinal diameter of orbit  $\frac{1}{2}$  length of head. Color black, abdomen brown, fins dusky; first dorsal with white extremity and 2 longitudinal black bars, 1 along the base; other fins with small black bars; [no] maxillary or caudal spot or ocellus. Length 5 inches. West Indies, south to Rio Janeiro. (Cope.) A specimen in our collection from Jamaica. It is close to *E. pisonis*, but the cheeks are not fully scaled. (*perniger*, very black.)

*Culius perniger*, COPE, Trans. Am. Philos. Soc. 1870, 473, St. Martins. (Coll. Dr. R. E. van Rijgersma.)

*Eleotris perniger*, EIGENMANN & EIGENMANN, l. c., 55.

2524. ELEOTRIS PICTUS, Kner & Steindachner.

(GUAVINA.)

Head 3 to  $3\frac{1}{2}$ ; depth 6. D. VI-I, 7 or 8; A. I, 7 or 8; lateral line 60; 24 scales in an oblique series from front of soft dorsal downward and back-

\* *Eleotris belizianus* is described as follows: Head 4 in total; depth 5. D. VI-I, 9; A. I, 8; scales 60; eye 5 in head. Preopercle with a spine turned downward; 16 rows of scales between soft dorsal and anal; scales of top of head a little smaller than those of body, extending forward nearly to front of eyes; cheeks scaly; scales elliptate. Interocular space flattened,  $\frac{1}{2}$  broader than eye; snout depressed a little longer than eye; lower jaw prominent; outer teeth enlarged; maxillary reaching front of eye. Dorsals contiguous. Color brownish, faint dark streaks on the fins. Belize; Cayenne. (Sauvage.) Length 100 mm. Evidently not different from *E. pisonis*.



ward to anal; about 20 in a vertical series. Body elongate, depressed anteriorly; head especially very broad and flat; mouth large, broad, very oblique, the maxillary reaching nearly or quite to opposite posterior margin of eye, its length  $2\frac{1}{2}$  to  $2\frac{3}{4}$  in head; lower jaw considerably projecting. Teeth in jaws all equal, in broad bands, the outer not at all enlarged. Eye small, anterior, its length in adult 2 in interorbital width, which width is about 3 in head; a conspicuous knob at upper anterior and posterior angles of orbit; preopercular spine well developed, strong, compressed, directed downward and forward. Scales on head very small, mostly cycloid, covering cheeks and opercles and upper part of head to the eyes; scales on body smaller and smoother than in most other species, those on belly much smaller than those on sides; scales on back and belly cycloid, only those on sides distinctly ctenoid. Pectoral fins moderate, reaching to near end of base of first dorsal,  $1\frac{1}{2}$  in head; ventrals inserted just behind axil, reaching halfway to vent, about 2 in head. Interspace between dorsals about equal to diameter of eye. Soft dorsal and anal short and high, very similar, coterminous; last ray of anal a little longer than  $\frac{1}{2}$  length of head; caudal peduncle long, a little shorter than head. Caudal fin rounded,  $1\frac{1}{2}$  in head. Color\* dark, dull olivaceous brown, paler below; younger individuals mottled below with bluish and speckled with dark brown; sides without longitudinal stripes; fins dusky, all of them finely mottled and speckled with darker, the dark markings on dorsal and anal forming undulated longitudinal stripes; on pectorals and ventrals forming dark bars. Distinguished from related species by the larger mouth with small, equal teeth, and the small, smoothish scales. Length about 18 inches. Streams of the Pacific Coast, from Sonora, south to Panama; abundant in Rio Presidio, at Mazatlan, where the types of *E. aequidens* were taken; not rare about Panama. (*pictus*, painted.)

*Eleotris pictus*, KNER & STEINDACHNER, Abh. Ab. Wiss. Wien 1864, 18, pl. 3, f. 1, Rio Bayano, near Panama; depth 6 to 7 in total length; scales 60.

*Culius aequidens*, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1881, 461, Rio Presidio, near Mazatlan. (Types, Nos. 28268 and 29240. Coll. Gilbert.)

*Eleotris aequidens* JORDAN & EIGENMANN, l. c., 483.

### 805. ALEXURUS, Jordan.

*Alexurus*, JORDAN, Proc. Cal. Ac. Sci. 1895, 512 (*arniger*).

Body elongate, covered with small cycloid scales; preopercle with a small, concealed, hooked spine at its angle, as in *Eleotris*; caudal fin broad, its base with many procurrent rays. In other respects similar to *Eleotris*. One species known; mariue. (ἀλέξω, to defend; ὄψα, tail, from the caudal fulcra.)

\* A young example shows the following details of coloration in life: Blackish everywhere, sides with faint whitish streaks, along rows of scales a broad, blackish lateral band occupying whole of side, back and belly paler, traces of faint dark cross bands; caudal black, with a pale margin and some dark cross shades; pectorals, dorsals, and ventrals more or less barred with black; preopercular spine well developed; a whitish bar at base of caudal with a darker one before it.

2525. ALEXURUS ARMIGER, Jordan & Richardson.

Head  $4\frac{1}{2}$ ; depth 8. D. VI-13; A. 11; V. I, 5; scales about 102-30; eye 8 in head; maxillary  $2\frac{1}{2}$ ; mandible  $2\frac{1}{2}$ ; snout  $5\frac{1}{2}$ ; interorbital  $4\frac{1}{2}$ ; pectoral  $1\frac{1}{2}$ ; caudal equals head; ventral 2; last dorsal ray  $1\frac{1}{2}$ . Body long and low, compressed posteriorly, depressed in front. Head flattish and broad above, the cheeks moderately tumid. Eyes small, high up, separated by a broad, flattish, interorbital space; snout short; mouth moderate, very oblique, the maxillary ceasing below the center of pupil; lower jaw very heavy, oblique, projecting beyond upper, its outline horseshoe-shaped, obtuse in front. Teeth in rather broad bands, the outer enlarged below, but scarcely so above; none of them canine-like. Top of head with very small scales; cheeks and opercles with rudimentary scales above; preopercle with a concealed antrorse hook below, as in *Eleotris*; scales on body very small, perfectly smooth, partially embedded; scales on nape and throat minute. Gill membranes extending a little forward below, so that the branchiostegals are free from the isthmus. Insertion of dorsal twice as far from middle of base of caudal as from tip of snout; the fin low, its slender rays slightly filamentous; soft dorsal low, its last ray highest; anal similar, beginning under second dorsal ray; caudal long, bluntly pointed behind, with strongly procurrent base above and below, the base above  $\frac{2}{3}$  length of head, formed of 14 short rays, that below a little shorter, of 12 rays, this procurrent portion forming an angle with the caudal proper where it joins it; pectorals and ventrals short, the ventrals inserted under pectorals. Color olive green, dusky above, paler below, but everywhere covered with fine black dots; both dorsals with the membranes pale, the rays each barred with black; caudal mesially blackish, all the rays barred or checkered in fine pattern; pectoral and anal pale, similarly speckled, base of pectoral dusky; ventral finely speckled. La Paz, Lower California; 1 specimen,  $6\frac{1}{2}$  inches long, taken by Mr. James A. Richardson. (*armiger*, bearing arms, from the concealed spine.)

*Alexurus armiger*, JORDAN & RICHARDSON, Proc. Cal. Ac. Sci. 1895, 511, pl. 48, La Paz. (Type in L. S. Jr. Univ. Mus. Coll. James A. Richardson.)

806. EROTELIS, Poey.

(ESMERALDAS DE MAR.)

*Erotelis*, POEY, Memorias, II, 273, 1861 (*valenciennesi* = *smaragdus*).

Body very slender, elongate, covered with minute cycloid scales. Ventrals separate, the rays I, 5. No teeth on vomer. Lower pharyngeals subtriangular, the teeth stiff, villiform, none of them lamelliform. Post-temporal bones short, strongly divergent, the distance between their insertions about equal to the narrow interorbital space; top of head with a strong median keel, highest on the occipital region; no supraoccipital crest; no preopercular spine. Mouth very oblique. One species known; strictly marine. (Name an anagram of *Eleotris*.)

2526. *EROTELIS SMARAGDUS* (Cuvier & Valenciennes).

(ESMERALDA NEGRA; ESMERALDA DE MAR.)

Head  $4\frac{1}{2}$  to  $5\frac{1}{2}$ ; depth 8 to 12. D. VI-I, 10; A. I, 9; V. I, 5; scales 100. Body very long and slender, compressed behind, the form much as in *Gobius oceanicus*. Head depressed, flattish above, the eyes mostly superior, not  $\frac{1}{2}$  the width of the interorbital area, which has a knob near its middle. Mouth very oblique, the lower jaw much projecting, the maxillary about reaching front of eyes; teeth rather small, in bands. Fins rather high; dorsal spines slender, lower than the highest soft rays, which are  $1\frac{1}{2}$  in head; caudal lanceolate,  $\frac{1}{2}$  longer than head; ventrals moderate, 2 in head. Scales very small, cycloid. Color very dark green, almost black; the fins mostly bluish, the dorsal with brown lines; some dark markings about eye and on base of pectoral above. Length 8 inches. Coral shores among green algae; known from Key West and Cuba; not common; not entering rivers. Here described from Key West specimens. (*σμαραγδός*, emerald.)

*Eleotris smaragdus*, CUVIER & VALENCIENNES, Hist. Nat. Poiss., XII, 231, 1887, Cuba; JORDAN & GILBERT, Proc. U. S. Nat. Mus., 1884, 143.

*Erotelis valenciennesi*, POEY, Memorias, II, 273, 1861, Cuba.

*Erotelis smaragdus*, JORDAN & EIGENMANN, l. c., 484.

807. *GYMNELEOTRIS*, Bleeker.

*Gymneleotris*, BLEEKER, Esquisse d'un Syst. Nat. des Gobioid., 304, 1874 (*seminuda*).

Body scaled only posteriorly, the anterior half and the head naked. Ventrals separate, I, 5. Vomer without teeth. Isthmus broad; skull without crests. Lower jaw with 4 large recurved teeth. Otherwise essentially as in *Eleotris*, the preopercle probably without spine. (*γυμνός*, naked; *Eleotris*.)

2527. *GYMNELEOTRIS SEMINUDUS* (Günther).

Head  $3\frac{1}{2}$ . D. VII-11; A. 9. Head depressed, broader than high, flat above. Snout rather obtuse, longer than eye, lower jaw somewhat prominent; cleft of mouth extending to below anterior margin of orbit. Teeth in upper jaw in a narrow band, the lower having 4 somewhat larger and recurved teeth in front, appearing to form a single series; palate toothless. None of the fin rays prolonged; pectoral not quite extending to origin of second dorsal; ventral much shorter than pectoral, its inner ray the longest, the others gradually decreasing in length outward; caudal fin rounded. Head and trunk naked; tail covered with small scales. Brown, with numerous well-defined white cross stripes on head as well as on body; vertical fins black. Panama. (Günther); known from the type only, a young example,  $1\frac{1}{2}$  inches long; not seen by us. (*seminudus*, half-naked.)

*Eleotris seminuda*, GÜNTHER, Proc. Zool. Soc. London 1864, 24, pl. 4, figs. 2, 2a, Panama; GÜNTHER, Fish. Centr. Amer., 441, 1869.

*Gymneleotris seminuda*, JORDAN & EIGENMANN, l. c., 484.

Chrio  
TH  
scale  
Head  
rate,  
rays.  
what  
in up  
on inn

Hea  
maxil  
but w  
diamet  
head;  
length  
lower;  
head; l  
uniform  
anal bl  
2825, of

Chrioepi  
Gulf

Sicydium,

Body  
oblong  
promin  
series of  
extremi  
ous mov  
behind t  
teeth hid  
out or br  
all simple  
of moder  
caudal q  
openings  
streams o  
gourd-sha

a. Body o  
b. Sc  
bb. Sc  
aa. Body

808. CHRIOLEPIS, Gilbert.

*Chriolepis*, GILBERT, Proc. U. S. Nat. Mus. 1891, 557 (*minutillus*).

This genus differs from *Gymneleotris*, Bleeker, in the total absence of scales, and the absence of enlarged canines in the front of the mandible. Head and body compressed, the former as deep as wide. Ventrals separate, near together, the inner rays longest, each with 1 spine and 5 soft rays. Teeth in a rather wide band in upper jaw, the outer series somewhat enlarged. Teeth in mandible in a single series, similar to outer row in upper jaw, none of them canine-like. Gill slits narrow; no dorsal flaps on inner edge of shoulder girdle. Size small. (*χρεια*, want; *λεπίς*, scale.)

2528. CHRIOLEPIS MINUTILLUS, Gilbert.

Head  $3\frac{1}{2}$ ; depth  $4\frac{1}{2}$  in length. D. VII-12; A. 11. Mouth oblique, the maxillary reaching to below middle of orbit,  $2\frac{1}{2}$  in head; eyes high up, but with lateral range, separated by a narrow interorbital space less than diameter of pupil; diameter of orbit nearly twice length of snout,  $3\frac{1}{2}$  in head; dorsal spines high and slender, but not filamentous, the longest  $\frac{1}{2}$  length of head; soft dorsal rays higher, nearly  $\frac{2}{3}$  length of head; the anal lower; caudal short, broadly rounded, the depth of peduncle  $\frac{1}{2}$  length of head; length of pectoral equaling that of head without snout. Color uniform light brown on head and body, above and below; fins dusky, the anal blackish. A single specimen, 1 inch long, from Albatross Station 2825, off the east coast of Lower California. (*minutillus*, very small.)

*Chriolepis minutillus*, GILBERT, Proc. U. S. Nat. Mus. 1891, 558, Albatross Station 2825, Gulf of California, in 79 fathoms.

809. SICYDIUM, Cuvier & Valenciennes.

*Sicydium*, CUVIER & VALENCIENNES, Hist. Nat. Poiss., XII, 168, 1837 (*plumieri*).

Body subcylindrical, covered with rather small ctenoid scales; head oblong and broad, with cleft of mouth nearly horizontal; upper jaw prominent; snout obtusely rounded; lips very thick, the lower with a series of numerous slender horizontal teeth, of which sometimes only the extremities are visible; upper jaw with a single uniform series of numerous movable small teeth attached by ligament to edge of maxillary; behind this outer visible series lie numerous other parallel series of young teeth hidden in the gum, which succeed the former as they become worn out or broken; lower jaw with a series of widely set conical teeth; teeth all simple, slender, the distal half bent inward nearly at a right angle; eyes of moderate size; 2 dorsal fins, the anterior with 6 (5 or 7) flexible spines; caudal quite free; ventrals united into a short cup-shaped disk; gill openings of moderate width; 4 branchiostegals. Species few in the streams of the West Indies. (*σικύδιον*, diminutive of *σικύα*, a gourd, or gourd-shaped cupping glass, from the ventral disk.)

- a. Body covered with small scales.
- b. Scales very small, about 84.
- bb. Scales moderate, about 68.
- aa. Body nearly naked.

PLUMIERI, 2529.  
ANTILLARUM, 2530.  
VINCENTE, 2531.

## 2520. SICYDIUM PLUMIERI (Bloch).

(SIRAJÓ.)

Head 4 to 4½; depth 4½; eye 6 to 7 in head, 2 to 3 in interorbital width. D. VI-I, 10; A. I, 10; scales 84. Teeth in upper jaw long, slender, bent inward at right angles, only the lips protruding from the gums. Front teeth of lower jaw not larger than those behind; a single row of inconspicuous papillæ on the gum beneath the upper lip, a large median papilla above the maxillary suture; a median cleft in the upper lip. Pectorals longer than head; third, fourth, and fifth dorsal spines produced into long ribbons, the fourth, which is the longest, being 2 to 3 times height of body. Body usually covered with small scales, reduced in size on neck and belly; frequently almost naked, the scales present only on posterior part of body. Caudal deeply emarginate. Color olive or violet brown, with about 7 more or less distinct dark vertical bars; a dark bar at base of pectoral; dorsal with irregular dark markings; anal fin with a dark marginal band, sometimes edged with white; an H-shaped figure on base of caudal fin, and a black bar on its posterior half. Fresh waters of the West Indies. (Named for Père Charles Plumier, who discovered the species at Martinique.)

*Gobius plumieri*, BLOCH, Ichth., 125, pl. 178, fig. 3, 1786, Martinique; on a drawing by PLUMIER.

*Sicydium siragus*, POEY, *Memorias*, II, 278, 1861, Santiago de Cuba.

*Sicydium plumierii*, CUVIER & VALENCIENNES, *Hist. Nat. Poiss.*, XII, 168, 1837; GILL, *Proc. Ac. Nat. Sci. Phila.* 1860, 101; GÜNTHER, *Cat.*, III, 92, 1861; OGILVIE-GRANT, *Proc. Zool. Soc. Lond.* 1884, 156, pl. 11, fig. 1; JORDAN & EIGENMANN, *L. c.*, 484; EIGENMANN & EIGENMANN, *L. c.*, 56.

## 2530. SICYDIUM ANTILLARUM, Ogilvie-Grant.

Head 4½; depth 6; width of head ½ length. D. VI-I, 10; A. I, 10; scales 68. Teeth in upper jaw long, slender, and bent inward over the gum at right angles. A row of small lamelliform transverse papillæ on the gum beneath upper lip, with a larger median lamelliform papilla above maxillary suture; a slight median cleft in upper lip; maxillæ at right angles to one another; horizontal teeth conspicuous. Scales on body and tail subequal and larger than those on neck and belly. Maxilla not extending to vertical from posterior margin of eye, the diameter of which is contained 6½ times in length of head and twice in interorbital space. Length of pectoral greater than that of head. The third, fourth, and fifth dorsal spines produced into long narrow ribands; the fourth, which is longest, nearly 3 times height of body; second dorsal considerably higher than body. Color uniform violet brown; dorsal fins with irregular wavy dark markings; anal with a black and white marginal band; caudal with a dark band on upper margin. One specimen, 4¼ inches long, from Barbados (Ogilvie-Grant); not seen by us. (*antillarum* of the Antilles.)

*Sicydium antillarum*, OGILVIE-GRANT, *Proc. Zool. Soc. Lond.* 1884, 157, Barbados.

A  
by  
men  
are  
erab  
thos  
Cauc  
bars  
spots  
half  
with  
types  
of St  
plain  
the b  
are pl  
Sicydiu  
nan

Cotylopo  
(acu  
Sicya, J

This  
in exte  
jaw; it  
triensp  
(usp) or  
(usp)

SICYA (a  
a. Teet  
l  
b.  
bb.

Head  
eye 6 in  
cuspid, t  
very soft  
papillæ  
papilla;  
an angle  
strongly

2531. *SICYDIUM VINCENTE*, Jordan & Evermann, new species.

Another species of *Sicydium* or of some related genus is thus mentioned by Dr. Eigenmann: "Mr. Samuel Garman collected several hundred specimens of this species at Kingston, St. Vincent. Most of these specimens are less than an inch in length, the longest  $1\frac{1}{2}$  inches; they differ considerably in coloration from the adult; most are entirely naked, a few of those examined having scales only on the posterior part of the body. Caudal deeply emarginate. There are traces of about 7 dark vertical bars; a black bar at base of pectoral; dorsals with several series of black spots; an H-shaped figure on base of caudal, a black bar on the posterior half of caudal fin; belly and lower part of body plain; everywhere else with black points. The specimens collected by Mr. Garman may be the types of a new species. No large specimens were collected at the Island of St. Vincent. Specimens  $1\frac{1}{2}$  inches in length from Hayti have the fins plain and a series of blotches along the middle of the posterior part of the body; the body, except the belly, is entirely covered with scales which are plainly ctenoid." (Eigenmann.) (Named for St. Vincent.)

*Sicydium vincente*, JORDAN & EVERMANN, Check-List Fishes, 456, 1896, St. Vincent Island; name only.

310. *COTYLOPUS*, Guichenot.

*Cotylopus*, GUICHENOT, in Maillard, Notes sur l'Isle de la Réunion, II, Addendum 9, 1864 (*acutipinnis*).

*Sicya*, JORDAN & EVERMANN, Check-List Fishes, 456, 1896 (*gymnogaster*).

This genus is closely allied to *Sicydium*, agreeing closely with the latter in external characters and in the absence of larger teeth in front of lower jaw; it differs chiefly in the form of the upper teeth which are curved, tricuspid, and trident-shaped, the middle cusp either permanent (*Cotylopus*) or else worn away leaving the tooth apparently bicuspid (*Sicya*). (*νοῦλον* cup; *πούς*, foot.)

*SICYA* (*σκύα*, a gourd, or gourd-shaped cup):

a. Teeth in upper jaw curved, tricuspid, trident-shaped, the lateral lobes long, the middle short and suspended between the outer lobes, and soon wearing away leaving the tooth apparently bicuspid.

b. Neck and belly naked; a double or triple row of small papillæ on the gum beneath the upper lip.

bb. Neck and belly covered with small scales; gum beneath the upper lip smooth.

Gymnogaster, 2532.  
SALVINI, 2533.

Subgenus *SICYA*, Jordan & Evermann.

2532. *COTYLOPUS GYMNOGASTER* (Ogilvie-Grant).

Head  $4\frac{1}{2}$  to 5; depth  $5\frac{1}{2}$  to 6. D. VI-I, 10; A. I, 10; scales 60 to 64; eye 6 in head, twice in interorbital space. Teeth in the upper jaw tricuspid, the middle cusp, which is situated at the anterior end of tooth, is very soft and soon becomes worn away. A double or treble row of small papillæ on the gum beneath the upper lip, without a larger median papilla; upper lip with a very slight median notch; maxillæ containing an angle of about  $75^{\circ}$ ; horizontal teeth more or less inconspicuous. Scales strongly ctenoid; neck and belly naked. Length of pectoral greater than

that of head. Second, third, and fourth dorsal spines produced into filaments; the third, which is the longest, twice height of body; second dorsal higher than body. Color violet brown, yellowish in young specimens, shaded with indistinct transverse bands of darker; irregular brown spots on axis of pectoral, and a broad dark band from base of pectoral to root of caudal, both more or less indistinct in adult specimens; fins violet, clouded with darker. Length  $4\frac{1}{2}$  inches. Streams about Mazatlan (Ogilvie-Grant); not seen by us. (*γυμνός*, naked; *γαστήρ*, belly.)

*Sicydium gymnogaster*, OGILVIE-GRANT, Proc. Zool. Soc. Lond. 1884, 158, pl. 11, fig. 2, and pl. 12, fig. 6, Mazatlan.

*Sicyopterus gymnogaster*, JORDAN & EIGENMANN, l. c., 485.

#### 2533. COTYLOPUS SALVINI (Ogilvie-Grant).

Head  $4\frac{1}{2}$ ; depth  $6\frac{1}{2}$ . D. VI-9 or 10; A. I, 10; scales 78; eye  $5\frac{1}{2}$  in head, twice in interorbital space. Teeth in upper jaw tricuspid; the middle cusp, which is situated at anterior end of tooth, very soft and soon becomes worn away. Gum beneath upper lip smooth; a median papillose tubercle above maxillary suture; upper lip with a small median notch; maxilla containing an angle of about  $75^\circ$ ; horizontal teeth conspicuous. Scales ctenoid, those on neck and belly smaller than those on body and tail. Length of pectoral rather greater than that of head. Second and third dorsal spines subequal and produced into short filaments,  $1\frac{1}{2}$  times height of body; second dorsal not so high as body. Color olive brown; anal yellow, with a black and white band along margin; membrane of second dorsal clear, spotted with brown; caudal with a dark and yellow band round the extremity. Length  $4\frac{3}{8}$  inches. Streams near Panama; 1 specimen known. (Ogilvie-Grant.) (Named for Osbert Salvin, who collected largely in Central America for the British Museum.)

*Sicydium salvini*, OGILVIE-GRANT, Proc. Zool. Soc. Lond. 1884, 159, pl. 12, fig. 2, Panama.

*Sicyopterus salvini*, JORDAN & EIGENMANN, l. c., 485.

#### 811. EVORTHODUS, Gill.

*Evorthodus*, GILL, Proc. Ac. Nat. Sci. Phila. 1859, 195 (*breviceps*).

Body elongate, covered with ctenoid scales of moderate size. Head thick, short, naked. Isthmus moderate. Teeth in a single series, with the crown emarginate, those of the lower jaw horizontal; no canines. First dorsal of 6 spines; ventral fins united, not adherent to the belly, otherwise as in *Gobius*, so far as known. (*εὖ*, well; *ὀρθός*, straight; *ὀδόν*, tooth.)

#### 2534. EVORTHODUS BREVICEPS, Gill.

Head  $4\frac{1}{2}$ , about as deep as wide; depth  $4\frac{1}{2}$ . D. VI-I, 10; A. I, 11; eye 3. Teeth emarginate, uniserial, those of lower jaw nearly horizontal. Snout blunt, profile evenly decurved; caudal rounded, 3 in length of body; some of the dorsal rays filamentous. Color light brown, with irregular blackish blotches along sides; 2 black spots at base of caudal fin, 1 above the other, alternating with 1 more anterior on the peduncle; first dorsal

wi  
rov  
not  
Evo

Loph

Do

muc

11 +

foran

ably

disti

area

Hea

9 or 10

little

naked

spinot

profile

obliqu

border

bands,

series

upper

produc

caudal

the up

median

at the

interor

the orb

large fe

dorsal l

plain;

souther

of Cuba

water a

*Gobius cy*

VALE

*Gobius cr*

1837,

*Lophogob*

ENNUN

MANN

with 2 bands parallel with its upper margin; second dorsal with 3 narrow longitudinal bands. (Gill.) Fresh waters of Trinidad and Surinam; not seen by us. (*brevis*, short; *-ceps*, head.)

*Eporthodus breviceps*, GILL, Proc. Ac. Nat. Sci. Phila. 1859, 105, Trinidad; JORDAN & EIGENMANN, l. c., 486.

812. LOPHOGOBIUS, Gill.

(CRESTED GOBIES.)

*Lophogobius*, GILL, Proc. Ac. Nat. Sci. Phila. 1862, 240 (*crisitagalli*=*cyprinoides*).

Dorsal spines 6; scales evidently ctenoid. Body short, compressed, form much as in *Dormitator*; nape with fleshy crest; scales large. Vertebrae 11 + 15. Interorbital area of cranium anteriorly elevated, with a large foramen-like depression in front of eye. One species, differing considerably in form from the other Gobies. The study of its skeleton shows no distinction of much importance unless the peculiar form of its interorbital area be regarded as such. (*λόφος*, crest; *gobius*.)

2535. LOPHOGOBIUS CYPRINOIDES (Pallas).

Head  $3\frac{3}{4}$ ; depth  $3\frac{3}{4}$ ; greatest width  $5\frac{1}{2}$  to  $6\frac{1}{2}$ . D. VI or VII-10 or 11; A. 9 or 10; scales 26 to 30; vertebrae 11 + 15; eye  $3\frac{1}{2}$  to 4. Body short and deep, little compressed, formed much as in *Cyprinodon*; head naked, a prominent naked dermal crest extending from above middle of eye to near front of spinous dorsal; interorbital width slightly less than diameter of eye; profile convex; snout short, bluntish, about as long as eye; mouth very oblique, the gape slightly curved; front of upper lip on level of lower border of eye; lower jaw somewhat projecting; teeth in both jaws in bands, the outer series erect and somewhat enlarged, those of the inner series small; scales large, reduced on breast and nape; a few scales on upper part of opercle; median line before dorsal naked; dorsal spines produced in short filaments; last rays of soft dorsal reaching caudal; caudal rounded; pectorals lanceolate, reaching beyond insertion of anal, the upper rays not silk-like; skull very broad and short, with low, median crest, highest behind; double crests of temporal region joining at the upper posterior angles of the eyes and forming a bridge over the interorbital area, the crests ending abruptly above the anterior part of the orbit, forming a decided angle, the bridged interorbital leaving a large foramen in front of this angle. Color blackish green in life; spinous dorsal black; soft dorsal, ventrals, and anal dark, plain; pectorals lightish, plain; caudal finely mottled. Length 2 inches. West Indies, north to southern Florida; generally common in the streams and brackish waters of Cuba and other islands. Recently taken by Dr. Evermann in brackish water at Biscayne Bay, Florida. (*κυπρίνος*, carp; *εἶδος*, resemblance.)

*Gobius cyprinoides*, PALLAS, Spicilegia, Zool., VIII, 17, pl. 1, fig. 5, 1770, Amboina; CUVIER & VALENCIENNES, Hist. Nat. Poiss., XII, 129, 1837; GÜNTHER, Cat. Fish., III, 8, 1861.

*Gobius crisitagalli*, VALENCIENNES, in CUVIER & VALENCIENNES, Hist. Nat. Poiss., XII, 130, 1837, Havana; GUICHENOT, in Ramon de la Sagra, Hist. Cuba, 128, pl. 3, fig. 3, 1850.

*Lophogobius cyprinoides*, POEY, Repertorio, I, 335, 1867; POEY, Synopsis, 393, 1868; POEY, Enumeratio, 125, 1876; JORDAN & EIGENMANN, Proc. U. S. Nat. Mus. 1886, 487; EVERMANN & KENDALL, Bull. U. S. Fish Comm. 1897, 131, plate 9, fig. 13.



## 813. GOBIUS (Artedi) Linnaeus.

## (GOBIES.)

- Gobius*, ARTEDI, Genora, 28, 1738 (*Gobius ex nigricante varius*, etc., = *niger*).  
*Gobius*, LINNÆUS, Syst. Nat., Ed. x, 262, 1758 (*niger*, etc.), and of authors generally.  
*Gobionellus*, GIRARD, Proc. Ac. Nat. Sci. Phila. 1858, 168 (*hastatus* = *oceanicus*).  
*Ctenogobius*, GILL, Fish. Trinidad, 374, 1858 (*fasciatus*).  
*Euctenogobius*, GILL, Annals Lyc. Nat. Hist. New York 1859, 45 (*badius*).  
*Smaragdus*, POEY, Memoriae, II, 279, 1861 (*smaragdus*).  
? *Pomatoschistus*, GILL, Proc. Ac. Nat. Sci. Phila. 1863, 263, footnote (*minutus*).  
*Coryphopterus*, GILL, Proc. Ac. Nat. Sci. Phila. 1863, 263 (*glaucofrœnum*).  
? *Deltentosteus*, GILL, Proc. Ac. Nat. Sci. Phila. 1863, 263, footnote (*quadrimaculatus*).  
? *Gobiichthys*, KLUNZINGER, Fisch. Rothen Meer, 179, 1871 (*petersii*).  
? *Mesogobius*, BLEEKER, Esquisse d'un Syst. Nat. Gobioid., 317, 1874 (*guavina*).  
? *Stenogobius*, BLEEKER, l. c., 317 (*gymnopomus*).  
? *Oligolepis*, BLEEKER, l. c., 318 (*melanostigma*).  
? *Gnatholepis*, BLEEKER, l. c., 318 (*anjerensis*).  
? *Callogobius*, BLEEKER, l. c., 318 (*hasselti*).  
? *Hypogymnogobius*, BLEEKER, l. c., 318 (*xanthozona*).  
? *Hemigobius*, BLEEKER, l. c., 318 (*melanurus*).  
? *Cephalogobius*, BLEEKER, l. c., 320 (*sublitus*).  
? *Acentrogobius*, BLEEKER, l. c., 321 (*chlorostigma*).  
? *Porogobius*, BLEEKER, l. c., 321 (*schleyeri*).  
? *Amblygobius*, BLEEKER, l. c., 322 (*sphinx*).  
*Zonogobius*, BLEEKER, l. c., 323 (*semifasciatus*).  
? *Odontogobius*, BLEEKER, l. c., 323 (*bynoënsis*).  
? *Stigmatogobius*, BLEEKER, l. c., 323 (*pleurostigma*).  
? *Ozyurichthys*, BLEEKER, l. c., 324 (*beloaso*).  
*Lythrypnus*, JORDAN & EVERMANN, Check-List Fishes, 458, 1896 (*dallii*).

Body oblong or elongate, compressed behind. Head oblong, more or less depressed. Eyes high, anterior, close together; opercles unarmed. Mouth moderate. Teeth on jaws only, conical, in several series, those in the outer row enlarged; no canines. Isthmus broad. Shoulder girdle without fleshy flaps or papillæ. Skull depressed, abruptly widened behind the eyes and without distinct median keel. Scales moderate, tenoid, permanently covering the body; cheeks usually naked; belly generally scaly. Dorsal with 6 rather weak spines; pectorals well developed, the upper rays sometimes very slender and silky; ventrals completely united, not adnate to the belly; caudal fin usually obtuse. Species very numerous. The genus *Gobius*, as here understood, comprises a very large number of species more or less closely related to the European type of the genus, *Gobius niger*, and its American relative, *Gobius saporator*. An examination of skulls or skeletons of numerous European and American species shows a remarkable uniformity in most respects. The general form and structure of the cranium is the same in all, the only differences being very minor ones in the height of certain crests. *Gobius oceanicus* seems the most aberrant, but seems to be inseparable generically on account of intermediate forms. Probably several of the many genera indicated by Bleeker will prove valid, but only a thorough study of skeletons can establish them. It is not unlikely that *Ctenogobius*, to which group most of our species belong, may be separable from *Gobius*. (*κωβίδος*; Latin, *Gobius* or *Gobio*, a name applied to the gudgeon (*Gobio gobio*) and

to other small fishes; allied to *Cobitis*, chub, etc. According to Apostolides *κωβιδός* and *γωβιδός* are common names in modern Greek for all species of the genus *Gobius*. Aristotle *κωβιδός*, 610b, 4, 598a 11 16, 508b 16, 569b 23, 621b 13 19, 567b 11, 591b 13, 601b 22, 835b 14. The *κωβιδός* has many pyloric appendages above the stomach, spawns near the land on the rocks, the bunches of eggs are flat and crumbling; it feeds on mud, seaweed, sea moss, etc.; lives near the land, gets fat in the rivers, and is found in schools. The white *κωβιδός*, found in the Euripus of Lesbos, never leaves that lagoon for the open sea as the other fishes found there do. Latin *Gobio* and *Cobio*, Plin. *Gobius*, Ovid., Hal. 12, 8. Martial 13, 88. Horace A. Hoffman.)

GObIUS:

a. Upper rays of pectoral fin silk-like; i. e., short and very slender and flexible, free for nearly their whole length.

b. Body robust, compressed posteriorly; depth  $4\frac{1}{2}$  to  $5\frac{1}{2}$  in length; head broad, low, rounded in profile, its length  $3\frac{1}{6}$  to  $3\frac{1}{3}$  in body; eye 4 to 5 in head; mouth large, little oblique; lips thick; teeth in both jaws in bands, the outer series enlarged; those on lower jaw subequal; scales large, strongly ctenoid, smaller on nape and belly; dorsal spines short, none filamentous. Color olivaceous, light or dark, varying from sand color to greenish black, everywhere mottled and marbled with dark and paler; a faint dusky spot behind eye. D. VI-9 or 10; A. I, 7 to 9. Scales 36 to 41. SOFORATOR, 2536.

CTENOGObIUS (*kreis*, comb; *Gobius*):

aa. Upper rays of pectoral normal, not silk-like, similar to the others.

c. Scales large, 25 to 33.

d. Color in life olivaceous, more or less spotted, never red.

e. Dorsal soft rays 12 to 14; vertex and nape with a slight median fold of skin.

f. Body compressed, its depth 5 in length; head  $3\frac{1}{2}$  to  $3\frac{1}{3}$ ; eye 3 to  $3\frac{1}{4}$  in head; vertex and nape with a slight median fold of skin; maxillary reaching about to front of pupil; lower jaw very slightly produced; teeth in bands, the outer slightly enlarged. Olivaceous; spinous dorsal black at tip; second dorsal finely checkered in adult. D. VI-14; A. I, 11. Scales 25 or 26-10. NICHOLSON, 2537.

ff. Body long, not much compressed; head  $3\frac{1}{2}$ ; eye 3 in head; no median fold on vertex and nape; a dark spot on first dorsal.

EIGENMANNI, 2538.

ee. Dorsal soft rays 10 to 12; no median fold of skin on vertex and nape.

g. Caudal with 2 spots at its base; jaws unequal, the lower slightly produced; body robust, compressed behind, the depth 5 in total length; head  $4\frac{1}{2}$ ; eye longer than snout,  $3\frac{1}{2}$  in head; maxillary reaching pupil; teeth in a band, the outer enlarged and distant, the inner enlarged and bent backward. Brownish; a faint blue spot on each scale; six spots along middle of back; similar spots on scapular region and middle of sides; 2 spots on base of caudal; a dark spot above opercle; blue dots on head; a straight blue line crossing cheek above and continued on opercle; dorsals faintly spotted. D. VI-10; A. 10. Scales 25-7. (Gill.)

OLAUCOPRENUM, 2539.

gg. Caudal plain or with but a single spot at its base.

h. Dorsal spines low, the highest little longer than head.

i. Region from nape to dorsal entirely scaled.

j. Pores on preopercle not very conspicuous; no canine teeth.

k. Body very slender, compressed, the depth  $5\frac{1}{2}$  in length; caudal much longer than head; mouth rather large, the lower jaw projecting; teeth unequal, rather strong; yellowish, much spotted with darker. D. VI-12; A. 12. Scales 35.

MANGLICOLA, 2540.

kk. Body subfusiform, little compressed; depth  $4\frac{1}{2}$  in length; head blunt, 4 in length, rounded in profile; eye equal to snout, 4 in head. Mouth small, horizontal, the lower jaw included; maxillary 3 in head, reaching to below eye. Teeth small, in bands in both jaws, the outer enlarged, those of the upper jaw very slender. Scales large, ctenoid, those of nape and belly little reduced. Longest dorsal spine shorter than head. Caudal scarcely pointed, about as long as head. Color whitish gray, middle of sides with 4 or 5 dark blotches, from each of which a narrow dark bar extends downward and forward; a large black blotch above pectorals, obsolete in female; a small black spot at base of caudal; a dark mark below eye; vertical fins barred. D. VI-12; A. 11 or 12. Scales 33.

STIGMATURUS, 2541.

jj. Pores on preopercle very conspicuous; lower jaw with small canines. D. VI-I, 9; A. I, 9.

QUADRIPORUS, 2542.

ii. Region between nape and dorsal with a narrow naked median strip. Body moderately elongate, subfusiform, the depth  $5\frac{1}{2}$  in length. Head large, not so blunt as in *G. boleosoma*, its length  $3\frac{1}{2}$  to  $3\frac{3}{4}$  in length; anterior profile gently decurved; snout  $3\frac{1}{2}$  to  $3\frac{3}{4}$  in head; eye 4; mouth large, slightly oblique; maxillary extending to front of pupil,  $2\frac{1}{2}$  in head. Teeth small, slender and curved, in moderate bands. Scales moderate, ctenoid, those in front much reduced in size; breast naked. Longest dorsal spine  $1\frac{1}{2}$  in head. Caudal as long as head, somewhat pointed. Olivaceous, mottled with gray; about 5 rounded dark blotches along middle of sides, the last forming a spot at base of caudal; no dark spot on side of nape; some dark marks on head; vertical fins barred. D. VI-12; A. 13. Scales 33 to 35.

SHUFELDTI, 2543.

iii. Region between nape and dorsal entirely naked.

l. Highest rays of second dorsal little more than  $\frac{1}{2}$  head, none of them reaching base of caudal.

m. Profile much decurved, skull rounded behind, without distinct median ridge; mouth horizontal. Body elongate, deepest below front of dorsal, tapering regularly backward, the greatest depth  $5\frac{1}{2}$  in length. Head short, blunt, pro-

file anteriorly abruptly decurved, cheek somewhat swollen. Length of head  $3\frac{1}{2}$  in body. Snout about equal eye,  $3\frac{1}{2}$  in head. Mouth horizontal, maxillary reaching to below pupil (in male); lower jaw included. Teeth in each jaw in a band, the outer row of the upper jaw large, recurved. Scales large, ctenoid, somewhat reduced anteriorly. Nape, breast, and belly naked. Dorsal spines about  $\frac{2}{3}$  of head. Caudal pointed,  $2\frac{1}{2}$  to  $3\frac{1}{2}$  in body. Color olivaceous, with numerous dark reticulations on the back; 5 black spots along the sides, the last forming a spot on base of caudal, sometimes with V-shaped dark bars extending from them to dorsal; breast and sides of belly with numerous dark specks in male; a dark line between eyes; a dark line from eye to middle of premaxillary, some dark spots below eye, sometimes forming bars, sometimes a stripe; a large oblique spot above pectorals, continued on opercle; a black spot at base of pectoral; dorsals and caudal barred, anal uniform dusky, ventrals and pectorals black in male, white in female. D. VI-11; A. 10 to 12. Scales 25 to 30.

BOLEOSOMA, 2544.

*mm.* Profile moderately decurved; eye longer than snout,  $3\frac{1}{2}$  in head. Color yellowish, oblong dark blotches on middle of sides; dorsal and caudal barred. Head 4; depth 6. D. VI-12; A. 10.

FASCIATUS, 2545.

*II.* Highest rays of second dorsal as long as head, the last reaching base of caudal. Body elongate, the back not arched; depth 6 in length; head 4, not compressed, the cheeks tumid. Profile abruptly decurved, the snout  $3\frac{1}{2}$  in head. Mouth large, nearly horizontal, the maxillary reaching posterior edge of eye in males, middle of eye in females. Teeth in narrow bands in each jaw, the outer somewhat enlarged, the outer in some (males?) much enlarged above and recurved, the enlarged teeth fixed, the others movable. Scales large, ctenoid, reduced anteriorly; belly naked. Dorsal spines little filamentous, the longest about equal to head; caudal  $2\frac{1}{2}$  to 3 in body. Males dark olive, with 4 oblong dark blotches along middle of sides; a dark caudal spot; a black blotch larger than eye on each side of shoulder; dorsal spotted;

caudal reddish above, dusky below; females with 5 oblong dark blotches on sides, the last on base of caudal; from each of the middle blotches a V-shaped bar runs to the back; a black shoulder blotch; a dark bar from eye to mouth; ventrals pale, with 2 dark streaks. D. VI-11; A. 12. Scales 30 (27 to 33) at least in males.

ENCEOMUS, 2546.

ll. Dorsal spines high, the highest reaching past middle of second dorsal. Nape scaly. Body elongate, moderately compressed.

n. Depth 5 to 6 in length. Profile little decurved, skull flattish behind, much broader than in *Gobius boleosoma*, with an evident median ridge; mouth very oblique, much larger than in *G. boleosoma*; lower jaw thin and flat. Back slightly arched. Body a little deeper and rather less compressed than in *G. enceomus*, the depth 5 to 6 in length. Head 4. Anterior profile moderately decurved. Eye  $3\frac{1}{2}$  in head. Mouth large, oblique; maxillary reaching to below pupil in both sexes. Teeth above uniserial, some of them enlarged and recurved; lower teeth in a narrow band; males sometimes with the hindmost of the outer series a strong, exerted, recurved canine; belly naked. Longest dorsal spine  $\frac{3}{4}$  head in females, elevated in males; soft dorsal elevated in males; caudal  $3\frac{1}{2}$  in body. Color light greenish; sides of male with 5 or 6 narrow, straight, rather sharply defined whitish or yellowish cross bars, regularly placed; 4 dark bars, 3 below eye and 1 on opercle; a small dark spot behind and above opercle; vertical fins barred; female with a row of irregular dark spots connected by a dusky streak, and with the pale cross bars obsolete. D. VI-12; A. 13. Scales 27.

STIGMATICUS, 2547.

nn. Depth  $4\frac{1}{2}$  in length; the profile very obtuse anteriorly; mouth nearly horizontal, the maxillary extending beyond pupil,  $2\frac{3}{4}$  in head. Teeth strong, uniserial, 4 shortish canines in lower jaw behind the other teeth; upper teeth largest. Some of the dorsal spines filamentous, reaching (in male) past middle of second dorsal; caudal  $\frac{1}{2}$  longer than head; scales large, ctenoid, those on nape and belly much reduced in size. Dark olive, with 4 or 5 irregular, confluent, blackish cross bands, besides irregular, dark blotches; head marked with darker; fins mostly dusky; caudal dark blue with 2 red longitudinal stripes. D. VI-11; A. 16. Scales 27.

LYRICUS, 2548.

nnn. Depth 4 in length; mouth nearly horizontal; teeth short and thick, uniserial; yellowish, much mottled and blotched. D. VI-11; A. 11. Scales 30.

GARMANI, 2549.

dd. Color in life, cherry red, with many bluish cross bars; body stout; depth  $4\frac{1}{2}$  in length. D. VI-11; A. 9. Scales large. ZEBRA, 2550.  
 cc. Scales moderate or small, 40 to 90.

o. Soft dorsal and anal short, each of 10 to 14 rays; body more or less elongate.

EUCTENOGOBIOUS (ev, well; κρείς, comb; *Gobius*):

p. Caudal rounded, not much longer than head.

q. Scales 40; dorsal with 9 soft rays only; anal with 9; depth  $6\frac{1}{2}$  in total length; head broad, flattish; snout short, decurved; eye  $4\frac{1}{2}$  in head,  $1\frac{1}{2}$  in interorbital area, longer than snout; maxillary extending to below middle of eye. Some of the dorsal spines produced in filaments, the third  $1\frac{1}{2}$  times depth of body; caudal short, rounded. Two rows of ill-defined blotches on upper half of body; 2 rows of brownish spots on second dorsal, the upper strongly marked

POEYI, 2551.

qq. Scales 50; dorsal and anal with 10 soft rays each; profile very oblique. Color dark brown. BADIUS, 2552.

GOBIONELLUS\* (diminutive of *Gobius*):

pp. Caudal lanceolate, much longer than head; lower jaw thin; usually a green spot on roof of mouth in life.

r. Body rather deep, the depth about 5 in length.

s. Teeth minute, seen with a lens only. Dorsal spines filamentous; scales much reduced below. D. VI-13; A. 14. Scales 62. MICRODON, 2553.

ss. Teeth well developed.

t. Scales rather large, 39 to 42; body moderately elongate, compressed; depth  $5\frac{1}{2}$ ; head 4. Head not compressed, the cheeks tumid, the snout short, abruptly decurved; mouth large, little oblique, the jaws equal, the maxillary  $2\frac{1}{2}$  in head, reaching to below pupil; eye 5 in head; teeth above large, unequal, uniserial, some of them fixed, those below small, in a band. Scales anteriorly cycloid, becoming larger posteriorly, and ctenoid; dorsal spines scarcely filamentous, none of them as high as body; caudal  $2\frac{1}{2}$  in body. Light olive, with dark olive blotches; body and head with many conspicuous round spots of cream color, each surrounded by a dusky ring, these most distinct on the head, all smaller than pupil; snout with dusky streaks; dorsals and caudal sharply barred; anal and ventrals dusky (in male); a small round spot at base of caudal. D. VI-11; A. 11.

SMARAGDUS, 2554.

tt. Scales comparatively small (53). Body elongate, compressed behind; head a little compressed,  $3\frac{1}{2}$  in length; depth 5; eye  $3\frac{1}{2}$  in head, shorter than the rounded snout; maxillary reaching to below middle of eye; teeth small, the outer a little enlarged; dorsal spines all shorter than head, not filamentous. Nape scaly, its scales much reduced in size; scales ctenoid. Two violet stripes from eye to mouth; 8 or 9 violet bars on sides; 3 or 4 bars on caudal; second dorsal spotted. D. VI-12; A. 11 or 12.

STRIGATUS, 2555.

rr. Body elongate, the depth  $6\frac{1}{2}$  to 9 in length; head  $4\frac{1}{2}$ ; teeth well developed; caudal very long.

\* *Gobionellus* is probably generically distinct from *Gobius* and *Ctenogobius*, but at present we do not know how to limit it, and therefore we are unable to define it.

- u. Scales 55 to 60; eye longer than snout, 4 in head; mouth slightly oblique, the jaws equal, the maxillary not reaching center of eye; teeth in a narrow band, the outer much enlarged and separated from the others by a narrow interspace. Second dorsal spine not equal to depth of body. Caudal  $3\frac{1}{2}$  in body. Scales on nape and axil very small, those on posterior part of body much larger. Light olive green; a series of brown spots along middle of tail; sides of head with dusky blotches, vertical fins dotted with black. D. VI-13; A. 14. Scales 58-20. SAGITTULA, 2556.
- uu. Scales very small (60 to 90); caudal more than twice as long as head in adult. Body compressed, extremely elongate, the depth 6 to 9 in length; head higher than wide, short, compressed,  $4\frac{1}{2}$  to 5 in length; mouth wide, oblique; maxillary in adult reaching to below posterior border of eye. Lower jaw very thin and flat; teeth in both jaws small, subequal, those in the upper jaw in a single series, those of the lower in a narrow band; outer teeth somewhat movable. Scales anteriorly small, cycloid, embedded, those behind larger and ctenoid; a few scales on upper anterior corner of opercle; dorsal fins high, some of the spines filamentous, longer than head. Caudal very long filamentous, 2 to  $2\frac{1}{2}$  in body. Light olive; fins dusky in male; a round, black spot on side, a little larger than eye, below spinous dorsal; first dorsal spine with 2 or 3 black spots; a small dusky spot at base of caudal; emerald spot on tongue conspicuous, fading in spirits. D. VI-14; A. 14 or 15.
- v. Head  $5\frac{1}{2}$  to 6 in length; scales 60 to 70; patch of scales on opercle obsolete. HASTATUS, 2557.
- vv. Head 7 to 8 in length; scales about 90; patch of scales on opercle well developed.

OCEANICUS, 2558.

LYTHRYPNUS (*λύθρον*, gore; *ύπνός*, slumberer; a red sleeper):

- oo. Soft dorsal and anal very long; D. VI-17; A. 14. Body short, compressed; mouth very oblique; jaws with distant canine-like teeth. Coral red, with bluish crossbands and markings. Scales 40. DALLI, 2559.

## Subgenus GOBIUS.

## 2536. GOBIUS SOPORATOR,\* Cuvier &amp; Valenciennes.

(SLEEPER; MAPO; CAIMAN.)

Head  $3\frac{1}{10}$  to  $3\frac{3}{8}$ ; depth  $4\frac{1}{2}$  to  $5\frac{1}{2}$ ; eye 4 to 5. D. VI-I, 9 or 10; A. I, 7 to 9; scales 35 to 41—13 to 15. Vert. 11 + 16. Body robust, compressed

\*The specimens examined are from Panama, Barbados, Parí, Itapua, Cuba, Galapagos, Sambara, Bahia, Orange Key, Bahamas, Pernambuco, St. Thomas, Tortugas, Florida Keys, Martinique, Sao Matheas, Curuca, Rio de Janeiro, Rio Doce. "The color variations among examples of this species are very great, specimens from one locality varying from plain sand color, or gray, to greenish black; some dark brown specimens have light bars across the back; in others the scales have light centers forming horizontal series of light lines; sometimes there are light spots on sides of head and cheek; some specimens are conspicuously marbled with light and dark brown, and white spots occur in the centers of some of the scales on specimens of any ground color, these white spots being brighter on some of the scales than on the others, forming interrupted longitudinal lines. If any value could be placed upon the coloration, almost every specimen would be a distinct species. The color variation is irrespective of locality, some localities having all the above-described variations. The types of Poey's *mapo*, *lacertus*, and *brunneus* prove to be color varieties of *Gobius soporator*." (Eigenmann.)

teriorly; head broad, low, rounded in profile; mouth large, little oblique; lips thick; teeth on upper jaw in a broad band, those of outer series enlarged, the inner ones minute; teeth on lower jaw in a broad band, the outer row enlarged, but not quite as large as the outer series on upper jaw. Anterior half of trunk scaled, head naked; scales large, strongly ctenoid, smaller on nape and belly. Dorsal spines short, not filamentous; upper rays of pectoral fin silk-like, short, and very slender and flexible, free for nearly their whole length; caudal short. Skull posteriorly much as in *Lophogobius cyprinoides*, but the median crest reduced to a slight ridge. Lateral crests very high and closely approximated, rising obliquely outward; the inner crests meeting behind eye, the outer ones forming a very high border about the orbit. Interorbital very narrow and deep, with a median ridge. Coloration that of the rocks, usually granite gray or olivaceous, light or dark, varying from sand color to greenish black, everywhere mottled and marbled with darker and paler, often with brassy or greenish; a faint dusky spot behind eye; coloration varying indefinitely with the surroundings; pectorals, dorsals, and caudal generally mottled; anal and ventrals usually plain. Length 3 to 6 inches. Specimens from Pensacola show the following characters: Head  $3\frac{1}{2}$  (4 in total); depth 4 (5). D. VI-10; A. I, 9; scales 30 to 38; 12 rows of scales from first dorsal downward and backward to anal. Scales on nape extremely small, those on sides firm, ctenoid; first dorsal with an oblique median shade of blackish, the base in front and the distal part light orange; second dorsal dusky at the base, with some spots, its margin light orange; caudal reddish, with dusky cross lines or spots; anal and ventral dusky, yellowish at base in the female; pectoral olivaceous, yellowish at base, reddish at tip, 2 dark spots on base of pectorals. Form robust. Head rather blunt and heavy, the snout less abruptly decurved than in *G. lyricus*. Mouth moderate, the jaws equal, the maxillary reaching about to front of pupil,  $2\frac{3}{8}$  in head. Teeth in moderate bands, the outer series somewhat enlarged. Cheeks full, tumid. Eyes moderate, placed rather high, much broader than the interorbital space. Dorsal spines slender, the first longer than the other, but not filamentous,  $1\frac{3}{8}$  in head; caudal rounded,  $1\frac{1}{2}$  in head; upper rays of pectorals silk-like, the fin somewhat longer than the ventral,  $1\frac{1}{2}$  in head. Color in life, very deep olive green, the back and sides obscurely barred and much marbled with different shades of olive green; cheeks with dark markings, forming reticulations around pale spots; whole under part of head blackish in the males, yellowish in the females. Tropical seas; universally distributed and almost everywhere common, lurking among stones or on sand in shallow water, or in rock pools, moving very quickly when disturbed, north on our coast to Carolina and Gulf of California. The commonest of all shore fishes in tropical America. Among our species it seems to be the one most nearly related to the European *Gobius niger*, and it may, therefore, be held to represent the subgenus *Gobius*, if our other species be placed in different subgenera. Perhaps all the others will ultimately be removed from *Gobius*.

*Gobius saporator*, CUVIER & VALENCIENNES, Hist. Nat. Poiss., XII, 56, 1837, Martinique; GÜNTHER, Cat. Fish., III, 26, 549, 1861; POEY, Enumeratio, 124, 1876; JORDAN & GILBERT, Synopsis, 634, 1883.



- Gobius lineatus*, JENYNS, Zool. Voy. Beagle, 95, pl. 19, fig. 2, 1842, Galapagos Archipelago. (Coll. Charles Darwin.)
- Gobius catulus*, GIRARD, Proc. Ac. Nat. Sci. Phila. 1858, 169, St. Joseph Island, Texas; GIRARD, U. S. and Mex. Bound. Survey, Zool., 26, pl. 12, figs. 9 and 10, 1859; JORDAN & EIGENMANN, *l.c.*, 493.
- Gobius nipo*, POEY, Memorias, II, 277, 1861, Cuba; POEY, Synopsis, 392, 1868.
- Gobius lacertus*, POEY, Memorias, II, 278, 1861, Cuba; POEY, Synopsis, 392, 1868; POEY, Enumeratio, 125, 1876.
- Gobius andrei*, SAUVAGE, Bull. Soc. Philom., Ser. 7, IV, 44, 1880, Rio Guayas, Ecuador. (Coll. André.)
- Gobius carolinensis*, GILL, Proc. Ac. Nat. Sci. Phila. 1863, 268, Charleston, South Carolina; JORDAN & GILBERT, Synopsis, 634, 1883.
- Gobius brunneus*, POEY, Synopsis, 393, 1868, Havana; name preoccupied.
- Evorthodus catulus*, JORDAN & GILBERT, Synopsis, 632, 1883.

According to Dr. Eigenmann, *Gobius albopunctatus* of the Western Pacific can not be separated from *Gobius saporator*. In this case several other synonyms should be added.

Subgenus CTENOGOBIUS, Gill.

2537. GOBIUS NICHOLSII, Bean.

Head  $3\frac{1}{2}$  to  $3\frac{3}{4}$ ; depth 5 to  $5\frac{1}{2}$ . D. VI-I, 12 to 14; A. I, 11; scales 25 or 26-10. Body compressed; width of head about twice in its length. Mouth oblique, the maxillary reaching to front of pupil,  $2\frac{1}{2}$  to 3 in head; lower jaw very slightly produced. Teeth present on both jaws; the outer series of long, conical teeth, placed at a considerable distance apart; the enlarged teeth on lower jaw not extending on the sides; the inner series of a band of small teeth. Interorbital space very narrow, equaling pupil. Snout 4 to  $4\frac{1}{2}$  in head. Eyes large, placed high, 3 to  $3\frac{1}{2}$  in head. Caudal peduncle  $2\frac{3}{4}$  to  $3\frac{1}{2}$  in head. Scales large, caducous, ctenoid; lacking on head, nape, and fins. Dorsal spines slender, flexible; base of first dorsal  $1\frac{3}{4}$  to 2 in head; soft dorsal and anal similar; base of anal  $1\frac{1}{2}$  to  $1\frac{3}{4}$  in base of soft dorsal, and  $1\frac{1}{2}$  to  $1\frac{3}{4}$  in head; ventrals  $1\frac{1}{2}$  to  $1\frac{3}{4}$  in head, inserted below or slightly behind origin of pectorals; pectorals reaching a considerable distance beyond ventrals  $1\frac{1}{2}$  to  $1\frac{3}{4}$  in head; caudal rounded, not equaling head. Color in spirits, light yellowish brown, with traces of darker, lighter below; ventrals usually dusky; spinous dorsal narrowly margined with black; second dorsal finely checkered in adult; other fins yellowish, not distinctly marked. Length 2 to  $3\frac{1}{2}$  inches. Coast of British Columbia; not rare. Here described from 5 specimens from Albatross Station 2944, numbered 66 in the L. S. Jr. Univ. Museum. (Named for Capt. Henry E. Nichols, U. S. N., its discoverer.)

*Gobius nicholsii*, BEAN, Proc. U. S. Nat. Mus. 1881, 469, Departure Bay, British Columbia; JORDAN & GILBERT, Synopsis, 946, 1883; JORDAN & EIGENMANN, *l.c.*, 494.

2538. GOBIUS EIGENMANNI, Garman.

D. VII-12; A. 13; P. 19; scales 27-7. Body rather stout, body cavity more than  $\frac{1}{2}$  the length from snout to base of caudal. Head  $\frac{2}{3}$  of the total length or  $\frac{2}{3}$  of the distance to the caudal base, blunt and rounded anteriorly, very narrow between the eyes, slightly compressed. Eyes large,  $\frac{1}{2}$  of the

hea  
the  
the  
long  
of t  
to l  
as h  
2 row  
near  
halv  
strea  
the la  
tingu  
West,  
guish  
Ameri  
*Gobius*  
of T

Head  
robust,  
project  
series,  
upper r  
and ped  
a faint  
formed  
the seco  
similar l  
along th  
a straight  
Length  
Washing  
taken th  
(*glaucus*,

\* Dr. Eig  
*Gobius gla*  
scales in a  
jaws equal,  
outer row o  
and equalin  
lar to soft d  
ing past ve  
in spirits, 1  
specimens ne  
middle of st  
ward from  
allel to it ac  
triangular d  
cle and sno  
differing fro  
1 $\frac{1}{2}$ , 1 $\frac{1}{2}$ , 1 $\frac{1}{2}$ , 1 $\frac{1}{2}$

head, very close together. Snout short, little more than  $\frac{1}{2}$  as long as the eye. Mouth wide; maxillary reaching a vertical from the middle of the eye, moderately oblique. First dorsal higher, anterior 3 rays prolonged in the filaments, third ray longest and reaching to the eighth ray of the second dorsal; origin of anal fin midway from edge of preopercle to base of caudal; pectorals nearly as long as the head; caudal as long as head, pointed. Scales large, thin, deciduous, 27 in a longitudinal series, 2 rows above the lateral line. Yellowish, with a few punctulations of black near the bases of the caudal rays, with a light-edged black spot on the outer halves of the fourth to the sixth rays of the first dorsal and with a black streak around the mouth immediately above the maxillary. The long body, the large eye, the dorsal spot, and the streak above the mouth serve to distinguish this species from its nearest allies of the same locality. Off Key West, in 60 fathoms. ("The specific name is given in honor of the distinguished ichthyologists who have added so much to our knowledge of the American Gobiidae, C. H. and R. S. Eigenmann.")

*Gobius eigenmanni*, GARMAN, Bull. Lab. Nat. Sci. State Univ. Iowa, vol. IV, No. 1, 88, 1896; off Key West in 60 fathoms. (Coll. Iowa Univ. Bahama Expedition.)

2539. GOBIUS GLAUCOFRENUM (Gill).

Head  $3\frac{1}{2}$ ; depth  $4\frac{1}{2}$ . D. VI-10; A. I, 9; P. 18; lateral transverse 7. Body robust, compressed; head naked; mouth oblique, the lower jaw slightly projecting, the maxillaries extending to below pupil; teeth long, in many series, the outer curved; scales ctenoid, large. Pectoral fin with the upper rays little branched, not silk-like; cheeks scarcely tumid; caudal and pectoral longer than ventrals, about as long as head. Tawny, with a faint blue spot in the center of each scale, and with 6 spots, each formed by aggregation of dark dots, on the ridge of the back between the second dorsal spine and the axil of the soft dorsal fin; another row of similar but fainter spots runs from the scapular region, and a third row along the middle of the sides; head tawny, with dark spots and blue dots; a straight blue line across the cheek; dorsal fins with faint blue spots. Length  $1\frac{1}{2}$  inches. Florida Keys; said to have come from the coast of Washington, but this is probably an error, as the species has not since been taken there, while 1 apparently identical has been taken at Tortugas.\* (*glaucus*, glaucous; *frænum*, bridle.)

\* Dr. Eigenmann thus describes the specimens from the Tortugas examined by him: *Gobius glaucofrenum* (Gill). Head  $3\frac{1}{2}$  ( $4\frac{1}{2}$  in total); depth  $4\frac{1}{2}$  ( $5\frac{1}{2}$ ). D. VI-10; A. 10; scales in a median series, 23, in a transverse series, 8; eye as long as snout,  $3\frac{1}{2}$  in head, jaws equal, maxillary barely reaching pupil. Teeth in bands in both jaws, those of the outer row of lower jaw enlarged. Dorsal spines scarcely filamentous, the third highest and equalling depth of body. Posterior dorsal rays highest, as high as spines; anal similar to soft dorsal; pectoral long and narrow, longer than head,  $3\frac{1}{2}$  in body; ventral reaching past vent; scales large, thin, finely toothed, reduced on breast; nape naked. Color in spirits, light yellowish brown; a light spot on each scale, the spots especially conspicuous near shoulder; 6 dark spots on middle of back; fainter but similar spots along middle of sides; a conspicuous dark spot above opercle; a wavy light line extending forward from it through lower rim of eye to snout; a straight pale-blue bar extending parallel to it across preopercle and cheek to corner of mouth; a narrow faint bar below it; a triangular dark spot at corner of mouth; cheeks and preopercle purplish chocolate; opercle and snout plain yellowish; 2 brown spots at base of caudal; the smaller specimens differing from this in having the markings more distinct. Length of 4 specimens examined, 1 $\frac{1}{2}$ , 1 $\frac{1}{2}$ , 1 $\frac{1}{2}$ , 1 $\frac{1}{2}$  inches. (Eigenmann.)

*Coryphopterus glaucofrœnum*, GILL, Proc. Ac. Nat. Sci. Phila. 1863, 263, Coast of Washington (evidently an error).

*Gobius glaucofrœnum*, JORDAN & GILBERT, Synops. 635, 1883; JORDAN & EIGENMANN, l. c., 494; EIGENMANN & EIGENMANN, Proc. Cal. Ac. Sci. 1888, 59.

2540. *GOBIUS MANGLICOLA*, Jordan & Starks.

Head  $4\frac{1}{2}$ ; depth  $5\frac{1}{2}$ . D. VI-12; A. 12; scales about 35, not to be exactly counted; caudal lanceolate,  $2\frac{3}{8}$  in body; pectoral about equal to head; dorsal spine slender, not filamentous,  $1\frac{1}{2}$  in head; eyes large, close together, the range partly vertical, the narrow interorbital deeply furrowed; no flaps on shoulder girdle; scales moderate, ctenoid anteriorly, becoming smooth behind; median keel on head slight; head naked. Body long, compressed, the head depressed, the cheeks tumid; snout bluntly truncate; mouth large, the maxillary reaching the middle of eye, not produced backward, truncated behind, somewhat oblique, the lower jaw a little the longer; lower jaw flat; teeth strong, the outer in both jaws enlarged; cranium without median crest, abruptly widened behind eyes. Color light olive, mottled with darker; 6 oblong blotches of blackish on sides as in *Gobius boleosoma*, the last at base of caudal; dorsals and caudal finely checkered and barred with dark brownish orange and blackish; anal mottled; a dark shoulder spot; a dark bar before eye and 1 below eye; ventrals dusky, the edge pale. One specimen,  $1\frac{1}{2}$  inches long. Mazatlan; found in the mud of the Astillero among the roots of mangrove bushes (*Rhizophora mangle*), (whence the name *mangle*; *colo*, I inhabit).

*Gobius manglicola*, JORDAN & STARKS, Proc. Cal. Ac. Sci. 1895, 496, Mazatlan. (Coll. Hopkins Expedition to Mazatlan. Type, 3095, L. S. Jr. Univ. Mus.)

2541. *GOBIUS STIGMATURUS*, Goode & Bean.

Head 4; depth  $4\frac{1}{2}$ ; eye 4, about equal to snout. D. VI-12; A. 11 or 12; scales 33. Body subfusiform, little compressed; head blunt, the profile rounded. Mouth small, horizontal, the lower jaw included; maxillary 3 in head, reaching eye. Teeth small, in bands in both jaws, the outer enlarged, those of the upper jaw very slender. Region from nape to dorsal entirely scaled, the scales large, ctenoid, those on nape and belly little reduced. Dorsal spines short, the longest shorter than the head; caudal fin scarcely pointed, about as long as head. Grayish white, middle of sides with 4 or 5 dark blotches, from each of which a dark bar extends downward and forward; a large black blotch above pectoral, obsolete in the female; a small black spot at base of caudal, and a dark mark below the eye; vertical fins barred. Two specimens taken in a shallow bay at Key West are thus described: Very pale olive, everywhere freckled and spotted; lower part of sides silvery, crossed by faint and narrow cross streaks of light brown; sides with about 5 faint dark blotches; a dark blotch below eye and 1 on opercle; a round black spot at base of caudal; bars on verticle fins light olive. Numerous other specimens are less freckled in coloration, and have a more diffuse caudal spot as well as a vague dark spot at the shoulder. The dusky marks on the sides are larger.

We find no other differences, and refer all of them to *G. stigmaturus*. The relations of *G. boleosoma*, *G. stigmaturus*, and *G. encromus* are certainly very intimate. Florida Keys, not very common, our specimens from Key West. (*στρυμα*, spot; *ὄψα*, tail.)

*Gobius stigmaturus*, GOODE & BEAN, Proc. U. S. Nat. Mus. 1882, 418, no type locality given, but specimens probably from Florida Keys; JORDAN & GILBERT, Synopsis, 940, 1883; JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1884, 140; JORDAN & EIGENMANN, l. c., 495.

2542. GOBIUS QUADRIPORUS, Cuvier & Valenciennes.

D. VI-I, 9; A. I, 9; scales as in *Gobius caninus*. The 2 pores on the vertical arm of preopercle very open; 2 smaller ones above them; teeth of outer series small; 2 small canines on each side of lower jaw; dorsal spines not prolonged as filaments. Color yellowish, with lighter lines which follow the rows of scales; brown spots on dorsal; 2 lines on cheek. Surinam. (Cuvier & Valenciennes.) Not seen by us. (*quatuor*, four; *porus*, pore.)

*Gobius quadriporus*, CUVIER & VALENCIENNES, Hist. Nat. Poiss., XII, 87, 1837, Surinam; EIGENMANN & EIGENMANN, Proc. Cal. Ac. Sci. 1888, 61.

2543. GOBIUS SHUFELDTI, Jordan & Eigenmann.

Head  $3\frac{3}{4}$  to  $3\frac{1}{2}$ ; depth  $5\frac{1}{4}$ ; eye 4; snout  $3\frac{1}{4}$  to  $3\frac{1}{2}$ . D. VI-12; A. 13; scales 33 to 35. Body moderately elongate, subfusiform; head less blunt than in *Gobius boleosoma*, the anterior profile gently deurved; mouth large, slightly oblique; maxillary extending to front of pupil,  $2\frac{1}{4}$  in head. Teeth small, slender, and curved, in moderate bands; scales covering anterior half of trunk; head and breast naked; scales moderate, ctenoid, those in front much reduced. Longest dorsal spine  $1\frac{1}{2}$  in head; caudal fin as long as head, somewhat pointed. Olivaceous, mottled with gray; about 5 round dark blotches along middle of side, the last at base of caudal; no dark spot on side of nape; some dark marks on head; vertical fins barred. Gulf coast of the United States, known as yet only from fresh waters about New Orleans. (Named for Dr. Robert Wilson Shufeldt, U. S. A., who collected the types.)

*Gobius shufeldti*, JORDAN & EIGENMANN, Proc. U. S. Nat. Mus. 1886, 495, New Orleans. (Type, No. 35202.)

2544. GOBIUS BOLEOSOMA, Jordan & Gilbert.

Head 4 (5 in total); depth  $4\frac{1}{2}$  to  $5\frac{1}{2}$ . D. VI-12; A. I, 10 to 12; scales 25 to 30. Body slender, subfusiform, little compressed; head moderate, not very blunt, the anterior profile somewhat evenly deurved, the snout not very short, scarcely shorter than the large eye; mouth not very large, horizontal, the lower jaw included, the maxillary extending slightly beyond front of pupil, its length about 3 in head; teeth small, slender, in narrow bands, those of the outer series longer than the others; eyes placed high, about 4 in head; interorbital space not wider than pupil; scales moderate, ctenoid, those on nape and belly not much reduced in size; gill opening not continued forward above opercle; first dorsal with

the spines slender but rather firm, none of them filamentous, the longest about  $\frac{3}{4}$  head; second dorsal and anal rather large; caudal long, pointed, slightly longer than head; pectorals large, slightly longer than head, none of the upper rays silk-like; ventrals slightly shorter than head, inserted below axil of pectorals; skull rounded behind, no ridges nor crests; crossts at side minute; interorbital very narrow. Color in life: Male, deep olive green, mottled with darker; middle of side with 4 or 5 vague darker blotches; a jet-black spot above gill opening, on side of back; head mottled, dusky below; usually a dark bar below eye; dorsals tipped with bright yellowish, each crossed by numerous narrow, somewhat oblique, interrupted bars or series of spots, these being of a rich reddish brown color; caudal barred with black, its upper edge tinged with orange; anal nearly plain, with a slight orange tinge; ventrals bluish black, their edges whitish. Female, paler and duller in color, more mottled, the black spot above gill opening obsolete or nearly so; a dark spot at base of caudal; upper fins barred, as in the male; lower fins mostly pale, tinged with orange. Many specimens of this species, the largest about 2 inches in length, were obtained in the Laguna Grande at Pensacola. It lurks in sea wrack on muddy bottoms in very shallow water (6 to 12 inches). In form, size, coloration, and movements this little fish bears a remarkable resemblance to the percid, *Boleosoma olmstedii*. Gulf of Mexico, Pensacola to Key West; common in shallow sandy bays, lurking in sea wrack at the depth of a foot (whence the name *βολις*, dart; *σῶμα*, body).

*Gobius boleosoma*, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1882, 295, Laguna Grande, Pensacola; *ibid.*, Synopsis, 946, 1883; JORDAN & EOENMANN, *l. c.*, 495.

#### 2545. GOBIUS FASCIATUS (Gill).

Head 4 in length ( $4\frac{1}{2}$  in total); depth 6 (7). D. VI-12; A. 10; scales 30-7. Body slender, elongate; head somewhat pointed; profile rounded, not as much as in *Gobius garmani* and *Gobius boleosoma*; eye large, slightly longer than snout,  $3\frac{1}{2}$  in head; interorbital area scarcely wider than pupil; mouth slightly oblique, maxillary extending to below anterior margin of pupil,  $3\frac{1}{2}$  in head; lower jaw thin and flat; teeth strong, recurved, in a band in each jaw, the teeth of the outer series of the upper jaw enlarged, several times as large as those of the inner series. Scales finely ctenoid (fallen off anteriorly in specimen examined); antedorsal region and breast naked. Dorsal spines slender, filamentous near tip, not reaching second dorsal,  $1\frac{1}{2}$  in length of head; second dorsal of moderate height; caudal (tips broken) about 5 in length,  $1\frac{1}{2}$  in length of head; ventral not reaching vent,  $1\frac{1}{2}$  in head; pectorals pointed, equaling the head in length. Color yellowish, marbled with darker above; 4 oblong dark blotches along middle of sides; a darker spot at base of caudal; narrow dark stripes across nape; a faint dark stripe along upper margin of opercle, through lower margin of eye to snout; another extending from angle of mouth to edge of preopercle, then extending down along the margin of the preopercle and ending in a dark blotch on the lower part of the cheek; a dark spot on opercle; first dorsal with 2 curved bars; caudal with 3 rather broad dark bars; anal

dusky; connecting membrane of ventral white, its first rays blackish, outer rays yellowish; lower parts yellowish. West Indies; not seen by us. This description by Eigenmann, from a specimen  $1\frac{1}{4}$  inches long, No. 13231, M. C. Z., collected in Hayti by Dr. Weinland. (*fasciatus*, branded.)

*Ctenogobius fasciatus*, GILL, Fishes Trinidad, 378, 1858, Trinidad.

*Gobius fasciatus*, GÜNTHER, Cat., III, 34, 1861; JORDAN & EIGENMANN, Proc. U. S. Nat. Mus. 1886, 495; EIGENMANN & EIGENMANN, Proc. Cal. Ac. Sci. 1888, 62.

2546. GOBIUS ENCEOMUS, Jordan & Gilbert.

Head 4; depth 6; snout  $3\frac{1}{2}$ . D. VI-11; A. 12; scales 27 to 33. Body very elongate, much tapering backward; head compressed, the cheeks high and vertical; snout very short, compressed, obtusely rounded vertically. Mouth nearly horizontal, low, large, the maxillary 2 in head, nearly reaching vertical from posterior margin of orbit. Teeth in very narrow bands in both jaws, those of the outer series in the upper jaw much enlarged and recurved in some specimens; eyes inserted high, the interorbital space very narrow, about as wide as pupil; diameter of orbit much greater than snout, nearly  $\frac{1}{2}$  of head. Gill opening  $2\frac{1}{2}$  in head, the isthmus wide. Dorsals contiguous, the membrane of spinous dorsal reaching nearly to base of soft dorsal; dorsal spines high, of nearly uniform length, the last reaching well beyond origin of soft dorsal when depressed; the longest spine about  $\frac{1}{2}$  length of head; soft dorsal and anal long and high, the posterior rays of both fins reaching at least to base of caudal when depressed; caudal lanceolate, the middle rays produced,  $2\frac{3}{4}$  in body; ventrals reaching vent, somewhat longer than pectorals, which about equal length of head; ventral sheath well developed, its length  $\frac{2}{3}$  that of fin. Body wholly covered with large, strongly ctenoid scales, which are much reduced in size anteriorly; head, antedorsal region, and breast naked. In female specimens the mouth is evidently smaller, and the caudal less elongate. Colors in life: Male, light olivaceous, mottled above with darker olive brown; a series of about 4 obscure oblong dark blotches along middle of sides; a dark spot at base of caudal; each side of nape with an intense blue-black spot larger than eye; an obscure dusky streak from eye forward to mouth; a small dusky spot sometimes present on upper portion of base of pectorals; both dorsals translucent, with a series of bright reddish-brown spots as large as pupil; upper lobe of caudal light reddish, the lower lobe blue black; anal and ventrals dusky bluish, pectorals slightly dusky, with a narrow, bright pink border behind. Female, without bright markings; body light olive, with 5 oblong dark blotches on sides, the last on base of caudal; from each of the 3 middle blotches a V-shaped bar runs to the back (these visible also in males); back somewhat mottled with dusky; a black blotch on scapula; a small one on opercle; a dark bar from eye forward to mouth. Vertical fins with dusky streaks, these appearing on caudal in the form of cross bars; ventrals light, with 2 lengthwise dark streaks; pectorals plain. South Carolina to Key West, in sandy bays; scarce. Length 2 inches. (*ἐγκάτω*, brand; *ῶμος*, shoulder.)

*Gobius encraomus*,\* JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1882, 611, Charleston, South Carolina (Type, No. 29673, 3 specimens. Coll. C. H. Gilbert); JORDAN & GILBERT, Synopsis, 945, 1883; JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1884, 142; JORDAN & EIGENMANN, l. c., 496.

## 2547. GOBIUS STIGMATICUS (Poey).

Head 4; depth 5 to 6; eye  $3\frac{1}{2}$ . D. VI-12; A. 12 or 13; scales 27. Body a little deeper and less compressed than in *Gobius encraomus*. Anterior profile moderately decurved; back slightly arched; skull flattish behind, much broader than in *G. boleosoma*, with an evident median ridge; mouth oblique, large, lower jaw thin and flat, maxillary reaching to below pupil. Teeth above uniserial, some of them enlarged and recurved; lower teeth in a narrow band, males sometimes with the hindmost of the outer series a strong, exerted, recurved canine (present in Poey's type). Anterior half of body scaled except region between nape and dorsal, which is naked; breast naked. Longest dorsal spine  $\frac{2}{3}$  head, sometimes elongate; caudal  $3\frac{1}{2}$  in body. Light greenish, sides of male with 5 or 6 narrow, straight, whitish or yellowish cross bars, regularly placed; 4 dark bars on head, 3 below the eye and 1 on opercle; a small dark spot behind and above opercle; ventral fins barred; female with a row of irregular dark spots connected by a dusky streak, the pale cross bars obsolete. Coast of North Carolina, Florida Keys, the West Indies, southward to Rio Janeiro; common at Havana. Subject to considerable variation. Brazilian specimens said by Eigenmann to be darker, the bars on cheek conspicuous; third dorsal spine often much elongate, reaching fifth dorsal ray, last soft ray sometimes reaching caudal. (*stigmaticus*, spotty.)

*Smaragdus stigmaticus*, POEY, Memoriae, II, 281, 1861, Cuba.

*Gobionellus stigmaticus*, POEY, Synopsis, 394, 1862. POEY, Enumeratio, 126, 1876; JORDAN & GILBERT, Synopsis, 947, 1883.

*Gobius stigmaticus*, JORDAN, Proc. U. S. Nat. Mus. 1880, 49; JORDAN & EIGENMANN, l. c., 496.

## 2548. GOBIUS LYRICUS, Girard.

Head  $4\frac{1}{2}$ ; depth  $4\frac{1}{2}$ . D. VI-11; A. I, 10; scales 27. Body rather elongate, moderately compressed; head rather short, the profile very obtuse, descending abruptly from before the front of the eye to the snout; eyes small, placed high, about as long as snout, and about  $4\frac{1}{2}$  in head; mouth nearly horizontal, much below level of eye, the maxillary extending to beyond pupil,  $2\frac{3}{4}$  in head; jaws subequal; teeth strong, in 1 series in each jaw; in the lower jaw about 4 shortish, canine-like teeth behind the other teeth; anterior teeth of lower jaw small, of upper jaw rather large; gill

\* One small specimen, taken with the seine in a shallow bay, at Key West, is described as follows:

Light green, with 5 diffuse spots of darker green on sides, the posterior one most conspicuous; pectorals, both dorsals, and caudal edged above with pale orange; ventrals mostly black, edged with paler; anal dark; a conspicuous dusky shoulder spot; maxillary reaching to below middle of eye; caudal about  $\frac{1}{2}$  longer than head. Lateral line about 30. This little specimen appears to be identical with that described by us from Charleston under the name *Gobius encraomus*. The species is allied to *G. stigmaticus*, but has a much slenderer body. The number of scales in a lateral series is less than 37, the number originally stated by us. There are about 33 in this specimen. (Jordan & Gilbert.)

opening  
spines fi  
dorsal, w  
and poin  
reaching  
somewha  
scales lau  
size; hea  
blackish  
marbled  
first dors  
and anal  
bright red  
head and  
Mexico, fr  
(*lyricus*, pe

*Gobius lyricus*  
GIRARD,  
550, 1861;  
*Smaragdus e*  
Coll. Fel  
*Gobius vour*  
probably  
*Eucetengobius*

Head 4 in  
30-7. Body  
decurved. r  
tal; lower j  
lips thin; te  
contiguous;  
the rest, rea  
ing base of  
broad, 5 in l  
slightly red  
brown; a ser

A specimen  
4. The second  
length of head,  
the head,  $2\frac{1}{2}$  in  
with minute da  
which are jet-b  
teriorly, the bas  
with dark point  
caudal dusky,  
edged with whi  
and short, whit  
1 *Gobius vour*  
securely barred  
scales smaller t  
third dorsal t  
(Named for Dr.

opening not continued forward above opercle; first dorsal with 2 or 3 spines filamentous, the longest reaching past the middle of the second dorsal, which is of moderate height and similar to the anal; caudal long and pointed,  $\frac{1}{2}$  longer than the head; pectoral as long as head, about reaching front of anal; upper rays of pectorals not silk-like; ventrals somewhat shorter than head, their insertion below front of pectorals; scales large, rough, those on nape, pectoral region, and belly reduced in size; head naked. Color in life, dark olive, with 4 or 5 irregular confluent blackish cross bands, besides dark blotches and irregular markings; head marbled with darker, the jaws, opercles, and branchiostegals blackish; first dorsal mostly dusky translucent, somewhat barred; second dorsal and anal plain dusky; caudal dark blue, with 2 longitudinal stripes of bright red; pectoral finely barred or reticulated with blackish and pale; head and belly yellowish. Female specimens duller and paler. Gulf of Mexico, from Galveston to Cuba and the Lesser Antilles; rather common. (*lyricus*, pertaining to a lyre, apparently an allusion to the dorsal spines.)

*Gobius lyricus*, GIRARD, Proc. Ac. Nat. Sci. Phila. 1858, 169, Brazos Santiago, Texas; GIRARD, U. S. and Mex. Bound. Surv., 25, pl. 12, figs. 4 and 5, 1859; GÜNTHER, Cat., III, 550, 1861; JORDAN & EIGENMANN, l. c., 496; EIGENMANN & EIGENMANN, l. c., 63.

*Smaragdus costalei*, POEY, Memorias, II, 280, 1861, Havana. (Type, No. 13109, M. C. Z. Coll. Felipe Poey.)

*Gobius wurdemanni*, GIRARD, Proc. Ac. Nat. Sci. Phila. 1858, 169, Brazos Santiago; probably the female; JORDAN & GILBERT, Synopsis, 634.f

*Eutengobius lyricus*, JORDAN & GILBERT, Synopsis, 633, 1883.

#### 2549. GOBIUS GARMANI, Eigenmann & Eigenmann.

Head 4 in length ( $5\frac{1}{2}$  in total); depth 4 ( $5\frac{1}{2}$ ). D. VI-11; A. 11; scales 30-7. Body robust, head short and blunt; profile in front of eye abruptly decurved, rounded much as in *Gobius boleosoma*; mouth inferior, horizontal; lower jaw included; maxillary extending to below pupil,  $2\frac{1}{2}$  in head; lips thin; teeth short and thick, in a single series in each jaw. Dorsals contiguous; dorsal spines filamentous, the second and third longer than the rest, reaching past first third of second dorsal; last dorsal rays reaching base of caudal; pectorals equaling head in length; ventral short and broad, 5 in body; caudal rather long and pointed, 3 in body. Scales large, slightly reduced and cycloid on nape. Color yellowish, marbled with brown; a series of irregular blotches along the sides; a light spot at base

A specimen from St. Kitts is thus described by Eigenmann: "Depth 5 in length; head 4. The second and third dorsal spines extend to base of caudal; dorsal scarcely less than length of head, the last rays reaching past base of caudal; the caudal fin is  $\frac{1}{2}$  longer than the head,  $2\frac{1}{2}$  in body. Color light brown, faintly marked with darker; the first dorsal with minute dark points, the lower fourth of the spines with simple dark spots, above which are jet-black spots ocellated with white; the second dorsal fin dusky, darker posteriorly, the basal portion of the last half of the fin evenly black, the anterior 4 rays marked with dark points similar to the spots on the lower parts of the spines of the first dorsal; caudal dusky, with 2 light bars; anal plain, darker than body; ventral fins blackish, edged with white; pectorals blackish, with many series of white spots on the membrane, and short, white bars at base; branchiostegal membrane black, with a light margin.

*Gobius wurdemanni*, Girard. Appearance of *Gobius lyricus*. Reddish brown, obscurely barred with dusky. Head larger; caudal shorter; ventrals shorter; anal lower; scales smaller than in *G. lyricus*; teeth very slender, much smaller than in *G. lyricus*; third dorsal fine filamentous. D. VI-11; A. 12. Brazos Santiago, Texas. (Girard.) (Named for Dr. Gustav Wurdemann, its collector.)



of caudal, partly or wholly surrounded by a broad ring of dark brown; head slate color, white below; 3 dark bars extending forward and downward from eye to mouth; a triangular spot on opercle; dorsals, caudal, and pectorals finely barred with black; a chocolate bar on base of ventral; anal margined with white; an irregular black bar on shoulder and upper half of pectoral; everywhere more or less blotched with darker, the blotches at times forming numerous bars across the back. Dominica, Fort de France, Martinique, St. Kitts. (Eigermann & Eigenmann.) Not seen by us. Apparently very close to *Gobius lyricus*, if not the same. (Named for its discoverer, Prof. Samuel Garman.)

*Gobius garmani*, EIGERMANN & EIGEMANN, Bull. Cal. Ac. Sci. 1888, 61, Dominica, Fort de France, Martinique, St. Kitts. (Coll. Samuel Garman.)

#### 2550. GOBIUS ZEBRA, Gilbert.

Head 3; depth  $4\frac{1}{2}$ ; eye  $3\frac{1}{2}$  in head. D. VI-11 or 12; A. 9. Body not elongate, the snout short, the mouth oblique, with maxillary reaching below middle of orbit. Mouth small, the maxillary  $2\frac{1}{2}$  in head. Interorbital space very narrow. Teeth in upper jaw in a narrow band or double series, the outer row enlarged and spaced; lower jaw apparently with a single series, similar to the outer row in the upper jaw. Scales cycloid, large, wanting on nape and a narrow strip along base of spinous dorsal. Color cherry red, head and sides with 15 blue cross bars, a little narrower than interspaces, encircling the body posteriorly, lacking for a short distance on belly and under side of head; on upper side of head and nape these bars run obliquely forward and downward, but elsewhere vertical; on middle of each interspace a very narrow blue line, becoming indistinct on lower part of sides; on cheeks the blue bars are connected by narrow cross lines, forming blue reticulations surrounding round spots of the ground color. Length of types  $\frac{1}{2}$  inch. Two specimens from Albatross Station 2989, west coast of Mexico, in 36 fathoms. (*zebra*, *zebra*, from the stripes.)

*Gobius zebra*, GILBERT, Proc. U. S. Nat. Mus. 1890, 73, Albatross Station 2989, west Coast of Mexico.

#### Subgenus EUCTENOGOBIUS, Gill.

#### 2551. GOBIUS POEYI, Steindachner.

Head broad and flattish; depth  $6\frac{1}{2}$  in total length; eye  $4\frac{1}{2}$ ,  $1\frac{1}{2}$  in interorbital width, longer than snout; snout short and decurved. D. VI-9; A. 9; scales 40. Maxillary extending to below middle of eye. Some of the dorsal spines produced and filamentous, the third  $1\frac{1}{2}$  times depth of body; caudal short, rounded. Two rows of ill-defined blotches on upper half of body; dorsals and caudal sharply barred, anal and ventrals dusky (male). A small round dark spot at base of caudal. (Steindachner.) Barbados; not seen by us. (Named for Prof. Felipe Poey.)

*Gobius poeyi*, STEINDACHNER, Ichthyol. Notizen, VI, 44, 1867, Barbados; JORDAN & EIGERMANN, l. c., 497.

Head  
profile  
margin  
5 in to  
heavy  
plumbe  
mouth c

*Euctenogobius*  
*Gobius bo*  
*Gobius ba*

Head  
pressed,  
nearly h  
riorly; m  
 $\frac{1}{2}$  length  
a lens, th  
outer seri  
of still s  
pupil. Ie  
rals. Sea  
posterior  
spinous d  
orbits, bu  
dorsal; h  
First 4 sp  
depressed  
lar. not h  
caudal lar  
equal, reac  
punctulati  
oblique du  
spinous do  
2 inches.  
small;  $\delta\delta$

*Gobius micro*  
Rio Ahor

Head 4;  
39 to 42. I  
the cheeks  
little obliqu

2552. *GOBIUS BADIUS* (Gill).

Head 6 in total; depth 7. D. VI-I, 10; A. I, 10; scales 50-18. Anterior profile very oblique; a line of pores above each eye; 2 on upper ascending margin of preopercle; eye 4 in head; interorbital space  $3\frac{1}{2}$  in eye; caudal 5 in total length; pectoral 6. Color dark bay with a posteriorly straight heavy dot in the center of each scale on back and sides above; head plumbeous, with 2 livid blue bands from eye to upper jaw. (Gill). About mouth of Amazon; not seen by us. (*badius*, bay color, dark red.)

*Euctenogobius badius*, GILL, Ann. Lyc. Nat. Hist. N. Y., VII, 1857, 47, Amazon.

*Gobius bosci*, SAUVAGE, Bull. Soc. Philom. Paris, IV, 44, 1880.

*Gobius badius*, EIGENMANN & EIGENMANN, l. c., 65.

Subgenus *GOBIONELLUS*, Girard.

2553. *GOBIUS MICRODON*, Gilbert.

Head  $4\frac{1}{2}$ ; depth 5. D. VI-13; A. 14; scales 62. Head and body compressed, everywhere deeper than wide. Mouth at lower profile of snout, nearly horizontal, the lower jaw extremely weak, broadly rounded anteriorly; maxillary reaching vertical from hinder margin of pupil, nearly  $\frac{1}{2}$  length of head. Teeth minute scarcely perceptible without the use of a lens, those in upper jaw in a single series. Mandible with a close set outer series of teeth, separated by an interval from an inner narrow band of still smaller teeth. Interorbital space narrow, less than diameter of pupil. Isthmus wide, the gill slits extending little below base of pectorals. Scales minute and cycloid anteriorly and on belly, becoming larger posteriorly; on sides they are everywhere ctenoid behind the middle of spinous dorsal; belly wholly scaled; nape scaled forward nearly to orbits, but with a narrow median naked streak running back to front of dorsal; breast and sides of head naked. Dorsal fins not connected. First 4 spines filamentous, the longest longer than head, reaching when depressed to base of third ray of soft dorsal. Soft dorsal similar, not high, the last rays not extending beyond the base of caudal; caudal lanceolate, much longer than head; pectorals and ventrals about equal, reaching vent. Color nearly uniform light olivo, with minute darker punctulations which sometimes form darker margins to the scales; an oblique dusky streak on opercle; 3 or 4 oblique obscure dark cross bars on spinous dorsal, and 4 or 5 on tail; ventrals with white pigment. Length 2 inches. San Juan Lagoon, west coast of Mexico. (Gilbert.) (*μικρός*, small; *ὄδους*, tooth.)

*Gobius microdon*, GILBERT, Proc. U. S. Nat. Mus. 1891, 554, San Juan Lagoon, north of Rio Ahomè, Mexico. (Coll. Gilbert.)

2554. *GOBIUS SMARAGDUS*, Cuvier & Valenciennes.

(ESMERALDA.)

Head 4; depth  $5\frac{1}{2}$  to  $5\frac{3}{4}$ ; eye 4 to 5. D. VI-11 or 12; A. 11 or 12; scales 39 to 42. Body moderately elongate, compressed; head not compressed; the cheeks tumid; the snout short and abruptly decurved; mouth large, little oblique; lower jaw slightly inferior; maxillary reaching to below

pupil or to posterior margin of orbit,  $2\frac{1}{2}$  to  $2\frac{3}{4}$  in head; outer row of teeth on upper jaw enlarged; the narrow band of teeth back of this row separated from it by a space; teeth on lower jaw in a band, subequal. Scales cycloid anteriorly, becoming larger and ctenoid posteriorly. Caudal  $2\frac{1}{2}$  to  $2\frac{3}{4}$  in body. Male, light olive, with dark-olive blotches; body and head with many conspicuous round cream-colored spots, each surrounded by a dusky ring, these smaller than pupil and most distinct on head; snout with dusky streaks; dorsal and caudal plainly barred; pectoral crossed with dark wavy lines, dusky at base; anal and ventrals dusky; a small dark spot at base of caudal; a shining deep-green spot inside the mouth in life. Female, plain olivaceous, nearly or quite immaculate. West Indies, south to Rio Janeiro north to St. Augustine, Florida (Dr. Oliver P. Hay), and to Charleston (C. H. Gilbert); specimens before us from Marco Island, Florida (J. A. Henshall). (*Μαράγδος*, emerald, from the bright-green spot on the tongue.)

*Gobius smaragdus*, CUVIER & VALENCIENNES, Hist. Nat. Poiss., XII, 120, 1837, Cuba; JORDAN & EIGENMANN, l. c., 497.

*Smaragdus valenciennesi*, POEY, Memorias, II, 280, 1861, Cuba.

*Gobionellus smaragdus*, POEY, Synopsis, 394, 1868; POEY, Enumeratio, 126, 1876.

2555. GOBIUS STRIGATUS, O'Shaughnessy.

Head  $3\frac{1}{2}$ ; depth 5; eye  $3\frac{1}{2}$ , shorter than the rounded snout. D. VI-12; A. 11 or 12; scales 53-13. Body elongate, compressed posteriorly; head little compressed; maxillary reaching to below middle of eye; teeth small, the outer a little enlarged; dorsal spines all shorter than head, not filamentous. Head naked; anterior half of body covered with ctenoid scales, those on nape much reduced in size. Two violet stripes from mouth to eye, 8 or 9 violet bars on side, 3 or 4 on caudal; second dorsal spotted. (Steindachner.) Coast of Surinam. (*strigatus*, striped.)

*Gobius strigatus*, O'SHAUGHNESSY, Ann. Mag. Nat. Hist., series 4, XV, 1875, 145, Surinam.

*Gobius kraussi*, STEINDACHNER, Ichth. Beiträge, VIII, 16, 1879, Surinam; JORDAN & EIGENMANN, l. c., 497.

2556. GOBIUS SAGITTULA (Günther).

Head  $4\frac{1}{2}$  to 5 in length to base of caudal; depth 6 to 8; eye  $4\frac{1}{2}$  to  $5\frac{1}{2}$ . D. VI-13 or 14; A. 13; scales about 66 in longitudinal, 15 in transverse series, counted just below space between the two dorsals. Body slender, tapering pretty regularly from middle of first dorsal to caudal, most compressed posteriorly, depth about uniform from head to origin of second dorsal. Head short, depressed, and broad; mouth large, nearly horizontal, the maxillary in adults  $2\frac{3}{4}$  in head, reaching beyond middle of eye; distance between maxillaries at their posterior ends greater than their length; eye about  $\frac{2}{3}$  the bony interorbital space. Teeth in a narrow band in each jaw, those in lower jaw uniform, the outer series in upper jaw considerably enlarged and separated by an interspace from the inner band. Pseudobranchiae well developed. Gill rakers short and flexible. Longest dorsal spine about  $\frac{2}{3}$  head; distance between dorsals less than diameter of eye; pectorals  $1\frac{1}{2}$  in head, their tips reaching past middle of spinous dorsal; ventrals about equaling pectorals, reaching more than halfway to origin of anal; anal equal and opposite to the second dorsal, but slightly lower; caudal fin greatly elongate, more than  $\frac{1}{2}$  head and

body in  
with ag  
ered wi  
toward  
General  
sides wi  
under fi  
is somet  
near its  
upon ea  
and ma  
dorsal fi  
with cro  
ous narro  
and neig  
mouths o  
of longie  
arrow.)

*Euctenogobius*  
Americ  
of Centr  
*Gobius longi*  
Guayma  
*Gobius sagitt*

Head 4  
11-15. Bo  
out; head  
jaws equal  
lower jaw  
the upper j  
outer teeth  
embedded,  
*Gobius ocean*  
without the  
spines filam  
mentous, 2  
trials, none  
to this speci  
length, 7 in  
obsolescence  
ent coloratio  
*oceanicus*, wi  
The two need

*Gobionellus ha*  
Texas; GIB  
*Gobius lanceola*

body in largest specimens,  $2\frac{1}{2}$  in smaller ones, its relative length increasing with age. Head scaleless, predorsal region with small scales; body covered with close-set etenoid scales, small and greatly crowded anteriorly, toward the caudal fin growing gradually larger and more strongly etenoid. General color light yellowish, palest below, upper parts darker; middle of sides with 5 elongate black blotches, most distinct in the young; the first under first dorsal, second under origin of second dorsal, the third, which is sometimes almost double, at about middle of second dorsal, the fourth near its posterior end, and the last at base of caudal; a large black spot upon each shoulder just above origin of pectoral fin; head plain; lips and maxillary dark; opercle with a dark blotch; basal portion of dorsal fins with dark lines formed of spots; anal unmarked; pectorals with cross lines formed of dots; ventrals plain; caudal crossed by numerous narrow dark bars. Reaching a length of 8 inches. Gulf of California and neighboring waters south to Panama; very common in lagoons and mouths of rivers. The types of *sagittula* are evidently the young, those of *longicauda* the adults of the same species. (diminutive of *sagitta*, arrow.)

*Euctenogobius sagittula*, GÜNTHER, Proc. Zool. Soc. London 1861, 3, West coast Central America, young individuals; GÜNTHER, Cat. Fishes, III, 555, 1861; GÜNTHER, Fishes of Centr. Amer., 389, 1869.

*Gobius longicauda*, JENKINS & EVERMANN, Proc. U. S. Nat. Mus. 1888, 146, adult examples, Guaymas. (Coll. Evermann & Jenkins. Type, No. 39636.)

*Gobius sagittula*, JORDAN & EIGENMANN, l. c., 497.

2557. GOBIUS HASTATUS, Girard.

(EMERALD FISH; SHARP-TAILED GOBY.)

Head  $4\frac{1}{2}$  to 5; depth 6 to  $7\frac{1}{2}$ . D. VI-14; A. 14 or 15; scales 60; vertebrae 11+15. Body compressed, extremely elongate; depth nearly equal throughout; head short, compressed, deeper than wide; mouth wide, oblique, the jaws equal; maxillary in adult reaching to below posterior border of eye; lower jaw very thin and flat; teeth in each jaw small, subequal, those in the upper jaw in a single series, those in the lower jaw in a narrow band; outer teeth somewhat movable; scales anteriorly small, cycloid, and embedded, those behind larger and etenoid; the scales larger than in *Gobius oceanicus*; a few scales on upper anterior corner of opercle, but without the large patch seen in *G. oceanicus*; dorsal fins high, some of the spines filamentous and longer than the head; caudal very long and filamentous, 2 to  $2\frac{1}{2}$  in body; pectoral slightly longer than head or than ventrals, none of its rays silk-like. A single specimen from Ceylon belongs to this species, which appears to be characterized by a longer head (5 in length, 7 in total), by the much larger scales (60 in a lateral line), by the obsolescence of the patch of scales on opercles, and by slightly different coloration. This may be really only the extreme of variation of *G. oceanicus*, with which species most authors have hitherto confounded it. The two need detailed comparison. Coast of Texas. (*hastatus*, spear-like.)

*Gobionellus hastatus*, GIRARD, Proc. Ac. Nat. Sci. Phila., 1858, 168, St. Josephs Island, Texas; GIRARD, U. S. and Mex. Bound. Surv., 25, pl. 12, figs. 7 and 8, 1859.

*Gobius lanceolatus*, GÜNTHER, Cat., III, 50, 1861, and of authors; not of BLOCH.

2558. *Gobius oceanicus*, Pallas.

(ESMERALDA; ENDORMI ÉMERAUDE; BACALIAO SABARA.)

Head  $4\frac{1}{2}$  to 6; depth  $6\frac{1}{2}$  to  $8\frac{1}{2}$ ; eye 4 to 5 in head; ventral 6 to  $6\frac{1}{2}$ ; pectorals  $5\frac{1}{2}$  to  $6\frac{1}{2}$ . D. VI-14. A. I, 14 or 15; scales about 65. Body extremely elongate; head very short; upper part of opercle scaled, head otherwise naked. Scales on body very small, becoming much larger behind. All the dorsal spines more or less filamentous; caudal fin nearly half length of rest of body. Skull behind eye broad and short, its length  $1\frac{1}{2}$  in width, no decided ridges nor crests; lateral crests large and stout behind, minute forward; interorbital area narrow, deeply grooved, with a median ridge. Color in spirits, reddish olive; a distinct, round, blackish blotch below spinous dorsal, twice as large as orbit; an indistinct dusky shade along middle of sides, terminating in a dusky blotch on base of caudal; middle of sides with a series of marks, formed by very voiny lines widely diverging backward; a similar narrow line from eye to maxillary, and 1 from eye backward to upper angle of preopercle; evident traces of the emerald spot at base of tongue; 2 small dark spots on first dorsal spine; spinous dorsal dusky, with a light and dusky streak at base; soft dorsal dusky, a light (bluish in life) area behind each ray; anterior rays barred with light and dark; anal and ventrals whitish (probably blue in life), the ventrals without dark markings; pectorals dusky, the base lighter, and with some indistinct dusky bars; a dusky half bar on the upper part of the axil; base of tongue tuberculate, and shining with bright blue and green reflections like a precious stone (hence the names *smaragdus*, *esmeralda*, etc.), this color fading in spirits. Vertebrae elongate,  $11 + 15 = 26$ . Length a foot. South Atlantic and Gulf coasts of the United States and southward through the West Indies; not rare, perhaps intergrading with the preceding. Here described from a specimen 11 inches long, taken by Dr. Gilbert in Charleston Harbor. (*oceanicus*, ocean.)

*Gobius cauda longissima acuminata*, GRONOW, Zooph., 82, No. 277, pl. 4, fig. 4, 1763, locality unknown.

*Gobius oceanicus*, PALLAS, Spicilegia, VIII, 4, 1769, locality unknown; after GRONOW; JORDAN & EIGENMANN, L. c. 497.

*Gobius lanceolatus*, BLOCH, Fische Deutschlands, II, 8, pl. 38, fig. 1, 1783, Martinique, figure probably from PLUMIER; CUVIER & VALENCIENNES, Hist. Nat. Poiss., XII, 114, 1837; POEY, Synopses, 393, 1868.

*Gobius bacalans*, CUVIER & VALENCIENNES, Hist. Nat. Poiss., XII, 119, 1837, Surinam (Coll. Le Valliant); Cayenne (Coll. Richard); Cuba (Coll. Poey).

*Gobionellus oceanicus*, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1882, 613; JORDAN & GILBERT, Synopses, 636, 1883.

## Subgenus LYTHRYPNUS, Jordan &amp; Evermann.

2559. *Gobius dalli*, Gilbert.

Head  $3\frac{3}{4}$ ; depth  $4\frac{1}{2}$ . D. VI-17; A. 14; scales 40. Body short, compressed, resembling *Microgobius*. Head high, mouth moderate, very oblique; upper pectoral rays normal; scales ctenoid, of moderate size; anterior dorsal spines much produced. Mouth very oblique, the maxillary

reaching  
diameter  
distant, c  
teeth. D  
largest sp  
last dorsa  
dorsal rat  
head; ven  
rays not fr  
trunk, wit  
cous, and  
our specim  
bands not  
and fainter  
uniting on  
encircles t  
portions, 1  
and backw  
along profi  
obliquely d  
cally down  
spinous dor  
Several sm  
3001, in 33 f  
Dall, in abo  
type of a dis  
of the body  
*Gobius dalli*, G  
fornia (Col

*Garmannia*, Jo  
*Euppnias*, Jo

Anterior h  
small scales;  
in front. Ven  
for Mr. Samu  
Comparative  
contribution

GARMANNIA:

a. Scales mod

b. Scales

11

16

bb. Scale

d

reaching vertical from front of pupil,  $2\frac{1}{2}$  in length of head. Snout short,  $\frac{3}{8}$  diameter of orbit, which is 3 in head. Jaws with an outer series of long, distant, canine-like teeth, and an inner series or a narrow band of minute teeth. Dorsal spines 6, the 2 anterior greatly elongate, not free, in our largest specimen extending beyond middle of soft dorsal; membrane from last dorsal spine reaching to, or nearly to, base of first soft ray; soft dorsal rather high, the fin long; caudal rounded, less than length of head; ventrals free from belly, fully united; pectorals short, the upper rays not free nor silk-like. Scales of moderate size, ctenoid, covering entire trunk, with possible exception of the nape; the scales are readily caducous, and are lacking on nape and frequently on anterior third of body in our specimens. Color light coral red, anteriorly with 4 to 6 narrow blue bands not reaching ventral outline, the posterior ones growing narrower and fainter; a blue streak upward and backward from each orbit, the 2 uniting on occiput; a transverse interorbital bar, a continuation of which encircles the orbit anteriorly; below orbit, a blue bar consisting of 2 portions, 1 running downward and obliquely backward, the other upward and backward; in the largest specimen a blue streak runs from occiput along profile to front of dorsal; the first blue bar runs from nape obliquely downward and forward, ending on opercle; the second vertically downward from front of spinous dorsal, the third under middle of spinous dorsal, the remaining bars under soft dorsal; fins unmarked. Several small specimens, the largest 1 inch long, from *Albatross Station* 3001, in 33 fathoms. A single slightly larger example dredged by Dr. W. H. Dall, in about 35 fathoms, off Catalina Harbor, California. Probably the type of a distinct genus distinguished by the many-rayed fins and the form of the body and head. (Named for its discoverer, William Healey Dall.)

*Gobius dalli*, GILBERT, Proc. U. S. Nat. Mus. 1890, 73, *Albatross Station* 3001, Lower California (Coll. *Albatross*); Catalina Harbor (Coll. W. H. Dall).

#### 814. GARMANNIA, Jordan & Evermann.

(HALF-NAKED GOBIES.)

*Garmannia*, JORDAN & EVERMANN, Proc. Cal. Ac. Sci. 1895, 495, pl. 49 (*paradoxus*).  
*Eupinnias*, JORDAN & EVERMANN, new subgenus (*seminudus*).

Anterior half of body naked; posterior half covered with moderate or small scales; teeth rather strong, unequal, usually 2 small curved canines in front. Very small gobies. Otherwise essentially as in *Gobius*. ("Named for Mr. Samuel Garman, the accomplished ichthyologist of the Museum of Comparative Zoology at Cambridge, Mass., in recognition of his important contributions to ichthyology.")

GARMANNIA:

a. Scales moderate.

b. Scales ctenoid, 13 or 14 series developed; first dorsal spine filamentous; D. VI-11; A. 9. Body rather robust, the depth about  $4\frac{3}{8}$  in length; the head  $3\frac{3}{8}$ ; lower jaw with 2 curved canines. PARADOXA, 2560.

bb. Scales smaller, 17 series developed; depth  $4\frac{3}{8}$  in length. D. VI-10; A. 8; first dorsal spine not filamentous; lower jaw with small canines.

HEMIGYMA, 2561.

## ENYPNIAS (ένύπνιος, in one's sleep):

aa. Scales excessively minute; body slender, the depth 6 in length. D. VI-15; A. 10; dorsal spines not filamentous; lower jaw with 2 small curved canines in front. SEMINUDA, 2562.

## Subgenus GARMANNIA.

## 2560. GARMANNIA PARADOXA (Günther).

Head about  $3\frac{1}{2}$  ( $4\frac{1}{2}$  in total); depth about  $4\frac{2}{3}$  ( $5\frac{2}{3}$  with caudal). D. VI-11; A. 9; scales 14. Head nearly as broad as high, its width being rather more than  $\frac{1}{2}$  of its length. Eyes rather close together, of moderate size. Snout obtuse, rounded, as long as the eye; cleft of the mouth slightly oblique, with the jaws equal in length, and with maxillary extending to below middle of the eye. Teeth in villiform bands; 2 curved canine teeth on each side of the lower jaw. Head and trunk entirely naked to between second dorsal and anal, the remainder covered with etenoid scales of moderate size, 9 or 10 of them in 1 of the anterior transverse series. First dorsal spine elongate, filiform, sometimes extending to the base of the caudal; caudal rounded, shorter than head; none of pectoral rays silk-like; ventral terminating at a great distance from vent. Blackish in spirits; caudal and ventral fins black, dorsal filament whitish. (Günther.) Panama to Mazatlan; scarce. Our single specimen from the estuary at Mazatlan differs somewhat from Dr. Günther's account. It is thus described: Head  $3\frac{1}{2}$ ; depth  $4\frac{1}{3}$ . D. VI-11; A. 9; scales 12; eye 4 in head; snout  $4\frac{1}{2}$ ; pectoral  $1\frac{1}{2}$  in head; dorsal spine  $1\frac{1}{2}$ . Form of *Gobiosoma bosci*. Body compressed; head broad and depressed, with tumid cheeks; snout not very blunt, short, oblique-truncate; eyes rather large, high, the maxillary not produced, extending to their posterior margin; mouth large, oblique; lower jaw heavy, slightly projecting; teeth strong; gill openings narrow, not wider than base of pectoral. First dorsal rather high, the first spine filamentous, reaching past soft dorsal; other fins low. Head and anterior half of body to front of soft dorsal naked; scattering scales coming in above, 12 rows of imbricated slightly etenoid scales along median line of caudal peduncle and forward to middle of soft dorsal, the scaled area about as long as head, the upper parts better scaled than lower. No flaps on shoulder girdle. Olivaceous, with 7 or 8 dark cross shades, 2 on head, 1 across gill openings, 1 behind pectoral, and a broad 1 below soft dorsal; dorsals dusky, the filamentous ray pink; lower half of soft dorsal yellowish, upper dusky; lower fins black; caudal dusky; a dark speck at angle of opercle; skin everywhere punctate with black; a pale olive bar at base of caudal. Skull without median crest; interorbital space not concave; head not very abruptly widened behind eyes. Pacific coast of Mexico and Central America. One specimen,  $1\frac{1}{2}$  inches long, recently obtained on muddy bottoms among the mangroves lining the estuary at Mazatlan. (*paradoxus*, paradox.)

*Gobius paradoxus*, GÜNTHER, Proc. Zool. Soc. Lond. 1861, 3, west coast Central America; GÜNTHER, Cat., III, 549, 1861; JORDAN & EIGENMANN, Proc. U. S. Nat. Mus. 1886, 498. *Garmannia paradoxa*, JORDAN, Proc. Cal. Ac. Sci. 1895, 497, pl. 59.

Head  
Garmann  
greatest  
fins. He  
to mouth  
 $1\frac{1}{2}$  in rou  
orbit; m  
ing beyon  
teeth in v  
times as l  
jaw simil  
weakly et  
extending  
tion; the r  
scales, the  
ward to th  
smallest.  
head, the t  
height, equ  
than the an  
of body,  $1\frac{1}{2}$   
rounded, ra  
out distinct  
faint cross l  
fins, are for  
last at base  
anterior hal  
of caudal; i  
eye to upper  
tions closely  
the dredge.

*Gobius henigmyi*  
the West I

Head 4; dep  
naked; sides  
posteriorly.  
its width  $\frac{2}{3}$  le  
size; snout ob  
with the jaws  
the middle of  
jaw slightly e  
jaw. Dorsal f  
lower than ant

## 2561. GARMANNIA HEMIGYMNA (Eigenmann &amp; Eigenmann).

Head  $3\frac{3}{8}$  ( $4\frac{1}{2}$  in total); depth  $4\frac{3}{8}$ . D. VI-10; A. 8; scales smaller than in *Garmannia paradoxa*, 17-7. Body compressed, depressed anteriorly, the greatest depth in this specimen being at origin of anal and second dorsal fins. Head wider than deep, rounded; profile much decurved from eye to mouth as in *paradoxa*; eye perfectly round, smaller than in *paradoxa*,  $1\frac{1}{8}$  in rounded snout, 5 in head; interorbital space scarcely wider than orbit; mouth somewhat oblique, larger than in *paradoxa*; maxillary reaching beyond posterior rim of orbit; lower jaw slightly shorter than upper; teeth in upper jaw in a band, the outer series remote, and the teeth several times as large as in the inner row, all more or less movable; teeth in lower jaw similar, a recurved canine on each side near the front. Scales very weakly ctenoid, covering only the sides of the posterior half of body, not extending quite to base of dorsal or anal fins even at their posterior insertion; the upper and lower edges of the caudal peduncle likewise free from scales, the scaly region, however, widest on peduncle and tapering forward to the central point opposite beginning of anal, where the scales are smallest. First spine of the dorsal not elongate as in *G. paradoxa*,  $1\frac{1}{2}$  in head, the third, fourth, and fifth spines slightly exceeding the first in height, equaling the posterior rays of soft dorsal, which are little higher than the anterior rays of the soft dorsal; caudal rounded, about 4 in length of body,  $1\frac{1}{2}$  in head; ventral not reaching vent,  $1\frac{1}{2}$  in head; pectorals rounded, rather short and broad,  $1\frac{1}{2}$  in head. Color light olivaceous, without distinct markings, everywhere with minute dark punctulations; 8 faint cross bars from dorsal to middle of sides, which, close under dorsal fins, are formed of 2 blackish dots; 8 black dots along lateral line, the last at base of caudal; fins all smutty, the pectoral lightest, white on its anterior half, 2 dusky spots at its base; opercle ashy; a light bar at base of caudal; iris blackish blue, a short straight streak of same color from eye to upper lip; an irregular bluish mark on cheeks formed of punctulations closely crowded. West Indies, exact locality unknown; taken with the dredge. (*ἡμί*, half; *γυμνός*, naked.)

*Gobius hemigymnus*, EIGENMANN & EIGENMANN, Proc. Cal. Ac. Sci. 1888, 66, dredged in the West Indies.

Subgenus ENYPNIAS, Jordan & Evermann.

## 2562. GARMANNIA SEMINUDA (Günther).

Head 4; depth 6. D. VI-15; A. 10. Head and anterior portion of trunk naked; sides with exceedingly small scales, becoming somewhat larger posteriorly. Head with the cheeks swollen, depressed, broader than high, its width  $\frac{3}{4}$  length. Eyes close together, directed upward, of moderate size; snout obtuse, as long as the eye; cleft of the mouth slightly oblique, with the jaws equal anteriorly, and with the maxillary extending to below the middle of the eye. Teeth in villiform bands, the anterior of the lower jaw slightly enlarged; 2 small curved canine teeth on each side of lower jaw. Dorsal fins rather low, the hind part of the spinous dorsal scarcely lower than anterior; caudal rounded, as long as pectoral; none of pectoral



rays silk-like; ventral rather short, terminating at a great distance from vent. Blackish; fins and sides of head dotted with black; ventrals black. (Günther.) Panama; not seen by us; probably the type of a distinct genus. (*seminudus*, half-naked.)

*Gobius seminudus*, GÜNTHER, Proc. Zool. Soc. London 1861, 3, west coast Central America; GÜNTHER, Cat., III, 554, 1861; JORDAN & EIGENMANN, Proc. U. S. Nat. Mus. 1886, 498.

### 815. AWAOUS, Steindachner.

*Awaous*,\* STEINDACHNER, Verh. Mat. Phys. Naturw. 1860, 289; after *les Awaous* of CUVIER & VALENCIENNES (*ocellaris*, etc).

*Chonophorus*, POEY, Mémoires, II, 274, 1861 (*buccelentus* = *taiasica*).

*Awaous*, BLEEKER, Esquisse d'un Syst. Nat. Gobioides, 320, 1874 (*ocellaris*); after *les Awaous* of CUVIER & VALENCIENNES.

Inner edge of shoulder girdle with 2 or more conspicuous dermal flaps; preorbital region very long; premaxillary and maxillary strong; lips thick; scales rather small, etenoid, 40 to 80 in a longitudinal series; interorbital groove with a conspicuous median crest; otherwise essentially as in *Gobius*. The species reach a large size and are confined to the fresh waters of the tropics of America and the Hawaiian Islands. The Asiatic species of similar habit have much larger scales and seem to form a distinct genus, *Rhinogobius*, Gill. The physiognomy in each is peculiar, the snout being long and convex. (*Awaou*, a Hawaiian name.)

a. Scales about 53, little crowded anteriorly, 21 before dorsal on nape; depth  $5\frac{1}{2}$  in length; head 4; eyes placed high, interorbital area equal to diameter of eye; mouth horizontal; maxillary extending to middle of eye. In head, lower jaw more flat than in *A. taiasica*; teeth small, in narrow bands, those of the outer row above enlarged, some large teeth in band of lower jaw. D. VI-I, 12; A. I, 10. Uniform yellowish in spirits. FLAVUS, 2563.

aa. Scales 60 to 70, crowded anteriorly, about 30 scales before the dorsal on nape; body compressed posteriorly, rather depressed anteriorly; greatest depth  $5\frac{1}{2}$  in length; head  $3\frac{1}{2}$  in length. Olivaceous, a series of irregular, roundish blotches along middle of sides; narrow dark streaks radiating from eye; a blackish streak running across upper margin of opercle and extending obliquely across base of upper pectoral rays; belly white; dorsal and caudal more or less distinctly barred with wavy blackish lines.

b. About 15 scales between second dorsal and base of anal. NELSONI, 2564.

bb. About 21 scales between second dorsal and base of anal. TAIASICA, 2565.

aaa. Scales 76 to 82; 24 scales between second dorsal and anal; head as broad as high; depth of body  $6\frac{1}{2}$  in length; head 4; head flat above, snout elongate, upper profile oblique; eye  $\frac{1}{2}$  of head, equals interorbital area (in adult); mouth horizontal; lower jaw included; maxillary reaching to below anterior margin of eye; teeth of the outer series enlarged; canine teeth none; scales etenoid, those on nape and anterior part of body very small; head naked; dorsal fins lower than body, none of the spines produced; caudal rounded, 7 in length of body. Yellowish olive; back and sides reticulated with blackish; head, dorsal, caudal, and pectoral fins dotted with blackish, the spots forming streaks on second dorsal; 6 cross series of dots on the caudal; an irregular small blackish spot on the upper part of the root of pectoral. D. VI-11; A. 11; scales about 80. MEXICANUS, 2566.

\*The name "Les Awaous," given to this group by Valenciennes, was a French plural, not a generic appellation, and if used as the name of a genus must be dated from its use in that sense by Steindachner or Bleeker. The Hawaiian type of "*Awaous*" agrees with the American species (*Chonophorus*) in the character of the flaps on the shoulder girdle, as well as in general appearance. The Asiatic genus, *Rhinogobius*, Gill (*similis*), seems to be very close to *Chonophorus*, but the scales are larger, 28 in the lateral series.

Head  
D. VI-1,  
before t  
eye,  $2\frac{1}{2}$  i  
narrow l  
of lower  
of sides;  
line alon  
Rivers of  
*Gobius flavu*  
Cat. Fis  
*Chonophoru*

Head  $3\frac{1}{2}$   
11; scales a  
large, qua  
abruptly de  
groove with  
vertical of  
the outer s  
head; ventr  
Dorsal fins  
slender, wea  
from caudal  
 $1\frac{1}{2}$  in head.  
erate, high t  
etenoid, ver  
teriorly, abo  
backward to  
few minnte  
and blotche  
under middle  
spots; cauda  
pale; pectora  
blotch at bas  
orbital, longe  
4 inches. Kn  
mens were ol  
Edward Will  
his work upo

*Awaous nelsoni*,  
pools at Ros  
533 U. S. Fish

2563. AWAOUS FLAVUS (Cuvier & Valenciennes).

Head 4; depth  $5\frac{3}{8}$  to  $6\frac{1}{2}$ ; eye equal to the interorbital width, placed high. D. VI-I, 12; A. I, 10; scales about 53 to 55, little crowded anteriorly, 21 before the dorsal. Mouth horizontal, maxillary extending to middle of eye,  $2\frac{1}{4}$  in head; lower jaw flatter than in *Awaous taiasica*; teeth small, in narrow bands, those of the outer row enlarged; some large teeth in band of lower jaw. Yellowish, with a row of faint ocellated spots along middle of sides; dorsal and caudal faintly barred; lines radiating from eye, a line along opercle halfway to pectoral, sometimes uniform blue-black. Rivers of Surinam and Brazil, south to Bahia. (*flavus*, yellow.)

*Gobius flavus*, CUVIER & VALENCIENNES, Hist. Nat. Poiss., XII, 60, 1837, Surinam; GÜNTHER, Cat. Fish., III, 13, 1861.

*Chonophorus flavus*, JORDAN & EIGENMANN, l. c., 500; EIGENMANN & EIGENMANN, l. c., 67.

2564. AWAOUS NELSONI, Evermann.

Head  $3\frac{1}{2}$ ; depth 6; eye  $5\frac{1}{2}$  in head; snout 3; maxillary  $2\frac{1}{4}$ . D. VI-11; A. 11; scales about 63. Body long, compressed and tapering posteriorly; head large, quadrate, mouth nearly horizontal, lower jaw included; snout abruptly decurved; top of head flat, the interorbital with a slight median groove with a thin, raised edge on each side; maxillary reaching about to vertical of anterior edge of pupil; teeth in bands on jaws very small, the outer somewhat enlarged; pectoral rays normal, the longest  $1\frac{1}{2}$  in head; ventrals completely united, the disk free from belly,  $1\frac{1}{2}$  in head. Dorsal fins separated by a space about  $\frac{2}{3}$  diameter of eye; dorsal spines slender, weak, about  $1\frac{1}{2}$  in head; soft dorsal and anal similar, each free from caudal; caudal fin rather short and rounded, its middle rays about  $1\frac{1}{2}$  in head. Gill membranes broadly united to the isthmus; eyes moderate, high up, the interorbital width equal to the eye's diameter. Scales ctenoid, very small and irregularly crowded anteriorly, much larger posteriorly, about 15 rows counting from origin of soft dorsal downward and backward to the anal fin; head naked, but with slight indication of a few minute embedded scales on opercles. Color grayish; head mottled and blotched with dark; side with 7 or 8 black blotches, the largest under middle of pectoral fin; dorsals pale, crossed by several lines of black spots; caudal pale, with about 6 or 7 dark cross bars; ventrals and anal pale; pectorals pale, dusted with dark specks and with a small dark blotch at base of upper rays. Close to *A. taiasica*, but with broader interorbital, longer snout and larger scales on posterior part of body. Length 4 inches. Known only from fresh water at Rosario, Sinaloa, where 8 specimens were obtained July 27, 1897, by Mr. E. W. Nelson. (Named for Mr. Edward William Nelson, the well-known ornithologist, in recognition of his work upon the fishes of Illinois in 1876.)

*Awaous nelsoni*, EVERMANN, Proc. Biol. Soc. Washington, vol. XII, 1898, 3, fresh-water pools at Rosario, Sinaloa, Mexico. (Type, No. 48836, U. S. Nat. Mus.; cotypes, No. 533 U. S. Fish Comm., 5793 L. S. Jr. Univ. Mus., and 48837 U. S. Nat. Mus.)

## 2565. AWAOUS TAIASICA (Lichtenstein).

(GUAVINA HOYERA; ANOMA DE RIO.)

Head  $3\frac{1}{2}$ ; depth  $5\frac{1}{2}$ ; eye small, less than interorbital width (in adult), 3 in snout (twice in young), and about 7 in head. D. VI-11; A. 11; scales 60 to 70, crowded anteriorly, about 30 before dorsal fin, 21 between second dorsal and anal. Body compressed posteriorly, rather depressed anteriorly; head broader than deep. Distance from eye to mouth  $3\frac{1}{2}$  in head, the preorbital being much enlarged; mouth large, horizontal, maxillary extending to below anterior part of orbit in adult male, shorter in young; lower jaw included. Teeth of the upper jaw in 2 series, those in anterior series much enlarged and recurved; teeth of lower jaw in a narrow band, the outer series scarcely enlarged. Inner edge of the shoulder girdle with 2 or 3 rather long papillae. Body covered with ctenoid scales, much reduced in size anteriorly; nape closely scaled, breast scaly, head naked. Dorsal fins less than depth of body, the spines scarcely filamentous, not as long as the soft rays; caudal rounded, shorter than the head; ventrals very broad and short,  $1\frac{1}{2}$  to  $1\frac{3}{4}$  in head, the rays very much branched. Skull rounded behind, with a very short crest in its middle; lateral crests high and thin, converging into 1 opposite the insertion of suprascapula, inner crests not meeting behind eye, the outer ones extending around orbit. A low, blunt ridge between the posterior corners of orbit, becoming much higher forward, continued as the ethmoid and ending abruptly some distance in advance of orbit. Teeth in upper jaw in a few series, those of outer series many times larger than the others, which are minute; those of lower jaw all alike small, in a band. Olivaceous, with a series of irregular, roundish blotches along middle of side, and narrow dark streaks radiating from eye; a blackish streak running across upper margin of opercle and extending obliquely across base of upper pectoral ray; belly white; dorsal and caudal more or less distinctly barred with wavy blackish lines. Length a foot or more. Extremely variable in form and coloration, as is the case with most widely distributed fresh-water fishes. Fresh waters of the West Indies and both coasts of Mexico, south to Brazil; common in Cuba, in Sinaloa, and about La Paz in Lower California, thence southward to Panama. (*taiasica*, Brazilian name of some other goby.)

*Awaous guacu*, MARCGRAVE, Hist. Brasil., 166, 1648, Brazil.

*Gobius taiasica*, LICHTENSTEIN, Berl. Abhandl., 273, 1822, Brazil; not *Tajasica* MARCGRAVE. *Gobius banana*, CUVIER & VALENCIENNES, Hist. Nat. Poiss., XII, 103, 1837, San Domingo; GÜNTHER, Cat., III, 59, 1861.

*Gobius martinicus*, CUVIER & VALENCIENNES, Hist. Nat. Poiss., XII, 105, 1837, Martinique. *Chonophorus bucculentus*, POEY, Memorias, II, 275, 1861, Cuba.

*Rhinogobius contractus*, \* POEY, Memorias, II, 424, 1861, Cuba; POEY, Enumeratio, 125, 1875

\* The following are the characters assigned to *Awaous contractus* (Poey): Head 4; depth  $5\frac{1}{2}$ ; D. VI-11; A. 11; eye 7 in head; maxillary ceasing  $\frac{1}{2}$  an eye's diameter before eye. Head smaller than in *A. taiasica*. Greenish brown; the cheeks with brown lines; body with brown points; dorsals brownish, with brown longitudinal bands more numerous on the second; caudal with 7 brown bands, made of lanceolate spots on the rays; pectorals speckled; ventrals and anal rose color. Cuba (Poey); probably not different from *A. taiasica*; said to differ in the small mouth, which probably varies with age and sex.

*Gobius a*  
Mex  
Enclenog  
(Coll  
Chonopho

Head  
second d  
gate, up  
illary re  
series en  
anterior p  
depth of  
of body.  
head, dor  
forming s  
an irregul  
toral. (G  
known to

*Gobius mezi*  
*Chonophoru*

*Bollmannia*,

This geni  
inner edge d  
without tra  
*Gobius prop*  
the presenc  
of the Pacif  
gobies. ("I  
Charles Har  
exploration  
ing through

a. \* A conspi  
least  
b. Filan  
o  
bb. Fila

aa. No black  
depth  
c. Head

cc. Head

\* This anal

*Gobius dolichocephalus*, COPE, Trans. Amer. Philos. Soc. Phila. 1869, 403, near Orizaba, Mexico.

*Euctenogobius latus*, O'SHAUGHNESSY, Ann. Mag. Nat. Hist., Series 4, xv, 1875, 146, Bahia. (Coll. Dr. Wucherer.)

*Chronophorus taiasica*, JORDAN & EIGENMANN, l. c., 500.

2566. AWAOUS MEXICANUS (Günther).

Head 4; depth 6½; eye 8. D. VI-11; A. 11; scales 76 to 82, 24 between second dorsal and anal. Head as broad as deep, flat above, snout elongate, upper profile oblique; mouth horizontal, lower jaw included, maxillary reaching to below anterior margin of eye. Teeth of the outer series enlarged; no canine teeth. Scales ctenoid, those on nape and anterior part of body very small; head naked. Dorsal fins lower than depth of body, none of the spines produced; caudal rounded, 7 in length of body. Yellowish olive; back and sides reticulated with blackish; head, dorsal, caudal, and pectoral fins dotted with blackish, the spots forming streaks on second dorsal; 6 cross series of dots on caudal; an irregular, small blackish spot on the upper part of the base of the pectoral. (Günther.) Fresh-water streams of the eastern slope of Mexico; known to us only from Dr. Günther's description.

*Gobius mexicanus*, GÜNTHER, Cat., III, 61, 1861, Mexico.\*

*Chronophorus mexicanus*, JORDAN & EIGENMANN, l. c., 501.

816. BOLLMANNIA, Jordan.

*Bollmannia*, JORDAN, Proc. U. S. Nat. Mus. 1889, 164 (*chlamydes*).

This genus differs from *Lepidogobius* by having no fleshy processes on inner edge of shoulder girdle, the interorbital area of skull narrower and without trace of median keel, and by very large ctenoid scales. From *Gobius* proper it is distinguished by the presence of 7 dorsal spines and by the presence of large scales on the cheeks. Species inhabiting the depths of the Pacific; not found in shoal waters as is the case with most other gobies. ("I have named this species in honor of my late colleague, Mr. Charles Harvey Bollman, whose untimely death, while engaged in the exploration of the rivers of Georgia, took place while this paper was passing through the press."—Jordan.)

a. \* A conspicuous black spot on posterior portion of spinous dorsal. Body deep, the least depth of caudal peduncle greater than diameter of orbit.

b. Filamentous dorsal spines very long, reaching beyond middle of soft dorsal when depressed. Lower caudal rays black; dorsal spot conspicuously ocellated. Eye large, 3 to 3½ in head. OCELLATA, 2567.

bb. Filamentous dorsal spines shorter. Lower caudal rays not black, and dorsal spot not ocellated. Eye smaller, 3¾ to 4 in head. CHLAMYDES, 2568.

aa. No black spot on spinous dorsal. Body slender, the depth ½ the length. Least depth of caudal peduncle not greater than diameter of orbit.

c. Head large, 3 to 3½ in length. No black spot at base of caudal. Fins low.

MACROPOMA, 2569.

cc. Head smaller, 2½ in length. A black spot at base of caudal. Fins higher.

STIGMATURA, 2570.

\* This analysis of species is taken from Gilbert, Proc. U. S. Nat. Mus. 1891, 555.

2567. *BOLLMANNIA OCELLATA*, Gilbert.

Head  $3\frac{1}{2}$  to  $3\frac{3}{8}$  in length; depth  $4\frac{1}{2}$ . D. VII-14 or 15; A. 14; scales 27. Very close to *Bollmannia chlamydes*, differing from the latter constantly in the following respects: The eye is larger, 3 to  $3\frac{1}{2}$  in head ( $3\frac{1}{2}$  to 4 in *chlamydes*); the filamentous rays of spinous dorsal are much longer, reaching in adults, when laid back, to or nearly to end of base of soft dorsal,  $1\frac{1}{2}$  to  $1\frac{3}{4}$  times length of head; rarely the filamentous dorsal rays are little more elongate than in *chlamydes*. Teeth in a narrow band in each jaw, the outer series in upper jaw, and both outer and inner series in lower jaw enlarged, but not canine-like; maxillary not reaching vertical from middle of pupil,  $\frac{1}{2}$  length of head; interorbital width less than  $\frac{1}{2}$  diameter of pupil; opercle short, its length being less than the diameter of the eye; pectorals nearly as long as head, a trifle more than length of ventrals, which scarcely reach vent; caudal much longer than head in adults, 7 or 8 scales before dorsal. Black spot on posterior part of spinous dorsal jet black, conspicuously ocellated with white; a black streak along lower margin of caudal, including several of the lower rays, and running from base to tip of fin; no dusky bars visible on sides in any of the types; fins dusky; membranes uniting outer rays of ventrals white instead of black, as in *chlamydes*; no black spot at base of caudal; branchiostegal membrane with a medial black streak; anal blackish. Numerous specimens from the northern part of the Gulf of California, at Albatross Stations 3031 and 3035, in 30 and 33 fathoms. This species may vary into the typical *chlamydes* but the material before us does not justify us in so identifying it. (Gilbert.) (*ocellatus*, ocellated.)

*Bollmannia ocellata*, GILBERT, Proc. U. S. Nat. Mus. 1891, 555, Gulf of California.

2568. *BOLLMANNIA CHLAMYDES*, Jordan.

Head  $3\frac{1}{2}$  (5 to  $5\frac{1}{2}$  in total); depth  $4\frac{1}{2}$  ( $6\frac{3}{4}$  to 7). D. VII-15; A. 15; scales in a longitudinal series about 28, 8 or 9 in a cross series at vent. Body rather robust, compressed; head large and heavy, its profile evenly curved; mouth very large, oblique, the lower jaw projecting; maxillary reaching to opposite pupil,  $2\frac{1}{4}$  to  $2\frac{3}{4}$  in head; teeth small, sharp, in several series, the outer, especially in lower jaw, somewhat enlarged; eye longer than snout,  $5\frac{1}{4}$  to 4 in head; interorbital area very narrow, concave, its least width about  $\frac{1}{3}$  of eye or almost equal to pupil; scales very large, eonoid; little reduced on breast and nape; about 8 before dorsal, where they are little smaller than on body; top and sides of head with large scales; scales on cheek in 4 rows; 2 rows on upper part of opercle; the scales on head lost in some of the specimens; dorsal spines slender, filamentous, fifth longest,  $1\frac{1}{2}$  in head; first 2 in head, last  $3\frac{1}{2}$  to 4; first soft dorsal ray  $2\frac{3}{4}$  in head, the antepenultimate longest and about equal to head; first anal ray equal to snout, the antepenultimate  $1\frac{1}{2}$  in head; middle caudal rays very long, somewhat more than  $\frac{1}{2}$  length of body; pectorals  $1\frac{1}{2}$  in head; ventrals  $1\frac{1}{4}$ . Color olivaceous, darkest above; scales with a few black dots, some of the posterior occasionally dark edged; sides with 8 or 10 obscure dusky vertical bars, which are narrower than the inter-

spaces  
with a  
dark i  
dark-e  
soft d  
dark-e  
some o  
dorsal  
edged  
specim  
2800 in

*Bollman*  
of Ca  
47' 30"

Head  
its slen  
tion. C  
diamete  
larger t  
*B. chlam*  
to  $3\frac{1}{2}$  in  
longest  
never b  
low, the  
when de  
front of  
duced as  
in spirit  
sal dusk  
rather la  
black an  
membran  
branchio  
specimen  
Station 2

*Bollmanni*  
near L

Head s  
slightly  
A. 14; lat  
*poma*, the  
posterior  
rays wher

spaces, and in some specimens wholly obsolete; snout bluish; opercles with a dark shade; lips, gular region, and anterior branchiostegals very dark in males; upper part of spinous dorsal darkest, with a few lighter dark-edged oval spots, a well-marked black blotch between last 2 spines; soft dorsal dusky, usually with about 3 well-developed rows of lighter, dark-edged oval spots; anal dusky, crossed by 2 narrow bluish streaks; some of the last rays occasionally with a few spots similar to those on dorsal; caudal, pectorals, and ventrals dusky, tinged with blue; ventrals edged with pale. Length  $4\frac{1}{2}$  inches. West coast of Colombia. Many specimens of this abundant species were dredged at *Albatross* Stations 2800 in 7 fathoms and 2805 in  $51\frac{1}{2}$  fathoms. (*χλαμύδης*, cloaked.)

*Bollmannia chlamydes*, JORDAN, Proc. U. S. Nat. Mus. 1889, 164, Pacific Ocean, off coast of Colombia, Station 2800,  $8^{\circ} 51' N.$ ,  $79^{\circ} 41' 30'' W.$ , and Station 2805,  $7^{\circ} 56' N.$ ,  $79^{\circ} 41' 30'' W.$  (Type, No. 41158, U. S. Nat. Mus. Coll. *Albatross*.)

2569. BOLLMANNIA MACROPOMA, Gilbert.

Head 3 to  $3\frac{1}{2}$ ; depth 5. D. VII-14; A. 14; scales 28. Characterized by its slender form, low fins, large opercle, and comparatively plain coloration. Caudal peduncle correspondingly slender, its least height equaling diameter of eye. Head very large and heavy; opercle conspicuously larger than in *B. ocellata*, agreeing in this respect more nearly with *B. chlamydes*; dentition as in other species of the genus; eye large,  $3\frac{1}{2}$  to  $3\frac{3}{4}$  in the head. Dorsal spines slender, comparatively little produced, the longest usually not reaching the base of the first ray of second dorsal, and never beyond the base of the second or third ray; soft dorsal and anal low, the posterior rays usually not reaching the rudimentary caudal rays when depressed, about  $\frac{1}{2}$  length of head; pectoral long, extending beyond front of anal; the ventrals to or nearly to vent; middle caudal rays produced as usual, varying in length; scales 8 to 10 in front of dorsal. Color in spirits, light brownish, the sides with 3 vertical dusky bars; spinous dorsal dusky, but without distinct black spot; caudal slightly dusky, with rather large elliptical light spots, as in *B. chlamydes*, the lower rays not black and no black spot at its base; ventrals blackish, including anterior membrane; second dorsal and anal dusky, without evident light spots; branchiostegal membranes sometimes slightly dusky, but not black. Many specimens from the Gulf of California just north of La Paz Bay, at *Albatross* Station 2996, in 112 fathoms. (Gilbert.) (*μακρός*, large; *πίσμα*, opercle.)

*Bollmannia macropoma*, GILBERT, Proc. U. S. Nat. Mus. 1891, 556, *Albatross* Station 2996, near La Paz, Lower California.

2570. BOLLMANNIA STIGMATURA, Gilbert.

Head short,  $3\frac{3}{4}$  in length; depth 5; least depth of caudal peduncle slightly less than diameter of eye; eye large,  $2\frac{1}{2}$  in head. D. VII-15; A. 14; lateral line 28. Dorsal spines filamentous, longer than in *B. macropoma*, the longest reaching base of fifth to seventh ray of second dorsal; posterior rays of second dorsal and anal often reaching base of median rays when depressed; pectorals not reaching beyond front of anal. Color

almost uniform light brownish; lips black, the fins only slightly dusky, the caudal with elliptical light spots; a roundish dusky spot at base of caudal; branchiostegal membranes not black. Many specimens from the northern part of the Gulf of California, at Albatross Stations 3016 and 3017, in 76 and 58 fathoms. This species agrees with *Bollmannia macropoma* in its elongate form, comparatively low fins, and in the absence of a black spot on the spinous dorsal. It differs conspicuously in the very short head and narrow opercle, and in the presence of a black spot at base of tail. The eye is also larger and the fins higher. None of the specimens shows dusky bars on the sides, a conspicuous feature in *B. macropoma*. (Gilbert.) (*στρυμνα*, spot; *οὐρά*, tail.)

*Bollmannia stigmatura*, GILBERT, Proc. U. S. Nat. Mus. 1891, 556, Gulf of California, Albatross Stations 3016, 3017.

### 817. ABOMA, Jordan & Starks.

*Aboma*, JORDAN & STARKS, Proc. Cal. Ac. Sci. 1895, 497 (*etheostoma*).

This genus, allied to *Microgobius*, is distinguished by the large, ctenoid scales, which cover the body; head naked, rather long, pointed in profile; the mouth moderate, not very oblique; teeth rather strong. Dorsal spines more than 6, none of them filamentous; soft dorsal and anal short; no flaps on shoulder girdle. Cranium with a slight median crest. (The name *Aboma* is used by the Mexicans in Sinaloa as synonymous with *goby*.)

- a. Scales very large, 26 or 27; profile not very steep, the snout rather pointed.  
 b. Sides with a jet-black lateral band; caudal with dark cross bars; maxillary 3 in head. ETHEOSTOMA, 2571.  
 bb. Sides with 4 oblique dark cross bars; a large dark spot at base of caudal; mouth larger, the maxillary 2 in head. LUCETIAE, 2572.  
 aa. Scales smaller, about 37; profile very steep, the snout rounded; sides with numerous pale cross bands with darker spots. CHIQUITA, 2573.

### 2571. ABOMA ETHEOSTOMA, Jordan & Starks.

Head  $3\frac{1}{2}$ ; depth 5. D. VIII-11; A. 10; scales 26; longest dorsal spine  $1\frac{1}{2}$  in head; eye 3; snout 4; maxillary 3. Body long and low, moderately depressed and pointed forward. Scales large, ctenoid behind, none on head, those on nape and belly much reduced. Mouth moderate, terminal, moderately oblique; the maxillary reaching middle of pupil; jaws subequal, or the lower a little the longer; teeth rather strong. No flaps on shoulder girdle. Cranium with a slight median crest. Interorbital ridge not hollowed out; skull not abruptly widened behind. Color olivaceous, side with a very broad jet-black lateral band, 3 times interrupted by silvery; caudal white, with 4 < shaped bands, growing progressively fainter behind; pectoral mottled gray, with a jet-black oblique crescent toward its base surrounding a large yellow spot; side of head with 4 round gray spots separated by black, the largest below eye, with a black streak before it; first dorsal jet-black, second mottled, the produced spine with yellowish; ventral and anal pale. A single small specimen,  $1\frac{1}{2}$  inches long,

found in the mud on a shallow bottom in the Astillero at Mazatlan. (*Etheostoma*, a darter, which this species strongly resembles.)

*Aboma etheostoma*, JORDAN & STARRS, Proc. Cal. Ac. Sci. 1895, 498, pl. 50, Mazatlan. (Coll. Hopkins Expedition to Mazatlan.)

2572. ABOMA LUCRETIE (Eigenmann & Eigenmann).

Head  $3\frac{1}{2}$  in length; depth  $5\frac{1}{2}$ . D. VII-10; A. 12; scales 28-8. Body slightly compressed posteriorly; head little wider than high; eye placed high, its diameter equaling length of snout,  $4\frac{1}{2}$  in head; profile little decurved; mouth large, oblique; maxillary extending below posterior margin of orbit, 2 in head; intermaxillary anteriorly on a level with center of pupil; teeth all recurved, large, those of upper jaw in a narrow band; teeth of outer and inner series enlarged, those of lower jaw similar, largest in front. No dermal flaps on shoulder girdle. Scales large, very weakly ctenoid, becoming cycloid and very much crowded above and below pectoral; head, breast, and anterior part of nape naked. As seen through a lens, these regions seem to be covered with minute embedded scales; this effect is, no doubt, due in light reticulations on a darker ground. Dorsal spines slender, not filamentous; caudal pointed, 3 in length of body; ventrals  $1\frac{1}{2}$  in head; pectorals longer than head. Color light brownish, with 4 oblique dark cross bars as wide as interspaces; 4 narrower transverse bars on nape and back; a large dusky spot at base of caudal; upper half of base of pectoral black; a black spot on opercle, margined below and behind with silvery; fins dusky. Pearl Island, Gulf of Panama; only 1 specimen known. (Named for Mrs. Lucretia M. Smith of San Diego, mother of Mrs. Eigenmann.)

2573. ABOMA CHIQUITA (Jenkins & Evermann).

Head  $3\frac{1}{2}$  to  $3\frac{3}{4}$ ; depth  $4\frac{1}{2}$  to  $4\frac{3}{4}$ . D. VII-11; A. 10; eye  $4\frac{1}{2}$  in head in adult, 4 in young; scales 67-17. Body rather stout, compressed; head short, somewhat depressed, widened behind orbits; snout short and narrowly rounded; profile in front of eye very steep, less so to occiput, and nearly straight from there to caudal fin; eyes moderate, well up; interorbital space very narrow, less than eye; greatest width of head equaling greatest depth of body. Top of head, opercles, and space in front of dorsal naked, rest of body covered with small, strongly ctenoid scales, which increase in size upon the caudal peduncle. Spinous dorsal with its first spine filamentous in adult, much longer than head and reaching middle of soft dorsal (this filament wanting in young; distance from snout to origin of spinous dorsal a little more than  $\frac{1}{2}$  distance to base of caudal; second dorsal but slightly separated from spinous, its origin about midway of total length of fish; anal of about the same shape and size as soft dorsal, but beginning a little behind it; pectorals tapering, about equaling head in length, their tips not reaching origin of anal, but to origin of soft dorsal; ventrals united, free from belly, inserted behind pectorals, but their tips not reaching tips of pectorals. Teeth apparently in a single series, small and weak. Ground color pale yellowish, thickly mottled with fine pum-



tulations of dark; about 7 pretty well-defined larger spots of dark brown along middle of side; 8 or 9 faint cross bars of lighter, a number of small light spots scattered irregularly over the sides; head dark; dorsal, anal, and ventral fins covered with fine black points; in some specimens the dorsals and anal quite dark; pectorals plainer; caudal similar to ventrals; "the cranium is depressed and flattish behind the orbits, without distinct median keel on occiput or on interorbital area. The form of the head is as in typical *Gobius*, the occiput abruptly widened behind the eyes; the ridges also similar, the orbital ridge bounding the orbit behind as well as above the eye and joining the temporal ridge laterally." (Gilbert MS.) Length 1 to 2 inches. Gulf of California; abundant. The original description from young examples, here corrected in accordance with Dr. Gilbert's notes on many adults taken by him at La Paz. (Spanish, *chiquito*; a diminutive of *chico*, a little one.)

*Gobius chiquita*, JENKINS & EVERMANN, Proc. U. S. Nat. Mus. 1888, 146, Guaymas, Sonora. (Type, No. 39634. Coll. Jenkins & Evermann.)

### 818. MICROGOBIUS, Poey.

*Microgobius*, POEY, Enumeratio, 127, 1875 (*signatus*).

Dorsal spines 7 or 8; scales very small, cycloid or weakly ctenoid, the body scaled anteriorly as well as posteriorly, the head naked, the nape, belly, and breast usually so. Inner edge of shoulder girdle without fleshy processes; body more or less compressed; mouth large, very oblique; the lower jaw conspicuous, teeth strong; interorbital groove with or without a median ridge. Vertebrae 11-15 or 16. (*μικρός*, small; *Gobius*.)

a. Scales about 42. Body elongate, moderately compressed, the depth 4 to 5 in length; head long and large, rather sharp in profile, 3 to 3½ in body; eye longer than snout, 4 in head; mouth large, very oblique, the lower jaw strongly projecting; maxillary 1½ to 2½ in head, extending to opposite middle of eye, or much beyond front of orbit; teeth in few series, the outer very long and slender, curved, the lower longest, none canine-like; scales small, some of them with short, thick teeth, those of anterior part of body not well developed; dorsal spines more or less filamentous, the third and fourth or fourth and fifth sometimes with long filaments; caudal pointed, about as long as head. Grayish olive, with rather sharply-defined markings of darker brown overlaid with orange in life; head with a pale bluish or gilt stripe from maxillary backward across suborbital region to upper edge of gill opening; another pale gilt streak from snout along lower part of eye, another from angle of mouth upward and backward; rest of head dark; opercle with an oblique blackish bar; top of head and nape with dark marbling surrounded by paler reticulations; back with a series of black cross blotches mostly separated on the median line; 2 narrower dark vertical bars behind pectoral; middle line of side posteriorly with longitudinally oblong black blotches; besides these, numerous other blotches not regularly arranged; first dorsal with 2 or 3 oblique black bands; second dorsal pale, with about 4 series of black dots; caudal spotted with black; pectoral yellowish; ventral black, its center yellowish (male); anal pale. D. VII-15; A. 16 or 17. OULOSUS, 2574.

aa. Scales about 50; snout not pointed; depth 5½ in length; mouth large, the maxillary 2½ in head; teeth strong. Color yellowish, much dotted, but without bars.

EULPIS, 2575.

aaa. Sca  
b

bb

Head 3  
15; A. 16  
ately com  
large, ver  
in head,  
outer very  
Body enti  
naked; se  
rior part  
tous, the  
long as he  
flattened  
skull. Do  
meeting i  
grooved,

aaa. Scales 65 or more.

b. Caudal fin more than  $\frac{1}{3}$  ( $\frac{2}{3}$ ) length of body. Scales very small, cycloid, deciduous. Body elongate, much compressed, highest in front of ventrals, tapering regularly to the very narrow, short caudal peduncle; greatest depth  $4\frac{1}{2}$  in length; head  $3\frac{1}{2}$ . Head compressed, much higher than wide; snout very short, acute, preorbital not as wide as pupil; mouth terminal, very wide and oblique; jaws equal; maxillary reaching vertical from middle of orbit, 2 in head. Outer series of teeth enlarged. Eye 3 in head. Dorsals closely contiguous; spines very slender, the fifth slightly produced and filamentous; pectorals as long as head. Head and body translucent, overlaid by brilliant green luster, formed by minute, close-set green points; 3 conspicuous translucent bars wider than the interspaces, crossing body close behind head; head with 2 brilliant narrow blue and green lines running obliquely across cheek below eye; dorsal whitish, with 2 or 3 lengthwise series of large reddish-brown spots; spinous dorsal blackish at base, upper caudal rays marked with red, the lower portion of caudal and most of the anal fin blackish, anal whitish at base, the anterior rays tipped with white. In spirits, body dusted with dark points; 2 light cross bars toward head; lower part of caudal and anal black. D. VII-16; A. 15.

THALASSINUS, 2576.

bb. Caudal fin less than  $\frac{1}{3}$  length of body. Scales small, cycloid, embedded. Body very much compressed, more or less elongate, greatest depth at ventrals 4 (female) to  $6\frac{1}{2}$  (male) in length; head  $3\frac{1}{2}$  to 4. Head much compressed, much deeper than wide. Snout very short, acute, the anterior profile not recurved, not steep; preorbital not as wide as pupil; mouth very large, very oblique or almost vertical; maxillary extending to below pupil, 2 in head (in male,  $2\frac{1}{2}$  in female). Lower jaw projecting, the teeth of the outer series enlarged, recurved. Eye  $3\frac{1}{2}$  to 4 in head. Dorsals contiguous, spines very fine, produced in filaments, the third highest, a little longer than head; second dorsal and anal high. Head and nape naked. In the female the depth is greater, mouth less oblique, smaller; profile from spinous dorsal oblique. First dorsal spine highest,  $3\frac{1}{2}$  in length. Ventrals much shorter than in males. Dark gray; female with a short bright blue bar bordered by blackish above pectorals; a blotch of sky blue and orange below eye; fins dusky, the ventrals pale in female, dusky in males. Males with the body plain bluish gray. D. VII-17 to 20; A. 18 to 21; scales 68 to 70.

SIGNATUS, 2577.

2574. MICROGOBIUS GULOSUS (Girard).

Head 3 to  $3\frac{1}{2}$ ; depth 4 to 5; eye 4 in head, longer than snout. D. VII-15; A. 16 or 17; scales about 42; vertebrae 11 + 15. Body elongate, moderately compressed; head long and large, rather sharp in profile; mouth large, very oblique, the lower jaw strongly projecting; maxillary  $1\frac{1}{2}$  to  $2\frac{1}{2}$  in head, extending to opposite middle of eye. Teeth in few series, the outer very long and slender, curved, the lower longest, none canine-like. Body entirely scaled, except the nape, belly, breast, and head, which are naked; scales small, some of them with short thick teeth, those on anterior part of body not well developed. Dorsal spines more or less filamentous, the third to fifth sometimes with long filaments; caudal pointed, as long as head. Ventrals as long as pectorals, which are  $1\frac{1}{2}$  in head. Skull flattened behind, with a median ridge extending from eyes back to end of skull. Double crests bordering skull in front and on sides, the inner ones meeting in front of median crest. Interorbital very narrow and deeply grooved, with a median ridge. Frontal bones very thin and fragile.

Teeth on each jaw in narrow bands, all alike. Coloration in life, light grayish olive, with rather sharply defined markings of darker brown; head with a pale bluish stripe from behind the angle of the mouth upward and forward parallel with the gape to below front of eye, then turning abruptly backward across suborbital region to upper edge of gill opening; another pale streak from snout along lower part of eye; between this and the first streak a dusky area; below the first-mentioned streak a dusky region on cheek; opercle with an oblique blackish bar; top of head with dark marblings surrounded by paler reticulations; back with a series of black cross blotches, mostly separated on the median line; 2 narrow vertical dark bars behind pectoral; middle line of side posteriorly with longitudinally oblong black blotches; besides these numerous other blotches not regularly arranged; first dorsal with 2 or 3 oblique black bands; second dorsal pale, with about 4 series of black dots; caudal spotted with black, pectoral yellowish, ventral black, its center yellowish; anal pale; lower side of head pale; jaws dusky. Coast of Florida to Texas, in sandy or weedy bays, common north to Indian River. A strongly marked species with no near relative among our other gobies. The specimens here described from Pensacola. (*gulosus*, large-mouthed.)

*Gobius gulosus*, GIRARD, Proc. Ac. Nat. Sci. Phila. 1858, 169, Indianola, Texas; GIRARD, U. S. and Mex. Bound. Surv., Zool., 26, 1859; JORDAN & GILBERT, Synopsis, 634, 1883.

*Lepidogobius gulosus*, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1882, 294; JORDAN & GILBERT, Synopsis, 945, 1883.

*Microgobius gulosus*, JORDAN & EIGENMANN, l. c., 505.

2575. MICROGOBIUS EULEPIS, Eigenmann & Eigenmann.

Head 4 in length ( $5\frac{1}{2}$  in total); depth  $5\frac{1}{2}$  (7). D. VII-15; A. 16; scales 50-14. Body elongate, scarcely compressed; head slightly higher than wide, the depth  $1\frac{1}{2}$  in its length; eye large, longer than snout,  $3\frac{1}{2}$  in head; snout 5 in head, rather broad, not pointed as in *M. thalassinus*; preorbital narrower than pupil; mouth very oblique, maxillary not extending beyond anterior margin of pupil,  $2\frac{1}{2}$  in head; teeth in upper jaw in a very narrow band, slightly enlarged in outer series, largest toward angle of mouth; teeth of lower jaw in a similar band, some of outer ones in front long and slender. Scales cycloid, rather large, crowded anteriorly, regularly arranged, not embedded as in *M. signatus*, not deciduous as in *M. thalassinus*; breast, nape, and region along spinous dorsal naked. First dorsal spine equidistant from tip of snout and first anal ray; longest dorsal spine  $1\frac{1}{2}$  in head; caudal fin about 4 in body; ventral not reaching vent, equaling length of head, the basal membrane  $\frac{1}{2}$  of its actual length; pectoral equaling length of head. Color yellow or very light brown, dotted with minute dark points above; scales along back with a dark margin; head and nape with minute points; spinous dorsal transparent, a marked black spot on upper part of membrane between fourth and fifth dorsal spines; other fins plain; a light vertical bar on posterior margin of preopercle;

no other  
from a sp

*Microgobius*  
Monro

Head 3  
elongate,  
regularly  
liar, tran  
Head com  
preorbital  
the jaws e  
of head; t  
canine-like  
series seen  
verify it w  
the diamet  
spines very  
(in our spe  
late, very l  
torals as lo  
membranes  
front of ar  
small eye  
as they hav  
Head and  
formed by c  
toward the  
able on eam  
interspaces  
liant narrow  
eye; opercl  
sals whitish  
spinous dor  
lower portio  
at base, the  
pectorals tr  
points; 2 li  
black. Coa  
long (No. 2  
Charleston  
sea.)

*Gobius thalass*  
bor, South  
*Lepidogobius t*  
*Microgobius th*

no other bars or stripes anywhere. Fortress Monroe, Virginia; known from a specimen  $1\frac{1}{2}$  inches long. ( $\epsilon\psi$ , well;  $\lambda\epsilon\pi\iota\zeta$ , sealed.)

*Microgobius eulepis*, EIGENMANN & EIGENMANN, Proc. Cal. Ac. Sci. 1888, 69, Fortress Monroe, Virginia. (Type, No. 27123, M. C. Z. Coll. Mrs. C. N. Willard.)

2576. MICROGOBIUS THALASSINUS, Jordan & Gilbert.

Head  $3\frac{1}{2}$  in length; depth  $4\frac{1}{4}$ . D. VII-16; A. 15; eye 3 in head. Body elongate, much compressed, highest in front of ventrals, thence tapering regularly to a very narrow, short caudal peduncle; the body with a peculiar, translucent, fragile appearance, common also to *Z. emblematicus*. Head compressed, much higher than wide; snout very short, acute, the preorbital not as wide as pupil; mouth terminal, very wide and oblique, the jaws equal; maxillary reaching vertical from middle of orbit,  $\frac{1}{2}$  length of head; teeth in a narrow band in each jaw, the outer series enlarged, canine-like (under a microscope the band of small teeth behind the outer series seems evident, but the size of our specimens does not enable us to verify it with certainty); eyes placed high, separated by a narrow ridge, the diameter about  $\frac{1}{2}$  length of head. Dorsals very closely contiguous; spines very slender, the fifth slightly produced and filamentous, reaching (in our specimens) to base of third soft ray when depressed; caudal lanceolate, very long and pointed, the middle rays produced,  $2\frac{2}{3}$  in body; pectorals as long as head; the upper rays not silk-like; ventrals with basal membranes well developed; the fin long, reaching to or slightly beyond front of anal, somewhat longer than head. Body covered with rather small cycloid scales; head naked; the scales very readily deciduous; as they have in our specimens mostly fallen off, the count can not be given. Head and body translucent, overlaid by brilliant green luster, which is formed by exceedingly minute close-set green points; the luster is intense toward the head, where it assumes a blue tint, and becomes hardly noticeable on caudal peduncle; 3 conspicuous translucent bars, wider than the interspaces, crossing body immediately behind head; head with 2 brilliant narrow blue or green lines running obliquely across cheek below eye; opercle with greenish luster; branchiostegal membrane white; dorsals whitish, with 2 or 3 lengthwise series of large reddish-brown spots; spinous dorsal blackish at base; upper caudal rays marked with red, the lower portion of caudal and the most of the anal fin blackish, anal whitish at base, the anterior rays tipped with brilliant white; ventrals light buff; pectorals translucent. In spirits, the body appears dusted with dark points; 2 light cross bars toward head; lower part of caudal and anal black. Coast of South Carolina; two specimens, the largest  $1\frac{1}{2}$  inches long (No. 29674, U. S. Nat. Mus.), were taken in muddy tide pools in Charleston Harbor. ( $\theta\alpha\lambda\alpha\sigma\sigma\iota\nu\acute{o}\varsigma$ , thalassinus, sea-green;  $\theta\acute{\alpha}\lambda\lambda\alpha\sigma\sigma\alpha$ , the sea.)

*Gobius thalassinus*, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1882, 612, Charleston Harbor, South Carolina. (Coll. C. H. Gilbert.)

*Lepidogobius thalassinus*, JORDAN & GILBERT, Synopses, 947, 1883.

*Microgobius thalassinus*, JORDAN & EIGENMANN, l. c., 505.

## 2577. MICROGOBIUS SIGNATUS, Poey.

Head  $3\frac{1}{2}$  to 4; depth 4 (female) to  $6\frac{1}{2}$  (male); eye  $3\frac{1}{4}$  to 4. D. VII-17 to 20; A. 18 to 21; scales 68 to 70; vertebrae 14 + 15. Body very much compressed, more or less elongate; head much compressed, deeper than wide; snout very short, acute, the anterior profile not decurved, not steep; preorbital not as wide as pupil; mouth very large, almost vertical; maxillary extending to below pupil, 2 in head in male,  $2\frac{1}{4}$  in female; lower jaw projecting; teeth of the outer series enlarged and recurved. Dorsals contiguous, spines very fine, produced in filaments, the third longest, a little longer than head; second dorsal and anal high. Scales as in *M. gulosus*. Skull rounded, very fragile; a median crest which is highest between eyes; lateral crests developed, the inner ones meeting above posterior part of eye; interorbital comparatively broad, the median crest ending above anterior part of the orbit. Teeth in each jaw in 2 or 3 series; outer series of the upper jaw enlarged and recurved, the inner ones minute; outer series of lower jaw smaller than those of upper jaw, the one nearest angle of mouth an enlarged canine. Dark gray; female with a short bright blue bar, bordered by blackish above pectoral; a blotch of sky blue and orange below eye; fins dusky, the ventrals pale in female; males with the body plain bluish gray. The sexual differences in this species are very strongly marked. West Indies, in salt water; common in Cuba; one of the smallest gobies, barely 2 inches long. Here described from Havana examples collected by Dr. Jordan. (*signatus*, marked.)

*Microgobius signatus*, POEY, Enumeratio, 127, pl. 5, fig. 3, 1875, Cuba (Type in M. C. Z. Coll. Poey); JORDAN, Proc. U. S. Nat. Mus. 1886, 49; JORDAN & EIGENMANN, l. c., 505.

## 819. ZALYPNUS, Jordan &amp; Evermann.

*Zalypnus*, JORDAN & EVERMANN, Check-List Fishes, 459, 1896 (*emblematicus*).

This genus differs from *Microgobius* in having the anterior half of the body naked. Soft dorsal and anal long, of 16 or 17 rays. Two species known. (*ζάλη*, surf; *ὑπνος*, slumber.)

a. Scales 48; shoulder with a round black spot; none of the dorsal spines elongate.

CYCLOLEPIS, 2578.

aa. Scales 65; a silvery cross bar behind pectorals; some of the dorsal spines usually elongate.

EMBLEMATICUS, 2579.

## 2578. ZALYPNUS CYCLOLEPIS (Gilbert).

D. VII-16; A. 17; scales 48. Body somewhat elongate, compressed, the mouth very large, narrow, and oblique; maxillary produced beyond the rictus for a distance equaling  $\frac{3}{8}$  diameter of orbit, reaching vertical from posterior margin of pupil,  $1\frac{1}{3}$  in head; snout short, 5 in head; eye larger,  $3\frac{1}{4}$  in head; interorbital width  $\frac{1}{2}$  orbit; teeth in upper jaw in 2 series, the outer enlarged and distant; in lower jaw apparently in a single series, similar to outer series of upper jaw, with 2 stronger canines anteriorly. Inner edge of shoulder girdle without fleshy prominences. Dorsal spines

7, none of soft ray; rays reaching caudal lobe in our specimens small, abundant. Color in spot on side, resembling *Zalypnus*. A 7 fathoms.

*Microgobius*  
Lower

Head  $3\frac{1}{2}$  of body compressed, rather broader than its length. Lower jaw forming a curved. Extremely slender; head a naked strip reaching to dorsal and compressed near longer than dots; sides pairs of gill large as pectorals and bars along longitudinal dorsal; those about 3 series low, dusky; lower end of thus crossing olive, with known only dorsal.)

*Gobius emblematicus*  
*Lepidogobius*

7, none of them elongate, the membrane of last spine reaching base of first soft ray; soft anal rays of moderate height,  $1\frac{1}{2}$  in head, the tips of last rays reaching base of caudal, the fin similar to soft dorsal but lower; caudal long, apparently rounded posteriorly, longer than head (mutilated in our specimen); ventrals and pectorals reaching vent. Scales cycloid, small, absent on belly, nape, and on sides in front of fourth dorsal spine. Color in spirits, light olive, the fins dusky; a conspicuous round black spot on shoulder,  $\frac{1}{2}$  size of eye, its posterior margin denser black. Resembling *Zalypnus emblematicus*, differing in its larger scales and different coloration. A single specimen, about 2 inches long, from Lower California, in 7 fathoms. (Gilbert.) (κύκλος, circle, cycloid; λεπής, scale.)

*Microgobius cyclolepis*, GILBERT, Proc. U. S. Nat. Mus. 1891, 74, Albatross Station 3020, Lower California.

2579. ZALYPNUS EMBLEMATICUS (Jordan & Gilbert).

Head  $3\frac{2}{3}$ ; depth 5. D. VII-16; A. 17; scales about 65. Anterior part of body naked; teeth of upper jaw in one series; body elongate, compressed, heaviest forward; depth 5 in length; head  $3\frac{2}{3}$ ; snout short, rather broad, acute in profile; mouth terminal, very oblique; gape wide, its length nearly  $\frac{1}{2}$  head; maxillary reaching to opposite middle of pupil; lower jaw projecting. Teeth in lower jaw partly in 2 series in front, forming a single row laterally; anterior teeth in both jaws strong, incurved. Eyes very large, about  $\frac{1}{3}$  of head; snout less than orbit. Scales extremely small, cycloid, scarcely increasing in size toward caudal peduncle; head and anterior part of body to front of dorsal fin naked; a narrow naked strip along base of anterior  $\frac{1}{2}$  of spinous dorsal. Dorsal spines very slender and weak, some of the middle ones usually prolonged, sometimes reaching nearly to the base of caudal, sometimes little elevated; second dorsal and anal similar to each other, the rays high, the last when depressed nearly reaching to the base of caudal; caudal pointed, a little longer than head. Light olivaceous; above thickly punctate with pale dots; sides very thickly covered with golden-green specks; back with 6 pairs of golden-green spots on each side of the dorsal fin, each nearly as large as pupil; sides of head and anterior half of body with wide streaks and bars alternately of purplish blue and golden bronze; those on cheek longitudinal; those on opercle extending obliquely upward and backward, those on body vertical; first dorsal dusky, second dorsal with about 3 series of light-blue spots; anal pale; caudal yellowish green below, dusky above, a very conspicuous narrow bright-red streak from the lower end of the base to the tip of the fifth or sixth ray from the bottom, thus crossing the rays obliquely; ventrals bluish. In spirits, plain light olive, with a silvery cross bar behind pectorals. Length  $3\frac{1}{2}$  inches. Panama; known only from the original types. (ἔμβλημα, a banner, from the high dorsal.)

*Gobius emblematicus*, JORDAN & GILBERT, Bull. U. S. Fish Comm. 1881, 330, Bay of Panama.  
*Lepidogobius emblematicus*, JORDAN & EIGENMANN, l. c., 505.

## 820. EUCYCLOGOBIUS, Gill.

*Eucyclogobius*, GILL, Proc. Ac. Nat. Sci. Phila. 1862, 279 (*newberryi*).

This genus is allied to *Lepidogobius*, differing chiefly in the naked head and short, chubby body; shoulder girdle with a few dermal flaps; opercle adnate to shoulder girdle from the angle upward; dorsal spines 6 or 7; soft dorsal short; scales all cycloid; cranium depressed behind the parietal region, somewhat excavated, the supraoccipital crest rather high, not extending so far forward as the orbit. Species small, in fresh or brackish waters of California. (*εὖ*, well; *κύκλος*, circle (cycloid); *Gobius*.)

## 2580. EUCYCLOGOBIUS NEWBERRYI (Girard).

Head  $3\frac{3}{4}$  to  $3\frac{1}{2}$ ; depth  $4\frac{1}{2}$  to  $5\frac{1}{2}$ . D. VI or VII\*—11; A. 10 or 11 (8 in one specimen, perhaps abnormal); scales about 60 to 70, too irregular for exact counting. Body moderately elongate, somewhat compressed, tapering posteriorly; head rounded above, its width  $2\frac{1}{2}$  in its length; mouth large, oblique, the maxillary reaching to or beyond posterior margin of orbit, 2 to  $2\frac{1}{2}$  in head; interorbital space wide, 4 to  $4\frac{1}{2}$  in head; snout bluntish, broad, a little longer than interorbital width; eye small, 5 in head; teeth present on both jaws, slender, canine-like, arranged in series, the outer row enlarged; caudal peduncle 3 to  $3\frac{1}{2}$  in head; gill slit about  $2\frac{1}{2}$  in head, its upper edge opposite or slightly above uppermost ray of pectoral; scales minute, cycloid, inconspicuous, wanting on head, nape, and fins; shoulder girdle with 2 or 3 small dermal flaps; dorsals separated by a narrow space; dorsal spines very slender; base of spinous dorsal  $2\frac{1}{2}$  to  $2\frac{3}{4}$  in head; anal similar to soft dorsal, its base about  $1\frac{1}{2}$  in head; caudal subtruncate,  $1\frac{1}{2}$  to  $1\frac{1}{4}$  in head; ventrals inserted under or slightly behind lower edge of base of pectorals,  $1\frac{3}{4}$  to 2 in head; pectorals  $1\frac{1}{2}$  to  $1\frac{3}{4}$  in head. Dark olivaceous, mottled with darker; head with some dusky markings; the sides and back with irregular dark markings as in species of *Ethcostominae*; dorsals distinctly mottled; the first 3 or 4 dorsal spines margined with paler; caudal with faint, broad, wavy cross bars, a faint spot at its base; anal dusky; ventrals yellowish, dusky in males; pectorals plain. Length about 2 inches. Streams of California, in small clear brooks near the sea; locally common in San Luis Obispo Creek, where the specimens here described were taken; probably confined to fresh waters. (Named for Dr. John Strong Newberry of Columbia College, then also on the U. S. Geological Survey.)

*Gobius newberryi*, GIRARD, Proc. Ac. Nat. Sci. Phila. 1856, 136, Tomales Bay (Coll. E. Samuels); GIRARD, Jour. Bost. Soc. Nat. Hist. 1857, 530, pl. 25, figs. 5 to 8; GIRARD, Pac. R. R. Surv., x, 128, 1858.

*Lepidogobius newberryi*, JORDAN & GILBERT, Synopsis, 637, 1883; JORDAN & EIGENMANN, l. c., 503.

\* Of the nine specimens examined from San Luis Obispo Creek, five have 7 dorsal spines and the other four 6. Girard gives the fin rays as D. VIII-13; A. 12; but we have seen no specimens either with 8 spines or 13 rays. Six specimens from Wadell Creek, Santa Cruz County, California, show the following fin variation: D. VI in 4; D. VII in 1; D. V (?) in 1; D. rays 10 in 4; D. rays 9 in 2; A. 10 in 2; A. 8 in 1; A. 9 in 3.

*Lepidogobius*  
*Cyclogobius*

This g  
small eye  
2 or 3 do  
little de  
*Gobius*, t  
abruptly  
(*Αεπίς*, so

Head 4  
long as d  
subinfurc  
space nar  
maxillary  
small, all  
set, in a b  
very much  
head, and  
of head se  
head; sof  
somewhat  
rusty red  
half of all  
"This spec  
snout, and  
more depre  
low median  
ous behind  
coast of Ne  
rather deep  
and sold in

*Gobius gracilis*  
*Gobius gr*  
*Gobius lepidu*  
*gracilis*;  
*Lepidogobius*  
opsis, 637.

*Gillichthys*, C  
Gillie, GÜNT  
*Saccostoma* (G  
name prec

Body mo  
embedded a

821. LEPIDOGOBIUS, Gill.

*Lepidogobius*, GILL, Ann. Lye. Nat. Hist. N. Y. 1859, 14 (*lepidus*).  
*Cyclogobius*, STEINDACHNER, S. B. K. Ak. Wiss. Wien, XLII, 1860, 284 (*lepidus*).

This genus contains small gobies with the head and body covered with small cycloid scales; dorsal spines 7; inner edge of shoulder girdle with 2 or 3 dermal flaps; interorbital groove with the median ridge of skull little developed; body elongate, subterete; otherwise essentially as in *Gobius*, the skull nearly as in *Gillichthys*, with a median keel and not abruptly widened behind the eye. Pacific Ocean; not entering rivers. (Αεπίς, scale; *Gobius*.)

2581. LEPIDOGOBIUS LEPIDUS (Girard).

Head  $4\frac{1}{2}$ , regularly conical; depth 7; eye 4, equal to snout, twice as long as deep. D. VII-16 to 18; A. 15; scales about 86. Body elongate, subfusiform, little compressed. Snout not obtuse in profile; interorbital space narrow, about equal to diameter of pupil. Mouth rather large, maxillary reaching to below posterior edge of pupil,  $2\frac{1}{2}$  in head; teeth small, all similar, those of upper jaw in 2 or 3 series, those of lower jaw close set, in a broad band. Body covered with small cycloid scales which are very much reduced anteriorly, especially on the nape; cheeks, sides of head, and upper posterior part of opercles covered with small scales; top of head scaly to eye; breast scaled. Dorsal spines weak, the longest 2 in head; soft dorsal low, none of the rays reaching caudal; caudal long, somewhat pointed. Color very pale olive, with roundish blotches of rusty red on back and sides; vertical fins mottled with reddish; distal half of all fins and under side of head blackish, especially in the males. "This species is remarkable for numerous lines of papillæ on mandible, snout, and sides of head. The occipital region of the skull is somewhat more depressed than in *Gobius saporator*, and has much lower ridges. A low median carina is present and the low supraorbital ridges are continuous behind the eyes with the temporal crests." (Gilbert MS.) Pacific coast of North America, from Vancouver Island to Lower California; in rather deep water off San Francisco Bay; often scined in great numbers and sold in restaurants as "whitebait." (*lepidus*, pretty.)

*Gobius gracilis*, GIRARD, Proc. Ac. Nat. Sci. Phila. 1854, 134, San Francisco; preoccupied by *Gobius gracilis*, JENYNS.  
*Gobius lepidus*, GIRARD, Pac. R. R. Surv., x, 127, pl. 25a, figs. 5 and 6, 1858; substitute for *gracilis*; GÜNTHER, Cat., III, 78, 1861.  
*Lepidogobius gracilis*, GILL, Ann. Lye. Nat. Hist. N. Y. 1859, 14; JORDAN & GILBERT, Synopsis, 637, 1883; JORDAN & EUGENMANN, l. c., 502.



822. GILLICHTHYS, Cooper.

*Gillichthys*, COOPER, Proc. Cal. Ac. Sci. 1863, 109 (*mirabilis*).  
*Gillia*, GÜNTHER, Zool. Record 1864, 157 (*mirabilis*); name preoccupied.  
*Saccostoma* (GUICHENOT MS.) SUAVAGE, Bull. Soc. Philom. Paris 1882, 171 (*gulosum*); name preoccupied.



Body moderately elongate, compressed, covered with small, cycloid, embedded scales; belly and head naked. Scales of the young more or







**IMAGE EVALUATION  
TEST TARGET (MT-3)**



**Photographic  
Sciences  
Corporation**

23 WEST MAIN STREET  
WEBSTER, N.Y. 14580  
(716) 872-4503

18  
19  
20  
22  
25  
28  
32  
36

10  
11  
12  
13  
14  
15  
16  
17

less ciliated. Eyes small, almost superior. Gape wide, the maxillary in the adult inordinately developed, prolonged backward to the base of the pectorals, its posterior part a cartilaginous expansion, connected to an expansion of the skin of the lower jaw, thus forming a channel backward from the mouth, almost exactly as in *Neoclinus* and *Opisthognathus*, genera otherwise very different. Teeth small, even, in broad bands. Skull in adult with a strong median keel, not abruptly widened behind the eye, triangular behind; young with the keel obsolete. Dorsal fins 2, the second high, the first of 6 very weak spines, none of which is exerted; soft dorsal and anal short; caudal less rounded; pectorals large; isthmus broad. Singular little fishes, in brackish waters, burrowing in the mud; confined to the Pacific. (Named for Theodore Gill.)

a. Head moderately depressed; dorsal fins close together.

MIRABILIS, 2582.

aa. Head very broad and depressed; distance between dorsals  $\frac{1}{2}$  length of first dorsal.

DETRUSUS, 2583.

2582. GILLICHTHYS MIRABILIS, Cooper.

(LONG-JAWED GOBY.)

Head  $3\frac{1}{2}$ ; depth 5; eye 6 to 7; snout longer than eye, low, little decurved. D. VI-12; A. 10; vertebrae 15+17. Body stout, somewhat compressed behind, broad and depressed anteriorly; head broader than deep, its width  $1\frac{1}{2}$ , its depth 2 or more in its length; interorbital space greater than eye. Mouth very large; maxillary variable, extending to base of pectoral in adult, broadened behind; fold of lower lip extending its full length. Teeth all alike, small, fixed, and in bands, the band of the lower jaw broader than that of the upper. Scales small, cycloid, irregularly placed, largest from front of dorsal backward, decreasing in size anteriorly; head, breast, belly, and  $\frac{1}{2}$  of nape naked. Dorsal spines not filamentous, not as long as the soft rays which are little more than  $\frac{1}{2}$  depth of body; caudal broad, short, rounded; pectorals broad and rounded, longer than ventrals, 2 in head. Skull not abruptly widened behind eye, as in *Gobius*, being triangular posteriorly. No lateral ridges; a strong median keel; a short transverse crest behind orbit. Interorbital not deeply grooved, with a blunt median ridge. Orbit not bordered by any prominent ridges. Teeth in both jaws, close set, in bands, all alike. Dull olive, very finely marbled with darker; sides of head and maxillary finely punctuate; fins olive; belly yellowish. Length 8 inches. Pacific coast of North America, from San Francisco to Cerros Island; a most remarkable little fish; very abundant in the mud flats in shallow water along the California coast, burrowing in hole in the mud like a crawfish, and readily taking the hook baited with flesh or worm when dropped into the mouth of the burrow. (*mirabilis*, wonderful.)

*Gillichthys mirabilis*, COOPER, Proc. Cal. Ac. Sci. 1863, 100, San Diego Bay; LOCKINGTON, Amer. Nat. 1879; JORDAN & GILBERT, Synopsis, 630, 1883; JORDAN & EIGENMANN, l. c., 510; EVERMANN & JENKINS, Proc. U. S. Nat. Mus. 1891, 162.

*Gobius townsendi*, EIGENMANN & EIGENMANN, Proc. U. S. Nat. Mus. 1888, 463, San Diego; young.

Allied to  
more depressed  
the 2 dorsals  
D. VI-13;  
riorly but  
pectoral to  
and front  
the shortest  
in *G. mirabilis*  
differing from  
wing-like pectoral  
head; maxillary  
peduncle 2  
dorsal; length  
length of 10  
with dark pectoral  
probably due  
Gulf of California  
about 5 inches  
near the mouth  
abundant.  
depressed.)

*Gillichthys detrusus*  
shoe Bend,

*Quietula*, JORDAN

This genus  
presence of 2  
girdle. Maxillary  
cranium essential  
lagoons and r

Head  $3\frac{1}{2}$  (4  
50-18; B. 5.  
from shoulder  
than body, its  
diameter of eye  
of eye; mouth  
terior margin  
usually prolonged  
eye. Scales small  
verse. Teeth  
blunt, and curved

## 2583. GILlichTHYS DETRUSUS, Gilbert &amp; Scofield.

Allied to *Gillichthys mirabilis*, Cooper, differing in the broader and more depressed head, the larger anal fin, and greater distance between the 2 dorsals. Head  $3\frac{1}{2}$ ; depth 5; eye 7 in head; snout 4; interorbital  $5\frac{1}{2}$ . D. VI-13; A. 11 developed rays (10 in *G. mirabilis*); scales very fine anteriorly but becoming much larger posteriorly; about 75 scales from base of pectoral to caudal, and about 25 longitudinal rows between front of anal and front of second dorsal. The head is depressed, the frontals broad, the shortest distance across being contained in the head 8 times (11 times in *G. mirabilis*.) The postfrontals are small and project but very little, differing from *G. mirabilis*, where the postfrontals project into an elevated wing-like process. The width of the isthmus contained 3 times in the head; maxillary  $1\frac{1}{2}$  and mandible  $1\frac{3}{4}$  in head. Least depth of caudal peduncle  $2\frac{3}{4}$  in head. Distance between dorsals equal to  $\frac{1}{2}$  length of first dorsal; length of first dorsal  $2\frac{1}{2}$  in head; second dorsal  $1\frac{1}{2}$ ; anal 2 in head; length of longest pectoral ray  $1\frac{1}{2}$  in head. Color a very pale olive, some with dark punctulations about the head and fins. The pale coloration is probably due to their life in shallow water on bottom of pale sand. Gulf of California. The types and numerous other specimens, the longest about 5 inches long, were taken by Dr. C. H. Gilbert at Horseshoe Bend, near the mouth of the Colorado River, in Mexico, where they are quite abundant. These are numbered 3836 in L. S. Jr. Univ. Mus. (*detrusus*, depressed.)

*Gillichthys detrusus*, GILBERT & SCOFIELD, Proc. U. S. Nat. Mus., xx, 1897, 498, pl. 38, Horseshoe Bend, mouth of Colorado River. (Type, No. 48127. Coll. Gilbert & Alexander.)

## 823. QUIETULA, Jordan &amp; Evermann.

*Quietula*, JORDAN & EVERMANN, Proc. Cal. Ac. Sci. 1895, 839 (*y-cauda*).

This genus is closely related to *Gillichthys*, from which it differs in the presence of 2 or 3 cutaneous flaps on the inner edge of the shoulder girdle. Maxillary elongate, as in *Gillichthys*; scales rather small, cycloid; cranium essentially as in *Gillichthys*. Small gobies living in the mud of lagoons and river mouths. (A diminutive, from *quies*, quiet.)

## 2584. QUIETULA Y-CAUDA (Jenkins &amp; Evermann).

Head  $3\frac{1}{2}$  (4); depth 7 (8); eye  $3\frac{1}{2}$ . D. V-14 or 15; A. 15; scales about 50-18; B. 5. Body moderately elongate, compressed, narrowing regularly from shoulder girdle to caudal fin; head not greatly depressed, broader than body, its length 4 in body; snout rounded, short, about equal to diameter of eye; interorbital space narrow, not greater than  $\frac{1}{2}$  diameter of eye; mouth rather large, its gape extending nearly to vertical of posterior margin of orbit; maxillary somewhat variable in length, but usually prolonged behind eye for a distance nearly equal to diameter of eye. Scales small, cycloid, about 50 in longitudinal series, 18 in transverse. Teeth in a narrow band on premaxillaries and mandible, short, blunt, and curved slightly backward, most closely set and most numerous

ou premaxillaries. Shoulder girdle with 2 or 3 small cutaneous flaps on its inner edge. Fins moderate; dorsal of 5 spines and 16 soft rays, the spines unconnected with the rayed portion, the space between them about equal to  $\frac{1}{2}$  diameter of eye; the spines weak and flexible, their length  $\frac{1}{2}$  that of head; soft dorsal beginning at a point a little nearer end of snout than tip of caudal and extending nearly to caudal, its height about equal to that of spinous portion, the first few rays slightly graduated; anal having 15 rays and beginning a little behind origin of soft dorsal, the rays about equaling those of dorsal in length; pectorals moderate, inserted a little below axis of the body, their length greater than depth of body, their tips reaching a vertical from posterior part of spinous dorsal; ventrals united, but not adnate to belly, inserted slightly in front of pectorals and their tips not quite reaching those of pectorals. Ground color light; head and body pretty uniformly covered with dark punctulations; an irregular dark bar across occiput; breast and belly pale; a row of 9 or 10 small dark blotches along middle of side, the one at base of caudal plainest and having a shape something like the Greek letter T; about 6 dark blotches along median line of back; peritoneum dark. Length about  $1\frac{3}{4}$  inches. Pacific coast of North America, from Guaymas to Vancouver Island; excessively abundant from San Diego southward in mud flats; specimens recorded from Saanich Arm, Vancouver Island, San Diego, mouth of Colorado River, San Luis Gonzales Bay, St. Georges Bay, Concepcion Bay, Guaymas, and La Paz. It was at first confounded with the young of *Gillichthys mirabilis*, from which genus it differs in the presence of dermal flaps on the shoulder girdle.\* (caudal, tail, which has a Y-like mark.)

*Gillichthys y-cauda*, JENKINS & EVERMANN, Proc. U. S. Nat. Mus. 1888, 147, Guaymas, Sonora. (Type, No. 39637. Coll. Jenkins & Evermann.)

*Quietula y-cauda*, JORDAN & STARKS, Proc. Cal. Ac. Sci. 1895, 839.

*Gillichthys guaymasiae*, JENKINS & EVERMANN, Proc. U. S. Nat. Mus. 1888, 148, Guaymas, Sonora; young specimens  $2\frac{1}{2}$  inches long. (Type, No. 39627. Coll. Jenkins & Evermann.)

\* "The cranium is similar to that of *Gillichthys mirabilis*, the occiput being depressed, wedge-shaped, narrowed anteriorly with a blunt median carina, the supraorbital and temporal ridges not continuous behind the eye. As in *Gillichthys mirabilis*, the supraorbital ridges end in wing-like expansions immediately behind the interorbital space." (Gilbert MS.)

† *Gillichthys guaymasiae* is thus described: Head 3 ( $3\frac{1}{2}$  in total); depth 6 (7). D. V-14; A. 13; eye 5. Body quite slender, elongate, but little compressed; head long, narrow, not much widened behind the eyes, not depressed, forming  $\frac{1}{2}$  the length to base of caudal. Profile gently arched from snout to  $\frac{1}{2}$  the distance to dorsal fin, from there nearly straight to dorsal, and then gently curved to caudal peduncle; ventral outline nearly straight; a considerable prominence on the snout made by the enlarged end of the turbinal bone. Eye somewhat above the median line, not quite equaling the snout in length; interorbital space narrow,  $1\frac{1}{2}$  times in the eye. The maxillaries are much produced, in some specimens nearly reaching the gill openings, broadest at the middle, and tapering to a blunt point posteriorly; premaxillaries not protractile, but little movable at the symphysis, more than  $\frac{1}{2}$  as long as the maxillaries. Gill rakers 2 above the angle, 10 below, short and blunt, the first 4 the largest, those on the second arch but little developed. Teeth well developed, in a single series, on mandible and premaxillaries, all slightly curved backward. Tongue not so broad as in *Gillichthys mirabilis*, Cooper; it is gently rounded at the tip, which is free for a much greater length than in *Gillichthys mirabilis*. Peritoneum black or blackish, and the intestine short, but little longer than the head, and not at all convoluted. Scales small, embedded, and scarcely perceptible except on sides; no pores appear to be developed. First dorsal of fine flexible spines, distance of origin from snout  $2\frac{1}{2}$  length of body, and separated from the second dorsal by a distance but little greater than length of snout; second dorsal of 14 rays of nearly equal length, which equals the

*Thynnus*, J.

This g...  
presence...  
minute, e...  
rounded, ...  
iting muc...

Head 3

14 to 16; I...  
subconica...  
nearly str...  
of eyes. ...  
head,  $\frac{1}{2}$  in...  
below mid...  
broad ban...  
One, rarel...  
eyeloid, en...  
from tip o...  
dorsal spin...  
area about...  
caudal; ca...  
dorsal fin;...  
long. Pect...  
head and b...  
the punctu...  
hyaline at...  
with white...  
of black do...  
dal margin...  
dian portio...  
dotted with...  
dotted, its...  
ventrals mi...  
a large, co...  
blackish; l...

distance from...  
equaling leng...  
between the 2...  
to middle of t...  
length of fish...  
head; ventral...  
Color in life w...  
back, alternat...  
black with 2 d...  
ber of dark sp...  
marked being...  
about 4 series...  
dark spots or...  
less plain, esp...

## 824. ILYPNUS, Jordan &amp; Evermann.

*Ilypnus*, JORDAN & EVERMANN, Check-List Fishes, 460, 1896 (*gilberti*).

This genus is allied to *Clevelandia*, from which it differs chiefly in the presence of dermal flaps on the inner edge of the shoulder girdle; scales minute, embedded, cycloid; dorsal with 5 spines; occiput transversely rounded, without median keel; maxillary moderate. Small gobies, inhabiting mud flats. (*ἰλύς*, mud; *ἕπνος*, slumber.)

## 2585. ILYPNUS GILBERTI (Eigenmann &amp; Eigenmann.)

Head 3 to 3½ (3½ to 4 in total); depth 5 to 5½ (6 to 7). D. V-15 to 17; A. 14 to 16; B. 5; vertebrae 15 + 19. Form elongate, compressed. Head long, subconical, about as high as wide, its width 2½ in its length. Profile nearly straight from eyes to spinous dorsal, decidedly decurved in front of eyes. Eye entirely above the premaxillary level, 1 in snout, 4½ in head, ½ in interorbital. Mouth slightly oblique; maxillary extending to below middle of eye, lower jaw slightly included. Teeth villiform, in a broad band in each jaw, the outer series of lower jaw somewhat enlarged. One, rarely 2, dermal flaps on inner edge of shoulder girdle. Scales cycloid, embedded, very small; head, nape, and breast naked. Distance from tip of snout to insertion of spinous dorsal 2½ in length; highest dorsal spine about ¾ length of head; soft dorsal rays lower; interdorsal area about ½ orbital diameter; tip of last dorsal ray not reaching base of caudal; caudal broad and rounded when expanded; anal similar to soft dorsal fin; ventral fins large, nearly reaching vent in specimens 1¾ inches long. Pectorals usually shorter than ventrals. Color in life, and color; head and body with small rust-colored spots, which are dotted with black, the punctulations forming a more or less regular network; dorsal fins hyaline at base, bright rust-colored above, and rather broadly margined with white, everywhere black punctate except on margins; about 3 groups of black dots on each ray, giving a barred appearance to these fins; caudal margined with white, upper and lower parts of fin rust colored, median portion dark gray; about 5 wavy, rustlike, vertical bars; entire fin dotted with black except its margin; anal fin hyaline at base, sparsely dotted, its middle third jet-black, margined with white; pectorals and ventrals milky white, yellowish, sparingly black dotted and white edged; a large, conspicuous, metallic blue-black spot on opercle; top of head blackish; belly white or yellowish; chin and throat white, sometimes

distance from end of snout to middle of pupil; length of base of soft dorsal not quite equaling length of head; distance of posterior end from caudal fin equaling distance between the 2 dorsal fins. Origin of anal behind that of soft dorsal and a little posterior to middle of total length of fish; its base 1½ times in base of soft dorsal, or about 4 in length of fish to base of caudal fin; pectorals moderate, a little more than ½ length of head; ventrals inserted slightly behind the pectorals and about equaling them in length. Color in life whitish beneath, grayish or mottled above; 6 double white spots along the back, alternating with fine blackish areas; a white spot behind each eye on top of head; cheek with 2 dark bands extending obliquely backward and downward from eye; a number of dark blotches on opercles; about 7 dusky areas along the side, the last and most marked being upon the base of the caudal fin; dorsal fins finely marked lengthwise by about 4 series of small dark spots; caudal crossed by 5 or 6 wavy vertical bars of very fine dark spots or points; anal, pectorals, and ventrals plain. In alcohol these markings are less plain, especially the white and black areas upon the back. Length 2½ inches.

punctate. Young lighter, showing the reticulations, but the other markings faint or undeveloped. Length about  $2\frac{1}{2}$  inches. (Eigenmann.) "This species agrees with *Lepidogobius* in the presence of papillae on the inner edge of shoulder girdle. It differs decidedly in the shape of the occipital region of the cranium, which is transversely evenly convex as in *Clevelandia*; not abruptly widened behind the orbits, not continuous laterally with the temporal ridge as in *Gobius*, *Lepidogobius*, etc. From *Clevelandia* and *Gillichthys*, *Lepidogobius gilberti* differs in the presence of papillae on the shoulder girdle, and from *Gillichthys y-cauda* in the shape of the cranium." (Gilbert MS.) San Diego Bay and southward; found by Dr. Gilbert abundant at Magdalena Bay, at Concepcion Bay, and St. Georges Bay, in the Gulf of California. (Named for Charles Henry Gilbert, professor of Zoology in the Leland Stanford Junior University.)

*Lepidogobius gilberti*, EIGENMANN & EIGENMANN, Proc. U. S. Nat. Mus. 1888, 464, San Diego Bay. (Type, No. 40128, U. S. Nat. Mus. Coll. C. H. Eigenmann.)

### 825. CLEVELANDIA, Eigenmann & Eigenmann.

*Clevelandia*, EIGENMANN & EIGENMANN, Proc. Cal. Ac. Sci. 1888, 73 (*longipinnis*, EIGENMANN & EIGENMANN, = *rosæ*).

This genus is closely allied to *Gillichthys*, differing chiefly in the form of the skull, which is rounded above, strongly convex in transverse profile, perfectly smooth, without ridges or crests. Body long and slender; maxillary much produced, but not extending to the gill opening; mouth horizontal; dorsal spines 4 or 5, very weak; body covered with minute cycloid embedded scales; soft dorsal and anal long, each of 14 to 17 rays. (Named for Daniel Cleveland, esq., president of the San Diego Society of Natural History, a gentleman deeply interested in scientific matters.)

a. Caudal short, rounded; dorsal spines 5.

IOS, 2586.

aa. Caudal pointed, scarcely shorter than head; dorsal spines 4.

ROSÆ, 2587.

### 2586. CLEVELANDIA IOS (Jordan & Gilbert).

Head  $3\frac{1}{2}$  in length of body; depth 6. D. V-16; A. 14; eye  $6\frac{1}{2}$  in head; maxillary  $1\frac{1}{2}$ ; pectoral  $1\frac{1}{2}$ ; ventrals  $1\frac{1}{2}$ ; caudal  $1\frac{1}{2}$ ; base of soft dorsal 3 in length of body; base of anal  $3\frac{1}{2}$ . Body long and slender, compressed, the back not elevated; caudal peduncle moderately wide; head long, profile steep to within a short distance of the front of the eye, thence horizontal; mouth very large, not very oblique, the maxillary projecting to opposite the middle of the cheek; jaws subequal; teeth in narrow villiform bands; eye small, longer than wide, set high in head; interorbital space narrow, about as wide as eye. Body covered with very small cycloid scales, too small to count; spinous dorsal well separated from soft dorsal, the spines slender; soft dorsal the higher, its origin a little nearer base of caudal fin than tip of snout; anal about equal to soft dorsal in height, its origin a little behind first dorsal ray, ending at about the same comparative place as soft dorsal; ventrals inserted slightly behind pectorals, reaching midway between their base and front of anal; caudal short, its end rounded.

Color light  
form mot  
head black  
some dark  
caudal w  
waters.

off Port  
imperfect  
grammos

*Gobiosoma*  
cover  
JORDAN  
*Clevelandia*

Head 4

Body very  
riorly muc  
than snout

anteorbital  
extending  
anteriorly

in each jaw  
others. S

anterior p  
dorsal are  
2 $\frac{1}{2}$  in body

equals snou  
not extend  
head; vent

in head; v  
numerous

body; belly  
cle white;

blotch at t  
light areas

illaries pla  
dark bars;  
irregularly

Diego Bay  
Mr. and M  
species simi  
Eigenmann

*Clevelandia*  
*osoma long*  
*Clevelandia*  
R. S. Eige



Color light olivaceous, the cheeks and sides with many dark points which form mottlings; snout dark; a dark spot on upper part of opercle; top of head black; dorsals light, with 3 or 4 dark lines running across the rays; some dark spots on base of anal; pectorals crossed with dark wavy lines; caudal with about 5 irregular cross bars. Puget Sound and neighboring waters. Here described from 2 specimens, each 2 inches in length, dredged off Port Orchard by Mr. Edwin C. Starks. The original description is imperfect and partly incorrect, the single type, from the stomach of *Heterogrammos asper*, being in bad condition. (Iós, arrow.)

*Gobiosoma ios*, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1882, 437, Saanich Arm, Vancouver Island (Coll. Jordan & Gilbert); JORDAN & GILBERT, Synopsis, 948, 1883; JORDAN & EIGENMANN, l. c., 509.

*Clevelandia ios*, JORDAN & STARKS, Proc. Cal. Ac. Sci. 1895, 839, pl. 100.

2587. CLEVELANDIA ROSE, Jordan & Evermann.

Head 4 (4½ in total); depth 6½ (7). D. IV-16; A. 17; scales 70-18. Body very much elongate, slender; head long and slender, depressed anteriorly much as in *Lucius*; profile straight; eye moderate, slightly shorter than snout, 4½ in length of head; interorbital area about as wide as pupil; anteorbital area scarcely ½ diameter of eye; mouth large, maxillary extending much beyond orbit; lower jaw flat, slightly curved upward anteriorly; mouth very much as in *Lucius*; teeth all small, in narrow bands in each jaw; the outer ones of the upper jaw slightly larger than the others. Scales minute, slightly enlarged posteriorly; the margins plain, anterior part of the exposed area lengthwise striated; breast and antedorsal area naked. Distance from snout to insertion of first dorsal spine 2½ in body; the spines slender and short, 3 in head; interdorsal area equals snout and eye; dorsal rays slightly longer than spines, the last ray not extending halfway to caudal; caudal pointed, scarcely shorter than head; ventrals not reaching halfway to vent, 1½ in head; pectoral 1½ in head; vent slightly behind middle of body. Color light brownish; numerous darker spots of aggregated points along nape and upper half of body; belly white; head slightly darker than body; posterior edge of opercle white; an oblique silvery bar on the lower half of opercle, and a light blotch at the upper corner of opercle; cheek with black points; some light areas below eye; lower surface of head and posterior part of maxillaries plain; 2 dark bars on spinous dorsal; second dorsal with 3 or 4 dark bars; a curved black bar at base of caudal; remainder of caudal irregularly barred with dark; other fins plain. Length 1½ inches. San Diego Bay (Eigenmann & Eigenmann); at first incorrectly identified by Mr. and Mrs. Eigenmann with *Evermannia longipinne* (Steindachner), a species similar in habit but wholly scaleless. (Named for Mrs. Rosa Smith Eigenmann.)

*Clevelandia longipinnis*, EIGENMANN & EIGENMANN, Proc. Cal. Ac. Sci. 1888, 73; not *Gobiosoma longipinne*, STEINDACHNER.

*Clevelandia roseae*, JORDAN & EVERMANN, Proc. Cal. Ac. Sci. 1896, 229, San Diego. (Coll. R. S. Eigenmann.)

## 826. EVERMANNIA, Jordan.

*Evermannia*, JORDAN, Proc. Cal. Ac. Sci., IV, 1895, series 2, 592 (*zosterura*).

Body slender, compressed behind, entirely naked. Head long, slender. Snout rather pointed; mouth moderate, terminal, the maxillary more or less produced backward; teeth small and slender, the outer above slightly enlarged. Skull with a small median crest, not much widened behind. Interorbital space very narrow, channeled; no dermal flaps on shoulder girdle; first dorsal of 4 to 6 spines; second dorsal and anal long, of 14 or 15 rays. Caudal lanceolate. Ventrals formed as in *Gobius* and *Gobiosoma*. Size small, the sexes not colored alike. Species living in holes in sand and mud between tide marks. (Named for "my former student and later scientific associate, Dr. Barton Warren Evermann, now ichthyologist of the United States Fish Commission, in recognition of his work on the fishes of the Gulf of California."—Jordan.)

a. Head  $3\frac{1}{2}$ ; depth  $5\frac{1}{2}$ ; body and fins dotted.

LONGIPINNIS, 2588.

aa. Head 3 $\frac{1}{2}$ ; depth 6 $\frac{1}{2}$ ; vertical fins in males banded with black and with white edgings.

ZOSTEBURA, 2589.

## 2588. EVERMANNIA LONGIPINNIS (Steindachner).

Head  $3\frac{1}{2}$ ; depth 5 to  $5\frac{1}{2}$ . D. IV to VI-16 or 17; A. 16 or 17; snout slightly decurved in profile,  $3\frac{1}{2}$  in head; eye 6, greater than interorbital width. Body very slender. Mouth somewhat oblique, the jaws equal; maxillary extending beyond middle of head to a distance behind eye equal to diameter of eye. Teeth in each jaw in 2 series laterally and 3 in front, those of the outer series somewhat enlarged. Fins low, the longest dorsal spine 2 in head; pectoral a little shorter than caudal, scarcely longer than ventrals. Caudal rounded, shorter than head, probably  $4\frac{1}{2}$  in body. Body and head completely naked.\* Brownish yellow; upper parts of head and body with small, irregularly placed brown spots and streaks; dorsals and caudal finely barred with dark specks. (Steindachner). Gulf of California; not seen by us; known from 3 specimens 37 $\frac{1}{2}$  mm. long. We refer this species provisionally to *Evermannia*, with which genus it agrees in external respects, although the mouth is much larger. It may be the type of a distinct genus. It differs from *Clevelandia* in the entire absence of scales. (*longus*, long; *pinna*, fin.)

*Gobiosoma longipinne*, STEINDACHNER, Ich. Beitr., VIII, 27, 1879, Las Animas Bay, Gulf of California; JORDAN & EIGENMANN, l. c., 509.

*Evermannia longipinnis*, JORDAN, Proc. Cal. Ac. Sci. 1896, 229.

## 2589. EVERMANNIA ZOSTEBURA (Jordan &amp; Gilbert).

Head  $3\frac{1}{2}$ ; depth 6. D. IV-15; A. 14; eye equals snout, 5 in head; P. 1 $\frac{1}{2}$ ; C. 1 $\frac{1}{2}$ . Body compressed, profile convex; snout short, not very blunt; eyes high, the maxillary reaching to their posterior margin; mouth oblique, jaws equal; first spine of dorsal filamentous, reaching to middle

\* At our request Dr. Steindachner has reexamined the types of this species. He still finds them "vollkommen schuppenlos."

of soft do  
with dots  
bands; fir  
broad med  
caudal ye  
edge abov  
male with  
and anal  
fins with p  
2 inches.  
of Mazatla  
sand. It i  
a very han  
than any o

*Gobiosoma zo*  
(fin rays  
& EIGENMANN  
*Evermannia*

*Gobiosoma, G*

Body ent  
in several s  
mally 7, rar  
head; shoul  
 $\sigma\mu\alpha$ , body

a. Coloration

b. Maxi

d

h

le

n

er

n

b

bb. Max

c.

of soft dorsal (male); body entirely naked. Body everywhere speckled with dots of dark brown. Male sometimes with traces of 8 olive cross bands; fins very ornate, the dorsal and anal yellowish at base, then a broad median band of jet black, then a broad white margin; middle of caudal yellow to the tip, with a black band above and below, and a white edge above and below this as in dorsal and anal; no bands on tail. Female with dorsal filament short, reaching about to first soft ray; dorsals and anal checkered with blackish; caudal faintly barred; all vertical fins with pale edgings, but without the black stripe of the males. Length 2 inches. Very common on sandy bottoms, everywhere about the estuary of Mazatlan, the numerous specimens here described being dug out of the sand. It is seldom found much, if any, below the mark of low tide. It is a very handsomely colored species, the male being more strikingly marked than any other of our gobies. (ζωστῆρ, band; οὐρά, tail.)

*Gobiosoma zosterurum*, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1881, 361, Mazatlan (fin rays incorrect), (Type, No. 29245, U. S. Nat. Mus. Coll. C. H. Gilbert); JORDAN & EIGENMANN, l. c., 509.

*Evermannia zosterura*, JORDAN, Proc. Cal. Ac. Sci., 2d ser., vol. iv, 1895, 498, pl. 51.

## 827. GOBIOSOMA, Girard.

(NAKED GOBIES.)

*Gobiosoma*, GIRARD, Proc. Ac. Nat. Sci. Phila. 1858, 169 (*alepidotus*).

Body entirely naked; mouth moderate, horizontal; snout blunt; teeth in several series, the outer row enlarged; no canines; dorsal spines normally 7, rarely 5 or 6; second dorsal and anal short; no barbels about head; shoulder girdle without flaps. Species chiefly American. (*Gobius*: βῶμα, body.)

a. Coloration olivaceous, mottled with darker; no red nor blue.

b. Maxillary extending to beyond pupil,  $4\frac{1}{2}$  in head; color blackish, with sharply defined cross bars of whitish. Body rather short, the depth  $5\frac{1}{2}$  in length; head  $3\frac{1}{2}$ ; snout low, little obtuse; mouth large, rather oblique, the maxillary  $2\frac{3}{4}$  in head; teeth small, in few series above, in a band below, the outer enlarged; fins low; caudal  $1\frac{1}{2}$  in head. Cross bands on body as wide as eye, not quite meeting below; a dark blotch on base of pectoral, a fainter one on base of caudal; fins nearly plain. D. VII-13; A. 12. HISTRIO, 2590.

bb. Maxillary extending to below posterior part of orbit; coloration not sharply defined, the body usually with dark cross streaks.

c. Body rather short, chubby, the depth about 4 in length; head about  $3\frac{1}{2}$ ; head rounded above; teeth in several series, slender, the outer ones somewhat elongate, none of the inner ones specially enlarged. Color olivaceous, with dark points; sides with narrow, alternating light and dark bars; a row of small linear dark spots along middle of sides; first dorsal with 3 oblique dark bars; second dorsal, caudal, and pectorals finely barred, base and edge of anal light, middle dark; breast with many well-defined spots; a dark line running forward and downward from eye to angle of mouth, another extending straight down; a black bar on edge of preopercle, a black spot on upper edge of opercle. D. VII-13; A. 10. MOLESTUM, 2591.

cc. Body more elongate, depth 5 to 6 in body; head very broad, flattish above, with tumid cheeks, its length  $3\frac{1}{2}$  in body; eye small, longer than snout, 5 in head; mouth large, little oblique, the jaws subequal.

d. Soft dorsal with 14 rays; no crescent at base of caudal; maxillary extending to below posterior part of orbit (at least in male),  $2\frac{1}{2}$  in head; teeth in few series, the outer considerably enlarged; 2 teeth on each side of inner series of lower jaw especially large canines; dorsal spines slender, none filamentous; caudal rounded. Olivaceous, with darker cross shades of rounded spots; vertical fins dusky, faintly barred. Teeth of the female smaller than those of the male but smaller; head narrower; more slender. D. VII-14; A. 10. no. 2592.

dd. Soft dorsal with 12 rays; a brown crescent at base of caudal.

CRESCENTALE, 2503.

aa. Coloration not plain olivaceous; head with a red bar; anterior dorsal rays not produced in filaments; head and body compressed; groates: depth  $5\frac{1}{2}$  in total length, head about 4; angle of mouth little behind center of eye; eye 4 in head; teeth pointed, in several series, those of the outer series a little enlarged; caudal rounded. Head light yellow; a carmine-red bar extending along upper edge of head, from upper corner of gill opening to snout, where it joins its fellow, ending behind over the pectoral in a small indigo-blue spot; body with 16 or 17 light green, well-defined cross bars, separated by narrow white stripes; fins chiefly greenish. D. VII (VI)-11 or 12; A. 10.

MULTIFASCIATUM, 2594.

2590. *GOBIOSOMA HISTRIO*, Jordan.

Head  $3\frac{1}{2}$ ; depth  $5\frac{1}{2}$ . D. VII-12 or 13; A. 11 or 12; maxillary  $2\frac{1}{2}$  in head; caudal 14. Body rather short; snout depressed, little obtuse; mouth large, rather oblique, maxillary reaching to below posterior part of orbit; chin without barbels; many series of minute papillae along mucous pores of head. Teeth small, in few series above, in a band below, the outer enlarged. Fins low. Cross bands on body whitish, as wide as eye, not quite meeting below; a dark blotch on base of pectoral, a fainter one on base of caudal; fins nearly plain. Length 2 inches. Gulf of California; known only from the Gulf of California at Guaymas (Emerie; Evermann & Jenkins) and La Paz (Gilbert). (*histrío*, a harlequin.)

*Gobiosoma histrío*, JORDAN, Proc. U. S. Nat. Mus. 1884, 260, Guaymas, Mexico (Coll. H. F. Emerie); JORDAN & EIGENMANN, l. c., 508; EVERMANN & JENKINS, Proc. U. S. Nat. Mus. 1891, 162.

2591. *GOBIOSOMA MOLESTUM*, Girard.

Head about  $3\frac{1}{2}$ ; depth 4. D. VII-13; A. 10; vertebrae 12 + 15. Body rather short, maxillary extending to below posterior part of orbit. Teeth in several series, slender, the outer ones somewhat elongate, none of the inner ones specially enlarged. Skull flattish, with a slight median keel; lateral crests developed, lower and stronger than in *Gobius*; interorbital very narrow, bounded by 2 minute crests; bones of the skull very weak and fragile. Teeth in both jaws recurved, in 2 or 3 series. Olivaceous, with dark points; sides with narrow, alternating light and dark bars; a row of small dark spots along middle of side; first dorsal with 3 oblique dark bars; second dorsal, caudal, and pectorals finely barred; base and

edge of  
dark line  
another  
prooperc  
at Key W  
of dark  
mottled  
plain. I  
shallow v  
(*molestus*,

*Gobiosoma*  
(GIRARD)  
JORDAN  
*Gobiosoma*  
Pensac

Head  $3\frac{1}{2}$   
Body mor  
Eye smal  
extending  
Teeth in fe  
of inner se  
not filam  
rounded sp  
United Sta  
ward in sl  
Charleston

*Gobius bosci*,  
Carolina  
*Gobius alepic*  
KAY, N.  
*Gobius viridi*  
New Yor  
*Gobiosoma bo*  
l. c., 508.  
*Gobiosoma al*

Head  $3\frac{1}{2}$   
11. Body v  
the head a  
oblique, the  
border of or  
space. Tee  
like, and di  
broad base,

edge of anal light, middle dark; breast with many well-defined spots; a dark line running forward and downward from eye to angle of mouth, another extending straight downward from eye; a black bar on edge of preopercle, and a black spot on upper edge of opercle. A specimen taken at Key West is thus described: Pale olive, with darker cross bands formed of dark dots; a row of dark dots along middle of side; vertical fins all mottled and faintly barred with dark olive; pectorals and ventrals nearly plain. Length  $2\frac{1}{2}$  inches. Gulf coast of the United States; common in shallow waters along the coast from Key West to Texas and south to Bahia. (*molestus*, disturbed.)

*Gobiosoma molestum*, GIRAUD, Proc. Ac. Nat. Sci. Phila. 1858, 169, Indianola, Texas; GIRAUD, U. S. and Mex. Bound. Surv., 27, pl. 12, fig. 14, 1859; GÜNTHER, Cat., III, 550, 1861; JORDAN & GILBERT, Synopsis, 638, 1883; JORDAN & EIGENMANN, l. c., 508.  
*Gobiosoma alepidotum*, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1882, 297, Laguna Grande, Pensacola. (Coll. Dr. Jordan.)

2592. GOBIOSOMA BOSCI (Lacépède).

(NAKED GOBY.)

Head  $3\frac{1}{2}$ ; depth 5 to 6. D. VII-14; A. 10; eye 5, longer than snout. Body more elongate; head very broad, flattish above, with tumid cheeks. Eye small. Mouth large, little oblique, jaws subequal, the maxillary extending to below posterior part of orbit (at least in male),  $2\frac{1}{2}$  in head. Teeth in few series, the outer considerably enlarged; 2 teeth on each side of inner series of lower jaw especially large canines. Dorsal spines slender, not filamentous; caudal rounded. Olivaceous, with darker cross shades of rounded spots; vertical fins dusky, faintly barred. Atlantic coast of the United States, Cape Cod to Florida; generally common, especially southward in shallow grassy bays. (Named for M. Bosc, French consul at Charleston in the last century; an ardent naturalist.)

*Gobius bosci*, LACÉPÈDE, Hist. Nat. Poiss., II, 555, pl. 16, fig. 1, 1798, Charleston, South Carolina. (Coll. M. Bosc.)  
*Gobius alepidotus*, BLOCH & SCHNEIDER, Syst. Ichthyol., 547, 1801, after LACÉPÈDE; DE KAY, N. Y. Fauna: Fishes, 160, pl. 23, fig. 70, 1842.  
*Gobius viridipallidus*, MITCHILL, Trans. Lit. and Philos. Soc. N. Y., I, 1814, 379, pl. 1, fig. 8, New York.  
*Gobiosoma bosci*, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1882, 613; JORDAN & EIGENMANN, l. c., 508.  
*Gobiosoma alepidotum*, GÜNTHER, Cat., III, 85, 1861; JORDAN & GILBERT, Synopsis, 638, 1883.

2593. GOBIOSOMA CRESCENTALE, Gilbert.

Head  $3\frac{1}{2}$  in length; depth  $6\frac{1}{2}$ ; eye  $5\frac{1}{2}$  in head; snout  $5\frac{1}{2}$ . D. VII-12; A. 11. Body very slender, the head depressed, broad and flattened above, the head and body of nearly equal depth throughout. Mouth small, oblique, the maxillary not extending beyond the vertical from posterior border of orbit,  $2\frac{1}{2}$  in head; eyes small,  $1\frac{1}{2}$  in the rather broad interorbital space. Teeth in bands in both jaws, the outer series enlarged, canine-like, and distant. Fins all small, the caudal short and rounded from a broad base, pectoral as long as head without snout; ventrals short, not

reaching  $\frac{3}{4}$  the distance from their base to vent; dorsal spines not filamentous; skin wholly naked. Color in spirits, lower half of head and body uniform warm brown, the back much lighter, the two areas separated by a well-defined line along middle of sides; this line passing through orbit and through the middle of the base of the pectoral fin; back light grayish, with brownish reticulations, which tend to form 5 or 6 indistinct darker bars uniting with the darker area below the lateral line; a conspicuous brown crescent at base of caudal and pectorals, broad below, narrowing above, margined in front with whitish; anal brown at base; dorsal and caudal with small brown spots forming faint cross series. A single specimen known. Off coast of Lower California. (Gilbert.) (*erescentalis*, pertaining to a crescent.)

*Gobiosoma erescentalis*, GILBERT, Proc. U. S. Nat. Mus. 1891, 557, off coast of Lower California, at Albatross Station 2825, 24° 32' N., 110° 19' 15" W., in 79 fathoms.

#### 2594. GOBIOSOMA MULTIFASCIATUM, Steindachner.

Head about 4; depth 5 $\frac{1}{2}$ . D. VII-12 (Poey), VI-11 (Steindachner); eye 4 in head. Body and head compressed. Angle of mouth little behind center of eye. Teeth pointed, in several series, those of outer series somewhat enlarged. Dorsal rays not filamentous; caudal fin rounded. Head light yellow; a carmine-red bar extending along upper edge of head, from upper corner of gill opening to snout, where it joins its fellow, ending behind over the pectoral in a small indigo-blue spot; body with 16 or 17 light-green, well-defined cross bars, separated by narrow white stripes. (Steindachner.) West Indies; known from Cuba, St. Thomas, and the Lesser Antilles; not seen by us. Its coloration is very different from that of *Gobiosoma*, and it may belong to a distinct genus. (*multus*, many; *fasciatus*, banded.)

*Gobius lineatus*,\* POEY, Memorias, II, 424, 1861, Cuba; name preoccupied by *Gobius lineatus*, JENYNS.

*Gobiosoma multifasciatum*, STEINDACHNER, Ichth. Beitrage, V, 183, 1870, Lesser Antilles; JORDAN & EIGENMANN, l. c., 509; EIGENMANN & EIGENMANN, l. c., 73.

#### 828. BARBULIFER, Eigenmann & Eigenmann.

*Barbulifer*, EIGENMANN & EIGENMANN, Proc. Cal. Ac. Sci. 1888, 70 (*papillosus*).

A series of numerous minute barbels around the mouth and chin; otherwise as in *Gobiosoma*; body naked, the dorsal spines 7; second dorsal and anal very short. (*barbula*, a small barbel; *fero*, I bear.)

#### 2595. BARBULIFER CEUTHÆCUS (Jordan & Gilbert).

Head 3 $\frac{1}{2}$ ; depth 7. D. VII-10; eye 4; A. 10. Body slender; head narrow and slender, depressed; snout not blunt; mouth terminal, oblique, the maxillary reaching to below eye, 3 in head; eyes close together; chin

\* *Gobius lineatus* is thus described: Head 3 $\frac{1}{2}$ ; depth of body 6 in length. D. VII-12; eye 6 in head. Body elongate, subcylindrical, maxillary extending almost to below middle of eye; pectorals rounded; dorsals high. Yellowish green; the body with 20 vertical yellow bands; a red band extending from snout to point of opercle; fins yellowish. Cuba. (Poey.) Type .43 mm. in length.

with a f  
Upper h  
areas al  
lower pa  
middle o  
jections;  
pectoral,  
(κεθνος,  
cavity of

*Gobiosoma*  
(Type)  
*Barbulifer*  
Florida

*Typhlogobius*  
*Othonops*, R

Body m  
naked ski  
the maxill  
row band  
capable of  
to mere ven  
greatly mo  
ble spines  
ventral di  
known; si  
marks. (z

\* This spec  
scribed by D  
short and rob  
dorsal spine  
orbital area  
reaching to  
orbital diam  
angle of the  
colored (2 of  
mouth longer  
chin, behind  
barbels on ea  
gin of the ey  
straight down  
another series  
the opercle;  
the opercle; a  
larger and esp  
eye, extending  
rounding mou  
posteriorly, oth  
higher than fin  
anal lower th  
Color yellow;  
spots of darke  
upper part of  
oblique bar of  
on base of cau  
mann.)

with a fringe of short barbules; vertical fins high, rays not filamentous. Upper half of head and body brown, finely speckled; 4 oblong, colorless areas along base of dorsals and a smaller one on back of caudal peduncle; lower parts abruptly pale; back with 5 or 6 blackish cross bars reaching to middle of sides, below which they extend as 5 or 6 short V-shaped projections; a brownish streak below eye; a small brown bar on base of pectoral, and a jet black bar at base of caudal. About Key West; scarce. (*κεῦθος*, a cavity; *οικέω*, to inhabit; the type specimen taken from the cavity of a sponge.)

*Gobiooma ceuthæctum*, JORDAN & GILBERT, Proc. U.S. Nat. Mus. 1884, 29, Key West; young (Type in U. S. N. M.); JORDAN & EIGENMANN, *l. c.*, 508.

*Barbulifer papillosus*,\* EIGENMANN & EIGENMANN, Proc. Cal. Ac. Sci. 1888, 70, Key West, Florida; adult.

### 829. TYPHLOGOBIUS, Steindachner.

(BLIND GOBIES.)

*Typhlogobius*, STEINDACHNER, Ichth. Beitrage, VIII, 24, 1879 (*californiensis*).

*Othonops*, ROSA SMITH, Proc. U. S. Nat. Mus. 1881, 19 (*eos = californiensis*).

Body moderately elongate, compressed, covered with loose, smooth, naked skin. Head large, depressed, with tumid cheeks. Mouth large, the maxillary reaching to beyond the orbit; jaws equal, each with a narrow band of villiform teeth, the outer teeth slightly enlarged; lower jaw capable of little motion; snout rounded; no cirri. Eyes very small, reduced to mere vestiges, covered by skin, and functional only in the young. Skull greatly modified, the brain case quadrate. Fins low; first dorsal of 2 flexible spines; second dorsal moderate; anal very short; caudal rounded; ventral disk as in *Gobius*. Gill openings rather narrow. One species known; singular blind gobies, living like slugs under rocks between tide marks. (*τυφλός*, blind; *Gobius*.)

\*This species, which we suppose to be the adult of *Barbulifer ceuthæctus*, is thus described by Dr. Eigenmann: Head  $3\frac{1}{2}$  ( $4\frac{1}{2}$  in total); depth  $4\frac{1}{2}$  ( $5\frac{1}{2}$ ). D. VII-9; A. 9. Body short and robust, deepest below first dorsal spine; head blunt, profile straight from first dorsal spine to eye, much curved in front of eye; eye longer than snout,  $\frac{3}{4}$  in head; inter-orbital area  $\frac{2}{3}$  diameter of eye; snout blunt; mouth small, oblique; maxillary 3 in head, reaching to below anterior margin of pupil; lips thick. About 21 barbels, in length  $\frac{1}{2}$  orbital diameter or longer, arranged as follows: A series of 7 cross the snout from one angle of the mouth to the opposite angle, the anterior 3 on the snout rather thick and colored (2 of them nasal), all the others yellowish, the barbel nearest each angle of the mouth longer than any of the others; on the lower jaw a barbel near each rictus, 2 on the chin, behind which are 2 pairs of barbels; posterior to these and below the rictus are 2 barbels on each side; 1 slender barbel on each side of preopercle below the posterior margin of the eye. Numerous rows of pores or papillæ on the head; 1 series extending straight downward on the anterior part of the opercle, from the upper end of which another series extends perpendicularly backward; other pores irregularly scattered on the opercle; a double series extending along edge of preopercle, the pores becoming larger and especially conspicuous below, meeting on the chin; 6 or 7 series radiating from eye, extending to snout, maxillary, and opercular series below; a row of pores nearly surrounding mouth, curving backward, encircling the nasal opening; 1 series about the eye posteriorly, otherwise none on top of head or nape; fins high and rounded; second dorsal higher than first,  $1\frac{1}{2}$  in head, caudal very broad and rounded, equal to the head in length; anal lower than soft dorsal; ventral reaching  $\frac{2}{3}$  to vent,  $1\frac{1}{2}$  in head; pectoral  $1\frac{1}{2}$  in head. Color yellow; upper half of body with a broad band of purplish spots; 6 diamond-shaped spots of darker cross the band, extending above and below it; nape, top of head, and upper part of cheek covered with dark points; opercle light yellow, cheeks darker; an oblique bar of black points on upper half of pectoral base, a curved bar of fainter spots on base of caudal; fins otherwise colorless and transparent. Length  $1\frac{1}{2}$  inches. (Eigenmann.)

## 2590. TYPHLOGOBIOUS CALIFORNIENSIS, Steindachner.

(BLIND GOBY OF POINT LOMA; PINK-FISH.)

Head  $3\frac{1}{2}$ ; depth 5; eye 6; eye concealed, very small; D. II-12; A. 12. Vertebrae 17+13. Body subcylindrical, the males more compressed behind; head very broad behind, its greatest width  $\frac{2}{3}$  its length. Interorbital space a mere ridge; skin about mouth and eye very loose; a snail papilla in front of nasal opening. Lower lip developed as a fold; another fold of skin behind it, bordered with fine cilia; behind this fold is a row of short, thick papillae; edge of jaw rounded. Spinous dorsal remote from the soft dorsal in the male, but connected with it by a low membrane, this membrane absent in the female; soft dorsal much higher than the spinous; caudal broad, rounded; anal very short, inserted under sixth dorsal ray, and coterminous with dorsal; pectorals little longer than ventrals, 2 in head. Body naked; males with small tubercular plates irregularly placed. Skull highest at its posterior part, depressed forward; the bones all thick and strong. No lateral crests; a median keel which is lowest behind. Orbit not bounded by any ridges. Two keels diverge from the posterior end of the median keel to the insertion of the suprascapula. Premaxillaries and mandible very long. Teeth of the upper jaw all alike, long, close-set, in a broad band, those of the lower jaw in a narrow band, the inner ones apparently larger. Color uniform light pink. Length 2 inches. Coast of Lower California, from San Diego southward to Cerros Island; an extraordinary fish, found attached to the lower side of rocks in shallow water or surf; especially common at Point Loma.

*Typhlogobius californiensis*, STEINDACHNER, Ichth. Beiträge, VIII, 24, 1879, False Bay, San Diego, California (Coll. Prof. Eesmark); JORDAN & GILBERT, Synopsis, 639, 1883; JORDAN & EIGENMANN, l. c., 511.

*Othonops eos*, ROSA SMITH, Proc. U. S. Nat. Mus. 1881, 53, Point Loma, California.

## 830. TYNTLASTES, Günther.

*Tyntlastes*, GÜNTHER, Proc. Zool. Soc. London 1862, 193 (*sagitta*).

Body elongate, compressed, covered with small, imbricate, cycloid scales. Head elongate, quadrangular. Mouth wide, oblique, the lower jaw projecting; teeth small, in single series, none on vomer or palatines. Eyes very small, or rudimentary. Dorsal fin single, continuous, about 6 of its anterior rays simple; caudal fin pointed, more or less joined to the dorsal and anal; ventral fins united. Air bladder very small or absent. No pseudobranchiae. Vertebrae 11+20. Pacific Ocean. (*τυντλάστης*, a mud-dabber.)

a. Dorsal and anal each with 15 soft rays; head  $4\frac{1}{2}$  in length.

BREVIS, 2597.

aa. Soft dorsal and anal each with 21 unbranched or soft rays; head  $5\frac{1}{2}$  in length.

SAGITTA, 2598.

## 2597. TYNTLASTES BREVIS (Günther).

Head  $4\frac{1}{2}$ ; depth 8. D. VI, 15; \* A. 15. Eyes minute. Jaws each with a

\* The dorsal formula is apparently VIII, 14 in 2 half-digested specimens taken from the stomach of a *Centroponus* at Panama. (Gilbert.)

series of  
seen by u

*Amblyopus*  
Centr. A  
*Tyntlastes b*

Head 5  
pressed.  
small. S  
4 in body  
under par  
grayish.  
exact loca

*Amblyopus*  
from Lo  
*Tyntlastes s*

*Gobioides*, L.  
*Plecopodus*,  
*Gobioides*  
*Ognichodes*, S

Body gr  
head small  
ing; gill o  
being very

Dorsal fin l  
widely sep  
del; ventr  
air bladd  
genus *Gobi*  
of scales, a  
Brackish w  
*είδος*, rese

a. Eye sma  
aa. Eye mi

Head  $5\frac{1}{2}$  (  
to 10 in he  
l, 16. Body  
teeth in ba  
*Gobioides pe*  
part of bod



series of wide-set teeth. Caudal fin black. (Günther.) Panama; not seen by us. (*brevis*, short.)

*Amblyopus brevis*, GÜNTHER, Proc. Zool. Soc. London 1864, 151, Panama; GÜNTHER, Fishes Centr. Amer., 441, 1869.

*Tyntlastes brevis* JORDAN & EIGENMANN, l. c., 512.

2598. TYNTLASTES SAGITTA (Günther).

Head  $5\frac{3}{4}$ ; depth  $9\frac{3}{4}$ . D. VI, 21; A. 21. Body and head elongate, compressed. Maxillary reaching to behind eye; teeth subhorizontal, very small. Scales becoming larger posteriorly. Caudal arrow-shaped, about 4 in body; pectorals as long as ventrals, 2 in head. Grayish, sides and under parts silvery; an ovate gray spot before each dorsal ray; caudal grayish. (Günther.) Length  $9\frac{1}{2}$  inches. Coast of Lower California; exact locality unknown. (*sagitta*, arrow.)

*Amblyopus sagitta*, GÜNTHER, Proc. Zool. Soc. London 1862, 193, "California," probably from Lower California.

*Tyntlastes sagitta*, JORDAN & GILBERT, Synopsis, 639, 1883; JORDAN & EIGENMANN, l. c., 512.

831. GOBIOIDES, Lacépède.

(BARRETOS.)

*Gobioides*, LACÉPÈDE, Hist. Nat. Poiss., II, 580, 1798 (*broussonnetii*).

*Plecopodus*, RAFINESQUE, Analyse de la Nature, 87, 1815 (*broussonnetii*); substitute for *Gobioides*, regarded as objectionable.

*Ognichodes*, SWAINSON, Nat. Hist. Class'n Animals, II, 183 and 276, 1839 (*broussonnetii*).

Body greatly elongate, compressed behind, the scales very minute; head small; eyes very small; mouth large, oblique, the lower jaw projecting; gill openings moderate. Teeth in a band, those in the outer series being very strong. Dorsal rays V to VII, 15 to 23; anal rays 16 to 23. Dorsal fin low, continuous, the spines similar to the soft rays, but more widely separated; the soft dorsal and the anal are joined to base of caudal; ventrals 45, united in a disk which is formed much as in *Gobius*. No air bladder; no pseudobranchiae. From *Tenioides* (= *Amblyopus*) the genus *Gobioides* is distinguished by the absence of barbels, the presence of scales, and by the much smaller number of rays in its vertical fins. Brackish waters of the Tropics, reaching a considerable size. (*Gobius*; *είδος*, resemblance.)

a. Eye small, but evident; scales evident, larger behind.

BROUSSONNETII, 2599.

aa. Eye minute, not evident; scales minute.

PERUANUS, 2600.

2599. GOBIOIDES BROUSSONNETII, Lacépède.

Head  $5\frac{1}{2}$  (young) to 7 (adult); caudal  $3\frac{1}{2}$  to 5; eye small but evident, 7 to 10 in head; interorbital space 1 to  $1\frac{1}{2}$  diameter of eye. D. VII, 16; A. I, 16. Body elongate, mouth oblique, maxillary extending beyond eye; teeth in bands, the outer series enlarged, shorter, and closer set than in *Gobioides peruanus*; scales twice as large as in *peruanus*, those on anterior part of body not imbricated, much smaller than those on posterior part,

which are elongate oval in form. Violet bars extending downward and forward on the upper part of body; sometimes a violet spot with a lighter or darker dot at end of the bars; head marbled or spotted with dark violet or brown. (Steindachner.) Length 20 inches or more. West Indies to Brazil; common southward, ascending rivers; once taken near New Orleans (Bean & Bean). (Named for Dr. Augustin Broussonnet, professor in the University of Montpellier.)

*Gobioides broussonnetii*, LACÉPÈDE, Hist. Nat. Poiss., II, 580, 1798, probably from Surinam, "given by Holland to France."

*Amblyopus brasiliensis*, BLOCH & SCHNEIDER, Syst. Ichth., 69, 1801, Brazil; on drawing made by Prince Maurice; CUVIER & VALENCIENNES, Hist. Nat. Poiss., XII, 121, 1837.

*Gobius oblongus*, BLOCH & SCHNEIDER, Syst. Ichth., 548, 1801; based on LACÉPÈDE.

*Gobioides burreto*, POEY, Memorias, II, 282, 1861, Cuba; POEY, Synopsis, 394, 1868; POEY, Enumeratio, 125, 1876.

*Amblyopus mexicanus*, O'SHAUGHNESSY, Ann. Mag. Nat. Hist., series IV, vol. XV, 1875, 147, Mexico.\*

*Gobioides broussonneti*, JORDAN & EIGENMANN, l. c., 512; BEAN & BEAN, † Proc. U. S. Nat. Mus. 1895, 631.

#### 2600. GOBIOIDES PERUANUS (Steindachner).

Head 5; depth 11. D. VII, 17; A. I, 16. Eye scarcely visible, much smaller than in *G. broussonneti*; scales very minute; snout  $2\frac{1}{2}$  in post-orbital part of head; interorbital 5 in head; lower jaw slightly projecting; maxillary  $2\frac{3}{4}$  in head; a series of large slender teeth in each jaw, behind which, in each jaw, is a narrow band of fine teeth; caudal  $4\frac{1}{2}$  in

\* The following is Mr. O'Shaughnessy's description of *Amblyopus mexicanus*: D. VII, 15; A. I, 15. Depth 13 in total length. Body covered all over with scale-shaped crypts. Head naked. Dorsal  $\frac{3}{4}$  height of body. Eye small, but distinct. Snout obtuse; lower jaw extending a little beyond upper. Teeth small, close set, the outer series much smaller and more closely set than in *G. broussonnetii*. Dorsal and anal connected with the caudal. Upper parts dark brown, with a series of white spots along the whole length of the side; lower parts of sides and belly white. One specimen in the British Museum, from Mexico, purchased. Length  $20\frac{1}{2}$  inches. (O'Shaughnessy.)

† This seems to differ from *G. broussonnetii* in color only.  
‡ The following description is given by Bean & Bean of *Gobioides broussonnetii* (Lacépède): Head 7; depth 14. D. VI, 17; A. I, 16. The greatest depth of the head equals the length of the upper jaw, or about  $\frac{1}{2}$  the length of head without snout. The body is compressed. Its greatest thickness is contained  $1\frac{1}{2}$  times in its greatest depth. The teeth are in narrow bands in each jaw, some of those in the outer row enlarged, canine-like, and curved inward. All of the teeth are more or less curved inward and depressible. The vomer and palate are toothless. The mouth is oblique, the lower jaw projecting slightly beyond the upper. The maxilla extends well behind the eye, its length is slightly more than  $\frac{1}{2}$  that of head without the snout. It is not much expanded posteriorly. Eyes very small, their diameter equaling  $\frac{1}{2}$  length of snout, about equal to width of interorbital space. The snout scarcely equals more than  $\frac{1}{2}$  of the head's length. Gill openings wide, the membranes wholly joined to the isthmus. Branchiostegals much curved, 4 in number. The dorsal begins at a distance from the nape equal to the postorbital part of the head, the origin being about over the end of the extended pectoral. The ventral reaches farther back than the pectoral, and is longer than that fin, its length equaling postorbital part of head. The distance of the vent from the tip of the snout equals somewhat more than 3 times the length of the head; it is under the interspace between the last spine and first ray of the dorsal, with a small genital papilla behind it. The caudal is very long and tapering,  $1\frac{1}{2}$  times as long as the head. The dorsal spines are long and slender, the fifth nearly as long as the postorbital part of the head. The second dorsal ray is slightly longer. The anal rays are about as long as those of the dorsal. The scales are thin, not imbricated, except on the posterior part of the head, where they are long and elliptical in shape. The head and breast are naked. The colors have faded out in alcohol; the ground color appears to have been light brown, with darker blotches on the median line of the body under the spinous portion of the dorsal and the anterior part of the soft dorsal. (Bean & Bean.) Here described from a specimen obtained in the Gulf of Mexico by Mr. Robert S. Day, of New Orleans, Louisiana, and is No. 38220, U. S. Nat. Mus.

body, co-  
series of  
the mem  
ascendin

*Amblyopu*  
42, 188  
*Gobioides*

*Cayennia*,  
Body n  
ventrals  
anterior  
Otherwis  
(Name fro

Head 9;  
than wide  
posterior  
caudal 7 i  
black ant  
(Guichenot  
Paris.)

*Cayennia g*  
MANN &

Bony fis  
upper surr  
ventral fin  
bones flat  
with the  
appressed  
scapula) I  
(carpals)."

This rem  
disk; κερα

Body fus  
wide, with  
tongue. P  
upper. Spi

3030-

body, connected by membrane to dorsal and anal; sides with regular cross series of pores. Body with narrow angular cross bars; dorsal rays violet, the membrane yellowish. (Steindachner.) Shores of Ecuador and Peru, ascending rivers.

*Amblyopus peruanus*, STEINDACHER, Fisch-Fauna des Cauca und Flüsse bei Guayaquil, 42, 1880, Guayaquil.

*Gobioides peruanus*, EIGENMANN & EIGENMANN, Proc. Cal. Ac. Sci., 2d ser., 1, 1888, 75.

### 832. CAYENNIA, Sauvage.

*Cayennia*, SAUVAGE, Bull. Sci. Philom., ser. 7, IV, 1880, 57 (*guichenoti*).

Body much elongate; dorsals united, caudal free from dorsal and anal; ventrals united, not adhering to belly; teeth small, the outer enlarged; anterior part of body naked, posterior part covered with cycloid scales. Otherwise as in *Gobioides*, from which the genus may not be separable. (Name from Cayenne.)

#### 2601. CAYENNIA GUICHENOTI, Sauvage.

Head 9; depth 17. D. VI, 17; A. I, 16; vertebrae about 36. Head deeper than wide; eye small, placed well forward; maxillary reaching to below posterior margin of eye; a low membrane connecting dorsal and caudal; caudal 7 in length; ventrals  $1\frac{1}{2}$  in head. Color brownish, marbled with black anteriorly. Cayenne (Sauvage); not seen by us. (Named for A. Guichenot, formerly ichthyologist of the Muséum d'Histoire Naturelle at Paris.)

*Cayennia guichenoti*, SAUVAGE, Bull. Soc. Philom., ser. 7, IV, 1880, 57 Cayenne; EIGENMANN & EIGENMANN, Proc. Cal. Ac. Sci. 1888, 76.

## Suborder DISCOCEPHALI.

Bony fishes "with a suctorial transversely laminated oval disk on the upper surface of the head (homologous with a flat dorsal fin), thoracic ventral fins with external spines, a simple basis cranii, intermaxillary bones flattened, with the ascending processes deflected sideways, and with the supramaxillary bones attenuated backward, flattened, and appressed to the dorsal surface of the intermaxillaries; hypercoracoid (or scapula) perforated nearly in the center, and with 4 short actinosts (carpals)." (Gill.)

This remarkable group consists of a single family, *Echeneididae*. (*δίσκος*, disk; *κεφαλή*, head.)

### Family CLXXXIX. ECHENEIDIDÆ.

#### (THE REMORAS.)

Body fusiform, elongate, covered with minute, cycloid scales. Mouth wide, with villiform teeth on jaws, vomer, palatines, and usually on tongue. Premaxillaries not protractile. Lower jaw projecting beyond upper. Spinous dorsal modified into a sucking disk, which is placed on

the top of the head and neck, and is composed of a double series of transverse, movable, cartilaginous plates, serrated on their posterior or free edges. By means of this disk these fishes attach themselves to other fishes or to floating objects, and are carried for great distances in the sea. Opercles unarmed. Pectoral fins placed high; ventral fins present, thoracic and close together, I, 5; dorsal and anal fins long, without spines, opposite each other; caudal fin emarginate or rounded. Branchiostegals 7. Gills 4, a slit behind the fourth; gill rakers short; gill membranes not united, free from the isthmus. Pseudobranchiae obsolete. Several pyloric appendages. No air bladder. No finlets. No caudal keel. Vertebrae more than 10 + 14. Genera 4; species about 10, found in all seas, all having a very wide range. The species of this group are apparently descended from a fossil genus, *Opisthomyzon*,\* Cope (*glaronensis*), characterized by the small posterior disk and slender body.

The following description of this family is given by Dr. Gill: Body elongated, subcylindrical, diminishing backward gradually from the head and into the slender caudal peduncle. Anus subcentral. Scales cycloid, very small, and not, or scarcely, imbricated. Lateral line nearly straight and very faint. Head above oblong and with a flattened straight upper surface, furnished with an adhesive oblong or elongated, laminated disk. The eyes are rather small, submedian, and overlunged by the disk. Suborbital bones forming a slender infraorbital chain; the first or preorbital triangular and thick. Opercular apparatus normally developed and unarmed. Nostrils double, close together. Mouth terminal or, rather, superior, the lower jaw projecting, but with the cleft nearly horizontal and not extending laterally to the eyes. Teeth present on the jaws and palate. Branchial apertures ample and fissured forward. Branchiostegal rays 7 (or 8) on each side. The adhesive disk on the upper surface of the head is a modified first dorsal fin, and from the snout generally extends more or less posteriorly on the nape and back; it is oblong or elongated and of an oval or elliptical form, divided into equal halves by a longitudinal septum, and with more or less numerous transverse laminae in each division, the laminae being slightly erectile and depressible. Dorsal fin oblong or elongated on the posterior half of the body (including head), ending some distance from the caudal. Anal fin opposite and similar to the dorsal. Caudal fin rather small, variable in outline, but never deeply forked. Pectoral fins moderate, inserted high on the sides. Ventral fins thoracic, each with a spine and 5 branched rays. The vertebral column has vertebrae in slightly increased numbers, the abdominal vertebrae being about 12 to 14 and the caudal 15 or 16. The stomach is cecal and the pyloric caeca are present in moderate numbers. The air bladder is obsolete.

\* "A careful comparison of the proportions of all the parts of the skeleton of the fossil *Echeneis* with those of the living forms, such as *Echeneis naucrates* or *Echeneis renora*, shows that the fossil differs nearly equally from both, and that it was a more normally shaped fish than either of these forms. The head was narrower and less flattened, the preoperculum wider, but its two jaws had nearly the same length. The ribs, as also the neural and humeral spines, were longer, the tail more forked, and the soft dorsal fin much longer. In fact, it was a more compressed type, probably a far better swimmer than its living congeners, as might be expected, if the smallness of the adhesive disk is taken into account." (Storrs.) This form (*Echeneis glaronensis*, Wellstein) is made the type of the genus *Opisthomyzon*, Cope, the name referring to the posterior portion of the small disk. The vertebrae in *Opisthomyzon* are 10 + 13 = 23.

Conce  
pertinen  
"The  
of know  
*Caranx*,  
of its st  
also had  
tegratin  
*Scombrid*  
just 24.  
was rep  
25 = 12 +  
with the  
colored l  
flattened  
consider  
The very  
ignored,  
to assum  
longer de  
disk in pl  
of the ge  
almost in  
and in a l  
probably  
it departs  
and rather  
*noidea*.  
head, of a  
Neverthel  
extreme n  
view of al  
constitut  
sistently o  
order of te  
ment of 3  
is sufficien  
1 or 2 such  
and any r  
atrophy of  
depression  
or left alte  
are manife  
torial disk  
resist the  
vaguely re  
modificatio  
related wit  
But whatev

Concerning the relations of this family, Dr. Gill has the following pertinent remarks:

“The family of *Scomberoides* was constituted by Cuvier for certain forms of known organization, among which were fishes evidently related to *Caranx*, but which had free dorsal spines. In the absence of knowledge of its structure, the genus *Elacate* was approximated to such because it also had free dorsal spines. Dr. Günther conceived the idea of disintegrating this family, because, inter alia, the typical *Scomberoides* (family *Scombridae*) had more than 24 vertebrae and others (family *Carangidae*) had just 24. The assumption of Cuvier as to the relationship of *Elacate* was repeated, but inasmuch as it has ‘more than 24 vertebrae’ (it has  $25 = 12 + 13$ ) it was severed from the free-spined *Carangidae* and associated with the *Scombridae*. *Elacate* has an elongated body, flattish head, and a colored longitudinal lateral band; *Echeneis* has also an elongated body, flattened head, and a longitudinal lateral band; therefore *Echeneis* was considered to be next allied to *Elacate* and to belong to the same family. The very numerous differences in structure between the two were entirely ignored, and the reference of the *Echeneis* to the *Scombridae* is simply due to assumption piled on assumption. The collocation need not, therefore, longer detain us. The possession by *Echeneis* of the anterior oval cephalic disk in place of a spinous dorsal fin would alone necessitate the isolation of the genus as a peculiar family. But that difference is associated with almost innumerable other peculiarities of the skeleton and other parts, and in a logical system it must be removed far from the *Scombridae*, and probably be endowed with subordinal distinction. In all essential respects it departs greatly from the type of structure manifested in the *Scombridae* and rather approximates—but very distantly—the *Gobioidea* and *Blenioidea*. In those types we have in some a tendency to flattening of the head, of anterior development of the dorsal fin, a simple basis cranii, etc. Nevertheless, there is no close affinity nor even any tendency to the extreme modification of the spinous dorsal exhibited by *Echeneis*. In view of all these facts *Echeneis*, with its subdivisions, may be regarded as constituting not only a family but a suborder. \* \* \* Who can consistently object to the proposition to segregate the *Echeneididae* as a suborder of teleocephaleous fishes? Not those who consider that the development of 3 or 4 inarticulate rays (or even less) in the front of the dorsal fin is sufficient to ordinarily differentiate a given form from another with only 1 or 2 such. Certainly the difference between the constituents of a disk and any rays or spines is much greater than the mere development or atrophy of articulations. Not those who consider that the manner of depression of spines, whether directly over the following, or to the right or left alternately, are of ordinal importance; for such differences again are manifestly of less morphological significance than the factors of a suctorial disk. Nevertheless, there are doubtless many who will passively resist the proposition because of a conservative spirit, and who will vaguely recur to the development of the disk as being a ‘teleological modification,’ and as if it were not an actual fact and a development correlated with radical modifications of all parts of the skeleton at least. But whatever may be the closest relations of *Echeneis*, or the systematic

value of its peculiarities, it is certain that it is not allied to *Elacate* any more than to others of the hosts of Scombroid, Percoid, and kindred fishes, and that it differs in toto from it, notwithstanding the claims that have been made otherwise. It is true that there is a striking resemblance, especially between the young—almos' as great, for example, as that between the placental mouse and the marsupial *Antechinomys*—but the like is entirely superficial, and the scientific ichthyologist should be no more misled in the case than would the scientific therologist by the likeness of the marsupial and placental mammals."

- a. Body very slender, the vertebrae 14 + 16 = 30; ventrals narrowly adnate to abdomen; lower jaw produced in a flap; pectorals acute, with flexible rays. PHTHEIRICHTHYS, 833.  
 b. Laminae 10 only. ECHENEIS, 834.  
 bb. Laminae 20 to 28. ECHENEIS, 834.  
 aa. Body rather robust, the vertebrae 12 + 15 = 27; ventrals broadly adnate to abdomen; lower jaw not produced; pectorals rounded. REMILEGIA, 835.  
 c. Laminae 24 to 27. REMILEGIA, 835.  
 cc. Laminae 16 to 20. REMILEGIA, 835.  
 d. Pectoral rays soft and flexible. REMORA, 836.  
 dd. Pectoral rays stiff and ossified. RHOMBOCHIRUS, 837.

### 833. PHTHEIRICHTHYS, Gill.

*Phtheirichthys*, GILL, Proc. Ac. Nat. Sci. Phila. 1862, 239 (*lineata*).

Disk with 10 laminae; palatines with sharp teeth; teeth in pairs, uniform in all ages; otherwise as in *Echeneis*. A single species, found attached to spearfishes and Barracudas. (*ψθεῖρ*, a louse; *ἰχθύς*, fish.)

### 2602. PHTHEIRICHTHYS LINEATUS (Menzies).

Head 5; disk twice as long as broad, its length  $4\frac{1}{2}$  in body. D. X-33; A. 33. Lower jaw very narrow, much projecting. Body blackish, with 2 whitish lateral bands; all the fins white-margined. Tropical seas, ranging north to South Carolina and Pensacola; rather rare. (*lineatus*, striped.)

*Echeneis lineata*, MENZIES, Trans. Linn. Soc. London, 1, 1791, 187, pl. 17, fig. 1, Pacific Ocean between the tropics; GÜNTHER, Cat., II, 382, 1860.

*Echeneis tropica*, EUPHRASEN, Nya Handl., XII, 317, 1791, Atlantic between the Tropics.

*Echeneis apicalis*, POEY, Memorias, II, 254, 1861, Cuba. (Coll. Poey.)

*Echeneis sphyraenarum*, POEY, Memorias, II, 255, 1861, Cuba, on Barracudas. (Coll. Poey.)

*Phtheirichthys lineatus*, JORDAN & GILBERT, Synopsis, 969, 1883.

### 834. ECHENEIS (Artedi) Linnaeus.

*Echeneis* (ARTEDI) LINNAEUS, Syst. Nat., Ed. x, 260, 1758 (*naucrates*).

*Leptecheneis*, GILL, Proc. Ac. Nat. Sci. Phila. 1864, 60 (*naucrates*); the name *Echeneis* being transferred to *E. remora*, the only species known to Artedi.

Body comparatively elongate, the vertebrae 14 + 16 = 30; disk long, of 20 to 28 laminae; pectoral pointed, its rays soft and flexible; soft dorsal and anal long, of 30 to 41 rays each; caudal lunate in the adult, convex in the young. Species of wide distribution, attaching themselves mainly to sea turtles and large fishes. (*ἐχενήϊς*, an ancient name, from *ἔχω*, to hold back; *ράβδος*, a ship.)

a. Disk of 22 to 26 laminae (rarely 21 or 28), its length less than  $\frac{1}{2}$  body.

NAUCRATES, 2603.

aa. Disk of 20 or 21 laminae, its length more than  $\frac{1}{2}$  body.

NAUCRATEOIDES, 2604.

2603. *ECHENEIS NAUCRATES*, Linnaeus.

(SHARK-SUCKER; PEGA; PEGADOR; SUCKING-FISIL.)

Head  $5\frac{1}{2}$ ; depth 11 to 12. D. XXII to XXVIII (rarely XXI)—32 to 41; A. 31 to 38. Breadth between pectorals  $7\frac{1}{2}$ ; disk 4 to 5 in body; eye 5 in head; snout  $2\frac{1}{2}$ ; maxillary 3; from angle of mouth to tip of lower jaw  $2\frac{3}{4}$ ; pectoral  $1\frac{3}{4}$ ; ventrals  $1\frac{1}{2}$ ; middle caudal rays  $1\frac{3}{4}$ ; highest anal ray 2; highest dorsal ray  $2\frac{1}{2}$ ; width of disk  $2\frac{1}{2}$  in its length; base of dorsal  $2\frac{1}{2}$ , anal  $2\frac{1}{2}$ , in body. Body elongate, subterete, slender. Lower jaw strongly projecting, the tip flexible; maxillary reaching nostril; teeth uniform in the adult, the young with series of small slender teeth in advance of the others; gill rakers short and slender, about equal to pupil; vertical fins low. Anal rays higher than dorsal anteriorly; pectorals reaching very slightly past tips of ventrals; origin of ventral spine under middle of pectoral base; inner rays of ventral fins narrowly adnate to the abdomen; dorsal and anal commencing and ending opposite each other; caudal with the middle rays produced in the young, the fin becoming emarginate or lunate with age. Color brownish; belly dark, like the back, as usual in this family; sides with a broad stripe of darker edged with whitish extending through eye to snout; caudal black, its outer angles whitish; pectorals and ventrals black, sometimes bordered with pale; dorsal and anal broadly edged with white anteriorly; adult nearly uniform dark brown, not paler below. Warm seas, universally distributed; common north to Cape Cod and occasionally to San Francisco, attaching itself to turtles and to large fishes. This species is very common in the tropics, being found attached to sharks, groupers, or any other large fish, without regard to species. Few large sharks at Key West are without them. They are often caught with hook and line from the wharf, where they frequently forsake their host to take the bait. Lütken's remark that only *Remora remora* has been recorded from sharks is no longer true. Several writers have recognized 2 species of *Echeneis* proper—*naucrates*, with 22 to 26 laminae, the disk 4 to 5 in body, and *naucrateoides* (= *albicauda* = *holbrookii* = *lineatus*), in which the disk is longer,  $3\frac{3}{4}$  to 4 in body, but composed of fewer, 20 or 21, laminae. The latter form is rather common on our coast, the specimens from Key West above mentioned having 21. We doubt the existence of any permanent difference between the two, but provisionally retain *Echeneis naucrateoides* as a species distinct from *Echeneis naucrates* until more complete comparison can be made. (*naucrates*, a pilot; *ναύς*, ship; *ἡγάρτω*, to govern, guide.)

*Echeneis naucrates* (misprint for *naucrates*), LINNÆUS, Syst. Nat., Ed. x, 261, 1758, "in Pelago Indico;" GÜNTHER, Cat., II, 384, 1860; JORDAN & GILBERT, Synopsis, 416, 1883.

*Echeneis albicauda*, MITCHELL, Amer. Monthly Mag., II, 1817, 244, New York.

*Echeneis lunata*, BANCROFT, Proc. Comm. Zool. Soc., I, 1830, 134, Kingston, Jamaica.

† *Echeneis vittata*, LOWE, Proc. Zool. Soc. Lond. 1839, 89, Madeira.

*Echeneis fasciata*, GRONOW, Ed. Gray, 92, 1854, Mediterranean Sea.

*Leptecheneis naucrates*, GILL, Proc. Ac. Nat. Sci. Phila. 1864, 60.

*Echeneis vittata*, RÜPPELL, Neue Wirb. Fische, 82, 1855, Red Sea.

*Echeneis guaiacan*, POEY, Memorias, II, 248, 1861, Cuba; young. (Coll. Poey.)

*Echeneis verticalis*, POEY, Memorias, II, 253, 1861, Cuba; young.

*Echeneis metallica*, POEY, Memorias, II, 252, 1861, Cuba; D. XXIII, 40; A. 37; large specimens, metallic green, the bands faint. (Coll. Poey.)

*Echeneis fusca*, GRONOW, Cat. Fish., 92, 1854; after *E. naucrates*, L.

#### 2604. ECHENEIS NAUCRATEOIDES, Zuiew.

Head 5; depth 11. D. XX or XXI-32 to 35; A. 33 to 35. Disk  $3\frac{1}{2}$  to  $3\frac{3}{4}$  in total, twice width of body between pectorals. In all other respects essentially as in *Echeneis naucrates*, the disk longer, but composed of fewer laminae, the laminae being farther apart. Color of *Echeneis naucrates*. Cape Cod to West Indies, common on our south Atlantic coast; specimens before us from Key West. (*naucrates*, *ναυκράτης*, a pilot; *εἶδος*, resemblance.)

*Echeneis naucrateoides*, ZUIEW, Nova Acta Acad. Sci. Imp. Petropol., IV, 1789, 279, no locality.

*Echeneis lineata*, HOLBROOK, Ichth. S. C., 102, 1860, Charleston, South Carolina; not of MENZIES.

*Echeneis holbrooki*, GÜNTHER, Cat., II, 382, 1860, Jamaica; D. XIII, 35; A. 33.

*Leptecheneis naucrateoides*, GILL, l. c., 61.

#### 835. REMILEGIA, Gill.

*Remilegia*, GILL, Proc. Ac. Nat. Sci. Phila. 1864, 61 (*australis*).

This genus differs from *Remora* chiefly in the length of the sucking disk, which has 24 to 27 laminae; the soft dorsal and anal are proportionately short. (A metathesis for *remeligo*, the delayer or hinderer.)

#### 2605. REMILEGIA AUSTRALIS (Bennett).

D. XXVII-22; A. 21 to 23. The length of the disk is  $2\frac{1}{2}$  in the total, the width of the body between the pectorals  $5\frac{1}{2}$ . Caudal truncated; dorsal and anal fins not continued to the caudal. Color brown. This species has the general habit of *E. remora*, but may be readily distinguished from all the others by the extraordinary size of the disk, which is elongate, subelliptical, obtusely rounded anteriorly and posteriorly, and formed by 27 pairs of laminae; it extends backward beyond the vertical from the tip of the ventrals, and its length is  $2\frac{1}{2}$  in the total. The spines with which the single laminae are armed are less conspicuous than in the other species, and do not offer the same resistance to the touch. There is a large posterior portion of the disk which is not provided with laminae, but quite smooth. The width of the disk, taken between the extremities of the bony laminae, is  $\frac{1}{2}$  of its length; the membranaceous margin is bent upward. The head and the body below the disk are depressed, and their height is  $9\frac{1}{2}$  in the total length, whilst the width between the pectorals is  $5\frac{1}{2}$  in it. The body between the disk and the vertical fins is quadrangular, tapering posteriorly. The upper jaw is subtruncated, and overreached by the lower, which is much narrower; both are armed with a broad band of villiform teeth, and with an outer series of larger ones on

the s  
teeth  
denti  
from  
length  
all th  
and 4  
the p  
low g  
disk i  
a thic  
anal i  
tically  
ceived  
caviti  
seas;  
Hygo  
*Echene*  
*Echene*  
bal

*Remora*  
*Echeneis*  
rest  
*Remora*  
*Remora*

Bod  
lamina  
moder  
selves  
"hold

REM  
a  
aa

Head  
tebrae  
fins ro



the sides; the vomerine and palatine bones have a continuous band of teeth, narrowest on the vomer; the tongue is hard, cartilaginous, and destitute of teeth. The cleft of the mouth reaches only to the vertical from the nostril; the eye is small. The pectoral is rounded and small, its length being  $\frac{1}{3}$  of the total; the ventrals are slightly pointed, and, as in all the species of the genus, composed of 1 spine, hidden in the skin, and 4 soft rays; they are inserted immediately behind the vertical from the pectoral, which they equal in length; they can be received in a shallow groove on the abdomen. The distance between the dorsal and the disk is  $3\frac{1}{2}$  in the length of the latter; the dorsal is low, and enveloped in a thick membrane. The caudal is truncated when stretched out. The anal is very similar to the dorsal, and its origin and termination fall vertically below those of the latter. The scales are minute, and can be perceived only by the aid of a magnifier; they are embedded in pore-like cavities. (Günther: description of type of *Echeneis scutata*.) Tropical seas; rare; recorded by Dr. Lütken from  $10^{\circ}$  N.,  $39^{\circ}$  W. (Coll. Capt. V. Hygom) from a dolphin; not seen by us. (*australis*, southern.)

*Echeneis australis*, BENNETT, Narr. Whaling Voyage, II, 273, pls. 24-26, 1840.

*Echeneis scutata*, GÜNTHER, Ann. Mag. Nat. Hist. 1860, 401, pl. 10, f. B, Ceylon (Coll. Dr. Sibald); GÜNTHER, Cat. Fish., II, 381, 1860; LÜTKEN, Vid. Medd. Kjöbenh. 1875, 42.

### 836. REMORA, Gill.

(REMORAS.)

*Remora*, GILL, Proc. Ac. Nat. Sci. Phila. 1862, 239 (*remora*).

*Echeneis*, GILL, Proc. Ac. Nat. Sci. Phila. 1864, 60 (*remora*); not *Echeneis*, GILL, 1862, restricted to *naucrates*.

*Remoropsis*, GILL, Proc. Ac. Nat. Sci. Phila. 1864, 60 (*brachypterus*).

*Remorina*, JORDAN & EVERMANN, Check-List Fishes, 490, 1896 (*albescens*).

Body rather robust, the vertebrae  $12 + 15 = 27$ ; disk shortish, of 13 to 18 laminae; pectoral rounded, its rays soft and flexible; soft dorsal and anal moderate, of 20 to 30 rays; caudal subtruncate. Species attaching themselves to large fishes, especially to sharks. (*Remora*, an ancient name, "holding back.")

REMORA:

a. Laminae about 18; soft dorsal with 23 rays.

REMORA, 2606.

aa. Laminae 13 to 16.

REMORINA:

b. Dorsal rays XIII, 22.

ALBESCENS, 2607.

REMOROPSIS (*Remora*; 5ψis, appearance):

bb. Dorsal rays XIV, XVI, 29 to 32.

BRACHYPTERA, 2608.

#### Subgenus REMORA.

2606. REMORA REMORA (Linnaeus).

(REMORA.)

Head 4; disk  $2\frac{1}{2}$ ; width between pectorals  $5\frac{1}{2}$ . D. XVIII-23; A. 25; vertebrae  $12 + 15$ . Body comparatively robust, compressed behind. Pectoral fins rounded, short, and broad, their rays short and flexible; ventral fins

adnate to the abdomen for more than  $\frac{1}{2}$  the length of their inner edge. Tip of lower jaw not produced into a flap; head broad, depressed; disk longer than the dorsal or the anal fin; maxillary scarcely reaching front of orbit. Caudal lunate; vertical fins rather high; pectoral  $\frac{3}{4}$  length of head. Color blackish, nearly uniform above and below. Length 15 inches. Warm seas, north to New York and San Francisco, where it is not rare; usually found attached to large sharks; very common in the West Indies; more robust than *Echeneis naucrates*, and reaching a smaller size.

*Echeneis remora*, LINNÆUS, Syst. Nat., Ed. x, 260, 1758, "in Pelago Indico;" GÜNTHER, Cat., II, 378, 1860; LÜTKEN, Vid. Medd. Kjöbenh. 1875, 38; JORDAN & GILBERT, Synopsis, 417, 1883.

*Echeneis equalipeta*, DALDORF, Skrivt. Naturh. Selsk., II, 1797, 157, Atlantic Ocean between the tropics; GÜNTHER, Cat., II, 377, 1860.

*Echeneis jacobæa*, LOWE, Proc. Zool. Soc. London 1839, 89, Madeira.

*Echeneis remoroides*, BLEEKER, Batöi, II, 70, Batoe.

*Echeneis parva*, GRONOW, Cat. Fish., Ed. Gray, 92, 1854, no locality; after *E. remora*, L.

*Echeneis postica*, POEY, Memorias, II, 255, 1861, Havana. (Coll. Poey.)

*Remora jacobæa*, GILL, Proc. Ac. Nat. Sci. Phila. 1862, 239.

#### Subgenus REMORINA, Jordan & Evermann.

#### 2607. REMORA ALBESCENS (Temminck & Schlegel).

Length of disk  $3\frac{1}{2}$  to  $3\frac{1}{2}$  in total length; width between pectorals 5 to  $5\frac{1}{2}$ ; number of laminae on disk 13 or 14. D. XIII-22; A. 22. Angle of mouth in the vertical from the third lamina of the disk. Length of ventral fins equal to the distance between root of pectoral and posterior margin of eye. Color uniform grayish brown. (Günther.) Tropical Pacific, straying to America; a specimen taken at La Paz, Gulf of California (Streets), and 1 in the Gulf of Mexico (Bean). (*albescens*, whitish.)

*Echeneis albescens*, TEMMINCK & SCHLEGEL, Fauna Japonica, Poiss., 272, pl. 120, fig. 3, 1842, Japan; GÜNTHER, Cat., II, 377, 1860; STREETS, Bull. U. S. Nat. Mus., VII, 54, 1877.

*Remora albescens*, JORDAN, Cat. Fishes, 66, 1885.

#### Subgenus REMOROPSIS, Gill.

#### 2608. REMORA BRACHYPTERA (Lowe).

Head nearly 4; width between pectorals  $6\frac{1}{2}$ . D. XIV to XVI-29 to 32; A. 25 to 30. Body robust, the greatest depth nearly twice the length of the short pectoral fins; disk shorter than base of dorsal, rather broad; upper jaw angular. Caudal nearly truncate. Light brown, darker below, fins paler. Warm seas, occasionally north to Cape Cod. (*βραχύς*, short; *πτερόν*, fin.)

*Echeneis brachyptera*, LOWE, Proc. Zool. Soc. London 1839, 89, Madeira; GÜNTHER, Cat., II, 378, 1860; JORDAN & GILBERT, Synopsis, 417, 1883.

*Echeneis sexdecimlamellata*, EYDOUX & GERVAIS, Voy. Favorite, v, 77, pl. 31, 1839, Indian Ocean?

*Echeneis quatuordecimlaminatus*, STORER, Rept. Fishes Mass., 155, 1839, Holmes Hole.

*Echeneis pallida*, TEMMINCK & SCHLEGEL, Fauna Japonica, Poiss., 271, pl. 120, fig. 2, 3, 1842, Japan.

*Echeneis nieuhofti*, BLEEKER, Sumatra, II, 279, Sumatra.

*Remoropsis brachypterus*, GILL, Proc. Ac. Nat. Sci. Phila. 1864, 60.

Rhomboid

This g  
of the pe  
rays all  
the usua  
One spec

Head 4  
to 23; A.  
ing to the  
Disk very  
forward b  
angles. I  
and a spot  
on species

*Echeneis or*  
GÜNTHE  
*Echeneis tet*

A large  
ships, show  
*Batrachoid*  
short or w  
tion is less  
general, de  
is wanting  
The bones  
often inse  
The group  
*Tranoscopo*  
Dr. Gill ob  
group and r  
don, and L  
of this gro  
*Trachinida*

a. Mouth h  
rays  
gills  
b. Snow  
c.

837. RHOMBOCHIRUS, Gill.

*Rhombochirus*, GILL, Proc. Ac. Nat. Sci. Phila. 1863, 88 (*ostochir*).

This genus agrees with *Remora* in every respect excepting the structure of the pectoral fins. These are short and broad, rhombic in outline, the rays all flat, broad and stiff, being partially ossified, although showing the usual articulation; upper rays of pectoral broader than the others. One species known. (*ῥόμβος*, rhoub; *χείρ*, hand.)

2609. RHOMBOCHIRUS OSTEOCHIR (Cuvier).

Head  $4\frac{1}{2}$  in length; disk  $2\frac{1}{2}$ ; width between pectorals 5. D. XVIII-21 to 23; A. 20 or 21; P. 20. Mouth very small, maxillary not nearly reaching to the line of the orbit; outer series of teeth longer than the others. Disk very large, broader and rougher than in *Remora remora*, extending forward beyond the tip of the snout. Caudal fin emarginate, with rounded angles. Light brown; underside of head, ventral line, part of ventrals and a spot on pectorals pale. West Indies north to Cape Cod; parasitic on species of *Tetrapturus*; rather rare. (*ὀστρέον*, bone; *χείρ*, hand.)

*Echeneis ostochir*, CUVIER, Règne Animal Ed. 2, vol. II, 348, 1829, no locality given; GÜNTHER, Cat., II, 381, 1860; JORDAN & GILBERT, Synopsis, 418, 1883.

*Echeneis tetrapturorum*, POEY, Memorias, II, 256, 1858, Cuba. (Coll. Poey.)

Group TRACHINOIDEA.

(THE TRACHINOID FISHES.)

A large group of transitional forms, some of them of doubtful relationships, showing affinities with the *Percoidae* on the one hand and with the *Batrachoididae* and *Blennoidea* on the other. In general, the spinous dorsal is short or weak, the soft dorsal long and similar to the anal, and the squamation is less complete and less ctenoid than in the *Percoidae*. The skull is, in general, depressed, with the supraocular crest low, and the suborbital stay is wanting, although in some genera the suborbital bones are enlarged. The bones of the skull are not strongly armed, and the ventral fins are often inserted well forward, and they are sometimes reduced in size. The group is divided by Dr. Gill into *Percophidoidea*, *Trachinoidea*, and *Tranoscopoidea*. The two latter groups are natural and related, but, as Dr. Gill observes, "the *Percophidoidea* are undoubtedly a heterogeneous group and need a thorough revision." The relations of *Bathymaster*, *Trichodon*, and *Latilus* especially are uncertain. Several of the leading families of this group are confined to the South Temperate Zone, and none of the *Trachinidae* occurs within our limits.

- a. Mouth horizontal or moderately oblique, the lips not fringed; eyes lateral; ventral rays I, 5, their insertion more or less before the pectorals; suborbitals moderate; gills 4, a slit behind the fourth.
- b. Snout subconic, not prolonged and spatulate; ventrals not widely separated.
- c. Body covered with scales; dorsal spines flexible.
- d. Lateral line complete; caudal fin forked; vertebrae 24 to 27.

MALACANTHIDÆ, CXO.

- dd. Lateral line incomplete, running close to the back; caudal rounded or lanceolate; dorsal fin continuous.
- e. Vertebrae about 27; scales cycloid; maxillary more or less dilated behind, with a supplemental bone; middle rays of ventrals longest. *OPISTHOGNATHIDÆ, CXCII.*
- ee. Vertebrae about 50; scales ctenoid; maxillary not dilated, without supplement bone; inner rays of ventrals longest. *BATHYMASTERIDÆ, CXCIII.*
- cc. Body naked; snout short; mouth very large, the maxillary much produced behind; jaws with sharp canines; lateral line well developed; dorsals 2; caudal forked. *CHIASSODONTIDÆ, CXCIII.*
- bb. Snout much prolonged and spatulate; ventrals widely separated; body scaly or naked; lateral line near the back; dorsal usually divided. *CHLENICHTHYIDÆ, CXCIV.*
- aa. Mouth vertical, the lips fringed.
- f. Eyes lateral; gills 4, a slit behind the last; preopercle armed; body naked, compressed; caudal lunate, on a slender peduncle; vertebrae about 48. *TRICHODONTIDÆ, CXCIV.*
- ff. Eyes superior; gills more or less reduced, usually  $3\frac{1}{2}$ , the last slit small or wanting; suborbitals more or less dilated; body scaly or naked.
- g. Lateral line well developed, concurrent with the back anteriorly; dorsal spines slender, not pungent; vertebrae about 25 to 30.
- h. Ventral rays I, 3. *DACTYLOSCOPIDÆ, CXCVI.*
- gg. Lateral line obscure; dorsal spines few, more or less pungent, sometimes obsolete. *URANOSCOPIDÆ, CXCVII.*

## Family CXC. MALACANTHIDÆ.

## (THE BLANQUILLOS.)

Body more or less elongate, fusiform or compressed. Head subconical, the anterior profile usually convex; suborbital without bony stay; the bones not greatly developed; cranial bones not cavernous; opercular bones mostly unarmed. Mouth rather terminal, little oblique; teeth rather strong; no teeth on vomer or palatines; the premaxillary usually with a blunt posterior canine, somewhat as in the *Labridæ*; premaxillaries protractile; maxillary without supplemental bone, not slipping under the edge of the preorbital. Gills 4, a long slit behind the fourth; pseudo-branchiæ well developed; gill membranes separate, or more or less united, often adherent to the isthmus; lower pharyngeals separate. Scales small, ctenoid; lateral line present, complete, more or less concurrent with the back; dorsal fin long and low, usually continuous, the spinous portion always much less developed than the soft portion, but never obsolete; anal fin very long, its spines feeble and few; caudal fin forked; tail diphyccercal; ventrals thoracic or subjugular, I, 5, close together; pectoral fins not very broad, the rays all branched; vertebrae in normal or slightly increased number (24 to 30). Pyloric cæca few or none. Fishes of the temperate and tropical seas, some of them reaching a large size. Genera about 6; species about 8 to 10, mostly American. The relationships of the family are obscure, and it may be that the genera here associated are not really closely allied. (*Malacanthidæ*, Günther, Cat., III, 359, 1861; *Trachinidæ*, part, Günther, Cat., II, 225-264, 1860.)

MALACA  
a. Verte  
b. Soft

aa. Vert  
CAULO  
c. So

d.

LA

cc. S

e.

Malacant

Body  
the jaw  
continu  
anal ver  
outer ser  
at the p  
preoper  
tebrae in  
America

Head.  
illary 2  
caudal  
long and  
file of h  
there is  
dorsal.  
posterior  
growing  
at the si  
largest;  
lower ja  
jaw ante  
a single  
5 + 7. T  
lower ja  
and low,

MALACANTHINÆ:

a. Vertebrae 24; preopercle entire.

b. Soft dorsal and anal extremely long, each with more than 40 rays; preopercle entire; form slender; scales very small. MALACANTHUS, 838.

aa. Vertebrae more than 24; preopercle more or less serrate.

CAULOLATILINÆ:

c. Soft dorsal and anal moderate, each with 22 to 27 soft rays; preopercle serrate; scales rather small; form robust.

d. Upper jaw with posterior canines; dorsal spines graduated. CAULOLATILUS, 839.

LATHLINÆ:

cc. Soft dorsal and anal short, each of 13 to 15 soft rays; preopercle denticulate; scales small; form robust.

e. Nape with a large adipose appendage; a fleshy prolongation on each side of the labial fold, extending forward behind angle of mouth. LOPHOLATILUS, 840.

838. MALACANTHUS, Cuvier.

(MATAJUELO BLANCO.)

*Malacanthus*, CUVIER, Règne Animal, Ed. 2, vol. II, 205, 1829 (*plumieri*).

Body elongate, slightly compressed; cleft of mouth horizontal, with the jaws equal; eyes lateral; scales very small, minutely ciliated; one continuous dorsal, with the first 4 to 6 rays not articulated; dorsal and anal very long; pectoral rays all branched; jaws with villiform teeth; an outer series of stronger teeth, some of them canine-like, and with a canine at the posterior extremity of the intermaxillary; no teeth on the palate; preopercle entire; opercle with a spine; gill rakers little developed; vertebrae in small number,  $10 + 14 = 24$ . One species, a shore fish of tropical America. (*μαλακός*, soft; *ἄκανθα*, spine.)

2610. MALACANTHUS PLUMIERI (Bloch).

(MATAJUELO BLANCO.)

Head  $3\frac{1}{2}$ ; depth  $6\frac{1}{2}$ . D. VI, 49; A. 48; scales 14–130–30; eye  $5\frac{1}{2}$  in head; maxillary  $2\frac{1}{2}$ ; snout  $2\frac{1}{2}$ ; P. 2; longest dorsal rays 3, equal to anal ray; upper caudal lobe  $1\frac{1}{2}$ . Body elongate, little compressed. Head moderately long and pointed; eye placed high; interorbital flat, as wide as eye; profile of head obliquely straight from tip of snout to above nostril, where there is a slight angle formed, thence nearly horizontally straight to dorsal. Mouth large, maxillary reaching slightly past the vertical from posterior nostril; jaws equal; a band of villiform teeth in upper jaw growing broader anteriorly, and another row of small, even, conical teeth at the sides, and 6 well-developed canines in front, the 2 outer ones the largest; a canine on premaxillary at angle of mouth; villiform teeth in lower jaw not extending very far back; large recurved canines on side of jaw anteriorly, small conical teeth in front and on sides posteriorly, with a single large canine at angle of mouth; gill rakers rudimentary, about  $5 + 7$ . Top of head forward from above middle of eye, preorbital, and lower jaw, naked; fins without scales. Dorsal and anal similar, long and low, continuous; pectoral reaching past tips of ventrals to front of

anal; ventrals not reaching to vent, origin of ventral spine slightly behind base of pectoral; caudal forked, the lobes elongate, sometimes produced into a filament. Color in spirits, uniform, pale olive brown above, white below; fins light brownish; no distinct markings. Length 15 inches. West Indies, rather common; used as food. Here described from specimens from Havana. (Named for Père Plumier, of Martinique.)

*Mateuelo blanco*, PARRA, Dif. Piezas Hist. Nat. Cuba, 22, tab. 13, f. r. 1787, Cuba.

*Coryphæna plumieri*, BLOCH, Ichthyol., v, 119, pl. 175, 1787, Martinique; from a drawing by PLUMIER.

*Malacanthus truchinus*, VALENCIENNES, in CUVIER, Règne Animal, pl. 90, fig. 3.

*Sparus oblongus*, BLOCH & SCHNEIDER, Syst. Ichth., 283, 1801; after PARRA.

*Malacanthus plumieri*, CUVIER & VALENCIENNES, Hist. Nat. Poiss., XIII, 319, 380, 1839, specimens from San Domingo; GÜNTHER, Cat., III, 358, 1861.

### 839. CAULOLATILUS, GILL.

(BLANQUILLOS.)

*Caulolatilus*, GILL, Proc. Ac. Nat. Sci. Phila. 1862, 240 (no diagnosis), and GILL, Proc. Ac. Nat. Sci. Phila. 1865, 66 (*chrysops*).

*Dekaya*, COOPER, Proc. Cal. Ac. Sci. 1864, 70 (*princeps*), not *Dekaya*, MILNE-EDWARDS & HAIME, 1851, a genus of corals.

Body elongate, subfusiform, not strongly compressed, heavy forward, tapering to a rather slender caudal peduncle; profile of head strongly arched; mouth moderate, little oblique, the jaws nearly equal; lips thick; maxillary narrow, not slipping under the preorbital; teeth in villiform bands, preceded by a row of stronger acute teeth; posterior teeth in each jaw canine-like, directed forward; posterior canines of upper jaw largest; no teeth on vomer or palatines; preopercle pectinate, the teeth nearly even; opercle with a blunt, flat spine; eyes large, lateral; gill membranes slightly connected, forming a fold across the isthmus, with which they are narrowly joined; branchiostegals 6; gill rakers short and stout; nostrils double, round, close together; scales small, firm, ctenoid; lateral line continuous, concurrent with the back; dorsal with 7 to 9 slender, pointed, graduated spines and 22 to 27 soft rays; anal similar to soft dorsal, with 1 or 2 small spines and more than 20 soft rays; caudal fin forked; ventral fins thoracic; no adipose appendage at the nape; vertebrae 12+15=27. Large fishes of the warm seas of America; valued as food. (*καυλός*, stem; *Latilus*; being distinguished from *Latilus* by the many rays.)

a. Scales small, about 125 in the lateral line, about 50 in a transverse series.

b. Eye large,  $4\frac{1}{2}$  in the head; depth 4 in length; scales 16-125-40.

PRINCEPS, 2611.

bb. Eye small, 6 in head; depth  $3\frac{1}{2}$  in body; scales 13-120-35.

MICROPS, 2612.

ac. Scales larger, about 108 in the lateral line, about 25 in a transverse series; scales 12-108-25.

CYANOPS, 2613.

### 2611. CAULOLATILUS PRINCEPS (Jenyns).

(BLANQUILLO; WHITE-FISH.)

Head  $3\frac{1}{2}$ ; depth 4. D. IX, 24; A. II, 23; scales 16-125-40. Flesh of the occiput becoming thick with age, as in *Harpe*. Eye large, about  $\frac{1}{2}$  the convex interorbital space,  $4\frac{1}{2}$  in head; maxillaries reaching front of eye;

teeth rather nearly equal; cheeks and between the torals, the the longer torals false bluish reflection, olive, tinged anal green of the Pacific abundant about tance. Length which the *Caulolatilus* *Latilus princeps* Archipelago *Dekaya anona* *Caulolatilus* *Xantus* *Caulolatilus* *Caulolatilus* GILBERT.

Head  $3\frac{1}{2}$ ; series, 120 eye  $5\frac{1}{2}$  in head dorsal spine robust; upper snout divided separated lateral scarcely included; like a large opercle fin opercle; similar anal similar vent; original from slightly the below the at base, d Mexico, in from the full species small; *Caulolatilus* Florida. *Caulolatilus* CUVIER

teeth rather strong; preopercle finely, evenly, and acutely serrate behind, nearly entire below; preopercle, interopercle, and preorbital naked; cheeks and opercles scaly; top of head scaled on the median line to between the eyes; dorsal spines flexible; ventrals slightly behind the pectorals, the outer rays longest; caudal moderately forked, the upper lobe the longer; caudal peduncle short and slender, abruptly contracted; pectorals falcate, longer than caudal,  $\frac{5}{8}$  length of the head. Olivaceous, with bluish reflections; brownish above, greenish below; fins light greenish olive, tinged with bluish and orange, the colors always pale; dorsal and anal greenish, with a bluish band near the tip; axil dusky. Rocky islands of the Pacific coast from Monterey southward to the Galapagos; abundant about the Santa Barbara Islands; a food fish of considerable importance. Length 40 inches. We are unable to detect any differences by which the Californian form, *Caulolatilus anomalus*, can be separated from *Caulolatilus princeps*. (*princeps*, a leader.)

*Latilus princeps*, JENYNS, Zool. Beagle, Fishes, 52, pl. 11, 1840, Chatham Island, Galapagos Archipelago (Coll. Charles Darwin); GÜNTHER, Cat., 11, 253, 1860.

*Dekaya anomalus*, COOPER, Proc. Cal. A. C. Sci. 1864, 70, coast of Southern California.

*Caulolatilus affinis*, GILL, Proc. Ac. Nat. Sci. Phila. 1865, 68, Cape St. Lucas. (Coll. John Xantus.)

*Caulolatilus princeps*, GILL, l. c. 68.

*Caulolatilus anomalus*, GILL, l. c. 68; STREETS, Bull. U. S. Nat. Mus., VII, 48, 1877; JORDAN & GILBERT, Synopsis, 625, 1883.

#### 2612. CAULOLATILUS MICROPS, Goode & Bean.

Head  $3\frac{1}{2}$ ; depth  $3\frac{1}{2}$ . D. VII, 25; A. I, 23; scales 105 counting the oblique series, 120 counting the row above lateral line; transverse rows 12 + 30; eye  $5\frac{1}{2}$  in head; snout 2; maxillary  $2\frac{1}{2}$ ; pectoral  $1\frac{1}{2}$ ; ventral  $2\frac{1}{4}$ ; highest dorsal spine  $3\frac{1}{2}$ ; highest anal rays  $3\frac{1}{2}$ ; upper caudal lobe  $1\frac{1}{2}$ . Body rather robust; upper profile of head rather steep, evenly rounded from tip of snout to dorsal; nostrils small, midway between eye and tip of snout, separated by a distance equal to  $\frac{1}{2}$  diameter of pupil; mouth large, maxillary scarcely reaching to anterior margin of eye; lips thick; lower jaw included; jaws with small conical teeth, the outer row enlarged, canine-like, a large tooth on posterior end of maxillary at angle of mouth; preopercle finely and evenly serrate on its vertical limb; a broad flat spine on opercle; snout, preorbital and lower jaw naked; fins scaleless; dorsal and anal similar, long and low; pectorals reaching far past tips of ventrals to vent; origin of ventral spine about the length of 2 scales behind the vertical from pectoral base; caudal fin lunate when spread, its upper lobe slightly the longer. Color reddish, marked with yellow; a yellow band below the eye; a dark blotch in and above axil of pectoral; dorsal light at base, darker above, with many indistinct brownish spots. Gulf of Mexico, in rather deep water; not rare. Here described from a specimen from the Pensacola Snapper Banks, 26 inches in length. A rather doubtful species, perhaps not distinct from *C. cyanops* or *C. chrysops*. (*μικρός*, small; *ὄψ*, eye.)

*Caulolatilus microps*, GOODE & BEAN, Proc. U. S. Nat. Mus. 1878, 43, off Pensacola, Florida. (Coll. SILLAS STRAUSS.)

*Caulolatilus chrysops*, JORDAN & GILBERT, Synopsis, 626, 1883; not *Latilus chrysops*, CUVIER & VALENCIENNES.

## 2613. CAULOLATILUS CYANOPS,\* Poey.

(BLANQUILLO.)

Head 4 in total length. D. VII, 24; A. I, 22 (scales 10-108-25. Bean). Profile convex before the eye, not ascending to the nape; no scales on the fins; soft rays little divided; caudal slightly lunate; first caudal vertebra spoon-like, its cavity receiving the air bladder; vertebrae 12 + 15; no pyloric caeca, stomach short, air bladder large. Color greenish above, a faint, broad, interrupted brown band above the lateral line; some small brown spots above and below it; region below the eye clear blue, not very different from the color of the belly; soft dorsal brown, paler at its base, edged with orange; spinous dorsal orange. (Poey.) Coast of Cuba; not seen by us. Both this and the preceding species may be identical with *Caulolatilus chrysope*, a species described from the coast of Brazil. (καίρεος, blue; ὠψ, eye.)

1 *Latilus chrysope*, CUVIER & VALENCIENNES, Hist. Nat. Poiss., IX, 496, 1833, Brazil (Coll. M. Gay); GÜNTHER, Cat., II, 253, 1860.

*Caulolatilus cyanops*, POEY, Repertorio, I, 312, 1867, Cuba. (Coll. Poey.)

## 840. LOPHOLATILUS, Goode &amp; Bean.

(TILE-FISHES.)

*Lopholatilus*, GOODE & BEAN, Proc. U. S. Nat. Mus. 1879, 205 (*chameleonticeps*).

Body stout, somewhat compressed; mouth moderate, maxillary reaching anterior margin of the orbit; opercle and preopercle scaly, the latter finely denticulate; upper jaw with outer series of stronger teeth, behind which is a band of villiform teeth; lower jaw with a few large canines, and an inner series of small conical teeth; vomer and palatines toothless; nape with a large adipose appendage; a fleshy prolongation upon each side of the labial fold, extending backward beyond the angle of the mouth; stomach small, siphonal, barely more than a loop in the very large intestine; alimentary canal short, less than total length of the body; air bladder simple, with thick muscular walls, strongly attached to the roof of the abdominal cavity by numerous root-like appendages, resembling somewhat that of *Pogonias*. Deep-sea fishes. (λόφος, crest; *Latilus*.)

## 2614. LOPHOLATILUS CHAMELEONTICEPS, Goode &amp; Bean.

(TILE-FISH.)

Head 3; depth 3½. D. VII, 15; A. II, 13; scales 8-93-30. Body stout, somewhat compressed, its greatest width equaling length of caudal peduncle; intermaxillaries supplied with a series of from 19 to 23 canine teeth, behind which is a band of villiform teeth, widest at the symphysis; mandible with about 12 large canines; eye rather small, its diameter 6¼ in

\* The characters distinguishing *Caulolatilus chrysope* are thus given by Poey: Head 4½ in total length. D. VIII, 24; A. II, 22. Profile most gibbous behind the eye; a very bright gilded band below the eye, broader anteriorly; dorsal fin brown with irregular blue spots; axillary spot green. Coast of Brazil.

head, and  
posterior no  
of snout to  
outer rays;  
Back bluish  
blue and blu  
line of belly  
head, blue u  
of head pear  
nounced in f  
with numerou  
with blue an  
iridescent, li  
more or less  
lower border  
of this gener  
rays; dorsal g  
to the upper  
on or near th  
of bright yel  
with rosy and  
the western A  
"The tilefish  
Nantucket Sh  
some economi  
at New York, l  
bers of dead an  
Collins estima  
5,000 to 7,500 s  
Several visits v  
where these fis  
and it was tho  
were taken by  
to 73° W. The  
Colonel McDon  
(*chameleon*, χα  
*Lopholatilus cham  
tucket Shoals  
Comm. (1882) p  
647, with plate:*

Body oblong  
small eyeloid se  
dorsal fin, not ex  
the anterior pro



head, and about twice length of labial appendages; distance between posterior nostril and eye equal first anal spine, and  $\frac{1}{2}$  distance from tip of snout to anterior nostril. Caudal fin emarginate, middle rays  $1\frac{1}{2}$  in outer rays; vent under interval between fourth and fifth dorsal rays. Back bluish, with a green tinge, iridescent, changing through purplish blue and bluish gray to rosy white below, and milky white toward median line of belly; head rosy, iridescent, with red tints most abundant on forehead, blue under the eyes, cheeks fawn-colored; throat and under side of head pearly white, with an occasional tint of lemon yellow, most pronounced in front of ventrals and on anterior portion of ventral fins; back with numerous maculations of bright yellow or golden; anal purplish, with blue and rose tints, iridescent; margin of anal rich purplish blue, iridescent, like the most beautiful mother-of-pearl, this color pervading more or less the whole fin, which has large yellow maculations, the lower border rose-colored, like the belly, base of the fin also partaking of this general hue; dashes of milk white on base of anal between the rays; dorsal gray; in front of the seventh dorsal the upper third posterior to the upper two-thirds dark brown; spots of yellow, large, elongate, on or near the rays; adipose fin whitish brown or yellow, a large group of bright yellow confluent spots at the base; pectorals sepia-colored, with rosy and purplish iridescence. (Goode & Bean.) Deep waters of the western Atlantic, at times very abundant; now rare or almost extinct. "The tilefish was first observed in 1879 by fishermen fishing for cod on Nantucket Shoals. From its abundance it was thought to become of some economic importance. In March and April, 1882, vessels arriving at New York, Philadelphia, and Boston reported having passed large numbers of dead and dying fish, the majority of which were tilefish. Captain Collins estimated the area covered by dead and dying fish to be from 5,000 to 7,500 square statute miles, the number of fish to be 1,000,000,000. Several visits were made by the Fish Commission vessels to the grounds where these fishes were formerly abundant, but no specimen was obtained, and it was thought to have become extinct. In 1892 several specimens were taken by the *Grampus* in latitude  $38^{\circ}$  to  $40^{\circ}$  N., and longitude  $71^{\circ}$  to  $73^{\circ}$  W. The wholesale destruction of the tilefish in 1882 is thought by Colonel McDonald to be due to climatic causes." (Goode & Bean.) (*chamaeleon*,  $\chi\alpha\mu\alpha\iota\lambda\acute{\epsilon}\omega\nu$ ; - *ceps*, head.)

*Lopholatilus chamaeleonticeps*, GOODE & BEAN, Proc. U. S. Nat. Mus. 1879, 205, Nantucket Shoals; JORDAN & GILBERT, Synopsis, 624, 1883; COLLINS, Rept. U. S. Fish Comm. (1882) 1884, 237; LUCAS, Rept. U. S. Nat. Mus. (Smithsonian Report) 1889, 647, with plate; GOODE & BEAN, Oceanic Ichthyology, 284, 1896.

#### Family CXCI. OPISTHOGNATHIDÆ.

##### (THE JAW-FISHES.)

Body oblong or elongate, low, moderately compressed, covered with small cycloid scales; lateral line present, straight, running close to the dorsal fin, not extending much behind middle of body. Head large, naked, the anterior profile decurved, no ridges, spines, or crests above. Mouth

terminal, horizontal, its cleft usually very wide, the maxillary sometimes greatly dilated; supplemental maxillary present; premaxillaries protractile; jaws subequal, with conical or cardiform teeth; vomer usually with a few teeth; palatines toothless; opercles unarmed; no suborbital stay. Pseudobranchiae present. Gill rakers rather long; gills 4, a slit behind the fourth; gill membranes somewhat united, free from the isthmus. Branchiostegals 6. Air bladder present. No pyloric caeca. Vertebrae large, about 27 in number. Dorsal fin long, continuous, its anterior half composed of slender, flexible spines, which pass gradually into soft rays; caudal distinct, rounded or lanceolate; tail not isocercal, the last vertebra expanded (27 to 34); anal long and low, without distinct spines; ventrals separate, jugular, I, 5, the middle rays longest; pectorals fan-shaped. Three genera, of about 15 species; small fishes inhabiting rocky bottoms in tropical seas, many of them with bright markings. The American species are all rarities, living about rocks in deep or shallow water; nowhere abundant and none of the species well represented in collections. (*Trachinida*, genus *Opisthognathus*, Günther, Cat., II, 254-256.)

a. Maxillary of great length, nearly as long as head, produced behind in a flexible lamina. OPISTHOGNATHUS, 841.

aa. Maxillary normal, truncate behind, much shorter than head.

b. Caudal moderate, rounded behind; body oblong, moderately compressed.

GNATHYPOPS, 242.

bb. Caudal lanceolate, long and pointed; body elongate. LONCHOPISTHUS, 843.

#### 841. OPISTHOGNATHUS, Cuvier.

*Opisthognathus*, CUVIER, Règne Anim., Ed. 2, vol. II, 240, 1829 (*sonnerati*).

Maxillary prolonged backward in a long flexible lamina, which reaches about to base of pectoral. Characters of the genus otherwise included above. It has been suggested that the species of *Gnathypops* are females of analogous species of *Opisthognathus*, the long maxillary being a character of the male. This seems impossible, but deserves an investigation. The fact that *Gnathypops marillosa* has but 27 vertebrae, while its long-jawed cognate, *Opisthognathus macrognathum*, is said by Poey to have 34 vertebrae, is opposed to this view, as is also the fact that the analogous species do not in other respects exactly correspond, as in *Gnathypops mystacina*, the scales are smaller than in *Opisthognathus lonchurum*; *Gnathypops rhomalca* has fewer fin rays than *Opisthognathus punctatum*, etc. But the parallelism of species in the two genera living in the same waters is remarkable. (ὀπίσθε, behind; γνάθος, jaw.)

a. Scales moderate, about 67; D. X, 15; A. II, 13; body nearly plain olivaceous, the maxillary not distinctly striped within. LONCHURUM, 2615.

aa. Scales very small, 100 to 150 in longitudinal series; dorsal rays about XI, 17; A. II, 16; body and fins much variegated, the maxillary within with 2 ink-black stripes on a milk-white ground.

b. Dorsal without large black spot in front; scales 120. PUNCTATUM, 2616.

bb. Dorsal with a large black spot more or less ocellated.

c. Scales about 100.

MACROGNATHUM, 2617.

cc. Scales about 140.

OMMATUM, 2618.

Head  
very sh  
rather  
Longes  
caudal  
1½ in h  
ceous; i  
dusky  
water.  
at Pens  
found i  
(Λόγχο),  
*Opisthogn*  
Bank  
& G-

D. 28;  
in latera  
teeth.  
everywh  
irregular  
ventral f  
finely spe  
ocelli, th  
upper pa  
edge, lik  
broad, bl  
largest; l  
inky bla  
descriptio  
considera  
same wat  
(*punctatus*  
*Opisthognat*  
DAN, Pr

Head 3;  
moderate  
snout very  
reaching  
length con  
set, formin  
jaw; later  
rakers lon  
3030-

2615. OPISTHOGNATHUS LONCHURUM, Jordan & Gilbert.

Head  $3\frac{1}{2}$ ; depth  $4\frac{1}{2}$ . D. 25; A. 15; scales 67. Head moderate; snout very short, shorter than pupil; eye  $3\frac{1}{2}$  in head; maxillary  $1\frac{1}{2}$  in head, rather narrow; lower jaw included; vomer with 5 rather large teeth. Longest dorsal spine about as long as head, slightly higher than soft rays; caudal long, the middle rays scarcely shorter than head; longest anal rays  $1\frac{1}{2}$  in head; pectoral little more than  $\frac{1}{2}$  head. Scales moderate. Olivaceous; margin of upper lip with a narrow black stripe; caudal with 3 dusky bars; color of rest of body uniform. Gulf of Mexico, in deep water. Two specimens known, taken from the stomach of a Red Snapper at Pensacola, Florida. The species resembles *Gnathypops mystacinus*, found in the same waters, but the latter species has smaller scales. (*λόγχη*, lance; *ὄψρα*, tail.)

*Opisthognathus lonchurus*, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1882, 290, Snapper Banks, off Pensacola, Florida (Type, No. 29671. Coll. Jordan & Stearns); JORDAN & GILBERT, Synopsis, 943, 1883.

2616. OPISTHOGNATHUS PUNCTATUM, Peters.

D. 28; A. 18. Body moderately elongate; scales very small, about 125 in lateral line. Dorsal spines continuous with the soft rays. No vomerine teeth. Maxillary very long, extending slightly beyond head. Head everywhere finely speckled with black, the body more coarsely and irregularly spotted; pectoral finely and closely speckled, its edge plain; ventral fin dusky, similarly marked; dorsal without large black blotch, finely spotted, the spots behind gradually forming the boundaries of white ocelli, the base of the fins having rings of white around black spots, the upper part with dark rings around pale spots; caudal with pale spots, its edge, like that of the dorsal, somewhat dusky, not black; anal with a broad, blackish edge, and with dark spots, those near the base of the fin largest; lining membrane of maxillary with the usual bands of white and inky black. Mazatlan. Only the type of this species is yet known, its description having been taken by us from the original specimen. It bears considerable resemblance to *Gnathypops rhomatea*, which is found in the same waters, differing in the generic character of the dilated maxillary. (*punctatus*, spotted.)

*Opisthognathus punctatus*, PETERS, Berliner Monatsberichte 1869, 708, Mazatlan; JORDAN, Proc. Ac. Nat. Sci. Phila. 1883, 290; JORDAN, Cat. Fish. N. A., 118, 1885.

2617. OPISTHOGNATHUS MACROGNATHUM, Poey.

Head  $3\frac{1}{2}$ ; depth 5. D. XI, 16; A. II, 16 or 17; P. 17; scales 100. Body moderately elongate, somewhat compressed. Head blunt anteriorly; snout very short, about as long as pupil; eye large, 4 in head; maxillary reaching slightly past edge of preopercle, but not to end of head, its length contained  $3\frac{1}{2}$  times in length of body. Teeth rather strong, wide set, forming 2 distinct series, directed backward, especially in the upper jaw; lateral teeth of lower jaw largest; a single vomerine tooth. Gill rakers long and slender, nearly 20 below angle. Scales very small.

Dorsal fin low, continuous, the soft rays but little higher than the spines, which are slender and flexible, the longest  $3\frac{1}{2}$  in head; caudal short, rounded, its length  $5\frac{1}{2}$  in body; anal similar to soft dorsal; pectoral  $\frac{1}{2}$  as long as head. Grayish olive, much variegated with yellowish and dark olive; about 6 irregular dusky bands on the body, which extend on the dorsal fin; whitish markings on body forming roundish spots, surrounded by reticulations of grayish olive; head marbled, its posterior part, as well as the sides of the back and pectoral base, with small blackish dots; membrane lining inside of maxillary with 2 curved inky-black bands on a white ground; angle of mouth with a black spot; lining of opercle black; fins all variegated like the body. Florida Keys to Cuba. Here described from the type of *O. scaphiurum*, from Garden Key, but *O. macrognathum* seems to be the same. (*μακρὸς*, long; *γναθός*, jaw.)

*Opisthognathus macrognathus*,\* POEY, Memorias, II, 284, July, 1860, Cuba. (Coll. Poey.)

*Opisthognathus megastoma*, GÜNTHER, Cat., II, 255, September, 1860, Gulf of Mexico. (Haslar Collection.)

*Opisthognathus scaphiurus*, GOODE & BEAN, Proc. U. S. Nat. Mus. 1882, 417, Garden Key, Florida (Type, No. 5938, U. S. Nat. Mus. Coll. Dr. Whitehurst); JORDAN & GILBERT, Synopsis, 943, 1883.

#### 2618. OPISTHOGNATHUS OMMATUM, Jenkins & Evermann.

Head 3; width of head 5; depth 5; eye 3 in head. D. 28; A. 18; scales about 140. Body moderate, compressed, depth  $4\frac{1}{2}$ ; width behind the head  $8\frac{3}{5}$  in length of body. Head large, its breadth equaling its depth, being 5 in length of body. Scales small, embedded; head naked, lateral line extending past middle of dorsal fin. Mouth large. Maxillary long,  $1\frac{1}{2}$  in head; postorbital portion  $2\frac{3}{5}$  in head, not extending beyond head; snout short, its length less than  $\frac{1}{2}$  diameter of eye; distance from tip of snout to end of maxillary  $3\frac{3}{5}$  in length of body and  $1\frac{1}{2}$  in head. Teeth in front part of each jaw in several series, on sides of jaws reduced to a single series, the outer series strong; a tooth on the vomer; gill membranes connected; the interorbital space very narrow,  $11\frac{1}{2}$  in head. Distance from snout to origin of dorsal but little greater than length of head; space between dorsal and caudal fins  $\frac{1}{2}$  greater than length of snout; no depression between spinous and soft rays of dorsal fin, the dorsal equaling the anal in height, its longest ray  $1\frac{3}{5}$  times the eye; pectorals slightly longer than ventrals, being 2 in head; breadth of pectorals 3 in head; ventrals inserted slightly in front of pectorals; caudal rounded and narrow. Coloration: Body irregularly mottled with dark, head evenly blackish; dorsal fin blackish on the posterior portion, with 2 rows of 4 or 5 pale spots well separated; a large ocellated spot from the third to the sixth spines, including them, greater than diameter of eye; anal fin black, with a series of pale spots on the rays, the base pale; caudal black, with 2 pale

\* Poey thus describes his specimens of *Opisthognathus macrognathum*: "Head  $3\frac{1}{2}$  in total; depth  $5\frac{1}{2}$ ; eye nearly 4, twice length of snout, 3 times interorbital width. D. XI, 16; A. II, 16. Vomer with 2 teeth; spines not pungent. Body covered with large yellowish points on a brown ground; 7 broad brown bands on sides, not reaching belly, but extending to middle of dorsal, which, like the anal, has yellow points; a large black ocellus between sixth and ninth spines of dorsal; maxillary with 2 ink-black bands on a milk-white ground; pectorals, ventrals, and caudal yellowish with black points. No pyloric caeca; vertebrae  $10 + 24 = 34$ , the first 5 strong." (Poey.)

spots at the base and a row of spots across the middle; lining of maxillary with bands of black and white. Bay of Guaymas; 3 specimens known. (*ὀμμάτος*, eyed, from its ocellate dorsal.)

*Opisthognathus ommata*, JENKINS & EVERMANN, Proc. U. S. Nat. Mus. 1888, 153, Guaymas. (Type, No. 39640. Coll. Jenkins & Evermann.)

#### 842. GNATHYPOPS, Gill.

*Gnathypops*, GILL, Proc. Ac. Nat. Sci. Phila. 1862, 241 (*maxillosus*).

This genus differs from *Opisthognathus* in having the maxillary of medium length and truncate behind, not extending to edge of opercle; caudal moderate, rounded behind. Species in form and habit agreeing closely with those of *Opisthognathus*. (*γνάθος*, jaw; *ὑπο*, below; *ὤψ*, eye.)

- a. Body and fins spotted with black and often with pale.
- b. Dorsal fin with a conspicuous dusky blotch in front.
- c. Scales very small, about 120; dorsal rays 26. SCOPS, 2619.
- cc. Scales moderate, about 65; dorsal rays 15. MAXILLOSA, 2620.
- bb. Dorsal fin without distinct blotch in front; scales small, about 100.
- d. Dorsal rays about XI, 16; dark spots on head and body few. MACROPS, 2621.
- dd. Dorsal rays about XI, 13; dark spots on head and body numerous.
- e. Dorsal fin distinctly notched; lateral line not reaching middle of dorsal. RHOMALEA, 2622.
- ce. Dorsal fin not notched; lateral line reaching middle of dorsal. SNYDERI, 2623.
- aa. Body and fins nearly uniform olive; the spots few and sparse; dorsal rays 24; scales 100; no black on membrane of maxillary. MYSTACINA, 2624.

#### 2619. GNATHYPOPS SCOPS, Jenkins & Evermann.

Head  $3\frac{1}{2}$ ; width of head  $5\frac{1}{2}$ , its depth  $4\frac{1}{2}$ . D. 26 (X, 16); A. 19 (II, 17); scales 3-122-40. Scales small, none on head; lateral line extending to about middle of dorsal fin; mouth large; maxillary extending beyond eye a distance  $4\frac{1}{6}$  in head; snout  $6\frac{1}{2}$  in head; teeth in bands, outer series on upper jaw rather strong; a single tooth on vomer; gill membranes connected. Opercle ending in a long flap, which extends upward and backward, nearly meeting over the back in front of the dorsal fin. Eye large,  $2\frac{1}{2}$  in head; interorbital space narrow, 11 in head. No depression between the dorsal spines and the soft rays, which are scarcely distinguishable; height of dorsal equal to that of anal; ventrals inserted in front of pectorals; pectorals equal to ventrals in length, 7 in body; caudal rounded. Coloration, in alcohol: Body pale, covered with many dark spots about the size of 3 to 6 scales; top of head with smaller dark spots; sides of head with whitish spots; dorsal fin with a black ocellated spot equal to eye on the space between second and fifth spines; remainder of fin dark, with many white spots running into each other on some portions, so as to form irregular lines; base of anal pale, the outer edge black; caudal dark with 2 whitish spots at the base, and a row of 6 white spots across the middle on alternate rays; pectorals lighter, with small whitish specks; ventrals dusky; belly pale. Guaymas; 3 specimens known, respectively 115 cm., 10 cm., and 7 cm. in length to base of caudal. This species is the

analogue of *Opisthognathus ommatum*, also from Guaymas. (*scops*, the screech owl; *ὄκωψ*, from *ὄκοπέω*, to look, in allusion to the large eyes.)

*Gnathypops scops*, JENKINS & EVERMANN, Proc. U. S. Nat. Mus. 1888, 152, Guaymas. (Type, No. 39641, U. S. Nat. Mus. Coll. Jenkins & Evermann.)

2620. GNATHYPOPS MAXILLOSA (Poey).

Head  $3\frac{1}{2}$ ; depth  $4\frac{1}{2}$ . D. VIII, 17; A. II, 13; scales 65. Body moderately compressed; head not very large; maxillary truncate behind, extending behind eye for a distance for about  $\frac{2}{3}$  diameter of eye, its length  $1\frac{1}{2}$  in head; eye  $3\frac{1}{2}$  in head. Teeth conical, curved, well separated, mostly in a single series; no teeth on vomer. Fins moderate; dorsal continuous, its spines slender; caudal short, its length  $\frac{2}{3}$  head. Color grayish olive, with 7 irregular A-shaped bars of darker, everywhere much marbled and variegated; fins all similarly marked, the ventrals dusky, the dorsal with a dusky blotch in front. Cuba, north to Florida, from which locality the specimen here described was taken. Evidently very close to the Brazilian species *G. curieri*, the eye perhaps smaller. According to Poey, *Gnathypops maxillosa* has the eye 4 in head; D. VIII, 18; A. II, 15; 2 teeth on vomer; spinous dorsal lower than soft dorsal; body covered with large yellow spots on a ground color of clear brown; 6 brown cross bands reaching middle of dorsal, which is variegated with yellow and reddish, as is the anal; maxillary yellowish on its posterior border, the middle blackish; other spots on the jaws; ventrals, pectoral, and caudal yellowish, the ventrals finely spotted with brown, the caudal with 5 brown bands; base of pectorals with dark spots. Vertebrae  $10 + 17 = 27$ . (*maxillosus*, pertaining to the jaw.)

*Opisthognathus maxillosus*, POEY, Memorias, II, 286, 1860, Cuba. (Coll. Poey.)

*Gnathypops maxillosus*, GILL, Proc. Ac. Nat. Sci. Phila. 1862, 241; POEY, Synopsis, 400; JORDAN & GILBERT, Synopsis, 942, 1883.

2621. GNATHYPOPS MACROPS (Poey).

Head  $3\frac{1}{2}$  in total (with caudal?); depth about  $4\frac{1}{2}$ . D. XI, 16; A. II, 15 or 16; scales 100. Eye  $3\frac{1}{2}$  in head; maxillary extending beyond eye  $\frac{1}{2}$  of a diameter,  $5\frac{1}{2}$  in head; vomer with 6 conical teeth. Color (faded in the type) reddish olive, with round, yellowish spots and vestiges of vertical bands; dorsal and anal plain, pectorals with brown bands; jaws not spotted with brown and white. Coast of Cuba (Poey); known from 1 specimen 132 mm. long; not seen by us. According to Poey, it may not be distinct from *G. maxillosa*, which in turn may possibly be the female of *Opisthognathus macrognathum*. This species may also be identical with the Brazilian species *Gnathypops curieri*,\* but the latter has a dorsal ocellus and apparently larger scales. (*μακρός*, large; *ὄψ*, eye.)

*Opisthognathus macrops*, POEY, Memorias, II, 287, 1860, Cuba. (Coll. Poey.)

\* *Gnathypops curieri* (Valenciennes). Head  $3\frac{1}{2}$ ; depth  $4\frac{1}{2}$ ; eye  $3\frac{1}{2}$  in head. D. X, 18; A. II, 16; scales 70. Maxillary reaching beyond the vertical from posterior margin of orbit,  $1\frac{1}{2}$  in head; eye  $3\frac{1}{2}$  in head, dorsal fin not notched. Olivaceous; a large dark-blue ovate ocellate spot between the fourth and eighth dorsal spines; dorsal and anal mottled, the edge dusky posteriorly; caudal with 3 dark-bluish bands. Bahia (Valenciennes). (Named for Georges Dagobert Cuvier.)

*Opisthognathus curieri*, CUVIER & VALENCIENNES, Hist. Nat. Poiss., XI, 504, 1836, Bahia (Coll. Blanchet); GÜNTHER, Cat., II, 256, 1860.

Head  
rather  
than b  
somewh  
greatest  
not very  
of the fi  
well bey  
mandibul  
small, b  
Teeth n  
on the  
upper ja  
branches s  
of eye, 9  
Head na  
naked; l  
scales in  
rather d  
spines 3  
scarcely  
bony op  
last rays  
fin round  
est rays  
pectoral,  
spirits, o  
where m  
thickly a  
above an  
on lips a  
eye thick  
upper part  
and disap  
first 3 or  
clouds, an  
and anal  
greenish  
water; 1  
This is pe  
*Opisthognathus*  
Maria  
Head 3  
on vomer  
sal; no n

## 2022. GNATHYPOPS RHOMALEA (Jordan &amp; Gilbert).

Head  $2\frac{1}{2}$ ; depth 4. D. XI, 13; A. II, 13; scales 103 (pores fewer). Body rather robust, compressed; head very large, ovoid, thicker and deeper than body, with swollen cheeks, the occipital region high, the snout somewhat truncate, the intermediate profile forming a nearly even curve; greatest depth of head equal to its thickness and  $\frac{2}{3}$  its length. Eye not very large, 6 in head, longer than snout, about equal to the width of the flattish interorbital space. Mouth large, the maxillary extending well beyond the eye, but not to the margin of the preopercle nor to the mandibular joint, its posterior margin truncate; supplemental bone small, but distinct; length of maxillary from end of snout  $1\frac{1}{2}$  in head. Teeth moderate, in both jaws, in broad bands which become narrow on the sides; outer series of teeth somewhat enlarged, especially in upper jaw; 1 rather small, blunt tooth on middle of vomer. Gill membranes scarcely connected; gill rakers long and slender, about  $\frac{1}{2}$  diameter of eye, 9 + 19; pseudobranchiae situated in a cavity above the gill arches. Head naked; scales on body small, smooth, somewhat embedded; breast naked; lateral line ceasing opposite anterior third of second dorsal; 103 scales in a longitudinal series from head to caudal. Dorsal fin high; a rather deep notch separating the spines from the soft rays; the longest spines 3 in length of head, more than  $\frac{1}{2}$  longer than the last spine, and scarcely lower than the soft rays. Insertion of dorsal opposite tip of the bony opercle, the opercular flap extending to opposite the third spine, last rays of dorsal and anal reaching past the base of caudal rays; caudal fin rounded, about  $\frac{1}{2}$  length of head; anal higher than soft dorsal, its longest rays  $2\frac{1}{2}$  in head; ventrals large, close together, inserted in front of pectoral,  $1\frac{1}{2}$  in head; pectorals short and broad,  $1\frac{1}{2}$  in head. Color in spirits, olivaceous, slightly brownish above, scarcely paler below, everywhere more or less tinged and mottled with greenish; head everywhere thickly and closely covered with small rounded dark-brown spots, largest above and on cheeks, where they are about as large as pin heads; smaller on lips and opercles, most thickly set on the anterior part of the head; eye thickly spotted; spots similar to those on the head extending along upper part of back, forming a vague band, which grows narrower backward and disappears opposite front of second dorsal; front side of pectoral and first 3 or 4 dorsal spines with dark spots; dorsal dusky olive, with darker clouds, and with some dark spots, especially on the spinous part; caudal and anal plain dusky or faintly marbled with paler; ventrals blackish, greenish at base; pectorals dusky green. Gulf of California, in shallow water; 1 specimen known, the type (above described) 16 inches long. This is perhaps the largest species of the genus. (*ρομαλέος*, robust.)

*Opisthognathus rhomaleus*, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1881, 276, Santa Maria Cove, Lower California. (Type, No. 29382. Coll. Lieut. Henry L. Nichols.)

## 2023. GNATHYPOPS SNYDERI, Jordan &amp; Evermann, new species.

Head  $3\frac{1}{2}$ ; depth  $4\frac{1}{2}$ . D. X, 14; A. II, 13; scales 93; 60 pores; 3 or 4 teeth on vomer; lateral line very distinct, extending to fourteenth ray of dorsal; no notch separating the dorsals, the spines and soft rays not sepa-

table, the last spine not much shorter than the longest, which is  $2\frac{1}{2}$  head; longest anal ray 2, pectoral  $2\frac{1}{2}$ . Vertebrae  $10 + 17 = 27$ . Body olivaceous, with 5 broad faint dusky cross shades; head with many round black spots of varying sizes, some as large as pin heads, the largest below and between eyes, covering both jaws and the membrane of the maxillary, few on cheeks, most numerous on forehead; similar spots extending along side of back to end of lateral line; dorsal dusky, with 6 round dusky blotches at its base, corresponding to the dark shades on body; caudal, anal, and pectorals plain dusky olive; ventrals blackish. Gulf of California; known from 1 specimen collected by Dr. Gilbert in San Luis Gonzales Bay. The species is close to *G. rhomalea*, but has a different dorsal fin and lateral line. Type, No. 2014 L. S. Jr. Univ. Mus., about 8 inches long. (Named for John O. Snyder, curator of fishes in Leland Stanford Junior University.)

2624. GNATHYPOPS MYSTACINA, Jordan.

Head  $3\frac{1}{2}$  in length ( $3\frac{5}{8}$  to tip of caudal); depth  $4\frac{1}{4}$  ( $5\frac{1}{8}$ ). D. 23 or 24 (X, 14); A. II, 11; lateral line with about 54 tubes; 100 scales between gill opening and caudal. Head rather elongate, very blunt in profile; snout very short, not longer than pupil; eye large, about  $3\frac{1}{4}$  in length; maxillary  $1\frac{3}{8}$  in length of head, 5 in length to base of caudal,  $6\frac{1}{4}$  in total length to tip of caudal; end of maxillary abruptly truncate, not ending in a flexible lamina, the supplemental bone well developed; lower jaw slightly included. Teeth in each jaw in a narrow band, the outer slender, enlarged; vomer with about 4 slender teeth; palatines toothless. Gill rakers long and slender. Gill membranes nearly separate, free from the isthmus. Scales very small; lateral line extending to below anterior part of soft dorsal, its length  $\frac{1}{2}$  that of head. Dorsal spines not distinguishable from the soft rays, the rays apparently fewer than usual, none of them very high, the last ray  $2\frac{1}{4}$  in head; caudal short, apparently truncate,  $1\frac{1}{2}$  in head; anal rather low; pectorals 2 in head; ventrals  $1\frac{1}{4}$ . Color nearly plain olive green, without bands or spots on body or fins; vertical fins tipped with blackish; maxillary with a faint median blackish stripe; pectoral with 2 dusky cross shades; no black or white on lining membrane of jaws. Length  $3\frac{1}{2}$  inches. Deep waters of Gulf of Mexico; the few specimens known from the stomachs of Red Snapper (*Neomanis aya*) from the Pensacola Snapper Banks. It resembles *Opisthognathus lonchurum*, but the scales are smaller. ( $\mu\acute{\upsilon}\sigma\tau\alpha\zeta$ , mustache, from the maxillary stripe).

*Gnathypops mystacinus*, JORDAN, Proc. U. S. Nat. Mus. 1884, 37, Snapper Banks off Pensacola. (Coll. Jordan & Stearns. Type, 34976, U. S. Nat. Mus.)

843. LONCHOPISTHUS, Gill.

*Lonchopisthus*, GILL, Proc. Ac. Nat. Sci. Phila. 1862, 241 (*micrognathus*).

This genus differs from *Gnathypops* in the slender, compressed body, the still smaller maxillary and the lanceolate caudal fin. The single species is very rare. ( $\lambda\acute{o}\gamma\chi\eta$ , lance;  $\delta\pi\iota\theta\epsilon$ , behind, from the form of the caudal.)

Head  
Body e  
diamete  
jaws; n  
profile  
brown,  
the first  
the pect  
Length  
us in th  
*Opisthogn*  
*Lonchopis*

Body  
oid scale  
nearly h  
laries pr  
out supp  
moderate  
bands of  
on head.  
nected,  
branchia  
top and  
line cons  
sometimes  
few of th  
the poste  
convex;  
ward and  
pectorals  
fied. Py  
numbers,  
Pacific; I  
tain; ext  
not be cl

a. Gill m  
b. H

bb. I

aa. Gill  
n



2625. LONCHOPISTHUS MICROGNATHUS (Poey).

Head 5 in total; depth 6. D. X, 17; A. II, 16; scales 80; eye 3 in head. Body elongate, compressed, snout short; maxillary reaching  $\frac{1}{2}$  an eye's diameter behind the eye; no teeth on vomer; no second row of teeth in jaws; no scales on head except on cheek; lateral line almost touching profile of back. Caudal long and pointed as in *Gobius oceanicus*. Dark brown, paler below; 20 narrow vertical whitish bands from back to belly, the first 2 on cheek, the third on opercle; fins colored like body, except the pectorals, which are yellow, the edge orange. Vertebrae 10 + 18 = 28. Length 4 inches. Cuba; rare (Poey); only the types known, examined by us in the National Museum. (*μικρός*, small; *γνάθος*, jaw.)

*Opisthognathus micrognathus*, POEY, *Memorias*, II, 287, 1860, Cuba. (Coll. Poey.)

*Lonchopisthus micrognathus*, GILL, *Proc. Ac. Nat. Sci. Phila.* 1862, 241.

Family CXCII. BATHYMASTERIDÆ.

(THE RONQUILS.)

Body rather elongate, moderately compressed, covered with small, ctenoid scales. Head rather large, subconic. Eyes large. Mouth moderate, nearly horizontal, the lower jaw slightly projecting; lips full; premaxillaries protractile, not extending to angle of the mouth; maxillary without supplemental bone, not slipping under the narrow preorbital. Teeth moderate, in a cardiform band in each jaw, the outer somewhat enlarged; bands of teeth on vomer and palatines. No barbels; no crests or spines on head. Branchiostegals 6. Gill membranes scarcely or broadly connected, free from the isthmus; gill rakers few, very short. Pseudo-branchiae large. Opercular bones unarmed. Mucous pores numerous on top and sides of head, sometimes provided with fringed flaps. Lateral line conspicuous, placed high, not quite reaching the caudal fin, its scales sometimes enlarged. Dorsal fin long, continuous, moderately high, a few of the foremost rays inarticulate, none of them pungent or spine-like; the posterior rays branched; anal fin long, similar to the dorsal; caudal convex; pectorals rather broad, their bases extending obliquely downward and backward, their rays all branched; ventrals slightly in front of pectorals, 1, 5, close together, the inner rays longest. Skeleton well ossified. Pyloric caeca few (2 or 3). No anal papilla. Vertebrae in large numbers, about 14 + 35. Three species known, from the Northern Pacific; here referred to 3 genera. The relations of the group are uncertain; externally they resemble the *Opisthognathida*, but the relation can not be close, and the number of vertebrae is greatly increased.

a. Gill membranes not connected below.

b. Head naked; scales in lateral line not enlarged; only first 3 or 4 rays in dorsal fin unbranched; pores of head with small flaps. BATHYMASTER, 844.

bb. Head scaly on cheeks; scales in lateral line enlarged; anterior 20 to 30 rays of dorsal fin unbranched; pores of head mostly without flaps.

RONQUILUS, 845.

aa. Gill membranes broadly connected; cheeks scaly; scales in lateral line enlarged; about 15 of anterior rays of dorsal simple. RATHSUNELLA, 846.

## 844. BATHYMASTER, Cope.

*Bathymaster*, COPE, Proc. Amer. Phil. Soc. 1873, 31 (*signatus*).

Head naked; pores of head large, many of them with dermal flaps; gill membranes scarcely connected; scales of lateral line similar to the others; dorsal fin with but 3 or 4 of its anterior rays unbranched; characters otherwise included above. (*βαθύς*, deep; *μαστήρ*, searcher.)

## 2626. BATHYMASTER SIGNATUS, Cope.

Head  $3\frac{1}{2}$ ; depth 5. D. 47; A. 34; scales 6-95-19; eye  $4\frac{1}{2}$  in head; maxillary  $2\frac{1}{2}$ ; snout  $4\frac{1}{2}$ ; pectoral  $1\frac{1}{2}$ ; ventral  $2\frac{1}{2}$ ; highest dorsal ray  $2\frac{1}{2}$ ; highest anal ray  $3\frac{1}{2}$ ; caudal  $2\frac{1}{2}$ . Body compressed, elongate, anterior profile convex from tip of snout to dorsal; mouth not very oblique, the maxillary reaching the vertical from posterior edge of orbit; snout about equal to eye; jaws equal, with bands of small conical teeth, outer row enlarged; lower jaw with a single row at the sides; well developed conical teeth on vomer and palatines. Branchiostegal membranes not united; margin of preopercle free, furnished with 5 or 6 conspicuous mucous pores; large pores on top and sides of head, each with a small flap; opercle ending in a flap behind; gill rakers moderately long and slender,  $\frac{2}{3}$  eye, about 7 + 18; many mucous pores on top of head and under eye; head entirely naked; dorsal and pectoral with fine scales running about halfway up the fin; anal naked; a naked strip from nape to dorsal; pectoral broad and fan-shaped, its lower rays smaller, reaching to front of anal; origin of ventral spine about the diameter in front of the lower end of pectoral base; dorsal about uniform in height for nearly its entire length, higher than anal; dorsal and anal rays about reaching to base of caudal rays; first 3 or 4 rays of dorsal simple, the others branched; caudal truncate or slightly rounded. Color almost uniform warm brown with darker shades, the fins somewhat mottled with yellowish, the anal and ventrals blackish, other fins dusky; a conspicuous black ocellated blotch on front of dorsal, covering tips of 4 or 5 spines.\* Shores of southern Alaska, from Unimak

\* Concerning this species, Dr. Gilbert has the following note: "*Bathymaster signatus* is taken very abundantly in our series of shallow-water dredgings along the southern shore of the Alaskan Peninsula, and northward through Unimak Pass. The stations at which it was obtained are numbered 3211, 3212, 3213, 3214, 3215, 3217, 3220, 3222, and 3223, and the depth range from 34 to 58 fathoms. In addition, a very few small specimens were secured at Stations 3202, 3309, 3221 and 3353, north of the Aleutian Islands, in depths of 19 to 71 fathoms, but the species is evidently not abundant in Bering Sea. No examples were taken in any of the very numerous dredgings made in Bristol Bay. In life the sides are olive brown, and the upper parts show faint traces of 6 or 7 broad dusky cross bars, which correspond to or alternate with an equal number below the lateral line; the anal and ventral fins, the branchiostegal and gular membranes, the lower pectoral rays, and the snout blue black; anterior edge of orbit and front edge of preorbital light yellow; the pores on edge of preopercle, 2 pores above and behind maxillary, and 3 at upper edge of opercle, bright scarlet; a large black blotch on anterior dorsal rays; distal half of anterior portion of dorsal fin and the upper pectoral rays yellow. Outer ventral ray simple and inarticulate, followed by 5 branched rays. Only the first 2 dorsal rays spinous, being soft and flexible, but unjointed. The third and all following rays jointed and forked. All of the anal rays jointed. A specimen from *Albatross Station* 3211, 35 mm. in length to base of caudal, shows that the ventrals occupy very different positions in adults and in young. In the latter they are truly thoracic in position, and are inserted as much behind base of pectorals as they are located in advance of this point in adults. A specimen 65 mm. long is entirely similar to adults in this respect."

Pass to Sitka; not uncommon in water of moderate depths. Here described from a specimen collected by the *Albatross* (No. 2143, L. S. Jr. Univ. Mus.), Station 3214, 11 inches in length. Other specimens taken in rock pools at Sitka are dark green, almost black. (*signatus*, marked.)

*Bathymaster signatus*, COPE, Proc. Amer. Philos. Soc. 1873, 31, Sitka (Coll. Prof. George Davidson); GILBERT, Proc. U. S. Nat. Mus. 1888, 554.

845. RONQUILUS, Jordan & Starks.

*Ronquilus*, JORDAN & STARKS, Proc. Cal. Ac. Sci. 1895, 838 (*Jordani*).

Checks scaly; scales of lateral line enlarged; anterior half of dorsal fin more or less composed of unbranched rays; mucous pores on head without conspicuous flaps; gill membranes separate. One species. (*Ronquil*, a Spanish name of the typical species, possibly from *ρόγχος*, one who grunts.)

2627. RONQUILUS JORDANI (Gilbert).

(RONQUIL.)

Head  $4\frac{1}{2}$ ; depth  $6\frac{1}{2}$ . D. 41; A. 33; V. I, 5; P. 18; scales 92+6 (tubes), about 200 transverse. Body rather elongate, moderately compressed. Eye large, about as long as snout, 4 in head, its diameter much more than the interocular space; maxillary extending to below front of pupil; cheeks closely scaly; rest of head entirely naked; a narrow, naked area in front of dorsal, bounded by rows of mucous pores; skull with large mucous cavities behind the eyes, which are translucent in life; scales of lateral line enlarged, twice as far apart as the others. Dorsal fin inserted at a distance behind the occiput, less than the diameter of the eye; pectorals  $\frac{1}{2}$  the length of the head; fourth ray of ventrals longest; vent much nearer snout than root of caudal; vertebrae 14 + 35 = 49; olivaceous, tinged with brown; about 8 round, faint-bluish blotches along the sides, each surrounded by rings of yellow spots; a yellow ring around the eye and a yellow band along the cheek; fins translucent, the anal with a yellowish strip and a deep-bluish or black edging; dorsal reddish or yellow, with a dusky blotch in front; ventrals dusky; pectorals with the lower rays blackish or dark blue, larger specimens nearly uniformly dark, the color varying with the surroundings. Bristol Bay to Puget Sound, about rocks, in water of moderate depth. Length 6 to 10 inches. Known from Seattle and from Wrangel and Bristol Bay, the latter specimen in 32 fathoms. (Named for its discoverer, David Starr Jordan.)

*Bathymaster signatus*, JORDAN & GILBERT, Synopsis, 623, 1883; not *B. signatus*, COPE.

*Bathymaster jordani*, GILBERT, Proc. U. S. Nat. Mus. 1888, 554, Elliott Bay at Seattle (Coll. Jordan) and Fort Wrangel, Alaska. (Coll. *Albatross*.)

*Ronquilus jordani*, JORDAN & STARKS, Proc. Cal. Ac. Sci. 1895, 838, pl. 99.

846. RATHBUNELLA, Jordan & Evermann.

*Rathbunella*, JORDAN & EVERMANN, Check-List Fishes, 463, 1896 (*hypoplectus*).

This genus differs from *Ronquilus* in having the gill membranes broadly united across the isthmus. The unbranched anterior rays form about  $\frac{1}{2}$

of the dorsal fin. (Named for Mr. Richard Rathbun, then chief of the Division of Scientific Inquiry in the U. S. Fish Commission, in recognition of his many services to science.)

2628. *RATHBUNELLA HYPOPLECTA* (Gilbert).

Head  $4\frac{2}{3}$  in length; depth 7. D. 46; A. 33. Head and body compressed, elongate, the anterior profile of head compressed, declivous; mouth somewhat oblique, at lower side of snout, small, the maxillary reaching vertical from middle of pupil,  $3\frac{1}{8}$  in head; snout very slightly shorter than orbit,  $4\frac{1}{2}$  in head; diameter of orbit 4 in head; teeth well developed, in broad bands on jaws, vomer and palatines, the vomer and palatine patches nearly continuous; branchiostegal membranes broadly united, free from isthmus, forming a fold whose depth exceeds  $\frac{1}{2}$  diameter of orbit. Margin of preopercle adnate behind, slightly free below, furnished with a series of 6 conspicuous mucous pores; head without spines, ridges or filaments; inner margin of shoulder girdle conspicuously notched above and below, but without hook; gill rakers tubercular, few in number; a well-marked slit behind last gill. Distance from nape to front of dorsal fin equals its distance from posterior border of eye; anterior 10 or 12 dorsal rays simple and apparently not articulate, but flexible and not spine-like; distance from front of anal to base of ventrals  $2\frac{1}{2}$  in its distance from base of caudal, all but first  $\frac{1}{2}$  of dorsal rays, and all of anal rays forked at tip; dorsal not high, the longest rays  $\frac{1}{2}$  head; highest anal ray equals snout and  $\frac{1}{2}$  eye; last dorsal and anal rays entirely disconnected from caudal, leaving a free space on caudal peduncle  $\frac{1}{2}$  diameter of orbit; ventrals I, 5, in advance of base of pectorals, narrowly triangular, the inner rays longest; pectorals with curved base running backward and downward, the rays all branched, 18 in number, the width of base of fin  $3\frac{1}{2}$  in head, the longest ray  $1\frac{1}{2}$  in head; caudal rounded,  $\frac{2}{3}$  length of head. Body covered with small, partially embedded, cycloid scales, including antedorsal region, belly, breast, and area in front of base of pectorals; checks covered with similar but smaller scales, the opercles and rest of head naked. Lateral line running high, parallel with back, on a series of enlarged scales, which are also partly embedded in the thick skin; the lateral line fails to reach base of caudal by a distance equaling  $\frac{1}{2}$  of head, and is present on 82 scales. Color, dark olive-brown above, lighter below; a series of about 12 quadrate dark blotches below lateral line, connected more or less by dusky streaks with an alternating series along base of dorsal; no bright colors; dorsal, pectorals, ventrals and branchiostegal membranes dusky straw color; anal black, the rays white tipped; caudal blackish; peritoneum white. A single specimen, 8 inches long, from *Albatross Station* 2944, off Santa Barbara Islands, in 30 fathoms.

*Bathymaster hypoplectus*, GILBERT, Proc. U. S. Nat. Mus. 1890, 97, off Santa Barbara Islands, California, at Albatross Station 2944. (Coll. *Albatross*.)

Body  
skin na  
rather sh  
normal, t  
narrow;  
ous long  
palatines  
Opercula  
caudal fi  
sharp tee  
dar & G

a. Jaws  
er  
aa. Jaws  
p

*Chiasmodes*  
*Chiasmodes*

Body el  
ent, its w  
large; lo  
pointed t  
ine teeth  
No pseud  
the isthm  
inserted  
caudal.  
deep sea,  
their own  
body. (Ch  
anterior c

Head 3  
crown fla  
preopercle  
ail of wh  
ing each

Family CXCIII. CHIASMODONTIDÆ.

(THE BLACK SWALLOWERS.)

Body elongate, subcylindrical, or slightly tapering; head subconic. Skin naked; lateral line continuous, placed low; 2 dorsal fins, the first rather short, of slender spines, the second dorsal and anal long; ventrals normal, thoracic, inserted before pectorals, the rays I, 5; pectorals long and narrow; mouth very deeply cleft, reaching beyond the eyes, with numerous long, sharp, movable teeth, the anterior canines movable; teeth on palatines; upper jaw not protractile, the maxillary produced backward. Opercular apparatus very oblique and reduced; no spines or cirri on head; caudal fin forked. Genera 2; species 2; deep-sea fishes, notable for the sharp teeth and for the extensible stomach. (*Chiasmodontidae*, Gill, in Jordan & Gilbert, Synopsis, 964, 1883.)

- a. Jaws with some of the anterior canines extremely long and movable, the 2 anterior crossing each other; lower jaw projecting. CHIASMODON, 847.  
aa. Jaws with slender, close-set teeth, none of them greatly produced; lower jaw not prominent. PSEUDOSCOPELUS, 848.

847. CHIASMODON, Johnson.

(BLACK SWALLOWERS.)

*Chiasmodon*, JOHNSON, Proc. Zool. Soc. London 1863, 408 (*niger*).  
*Chiasmodus*, GÜNTHER, change of spelling.

Body elongate, compressed, and tapering posteriorly, naked; belly pendant, its walls membranaceous, capable of great dilation. Mouth very large; lower jaw longer than upper; each jaw with 2 series of large, pointed teeth, some of the anterior being very large and movable; vomerine teeth none; palatines with teeth similar to those in the jaws. Gills 4. No pseudobranchia. Gill openings very wide, the membranes joined to the isthmus for a short distance. Dorsal fins 2; anal single; ventrals inserted below pectorals, each of 5 soft rays. Tail truncate at base of caudal. Caudal forked, free from dorsal and anal. Singular fishes of the deep sea, remarkable for their ability to swallow fishes of many times their own size by means of the great distensibility of the walls of the body. (*χίασμα*, a mark of the form of the letter  $\chi$ ; *ὄδους*, tooth; the 2 anterior canines crossing each other when depressed.)

2629. CHIASMODON NIGER, Johnson.

Head  $3\frac{1}{2}$ . D. XI-28; A. 27; P. 13; V. 5. Head compressed, elongate, the crown flat, its depth less than  $\frac{1}{2}$  its length; maxillary reaching angle of preopercle; both jaws armed with long, pointed, wide-set teeth, nearly all of which are movable; 2 anterior teeth of upper jaw very long, crossing each other when depressed; 3 anterior pairs of teeth in lower jaw

likewise prolonged, the third pair the longest; palatines with a longer, fixed tooth in front. Eye moderate, above the anterior part of maxillary,  $4\frac{1}{2}$  in head, shorter than snout, as wide as interorbital space. Lateral line in a longitudinal groove. First dorsal of slender rays, its base  $2\frac{1}{2}$  in that of second dorsal; anal commencing behind second dorsal, its anterior rays without connection with vertebral column; posterior rays of anal and dorsal very feeble; pectoral as long as head without snout; ventral  $\frac{1}{2}$  as long as pectoral. Color entirely black. Length 12 inches. (Günther.) Deep waters of the Atlantic; a remarkable fish, the walls of the body inordinately extensible; taken at Madeira, in the mid-Atlantic, near the island of Dominica, and off the coast of Massachusetts.\* (*niger*, black.)

*Chiasmodon niger*, JOHNSON, Proc. Zool. Soc. London 1863, 408, Madeira; JORDAN & GILBERT, Synopsis, 964; GOODE & BEAN, Oceanic Ichthyology, 292, 1896.

*Chiasmodus niger*, GÜNTHER, Cat., v, 435, 1864; CARTEB, Proc. Zool. Soc. 1866, 38; GÜNTHER, Challenger Report, Deep Sea Fishes, xxii, 99, 1887.

#### 848. PSEUDOSCOPELUS, Lütken.

*Pseudoscopus*, LÜTKEN, Spolia Atlantica, Scopelini, 64, 1892 (*scriptus*).

Body perciform, scaleless, naked; mouth very large; eyes moderate; the slender maxillary reaching far beyond eye; jaws and palate with slender, close-set teeth; ventral fins short, subthoracic, of 1 spine and 5 rays; first dorsal short, of about 8 slender spines; posterior dorsal long, similar to the anal. Each jaw with a distinct line of pores, a median line of pores before ventrals, a cross line connecting ventrals, a series of pores from the vent passing around anal on each side. Lateral line well developed, running high. Head without spines. Gill openings very broad. Pectorals long; caudal short, forked. One species known, in deep water. (*ψευδής*, false; *Scopelus*.)

#### 2630. PSEUDOSCOPELUS SCRIPTUS, Lütken.

Head  $3\frac{1}{2}$ ; depth  $4\frac{1}{2}$ . D. VIII-22; A. 22; V. I, 5. Body subfusiform, somewhat compressed. Head large, the snout short and pointed,  $4\frac{1}{2}$  in head, the small eye, about 5. Jaws subequal, maxillary  $1\frac{1}{2}$  in head; cheek V-shaped, very oblique; bones of head not serrate. Form of head and mouth much as in *Engraulis* or *Scopelus*. Pectoral nearly as long as head, reaching past front of anal; soft dorsal higher than spinous, the anterior rays of soft dorsal and anal elevated. Pores as above described.

\* The first specimen of this remarkable fish was obtained at Magdalena (Madeira), at a depth of 312 fathoms, in 1850, by Lowe, who, however, omitted to give a description of it. The species was rediscovered at the same locality by Johnson twelve years later. A third specimen was picked up from the surface, near the island of Dominica. A fourth example was obtained by the *Challenger* in mid-Atlantic, at Station 107, in 1,500 fathoms, on August 26, 1873. A fifth was obtained by the U. S. National Museum from Capt. Thomas F. Hodgdon of the Gloucester schooner *Beane W. Somers*. It was found on Le Have Bank, floating on the surface, in June, 1880. (Goode & Bean.)

One spec  
uncertain  
(*scriptus*,  
*Pseudoscop*

Body r  
to the ca  
naked or  
near the  
depressed  
ridges. I  
ulated wi  
and subo  
lateral an  
its border  
processes  
posterior  
their ext  
wide; gill  
stegals 6.  
tion short  
often sim  
than the  
caudal fin  
rays divid  
boid area  
thickened  
oped or o  
elongated  
area in ve  
family is  
hined with  
distinguis  
allied to t  
set by t  
rate, by  
eat, with  
Tropics.

a. Body co  
derm

*Hypsicoete*

Body elc  
much depr

One specimen from Old Bahama Straits. (Lütken.) A singular fish of uncertain relationships, remarkable for the development of mucous pores. (*scriptus*, written.)

*Pseudoscopelus scriptus*, LÜTKEN, Spolia Atlantica, Scopelini, 64, 1892, Old Bahama Straits.

Family CXCIV. CHANICHTHYIDÆ.

Body rather elongated, gradually and regularly declining from the nape to the caudal fin; anteriorly subcylindrical or scarcely compressed. Skin naked or covered with small scales. Lateral line high on the sides and near the dorsal fin. Head moderate or large, with the snout prolonged, depressed, and spatuliform. Crown depressed, not relieved by crests or ridges. Preorbital bones large; suborbital chain very narrow, not articulated with the preopercle. Opercular bones all present, the interopercle and subopercle moderately developed. Mouth terminal, with the cleft lateral and large, extending to the vertical of the eye; upper jaw with its border formed almost entirely by the premaxillaries, whose posterior processes are very short; maxillaries with their articulations entirely posterior to the premaxillaries, slender and gradually enlarged toward their extremities. Teeth on the jaws; palate unarmed. Gill openings wide; gill membranes inferiorly deeply emarginated behind. Branchiostegals 6. Pseudobranchiæ developed. Dorsal fin with its spinous portion short, and usually distinct from the soft, the rays of the latter often simply articulated and not branched; anal fin a little shorter than the dorsal, its rays divided, the membrane notched behind each; caudal fin not forked; pectoral fins well developed, with their inferior rays divided; ventral fins jugular or subjugular, separated by a rhomboid area, each with a spine and 5 rays, the first of which is frequently thickened and entire. Cranium flattened behind, the crests little developed or obsolete. The spatuliform snout is principally formed by the elongated frontal bones. Stomach of moderate size and caecal. Pyloric caeca in very small number. The chief distinctive characteristic of this family is doubtless the spatuliform extensions of the snout. This, combined with the extent of the fins, structure of the head, and general form, distinguish the group from all others. It appears to be most closely allied to the *Harpagiferidæ* and *Notothenidæ*. From the former it is separated by the form of the head, as well as by that of the body. From the latter, by the same features, and also by the naked skin. (Gill.) Genera 1, with about 6 species; inhabiting rather deep waters, mostly in the Tropics. (*Chanichthyoidæ*, Gill, Proc. Ac. Nat. Sci. Phila. 1861, 507.)

a. Body covered with cycloid, deciduous scales; maxillary with a flap; opercle with a dermal flap. HYPsicOMETES, 849.

849. HYPsicOMETES, Goode.

*Hypsicometes*, GOODE, Proc. U. S. Nat. Mus. 1880, 347 (*goboides*).

Body elongate, subcylindrical, tapering posteriorly. Head very large, much depressed, with snout elongate, spatulate; cleft of mouth very wide,

horizontal, with lower jaw much the longer; the posterior margin of the maxillary wide, free, and with a long cutaneous flap. Eyes very large, close together, subvertical. Scales large, cycloid, deciduous; lateral line conspicuous and continuous, descending abruptly behind pectorals, its scales smaller than those of the body adjoining. Teeth acicular, in bands on the jaws, vomer, and palatines, the largest being upon the palatines, the vomer, and upon 2 pads on either side of the symphysis of the maxillaries. A sharp, short, strong scapular spine. Opercle with 3 feeble, sharp spines, each at the end of a strong feeble ridge; a long, skinny opercular flap extending far beyond the bony portion, and covered with scales. Branchiostegals 6. Gill membranes free from the isthmus, except far in front, where they are united to it, the left-hand flap overlapping the right at the point of junction. Pseudobranchiæ present. Gill rakers short. (*ὄψι*, below, i. e., in deep water; *καμῆτης*, dweller.)

## 2631. HYPsicOMETES GObOIDES, Goode.

Head about  $2\frac{1}{2}$ ; depth  $7\frac{1}{2}$ ; orbit  $4\frac{1}{2}$  in head, or  $1\frac{1}{2}$  in snout. D. VI-15 to 17; A. 16 to 18; V. I, 5; P. 26; scales 65. Mouth very wide, horizontal, the maxillary, which is expanded spoon-like posteriorly, reaching considerably beyond vertical from anterior margin of orbit; eye considerably nearer tip of snout than end of flap, and equidistant between tip of snout and tip of uppermost spine of operculum; entire upper surface of head, cheeks, and opercula covered with scales, except upon bony portion of snout; first dorsal fin placed far forward, not far behind vertical from axil of pectoral; interspace between termination of first dorsal and beginning of second equal to diameter of the orbit, this fin composed of 6 spines, the first and second of which are longest, equal to distance from anterior margin of orbit to tip of lower jaw, and triangular in form; origin of second dorsal almost vertical from that of anal, and terminating a little in advance of the latter; second dorsal fin highest in front and low behind; length of caudal peduncle a little less than length of snout; caudal rounded; pectoral very broad at base, rounded, extending beyond vent and nearly to vertical from origin of anal; lower rays branched; ventrals far apart, horizontal, *Trigla*-like, composed of 1 flexible spine and 5 branched rays, their insertion far forward and far in advance of base of pectorals. Color grayish brown; lighter and yellowish below. Known only from a very small specimen, in which many of the important characters were not discernible. This specimen (No. 26007, U. S. Nat. Mus.) was taken by the *Fish Hawk* from Station 871, in  $40^{\circ} 02' 54''$  N. lat.,  $70^{\circ} 23' 40''$  W. lon., at a depth of 115 fathoms, and is much contracted and distorted from immersion in strong alcohol. (Goode.) (*Gobius*; *εἶδος*, resemblance.)

*Hypsicometes goboides*, GOODE, Proc. U. S. Nat. Mus. 1880, 348, lat.  $40^{\circ}$ ,  $02'$ ,  $54''$  N., lon.  $70^{\circ} 23' 40''$  W., in 115 fathoms (Coll. *Fish Hawk*): JORDAN & GILBERT, Synopsis, 808, 1883; GOODE & BEAN, Oceanic Ichthyology, 290, fig. 263, 1896.

Body  
sides ve  
almost v  
fringed;  
plement  
moderate  
vomer; p  
short, sle  
branchio  
large. P  
upward.  
no ! arbe  
spines; n  
the first  
out distin  
than the  
pectorals  
area betw  
soft rays  
together,  
caudal lu  
bra nume  
from the l  
and other  
*Uranoscop*  
a. First d  
aa. First e

*Trichodon*, S  
*Trichodon*, C

Character  
low, of 15 s

Head from  
eye  $4\frac{1}{2}$  in he  
 $1\frac{1}{2}$ ; height  
dorsal outli  
dorsal, then



## Family CXC. TRICHODONTIDÆ.

(THE SAND-FISHES.)

Body rather elongate, compressed, naked. Head short, flat on top, the sides vertical. Eyes large, high up, but not superior. Mouth large, almost vertical; lower jaw projecting, its tip entering the profile; lips fringed; premaxillaries protractile; maxillary very broad, without supplemental bone, not slipping under the very narrow preorbital. Teeth moderate, slender and sharp, but not setiform, in bands on jaws and vomer; palatines toothless; inner teeth of jaws depressible. Gill rakers short, slender; gill membranes narrowly united, free from the isthmus. Branchiostegals 5. Gills 4, a slit behind the fourth. Pseudobranchia large. Preopercle with 5 prominent spines, the 2 upper directed strongly upward, the 2 lower downward, the middle 1 downward and backward; no barbels; opercle small, strongly striate, unarmed; preorbital with spines; no suborbital stay. Lateral line obsolete. Dorsal fins separate, the first the larger, of numerous slender spines; anal fin elongate, without distinct spines, the rays of anterior third of the fin much shorter than the others, the beginning of the fin below middle of spinous dorsal; pectorals with a very broad, curved, procurrent base; a broad lunate area between pectoral and gill opening, nearly covered by the opercle; soft rays of dorsal, anal, and pectoral fins all simple; ventrals I, 5, close together, thoracic, but behind the pectorals, the middle rays longest; caudal lunate, with many accessory rays, on a slender peduncle. Vertebrae numerous, 48 in typical species. Two genera and 2 species known; from the North Pacific; living in sand near the shore. The fringed lips and other characters indicate the relationship of these fishes with the *Uranoscopida*. (*Trachinida*, genus *Trichodon*, Günther, Cat., II, 250.)

- a. First dorsal long and rather low, of 14 or 15 spines.  
aa. First dorsal short and high, of 10 spines.

TRICHODON, 850.  
ARCTOSCOPIUS, 851.

## 850. TRICHODON (Steller) Cuvier.

(SAND-FISHES.)

*Trichodon*, STELLER, in Tilesius, Mem. Acad. St. Petersburg, IV, 1811, 468 (*trichodon*).  
*Trichodon*, CUVIER, Règne Animal, Ed. II, vol. 2, 149, 1829 (*trichodon*).

Characters of the genus included above, the first dorsal long and rather low, of 15 spines. One species. (*θριζ*, hair; *ὀδόν*, tooth.)

## 2632. TRICHODON TRICHODON (Tilesius).

(SAND-FISH.)

Head from tip of upper jaw,  $3\frac{1}{2}$ ; depth  $3\frac{1}{2}$ . D. XIII-I, 18; A. 28; P. 22; eye  $4\frac{1}{2}$  in head, snout  $4\frac{1}{2}$ ; maxillary 2; interorbital 3; pectoral  $1\frac{1}{2}$ ; ventral  $1\frac{1}{2}$ ; height of spinous dorsal  $3\frac{1}{2}$ . Body moderately elongate, compressed; dorsal outline slightly concave and sloping gently upward from snout to dorsal, thence turning at a very slight angle nearly straight to caudal;

ventral outline well rounded from chin to caudal peduncle, the curve much more gradual posteriorly; head and body everywhere covered with thin naked skin. Mouth large, superior, nearly vertical, the lower jaw projecting, its tips entering the profile; lips fringed; maxillary reaching to middle of pupil; teeth in 2 or 3 rows, small, sharp and recurved; teeth on vomer; palatines toothless. Eyes placed high, their diameter equal to length of snout; interorbital wide and flat, a third wider than eye; top of head smooth, sometimes rugose in younger individuals, covered with thin smooth skin; anterior nostril ending in a tube; preopercle with 5 spines, the 1 at angle largest, the 2 upper ones pointing upward and backward, the middle one pointing downward and backward, the 2 lower ones pointing downward and forward; opercle with radiating ridges; gill rakers short and slender, numerous. Origin of spinous dorsal behind base of pectoral, its distance from snout 3 in body, the spines not varying greatly in length, the last one connected by a membrane to the back; soft dorsal well separated from spinous, its rays about equal to spines in length, highest in front; anal long, its origin nearer to the snout than base of caudal by a distance equal to the length of the eye. Pectoral, when spread, broadly rounded behind, its lower rays rapidly decreasing in size below, reaching well past front of anal; ventrals inserted behind base of pectorals a distance equal to  $\frac{2}{3}$  eye, their tips reaching to vent. Lateral line running high. Vertebrae  $17 + 30 = 47$ . Color silvery, light brown above; a dark brown streak following the lateral line, broken up into spots anteriorly; quadrangular, dark brown marks along the back at base of dorsals, chain-like markings in front of dorsal on nape; snout and tip of lower jaw dark; a dark line at lower part of eye; dorsals light, a dark streak along upper part of spinous dorsal; pectorals dusky; ventrals and anal colorless. Length 8 to 10 inches. North Pacific, on sandy shores, from Bering Sea to Monterey; very abundant northward; burying in the sand. Here described from a specimen,  $8\frac{1}{2}$  inches in length, from Herendeen Bay, Alaska (*Albatross* collection). Possibly detailed comparison may show a difference between California specimens and those from Bering Sea.

*Trachinus trichodon*, TILESIIUS,\* Mem. Acad. St. Petersburg, IV, 1811, pl. 15, fig. 8, 473, Kamchatka; PALLAS, Zoographia Rosso-Asiatica, III, 235, 1811.†

\* The specific name *trichodon* should apparently date from Tilesius, 1811. Although Vol. IV, of the Mem. Acad. St. Petersburg bears the date 1813 it was for the year 1811, and it is evident that the plate containing the figure of this species was accessible to Pallas as early as 1811, for, in his "Zoographia," printed in 1811, though not published until 1831, Pallas refers to the plate of Tilesius in very definite terms. The fact that Pallas was, in 1811, thus able to refer definitely to Tilesius's plate of *Trachinus trichodon*, fixes the date of publication of that plate at least as early as 1811. That this plate appeared in the volume of Memoirs for 1811 (though the volume was not published until 1813), fixes 1811 as the date for the name. Though the "Zoographia" of Pallas was not formally published until 1831, it was printed in 1811, and Cuvier & Valenciennes evidently had a copy in 1829, as they refer to it.

† Tilesius confused matters greatly by using, in one and the same article, three different names or combinations of names for this fish. At the beginning of this article (p. 406) in a bald list of the species discussed in the paper, he has "*Drachinus trichodon*." On page 466 he has "*Trachinus gasteropelecus*," accompanied by a full description of the species. In a footnote on page 473, he has "*Trachinus trichodon*" together with a description which he says applies to the young, and finally his pl. 15, fig. 8, is marked "*Trachinus trichodon*."

*Drachinus*  
*Trachinus*  
*Trichodon*  
*Trachin*  
sis, 627,  
*Trichodon* li  
A. 28; I

*Arctoscopus*,

This gen  
dorsal whi  
for *Uranos*

Head  $3\frac{1}{2}$ ;  
and color  
spines slen  
opercle wit  
Pectoral w  
the fin cons  
of the dors  
tion as in *T*  
 $4\frac{1}{2}$  inches.  
Japan, and  
Iturup Islan

*Trichodon* japo  
Fishes N.  
*Arctoscopus* japo

Body obl  
cycloid, imbr  
side of back,  
small, superi  
bony stay c  
fringed; mou  
the entire ed  
openings very  
psendobranch  
divided, sever  
close behind t  
from dorsal an  
ventrals jugu  
Genera 4; spe

3030—

- Drachinus trichodon*, TILESIIUS, Mem. Acad. St. Petersburg, IV, 1811, 406; name only.  
*Trachinus gasteropeleus*, TILESIIUS, l. c., 466, 1811, Kamchatka.  
*Trichodon stelleri*, CUVIER & VALENCIENNES, Hist. Nat. Poiss., III, 154, pl. 57, 1820; based on  
*Trachinus trichodon* PALLAS; GÜNTHER, Cat., II, 251, 1860; JORDAN & GILBERT, Synop-  
 sis, 627, 1883.  
*Trichodon lineatus*, AYERS, Proc. Ac. Nat. Sci. Phila. 1860, 60, San Francisco; D. XV-18;  
 A. 28; P. 23.

851. ARCTOSCOPUS, Jordan & Evermann.

*Arctoscopus*, JORDAN & EVERMANN, Check-List Fishes, 464, 1896 (*japonicus*).

This genus differs from *Trichodon* in the short, high, triangular spinous dorsal which is composed of 10 spines. (*ἄρκτος*, northern; *σκοπός*, gazer; for *Uranoscopus*.)

2633. ARCTOSCOPUS JAPONICUS (Steindachner).

Head 3½; depth 3½. D. X or XI-13; A. 30 or 31; P. 25. Form of body and coloration of *Trichodon trichodon*. First dorsal high, triangular, the spines slender, separated by a long interval from the second dorsal. Preopercle with 5 sharp spines; the 2 spines on the preorbital very small. Pectoral well developed, all its rays simple, the lower a little thickened, the fin considerably longer than the head and reaching past the last spine of the dorsal; anal fin with its rays gradually longer posteriorly. Dentition as in *Trichodon trichodon*, but the mouth rather more oblique. Length 4½ inches. North Pacific; scarce. Recorded from Strietok, in the Sea of Japan, and Sitka, Alaska, by Steindachner, and by Jordan & Gilbert from Iturup Island (Kurils). (*japonicus*, from Japan.)

*Trichodon japonicus*, STEINDACHNER, Ichth. Beitr., x, 4, 1881, Strietok; Sitka; JORDAN, Cat. Fishes N. A., 117, 1885.

*Arctoscopus japonicus*, JORDAN & GILBERT, Rept. Fur Seal Investig., 1898.

Family CXCVI. DACTYLOSCOPIDÆ.

(THE SAND STAR-GAZERS.)

Body oblong, low, compressed posteriorly, covered with moderate, cycloid, imbricated scales; lateral line complete, anteriorly running along side of back, posteriorly median; head oblong, nearly plane above; eyes small, superior, well forward; suborbital bones enlarged, but without bony stay connecting with the preopercle; nostrils double; opercles fringed; mouth nearly vertical; premaxillaries protractile, not forming the entire edge of the upper jaw; lips fringed as in *Uranoscopida*; gill openings very broad, the membranes separated and free from the isthmus, pseudobranchiæ present or obsolete. Dorsal fin very long, continuous or divided, several of the anterior rays spinous; anal very long, commencing close behind the vent, which is near the breast; caudal diphyercal, free from dorsal and anal; pectorals variable, the base broad and procurrent; ventrals jugular, I, 3; vertebrae more than 10 + 14; pyloric caeca none. Genera 4; species about 10; small fishes living on sandy shores of tropical

America. This family is nearly related to *Uranoscopidae*, of which group it seems to be a reduced or degenerate branch. Its relations with the Asiatic family *Leptoscopidae* are most intimate, the incomplete ventrals and simple pectoral rays of *Dactyloscopidae* being the chief distinctive features. (*Dactyloscopidae*, Gill, Arrangm. Families Fishes, 1872.)

- a. Dorsal fin divided, the first dorsal composed of 3 spines inserted on the nape; head not cuboid; chin without flap; fringes of lips small. GILLELLUS, 852.
- aa. Dorsal fin continuous.
- b. Dorsal fin commencing at the nape; pseudobranchiae very small or obsolete; head cuboid. DACTYLOSCOPUS, 853.
- bb. Dorsal fin commencing far behind the nape; pseudobranchiae well developed.
- c. Head cuboid, formed as in *Dactyloscopus*; the mouth vertical. DACTYLAGNUS, 854.
- cc. Head elongate-conoid, the lower jaw projecting, with a fleshy flap at tip. MYXODAGNUS, 855.

### 852. GILLELLUS, Gilbert.

*Gillellus*, GILBERT, Proc. U. S. Nat. Mus. 1890, 98 (*semicinctus*).

A separate dorsal fin on the nape composed of 3 spines. Lateral line descending posteriorly, its dorsal and median portions about equal. Fringes of upper lip obsolete, those of lower lip little evident. Head not cuboid, the mouth moderately oblique, the lower jaw rounded in front and without symphyseal flap. The physiognomy is intermediate between *Dactyloscopus* and *Myxodagnus*, from each of which the genus is well separated by the characters of the dorsal fin and the lateral line. ("Named in honor of Dr. Theodore Gill, to whom we owe our knowledge of the previously described members of this most interesting group." Gilbert.)

- a. Tip of lower jaw projecting.
- b. Anterior portion of lateral line longer than posterior portion; the scales 25 to 28 + 3 + 15 to 18 = 43 to 49. D. III-IX, 28; A. II, 30 or 31. SEMICINCTUS, 2634.
- bb. Anterior portion of lateral line much shorter than posterior portion, 2½ times in the latter; scales 18 + 3 + 27 = 48. D. II-IX, 31; A. II, 35. ARENICOLA, 2635.
- aa. Tip of lower jaw scarcely projecting; anterior portion of lateral line 1½ times in posterior. D. I-IX, 31; A. II, 34. ORNATUS, 2636.

### 2634. GILLELLUS SEMICINCTUS, Gilbert.

Head 3½; depth 5½. D. III-IX or X, 28; A. II, 30 or 31; scales 25 to 28-3-15 to 18 (43 to 49 scales in all). Body deep, tapering rapidly either way from front of dorsal. Mouth moderately oblique, the maxillary extending beyond orbit, 3 in head; tip of lower jaw projecting; teeth in a narrow band in front of jaws, becoming a single series laterally; none of the teeth enlarged. Opercular fringes well developed, 8 or 9 in number; fold of membrane between rami of lower jaw well developed; pseudobranchiae apparently not developed; gill rakers obsolete. Dorsal fin beginning at a distance from occiput less than diameter of eye, the first 3 rays entirely detached from the rest of the fin, the first ray the highest, the second and third shortened; of the remaining part of the fin the first 9 or

10 rays caudal a along th and base portion c ceons, th hind and the lower duncle, a of sides 1 cross bars covering ward and and anter obtained 2829, and the coast Atlantic fo

*Gillellus semicinctus* Gilbert  
Gulf of C

Head 4½ in v very slende mandible p fringes app orbit; eye fringes near terior dorsal rated by a s Lateral line separated by the posterior ish blotches the alternat sides, as do median porti lncent. A s bert.) (*arenicola*)

*Gillellus arenicola*

Head 4½ in l With the elor differing in th eral line. He slightly longe

10 rays are unarticulated and spinous; first 2 anal rays not articulated; caudal about  $1\frac{1}{2}$  in head; pectorals  $1\frac{1}{2}$ . Lateral line running anteriorly along the very base of spinous dorsal, no scales intervening between it and base of fin; it descends to middle of sides posteriorly, the median portion of its length shorter than the dorsal portion. Color light olivaceous, the back with 6 broad cross bars of pink, narrowly margined behind and in front with blackish, terminating below on middle of sides; the lower of these bars frequently black; a black bar across caudal peduncle, and sometimes a black line at base of caudal; along median line of sides frequently a series of small black spots alternating with the cross bars; a similar series along median dorsal line; a large pink blotch covering occiput; a dusky bar across interorbital space, running downward and backward across cheek; silvery spots and blotches on cheeks and anterior portions of opercles; fins unmarked. Specimens have been obtained in the Gulf of California by the *Albatross*, at Stations 2827 and 2829, and by the *Grampus* in the Atlantic, at Stations 5108 and 5112, off the coast of Florida; no specific difference among them noticed, but the Atlantic form needs further study. (Gilbert.) (*semi*, half; *cinctus*, belted.)

*Gillellus semicinctus*, GILBERT, Proc. U. S. Nat. Mus. 1890, 98, Albatross Stations 2827, 2829, Gulf of California (Coll. *Albatross*); JORDAN, Proc. Cal. A. C. Sci. 1896, 229, pl. 32.

2635. GILLELLUS ARENICOLA, Gilbert.

Head  $4\frac{3}{4}$  in length; depth  $8\frac{1}{4}$ . D. II-IX, 31; A. II, 35; scales 18-3-27. Body very slender and elongate, much as in *Myxodagmus*, the snout sharp, the mandible produced at symphysis and conspicuously projecting; labial fringes apparently obsolete; maxillary reaching vertical from middle of orbit; eye small, about equaling length of snout, 6 in head; opercular fringes nearly obsolete, 3 or 4 small ones at upper edge of opercle. Anterior dorsal inserted close behind occiput, composed of 3 rays, and separated by a short interspace from rest of fin; pectorals longer than head. Lateral line anteriorly running along base of dorsal, from which it is not separated by intervening scales, the anterior portion contained  $2\frac{1}{2}$  times in the posterior median portion. Color light olivaceous, the head with grayish blotches and small pearly spots; 11 dark bars downward from back, the alternate ones narrower and fainter and not extending to middle of sides, as do the others; the margins of the larger bars darker than the median portion, the bars not continued onto dorsal fin; all the fins translucent. A single specimen  $1\frac{1}{2}$  inches long, from Cape San Lucas. (Gilbert.) (*arena*, sand; *colo*, I inhabit.)

*Gillellus arenicola*, GILBERT, Proc. U. S. Nat. Mus. 1890, 99, Cape San Lucas. (Coll. Gilbert.)

2636. GILLELLUS ORNATUS, Gilbert.

Head  $4\frac{1}{2}$  in length; depth 8. D. III-IX, 31; A. II, 34; scales not counted. With the elongate form and general appearance of *Gillellus arenicola*, but differing in the subequal jaws and in the long anterior portion of the lateral line. Head conical, acute, very small; jaws nearly equal, the lower slightly longer than the upper, but not noticeably protruding. In this

respect the species resembles most strongly *G. semicinctus*, from which it varies widely in the general form and proportions. Snout extremely short, scarcely equaling diameter of the minute eye; diameter of orbit about 7 in head. Mouth oblique, the maxillary 4 in head, reaching nearly to vertical from posterior margin of orbit. Lips without fringes. Eyes separated by a narrow septum, the interorbital width being less than the diameter of the pupil. Opercular fringes few and small, flat, and not terminating evident ridges as in *Dactyloscopus*. Dorsal beginning well forward, its origin less than diameter of orbit behind the posterior line of occiput; anterior detached part of fin consisting apparently of 3 rays, the first of which is the longest, the second and third equal and short; fourth spine again longer; spines as usual slenderer than the rays, and showing no articulations, but with some difficulty discriminated from them; pectoral as long as head. Anterior part of lateral line running immediately along base of dorsal, without intervening scales, as in other members of this genus. It is much longer than in *G. arenicola* and is contained  $1\frac{1}{2}$  times in the posterior median portion. There are 3 scales between the posterior part of the lateral line and the base of the dorsal. Color similar to that of *G. arenicola* and *G. semicinctus*, light olivaceous, unmarked below the middle of the sides, the back and upper half of sides with 8 brown bars which extend downward to lateral line; the upper part of each bar with a lighter central area, the light areas between the bars marked more or less with brown, which sometimes forms indistinct secondary bars; a blackish bar at base of caudal, and a faint streak below eye; a large pearly blotch on opercle. A single specimen, about 2 inches long, from Albatross Station 2828 in the Gulf of California. (Gilbert.) (*ornatus*, adorned.)

*Gillulus ornatus*, GILBERT, Proc. U. S. Nat. Mus. 1891, 558, Gulf of California. (Coll. Gilbert.)

### 853. DACTYLOSCOPUS,\* Gill.

*Dactyloscopus*, GILL, Proc. Ac. Nat. Sci. Phila. 1859, 132 (*tridigitatus*).

*Esloscopus*, JORDAN & EVERMANN, Check-List Fishes, 465, 1896 (*zelotes*).

Body moderately elongate, covered with rather large, cycloid scales; head cuboid, oblong, and nearly flat above; eyes small; interorbital space broad; mouth nearly vertical; lower jaw not dilated beneath nor emarginate in front, without barbels; no intralabial filament; teeth villiform, on jaws only; pseudobranchiæ very small or obsolete. Dorsal commencing at the nape, with 6 or 12 slender spines, the soft rays numerous; anal

\*This genus is thus defined by Dr. Gill: "Body elongate with the dorsal and abdominal outlines slowly converging to the caudal fin. Scales large, regularly imbricated. Lateral line straight, and running along the middle of the side. Head oblong, subcubical, and smooth. Preopercle entire, opercle radially fringed behind. Mouth nearly vertical. Tongue thick, narrowed anteriorly, attached to the floor of the mouth. Labial velum without a barbel. Anus a short distance behind the base of the pectoral fins. Dorsal fin subequal, single, and very long, commencing above or before the anus and continued almost to the base of the caudal. Anal fin commencing behind the anus, and with the same form and termination as the dorsal. Caudal fin small and narrow, posteriorly subtruncated. Pectoral fins subangular. Ventral fins jugular, closely approximated, and each with 3 stout simple and articulated rays."

inserted be  
gazer, shor

DACTYLOSCOP

a. Dorsal r

b. Soft

bb. Sof

c.

co

ESLOSOPUS (E

aa. Dorsal r

Head abou

XII, 22; A. 1

total length

diameter of e

*tridigitatus*;

ments; origi

origin of ana

2 anal rays

yellow, with

lines of abou

lines proceed

angles of ad

often present

above. Each

from the ant

sinuated neck

by a dark lin

*toralis*, pertai

*Dactyloscopus pe*

John Xantus

Head 5 (in

V. I. 3; scales

only; opercula

the lower ang

margin of the

length of bod

evident in liv

part whitish;

inserted behind dorsal; ventral rays I, 3. (δάκτυλος, finger; σκοπός, gazer, short for *Uranoscopus*.)

DACTYLOSCOPUS:

- a. Dorsal rays X to XII, 22 to 31; anal rays less than 35.
- b. Soft dorsal with 22 soft rays; anal with 26. PECTORALIS, 2637.
- bb. Soft dorsal with 28 to 31 rays; anal with 32 or 33; scales about 45.
- c. Body rather slender, the depth about 6 in length (7 with caudal); opercular fringe of 15 filaments. TRIDIGITATUS, 2638.
- cc. Body rather stout, the depth  $5\frac{1}{2}$  in length (6 in total with caudal); opercular fringe of 18 filaments.
- d. Back not barred; head blotched and dotted. POEYI, 2639.
- dd. Back with about 10 pale cross bars; head marked with whitish; a dark bar at base of caudal. LUNATICUS, 2640.

ESLOSCOPUS (έσλός, good; σκοπός for *Uranoscopus*):

- aa. Dorsal rays VI, 38; anal rays II, 37; scales 6-51-5. ZELOTES, 2641.

Subgenus DACTYLOSCOPUS.

2637. DACTYLOSCOPUS PECTORALIS, GILL.

Head about 5 in total length with caudal; depth about 7 (in total). D. XII, 22; A. II, 26; P. 12; V. I, 3. Width of head behind operculum 7 in total length with caudal; eye small, 10 in head; interorbital space equals diameter of eye; preoperculum broader at the angle than in *Dactyloscopus tridigitatus*; pores well developed; opercular fringe of 11 or 12 free filaments; origin of dorsal between  $\frac{1}{2}$  and  $\frac{1}{3}$  length of fish from tip of snout; origin of anal under sixth or seventh dorsal ray, the first 12 dorsal and 2 anal rays simple. Pseudobranchiæ obsolete. Color light brownish yellow, with dark spots on the back, arranged in lines forming the outlines of about 6 quadrangular areas, from the angles of which irregular lines proceed downward, converging toward those departing from the angles of adjoining areas; more scattered and irregular spots and dots often present below the lateral line; head lighter, diffused with pink above. Each orbit with 4 diverging bands, 1 in front, a bifurcated one from the antero-inferior angle, and 2 from posterior border, a transverse sinuated nuchal line; upper angle of operculum whitish, bounded in front by a dark line or spot. (Gill.) Cape San Lucas; not seen by us. (*pectoralis*, pertaining to the breast.)

*Dactyloscopus pectoralis*, GILL, Proc. Ac. Nat. Sci. Phila. 1861, 267, Cape San Lucas. (Coll. John Xantus.)

2638. DACTYLOSCOPUS TRIDIGITATUS, GILL

Head 5 (in total) with caudal; depth 7. D. XII, 28; A. II, 32; P. 13; V. I, 3; scales  $11 + 4 + 30 = 45$ . Body slender, much compressed posteriorly; opercular fringe of 15 separate filaments. Origin of dorsal fin over the lower angle of the base of the pectorals, or immediately before the margin of the operculum, its distance from snout to dorsal 5 in total length of body. Pseudobranchiæ very small (overlooked by Dr. Gill, but evident in living specimens). In life, pale sand color above, the lower part whitish; above 12 narrow cross bands of whitish on the back, not

extending down far on the sides; head mottled above; fins all pale. West Indies, north to Key West; rather common in coral sand in shallow water about Key West. (*tres*, three; *digitus*, finger, from the 3 ventral rays.)

*Dactyloscopus tridigitatus*, GILL, Proc. Ac. Nat. Sci. Phila. 1859, 132, Barbados (Coll. Dr. Gill); GILL, *I. c.*, 1861, 264; GILL, *I. c.*, 1862, 505; GÜNTHER, Cat., III, 279, 1861; JORDAN & GILBERT, Synopsis, 753, 1883; JORDAN, Proc. U. S. Nat. Mus. 1884, 140.

2639. DACTYLOSCOPUS POEYI, GILL.

Head 5 in total length; depth  $6\frac{1}{2}$  in total. D. XI, 31; A. II, 32. Body more robust than in *D. tridigitatus*; head plane above and obtusely angulated at the sides of the plane; thickness of the head behind the preoperculum exceeding  $\frac{1}{2}$  of its length; interorbital space  $\frac{2}{3}$  diameter of eye. Eye about 7 in head; preopercle as in *D. tridigitatus*, pores indistinct or obsolete; opercular fringe of about 18 filaments, the lowest of which are scarcely extended beyond the margin; origin of dorsal fin  $\frac{1}{2}$  distance from tip of snout; origin of anal fin under sixth dorsal ray; scales of moderate size and regularly imbricated. Color reddish brown, dotted with darker above the lateral line; head blotched and dotted with darker; opercles variegated; opercular bones nearly immaculate. (Gill.) Cuba; not seen by us. (Named for Prof. Felipe Poey.)

*Dactyloscopus poeyi*, GILL, Proc. Ac. Nat. Sci. Phila. 1861, 266, Cuba. (Coll. Felipe Poey.)

2640. DACTYLOSCOPUS LUNATICUS, Gilbert.

Head (to end of opercular fringes)  $3\frac{2}{3}$ , from tip of lower jaw to 1 fringes 4; depth greater than in related species,  $5\frac{1}{2}$  in length. D. X or XI, 29 or 30; A. II, 32 or 33; scales about  $11 + 4 + 30 =$  about 45. Head cuboid, narrowed forward, the vertex gently convex; width at occiput  $\frac{1}{2}$  length of head (to base of fringes on opercle). Mouth nearly vertical, maxillary  $2\frac{2}{3}$  in head. Labial fringes short but evident. A short nasal filament. Teeth in a rather broad cardiform band on front of upper jaw, becoming narrow laterally; in lower jaw a single series, or an irregular double series anteriorly; vomer and palatines toothless. Eyes small, very close together, the interorbital width about  $\frac{1}{2}$  their diameter, which equals length of snout, or about  $\frac{1}{3}$  head. Gill laminae much reduced in size; a small round pore behind inner arch. Gill rakers obsolete; pseudo-branchiae small but evident. Opercular fringes composed of 18 filaments. Dorsal beginning at a distance behind occiput equaling diameter of orbit, its anterior rays but partly joined by membrane, the first 10 or 11 slender and not articulated, the last ray distant from base of caudal about a diameter of orbit; origin of anal under sixth dorsal spine, the 2 anterior rays not articulated; pectorals short,  $1\frac{2}{3}$  in head, containing 14 or 15 rays; caudal very small, with 10 developed rays, its length  $2\frac{2}{3}$  in head. Lateral line running high in its anterior portion, declining on 3 or 4 scales, the posterior portion on middle of sides with 29 or 30 tubes; 4 scales between median portions of lateral line and base of dorsal. Color light olivaceous, a dark streak along back, 1

along n  
darker r  
less evi  
mandibl  
shape, l  
black; r  
a narrow  
longest t  
fathoms.

*Dactylosco*  
*Albatro*

Head 4  
B. 6. He  
 $\frac{1}{2}$  length  
anal fin,  
compress  
very small  
length of  
in a short  
as in *Ura*

behind or  
teeth on t  
wide, flex  
fins, the f  
opercle to  
branchios  
forming a  
round pore  
from snou  
those follo  
tion can be  
differentia  
fourth dor  
under ante  
Scales larg  
behind pec  
opercle, ru  
downward  
ings of the  
fringes ele  
base; eyes  
present des  
(ζηλώτης, a

*Dactyloscopus*  
*Dactyloscopus*  
Dow).



along middle of sides, and a fainter one along base of anal, formed by darker margins to the scales; median dorsal line with 10 or 11 more or less evident narrow pearly white cross bars; top of head and front of mandible colored like the back, the pearly blotches varying in size and shape, but symmetrically arranged, many of them narrowly edged with black; nasal tentacle white; white streaks on preopercle; caudal with a narrow black bar at base. Gulf of California. Three specimens, the longest 3 inches, from *Albatross* Stations 2797 and 3012, the latter in 22 fathoms. (Gilbe t.) (*lunaticus*, moon-struck.)

*Dactyloscopus lunaticus*, GILBERT, Proc. U. S. Nat. Mus. 1890, 99, Gulf of California. (Coll. *Albatross*.)

Subgenus *ESLOSCOPUS*, Jordan & Evermann.

2641. *DACTYLOSCOPUS ZELOTES*, Jordan & Gilbert, new species.

Head  $4\frac{1}{2}$  in length; depth  $6\frac{3}{4}$ . D. VI, 38; A. II, 37; V. 3; scales 6-51-5; B. 6. Head and body slender, compressed, the greatest width at occiput,  $\frac{1}{4}$  length of head; the greatest depth immediately behind insertion of anal fin, thence tapering to a very narrow tail. Head narrow, enboid, compressed, the upper surface nearly plane, the cheeks vertical. Eyes very small, superior, with little lateral range; diameter of orbit about  $\frac{1}{5}$  length of head; snout very short, about equaling orbit; anterior nostril in a short tube; gape subvertical, the lower jaw very heavy, projecting, as in *Uranoscopus*; premaxillaries protractile, the processes reaching far behind orbits; lips fringed; both jaws with bands of villiform teeth; no teeth on tongue, vomer, or palatines. Subopercle and interopercle very wide, flexible, striate, the latter overlapping throat and base of ventral fins, the former wholly covering base of pectoral fins; the striations of opercle terminate posteriorly in a wide, coarse, membranaceous fringe; branchiostegal membranes not united, free from the isthmus; pubic bones forming a sharp projection at throat; no pseudobranchiae; gills small, a round pore behind the fourth. Dorsal beginning on the nape, its distance from snout about equaling depth of body, the first 6 rays shorter than those following and not connected by membrane; as no traces of articulation can be found, they are probably flexible spines, but are not clearly differentiated from those immediately following; origin of anal under fourth dorsal spine; caudal distinct, narrow, short; ventrals inserted under anterior margin of preopercle; ventrals 2 in head; pectorals  $1\frac{1}{2}$ . Scales large, with entire edges, wanting on head, breast, and region behind pectoral fins. Lateral line beginning at upper posterior angle of opercle, running parallel with the back on about 12 scales, then obliquely downward to middle of body. Color in spirits, light olivaceous, the edgings of the scales, some vermiculations on top of head, and the labial fringes clear brown; fins translucent, the caudal with a brown bar at base; eyes dark. Length  $3\frac{1}{2}$  inches. Panama; 1 specimen known. The present description copied from the original in Proc. Nat. Mus. 1882, 628. (*Ζηλότης*, an imitator, from its resemblance to *Dactylagnus mundus*.)

*Dactyloscopus*, sp. nov., JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1882, 628, Panama.

*Dactyloscopus zelotes*, JORDAN & GILBERT, new species (MS. 1882), Panama (Coll. Capt. Dow).

## 854. DACTYLAGNUS, Gill.

*Dactylagnus*, GILL, Proc. Ac. Nat. Sci. Phila. 1862, 505 (*mundus*).

Body moderately elongated, covered with rather large and uniform scales. Head cuboid, oblong, scarcely convex transversely above. Eyes small, directed obliquely upward, and situated near the snout on the upper surface of the head. Interorbital area moderate and channeled. Mouth very oblique or subvertical, the snout truncated in front; lower jaw transversely convex in front and with no barbel; teeth acute, in a narrow band along each jaw; palate smooth. Dorsal fin perfectly entire, commencing rather farther behind than the anal, and with its anterior portion armed with about 10 slender spines; anal fin longer than the dorsal. This genus closely resembles *Dactyloscopus* externally. It differs from the latter genus chiefly in the structure of the dorsal fin and the well-developed pseudobranchiæ. (*δάκτυλος*, finger; *ἄγνος*, *Agnus*, an old name of *Uranoscopus scaber*.)

## 2642. DACTYLAGNUS MUNDUS, Gill.

Head  $4\frac{1}{2}$ ; depth  $6\frac{1}{2}$ . D. X, 31; A. II, 38; scales 2-48-10; eye 6 in head; maxillary  $2\frac{1}{2}$ ; snout equals eye; highest dorsal spine 3; highest anal ray  $2\frac{1}{2}$ ; pectoral equals head; caudal  $1\frac{1}{2}$ . Body elongate, compressed, tapering posteriorly; upper profile of head nearly horizontal, slightly convex; eyes superior, looking upward; interorbital narrow, concave; lower jaw strongly projecting, mouth nearly vertical; teeth small and conical, in narrow bands, widest in front; vomer and palatines toothless; lips furnished with labial fringes about as long as diameter of eye; nostril ending in a tube; preopercle entire; opercle fringed on its upper edge, a flap of skin downward from opercle covers the branchiostegals; pseudobranchiæ present; gill rakers not developed; head and belly naked; fins naked. Lateral line running near the back through 14 scales, deflected on 4, and thence continued along the middle through 36. Dorsal low, long, and continuous, distance from its origin to tip of snout  $3\frac{1}{2}$  in body; anal similar, slightly higher and longer; posterior rays of dorsal and anal reaching to base of caudal rays; upper rays of pectoral the longest, reaching to the vertical from tenth anal ray, the lower rays short, graduated, tip of fin slightly curved up; origin of ventrals in front of pectorals, the inner rays the longest, reaching about to vent; caudal truncate, or very slightly rounded. Color in spirits, light brown above, white below, each scale on back with a dark brown spot; top of head with a few brown spots; fins colorless. Length  $4\frac{1}{2}$  inches. Gulf of California. Here described from specimens collected by the *Albatross* at Carmen Island, Gulf of California; the type from Cape San Lucas. (*mundus*, neat.)

*Dactylagnus mundus*, GILL, Proc. Ac. Nat. Sci. Phila. 1862, 505, Cape San Lucas. (Coll. Xantus.)

*Myzodagnus*

This genus is elongate, short fin dorsal fin a genus old name

Head pectoral front of acutely the eyes between

rated by tical; op as long a 6 or 7 sh continuo mencing reaching curve in caudal t arranged on the si along the brown, re the scales operculum Described collectors

*Myzodagnus*  
Xantus

Head l conic, sub either nak

This gen tained abou as long as l extremely r furnished w uniform te anns, and fr the dorsal.

855. MYXODAGNUS,\* Gill.

*Myxodagnus*, GILL, Proc. Ac. Nat. Sci. Phila. 1861, 269 (*opercularis*).

This genus differs from *Dactyloscopus* in the form of the head, which is elongate-conoid, the lower jaw obtusely pointed and provided with a short flap in front. The pseudobranchia are well developed and the dorsal fin commences far behind the nape. One species known. (*Myxodes*, a genus of blennies, which this fish resembles in form; *Agnus*, ἄγνος, an old name of *Uranoscopus scaber*.)

2643. MYXODAGNUS OPERCULARIS, Gill.

Head 5 without lower jaw; depth 7. D. 36; A. II, 36; scales 2-44-9; pectorals equal head; ventrals 1½; caudal 1½. The body is deepest at front of dorsal fin, tapering regularly to the caudal fin. Head elongated, acutely conical; profile nearly straight, but slightly concave in front of the eyes; the crown is transversely arched and smooth; the frontal bones between the eyes are exceedingly narrow, so that the orbits appear separated by little more than a mere septum; eyes large, longitudinally elliptical; opercular pores obsolete; the postorbital or temporal ridge is nearly as long as the diameter of the orbit; the opercular fringe is composed of 6 or 7 short filaments; origin of dorsal above vent, the fin very low and continuous, its last rays not reaching to base of caudal rays; anal commencing slightly in front of dorsal, similar to it but higher, its last ray reaching to base of caudal rays; pectoral large and pointed, reaching to curve in lateral line; rays of ventral subequal, reaching about to vent; caudal truncate; scales moderate, finely striated concentrically and arranged in 11 rows on each side; the lateral line runs through 12 scales on the sides of the back, is then deflected through 3, and thence runs along the fifth row from the back through 36. Color light yellowish brown, rendered darker on the back by congregations of dark spots on the scales; there is a pearly patch behind and beneath the eye, and the operculum is also colored in the same manner. (Gill.) Cape San Lucas. Described from a specimen 2½ inches in length. Not obtained by recent collectors. (*opercularis*, pertaining to the gill cover.)

*Myxodagnus opercularis*, GILL, Proc. Ac. Nat. Sci. Phila. 1861, 270, Cape San Lucas. (Coll. Xantus.)

Family CXCVII. URANOSCOPIDÆ.

(THE STAR-GAZERS.)

Head large, broad, partly covered with bony plates. Body elongate, conic, subcompressed, widest and usually deepest at the occiput. Body either naked or covered with very small, smooth, adherent scales, which

\* This genus is thus defined by Dr. Gill: Body quite slender, the greatest height contained about 10 times in length. Head rather elongated and acutely conical, about twice as long as high; eyes large and elliptical, and very closely approximated; frontal bones extremely narrow. Mouth oblique; lower jaw projecting much beyond the upper and furnished with a short, compressed, and wide flap or barbel in front of the symphysis; villiform teeth present only on the jaws. Dorsal fin inserted behind the vertical of the anus, and furnished with simple and articulated rays; anal fin as long as or longer than the dorsal.

are arranged in very oblique series running downward and backward; the scales on the belly inconspicuous or obsolete. Lateral line little developed, running high. Eyes small, on anterior and upper portion of head, with vertical rings. Mouth vertical, with strong and prominent mandible; teeth moderate, on jaws, vomer, and palatines. Premaxillaries freely protractile; maxillary broad, without supplemental bones, not slipping under the preorbital. Gill openings wide, continued forward; gill membranes nearly separate, free from isthmus. Pseudobranchiae present; 6 branchiostegals;  $3\frac{1}{2}$  gills, a slit behind the last; no anal papilla. Spinous dorsal very short or wanting; second dorsal long. Anal and pectorals large, the latter with broad oblique bases, the lower rays rapidly shortened, most of them branched; ventrals jugular, close together, 1, 5, the spine very short, innermost ray longest; caudal not forked. Air bladder generally absent; pyloric caeca in moderate number. Vertebrae 24 to 26. Carnivorous fishes, living on the bottom of the shores of most warm regions. Genera 8; species 25.

## URANOSCOPINÆ:

a. Spinous dorsal of 4 or 5 well developed spines; scales present.

b. Head above not entirely covered with bone, the occipital plate ceasing much behind the orbits; from the middle line anteriorly a Y-shaped bony process extends forward, the tips of the fork between the eyes; a trapezoidal space on either side of the Y, covered by naked skin, bounded by the Y, the eyes, the suborbitals, and the occipital plate. A covered furrow behind and on the inner side of each eye terminating near front of orbits, its edges fringed. Head without spines; humeral spine obsolete; lips and nostrils fringed; no retractile tentacle in mouth. ASTROSCOPUS, 856.

## KATHETOSTOMATINÆ:

aa. Spinous dorsal obsolete; no scales; head above covered with bone except the groove of the premaxillary spine; the bony occipital plate coalescing with the orbital rims; humeral spine well developed; no distinct protuberances on top of head; no spine in front of humeral spine; 2 small forward-directed spines in front of eye; 3 small spines on lower margin of preopercle; upper lip scarcely fringed; no retractile tentacle in mouth; ventral fin not largely adnate to abdomen. KATHETOSTOMA, 857.

## 856. ASTROSCOPUS, Brevoort.

(ELECTRIC STAR-GAZERS.)

*Astroscopus* (BREVOORT) GILL, Proc. Ac. Nat. Sci. Phila. 1860, 20 (*anoplos*; young).

*Agnus*, GÜNTHER, Cat. Fishes, II, 229, 1860 (*anoplos*).

*Upsilonphorus*, GILL, Proc. Ac. Nat. Sci. Phila. 1861, 113 (misprint for *Upsilonphorus*) (*græcum*; adult).

Body robust. Head above not entirely covered with bone, the occipital plate ceasing much behind the orbits; from the middle line anteriorly a Y-shaped bony process extends forward, the tips of the fork between the eyes; a trapezoidal space on either side of the Y, covered by naked skin, bounded by the Y, the eyes, the suborbitals, and the occipital plate. A covered furrow behind and on the inner side of each eye terminating near front of orbits, its edges fringed. Head without spines; humeral spine obsolete; lips and nostrils fringed; no retractile tentacle in mouth. Young individuals with top of head largely covered by bone. Head scaleless; back and sides covered with close-set scales; belly mostly naked. Humeral

spin  
low,  
whic  
Ame  
by t

a. I

aa.

He  
eye I  
dorsa  
very  
post  
verti  
eter  
mem  
ical,  
teeth  
very  
times  
ridg  
area;  
but sh  
trils fr  
ular r  
obliqu  
forwar  
spines  
and lu  
brauel  
from fi  
scales  
less th  
very sl  
pointe  
of sixt  
\* The  
spine w  
directe  
stout p  
extendi  
tentacle  
scales w

spine obsolete; no spine before the ventrals. First dorsal small, of 4 or 5 low, stout, pungent spines, connected by membrane to the second dorsal, which is rather high and long; pectorals and ventrals large. Species American, distinguished from the Old World genus, *Uranoscopus*,\* chiefly by the unarmed head. (*ἀστρον*, star; *βιοπέω*, v. look.)

a. Naked space between forks of the Y on top of head long and narrow, but shorter than the vertical limb of the Y; no distinct spines before eye; sides with round pale spots, each with a dark ring.

b. Dorsal spines 4, rather high; scales normal.

Y-GRECUM, 2644.

bb. Dorsal spines 5, lower than in *y-græcum*; scales of sides cohering in oblique series.

ZEPHYREUS, 2645.

aa. Naked space between the forks of the Y short and broad, but longer than the very short vertical limb of the Y; 2 distinct spines directed forward before the eye; sides with small pale spots, not dark-edged.

GUTTATUS, 2646.

2644. ASTROSCOPUS Y-GRECUM (Cuvier & Valenciennes).

Head, without lower jaw,  $2\frac{2}{3}$ ; depth  $3\frac{1}{2}$ . D. IV-I, 12; A. 13; scales 80; eye  $1\frac{1}{2}$  in head; maxillary 2; pectoral  $1\frac{1}{10}$ ; second dorsal spine 4; highest dorsal ray 2; highest anal ray 3; caudal  $1\frac{1}{2}$ . Body moderately elongate, very robust forward, greatest depth at occiput; anteriorly subcylindrical, posteriorly somewhat compressed. Head large and broad; mouth large, vertical, a fringe of barbels on each jaw, slightly longer than the diameter of the eye; tongue extremely large and fleshy, forming a pad under membrane of lower jaw which projects forward somewhat. Teeth conical, small and movable, in many bands in upper jaw, in lower jaw the teeth are larger and in fewer bands; teeth on vomer and palatines. Eyes very small but prominent, set on top of head; interorbital very wide,  $3\frac{1}{2}$  times wider than the eye; bones on top of head coarsely granular; Y-shaped ridge on top of head conspicuous, on each side of which is a broad naked area; naked space between forks of Y on top of head long and narrow, but shorter than vertical limb of the Y which is very long; edges of nostrils fringed, anterior nostril round, separated from the eye by a high granular ridge; posterior nostril ending in a long curved furrow, which runs obliquely across the naked area behind eye, its posterior end not curved forward, its length  $2\frac{1}{2}$  times the diameter of the eye; 2 or 3 small blunt spines in front of the eye; surface of the bones of opercle, preopercle, and humeral process coarsely granular; gill rakers not developed; pseudo-branchiae small. Head entirely scaleless; belly naked below a line drawn from fifth anal ray to upper end of pectoral base; fins without scales; scales very small and somewhat embedded. Width of pectoral at base less than  $\frac{1}{2}$  length of the head, the upper rays longest, the lower rays very short, graduated from the lower side to the upper; fin somewhat pointed behind and curved up, its tip reaching to the vertical from base of sixth dorsal ray; the rays of ventrals very thick and swollen, the inner

\* The following are the characters of *Uranoscopus*: Head with spines; humeral spine well developed; 1 strong spine on subopercle, 4 smaller ones on preopercle, all directed downward; 1 small spine directly above and in front of humeral spine; 4 low, stout protuberances on top of head pointing backward; naked space between eyes extending back to posterior part of orbits; upper lip and nostrils not fringed; retractile tentacle in mouth more or less developed. First dorsal with about 4 pungent spines; scales well developed.

rays the longest, reaching midway from their base to end of pectorals; origin of fin a distance of the width of pectoral in front of the lower edge of pectoral base; soft dorsal much higher than anal; posterior rays reaching slightly past the vertical from base of the last anal ray; end of the last anal ray about reaching to base of caudal rays; caudal truncate or slightly rounded; a ridge of skin along middle of belly from the ventrals to vent. Dark brown above, paler below; upper parts densely covered with small rounded white spots, each surrounded by a black ring; lower jaw and labial fringes similarly spotted; spinous dorsal black, white posteriorly; soft dorsal brown anteriorly with a horizontal white and black band, then tipped with white; posteriorly with 2 vertical black stripes and a white one between them; caudal black, tipped with white, with 2 to 4 white longitudinal stripes, its upper and lower edges narrowly white; the anal white at base and tip, with a black median band,  $\frac{1}{2}$  depth of fin, darkest posteriorly; pectorals brown, with a black band below, the lower edge white, the upper ray spotted; ventrals white with a black lengthwise streak. Old examples lose the black ring around the spots, and the edges of the spots are blended into the dark brown of the back; a dark stripe running from the upper angle of gill opening to caudal. South Atlantic coast from Cape Hatteras to the Caribbean Sea, in sandy bays, rather common in shallow water, varying much with age. Here described from a specimen, 15 inches in length, from Charleston, South Carolina. It is recorded from Charleston, Beaufort, Matanzas River, St. Johns River, Pensacola, Key West, and "the Caribbean Sea." According to Dr. James A. Henshall, the naked area on top of the head in *Astroscoptes* is the seat of electric power. This interesting statement needs verification. (Named from the armature of the head, in the form of the Greek  $\tau$ .)

*Uranoscopus y-græcum*, CUVIER & VALENCIENNES, Hist. Nat. Poiss., III, 308, 1829, origin unknown; GÜNTHER, Cat., II, 229, 1860.

*Uranoscopus anoplos*,\* CUVIER & VALENCIENNES, Hist. Nat. Poiss., VIII, 493, 1831 (young examples), Charleston, South Carolina.

*Upsilonphorus y-græcum*, GILL, Proc. Ac. Nat. Sci. Phila. 1861, 113; KIRSCH, l. c., 263, 1889.

*Astroscoptes y-græcum*, BEAN, Proc. U. S. Nat. Mus. 1879, 58; JORDAN & GILBERT, Synopsis, 628, 1883.

*Agnus anoplos*, GÜNTHER, Cat., II, 229, 1860.

*Astroscoptes anoplos*, JORDAN & GILBERT, Synopsis, 629, 1883.

*Astroscoptes anoplos*, KIRSCH, Proc. Ac. Nat. Sci. Phila. 1889, 262.

\* The genus *Astroscoptes* was based on small specimens which, in our present opinion are simply immature examples of the species *y-græcum*. The supposed genus is thus described in distinction from *Upsilonphorus*, which seems to us the adult of the same type: Head covered above with bone except a small region between and in front of the eyes, the bony occipital plate coalescing with the orbital fins; no spines on head; humeral spines obsolete; occipital region with bluntish projections; naked space between eyes extending back to near middle of orbits; lips and nostrils fringed; no retractile tentacle in mouth.

The following characters are assigned by Dr. Kirsch to *Astroscoptes anoplos*: Head 24; depth 34. D. IV-1, 13; A. 13. Pectorals rather large, their longest ray equaling in length base of second dorsal and extending to front of that fin; ventrals equaling pectorals in length, and extending to front of that fin; the second dorsal equaling anal but its anterior insertion slightly posterior to that; anal rays reaching base of caudal; vent much nearer base of caudal than to tip of snout. Color dark brown above, yellowish below; lighter portions of body covered with small white specks; chin jet-black; all the fins whitish. Length 2 inches. (Specimen from Key West). Small individuals are found along the coast from Cape Hatteras to Florida wherever *A. y-græcum* is found. The adult differs mainly in the armature of the top of the head, a characteristic which is developed at different ages in different individuals.

2645. ASTROSCOPUS ZEPHYREUS, Gilbert & Starks.

Head, without lower jaw,  $2\frac{2}{3}$ ; depth  $3\frac{1}{3}$ . D. V, 13; A. 14; scales 84; eye 12 in head; maxillary  $2\frac{1}{4}$ ; pectoral  $1\frac{1}{4}$ ; second dorsal spine 7; highest dorsal ray  $2\frac{1}{4}$ ; highest anal ray  $3\frac{1}{4}$ ; caudal  $1\frac{1}{4}$ . Body robust, widest at occiput, slightly compressed posteriorly; anteriorly subcylindrical. Head very large and broad, wider than the body; mouth large, vertical, a fringe of barbels curving over mouth on each jaw; length a little greater than the diameter of the eye; tongue very large and fleshy, forming a pad under the membrane of lower jaw, which projects forward somewhat; teeth conical, small and movable, in many bands in upper jaw; in lower jaw the teeth are larger and in 2 or 3 rows; vomer and palatines with teeth; eyes very small but prominent, set on top of head; interorbital very wide, 4 times as wide as the eye; bones on top of head coarsely granular; Y-shaped ridge on top of head conspicuous, on each side of which is a broad naked area, the form of these and other bones of the head exactly as in *A. y-gracum*; edges of nostrils closely fringed, anterior nostril round, the ridge between it and eye not very high or conspicuous; posterior nostril ending in a long curved furrow which runs obliquely across the naked area behind eyes; at its posterior end it turns sharply forward, its length  $2\frac{3}{8}$  times the diameter of the eye; 2 very short blunt spines in front of the eye; surface of the opercle, preopercle, and humeral process granular, not so rough as in *Astrosopus y-gracum*; gill rakers not developed; pseudobranchiae very small. Head entirely scaleless; belly naked below a line drawn from first anal ray to the middle of the pectoral base; fins without scales; scales small and nearly square, grown together side by side, forming series of oblique plates. Width of pectoral at base slightly less than  $\frac{1}{2}$  length of head, the lower rays very short and graduated to the long upper rays, the fin pointed and slightly turned up, its tip reaching to the vertical from base of the third dorsal ray; the ventral rays thick and swollen, the inner rays the longest, its tip reaching about midway between its base and tips of pectorals; origin of fin in front of pectorals a distance equal to the width of pectoral base; soft dorsal somewhat higher than anal, its posterior rays reaching to the vertical from base of last anal ray; tip of last anal ray nearly reaching to the base of caudal rays; caudal truncate or slightly rounded; a fold of skin along middle line of belly from ventrals to vent. Color dark brown above, paler below; upper parts with many round white spots of various sizes, edged with rings of dark brown; spinous dorsal black, light posteriorly; soft dorsal light at base, the ends of the rays with black and white stripes; pectoral and anal dusky with light edge; caudal with longitudinal black and white stripes. Pacific coast of Mexico. One specimen, numbered 333, in the Leland Stanford Junior University Museum, collected by the *Albatross* at Magdalena Bay, Lower California. It is 12 inches in length. A distinct electric shock was given by this fish when alive, the electric organs being in the fleshy areas on top of head behind

eyes. (Gilbert.) A second large specimen was sent from Mazatlan by Dr. George W. Rogers, having been taken by Ygnacio Moreno in January, 1896. (*ζεύριος*, western; *ζεύρος*, the west wind.)

*Astroscopus zephyreus*, (GILBERT & STARKS, Proc. U. S. Nat. Mus. 1896, 453, pl. 53, fig. 2, and pl. 54, Magdalena Bay, Lower California (Type No. 47743. Coll. *Albatross*).

2646. *ASTROSCOPUS GUTTATUS* (Abbott).

Depth 4 in length in young and  $3\frac{1}{2}$  in adult. D. IV or V-13 or 14; A. 13; V. I, 5. Eye  $5\frac{1}{2}$  in interorbital space. Naked space between forks of Y on top of head short and broad, but longer than the vertical limb of the Y, which is very short; 2 distinct spinules directed forward before eye; white spots on body very small and irregular without dark rings; base of dorsals equaling in length the distance from front of first dorsal to tip of snout; base of first dorsal twice length of its longest spine; first spine equaling second in length, and 3 times length of last; length of middle caudal rays a little less than that of ventrals; pectorals slightly longer than ventrals,  $3\frac{1}{2}$  in total length, and extending to fifth anal ray. Color of upper parts of body and lower jaw bright chocolate; belly and throat white; darker portions covered with numerous circular spots much lighter than ground color; membrane of first dorsal black; second dorsal white with 3 irregular bands of dull black obliquely across it; the caudal with 3 parallel bands of blackish brown, the middle of which appears to be the continuation of a variable longitudinal band on the center of each side; the anal having a variable band of dull brown, darker upon the posterior termination. Length 12 inches. Atlantic coast of the United States, from Long Island to Virginia; apparently scarce. Recorded from Cape May; Tompkinsville, New York; Norfolk, Virginia; Somers Point, New Jersey, etc.; not known south of Cape Hatteras. In *Astroscopus guttatus* the pale spots are much smaller, less sharply defined, and occupy a smaller area than in *A. y-gracum*; the lower part of the head has 2 black blotches in each species; the second dorsal, anal, and ventrals are nearly or quite plain. The naked area behind each eye is (in *A. guttatus*) lunate, its length barely twice that of the snout; the bony Y-shaped plate is short and broad, concave on the median line, and forked for about  $\frac{1}{2}$  its length, the posterior undivided portion broader than long; the bony bridge across the occiput but little shorter than the part of the head which precedes it. In *A. y-gracum* the naked area is trapezoidal, longer than broad, and about 4 times the length of the snout; the Y is forked for more than  $\frac{1}{2}$  its length, its undivided part more than twice as long as broad, and not concave; the occipital plate is not  $\frac{1}{2}$  as long as the part of the head which precedes it. (*guttatus*, spotted, as with rain drops.)

*Astroscopus guttatus*, ABBOTT, Proc. Ac. Nat. Sci. Phila. 1860, 365, Cape May, New Jersey.  
*Epsilonphorus guttatus*, BEAN, Proc. U. S. Nat. Mus. 1879, 60; KISSCH, *l. c.*, 264, 1889.



## 857. KATHETOSTOMA, Günther.

*Kathetostoma*, GÜNTHER, Cat. Fish., 11, 231, 1860 (*læve*).

Body robust, formed as in *Astroscopus* and *Uranoscopus*. Scales none. One continuous dorsal without spines; ventrals jugular not adnate to the abdomen; pectoral rays branched; some bones of the head armed. Cavity of the gills without superior opening; 6 branchiostegals; pseudo-branchiae present. Air bladder none. Three species known, the type, *Kathetostoma læve*, being from Australia. (*κάθετος*, vertical; *στόμα*, mouth.)

*a.* Dorsal rays 13; anal 13; body shaded and dotted with blackish. AVERRUNCUS, 2647.

*aa.* Dorsal rays 10; anal 12; body spotted with white. ALBIGUTTA, 2648.

## 2647. KATHETOSTOMA AVERRUNCUS, Jordan &amp; Bollman.

Head  $2\frac{2}{3}$ ,  $3\frac{1}{2}$  with caudal; depth  $3\frac{1}{4}$ . D. 13; A. 13. Body short and robust, its width behind base of pectorals equal to length of top of head. Head very large, its width at opercle less than its length by  $\frac{1}{2}$  length of eye. Mouth large, vertical; maxillary 2 in head. Snout  $1\frac{2}{3}$  in eye. Eye rather small, 5 in head. Teeth of lower jaw largest, inner row of each jaw enlarged and movable; vomer and palatines with a few large, conical teeth. Lower jaw without tentacle. Interorbital space lightly concave,  $1\frac{1}{2}$  times length of eye. Premaxillary groove as broad as long,  $1\frac{1}{2}$  in eye, obtuse behind, extending backward just past middle of pupil. Distance between bases of humeral spines  $1\frac{1}{2}$  in top of head. Preorbital with 3 spines in front directed forward and downward. Preopercle with 3 spines below angle directed downward and forward. Two antrorse spines on mandible, and 2 on breast before ventrals. Bones of top of head coarsely granular, striate, no naked area above except premaxillary groove; 2 points on occipital region whence granular ridges radiate; opercles and orbital bones coarsely granular, but not striate. No trace of scales or of spinous dorsal. Base of dorsal equal to base of anal,  $1\frac{2}{3}$  in head, longest ray equal to depth of cheek; pectorals  $\frac{1}{2}$  eye, length greater than that of top of head; ventrals reaching more than halfway to vent, their length equal to that of top of head. A few small depressions resembling embedded scales on region before dorsal and above head. Color blackish brown, mottled with pale; lower parts pale, dusted with brown; lips and gular region black; dorsal dusty, with 5 indistinct, partly confluent, whitish spots along its base; anterior part of anal pale, posterior thickly dusted with blackish, tips of rays pale; pectorals blackish, faintly barred; axil dusted outside, inner part very pale; ventrals pale; caudal with 3 irregular oblique dark bars; floor of mouth pinkish; tongue dusted with dark specks. Length  $4\frac{1}{2}$  inches. Pacific Ocean, off coast of Colombia; a single specimen dredged at a depth of 7 fathoms; a most singular fish. (*averruncus*, a deity which wards off; from the mailed head.)

*Kathetostoma averruncus*, JORDAN & BOLLMAN, Proc. U. S. Nat. Mus. 1889, 163, off coast of Colombia, at Albatross Station 2800,  $8^{\circ} 57' N.$ ,  $79^{\circ} 31' 30'' W.$ ; KIRSCH, l. c., 259, 1889; JORDAN, Proc. Cal. Ac. Sci. 1896, 229, pl. 31.

## 2648. KATHETOSTOMA ALBIQUTTA, Bean.

Head 3; greatest width 3; depth  $3\frac{1}{2}$ . D. 10; A. 12; interorbital space 4 in head, containing a deep groove, the length of which is slightly greater than its width and nearly equaling length of eye. Mouth nearly vertical when closed; intermaxillary slightly protractile, the length of its tooth-bearing surface  $\frac{2}{3}$  length of head; maxillary very broadly expanded behind, its greatest width about 3 in length, extending almost to vertical from middle of eye; end of mandible not much farther back; length of mandible  $4\frac{3}{4}$  in length; mandible having 2 blunt prominences at its posterior end; the exposed portion of the maxillary traversed by radiating striae. The lower limb of preoperculum with 3 stout spines along its lower border; length of humeral spine 3 in head; humerus very strongly rugose on its upper border; 3 short spines on the anterior edge of preorbital. Teeth in villiform bands in the intermaxillary and mandible, and on vomer; palatines in a very short band; a cavity between head of vomer and the processes of the intermaxillary ending in a semi-circular canal behind, which is separated from the anterior cavity by a flap of skin. Gill openings very wide and only narrowly attached to the isthmus, leaving a free posterior border. Pseudobranchiae present, small; a small, narrow slit behind the last gill, its length about  $\frac{2}{3}$  that of eye; gill rakers tubercular, none on anterior arch. A pair of short but stout spines in front of ventrals. The origin of dorsal a little nearer to root of caudal than to tip of snout, midway between base of caudal and middle of eye; length of dorsal base about 3 in length, the third ray the longest, its length nearly  $\frac{1}{2}$  length of base of fin, the last ray about as long as eye, and the first scarcely longer than this. The anal origin directly under that of dorsal, the base of fin slightly longer than that of dorsal; the seventh, eighth, and ninth anal rays about the longest, their length equaling about  $\frac{2}{3}$  that of middle caudal rays; the first ray not much more than  $\frac{1}{2}$  as long as the longest and the rays gradually increasing in size to the ninth; length of pectoral  $3\frac{1}{2}$  in body; length of lowermost ray less than  $\frac{1}{2}$  length of head; only the first ray simple, the rest divided. Ventral origin under eye; the longest ray of ventral slightly shorter than mandible. Caudal slightly rounded when expanded, the middle rays as long as head without snout. The lateral line beginning near the root of humeral spine, curving upward slightly and running along back to end of dorsal, then curving downward to near the middle of the caudal base; skin naked. Color, upper parts light brown, the upper surface of the head minutely dotted with white; the back with numerous roundish spots and oblong blotches of whitish; lower parts pale; the dorsal with 2 or three dark blotches near the margin, in some cases not much larger than eye, in others fully twice as long; caudal with 9 black blotches, those on outer rays largest, differing in size in different specimens, these blotches distributed over the greater portion of the fin; anal pale, with the exception of a brownish blotch on the membrane of the last 3 rays; pectoral with a brownish submarginal band on its outer half, this band sometimes broken up on the mem-

brane  
88 fathCathetos  
tros  
NatThis  
reduct  
liaritie  
shaft oBody  
large, d  
large, t  
slit beh  
sides, t  
mostly G  
post-ten  
dorsal fi  
anal fin  
pectoral  
ceval, th  
(Carnivor  
ing river  
by mean  
species t  
glands.  
166-177.)a. Dorsal  
st

b. B

bb. 1

aa. Dorsal  
st  
sc  
c. S

cc. 1

3030

brane; ventrals pale. Length about 6½ inches. Gulf of Mexico, in 27 to 88 fathoms; 1 specimen known. (*albus*, white; *gutta*, spot.)

*Cathostoma albigutta*, BEAN, Proc. U. S. Nat. Mus. 1892, 121, Gulf of Mexico, at Albartross Station 2403, Lat. 28° 42' 30" N., Lon. 85° 29' 00" W. (Type, No. 39304, U. S. Nat. Mus.)

### Suborder HAPLODOCI.

This group is distinguished mainly by the undivided post-temporal, the reduction in the number of gill arches to 3, and by the absence of peculiarities shown by related forms. One family. (*ἀπι.όος*, simple; *δοκος*, a shaft or beam, from the form of the post-temporal.)

#### Family CXCVIII. BATRACHOIDIDÆ.

(THE TOAD-FISHES.)

Body more or less robust, depressed anteriorly, compressed behind; head large, depressed, its muciferous channels well developed; mouth very large, the teeth generally strong; premaxillaries protractile; gills 3, a slit behind the last; pseudobranchiæ none; gill openings restricted to the sides, the membranes broadly united to the isthmus; branchiostegals mostly 6; gill rakers present, moderate; suborbital without bony stay; post-temporal bone simple, undivided; scales small, cycloid, or wanting; dorsal fins 2, the first of 2 or 3 low, stout spines; soft dorsal very long; anal fin similar, but shorter; ventrals rather large, jugular, I, 2 or I, 3; pectorals very broad, the rays branched; pyloric caeca none; tail diphy-cercal, the caudal fin distinct, rounded; vertebrae in large number, 32 to 45. (Carnivorous coast fishes, mostly of the warm seas, some of them ascending rivers; the young of some or all the species fasten themselves to rocks by means of an adhesive ventral disk, which soon disappears. In some species the spines of the head and dorsal fin are provided with poison glands. Genera 7; species about 15. (*Batrachide*, Günther, Cat., III, 166-177.)

- a. Dorsal spines 3; opercle developed as 2 strong diverging spines; subopercle rather strong, with 2 spines similar to those of opercle; no venom glands.
- b. Body scaly; branches of subopercular spine subequal and diverging; frontal region broad, flat, and slightly depressed, its median ridge rather prominent. BATRACHOIDES, 858.
- bb. Body scaleless; branches of subopercular spine parallel, the lower branch much the shorter; vertebrae 10 + 22; frontal region not depressed, its median ridge prominent. OPSANUS, 859.
- aa. Dorsal spines 2; opercle very small, its posterior part developed as a single strong spine; subopercle feebly developed, narrowed, and not ending in a spine; body scaleless.
- c. Spines solid, without venom glands; several lateral lines on sides of head and body, composed of pores and shining spots, some of these accompanied by cirri; canine teeth present; vertebrae 12 + 31; frontal region depressed, forming a triangular area below level of temporal region, its median ridge very low. PORICHTHYS, 860.
- cc. Spines of dorsal fin and operculum hollow and connected with venom glands; lateral line on sides of body single; no canine teeth.
- d. Dorsal and anal free from caudal. THALASSOPHRYNE, 861.
- dd. Dorsal and anal fully joined to caudal. DECTOR, 862.

## 858. BATRACHOIDES, Lacépède.

*Batrachoides*, LACÉPÈDE, Hist. Nat. Poiss., III, 306, 1798 ("tau," Lacépède\*=*surinamensis*).  
*Batrachus* (KLEIN), BLOCH & SCHNEIDER Syst. Ichth., 42, 1801 ("tau," *didactylus, surinamensis*, etc.; substitute for *Batrachoides*).

*Batrachius*, RAFINESQUE, Anal. Nat. 1815, 82 (substitute for *Batrachoides*).

Body robust, formed as in *Opsanus*. Dorsal spines 3; opercle developed as 2 strong diverging spines; subopercle strongly developed; branches of subopercular spine subequal and diverging; body covered with small ctenoid scales; frontal region broad, flat, and slightly depressed, its median ridge rather prominent. Mucous pores of sides not greatly developed. No poison glands. Shore fishes of warm regions. (*βάτραχος*, frog; *εἶδος*, resemblance.)

a. Teeth small, about 14 on the vomer; anterior teeth of lower jaw in a band; lateral teeth of palatine enlarged and canine-like; irregularly arranged.

SURINAMENSIS, 2649.

aa. Teeth larger, about 8 on vomer; anterior teeth of lower jaw in 2 rows; 3 teeth on middle of palatines enlarged and canine-like, the middle one the smallest.

PACIFICI, 2650.

## 2649. BATRACHOIDES SURINAMENSIS (Bloch &amp; Schneider).

(SAPO.)

Head  $3\frac{1}{2}$  in length of body; depth 6. D. III-29; A. 26. Teeth small, about 14 on vomer; anterior teeth on lower jaw in a band; lateral teeth on palatines enlarged and canine-like, irregularly arranged; pectoral without pores on its inner surface. Color grayish, darker on sides and head; base of soft dorsal pale, with a dark, irregular line above; upper part of fin lighter; caudal nearly black; anal fin light with some dark markings. Coasts of Guiana and Brazil; not rare on sandy shores; our specimen from Curaçao.

*Batrachoides tau*, LACÉPÈDE, Hist. Nat. Poiss., 306, pl. 12, fig. 1, 1798; not *Gadus tau*, LINNÆUS.

*Batrachus surinamensis*, BLOCH & SCHNEIDER, Syst. Ichth., 43, 1801, Surinam; from a specimen in the Museum of Vaillant in Paris; GÜNTHER, Cat., III, 173, 1861; MEEK & HALL, Proc. Ac. Nat. Sci. Phila. 1885, 61.

## 2650. BATRACHOIDES PACIFICI (Günther).

Head 3 in length; depth about 6. D. III-26; A. 22. Teeth rather large, about 8 on vomer; anterior teeth on lower jaw in 2 rows; lateral teeth on lower jaw gradually increasing to middle of jaw, behind which they become abruptly smaller and then gradually increase to end of jaw; 3 teeth on middle of palatines enlarged and canine-like, the middle one the smallest; pectoral with a row of pores on inner surface. Color olivaceous brown; some indistinct dark cross bands on body; dorsal with about 7 very irregular oblique dark bars, anal with about 5; pectorals and caudal

\* "Il est revêtu d'écaillés molles, petites, minces, rondes, brunes, bordées de blanc, et arrosées par une mucosité très abondante, comme celles de la lote et de la mustelle." (Lacépède.) Lacépède's specimen was therefore one of the scaly species, not an *Opsanus*. No species of the latter group seems to have been known to Lacépède or to Schneider.

dark, w  
proceedi  
ined by

*Batrachu*  
435, 18  
*Batrachoi*  
Nat. S

*Opsanus*,  
*Batrachus*  
SCUNE

Body o  
jaws, vor  
mandible  
jaw small  
upper an  
the skin  
second th  
obscure,  
cirri on b  
mostly of  
feeding o  
(ωψ, eye

a. Nostri  
tle  
ex  
les

aa. Nostri  
st  
u  
de  
ne  
fo  
un

Head 2  
head broa  
teeth blur

\* The nam  
merely as a  
wrongly sup  
Schneider.  
any use of  
nesque had g  
for *Batrachu*  
† The Braz  
not been recd

dark, with few light cross bands. Panama; locally common, close to the preceding but with smaller teeth and fewer fin rays. The specimen examined by us collected by Dr. Gilbert.

*Batrachus pacifici*, GÜNTHER, Cat., III, 173, 1861, Panama; GÜNTHER, Fishes Centr. Amer., 435, 1860.

*Batrachoides pacifici*, GILL, Proc. Ac. Nat. Sci. Phila. 1863, 170; MEEK & HALL, Proc. Ac. Nat. Sci. Phila. 1885, 62.

859. OPSANUS,\* Rafinesque.

(TOAD-FISHES.)

*Opsanus*, RAFINESQUE, Amer. Monthly Mag. 1817, 203 (*cerapatus*).

*Batrachus*, JORDAN & GILBERT, Synopsis, 751, 1883, and of authors; not of BLOCH & SCHNEIDER.

Body comparatively short and robust, scaleless; head large, depressed; jaws, vomer, and palatines each with a single series of strong blunt teeth; mandible with an additional external series at symphysis; teeth of upper jaw small; dentary bones forming an acute angle at symphysis; lips fleshy; upper angle of opercle with 2 diverging spines, more or less concealed in the skin; no poison glands; spinous dorsal of 3 stout, short spines, the second the longest; axil of pectoral with a large foramen; † lateral line obscure, its pores not conspicuous; young with a series of small, tufted cirri on back and sides; branchiostegals 6; vertebrae 12 + 22. Shore fishes, mostly of temperate regions; voracious creatures, living on the bottoms, feeding on mollusks and crustacea, and having great strength of jaw. (αψ, eye; ἄνω, upward; "the name means looking up." Rafinesque.)

a. Nostrils with fleshy tentacle between them. Color brownish or dusky greenish, mottled with darker and lighter, the dark on sides of body in large irregular blotches extending from base of dorsal to about  $\frac{3}{4}$  distance to base of anal, and more or less covered with small pale spots; belly and chin plain white or yellowish.

TAU, 2651.

aa. Nostrils without fleshy tentacle. Color whitish or gray, everywhere blotched or spotted with brownish yellow and black, the black spots on top of head smaller and more numerous than on rest of body; a large black blotch at base of spinous dorsal, running up on fin; 3 black blotches along base of soft dorsal, which do not extend  $\frac{3}{4}$  the distance to base of anal; pectoral with black spots which do not form cross bands; ventrals with more dark markings than in tau; dorsal, anal, and caudal marked nearly as in tau.

PARDUS, 2652.

2651. OPSANUS TAU (Linnaeus).

(TOADFISH; SAPO; SLIMER; OYSTER-FISH.)

Head  $2\frac{3}{4}$ ; depth  $4\frac{1}{4}$ . D. III-26 to 28; A. 24. Body robust, naked, the head broad; mouth large, the very strong jaws closing with great force; teeth blunt, those on mandible small anteriorly, regularly increasing in

\* The name *Batrachus* should not be used for this genus, as it was originally given merely as a substitute for *Batrachoides*, having properly the same type, *surinamensis*, wrongly supposed to be *tau* of Linnaeus, a species unknown to Lacépède and Bloch & Schneider. No congener of *tau* was placed in *Batrachus* by Bloch & Schneider. Prior to any use of *Batrachus* as the generic name of the naked toadfishes, allied to *tau*, Rafinesque had given to one of the latter the generic name *Opsanus*, which can not be set aside for *Batrachus*, the latter being an unnecessary synonym of *Batrachoides*.

† The Brazilian genus, *Maregravia* (*cryptocentra*), in which this foramen is wanting, has not been recorded from north of the equator.

size backward, those on vomer prominent; a broad flap above orbit; tip of maxillary and lower side of mandible with conspicuous cirri; a series of smaller cirri along margin of preopercle; subopercle ending in a long, sharp spine; orbit about equaling interorbital width or length of snout; pectoral with a large foramen in the axil. Dusky olive, with black markings confluent on the sides and forming irregular, indistinct bars; belly and under side of head lighter; sides often with many pale yellow or whitish spots; soft dorsal with 6 to 9 oblique light bands; anal with 5 to 9; caudal and pectoral fins with 5 to 7 light cross bands, these formed chiefly from light spots; ventrals with some dark markings. In specimens from shallow water or algae, the brown becomes nearly black and more extended, the belly and chin spotted with darker, and top of the head has no distinct markings. The deeper-water specimens are lighter in coloration than those from near the surface, and those from the coral reefs (var. *beta*, Goode & Bean) are paler than those from the green algae and sea wrack; otherwise no differences seem to exist. In young individuals the head is more narrow and rounded, and the lower branch of the subopercular spine proportionally larger than in the adult. Cape Cod to Cuba; very abundant among rocks and weeds close to the shore northward, in deeper water southward; the young clinging to rocks by a ventral sucking disk, which is soon lost. Length 15 inches. Not valued as food. (*tau*, T, the bones on the head when dried showing a T-shaped figure.)

*Gadus tau*, LINNÆUS, Syst. Nat., Ed. XII, 440, 1766, Carolina. (Coll. Dr. Garden.)

*Cottus glaber*, SCHÖPF, Schrift. Naturf. Freunde, VIII, 1788, 146, Long Island; D. 25; V. 3; A. 21; short cirri below mouth.

*Cottus chaetodon*, BLOCH & SCHNEIDER, Syst. Ichth., 62, 1801, New York; after SCHÖPF.

*Lophius bufo*, MITCHILL, Trans. Lit. and Phil. Soc. 1815, 463, New York.

*Opsanus cerapalus*, RAFINESQUE, Amer. Monthly Mag., Jan., 1817, 204, south coast of Long Island. (Coll. C. S. Rafinesque.)

*Batrachoides vernullas*, LE SUEUR, Mém. Mus., v, 1819, 157, pl. 17, coast of Rhode Island.

*Batrachoides variegatus*, LE SUEUR, Jour. Ac. Nat. Sci. Phila., III, 1823, 399 and 401, Egg Harbor, New Jersey.

*Batrachus celatus*, DE KAY, New York Fauna: Fishes, 170, pl. 50, f. 161, 1842, New York.

*Batrachus tau beta*, GOODE & BEAN, Proc. U. S. Nat. Mus. 1882, 236, Gulf of Mexico.

*Cottus glaber*, WALBAUM, Artedi Piscum, III, 392, 1792; after SCHÖPF.

*Batrachus tau*, CUVIER & VALENCIENNES, Hist. Nat. Poiss., XII, 478, 1837; DE KAY, N. Y.

Fauna: Fishes, 168, pl. 28, fig. 26, 1842; GÜNTHER, Cat., III, 167, 1861; JORDAN & GILBERT, Synopsis, 751, 1883; MEEK & HALL, Proc. Ac. Nat. Sci. Phila. 1885, 59.

*Batrachus variegatus*, CUVIER & VALENCIENNES, Hist. Nat. Poiss., XII, 484, 1837.

#### 2652. OPSANUS PARDUS (Goode & Bean).

(SAPO.)

Head to end of opercular spine 3; depth 4. D. III-26; A. 22; maxillary  $1\frac{1}{2}$  in head; pectoral  $2\frac{1}{2}$ ; ventral 2; highest dorsal ray  $2\frac{1}{2}$ ; highest anal ray  $3\frac{1}{2}$ ; caudal 2. Body short and robust, compressed posteriorly; head large, somewhat depressed, wider than the body; eyes placed high, not so wide as the slightly concave interorbital space; mouth large, the maxillary reaching far beyond the eye, the lower jaw slightly projecting; a double row of small blunt teeth in upper jaw, not running very far back at the sides; lower jaw with a single row of much larger pebble-like teeth running well back and biting against a single row of similar teeth on pala-

tines; teeth; many of maxillary spines, the lower very short, covering longer of caudal behind, ventral posterior spread. spots of the large round skin and it is rat

*Batrachus*  
Bank  
Phila.

*Porichthys*

Body projecting opercle ventral subopercle scales on sides these are frontal regional area shallowly deep American some of origin and the extra

NOTE.— phosphorescence has made a "Porichthys" over the surface although

tines; a few teeth in front of jaw which bite against the premaxillary teeth; vomer with 1 or 2 irregular rows of large blunt teeth; head with many fleshy tentacles, 1 over each eye, a row around lower jaw, 1 on end of maxillary, and a row around preopercle; opercle ending in 2 diverging spines, the lower shorter; subopercle ending in a spine, its tip equal with the lower opercular spine, these spines not piercing the skin; gill rakers very short, scarcely developed. Body and fins covered with a soft smooth skin, which is exceedingly loose nearly to the ends of fin rays, and entirely covering the dorsal spines. Dorsal spines very short but stout; soft dorsal longer and higher than anal, but in other ways similar, reaching past base of caudal rays; pectoral short, as wide as long, round and fan-shaped behind, reaching to vertical from base of fourth dorsal ray; origin of ventral far in front of pectorals, the fins reaching to the vertical from the posterior edge of spinous dorsal; caudal well rounded, fan-shaped when spread. Color very pale yellowish brown, thickly covered with round spots of dark brown, those on head smaller; belly with numerous spots, the largest as large as eye; back with many oblong blotches, besides small round spots; fins blotched and banded. Gulf of Mexico, in deep water. This form has a very different coloration from *O. tau* and the texture of its skin and flesh is also less firm, but the technical differences are slight and it is rather a deep-water variety than a species. (*pardus*, leopard.)

*Batrachus tau pardus*, GOODE & BEAN, Proc. U. S. Nat. Mus. 1879, 336, Pensacola Snapper Banks; JORDAN & GILBERT, Synopsis, 751, 1883; MEEK & HALL, Proc. Ac. Nat. Sci. Phila. 1885, 60.

### 86c. PORICHTHYS, Girard.

(MIDSHIPMEN.)

*Porichthys*, GIRARD, Proc. Ac. Nat. Sci. Phila. 1854, 141 (*notatus*).

Body rather elongate; head not very broad, depressed, the lower jaw projecting. Dorsal spines 2; pectoral broad, without foramen in axil; opercle very small, its posterior part developed as a strong, single spine; suboperculum feebly developed, narrowed and not ending in a spine; no scales on body; spines solid, without venom glands; several lateral lines on sides of head and body, composed of pores and shining spots, some of these accompanied by cirri; canine teeth present; vertebrae 12 + 31; frontal region depressed, forming a triangular area below level of temporal region, its median ridge very low. Branchiostegals 6; interorbital area short, wide, and with shallow grooves. Air bladder more or less deeply divided into 2 lateral parts. Pyloric appendages none. Species American; remarkable for the very great development of mucous pores, some of which simulate the photophores of *Myctophum*, but are different in origin and not at all luminous. (*πόρος*, pore; *ἰχθύς*, fish; in allusion to the extraordinary development of the mucous system.)

NOTE.—The following account of the distribution, structure, and development of the phosphorescent organs of *Porichthys* is furnished us by Prof. Charles Wilson Greene, who has made a careful study of these organs:

"*Porichthys* has numerous lines of conspicuous bright silvery spots distributed in rows over the surface of the body. These spots have been called phosphorescent organs, although no such function has yet been observed, the name arising out of a superficial

resemblance. These so-called phosphorescent organs are arranged in rows over the body, and are definite and characteristic and quite constant in location in different individuals. They are accompanied by rows of epidermal sense organs, the two having an intimate relation in distribution over the surface of the fish. In surface view the shining organs have a bright silvery appearance, are more or less round in outline, size from a mere dot to 0.8 mm. in diameter, and surrounded or bordered on one side by an increased amount of pigment. The end buds present a round, transparent, or pellucid, and usually slightly raised, point. Each end bud is bordered by a pair of papillae. There are about 20 well-defined lines as follows: The *lateral* row, from posterior upper border of pectoral straight alongside to upper third of base of caudal, 35 pairs with an end bud between each pair, upper series small or rudimentary, segmentally arranged and between myomeres. The *pleural* row, from middle of base of pectoral, curves backward and downward to a point above first anal ray then straight nearly to base of caudal, 43 to 62 organs. End buds below each organ to above middle of caudal, 31 organs. The *caudal* rows, end buds only, 2 longitudinal rows on upper and lower thirds of fin. The *anal* row, on either side base of anal fin from third anal ray to base of caudal. Phosphorescent organs in pairs, a pair for each anal ray, 1 end bud for each pair. The *gastric* row, from front around lower edge of pectoral and along side of belly to opposite anal papilla, 30 phosphorescent organs. The *gular* row, from isthmus along ventral side of ventral fin then outward to join gastric row, spur runs forward along external side of ventral fin, 27 organs. A parallel line of 50 end buds follows the gular row and posterior end of gastric. The *ventral* row with its fellow forms a parenthesis on the stomach from the side of the anus  $\frac{1}{2}$  the distance to the ventral fin, 34 organs; no end buds. The *branchiostegal* row, from the isthmus outward over branchiostegal membrane and between first and second rays, no end buds. The *mandibular* row of phosphorescent organs extends around inner edge of ridge formed by the dentary bones; the row of end buds along the outer rim of the same ridge. The *opercular* rows, upper and lower, extend backward and upward across opercle. The *scapular* row, from above opercular spine straight back above pectoral fin, the curves in toward the base of the dorsal fin opposite the third dorsal ray. The *dorsal* row, along base of dorsal fin to base of caudal. This row and the scapular row consist of well developed end buds and rudimentary phosphorescent organs. The *occipital* and *frontal* rows, along the occipital and frontal regions, short rows of small and poorly developed organs. The *nasal*, from the posterior nasal tube to base of anterior tube. The *suborbital* and *post-orbital*, from posterior nasal opening around under eye backward and downward to opercle. A *malar* row, from the suborbital down across the cheek. A *maxillary* across the posterior end of maxillary bone. The rows on the head consist of well-developed end buds with rudimentary and irregularly placed phosphorescent organs. The phosphorescent organs are embedded in the connective tissue dermis of the skin, and in section show a uniform general structure throughout the body. A typical organ from the anal or ventral rows consists of an outer spherical group of cells called a lens, resting in a deeper cup-like structure, the capsule, and this in turn in a cup of fibrillar connective tissue called the reflector. The lens consists of cells, polygonal in the center of the group and flattened or fusiform around the periphery. They have a large conspicuous nucleus and a dense, homogeneous, highly refracting cell body. The outlines of the cells are very distinct. In the cells of the capsule the nuclei stain readily, but the granular protoplasm with difficulty, and the cell boundaries are indistinct and usually obliterated. In some specimens connective tissue septa penetrate the capsule. Blood capillaries are always present. The reflector extends well up around the sides of the lens; it consists of fibrillar connective tissue which strongly reflects light. Much pigment is embedded in its meshes. No nerves have yet been traced to the organ. The developing phosphorescent organs do not appear in the embryo fish until it is 15 to 16 mm. long. Then a bud appears in the lower layer of the epidermis, which soon becomes constricted off as a spherical mass of cells lying in the subepidermal connective tissue. This mass later slightly elongates and gives rise by constriction to the lens and the capsule. The reflector is developed from the surrounding connective tissue, so also the pigment cells. Mature organs are not found until the fish reaches a length of over 20 mm. The end buds appear much earlier, 9 to 10 mm." (Charles Wilson Greene.)

a. Ab

b.

bb

Head  
gate, ta  
long an  
illary re  
series on  
of the ar  
ter stron  
than in  
inward;  
inward;  
ward an  
or 5); an  
much lar  
In *P. ma*  
enlarged  
much lar  
the upper  
axillary  
caudal n  
trals, 2 $\frac{1}{2}$ .  
and clou  
along mi  
long, the  
or less c  
crescent-  
blue-bla  
suborbital  
pores; or  
behind a  
cheek ste



a. Abdomen with 4 longitudinal series of pores, each of which is accompanied by a shining silvery body; 4 rows of shining spots on sides of body; a white blotch below eye, with a black crescent below it.

b. Teeth on palatines few (4 or 5), 1 to 3 of them developed as very strong canines, as large as canines on vomer; dorsal fin with distinct black blotches; back with dark saddles; third lateral line extending nearly to base of caudal.

POROSISSIMUS, 2653.

bb. Teeth on palatines numerous, none of them canine, and all much smaller than canines on vomer.

c. Third lateral line ceasing at second third of anal; cross bands on back and dorsal fin very faint or wanting; dorsal fin with a faint dark edge; sides of head and shoulder without distinct spots; body rather elongate.

NOTATUS, 2654.

cc. Third lateral line extending nearly to end of anal; cross bands on back and dorsal fin very distinct, appearing as roundish blotches, those on the dorsal fin along the margin; sides of head and humeral region much spotted with brown; body robust.

MARGARITATUS, 2655.

2653. PORICHTHYS POROSISSIMUS (Cuvier & Valenciennes).

(BAGRE SAPO.)

Head  $3\frac{1}{2}$  ( $4\frac{1}{2}$  in total); depth  $5\frac{1}{2}$  (6). D. II-37; A. 34. Body rather elongate, tapering and compressed behind. Head depressed,  $\frac{3}{4}$  as broad as long and  $\frac{1}{2}$  wider than deep; lower jaw considerably projecting, maxillary reaching to well behind eye, its length  $1\frac{1}{2}$  in head. Teeth in single series on jaws, vomer, and palatines, those of upper jaw very small, a few of the anterior and 2 or 3 of the lateral teeth somewhat enlarged, the latter strongly hooked forward; teeth in lower jaw strong, rather weaker than in *P. margaritatus*, those in the front of the jaw hooked strongly inward; the lateral teeth, which are larger, hooked backward and inward; 1 or 2 strong canines on each side of vomer, these curved backward and outward; teeth on palatines distant, few in number (usually 4 or 5); among these are 1 to 3 very strong canines (usually, but not always, much larger than canines on vomer), strongly curved forward and inward. In *P. margaritatus* and *P. notatus*, the palatine teeth are not especially enlarged, subequal and more numerous, the canines on the vomer being much larger than any of the other teeth. Gill openings extending from the upper edge of pectoral to just below lower edge. Pectoral without axillary foramen; height of soft dorsal about 3 in head; length of caudal nearly 2; height of anal  $3\frac{1}{2}$ ; length of pectorals  $1\frac{3}{4}$ ; of ventrals,  $2\frac{3}{4}$ . Color in life, light brown above, the top of head much darker and clouded with dark brown; a row of about 10 bar-like dark blotches along middle of side, each larger than eye, those anteriorly deeper than long, the others longer than deep; each of these blotches usually more or less confluent with a saddle-like dark blotch across the back; a crescent-shaped pale translucent area below the eye; below this a larger blue-black area, irregularly crescent-shaped, covering the preorbital and suborbital region, bounded below and behind by a row of shining mucous pores; on it are about 4 large pores, and above and behind it, close behind and below eye, is a large shining pore bordered with black; cheek steel bluish; sides of body silvery, becoming golden below; lower

part of head and belly bright golden; a dark stripe along base of dorsal; soft dorsal with 2 or 3 rows of small round dark olive spots, the upper row posteriorly becoming a dark edging to the fin; caudal, dull red, edged with dusky; anal very pale, edged with blackish; pectorals light orange, usually with some small dark spots above; ventrals orange, slightly darker anteriorly. Numerous series of pores on the body, those of the lateral line accompanied by shining golden bodies, as in other species of the genus. According to fishermen, these bodies are phosphorescent, shining at night; a statement which is probably true, although we have been unable to verify it; pores on sides of back not shining. Most of the pores, as in other species, accompanied by numerous small cirri or cilia; the arrangement of the lines of pores and shining bodies not very different from that found in *P. notatus*. It may be thus described in detail: A series of pores beginning at tip of snout, extending down around preorbital region, bounding the dark subocular blotch and joining almost at a right angle with a series of pores which extends downward from lower posterior corner of eye to angle of mouth. Another series diverges from the first in front of eye, passing close below eye, then upward above cheek, ending in a large pore behind preopercle. A curved series of pores extending backward along opercle, and another parallel with it along subopercle. Two obscure series from front of eye along top of head, becoming wide apart at the vertex, converging at the nape, then slightly diverging, converging in front of spinous dorsal, then again diverging to pass around the fin, each at last becoming straight at front of soft dorsal, extending close to its base to its last ray, there being about 2 pores to each ray. Just below this series, at front of soft dorsal on each side, begins a second series, with the pores wider apart and somewhat irregular, ceasing near the middle of the soft dorsal fin. The lateral line proper next begins above upper posterior angle of preopercle, whence a short branch passes directly upward. Opposite front of soft dorsal, the lateral line is interrupted for a distance a little more than diameter of eye. A short branch arises at this interruption and passes upward and backward at an angle from the end of the anterior part; thence the lateral line passes straight to base of caudal. The next series arises just behind axil of pectoral, then curves abruptly downward and backward, becoming straight opposite third ray of anal, thence proceeding to base of caudal, the pores small and close-set, anteriorly bead-like and shining, becoming dull toward the tail. Next comes a double series on each side of base of anal, the 2 series converging behind and finally coalescing. Another series begins at the middle of the base of the pectoral in front, curves downward, around the base of the fin, and, proceeding directly backward, ceases opposite vent. A series begins midway between gill opening and ventral and, extending straight backward, ceases opposite base of pectoral. Another begins, on each side, on lower side of head, directly below angle of mouth, the two diverging slightly between ventrals, then converging a little behind ventrals, then abruptly diverging, joining the series last mentioned, on each side, just in front of base of pectoral. A cross series of pores extends straight across belly, between

vent  
abrup  
in fro  
on ea  
obtus  
the m  
mand  
the br  
large  
Lengt  
on san  
rare a  
descri  
types  
other  
differe  
*P. por*  
eiro 32  
repres  
*mus* ha  
unequ  
grayish  
with b  
outer e  
with d

*Batrache*  
nam  
Delal  
*Porichthy*  
Texa  
*Porichthy*  
1883,  
*Porichthy*  
Ac. N

Head  
maxilla  
opercle  
lower ja  
is a pat  
lar to t  
small te  
canines  
lower li  
this br

vent and anal fin. At each end of this cross series a series of pores turns abruptly forward, the two meeting in an acute angle on the belly just in front of a vertical from base of pectorals. Finally, 3 parallel series on each side of lower parts of head meet in front, the two anterior in obtuse curves, the posterior in an acute angle. The anterior series along the mandible ends at the corner of the mouth. The next just behind the mandible ends just below the corner of the mouth. The next passes along the branchiostegal region, ending at the gill opening. Mandible with 2 large foramina. A series of dark-colored pores along each side of tongue. Length 8 inches. South Carolina to Texas, and southward to Argentina, on sandy shores; not very common, and found in rather deep water. Not rare about Galveston, but unknown to fishermen at Pensacola. Here described from the types of *P. plectrodon*, the North American form. The types of *P. porosissimus* examined by us in Paris agree in dentition and other respects. Except for the remote locality there is no suggestion of differences. We are informed by Dr. Vaillant that the type specimen of *P. porosissimus* from St. Catherine, has 33 anal rays, that from Rio Janeiro 32, and that the number 27, given by Valenciennes for this species, represents an error in counting. According to Valenciennes, *P. porosissimus* has D. II-36; A. 27; each palatine bone with a row of small, pointed, unequal teeth; row of pores above anal reaching base of caudal. Color grayish brown above, silvery white below; dorsal and anal whitish, edged with brown; pectoral with longitudinal lines; ventrals brownish on the outer edge; caudal whitish at base, the rest brownish; some specimens with dark cross bands. (*porosissimus*, most porous.)

*Batrachus porosissimus*, CUVIER & VALENCIENNES, Hist. Nat. Poiss., XII, 501, 1837, Surinam (Coll. Leschenault & Douméro), Cayenne (Coll. Poiteau), Rio Janeiro (Coll. Delalande), St. Catherine (Coll. Lesson & Garnot).

*Porichthys plectrodon*, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1882, 291, Galveston, Texas (Type, No. 30894. Coll. D. S. Jordan); JORDAN & EVERMANN, Synopsis, 958, 1883.

*Porichthys porosissimus*, GÜNTHER, Cat., III, 176, 1861; JORDAN, Proc. Ac. Nat. Sci. Phila. 1883, 291; BERG, An. Mus. Nac. Buenos Aires, 1895, 70.

*Porichthys porosissimus*, JORDAN & GILBERT, Synopsis, 751, 1883; MEEF & HALL, Proc. Ac. Nat. Sci. Phila. 1885, 57.

#### 2654. PORICHTHYS NOTATUS, Girard.

(SINGING FISH; MIDSHIPMAN; CABEZON; SAPO.)

Head  $3\frac{1}{2}$ ; depth 2. D. II-37; A. 33; V. I, 2; P. 18; eye 8 in head; maxillary 2; pectoral  $1\frac{1}{2}$ ; ventral  $2\frac{1}{2}$ ; caudal  $2\frac{1}{2}$ . Head narrowed forward; opercle developed as a strong spine; maxillary reaching beyond orbit; lower jaw with a single row of about 10 large, recurved teeth, behind which is a patch of small teeth; sides of jaw with a single series of canines similar to those in front, but larger; upper jaw with an irregular series of small teeth; palatines with a single series of conical teeth; 2 large curved canines on vomer; head with several rows of fringed pores; 1 row along lower line of opercle and subopercle; another along upper edge of cheek, this branching behind and below the orbit, 1 branch running forward

below the orbit and around the snout, the other vertically downward behind the maxillary; a series of fringes behind the lower lip; behind this a series of pores without fringes; a short straight series of pores on each side of vertex; a row of pores along the base of the dorsal fin, curving at front of dorsal, and terminating at upper angle of opercle; a row below this, not reaching base of pectoral; the third row not reaching base of caudal, but ceasing at second third of anal to about its twentieth ray, and is anteriorly strongly curved upward to base of pectoral; 2 concentric series on the abdomen, the outer extending forward between bases of ventrals. The so-called "shining pores" on the sides are not pores, but bright round pieces of shiny membrane, showing through a translucent skin; each of the spots has above it a pair of fringed flaps with a small pore between them; the rows of flaps along dorsal and anal similar, long and low, their last rays reaching base of caudal rays; pectoral broad, somewhat pointed behind, reaching to the vertical from the fifth anal ray; origin of ventrals in advance of pectorals, in distance equal to length of maxillary, their tips not reaching to pectoral base; caudal well rounded. Olive brown above, with coppery reflections, the belly brassy-yellow; sides with irregular broad vertical cross blotches, most distinct in the young; dorsal grayish, with oblique dark bars; vertical fins sometimes margined with black; pores of lateral line head-like, shining silvery; a white space below eye, with a black crescent below it; head yellowish brown, with no dark spots on opercle and shoulder; peritoneum black. Length 15 inches. Pacific coast; very abundant from Lower California to Puget Sound; living under stones, near the shore northward, in deeper water southward. It makes a peculiar humming noise with its air bladder, hence the name singing fish. (*notatus*, spotted; noted.)

*Porichthys notatus*, GIRARD, Proc. Ac. Nat. Sci. Phila. 1854, 141, San Francisco; GIRARD, Pac. R. R. Surv., X, Fishes, 134, 1858.

*Porichthys margaritatus*, MEEK & HALL, Proc. Ac. Nat. Sci. Phila. 1885, 56; not of RICHARDSON.

*Porichthys porosissimus*, JORDAN & GILBERT, Synopsis, 751, 1883 (not of CUVIER & VALENNES); GÜNTHER, Cat., III, 176, 1861 (in part).

#### 2655. PORICHTHYS MARGARITATUS (Richardson).

Head  $3\frac{1}{2}$  to  $3\frac{3}{8}$ ; depth  $4\frac{1}{2}$  to  $5\frac{1}{2}$ . D. II-37; A. 33. Similar to *Porichthys notatus*, differing chiefly in color. Top and sides of head and space above pectorals with numerous round dark brown spots and freckles, behind pectorals 6 to 8 vertical  $\frac{1}{2}$  cross bars; dorsal not margined with black, but with 8 to 10 black submarginal spots; anal, with the exception of a few posterior rays, pale; caudal black at base and tip; pectorals with a few dots at base and on upper rays; a roundish white blotch below eye, below this a jet-black crescent. Palatine teeth small, 1 or 2 slightly enlarged. Series of shining spots arranged as in *P. notatus*, except that the third series extends almost to end of anal, to about its thirtieth ray. Pacific coast of tropical America. This species was obtained by the *Albatross* in large numbers off the west coast of Colombia, at Station 2795 at

a de  
larg  
Por  
por  
Batr  
C  
Por  
C  
it  
  
Thala  
  
Bo  
soft  
its p  
devel  
Spine  
of bo  
some  
spine

\* vac  
allusio  
† In  
spines  
; The  
" In th  
fishes  
The op  
with a  
fang of  
it has  
closed,  
canal r  
situate  
ovate s  
preserv  
sistency  
in the c  
or fluid  
neighb  
free in  
tube.  
appears  
thru a  
which i  
their sl  
front o  
somewh  
diate n  
base; th  
which i  
these m  
secreted  
of the r  
fluid sec  
organs,  
opulon  
found i  
qualitie

a depth of 33 fathoms, and at Station 2802 at a depth of 16 fathoms. The largest specimens are about  $4\frac{1}{2}$  inches long. In dentition it agrees with *Porichthys notatus*, but in color and arrangement of spots it resembles *P. porosissimus*. (*margaritatus*, hearing pearls; *μάργαρος*.)

*Batrachus margaritatus*, RICHARDSON, Voyage Sulphur, Fishes, 67, 1845, Pacific coast of Central America; coloration and arrangement of lines identical with *porosissimus*.

*Porichthys nautopedium*,\* JORDAN & BOLLMAN, Proc. U. S. Nat. Mus. 1889, 171, Pacific Ocean, off coast of Colombia, Albatross Station, No. 2802, 8° 38' N., 78° 31' 30" W., in 16 fathoms. (Type, No. 41145, U. S. Nat. Mus. Coll. Albatross.)

### 861. THALASSOPHRYNE, Günther.

(POISON TOAD-FISHES.)

*Thalassophryne*, GÜNTHER, Cat. Fishes, III, 174, 1861 (*maculosa*).

Body rather elongate, compressed; head moderate. Dorsal spines 2;† soft dorsal and anal rather short, free from caudal; opercle very small, its posterior part developed as a single strong spine; subopercle feebly developed, narrowed and not ending in a spine; no scales on body. Spines hollow, and connected with venom glands. Lateral line on sides of body single; jaws without canine teeth. Species all South American, some of them ascending rivers; all of them noted for their venomous spines.‡ (*θαλάσσια*, the sea; *ὄρνυη*, toad.)

\* *ναυτοπαίδιον*, sailor-boy, from the common name "midshipman," a name given in allusion to the "buttons" on the belly of the fish.

† In *Thalassothia*, Berg, a South American genus, likewise with poison glands, 4 dorsal spines are present.

‡ The poison organs of *Thalassophryne reticulata* are thus described by Dr. Günther: "In this species I first observed and closely examined the poison organ with which the fishes of this genus are provided. Its structure is as follows: (1) The opercular part: The operculum is very narrow, vertically styliform, and very mobile; it is armed behind with a spine, 8 lines long in a specimen of 10½ inches, and of the same form as the venom fang of a snake; it is, however, somewhat less curved, being only slightly bent upward; it has a longish slit at the outer side of its extremity, which leads into a canal perfectly closed, and running along the whole length of its interior; a bristle introduced into the canal reappears through another opening at the base of the spine, entering into a sac situated on the opercle and along the basal half of the spine; the sac is of an oblong-ovate shape, and about double the size of an oat grain. Though the specimen had been preserved in spirits for about 9 months, it still contained a whitish substance of the consistency of thick cream, which on the slightest pressure freely flowed from the opening in the extremity of the spine. On the other hand, the sac could be easily filled with air or fluid from the foramen of the spine. No gland could be discovered in the immediate neighborhood of the sac; but on a more careful inspection I found a minute tube floating free in the sac, whilst on the left-hand side there is only a small opening instead of the tube. The attempts to introduce a bristle into this opening for any distance failed, as it appears to lead into the interior of the basal portion of the operculum, to which the sac firmly adheres at this spot. (2) The dorsal part is composed of the 2 dorsal spines, each of which is 10 lines long. The whole arrangement is the same as in the opercular spines; their slit is at the front side of the point; each has a separate sac, which occupies the front of the basal portion; the contents were the same as in the opercular sacs, but in somewhat greater quantity. A strong branch of the lateral line ascends to the immediate neighborhood of their base. Thus we have 4 poison spines, each with a sac at its base; the walls of the sacs are thin, composed of a fibrous membrane, the interior of which is coated over with mucous. There are no secretory glands embedded between these membranes, and these sacs are probably merely the reservoirs in which the fluid secreted accumulates. The absence of a secretory organ in the immediate neighborhood of the reservoirs (an organ the size of which would be in accordance with the quantity of fluid secreted), the diversity of the osseous spines which have been modified into poison organs, and the actual communication indicated by the foramen in the sac, lead me to the opinion that the organ of secretion is either that system of nucleiferous channels which is found in nearly the whole class of fishes, and the secretion of which has poisonous qualities in a few of them, or at least an independent portion of it. This description was

a. Dorsal and anal fins not joined to the caudal.

b. Dorsal and anal fins rather short; D. II-19; A. 18; pectoral fins short, their tips reaching to origin of anal. Color brown, marbled with darker; pectoral fins and sides of body with some round black spots; chin and ventrals brownish; belly white. MACULOSA, 2656.

bb. Dorsal and anal fins longer; D. II-24; A. 24; pectoral fins longer, their tips reaching to sixth anal ray. Color of head, body, and fins brown, with a network of yellowish lines; dorsal, anal, caudal, and pectoral fins with white margins. RETICULATA, 2657.

2656. THALASSOPHYRYNE MACULOSA, Günther.

D. II-19; A. 18; V. I, 2. The head is somewhat longer than broad, its length being contained  $3\frac{1}{2}$  in the total; it is moderately depressed. The snout is short, obtuse, with the cleft of the mouth ascending obliquely upward, and with the chin prominent. The maxillary extends to the vertical from the posterior margin of the orbit. The teeth are obtusely conical, standing in single series, except anteriorly in the lower jaw, where they form 2 series, and in the upper, where they are cardiform, in a narrow band. The eyes are directed upward and very small, their width being  $\frac{1}{2}$  of that of the bony bridge between the orbits. Gill covers with a single spine; it is long, slender, cylindrical, like one of the dorsal spines, and has the operculum for its base. Gill opening not very narrow; it extends from the upper

made from the first example; through the kindness of Captain Dow I received 2 other specimens, and in the hope of proving the connection of the poison bags with the lateral-line system, I asked Dr. Pettigrew, of the Royal College of Surgeons, a gentleman whose great skill has enriched that collection with a series of the most admirable anatomical preparations, to lend me his assistance in injecting the canals. The injection of the bags through the opening of the spine was easily accomplished; but we failed to drive the fluid beyond the bag, or to fill with it any other part of the system of muciferous canals. This, however, does not disprove the connection of the poison bags with that system, inasmuch as it became apparent that, if there be minute openings they are so contracted by the action of the spirit in which the specimens were preserved, as to be impassable to the fluid of injection. A great part of the lateral-line system consists of open canals; however, on some parts of the body, these canals are entirely covered by the skin; thus, for instance, the open lateral line ceases apparently in the suprascapular region, being continued in the parietal region. We could not discover any trace of an opening by which the open canal leads to below the skin; yet we could distinctly trace the existence of the continuation of the canal by a depressed line, so that it is quite evident that such openings do exist, although they may be passable only in fresh specimens. Thus, likewise, the existence of openings in the bags, as I believed to have found in the first specimen dissected, may be proved by examination of fresh examples. The sacs are without an external muscular layer, and situated immediately below the loose, thick skin which envelops their spines to their extremity; the ejection of the poison into a living animal, therefore, can only be effected by the pressure to which the sac is subjected the moment the spine enters another body. Nobody will suppose that a complicated apparatus like the one described can be intended for conveying an innocuous substance; and therefore I have not hesitated to designate it as poisonous; and, Captain Dow informs me in a letter lately received, 'the natives of Panama seemed quite familiar with the existence of the spines and of the emission from them of a poison which, when introduced into a wound, caused fever, an effect somewhat similar to that produced by the sting of a scorpion; but in no case was a wound caused by one of them known to result seriously. The slightest pressure of the finger at the base of the spine caused the poison to jet a foot or more from the opening of the spine.' The greatest importance must be attached to this fact, inasmuch as it assists us in our inquiries into the nature of the functions of the muciferous system, the idea of its being a secretory organ having lately been superseded by the notion that it serves merely as a stratum for the distribution of peripheral nerves. Also the objection that the Stingrays and many Silurid fishes are not poisonous, because they have no poison organ, can not be maintained, although the organs conveying their poison are neither so well adapted for this purpose nor in such a perfect connection with the secretory mucous system as in *Thalassophryne*. The poison organ serves merely as a weapon of defense. All the Batrachoids with obtuse teeth on the palate and in the lower jaw feed on Mollusca and Crustaceans.' (Günther.)

base of the pectoral obliquely downward and forward to the level of the inferior base of the pectoral. The 2 dorsal spines are slender, pungent, about  $\frac{1}{2}$  the length of the head. Dorsal and anal fins terminate immediately before the root of the caudal, the length of which is  $\frac{1}{2}$  the total; pectoral obliquely rounded, extending to the origin of the anal; ventral rather short, not quite  $\frac{1}{2}$  the length of the head, extending to the base of the pectoral. Skin perfectly smooth, with some very short tentacles at the lower jaw. Two short horizontal muciferous channels on the cheek and the lateral line are very distinct; they are not, as usually, composed of a series of distant pores, but the pores are confluent, forming 1 continuous groove of a white color. Other muciferous channels, as for instance along the base of the anal, are composed of separate indistinct pores. Color brown, marbled with darker; pectoral fins and sides of the body with some round black spots; chin and ventrals brownish; belly white. The general habit is that of a *Batrachus* [*Opsanus*]. One specimen, from Puerto Cabello, Caribbean Sea. (Günther.) (*maculosus*, spotted.)

*Thalassophryne maculosa*, GÜNTHER, Cat., III, 175, 1861, Puerto Cabello; GÜNTHER, Fishes of Centr. Amer., 436, pl. 68, fig. 1, 1869; MEEK & HALL, Proc. Ac. Nat. Sci. Phila. 1885, 54.

2657. THALASSOPHRYNE RETICULATA, Günther.

D. II-24; A. 24; V. I, 2; P. 16. The length of the head is  $\frac{2}{3}$  of the total length (without caudal). The teeth on the palate are in a single series, very short, obtuse, incisor-like. Pectoral very large, extending back to the sixth anal ray. Head, body, and fins brown, with a network of yellowish lines; vertical and pectoral fins with a white margin. In other respects this species agrees with *T. maculosa*. Length 13 inches. Panama; not rare. (*reticulatus*, netted.)

*Thalassophryne reticulata*, GÜNTHER, Proc. Zool. Soc. London 1864, 150, 155, Panama; GÜNTHER, Fish. Centr. Amer., 437, pl. 68, fig. 2, 1869; JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1882, 62; MEEK & HALL, Proc. Ac. Nat. Sci. Phila. 1885, 7.

862. DÆCTOR, Jordan & Evermann, new genus.

(POISON TOAD-FISHES.)

*Dæctor*, JORDAN & EVERMANN, new genus (*dowi*).

This genus differs from *Thalassophryne* in the more elongate body and the many-rayed soft dorsal and anal fins, the last rays of which are fully joined to the caudal. (*δαίκτηρ*, slayer; from *δαίζω*, to slay.)

2658. DÆCTOR DOWI (Jordan & Gilbert).

Head 4 in length ( $4\frac{1}{2}$  with caudal); depth  $5\frac{1}{2}$  ( $6\frac{2}{3}$ ). D. II-33; A. 30. Body comparatively elongate, compressed behind. Head low and rather narrow, its width  $1\frac{1}{2}$  in its length. Eye very small, the diameter not  $\frac{1}{2}$  the interorbital space, and about as long as snout, 8 in head. Interorbital width about  $5\frac{1}{2}$  in head. Opercular spine short, nearly 4 in head. Mouth oblique, the lower jaw much projecting. Maxillary 2 in head, extending

to beyond eye. Teeth small, those on the palatine largest; teeth of upper jaw smaller than those of the lower; anterior teeth of the lower jaw in about 2 series. Pectoral fins long,  $1\frac{1}{2}$  in head, reaching about to fifth anal ray; last rays of dorsal and anal fully joined to the caudal. Color olivaceous, with darker blotches; first dorsal black; under parts pale; posterior portion of anal edged with dark. Pacific coast of North America, from Punta Arenas to Panama; rare. (Named for Capt. John M. Dow, who obtained a fine specimen (now destroyed) from Panama.)

*Thalassophryne dowi*, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1887, 388, Punta Arenas (Type, No. 39085, U. S. Nat. Mus. Coll. Cornell University); JORDAN, Proc. Cal. Ac. Sci. 1896, 231, pl. 38.

### Suborder XENOPTERYGII.

#### (THE CLING-FISHES.)

Breast with a broad sucking disk, between the wide-set ventral fins, this formed from the skin of the breast, not from the ventral fins themselves. Ventral rays I, 4 or I, 5; no scales; no spinous dorsal; no suborbital ring; palatine arcade materially modified; no air bladder; vertebrae in increased numbers; gill arches reduced. A well-marked group of small fishes, constituting a single family. (*ξέρος*, strange; *πρέρυξ*, fin.)

### Family CXCIX. GOBIESOCIDÆ.

#### (CLING-FISHES.)

Body rather elongate, tadpole-shaped, broad and depressed in front, covered by smooth, naked skin; mouth moderate; upper jaw protractile; teeth usually rather strong, the anterior conical or incisor-like; posterior canines sometimes present; suborbital ring wanting; no bony stay from suborbital across cheek; opercle reduced to a spine-like projection concealed in the skin, behind the angle of the large preopercle, this spine sometimes obsolete; pseudobranchiae small or wanting, gills 3 or  $2\frac{1}{2}$ ; gill membranes broadly united, free or united to the isthmus; dorsal fin on the posterior part of the body, opposite to the anal and similar to it, both fins without spines; ventral fins wide apart, each with 1 concealed spine and 4 or 5 soft rays. Between and behind the ventrals is a large sucking disk, the ventrals usually forming part of it. This sucking disk, which is wholly different in structure from that of *Cyclopterus* and *Liparis*, is thus described by Dr. Günther: "The whole disk is exceedingly large, subcircular, longer than broad, its length being (often)  $\frac{1}{2}$  of the whole length of the fish. The central portion is formed merely by skin, which is separated from the pelvic or pubic bones by several layers of muscles. The peripheric portion is divided into an anterior and posterior part by a deep notch behind the ventrals. The anterior peripheric portion is formed by the ventral rays, the membrane between them and a broad fringe, which extends anteriorly from one ventral to the other. This fringe is a fold of the skin containing on one side the rudimentary ventral spine, but

no ear  
on the  
a free,  
coid is  
extend  
runs th  
contint  
posteri  
the sol  
many p  
the ven  
short;  
22=26  
usually  
to them  
but the  
or *Batra*  
*Gobiesoc*  
GOBIESOC  
a. Gill r  
n  
b. I

bb. I

*Caularchus*,

This gen  
the only s  
the dorsal  
size and i  
accord wit  
anus; from



no cartilage. The posterior peripheric portion is suspended on each side on the coracoid, the upper bone of which is exceedingly broad, becoming a free, movable plate behind the pectoral. The lower bone of the coracoid is of a triangular form, and supports a very broad fold of the skin, extending from one side to the other, and containing a cartilage which runs through the whole of that fold. Fine processes of the cartilago are continued into the soft striated margin, in which the disk terminates posteriorly. The face of the disk is coated with a thick epidermis, like the sole of the foot in higher animals. The epidermis is divided into many polygonal plates. There are no such plates between the roots of the ventral fins." (Günther, Cat., III, 495.) No air bladder; intestines short; pyloric caeca few or none; skeleton firm; vertebrae 13 or 14 + 13 to 22 = 26 to 36. Carnivorous fishes of small size, chiefly of the warm seas, usually living among loose stones between the tide marks and clinging to them firmly by means of the adhesive disk. Their relations are obscure, but they are probably descended from allies or ancestors of the *Cottidae* or *Batrachoididae*. Genera about 15; species 50. The principal genus is *Gobiesox*. (*Gobiesocidae*, Günther, Cat., III, 489-515.)

GOMESOCINÆ:

- a. Gill membranes free from the isthmus; gills 3; posterior part of sucking disk with no free anterior margin.
- b. Incisors of lower jaw with entire edges.
  - c. Vertebrae about 32; anal fin long, nearly as long as dorsal. CAULARCHUS, 863.
  - cc. Vertebrae about 26; anal fin short.
    - d. Dorsal fin very long, of about 17 rays, twice as long as the moderate anal, which has 8 or 9 rays; disk broad; upper teeth in several rows. BRYSSETERES, 864.
    - dd. Dorsal fin moderate or short, of 4 to 13 rays.
      - e. Disk more or less broad, its length  $2\frac{1}{2}$  to 3 in body; dorsal and anal not very short, their rays 6 or more; body tapering rapidly backward; opercular spine strong. GOBIESOX, 865.
      - ee. Disk very narrow, its width 4 to 5 in body; head short,  $3\frac{1}{2}$  to 4 in body; dorsal and anal very short and small; a patch of teeth in each jaw behind the large teeth; sucking disk small. RIMICOLA, 866.
  - db. Incisors of lower jaw tricuspid or serrate; dorsal and anal fins short; vertebrae about 28. ARBACIOSA, 867.

863. CAULARCHUS, Gill.

(*Caularchus*, GILL, Proc. Ac. Nat. Sci. Phila. 1862, 330 (*meandricus*)).

This genus differs from *Gobiesox* chiefly in the numerous vertebrae, 32 in the only species known. The incisors are entire, the anal fin similar to the dorsal, each having 12 or 13 rays. The single species reaches a large size and is found farther north than any other of the group, a fact in accord with the increased number of vertebrae. (*καυλός*, stem; *ἀρχός*, anus; from the many-rayed anal.)

## 2659. CAULARCHUS MEANDRICUS (Girard).

(SUCK-FISH.)

Head  $2\frac{1}{2}$ ; depth  $6\frac{1}{2}$ . D. 13; A. 12; V. I, 4; vertebrae  $13 + 19 = 32$ ; eye  $7\frac{1}{2}$  in head; distance from vent to caudal  $2\frac{1}{2}$  in length of body; sucking disk as broad as long,  $3\frac{1}{2}$  in length. Head broad, nearly circular when viewed from above; interorbital width 3 in head; mouth wide, its width more than  $\frac{1}{2}$  length of head; maxillary extending to below eye; outer teeth of upper jaw rather strong, close set, vertical, conical, or slightly compressed, a narrow band of small, conical teeth behind them; lower jaw with larger teeth, 6 or 8 of the anterior broad, incisor-like, with entire edges, placed nearly horizontally; lateral and posterior teeth small, as in upper jaw; nostrils ending in tubes; spine on opercle sharp, but not projecting through the skin; origin of dorsal fin a little in advance of vent, the fin much higher than the anal; vent midway between anal and posterior edge of disk; pectorals short and broad, not extending back past the margin of the ventral disk, the 3 lower rays forming part of disk; caudal rounded. Color light olive, everywhere reticulated with brownish orange; middle of upper lip black; a light bar between eyes and 1 across cheek; vertical fins dusky; caudal with 2 faint brownish bars near its base. Specimens from red algae are light pink, mottled with darker, the pale band between eyes very distinct. Length 6 inches. Pacific coast of United States, from Vancouver Island to Point Conception; everywhere very abundant in rock pools; the largest specimen of *Gobiesocidae*. (*meandricus*, meandering, in allusion to the reticulated streaks.)

*Lepidogaster reticulatus*, GIRARD, Proc. Ac. Nat. Sci. Phila. 1854, 155, San Luis Obispo, California; name preoccupied.

*Lepidogaster meandricus*, GIRARD, Pacific R. R. Surv., x, Fishes, 130, 1858, San Luis Obispo, California; substitute for *reticulatus*, preoccupied in *Lepidogaster*; GÜNTHER, Cat., III, 505, 1861.

*Gobiesox reticulatus*, JORDAN & GILBERT, Synopsis, 749, 1883.

## 864. BRYSSETÆRES, Jordan &amp; Evermann.

*Brysseteres*, JORDAN & EVERMANN, Proc. Cal. Ac. Sci. 1896, 230 (*pinniger*).

This genus differs from *Gobiesox* solely in the great development of the dorsal fin, which has 17 rays, the moderate anal having but 8 or 9; the vertebrae  $10 + 16$ , as usual in *Gobiesox*. One species known. (*βρύστος*, sea-urchin; *ἔταπος*, comrade, the species living in rock pools with the sea-urchins.)

## 2660. BRYSSETERES PINNIGER (Gilbert).

Head  $2\frac{3}{4}$  to  $2\frac{1}{2}$ ; width of body  $4\frac{1}{2}$ ; of head 3 in length. D. 16 or 17; A. 8 or 9. Interorbital width  $3\frac{1}{2}$  in head; eye  $\frac{1}{2}$  interorbital width; width of mouth  $1\frac{1}{2}$  to  $1\frac{1}{4}$  in head. Teeth in upper jaw conic, acute, in several series, the anterior row in front enlarged, unequal; in lower jaw the teeth mesially in 2 distinct series, those in middle of anterior row narrow, entire incisors, those laterally conic, canine-like. No evident oper-

cular  
Front  
base  
betwe  
Caud  
at ba  
surro  
and 1  
obliqu  
freque  
vertic  
out di  
with  
(Gilbe  
Refugi  
tinguis

*Gobiesox*  
form  
*Brysseteres*

*Gobiesox*,  
*Megaphal*  
*sox*, r  
*Sicyoptes*,  
cies, v  
*Tomiodon*  
*Sicyoptes*  
*Bryssophis*

Body  
with tou  
in front  
strong in  
a series  
teeth on  
the throa  
portion v  
sal rays  
Species n  
the shore  
being fev

BRYSSOPH  
a. Dorsa  
in

aa. Dors  
305

cular spine. Disk about as broad as long, its length about that of head. Front of dorsal varying in position, about midway between snout and base of caudal, its length about  $\frac{1}{2}$  that of body. Vent nearly equidistant between disk and front of anal, the base of the latter  $3\frac{1}{2}$  to  $3\frac{3}{4}$  in body. Caudal  $1\frac{1}{2}$  to  $1\frac{3}{4}$  in head. Pectorals about  $\frac{1}{4}$  of head, with a distinct fold at base. Color variable; anteriorly usually with reticulating dark lines surrounding yellowish spots; a narrow dark streak forward, 1 downward, and 1 backward from orbit; below dorsal fin about 6 dark bars running obliquely downward and backward, these sometimes in greater number, frequently more or less irregular and interconnected, often divided by vertical streaks or series of dots; body sometimes light in spirits, with out distinctive markings; vertical fins usually dusky, narrowly margined with white, sometimes lighter with dark margins. Length  $2\frac{1}{2}$  inches. (Gilbert.) Gulf of California, abundant; specimens known from Puerto Refugio (Angel Island), San Luis Gonzales Bay, and La Paz. Well distinguished by its long dorsal. (*pinniger*, fin-bearing.)

*Gobiesox pinniger*, GILBERT, Proc. U. S. Nat. Mus. 1890, 94, Puerto Refugio, Gulf of California. (Coll. Albatross.)

*Bryssotæres pinniger*, JORDAN, Proc. Cal. Ac. Sci. 1896, 230, pl. 34.

### 865. GOBIESOX, Lacépède.

(CLING-FISHES.)

*Gobiesox*, LACÉPÈDE, Hist. Nat. Poiss., II, 595, 1799 (*cephalus*).

*Megaphalus*, RAFINESQUE, Analyse de la Nature 1815, 86 (*cephalus*, substitute for *Gobiesox*, regarded as an objectionable compound).

*Sicyases*, MÜLLER & TROSCHEL, Archiv für Naturgesch. 1843, 298 (*sanguineus*; small species, with upper teeth uniserial).

*Tomiscodon*, BRISOUT DE BARNEVILLE, Rev. Zool., 144, 1846 (*chilensis* = *Sicyases*).

*Sicyogaster*, BRISOUT DE BARNEVILLE, Rev. Zool., 144, 1846 (*marmoratus* = *Gobiesox*).

*Bryssophilus*, JORDAN & EVERMANN, new subgenus (*papillifer*).

Body anteriorly very broad and depressed, posteriorly slender, covered with tough, smooth skin; opercle with a strong spine; head large, rounded in front; mouth terminal, crescent-shaped; lower jaw with a series of strong incisors in front, their edges rounded or truncate; upper jaw with a series of strong teeth, behind which are sometimes smaller teeth; no teeth on vomer or palatines; gills 3; gill membranes broadly united under the throat, not attached to the isthmus; sucking disk large, the posterior portion without anterior free margin. Dorsal and anal moderate, the dorsal rays 6 to 12, the anal rays 6 to 10. Vertebrae about 26, as far as known. Species numerous, all American; mostly tropical, clinging to rocks near the shore. (*Gobius*; *Esox*; the resemblances either to the goby or the pike being few or remote.)

*BRYSSOPHILUS* (*βρύσσοσ*, sea urchin; *φιλέω*, to love):

a. Dorsal fin comparatively long, of about 13 rays; anal rays 9; disk broad; upper teeth in several rows; lower incisors narrow; papillae below chin; color olivaceous.

PAPILLIFER, 2661.

aa. Dorsal fin moderate or short, its rays 6 to 11.

## GOBIESOX:

c. Upper teeth in more than 1 series (character not verified in a few species); head broad.

d. Coloration in life chiefly olivaceous, without red, sometimes banded with darker or paler.

e. Dorsal rays 12; anal rays 7.

OYRINUS, 2662.

ee. Dorsal rays 11; anal rays 6; fins black.

NIGRIPINNIS, 2663.

eee. Dorsal rays 9 or 10; anal rays 6.

CEPHALUS, 2664.

eeee. Dorsal rays 8; anal rays 6.

TUDES, 2665.

eeeee. Dorsal rays 11; anal rays 10.

STRUMOSUS, 2666.

eeeee. Dorsal rays 10; anal rays 8.

VIRGATULUS, 2667.

eeeeeee. Dorsal rays 9; anal rays 7.

f. Width of head  $3\frac{1}{2}$  in length; color plain brown.

ADUSTUS, 2668.

g. Width of head 5 in length; color blackish, with yellow vermiculations.

FUNEBSIS, 2669.

eeeeeee. Dorsal rays 7; anal rays 7; eyes variegated.

PCEILOPHTHALMUS, 2670.

dd. Coloration in life chiefly bright red, or else with red spots or bands, the color not fading in spirits.

g. Color red, with deep red spots. D. 6; A. 5.

RHODOSPILUS, 2671.

gg. Color uniform red, unspotted, the color not fading in spirits; dorsal rays 6 to 8; anal rays 6.

h. Lower jaw with short incisors on each side, followed by canines.

MACROPHTHALMUS, 2672.

hh. Lower jaw with 2 horizontal incisors on each side, the third horizontal tooth not incisor-like; no distinct canines.

CERASINUS, 2673.

## SICYASES (σικύα, a sucking cup made of a gourd):

ee. Upper teeth in a single series (character not verified on some species); dorsal and anal short.

i. Color chiefly red.

j. Body with cross bands of deep red; iris red; dorsal rays 6; anal rays 5; head broad, the eyes very large.

ERYTHROPS, 2674.

jj. Body with dark cross bands and with spots of clear blue; body rather slender. D. 6 or 7; A. 6.

RUBIGINOSUS, 2675.

jjj. Body plain, light red; form rather slender.

CARNEUS, 2676.

ii. Color olivaceous or brownish, not red.

k. Dorsal rays 9; anal rays 6.

l. Color olivaceous, without bands.

HERES, 2677.

ll. Color greenish, with 3 dark cross bands and many dots.

PUNCTULATUS, 2678.

kk. Dorsal rays 7; anal rays 7; body with dark cross bands.

FASCIATUS, 2679.

## Subgenus BRYSSOPHILUS, Jordan &amp; Evermann.

## 2661. GOBIESOX PAPILLIFER, Gilbert.

Head  $2\frac{3}{4}$ ; width of body  $3\frac{3}{4}$ ; width of head  $2\frac{1}{4}$ . D. 13; A. 9. Width of mouth  $1\frac{1}{2}$  in head; interorbital width 3; eye  $\frac{1}{4}$  interorbital width; teeth in upper jaw conic, acute, very small, in 2 or more series, 2 of them slightly enlarged, canine-like; teeth in lower jaw in 2 series, the outer anteriorly, narrow entire incisors, with rounded tips, becoming conical laterally; opercular spine sharp, evident, though not projecting through the integument; lips and lower side of head anteriorly with fleshy papillæ; disk about as broad as long, its length  $1\frac{1}{2}$  in head; distance from

front  
except  
in head  
fleshy  
head  
base,  
margin  
having  
inches  
to the

Gobiesox  
form

B. G.  
coraco  
jaw ca  
are as  
lar; he  
width  
or 3 tim  
gin of  
distanc  
blotch  
Indies;  
apparar  
undus,  
tadpole

Lepadog  
from  
Cotylis n  
Gobiesox  
Skin  
Gobiesox  
GUNT

D. 11  
rays as  
Light b  
fier); v  
doubtful  
black; 2

Cotylis ni  
Gobiesox  
Amer

front of dorsal to base of caudal  $1\frac{1}{2}$  in its distance from tip of snout; vent exceptional in position, immediately in front of anal fin; base of anal  $1\frac{1}{2}$  in head; caudal rather acute,  $1\frac{1}{2}$  in head; pectorals  $2\frac{1}{2}$  in head, a distinct fleshy fold at base. In spirits, uniform dark olivaceous, lower side of head and disk light; pectorals dusky; vertical fins with a black bar at base, then a white bar, followed by a wide, dusky area, and narrowly margined with white; caudal with all these marks except the black bar, having the posterior outlines curved, following margin of fin. Length  $1\frac{1}{2}$  inches. Magdalena Bay, Lower California. (Gilbert.) Possibly related to the genus *Caularchus*. (*papilla*; *fero*, I bear.)

*Gobiesox papillifer*, GILBERT, Proc. U. S. Nat. Mus. 1890, 96, Magdalena Bay, Lower California. (Coll. Albatross.)

Subgenus GOBIESOX.

2662. GOBIESOX GYRINUS, Jordan & Evermann, new species.

B. 6; D. 12; A. 7; V. I, 4; P. 20. A vertical fold of skin at base of pectoral; coracoid distinctly below level of upper margin of pectoral; teeth of upper jaw cardiform, lower jaw with very narrow but compressed incisors, which are as short as the other teeth. Lateral profile of head nearly semicircular; head much depressed, as long as broad, its length being  $\frac{2}{3}$  of the total; width of interorbital space somewhat less than  $\frac{1}{2}$  greatest width of head, or 3 times diameter of eye; cleft of mouth extending beyond anterior margin of eye; distance of origin of dorsal from caudal more than  $\frac{1}{2}$  of its distance from snout. Brownish, with scattered dark spots; a black blotch anteriorly on the dorsal fin. Length 3 inches. (Günther.) West Indies; not seen by us. A valid species, according to Dr. Günther, but apparently as yet without tenable specific name, as the original *Cyclopterus nudus*, Linnæus, must have been some other fish. (*gyrinus*; *yupivos*, a tadpole.)

*Lepadogaster nudus*, BLOCH & SCHNEIDER, Syst. Ichth., 2, 1801; in part, description taken from *Cyclopterus nudus*, Linnæus, except the count of fin rays. D. 12; A. 6.

*Cotylis nuda*, MÜLLER & TROSCHEL, Hor. Ichth., III, 18, pl. 3, f. 2,

*Gobiesox nudus*, GÜNTHER, Cat. Fish., III, 502, 1861, Island of Cordova. (Coll. G. U. Skinner.)

*Gobiesox gyrinus*, JORDAN & EVERMANN, Check-List Fishes, 491, 1896, Cordova; after GÜNTHER; name only.

2663. GOBIESOX NIGRIPINNIS (Peters).

D. 11; A. 6; P. 22. "Nostrils, mouth, teeth, opercular spine, and fin rays as in *Cotylis stannii* (*Gobiesox cephalus*), but the dorsal fin longer. Light brown above (minutely dotted with black, if viewed by a magnifier); vertical fins black." Puerto Cabello (Peters); not seen by us; a doubtful species, perhaps identical with *G. cephalus* or *G. nudus*. (*niger*, black; *pinna*, fin.)

*Cotylis nigripinnis*, PETERS, Berl. Monatsber. 1859, 412, Puerto Cabello.

*Gobiesox nigripinnis*, GÜNTHER, Cat., III, 502, 1861; after PETERS; GÜNTHER, Fish. Centr. Amer., 390, 1869.

2664. *GOBIESOX CEPHALUS*, Lacépède.

(TÉTARD; TESTAR.)

D. 9 or 10; A. 6; C. 12; P. 19 or 20. Head and anterior part of body very broad, much depressed; skin tough, naked, and smooth; head nearly as broad as long, with its profile semielliptical, the snout being very obtuse and rounded. The upper surface of the head is quite flat, gently sloping downward in a straight line from the nape to the snout. The greatest width of the interorbital space is  $\frac{1}{2}$  of that of the head, or 4 times the diameter of the eye. The cleft of the mouth is horizontal, curved, wide, extending to below the center of the eye; the lips are thick, the lower being divided into 5 portions by 4 vertical grooves, the central portion being the smallest, the lateral ones the largest and hanging downward. The upper jaw is slightly protractile, and there is a broad velum behind the teeth in each jaw. A band of short conical teeth in the upper jaw; a single series in the lower, the anterior ones being slightly compressed incisors, and small like the lateral teeth, which are conical. The eye is small, situated immediately below the upper profile of the head. Two nostrils, close together, opposite the upper angle of the orbit, their margins being slightly raised. The lower angle of the opercular apparatus terminates posteriorly in an obtuse movable point enveloped in skin and directed backward. The gill openings are somewhat narrow in consequence of the small degree of expansibility of the gill covers, but the gill membranes have the margin quite free, being united together under the throat, and not attached to the isthmus. There are only 3 gills; the pseudobranchiae are quite rudimentary, indicated by 2 or 3 short lamellae. The distance of the origin of the dorsal fin from the caudal is nearly  $\frac{1}{2}$  of its distance from the snout, its first ray is much shorter than the others, and apparently without articulations. The caudal rounded and of moderate length; the anal is only  $\frac{1}{2}$  as long as the dorsal, commencing below its middle and terminating in the same vertical. The pectoral is broad and short, its lower  $\frac{1}{2}$  being longer than the upper; it is slightly connected with the ventral. A vertical fold of the skin at the base of the pectoral; the coracoid is so high as to reach to the upper margin of the pectoral. The adhesive apparatus as broad as long, its length being contained  $3\frac{1}{2}$  times in the total. The vent and the porus urogenitalis are close together, situated midway between the margin of the ventral disk and the anal. The anal papilla is small. The color is brown (in spirits), whitish inferiorly. Length of adult, 7 inches. (Günther.) Caribbean Sea, said to be common; not seen by us. The original *G. cephalus* seems nearer the next species, if the 2 are really different. If that be the case the present species may stand as *Gobiesox stannii*. But we have no material adequate to settle this question. (*cephalus*, big-headed; *κεφαλή*, head.)

*Gobiesox cephalus*, LACEPÈDE, Hist. Nat. Poiss., II, 505, 1798, Martinique; on a drawing by PLUMIER; D. 8; A. 4 or 5; color plain reddish; anal inserted behind dorsal; head broad; eyes blue; GÜNTHER, Cat., III, 499, 1861.

*Lepadogaster testar*, BLOCH & SCHNEIDER, Syst. Ichth., 445, 1801, Martinique; after PLUMIER.

*Cotylin stannii*, MÜLLER & TROSCHEL, Her. Ichthyol., III, 18, taf. 3, fig. 3, 1845.

Head  
descri  
Head  
large,  
entire  
in snout  
of head  
shorter  
out sp  
to be f  
Chines  
fers fr  
shorter

*Gobiesox*  
unkn

D. 11  
length;  
end of m  
ing ab  
on each  
which t  
11 on ea  
canine.  
Carolina  
ville by  
by its lo  
to the sw

*Gobiesox*  
lina;  
River.

Head  
10; A. 8  
rather b  
in head,  
convex.  
mouth ex  
than upp  
teeth of  
middle te  
entire or  
Distance

2665. *GOBIESOX TUDES*, Richardson.

Head  $2\frac{1}{2}$ ; depth  $4\frac{1}{2}$ ; width of head  $2\frac{1}{2}$ . D. 8; A. 6 in plate (5 in the description, the first short ray apparently not counted by Richardson). Head very broad, as broad as long, abruptly truncated anteriorly; mouth large, the maxillary reaching front of eye; lower jaw included; teeth entire; eye large,  $4\frac{1}{2}$  in head, a little more than  $\frac{1}{2}$  interorbital width,  $1\frac{1}{2}$  in snout. Distance from front of dorsal to caudal about equal to length of head; insertion of dorsal before vent; the anal behind dorsal and much shorter than it; pectorals short. Color uniform, probably greenish, without spots or stripes. Length 5 inches. Locality "unknown, but supposed to be from China." (Richardson.) The species is, however, certainly not Chinese and is more likely to be from the West Indies. This species differs from *Gobiesox cephalus*, as described by Günther, in the larger eye and shorter dorsal. It is probably the same species. (*tudes*, hammer.)

*Gobiesox tudes*, RICHARDSON, Voy. Sulphur, Fish., 103, pl. 46, figs. 1-3, 1845, habitat unknown, erroneously supposed to be China.

2666. *GOBIESOX STRUMOSUS*, Cope.

D. 11; A. 10; C. 16; P. 21. Head extremely wide, its width  $2\frac{3}{4}$  in total length; this width partly produced by a large fleshy mass extending from end of maxillary to end of interopercle; eye small; profile of head descending abruptly from posterior line of orbits. Superior dental series 12 on each side, externally, but the 3 median teeth conceal some series of which the second 3 external teeth are a continuation; inferior teeth 11 on each side; 4 median incisors horizontal and subequal; no marked canine. Bluish pigmented, fins blackish. (Cope.) Hilton Head, South Carolina, and Indian River, Florida; 4 specimens recently taken at Titusville by Evermann & Bean; apparently distinguished from *G. virgatulus* by its longer anal. (*strumosus*, from struma, a scrofulous tumor, alluding to the swollen cheek.)

*Gobiesox strumosus*, COPE, Proc. Ac. Nat. Sci. Phila. 1870, 121, Hilton Head, South Carolina; JORDAN & GILBERT, Synopsis, 749, 1883; EVERMANN & BEAN, Fishes of Indian River, Florida, in Rept. U. S. Fish Comm. 1896, 248.

2667. *GOBIESOX VIRGATULUS*, Jordan & Gilbert.

Head  $2\frac{1}{2}$  ( $3\frac{1}{2}$  with caudal); width of head  $3\frac{1}{2}$ ; depth 6 (7 in total). D. 10; A. 8 or 9; vertebrae  $10 + 16 = 26$ . Body rather slender, the head low and rather broad, broadly rounded anteriorly; eyes very small, about 4 to 6 in head, about  $2\frac{1}{2}$  in interorbital width; interorbital space broad, slightly convex. Cheeks prominent; opercle ending in a sharp spine. Cleft of mouth extending to below front of orbit; lower jaw somewhat shorter than upper. Teeth of upper jaw in a narrow band of about 2 series; 4 teeth of outer series a little larger than the rest, somewhat canine-like; middle teeth of lower jaw incisor-like and partly horizontal, their edges entire or somewhat concave. Ventral disk considerably shorter than head. Distance from root of caudal to front of dorsal  $2\frac{1}{2}$  in length. Pectorals

short, about  $2\frac{2}{3}$  in head. Color in life olivaceous, with numerous paler spots and broad diffuse dark bars; the whole body covered with rather faint, wavy, longitudinal stripes or lines of a light orange-brown color, about as wide as the interspace, much as in some species of *Liparis*, these entirely disappearing in alcohol; skin everywhere with dark punctulations; caudal dusky, slightly barred with paler, its tip abruptly yellowish; dorsal and anal dusky, the darker parts corresponding to dark bars on the body, barred. A rather large species. Length 2 to 4 inches. Common among ballast rocks, from Pensacola Bay north to Charleston. Our specimens from Pensacola and Charleston. (*virgatulus*, narrowly striped.)

*Gobiesox virgatulus*, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1882, 293, Pensacola, Florida (Coll. Jordan & Stearn.); JORDAN & GILBERT, Synopsis, 958, 1883; GOODE & BEAN, Proc. U. S. Nat. Mus. 1882, 236; JORDAN, Proc. U. S. Nat. Mus. 1884, 149.

2668. *GOBIESOX ADUSTUS*, Jordan & Gilbert.

Head 3; depth  $5\frac{1}{2}$ . D. 9; A. 7. Head and body broad and flat, much depressed; width of head nearly equal to its length,  $3\frac{2}{3}$  in body. Incisors in middle of lower jaw entire, broad; those in upper jaw narrow, blunt, little compressed, entire, shorter than the lateral teeth; behind these 2 or 3 series of smaller teeth. Eyes rather large, separated by a broad interorbital space, which is  $\frac{1}{3}$  length of head and about  $\frac{1}{2}$  greater than diameter of eye. Opercular spine sharp. Pectoral short, about  $\frac{1}{2}$  length of head; ventral disk as long as head; distance from base of caudal to front of dorsal equaling  $\frac{3}{10}$  of the length; caudal rounded behind. Brown, banded with blackish on body, head marbled with darker brown; front of dorsal black, the fins dusky with darker points. Pacific Coast of Mexico. Three specimens, the largest about 2 inches long, were obtained in a tide pool at Mazatlan. (*adustus*, scorched; brown.)

*Gobiesox adustus*, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1881, 360, Mazatlan, Mexico; JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1882, 627; JORDAN & GILBERT, Bull. U. S. Fish Comm. 1882, 108.

2669. *GOBIESOX FUNEBRIS*, Gilbert.

Body rather slender, its width 5 in length; width of head  $3\frac{1}{2}$  to nearly 4; head  $2\frac{3}{4}$  to 3 in length; depth  $\frac{1}{2}$  head. D. 9; A. 6 or 7. Teeth in upper jaw conical, in several series, unequal but without canines; in lower jaw mesially in 2 series, the outer of narrow, entire incisors, truncate or rounded, without lateral canines. Interorbital space wide, 3 in head, the eye small,  $\frac{2}{3}$  interorbital width. Mouth very wide,  $\frac{1}{2}$  or more than  $\frac{1}{2}$  length of head. Ventral disk wider than long, its length  $1\frac{1}{2}$  to  $1\frac{2}{3}$  in head. No evident opercular spine. Distance from front of dorsal to base of caudal  $2\frac{2}{3}$  to 3 in length anterior to dorsal; distance from vent to front of anal fin  $1\frac{1}{2}$  to  $1\frac{2}{3}$  in distance from vent to disk; base of dorsal from  $1\frac{1}{2}$  to  $1\frac{1}{4}$  in head; base of anal about  $\frac{1}{2}$  head; caudal rounded,  $1\frac{2}{3}$  to  $1\frac{1}{2}$  in head; pectorals  $3\frac{1}{4}$  in head. Color varying from dark olive brown to black, everywhere covered with fine, yellowish vermiculations, usually arranged to



form narrow lighter bars on the sides; 3 or 4 obscure dark streaks radiating from the eye; blackish below, the fins varying from blackish to straw color. Length  $2\frac{1}{2}$  inches. (Gilbert.) Gulf of California; abundant at Puerto Refugio (Angel Island) and La Paz. (*funebria*, funereal, from the dark color.)

*Gobiesox funebria*, GILBERT, Proc. U. S. Nat. Mus. 1890, 95, Puerto Refugio, Gulf of California. (Coll. Albatross.)

2670. *GOBIESOX PARCLOPHthalmus*, Jenyns.

Head 3, as wide as long. D. 7; A. 7. Opercular spine long and slender; teeth strong, somewhat crowded in front, the anterior in both jaws incisor-like; upper teeth conical, with smaller ones behind; 6 middle teeth of lower jaw incisor-like, projecting forward, their form not described (probably entire). Eyes large, close together, less than a diameter apart. General color olivaceous or brownish white, unmarked; iris golden, with pink and blue. Length  $1\frac{1}{2}$  inches. (Jenyns.) Chatham Island, Galapagos; only the single type known. (*ποικίλος*, variegated; *ὄφθαλμός*, eye.)

*Gobiesox parclphthalmus*, JENYNS, Voy. Beagle, Fishes, 141, pl. 27, figs. 2, 2a, 2b, 1842, Chatham Island (Coll. Darwin); GÜNTHER, Cat., III, 503, 1861.

2671. *GOBIESOX RHODOSPILUS*, Günther.

D. 6; A. 5; C. 8 or 9; P. 16. A vertical fold of skin along lower half of base of pectoral. Distance from front of dorsal to caudal  $2\frac{3}{8}$  in its distance from snout; anal before third dorsal ray. A very narrow band of short conical teeth in upper jaw, 1 lateral tooth larger than the others, recurved, canine-like; lower jaw with 1 series of teeth, the anterior narrow incisors, the outer distinctly canine, like the outer above. Rose-colored with rose-red transverse spots, each with an edge of deep-red dots. Panama. (Günther.) Not seen by us; known from 2 specimens, each  $1\frac{1}{2}$  inches long. (*ρόδον*, rose; *σπίλος*, spot.)

*Gobiesox rhodospilus*, GÜNTHER, Proc. Zool. Soc. Lond. 1864, 25, Panama (Coll. Captain Dow); GÜNTHER, Fish. Centr. Amer., 445, 1860.

2672. *GOBIESOX MACROPHthalmus*, Günther.

Eye  $4\frac{1}{2}$  in head. D. 8; A. 6; C. 12; P. 22. Head and anterior part of body very broad and much depressed, the head as broad as long, its profile semi-elliptical, the snout obtuse and rounded; top of head quite flat; interorbital width equal to eye. Mouth horizontal, curved, moderate, the cleft reaching beyond anterior margin of eye; an acute spine at lower angle of opercle; 4 short incisors on each side in lower jaw, separated from the conical lateral teeth by a larger canine-like tooth. Insertion of dorsal nearer caudal than snout; caudal rounded. Color, uniform reddish. (Günther.) Locality unknown; probably West Indies. (*μακρός*, large; *ὄφθαλμός*, eye.)

Dr. Eigenmann gives the following notes on a small specimen from St.

Thomas, which seems referable to *Gobiesox macrophthalmus*, differing in the slightly shorter dorsal and larger eye:

"Dorsal 6 or 7; anal about 7; head about 3; width of body  $3\frac{1}{2}$ ; width of mouth 2 in head; interorbital width  $4\frac{1}{2}$ , equal to snout. Eye large,  $2\frac{1}{2}$  in head. Teeth in the upper jaw conic, in more than 1 series in front, some in the outer row enlarged; teeth on the lower jaw in a single series, about 4 blunt incisors on each side followed by the canines, behind which the teeth are much smaller and conic. Width of disk  $1\frac{1}{2}$  in its length,  $2\frac{1}{2}$  in head; opercular spine strong; distance of origin of dorsal from caudal  $2\frac{1}{2}$  in its distance from tip of snout. Sides and back uniform bright red; eye black, iris bright red; lower surface yellow, dotted with bright red. One specimen 23 mm. long in the Museum of the University of Indiana from St. Thomas (Coll. Edward W. Brigham), much shrunken and fins hardened by strong alcohol." (Eigenmann, in lit.)

*Gobiesox macrophthalmus*, GÜNTHER, Cat., III, 502, 1861, locality unknown.

#### 2673. GOBIESOX CERASINUS, Cope.

Head 3 in total with caudal. D. 6; A. 6; C. 12; P. 24; V. 4. Head very wide, ovate, as broad as long to upper base of pectoral. Eye large,  $3\frac{1}{2}$  in head, equal to frontal width. Ten teeth on each side of each jaw, none of the upper being incisors, the 2 median on each side larger than the others; 3 teeth on each side in lower jaw horizontal, the others vertical, 2 of the horizontal teeth incisors, the median one on each side of these much the larger; each horizontal tooth with a small one behind it; no canines. Profile regularly descending from supraoccipital; a long subopercular spine. Dorsal beginning with last fourth of distance between tip of snout and base of caudal. Body and fins light crimson lake above, whitish below; no spots. One specimen,  $2\frac{1}{2}$  inches long, from St. Martins. (Cope.) (*κεράσινος*, *cerasinus*, cherry color.)

? *Cyclopterus nudus*, LINNÆUS, Syst. Nat., Ed. x, 260, 1758, "India;" from a specimen in Mus. Adolph Fred. (tab. 27), said to be 2 inches in length; the head broad with a sharp spine behind; dorsal rays 6; not *Lepadogaster nudus*, BLOCH & SCHNEIDER, Syst. Ichth., 2, 1801, who give "D. 12, A. 6,"\* the description otherwise that of Linnæus: not *Gobiesox nudus* of recent authors, which is a species (*G. gyrinus*) allied to *G. virgatulus*.

*Gobiesox cerasinus*, COPE, Trans. Am. Phil. Soc., XIV, 1871, 473, St. Martins, West Indies. (Coll. Dr. R. E. Van Rijgersma. Type in Ac. Sc. Phila.)

Subgenus SICYASES,† Müller & Troschel.

#### 2674. GOBIESOX ERYTHROPS, Jordan & Gilbert.

Head  $2\frac{1}{2}$ ; depth 6. D. 6; A. 5. Head scarcely longer than broad, proportionately very broad and depressed, its breadth 3 times in total.

\* We do not know by what authority the number of fin rays given by Linnæus (D. 6) was altered to "D. 12, A. 6" by Schneider (Syst. Ichth.). The last-named figures agree with *nudus*, as described by Dr. Günther, that is, with our *G. gyrinus*. If the Linnæan type of *nudus* really had D. 6, it must have been *cerasinus* or *macrophthalmus* or some very similar species. The scanty Linnæan description agrees best with *cerasinus*. The name *nudus*, if used at all, must be taken for a species to which the Linnæan description may be applied. In our judgment the uncertainty is too great to justify the substitution of *nudus* for either *cerasinus* or *macrophthalmus*. It could be no other known species, however.

† This subgenus is composed of small species with the upper teeth in 1 series. This character should be verified on all our species, as perhaps none of them belongs to it.

Incisors in both jaws, entire and rather broad, the lateral teeth, as usual, pointed; no canines. Eyes very large, considerably wider than the narrow interorbital area,  $3\frac{1}{2}$  in head; interorbital area nearly 5 in head. Ventral disk a little longer than head,  $2\frac{3}{8}$  in body. Pectoral about  $\frac{1}{2}$  length of head. Distance from front of dorsal to caudal,  $3\frac{1}{2}$  in body. Caudal truncate, with rounded edges. Light olivaceous; body with 3 or 4 bars of cherry red; head marbled with red; eyes intensely cherry red, their upper border blackish; fins pale, the upper mottled with reddish; caudal barred with red. Two specimens,  $1\frac{1}{2}$  inches long, taken in a rock pool at Mazatlan; also recorded from the Tres Marias Islands. (*έρυθρός*, red; *ὤψ*, eye.)

*Gobiesox erythroptus*, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1881, 320, Mazatlan, Mexico (Type, No. 29248. Coll. Gilbert); JORDAN & GILBERT, Bull. U. S. Fish Comm. 1882, 108; JORDAN, Fishes of Sinaloa, in Proc. Cal. Ac. Sci. 1895, 499.

2675. *GOBIESOX RUBIGINOSUS* (Poey).

D. 6 or 7; A. 6; V. 4; P. 25. Head 3, including caudal, its greatest width twice its height; eye 2 in interorbital width. Body slender; head semi-oval, obtuse; distance from front of dorsal to caudal  $1\frac{1}{2}$  in length, including caudal; dorsal opposite anal, beginning at fourth seventh of total length; mouth terminal; teeth not examined by Poey. Color red, with 12 dark bands and with many scattered spots of clear blue on the body; eyes with a red circle. Length 22 mm. Cuba (Poey); not seen by us; locally common at Matanzas; perhaps a species of *Arbacia*. (*rubiginosus*, reddish.)

*Sicyases rubiginosus*, POEY, Synopsis, 391, 1868, wharves of Palmasola, Matanzas, Cuba (Coll. Poey); POEY, Enumeratio, 124, 1875.

2676. *GOBIESOX CARNEUS* (Poey).

Head rounded; body very slender; eyes large, as wide as interorbital space; mouth inferior; teeth not examined by Poey. Color pale red, with some white specks and bands. Length 22 mm. Otherwise essentially as in *Gobiesox rubiginosus*. (Poey.) Matanzas; a doubtful species. (*carneus*, flesh-colored.)

*Sicyases carneus*, POEY, Synopsis, 392, 1868, wharves of Palmasola, Matanzas, Cuba (Coll. Poey); POEY, Enumeratio, 124, 1875.

2677. *GOBIESOX HERES*, Jordan & Bollman.

Head  $2\frac{3}{4}$  ( $3\frac{1}{4}$  in total); depth 6 ( $7\frac{1}{2}$ ). D. 9; A. 6. Body rather slender; head low and broad, greatest breadth not quite equal to length, its anterior margin not so broadly rounded as in *G. virgatulus*. Eyes very small,  $1\frac{1}{2}$  in interorbital space, 5 in head; interorbital bone appearing convex, least width  $3\frac{1}{4}$  in head and about equal to length of snout; cleft of mouth extending to beyond middle of eye; lower jaw included; teeth uniserial, those of upper jaw all canines, the first 3 on each side small, but becoming larger outward, next 3 or 4 much larger, rest smaller than those in front; anterior teeth of lower jaw entire incisors, which have on each side

about 6 large graduated canines and behind these a few smaller ones; teeth of lower jaw slightly oblique. Distance from front of dorsal to root of caudal about  $2\frac{5}{8}$  in body ( $3\frac{1}{2}$  in total). Pectorals moderate, 2 in head; ventral disk  $1\frac{1}{4}$  in head. Color olivaceous, without any distinct bands; the occipital region and the caudal peduncle darker; body irregularly mottled with groups of darker spots; nape, preopercle, cheeks, and snout with numerous dark points; indistinct dark lines radiating from eye; lips dark; fins dusky; dorsal and anal with the first rays black; a pale spot near base of caudal; axil of pectoral dusky. Green Turtle Cay, Bahamas; a single specimen known,  $2\frac{1}{4}$  inches in length. (*heres*, one who clings.)

*Gobiesox haeres*, JORDAN & BOLLMAN, Proc. U. S. Nat. Mus. 1888, 552, Green Turtle Cay, Bahamas. (Coll. Dr. Charles L. Edwards.)

2678. *GOBIESOX PUNCTULATUS* (Poey).

Head very broad, 3 in total length with caudal. D. 9; A. 6. Color brown, covered with black points; 3 dark transverse bands; none on the head. Teeth not described, the incisors probably entire. Length 38 mm. Cuba (Poey); not seen by us. (*punctulatus*, speckled.)

*Sicyases punctulatus*, POEY, Enumeratio, 124, 1875, Havana.

2679. *GOBIESOX FASCIATUS* (Peters).

D. 7; A. 7; head and body with alternate dark green and yellowish cross bands. Commencement of dorsal before that of anal, its distance from caudal equal to length of its base. Type, 50 mm. long. (Peters); not seen by us; teeth not described. Puerto Cabello. (*fasciatus*, banded.)

*Sicyases fasciatus*, PETERS, Monatsber. Berl. Acad. 1859, 412, Puerto Cabello; GÜNTHER, Cat., III, 497, 1861; GÜNTHER, Fishes Centr. Amer., 390, 1869.

866. *RIMICOLA*, Jordan & Evermann.

*Rimicola*, JORDAN & EVERMANN, Proc. Cal. Ac. Sci. 1896, 231 (*muscarum*).

This genus differs from *Gobiesox* mainly in the very slender body and head. Head  $3\frac{1}{4}$  to 4 in length, its width less than its length; dorsal and anal very short, of 4 to 6 rays each; incisors entire; a crescent-shaped patch of teeth in each jaw behind the large teeth; opercular spine weak or obsolete; sucking disk small. Species of small size; living below tide marks. (*rima*, a crevice; *colo*, I inhabit.)

a. Dorsal rays 6; anal 5; color yellowish, with a brown lateral band and numerous brownish spots. MUSCARUM, 2680.

aa. Dorsal rays 4; anal 5; color uniform light green. EIGENMANNI, 2681.

2680. *RIMICOLA MUSCARUM* (Meek & Pierson).

Head  $3\frac{1}{2}$  in length; depth  $8\frac{1}{4}$ ; D. 6; A. 5. Body elongate, slender, depressed anteriorly, but very narrow, slightly compressed posteriorly, the greatest width of body immediately behind head, 7 in length. Head narrow, much depressed, wider posteriorly. Eye small, its diameter  $2\frac{1}{2}$  in interorbital width, 5 in head. Maxillary reaching to the front of the eye,

its length less than 3 in head. Teeth in upper jaw conical, acute, curved, forming a crescent-shaped patch, those of the anterior row enlarged; in the lower jaw an anterior row of about 5 broad, entire incisors placed nearly horizontally; behind these a crescent-shaped patch of teeth, similar to those in the upper jaw, becoming canine-like laterally. No evident opercular spine. Ventral disk longer than broad, its length  $1\frac{1}{2}$  in head  $6\frac{1}{2}$  in length; distance from vent to front of anal  $2\frac{1}{2}$  in the distance from vent to disk; pectoral fin broad, short,  $2\frac{1}{2}$  in head; dorsal and anal fins small, the anal slightly in advance; caudal fin rounded. Ground color, in alcohol, light yellowish, paler below; above everywhere sparsely covered with distinct brownish-red spots about as large as pupil; a lateral band of the same color begins on the front of the snout, where it joins the one on the opposite side, extends through the eye across the opercle to the caudal, becoming very indistinct posteriorly; this lateral stripe is in strong contrast with the uniform pale ventral surface. Coast of California. Two specimens were dredged in Monterey Bay at a depth of about 10 fathoms. One of these, the type, is  $1\frac{1}{2}$  inches long. The second specimen ( $1\frac{1}{10}$  inches long) has the dorsal spots confined to the top of the head and nuchal region and the lateral stripe disappearing slightly behind middle of body, and having the ventral surface marked posteriorly with brownish-red spots like the spots on the dorsal surface. (*muscarum*, of the flies, from the fly-speck markings.)

*Gobiesox muscarum*, MEEK & PIERSON, Proc. Cal. Ac. Sci. 1895, 571, with colored plate, Monterey Bay. (Coll. S. E. Meek and Charles J. Pierson. Type in L. S. Jr. Univ. Mus.)

2681. RIMICOLA EIGENMANNI (Gilbert).

D. 4; A. 5. Head  $3\frac{2}{3}$  in length; depth about  $\frac{1}{2}$  head. Body very slender and narrow, the width of head  $4\frac{1}{3}$  in length; width of body 6. Mouth wide, the distance between its angles  $\frac{1}{2}$  length of head, the maxillary scarcely reaching vertical from front of orbit. Interorbital space wide, about  $\frac{1}{2}$  head. Eye very small, about 3 in interorbital width. Teeth in upper jaw conic, acute, in several series, the anterior in upper jaw enlarged; teeth in lower jaw also in several series, those of front row narrow incisors, entire, with rounded or truncate edges; disk very small and narrow, its width about  $\frac{2}{3}$  its length, the latter  $1\frac{2}{3}$  in length of head. Fins all small, the base of dorsal  $\frac{2}{3}$  length of head, less than free portion of caudal peduncle; distance from origin of dorsal to base of caudal  $3\frac{1}{4}$  in length before dorsal; distance from vent to front of anal fin  $1\frac{3}{8}$  in its distance from disk; caudal broadly rounded, its length  $1\frac{1}{2}$  in head; pectoral somewhat pointed, about  $\frac{1}{2}$  head; coracoid plate small, about  $\frac{1}{2}$  height of pectoral and less than  $\frac{1}{2}$  its length. Color uniform light olive green, without distinctive markings. Type, a single specimen, about 1 inch long, taken at Point Loma, near San Diego, California. Other specimens were taken some years since at San Cristobal Bay by Mr. Charles H. Townsend, and were referred to as *Gobiesox rhessodon* by Mrs. Eigenmann, Proc. U. S. Nat. Mus. 1884, page 553. (Named for Dr. Carl H. Eigenmann.)

*Gobiesox eigenmanni*, GILBERT, Proc. U. S. Nat. Mus. 1890, 96, Point Loma, near San Diego, California. (Coll. Gilbert.)

## 867. ARBACIOSA, Jordan &amp; Evermann.

*Arbaciosa*, JORDAN & EVERMANN, Proc. Cal. Ac. Sci. 1896, 290 (*humeralis*.)

This genus differs from *Gobiesox* chiefly in the character of the incisor teeth of the lower jaw; these are strongly serrate, or trienspid, making a ragged cutting edge. Size small; dorsal and anal comparatively short; head not very broad, the jaws contracted; vertebræ (in *Arbaciosa zebra*) 28. Some species provisionally referred to the section *Sicyases* of *Gobiesox* may prove to belong to *Arbaciosa*; small species, living in rock pools, among the sea urchins, by whose spines they are protected. This relation of *Arbaciosa zebra* with the Echinoid *Arbacia stellata* is especially constant. (*Arbacia*, a sea urchin.)

- a. Anal fin long, about 10 rays; dorsal rays 11; teeth above in 1 series; color olivaceous. RHESODON, 2682.
- aa. Anal fin of 5 to 7 rays.
- b. Dorsal fin of 8 or 9 rays; teeth in single series; color brownish, with red bars and a large black humeral spot. HUMERALIS, 2683.
- bb. Dorsal fin of 7 rays; color greenish, with pale spots and numerous pale cross bands; no red; body slender. RUPESTRIS, 2684.
- bbb. Dorsal fin of 6 rays; color chiefly red.
- c. General color pinkish olivaceous, with some bright red; back with 5 reddish-brown or blackish bars. Upper teeth in more than 1 series. Body comparatively slender, the depth nearly 8 in length. ZEBRA, 2685.
- cc. General color bright rosy red, black, with 1 to 3 faint dark bars. Upper teeth nearly uniserial. Body comparatively stout, the depth  $5\frac{1}{2}$  in length. EOS, 2686.

## 2682. ARBACIOSA RHESODON (Rosa Smith).

Head  $3\frac{1}{2}$ ; depth  $6\frac{1}{2}$ . D. 11; A. 10; eye  $4\frac{1}{2}$  in head,  $\frac{3}{8}$  in interorbital space; ventral disk  $1\frac{1}{2}$  in head; pectoral 2; caudal  $2\frac{1}{2}$ . Form much as in *Gobiesox mandricus*; snout bluntly and evenly decurved; the greatest height of the body across the pectoral fins; head broader than body but less deep; maxillary extending to below the eye; incisors of lower jaw not much declined, each of them trienspid, the central cusp longest; teeth of upper jaw conical, in an irregular series of 7 to 9; teeth in each jaw in single series; opercular spine sharp; distance from vent to caudal  $2\frac{3}{8}$  in length of body; dorsal a little longer than the anal, having its origin in advance of the anal and terminating opposite it; caudal rounded. Color dark olivaceous, usually with 3 broad yellowish cross bands above, the first across interorbital space and cheek, the second very wide, across back and front of dorsal fin, the third below middle of dorsal, some or all of these sometimes wanting; a dark bar at base of caudal; belly yellowish. Length  $2\frac{1}{2}$  inches. San Diego to the northern part of the Gulf of California; locally abundant in rock pools. (*ῥῥῥῥ*, to make ragged; *ὀδόν*, tooth.)

*Gobiesox rhesodon*, ROSA SMITH, Proc. U. S. Nat. Mus. 1881, 140, San Diego, California; JORDAN & GILBERT, Synopsis, 749, 1883; ROSA SMITH, Proc. U. S. Nat. Mus. 1883, 235.

2683. ARBACIOSA HUMERALIS (Gilbert).

Head 3 to  $3\frac{1}{2}$ ; width of body  $4\frac{1}{2}$ ; width of head  $3\frac{3}{4}$ ; eye very small, 3 in interorbital width. D. 8 or 9; A. 7. Body of moderate width, the head not evenly rounded anteriorly, becoming contracted opposite eyes, the snout forming a quadrate projection beyond the profile, as seen from above. Teeth in a single series in each jaw, the anterior narrow incisors, trilobate at tip, the 2 posterior teeth on each side strong, conical canines, somewhat recurved; about 12 incisors in the upper jaw. Interorbital space very wide, about equaling width of mouth,  $2\frac{1}{2}$  to  $2\frac{1}{2}$  in length of head. Ventral disk about as wide as long, its length  $1\frac{1}{2}$  in head. Opercular spine large and strong, but not exposed. Distance from front of dorsal to base of caudal 3 in length anterior to dorsal; base of dorsal  $1\frac{1}{2}$  in head; base of anal about equals base of dorsal; distance from vent to front of anal half its distance from disk; caudal broadly rounded,  $1\frac{1}{4}$  in head; pectoral  $\frac{1}{2}$  head, without distinct fold of skin across it. Ground-color dark olive brownish, crossed by many carmine-red bars, these somewhat broken anteriorly and above, to form reticulating lines, posteriorly and on lower part of sides more regular and running obliquely downward and backward; a conspicuous round humeral spot, larger than eye, in life black with golden-green reflections; numerous streaks from eye backward across cheek and opercles. (Gilbert.) Gulf of California; abundant at Puerto Refugio (Angel Island); also known from La Paz. (*humeralis*, pertaining to the shoulder, *humerus*.)

*Gobiesox humeralis*, GILBERT, Proc. U. S. Nat. Mus. 1890, 95, Puerto Refugio, Gulf of California. (Coll. *Albatross*.)

*Arbacirosa humeralis*, JORDAN, Proc. Cal. Ac. Sci. 1896, 230, pl. 35.

2684. ARBACIOSA RUPESTRIS (Poey).

Head 4 in total length with caudal; depth 6; eye 4 in head; snout less than eye. D. 7; A. 7. Forehead little decurved; eyes well separated; mouth small, with 1 row of compressed, close-set incisors with denticulated edges, 6 on each side in each jaw; snout truncate, as seen from above. Pectorals short, rounded. Dorsal and anal alike, opposed, highest in front. Caudal rounded. Color greenish ash, each side with 6 large oval spots, those behind touching; sides with about a dozen vertical bands of straw yellow or whitish, these bands sometimes interrupted, forming 2 series of points; 2 small similar bands from the eye, another toward tip of snout; a brown pale-edged band between eyes; some white spots on sides of head. Length  $1\frac{1}{2}$  inches. Coral reefs of Cuba; not rare. (Poey); not seen by us; said to be distinguished from other Cuban species by the slender body and narrow head. (*rupestris*, living among rocks.)

*Gobiesox rupestris*, POEY, Memorias, II, 283, 1861, Cuba.

*Sicyases rupestris*, POEY, Synopsis, 391, 1868; POEY, Enumeratio, 124, 1875.

2685. ARBACIOSA ZEBRA (Jordan & Gilbert).

Head  $3\frac{3}{4}$  in length; depth nearly 8. D. 6 or 7; A. 5 or 6; vertebrae 11 + 17 = 28. Body comparatively very long and narrow, the greatest width about  $\frac{1}{2}$  the total length. Head narrow, depressed, its width about  $4\frac{1}{2}$

times in length of body; eye small, its diameter about  $\frac{1}{2}$  interorbital width; opercular spine well developed; ventral disk nearly as long as head; mouth rather small, anterior, maxillary reaching front of eye; incisors of lower jaw nearly horizontal, rather broad, 3-lobed at tip, the middle cusp the longest; upper teeth much smaller, the median ones compressed, blunt, close set, a little shorter than the lateral teeth and with dentate edges, 1 or 2 series of small teeth close behind them; anal beginning under middle of dorsal; the distance from insertion of dorsal to base of caudal contained  $3\frac{1}{2}$  in length; pectoral  $\frac{1}{2}$  as long as head; caudal truncate, with rounded angles. Back with 5 dark cross bars about as wide as the interspaces, 3 of them in front of dorsal fin, the 2 anterior much broader and more distinct than the others; these bars all distinct on back, fading on sides, which are often vaguely clouded with dark; the color of these dark bars varies from reddish brown to black, and that of the interspaces from olivaceous to light pink and bright rose red; top of head bright red, marbled with light slaty bluish; a black blotch on opercle, and 2 very distinct black cross spots, 1 on each side of median line, forming the front of first dorsal bar; cheek sometimes with 2 or 3 pale bluish streaks; dorsal, pectoral, and caudal more or less shaded with dusky; lower fins pale; usually a dark bar at base of caudal and 1 across middle of fin; shade of ground color extremely variable.\* Very abundant in the rocky tide pools around Mazatlan, hiding everywhere under the numerous sea-urchins, especially *Arbacia stellata*, the protective coloration of both being that of the *Corallina*, which lines the rock pools. Length 2 to 3 inches. (*zebra*, from the banded coloration.)

*Gobiesox zebra*, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1881, 359, Mazatlan, Mexico (Type, No. 29250. Coll. Gilbert); JORDAN & GILBERT, Bull. U. S. Fish Comm. 1882, 108; JORDAN, Proc. Cal. Ac. Sci. 1895, 499.

\* The following note on the variations is furnished by Miss Susan B. Bristol:

"I find 4 specimens of this species which differ considerably from the typical form. These may represent a distinct species, but at present we are inclined to think that all these forms are modifications of one species, *Arbaciola zebra*. The following is a description of a specimen  $1\frac{1}{2}$  inches long, taken at Mazatlan (No. 4166 in the L. S. Jr. Univ. Museum): Head  $3\frac{1}{2}$ ; depth 9. D. 5 or 6; A. 6. Body slender, much depressed, compressed posteriorly, the greatest width  $4\frac{1}{2}$  in length. Head depressed, its width  $\frac{1}{2}$  in its length. Eye very small, about  $\frac{1}{2}$  in interorbital width. Snout rather rounded,  $3\frac{1}{2}$  in head. Opercular spine present. Interorbital width  $2\frac{1}{2}$  in head. Ventral disk  $\frac{1}{2}$  in head. Mouth small, the lower jaw inferior; outer teeth in both jaws serrate. Anal beginning at end of the first  $\frac{2}{3}$  of dorsal. Distance from front of dorsal to base of caudal  $1\frac{1}{2}$  in head. Caudal terminate. Pectoral  $2\frac{1}{2}$  in head. Color bright red, with very irregular yellow mottlings on back and sides, light yellow below; back with 4 irregular dark-red cross bars, the posterior 3 of which are wider than the interspaces; 3 of the cross bars in front of the dorsal fin, and the fourth on either side of the dorsal; 2 conspicuous black spots about  $\frac{1}{2}$  as large as eye, 1 on either side of median line on back above the pectorals a short distance behind their origin; snout plain, dark red; pupil white; 2 yellow parallel stripes extending from eye backward and downward, the second ending at a point about  $\frac{1}{2}$  the distance from tip of snout to end of opercle; dorsal, caudal, and anal dusky; ventrals and pectorals paler; a large red blotch at base of pectorals extending for a considerable distance on the fin. Another specimen from Mazatlan, bright red in color, about  $\frac{3}{4}$  of an inch long (also in bottle No. 4166, L. S. Jr. Univ. Museum), differs from the preceding form in the following respects: In the greater depth, which is  $6\frac{1}{2}$  in length, in the smaller ventral disk, which is  $1\frac{1}{2}$  in head; in the more pointed snout; in the absence of the 2 black spots above pectorals; and in having the 4 dark red bands on the back more distinctly marked. Two specimens from Guaymas, Mexico, 1 and  $1\frac{1}{2}$  inches long, No. 92 in the L. S. Jr. Univ. Museum, are chocolate brown in color, the shorter having on its back, including the bar at base of caudal, 7 dark brown cross bars and no dark spots above the pectoral, while in the longer there are no cross bars but a dark brown spot about  $\frac{1}{2}$  as large as the eye is present above the pectoral; also, in the longer one, the dorsal begins at the end of first third of anal. The eye in the larger specimens of *zebra* is larger than in these 4 specimens, but some of the smaller specimens seem to be intermediate in this regard between the typical form and these forms."



2686. ARBACIOSA EOS (Jordan & Gilbert).

Head 3; depth  $5\frac{1}{2}$ ; eye moderate,  $1\frac{1}{2}$  in interorbital width, which is about  $3\frac{1}{2}$  in head. D. 6; A. 6. Body comparatively short, stout, and narrow; the head rather broad, but, like the body, much less depressed than in *G. erythroptus*; width of head less than its length, or  $3\frac{2}{3}$  in body. Incisors serrate or tricuspid. Pectorals about 4 in head; ventral disk shorter than head. Distance from base of caudal to front of dorsal  $3\frac{1}{2}$  in total length; caudal truncate. Bright rosy red, sometimes dusky above with black points; back with 1 to 3 faint dark bars; 3 dark lines downward and backward from orbit, and usually 1 or 2 more on opercle; caudal usually with a reddish bar at base and a dusky one toward tip; fins otherwise nearly plain. Pacific coast of Mexico; abundant in rock pools about Mazatlan in company with *Arbaciosa zebra*, hiding under sea-urchins, especially with *Arbacia stellata*. Length  $1\frac{1}{2}$  inches. (*ηδς*, sunrise; from the red colors.)

*Gobiosox eos*, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1881, 360, Mazatlan, Mexico (Coll. Gilbert. Type, No. 29247); JORDAN & GILBERT, Bull. U. S. Fish Com. 1882, 108; JORDAN, Proc. Cal. Ac. Sci. 1895, 499.

Group BLENNIODEA.

(BLENNIROID FISHES.)

Body more or less elongate, naked or with scales, large or small; ventral fins small, more or less advanced in position, often wanting, the number of soft rays always less than 5; hypercoracoid perforate, the shoulder girdle normally formed; skull not armed with spines; suborbital not developed as a bony stay articulating with the preopercle; pseudobranchiae present; dorsal fin long, its anterior half, and sometimes the whole fin, composed of spines; anal long; tail homocercal, the caudal usually rounded, rarely forked; vertebrae numerous, especially in the arctic species. A large group, with ill-defined boundaries, the more primitive forms showing affinities with the *Trachinoidea*, *Cirrhitidae*, and other more typical fishes, the extremes very aberrant and passing directly into the *Ophidoidea*, and other forms lacking spines in the fins. We begin the series with the least modified of the type, the *Clinina*, from ancestors of which group the others have doubtless descended.

a. Caudal fin present, sometimes united to dorsal and anal; dorsal spines connected by membrane.

b. Gill openings not reduced to horizontal slits below the pectoral fins.

c. Teeth not developed as coarse molars.

d. Mouth not vertical.


BLENNIIDÆ, CC.



dd. Mouth nearly vertical; scales small or wanting; no lateral line; no ventral fins; dorsal composed entirely of slender spines; gill membranes attached to the isthmus; teeth strong.

CRYPTACANTHODIDÆ, CCI.



cc. Teeth developed as coarse molars on vomer, palatines, and sides of lower jaw; dorsal of flexible spines only; scales minute; gill membranes joined to the isthmus; no ventral fins; air bladder present; no lateral line.

ANARRICHADIDÆ, CCII.





**IMAGE EVALUATION  
TEST TARGET (MT-3)**



**Photographic  
Sciences  
Corporation**

23 WEST MAIN STREET  
WEBSTER, N.Y. 14580  
(716) 872-4503

1.5 1.8  
1.8 2.0  
2.0 2.2  
2.2 2.5  
2.5 2.8  
2.8 3.2  
3.2 3.6  
3.6 4.0

1.5 1.8  
1.8 2.0  
2.0 2.2  
2.2 2.5  
2.5 2.8  
2.8 3.2  
3.2 3.6  
3.6 4.0

- bb. Gill openings reduced to separate, narrow, nearly horizontal slits below and in front of the pectoral fin; ventrals small; dorsal fin long and low, anteriorly of slender spines; vertical fins connected. CERDALIDÆ, CCIII.
- aa. Caudal fin none, the tail tapering to a point; no ventral fins; no lateral line; scales rudimentary; anterior part of dorsal of low free-hooked spines, the posterior part of many slender soft rays; teeth in jaws only, close set in 1 row. PTILICHTHYIDÆ, CCIV.

## Family CC. BLENNIIDÆ.

## (THE BLENNIES.)

Body oblong or elongate, naked or covered with moderate or small scales which are ctenoid or cycloid; lateral line variously developed, often wanting, often duplicated; mouth large or small, the teeth various; gill membranes free from isthmus or more or less attached to it; pseudo-branchiæ present; ventrals jugular or subthoracic, of 1 spine and 1 to 3 soft rays, often wanting; dorsal fin of spines anteriorly, with or without soft rays; anal fin long, similar to soft dorsal; caudal well developed. Vertebrae in moderate or large number, 30 to 80. Carnivorous fishes of moderate or small size, mostly living near the shore in the tropical and temperate or arctic seas; most of them are carnivorous, the *Clininae*, so far as known, ovoviviparous, the rest mostly oviparous. Genera, about 80; species, about 400; chiefly of the rock pools and algae; some species in the lakes of Italy. Dr. Gill divides the tropical Blennies into 3 families, *Cliniida*, *Blenniida*, and *Chanopsida*. The first and second of these are fairly well defined. The third is now heterogeneous, and some of its members are intermediate between the other two. The arctic Blennies he again divides into *Xiphidiida*, *Cebedichthyidae*, and *Stichaida*, but the first and last of these groups intergrade, the *Xiphidiinae* are modified *Clininae*, and there are other forms as well entitled to separate rank as *Cebedichthys*. It seems to us better to treat the group as a single family with many subfamilies. (*Blenniida*, Günther, Cat., III, 206-297.)

I. Tropical Blennies, with the vertebrae mostly in moderate number, usually fewer than 45; lateral line usually arched high above the pectoral, if present; dorsal fin with soft rays, at least 1 being present; anal spines little developed; ventrals well developed, usually I, 3.

## a. Body scaly.

## CLININÆ:

- b. Lateral line present, arched anteriorly over the pectoral, becoming posteriorly median in position, or else obsolete; species ovoviviparous.
- c. Scales ctenoid, very rough, 35 to 40 in lateral line; dorsal divided into 3 fins; no cirri above eye. ENNEANECTES, 868.
- cc. Scales cycloid; dorsal fin not divided into 3 fins.
- d. Dorsal with 6 to 20 soft rays.
- e. Shoulder girdle with a small upturned hook on its inner edge.
- f. Scales along lateral line anteriorly not enlarged; snout sharp; first 5 spines of dorsal more or less modified.
- g. Caudal fin forked; air bladder present; scales minute; teeth in jaws in more than 1 series, on vomer and palatines; first 5 dorsal spines lengthened, and partly separated. HETEROSTICHUS, 869.

- gg.* Caudal fin truncate; air bladder wanting; scales minute; teeth in jaws in more than 1 series; teeth on vomer, none on palatines; first 5 dorsal spines lengthened and partly separated. GIBBONIA, 870.
- ee.* Shoulder girdle without upturned hook on its inner edge above.
- h.* Maxillary greatly developed, reaching much beyond eye; teeth on vomer and palatines; scales minute; soft dorsal long. NEOCLINUS, 871.
- hh.* Maxillary normal, not greatly expanded.
- i.* Anterior part of lateral line normally formed; usually a comb of filaments at the nape.
- j.* Palatines without teeth; scales moderate or small, 38 to 110 in lateral line.
- k.* Teeth in jaws in 1 row only; teeth usually on vomer, none on palatines; usually a comb of filaments at the nape. MALACOTENUS, 872.
- kk.* Teeth in jaws in more than 1 row, a band of villiform teeth behind the others; teeth on vomer, none on palatines.
- l.* Body oblong, the depth  $3\frac{1}{2}$  to  $4\frac{1}{2}$  in length; 2 filament above the eye. LABRISOMUS, 873.
- ll.* Body elongate, the depth about 6 in length; no filaments above the eye. MNIERPES, 874.
- jj.* Palatines with teeth, those in jaws in more than 1 series; scales large, 30 to 37 in lateral line; no nuchal filaments.
- m.* Head very broad, depressed; soft dorsal of about 20 rays. GOBIOCLINUS, 875.
- mm.* Head moderate, not depressed; soft dorsal of about 8 rays. STARKSIA, 876.
- ii.* Anterior part of lateral line running on a series of enlarged scales without visible pores; teeth in more than 1 series in jaws; teeth on vomer and front of palatines. CRYPTOTREMA, 877.
- dd.* Dorsal with 1 short soft ray only; scales large; teeth in jaws in more than 1 series; teeth on vomer, none on palatines.
- n.* Dorsal fin more or less deeply notched behind the third spine.
- o.* First 3 spines of dorsal very slender, close set, forming a separate ribbon-shaped fin, which is much higher than any of the spines in the second dorsal; anal spines rather high; body strongly compressed, the snout very sharp. EXERPES, 878.
- oo.* First 3 dorsal spines stiff, wide set, not remote from rest of fin behind dorsal notch; anal spines short; body more elongate, the snout less acute. AUCHENOPTERUS, 879.
- m.* Dorsal fin continuous, not notched. PARACLINUS, 880.

EMMNIONÆ:

- bb.* Lateral line straight, close to the dorsal fin; scales small, cycloid; dorsal notched, its anterior half of slender spines; no cirri on head; ventrals thickish, inserted slightly before pectorals; teeth in bands, the outer enlarged. EMMNION, 881.

aa. Body scaleless; species oviparous, so far as known.

p. Teeth comb-shaped, in a single row in each jaw, behind which are sometimes long canines; vomer and palatines usually toothless; lateral line usually single, with a strong arch anteriorly; dorsal fin long, continuous, or divided into 2 fins, the anterior portion composed of spines, which are stiff or flexible; anal fin long, usually with 1 or 2 small spines; ventrals well-developed, jugular, of 2 or 3 rays.

q. Teeth all fixed, attached to the bone of the jaws and not movable.

RUNULINÆ:

r. Caudal fin innate or forked; teeth compressed; spines and soft rays of dorsal indistinguishable.

s. Ventral fins very long, each of a spine and a soft ray.

ATOPOCLINUS, 882.

ss. Ventral fins not  $\frac{1}{2}$  length of head, each with about 2 soft rays; gill opening reduced to a small slit above pectoral.

RUNULA, 883.

BLENNIINÆ:

rr. Caudal fin rounded; teeth slender; gill membranes not reduced to a small slit.

t. Teeth all fixed, attached to the bone of the jaws.

v. Gill membranes free from the isthmus, or at least forming a distinct fold across it.

w. Jaws one or both with a posterior fang-like canine, much longer than the anterior teeth.

BLENNIUS, 884.

ww. Jaws without canines, the teeth all equal.

SCARTELLA, 885.

vv. Gill membranes broadly united to the isthmus, the gill openings restricted to the sides.

x. Jaws one or both with posterior fang-like canines.

HYPLEUROCHILUS, 886.

xx. Jaws without posterior canines; the teeth equal.

y. Three articulated ventral rays.

z. Mouth small, the maxillary extending scarcely beyond front of eye; the head decurved in profile.

HYPUBLENNIUS, 887.

zz. Mouth large, the maxillary extending beyond vertical from middle of eye; the head rather pointed in profile.

CHASMODES, 888.

yy. Four articulated ventral rays.

HONESTHES, 889.

SALARIINÆ:

qq. Teeth of front of jaws all movable, implanted on the skin of the lips.

a'. Vomer toothless.

b'. Jaws without posterior canines; dorsal fin deeply notched.

SCARTICHTHYS, 890.

bb'. Jaws one or both with posterior fang-like canines.

x. Dorsal fin continuous.

RUPISCARTES, 891.

xx. Dorsal fin divided.

ENTOMACRODUS, 892.

aa'. Vomer with a few teeth; posterior canines small.

SALARICHTHYS, 893.

pp. Teeth unequal, not comb-like; body oblong or elongate, more or less eel-shaped, naked, or rarely with rudimentary scales; supraocular flap sometimes present. Gill membranes united, free from the isthmus; dorsal fin very long, sometimes divided into 2 fins; formed of flexible spines, which often pass gradually into soft rays; anal fin long; ventral

II. Ble  
eral line  
cycloid, i  
j'. Gill  
k'.

fins thoracic or subjugular, usually, not much, if any, before the pectorals, composed of 2 soft rays each, the spine rudimentary; caudal well developed, the dorsal and anal usually more or less joined to it at base.

**OPHIUBLENNIINÆ:**

*e'*. Jaws each with 4 strong hooked canines in front; a hooked posterior canine below; a cirrus above eye and 1 above nostril; body scaleless; caudal fin forked; dorsal fin notched; body not eel-shaped; dorsal and anal free from caudal; ventrals small.

OPHIUBLENNIUS, 894.

*cc'*. Jaws with numerous teeth, not as above; caudal fin not forked.

**EMBLEMARIINÆ:**

*d'*. Body not eel-shaped; dorsal and anal not joined to caudal; no scales; no cirri; no lateral line; ventrals before pectorals; teeth on palatines; caudal fin rounded.

*e'*. Dorsal fin very high, not notched, the spines passing gradually into the soft rays; jaws long, sharp at tip.

EMBLEMARIA, 895.

*dd'*. Body elongate or eel-shaped; the dorsal and anal low, joined to base of caudal.

**CHELENOPSISINÆ:**

*f'*. Ventrals subjugular, more or less before pectorals; palatines with teeth; jaws long and sharp.

*g'*. Jaws with strong teeth, not as above described; dorsal fin with its anterior half of flexible spine, the posterior half of soft rays, the former gradually passing into the latter; jaws long, pike-like; ventrals inserted slightly before pectorals; anal with 2 spines; a villiform band of teeth in each jaw behind anterior teeth.

*h'*. Vomer toothless. Dorsal rays about XVIII, 38; anal II, 38. CHELENOPSIS, 896.

*hh'*. Vomer with a few teeth. Dorsal rays XVIII, 32; anal II, 30. LUCIOBLENNIUS, 897.

**PHOLIDICTHYINÆ:**

*f'*. Ventrals subthoracic, inserted below pectorals; teeth in jaws uniserial; anal fin without spines.

*i'*. Dorsal fin continuous, its spines indistinguishable from the soft rays. PHOLIDICTHYS, 898.

*ii'*. Dorsal divided into 2 fins, the anterior portion of 3 flexible spines behind the nape.

PSEDNOBLENNIUS, 899.

II. Blennies arctic or subarctic; the vertebrae in large number, usually 50 or more; lateral line various, usually median; dorsal fin usually without soft rays; scales small, cycloid, rarely wanting.

*j'*. Gill openings not continued forward below, the membranes broadly united, sometimes joined to the isthmus; ventral fins small or obsolete; scales small, cycloid.

*k'*. Pectoral fins short or wanting, never pointed, and never more than  $\frac{1}{4}$  head; pyloric caeca usually, but not always, obsolete.

*l'*. Body not covered with crosswise tubes at right angles to the lateral line.

*m'*. Dorsal fin composed of spines only.

**STATHMONOTINÆ:**

*n'*. Body scaleless; ventrals moderately developed; anal spines 2; no lateral line; no pseudobranchiae. STATMONOTUS, 900.

*nn'*. Body covered with small smooth scales.

**CHIROLOPHINÆ:**

*o'*. Ventral fins well developed, of 1 spine and 3 rays; no anal spines; top of head with many cirri; a row of large pores above base of pectorals; gill membranes free from isthmus; no pyloric caeca.



- p'*. Lateral line obsolete, only the row of pores being present. BRYOSTEMMA, 901.  
*oo'*. Ventral fins rudimentary or wanting, not more than 1 soft ray present; dorsal spines all short and rigid.

## PHOLIDINÆ:

*q'*. Lateral line obsolete.

*r'*. Gill membranes broadly united, free from the isthmus; no pyloric caeca; carnivorous.

*s'*. Anal fin with a large sheathed spine; ventrals wanting.

*t'*. Anal spine very long, pen-shaped, its anterior surface channelled; pectoral fins moderate.

APODICHTHYS, 902.

*u'*. Anal spine moderate or small, not pen-shaped, its anterior edge convex, not channelled.

*w'*. Pectoral fins very small; anal spine moderate.

XERERPES, 903.

*uw'*. Pectoral fins wholly wanting; anal spine small.

ULVICOLA, 904.

*as'*. Anal fin with 2 small spines or with none.

*v'*. Ventral fins reduced to a short spine, followed by a rudimentary ray.

*w'*. Caudal fin well developed.

PHOLIS, 905.

*ww'*. Caudal fin very narrow, the dorsal and anal united around the tapering tail.

GUNNELLOPS, 906.

*vv'*. Ventral fins entirely wanting; caudal as in *Pholis*.

ASTERNOPTERYX, 907.

*rr'*. Gill membranes joined to the isthmus, sometimes forming a fold across it; no ventral fins; no anal spines; top of head with fleshy crests; pyloric caeca present; body naked anteriorly, with small scales posteriorly. ANOPLARCHUS, 908.

## XIPHIDIINÆ:

*qq'*. Lateral lines several, each with many short cross branches; pyloric caeca present; gill membranes free from isthmus; ventrals none; anal spines 2 or 3, small; herbivorous.

*jj'*. Gill

LUC  
a

STIC  
aa

*Enneanect*

Body r  
line almo  
the eye e  
divided i  
10, the so  
toral lon

*w'*. Pectorals small but well developed, much longer than eye.

XIPHISTES, 909.

*ww'*. Pectorals minute, not longer than eye.

XIPHIDIUM, 910.

CEBEDICHTHYINÆ:

*mm'*. Dorsal fin with its posterior half composed of soft rays; gill membranes broadly united, free from isthmus; ventrals wanting; lateral line single, high; pyloric caeca present; herbivorous.

CEBEDICHTHYS, 911.

DICTYOSOMATINÆ:

*W*. Body covered with crosswise tubes at right angles with the lateral line and forming a network with it.

*x'*. Dorsal fin of spines only; teeth strong; ventral fins present, well developed; gill membranes broadly united, free from the isthmus.

PLAGIOGRAMMUS, 912.

*kk'*. Pectoral fins long and rounded or pointed, nearly as long as head; dorsal fin high; gill membranes broadly united, free from the isthmus; no lateral line; species probably all herbivorous.

OPISTHOCENTRINÆ:

*y'*. Ventral fins wanting.

*z'*. Dorsal with its posterior spines rigid and sharp; head scaly.

OPISTHOCENTRUS, 913.

*zz'*. Dorsal with its spines all flexible; head naked.

PHOLIDAPUS, 914.

PLECTOBANCHINÆ:

*yy'*. Ventral fins well developed; dorsal spines all pungent; body greatly elongate.

PLECTOBANCHUS, 915.

*jj'*. Gill openings continued forward below, the membranes separate or nearly so, scarcely joined to the isthmus; pectorals and ventrals well developed; dorsal spines slender, pungent, the fin without soft rays; herbivorous species.

LUMPENINÆ:

*a''*. Lateral line obsolete or obscure; body greatly elongate.

*b''*. Pectorals with the upper and middle rays shortened, shorter than lower; teeth on vomer and palatines.

LEPTOCLINUS, 916.

*bb''*. Pectorals with the middle rays longest.

*c''*. Lateral line not wholly obsolete, a series of distant pores along sides; teeth on vomer and palatines.

POROCLINUS, 917.

*cc''*. Lateral line obsolete, only a few small pores being traceable; no teeth on vomer; palatine teeth small or wanting.

LUMPENUS, 918.

STICHÆINÆ:

*aa''*. Lateral line present, single, double, or triple; body moderately elongate; teeth on jaws, vomer, and palatines.

*d''*. Lateral line simple, one on each side of back.

STICHÆUS, 919.

*dd''*. Lateral lines 2, or dividing into 2 on each side.

ULVARIUS, 920.

*ddd''*. Lateral line forking, forming 3 on each side.

EUMESOGRAMMUS, 921.

868. ENNEANECTES, Jordan & Evermann.

*Enneanectes*, JORDAN & EVERMANN, Proc. Cal. Ac. Sci. 1895, 501 (*carminatis*).

Body rather robust, covered with large, rough ctenoid scales; lateral line almost obsolete; mouth moderate, the jaws equal; no tentacle above the eye or on nape; no hook on shoulder girdle; eye large; dorsal fin divided into 3 fins, the first of 3 or 4 slender spines, the second of about 10, the soft dorsal of about 7 rays; caudal rounded; anal fin long; pectoral long, the lower rays simple and thickened. Small fish of the rock

pools, closely allied to the Old World genus, *Tripterygion*, Risso, but distinguished by the chubby body, short fins, and large, rough scales. (*ἐννέα*, nine; *ῥηκτηρ*, swimmer, there being 9 fins.)

## 2687. ENNEANECTES CARMINALIS (Jordan &amp; Gilbert).

Head  $3\frac{3}{8}$ ; depth  $4\frac{1}{2}$  to  $5\frac{1}{2}$ . D. III-XII, 9 (IV-X, 8 in the specimen before us); A. II, 11 (misprinted II, 17) scales 33 to 40. Body rather stout, heavy forward, rapidly tapering behind. Head short, the snout low and rather pointed, the profile straight and steep from the snout to opposite the front of the eyes, there forming an angle and extending backward nearly in a straight line; eyes very large, longer than snout, 3 in head, high up, and close together; mouth wide, the jaws subequal, the maxillary extending backward to front of pupil; teeth moderate, essentially as in species of *Labrisomus*, those of the outer series enlarged; no evident cirri on the head; scales on body of moderate size, ctenoid, the edges strongly pectinate; belly naked; lateral line extending to opposite last ray of soft dorsal, ascending anteriorly, but without convex curve; dorsals 3, the first and second contiguous, the second and third well separated; first dorsal of 3 spines, the first of which is the highest and about as long as diameter of eye; the second dorsal of higher and slenderer spines, the anterior the highest, the longest about equaling greatest depth of body; soft dorsal shorter and a little lower than second spinous dorsal; caudal small; anal long, beginning nearly under middle of spinous dorsal; pectoral long, longer than head, reaching much past front of anal; ventrals  $\frac{3}{4}$  length of head. Color light brownish, with 4 dark-brown cross bars on sides, about as wide as the interspaces, which are marked with more or less reddish and with some lighter spots; belly pale; space behind pectoral dark; a dark bar downward and 1 forward from eye; first dorsal mottled with darker, second and third dorsals nearly plain; a narrow, dark bar at base of caudal and a broader one toward the tip, the fin sometimes entirely black; pectorals somewhat barred; lower fins plain. Mazatlan, in tide pools; the types, 4 specimens, each about  $1\frac{1}{2}$  inches long. Another from the same locality, since figured by Dr. Jordan, differs somewhat in the count of the fin rays; but the very small size of the specimen prevents us from being entirely sure of its correctness. (*carmen*, a hetchel, from the rough scales.)

*Tripterygion carminale*, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1881, 362, Mazatlan. (Type, No. 28118. Coll. Gilbert.)

*Enneanectes carminalis*, JORDAN, Proc. Cal. Ac. Sci. 1895, 510, with plate of young example.

## 869. HETEROSTICHUS, Girard.

*Heterostichus*, GIRARD, Proc. Ac. Nat. Sci. Phila. 1854, 143 (*rostratus*).

Body rather elongate, compressed, covered with very small, smooth scales, those along lateral line not enlarged; head long and low, the snout conic, produced, very acute; premaxillaries protractile; mouth moderate, terminal; each jaw with a row of conical teeth, behind which anteriorly is a broad patch of villiform teeth; vomer and palatines with villiform teeth; gill rakers feeble; gill membranes broadly united, free from the

isthmus  
long, th  
rest an  
est, rat  
pectora  
pectora  
from wh  
separat  
tiation

Head  
maxilla  
soft dor  
deepest  
strongly  
interorb  
entirely  
of head  
opening  
or twelf  
length o  
of the p  
to the a  
transluc  
irregular  
tinct lig  
similar s  
outline o  
dorsal to  
color var  
spot beh  
whole le  
Clinoid  
tion. H  
Francisc  
*Heterostich*  
(Type  
GÜNT

*Gibbonsia*,  
*Blakea*, ST

Body l  
those al  
decurved  
conical t

isthmus; orbital cirri minute or wanting; checks scaly. Dorsal fin very long, the posterior rays soft, the 5 anterior spines wider apart than the rest and separated from them by a notch, the first and second spines longest, rather flexible, the other spines stiff; caudal fin forked; ventrals I, 3; pectorals moderate; lateral line simple, complete, abruptly curved behind pectorals; air bladder present, large. Size large. Close to *Gibbonsia*, from which the presence of the air bladder and the form of the caudal separate it. (*ἔρεπος*, different; *στίχος*, rank; in allusion to the differentiation of the anterior dorsal spines.)

## 2688. HETEROSTICHUS ROSTRATUS, Girard.

(KELPFISH.)

Head  $3\frac{1}{2}$  in body; depth  $4\frac{1}{3}$ . D. V-XXXIII, 13; A. II, 34; eye 7 in head; maxillary  $2\frac{1}{2}$ ; pectoral  $1\frac{1}{3}$ ; ventral  $2\frac{1}{2}$ ; first dorsal spine  $4\frac{1}{3}$ ; highest ray of soft dorsal  $2\frac{1}{2}$ ; third anal ray  $2\frac{5}{8}$ ; caudal  $2\frac{1}{2}$ . Body much compressed, deepest anteriorly; head slender, compressed and pointed; lower jaw strongly projecting, with thick lip; maxillary reaching pupil; width of interorbital a little greater than eye; orbital cirrus minute, usually entirely wanting; cheek and upper edge of opercle with small scales, rest of head naked. Origin of dorsal a little in front of the vertical from gill opening; pectoral under third dorsal spine, reaching to below the eleventh or twelfth; ventrals inserted in front of pectorals in distance equal to length of snout, their tips reaching about  $\frac{1}{2}$  of their length beyond base of the pectoral; soft dorsal higher than spinous, ending slightly anterior to the anal; caudal furcate, the middle rays  $\frac{3}{8}$  length of outer. Color translucent, reddish brown, varying to blackish or olive, a series of large irregular light spots along sides below lateral line, continuous with a distinct light bar from eye to edge of opercle, bordered with black above, a similar spot on base of pectoral; an irregular line of large spots following outline of body under dorsal and above anal; a clear cut white streak from dorsal to tip of snout and continued on lower lip, the hue and pattern of color varying greatly; young examples most variegated; a translucent spot behind third dorsal spine, generally followed by similar spots for the whole length of the fin. San Francisco to San Diego. The largest of the Clinoid blennies, very abundant in the kelp, with which it agrees in coloration. Here described from a specimen, 16 inches in length, from San Francisco market. (*rostratus*, long-nosed.)

*Heterostichus rostratus*, GIRARD, Proc. Ac. Nat. Sci. Phila. 1854, 143, San Diego, California (Type, No. 284. Coll. A. Cassidy); GIRARD, Pac. R. R. Surv., x, Fishes, 26, pl. 13, 1858; GÜNTHER, Cat., 261, 1861; JORDAN & GILBERT, Synopsis, 764, 1883.

## 870. GIBBONSIA, Cooper.

*Gibbonsia*, COOPER, Proc. Cal. Ac. Nat. Sci., III, 1864, 109 (*elegans*).

*Ilakea*, STEINDACHNER, Ichth. Beiträge, v, 148, 1876 (*elegans*).

Body less elongate and compressed, covered with minute cycloid scales, those along lateral line not enlarged; lateral line complete, abruptly decurved behind the pectoral; head somewhat pointed; snout unequal; conical teeth on jaws and vomer, the teeth mostly in single series, except

in front, where there is a narrow villiform band; no conspicuous posterior canines; maxillary not produced backward from angle of mouth; a tentacle above eye, none at nape; gill membranes united, free from the isthmus. Shoulder girdle with an upturned hook on its inner edge as in *Clinus*. Dorsal fin long and low, chiefly composed of spines, 5 of the anterior spines different from the others, longer and set farther apart; anal fin low, with 2 spines; ventral fins jugular, of 1 spine and 2 or 3 rays; caudal fin truncate; branchiostegals 6; no air bladder; pyloric caeca absent. Viviparous. Pacific coast; bright-colored fishes, inhabiting rock pools among algae. This genus is very close to *Clinus* (type *C. acuminatus*, Cuvier & Valenciennes), differing chiefly in the form of the dorsal fin and in the pointed snout. In *Clinus* the first 3 dorsal spines are shorter than the others. (Named for Dr. William Peters Gibbons, of Alameda, California, who was one of the early naturalists in the California Academy of Sciences.)

- a. Dorsal rays about V-XXXI, 10; anal rays about II, 26; soft dorsal low; coloration comparatively plain, the soft dorsal without pellucid area. EVIDES, 2689.  
 aa. Dorsal rays about V-XXVIII, 7; anal rays II, 24; soft dorsal high; coloration more or less highly variegated; soft dorsal with a large pellucid blotch posteriorly.

ELEGANS, 2690.

2689. GIBBONSIA EVIDES (Jordan & Gilbert).

(KELPFISH; SEÑORITA.)

Head  $4\frac{3}{4}$ ; depth  $4\frac{1}{2}$ . D. V-XXX or XXXI, 10 or 11; A. II, 26 or 27. Body elongate, compressed; head small, rather pointed; mouth quite small, terminal, the maxillary about reaching pupil,  $3\frac{1}{4}$  in head; lower jaw projecting, vomer with teeth; no teeth on palatines; posterior teeth not recurved; eye moderate, shorter than snout, 5 to 6 in head; a small supra-ocular flap, not higher than pupil; nasal cirrus very small; first spine of dorsal inserted over preopercle, its length more than  $\frac{1}{2}$  that of head, the second nearly equal; the third, fourth, and fifth progressively shorter; the sixth about as long as the fourth; the seventh longer; the rest nearly equal to the last, which is lower than the soft rays; the soft dorsal lower and more rounded than in *G. elegans*, the longest ray  $2\frac{3}{4}$  in head; pectorals moderate, not reaching vent; ventrals moderate; scales very small, smooth; head naked; no air bladder. Usual color of adult, translucent, reddish or orange, nearly plain or with oblong dark clouds below middle of sides anteriorly; often scattered blackish spots on sides, irregularly placed, forming a broken lateral band, most distinct in the young; a large pellucid spot on the membrane behind third dorsal spine, sometimes some small ones behind it; pectorals nearly plain; dorsal and anal plain, reddish, with a broad dusky shade distally; soft dorsal without pellucid area; caudal plain; a dark streak backward from eye; young examples often variegated, with light and dark shades of red, brown, and white, sometimes with 6 to 8 dark cross bars, sometimes with 4 or 5 lengthwise stripes alternating with paler ones, the hue varying exceedingly and dependent on the surroundings, but never so extravagantly spotted as in *Gibbonsia elegans*. Length 9 inches. Coast of California south to Point Concepcion; abundant in the kelp, rarely in rock pools. Here described from specimens from Monterey. (*εὐεῖδης*, comely; *εὐ*, well; *εἶδος*, appearance.)

Blake,  
cisc  
*Clinus*  
oxe

Head  
compre  
obliqu  
jecting  
as snou  
nasal  
dorsal  
torals  
a slend  
or red,  
darker  
brown  
or more  
usually  
and ca  
(*Macro*  
soft dor  
or obse  
pools l  
specime  
smaller  
though  
other in

*Myxodes*  
*Clinus* o  
*Clinus* e  
not o

\* The 1  
dan & G  
cias. It  
genus *C*

† The f  
Head 4  
pressed,  
orbital s  
tentacle  
Three fir  
interval  
longer th  
well sepa  
below la  
ocellus a  
pale brow  
than tha  
terior op  
extremif  
the right  
rounded  
suboperc  
of Califor

*Blakea elegans*, STEINDACHNER, Ichth. Beiträge, v, 148, 1876, specimens from San Francisco; not *Myxodes elegans*, COOPER.

*Clinus evides*,\* JORDAN & GILBERT, Synopsis, 763, 1883; specimens from Monterey, exclusive of part of synonymy; name a substitute for *elegans*, preoccupied in *Clinus*.

2690. GIBBONSIA ELEGANS (Cooper).

(SPOTTED KELPFISH.)

Head  $4\frac{1}{2}$ ; depth  $4\frac{1}{2}$ . D. V-XXVIII, 7; A. II, 24. Body rather strongly compressed; head short, rather pointed, mouth small, terminal, rather oblique, the maxillary barely reaching pupil,  $3\frac{1}{2}$  in head; lower jaw projecting; teeth as in *Gibbonsia evides*; eye rather large,  $4\frac{1}{2}$  in head, as long as snout; a small fringed supraocular flap, as long as pupil; a slender nasal cirrus; first dorsal spine  $2\frac{1}{2}$  in head; fins as in *G. evides*, the soft dorsal shorter, higher, and less rounded, its longest ray  $2\frac{1}{2}$  in head; pectorals and ventrals moderate, about as in *G. evides*; caudal fan-shaped on a slender peduncle; scales small and smooth; head naked. Color brown or red, agreeing with rocks or with *Corallina*, usually with eight irregular darker cross bars extending on the dorsal and anal, sometimes nearly plain brown; a dark spot probably always present behind head, and some, 1 or more, along lateral line posteriorly; spinous dorsal with a pellucid spot; usually many pale and dark spots and freckles on head and fins; pectoral and caudal usually barred, but plain in specimens taken in the kelp (*Macrocystis*), these latter much less variegated than tide-pool specimens; soft dorsal always with a large pellucid blotch posteriorly, this wanting or obscure in *G. evides*. Coast of southern California; abundant in rock pools lined with *Corallina* from Point Concepcion to Todos Santos; the specimens here described from Point Loma. Close to the preceding, but smaller and more brightly colored, the fin rays fewer. These differences, though small, seem to be constant; whether the 2 species overlap each other in geographical range is not known. (*elegans*, elegant.)

*Myxodes elegans*, COOPER, Proc. Cal. Ac. Sci., III, 1864, 109, San Diego and Santa Barbara.

*Clinus ocellifer*, † MOCQUARD, Bull. Soc. Philom. Paris 1886, 44, California.

*Clinus evides*, ROSA SMITH, Proc. U. S. Nat. Mus. 1883, 235, specimens from Todos Santos; not of JORDAN & GILBERT.

\* The name *evides* may apparently be retained for this species, as the description of Jordan & Gilbert (Synopsis, 763) is based entirely on Monterey specimens, typical of this species. It was intended, however, as a substitute for the name *elegans*, already used in the genus *Clinus*, to which these species were then referred.

† The following is a translation of the description of *Clinus ocellifer* (Mocquard):

Head  $4\frac{1}{2}$ ; depth  $4\frac{1}{2}$ . D. IPI-XXX, 8; A. II, 24; C. 13; P. 12; V. I, 3. Body strongly compressed, tapering rapidly behind. Eye a little longer than snout,  $3\frac{1}{2}$  in head, twice inter-orbital space; lower jaw a little longer than maxillary, reaching front of eye; a little tentacle on anterior nostril, elongate, with 4 or 5 unequal branches; a tentacle over eye. Three first dorsal spines nearly double length of those which follow, and separated by an interval equal to that which separates the first spine from the third; last dorsal spines longer than those that precede and stronger than any of the others; dorsal and caudal well separated. Scales very small. Opposite fifth and sixth dorsal spines immediately below lateral line is a lens-shaped spot of brownish black with a dull border; a second ocellus a little before the posterior extremity of the spinous dorsal; this surrounded by a pale brown circle in 1 specimen; the sides also with 5 irregular bands of a paler brown than that of the spots; the anterior is a little behind the corresponding spot; the posterior opposite the third or fourth soft ray of the dorsal; the posterior spot at the upper extremity of the fourth band; other spots of the same color at the base of the dorsal on the right of the caudal; sides with a longitudinal series of small white spots, not surrounded by black circle; other spots on the anal in 6 transverse lines; larger spots on snopercle and about the ventrals. Teeth on the vomer, none on the palatines. Coast of California. Two specimens, 83 mm. long. (Mocquard.)

## 871. NEOCLINUS, Girard.

*Neoclinus*, GIRARD, U. S. Pac. R. R. Surv., X, Fish., 114, 1858 (*blanchardi*).  
*Pterognathus*,\* GIRARD, Proc. Ac. Nat. Sci. Phila. 1859, 57 (*satiricus*).

Body compressed, rather elongate, covered with minute cycloid scales; lateral line present, incomplete, high anteriorly; head naked, the cheeks tumid; upper jaw protractile; maxillary greatly produced backward, more than  $\frac{2}{3}$  length of head, reaching far beyond the eye; both jaws, vomer, and palatines with stout, unequal, conical teeth in a single series, besides which, in the front of the jaws, are smaller teeth; nasal and supra-ocular region with fringed tentacles; gill membranes broadly united, free from the isthmus; gill rakers weak. Dorsal fin long, scarcely emarginate, its anterior  $\frac{2}{3}$  composed of slender, flexible spines, which are similar to the soft rays, all of which are simple; anal long, its rays all simple; ventrals moderate, 1, 3; caudal fin distinct; pectorals rather broad, rounded; no air bladder; no pyloric caeca. Pacific coast, in shallow water; remarkable for the great development of the maxillary, as in *Opisthognathus* and *Gillichthys*. (*νέος*, new; *κλίνοσ*, *Clinus*.)

## NEOCLINUS:

a. Maxillary long, but not reaching beyond head; membrane of jaws white,

BLANCHARDI, 2691.

PTEROGNATHUS (*πτερόν*, wing; *γνάθος*, jaw):

aa. Maxillary inordinately developed, reaching gill opening in the adult; maxillary flap blackish, edged with bright yellow. SATIRICUS, 2692.

## Subgenus NEOCLINUS.

## 2691. NEOCLINUS BLANCHARDI, Girard.

Head 4; depth 5 $\frac{1}{2}$ . D. XXIV, 17; A. II, 30; eye 5 in head; maxillary variable, about 1 $\frac{1}{2}$ ; pectoral 2; caudal 1 $\frac{1}{2}$  to 2. Upper profile of head convex, snout rather steep; jaws subequal; teeth on jaws, vomer, and palatines, subequal, canine-like; eye set high in head, equal to length of snout. Males with a long thick cirrus over front of middle of eye, twice as long as eye, its end multifid, 3 or 4 short, slender ones behind it over posterior half of eye; females with a much smaller cirrus in front, seldom as long as eye, the posterior ones similar to those of male; both with a multifid flap at anterior nostril; maxillary never reaching past preopercle (in specimens from 6 to 8 inches in length), not longer in males than in females. Head naked; scales on body very small, somewhat embedded; no scales on fins; origin of dorsal directly behind occiput, no notch between spinous and soft dorsals; the tips of last dorsal and anal rays reaching to base of caudal fin; pectorals broad, scarcely reaching to vent; about  $\frac{1}{2}$  the length of ventrals in front of base of pectoral. Color varying from dark red or

\*"It is more than probable that had we been acquainted with this second species of *Neoclinus* first, we would have been misled as to its real generic characters, and framed a name in allusion to the condition of the upper jaw, such as *Pterognathus*, for example, which would have been most characteristic, for that upper jaw is as truly winged as the upper members of the flying squirrels. We can not help thinking that Cuvier himself would not have coined the name of *Opisthognathus* had he had before him the species which bears his name instead of that which he dedicated to Sonnerat. These two genera (*Opisthognathus* and *Neoclinus*) will furnish one of the best themes to ichthyological studies, as they exemplify the fact that specific characters may be developed to exaggeration, and become more conspicuous than the generic characters themselves." (Girard.)

plum  
spot,  
dorsal  
white  
entire  
Coast  
able fr  
Pacific  
in ma  
differs  
dorsal  
entire  
slight  
for its  
*Neoclin*  
No.  
Syn

Head  
2; can  
equal;  
equal t  
4 small  
times a  
multifid  
preoper  
longer  
embedd  
notch b  
reaching  
anterior  
reddish  
spot oce  
lar one  
ventrals  
mens; a  
dal rays  
broadly  
of Calif  
ing spec  
Pacific C  
lary; sli  
dorsal; r  
blackish  
*Neoclinus*  
30 fact  
Synop

plum color to olive green; sides mottled and spotted with darker; a dark spot, ocellated with yellow, generally present between first and second dorsal spines; dorsal blackish toward ends of rays; pectorals and anal white in female, slightly dusky in male; unexposed portion of lower lip entirely white; a yellow spot on base of caudal rays below and above. Coast of California, from Monterey to Santa Barbara; not rare; a remarkable fish. Here described from specimens from 6 to 8 inches in length, from Pacific Grove, California. We do not know what variation there may be in maxillary and barbels in larger or smaller specimens. This species differs from *N. satiricus* in having no second spot behind seventh spine of dorsal; in having that part of lower lip which is covered by the maxillary entirely white; barbels in male much longer; maxillary shorter; head slightly shorter; and in having the pectorals and anal lighter. (Named for its discoverer, Dr. S. B. Blanchard.)

*Neoclinus blanchardi*, GIRARD, U. S. Pac. R. R. Surv., x, Fish., 114, 1858, San Diego (Type, No. 691. Coll. Dr. S. B. Blanchard); GÜNTHER, Cat., III, 253, 1861; JORDAN & GILBERT, Synopsis, 761, 1883.

Subgenus PTEROGNATHUS, Girard.

2692. NEOCLINUS SATIRICUS, Girard.

Head  $3\frac{1}{2}$  in body; depth 6. D. XVI, 17; A. 30; eye 5 in head; pectoral 2; caudal  $2\frac{1}{2}$ . Head bluntish, convex in profile; snout steep; jaws subequal; unequal, small canines on jaws, vomer, and palatines; eye about equal to length of snout, interorbital flatish, about  $\frac{1}{2}$  eye in width; 3 or 4 small barbels above eye, seldom as long as eye, the anterior one sometimes absent on one or both sides; cirri not differentiated in the female; a multifid flap on anterior nostril; maxillary always reaching past edge of preopercle (in examples 6 to 9 inches in length), just past in females, longer than head in males. Head naked, scales on body small, partly embedded; no scales on fins. Origin of dorsal directly behind occiput; no notch between spinous and soft dorsals; pectorals in the larger examples reaching to vent; last rays of dorsal and anal reaching base of caudal fin; anterior half of ventrals in front of base of pectorals. Color in spirits, reddish brown or olive green, mottled and spotted with darker; a dark spot ocellated with yellow between first and second dorsal spines, a similar one between seventh and ninth; dorsal blackish, pectoral, anal and ventrals varying from dusky to black, in no case light in our specimens; a yellow spot sometimes present, below and above, on base of caudal rays; the membrane connecting maxillary with lower jaw blackish, broadly and abruptly edged with white (probably yellow in life). Coast of California, from Monterey to Santa Barbara; a rare and most interesting species. Here described from specimens, 6 to 9 inches in length, from Pacific Grove, California. Differing from *N. blanchardi* in length of maxillary; slightly larger head; males without long cirri; a second spot on dorsal; fins darker, and especially in having the membrane of lower lip blackish, edged with white. (*satiricus*, satirical.)

*Neoclinus satiricus*, GIRARD, Proc. Ac. Nat. Sci. Phila. 1859, 57, Monterey, California, in 30 fathoms (Coll. A. S. Taylor); GÜNTHER, Cat., III, 260, 1861; JORDAN & GILBERT, Synopsis, 761, 1883.



## 872. MALACOTENUS, Gill.

*Malacotenus*, GILL, Proc. Ac. Nat. Sci. Phila. 1860, 103 (*delalandi*).

This genus is very close to *Labrisomus*, differing in the dentition, the teeth in the jaws being in single series; vomer with a few teeth or with none, none on palatines. The form of the dorsal fin in some species is different, there being usually a notch behind the fourth spine as well as at front of soft dorsal. Most of the species are not well known, and perhaps more than 1 genus is here included. (*μαλακός*, soft; *κνείς*, comb, in reference to the comb of filaments at the nape in the typical species.)

a. Nape without filaments.

b. Orbital tentacle present.

c. D. XXI, 8; spinous dorsal not notched, the first rays shortest; body elongate; snout pointed; scales large, about 38. OCELLATUS, 2693.

cc. D. XX, 12; spinous dorsal weakly notched; body rather robust.

VARIUS, 2694.

bb. Orbital tentacle wanting; dorsal rays XXI, 11; spinous dorsal weakly notched; ventrals long. MACROPUS, 2695.

aa. Nape with a single tentacle. D. XVIII, 9; a tentacle above eye. LUGUBRIS, 2696.

aaa. Nape with a comb of slender tentacles; spinous dorsal more or less notched behind fourth or fifth spine.

d. Orbital tentacle present. D. XVIII to XX, 11 or 12 vomer with teeth.

e. Scales 43 or 44.

f. Highest soft ray of dorsal  $1\frac{1}{2}$  in head; dorsal without ocelli.

GILLI, 2697.

ff. Soft rays of dorsal  $1\frac{1}{2}$  in head; dorsal fin with 2 large black ocelli; ventral fins long, as long as head. BIMACULATUS, 2698.

ee. Scales 55; ventrals moderate, shorter than head. DELALANDI, 2699.

dd. Orbital tentacle wanting; (no vomerine teeth f.).

VERSICOLOR, 2700.

aaaa. Nuchal and other filaments undescribed, a black ocellus on front of dorsal.

D. XX, 11; scales 46.

BIGUTTATUS, 2701.

## 7693. MALACOTENUS OCELLATUS (Steindachner).

Head 4 to  $4\frac{1}{2}$ ; depth 5 to  $5\frac{1}{2}$ . D. XXI, 8; A. II, 8; scales 38; eye  $4\frac{1}{2}$  to 5 in head; snout  $5\frac{1}{2}$ ; interorbital width 10. Body elongate; the snout short; profile not steep; tentacle above eye very slender, none on nape. Maxillary  $\frac{1}{2}$  long as head, reaching posterior margin of eye. Teeth on jaws and vomer in 1 row, none on palatines. Dorsal with a notch between the spines and soft rays; spines all short, the longest not  $\frac{1}{2}$  head, the anterior shortest; the longest soft rays  $1\frac{1}{2}$  in head; dorsal slightly joined to base of caudal; ventral and caudal each  $1\frac{1}{2}$  in head; pectoral almost as long as head. Lateral line complete, strongly arched anteriorly. Color brownish; 8 pairs of narrow dark-brown cross bands on the body, most distinct above, sometimes broken up into cross spots; first membrane of the dorsal fin with black spot behind, sometimes a similar one, oval and indigo, behind eye; numerous sky-blue spots bordered with darker on sides of head and part of body; anal pale violet, edged with white, sometimes spotted; caudal gray, with darker spots in cross rows. Bahama Islands. Length 2 inches. (Steindachner.) Not seen by

us.  
dorsal  
howe  
genus  
Clinus

Head  
in head  
opposi  
Teeth  
a tent  
eral li  
part fo  
which  
dorsal  
deeply  
isolated  
dotted  
(Poey.)

*Myzodes*

Head  
P. 17; e  
of eye.  
palatine  
plete; ve  
est, the  
brown.  
species e  
on the s  
(*μακρός*,

*Myzodes m*

D. XVI  
side of th  
Dorsal fin  
first spine  
brown poi  
of the first  
men 55 mm  
color.)

*Myzodes lug*

us. Perhaps not a member of this genus; the large scales, entire spinous dorsal, and short soft dorsal, indicating affinities with *Starksia*, which has, however, a different dentition. It may prove to be the type of a distinct genus. (*ocellatus*, with eye-like spots.)

*Clinus ocellatus*, STEINDACHNER, Ichth. Beitr., v, 182, 1876, Bahama Islands.

2694. MALACOTENUS VARIUS (Poey).

Head  $3\frac{3}{4}$  in total length; depth  $4\frac{1}{4}$ . D. XX, 12; A. 18; C. 14; pectoral  $1\frac{3}{4}$  in head; eye  $3\frac{3}{4}$  in head, equal to snout. Mouth small; maxillary reaching opposite front of eye; profile prolonged; nostrils small, not tubular. Teeth firm, in 1 row, the points sharp and incurved; no teeth on vomer; a tentacle over eye, none at the nape; head naked; body scaly; lateral line short; dorsal beginning over middle of opercle, the spinous part forming a sinuous curve; the spines firm; the first higher than the 4 which follow; the last low; the next to the last lower than the last; soft dorsal higher than the spines; ventral rays apparently 2, the last one deeply divided. Color clear yellowish; the body spotted with black; an isolated spot at the end of the dorsal fin; vertical fins with all the rays dotted with black; pectoral pale, without specks. Length 52 mm. Cuba. (Poey.) Not seen by us. (*varius*, variegated.)

*Myxodes varius*, POEY, Enumeratio, 132, pl. 5, f. 2, 1875, Havana. (Coll. Rafael Arango.)

2695. MALACOTENUS MACROPUS (Poey).

Head  $4\frac{1}{2}$  in total length with caudal; depth  $5\frac{1}{2}$ . D. XXI, 11; A. I, 20; P. 17; eye  $\frac{1}{2}$  longer than snout,  $3\frac{1}{2}$  in head. Maxillary reaching front of eye. Teeth in 1 series, acute, not close-set; none on the vomer or palatines; no cilia over the eye nor on the nape; lateral line almost complete; ventrals as long as the depth of the trunk; first dorsal spine longest, the others forming a weak curve. Color uniform metallic coppery brown. Cuba (Poey); one specimen 35 mm. long. The type of this species examined by us in the Mus. Comp. Zool. It has scales 35; no hook on the shoulder girdle, and apparently no teeth on vomer or palatines. (*μακρός*, long; *πούς*, foot.)

*Myxodes macropus*, POEY, Synopsis, III, 90, 1868, Havana. (Coll. Poey.)

2696. MALACOTENUS LUGUBRIS (Poey).

D. XVIII, 9; A. 20. Tentacle over eye; a filiform appendage on the side of the neck. Ventral very long, extending much beyond the vent. Dorsal fin with 2 depressions, the soft part short and very high, the first spine moderate. Color dark brown, with oblique vertical bands and brown points scattered over the head and trunk; a black spot at the base of the first 3 dorsal membranes; ventral entirely white. Cuba. One specimen 55 mm. long. (Poey.) Not seen by us. (*lugubris*, dismal, from the dark color.)

*Myxodes lugubris*, POEY, Enumeratio, 131, 1875, Cuba. (Coll. Poey.)

## 2697. MALACOTENUS GILLII (Steindachner).

Head  $4\frac{1}{2}$ ; depth  $4\frac{1}{2}$ ; eye 4 in head; snout  $3\frac{1}{2}$ ; interorbital 6 in head. D. XIX, 11; A. II, 17; P. 14; scales 43. Head pointed, conic anteriorly; snout longer than eye; a rather high tentacle above the eye, slender and split to the base, numerous others on the side of the nape; teeth not described. First three dorsal spines wider apart than others, first longest; eighth to tenth spines highest,  $\frac{1}{2}$  head; highest soft ray  $1\frac{1}{2}$  in head; dorsal deeply notched. Body greenish gray, with brown spots or faint cross bands; head and dorsal marbled with darker; ventrals white, the longest ray a little longer than head, reaching anal; anal edged with dark; pectoral as long as head. Barbados. Two specimens, the larger 2 inches long. (Steindachner.) This species may be a *Labrisomus*. (Named for Dr. Theodore Gill.)

*Clinus gillii*, STEINDACHNER, Ichth. Notizen, VI, 46, 1867, Barbados.

## 2698. MALACOTENUS BIMACULATUS (Steindachner).

Head  $4\frac{2}{3}$ ; depth  $4\frac{1}{2}$ . D. XX, 10; A. II, 19; ventral 3; scales 44. Near to *M. delalandi*, but the body deeper (said to be  $5\frac{1}{2}$  in the latter species, which is not the case). Profile to snout steep; eye a little shorter than snout,  $3\frac{1}{2}$  in head; jaws equal, each like the vomer with 1 row of teeth; maxillary reaching about to front of pupil. Interorbital space narrow, more than  $\frac{1}{2}$  width of eye; a very slender, rather long, bifid tentacle above eye; tentacles on the nape, upper 1 almost as long as tentacle above eye. Upper margin of dorsal weakly notched between first and fifth spines, more deeply between spines and soft rays, the former as in *M. delalandi*; longest soft rays  $1\frac{1}{2}$  in head; longest spines 2 in head; first 4 spines more widely separated than the others; pectoral and ventral as long as head; caudal a little shorter. Body brown, with dark-brown bands and numerous blackish spots, only the cross bands on the head strongly marked; tips of the anal rays whitish; above these a bluish violet streak; pectoral with 2 milk-white spots at base; a large black isolated spot at the base of the first 4 dorsal spines, a second on the last 4 spines, extending on the body; anal and caudal thickly spotted with brown. Small rocky islands to the north of Cuba. (Steindachner.) Not seen by us; evidently close to *Malacotenus delalandi*, but the scales larger. (*bis*, two; *maculatus*, spotted.)

*Clinus bimaculatus*, STEINDACHNER, Ichth. Beitr., v, 180, 1876, small, rocky islands north of Cuba.

## 2699. MALACOTENUS DELALANDI (Cuvier &amp; Valenciennes).

Head  $3\frac{1}{2}$ ; depth  $3\frac{1}{2}$  to  $4\frac{1}{2}$ . D. IV-XVI, 11; A. II, 18; scales 55. Form rather stout, compressed; snout not very short, rather pointed, the profile gibbous above the eyes, thence declining straight to the tip of the snout; mouth rather small, the maxillary reaching front of eye; teeth in a single series in each jaw; vomer with a few teeth, none on palatines; eye large,  $3\frac{1}{2}$  in head, as long as snout; small slender cirri above the eyes, and a fringe of moderately long filaments at the nape rather longer than the orbital cirri. Outline of spinous dorsal emarginate; first spine a little longer than eye, the second, third, and fourth progressively shortened, the

fifth  
decr  
soft  
long  
betw  
of h  
nake  
oliva  
brow  
base  
necte  
occip  
of pe  
with  
brow  
barre  
Coast  
from  
the r  
sopora  
*M. zon*  
accou  
lected  
*Clinus*  
Dell  
*Clinus*  
Gill  
*Clinus*  
*Labriso*

Head  
erate;  
long a  
not rea  
and sl  
side [u  
of filif  
lateral  
the 4 v  
pector  
trunk,  
ventra  
body a  
in leng  
the orb  
color, v  
*Myzoda*

\* This  
genus al

fifth again longer; the eighth to eleventh spines longest, thence gradually decreasing to the next to the last, which is much shorter than the last; soft dorsal rays considerably higher than the spines, the longest about  $\frac{1}{2}$  length of head; anal long, not very high, the membrane deeply notched between all but the last 6 rays, which are the highest. Pectorals  $\frac{3}{4}$  length of head; ventrals as long as from snout to edge of preopercle. Belly naked anteriorly; the scales small, cycloid; lateral line complete. Color olivaceous, darker above, much mottled and speckled with clear dark brown; sides with 5 distinct irregular dark-brown bars, extending from base of dorsal to level of lower margin of pectoral, their lower edges connected by a vague undulating longitudinal band; a blackish blotch on occipital region, and black blotches on cheeks, opercles, and before base of pectoral; opercle with several narrow pinkish streaks; head below with narrow streaks formed by series of dark-brown spots; an interrupted brown bar across lower jaw; belly unspotted; ventrals pale; other fins all barred with narrow series of dark-brown dots; anal somewhat dusky. Coast of Brazil and the west coast of Mexico; common. Here described from the types of *Clinus zonifer*. This is the most abundant denizen of the rock pools around Mazatlan, with the single exception of *Gobius soporator*, reaching a length of 3 to 5 inches. We are unable to separate *M. zonifer* from Mazatlan from Bahia examples of *M. delalandi*, and take our account from specimens of the former. (Named for Delalande, who collected for Cuvier in Brazil.)

*Clinus delalandii*, CUVIER & VALENCIENNES, Hist. Nat. Poiss., XI, 378, 1836, Brazil (Coll. Delalande); GÜNTHER, Cat., III, 264, 1861.

*Clinus zonifer*, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1881, 361, Mazatlan. (Coll. C. H. Gilbert.)

*Clinus philippi*, LOCKINGTON, Proc. Ac. Nat. Sci. Phila. 1881, 114; not of STEINDACHNER.

*Labrisomus delalandi*, JORDAN, Proc. U. S. Nat. Mus. 1888, 333.

#### 2700. MALACOTENUS VERSICOLOR (Poey).

Head  $3\frac{1}{2}$ ; depth  $3\frac{1}{2}$ . D. XVIII, 12; A. 20. Body compressed; head moderate; snout prolonged; pectoral  $1\frac{1}{2}$  in head; ventral  $1\frac{1}{2}$ ; eye large, as long as snout,  $3\frac{1}{2}$  in head; nostrils not tubular; mouth small; maxillary not reaching so far as eye; teeth firm, in 1 row, those above much larger and slightly curved backward; 12 teeth above and 9 below on each side [no teeth on vomer]; no tentacle over eye; no anal papilla; a comb of filiform tentacles on each side of the neck; head naked; body scaly; lateral line short; dorsal with 2 depressions, the first spine higher than the 4 which follow, the depressions much more marked than in *M. varius*; pectoral reaching beyond front of anal. Color yellowish brown; head, trunk, and fins varied with vertical brown bands and large brown spots; ventrals yellowish. On the figure the 5 bands behind the anal cross the body and extend on the vertical fins. Cuba. One specimen known, 53 mm. in length. (Poey.) Apparently very close to *M. delalandi*, but lacking the orbital tentacle, and, according to Poey, vomerine teeth also. (*versicolor*, variegated.)

*Myxodes versicolor*,\* POEY, Enumeratio, 131, pl. 5, f. 1, 1875, Cuba. (Coll. Poey.)

\* This species and its affines were referred by Poey to *Myxodes*, a South American genus allied to *Clinus* and *Gibbonsia*, but differing from the latter in its uniserial teeth.

## 2701. MALACOTENUS BIGUTTATUS (Cope).

Dorsal XIX-I, 11; anal II, 16; The first dorsal spines the longest, last spine longer than penultimate; length of head without opercular flap,  $3\frac{1}{2}$  times in length (exclusive of caudal fin); eye a little less than  $\frac{1}{2}$  length of head,  $\frac{2}{3}$  greater than interorbital width; pectoral fin reaching to fifth anal; scales large, 4-16-10 [cirri and teeth not described]. Pale reddish brown, humeral red-veined; rufous specks on anterior part of sides; 7 subquadrate brown blotches from nape to caudal fin, continued with interruptions as lateral bands, the fourth near end of spinous dorsal black; a black spot at base of membrane between first to third dorsal spines; 2 small brown spots behind orbit, the posterior on operculum. Length 2.25 inches. This species is well distinguished from *Labrisomus nuchipinnis* by the large scales, form of dorsal fin, coloration, etc. From New Providence, Bahamas; Dr. H. C. Wood's collection. Also a very small specimen from Dr. Rijgersma, St. Martins. (Cope.) Not seen by us. (*bis*, two; *guttatus*, spotted.)

*Labrisomus biguttatus*, COPE, Trans. Am. Philos. Soc. Phila. 1873, 473, New Providence, Bahama Islands. (Coll. Dr. H. C. Wood.)

## 873. LABRISOMUS,\* Swainson.

*Labrisomus*, SWAINSON, Nat. Hist. Class'n Fishes, II, 277, 1839 (*pectinifer*).

*Lepisoma*, DE KAY, New York Fauna: Fishes, 41, 1842 (*cirrhosum*).

*Labrosomus*, GILL, amended spelling.

? *Blennioclinus*, GILL, Proc. Ac. Nat. Sci. Phila. 1860, 103 (*brachycephalus*).

? *Auchenionchus* (misprinted *Anchenionchus*), GILL, Proc. Ac. Nat. Sci. Phila. 1860, 103 (*variolosus*).

? *Calliclinus*, GILL, Proc. Ac. Nat. Sci. Phila. 1860, 103 (*geniguttatus*).

? *Ophthalmolophus*, GILL, Proc. Ac. Nat. Sci. Phila. 1860, 104 (*latipinnis*).

\* Concerning this genus and its affines, Dr. Gill remarks:

"The name *Labrosomus* (or *Labrisomus*) was first published in 1839, in the second volume of the 'Natural History of Fishes, Amphibians, and Reptiles.' At the seventy-fifth page of that volume, Swainson has divided the Cuvierian genus *Clinus* into 5 genera: *Olinus*, of which the *Olinus acuminatus*, Cuvier, is taken as the type; *Labrisomus* with *Clinus pectinifer*, Valenciennes, as type; *Tripterygion*, Risso, *Olinitrachus*, Reese, which is typified by *Blennius variabilis* of Rafinesque, and *Blennopsis*, of which the *Clinus anguillarum*, Valenciennes is the only true species. Of these genera, *Clinus* Swainson, and *Olinitrachus* Swainson, are distinguished by false or illusory characters, and cannot be regarded as distinct. The others are valid, but their characters require revision. The only claim to distinction of the genus *Labrosomus* given by Swainson, is founded on the strong, conic, and pointed row of front teeth, behind which are villiform ones; a thicker body than in *Clinus*, and the dorsal fin distinctly emarginate toward the caudal. The genus resting on these characters alone is composed of very incongruous elements. To it are referred, at page 277 of the second volume, the following species, all of which are described as species of *Clinus* by Valenciennes: *Labrosomus gobio*, *L. pectinifer*, *L. capillatus*, *L. delalandii*, *L. lineatus*, *L. variolosus*, *L. peruvianus*, *L. microcirrhis*, *L. ? geniguttatus*, *L. elegans*, *L. ? litoreus* and *L. latipinnis*. Of these species, not more than 3 can, with propriety, be regarded as congeners, if the *Labrosomus pectinifer* is taken as the type. These are *Labrosomus pectinifer*, *L. capillatus*, and perhaps *L. delalandii*. The latter is more probably the representative of a distinct genus. That genus is distinguished from *Labrosomus* by the smaller mouth, the presence of only 2 rays to the ventral fins, and perhaps by the undulating margin of the spinous portion of the dorsal fin. It may be named *Malacotenus*, in allusion to the pectiniform row of filaments. This genus is the nearest ally of *Labrosomus*. All the others are very distinct. *Labrisomus gobio* Swainson, is the type of quite a distinct genus, whose characters consist of a broad, depressed head, with a very short muzzle, large approximated eyes, superciliary and nasal tentacles, 2 ventral rays, and a comparatively short spinous dorsal. The genus may be called *Gobioclinus*. The only species, *Gobioclinus gobio*, is found in the West Indies, and has but 18 dorsal spines. *Labrisomus linearis* Swainson is synonymous with *Clinus brachycephalus*, Valenciennes. This, also, is the type of a

Bod  
rather  
behind  
vomere  
with a  
ous sle  
torals  
should  
Intesti  
not we  
bly be  
chiefly  
edge of  
the typ

a. Sea

b.

bb

bb

aa. Sea

D. XV

on nostr  
teeth, b  
on the b  
from the  
brown w

distinct g  
very conv  
portion of  
name *Ble  
ciennes m  
with later  
large; the  
form ones  
on each pe  
each 3 ray  
named *A  
gans* and  
logical pr  
more appr  
small size  
of the ver  
ciliary, n  
proposed.  
species of  
*Ophthalm**

30

Body oblong, robust; head naked, short, compressed above; mouth rather large, with a row of stout, bluntish teeth in front of each jaw, behind which is a band of smaller teeth, broadest in lower jaw; teeth on vomer, no teeth on palatines; a tentacle above the eye; sides of neck with a tuft or series of fine filaments; dorsal fin continuous, with numerous slender spines and many soft rays, the spines not very unequal; pectorals long; lateral line continuous; scales moderate or small, cycloid; shoulder girdle without upturned hook-like process on its inner edge. Intestinal canal short, shorter than body. The limits of this genus are not well defined, and most of the nominal genera above named will probably be found worthy of recognition. This genus differs from *Clinus* chiefly in the absence of the upturned spine-like processes on the inner edge of the shoulder girdle. This process is found on *Clinus acuminatus*, the type of the genus *Clinus*. (*Labrus*;  $\delta\epsilon\upsilon\mu\alpha$ , body.)

- a. Scales moderate, about 70 in lateral line (so far as known); soft dorsal with 11 to 13 rays.  
 b. Dorsal spines 16; anal rays 20; tentacles on nape. HERMINIER, 2702.  
 bb. Dorsal spines 18; no teeth on palatines; first ray of dorsal not longest; orbital tentacle well developed; nape with a conspicuous comb of fringes.  
 d. Vomer with a cluster of small teeth. NUCHIPINNIS, 2703.  
 dd. Vomer with 3 to 5 large blunt teeth arranged in the form of a  $\Lambda$ .  
 XANTI, 2704.  
 bbb. Dorsal spines 20; teeth on palatines (?); first dorsal spine longest.  
 BUCCIFERUS, 2705.  
 aa. Scales very small, about 110; a comb of fringes at nape; first dorsal spines low; head with yellow spots.  
 MICROLEPIDOTUS, 2706.

2702. LABRISOMUS HERMINIER (Le Sueur).

D. XVI, 11; A. 20; C. 14; P. 16; V. 3. Body slender, compressed. Cilia on nostrils, above the eye, and on the nape; lips thick, concealing conical teeth, behind each band of smaller teeth; teeth also on the palate and on the base of the gill arches. Scales rather large. Lateral line curved from the pectoral, becoming straight thence to the tail. Color reddish brown with numerous spots; a black spot at front of spinous dorsal.

distinct genus distinguished by its abbreviated and blenniform head, the profile being very convex; by the villiform teeth, the absence of superciliary tentacles, the spinous portion of the dorsal long, and the presence of only 2 rays to the ventral fins. The name *Blennioctinus* is conferred on it; for the species, the specific name of Valenciennes must be retained. *Labrisomus variolosus* is distinguished by a large thick head, with lateral eyes, short superciliary tentacles, and a small nuchal one. The mouth is large; the teeth of the jaws in an outer row strong and conical, behind which are villiform ones; those of the vomer and palate are villiform, in 3 patches, 1 on the vomer and 1 on each palatine bone. The spinous portion of the dorsal is long, and the ventrals have each 3 rays. The species thus characterized is the type of a new genus which may be named *Anchenionchus* (misprint for *Auchenionchus*). *Labrisomus microcirrhia*, *L. elegans* and *L. peruvianus* are nearly related to *Anchenionchus*, and are from the same zoological province. *Labrosomus ? geniguttatus* is distinguished from *Anchenionchus* by the more approximated eyes and by the disposition of the vomero-palatine teeth, as well as the small size of the anterior row of maxillary teeth. The dorsal is moderately long, and each of the ventrals has 3 rays. The mouth is comparatively small, and there are superciliary, nasal, and nuchal tentacles. For this species the generic name *Callickinus* is proposed. *Labrisomus latipinnis* is related to *Blennioctinus*, but is distinguished from the species of that genus by the presence of superciliary tentacles. The generic name of *Ophthalmostophus* may be retained for it." (Proc. Ac. Nat. Sci. Phila. 1860, 102, 103.)

St. Bartholomew, West Indies; known from one specimen taken among madreporic rocks. (Le Sueur.) Not recognized by any recent author; perhaps not distinct from *L. nuchipinnis*.

*Blennius herminier*, LE SUEUR, Jour. Ac. Nat. Sci. Phila., iv, 1824, 361, St. Bartholomew.  
*Olinus herminieri*, CUVIER & VALENCIENNES, Hist. Nat. Poiss., xi, 380, 1836.

2703. LABRISOMUS NUCHIPPINNIS (Quoy & Gaimard).

Head  $3\frac{1}{2}$ ; depth  $3\frac{1}{4}$ . D. XVIII, 12; A. II, 17; scales 70. Body oblong, rather robust; head naked, thick, short, not very obtuse anteriorly, compressed above; mouth rather large, the maxillaries not prolonged backward, extending to opposite the posterior part of eye,  $2\frac{1}{4}$  in head; teeth on vomer and palatines; front teeth of jaws conic, strong, behind them a band of villiform teeth, broadest in lower jaw; vomer with a patch of smallish teeth; eyes large; interorbital space very narrow; each side of neck with a long series of hair-like filaments, nearly as long as eye; orbital tentacle short and broad, multifold; nostril with a tufted barbel; lower jaw slightly projecting, its posterior teeth sometimes recurved; pectorals a little shorter than head, reaching vent. Dorsal spines rather slender, the 3 anterior spines scarcely shorter than the others, all the spines lower than the soft rays; dorsal fin commencing near the nape, the spinous portion long; soft rays higher than the spines; caudal small; pectorals rather large; ventrals moderate; gill-membranes broadly united, free from the isthmus; lateral line complete, high anteriorly, then abruptly decurved; membranes of vertical fins scaly; scales not very small, cycloid. Reddish brown, sometimes with vertical bands; a black spot on opercle, which is often edged with white; cheeks and fins reticulate or dotted. Length 6 to 8 inches. West Indies, north to Florida Keys, south to Brazil; generally common in rock pools; also recorded from the Canary Islands. (*nucha*, nape; *pinna*, fin.)

*Olinus nuchipinnis*, QUOY & GAIMARD, Voy. Uranie et Physicienne, Zool., 255, 1824, Brazil (Coll. M. Freycinet & M. Gay); GÜNTHER, Cat., iii, 262, 1861; JORDAN & GILBERT, Synopsis, 762, 1883.

*Olinus pectinifer*, CUVIER & VALENCIENNES, Hist. Nat. Poiss., xi, 374, 1836, Bahia.

*Lepisoma cirrhosum*, DE KAY, N. Y. Fauna: Fishes, 41, 1842, Florida.

*Olinus canariensis*, VALENCIENNES, in WEBB & BERTHELOT, Poiss. Iles Canaries, 60, 17, f. 3, Canary Islands.

*Olinus capillatus*, CUVIER & VALENCIENNES, Hist. Nat. Poiss., xi, 377, 1836, Martinique.

*Labrosomus pectinifer*, GILL, Proc. Ac. Nat. Sci. Phila. 1860, 105.

*Labrisomus capillatus*, GILL, Proc. Ac. Nat. Sci. Phila. 1860, 107.

2704. LABRISOMUS XANTI, GILL.

Head  $3\frac{1}{2}$  in body; depth  $3\frac{1}{4}$ . D. XVIII, 12; A. II, 18; scales 10-64 (pores)-12 (from front of straight portion of lateral line to anal); eye  $4\frac{1}{4}$  in head, maxillary 2; highest dorsal spine  $2\frac{5}{8}$ ; pectoral  $1\frac{1}{4}$ ; caudal  $1\frac{3}{4}$ . Body not greatly elongate, compressed, anterior profile well rounded from snout to nape; mouth rather large, the maxillary reaching to below middle of eye; teeth small, canine-like, growing gradually larger toward

front of upper jaw; side teeth on lower jaw very small, abruptly enlarged on front half of jaw; teeth on vomer A-shaped, in a single row, the ones at the angles enlarged, 1 or 2 small ones between them at the sides; small multifid dermal flaps at nape, over eye, and above nostril; interorbital concave at the middle,  $\frac{2}{3}$  the diameter of eye; gill rakers small and short, 3 + 6 in number. First dorsal spine inserted behind eye a distance equal to diameter of eye, about  $\frac{1}{2}$  shorter than longest spine; soft dorsal the higher; origin of anal midway between snout and base of caudal, not running as far back as dorsal; pectoral reaching a little past front of anal; ventrals long and slender, inserted a little in front of pectorals, their ends not reaching vent; caudal rounded. Color in spirits, brownish gray, with about 6 wide irregular cross bars which are darker toward their edges, 2 black streaks running downward and backward from eye; cheeks and opercles with many small light blue spots; spinous dorsal mottled and spotted with darker, other fins with small irregular dark lines running across the rays; ventrals dusky; tentacles on head black. Described from a specimen 5 inches in length from La Paz, Lower California. Pacific coast of Tropical America from Gulf of California to Panama; common in rock pools; representing on the Pacific coast the scarcely different *L. nuchipinnis*. (Named for John Xantus.)

*Labrosomus xanti*, GILL, Proc. Ac. Nat. Sci. Phila. 1860, 107, Cerro Blanco (Type, Nos. 2334, 2335, 2478. Coll. J. Xantus); JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1882, 368.

2705. *LABRISOMUS BUCCIFERUS*, Poey.

Head  $3\frac{1}{2}$  in total length with caudal; depth 5. D. XX, 11; A. II, 19; eye 4 in head, a little longer than snout; anterior nostril with a little tube; lower jaw longer; forehead convex, the snout short; mouth large, reaching beyond middle of eye; a few filaments on nape and 1 above preopercle. Teeth cardiform, the outer ones large; teeth on vomer (and palatines). Body scaly. Lateral line complete. First ray of dorsal longest, the others forming a convex curve up to the 19, which is shortest; pectoral moderate, of 12 rays, the lower thickened. Color brownish yellow, with vertical brown points extending on fins; a series of pale points along sides; the head gray, cheek dark brown. One specimen 55 mm. long. Cuba. (Poey.) Not seen by us. (*bucca*, cheek; *fero*, I bear).

*Labrosomus bucciferus*, POEY, Synopsis, 399, 1868, Cuba. (Coll. Poey.)

2706. *LABRISOMUS MICROLEPIDOTUS*, Poey.

Length of head equal to depth; pectoral  $1\frac{1}{2}$  in head; eye  $1\frac{1}{2}$  in snout,  $4\frac{1}{2}$  in head. Maxillary reaching to base of middle of eye,  $2\frac{1}{2}$  in head. Mouth oblique, with strong teeth; the lower jaw the longer; small fringe of tentacles at anterior nostril above eye, and a comb of fringes at the nape. First 3 dorsal spines subequal, considerably lower than the second 3, which become progressively longer. Scales in lateral line about 110. Head brown, with small yellow spots scattered over its lower part and on the gill membranes. Pectoral and caudal with some black points. Cuba.



(Poey.) Known from an imperfect description, with a drawing of the head of a specimen 180 mm. long. (*μικρός*, small; *λεπιδωρός*, scaly.)

*Labrosomus microlepidotus*, POEY, Anal. Soc. Esp. Hist. Nat., XIX, 1880, 246, 1, 8, f. 2, Cuba. (Coll. Poey.)

#### 874. MNIERPES, Jordan & Evermann.

*Mnierpes*, JORDAN & EVERMANN, Check-List Fishes, 468, 1896 (*macrocephalus*).

This genus is close to *Labrisomus*, from which it differs chiefly in the very elongate body and in the absence of an orbital tentacle. The dorsal spines are more numerous, and probably the vertebrae also. The lips are thick and there is no trace of hook on the shoulder girdle. A band of filliform teeth in the jaws behind the anterior series; teeth on vomer, none on palatines. (*μύριον*, moss; *ἔρπηξ*, creeper.)

#### 2707. MNIERPES MACROCEPHALUS (Günther).

Head  $4\frac{1}{2}$ ; depth 6 to  $6\frac{1}{2}$  ( $7\frac{1}{2}$  in total). D. XXII, 12; A. II, 24; C. 13; P. 13; V. I, 3; scales about 70. The head is depressed, rather short, nearly as broad as long; crown of the head broad and flat; interorbital space concave, narrower than the orbit. Snout very short, obtuse, rounded; the maxillary not extending to behind the posterior margin of the orbit; lips thick. Teeth in jaws forming a band with an outer series of stronger ones; vomerine teeth in a narrow band; palatine teeth none. No orbital tentacles, those at the nostril and on the neck very small. Gill openings wide, the gill membranes being united at the throat. Head naked; scales on the body not very small, cycloid. Dorsal fin commencing at occiput, and terminating near base of caudal, the spines flexible, and much lower than the soft rays; the 3 anterior ones rather more remote from one another than the following; none of the rays of this or of the other fins branched; caudal rounded; anal higher posteriorly than anteriorly, about as high as the spinous dorsal; pectorals rounded, with the middle rays longest, shorter than the head; ventrals jugular,  $\frac{1}{2}$  as long as the pectoral, with the spine and the outer ray enveloped in a common thick membrane. Dark grayish olive; head and fins blackish; head, base of the pectoral, anterior part of the body, and dorsal dotted with white. Pacific coast of Central America. (Günther.) Known from a few specimens from Panama. Those examined by us (*Mus. Comp. Zool.*) have the sides much freckled and mottled with pale. (*μακρός*, long; *κεφαλή*, head.)

*Clinus macrocephalus*, GÜNTHER, Cat., III, 267, 1861, Pacific coast of Central America (Coll. Capt. John M. Dow); GÜNTHER, Fish. Centr. Amer., 442, pl. 69, fig. 2, 1869.  
*Labrosomus macrocephalus*, JORDAN, Proc. U. S. Nat. Mus. 1885, 389.

#### 875. GOBIOCLINUS, Gill.

*Gobioclinus*, GILL, Proc. Ac. Nat. Sci. Phila. 1860, 102 (*gobio*).

Body robust; head broad, depressed, with a very short muzzle. Eyes large, approximated, close together; palatine teeth present; a tentacle above eye; no nuchal filaments. Scales very large, about 30 in the lateral

line.  
risom  
nuch

Hea  
14; V  
less.  
over t  
desce  
the sk  
what  
teeth,  
vomer  
free fr  
betwe  
total l  
opposi  
the cro  
Known  
ciennes  
the gud  
*Clinus g*  
(Coll.  
*Gobioclin*

*Starksia*,

This  
presenc  
the com  
recognit

Head  
In appe  
*opterus*.  
the max  
Teeth s  
series en  
in front  
4 in hea  
minating  
gill mem  
of shoul  
eye and

line. Spinous dorsal of 18 spines. This genus seems to differ from *Labrisomus* in the large scales, differently formed head, and in the absence of nuchal filaments. (*Gobio*, the gudgeon; *Clinus*.)

2708. *GOBIOCLINUS GOBIO* (Cuvier & Valenciennes).

Head  $3\frac{1}{2}$  in total length; depth  $4\frac{1}{2}$ . D. XVIII, 19; A. II, 17; C. 15; P. 14; V. 2; scales 30–10. Head nearly as broad as long, its height a third less. Eye large,  $2\frac{1}{2}$  in head, twice interorbital space; a very small tentacle over the eye, another on the nostril. Profile rounded between the eyes, descending vertically to the snout, which is very short. Cheeks inflated; the skull a little rough. Mouth reaching to opposite middle of eye, somewhat black; teeth small, conic, and pointed; upper jaw with 26 equal teeth, the lower with 16, the last 2 larger and more curved; teeth on vomer and palatines, simple, in 2 irregular rows; gill membranes united, free from isthmus. Body posteriorly compressed. Dorsal slightly notched between spines and soft rays of anal; pectorals equal to ventrals, 5 in total length; caudal obtuse, 6 in total length. Lateral line disappearing opposite tip of ventral. Color greenish, with traces of cloudy brownish; the cross bands a deep brown, pointed at base of caudal. Lesser Antilles. Known from several specimens, one 2 inches in length. (Cuvier & Valenciennes.) Not seen by us; apparently a strongly marked species. (*Gobio*, the gudgeon, from its resemblance to *Cottus gobio*, the miller's thumb.)

*Clinus gobio*, CUVIER & VALENCIENNES, Hist. Nat. Poiss., XI, 395, 1836, Lesser Antilles. (Coll. Plée.)

*Gobioclinus gobio*, GILL, Proc. Ac. Nat. Sci. Phila. 1860, 102.

876. *STARKSIA*, Jordan & Evermann.

*Starksia*, JORDAN & EVERMANN, Proc. Cal. Ac. Sci. 1896, 231 (*cremnobates*).

This genus is related to *Labrisomus*, differing in the large scales, the presence of palatine teeth, the short soft dorsal fin, and the absence of the comb of nuchal filaments. (Named for Mr. Edwin Chapin Starks, in recognition of his work on the fishes of Western America.)

2709. *STARKSIA CREMNOBATES* (Gilbert).

Head  $3\frac{1}{2}$  in length; depth  $4\frac{1}{2}$ . D. XXI or XXII, 8; A. II, 19; scales 37. In appearance resembling very strongly the species of the genus *Auchenopterus*. Body slender, snout sharp, the jaws equal; mouth wide, oblique, the maxillary reaching vertical from posterior margin of orbit,  $2\frac{1}{2}$  in head. Teeth small, villiform, forming a band in front of upper jaw, the outer series enlarged; in lower jaw a single series laterally, becoming double in front; similar teeth on vomer and palatines. Eye longer than snout, 4 in head; interorbital width less than diameter of pupil; opercle terminating in an evenly convex process behind, without spinous points; gill membranes broadly united, free from isthmus; no hook on inner edge of shoulder girdle; nostrils with a flap; a single slender filament above eye and 1 or more on each side of the nape. A slight notch between first

and third dorsal spines and another between the eighteenth and twenty-first spines; the spines are low and strong, the highest equaling the snout and  $\frac{1}{2}$  eye; soft rays higher, the longest equaling  $\frac{1}{2}$  head; caudal short, rounded, entirely free from dorsal and anal; anal similar to soft dorsal, the first 2 rays spinous; ventrals inserted well in advance of pectorals, each consisting of 1 spine and 2 soft rays, which are joined only at base; pectorals pointed, the lower rays the longest,  $1\frac{1}{2}$  in head. Scales large, cycloid, the lateral line running high in front, descending to middle of sides immediately behind pectorals, thence running straight to tail. In the types, which are probably immature, the pores are not developed on posterior part of body. Color in spirits, uniform light olivaceous, a small dusky spot behind orbit and 1 below and behind it; opercle dusky. In 1 specimen the rays of soft dorsal, anal, and caudal are finely barred with dusky. (Gilbert.) Length  $1\frac{1}{2}$  inches. Gulf of California. Two specimens known, from Albatross Station 3001, in 71 fathoms. (*Cremnobates*; κρημνοβάτης, one that haunts rocks; a synonym of *Auchenopterus*.)

*Labrosomus cremnobates*, GILBERT, Proc. U. S. Nat. Mus. 1890, 100, Gulf of California. (Coll. Albatross).

*Starksia cremnobates*, JORDAN, Proc. Cal. Ac. Sci. 1896, 231.

#### 877. CRYPTOTREMA, Gilbert.

*Cryptotrema*, GILBERT, Proc. U. S. Nat. Mus. 1890, 101 (*corallinum*).

This genus differs from *Labrosomus* chiefly in the absence of nuchal filaments and in the modified anterior portion of the lateral line, which runs on a series of enlarged scales having no externally visible pores. (κρυπτός, concealed; τρήμα, pore.)

#### 2710. CRYPTOTREMA CORALLINUM, Gilbert.

Head  $3\frac{1}{2}$  to 4 in length; depth  $5\frac{1}{2}$ . D. XXVII, 12; A. II, 27. Body elongate, regularly tapering backward to caudal peduncle, whose depth equals length of snout, which is sharp; mouth nearly horizontal; maxillary reaching middle of eye or beyond,  $2\frac{1}{2}$  to  $2\frac{1}{2}$  in head; teeth strong, but none of them enlarged, in a single series in jaws laterally, becoming double anteriorly; teeth on vomer and in a small distinct patch on front of palatines; eyes large, the interorbital space flat, nearly  $\frac{1}{2}$  diameter of orbit; orbit slightly exceeding length of snout,  $3\frac{1}{2}$  in head; branchiostegial membranes broadly united, free from isthmus, the posterior edge on vertical from preopercular margin; anterior nostril in a short tube, a slender flap arising from its posterior margin; a pair of simple slender filaments arising from the upper edge of each orbit, 1 on each side of nape, none others on head; gill rakers very short and weak; shoulder girdle without hook on its inner edge. Scales rather large, cycloid, the head alone naked; lateral line in its upper anterior portion without externally visible tubes, its position shown by a series of enlarged scales twice the size of the others; on these the tubes are wholly on the under side, each opening anteriorly by a single pore under the edge of the pre-

ceding scale; anteriorly the lateral line runs near the back and parallel with it, becoming suddenly declined behind middle of trunk, thence running on middle of side; the oblique portion of lateral line rests on about 7 scales, and the externally visible tubes of lateral line begin at this point; posterior portion of lateral line contained  $1\frac{1}{2}$  to  $1\frac{3}{4}$  in dorsal portion; scales of lateral line, 45 in dorsal portion, 7 in oblique portion, and 18 in posterior portion. A slight notch behind fourth dorsal spine, the second and third spines slightly longer than those following, the first little longer than the fourth, the longest spine about  $2\frac{1}{2}$  in head; first 2 anal rays spinous, but weak and flexible; last dorsal and anal rays not joined by membrane to caudal peduncle, the depth of the latter equaling the length of its free portion; ventrals long and narrow, nearly reaching vent in males, consisting of 1 spine and 3 simple rays; pectorals with some of the lower rays longest,  $1\frac{1}{2}$  in head; all of pectoral rays simple, 14 in number; caudal fin truncate,  $1\frac{3}{4}$  to  $1\frac{1}{2}$  in head. Length 5 inches. Color dusky olive above, with irregular narrow longitudinal streaks of bright coral red, and 7 round black blotches above middle of sides; reticulating red lines and spots on top and sides of head and snout; branchiostegal membranes dusky in males; 2 red streaks on base of pectorals; dorsal somewhat dusky, marked with lines of red spots; caudal with 3 rather faint cross bars; pectorals, ventrals, and anal largely black in males, pale in females; the red shades persistent in alcohol. Santa Barbara Islands. Three specimens from Albatross Station 2945, in 30 fathoms. (Gilbert.) (*Corallina*, a calcareous alga, among which it lives.)

*Cryptotrema corallinum*, GILBERT, Proc. U. S. Nat. Mus. 1890, 101, off Santa Barbara Islands. (Coll. C. H. Gilbert.)

### 878. EXERPES, Jordan & Evermann.

*Exerpes*, JORDAN & EVERMANN, Proc. Cal. Ac. Sci. 1896, 232 (*asper*).

Body slender, much compressed; the snout long, sharp in profile; first dorsal ribbon-shaped, the 3 slender spines close together, inserted at the nape, much in advance of the rest of the fin; ventrals very long and slender. Otherwise as in *Auchenopterus*, the scales large, and but 1 soft ray in the dorsal fin. ( $\epsilon\xi\omega$ , without;  $\epsilon\rho\eta\varsigma$ , creeper.)

### 2711. EXERPES ASPER (Jenkins & Evermann).

Head 3 ( $3\frac{3}{8}$  in total); depth  $5\frac{1}{2}$  ( $6\frac{1}{8}$ ); eye  $4\frac{1}{2}$  in head; scales 6-13-7, about 40 pores. D. III-XXV, 1; A. II, 20. Body compressed; head narrow, pointed; snout long, lower jaw slightly the longer; mouth a little oblique, cleft moderate, maxillary not reaching nearly to vertical at front of orbit. Teeth in 1 well-defined outer series and a broken inner one, those in the outer series strongest and of pretty uniform size, short and broad; vomerine teeth in a single patch; no palatine teeth. No tentacles of any kind about the head. Profile nearly straight from snout to origin of first dorsal, but very slightly arched from there to base of caudal fin. Scales rather large, cycloid, about 6 rows between origin of second dorsal and lateral line just behind its angle, and about 7 from there to mid-

dle of ventral surface; 9 rows from origin of second dorsal to upper limb of opercle; entire head, opercles, and fins naked. Lateral line beginning at upper limb of opercle on a level with the pupil, almost exactly under the middle of the first dorsal fin, and a little more than  $\frac{1}{2}$  the distance from top of nape to the under side of the throat, arching gently for 7 or 8 scales, leaving but 1 row of scales between it and the first spines of the second dorsal; on the ninth, tenth, and eleventh scales it bears slightly downward until 2 rows are left between it and the dorsal, then a sharp turn is made which puts it 4 scales further down, and from there it pursues a nearly direct line to middle of base of caudal fin. Dorsal fins separate, the first of 3 slender, very close-set, flexible spines, their length about twice in that of head, the fin ribbon-shaped; second dorsal separated from first by a distance somewhat greater than diameter of eye, and composed of 25 rather stout, sharp spines and 1 terminal soft ray; the first 3 are graduated, the first being contained  $1\frac{1}{2}$  times in distance between the 2 fins, the second is about  $\frac{1}{2}$  longer, and the third still a little longer; the remaining 22 are of approximately equal length, about equaling distance from origin of first dorsal to that of second; the 1 soft ray somewhat shorter than spines, well separated from caudal by a space equal to that between dorsals; pectorals inserted under middle of space separating dorsals, composed of 14 rays, equaling eye and snout in length, and reaching slightly past origin of anal; ventral of 2 rays inserted directly under origin of first dorsal and considerably in front of pectorals, which they somewhat exceed in length, in some specimens reaching vent; anal fin beginning slightly in front of posterior end of pectorals, a little lower than second dorsal and reaching a trifle nearer to caudal fin; first spine longer and more slender than the first regular dorsal spine, while the second equals the third dorsal in length. Caudal rounded, equaling in length the greatest depth of fish. Coloration in alcohol, pale, pretty regularly covered with very fine dark punctulations, thickest on back, palest below; a large dark opercular blotch, 2 similar postocular blotches, and usually a darkish bar extends downward from eye; upper half of preorbital region dark, outer margin of jaws dark; breast and under parts of head pale, top of head and nape dark; first dorsal quite dark, almost black; second dorsal pale, obscurely mottled with brown, which is disposed in about 5 indistinct areas; a large black ocellus upon the twelfth and thirteenth spines of second dorsal, and a similar one upon the twenty-third and twenty-fourth spines; each ocellus is surrounded by a narrow circle of white or pale orange. In the 6 specimens before us there is a slight variation as to the exact position of the 2 ocelli; in 1 example the second ocellus extends back upon the twenty-fifth spine also, but in every case the twelfth and thirteenth and the twenty-third and twenty-fourth are the spines which most evidently locate the spots; pectorals and ventrals plain; anal paler than dorsal, sparsely covered with fine dark points, so grouped as to form 3 or 4 darker areas. Length  $2\frac{1}{2}$  inches. Gulf of California. Known from 6 specimens taken from masses of kelp hauled out by the seine from the bay of Guaymas. (Jenkins & Evermann.) (*asper*, rough.)

*Auchenopterus asper*, JENKINS & EVERMANN, Proc. U. S. Nat. Mus. 1888, 164, Guaymas, Mexico. (Type, No. 39643. Coll. Jenkins & Evermann.)

Auch  
Oren

Coral

Bo

scale

mod

the v

unite

Dors

lowe

stiff

with

later

differ

the la

CORAL

a. F

AUCHE

aa. F

Head

snout 4

spines

879. AUCHENOPTERUS, Günther.

*Auchenopterus*, GÜNTHER, Cat., III, 275, 1861 (*monophthalmus*).

*Oreannobates*, GÜNTHER, Proc. Zool. Soc. Lond. 1861, 374 (*monophthalmus*). Substitute for *Auchenopterus*, regarded as preoccupied on account of its similarity to *Auchenipterus*, a genus of *Siluridae*.

*Corallicola*, JORDAN & EVERMANN, new subgenus (*marmoratus*).

Body moderately elongate, compressed, covered with rather large, cycloid scales; head shortish, naked, the snout rather pointed; cheeks full; mouth moderate, with a band of conical teeth in the jaws and about 1 series on the vomer, none on the palatines; lower jaw prominent; gill membranes united, free from the isthmus; upper surface of head with tentacles. Dorsal fin composed of stiff spines, with but a single soft ray, which is lower than the spines; first 3 spines more or less separated from the others, stiff and rather wider set, sometimes higher than the others; anal fin low, with 2 short spines; ventrals jugular, well developed; pectorals broad; lateral line complete, strongly curved anteriorly. Warm seas. This genus differs from *Cristiceps* in having but 1 soft ray in the dorsal fin, and in the large scales. (*αύχην*, nape; *πτερόν*, fin.)

CORALLICOLA (*Corallus*, coral; *colo*, I inhabit):

a. First 3 or 4 spines of dorsal forming a separate fin, being much higher than any of the spines in the posterior part of the fin; snout rather acute.

b. Scales 33; dorsal with 1 ocellus, anal with none; a black cross bar at base of caudal; a yellow spot behind eye; snout pointed. NIGRIPINNIS, 2712.

bb. Scales 27 or 38.

c. First dorsal spine longer than second; dorsal with 2 ocelli; anal blackish; D. IV-XXIV, 1. ALTIVELIS, 2713.

cc. First dorsal spine shorter than second; snout slender, very acute; caudal pale; dorsal with 2 ocelli, anal with 1; D. III-XXII, 1. MARMORATUS, 2714.

AUCHENOPTERUS:

aa. First 3 spines of dorsal scarcely forming a separate fin, none of them higher than the posterior spines; snout not very acute; anal without ocellus.

d. Caudal fin pale, usually with a dark bar at its base; a notch between third and fourth dorsal spines.

e. Dorsal spines about 31.

f. Scales 34 to 36; membrane of third spine joining fourth at its base; dorsal and anal plain dusky. AFFINIS, 2715.

ff. Scales 38.

g. Membrane of third spine joining fourth slightly above its base. MONOPHTHALMUS, 2716.

gg. Membrane of third spine joining fourth spine much above its base. INTEGRIPINNIS, 2717.

ee. Dorsal spines about 28; membrane of third spine joining fourth above its base; scales 38; body with distinct cross bars; dorsal with 1 ocellus. FASCIATUS, 2718.

dd. Caudal fin black; body chiefly black; head mottled with whitish; membrane of third dorsal fin joining fourth near its summit, the fin not notched; dorsal spines 30; dorsal with 2 ocelli. NOX, 2719.

Subgenus CORALLICOLA, Jordan & Evermann.

2712. AUCHENOPTERUS NIGRIPINNIS (Steindachner).

Head 4; depth 5½. D. XXVIII, 1; A. II, 27; scales 33; eye 4½ in head; snout 4½, equal to interorbital space; snout pointed. Three first dorsal spines higher than the others and further apart. A tentacle over eye.

Scales of body much largest anteriorly; lateral line arched. A deep black spot with a white ring between the twenty-second and twenty-fourth spines; anal edged with white; black cross band at base of caudal with silvery point at upper base of pectoral; a diffuse yellowish spot below and behind eye. Barbados. One specimen 1 inch and 7 lines long. (Steindachner); not seen by us. (*niger*, black; *pinnis*, fin.)

*Olinus nigripinnis*, STEINDACHNER, Ich. Notizen, VI, 46, 1867, Barbados.

2713. AUCHENOPTERUS ALTIVELIS (Lockington).

D. IV-XXIV, 1; A. 21; P. 13; C. 13; V. 2; scales 37. Body compressed, greatest depth a little behind pectoral axil; greatest thickness at gill covers; dorsal and abdominal profiles of similar curvature, decreasing regularly to the caudal fin; profile of occiput and superorbital regions convex; snout somewhat produced, its upper outline slightly concave. Head 4 in total length; greatest depth a little less than length of head; caudal peduncle about  $\frac{1}{2}$  of the greatest depth. Eye round, lateral, with a slight direction upward, its diameter less than the length of the snout; interorbital area nearly equal in width to the diameter of the eye, concave transversely, upper orbital borders slightly raised. A short nasal tentacle slightly anterior to the front margin of the eye; a large fimbriated tentacle on each side of the first dorsal ray. Cleft of mouth oblique, the lower jaw the longer; the posterior convex extremity of the club-shaped maxillary about vertical with the center of the pupil. Teeth of the outer row regular, sharp, incurved, the largest in front, gradually decreasing along the lateral portions of the jaws, and not extending much past the middle of their length; a narrow band of small teeth in the rear of the outer row; vomerine teeth present. Branchiostegals 6; gill openings continuous, membranes not attached to the isthmus. Distance from first ray of dorsal to posterior margin of eye equal to length of snout; first 2 rays of dorsal much developed, the first slightly the longer, and nearly equal in height to the distance of its base from the tip of the upper jaw; third ray about  $\frac{1}{2}$  the length of the first; fourth very short; succeeding rays to the twenty-sixth longer than the third, the last 3 somewhat decreasing. Anal commencing under eleventh dorsal ray, coterminous with, and equal in height to, the dorsal. Caudal with 13 simple jointed rays, the longest in the center, posterior margin convex. Pectorals narrow, lanceolate, the fifth and sixth rays longest and  $\frac{1}{2}$  the length of the head. Ventrals inserted in advance of the pectorals. Lateral line with 37 simple pores, parallel with dorsal outline to opposite the origin of the anal, where it is deflected almost perpendicularly downward to the middle of the side of the body, along which it continues to its termination. Scales rather large, about 10 in a transverse row in the central part of the body, their posterior margin membranaceous; no scales on fins; a line of pores around the margin of the orbit, another along the posterior margin of the preoperculum, connected with each other and with the lateral line by a line from the center of the hinder border of the eye. Color in alcohol, bright pink above, becoming dusky below; underside of head light olivaceous, lower lip

blackish; dorsal pink, dusky on its margin, a black spot on the fourth ray, and another on its hinder part upon the twenty-fourth and twenty-fifth rays, the latter spot extending on to the body; membrane of anal black; occipital tentacles black. La Paz, Lower California. A single specimen,  $1\frac{3}{10}$  inches long, dredged at a depth of 22 fathoms. (Lockington.) (*altus*, high; *velum*, sail.)

*Cremnobates altivelis*, LOCKINGTON, Proc. Ac. Nat. Sci. Phila. 1881, 116, La Paz, Lower California. (Coll. W. J. Fisher.)

2714. **AUCHENOPTERUS MARMORATUS** (Steindachner).

Head  $3\frac{3}{8}$  to  $3\frac{1}{2}$  in body; depth  $3\frac{1}{2}$  to  $3\frac{5}{8}$ . D. III-XXII, 1; A. II, 19; scales 2-36-9 (28 or 29 anteriorly); eye 4 to 5 in head; first dorsal  $1\frac{1}{4}$ ; pectoral  $1\frac{1}{4}$ . Body comparatively deep, compressed, the back somewhat arched; head pointed; mouth large, the maxillary extending to behind the eye, 2 in head; opercle with a sharp spine; jaws equal; teeth pointed, in narrow bands, the outer larger; vomerine teeth in 1 row; supraocular tentacle small, about as large as nuchal tentacle; no nasal tentacle. Pectoral a little shorter than head; dorsals separate, the first dorsal higher than second dorsal, the spines of which are about  $\frac{1}{2}$  head. Color in life of varying shades of olive gray or sand color, with a series of whitish blotches on head and along sides; markings on dorsal and anal whitish; 2 dark-blue ocelli on dorsal and 1 on anal, these edged with orange and interiorly with black; ventrals, pectorals, and caudal whitish, barred with clear orange red; first dorsal black at tip; a curved blackish line at base of caudal; lower side of head yellowish brown, with whitish bands; specimens from coral reefs more spotted. Florida Keys to Cuba; common in the eelgrass at Key West. Our specimens, 2 to  $2\frac{1}{2}$  inches long, taken at Key West and Havana. (*marmoratus*, marbled.)

*Cremnobates marmoratus*, STEINDACHNER, Ichth. Beiträge, v, 174, pl. 12, f. 6, 1876, a small rocky island north of Cuba; JORDAN & GILBERT, Synopsis, 962, 1883; JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1884, 142.

Subgenus **AUCHENOPTERUS**.

2715. **AUCHENOPTERUS AFFINIS** (Steindachner).

Head 4; depth  $4\frac{3}{8}$ . D. III-XXVII, 1; A. II, 19; V. I, 2; scales 33 to 35. Form of *A. integripinnis*; maxillary reaching to below posterior margin of eye; a fringed tentacle above eye and 1 on each side of occiput. First dorsal low, its longest (second) ray shorter than the highest of second dorsal; membrane of third spine joining the fourth spine just above its base; last ray of second dorsal joined by membrane to base of caudal. Dark brown, paler than in *A. nox*, but darker and more uniform than in *Auchenopterus fasciatus*; lower side of head pearly gray, thickly speckled with darker; sides with 5 very faint darker cross bands; dorsal and anal dusky, the latter with a pale edge; between the eighteenth and twenty-second dorsal spines a large dark spot ocellated with yellowish; caudal yellowish white, with darker cross streaks, a blackish band at its base; pectoral



dusky at base, its posterior half yellowish, with darker cross streaks; ventral similar; a wedge-shaped, whitish band extending backward from eye to opercle. West Indies; recorded from Key West and St. Thomas. Here described from specimens from Key West. (*affinis*, related,—to *A. monophthalmus*.)

*Cremnobates affinis*, STEINDACHNER, Ichth. Beiträge, v, 178, 1876, St. Thomas; JORDAN, Proc. U. S. Nat. Mus. 1884, 142; JORDAN, Cat. Fishes N. A., 121, 1885.

2716. *AUCHENOPTERUS MONOPHTHALMUS*, Günther.

Head  $3\frac{1}{2}$ ; depth 4. D. III-XXVI, 1; A. II, 18; scales 2-32-9; eye 5 in head; maxillary  $1\frac{1}{2}$ ; pectoral  $1\frac{1}{2}$ ; caudal  $1\frac{1}{2}$ . Body compressed, deepest at middle of pectorals; head moderately pointed, the upper profile slightly and evenly convex; mouth large, maxillary reaching past eye; jaws subequal; teeth villiform, in bands on jaws, vomer, and palatines; interorbital space flat, as wide as eye; a multifold dermal flap over posterior edge of eye, and a smaller one on each side of nape; head naked; body with rather large, regular scales; fins naked. Origin of dorsal over edge of preopercle, the first 3 spines separated from rest of fin by a rather deep notch, the membrane from third spine joining fourth spine at about its middle; spines of posterior part of dorsal the highest; front of anal midway between tip of snout and base of caudal, tips of last rays reaching slightly beyond base of caudal and tips of last dorsal rays; pectorals reaching front of anal; ventrals long and slender, inserted in front of base of pectorals a distance equal to  $1\frac{1}{2}$  eye; caudal rounded. Color light grayish red or brown, with about 6 cross bars of darker brown, running up on dorsal; between the bars are scattered milky white irregular spots; a black spot, ocellated with white, on front of dorsal, a similar spot near posterior end, sometimes duplicated; narrow cross bars on anal; a dark bar on base of caudal, and a dark blotch on base of pectoral. Here described from specimens, a couple of inches in length, from La Paz, Lower California. Gulf of California to Panama, abundant in rock pools, creeping about among *Corallina*; close to *A. integripinnis*, but the first dorsal higher and more separate from rest of fin. (*μόνος*, one; *ὀφθαλμός*, eye, from the dorsal ocellus.)

*Auchenopterus monophthalmus*, GÜNTHER, Cat. Fish., III, 275, 1861, Panama; JORDAN, Proc. Cal. Ac. Sci. 1895, 501.

*Cremnobates monophthalmus*, GÜNTHER, Proc. Zool. Soc. Lond. 1861, 374; GÜNTHER, Fish. Centr. Amer., 4, 2, pl. 69, fig. 1, 1869.

2717. *AUCHENOPTERUS INTEGRIPINNIS* (Rosa Smith).

Head  $3\frac{1}{2}$ ; depth  $4\frac{1}{2}$ ; eye 4 in head. D. III-XXVII, 1; A. II, 20; scales 2-36-9; pectoral  $1\frac{1}{2}$ ; caudal  $1\frac{1}{2}$ . Head stout, broad, conical; mouth little oblique, maxillary reaching posterior margin of eye; eyes large; nasal, supraocular and nuchal regions with fringed cirri, those at the nape flap-like. First and second dorsal spines low, a little higher than the third, which, in turn, is higher than the fourth and separated from it by an interspace, the membrane between the third and fourth spines not deeply

emarginate, membrane from third spine attached to the lower  $\frac{2}{3}$  of fourth; anterior spines not forming a separate fin; highest anterior spine not higher than the highest of the posterior part of fin. Color dark brown, variegated with different shades of brown and reddish; about 5 indistinct dark cross bars; a distinct ocellated black spot on posterior part of dorsal fin; caudal fin abruptly translucent, speckled, a black bar at its base; base of pectorals violet, bordered with black, the rest of the fin checkered; ventrals barred. Length  $2\frac{1}{2}$  inches. Coast of California and southward to Todos Santos; abundant in rock pools among *Corallina*. Here described from a specimen,  $1\frac{1}{2}$  inches in length, from San Cristobal, Lower California. (*integer*, entire; *pinna*, fin.)

*Oremonobates integripinna*, ROSA SMITH, Proc. U. S. Nat. Mus. 1880, 147, La Jolla, near San Diego (Coll. Rosa Smith); JORDAN & GILBERT, Synopsis, 764, 1883.

2718. *AUCHENOPTERUS FASCIATUS* (Steindachner).

Head 4; depth  $4\frac{1}{2}$ . D. III-XXIV, 1; A. II, 18; scales 37. Body rather slender, a little deeper than in *A. integripinna*, the snout less acute than in *A. marmoratus*. First dorsal spine rather higher than second and lower than the spines of posterior part of fin; membrane of third spine joining second dorsal at a point above its base, the two parts of the fin therefore separated only by an emargination. Tentacle above eye slender, small; cirri on side of occiput bluish. In life, light pinkish brown, much mottled, and with traces of 6 to 8 faint darker bars; head and its cirri above whitish; 3 blackish spots behind eye, radiating from it, the lower one largest; preopercle with 3 dark dots; dorsal pale, with 9 blackish blotches, in the next to the last of which is a large blue-black ocellus, edged with orange; anal with 5 dark blotches and no ocellus; a blackish bar across base of caudal; rest of caudal and pale part of anal with dark dots; ventrals whitish, barred with black; pectoral similar, its base with a whitish area, which has a brown center, below which is a small black spot. Length 2 inches. Florida Straits; north to Key West. Here described from specimens from Key West. (*fasciatus*, banded.)

*Oremonobates fasciatus*, STEINDACHNER, Ichth. Beiträge, v, 176, 1876, Florida Straits; JORDAN, Proc. U. S. Nat. Mus. 1884, 142; JORDAN, Cat. Fishes N. A., 121, 1885.

2719. *AUCHENOPTERUS NOX* (Jordan & Gilbert).

Head  $3\frac{1}{2}$ ; depth  $3\frac{1}{2}$ . D. II-XXVII; A. II, 18; lateral line with 34 tubes. Snout not very acute, the upper and lower profiles of head nearly evenly convex; mouth large, maxillary reaching slightly beyond eye,  $\frac{1}{2}$  length of head; eye large, equaling length of snout, greater than interorbital width, 4 in head (to end of opercular spine); interorbital width  $4\frac{1}{2}$  in head; nasal, supraorbital, and occipital tentacles present, those on snout and above the orbits simple, slender filaments, the latter about as long as diameter of orbit, 1 of them divided to the base, the other simple; the tentacle on each side of nape a compressed slip of skin higher than wide, the margin uneven, but not fringed. Anterior dorsal spines not much elevated,

not higher than some of the posterior spines; the first and second spines about equal,  $2\frac{1}{2}$  in head; the third spine shorter, about equal in length to the fourth, from which it is separated by a wide membrane, which is, however, not at all notched; the spines thence increase in length toward the last; caudal  $1\frac{1}{2}$  in head; pectorals reaching anal, nearly equaling length of head; ventrals not reaching vent,  $1\frac{1}{2}$  in head. Scales large, 4 series above lateral line and 4 below. Color, body and fins uniform blackish brown; a few small silvery-white specks on dorsal region, mostly along base of dorsal fin; head and base of pectoral fin with light pink areas and mottlings; snout pink above; nape with a pink cross bar; a dark streak upward and backward from eye to nape; a light streak from eye backward to opercle and 1 backward and downward; lower jaw mottled with light and dark; a small round, black spot near base of dorsal between twenty-third and twenty-fifth spines, and 1 between twenty-eighth and thirtieth, both very faintly ocellated with lighter; slight whitish tips on ventrals and lower edge of caudal. Key West; known from a single specimen,  $1\frac{1}{2}$  inches long, taken with the seine in algae on a rocky bottom at Key West. Its congeners, *A. marmoratus*, *A. fasciatus*, and *A. affinis*, were found in the same waters, *A. marmoratus* being much the most abundant of the 4, and reaching the largest size. (*nox*, night.)

*Cremnobates nox*, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1884, 30, Key West. (Coll. Jordan.)

#### 880. PARACLINUS, Mocquard.

*Acanthoclinus*, MOCQUARD, Bull. Soc. Philom. Paris 1885, 18 (*chaperi*); name preoccupied.  
*Paraclinus*, MOCQUARD, Bull. Soc. Philom. Paris 1886, 11 (*chaperi*).

Body elongate, compressed, covered with cycloid scales; mouth large, each jaw armed with an external row of conical teeth, with some teeth behind; teeth on the palate; dorsal very long, continuous, composed entirely of spines, anal with 2 spines; ventrals jugular, with few rays; tentacles on head; gill opening very broad; 6 branchiostegals; lateral line interrupted. Evidently very close to *Auchenopterus*, from which it may be distinguished by the continuous dorsal fin, a character which needs verification. (*παρά*, near; *Clinus*.)

#### 2720. PARACLINUS CHAPERI, Mocquard.

Head  $4\frac{1}{2}$ ; depth  $4\frac{1}{2}$ . D. XXXI; A. II, 19; P. 13; V. 2; scales 35. Body elongate, very strongly compressed; eye large, equal to snout or inter-orbital width; lower jaw slightly the longer; mouth oblique, reaching front of eye; outer row of teeth strong, canine-like, slender and more close set above, below diminishing rapidly in length, the bands of small teeth limited to front of each jaw, a curved group of teeth on palate; dorsal beginning over preopercle, not notched, composed entirely of stout spines; anal equally long; ventrals very narrow, of 2 soft rays, well separated, the inner slightly longer than outer; head with 3 pairs of tentacles, 1 at the nape, filiform, small,  $\frac{1}{2}$  as long as eye; the second below the orbit, broadened at base, separated into 3 or 4 branches, progressively

long  
in fr  
its fr  
rupte  
rows  
bran  
brow  
in Ve  
Not s

Acanth  
zu  
Paract

Emmm

Bod  
eral li  
decurv  
bands,  
its ant  
thickis  
agos la

Head  
3-50-11  
the dor  
ray, fro  
about s  
length;  
thence  
jaw inc  
about 2  
with 8 e  
behind  
mouth;  
greatly  
front te  
tractile  
3 $\frac{1}{2}$  in he  
equal.  
united,  
cirri ab  
cycloid s  
the body  
very dist

longer from the inner outward, longer than eye; nuchal filament a little in front of dorsal, in form, oblong, entire, laminated, a little broader at its free edge,  $\frac{3}{4}$  as long as eye; scales large, cycloid; lateral line interrupted before front of anal, anterior part rounding over eye with only 2 rows of scales between it and the dorsal, posterior part median; gill membranes broadly united, free from isthmus. Body brownish yellow, fins brown, the base and the caudal darker. Bay of Guanta, near Barcelona, in Venezuela; 1 specimen, 33 mm. long to base of caudal. (Mocquard.) Not seen by us. (Named for its collector, M. Chaper.)

*Acanthoclinus chaperi*, MOCQUARD, Bull. Soc. Philom. Paris 1885, 19, Bay of Guanta, Venezuela.

*Paraclinus chaperi*, MOCQUARD, Bull. Soc. Philom. Paris 1886, 41.

881. EMMNION, Jordan.

*Emmnion*, JORDAN, in Gilbert. Proc. U. S. Nat. Mus. 1896, 454 (*bristolae*).

Body elongate, covered with caducous, cycloid scales of small size; lateral line straight, ending near base of last dorsal ray. Head moderate, decurved anteriorly, without cirri; mouth moderate; teeth in jaws in bands, the outer enlarged; no teeth on vomer or palatines; dorsal notched, its anterior  $\frac{3}{4}$  of flexible spines of moderate height; ventrals I, 3, the rays thickish, the fin inserted slightly before pectorals; caudal free. Galapagos Islands. ( $\epsilon\gamma$ , in;  $\mu\nu\iota\omicron\nu$ , sea moss, or alga.)

2721. EMMNION BRISTOLE, Jordan.

Head  $5\frac{1}{2}$ ; depth  $7\frac{1}{2}$ . D. XXV, 13; A. I, 27; P. 13; V. I, 3; Br. 5; scales 3-50-11, the count not certain. Body slender, moderately compressed, the dorsal profile forming a nearly straight line from occiput to first dorsal ray, from thence descending very gently to base of caudal; ventral profile about straight. Head broad, slightly convex above, its width  $1\frac{1}{4}$  in its length; anterior profile from first dorsal spine to a point above eye straight, thence abruptly descending to tip of snout; mouth horizontal, the lower jaw included; maxillary reaching nearly to posterior margin of eye, about  $2\frac{1}{2}$  in head. Teeth present on both jaws, canine-like; upper jaw with 8 enlarged teeth in front, about 2 or 3 series of much smaller teeth behind these, only 1 series of which extends into posterior region of mouth; lower jaw with a series of teeth in front and on sides which is greatly enlarged in front; a patch of very small teeth behind the enlarged front teeth; no teeth on vomer or palatines. Premaxillary very protractile; snout blunt,  $4\frac{1}{2}$  in head; eyes large, round, placed close together,  $3\frac{1}{2}$  in head; interorbital region very narrow, less than pupil; nostrils equal. Caudal peduncle  $2\frac{1}{2}$  in head; branchiostegal membranes deeply united, free from isthmus; gills 4, a small slit behind the fourth; no cirri above eyes, nor filaments on nape; head naked, body covered with cycloid scales, those on nape much smaller; belly naked. The scales on the body are apparently caducous as all have fallen, but the points are very distinct and they seem to have been embedded on their anterior edge,

as the sac-like fold of skin is prominent. Lateral line simple, straight, running from upper edge of gill opening to last ray of dorsal when it is lost, not reaching the caudal; it is placed very high, and gradually approaches the dorsal fin, from which it is separated only by a very small distance. Dorsal extending from a point a short distance behind occiput nearly to base of caudal, emarginate; last spine shortest, about  $2\frac{1}{2}$  in first soft ray, which is  $2\frac{1}{2}$  in head; the longest spines about 3 in head, all the spines slender and flexible. Anal extending from behind vent nearly to base of caudal; similar to soft dorsal, its rays lower. Ventrals well developed with broad base, the rays thickish, inserted very slightly in front of base of pectorals,  $1\frac{1}{2}$  in head, reaching  $\frac{2}{3}$  the distance to vent. Caudal subtruncate. Pectorals reaching past vent, about as long as head. Dorsal and anal free from caudal. Color in spirits, dark dull reddish-brown, lighter below; head very dark; dorsals, pectorals, and caudal blackish, pectorals and caudal with lighter blotches; anal and ventrals dusky, anal margined with darker. Length about 3 inches. Galapagos Islands; one specimen known, evidently a rock-pool species. (Named for Miss Susan Brown Bristol, of the department of zoology in Leland Stanford Junior University.)

*Emmation bristolae*, JORDAN, in GILBERT, Proc. U. S. Nat. Mus. 1896, 454, pl. 55, fig. 1, Galapagos Islands. (Coll. Albatross.)

#### 882. ATOPOCLINUS, Vaillant.

*Atopoclinus*, VAILLANT, Bull. Sci. Philom. Paris, serie 8, tome vi, 1894, 73 (*ringens*).

Body elongate, subcylindrical, without visible scales. Head obtuse, the snout short, rounded; mouth inferior, transverse, with compressed trenchant teeth in each jaw, those above at least in a single row, solidly fastened to the skeleton; teeth on vomer and palatines uncertain. Dorsal continuous, extending the whole length of the back, from the nape to the caudal peduncle, its rays mostly simple, only the posterior articulate; anal occupying nearly  $\frac{1}{2}$  the length, touching the caudal, which is, nevertheless, distinct; caudal deeply forked; ventrals distinctly jugular, very long, of a spine and a ray; no tentacles; gill membranes apparently rounded at the isthmus. Gulf of California; a singular genus evidently closely allied to *Runula*. (*ἄτοπος*, strange; *Clinus*.)

#### 2722. ATOPOCLINUS RINGENS, Vaillant.

Head 5; depth 7. D. 24; A. 18; P. 15; V. I, 1. Eye large, 7 in head; interorbital space broad, 3 in head. Caudal a little longer than head. Color clear chamois brown, the belly pale; a brown band before the snout, across the eye to the caudal, on which it extends; a silvery stripe bordering this band above, and below for part of its length. Gulf of California. (Vaillant); known from 1 specimen badly shriveled, 39 mm. in length. (*ringens*, gaping.)

*Atopoclinus ringens*, VAILLANT, Bull. Sci. Philom. Paris, serie 8, tome vi, No. 2, February 25, 1894, 74, Gulf of California. (Coll. Léon Diguët.)

883. RUNULA, Jordan & Bollman.

*Runula*, JORDAN & BOLLMAN, Proc. U. S. Nat. Mus. 1889, 171 (*azalea*).

Body slender, its back not elevated; mouth small, inferior, destitute of canines; teeth fixed, upper largest; dorsal fin continuous, its spines and soft rays indistinguishable, most of them articulate; caudal fin lunate; gill openings reduced to a vertical slit in front of pectoral; scales none. This genus is remotely allied to the East Indian genus *Petroskirtes*, but has the mouth and dentition different, and the caudal fin, unlike that of most blennioid fishes, is forked. (Diminutive of *runa*, a dart or javelin.)

2723. RUNULA AZALEA, Jordan & Bollman.

Head  $4\frac{1}{2}$ ; depth  $6\frac{1}{2}$ . D. 42; A. 26 or 27; V. I, 2. Body moderately elongate, not much compressed; head rather long, its upper outlines convex; snout short and very blunt; mouth entirely inferior, transverse, each jaw provided with long, slender, close-set curved teeth; no evident posterior canines; eye moderate, equal to snout and nearly equal to inter-orbital width, 4 in head; no tentacles on head; gill membranes fully united to the isthmus, the gill opening reduced to a vertical slit, its lower edge opposite middle of base of pectoral; no scales; lateral line very high, concurrent with the back; dorsal fin very low, continuous; the feeble spines and soft rays indistinguishable, the fin beginning at occiput; anal similar to soft dorsal; caudal lunate behind, well separated from dorsal and anal; pectorals small, rounded, about  $1\frac{1}{2}$  in head; ventrals short, before pectorals, about 2 in head. Color reddish brown, silvery below, about 5 dusky cross shades; a dusky lateral streak; a black spot surrounded by paler at base of caudal; dorsal with about 6 black cross-bars; anal with 4; other fins pale; lower half of head abruptly pale. Galapagos Archipelago. The type, 2 inches long, taken at Indefatigable Island; 3 more specimens have since been obtained from the same island. (*ἀζαλέος*, parched, from the brown color.)

*Runula azalea*, JORDAN & BOLLMAN, Proc. U. S. Nat. Mus. 1889, 171, Indefatigable Island, Galapagos Archipelago (Coll. Albatross); JORDAN, Proc. Cal. Ac. Sci. 1896, 233, pl. 37.

884. BLENNIUS (Artedi) Linnaeus.

(BLENNIES.)

*Blennius*, ARTEDI, Genera Piscium, 27, 1738.

*Blennius*, LINNÆUS, Syst. Nat., Ed. x, 256, 1758 (*galerita*).

*Salaria*, FORSKÅL, Descr. Anim., 22, 1777 (*basiliæus*).

*Pholis*, FLEMING, Brit. Anim., 207, 1828 (*lævis* = *pholis*); not *Pholis* SCOPOLI, 1777.

*Adonis*, CRONOW, Cat. Fish., Ed. Gray, 93, 1754 (*pavoninus* = *ocellaris*).

*Lipophrys*, GILL, American Naturalist, June, 1896, 498 (*pholis*).

Body oblong, compressed, naked; head short, the profile usually bluntly rounded; mouth small, horizontal, with a single series of long, slender, curved, close-set teeth in each jaw, besides which, in the lower jaw at least, is a rather short and stout fang-like canine tooth on each side;

premaxillaries not protractile; gill openings wide, extending forward below, the membranes free from the isthmus, or at least forming a broad fold across it. Dorsal fin entire, or more or less emarginate, the spines slender; pectorals moderate; ventrals well developed, 1, 3; no pyloric caeca; lateral line developed anteriorly. Species numerous, lurking under rocks and algae in most warm seas; some species in the lakes of northern Italy. The European species in general are larger in size than ours, with higher fins. (*Blennius*, the ancient name, from *βλέννα*, slime.)

**LIPOPHRYS** (*ἀέτω*, to disappear; *ὄφρυς*, eye-brow):

a. Supraorbital cirrus wanting; snout not very blunt in profile.

b. Posterior canine present in each jaw; dorsal slightly emarginate; D. XII, 18.

CAROLINUS, 2724.

**BLENNIUS**:

aa. Supraorbital cirrus present; profile of snout more or less blunt.

c. Canines strong, present in both jaws; no nuchal cirri.

d. Dorsal rays XI or XII, 17 or 18.

e. Supraorbital cirrus bifid; dorsal free from caudal.

f. Supraorbital cirrus as long as head; dorsal emarginate; sides spotted; D. XI, 17.

FUCORUM, 2725.

ff. Supraorbital cirrus as long as eye and snout; dorsal continuous; color olivaceous, with dark bars; D. XI, 18.

STEARNSI, 2726.

ee. Supraorbital cirrus bifid, nearly as long as head; last ray of dorsal joined to caudal; sides with a network of blue lines; D. XII, 18.

FAVOSUS, 2727.

dd. Dorsal rays XII, 21 or 22; supraorbital cirrus long, fringed; dorsal free from caudal; cheeks with network of lines; body nearly plain.

PILICORNIS, 2728.

cc. Canines short and stoutish, present in lower jaw only (undescribed in *truncatus* and in *marmoreus*.)

g. Nape without cirrus; snout abruptly decurved; body robust, marbled; D. XII, 20.

MARMOREUS, 2729.

gg. Nape with a cirrus on each side.

h. Dorsal and anal free from caudal. Nape with a filiform bifid tentacle on each side; teeth undescribed; supraorbital tentacle simple; color olive, with bright spots.

TRUNCATUS, 2730.

hh. Dorsal and anal with the last ray largely joined by membrane to caudal; nape with a small cirrus; posterior canines strong, in lower jaw only; dorsal not notched; color uniform brown; D. XII, 13.

VINCTUS, 2731.

ggg. Nape with a comb of many close-set cirri on a fleshy crest; lower jaw only with short posterior canines; dorsal fin continuous, free from caudal; D. XII, 16 or 17.

CRISTATUS, 2732.

**Subgenus LIPOPHRYS, Gill.**

**2724. BLENNIUS CAROLINUS** (Cuvier & Valenciennes).

D. XII, 18; A. 17. Body rather long and slender, more elongate than in the European species, *Blennius pholis*, more compressed, the head longer; maxillary extending to opposite middle of eye; teeth  $\frac{1}{4}$ , with strong canines on both jaws; gill membranes free from isthmus; no trace of tentacles above eye; dorsal spines slender, a little lower than the soft

ray  
Gr  
do  
lin  
the  
the  
Pho  
Ble

H  
as  
h  
gin  
stee  
very  
Oliv  
chee  
spot  
with  
type  
New

Blenn  
of  
F  
Blenn  
N

He  
21.  
profil  
the j  
times  
a spa  
side  
canin  
space  
a lon  
tip of  
brane  
acros  
ous h  
portio

\* Ven  
dorsal  
body a

rays, the fin little emarginate; dorsal and anal not joined to the caudal. Greenish, with 4 or 5 irregular dark spots or shades along the back; dorsal with a large black spot in front; anal brown-edged. South Carolina. Only the original type in the museum at Paris known; from this the present description was taken. No later collector has recognized the species and it may not be American.

*Pholis carolinus*, CUVIER & VALENCIENNES, Hist. Nat. Poiss., xi, 276, 1836, Carolina. (Coll. M. Bosc.)

*Blennius carolinus*, JORDAN & GILBERT, Synopsis, 760, 1883.

2725. *BLENNIUS FUCORUM*, Cuvier & Valenciennes.

Head 5 in total length. D. XI, 17; A. 18. Orbital cirri nearly as long as head, bifid at tip, and fringed at the base. Dorsal fin slightly emarginate, free from the caudal, the spines rather stiff. Head very short and steep, its profile nearly vertical; 24 teeth in each jaw; each jaw with very strong canines; gill membranes free from the isthmus posteriorly. Olive green, becoming darker above, with numerous brown spots on the cheeks and sides of the body; below reddish; dorsal with a large black spot in front, behind which are smaller spots; spinous dorsal edged with paler. (Cuvier & Valenciennes.) Open ocean in floating *Fucus*; the type from near the Azores; recorded by De Kay from the open sea, off New York, in floating seaweed. (*fucorum*, of the seaweed, *Fucus*.)

*Blennius fucorum*, CUVIER & VALENCIENNES, Hist. Nat. Poiss., xi, 263, 1836, 240 miles south of the Azores (Coll. Claude Gay); GÜNTHER, Cat., iii, 217, 1861; DE KAY, N. Y. Fauna: Fishes, 149, pl. 22, fig. 66, 1842; JORDAN & GILBERT, Synopsis, 710, 1883.

*Blennius oceanicus*,\* CUVIER & VALENCIENNES, Hist. Nat. Poiss., xi, 265, 1836, open sea, 29° N., 50° W.; on a drawing by CLAUDE GAY.

2726. *BLENNIUS STEARNSI*, Jordan & Gilbert.

Head  $3\frac{1}{2}$  ( $4\frac{1}{2}$  in total); depth  $4\frac{1}{2}$  ( $5\frac{1}{2}$ ); eye  $4\frac{1}{2}$ ; snout  $4\frac{1}{2}$ . D. XI, 18; A. II, 21. Body much elongate, compressed, tapering regularly behind; anterior profile moderately decurved; snout short and blunt; mouth large, oblique, the jaws even; maxillary reaching slightly beyond middle of orbit,  $2\frac{1}{2}$  times in head; teeth in the front of the jaws only, occupying on each side a space equal to  $\frac{1}{2}$  length of maxillary; teeth  $\frac{3}{4}$ , the lateral one on each side much enlarged and canine-like, rather short but strongly curved; canine in upper jaw equaling about  $\frac{1}{2}$  diameter of pupil; interorbital space very narrow, not as wide as pupil; upper posterior rim of orbit with a long slender filament, forked at base, its length equaling distance from tip of snout to posterior rim of orbit; no filaments at the nape; gill membranes somewhat united to the isthmus in front, but forming a broad fold across it posteriorly, the gill openings of the two sides therefore continuous below. Dorsals rather high; no notch between the spines and soft portion, the membrane of last ray not reaching base of caudal; spines of

\* Very near *Blennius fucorum*, the profile more oblique, the cirri shorter, the spinous dorsal lower, the caudal more truncate; anal shorter. Color brown with brown spots on body and fins; sides clear green; belly silvery. Length 2 inches. (Cuvier & Valenciennes.)



nearly uniform height, all very slender and flexible, the tips almost filamentous; highest spine  $\frac{1}{2}$  length of head; highest soft ray  $1\frac{1}{2}$  in head; anal lower than dorsal, its longest ray very slightly less than  $\frac{1}{2}$  length of head; length of caudal peduncle more than  $\frac{1}{2}$  its height, about equaling the diameter of orbit; caudal about equaling pectoral,  $1\frac{1}{2}$  in head; ventrals long, the inner ray much the longest,  $1\frac{1}{2}$  in head, not quite reaching vent. Color light greenish olive, somewhat mottled; sides with irregular dark bars formed of spots, these extending on the fin; skin everywhere finely punctate; dorsal dark olive, the spinous part darker at tip; anal blackish, with paler edge; ventrals dusky; pectorals and caudal olive. Gulf of Mexico, in deep water. Three specimens known, the largest 3 inches long, taken from the stomach of a Red Snapper, at Pensacola. (Named for Silas Stearns.)

*Blennius stearnsi*, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1882, 300, Pensacola Snapper Banks. (Type, No. 29669, U. S. Nat. Mus. Coll. Jordan & Stearns.)

2727. *BLENNIUS FAVOSUS*, Goode & Bean.

Head  $3\frac{1}{2}$ ; depth  $4\frac{1}{2}$ . D. XII, 18; A. II, 20. Body comparatively elongate and compressed; anterior profile moderately decurved; head nearly  $\frac{1}{2}$  longer than deep; snout very short and blunt; mouth large, horizontal; jaws even; maxillary reaching posterior margin of orbit, its length  $2\frac{1}{2}$  in head. Each jaw with a long, curved, posterior canine; the canines of lower jaw largest. Preorbital  $\frac{2}{3}$  diameter of eye, which is  $3\frac{1}{2}$  in head, and equals more than twice interorbital width. An extremely long and slender supraocular cirrus, trifid to the base, the longest branch nearly as long as the head; no nuchal cirri. Gill membranes forming a rather narrow fold across the isthmus. Dorsal low, continuous, the spines very slender and flexible, the longest  $\frac{1}{2}$  as long as the head; the longest soft ray  $\frac{3}{4}$  as long as head; the last ray slightly joined to base of caudal; caudal  $\frac{3}{4}$  as long as head; anal rather high; pectorals  $\frac{1}{2}$  as long as head; only the straight part of lateral line developed. Color faded, brownish, finely reticulated, a series of obscure bluish blotches along the sides; front and sides of head marked with very distinct blue, reticulating lines surrounding honeycomb-like hexagonal interspaces; top of head with many small blue spots; dorsal with black dots and streaks; a black spot bordered with whitish between the first and second dorsal spines; anal with oblique blue streaks, the fin margined with dusky, tips of rays whitish; base of pectorals with blue reticulations. The whole body was probably reticulated with blue in life. Gulf of Mexico. Known from 2 specimens collected at Garden Key, Florida, by Gustav Würdemann; they are  $3\frac{1}{2}$  inches and 3 inches long, respectively. (*favosus*, honeycombed.)

*Blennius favosus*, GOODE & BEAN, Proc. U. S. Nat. Mus. 1882, 416, Garden Key, Florida (Type, No. 2629, U. S. Nat. Mus. Coll. Gustav Würdemann); JORDAN & GILBERT, Synopsis, 961, 1883.

2728. *BLENNIUS PILICORNIS*, Cuvier & Valenciennes.

Head  $4\frac{1}{2}$  with caudal; depth  $5\frac{1}{2}$ . D. XII, 21 or 22; A. 23 or 24. Snout obtuse, the upper profile very oblique. A strong curved canine in each jaw. Orbital tentacle filiform, with several smaller ones at base. Inter-

orbital space flat, its width  $\frac{1}{2}$  vertical diameter of eye; no groove or crest on the neck. Dorsal slightly notched, the spines flexible; caudal separate. Brown, dorsal and caudal spotted with darker. Length 5 to 6 inches. (Günther.) Coast of Brazil north to the West Indies, recorded from Rio Janeiro, Bahia, and the Tortugas, and off the coast of Florida. Mr. Garman gives the following color note on Tortugas specimens, collected by Prof. C. C. Nutting: Small, hexagonal reticulations on cheeks, resembling scales; anal darker toward ends of rays, the tips white; dorsal darker in outer half; basal part of dorsal and anal pale, sides with a few scattered black dots; median rays of caudal longer, the outer margin dark; caudal, pectorals, and ventrals paler than dorsal. (*pilicornis*, with downy horns.)

*Blennius pilicornis*, CUVIER & VALENCIENNES, Hist. Nat. Poiss., XI, 254, 1836, Rio Janeiro (Coll. Delalande and Gay); CASTELNAU, Anim. Nonv., etc., Amer. Snd, 25, 1855; GARMAN, Bull. Iowa Lab. Nat. Sci. 1896, 89.

*Blennius pilicornis*, GÜNTHER, Cat., III, 216, 1861. (Coll. M. Parzudaki.)

2729. *BLENNIUS MARMOREUS*, Poey.

Head  $4\frac{1}{2}$  in total length with caudal; depth 5. D. XII, 20; A. 16; P. 13. Eye very high, near the profile, twice length of snout. Snout round, falling off abruptly, but less so than in *Blennius truncatus*; posterior nostrils with a distinct tube; superciliary tentacle divided into 3 branches; no cilia at the nape. Teeth undescribed. Gill membranes not described. Pectoral and caudal round; dorsal low, the median spines highest, the soft rays a little higher, the difference slight. Color yellowish brown, darker medially, paler below; under the lens covered with small dots; fins below yellowish. This species differs from *Scartella microstoma* in the stout trunk, the more blunt head, the cilia on the head and in the tube of the nostril. Cuba; 1 specimen 2 inches long. (Poey); not seen by us; perhaps not a *Blennius* as here understood. (*marmoreus*, marbled.)

*Blennius marmoreus*, POEY, Enumeratio, 130, 1875, Cuba. (Coll. Rafael Arango.)

2730. *BLENNIUS TRUNCATUS* (Poey).

Head  $5\frac{1}{2}$  in total length with caudal; depth  $5\frac{1}{2}$ . D. XII, 19; A. I, 20. Eyes placed very high, profile before them vertical, suggesting the forehead of a bull without horns; mouth small, maxillary reaching below posterior border of eye; anterior nostril divided into 5 at tip; 2 filiform tentacles with a common base on each side of nape; a simple tentacle behind eye; some pores on the head, which is compressed; teeth undescribed; gill membranes undescribed; gill membranes united and free from isthmus; dorsal notched medially; caudal truncate, with 2 faint angles; lateral line long, reaching beyond the point of the pectorals. Color olive, with some bright spots on trunk; the vertical fins darker. Cuba; 1 specimen  $3\frac{1}{2}$  inches long. (Poey); not seen by us; perhaps not a species of *Blennius*. (*truncatus*, cut off short.)

*Blennius truncatus*, POEY, Memorias, II, 424, 1861, Cuba. (Coll. Poey.)

2731. *BLENNIUS VINCTUS*, Poey.

Head  $3\frac{1}{2}$  to base of caudal; depth 4. D. XII, 13; A. I, 8; V. 3. Eye high, 4 in head, as long as snout. Anterior nostril in a short tube. Jaws equal; 4 pores on the side of the lower jaw; 1 on the opercle; 4 on the suborbital; 4 below eye. A long tentacle above eye; another very small one on the nape. Maxillary reaching to below front of pupil. Teeth large, not pointed, compressed, in 1 series of 10 to 12 on each side of each jaw, feeble, somewhat moveable; gill membranes united, free from isthmus. Dorsal elevated backward, connected by a membrane to the first third of the caudal, as is also the anal, twenty second ray highest, its height  $\frac{1}{2}$  depth of body and double length of the dorsal ray above tip of pectoral; anal similar,  $\frac{1}{2}$  also of the rays of the dorsal and anal simple; the spines flexible, differing from the others in not being articulate; pectoral pointed, its middle rays longest, and also more robust, all simple; ventral not very short; caudal rounded. Lateral line forming a curve anteriorly. Color uniform brown. Cuba. (*vinctus*, bound.)

*Blennius vinctus*, POEY, Repertorio, 243, 1867, Havana. (Coli. Felipe Poey. Type\* No. 12647, Mus. Comp. Zool.)

2732. *BLENNIUS CRISTATUS*, Linnaeus.

Head 4; depth 4. D. XI, 16; A. 19; maxillary 3. Body moderately elongate, compressed; the head very blunt and deep, almost as deep as long, its anterior profile straight or slightly concave, and nearly vertical. Mouth moderate, the maxillary reaching to past front of eye; lower jaw with 2 short stoutish posterior canines, scarcely longer than the front teeth; upper jaw without canines. Teeth about  $\frac{3}{2}$ . Preorbital deep, its depth equal to diameter of eye and contained  $4\frac{1}{2}$  times in length of head. Interorbital space flat, narrow,  $\frac{2}{3}$  width of eye. Supraocular cirri small, fringed, their length about equal to that of pupil. Nape with a longitudinal dermal crest reaching to front of dorsal, provided with a series of about 20 filaments, the longest about as long as the eye. Gill membranes forming a broad fold across the isthmus, as in all species of *Blennius*. Dorsal nearly continuous, the last spine a little lower than the first soft ray, not very high, beginning on the nape in front of the vertical of the preopercle, the spines all slender and flexible, the longest  $\frac{3}{4}$  as long as the head, the longest soft ray  $\frac{1}{2}$  as long as the head; caudal free from dorsal and anal,  $\frac{1}{2}$  as long as head; anal moderate,  $\frac{2}{3}$  length of head; pectoral somewhat shorter than head; ventral a little more than  $\frac{1}{2}$  length of head. Lateral line forming the usual arch above pectoral, and continued backward on median line to base of caudal, becoming indistinct posteriorly.

\* On the type of *Blennius vinctus* we have the following notes: "No. 12647, M. C. Z. Cuba. (Poey.) One and a half inches long, in poor condition. Head ca  $3\frac{1}{2}$ ; depth ca 4. D. XII, 13; A. II, 13. Dorsal joined to caudal as far as tips of the rays, which are high. Dorsal spines high and stiff, the fin not notched, the soft rays higher. A thick scale-like fringed cirrus above each eye, nearly as long as eye, which is small. Gill membrane free. Head blunt. Maxillary to front of pupil. Lower jaw with very strong canines; upper jaw with none. No nuchal cirri."

Color faded, apparently olivaceous, with about 6 dark cross bars, which extend on the dorsal fin; anal and posterior  $\frac{1}{4}$  of body with numerous round, whitish, stellate spots, probably bluish in life; bluish streaks from eye across the cheeks; anal edged with dusky; the other fins vaguely marked. Length  $2\frac{1}{2}$  to 4 inches. Tropical parts of the Atlantic, among rocks, widely diffused and variable. The above description from the type of *Blennius asterias*, from Garden Key, Florida. We have the following notes on numerous specimens from Abrolhos Islands, off the Coast of Brazil (Coll. Albatross): D. XII, 15. Nape with a fringed crest of 10 to 18 filaments. A small trifold tentacle above eye; posterior canines in lower jaw only, short and small; gill membranes broadly united, nearly free from the isthmus. Dorsal slightly notched; nasal tentacle present. Color excessively variable, mostly grayish, with 5 or 6 cross blotches on the back, extending to form quadrate blotches on the side; body mottled; fins also mottled; the anal dark, with a pale edge. Some specimens highly variegated, the caudal banded and with black and white spots; pale streaks from the eye across the cheek; dark bars on sides, extending on dorsal. Most specimens have the region above anal with numerous round whitish spots and some dark ones. These spots sometimes nearly obsolete, most evident on the paler specimens.

The following notes are taken from a specimen, No. 4635, M. C. Z., from Para, Brazil (Coll. Agassiz and Bourgeot): Head 4; depth  $4\frac{1}{2}$ . D. XII, 14; A. I, 16. Maxillary to front of eye, about equal to eye. Gill membranes free. Lower jaw with a very small canine, not twice the length of the upper teeth. Orbital cirrus quite small; a row of cirri along the nape, longer than the orbital cirrus. Head not very blunt, the anterior profile forming an angle above eye, thence straight and steep. Dorsal spines rather low and flexible, the fin scarcely notched. Color nearly lost; dark marblings on sides and on dorsal fin. This species is evidently the *Blennius crinitus* of Günther and the *B. asterias* of Goode & Bean, probably the *nuchifilis* of Cuvier and Valenciennes, and in all probability the *cristatus* of Linnaeus, also. These nominal species are from various localities in the Atlantic. If our specimens are all alike, all these forms most likely belong to 1 species. For this *cristatus* is the oldest name. The very small canines show considerable divergence from the type of *Blennius*, approaching *Scartella*. (Eu.) (*cristatus*, crested.)

*Blennius crista setacea longitudinale inter oculos*, GRONOW, Museum, I, No. 75; D. 26; A. 16; locality unknown. (Coll. Vosmaer.)

*Blennius cristatus*, LINNÆUS, Syst. Nat., Ed. x, 1, 256, 1758, Indies, after GRONOW; GÜNTHER, Cat. Fish., III, 223, 1861; JORDAN, Proc. U. S. Nat. Mus. 1890, 329.

*Blennius crinitus*, CUVIER & VALENCIENNES, Hist. Nat. Poiss., XI, 237, 1836, La Rochelle, France (Coll. D'Orbigny); GÜNTHER, Cat., III, 224, 1861.

*Blennius nuchifilis*, CUVIER & VALENCIENNES, Hist. Nat. Poiss., XI, 253, 1836, Isle of Ascension. (Coll. Quoy & Gaimard.)

*Blennius asterias*, GOODE & BEAN, Proc. U. S. Nat. Mus. 1882, 416, Garden Key, Florida (Type, No. 2620. Coll. G. Würdemann); Garden Key, Florida (Type, No. 2625. Coll. Dr. Whitehurst); Tortugas (Type, No. 6596. Coll. Dr. J. B. Holder); JORDAN & GILBERT, Synopsis, 961, 1883.

*Adonis cristatus*, GRONOW, Cat. Fish., Ed. Gray, 95, 1854.

## 885. SCARTELLA, Jordan.

*Scartella*, JORDAN, Proc. U. S. Nat. Mus. 1886, 50 (*microstoma*).

This genus differs from *Blennius* only in the entire absence of the posterior canine. The relations of this genus with such species of *Blennius* as *Blennius cristatus* are very close. It may be that the groups should be reunited, or that several species here referred to *Blennius* should be placed in *Scartella*. (*ὁ κάρτης*, one who leaps.)

## 2733. SCARTELLA MICROSTOMA (Poey).

Head 4 in length (5 with caudal); depth  $3\frac{5}{8}$  ( $4\frac{3}{8}$ ); eye  $3\frac{1}{4}$ . D. XI, 14; A. 15 or 16. Body rather stout, compressed posteriorly; head short, the anterior profile straight and very steep, almost vertical from tip of snout to above eye, where a sharp angle is formed with the straight line of the back. Eye large, longer than snout. Mouth moderate, the maxillary reaching to below front of pupil, its length  $3\frac{1}{2}$  in head. Teeth uniform; no posterior canines in either jaw. A small tufted or multifid cirrus over each eye, its length less than diameter of pupil; a row of about 3 short, slender cirri along each side of nape. Gill membranes broadly united, free from isthmus. Lateral line extending about to end of pectoral, each pore with a short, simple branch above and below, directed outward and backward; some conspicuous pores radiating from the eye. Dorsal fin low, subcontinuous, the spines rather slender, lower than the soft rays, the middle spines not much higher than the last; longest rays of dorsal about  $\frac{1}{2}$  as long as head; caudal free from dorsal and anal, a little shorter than head; anal low; pectorals slightly longer than head; ventrals  $1\frac{1}{2}$  in head. The fins are somewhat shriveled, so that the count of the rays is made with difficulty and may not be perfectly exact. Color very dark olive brown, paler below; head and anterior half of body plain, posterior half sprinkled with sharply defined dots of a vivid sky-blue color, becoming white in alcohol; about 6 obscure round darker blotches in a longitudinal series along sides posteriorly; fins dusky olive, mottled with darker, the caudal obscurely barred, the anal with a pale edge; spinous dorsal nearly black. Length  $3\frac{1}{2}$  inches. Cuba. Here described from a specimen taken by Dr. Jordan in Havana. We have also the following notes on Poey's type in the museum at Cambridge: D. XI, 15; A. 17. Dorsal and anal free from caudal. Body rather robust, the head blunt. Last tooth in each jaw a shade longer than its neighbor, but not canine-like. Gill membrane free from isthmus. Dorsal spines low, rather stiff, the fin deeply notched. Color much mottled, with some white spots on posterior half of body; a black ocellus behind first dorsal spine; 5 dark bars along back.

The following is Poey's description: Head  $4\frac{1}{2}$  in total length with caudal; depth  $5\frac{1}{4}$ . D. XII, 15; A. I, 17; P. 14. Snout short; profile falling abruptly; mouth small; eye 3 in head, twice interorbital space. Teeth 15 on each side in each jaw. Gill membrane broadly united, free from isthmus. Lat-

eral line disappearing on middle of back; a row of 6 filaments arranged in pairs on each side of the nape. Membranaceous tentacles over the eye; dorsal somewhat notched, pectoral strongly developed at base. Color brown, with 5 or 6 darker points which form on the back and reach base of the dorsal; pearly spots along sides and some below of the same color; caudal with 3 brown points. Cuba. One specimen, 46 mm. long. (Poey.) (*μικρός*, small; *στόμα*, mouth.)

*Blennius microstomus*, POEY, Memorias, II, 288, 1861, Cuba. (Coll. Poey.)

*Scartella microstoma*, JORDAN, Proc. U. S. Nat. Mus. 1886, 50.

### 886. HYPLEUROCHILUS, Gill.

*Hypleurochilus*, GILL, Proc. Ac. Nat. Sci. Phila. 1861, 168 (*geminatus*).

This genus differs from *Blennius* in the restriction of the gill-openings to the sides, the gill-membranes being broadly and fully joined to the isthmus; canines well developed. (*υ*, upsilon; *πλευρόον*, side; *χείλος*, lip; in allusion to the V-shaped lateral lips.)

### 2734. HYPLEUROCHILUS GEMINATUS (Wood).

Head  $3\frac{1}{2}$  to  $3\frac{3}{4}$ ; depth  $3\frac{1}{2}$  to 4. D. XI, 15 to XIII, 14; A. II, 18. Head not very blunt, the anterior profile straight, oblique; male (*multifilis*) with the supraocular cirrus very large, each with 4 smaller ones at base; supraocular cirrus in female (*geminatus*) low, shorter than eye, branched at tip; interorbital space concave, not  $\frac{1}{2}$  diameter of eye; a slight transverse groove behind eye; canines in both jaws, very strong, hooked backward, the lower considerably stronger than upper; gill openings extending downward to opposite or slightly below lower edge of pectoral. Dorsal fin not emarginate, the spines slender, but rather stiff, lower than the soft rays; pectorals shortish, ventrals rather long. Olive brown, faintly barred with darker; sides plain, or with several pairs of spots of a reddish-brown color, arranged pretty regularly in a double row; vertical fins edged with darker, especially the anal; dorsal black in front. Length  $2\frac{1}{2}$  inches. South Atlantic and Gulf coast of the United States, in shallow water; abundant in empty shells and clusters of tunicates. The sexes quite unlike, the male (*multifilis*) distinguished by the high suborbital crest. (*geminatus*, twin.)

*Blennius geminatus*, WOOD, Journ. Ac. Nat. Sci. Phila., IV, 1824, 278, Charleston, South Carolina, female (Coll. Prof. Bache); CUVIER & VALENCIENNES, Hist. Nat. Poiss., XI, 265, 1836.

*Blennius multifilis*, GIRARD, Proc. Ac. Nat. Sci. Phila. 1858, 169, St. Josephs Island, Texas, male (Coll. Gustav Würdemann); GIRARD, U. S. and Mex. Bound. Surv., Zool., pl. 12, fig. 6, 27, 1859; GÜNTHER, Cat., III, 562, 1861

*Hypleurochilus multifilis*, GILL, Proc. Ac. Nat. Sci. Phila. 1861, 168; JORDAN & GILBERT, Synopsis, 758, 1883.

*Hypleurochilus geminatus*, JORDAN & GILBERT, Synopsis, 759, 1883.

?*Blennius geminatus*, GÜNTHER, Cat., III, 288, 1861.

## 887. HYPSOBLENNIUS,\* Gill.

*Hypsoblennius*, GILL, Cat. Fish. East Coast U. S., 20, 1861 (*hentz*; no diagnosis).

*Isestes*, JORDAN & GILBERT, Synopsis, 757, 1883 (*gentilis*).

*Blenniolus*, JORDAN & EVERMANN, new subgenus (*brevipinnis*).

This genus differs from *Blennius* in the absence of canine teeth and in the restriction of the gill openings to the sides, the gill membranes being fully united to the isthmus as far upward as the base of the pectorals; ventral with 1 short, strong spine and 3 simple, articulated rays. The known species are American. ( $\psi\psi$ , high; *Blennius*.)

## HYPSOBLENNIUS:

a. Dorsal fin continuous, its margin entire or slightly notched.

b. Dorsal rays XI to XIII, 17 to 19. Pacific species.

c. Orbital cirrus multifid; spines of dorsal stiff; sides blotched or freckled. GILBERTI, 2735.

cc. Orbital cirrus simple or fringed.

d. Spines of dorsal slender and flexible; sides with round dark spots; anal rays 21. GENTILIS, 2736.

dd. Spines of dorsal rather stiff; sides with irregular dark cross bands rather than spots; anal rays 19. STRIATUS, 2737.

bb. Dorsal rays XII, 14 or XII, 15. Atlantic species.

e. Orbital cirrus simple, large or small; body everywhere with dark spots; dorsal spines rather low, stiffish. IONTHAS, 2738.

ee. Orbital tentacle forked at tip, long in males; dorsal spines stiff; body spotted. HENTZ, 2739.

BLENNIOLUS, (diminutive form, from *Blennius*):

aa. Dorsal fin deeply notched, very short, its rays XI, 12 or XII, 12; orbital tentacle slender, fringed; a dark lateral shade. BREVIPINNIS, 2740.

## Subgenus HYPSOBLENNIUS, Gill.

## 2735. HYPSOBLENNIUS GILBERTI (Jordan).

Head 4 in length ( $4\frac{3}{8}$  with caudal); depth 4 ( $4\frac{3}{8}$ ). D. XII, 19; A. II, 21. Body comparatively robust, deep, and compressed. Head large, rounded, the anterior profile less blunt than in *H. gentilis* and less rounded, nearly straight from tip of snout to above eye, thence again nearly straight to front of dorsal. Length of snout about equal to diameter of eye,  $4\frac{1}{2}$  in head. Mouth rather small, terminal, the maxillary reaching to opposite middle of eye,  $2\frac{3}{8}$  in head. Teeth subequal, with no trace of posterior canines. Superciliary tentacle large, multifid, much branched from near the base, the principal division  $3\frac{3}{8}$  in head. Gill openings larger than in *H. gentilis*, extending downward to the level of lower edge of pectoral, the length of the slit  $1\frac{1}{2}$  in head. Lateral line developed beyond the straight part, its posterior portion curved downward. Dorsal fin continuous, with a slight but distinct depression between the spinous and soft parts, the spines somewhat curved, but stiff and strong, the longest spine about  $2\frac{1}{2}$  in head; longest soft rays 2 in head. Caudal fin free from dorsal and anal,  $1\frac{1}{2}$  in head; ventrals  $1\frac{1}{2}$  in head; pectorals about as long as head. Males,

\* The recent identification of *Blennius hentz* with *Isestes punctatus* enables us to understand the undefined genus *Hypsoblennius*, and to substitute it for the later *Isestes*. Our judgment is opposed to the recognition of such unexplained "typonyms," but we defer to the custom of the American Ornithologists' Union.

as usual in this genus, with the anal spines partly detached, and provided with fleshy tips. Coloration olivaceous, the body and fins everywhere profusely mottled and reticulated with darker; obscure dark shades extending downward from eye across, or partly across, lower side of head; head without distinct spots or other sharply defined markings, except faint streaks radiating from eye; no pale bars on side of head in either sex; some yellowish markings on anterior part of dorsal. Length 5 inches. California, from Point Concepcion southward to Todos Santos or beyond; common among rocks in the kelp; our specimens from Santa Barbara and Point Loma. (Named for Charles Henry Gilbert.)

*Isoetes gilberti*, JORDAN, Proc. U. S. Nat. Mus. 1882, 349, Santa Barbara, California (Type, Nos. 26916 and 26917. Coll. Jordan & Gilbert); ROSA SMITH, Proc. U. S. Nat. Mus. 1883, 235, specimens from Todos Santos Bay; D. XI or XII, 16 to 21; A. 19 to 21; head  $4\frac{1}{2}$ ; depth  $4\frac{1}{2}$ .

2736. *HYPSOBLENNIUS GENTILIS* (Girard).

Head  $3\frac{3}{8}$  in length ( $4\frac{1}{2}$  with caudal); depth 4 ( $4\frac{1}{2}$ ). D. XIII, 17; A. II, 19. Body rather robust, deep and compressed, the head large, very bluntly and evenly rounded in profile, more obtuse and more evenly curved than in *H. gilberti*, the snout shorter, about equal to eye,  $4\frac{1}{2}$  in head. Mouth rather small, terminal, the maxillary reaching to opposite middle of eye, its length 3 in head. Teeth subequal, the hindmost on each side of upper jaw shorter than the others, and a little apart from them but not forming "a small canine," as stated by Girard. Superciliary tentacle long and simple in the male, its edge fringed with short branchlets, its length about 3 in head; tentacles much smaller in the female, where they are scarcely visible. Gill opening extending downward not quite to lower edge of pectoral, its length (vertical)  $2\frac{1}{2}$  in head. Lateral line with only the straight anterior portion developed, not curved downward posteriorly. Dorsal fins continuous, with scarcely a trace of emargination between the spinous and soft parts. Dorsal spines comparatively low and flexible, much less strong than in *H. gilberti*, the longest spines 3 in head; longest soft rays  $1\frac{3}{8}$ . Caudal free from dorsal and anal,  $1\frac{1}{2}$  in head; ventrals  $1\frac{3}{8}$  in head; pectorals  $1\frac{1}{2}$ . Coloration in spirits, brown, the whole body closely mottled and blotched with darker brown, so that the light ground color forms, especially anteriorly, light reticulations around darker spots; on the head the dark spots are small and close together, smallest anteriorly, the lower parts of the head being immaculate, extending from the curve of the preopercle downward, across the interopercle and branchiostegals, in a sharply defined white bar (said to be golden yellow in life), edged with black; behind this and parallel with it across subopercle and isthmus is a similar bar, these bars present only in the males; a few pale spots or bars in front of these; back with about 6 dusky cross shades, below each of these is an oblong dark blotch, the anterior placed along the lateral line, altogether forming an interrupted dark stripe; a similar dark stripe near the median line of the body, interrupted by some pale blotches. Fins all blotched and spotted by light and dark colors, but without distinct markings (a bluish spot in front of dorsal in life);



ventrals and anal nearly blackish in males, the base of the anal with a pale streak. Females more distinctly blotched, with a black spot in front of dorsal and white spots on middle of sides; head lacking the pale bars and black spots, but much mottled with brown and whitish; a very distinct blackish blotch on front of spinal dorsal; pectoral and caudal pale, a dark blotch on base of pectoral. Length about 4 inches. Monterey to Cape San Lucas; common southward in rock pools. Here described from specimens from Angel Island, Gulf of California, from Cape San Lucas, and from Monterey and San Diego. (*gentilis*, related.)

*Blennius gentilis*, GIRARD, Proc. Ac. Nat. Sci. Phil. 1854, 149, Monterey, California. Types, Nos. 690 and 785 (Coll. A. Cassidy; No. 489, Lieut. Trowbridge); GIRARD, Pac. R. R. Surv., x, Fishes, 113, pl. 25a, fig. 4, 1858; GÜNTHER, Cat., III, 217, 1861.

*Isestes gentilis*, STEINDACHNER, Ichth. Beiträge, v, 150, 1876; JORDAN, Proc. U. S. Nat. Mus. 1882, 350; JORDAN & GILBERT, Synopsis, 956, 1883; JORDAN, Proc. U. S. Nat. Mus. 1882, 349.

#### 2737. HYPSOBLENNIUS STRIATUS (Steindachner).

Head 4 to  $4\frac{1}{2}$ ; depth  $4\frac{3}{8}$  to 5. D. XI or XII, 17; A. 19; P. 15; V. I, 3. Snout steep, and slightly concave in older examples; interorbital narrow, equal to  $\frac{1}{2}$  eye; origin of dorsal a little before the edge of preopercle; second and third dorsal spines equal to the distance from tip of snout to edge of preopercle; dorsal and anal free from caudal; pectoral reaching nearly to front of anal. Color yellowish below, sides brownish, irregular dark-brown cross bars on back and sides; toward the caudal are rows of spots, 4 or 5 wider cross bars of dark brown or violet; a dark blotch from the third to the fifth dorsal spine, behind which are irregular longitudinal dark stripes: anal edged with white, behind which runs a violet line; pectoral and caudal spotted; a dark oval spot behind eye; a brown line from first dorsal spine to eye. Panama (Steindachner), where specimens were also taken by Dr. Gilbert, none of these showing posterior canines, although Steindachner notes the presence of a small canine in 1 specimen. (*striatus*, striped.)

*Blennius striatus*, STEINDACHNER, Ichth. Beiträge, v, 15, 1876, with plates, Panama.

*Isestes striatus*, JORDAN & GILBERT, Bull. U. S. Fish Comm. 1882, 111.

#### 2738. HYPSOBLENNIUS IONTHAS (Jordan & Gilbert).

Head  $3\frac{3}{8}$  to 4 ( $4\frac{1}{8}$  to  $4\frac{3}{8}$  in total); depth  $3\frac{1}{8}$  to  $3\frac{1}{2}$  ( $3\frac{1}{8}$  to  $4\frac{7}{8}$ ). D. XII, 13, or XII, 14; A. II, 13, or II, 14. Body rather deep, moderately compressed, the back little elevated. Head short, blunt, but less so than in *H. punctatus*; the profile prominent above the eye, thence descending abruptly but not vertically to the tip of the snout; length of snout  $3\frac{1}{8}$  in head. Mouth small, low, its cleft largely anterior, the short maxillary scarcely reaching past the front of the eye, 4 in head. Eyes large, placed high, 5 in head, the interorbital space about  $\frac{1}{4}$  their diameter. Female (*ionthas*) with the orbital cirrus low, scarcely larger than nasal cirrus, which is about equal to diameter of pupil. Teeth moderate, equal; no posterior canines. Gill opening extending

dow  
of  
roa  
stifi  
the  
com  
as h  
with  
sma  
part  
sepa  
ish  
olive  
dusk  
dusk  
Th  
D. X  
not c  
abru  
very  
in he  
ing v  
head  
Nasa  
bital  
midd  
scarc  
longe  
in he  
ventr  
tip o  
faint  
whic  
a yel  
front  
nume  
hook

*Isest*  
(T  
S;  
*Isest*  
N  
G  
\* TI  
excep  
both  
separ  
sion t

downward to a point varying from a little above to a little below middle of base of pectoral, the height of the slit 3 in head. Lateral line not reaching tip of pectoral. Dorsal fin continuous, the spines low and rather stiff, slenderer than in *H. punctatus*, the longest spines a little lower than the soft rays, which are about  $1\frac{1}{2}$  in head. Caudal free from anal, slightly connected with dorsal; a little shorter than head; pectoral about as long as head; ventrals shorter than head. Color of female clear olive green, with only traces of darker bars; body everywhere densely freckled with small round blackish spots, smaller than the pupil; on the sides and lower part of head these spots are reduced to close set dots; 2 dark lines, separated by a golden area, downward from eye; a vertical curved blackish patch behind eye, in front of which is a golden area; vertical fins olive green, dorsal and caudal usually mottled with dusky; paired fins dusky olive; lower parts of head tinged with golden, sometimes with dusky cross bars; cirri green.

The male (*scrutator*) is thus described: Head 4 ( $4\frac{1}{2}$  in total); depth  $3\frac{1}{2}$  ( $4\frac{1}{2}$ ). D. XII, 14 or 15; A. II, 15 or 16. Body rather deep, compressed, the back not elevated. Head short, very blunt, almost as deep as long, the profile abruptly descending before eye, the snout about  $\frac{1}{2}$  length of head. Mouth very small, anterior, the maxillary extending to opposite front of eye,  $3\frac{1}{2}$  in head; teeth subequal, without canines. Orbital cirri very long, reaching when depressed about to the front of dorsal, their length more than  $\frac{1}{2}$  head in adult, somewhat shorter in young; a short branch near its middle. Nasal barbel minute. Eye large, much broader than the concave interorbital space, about  $4\frac{1}{2}$  in head. Lower edge of gill opening a little below middle of base of pectoral, the depth of the slit  $2\frac{1}{2}$  to 3 in head. Dorsal fin scarcely emarginate, the spines rather stiff, lower than the soft rays, the longest spine 2 in head. Caudal slightly connected at base with dorsal,  $1\frac{1}{2}$  in head; pectoral about as long as head, reaching past front of anal; ventrals  $1\frac{1}{2}$  in head. Lateral line extending to base of eighth spine, not to tip of pectoral. Color in life, deep olive green, almost immaculate, or with faint traces of darker vertical bars; a golden blotch behind eye, behind which is a dusky crescent; 2 dark bars downward from eye, separated by a yellowish area; fins all dusky greenish, nearly or quite immaculate; front of spinous dorsal blackish. South Carolina to Texas, in rock pools; numerous specimens, the largest about  $2\frac{1}{2}$  inches long, were obtained with hook and line from the wharves at Pensacola. (*ἰορθός*, freckled.)

*Isestes ionthas*, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1882, 299, Pensacola, Florida (Type, No. 30856, U. S. Nat. Mus. Coll. Jordan & Stearns), female; JORDAN & GILBERT, Synopsis, 960, 1883.

*Isestes scrutator*,\* JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1882, 300, Pensacola (Type, No. 30850. (Coll. Jordan & Stearns); Galveston (Coll. Dr. August Galny); JORDAN & GILBERT, Synopsis, 960, 1883.

\* The form called *scrutator* agrees very closely with *Hypsoblennius ionthas* in all respects except the great length of the orbital cirrus and the different coloration of the body. In both the golden blotch and dark crescent behind the eye are distinct, as also the 2 dark bars separated by a yellow one below the eye. Renewed comparison strengthens our impression that *Hypsoblennius scrutator* is the male of *Hypsoblennius ionthas*.

## 2739. HYPSOBLENNIUS HENTZ (Le Sueur).

Head  $3\frac{3}{8}$ ; depth 3. D. XII, 15; A. 18; pectoral  $1\frac{1}{2}$  in head; ventral  $1\frac{1}{2}$ ; gill slit  $2\frac{1}{2}$ ; eye  $4\frac{1}{2}$ ; maxillary  $2\frac{3}{8}$ . Orbital tentacle very slender, once forked, 3 in head. Body rather deep; head large, obtuse; interorbital space concave,  $\frac{1}{2}$  the diameter of orbit; orbital cirrus as long as dorsal spines, bifid at tip, branched below; a minute nasal cirrus; no canines; gill openings extending to about lower fourth of base of pectoral, thus narrower than in most related species. Dorsal fin high, little notched, the soft part highest, the spines stiff,  $2\frac{3}{8}$  in head. Tip of each dorsal spine with a filiform, articulated, ray-like appendage. Color in spirits, olivaceous, back and sides of head and body everywhere covered with brown spots, very irregular in size and shape; on posterior part of body the spots are larger, and show a tendency to form vertical bars; cheeks dark; lower side of head with traces of 3 cross bars; spinous dorsal with an elliptical black spot on membrane of first 3 spines; soft dorsal and caudal obscurely barred; anal, ventrals, and lower rays of pectorals dusky; pectorals olivaceous, spotted with brown. Coasts of North and South Carolina, south to Indian River, Florida; locally common. (Named for its collector, Dr. Nicholas Marcellus Hentz, "the father of American Araneology.")

*Blennius punctatus*, WOOD, Journ. Ac. Nat. Sci. Phila., IV, 1825, 278, Charleston, South Carolina (Coll. Prof. Bache); CUVIER & VALENCIENNES, Hist. Nat. Poiss., XI, 267, 1836; GÜNTHER, Cat., III, 228, 1861; not *Blennius punctatus*, Fabricius, 1780, which is a *Stichæus*.

*Blennius hentz*,\* LE SUEUR, Journ. Ac. Nat. Sci. Phila., IV, 1825, 363, Charleston, South Carolina. (Coll. Dr. Hentz.)

*Hypsoblennius hentzi*, GILL, Cat. Fish. East Coast N. A., 1861 (*nomen nudum*).

*Hypoleurochilus punctatus*, GILL, Cat. Fish. East Coast N. A., 20, 1873.

*Isesthes punctatus*, JORDAN & GILBERT, Synopsis, 758, 1883; JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1883, 616.

*Isesthes hentzi*, JORDAN & GILBERT, Synopsis, 960, 1883.

## Subgenus BLENNIOLUS, Jordan &amp; Evermann.

## 2740. HYPSOBLENNIUS BREVIPINNIS (Günther).

Head  $3\frac{1}{2}$ ; depth 4. D. XII, 12; A. II, 14; pectoral  $1\frac{1}{2}$  in head; ventral  $1\frac{1}{2}$ ; gill slit  $2\frac{1}{2}$ ; eye  $3\frac{1}{2}$  in head; snout  $2\frac{1}{2}$  in head; maxillary  $2\frac{1}{2}$ . Orbital tentacle slender, less than eye. Body rather deep, compressed, back not elevated; anterior profile from first dorsal spine to above eye almost horizontal or slightly decurved, from thence to tip of snout abruptly decurved; head large, its width not quite 2 in its length; interorbital space narrow, grooved, about equaling pupil; eyes large, placed high and close together. Mouth small, low, the maxillary reaching to pupil; teeth subequal, pectinate; no canines; dorsal fin continuous, deeply emarginate, the spines lower than the soft rays, the longest spine about  $2\frac{1}{2}$  in head; caudal free

\* The following is the substance of the account of "*Blennius hentz*:" Depth  $3\frac{1}{2}$  (in total). D. XI, 14; A. 16. Body little elongate; snout very short, but not vertically truncate; eyes above angle of mouth, placed high; gill slit extending from level of base of pectoral fin to height of eye; teeth equal; dorsal slightly depressed in the middle; pectorals large; a short cirrus above each eye and a smaller one over each nostril. Light bluish ash, mixed with rufous, with numerous irregular black and rufous spots; dorsal black, with whitish spots; soft dorsal with 5 dark bands; ventrals blackish, with pale bands; caudal with 3 or 4 dark bands. Charleston Harbor, South Carolina. (Le Sueur.)

from anal and dorsal; lateral line not reaching soft dorsal. Olive brown, lighter below; back and upper half of sides irregularly marked with about 6 distinct dark-brown cross bars, these uniting at their lower edges and forming a continuous line from head to base of caudal; the bars nearly confluent on the back at base of dorsal fin; a dark lateral band nearly as wide as eye from opercle to base of caudal, containing 5 or 6 light-yellowish spots corresponding to the pale interspaces along the back; fins dusky, anal margined with black; head with a dark spot behind each eye, and 2 smaller blotches in the median line, 1 immediately behind the eyes, the other a short distance in front of dorsal. Pacific coast of Mexico, from Mazatlan to Panama; rather common. The specimens here described from Mazatlan. (*brevis*, short; *pinna*, a fin.)

*Blennius brevipinnis*, GÜNTHER, Cat. Fish., III, 226, 1861, Pacific coast Central America (Coll. Capt. John M. Dow); one specimen wrongly attributed to Hawaiian Islands.

888. CHASMODES, Cuvier & Valenciennes.

*Chasmodes*, CUVIER & VALENCIENNES, Hist. Nat. Poiss., XI, 295, 1836 (*bosquianus*).

*Blenitrachus*, SWAINSON, Class'n Fishes, etc., II, 78, 274, 1839 (*quadrifasciatus*).

Body oblong, compressed, naked; head triangular in profile, the snout somewhat pointed; mouth large, with lateral cleft, the maxillary usually, but not always, extending to beyond eye; premaxillaries not protractile; teeth in a single series, long and slender, comb-like, confined to the front of each jaw; no canines; cirri very small or wanting; gill openings very small, their lower edge above the middle of the base of the pectorals; lateral line incomplete. Fins as in *Blennius*. American. The species with smaller mouth approach *Hypsoblennius*, which genus is not far separated from *Chasmodes*. (*χασμώδη*, yawning.)

a. Dorsal and anal free from caudal.

b. Anal rays 18 or 19; body not banded.

JENKINSI, 2741.

bb. Anal rays 15; body with 4 dark cross bands.

QUADRIFASCIATUS, 2742.

aa. Dorsal joined to base of caudal.

c. Mouth moderate, the maxillary not extending to posterior border of eye,  $2\frac{1}{2}$  in head.

SABURRÆ, 2743.

cc. Mouth large, maxillary reaching posterior border of eye.

NOVEMLINEATUS, 2744.

ccc. Mouth very large, the maxillary extending to beyond eye.

BOSQUIANUS, 2745.

2741. CHASMODES JENKINSI (Jordan & Evermann).

Head  $3\frac{1}{2}$  (4 in total); depth 4 (5). D. XII, 17; A. 18 or 19; eye 4 to 5 in head. Body more robust than in related species, resembling *Hypsoblennius*; head large, gently rounded in profile, the snout steep, 4 in head; interorbital space narrow, grooved; orbital tentacle (male) much as in *Hypsoblennius gilberti*, about 3 in head, branched, the branches usually 4; mouth much larger than in *Hypsoblennius*, the maxillary  $2\frac{1}{2}$  to 3 in head, reaching to below posterior margin of eye; teeth even, comb-like; gill opening 2 in head, extending downward nearly to lower edge of pectoral, much larger than in *Chasmodes saburræ*. Dorsal little notched, the spines

slender,  $2\frac{1}{2}$  in head, the rays a little higher; anal lower, the rays  $3\frac{1}{2}$  to 4 in head; pectorals reaching anal,  $1\frac{1}{2}$  in head; ventrals  $2\frac{1}{2}$ ; dorsal and anal free from caudal. Color in life, according to Evermann & Jenkins, yellowish; 5 quadrate spots of darker extending from dorsal to a line drawn from middle of eye to lower base of caudal, the anterior one above tip of pectoral; median line of side with a more or less distinct series of small spots; a short dark vertical line behind the eye; a dark blotch in front of origin of dorsal fin and another on humeral region; underside of head with 2 ill-defined dark bands; dorsal fin more or less speckled with black, the anal with a narrow white border above which is a broader band of deep brown. Six specimens, the largest about 3 inches long, were obtained at Guaymas, Sonora, by Drs. Evermann & Jenkins, in 1887. One of these, (No. 412, L. S. Jr. Univ. Mus.), examined by us, is the type of the present description. The large mouth distinguishes this species at once from *Hypsoblennius striatus*, with which it has been identified. The species is intermediate between typical *Chasmodes* and *Hypsoblennius*, and its discovery may make it necessary to merge the latter in *Chasmodes*. (Named for Dr. Oliver Peebles Jenkins.)

*Hypsoblennius striatus*, EVERMANN & JENKINS, Proc. U. S. Nat. Mus. 1891, 163; not of STEINDACHNER.

*Chasmodes jenkinsi*, JORDAN & EVERMANN, Proc. Cal. Ac. Sci. 1896, 232, pl. 39, Guaymas. (Coll. Evermann & Jenkins.)

#### 2742. CHASMODES QUADRIFASCIATUS (Wood).

D. 27; A. 15. Form of *Chasmodes bosquianus*: Lower jaw slightly longer than the upper. Dorsal and anal free from caudal; anal fin highest anteriorly. Body with 4 distinct brownish bands, a fifth broader and less marked on the neck; 4 round yellowish spots along base of anal; head spotted with blackish. (Wood.) *Habitat* uncertain, probably South Atlantic coast of the United States; not recognized by recent collectors; very likely based on the female of *C. bosquianus*, with the caudal torn from the other vertical fins. (*quadri-*, four; *fasciatus*, banded).

*Pholis quadrifasciatus*, WOOD, Journ. Ac. Nat. Sci. Phila., IV, 1825, 282; locality unknown, probably South Carolina. (Coll. Rubens Peale.)

*Chasmodes quadrifasciatus*, CUVIER & VALENCIENNES, Hist. Nat. Poiss., XI, 298, 1836; GÜNTHER, Cat. Fish., III, 229, 1861; JORDAN & GILBERT, Synopsis, 757, 1883.

#### 2743. CHASMODES SABURRE, Jordan & Gilbert.

Head  $3\frac{1}{2}$  to  $3\frac{3}{4}$ ; depth  $3\frac{1}{2}$  to  $3\frac{3}{4}$ . D. XI or XII, 17 to 19; A. II, 18 or 19. Body rather deep and compressed, less elongate than in *C. bosquianus*; the back somewhat arched. Head comparatively short, much shorter than in *C. bosquianus*, not  $\frac{1}{2}$  longer than deep; profile forming a nearly even curve from the base of the dorsal to the tip of the snout; mouth notably smaller than in *C. bosquianus*; maxillary not reaching posterior margin of eye, its length  $2\frac{1}{2}$  in head; teeth occupying about  $\frac{1}{2}$  of lower jaw; height of gill slit  $3\frac{1}{2}$  in head, its lower ray opposite third ray of pectoral. A minute cirrus, shorter than pupil, above each eye and each nostril. Dorsal con-

tin  
ray  
dar  
lati  
fron  
spot  
mos  
and  
obsc  
ish;  
tips  
with  
blac  
with  
cola  
wate  
but v

Chasm  
F  
188

Hea  
borde  
body  
nearly  
what  
anal h  
nearly  
dark,  
separa  
with s  
tled w  
just be  
are 2  
ward  
whole  
ously  
dark a  
ventra  
the Ur  
Florid  
mann

Pholis n  
Sou

\* The  
with 9 v  
first and  
somewh

tinuous, with slender rays, the last one joined to the caudal. First two rays of anal short, thick, and fleshy in the males. Male deep olive, with dark cross shades; numerous pale spots on the sides which form undulating lines converging backwards; dark stripes downward and forward from eye; top of head and upper part of dorsal fin usually with fine black spots; spinous dorsal with a median orange longitudinal band; other fins mostly dusky olive. Some specimens with the outer part of both dorsals and the top of head dusted with black spots, others with these spots obsolete; soft dorsal and caudal light orange, barred with light greenish; anal dull orange, with an obscure blackish median band, the exerted tips of the rays abruptly whitish; pectorals dusky olive, strongly tinged with orange; ventrals blackish, orange at tip. Female with about 8 blackish cross bands extending on the dorsal fin; the body everywhere with pale spots; fins all sharply barred with blackish and olive. Pensacola Bay, Florida; common about the wharves and ballast rocks in shallow water; taken with seines and pinhooks. Allied to *Chasmodes bosquianus*, but with the mouth smaller, the form less elongate. (*saburra*, ballast.)

*Chasmodes saburrae*, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1882, 298, Pensacola, Florida (Type, No. 30824. Coll. Jordan & Stearns); JORDAN & GILBERT, Synopsis, 958, 1883.

2744. CHASMODES NOVELLINEATUS (Wood).

Head  $3\frac{1}{2}$ ; depth  $3\frac{1}{2}$ ; eye  $4\frac{1}{2}$ ; snout  $3\frac{1}{2}$ ; maxillary reaching posterior border of eye. D. XI, 18; A. III, 17. Head and shoulders heavy, the body lance-shaped, tapering gradually to tail; snout short, blunt, profile nearly vertical to eye, thence gently rounded; mouth rather large, somewhat oblique, the maxillary reaching posterior border of eye; dorsal and anal high, longest dorsal rays 2 in head; anal considerably lower; pectoral nearly as long as head; ventrals  $1\frac{1}{2}$  in head. Color, side with 6 broad, dark, vertical bars, the anterior 4 extending on the dorsal fin, these bars separated by irregular narrow pale spaces; entire side profusely covered with small white spots; a small black spot at base of caudal; head mottled with light and dark; 2 small dark spots on under side of lower jaw; just behind these and extending downward from the angles of the mouth are 2 other larger, blacker spots, while behind these, extending downward and backward from middle of cheek, is an irregular black line; whole head with numerous fine dark punctulations; dorsal and anal variously spotted or barred with light and dark; spinous dorsal with a large dark area at top of anterior spines; caudal faintly barred; pectorals and ventrals more plainly barred. Length 2 inches. South Atlantic coast of the United States, South Carolina to Florida; abundant in Indian River, Florida, where numerous specimens were taken in January, 1896, by Evermann & Bean. (*novem*, nine; *lineatus*, lined.)

*Pholis novemlineatus*, WOOD,\* Journ. Ac. Nat. Sci. Phila., IV, 1825, 280, Charleston Harbor, South Carolina.

\* The following is the substance of Wood's original description of this species: "Body with 9 whitish longitudinal bands; dorsal fin with an irregular blackish spot between the first and second rays; remainder of the fin clouded with dusky brown. Head descending somewhat abruptly, tuberculated anteriorly; nostrils with a small appendage; head, lips,

*Chasmodes saburrae*, EVERMANN & BEAN, Fishes of Indian River, Florida, in Rept. U. S. Fish Comm. 1896, 247; not of JORDAN & GILBERT.

*Chasmodes novemlineatus*, GÜNTHER, Cat., III, 229, 1861.

2745. *CHASMODES BOSQUIANUS* (Lacépède).

Head  $3\frac{1}{2}$ ; depth  $3\frac{1}{2}$ . D. XI, 19; A. II, 19. Orbital tentacle very minute or wanting; maxillary extending to rather beyond eye; interocular space very narrow, not concave. Dorsal fin not emarginate, the spines slender. Dorsal joined to base of caudal; anal free. Color (in male) olive green, with about 9 horizontal narrow blue lines, these somewhat irregular and interrupted, converging backward; opercular membrane and a broad stripe through middle of spinous dorsal deep orange yellow; anal fin dark, the rays with white membranaceous tips; female dark olive green, reticulated with narrow, pale green lines, and with several broad dark bars, which are more distinct posteriorly; vertical fins similarly marked; head finely dotted with black; a dusky spot at base of caudal in both sexes. New York to Florida; common southward in shallow water. (Named for M. Bose, who collected at Charleston for Lacépède.)

*Blennius bosquianus*, LACÉPÈDE, Nat. Hist. Poiss., II, 403, 1800 (female), South Carolina. (Coll. Bose.)

*Pholis quadrifasciatus*, WOOD, Journ. Ac. Nat. Sci. Phila., IV, 1824, 282, locality unknown, probably South Carolina. (Coll. Rubens Peale.)

*Chasmodes bosquianus*, GÜNTHER, Cat., III, 229, 1861.

*Chasmodes bosquianus*, JORDAN & GILBERT, Synopsis, 756, 1883.

889. *HOMESTHES*, Gilbert, new genus.

*Homesthes*, GILBERT, new genus (*caulopus*).

Differing from *Hypsoblennius* chiefly in the presence of 4 articulated ventral rays instead of 3, as usual in *Blenninae*. We have examined the ventrals of *Hypsoblennius striatus*, *punctatus*, *ionthas*, *gentilis*, and *gilberti*, and have found them to consist constantly of 1 short, strong spine and 3 simple articulated rays. In *Homesthes caulopus* there is 1 strong, short spine and 4 well-developed simple jointed rays. (*ὁμός*, uniform; *ἔσθια*, to eat.)

2746. *HOMESTHES CAULOPUS*, Gilbert, new species.

Head  $3\frac{3}{8}$  in length; depth at base of ventrals 4, at middle of abdomen  $3\frac{3}{8}$ ; least depth of caudal peduncle  $\frac{1}{2}$  length of head; snout 4; eye 4 to  $4\frac{1}{5}$ . D. XII, 15 or 16; A. II, 17; P. 14. Longest dorsal spine  $2\frac{3}{8}$ ; last dorsal

opercula, etc., and base of the pectoral fins, finely spotted with bluish black, the spots being larger on the front and opercula; branchial opening extremely small, extending  $\frac{1}{2}$  of the length of the external curve of the operculum; mouth descending little; gape moderate; sides of the head fleshy; body compressed; rib spaces evident; sides with 9 longitudinal whitish lines, some of which are interrupted; behind the eye and under the dorsal fin are 2 irregular whitish patches; dorsal fin commencing before the pectoral fins; between the first and second rays is an irregular blackish spot, several of the following rays are also spotted, the color of the spots becoming lighter as they recede toward the tail, where they mingle with the dusky color of the fin and are lost; fin rising posteriorly, and joining the caudal fin at about  $\frac{1}{2}$  the distance from its extremity; anal fin commencing under the termination of the pectoral fin, and extending nearly to the tail; caudal fin rounded; ventral fins 2-rayed; pectoral fins rather large, the base thick and fleshy, finely spotted with bluish black; anus small, tubercle small; color brownish, fins dusky. D. 30; C. 123; A. 20; V. 2; P. 13. Length  $3\frac{1}{2}$  inches; depth, exclusive of the dorsal fin, hardly 1 inch.

spine  $3\frac{1}{2}$ ; longest (tenth) dorsal ray 2; longest (fifteenth) anal ray  $2\frac{1}{2}$ ; ventrals  $1\frac{1}{2}$ ; longest pectoral ray  $1\frac{1}{2}$  to  $1\frac{3}{4}$ ; caudal  $1\frac{1}{2}$ . Robust, moderately compressed, with wide heavy head and short, bluntly rounded snout, the anterior profile of which is nearly vertical. In shape and general appearance much resembling *Hypsoblennius gilberti*. Mouth very wide, horizontal, short, the maxillaries reaching vertical from hinder edge of pupil, 3 to  $3\frac{1}{2}$  in head. Teeth, as usual in this group, the posterior not enlarged or canine-like. Nostrils with slightly elevated margins, scarcely tubular, the hinder edge of anterior nostril produced into a conspicuous laciniate flap, about  $\frac{2}{3}$  as long as the diameter of orbit. A similar but larger orbital cirrus, divided nearly to the base into 6 or 8 slender filaments. Interorbital space deeply grooved, without median ridge, opening posteriorly into the deep transverse groove which separates the orbital region from the somewhat swollen occiput, its width  $1\frac{1}{2}$  eye. The mucous canals of head give off transverse branches which open by numerous pores. These thickly beset the snout, subocular region, top of head, preopercle, and upper portion of opercle. Width of gill slit equaling or slightly exceeding  $\frac{1}{2}$  length of head, confined to area above lower base of pectorals. First dorsal spine over margin of preopercle; spinous dorsal low, of nearly uniform height, much lower than second dorsal, the spines rather strong at base, with weak reflexed tips; membrane of last dorsal ray joined to extreme base of rudimentary caudal rays; anal low, rising slightly posteriorly, leaving a short free interval between its last ray and caudal. Lateral line strongly developed anteriorly for a distance equaling length of head; from that point it is only faintly visible, declining abruptly to middle of sides, along which it may be traced to base of caudal; the anterior portion gives off numerous pairs of short transverse lines, each of which ends in a pore; no pores or lines are visible posteriorly. Blackish, without sharp markings, the sides with irregular light blotches, some of which are subcircular in outline and contain 1 or more black central specks; the light markings near the back elongate and vertically placed, faintly outlining dark bars of the ground color; a vertical black blotch on cheek behind eye; lower parts lighter; no distinct bars on head; fins all blackish, the anal, the ventrals, the lower caudal and pectoral rays deeper black; anal and caudal margined with white, some of the dorsal rays narrowly tipped with white; tentacles whitish. Two specimens, 4 and  $4\frac{1}{2}$  inches long, from Panama Bay. (Gilbert.) (*καυλός*, stem; *πύς*, foot.)

*Hemesthes caulopus*, GILBERT MS., Fishes of Panama, Panama. (Coll. Gilbert. Type No. 5623, L. S. Jr. Univ. Mus.)

### 890. SCARTICHTHYS, Jordan & Evermann, new genus.

*Scartes*, JORDAN & EVERMANN, Check-List Fishes, 471, 1896 (*rubropunctatus*); preoccupied by *Scartes*, Swainson, a genus of mammals.

*Scartichthys*, JORDAN & EVERMANN, new genus (*rubripunctatus*).

Body elongate, slowly declining to the caudal. Head obliquely compressed, oblong, the profile more or less vertical. Eyes lateral, closely approximated, situated at the angle of the profile with the postocular



region. Gill apertures continuous under the throat, gill membrane free from isthmus. Branchiostegals 6. Mouth moderate, the contour of the upper jaw semicircular; upper jaw protruding beyond the lower; lips moderate, uniform and free, concealing the teeth. Teeth labial and movable, very slender and recurved, contiguous and uniserial; no posterior canines. Dorsal fin divided; anal similar to soft dorsal; caudal obtusely rounded; pectorals moderate, angularly rounded; ventrals approximated, each with 3 simple rays, the internal of which is smallest. This genus is very close to the Old World genus, *Salarias*,\* Cuvier, which differs in having the dorsal fin continuous, as in *Rupiscartes*. (σκάρτης, one who leaps; ἰχθύς, fish.)

2747. SCARTICHTHYS RUBROPUNCTATUS (Cuvier & Valenciennes).

Head 4; depth 4 (5 with caudal); D. XI-16; A. 20; eye  $4\frac{1}{2}$  in head; teeth less flexible than in *Rupiscartes atlanticus*; no canine teeth; the forehead not projecting beyond the mouth; a very small tentacle on the neck, a longer fringed one above the orbit; dorsal fin deeply notched, not extending on to the caudal. Color brown, marbled with black, and dotted with reddish; a black spot on the anterior part of the dorsal; throat with 2 or 3 brownish cross bands; a jet-black spot behind eye, with a narrow edge posteriorly. (Günther.) Coast of Peru and Chile, north to Panama. Specimens examined by us collected by Prof. Frank H. Bradley at Pearl Islands, near Panama, and at Callao. Length 3 inches. (*ruber*, red; *punctatus*, spotted.)

*Salarias rubropunctatus*, CUVIER & VALENCIENNES, Hist. Nat. Poiss., XI, 348, 1836, Juan Fernandez (Coll. Claude Gatz); GÜNTHER, Cat., III, 249, 1861; JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1882, 628; not of KNER, Novara-Fische, 198.

891. RUPISCARTES, Swainson.

*Alticus*,† COMMERSON, in LACÉPÈDE, Hist. Nat. Poiss., II, 458, 1800 (*saliens*).

*Alticus*,‡ CUVIER & VALENCIENNES, Hist. Nat. Poiss., XI, 337, 1836 (*alticus*).

*Rupiscartes*, SWAINSON, Nat. Hist. Class'n Anim., II, 275, 1839 (*alticus*).

This genus, as here understood, differs from *Salarias* only in the presence in 1 or both jaws of posterior canines. Dorsal fin continuous, without deep notch. Vertebrae  $12 + 22 = 34$  (*atlanticus*). (*rupes*, rock; σκάρτης, one who leaps; *Rupiscartes tridactylus* (*alticus*), "said to jump on the sea rocks like a lizard." Swainson.)

\* *Salarias*, CUVIER, Règne Anim., Ed. 2, II, 175, 1829 (*quadripinnis*). *Erypichthys*, SWAINSON, Nat. Hist. Class'n. Anim. II, 275, 1839 (*quadripinnis*, etc.). (σαλάρια, a modern Greek name of *Blennius basilius*.)

† We do not think that the name *Alticus* can be substituted for *Rupiscartes*, because Lacépède does not adopt this genus of Commerson, but merges it in *Blennius*, quoting Commerson's account as a footnote. This is as follows: "*Alticus saltatorius*, pinna spuria in capitis vertice; seu pinnula longitudinali pone oculos cartilaginea; seu alticus desultor, occipite cristato, ore circulare deorsum patulo." Apparently this quotation of a generic description not approved, does not give priority to the latter.

‡ This genus *Alticus* is not adopted by Cuvier & Valenciennes. Valenciennes speaks of "un petit *Salarias* que nous paraît être celui-là même sur lequel Commerson avait établi son genre *Alticus*." But a genus is not established until it is accepted by some authority as well as defined.

b  
H  
fr  
to  
M  
be  
an  
op  
fr  
sal  
to  
rea  
int  
tur  
ho  
dee  
bra  
Sou  
ver  
exc  
nea  
the  
dar  
fins  
one  
Trop  
Wes  
Maz

Pun  
Salar  
(  
Rupi

Salar  
Ento

Th  
fin is  
shar

a.

\* Th  
based  
was n

2748. *RUPISCARTES ATLANTICUS* (Cuvier & Valenciennes).

Head 4 to 4½; depth 3½ to 3¾. D. XII, 21 or XIII, 20; A. 24 or 25; vertebrae 12 + 22 = 34; eye 4 to 4½ in head. Body rather high, compressed. Head short, very blunt, its width about 2 in the length; anterior profile from first dorsal spine to above eye straight or slightly convex; from thence to tip of snout abruptly decurved, in some specimens nearly vertical. Mouth inferior, lower jaw included; maxillary about reaching posterior border of eye. Teeth small, pectinate, the lower canines exceedingly large and entering the cavity in the palate. Supraorbital tentacle well developed, slender; a group of 5 or 6 short tentacles on either side of head in front of nostrils and on either side of neck, these shorter than pupil. Dorsal fin not emarginate, extending from a point above middle of operculum to base of caudal; anal lower than soft dorsal, 1½ to 2 in dorsal; pectorals reaching past vent, about equaling head; ventrals about 2 in head. "The intestinal tract is more than 3 times as long as the entire body. The structure of the skeleton is very similar to that of the Blennies; the jaw bones, however, are still shorter, and the intermaxillary and mandibular are deeply concave anteriorly. There are 12 abdominal and 22 caudal vertebrae, the former portion being only ½ as long as the caudal." (Günther.) Some specimens, apparently males, with the anterior profile vertical and very high; fins high; caudal lanceolate, the black median rays much exceeding the outer pale ones. Females with the anterior profile a nearly even curve, the caudal lunate, its median black rays shorter than the outer pale ones. Body liver brown, paler below, with usually 5 or 6 darker cross bars extending on the dorsal; a black spot behind eye in all; fins mostly blackish, an orange area on upper edge of caudal; a yellow one tinged reddish below; eye red posteriorly. Length 6 to 8 inches. Tropical America, on both coasts, very abundant in rock pools, north to West Indies and to Todos Santos. Here described from specimens from Mazatlan.

*Punaru*, MARCGRABE, Hist. Brazil, 165, 1648, Brazil.

*Salaria atlanticus*, CUVIER & VALENCIENNES, Hist. Nat. Poiss., XI, 321, 1836, Madeira (Coll. Richardson). Antilles (Coll. Plée); GÜNTHER, Cat. Fish., III, 242, 1861.

*Rupiscartes atlanticus*, JORDAN, Proc. U. S. Nat. Mus. 1898, 333.

892. *ENTOMACRODUS*,\* Gill.

*Salaria*, SWAINSON, Nat. Hist. Class'n Fishes, II, 274, 1839 (*vermicularis*; not of CUVIER).

*Entomacrodus*, GILL, Proc. Ac. Nat. Sci. Phila. 1859, 168 (*nigricans*).

This genus has large posterior canines as in *Rupiscartes*, but the dorsal fin is divided into 2 fins as in *Scartichthys*. (ἐν, in; τόμος, cutting; ἀκρός, sharp; ὀδούς, tooth.)

a. Orbital cirrus present; dorsal rays XII or XIII-15; canines small.

b. Cirrus above eye divided; anal rays 15.

CHIOSTICTUS, 2749.

bb. Cirrus above eye simple or nearly so; anal rays 18; body with pearly spots.

MARGARTACEUS, 2750.

\* This genus is equivalent to *Salaria* of Swainson, but the generic name *Salaria* was based on *Salaria quadripinnis*, before either of the species referred to it by Swainson was made known.

aa. Orbital cirrus wanting; no cirri at nape.

c. Dorsal rays XII-19; anal 15; body rather slender, the depth about 5 in length; body with bands and spots. DECORATUS, 2751.

cc. Dorsal rays XI-15; anal 17; body very slender, the depth about 6 in length; color blackish, nearly plain. NIGRICANS, 2752.

2749. ENTOMACRODUS CHIOSTICTUS (Jordan & Gilbert).

Head  $4\frac{1}{2}$  in length; depth  $5\frac{1}{4}$ . D. XII-15; A. 15; eye  $3\frac{1}{4}$  to  $4\frac{1}{4}$  in head, varying with age. Body moderately elongate, compressed, the head short, blunt, almost globular, about as broad as deep, and a little longer than broad. Mouth inferior, with little lateral cleft, the lower jaw included; width of cleft of mouth  $\frac{2}{3}$  length of head. Teeth small, weak, finely pectinate; canine teeth small, not so long as diameter of pupil. Supraorbital cirrus divided into 4, its height  $\frac{2}{3}$  that of eye; a few minute slips at the nape. Interorbital space channeled, narrower than eye. Maxillary extending to behind middle of eye. No crest on top of head. First dorsal low and even, its spines rather slender, the last spines short, scarcely connected by membrane with the soft rays; soft dorsal well separated from caudal; caudal subtruncate, with rounded angles; anal lower than soft dorsal, with a little longer base; pectorals a little longer than head; ventrals about  $\frac{1}{2}$  as long. Color in life, olive brown above, lighter below; 5 broad, dark bars from dorsal fin to middle of sides, each terminating above on the fin, and below on sides in a pair of black spots; sometimes only the dark spots are distinguishable, the bars being obscure; sides below spinous dorsal with numerous black specks, and with numerous oblong spots of bright silvery; sometimes a silvery streak from upper portion of base of pectorals to base of caudal; a broad salmon-colored streak on each side of ventral line; sometimes the space between the silvery lateral band and the base of the anal is darker, the vertical bars again appearing as pairs of black, vertical blotches; head yellowish olive, darker above, and reticulating with narrow brown lines, these appearing as parallel bars on the upper lip, and radiating from the median line on the upper side of the head; vertical fins light grayish, with black spots, which appear as wavy bars on the caudal fin; pectorals and ventrals pale, the former with a yellowish shade at base; orbital tentacles bright red. Pacific coast of Mexico. Known from 4 specimens (the largest  $2\frac{1}{4}$  inches in length), taken in a deep rock pool at Mazatlan. Two others taken by the *Albatross* from Clarion Island. (*χιών*, snow; *στριχτός*, spotted.)

*Salaria chiostictus*, JORDAN & GILBERT, Synopsis, 363, 1883, Mazatlan, Mexico. (Coll. Jordan & Gilbert.)

2750. ENTOMACRODUS MARGARITACEUS (Poey).

Head 5 in total length with caudal; depth  $6\frac{1}{2}$ . D. XII-14; A. I, 14; eye  $4\frac{1}{2}$  in head, well forward. Body large, snout abruptly decurved; mouth very low, maxillary reaching anterior nostril, which has a little tentacle; (canines small); small tentacle over eye; gill membranes broadly connected, free from isthmus; dorsal deeply emarginate, almost divided; anal beginning under middle of body without caudal, and anal papillae and caudal

rounded; ventrals short; lateral line present anteriorly, no tentacles on nape. Color brown, with 2 vertical bands of a dusky silvery; a central point in each band shining bright. One specimen,  $2\frac{1}{2}$  inches long. Cuba. (Poey.) Perhaps a *Salarias*.

We have the following notes on a specimen, possibly the type of this species, sent by Poey to the museum at Cambridge: Head  $4\frac{2}{3}$ ; depth 5. D. XII-15; A. 18. Body slender. Interorbital concave. Head short, blunt, almost round; a small cirrus over the eye, none on nape. Canines present, small. Body with about 6 dark cross bars besides pearly spots and various markings. Dorsal divided nearly to base. Closely resembles *Salariichthys textilis*. (*margarita*, *μργαρίτης*, pearl.)

*Salarias margaritaceus*, POEY, Memorias, II, 289, 1861, Cuba. (Coll. Poey.)

2751. ENTOMACRODUS DECORATUS, Poey.

Head 5 in total length with caudal; depth 5. D. XII-19; A. 15; P. 14. Eye very high; anterior nostril prolonged in a tube; nape following a straight line to the posterior nostril, profile thin, following a straight and oblique line to mouth, which is very low and short, the maxillary reaching posterior nostril. Lower jaw shorter. Teeth movable, numerous, incurved, close set, in 1 row. (Canines not described.) No cilia on head. Dorsals of equal length, the soft rays more elevated; anal similar to second dorsal; pectoral broad, its lower rays thickened; caudal rounded. Color brownish yellow; the body with darker cross bands, which begin below the middle of the first dorsal, alternating with narrower spaces of the ground color; along the middle and edges of the bands vertical rows of sky-blue spots; in the pale interspaces below the lateral line, which is much curved, a white spot; 3 pale spots placed obliquely below the eye; rays of dorsal and caudal dotted with black. One specimen, 2 inches long. Cuba. (Poey.) Not seen by us; perhaps a *Salarias*. (*decoratus*, decorated.)

*Entomacrodus decoratus*, POEY, Synopsis, 398, 1868, Cuba. (Coll. Poey.)

2752. ENTOMACRODUS NIGRICANS, GILL.

The elongated body, from the snout to the end of the caudal fin, is between 7 and 8 times longer than it is high at the pectorals. Its height at the caudal is about  $\frac{1}{3}$  of the same length. The head is subquadrate, and forms  $\frac{1}{3}$  of the total length. Its greatest height equals  $\frac{1}{3}$  of its length. Its sides decline obliquely outward and downward. The first dorsal commences near the nape, and 2 of its rays are in advance of the pectorals. The second dorsal commences immediately behind the first, and nearly over the fourth ray of the anal, it ceases some distance from the base of the caudal. The anal is more uniform in height than the dorsal, and ceases before it does. The caudal forms less than  $\frac{1}{3}$  of the total length. D. XI-15; A. 17; P. 15; V. 3. The general color of the body and fins is blackish. West Indies. A single specimen was caught in shallow water, at the island of Barbados, near Bridgetown. (Gill.) Not seen by us. (*nigricans*, blackish.)

*Entomacrodus nigricans*, GILL, Proc. Ac. Nat. Sci. Phila. 1850, 168, Barbados. (Coll. Dr. Gill.)

## 893. SALARIICHTHYS, Guichenot.

*Salariichthys*, GUICHENOT, Mém. Soc. Sci. Nat. Cherbourg, XIII, 1867, 96 (*textilis*).

This genus differs from *Entomacrodus* in the presence of teeth on the vomer; dorsal deeply notched; cirri present over eye and on nape; posterior canines small. (*Salaris*; ἰχθύς, fish.)

## 2753. SALARIICHTHYS TEXTILIS (Quoy &amp; Gaimard).

D. XII, 16; A. 18. A few bluntish teeth on vomer; tentacles very small, fringed over nostril and eye, simple on neck; canines quite short; depth  $4\frac{3}{4}$ ; head  $4\frac{3}{4}$ ; pectoral short, little longer than head; gill membranes broadly united, free from isthmus; dorsal notched almost to base, free from caudal; orbital filament  $\frac{1}{2}$  eye. Olive, with 13 silvery cross streaks, not  $\frac{1}{2}$  as wide as the dark interspaces, some of the cross streaks Y-shaped; both dorsals with cross markings, the second with 12 or 13 streaks of dark obliquely upward and backward, alternately with similar pale streaks; cross bars on sides bent in middle, extending up and back and down and back from middle line parallel with muscular impressions; sides with some obscure pale dots; caudal barred with 7 dark bars; anal darkest mesially; lower side of head with dark streaks radiating from the isthmus; bars at chin Y-shaped, upper part of head with darker markings; pectoral nearly plain; a dusky area at base below which is a dusky spot; marblings at base of dorsal. West Indies, from Bermudas to Brazil. Here described from a specimen from Abrolhos Islands (Coll. *Albatross*). This specimen agrees fairly with the account given by Jenyns, but Jenyns describes 5 bars on the tail. It also agrees fairly with the account of the Bermuda specimens given by Goode. It is evidently the *Salaris vomerinus* of Cuvier & Valenciennes, and probably their *textilis* also; but their description of the latter does not apply very well to the coloration of our specimen. (*textilis*, woven.)

*Salaris textilis*, QUOY & GAIMARD MS., CUVIER & VALENCIENNES, Hist. Nat. Poiss., xi, 307, 1836, Ascension Island (Coll. Quoy & Gaimard); GÜNTHER, Cat., iii, 248, 1861; GOODE, Bull. U. S. Nat. Mus., v, 29, 1876.

*Salaris vomerinus*, CUVIER & VALENCIENNES, Hist. Nat. Poiss., xi, 349, 1836, Bahia. (Coll. Blanchet.)

*Salariichthys textilis*, JORDAN, Proc. U. S. Nat. Mus. 1890, 320.

## 894. OPHIOBLENNIUS, Gill.

*Blennophis*, VALENCIENNES, in WEBB & BERTHELOT, Poiss. Îles Canar., 60, 1844 (*webbii*; not

*Blennophis*, CUVIER & VALENCIENNES, a genus of *Olininae*).

*Ophioblennius*, GILL, Proc. Ac. Nat. Sci. Phila. 1860, 103 (*webbii*; substitute for *Blennophis*).

Body oblong, strongly compressed, scaleless; snout short, high, abruptly decurved anteriorly; symphysis of lower jaw of 4 hooked canines, the outer strongest and bent backward, almost forming a right angle; sides of lower jaw with 2 or 3 still larger canines, the hindmost very large and bent backward; upper jaw with 4 slender canines in front, followed by a long row of shorter, slender, movable teeth, which are set close together; nasal tentacle digitate; a low, simple tentacle above eye; gill openings wide. Dorsal fin long, the spines slender, separated by a slight notch from

the soft rays; caudal lunate or forked, free from dorsal and anal; ventrals small, I, 2; lateral line incomplete; pectorals large. A strongly marked genus, perhaps more nearly allied to *Blennius* than to *Emblemaria* or *Chenopsis*. (*ὄφις*, snake; *Blennius*, in allusion to the fang-like teeth.)

a. D. X, 20; A. 20; depth  $5\frac{1}{2}$  in length.

WEBBII, 2754.

aa. D. XI, 22; A. II, 23; depth  $4\frac{1}{2}$  in length.

STEINDACHNERI, 2755.

2754. OPHIOBLENNIUS WEBBII (Valenciennes).

Head 5; depth  $5\frac{1}{2}$ . D. X, 20; A. 20; P. 16. A slender tentacle above eye in front, and a much broader one, divided into 4 to the base, above the nostril. Snout obtuse, nearly vertical at tip; eye large; 4 teeth at end of upper jaw, strongly pointed, curved backward like hooks; lower jaw with 4 teeth at tip, the two middle ones like upper teeth, the two outer hidden and turned backward; a little recurved tooth on side of lower jaw; caudal fin forked; dorsal somewhat notched at the last spine; lateral line ending near middle of body. Olive green, light or dark; dorsal and anal dusky violet, the base pale; back and sides often with fine points; a dark spot behind eye; the silvery swim bladder showing through sides of belly. (Steindachner). Tropical Atlantic; known only from the Canaries and Barbados; not seen by us. (Named for P. B. Webb, one of the explorers of the Canary Islands.)

*Blennophis webbii*, VALENCIENNES, in WEBB & BERTHELOT, Îles Canar., Poiss., 60, pl. 20, f. 1, 1844, Fortaventura, Canary Islands (Coll. Webb); "caught in myriads at Puerto de Cabras in August, eaten as Anchovias" (Webb); GÜNTHER, Cat., III, 259, 1861; STEINDACHNER, Ichth. Notizen, VI, 48, 1867.

*Ophioblennius webbi*, JORDAN & GILBERT, Synopsis, 756, 1883.

2755. OPHIOBLENNIUS STEINDACHNERI, Jordan & Evermann, new species.

Head 4 to  $4\frac{1}{2}$ ; depth 4 to  $4\frac{1}{2}$ . D. XI, 22; A. II, 23; V. I, 2; P. 15. Head much compressed; eye  $3\frac{1}{2}$  in head; snout  $4\frac{1}{2}$ . Dorsal beginning above gill opening, ending just before caudal, its soft rays somewhat higher than the spines, the highest spine  $1\frac{1}{2}$  in head; caudal and pectorals each about as long as head; ventrals  $1\frac{1}{2}$  in head. Dark golden brown, sometimes with a broad cross band of dusky violet on back and dorsal fin; caudal with 2 dark longitudinal stripes; dorsal and anal purplish or orange; an intense, round, dark, ocellated spot behind eye. (Steindachner.) West coast of Mexico; not seen by us; recorded from near Mazatlan and the Tres Marias Islands. (Named for Dr. Franz Steindachner.)

*Blennophis (Ophioblennius) webbi*, STEINDACHNER, Ich. Beitr., VIII, 41, 1870, 5 specimens 70 mm. long, from Navidad near Mazatlan and the Tres Marias Islands.

*Ophioblennius steindachneri*, JORDAN & EVERMANN, Check-List Fishes N. and M. A., 472, 1896, name only, Tres Marias Islands; after STEINDACHNER.

895. EMBLEMARIA, Jordan & Gilbert.

*Emblemaria*, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1883, 627 (*nivipes*).

Body slender, not eel-shaped, compressed, scaleless. Ventrals present, jugular, each of 1 spine and 2 soft rays. A single high dorsal fin beginning on the nape and extending to the caudal, with which it is not conflu-

ent; no notch between spinous and soft parts. Head cuboid, compressed, narrowed anteriorly. Symphysis of lower jaw forming a very acute angle. A single series of strong, blunt, conical teeth on each jaw, and on vomer and palatines. Vomer and palatine teeth larger, their series continuous, parallel to the series in upper jaw. No cirri at the nape; sometimes a cirrus on upper part of eyeball. Gill openings very wide, the membranes broadly united below, free from the isthmus. Lateral line obsolete. This genus bears some resemblance to *Blennius*, but the dentition is entirely different, approaching that of *Chonopsis*. Tropical America, in rather deep water. (*Emblema*; ἐμβλημα, a banner.)

a. Eye without cirrus.

b. Depth 5 in length; dorsal rays 33; ventrals not pure white. ATLANTICA, 2756.

bb. Depth 7 in length; dorsal rays 37; ventrals pure white. NIVIPES, 2757.

aa. Eye with a long cirrus on eyeball above pupil; ventrals dusky; maxillary not extending beyond eye. OCULOCIRRIS, 2758.

2756. EMBLEMARIA ATLANTICA, Jordan & Evermann, new species.

Head  $3\frac{3}{4}$ ; depth 5. D. 35; A. 24; P. 15; V. 3. Body slender, compressed; head heavy; snout evenly decurved; mouth large, horizontal, reaching back of eye. Jaws with short, strong, incurved conical teeth. Fin rays long and filamentous, the longest dorsal rays as long as head; anal rays shorter. Coloration faded in the type, but traces of about 7 broad brown vertical bars as broad as eye and twice as broad as the pale interspaces, the dark bars extending upon dorsal fin; ventrals pale. Gulf of Mexico. Known from 1 specimen,  $3\frac{3}{4}$  inches long, taken from the stomach of *Neomantis aya*, on the Snapper Banks off Pensacola, Florida; very close to *E. nivipes*, but more robust, with fewer dorsal rays.

*Emblemaria atlantica*, JORDAN & EVERMANN, Check-List Fishes, 472, 1896, name only, Snapper Banks off Pensacola, Florida. (Type, No. 33915. Coll. Silas Stearns.)

2757. EMBLEMARIA NIVIPES, Jordan & Gilbert.

Head  $3\frac{3}{4}$  in length; depth 7. D. XXIII, 14; A. 25. Body everywhere equally compressed, posteriorly tapering; head wider than body, of about equal depth, with very short, subvertical, sharply compressed snout; eyes very large, approximated above, with some vertical range; orbital ridges sharply raised above, the interorbital region very narrow, channeled, about equaling diameter of pupil; eye  $3\frac{3}{4}$  in head. Gape very wide, horizontal, low, reaching much beyond eye, the maxillary about  $\frac{1}{2}$  head, not produced beyond angle of mouth; intermaxillaries separated by a groove from the snout, this groove continuous for the entire length of the upper jaw, maxillary not evident, apparently adnate to the skin of the preorbital. First dorsal spine inserted over margin of preopercle; spines all very slender and flexible, the posterior but weakly differentiated from the soft rays, the anterior portion of fin very high, the spines filiform, not exerted beyond the membrane; the longest dorsal spine about  $\frac{1}{2}$  length of body, the last spine about  $\frac{1}{2}$  head; membranes of last rays of both dorsal and anal slightly joined to base of caudal. Front of anal nearer snout than

base of caudal by a distance equaling  $\frac{1}{4}$  length of head. Caudal  $\frac{3}{4}$  length of head; ventrals and pectorals slightly less. Color in spirits, sides dark brown, with 8 to 10 lighter vertical bars of variable width; body lighter below; obscure cross bands on lower side of head; dorsal blackish anteriorly, whitish behind, with membrane at intervals of every second, third, or fourth ray dusky; caudal light at base, its tip blackish; anal dusky translucent; ventrals bright white, the basal portion dusky. Pearl Islands, near Panama. A specimen 2 inches long is the type of the species. Numerous smaller specimens were obtained at the same time. (*nix, nixis, snow; pes, foot.*)

*Emblemaria nivipes*, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1883, 627, Pearl Islands, near Panama. (Type, No. 29676. Coll. Prof. Frank H. Bradley.)

2758. EMBLEMARIA OCULOCIRRIS, Jordan.

Head  $3\frac{3}{4}$ ; depth  $6\frac{3}{4}$ . D. about 35; A. 25. Upper part of eyeball above pupil (sclerotica) with a slender cirrus tipped with black, this nearly as long as eye; eye longer than snout, about  $3\frac{3}{4}$  in head, the maxillary extending to below posterior part of pupil; snout sharper than in *Emblemaria nivipes*,  $\frac{3}{4}$  eye; teeth small, rather sharp, directed backward; longest dorsal spines as long as head; pectorals  $1\frac{1}{2}$  in head; ventrals  $1\frac{2}{3}$ , inserted before pectorals. Color in spirits, brown, with traces of about 9 blackish cross bars, which are separated on the back by whitish, quadrate interspaces; a white spot at nape; some dusky below eye; dorsal dusky, the pale bars of back extending on its base; anal dusky; ventrals blackish; caudal pale, its tip black; pectorals pale. Gulf of California. Known from 1 specimen,  $1\frac{1}{2}$  inches long, from La Paz. It is shriveled and in poor condition. It seems to be very close to *Emblemaria nivipes*, but differs in the presence of an ocular cirrus, in the sharper snout, smaller mouth, and dusky ventrals. The teeth seem rather more slender, but can not be well examined. (*oculus, eye; cirrus, filament.*)

*Emblemaria oculocirris*, JORDAN, in GILBERT, Proc. U. S. Nat. Mus. 1896, 456, La Paz. (Type, No. 47749. Coll. Albatross.)

896. CHÆNOPSIS, Gill.

*Chænopsis*, GILL, Ann. Lyc. Nat. Hist. N. Y., VIII, 1865, 141 (*ocellatus*).

Body naked, eel-like. Head much elongate, quadrate behind, conic in front, profile straight; snout acute, jaws produced; no teeth on vomer, teeth in front of jaws strong, with villiform teeth behind them. Dorsal and anal long, continuous, confluent with the caudal. Dorsal rays about XVIII, 38; anal II, 38. Ventrals inserted slightly before pectorals. West Indies. (*χαίρω, to yawn; ὄψις, face.*)

2759. CHÆNOPSIS OCELLATUS, Poey.

D. XVIII, 38; A. II, 38; C. 15. Body naked, eel-like; anus submedian. Head much elongate, quadrate behind at the opercular region, conic in front, with the profile rectilinear and the snout acute; eyes moderate;



mouth large, with the cleft wide and nearly horizontal. Teeth subcylindrical, in a uniform row, behind which, in front, there is a broad band of villiform teeth; on the palatine bones, uniserial and obtusely subcylindrical like those of the jaws; the palatine rows are parallel; vomer edentulous. Gill membranes confluent below, free from the isthmus. Dorsal and anal long, confluent with caudal; ventrals slightly in advance of pectorals, with 2 or 3 rays. (Gill.) Matanzas, Cuba; 1 specimen, examined by us in the National Museum. (*ocellatus*, having eye-like spots.)

*Chaenopsis ocellatus*, POEY, in GILL, Ann. Lyc. Nat. Hist. N. Y., VIII, 1867, 143, Matanzas, Cuba. (Coll. Poey.)

#### 897. LUCIOBLENNIUS, Gilbert.

*Lucioblennius*, GILBERT, Proc. U. S. Nat. Mus. 1890, 103 (*alepidotus*).

Body very elongate, wholly naked; gill membranes broadly united, free from isthmus; dorsal fin single, extending along the entire back, its anterior half spinous. Ventrals in front of pectorals, I, 2. First two anal rays spinous. Last rays of dorsal and anal joined to caudal. Teeth conic, not movable, in jaws and on vomer and palatines. Lateral line not described. A strange genus, evidently very close to *Chaenopsis*. (*Lucius*, pike; *Blennius*, blenny.)

#### 2760. LUCIOBLENNIUS ALEPIDOTUS, Gilbert.

Head 3 in length; depth  $3\frac{1}{2}$  in head. D. XVIII, 32; A. II, 30. Body much compressed, slender throughout, the head rather deeper and wider than body. Snout long, depressed, and rather wide, the anterior profile descending very gradually. Mouth nearly horizontal, the lower jaw protruding, the gape extending to much behind orbit, the entire physiognomy remarkably pike-like. Snout 4 in head; maxillary  $1\frac{1}{2}$ ; eye  $4\frac{2}{3}$  to 5. Teeth in a villiform band in upper jaw, the outer series slightly larger; in lower jaw in a single series laterally, widening into a patch anteriorly, the outer enlarged; a few teeth only on vomer; palatines with a long and rather broad patch similar to those in jaws. Dorsal fin beginning on the nape in advance of middle of opercle, the fin uniformly low, extending the whole length of back, the posterior ray joined by membrane with the caudal; the spines and rays are similar in appearance, flexible and simple, none of the soft rays branched; the spines are more slender, and show no joints, the articulations being present in small number on all the soft rays; the highest ray is less than diameter of orbit; anal and caudal rays similar to those of soft dorsal; caudal short, rounded; origin of anal midway between tip of snout and end of caudal fin, its first 2 rays spinous; ventrals under opercular margin, of 1 spine and 2 well-developed rays, nearly  $\frac{1}{2}$  as long as head; pectorals narrow, of apparently unbranched rays, about  $\frac{1}{2}$  as long as head. Color light olivaceous, with 11 vertical dark blotches on sides, most of which divide to form on middle of sides double vertical bars; top and sides of head with dark cloudings, and with numerous black specks of varying size; middle of sides and base

of dorsal with numerous pearly dots nearly as large as pupil; branchiostegal membrane black posteriorly; the lateral bars extended to base of dorsal, the anterior ones usually forming conspicuous black blotches which extend well up on the fin; other fins unmarked. Length  $1\frac{1}{2}$  inches. Gulf of California; two specimens from Albatross Station 3005, in 21 fathoms. (*ἀλεπίδωρός*, scaleless.)

*Lucioblennius alepidotus*, GILBERT, Proc. U. S. Nat. Mus. 1890, 103, Lower California (Coll. Albatross); JORDAN, Proc. Cal. Ac. Sci. 1896, 233, pl. 37.

898. PHOLIDICHTHYS, Bleeker.

*Pholidichthys*, BLEEKER, Boeroe, 406, 1856 (*leucotenia*).

Body elongate, tapering, naked; snout obtuse; no cirri; teeth unequal, on jaws only; dorsal, anal, and caudal fins distinct, but connected by a membrane; the dorsal formed of flexible spines; the soft rays, if present, not distinguishable from them; ventrals inserted scarcely before the pectorals, of 2 rays. Lateral line and vertebrae undescribed. Tropical parts of the Pacific. (*Pholis*; *ιχθύς*, fish.)

2761. PHOLIDICHTHYS ANGUILLIFORMIS, Lockington.

Head  $6\frac{1}{2}$  in total length with caudal; depth 16. Body exceedingly elongate, much compressed, naked; upper profile of head forming a continuous convex curve to the tip of the snout, which is about equal in length to the eye. Eye lateral, round; interorbital space about  $\frac{1}{3}$  of the diameter of the eye, convex transversely. Posterior extremity of maxillary vertical with the hinder margin of the eye. Tip of snout a little below the level from the center of the eye; mouth moderately oblique, lower jaw slightly the longer. Teeth of lower jaw in a close-set row, the largest in front, diminishing along the sides; teeth of upper jaw similar, but smaller; palate smooth. Vertical fins continuous, but distinct; dorsal entirely spinous; anal commencing a little behind the middle of the entire length of the fish; ventrals 2-rayed, very slightly in advance of the pectorals, which are about equal in length to the distance of their base from the eye. Color in spirits, dark blackish brown mingled with white upon top, sides, and lower parts of head; interorbital area and top of snout white. Gulf of California; a single specimen dredged off San Jose Island, Amortiguado Bay. Total length  $1\frac{1}{2}$  inches. Head  $\frac{1}{4}$  inch. The example is broken across, the branchiostegals are defective, the caudal fin broken and some fin rays missing, so that the fin formula can not be exactly given. The dorsal fin has above 60 rays. The body is much more slender than that of *P. leucotenia*, Bleeker, and there is no trace of the longitudinal bluish-white band of that species. (Lockington.) (*Anguilla*, eel; *forma*, shape.)

*Pholidichthys anguilliformis*, LOCKINGTON, Proc. Ac. Nat. Sci. Phila. 1881, 118, San Jose Island, Lower California. (Coll. W. J. Fisher.)

## 899. PSEDNOBLENNIUS, Jenkins &amp; Evermann.

*Pseudobleennius*, JENKINS & EVERMANN, Proc. U. S. Nat. Mus. 1888, 156 (*hypacanthus*).

Body compressed, elongate, naked; head short, blunt; no cirri; mouth large, the jaws subequal; teeth in a single series in each jaw, none on vomer or palatines; lateral line not developed. Dorsal fins 2, the first at the nape, of three flexible spines; second dorsal with a few slender spines which pass into the soft rays; anal much shorter than second dorsal, both fins joined to base of caudal; dorsal rays III-34; anal 27; ventral rays 2, the fin directly below pectorals. Apparently close to *Pholidichthys*, but with the dorsal divided and changing gradually from spines to soft rays. ( $\psi\epsilon\delta\nu\acute{o}\varsigma$ , naked; *Blennius*.)

## 2762. PSEDNOBLENNIUS HYPACANTHUS, Jenkins &amp; Evermann.

Head  $4\frac{1}{2}$  (5 in total); depth 7 (8); eye 4, equal to snout; B. G. D. III-34; A. 27. Body greatly compressed, elongate; head short, snout blunt, about equal to eye; anteorbital profile very steep, gently rounded from front of eye to first dorsal, from there nearly straight to caudal; ventral line nearly straight. Body naked, no membranaceous appendages. Mouth large, horizontal, jaws subequal, extending to beyond middle of eye. Teeth in a single series in each jaw, well developed, pretty uniform in size, slightly projecting backward; vomer and palatines apparently smooth. Eye large, equal to twice interorbital space, high up. Dorsal fin 2, the first of 3 very slender, flexible spines, hard to distinguish from soft rays, but they do not appear to be at all jointed. This fin is inserted upon the nape immediately above the posterior edge of the preopercle, and a distance in front of second dorsal nearly equal to length of snout, its very soft spines equal distance from end of snout to posterior rim of orbit; second dorsal begins directly over origin of pectorals and extends to caudal, with which it is slightly connected; first few rays of second dorsal very weak, flexible spines, the last few pretty evidently soft, jointed rays, while the intermediate ones are not distinguishable as definite spines or soft rays—in short, there seems to be a gradual change from spines to soft rays from the anterior to the posterior part of the fin. This character, if we mistake not, is entirely unique. The fin is of nearly uniform height, the rays about equaling those of the first dorsal in length; anal similar to second dorsal in shape and height, but much shorter, its origin being much behind that of the second dorsal or nearly halfway from the snout to base of caudal; posteriorly it extends coterminously with the dorsal, and, like it, is slightly joined to the caudal fin; caudal fin apparently rounded, fan-shaped, but its shape can not be exactly made out, as some of its rays are broken off; pectorals inserted below axis of body, directly over ventrals, their length about  $\frac{1}{2}$  that of head; ventrals of 2 rays, inserted under pectorals, about equal to pectorals in length; body entirely scaleless. Coloration in alcohol, pale, mottled with five dark points so arranged as to inclose circular areas with fewer spots; a long dark blotch behind the axil, inclining downward and backward; head covered with similar punctulations; opercles dusky; chin with 2

dark cross lines, separated by 1 of white, extending onto upper jaw on each side; top of head with a purple spot; sides with a series of about 6 short black lines, the last broadest and plainest; base of caudal with a distinct black blotch; first dorsal quite dark, almost black; second dorsal with about 8 pretty well-defined dark blotches at its base, rest of fin with numerous dark spots of different sizes; anal with about 12 dark blotches extending somewhat regularly from the base slightly forward, these separated by plain unmarked spaces of a little greater width; caudal sparingly marked with dark points arranged in wavy cross bars; pectorals and ventrals unmarked. Gulf of California at Guaymas. A single specimen, 1½ inches long, obtained from a shallow arm of the bay. (Jenkins & Evermann.) (*ὄπρo*, below (imperfect); *ἀκάρθα*, spine.)

*Pseudoblennius hypacanthus*, JENKINS & EVERMANN, Proc. U. S. Nat. Mus. 1888, 156, Guaymas, Mexico. (Type, No. 39638. Coll. Jenkins & Evermann.)

900. STATHMONOTUS, Bean.

*Stathmonotus*, BEAN, Proc. U. S. Nat. Mus. 1885, 191 (*hemphillii*).

Body moderately long and low, much compressed; head small, compressed, naked; mouth small, oblique; conical teeth in both jaws, in 2 series, the outer slightly enlarged and, in the upper jaw, somewhat recurved; a few teeth on the vomer. Gill membranes, as in *Pholis*, broadly united, free from the isthmus. Scales none. No lateral line. Dorsal fin long and low, beginning near the head, and consisting entirely of stiff, sharp spines, which are very short anteriorly and gradually increase in size posteriorly. Anal similar to dorsal, with 2 spines and many soft rays. Caudal short, rounded, scarcely separated from the dorsal and anal; pectorals small, much smaller than in *Pholis*, containing only a few rays; ventrals better developed than in *Pholis*, their position more anterior, consisting of a spine and 2 rays. Pseudobranchiae absent. Branchiostegals 5. Coast of Florida. (*στᾶθμῆ*, a carpenter's rule; *ὄπρo*, back.)

2763. STATHMONOTUS HEMPHILLII, Bean.

Head 7; depth 8 to 8½; D. LI; A. II, 27; V. I, 2; P. 5 or 6; eye 6 in head. Maxillary extending about to vertical through hind margin of eye; jaws subequal, or the lower projecting very slightly beyond upper; eyes small, separated by an interspace about equal to their own length, and very slightly greater than length of snout; pectoral very little more than ½ as long as head, and scarcely as long as ventral; dorsal beginning over posterior end of pectoral, its anterior spines very much shorter than the posterior ones; length of caudal about equal to length of postorbital part of head; vent slightly in advance of middle of total length to base of caudal, and about under the twentieth dorsal spine. Colors from the alcoholic specimen: A white line extending from tip of snout to caudal, divided into small segments by short cross bars, the first 2 on the head, and the last at origin of caudal; posteriorly, these short bars extend downward, terminating slightly below the base of the dorsal fin; several white blotches, simulating bars, on posterior half of anal fin; edge of

caudal white; sides and under surface of head with several whitish oblique bands forming V-shaped markings; a few roundish white blotches on sides of head, the most conspicuous behind eye; general color darkish brown, nearly black. Length about 2 inches. Key West; 2 specimens known. (Bean.) (Named for the collector, Henry Hemphill.)

*Stathmonotus hemphilli*, BEAN, Proc. U. S. Nat. Mus. 1885, 191, pl. 13, Key West, Florida. (Coll. Henry Hemphill. Type, No. 37193, U. S. Nat. Mus.)

#### 901. BRYOSTEMMA, Jordan & Starks.

*Bryostemma*, JORDAN & STARKS, Proc. Cal. Ac. Sci. 1895, 841 (*polyactocephalum*).

Body moderately elongate, covered with small scales; snout short; no teeth on vomer or palatines; teeth in jaws small; gill membranes united, free from the isthmus; nostrils, orbital regions, and neck with dermal flaps, the supraorbital flaps high. Dorsal fin long, of spines only; pectorals well developed, more than half length of head; ventrals well developed, jugular; caudal fin distinct. No air bladder or pyloric caeca. No true lateral line; a short series of large pores above pectoral. North Pacific, representing *Chirolophis* of the Atlantic. This genus differs from the European genus, *Chirolophis*, Swainson (*Blenniois*, Nilsson), in the absence of a true lateral line. Dr. Boulenger informs us that a true median lateral line is developed in *Chirolophis ascanii*. (*βρύον*, moss; *στέμμα*, crown.)

a. Dorsal with about 60 spines; anal with about 55 soft rays; a black spot on anterior part of dorsal, but no ocelli posteriorly. POLYACTOCEPHALUM, 2764.

aa. Dorsal with about 54 spines; anal with 40 soft rays; dorsal with several black ocelli, most distinct posteriorly. NUGATOR, 2765.

#### 2764. BRYOSTEMMA POLYACTOCEPHALUM (Pallas).

Head  $6\frac{1}{2}$ ; depth 6. D. LXI; A. 55 (51 to 57); P. 14; V. I, 3; lateral series with 9 to 15 pores. Body elongate, much compressed, covered with small, smooth, embedded scales. Head very short, blunt in profile; mouth short, terminal, the maxillary 3 in head; lower jaw heavy, projecting, its tip with 2 small slender cirri, which are pale in color; teeth subequal, small, bluntish, close set, in 1 row in each jaw; eyes 4 in head, near together; the snout 4; supraorbital cirri  $2\frac{1}{2}$  in head; interorbital space flat; a flat fringed cirrus over front of eye, these 2 joined at base, about 3 in head; a small cirrus about  $\frac{1}{2}$  length of this over posterior part of each eye, these 5 to 6 in head; top of head and nape covered with series of erect cirri, the longest nearly as long as eye; about 15 minute cirri along dorsal edge of lateral line, 1 on each pore. Rows of pores running around eye, under preopercle, and along entire length of the short lateral line; lateral series of pores  $\frac{1}{2}$  length of head; gill rakers not developed. Dorsal fin beginning over pectoral and running to caudal; anterior rays fringed with fleshy cirri; first ray, including cirri, 2 in length of head; anal beginning close behind vent and running to caudal, to which it is joined at base; distance from tip of snout to vent nearly 3 in body; pectoral fin but little shorter than head, its breadth at base not  $\frac{1}{2}$  its length. Color in spirits, pale

brownish, plain or mottled with darker, with about 13 dark blotches along dorsal and anal fins, more distinct on dorsal; a black spot on fourth to sixth dorsal spines very distinct; a faint one on anterior part of anal; a few dark markings about head and nape; cirri mostly pale. Bering Sea, south to Puget Sound and Yezo. Here described from a fine specimen,  $6\frac{1}{2}$  inches long, from Port Orchard, near Seattle, collected by Prof. O. B. Johnson. Other specimens before us from St. Paul (Pribilof Islands), from *Albatross* Stations 3213 and 3274, south and north of the Peninsula of Alaska, and from Petropaulski Harbor, Kamchatka. These specimens show a great deal of variation, and possibly represent 3 different species. It is more likely, however, that they represent extremes of variation. Young examples, collected by the *Albatross* in eastern Bering Sea, are more elongate and less compressed; body much mottled and vaguely barred; ventral fins checkered in fine pattern; head sand color; a black blotch on fourth to sixth dorsal spine; anterior dorsal spine little elevated and with few fringes; sides of head without cirri; anterior cirri joined almost to the tip, a little shorter than the posterior cirri, which are long and very slender. In 1 specimen of these, however, the cheeks are covered with densely matted cirri extending from the angle of the mouth to the dorsal. In these examples the anterior cirri are short and separate, about as long as the posterior cirri. The larger example, 75 cm. long, from Petropaulski, is evidently the typical *polyactcephalum*, and corresponds perfectly to Herzenstein's account of *B. japonicum*. It shows the following characters: Head  $6\frac{1}{2}$ ; depth  $5\frac{1}{2}$ . D. LXI; A. 45; P. 14; V. I, 3; lateral series with 6 pores. Body a little deeper than in Puget Sound examples; head short, blunt in profile; mouth short, terminal, oblique, the maxillary  $2\frac{3}{4}$  in head; lower jaw heavy, projecting, its tip with 2 broad fringed flaps of a dark color; eyes 4 in head, close together, the interorbital space concave; a fringed cirrus above each eye in front, the 2 connected with each other only in the thickened skin at base; a similar cirrus over each eye behind; the posterior cirri  $\frac{1}{2}$  longer than the anterior ones,  $2\frac{1}{2}$  in head; top of head and nape with similar cirri, none of them longer than pupil; a few small cirri on cheeks and opercles; some along lateral series of pores, which is  $2\frac{1}{2}$  in head; anterior rays of dorsal fringed with fleshy cirri, the first 2 in head; distance from snout to vent  $2\frac{3}{4}$  in body; pectorals nearly as long as head, the rays thickened in the adult, the base of the fin about  $\frac{1}{2}$  its length. Color very dark brown, with vague cross bands and many spots; dorsal and anal each with a broad black edge; other fins all black, the caudal barred. Perhaps the dark coloration and long cirri are characters of the adult male. (*πολύς*, many; *κύρις*, ray; *κεφαλή*, head.)

*Blennius polyactcephalus*, PALLAS, Zool. Rosso-Asiat., III, 179, 1811, Kamchatka.

*Chirolophus japonicus*, HERZENSTEIN, Mélanges Biologiques Soc. Sci. Petersb., XIII, 1890, 123, Yezo.

*Chirolophis polyactcephalus*, JORDAN & GILBERT, Synopsis, 765, 1883; BEAN in Nelson, Rept. Nat. Hist. Coll. Alaska, 305, pl. 15, f. 2, 1887.

*Bryostemma polyactcephalum*, JORDAN & STARKS, Proc. Cal. Ac. Sci. 1895, 841; JORDAN & GILBERT, Rept. Fur Seal Invest., 1898.

## 2765. BRYOSTEMMA NUGATOR, Jordan &amp; Williams.

Head  $5\frac{1}{2}$ ; depth  $5\frac{1}{2}$ ; D. LIV; A. 41; V. I, 3; pores of lateral line 25. Body elongate, less compressed than in *Bryostemma polyactoecephalum*, covered with small, smooth, embedded scales. Head short, very obtuse, almost truncate; top of head from nostrils to near front of dorsal covered with fleshy cirri, much smaller than in *B. polyactoecephalum*; only 2 or 3 small ones extending on first dorsal spine; supraorbital cirrus short, 4 to 5 in head; 2 small cirri placed at the sides of snout with a larger median one behind them, forming a triangle; jaws equal; mouth horizontal, the angle extending to below pupil; eyes small, 4 in head; snout very short, almost vertically truncate,  $\frac{3}{8}$  in eye; teeth of both jaws subequal, short, bluntish, and close set. Lateral line short,  $7\frac{1}{2}$  in length of body, concurrent with the dorsal outline of body. A line of pores begins in front of eye on a level with pupil, runs under eye and to a level with pupil again, then back to and along the entire length of the short lateral line. Gill rakers not developed; gill membranes free from isthmus. Vent  $\frac{1}{4}$  distance from tip of snout to tip of caudal; distance from origin of ventral to anus  $4\frac{1}{2}$  in length of body. Pectoral fin  $5\frac{1}{2}$  in body, as long as head. Dorsal fin beginning in front of the pectoral, highest along the posterior half, the longest spine  $2\frac{3}{8}$  in head, the fin higher than anal; dorsal slightly joined to caudal; anal separated from caudal; caudal rounded,  $1\frac{1}{2}$  in head; first dorsal spine  $4\frac{1}{2}$  in head, its surface with 2 or 3 small cirri. Color in spir. its of 1 specimen, probably male, dark brown, with 13 pale cross bars along back, extending on dorsal fin; along sides these become obsolete; on belly they become increased in number and broadened below; dorsal fin with 13 large, very distinct black ocelli with yellowish rings, 1 between each pair of the pale blotches; anal with about 7 small blackish spots at base on posterior part, the fin otherwise nearly plain; caudal faintly barred with light and dark; pectorals pale, with 2 dark pale-edged oblique bars before it; sides of head with irregular dark vertical bars, 1 of them forming an inverted A below eye, this and others extending across lower jaw; cirri mostly black. The other specimen, probably the female, has the body nearly plain brown, the dorsal with but 4 ocelli, the anterior 9 being replaced by dark bars on the fin; anal with dark oblique cross bars; pectorals barred with black; markings on head more sharply defined, coloration otherwise similar. This second specimen is  $4\frac{1}{2}$  inches in length, the other 4. Puget Sound; the above account from the 2 original types from near Seattle. Three others since obtained near Channel Rocks, Port Orchard, show the following life coloration: Dark red above, orange brown below, belly cream color; sides below with cream-colored cross bars, wider than eye, running from the axis of body downward and fading into the general color below; a A-shaped mark downward from eye across branchiostegals to isthmus, a similar mark behind eye across edge of preopercle, this last sometimes broken up and chain-like; top of head dark; snout light; 2 oblique dark bars at base of pectoral; dorsal with 12 or 13 sharp dark brown spots as large as eye, edged with bright red, these arranged regularly along the whole length of fin; pectorals and caudal bright red with wavy, irregular, brown

lines running across the rays; anal red, with dark brown bars as wide as the interspaces running obliquely downward and forward; ventrals light brown. (*nugator*, a fop.)

*Bryostemma nugator*, JORDAN & WILLIAMS, Proc. Cal. Ac. Sci. 1895, 843, pl. 101, Seattle, Washington. (Coll. Young Nat. Soc. Type, No. 3134, L. S. Jr. Univ.)

902. APODICHTHYS, Girard.

*Apodichthys*, GIRARD, Proc. Ac. Nat. Sci. Phila. 1854, 150 (*flavidus*).

Body elongate, compressed, covered with very small scales; no lateral line; snout short; mouth moderate, oblique; teeth in the jaws moderate, stouter anteriorly; vomer with teeth; gill membranes united, free from the isthmus. Dorsal fin long, low, even, of spines only; anal fin similar, preceded by a very large pen-shaped spine channeled along its anterior surface and hidden in a pouch of skin; caudal fin short, connected with dorsal and anal; no ventral fins; pectoral fins moderate; intestinal canal short, without pyloric caeca. Small, bright-colored fishes of the Pacific, living among rocks near shore. (*ἄποιος*, without feet; *ἰχθὺς*, fish; in allusion to the want of ventral fins.)

- a. Color various, green, olive, or scarlet; sides of head without silvery band; depth 7 to 8 in length; head 9. FLAVIDUS, 2766.  
 b. General color olivaceous. var. *flavidus*, 2766a.  
 bb. General color scarlet. var. *sanguineus*, 2766b.  
 bbb. General color grass green. var. *virescens*, 2766c.  
 aa. Color reddish; a bluish silvery stripe on side of head; depth 9 to 10 in length; head 7. UNIVITTATUS, 2767.

2766. APODICHTHYS FLAVIDUS, Girard.

Head  $9\frac{1}{4}$ ; depth  $7\frac{1}{4}$ . D. XCIII; A. I, 40. Head short; mouth very oblique; maxillary reaching pupil; upper jaw with a series of conical teeth, behind which is a patch of smaller teeth; sides of mandible with conical teeth in a single series, forming a patch in front; vomer with 3 conical teeth; palatines toothless; nape equidistant between front of dorsal and pupil. Anal spine very large,  $\frac{2}{3}$  length of head, shaped like a pen, deeply excavated on its anterior side, and very convex behind, very thin, flexible, and with sharp edges, entirely included in a pouch of skin; pectoral fins about  $\frac{2}{3}$  length of head. Color orange, varying with the surroundings to intense grass-green (var. *virescens*), yellowish brown (var. *flavidus*), crimson and dark purple (var. *sanguineus*); a few light round spots along axis of body posteriorly; a narrow black bar downward and backward from eye; a shorter, less distinct bar from upper margin of orbit backward to occiput; anal fin obliquely barred with brownish. Length 18 inches. Pacific coast, Vancouver Island to the Santa Barbara Islands; abundant; usually found below low tide mark. The following color notes are from specimens taken in Puget Sound belonging to the green form (var. *virescens*), the larger 10 inches in length, the smaller 3 inches. The large one is a bright grass-green, mottled with light gray; a series of blended white spots, as large as eye, along the axis of body



from the pectoral fin to the middle of caudal peduncle; belly with many similar spots smaller in size and somewhat sharper in outline; a row of conspicuous black spots, irregular in size, shape, and position, along back at the base of dorsal spines; a black line as wide as pupil from nape to eye, a similar line from eye to posterior end of maxillary; a faint light streak across cheek posteriorly; cheek and base of pectoral dusted with fine dark points. The small one is bright green without distinct markings on body; a silvery bar, running posteriorly from tip of snout through eye, across cheek, to the middle of opercle; no bar downward from eye to maxillary, or from eye to nape as in the large one. (*flavidus*, yellowish.)

*Apodichthys flavidus*, GIRARD, Proc. Ac. Nat. Sci. 1854, 150, Presidio, San Francisco Bay (Coll. Dr. Kennerly. Type, No. 494, U. S. Nat. Mus.); GIRARD, Pac. R. R. Surv., x, Fishes, 117, 1858; GÜNTHER, Cat., 290, 1861; JORDAN & GILBERT, Synopsis, 769, 1883.

*Apodichthys virescens*, AYRES, Proc. Cal. Ac. Nat. Sci. 1855, 55, San Francisco; GIRARD, Pac. R. R. Surv., x, Fishes, 118, 1858.

*Apodichthys inornatus*, GILL, Proc. Ac. Nat. Sci. Phila. 1862, 279, Puget Sound, probably (Coll. Northwestern Boundary Commission); D. XC; A. 38.

*Apodichthys sanguineus*, GILL, Proc. Ac. Nat. Sci. Phila. 1862, 279, California. (Coll. Dr. Samuel Hubbard.)

#### 2767. *APODICHTHYS UNIVITTATUS*, Lockington.

D. about XCV; A. about I, 40. Body elongate, much compressed, band-like, preserving almost same depth to about posterior fifth of body, thence tapering more rapidly to caudal fin. Head 7; depth nearly 10 times in total length; depth of caudal peduncle about  $\frac{1}{2}$  of that of body; snout obtuse, about  $\frac{3}{4}$  as long as diameter of eye, upper profile of head a continuous curve from snout to occiput. Interorbital area highly convex transversely, about equal in width to  $\frac{1}{2}$  diameter of eye. Eye entirely lateral, round, contained entirely in anterior half of head; iris golden. Mouth small, posterior extremity of the maxillary reaching to anterior margin of eye. Teeth small. Branchiostegals 5. Dorsal continuous with, but distinct from, anal, arising vertically from tip of operculum, and composed of spines only. Anal preceded by a long, sharp, slender spine of V-shaped transverse section, hollow side anterior, length of spine equal to about  $\frac{1}{2}$  depth of fish. Distance from anal spine to tip of operculum a little more than to tip of caudal. Caudal with numerous accessory rays, so that its sides are almost straight, posterior margin broken in the type, all rays simple. General color in spirits, light reddish, vertical fins rather bright, and top of head reddish brown; tip of snout brown; a silvery band (possibly bluish in life) from tip of snout, across lower part of eye, cheek, and opercles, terminating at about middle of length of operculum, this band bordered above by a narrower brown band. Lower California, probably from the gulf. A single specimen. Length 1.88 inches. The peculiar vitta upon each side of the head at once distinguishes this species from the other described forms. (Lockington.) Not seen by us. (*uni-vittatus*, having one band.)

*Apodichthys univittatus*, LOCKINGTON, Proc. Ac. Nat. Sci. Phila. 1881, 118, Gulf of California.

903. XERERPES, Jordan & Gilbert.

*Xererpes*, JORDAN & GILBERT, Proc. Cal. Ac. Sci. 1895, 846 (*fucorum*).

This genus differs from *Apodichthys* in the moderate size of the anal spine, which is rounded and not channeled on its anterior edge, and in the small size of its pectoral fins. The single known species lives in *Fucus* chiefly above low-tide mark and may often be shaken out of half-dry mats of seaweed on rocks well above the water. (*ξηρός*, dry; *ἔρπης*, creeper.)

2768. XERERPES FUCORUM (Jordan & Gilbert).

Head 10; depth  $9\frac{1}{2}$ . D. LXXXIII; A. 35. Form and dentition as in *Apodichthys flavidus*. Mouth very oblique, the maxillary reaching center of pupil; nape nearer front of dorsal than end of snout. Anal spine comparatively small, about  $\frac{1}{2}$  length of head, transversely very convex in front, and slightly concave or grooved behind, the pouch of skin at its base little developed; pectorals very small, shorter than eye; anal fin beginning nearer tip of caudal than tip of snout by about 3 times length of head. Bright olive green or deep red, the color varying with the surroundings; a row of dark spots along axis of body, these sometimes with light-bluish center, and connected by a very narrow dark streak; generally a dark streak downward from eye, but no other markings about head. Length 6 inches. Monterey to Puget Sound; abundant in rock pools and bunches of *Fucus*; remarkable for its active movements. It is found mostly in masses of *Fucus* attached to rocks between tide marks, and it is often found at low tide at a considerable distance from any water, kept damp by the masses of algae. Sometimes a dozen of them can be shaken from a bunch of algae attached to a dry rock. It is, like the species of *Xiphidion*, very active, moving over stones or sand, and showing less anxiety about the presence of its native element than any other fish known to us. (*fucorum*; of the *Fucus* or seaweed.)

*Apodichthys fucorum*, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1880, 139, Monterey (Coll. Jordan & Gilbert); JORDAN & GILBERT, Synopsis, 770, 1883.

*Xererpes fucorum*, JORDAN & STARKS, Proc. Cal. Ac. Sci. 1895, 846.

904. ULVICOLA, Gilbert & Starks.

*Ulvicola*, GILBERT & STARKS, Proc. U. S. Nat. Mus. 1896, 455 (*sanctæ-rosæ*).

This genus is allied to *Xererpes*, but differs in having the opercle above angle adnate to shoulder girdle, in the smaller size of the anal spine, and especially in the entire absence of pectoral fins. (*Ulva*, sea lettuce; *colo*, inhabit.)

2769. ULVICOLA SANCTÆ-ROSE, Gilbert & Starks.

Head 10 in body; depth 13. D. XCVII; A. I, 40; eye  $4\frac{1}{2}$  in head; caudal  $1\frac{1}{2}$ . Body elongate, as in *Apodichthys*, strongly compressed, upper profile of head slightly convex, no constriction at nape; mouth very small, oblique, the maxillary reaching about to front of eye; teeth very small in

a single row on jaws; vomer with teeth; interorbital a narrow, sharp ridge; snout about equal to length of eye; gill opening short, limited to the part below angle of opercle, above adnate to shoulder girdle. Origin of dorsal above upper end of gill opening, much nearer occiput than tip of snout; anal spine small, not channeled as in *Apodichthys flavidus*; origin of anal nearer base of caudal than tip of snout by a distance equal to twice length of head; pectorals and ventrals obsolete; caudal rather long, confluent with dorsal and anal. Color in spirits, light brown, slightly lighter under head and on belly; no markings. The type is a specimen  $4\frac{1}{2}$  inches in length, collected by the *Albatross* at Santa Rosa Island, off Santa Barbara, January 6, 1889. (Type, No. 47579. Coll. *Albatross*.)

*Utricola sanctæ-roseæ*, GILBERT & STARKS, Proc. U. S. Nat. Mus. 1896, 455, pl. 55, fig. 2, Santa Rosa Island, California.

## 905. PHOLIS (Gronow) Scopoli.

(GUNNELS.)

*Pholis*, GRONOW, Zoophylacæum, 78, 1765 (not binomial).

*Pholis*, SCOPOLI, Introd. Hist. Nat., 456, 1777 (*gunnellus*).

*Murænoïdes*, LACÉPÈDE, Hist. Nat. Poiss., II, 324, 1800 (*sujef*).

*Centronotus*, BLOCH & SCHNEIDER, Syst. Ichth., 165, 1801 (*fasciatus*).

*Dactyleptus*, Rafinesque Anal. de la Nature 1815, 82; substitute for *Murænoïdes*.

*Centronotus*, CUVIER, Règne Animal, Ed. 2, II, 239, 1829 (*gunnellus*).

*Ophisomus*,\* SWAINSON, Nat. Hist. Class'n. Anim., II, 277, 1839 (*gunnellus*).

*Urocentrus*, KNER, Sitzber. K. Akad. Wiss. Wien, LVIII, 1868, 51 (*pictus*).

*Rhodymenichthys*, JORDAN & EVERMANN, Check-List Fishes, 474, 1896 (*ruberrimus* = *dolichogaster*).

Body long and low, considerably compressed, somewhat band-shaped, the tail slowly tapering; head small, compressed, naked;† mouth rather small, oblique; jaws with rather small teeth in narrow bands or single series; vomer and palatines usually toothless; gill membranes broadly united, free from the isthmus; scales very small, smooth; no lateral line. Dorsal fin long and low, beginning near the head, composed entirely of stiff, sharp, subequal spines; anal similar in form, of 2 spines and many

\* Substitute for *Gunnellus*, the latter being a barbarous word derived from "gunwale." "Nomina generica quæ ex Græca vel Latina lingua radicem non habent, rogienda sunt." This rule has never been generally adopted.

† In *Pholis nebulosus*, a Japanese species, the head is scaly. This species is the type of a distinct genus; which may be called

## ENEDRIAS, Jordan &amp; Gilbert, new genus.

*Enedrias*, JORDAN & GILBERT, new genus (*nebulosus*).

This genus differs from *Pholis* in the scaly head. (*ἐνέδρα*, lurking place.)

*Enedrias nebulosa* (SCHLEGEL).

Head  $7\frac{1}{2}$  to 8; depth  $8\frac{1}{2}$  to  $9\frac{1}{2}$ . D. LXXX; A. II, 39. Dorsal and anal somewhat connected to caudal; pectoral  $2\frac{1}{2}$  to  $2\frac{3}{4}$  in head. Head small. Body everywhere freckled with dark blotches; 12 dark triangular blotches along base of dorsal; a row of dusky blotches on middle of side posteriorly; 10 or 12 dark blotches on base of anal; caudal dusky, edged with pale, 2 pale cross streaks on top of head; pectoral pale. Northern Japan to Okhotsk Sea, Gulf of Strietok; our specimens from Hakodate. (*nebulosus*, clouded.)

*Gunnellus nebulosus*, SCHLEGEL, Fauna Japonica, Poiss., 138, 1850, Bay of Magi, Japan.

*Centronotus nebulosus*, STEINDACHNER, Ichth. Beitr., IX, 24, 1880.

*Enedrias nebulosus*, JORDAN & GILBERT, Rept. Fur Seal Invest., 1898, with plate.

soft rays; caudal fin short and small, more or less joined to dorsal and anal; pectorals short, rather shorter than head; ventrals very small, of 1 spine and a rudimentary ray; intestinal canal short, without caeca. Shore fishes of the Northern seas. (*φωλίς*, name of some fish said to shelter itself when lying in wait by producing a cloud of mucus; *φωλίς*, one who lies in wait.)

UROCENTRUS (*ούρά*, tail; *κέντρον*, spine):

- a. Pectoral fin small,  $3\frac{1}{2}$  to 4 times in length of head; dorsal spines about 93; anal rays 48; body with 2 rows of dark blotches; fins nearly plain. PICTUS, 2770.  
 aa. Pectoral fin moderate, 2 to  $2\frac{1}{2}$  times in length of head.

RHODYMENICHTHYS (*Rhodymenia*, a large red alga; *ρόδον*, rose; *ύμην*, membrane; *ιχθύς*, fish).

- b. Dorsal and anal joined to the caudal to the full height of the spines, without constriction at base of caudal; body greatly compressed, ribbon-like. Dorsal spines about 93; anal about 47; pectorals short,  $2\frac{1}{2}$  in head; no ocelli along base of dorsal. DOLICHOGASTER, 2771.

PHOLIS:

- bb. Dorsal and anal slightly connected with caudal, leaving a constriction of outline at base of caudal; body less compressed; dorsal fin with dark blotches or ocelli.

c. Pectoral fins well developed, about  $\frac{1}{2}$  length of head. Dorsal spines about 88; anal rays about 42; pectoral  $2\frac{1}{2}$  in head; dorsal fin with dark quadrate blotches rather than ocelli; sides scarlet in adult, bounded with black. FASCIATUS, 2772.

cc. Dorsal spines about 80 (76 to 85); anal rays about 40; pectoral 2 in head; dorsal fin with small rounded black blotches. GUNNELUS, 2773.

ccc. Dorsal spines about 77; anal rays about 35; pectoral 2 in head; dorsal fin with ocelli, or lunate, dark blotches. ORNATUS, 2774.

Subgenus UROCENTRUS, Kner.

2770. PHOLIS PICTUS (Kner)

Head  $9\frac{1}{2}$  to  $10\frac{1}{2}$ ; depth 8 to 10. D. XCIII or XCIV; A. II, 46 to 48 (misprinted 40 in Kner's account). Eye as long as snout; mouth oblique, the upper jaw the longer, reaching to front of eye; pectorals very short, scarcely longer than eye, 3 to 4 in head; anal said to have an isolated channeled spine hidden in the skin, but our specimens show no peculiar structure. Color yellowish, with 2 lengthwise series of large oblong blackish blotches, the one along base of dorsal, but not on the fin, of 21 or 22 blotches, the other on lower part of sides, of about 25; a series of fainter blotches along base of anal; in other specimens the lower row becomes obscure, the upper more distinct, and the series above anal disappears; a black bar downward from eye, a whitish band behind it; opercles dusky. West side of Bering Sea; our specimens from Shana Bay, Iturup Island, Kuril Group.

As already shown by Steindachner, this is a typical *Pholis*, Kner having been in error in ascribing to it an isolated and channeled first anal spine. The ventral spines are bound down by the integument more closely than usual, but they are in other respects not peculiar. Each is accompanied by 2 short spinous rays concealed in the membrane, and difficult to detect.

The latter are stiff and pungent, and seem to be not articulated. The ventrals of *P. ornatus* show the same structure. Kner gives the anal formula as II, 40. This must be a misprint for II, 49, as the artist figures 51 rays in the fin, not differentiating the 2 anterior ones. (*pictus*, painted.)

*Urocentrus pictus*, KNER, Sitzungsber. d. k. Akad. D. Wissensch., LVIII, 1868, 51, taf. 7, fig. 21, Singapore; an error.

*Centronotus pictus*, STEINDACHNER, Ichth. Beiträge, IX, 25, 1880.

*Pholis pictus*, JORDAN & GILBERT, Rept. Fur. Seal Invest., 1898.

Subgenus RHODYMENICHTHYS, Jordan & Evermann.

2771. PHOLIS DOLICHOGASTER\* (Pallas).

(BUTTER-FISH.)

Head  $9\frac{1}{2}$  in length; depth 8. D. XCII; A. II, 44; pectoral 14; eye 5 in head; maxillary  $2\frac{1}{2}$ ; pectoral  $2\frac{1}{2}$ ; caudal 2; ventral spines  $1\frac{1}{2}$  in eye. Body elongate, much compressed; head small, its upper profile convex; mouth moderate, very oblique, the maxillary reaching to below middle of eye; teeth rather large and blunt, arranged in a single row, the anterior one not enlarged; interorbital space narrow, without a sharp ridge, its width less than eye; snout equal in length to eye; distance from tip of snout to occiput  $1\frac{1}{2}$  in head. Head entirely naked; body covered with small, cycloid, inconspicuous scales. Origin of dorsal over upper end of gill slit, its distance from nape equal to distance from nape to front of eye, the spines toward the anterior end of the fin the highest; origin of anal a little nearer tip of caudal than snout; dorsal and anal confluent with caudal, the anal more broadly connected than dorsal; pectorals small, rounded behind; ventral spines inserted directly under base of pectorals, their length little greater than their distance apart; caudal short and broad, well rounded in outline. Bering Sea; recorded from the Kurils, and from Robben, Bering, and Medni islands, and from Kigiktoiwik Bay. The specimen above described was taken at Robben Island by Capt. J. G. Blair, then in command of the guardship *Leon*. It is 9 inches long and is uniform red in color, with a few pale dots. Another specimen, 18 cm. long, taken by Mr. Gerald E. H. Barrett-Hamilton at Bering Island, shows the following characters: The color is cherry red on the body and fins, lighter on belly, lower half of cheek and under side of head; lips blackish anteriorly, a narrow black streak running from them along snout to eye and from eye across cheek and opercles toward upper edge of pectoral base; this line separates the deep-red upper part of the head from the lighter area below;

\* The following species is allied to *Pholis dolichogaster*:

*Pholis taczanowskii* (Steindachner).

Head 9; depth 10; D. LXXXII; A. II, 45; teeth bluntly conical; dorsal very low, joined to the caudal without constriction. Snout scarcely longer than eye, which is  $5\frac{1}{2}$  in head. Pectoral 3 in head. Scales very small, the head naked. Clear, yellowish gray, finely dotted, fins grayish, the pectoral yellowish; a yellowish streak edged with darker from eye to axil. Gulf of Strietok. (Steindachner.) (A personal name.)

*Centronotus taczanowskii*, STEINDACHNER, Ichth. Beitr., IX, 24, pl. 3, fig. 1, 1880, Gulf of Strietok, Okhotsk Sea. (Coll. Prof. Dybowsky.)

sides  
midd  
gin of  
dorsa  
specie  
The d  
15 ray  
torals  
las, is  
cienn  
II, 50  
havin  
spotte  
pector  
Compu  
*P. rub*  
olive  
men,  
streak  
is evid

*Bienni*  
Mu  
Gunnell  
Island  
Gunnell  
Centron  
Muræne  
*Pholis d*  
*Rhodym*  
Am  
*Pholis r*

Head  
V. I, 1.  
level o  
to snout  
length  
slight  
a brill  
of da  
each c  
areas  
speckl

\* Brig  
project  
Muræne  
305, pl.

sides of body with a number of minute scattered black spots; along middle of sides is a distant series of light spots as large as pupil, the margin of each with 2 to 4 black specks like those scattered over sides. The dorsal and anal are more widely joined to the caudal than in other species, the fins being higher posteriorly and without perceptible notch. The dorsal contains 93 spines, the anal 2 spines and 47 rays, the pectorals 15 rays. Head  $9\frac{1}{2}$  in length; depth  $7\frac{2}{3}$ . Eye 5 in head; maxillary  $3\frac{1}{2}$ ; pectorals  $2\frac{1}{2}$ ; caudal  $2\frac{1}{2}$ ; ventral spine  $2\frac{1}{2}$  in eye. *Blennius dolichogaster*, Pallas, is undoubtedly identical with *Gunnellus ruberrimus*, Cuvier & Valenciennes. They agree in the very long dorsal and anal fins (D. XCIII, A. II, 50 in *dolichogaster*), and in the color. *P. dolichogaster* is described as having the color brownish olive, shaded with greenish and yellowish, spotted with green above the lateral line; belly yellow; anal, caudal, and pectorals yellowish; dorsal and anal dusky, with transverse pale bars. Compare with this, details of coloration recently published concerning *P. ruberrimus* by Bean & Bean (Proc. U. S. Nat. Mus. 1896, 248): "Color olive brown, with minute black spots; belly yellowish." In another specimen, "Across the spinous dorsal there are 20 narrow, nearly vertical pale streaks. Similar streaks to the number of 12 cross the anal." The species is evidently not always red in life. (*δολιχός*, long; *γαστήρ*, belly.)

*Blennius dolichogaster*, PALLAS, Zoogr. Rosso-Asiat., III, 175, 1811, Kamchatka. (Type in Mus. Berlin.)

*Gunnellus ruberrimus*, CUVIER & VALENCIENNES, Hist. Nat. Poiss., XIV, 440, 1839, Kuril Islands; after notes of PALLAS,\* Zoogr. Rosso-Asiat., III, 178, 1811.

*Gunnellus dolichogaster*, CUVIER & VALENCIENNES, Hist. Nat. Poiss., XI, 436, 1836.

*Centronotus dolichogaster*, GÜNTHER, Cat., 288, 1861.

*Muraenoides dolichogaster*, JORDAN & GILBERT, Synopsis, 768, 1883.

*Pholis dolichogaster*, JORDAN & GILBERT, Rept. Fur Seal Invest., 1893.

*Rhodymenichthys ruberrimus*, JORDAN & EVERMANN, Check-List Fishes North and Middle America, 474, 1896.

*Pholis ruberrimus*, BEAN & BEAN, Proc. U. S. Nat. Mus. 1896, 248.

#### Subgenus PHOLIS.

##### 2772. PHOLIS FASCIATUS (Bloch & Schneider).

Head 8 to  $9\frac{1}{2}$ ; depth 7 to 9. D. LXXXVI to LXXXIX; A. II, 42 to 44; V. I, 1. Head scaleless; mouth decidedly oblique, the tip of lower jaw on level of middle of the eye; width of mouth nearly  $\frac{1}{2}$  head. Eye equal to snout, a little more than interorbital width; ventral spine  $\frac{2}{3}$  eye,  $\frac{1}{2}$  length of mandible; caudal  $\frac{1}{2}$  head; pectoral  $2\frac{1}{2}$  in head; vertical fins slightly joined at base. Ground color yellowish gray in life, the sides of a brilliant scarlet; base of dorsal occupied by 10 or 11 oblong blotches of dark brown, which extend to the tips of the fins; these blotches each divided upon the fin by a median spot of the ground color; the areas of the ground color alternating with these blotches are finely speckled with brown, a large spot of brown usually occupying a median

\* Bright red. Form of *tenia*; scales inconspicuous; ventrals each a single scarcely projecting spine; caudal broad, rounded, distinct. D. CXV. Kuril Islands. (Pallas.) *Muraenoides ruberrimus*, BEAN, in Nelson, Rept. Nat. Hist. Collections made in Alaska, 305, pl. XIV, fig. 1, 1887.

position upon the fin; middle and lower part of sides occupied by vermiculating brown lines on the ground color, these vermiculations arranged in more or less distinct cross bars, about 20 in number, reaching to or nearly to the midventral line, the posterior ones often continued faintly onto the anal fin; pectoral and caudal fins yellow, unmarked; a brown blotch across snout and tip of mandible, followed by a narrow yellowish bar descending to front of eye; interorbital space crossed by a broad brown bar with blackish margins, which become much narrower below and traverse the eye and the cheek; behind this a broader yellow bar, margined behind with a narrow brown line. In life the coloration is extremely brilliant, the pale markings being bright orange or scarlet. Bering Sea and Arctic Ocean, from Greenland to the Kurils; locally abundant; numerous fine large specimens taken from the stomachs of cormorants on St. Paul Island, Pribilof Group; others dredged in shallow waters. Our specimens from St. Paul, Bristol Bay, and Upernavik, Greenland. Three large specimens from St. Paul Island, the type locality of *P. maxillaris*, have been compared with a number of individuals of *P. fasciatus* from Upernavik, Greenland. We can appreciate no differences between the two. The size of the mouth and the length of the head are the same in specimens of equal length, and no difference exists in the development of the ventrals. The agreement seems to be perfect in the fin rays, relative proportion and coloration. Pallas's short account of *Blennius tænia* contains nothing distinctive except the number of fin rays and the statement that the body is banded. As both of these items agree with the present species, we may safely follow Bean & Bean in making the identification. In a specimen from St. Paul, 29 cm. long, the length of the maxillary is contained  $2\frac{3}{4}$  times in distance from tip of snout to origin of dorsal; the mandible equals the length of the pectoral. In a younger example, 15 cm. long, from Bristol Bay, the maxillary is contained  $3\frac{1}{2}$  in predorsal length; the mandible approximately equals length of pectoral. (*fasciatus*, banded.)

*Centronotus fasciatus*, BLOCH & SCHNEIDER, Syst. Ichth., 165, pl. 37, fig. 1, 1801, Tranquebar; an error? GÜNTHER, Cat., III, 287, 1861.

*Gunnellus grœnlandicus*, CUVIER & VALENCIENNES, Hist. Nat. Poiss., XI, 442, 1836, Greenland, after BLOCH & SCHNEIDER; REINHARDT, Dansk. Vidensk. Selsk. Nat. og Mathem. Aft., VII, 122, 1838.

*Gunnellus murænoïdes*, VALENCIENNES, in CUVIER, Règne Animal, Poiss., pl. 78, fig. 2, 916; after BLOCH & SCHNEIDER.

*Blennius tænia*,\* PALLAS, Zoogr. Rosso-Asiat., III, 1811, 178, Kuril Islands.

*Murænoïdes maxillaris*, BEAN, Proc. U. S. Nat. Mus. 1881, 147, St. Paul Island, Alaska (Type, No. 23999. Coll. Henry W. Elliott); JORDAN & GILBERT, Synopsis, 768, 1883.

*Gunnellus fasciatus*, CUVIER & VALENCIENNES, Hist. Nat. Poiss., XI, 441, 1836.

*Murænoïdes fasciatus*, JORDAN & GILBERT, Synopsis, 767, 1883.

*Murænoïdes tænia*, JORDAN & GILBERT, Synopsis, 766, 1883.

*Pholis fasciatus*, GILBERT, Rept. U. S. Fish Comm. 1893, 449; JORDAN & GILBERT, Rept. Fur Seal Invest., 1898.

\**Pholis tænia* is thus described: Body banded; teeth obtuse, subdistinct; head subtriangular, compressed; body ensiform, covered with minute embedded scales; vent median. Dorsal fin extending from near the head to the tail, the spines subequal; caudal subdistinct; pectorals small; ventrals represented by 2 recurved spines. Body banded. D. LXXXVII; A. 47. Kuril Islands. (Pallas.)

H  
38 to  
nake  
blun  
row  
dist  
head  
with  
gene  
war  
This  
in c  
orna  
enci  
12 in  
way  
and  
(En.

Blenn  
Blenn  
p  
Ophid  
A  
Centr  
af  
Muræ  
Blenn  
Blenn  
G  
Muræ  
Ophid  
N  
Gunnel  
Muræ  
Gunnel  
(C  
Gunnel  
C  
26

He  
very  
front  
of pe  
yellow

\* A  
is pur  
of dor

2773. PHOLIS GUNNELLUS (Linnaeus).

(GUNNEL; BUTTER FISH.)

Head from 7 to 8 in body; depth 7 to 8; D. LXXVI to LXXXV; A. II, 38 to 44; V. I, 1; eye 5 in head; maxillary 3; P. 2; C. 1½. Head compressed, naked; mouth oblique, the maxillary reaching to front of pupil; teeth blunt, in a single row, somewhat enlarged anteriorly; interorbital a narrow ridge about ¼ eye. Distance from origin of dorsal to nape equal to distance from nape to middle of eye; pectoral rather large, about 2 in head, inserted directly under front of dorsal. Color olive brown, sides with numerous obscure darker bars; base of dorsal with blackish spots, generally bordered with a narrow yellow line, a dark bar running downward and backward from eye; anal with dusky bars across the rays. This species differs from *Pholis ornatus* in the more numerous fin rays and in coloration; the spots on dorsal are black, edged with yellow; in *P. ornatus* they are yellow with a black bar before and behind, each partly encircling it; no black bordered light streak from eye to occiput. Length 12 inches. North Atlantic, from Labrador south to Woods Hole and Norway to France; abundant on rocky shores among algae, both in America and Europe. Here described from specimens from Salem, Massachusetts. (Eu.) (*grunnellus*, English gunnel, said to be corrupted from *gunwale*.)

*Blennius pinna dorsalis ocellis X nigris*, LINNÆUS, Mus. Adolph-Fred., 1, 69.

*Blennius gunnellus*, LINNÆUS, Syst. Nat., Ed. x, 257, 1758, Atlantic Ocean; after *Blennius pinna dorsalis*, etc.

*Ophidium imberbe*, LINNÆUS, Syst. Nat., Ed. x, 259, 1758, Europe; after *Oph. cirris careus*, ARTEDI.

*Centronotus gunnellus*, BLOCH & SCHNEIDER, Syst. Ichth., 167, 1801; GÜNTHER, Cat., III, 285, after RÜPPELL'S type.

*Muraenoides gunnellus*, JORDAN & GILBERT, Synopsis, 767, 1883.

*Blennius europæus*, OLAFSEN, Røselv Island, 1, 81, 1772, Iceland.

*Blennius muraenoides*, SUJEF, Act. Petrop. II, 1779, 195, no locality, probably the Baltic; GMELIN, Syst. Nat., 1184, 1788.

*Muraenoides sujef*, LACÉPÈDE, Hist. Nat. Poiss., II, 324, 1800; after SUJEF.

*Ophidium muraenatum*, MITCHILL, Trans. Lit. & Phil. Soc. N. Y., II, 1815, 361, pl. 1, f. 1. New York; earliest American name.

*Gunellus vulgaris*, FLEMING, British Anim., 207, 1828, England.

*Muraenoides guttatus*, LACÉPÈDE, Hist. Nat. Poiss., II, 324, 1800; YARRELL, Brit. Fish., I, 269.

*Gunellus ingens*, H. R. STORER, Bost. Journ. Nat. Hist., VI, 1850, 261, pl. 8, f. 1, Labrador. (Coll. H. R. Storer.)

*Gunellus macrocephalus*, GIRARD, in H. R. STORER, Bost. Journ. Nat. Hist., VI, 1850, 263, Chelsea Beach, Massachusetts (Coll. Chas. Girard); D. H. STORER, Rept. Fish. Mass., 281, pl. 17, f. 3.

2774. PHOLIS ORNATUS (Girard).

Head 8; depth 8. D. LXXVII to LXXIX; A. II, 35 to 37. Head naked, very narrow above; nape nearly equidistant between origin of dorsal and front of orbit; origin of anal equidistant between base of caudal and base of pectoral; pectoral 2 in head. Coloration, usually olive green above, yellow or orange below,\* but varying with the surroundings to brown

\* A specimen from near Seattle varies much in color from all the others before us. It is purplish red, paler below; 2 conspicuous white spots bordered with white on front of dorsal; a pale streak bordered with black from eye to nape.



and cherry red; traces of about 20 darker bars along sides; a dark bar downward from eye; fins reddish; a V-shaped mark from eye to occiput, grayish, bordered by jet-black; the common form with about 14 red spots along base of dorsal, each with a curved black bar in front and behind, partly encircling it; others with about as many broad  $\wedge$ -shaped darker blotches, which extend on the fin, the first one or two blotches often shaped as in the former case; anal white, unmarked. Length 12 inches. San Francisco to Bering Sea; very common northward, its range extending to Kamchatka; \* very common at Unalaska; always in shallow water. (*ornatus*, ornamented.)

*Gunnellus ornatus*, GIRARD, Proc. Ac. Nat. Sci. Phila. 1854, 149; GIRARD, Pac. R. R. Surv., x, Fishes, 116; pl. 25b, figs. 6 and 7, 1858 (Type, No. 490, Presidio, California, Coll. Lieut. Trowbridge; No. 491, Shoalwater Bay, Washington, Coll. Dr. J. G. Cooper; No. 492, Fort Steilacoom, Washington, Coll. Dr. Geo. Suckley).

*Centronotus letus*, COPE, Proc. Amer. Phil. Soc. Phila. 1873, 27, Sitka or Unalaska (Coll. George Davidson); A. II, 33.

*Muraenoides ornatus*, JORDAN & GILBERT, Synopsis, 767, 1883.

*Pholis ornatus*, GILBERT, Rept. U. S. Fish Comm. 1893, 450.

#### 906. GUNNELLOPS, Bleeker.

*Gunnellops*, BLEEKER, Versl. Ak. Amst., 2, VIII, 1874, 368 (*roseus*).

This genus is apparently distinguished from *Pholis* by the tapering tail, around which the vertical fins are confluent; palatine teeth present. (*Gunnellus*, Gunnell, an old name of *Pholis gunnellus*;  $\acute{\omega}\psi$ , appearance.)

#### 2775. GUNNELLOPS ROSEUS (Pallas).

D. ca. C; A. ca. 90; P. 9; V. I. Head obtuse, the lower jaw projecting; eyes large; body very long, compressed, tapering into a slender tail; pectorals small, ovate, hyaline; 2 spines in place of ventrals; dorsal extending from the nape to the end of the tail; anal joined to caudal. Color intensely red. Kuril Islands. (Pallas.) Not seen by any recent collector. (*roseus*, rosy.)

*Blennius roseus*, PALLAS, Zoogr. Rosso-Asiat., III, 177, 1811, Kuril Islands.

*Centronotus roseus*, GÜNTHER, Cat., III, 290, 1861.

*Gunnellops roseus*, JORDAN & EVERMANN, Check-List Fishes N. and M. A., 474, 1896.

#### 907. ASTERNOPTERYX, Rüppell.

*Asternopteryx* (RÜPPELL MS.) GÜNTHER, Cat. Fishes Brit. Mus., III, 288, 1861, name only; JORDAN & GILBERT, Synopsis, 769, 1883 (*gunnelliformis*).

This genus is closely allied to *Pholis*, differing chiefly in the entire absence of ventral fins. From *Pholidapus* it is distinguished by the shorter pectorals and by the more broadly united gill membranes. Greenland. A single species known. ( $\acute{\alpha}$ -, without;  $\sigma\tau\acute{\epsilon}\rho\nu\nu$ , breast;  $\pi\acute{\epsilon}\rho\nu\acute{\xi}$ , fin.)

#### 2776. ASTERNOPTERYX GUNELLIFORMIS, Rüppell.

Head 9; depth  $8\frac{1}{2}$ . D. LXXXVII (LXXXI, according to Günther); A. II, 40. Head and body strongly compressed; head bluntish, snout short,

\* We have specimens collected at Tareinsky Bay by Mr. Barrott-Hamilton.

jaws equal; maxillary reaching pupil, 3 in head; eye  $5\frac{1}{2}$ ; gill membranes broadly united, their outline not notched; no trace of ventral fins; pectoral large, 2 in head (3 according to Günther). Dorsal and anal joined to the caudal, the anal with a slight notch behind the last ray; dorsal spines short and all pungent. Color dark brown, clouded with darker; about 11 quadrate pale areas along dorsal fin extending on the sides, these areas each with a black central spot at tip and faintly marked with dark blotches; dorsal with dark spots; a dark band from eye downward, a pale band behind it; lips dark; anal fin bright orange; pectorals and gill membranes pale orange; caudal orange. Greenland. Here described from a fine specimen,  $9\frac{1}{2}$  inches long, in the U. S. National Museum, from Omanak Fjord, Karsak, Noursoak Peninsula, taken in 1897 by Schuchert and White; only the original type in the Senckenburg Museum hitherto known. (*Gunellus*; *forma*, shape.)

*Asternopteryx gunelliformis*, RÜPPELL MS.: type (in Senckenburg Museum) from Greenland.

*Centronotus gunelliformis*, GÜNTHER, Cat., III, 288, 1861.

*Muraenoides gunelliformis*, JORDAN & GILBERT, Synopsis, 769, 1883.

#### 908. ANOPLARCHUS, Gill.

*Anoplarchus*, GILL, Proc. Ac. Nat. Sci. Phila. 1861, 261 (*atropurpureus*).

Body elongated, compressed, covered with very small, embedded scales which are obsolete or concealed anteriorly; lateral line obsolete. Head small, compressed; eyes small; mouth oblique; teeth in each jaw in a narrow band, the outer somewhat enlarged; narrow bands of teeth on vomer and palatines; gill membranes attached to the isthmus; sometimes with a free fold behind; branchiostegals 5. Dorsal fin not very low; no anal spines; ventrals wanting; caudal fin small, entire; pectoral fins moderate or small; pyloric caeca present, few. Pacific. (*ἀνοπλος*, unarmed; *ἄρχος*, anus; the anal fin being without spines.)

a. Gill membranes narrowly joined to the isthmus, with a free fold behind; dorsal with about 63 spines. ELECTROLOPHUS, 2777.

aa. Gill membranes broadly joined to the isthmus, without free fold behind; dorsal with 54 to 57 spines. ATROPURPUREUS, 2778.

#### 2777. ANOPLARCHUS ELECTROLOPHUS (Pallas).

Head  $6\frac{3}{8}$  in length; depth  $7\frac{3}{8}$ . D. LXII or LXIII; A. 43. Mouth oblique, maxillary reaching vertical behind pupil,  $2\frac{1}{4}$  in head. Teeth in narrow bands on the jaws, the outer series in upper jaw somewhat enlarged; vomer and palatines with narrow bands of teeth; dentition similar to that in *A. atropurpureus*, which has been erroneously described as having the teeth in the jaws in single series and the vomer and palate toothless; gill membranes rather narrowly joined to the isthmus and with a free posterior edge slightly wider than pupil. *A. atropurpureus* has the gill opening somewhat more restricted and the gill membranes without free fold. Large pores on head arranged similarly in the two species. Spinous dorsal beginning slightly in advance of base of pectoral, its distance

from snout less than length of head; distance from origin of anal to tip of snout  $2\frac{3}{4}$  in length to base of caudal; pectoral short and broad, rounded,  $2\frac{1}{2}$  in head. Scales small, embedded, those on the anterior part of the body concealed by the thickened integument, as in *A. atropurpureus*. Coloration in our specimens nearly uniform dark olive, with obscure dusky mottlings on the side. In 1 specimen there is a light bar extending obliquely downward and backward from eye, with a dark bar above and below it, the 3 separated by narrow light gray lines; caudal narrowly cross-banded with light and dark as in *A. atropurpureus*, and the anal obliquely barred with the same. In the smallest specimen is a series of roundish spots about as large as eye along back just below dorsal fin; each spot seems to have a narrow dark margin, a light ring, a dusky ring, and a light center; a series of similar but smaller spots along middle of sides posteriorly; the colors were probably brighter and more varied in life. Western part of Bering Sea and Sea of Okhotsk. Here described from 3 small specimens,  $3\frac{1}{2}$  to 9 inches long, taken at Tareinsky Bay, Kamchatka, by Mr. Barrett-Hamilton; 2 other fine specimens since taken by Arthur W. Greeley in Monterey Bay; the only ones recorded since Pallas. They differ from specimens of *A. atropurpureus* in the higher crest, the more numerous fin rays, and in having the gill membranes with a distinct free margin. (*ἀλέκτωρ*, cock; *λόφος*, crest.)

*Blennius alectrolophus*, PALLAS, Zoogr. Rosso-Asiat., III, 174, 1811, Island of Talek, Gulf of Penschin, Okhotsk Sea.

*Gunnellus alectrolophus*, CUVIER & VALENCIENNES, Hist. Nat. Poiss., XI, 447, 1836.

*Centronotus alectrolophus*, GÜNTHER, Cat., III, 289, 1861.

*Anoplarchus alectrolophus*, JORDAN & GILBERT, Rept. Fur Seal Invest., 1898.

#### 2778. ANOPLARCHUS ATROPURPUREUS (Kittlitz)

Head  $6\frac{1}{2}$  in body; depth 7. D. LV; A. 40; eye  $5\frac{1}{2}$  in head; maxillary  $2\frac{1}{2}$ ; pectoral  $2\frac{1}{2}$ ; caudal  $1\frac{3}{4}$ . Head with a fleshy crest, which rests on a ridge of bone, its height in older examples about equal to eye; mouth rather large, the maxillary reaching beyond the orbit. Dorsal and anal comparatively high, barely connected with the base of caudal; nape midway between origin of dorsal and pupil. Body naked anteriorly, scaled behind. Color grayish olive, varying to brown; everywhere above finely marked with blackish reticulations; along each side of back a series of small, irregular, sharply defined grayish spots; a series of small pale spots along lateral line; belly pale; crest and middle line of back rather pale; under parts of head yellowish; an oblique, wedge-shaped, pale streak extending downward and backward from the eye, bounded on each side by a sharp light-red line, and then by a dusky area; lower jaw mottled; dorsal olivaceous, speckled, a blackish spot on front; anal olive, tinged with red; pectorals dull orange, barred at base; caudal reddish, with narrow pale streaks, and a light bar at base; color sometimes nearly plain purplish, but more often grayish and mottled. Alaska to San Francisco; abundant northward; common in Bering Sea. Here described from specimens from Neah Bay, Straits of Fuca, Washington. We have also specimens from the Pribilof Islands. (*ater*, black; *purpureus*, purple.)

*Ophidium atropurpureum*, KITTITZ, Denkwürd einer Reise Russ.-Amer., 1, 225, 1858, Alaska.

*Centronotus cristagalli*, GÜNTHER, Cat., III, 280, 1861, Vancouver Island.

*Anoplarchus purpuraceus*, GILL, Proc. Ac. Nat. Sci. Phila. 1861, 261, Washington Territory. (Coll. Dr Kennerly.)

*Anoplarchus cristagalli*, GÜNTHER, Cat., III, 564, 1861.

*Anoplarchus atropurpureus*, JORDAN & GILBERT, Synopsis, 771, 1883; JORDAN & STARKS, Proc. Cal. Ac. Sci. 1895, 846.

909. XIPHISTES, Jordan & Starks.

*Xiphistes*, JORDAN & STARKS, Proc. Cal. Ac. Sci. 1895, 846 (*chirus*).

This genus is very close to *Xiphidion*, differing in the well-developed pectoral fins, which are longer than eye; lower lateral line not connected with abdominal line. ( $\xi\psi\iota\sigma\tau\acute{\eta}\varsigma$ , a sword belt.)

a. Anal spines 3, branches of upper lateral line extending on dorsal fin; color grass-green. ULYE, 2779.

aa. Anal spines 2; branches of upper lateral line shorter; color brownish, marbled, and with red blotches. CHIRUS, 2780.

2779. XIPHISTES ULYE, Jordan & Starks.

Head 8; depth 10. D. LXXIV; A. III, 48; eye 5 in head; maxillary  $2\frac{1}{2}$ ; pectoral  $3\frac{1}{2}$ . Body eel-shaped, as in the related species; head short; mouth small, oblique, maxillary extending to below posterior margin of eye; jaws subequal, with canine teeth; 4 enlarged canines in front of lower jaw; teeth in upper jaw gradually enlarged from behind forward; eye moderate, equal to length of snout; interorbital space prominent, sharply convex, narrower than width of eye; nape not constricted. Five mucous canals radiating downward and backward from eye, not reaching to edge of preopercle, the branches running upward from upper lateral line ending on the membrane of dorsal, the lower lateral line not connected with the abdominal line. Lateral line otherwise as in *Xiphistes chirus*. Origin of dorsal at a distance behind nape equal to distance from nape to middle of eye, the fin posteriorly barely connecting with caudal; anal with 3 spines, its origin about a head's length nearer snout than base of caudal, connected with caudal posteriorly; pectorals equal in length to snout and  $\frac{1}{2}$  eye, slightly shorter than caudal; caudal rounded, fan-shaped. Color olive-green above, very bright green below; middle and lateral line posteriorly, with conspicuous white spots,  $\frac{1}{2}$  as large as pupil, each with a black spot before and behind it; a black streak from tip of snout, through eye, to nape, a streak starting from eye behind quickly fading out; dorsal darker than body, unmarked; the anterior third of anal green, without markings, behind this, faint cross bars of brown appear, growing broader and darker posteriorly; caudal olive green, with a light bar across base; pectorals green, without markings. One specimen obtained at Waadda Island, Neah Bay. It was found high on the rocks, among algae, just below high water mark. Length 5 inches. This species is very closely related to *Xiphistes chirus*; it differs from it chiefly in having 3 anal

spines, in the branches of the upper lateral line running higher, and in coloration. (*Utra*, the green sea lettuce.)

*Xiphidion ulva*, JORDAN & STARKS, Fishes of Puget Sound, 847, 1895, Waadda Island, Neah Bay. (Type, No. 3132, L. S. Jr. Univ. Mus. Coll. E. C. Starks.)

2780. XIPHISTES CHIRUS (Jordan & Gilbert).

Head 7; depth 9. D. LXX; A. II, 50. Head short; nape not constricted; mouth small; maxillary extending to middle of pupil; teeth strong, the anterior canine-like, bluntish; about 4 canines in lower jaw, 5 or 6 in the upper, similar to the teeth behind them, but somewhat larger. Abdominal lines meeting on the breast, but not connected with the lower lateral line. Dorsal fin beginning close behind pectoral; nape midway between middle of eye and front of dorsal; anal beginning about a head's length nearer snout than base of caudal; pectoral fin comparatively large, longer than the eye, its length about equal to distance between middle and lower lateral lines. Color olive brown, yellowish below; sides with marblings of different shades of brown, sometimes with short blackish vertical bars; some round black spots along the back and sides; a black spot behind opercles; numerous black spots on sides of head, forming in older individuals light and dark streaks, which radiate from eye across cheek and opercles, the pale streaks forming reticulations; dorsal with black spots and a series of bright reddish brown cross blotches; pectorals and caudal plain. Monterey to Alaska; smaller than the other species, and living in deeper water; abundant about Cape Flattery. (Jordan & Gilbert.) (*χειρ*, hand.)

*Xiphister chirus*, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1880, 135, Point of Los Pinos, near Monterey, California (Coll. Jordan & Gilbert); JORDAN & GILBERT, Synopsis, 772, 1883; JORDAN & STARKS, Fishes of Puget Sound, 846, 1895.

910. XIPHIDION, Girard.

*Xiphidion*, GIRARD, Pac. R. R. Surv., x, Fishes, 119, 1858 (*mucosum*); not *Xiphidium*, Serv., a genus of Grasshoppers.

*Xiphister*, JORDAN, Proc. U. S. Nat. Mus. 1879, 241 (*mucosum*); substitute for *Xiphidion*, regarded as preoccupied by *Xiphidium*.

Body elongate, eel-shaped, covered with small scales; lateral lines several: 1 along the median line of the side, 1 above this, and 1 below it; 1 on each side of the abdomen, the 2 meeting in front, and 1 from the occiput toward the base of the dorsal fin. Each of these has on each side series of short branches, placed at right angles to the main line, those on opposite sides alternating. Each of these branches has about 2 open mucous pores. Lower lateral line connected with the abdominal line. Head short, bluntish, scaleless; mouth moderate, oblique; jaws with rather strong teeth, the anterior canine-like; no teeth on vomer or palatines. Branchiostegals 6; gill membranes separate, free from the isthmus. A single long, low, uniform dorsal fin, consisting of spines only; anal fin similar in form, with small spines, indistinct or obsolete; caudal short, joined to dorsal and anal; no ventral fins; pectoral fins very small, shorter

the  
py  
tid  
sw

a

ad

H

lary

seri

simi

line

sal

that

caud

side

tran

radi

below

thes

strea

ing

yellow

rock

from

Xiph

C

m

Xiph

D

Xiph

\* T

length

Teeth

the an

canals

bran

anal;

simila

and n

but no

each

rior la

Color

which

from

which

Girard

head,

than eye. Intestinal canal moderately elongate, with 4 to 6 well-developed pyloric caeca. Herbivorous, feeding on algae. Active fishes, inhabiting tide pools and crevices among rocks in the North Pacific. (*ξιφιδιον*, a small sword.)

- a. Distance from origin of dorsal to occiput less than that from occiput to tip of snout; streaks radiating from eye paler in the center, edged above and below with blackish. MUCOSUM, 2781.
- aa. Distance from origin of dorsal to occiput greater than that from occiput to snout; streaks radiating from eye black, abruptly margined with pale olive. RUPESTRE, 2782.

2781. XIPHIDION MUCOSUM, Girard.

Head 7 in body; depth  $8\frac{1}{4}$ . D. LXXIV; A. 46; eye  $7\frac{1}{2}$  in head; maxillary  $2\frac{3}{4}$ ; caudal  $2\frac{3}{4}$ ; pectoral a little longer than eye. Lower jaw with a series of short, stout conical teeth; upper jaw with a narrow band of similar teeth; 2 strong canines in upper jaw, 4 in the lower. Lower lateral line sending a branch to the abdominal line; nape not constricted. Dorsal beginning anteriorly, distance from its origin to occiput less than that from occiput to tip of snout; origin of anal nearer snout than tip of caudal by about  $\frac{1}{2}$  length of head. Blackish green, pale on belly and sides of head, marked posteriorly with olive green in various pattern; a transverse light-greenish bar at base of caudal; 3 olive-brown streaks radiating backward from eye, paler in the center and edged above and below with blackish, outside of which is sometimes a streak of pale olive; these streaks all merge backward into the color of the head; middle streak broadly wedge-shaped, the third streak terminating before reaching margin of preopercle; old individuals often coarsely blotched with yellow. Length 18 inches. Monterey to Alaska; very abundant among rocks and algae. Here described from specimens, 9 or 10 inches in length, from Neah Bay, Straits of Fuca, Washington. (*μυκοςυς*, slimy.)

- Xiphidion mucosum*, GIRARD, Pac. R. R. Surv., x, Fishes, 119, 1858, South Farallones, California (Coll. Lieut. Trowbridge. Type, No. 493, U. S. Nat. Mus.); GÜNTHER, Cat., III, 291, 1861; JORDAN & STARKS, Fishes Puget Sound, 848, 1895.
- Xiphidion cruoreum*,\* COPE, Proc. Amer. Phil. Soc. Phila. 1873, 27, Sitka (Coll. Prof. George Davidson); JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1880, 137.
- Xiphister mucosum*, JORDAN & GILBERT, Synopsis, 772, 1883.

\* The following is the original description of *Xiphidion cruoreum*: Head  $8\frac{1}{2}$  in total length; depth  $9\frac{1}{4}$ ; eye 7 in head, equal to length of pectoral fin. D. about 70; A. 48; Br. 5. Teeth, 2 canines above, 4 below, subequal. Dorsal spines not commencing near the head, the anterior buried in a soft fold of skin; caudal fin not distinct. Three lateral mucous canals extending entire length of caudal fin, which have numerous alternating transverse branches, those of the superior reaching base of dorsal, those of inferior reaching base of anal; each of the cross branches with several excretory pores, none on the main stem; a similar but short tube extending from near base of dorsal fin to supra-occipital region, and not branching anteriorly; the superior lateral canal descending to near the median, but not joining it, nor does the latter extend into the inferior; another tubular line on each side of abdomen, these uniting on jugular region by a continuation of the inferior lateral tube. Vent nearer end of muzzle than end of caudal fin, by length of head. Color maroon, more reddish below; a vertical, broad, reddish bar at base of tail, beyond which is a dark spot; 2 brown radii, black-edged, extending backward and downward from eye. Body covered with small scales, except on the jugular and abdominal regions, which are naked. Length 8 inches. This fish is not very different from *X. mucosum*, Girard. It differs in the smaller eye, the more remote origin of the dorsal fin from the head, the lack of anterior union of the mucous canals, and the coloration. (Cope.)

## 2782. XIPHIDION RUPESTRE (Jordan &amp; Gilbert).

Head  $7\frac{1}{2}$  in body; depth 9. D. LXVIII; A. 50; eye 6 in head; maxillary  $2\frac{1}{2}$ ; caudal  $2\frac{1}{2}$ . Teeth essentially as in *X. mucosum*. The lower lateral line sends a branch to the abdominal line; a constriction at the nape. Distance from origin of dorsal to the occiput greater than the distance from the occiput to the snout. Anal fin beginning much in advance of middle of body, the distance from the first ray to tip of caudal exceeding the distance to snout by nearly twice length of head; pectoral very short, its length less than diameter of eye. Reddish brown, uniform or slightly shaded with lighter; a light olivaceous bar at base of caudal, extending on dorsal and anal, behind this a blackish area; tip of caudal usually pale; 3 long, well-defined stripes radiating backward from eye, these stripes uniform black, abruptly margined with very light olive; the central stripe proceeding straight backward from the eye,  $\frac{1}{2}$  breadth of cheek, at which point it is broadest; it is then narrowed and bent abruptly downward; both the middle and lower stripes reach the margin of preopercle. Length 12 inches. Smaller than the preceding, and equally abundant; among rocks and algae, from Vancouver Island to Monterey. Here described from specimens, 6 or 7 inches in length, from Neah Bay, Straits of Fuca, Washington. (*rupestris*, living among rocks.)

*Xiphister rupestris*, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1880, 137, Monterey Bay, California (Coll. Jordan & Gilbert); JORDAN & GILBERT, Synopsis, 773, 1883.  
*Xiphidion rupestris*, JORDAN & STARKS, Fishes Puget Sound, 848, 1895.

## 9.1. CEBEDICHTHYS,\* Ayres.

*Cebedichthys*, AYRES, Proc. Cal. Ac. Nat. Sci., 1, 1855, 59 (*violaceus*).

Body comparatively short, compressed, covered with minute scales; lateral line distinct, running very high, with very short branches, each ending in a pore, as in *Xiphistes*, but the branches more oblique and less regular. Head short; crown with a conspicuous fleshy longitudinal crest in the adult; jaws subequal, with conical teeth; villiform teeth on vomer and palatines; gill membranes united, free from the isthmus. Dorsal fin continuous, long and low, the anterior part composed of sharp spines, which are rather lower than the soft rays; caudal fin rounded, connected with dorsal and anal; anal fin similar to soft dorsal, with 1 or 2 small spines; pectorals small; ventrals wanting. Intestinal canal elongate, with several pyloric caeca. Pacific Ocean. Herbivorous; similar in habits

\* The following remarkable genus may be allied to *Cebedichthys*:

## NEOZOARCES, Steindachner.

*Neozoarces*, STEINDACHNER, Ichth. Beitr., IX, 26, 1880 (*pulcher*).

## NEOZOARCES PULCHER, Steindachner.

Body elongate, tapering backward, the dorsal and anal united at the tail without distinct caudal. Scales small, embedded, no lateral line. Mouth very large, the maxillary extending far beyond eye; lower jaw slightly longer than upper; blunt, conical teeth in many rows on jaws, vomer, and palatines. A thick tentacle above nostril; gill membranes united, free from isthmus. Dorsal low, the anterior portion of short, stiffish spines; no anal spine; ventrals wanting; pectorals moderate; pseudobranchiae present. Head 6; depth 9. D. XLI, 50; A. I, 75. Color highly variegated. Gulf of Strietok, Okhotsk Sea. (*vés*, new; *Zoarces*; but it has little affinity with the latter genus.)

*Neozoarces pulcher*, STEINDACHNER, Ichth. Beitr., IX, 27, taf. 6, f. 2, 1880, Gulf of Strietok. (Coll. Professor Dybowski.)

to the species of *Niphidion*. (*κηβος*, the Sapajou, a kind of monkey; *ιχθύς*, fish; in allusion to the "peculiar monkey-like" physiognomy as seen from the front.)

2783. *CEBEDICHTHYS VIOLACEUS* (Ayres).

Head  $6\frac{1}{2}$ ; depth 6. D. XXIII, 41; A. I, 41. Maxillary extending to or beyond orbit. Dorsal scaly at base; vent nearer snout than base of caudal; pectoral  $\frac{2}{3}$  length of head; nape midway between dorsal and eye. Dull olive grayish, mottled with lighter, sometimes reddish tinged; vertical fins all edged with reddish; cheek with 3 darker stripes, edged with paler, 1 downward and backward from the eye, close behind angle of mouth; another above it to root of pectoral; another running upward and backward from the eye, and meeting its fellow over the crest. Length 30 inches. San Francisco to Point Concepcion; abundant; often brought into the markets. (*violaceus*, violet.)

*Apodichthys violaceus*, GIRARD, Proc. Ac. Nat. Sci. Phila. 1854, 150, San Luis Obispo, California. (Coll. Dr. Kennerly.)

*Cebedichthys cristagalli*, AYRES, Proc. Cal. Ac. Nat. Sci., 1, 1855, 58, San Francisco.

*Cebedichthys violaceus*, GIRARD, Pac. R. R. Surv., x, Fishes, 121, pl. 26, figs. 4 and 5, 1858; JORDAN & GILBERT, Synopsis, 774, 1883.

912. *PLAGIOGRAMMUS*, Bean.

*Plagiogrammus*, BEAN, Proc. U. S. Nat. Mus. 1893, 699 (*hopkinsi*).

Body moderately elongate, compressed, covered with very small scales; lateral lines 2, viz, 1 beginning above and slightly in advance of the upper angle of the gill opening and extending along the upper part of the body, but not reaching the tail, and 1 beginning in advance of the end of this and reaching to the caudal; numerous lateral ridges on the sides, similar to those on *Dictyosoma* of Temminck & Schlegel; a series of sub-pentagonal plate-like bodies along the abdominal edge on each side between the ventral and the anal. Head moderately long, naked, with pointed snout; mouth oblique and rather large; jaws subequal, or the lower slightly projecting; jaws with strong teeth in broad bands, the intermaxillaries with an outer series of enlarged canine-like teeth; teeth on vomer and palate; a pair of large canines near the symphysis in each jaw, the canines of the upper jaw fitting into an interspace behind the mandibular canines. A series of pores on the ramus of the mandibula continuing around the preopercular edge; a series of similar pores along the lower margin of the preorbital continued backward and upward toward the nape. Anterior nostril tubular; posterior without tube. Maxillary broadly expanded posteriorly; lips well developed. Branchiostegals 5; gill membranes partly united, but free from the isthmus behind. Gill rakers minute, tubercular, in moderate number. A single long dorsal fin consisting of spines only, the spines longest in the posterior portion; anal fin lower than the dorsal, but similar in shape. Pectoral large, entirely below median line. Ventrals well developed, in advance of pectorals; caudal rounded, distinct. Intestinal canal short, with 5 small pyloric caeca. (*πλάγιος*, oblique; *γραμμή*, line.)



2784. *PLAGIOGRAMMUS HOPKINSI*, Bean.

Head 4; depth  $5\frac{1}{2}$ ; eye 5 D. XI; A. II, 29; V. 1, 5; B. 5; scales about 95; ridges on side 32. Snout *acute*; anterior nostril tubular and nearer eye than tip of snout; posterior nostril close to upper anterior margin of eye; maxillary extending almost to vertical through hind margin of eye; intermaxillary long, slender, and reaching nearly as far back as maxillary; intermaxillary teeth in broad bands, with an outer series of 5 or 6 large canines, those near the symphysis largest; teeth in mandible in broad bands in front, followed by several enlarged canine-like teeth; a large canine on each side of symphysis, the interspace between the 2 mandibular canines receiving the canines of the intermaxillary when the jaws are closed. A row of 8 pores along ramus of mandible and edge of preopercle; another ser. around lower margin of preorbital bone as described for the genus; about 8 gill rakers on first arch below angle. Distance of dorsal origin from snout nearly equal to length of head; spines lowest in front, the longest spine  $\frac{2}{3}$  length of head; longest rays of anal near end of fin and scarcely exceed length of eye; length of pectoral equaling that of postorbital part of head; ventrals close together; inner rays longest,  $\frac{2}{3}$  as long as head; caudal rounded, its length nearly  $\frac{1}{2}$  that of head; vent under eleventh spine of dorsal. Upper lateral line beginning above and slightly in advance of upper angle of gill opening, curving very slightly over pectoral and extending to below twenty-fifth spine of dorsal, its distance from dorsal edge equal to diameter of eye and also equal to its distance from lower lateral line; lower lateral line beginning under sixteenth spine of dorsal and extending to caudal. On each side of the abdominal ridge, between the ventrals and the vent, are about 10 sub-pentagonal plate-like bodies, the largest about  $\frac{1}{2}$  as long as eye. Color dusky brown, the fins black. Monterey, California; a few specimens dredged among rocks. Little is known about the habits of the species, beyond the fact that in the aquarium it hides in rock crevices and seldom ventures from its hiding place. (I take pleasure in associating with this blenny the name of Mr. Timothy Hopkins, of Menlo Park, California, the founder of the Seaside Laboratory at Pacific Grove, Monterey Bay, in commemoration of his services in behalf of science. Bean.)

*Plagiogrammus hopkinsi*, BEAN, Proc. U. S. Nat. Mus. 1893, 699, Monterey Bay, California. (Type, No. 44721, U. S. Nat. Mus.)

913. *OPISTHOCENTRUS*, Kner.

*Opisthocentrus*, KNER, Sitzber. Akad. Wiss. Wien 1868, 49 (*quinquemaculatus*).  
*Blenniophidium*, BOULENGER, Proc. Zool. Soc. Lond. 1892, 583 (*netropauli*).

Body moderately elongate, compressed, covered with very small cycloid scales. Mouth small, horizontal, protractile, with fleshy lips; small conical teeth in jaws and on vomer and palatines. No cirri. Gill membranes broadly connected, but free from isthmus; branchiostegals 4. Dorsal fin very long, extending from the nape to the caudal, with which it is sub-continuous; a few of the posterior rays are stiff spines, the rest being

simple and not articulate, but flexible; anal fin extending from the anus, which is a little nearer the anterior than the posterior extremity, to the caudal, formed exclusively of soft rays; no ventrals. No lateral line. No prominent anal papillæ. Pyloric appendages present. A remarkable genus, allied to *Lumpenus*, or rather to *Plectobranchus*, distinguished by having only the posterior spines rigid. North Pacific. (*ὀπίθεε*, behind; *κέρτρον*, spine.)

## 2785. OPISTHOCENTRUS OCELLATUS (Tilesius).

Head  $6\frac{1}{2}$ ; depth  $6\frac{1}{2}$  (without caudal). D. LV to LXI, usually LIX; A. 36 to 39; 5 to 7 of the posterior dorsal spines rigid; Eye as long as snout, 1 in head, and a little more than interorbital width; maxillary extending to below anterior fourth of eye; some wide pores on the head; checks, opercles, and occiput closely scaled; strips of small scales on the branchiostegal membrane between the rays. Dorsal rays continuous and subequal in depth, the longest spine  $2\frac{1}{2}$  in head in females,  $1\frac{1}{2}$  in males; pectoral  $1\frac{1}{2}$  in head, about as long as caudal. Anus twice as far from caudal as from base of pectoral. Yellowish brown, with ill-defined darker marblings; a crescentic black line on the top of the head from eye to eye; a black line, obliquely directed forward, below the eye, and another, in opposite direction, from the eye to the opercle; 2 dark-brown streaks across the nape, the second crossing the origin of the dorsal fin and extending to the base of the pectoral; dorsal and caudal fins grayish olive, lighter at the base, the dorsal with 5 to 9 (usually 6) large black spots at regular intervals, these wanting in the males; pectorals and anal colorless. Numerous specimens are from Tarensky Bay, Kamchatka; Petropaulski Harbor, and Shana Bay, Iturup Island. The number of dorsal ocelli varies from 5 to 9 in our specimens, 6 being the prevailing number. Of 24 specimens whose fins we have enumerated, 4 have 58 dorsal spines, 10 have 59, 5 have 60, and 5 have 61. In addition, 1 specimen has but 55 spines. The latter is the only male in the collection and is conspicuous by the absence of distinct dorsal ocelli and the great height of the vertical fins, the longest dorsal spine exceeding the length of the pectoral and contained  $1\frac{1}{2}$  times in head. In females the longest spine is  $2\frac{1}{2}$  in head. The anal contains 36 to 39 rays in all our specimens. The dorsal fin is composed exclusively of spines, the anterior flexible ones passing into the strong pungent ones near the posterior end. The stronger spines vary from 7 to 12 in number in our specimens. Our material answers the description of the type of *O. quinque-maculatus* which had 57 dorsal spines and 36 anal rays. It also agrees with specimens from Petropaulski, reported on by Bean & Bean (Proc. U. S. Nat. Mus. 1896, 391), with dorsal spines 58 in number. *Bleculophidium petropauli*, Boulenger, has but 52 dorsal spines, but it is otherwise not to be distinguished from *O. ocellatus*. Still more aberrant are 2 specimens from Gulf of Strietok, northern Japan, mentioned by Steindachner (Ichth. Beiträge, IX, 25), with but 50 to 53 spines and 32 to 34 anal rays. These may represent a distinct species. *Ophidium ocellatum* of Tilesius must be this species, but the count of fin rays is incorrect and may be taken from

the rough figure. *Opisthocentrus tenuis* is probably also identical with *O. ocellatus*, though the writers did not think so until after examination of the present large material. Coast of Kamchatka, southwestward to Okhotsk Sea, generally common from Komandorski Islands to Yezo. (*ocellatus*, with eye-like spots.)

*Ophidium ocellatum*, THESLIUS, Mém. Ac. St. Petersb., II, 1811, 237, Kamchatka. D. 80; A. 50; evidently an error. The rude figure shows D. 73; A. 50, the spines low; the dorsal with 5 ocelli.

*Centronotus* (*Opisthocentrus*) *quinquemaculatus*, KNER, Sitzber. Akad. Wiss. Wien 1868, 48, taf. 7, f. 20, "Pinang." Described from a young specimen 2 inches long, No. 6353, Mus. Wien.

*Gunnellus apus*, CUVIER & VALENCIENNES, Hist. Nat. Poiss., XIV, 426, 1839; after THESLIUS. *Centronotus apus*, GÜNTHER, Cat., III, 288, 1861.

*Blenniophidium petropauli*, BOULENGER, Proc. Zool. Soc. London 1892, 584, with plate, Petropaulski (Coll. George Baden-Powell); D. 52; A. 37; 5 ocelli.

*Opisthocentrus tenuis*,\* BEAN & BEAN, Proc. U. S. Nat. Mus. 1897, Volcano Bay, Port Morusan, Japan. (Coll. Col. Nicolai A. Grelbittski. Type, No. 47565, U. S. Nat. Mus.)

*Opisthocentrus quinquemaculatus*, STEINDACHNER, Ichth. Beitr., IX, 25, 1880; BEAN & BEAN, Proc. U. S. Nat. Mus. 1896, 381, 392.

*Opisthocentrus ocellatus*, JORDAN & GILBERT, Rept. Fur Seal Invest., 1898.

#### 914. PHOLIDAPUS, Bean & Bean.

*Pholidapus*, BEAN & BEAN, Proc. U. S. Nat. Mus. 1896, 389 (*grebnitskii*).

Body moderately elongate, compressed, covered with very small, smooth scales. Mouth small, horizontal; bands of small teeth on jaws and vomer, none on palatines. Head naked; gill membranes broadly connected, free from the isthmus; dorsal very long, composed entirely of flexible spines; anal of soft rays; caudal short, rounded, separate; no ventral fins; no lateral line; pyloric caeca present. This genus is close to *Opisthocentrus*, but has no pungent spines, and the head is naked. Okhotsk Sea. (*φολίς*, *Pholis*; *ἄππος*, without feet, i. e., ventral fins.)

#### 2786. PHOLIDAPUS DYBOWSKII (Steindachner).

Head  $5\frac{1}{2}$  to  $6\frac{3}{4}$ ; depth 6 to  $6\frac{1}{2}$ . D. LXII or LXIII; A. II, 39. Eye  $3\frac{3}{4}$  to  $4\frac{1}{2}$  in head; snout a little longer than eye; lower jaw scarcely included; 1 or 2 strong conical teeth on each side behind the narrow premaxillary band of teeth; teeth on vomer, none on palatines; no cirri; large pores

\* *Opisthocentrus tenuis* is thus described:

D. 39, XV; A. 38. Length of fish to caudal base  $5\frac{1}{2}$  inches; length of head 1; depth of body  $\frac{1}{2}$ ; the greatest width of the body is contained  $2\frac{1}{2}$  times in the length of the head. The diameter of the eye is nearly equal to the length of the snout and is contained 4 times in the length of the head; the width of the interorbital space is almost equal to the long diameter of eye. The maxilla reaches to the vertical part front of eye. Teeth bluntly rounded, embedded in flesh; vomerine teeth present; palatines none. The origin of the dorsal fin is over the end of the gill cover, its first 39 rays are simple and flexible, the last 15 are strong spines and end slightly above the membrane in stiff points, the longest spine is almost  $\frac{1}{2}$  as long as the head. The anal originates under the twentieth ray of the dorsal; its rays are divided and articulated; the longest ray is  $\frac{1}{2}$  as long as the head. The general color is brown with cross retentions of black. Sides of head and body along base of anal, orange; anal, caudal, and pectorals light with dusky shadings; dorsal finely mottled with black and bearing 6 black spots on areas of white, the first of these spots being on the sixth ray and the last on the next to last spine; a black bar from front of eye downward, and another from posterior margin obliquely down and backward. This species differs from the typical form in its greater compression of the body and its increased number of dorsal spines. (Bean & Bean.)

about eye and on opercles; longest dorsal spines  $2\frac{1}{2}$  to 3 in depth of body, last spines shorter and stiffer than others; dorsal and anal slightly joined to caudal; pectoral as long as caudal, about  $1\frac{1}{2}$  in head. Head naked. Brown or grayish, with faint spots or marblings; 1 or 2, rarely 3, dark ocelli on the dorsal; 3 or 4 dark streaks radiating from eye, the uppermost joining its fellow. Length 10 to 15 inches. Coast of northern Japan and sea of Okhotsk, north to the Kuril Islands. Our specimens, 5 in number, the largest 25 cm. long, from Shana Bay, Iturup Island. Steindachner's excellent and detailed description leaves nothing to be desired, and corresponds perfectly with our material except in the character of the scales. A careful examination of these under high magnification fails to show that they are "am hinteren Rande mit kurzen Zähnchen bewaffnet." The posterior border is entire and the scales strongly marked with concentric striae. Dorsal spines number 62, 63, 63, 64, 64. Dorsal ocelli present in all our specimens, 2 of them being faintly visible even in the youngest, 55 mm. long. *Pholidapus grebnitskii* seems to differ only in the shorter dorsal fin (57 spines). (Named for Professor Dybowski, its first collector.)

*Centronotus dybowskii*, STEINDACHNER, Ichth. Beiträge, IX, 22, 1880, Gulf of Strietok, northern Japan (Coll. Prof. Dybowski); JORDAN & GILBERT, Rept. Fur Seal Invest., 1898.

*Pholidapus grebnitskii*,\* BEAN & BEAN, Proc. U. S. Nat. Mus. 1896, pl. 34, 390, Yezo, Japan. (Coll. Col. Nicolai A. Grebnitski.)

#### 915. PLECTOBRANCHUS, Gilbert.

*Plectobranchnus*, GILBERT, Proc. U. S. Nat. Mus. 1890, 102 (*evides*).

Teeth conic, on jaws, vomer, and palatines, some of them canine-like. Body scaly; lateral line obsolete, its course indicated by a lighter streak on middle of sides. Gill slits not continued far forward, the membranes

\**Pholidapus grebnitskii*, Bean & Bean, is thus described: The specimens are 141 mm. long, including caudal; 126 mm. to base of caudal. The head (22 mm.) is equal to the greatest depth of body. The eye is slightly longer than the snout and  $\frac{2}{3}$  as long as the head. The interorbital space is narrow,  $\frac{2}{3}$  of the length of the eye. The naked head resembles that of *Pholis*, its length is contained about  $5\frac{1}{2}$  times in total length without the caudal. The mouth is small and very oblique; the mandible is slightly included and has a well developed lip. The maxilla is partly concealed under the preorbital bone; it does not quite reach to below the anterior margin of the pupil. The anterior nostril is midway between the eye and the tip of the intermaxilla. Seven mucous pores around the orbit; 3 on the preorbital bone. The pore in the origin of the semicircular dark band around the nape is continued backward by a series of 6 similar ones ending near the upper angle of the gill opening. A series of 10 or 11 pores beginning near the front of the chin on each side, extending backward and curving upward to the upper anterior edge of the operculum. The gill membranes are broadly united, but they are not joined to the isthmus. The dorsal origin is over the end of the head; the fin is low, and consists of spines, the longest and strongest in the posterior third being slightly longer than the eye. The distance of the vent from the tip of the snout contains the head length  $2\frac{2}{3}$  times. The anal is slightly lower than the dorsal, the rays longest posteriorly. The caudal is rounded, and is barely separated from the dorsal and anal. The pectoral base is broad, and the fin is  $\frac{3}{4}$  as long as the head. The intestine is slender, and is more than twice as long as the head. Stomach short, pear-shaped, with 6 slender pyloric caeca of unequal length, the longest about twice as long as the eye. The body is completely scaled, the scales very small, cycloid, closely imbricated, with numerous concentric striae, and they extend halfway up the membrane connecting the dorsal spines. The general body color is brown, the sides sparsely and vaguely mottled. The pectorals are pale. A narrow, dark band extends from the middle of the eye downward and forward, a similar band running backward from the eye on the preopercle; an interrupted semicircular band from eye to eye across the nape. D. LVII; A. II, 39 or 40. (The species is named for Mr. N. Grebnitski, to whose industry and zeal the Museum is indebted for many valuable collections. Bean & Bean.)

broadly united, wholly free from isthmus. Dorsal of spines only. Anal with 2 spines. Ventral with 1 spine and 3 well-developed rays. Lower pectoral rays longest, as in *Leptoclinus*. North Pacific. (*πλεκτός*, enfolded; *βράγχος*, gill.)

2787. *PLECTOBANCHUS EVIDES*, Gilbert.

Head rather long,  $4\frac{1}{2}$  in length, extending well beyond origin of dorsal fin; depth about 11. D. LVI; A. II, 34. Body very slender, the depth nearly constant throughout. Caudal peduncle without free portion, its depth  $2\frac{1}{2}$  in that of body. Upper jaw with a broad inner band of minute teeth in front and on the sides, the outer series enlarged, 2 in the front of the jaw distinctly canine-like; teeth in the lower jaw similar to the outer series above, in a single series laterally, forming a patch in front of jaw, where 2 of them are much enlarged canines, the largest teeth in the jaws; vomer and palatines with bands of small but very evident teeth. Eyes large, close together, the interorbital space  $\frac{1}{2}$  pupil. Orbit  $3\frac{1}{2}$  in head, longer than snout. Posterior nostril with a short flap, the tube obsolete. Mouth large, somewhat oblique, maxillary reaching middle of orbit,  $2\frac{1}{2}$  in head. Top of head with very large pores, a series running backward from each eye, the two joined by a cross series on occiput. Body covered with very small cycloid scales, including belly, nape, breast, and cheeks, those on breast and cheeks not imbricated; lateral line without visible pores. Spinous dorsal beginning well forward, the distance from its origin to nape less than from latter to posterior margin of orbit. Anterior spines short, but fully united by membrane, the longest spine  $3\frac{1}{2}$  in head; membrane of last spine reaching base of upper caudal rays; origin of anal very slightly in advance of middle of body; anal with 2 short, sharp spines, the rays longer, their terminal  $\frac{1}{2}$  free from membrane; last anal ray connected with base of lower caudal ray; ventrals well developed, nearly  $\frac{1}{2}$  head; pectorals with lowermost rays abruptly lengthened,  $\frac{2}{3}$  head; caudal short, rounded, little more than  $\frac{1}{2}$  head. Color dusky olive above, lighter below; sides crossed by about 25 narrow white bars, narrower than interspaces; 3 equidistant dark blotches near back, each double, the two halves occupying contiguous interspaces between white bars; branchiostegal membrane black; head without markings; pectorals white at base, the distal half black, margined with white; ventrals white; dorsal with alternating oblique bars of white and blackish, 2 jet-black roundish spots on its posterior portion; caudal whitish at base, then dusky, margined with white, its upper ray jet-black; anal light at base, becoming black at edge of membrane, the free tips of rays white. Coast of Oregon. A single specimen, 4 inches long, from *Albatross Station 3064*, in 46 fathoms. (*εὐειδής*, comely.)

*Plectobanchus evides*, GILBERT, Proc. U. S. Nat. Mus. 1890, 102, coast of Oregon, at *Albatross Station 3064*. (Coll. *Albatross*.)

916. *LEPTOCLINUS*, Gill.

*Otenodon*, NILSSON, Skandinav. Faun., IV, 190, 1853 (*maculatus*) (name three times\* pre-occupied).

*Leptoclinus*, GILL, Proc. Ac. Nat. Sci. Phila. 1864, 209 (*aculeatus*).

\* *Otenodon*, Wagler, 1830, a lizard; Ehrenberg, 1838, a rotifer, and Swainson, 1839, a fish.

Body much elongated; lateral line obsolete; teeth on jaws, vomer, and palatines; pectoral fins with the upper rays shortened; caudal fin subtruncate. Arctic seas. This genus is close to *Lumpenus*, differing mainly in the form of the pectoral. (*λεπρός*, slender; *Clínus*.)

2788. LEPTOCLINUS MACULATUS (Fries).

(LANGBARN.)

Head 5; depth 8. D. LX (LVIII to LX); A. 36 (35 to 38). Eye large,  $3\frac{1}{2}$  in head; snout short and blunt,  $4\frac{1}{2}$  in head, maxillary reaching past middle of eye,  $2\frac{1}{2}$  in head. Teeth in jaws, vomer, and palatines; jaws each having 2 strong canines in front. Scales small, cycloid. First 3 or 4 dorsal spines short and free; longest dorsal spines as long as eye; caudal fin free from dorsal and anal; ventrals 3 in head; pectorals rather large,  $1\frac{1}{2}$  in head. Color yellowish, irregularly marked with dark spots; a series of about 6 of these spots extending along sides close to base of dorsal fin; a series of smaller spots extending along center of sides from upper base of pectoral to caudal; dorsal irregularly covered with dark spots; caudal with 4 dark cross bands; anal, ventral, and pectorals plain yellowish. Bering Sea to Spitzbergen, south to Aleutian Islands and the coasts of Sweden and Norway. This description is taken from a specimen,  $5\frac{1}{2}$  inches long, from Alaska, near Unimak Pass (*Albatross Station 3309*). A few young individuals of this species, hitherto known only from the North Atlantic, were taken in Unimak Pass and in Bristol Bay, in  $29\frac{1}{2}$  to 70 fathoms. Three small specimens were also taken off Robben Reef, near the Kanchatka coast, in 28 fathoms, and one off Karluk, Kadiak Island. Having no Atlantic specimens of this species, we are unable to satisfy ourselves of the identity of the two, but no difference is evident from descriptions. The lateral line is much more distinct than in our specimens of *Lumpenus medius*, in which it can be made out with difficulty on scattered scales along middle of sides. (Eu.) (*maculatus*, spotted.)

*Clínus maculatus*, FRIES, Kgl. Vet. Ak. Handl. 49, 1837, Bohuslän, Sweden.

*Lumpenus aculeatus*, REINHARDT, Kong. Dansk. Vid. Selsk., vi, 1837, 190, no description.

*Clínus aculeatus*, REINHARDT, Dansk. Vidensk. Selsk. Natur. Afh., vii, 1838, 114, 122, 194, Spitzbergen.

*Otenodon maculatus*, NILSSON, Skand. Fauna, iv, 190, 1853.

*Stichæus maculatus*, GÜNTHER, Cat., iii, 281, 1861.

*Lumpenus aculeatus*, KRÖYER, Naturhist. Tidsskr., i, 377, 1862.

*Stichæus aculeatus*, GÜNTHER, Cat., iii, 282, 1861; COLLETT, Norske Nord-Havs Exp., 67, 1880.

*Lumpenus maculatus*, JORDAN & GILBERT, Synopsis, 777, 1883; Lilljeborg, Sveriges Och Norges Fish., 500, 1891.

*Leptoclinus maculatus*, GILBERT, Rept. U. S. Fish Comm. 1893, 450.

917. POROCLINUS, Bean.

*Poroclinus*, BEAN, Proc. U. S. Nat. Mus. 1890, 40 (*rothrocki*).

Body elongate, moderately compressed, covered with small scales; lateral line obsolete. Head moderately long; snout short; eyes large; interorbital space narrow. Mouth small, lower jaw slightly included; teeth on vomer and palate; narrow bands of teeth in jaws, the outer series

enlarged. Gill openings slightly prolonged forward below, narrowly attached to the isthmus anteriorly. Dorsal composed of many sharp, flexible spines, diminished in length anteriorly. Caudal long, pointed; anal with 3 spines and many rays; pectorals large, the middle rays longest; ventrals jugular, with 1 spine and 3 rays. Intestine short; pyloric caeca 1 or 2; no air bladder. Northern Seas. (*πόρος*, pore; *Clinus*.)

2789. *POROCLINUS ROTHROCKI*, Bean.

Head  $6\frac{1}{2}$  in length; depth at nape 12. D. LVII to LX; A. III, 40 to 42. Body tapering uniformly backward. Vent placed anteriorly, its distance from snout  $1\frac{3}{8}$  to  $1\frac{1}{2}$  in its distance from base of caudal. Snout compressed, slightly projecting, the lower jaw included; maxillary reaching vertical from front of pupil,  $3\frac{1}{2}$  to  $3\frac{3}{4}$  in head. Teeth acute, in narrow bands in the jaws, a single well-marked series on vomer, and a patch on front of palatines, those on vomer and palatines fully as large as those on jaws, and equally developed in young and adults. Eyes large, close together, the interorbital space convex, its width about  $\frac{1}{2}$  pupil. Diameter of orbit equaling length of maxillary, about  $3\frac{1}{2}$  in head. Nostril tubes well developed,  $\frac{1}{2}$  diameter of pupil. Gill openings narrower than in other described members of this group, extending forward below the vertical from posterior part of cheek, where they are firmly joined to isthmus, across which they do not form a fold. Gill rakers obsolete. Dorsal beginning over end of opercular flap, its distance from nape equaling distance of latter from posterior margin of pupil; membrane of last spine slightly joined to base of caudal; anterior dorsal spines short, but well connected by membrane; anal with 3 distinct spines, shorter than the rays that follow, the second the longest, all as strong as dorsal spines, and fully connected by membrane, rays all branched at tip, membrane of last ray joined only slightly to base of caudal; caudal sharply pointed in all our specimens, the median rays longest, about as long as head; pectorals evenly rounded, the median rays longest, 14 or 15 in number, all branched; ventrals well developed, about  $\frac{2}{3}$  as long as head, consisting of 1 short, sharp spine and 3 rays, the spine not closely joined to rays. Lateral line indistinct, usually appearing obsolete, more evident toward head, consisting of a series of distant pores along median line; scales very small, cycloid, imbricated, covering body, inclosing abdomen, breast, and nape; cheeks scaled, the head otherwise naked, or sometimes with a small patch of scales on upper part of opercles. Color, sides with a series of 10 to 12 narrow white cross bars, the first in front of dorsal fin, the last under last dorsal spine, the bars about  $\frac{1}{2}$  interspaces; above lateral line scales conspicuously margined with darker, below lateral line they broaden out and become forked; upper caudal rays at base with an oval white ring inclosing a darker area, this mark more conspicuous in the young; belly and ventrals white, other fins dusky, but without definite markings. (Gilbert.) Bering Sea. Known from 2 specimens; the type, 7 inches long, was taken August 4, 1888, at Albatross Station 2852, north latitude  $55^{\circ} 15'$ , west longitude  $159^{\circ} 37'$ , at a depth of 58 fathoms, between Nagai and Big Kouinshi Islands. The spec-

imen here described from Unalaska. (Named for Dr. J. T. Rothrock, professor of botany, University of Pennsylvania.)

*Poroclinus rothrocki*, BEAN, Proc. U. S. Nat. Mus. 1890, 40, 55° 15' N., 159° 37' W., between Nagai Island and Koniushi Islands. (Coll. Albatross.)

918. LUMPENUS, Reinhardt.

(SNAKE BLENNIES.)

*Lumpenus*, REINHARDT, Dansk. Vidensk. Selsk. Natur., VI, 1837, 110 (*lumpenus* = *fabricii*).

*Leptogummellus*, AYRES, Proc. Cal. Ac. Nat. Sci., I, 1854, 26 (*gracilis*).

*Centroblennius*, GILL, Proc. Ac. Nat. Sci. Phila. 1864, 209 (*nubilus*).

*Leptoblennius*, GILL, Proc. Ac. Nat. Sci. Phila. 1864, 209 (*serpentinus*).

*Anisarchus*, GILL, Proc. Ac. Nat. Sci. Phila. 1864, 209 (*medius*).

Body greatly elongate, moderately compressed, covered with small scales; lateral line indistinct or obsolete. Head long; snout short; no cirri; eyes large, placed high; mouth moderate, with a single row of rather small conical teeth on each jaw; palatine teeth present or absent; gill openings prolonged forward below, very narrowly united anteriorly to the isthmus, not forming a free fold across it. Dorsal composed of numerous, sharp, flexible, rather high, spines; caudal fin long; anal many-rayed; pectorals large, more than  $\frac{1}{2}$  length of head, the middle rays longest; ventrals well developed, jugular, I, 3 or I, 4; intestinal canal long; pyloric caeca present; no air bladder. Chiefly herbivorous. Northern seas. (*Lumpen*, a Danish name of *Zoarces viviparus*, with which these fishes were at first confounded.)

ANISARCHUS (*ἀνίσκος*, unequal; *ἄρχος*, for anal):

a. Anal fin very low in front, the rays gradually lengthened; dorsal spines 61; anal rays 42. MEDIUS, 2790.

aa. Anal fin not much lower in front than behind.

LUMPENUS:

b. Teeth on palatines more or less developed, at least in the adult; anal rays 40 to 46; dorsal spines 63 to 71.

c. Dorsal spines 69 to 71.

d. Anal rays 46; dorsal separate from caudal. ANGULARIS, 2791.

dd. Anal rays 41; dorsal slightly joined to caudal. MACKAYI, 2792.

cc. Dorsal spines about 63; anal rays 43. FABRICII, 2793.

LEPTOBLENNIUS (*λεπτός*, slender; *Blennius*).

bb. Teeth on palatines wanting; dorsal spines 72 to 75; anal rays about 50.

LAMPETREFORMIS, 2794.

Subgenus ANISARCHUS, Gill.

2790. LUMPENUS MEDIUS (Reinhardt).

Head 6; depth 10. D. LXI; A. 42; V. I, 3. Lower jaw scarcely included, the maxillary reaching front of eye; teeth on palatines, none on vomer; ventrals slender,  $\frac{1}{2}$  length of head; lower rays of pectoral shorter than middle ones, the fin shorter than head. Dorsal and anal slightly joined to the truncate caudal; anterior half of anal with the rays shortened. (Collett.) Yellowish, nearly plain. Greenland to Norway and Spitzbergen and westward to Bering Sea and Kamchatka. Specimens from the



coast of Kamchatka are not evidently different from the current figures and descriptions of Atlantic specimens. (Eu.) (*medius*, middle.)

*Clonus medius*, REINHARDT, Dansk. Vidensk. Aftn., VII, 1838, 194, Greenland.

*Lumpenus medius*, KRÖYER, Naturh. Tidsskr., I, 377, 1837; JORDAN & GILBERT, Synopsis, 777, 1883.

*Stichæus medius*, GÜNTHER, Cat., III, 281, 1861.

*Anisarchus medius*, GILL, Proc. Ac. Nat. Sci. Phila. 1864, 210.

*Lumpenus medius*, COLLETT, Norskø Nord-Hava Exp., 62, 1880; JORDAN & GILBERT, Rept. Fur Seal Invest., 1898.

#### Subgenus LUMPENUS.

##### 2791. LUMPENUS ANGUILLARIS (Pallas).

Head 8; depth 14. D. LXXI; A. 46 (45 to 50); V. I, 4; B. 7. Cheeks scaly; mouth somewhat oblique, the lower jaw included; maxillary reaching front of pupil; teeth on palatines, none on the vomer; a single series of rather long, conical, and not very closely-set teeth in each jaw. Gill openings prolonged forward a distance greater than length of snout; pyloric caeca 4, unequal. Fins all comparatively high, pectorals  $\frac{2}{3}$  length of head, the middle rays longest; ventrals  $\frac{1}{2}$  length of head; dorsal and anal distinct from the pointed caudal, which is nearly as long as head. Olive green above, pale below; sides marked above with dark olive brown; a series of more or less distinct oblong blotches of olive brown along middle of sides; dorsal barred or spotted; anal pale; opercle with a dark blotch; head dusky above. Length 18 inches. San Francisco to Alaska; very abundant northward to Sitka and Unalaska; originally recorded from Kamchatka. (*anguillaris*, eel-like.)

*Bleinnius anguillaris*, PALLAS, Zoogr. Rosso-Asiat., II, 176, 1811, Kamchatha and Aleutian Islands. (Coll. Billings and Merk.)

*Septogunnellus gracilis*, AYRES, Proc. Cal. Ac. Nat. Sci., I, 1855, 26, San Francisco.

*Gunnellus anguillaris*, CUVIER & VALENCIENNES, Hist. Nat. Poiss., XI, 434, 1836.

*Lumpenus anguillaris*, GIRARD, Pac. R. R. Surv., X, Fishes, 123, pl. 25b, figs. 1 to 3, 1858; STORER, Synopsis, 121, 1846; JORDAN & GILBERT, Synopsis, 777, 1883; JORDAN & STARKS, Fishes Puget Sound, 848, 1895.

*Stichæus anguillaris*, GÜNTHER, Cat., III, 282, 1861.

##### 2792. LUMPENUS MACKAYI (Gilbert).

Head  $6\frac{2}{3}$ ; depth 13 or 14; eye 8 in head; snout 4. D. LXIX; A. II, 41. Very elongate. Head compressed and high, especially anteriorly, the upper profile of snout very convex, the upper jaw decidedly longer than the lower. Mouth nearly horizontal. Maxillary reaching vertical from front or middle of pupil, its length  $3\frac{1}{2}$  to  $3\frac{3}{4}$  in head. Teeth small, in a narrow band in jaws; a single series of weak teeth on palatines; vomer toothless. Gill openings continued forward to below middle of cheeks, the membranes then narrowly joined to isthmus; gill rakers short and weak, about 10 on horizontal limb of arch. Eye small, its horizontal diameter  $\frac{1}{2}$  longer than its vertical, slightly longer than interorbital width. Distance from snout to nape equaling length of postorbital part of head. Opercles large, continued to beyond base of pectorals. Dorsal

beginning immediately above upper end of gill slit, the spines short, strong, and pungent, not flexible; some of the anterior spines short, but not free, the fin increasing in height to opposite front of anal, the longest spine equaling length of snout, the membrane of last spine joining base of upper rays of caudal; anal with 2 strong spines similar to those of the dorsal fin, the second twice length of first and  $\frac{1}{4}$  that of highest dorsal spines; anal rays all forked, the posterior longest, equaling length of snout and eye, free from the caudal; caudal fin rounded in younger specimens, lanceolate in adults, becoming in the latter  $\frac{1}{2}$  as long as head; ventrals short, of 1 short spine and 3 simple rays, the fin  $\frac{1}{2}$  length of head; pectorals large, the middle rays longest,  $\frac{2}{3}$  length of head. Scales small, smooth, elongate, imperfectly imbricated, partially embedded or altogether wanting on anterior part of back; cheeks scaled, head otherwise naked; faint traces of a lateral line sometimes visible on middle of sides anteriorly. Color in spirits, light olivaceous (light yellowish in life); a continuous jet-black streak from occiput along each side of dorsal to base of caudal, with 2 interrupted black streaks below it, the lowermost running on middle of side; top and sides of head darker, variously marked with anastomosing black lines and spots; opercles blackish; dorsal and caudal fins dusky translucent, without distinctive markings; anal and ventrals white; pectorals white or dusky; roof of mouth black; peritoneum black dorsally, white ventrally. Bering Sea. Several specimens were seined near the mouth of the Nushagak River, Alaska. (Gilbert.) (Named for Charles Lesley McKay, of Appleton, Wisconsin, a very able young ichthyologist, who was drowned at Nushagak, in Bristol Bay, in 1883.)



*Lumpenus mackayi*, GILBERT, Rept. U. S. Fish Comm. 1893 (1896), 450, pl. 32, mouth of Nushagak River, Bristol Bay. (Coll. Gilbert.)

#### 2793. LUMPENUS FABRICII (Cuvier & Valenciennes).




Head 8 or 9; depth 11 to 15. D. LXIII to LXV; A. 41 to 43; V. I, 3; P. 15. Upper jaw scarcely longer than lower; teeth on palatines few and small, often really or apparently wanting, especially in the young; maxillary not reaching eye; vertical fins distinct; pectorals large, ovate. Color light brown, with large pale rounded blotches separated by brown shades; head yellowish; pectorals yellowish mottled, with a dusky spot at base. Arctic seas; recorded from Spitzbergen, Greenland, the Gulf of St. Lawrence, Wellington Sound, Bristol Bay, and other localities in Bering Sea (Petropaulski and Plover Bay, as *L. auquillaris*). We have specimens from Bristol Bay, Disco, Upernavik, and the Gulf of St. Lawrence. These are apparently identical, and they show clearly the identity of *L. uubilus* with *L. fabricii*.

The following notes are from specimens taken in Bristol Bay, in  $4\frac{1}{2}$  to 14 fathoms: These specimens seem to agree in structural details with specimens of *Lumpenus fabricii* from the North Atlantic. The Pacific specimens are lighter in color, with the dusky mottlings confined to the dorsal region, and with a very distinct series of oblong brown blotches along lateral line, alternating with a lower series of small, faint, round spots.





**IMAGE EVALUATION  
TEST TARGET (MT-3)**



**Photographic  
Sciences  
Corporation**

23 WEST MAIN STREET  
WEBSTER, N.Y. 14580  
(716) 872-4503

15 128  
16 32 125  
17 36 122  
18 20

19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100

Under parts immaculate; the mottlings along base of dorsal frequently uniting to form a series of oblong blotches alternating with those of lateral line; other specimens show no traces of dorsal blotches; dorsal fin translucent, faintly mottled with darker; caudal with brownish cross bars; pectoral with a round dusky shade at base; fins otherwise unmarked. Mandible with a single series of conical teeth, which widens at symphysis into an irregular double series or narrow patch; a similar series of conical teeth in premaxillaries, within which is a band of fine villiform teeth. A number of small specimens from Disco, Greenland, are entirely similar except for the darker coloration. This species is near *L. anguillaris*, but the latter has a larger mouth, larger teeth, and more numerous fin rays. (Named for Otho Fabricius, the first student of the fishes of Greenland.)

*Blennius lumpenus*, FABRICIUS, Fauna Grönlundica, 151, 1780, Greenland; not *Blennius lumpenus*, LINNÆUS, which is a species of *Gaidropsarus*, with 2 barbels at the chin. *Gunnellus fabricii*, CUVIER & VALENCIENNES, Hist. Nat. Poiss., XI, 431, 1836, Greenland, after FABRICIUS; KRÖYER, Naturhist. Tidsskr., 1, 377, 1837; GAIMARD, Voy. Scand., Zool., Poiss., pl. 14, fig. 1.

*Lumpenus nubilus*, RICHARDSON, Last Arctic Voyage, Fishes, 13, pl. 28, 1855, Wellington Sound. (Coll. Edward Belcher.)

*Blennius (Olinus) lumpenus*, RICHARDSON, Fauna Bor.-Amer., 90, 1836.

*Olinus lumpenus*, REINHARDT, Dans. Vidensk. Selsk. Nat. Afh., VII, 194, 1838.

*Stichæus lumpenus*, GÜNTHER, Cat., III, 280, 1861.

*Stichæus nubilus*, GÜNTHER, Cat., III, 564, 1861.

*Centroblennius nubilus*, GILL, Proc. Ac. Nat. Sci. Phila. 1864, 209.

*Lumpenus fabricii*, JORDAN & GILBERT, Synopsis, 778, 1883.

*Leptoblennius nubilus*, JORDAN & GILBERT, Synopsis, 778, 1883; GILBERT, Rept. U. S. Fish Comm. 1893 (1896), 451.

#### 2794. LUMPENUS LAMPETREFORMIS (Walbaum).

(SNAKE BLENNY; TANGHROSME.)

Head 9; depth about 15. D. LXXIII (LXVIII to LXXIV); A. 50 (49 to 52); V. I, 3. Body elongate, head slender; lower jaw little shorter than upper; maxillary reaching front of eye. Vent well forward, near end of first third of body; pectoral convex, somewhat shorter than head; first 3 or 4 rays of dorsal short, little connected; caudal acuminate, free from dorsal and anal. Yellowish or greenish, with numerous (about 20) faint brown blotches of different sizes, some of them confluent and extending obliquely upward on dorsal; caudal with transverse dark shades. (Collett.) North Atlantic and Arctic on both shores, south to Sweden and Norway, east to Spitzbergen; rare south to Cape Cod, if *L. serpentinus* is the same. We can find no difference on a comparison of our notes with published figures and descriptions, except that Storer describes *serpentinus* as having the caudal plain yellowish. (Eu.) (*Lampetra*, lamprey; *forma*, form.)

*Blennius capiti laevi*, etc., MOHR, Hist. Nat. Islandiæ, 85, taf. 4, 1786, Iceland; D. 72; A. 54.

*Blennius lampetraformis*, WALBAUM, Artedi Piscium, III, 184, 1792, Iceland; after MOHR.

*Centronotus islandicus*, BLOCH & SCHNEIDER, Syst. Ichth., 157, 1801, Iceland; after MOHR.

*Olinus nebulosus*, FRIES, Vet. Akad. Handl., 55, 1837, Bohuslän, Sweden.

*Olinus mohri*, KRÖYER, Naturh. Tidsskr., 1 R, 1837, 32, Iceland.

*Blennius gracilis*, STUVITZ, Nye Mag., Naturvid., 1, 406, 1838, west coast of Norway.

*Blennius serpentinus*, STORER, Proc. Bost. Soc. Nat. Hist., III, 1848, 30, Massachusetts Bay, from the stomach of a codfish (Ccil. Capt. Nathaniel E. Atwood); STORER, Hist. Fish. Mass. 169, pl. 18, f. 1, 1867.

*Gunnellus islandicus*, CUVIER & VALENCIENNES, Hist. Nat. Poiss., XI, 433, 1836.

*Stichæus islandicus*, GÜNTHER, Cat., III, 281, 1861.

*Lumpenus lampetraformis*, COLLETT, Norske Nord-Havs Exp., 71, 1880; JORDAN & GILBERT, Synopsis, 778, 1883.

*Leptoblennius serpentinus*, JORDAN & GILBERT, Synopsis, 778, 1883.

### 919. STICHÆUS, Reinhardt.

*Stichæus*, REINHARDT, Dansk. Vidensk. Natur. og Math. Afhandl. 1837, 109 (*punctatus*).

*Notogrammus*, BEAN, Proc. U. S. Nat. Mus., IV, 1881, 147 (*rothrocki*); young.

Body moderately elongate, covered with small scales; teeth on jaws, vomer, and palatines. Lateral line present, single, running along side of back; pectorals and ventrals well developed. Dorsal moderately high, of spines only; gill openings continued forward below, the membranes scarcely united to the isthmus; pyloric caeca present. Arctic seas. (*στῆχάω*, to set in rows.)

### 2795. STICHEUS PUNCTATUS (Fabricius).

Head  $4\frac{1}{2}$ ; depth about 7. D. XLVIII or XLVIX; A. 32 to 35; eye twice interorbital width,  $4\frac{1}{2}$  in head; snout subconical, 4 in head. Maxillary about equal to snout,  $3\frac{2}{3}$  in head, reaching slightly beyond front of eye. Narrow bands of teeth in the jaws and present on vomer and palatines, the outer series in the upper jaw and the inner series in lower jaw enlarged. Scales small, cycloid; head and cheeks scaleless; longest dorsal spines slightly longer than snout. The membrane from last dorsal spine joining extreme base of upper caudal ray; anal wholly distinct; pectorals rather long, reaching vent,  $1\frac{1}{2}$  in head; ventrals  $2\frac{1}{2}$  in head. Numerous large pores scattered over top and sides of head. Lateral line rather close to back, running along the upper fourth of height of body and ending abruptly at about  $\frac{2}{3}$  the length of body. Color bright scarlet, the head marked below with 5 or 6 brown reticulations; a brown streak from snout through eye; fins irregularly marked by dark bars or spots; a narrow row of 5 large round black spots, each with a white band near its posterior margin, occurring at regular instances along dorsal fin; a row of about 8 large dark spots on anal. Arctic seas, from Greenland to northern Siberia, south to Bristol Bay and Newfoundland. Our description (from Dr. Gilbert) taken from a specimen, about 5 inches in length, from Karta Bay, Alaska. It agrees very closely with the account by Ensign H. G. Dresel, of 2 examples from Godhavn, Disco Island, Greenland. The Alaska species must be the same as the other. Dresel finds the depth  $7\frac{1}{2}$  in length. Dr. Gilbert further observes: A single specimen, 86 mm. long, was dredged in Bristol Bay, Alaska, Station 3239, depth  $11\frac{1}{2}$  fathoms. Several larger individuals were seized in Karta Bay, Prince of Wales Island, Alaska, July 12, 1889. The position of the lateral line in this species is incorrectly given as "median" by Jordan & Gilbert in the Synopsis, pp. 755 and 775. Cuvier and Valenciennes, in their description, drawn from the writings of Fabricius, state

that the lateral line runs along the upper fifth of the height of the body and terminates at about the middle of the length. This correctly describes its position in all our specimens, where it originates immediately above the opercle, exhibiting at first rather a strong upward convex curve, then running nearly parallel to the back, separated from base of dorsal fin by  $\frac{1}{4}$  height of body. It is very distinct throughout its course, and terminates at about the middle of the length. The narrow brown streak bounding the lateral line above, in *Notogrammus rothrocki*, is conspicuous in our smallest specimen (86 mm.). Branchiostegal membranes very narrowly joined anteriorly, forming a narrow free fold across the isthmus, from which they are entirely distinct. Narrow bands of teeth in the jaws, and distinctly present on vomer and palatines; the outer series in upper jaw and the inner series in the lower jaw enlarged. D. XLVII or XLVIII; A. I, 32 to 35. The membranes from last dorsal spine join extreme base of upper caudal ray; anal wholly distinct. We have not the material for a comparison of Pacific with Atlantic representatives of this species, and the published descriptions of the latter lack detail. (*punctatus*, spotted.)

*Blennius punctatus*, FABRICIUS, Fauna Grönlandica, 153, 1780, Greenland; REINHARDT, Naturhist. Selsk. Skrift., II, pt. 2, pl. 10, fig. 3.

*Notogrammus rothrocki*, BEAN, Proc. U. S. Nat. Mus., IV, 1881, 146, Plover Bay and Cape Lisburne, Siberia; young. (Types, Nos. 27565, 27580, and 27573. Coll. Dr. Bean.)

*Clinus punctatus*, RICHARDSON, Fauna Bor.-Amer., III, 88, 1836.

*Gunnellus punctatus*, CUVIER & VALENCIENNES, Hist. Nat. Poiss., XI, 428, 1836.

*Stichæus punctatus*, KRÖYER, Naturhist. Tidsskr., I, 377, 1837; GAIMARD, Voy. en Scand. et Japon., Zool., Poiss., pl. 20, fig. 2; GÜNTHER, Cat., III, 283, 1861; JORDAN & GILBERT, Synopsis, 775, 1883; DRESEL, Proc. U. S. Nat. Mus. 1884, 249; GILBERT, Rept. U. S. Fish Comm. 1893 (1896), 450.

#### 920. ULVARIA, Jordan & Evermann.

*Ulvaria*, JORDAN & EVERMANN, Check-List Fishes N. and M. A., 475, 1896 (*subbifurcatus*).

This genus is very close to *Eumesogrammus*, from which it differs in the absence of the lowermost or third lateral line, the median line being bifurcate. (*Ulva*, the sea lettuce, in which many Blennioid fishes live.)

#### 2796. ULVARIA SUBBIFURCATA (Storer).

Head  $4\frac{1}{2}$ ; depth nearly 5. D. XLIV; A. 30. Mouth rather large; maxillary reaching to below orbit; back somewhat arched; ventral outline nearly straight; eyes large; lateral lines 2 (the lowermost lateral line wanting); median lateral line forked; upper branch of median lateral line about  $\frac{2}{3}$  length of the head. Brownish, with several round paler blotches above at the base of the dorsal fin; spaces between these blotches darker, appearing like bars; a broad black bar crossing the opercle obliquely from below the orbit, and 2 parallel dark bars running backwards from orbit; belly yellowish white; dorsal fin with numerous black dots. North Atlantic, south to Cape Cod; very rare. (*subbifurcatus*, somewhat forked.)

*Pholis subbifurcatus*, STORER, Rep. Fish Mass., 63, 1839, Nahant, Mass. (Coll. Dr. Thos. M. Brewer); DE KAY, N. Y. Fauna: Fishes, 150, 1842; STORER, Hist. Fish. Mass., 258, 1867.

*Eumesogrammus subbifurcatus*, JORDAN & GILBERT, Synopsis, 775, 1883.

Eu  
I  
lar  
Sc  
ven  
or  
for  
cre  
the

H  
of n  
line  
ven  
dors  
edge  
land  
Olin  
Olin  
Stich  
Eum

\* T  
gran

Hea  
Mout  
profil  
Head  
long,  
each  
dorsal  
ing at  
small  
dark  
bars  
Bay.  
distin  
Sticha  
B

Hea  
illary  
like te  
joined  
right  
along  
placed  
trials  
e  
script  
differs  
The n  
its lat  
Sticha  
Si  
K  
† Th  
1864, 2



## 921. EUMESOGRAMMUS,\* Gill.

*Eumesogrammus*, GILL, Proc. Ac. Nat. Sci. Phila. 1864, 210 (*precisus*).

Body comparatively short, the back somewhat arched; mouth rather large, the jaws with villiform teeth; teeth on vomer and palatines. Scales small; lateral lines 3, without accessory branches; pectorals and ventrals well developed. Dorsal moderately high, of spines only, free or slightly connected with the rounded caudal; gill openings continued forward below, the membranes narrowly joined to the isthmus; pyloric caeca present. Northern seas. ( $\epsilon\upsilon$ , well;  $\mu\acute{\epsilon}\sigma\omicron\varsigma$ , middle;  $\gamma\rho\alpha\mu\mu\acute{\eta}$ , line; the longest lateral line being the middle one.)

## 2797. EUMESOGRAMMUS PRECISUS (Kröyer).

Head 4; depth nearly 6. D. XLIX; A. 34; V. 3. Snout subconical; cleft of mouth slightly oblique; vomerine and palatine teeth present; 3 lateral lines on each side, the median one continued to the base of the caudal; ventral fin  $\frac{1}{2}$  as long as the pectoral, which is much shorter than head; dorsal fin terminating just at root of caudal. An ovate, black, white-edged spot between the sixth and tenth dorsal spines. Coasts of Greenland. (Günther.) (*precisus*, exact.)

*Olinus precisus*, KROEYER, Naturh. Tidskr., t. 25, August, 1836, † Greenland.*Olinus unimaculatus*, REINHARDT, Dansk. Vidensk. Selsk., vii, 114, Feb., 1837, Greenland.*Sticheus unimaculatus*, REINHARDT, Dansk. Vidensk., 109, 1837; GÜNTHER, Cat., 283, 1861.*Eumesogrammus precisus*, JORDAN & GILBERT, Synopsis, 774, 1883.

\* The 2 following species from the Okhotsk Sea seem to represent 2 new genera (*Erno-grammus* and *Ozorthes*) closely related to *Eumesogrammus*:

## ERNOGRAMMUS ENNEAGRAMMUS (Kner).

Head 3 $\frac{1}{2}$ ; depth 6 $\frac{1}{2}$ . D. XLI; A. 33 or 34; P. 14 or 15. Eye 4 in head, as long as snout. Mouth large, nearly horizontal, the maxillary reaching middle of eye; lower jaw projecting; profile of snout nearly horizontal; fine pointed teeth in bands on jaws and across the vomer. Head naked; dorsal of high, slender spines; caudal separate, rounded; anal high; pectoral long,  $1\frac{1}{2}$  in head; ventrals  $\frac{1}{2}$  as long as pectorals; scales very small, smooth; lateral lines each with short oblique branches, each ending in a wide pore; 1 lateral line along base of dorsal from head to caudal, 1 along middle of side, 1 along base of anal to caudal, this forking at the vent and sending 2 parallel branches forward to the breast; brownish, 2 rows of small dark spots along middle lateral line; dorsal and anal with dark spots and a broad dark margin; pectorals with 3 black cross bands; a dark bar at base of caudal; 3 black bars from eye. Okhotsk Sea. Known from a specimen,  $1\frac{1}{2}$  inches long, from Decastris Bay. (Kner.) (*ιννα*, nine;  $\gamma\rho\alpha\mu\mu\acute{\eta}$ , line.) *Ernoagrammus*, new genus (*εννος*, branch), is distinguished from *Eumesogrammus* by the branching lateral line.

*Sticheus enneagrammus*, KNER, Sitzber. Akad. Wiss. Wien 1868, 16, taf. vi, f. 19 Decastris Bay. (No. 1401c Mus. Wien.)

## OZORTHE HEXAORAMMA (Schlegel).

Head 5 $\frac{1}{2}$ ; depth 5 $\frac{1}{2}$ . D. XLIII; A. 24. Snout pointed; mouth little oblique; the maxillary reaching front of eye; bands of fine teeth on vomer and palatines; a few large canine-like teeth in front; eye 5 in head; dorsal spines all stiff, the middle ones longest; dorsal joined to caudal by membrane; lateral lines 3, the upper partly interrupted, sending at right angles upward and downward lines which join the middle line; third lateral line along base of anal only. Scales small. Dorsal with large dark brown spots obliquely placed; 3 brown stripes across cheek; anal colored like dorsal; caudal pectoral and ventrals each with 3 dark cross bands. Northern coast of Japan to Okhotsk Sea. This description (after Kner) from a specimen from Decastris Bay (No. 5575, Mus. Wien.). This differs somewhat from the type of the species and may be different. ( $\acute{\omicron}\zeta$ , six;  $\gamma\rho\alpha\mu\mu\acute{\eta}$ , line.) The new genus *Ozorthes* ( $\acute{\omicron}\zeta\omicron\varsigma$ , branch;  $\acute{\omicron}\rho\theta\eta$ , right angle) is distinguished by the form of its lateral lines as above described.

*Sticheus hexagrammus*, SCHLEGEL, Fauna Japonica, Pisces, 136, pl. 3, f. 1., 1850. Bay of Simabara, Japan. Head 4 $\frac{1}{2}$ ; depth 6 $\frac{1}{2}$ . D. XL; A. 29. GÜNTHER, Cat., iii, 284, 1861; KNER, Sitzber. Akad. Wiss. Wien 1868, 45.

† These dates are thus given by Kröyer, as quoted by Dr. Gill, Proc. Ac. Nat. Sci. Phila. 1864, 210. We have been unable to verify them.

## Family CCI. CRYPTACANTHODIDÆ.

## (THE WRY-MOUTHS.)

Body very long and slender, compressed, naked or covered with small, cycloid scales; lateral line obsolete or composed of open pores without tubes; head oblong, cuboid, with vertical cheeks; conspicuous muciferous channels in mandible and preopercle; head flattish above, with deep rounded pits between and behind eyes; mouth large, very oblique; lower jaw very heavy, its tip projecting; premaxillary not protractile; jaws with rather sharp, conical teeth; larger teeth on the vomer and sometimes on palatines. Gill membranes joined to the isthmus, the gill openings prolonged forward below. Pyloric cæca 5. Pseudobranchiæ small. Dorsal fin long, composed entirely of spines, which are rather strong, but enveloped in the skin; dorsal and anal joined to the caudal; no ventral fins; pectorals short. Blennies of large size, of the Northern shores of America. Three species known, forming 3 genera. (*Blenniida*, genus *Cryptacanthodes*, Günther, Cat., III, 291, 1861.)

a. Body scaly; lateral line present, composed of open pores; isthmus narrow; teeth on palatines. DELOLEPIS, 922.

aa. Body naked; lateral line obsolete.

c. Palatines with teeth; isthmus narrow.

CRYPTACANTHODES, 923.

cc. Palatines toothless; isthmus rather broad.

LYCONECTES, 924.

## 922. DELOLEPIS, Bean.

*Deiolepis*, BEAN, Proc. U. S. Nat. Mus. 1882, 465 (*virgatus*).

Body anguilliform, moderately compressed posteriorly, covered with small, imbricated, cycloid scales; vent nearly median; a small anal papilla; lateral line continuous, straight, nearly median, composed of open pores, without prominent tubes. Head oblong, subquadrangular, naked, the muciferous channels well developed, the vertex shallow concave; snout short, obtuse; nostril tubular, close behind premaxillary; eyes small, high, separated by an interspace of moderate width, surrounded by a series of shallow pits; mouth wide, oblique, terminal, the lower jaw projecting beyond the upper; lips fleshy; premaxillaries slightly protractile, with 2 rows of small conical teeth; a few larger teeth at the symphysis; vomer and palatines with a few rather large teeth; tongue smooth, adherent; mandible with a few shallow pits, the series continued on the posterior border of preopercle; opercles unarmed. Gill membranes attached to a narrow isthmus; gill rakers very short; pseudobranchiæ present. Branchiostegals 6. Pectorals short, placed low, their bases vertical; ventrals none; dorsal beginning above gill opening, composed entirely of spines; anal with 2 spines and many split rays; dorsal and anal continuous with the caudal, which is rather long and pointed. Intestine short, with a few pyloric cæca. (*δῆλος*, visible; *λεπίς*, scale.)

## 2798. DELOLEPIS VIRGATUS, Bean.

Head 6; depth 10. D. LXXVI; A. II, 46; P. 13; cæca 6. Width of head equal to greatest depth of body; interorbital area equal to snout, or  $\frac{1}{2}$  length of mandible; maxillary reaching a little behind eye, its length 3 in

distance from snout to front of dorsal; eye 2 in snout, 11 in head. Beginning at a short distance behind origin of dorsal, small, oblong, cycloid scales, closely imbricated, cover a strip of the body along the lateral line; the scaled area gradually widens backward until, behind the vent, only a very narrow strip along bases of dorsal and anal is naked. Dorsal beginning over upper angle of gill opening; first spine  $\frac{1}{2}$  as long as the seventy-first or longest; caudal 11 in length; pectoral 3 in head. Brownish yellow; a brown stripe along lateral line, another along back, a third along base of anal. Length 30 inches. Coast of southern Alaska to Puget Sound; not rare about Seattle. (*virgatus*, striped.)

*Delolepis virgatus*, BEAN, Proc. U. S. Nat. Mus. 1881, 466, Kingcombe Inlet, British Columbia; Port Wrangel, Alaska (Coll. Capt. H. E. Nichois. Types, Nos. 29149 and 29150, U. S. Nat. Mus.); JORDAN & STARKS, Fishes Puget Sound, in Proc. Cal. Ac. Sci. 1895, 848.

923. CRYPTACANTHODES, Storer.

*Cryptacanthodes*, STORER, Rept. Fish. Mass., 28, 1839 (*maculatus*).

Body long and slender, compressed, naked, without lateral line; head cuboid, with vertical cheeks and conspicuous muciferous cavities; eyes small, placed high; mouth large, very oblique, the very heavy lower jaw prominent in front; jaws, vomer, and palatines with stoutish conical teeth, in few series. Gill openings prolonged forward below, narrowly attached to the isthmus. Dorsal fin of stoutish spines, hidden in the skin; dorsal and anal joined to caudal; pectorals short; ventrals wanting. (*κρυπτός*, hidden; *ἀκαρῶδης*, spineless.)

2799. CRYPTACANTHODES MACULATUS, Storer.

(WRY-MOUTH; GHOST-FISH.)

Head  $6\frac{1}{2}$ ; depth 13. D. LXXIII; A. 50. Eyes small, placed high, not so wide as the interorbital space, which has 2 ridges and 3 pits; orbital rim raised; 2 deep pits behind each eye at the temples; a deeper pit on the top of head between them; a raised ridge continued backward on each side of head behind orbital rim; maxillary extending to beyond eye; pseudo-branchiæ small; pectorals short, 3 in head, their tips reaching beyond front of dorsal; vent a little in front of middle of body. Light brownish, with several series of smallish dark spots, arranged in more or less regular rows, from head to base of caudal; vertical fins closely spotted with darker; head above thickly speckled; body sometimes ("*inornatus*") entirely immaculate. Length 24 inches. Labrador to Long Island Sound; not very common; a few specimens have been taken at Woods Hole. The ghost-fish form (*inornatus*) occasionally seen, is doubtless an albino. (*maculatus*, spotted.)

*Cryptacanthodes maculatus*, STORER, Rept. Fish. Mass., 28, 1839, coast of Massachusetts; DE KAY, N. Y. Fauna: Fishes, 63, pl. 18, fig. 50, 1842; GÜNTHER, Cat., III, 291, 1861; JORDAN & GILBERT, Synopsis, 780, 1883.

*Ophidium imberbe*, PECK, Amer. Acad., 2d part, II, 1804, 46, pl. 4, New Hampshire; A. 49; P. 14; C. 22; not of LINNÆUS.

*Fierasfer borealis*, DEKAY, New York Fauna: Fishes, 316, 1842, New York; after PECK. *Cryptacanthodes inornatus*, GILL, Proc. Ac. Nat. Sci. Phila. 1863, 332, Coast of Massachusetts; albino form.

## 924. LYCONNECTES, Gilbert.

*Lyconectes*, GILBERT, Rept. U. S. Fish Comm. 1893 (1896), 452 (*aleutensis*).

Mouth subvertical; lower jaw projecting; premaxillary protractile. Teeth strong, conic, wide set, in more than 1 series. Mucous pits prominent on head. Gill opening narrow, ceasing opposite middle of base of pectorals, the membranes widely joined to isthmus. Dorsal and anal wholly joined to caudal, the latter extending well beyond them; dorsal fin composed of spines only; no ventral fins. Body naked; no lateral line. This genus differs from *Cryptacanthodes* principally in the absence of palatine teeth, agreeing with it in general appearance and in most details of structure. Alaska. (*λύκος*, wolf; *νηκτήρ*, swimmer.)

## 2900. LYCONNECTES ALEUTENSIS, Gilbert.

Head  $7\frac{1}{2}$ ; depth  $14\frac{1}{2}$ . D. LXIX; A. 49; P. 13; caudal 18. Head square in cross section, the upper and lower surfaces plain, the cheeks vertical, the depth and width equal. Mouth still more oblique than in *Cryptacanthodes maculatus*, with much heavier mandible and less expanded maxillary, the exposed portion of the latter lying vertically, and not extending beyond vertical from middle of eye. Teeth all similar, few in number, those in premaxillary arranged in 2 series, the inner of which are smaller than the outer, from which they are separated by a wide interspace; teeth in mandible in a single series laterally, becoming a sparsely filled patch toward symphysis; 4 or 5 similar conical teeth on head of vomer; palatines toothless. A long nostril tube overhangs the upper lip. Upper lip separated by a fold from forehead, the upper jaw protractile. Eye extremely small, sunken in the socket, which it does not nearly fill, its diameter slightly less than  $\frac{1}{2}$  interorbital width; supraorbital rim not elevated, and containing no conspicuous projections; suborbital rim swollen, with an enlarged mucous channel; a conspicuous series of mucous pits along each mandible and the margin of preopercle; 2 series on top of head diverging backward from above the eyes; otherwise no pits or projections on head; a shallow triangular depression on occiput. Gill slit much less oblique than margin of preopercle, its length  $1\frac{1}{2}$  times the distance between lower ends of gill slits, the latter reaching the vertical from middle of opercles. Dorsal fin of rather flexible spines, not concealed in heavy fin membranes; origin of dorsal immediately behind axil of pectorals. Hinder margin of occiput midway between front of dorsal and middle of eye. Origin of anal well in advance of middle of length, its distance from tip of snout contained  $1\frac{3}{4}$  times in its distance from base of caudal. Pectoral short, rounded, its base separated by a wide prepectoral area from gill slit, the width of area  $\frac{2}{3}$  length of fin, the latter equaling distance from tip of snout to middle of eye. No ventrals. Body covered with lax naked skin, which also covers but does not obscure rays of anal fin; no pores to lateral line. Color in life, reddish on head, body, and fins, due to the blood vessels in the skin. Aleutian Islands. A single specimen, 180 mm. long, known. (Gilbert.)

*Lyconectes aleutensis*, GILBERT, Rept. U. S. Fish Comm. 1893 (1896), 452, pl. 34, fig. 3, Albatross Station 3312, north of Unalaska Island, in 45 fathoms. (Coll. Albatross.)

Family CCII. ANARHICHADIDÆ.

(THE WOLF-FISHES.)

Body oblong or elongate, covered with rudimentary scales; no lateral line. Head scaleless, without cirri, its bones very thick and strong, the profile strongly decurved. Mouth very large, oblique, the jaws anteriorly with very strong conical canines; sides of lower jaw with very strong molar teeth, which shut against a series of very coarse molars on the palatines; vomer solid, armed with strong molar teeth, the dentition adapted for crushing sea-urchins and mollusks. Gill membranes broadly united to the isthmus; no pyloric cæca. Dorsal fin high, composed entirely of flexible spines; no ventral fins; pectoral fins broad, placed low. Large carnivorous fishes of the northern seas. Two genera and about 6 species known. (*Blenniide*, pt., Günther, Cat., III, 208-211, 1861.)

ANARHICHADINÆ:

a. Body moderately elongate, the tail not tapering to a point; dorsal and anal separate from the caudal. ANARHICHAS, 925.

ANARRHICHTHYINÆ:

aa. Body eel-shaped, excessively elongate; the dorsal and anal joined with the caudal at the end of the long and tapering tail. ANARRHICHTHYS, 926.

925. ANARHICHAS (Artedi) Linnæus.

(WOLF-FISHES.)

*Anarhichas* (ARTEDI) LINNÆUS, Syst. Nat., Ed. x, 247, 1758 (*lupus*).

Body moderately elongate, covered with rudimentary scales; head scaleless, without cirri, compressed, narrowed above, the profile strongly decurved; mouth wide, oblique; premaxillary not protractile; jaws with very strong conical canines anteriorly; lateral teeth of lower jaw either molar or with pointed tubercles; upper jaw without lateral teeth; vomer extremely thick and solid, with 2 series of coarse molar teeth; palatines with 1 or 2 similar series. Gill membranes broadly joined to the isthmus; no lateral line. Dorsal fin rather high, composed entirely of flexible spines, which are enveloped in the skin; anal fin lower; caudal fin developed, free from dorsal and anal; no ventral fins; pectoral fins broad, placed low; air-bladder present; no pyloric cæca. Northern seas. (*Anarhichas* (or *Scansor*), the climber; an ancient name of *Anarhichas lupus*; from ἀναρριχάομαι, to climb or scramble up; the allusion not evident, the word spelled with a single *r* by Artedi and Linnæus.)

a. Dorsal spines 60 to 70.

b. Vomerine teeth not extending farther backward than the palatine teeth.

c. Back and sides vaguely mottled, without spots or bands; vomerine teeth not extending nearly as far backward as palatine teeth. LATIFRONS, 2801.

cc. Back and sides profusely covered with roundish black spots; vomerine teeth extending nearly as far backward as palatine teeth. MINOR, 2802.

bb. Vomerine teeth extending much farther backward than the short band of palatine teeth; sides of body with 9 to 12 darker cross bars; nape and shoulder with dark spots. LUPUS, 2803.

aa. Dorsal spines 80 to 85; body without bands or spots; vomerine teeth extending farther backward than palatine band.

d. Head moderate,  $4\frac{1}{2}$  in length; caudal rays 20; upper canines 4.

LEPTURUS, 2804.

dd. Head very large; caudal rays 17; upper canines 6.

ORIENTALIS, 2805.

2801. ANARRHICHAS LATIFRONS, Steenstrup & Hallgrímsson.

Head 5; depth 4. D. LXVII; A. 45. Body more robust than *A. lupus*, the dorsal fin lower. Head broad, the profile not strongly decurved; teeth much smaller than in *A. lupus*; vomerine teeth not extending nearly as far back as the palatine series. Pectorals  $\frac{3}{4}$  length of head; dorsal fin not very high, beginning above the gill opening, the longest spine less than  $\frac{1}{2}$  head; caudal  $2\frac{1}{2}$  in head. Brown, obscurely spotted with darker; the sides without dark bars or black spots. (Collett.) North Atlantic on both coasts, chiefly north of the Arctic Circle, south to Banquereau on our coast. (Eu.) (*latus*, broad; *frons*, forehead.)

*Anarrhichas latifrons*, STEENSTRUP & HALLGRÍMSSON, Förh. Skand. Naturf. 3 die Mite 1842, 647; BEAN, Proc. U. S. Nat. Mus., 11, 1879, 218; COLLETT, Meddelsk. Norges Fiske 1879, 46; JORDAN & GILBERT, Synopsis, 782, 1883; GOODE & BEAN, Oceanic Ichthyology, 301, fig. 271, 1896.

*Anarrhichas denticulatus*, KRÜYER, Overs. Vidensk. Selsk. Kjöb. 1844, 140.

2802. ANARRHICHAS MINOR, Olafsen.

Head  $5\frac{1}{2}$ ; depth  $5\frac{1}{2}$ . D. LXXVIII; A. 46. Form of *Anarrhichas lupus* or a little more slender; fins similarly formed, the dorsal a little lower. Vomerine teeth extending nearly or quite as far back as the palatines. Body pale olivaceous or yellowish; sides without vertical bars; round, black spots covering dorsal and caudal fins as well as back and sides down to the level of the pectoral; head spotted; belly immaculate. North Atlantic, on both coasts, chiefly north of the Arctic Circle, south to Eastport, Maine; Gloucester; and Norway. (Eu.) (*minor*, smaller.)

*Anarrhichas minor*, OLAFSEN, Reise i Island, 592, 1772, Iceland.

*Anarrhichas pantherinus*, ZUIEW, Nov. Act. Petrop. 1781, 271; BEAN, Proc. U. S. Nat. Mus., 11, 1879, 217; JORDAN & GILBERT, Synopsis, 781, 1883; GOODE & BEAN, Oceanic Ichthyology, 301, fig. 270, 1896.

*Anarrhichas karrak*, BONNATERRE, Encycl. Ichth., 38, 1788, Iceland; after OLAFSEN.

*Anarrhichas maculatus*, BLOCH & SCHNEIDER, Syst. Ichth., 498, 1801, Iceland; after OLAFSEN.

*Anarrhichas leopardus*, AGASSIZ in SPIX, Pisc. Brasil., tab. 51, 1829, "Atlantic Ocean."

2803. ANARRHICHAS LUPUS, Linnæus.

(WOLF-FISH.)

Head 6; depth  $5\frac{1}{2}$ . D. LXII; A. 42. Maxillary reaching beyond orbit; band of vomerine teeth extending much farther back than the short palatine band. Pectorals large, rounded,  $\frac{3}{4}$  length of head. Dorsal high, beginning over the gill opening, its longest rays about  $\frac{1}{2}$  length of head. Brownish or bluish gray; sides with numerous (9 to 12) very dark transverse bars, which are continued on the dorsal fin; besides these

numerous dark spots and reticulations, the spots most distinct below front of dorsal; fins dark; caudal tipped with reddish. Length 3 to 4 feet. North Atlantic, south to Cape Cod and France; rather common both in America and Europe. A large voracious fish, not valued as food. The American form, *vomerinus*, seems to be fully identical with the European. (Eu.) (*Lupus*, a wolf.)

*Anarhichastopus*, LINNÆUS, Syst. Nat., Ed. x, 247, 1758, no definite locality; after ARTEDI: GÜNTHER, Cat., III, 208, 1861; JORDAN & GILBERT, Synopsis, 781, 1883; GOODE & BEAN, Oceanic Ichthyology, 299, 1896.

*Anarhichas strigosus*, GÜBELIN, Syst. Nat., I, 1144, 1788, British Sea.

*Anarrhichas vomerinus*, AGASSIZ in STOREY, Hist. Fish. Mass., 205, pl. 18, fig. 1, 1867, Cusk Rocks, between Boston and Cape Ann.

2804. ANARHICHAS LEPTURUS, Bean.

(ALASKA WOLF-FISH.)

Head  $4\frac{1}{2}$ ; depth 5. D. LXXXI; A. 52; C. 20 or 21. Head moderate; maxillary  $\frac{1}{2}$  as long as head; nostril nearer eye than mouth. Four large canines in the upper jaw and 5 in the lower, all of them strongly recurved; behind the canines in each jaw are a few sharp, conical teeth, also recurved; palatine teeth in 2 series, 4 in the outer and 5 in the inner series, those in the outer series the longer; vomerine teeth in 2 series, the vomerine patch beginning in advance of the palatine, and extending farther back than the latter; head and fins scaleless; median line of body and all of tail with small, widely separated scales. Dark brown, without bands or spots; belly pale, clouded with very dark brown. (Bean.) Coasts of Alaska, south to Vancouver Island; common about the Aleutian Islands, and perhaps identical with *Anarhichas orientalis*. (*λεπτός*, slender; *ὄψα*, tail.)

*Anarrhichas lepturus*, BEAN, Proc. U. S. Nat. Mus., II, 1879, 212, St. Michaels, Alaska; JORDAN & GILBERT, Synopsis, 782, 1883; GOODE & BEAN, Ocean. Ichth., 299, 1896.

? *Anarrhichas orientalis*, PALLAS, Zoogr. Rosso-Asiat., III, 77, 1811, Kamchatka.

2805. ANARHICHAS ORIENTALIS, Pallas.

This species, if correctly described, would differ from *Anarhichas lepturus* in the very large head,  $2\frac{1}{2}$  times in total length of body; in the absence of scales; in having the nostril midway between eye and mouth, and in having 6 canines in the upper jaw. Color plain brown. D. LXXXIV; C. 17. Coast of Kamchatka. (Pallas.) As the first of these characters is certainly erroneous, it is likely that the others are also, and that this species is not distinct from *Anarhichas lepturus*. (*orientalis*, eastern.)

*Anarrhichas orientalis*, PALLAS, Zoogr. Rosso-Asiat., III, 77, 1811, Kamchatka.

? *Anarrhichas lepturus*, BEAN, Proc. U. S. Nat. Mus., II, 1879, 212, St. Michaels.

926. ANARRHICHTHYS, Ayres.

*Anarrhichthys*, AYRES, Proc. Cal. Ac. Nat. Sci., I, 1855, 32 (*ocellatus*).

Body elongate, tapering backward into a very long and compressed tail, around which the dorsal and anal are confluent with the caudal. Scales rudimentary; no lateral line. Dorsal high, composed entirely of

flexible spines; pectoral fins broad, placed low; no ventral fins. Head very large, compressed, the snout rather short; mouth large; jaws with very strong, conical canines anteriorly; vomer and palatines each with about 2 rows of coarse molars, the palatine band shutting against similar teeth on the sides of the lower jaw. Gill membranes broadly united to the isthmus. No pyloric caeca. Large, eel-shaped fishes of the North Pacific, remarkable for the tremendous dentition, the head essentially as in *Anarrhichas*, the body strikingly different. (*Anarrhichas*; *Ιχθύς*, fish.)

## 2906. ANARRHICHTHYS OCELLATUS, Ayres.

(WOLF-EEL.)

Head 11; depth 15. D. CCL; A. 233; P. 19. Body elongate, formed as in an eel; the head and jaws very strong. Pectorals broad, more than  $\frac{1}{2}$  head; longest dorsal spine  $\frac{1}{2}$  head. Color dark greenish, the body and dorsal fin everywhere covered with round, ocellated black spots of various sizes, the light markings forming reticulations around the spots; head paler, with the reticulations in much finer pattern; anal pale-edged. Length 5 to 8 feet. Pacific coast, from Monterey north to Puget Sound; generally common. One of our most remarkable fishes; rarely used as food. It feeds chiefly on sea-urchins and sand dollars. (*ocellatus*, with eye-like spots.)

*Anarrhichthys ocellatus*, AYRES, Proc. Cal. Ac. Nat. Sci., I, 1855, 31, San Francisco; JORDAN & GILBERT, Synopsis, 782, 1893; JORDAN & STARKS, Fishes Puget Sound, 848, 1895.

*Anarrhichthys felis*, GIRARD, Proc. Ac. Nat. Sci. Phila. 1854, 150, San Francisco (Type, No. 511. Coll. W. O. Ayres), name only, no description; GIRARD, U. S. Pac. R. R. Surv., x, Fish., 125, pl. 25a, figs. 1 to 3, 1858; GÜNTHER, Cat., III, 211, 1861.

## Family CCIII. CERDALIDÆ.

Body elongate, compressed, covered with small scales; no lateral line; head small; gill openings reduced to small slit-like openings, more or less horizontal in position; dorsal fin very long and low, anteriorly of slender spines, which pass gradually into the soft rays; no free spines; no cirri; tail not isocercal; pseudobranchia well developed. Three species known, from the west coast of tropical America in rock pools near the shore. The presence of some spines in the dorsal separates them from the *Scytalinidæ*, while the small gill openings distinguish them from the *Blenniidæ*, to which they are more nearly allied.

a. Ventral fins each with 2 rays; dorsal rays 41; body moderately elongate; greatest depth  $10\frac{1}{2}$  in length; distance from insertion of dorsal to occiput equal to length of head. CERDALE, 927.

aa. Ventral fins each with 1 ray; dorsal rays 48 to 55; body very elongate, eel-like, its depth 15 to 18 in length. MICRODESMUS, 928.

## 927. CERDALE, Jordan &amp; Gilbert.

*Cerdale*, JORDAN & GILBERT, Bull. U. S. Fish Comm. 1881, 332 (*ionthas*).

This genus differs from *Microdesmus* in the presence of 2 rays in the ventral fin. Its body is much less elongate than in *Microdesmus*. The gill openings are reduced to small, nearly horizontal slits below and in front of the pectoral fins. (*κερδαλή*), the wary one, the fox-like.)



2807. CERDALE IONTHAS, Jordan & Gilbert.

Head  $7\frac{1}{2}$  in length; depth  $10\frac{1}{2}$ . D. 41; A. 36 to 38; C. 4-17-4; P. 12; V. 2. Body considerably elongate, compressed, of nearly equal depth throughout, the head tapering rapidly from occiput to snout; snout short, not obtuse, but the lower jaw heavy and blunt, much projecting beyond the premaxillaries; gape very short and oblique, the tip of the premaxillary not reaching ventral from orbit. Margin of upper jaw formed entirely by the premaxillaries, which are free laterally, but scarcely movable mesially. Maxillary not distinguishable, probably enveloped in the integument of the snout. Teeth rather strong, short, blunt, in a double series in each jaw, apparently wanting on the vomer and palatines. Lips developed laterally, where they form a fold around the angle of the mouth; lower lip adnate mesially, the upper reduced to an obsolete fold. Length of gape  $\frac{1}{2}$  length of head. Nostrils 2, distant, the anterior at the end of the snout, almost labial, the posterior above front of orbit, both circular. Eye very small, somewhat less than interorbital width or than length of snout. Distance from snout to post margin of orbit contained  $2\frac{1}{2}$  times in length of head. Pseudobranchiae well developed. Gill openings very narrow, reduced to a short, nearly horizontal slit, extending forward from a point just below the lower base of the pectoral fin. Branchiostegals evident, apparently 4 in number. Vertical fins well developed; dorsal and anal both long, the membrane of the last ray of each joining the base of the rudimentary rays of the caudal. Distance from occiput to the origin of dorsal fin equal to the length of the head; rays of dorsal fin very slender, distinct, the membrane thin and transparent, the rays all, or nearly all, articulate, the anterior simple, the posterior bifid at tip. Vent slightly in advance of middle of length of body, the anal fin beginning immediately behind it; anal rays bifid at tip, excepting the first 2, which appear simple; tail not isocercal, truncate at base of caudal, most of the rays of the caudal springing from the expanding last vertebra; caudal fin rounded,  $\frac{1}{2}$  length of head, its rays much branched, more closely set than the rays of the dorsal and anal; rudimentary rays very numerous; ventral fins small, close together, inserted slightly in advance of the lower end of the pectoral, each fin composed of 2 rays, the inner prolonged beyond the outer, and bifid at tip, about as long as pectoral fin and  $\frac{1}{2}$  length of head; pectorals well developed, broad, the rays branched at tip. Head and body entirely covered with small scales, which are close set but hardly imbricate, not arranged in series; mandible, snout, and gill membrane scaly; scales on belly and breast smaller than the others and more thickly set; base of caudal and pectoral fins scaled. Coloration in life, body translucent light olive, immaculate below; back and sides very finely marked with clusters of fine dots, the ground color appearing as reticulations between the clusters, which are of irregular size and form; on the sides of the head these dots form bars, which radiate from the eye to the snout and lower side of the head. This species is known from 3 specimens,  $2\frac{1}{2}$  to 3 inches in length, taken in a rock pool at Panama. (*lorbás*, freckled.)

*Cerdale ionthas*, JORDAN & GILBERT, Bull. U. S. Fish Comm. 1881, 332, Panama. (Coll. Chas. H. Gilbert.)

## 928. MICRODESMUS, Günther.

*Microdesmus*, GÜNTHER, Proc. Zool. Soc. London 1864, 26 (*dipus*).

Body anguilliform, covered with rudimentary scales; head small, with short, obtuse snout and small mouth; lower jaw projecting; teeth minute, in jaws only; eyes very small; gill opening reduced to a very narrow, somewhat oblique slit, in front of lower part of pectorals; vertical fins well developed, the dorsal and anal joined to the caudal by a thin membrane; rays of dorsal mostly articulate, all but a few of the last simple; ventral fins very small, reduced to a single ray; pectorals moderate; vent normal, in middle of body. Pacific coast of tropical America. (*μικρός*, small; *δέσμος*, a band.)

a. Dorsal rays 55, the fin beginning less than a head's length behind occiput.

DIPUS, 2808.

aa. Dorsal rays 48, the fin beginning more than a head's length behind the occiput.

RETROPINNIS, 2809.

## 2808. MICRODESMUS DIPUS, Günther.

Head about 11 in total length; depth about 18. D. 55; A. 34; C. 16; P. 12; V. 1. Head rather compressed, snout short, mouth very narrow, lower jaw very prominent. Eye minute, lateral, and in anterior third of head. Dorsal fin commencing at a distance from occiput which is somewhat less than length of head, nearly even, the rays very distinct, the interradial membrane being thin and transparent; anal fin commencing immediately behind vent. Caudal rays much more slender and more closely set than those of dorsal and anal; caudal fin rounded,  $\frac{3}{4}$  length of head; pectorals as long as ventrals, and  $\frac{1}{4}$  as long as head; ventrals close together, and inserted a little behind root of pectoral. Upper parts uniformly brownish olive. Panama. Known from a single specimen,  $4\frac{1}{2}$  inches long. (Günther.) (*δύς*, two; *πούς*, foot.)

*Microdesmus dipus*, GÜNTHER, Proc. Zool. Soc. London, January 26, 1864, 4, pl. 3, fig. 2, Central America (Coll. Capt. Dow); JORDAN, Cat., 126, 1885.

## 2809. MICRODESMUS RETROPINNIS, Jordan &amp; Gilbert.

Head  $14\frac{1}{2}$  in length; greatest depth  $15\frac{1}{2}$ . D. 48; A. 29; C. 3-17-3; P. 13; V. 1. Body very elongate, compressed, tapering somewhat from front of dorsal to caudal peduncle. Head very small, rapidly tapering forward from occiput; upper profile with a noticeable depression behind the orbits, the outline thence to snout strongly convex. Mouth very small, somewhat oblique, the fleshy tip at symphysis of lower jaw projecting much beyond the premaxillaries; gape scarcely reaching vertical from orbit. Teeth small, apparently in a single series in each jaw only. Nostrils double, distant, the anterior near the end of snout, the posterior above anterior margin of orbit. Gill openings a very narrow, somewhat oblique slit, from front of lower third of pectoral fin downward and forward. Branchiostegals evident, 4 or 5 in number. Eye very small, lateral, situated near the upper profile of the head, its diameter nearly  $\frac{1}{4}$  the length of the short snout. Vertical fins well developed; dorsal and anal

connected with the caudal by a very delicate membrane. Distance from origin of dorsal fin to occiput 3 times the length of the head, its rays distinct, connected by thin transparent membrane, as are the rays of the anal; most of the rays simple and undivided (but mostly articulate), a few of the posterior only forked at tip; origin of anal fin nearly equidistant between gill rakers and tip of caudal, its rays mostly forked at tip; caudal rays much divided and more closely set than those of dorsal and anal, the fin somewhat pointed in outline, as long as the head; tail not isocercal, truncate at base of caudal fin; ventral fins very small, close together, inserted slightly behind base of pectorals; each fin reduced to a single undivided filament; pectoral fin small, pointed, the middle rays longest, much shorter than the ventrals, and  $\frac{1}{2}$  the length of the head. Vent considerably behind middle of total length of the fish (with caudal). Head and body covered with scattered rudimentary scales. Color in life, translucent light olive, with a series of irregular quadrate dark blotches along the back and a series along each side, these blotches formed of clusters of dark points. One specimen, nearly 4 inches in length, was taken in a rock pool at Panama; others since taken by Dr. Gilbert. This species differs from the description of the previously known *Microdesmus dipus*, Günther, in the posterior insertion of the dorsal and the posterior position of the vent, the smaller number of fin rays, the shorter head, longer ventrals, and mottled coloration. (*retro*, backward; *pinna*, fin.)

*Microdesmus retropinnis*, JORDAN & GILBERT, Bull. U. S. Fish Comm. 1881, 331, Panama. (Col. C. H. Gilbert.)

Family CCIV. PTLICHTHYIDÆ.

(THE QUILL-FISHES.)

Body extremely elongate, serpentiform, little compressed, the tail tapering to a point. Skin with a few thin, loose, scattered scales; no lateral line. Head unarmed, rather small; upper jaw not protractile; snout short; mouth oblique; lower jaw projecting considerably beyond the upper, with a protruding fleshy appendage at tip. Maxillary reaching front of eye. Mandible little movable. Both jaws with fine, close-set, sharp teeth, in 1 row, the posterior teeth a little the largest; no evident teeth on vomer or palatines. Gill openings restricted to below the most convex part of the opercle, the membranes broadly united below, free from the isthmus. Gills 4, a slit behind the fourth. Pseudobranchiæ very small, almost obsolete. Gill rakers short and stout. Pectorals short; ventrals wanting; dorsal beginning close behind the nape, the anterior portion for about  $\frac{1}{2}$  the length of the body composed of very low, stiff, free spines, hooked backward, the posterior portion higher, of slender soft rays connected by thin membrane. No caudal fin, the tip of the tail free. Anal similar to the soft dorsal, without spines. Vent at considerable distance from the head. North Pacific. A single species known.

Concerning the relationships of this interesting group, Dr. Gilbert observes:

"The genus *Ptilichthys*, of which this species [*P. goodii*] is the sole representative, has been doubtfully referred by Dr. Bean to the *Mastacembelidæ*, a

family of fresh-water fishes inhabiting the East Indies, characterized by having the shoulder girdle posteriorly placed and not articulating with the cranium (Order *Opisthomi*, Gill). The necessity for preserving intact the unique type of the species prevented Dr. Bean from making any anatomical examination of *Ptilichthys*, and it was reserved for Dr. Theodore Gill, in the Standard Natural History, III, 259, 1885, to express his disbelief in the relationships which have been suggested, and to make the fish the type of a peculiar family, the *Ptilichthyidae*, to be placed provisionally among the Blennioid series. His adherence to this view is again expressed in his list of 'Families and Subfamilies of Fishes,' appearing as the Sixth Memoir of Volume VI, of the National Academy of Sciences. He has doubtless indicated the proper position of this peculiar fish as nearly as we are now able to determine it. An examination of its shoulder girdle shows it to be entirely normal. The post-temporal is not furcate, but is a very slender bony rod attaching to the epiotic region of the skull, and giving loose attachment posteriorly to the almost equally slender postero-temporal. The latter overlaps the upper end of the clavicle in the usual manner. A postclavicle was not detected. The coracoid portion consists of a roundish, oblong, perforated hypercoracoid meeting the hypocoracoid directly, without intervening cartilage. The curved line separating the two bones corresponds distally with the interspace between the first (upper) and second actinosts. The hypocoracoid is broad and short; its mesially directed (i. e., inferior) process joins at its tip the clavicle, but is elsewhere separated from the latter by the usual elongate membranaceous interspace. The actinosts are 4 in number, of large size, hourglass-shaped. The jaws are normal, the premaxillary alone occupying the front and sides of upper jaw and bearing the teeth, while the maxillary is a broad bone lying behind it, overlapped proximally by the maxillary process of the palatines. Both vomer and palatines seem to be toothless. The alimentary canal is almost perfectly straight, with the anterior portion entirely enveloped in the long, narrow liver. At the pylorus occurs a short and abrupt U-shaped flexure, scarcely noticeable on account of the closeness with which the sides are joined, and the fact that the width of the flexure is no greater than the cross diameter of the tube. Pyloric caeca are not evident. Air bladder is entirely wanting. The ovary is single, apparently without viaduct, and contains in our specimen eggs which are comparatively very large." (Gilbert.) (*Ptilichthyidae*, Gill, Standard Nat. Hist., III, 259, 1885.)

#### 929. PTILICHTHYS, Bean.

*Ptilichthys*, BEAN, Proc. U. S. Nat. Mus., IV, 1881, 157 (*goodiei*).

Characters of the genus included above. (*πριλον*, quill; *ιχθυσ*, fish.)

#### 9810. PTILICHTHYS GOODEI, Bean.

D. XC, 145; A. about 185; P. 12. Eye rather large, as long as snout, 5 in head; cheeks and opercles long; pectoral fin  $\frac{1}{2}$  as long as head; soft dorsal and anal deeper than body posteriorly, anal a little lower than dorsal. Vent near end of anterior third of body; distance from vent to

beginning of soft dorsal  $3\frac{1}{2}$  times length of head; length of head twice its greatest depth,  $5\frac{1}{2}$  in distance to vent; appendage of mandible  $\frac{1}{2}$  as long as eye; free tip of caudal  $\frac{1}{2}$  eye. Orange or yellowish, body with a blackish longitudinal stripe; anal darker in color than dorsal. Length about 12 inches. Aleutian Islands; rare; in water of moderate depth. Here described from the original type from Unalaska; 2 other specimens known, the one studied by Dr. Gilbert taken in the entrance to the harbor of Unalaska. (Named for Dr. George Brown Goode.)

*Ptilichthys goodii*, BEAN, Proc. U. S. Nat. Mus., IV, 1881, 157, Port Levachef, Unalaska (Col. Sylvanus Bailey. Type, No. 26619, U. S. Nat. Mus.); JORDAN & GILBERT, Synopsis, 369, 1883; GILBERT, Rept. U. S. Fish Comm. 1893 (1896), 453.

### Group OPHIDIOIDEA.

#### (THE EEL-POUTS.)

This group, as a whole, agrees with the *Blennioidea* in all respects, except that no spines are developed in any of the fins, save sometimes in the posterior part of the dorsal. From the *Anacanthini*, with which the *Ophidioidea* agree in the jugular ventrals and in the absence of spines, they are separated by the form of the hypercoracoid, which is perforate, as in ordinary fishes. The group is a very large and varied one, widely distributed in all seas.

- a. Pseudobranchiæ well developed, very rarely small or obsolete.
  - b. Ventral fins jugular, inserted much behind the eye, often wanting, never filamentous.
    - c. Gill membranes broadly united, free from the isthmus; ventrals wanting; no scales. SCYTALINIDÆ, CCV.
    - cc. Gill membranes united to the isthmus, the gill openings lateral. ZOARCIDÆ, CCVI.
  - bb. Ventral fins developed as slender filaments attached at the throat not far behind eye.
    - e. Gill membranes broadly attached to the isthmus; no scales. DEREPODICHTHYIDÆ, CCVII.
    - ee. Gill membranes nearly separate, free from the isthmus; body scaly. OPHIDIDÆ, CCVIII.
- aa. Pseudobranchiæ absent (or rudimentary in some *Brotulidæ*).
  - f. Ventral fins wanting; no scales.
    - g. Vent normal, well behind pectorals. LYCODAPODIDÆ, CCIX.
    - gg. Vent at the throat. FIERASFERIDÆ, CCX.
  - f. Ventral fins well developed; vent posterior, normal.
    - h. Dorsal fin single, low; ventral fins short. BROTULIDÆ, CCXI.
    - hh. Dorsal fins 2, the anterior, at the nape, of a single long ray; ventral fins elongate. BREGMACEROTIDÆ, CCXII.

#### Family CCV. SCYTALINIDÆ.

Body elongate, compressed, eel-shaped, naked. Head depressed, with tumid cheeks, like the head of a snake. Mouth moderate, horizontal, the lower jaw the longer; teeth in a single series in the jaws, vomer, and palatines; no barbels. Gills 4, a slit behind the fourth; pseudobranchiæ present. Gill membranes broadly connected, free from the isthmus. Dor-

sal fin long and low, beginning near middle of body, of slender rays embedded in the skin; anal similar to dorsal, both connected to the caudal fin; tail diphyccercal; pectoral fins small, ventral fins wanting. Vent remote from the head, without papilla. Air bladder none; caeca none. Vertebrae numerous, small. The skeleton does not differ essentially from that of *Lycodopsis paciflous*, with which it has been compared. The skull is not at all depressed, the wide depressed form of the head of the fish is due to the fleshy cheeks. The frontal takes up the greater part of the top of the skull, the parietals are separated by the supraoccipital, which extends forward to the frontals. Opercles all present. Lower jaw large and strong. Post-temporal scarcely so firmly attached as in *Lycodes*; the clavicle long and slender. As here understood, this family consists of a single species, a shore fish of the Northern Pacific, living in the gravel between tide marks, and diving with great activity into the wet gravel when disturbed. Its relations are apparently with the *Zoarcidae*. It is not certain that *Scytalina* has any special affinity with the *Congrogadidae*, in which group it was at first placed by Jordan & Gilbert.

#### 930. SCYTALINA, Jordan & Gilbert.

*Scytalina*, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1880, 266 (*cerdale*).

*Scytaliscus*, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1883, 111 (*cerdale*); substitute for *Scytalina* on account of the earlier *Scytalinus*, Erichson, a genus of Coleoptera.

Body very long and slender, covered with small scales. Head depressed, shaped like the head of a snake, with tumid cheeks and a distinct neck. Eyes small, superior. Mouth rather large, the lower jaw slightly projecting. Teeth conic, in single series on jaws, vomer, and palatines. Each jaw with 2 canines in front. No lateral line; pseudobranchiae small. Gill rakers almost obsolete. Dorsal fin very low, its first ray near the middle of the body. Anal fin similar to dorsal, nearly as long. Tail diphyccercal, the caudal well developed. (Diminutive of *Scytale*, from *σκυτάλη*, a viper.)

#### 2811. SCYTALINA CERDALE, Jordan & Gilbert.

Head 8; depth 14. D. 41; A. 36. Head broader than body; body much deeper behind vent than anteriorly; snout depressed, rounded at tip; cheeks very long; opercle short; interorbital space rather broad, concave posteriorly; eyes very small, anterior and superior, 10 to 12 in head, 2 in snout, 3 to 4 in interorbital width; upper lip separated by a crease from the skin of the forehead; lower jaw scarcely projecting; edge of lower lip with pores, and small dermal flaps and fringes; maxillary extending somewhat beyond eye; anterior nostrils with small flaps. Lower jaw with a series of close-set, oven, conical teeth, besides 2 divergent canines in front; upper jaw with similar teeth in several series in front, the canines smaller and closer together. Pectorals inserted high, little longer than eye; insertion of dorsal slightly in front of anal, a little in front of middle of body; rays of vertical fins low and weak, those of caudal most developed; dorsal and anal joined to caudal; vent close in front of anal,

which is similar to dorsal. Flesh colored, with much mottling of purplish in fine pattern; belly nearly plain; caudal reddish-edged. Length 6 inches. Straits of Juan de Fuca; burrowing among rocks near tide mark. The 2 original types came from the shore of Waadda Island, near Cape Flattery, where the species lives in wet shingle and shows extraordinary activity in hiding among rocks when disturbed. In the same locality 25 additional specimens have been dug out of the gravel by Mr. E. C. Starks in 1895. The species is still unknown from any other locality. (*περδαλή*, the wary one, the fox.)

*Scytalina cerdale*, JORDAN & GILBERT, Proc. U. S. Nat. Mus., III, 1880, 266, Waadda Island (Type No. 27400. Coll. Jordan & Gilbert); JORDAN & GILBERT, Synopsis, 701, 1883; JORDAN & STARKS, Fishes of Puget Sound, in Proc. Cal. Ac. Sci. 1895, 849, pl. 104.

Family CCVI. ZOARCIDÆ.

(THE EEL-POUTS.)

Body elongate, more or less eel-shaped, naked or covered with very small, embedded, cycloid scales; head large; mouth large, with conical teeth in jaws, and sometimes on vomer and palatines; bones of head unarmed. Gill membranes broadly united to the isthmus, the gill opening reduced to a vertical slit; pseudobranchiæ present; gills 4, a slit behind the fourth. Dorsal and anal fins very long, of soft rays only, or the dorsal with a few spines in its posterior portion; vertical fins sometimes confluent around the tail; pectorals small; ventrals jugular, very small or wanting, if present, inserted behind the eye. Lateral line obsolete or little developed, sometimes bent downward behind pectorals, sometimes sending a branch on median line backward. Gill rakers small; pyloric caeca rudimentary; vent not near head. Pseudobranchiæ present. Genera about 15; species 50. Bottom fishes, chiefly of the Arctic and Antarctic seas; some of them, at least, are viviparous, and some descend to considerable depths. Dr. Gill thus enumerates the skeletal characters of the *Zoarcidæ*:

Orbito-rostral portion of the cranium contracted and shorter than the posterior, the cranial cavity open in front, but bounded laterally by the expansion of the annectant parasphenoid and frontals, with the supraoccipital declivous and tectiform behind, the occipitals above inclined forward along the sides of the supraoccipital, and the exoccipital condyles distant, with the hypercoracoid foraminat about its center and the hypocoracoid with an inferior process convergent to the proscapula. These characters are formulated from the skeleton of *Zoarces anguillaris*. (Gill, Proc. Ac. Nat. Sci. Phila. 1884, 179.) *Zoarchidæ*,\* SWAINSON, Nat. Hist. Class. Fishes, II, 82, 184, and 283, 1839. *Lycoodidæ*, GÜNTHER, Cat., IV, 319-326, 1862; genus *Zoarces*, GÜNTHER, Cat., III, 295. 1861.

ZOARCINÆ:

- I. Dorsal fin low behind, some of its posterior rays short and spine-like; ventrals small; scales present; teeth strong, conic, in jaws only; lateral line present, along middle of side. ZOARCES, 931.
- II. Dorsal fin continuous, of soft rays only.

\* The name *Zoarchidæ* or *Zoarcidæ* is prior to that of *Lycoodidæ*.

## LYCODINÆ:

- a. Ventral fins present.
- b. Vomer without teeth; body scaly.
- c. Palatines without teeth.
- d. Body very slender, the depth 12 to 16 times in length; lateral line short and faint, ventral in position. EMBRYX, 932.
- dd. Body rather robust, the depth 8 to 9 in length; lateral line rather faint, lateral in position. LYCODOPSIS, 933.
- cc. Palatines with teeth; lateral line distinct, running along middle of side. APRODON, 934.
- bb. Vomer and palatines with teeth.
- e. Lower jaw without barbels.
- f. Dorsal fin without sculptured scutes at base.
- g. Body rather deep, the depth 6 to 8 times in the length.
- h. Body more or less scaly. LYCODES, 935.
- hh. Body entirely naked, or with a few scales on tail only; none on body or fins. LYCODALEPIS, 936.
- gg. Body more slender, the depth 12 to 20 in the length; lateral line lateral in position.
- i. Pectoral fin with rounded outlines, the lower rays not greatly produced. LYCENCHELYS, 937.
- ii. Pectoral fin deeply notched, the lower rays much produced; lateral line ventral in position. FURCIMANUS, 938.
- ff. Dorsal fin with the rays each provided with a sculptured scute or appendage at base; no lateral line; body elongate. LYCODONUS, 939.
- ee. Lower jaw with many barbels; body slender, scaly. LYCONEMA, 940.
- aa. Ventral fins entirely wanting.

## GYMNELINÆ:

- j. Teeth moderate, nearly uniform, on jaws, vomer, and palatines.
- k. Body scaly; vomer and palatines with teeth; body compressed, not very slender; skull cavernous. BOTHROCARA, 941.
- kk. Body scaleless.
- l. Lower jaw not very prominent; body very slender; gill openings very narrow. GYMNELIS, 942.
- ll. Lower jaw very prominent; body slender, tapering behind; scales undescribed. LYCOCARA, 943.

## MELANOSTIGMATINÆ:

- jj. Teeth long, unequal, on jaws, vomer, and palatines; skin lax; gill openings reduced to a small foramen; body very slender; scales obsolete. MELANOSTIGMA, 944.

## 931. ZOARCES, Gill.

## (EEL-POUTS.)

*Enchelyopus*, KLEIN, Ichthyologia Missus, IV, 52, 1747; not as restricted by BLOCH & SCHNEIDER.

*Zoarces*, CUVIER, Règne Animal, Ed. 2, II, 240, 1829 (*viviparus*).

*Zoarchus*, SWAINSON, Nat. Hist. Class'n Fishes, II, 283, 1839 (*viviparus*).

*Enchelyopus*, GILL, Proc. Ac. Nat. Sci. Phila. 1863, 258 (*viviparus*); not of BLOCH & SCHNEIDER.

*Macrozoarces*, GILL, Proc. Ac. Nat. Sci. Phila. 1863, 258 (*anguillaris*).

Body elongate, compressed, tapering posteriorly; head oblong, heavy, narrowed above, the profile decurved; mouth large; teeth strong, conic, bluntish, in 2 series in the front of each jaw and 1 series on the sides;



teeth in outer series larger; no teeth on vomer or palatines; dorsal fin very long, low, some of its posterior rays much lower than the others, developed as sharp spines; pectoral fins broad; ventrals jugular, of 3 or 4 soft rays; scales small, not imbricated, embedded in the skin; lateral line slender, lateral in position; size large; species viviparous. The American and Asiatic species (subgenus *Macrozoarces*) differ from the European type of *Zoarces*, Cuvier, in the increased number of fin rays and vertebrae. In *Zoarces viviparus* (Linnæus), the European eelpout, the dorsal rays are about 100, the anal about 85, and the number of vertebrae is proportionally diminished. (ζωαρκής, viviparous.)

Subgenus MACROZOARCES, Gill.

2812. ZOARCES ANGUILLARIS\* (Peck).

(EEL-POUT; MUTTON-FISH; MOTHER OF EELS.)

Head 6; depth 7. D. 95, XVIII, 17; A. 105. Mouth moderate, lower jaw included; maxillary reaching beyond orbit; pectoral long, about  $\frac{2}{3}$  length of head; ventrals  $\frac{1}{2}$  head; highest ray of dorsal about equal to snout, the posterior spines about  $\frac{1}{2}$  length of eye; first ray of dorsal above preopercle. Reddish brown, mottled with olive, the scales paler than the skin about them; dorsal fin marked with darker; a dark streak from eye across cheek and opercles. Length 20 inches. Delaware to Labrador; rather common north of Cape Cod. Two forms occur, distinguished by the size of the jaws. These have been regarded as distinct species, but the large-mouthed form (*ciliatus*; *labrosus*) is doubtless the male, as a similar variation occurs in *Lycodopsis pacificus*, and exists in some degree in species of *Lycodes*. (*anguillaris*, eel-like.)

? *Encheliopus*, GRONOW, Zoophyl., 77, No. 266, 1763, America (*unicolor*); dorsal and anal united with the caudal.

? *Blennius americanus*, BLOCH & SCHNEIDER, Syst. Ichth., 171, 1801, America; after Gronow.

*Blennius anguillaris*, PECK, Mem. Amer. Ac. Sci., II, 1804, 46, New Hampshire.

*Blennius fimbriatus*, MITCHILL, Trans. Lit. and Phil. Soc. N. Y. 1815, 374, pl. 1, fig. 6, New York.

*Blennius ciliatus*, MITCHILL, Trans. Lit. and Phil. Soc. N. Y. 1815, 374, pl. 1, fig. 7, New York.

*Zoarces labrosus*, CUVIER, Règne Anim., Ed. II, vol. 2, 240, 1829, America; CUVIER & VALENCIENNES, Hist. Nat. Poiss., XI, 466, 1836.

*Zoarces gronovii*, CUVIER & VALENCIENNES, Hist. Nat. Poiss., XI, 469, 1836; after Gronow.

? *Encheliopus americanus*, GRONOW, Cat. Fishes, Ed. Gray, 102, 1854, American Ocean.

*Zoarces fimbriatus*, CUVIER & VALENCIENNES, Hist. Nat. Poiss., XI, 468, 1836.

*Blennius labrosus*, MITCHILL, Trans. Lit. and Phil. Soc. N. Y. 1815, 375.

*Zoarces anguillaris*, STORER, Fishes Mass., 66, 1839; STORER, Synopsis Fishes N. A., 375, 1845; GÜNTHER, Cat., III, 296, 1861; JORDAN & GILBERT, Synopsis, 784, 1883.

\* Allied to *Zoarces anguillaris* is the following species from the Ochotsk Sea:

ZOARCES ELONGATUS, Kner.

Head 5 $\frac{1}{2}$ ; depth 11 $\frac{1}{2}$ . D. 80, XII, 22. A. 90 or more. Lateral line extending somewhat beyond pectorals. Color brownish, no brown streak behind eye; dorsal with 12 to 14 large dark spots which extend on the back as faint bands, between which are smaller ones. Known from 1 specimen, 10 $\frac{1}{2}$  inches long, from Decastria Bay, near the mouth of the Amur. (Kner). (*elongatus*, elongate.)

*Zoarces elongatus*, KNER, Sitzber. k. k. Akad. Wien 1868, 52, taf. 7, f. 2, Ochotsk Sea. (No. 1502, Wien Mus.)

## 932. EMBRYX, Jordan &amp; Evermann.

*Embryx*, JORDAN & EVERMANN, new genus (*crotalinus*).

This genus differs from *Lycodopsis* in the very slender body, the depth being 12 to 16 times in the length, and especially in the ventral position of the lateral line which is faint and incomplete, only the anterior descending portion developed. Deep seas. (*ἔν, in; βρῦξ, abyss.*)

a. Ventrals nearly as long as eye; head  $6\frac{1}{2}$  in length; no scales on head.

CRASSILABRIS, 2813.

aa. Ventrals shorter than pupil; head  $5\frac{1}{2}$  in length; head with some scales.

CROTALINUS, 2814.

## 2813. EMBRYX CRASSILABRIS (Gilbert).

Head  $6\frac{1}{2}$ ; depth 16; maxillary reaching vertical from front of pupil, 3 in head; exposed portion of eye 6; snout 4; width of snout 3. Body exceedingly slender. Occiput flat, forming a right angle with the descending cheeks, the snout short and wide, the upper lip conspicuously thickened and fleshy on the sides. Upper jaw with a single series of rather large, distant teeth; mandible with a broad patch of cardiform teeth anteriorly, which becomes abruptly constricted on middle of lateral portion of jaw, the inner series alone continued backward toward angle. Palate smooth. Head not conspicuously excavated with mucous canals; series of pores present on mandible and sides of head. Gill openings continued forward to below pectorals, and about to vertical from middle of opercle; the width of the isthmus  $\frac{1}{2}$  the length of slit. Opercular flap with a wide membranaceous border, produced backward and largely covering base of pectorals. Gill rakers very little developed, about 12 movable rudiments on horizontal limb of arch. Origin of dorsal in front of middle of pectorals, slightly farther from occiput than is the latter from front of eye; distance from origin of anal to tip of snout  $3\frac{1}{2}$  in total length; ventrals nearly as long as eye, inserted under middle of opercle; pectorals with 14 or 15 rays, the upper portion of fin longest, the lower rays rapidly shortened, the longest rays  $\frac{1}{2}$  as long as head. Scales small, circular, covering nape, breast, and under side of pectorals, but absent on head. Lateral line single, inconspicuous, running below middle of sides, ventral in position, the pores not developed on the scales. Color light brownish above, dark below; lower side of head, margins of snout, gill membranes, part of opercles, and margins of vertical fins jet black; ventrals and posterior face of pectorals black; anterior face of pectorals light glaucous blue, margined with black; lining of mouth and gill cavity and peritoneum black. Pacific coast of southern California. A single specimen, 12 inches long, from *Albatross* Station 2839. (Gilbert.) (*crassus*, thick; *labrum*, lip.)

*Lycodopsis crassilabris*, GILBERT, Proc. U. S. Nat. Mus. 1890, 106, off southern California. (Type, No. 44280. Coll. *Albatross*.)

## 2814. EMBRYX CROTALINUS (Gilbert).

Head  $5\frac{1}{2}$ ; depth 12; maxillary reaching to behind middle of pupil,  $2\frac{1}{2}$  in head; eye 6; interorbital width 14; snout 4. Body very slender, with

much the appearance of *Lycenchelys paillus*, the cheeks tumid, much projecting laterally, the greatest width of head more than  $\frac{1}{2}$  its length. Snout short and broad, much depressed, the head scarcely constricted opposite orbits. Eyes with little lateral range. In the single type specimen the upper jaw greatly overlaps the lower, the mandibular band of teeth shutting entirely within those on premaxillaries. Teeth in upper jaw in a single series, 2 or 3 small teeth sometimes present anteriorly, giving traces of an inner series. In lower jaw the teeth are sparsely set in a broad band anteriorly, becoming suddenly contracted to a single series on middle of sides. None of the anterior mandibular teeth enlarged, 2 or 3 of posterior teeth on sides larger and hooked backward. No teeth on vomer or palatines. Nostril in a short tube. Gill slits wide, reaching to below pectorals, but not extending farther forward below than above. Width of isthmus  $\frac{1}{2}$  length of slit. A series of 7 pores along mandible and preopercle; a second series of 7 or 8 extending from snout along sides of head above premaxillaries. Lateral line faint, descending, its position ventral. Dorsal inserted over middle of pectorals, its origin as far from occiput as is the latter from front of pupil; distance of front of anal from snout equals  $\frac{1}{2}$  length of body; ventrals short, less than length of pupil; pectorals with posterior margin obliquely truncate, the upper rays longest, the lower growing regularly shorter, thickened at tips, the rays 15 or 16, the longest  $2\frac{3}{4}$  in head. Scales small, embedded, covering body and most of vertical fins. A few very small, scattered scales on nape, posterior part of occiput, and contiguous parts of cheeks and opercles. Lateral line single, indistinct, running obliquely downward to near base of anal, thence backward, not reaching base of caudal fin. Color dark brown, black on opercles, sides of snout, fins, and lower parts generally; a broad light bar across head behind eyes, extending down on cheeks; some light mottling on mandible and gular membrane; lower rays of pectorals margined with whitish; lining of mouth, gill cavity, and peritoneum jet black. North Pacific. Two specimens known; the type above described from Santa Barbara Islands, the second from Albatross Station 3210, south of Saanak Islands, Alaska, depth 483 fathoms.

On this Dr. Gilbert has the following notes:

"The stomach contained remains of Crustacea. Colors in life, head and body light brown, the lower parts darker; snout, suborbital region, and a band across pectorals greenish gilt; no light bar on head. Depth  $12\frac{1}{2}$  in length; head  $5\frac{1}{2}$ ; maxillary  $2\frac{1}{2}$  in head; eye 7, equal to interorbital width. Width of bone between orbits 17 in head. Snout  $3\frac{1}{2}$  in head. Teeth above in a narrow band, reaching only about halfway of gape. In the mandible, teeth are absent on posterior  $\frac{2}{3}$  of gape. The gill slit extends a little farther forward below than above. Ventrals as long as pupil. Longest pectoral ray  $2\frac{1}{2}$  in head. Head wholly scaled behind eyes. Lateral line not evident."

(*crotalinus*, from *Crotalus*, κρόταλος, a rattlesnake.)

*Lycodopsis crotalinus*, GILBERT, Proc. U. S. Nat. Mus. 1890, 105, Albatross Station, 2980, off Santa Barbara Islands. (Coll. Albatross.)

## 933. LYCODOPSIS, Collett.

*Lycodopsis*, COLLETT, Proc. Zool. Soc. London 1879, 381 (*pacificus*).

*Leurygnis*, LOCKINGTON, Proc. U. S. Nat. Mus. 1879, 326 (*paucidens*).

Body moderately elongate, the depth 8 to 9 times in length, covered with small, smooth, embedded scales. Lateral line rather faint, extending along middle of side. Head large; snout broad and long; interorbital space very narrow; mouth large, horizontal; teeth conical, those of the upper jaw in a single row; those of the lower in a band in front, the inner series enlarged, larger than the upper teeth; no teeth on vomer or palatines. Ventral fins very small; vertical fins continuous, without spines. Sexes more or less unlike, the mouth larger in the male. Pacific Ocean. (*Lycodes*; *opsis*, appearance.)

## 2815. LYCODOPSIS PACIFICUS (Collett).

Head  $4\frac{1}{2}$  (male) to  $5\frac{1}{2}$  (female); depth 8 (male) to  $8\frac{1}{2}$  (female). D. 100; A. 85. Female (*pacificus*), head comparatively short; orbital region not restricted, nor cheeks tumid; mouth comparatively small, the maxillary reaching center of pupil. Male (*paucidens*), with the head and mouth large, the snout very broad, the interorbital region constricted; maxillary reaching posterior edge of orbit. Head, nape, and axil of pectoral naked. Dorsal and anal fins enveloped in thick skin, which is covered with embedded scales like those on the body; pectoral  $\frac{1}{2}$  the length of head in female,  $\frac{3}{8}$  in male; ventrals  $\frac{1}{2}$  length of orbit; mandible  $\frac{1}{2}$  length of head in female,  $\frac{3}{8}$  in male; distance from snout to base of dorsal  $4\frac{1}{2}$  in length in female,  $3\frac{1}{2}$  in male. Lateral line lateral in portion. Light reddish olive, becoming lighter below; vertical fins margined with black; the scales paler than skin, forming light spots; pectorals dusky. Length 12 to 18 inches. San Francisco to Puget Sound; rather common in water of moderate depth offshore. Sexes markedly different.

*Lycodopsis pacificus*,\* COLLETT, Proc. Zool. Soc. London 1879, 381, female, Japan. (Coll. Peters) the locality given probably an error.

*Leurygnis paucidens*, LOCKINGTON, Proc. U. S. Nat. Mus. 1879, 326, off San Francisco, California, male (Type, No. 23502, U. S. Nat. Mus. Coll. W. N. Lockington); JORDAN & GILBERT, Synopsis, 785, 1883.

*Lycodopsis paucidens*, GILL, Proc. U. S. Nat. Mus. 1880, 248.

*Lycodopsis pacificus*, JORDAN & GILBERT, Synopsis, 785, 1883.

## 934. APRODON, Gilbert.

*Aprodon*, GILBERT, Proc. U. S. Nat. Mus. 1890, 106 (*cortezianus*).

This genus differs from *Lycodes* only in the dentition, the teeth being present in a single strong series on the palatines, but none on the vomer.

\* In regard to the type specimen of *Lycodes pacificus*, Professor Collett writes us as follows (December 2, 1895):

"I got the specimen for describing from the Museum of Berlin from the hands of Professor Peters himself, and he told me that the specimen was from Japan. It is not impossible that he was mistaken; but I can not have any opinion about that."

In view of the fact that the species is abundant off the California coast, whence Professor Peters had obtained collections, that it has not been found in Japan nor in Alaska, we have no doubt that the locality given by Professor Peters is erroneous, and that the fish really came from California.

† The examination of many specimens leaves no room for doubt that *L. pacificus* is the female and *L. paucidens* the male of the same species.

The genus is thus intermediate between *Lycodes* and *Lycodopsis*. ( $\acute{\alpha}$ , without;  $\pi\rho\acute{o}$ , before;  $\delta\delta\acute{o}\upsilon\varsigma$ , tooth.)

2816. APRODON CORTEZIANUS, Gilbert.

Head  $4\frac{1}{2}$  to  $4\frac{3}{4}$ ; depth  $7\frac{1}{2}$  to 9 in length; head high and narrow, snout broader, but long and very convex. Mouth large, maxillary reaching vertical from middle of orbit,  $2\frac{1}{2}$  in head; eye  $4\frac{1}{2}$ ; snout 3; depth of head 2. Teeth in premaxillaries strong, conical, in a single series; lower jaw with the teeth mainly in 2 series, an outer row of slightly enlarged teeth, and an inner row directed backward, a wide interspace between the two series with occasional scattered teeth only posteriorly; on sides of mandible a single series of teeth similar to those in upper jaw; vomer toothless; palatines with a single series of strong conical teeth. Head without conspicuous mucous pores; a strong ridge on middle of occiput anteriorly; gill slit wide, continued forward to vertical from preopercle, the width of isthmus 5 times in length of slit; gill rakers short, better developed than usual, 15 on horizontal limb of outer arch. The vertical limb of arches joined to gill cover by a fold of the lining membrane of the latter, as in *Macrourus*. Pseudobranchiae well developed. Origin of dorsal but little behind base of pectorals; the hinder margin of occiput midway between dorsal and front or middle of eye; distance from snout to origin of anal  $2\frac{2}{3}$  in total length; ventrals inserted under front of opercles, their length about  $\frac{2}{3}$  of orbit; pectorals very large, broadly rounded, the upper portion of fin longest, the lower rays rapidly shortened, the lowermost with broad, fleshy tips; rays 20 or 21 in number; scales of the usual type, those on abdomen so deeply embedded as to be almost invisible; head, anterior half of nape, breast, and base of pectorals naked; pectorals and ventrals not scaled, other fins partly covered; lateral line little developed, running along middle of sides and tail. Color light brownish, lighter below; vertical fins broadly margined with black, becoming almost wholly black behind; pectorals light at base, black distally, with a conspicuous white edge; ventrals white; lining of mouth white, of gill cavity dusky; peritoneum black. Cortez Banks, near San Diego, California. The types, 6 specimens, the longest 15 inches, from *Albatross* Stations 2925 and 2948, in 339 and 266 fathoms. Dr. Gilbert also records 1 specimen from *Albatross* Station 3349, off the coast of northern California, depth 239 fathoms. (*cortezianus*, from Cortez Banks.)

*Aprodon corteziana*, GILBERT, Proc. U. S. Nat. Mus. 1890, 107, Cortez Banks, off San Diego. (Type, No. 46457. Coll. *Albatross*.)

935. LYCODES, Reinhardt.

*Lycodes*, REINHARDT, Kongl. Dansk. Vidensk. Selsk. Naturv., VII, 1838, 153 (*vahlí*).  
*Lycias*, JORDAN & EVERMANN, new subgenus (*seminudus*).

Body moderately elongate, more or less eel-shaped, tapering behind, the depth from 6 to 10 times in the length; head oblong; mouth nearly horizontal; lower jaw included; conical teeth on jaws, vomer, and palatines, those on jaws and palatines mostly in a single series. Dorsal fin beginning behind base of pectoral, without any spines; the rays all soft and articu-

late; pectorals moderate, inserted rather high, its outline rounded; ventral fins small, of 3 or 4 rays. Scales small and embedded, present on part or all of the body, the scaly area more extensive in the adult than in the young. Lateral line faint, sometimes obsolete, normally bent downward behind pectorals and following ventral outline, sometimes with an accessory branch following middle of side; the median branch usually wanting. No air bladder; no anal papilla; pyloric caeca 2 or none. Specles numerous, chiefly of the northern seas, inhabiting considerable depths. In general, the male has the head and mouth larger than the female, and the lips thickened. (*λυκώδης*, wolfish.)

## LYCODES:\*

## I. Trunk more or less completely scaled.

a. Dorsal rays about 115; anal rays 90 to 105.

b. Head  $4\frac{1}{2}$  to 5 in length; depth 7 to 8.

c. Nape wholly scaly.

d. Lateral line double, with a median and a ventral branch; pectoral rays 22; body blackish with yellowish cross bands or series of spots. ESMARKII, 2817.

dd. Lateral line simple, ventral; body blackish, the young with 6 darker cross bands. VAHLII, 2818.

cc. Nape naked; lateral line obsolete; color plain brown, the fins edged with darker, pectoral rays 21; ventrals short. CONCOLOR, 2810.

bb. Head  $5\frac{1}{2}$  in length; depth 9; a naked area around dorsal; pectoral rays 19; lateral line ventral; color brownish mottled, the young barred; a black blotch at front of dorsal. ZOARCHUS, 2820.

aa. Dorsal rays 85 to 105; anal rays 68 to 83.

e. Head large,  $3\frac{3}{4}$  to  $4\frac{1}{2}$  in length; ventrals about as long as eye; depth 8 to  $9\frac{1}{2}$  in length; body chiefly scaly, the fins naked.

f. Body brownish, with a fine network of black lines on head and body, those on body in 5 groups; dorsal edged with black; lateral line probably developed anteriorly only, figured as median; pectorals broad, of about 23 rays. RETICULATUS, 2821.

ff. Body not covered with a network of black lines.

g. Color pale, with dark bands and 2 ocellated spots on the forehead; pectoral rays about 17; lateral line figured as lateral.

PERSPICILLUM 2822.

gg. Color grayish, without bands or spots; pectoral rays about 20; lateral line single, ventral.

FRIGIDUS, 2823.

ee. Head short, 5 to  $5\frac{1}{2}$  in length.

h. Pectoral broad, of 23 or 24 rays; lateral line single, ventral; color plain. TERRE-NOVÆ, 2824.

hh. Pectorals narrow, of about 18 rays; ventral fins shorter than eye; lateral line obsolete, or nearly so.

i. Dorsal rays 101 to 105; anal rays 81 to 90; dorsal and anal without dark markings; ventrals more than  $\frac{1}{2}$  length of eye; jaws with enlarged flaps of skin.

j. Body in adult not barred, but with 4 dark longitudinal stripes. DIGITATUS, 2825.

jj. Body with 14 to 16 pale crossbars above, which disappear in the adult. PALEARIS, 2826.

ii. Dorsal rays 85; anal 74; ventrals minute, not  $\frac{1}{2}$  length of eye; flaps of jaws narrow or obsolete. BREVIPES, 2827.

\* The analytical key to the species here given is far from satisfactory. The species should be divided into groups distinguished by the development of the lateral line and the breadth of the pectoral; unfortunately the last-named character has been neglected in most of the current descriptions; we have examined all the species accessible to us.

LYCIAS (λύκος, wolf):

II. Trunk naked anteriorly, scaled only on the tail or posterior half.

k. Dorsal fin scaled posteriorly; color brown, with faint yellow transverse bands on back. NEBULOSUS, 2828.

kk. Dorsal fin naked; color uniform pale grayish brown without spots or bands; pectoral rays 21; lateral line single, median. SEMINUDUS, 2829.

Subgenus LYCODES.

2817. LYCODES ESMARKII, Collett.

Head  $4\frac{1}{2}$ ; depth 8. D. 110 to 116; P. 22; A. 95; V. 4. Body behind front of dorsal scaled; vertical fins scaly; nape scaly; snout obtuse; maxillary not more than  $\frac{1}{2}$  head; lateral line indistinct, divided, having a median branch besides the ventral series of pores, the median series faint, soon obsolete; pectorals 8 in length; vertebrae 25 + 87. Brownish black, with a whitish-yellow patch on the nape, and 5 to 8 transverse bands of the same color across the dorsal and posteriorly across the anal, these bands becoming broken into annular spots with age. North Atlantic; recorded from Finmark and Spitzbergen. American specimens from the Gulf Stream in about lat. 40°. (Collett.) (Named for Professor Lauritz Esmark, of Copenhagen.)

*Lycodes esmarkii*, COLLETT, Norges Fiske, 95, 1874, Varanger Fjord, Finmark (Coll. Lensmand Klerk and Prof. Esmark); COLLETT, Norske Nord-Havs Exp., Fiske, 84, pl. 3, fig. 22, 1880.

*Lycodes vahlii*, GOODE & BEAN, Proc. U. S. Nat. Mus. 1879, 209, not of REINHARDT; JORDAN & GILBERT, Synopsis, 786, 1883; GOODE & BEAN, Oceanic Ichthyology, 303, 1896.

2818. LYCODES VAHLII, Reinhardt.

Head  $4\frac{1}{2}$ ; depth 8. D. 116; A. 93; V. 4. Head nearly twice as long as high; snout long, maxillary reaching to opposite middle of eye; distance of vent from ventrals nearly equal to length of head; ventral fins less than  $\frac{1}{2}$  as long as pectorals; vertical fins scaly; body wholly scaly; lateral line distinct, ventral in position; vertebrae 25 + 87. Brownish yellow, with 6 blackish cross bands extending on the dorsal fin and confluent on the belly, the first cross band on and below the anterior dorsal rays, the second above the vent; adults nearly uniform blackish. Coast of Greenland. (Günther.) (Named for Martin Vahl, an early Danish naturalist.)

*Lycodes vahlii*, REINHARDT, Kon. Dan. Vidensk. Selsk. Nat. Math. Afh., VII, 1838, 153, tab. V, Greenland; GILL, Cat. Fishes East Coast N. A., 46, 1861; GÜNTHER, Cat., IV, 319, 1862; JORDAN & GILBERT, Synopsis, 786, 1883; GOODE & BEAN, Oceanic Ichthyology, 303, 1896.

2819. LYCODES CONCOLOR, Gill & Townsend.

Head 5 in total; depth about  $7\frac{1}{2}$ ; eye  $7\frac{1}{4}$  in head; snout 3; ventral fin 2 in eye; pectoral 2 in head. D. 118; A. 98; P. 21. Body rather elongate, covered with very small, entirely separated embedded scales which become more distinct anteriorly and extend in advance of the dorsal fin and scapular region, as well as on the vertical fins; lateral line obsolete; pectorals with scattered scales on external and internal surfaces near base; a specialized area of smaller scales behind base of pectoral and a naked area around

upper axilla of pectoral; head moderate, entirely naked; nape naked. Upper teeth in a cardiform band in front, thinning out behind. Lips rather thin. Color nearly uniform; only relieved by the apparently lighter hue of the scales and the somewhat darker margins of the fins; the scales paler than the ground color, which is thus covered with whitish or silvery specks. Bering Sea. Only the type known, its length 22 inches, from which we have taken the above description. (*concolor*, uniformly colored.)

*Lycodes concolor*, GILL & TOWNSEND, Proc. Biol. Soc. Wash., XI, 1897 (Sept. 17, 1897), 233, Bering Sea, lat.  $55^{\circ} 19' N.$ , long.  $168^{\circ} 11' W.$ , Albatross Station 3608, (Aug. 12, 1895), in 276 fathoms. (Type, No. 48764, U. S. Nat. Mus. Coll. *Albatross*.)

2820. *LYCODES ZOARCHUS*, Goode & Bean.

Head nearly  $5\frac{1}{2}$  in total length; depth 9; eye 4 in head = snout. D. 116; A. 102; P. 19. Body covered with conspicuous embedded scales which extend behind the dorsal and anal, leaving only a narrow naked margin around these fins; head and pectorals naked. A lateral line begins slightly above the upper angle of the gill opening, rapidly curving downward and extending along the lower part of the body not far from base of anal fin; it can be traced above the anterior  $\frac{2}{3}$  of the anal. Interorbital distance, measured on the bone, 4 in eye; nostrils placed close to upper lip and as far from each other as from the eye; maxillary reaching to vertical through middle of eye; upper jaw  $2\frac{1}{2}$  in head; mandible nearly  $\frac{1}{2}$  head; mandible with a conspicuous flap on each side, about as long as eye, beginning at a distance from the symphysis equal to  $\frac{1}{2}$  length of eye; inner edge of mandible also with a slightly elevated ridge of skin. Length of intermaxillary series of teeth equal to  $\frac{1}{2}$  length of head; length of palatine series nearly equal to that of intermaxillary; vomerines in a round patch; mandibular teeth in 3 series; width of gill opening  $\frac{2}{3}$  length of head; ventrals in front of base of pectorals, their length 8 in head. Distance between lower angles of gill opening nearly  $\frac{1}{2}$  length of head; origin of dorsal distant from the head a space equal to  $\frac{1}{2}$  length of head, slightly behind middle of pectoral; pectoral, when extended, reaching to about vertical from sixth dorsal ray; longest ray of dorsal about  $\frac{1}{2}$  length of head; anal origin under seventeenth ray of dorsal; vent under fifteenth ray of dorsal; longest pectoral ray contained about  $9\frac{1}{2}$  times in total length. Lateral line distinct, ventral in position, the median pores absent. Color grayish brown, lighter on the belly and under surface of the head; sides irregularly mottled with darker, a narrow dark edge at tip of first 4 dorsal rays. In a young example (No. 39299, U. S. Nat. Mus.) the mottlings on the sides are band-like, the bands not extending below the middle of the body entirely. This example is from lat.  $44^{\circ} 26' N.$ , long.  $57^{\circ} 11' 15'' W.$ , 190 fathoms. The type of the description is a specimen 366 mm. long, obtained by the *Albatross* in lat.  $44^{\circ} 46' 30'' N.$ , 130 fathoms, off Nova Scotia. (*Zoarchus*, a synonym of *Zoarcis*; from *ζωαρχίς*, viviparous.)

*Lycodes zoarchus*, GOODE & BEAN, Oceanic Ichthyology, 308, 1896, off Nova Scotia, in 130 fathoms. (Type, No. 39298. Coll. *Albatross*.)



2821. LYCODES RETICULATUS, Reinhardt.

Head 4; depth about 8. D. 94; A. 75; V. 4. Body entirely scaly; lateral line faint, developed anteriorly (fide Günther's plate), probably becoming ventral; vertical fins naked. Head twice as long as high; snout long; maxillary extending to behind middle of eye; distance from vent to ventrals more than length of head; caeca 2. Brownish, with reticulated black lines on the head and body, those on the body disposed in 5 groups or cross bands, the 3 anterior of which emit 1 or 2 vertical streaks on the dorsal fin; dorsal dark edged. Length 14 inches. North Atlantic, from Greenland south to Narragansett Bay, in 17 to 140 fathoms; abundant also in northern Europe. (Eu.) (*reticulatus*, netted.)

*Lycodes reticulatus*, REINHARDT, Kong. Dansk. Vid. Afh., VII, 1838, 167, Greenland; GÜNTHER, Cat., IV, 320; GILL, l. c., 260; COLLETT, Nord-Hava Exp., 84; JORDAN & GILBERT, Synopsis, 787, 1883; GOODE & BEAN, Oceanic Ichthyology, 305, 1896.

*Lycodes rossi*, MALMGREN, Om Spetsbergen Fiskfauna, 516, 1864, Spitzbergen.  
*Lycodes gracilis*, SABS, Christ. Vid. Selsk. Forh. 1866, Dröbak.

2822. LYCODES PERSPICILLUM, Kröyer.

This species is distinguished by a light body color and dark bands, also 2 ocellated spots on the forehead, which have suggested the specific name. Still further separated from the previously known species of *Lycodes* by the smaller number of fin rays, larger eye, etc. (Kröyer.) Greenland and southward in deep water. Specimens were obtained by the *Albatross* from Station 2491, in 45° 24' 30" N. lat., 58° 35' 15" W. long., at a depth of 59 fathoms, and from Station 2456, in 47° 29' N. lat., 52° 18' W. long., at a depth of 86 fathoms.

The following is the substance of Dr. Günther's description:

Head 4 in total length; depth nearly 8. Head not quite twice as long as high; snout long; upper maxillary extending to below middle of eye. Distance of vent from ventrals nearly equal to length of head. Yellowish, with 9 or 10 brownish cross bands, edged with dark brown, and broader than the interspaces, the first occupying the upper parts of the head and inclosing a pair of roundish, yellowish spots situated behind the level of the eyes; the second cross band is on and before the anterior dorsal rays.

(In the figure of Goode & Bean the lateral line is represented as median, which is probably not correct.) (*perspicillum*, eyebrow, from the spot above the eye.)

*Lycodes perspicillum*, KRÖYER, Dansk. Vidensk. Selsk. Afhandl., XI, 1845, 233, Greenland; GÜNTHER, Cat., IV, 320, 1862; GILL, Proc. Ac. Nat. Sci. Phila. 1863, 260.

2823. LYCODES FRIGIDUS, Collott.

Head 4 to 4½ in total length; depth 6½ (to 9½, young). D. 93 to 98 (including ½ of the caudal, 99 to 104); A. 80 to 85 (including ½ of the caudal 86 to 90); P. 20 to 21; V. 3. Head wide and flat. Scales with very conspicuous mucous cavities below, small, covering the entire body, but not the head, nor the base of the dorsal and anal fins. In the young the middle of the belly, the base of the fins, and the fins themselves are usually naked.

Teeth present on intermaxillary, mandible, palatines, and vomer; lateral line low, extending from upper end of gill opening in a curved direction down toward vent from which it runs close along anal to end of tail. (Goode & Bean.) Pectoral fin obliquely truncate at tip, appearing furcate when not spread open. North Atlantic and Arctic Ocean, from Spitzbergen south to the New England coast, where many specimens were taken in 516 to 1,423 fathoms. (En.) One of Collett's specimens from Hammerfest, examined by us.) (*frigidus*, frozen.)

*Lycodes frigidus*, COLLETT, Forh. Selsk. Christ. 1878, Nos. 14 and 15, Beeren Island and Spitzbergen; COLLETT, Norske Nord-Havs Exp., 96, pl. 3, f. 23, 24, 1880; GOODE & BEAN, Oceanic Ichthyology, 305, 1896.

2824. LYCODES TERRE-NOVE, Collett.

Head 5 to 5½; depth 8 to 11. D. 106 to 108; A. 89 to 93; P. 23 or 24. Body slender, head small; pectorals broad; maxillary reaching to middle of eye; band of palatine teeth very short, scarcely ½ length of maxillary band; body entirely scaly, head naked; lateral line ventral, extending along edge of belly, the median branch wanting; vent before middle of body. Color lost in type, the only specimen known. Banks of Newfoundland, in 155 fathoms. (Collett.) (*terra*, land; *novus*, new, from Newfoundland.)

*Lycodes terre-nove*, COLLETT, Campagnes Scientifiques, L'Hirondelle, x, 1896, 54, Bank of Newfoundland, Hirondelle Station 162, in 155 fathoms. (Coll. Albert, Prince of Monaco.)

2825. LYCODES DIGITATUS, Gill & Townsend.

Head 5 in total; depth about 8½; eye 6½ in head; snout 3; ventral fin 1½ in eye; pectoral 1½ in head. D. 101; A. 81; P. 18. Body moderately elongate; covered with small, entirely separated embedded scales, which become nearer anteriorly and extend in advance of the dorsal fin as well as on the vertical fins; no specialized area of smaller scales behind base of pectorals; pectorals scaleless; head moderate, entirely naked; nape naked; upper jaw with outer row of close-set teeth, broader in front; teeth on vomer and palatines; lips rather thick. Color in alcohol, brownish yellow, suffused with reddish in front, variegated, darker anteriorly, with 4 dark longitudinal stripes most distinct about middle of body, fading out backward; fins light and without dark margins; head dark above and laterally light below. Bering Sea. Only the type known, from which we have taken this description, its length 18 inches; possibly the adult of *L. palearis*, but the pectoral fins are shorter than in the latter. (*digitatus*, fingered.)

*Lycodes digitatus*, GILL & TOWNSEND, Proc. Biol. Soc. Wash., XI, 1897 (Sept. 17, 1897), 232, Bering Sea, lat. 56° 14' N., long. 164° 8' W., at Albatross Station 3541, in 49 fathoms. (Type, No. 48765, U. S. Nat. Mus. Coll. Albatross.)

2826. LYCODES PALEARIS, Gilbert.

Head 5½ in length; depth 9½ to 11 in length, 2½ in head; eye 5 to 6 in head, 1½ to 2 in snout. Dorsal with about 105 rays, counted to middle of caudal; anal about 90; pectoral 18; ventrals 1½ to 1½ in eye, twice as long

as in *L. brevipes*; pectorals  $1\frac{1}{2}$  in head. Head naked; nape more or less naked, the scaleless area variable in extent, sometimes confined to its anterior third, sometimes reaching nearly to front of dorsal; body sparsely covered with embedded scales; axil naked; lateral line short, decurved, extending scarcely beyond middle of pectorals. Anal origin under eighteenth dorsal ray. Teeth present in jaws, vomer and palatines, those in premaxillaries laterally in a single series which widens anteriorly into a rather broad patch, the outer teeth somewhat enlarged, especially in front; all the premaxillary teeth shut outside on the mandibular series which are opposed to those on vomer and palatines; mandibular teeth arranged similarly to those in upper jaw, the lateral series somewhat enlarged, continuous with the inner edge of the symphyseal patch; vomerine teeth bluntly conic, 3 or 4 in number; palatines in a single series. Snout long, prominent, the upper jaw projecting beyond the lower for a distance equaling  $\frac{2}{3}$  of orbit; upper lip thin, much expanded laterally, continuous posteriorly with the lower lip which forms a wide free membranaceous lobe opposite middle of each mandible; anteriorly the lower lip becomes abruptly contracted and adnate to the jaw, leaving the symphyseal portion without free margin; inner edge of mandible with wide membranaceous borders, which increase in width anteriorly where they terminate in a pair of acutely pointed free flaps; these and the membranaceous margins very conspicuous in both young and old individuals. In *L. brevipes* they are very inconspicuous, becoming evident in adults only. General color brownish olive, growing lighter on the lower parts; dorsal with 14 to 16 white vertical bars, extending in young examples across back and sides and onto anal fin, in adults confined to the fins, and frequently indistinct or wanting; anterior dorsal angle frequently black, separated from remainder of fin by a curved white bar; dorsal and anal not black margined as in *L. brevipes*; in the latter, the white lateral bars are 9 to 12 in number, and are usually confined to upper half of body; there is also no black spot on anterior dorsal rays. This species is very close to *L. brevipes* Bean, differing constantly in the longer ventrals, the greater development of mandibular and labial folds, the more numerous white bars, and the smaller eye. Bering Sea. Three specimens, 113 to 166 mm. long, from Albatross Stations 3253 and 3254, in Bristol Bay, in 36 and 46 fathoms. (Gilbert.) (*palea*, the wattles of a cock.)

*Lycodes palearis*, GILBERT, Rept. U. S. Fish Comm. 1893 (1896), 454, Bristol Bay, Alaska. (Coll. Albatross.)

2827. LYCODES BREVIPES, Bean.

Head 5 in total length; depth 10. D. 85 to middle of caudal; A. 74; P. 21. Body covered with scales except immediately behind pectoral fins; head naked; dorsal and anal fins minutely scaled; diameter of eye equals the length of the snout, 4 in head; dorsal origin nearly over middle of pectoral; anal origin under eighteenth ray of dorsal; ventrals minute, scarcely more than  $\frac{1}{2}$  diameter of eye; pectorals 9 in length of the body; lateral line single, very faint, ventral in position, abruptly decurved and becoming obsolete over about the tenth anal ray. A narrow light band

across the nape and from 9 to 11 across the back extending downward about to median line and becoming obscure in adults; dorsal and anal each with a narrow dark margin. (Bean.) Aleutian Islands to Kadiak; abundant; taken by us in large numbers off Karluk in 1897. (*brevis*, short; *pes*, foot.)

*Lycodes brevipes*, BEAN, Proc. U. S. Nat. Mus. 1890, 38, between Unga and Nagai islands, at Albatross Station 2848, in 110 fathoms. (Type, No. 45382. Coll. *Albatross*.)

Subgenus LYCIAS, Jordan & Evermann.

2828. LYCODES NEBULOSUS, Kröyer.

D. 87; A. 68; P. 19; V. 3. Body naked anteriorly, the posterior part of dorsal fin scaly; the anal naked or nearly so. Brown, with small, faint, yellow, transverse bands across the back. Greenland. (Kröyer.) An imperfectly described species, not recognized by any recent writer. This species and the next should perhaps be placed in *Lycodalepis*. (*nebulosus*, clouded.)

*Lycodes nebulosus*, KRÖYER, Kong. Dan. Vidensk. Sel. 1844, 140, Greenland; GILL, Proc. Ac. Nat. Sci. Phila. 1863, 261; JORDAN & GILBERT, Synopsis, 787, 1883.

2829. LYCODES SEMINUDUS, Reinhardt.

Head  $3\frac{1}{2}$ ; depth 7. D. 91; A. 74; P. 21. Body naked in front of vent, scaly behind; fins naked. Head large. Distance of ventrals from vent somewhat more than length of head; caeca 2. Color uniform pale grayish brown, without spots or bands. North Atlantic, from Greenland to Spitzbergen; rare. (Collett.) (*semi*-, half; *nudus*, naked.)

*Lycodes seminudus*, REINHARDT, Kong. Dansk. Selsk., etc., 1838, 221, Omenak, Greenland; GÜNTHER, Cat., IV, 320, 1862; JORDAN & GILBERT, Synopsis, 787, 1883; GOODE & BEAN, Oceanic Ichthyology, 307, 1896.

936. LYCODALEPIS, Bleeker.

*Lycodalepis*, BLEEKER, Verl. Akad. Amst., Ed. 2, VIII, 1874, 369 (*mucosus*).

This genus differs from *Lycodes* in the absence of scales on trunk and fins; scattered scales sometimes present on the tail only. (*λυκόδης*, *Lycodes*; *ἀλεπίς*, without scales.)

a. Color brownish, with many cross bands and streaks of cream color; head  $4\frac{1}{2}$  in length; depth 8; lateral line obsolete; tail sometimes with a few scales. POLARIS, 2830.

aa. Color blackish, with about 5 narrow pale cross bars on back; head  $3\frac{1}{2}$  in length; depth 8; lateral line double, a median and a ventral series of pores being faintly developed. MUCOSUS, 2831.

2830. LYCODALEPIS POLARIS (Sabine).

Head  $4\frac{1}{2}$ ; depth 8. D. 85; A. 67; P. 18; V. 3; Br. 6. Head depressed, its greatest width  $\frac{2}{3}$  of its length; distance from tip of snout to nape nearly equaling greatest width of head, 6 in length; upper jaw  $1\frac{1}{2}$  to 2 in head, extending to vertical of hind margin of orbit, larger in male than in the female; a full series of teeth on premaxillaries, and in front of these a few smaller teeth form an outer imperfect series; a toothless space at symphysis,

first tooth on each side of this larger than any of the rest; 1 complete series of teeth on mandible, and in front of it, about the symphysis, 2 irregular short series; a few teeth in a cluster on head of vomer; palatines with a short single series; teeth all slender and slightly recurved; long diameter of eye 9 in head. Pectoral  $1\frac{1}{2}$  in head; ventral about as long as eye; longest dorsal ray  $3\frac{7}{8}$  in head; vent in middle of total length, immediately behind third cross band; longest anal ray  $4\frac{1}{2}$  in head; scattered scales present on posterior two-thirds of tail in 1 specimen (type of *L. coccineus*), wholly wanting in the others, typical of *L. turneri*; no scales on the fins; no trace of lateral line. Color light brown; abdomen grayish brown; lower parts of head cream; a band of cream on the anal from origin of rays to about their middle; a crescentic V-shaped band of same color, mottled with amber, crossing nape and continuing behind pectorals, extending backward to the first cross bar; a streak of cream, more or less interrupted by amber, extending backward from eye across cheek almost to end of operculum; 10 bands of cream color, bordered with dark amber, from tips of dorsal rays extending on lower half of body, becoming wider and somewhat broken below middle of body; a very indistinct caudal tip of cream color. In young examples these markings are very distinct; in older ones they grow progressively more obscure, the oldest having scattering blotches of cream color instead of bands, the V-shaped nuchal band persisting longest. The type of *Lycodes coccineus* is described as brown, red below; pectorals reddish brown above, carmine below; 9 bluish-white bands on the dorsal; a few whitish blotches on sides and on head; anal brownish red; head white below; a whitish blotch as large as eye at upper angle of gill opening. Length 18 to 20 inches. Arctic Ocean, Bering Straits, and adjacent waters south to St. Michaels. Here described from the type of *Lycodes coccineus* and from a number of specimens from Point Barrow referred to *Lycodes turneri*. Evidently all belong to the same species, but 1 has a scaly tail while the others are wholly naked. In 2 large examples, supposed to be males, the head is very much depressed, broad and flat, and the maxillary is more than  $\frac{1}{2}$  head. In the others the head is smaller, less flattened, with smaller mouth, the maxillary 2 in head. These are doubtless females and young. The species should probably stand as *Lycodalepis polaris*. (*polaris*, polar.)

*Blennius polaris*,\* SABINE, Parry's Journal, Voyage 1810-20, Supplement, 212, North Georgia.

*Lycodes turneri*, BEAN, Proc. U. S. Nat. Mus. 1878, 464, St. Michaels, Alaska (Type, No. 21520. Coll. Dr. Lucien M. Turner); TURNER, Contr. Nat. Hist. Alaska, 93, pl. 4, 1880.

*Lycodes coccineus*, BEAN, Proc. U. S. Nat. Mus., IV, 1881, 144, Big Diomedes Island, Bering Strait (Coll. Dr. Bean. Type, No. 27748, 20 inches long, with scales on the tail); JORDAN & GILBERT, Synopsis, 787, 1883.

*Blennius (Zoarches?) polaris*, RICHARDSON, Fauna Bor.-Amer., III, 94, 1836.

*Lycodes polaris*, GÜNTHER, Cat., IV, 321, 1862.

*Lycodalepis turneri*, JORDAN & GILBERT, Synopsis, 788, 1883; SCOFIELD, in JORDAN & GILBERT, Fur Seal Invest., 1898.

*Lycodalepis polaris*, JORDAN & GILBERT, Synopsis, 788, 1883.

\* *Blennius polaris* is thus described: Without any scales; length of the pectoral exceeding twice its breadth, having 15 rays. Yellowish, lighter on the belly, with 11 large saddle-like markings across the back, the middle of these markings being much lighter than their edges; the whole back and the sides marbled. (Sabine.) Coast of North Georgia.

## 2831. LYCODALEPIS MUCOSUS (Richardson).

Head  $3\frac{1}{2}$ ; depth 8. D. (including  $\frac{1}{2}$  of caudal) 90; A. (including  $\frac{1}{2}$  of caudal) 71; P. 18; V. 3. Body robust, head very large; snout 3 in head; interorbital area 6 in head; nostrils much farther from eyes than from each other, their distance from eyes  $4\frac{1}{2}$  in head; upper jaw  $6\frac{1}{2}$  in total length; lower jaw  $6\frac{1}{2}$ ; eyes small, close together, their long diameter 11 in the head; distance from tip of snout to base of pectoral fin  $3\frac{1}{2}$  in total length; pectoral fin  $6\frac{1}{2}$ ; length of ventrals equaling long diameter of eye. (Goode & Bean.) Lateral line (in specimens from Cumberland Gulf) very faint, but with both median and ventral branch. Blackish, with irregular white markings in the form of 5 faint and narrow bars across the back. Arctic America. (*mucosus*, slimy.)

*Lycodes mucosus*, RICHARDSON, Last Arctic Voyage, 362, pl. 26, 1855, Northumberland Sound; BEAN, Bull. U. S. Nat. Mus., No. 15, 112, 1879; GOODE & BEAN, Oceanic Ichthyology, 306, 1896.

*Lycodalepis mucosus*, JORDAN & GILBERT, Synopsis, 788, 1883.

## 937. LYCENCHELYS, Gill.

*Lycenchelys*, GILL, Proc. Ac. Nat. Sci. Phila. 1884, 110 (*muræna*).

This genus contains small and very slender species differing from *Lycodes* in the elongation of the body, the depth being from 10 to 20 times in the length. The lateral line is single and median in all known species. The genus is very close to *Lycodes*, but the position of the lateral line sufficiently defines it, especially in connection with the slender eel-like form. (*λύκος*, wolf; *ἔγχελυς*, eel.)

- a. Lower half of pectoral not notably longer than upper; depth 12 to 16 in length.  
b. Dorsal rays 92; anal 88; color grayish, with irregular brown patches.

VERRILLII, 2832.

- bb. Dorsal rays 118; anal 110; color brown, the head darker.

PAXILLUS, 2833.

- aa. Lower half of pectoral considerably longer than upper; head with large pores; depth 14 times in length; color dusky brown.

FORIFER, 2834.

## 2832. LYCENCHELYS VERRILLII (Goode &amp; Bean).

Head  $5\frac{1}{2}$ ; depth about 13; eye 2 in snout. D. 92; A. 88; P. 15; V. 5. Body elongate; head much depressed. Distance of vent from ventrals slightly greater than head, its distance from snout about 3 in body; distance of dorsal fin from snout  $\frac{1}{2}$  greater than head; distance of anal from snout twice head; dorsal and anal fins about equal in height, with even margins, not differentiated from caudal, the rays increasing somewhat in length posteriorly; distance of pectoral from snout about equal to head, twice length of pectoral; pectoral reaching vertical from base of second dorsal ray; distance of ventrals from snout less than head, their length less than  $\frac{1}{2}$  that of pectorals. Head, body, and fins enveloped in tough, lax skin. Scales cycloid, circular, and ovate, with numerous concentric striae, and about 18 lobes on margin, the whole perimeter being lobed; scales deeply embedded in the skin at distances from each other equal to their own diameters, most numerous on upper part of body and

extending upon base of dorsal; very few scales upon lower half of body, none on anal fin. Upper jaw far overlapping the lower; gape reaching orbit. A series of 6 large pores on each side, extending backward from nostril toward angle of opercle, the fourth of the series under center of orbit; a similar series, 7 on each side, along line of lower jaw from its symphysis to angle of opercle, all slit-like, the others circular. Nostrils at extremities of fleshy tubes. Teeth in lower jaw in 2 rows, nearly uniform in size; teeth of upper jaw in a single series, somewhat enlarged near the symphysis; patches of smaller teeth behind; about 7 teeth on vomer; a single row on palatines; all the teeth curved. Gill opening narrow, the membranes attached to the isthmus. Color, body above lateral line light grayish brown with numerous minute circular dots marking the position of the scales; pearly white below lateral line; brown irregular patches upon sides, bisected by lateral line, the lower half color of dorsal, that above darker and with the white dots, these brown patches 7 to 10 in number; a brown spot on tip of tail; abdominal region livid blue. Coast of Massachusetts, in deep water; a dwarf species very small in size. (Named for Prof. Addison E. Verrill of Yale University.)

*Lycodes verrillii*, GOODE & BEAN, Amer. Journ. Sci. Arts, XIV, 1877, 474, off coast of New England in the Gulf Stream; JORDAN & GILBERT, Synopsis, 786, 1883.

*Lycenchelys verrilli*, JORDAN, Cat., 124, 1885; GOODE & BEAN, Oceanic Ichthyology, 309, figs. 277 and 277 A, 1896.

2833. LYCENCHELYS PAXILLUS (Goode & Bean).

Head 8; depth 16; eye  $3\frac{1}{2}$  to 4 in head, equal to snout, which is 4 times interorbital width. D. (with  $\frac{1}{2}$  of caudal) 118; A. 110; P. 16; V. 3. Body attenuate, head broad, flat above, with declivous profile; cheeks full and protuberant; teeth stout, recurved, and sharply pointed, in a single series in each jaw, except at the symphysis; a few teeth clustered at the head of the vomer; palatines with a single series; the tubular nostril much nearer tip of snout than eye. Lateral line median, faint and short (in specimens examined by us). Dorsal beginning over tip of pectoral; ventral little longer than pupil. Scales very small, present everywhere except on head and pectorals, nearly covering vertical fins. Light brown, the head somewhat darker. Gulf stream, lat.  $35^{\circ}$  to  $41^{\circ}$  N., in deep water, 263 to 904 fathoms. (Goode & Bean.) (*paxillus*, a peg.)

*Lycodes paxillus*, GOODE & BEAN, Proc. U. S. Nat. Mus. 1879, 44, between LaHave and Sable Island Banks (Type, No. 22177. Coll. Capt. J. W. Collins), a male in breeding form; JORDAN & GILBERT, Synopsis, 785, 1883.

*Lycodes paxilloides*, GOODE & BEAN, Bull. Mus. Comp. Zool., X, No. 5, 207, 1883, off Newfoundland (Type in M. C. Z. Coll. *The Blake*); a normal, not sexually distorted individual.

*Lycenchelys paxillus*, JORDAN, Cat., 124, 1885; GOODE & BEAN, Oceanic Ichthyology, 311, figs. 279 and 282, 1896.

2834. LYCENCHELYS PORIFER (Gilbert).

Head  $5\frac{3}{4}$ ; depth 14. Body very slender. Head much contracted opposite orbits, the snout expanded, as in *Lycodopsis paucidens*. Mouth moderate, the maxillary reaching vertical from front of pupil,  $3\frac{1}{2}$  in head; eye  $5\frac{1}{2}$ ; snout  $3\frac{1}{2}$ ; interorbital width  $\frac{1}{2}$  eye. Teeth in front of premaxillaries

in 2 series, merging into 1 laterally, the outer series anteriorly somewhat enlarged; teeth in front of mandible in a broad band, narrowing laterally to a single series, none of them enlarged; vomer and palatines with single series. Head with 2 series of large and very conspicuous elongate pores, 1 series on mandible and subopercle, the second parallel with it on level of snout. Gill openings wide, extending forward beyond preopercular margin, the width of isthmus less than  $\frac{1}{2}$  length of slit. Distance from origin of dorsal to tip of snout  $4\frac{1}{2}$  in length. Median dorsal rays simply forked near base, those posteriorly in both dorsal and anal repeatedly subdividing. Distance of anal from snout  $2\frac{1}{2}$  in length; pectorals rounded, the lower half of fin longer than the upper, the rays thickened, the fin containing 15 or 16 rays, its length less than  $\frac{1}{2}$  head; ventrals longer and slenderer than usual, each apparently composed of 2 rays closely joined, their length  $\frac{1}{2}$  orbit, inserted unusually far forward, being in advance of preopercular margin. Scales very small, circular, partially embedded, covering body and vertical fins; head, antedorsal region, breast, and a strip connecting the two latter embracing base and axil of pectorals, naked; lateral line median. Color dusky brown, the fins, sides of head, and belly blackish; lining of mouth and gill cavity and peritoneum black. Off Lower California. A single specimen, 12 inches long, from Albatross Station 3009, in 857 fathoms. A transitional species approaching *Furcella*. (*porus*, pore; *fero*, I bear.)

*Lycodes porifer*, GILBERT, Proc. U. S. Nat. Mus. 1890, 104, off Lower California, in 857 fathoms. (Type, No. 44384. Coll. Dr. Gilbert.)

#### 938. FURCIMANUS, Jordan & Evermann.

*Furcimanus*, new genus (*diapterus*); JORDAN & EVERMANN, Check-List Fishes, 480, 1896 (*diapterus*); preoccupied by *Furcella*, Lamarck, 1801, a genus of mollusca.

This genus differs from *Lycenchelys* in the forked pectorals, the upper and lower rays being much longer than the middle ones. The lateral line is single and ventral in position (not lateral as in *Lycenchelys*). (*furca*, a fork; *manus*, hand.)

#### 2885. FURCIMANUS DIAPTERA (Gilbert).

Head  $5\frac{1}{2}$  to 6; depth 12; eye large, usually longer than snout, 3 to  $3\frac{1}{2}$  in head; snout  $3\frac{1}{2}$  to  $3\frac{3}{4}$ ; interorbital width about 10. Body slender. Mouth small, somewhat variable in length, the maxillary reaching vertical from between front and middle of pupil,  $2\frac{1}{2}$  to 3 in head. Teeth in premaxillaries in a double row throughout, the 2 series well separated, rarely with 1 or 2 teeth intercalated, showing traces of a third row; the teeth of inner series small and directed obliquely inward; those of outer series anteriorly enlarged, becoming smaller on sides of jaw; on front of mandible the teeth are in a broad band, in which traces of 3 or 4 irregular series can be made out; none of these enlarged; laterally the teeth are arranged in a single series, those opposite middle of cleft considerably enlarged; a small patch of from 2 to 5 teeth on vomer; palatines with a single row much shorter than premaxillary patch. Nostril with a short inconspicuous tube. Mandible and preopercular border with deep pit-like excavations, which are not evident in fresh specimens; no evident mucous pores on the head.



Gill openings wide, extending below the base of the pectorals; the gill membranes joined to isthmus for a distance equaling  $\frac{3}{4}$  length of slit; gill rakers very short, almost tubercular, but compressed and slightly movable, about 15 present on anterior limb of outer arch; a wide slit behind fourth gill. Ventrals short, inserted under middle of opercle. Pectorals deeply notched in both young and adults, the median rays much shorter than either upper or lower, the lobe produced by the elongate lower rays varying in length, being sometimes shorter than upper lobe, sometimes longer; the rays of lower lobe are thickened, and undoubtedly serve as a support to the fish when resting on the bottom, as has been observed in so many other forms; the pectorals contain 20 or 21 rays; in the structure of this fin the present species seems to differ from all previously described forms, with the exception of *L. esmarkii*, in which the notched condition of the fin does not persist in the adults. Scales small, embedded, covering entire body and vertical fins; the scales on nape are much reduced in size, and in 2 specimens ( $11\frac{1}{2}$  and  $7\frac{1}{2}$  inches long) are continued onto occiput, which they entirely cover; in another specimen, 9 inches long, the occiput is naked, and in another, 5 inches long, the anterior part of nape is likewise naked; in the latter, as in other specimens, the dorsal and anal are well scaled. Lateral line single, wavy, ventral in position, extending from above gill slit obliquely downward to near base of anal, along which it is continued for a variable distance, not reaching base of caudal. Color dusky brownish, blue-black on belly and along anterior portion of base of anal; 8 or 9 narrow white bars on sides, most conspicuous in the young, in which they are continued up on dorsal fin and become forked below on middle of sides, forming A-shaped marks; in adults these bars become faint or wholly disappear; when present, they are not continued on dorsal, and are usually vertically divided by a streak of the ground color; in the small specimen there is a distinct black blotch on margin of anterior dorsal rays; in adults, the vertical fins are brownish on basal portion, their distal half black; pectorals and ventrals deep blue-black; mouth, gill cavity, and peritoneum dusky or black. Several specimens, from *Albatross* Stations 2892, 2896, 3067, and 3077, in depths from 82 to 376 fathoms, off the coasts of California and Oregon. (Gilbert.) A remarkable species. (*δίá*, divided; *πρεπόv*, fin.)

*Lycodes diapterus*, GILBERT, Proc. U. S. Nat. Mus. 1891, 564, off the coast of Oregon, in 685 to 877 fathoms. (Type, No. 44385. Coll. Dr. Gilbert.)

### 939. LYCODONUS, Goode & Bean.

*Lycodonus*, GOODE & BEAN, Bull. Mus. Comp. Zool., x, No. 5, 208, 1883 (*mirabilis*).

Body elongate, formed as in *Lycenchelys* and *Lyciscus*; scales small, circular, embedded in the skin; lateral line very short or obsolete; jaws without fringes, lower jaw included; fin rays all articulated, each ray of dorsal and anal supported laterally by a pair of sculptured scutes; caudal distinct, not fully connate with dorsal and anal; ventrals present; gill opening narrow; teeth as in *Lycodes*. Deep water. (*Lycodes*, with a meaningless change of termination.)

2836. *LYCODONUS MIRABILIS*, Goode & Bean.

Head 7 in total length; depth about 18. D. about 80; A. about 70; C. 9; P. 18; V. 3; scales as in *Lycodes*, the scales not extending out upon the fins; no scales on head and nape. Lateral line apparently obsolete posteriorly; not extending back of the extremity of the pectoral, its position median; eye high up,  $2\frac{1}{2}$  in head, equal to postorbital portion of the head; the width of interorbital space less than diameter of pupil,  $3\frac{1}{2}$  times in long diameter of eye; nostrils immediately in front of eye; maxillary extending to vertical through anterior margin of pupil; mandible, to a little behind vertical through posterior margin of the pupil; dorsal fin inserted slightly behind vertical through base of pectoral (the portion of the fin present in the mutilated specimen before us contains 80 articulated rays; the first 10 or 11 scutes do not support rays, but whether rays were originally present or not can not be ascertained); longest dorsal ray about equal to longest anal ray, its length about 3 in head; distance of vent from snout twice length of head; anal beginning immediately behind vent, of about 70 articulated rays; caudal rays extending beyond tips of ultimate dorsal and anal rays, about 9 in number; distance of ventral from snout equal to twice length of upper jaw; middle ventral ray longest, it being  $\frac{1}{2}$  as long as postorbital part of head; length of pectoral equaling 3 times that of snout. Off the New England coast, in depths of 721 to 1,309 fathoms; a most remarkable little fish. (*mirabilis*, wonderful.)

*Lycodonus mirabilis*, GOODE & BEAN, Bull. Mus. Comp. Zool., x, No. 5, 208, 1883, New England Coast, lat.  $38^{\circ} 20' 8''$  N., long.  $73^{\circ} 23' 20''$  W., in 740 fathoms (Type in M. C. Z.); JORDAN, Cat. Fishes, 124, 1885; GOODE & BEAN, Oceanic Ichthyology, 312, 1896.

940. *LYCONEMA*, Gilbert.

*Lycinema*, GILBERT, Rept. U. S. Fish Comm. 1893 (1896), 471 (*barbatum*).

Generic characters as in *Lycodes*, but the lower jaw covered with a dense mass of slender filaments or barbels, between which can be seen the mucous pores of the mandible. In *Iluocates*, a related genus from the Antarctic, the mandible is provided with a series of hollow tubes, which are doubtless the produced margins of the pores. Alaska. (*λύκος*, wolf; *νήμα*, thread.)

2837. *LYCONEMA BARBATUM*, Gilbert.

Head  $6\frac{1}{2}$ ; depth  $11\frac{1}{2}$ ; maxillary 3 in head; eye  $3\frac{1}{2}$ ; snout  $4\frac{1}{2}$ . D. 103; A. 90 (each counted to middle of caudal); P. 15, its length  $1\frac{1}{10}$  in head; ventrals very short,  $\frac{1}{2}$  to  $\frac{2}{3}$  diameter of orbit. A dense fringe of filaments covers the entire under surface of lower jaw, extending to behind angle of mouth; another series laterally on the throat, and a few scattering ones sometimes present on the branchiostegal membranes; upper jaw without barbels. Body slender; upper jaw overlapping the lower; mouth small, maxillary reaching vertical from front of pupil; teeth all conical, none of them much enlarged, those in lower jaw in a patch or irregular double series, narrowing to a single series laterally; in upper jaw, a single series, the teeth of which increase in size toward the middle line, the mid-

the teeth being almost canine-like; behind the latter, a short inner series of small teeth directed backward; teeth on vomer and palatines in a single series. Gill slits continued forward to slightly beyond bases of ventrals, and to level of lower edge of base of pectorals; width between gill slits  $\frac{1}{2}$  diameter of eye; pseudobranchia well developed; posterior line of occiput midway between origin of dorsal and front of pupil or front of eye; origin of anal fin at end of first third of length of body; pectorals broad, with the posterior edge emarginate, some of the upper and the lower rays longer than the intermediate ones. Scales showing traces of definite arrangement in series, widely separated anteriorly, becoming crowded toward end of tail, continued up on the vertical fins, but not on head, on anterior half of snout, nor on the pectoral fins; lateral line very faintly shown, and for only a short distance behind head, where its course is obliquely downward; the usual series of mucous pores present, but not conspicuous. In spirits this species has an olive-brown ground color, becoming white on underside of head and on abdomen; a series of 8 or 9 brown spots  $\frac{1}{2}$  as large as eye, along middle of sides, those posteriorly continued downward onto base of anal, the last 2 or 3 reaching edge of fin and there developing into intense black blotches; a similar series of smaller spots corresponding in position to those just described occurs along the base of dorsal, these continued as faint bars on the fin, at the margin of which they develop into a black blotch, those posteriorly wider and more intense; an intermediate series of spots alternating with the 2 just described; an elliptical jet-black spot occupies the greater part of caudal fin, and is narrowly margined all around with white; peritoneum jet-black; the mouth and gill cavities white. Coast of Alaska, in rather deep water; known from 12 specimens, the longest  $6\frac{1}{2}$  inches; depth 204 fathoms. (*barbatus*, bearded.)

*Lycconema barbatus*, GILBERT, Rept. U. S. Fish Comm. 1893 (1896), 471, coast of Alaska, at Albatross Station 3129, lat.  $36^{\circ} 39' 40''$  N., long.  $122^{\circ} 01' W.$ , in 204 fathoms.

#### 941. BOTHROCARA, Bean.

*Bothrocara*, BEAN, Proc. U. S. Nat. Mus. 1890, 38 (*mollis*).

Body elongate, compressed, semitranslucent, covered with small scales; small teeth in jaws and on vomer and palatines; mucous pores about head largely developed. No ventral fins; dorsal and anal joined to caudal. Deep-sea fishes, allied to *Lycodes*, but lacking ventrals. The species have been referred to the Antarctic genus *Maynea*, CUNNINGHAM. From the latter, however, *Bothrocara mollis* seems to be distinct, differing in the larger mouth, more cavernous head, and lower dorsal. In some regards *B. pusilla* is intermediate, and it may belong to *Maynea*. ( $\beta\theta\theta\rho\sigma$ , cavity;  $\kappa\acute{\alpha}\rho\alpha$ , head.)

- a. Body elongate, with the head short, 6 in length; depth 9; mouth small; mucous cavities small; color light brown, the dorsal dark-edged. PUSILLA, 2838.  
 aa. Body deeper and more compressed; the large head  $4\frac{1}{2}$  in length ( $5\frac{1}{2}$  in young); depth  $6\frac{1}{2}$ ; mouth large; mucous cavities large; color uniform brown, the vertical fins dark-edged. Size large. MOLLIS, 2839.

2838. *BOTHROCARA PUSILLA* (Bean).

Head 6 in the total length; depth 9. D. 95, including  $\frac{1}{2}$  caudal; A. 81, including  $\frac{1}{2}$  of caudal; P. 17; eye 3 in head; snout 4. Body elongate, little compressed; head short; mouth small; maxillary extending to below front of pupil; gill clefts narrow, the anterior end below margin of preopercle; width of isthmus rather less than  $\frac{1}{2}$  of orbit; the low dorsal beginning nearly over axil of pectoral; pectoral nearly  $\frac{3}{4}$  as long as head; vent as far from end of head as dorsal origin from tip of snout. Color light brown; dorsal and anal with a narrow dark margin. Size small; length  $6\frac{1}{2}$  inches. Eastern parts of Bering Sea, and about the Alaskan Peninsula, in rather deep water. Besides the original types Dr. Gilbert records a few specimens from north of Unalaska, at depths of 121 to 351 fathoms. (*pusillus*, weak.)

*Maynea pusilla*, BEAN, Proc. U. S. Nat. Mus. 1890, 39, off Nagai Island, lat.  $55^{\circ} 10' N.$ , lon.  $160^{\circ} 18' W.$ , in 110 fathoms (Type, No. 45360. Coll. *Albatross*); GILBERT, Rept. U. S. Fish Comm. 1898 (1895), 455.

2839. *BOTHROCARA MOLLIS*, Bean.

Head  $4\frac{1}{2}$  in total length in adult,  $5\frac{1}{2}$  to  $5\frac{3}{4}$  in young; depth  $6\frac{3}{4}$  in adult, 10 in young. D. 100 to 105 to middle of caudal; A. 89 to 95 to middle of caudal; eye 4 in head in adult,  $3\frac{1}{2}$  in young. Body covered with embedded scales, which extend on dorsal and anal fins. Head naked, breast and nape scaly; snout blunt, the lower jaw included; maxillary reaching middle of pupil,  $2\frac{1}{2}$  in head; large mucous cavities conspicuous along mandible, suborbital ring, and top of head; vomerine and palatine teeth present, the latter in a narrow band, obscure in the young. Pectoral 2 in head; origin of dorsal slightly behind base of pectoral, its distance from tip of snout 4 in total length; origin of anal under seventeenth dorsal ray; longest dorsal ray 5 in head; longest anal ray  $8\frac{1}{2}$  in head. Gill openings wider than in *Bothrocara pusilla*, the anterior end of the cleft under posterior margin of eye, the width of the isthmus less than  $\frac{1}{2}$  diameter of pupil. Color uniform brown, fins lighter; dorsal and anal margined with black, more prominent posteriorly where it covers the entire fins. North Pacific. Adult examples from southern California, 18 inches long, were described as *Maynea brunnea*, while a young individual,  $5\frac{1}{2}$  inches long, from Queen Charlotte Islands, with the vomerine and palatine teeth not evident, was made the type of a distinct genus, as *Bothrocara mollis*. The two are identical and apparently belong to the same genus as *B. pusilla*. Similar specimens, 1 adult and 2 young, were dredged by us (*Albatross*) off Bogoslof Island in 664 fathoms. Dr. Gilbert records also specimens from near Unalaska, depth 316 fathoms. The teeth on the palatines are in a single series instead of a wide band, as stated in the original description. (*mollis*, soft.)

*Bothrocara mollis*, BEAN, Proc. U. S. Nat. Mus. 1890, 38, off Queen Charlotte Islands, in 876 fathoms (Type, No. 45359. Coll. *Albatross*); JORDAN & GILBERT, Rept. Fur Seal Invest., 1898.

*Maynea brunnea*, BEAN, Proc. U. S. Nat. Mus. 1890, 39, lat.  $33^{\circ} 8' N.$ , lon.  $118^{\circ} 40' W.$ , off San Clemente Island, southern California, in 414 fathoms. (Coll. *Albatross*.)

942. GYMNELIS, Reinhardt.

*Gymnelis*, REINHARDT, Dansk. Vidensk. Selsk. Afhandl., VII, 131, 1838 (*viride*).  
*Cepolophis*, KAUP, in Archiv für Naturgesch. 1856, 96 (*viridis*).

Body elongate, naked. Vertical fins without spines; ventral fins none. Small conical teeth on the jaws, vomer, and palatines. Gill openings very narrow. No air bladder; pyloric caeca none; no anal papilla. Size small. Cold seas. Two or 3 species known: *G. pictus*, from the Antarctic, and *G. viridis*, which ranges widely in Arctic waters, and with which the very dubious *G. stigma* is probably identical. (*γυμνός*, naked; *ἔρχελυς*, eel.)

a. Dorsal fin inserted close behind pectoral, its distance from it much less than diameter of eye; no ocellus on dorsal fin. VIRIDIS, 2840.

aa. Dorsal fin inserted an eye's diameter behind pectoral; a large black spot, ocellated with white, on dorsal fin above vent; other ocelli sometimes present.

STIGMA, 2841.

2840. GYMNELIS VIRIDIS (Fabricius).

Head about  $6\frac{1}{2}$ ; depth about 13; eye 7 in head. D. 100; A. 80. Snout subconical, longer than the eye; jaws equal; mouth oblique; maxillary reaching beyond eye; teeth rather small, conical, in a single series on each side, forming a patch anteriorly; distance from snout to vent  $2\frac{1}{2}$  times length of head. Pectoral rounded, inserted low, its length less than  $\frac{1}{4}$  that of head. Dorsal fin inserted close behind pectoral, its distance from it much less than diameter of eye. Body pale, with faint dark cross shades; dorsal clouded but without black spot; anal dusky. Arctic seas, Alaska to Greenland and Nova Scotia; abundant in the Arctic waters south to Unalaska and Bristol Bay, where specimens were taken in shallow water; our specimens from Bristol Bay.

Ensign H. G. Dresel records 1 small specimen (No. 28636, U. S. Nat. Mus.), badly preserved, obtained by Mr. Newton Pratt Scudder in Davis Straits, July, 1879. Length 100 mm. D. ca. 97; A. ca. 80. In this specimen the maxillary does not extend to the posterior margin of the eye, which is comparatively very large. Its diameter is longer than distance from tip of snout to orbit, and is contained 4 times in head. Head 7 in total length; depth 12. Pectoral 2 in head. (*viridis*, green.)

*Ophidium viride*, FABRICIUS, Faun. Græn., 141, 1780, Greenland.

*Ophidium unernak*, LACEPÈDE, Hist. Nat. Poiss., II, 280, 1800, Greenland; after FABRICIUS.

*Gymnelis viridis*, RICHARDSON, Last Arctic Voyage, 321, pl. 29, 1854.

*Gymnelis viridis*, REINHARDT, Dansk. Vidensk. Selsk. Afh., VII, 1838, 131; GÜNTHER, Cat.,

IV, 323, 1862; KRÖYER, Poissons du Nord, Voy. en Scand. et Lap., pl. 15, a-f; COLLETT,

Norske Nordh. Exped., Flske, 123, pl. 4, fig. 32, 1880; JORDAN & GILBERT, Synopsis,

789; GILBERT, Rept. U. S. Fish Comm. 1893 (1896), 455.

? *Gymnelis pictus*, GÜNTHER, Cat., IV, 324, 1862, no locality.

2841. GYMNELIS STIGMA (Lay & Bennett).

Head 6; depth 11. D. 90; A. 70. Form, size, and general appearance of *G. viridis*, the dorsal inserted farther back, an eye's diameter behind pectoral. A large, round black ocellus, ringed with white, on dorsal fin above vent; 2 or 3 other ocelli sometimes present; head and nape with small white spots; body with faint dark shades and bands. Otherwise

as in *G. viridis*, from which it may not be distinct; but the above characters appear in our specimens (from near the Pribilof Islands) and in Richardson's figure of *G. unimaculatus*. The white spots on the head were mistaken for "very small scales" in the original description of *G. stigma* from Dr. Collie's notes. This description is, in substance, as follows: No trace of ventral fins; dorsal, caudal, and anal fins united into a transparent ridge; rays of branchial covering distinct; scales very small. Color dilute brown, with void swathes and spots; a purplish spot near beginning of dorsal fin. Snout obtuse; chin with a large gibbosity; teeth small. Length about 5 inches. (Lay & Bennett.) Arctic regions, Greenland to Bering Sea, with the preceding, and apparently equally common. (*στρυμα*, spot.)

*Ophidium stigma*, LAY & BENNETT, Zool. Beechey's Voy., 67, pl. 20, fig. 1, 1839, Kotzebue Sound. (Coll. Dr. Collie.)

*Gymnelis viridis* var. *unimaculatus*, RICHARDSON, Last Arctic Voyage, 367, 1854, Northumberland Sound. (Coll. Edward Belcher.)

*Gymnelis stigma*, GÜNTHER, Cat., IV, 325, 1862; JORDAN & GILBERT, Synopsis, 789, 1883.

#### 943. LYCOCARA, Gill.

*Uronectes*, GÜNTHER, Cat., IV, 325, 1862 (*parrii*); name preoccupied in Crustacea.

*Lycocara*, GILL, Proc. Ac. Nat. Sci. Phila. 1884, 180 (*parrii*).

Body ensiform, compressed; tail long and tapering; ventrals none; vent not far distant from the head; numerous minute teeth in jaws and on palate; lower jaw the longer; no barbel; scales and gill openings unknown. One species, very imperfectly known, no specimens having been obtained by any recent collector. (*λύκος*, wolf; *κάρρα*, head.)

#### 2842. LYCOCARA PARRII (Ross).

Head 4. D. 50; A. 45; P. 37. Head very obtuse, its length, depth, and breadth equal; head broader than the body, flattened and grooved between the eyes, which are lateral and rather large; lower jaw the longer; jaws and palate with minute teeth; greatest depth of body somewhat more than length of head; neck much arched. Dorsal inserted just behind head; pectoral extending beyond vent. Vent not far distant from head. Color uniform. Baffins Bay. (Günther.) (Named for Capt. William Edward Parry, the Arctic explorer.)

*Ophidium parrii*, ROSS, Parry's Third Voyage, App., 109, 1826, Baffins Bay.

*Uronectes parrii*, GÜNTHER, Cat., IV, 328, 1862; JORDAN & GILBERT, Synopsis, 789, 1883.

#### 944. MELANOSTIGMA, Günther.

*Melanostigma*, GÜNTHER, Proc. Zool. Soc. Lond. 1881, 21 (*gelatinosum*).

This genus is distinguished from *Bohrocara* by the much more elongate teeth, which in the jaws, as well as on the vomer and palatines, stand in single series. Gill openings much smaller than in related forms, reduced to a small foramen above the base of the pectoral. Skin loose and movable, as in *Liparis*, enveloping the vertical fins; pectorals very small;

ventrals none. Body tapering very rapidly backward; the tail very slender. Deep-sea fishes, of soft substance, allied to *Lothrocara*, but with stronger teeth. (*μέλας*, black; *στίγμα*, spot.)

a. Maxillary reaching beyond front of pupil; color purplish gray, becoming black on the tail. GELATINOSUM, 2843.

aa. Maxillary not reaching beyond vertical from front of pupil; color uniform deep black. PAMMELAS, 2844.

2843. MELANOSTIGMA GELATINOSUM, Günther.

Body enveloped in a loose, delicate skin, as in *Iiparis*. Head large, deep, compressed, with obtuse snout. Eye large,  $3\frac{1}{2}$  in head, and longer than snout. Cleft of mouth rather oblique, but lower jaw not projecting beyond upper; lips not fleshy; gill opening reduced to a very narrow foramen above base of pectoral fin; origin of dorsal fin and root of pectoral fin enveloped in loose skin of body; dorsal fin probably commencing above middle of pectoral, low at first, but becoming considerably higher posteriorly; pectorals very narrow, consisting of a few rays only. Upper parts tinged with a purplish-gray; sides mottled with same color, which toward end of tail becomes more intense, almost black; inside of mouth, gill openings, and vent black. Total length of the type specimen  $5\frac{1}{2}$  inches; distance of the snout from the gill opening  $\frac{3}{4}$  inches, from the vent  $1\frac{1}{4}$  inches. (Günther.) Deep waters of the western Atlantic; originally known from the Straits of Magellan, but since obtained at various localities from Cape Cod to West Indies, in 500 to 1,000 fathoms. The identity of these specimens with the original types from South America may be questionable. (*gelatinosus*, jelly-like.)

*Melanostigma gelatinosum*, GÜNTHER, Proc. Zool. Soc. London 1881, 21, Tilly Bay, Straits of Magellan, in 24 fathoms (Coll. H. M. S. *Alert*, Dr. Coppinger); GÜNTHER, Challenger Report, XXII, 82, 1887; GOODE & BEAN, Bull. Mus. Comp. Zool., x, No. 5, 209, 1893; JORDAN, Cat. Fish. N. A., 125, 1885; GOODE & BEAN, Oceanic Ichthyology, 314, 1896.

2844. MELANOSTIGMA PAMMELAS, Gilbert.

Head 8 in total length; depth  $12\frac{1}{4}$ ; pectoral narrow, its length  $2\frac{3}{4}$  in head; eye large,  $3\frac{1}{2}$  in head; snout short and broad, 7 in head. Well distinguished from *M. gelatinosum* by the wider, blunter head, the smaller, less oblique mouth, the uniform black coloration, and the arrangement of the teeth in the jaws in 2 series. As in *M. gelatinosum*, the head and body are enveloped in a loose, thin skin, which is thrown into folds in alcoholic specimens, and entirely conceals anterior portion of dorsal and anal fins. On dissection the dorsal is seen to have its origin close behind the head, at a point over middle of pectoral fin; anal beginning immediately behind vent, the rays of both fins enveloped in a gelatinous, subcutaneous tissue. Head broad, with its greatest width equaling its greatest depth; mouth broad, somewhat oblique, with equal jaws, the maxillary reaching vertical from front of pupil; each jaw with teeth in two distinct series in front, in a single series laterally in lower jaw, the outer teeth in front enlarged, almost canine-like. Gill opening a small pore above base of pectoral, its diameter about  $\frac{1}{4}$  that of eye. Color intense black on head and abdomen, brownish black elsewhere. Length

of type  $4\frac{1}{2}$  inches. Coast of southern Alaska. Three other specimens are at hand from *Albatross Station* 3126 (lat.  $36^{\circ} 49' 20''$  N., long.  $122^{\circ} 12' 30''$  W.; depth 456 fathoms). In the smallest,  $2\frac{1}{2}$  inches long, the head and abdomen are jet-black, but the rest of the body is only slightly dusky. (Gilbert.) ( $\pi\acute{\alpha}\varsigma$ , all;  $\mu\acute{\epsilon}\lambda\alpha\varsigma$ , black.)

*Melanostigma pammelae*, GILBERT, Rept. U. S. Fish Comm. 1893 (1896), 472, pl. 35, coast of southern Alaska, at *Albatross Station* 3202, lat.  $36^{\circ} 46' 10''$  N., long.  $121^{\circ} 58' 45''$  W., in 322 fathoms.

#### Family CCVII. DEREPODICHTHYIDÆ.

Deep-sea fishes of slender body, scaleless, and without lateral line, somewhat resembling the *Zoarcidæ*, but with each ventral fin reduced to a slender, unbranched filament, the two very closely approximate, and springing from a common projecting base located far forward, below the eye. Gill opening a narrow, vertical slit. Character otherwise given below. A single species known; apparently intermediate between the *Zoarcidæ* and the *Ophidiidæ*.

#### 945. DEREPODICHTHYS, Gilbert.

*Derepodichthys*, GILBERT, Rept. U. S. Fish Comm. 1893 (1896), 456 (*alepidotus*).

Body slender; no scales; no lateral line; ventral fins reduced each to a slender, unbranched filament, the two very closely approximate, and springing from a common projecting base, which is located far forward below the eye, as *Ophidion*. Gill opening a narrow, vertical slit, little wider than base of pectorals. Teeth cardiform, curved, few in number, in narrow bands or irregular single series on jaws, vomer, and palatines. ( $\delta\acute{\epsilon}\rho\eta$ , throat;  $\pi\acute{o}\upsilon\varsigma$ , foot;  $\iota\chi\theta\acute{\upsilon}\varsigma$ , fish.)

#### 2845. DEREPODICHTHYS ALEPIDOTUS, Gilbert.

Head  $8\frac{1}{2}$  in total length; depth of head and body  $2\frac{1}{2}$  in head; width of head  $2\frac{3}{4}$  in head; distance from tip of snout to base of ventrals  $2\frac{1}{2}$  in length of head. Distance from tip of snout to front of dorsal  $5\frac{1}{4}$  in total length, from tip of snout to vent  $3\frac{3}{4}$  in total. Head and body very long and slender, the former resembling a *Lycodes* in appearance, being moderately compressed, with a flattish occiput and a gentle rounded decurved rostral profile. Mouth slightly oblique, quite at lower side of snout; the lower jaw shorter, fitting within the upper; maxillary and premaxillary entirely concealed within the thick skin of the upper lip, which is directly continuous with that of the forehead, the upper jaw being therefore nonprotractile; angle of mouth under front of pupil, its distance from tip of snout  $2\frac{3}{4}$  in head. Teeth cardiform, curved, few in number, in narrow bands or irregular single series on jaws, vomer, and palatines. Eye small, not filling the elongate orbit, the diameter of exposed portion of eyeball slightly less than  $\frac{1}{2}$  length of snout, the latter  $3\frac{1}{2}$  in head. A series of large mucous pores on snout and lower part of cheeks; a second series on mandible; no pores on body. Gill slit vertical, not continued forward, its lower end slightly above base of lower pectoral rays; length of slit  $\frac{1}{2}$  length of



head, slightly less than distance between slits. Pectorals long and slender, reaching halfway to vent,  $1\frac{1}{2}$  in head; dorsal and anal confluent with the caudal, concealed in the thick integument, so that the rays can not be counted. Color in spirits, light brownish, the dorsal and pectorals whitish, the anal with a dark margin which becomes black posteriorly; lips dusky; abdominal region blue black. Coast of British Columbia. A single specimen,  $4\frac{1}{2}$  inches long, dredged off Queen Charlotte Island. (Gilbert.) (*ἀλεπίδωρος*, scaleless.)

*Derepodichthys alepidotus*, GILBERT, Rept. U. S. Fish Comm. 1893 (1896), 456, Queen Charlotte Island, at Albatross Station 3342, in 1,588 fathoms.

Family CCVIII. OPHIDIIDÆ.

(THE CUSK EELS.)

Body elongate, compressed, more or less eel-shaped, usually covered with very small scales, which are not imbricated, but placed in oblique series at right angles with each other; head large, lower jaw included; both jaws, and usually vomer and palatines also, with villiform or cardiform teeth; premaxillaries protractile; gill openings very wide, the gill membranes separate, anteriorly narrowly joined to the isthmus behind the ventrals; pseudobranchiæ small. Gills 4, a slit behind the fourth; vent more or less posterior. Vertical fins low, without spines, confluent around the tail; tail isocercal; ventral fins at the throat, each developed as a long, forked barbel. Air bladder and pyloric cæca present. To this Dr. Gill adds the following characters, shared more or less by related families: "Orbito-rostral portion of cranium contracted and shorter than the posterior, the cranial cavity closed in part by the expansion and junction of the parasphenoid and frontals, the supraoccipital horizontal and cariniform posteriorly, the exoccipitals expanded backward and upward behind the supraoccipital, the exoccipital condyles contiguous, and with the hypercoracoid (scapula, Parker) fenestrate (or foraminated) about its center, and the hypercoracoid with its inferior process divergent from the proscapula." Genera 7, species about 25. Carnivorous fishes; found in most warm seas, some of them descending to considerable depths, the group especially well represented in tropical America. (*Ophidiida*, group *Ophidiina*, Günther, Cat., IV, 376-380, 1862.)

- a. Head scaly, at least above; body covered with scales imbricated in quincunx; snout usually with a spine at tip; opercle with or without spinous tip; air bladder, so far as known, ovate, without posterior foramen. LEPOPHIDIUM, 946.
- aa. Head scaleless; scales of body rudimentary, scarcely embedded.
  - b. Air bladder oblong-ovate, not contracted behind, and without posterior foramen.
  - c. Opercle ending in a flat point behind, without spine. OPHIDIUM, 947.
  - cc. Opercle ending behind in a strong spine concealed in the skin. CHILARA, 948.
- bb. Air bladder short, thick, reniform or orbicular, with a large foramen behind.
  - d. Opercle ending in a flat point, without spine. RISSOLA, 949.
  - dd. Opercle ending behind in a spine concealed in the skin. OTOPHIDIUM, 950.

946. *LEPOPHIDIUM*, Gill.

*Leptophidium*, GILL, Proc. Ac. Nat. Sci. Phila. 1863, 210 (*profundorum*); name preoccupied in Serpents by *Leptophidium*, HALLOWELL, 1860.

*Lepophidium*, GILL, Amer. Nat., Feb., 1895, 16 (*profundorum*).

Body much elongate, moderately compressed, with back and abdominal regions arched, more compressed and slowly decreasing in height backward to an abruptly rounded point; scales regularly imbricated in quincunx oval, and with striæ radiating backward; head with imbricated scales, extending to forehead; snout high, projecting forward, and obtusely rounded, armed above with a short, nearly concealed spine directed forward and somewhat downward, obsolete in 1 species; mouth moderate, oblique; teeth of jaws villiform, immersed in a mucous membrane, separated by an interval from the longer ones in the outer row, which are pointed and usually movable; vomer and palatines with teeth. Deep waters of America on both coasts. Perhaps a fuller knowledge of the species of this genus will lead to its subdivision. (*λέπος*, scale; *Ophidium*, from the squamation.)

a. Snout without decurved hook or spine; gill rakers 8; head 5 in length; depth 7; pectorals 10; body marbled, the vertical fins edged with black.

MARMORATUM, 2846.

aa. Snout with a decurved hook or spine at tip, sometimes more or less concealed in the skin.

b. Gill rakers 7 to 9 in number.

c. Head large,  $3\frac{3}{4}$  to  $4\frac{1}{2}$  in length.

d. Body stoutish, the depth 6 in length; scales 125; no black blotch on front of dorsal.

EMMELAS, 2847.

dd. Body slender, the depth  $9\frac{1}{2}$  in length; dorsal with a black blotch in front; scales 180.

STIGMATISTIUM, 2848.

cc. Head moderate, 6 in length; depth 10; vertical fins black-edged.

e. Anterior teeth in jaws movable; pectoral 11 in body; body without white spots.

PROFUNDORUM, 2849.

ee. Anterior teeth in jaws not movable; pectoral 13 to 14 in body; body with whitish spots.

CERVINUM, 2850.

bb. Gill rakers 4 in number.

f. Scales moderate, 175 to 200 in lateral line.

g. Body without dark cross bars; dorsal and anal margined with black; air bladder oblong.

h. Head  $4\frac{1}{2}$  in length; depth 8; pectoral 10 $\frac{1}{2}$ . Pacific species.

PRORATES, 2851.

hh. Atlantic species imperfectly described.

BREVI BARBE, 2852.

gg. Body with dark cross bars; dorsal spotted with black; anal wholly black; head  $5\frac{1}{2}$  in length; depth  $8\frac{1}{2}$ .

PARDALE, 2853.

ff. Scales minute, about 250 in lateral line; head  $4\frac{3}{4}$ ; depth  $7\frac{1}{2}$  to 8; color nearly plain, the fins dark edged.

MICROLEPIS, 2854.

2846. *LEPOPHIDIUM MARMORATUM* (Goode & Bean).

Head 5; depth  $7\frac{1}{2}$ ; eye 4 in head; snout about 5. Body somewhat elongate, stontish anteriorly, gradually tapering; head thickish; inter-orbital area broad, convex, its width nearly equal to length of snout, which is blunt, spineless; eye circular, somewhat exceeding length of snout. Maxillary extending to vertical through posterior margin of orbit,

the mandible far beyond, its length equal to that of postorbital portion of head. Teeth on vomer and in jaws in villiform bands, the outer series in the latter slightly enlarged. Pseudobranchiæ present; gill rakers short, 8 below angle of first arch, the longest less than  $\frac{1}{2}$  diameter of eye. Branchiostegals 7. Ventrals as long as postorbital part of head. Dorsal origin at distance from snout contained  $4\frac{1}{2}$  in total length, with 28 rays in a space equal to length of head, counting from the origin of the fin; anal origin separated from snout by distance  $2\frac{3}{4}$  in total length; length of pectoral 2 in head, or 10 in total. Scales closely imbricated, ornamented with delicate concentric striæ; lateral line apparently complete, located about  $\frac{1}{2}$  distance from dorsal to ventral outline. Color yellowish gray, marbled along the upper half of head and body with olive brown; dorsal and anal fins with black margins. Gulf Stream, in 213 fathoms. (Goode & Bean.) (*marmoratus*, marbled.)

*Leptophidium marmoratum*, GOODE & BEAN, Proc. U. S. Nat. Mus. 1885, 423, lat.  $23^{\circ} 10' 39''$  N., long.  $82^{\circ} 20' 21''$  W., in 213 fathoms (Type, No. 37237, U. S. Nat. Mus. Coll. Albatross); GOODE & BEAN, Oceanic Ichthyology, 348, 1896.

2847. *LEPOPHIDIUM EMMELAS* (Gilbert).

Head  $3\frac{1}{2}$  to 4; depth  $5\frac{1}{2}$  to 6; eye  $4\frac{1}{2}$  in head; snout  $4\frac{1}{2}$ ; interorbital width 7; vertebrae  $13 + 44 = 57$ ; maxillary  $2\frac{3}{4}$  to  $2\frac{1}{2}$  in head; ventral filament  $2\frac{3}{4}$ ; pectoral 2 in head; scales 8-125-18 or 20 before dorsal. Body deep, compressed. Maxillary reaching slightly beyond orbit. Jaws slender and weak, the teeth in very narrow bands, the outer not enlarged. Rostral ridge very sharp, bearing a flat spine at its base directed upward and backward, terminating in a very slender sharp spine anteriorly. Opercle ending in a weak spinous point behind. Gill rakers short and slender, the longest  $\frac{1}{2}$  pupil, 8 or 9 movable ones developed. Skull and all bones of head very thin and papery. Dorsal beginning over base of pectorals, the nape midway between its origin and middle of orbit. Distance from snout to origin of anal  $1\frac{1}{2}$  in distance from latter to end of tail. Scales large, covering cheeks, opercles and top of head forward to middle of interorbital space. Color brownish, much dusted with minute specks; fins blackish, the vertical fins with an indistinct narrow whitish margin; inside of mouth dusky; the roof of mouth, lining of gill cavity, and peritoneum jet-black. Coast of Lower California. Many specimens, the longest 9 inches, from Albatross Stations 3007 and 3008, in 362 and 306 fathoms. (Gilbert.) (*έρ*, within; *μέλας*, black.)

*Leptophidium emmelas*, GILBERT, Proc. U. S. Nat. Mus. 1890, 110, coast of Lower California. (Coll. Albatross.)

2848. *LEPOPHIDIUM STIGMATISTIUM* (Gilbert).

Head  $4\frac{1}{2}$  in length; depth  $9\frac{1}{2}$ ; eye  $4\frac{1}{2}$  in head; snout 6; interorbital width  $6\frac{1}{2}$ ; maxillary reaching slightly beyond posterior border of eyes,  $2\frac{1}{2}$  in head. A strong rostral spine. Outer teeth scarcely enlarged, evidently so only in front of upper jaw. Gill rakers long and slender, strongly curved forward at tip, the longest equaling  $\frac{1}{2}$  eye; 7 well-developed gill-rakers present. Opercle ending in a rounded process, a broad soft flap

projecting beyond it. Dorsal inserted behind middle of pectorals, the nape equidistant from front of dorsal and base of rostral spine; pectorals  $2\frac{2}{3}$  in head; longest ventral filament  $2\frac{1}{4}$  in head; scales small, about as in *L. prorates*, 180 transverse series, 28 in front of dorsal, continued forward on top of head to front of pupil; cheeks and opercles scaly. Color dusky olivaceous, lighter below; dorsal with a large black blotch on anterior rays, the margin obscurely dusky; anal broadly margined with jet-black; caudal with median rays black at base, the outer rays and the margin light; lining of gill cavity jet-black; inside of mouth white; peritoneum bright silvery. A single specimen 10 inches long. Coast of Lower California. (Gilbert.) Much resembling *L. prorates*, differing in dentition, in gill rakers, and in color. (*στρυμα*, brand; *ιστιορ*, sail.)

*Leptophidium stigmatistium*, GILBERT, Proc. U. S. Nat. Mus. 1890, 109, off Lower California, at Albatross Station 2996, in 112 fathoms.

#### 2849. LEPOPHIDIUM PROFUNDORUM (Gill).

Head 6; depth 10. Body very slender; scales regularly arranged in quincunx order, those on head extending to forehead, opercles, and cheeks; snout high, projecting, armed with a concealed spinous hook; teeth villiform, separated by an interval from an outer row of longer, slender, movable teeth; eye longer than snout,  $3\frac{1}{4}$  in head; lateral line obsolete behind; vent toward end of first third of length; ventral fins short; gill rakers 8. Light rufous; vertical fins margined with black. Gulf Stream, off the coast of Florida. (Gill.) One specimen known. (*profundorum*, of the depths.)

*Leptophidium profundorum*, GILL, Proc. Ac. Nat. Sci. Phila. 1863, 211, Gulf Stream, off the Coast of Florida (Coll. Commodore Rodgers); GOODE & BEAN, Oceanic Ichthyology, 347, 1896.

*Ophidium profundorum*, JORDAN & GILBERT, Synopsis, 793, 1883.

#### 2850. LEPOPHIDIUM CERVINUM (Goode & Bean).

Head about  $6\frac{1}{4}$ ; depth about  $10\frac{1}{4}$ ; eye 4 in head; ventrals 3 in head. Body elongate, slender; head slender, somewhat compressed; interorbital area broad, convex, its width equal to length of snout, and  $5\frac{2}{3}$  in head; snout sharp, conical, armed with a short but sharp spine, and somewhat overhanging mouth; eye much exceeding length of snout; maxillary extending nearly to vertical through posterior margin of orbit,  $2\frac{2}{3}$  in head; mandible extending behind same vertical, its length equal to that of head without postorbital portion. Jaws, vomer, and palatines with narrow bands of villiform teeth, some of which are noticeably enlarged (rot movable). Pseudobranchiae present. Gill rakers short, 8 below angle of first arch, 4 of which are rudimentary, the longest 5 in diameter of eye. (In *L. profundorum* the gill rakers are slenderer and longer, though about equally numerous on the first arch.) Scales in about 11 rows from the origin of the dorsal to the median line of the body. Dorsal origin far back, at a distance from the snout  $4\frac{1}{4}$  in total length; at a distance from the eye equal to the head's length. (In *L. profundorum* this distance is  $\frac{2}{3}$  of the head's length and the first ray of the dorsal is nearly over the

middle of the extended pectoral; in *L. cervinum*, over its tip, or nearly so.) Distance of anal origin with snout 3 in total length. Length of pectoral 2 in head's length and 13 to 14 in that of body (10 in *L. marmoratum*, 11 in *L. profundorum*). Scales ornamented with radiating striae, densely covering all parts of the fish except snout, under surface of head, and the fins; lateral line continued almost to end of tail. Color brownish yellow, with numerous subcircular spots of white, with diameter  $\frac{1}{2}$  that of eye, along the upper half of body; vertical fins with narrow black margin. Gulf Stream. (Goode & Bean.) A specimen from off Sand Key Light, Florida, recorded by Mr. Garman. (*cervinus*, deer-like, from the faun-color.)

*Leptophidium cervinum*, GOODE & BEAN, Proc. U. S. Nat. Mus. 1385, 422, lat.  $40^{\circ}$  1' N., long.  $69^{\circ}$  56' W., depth 76 fathoms (Type, No. 28764. Coll. *Fish Hawk*); GOODE & BEAN, Oceanic Ichthyology, 346, 1896.

*Leptophidium cervinum*, GARMAN, Bull. Iowa Lab. Nat. His. 1896, 91.

2851. *LEPOPHIDIUM PROBATES* (Jordan & Bollman).

Head  $4\frac{1}{2}$  to  $4\frac{3}{8}$  ( $4\frac{3}{8}$  to  $4\frac{1}{2}$  in total); depth  $7\frac{1}{8}$  to  $8\frac{1}{8}$  ( $7\frac{1}{8}$  to  $8\frac{1}{8}$ ); eye  $4\frac{1}{2}$  in head; snout 5; maxillary  $2\frac{1}{2}$ ; interorbital  $1\frac{1}{8}$  in eye; pectoral  $2\frac{1}{2}$  in head; inner ventral filament shortest, the longer  $2\frac{3}{8}$  in head. Body moderately elongate, compressed, considerably stouter than in *L. profundorum*. Mouth large, maxillary reaching about  $\frac{1}{2}$  pupil's length beyond posterior border of eye. Outer teeth slightly enlarged, a little movable, those of upper jaw largest. Gill rakers rather long and slender,  $\frac{1}{2}$  length of eye, 4 developed. Tip of snout with a strong spine directed forward and slightly downward; opercle without spine, ending in a flat projection covered by skin. Dorsal beginning over middle of pectorals, longest ray 4 in head. Scales regularly imbricated, but very small, about 225 in a longitudinal series; scales on top of head extending forward to base of ethmoidal spine; sides of head covered with small scales; lateral line not reaching end of tail. Air bladder oblong-lanceolate. Color olivaceous, paler below; scales rather profusely dotted with black; a pale shade across opercles; lower jaw, gular region, and anterior branchiostegals dusted; dorsal and anal margined with black, the band on anal the broader; pectorals pale. Specimens of this species were obtained at Panama and at *Albatross Station* 2801, south of Panama. Length of type 10 inches. (*πρωπαρτης*, prow-bearing, from the rostral spine.)

*Leptophidium prorates*, JORDAN & BOLLMAN, Proc. U. S. Nat. Mus. 1889, 172, Panama. (Type, No 41149, U. S. Nat. Mus. Coll. *Albatross*.)

2852. *LEPOPHIDIUM BREVIBARBE* (Cuvier).

A short decurved spine at tip of snout; teeth strong; occiput and opercles scaly. Vertical fins edged with black. (Kaup.) Air bladder oblong ovate, without contracted portion and without posterior foramen; no single anterior bone replaced by cartilage. (Müller.) West Indies and Brazil; a scarcely known species; apparently close to *L. prorates*, but very insufficiently described. (*brevis*, short; *barba*, beard.)

*Ophidion brevibarbe*, CUVIER, Règne Animal, Ed. 2, vol. II, 358, 1829, Brazil; MÜLLER, Abhandl. Berl. Acad. 1843, 153, pl. 4, f. 4; KAUP, Apodes, 154, pl. 16, f. 1; GÜNTHER, Cat. Fish., IV, 379, 1862.

2853. *LEPOPHIDIUM PARDALE* (Gilbert).

Head  $5\frac{1}{2}$  in length; depth  $8\frac{1}{2}$ ; eye  $3\frac{3}{8}$  in head; snout  $4\frac{3}{8}$ ; interorbital  $1\frac{1}{2}$  in eye. Body very slender, with a short head and small mouth; maxillary scarcely reaching vertical from posterior border of orbit, its length  $2\frac{1}{8}$  in head; outer teeth very little enlarged, not movable; teeth present on jaws, vomer, and palatines. Gill rakers slender, the longest  $\frac{1}{2}$  eye, 4 developed. Tip of snout with a strong, concealed spine, as in *L. prorates*. Opercle ending in a short spine. Nape midway between front of dorsal and front of pupil; dorsal beginning over middle of pectorals, which are  $2\frac{1}{2}$  in head; ventral filaments very short; the inner the longer,  $4\frac{3}{8}$  in head. Scales very small, about 200 in a longitudinal series, extending forward on top of head to middle of interorbital space; cheeks and opercles scaly. Light olive, a series of 8 black bars downward from back, scarcely reaching lateral line, sometimes continuous with their fellows of the other side, and alternating with smaller black spots on dorsal outline; below the smaller spots a series of round spots nearly as large as eye along middle line of sides; sides and lower parts of head and body dusted with rather coarse black specks; dorsal light, the margin with 10 elongate black blotches, usually longer than the interspaces; caudal dusky at base, its distal half white; anal wholly black; peritoneum and lining of gill cavity white. Lower California. A single specimen, length  $7\frac{1}{8}$  inches, from *Albatross* Station 3014, in 29 fathoms. (Gilbert.) (*πάρδαλις*, leopard.)

*Leptophidium pardale*, GILBERT, Proc. U. S. Nat. Mus. 1890, 108, off Lower California. (Type, No. 44382. Coll. Dr. Gilbert.)

2854. *LEPOPHIDIUM MICROLEPIS* (Gilbert).

Head  $4\frac{3}{8}$  in length; depth  $7\frac{1}{2}$  to 8; eye  $4\frac{1}{2}$  to 5 in head; snout 5; interorbital width  $6\frac{1}{2}$ ; maxillary extending beyond orbit,  $2\frac{1}{2}$  to  $2\frac{1}{2}$ . Rostral spine very strong, as in *L. prorates*. Outer teeth enlarged, not at all movable, those in upper jaw largest. Four gill rakers developed, the longest  $3\frac{1}{2}$  in eye. Opercle ending in a short concealed spinous point. Dorsal inserted in front of middle of pectorals, the distance from nape to front of dorsal usually less than from nape to middle of eye; longest ventral filament  $3\frac{1}{8}$  to  $3\frac{1}{2}$  in length of head; pectorals  $2\frac{1}{2}$  to  $2\frac{1}{2}$  in head. Scales exceedingly small, regularly imbricated, in about 250 transverse series, 35 transverse series between nape and dorsal (about 175 transverse rows in *L. prorates*, 25 series between nape and dorsal). Top of head scaly as far as front of eyes. Cheeks and opercles scaly. Color as in *L. prorates*, the lining of peritoneum and gill cavity silvery white, the former with little or no black specking. Closely related to *L. prorates*, differing principally in the much smaller scales. Gulf of California. Many specimens, the longest 14 inches, from *Albatross* Stations 3015 and 3016, in 145 and 76 fathoms. (Gilbert.) (*μικρός*, small; *λεπίς*, scale.)

*Leptophidium microlepis*, GILBERT, Proc. U. S. Nat. Mus. 1890, 109, Gulf of California. (Coll. Dr. Gilbert.)

947. OPHIDION (Artedi) Linnæus.

(Cusk Eels.)

*Ophidion* (ARTEDI) LINNÆUS, Syst. Nat., Ed. x, 259, 1758 (*barbatum*).

*Ophidium*, LINNÆUS, Syst. Nat., Ed. XII, 431, 1766, and of most recent authors; changed spelling.

Body moderately elongate, compressed; scales small, usually not imbricated, but arranged in short, oblique series, often placed at right angles with each other, much as in *Anguilla*. Head naked; teeth villiform, those of the outer series more or less enlarged, none of them movable; teeth on vomer and palatines bluntish, some of them enlarged. Vent well behind pectorals. Opercle without distinct spine; sometimes (*O. barbatum*) a distinct spine at tip of snout. Air bladder oblong-ovate, tapering behind, without foramen. Shore species mostly European. (*Ophidium*, an ancient name, from *ὄφιδιον*, a small snake.)

a. Gill rakers 4.

b. Head  $4\frac{1}{2}$  in length; depth 7; fins not dark edged.

BEANI, 2855.

bb. Head 6 in length; inner ray of ventral  $1\frac{1}{2}$  in length of outer, which is shorter than head; fins dark-edged.

HOLBROOKI, 2856.

aa. Gill rakers 6 or 7; head  $5\frac{1}{2}$  to 6 in length; depth 8 to 10; color silvery, unspotted; fins not dark-edged; ventrals nearly as long as head.

GRAELLSI, 2857.

2855. OPHIDION BEANI, Jordan & Gilbert.

Head  $4\frac{1}{2}$  in length; depth about 7. Head small, the profile not very obtuse; snout  $4\frac{1}{2}$  in head; eye  $3\frac{1}{2}$ , more than twice the narrow interorbital space; mouth oblique, the maxillary reaching to posterior border of pupil, 2 in head; lower jaw slightly included; teeth small, in narrow bands in the jaws, the outer series in upper jaw somewhat enlarged; vomerine and palatine teeth small, subequal; head naked; snout spineless; opercle without spine; no evident pseudobranchiæ; gill rakers rather long and strong, 4 below angle of arch; occiput nearly midway between origin of dorsal and front of eye. Air bladder long and slender, occupying nearly the whole length of abdominal cavity, tapering backward. Very light olive, somewhat punctate above, slightly silvery below; fins without trace of dark edging (but being mutilated they may have been dark-edged in life). Gulf of Mexico. Two specimens, 1 of which is in good condition and about 4 inches long, were taken from the stomach of a red snapper, at Pensacola. (Named for Dr. Tarleton Hoffman Bean.)

*Ophidium graellsii*, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1882, 301; JORDAN & GILBERT, Synopsis, 963, 1883; not of POEY.

*Ophidion beani*, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1883, 43, Snapper Banks off Pensacola (Coll. Jordan & Stearns. Type, No. 30868, U. S. Nat. Mus.); JORDAN, Cat. Fishes N. A., 126, 1885.

2856. OPHIDION HOLBROOKI (Pinnam).

Head 6 in total length. Inner barbel nearly  $\frac{2}{3}$  length of the outer; outer barbel equal to distance from center of eye to point of operculum; maxillary reaching to posterior border of eye. Length of eye equal to distance from its posterior margin to ridge of preoperculum. Dorsal and anal

with a black margin. Gill rakers 4; air bladder long, pointed, with a foramen. (Putnam.) Length 6 inches. Gulf of Mexico; recorded from Key West, Florida; not seen by us. (Named for Dr. John Edwards Holbrook, the distinguished ichthyologist of Charleston.)

*Ophidion josephi*, GIRARD, U. S. and Mex. Bound. Surv., Ichth., 29, 1859, St. Joseph Island, Texas; JORDAN & GILBERT, Synopsis 793, 1883; quite as likely to be *Rissola marginata*.

*Ophidium holbrooki*, PUTNAM, Proc. Bost. Soc. Nat. Hist. 1874, 342, Key West, Florida; JORDAN & GILBERT, Synopsis, 793, 1883.

#### 2857. OPHIDION GRAELLSI, Poey.

Head  $5\frac{1}{2}$  in body; depth 10; eye 3 in head; pectoral  $2\frac{1}{2}$ ; ventral scarcely as long as head. Body elongate, compressed; mouth large, the maxillary reaching to posterior margin of pupil; small teeth on jaws, vomer, and palatines; eye very large, greater than length of snout; interorbital space  $\frac{3}{4}$  of eye, a sharp ridge along its middle to tip of snout, where it ends in a sharp spine; opercles unarmed; about 6 gill rakers developed on lower part of gill arch, apparently none above; pseudobranchie small, if present. Air bladder, injured in specimen examined, apparently lanceolate; dorsal and anal low, confluent with caudal, which ends in a point; pectorals small, their ends scarcely reaching midway from their base to front of anal; ventrals with 2 filamentous rays, the outer scarcely as long as head, the inner  $\frac{1}{2}$  as long. Color in spirits, reddish brown, with silvery reflections on sides; head silvery, upper part of eye black; fins the color of the body, with no dark edgings. Coasts of Cuba; rare. Here described from a specimen from Havana, Cuba,  $2\frac{1}{2}$  inches in length, sent by Professor Poey. Poey has also sent a drawing of his original type, a much larger specimen, which he describes as follows:

Head  $5\frac{1}{2}$ ; depth 8; snout rounded; eye large,  $4\frac{1}{2}$  in head; maxillary reaching posterior border of eye; teeth small, slender, with a villiform band behind them; teeth on vomer and palatines; scales small; head scaly, except on snout; lateral line high; branchiostegals 7; dorsal beginning over second third of pectoral, joining anal behind; about 100 rays in each fin; vent a little behind first third of length. Yellowish brown, silvery on side of head; no black on fins. Air bladder distinct; no pyloric caeca. Intestine with 2 short turns. Cuba. (Poey.) Air bladder, gill rakers, and ventral not described. Length 230 mm. Rare; not reaching a foot in length. (Named for Mariano de la P. Graells, director of the Botanic Garden at Madrid, "comme témoignage de mon estime pour ses travaux scientifiques, et pour la zèle qu'il déploie . . . pour l'acquisition des objets et l'acclimatation des espèces.")

*Ophidium graellsii*, POEY, Memorias, II, 425, 1860, Havana (Coll. Poey); POEY, Synopsis, 402, 1867.

#### 948. CHILARA, Jordan & Evermann.

*Chilara*, JORDAN & EVERMANN, Check-List Fish. N. and M. A., 482, 1896 (*taylori*).

This genus contains a single robust species which differs from *Ophidium* only in the presence of a stout concealed spine at tip of opercle; the air bladder



is oblong-ovate, the head naked and the snout without spine. (*χιλάρι*, the modern Greek name of the species of *Ophidion* and *Rissola*.)

2858. CHILABA TAYLOBI (Girard).

Head 6; depth 8; head large, little compressed, naked; top of head with conspicuous mucous pores; dorsal fin beginning over the pectorals; outer ray of ventral little more than  $\frac{1}{2}$  length of head, inner about  $\frac{1}{3}$ ; air bladder ovate, not contracted; 7 gill rakers below the angle of the arch; pseudobranchia developed; no spine on the end of the snout; opercle with a flat spine concealed in its membranes; outer teeth in both jaws considerably enlarged, the upper largest. Color light olive; head and upper parts covered with conspicuous round dark, olive-brown spots; chin dusky; vertical fins edged with black. Length 12 inches. Coast of California, from Monterey to San Diego; not rare in waters of moderate depth. (Named for A. S. Taylor, its discoverer.)

*Ophidium taylori*, GIRARD, Pac. R. R. Surv., x, Fishes, 138, 1858, Monterey, California (Type, No. 867. Coll. A. S. Taylor); JORDAN & GILBERT, Synopsis, 793, 1883.

949. RISSOLA, Jordan & Evermann.

*Rissola*, JORDAN & EVERMANN, Check-List Fish. N. and M. A., 483, 1896 (*marginatum*).

This genus contains species agreeing with *Ophidion* in general characters, but with the air bladder short, broad, spherical or kidney-shaped, with a posterior foramen. Species chiefly of the Mediterranean. (Named for Anastase Risso, apothecary at Nice, author of the *Ichthyologie de Nice*, 1810, and *Histoire Naturelle de l'Europe Méridionale*, 1826, two of the very best of local faunal works, the foundation of our knowledge of the fishes of the Mediterranean.)

2859. RISSOLA MARGINATA (DeKay).

Head  $6\frac{1}{2}$ ; depth  $7\frac{1}{2}$ ; eye 4 in head; maxillary reaching posterior margin of orbit; air bladder short and broad, with a foramen on the under side; upper ray of ventral about equaling length of head; inner ray  $\frac{1}{2}$  length of outer; gill rakers 4; color nearly plain brownish; dorsal and anal fins margined with black. Coast of the United States, from New York south to Pensacola and the coast of Texas; not very common; very similar to the Mediterranean species *Rissola rochii* (Müller), but probably distinct. (*marginatus*, margined.)

*Ophidium marginatum*, DE KAY, N. Y. Fauna: Fish., 315, 1842, New York Harbor; PUTNAM, Proc. Bost. Soc. Nat. Hist. 1874, 342; JORDAN & GILBERT, Synopsis, 792, 1883.

*Ophidium josephi*, \* GIRARD, U. S. and Mex. Bound. Surv., Zool., 29, 1859, Saint Joseph Island, Texas; JORDAN & GILBERT, Synopsis, 793, 1893.

\* The scanty description of *Ophidium josephi* agrees fairly with either *Rissola marginata* or *Ophidion holbrooki*, and may be either. The following is the substance of Girard's account:

"Head 6 in length; eye moderate, 4 in head; maxillary extending to opposite its posterior margin; origin of dorsal at some distance behind base of pectorals. Body shorter and pectorals more elongate than in *O. taylori*. Pale olive, sprinkled all over with brownish specks; belly and sides of head plain; vertical fins edged with black."

950. *OTOPHIDIUM*, Gill.*Otophidium*, GILL, in JORDAN, Cat. Fish. N. A., 126, 1885 (*omostigma*).

This genus differs from *Ophidion*, in the form of the air bladder, which is short, thick, and with a large foramen (not examined in *O. galeoides*). The opercle ends in a concealed spine as in *Chilara*. Species American, so far as known. ( $\sigma\upsilon\varsigma$   $\acute{\alpha}\tau\rho\acute{\omicron}\varsigma$ , ear; *Ophidium*.)

## a. Gill rakers 4.

b. Head long,  $4\frac{1}{2}$  to  $4\frac{3}{4}$  in length; depth  $5\frac{1}{4}$  to 6.

c. Scapular region with a jet-black spot; pseudobranchiæ little developed; ventrals  $\frac{1}{2}$  length of head; maxillary  $1\frac{1}{2}$  in head. **OMOSTIGMUM**, 2860.

cc. Scapular region without jet-black spot; pseudobranchiæ well developed; ventrals with the inner ray longest,  $\frac{1}{2}$  head; body with dark cross bands. **INDEFATIGABILE**, 2861.

bb. Head moderate,  $5\frac{1}{2}$  in body; depth 6; a pale spot before dorsal; pale spots along lateral line; ventral  $\frac{1}{2}$  head. **GALEOIDES**, 2862.

2860. *OTOPHIDIUM OMOSTIGMUM* (Jordan & Gilbert).

Head  $4\frac{1}{2}$  in length; depth about 6. Body comparatively short, highest at occiput, thence tapering rapidly to tip of tail; upper profile of head very convex; snout blunt; mouth horizontal, the lower jaw included; maxillary not quite reaching posterior border of orbit; teeth in jaws uniform, strongly incurved, in rather broad bands; a single series of small teeth on vomer, those on palatines minute; maxillary  $1\frac{1}{2}$  in head; eye large, 3 in head, much larger than snout, equaling twice interorbital width; opercle terminating in a strong, compressed spine, the length of which is about  $\frac{2}{3}$  diameter of pupil; gill rakers very small, 4 below on anterior arch. Longest ventral filament  $\frac{1}{2}$  length of head; the shorter  $\frac{2}{3}$  length of longer. Distance from origin of dorsal to tip of snout  $3\frac{1}{2}$  in total length; distance from origin of anal to snout  $2\frac{1}{2}$  in total length. Scales minute, embedded. Pseudobranchiæ probably present (type reexamined by us). Air bladder short, thick, with a large posterior foramen. Color light olive green, silvery on belly, cheeks, and lower side of head; sides above with a few irregular, large, scattered, dark blotches, about 9 of these along base of dorsal fin; an intensely black, round blotch on scapular region, rather larger than pupil; dorsal with black blotches; anal largely black; upper half of eye black, lower half bright silvery. Gulf of Mexico. A single specimen,  $3\frac{1}{2}$  inches long, taken from the stomach of a red snapper, at the Snapper Banks off Pensacola. ( $\acute{\omega}\mu\omicron\varsigma$ , shoulder;  $\sigma\tau\iota\gamma\mu\alpha$ , spot.)

*Genypterus o. nostigma*, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1882, 301, Pensacola Snapper Banks (Coll. Jordan & Stearns. Type, 29670, U. S. Nat. Mus.); JORDAN & GILBERT, Synopsis, 963, 1883.

*Otophidium omostigma*, GOODE & BEAN, Oceanic Ichthyology, 345, fig. 305, 1896.

2861. *OTOPHIDIUM INDEFATIGABILE*, Jordan & Bollman.

Head  $4\frac{2}{3}$  ( $4\frac{1}{2}$  in total); depth  $5\frac{2}{3}$  ( $5\frac{1}{2}$ ); eye large, 3 in head; snout 4. Body rather short, compressed, width of nape  $2\frac{1}{2}$  in head. Mouth large; maxillary reaching to opposite posterior margin of pupil,  $1\frac{1}{2}$  in head; outer row of teeth of each jaw very slightly enlarged. Interorbital space 2 in eye; interorbital area with a thin crest under the skin, this ending in 2 com-

pressed spines, 1 turning forward, the other backward over front of eye, these spines concealed by the skin. Gill rakers short and thick, less than  $\frac{1}{2}$  pupil, 4 developed. Dorsal beginning at end of anterior third of pectorals, longest ray  $3\frac{1}{2}$  in head; pectorals 2 in head; inner ventral filament longest, 2 in head. Air bladder short and thick, with a foramen. Scales very small, more or less imbricated on body; head naked. Opercle with a sharp, partly concealed spine. Pseudobranchiæ present. Color pale yellowish brown, silvery on belly and sides of head; back with about 12 irregular dark cross bands, the alternate ones being narrower and broken up into spots, 2 before dorsal; a few scattered spots about as large as pupil on sides, these most distinct about the shoulder; dorsal pale, first rays black, and with 3 or 4 other black blotches on upper part; anal black, margined with white; pectorals pale, axil dusky; caudal and posterior part of anal pale; chin pale. A single specimen obtained at Indefatigable Island, Galapagos Archipelago. Length 4 inches. (*indefatigabilis*, tireless.)

*Otophidium indefatigabile*, JORDAN & BOLLMAN, Proc. U. S. Nat. Mus. 1889, 172, Indefatigable Island, in the Galapagos Archipelago. (Type, No. 44393. Coll. Albatross.)

2862. *OTOPHIDIUM GALEOIDES* (Gilbert).

Head  $5\frac{1}{2}$  in length; depth 6. D. 125. Maxillary reaching beyond pupil,  $2\frac{1}{2}$  in head; snout  $4\frac{1}{2}$ ; eye  $3\frac{1}{2}$ . Gill rakers short and broad, 4 of them developed. Opercle ending in a sharp concealed spine. Outer teeth little enlarged. Dorsal beginning over middle of pectorals, the nape equidistant between front of dorsal and tip of snout. Caudal very short and bluntly rounded, as in *Chilara taylori*, the rays not projecting beyond dorsal and anal; pectorals  $1\frac{1}{2}$  in head; ventral filament  $\frac{1}{2}$  head. Scales as in *C. taylori*, not at all imbricated, arranged with their long axes frequently at right angles to each other; head naked. Color light olive, without bars, a narrow dusky streak along base of dorsal, and a round light spot at origin of dorsal; a series of small olive-brown spots along lateral line, with a few scattering spots below it but none above; nape and head without spots; vertical fins translucent; dorsal with a large black blotch on tip of anterior rays, the fin behind this narrowly edged with black, which does not surround the caudal; anal with much silvery-white pigment anteriorly on distal portion, becoming dusky behind; pectorals translucent, edged with white below; peritoneum, buccal, and gill cavities white. Closely related to *Otophidium indefatigabile*, differing in color, and in the much shorter head, smaller mouth, less imbricated scales, the more posterior insertion of dorsal, and the absence of spines on head. Air bladder not examined. Pseudobranchiæ present. Gulf of California. One specimen,  $5\frac{1}{2}$  inches long, from Albatross Station 3025, in  $9\frac{1}{2}$  fathoms. (Gilbert.) ( $\gamma\alpha\lambda\eta$ , shark;  $\epsilon\acute{\iota}\delta\omicron\varsigma$ , appearance.)

*Otophidium galeoides*, GILBERT, Proc. U. S. Nat. Mus. 1890, 110, Gulf of California, lat.  $31^{\circ} 21' 15''$  N., long.  $113^{\circ} 59'$  W. (Type, No. 44381. Coll. Albatross.)

Family CCIX. LYCODAPODIDÆ.

Deep-sea fishes allied to the *Fierasferide*, differing chiefly in the normal position of the vent, which is remote from the head, and just before the

anal fin; gill openings large, the membranes united anteriorly only, free from the isthmus, as in *Fierasfer*. Pseudobranchia wanting; no scales; no lateral line; no ventral fins. One genus with 4 known species, from the North Pacific.

#### 951. LYCODAPUS, Gilbert.

*Lycodapus*, GILBERT, Proc. U. S. Nat. Mus. 1890, 107 (*fierasfer*).

Body naked. Ventrals wanting. Vertical fins united around the tail. Gill openings wide, continued forward under the throat; the gill membranes anteriorly narrowly united, loosely joined to the isthmus by a fold of lax skin. Branchiostegals 6. No pseudobranchia. Gills 4, a wide slit behind inner arch. Gill rakers developed. Teeth present in jaws and on vomer and palatines, none of them enlarged. Vent remote from the throat. (*Lycodes*; *ἄπους*, footless.)

a. Body slender, the depth 8 to 11 in length.

b. Head rather large,  $4\frac{1}{2}$  to  $5\frac{1}{2}$  in length.

c. Head, body, and fins with very many mucous pores; dorsal rays 70; anal 60.

DERMATINUS, 2863.

cc. Head, body, and fins with very few mucous pores; dorsal rays 82; anal 70.

FIERASFER, 2864.

bb. Head small,  $7\frac{1}{2}$  in length; gill openings not extending above base of pectorals.

PARVICEPS, 2865.

aa. Body very slender, the depth about 15 in length; dorsal rays about 100.

EXTENSUS, 2866.

#### 2863. LYCODAPUS DERMATINUS, Gilbert.

Head  $4\frac{1}{2}$ ; depth  $1\frac{1}{2}$  in head; eye 5 in head; snout 4; maxillary  $2\frac{1}{2}$ . D. 70; A. 60. Very similar to *L. fierasfer*, but the head, body and fins covered with a thick loose skin which contains numerous pores, or openings for the mucous canals. One series of these runs along middle of sides and forms the lateral line; it rises anteriorly above the gill opening, and is continued forward on top of head, the two meeting between eyes; a second series runs between eye and upper lip, and curves around on middle of cheek, running upward to behind eye: one series runs along a fold bordering mandible, 1 along preopercular margin, and 1 on opercle. In *L. fierasfer* a few pores are visible on mandible, and 1 or 2 can frequently be made out on preopercular margins. The skin is very thin and delicate, and the fin rays are very evident through the membrane. The general proportions and the dentition of the type are essentially as in *L. fierasfer*, but the vomerine teeth are long and hooked backward. Mandible heavier than in *L. fierasfer*. Origin of dorsal vertically above axil of pectorals. Length of head and trunk  $\frac{1}{2}$  total length. Teeth in narrow bands in the jaws, a single series on vomer and palatines. Gill membranes very narrowly joined below and free from the isthmus, as in *L. fierasfer*. Pectorals much longer than in *L. fierasfer*. General color in spirits light brownish yellow, made somewhat dusky by the pigment spots in the skin; body, and especially the fins, darker posteriorly. Aleutian

Islands, in deep water. Only the type known, an example  $4\frac{1}{2}$  inches long. (*δέρμα*, skin; *dermatinus*, skinny.)

*Lycodapus dermatinus*, GILBERT, Rept. U. S. Fish Comm. 1893 (1896), 471, pl. 35, Aleutian Islands, lat.  $37^{\circ} 54' 10''$  N., long.  $123^{\circ} 30'$  W., at Albatross Station 3162, in 552 fathoms.

2864. *LYCODAPUS FIERASFER*, Gilbert.

Head  $5\frac{1}{2}$ ; depth 10; eye  $4\frac{1}{2}$  in head; snout  $3\frac{1}{2}$ ; maxillary 2 to  $2\frac{1}{2}$ . D. 82; A. 70. Body compressed, elongate, tapering rather rapidly backward, the tail not produced to a filament; head flat above, the cheeks deep, vertical, the mouth very oblique, with the lower jaw slightly the longer and nearly entering the upper profile; skull very thin and papery, translucent; jaws weak; gape of mouth wide, the maxillary reaching vertical from behind front of pupil; teeth all small, in a very narrow band in jaws, in a single series on vomer and palatines; interorbital width  $\frac{2}{3}$  of eye; snout broad, depressed, spatulate, its tip prominent, turned upward, the upper profile thus longitudinally concave; an evident median ridge on snout and interorbital space; gill slits continued forward below to vertical from middle of eye, the membranes united for a distance equaling diameter of pupil; gill rakers short, less than diameter of pupil, strongly toothed, about 10 on horizontal limb of arch; head without conspicuous mucous pores or cavities. Dorsal beginning well forward, its distance from occiput slightly less than that from occiput to nostril. Dorsal and anal rays slender, all articulated, branched only at tips; caudal not distinct, the rays springing from end of tail not projecting beyond the others; origin of anal immediately behind vent, its distance from snout nearly equaling  $\frac{1}{2}$  total length; pectorals narrow, varying in length, about  $2\frac{1}{2}$  in head. Body and fins invested in a rather lax transparent skin, without traces of scales. Color, body translucent, dusted with black specks; abdomen blackish; lips, inside of mouth, lining of gill cavity, and peritoneum jet-black; iris silvery. (Gilbert.) North Pacific. The types, several specimens, the longest  $5\frac{1}{2}$  inches, from *Albatross* Stations 2980, 3010, 3072, off Lower California, in 610 to 1,005 fathoms. Also taken near Unalaska in 109 fathoms. (*Fierasfer*, the pearfish.)

*Lycodapus fierasfer*, GILBERT, Proc. U. S. Nat. Mus. 1890, 108, off Lower California, in 610 to 1,005 fathoms (Coll. Dr. Gilbert); JORDAN, Proc. Cal. Ae. Sci. 1896, 234, pl. 23.

2865. *LYCODAPUS PARVICEPS*, Gilbert.

Head  $7\frac{3}{4}$ ; depth 11; eye  $4\frac{1}{2}$  in head; snout  $3\frac{1}{2}$ ; least interorbital width 5; maxillary  $2\frac{1}{2}$ ; pectoral  $2\frac{1}{2}$ . D. 100; A. about 85 (both counted to middle of caudal); P. 9; no ventrals. Upper profile of head nearly straight, not longitudinally concave as in *L. fierasfer*; head deeper and narrower, the snout less spatulate; skin thicker. A conspicuous series of pores on mandible and along preopercular margin; gill slit very oblique, extending anteriorly as far as vertical from eye, the membranes then narrowly united, free from the isthmus except at extreme front; gill slit superiorly much more restricted than in *L. fierasfer*, not extending above base of pectorals, while in the latter it extends above them for  $\frac{1}{2}$  diameter of eye. Mouth

oblique, maxillary reaching vertical from middle of eye; jaws even at tip, the mandible slightly included laterally; mandibular teeth in a moderate band anteriorly, the inner series enlarged, narrowing posteriorly to a single row; premaxillary teeth of uniform size, in a narrow band throughout; vomer with 4 canine-like teeth; palatine teeth small, in a single close-set series. Distance from origin of dorsal to occiput slightly less than that from occiput to posterior nostril; head and trunk contained  $3\frac{1}{2}$  in tail. Body brownish in spirits, fins whitish, translucent; everywhere dusted with black specks; tail and fins distinctly blackish posteriorly; orbit blackish above; gill cavity silvery, blackish anteriorly; mouth blackish, except anteriorly; peritoneum black, the color not showing through the abdominal wall. Similar to *L. fierasfer*, differing in the much smaller head, longer, slenderer body, the thicker skin with more evident mucous pores, and in the more restricted gill openings. Aleutian Islands, in moderately deep water. Only the type known, a specimen about 5 inches long. (Gilbert.) (*parvus*, small; *-ceps*, head.)

*Lycodapus parviceps*, GILBERT, Rept. U. S. Fish Comm. 1893 (1896), 455, north of Unalaska Island at Albatross Station 3324, in 109 fathoms. (Coll. Dr. Gilbert.)

2866. *LYCODAPUS EXTENSUS* (Gilbert).

Head  $6\frac{1}{2}$ ; depth  $15\frac{1}{2}$ ; eye  $4\frac{1}{2}$  in head; snout  $3\frac{1}{2}$ ; interorbital width  $1\frac{1}{2}$  in eye; pectoral  $2\frac{3}{4}$  in head. D. 96 (the extreme end of the tail wanting). Gill openings as in *L. fierasfer*, extending well above base of ventrals. Skin thin, the mucous pores inconspicuous, evident on mandible and along margin of preopercle. Upper profile of head longitudinally concave, shaped as in *L. fierasfer*, but slenderer, its depth greater than that of body. Mouth oblique, the maxillary reaching vertical from middle of eye,  $2\frac{1}{2}$  in head. Teeth in narrow bands in each jaw, tapering laterally to single series; vomerine teeth more numerous than in *L. parviceps* or *L. fierasfer*, small, not canine-like, in a single series; palatine teeth wanting, as in some individuals of *L. fierasfer*. Occiput midway between front of dorsal and anterior nostril; pectorals slenderer and longer than in *L. fierasfer*. Head and trunk contained  $2\frac{3}{4}$  times in tail. Color light brownish, the black peritoneum visible through the skin of the abdomen; mouth and gill cavity largely dusky; a narrow dark-brown streak along base of dorsal and anal, occupying, toward tip of tail, the entire height of both fins. An extremely slender elongate form, with head smaller than *L. fierasfer*, but otherwise resembling that species more than *L. parviceps*. Aleutian Islands, in rather deep water. Only the type, a specimen 4 inches long, known. (Gilbert.) (*extensus*, stretched out.)

*Lycodalepis extensus*, GILBERT, Rept. U. S. Fish Comm. 1893 (1896), 455, north of Unalaska, at Albatross Station 3324, in 109 fathoms. (Coll. Dr. Gilbert.)

Family CCX. FIERASFERIDÆ.

(THE PEARL-FISHES.)

Body elongate, compressed, tapering into a long and slender tail; no scales; teeth cardiform, on jaws, vomer, and palatines; canine teeth often present; no barbels; lower jaw included; vent at the throat; gill mem-

branes somewhat united, free from the isthmus; no pseudobranchiæ; no pyloric cæca; vertical fins very low, confluent, without spines; no ventral fins; pectoral fins present or absent. Small shore fishes of tropical seas, often living in shells of mollusks, echinoderms, etc., being especially often commensal with the pearl oyster and with the larger *Holothuria*. Genera 3; species 12. (*Ophidiidæ*, group *Fierasferina*, Günther, Cat., IV, 381-384, 1862.)

a. Pectoral fins present; no distinct caudal fin; gill membranes connected anteriorly only. FIERASFER, 952.

952. FIERASFER, Cuvier.

- Fierasfer*, CUVIER, Règne Anim., Ed. 1, II, 239, 1817 (*imberbe=acus*).  
*Echiodon*, THOMPSON, Proc. Zool. Soc. London 1837, 55 (*drummondii*).  
*Diaphasia*, LOWE, Proc. Zool. Soc. London 1843, 92 (*acus*).  
*Oxybeles*, RICHARDSON, Voy. Erebus and Terror, Fishes, 74, 1844-48 (*homei*).  
*Porobronchus*, KAUF, Ann. Mag. Nat. Hist. 1860, 272 (larva of *Fierasfer acus*).  
*Carapus*,\* GILL, Proc. Ac. Nat. Sci. Phila. 1864, 152 (after RAFINESQUE, 1810; not type).  
*Vezillifer*, GASCO, Bull. Assoc. Nat. Med. Napoli 1870, 59 (larva of *Fierasfer acus*).  
*Lefroyia*, JONES, Zoologist, IX, 1874, 3838 (*bermudensis*).

Gill membranes little connected, leaving the isthmus bare. No distinct caudal fin; pectoral fins developed. The species of this genus are not well known, and their characters and nomenclatures are uncertain. It is not unlikely that the American species are all reducible to one, *Fierasfer affinis* or *dubius*, but our scanty material will not justify us in taking this view. (*Fierasfer*, the ancient name, from *φειρός*, sleek and shining.)

- a. Vomer with canine teeth; pectoral about  $\frac{1}{2}$  length of head.  
 b. Front teeth of upper jaw enlarged; head 7 to 8 in length; depth  $11\frac{1}{4}$  to 15 times in length of body. AFFINIS, 2867.  
 bb. Front teeth of upper jaw not enlarged; head  $6\frac{1}{2}$  in length; depth about 10 $\frac{1}{2}$  times in length of body. ARENICOLA, 2868.  
 aa. Vomer with small teeth, scarcely canine-like; pectoral about  $2\frac{1}{2}$  in head; head 7 to  $8\frac{1}{2}$  in body. BERMUDENSIS, 2869.

2867. FIERASFER AFFINIS† (Günther).

(PEARL-FISH.)

Head  $7\frac{1}{2}$ ; depth of head 15. Maxillary extending slightly beyond orbit; lower teeth larger than the upper, except 2 to 4 front teeth of upper jaw, which are about equal to lower teeth; vomer with 3 to 6 teeth, 2 or 3 of

\* The name *Carapus*, Rafinesque, has been substituted for *Fierasfer* by Gill and Poey. This change seems to us not justifiable, as it is certainly not desirable. The name *Carapus* first appears in Rafinesque's *Indice d'Istiologia Siciliana*, 57, 1810. No type is mentioned by Rafinesque, but the diagnosis is taken from that of Lacépède's second subgenus under *Gymnotus*, which contains the three species, *carapo*, *fierasfer*, and *longirostratus*. Of these species, *carapo* is the original Linnæan type (Ed. x) of the genus *Gymnotus*. *Carapus* should therefore be regarded as a synonym of *Gymnotus*. The Brazilian name *carapo* evidently suggested the word *Carapus*, although Dr. Gill derives the name from *καπα*, head; *αροος*, footless, an ex post facto distinction from *Ophedion*. In a list of Sicilian fishes, on page 37 of Rafinesque's *Indice*, published somewhat later, the name *Carapus acus* appears for *Fierasfer acus*. This reference of a species of *Fierasfer* to *Gymnotus* or *Carapus* was due to Rafinesque's ignorance of its relations.

† In the Museum of Comparative Zoology is "one valve of a pearl oyster, in which a specimen of *Fierasfer dubius* is beautifully inclosed in a pearly covering, deposited on it by the oyster." (Putnam.)

these canine-like. Pectoral  $\frac{1}{2}$  head; vent under base of pectoral. Dorsal fin low, but distinct; anal much more developed than dorsal, its longest rays about in the middle of the fish. Air bladder long, slightly constricted behind. Gill membranes not covering isthmus. Color in spirits, uniform light brown, with a short silvery band along the sides of the abdomen made by confluent spots. (Putnam, description of *F. dubius*.) Panama; especially common among the Pearl Islands, chiefly in shells of pearl oysters. This species should probably stand as *Fierasfer affinis*.

The following notes are from numerous specimens, 3 to 4 inches long, from Pearl Islands, collected by Prof. Bradley, these also being types of *Fierasfer dubius*: Head  $6\frac{1}{2}$  to  $7\frac{1}{2}$ ; eye  $4\frac{1}{2}$  to 5 in head. Teeth in upper jaw small, acute, in a rather narrow band; sometimes a few in the front of the jaw inconspicuously enlarged; those in lower jaw and on palatines conic, blunt, in somewhat wider bands, the outer series of lower jaw enlarged, canine-like; vomer with a narrowly oblong patch of small, blunt teeth, surrounding a median series of 3 to 6 conspicuously enlarged, retrorsely curved canines, which are usually much the largest teeth in the mouth. Two specimens from Albatross Station 3021, Lower California, agree in general with the above account: Head  $7\frac{1}{2}$ ; depth  $11\frac{1}{2}$ ; eye 4; 2 upper teeth on each side somewhat enlarged, about as large as lateral teeth on mandibles; vomerine canines larger. Professor Putnam refers also to *Fierasfer dubius* specimens from Key Biscayne, Florida (Coll. Theodore Lyman); Tortugas (Coll. Gustav Würdemann); Cape Florida (Coll. Würdemann), and New Providence, Bahama (Coll. F. G. Shaw). These specimens apparently belong rather to *Fierasfer bermudensis*, if that species be different. (*affinis*, related, to *Fierasfer acus*.)

? *Fierasfer affinis*,\* GÜNTHER, Cat., IV, 381, 1862, no locality given.

*Fierasfer dubius*, PUTNAM, Proc. Bost. Soc. Nat. Hist. 1874, 344, Pearl Islands (Coll. Prof. Frank H. Bradley); JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1882, 629; JORDAN & GILBERT, Synopsis, 791, 1883.

#### 2868. FIERASFER ARENICOLA, Jordan & Gilbert.

Head  $6\frac{1}{2}$  in length; depth  $10\frac{1}{2}$ ; eye 5 in head; snout 5. Body with nape slightly elevated, thence tapering regularly to the tail. Snout blunt, rounded, protruding; mouth subinferior, nearly horizontal, large, the lower jaw included; gape wide, the maxillary  $\frac{1}{2}$  length of head, extending beyond vertical from orbit; teeth in upper jaw very small, acute, in a narrow band, none of them enlarged; those in lower jaw and on vomer blunt, conic, in a wide band; those in outer series acute; a few on each side of mandible and 2 or 3 anteriorly on vomer, enlarged, canine-like. Gill openings very wide, the branchiostegal membranes little united, leaving nearly all of isthmus uncovered; the membranes

\* *Fierasfer affinis*, Günther, is thus described:

"The length of the head is  $\frac{1}{2}$  of the total; its greatest width is rather less than  $\frac{1}{2}$  of its length. Gill openings rather wide, the united gill membranes leaving the greater portion of the isthmus uncovered. Teeth cardiform; a pair in front of the upper jaw, a series on the side of the lower, and several others on the vomer larger than the rest. Dorsal fin low but very distinct. The length of the pectoral nearly  $\frac{1}{2}$  that of the head. (This species is) similar to *F. acus*, but with a very different dentition." (Günther.) Described from a specimen 8 inches long, from unknown locality. This description, so far as it goes, agrees with *Fierasfer dubius*, but the specimen may not be American.



united as far back only as vertical from end of maxillary; opercle adherent above the upper angle, which is produced in a point extending above the base of pectorals; below the angle the opercular margin runs very obliquely forward. Eye large, greater than interorbital width. Origin of dorsal fin distant from nape by the length of the head, the fin a very inconspicuous fold anteriorly, becoming higher posteriorly, where the rays are evident; anal well developed along entire length, beginning immediately behind vent and running to tail, its rays visible; caudal exceedingly short; pectorals very well developed, more than  $\frac{1}{2}$  length of head; vent just in front of base of pectorals. Head and body perfectly translucent; a faint silvery luster on middle of sides anteriorly; a few inconspicuous small light yellowish spots along middle of sides (disappearing in alcohol); tip of tail dusky; upper margin of orbit black. Pacific coast of Mexico. A single specimen,  $3\frac{1}{2}$  inches long, was found buried in the sand at low tide on the beach at Mazatlan. This specimen may be identical with *Fierasfer dubius*, but it is more robust than Putnam's types, with longer head and without enlarged teeth in upper jaw. It may be regarded as distinct, pending investigation. (*arena*, sand; *colo*, I inhabit.)

*Fierasfer arenicola*, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1881, 363, Mazatlan. (Type, No. 29244. Coll. C. H. Gilbert.)

2869. FIERASFER BERMUDENSIS (Jones).

Head  $8\frac{1}{2}$  in length; eye 4, longer than snout; mouth large, the maxillary reaching beyond orbit; pectoral  $2\frac{1}{2}$  in head. Teeth small, acute, uniserial, 3 in a line on the vomer; palatine teeth small.\* Color pale brownish, a bluish streak crossing the nape between the opercles, 4 pale points on the back. Vertebrae 100. Length 140 mm. West Indies. This description (by Poey) from a specimen taken in the stomach of a holothurian at Havana. Others are recorded from Key West and St. Thomas, the latter from an oyster; not seen by us; doubtfully distinct from *Fierasfer dubius* or *affinis*, but the vomerine teeth said to be smaller.

†*Carapus affinis*, POEY, Synopsis, 402, 1867; not *Fierasfer affinis*, GÜNTHER.

*Lefroyia bermudensis*, JONES, Zoologist, ix, 1874, 3838, Bermuda.† (Coll. General Lefroy.)

\* In another specimen, according to Poey, the teeth are villiform, with an enlarged series outside, the lower teeth largest, the ninth, tenth, and eleventh largest; teeth on vomer small, acute, in a row; palatine teeth bluntish.

† *Fierasfer bermudensis* (Jones) was thus originally described:

"Total length rather more than  $4\frac{1}{2}$  inches. Greatest depth at the vertical of the pectorals  $3\frac{1}{2}$  lines. The length of the head is slightly more than  $\frac{1}{2}$  of the total length. The greatest width of the head rather less than  $\frac{1}{2}$  of its length. Body naked, attenuate, compressed. Facial outline angose. Eye moderate, horizontal diameter of eyeball  $1\frac{1}{2}$  lines; vertical diameter  $1\frac{1}{2}$  lines. Gape of the mouth ovoid. Lower jaw shorter and received within the upper. Cardiform teeth of irregular size in both jaws, vomer and palatines, those of the latter largest. Branchiostegals 7, inflated, united below. Vent thornlike. Pectorals originating at the upper angle of the operculum, 3 lines in extent, and composed of very delicate soft rays. Dorsal indistinct, commencing in a groove about the vertical of the twentieth anal ray, continuous to caudal extreme, where, in conjunction with the anal, it forms a small filamentous tip. Anal prominent, commencing immediately behind the vent in advance of the vertical of the upper angle of the operculum, and extending to the caudal extreme. About its center it is equal in depth to that of the body at same position. Owing to the delicate texture of the fins it is impossible to ascertain certainly the number of rays, but those of the anal exceed 140. Color, when dried out of spirits, golden yellow; the body transparent, showing the vertebrae within; a condition, according to Lefroy, equally observable in life.

"I propose to publish it as *Lefroyia bermudensis*, in compliment to the gallant officer to whom I am indebted for the specimen." (J. Matthew Jones.)

This species is probably identical with the one called *affinis* by Poey.

## Family CCXI. BROTULIDÆ.

## (THE BROTULOID FISHES.)

Body elongated, compressed, regularly tapering behind, the tail generally subtruncate at base of caudal fin, not isocercal; vent submedian; scales cycloid and minute, embedded in the lax skin, which more or less envelops the fins, sometimes wanting; gill openings very large, the membranes mostly free from the isthmus; vertical fins united or contiguous at base of caudal; dorsal fin commencing not far from nape; caudal narrow or pointed; ventral fins small, few-rayed, attached to the humeral arch and more or less in advance of pectoral. Pyloric cæca few (1 or 2), rarely obsolete or in increased number (12); maxillaries generally enlarged behind and produced toward their upper angle. (Gill.) Pseudobranchiæ small or wanting, hypercoracoid with the usual foramen, as in Blennioid fishes. These fishes are closely related to the *Zoarcidae*. In spite of various external resemblances to the *Gadida*, their affinities are rather with the Blennioid forms than with the latter. Genera about 45, species about 100; largely of the depths of the sea; 2 species degenerated into blind cave fishes. We have not had material for any elaborate study of these fishes and follow closely the arrangement given by Goode & Bean. (*Brotuloidæ*, Gill, Proc. Ac. Nat. Sci. Phila. 1863, 232, and 1884, 175.)

## BROTULINÆ:

- a. Snout and lower jaw each with well-developed barbels; vertical fins united; teeth on vomer and palatines.  
 b. Ventrals each reduced to a bifid filament. BROTULA, 953.  
 aa. Snout and lower jaw without barbels.

## LUCIFUGINÆ:

- c. Species blind, dwelling in fresh-water streams in caves; barbels replaced by cilia.  
 d. Palatines with strong teeth; teeth in lower jaw strong. STYGCOLZ, 954.  
 dd. Palatines toothless; teeth in jaws villiform. LUCIFUGA, 955.  
 cc. Species marine, the eyes usually well developed.

## BROSOMOPHYCINÆ:

- e. Caudal fin differentiated, on a distinct caudal peduncle.  
 f. Snout and lower jaw with small cilia; head naked, or nearly so. BROSOMOPHYCIS, 956.  
 ff. Snout and lower jaw without cilia; head more or less scaly. OGILBIA, 957.  
 ee. Caudal fin not differentiated, without distinct peduncle.

## BYTHITINÆ:

- g. Ventrals inserted on the isthmus, not far from the humeral symphysis.  
 h. Pectorals normal, simple; eyes present.  
 i. Lateral line present posteriorly, but broken in the middle; palatines with teeth; ventrals a pair of filaments each of 2 closely united rays. BYTHITES, 958.  
 ii. Lateral line obsolete posteriorly.  
 j. Ventrals each of a single ray.  
 k. Lateral line distinct on front of body.  
 l. Preopercle without spines; head scaly (except snout); opercle with a single spine; vent median. CATÆTYX, 959.

- ii. Preopercle with 3 or 4 spines, opercle with a single one; head partially naked.  
DICROMITA, 960.
- kk. Lateral line obsolescent, almost, or quite invisible; opercle with a feeble spine; head smooth; eyes small.
- m. Ventral consisting of a single ray.  
BASSOZETUS, 961.
- mm. Ventral bifid.  
MCEBIA, 962.
- jj. Ventrals each of a pair of rays.
- n. Caudal fin exerted, but confluent with anal and dorsal.
- o. Head scaly.
- p. Preopercle with small spines at its angle, opercle with 1 spine.  
NEORHYTHMIS, 963.
- pp. Preopercle unarmed.
- q. Opercle with 2 spines; ventrals close together.  
BENTHOCOMETES, 964.
- qq. Opercle with 1 strong spine; ventrals far apart.  
BASSOGIGAS, 965.
- nn. Caudal not confluent with vertical fins, but without distinct peduncle; teeth on jaws, vomer, and palatines in villiform bands; preopercle unarmed; head scaly.
- r. Opercle with a flat spine; snout much produced and dilated; lateral line very indistinct (or absent?).  
BARATHRODEMUS, 966.
- rr. Opercle a triangular flap, unarmed; lower pectoral rays prolonged, the lowest filamentous.  
NELATONUS, 967.
- iii. Lateral line represented by 3 rows of pores—dorsal, lateral, and ventral; head with spines.
- s. Ventrals of 2 distinct rays; opercular spine moderate, straight.  
POROGADUS, 968.
- ss. Ventrals each of 2 united rays, opercular spines strong, curved.  
PENOPUS, 969.
- hh. Pectorals with the lower rays differentiated.
- t. Preopercle armed with 3 spines; opercle armed with 1 spine; lateral line obsolete posteriorly; ventrals bifid.  
DICROLENE, 970.
- tt. Preopercle unarmed; a single spine on opercle; lateral line absent (?); ventrals each a pair of filaments, closely united throughout.  
MIXONUS, 971.
- APHYONINÆ:
- gg. Ventrals inserted on humeral symphysis; lateral line obsolete (in almost every case); ventrals each of a single filament; body naked; notochord persistent.
- u. Eye visible through the skin; a few fang-like teeth on vomer and mandible.  
BARATHRONUS, 972.
- uu. Eye not visible; no teeth on maxillary or palatines; teeth on vomer rudimentary, those on mandible small.  
APHYONUS, 973.

## 953. BROTULA, Cuvier.

(BRÓTULAS.)

*Brotula*, CUVIER, Règne, Anim., Ed. 2, II, 296, 1829 (*barbata*).

Body elongate, compressed, covered with minute smooth scales; eye moderate; mouth medium, with villiform teeth on jaws, vomer, and palatines; lower jaw included; each jaw with 3 barbels on each side. Dorsal fin long and low, the dorsal and anal joined to the caudal. Ventral fins each reduced to a single filament of 1 ray. Eight branchiostegals. Air bladder large, with 2 horns posteriorly. One pyloric caeca. Vertebrae  $16 + 39 = 55$ . Tropical seas, in water of moderate depth. (*Brótula*, Spanish name of *Brotula barbata*.)

## 2870. BROTULA BARBATA (Bloch &amp; Schneider).

(BRÓTULA.)

Head  $4\frac{1}{2}$ ; depth about 5. D. 123; A. 93; V. 1. Upper jaw the longer. Ventral fin  $\frac{1}{2}$  as long as head. Dorsal commencing behind vertical from root of pectoral; vertical fins covered with thick skin. Color nearly uniform brown. Length 12 to 18 inches. West Indies, rare; in water of moderate depth. One specimen obtained by us in the market of Havana. (*barbatus*, bearded.)

*Brótula*, PARRA, Dif. Piezas Hist. Nat., 70, lam. 31, fig. 2, 1780, Havana.*Enchelyopus barbatus*, BLOCH & SCHNEIDER, Syst. Ichth., 52, 1801; after PARRA.*Brotula barbata*, CUVIER, Règne Anim., Ed. 2, II, 296, 1829; POEY, Memorias, II, 102, lam. 9, fig. 2, 1860; GÜNTHER, Cat., IV, 371, 1862.

## 954. STYGICOLA, Gill.

*Stygicola*, GILL, Proc. Ac. Nat. Sci. Phila. 1863, 252 (*dentatus*).

This genus differs from *Lucifuga* in the presence of palatine teeth. The teeth in the jaws are larger. As in *Lucifuga*, the single known species inhabits cave streams in Cuba. (*στυγῆ*, *Styx*, the river of the lower regions; *colo*, I inhabit.)

## 2871. STYGICOLA DENTATUS (Poey).

Head  $2\frac{3}{4}$ ; depth  $3\frac{3}{4}$ . D. 90; A. 70; P. 17; V. 1. Vertebrae  $11 + 37 = 48$ . Eyes usually wanting, occasionally represented by a rudiment; head elevated at the nape, the general form less slender than in *Lucifuga*, the belly more prominent; no scales on the nape; strong teeth, well separated, on the palatines as well as the vomer; teeth in the jaws larger than in *Lucifuga*; posterior with a large apophysis. Color translucent violet, with darker areas on nape and throat. Cases of the province of San Antonio, in southern Cuba. Largest specimen 5 inches long. (Poey.) (*dentatus*, toothed.)

*Lucifuga dentatus*, POEY, Memorias, II, 102, 1860, Cave of Cajfo (Coll. Noda), Cave of Castle La Industria (Coll. Dubrocá), Cave of Ashton (Coll. Fabre); GÜNTHER, Cat., IV, 373, 1862.*Stygicola dentata*, GILL, Proc. Ac. Nat. Sci. Phila. 1863, 252.

955. LUCIFUGA, Poey.

(CUBAN BLINDFISH.)

*Lucifuga*, POEY, Memorias, II, 95, 1860 (*subterraneus*).

Body moderately elongate, translucent pinkish, covered with minute scales. Eye rudimentary, covered by the skin; bands of villiform teeth in the jaws and vomer, none on the palatines; nostrils 2 on each side; no barbels; head with small tactile cirri; no spines on head; gills 4; no pseudobranchiæ; gill opening large, extending forward nearly to the symphysis, the gill membranes not united; branchiostegals 7 or 8; vertical fins low, united around the tail; ventrals each reduced to a short thin filament; male with an anal papilla, no pyloric caeca; air bladder large, rounded behind, joined to the base of the skull. Cave streams of Cuba; the eyes having undergone a degeneration similar to that seen in *Amblyopsis*. These fishes have no relation to the blind cave fishes of North America, but are derived from marine types, their ancestors being evidently allies of *Ogilbia* and *Brotula*. It is known that blindfishes are found also in caves of the islands of Jamaica, but no specimens have been seen by naturalists. (*lux*, light; *fugo*, I flee.)

2372. LUCIFUGA SUBTERRANEUS, Poey.

(PEZ CIEGO.)

Head  $2\frac{3}{4}$ ; depth  $3\frac{3}{4}$ . Branchiostegals 7. D. 70; A. 70; P. 51; V. 1; C. 9. Vertebrae  $11 + 36 = 47$ . Body elongate, compressed, tapering, pointed; head low at nape, much depressed anteriorly, broad, covered with soft, white, wrinkled skin, with microscopic cirri, having firm and conical tubes; no barbels on lips or chin; skin of head with many pores; scales not ciliate, present on body and top of head and on opercles; lateral line median, marked by a series of microscopic cirri like those on head, these wanting posteriorly; eyes wanting; nasal openings double; mouth large; lower jaw shorter; lips fleshy; maxillary broad at tip,  $\frac{2}{3}$  length of head; teeth in jaws very short and sharp, in a band; vomerine teeth larger; no palatine teeth; pharyngeal teeth slender; tongue smooth; gill openings large; males with an anal papilla; fin rays simple, flexible, jointed but not branched; dorsal beginning at a point about  $\frac{1}{2}$  nearer tip of snout than tip of caudal; anal smaller, beginning farther back, the 2 fins fully joined to the pointed caudal; pectoral fin short, falcate, nearly  $\frac{1}{2}$  head; ventral in front of pectoral, a slender ray not  $\frac{1}{2}$  length of maxillary. Color transparent rosy, head reddish, becoming darker in alcohol. No pyloric caeca; intestines short; air bladder large. Described from 12 specimens, the longest about  $4\frac{1}{2}$  inches; found in caves of the jurisdiction of San Antonio, in the southern part of Cuba. (Poey.) (*sub*, under; *terra*, earth.)

*Lucifuga subterraneus*, POEY, Memorias, II, 96, 1860, San Antonio, Cuba (Coll. D. Tranquillino); Sandalio de Noda (Coll. D. Juan Antonio Fabre); first coll. from Cajío Cave, 1831 (Noda); second, Cave at La Industria (Coll. Dubrocá); third, Ashton Cave, San Andreas (Coll. Fabre); fourth, Cave of the Dragon (Coll. Fabre); fifth, Cave at the Castle of Concord (Coll. Layunta).

## 956. BROSOMPHYCIS, Gill.

*Brosomphycis*, GILL, Proc. Ac. Nat. Sci. Phila. 1861, 168 (*marginatus*).

*Halias*, AYRES, Proc. Cal. Ac. Sci. 1861, 52 (*marginatus*); preoccupied.

Body elongate, moderately compressed; head unarmed; snout not long; teeth sharp, curved, in bands on jaws, vomer, and palatines; small cilia above snout and on anterior part of lower jaw. Body covered with thin cycloid scales; scales on head rudimentary or wanting. Caudal fin differentiated, entirely separated from the dorsal and anal; caudal peduncle slender. California. This genus is very close to *Ogilbia*, differing in the ciliated lips. Its species reaches a larger size. (*Brosomius*; *Phycis*.)

## 2873. BROSOMPHYCIS MARGINATUS (Ayres).

Head  $4\frac{1}{2}$  in body; depth  $6\frac{1}{2}$ . D. 92; A. 70; eye  $7\frac{1}{2}$  in head; snout  $4\frac{1}{2}$ ; maxillary 2; pectoral  $1\frac{1}{2}$ ; caudal 3; body elongate, moderately compressed; snout blunt; profile of head straight from snout; snout scarcely overhanging mouth; jaws subequal, the teeth conical, sharp, and slightly curved back, in bands on jaws, vomer, and palatines; maxillary reaching  $\frac{1}{2}$  the eye's diameter beyond eye; snout and lower jaw thickly covered with small cilia; head naked with the exception of small scales above; 2 large pores at tip of chin, a few large ones around preopercle and preorbital, 1 around gill opening, behind which is a pocket in the skin; about 3 short gill rakers developed below the angle of first arch, with many rough plates, not differentiated from those on the other arches. Dorsal and anal long and low, the rays embedded in the skin; tips of last rays each beyond the base of the caudal about  $\frac{1}{2}$  the length of caudal rays; distance of front of dorsal from snout  $3\frac{1}{2}$  in length of body; origin of anal a little nearer base of caudal than tip of snout; pectoral reaching about half way from its base to front of anal; ventrals developed as long filaments; caudal slender and rounded behind. Color bright reddish brown; fins edged with bright rose-red. Coast of California, in water of moderate depth; rare. Here described from a specimen, 12 inches in length, collected off San Francisco by Mr. W. G. W. Harford. (*marginatus*, edged.)

*Brosomius marginatus*, AYRES, Proc. Cal. Ac. Nat. Sci., 1, 1854, 13, San Francisco (Coll. W. O. Ayres); GIRARD, Pac. R. R. Surv., X, Fishes, 141, 1858.

*Brosomphycis marginatus*, GILL, Proc. Ac. Nat. Sci. Phila. 1861, 168; GILL, Proc. Ac. Nat. Sci. Phila. 1862, 280.

*Halias marginatus*, AYRES, Proc. Cal. Ac. Nat. Sci., pt. 2, 1861, 52.

*Dinematichthys marginatus*, GÜNTHER, Cat. Fishes Brit. Mus., IV, 375; JORDAN & GILBERT, Synopsis, 796, 1883.

## 957. OGILBIA, Jordan &amp; Evermann.

*Ogilbia*, JORDAN & EVERMANN, in EVERMANN & KENDALL, Bull. U. S. Fish Comm. 1897 (February 9, 1898), 132 (*cayorum*).

Body moderately elongate, covered with minute, smooth, embedded scales; sides of head with similar scales; lateral line inconspicuous; opercle with a very small spine, preopercle unarmed; no strong hook on maxillary; no barbels nor cilia; teeth in jaws in bands, similar teeth on

vomer and palatines; caudal free from the dorsal and anal; lower lip without cirri; dorsal and anal rays covered by the skin, ventrals each reduced to a filament of 2 rays; anal papilla of the male without horny claspers. Small fishes of the tropical shores of America, living in rock pools and shallows among algae. This genus is closely allied to the East Indian genus *Dinematichthys*, differing in the absence of anal papilla and claspers and in the shorter vertical fins. (Named for J. Douglas Ogilby, the accomplished naturalist of the museum of Sydney, in recognition of his excellent work on the fishes of Australia.)

a. Snout very short, about 7 in head; eye small, 10 or 11 in head; scales small, obscure, snout very short,  $6\frac{1}{2}$  in head. VENTRALIS, 2874.

aa. Snout longer, about 4 in head; eye about  $8\frac{1}{2}$  in head; scales larger, distinct. CAYORUM, 2875.

2874. OGILBIA VENTRALIS (Gill).

Head  $4\frac{1}{2}$  in body; depth  $5\frac{1}{2}$ . D. 64; A. 50; scales about 100; eye 10 or 11 in head; maxillary 2; pectoral  $1\frac{1}{2}$ ; caudal  $1\frac{1}{2}$ . Body elongate, moderately compressed; snout blunt, the profile behind snout nearly straight to occiput; mouth large, the maxillary extending 2 or 3 times the eye's diameter behind eye; teeth small, in bands on jaws, vomer, and palatines; eye very small, nearer snout than posterior end of maxillary; no cilia on snout and chin; body apparently naked to the unaided eye; but body and top of the head covered with small scales, which can be seen by the aid of a lens. Origin of dorsal distant from tip of snout by a space contained  $3\frac{1}{2}$  times in body; front of anal about midway between tip of snout and base of caudal; tips of last dorsal and anal rays reaching about to the middle of caudal rays, but not connected; pectorals scarcely reaching midway between their base and the front of anal; ventrals filamentous; caudal slender and rounded behind. Color in spirits, light brown above, lighter below; fins all colorless; without distinct marking anywhere. Gulf of California; not rare in rock pools; several specimens, 2 to 4 inches in length taken by us at Mazatlan. Here described from a specimen, 2 inches in length, from La Paz Harbor, Lower California. (*ventralis*, pertaining to the belly.)

*Bromophycis ventralis*, GILL, Proc. Ac. Sci. Phila. 1863, 253, Cape San Lucas. (Coll. Xantus.)

*Dinematichthys ventralis*, JORDAN, Proc. Cal. Ac. Sci. 1895, 502, pl. 54.

2875. OGILBIA CAYORUM, Evermann & Kendall.

Head 4; depth  $4\frac{1}{2}$ ; eye  $8\frac{1}{2}$ ; snout 4. D. about 68; A. about 50; scales about 14-87-13; maxillary  $1\frac{1}{2}$ ; pectoral  $1\frac{1}{2}$ ; ventral  $1\frac{1}{2}$ ; caudal  $2\frac{1}{2}$ . Body moderately elongate, compressed; head moderate, snout blunt; mouth large, jaws subequal, maxillary extending beyond vertical of eye a distance nearly equal to length of snout; eye very small, high up, situated in anterior third of head; nostril small, close to eye; teeth small, in bands on jaws, vomer, and palatines; back elevated, strongly arched from snout to origin of dorsal fin, thence descending in a nearly straight line to base of caudal; ventral outline comparatively straight, slightly concave at front of anal. Dorsal and anal long and low, distinct from caudal, the

posterior rays longest, about  $3\frac{1}{2}$  in head, base of each scaled; distance from tip of snout to origin of dorsal about 3 in length of body; origin of anal under about twenty-second dorsal ray, equidistant between tip of snout and base of caudal; scales very small, embedded, but showing distinctly under a lens; cheek and opercles partially covered with minute, embedded scales; top of head naked; opercle with a large, flat, flexible spine on level with eye. No barbels, cilia, nor tubercles; 2 large mucous pores at symphysis of lower jaw and 2 on preorbitals near anterior edge on each side; a row of 5 or 6 pores on lower jaw and edge of preopercle. Color uniform pale olivaceous or light brown, finely punctate with minute brown specks. Key West. Only the type known, an example,  $2\frac{1}{2}$  inches long, seined on a shoal covered with algae at Key West. (Cayo Hueso, or Bone Key, the original Spanish name for the Island of Key West, whence the name *cayorum*, of the keys.)

*Ogilbia cayorum*, EVERMANN & KENDALL, Bull. U. S. Fish Comm. 1897 (Feb. 9, 1898), 132, pl. 9, fig. 14, Key West, Florida. (Type, No. 48792. Coll. Evermann & Kendall.)

#### 958. BYTHITES, Reinhardt.

*Bythites*, REINHARDT, Dansk. Vidensk. Selsk. Afhandl., VII, 1838, 178 (*fuscus*).

Body elongate, covered with minute scales. Head large, thick; mouth large; jaws equal; no barbel; bands of teeth in the jaws and on vomer and palatines. Branchiostegals 8; gill membranes united, free from the isthmus; eyes moderate. Lateral line interrupted. Vertical fins united; ventral fins reduced to simple filaments, each composed of 2 rays closely united. Air bladder large; 2 pyloric caeca. A thick, conical, anal papilla (in the male). Greenland. (*βυθίτης*, an animal of the depths, from *βυθίος*, the deep.)

#### 2876. BYTHITES FUSCUS, Reinhardt.

Head about 4; depth  $4\frac{1}{2}$ . Body somewhat compressed, lipariform; snout obtuse, naked, with minute cirri. Mandible long, curved, extending far behind vertical from posterior margin of orbit; eye small; scales moderate on body; lateral line complete, but interrupted over vent, the two parts slightly overlapping the same vertical; vertical fins confluent, enveloped in thick skin; pectorals broad, lanceolate, with broad base; ventrals filiform, reaching behind origin of pectoral, as long as pectoral and  $\frac{2}{3}$  as long as head; a conspicuous anal papilla in the male. The only known specimen, now in the museum at Copenhagen, was obtained in Greenland half a century ago. (Goode & Bean.) (*fuscus*, dusky.)

*Bythites fuscus*, REINHARDT, Dansk. Vidensk. Selsk. Afh., VII, 1838, 178, Greenland; GÜNTHER, Cat., IV, 375, 1863; JORDAN & GILBERT, Synopsis, 795, 1883; GOODE & BEAN, Oceanic Ichthyology, 316, 1896.

#### 959. CATÆTYX, Günther.

*Catætyx*, GÜNTHER, Challenger Report, XXII, 104, 1887 (*messieri*).

Body compressed, elongate, covered with very small and thin scales; lateral line indistinct, interrupted. Head oblong, with somewhat pointed snout, covered with very small scales, only the anterior part of the snout



naked; bones of the head rather firm, but with the muciferous system well developed, the canals having wide openings along the infraorbital, and on the lower limb of the preoperculum; eye rather small; nostrils far apart, the posterior in front of the eye and the anterior at the extremity of the snout; operculum with a spine behind; no other armature on the head; snout not swollen, but the upper jaw slightly overlapping the lower; barbels none; mouth wide; bands of villiform teeth in the jaws, on the vomer, and the palatine bones; a series of larger teeth along the sides of the lower jaw; tail not much attenuated; vertical fins confluent; ventrals close together, reduced to a pair of fine, simple filaments, and inserted somewhat behind the isthmus, below the middle of the operculum. Gills 4, with short, broad gill rakers and well developed laminae; pseudobranchiae none; branchiostegals 8; pyloric appendages. Deep seas. Two species known. (*καται*, at the bottom; *τύξις*, τυγχάνω, find.)

2877. CATETYX RUBRIROSTRIS, Gilbert.

Depth of body below origin of dorsal equals  $\frac{1}{2}$  distance from end of snout to vent, 7 in length; head 4; distance from snout to origin of dorsal  $3\frac{1}{2}$ ; from snout to vent  $2\frac{7}{8}$ ; maxillary extending beyond eye,  $2\frac{3}{8}$  in head; eye equaling snout,  $5\frac{1}{2}$ ; interorbital width 7; width of snout  $3\frac{1}{2}$ . Teeth in upper jaw in a narrow band, minute, compressed, narrowly triangular, none of them enlarged; in the lower jaw a still narrower band of similar teeth, the posterior row slightly enlarged and increasing a little in size on sides of jaw, where it is accompanied by a single series only of the smaller teeth; this lateral series is continued backward far beyond premaxillary band; teeth on vomer and palatines similar to those in sides of lower jaw, the former in a V-shaped patch, the latter in a long and very narrow band. Anterior nostril in a short tube at tip of snout, the posterior large, without tube, immediately in front of eye; system of mucous pores well developed but not conspicuous, the pores collapsing on account of the thinness of the skin covering head; large mucous tube below eye, extending around front of snout and opening by slit-like pores along edge of snout and lower margin of infraorbital flap, opening posteriorly by a vertical slit  $\frac{1}{2}$  as long as pupil, immediately above end of maxillary; another series of pores along mandible and at edge of expanded limb of preopercle; no other evident pores. Angle of preopercle much expanded, its width equaling diameter of pupil; a sharp, strong spine arising from anterior portion of opercle, the structure of the gill flap apparently like that of *Bassogigas stelliferoides*; a short, sharp spine directed backward immediately behind posterior nostril; no other spines on head; gill openings wide, continuing forward to below posterior margin of orbit, the membranes wholly free from the isthmus; gill rakers short but not very broad, about  $\frac{1}{2}$  length of pupil, only 2 or 3 developed immediately in front of angle of arch. Dorsal beginning over or slightly behind middle of pectorals, the distance from its origin to occiput equaling or somewhat exceeding distance of latter from tip of snout; dorsal and anal fully united to caudal. The caudal has a base of appreciable width, bearing about 12 close-set rays, which extend much beyond tips of last dorsal and

anal rays; origin of anal nearer snout than base of caudal; ventrals slender, each consisting of a single ray, inserted very near together, under anterior portion of opercle, their length about equaling that of maxillary; pectorals with about 23 rays, evenly rounded behind, their length  $1\frac{1}{2}$  in head. Scales very small, cycloid, regularly imbricated, in about 135 transverse series; nape and belly scaled, as is also the head, excepting snout, mandible, suborbital, and sometimes interorbital areas. Color dusky olive, the ventrals white, the other fins black, at least on distal portions; opercles, gill membranes, sides and top of snout, and posterior portion of abdomen blue black; snout flushed with dark ruby red in life; lining membrane of mouth and gill cavity, and peritoneum jet-black. Closely related to *C. messieri*, differing in the shorter, broader snout, the wider preopercle, the more anterior origin of anal, and apparently in the gill rakers and pores on head. Off coast of California. Four specimens known, the longest  $4\frac{1}{2}$  inches in length. (Gilbert.) (*ruber*, red; *rostrum*, snout.)

*Catatyx rubrirostris*, GILBERT, Proc. U. S. Nat. Mus. 1890, 111, off coast of California, at Albatross Stations 2909, 2925, and 2936, in 205 to 359 fathoms. (Type, No. 44379.)

#### 960. DICROMITA, Goode & Bean.

*Dicromita*, GOOD & BEAN, Oceanic Ichthyology, 319, 1896 (*agassizii*).

Brotulids resembling in form and general appearance *Catatyx* and *Diaplanacanthopoma*, having the lateral line obsolete, or interrupted posteriorly; ventrals a pair of simple, fine filaments, and with teeth upon the palatines. It has, however, 3 or 4 small spines upon the preoperculum, as well as a sharp spine upon the upper angle of the operculum; and the lateral line, though indistinct, is traceable for  $\frac{1}{2}$  or  $\frac{2}{3}$  the length of the body, which, like the upper part of the head, is covered with small, deciduous scales, the opercular region being apparently scaleless, and the boxes of the suborbital region almost uncovered, with conspicuous sinuses, which show through the transparent texture of the surface. Head oblong; snout somewhat produced, depressed, and turgid, resembling, though in a less degree, that of *Barathrodemus*. Eye moderate, conspicuous. Mouth wide; teeth villiform, in bands on the jaws and palatines, and very minute upon the vomer, which has a roughened, knob-like enlargement at its angle. Vent premedian. Ventral fins confluent; ventrals rooted very close together, each reduced to a fine, flexible, simple filament, planted somewhat behind the isthmus and below the middle of the operculum. Gills 4, with well-developed laminae and rather long, slender gill rakers. Branchiostegals 8. Pseudobranchiae apparently absent. (*δίκρος*, forked; *μίτρος*, thread.)

#### 2878. DICROMITA AGASSIZII, Goode & Bean.

Body elongate, much compressed, its height about  $\frac{1}{3}$  of its total length, its width about  $\frac{1}{2}$  its greatest height; head slightly greater than height of body, about twice its own width; mouth very large, the maxillary curved and much dilated at its extremity, reaching far behind the vertical from the posterior margin of orbit; jaws nearly equal, the snout considerably

produced and dilated, its length equal to diameter of eye and  $\frac{1}{2}$  length of head. Teeth very fine, villiform, in bands on jaws and palatines, and also present on vomer, though very small, especially upon the rounded, globular process of the angle. Lateral line very indistinct, interrupted, but extending behind the vent at least  $\frac{1}{2}$  of the way to tip of tail. Dorsal origin nearly in vertical from the axil of the pectoral; ventrals very slender, villiform, closely approximate at their roots, and less than  $\frac{1}{2}$  as long as the head. Color brownish. A specimen was obtained by the *Blake* off Granada, Station XCIII, at a depth of 291 fathoms. The collateral type was obtained by the *Albatross* at Station 2374, in lat.  $29^{\circ} 11' 30''$  N., long.  $85^{\circ} 29' W.$ , at a depth of 26 fathoms. (Goode & Bean.) (Named for Prof. Alexander Agassiz.)

*Dicromita agassizii*, GOODE & BEAN, Oceanic Ichthyology, 319, fig. 285, 1896, off Granada, in 291 fathoms (Coll. the *Blake*); Lat.  $29^{\circ} 11' 30''$  N., Long.  $85^{\circ} 29' W.$ , in 26 fathoms. (Coll. *Albatross*.) (Type in M. C. Z.)

#### 961. BASSOZETUS, Gill.

*Bathynectes*, GÜNTHER, Ann. and Mag. Nat. Hist., 11, 1878, 20 (*compressus*); name preoccupied in Crustacea.

*Bassozetus*, GILL, Proc. U. S. Nat. Mus., VI, 1883, 59 (*normalis*).

*Bathynus*, GOODE & BEAN, Proc. U. S. Nat. Mus., VIII, 1886, 663 (*catena*).

Body compressed, with long tapering tail, covered with deciduous thin scales of moderate size. Bones of the head very soft and cavernous, the upper opercular spine very feeble, ridge-like; no other armature of the head. Head scaly, except the snout, which is obtusely rounded off, with the jaws equal or nearly equal in front. Mouth very wide; bands of villiform teeth in the jaws, on the vomer, and palatine bones. Barbels none. Eye small; anterior nostril about midway between the posterior and the extremity of the snout. Vertical fins confluent; ventrals close together, reduced to a pair of simple filaments, and inserted below the rounded angle of the preoperculum. Gills 4, with short gill laminae, but with long stiff gill rakers on the first branchial arch. Pseudobranchiae none. Branchiostegals 8. Pyloric appendages none. ( $\beta\acute{\alpha}\sigma\sigma\omega\nu$ , for  $\beta\alpha\theta\acute{\upsilon}\varsigma$ , deep;  $\zeta\eta\rho\acute{\epsilon}\omega\nu$ , seek.)

a. Body moderately elongate, the depth 9 to 10 in length; dorsal rays 116; anal 92 to 96.

b. Head 6 in length.

NORMALIS, 2879.

bb. Head about  $7\frac{1}{2}$  in length.

COMPRESSUS, 2880.

aa. Body more elongate, the depth  $12\frac{1}{2}$  in length; head 83; head with conspicuous, chain-like rows of pores.

CATENA, 2881.

aaa. Body excessively attenuate, the depth more than 16 times in length; dorsal rays 138; anal 115; distance from snout to vent nearly 4 times in body.

TEENIA, 2882.

#### 2879. BASSOZETUS NORMALIS, Gill.

Head 6 in total length; eye 4 in snout. D. 116; A. 96. Body much compressed, its width in the region of vent not more than  $\frac{1}{2}$  of its height, which at the same point is about  $\frac{1}{3}$  of the total length; greatest height of body, over the origin of the pectorals, about  $\frac{2}{3}$  the distance from base of pectorals to vent, the vent being about twice as distant from base of

caudal rays as from snout. Head moderately compressed, flat above; snout obtuse rounded, turgid; lower jaw considerably included. Bones of head not completely ossified, very cavernous in the alcoholic specimen, the head showing many deep sinuosities and depressions. Eye very small, situated about midway between the tip of the snout and the vertical from the posterior end of maxillary. Teeth all small and short, densely set, forming narrow, villiform bands; vomerine band open V-shaped. Dorsal fin beginning far in advance of origin of pectoral and above upper angle of gill opening, rays longest in region of vent; anal beginning immediately behind vent, its rays not quite so long as those of dorsal; pectoral with broad base, short, not exceeding much more than halfway to vertical from vent, its length considerably less than that of postorbital portion of head. Ventral rays very slender, villiform, reaching almost to vent, far beyond pectoral, their length almost equal to that of head. Scales moderate, very deciduous, extending upon cheeks and on top of head almost to tip of snout; no evidence of a lateral line. Color light, the head and abdomen blackish; inside of mouth purplish brown. The *Blake* secured specimens from Station CCIV, in lat.  $24^{\circ} 33' N.$ , at a depth of 1,920 fathoms, and from Station LXXXIV, off Dominica, in 1,131 fathoms. The *Albatross* also obtained examples (No. 49416, U. S. Nat. Mus.) from Station 2380, in Lat.  $28^{\circ} 02' 30'' N.$ , Long.  $87^{\circ} 43' 45'' W.$ , at a depth of 1,430 fathoms; (No. 33306, U. S. Nat. Mus.) from Station 2042, in lat.  $39^{\circ} 33' N.$ , Long.  $68^{\circ} 26' 45'' W.$ , at a depth of 1,555 fathoms. (Goode & Bean.) West Indies, Gulf of Mexico, and to lat.  $40^{\circ} N.$ , in region of the Gulf Stream. (*normalis*, normal.)

*Bassozetus normalis*, GILL, Proc. U. S. Nat. Mus. 1883, 259, Lat.  $39^{\circ} 33' N.$ , Long.  $68^{\circ} 26' 45'' W.$ , in 1,555 fathoms (Type, No. 33306. Coll. *Albatross*); GOODE & BEAN, Oceanic Ichthyology, 322, fig. 287, 1896.

#### 2880. BASSOZETUS COMPRESSUS (Günther).

D. 116; A. 92; P. 23; V. 1. The greatest depth of the body is above the end of the gill cover and about  $\frac{1}{2}$  length of trunk; vent twice as distant from extremity of tail as from snout, consequently the tail is more moderately attenuated. Head compressed like the body, and about  $\frac{3}{4}$  length of trunk; superficial bones form large muciferous cavities which, when full, must give to the head a much more evenly rounded appearance than in the preserved state, when the supporting bony ridges project more or less from under the skin. Snout slightly swollen, but the jaws nearly even in front, the wide mouth slightly ascending forward; maxillary with the form usual in these Gadoid fishes, dilated behind, and extending far behind the eye. Eye very small,  $\frac{1}{2}$  length of snout, and  $\frac{1}{4}$  that of head, placed high up on the side, and not possessing an orbital fold of integument; interorbital space rather convex and equal in width to 3 diameters of eye. Teeth all very small, short, densely set, and forming villiform bands, the broadest on maxillary bone and quite uncovered on the sides, no labial folds being developed; palatine band broader than the mandibular, and the vomerine band V-shaped, each arm being bent with the convexity inward. Gill opening and cavity very wide and of an intense black; gill rakers much longer than the laminae, 15 in number on

the anterior arch, besides some rudimentary ones above. Dorsal fin commencing above upper end of gill opening, with short rays partly hidden in the skin, becoming longer in middle of fin, but remaining of moderate length; anal shorter; pectoral with a rather narrow base, quite free, and composed of feeble rays, its length only  $\frac{1}{2}$  that of head; ventral rays very feeble, reaching somewhat beyond the root of pectoral. In the specimens examined only very few of the thin, cycloid scales have been preserved; they are of moderate size, there being about 16 in a transverse series running from the vent to the dorsal fin; the lateral line, if it was developed, can no longer be traced. Blackish, with the fins, head, and abdomen black. Specimens of this very fine and truly bathybial fish were obtained at great depths on the southeast of New Guinea, off the Philippine islands, and in the mid-Atlantic; the exact localities being 75 miles east-southeast of Raine Island, Station 184, depth 1,400 fathoms; two specimens, 17 and  $4\frac{1}{2}$  inches long. Philippine Islands, Station 205, depth 1,050 fathoms; one specimen,  $5\frac{1}{2}$  inches long. Mid-Atlantic, Station 107, depth 1,500 fathoms; one specimen,  $5\frac{1}{2}$  inches long. The young are extremely similar to the old, but have a larger eye, which is  $\frac{1}{3}$  of length of head. The specimen from Station 205 (Philippine Islands) has longer ventral filaments, extending nearly to the vent. (Günther.) (*compressus*, compressed.)

*Bathynectes compressus*, GÜNTHER, Ann. Mag. Nat. Hist., II, 1878, 20, Challenger Station 107, mid-Atlantic, in 1,500 fathoms. (Coll. Challenger.)

*Bathyonus compressus*, GÜNTHER, Challenger Report, XXII, 109, 1887.

*Bassozetus compressus*, GOODE & BEAN, Oceanic Ichthyology, 322, 1896.

2881. BASSOZETUS CATENA, Goode & Bean.

Head  $8\frac{3}{4}$ ; depth  $1\frac{1}{2}$  in head or  $12\frac{1}{2}$  in body; eye 5; snout 5; interorbital width 5. Body very elongate, much compressed, and tapering into a slender, whip-like tail. Head without spines, vertex convex, not much compressed, higher than body. Interorbital area somewhat convex. The muciferous channel upon the infraorbital ring shows in its course several wide subcircular sinuses, closely approximated; a similar row upon the posterior edge of the preoperculum and continued forward upon the under surface of the mandible; the vertex also has a semicircle of similar sinuses. Maxillary extending beyond vertical through posterior margin of orbit, its length equal to that of postorbital part of head; mandible  $\frac{2}{3}$  as long as head and equal in length to height of body; jaws, vomer, and palate with bands of villiform teeth, the vomerine band V-shaped. Nostrils in front of middle of eye, separated by a slight interspace, the anterior nearer to its mate than to tip of snout. Branchiostegals 8; pseudobranchiae absent. Gill rakers long and numerous, the longest slightly exceeding diameter of eye, 15 developed below angle of first arch, besides several rudiments; dorsal origin slightly behind that of pectoral, its distance from tip of snout about  $7\frac{1}{2}$  in total, rays well developed; in the anterior  $\frac{1}{3}$  of the fin, in a space equal to length of head, were counted 20 rays, the longest of which is  $\frac{2}{3}$  as long as head; anal origin under twenty-first dorsal ray, its rays shorter than those of dorsal; pectoral extending to vertical from eighteenth ray of dorsal,  $\frac{1}{2}$  as long as head; ventrals composed each

of a simple filament, the origin slightly in advance of vertical through pectoral origin, the length  $\frac{3}{4}$  that of head, not reaching nearly to vent, the distance of which from origin of ventrals is slightly greater than length of head. Color brownish yellow; head and abdomen blackish. Gulf of Mexico, in great depths. Only the type known, 237 mm. long. (*catena*, chain, from the arrangement of the mucous cavities on the head.)

*Bassozetus catena*, GOODE & BEAN, Proc. U. S. Nat. Mus. 1885, 603, Lat. 28° 00' 15" N., Long. 87° 42' W., in 1,467 fathoms (Type, No. 37341. Coll. *Albatross*); GÜNTHER, Challenger Report, XXII, 111, 1887; GOODE & BEAN, Oceanic Ichthyology, 323, fig. 286, 1896.

2882. BASSOZETUS TENIA (Günther).

D. 138; A. 115; P. 30; V. 1. The greatest depth of the body is below the origin of the dorsal fin and about  $\frac{1}{3}$  of the length of the trunk, the vent being not quite thrice as distant from the extremity of the tail as from the snout; therefore, the whole of the fish, and especially the tail, is much attenuated. Head not compressed, low and long, forming  $\frac{2}{3}$  length of trunk. Structure of the bones of the head as in *B. compressus*. Snout rather swollen and broad, the upper jaw but slightly overlapping the lower; maxillary extending far behind the eye, which is very small,  $\frac{1}{4}$  length of snout, about  $\frac{1}{4}$  that of head, and  $\frac{1}{4}$  width of interorbital space. Teeth very small and short, densely set, forming narrow, villiform bands; vomerine bands open A-shaped. Gill cavity deep black; gill rakers long and slender, 16 in number, with some rudimentary ones in front and behind. Dorsal fin commencing above upper end of gill opening, with short rays partly hidden in the skin, the rays becoming longer on the anterior third of tail, but remaining of moderate length, the anal rays still shorter; pectoral with a broad base, quite free, and composed of rather feeble rays, its length equal to that of postorbital portion of head; ventral rays very feeble, reaching nearly to the middle of the pectoral. The scales must have been extremely thin and rather small; there were probably about 20 in a transverse series running from the vent to the dorsal fin. The lateral line can not be made out. Light colored (possibly pink in life), with the head and abdomen black. Only 1 specimen known of this eminently bathybial fish, obtained in mid-Atlantic (*Challenger* Station 104) at a depth of 2,500 fathoms. Its total length is 10 inches. (*ταυρία*, ribbon.)

*Bathyonus tenia*, GÜNTHER, Challenger Report, XXII, 110, 1887, pl. 23, fig. A, mid-Atlantic, Station 104, at a depth of 2,500 fathoms; GOODE & BEAN, Oceanic Ichth., 323, 1896.

962. MCEBIA, Goode & Bean.

*Mobia*, GOODE & BEAN, Oceanic Ichthyology, 331, 1896 (*gracilis*).

Brotulids resembling *Bassozetus* in general form, excepting that the tail is prolonged in a very slender filament, the dorsal and anal rays being extremely short posteriorly, but positively confluent with the caudal rays, which are much longer and much exerted; ventrals each bifid, instead of a single ray, as in *Bassozetus*. Head very cavernous, the sinuses large and conspicuous on the infraorbital ring, on the mandible, and the pre-

operculum. A single, short, feeble spine on the shoulder, but none upon the operculum or preoperculum, though certain projections seem to show above the eye, doubtless due to the shrinkage of the integument upon the underlying projections of the bone. Mouth very wide, the extremity of the maxillary much dilated; posterior nostrils very wide and separated from the eye by a small, spinous projection of bone; teeth in narrow bands, that on the vomer V-shaped, with the 2 arms straight. A few large scales in a row starting from the upper angle of the gill opening and terminating over the axle of the pectoral. Gill rakers on outer arch rather numerous, long and slender. Pseudobranchiae represented by 2 minute globules. Deep sea; 2 species known. ("Named in honor of Prof. Karl Möbius, director of the Royal Zoological Museum in Berlin, who has added much to our knowledge of marine life by his noble work, Die Fauna der Kielerbucht, and by numerous other writings.")

2883. *MÆBIA PROMELAS* (Gilbert).

Head 2 in trunk; depth 3. Body  $3\frac{1}{2}$  to  $3\frac{3}{4}$  in tail. Tail produced into a filament, the caudal basis extremely narrow, supporting 5 long slender rays which are firmly bound together. Mouth terminal, large, the maxillary much dilated at tip, reaching well behind the eye,  $1\frac{1}{2}$  in head; lower jaw included, the tip slightly produced. Teeth in villiform bands on mandible, premaxillary, vomer, and palatines, the band on mandible very narrow, that on vomer with the diverging arms much incurved, the anterior angle rounded. Tongue toothless, some of the basibranchials forming a sharply elevated denticerous crest. Gill laminae extremely narrow, the gill rakers of outer arch very long and slender, 1 (with 4 rudiments) above angle, 15 below. Infraorbital chain with 6 mucous sinuses, the mandible with 5, preopercle with 5, and a number on top of head; these are all bridged over with very delicate membrane which is easily ruptured. A row of low, strong spinous points directed posteriorly on the ridge running backward from the eye; no other spines on head, though a number of short spinous points are made evident when the skin is removed; opercular spine rather weak. A distinct membranaceous flap runs along the projecting edge of shoulder girdle, connecting pectorals with upper end of gill flap; pectorals slender, equaling postorbital part of head; ventrals each of a bifid filament, the two branches joined at the base for a very short distance, variable in length, reaching to or nearly to tips of pectorals, usually contained about  $1\frac{1}{2}$  times in head; dorsal beginning a trifle behind base of pectorals, its distance from occiput equaling distance of latter from front of eye. Scales very small, apparently covering a part of top of head; 3 series of large pores on sides; 1 from upper end of gill slit backward parallel with dorsal outline; a second along middle of sides; the third beginning halfway between base of pectorals and ventral outline, extending backward on belly and along base of anal fin, these lines all somewhat indistinct, and it can not be determined how far they extend backward. Color light brown; head (except occiput), mouth, gill cavity, and abdomen jet-black; fins dusky. This species closely resembles *Mæbia gracilis* (Günther), from New Guinea, dif-

fering in the following respects: Depth 3 in trunk (in *gracilis*  $3\frac{1}{2}$ ); eye 6 in head (in *gracilis*  $5\frac{1}{2}$ ); vomerine patch of teeth with the two arms incurved (U-shaped in *gracilis*); dorsal fin beginning behind the pectoral (over root of pectoral in *gracilis*); an additional series of large scales (lateral line) along middle of sides, and another along ventral outline; trunk  $2\frac{1}{2}$  in tail ( $2\frac{3}{8}$  in *gracilis*); ventrals shorter, not reaching past tips of pectorals. Five specimens, the largest  $9\frac{1}{2}$  inches long, from *Albatross* Station 3010, at a depth of 1,005 fathoms, in the Gulf of California. (Gilbert.) (*πρό*, before; *μέλας*, black.)

*Porogadus promelas*, GILBERT, Proc. U. S. Nat. Mus. 1891, 547, Gulf of California, in 1,005 fathoms. (Coll. *Albatross*.)

### 963. NEOBYTHITES, Goode & Bean.

*Neobythites*, GOODE & BEAN, Proc. U. S. Nat. Mus. 1885, 600 (*gilli*).

Brotulids having the body elongate, compressed, covered with small scales, and the head also scaled; lateral line incomplete, obsolete posteriorly. Eye moderate; snout moderate, rounded, slightly produced, the lower jaw slightly included; no barbel. Teeth villiform, in narrow bands in jaws and palatines; vomerine teeth in a V-shaped patch; 2 weak spines at angle of preoperculum, and a stronger one at angle of operculum. Gill openings wide, the membranes deeply cleft and not attached to the isthmus; vertical fins united; ventrals reduced each to a bifid ray. Branchiostegals 8. Pseudobranchiæ present, but small. Air bladder present. (*νέος*, new; *Bythites*.)

a. Scales about 88 in longitudinal series; depth  $4\frac{1}{2}$  in length.

GILLI, 2884.

aa. Scales about 123 in longitudinal series; depth  $5\frac{1}{2}$  in length; dorsal rays 101.

MARGINATUS, 2885.

### 2884. NEOBYTHITES GILLI, Goode & Bean.

Body compressed, its height contained  $4\frac{2}{3}$  times in total length, and less than length of head; interorbital area convex, its width equal to diameter of circular eye,  $3\frac{2}{3}$  in length of head, and  $1\frac{1}{2}$  in length of snout in young. Head compressed, deeper than broad, with wide sinuses, its length contained  $4\frac{1}{2}$  times in that of the body; snout obtusely rounded, slightly produced; mouth large, the maxillary extending considerably behind the vertical through posterior margin of eye, expanded posteriorly; mandible still longer, its length about  $2\frac{1}{2}$  times in height of body; interorbital space convex. Teeth in villiform bands in jaws and on palatines; vomerine patch subcircular, with angles extended posteriorly. Gill rakers moderately long and slender, somewhat numerous, the longest about  $\frac{2}{3}$  diameter of eye, 11 developed and 3 rudiments below the angle. Pseudobranchiæ absent; gill opening wide, the membrane deeply cleft, free from the isthmus behind. A single long, flat spine attached to posterior portion of operculum, high up, extending back to its edge; a small hidden spine at lower angle of preoperculum. Nostrils small, the anterior one in a very short tube, almost upon tip of snout; posterior nostril slightly larger, not tubular, immediately in front of middle of eye. Scales mod-



crate, upon head and body, in 88 vertical rows, 7 rows between dorsal origin and lateral line, which becomes obsolete in its posterior half, 16 or 17 from vent forward to lateral line; dorsal origin behind that of ventral and pectoral, its distance from snout contained 4 times in total length, its rays moderately long; anal origin under eighteenth dorsal ray, its distance from snout contained  $2\frac{1}{2}$  times in body length, rays rather slenderer than those in the dorsal; caudal rays 6 or 7 in number, their length contained 9 times in total length, not differentiated from those of the adjacent fins; pectoral origin well forward, its base somewhat concealed by the flap of the operculum, its length about equal to  $\frac{2}{3}$  that of head; ventrals each a bifid ray, the inner filament the longer, inserted slightly in advance of the base of the pectoral, not far from humeral symphysis, and reaching nearly to vent, its length nearly equal to height of body; distance from origin of ventral to vent slightly greater than height of body; color light yellow, with silvery reflections, with cloudings of brown above lateral line and numerous black chromatophores; a series of irregular brown blotches above the lateral line, with 1 or 2 much darker, extending upon the dorsal fin. In many specimens the color is uniform yellow, with simply the dark ocelli showing. (Goode & Bean.) Atlantic, in rather deep water, from Gulf Stream to the coast of Brazil. (Named for Dr. Theodore Gill.)

*Neobythites gillii*, GOODE & BEAN, Proc. U. S. Nat. Mus. 1885, 601, Lat.  $28^{\circ} 36' N.$ , Long.  $85^{\circ} 33' W.$ , in 111 fathoms (Type, No. 37340. Coll. *Albatross*); GÜNTHER, Challenger Report, XXII, 103, 1887; GOODE & BEAN, Oceanic Ichth., 325, fig. 289, 1896.

*Neobythites ocellatus*, GÜNTHER, Challenger Report, XXII, 103, pl. 21, fig. B, 1887, off Pernambuco, in 350 fathoms.

2885. NEOBYTHITES MARGINATUS, Goode & Bean.

Head  $4\frac{1}{2}$  in total length; depth  $5\frac{1}{2}$ . D. 101; scales 7-123-29. Body compressed, somewhat elongate; interorbital area convex, its width greater than the diameter of the circular eye. Mouth large, the maxillary extending considerably behind vertical through posterior margin of orbit, its length 2 in head; mandible slightly more than  $\frac{2}{3}$  height of body. Teeth as in *N. gillii*. Gill rakers slightly longer than  $\frac{1}{2}$  the diameter of eye, 7 and 3 rudiments below the angle of the anterior arch. Pseudobranchiae absent. A long flat spine upon the upper edge of the operculum, extending back nearly to its margin; 2 short, flat spines upon the angle of the preoperculum. Nostrils as in *N. gillii*. Scales small, very closely imbricated, the lateral line obsolete in its posterior half. Distance of dorsal origin from snout 4 times in total length; anal origin under fourteenth dorsal ray, at a distance from the snout  $2\frac{2}{3}$  times in total length. Caudal of about 8 or 9 rays, very closely placed, about  $10\frac{1}{2}$  times in total length; pectoral placed much as in *Benthocometes*, its length about  $2\frac{1}{2}$  times that of the head, extending to vertical through the vent; ventral a bifid ray inserted in advance of base of pectoral, not reaching to the vent, its length considerably less than height of body; distance of ventral origin from vent slightly more than height of body. Color light yellowish brown, an obscure narrow band of darker brown commencing on the snout, inter-

rupted by the eye, and extending backward  $\frac{2}{3}$  distance to tail; another beginning on the snout, extending over eye and back as far as first described, interrupted posteriorly; dorsal fin milky white at base in its anterior third; above this a blackish band extending whole length of fin; a narrow white margin above. The type is from Blake Station LXXIX, off Barbados, in 209 fathoms. (*marginatus*, edged.)

*Neobythites marginatus*, GOODE & BEAN, Bull. Mus. Comp. Zool., x, No. 5, 162, 1883 off Barbados, in 209 fathoms (Coll. Blake); GOODE & BEAN, Oceanic Ichthyology, 326, fig. 290, 1896.

#### 964. BENTHOCOMETES, Goode & Bean.

*Benthocometes*, GOODE & BEAN, Oceanic Ichthyology, 327, 1896 (*robustus*).

Brotulids, similar in appearance and structure to *Neobythites* and *Bassogigas*, distinguished by 2 short flat spines upon the anterior portion of the operculum, placed at some distance from each other, and by the absence of spines on the preoperculum. The lateral line is complete, and extends without interruption to the posterior fourth of the body, where it becomes obsolete. The vomerine teeth are bunched in a circular patch instead of being arranged in triangular form. The head is comparatively short, with the jaws in front nearly equal; the snout not produced, but obtuse, rounded, and almost declivous in its outline. Deep sea. Two species known. (*βένθος*, the depths; *κατοικήτης*, inhabitant.)

#### 2886. BENTHOCOMETES ROBUSTUS, Goode & Bean.

Body rather short and deep, its greatest height nearly  $4\frac{1}{2}$  in total length and about equal to length of head; interorbital area convex, its width greater than diameter of the circular eye, and  $1\frac{1}{2}$  times length of snout; head about 4 times diameter of eye; mouth moderate, the maxillary extending to vertical through posterior margin of eye, the mandible a little beyond, its length equal to that of postorbital part of head. Teeth in villiform bands in jaws and on palatines; vomerine teeth bunched in a circular patch. Gill rakers moderate, the longest a little more than twice in diameter of eye, 4 above angle of first arch, 11 below. Pseudobranchiae rudimentary. Gill opening wide, the membrane deeply cleft behind, free from the isthmus. A pair of short flat spines upon the anterior portion of the operculum. Nostrils small, the anterior as close to the snout as the posterior ones are to the eyes; no apparent cirri. Scales minute; lateral line obsolete on the last fourth of body. Dorsal origin behind that of ventral and pectoral, its distance from snout  $3\frac{2}{3}$  times in body; height of dorsal fin moderate, the longest ray about 3 times in head; anal origin under eighteenth ray of dorsal, the height of fin about equaling that of dorsal; vertical fins not connate with the caudal, which consists of 12 or 13 very slender rays, its length nearly equal to  $\frac{1}{2}$  head; pectoral with a broad base, close to gill opening, its length nearly  $\frac{2}{3}$  that of head; ventral a single bifid ray, inserted in advance of vertical through base of pectorals, and not far from humeral symphysis, reaching nearly halfway to vent, the distance of which from the origin of the ven-

tral is equal to length of head. Color yellowish brown. The type of this species, a specimen 88 mm. long, was taken by the *Blake* from Station XCIV, off Moro Castle, Cuba, at a depth of from 250 to 400 fathoms. A collateral type specimen (No. 29057) was obtained by the *Fish Hawk* from Station 1043 in Lat. 38° 39' N., Long. 73° 11' W., at a depth of 130 fathoms. (Goode & Bean.) West Indies, to lat. 39° N., in Gulf stream. (*robustus*, stout.)

*Neobythites robustus*, GOODE & BEAN, Bull. Mus. Comp. Zool., X, No. 5, 161, 1883, off Moro Castle, Cuba, in from 250 to 400 fathoms. (Type in M. C. Z. Coll. *Blake*.)

*Benthocometes robustus*, GOODE & BEAN, Oceanic Ichthyology, 327, fig. 288, 1896.

### 965. BASSOGIGAS, Gill.

*Bassogigas*, GILL MS. in GOODE & BEAN, Oceanic Ichthyology, 328, 1896 (*gillii*).

Brotulids having the body elongate, compressed, covered with a thick, heavy skin, which upon the head covers and obscures all the angles of the skull; scales small, covering body and head completely; lateral line indistinct for the greater part of the course, but apparently extending at least  $\frac{3}{4}$  of the way from the operculum to the tail; eye moderate; vertical fins completely united; ventrals a pair of bifid filaments inserted behind the humeral symphysis and remote at their bases, short, rather stout; snout without barbels, slightly produced, the lower jaw being barely included; villiform teeth in the jaws, on the vomer and palatines; vomerine patch V-shaped, but with its arms broadly expanded and thicker at the angle, so that it is almost triangular; operculum with a long, sharp spine; preoperculum unarmed; branchiostegals 8; air bladder present; pseudo-branchiæ small. Deep sea. (*βάσσων*, for *βαθύς*, deep; *γίγας*, giant.)

a. Dorsal fin with 83 rays; anal 67.

GILLII, 2887.

aa. Dorsal fin with 95 rays; anal 82.

STELLIFEROIDES, 2888.

### 2887. BASSOGIGAS GILLII, Goode & Bean.

D. 83; C. 6; A. 67. Head rather short and broad, with snout slightly overlapping the lower jaw; diameter of the eye scarcely  $\frac{1}{3}$  of the length of the snout and about  $\frac{1}{2}$  that of head; maxillary extending far behind eye, the vertical from the anterior margin of orbit nearly bisecting it, its length  $\frac{1}{2}$  that of head, and its posterior margin ending in a broad triangular dilation; teeth normal; anterior and posterior nostrils separated by a space greater than diameter of eye; preoperculum with a square, rounded angle; no armature; operculum with a strong, sharp spine above, the tip of which projects slightly beyond the opercular flap; distance of vent from root of pectoral slightly more than length of head, as far removed from this point as is the anterior nostril; scales moderate, covering the entire head; lateral line somewhat conspicuous, obsolete in its posterior third. Dorsal and anal fins enveloped in thick scaly skin; origin of dorsal in advance of middle of pectoral; pectorals rounded, broad, and very short; less than  $\frac{1}{2}$  as long as head and extending about  $\frac{1}{2}$  distance from origin to vertical from vent; ventrals inserted somewhat behind angle of preoperculum, extending to vertical from axil of pectoral, and about  $\frac{1}{2}$  of distance from origin to vent; each ventral filament bifid, the inner

part being the longer. Color uniform grayish brown; fins darker. The type of this species was obtained by the *Albatross* from Station 2684, off Cape Henlopen, Delaware, in Lat. 39° 35' N., Long. 70° 54' W., at a depth of 1,106 fathoms. (Goode & Bean.) (Named for Dr. Theodore Gill.)

*Bassogigas gillii*, GOODE & BEAN, Oceanic Ichthyology, 328, fig. 291, 1896, off Cape Henlopen, Delaware, in 1,106 fathoms. (Type, No. 39417. Coll. *Albatross*.)

288°. *BASSOGIGAS STELLIFEROIDES* (Gilbert).

Head 4 to 4½ in length; depth 5 to 5½. D. 95; A. 82; scales 110. Physiognomy strikingly like that of the Sciaenoid genus *Stellifer*. Mouth large, oblique, the lower jaw included, maxillary reaching well beyond orbit, ¼ length of head. Teeth uniform, small, in narrow bands, those on vomer in a ∩-shaped patch; a well-developed band on palatines; tongue smooth, a well-developed dentigerous crest on median line behind it; no barbel at symphysis. Snout short, bluntly rounded, about equaling diameter of orbit, slightly overhanging mouth, 5 in head; interorbital width 4; upper limb of preopercle extending obliquely downward and backward, largely adnate, the angle produced into a free membranaceous flap which entirely conceals the narrow interopercle, and bears no spines. The structure of the gill flap does not appear to have been correctly interpreted. The opercle is strong, but of small extent, forking at its base, 1 branch continued straight backward as a strong spine, the second a narrow flat process downward and somewhat backward, parallel with and little distant from margin of preopercle. Filling the deep notch between these 2 processes, and forming the greater portion of the gill flap, is the thin membranaceous subopercle. Branchiostegal rays 7. Gill rakers long and slender, the longest ¾ diameter of orbit, 7 above angle, 13 and about 5 rudiments below. Nape midway between front of dorsal and front of eye; dorsal and anal similar, uniform, low, joined to base of caudal, the latter truncate, projecting well beyond them; ventrals inserted under angle of preopercle, each of a single ray forked to the very base, the 2 branches united by membrane for a distance equaling ¾ orbit, the inner filaments being longest, ½ longer than head, and extending well beyond front of anal; pectorals long and narrow, 1½ in head; a narrow membranaceous flap connecting base of pectorals with upper angle of opercular flap. Scales small, well imbricated, entirely investing body and head, including gular membrane and part of gill membranes; lateral line nearly complete, lacking for about ¼ length of body, running high, parallel with dorsal outline. Color silvery gray, dusted with coarse black specks, darker along dorsal outline; dorsal and anal with a narrow light streak at base, otherwise dusky, becoming black posteriorly, and with a narrow white margin; caudal black, with a broad white terminal bar; pectorals and ventrals white, with few black specks; peritoneum silvery white; mouth white anteriorly, its posterior portion and gill cavity jet-black. Pacific Ocean, off coast of Lower California. Many specimens from *Albatross* Station 2996, in 112 fathoms. Length 7 inches. (*Stellifer*, a genus of *Sciaenidae*; εἶδος, resemblance.)

*Neobythites stelliferoides*, GILBERT, Proc. U. S. Nat. Mus. 1891, 112, off Lower California. (Type, No. 44383. Coll. Dr. Gilbert.)

966. BARATHRODEMUS, Goode & Bean.

*Barathrodemus*, GOODE & BEAN, Bull. Mus. Comp. Zool., x, No. 5, 200, 1883 (*manatinus*).

Body brotuliform, much compressed; head compressed; mouth moderate. Head unarmed, except for a short flattened spine at upper angle of opercle. Snout long, projecting far beyond premaxillaries, its tip much swollen; jaws subequal in front. Teeth minute, in villiform bands on jaws, vomer, and palatines. No barbels. Anterior nostrils on the outer angles of the dilated snout, circular, each surrounded by a cluster of mucous tubes. Posterior nostrils above front of eye. Gill openings wide, the membranes not united. Gill rakers rather few. Body and head covered with small, thin, scarcely imbricated scales. Dorsal and anal long. Caudal fin separate, long, and slender. Ventrals close together, far in front of pectorals, each reduced to a single bifid ray. Deep-sea fishes. (*βάραθρον*, a gulf or deep abyss; *δημιος*, people.)

2889. BARATHRODEMUS MANATINUS, Goode & Bean.

Head about 6 in total length; depth  $7\frac{1}{2}$ . D. 106; A. 86; C. 2+5+2; P. 18 to 20; V. 1/1; scales about 175. Body much compressed. Dorsal and anal outline approaching at an equal angle the horizontal axis. Scales small, about 175 rows between the branchial opening and the tail, and about 34 rows, counting upward and forward obliquely from the origin of the anal to the dorsal line; lateral line apparently absent. Head considerably compressed, with rounded upper surface, its width contained  $2\frac{1}{2}$  times in its length, its greatest height equaling  $\frac{2}{3}$  its length. Snout slightly longer than the horizontal diameter of the eye, and projecting beyond tip of upper jaw a distance equal to vertical diameter of eye, much dilated and swollen, the anterior pair of nostrils being situated at the most salient angles; snout in general form resembling that of a manatee, whence the specific name. Mouth moderate, its cleft extending to the vertical from the center of the orbit; length of upper jaw equal to twice horizontal diameter of eye, and contained  $2\frac{1}{2}$  times in length of head; posterior portion of maxillary considerably expanded; maxillary largely included within a skinny sheath; when the mouth is closed the lower jaw is entirely included within the upper. Vomer and palatine with bands of teeth more than twice as broad as the bands of the intermaxillaries and on the mandible. Eye elliptical in form, its vertical diameter  $\frac{2}{3}$  of its horizontal, the latter being equal to distance from tip of snout to posterior nostril, and contained  $5\frac{1}{2}$  times in length of head; distance of eye from dorsal outline equal to  $\frac{1}{2}$  its horizontal diameter, and to  $\frac{1}{2}$  height of head in a perpendicular through center of eye; interorbital space rounded, its width equal to horizontal diameter of eye. Dorsal fin inserted in the vertical above insertion of pectoral, at a distance from end of snout equal to that of insertion of pectoral; anal inserted under twenty-first to twenty-third dorsal ray, and at a distance from snout about equal to  $\frac{1}{2}$  body length; height of dorsal and anal fins about equal to  $\frac{1}{2}$  height of body at insertion of anal, their bases extending almost to insertion of caudal; caudal composed of 9 rays, the 5 medial ones almost

equal in length, though the tip of the tail is slightly rounded, about equal to height of body midway between branchial opening and base of tail; ventrals inserted almost under middle of operculum, in length about equal to  $\frac{1}{2}$  length of head; pectorals inserted under origin of dorsal, and at a distance behind branchial opening equal to  $\frac{1}{3}$  vertical diameter of eye, its length equal to greatest height of the body. Color grayish brown; abdominal region black. (Goode & Bean.) Gulf stream, north of the Bermudas, in 647 to 1,395 fathoms. (*Manati*, like the manatee or sea cow.)

*Barthrodemus manatinus*, GOODE & BEAN, Bull. Mus. Comp. Zool., x, No. 5, 200, 1883, Lat. 33° 35' 20" N., Long. 76° W., in 647 fathoms (Type in M. C. Z. Coll. Blake); JORDAN, Cat., 127, 1885; GÜNTHER, Challenger Report, xxii, 100, 1887; GOODE & BEAN, Oceanic Ichthyology, 332, fig. 204, 1896.

#### 967. NEMATONUS, Günther.

*Nematonus*, GÜNTHER, Challenger Report, xxii, 114, 1887 (*pectoralis*).

Body compressed, with long tapering tail. Bones of head soft, muciferous channels moderately developed, and with the integument very thin or absent on the upper portion and snout. Operculum cartilaginous and flat; a broad process near its upper angle corresponding to the opercular spine in some of the related genera, the head otherwise unarmed, though irregular by reason of the cranial bones. Snout much depressed, broad, rounded; jaws equal in front; mouth very wide; bands of villiform teeth in jaws, on vomer and palatines. Barbel none. Eyes small. Vertical fins confluent; ventrals a pair of bifid filaments close together, on the isthmus, close to the humeral symphysis. Gills 4, with very short laminae and rather short, incurved, acicular gill rakers on the first arch, and much shorter, less numerous, spatulate ones on the 3 other arches. Pseudobranchiae rudimentary. No traces of a lateral line, though the body is covered with scales of considerable size, almost as large as the eye, and the cheek with others still larger. *Nematonus* differs from *Porogadus* not only in the absence of spines upon the head, as Günther has indicated, but in the much less ossified opercular apparatus, in the shorter and thicker head, in the absence of the 3 series of pores simulating lateral lines, and in the tendency to prolongation in the lower rays of the pectoral, which increase from the uppermost to the lowermost in *Nematonus*, while *Porogadus* has a lanceolate fin, and also in the extreme exertion of the caudal rays. (*νήμα*, thread; *Onus*, the rockling.)

#### 2890. NEMATONUS PECTORALIS, Goode & Bean.

D. 93; A. 73; P. 17; V. 2. Body moderately elongate, much compressed, the tail much shorter and more robust than in *Bassoctes catena*, its height equaling  $1\frac{1}{2}$  times length of head and  $\frac{1}{2}$  that of body. Head stoutish, not much compressed, lower than body, its length contained  $5\frac{1}{2}$  times in the body; snout compressed, broad at its tip, its length exceeding diameter of the circular eye; interorbital area slightly convex, its width slightly exceeding twice diameter of eye, 3 times in head. Maxillary

extending far behind eye, its length less than that of preorbital portion of head; mandible as long as postorbital portion of head; jaws, vomer, and palatines with narrow bands of villiform teeth, normally arranged. Branchiostegals 8. Gill lamellae very short; gill rakers long and numerous, 18 on first arch below the angle, 5 above, 4 of which are rudimentary. Pseudobranchiae present, but very rudimentary. Anterior nostrils on the top of the snout and near the median line of the head, near its tip, separated by a space about equal to diameter of eye; posterior nostrils in front of eye. Muciferous pores large, arranged much as in *B. catena*. Dorsal origin in the same vertical with that of pectorals, its distance from tip of snout contained 5 times in total and equaling twice length of maxillary. Rays well developed in anterior third, the longest  $\frac{2}{3}$  of head; anal origin under twentieth dorsal ray, its rays nearly as long as those of dorsal; pectoral with its penultimate ray produced, extending to thirteenth ray of anal, nearly twice as long as head; ventrals originating in advance of vertical through pectorals, and each a bifid filament; distance of ventral origin from tip of snout equaling length of ventral and about  $\frac{3}{4}$  as long as head; distance of ventral origin from vent considerably greater than length of head; distance from tip of ventral to vent equal to  $\frac{1}{2}$  the length of the head. Number of scales in transverse series from vent to dorsal about 23; from the upper angle of the gill opening to the vertical through origin of anal 32. Color brownish yellow; head and abdomen blackish. The type (No. 37342, U. S. Nat. Mus.) was taken at *Albatross Station* 2380, Lat.  $28^{\circ} 02' 30''$  N., Long.  $87^{\circ} 43' 45''$  W. in 1,430 fathoms. It is 183 mm. long to the caudal base, 215 with caudal. Another young specimen, 70 mm. long, was taken at *Blake Station* XCV, off *Dominica*, in 330 fathoms. (Goode & Bean.) (*pectoralis*, pertaining to the breast.)

*Nematonus pectoralis*, GOODE & BEAN, Proc. U. S. Nat. Mus. 1885, 604, Lat.  $28^{\circ} 02' 30''$  N., Long.  $87^{\circ} 43' 45''$  W., in 1,430 fathoms (Type, No. 37342. Coll. *Albatross*); GÜNTHER, Challenger Report, XXII, 114, 1887; GOODE & BEAN, *Oceanic Ichth.*, 333, fig. 295, 1896.

#### 968. POROGADUS, Goode & Bean.

*Porogadus*, GOODE & BEAN, Proc. U. S. Nat. Mus. 1885, 682 (*miles*).

Body brotuliform, much compressed; head with numerous spines on interorbital space, 2 pairs on the shoulders, 1 at angle of operculum and a double series on angle of preoperculum; head with numerous mucous pores, as in *Bassozetus*; mouth large; snout moderate, not projecting much beyond the upper jaw; jaws nearly equal in front; teeth in villiform bands in jaws and on vomer and palatines; barbel none; gill openings wide, membranes narrowly united, not attached to the isthmus; gills 4; gill laminae short; gill rakers moderate, numerous; pseudo-branchiae absent; caudal fin of few rays, on a very narrow base, not prolonged, scarcely differentiated from the vertical fins; dorsal and anal fins well developed; pectorals simple, moderate; each ventral a single bifid ray close to the humeral symphysis; branchiostegals 8; scales small; lateral line apparently triple, or replaced by 3 series of pores—1 close to ventral outline, 1 median, and another along base of dorsal. (*πόρος*, pore; *Gadus*, the codfish.)

## 2891. POROGADUS MILES, Goode &amp; Bean.

Head  $6\frac{1}{2}$ ; depth 10; eye  $5\frac{1}{2}$  in head; body much compressed, elongate, tapering to a very slender tail; head long, moderately compressed, subconical, the profile gradually ascending in nearly a straight line from tip of snout to origin of dorsal; interorbital space slightly convex, spiny, its width  $4\frac{1}{2}$  times in length of head, and slightly greater than diameter of eye; opercles and head generally covered with numerous and strong spines, as described in the generic diagnosis; mouth very large and wide; maxillary extending far behind eye and much expanded at its tip, its length more than  $\frac{1}{2}$  that of head; length of mandible equal to greatest height of body; jaws, vomer, and palatines with narrow bands of villiform teeth, none of which is enlarged; gill rakers 15 on anterior arch below the angle, 3 rudimentary ones above. Anterior pair of nostrils nearly on top of snout and somewhat nearer its tip than to eye, separated by a narrow space and placed immediately in front of middle of eye; behind each posterior nostril a strong spine projecting outward and upward; pores of the head arranged much as in *Bassocetus*; scales minute; lateral line not to be clearly made out; 3 rows of minute pores on each side of dorsal, median, and ventral, beginning near head and extending well toward extremity of tail. Dorsal origin slightly behind vertical through pectoral base, its distance from snout nearly 6 times in length of body, its rays moderately long, the longest about as long as snout, and very numerous; anal origin in vertical from twenty-second or twenty-third dorsal ray, its distance from snout  $3\frac{1}{2}$  times in length of body, its rays about as long as those of dorsal; pectoral imperfect, its length in the type equaling  $\frac{1}{2}$  that of head; ventrals a bifid filament, placed close to the humeral symphysis, well in advance of pectoral, its length equal to height of body; distance from origin of ventrals to vent nearly equal to length of head; ventral not reaching vent by a distance equal to length of snout. Color blackish brown. The type (No. 35625, U. S. Nat. Mus.) is 153 mm. in length, from *Albatross* Station 2230, lat.  $38^{\circ} 27' N.$ , long.  $73^{\circ} 02' W.$ , at a depth of 1,168 fathoms. (Goode & Bean.) (*miles*, a soldier.)

*Porogadus miles*, GOODE & BEAN, Proc. U. S. Nat. Mus. 1885, 602, Lat.  $38^{\circ} 27' N.$ , Long.  $73^{\circ} 02' W.$ , in 1,168 fathoms; GOODE & BEAN, Oceanic Ichthyology, 334, fig. 292, 1896.

## 969. PENOPUS, Goode &amp; Bean.

*Penopus*, GOODE & BEAN, Oceanic Ichthyology, 335, 1896 (*maedonaldii*).

Body stout in front, tapering behind; tail not greatly exceeding the length of the rest of the fish; head scaly, thick, its top surface flat, with depressed and moderately projecting snout; a pair of minute postnasal spines; a strong and much curved spine on the operculum; several weak spines on the angle of the preoperculum, and several at the posterior angle of the suboperculum; mouth moderately large, the lower jaw included; several narrow slit-like pores along the margin of the preorbital and suborbital; 2 minute pores on under surface of mandible near its symphysis, and not far behind them 2 long slit-like pores; the



anterior nostril in a long slit, the posterior larger, oblong in shape, and  $\frac{1}{2}$  concealed by a fold of skin; eye small; the teeth appear only in minute asperities, the intermaxillary band much wider in front than behind; mandibular band narrow throughout; vomerine band very narrow, V-shaped; palatines in a long, broad band; gill openings wide, deeply cleft in front, narrowly joined to the isthmus; branchiostegals 8; no pseudobranchia; gill rakers long and slender, not numerous; gill laminae moderately long, a long slit behind the fourth gill; scales very small; lateral lines 3; caudal fin consisting of few rays, well differentiated from the dorsal and anal; dorsal beginning not far behind head; ventrals slightly in advance of the pectorals and composed of 2 rays, united by membrane, which forms a margin around them; pectoral normal, several of its upper rays simple; vent not much in advance of middle of total length. This genus agrees with *Porogadus* in nearly every respect except in the scarcity of spines on the head and in the structure of the ventrals. *Porogadus* has the ventrals composed of 2 distinct rays which are separated throughout their entire length, but in *Penopus* the 2 rays are inclosed in a membrane which connects them and forms a margin around them. In *Porogadus*, also, the suboperculum has a smooth margin and the opercular spine is weaker than in *Penopus*, and is not curved. Deep seas. (*νήρη*, thread; *πούς*, foot.)

2892. *PENOPUS MACDONALDI*, Goode & Bean.

D. 137; A. 102. Greatest height of body equaling length of postorbital part of head and about  $\frac{1}{3}$  of total without caudal; greatest width of body anteriorly about  $\frac{3}{4}$  of its greatest height; head stout, its greatest width equaling  $\frac{3}{4}$  of its greatest depth and more than  $\frac{1}{2}$  of its length; width of interorbital space about  $\frac{1}{3}$  length of head; eye very small, its length less than  $\frac{1}{2}$  width of interorbital space; distance from eye to tip of snout equaling length of intermaxillary; distance of anterior nostril from tip of snout equaling length of eye; distance of posterior nostril from eye slightly less than its distance from tip of snout; maxillary expanded behind and reaching somewhat behind eye, its length equaling that of snout; mandible extending much behind eye, its length equal to postorbital part of head. Dorsal beginning over middle of pectoral, its rays well developed, those in middle of fin longer than anterior ones; anal beginning under twenty-seventh ray of dorsal, middle rays longest; pectoral nearly  $\frac{1}{2}$  length of head and about equal to distance of its tip from vent; ventral about  $\frac{1}{3}$  distance of its origin from origin of anal. Lateral lines 3, the uppermost beginning at the upper angle of the gill opening, quickly approaching top of body near base of dorsal and merging into dorsal base about middle of tail; median lateral line beginning a little behind head and extending almost to root of caudal, becoming very faint posteriorly; lowermost lateral line with its origin under and not far from base of pectoral, extending along lower side of tail and merging into base of anal fin somewhat beyond middle of length of tail. Color yellowish brown; operculum, opercular flap and branchiostegal membrane, pectoral, and ventral dusky. Only a single specimen, 315 mm. long, known; obtained by the *Albatross* September 18, 1886, at Station 2716, Lat. 38°

29° 30' N., Long. 70° 57' W., in 1,631 fathoms. (Goode & Bean.) (Named for Hon. Marshall McDonald.)

*Penopus mardonaldi*, GOODE & BEAN, *Océanide* 10 (11) (12) (13) (14) (15) (16) (17), 336, fig. 293, 1896, Lat. 38° 29' 30' N., Long. 70° 57' W., in 1,631 fathoms. (Type, No. 39413.)

970. *DICROLENE*, Goode & Bean.

*Dicrolene*, GOODE & BEAN, Bull. Mus. Comp. Zool., 8, No. 5, 202, 1883 (*Intronigra*).

Brotulids with body moderately compressed; head somewhat compressed; mouth large; tip of maxillary much dilated; eye large, placed close to dorsal profile. Head with supraorbital spines; several strong spines on preopercle and 1 long spine at upper angle of opercle. Snout short, not projecting beyond the upper jaw; jaws subequal. Teeth in narrow villiform bands in each jaw, on head of vomer, and on palatines. No barbel. Gill openings wide, membranes not united; gills 4; gill laminae of moderate length; gill rakers rather long, not numerous; pseudobranchiae absent. Caudal not confluent with dorsal and anal, but without a distinct peduncle. Dorsal and anal fins long; pectoral rays in 2 groups, several of the lower ones being separated and much produced; ventrals a pair of bifid rays, close together on the isthmus. Branchiostegals 8. Body and head covered with small scales; lateral line close to base of dorsal fin, apparently becoming obsolete on posterior third of body. Stomach siphonal; pyloric caeca few and rudimentary; intestine shorter than body. Deep sea; a single species known. (*δικρός*, forked; *ὄλεν*, limb.)

2893. *DICROLENE INTRONIGRA*, Goode & Bean.

Head 5; eye large, 4 in head; interorbital width 4. D. 100; A. about 85; C. 6 or 7; V. 1; P. 19+7 or 8; scales 110 to 120. Body moderately compressed, its dorsal and ventral outlines approaching at an equal angle the horizontal axis, and tapering to a narrow point. Head somewhat compressed, with flattish upper surface, which is encroached upon by the upper margin of orbit; a strong spine at posterior upper margin of orbit, pointing backward and upward; a long, sharp spine at upper angle of opercle, its exposed portion 2 in eye; 3 equidistant spines on lower posterior border of preopercle, much weaker than that on opercle. Large muciferous cavities in bones of head; a row of large cavities extending backward from upper angle of orbit, and continuous with those on lateral line. Mouth large, its cleft considerably more than  $\frac{1}{2}$  head, the maxillary extending beyond eye and with scales upon its expanded tip. Distance from snout to origin of dorsal fin  $\frac{3}{4}$  total length; anal inserted under twenty-fifth or twenty-sixth dorsal ray; height of dorsal and anal fins each about equal to eye; length of caudal fin 2 in distance from snout to dorsal; ventrals about equal to upper jaw; pectorals inserted close to branchial aperture, the 8 lower rays free and much prolonged, the longest and most anterior being about 3 in body, and more than 3 times as long as the contiguous posterior ray of the normally constructed portion of the fin, which is, however, about equal to the last free rays. West Indies, Gulf of

Mes  
ou t  
Dicr  
:  
:

Mix

In  
on t  
stro  
cove  
comp  
exce  
chee  
I sm  
small  
short  
consi  
inser  
thin  
teeth  
gill  
intra

Hea  
below  
extre  
with  
exam  
length  
maxil  
vent  
root  
anal  
its ray  
vent  
10, m  
appa  
inches  
enorm  
was tr  
Bathym  
oms  
Siro  
Cap  
Mixou  
Oce

Mexico, and Gulf Stream in various localities, and off coast of Soudan and on the bank d'Arguin, in deep water. (*intro*, within; *niger*, black.)

*Dicrolene intronigra*, GOODE & BEAN, Bull. Mus. Comp. Zool., x, No. 5, 202, 1883, Gulf Stream, Lat. 39° 59' 45" N., Long. 68° 54' W. (Coll. Blake); GÜNTHER, Challenger Report, XXII, 107, 1887; VAILLANT, Exp. Sci. Travailleur et Talisman, 258, pl. 23, fig. 2, 1888; GOODE & BEAN, Oceanic Ichthyology, 338, fig. 297 A and B, 1896.

971. MIXONUS, Günther.

*Mixonus*, GÜNTHER, Challenger Report, XXII, 108, 1887 (*laticeps*).

Lower pectoral rays free, not united by membrane with, but inserted on the same base as, the upper part of the fin; they are but slightly stronger than the other rays and prolonged. Body elongate, compressed, covered with small, very thin and deciduous scales. Head slightly compressed, broad and flat above, depressed in front, naked (with the exception of the parts between the mandibles, and, perhaps, of the cheeks). Bones thin, with muciferous system moderately developed; only 1 small spine above on the operculum; preoperculum without spine. Eye small. Vertical fins united, but the narrow caudal projecting beyond the short dorsal and anal rays. Ventrals each reduced to a filament, which consists of 2 rays firmly bound together in their whole length; they are inserted behind the humeral symphysis and close together. Snout broad, rounded, scarcely overlapping the lower jaw. Mouth very wide; villiform teeth in the jaws, on the vomer, and palatine bones. Gill laminae short; gill rakers long, not very closely set. Pseudobranchiae none. (*mitis*, *mixure*, half; *Onus*, a synonym of *Haidropsarns*, the rockling.)

2894. MIXONUS LATICEPS (Günther).

Head 2; depth 3; eye 8 in head; snout 1. P. 17. Greatest depth of body below origin of dorsal fin; distance of vent from snout  $\frac{2}{3}$  its distance from extremity of spinal column. Crown of head remarkably convex, covered with an extremely thin and transparent skin, which, perhaps, in older examples is senile; interorbital space less convex, and equaling in width the length of snout including the eye; eye small, above middle of length of the maxillary; posterior nostrils wide, open, in front of the eye. Distance of vent from ventrals exceeds length of head; origin of dorsal fin above root of pectorals, its rays of moderate length, but longer than those of anal; pectoral with a rather narrow base, as long as head without snout, its rays feeble, 3 or 4 lower ones a little stouter, detached, and prolonged; ventral filaments not reaching as far backward as pectoral. Gill rakers 10, much longer than the laminae. Whitish, with the abdomen and gill apparatus black. Mid-Atlantic, in profound depths. One specimen, 5½ inches long, was obtained in mid-Atlantic (*Challenger* Station 104), at the enormous depth of 2,500 fathoms. The second (type of *Sirembu guntheri*) was taken off Cape Verde, in 3,200 meters. (*latus*, broad; *-ceps*, head.)

*Bathyneetes laticeps*, GÜNTHER, Ann. Mag. Nat. Hist. 1878, 20, mid-Atlantic, in 2,500 fathoms. (Coll. Challenger.)

*Sirembu guntheri*, VAILLANT, Exp. Sci. Trav. et Talisman, 268, p. XXIV, fig. 5, 1889, off the Cape Verde Islands, at a depth of 3,200 meters.

*Mixonus laticeps*, GÜNTHER, Challenger Report, XXII, 108, pl. 25, fig. 8, 1887; GOODE & BEAN, Oceanic Ichthyology, 339, fig. 297 A, 1896.

## 972. BARATHRONUS, Goode &amp; Bean.

*Barathronus*, GOODE & BEAN, Bull. Mus. Comp. Zool., x, No. 5, 164, 1883 (*bicolor*).

Brotnlids having the head stout, body and tail compressed, covered closely by skin; scaleless; vent far behind pectoral, included in a cleft; mouth wide, oblique, the lower jaw projecting; intermaxillary teeth rudimentary; several fang-like teeth on the head of the vomer, none on palatines, a few rather large, recurved, separated teeth in the mandible; nostrils close together and small; eye visible through the skin, partly upon the top of the head, with or without dark pigment in the iris; barbel none; gill rakers very numerous and slender, and rather long; gill laminae well developed on all the arches; no pseudobranchiae; head full of muciferous channels; gill membranes not united, but covered by a fold of skin; ventrals reduced to single simple rays, placed in advance of the pectorals and close to the humeral symphysis; dorsal and anal placed far back; caudal scarcely differentiated, composed of rather numerous, very slender rays upon a somewhat narrow base. (*βάραθρον*, the abyss; *ὄνος*, *Onus*, the rockling.)

## 2895. BARATHRONUS BICOLOR, Goode &amp; Bean.

Head  $5\frac{1}{2}$  in total, its width  $\frac{3}{4}$  its length; depth  $6\frac{1}{2}$ ; orbit  $4\frac{1}{4}$  in head; interorbital width  $4\frac{1}{4}$ . D. about 70; A. 57. Body much compressed; eye concealed by the skin; maxillary extending slightly beyond the perpendicular through posterior margin of orbit, almost entirely concealed under the preorbital, and much expanded at tip, where its width is rather greater than that of eye. Intermaxillary very thin, broad, and slightly protractile; vomer very close to intermaxillary symphysis, its head somewhat raised and bearing 3 fang-like teeth (2 of which are off one side and 1 on the other in the type), separated by a moderately wide interspace; mandible with 5 enlarged, separate, recurved teeth upon each side, which increase in size posteriorly, its upper edge, posteriorly, produced above the level of the tooth-bearing surface, and received under the expanded maxillary; longest gill raker about as long as eye. Dorsal origin distant from the snout about  $\frac{1}{2}$  total length; dorsal rays well developed, numerous, long, and slender, the longest about 3 times in length of head; anal originating in vertical from fourteenth dorsal ray, equidistant from eye and base of caudal, longest rays about as long as those in the dorsal; pectoral with a fleshy base, its length a little less than height of body. Ventral well in advance of pectoral, close to humeral symphysis, the rays being placed very close together at their origin, the length of the fin contained about 9 times in the total length, about 3 times in the distance from its origin to the vent. Caudal with about 10 rays, its length about 8 times in total length. Color yellowish white, with a broad vertical band of black from origin of ventral nearly to vent; another similar and narrower band above it upon each side. The type, 120 mm. long, from Blake Station LXXI, off Guadalupe, at a depth of 769 fathoms. (Goode & Bean.) (*bicolor*, two-colored.)

*Barathronus bicolor*, GOODE & BEAN, Bull. Mus. Comp. Zool., x, No. 5, 164, 1883, off Guadalupe, in 769 fathoms (Coll. Blake); GOODE & BEAN, Oceanic Ichthyology, 341, fig. 298, 1896.

Aph  
H  
sca  
of t  
is w  
plu  
men  
exte  
men  
lam  
gill  
sinu  
are  
supe  
line  
the r

Bo  
thick  
4½ in  
not c  
abon  
marg  
equal  
mand  
but a  
gill r  
almos  
total  
head  
its ra  
origin  
head  
symp  
pecto  
Skin  
Gulf o  
gelatin

Aphyon  
N.,  
olo

Bod  
almost

## 973. APHYONUS, Günther.

*Aphyonus*, GÜNTHER, Ann. Mag. Nat. Hist. 1878, 22 (*gelatinosus*).

Head, body, and tapering tail strongly compressed, enveloped in a thin, scaleless, loose skin. Vent far behind the pectoral, at nearly the middle of the total length. Snout swollen, projecting beyond the mouth, which is wide. No teeth in the upper jaw; small conical teeth in the lower, pluriserial in front and uniserial on the side. Vomer with a few rudimentary teeth; palatine teeth none. Nostrils close together, small. No externally visible eye. Barbel none. Ventrals reduced to simple filaments, placed close together and near to the humeral symphysis. Gill membranes not united. Four branchial arches, the posterior without gill laminae, the anterior with very short gill rakers and with rather short gill laminae. Head covered with a system of wide muciferous channels and sinuses, the dermal bones being almost membranaceous, while the others are in a semicartilaginous condition. Notochord persistent, but with a superficial indication of the vertebral segments, as in some Leptocephaline forms. (Günther.) (*ἀφύνη*, anchovy, a small translucent fish; *Onus*, the rockling.)

## 2896. APHYONUS MOLLIS, Goode &amp; Bean.

Body much compressed, its greatest height 6 in its total length. Head thicker than body, its height slightly greater. Length of head about  $4\frac{1}{2}$  in total, width over  $\frac{1}{2}$  its length. Snout  $3\frac{1}{2}$  in length of head. Eye not externally visible. Diameter of orbit, as seen through the skin, about  $\frac{1}{2}$  length of head. Maxillary extending to vertical through posterior margin of orbit, the mandible somewhat farther back, its length nearly equal to height of body. A few weak teeth on vomer, palatines, and mandible, and very rudimentary ones in maxillary, not visible to the eye, but appreciable to the touch. Gill laminae on the fourth and rudimentary gill rakers, 8 rudiments and 4 developed below the angle. Dorsal origin almost over posterior edge of operculum, its distance from the snout  $\frac{1}{2}$  of total length, dorsal rays more than 110, well developed, the longest 3 in head; anal origin slightly nearer base of caudal than to the tip of snout, its rays shorter than those in the dorsal; pectoral with a fleshy base, its origin somewhat behind that of the dorsal, its length equal to width of head; ventral origin in advance of that of pectoral, close to humeral symphysis, the fin a single simple ray, whose length equals that of the pectoral, its tip not reaching vent by a space equal to height of head. Skin not loose. Texture of body rather firm, not transparent; whitish. Gulf of Mexico, in deep water. This species is closely allied to *Aphyonus gelatinosus*. (Goode & Bean.) (*mollis*, soft.)

*Aphyonus mollis*, GOODE & BEAN, Bull. Mus. Comp. Zool., x, No. 5, 163, 1883, Lat.  $24^{\circ} 36'$  N., Long.  $84^{\circ} 5'$  W., in 955 fathoms (Coll. Blake); GOODE & BEAN, Oceanic Ichthyology, 342, fig. 299, 1896.

## Family CCXII. BREGMACEROTIDÆ.

Body stout, with robust caudal portion, truncate or convex behind, almost without procurent caudal rays above or below; vent before mid-

dle of body; suborbitals moderate; no barbels, spines, nor cirri on head; mouth terminal, with minute teeth on jaws and vomer, none on palatines; ventrals jugular, extremely long, few-rayed, the rays dilated and separate nearly to base. Dorsal fins 2, the first an elongate, slender occipital ray; second dorsal on posterior half of body, of soft rays, depressed medially, so that it forms 2 lobes; no spines in fins. Anal nearly similar to the soft dorsal and similarly depressed in the middle; dorsal and anal depressible in a groove of scales. Hypercoracoid perforate; no pseudo-branchiæ; gill openings wide, the membranes free from the isthmus. A single genus with 2 or 3 species found in the open sea, probably near the surface; widely distributed. The presence of the hypercoracoid foramen shows that this family is allied to the *Brotulidæ* rather than to the *Gadidæ*. From the *Brotulidæ* it is mainly distinguished by the development of its dorsal and ventral fins.

#### 974. BREGMACEROS, Thompson.

*Bregmaceros*, THOMPSON, in Charlesworth's Mag. Nat. Hist., IV, 1840, 184 (*maccellandii*).  
*Calloptilum*, RICHARDSON, Voy. Sulph., Fish., 94, pl. 48, figs. 4-7, 1843 (*mirum*).  
*Asthenurus*, TICKELL, Journ. Asiat. Soc. Bengal 1865, 32 (*atripinnis*).

Characters of the genus included above. (*βρέγμα*, the upper part of the head, the nape; *κέρας*, horn.)

a. Scales in transverse series 14; scales in lateral series 58 to 64.

MACCELLANDII, 2897.

aa. Scales in transverse series 10; in lateral series 65; anterior lobes of dorsal and anal lower than in *B. maccellandii*.

ATLANTICUS, 2898.

#### 2897. BREGMACEROS MACCELLANDII, Thompson.

Head  $5\frac{2}{3}$ ; depth  $6\frac{3}{8}$ . D. about I, 18-X-22 (16 + X-15); A. about 18, X, 22 (22, X, 20); V. 4 or 5; scales 58-14 (64-14). Body moderately elongate, compressed, the form somewhat as in *Ophidion*, the back not elevated. Head short and small, moderately compressed; bones of head thin, without serrature or spine; eye moderate, 3 in head; interorbital space ridged, about as broad as eye; snout blunt, rather shorter than eye; mouth very oblique, the jaws subequal; maxillary reaching to beyond middle of eye,  $2\frac{1}{2}$  in head; lower jaw flattish, curved upward; teeth in both jaws moderate, slender, close set, recurved, apparently in a single series. Tongue conspicuous; no teeth evident on vomer or palatines; branchiostegals 7 or 8; gill membranes separate, free from the isthmus; no evident pseudobranchiæ; gill rakers obsolete; no barbels about jaws. Body with rather large, thin, caducous scales (nearly all of them fallen in the typical specimens so that they can not be counted). Dorsal fin beginning with a single long and very slender spine on occiput, this nearly  $\frac{1}{2}$  longer than head. Behind this, for a distance about equal to its length, the rudimentary rays, if present, do not rise above the sheath on each side. Nearly opposite the vent begins the dorsal proper, the distance of its first ray from snout being about  $\frac{2}{3}$  length of body; about 12 rays are moderately elevated, about  $\frac{1}{2}$  length of head. The others are gradually shorter and more slender, becoming too

small to count, until just before caudal, where the fin becomes conspicuous again, this posterior lobe not  $\frac{1}{2}$  so high as the anterior. Anal opposite dorsal and similar to it, the first ray close behind vent; caudal free from dorsal and anal, the caudal peduncle truncate at its base. Ventrals of 3 long rays, with a fourth at the inner base of the third; this fourth is probably a rudiment of 2. The ventrals are jugular in position, the rays very long and filamentous, the longest about  $\frac{1}{2}$  the body, reaching to the middle of anal fin. Pectorals inserted high, somewhat shorter than head. Vent slightly behind end of anterior  $\frac{1}{2}$  of total length. Color brown above, sides and below silvery; back and base of anal closely dotted with dusky; dorsal mostly dusky; caudal pale, dusky at base, with a narrow white cross bar; lower fins pale; the dark marking on front of back assume something of the form of lengthwise streaks. Tropical Pacific; Bay of Bengal; Philippine Islands; coast of China, etc., east to the coast of Central America, living near the surface in the open sea. Here described from the types of *Bregmaceros bathymaster*, two specimens, 1 $\frac{1}{2}$  and 2 inches in length, dredged at *Albatross* Station 2804, south of Panama, in 47 fathoms depth. Two others, 4 inches long, found later off the coast of Panama. A recomparison of these latter specimens with Günther's\* detailed account of *B. maccllellandii* shows no difference whatever, and we regard *B. bathymaster* as identical with the latter. Günther counts the scales 64-14; we find 58-14. In our largest specimens the ventrals reach middle of anal. (Named for Dr. John McClelland, of the Bengal Medical Service, who first studied the fishes of the Ganges.)

*Bregmaceros maccllellandii*, THOMPSON, in Charlesworth's Mag. Nat. Hist., IV, 1840, 184, mouth of the Ganges; GÜNTHER, Cat., IV, 368, 1862.

*Calloptilum mirum*, RICHARDSON, Voyage Sulphur, Fish., 95, pl. 46, figs. 4-7, 1843.

*Athenurus atripinnis*, TICKELL, Journ. Asiat. Soc. Bengal 1865, 32, with plate, Bay of Bengal.

*Bregmaceros bathymaster*, JORDAN & BOLLMAN, Proc. U. S. Nat. Mns. 1889, Lat. 8° 13' 30" N., Long. 79° 37' 45" W., southwest of Panama. (Type, No. 41137. Coll. *Albatross*.)

*Bregmaceros atripinnis*, DAY, Proc. Zool. Soc. Lond. 1869, 522, Bay of Bengal; types, same specimens described by TICKELL.

#### 2598. BREGMACEROS ATLANTICUS, Goode & Bean.

Head  $5\frac{1}{2}$ ; depth  $7\frac{1}{2}$  in total length. D. I-15, X, 16; A. 15 or 16 + X (7 or 8) + 21 or 22; scales 65-10. Length 46 mm. Body compressed, moderately elongate. Interorbital area convex, its width greater than eye, which is 4 in head; jaws even in front; maxillary reaching to vertical through middle of eye; mandible to vertical through posterior margin of eye; teeth in intermaxillary minute, apparently in a single series; mandibu-

\* The following is Dr. Günther's account of *Bregmaceros maccllellandii*, taken from specimens from the China Sea:

"B. 7; D. I, 16 + X + 15; A. 22 + X + 20; V. 5 or 6; scales 64-14. Occipital ray very slender, longer than head; dorsal and anal fins depressible in a groove formed by the scales along the bases of these fins; anterior portions of dorsal and anal elevated, connected with the posterior lower portion by a series of very short extremely feeble rays. Vent at end of anterior third of total length. Three outer rays of ventral fins dilated, compressed, simple, much elongate, reaching to or nearly to middle of anal; the second and third rays sometimes united at base. Silvery, minutely dotted with brown."

lary teeth biserial, the inner teeth enlarged. Cephalic appendage reaching nearly to base of first dorsal, its length  $4\frac{1}{2}$  in total. Distance of dorsal from snout  $2\frac{1}{2}$  in total, that of the anal the same; the dorsal and anal fins received in a groove formed by the scales along their bases; anterior portion of second dorsal and second anal less elevated than in *B. maclellandii*. The differentiations between the developed and undeveloped rays of the anal are so slight that the limits of the so-called anterior and posterior sections of the fin can not be determined. Length of the longest anal ray about 2 in body length. Specimens were obtained by the Blake at the following stations: XCIX, off Granada, 90 fathoms; CXIII, off Neris, 305 fathoms; CLXXXV, Lat.  $25^{\circ} 33' N.$ , Long.  $84^{\circ} 21' W.$ , 101 fathoms. (Goode & Bean.) This species seems doubtfully distinct from *B. maclellandii*. (*atlanticus*, of the Atlantic.)

*Bregmaceros atlanticus*, GOODE & BEAN, Bull. Mus. Comp. Zool., XII, No. 5, 165, 1886, West Indies, off Granada and Neris (Coll. Blake); GOODE & BEAN, Oceanic Ichthyology, 388, fig. 331, 1896.

### Suborder ANACANTHINI.

#### (THE JUGULAR FISHES.)

Vertical fins very long, destitute of true spines; tail isocercal, the posterior vertebra progressively smaller; ventrals jugular, without spines; hypercoracoid without perforation or foramen; no pseudobranchiae. The osteological characters of this group, called by him *Gadoidea*, are thus given by Dr. Gill:

"Jugulares with the orbito-rostral portion of the cranium longer than the posterior portion, the cranial cavity widely open in front; the supra-occipital well developed, horizontal and cariniform behind, with the exoccipitals contracted forward and overhung by the supraoccipital, the exoccipital condyles distant and feebly developed, with the hypercoracoid entire, the hypocoracoid with its inferior process convergent toward the proscapula, and the fenestra between the hypercoracoid and hypocoracoid." (Gill, Proc. Ac. Nat. Sci. Phila. 1884, 170.)

A large and important group, chiefly confined to the cold depths of the ocean and the northern seas. From all other typical fishes they are separated by the entire hypercoracoid. (*ἀν-* privative, without; *ἀκανθα*, spine.)

a. Caudal fin present; tail not greatly elongate; body tapering or coniform behind, with many procurent caudal rays above and below; suborbitals moderate.

b. Frontal bones paired, with a triangular excavated area above, the divergent frontal crests continuous from the forked occipital crest; ribs wide, approximated, channeled below or with inflected sides; no barbels.

MERLUCCIDÆ, CCXIII.

bb. Frontal bones normal, not forming a triangular excavated area above; ribs normal; chin with a barbel (rarely obsolete).

GADIDÆ, CCXIV.

aa. Caudal fin wanting; tail very long, tapering behind; suborbitals very broad.

MACROURIDÆ, CCXV.



Family CCXIII. MERLUCCIIDÆ.

(THE HAKES.)

Body moderately elongate, covered with small, smooth, deciduous scales; posterior part of body coniform and with the caudal rays procurrent forward; vent submedian. Head elongate, depressed, pike-like; suborbital bones moderate; mouth terminal, with strong teeth; no barbels; ventrals subjugular; dorsal fins 2, a short anterior and long posterior one, a long anal corresponding to the second dorsal; ribs wide, approximated, and channeled below or with inflected sides; frontal bones paired, excavated, with divergent crests continuous from the forked occipital crest. A single genus, with about 4 species; large cod-like fishes, of voracious habit, inhabiting moderate depths, and distinguished from the *Gadida* mainly by the structure of the frontal bones and the ribs. (*Merlucciidae*, Gill, Proc. Ac. Nat. Sci. Phila. 1884, 772.)

975. MERLUCCIUS, Rafinesque.

(HAKES.)

*Merluccius*, RAFINESQUE, Caratteri di Alcuni Nuovi Generi, etc., 26, 1810 (*merluccius*).

*Onus*, RAFINESQUE, Indice d'Ittol. Sicil., 12, 1810 (*riali* = *merluccius*); substitute for *Merluccius*.

*Merlangus*, RAFINESQUE, Indico d'Ittol. Sicil., 30, 1810 (*riali*); substitute for *Onus*.

*Merlus*, GUICHENOT, in Gay, Hist. Nat. Chili, Zool., II, 328, 1847 (*gayi*).

*Stomodon*, MITCHILL, Rept. Fish. N. Y. 1814, 7 (*bilinearis*).

*Homalopomus*, GIRARD, Proc. Ac. Nat. Sci. Phila. 1856, 132 (*trowbridgei*).

*Epicopus*, GÜNTHER, Cat. Fish. Brit. Mus., II, 248, 1860 (*gayi*).

Body elongate, covered with small, deciduous scales. Head slender, conical, the snout long, depressed; a well-defined, oblong, triangular excavation at the forehead, bounded by the ridges on the separated frontal bones, these ridges converging backward into the low occipital crest; eye rather large; edge of preopercle free; preopercle with a channel behind its crest, crossed by short radiating ridges; mouth large, oblique; maxillary extending to opposite the eye; lower jaw longer; no barbels; jaws with slender teeth, of various sizes, in about 2 series, those of the inner row longer and movable; vomer with similar teeth; palatines toothless. Branchiostegals 7. Gill rakers long; gill membranes not united. Dorsal fins 2, well separated, the first short, the second long, with a deep emargination; anal emarginate, similar to second dorsal; ventral fins well developed, with about 7 rays; vertebrae peculiarly modified, the neural spines well developed and wedged into one another; frontal bone double and the skull otherwise peculiar in several respects. Species several, very similar in appearance; ill-favored fishes of soft flesh and fragile fins, inhabiting water of some depth. Large voracious fishes, little valued as food. (*Merluccius*, the ancient name, meaning sea pike.)

a. Scales moderate, about 110 in lateral line; teeth very strong. D. 19-36; A. 36.

MERLUCCIUS, 2899.

aa. Scales small, 135 to 150 in lateral line; teeth moderate. D. 11 to 13-41; A. 41.

b. Ventrals long, about  $1\frac{3}{4}$  in head.

BILINEARIS, 2900.

bb. Ventrals short, about  $2\frac{1}{2}$  in head.

PRODUCTUS, 2901.

## 2899. MERLUCCIUS MERLUCCIUS (Linneus).

(EUROPEAN HAKE.)

Head large,  $3\frac{1}{2}$ ; depth  $6\frac{1}{2}$ . D. 10-36; A. 36; vertebrae 25+26; scales 150. Ventrals a little more than  $\frac{1}{2}$  head; teeth very long. Dusky above, silvery below; dorsal, caudal, and distal part of pectoral blackish; inside of opercle black; inside of mouth black posteriorly, pale in front; peritoneum black. Coasts of Europe, generally abundant, south to Madeira and Italy, straying to Greenland.\* Here described from specimens taken at Genoa. The identity of the Greenland Hake with *M. merluccius*† is perhaps uncertain. (Eu.) (*merluccius*, ancient name; *mare*, sea; *Lucius*, pike.)

*Gadus merluccius*, LINNÆUS, Syst. Nat., Ed. x, 254, 1758, Europe; after authors.

*Merluccius smiridus*, RAFINESQUE, Caratteri, etc., 26, 1810; JORDAN & GILBERT, Synopsis, 809, 1883; LILLJERORG, Sveriges Fiske, II, 121, 1891.

*Gadus ruber*, LACÉPÈDE, Hist. Nat. Poiss., v, 673, 1803, Scotland; Dieppe; on notes by M. NÜEL; young.

*Gadus merlus*, RISSO, Ichth. Nice, 122, 1810, Nice.

*Onus riali*, RAFINESQUE, Indice d'Ittiol. Sicil., 26, 1810; substitute for *merluccius*.

*Merluccius vulgaris*, FLEMING, Brit. Anim., 195, 1828; GÜNTHER, Cat., IV, 344, 1862.

*Merluccius esculentus*, RISSO, Eur. Mérid., III, 1826, 220, Nice.

! *Merluccius ambiguus*, LOWE, Proc. Zool. Soc. Lond. 1840, 37, Madeira.

*Merluccius sinuatus*, SWAINSON, in Lowe, Proc. Zool. Soc. 1840, 38.

*Merluccius lanatus*, GRONOW, Cat. Fish., Ed. Gray, 130, 1854, Mediterranean.

*Epicopus gayi*, GÜNTHER, Cat., II, 248, 1860, no locality; not *M. gayi*, GUICHENOT, which is the Chilean Hake.

*Merluccius linnei*, MALM, Götheborgs och Bohusläna Fauna, 489, 1877.

## 2900. MERLUCCIUS BILINEARIS (Mitchill).

(SILVER HAKE; NEW ENGLAND HAKE; WHITING.)

Head  $3\frac{1}{2}$ ; depth  $6\frac{1}{2}$ . D. 13-41; A. 40; scales 100 to 110. Top of head with W-shaped ridges very conspicuous; eye shorter than snout and less than interorbital width; maxillary reaching posterior border of pupil; teeth not very large, smaller than in the European species, *Merluccius merluccius*. Scales larger than in other species; pectorals and ventrals long, the latter reaching  $\frac{3}{4}$  distance to vent, their length about  $\frac{2}{3}$  that of head. Grayish, darker above, dull silvery below; axil and edge of pectoral somewhat blackish; inside of opercle dusky silvery; inside of mouth dusky bluish; peritoneum nearly black. Coasts of New England and northward to Straits of Belle Isle; south, in deep water, to the Bahamas; rather common; used as food; breeding in deep water, though often taken near shore, northward. This species resembles the European Hake, *Merluccius merluccius*, but the latter has smaller scales, about 150, and larger teeth. (*bilinearis*, two-lined.)

\* The Iceland Hake has been described as *Merluccius argentatus* (Faber). According to Faber, it has large teeth, the mouth white within, and the rays D. 15-43; A. 51; the fins deeply notched. It is perhaps a valid species, and, if so, it doubtless occurs in Greenland. (*argentatus*, silvered.)

*Gadus merluccius (argentatus)*, FABER, Fische Islands, 90, 1829, Iceland.

*Merluccius argentatus*, GÜNTHER, Cat., IV, 346, 1862.

† "Dans l'Amérique du Nord, on cite ce poisson de Grønland, mais l'exactitude de cette indication paraît douteuse." (Collett, Comp. Sci. Hirondelle, 1896, 58.)

*Stomodon bilinearis*, MITCHILL, Rept. Fish. N. Y., 7, 1814, New York.

*Gadus albidus*, MITCHILL, Journ. Ac. Nat. Sci. Phila., 1, 1817, 409, New York.

*Mertucius albidus*, STORER, Hist. Fish. Mass., 363.

*Mertucius bilinearis*, GOODE & BEAN, Bull. Essex Inst., XI, 9, 1879, JORDAN & GILBERT, Synopsis, 809, 1883; GOODE & BEAN, Oceanic Ichthyology, 386, fig. 330, 1896.

2901. MERLUCCIUS PRODUCTUS (Ayres).

Head  $3\frac{1}{2}$ ; depth 7. D. 11-41; A. 43; V. 7; scales 136. Head with the W-shaped ridges less strongly marked; maxillary reaching center of pupil; eye large; pectorals long and narrow, reaching vent; ventrals much smaller than in *M. bilinearis*, reaching halfway to vent, their length about  $\frac{2}{3}$  that of head; caudal somewhat forked. Scales quite small, deciduous. Teeth moderate. Silver gray; head dusted with coarse black dots; inside of mouth and opercle jet-black; peritoneum silvery, with black specks. Length 3 feet. Pacific coast of America, from Santa Catalina Island northward to Puget Sound; everywhere abundant at moderate depths; used as food. (*productus*, drawn out.)

*Merlangus productus*, AYRES, Proc. Cal. Ac. Nat. Sci. 1855, 64, San Francisco.

*Homalopomus trowbridgii*, GIRARD, Proc. Ac. Nat. Sci. Phila. 1856, 132, Astoria, Oregon. (Coll. Lieut. W. P. Trowbridge.)

*Gadus productus*, GÜNTHER, Cat., IV, 338, 1862.

*Mertucius productus*, GILL, Proc. Ac. Nat. Sci. Phila. 1863, 247; JORDAN & GILBERT, Synopsis, 809, 1883.

Family CCXIV. GADIDÆ.

(THE CODFISHES.)

Body more or less elongate, the caudal region moderate, coniform behind, and with the caudal rays procurrent above and below; vent submedian; suborbital bones moderate; scales small, cycloid; mouth large, terminal; chin with a barbel, more or less developed. Gill openings very wide; gill membranes separated or somewhat united, commonly free from the isthmus; no spines, the fin rays all articulated. Dorsal fin extending almost the length of the back, forming 1, 2, or 3 fins; anal fin long, single or divided; caudal fin distinct, or confluent with the dorsal and anal; ventral fins jugular, but attached to the pubic bone, each of 1 to 8 branched rays. Gills 4, a slit behind the fourth. No pseudobranchia. Edge of preopercle usually covered by skin of head. Pyloric caeca usually numerous, but sometimes few or none. Air bladder generally well developed. Genera about 25, species about 140; an important family, many of its members being highly valued as food. They inhabit chiefly the northern seas, sometimes venturing into the oceanic abysses. One genus (*Lota*) is confined to the fresh waters. (*Gadidæ*, Günther, Cat., IV, 326-369.)

GADINÆ:

a. Anal divided into 2 separate fins; dorsal fin divided into 3.



b. Lower jaw distinctly projecting; barbel small or obsolete; caudal concavo behind.

c. Teeth in upper jaw slender, wide, set in 1 or 2 series; caudal forked.




BOREOGADUS, 976.

cc. Teeth in upper jaw in a-villiform band, the outer somewhat larger; caudal lunate.





**IMAGE EVALUATION  
TEST TARGET (MT-3)**



**Photographic  
Sciences  
Corporation**

23 WEST MAIN STREET  
WEBSTER, N.Y. 14580  
(716) 872-4503

28  
32  
36  
40  
44  
48  
52  
56  
60  
64  
68  
72  
76  
80  
84  
88  
92  
96  
100

10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100

- d. Subopercle and postclavicle normal, both thin and flat, not enlarged and ivory-like. POLLACHIUS, 977.
- dd. Subopercle and postclavicle enlarged, the bone dense and smooth, like ivory. THERAGRA, 978.
- bb. Lower jaw included; barbel well developed; caudal not concave behind.
- e. Hypocoracoid not swollen and ivory-like; lateral line pale; supraoccipital crest moderate.
- f. Transverse processes of vertebræ thickened, swollen, and ivory-like at tip; small codfishes of the Arctic. ELEGINUS, 979.
- ff. Transverse processes of vertebræ not swollen at tip.
- g. Vent in front of second dorsal; size very small. MICROGADUS, 980.
- gg. Vent below second dorsal; typical codfishes of large size. GADUS, 981.
- ee. Hypocoracoid much swollen and ivory-like; lateral line black; mouth small, the maxillary not reaching to opposite eye; supraoccipital crest very high. MELANOGRAMMUS, 982.
- aa. Anal fin forming a continuous fin or sometimes deeply notched; dorsal not divided into 3 fins.
- h. Dorsal fin divided into 2 fins.
- i. Anterior dorsal composed of distinct rays, similar to those in second dorsal.
- j. Ventral fins rather broad, each of about 6 rays.
- MORINÆ:
- k. Anal fin with a deep notch.
- l. Snout not much depressed, its edge without keel; tail slender. LEPIDION, 983.
- u. Snout flat, depressed, keeled on the edge; tail attenuate. ANTIMORA, 984.
- LOTINÆ:
- kk. Anal fin not notched; mouth terminal.
- m. Vomer toothless.
- n. Teeth in jaws unequal, outer series enlarged.
- o. Barbel obsolete. URALOPTUS, 985.
- oo. Barbel well developed. LOTELLA, 986.
- nn. Teeth in jaws all villiform; barbel developed. PHYSICULUS, 987.
- mm. Vomer with teeth; head not compressed.
- p. Vomer and mandible without canines. Fresh-water species. LOTA, 988.
- pp. Vomer and mandible armed with canines. Deep-water species. MOLVA, 989.
- PHYCINÆ:
- jj. Ventral rays very slender, each of 1 or 2 rays.
- q. Ventrals each of 2 or 3 slender rays. UROPHYCIS, 990.
- qq. Ventrals each of a single bifid ray. LÆMONEMA, 991.
- GAIDROPSARINÆ:
- ii. Anterior dorsal formed of a single slender ray, followed by a band of fringes; ventrals each of 5 to 7 rays.
- r. Barbels 3; snout with 2 barbels, 1 at each nostril, none at tip; chin with 1 barbel. GAIDROPSARUS, 992.
- rr. Barbels 4; snout with 3 barbels, 1 at tip of snout and 1 on each nostril; chin with 1 barbel, head high and compressed; no canines. ENCHELYOPUS, 993.
- BROSMINÆ:
- hh. Dorsal fin continuous, undivided; ventrals several-rayed; teeth on jaws, vomer, and palatines; mouth large; frontal bone. BROSME, 994.

976. BOREOGADUS, Günther.

*Boreogadus*, GÜNTHER, Cat. Fish. Brit. Mus., IV, 336, 1862 (*fabricii*).

This genus is closely allied to *Pollachius*, the body more slender, the caudal fin more deeply forked, and the teeth in both jaws slender, sharp, wide set, in 1 or 2 series. Small codfishes of the Arctic. (*βόρεος*, northern; *Gadus*.)

2902. BOREOGADUS SAIDA (Lepechin).

Head  $3\frac{1}{2}$ ; depth  $5\frac{1}{2}$ ; eye 4 in head; snout  $3\frac{1}{2}$ ; interorbital space  $4\frac{1}{2}$ ; gill rakers 9 to 13 + 30 to 32. D. 13-14-20; A. 16-21. Body slender, little compressed; head long, rather pointed, the lower jaw projecting; barbel minute; maxillary reaching middle of pupil; mandible 2 in head; teeth in upper jaw in 1 series, except in front, when the row is double; teeth in lower jaw uniserial; teeth nearly uniform in size, sharp, and wide set; teeth on vomer few, similar to those in jaws. Gill rakers numerous, long and slender, the longest  $\frac{1}{2}$  eye; vent slightly before second dorsal; caudal peduncle slender, rounded, its depth scarcely more than  $\frac{1}{2}$  eye. Pectorals reaching vent,  $1\frac{1}{2}$  in head; ventrals  $1\frac{1}{2}$ , the second ray exerted for  $\frac{2}{3}$  its length; first dorsal highest; front of second dorsal midway between tip of snout and base of caudal. Caudal forked for a distance equal to  $\frac{1}{2}$  eye, the tips rounded. Color plain brownish, silvery below, the body with fine black points, most numerous above; dorsals and caudals dusky, the rays blackish distally, their edge narrowly white; anal similarly colored, pale at base; pectoral uniform dusky, pale-edged; ventrals somewhat dusky; peritoneum blackish. Length 6 to 8 inches. Arctic seas of Asia and America, from Greenland to Siberia; generally common in the far North, but rare in Bering Sea and south of Greenland. Here described (by Mr. Norman B. Scofield) from specimens from Davis Straits and Melville Bay, Greenland, the largest  $6\frac{1}{2}$  inches long, and from specimens taken by Mr. Scofield at Point Barrow, Port Clarence, and Herschel Island. There is no difference between Greenland and Alaskan specimens. The range of fin rays is D. 12 to 15-12 to 15-18 to 22; A. 15 to 18-20 to 22. Concerning its habits Mr. Scofield observes:

"This fish appears to be quite abundant north of Bering Straits. It was especially brought to our notice by its habit of hiding in small holes in the floating ice, from which it was dislodged by our steamer striking and turning over the blocks of ice. This floating ice was usually in 7 fathoms of water and 1 or 2 miles from the coast. At Herschel Island we took it with the seine in shallow water along the beach. Lucien H. Turner reports it from St Michaels, where he took it through the ice in February, and was told by the natives that it appeared there only in winter. According to Richardson it spawns in Greenland in February, laying its eggs in the seaweeds along the shore under the ice." According to Richardson, in Northumberland Sound, "when hotly pursued by the Boluga or white whale, it has been observed, in its endeavors to escape, to leap by hundreds on the ice." (Eu.) (*saida*, Russian name.)



- Gadus saida*, LEPECHIN, N. v. Comm. Ac. Sci. Petrop. 1774, 512, White Sea; PALLAS, Zoogr. Rosso Asiat., III, 199, 1811; GÜNTHER, Cat., IV, 337, 1862; COLLETT, Norske Nord-Havs Exped., 126, 1880; JORDAN & GILBERT, Synopsis, 307, 1883.
- Merlangus polaris*, SABINE, Supp. Parry's Voyage, CCXI, 1824, Baffins Bay; RICHARDSON, Last Arctic Voyage, 27, 1824.
- Gadus fabricii*, RICHARDSON, Fauna Bor.-Amer., III, 245, 1836, northern bays of Greenland; after *Gadus aeglefinus* of Fabricius.
- Gadus agilis*, REINHARDT, Dansko Vid. Selsk. Aft., VII, 126, 1838, Greenland.
- Gadus glacialis*, PETERS, Nord Pol. Exped., II, 172, 1874.
- Pollachius polaris*, GILL, Cat. Fish. East Coast N. A., 218, 1861.
- Boreogadus polaris*, GILL, Proc. Ac. Nat. Sci. Phila. 1863, 233.
- Boreogadus saida*, BEAN, Bull. U. S. Nat. Mus., IV, 108, 1879; SCOFIELD, in Jordan & Gilbert, Rept. Fur Seal Invest., 1898.

## 977. POLLACHIUS, Nilsson.

(POLLACKS.)

*Pollachius*, NILSSON, in Bonaparte, Catalogo Metodico Pesci Europ., 45, 1846 (*pollachius*).

Body rather elongate, covered with minute scales; mouth moderate or large, the lower jaw projecting; barbel very small or obsolete; villiform teeth on vomer, none on palatines; teeth in jaws equal or the outer slightly enlarged; gill membranes more or less united; subopercle and postclavicle not enlarged and not ivory-like; dorsal fins 3; anal 2; caudal lunate; vent under first dorsal. Large fishes of the northern seas. (Plog or Pollack, the English vernacular name, latinized as *Pollachius*, as though derived from  $\pi\omicron\lambda\lambda\alpha\chi\tilde{\eta}$ , many fashioned.)

## 2903. POLLACHIUS VIRENS (Linnaeus).

(POLLACK; COAL-FISH; GREEN COD.)

Head 4; depth 4½. D. 13-22-20; A. 25-20; scales about 150; vertebrae 54. Body rather elongate, compressed; snout sharp and conic; mouth rather small, oblique; maxillary reaching beyond front of orbit; lower jaw slightly the longer; teeth in the upper jaw nearly equal, the outer series not being especially enlarged; barbel rudimentary or obsolete; gill membranes considerably united, free from isthmus; vent under first dorsal; caudal fin lunate; pectorals short, scarcely reaching anal; ventrals short, their origin in front of base of pectoral a distance about equal to diameter of eye. Greenish brown above; sides and below somewhat silvery; lateral line pale; fins mostly pale; sometimes a black spot in the axil. North Atlantic; common northward on both coasts, south to Cape Cod and France. (Eu.) (*virens*, green.)

*Gadus virens*, LINNÆUS, Syst. Nat., Ed. x, 253, 1758, Seas of Europe; after *Gadus tripterygius imberbis* of the Fauna Suecica; GÜNTHER, Cat., IV, 339; JORDAN & GILBERT, Synopsis, 807, 1882.

*Gadus carbonarius*, LINNÆUS, Syst. Nat., Ed. x, 254, 1758, seas of Europe; after *Gadus dorso tripterygius imberbi* of ARTEDI.

*Gadus colinus*, LACÉPÈDE, Hist. Nat. Poiss., II, 416, 1800, England, etc.; after LE COLIN of Danberton.

*Gadus virens*, LACÉPÈDE, Hist. Nat. Poiss., II, 417, 1800.

*Merlangus purpureus*, MITCHELL, Trans. Lit. and Phil. Soc., I, 1815, 370, New York.

*Merlangus leptocephalus*, DE KAY, New York Fauna: Fishes, 288, pl. 45, fig. 146, 1842, Long Island.

*Merlangus purpureus*, STORER, Rept. Fish. Mass., 130, 1830.

*Pollachius carbonarius*, GILL, Proc. Ac. Nat. Sci. Phila. 1863, 233.

978. THERAGRA, Lucas, a new genus.

(ALASKAN POLLACKS.)

*Theragra*, LUCAS, in Jordan & Gilbert, Rept. Fur Seal Invest. 1896 (1898) (*chalcogrammus*).

This genus is closely allied to *Pollachius*, differing in the following respects: Suboperculum thick, smooth and dense instead of being thin and squamous as in *Pollachius*; the postclavicle is also similar in structure while its proximal portion is subcircular in *Theragra* and rhomboidal in *Pollachius*; this ivory-like character of the suboperculum and postclavicle is so marked that it serves to distinguish these bones at a glance, being entirely different from what is found in the corresponding bones of other gadoids. The Alaskan Pollack farther differs from the Atlantic Pollack in having 19 precaudal vertebrae and 33 caudal, instead of 23 precaudals and 32 caudals; the bodies of the vertebrae are also slightly longer and more deeply sculptured in the Alaskan fish and the spinous process of the anterior dorsals less elevated. The vertebral differences between the 2 genera are merely differences of degree and of specific value only, but the differences between the subopercula and postclavicles are different in kind, distinguishing the Alaskan Pollack not only from the Atlantic Pollack, but from other gadoids. (*θύρα*, best; *ἄγρᾱ*, prey or food; the Alaskan Pollack being a chief food of the fur seal, *Callorhinus*.)

a. Dorsal rays about 13 or 14-17-18 or 19; anal rays 20-20; side with 2 interrupted dark longitudinal bands. CHALCOGRAMMA, 2904.

aa. Dorsal rays 10 or 11-13 to 15-16; anal rays 16 to 19-16 to 19; sides plain dusky; body less elongate, the snout blunter, the fins lower. FUCENSIS, 2905.

2904. THERAGRA CHALCOGRAMMA (Pallas).

(ALASKA POLLACK.)

Head 4; depth 6. D. 12-14-18; A. 20-20. Eye 5 in head; snout  $3\frac{1}{2}$ ; maxillary  $2\frac{1}{2}$ ; snout conic, sharp, rounded in profile; mouth oblique; maxillary reaching middle of pupil; chin with a minute barbel; teeth small, those of the outer row above slightly enlarged; eye large, wider than the flat interorbital space, 4 in head. Gill membranes somewhat united, the posterior outline deeply emarginate; vent under interspace between first and second dorsal; first dorsal higher than the others, the second lowest; ventrals filamentous, reaching nearly to vent; pectorals long, reaching past front of anal,  $1\frac{1}{2}$  in head; caudal somewhat concave. Olivaceous above, sides silvery, with 2 interrupted stripes of dark, brassy,

olive along sides, these irregular on their edges, about  $\frac{1}{2}$  width of eye, with uneven edges; a trace of a third similar stripe below anteriorly, the stripes very irregular; back mottled. Dorsal plain dark olive; pectoral quite dark; lower fins ashy; caudal ashy olive. Bering Sea and neighboring waters, probably south to Sitka and the Kurils. Our specimens from Unalaska, Robben Reef, Komandorski and Pribilof islands and Bristol Bay. Excessively common throughout Bering Sea, swimming near the surface, and furnishing the greater part of the food of the fur seal. This animal rarely catches the true codfish, which swims nearer the bottom. Length 3 feet. (*χαλκός*, brass; *γραμμή*, line.)

*Gadus chalcogrammus*, PALLAS, Zoogr. Rosso-Asiat., III, 198, 1811, Kamchatka; GÜNTHER, Cat., IV, 340, 1862; JORDAN & GILBERT, Synopsis, 807, 1883.

*Gadus periscopus*, COPE, Proc. Am. Philos. Soc. Phila. 1873, 30, Unalaska (Coll. George Davidson).

*Pollachius chalcogrammus*, JORDAN, Cat. Fish. N. A. (130) 918, 1885.

*Theragra chalcogramma*, JORDAN & GILBERT, Rept. Fur Seal Invest., 1898.

2905. *THERAGRA FUCENSIS* (Jordan & Gilbert).

(WALL-EYED POLLACK; PUGET SOUND POLLACK.)

Head  $3\frac{1}{2}$  in body; depth  $5\frac{1}{2}$ . D. 10-13-16 to 11-15-16; A. 16-19 to 19-19; eye  $4\frac{1}{2}$  in head; maxillary  $2\frac{3}{4}$ ; pectoral  $1\frac{3}{4}$ ; longest caudal ray 2. Body elongate, not greatly compressed; mouth large, the maxillary reaching to below middle of eye; jaws with minute, sharp, curved teeth, the outer series enlarged; teeth on vomer, palatines toothless; lower jaw projecting, a very small barbel under its tip; interorbital space wide, very slightly and evenly convex, wider than the diameter of eye; nostrils much nearer eye than tip of snout, the posterior much the larger; head almost entirely covered with small scales; gill rakers numerous, the longest as long as pupil, about  $5+27$  in number. Distance of origin of first dorsal from snout  $3\frac{1}{2}$  in body; first rays of first dorsal reaching far past the ends of last rays where fin is depressed; first rays of other dorsals and anals scarcely reaching the base of last rays; caudal slightly forked or subtruncate when spread, the lobes subequal; end of pectoral reaching to front of anal; ventrals inserted in front of base of pectoral in distance a little more than diameter of eye, ending in a filamentous point. Color nearly plain sooty, with no distinct lateral bands, and with generally only a trace of a pale lateral streak along the side; on the head some diffuse dark spots; fins all dusky. The band of teeth in the premaxillary is wider than in *Theragra chalcogramma*, and the band is widened at the anterior end; the body is shorter; eye smaller; color darker; fins not so high; caudal not so deeply forked. Pacific coast, from Vancouver Island to Monterey, abundant in Puget Sound; probably northward to Kadiak, replacing *T. chalcogramma* to the southward. This form may intergrade with *Theragra chalcogramma*, though the original types seem well separated. Little is known of its range to the northward. Scofield and Seale took a specimen in Chignik Bay in northern Alaska, which

seems as near *T. fucensis* as *T. chalcogramma*. Its rays\* are D. 11-16-17; A. 18-17; ventrals reaching  $\frac{3}{4}$  distance to vent; interorbital space wider than eye; coloration dark. But its body is as slender as in *T. chalcogramma*. (*fucensis*, from the straits of Juan de Fuca.)

*Pollachius chalcogrammus fucensis*, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1893, 315, Puget Sound at Tacoma. (Type, No. 44455. Coll. David H. Hume.)

979. ELEGINUS, Fischer.

*Eleginus*, FISCHER, Mem. Soc. Nat. Moscow, v, 4, 2d Ed., 252-257, 1813 (*navaga*); not *Eleginus* of later authors.

*Tilesia*, SWAINSON, Nat. Hist. Class'n Fishes, II, 300, 1839 (*gracilis*); name preoccupied.

*Pleurogadus*, BEAN, in JORDAN, Cat. Fish. N. A., 130, 1885 (*gracilis*); substitute for *Tilesia* preoccupied.

This genus differs from the other codfishes in the structure of the transverse processes of the vertebrae, which are club-shaped, narrow at base, but expanding distally into a rounded hollow bulb at their tips. Skeleton otherwise essentially as in *Microgadus*, the skull similar. Small codfishes of the Arctic seas. (*ἐλεγιῖνος*, a social fish mentioned by Aristotle.)

2906. ELEGINUS NAVAGA (Kölræuter).

(WACHNA COD.)

Head  $3\frac{1}{2}$  in length of body; depth 6; eye  $5\frac{1}{2}$  in head; snout 3; interorbital space  $4\frac{1}{2}$ ; gill rakers 20 or 21; barbel small, equal to pupil; dorsal 13-18-18; anal 22-20; scales small, 157 transverse rows above lateral line from gill opening to first rudimentary caudal rays. Body slender and rounded with a rather long head; snout viewed from above rounded, but running to a rather sharp point when viewed from the side; lower jaw included, the fleshy snout projecting beyond the maxillary, its length slightly greater than that of the snout; tip of maxillary on a vertical with the front of the pupil; articulation of mandible with quadrate bone on a vertical running midway between pupil and posterior edge of eye; teeth

\* The following is the count of fin rays in 13 specimens of *Theragra* of the two species:

Dorsal.	Anal.	Locality.
<i>T. chalcogramma</i> :		
13-15-20.....	19-20	Kamchatka.
13-15-20.....	19-21	Unalaska.
14-19-23.....	24-22	Pribilof Islands.
14-16-21.....	21-23	Do.
14-17-18.....	23-21	St. Paul Island.
13-15-19.....	21-22	Do.
14-17-19.....	21-20	Do.
14-17-18.....	22-20	Kamchatka.
11-16-17.....	19-17	Chignik Bay.
<i>T. fucensis</i> :		
10-15-17.....	18-16	Puget Sound.
11-15-16.....	19-18	Do.
10-14-16.....	16-19	Do.
12-13-17.....	19-19	Do.

all slender and curved backward, those in upper jaw in several irregular rows, the outer row regular and with slightly larger teeth; teeth in lower jaw in a single row except in front where they are in a double row; teeth on vomer few and about the size of the smaller teeth in the upper jaw; gill rakers moderate, the longest not quite equal to diameter of pupil; caudal peduncle compressed, its depth equal to diameter of eye; vent under front of second dorsal; pectoral fin not reaching vent, its length  $1\frac{1}{2}$  times in head; ventrals reaching halfway to vent, the second ray moderately produced; first dorsal highest; distance between second and third dorsals twice distance between first and second; caudal fin very slightly concave; third ray of second dorsal midway between tip of snout and base of middle caudal rays. Color somewhat mottled, grayish brown above, light silvery below; the 3 dorsals and caudal dusky and edged with white; pectorals uniform dusky; ventrals but slightly dusted with black; anal with a few punctulations at their anterior ends; peritoneum pale. Arctic shores of Asia and North America, south to Bering Sea, locally abundant. It reaches the length of about a foot. Here described (by Norman B. Scofield) from numerous specimens, the largest 11 inches long, taken at Port Clarence by Scofield and Seale, and at Petropaulski by the *Albatross* (Fur Seal Invest. of 1896). The range of the fin rays is D. 12 to 15—18 to 21—18 to 21; A. 20 to 23—20 to 23. Mr. Scofield has prepared a skeleton of this species for comparison with that of *Microgadus proximus* from San Francisco. There is very little difference in the skulls. There is no difference in the neural spines of the vertebrae. The transverse processes of the vertebrae in *Microgadus proximus* are flattened and plate-like, while in *Eleginus navaga* they are club-shaped, narrow at base where they leave the centrum, but expanding into a rounded hollow bulb at the distal end. This character defines the genus *Eleginus*. (*navaga*, a Russian name.)

*Gadus navaga*, KÖLBEUTER, Nov. Comm. Ac. Petrop., XIV, 1770, 484, pl. 12, coast of northern Russia; PALLAS, Zoogr. Rosso-Asiat., III, 196, 1811.

*Gadus gracilis*, TILESIIUS, Mém. Ac. Imp. Petersb., II, 1810, 354, Kamchatka; JORDAN & GILBERT, Synopsis, 804, 1883.

*Gadus wachna*, PALLAS, Zoogr. Rosso-Asiat., III, 182, 1811, Kamchatka.

*Tilesia gracilis*, SWAINSON, Nat. Hist. Fish., II, 300, 1839; BEAN, Proc. U. S. Nat. Mus. 1881, 243.

*Pleurogadus gracilis*, BEAN, in JORDAN, Cat. Fish. N. A., 130, 1885.

*Eleginus navaga*, GILL, Proc. U. S. Nat. Mus. 1890, 303.

### 980. MICROGADUS, Gill.

(TOMCODS.)

*Microgadus*, GILL, Proc. Ac. Nat. Sci. Phila. 1865, 69 (*proximus*).

Very small codfishes allied to *Gadus*, but with the vent placed before the second dorsal and with a different structure of the cranium. The following is Professor Gill's account of the skull of *Microgadus proximus*, the italicised portions indicating the differences from *Gadus*:

The cranium is proportionally broader toward the front and less flattened, while the brain case is flattened below, *decidedly swollen* on each

side of a depressed *sphenoidal groove*, and has an ovate cardiform shape; the *paraoccipital* or *epiotic* is not produced into an angle behind, but is obtusely rounded, and its posterior or *outwardly descending ridge blunt*; the *opisthotic* is well developed, oblong, and with its recentering angle *high up*, and, on a line with it, the surface is divided into 2 parts—a narrow and a flattened one, and a lower expanded one, much swollen; the *alisphenoid* or *prootic* is *oblong*, acutely emarginate in front, swollen from the region of the high anterior sinus, and above a little produced forward; the great *frontal* is a little longer than broad, with supraoccipital crest *continued forward* on the bone, and near the front expanded upward, and with the *expanded portion* behind dividing into narrow *lateral wings*; the lateral testiform ridges of the frontal are continued forward and *curved outward* toward the antero-lateral angles; the anterior frontals are *mostly covered in front* by the great frontal, and are much *developed* in the direction of the antero-lateral angles, the inferior expanded axillary portion being very narrow; the nasal has a rounded ridge in front, continued well below, and its posterior crest is *laminar* and trenchant.

Species American; valued as food. (*μικρός*, small; *γάδος*, *Gadus*.)

a. Second anal with 21 or 22 rays; snout rather long; body semitranslucent; first anal and ventrals pale; body scarcely blotched with blackish. PROXIMUS, 2907.

aa. Second anal with 16 to 20 rays; snout shorter; body opaque; first anal and ventrals dusky; body blotched above with blackish. TOMCOD, 2908.

2907. MICROGADUS PROXIMUS (Girard).

(CALIFORNIA TOMCOD.)

Head  $3\frac{1}{2}$  in body; depth 5. D. 14–18–18 to 21; A. 21 or 22–21 or 22; V. 6–7; eye 5 in head; maxillary  $2\frac{1}{2}$ ; pectoral 2; highest dorsal spine 2; middle caudal rays equal to snout. Head long, convex above, somewhat compressed, with vertical sides; eye moderate; mouth rather large; maxillary reaching to below pupil; barbel small; teeth in each jaw in a band, the outer row a little enlarged. Gill membranes a little connected, free from the isthmus. First dorsal highest, somewhat falcate; first anal longer and higher than second; pectorals moderate, reaching anal; ventrals filamentous, scarcely reaching anal; caudal slightly emarginate or subtruncate when fin is spread. Lateral line very distinct, wavy, high anteriorly, slightly interrupted posteriorly. Vent below first dorsal. Color olivaceous above, pale, or slightly translucent white below; dorsal fins dusky, paler at base; first anal and ventrals uncolored; second anal dusted with dark points. Monterey to Unalaska; abundant; a food-fish of considerable importance, the flesh delicate but without much flavor. Here described from a specimen, 8 inches in length, from Alaska, Albatross Station 3213. It reaches the length of about a foot. (*proximus*, near, to *Microgadus tomcod*.)

*Gadus proximus*, GIRARD, Proc. Ac. Nat. Sci. Phila. 1854, 141, San Francisco; GIRARD, U. S.

Pac. R. R. Surv., X, Fishes, 142, 1858; JORDAN & GILBERT, Synopsis, 805, 1883.

*Morrhua californica*, AYRES, Proc. Cal. Ac. Nat. Sci. 1854, 9, San Francisco.

*Gadus californicus*, GÜNTHER, Cat., IV, 332, 1862.

*Microgadus proximus*, GILL, Proc. Ac. Nat. Sci. Phila. 1865, 69.

## 2908. MICROGADUS TOMCOD (Walbaum).

(TOMCOD; FROSTFISH.)

Head 4 in body; depth 5. D. 13 to 15-15 to 19-16 to 18; A. 17 to 21-16 to 20; eye 5 in head; maxillary  $2\frac{1}{2}$ ; pectoral 13; middle caudal rays  $2\frac{1}{2}$ ; first dorsal rays 13. Snout rounded, less produced than in *Microgadus proximus*; mouth short; maxillary  $2\frac{1}{2}$  in head, reaching pupil; eye large,  $3\frac{1}{2}$  in head; barbel small; pectorals reaching vent; ventrals filamentous, not reaching vent; vent under interval between first and second dorsals. Color olive brown, distinctly blotched and spotted with darker, lighter on the belly; more opaque than in *M. proximus*; back and sides profusely punctulate; dorsals and caudal blotched with darker; anals coarsely punctulate anteriorly, colorless posteriorly; ventrals and pectorals dusky. Virginia to Labrador; very common northward, and valued as a food-fish. Here described from a specimen, 9 inches in length, from Boston, Massachusetts. Length 1 foot. (*tomcod*, a vernacular name.)

*Tomcod*, SCHÖPF, Schrift. Naturf. Freunde, VIII, 140, 1780, New York.

*Gadus tomcod*, WALBAUM, Artedi Piscium, III, 133, 1792, after SCHÖPF; JORDAN & GILBERT, Synopsis, 806, 1883.

*Gadus frost*, WALBAUM, Artedi Piscium, III, 134, 1792, North America; after Frost-fish of Pennant.

*Gadus tomcodus*, MITCHILL, Trans. Lit. and Phil. Soc., I, 1815, 368, New York; GÜNTHER, Cat., IV, 331, 1862.

*Gadus pruinus*, MITCHILL, Trans. Lit. and Phil. Soc., I, 1815, 368, New York.

*Gadus tomcodus fuscus*, MITCHILL, Trans. Lit. and Philos. Soc., I, 1815, 369, New York.

*Gadus tomcodus luteus*, MITCHILL, Trans. Lit. and Philos. Soc., I, 1815, 369, New York.

*Gadus tomcodus mixtus*, MITCHILL, Trans. Lit. and Philos. Soc., I, 1815, 369, New York.

*Gadus polymorphus*, MITCHILL, Trans. Lit. and Philos. Soc., I, 1815, 369, New York.

*Morrhua americana*, STORER, Rept. Fish. Mass., 120, 1839, coast of Massachusetts.

## 981. GADUS (Artedi) Linnæus.

(CODFISHES.)

*Gadus*, LINNÆUS, Syst. Nat., Ed. x, 251, 1758 (*morrhua*); after ARTEDI.

*Morrhua*, OKEN, Isis 1817, 1182 (*morrhua*; on *les Morrhues* of CUVIER).

*Cephus*, SWAINSON, Nat. Hist. Class'n Fishes, II, 300, 1839 (*macrocephalus*).

Body moderately elongate, compressed and tapering behind. Scales very small; lateral line present, pale. Head narrowed anteriorly; mouth moderate, the maxillary reaching past front of eye; chin with a barbel; teeth in jaws cardiform, subequal; vomer with teeth; none on the palatines; cranium without the expanded crests seen in *Melanogrammus*; no part of the skeleton expanded and ivory-like. Dorsal fins 3, well separated; anal fins 2; ventral fins well developed, of about 7 rays. Species of the Northern Seas; highly valued as food. (*Gadus*, the Latin name, akin to the English word cod.)

a. Eye moderate, about  $\frac{1}{2}$  snout in adult; axil without dusky spot.

b. Air bladder large. Atlantic codfish.

bb. Air bladder small. Pacific codfish.

aa. Eye large, more than  $\frac{1}{2}$  length of snout; axil with a dusky spot; caudal peduncle slender.

CALLARIAS, 2909.

MACROCEPHALUS, 2910.

OGAC, 2911.

B  
ma  
Hea  
helo  
teet  
bita  
Gill  
Pec  
reac  
base  
seco  
and  
anal  
  
\* W  
mon  
name  
the n  
are o  
but f

2900. GADUS CALLARIAS,\* Linnæus.

(COMMON CODFISH.)

Head  $3\frac{1}{2}$  to  $4\frac{1}{2}$ ; depth about 4. D. 14–21–19; A. 20–18. Head large, but varying much in size; maxillary about reaching middle of orbit; occipital keel not greatly developed; teeth strong, cardiform, in narrow bands, those of the outer row in the upper jaw and of the inner row in the lower jaw somewhat enlarged. Eye moderate, about  $\frac{1}{2}$  length of snout. First dorsal little elevated, its height about  $\frac{1}{2}$  length of head; vent under front of second dorsal; caudal slightly emarginate; pectorals  $\frac{1}{2}$  length of head. Greenish or brownish, subject to many variations, sometimes yellowish or reddish; back and sides with numerous rounded brownish spots; lateral line pale; fins dark. North Atlantic, south to Virginia, and France; one of the most important of food-fishes. (Eu.) (*Callarias*, an old name of the codfish.)

*Gadus callarias*, LINNÆUS, Syst. Nat., Ed. x, 252, 1758, young examples, Baltic Sea and oceans of Europe, after *Gadus*, etc., *cauda integra* of the Fauna Suecica; CUVIER, Règne Animal, Ed. 2, vol. II, 332, 1829; JORDAN & GILBERT, Synopsis, 304, 1883.

*Gadus morhua*, LINNÆUS, Syst. Nat., Ed. x, 252, 1758, seas of Europe, after *Gadus*, etc., *cauda subæquali* of the Fauna Suecica; RICHARDSON, Fauna Bor.-Amer., 242, 1836.

*Gadus barbatus*, LINNÆUS, Syst. Nat., Ed. x, i, 252, 1758.

*Gadus vertagus*, WALBAUM, Artedi Pisc., III, 143, 1792; after Jägershen, KLEIN, Hist. Nat. Pisc., v, 7, pl. 2, fig. 1, 1749.

?*Gadus heteroglossus*, WALBAUM, l. c., 144; after Hornbogen of KLEIN.

*Gadus arenosus*, MITCHILL, Trans. Lit. and Philos. Soc., I, 1815, 368, New York.

*Gadus rupestris*, MITCHILL, Trans. Lit. and Philos. Soc., I, 1815, 368, New York.

?*Gadus nanus*, FABER, Fische Islands, 113, Iceland.

*Morrhua americana*, STORER, Hist. Fish Mass., 343, 1867.

*Gadus morrhua*, GÜNTHER, Cat., IV, 328, 1862; GOODE & BEAN, Oceanic Ichthyology, 354.

2910. GADUS MACROCEPHALUS, Tilesius.

(ALASKA CODFISH.)

Head 3 in body; depth  $4\frac{1}{2}$ . D. 13–18–16. A. 21–17; eye 6 in head; maxillary  $2\frac{1}{2}$ ; highest dorsal ray 3; pectoral  $2\frac{1}{2}$ ; middle caudal rays 4. Head large, the snout blunt; mouth large, the maxillary reaching to below front of pupil, snout projecting beyond mouth, lower jaw included; teeth strong, cardiform, in narrow bands on jaws and vomer; interorbital wide,  $1\frac{1}{2}$  times wider than diameter of eye, very slightly convex. Gill rakers moderate, about equal to pupil in length,  $3 + 17$  in number. Pectoral reaching to below end of first dorsal; ends of first dorsal rays reaching second dorsal when fin is depressed; ventrals inserted in front of base of pectorals in distance equal to diameter of eye; veins under front of second dorsal; caudal subtruncate. Color brownish, lighter below, back and sides with numerous brownish spots; fins, with the exception of first anal and ventrals, dusky. This species is very abundant in Bering Sea,

\* We retain the name *Gadus callarias*, Linnæus for the codfish, instead of the commonly used name *Gadus morhua*, applied by Linnæus to the same species, because the name *Gadus callarias* stands first on the page on which it occurs. To accord priority to the name standing first is essential to fixity, and not the less so if the competing names are of the same actual date, published by the same author. It is not justice nor elegance, but fixity, which the rules of nomenclature aim to secure.



on both shores, in 15 to 130 fathoms, forming an important article of commerce. Its range southward extends to the offshore banks of Oregon. In external respects we recognize no distinction between this species and the common eastern codfish, except that the head seems larger. Here described from a specimen 20 inches long, taken in the Straits of Fuca by the *Albatross*. Concerning this species Dr. Gilbert observes:

It has been frequently pointed out, and is well known to fishermen that the Pacific codfish has a smaller air bladder or sound than the Atlantic cod. Pending an examination of this question, which we are not now in a position to make, we propose to recognize the Pacific fish as a distinct species. (*μακρός*, long; *κεφαλή*, head.)

*Gadus macrocephalus*, TILESIIUS, Mém. Acad. Sci. St. Petersburg, II, 1810, 360, Kamchatka; GÜNTHER, Cat., IV, 330, 1862.

*Gadus pygmaeus*, PALLAS, Zoogr. Rosso-Asiat., III, 1811, Kamchatka.

*Gadus auratus*, COPE, Proc. Am. Philos. Soc., 1873, 30, Unalaska.

#### 2911. GADUS OGAC, Richardson.

(GREENLAND CODFISH.)

Head  $3\frac{1}{2}$ . D. 14 or 15-18 to 20-17 to 20; A. 20 to 22-18 to 19; V-6. This species resembles the common cod (*Gadus callarias*), but differs from it as follows: It has a more slender caudal peduncle, larger eye, greater interorbital width, longer barbel, more advanced position of ventral fins, and a longer pectoral fin. Color dark, blackish brown above, lighter below, with yellowish marblings; the tip of the dorsal, anal, and caudal fins black; ventral and pectorals dark brown or black, a dusky spot on the axil; barbel black. Coast of Greenland; not seen by us. The above notes from specimens collected at Godhavn, Greenland, examined by Ensign Dresel. (*ogac*, a native name.)

*Gadus ogac*, RICHARDSON, Fauna Bor.-Amer., 246, 1836, Greenland; REINHARDT, Vid. Selsk. Naturvid. Math. Afh. 1838; DRESEL, Proc. U. S. Nat. Mus. 1884, 246.

*Gadus ogat*, KRÖYER, Voy. Scand. et Lap., pl. 19.

#### 982. MELANOGRAMMUS, Gill.

(HADDOCKS.)

*Melanogrammus*, GILL, Proc. Ac. Nat. Sci. Phila. 1862, 280 (*eglefinus*).

*Eglefinus*, MALM, Götheborgs och Bohusläns Fauna, 481, 1877 (*eglefinus*).

This genus is distinguished from *Gadus* by its smaller mouth, the produced first dorsal fin, black lateral line, and especially by the great enlargement of the hypocoracoid, which is dense and ivory-like. The lateral line is always black, and the supraoccipital and other crests on the head are largely developed. Food fishes of large size. (*μέλας*, black: *γραμμή*, line.)

#### 2912. MELANOGRAMMUS EGLEFINUS, Linnæus.

(HAI DOCK.)

Head  $3\frac{1}{2}$ ; depth  $4\frac{1}{2}$ . D. 15-24-21; A. 23-21. Snout long and narrow, overlapping the small mouth; maxillary barely reaching front of orbit;

teeth subequal, large, in a cardiform band in upper jaw; in a single series on lower jaw and on vomer; occiput carinated; a ridge extending backward from each orbit; eye very large,  $\frac{3}{4}$  length of snout, 4 in head. Anterior rays of first dorsal elevated,  $\frac{3}{4}$  length of head, the fin pointed, higher than second and third dorsals; caudal lunate; vent below front of second dorsal. The skull in this species is more depressed than in *Gadus callarias*, broader, and thinner in texture; occipital crest exceedingly high, much higher than in *Gadus*, the wing-like projections at its base anteriorly spreading widely, raised above the surface of the skull. Dark gray above, whitish below; lateral line black; a large dark blotch above the pectorals; dorsals and caudal dusky. North Atlantic, on both coasts, south to France and North Carolina; in deeper water to Cape Hatteras; an important food-fish, reaching a considerable size. (Eu.) (*eglefinus*, an old name of the haddock, from the French Aiglefin or Aigrefin, according to Bellon; perhaps from *aigre faim*, extremely hungry, voracious.)

*Gadus eglefinus*, LINNÆUS, Syst. Nat., Ed. x, 251, 1759, seas of Europe, after *Gadus*, etc., *cauda biloba*, of the Fauna Suecica; JORDAN & GILBERT, Synopsis, 803, 1883.

*Morrhua eglefinus*, FLEMING, British Animals, 191, 1828.

*Morrhua punctatus*, FLEMING, British Animals, 192, 1828.

*Melanogrammus eglefinus*, GILL, Proc. Ac. Nat. Sci. Phila. 1862, 280; *ibid.* 1863, 237; GOODE & BEAN, Oceanic Ichthyology, 354, 1896.

*Eglefinus linnæi*, MALM, Götteborgs och Bohusläns Fauna, 481, 1877.

### 983. LEPIDION, Swainson

*Lepidion*,\* SWAINSON, Nat. Hist. Class'n Anim., I, 318, 1838, and II, 300, 1839 (*lepidion*).

*Haloporphyrus*, GÜNTHER, Cat. Fish. Brit. Mus., IV, 358, 1862 (*lepidion*).

Body elongate, covered with small scales; head not greatly depressed, higher than broad; the snout subconical, obtusely rounded; tail tapering behind; jaws with bands of villiform teeth; a roundish patch of teeth on vomer; no teeth on palatines; chin with a barbel; branchiostegals 7. Caudal fin separate; 2 dorsal fins and 1 anal; the first dorsal short; ventrals narrow, of 6 rays. Deep waters. The American species distinguished from the *Lepidion lepidion* (Risso), of the Mediterranean, by its non-filamentous first dorsal. (*λεπίδιον*, diminutive of *λεπίς*, scale:—small-scaled.)

### 2913. LEPIDION VERECUNDUM, Jordan & Cramer.

Head  $3\frac{1}{2}$ ; depth  $4\frac{1}{4}$ . D. VIII-40; A. 37; V. apparently 4 (some rays broken on each side); scales about 75, not to be exactly counted. Body robust, compressed, tapering from the large head to the very slender, attenuate tail, which is not so broad as pupil; head large, not greatly compressed, not keeled above, its sides scaly; lower jaw with some scales; interorbital space depressed,  $5\frac{1}{2}$  in head; eye very large (in young),  $2\frac{3}{4}$  in head; snout short, depressed, not pointed, and with lateral keel,  $5\frac{1}{2}$  in head; preorbital very narrow; mouth rather large, oblique, the maxillary reaching to below front of pupil,  $2\frac{3}{4}$  in head; lower jaw slightly longer, its tip with a stiffish pointed projection representing the barbel; teeth small, in bands, a few

\* *Lepidion* is sufficiently distinct from *Lepidia*, Savigny, 1817.

on vomer. No spines on snout or opercles. Gill membranes somewhat united, free from isthmus. Gill rakers slender, rather long, 10 to 12 on lower part of arch. Scales very small, mostly lost posteriorly (in our specimen) and not to be exactly counted; lateral line not evident. First dorsal rather low and long, none of its rays produced, the longest about  $\frac{1}{2}$  head; ventrals filamentous,  $\frac{1}{2}$  head; pectorals about  $\frac{1}{2}$  head; caudal  $2\frac{1}{2}$  head; anal deeply notched behind the middle, its posterior lobe highest. Color uniform purplish black, the fins paler. One young individual,  $2\frac{1}{2}$  inches long, from *Albatross Station* 2995, off the Revillagigedo Islands. (*verecundus*, modest.)

*Lepidion verecundum*, JORDAN & CRAMER, Proc. U. S. Nat. Mus. 1896, 456, Revillagigedo Islands, at Albatross Station 2993. (Coll. *Albatross*.)

#### 984. *ANTIMORA*, Günther.

*Antimora*, GÜNTHER, Ann. Mag. Nat. Hist. 1876, 2 (*rostrata*).

This group differs from *Lepidion* in the form of the snout, the backward position of the vent, the imperfect division of the anal, in which latter respect it approaches *Mora*. In *Lepidion* the snout is subconical, obtusely rounded; in *Antimora* it forms a flat, triangular lamina, sharply keeled at the sides, resembling the snout of *Macrourus*. Body elongate, compressed, tapering into a slender tail. Scales very small. Head entirely scaly, even to the gill membranes. Snout depressed, thin and flat, projecting beyond the mouth; mouth rather large; chin with a barbel; jaws with bands of villiform teeth; a small roundish patch of teeth on vomer, none on palatines. Dorsal fins 2, the first short, its anterior ray produced into a long filament; anal fin deeply notched, almost separated into 2 fins; ventral fins with 6 rays, 1 of them filamentous; caudal truncate. Branchiostegals 7. Deep-water fishes. (*avri*, opposite; *Mora*, a related genus.)

a. Head rather small,  $4\frac{1}{2}$  in length; scales 115.

VIOLA, 2914.

aa. Head rather large, about  $5\frac{1}{2}$  in length; scales 130.

MICROLEPIS, 2915.

#### 2914. *ANTIMORA VIOLA* (Goode & Bean).

Head  $4\frac{1}{2}$  in body; depth 5. D. 4-53; A. 40; V. 6; scales 11-115-27. Snout broad, pointed at tip, much depressed, forming a roof-like projection above mouth; a conspicuous keel extending backward from tip of snout along the suborbital to the posterior margin of the eye. Mouth U-shaped, wholly inferior; maxillary nearly reaching posterior margin of orbit; interorbital space flat, as wide as the large eye, the orbital ridges somewhat elevated; barbel about  $\frac{1}{2}$  diameter of orbit. First dorsal with its first ray much produced, longer than head; anal fin deeply notched near its middle. Caudal peduncle as long as eye, its depth more than  $\frac{1}{2}$  its length; longest ray of ventrals reaching about halfway to vent; pectoral  $1\frac{1}{2}$  in head. Color deep violet or blue black; inside of mouth and opercles blue black. Banks of Newfoundland and southward, in deep water. (Goode & Bean.) (*viola*, violet.)

*Haloporphyrus viola*, GOODE & BEAN, Proc. U. S. Nat. Mus., 1, 1878, 256, La Have Bank, 400 to 500 fathoms; JORDAN & GILBERT, Synopsis, 800, 1883.

*Antimora viola*, GOODE & BEAN, Oceanic Ichthyology, 372, fig. 324, 1896.

## 2915. ANTIMORA MICROLEPIS, Bean.

Head about 4 in total length with caudal; depth  $5\frac{3}{4}$  without caudal; eye 4 in head, nearly equal to snout. D. 4 or 5-51; A. 41; barbel very slender, 2 in eye; gill rakers short, slender, 4 + 11. Maxillary reaching to nearly below posterior edge of eye; longest ray of first dorsal about  $\frac{1}{2}$  as long as head; anal deeply emarginate, beginning under twentieth ray of second dorsal; second ventral ray  $1\frac{1}{6}$  in head. Scales very small, about 9 rows between origin of second dorsal and lateral line, and about 130 in lateral line. Color olivaceous, deeper on opercles and branchiostegal membranes and on inside of mouth. Off Queen Charlotte Islands. Several specimens taken by the *Albatross* at different stations in Bering Sea, at depths of 350 and 351 fathoms, and off the coasts of the Queen Charlotte Islands and California, at depths of 1,588 and 455 fathoms. One large specimen, from off Bogoslof Island, has the filamentous ray of first dorsal  $\frac{3}{4}$  length of head, and the eye is shorter than the snout. (*μικρός*, small; *λεπίς*, scale.)

*Antimora microlepis*, BEAN, Proc. U. S. Nat. Mus. 1890, 38, off Cape St. James, Queen Charlotte Island, at Albatross Station 2860, in 876 fathoms (Type, No. 45361); GILBERT, Rept. U. S. Fish Comm. 1893 (1896), 456 and 473; JORDAN & GILBERT, Rept. Fur Seal Invest., 1898.

## 985. URALEPTUS, Costa.

*Uraleptus*, COSTA, Archiv fur Naturgesch. 1858, 87 (*maraldi*).  
*Gadella*, LOWE, Proc. Zool. Soc. Lon<sup>n</sup>. 1843, 91 (*gracilis*).

Body elongate, compressed, and tapering posteriorly, covered with small scales. A separate caudal; 2 dorsal fins and 1 anal; ventral fins narrow, with flat base, composed of 6 rays. Upper and lower jaw with an outer series of strong curved teeth; vomerine and palatine teeth none; chin without barbel. Branchiostegals 7. Deep waters of the Atlantic. (*ὄψα*, tail; *λεπρός*, slender.)

## 2916. URALEPTUS MARALDI (Risso).

Head 4; depth 6 Head rather thick, its greatest width equal to its height, which is somewhat more than  $\frac{1}{2}$  its length; cleft of mouth oblique, wide, the maxillary extending to below posterior margin of orbit; lower jaw received within the upper, but both nearly equal in length anteriorly, each armed with a series of rather large, curved, widely set teeth; another series of small teeth within the outer in the upper jaw. Snout rather broad, obtusely rounder, scarcely longer than eye, which is  $4\frac{1}{2}$  in head; interorbital space emarginate on each side of upper part of orbit, its width somewhat more than diameter of eye. Nape broad, scarcely elevated, with a spine on each side pointing outwards and covered by skin. Operculum small, with a slender horizontal spine posteriorly, the part below the spine being emarginate; gill membranes united below the throat by a rather narrow cutaneous bridge, not attached to the isthmus; gill openings wide; gills 4, a slit behind the fourth; pseudobranchiae glandular. Trunk rather low; tail tapering into a very narrow band;

first dorsal fin commencing behind vertical from base of pectoral, somewhat higher than long, and not higher than second; second dorsal commencing immediately behind the first, its rays increasing somewhat in length posteriorly, one of the longest being  $\frac{1}{2}$  as long as head, the whole fin naked; caudal fin slender, slightly rounded, entirely free from dorsal and anal, and nearly  $\frac{1}{2}$  as long as head; anal fin commencing at some distance behind the vent, which is situated below the origin of the first dorsal, very similar to the second dorsal; pectoral inserted somewhat below middle of body, its length equaling distance between front margin of eye and end of operculum; ventrals narrow, slender, the outer ray produced into a filament shorter than the pectoral. Scales extending over the whole head, the chin and the thin lips being naked. (Günther.) Tropical Atlantic. This form, originally described from Nice, has since been found at Madeira by Johnson, and at Naples and Catania by Giglioli. The *Blake* obtained a poor specimen, apparently of this form, at station LXXXI, off the Island of Nevis, in the West Indies. (Goode & Bean.) (Eu.) (A personal name for one of "quelques hommes que les talens, le mérite, la gloire ou l'amitié m'ont désignés.")

*Gadus maraldi*, RISSO, Ichth. Nice, 123, pl. 6, fig. 13, 1810, Nice.

*Merluccius attenuatus*, COCCO.

*Gadella gracilis*, LOWE, Proc. Zool. Soc. Lond. 1843, 91, Madeira. (Type in University of Cambridge.)

*Merluccius maraldi*, RISSO, Eur. Mérid., III, 220, 1826.

*Uraletus maraldi*, GÜNTHER, Cat., IV, 349, 1862; GÜNTHER, Challenger Report, XXII, 87, 1887; GOODE & BEAN, Oceanic Ichthyology, 367, fig. 320, 1896.

#### 986. *LOTELLA*, Kaup.

*Lotella*, KAUP, Archiv für Naturgesch. 1858, 28 (*schlegeli*).

This genus differs from *Physiculus* chiefly in the presence in both jaws of an outer row of large teeth. Deep sea. (Name, a diminutive of *Lotø*.)

#### 2917. *LOTELLA MAXILLARIS*, Bean.

Head about  $4\frac{1}{2}$ ; depth 5. D. 5-55; A. 44; V. 10; scales about 7 or 8-115-14 or 15. Snout short; eye 3 in head; maxillary reaching vertical through anterior margin of pupil, its length equaling that of postorbital part of head. Teeth in narrow bands in jaws, the outer series enlarged; vomer and palute apparently without teeth. Vent situated about under eighth ray of second dorsal; distance of first dorsal from tip of snout 4 times in total length including caudal; ventrals extending to about vertical from origin of second dorsal, not reaching nearly to vent; longest ray of first dorsal a little more than  $\frac{1}{2}$  as long as head; none of the rays of second dorsal or of anal as long as first ray of first dorsal; longest ray of second dorsal not much exceeding  $\frac{1}{2}$  of height of body; longest ray of anal about  $\frac{1}{2}$  length of ventral; origin of anal about under tenth ray of second dorsal; ventrals situated about under beginning of posterior third of head, their length  $\frac{1}{2}$  that of second dorsal base; origin of pectoral somewhat in advance of that of first dorsal, the fin imperfect, but its length probably slightly exceeding that of ventral; caudal

rounded. Color very light brown; the margins of the dorsal and anal, in their posterior portions, blackish. (Goode & Bean.) Gulf stream, Lat. 40° N., in 396 fathoms. (*maxillaris*, pertaining to the upper jaw.)

*Lotella maxillaris*, BEAN, Proc. U. S. Nat. Mus. 1884, 241, Lat. 39° 55' N., Long. 70° 28' W. in 396 fathoms (Type, No. 29832. Coll. Fish Hawk); GOODE & BEAN, Oceanic Ichthyology, 368, 1896.

987. **PHYSICULUS**, Kaup.

*Physiculus*, KAUP, Archiv für Naturgesch. 1858, 88 (*dabwigkii*).

Body elongate, covered with small scales; head entirely scaly; snout broad, obtusely rounded, projecting beyond the mouth; mouth of moderate size; chin with a barbel; jaws with bands of villiform teeth; vomer and palatines toothless. Dorsals 2; anal fin single, not notched; ventral fin with 5 rays, the outer ray filamentous; caudal rounded, slender, free; branchiostegals 7. (*φυκίς*, an ancient name of some fish living in the *Fucus*, *φύκος*, probably a species of *Gobius*.)

a. Scales moderate, about 62 in a longitudinal series; gill rakers few; dorsal rays 10-49; anal 54; ventral reaching fourth anal ray. FULVUS, 2918.

aa. Scales very small, about 100 in a longitudinal series.

b. Gill rakers few, about 11 below arch; head 4 in length; depth 5; ventrals filamentous at tip, as long as head, reaching tenth ray of anal.

NEMATOPUS, 2919.

bb. Gill rakers undescribed, probably few; head 4 in length; depth 6; ventrals shorter than head, reaching front of anal.

KAUPI, 2920.

bbb. Gill rakers very numerous, 7+18; head 3½ in length; depth 4½; ventrals reaching seventh anal ray.

RASTRELLIGER, 2921.

2918. **PHYSICULUS FULVUS**, Bean.

Head about 4; depth 4½. D. 10-49; A. 54; V. 7; scales 6-61 to 62-16. Head broad and depressed; snout short; eye 3½ in head; the length of the upper jaw 2½ in head, about equal to space between ventrals and anal origin; maxillary not quite reaching vertical through hind margin of eye; barbel 6 in head. Teeth in narrow bands in jaw; no outer series of enlarged teeth, but a few in the middle of the bands in each jaw are slightly larger than the others; all of the teeth, however, inconspicuous; vomer and palate smooth. Vent situated about under third ray of first dorsal; distance of first dorsal from tip of snout equaling 3 times length of its base, its longest ray twice length of snout, and slightly exceeding length of longest of second dorsal; length of second dorsal base 3 times length of pectoral, which is nearly 5½ times in total without caudal. Origin of anal about in a vertical let fall from base of fifth ray of first dorsal; distance of ventral from tip of snout about 5½ times in body. Tip of ventral when extended backward reaching base of fourth anal ray; length of middle caudal ray 3 in head. Lateral line very indistinct, situated rather high, following pretty closely the contour of back. Gill rakers moderately short and not numerous. General color a light yellowish brown; under surface of head, the abdomen, margins of dorsal and anal fins, lips, and axil of pectoral very dark brown; a dark brown blotch on the suboperculum; inside of mouth and gill membranes white. (Bean.)

Caribbean Sea, north to 40° in region of Gulf Stream, reaching a depth of 955 fathoms. (*fulvus*, brownish yellow.)

*Physiculus fulvus*, BEAN, Proc. U. S. Nat. Mus. 1884, 240, Lat. 40° 1' N., Long. 69° 56' W. in 79 fathoms (Type, No. 28766. Coll. *Fish Hawk*); GOODE & BEAN, Oceanic Ichthyology, 366, 1896.

2919. *PHYSICULUS NEMATOPUS*, Gilbert.

Head 4 in length; depth 5. D. 7 to 9—56 to 61; A. 59 to 64; scales 90 to 105. Length of caudal peduncle to base of median caudal rays  $5\frac{1}{2}$  in head. Snout very broadly rounded, its width twice its length, which is  $4\frac{3}{8}$  in head; eye  $3\frac{3}{8}$ ; interorbital 4; maxillary 2, reaching slightly beyond vertical from posterior margin of orbit. None of the teeth enlarged; palate smooth. Branchiostegal membranes more narrowly joined than in the *P. rastrelliger*, but wholly free from isthmus. Gill rakers short and slender, 11 movable ones on horizontal limb of arch. Origin of first dorsal over base of pectorals, its distance from tip of snout  $3\frac{3}{8}$  in length; base of first dorsal equaling snout and  $\frac{1}{2}$  eye, its highest ray  $2\frac{1}{2}$  in head; free portion of caudal peduncle  $\frac{1}{2}$  diameter of orbit; notch of dorsal and anal fins not conspicuous, the posterior dorsal rays little longer than those which precede,  $2\frac{1}{2}$  in head; caudal  $2\frac{3}{8}$ ; pectorals  $1\frac{1}{2}$ ; ventrals with broad base and 7 rays, the outer 2 filamentous, the second the longest, reaching base of tenth to twelfth anal rays, and as long as head; distance between bases of ventrals equals interorbital width; scales small, regularly imbricated, becoming minute on snout, which they completely invest, as well as mandible and gular membranes; lateral line present on anterior half of body only, 8 scales above it anteriorly. Color light olive brown, sprinkled with dark specks, the sides of head and trunk with silvery luster; snout, mandible, and gular membrane dusky; abdominal area, branchiostegal membranes, base of ventrals, axillary blotch, and front of anal, purplish black; posterior edge of gill membranes and opercular flap white; dorsals dusky, with an inconspicuous darker margin, which becomes more marked posteriorly; anal darker, margined with black; caudal blackish; pectorals and filamentous portion of ventral white. Inside of mouth and gill cavity white; peritoneum silvery, rendered black on sides by clusters of spots. Coast of southern California. Many specimens, the largest 7 inches long, from *Albatross Stations* 2997, 3011, 3015, and 3016, in 71 to 221 fathoms. (Gilbert.) (*νήμα*, thread; *πούς*, foot.)

*Physiculus nematopus*, GILBERT, Proc. U. S. Nat. Mus. 1890, 114, coast of southern California. (Types, No. 46488 and 46555. Coll. Dr. Gilbert.)

2920. *PHYSICULUS KAUPI*, Poey.

Head  $3\frac{1}{2}$ ; depth 4. D. 10—60; A. 60; P. 30; V. 8; C. 17; scales 12 to 15—over 100. Body and head short, swollen; tail regularly narrowed; vent below base of pectoral; eye high, equal to snout,  $4\frac{1}{2}$  in head; nostrils with valves; snout blunt; lower jaw the shorter; maxillary reaching slightly beyond eye; each jaw with a band of cardiform teeth, none on palate and tongue; maxillary sloping under skin of cheek; barbel a little longer than eye; opercular bones covered with skin, without spines; no spines at

nape; gill membranes somewhat united, free from isthmus; gills 4, a slit behind fourth; no pseudobranchiae; lateral line parallel with the back to beyond middle of body, then turning down suddenly, continuing to base of caudal. Scales small, cycloid, not easily counted; head scaly, even to the lips; vertical fins with small scales; ventral filamentous, equaling  $\frac{2}{3}$  length of head, all the rays except the first short; ventral with 2 filamentous rays, which reach to front of anal and are about  $1\frac{1}{2}$  in head; first dorsal as high as long, beginning behind base of pectoral, its longest rays about  $2\frac{1}{2}$  in head; soft dorsal and anal low, free from the small rounded caudal; pectoral falcate,  $1\frac{1}{2}$  in head. Color yellowish brown, bluish on belly; second dorsal and anal edged with darker brown. Type, 1 specimen, 250 mm. long. (Poey.) Deep waters of the Atlantic.

This species has constantly (as far as is shown in our specimens) a broader base to the ventral fins than *Physiculus dalwigkii*, and they are formed of 7 rays, of which the largest may or may not reach the anal fin; the fin rays vary within proportionate limits; they are, D. 9 or 10—60 to 66; A. 60 to 70; there are 13 scales between the anterior dorsal and lateral line; the caudal peduncle is shorter and less slender than in the Maderian form, but otherwise the species are so similar as to scarcely deserve specific separation. Poey obtained a specimen at Cuba, and Melliss 2 at St. Helena. These differ in no respect from 5 examples, 11 to 16 inches in length, found by the *Challenger* off Inosima in 345 fathoms. (Günther, *Challenger Report*, xxii, 88, pl. xvii, fig. A, 1887.) (Named for Dr. J. J. Kaup, author of a work on the Apodal fishes.)

*Physiculus kaupi*, POEY, Repertorio, 1, 186, 1865, Matanzas. (Coll. Don Cirilo Dulzaides.)  
? *Physiculus japonicus*, HILGENDORF, Sitz. Naturf. Freunde, Berlin, 1879, 80, Japan.

2921. *PHYSICULUS RASTRELLIGER*, Gilbert.

Head  $3\frac{1}{2}$  in length; depth  $4\frac{1}{2}$ . D. 8 or 9—53 to 61; A. 57; scales 100 to 110. Length of caudal peduncle to base of median caudal rays,  $2\frac{1}{2}$  in head. Snout short and broadly rounded,  $4\frac{1}{2}$  in head; eye  $3\frac{3}{8}$ ; interorbital width  $4\frac{1}{4}$  to  $4\frac{3}{8}$ ; maxillary  $2\frac{1}{2}$ , extending to vertical from posterior margin of pupil. Teeth in rather broad bands, none of them enlarged; width of patch on premaxillaries  $\frac{1}{2}$  pupil; vomer and palatines toothless. Branchiostegal membranes broadly united, joined to the isthmus anteriorly, the width of the free fold more than  $\frac{1}{2}$  pupil. Gill rakers numerous, slender, moderately long, the longest  $\frac{1}{4}$  diameter of orbit, about 7 above angle, 17 to 19 below, the anterior ones short but movable. Origin of first dorsal slightly in advance of base of pectorals, its distance from tip of snout  $3\frac{3}{8}$  in length; base of first dorsal equaling length of snout, its longest ray  $2\frac{3}{8}$  in head; free portion of caudal peduncle equaling diameter of eye; second dorsal notched, the median rays  $\frac{1}{4}$  the height of the highest anterior rays, the posterior highest, equaling first dorsal and longest caudal rays; anal similar to soft dorsal, but lower; ventrals under middle of opercle, the distance between their bases little less than interorbital width, equaling distance from vent to anal fin; ventrals with 7 rays, the outer 2 produced, the second the longest, reaching base of seventh or eighth anal ray; pectorals with broad base, covered with lax membrane,



containing 26 to 28 rays, their length  $1\frac{1}{2}$  to  $1\frac{3}{8}$  in head. Scales small, comparatively little reduced on top of head, a broad ring encircling snout in front of eyes naked, a very narrow patch of scales between this laterally and premaxillaries; scales in 100 to 110 transverse rows, 8 or 9 between lateral line and front of dorsal; lateral line wanting on posterior part of body, in the latter part of its course present on occasional scales only. Color uniform grayish olive on sides, each scale, or at least its marginal  $\frac{1}{2}$ , closely covered with minute dark specks; gular and branchiostegal membranes, ventral region, and axil of pectorals blue black; basal portion of vertical fins light bluish, margined with blackish; pectorals dusky; ventrals blue black at base, the distal portion white; lining membrane of mouth and gill cavity white; peritoneum silvery, but in places so filled with black specks as to appear black. Coast of southern California. Many specimens, the longest 8 inches, from *Albatross Stations* 3045 and 2987, in 184 and 171 fathoms. (Gilbert.) (*rastrelliger*, bearing small gill rakers; *rastrum*, rake; *gero*, I bear.)

*Physiulus rastrelliger*, GILBERT, Proc. U. S. Nat. Mus. 1890, 113, coast of southern California. (Type No. 48266. Coll. *Albatross*.)

### 988. LOTA (Cuvier) Oken.

(BURBOTS.)

*Les Lottes*, CUVIER, Règne Anim., Ed. 1, vol. 2, 215, 1817 (*lota*).

*Lota*, OKEN, Isis 1817, 1182 (*lota*).

Body long and low, compressed behind. Head small, depressed, rather broad; anterior nostrils each with a small barbel; chin with a long barbel; snout and lower parts of head naked; mouth moderate, the lower jaw included; each jaw with broad bands of equal, villiform teeth; vomer with a broad, crescent-shaped band of similar teeth; no teeth on palatines. Gill openings wide, the membrane somewhat connected, free from the isthmus. Scales very small, embedded; vertical fins scaly. Dorsal fins 2, the first short, the second long, similar to the anal; caudal rounded, its outer rays procurrent; ventrals of several rays. One or 2 species, living in fresh waters of northern regions. (*Lota*, the ancient name used by Rondelet, in French, *la Lotte*.)

### 2922. LOTA MACULOSA (Le Sueur).

(BURBOT; LAKE LAWYER; LING.)

Head  $4\frac{1}{2}$  in body; depth  $5\frac{1}{2}$ . D. 13-76; A. 68; ventral 7; eye 7 in head; pectoral  $1\frac{1}{2}$  in head; maxillary  $2\frac{3}{8}$ ; middle caudal rays  $2\frac{3}{8}$ . Body elongate, not much compressed anteriorly; head slightly depressed; mouth large, the maxillary reaching to posterior margin of eye; teeth villiform, in bands on jaws and vomer; barbel longer than the small eye; interorbital broad, nearly twice diameter of eye; gill rakers very short, about 3 + 6 in number; anterior nostrils with barbels; body covered with small embedded scales; pectorals scarcely reaching to below front of dorsal; ending of ventrals filamentous; caudal rounded; vertebrae 21 + 38 = 59; caeca 30. Dark olive, thickly marbled and reticulated with blackish;

yellowish or dusky beneath; young often sharply marked, the adult becoming dull grayish; vertical fins with dusky margins. Length 2 feet. Lakes and sluggish streams. New England and Great Lake region, north to the Arctic seas and west to the headwaters of the Missouri, the Frazer River basin, and Bering Straits; abundant northward; rare in the Ohio River and the Upper Mississippi; a rather coarse and tasteless fish, seldom used as food. Here described from a specimen, 18 inches long, from Lake Michigan at Michigan City, Indiana. The American Burbot is very close to the common species of northern Europe and Asia, *Lota lota* (Linnaeus) = *Lota vulgaris*, Cuvier = *Lota communis*, Rapp, and may prove wholly identical with the latter. In *Lota lota* the pectorals reach beyond front of dorsal, being  $1\frac{1}{2}$  in head. (*maculosus*, spotted.)

*Gadus maculosus*, LE SUEUR, Jour. Ac. Nat. Sci. Phila., 1, 1817, 33, Lake Eric.

*Molva maculosa*, LE SUEUR, Mém. Mus., v, 1819, pl. 16.

*Lota maculosa*, DE KAY, New York Fauna: Fishes, 284, pl. 52, fig. 168, 1842.

*Gadus compressus*, LE SUEUR, Jour. Ac. Nat. Sci. Phila., 1, 1817, 84, Connecticut River.

*Lota compressa*, DE KAY, New York, Fauna: Fishes, 285, pl. 78, figs. 244, 245, 1842.

*Gadus lacustris*, MITCHELL, Amer. Monthly Mag., II, 1818, 244, Sebago Pond, Maine (Coli. Henry A. S. Dearborn).

*Molva huntia*, LE SUEUR, Mém. Mus., v, 1819, 161, Connecticut River.

*Lota inornata*, DE KAY, New York Fauna: Fishes, 283, pl. 45, fig. 145, 1842, Hudson River, Lansingburgh, N. Y.

*Lota brosmiana*, STORER, Boston Journ. Nat. Hist., IV, 1839, pl. 5, fig. 1, New Hampshire.

#### 989. MOLVA, Fleming.

(LINGS.)

*Molva*, FLEMING, British Animals, 192, 1828 (*vulgaris*).

*Molva*, NILSSON, Skandinav. Fauna, IV, 573, 1832 (*molva*).

Body elongate, covered with very small scales. Chin with a barbel; lower jaw included; bands of teeth on jaws and vomer; lower jaw with large canines which are arrow-shaped and movable; vomer with a curved series of canines mixed with small teeth, these mostly fixed; no teeth on palatines. Gill membranes broadly united. Two dorsal fins, both well developed; 1 anal fin; ventrals with several rays. Northern seas. (An old name of the salt-water ling.)

#### 2923. MOLVA MOLVA (Linnaeus).

Head 5; depth 7 or 8. D. 13 to 16-63 to 70; A. 57 to 66 (vertebrae 27 + 37 = 64). Upper jaw the longer, the maxillary reaching to below middle of orbit. Teeth cardiform in the jaws, with an inner row of rather widely separated and larger ones on mandible; a semicircular band on vomer, among which a few larger ones are interspersed. First dorsal inserted over the latter half of pectoral, its greatest height  $\frac{3}{4}$  that of body below it; pectoral about  $\frac{1}{2}$  as long as head; anal insertion in vertical over seventh or eighth ray of second dorsal. Barbel longer than eye, which is about equal to width of interorbital space. Scales small, covering head and fins. Color black gray, lighter on the sides and beneath; vertical fins edged with white; a dark blotch at the posterior end of the first dorsal, and a more distinct one on the end of the second dorsal. Arctic parts of the

Atlantic, south in deep water. This fish, the "ling" of Europe, is found from Spitzbergen to the Gulf of Gascony, where specimens have been taken very exceptionally at Areachon and San Juan de Luz. It is very rare, however, south of the British Channel, and most abundant along the coast of northern Europe, especially in the German Ocean and off Norway. It is rare about Iceland, Greenland, and the Faroe Islands, and has never been found in the Baltic. It is said to have been found in the deep water off Newfoundland, but we have been unable to find the specific record. Collett states that on the Norwegian coast young examples rarely occur in less depth than 100 fathoms, and according to Lilljeborg the largest are caught in from 80 to 150 fathoms. (Goode & Bean.) (*molva*, an ancient name.)

*Gadus molva*, LINNÆUS, Syst. Nat., Ed. x, 254, 1858, seas of Europe; after *Gadus dorso dipterygia*, ARTEDI.

*Molva vulgaris*, FLEMING, British Animals, 192, 1828; GÜNTHER, Cat., IV, 361, 1862; GOODE & BEAN, Oceanic Ichthyology, 364, fig. 317, 1896.

*Gadus raptor*, NILSSON, Prodrömus, 46, Sweden.

*Molva linnaei*, MALM, Götheborgs och Bohusläns Fauna, 491, 1877.

### 990. UROPHYCIS, Gill.

(CODLINGS.)

*Phycis*, BLOCH & SCHNEIDER, Syst. Ichth., 56, 1801 (*tinea* = *blennioides*); not *Phycis*, Fabricius, 1798, a genus of Lepidoptera.

*Phycis*, RAFINESQUE, Amer. Monthly Mag. 1818, 243 (*marginata*).

*Urophycis*, GILL, Proc. Ac. Nat. Sci. Phila. 1863, 240 (*regius*).

*Emphycus*, JORDAN & EVERMANN, new subgenus (*tenuis*).

Body rather elongate; head subconic; mouth rather large, the maxillary reaching to below eye; lower jaw included; chin with a small barbel; jaws and vomer with broad bands of subequal, pointed teeth; palatines toothless. Dorsal fins 2, the first sometimes produced at tip; second dorsal long, similar to the anal; ventrals wide apart, filamentous, each of 3 slender rays, closely jointed, appearing like one befid filament. Gill membranes somewhat connected, narrowly joined to the isthmus. (*οὐρά*, tail; *Phycis*.)

#### UROPHYCIS:

a. First dorsal fin not elevated, none of its rays filamentous.

b. Scales moderate, 90 to 95 in a longitudinal series.

c. Dorsal rays 8-43; anal 45; sides with some pale spots.

REGIUS, 2924.

cc. Dorsal rays 10-66; anal rays 57; barbel minute.

CIRRATUS, 2925.

bb. Scales small, 120 to 155 in a longitudinal series.

d. Dorsal rays 13-57; anal about 50; scales 120; sides with some pale spots.

FLOBIDANUS, 2926.

dd. Dorsal rays 10-62; anal about 53; scales 155.

EARLI, 2927.

EMPHYCUS (*ἐμφύκος*, in the seaweed):

aa. First dorsal fin elevated, 1 or more of its rays filamentous.

e. Scales about 140; dorsal rays 9-57; anal 48; ventrals reaching vent.

TENIUS, 2928.

ee. Scales about 110; dorsal rays 9-57; anal 50; ventrals reaching beyond vent, not longer than head.

CHUSS, 2929.

eee. Scales about 90; dorsal rays 9-56; anal 50; second dorsal filamentous; ventrals very long, nearly 3 times length of head.

CHESTERI, 2930.

Subgenus UROPHYCIS.

2924. UROPHYCIS REGIUS (Walbaum).

Head  $4\frac{1}{2}$  in body; depth 5. D. 8-13; A. 40; scales about 90; eye  $4\frac{1}{2}$  in head; maxillary 2; pectoral  $1\frac{1}{2}$ ; caudal  $1\frac{1}{2}$ . Body rather elongate, compressed; mouth large, the maxillary reaching slightly past posterior margin of eye; lower jaw included; cardiform teeth on jaws and vomer; interorbital flattish, about equal in width to the diameter of eye; gill rakers short, 3 + 12 in number; origin of dorsal over base of pectorals; pectorals slender, barely reaching to front of anal; ventrals filamentous, composed of 2 rays each with the inner ray the larger, inserted in front of base of pectoral in distance equal to  $1\frac{1}{2}$  diameter of eye, their ends reaching beyond front of anal; front of anal nearer snout than base of caudal, by nearly a head's length; caudal subtruncate. Pate brownish tinged with yellowish, the lateral line dark brown, interrupted by white spots; inside of mouth white; first dorsal largely black, this color surrounded by white; second dorsal olivaceous, with irregular round dark spots; caudal, anal, and pectorals dusky; ventrals and lower edge of pectorals white; 2 vertical series of round dark spots on the sides of the head. North Atlantic, south to Cape Fear; ranging from shallow water to a depth of 167 fathoms. Here described from a specimen, 8 inches in length, from Charleston, South Carolina. The species is said to exhibit electric powers in life. (*regius*, royal.)

*Blennius*, sp., SCHÖPF, Schrift. Naturf. Freunde, Berlin, VIII, 1780, 142, New York.

*Blennius regius*, WALBAUM, Artedi, Pisc. III, 186, 1792; after SCHÖPF.

*Enchelyopus regalis*, BLOCH & SCHNEIDER, Syst. Ichth., 53, 1801, after SCHÖPF.

*Gadus punctatus*, MITCHILL, Trans. Lit. and Phil. Soc. N. Y., I, 1815, 372, New York.

*Urophycis regius*, GILL, Proc. Ac. Nat. Sci. Phila. 1863, 240.

*Phycis regius*, GOODE & BEAN, Oceanic Ichthyology, 357, 1896.

*Phycis regalis*, GÜNTHER, Cat., IV, 355, 1862.

*Phycis punctatus*, DE KAY, N. Y. Fauna: Fishes, 292, 1842.

2925. UROPHYCIS CIBRATUS, Goode & Bean.

Head 4; depth 5. D. 10-66; A. 57; scales 6-93-20. Body moderately stout; eye large, about 4 in head; interorbital space 2 in eye. Maxillary not reaching posterior margin of orbit in large specimens, but in smaller ones it extends fully to that vertical; mandible extending far beyond posterior margin of eye, its length about equal to postorbital part of head; barbel minute in all examples examined, its length usually about  $\frac{1}{2}$  that of eye. Teeth in villiform bands in both jaws, the intermaxillary bands being wider than those of mandible; vomerines in a narrow villiform band. Gill rakers 2 + 12, the largest club-shaped at end, the longest 4 in eye. Gill membranes attached to isthmus, but with a narrow, free posterior border. Length of pectorals about  $\frac{1}{2}$  distance from ventral to anal origin, reaching to about the twenty-sixth row of scales; ventral reaching in some specimens slightly beyond origin of anal; in 1 individual almost to middle of anal fin; none of the dorsal rays filamentous, the longest from  $2\frac{1}{2}$  to 3 times in head; base of first dorsal about equal to length of eye in most specimens; in smaller examples somewhat greater, about 3 in head;

vent under sixteenth ray of second dorsal. Color light brown; lower parts minutely dotted; dorsals with narrow dark margins; caudal with a broad dark margin; anal with a narrow dark margin in its posterior third; roof of mouth and interior of gill cavity dark brown. Deep water of the Gulf of Mexico. (Goode & Benn.) (*cirratus*, bearing cirri.)

*Phycis cirratus*, GOODE & BEAN, Oceanic Ichthyology, 358, 1896, Gulf of Mexico at Lat. 29° 03' 15" N., Long. 88° 16' W. (Type, No. 39059. Coll. Albatross.)

2926. *UROPHYCIS FLORIDANUS*, Bean & Dressel.

Head 4 in body; depth 6. D. 13-57; A. 49; scales about 120; eye 6 in head; maxillary 2; height of first dorsal 2½; middle caudal rays 2. Body rather elongate, compressed, head subconic; mouth large, the maxillary reaching to below posterior margin of orbit; upper jaw and snout somewhat projecting beyond lower; small cardiform teeth, in narrow bands on jaws and vomer; barbel very slender, small; interorbital space wide, slightly convex, nearly twice as wide as eye; gill rakers small, slender, 2+11 in number. Origin of dorsal a little behind the vertical from base of pectoral; first dorsal high, slightly falcate; second dorsal a little higher in its anterior end, higher than anal; origin of anal about midway between tip of snout and base of caudal; pectoral slender, reaching an eye's diameter beyond front of second dorsal; ventrals inserted twice diameter of eye in front of pectorals, 2-rayed, the inner ray the longer, not reaching to vent in larger examples, reaching to front of anal in small ones; caudal long and rounded. Color in spirits, reddish brown, light below, a small black spot above eye, a vertical series of 3 or 4 behind eye, and 2 on opercle, these spots less than ¼ pupil, distinct and clear cut; a dark streak from preorbital across cheek to edge of opercle, lateral line black, interrupted at short intervals by white spots; fins dusky, with the exception of pectorals and ventrals, dark toward the ends of the rays. Gulf of Mexico, in rather shallow water, coming to shore in abundance about Pensacola in cold weather. Here described from a specimen, 7½ inches in length, from Pensacola, Florida. (*floridanus*, from Florida.)

*Phycis floridanus*, BEAN & DRESEL, Proc. Biol. Soc. Wash. 1884, 100, Pensacola, Florida (Coll. Silas Stearns); JORDAN, Cat. Fish. N. A., 129, 1885.

2927. *UROPHYCIS EARLLI*, Bean.

Head 3½ in body; depth 5. D. 10-60; A. 53; scales 155; eye 6 in head; maxillary 2. Body moderately elongate, not much compressed anteriorly; mouth large, the maxillary reaching to below or very slightly past posterior margin of eye; snout and upper jaw projecting beyond lower jaw; teeth strong, cardiform in a narrow band on vomer and lower jaw, in a rather wide band in upper; interorbital wide, convex, about 1½ times eye; gill rakers short and blunt, about 2+9. Origin of dorsal slightly behind the vertical from base of pectoral; origin of anal about midway between snout and base of caudal; ventrals 2-rayed, the inner a little the longer, not reaching to vent. Color brown, with some light spots on the second dorsal fin and on the sides; anal and both dorsals margined with brown.

Atlantic coast of United States, southward in water of moderate depth; not common. Here described from a specimen, 17 inches in length, from Charleston, South Carolina. (Named for R. Edward Earll, then assistant to the United States Fish Commission.)

*Phycis earlli*, BEAN, Proc. U. S. Nat. Mus., III, 1880, 69, Charleston, S. C. (Coll. R. E. Earll. Type, Nos. 25207, 25208, and 25209); JORDAN & GILBERT, Synopsis, 798, 1883.

Subgenus EMPHYCUS, Jordan & Evermann.

2928. UROPHYCIS TENUIS (Mitchill).

(CODLING; WHITE HAKE; SQUIRREL-HAKE.)

Head  $4\frac{1}{2}$ ; depth  $5\frac{1}{2}$ . D. 9-57; A. 48; scales 138. Snout longer than eye, narrower and more pointed than in *P. chuss*. Eye large, usually wider than interorbital space; maxillary reaching beyond pupil. Filamentous dorsal ray about  $\frac{2}{3}$  length of head; ventral fins about reaching vent. Scales very small. Brownish, lighter and yellowish below; fins very dark. Banks of Newfoundland to Cape Hatteras; abundant northward in rather deep water, reaching a depth of 304 fathoms. The species resembles *Phycis chuss*, differing chiefly in the smaller scales. (*tenuis*, slender.)

*Gadus tenuis*, MITCHILL, Trans. Lit. and Phil. Soc. N. Y., 1815, 372, New York.

*Phycis dekayi*, KAUP, Archiv Natur. 1858, 89, North America.

*Phycis rostratus*, GÜNTHER, Cat. Fish. Brit. Mus., IV, 353, 1862, no locality; D. 9-59 to 62; A. 49 to 50; scales ca. 150; ventrals immaculate, reaching front of anal.

*Phycis tenuis*, DE KAY, N. Y. Fauna: Fishes, 203, 1842; GILL, Proc. Ac. Nat. Sci. Phila., 1863, 238; JORDAN & GILBERT, Synopsis, 709, 1883; GOODE & BEAN, Oceanic Ichthyology, 350, fig. 312, 1896.

2929. UROPHYCIS CHUSS (Walbaum).

(CODLING; SQUIRREL-HAKE.)

Head  $4\frac{1}{2}$ ; depth 5. D. 9-57; A. 50; scales 110. Body rather slender; head depressed; eye large, about equal to interorbital width; maxillary reaching posterior margin of pupil; filamentous dorsal ray about  $\frac{2}{3}$  length of body, when perfect; pectorals  $\frac{2}{3}$  length of head; ventral fins extending beyond the vent; scales comparatively large. Brownish above, sides lighter and tinged with yellowish; thickly punctulate with darker; below pale; inside of mouth white; vertical fins somewhat dusky; anal fin margined with pale; lateral line not dark. Atlantic coast, from Gulf of St. Lawrence to Virginia; common northward; reaching a depth of 300 fathoms. (*chuss*, a vernacular name now obsolete, apparently derived from cusk.)

*Chuss*, SCHÖPF, Schrift. Naturf. Freunde, Berlin, VIII, 1780, 143, New York.

*Blennius chuss*, WALBAUM, Arted. Pisc., 186, 1792; after SCHÖPF.

*Enchelyopus americanus*, BLOCH & SCHNEIDER, Syst. Ichth., 53, 1801; after SCHÖPF.

*Gadus longipes*, MITCHILL, Trans. Lit. and Phil. Soc. N. Y., 1, 372, pl. 1, fig. 4, 1815, New York.

*Phycis marginatus*, RAFINESQUE, Amer. Month. Mag., Jan., 1818, 205, Point Judith, Rhode Island. D. 10-60; A. 40; ventral reaching anal; tail black-edged.

*Phycis americanus*, STÖBER, Rept. Fish. Mass., 138, 1839; GÜNTHER, Cat., IV, 353, 1862.

*Phycis chuss*, GILL, Proc. Ac. Nat. Sci. Phila., 1863, 237; JORDAN & GILBERT, Synopsis, 799, 1883; GOODE & BEAN, Oceanic Ichthyology, 359, fig. 311, 1896.

## 2930. UROPHYCIS CHESTERI, Goode &amp; Bean.

Head  $4\frac{1}{2}$ ; depth 5; orbit  $3\frac{1}{2}$  in head; maxillary 2; barbel about 3 in orbit. D. 9 or 10-55 to 57; A. 56; C. 5, 18 to 21, 5; P. 17 or 18; V. 3; scales 7-90 or 91-28. Vent situated under the twelfth ray of second dorsal, and equidistant from tip of snout and end of second dorsal; distance of dorsal fin from snout equal to twice length of mandible; third ray of first dorsal extremely elongate, extending to a point (thirty-third ray of second dorsal)  $\frac{2}{3}$  of distance from snout to tip of caudal, its length more than twice that of head, and more than 4 times as long as the rays immediately preceding and following it; anal fin inserted immediately behind vent, its distance from root of ventrals equal to that of dorsal from snout; as in other species of the genus, ventral of 3 rays, the first 2 much prolonged, the first contained 3 times in length of body, the second almost 3 times as long as head, reaching to fortieth anal ray or  $\frac{1}{4}$  of distance from snout to tip of caudal, the third shorter than diameter of orbit; pectoral 4 times as long as operculum. Scales large and thin, easily wrinkling with the folding of the thick, loose skin, particularly in the median line of sides of body. Lateral line much broken on posterior half of body. (Goode & Bean.) Atlantic coast of United States, in 100 to 500 fathoms, with *Macrourus bairdi*, the most abundant fish on the continental slope, swarming everywhere below the 100-fathom line. (Named for Capt. Hubbard C. Chester.)

*Phycis chesteri*, GOODE & BEAN, Proc. U. S. Nat. Mus. 1878, 256, off Cape Ann, in 140 fathoms (Coll. Captain Chester); JOYDAN & GILBERT, Synopsis, 800, 1883; GOODE & BEAN, Oceanic Ichthyology, 360, fig. 313, 1896.

## 991. LÆMONEMA, Günther.

*Læmonema*, GÜNTHER, Cat. Fish. Brit. Mus., IV, 356, 1862 (*yarrellii*).

Body of moderate length, covered with small scales; fins naked. A separate caudal; 2 dorsal fins and 1 anal, the anterior dorsal composed of 5 rays; ventrals reduced to a single long ray, bifid at its end. Bands of villiform teeth in the jaws; a small group of vomerine teeth; none on the palatine bones. Chin with a barbel. Branchiostegals 7. Deep sea. (*λαζμός*, throat; *νήμα*, thread.)

a. Scales 13-140-31; barbel  $\frac{1}{2}$  eye; dorsal and anal with narrow black edgings.

BARBATULUM, 2931.

aa. Scales 16-160-38; barbel  $\frac{2}{3}$  eye; a large, triangular, black blotch on tall and adjacent parts of vertical fins.

MELANURUM, 2932.

## 2931. LEMONEMA BARBATULUM, Goode &amp; Bean.

Head  $4\frac{1}{2}$ ; depth  $4\frac{1}{2}$ ; orbit 3 in head; upper jaw more than 2; barbel about 2 in eye. D. 5-63; A. 59; P. 19; V. 2; scales 13-140-31. Vent situated under sixth or seventh ray of second dorsal. Distance of first dorsal from snout 4 in body; base of first dorsal  $\frac{1}{2}$  as long as middle caudal rays, that of second slightly more than 3 times length of head; first dorsal composed of 5 rays, the first of which is elongate, 3 times as long as middle caudal rays, extending to base of twenty-fourth ray of second dorsal; anal fin inserted at a distance from tip of snout equal to twice length of head,

its distance from insertion of ventrals being equal to length of head; length of ventrals equal to that of pectorals, their tip not extending to vent. Scales small, very thin, deciduous, crowded anteriorly; lateral line not well defined on posterior part of body. Color similar to that of the various species of *Phycis*; the dorsal and anal fins with narrow black margins. The length of the first dorsal ray is very variable, being shorter in younger individuals. This species differs from *L. yarrellii* by its much smaller scales, and from *L. robustum* by the greater number of rays in the dorsal and anal fins, and its much shorter ventrals. (Goode & Bean.) Gulf Stream, reaching a depth of 312 fathoms. (*barbatulus*, having small barbels.)

*Lemonema barbatula*, GOODE & BEAN, Bull. Mus. Com. Zool., x, 204, 1883, Gulf Stream, Lat. 32° 43' N., Long. 77° 20' W., in 230 fathoms, and Lat. 28° 35' N., Long. 73° 13' W.; GOODE & BEAN, Oceanic Ichthyology, 362, figs. 315 and 315A, 1896.

2932. LEMONEMA MELANURUM, Goode & Bean.

Head about 4½; depth 4½; eye 3 in head; snout 4; interorbital width 6. D. 6-57; A. 55; P. 25; V. 2; Br. 7; scales 16-160-38. Maxillary extending to below middle of eye; intermaxillary nearly ½ length of head; mandible slightly more than twice length of snout. Teeth in intermaxillary and mandible in villiform bands; vomerine teeth in a small circular patch on middle of head of bone. Barbel about as long as snout. Distance of first dorsal from tip of snout about 4 in snout; length of first ray of dorsal equaling that of head without snout; last ray of dorsal scarcely more than ½ as long as first; ventral consisting of a single bifid ray, its distance from tip of snout equal to length of head, its length nearly equal to that of dorsal or the pectoral when extended, not reaching vent by a distance equal to length of snout; pectoral equaling that of longest dorsal ray, and also equaling head without snout; second dorsal higher anteriorly, and posteriorly much higher than in middle; longest anterior ray ½ length of ventral; longest posterior ray ½ length of head. Vent under eighth ray of second dorsal. Gill rakers 5 + 15, the longest ½ as long as snout. Color very light brown, the dorsals and anal with a narrow dark margin; a conspicuous, large, triangular, dark blotch on last rays of dorsal and anal, and a dark blotch occupying almost the whole of caudal, leaving a margin of whitish around it. (Goode & Bean.) Caribbean Sea, north to New York; reaching a depth of 1,467 fathoms. (*μελας*, black; *οὐρά*, tail.)

*Lemonema melanurum*, GOODE & BEAN, Oceanic Ichthyology, 363, fig. 316, 1896, Gulf Stream, Lat. 30° 44' N., Long. 79° 26' W., in 440 fathoms. (Type, No. 38270. Coll. Albatross.)

992. GAIDROPSARUS, Rafinesque.

(THREE-BEARDED ROCKLINGS.)

*Gaidropsarus*, RAFINESQUE, Indice d'Ittiol. Siciliana, 1810 (*mustellaris* = *mediterraneus*); description from a rough figure of RONDELET.

*Les Mustèles*, CUVIER, Règne Anim., Ed. 1, vol. 2, 215, 1817 (*triacirrhatus* = *mediterraneus*).  
*Mustela*, OKEN, Isis, 1817 (for *les Mustèles*; not *Mustela*, a genus of mammals).



*Onos*, RISSO, Hist. Eur. Mérid., III, 214, 1826 (*mustella* = *mediterraneus*).

*Mustela*, STARK, Elem. Nat. Hist., I, 425, 1828 (after *les Mustèles*).

*Motella*, CUVIER, Règne Anim. Ed. 2, vol. II, 334, 1829 (*vulgaris* = *tricirratus*).

*Onus*, GÜNTHER, corrected spelling.

Body rather elongate, covered with minute scales; head not compressed, the upper jaw the longer; snout with 2 conspicuous barbels, the chin with 1; teeth on jaws and vomer in bands, palatines toothless; dorsals 2, the anterior of a single long ray followed by a series of short fringe-like rays concealed in a groove; second dorsal and anal long, similar to each other; caudal rounded or lanceolate; ventral rays 5 to 7. Small fishes of the northern seas, descending to deep water. We here regard the 5-bearded Rocklings (*Ciliata*, Couch, 1832) = *Couchia*, Thompson, 1856 = *Molcella*, Kaup, 1858, as a distinct genus, distinguished by the 5 barbels at the tip of the snout. (*γαϊδροψάρα*, a modern Greek name used by Rondelet for a species of this group.)

The name *γαϊδροψάρον* is now applied in Athens to the Pollack-like fish, *Micromesistius poutassou* (Risso).

According to Prof. Horace A. Hoffman "the name *γαϊδουροψάρον* is modern, meaning doukey fish. *Γαϊδουρος* = *γάδαρος* = ass, donkey. The ancients called a certain fish *όνος*, ass. Dorio, in Athenæus, VII, 99, says some persons call the *όνος* (i. e., the fish *όνος*) *γάδος*. Epicharmus, in his Marriage of Hebe, says: 'Wide-gaping *χάνναι* and monstrous-bellied *όνοι*.' (See Aristotle 599b 33, 601a 1, 620b 29, frag. 307, 1530a.) According to Aristotle the *όνος* has a mouth opening wide (literally, breaking back), like the *γαλεοί*. It leads a solitary life, is the only fish which has its heart in its belly, has stones in its brain like millstones in form, and is the only fish which lies torpid in the warmest days under the reign of the dog star, Sirius, the other fishes going into this torpid state in the wintriest days. The *όνος*, *βάτος*, *ψήττα*, and *ρίνη* bury themselves in the sand, and after they make themselves invisible they wave the things in their mouths which fishermen call little rods or little wands (*ράβδία*). (Hoffman & Jordan, Fishes of Athens, Proc. Ac. Nat. Sci. Phila. 1887, 146.)

a. First ray of first dorsal long, as long as head; head small,  $5\frac{1}{2}$  in length; teeth rather feeble, uniform. D. 59; A. 45; P. 25. Color uniform brick red.

ENSIS, 2933.

aa. First ray of first dorsal short, about as long as snout.

b. Pectoral rays 22 to 24; upper jaw without cirri or rudimentary barbels along the premaxillary; maxillary reaching posterior border of eye; head  $5\frac{1}{2}$  in length. D. 56; A. 45. Color reddish. ARGENTATUS, 2934.

bb. Pectoral rays 16; upper jaw with short cirri or barbels along the premaxillary; maxillary reaching far beyond eye. D. 50; A. 42. Head 4 in length. Color brownish. SEPTENTRIONALIS, 2935.

#### 2933. GAIDROPSARUS ENSIS (Reinhardt).

Head  $5\frac{1}{2}$ ; depth  $4\frac{1}{2}$ . D. 59; A. 44 to 46; P. 22 to 27; V. 8. Body unusually deep, being greatest at the vent; head small; eye rather large, nearly as long as snout, equaling interorbital area, and in anterior half of head; posterior margin of orbit nearly equidistant between tip of snout and posterior margin of operculum. Mouth normal; supramaxillary end-

ing under posterior margin of pupil. Teeth in a narrow band in each jaw, some of those at least in outer row of upper jaw slightly enlarged and brownish colored; teeth of vomer forming a short curved band in 2 rows. Nasal barbel about equaling diameter of eye. Chin barbel small and not much exceeding  $\frac{1}{2}$  diameter of eye. Foremost ray of first dorsal springing from back above opercular margin; second dorsal fin low in front, but rising rapidly to seventh or eighth ray, behind which it is nearly uniform for a long distance, and the highest at posterior portion; anal fin much lower than second dorsal; caudal slightly emarginate, almost truncate behind, its median rays about  $\frac{2}{3}$  as long as head; pectorals nearly  $\frac{2}{3}$  as long as head, produced toward the upper angles, the third ray being longest; ventral fins with their bases mostly in advance of pectorals, the longest ray filamentous and nearly equaling pectoral. Lateral line obsolete. (Goode & Bean.) Atlantic coast of North America, from Greenland to Cape Hatteras; in deep waters, reaching a depth in the Gulf Stream of 1,081 fathoms. (*ensis*, sword.)

*Motella ensis*, REINHARDT, Dansk. Vidensk. Selsk. Afhandl., vii, 15, 1838, Greenland.  
*Onos rufus*, GILL, Proc. U. S. Nat. Mus. 1883, 259, Gulf Stream; GILL, Proc. Ac. Nat. Sci. Phila. 1884, 172; JORDAN, Cat. Fish. N. A., 128, 1885.  
*Onos ensis*, GILL, Proc. Ac. Nat. Sci. Phila. 1863, 241; GILL, Cat. Fish. E. Coast U. S., 18, 1873; JORDAN & GILBERT, Synopsis, 797, 1883; JORDAN, Cat. Fish. N. A., 128, 1885; GOODE & BEAN, Oceanic Ichthyology, 381, fig. 327, 1896.

2934. GAIDROPSARUS ARGENTATUS (Reinhardt).

Head 5; depth  $5\frac{1}{2}$ . D. 54 to 59; A. 45 or 46. Body elongate; head small; teeth in several rows, 1 row more enlarged than the others; maxillary reaching posterior border of eye; eye large,  $5\frac{1}{2}$  in head; interorbital space scarcely exceeding the eye. Lateral line with about 27 enlarged pores along its entire length. First ray of first dorsal short, little longer than snout; vent near middle of length. Reddish gray, changing to bluish on the head and abdomen; tips of dorsal, anal, and caudal red, also the barbels and first ray of first dorsal; cavity of mouth pale. Coasts of Greenland (Collett), south to Faroë and Bear Islands; not seen by us. There can be no doubt that *Motella argentata* is the young of the species later called *Motella reinhardtii*. (Eu.) (*argentatus*, silvered.)

*Motella argentata*, REINHARDT, Dansk. Vidensk. Selsk. Afh., vii, 128, 1838; Greenland; young.  
*Motella reinhardtii*, KÖYER MS., 1852; COLLETT, Forh. Vid. Selsk. Chr., No. 14, 83, 1878, Greenland.  
*Couchia argentata*, GÜNTHER, Cat., iv, 365, 1862.  
*Oliata argentata*, GILL, Proc. Ac. Nat. Sci. Phila. 1863, 241; GILL, Cat. Fish. E. Coast U. S., 18, 1873.  
*Onos reinhardtii*, GILL, Proc. Ac. Nat. Sci. Phila. 1863, 241; GILL, Cat. Fish. E. Coast U. S., 18, 1873; COLLETT, Norske Nord-Havs Exp., 131, 1880; JORDAN & GILBERT, Synopsis, 797, 1883; JORDAN, Cat. Fish. N. A., 128, 1885; GOODE & BEAN, Oceanic Ichthyology, 383, 1896.

2935. GAIDROPSARUS SEPTENTRIONALIS (Collett).

Head 4; depth  $5\frac{1}{2}$ . D. 50; A. 42; P. 16. Three barbels, 2 at the nostrils, 1 at the chin, besides a row of about 8 shorter rudimentary barbels along the edge of the upper lip; eye small,  $\frac{1}{2}$  length of snout; cleft of mouth

extending far beyond eye, its length nearly equal to that of postorbital part of head; teeth rather small, unequal; outer teeth of upper jaw and some of the inner teeth of lower enlarged; first ray of first dorsal short, about as long as snout; vent midway between tip of snout and last anal ray; lateral line with about 20 large pores. Grayish brown, paler below; cavity of mouth white. Coast of Norway; 1 specimen known from Greenland. (Collett.) (Eu.) (*septentrionalis*, northern.)

*Motella septentrionalis*, COLLETT, Ann. Mag. Nat. Hist. 1874, 15, 82, Lofoten, Norway.  
*Onos septentrionalis*, COLLETT, Norske Nord-Havs Exped., 139, 1880; JORDAN, Cat. Fish. N. A., 128, 1885.

### 993. ENCHELYOPUS, Bloch & Schneider.

(FOUR-BEARDED ROCKLINGS)

*Enchelyopus*, BLOCH & SCHNEIDER, Syst. Ichth., 50, 1801 (*cimbrius*; the first species mentioned and the one left as type after elimination of the genera, defined prior to *Rhinonemus*).

*Rhinonemus*, GILL, Proc. Ac. Nat. Sci. Phila. 1883, 241 (*cimbrius*).

Barbels 4, 1 at each nostril, 1 at tip of snout, and 1 at the chin; head high and compressed anteriorly; teeth in narrow bands, some of them enlarged; otherwise essentially as in *Gaidropsarus*. North Atlantic. (*ἔγχελυπός*, resembling an eel; "*facie anguillaris*.")

### 2936. ENCHELYOPUS CIMBRIUS (Linnaeus).

(FOUR-BEARDED ROCKLING.)

Head  $5\frac{1}{2}$ ; depth 9. D. 45 to 50; A. 41 or 42; V. 5. Body slender, tapering from the shoulders back; caudal peduncle narrow, 4 in head; snout moderate, blunt, rounded, not depressed, a little shorter than the eye; eye large, subcircular, 4 in head; interorbital space narrow, equal to vertical diameter of eye, 6 in head; teeth villiform, those in the upper jaw unequal, small, with about 8 enlarged in front, those of the lower jaw long and slender, of equal length, a few somewhat enlarged in front; maxillary reaching beyond posterior border of eye, a barbel at each nostril, 1 on tip of snout and 1 on chin, stitch-like; lateral line with about 35 enlarged pores along its entire length; first (free) ray of dorsal nearly as long as head; ventral  $\frac{1}{2}$  head; caudal acute. Light clivaceous (salmon-red); first dorsal ray and posterior end of dorsal and anal abruptly black, as is lower half of caudal; pectorals and ventrals pale; sides of head somewhat silvery; cavity of mouth dark bluish. North Atlantic, on both coasts, south in deep water to the Gulf Stream; common in Massachusetts Bay; our specimens from Woods Hole; the young ("mackerel midges") silvery, unlike the adult in appearance. (Eu.) (*cimbrius*, Welsh.)

*Gadus cimbrius*, LINNÆUS, Syst. Nat., Ed. 12, 1, 440, 1766, Atlantic Ocean; Scania (Coll. Dr. Strussenfelt).

*Motella caudacuta*, STORER, Proc. Bost. Soc. Nat. Hist., III, 1848, 5, Cape Cod, Provincetown, Mass. (Coll. Herman M. Smith); STORER, Amer. Ac. Sci., 411, 1867; STORER, Hist. Fish. Mass., 183, 1867.

*Rhinonemus caudacuta*, GILL, Proc. Ac. Nat. Sci. Phila. 1863, 241; GOODE & BEAN, Amer. Journ. Sci. and Arts 1877, 476; JORDAN, Cat. Fish. N. A., 128, 1885.

*Motella cimbrius*, NILSSON, Prod. Ich. Scand., 48, 1832; BELL, Can. Nat. and Geol., IV, 209, 1850.

*Onos cimbricus*, GOODE & BEAN, Proc. U. S. Nat. Mus. 1878, 349; GOODE & BEAN, Bull. Essex Inst., XI, 1879; JORDAN & GILBERT, Synopsis, 797, 1883; GOODE & BEAN, Bull. Mus. Comp. Zool., X, No. 5, 217, 1883.

*Rhinonemus cimbricus*, JORDAN, Cat. Fish N. A., 128, 1885; GOODE & BEAN, Oceanic Ichthyology, 384, fig. 328, 1896.

*Enchelyopus cimbricus* (misprint for *cimbricus*), BLOCH & SCHNEIDER, Syst. Ich., 50, pl. 9, 1801.

994. BROSME (Cuvier) Oke.

(CUSKS.)

*Les Brosmes*, CUVIER, Règne Animal, Ed. 1, vol. 2, 216, 1817 (*brosme*).

*Brosme*, OKEN, Isis, 1817, 1182; after CUVIER.

*Brosmius*, CUVIER, Règne Animal Ed. 2, vol. II, 334, 1829 (*brosme*).

Body moderately elongate, covered with very small scales. Mouth rather large, with teeth in the jaws, vomer, and palatines, some of those on the vomer and palatines enlarged; chin with a barbel; branchiostegals 7. Dorsal fin single, continuous, not elevated, not notched; anal fin similar, but shorter; caudal fin rounded; ventral fin several-rayed. Northern seas. (From the Danish vernacular name, *brosme*.)

2937. BROSME BROSME (Müller).

(CUSK.)

D. 98; A. 71; P. 24; V. 5. Body cylindrical, posteriorly compressed; head flattened above. Mouth large, oblique, maxillary reaching beyond orbit; lower jaw included; several rows of sharp teeth on jaws, vomer, and palatines; barbel about 5 in head; interorbital greater than the diameter of eye. Origin of dorsal above anterior half of pectoral; pectoral round, 2½ in head; caudal rounded behind. Brownish above, the sides yellowish, sometimes mottled with brown; young uniform dark slate color, or with transverse yellow bands; vertical fins bordered with blackish, and with a white edge. (Storer.) North Atlantic, south to Cape Cod and Denmark; rare southward on our coasts. (*brosme*, a Danish name.)

*Gadus brosmæ*, MÜLLER, Prodr. Zool. Dan., 41, 1776, Denmark; FABRICIUS Fauna Grœnlandica, 140, 1780.

*Gadus lubb*, EUPHRASEN, Vet. Akad. Handl. 1794, 223, tab. 8.

*Gadus torsk*, BONNATERRE, Encycl. Meth., 51, 1788, Söndmøre, Norway; after Strom.

*Brosmius vulgaris*, FLEMING, British Anim., 194, 1828.

*Brosmius flavescens*, LE SUEUR, Mém. Mus., V, 1819, 158, Banks of Newfoundland; chin with 2 barbels; lower jaw longest.

*Brosmius flavescens*, GÜNTHER, Cat., IV, 360, 1862; STORER, Hist. Fish. Mass., 368, 1867.

*Enchelyopus brosmæ*, BLOCH & SCHNEIDER, Syst. Ichth., 51, 1801.

*Brosmius brosmæ*, GÜNTHER, Cat., IV, 360, 1862; JORDAN & GILBERT, Synopsis, 802, 1883;

GOODE & BEAN, Oceanic Ichthyology, 385, fig. 329, 1896.

*Blennius torsk*, LACÉPÈDE, Hist. Nat. Poiss., II, 508, 1800.

Family CCXV. MACROURIDÆ.

(THE GRENADIERS.)

Body elongate, tapering into a very long compressed tail, which ends in a point; scales moderate, usually keeled or spinous, sometimes smooth. Suborbital bones enlarged, sometimes cavernous. Teeth villiform or cardii-

form, in bands, on the jaws only; tip of lower jaw with a barbel; premaxillary protractile. Dorsal fins 2, the first short and high, of stiff, spine-like branched rays; the second dorsal very long, usually of very low feeble rays, continued to the end of the tail; anal fin similar to the second dorsal, but usually much higher; no caudal fin; ventrals small, subjugular, each of about 8 rays. Branchiostegals 6 or 7. Lateral line present. Gills  $3\frac{1}{2}$  or 4, a slit behind the fourth. Gill rakers small; gill membranes free or narrowly united to the isthmus, usually more or less connected; pseudo-branchiae wanting or rudimentary; pyloric caeca numerous; air bladder present. Genera 18; species about 50, chiefly of the northern seas, all in deep water. They differ from the codfishes chiefly in the elongate and degenerate condition of the posterior part of the body. Dr. Gill succinctly defines the group as "Gadoidea with an elongated tail tapering backward and destitute of a caudal fin, postpectoral anus, enlarged sub-orbital bones, inferior mouth, subbrachial ventrals, a distinct anterior dorsal, and a long second dorsal and anal converging on end of tail." We here follow Goode & Bean in the general arrangement of the genera of *Macrouridae*. Some of these can, however, be only provisionally adopted, as the characters of dentition, form of mouth, and character of the second dorsal spine or ray, are subject to much intergradation. These characters seem much more distinct on paper than they are in fact. Still, most of the genera here adopted will ultimately prove valid. (*Macruridae*, Günther, Cat., iv, 390-398, 1862.)

a. First branchial arch free, without fold of membrane across it; mouth large; second dorsal well developed, higher than the anal.

BATHYGADINÆ:

b. Gills  $3\frac{1}{2}$ ; snout short and blunt, the jaws even in front; teeth in villiform bands, sometimes obsolete; bones of head soft and cavernous; scales smooth; first dorsal low, its spine not produced. BATHYGADUS, 995.

TRACHYRINCHINÆ:

bb. Gills 4; snout rather long.

c. Teeth in upper jaw in 2 series, the outer enlarged, those in lower jaw in 1 series; mouth subterminal; barbel obsolete; nape without scaleless fossa; vomer with teeth; bones of head soft and cavernous; tail very long, flagelliform; anal fin with an elevated anterior lobe.

STRINDACHNERIA, 996.

cc. Teeth in both jaws in villiform bands; barbel developed; mouth inferior; a naked fossa on each side of nape; a row of armed scales along base of dorsal anteriorly; opercle very small; anal not elevated in front.

TRACHYRINCUS, 997.

MACROURINÆ:

aa. First branchial arch with a fold of membrane across its terminal portion; gills 4, a slit behind the fourth; barbel well developed.

d. Teeth not all in villiform bands, those of lower jaw in 1 series; mouth rather large, with more or less of lateral cleft.

e. Upper jaw without villiform band behind the enlarged anterior teeth, the inner teeth, if present chiefly uniserial, not in villiform bands.

f. Dorsal fins widely separated, the interspace greater than base of first.

g. First dorsal with the spine not serrate, its insertion over pectoral or nearly so; pectoral placed high, opposite upper angle of gill cleft; scales small, bristly; bones of head cavernous; ventrals short and weak. MALACOCEPHALUS, 998.

gg. First dorsal with its spine more or less strongly serrate; pectoral inserted below upper angle of gill cleft.

- k. Scales nearly smooth, with weak ridges which are not spinigerous. MOSELEYA, 999.
- kh. Scales rough with strong ridges. NEMATONURUS, 1000.
- ff. Dorsal fins near together, the interspace less than base of first; scales rough.
- i. Dorsal spine weak, unarmed or very nearly so; pectorals moderate. ALBATROSSIA, 1001.
- ii. Dorsal spine very strongly serrate; pectorals very long. BOGOS LOVIUS, 1002.
- ee. Upper jaw with a distinct villiform band behind the outer series of enlarged teeth; dorsal spine serrate; dorsal fins not widely separated. CHALINURA, 1003.
- dd. Teeth in villiform bands above and below, the outer scarcely enlarged and not separated from the rest; the lower band sometimes becoming a single series laterally; scales rough.
- j. Mouth wide, with considerable lateral cleft.
- k. Dorsal spine finely barbed; skull rather firm; dorsals moderately separated. CORYPHÆNOIDES, 1004.
- kk. Dorsal spine entirely smooth; bones of skull very thin and papery; dorsals well separated. HYMENOCEPHALUS, 1005.
- jj. Mouth inferior, small, with little lateral cleft; a more or less distinct ridge across the suborbital region.
- l. Scales spinous, very rough.
- m. Scales distinct, regularly imbricated.
- n. Long dorsal spine serrate in front; mouth subinferior, below the short snout. MACROURUS, 1006.
- nn. Long dorsal spine smooth; mouth wholly inferior, below the long sturgeon-like snout. CELOBRYNCHUS, 1007.
- mm. Scales indistinct, scarcely imbricated; the whole body rough-villous; dorsal spine smooth. TRACHONURUS, 1008.
- ll. Scales all thin and smooth, dorsal fin slightly serrulate. LIONURUS, 1009.

995. BATHYGADUS, Günther.

*Bathygadus*, GÜNTHER, Ann. Mag. Nat. Hist. 1878, 23 (*cottoides*).

Head large, fleshy, without prominent ridges, spiny armature or external depressions; nape elevated, hump-like. Snout broad, obtuse, not produced; mouth terminal very large, with small villiform teeth or none; suborbital ridge very low, not joined to the angle of the preoperculum. Maxillary entirely received within a groove under the prefrontal and suborbital bones, its tips narrowed and blade-like; premaxillaries protractile downward, separated anteriorly, rib-shaped, compressed vertically, very broad and without true teeth; provided posteriorly with a short flange, which is received under the maxillary; mandible received within the intermaxillary bones, without true teeth, but with minute asperities, similar to those in the upper jaw; vomer and palatines toothless. Barbel sometimes present. No pseudobranchie. Gill rakers numerous, moderate, lanceolate, with minute denticulations along their inner edge. Branchiostegal membrane free from the isthmus, deeply cleft. Branchiostegals 7, very stiff. Gill opening very wide; gills 3½; anterior gill arch free. Operculum with a blunt, spine-like prominence at its angle. Ventrals below the pectorals, many-rayed, the anterior rays produced; dorsal consisting for the most part of branched rays, higher than the anal, the

first dorsal low, without differentiated spine. Scales cycloid, unarmed; lateral line strongly arched over the pectoral. Deep seas. This genus differs from *Macrourus* and its allies in the structure of both the first and last gill arches. It is perhaps the most primitive of the family and as such is nearest allied to the *Gadidae*. (*Bathús*, deep; *Gadus*, codfish.)

- a. Pectoral and ventral fins moderate, not much, if any, longer than head.  
b. Jaws without teeth; pectoral fin broad, of 25 rays; depth  $5\frac{1}{2}$  in length.

ARCUATUS, 2938.

- bb. Jaws with small teeth; pectoral fin narrow; depth 6 to  $6\frac{1}{2}$  in length.  
c. Eye moderate, 5 in head; pectoral rays 14, the fin  $\frac{1}{2}$  as long as head.

FAVOSUS, 2939.

- cc. Eye very large,  $2\frac{1}{2}$  in head; pectoral as long as head without snout.

MACROPS, 2940.

- aa. Pectoral and ventral fins much produced, much longer than head, each reaching about halfway to tip of caudal; depth  $7\frac{1}{2}$  in length; pectoral rays 13.

LONGIFILIS, 2941.

2938. *BATHYGADUS ARCUATUS*, Goode & Bean.

Head 5 in total length; depth  $5\frac{1}{2}$ ; eye  $4\frac{1}{2}$  in head; snout  $4\frac{1}{2}$ . D. II, 9 or 10-135; A. 120; P. 25; V. 8; scales 8-140-13 or 14 (counting backward from vent to lateral line), 22 counting forward. Body shaped much as in *Chalinura simula*, but the nape still more convex; back gibbous, the dorsal outline rising rapidly from interorbital region to origin of first dorsal, whence it descends gradually to end of tail. Scales moderate, cycloid, subovate, without armature, those of abdominal region and those above pectorals the largest; lateral line strongly arched over the pectorals, length of the arched portion contained about  $3\frac{1}{2}$  times in straight portion, greatest height of arch about  $\frac{1}{2}$  its chord; scales covering all parts of head except jaws and chin. Interorbital area flat, its width 6 in head; post-orbital portion of head about  $2\frac{1}{2}$  times diameter of eye; operculum terminating in a flat obtuse spine, its length, including the flap, about equal to diameter of eye; preoperculum entire, with a prominent ridge in advance of its posterior edge; snout very broad, obtuse, the intermaxillaries extending beyond it, its width at nostrils equal to about twice length of eye; posterior extremities of intermaxillary processes elevated, producing a decided hump upon top of snout; ridge formed by prefrontal and sub-orbital bones terminating very slightly behind posterior margin of orbit, and not connected with angle of preoperculum. Nostrils immediately in front of lower part of eye, not tubular, the anterior one very small, pore-like, only about  $\frac{1}{2}$  as large as posterior one; distance of anterior nostril from tip of snout about  $\frac{1}{2}$  length of eye. Length of barbel  $6\frac{1}{2}$  in length of body, and equal to length of head without snout, more than 3 times as long as eye. No true teeth, the intermaxillaries and mandible being broad plates, covered with minute asperities; a naked space at the symphysis of intermaxillaries; distance of first dorsal from snout nearly  $3\frac{1}{2}$  times length of its base, the first spine minute, the second (in the type) somewhat mutilated, its length nearly 3 in length of head, not stouter than the branched rays, and entirely smooth; second dorsal fin separated from first by a very short interspace, equal to about  $\frac{1}{2}$  of length of eye, its rays long, subequal, the first slightly the longest, its length equal to that of base of

first dorsal; anal much lower than dorsal, the longest rays being in front, its third ray about  $\frac{1}{2}$  as long as first ray of second dorsal; this fin inserted under the seventh ray of second dorsal; about 3 of the terminal anal rays might be considered caudal rays; pectoral inserted slightly in advance of ventral, which is in about the same vertical with the origin of the first dorsal, second ray of pectoral slightly produced; length of pectoral equal to that of head without snout; ventral insertion distant from tip of snout a distance equal to that of first dorsal from snout, the first and second rays filamentous, the latter slightly the longer, and extending to the fifteenth or eighteenth ray of anal fin. Color brown; vertical fins bluish or black; peritoneum black; inside of gill covers and roof of mouth bluish. (Goode & Bean.) West Indies and Gulf of Mexico. Three specimens known; the type from near Martinique. (*arcuatus*, arched.)

*Bathygadus arcuatus*, GOODE & BEAN, Bull. Mus. Comp. Zool., XII, No. 5, 158, 1883, off Martinique, in 334 fathoms (Coll. Blake); GOODE & BEAN, Oceanic Ichthyology, 421, 1896.

2939. BATHYGADUS FAVOSUS, Goode & Bean.

Head  $5\frac{1}{2}$  in total length; depth about 6; eye 5 in head; snout about 4. D. II, 9-125; A. 110; V. 9; P. 14; B. 7; scales 10-135-16. Body heavy, stout, the profile descending gradually and in a slight curve from first dorsal to snout. Scales small, deciduous, cycloid, without armature; interorbital area slightly convex, its greatest width about 3 in head; the postorbital part of head  $2\frac{1}{2}$  times as long as eye; snout broad, oblique, its width at the nostrils a little more than that of interorbital area; nostrils close to and in front of middle of eye, the posterior somewhat the larger; no barbel. Teeth in both jaws in villiform bands, a naked space at symphysis of intermaxillaries; intermaxillary bands more than twice as wide as those of mandible; vomer and palatines toothless. Gill rakers 20 + 25, the longest on anterior arch slightly more than  $\frac{1}{2}$  eye; pseudobranchiae present, very rudimentary in some individuals, in others wanting or present only on one side; first dorsal distant from snout a distance slightly more than length of head, length of its base about equal to width of snout at nostrils, the fin consisting of 2 spines, the first minute, and 9 branched rays; length of longest dorsal spine, which is armed, 2 in head; second dorsal beginning immediately behind first, the membrane being continuous; anterior rays longest, apparently about  $\frac{1}{2}$  length of head; anal lower than second dorsal, its distance from snout about equal to  $\frac{1}{2}$  of total length; pectoral inserted under anterior rays of first dorsal and very slightly in advance of origin of ventral, its length more than  $\frac{1}{2}$  that of head; distance of ventral from snout 5 times in total length; this fin inserted nearly under base of pectoral; the first ray somewhat produced, its tip reaching to fourth ray of anal fin. Color bluish brown, darkest upon head and abdomen. West Indies. The type specimen, 350 mm. in length, was obtained by the Blake from Station LXXX, off Martinique, at a depth of 472 fathoms. (Goode & Bean.) (*favosus*, like honeycomb.)

*Bathygadus favosus*, GOODE & BEAN, Bull. Mus. Comp. Zool., XII, No. 5, 160, 1883, off Martinique in 472 fathoms (Coll. Blake); GOODE & BEAN, Oceanic Ichthyology, 420, fig. 352, 1896.



## 2940. BATHYGADUS MACROPS, Goode &amp; Bean.

Head  $5\frac{1}{2}$  in total length; depth  $6\frac{1}{2}$ ; eye  $2\frac{1}{2}$  in head; snout 5. D. II, 8—about 125; V. 8. Body somewhat compressed; scales small, deciduous, about 25 rows in an oblique line from the vent to the dorsal fin, 24 from the upper angle of operculum to the vertical through origin of the anal; interorbital area nearly flat, its width 4 in head; postorbital part of head somewhat longer than diameter of eye; snout broad, obtuse; nostrils close to eye, the posterior nearly twice as large as anterior one; maxillary extending to vertical through posterior margin of orbit, its length equal to that of head without its postorbital portion; length of mandible 3 times that of snout; intermaxillaries and mandible provided with narrow bands of villiform teeth, those of the mandible much shorter. A minute barbel, about  $\frac{1}{4}$  as long as snout. Vomer and palate toothless. Gill rakers lanceolate, elongate,  $7 + 26$ , the longest 7 in head; pseudobranchiae absent; distance of first dorsal from snout nearly 5 times in total length, second or longest ray in the typical specimen twice length of snout; second dorsal almost continuous with the first, its anterior rays the longest, about 4 times in length of head; anal inserted under fourteenth ray of second dorsal, its rays all very short; in a distance equal to length of head, counting back from insertion, there are 33 rays; pectoral inserted under first branched ray of first dorsal, its length in the most nearly perfect specimens equaling length of head without snout; ventral origin very slightly behind origin of pectoral under third branched ray of dorsal, reaching nearly to vent when laid back, its length equaling 3 times that of the snout. Branchiostegals 7. Color yellowish gray, lighter below. (Goode & Bean.) In deep waters of the Gulf of Mexico off the United States coast, in 321 to 347 fathoms. (*μακρός*, large; *ὤψ*, eye.)

*Bathygadus macrops*, GOODE & BEAN, Proc. U. S. Nat. Mus. 1885, 598, Gulf of Mexico, Lat.  $28^{\circ} 34' N.$ , Long.  $86^{\circ} 48' W.$ , in 335 fathoms (Type, No. 37339. Coll. Albatross); GÜNTHER, Challenger Report, XXII, 156, 1887; GOODE & BEAN, Oceanic Ichthyology, 423, 1896.

## 2941. BATHYGADUS LONGIFILIS, Goode &amp; Bean.

Head about  $5\frac{1}{2}$  in total length; depth  $7\frac{1}{2}$ ; eye 4 in head; snout 4. D. II, 8 or 9—about 140; P. 13; V. 8; scales about 142. Body more compressed than in *B. macrops*; scales small, cycloid, deciduous, about 25 rows from the vent upward and forward to the dorsal fin, interorbital area flattened, its greatest width  $3\frac{1}{2}$  times in total length of head; postorbital portion of head twice as long as eye; snout and nostrils normal; maxillary reaching somewhat beyond posterior margin of orbit, its length twice in distance from snout to origin of first dorsal; length of mandible  $2\frac{1}{2}$  times in snout; barbel slender, long, its length equal to  $1\frac{1}{2}$  times orbital diameter. Teeth in narrow villiform bands in each jaw, none on vomer or palatine bones; gill rakers very long and slender, numerous,  $17 + 35$ , the longest nearly 6 in head; pseudobranchiae absent; first dorsal of 2 stout spines, the first minute, the second elongate, and 8 or 9 branched rays, its distance from snout  $5\frac{1}{2}$  in total; second or longest simple ray nearly 8 times length of

snout, and reaching to or beyond the thirtieth ray of the second dorsal; second dorsal almost continuous with the first, its anterior rays longest and not diminishing rapidly in size toward tail; anal inserted under ninth ray of second dorsal, its rays much shorter than those of dorsal, and situated about same distance apart; pectorals inserted under anterior portion of first dorsal, first ray much produced, extending more than halfway from its insertion to tip of tail; ventral origin slightly behind origin of pectoral, under third branched ray of dorsal, its first ray much enlarged, extending more than halfway from its insertion to tip of caudal, its length  $2\frac{1}{2}$  times in total length; branchiostegals 7. Color yellowish gray, abdomen bluish. This form is closely allied to *B. multifilis*, described by Günther from off the Philippines (Challenger Report, XXII, 155, pl. 42, fig. B, 1887), which, however, appears to have a smaller eye, less elongate filaments, and ventrals inserted in advance of the first dorsal, while the anal appears to be further back, under the twelfth or thirteenth ray of second dorsal. Both species are provided with long, slender barbels; in other respects they are closer to *B. cottoides*, the typical species, than to *B. macrops*. (Goode & Bean.) Deep waters of the Gulf of Mexico, in 525 to 730 fathoms. (*longus*, long; *filum*, thread.)

*Bathygadus longifilis*, GOODE & BEAN, Proc. U. S. Nat. Mus. 1885, 599, Gulf of Mexico, Lat.  $28^{\circ} 47' 30''$  N., Long.  $87^{\circ} 27'$  W., in 724 fathoms (Type, No. 37338. Coll. *Albatross*); GÜNTHER, Challenger Report, XXII, 157, 1887; ALCOCK, Ann. Mag. Nat. Hist. 1890, 302; ALCOCK, l. c. 1891, 123; GOODE & BEAN, Oceanic Ichthyology, 422, 1896.  
*Hymenocephalus longifilis*, VAILLANT, Exp. Sci. Trav. et Tails, 218, pl. 23, fig. 1, 1888.

#### 996. STEINDACHNERIA, GOODE & BEAN.

*Steindachneria*, GOODE & BEAN, in AGASSIZ, Three Cruises of the *Blake*, II, 26, 1888 (no type; short diagnosis\*); not *Steindachneria*, EIGENMANN, Nematognathi, Occasional Papers, I, Cal. Ac. Sci. 1890, 100 and 202, a genus of Silurid fishes.  
*Steindachneria*, GOODE & BEAN, Oceanic Ichthyology, 419, 1896 (*argentea*).  
*Steindachnerella*, † EIGENMANN, American Naturalist, February, 1897, 159 (*argentea*).

Body compressed, with tapering tail. Mouth large, terminal. Dorsal fins continuous, both elevated anteriorly; anal divided, the anterior portion elevated, the posterior low. Teeth in each jaw biserial, the outer much enlarged, vomerine teeth present. Bones of head soft and cavernous. Eye large. Gill membranes connected anteriorly, free from the isthmus. Gill rakers slender, rather numerous; vent in anterior third of length. No pseudobranchiæ. Branchiostegals 7. No barbel. Pectorals and ventrals both below first dorsal. Scales thin, cycloid, deciduous. Deep seas. ("This remarkable genus is named in honor of Dr. Franz Steindachner, Custos of the Imperial Zoological Museum of Vienna," one of the ablest naturalists of the century.)

\* "*Steindachneria*, a Macrurid with a high differentiated first anal spine." (Goode & Bean.)

† As the original diagnosis of the Macrurid genus *Steindachneria*, although very short, is correct and sufficient for identification, the name in question should be retained for it rather than *Steindachnerella*, and the Silurid genus *Steindachneria*, Eigenmann should receive a new name.

9942. *STEINDACHNERIA ARGENTEA*, Goode & Bean.

Head  $5\frac{1}{2}$  in total; depth  $7\frac{1}{2}$ , at anal origin 8; eye  $3\frac{1}{2}$  in head; snout about  $5\frac{1}{2}$ ; interorbital width  $5\frac{1}{2}$ ; maxillary 2; premaxillary 2; mandible  $1\frac{1}{2}$ ; gill rakers 4 or  $5+19$ ; D. VIII,  $123+$ ; A.  $10+113$ ; P. 15; V. 8. Head and body compressed; tail tapering to a very fine point. Scales small, deciduous, cycloid, 6 rows between lateral line and origin of soft dorsal. Nostrils nearer eye than end of snout, the anterior nostril nearly circular, the posterior much longer and slightly concave; no barbel. Maxillary dilated at the extremity and somewhat produced downward into an obtuse point, reaching nearly to a vertical at posterior margin of orbit, and concealed by the preorbital; premaxillaries slightly protractile, much attenuated posteriorly; mandible reaching slightly behind eye. Premaxillary and mandibular teeth biserial, those of the outer series enlarged and rather widely set, some of the enlarged teeth slightly sagittate at tip; vomerine teeth well developed; upper pharyngeal teeth in 2 broad, well-developed patches. Gill rakers slender, the longest about 2 in eye. Distance from snout to first dorsal about  $\frac{1}{2}$  total length, the first spine elongate, filiform, and reaching fourteenth ray of second dorsal; base of first dorsal about 1 in head; longest ray of second dorsal about  $2\frac{1}{2}$  in head, the rays diminishing in size rapidly, the last minute; origin of anal under sixth ray of second dorsal, not far behind the vent, the anterior elevated portion consisting of 10 rays, all of which except the first are divided, the second ray longest, twice length of eye, the tenth ray only about  $\frac{1}{2}$  length of second, and separated by a small membrane from rest of fin which consists of very minute rays. Vent under fourth ray of second dorsal. Origin of ventrals under base of pectorals and about under third spine of first dorsal; first ventral ray filamentous, reaching origin of anal; pectoral reaching to below fifteenth ray of second dorsal. Gulf of Mexico. Only the type known. Length 233 mm. (*argenteus*, silvery.)

*Steindachneria argentea*, GOODE & BEAN, *Oceanic Ichthyology*, 419, fig. 351, 1896, off delta of Mississippi River, Lat.  $39^{\circ} 14' 30''$  N., Long.  $88^{\circ} 09' 30''$  W., in 68 fathoms. (Type, No. 37350. Coll. *Albatross*.)

997. *TRACHYRINCUS*, Giorna.

*Trachyrincus*, GIORNA, *Mem. Accad. Imp. Turin*, XVI, 1803, 178 (no type mentioned).

*Lepidoleprus*, RISSO, *Ichth. Nice*, 197, 1810 (*trachyrincus*).

*Oxycephas*, RAFINESQUE, *Caratteri*, 31, 1810 (*scabrus* = *trachyrincus*).

*Lepidosoma*, SWAINSON, *Nat. Hist. Class'n Fish.*, II, 281, 1839 (*trachyrhynchus*).

*Trachyrhynchus*, GÜNTHER, *Challenger Report*, XXII, 152, 1887; corrected spelling.

Snout produced in a long depressed process which is sharply pointed in front, with a sharp lateral edge, which is continued in a straight line across the suborbital region. Mouth inferior, horseshoe-shaped, placed like the mouth of a sturgeon. Teeth in both jaws in villiform bands; chin with a barbel; a scaleless fossa on each side of nape. Second dorsal well developed. Scales moderate, spinigerous; a series of larger scales, each armed with a projecting ridge, along each side of base of dorsal and anal anteriorly. Opercle small. Gill membranes scarcely united; gills 4; first gill arch free, with short, styliform gill rakers. Deep seas. This

genus and its allies differ from *Macrourus* in the important character of the structure of the first gill arch. (*τραχύς*, rough; *ρύγχος*, snout; hence properly, but not originally, spelled *Trachyrrhynchus*.)

2943. TRACHYRINCUS HELOLEPIS, Gilbert.

Head  $3\frac{1}{2}$  in total; depth 7; eye large, 4 in head, = interorbital width; snout  $2\frac{1}{2}$ , its greatest width  $1\frac{1}{2}$  in its length. D. 11. Snout depressed, flat, narrowly triangular, tapering to a sharp point, its lateral ridges continuous backward over suborbital chain and across cheek. Interorbital space wide and flat. Ethmoidal ridge not prominent. Mouth wholly inferior, U-shaped, overpassed by the snout by a distance contained  $3\frac{1}{2}$  in head. Barbel slender, short, less than  $\frac{1}{2}$  diameter of orbit. Teeth finely villiform, in very broad bands in each jaw, none of them enlarged. Maxillary reaching to or almost to vertical from hinder margin of orbit,  $3\frac{1}{2}$  in head. Opercle very small, triangular, its length behind preopercular margin scarcely more than  $\frac{1}{2}$  diameter of orbit; outer gill arch not adnate to the opercle, its lower limb with 17 short gill rakers, which are not tubercular. Distance of dorsal fin from nape  $3\frac{1}{2}$  in head, the 2 dorsal fins closely approximated; second dorsal ray not spine-like, soft and flexible, and not longer than the succeeding rays, its length  $\frac{2}{3}$  the diameter of orbit. Vent located immediately in front of origin of anal fin, its distance from ventrals  $1\frac{1}{2}$  in head. Ventrals short, inserted well in advance of base of pectorals, the outer ray little produced, its length  $1\frac{1}{2}$  in diameter of orbit. Scales all with their margins embedded, and therefore appearing non-imbriicated, the central portion of each projecting, tubercle-like, and bearing a single strong central spine, with sometimes 2 or 3 smaller ones; belly and breast sometimes covered with much smaller scales similarly armed; no naked area between bases of ventrals; enlarged plates along bases of dorsals and anal bearing each a strong compressed backwardly-curved spine, usually without distinct serrations; from the base of the central spine radiate lines of short spinous points; dorsal series of plates continued forward to the nape, the predorsal portion of the included groove covered with scales; ventral series scarcely extending beyond vent, but extending further posteriorly than do the dorsal plates; scales on top of head with a median serrated ridge; temporal fossæ small but evident, naked. Color apparently dark brown; gill cavity and peritoneum black. Pacific Ocean, off the coast of Central America, in deep water. Only the type known, a specimen 18 inches long. (*ήλος*, tubercle; *λεπίς*, scale.)

*Trachyrrhynchus helolepis*, GILBERT, Proc. U. S. Nat. Mus. 1891, 562, Pacific Coast of Central America in deep water. (Type, No. 48205.)

998. MALACOCEPHALUS, Günther.

*Malacocephalus*, GÜNTHER, Cat. Fish. Brit. Mus., IV, 396, 1862 (*levis*).

Intermaxillary teeth biserial, mandibular teeth uniserial. Mouth lateral; snout short, obtuse. Head without prominent ridges, with wide muciferous cavities. Dorsal fin over origin of pectorals, its longest spine

smooth; dorsal fins widely separated. Pectorals short, placed high, opposite upper angle of gill cleft. Scales small, bristly. Origin of lateral line at upper angle of gill cleft. (*μαλακός*, soft; *κεφαλή*, head.)

2944. *MALACOCEPHALUS OCCIDENTALIS*, Goode & Bean.

Eye  $2\frac{1}{2}$  in head; barbel slightly longer than eye; snout 4 in head; interorbital space 4. Agreeing with Günther's description of *M. lewis*, but differing in the position of the vent, the ventrals, and the anal fin, the last commencing at a distance behind the vent equal to length of snout; distance of vent from origin of ventrals less than its distance from origin of anal; ventrals originate under middle of first dorsal; origin of pectorals under that of first dorsal, the pectorals as long as head without postorbital flap; ventrals reaching to or slightly beyond origin of anal. Gill rakers rudimentary,  $x+11$ . Second dorsal spine nearly equal to length of head; first branched dorsal ray about as long as head. Atlantic Ocean, off Cape Hatteras, and Caribbean Sea. Length  $8\frac{1}{2}$  inches; a doubtful species, perhaps identical with *M. lewis*. (*occidentalis*, western.)

*Malaccocephalus occidentalis*, GOODE & BEAN, Proc. U. S. Nat. Mus. 1885, 597, off Cape Hatteras, at Albatross Station 2310, Lat.  $35^{\circ} 44'$  N., Long.  $79^{\circ} 51'$  W., in 132 fathoms. (Type, No. 37336.)

999. *MOSELEYA*, Goode & Bean.

*Moseleya*, GOODE & BEAN, Oceanic Ichthyology, 417, 1896 (*longifilis*).

This genus is near *Nematonurus*, having the mouth small, the upper teeth in 1 or 2 series, the dorsal spine weakly serrate, and the dorsal fins well separated. The chief difference lies in the scales, which are feebly ridged and nearly or quite smooth. The typical species, *M. longifilis* (Günther), is from off the coast of Japan. ("Named in honor of Prof. Henry N. Moseley, F. R. S., of Oxford University, whose contributions to natural history while naturalist of H. M. S. *Challenger* we desire to commemorate.")

2945. *MOSELEYA CYCLOLEPIS* (Gilbert).

Dorsal II-8 or 9; ventral 12; eye  $4\frac{1}{2}$  in head; snout  $3\frac{1}{2}$ ; maxillary  $2\frac{1}{2}$ . Head smooth, compressed, without conspicuous ridges; median and lateral rostral ridges terminating in slightly projecting points, the median process, a short portion of the median ridge, and the edge of the membrane connecting median with lateral processes, with spinous scales and points. Snout projecting beyond the premaxillaries for  $\frac{1}{2}$  its length. Eye small, less than snout, very slightly exceeding interorbital space; mouth small, wholly inferior, maxillary reaching vertical from posterior margin of pupil. Premaxillary teeth in 2 series, the outer similar to those in mandible, not enlarged or canine-like, the inner series smaller, directed obliquely backward; a single series of teeth in mandible, not widening into a patch at symphysis. Barbel thick at base,  $\frac{1}{2}$  length of snout. Preopercle incurved above the angle, the lower limb expanded, the marginal region striate. First dorsal inserted behind axil of pectoral (second spine broken in both specimens examined), the basal portion smooth, a single sharp

barb showing that the spine is serrate; base of first dorsal equals length of snout; interspace between dorsals exceeding length of first dorsal base by  $\frac{1}{4}$  to  $\frac{1}{2}$  length of latter. Vent immediately in advance of origin of anal, under middle of interspace between dorsals; dorsal low and inconspicuous and the anal higher, as usual in this group; pectorals very slender,  $1\frac{1}{10}$  in length of head; outer ventral ray filamentous, reaching third or fourth anal ray. Scales mostly lost, the few remaining on head either entirely smooth or bearing a single median keel with 1 or 2 low spinous points; those on body without spines, either entirely smooth or showing traces of a low median keel; 6 scales in an oblique series between lateral line and middle of base of dorsal. Color dark brown, the anterior portion of back and sides with small scattered black spots; opercles, lower side of head including gill membranes and ventral area black, as are also the mouth and gill cavity and the peritoneum. A species with the general appearance, including the protruding snout, the inferior mouth and comparatively weak dentition of *Nematonurus armatus* and *N. affinis*, but with the dorsals less widely separated, the vent anterior in position, and the scales unarmed, as in *Moseleya longifilis*. (Gilbert.) Coast of British Columbia. Two specimens, the longest 150 mm., from Station 3342, off Queen Charlotte Islands, depth 1,588 fathoms. (*κύκλος*, circle; *λεπίς*, scale.)

*Nematonurus cyclolepis*, GILBERT, Rept. U. S. Fish Comm. 1893 (1896), 458, off Queen Charlotte Islands, at Albatross Station 3342, in 1,588 fathoms.

#### 1000. NEMATONURUS, Günther.

*Nematonurus*, GÜNTHER, Challenger Report, Deep-Sea Fishes, XXII, 124, 150, 1887 (*armatus*).

Body rather robust, covered with rough, strongly-ridged scales. Head short; mouth small or moderate, more or less inferior; teeth in upper jaw rather strong, in 1 series or nearly so; lower teeth uniserial; mucous cavities small; pectoral fin inserted low, below upper angle of gill cleft; ventrals well developed, the outer ray filamentous; long ray of dorsal serrated; space between dorsals long, much greater than length of first dorsal. Deep seas. A well-marked genus, distinguished by its rough, firm scales and the wide space between dorsals. (*νήμα*, thread; *οὐρά*, tail.)

a. Depth  $6\frac{1}{2}$  in length; scales without distinct median keel.

GOODEI, 2946.

aa. Depth  $5\frac{1}{2}$  in length; scales with the median keel prominent; suborbital narrow, with well-marked mucous partitions.

SUBORBITALIS, 2947.

#### 2946. NEMATONURUS GOODEI\* (Günther).

Head  $5\frac{1}{2}$ ; depth  $6\frac{1}{2}$ ; eye 5 in head; snout  $4\frac{1}{2}$ ; interorbital width  $4\frac{1}{2}$ ; postorbital part of head  $8\frac{1}{2}$ ; first dorsal II, 8 or 9; second dorsal 105; A. 110; P. 20; V. 10; scales 7-150-18, small, strong, free portions covered by series of small vitreous spines arranged in about 6 rows; no specialization of the central row, though the median spine at margin of scale projects

\* By some inadvertence this species is recorded by Goode & Bean as a *Hymenocephalus* (Oceanic Ichth., 407). On p. 408 it is said to be a *Nematonurus*. It has obviously no affinity with *Hymenocephalus*, and is, in fact, an ally of *Nematonurus armatus*.

most strongly and is longest. Width of interorbital area a little greater than horizontal diameter of orbit and length of operculum; snout triangular, depressed, its tip in axis of body nearly on a level with lower margin of eye, its lower surface forming an angle with the body axis, about equal to that formed with same by its upper profile; superior ridge pronounced anteriorly, but ending in advance of concavity in interorbital space; lateral ridges prominent, continuing posteriorly to eye, with strong angular projections in front of nostrils; no ridges continued from supraorbital region; nostrils rather close to eye; barbel shorter than eye; tip of lower jaw under anterior nostril; cleft of mouth under posterior margin of orbit; under surface of head naked, with the exception of a few minute, spiny tubercles on under surface of mandible; suborbital ridge very slightly developed; the intermaxillary a long bone, nearly as long as the maxillary; mouth large; teeth on intermaxillary in a double series, those of the outer series much larger than the inner; teeth in mandible uniserial. Dorsal spine strongly serrated; distance of first dorsal from snout equal to nearly 4 times length of its base, its distance from anterior margin of orbit equal to length of head; first spine minute, second strongly serrated, nearly  $\frac{1}{2}$  length of head, when laid down is far from reaching origin of second dorsal; when the fin is erect its superior margin is nearly at right angles to plane of back and slightly convex; distance between dorsals twice length of base of first, the second beginning in the perpendicular from fifth ray of anal; anal about 3 times as high as second dorsal; vent under thirtieth scale of lateral line directly in advance of the anal and at a distance from ventral considerably greater than length of that fin; distance of pectoral from snout slightly more than length of head, its length less than that of dorsal spine, slightly more than  $\frac{1}{2}$  its distance from the snout, its insertion (upper axil) in middle line of body; insertion of ventral under that of pectoral, slightly in advance of that of dorsal, its first ray not greatly prolonged, about  $\frac{1}{2}$  length of distance of fin from snout; branchiostegal membrane narrowly attached to the isthmus, leaving no free margin behind; gill rakers very small tubercles, only 10 below angle on first arch. Color dark reddish brown, spines upon the scales with a metallic luster; young with 3 stellate bosses upon snout, 1 at tip, 1 at some distance upon each side. Length of specimen described 322 millimeters. (Goode & Bean.) Gulf Stream, from Cape Cod to Havana; generally abundant. (Named for George Brown Goode.)

*Macrurus asper*, GOODE & BEAN, Bull. Mus. Comp. Zool., x, No. 5, 196, 1883, Gulf Stream south of New England, Lat.  $41^{\circ} 24' 25''$  N., Long.  $65^{\circ} 33' 30''$  W., in 1,242 fathoms; name preoccupied by *Macrurus asper*, GÜNTHER; JORDAN, Cat., 131, 1885.

*Macrurus goodii*, GÜNTHER, Challenger Report, XXII, 136, 1887; substitute for *Macrurus asper*.

*Hymenocephalus goodii*, GOODE & BEAN, Oceanic Ichthyology, 407, fig. 340, 1896.

2947. NEMATONURUS SUBORBITALIS (Gill & Townsend).

Head  $5\frac{1}{2}$ ; depth  $5\frac{1}{2}$ ; eye 5 in head; snout  $4\frac{1}{2}$ ; maxillary  $2\frac{1}{2}$ . D. 12-85; A. 102; P. 19; V. 11. Mouth wholly inferior; scales closely adherent and rather large, mostly short and roundish, with considerable exposed sur-

faces, having radiating ridges beset with weak spines; head a little more than  $\frac{1}{2}$  of the entire length; snout projecting but little; median and lateral tubercles faintly developed; infraorbital narrow, divided into 2 well marked areas, an upper wider, distinguished by the glassy tubercular scales, and the narrow lower, almost skinny and scaleless; the ridge independently, is little marked; teeth biserial in the upper jaw, robust in the outer row, very weak in the inner; uniserial in lower jaw and scarcely incurved; dorsal spine strongly serrate,  $1\frac{1}{2}$  in head; pectoral  $1\frac{1}{2}$  in head; ventrals  $1\frac{1}{2}$  in head, with short filaments, reaching vent; interspace between dorsals  $\frac{1}{2}$  greater than base of first. Bering Sea. Only the type, 20 inches long, known, the above description taken from it by us. (*suborbitalis*, pertaining to the region below the eye.)

*Macrurus* (*Nematonurus*) *suborbitalis*, GILL & TOWNSEND, Proc. Biol. Soc. Wash., xi, 1897 (Sept. 17, 1897), 234, Bering Sea, southwest of Pribilof Islands, Albatross Station 3603, in 1,771 fathoms. (Type, No. 48773, U. S. Nat. Mus. Coll. Albatross.)

1001. ALBATROSSIA, Jordan & Evermann, new genus.

*Albatrossia*, JORDAN & EVERMANN, new genus (*pectoralis*).

This genus has the form and appearance of *Chalinura*, with the dentition of *Nematonurus*, and the dorsal spines of *Malacocephalus* and *Optonurus*; teeth in the upper jaw strong, in an irregular double series, the outer enlarged; the inner series growing double with age; lower teeth uniserial or nearly so; scales small, rather firm, rough; dorsal spine weak, smooth or very slightly serrate; dorsal fins close together; ventrals well developed; pectorals moderate. Size large. (Named for the good ship *Albatross*, in remembrance of her splendid contributions to our knowledge of the life of the deep seas.)

2048. ALBATROSSIA PECTORALIS (Gilbert).

Head 6 in total; depth  $1\frac{1}{2}$  in head; eye  $4\frac{1}{2}$  to 5 in head,  $1\frac{1}{2}$  in snout. D. X-128; A. 121; V. 7; P. 17; mouth wide, lateral, the short snout projecting beyond premaxillaries for a distance about equaling  $\frac{1}{2}$  diameter of orbit; suborbital ridge and lateral ridge on snout inconspicuous; a strong median ridge on snout and a pair of parallel ridges forward from above nostrils; maxillary reaching well behind vertical from posterior margin of orbit,  $2\frac{1}{2}$  in head; teeth in 2 somewhat irregular series in front of premaxillaries, the outer series enlarged, the inner directed obliquely inward, the two series merging into one laterally; mandible with a single row, similar to inner series of upper jaw; barbel short,  $\frac{2}{3}$  to  $\frac{7}{8}$  diameter of orbit; angle of preopercle bluntly rounded, not produced; outer gill arch adnate, as usual in *Macrurus*, 7 short tubercular gill rakers present on its free portion; first dorsal spine slender and weak, with 1 or 2 small retrorse prickles near its middle; distance between dorsals equal to  $\frac{2}{3}$  base of first; vent immediately in front of anal origin, its distance from base of ventrals slightly more than  $\frac{1}{2}$  head; pectorals long and narrow, reaching vertical from ninth or tenth ray of second dorsal, more than  $\frac{1}{2}$  length of head; outer ventral ray produced into a long slender filament, reaching  $\frac{1}{2}$  the



distance from its base to front of anal; scales rather small, 10 or 11 in a series between lateral line and origin of second dorsal or middle of first dorsal; scales on sides very thin and flexible, readily deciduous, each furnished with low diverging ridges, usually 3 in number, bearing few minute spinules, and projecting but little beyond the margins of the scales; entire head, including snout and mandibles, invested with much smaller scales irregularly imbricated, those on the opercles marked similarly to those on sides, the others usually each with a single median ridge terminating in a spinous point; no naked spots or pits on head or between ventral fins; a small narrow area behind and below axil of pectorals. Color light grayish, darker on belly and head; mouth, gill cavity, and peritoneum black; lateral line black; dorsals and ventrals dusky; anal lighter, edged with blackish; pectorals black. Bering Sea to Oregon. Specimens have been taken at *Albatross* Stations 3071, 3074, and 3075, in depths of 685 to 877 fathoms, off the coast of Oregon, and from near Bogoslof Island in Bering Sea in 664 fathoms. It is a large, firm-fleshed species, easily recognized. (*pectoralis*, pertaining to the pectoral.)

*Macrurus (Malacocephalus) pectoralis*, GILBERT, Proc. U. S. Nat. Mus. 1891, 563, off the coast of Oregon. (Coll. Dr. Gilbert.)

*Macrurus (Nematonurus) magnus*,\* GILL & TOWNSEND, Proc. Biol. Soc. Wash., XI, 1897 (Sept. 17, 1897), 234, Bering Sea, southwest of Pribilof Islands. (Types, No. 48770 and 48771, U. S. Nat. Mus. Coll. *Albatross*.)

*Albatrossia pectoralis*, JORDAN & GILBERT, Report Fur Seal Invest., 1898.

#### 1002. BOGOSLOVIUS, Jordan & Evermann, new genus.

*Bogoslovius*, JORDAN & EVERMANN, new genus (*clarki*).

This genus is close to *Chalinura*, from which it is distinguished by its dentition, having the teeth in the upper jaw in 2 series, the outer slender and sharp, slightly arrow-shaped; those of the inner small, close set, replacing the villiform band of *Chalinura*. Scales excessively rough; ventral filament produced; dorsal spine filamentous, sharply serrate; dorsal fins close together; pectorals inserted below upper angle of gill opening. Deep seas. (Named for the volcanic island, St. John Bogoslof, in Bering Sea, near which the typical species was dredged.)

*a.* Ventrals much longer than head, reaching far beyond front of anal. CLARKI, 2949.

*aa.* Ventrals shorter than head, scarcely reaching front of anal. FIRMISQUAMIS, 2950.

\* We have examined the type and cotypes of *Macrurus (Nematonurus) magnus*, Gill & Townsend, and find them to agree fully with *Albatrossia pectoralis* (Gilbert). The type may be redescribed as follows: Head 5½; depth 7½; eye 4½ in head; snout 4½ to 4¾; maxillary 2½; pectoral 2 in head; ventral with short filament, 2½ in head. Mouth large, with lateral cleft. Dorsals well separated, the interspace not ¼ base of first dorsal; long dorsal spine smooth, or with 1 or 2 roughnesses near its tip, its length 3½ in head; second dorsal low; pectoral inserted low, below angle of opercle. Scales moderately large, readily deciduous, decidedly oblong or long, with a small exposed surface which is beset with about 5 radiating ridges with conspicuous spinigerous ridges on dorsal surface, but not armed at tip; head regularly conical; snout rather long, projecting ¼ its length beyond mandible; tubercles feebly developed, plain, and continuous from 3 parallel ridges; infra-orbital flat, with the crest rather nearer the orbit than its lower margin; its entire surface senly; teeth in the upper jaw biserial or triserial in front, the outer series strongly hooked, the inner series considerably smaller and well separated from the outer series; an irregular series between in the type specimen; teeth in lower jaw uniserial or irregularly biserial. Three specimens, the largest (type of *M. magnus*) 43 inches long.

## 2949. BOGOSLOVIUS CLARKI, Jordan &amp; Gilbert.

Eye  $4\frac{1}{2}$  in head; maxillary  $2\frac{1}{2}$ . D. II, 12- ; P. 19; V. 10. Snout short, slightly exceeding diameter of eye,  $3\frac{5}{10}$  in head; median and nasal ridges very little projecting anteriorly, without radiating spines; tip of snout very little projecting beyond the mouth, for a distance not exceeding  $\frac{1}{2}$  the interspace between ends of median and nasal ridges. Suborbital ridge inconspicuous, scarcely extending beyond the eye; mucous pores on head prominent. Mouth large, oblique, the lower jaw included, the maxillary nearly reaching vertical from posterior edge of orbit. Outer premaxillary teeth slender, sharp, unequal, rather distant, not very strong, slightly widened and arrow-shaped near tip, becoming very small toward angle of mouth; within this, and well separated from it, a close-set series of short teeth directed inward. Mandibular teeth slender, unequal, in a single series corresponding to outer series in upper jaw, slightly widening at symphysis, which is not prominent. Barbel very short, less than  $\frac{1}{2}$  diameter of pupil. Eye of moderate size, equaling distance from tip of snout to middle of anterior nostril,  $1\frac{1}{10}$  in interorbital width. Preopercle broadly rounded, the angle little produced backward, leaving a strip of interopercle exposed along its entire length. Gill membranes joined to the isthmus, with a narrow free edge. Gill rakers very short and thick, 3 + 12 in number, including rudiments. Dorsal beginning above base of pectorals, the second spine long, filamentous at tip,  $1\frac{1}{2}$  in head, its anterior margin sharply serrate, except in basal third; base of first dorsal  $2\frac{1}{2}$  in head; interspace between dorsals very short, usually less than diameter of pupil. Pectorals very long and slender, equaling or exceeding length of head behind snout; insertion of pectorals below upper angle of gill opening. Outer ventral ray excessively produced, twice or more than twice length of head in uninjured adults, reaching base of fiftieth anal ray or beyond. Vent immediately before anal origin. Scales in a strip along the back firm and very rough, none others preserved in our specimens; scales with 3 to 5 sharp, radiating ridges, each ridge with several sharply projecting spines, the posterior of which project beyond the margin of the scale. Color very light gray, the vertical fins blackish posteriorly; mouth and gill cavity and peritoneum jet-black. Bering Sea. Known from 4 specimens, 24 to 41 cm. long, from *Albatross* Station 3634, off Bogoslof Island, in 664 fathoms. (Named for George Archibald Clark, secretary of the Fur Seal Commission for 1896 and 1897, in recognition of his researches on the mammalia of Bering Sea.)

*Bogoslavius clarki*, JORDAN & GILBERT, Report Fur Seal Invest., 1898, Bering Sea off Bogosicf Island, in 664 fathoms.

## 2950. BOGOSLOVIUS FIRMISQUAMIS (Gill &amp; Townsend).

Head 5 in total; depth  $6\frac{1}{2}$ ; eye  $4\frac{1}{2}$  in head; snout  $3\frac{1}{2}$ ; second dorsal spine  $1\frac{1}{2}$  in head; pectoral 2; ventral  $1\frac{1}{2}$ ; maxillary  $2\frac{1}{2}$ . D. II, 10-126; A. 105; P. 20; V. 8. Scales firmly affixed, oblong or rather short, and with considerable exposed surfaces, which have subequal radiating ridges beset with numerous acute spinelets, the ridges varying from 3 to 8 in number; head regularly convex in profile: rostral tubercles obsolete and infraor-

bital ridge rounded; barbel greater than pupil; teeth biserial or partly triserial above; second dorsal spine with short retrorse serrae, the lower fifth smooth; base of first dorsal  $3\frac{1}{2}$  in head; interspace between dorsal fins  $\frac{1}{2}$  base of first dorsal, greater than diameter of pupil. This species is distinguishable from most American *Macrouri* by the very firm scales, and from *B. clarki* by the much shorter ventral. Bering Sea. Only the type, 31 inches long, known. (*firmus*, firm; *squama*, scale.)

*Macrurus firmisquamis*, GILL & TOWNSEND, Proc. Biol. Soc. Wash., xi, 1897 (Sept. 17, 1897), 234, Bering Sea, southwest of Pribilof Islands. (Type, No. 48772, U. S. Nat. Mus. Coll. Albatross.)

### 1003. CHALINURA, Goode & Bean.

*Chalinura*, GOODE & BEAN, Bull. Mus. Comp. Zool., x, No. 5, 198, 1883 (*simula*).

*Chalinurus*,\* GÜNTHER, Challenger Report, xxii, 124, 144, 1887; change in spelling.

Scales cycloid, fluted longitudinally, with slightly radiating striae. Snout long, broad, truncate, not much produced. Mouth lateral, subterminal, very large. Head without prominent ridges, except the subocular ones and those upon the snout. Suborbital ridge not reaching angle of preopercle. Teeth in the upper jaw in a villiform band, with an outer series much enlarged, those of the lower jaw uniserial, large. No teeth on vomer or palatines; small pseudobranchiae present. Gill rakers spiny, strong, depressible, in double series on anterior arch. Ventrals below the pectorals; chin with a barbel. Dorsal spine serrate; soft dorsal much lower than anal. Deep sea fishes. Species numerous. This genus is allied to *Macrourus*, differing in the dentition; the genus *Optonurus*, with dorsal spine unarmed, is very close to *Chalinura*. (*χαλινός*, a strap or thong; *ὄψρά*, tail.)

a. Snout long, longer than eye, which is 5 in head; pectoral  $1\frac{1}{2}$  in head; dorsal spine  $1\frac{1}{2}$  in head; scales 130. SERRULA, 2951.

aa. Snout moderate, about as long as eye, which is 4 in head; dorsal, pectoral, and ventral produced, the pectoral  $1\frac{1}{2}$  in head, the dorsal spine and ventral filament each about as long as head. FILIFERA, 2952.

aaa. Snout very short, as long as eye, which is 5 in head; ventrals very long.

SIMULA, 2953.

### 2951. CHALINURA SERRULA, Bean.

Head  $5\frac{1}{2}$  in total length. D. II, 9-76 (?); scales 7 or 8-130-17; Br. 6. Cheeks and opercles scaly; snout with a median serrated keel on the nose; diameter of eye less than length of snout, 5 in head; maxillary reaching vertical from posterior margin of eye, its length  $2\frac{1}{2}$  in head; mandible about 2 in head, a row of 5 pores on its under surface and 6 pores on the edge of the suborbitals; branchiostegal membrane narrowly free from the isthmus, the first gill opening restricted as in *Macrourus*; gill rakers small tubercles, 11 below the angle of the first arch, and only 1 or 2 above the angle; length of pectoral equals postorbital part of head; ventrals

\* Goode & Bean rightly protest against the wanton "action of the English ichthyologists in changing the form of the generic name" *Chalinura*. *Chalinura* is perfectly correct, and should be used even if it were not so, as it is the original form, the only reason for changing it being that other generic names in the group end in *urus*.

about as long as head; longest dorsal spine strongly serrated and nearly equaling length of head without snout; dorsals separated by an interspace  $\frac{2}{3}$  as long as head. Color brown; head, abdomen, and inside of mouth purple, the purple areas less marked in the type specimen, which is  $12\frac{1}{2}$  inches long. Coast of British Columbia, east of Prince of Wales Island, in 1,569 fathoms. (*serrula*, a fine saw.)

*Chalinura serrula*, BEAN, Proc. U. S. Nat. Mus. 1890, 37, east of Prince of Wales Island, in 1,569 fathoms. (Coll. *Albatross*.)

2952. CHALINURA FILIFERA, Gilbert.

D. II, 12 to 14; P. 20 to 22; V. 9 or 10; eye 4 in head. Snout short, slightly exceeding diameter of eye,  $3\frac{1}{2}$  in head, median ridge and nasal ridges terminating each in a much projecting point, furnishing each with a short rosette of radiating spines and ridges, outline between these points concave; tip of snout projecting beyond premaxillaries for a distance equaling that which separates the central rosette from 1 of the lateral ones; infraorbital ridges inconspicuous, not reaching angle of preopercle behind or bony portion in front. Mouth large, slightly oblique, with extensive lateral cleft, the maxillary reaching vertical from posterior margin of pupil,  $2\frac{2}{3}$  in head, equaling distance from tip of snout to middle of eye. Outer series of teeth in premaxillary strong, succeeding from a narrow band of smaller cardiform teeth; mandibular teeth similar to inner band of upper jaw, the band becoming slightly wider at the prominent symphysis. Barbel short,  $\frac{1}{2}$  to  $\frac{2}{3}$  length of snout. Eye large, the diameter of orbit slightly less than interorbital width on snout. Angle of preopercle produced backward, concealing all but the extreme posterior angle of interopercle, the margin appearing serrulate when divested of skin; gill membranes joined to isthmus, with a posterior free margin; gill rakers very short and heavy, 1 + 11. Dorsal beginning vertically above base of pectorals, the second spine extremely long and slender, smooth basally, the terminal half rather strongly toothed, becoming very slender toward tip and terminating in a long membranaceous filament. (In 1 specimen it exceeds length of head, in the others it equals  $\frac{2}{3}$  that length.) Length of base of first dorsal equaling  $\frac{1}{2}$  length of head; interspace between dorsals short,  $\frac{2}{3}$  to  $\frac{1}{2}$  length of snout. Pectorals very long and slender, equaling the head without the snout; outer ventral rays very long and filamentous, equaling length of head; vent immediately in advance of anal origin. Scales rather thin, those on back and sides with above 5 diverging ridges, each of which bears a number of short rigid spinules directed very obliquely backward, the posterior projecting but little beyond the margin of the scale; 8 or 9 scales in an oblique series between the middle of first dorsal and the lateral line. Dark brown; the fins, gill membranes, lips, nostrils, and underside of snout black; anterior part of mouth and lining of gill cavity purple; peritoneum blackish brown. Related to *C. serrula*, Bean, from the same region and depth, differing in the larger eye, shorter mental barbel, longer snout, longer pectoral fins, shorter interspace between dorsals and the longer dorsal fin. Coast of British Colum-

bia; known from 3 specimens, 520 to 550 mm. long. (Gilbert.) (*flum*, thread; *fero*, I bear.)

*Chalinura flifera*, GILBERT, Rept. U. S. Fish Comm. 1893 (1896), 458, off Queen Charlotte Islands, at Albatross Station 3342, in 1,588 fathoms.

2953. *CHALINURA SIMULA*, Goode & Bean.

Head 5; depth 6; orbit 6 in head; snout 3; interorbital width greater than eye; postorbital part of head 3 times as long as eye; opercle 2 in upper jaw. D. II, 9-113; A. 118; P. 20; V. 9; Br. 6; scales 8-150-17 to 19. Body shaped much as in *Coryphænoides*, but rather stout; back more gibbous in profile, the dorsal outline rising quite rapidly from the interorbital region to origin of first dorsal, thence descending almost in a straight line to end of tail. Preopercle emarginate on its posterior limb. Snout broad, obtuse, scarcely projecting beyond the mouth, its width nearly as great at tip as its own length; median ridge very prominent, gibbons in outline when viewed laterally; lateral ridges starting almost at right angles with the median, and continued upon sides of head; no supraorbital ridges. Nostrils in front of middle of eye, and nearer its anterior margin than to tip of snout; barbel longer than eye; teeth in upper jaw in a broad villiform band, the outer series very much enlarged; lower jaw with teeth in a single series. Scales rather small, but with indications, particularly on the head, of radiating striae. Origin of first dorsal from snout  $4\frac{1}{2}$  in its base, or from anterior margin of orbit 1 in head; first dorsal spine very short, second rather stout,  $1\frac{1}{2}$  in head, and with a simple serration anteriorly, the serræ closely appressed to the spine; second dorsal separated from the first by a distance equal to length of upper jaw; anal high, its average rays about 3 times as long as those of dorsal, inserted slightly behind perpendicular from last ray of first dorsal; pectoral inserted over base of ventral; origin of ventral from snout less than its longest ray, which is produced in a filament extending to base of eighteenth anal ray. (Goode & Beau.) West Indies and Gulf Stream, in deep water. (*simulus*, png-nosed.)

*Chalinura simula*, GOODE & BEAN, Bull. Mus. Comp. Zool., x, No. 5, 199, 1883, Gulf Stream, at Blake Station 308, Lat.  $41^{\circ} 25' 45''$  N., Long.  $65^{\circ} 35' 30''$  W., in 1,242 fathoms; JORDAN, Cat., 132, 1885; GOODE & BEAN, Oceanic Ichthyology, 412, fig. 345, 1896. *Macrurus simulus*, GÜNTHER, Challenger Report, xxii, 148, 1887.

1004. *CORYPHÆNOIDES*, Gunner.

*Coryphænoides*, GUNNER, Trondhj. Selsk. Skrift., III, 50, 1765 (*rupestris*).  
*Branchiostegus*, RAFINESQUE, Analyse de la Nature 1810, 86 (substitute for *Coryphænoides*).

Snout short, obtuse, high, obliquely truncated, soft to the touch, except its bony center; mouth broad, terminal, its cleft lateral; head without prominent ridges, the membrane bones of the side of the head soft and papery; teeth villiform in both jaws, those in the outer series of upper jaws somewhat enlarged. Scales spinous, second or elongate dorsal ray finely serrated in front. Lower jaw with a barbel at tip. Deep Sea.

Close to *Macrourus*, differing in the larger terminal mouth. (*κορυφαίνα*, *Coryphæna*; *εἶδος*, resemblance.)

a. Head 4 in length; gill rakers 4+15=19.  
aa. Head 6 in length; gill rakers 3+11=14.

RUPESTRIS, 2954.

CARAPINUS, 2955.

2954. *CORYPHENOIDES RUPESTRIS*, Gunner.

D. 10; P. 19; V. 7; gill rakers 4+15=19. Head short, rather compressed; snout short, obliquely truncated in front; cleft of mouth wide, lateral, extending to beyond the center of eye; intermaxillary not much shorter than maxillary. Teeth in villiform bands in each jaw; barbel very small. Interorbital space convex, its width being considerably more than diameter of eye, which, in a specimen 3 feet long, is equal to the length of the snout and  $\frac{1}{2}$  of that of the head. Scales equally rough over the whole of their surface, all the spinelets being directed backward; 7 or 8 scales in a transverse series between the dorsal fin and the lateral line; head entirely covered with small scales. Anterior dorsal spine armed with numerous small closely set barbs; outer ventral ray produced into a long filament. Distance between the vent and isthmus  $\frac{2}{3}$  the length of the head. The gill membrane entirely free from the isthmus behind. Intermaxillary continues beyond its vertical process and extending almost as far back as the maxillary, these 2 bones being about equal in length; last third of intermaxillary toothless; intermaxillary teeth in a very narrow band, which is uniform in width, the outer teeth only slightly enlarged; mandible with villiform teeth in a broad bunch-like band at the symphysis and becoming uniserial behind. Eye nearly circular. Snout projecting slightly. Gill rakers longer and less tubercular in character than in *Macrourus berglax* and *M. acrolepis*. The suborbital ridge feebly developed and very abruptly curved upward and narrowed in front of the eye where it joins the nasal ridge. In *M. berglax* and *M. acrolepis* the suborbital ridge is very strong and is continued almost in a straight line toward the nasal ridge. (Goode & Bean.) Arctic seas and the north Atlantic, on both coasts south to the banks of Newfoundland and Norway, in deep water. (Eu.) (*rupestris*, living about rocks.)

*Coryphænoides rupestris*, GUNNER, Trondhjem Selsk. Skrift., III, 50, pl. 3, fig. 1, 1765, Norway; COLLETT, Norges Fiske, 131; JORDAN & GILBERT, Synopsis, 812, 1883; GOODE & BEAN, Oceanic Ichthyology, 402, 1896.

*Lepidoleprus norvegicus*, NILSSON, Prodr. Ichth. Scand., 51, 1832, Norway.

*Coryphænoides norvegicus*, GÜNTHER, Cat., IV, 396, 1862.

*Macrourus stromii*, REINHARDT, Dansk. Vidensk. Aftandl., VII, 129, 1828; (GAIMARD), Voy. Skand., Poiss., pl. 11.

*Macrurus rupestris*, GÜNTHER, Challenger Report, XXII, 138, 1887.

2955. *CORYPHENOIDES CARAPINUS*, Goode & Bean.

Head 6. D. II, 8-100; A. 117; V. 10; eye 4 in head. Snout acute, projecting beyond the mouth, its tip at a distance from the mouth equal to or greater than diameter of eye. Bones of head very soft and flexible, its surface very irregular, there being a very prominent subocular ridge, a prominent ridge extending from tip of snout to middle of interorbital space, and a curved ridge extending from upper anterior margin of orbit

over cavity containing nostrils to a prominent point at side of and slightly posterior to tip of snout; barbel  $\frac{3}{4}$  as long as eye. Interorbital space almost twice diameter of eye, equal to length of upper jaw; preoperculum crenulate; upper jaw extending to vertical through posterior margin of pupil, its length equaling  $\frac{1}{4}$  that of head without snout; mandible extending behind vertical through posterior margin of orbit, its length 3 times in distance from tip of snout to origin of first dorsal. Teeth in villiform bands on intermaxillary and mandible, the mandibular series uniserial in about the second half of its length. First ray of dorsal very short, second compressed anteriorly and serrated, with slender teeth closely appressed and bent upward, its length equaling length of head and greater than height of body; this fin seated upon a hump-like elevation of the back, its base as long as snout; second dorsal beginning over tenth or twelfth anal ray, and at a distance from end of first dorsal equal to length of head without snout; vent located not far behind vertical from end of first dorsal. Scales 22 to 24 in a transverse series (the position of the lateral line can not be determined, but there appear to be 4 above it); scales oval, membranaceous, showing several parallel ridges composed of small spines. Gill membrane very deeply cleft and attached to the isthmus; gill rakers short and stout, about 11 below the angle on the first arch. (Goode & Bean.) Gulf Stream, in deep water. (*carapinus*, formed as in *Carapus*.)

*Coryphaenoides carapinus*, GOODE & BEAN, Bull. Mus. Comp. Zool., x, No. 5, 195, 1883, Gulf Stream, Lat.  $41^{\circ} 24' 45''$  N., Long.  $65^{\circ} 35' 30''$  W., in 1,242 fathoms (Type in M. C. Z. Coll. Blake); GÜNTHER, Challenger Report, Deep-Sea Fishes, XXII, 139, 1887; GOODE & BEAN, Oceanic Ichthyology, 404, fig. 339, 1896.

#### 1005. HYMENOCEPHALUS, Giglioli.

*Hymenocephalus*, GIOLIOI, Pelagos, Genoa, 228, 1884 (*italicus*).

*Mystacourus*, GÜNTHER, Challenger Report, Deep-Sea Fishes, XXII, 124, 1887 (*italicus*).

This genus is closely allied to *Coryphaenoides*, differing in the smooth dorsal spine, and the membranaceous skull. First dorsal broad, placed far forward over base of pectoral; second dorsal and anal origins nearly opposite, and separated by a considerable space from the vertical from the end of first dorsal; vent far from ventrals. Head large, naked, soft, and cavernous; snout abrupt, perpendicular, or parabolic; mouth lateral, wide. Eye very large, orbital margin forming part of profile of head. Barbel long. Pectoral rather narrow (10 to 16 rays). Scales thin, deciduous, with fine short spines. Under parts in advance of ventral wholly or partly naked. Deep seas. Remarkable for the papery structure of the bones of the head. (*ὕμην*, membrane; *κεφαλή*, head.)

#### 2956. HYMENOCEPHALUS CAVERNOSUS (Goode & Bean).

Head about 6 in total length; depth 7. D. II, 10-133; A. 27 rays, in a space equal to length of head. Body stoutish, the bones of head very soft and cavernous, spongy, in many places without muscular covering; interorbital area doubly concave, with a spinous medial ridge, its greatest

width about  $2\frac{1}{2}$  in length of head; postorbital portion of head about  $\frac{1}{2}$  its length,  $1\frac{1}{2}$  as long as eye, which is circular, its diameter contained  $2\frac{1}{2}$  times in length of head. Snout broad, very obtuse, its width at nostril nearly equal to interorbital width, its length  $4\frac{1}{2}$  times in that of the head; nostrils normal. Teeth in each jaw in villiform bands, very small; a naked space at the symphysis of intermaxillaries; vomer and palatine toothless. Gill-rakers very short, minute, and rather numerous, about 18 below angle of anterior p. h. Pseudobranchiæ absent. Barbel  $\frac{2}{3}$  as long as eye. First dorsal composed of 2 spines, the first minute, inserted at a distance from the snout equal to length of head, the second as long as head without snout, and 10 branched rays, its base equal to diameter of eye; second dorsal almost rudimentary, its rays remarkably short, about 133 in number, its distance from first dorsal  $\frac{1}{2}$  length of head; anal much higher than second dorsal, its distance from snout contained about  $3\frac{1}{2}$  times in total length; anterior anal rays longest, in length about  $\frac{2}{3}$  diameter of eye; pectoral inserted under first branched ray of first dorsal, its length equal to twice that of eye and about  $\frac{2}{3}$  that of head. Scales (on type) mostly wanting, except a few on breast and nape, these being rough with small points, dentate behind. Ventral slightly behind the pectoral, its first ray filamentous, reaching to the base of the tenth anal ray, consisting of 11 rays. Color gray, with silvery tints on sides; abdomen and lips dark. (Goode & Bean.) Gulf of Mexico, in deep water. One young individual known. Length 162 mm. (*cavernosus*, cavernous.)

*Bathygadus cavernosus*, GOODE & BEAN, Proc. U. S. Nat. Mus. 1885, 598, Gulf of Mexico, at Albatross Station 2398, Lat.  $28^{\circ} 45' N$ , Long.  $86^{\circ} 26' W.$ , in 227 fathoms (Type, No. 37337. Coll. *Albatross*); GÜNTHER, Challenger Report, xxii, 156, 1887.

*Hymenocephalus cavernosus*, GOODE & BEAN, Oceanic Ichthyology, 408, fig. 341, 1896.

#### 1006. MACROURUS, Bloch.

*Macrourus*, BLOCH, Ichth., v, 152, 1787 (*rupestris* = *berglax*).

*Macruropus*, BLEEKER, Versl. Med. Akad. Welenth. Amsterd., viii, 1874, 369 (*serratus*).

*Macrurus*, GÜNTHER, Cat., iv, 392, 1862; corrected spelling.

Snout broadly conical, high, projecting beyond mouth; mouth moderate, its cleft horizontal, U-shaped, entirely inferior; teeth in both jaws in villiform bands, those of the outer series not enlarged; head with roughened bony ridges, one of which, on the suborbital and preorbital, simulates the suborbital stay of the Cottoids; eyes very large; scales imbricate, very rough, keeled. Dorsal spine long, serrated on the anterior edge. Deep water fishes. (*μακρός*, long; *ὄψα*, tail, hence correctly written *Macrurus*, but *Macrourus* is the original name as given by Bloch.)

a. Top of head with 4 to 6 distinct ridges; depth 6 to 7 in length; 5 scales between lateral line and dorsal.

b. Anal rays 148; scales each with a strong ridge. BERGLAX, 2957.

bb. Anal rays 121; scales each with 3 to 5 spinules, otherwise almost unarmed; ridges c top of head very rough. HOLOTRACHYS, 2958.

aa. Top of snout with indistinct ridges or with none.

c. Pectoral fin moderate,  $1\frac{1}{2}$  to 2 in head.

d. Body rather elongate, the depth 7 to 8 in length; bones of head rather firm; dorsal spine strongly serrated.



- e. Head short,  $6\frac{1}{2}$  in head; pectoral more than  $\frac{1}{2}$  head; snout with bony ridges above. BAIRDII, 2959.  
 ee. Head  $5\frac{1}{2}$  in length; ventrals 5 in body; pectoral  $1\frac{1}{2}$  in head. LEPTURUS, 2960.  
 eee. Head longer,  $4\frac{1}{2}$  in length; pectoral  $\frac{1}{2}$  as long as head; eye as long as snout, 4 in head; ventral 8 in body. ACROLEPIS, 2961.  
 dd. Body rather robust, the depth  $5\frac{1}{2}$  in length; head without ridges above; scales spinous, not ridged; dorsal spine  $1\frac{1}{2}$  in head. STELGIDOLEPIS, 2962.  
 ce. Pectoral fin elongate, about as long as head; head elevated, not ridged above, the bones soft; eye large; second dorsal spine rough, nearly as long as head; scales each with 7 to 9 ridges. CINEREUS, 2963.

## 2957. MACROURUS BERGLAX, Lacépède.

D. 12-124; A. 148; P. 18 or 19; V. 8. Short snout, subtriangular, pointed in front, much shorter than the large eye, which is  $\frac{1}{4}$  or  $\frac{1}{2}$  length of head in the adult. Internaxillary very short,  $\frac{1}{2}$  length of maxillary, and not continued beyond its expanded vertical process. Eye oblong. Whole under surface of head below suborbital and nasal ridge naked; axil of pectoral naked; space between ventrals scaled; body scales each with a single strong median keel, made up of 5 to 8 spines directed backward; some scales, particularly of head, have also 2 lateral keels; 6 longitudinal series of scales between first dorsal fin and lateral line; first dorsal spine indistinctly denticulated toward the point; length of pectoral nearly or quite  $\frac{1}{2}$  length of head; longest spine of dorsal very finely serrated along its anterior margin, the serrations becoming obsolete near its base. Vent situated behind origin of second dorsal fin. Gill rakers very small, tubercular, 9 to 11 on the first arch; gill membranes broadly joined, free from the isthmus behind. This form, originally discovered on the coast of Norway, has been found abundantly as far south as Georges Bank, where the halibut fishermen catch it, or some closely allied form, on their trawls. The first specimen seen by American naturalists was picked up floating at the surface off the mouth of New York Harbor. The *Albatross* obtained it from Station 2528, in Lat.  $41^{\circ} 47' N.$ , Long.  $65^{\circ} 37' 30'' W.$ , at a depth of 677 fathoms. Günther knew it from Finmark and Greenland, as well as from New England. He calls attention to remarkable individual variations in the specimens examined by him. (*berglax*, Norwegian name, from *berg*, cliff; *lax*, salmon.)

*Macrourus berglax*, LACÉPÈDE, Hist. Nat. Poiss., III, 170, 1800, Greenland, Søndmøre; JORDAN, Cat. Fish. N. A., 131, 1885.

*Macrourus fabricii*, SUNDEVALL, Vet. Akad. Handl. 1840, 6; COLLETT, Norges Fiske, 128, 1875; LILLJEBORG, Sverig. og. Norges Fiske, 242; GOODE & BEAN, Cat. Fish. Essex Co. and Mass. Bay, 7, 1879; GÜNTHER, Challenger Report, XXII, 130, 1887.

*Macrourus rupestris*, GÜNTHER, Cat. Fish. Brit. Mus., IV, 390, 1862 (not of Gunner).

*Macrurus berglax*, GOODE & BEAN, Oceanic Ichthyology, 391, fig. 334, 1896.

## 2958. MACROURUS HOLOTRACHYS, Günther.

Head  $4\frac{1}{2}$  in length; depth  $6\frac{1}{2}$ . D. 12-115 to 125; A. 121; P. 20 or 21; V. 5; eye large, round, as long as snout,  $2\frac{1}{2}$  in head, much wider than interorbital space. Snout triangular, each point with a tubercle, covered with

strong spines, this border continued as a strong ridge below eye, extending across opercle, this crest covered with coarse, spinous tubercles; mouth rather small, the maxillary reaching middle of eye; teeth very small, close set. Head with salient ridges above, covered with spinous scales; 1 ridge above eye, toward upper angle of gill opening, another ridge along the vertex, nearly parallel with this above it, besides a short temporal ridge; vent far back, under seventh ray of second dorsal. First dorsal not far behind eye, the long ray slightly serrulate; ventrals with a short filament. Scales each with a median crest of 3 to 5 spinules, otherwise almost unarmed. Five scales between lateral line and dorsal. (Collett.) Depths of the Atlantic. Known from 2 specimens, the type 9 inches long, from the mouth of Rio de la Plata, in 600 fathoms; the second, above described, about a foot long, from the banks of Newfoundland, in 1,267 fathoms. (*όλος*, wholly; *τραχύς*, rough.)

*Macrurus holotrachys*, GÜNTHER, Ann. Mag. Nat. Hist. 11, 1878, 24 mouth of Rio de la Plata in 600 fathoms; GÜNTHER, Challenger Report, xxii, pl. 28, fig. B, 1887; COLLETT, Compagnes Scient. de l'Hirondelle, 1896, 83, pl. 2, fig. 6; GOODE & BEAN, Oceanic Ichthyology, 396, 1896.

2950. MACROURUS BAIRDII, Goode & Bean.

(COMMON RAT-TAIL.)

Head  $6\frac{1}{2}$  in total length; depth 8; greatest width 13. D. II, 11-137; A. 120; P. 15; V. 7; scales 6-152-19 or 20. Body much compressed posteriorly, tapering from first dorsal to tip of tail; scales irregularly polygonal, the free portions covered with transparent vitreous spines, arranged in from 10 to 12 irregular longitudinal rows. On head and upper part of body, in advance of the first dorsal, the median row of spines most prominent, and presenting the appearance of a low median keel. Lateral line nearly straight, formed by a smooth groove, which replaces 2 or 3 median rows of spines of each scale; greatest height at posterior margin of orbit greater than width at same point,  $1\frac{1}{4}$  times in length of head; width of interorbital area equal to length of snout and length of maxillary; length of postorbital region about equal to horizontal diameter of orbit; length of operculum about  $\frac{1}{2}$  length of mandible. Snout sharp, a front view presenting 4 ridges radiating from tip at right angles to each other, the lower one being merely a fold in the skin of the under surface of the head, horizontal ridges continued into the ridges upon the suborbitals; ridge extending backward from tip of snout upon top of head lost in the interorbital space; branches of the horizontal ridges continued upon upper margins of orbit and there disappearing. Nostrils immediately in front of orbit, the posterior pair much the longer. Mouth situated entirely on lower side of head; symphysis of lower jaw in vertical from anterior margin of orbit, and articulations of mandibles in vertical from posterior margin of orbit; width of cleft of mouth equal to distance between symphysis of maxillaries and line connecting their articulations; upper jaw protractile vertically. Teeth conical, somewhat recurved, of nearly uniform size, arranged in villiform bands; palate smooth. Distance of first dorsal from snout about 4 times the length of its base, and from anterior margin of

orbit equal to length of head; first spine very short, not much longer than the teeth of the second spine; second spine in length twice horizontal diameter of orbit, stout, its anterior margin armed from base to tip with 15 teeth pointing upward, the uppermost slender; its length to tip of filament almost equal to distance from origin of second dorsal, this tip when laid back reaching almost to second dorsal; rays decreasing regularly in length so that, when the fin is upright, its shape approximates that of a right-angled triangle, the hypotenuse of which is the second dorsal spine, and its perpendicular side a line touching the tips of the rays; length of base of second dorsal less than that of the anal, its origin over the thirteenth scale of lateral line. Length of longest ray less than length of barbel; all rays very feeble; membrane scarcely perceptible; distance of anal from snout  $3\frac{1}{2}$  times in its length at base, its origin under eighteenth scale of lateral line; length of first ray  $\frac{1}{2}$  the length of tenth, and 3 times the length of last ray, the length of rays increasing to a point beneath anterior part of first dorsal, and thence gradually decreasing to tip of tail; distance of pectoral from snout 4 times width of interorbital area, its length twice length of mandible; insertion above the middle of depth of body, on a level with center of orbit, its third ray longest, its tip reaching to vertical from base of fourth ray; insertion of ventral behind pectoral and almost under that of first dorsal, its distance from snout slightly exceeding twice its length; tip of ventral filament reaching base of third anal ray. Ground color, light brownish gray; under parts silvery; belly darker, bluish; under surface of snout pink, as is also the first dorsal, except spines; spines of dorsal, ventral, and anterior anal rays blackish; throat, branchiostegal membrane, and isthmus rich copper violet; sclerotic coat green; eyes very dark blue. This species was first deep-sea fish obtained by the Fish Commission or described by an American ichthyologist. It ranges in depths from 9 to 1,255 fathoms. This species is distinguished by Günther from his *Macrourus equalis*, which it closely resembles, (1) by its longer snout, which is nearly equal to the diameter of the eye, and (2) by the smaller number of ventral rays (7). (Goode & Bean.) West Indies to Massachusetts Bay, usually in great depths; excessively abundant on the continental slope, with *Phycis chesteri*, far outnumbering all other deep-sea fishes in the region. (Named for Spencer Fullerton Baird.)

*Macrourus bairdii*, GOODE & BEAN, Amer. Journ. Sci. and Arts 1877, 471, Massachusetts Bay; GOODE, Proc. U. S. Nat. Mus. 1880, 337, 475; GÜNTHER, Challenger Report, xxii, 135, pl. 22, fig. B, 1887; GOODE & BEAN, Oceanic Ichthyology, 393, fig. 335, 1896.

#### 2960. MACROURUS LEPTURUS, Gill & Townsend.

Head  $5\frac{1}{2}$ ; depth 8; eye  $4\frac{1}{3}$  in head; snout 4; maxillary  $2\frac{1}{3}$ . D. XIV-122; A. 116; P. 20; V. 8. Scales deciduous and moderate, oblong or oval with reduced exposed surfaces, those on the back or above the lateral line with a few, 3 to 5, ridges beset with spines, but those below mostly unarmed; head regularly conical; snout moderately extended; median tubercle very projecting, the lateral well developed, connected with the median by a well-defined ridge; infraorbital vertical, with the ridge linear and near

the orbit; teeth cardiform in both jaws; the lower teeth beset the outer slope of the jaw. Ventral as long as head; pectoral  $1\frac{1}{2}$  in head; dorsal spine serrate,  $1\frac{1}{2}$  in head. Apparently close to *M. acrolepis*, but probably with shorter head, longer ventrals, and longer dorsal spine, the eye also larger. Length 22 to 26 inches. Bering Sea. Only 2 specimens known. (*λεπρός*, slender; *οὐρά*, tail.)

*Macrurus lepturus*, GILL & TOWNSEND, Proc. Biol. Soc. Wash., XI, 1897 (Sept. 17, 1897), 233, Bering Sea, southwest of Pribilof Islands, Albatross Station 3604, in 1,401 fathoms. (Type, No. 48767, U. S. Nat. Mus. Coll. Albatross.)

*Macrurus dorsalis*,\* GILL & TOWNSEND, Proc. Biol. Soc. Wash., XI, 1897 (Sept. 17, 1897), 233, Bering Sea, southwest of Pribilof Islands, Albatross Station 3604, in 1,401 fathoms. (Type, No. 48768, U. S. Nat. Mus. Coll. Albatross.)

#### 2961. MACROURUS ACROLEPIS, Bean.

Head about  $4\frac{1}{2}$ ; depth at ventrals 7; eye  $3\frac{1}{2}$  in head; snout 4; maxillary  $2\frac{1}{2}$ ; mandible  $2\frac{1}{2}$ ; pectoral about 2; ventral about  $1\frac{1}{2}$ . D. XI-111+; A. 94+; P. 20. Form of *M. bergylax*, width of head  $\frac{2}{3}$  its height; interorbital width  $\frac{1}{2}$  eye; snout moderate, pointed. Origin of first dorsal from snout a distance 3 times length of upper jaw; base of first dorsal  $3\frac{1}{2}$  in head, or about 3 times distance between dorsals; first dorsal spine very short, the second about  $1\frac{1}{2}$  in head, serrate in front. Distance of anal from snout  $2\frac{1}{2}$  times its length; distance of pectoral from snout slightly greater than head; distance of ventral origin from snout  $\frac{1}{2}$  its length. Length 2 feet or more. Coasts of Vancouver Island, Washington and Oregon, in deep water, in 345 to 786 fathoms; common. A small specimen taken by us off Bogoslof Island. Our specimens have 11 rays in the first dorsal, not 11, 11 or 13, as given by Bean. (*ἀκρος*, sharp; *λεπίς*, scale.)

*Macrurus acrolepis*, BEAN, Proc. U. S. Nat. Mus. 1883, 362, Straits of Juan de Fuca, near Neah Bay, Washington (Coll. James G. Swan, from the stomach of a fur seal); JORDAN, Cat. Fish. N. A., 131, 1885; GILBERT, Rept. U. S. Fish Comm. 1893 (1896), 457.

*Macrourus acrolepis*, JORDAN & GILBERT, Rept. Fur Seal Invest., 1898.

#### 2962. MACROURUS STELGIDOLEPIS, Gilbert.

Head  $4\frac{1}{2}$ ; depth  $5\frac{1}{2}$ ; eye small,  $3\frac{1}{2}$  to 4 in head; snout  $4\frac{1}{2}$ . D. II, 10 or 11; A. 130; scales 155; 5 or 6 scales between lateral line and base of first dorsal. Body deep, the lower profile rapidly rising along anterior portion of base of anal, the tail thus abruptly becoming slender. Head short and deep; snout heavy, little produced, acute at extreme tip; infraorbital ridge not prominent on sides of head or snout, not continued backward on preopercle. A pair of narrow, transverse naked strips on upper surface of snout near tip, separated on each side by a single scale from the naked

\* The following is Gill & Townsend's description of *Macrourus dorsalis*:

"Dorsal 15-120; anal 122; pectoral 21; ventral 9. Scales deciduous and rather small, diversiform, with small exposed surfaces; near the dorsal they have about 5 radiating spinigerous ridges, but below the lateral line these ridges are fewer and unarmed; snout short, projecting a considerable length beyond the eye and a little beyond the supramaxillary; median tubercle very prominent; connecting ridge well defined; infraorbital nearly vertical, with the ridge linear and near the orbit; teeth cardiform."

To this we add the following, from our examination of their type: Head  $5\frac{1}{2}$ ; depth 7; eye  $4\frac{1}{2}$ ; snout 4; interorbital width slightly greater than eye; maxillary  $2\frac{1}{2}$ ; ventral fin  $\frac{1}{2}$  longer than head,  $4\frac{1}{2}$  in body; pectoral  $1\frac{1}{2}$  in head. Dorsal spine strongly serrate, its length equal to that of head.

nostril fossa; a double series of scales intervenes between the nostrils and the orbit; lower side of snout wholly naked anteriorly, partly scaled laterally. Mouth large, overhung by premaxillaries for a distance about  $\frac{1}{2}$  diameter of orbit; premaxillaries in advance of nostrils; maxillaries reaching vertical from posterior margin of pupil,  $2\frac{1}{2}$  in head; snout about equaling interorbital width; barbel long,  $\frac{1}{2}$  orbit. Teeth in cardiform bands of equal width in both jaws, narrowed laterally, but not to a single series; anterior series in upper jaw enlarged, in lower jaw all the teeth of equal size. Preopercle broadly rounded, the angle but moderately produced, a narrow strip of the interopercle visible for its entire length; outer gill arch partially joined to cover, as usual; gill rakers obsolete; gill membrane united, forming a wide free fold across isthmus posteriorly. Scales without ridges, their exposed surfaces thickly beset with spines which are usually without definite arrangement; the marginal spine longest, thence decreasing in length to the base, about 40 present on each scale on middle of sides; scales on head crowded, the spines shorter and not directed backward as on the body; a rosette of short spines on tip of snout; no naked area between ventrals; mandible and gill membranes partly scaled; no considerable naked area in axil of pectorals. Dorsal inserted over base of pectorals, the length of its base slightly less than  $\frac{1}{2}$  the interspace between base of dorsals; second dorsal spine rather short and fragile, furnished anteriorly with a series of retrorse spinules, its length slightly exceeding  $\frac{1}{2}$  that of head, its tip not reaching origin of second dorsal; origin of anal fin well in advance of second dorsal; the vent unusually far forward, its distance from base of ventrals 2 to  $2\frac{1}{2}$  in its distance from anal fin; ventrals less widely separated than in *M. scaphopsis*, the outer ray produced, extending beyond front of anal; ventrals with 10 rays; pectorals with 22 to 24 rays; longest pectoral ray equals  $\frac{1}{2}$  head. Color very dark brownish, lighter on tail; lower side of head, breast, and abdominal region, including front of anal and base of pectorals, blue black; roof of mouth, valvular flap of membrane behind bands of teeth, gill membranes, and upper posterior portion of opercular lining, black; mouth and gill cavity otherwise white; peritoneum bright silvery, with little black specking; fins dusky. (Gilbert.) Coast of southern California. Two specimens, the longest 12 inches in length, from Albatross Station 2960, in 267 fathoms. (*στελγίς*, a scraper; *πίς*, scale.)

*Macrurus stelgidolepis*, GILBERT, Proc. U. S. Nat. Mus. 1890, 116, coast of southern California, at Albatross Station 2960, in 267 fathoms.

2963. MACROURUS CINEREUS, Gilbert.

(POP-EYE.)

D. II, 10 or 11; ventral 9; 7 scales between lateral line and first dorsal. Eye  $3\frac{1}{2}$  to 4 in head; snout about 4, high and blunt, but little overlapping the mouth, terminating in a pointed prolongation of the median ridge, which bears at its tip a bony tubercle furnished with radiating ridges; nasal ridges terminating in shorter and smaller, but similar, tubercles, the outline between them concave; tip of snout overpassing the premaxillaries for  $\frac{1}{2}$  its length; eye very large and protuberant; mouth of

moderate size, the maxillary reaching vertical from hinder margin of orbit, equaling length of snout and  $\frac{1}{2}$  of eye. Teeth finely villiform, in each jaw, the outer series not at all enlarged, the mandibular band narrow. Barbel short and slender, its length less than  $\frac{1}{2}$  diameter of pupil; interorbital width  $\frac{1}{2}$  diameter of orbit, equaling length of snout; preopercle greatly expanded, much overlapping the interorbital below, leaving exposed only the extreme posterior angle. Gill membranes narrowly joined, with a posterior fold, free from the isthmus; gill rakers short, compressed, almost tubercular, 2 + 12. Origin of dorsal well behind base of pectorals; second dorsal spine long and filamentous, strongly spinous except on extreme base and tip; length of spine  $\frac{2}{3}$  to  $\frac{1}{2}$  head; base of first dorsal equaling diameter of orbit; interspace between dorsals  $\frac{2}{3}$  to  $\frac{1}{2}$  base of first dorsal; pectoral long and slender, equaling length of head behind anterior nostril opening, about as long as the filamentous outer ventral ray; vent immediately in front of anal origin. Scales on sides well imbricated, each with 7 to 9 parallel ridges which bear short sharp spines directed very obliquely backward; 7 scales between lateral line and base of first dorsal. Color uniform light grayish on body and fins, with the exception of the blackish pectorals and ventrals; sides of head silvery; mouth, gill cavity, and peritoneum brownish or purplish black; gill membranes and gular membrane dusky. (Gilbert.) Bering Sea; excessively abundant in the depths, where it outnumbered all other fishes. Numerous specimens from north of Unalaska Island, at Albatross Stations 3307 and 3329, in 1,033 and 399 fathoms; and the North Pacific, south of Unimak Island, Albatross Station 3340, in 695 fathoms. Our many specimens from near Bogoslof Island, in 664 fathoms. (*cinereus*, ashy gray.)

*Macrourus cinereus*, GILBERT, Rept. U. S. Fish Comm. 1893 (1896), 457, near Unalaska and Unimak Islands, in 399 to 1,033 fathoms; JORDAN & GILBERT, Report Fur Seal Invest., 1898.

1007. CÆLORHYNCHUS, Giorna.

*Cælorhynchus*, GIORNA, Mém. Ac. Sci. Turin, XVI, 178, 1803 ("Cælorhynche la ville").

*Krohnius*, COCCO, Lettera al Sig. Augusto Krohn, Pesci del Mare de Messina, 1, 1844 (*filamentosus*; larva).

*Paramacrurus*, BLEEKER, Versl. Med. Ak. Wetensk. Amsterd. 1874, 103 (*australis*).

*Oxymacrurus*, BLEEKER, Versl. Med. Ak. Wetensk. Amsterd. 1874, 103 (*japonicus*).

This genus agrees with *Macrourus* in all essential respects, except that the small mouth is wholly below the long-pointed, sturgeon-like snout. Dorsal spine smooth in typical species, those with serrate spine having been lately separated under the generic name *Cwlocephalus*. (Gilbert & Cramer, Proc. U. S. Nat. Mus. 1896, 422) (*acipenserinus*). Species numerous. (*κοιλος*, hollow; *ρυγχος*, snout.)

a. Head large,  $3\frac{1}{2}$  in length; depth 7; eye  $\frac{1}{2}$  length of the long snout, 4 in head; dorsal spine moderate. OCCA, 2964.

aa. Head short,  $4\frac{1}{2}$  to 5 in length.

b. Body rather elongate, the depth 8 in length; eye as long as snout, 3 in head.

CARMINATUS, 2965.

bb. Body less elongate, the depth  $6\frac{1}{2}$  to  $6\frac{3}{4}$  in length.

c. Dorsal spine long; anal rays about 110; scales 124. CARIBLÆUS, 2966.

cc. Dorsal spine very short; anal rays 95; scales 98. SCAPHOPHIS, 2967.

2964. *CELORHYNCHUS OCCA* (Goode & Bean).

Head  $3\frac{1}{2}$  in total length; depth 7 in total length; snout exceedingly elongate, nearly twice as long as diameter of eye; a black flap between nostrils; angle of mouth nearly reaching vertical from posterior margin of the orbit; ridge of head very strong and continuous from snout to angle of preopercle, having, also, strong supraocular and occipital ridges; eye nearly round, its horizontal diameter 4 in head and equal to interorbital space; ventral originating under middle of first dorsal, and extending to fourth ray of anal; distance from ventral origin to vent  $3\frac{1}{2}$  in length of head; second spine of dorsal weak and smooth, its length equal to post-orbital part of head, its base slightly less than distance between first and second dorsals; squamation excessively rough, each scale bearing about 5 large spines besides many smaller ones, the median spine of the large series being much the largest; 5 rows of scales between origin of dorsal and lateral line, 19 from vent forward to lateral line and 12 backward; barbel  $\frac{1}{2}$  as long as snout. This species has scales similar to those of *Macrourus berglax*, there being a strong median keel formed by series of spines, of which the last is the largest; surface of each scale also with about 4 or more lateral ridges formed by series of short spines. In a much larger example (U. S. Nat. Mus. No. 37334), measuring 18 inches in length, the lateral series of keels have greatly increased in number, the individual spines having become more prominent, so that the median keel has become less conspicuous than in the type. In the larger specimen referred to, the nakedness of the under surface of the head is even more pronounced than in the smaller, in which the under surface of the head beneath the suborbital and nasal ridge is almost entirely naked. The intermaxillary has a very short bone similar in structure and dentition to that of *Macrourus berglax*, that is to say, the intermaxillary teeth are in a rather broad villiform band, and the outer teeth are not enlarged; mandibular teeth in a similar broad villiform band; mouth entirely inferior and small. Gill membranes attached across the isthmus, very little emarginate, and not deeply cleft; in the large example the gill membrane is attached to the isthmus and not deeply cleft, but there is a very narrow free margin behind. The gill rakers are very short, tubercular, and few in number, certainly not more numerous than in *M. berglax*; in the large example only 8 little tubercles can be seen on the first gill arch. Second spine of the dorsal in the type specimen is smooth, with the exception of 2 weak spines near its tip, but in the large example there is no trace of serrations on the dorsal spine. (Goode & Bean.) Length 450 mm. Gulf of Mexico and West Indies, in deep water. (*occa*, a harrow, from the rough scales.)

*Macrourus occa*, GOODE & BEAN, Proc. U. S. Nat. Mus. 1885, 595, Gulf of Mexico, Lat.  $28^{\circ} 34'$  N., Long.  $86^{\circ} 48'$  W., in 335 fathoms. (Type, No. 37334. Coll. *Albatross*.)

*Celorhynchus occa*, GOODE & BEAN, Oceanic Ichthyology, 400, figs. 332, 333, and 337, 1896.

2965. *CELORHYNCHUS CARMINATUS* (Goode).

Head about 5 in total length; depth 8; eye about 5 in head, equaling interorbital width; snout equaling eye or postorbital part of head; length of opercle about 2 in snout. Body less elongate than in *M. bairdii*. Snout

long, sharp, depressed, triangular, the lower surface more nearly parallel with the axis of body than in *M. bairdii*; lateral ridges more pronounced, continued in a straight line under eye and upon preopercle; strong horizontal ridges running from supraorbital margins to gill openings, parallel with subocular ridges; nostrils immediately in front of orbit; barbel very short. Teeth small, conical, somewhat recurved, arranged in villiform bands. Origin of first dorsal to snout  $4\frac{1}{2}$  times its base, its distance from anterior margin of orbit much less than length of head; first spine very short, hardly perceptible above the skin; second spine about 2 in head, slender and unarmed, when laid back its tip reaching to or beyond origin of second dorsal, the spines decreasing in length very gradually, the sixth being nearly as long as second, so that the fin is not so triangular as in *M. bairdii*; second dorsal beginning in a perpendicular from seventh anal ray; anal much higher than in *M. bairdii*, nearly equal to  $\frac{1}{2}$  interorbital width, its origin under eighteenth scale of lateral line, its longest rays as long as interorbital width; distance of pectoral from snout equaling twice its own length, which about equals longest dorsal spine; origin of pectoral below middle of depth of body and below level of middle of orbit, its tip not reaching origin of anal; insertion of ventrals behind pectoral, slightly in advance of first dorsal, its distance from snout greater than twice its length, the long filament not reaching anal. Color silvery gray. Length 250 mm. This species is extremely close to the common Mediterranean species, *C. caelorhynchus* (Risso), but the spines on the scales are a little larger. West Indies, Gulf of Mexico, and in the Gulf Stream in deep water; abundant; taken at many stations by the *Albatross*, the *Blake*, the *Fish Hawk*, and the *Challenger*, in 115 to 464 fathoms. (*carminatus*, from carmen, a wool card.)

*Macurus carminatus*, GOODE, Proc. U. S. Nat. Mus., III, 1880, 346 and 475, Gulf Stream off Rhode Island, Lat.  $40^{\circ} 02' 54''$  N., Long.  $70^{\circ} 23' 40''$ , at Fish Hawk Station 871, in 115 fathoms (Type, No. 26007); GOODE & BEAN, Bull. Mus. Comp. Zool., x, No. 5, 196, 1883. *Macurus (Caelorhynchus) carminatus*, GÜNTHER, Challenger Report, Deep-Sea Fishes, XXII, 129, pl. 5, fig. 13, 1887.

*Caelorhynchus carminatus*, GOODE & BEAN, Oceanic Ichthyology, 398, fig. 336, 1896.

2966. *CAELORHYNCHUS CARIBBEUS* (Goode & Bean).

Head  $4\frac{1}{2}$  in total length; depth  $6\frac{1}{2}$  in total length. 1 D. II, 8; 2 D. at least 110; A. 110+; scales 6-124-15 or 16. Body normal in shape; scales moderate, strong, densely covered with minute spines, without enlarged median keel; interorbital area flat, its greatest width about 5 times in length of head; postorbital portion of head about 3 in head, and just as long as eye, which is oval, and  $1\frac{2}{3}$  as long as its vertical diameter. Snout long, thin, diaphanous, with acuminate point, its general form resembling that of *C. carminatus*. The nostrils close to the orbit, the posterior one much the larger. Teeth in each jaw in villiform bands, minute. Barbel slender and short, its length  $\frac{1}{2}$  that of eye. Maxillary extending to vertical through middle of pupil; upper jaw about 3 in head; mandible  $2\frac{1}{2}$ ; intermaxillary short. Outer series of teeth on intermaxillary and mandible not enlarged, the teeth not becoming uniserial. Gill membranes narrowly attached to the isthmus; gill rakers minute, tubercular, about 10 on



first arch. Suborbital ridge very strong, continued almost in a straight line by the lateral ridge of the snout; under surface of head, except chin and branchiostegal region, densely covered with small, spiny tubercles; a naked space on underside of snout, occupying almost entire distance from front of mouth to tip of snout, widest anteriorly, the greatest width 5 in snout; intermaxillary protractile in a vertical direction; mouth distinctly inferior. Origin of second dorsal over seventh anal ray, about an eye's diameter behind first dorsal; length of anal rays about 4 in head; origin of pectoral in front of first dorsal, its length 2 in head, its tip reaching fifth anal ray. Color silvery gray, with yellowish and lavender tints. Length 290 mm. Caribbean Sea north to the Gulf of Mexico, in deep water. (*caribbaeus*, of the Caribbean Sea.)

*Macurus caribbaeus*, GOODE & BEAN, Proc. U. S. Nat. Mus. 1885, 594, Gulf of Mexico, at Albatross Station 2377, Lat. 29° 07' 30" N., Long. 83° 08' W., in 210 fathoms (Type, No. 37333); GÜNTHER, Challenger Report, Deep-Sea Fishes, XXII, 124, note 3, 1887.  
*Ceolorhynchus caribbaeus*, GOODE & BEAN, Oceanic Ichthyology, 401, fig. 338, 1896.

2967. *CEOLORHYNCHUS SCAPHOPSIS* (Gilbert).

Depth  $6\frac{1}{2}$  in total length; head  $4\frac{1}{2}$ ; scales 98. D. II, 8; A. ca. 95. Snout flattened, acute, the conspicuous infraorbital ridge forming a strong ridge along its sides, the two meeting at tip in a salient point; an evident keel extending from tip of snout to middle of interorbital area; supraorbital ridge dividing anteriorly, 1 branch running down in front of nostril, the other separating nostril fossa from orbit; between the ridges the head is covered with a soft, yielding integument, which is semitranslucent. Lower side of snout wholly naked below, and with a large naked area above on each side of tip; snout projecting beyond mouth for a distance equaling length of maxillary. Mouth of moderate size, the maxillary reaching vertical from posterior margin of pupil,  $3\frac{1}{2}$  in head. Teeth villiform, in a broad band in upper jaw, in a narrower band below, not reduced to a single series laterally in either jaw, and none of the teeth enlarged. Eye large, elliptical, equaling length of snout,  $3\frac{1}{2}$  in head; interorbital width  $4\frac{1}{2}$ . Barbel short, about  $\frac{1}{2}$  pupil; preopercular angle greatly produced backward, wholly concealing the interopercle, the strong infraorbital ridge failing to reach preopercular margin by only  $\frac{1}{2}$  diameter of pupil. Structure of gills as usual in this genus, the gill rakers obsolete; gill membranes broadly united, joined to isthmus, across which they form posteriorly a very narrow free fold. Besides the ridges already described on head, there are a pair on occiput, a pair from upper posterior margin of orbit to upper angle of gill opening, and a median ridge on nape reaching about halfway from occiput to dorsal. These ridges, as well as the interorbital space and the area between the occipital ridges, covered with scales compressed to a knife-like edge, which is provided with a single series of backward-directed spines; scales on infraorbital and rostral ridges bearing stellate spines or are similar to those on temporal region, sides of head, and body generally; scales on body large; 3 longitudinal series between lateral line and middle of first dorsal; each scale provided with a ridge bearing about 6 backward-directed spines, and from 2 to 4

pairs of lateral ridges also bearing spines, the lateral ridges sometimes extending the whole width of scales, sometimes confined to their basal portion; marginal spines longest; axil of pectoral naked, its base anteriorly with small cycloid scales; a naked, much depressed, elliptical area between bases of ventrals in all specimens; second dorsal spine smooth, weak, little exceeding length of soft rays, equaling length of snout and orbit; base of first dorsal  $1\frac{1}{2}$  in interspace between dorsals, which is  $2\frac{2}{3}$  in head; distance from front of anal to snout equaling  $\frac{1}{3}$  total length; ventrals with the outer ray produced, about reaching front of anal; pectorals reaching beyond anal  $\frac{1}{2}$  length of head; ventrals with 7 rays; pectorals with 15 to 17 rays. Color light olive brown, dusted with coarse black specks; axil of pectorals, belly, ventrals, and branchiostegal membranes blue black; lower side of head dusky; mouth anteriorly, including tongue and  $\frac{1}{2}$  of palate, white, its posterior part and most of lining of gill cavity jet-black; inner lining of cheeks abruptly white; lower part of iris silvery; peritoneum silvery, with coarse dusky specks; vertical fins dusky, the anterior portion of anal black. Coast of southern California. Many specimens, the longest 12 inches long, from *Albatross Station* 3015, in 145 fathoms. (Gilbert.) (*σκάφη*, spade; *ἄψις*, face.)

*Macrurus (Calorhynchus) scaphopsis*, GILBERT, Proc. U. S. Nat. Mus. 1890, 115. *Albatross Station* 3015, coast of southern California, in 145 fathoms.

#### 1008. TRACHONURUS, Günther.

*Trachonurus*, GÜNTHER, Challenger Report, Deep-Sea Fishes, xxii, 124, 1887 (*villosus*).

Scales not imbricated, separated by furrows, and densely covered with sharp spinules, so that the animal seems villous to the touch; dorsal spine smooth; dorsal much lower than anal; teeth in both jaws in villiform bands; snout obtuse, the mouth subinferior; suborbital ridge little developed. This genus is distinguished from *Calorhynchus* by the indistinct squamation. (*τραχὺς*, rough; *ὄψα*, tail.)

#### 2968. TRACHONURUS SULCATUS (Goode & Bean).

Head  $7\frac{1}{2}$  in total length; depth about  $9\frac{1}{2}$ ; eye  $3\frac{3}{8}$  in head; snout 4 to  $4\frac{1}{2}$ . D. II, 8 or 9, the second of numerous low rays; A. 120; V. 7; P. 13; scales 7-175 or more—33. Barbel  $2\frac{1}{2}$  to 2 in eye. Body elongate, rapidly contracted behind the abdomen; the tail long and whip-like. Scales moderate, strongly armed, each with 8 to 10 spinelets, irregularly placed, less numerous in the young, which feel bristly to the touch, separated by wide deep furrows; armature of head similar to that of body, but the scales upon snout, cheeks, and chin have very feeble spines. Interorbital area nearly flat, its length equaling diameter of eye or about 3 in head; postorbital part of head as long as eye; snout short, obtuse, scarcely overhanging the mouth; nostrils somewhat above level of middle of eye, the anterior one nearly upon the dorsal outline. Upper jaw with 2 series of teeth in villiform bands, the outer series slightly enlarged; teeth of lower jaw in a single series; maxillary reaching to vertical through hind margin of pupil in adult, shorter in younger individuals; length of upper jaw,

including maxillary, 3 in head; mandible 2 in depth of body; barbel 2 to 2½ in eye. Gill rakers very small, tubular, almost rudimentary, about 10 below angle of first arch; attachment of membrane to first arch very extensive, but free from isthmus; no pseudobranchiae. First dorsal comparatively low, the first spine rudimentary, the second elongate and smooth; insertion of first dorsal immediately over or somewhat behind base of pectoral, its distance from snout 1½ in head, its base equal to snout, its longest spine, when laid down, reaching behind origin of second dorsal, or 1½ to 2 in head; second dorsal very low, its distance from first 3 to 4 in head, 32 rays in a distance equal to length of head; 22 in same distance of anal; anal much higher than second dorsal, yet very low, its longest ray equal to eye; distance of anal origin from snout 4½ in total length, or nearly under origin of second dorsal; pectoral inserted under or somewhat in front of origin of first dorsal, its length about 2 in head; ventral inserted behind vertical from end of base of first dorsal, extending to origin of anal, its length about equaling eye; vent about midway between origin of ventrals and anal. Color brown; abdomen and lower parts of head blackish in the young. West Indies and Gulf of Mexico, in deep water; taken both by the *Albatross* and the *Blake*. (*sulcatus*, furrowed.)

*Coryphænoïdes sulcatus*, GOODE & BEAN, Proc. U. S. Nat. Mus. 1885, 596, Gulf of Mexico, at Albatross Station 2394, Lat. 28° 38' 30" N., Long. 87° 02' W., in 420 fathoms (Type, No. 37335); GOODE & BEAN, Oceanic Ichthyology, 403, 1896.

*Macrurus (Malacocephalus) sulcatus*, GÜNTHER, Challenger Report, Deep-Sea Fishes, XXII, 169, 1887.

*Trachonurus sulcatus*, GOODE & BEAN, Oceanic Ichthyology, 410, fig. 343, 1896.

#### 1009. LIONURUS, Günther.

*Lionurus*, GÜNTHER, Challenger Report, XXII, Deep-Sea Fishes, 124, 1887 (*filicauda*).

This genus is close to *Macrurus*, differing in the smooth, flaccid scales, and soft, cavernous skull, characters associated with its extreme bathybial degradation. (*λεῖος*, smooth; *ὄψα*, tail.)

a. Barbel minute, not ¼ pupil; eye small, 5 in head; tail very slender.

FILICAUDA, 2969.

aa. Barbel moderate, 1½ in eye; eye 3½ in head.

LIOLEPIS, 2970.

#### 2969. LIONURUS FILICAUDA (Günther).

D. 11; P. 20; V. 9; caeca 7. Snout considerably projecting beyond the mouth, pointed in the middle, twice as long as eye, which is unusually small, only ½ as wide as interorbital space. Mouth rather wide, extending beyond the center of the eye. Upper teeth villiform, in a very narrow band, those of mandible very small, biserial. Barbel minute. Preoperculum with the angle produced backward, broadly rounded and crenulated on the margin. Terminal portion of the tail prolonged into a long filament, more slender than in any of the other species. Bones of head soft. Scales of moderate size, thin, cycloid, and deciduous, 6 or 7 in a transverse series between the first dorsal spine and the lateral line; snout

and inferior half of the infraorbital region naked. Second dorsal spine slender, with the barbs in front very inconspicuous and sometimes entirely absent; distance between dorsal fins less than length of head; outer ventral ray produced into a small filament. Distance between vent and isthmus less than length of head. Head and trunk whitish, tail brownish, lower part of head and gill openings black. (Günther.) This species is clearly one of those in this family which extends to the greatest depths. The decrease in the size of the eye, the very soft bones, the concomitant want of firmness in the structure of the scales, and the tail, which tapers into a very fine filament, indicate its abyssal abode. The scales are nearly all gone in all the specimens obtained. The species appears to be abundant in individuals, and has, like a true deep-sea fish, a wide distribution. (Günther.) Antarctic Ocean and deep seas off both coasts of South America. (*filum*, thread; *cauda*, tail.)

*Coryphaenoides (Lionurus) flicauda*, GÜNTHER, Ann. and Mag. Nat. Hist., xx, 1878, 27, Deep seas on both sides of South America, in 1,375 to 2,650 fathoms.  
*Macrurus flicauda*, GÜNTHER, Challenger Report, Deep-Sea Fishes, 141, pl. 34, fig. B, 1887.  
*Lionurus flicauda*, GOODE & BEAN, Oceanic Ichthyology, 409, fig. 342, 1896.

## 2970. LIONURUS LIOLEPIS, Gilbert.

Head  $4\frac{3}{4}$  in length; depth  $6\frac{1}{2}$ ; maxillary nearly reaching vertical from posterior margin of orbit,  $2\frac{1}{2}$  in head; eye  $3\frac{3}{4}$ ; interorbital space concave, equaling snout,  $4\frac{1}{2}$ ; barbel  $\frac{2}{3}$  eye. D. II, 10. A. 120. Snout short and high, with well-marked lateral ridge, the extreme tip flattened; the median ethmoidal ridge is prominent, and the supraocular ridge is continued forward on the snout, meeting the lateral ridge in a projecting point. Top of snout wholly naked mesially, a narrow band of scales around each margin and in front. Lower side of head, including under side of snout, mandibles, gill membranes, and most of interopercles, naked; a very small patch of scales on posterior part of interopercle. Mouth moderate, the snout overhanging the premaxillaries for a distance equaling  $\frac{1}{2}$  maxillary. Teeth in rather narrow cardiform bands in each jaw, not, however, forming single series laterally; the outer series in upper jaw only is enlarged; angle of preopercle little produced, not concealing the interopercle; infraorbital ridge not continued on to it. Gill membranes forming posteriorly a rather wide free fold across isthmus; outer gill arch joined to gill cover as usual in this genus. Scales small, everywhere cycloid, very deciduous, lost in most specimens; no spines developed, but occasionally can be seen traces of a median ridge and a pair of lateral ridges; about 6 or 7 series of scales between lateral line and base of first dorsal. Origin of first dorsal over or in advance of base of pectorals; base of first dorsal  $1\frac{1}{2}$  to  $1\frac{3}{4}$  in interval between dorsals; second dorsal spine usually smooth, occasionally with from 1 to 3 weak prickles near the middle; length of spine  $1\frac{1}{2}$  in head. Origin of anal slightly behind first dorsal, the vent midway between base of ventrals and anal; ventrals short, the outer ray slightly produced, with from 10 to 12 rays; pectorals with 20 or 21 rays. Color very dark brown; snout, opercles, lower side of head, and abdominal region black or blue black; mouth and gill cavity

black; peritoneum dusky silvery. Coast of southern California. Many specimens taken at *Albatross Station* 2980, in 603 fathoms. (*λελος*, smooth; *λεπίς*, scale.) (Gilbert.)

*Macrurus* (*Lionurus*) *liolepis*, GILBERT, Proc. U. S. Nat. Mus. 1890, 117, coast of southern California, at *Albatross Station* 2980, in 603 fathoms.

Remotely related to the Scombriform fishes, and perhaps derived from the same ancestral stock as the *Trichiuridae*, is the singular

### Suborder TÆNIOSOMI.

#### (THE RIBBON-FISHES.)

This group is thus defined by Dr. Gill:

"Scapular arch subnormal, post-temporal undivided and closely applied to the back of the cranium, between the epiotic and pterotic, or upon the parietal; hypercoracoid perforate at or near the margin; cranium with the epiotics enlarged, encroaching backward and juxtaposed behind, intervening between the exoccipitals and supraoccipital; prootic and opisthotic represented chiefly by the enlarged prootic; suborbital chain imperfect; the ocular bones separated by intervening cartilaginous elements; the hypopharyngeals styliform and parallel with the branchial arches; epipharyngeals in full number (4 pairs), and mostly compressed; the dorsal fin composed of inarticulate rays or spines, separable into lateral halves, and the ventrals (when present) subbrachial. A myodome may be present or absent, none being developed in the *Regalecidae*, but 1 being distinct and supplemented by a dichost in the *Trachyteridae*." (Gill.)

"The ribbon-fishes," says Günther, "are true deep-sea fishes, met with in all parts of the oceans, generally found when floating dead on the surface or thrown ashore by the waves. Their body is like a band, specimens of from 15 to 20 feet long being from 10 to 12 inches deep and about an inch or two broad at their thickest part. The eye is large and lateral; the mouth small, armed with very feeble teeth; the head deep and short. A high dorsal fin runs along the whole length of the back, and is supported by extremely numerous rays, its foremost portion, on the head, is detached from the rest of the fin, and composed of very elongate flexible spines. The anal fin is absent. The caudal fin (if preserved, which is rarely the case in adults) has an extra-axial position, being directed upward like a fan. The ventrals are thoracic, either compressed of several rays or reduced to a single long filament. The coloration is generally silvery, with rosy fins. When these fishes reach the surface of the water the expansion of the gases within their bodies has so loosened all the parts of their muscular and bony system that they can be lifted out of the water with difficulty only, and nearly always portions of the body and fins are broken and lost. The bones contain very little bony matter, and are very porous, thin, and light. At what depth ribbon fishes live is not known; probably the depths vary for different species; but although none has yet been obtained by means of the deep-sea dredge, they must be abundant at the bottom of all oceans, as dead fishes or fragments of them are frequently obtained.

Some writers have supposed from the great length and narrow shape of these fishes that they have been mistaken for 'sea serpents,' but as these monsters of the sea are always represented by those who have had the good fortune of meeting with them as remarkably active, it is not likely that harmless ribbon-fishes, which are either dying or dead, have been the objects described as 'sea serpents.'" (*ταβία*, ribbon; *σῶμα*, body.)

FAMILIES OF TÆNIOSOMI.

a. Ventral fins reduced each to a single long filament, thickened at the tip; anterior rays of dorsal produced; mouth small; caudal fin short or wanting.

REGALECIDÆ, CCXVI.

aa. Ventral fins normally developed or else wanting.

b. Caudal fin short, fan-shaped, inserted at an angle with axis of body; the tail not much produced beyond it.

TRACHYPTERIDÆ, CCXVII.

bb. Caudal fin short, the tail beyond it ending in a long filament, longer than rest of body.

STYLEPORIDÆ, CCXVIII.

Family CCXVI. REGALECIDÆ.

(OAR-FISHES.)

Body very elongated and compressed, the head oblong, the opercular apparatus well developed (the operculum extended backward, the suboperculum obliquely behind it, and the interoperculum extended upward below the 2), the preorbital chain oblique and widest at the second bone; ventrals represented by single elongate rays, the cranium with the myodome atrophied and the dichost suppressed, the supraoccipital pushed forward by the extensive development of the epiotics which encroach forward on the roof as well as back and sides of the cranium, and with short ribs. (Gill.) Superficial characters are the very long dorsal, extending the whole length of the back and with the rays at the nape much produced; pectorals very short; caudal fin short or wanting; anal very low; head small; mouth very short; no air bladder; pyloric caeca numerous. One genus, with 2 or more species. Very large, surface-swimming fishes of the open seas; the great size, undulating motion and projecting mane causing them frequently to be taken for sea serpents. (*Regalecida*, Gill, Standard Nat. Hist., III, 1885; GILL, Amer. Nat. 1890, 482.)

1010. REGALECUS, Brinnich.

(OAR-FISHES.)

*Regalecus*, BRINNICH, Nya Sammling, III, 414, 1788 (*glesne*).  
*Gymnetrus*, BLOCH & SCHNEIDER, Syst. Ichth., 487, 1801 (*remipes*).  
*Xypterus*, RAFINESQUE, Indice, 59, 1810 (*imperati*).

Characters of the genus included above. "It is not certain that there is more than 1 species of *Regalecus*, although, as the synonymy which follows clearly shows, various names have been suggested in connection with the comparatively few individuals which, during the past century and a half, have been captured in the North Atlantic. There appears to be consider-

able possibility of individual variation in proportions of height to length, and in the number of rays in the dorsal fin, but it is a fact well known to ichthyologists that constancy is not to be expected in forms in which the number of vertebrae and fin rays has been extended far beyond the normal average. It should also be said that most of the individuals studied have been in very imperfect condition, and also that in many instances the observations have been made by untrained observers, so that it seems doubtful whether there is really more than 1 species to be assigned to the Atlantic fauna. At all events, Günther, Collett, Lütken, and Dry agree in the idea that it is impossible to discriminate between the forms already described, and we follow their lead in considering them all, for the present, as a single species. It is not impossible, of course, that should better material be obtained, it may be desirable to separate the group into more subspecies, but until this shall be done discrimination leads to confusion rather than to definite knowledge. The fishes belonging to the genus *Regalecus* are very remarkable, not only on account of their peculiar appearance and structure, but because of their enormous size. They have been known to attain the length of 20 feet, and it is more than probable that they grow very much longer, and that many of the creatures popularly identified with the "sea serpent" are only large individuals of this type. Indeed, it seems quite safe to assign to this group all the so-called "sea serpents" which have been described as swimming rapidly near the surface, with a horse-like head raised above the water, surmounted by a mane-like crest of red or brown. The individual which came ashore at Hungry Bay, in Bermuda, in 1860, and which was about 17 feet long, was described by the people who saw it before its capture as being very much larger, and as having a head of an immense horse with a flaming red mane." (Goode & Bean.) (*rex*, king; *halec*, herring. The species have long been known as "king of the herrings," as have those of *Trachipterus*.)

2971. REGALECUS GIESNE (Ascanius).

(OAR-FISH; SEA SERPENT.)

Head 16 to 20; depth 12 to 24; eye 4 to 6 in head; snout short, truncated. D. 275 to 400; P. 11 to 14; V. I. Body very elongate. Cleft of mouth vertical, the upper jaw very protractile; jaws minute or absent. Anterior 8 to 15 rays of dorsal forming an elevated crest, sometimes in 2 parts, the posterior rays with membranaceous tips; each ventral ray with a lobate membranaceous tip; skin with numerous bony tubercles; lateral line placed low. Color silvery gray, with a few spots or streaks of darker hue, most numerous anteriorly. Günther (Challenger Report, xxii, 73 to 76) has in the most painstaking manner brought together a list of the specimens taken in the North Atlantic, as far as they are known to science. He mentions 14 known upon the Scandinavian coasts from 1740 to 1852; 19 on the British coasts from 1759 to 1884; 1 in the Mediterranean (he states, however, that about  $\frac{1}{2}$  a dozen specimens have been observed in the Mediterranean); 1 in the Bermudas; 3 at the Cape of Good Hope; 1 in the Indian Ocean, and 5 off the coast of New Zealand. He calls attention to

the fact that of those observed on the British and Scandinavian coasts 4 were observed in the month of January, 5 in February, 8 in March, 2 in April, 1 in May, 1 in June, 1 in July, 2 in August, 1 in September, and 1 in October. He also calls attention to the fact that by far the greater proportion of their capture, in the Northern Hemisphere at least, is in the stormy season. This agrees with what we know of the capture of *Trachypterus*, which likewise seems to be brought to the surface only by great commotions of the ocean. The popular name of *Regalecus* is oarfish, in allusion to the blade-like expansion of the extremities of the 2 ventral fins. *Regalecus* is also called in the books the "king of the herrings." Strangely enough, no representative of this genus has been found on the coast of North America. Günther is of the opinion that the distribution of this fish in the depths of the sea is the same as that of *Trachypterus*. The similarity in their geographical distribution is quite remarkable. (Goode & Bean.) (Eu.) (*glesne*, from "Glesnes," a farm at Glesvær, near Bergen, where the type of the species was taken.)

- Spada marina*, IMPERATO, Hist. Nat., 679, 687, 1599, Naples.  
*Regalecus glesne*, ASCANIUS, Icones Rerum Nat., II, pl. 11, about 1788, Glesvær, Norway.  
*Ophidium glesne*, ASCANIUS, Nya Saml. Vid. Selsk. Skrift., III, 419, 1788.  
*Regalecus remipes*, BRÜNNICH, Nya Saml. Vid. Selsk. Skrift., III, 1788, 414, taf. B., figs. 4, 5;  
 WALBAUM, Artodi Piscium, III, 647, tab. 3, fig. 4, 1792.  
*Cepola gladius*, WALBAUM, Artodi Piscium, III, 617, 1792.  
*Gymnetrus hawkenii*, BLOCH, Ichthyol., XII, 88, 425, 1792.  
*Gymnetrus grillii*, LINDROTH, Vet. Akad. Handl. 1798, 291, pl. 8.  
*Gymnetrus ascanii*, SHAW, Gen. Zool., IV, 197, 1803; after Ascanius.  
*Xypterus imperati*, RAFINESQUE, Indice, 59, 1810; after Ferrante Imperato.  
*Gymnetrus longiradiatus*, RISSO, Ehr. Mérid., III, 296, 1826, Nice.  
*Gymnetrus telium*, CUVIER & VALENCIENNES, Hist. Nat. Poiss., X, 361, pl. 299, 1834, Nice.  
*Regalecus banksii*, CUVIER & VALENCIENNES, Hist. Nat. Poiss., X, 365, 1834, Filey Bay, Yorkshire.  
*Gymnetrus campensis*, CUVIER & VALENCIENNES, Hist. Nat. Poiss., X, 376, 1834, Cape of Good Hope.  
*Regalecus glesne*, ASCANIUS, Icones Rerum Naturalium, 1806, pl. 11; LACÉPÈDE, Hist. Nat. Poiss., II, 214, 215, 1800; GOODE & BEAN, Oceanic Ichthyology, 480, fig. 395, 1896.  
*Gymnetrus remipes*, BLOCH & SCHNEIDER, Syst. Ichth., 482, tab. 88, 1801; YARRELL, Brit. Fishes, Ed. 2, I, 223, and Ed. 3, II, 301.  
*Gymnetrus glesne*, CUVIER & VALENCIENNES, Hist. Nat. Poiss., X, 366.  
*Gymnetrus gladius*, CUVIER & VALENCIENNES, Hist. Nat. Poiss., X, 352, pl. 298, 1835.  
*Regalecus gladius*, GÜNTHER, Cat., III, 308, 1861.

#### Family CCXVII. TRACHYPTERIDÆ.

##### (THE KING OF THE HERRINGS.)

Body moderately elongate, strongly compressed, naked, the skin smooth or prickly. Lateral line present. Head short; the mouth rather small, terminal, with feeble teeth; premaxillaries protractile; opercles unarmed; opercular apparatus abbreviated (the operculum extended downward, the suboperculum below it, and the interoperculum contracted backward and bounded behind by the operculum and suboperculum); the cranium with a myodome and dichost, the supraoccipital continued behind into a prominence; the epiotics confined to the sides and back of the cranium, and without ribs. Eye large, lateral; branchiostegals 6; gill membranes



separate, free from the isthmus; gills 4, a slit behind the fourth. Pseudo-branchial well-developed, in a pouch formed by a fold of the mucous membrane. Dorsal fin single, extending from the head to the tail, its rays all technically spinous, being neither articulated nor branched, but all very soft, flexible, and fragile; anal fin wanting; pectorals short; ventrals thornlike, the rays elongate, less than 1, 5 in number, usually atrophied in the adult; caudal fin either rudimentary or else divided into 2 parts, the upper and larger fan-shaped, directed obliquely upward from the slender tip of the tail. Bones very soft, the muscles little coherent. Pyloric coeca very numerous. Vertebrae in large number. Deep-sea fishes, often of large size, found in most warm seas. Their extreme fragility renders them rare in collections, and the species are little known. One genus; species about 12. The ribbon-fishes are well known in the Eastern Atlantic and the Mediterranean, and have even been found as far west as Madeira [and Cuba]. Some few representatives have been found on the west coast of South America, and 1 or 2 examples have been taken in New Zealand. They are generally admitted to be true deep-sea fishes, which live at very great depths, and are only found when floating dead on the surface or washed ashore by the waves. Almost nothing is known of their habits except through Nilsson's observations in the Far North. This naturalist, as well as Olafsen, appears to have had the opportunity of observing them in life. They say that they approach the shore at flood tide on sandy shelving bottoms, and are often left by the retreating waves. Nilsson's opinion is that its habits resemble those of the flat fishes, and that they move with one side turned obliquely upward, the other toward the ground; and he says that they have been seen on the bottom in 2 or 3 fathoms of water, where the fisherman hook them up with the implements employed to raise dead seals, and that they are slow swimmers. This is not necessarily the case, however, for the removal of pressure and the rough treatment by which they were probably washed upon the shore would be demoralizing, to say the least. *Trichiurus*, a fish similar in form, is a very strong, swift swimmer, and so is *Regalecus*. Whether or not the habits of *Trachypterus arcticus*, on which these observations were made, are a safe guide in regard to the other forms is a matter of some doubt, but it is certain that they live far from the surface except near the Arctic Circle, and that they only come ashore accidentally. They have never been taken by the deep-sea dredge or trawl-net, and, indeed, perfect specimens are very rare, the bodies being very soft and brittle, the bones and fin rays exceedingly fragile. A considerable number of species have been described, but in most instances each was based upon 1 or 2 specimens. It is probable that future studies may be as fruitful as that of Emery, who, by means of a series of 23 specimens, succeeded in uniting at least 3 of the Mediterranean species, which for half a century or more had been regarded as distinct. The common species of the Eastern Atlantic, *Trachypterus atlanticus*, is not rare, 1 or more specimens, according to Günther, being secured along the coast of northern Europe after almost every severe gale. We desire to quote the recommendation of Dr. Günther, and to strongly urge upon any one who may be so fortun-

nate as to secure 1 of these fishes, that no attempt should be made to keep it entire, but that it should be cut into short lengths and preserved in the strongest spirits, each piece wrapped separately in muslin. (Goode & Bean.)

1011. TRACHYPTERUS, GOUAN.

(KING OF THE HERRINGS.)

*Trachypterus*, GOUAN, Hist. Poiss., 104, 153, 1770 (*trachypterus*).

*Bogmanus*, BLOCH & SCHNEIDER, Syst. Ichth., 518, 1801 (*islandicus* = *arcticus*).

Body elongate, compressed, ribbon-shaped, the dorsal fin extending the entire length of the back. Anal absent; each ventral well developed, if present, but sometimes absent. Caudal present and placed for the most part above the longitudinal axis of the body. No air bladder. Pyloric appendages numerous. Ventrals appearing to be absent in some individuals, but Day calls attention to the fact that most of the specimens of *T. arcticus* taken along the coast of Great Britain had no ventrals. In the very young, as has been shown by Emery, the fin rays commence to grow when it is about 6 mm. long, and continue to lengthen until it is about 24 mm. long, after which a partial shortening takes place. Ventrals very elongate in the young, and the caudal rays much longer than in the grown fish. Young individuals (from 2 to 4 inches) are not rarely met with near the surface; they possess the most extraordinary development of fin rays observed in the whole class of fishes, some of them being several times larger than the body, and provided with lappet-like dilatations. There is no doubt that fishes with such delicate appendages are bred and live in depths where the water is absolutely quiet, as a sojourn in the disturbed water of the surface would deprive them at once of organs which must be of some utility for their preservation. (Goode & Bean.) (*τραχύς*, rough; *πτερόν*, fin.)

a. Color bright metallic silvery, a jet-black blotch at base of dorsal; 3 dark spots on side, 2 smaller ones on belly; anterior profile, snout, and tip of mandible, jet black; caudal and ventral fins carmine red in life. REX-SALMONORUM, 2972.

aa. Color shining leaden gray; no black.

TRACHYPTERUS, 2973.

2972. TRACHYPTERUS REX-SALMONORUM, Jordan & Gilbert.

(KING OF THE SALMON.)

Head  $8\frac{1}{2}$ ; cross depth at nape 8. D. V-170; C. 8; V. 6; P. 11. Body long and slender, closely compressed and ribbon-shaped, as usual in the genus. Head short, deeper than long, the anterior profile steep and nearly straight to the base of the nuchal crest; dorsal fin beginning on the top of nuchal crest, which is directly over the second third of the diameter of eye; height of crest slightly more than diameter of eye, the latter greater than length of snout, and 3 in head. Mouth oblique; maxillary rugose and very broad, its width  $\frac{1}{2}$  its length; length of lower jaw greater than length of snout,  $2\frac{1}{2}$  in head, its angle under the front of the orbit. Opercular bones rugose, entirely covering the gills. Premaxillary covered with minute and feeble teeth, in addition to which in this specimen are 3 canines, 2 on one side and 1 on the other, directed very obliquely

backward. On the side having 2 canines, 1 is placed directly behind the other; lower jaw with 3 strong canines on one side and 2 strong and 1 weak canine on the other, all directed obliquely backward and inward. Dorsal fins slightly connected at base; the filamentous rays of the first dorsal not quite twice the length of head; ventrals inserted just below axil of pectorals, filamentous, about  $\frac{1}{2}$  longer than head; pectorals  $\frac{1}{2}$  longer than eye; caudal rays simple to near tip, where is sometimes a single fork, the longest filamentous rays about 3 times length of head; dorsal fin much lower than body, longest rays of second dorsal nearly  $\frac{3}{4}$  length of head; a series of spinules along base of dorsal, 1 pair for each ray. Lateral line well developed, with a series of small inconspicuous plates, each of which has a minute central prickle. Lower part of the body thickly beset with small spinous tubercles; rest of the skin naked; rays of all the fins accompanied by a series of small prickles. Coloration everywhere bright metallic silvery, an oblong jet-black blotch a little longer than eye lying close along base of dorsal and beginning  $1\frac{1}{2}$  diameters of eye behind eye; 3 larger spots, dusky but not black, lying behind this along side between lateral line and dorsal fin; 2 smaller dusky spots on belly, the one just behind base of ventrals, the other under the second of the 4 spots of back; these spots, except the first one mentioned, are all diffuse and a little less than twice the diameter of eye in length and about twice as long as deep; anterior profile below crest, including front of snout and tip of mandible, jet-black; caudal and ventral fins carmine red in life; other fins unmarked. Length 17 inches. This species bears some resemblance to *Trachypterus altivelis* described by Kner from Valparaiso. The latter species has, however, the nuchal crest much lower and farther back, the first dorsal and the ventrals much lower, the second dorsal fin higher, the skin rougher, the 4 black spots different in size and position from those found in our specimen, and the caudal rays divided near the base. It is probable that the 3 specimens of *Trachypterus* mentioned in the Synopsis of the Fishes of North America, p. 619, and referred with doubt to *Trachypterus altivelis*, really belong to the present species. Four specimens known; 1 from Santa Cruz, California, taken by Dr. C. L. Anderson; 2 from the Straits of Fuca, taken by Mr. J. G. Swan, and the type, obtained by a fisherman (Mr. Knox) in the open sea outside the bay of San Francisco. According to Mr. Swan the species is known by the Makah Indians west of the Straits of Fuca as "king of the salmon," and its destruction is believed to have a baneful influence on the salmon fishing. "When the king of the salmon is killed the salmon will cease to run." (*rex*, king; *salmonorum*, of the salmon.)

?*Trachypterus altivelis*, JOEDAN & GILBERT, Proc. U. S. Nat. Mus. 1381, 52; JORDAN & GILBERT, Synopsis, 618, 1883, specimen from Santa Cruz; not of KNER.

*Trachypterus rex-salmonorum*, JORDAN & GILBERT, Proc. Cal. Ac. Sci. 1894, 145, pl. 9, open sea outside Bay of San Francisco. (Type, No. 1382, L. S. Jr. Univ. Coll. Mr. Knox.)

#### 2073. TRACHYPTERUS TRACHYURUS, Poey.

D. 82; P. 15; V. 6. Eye  $2\frac{1}{2}$  in head, high, as long as snout. Mouth almost vertical. Bones of head thin as paper. Lateral line a little concave on the middle of trunk. No scales; pectorals small; ventrals behind

pectorals, very long, reaching past vent, which is at second third of length, including caudal; dorsal almost as high as body, without plume in front. Vertebrae 36 + 18. Shining leaden gray, a silvery band produced by the vertebral column showing through. Cuba. (Poey); not seen by us. (*τραχύς*, rough; *οὐρά*, tail.)

*Trachypterus trachyurus*, POEY, *Memorias*, II, 420, 1861, Cuba.

Family CCXVIII. STYLEPHORIDÆ.

Body elongate, compressed, ribbon-shaped; the dorsal extending from head nearly to end of tail; tail terminating in an exceedingly long, cord-like appendage, about twice as long as head. Anal absent; ventrals absent; caudal erected upward, having its rays connected by a rather firm membrane. Snout produced; mouth small, toothless; maxillary bones small, short, hidden behind premaxillaries; mandible long, extending far behind the eye. Eye large, turned forward; suborbital very large, covering nearly the whole of cheek and extending backward behind eye. Opercles small. Gill openings wide; gills 4. Vent premedian. Branchiostegals 4. (Goode & Bean.) This family is based on a single specimen obtained in the West Indies in 1790 and preserved in the British Museum. The relations of the fish are uncertain, and it may not belong to the *Tenisonomi*. Its nearest relations are, however, apparently with *Trachypterus*.

(*Stylephoridae*, SWAINSON, *Nat. Hist. Class'n Fishes*, II, 47, 1839.)

1012. STYLEPHORUS, Shaw.

*Stylephorus*, SHAW, *Trans. Linn. Soc. Lond.*, I, 1791, 90 (*chordatus*).

Characters of the genus included above. (*στῦλος*, a style or projecting part; *φορέω*, to bear.)

2974. STYLEPHORUS CHORDATUS, Shaw.

Head 6; depth 5. D. 110; C. 6; P. 13; B. 4. Snout produced, subcylindrical; mouth small and toothless; maxillary bones small, short, and hidden behind the intermaxillaries; mandible long, extending far behind eye; eyes large, close together, directed forward toward snout; suborbital very large, covering nearly the whole of cheek, and extending backward behind eye; opercles small; gill openings very wide; gills 4. Vents situated before middle of total length; pectorals pointed, directed upward, about  $\frac{1}{2}$  as long as head; dorsal extending from head nearly to end of tail; caudal directed upward, and having its rays connected by a rather firm membrane, the tail terminating in a narrow band-like appendage about twice as long as body. Color uniform silvery. (Günther.) This remarkable form is known only from a single specimen, 11 inches long, with the caudal appendage 22 inches in length, which was taken in the Atlantic, between Cuba and Martinique, about the year 1790, and is now in the British Museum. It is undoubtedly an inhabitant of great depths. (Goode & Bean.) (*chordatus*, with a chord; from *χόρδη*, string.)

*Stylephorus ch. datus*, SHAW, *Trans. Linn. Soc. London*, I, 1791, 90, pl. 6, between Cuba and Martinique; SHAW, *Zool.*, IV, 87; SHAW, *Naturalists' Miscellany*, VIII, pl. 274; BLAINVILLE, *Journ. Phys.*, LXXXVII, 30, pl. 1, fig. 1; CUVIER & VALENCIENNES, *Hist. Nat. Poiss.*, X, 381 GÜNTHER, *Cat.*, III, 306, 1861; GOODE & BEAN, *Oceanic Ichthyology*, 482, pl. 66, figs. 393 and 394, 1896.

## Suborder HETEROSOMATA.

## (THE FLATFISHES.)

"Cranium posteriorly normal; anteriorly with twisted vertex, to allow 2 orbits on the same side, or 1 vertical and 1 lateral; basis cranii not quite simple. Dorsal fin long, of jointed rays; superior pharyngeals 4, the third longest, much extended forward, the inferior separate." (Cope.) This suborder includes the 2 families, *Pleuronectidae* and *Soleidae*. Its nearest relationship is probably with the *Gadidae*, although the developed pseudobranchiae and the thoracic ventral fins, indicate an early differentiation from the anacanthine fishes. In the very young fishes the 2 sides of the body are alike and the eyes are 1 on each side, with normal cranium. (*ἕτερος*, different; *σῶμα*, body.) (*Anacanthini pleuronectoides*, Günther, Cat., IV, 399, 504.)

## FAMILIES OF HETEROSOMATA.

a. Preopercular margin more or less distinct, not hidden by the skin and scales of the head; eyes large, well separated; mouth moderate or large; teeth present.

## PLEURONECTIDÆ, CCXIX.

aa. Preopercular margin adnate, hidden by the skin and scales of the head; eyes small, close together; mouth very small, much twisted; teeth rudimentary or wanting.

## SOLEIDÆ, CCXX.

## Family CCXIX. PLEURONECTIDÆ.\*

## (THE FLOUNDERS.)

Body strongly compressed, oval or elliptical in outline; head unsymmetrical, the cranium twisted, both eyes being on the same side of the body, which is horizontal in life, the eyed side being uppermost and colored, the blind side lowermost and usually plain. In the very young fish the bones of the head are symmetrical, 1 eye on each side, and the body is vertical in the water. In most species the cranium becomes twisted, bringing the upper eye over with it. Eyes large, well separated. Mouth small or large, the dentition various, the teeth always present; premaxillaries protractile; no supplemental maxillary bone; pseudo-branchiae present. Gills 4, a slit behind the fourth; lower pharyngeals separate; no air bladder; preopercle with its margin usually distinct, not wholly adnate or hidden by the skin of the head; vent not far behind head, the viscera confined to the anterior part of the body. Scales various, rarely absent, usually small. Lateral line usually present, extending on the caudal fin, sometimes duplicated or wanting. Dorsal fin long, continuous, of soft rays only, beginning on the head; anal similar, shorter; caudal various, sometimes coalescent with dorsal and anal; pectorals inserted rather high, rarely wanting; ventrals under the pectorals, usually of several soft rays, one of them sometimes wanting. Fishes mostly carnivorous, inhabiting sandy bottoms in all seas, some species ascending rivers. Many of them are important food-fishes. Genera about 55; species

\* For complete synonymy and descriptions of the American species of this family of fishes, see "A review of the flounders and soles (*Pleuronectidae*) of America and Europe," by David Starr Jordan and David Kop Goss, in Report United States Fish Comm. for 1886, 225-342, pls. 1 to 9, first published in 1889.

nearly 500. The group "*Bibroniida*" recently recognized by some of the Italian ichthyologists as a separate family ("*Bibronidi*"), is composed entirely of larval forms in the early stages of their development. In this condition the eyes are symmetrical and the body translucent. Several generic names have been given to these peculiar forms (*Peloria*, *Bibronia*, *Coccolus*, *Charybdia*, *Bascanius*, *Delothyris*), but, of course, these genera can have no permanent place in the system. *Peloria* has been shown by Dr. Emery to be the young of *Platophrys*. The others seem to belong to the *Cynoglossinae* or to some allied group, but we are not yet certain as to the correct identification of any of them. We recognize among the *Pleuronectida* 6 subfamilies—*Hippoglossinae*, *Psettinae*, *Samarinae*, *Pleuronectinae*, *Oncopterinae*, and *Pelecanichthyinae*. These subfamilies are natural groups and are in most cases easily distinguished, although some few aberrant genera exist, which serve as links joining one group to another. Thus *Isopsetta* of the *Pleuronectinae* is certainly a near ally of *Psettichthys*, which is as certainly a genuine member of the *Hippoglossinae*. The *Hippoglossinae* and the *Pleuronectinae* are largely arctic in their distribution, few of the former group and none of the latter extending into the Tropics. The *Oncopterinae* seem to take the place of the *Pleuronectinae* in antarctic waters, but the species of this group are few in number. The *Psettinae* and the soles are, on the other hand, essentially warm-water fishes, their representatives in the north being comparatively few. The *Samarinae* are few in number and belong to the East Indian fauna, and the single species of *Pelecanichthyinae* belongs to the bassalian fauna of the Pacific. As the tropical *Hippoglossinae* and all the *Psettinae* are sinistral species, the eyes and color being on the left side of the body, it follows that the tropical flounders are nearly all left-sided species, while those of arctic and antarctic waters are chiefly dextral species, the eyes and color on the right. The *Hippoglossinae* are the most generalized of the flatfishes. From the northern representatives of this group, the *Solies* of *Hippoglossoides*, the *Pleuronectinae*, are certainly descended. The *Solies* are apparently derived from ancestors of the type of *Paratichthys*. The soles show closest affinities with the *Psettinae*, from ancestors of which group they have become degraded. Very remarkable is the relation between the number of vertebrae and the geographical distribution of the various species. It has been already noticed by Dr. Gill, Dr. Günther and others that in some groups of fishes northern representatives have the number of vertebrae increased. In no group is this more striking than in the flounders, as the following table, showing the numbers of the vertebrae in various species, will clearly show. The numbers inclosed in brackets are copied from Dr. Günther; the others represent our own count of specimens.

Numbers of vertebrae in flounders.

I.—HIPPOGLOSSINAE.

<i>Hippoglossus hippoglossus</i> .....	16 + 34 = 50
<i>Atheresthes stomias</i> .....	12 + 37 = 49
<i>Hippoglossoides platessoides</i> .....	13 + 32 = 45
<i>Lyopsetta exilis</i> .....	11 + 34 = 45
<i>Eopsetta jordani</i> .....	11 + 32 = 43

<i>Psettichthys melanostictus</i> .....	11 + 29 = 40
<i>Paralichthys oblongus</i> .....	11 + 30 = 41
<i>Paralichthys dentatus</i> .....	10 + 30 = 40
<i>Paralichthys lethostigmus</i> .....	10 + 27 = 37
<i>Paralichthys albignatus</i> .....	10 + 27 = 37
<i>Paralichthys californicus</i> .....	10 + 25 = 35
<i>Xystonryx liolepis</i> .....	12 + 25 = 37
<i>Ancylopsetta quadrocellata</i> .....	9 + 26 = 35

## II.—PLEURONECTINÆ.

<i>Glyptocephalus zachirus</i> .....	13 + 52 = 65
<i>Glyptocephalus cynoglossus</i> .....	[58]
<i>Microstomus pacificus</i> .....	12 + 40 = 52
<i>Microstomus kitt</i> .....	[13 + 35 = 48]
<i>Parophrys vetulus</i> .....	11 + 33 = 44
<i>Pleuronectes platessa</i> .....	[14 + 29 = 43]
<i>Isopsetta isolepis</i> .....	10 + 32 = 42
<i>Lepidopsetta bilineata</i> .....	11 + 29 = 40
<i>Limanda limanda</i> .....	[40]
<i>Liopsetta glacialis</i> .....	13 + 27 = 40
<i>Pleuronichthys decurrens</i> .....	14 + 26 = 40
<i>Pleuronichthys verticalis</i> .....	13 + 25 = 38
<i>Flesus glaber</i> .....	11 + 26 = 37
<i>Flesus flesus</i> .....	[12 + 24 = 36]
<i>Pseudopleuronectes americanus</i> .....	10 + 26 = 36
<i>Hypsopsetta guttulata</i> .....	11 + 24 = 35
<i>Platichthys stellatus</i> .....	12 + 23 = 35

## III.—PSETTINÆ.

<i>Monolene sessilicauda</i> .....	[43]
<i>Lepidorhombus whiff-iaonis</i> .....	[11 + 30 = 41]
<i>Citharichthys sordidus</i> .....	11 + 29 = 40
<i>Platophrys lunatus</i> .....	9 + 30 = 39
<i>Arnoglossus laterna</i> .....	10 + 28 = 38
<i>Arnoglossus grohmanni</i> .....	10 + 28 = 38
<i>Zenogopterus punctatus</i> .....	[12 + 25 = 37]
<i>Platophrys ocellatus</i> .....	10 + 27 = 37
<i>Lophopsetta maculata</i> .....	11 + 25 = 36
<i>Bothus rhombus</i> .....	12 + 24 = 36
<i>Syacium papillosum</i> .....	11 + 25 = 36
<i>Citharichthys arctifrons</i> .....	10 + 26 = 36
<i>Syacium micrurum</i> .....	10 + 25 = 35
<i>Phrynorhombus regius</i> .....	10 + 25 = 35
<i>Citharichthys spilopterus</i> .....	10 + 24 = 34
<i>Citharichthys macrops</i> .....	10 + 24 = 34
<i>Etropus microstomus</i> .....	10 + 24 = 34
<i>Etropus crossotus</i> .....	10 + 24 = 34
<i>Azevia panamensis</i> .....	33
<i>Psetta maxima</i> .....	12 + 19 = 31

The subdivision of the flounders into genera leaves room for considerable variety of opinion. Most of the species are well defined and easily recognized, but they do not fall readily into generic groups unless we regard almost every well-marked species as the type of a distinct genus. A natural result of an attempt at sharply defining the genera is to reach what seems an extreme degree of generic subdivision. On the other hand, attempts to unite these smaller groups to form larger ones often leave these larger ones at once unnatural and ill-defined.

It will probably appear to some that the process of generic subdivision has been in this paper carried too far. It is possible that this is true, but the arrangement which we have adopted seems to bring out the relations of the different forms better than can be done by a more conservative view of the genera. (*Pleuronectida*, Günther, Cat., IV, 1862.)

SUBFAMILIES OF PLEURONECTIDÆ.

- A. Ventral fins symmetrical, similar in position and in form of base, the ventral of the colored side not extended along the ridge of the abdomen.
- a. Mouth nearly symmetrical, the dentition nearly equally developed on both sides, the gape usually but not always wide. (Halibut tribe.) HIPPOGLOSSINÆ, I.
- aa. Mouth unsymmetrical, the jaws on the eyed side with nearly straight outline, the bones on the blind side strongly curved; teeth chiefly on the blind side.
- b. Eyes and color on the right side (with occasional exceptions). (Flounder tribe.) PLEURONECTINÆ, II.
- AA. Ventral fins unsymmetrical, dissimilar in position and usually also in form, the ventral fin of the eyed side being extended along the ridge of the abdomen. Eyes and color on the left side. (Turbot tribe.) PSETTINÆ, III.

ANALYSIS OF GENERA.

I. HIPPOGLOSSINÆ.

(HALIBUT TRIBE.)

*Large-mouthed flounders with the ventral fins symmetrical.*—Mouth symmetrical, the jaws and the dentition nearly equally developed on both sides; gape usually wide, the maxillary more than  $\frac{1}{2}$  length of head. Lower pharyngeals narrow, usually with but 1 or 2 rows of sharp teeth; teeth in jaws usually acute. Eyes large; edge of preopercle free. Pectoral and ventral fins well developed, *the ventral fins similar in position and in form of base, the ventral fin of the eyed side not being attached along the ridge of the abdomen.* Septum of gill cavity without foramen.

- a. Vertebrae and fin rays much increased in number (the vertebrae about 50; dorsal rays about 100, anal rays about 85); body comparatively elongate; caudal fin lunate; lateral line simple; anal spine mostly obsolete. Dextral species, arctic in distribution. (Genera allied to *Hippoglossus*.)
- b. Large teeth in both jaws arrow-shaped, biserial, some of them depressible; upper eye with vertical range; gill rakers short; scales deciduous, ciliated; lateral line without arch; flesh soft. Vertebrae (in *A. stomias*) 12 + 37 = 49. ATHERESTHES, 1013.
- bb. Large teeth not arrow-shaped, biserial above, uniserial below; scales very small, cycloid; gill rakers long and slender; eyes strictly lateral.
- c. Lateral line without anterior arch; lower pharyngeal teeth uniserial. REINHARDTIUS, 1014.
- cc. Lateral line with an anterior arch; lower pharyngeal teeth biserial; vertebrae (in *H. hippoglossus*) 16 + 34 = 50. HIPPOGLOSSUS, 1015.
- aa. Vertebrae and fin rays in moderate number (vertebrae less than 46; dorsal rays fewer than 95; anal rays fewer than 75); caudal fin double truncate or rounded, the median rays longest.
- d. Lateral line without distinct anterior arch; vertebrae 40 to 46; body normally dextral; caudal peduncle distinct; scales ciliated; anal spine usually strong. Species of subarctic distribution. (Genera allied to *Hippoglossoides*.)

\* Frequently sinistral in *Hippoglossoides classodon*.



e. Lateral line simple, without accessory dorsal branch; teeth sharp, those of lower jaw uniserial; dorsal beginning above eye.

f. Teeth in the upper jaw biserial.

g. Scales comparatively large, thin, and deciduous; lateral line 70; body slender, the flesh soft; vertebræ (in *L. exilis*) 11 + 34 = 45.  
LYOPSETTA, 1010.

gg. Scales small and adherent; lateral line 96; body robust, the flesh firm; vertebræ (in *E. jordani*) 11 + 32 = 43.  
EOPSETTA, 1017.

ff. Teeth in the upper jaw uniserial; scales small and flesh firm; vertebræ (in *H. platessoides*) 13 + 32 = 45. HIPPOGLOSSOIDES, 1018.

ee. Lateral line with an accessory dorsal branch; scales small, firm, ctenoid; dorsal fin beginning before the eye; teeth sharp, unequal, some of them canine-like; mouth not large; lower pharyngeal teeth sharp, uniserial; vertebræ (in *P. melanostictus*) 11 + 29 = 40. PSETTICHTHYS, 1019.

dd. Lateral line with an arch in front; no accessory branch; vertebræ in smaller number (35 to 41); anal spine usually obsolete; body normally sinistral. (Species chiefly of the temperate or subtropical seas, none of them Arctic and none European.) (Genera allied to *Paralichthys*.)

h. Dorsal fin beginning above the pupil; teeth rather small; no canines; body indifferently dextral or sinistral (in some species at least).

i. Scales ctenoid.

j. Teeth in upper jaw in 2 series; gill rakers broad.  
VERASPER, 1020.

jj. Teeth all uniserial; gill rakers slender. HIPPOGLOSSINA, 1021.

ii. Scales cycloid; teeth uniserial; gill rakers short and thick.

k. Teeth small, pointed, equal. LIPOGLOSSINA, 1022.

kk. Teeth unequal, blunt, conical; caudal fin subsessile, the caudal peduncle extremely short; skin of shoulder girdle with patches of cup-shaped scales; vertebræ (in *X. kiolepis*) 12 + 25 = 37. XYSTREURYS, 1023.

hh. Dorsal fin beginning in advance of eye; teeth sharp, uniserial or smooth.

l. Scales weakly ciliated; caudal fin with a distinct peduncle; mouth large; teeth unequal, some of the anterior canine like; gill rakers rather long and slender; no dorsal lobe nor produced ventral rays; vertebræ 35 to 41. PARALICHTHYS, 1024.

ll. Scales very strongly ctenoid on both sides of body; mouth smallish, with small, sharp teeth; anterior rays of dorsal more or less exerted, thus forming a more or less distinct lobe; gill membranes considerably united; gill rakers short and broad; caudal peduncle short; left ventral produced; vertebræ (in *A. quadrocellata*) 9 + 26 = 35.

m. Lateral line with its tubes much branched, covering parts of contiguous scales; dorsal lobe low; left ventral much produced. RAMULARIA, 1025.

mm. Lateral line with its tubes simple, not branched.

n. Body broad, ovate, the depth more than  $\frac{1}{2}$  length; dorsal lobe and left ventral moderately produced.  
ANCYLOPSETTA, 1026.

nn. Body elliptical, the depth not more than  $\frac{1}{2}$  length; dorsal lobe and left ventral greatly produced.  
NOTOSEMA, 1027.

lll. Scales entirely smooth; caudal peduncle short; mouth small; gill rakers short and thick; dorsal with an anterior lobe; left ventral elongate. GASTROPSETTA, 1028

II.—PLEURONECTINÆ.

(FLOUNDER TRIBE.)

Mouth small, unsymmetrical, the jaws on the eyed side with nearly straight outline, the bones on the blind side strongly curved; dentition chiefly developed on the blind side; eyes large; edge of preopercle not hidden by the scales; pectoral fins well developed; vertical fins well separated; ventral fins nearly or quite symmetrical, that of the eyed side not prolonged along the ridge of the abdomen; anal spine usually strong (obsolete in *Microstomus* and *Embassichthys*). Body dextral (except frequently in *Platichthys stellatus*). Species arctic or subarctic in distribution.

- a. Vertebrae in moderate number, from  $10 + 26 = 36$  to  $11 + 33 = 44$ ; dorsal rays 65 to 80; anal rays 45 to 60.
- b. Teeth small, acute, in several series; lateral line nearly straight, with an accessory dorsal branch; lower pharyngeals narrow, with small biserial teeth; scales cycloid.
- c. Lips thick, each with several longitudinal folds; dorsal fin beginning on the blind side; vertebrae 38 to 40. PLEURONICTHYS, 1029.
- cc. Lips simple; dorsal fin beginning on the median line; vertebrae (in *I. guttulata*)  $11 + 24 = 35$ . HYPSOPSETTA, 1030.
- bb. Teeth chiefly uniserial, all more or less blunt, conical or incisor-like.
- d. Lateral line with an accessory dorsal branch.
- e. Lateral line without distinct arch in front.
- f. Teeth compressed, incisor-like, close set.
- g. Scales closely imbricated, mostly cycloid; upper eye on median line; vertebrae (in *P. vetulus*)  $11 + 33 = 44$ . PAROPHRY, 1031.
- gg. Scales scarcely imbricated, all very strongly ctenoid; eyes both lateral. INOPSETTA, 1032.
- ff. Teeth conical, separated, not incisor-like; scales closely imbricated, all strongly ctenoid; mouth comparatively large (approaching that of *Psettichthys*); vertebrae (in *I. isolepis*)  $10 + 32 = 42$ . ISOPSETTA, 1033.
- ee. Lateral line with a distinct arch in front; scales imbricated, rough-ctenoid; vertebrae (in *L. bilineata*)  $11 + 29 = 40$ . LEPIDOPSETTA, 1034.
- dd. Lateral line without accessory dorsal branch.
- h. Lateral line with a distinct arch in front; scales imbricated, rough-ctenoid; vertebrae (in *L. limanda*) about 40. LIMANDA, 1035.
- hh. Lateral line without distinct arch in front.
- i. Scales regularly imbricate, all (on eyed side) ctenoid in both sexes; no stellate tubercles on head nor on bases of dorsal and anal fins; teeth, incisor-like, close set; lower pharyngeals very narrow, each with 2 rows of separate, conical teeth; fin rays scaly. PSEUDOPLEURONECTES, 1036.
- ii. Scales imperfectly imbricated, or else not all ctenoid.
- j. Scales chiefly cycloid in both sexes; lower pharyngeals small and narrow, separate, each with about 1 row of small, bluntish teeth; teeth incisor-like, close set, forming a cutting edge; no stellate scales at base of dorsal and anal. PLEURONECTES, 1037.
- jj. Scales rough-ctenoid in the male, more or less cycloid in the female (fin rays scaly in the male, naked in the female); lower pharyngeals very large, more or less united in the adult, their surface somewhat concave, the teeth in 5 or 6 rows, large, blunt, close set; teeth in jaws incisor-like; fin rays of dorsal and anal without tubercles at base. LIOPSETTA, 1038.

- jjj. Scales all in both sexes and on both sides of the body represented by coarse scattered stellate tubercles; similar tubercles between bases of dorsal and anal rays; lateral line without scales; lower pharyngeals broad, each with 3 rows of blunt, coarse teeth; teeth incisor-like. PLATICTHYS, 1039.
- aa. Vertebrae in increased number (varying from  $13 + 35 = 48$  to  $13 + 52 = 65$ ); dorsal rays 90 to 120; anal rays 70 to 100; teeth broad, incisor-like; scales small, all eyeloid. (Genera allied to *Glyptocephalus*.)
- k. Left side of skull normal; anal spine obsolete; vertebrae 48 to 52.
- l. Body elongate, the depth  $2\frac{1}{2}$  to 3 in length; vertebrae 48 to 52. MICROSTOMUS, 1040.
- ll. Body stenter, the depth 2 to  $2\frac{1}{2}$  in length; vertebrae more numerous, about 63. EMBASSICHTHYS, 1041.
- kk. Left side of skull with large mucous cavities; anal spine strong; vertebrae 58 to 65. GLYPTOCEPHALUS, 1042.

## III.—PSETTINÆ.

## (TURBOT TRIBE.)

*Large-mouthed flounders, with the ventral fins unsymmetrical.*—Mouth symmetrical, the dentition nearly equally developed on both sides; gape usually wide (narrow in *Platophrys*, *Etropus*, etc.), the maxillary commonly more than  $\frac{1}{2}$  length of head; lower pharyngeals narrow, each with one or more rows or a narrow band of small, sharp teeth; teeth in jaws acute; eyes not minute; pectorals and ventrals usually well developed; edge of preopercle free; ventral fins dissimilar in form or in position, that of the left or eyed side inserted on the ridge of the abdomen, its base extended along this ridge, its rays more or less wide apart; caudal fin rounded or subtruncate; no accessory lateral line; anal spine usually weak or obsolete; a pelvic spine sometimes developed; vertebrae in moderate or small number, 31 to 45. Body sinistral. Species chiefly tropical or subtropical in distribution.

- a. Pectoral fin of both sides present; septum of gill cavity below gill arches without foramen; a deep emargination near the isthmus; ventral fins free from anal.
- b. Vomer with teeth; lateral line with a strong arch in front; teeth subequal, in villiform bands; body broadly ovate; caudal fin subsessile; interorbital area broad; scales small, cycloid; gill rakers long and slender; anterior dorsal rays produced; vertebrae 36. LOPHOPSETTA, 1043.
- bb. Vomer toothless; ventral fins free from anal; caudal fin subsessile.
- c. Lateral line with a distinct arch in front; teeth small, uniserial, or imperfectly biserial.
- d. Interorbital space more or less broad, deeply concave, at least in the males; form broad ovate; gill rakers short and thick.
- e. Scales small, ctenoid, adherent, 75 to 100 or more; anterior rays of dorsal not elevated; pectoral of left side usually filamentous in the male; vertebrae (in *P. lunatus*)  $9 + 30 = 39$ . PLATOPHRYS, 1044.
- ee. Scales moderate, 60 to 70; anterior rays of dorsal greatly produced; no lateral line on blind side. PERISSIAS, 1045.
- dd. Interorbital space a narrow ridge; dorsal not elevated in front.
- f. Gill rakers obsolete; interorbital area armed with a spine; scales rough. ENOYOPHRYS, 1046.
- ff. Gill rakers slender; right ventral elongate; scales ctenoid. TRICHOPSETTA, 1047.

cc. Lateral line without arch in front.

g. Teeth in upper jaw biserial, in the lower uniserial, the front teeth of upper jaw enlarged; vertebrae 35 or 36; gill rakers short; interorbital space broad in the male. SYACIUM, 1048.

gg. Teeth in each jaw uniserial; interorbital space very narrow, the ridges coalescing between the eyes.

h. Mouth not very small, the maxillary more than  $\frac{1}{2}$  length of head.

i. Gill rakers very short and thick, tubercle-like.

j. Scales cycloid, small, and firm. CYCLOPSETTA, 1049.

ji. Scales small, firm, ctenoid. AZEVIA, 1050.

ii. Gill rakers slender, of moderate length; scales thin, deciduous, ciliated; vertebrae 34 to 40. CITHARICHTHYS, 1051.

hh. Mouth very small, the teeth subequal, the maxillary less than  $\frac{1}{2}$  length of head; scales thin; teeth uniserial; vertebrae 9 + 25 = 34. ETROPUS, 1052.

aa. Pectoral fin of blind side wanting; eyes very close together; caudal fin subsessile; teeth small, uniserial; mouth moderate; lateral line of eyed side arched, that of right side nearly straight; dorsal fin beginning on snout, its anterior rays not exerted, its rays all simple and very numerous; gill rakers few and feeble; scales small; body thin, very elongate; vertebrae (in *M. sessilicauda*) 43; (deep-sea flounders). MONOLENE, 1053.

### 1013. ATHERESTHES, Jordan & Gilbert.

*Atheresthes*, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1880, 51 (*stomias*).

Eyes and color on the right side. Body very long and slender, closely compressed, tapering into a long and slender caudal peduncle; head elongate, narrow; mouth extremely large, oblique; the long and narrow maxillary extending beyond the eye; each jaw with 2 irregular series of sharp, unequal, arrow-shaped teeth, some of them long and wide set, and others short and close set, sharp; the long teeth freely depressible. Gill rakers numerous, long, slender, and stiff, strongly dentate within. Scales rather large, thin and readily deciduous, slightly ciliated, those on the blind side similar, smooth; lateral line without arch. Fins low and fragile; dorsal commencing over the eye, its anterior rays low, the posterior rays somewhat forked; no anal spine; pectorals and ventrals small, both of the latter lateral; caudal lunate. The single species which constitutes this genus is one of the most remarkable of the flounders. Of all the group, it approaches in form and general characters most nearly to the Gadoid fishes, from ancestors of which we may presume the flounders to be descended, although Dr. Gill has suggested the possibility of their descent from Trachypteroid fishes. (*ἀθήρη*, the beard or spike of an ear of corn; *ἔσθια*, to eat; from the arrow-shaped teeth.)

### 2975. ATHERESTHES STOMIAS (Jordan & Gilbert).

(THE ARROW-TOOTHED HALIBUT.)

Head about  $3\frac{1}{2}$  in length; depth  $3\frac{1}{2}$ ; eye large,  $4\frac{1}{2}$  in head. D. 103; A. 86; scales 135; vertebrae 12 + 37 = 49. Head long, the snout protruding, somewhat truncate at tip; mouth excessively large; the maxillary more than  $\frac{1}{2}$  length of head, and reaching behind eye; premaxillary in front above the level of the lower eye; teeth in upper jaw anteriorly in a single

series, long, slender, and wide set, much smaller and closer set behind; on sides of jaw the teeth are very small and in 2 distinct series, the inner of which corresponds to the single series in front, the teeth thus gradually increasing in size forward; teeth in inner series of lower jaw very sharp and slender, longer than the upper teeth, wide set, alternating with shorter, depressed teeth; outside of these larger teeth is a series of fixed small teeth; all of the long teeth in both jaws depressible and conspicuously arrow-shaped toward their tips; inner series of small teeth in upper jaw also arrow-shaped, depressible; interorbital space scaly, ridged, not a third width of eye. Gill rakers long and strong, about 4 + 13 in number, the longest more than  $\frac{1}{2}$  diameter of eye. Upper eye with its range entirely vertical. Scales extremely thin, irregular in size, not evenly imbricated; lateral line very prominent. Dorsal fin beginning just behind the middle of the eye; caudal peduncle nearly as long as the pectoral fin, about  $\frac{2}{3}$  length of head. Plain olive brown, the margins of the scales darker; blind side dusted with black points. Length 2 feet. Bering Sea to San Francisco, common northward; not rare in deep water off San Francisco, and is brought in in considerable numbers from the sweep-nets (*parranzelle*) used in Drakes Bay. At Unalaska it occurs commonly in shallow water. In the north the flesh is firmer and the coloration more pronounced. Dr. Gilbert dredged it in abundance on both sides of the peninsula of Alaska and in Bristol Bay, in 32 to 406 fathoms. Mr. Scofield found it abundant in Chignik Bay, and it was taken by us in 1897 at Unga and Karluk. (*στρομιάς*, large mouthed.)

*Platysomatichthys stomias*, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1880, 51, 301, San Francisco. (Coll. Jordan & Gilbert.)

*Atheresthes stomias*, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1880, 57, 454; BEAN, Proc. U. S. Nat. Mus. 1881, 242; JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1881, 66; JORDAN & GILBERT, Synopsis, 820, 1883; BEAN, Proc. U. S. Nat. Mus. 1883, 354; JORDAN, Nat. Hist. Aquat. Anim., 188, pl. 53, 1884; JORDAN & GOSS, Review Flounders and Soles, 236, pl. 1, 1889; GILBERT, Rept. U. S. Fish Comm. 1893 (1896), 459.

#### 1014. REINHARDTIUS, GILL.

*Reinhardtius*, GILL, Cat. Fishes East Coast N. A., 50, 1861 (*hippoglossoides*; no description).

*Platysomatichthys*, BLEEKER, Comptes Rendus, Ac. Sci. Amsterdam, XIII, 1862, 426 (*pinniguis* = *hippoglossoides*).

*Reinhardtius*, GILL, Proc. Ac. Nat. Sci. Phila. 1864, 218 (*hippoglossoides*).

Eyes and color on right side. Body more or less elongate, compressed; head long and large; mouth large; maxillary reaching beyond eye; jaws with strong, unequal teeth, the upper with 2 series in front, these converging behind; lower jaw with a single series of strong, distant teeth; no teeth on vomer or palatines. Gill rakers few, short, stout, and rough. Fins rather low; caudal fin lunate. Lower pharyngeal teeth in 1 row. Scales small, cycloid; lateral line without anterior curve. One species known, an arctic fish, in some degree intermediate between the true halibut and *Atheresthes*. (Named for Prof. Johann Reinhardt, of the University of Copenhagen, an able investigator of the fishes of Greenland.)

2976. REINHARDTIUS HIPPOGLOSSOIDES (Walbaum).

(GREENLAND HALIBUT.)

Head  $3\frac{1}{2}$  in length; depth nearly 3. D. 100; A. 75; scales 160; orbit 8 in head; snout about  $3\frac{1}{2}$ , more than twice as long as orbit; eyes even in front; interorbital space flat, scaly, wider than the orbit; lower jaw prominent; length of maxillary  $2\frac{1}{2}$  in head; teeth conical, pointed; upper jaw with 2 series, convergent posteriorly, those of the outer series gradually smaller posteriorly; a pair of strong canine teeth anteriorly in the inner series, the other teeth of this series being very small; lower jaw with a series of strong, distant teeth. Gill rakers short, thick, and strongly dentate. Fins naked. Longest dorsal rays  $\frac{1}{2}$  length of head; no anal spine; dorsal and anal rays all simple, the dorsal beginning over posterior third of the eye. Scales very small, not ciliated. Yellowish brown. Reaching a very large size. Arctic parts of the Atlantic, south to Finland and the Grand Banks; not very common. (Eu.) (*ἰππόγλωσσός*, halibut; *εἶδος*, resemblance.)

*Pleuronectes cynoglossus*, FABRICIUS, Fauna Grœnlandica, 163, 1780, Greenland; not of LINNÆUS.

*Pleuronectes hippoglossoides*, WALBAUM, Artedi Piscium, 115, 1792; based on FABRICIUS.

*Pleuronectes pinguis*, FABRICIUS, Zoologiske Bidrag., 43, 1824, Greenland.

*Hippoglossus grœnlandicus*, GÜNTHER, Cat., IV, 404, 1862, Greenland.

*Reinhardtius hippoglossoides*, GILL, Cat. Fishes East Coast N. A., 50, 1861; GILL, Proc. Ac. Nat. Sci. Phila. 1864, 218.

*Platysomatichthys hippoglossoides*, GOODE & BEAN, Bull. Essex Inst., II, 7, 1879; COLLETT, Norske Nord-Havs Exped., 142, 1880; JORDAN & GILBERT, Synopsis, 819, 1883; GOODE, Nat. Hist. Aquat. Anim., 197, pl. 56, 1884; JORDAN & GOSS, Review Flounders and Soles, 237, pl. II, 1889; and of late American writers generally.

*Hippoglossus pinguis*, REINHARDT, Kgl. Dansk. Vidensk. Selsk., 116, 1838.

*Platysomatichthys pinguis*, BLEEKER, l. c., 426, 1862.

1015. HIPPOGLOSSUS, Cuvier.

(HALIBUT.)

*Hippoglossus*, CUVIER, Règne Animal, Ed. 1, II, 221, 1817 (*hippoglossus*).

Eyes and color on the right side. Form oblong, not strongly compressed. Mouth wide, oblique; teeth in the upper jaw in 2 series, those below in 1; anterior teeth in upper jaw, and lateral teeth in lower, strong; no teeth on vomer or palatines; lower pharyngeal teeth in 2 rows. Dorsal fin beginning above the eye, its middle rays elevated, the posterior rays of dorsal and anal bifid; caudal fin lunate; ventral fins both lateral. Scales very small, cycloid; lateral line with a strong curve in front. Gill rakers few, short, compressed, wide set. Vertebrae 16 + 34. Largest of the flounders. This genus contains but 1 species, the well-known halibut; abundant on both coasts of the North Atlantic and of the North Pacific. (*Hippoglossus*, the ancient name of the halibut, from ἵππος, horse; γλῶσσα, tongue.)

2977. HIPPOGLOSSUS HIPPOGLOSSUS (Linnæus).

(HALIBUT.)

Head  $3\frac{1}{2}$ ; depth about 3. D. 105; A. 78; scales 150 or more. Body comparatively elongate, not strongly compressed, deep mesially, thence rapidly tapering each way; head broad; eyes large, separated by a very broad

flattish area; lower eye slightly advanced; mouth large, the maxillary reaching middle of orbit. Nearly uniform dark brown; blind side white. One of our most important food-fishes, reaching a weight sometimes of 400 pounds. Found in all northern seas, southward in deep water to France, Sandy Hook, and occasionally to the Farallones off San Francisco; abundant throughout the North Atlantic as also the North Pacific and Bering Sea, in water of moderate depth; taken with hook and line on all cod banks.

*Pleuronectes hippoglossus*, LINNÆUS, *Systema Naturæ*, Ed. x, 269, 1758, European Ocean.  
*Hippoglossus vulgaris*, FLEMING, *British Animals*, 197, 1828; GÜNTHER, *Cat.*, IV, 403, 1862; DAY, *Fishes Great Britain*, II, 5, pl. 44; STORER, *Fish. Mass.*, 145, 1839; DE KAY, *New York Fauna: Fishes*, pl. 49, f. 157, 204, 1842; STORER, *Synopsis Fish. N. A.*, 475, 1847; LOCKINGTON, *Rep. Com. Fisheries California*, 39, 1878-79; LOCKINGTON, *Proc. U. S. Nat. Mus.* 1879, 71; BEAN, *Proc. U. S. Nat. Mus.* 1879, 63; JORDAN & GILBERT, *Proc. U. S. Nat. Mus.* 1880, 454; GOODE, *Proc. U. S. Nat. Mus.* 1880, 471; JORDAN & GILBERT, *Proc. U. S. Nat. Mus.* 1881, 66; BEAN, *Proc. U. S. Nat. Mus.* 1881, 242; JORDAN & GILBERT, *Synopsis*, 819, 1883; BEAN, *Cat. Col. Fish. U. S. Nat. Mus.* 1883, 20; DRESEL, *Proc. U. S. Nat. Mus.* 1884, 244; GOODE, *Nat. Hist. Aquatic Anim.*, 189, pl. 54, 1884; and of American writers generally.

*Hippoglossus maximus*, GOTTSCHKE, *Archiv für Naturgesch.* 1835, 164, no locality.

*Hippoglossus gigas*, SWAINSON, *Nat. Hist. Class'n Anim.*, II, 302, 1839, no locality.

*Hippoglossus ponticus*, BONAPARTE, *Catalogo Metodico*, 47, 1846, Black Sea; after PALLAS.

*Hippoglossus americanus*, GILL, *Proc. Ac. Nat. Sci. Phila.* 1864, 220.

*Hippoglossus hippoglossus*, JORDAN, *Cat. Fish. N. A.*, 133, 1885; JORDAN & GOSS, *Review Flounders and Soles*, 237, pl. 3, 1889.

#### 1016. LYOPSETTA, Jordan & Goss.

*Lyopsetta*, JORDAN & GOSS, in JORDAN, *Cat. Fish. N. A.*, 135, 1885 (*exilis*).

Teeth sharp, those of the lower jaw uniserial, the upper jaw biserial; lateral line simple (without accessory dorsal branch) and without distinct anterior arch. Scales comparatively large, thin, ciliated, and deciduous; body dextral; anal spine usually strong; vertebræ about 45; body slender, the flesh soft; dorsal fin beginning above eye. This genus contains but a single species, a small, soft-bodied flounder, of the waters of the North Pacific. In its technical characters *Lyopsetta* is very close to *Hippoglossoides*, but the species has the soft flesh of *Atheresthes*. (λύω, to loosen; ψήττα, flounder.)

#### 2978. LYOPSETTA EXILIS (Jordan & Gilbert).

Head 4; depth 3½. D. 78; A. 62; V. 6; scales 16-71-18. Body slender, compressed, the flesh soft; caudal peduncle slender; mouth not large, very oblique, the gape curved; lower jaw scarcely projecting, with a knob at symphysis; maxillary rather narrow, reaching middle of pupil, 2½ in length of head; teeth small, slender, close set, nearly uniform; above in 2 series, below in 1. Eyes large, separated by a sharp, scaly ridge; lower eye advanced. Scales comparatively large, thin and deciduous, ctenoid, but not so rough as in the other species, those on blind side similar, less rough. Lateral line prominent, rising anteriorly, without trace of arch. Fins low, fragile; anal preceded by a spine; caudal fin long, rather pointed; pectorals small, the right pectoral little more

than  $\frac{1}{2}$  length of head. Dorsal beginning immediately in front of pupil; anal higher than dorsal. Gill rakers short, slender, toothed, 9 below angle, the longest about  $\frac{1}{2}$  diameter of orbit. Pale olivaceous brown, with dark points, forming edgings on each scale; bronze spots sometimes present; fins mostly dusky; dorsal and anal edged anteriorly with yellowish; ventrals largely yellow. Length 12 inches. North Pacific, in rather deep water; San Francisco to Puget Sound. This small flounder is brought in in large quantities by the sweep nets off San Francisco. It is of little value as a food-fish. (*exilis*, slender.)

*Hippoglossoides exilis*, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1880, 154, off San Francisco (Type, No. 27121. Coll. Jordan & Gilbert); JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1880, 454; JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1881, 67; JORDAN & GILBERT, Synopsis, 827, 1883.

*Lyopsetta exilis*, JORDAN & GOSS, Review Flounders and Soles, 238, 1880.

1017. **EOPSETTA**, Jordan & Goss.

*Eopsetta*, JORDAN & GOSS, in JORDAN, Cat. Fish. N. A., 135, 1885 (*jordani*).

Teeth sharp, those of the lower jaw uniserial, the upper biserial; scales small, ciliated, and adherent; lateral line without accessory dorsal branch and without distinct anterior arch; anal spine usually strong; body normally dextral, robust, the flesh firm; dorsal fin beginning above eye; vertebrae about 43. This genus contains but a single species, a large flounder which is abundant on the coast of California. It is very close to the genus *Hippoglossoides*. (*ἔως*, morning; *ψῆττα*, flounder.)

2979. **EOPSETTA JORDANI** (Lockington).

(CALIFORNIA "SOLE.")

Head  $3\frac{1}{2}$ ; depth  $2\frac{1}{2}$ . D. 94; A. 72; scales 96. Body broadly elliptical. Dorsal and ventral outline equally and regularly curved. Mouth oblique, the jaws about even, the symphyseal knob but little projecting; gape curved; maxillary broad, reaching to behind pupil,  $2\frac{1}{2}$  in head; teeth in 2 series in the upper jaw, the inner series small and distant from the outer, which is considerably enlarged in front; lower jaw with a single series similar to the outer series in the upper jaw, but larger. Gill rakers roughish, strong, about 15 below angle, the longest about  $\frac{1}{2}$  as long as eye. Lower pharyngeals rather narrow, each with a single row of sharp teeth. Eyes large; interorbital space a narrow, blunt, scaly ridge. Dorsal beginning over anterior margin of pupil, the rays all simple; caudal fin with the middle rays slightly produced; anal preceded by a spine; pectoral  $\frac{1}{2}$  length of head. Scales of colored side small, firm, strongly ciliated, nearly uniform over head and body; lower jaw and snout scaleless; scales on blind side smooth. Olive brown, nearly uniform; membrane of dorsal and anal fins clouded with darker. Length 20 inches. Pacific Coast of the United States from Puget Sound to Point Concepcion. One of the commonest flatfishes of the California coast, being found in abundance in shallow water from Monterey northward. It is a good food-fish, and large numbers are dried each year by the Chinese. (Named for David Starr Jordan.)



*Hippoglossoides jordani*, LOCKINGTON, Proc. U. S. Nat. Mus. 1879, 73, San Francisco (Coll. W. N. Lockington); LOCKINGTON, Rep. Com. Fisheries California 1878-79, 40; LOCKINGTON, Scientific Press Supplement, April, 1879, 120; JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1880, 454; JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1881, 67; JORDAN & GILBERT, Synopsis, 826, 1883; JORDAN, Nat. Hist. Aquat. Anim., 187, 1884.  
*Eopsetta jordani*, JORDAN & GOSS, Review Flounders and Soles, 239, 1885.

## 1018. HIPPOGLOSSOIDES, Gottsche.

*Hippoglossoides*, GOTTSCHÉ, Archiv für Naturgesch. 1835, 164 ("*limanda*" = *platessoides*).  
*Citharus*, REINHARDT, Kong. Dansk. Vid. Selsk. 1838, 116 (*platessoides*); not *Citharus* BLEEKER, 1862.  
*Drepanopsetta*, GILL, Cat. Fish. East Coast N. A., 50, 1861 (*platessoides*).  
*Pomatopsetta*, GILL, Proc. Ac. Nat. Sci. Phila. 1864, 217 ("*dentata*" = *platessoides*).

Eyes and color on the right side (except sometimes in *H. classodon*). Body oblong, moderately compressed; mouth rather large, with 1 row of sharp teeth on each jaw; no teeth on vomer or palatines; gill rakers rather long and slender; scales ctenoid; lateral line nearly straight, simple; dorsal fin low in front, beginning over or before the eye; ventrals both lateral; caudal double truncate, produced behind. This genus, as here restricted, contains 3 closely related species, 2 of the North Pacific, 1 of the North Atlantic. All are essentially arctic species, inhabiting shallow waters in the regions where they are most abundant. (*ἰππόγλωσσος*, *Hippoglossus*; *εἶδος*, resemblance.)

a. Dorsal rays about 88; anal about 70; gill rakers  $x + 10$ ; interorbital space with an obtuse, prominent, rather broad ridge. PLATÉSSOIDES, 2980.

aa. Dors. rays about 82; anal about 61; gill rakers  $x + 12$  to 14; interorbital space with a narrow, nearly naked ridge. ELASSODON, 2981.

aaa. Dorsal rays 72 to 76; anal 56 to 60; gill rakers  $x + 12$ ; interorbital space moderate, with 2 rows of scales.

b. Depth  $2\frac{1}{2}$  in length; D. 76; A. 60; pectoral  $\frac{1}{2}$  length of head. ROBUSTUS, 2982.

bb. Depth  $2\frac{1}{2}$  in length; D. 73; A. 56; pectoral  $\frac{1}{3}$  in length of head.

HAMILTONI, 2983.

## 2980. HIPPOGLOSSOIDES PLATESSOIDES (Fabricius).

(SAND-DAB.)

Head  $3\frac{1}{2}$ ; depth  $2\frac{1}{2}$ . D. 88 (80 to 93); A. 70 (64 to 75); scales 90 (pores). Body ovate; mouth moderate, oblique; maxillary narrow, reaching to below pupil,  $2\frac{1}{2}$  in length of head; teeth rather small, conical, larger anteriorly, in 1 row in each jaw, those in the lower largest. Eyes rather large, the upper longer than snout,  $4\frac{1}{2}$  in head; lower jaw included, but with a projecting knob at the chin; snout thick, scaly; interorbital space narrow, with a raised obtuse ridge entirely covered with rough scales in about 6 series; mandible with a series of scales; gill rakers rather short and robust, not toothed, about 10 below angle, the longest less than  $\frac{1}{2}$  length of eye; fins with small, rough scales; a strong preanal spine; pectoral not quite  $\frac{1}{2}$  length of head. Reddish brown, nearly plain. The identity of the American and European representatives of this species (*platessoides* and *limandoides*) is now conceded by all writers. A little difference is recognizable between arctic and subarctic examples, the

former having a somewhat greater number of fin rays. Thus Greenland specimens, according to Collett, have D. 88, A. 69; specimens from Finmark have D. 92, A. 72; these representing the var. *platessoides*. Specimens from England (var. *limandoides*) have D. 80, A. 66, while those from intermediate localities present in general fin formula likewise intermediate, showing that no sharp division is possible. This is a rather common food-fish of the deep waters northward, on both sides of the ocean. North Atlantic, south to Cape Cod, and the coasts of England and Scandinavia. (Eu.) (*platessa*, the plaice; εἶδος, resemblance.)

- Pleuronectes linguatula*, MÜLLER, Zool. Dan. Prodrömus, 45, 1776; not of LINNÆUS.  
*Pleuronectes platessoides*, FABRICIUS, Fauna Greenlandica, 164, 1780, Greenland.  
*Pleuronectes limandoides*, BLOCH, Ansl. Fische, iii, 24 tab. 186, 1787, Europe, and of various copyists.  
*Pleuronectes limandanus*, PARNELL, Edinburgh New Phil. Journ. 1835, 210.  
*Citharus platessoides*, REINHARDT, Kongl. Dansk. Vid. Selsk., 116, 1838.  
*Drepanopsetta platessoides*, GILL, Cat. Fish. East Coast N. A., 50, 1861.  
*Hippoglossoides platessoides*, GILL, Proc. Ac. Nat. Sci. Phila. 1864, 217; COLLETT, Norskø Nord-Havs. Exped., 144, 1880; GOODE, Proc. U. S. Nat. Mus. 1880, 471; JORDAN & GILBERT, Synopsis, 826, 1883; STEARNS, Proc. U. S. Nat. Mus. 1883, 125; GOODE, Nat. Hist. Aquatic Ania., 197, pl. 55, 1884; JORDAN & GOSS, Review Flounders and Soles, 240, pl. 4, 1889; GOODE & BEAN, Ocean Ichthyology, 438, 1896, and of recent American writers generally.  
*Hippoglossoides limandoides*, GÜNTHER, Cat., iv, 405, 1862; DAY, Fishes Great Britain and Ireland, ii, 9, pl. 45, 1884.  
*Hippoglossoides limanda*, GOTTSCHÉ, Archiv für Naturgesch. 1835, 168; not *P. limanda*, LINNÆUS.  
*Platessa dentata*, STORER, Rept. Fish. Mass., 143, 1839; DE KAY, N. Y. Fauna: Fishes, 298, 1842; STORER, Synopsis, 476, 1846.  
*Hippoglossoides dentatus*, GÜNTHER, Cat., iv, 406, 1862; GÜNTHER, Challenger Report, xxii, Fishes, 3, 1887.  
*Pomatopsetta dentata*, GILL, Proc. Ac. Nat. Sci. Phila. 1864, 217.

2981. HIPPOGLOSSOIDES ELASSODON, Jordan & Gilbert.

Head  $3\frac{1}{2}$ ; depth  $2\frac{1}{2}$ ; eye 4 in head. D. 77 to 87; A. 59 to 67; V. 6; scales 45-100-40. Body oblong-elliptical; caudal peduncle about as long as deep; upper profile of head continuous with the outline of back; depression over eye slight; mouth rather large, the gape curved, considerably wider on the blind side; lower jaw projecting, with a symphyseal knob; maxillary narrow, reaching beyond middle of pupil,  $2\frac{1}{2}$  in head; teeth small, close set, nearly uniform, in a single row. Gill rakers slender, smooth, 14 to 16 below arch, the longest nearly  $\frac{1}{2}$  diameter of orbit. Eyes large, separated by a narrow, knife-like ridge, which is naked, or with a single series of scales. Scales small, firm, rough, those on tail roughest, those on blind side similar, mostly smooth anteriorly. Lateral line rising anteriorly, but without arch; dorsal beginning immediately in front of pupil; anal preceded by a spine; caudal long; pectoral of eyed side  $\frac{1}{2}$  length of head; ventral reaching past front of anal; pectoral and ventral of eyed side with prickle-like scales. Brownish, nearly uniform, sometimes spotted with darker; fins grayish, irregularly blotched with dusky. Body sometimes sinistral. Length 18 inches. Bering Sea south to Cape Fattery; a rather abundant shore fish in Puget Sound, and it

seems to be still more common northward, being, in Alaska, a food-fish of some importance. Abundant north and south of the Aleutian Islands and in Bristol Bay. Our specimens from Kamchatka agree in all respects; D. 77 to 84; A. 60 or 61. Pectoral not quite  $\frac{1}{2}$  head. Interorbital ridge sharp, with 1 series of scales; gill rakers  $x+11$ . (*Ἰπποβόω*, to diminish; *ὀδόν*, tooth.)

*Hippoglossoides classodon*, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1880, 178, Seattle; Tacoma (Type, No. 27263. Coll. D. S. Jordan); JOHNSON & GILBERT, Proc. U. S. Nat. Mus. 1880, 451; BEAN, Proc. U. S. Nat. Mus. 1881, 242; JORDAN & GILBERT, Synopsis, 826, 1883; BEAN, Proc. U. S. Nat. Mus. 1883, 20; JORDAN, Nat. Hist. Aquat. Anim., 188, pl. 52, 1884; JORDAN & GOSS, Review Flounders and Soles, 241, pl. 5, 1889; JORDAN & GILBERT, Rept. Fur Seal Invest., 1898.

2982. *HIPPOGLOSSOIDES ROBUSTUS*, Gill & Townsend

Head  $3\frac{3}{8}$ ; depth  $2\frac{1}{8}$ ; eye  $5\frac{1}{2}$  in head. D. 76; A. 60; scales 95 (pores). Interorbital space a broad, somewhat elevated ridge with 2 rows of scales. Body rather high, its greatest height nearly equaling  $\frac{1}{2}$  the length from snout to base of caudal; profile decurved above the eye; body thick; scales on head separate and rarely touch each other. Gill rakers long,  $x+11$ . Maxillary  $2\frac{1}{2}$  in head, directed upward anteriorly; teeth of the single row mostly separated from each other by intervals equal to width of teeth, curved inward, and uniform on the sides; toward front 4 or 5 enlarged, preceded by 2 smaller, leaving the middle toothless; in the lower jaw of nearly uniform size and inclining backward. Pectoral  $\frac{1}{2}$  head; ventrals reaching first or second anal ray. Scales on body ciliated or weakly ctenoid, those on cheek smoother; no ctenoid scales on blind side. Caudal shorter than in *H. hamiltoni*,  $1\frac{1}{2}$  in head. No exerted nasal tubes. Color plain brown. Bering Sea. Only the type known,  $12\frac{1}{2}$  inches long, from which we have taken the above description. (*robustus*, robust.)

*Hippoglossoides robustus*, GILL & TOWNSEND, Proc. Biol. Soc. Wash., XI, 1897 (sept. 17, 1897), 234, Bering Sea, Lat.  $56^{\circ} 14' N.$ , Long.  $164^{\circ} 08' W.$ , Albatross Station 3541, in 49 fathoms. (Type, No. 48766, U. S. Nat. Mus. Coll. Albatross.)

2983. *HIPPOGLOSSOIDES HAMILTONI*, Jordan & Gilbert, new species.

Head  $3\frac{1}{2}$  in length; depth  $2\frac{1}{2}$ ; longest diameter of upper eye  $3\frac{1}{2}$  in head; snout (measured from upper eye) 5 in head; maxillary of colored side  $2\frac{1}{2}$ , of blind side  $2\frac{1}{2}$ , in head; depth of caudal peduncle equaling its length,  $3\frac{1}{2}$  in head. D. 79; A. 56; P. 11; pores in lateral line 91. Upper profile of head continuing the dorsal curve without interruption, there being a slight depression above the eye and an increased convexity on the snout; mandible very heavy, projecting anteriorly, so that its symphyseal profile completes the curve of the snout; a very short prominence at symphysis directed vertically downward; gape strongly curved and the mouth narrowed anteriorly, so that the maxillary and premaxillary are almost wholly concealed along the middle of their length by the overarched prefrontal; teeth acute, in a single series in each jaw, all except the anterior teeth in each jaw short; at the symphysis of lower jaw the teeth are

longer and directed inward, while in the anterior end of each premaxillary the teeth are still more enlarged, and the series on each side describes a strong curve with its convex side toward the median line; maxillary reaching vertical from slightly behind middle of lower eye; nostril tubes conspicuous, the anterior in closest proximity to the upper lip, which it entirely overhangs; posterior nostril tube wider and slightly shorter; eyes of nearly equal size, and opposite, separated by a wider ridge than in *H. classodon*, the ridge bearing in its narrowest portion 2 well-defined rows of strongly spinous scales; a conspicuous series of pores joining lateral line with upper margin of upper eye, and another encircling the lower eye below and behind; a third series along mandible and preopercle; 1 large pore above posterior nostril; gill rakers slender, unarmed, 2 above the angle, 11 or 12 below it, the longest  $2\frac{1}{2}$  in eye; dorsal fin beginning above front of pupil, the longest ray  $2\frac{3}{8}$  in head; anal preceded by a strong spine, its height equaling that of dorsal; pectoral very long and slender,  $\frac{3}{4}$  length of head, that of blind side shorter,  $\frac{1}{2}$  length of head; ventrals reaching to base of fourth or fifth anal ray; caudal long, evenly rounded behind, the middle rays not longer than those adjacent, their length equaling distance from tip of snout to preopercular margin; scales on colored side strongly ctenoid except in a strip along middle of sides anteriorly; elsewhere each scale provided with 2 to 4 long spines; on blind side they are smooth except on nape and caudal peduncle; cheeks, opercles, and interorbital space covered with larger, rougher scales than those on sides; mandible and snout naked; a single series surrounding each eye anteriorly, and 1 on maxillary or colored side; blind side of head with maxillary naked; cheeks covered with minute smooth thin scales, the opercles with a few scattered spinous scales, the preopercle naked. Color nearly uniform brownish, without distinctive markings on body or fins. One specimen, 17 cm. long, from Albatross Station 3641, off Dalnoi Point, Kamchatka; depth 16 fathoms. Allied to *Hippoglossoides classodon*, from which it differs in the fewer fin rays and scales, the wider interorbital space, the longer caudal and pectoral fins and the much smaller symphyseal knob. The nasal tubes are larger, the scales rougher, and the anterior part of lateral line more arched. Its relations with *H. robustus* are much nearer but the species are apparently distinct. (Named for Gerald Edwin H. Barrett-Hamilton, of Dublin, member of the British Commission of Fur Seal Investigation, 1896 and 1897, who made valuable collections of Kamchatkan fishes.)

*Hippoglossoides hamiltoni*, JORDAN & GILBERT, Rept. Fur Seal Invest., 1898, Dalnoi Point, Kamchatka. (Coll. Albatross.)

#### 1019. PSETTICHTHYS, Girard.

*Psettichthys*, GIRARD, Proc. Ac. Nat. Sci. Phila. 1854. 140 (*melanostictus*).

Body dextral; teeth uniserial, sharp, unequal, some of them canine-like; mouth moderate, the lower pharyngeal teeth sharp, uniserial; scales small, ctenoid, ciliated, and firm; lateral line with an accessory dorsal branch and without distinct anterior arch; anal spine strong; dorsal fin

beginning before the eye; vertebrae about 40; flesh firm. This genus contains but 1 species, found on the coast of California. It is nearly related to *Hippoglossoides*, but possesses the peculiar accessory dorsal branch to the lateral line, characteristic of so many of the Pacific coast flounders. (*ψιττα*, the turbot; *ιχθύς*, fish.)

2984. *PSETTICHTHYS MELANOSTICTUS*, Girard.

Head 4; depth  $2\frac{1}{2}$ . D. 85; A. 60; scales 112. Body not very deep, elliptical; mouth rather small, the maxillary extending to below pupil,  $2\frac{2}{3}$  in head; teeth large, in a single series in each jaw, those in lower jaw largest; a few large canines in front of each jaw. Eyes very small, separated by a broad, flat, scaly space, without ridge; lower eye slightly in advance of upper; gill rakers rather stout, weak, hooked at tip, 14 below the angle; scales very small, etenoid on colored side; lateral line nearly straight, with a long accessory dorsal branch; dorsal commencing in advance of upper eye, the anterior rays elevated, slender and exerted, the longest about  $\frac{1}{2}$  length of head; first ray of dorsal nearly free from its membrane; pectoral fin short,  $2\frac{1}{2}$  in head; anal fin preceded by a spine; caudal large, strongly convex; lower pharyngeals very narrow, each with 1 row of sharp, recurved teeth. Grayish brown, finely speckled with darker on body and fins. Pacific coast of North America, from Sitka south to Monterey. This is one of the commoner flounders of the Pacific coast, being everywhere known by the name of "Sole." It lives near the shore, and reaches a length of about 20 inches. In color this species is quite unlike the species of *Hippoglossoides*, but in most other respects the two groups are closely allied. (*μέλας*, black; *στικτός*, spotted.)

*Psettichthys melanostictus*, GIRARD, Proc. Ac. Nat. Sci. Phila. 1854, 140, San Francisco; Astoria, Oregon; GIRARD, U. S. Pac. R. R. Surv., x, Fishes, 154, 1858; GÜNTHER, Cat., IV, 420, 1862; LOCKINGTON, Rep. Com. Fisheries Cal. 1878-79, 40; LOCKINGTON, Proc. U. S. Nat. Mus. 1879, 76; JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1880, 453; JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1881, 67; JORDAN, Nat. Hist. Acquatic Animals, 186, pl. 51, 1884; JORDAN & GOSS, Review Flounders and Soles, 241, pl. 6, 1889.

*Hippoglossoides melanostictus*, JORDAN & GILBERT, Synopsis, 828, 1883.

1020. *VERASPER*, Jordan & Gilbert, new genus.

*Verasper*, JORDAN & GILBERT, Report Fur Seal Invest., 1898 MS. (*mosevi*).

This genus is allied to *Xystreureys* and *Hippoglossina*, having few short gill rakers like the former and strongly etenoid scales like the latter. It differs strongly from all its congeners in having the premaxillary teeth in 2 series, teeth uniformly small, without canines. Body dextral; dorsal inserted above the front of pupil; lateral line strongly arched above the root of the pectoral, without recurrent dorsal branch; scales firm, extremely spinous; gill rakers short, thick, and triangular, few in number; none of the fin rays notably produced or exerted. Japan and Kuril Islands; 2 species known, the following and *V. variegatus* (Schlegel), a common food fish of Japan, the 2 very closely related. (*verus*, true; *asper*, rough, the word being suggested by *Veratrum*.)

2985. VERASPER MOSERI, Jordan &amp; Gilbert, new species.

Head  $3\frac{1}{4}$  in length to base of caudal; depth 2. D. 82; A. 58; pectoral 12; pores in lateral line 84. Depth of caudal peduncle 4 in greatest depth of body; length of caudal peduncle, measured axially,  $1\frac{1}{3}$  in its depth. Head much depressed, with rather wide, flat interorbital space, resembling in appearance *Psettichthys melanoostictus*, its thickness at interorbital space equaling distance between pupils of upper and lower eyes. Mouth small, very oblique, the gape strongly arched, the broad maxillary reaching a vertical behind middle of pupil,  $2\frac{1}{4}$  in head; mandible narrowing toward tip, with very rudimentary symphyseal knob. Teeth in upper jaw in 2 distinct series throughout, those of the outer series increasing slightly in size toward front of jaw, but none of them canine-like; mandibular teeth in 1 row, except at symphysis, where a few teeth form a short outer series. Nasal openings of eyed side approximated in front of middle of interorbital space, the anterior with a short tube, the posterior with a raised rim. Eyes small, their anterior margins opposite, the diameter of lower eye equaling distance from tip of snout to posterior nostral,  $6\frac{1}{2}$  in head. Interorbital space rather broad and flat, not ridge-like, its total width equaling  $\frac{1}{2}$  diameter of orbit. Gill rakers short, broad, triangular, minutely toothed on inner margin,  $\frac{1}{3}$  diameter of eye; 7 present on horizontal limb of outer arch. Lateral line with a short high anterior arch, the cord of which is  $\frac{1}{3}$  the straight portion; height of arch  $\frac{1}{3}$  its length; behind the arch lateral line descending in a gentle curve to middle of sides, the scales very rough, each possessing several long, sharp spines diverging from median portion of posterior margin; anterior and posterior portions of dorsal and anal fins naked, the rays of the middle portion each with a series of strongly ctenoid scales; caudal densely scaled to tip; pectorals and ventrals naked; head covered with strongly spinous scales, excepting snout, maxillary, and mandible; on blind side of head the snout, jaws, preopercle, subopercle, lower half of opercle, and all but a central strip on interopercle, scaleless; on blind side the scales are rough on head, ventral area, and along bases of ventral fins, largely smooth elsewhere. Dorsal beginning above front of pupil, the rays increasing in length to the forty-fifth, which is  $2\frac{2}{3}$  in head; longest anal ray (the seventeenth)  $2\frac{1}{2}$  in head. Caudal broadly rounded,  $1\frac{1}{2}$  in head; pectoral short and broad,  $2\frac{2}{3}$  in head; ventrals of nearly equal length, reaching origin of anal,  $3\frac{1}{2}$  in head; no anal spine. Color in spirits, centers of the scales light gray, the margins dark brown; fins light or dusky, the vertical fins with conspicuous black bars, parallel with the rays, these most evident on under side where the pigment seems principally to occur, and are seen through the fin more faintly on the colored side; lining of cheeks and gill cover of colored side dusky; peritoneum gray. Kuril Islands; 1 male 28 cm. long, from Shana Bay, Iturup Island; also taken at Hakodate. (Named for Jefferson Franklin Moser, U. S. N., Lieutenant-Commander, in charge of the U. S. Fish Commission Steamer *Albatross*, and a member of the United States Fur Seal Commission for 1896.)

*Verasper moseri*, JORDAN & GILBERT, Rept. Fur Seal Invest., 1898, U. S., Shana Bay, Iturup Island, Kuril Group. (Type No. 48797. Coll. *Albatross*, Capt. J. F. Moser.)

## 1021. HIPPOGLOSSINA, Steindachner.

*Hippoglossina*, STEINDACHNER, Ichth. Beiträge, v, 13, 1876 (*macrops*).

Teeth rather small, uniserial, no canines; lateral line with a strong arch in front, and with no accessory dorsal branch; anal spine obsolete; body indifferently dextral or sinistral (in some species at least). Scales ctenoid; dorsal fin beginning above pupil; gill rakers rather long and slender. This genus is intermediate between *Hippoglossoides* and *Paralichthys*, agreeing with the former in the insertion of the dorsal and in general appearance, and with the latter in the direction of the lateral line. Several species are now known. Some of them are dextral, and perhaps all of them are normally so, or perhaps, as in the case of *Xystreureys liolepis*, all are indifferently dextral or sinistral. (A diminutive of *Hippoglossus*, the halibut.)

- a. Mouth large, the maxillary extending to opposite posterior margin of eye, 2 in head; gill rakers numerous, 4 + 13; dorsal rays about 68; anal 53. STOMATA, 2986.  
 aa. Mouth moderate, the maxillary extending to opposite middle of pupil, about 2½ in head.  
 b. Dorsal rays about 66; anal 52; depth of body 2½ in length. MACROPS, 2987.  
 bb. Dorsal rays about 62; anal 48; depth of body 2½ in length; gill rakers 2 + 8 or 9. BOLLMANI, 2988.

## 2986. HIPPOGLOSSINA STOMATA, Eigenmann &amp; Eigenmann.

Head 2¾ to 3 in length; depth 2½ to 2¾. D. 67 to 70; A. 52 to 54; scales 80. Sinistral. Eye (not orbit) large, 5 in head; lower orbit slightly in advance of upper; interorbital a narrow ridge. Form, elongate elliptical, the profile depressed over the eye. Mouth large, maxillary extending to posterior margin of eye, as long as or longer than pectoral, 2 in head; lower jaw about 1¾ in head. Teeth small, uniserial; anterior nares of each side with long dermal flaps. Scales of left side all ctenoid, those of right side cycloid on anterior half or two-thirds of body; middle third of interorbital naked, anterior and posterior thirds scaled. Gill rakers 4 + 13 or 14. Dorsal beginning over middle of eye, anterior rays with but 1 or 2 scales, rest scaled to near tip, all but last 8 rays simple; anal similar to dorsal, with a strong procumbent spine; highest dorsal and anal rays about 3¼ in head; pectoral of colored side about 2 in head, that of blind side shorter; caudal double truncate, 5 to 5½ in length. Brown, strongly tinged in life with robin's-egg blue; numerous spots of light blue and light and dark brown; 5 pairs of large, dark-brown ocelli along dorsal and ventral parts of eyed side, the alternate ones longer and more conspicuous; fins colored like body, profusely mottled with light and dark; sinistral pectoral barred; a dark-brown spot above and below on caudal peduncle just in front of caudal, showing conspicuously on blind side. The eggs are probably pelagic; they are transparent, and measure 1.2 mm. in diameter; the single oil globule measures 0.16 mm. Coast of southern California; 2 specimens obtained in deep water off San Diego, November 7, 1889, both females, 1 with ripe eggs. (Eigenmann & Eigenmann.) (*στοματός*, large mouthed.)

*Hippoglossina stomata* EIGENMANN & EIGENMANN, Proc. Cal. Ac. Sci. 1890, 22, San Diego. (Coll. C. H. Eigenmann.)

## 2987. HIPPOGLOSSINA MACROPS, Steindachner.

Head  $2\frac{1}{2}$ ; depth  $2\frac{1}{2}$ . D. 66; A. 52; scales 75 to 80; upper orbit  $3\frac{1}{2}$  in head. Body elliptical, deeper than in related species; mouth moderate, the maxillary reaching to middle of eye; teeth small, sharp, uniserial; lower eye slightly in front of upper; eyes separated by a naked narrow ridge; nostrils close together, the anterior ending in a tube; horizontal limb of preopercle somewhat concave, the vertical convex. Dorsal beginning over middle of eye; pectoral of left side  $\frac{1}{2}$  head, much longer than maxillary, which is  $2\frac{3}{8}$  in head; interorbital space a narrow ridge; scales of left side all strongly ctenoid, those on blind side ciliated only on posterior third of body; no anal spine. Color brownish, with obscure darker blotches. Body sinistral (in the only specimen known). (Steindachner.) Pacific coast of Mexico. One specimen from Mazatlan; not seen by us. ( $\mu\alpha\kappa\rho\acute{o}\varsigma$ , large;  $\omega\psi$ , eye.)

*Hippoglossina macrops*, STEINDACHNER, Ichth. Beitr., v, 13, pl. 3, 1876, Mazatlan; JORDAN & GOSS, Review Flounders and Soles, 242, 1889.

## 2988. HIPPOGLOSSINA BOLLMANI, Gilbert.

Head 3 ( $3\frac{1}{2}$  to  $3\frac{3}{4}$ ) in length; depth  $2\frac{3}{8}$  to  $2\frac{1}{2}$  ( $3\frac{1}{4}$  to  $3\frac{1}{2}$ ); snout 5 in head. D. 60 to 63; A. 47 to 49; scales along lateral line 70 to 75. Body regularly elongate, elliptical; dorsal and ventral outlines equally curved; orbital rim entering anterior profile, which is equally curved before and behind eyes; greatest depth of body above pectorals. Mouth rather large, the maxillary reaching about to middle of pupil,  $2\frac{3}{8}$  to  $2\frac{3}{4}$  in head. Teeth equally developed on both sides, small and equal, uniserial. Premaxillary spine prominent. Interorbital space a narrow, sharp, naked ridge; eyes large, the lower slightly in advance of upper,  $3\frac{3}{8}$  to 4 in head. Gill rakers moderately long and slender, the longest 3 in length of ventral of eyed side; 2 + 8 or 9 developed, the last 2 much shorter. Scales small, firm, strongly ctenoid, those below pectoral much reduced, about 40 in a cross series; arch of lateral line strongly marked,  $2\frac{3}{8}$  to  $2\frac{1}{2}$  in straight part. Dorsal beginning above middle of pupil of upper eye, its anterior rays low, its longest rays  $2\frac{1}{2}$  in head; a strong antrorse spine before anal; pectoral of eyed side 2 in head, that of blind side  $2\frac{1}{2}$  to  $2\frac{3}{8}$  in head; ventrals subequal, each 6-rayed, 4 in head, extending more than  $\frac{1}{2}$  their length beyond anus; each is lateral, but that of eyed side nearest ridge of abdomen, and a little behind its fellow; last ray of left ventral joined to abdomen alongside of anal spine; caudal acute, its peduncle long. Color grayish brown, a row of 6 round, bluish spots, smaller than pupil, along base of dorsal, 4 similar spots along base of anal, and a few indistinct smaller ones on rest of body and head; body with 6 large black spots somewhat smaller than eye, these regularly 4 below dorsal and 2 above anal, the first of dorsal above arch of lateral line, the second above anterior third of straight part, the third at base of last rays and almost forming a cross bar with the 1 at base of anal rays. Dorsal, anal, and caudal dusky, with small whitish spots; a pale spot at base of last 4 dorsal and anal rays; a small black spot at base of outer caudal rays on peduncle; pectorals and ventrals dusky, but not spotted; right side immaculate. Length



7 inches. Pacific coast of Colombia. Numerous specimens were dredged at *Albatross Station* 2805, at a depth of 51½ fathoms. This species differs from *Hippoglossina stomata* in the gill rakers, which are shorter and fewer in number, and in the larger scales on sides. Scales in 16 rows between lateral line and back, instead of 21 or 22, as in *H. stomata*. Gill rakers somewhat shorter, 8 or 9 on anterior limb, 2 on upper limb. In *H. macrops* the gill rakers are slender, close set, 13 or 14 on anterior limb, 4 on vertical limb. In other respects of color, fin rays, and squamation agreeing perfectly with *H. stomata*. (Named for Charles Harvey Bollman.)

*Hippoglossina macrops*, JORDAN & BOLLMAN, Proc. U. S. Nat. Mus. 1889, 175; not of STEINDACHNER.

*Hippoglossina bollmani*, GILBERT, Proc. U. S. Nat. Mus. 1890, 122, *Albatross Station* 2805, southwest of Panama, in 51½ fathoms. (Type, No. 41143.)

#### 1022. LIGLOSSINA, Gilbert.

*Lioglossina*, GILBERT, Proc. U. S. Nat. Mus. 1890, 122 (*tetrophthalmus*).

This genus is allied to *Hippoglossina*, but its scales are all cycloid, the teeth are small, pointed, uniserial, and uniform, and the gill rakers short and thick. (λεῖδς, smooth; πλωβόα, tongue; for *Hippoglossina*.)

#### 2989. LIGLOSSINA TETROPTHALMA, Gilbert.

Head large, 3½ in length in a specimen 1 foot long. D. 76 to 83; A. 58 to 62; lateral line (pores) 97. Body of moderate height, the profile distinctly angulated above upper pupil, the snout projecting; length of caudal peduncle ½ its depth, its outlines diverging backward; depth of body 2½ in length; snout projecting beyond profile, bluntly rounded, the lower jaw included. Mouth large, the maxillary reaching nearly to vertical from posterior border of lower eye, 2½ in head; a blunt projecting process anteriorly from head of maxillary. Teeth small, pointed, in a single close-set series in each jaw, none of them enlarged; vomer toothless; lower eye slightly in advance of upper; vertical from front of upper falling midway between front of orbit and front of pupil of lower eye; vertical diameter of upper orbit but little more than ½ its longitudinal diameter, which is contained 3½ in head; interorbital space a blunt high ridge, entirely scaleless, its width ⅔ diameter of orbit. Anterior nostril of blind side with a very long flap, that of eyed side shorter; a well-marked cutaneous flap on lower eye above pupil. Gill rakers very large, broad, and strong, well toothed on inner edges, longest equaling diameter of pupil, the number on outer gill arch 10 or 11. First dorsal ray over anterior margin of pupil of upper eye, the fin not high, its highest ray 3 in head; anal similar; caudal sharply double truncate, the median rays produced; ventrals rounded, equal, barely reaching front of anal; no spine before anal fin; pectorals moderate, with 9 or 10 developed rays, ½ length of head; ventral 6. Scales rather small, growing distinctly larger posteriorly, everywhere smooth; head scaled, except snout, interorbital area, mandible, and part of maxillary, the latter with a patch of scales on posterior end of its expanded portion; on blind side an area around nostrils, and the greater part of exposed portion of preorbital, scaleless; fin rays of vertical fins,

all with bands of fine scales, those on caudal especially broad; lateral line with a broad arch in front, the cord of which is  $3\frac{1}{2}$  in straight portion. Color dusky brownish, with 2 conspicuous pairs of round black spots narrowly edged with gray, the anterior pair about  $\frac{1}{2}$  size of orbit, the posterior larger than pupil; the anterior pair under beginning of posterior third of dorsal, and about halfway between lateral line and dorsal and anal margins, respectively; the posterior pair nearer outline of body and about under the tenth before the last dorsal ray; vertical fins obscurely blotched with darker; ventral of eyed side with conspicuous black blotch margined with white, occupying the distal portion of its inner 2 rays; pectoral unmarked; membrane of gill cavity and peritoneum white. Two specimens, each about 12 inches long, from the Gulf of California, taken in 29 and 76 fathoms, at Albatross Stations 3014 and 3016. (Gilbert.) (*τετρα-*, four; *ὄφθαλμός*, eye, or eye-like spot.)

*Hippoglossina tetraphthalma*, GILBERT, Proc. U. S. Nat. Mus. 1890, 122, Gulf of California. (Coll. Dr. Gilbert.)

1023. XYSTREURYS, Jordan & Gilbert.

*Xystreurus*, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1880, 34 (*liolepis*).

Body broad, covered with small smooth scales. Teeth rather small, uniserial and bluntly conical, unequal; no canines; caudal fin subsessile, the caudal peduncle extremely short; skin of shoulder girdle with patches of cup-shaped scales; lateral line with a strong anterior arch, no accessory branch; vertebrae about 37; gill rakers short and thick. This genus is very close to *Hippoglossina*, differing chiefly in the subsessile caudal fin, the smooth scales, and the peculiar, short, thick gill rakers. The typical species, like some other Pacific coast flounders, is almost indifferently dextral or sinistral. (*ξίστρονον*, raker; *εἰρὺς*, wide, from the broad gill rakers.)

2990. XYSTREURYS LIOLEPIS, Jordan & Gilbert.

Head  $3\frac{1}{2}$ ; depth  $1\frac{1}{2}$ . D. 80; A. 62; scales 123. Vertebrae  $12 + 25 = 37$ . Body elliptical ovate, broad and compressed, its curves regular; the profile continuous with curve of back; mouth small, very oblique, the lower jaw included; maxillary reaching about to pupil,  $2\frac{1}{2}$  in head; eyes rather large,  $4\frac{1}{2}$  in head, separated by a very narrow, blunt scaly ridge; teeth small, conical, blunt, in a single row; those in lower jaw subequal, close set; those in upper jaw more distant, decreasing in size backward; teeth  $\begin{smallmatrix} 12 + 13 \\ 14 + 15 \end{smallmatrix}$  Gill rakers  $2 + 7$ , very short, broad, and strong, minutely serrate on inner margin, about 7 below angle, the longest scarcely  $\frac{1}{2}$  as long as the eye. Scales small, oblong, cycloid, the smaller accessory scales extremely numerous; lateral line without dorsal branch, with a broad curve above pectorals; branchial arches and skin of the shoulder girdle with small, cup-shaped, tubercular scales. Dorsal rather high, firm, low in front, beginning just in advance of middle of pupil, highest near the middle of the body; caudal peduncle very short and deep, its depth 4 times its length. Pectoral of eyed side falcate, usually much longer than head,

its length varying considerably. Caudal fin somewhat double truncate, with rounded angles, the middle rays being produced. Anterior nostril of blind side with a long flap. Color olive brown, mottled with darker, sometimes with very distinct round black blotches; vertical fins blotched with dark; pectoral of colored side with oblique bars. Length 15 inches. Southern California, rather common from Point Concepcion southward to San Diego. It is a very variable species, the coloration and the length of the pectoral fins having a wide range of variation. The body is indifferently dextral or sinistral. (*λεῖος*, smooth; *λεπίς*, scale.)

*Xystreurus liolepis*, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1880, 34, Santa Barbara; JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1880, 454; JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1881, 66; JORDAN & GOSSE, Review of Flounders and Soles, 243, 1889.  
*Paralichthys liolepis*, JORDAN & GILBERT, Synopsis, 825, 1883.

## 1024. PARALICHTHYS, Girard.

(BASTARD HALIBUTS.)

*Paralichthys*, GIRARD, U. S. Pac. R. R. Surv., x, 146, 1858 (*maculosus*—*californicus*).  
*Pseudorhombus*, BLEEKER, Comptes Rendus, Acad. Sci. Amsterd., XIII, 1862, 5, Notice sur quelques genres de la famille des Pleuronectidae (*polyspiloi*).  
*Uropsetta*, GILL, Proc. Ac. Nat. Sci. Phila. 1862, 336 (*californicus*—*maculosus*).  
*Chaenopsetta*, GILL, Proc. Ac. Nat. Sci. Phila. 1864, 218 (*ocellaris*—*dentatus*).

Eyes and color normally on the left side. Body oblong; mouth large, oblique; each jaw with a single row of usually slender and sharp teeth, which are more or less enlarged anteriorly; no teeth on vomer or palatines. Gill rakers slender. Scales small, weakly ctenoid or ciliated; lateral line simple, with a strong curve anteriorly. Dorsal fine beginning before the eye, its anterior rays not produced; both ventrals lateral; caudal fin double truncate, or double concave, its middle rays produced; no anal spine. Species numerous, found in all warm seas. This genus, as now restricted, contains a considerable number of species, inhabiting both coasts of America and the eastern and southern coasts of Asia. As indicated by the reduced number of vertebrae, the species range further southward than do those of the type of *Hippoglossoides*. (*παράλληλος*, parallel; *ἰχθύς*, fish.)

a. Gill rakers in large number, about 9 + 20.

b. Gill rakers as long as eye and very slender. D. 72; A. 55; depth  $2\frac{3}{4}$  in length. CALIFORNICUS, 2991.

bb. Gill rakers shorter, about  $\frac{2}{3}$  length of eye. D. 80; A. 61; depth  $2\frac{1}{4}$  in length. ESTUARIUS, 2992.

aa. Gill rakers in moderate number (5 + 11 to 6 + 21), rather long and slender.

c. Dorsal rays 70 to 75; anal rays 54 to 60.

d. Head small, lateral line  $4\frac{1}{2}$  in length; depth  $2\frac{1}{4}$ ; interorbital space rather broad and flatish,  $\frac{2}{3}$  diameter of eye; eyes small,  $5\frac{1}{2}$  in head; gill rakers rather short, 4 + 15, the longest about  $\frac{2}{3}$  eye. BRASILIENSIS, 2993.

dd. Head rather large,  $3\frac{1}{2}$  in length; depth 2 to  $2\frac{1}{4}$ ; eyes small.

e. Gill rakers 5 to 6 + 15 to 18; eyes wide apart. ADSPERSUS, 2994.

ee. Gill rakers 5 + 11; eyes close together. WOOLMANI, 2995.


cc. Dorsal rays 85 to 93; anal rays 67 to 73; gill rakers 5 + 15 or 16, long and slender, the longest  $\frac{2}{3}$  length of eye; body ovate, the depth about  $2\frac{1}{4}$  in length; head about  $3\frac{1}{4}$ . DENTATUS, 2996.



- aaa. Gill rakers few, shortish, wide set, the number 2 + 8 to 3 + 10.
- f. Body ovate, more or less compressed and opaque; depth about  $2\frac{1}{2}$  in length; no distinct, definitely placed ocelli; scales cycloid.
- g. Dorsal rays in large number (85 to 93, as in *P. dentatus*); anal rays 65 to 73; pores of the lateral line about 100; accessory scales few; gill rakers 2 + 10, lanceolate, dentate, wide set, and much shorter than the eye. LETHOSTIGMUS, 2907.
- gg. Dorsal rays in moderate number (70 to 80); anal rays 54 to 61.
- h. Scales very small, about 120 in lateral line; depth of body about  $\frac{1}{2}$  length; head  $3\frac{1}{2}$  in length; gill rakers roughly toothed, 3 + 9 in number. SQUAMILENTUS, 2998.
- hh. Scales moderate, 90 to 100 pores in the lateral line; interorbital width about equal to length of eye; dorsal rays 75 to 81; anal rays 59 to 61; gill rakers 2 or 3 + 9 or 10. Coloration, grayish brown with numerous (more or less distinct) whitish blotches, which are rarely obsolete; vertebrae 10 + 27 = 37. ALBIGUTTUS, 2999.
- ff. Body oblong, strongly compressed, semitranslucent; scales weakly ciliated; about 93 pores in lateral line. Coloration, light grayish, thickly mottled with darker; 4 large horizontally oblong, black ocelli, each surrounded by pinkish area; 1 just behind middle of the body, below the dorsal, 1 opposite this, above anal, and 2 similar smaller spots below last rays of dorsal and above last of anal; vertebrae 11 + 30 = 41. OBLONGUS, 3000.

2991. PARALICHTHYS CALIFORNICUS (Ayres).




(BASTARD HALIBUT; MONTEREY HALIBUT.)

Head  $3\frac{1}{2}$  to 4 $\frac{1}{2}$ ; depth  $2\frac{1}{2}$ . D. 70; A. 55; scales 100. Vertebrae 10 + 25 = 35. Body rather long and thickish; caudal peduncle long; head small; eye small, little wider than the broad, flattish interorbital space; maxillary as long as pectoral,  $\frac{1}{2}$  length of head, reaching beyond eye; teeth slender, sharp, rather long, the canines moderato. Scales small, finely ciliate, each scale surrounded by narrow accessory scales; scales on blind side similar; fins with ctenoid scales. Dorsal low, beginning over front of upper eye just past pectoral, pointed, reaching curve of lateral line,  $2\frac{1}{2}$  in head, that of blind side shorter and rounded behind; arch of lateral line  $3\frac{1}{2}$  or 4 in straight part. Gill rakers very long and slender, numerous, as long as eye, about 9 + 20; lower pharyngeals narrow, with small slender teeth. Anal spine small, concealed. Grayish brown, uniform, or mottled with blackish and pale, the head sometimes sprinkled with black dots; young brownish, with bluish spots. Coast of California, Tomales Bay to Cerros Island. This large flounder is one of the common food-fishes of the Pacific coast, where it takes the place occupied on the Atlantic side by *Paralichthys dentatus*. It reaches a length of 3 feet and a weight of 60 pounds. From its resemblance to the halibut, it usually goes by the name of bastard halibut. It is readily distinguished from the Atlantic members of the same genus by its fewer fin rays and by its more numerous gill rakers. As was first shown by Mr. Lockington, the small fish called *Paralichthys maculosus*, is simply the young of the larger fish, then called *Uropsetta californica*. Unlike other species of the genus, *Paralichthys californicus* is almost as frequently dextral as sinistral. (*californicus*, Californian.)





**IMAGE EVALUATION  
TEST TARGET (MT-3)**



**Photographic  
Sciences  
Corporation**

23 WEST MAIN STREET  
WEBSTER, N.Y. 14580  
(716) 872-4503

128  
132  
122  
120

10

- Pleuronectes maculosus*, GIRARD, Proc. Ac. Nat. Sci. Phila. 1854, 155, young, San Diego.  
*Paralichthys maculosus*, GIRARD, U. S. Pac. R. R. Surv., x, Fishes, 147, 1858, not *Rhombus maculosus*, CUVIER, also a species of *Paralichthys*; GÜNTHER, Cat., IV, 431, 1862; GILL, Proc. Ac. Nat. Sci. Phila. 1864, 197; LOCKINGTON, Rep. Com. Fisheries California 1878-79, 41; LOCKINGTON, Proc. U. S. Nat. Mus. 1879, 79; JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1880, 454; JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1931, 66; JORDAN, Nat. His. Aquat. Anim., 182, 1884.  
*Hippoglossus californicus*, AYRES, Proc. Cal. Ac. Nat. Sci. 1859, 29, and 1860, fig. 10, adult, San Francisco.  
*Pseudorhombus californicus*, GÜNTHER, Cat., IV, 426, 1862.  
*Uropsetta californica*, GILL, Proc. Ac. Nat. Sci. Phila. 1862, 330; GILL, Proc. Ac. Nat. Sci. Phila. 1864, 198.  
*Paralichthys californicus*, JORDAN & GILBERT, Synopsis, 821, 1885; JORDAN & GOSS, Review Flounders and Soles, 245, 1889.

2992. *PARALICHTHYS ESTUARIUS*, Gilbert & Seafield.

Head  $3\frac{2}{3}$ ; depth  $2\frac{1}{2}$ ; eye  $5\frac{1}{2}$ ; interorbital space flat, 12 in head,  $\frac{1}{2}$  diameter of eye; maxillary 2 in head, equal to pectoral fin; gill rakers 9 + 20, the longest  $\frac{2}{3}$  length of eye. D. 72 to 83; A. 58 to 64. (In 7 specimens examined the rays are: Dorsal 72, 79, 81, 81, 82, 83, 83; anal 58, 60, 60, 62, 63, 63, 64.) Vertebrae 10 + 28; scales weakly ciliated, with small accessory scales, 105 in the lateral line; length of the arch contained 4 times in straight part of lateral line, 2 in head; height of arch  $4\frac{1}{2}$  in head. Four of the 7 specimens are sinistral. Color pale chocolate brown. Specimens small, 6 to 9 inches in length. Taken at Shoal Point, at mouth of the Colorado River, Mexico, by the United States Fish Commission steamer *Albatross*. This species is distinguished from other members of the genus by its numerous fin rays and many gill rakers. It is nearest related to *Paralichthys californicus*. (*estuarium*, pertaining to the river mouth.)

*Paralichthys estuarium*, GILBERT & SCOFIELD, Proc. U. S. Nat. Mus. 1897, 499, pl. xxix, Gulf of California, at mouth of Colorado River, Sonora. (Type, No. 48128. Coll. C. H. Gilbert.)

2993. *PARALICHTHYS BRASILIENSIS* (Ranzani).

Head  $4\frac{1}{2}$ ; depth  $2\frac{1}{2}$ . D. 70 to 75; A. 54 to 60; scales not very small, about 100 in course of lateral line; interorbital space rather broad and flattish,  $\frac{2}{3}$  diameter of eye; eyes small,  $5\frac{1}{2}$  in head; gill rakers rather short, 4 + 15, the longest about  $\frac{2}{3}$  eye; pectoral  $1\frac{1}{2}$  in head; curve of lateral line high and short, 4 in straight part, its height  $1\frac{1}{2}$  in its length; mouth moderate, the maxillary  $2\frac{1}{2}$  in head; teeth rather few, the anterior canines large. Color dark brown, more or less mottled and spotted with paler. South America; said to range northward to Guatemala. Here described from numerous specimens from Rio Janeiro and from Maldonado, in the Museum of Comparative Zoology. The locality "Guatemala" given by Günther seems to be somewhat doubtful, and the species may not occur in West Indian waters at all. (*brasiliensis*, living in Brazil.)

- Hippoglossus brasiliensis*, RANZANI, Nov. Spec. Plac., 10, tab. 3, 1840, Brazil.  
*Platessa orbignyana*, VALENCIENNES, D'Orbigny Voy. S. Amer. Mérid. Poiss., pt. 5, pl. 16, fig. 1, 1847.  
*Rhombus aramaca*, CASTELNAU, Anlm. nouv. ou rares, Poiss., 78, pl. 40, fig. 3; not of CUVIER.  
*Pseudorhombus vorax*, GÜNTHER, Cat., IV, 420, 1862, South America.  
*Pseudorhombus brasiliensis*, GÜNTHER, Fishes Contr. Amer., 473, 1860.  
*Paralichthys brasiliensis*, JORDAN & GOSS, Review Flounders and Soles, 246, 1889.



## 2094. PARALICHTHYS ADSPERSUE (Steindachner).

Head  $3\frac{1}{2}$ ; depth  $2\frac{1}{2}$ . D. 75; A. 58; scales 106; eye 6 in head; interorbital  $\frac{3}{4}$  vertical diameter of eye; maxillary  $2\frac{1}{2}$ ; mandible  $1\frac{1}{2}$ ; pectoral 2; caudal  $1\frac{3}{4}$ . Body moderately elongate and compressed; mouth large, the maxillary reaching a little past eye; teeth large, sharp, and slightly recurved, larger in front of jaws; snout very slightly produced; interorbital moderately wide, its posterior half with scales; anterior nostril with a flap which reaches to middle of posterior nostril; gill rakers 3 to 6 + 15 to 17, hardly as long as eye. Snout and mandible naked; end of maxillary and rest of head with scales; the rays of all the fins with small scales; the membrane naked; each scale on body with a row of accessory scales around its posterior edge; scales cycloid, the accessory scales giving the fish a rough feeling; curve of lateral line nearly 5 in the straight part, pectoral reaching slightly past curve of lateral line, its tip pointed; pectoral of blind side shorter, not reaching to end of curve, its tip blunt; origin of dorsal over anterior edge of upper eye, bending slightly toward the blind side; caudal double lunate. Color brownish gray, thickly mottled with many larger and smaller spots, points, and rings; side with 3 or 4 larger spots of irregular form and ocellated with paler.

Specimens taken by Dr. Jordan at Mazatlan are described as follows: "Head  $3\frac{1}{2}$ ; depth about 2 in length of body. D. 73 (70 to 76); A. 57 (53 to 60); P. 12; V. 6. Scales on lateral line about  $106 + 8$  with 35 dorsally and 36 ventrally. Flesh firm. Body oblong, moderately compressed; mouth large, oblique, the mandible very heavy, slightly projecting; 4 canine teeth on each side of lower jaw in adult specimens, 8 in young, the 2 anterior teeth long; anterior teeth of upper jaw strong, but smaller than those in the lower jaw; the lateral teeth very small and close set. Eye small, shorter than snout, about 7 (6 to 8) in length of head; interorbital area smooth, flattish,  $\frac{3}{4}$  width of eye. Scales cycloid, small anteriorly and larger posteriorly; lateral line strongly arched anteriorly, arch about  $3\frac{1}{2}$  in straight part. Gill rakers of medium length, broad, retrorse-serrate on inner side, longest about  $\frac{3}{4}$  length of eye, from 4 + 13 to 5 + 14 in number, counted in 8 specimens; pectoral fin about as long as mandible, slightly more than  $\frac{1}{2}$  length of head. Dorsal low, anterior origin opposite anterior margin of eye; caudal barely double concave; caudal peduncle very strong; anal spine obsolete; ventral fins small, inserted symmetrically; fins all scaly. Color: Large specimens are dark brown, with blotches on fins; small specimens are covered with pearly white and very dark brown blotches; the brown blotches almost circular, larger and with less definite outlines near the center of the body, very dark and distinct on caudal. Seven specimens were taken by the Hopkins Expedition in the estuary at Mazatlan, where they reach a length of 44 cm. Several specimens were also taken at La Paz. These specimens seem to be identical with *Paralichthys adspersus*. The original types have on an average more gill rakers than we find in our Mazatlan specimens, but this character is subject to variation, and no other distinction appears. In one of Dr. Steindachner's types from Callao (No. 11,417, Mus. Comp. Zool.) we find the gill rakers longer, 6 + 17; depth  $2\frac{1}{2}$  in length; D. 67; A. 51; scales 120;

arch of lateral line barely twice as long as high, nearly 5 in straight part; maxillary  $2\frac{1}{2}$  in head. Mr. Garman has kindly examined for us 6 other specimens, with the following results:

"*Paralichthys adpersus* from Callao, has gill rakers—

" $\frac{1}{7}$  as long as eye;

" $\frac{5}{8}$  about  $\frac{2}{3}$  as long as the eye.

" $\frac{1}{8}$  nearly as long as the eye.

" $\frac{3}{4}$  about  $\frac{2}{3}$  as long as the eye.

" $\frac{5}{8}$  about  $\frac{2}{3}$  as long as the eye.

" $\frac{1}{7}$  near  $\frac{2}{3}$  as long as eye."\*

We are now disposed to regard these Mazatlan specimens as identical with *Paralichthys adpersus*, the range of variation in the number of gill rakers in the latter probably including the former. Pacific coast of tropical America, from Gulf of California to the coast of Peru; everywhere abundant and very variable. (*adpersus*, covered with spots.)

*Pseudorhombus adpersus*, STEINDACHNER, Ichthyol. Notizen, v, 9, pl. 2, 1867, Chinchas Islands.

*Paralichthys adpersus*, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1882, 370; JORDAN & GILBERT, Bull. U. S. Fish. Comm. 1882, 108 and 111; JORDAN, Cat. Fish N. A., 133, 1885; JORDAN & GOSS, Review Flounders and Soles, 246, 1889; JORDAN & WILLIAMS, Proc. Cal. Ac. Sci. 1895, 503.

#### 2995. PARALICHTHYS WOOLMANI, Jordan & Williams.

Head  $3\frac{1}{2}$ ; depth about 2; gill rakers 5+11. D. 74; A. 57; P. 12; V. 6; scales 100. Flesh firm; body oblong; mouth large, mandible heavy, not projecting; about 8 teeth on each side of lower jaw, the anterior ones long and slender; teeth in upper jaw smaller than those in lower jaw, the lateral teeth very small and close set. Eye small,  $5\frac{1}{2}$  in length of head; interorbital area moderately prominent, narrow, about  $\frac{2}{3}$  length of eye. Scales cycloid, small anteriorly and increasing in size posteriorly, covering head and fins; lateral line greatly arched anteriorly, arch about  $3\frac{1}{2}$  times in length of straight portion. Gill rakers slender, the longest about  $\frac{1}{2}$  in length of eye. Pectoral and ventral fins small; pectoral about  $\frac{1}{2}$  in length of head; origin of dorsal opposite anterior margin of eye; caudal ending in an obtuse angle, not double concave; caudal peduncle strong; anal spine obsolete. Body and fins blotched with deep brown and pearly white and specked with very dark brown, blotches more definite on median fins and especially on caudal where there are 3 indefinite lines of blotches crossing the fin. Galapagos Islands. One specimen taken by the *Albatross* in 1890, which was at first identified as *Paralichthys adpersus*, from which species it differs but little except in the number of gill rakers. (Named for Mr. Albert Jefferson Woolman, of Duluth, Minnesota, in recognition of his work on the fishes of Mexico and Florida.)

*Paralichthys woolmani*, JORDAN & WILLIAMS, Proc. U. S. Nat. Mus. 1896, 457, Galapagos Islands. (Type, No. 47575. Coll. *Albatross*.)

\* Garman, in lit., May 3, 1895.

## 2906. PARALICHTHYS DENTATUS (Linnaeus).

(SUMMER FLOUNDER.)

Head  $3\frac{1}{2}$  to 4; depth  $2\frac{2}{3}$ ; eye 6 in head; maxillary 2; pectoral  $2\frac{1}{2}$ ; ventral 3; caudal peduncle 4; caudal  $1\frac{1}{4}$ . D. 86 to 91; A. 65 to 71; lateral line 108 (tubes). Curve of lateral line  $3\frac{2}{3}$  to  $4\frac{1}{2}$  in straight portion; body ovate; maxillary about  $\frac{1}{2}$  head, reaching past posterior margin of eye; mouth large, oblique, the gape curved; canines large, conical, wide set; gill rakers comparatively long and slender, longest  $\frac{2}{3}$  eye, 5 + 15 to 6 + 18 in number; interorbital area a rather flattish ridge, in the adult about equal to vertical diameter of eye, narrower in the young, forming a bony ridge; scales cycloid, each with numerous small accessory scales; vertebrae 11 + 30 = 41. Color in life, light olive brown; adults with very numerous small white spots on body and vertical fins; sometimes a series of larger white spots along bases of dorsal and anal fins; about 14 ocellated dark spots on sides, these sometimes little conspicuous, but always present; a series of 4 or 5 along base of dorsal, and 3 or 4 along base of anal, those of the 2 series opposite, and forming pairs; 2 pairs of smaller less distinct spots midway between these basal series and lateral line anteriorly, with a small one on lateral line in the center between them; a large distinct spot on lateral line behind middle of straight portion; fins without the round dark blotches. Atlantic coast of United States, from Cape Cod to Florida; the common flounder of the coasts of the Northern States, its range apparently not extending much south of Charleston. Of the species found in that region it is the most important from a commercial point of view. It reaches a length of about 3 feet and a weight of about 15 pounds. It has been confounded by nearly all writers with the more southern species now called *P. lethostigmus*, from which it is best distinguished by its much greater number of gill rakers and by its mottled coloration. On account of this confusion it is impossible wholly to disentangle its synonymy from that of *P. lethostigmus*. So far as the proper nomenclature of the two is concerned, this confusion makes little difference. There is no doubt that this is the original *Pleuronectes dentatus* of Linnaeus, as the original Linnæan type is still preserved in London. This has been examined by Dr. Bean and its identity with the present species fully established. It seems also certain that this is the *Platessa ocellaris* of De Kay, who properly distinguishes his *ocellaris* from his *oblonga*, the latter being *P. lethostigmus*. A little doubt must be attached to the *P. melanogaster* of Mitchill, very scantily described from a doubled (black-bellied) example of this species or of *P. lethostigmus*. As the former species is much more common about New York than the latter it is probable that Mitchill's fish belonged to it. We have also received a doubled example from New York corresponding exactly to Mitchill's description. We may therefore regard the name *melanogaster* as a synonym of *dentatus*. The differences in the gill rakers of these species were first noticed by Jordan & Gilbert in 1883. These authors erroneously referred all these synonyms to the species with the few gill rakers and described the present one as new under the name *Paralichthys ophryas*. The discovery of the Linnæan type of *Pleuronectes dentatus* has rendered a reconsideration of this matter

necessary, and it is evident that to the "*P. ophryus*" belong also the prior names *dentatus*, *melanogaster*, and *ocellaris*. (*dentatus*, toothed.)

*Pleuronectes dentatus*, LINNÆUS, Syst. Nat., Ed. XII, 1, 458, 1766, and of numerous copyists; MITCHILL, Trans. Lit. and Phil. Soc. N. Y. 1815, 390.

*Pleuronectes melanogaster*, MITCHILL, Trans. Lit. and Phil. Soc. N. Y. 1815, 390, New Yprk; doubled example.

*Platessa ocellaris*, DE KAY, N. Y. Fauna: Fishes, 300, pl. 47, fig. 152, 1842, New York.

*Paralichthys ophryus*, JORDAN & GILBERT, Synopsis, 822, 1883, Charleston.

*Platessa dentata*, STORER, Rept. Fish. Mass., 143, 1839.

*Pseudorhombus dentatus*, GOODE & BEAN, Proc. U. S. Nat. Mus. 1879, 123.

*Paralichthys dentatus*, GOODE, Nat. Hist. Aquat. Anim., 178, 1884, detailed account; includes *P. lethostigma*; JORDAN, Cat. Fish. N. A., 134, 1885; JORDAN & GOSS, Review Flounders and Soles, 246, 1889.

*Pseudorhombus ocellaris*, GÜNTHER, Cat., IV, 430, 1862; JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1878, 370.

*Chaenopsetta ocellaris*, GILL, Proc. Ac. Nat. Sci. Phila. 1864, 218.

*Paralichthys ocellaris*, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1882, 617.

#### 2997. PARALICHTHYS LETHOSTIGMUS, Jordan & Gilbert.

(SOUTHERN FLOUNDER.)

Head  $3\frac{3}{4}$ ; depth  $2\frac{1}{2}$ . D. 85 to 92; A. 65 to 73; pores about 100. Body ovate, more or less compressed and opaque; no distinct, definitely placed ocelli; scales cycloid. Mouth wide, oblique, the mandible very heavy and much projecting; 8 to 10 teeth on each side of the lower jaw, the 2 anterior teeth very long; anterior teeth of upper jaw strong, but smaller than those in the lower jaw; the lateral teeth very small and close set; eyes small, shorter than the snout, about 6 in head; interorbital space in adult broad, flattish, and scaly, as wide as length of eye. Accessory scales few; gill rakers 2 + 10, lanceolate, dentate, wide set, and much shorter than the eye; caudal peduncle rather long; length of arch of lateral line nearly  $\frac{1}{2}$  that of straight part. Color dusky olive, darker than in *P. dentatus*, and with very few darker mottlings or spots. This species is the common large flounder of the South Atlantic and Gulf coasts of the United States, ranging as far north as New York. It very closely resembles *Paralichthys dentatus*, with which it has been repeatedly confounded. It is, however, sharply distinguished by the character of the gill rakers. It is also always darker in color, and almost uniform, while *P. dentatus* is usually profusely spotted. Its only tenable name is the recent one, *Paralichthys lethostigma*. South Atlantic and Gulf coasts of United States, north to New York. (*λίσθη*, forgetfulness; *στίγμα*, spot, from the absence of spots.)

*Platessa oblonga*, DE KAY, New York Fauna: Fishes, 299, pl. 48, fig. 150, 1842, New York, not *Pleuronectes oblongus*, MITCHILL; STORER, Syn. Fish. N. A., 477, 1846.

*Paralichthys lethostigma*, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1884, 237, Jacksonville, Florida; JORDAN & GOSS, Review Flounders and Soles, 247, 1889.

*Pseudorhombus oblongus*, GÜNTHER, Cat., IV, 426, 1862.

*Chaenopsetta dentata*, GILL, Proc. Ac. Nat. Sci. Phila. 1864, 218.

*Pseudorhombus dentatus*, GOODE, Proc. U. S. Nat. Mus. 1879, 110; GOODE & BEAN, Proc. U. S. Nat. Mus. 1879, 123.

*Paralichthys dentatus*, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1882, 302; JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1882, 617; BEAN, Cat. Coll. Fish. Proc. U. S. Nat. Mus. 1883, 45; JORDAN & GILBERT, Synopsis, 822, 1883.

## 2998. PARALICHTHYS SQUAMILENTUS, Jordan &amp; Gilbert.

Head  $3\frac{1}{2}$ ; depth 2. D. 78; A. 59; scales 123 (pores). Body deep, strongly compressed; caudal peduncle very short; profile angulated at front of upper eye. Head wide, the eyes large, wide apart. Mouth very large, oblique, the broad maxillary reaching well beyond pupil, its length more than  $\frac{1}{2}$  the head. Lower jaw projecting; mandible with a sharp compressed knob at symphysis; teeth few, unequal, in a single row, about 8 in each jaw canine-like, the 2 in front of lower jaw longest; lateral teeth of upper jaw minute. Interorbital space a narrow scaleless bony ridge, slightly concave anteriorly, scarcely  $\frac{1}{2}$  diameter of pupil. Scales very small, smooth, adherent; curve of lateral line  $4\frac{1}{2}$  in straight part; snout, jaws, and preopercle naked. Gill rakers short,  $3+9$  in number, triangular, roughly toothed, little higher than wide, the longest nearly  $\frac{1}{2}$  eye. Dorsal beginning over front of eye, the anterior rays  $4\frac{1}{2}$  in head; pectoral short, shorter than maxillary; anal spine weak; caudal double rounded. Brownish; body and fins spotted with darker; caudal mottled with white; pectorals banded, with dark spots. South Atlantic and Gulf coasts of United States. This species is very close to *Paralichthys albiguttus*, from which it differs chiefly in the small scales. It seems to be rather rare. Besides the original types from Pensacola, another referred to the same species is in the National Museum from Charleston. (*squamilentus*, scaly.)

*Paralichthys squamilentus*, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1882, 303, Pensacola (Type, No. 30862); JORDAN & GILBERT, Synopsis, 823, 1883; BEAN, Cat. Coll. Fish, U. S. Nat. Mus. 1883, 45; JORDAN & GOSS, Review Flounders and Soles, 248, 1889.

## 2999. PARALICHTHYS ALBIGUTTUS, Jordan &amp; Gilbert.

(GULF FLOUNDER.)

Head  $3\frac{1}{2}$ ; depth  $2\frac{1}{2}$ . D. 72 to 80; A. 59 to 61; scales 9 to 100 (pores); eye 6 or 7 in head; maxillary  $1\frac{1}{2}$ ; pectoral  $2\frac{1}{2}$ ; ventral 3; caudal  $1\frac{1}{2}$ ; curve of lateral line 3 in straight part. Body moderately elongate-elliptical; mouth large, the maxillary reaching past eye; jaws subequal; teeth strong, slender, and curved, about 7 on side of lower jaw, 4 or 5 moderate canines in front of upper jaw, the lateral teeth being minute, close set; interorbital space  $\frac{2}{3}$  length of eye, the upper ridge rather prominent behind upper eye, scaled posteriorly; mandibles naked; a small patch of scales on maxillary; gill rakers broad and toothed behind, the longest  $2\frac{1}{2}$  in eye,  $3+10$  in number. Fins low; anterior rays of dorsal not elevated nor exerted, the longest rays behind the middle,  $2\frac{1}{2}$  in head; pectoral not reaching to end of curve; caudal double lunate. Scales moderate, cycloid, covered with epidermis which bears small flaps about the borders of many of the scales. Dark olive, mottled with dusky, and marked by numerous more or less distinct pale spots, which are sometimes obsolete; three dark spots, bordered with white, sometimes present, particularly in the young, 1 on lateral line posteriorly and 1 above and below anterior end of straight part of lateral line. Vertebrae  $10+27=37$ . South Atlantic and Gulf coasts of the United States. This species is common on the South Atlantic and Gulf coasts. It has the few gill rakers of *P. lethostigmus*, the mottled coloration of *P. dentatus*, while from each it is distinguished by its smaller number

of dorsal and anal rays. In the number of its vertebrae it agrees with *P. lethostigmus*. It seems to reach a smaller size than either of these species. Here described from a specimen, 16 inches in length, collected at Cedar Key, Florida. (*albus*, white; *gutta*, spot.)

*Pseudorhombus dentatus*, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1878, 370; not of LINNÆUS.

*Paralichthys albigutta*, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1882, 302, Pensacola (Type, No. 30818. Coll. Dr. Jordan); JORDAN & GILBERT, Synopsis, 823, 1883; JORDAN & SWAIN, Proc. U. S. Nat. Mus. 1884, 233; JORDAN & GOSS, Review Flounders and Soles, 248, 1889.

### 3000. PARALICHTHYS OBLONGUS (Mitchill).

(FOUR-SPOTTED FLOUNDER.)

Head 4; depth  $2\frac{1}{2}$ . D. 72; A. 60; scales 93. Body comparatively elongate, strongly compressed. Eyes large, nearly 4 in head, separated by a prominent, narrow, sharp ridge. Upper jaw with very numerous small, close-set teeth laterally, and 4 or 5 canines in front; the lateral teeth abruptly smaller than the anterior; each side of lower jaw with 7 to 10 teeth. Chin prominent. Maxillary narrow, reaching past middle of pupil,  $2\frac{1}{2}$  in length of head. Gape curved; gill rakers short and toothed behind, 2 + 8. Scales weakly ctenoid or cycloid. Dorsal low, beginning over front of eye, some of the anterior rays exerted, but not elongate, the longest rays behind middle of fin, not quite  $\frac{1}{2}$  head; caudal  $1\frac{1}{2}$  in head; pectoral  $1\frac{3}{8}$ ; anal spine obsolete. Grayish, thickly mottled with darker and somewhat translucent; 4 large, horizontally oblong, black ocelli, each surrounded by a pinkish area, 1 just behind middle of the body below the dorsal, 1 opposite this above anal, 2 similar smaller spots below last rays of dorsal and above last of anal. Coasts of New England and New York. This species is rather common on the coast of Cape Cod and the neighboring islands, but it has been rarely noticed elsewhere. The limits of its range are not yet definitely known. It is a very strongly marked species. Its translucency of coloration indicates that it lives in deeper water than the other species of the genus. Here described from specimen from Woods Hole.

Another specimen in our collection from Woods Hole, Massachusetts, referred to this species, shows the following characters: Brownish, somewhat mottled, without traces of ocelli (possibly faded); fins similar. Body rather elongate, slenderer than in other species and more compressed; mouth rather large, oblique, the lower jaw not projecting, the maxillary  $2\frac{1}{2}$  in head, reaching to opposite posterior border of pupil; about 12 teeth on each side of lower jaw, the anterior rather long, about equal to anterior teeth of upper jaw; lateral teeth of upper jaw becoming gradually smaller posteriorly, much larger, less numerous, and more widely set than in other species of this genus. Eyes large, longer than snout, 4 to  $4\frac{1}{2}$  in head, separated by a narrow, elevated, bony ridge, narrower than pupil, anteriorly scaleless, and curved behind the upper eye posteriorly. Scales moderate, cycloid, rather thin; curve of lateral line  $4\frac{1}{2}$  in straight part. Gill rakers 2 + 8 in number, rather long and slender, about  $4\frac{1}{2}$  in maxillary. Dorsal beginning above middle of eye, its anterior rays not longer than

others, the middle rays a little longer than longest of anal, which are about  $\frac{1}{2}$  head; caudal as long as head; anal spine obsolete; ventrals small; pectoral  $1\frac{1}{2}$  in head. Head  $4\frac{1}{2}$ ; depth  $2\frac{3}{4}$ . D. 77; A. 63; scales 90. Length about 14 inches. (*oblongus*, oblong.)

*Pleuronectes oblongus*, MITCHILL, Trans. Lit. and Phil. Soc., 1, 1815, 391, New York.

*Platessa quadrocellata*, STORER, Proc. Boston Soc. Nat. Hist. 1847, 242; STORER, Hist. Fish. Mass., 397, pl. 31, fig. 3, Provincetown.

*Chenopsetta oblonga*, GILL, Proc. Ac. Nat. Sci. Phila. 1864, 218.

*Paralichthys oblongus*, GOODE, Proc. U. S. Nat. Mus. 1880, 472; JORDAN & GILBERT, Synopsis, 824, 1883; JORDAN & GOSS, Review Flounders and Soles, 249, pl. 8, 1889; GOODE & BEAN, Oceanic Ichthyology, 436, 1896.

### 1025. RAMULARIA, Jordan & Evermann.

*Ramularia*, JORDAN & EVERMANN, new genus (*dendriticus*).

This genus is close to *Ancylosetta*, differing mainly in the structure of the lateral line, the tubes of which are borne by series of smaller, concealed cycloid scales, the free edges of which are notched to the opening of the pore; these scales are concealed in the skin, and from the pores proceed backward membranaceous tubes which ramify over the bases of contiguous scales. Dorsal scarcely elevated in front; left ventral much produced. Body broad ovate, sinistral, with very rough scales. Gill rakers few, very broad. (*ramulus*, a branchlet, from the tubes of the lateral line.)

#### 3001. RAMULARIA DENDRITICA (Gilbert).

Head  $3\frac{1}{2}$  in length; depth  $1\frac{1}{2}$ . D. 84; A. 63; scales 100; 36 scales in a series upward and backward from lateral line. Body very broad, its depth  $1\frac{1}{2}$  in length, the two outlines equally curved; profile not very strongly angulated in front of upper eye. Lower eye slightly in advance of upper; interorbital space a rather broad, convex, scaly ridge, about  $\frac{1}{2}$  upper eye, which is contained about 5 times in head and is equal to snout. A blunt spine on snout on head of maxillary. Nostril openings very broad, without tube, the anterior with a narrow flap. Mouth moderate, very oblique, the gape curved, the maxillary reaching slightly beyond vertical from middle of lower eye, 3 in head. Teeth in a single, rather close-set series in each jaw, strong, conical, directed very obliquely inward, becoming gradually larger toward front of jaw, but not canine-like. Gill rakers very short, barely movable, as broad as long, strongly toothed, 6 on anterior limb. Dorsal beginning over middle of upper eye, the anterior rays partly free toward tips, but little, if any, elevated above those that follow, the first  $2\frac{1}{2}$  in head; dorsal highest in its posterior third, the longest ray  $2\frac{1}{2}$  in head; anal similar, the rays of posterior third of each fin slightly forked at tip; caudal peduncle deep and short, its depth about  $\frac{1}{2}$  head, its length  $\frac{1}{2}$  its depth; caudal rounded, almost double truncate; ventrals with narrow bases, the left one slightly in advance of the right; fin greatly produced, reaching far beyond front of anal, a trifle shorter than head; left pectoral  $1\frac{1}{2}$  in head. Scales very strongly ctenoid, the edge spinous, the entire exposed portion rough; width of anterior arch of lateral line  $3\frac{1}{2}$  in straight portion; tubes of lateral line borne by a series of smaller con-

cealed cycloid scales, the free edges of which are notched to the opening of the pore; these scales entirely covered by the integument, and from the pores there proceed backward membranaceous tubes, ramifying over the bases of contiguous scales; this is true also of lateral line of blind side; eyed side entirely sealed except snout and mandible. Vertical fins covered with thick skin, each ray accompanied by 1 or 2 series of ctenoid scales; left ventral also scaled. Color olive brown, with 3 large black ocellated spots larger than orbit, the posterior one on lateral line in front of caudal peduncle, the 2 anterior under middle of dorsal, halfway between lateral line and dorsal and anal outlines, respectively; each spot with a light center; distal portion of vertical fins more or less brown on right side. (Gilbert.) Gulf of California. A single specimen, 13 inches long, from Albatross Station 3022, in 11 fathoms. (*ἀενδρίτιου*, like a tree, branched; *δένδρον*, tree.)

*Ancylopsetta dendritica*, GILBERT, Proc. U. S. Nat. Mus. 1890, 121, Gulf of California at Albatross Station 3022, in 11 fathoms.

#### 1026. ANCYLOPSETTA, Gill.

*Ancylopsetta*, GILL, Proc. Ac. Nat. Sci. Phila. 1864, 224 (*quadrocellata*).

Body sinistral, broadly ovate, the depth more than  $\frac{1}{2}$  length; mouth moderate; teeth uniserial, unequal, some of the anterior enlarged; caudal fin with a very short peduncle; scales very strongly ctenoid on both sides of the body; anterior rays of dorsal notably exerted, the rays of the anterior part of the fin elongate, thus forming a distinct lobe; gill membranes considerably united; gill rakers short and broad, with rough teeth; left ventral produced; vertebrae about 35. This genus is very close to *Paralichthys*, differing in the sessile caudal fin, the short gill rakers, the rough scales, and in the prolongation of the anterior rays of the dorsal fin. (*ἀγκυλος*, hook; *ψῆττα*, turbot.)

#### 3002. ANCYLOPSETTA QUADROCELLATA, Gill.

Head  $3\frac{1}{2}$  to  $3\frac{3}{4}$ ; depth  $1\frac{1}{2}$ . D. 70 to 76; A. 57 to 59; pores in lateral line 83 to 90; vertical series of scales 70; fourth or fifth dorsal ray longest, nearly  $\frac{3}{4}$  length of head. Caudal  $1\frac{1}{2}$  in head; ventral of colored side  $1\frac{3}{4}$ . Body oval, compressed, very deep; an abrupt angle above eye; mouth very small, the maxillary reaching to below middle of orbit,  $2\frac{1}{2}$  in length of head; teeth comparatively small, about 14 on each side of lower jaw; no strongly differentiated canines in either jaw. Eyes moderate, separated by a very narrow, sharp, scaly ridge; gill rakers very short, thick, few in number, 2 + 6 or 7, the longest less than  $\frac{1}{2}$  diameter of pupil; scales rather small, very strongly ctenoid, those on blind side also rough; curve of lateral line rather low; tubes of lateral line simple; dorsal beginning in front of pupil, its anterior rays long and filiform, much exerted; caudal short and rounded,  $1\frac{1}{2}$  in head; ventral fin of colored side rather long, as long as pectoral,  $\frac{1}{2}$  length of head; anal spine wanting. Brownish olive, with 4 large, oblong, ocellated spots, the first above the arch of the lateral line; the 3 posterior forming an isosceles triangle, the hindmost



being on the lateral line; the ocellated spots are frequently furnished with a bright white center, and the sides and vertical fins have often a few scattered white spots; a small, indistinct, dark spot on middle of each eighth or tenth ray of dorsal and anal. Vertebrae  $9 + 26 = 35$ . South Atlantic and Gulf coasts of the United States; not rare; a very handsome species. (*quadrocellatus*, having 4 ocelli.)

*Ancylopsetta quadrocellata*, GILL, Proc. Ac. Nat. Sci. Phila. 1864, 224; not *Platesa quadrocellata*, STORER; JORDAN & GOSS, Review Flounders and Soles, 250, 1889.

*Paralichthys ommatus*, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1882, 616, Charleston; JORDAN & GILBERT, Synopsis, 824, 1883; JORDAN & SWAIN, Proc. U. S. Nat. Mus. 1884, 234; JORDAN & GOSS, Review Flounders and Soles, 250, 1889.

*Pseudorhombus quadrocellatus*, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1878, 370.

### 1027. NOTOSEMA, Goode & Bean.

*Notosema*, GOODE & BEAN, Bull. Mus. Comp. Zool., x, No. 5, 192, 1883 (*dilectum*).

Body sinistral, elliptical in form, the caudal fin pedunculate. Mouth moderate, beneath the central axis of the body. Eyes large, close together, the upper one nearly encroaching upon the profile, the lower slightly in advance of the upper. Teeth in a single series in the jaws, about equally developed on each side, largest in front, absent on vomer and palatines. Pectoral fins somewhat unequal, that upon the blind side  $\frac{3}{4}$  as large as the other; dorsal fin commencing slightly behind anterior margin of upper eye, the first 8 rays separated into a distinct subdivision of the fin, several of them being prolonged; caudal rounded, sinistral; ventral much elongated. Scales small, ctenoid on colored side of body; lateral line prominent, strongly arched, alike on both sides, the tubes simple. Gill rakers moderately numerous, rather stout, subtriangular, pectinate posteriorly. Pseudobranchiae well developed. Vertebrae 35. This genus is scarcely distinct from *Ancylopsetta*, the body more elongate, the dorsal and ventral rays more produced. (*νῶτος*, back; *σῆμα*, banner.)

### 3003. NOTOSEMA DILECTUM (Goode & Bean).

Head  $3\frac{1}{2}$ ; depth 2. D. 68; V. 6. A. 54 to 56; scales 48 (pores) on straight part of lateral line; width of interorbital area almost imperceptible; mandible reaching to middle of pupil of lower eye, its length 2 in head; upper jaw  $2\frac{1}{2}$  times length of head. Origin of dorsal over anterior margin of eye, second and third rays the longest, which are 2 in greatest depth of body; anal beginning close to vent, its posterior rays longest; caudal pedunculate, double truncate; pectoral of eyed side subtriangular, its length  $5\frac{1}{2}$  in length of body; ventral of eyed side much produced, its length more than 3 times that of its mate. Color dark brown, speckled with darker, 3 large subcircular ocellated spots nearly as large as eye, with white center, dark iris, narrow, light margin, and a brown encircling outline, these arranged in an isosceles triangle, the apex on the lateral line, the others before it and distant from the lateral line a distance equal to their own diameter; blind side white; fins blotched with dark brown. (Goode & Bean.) Gulf Stream. Known from the original

types obtained in the deep waters (75 fathoms) of the Gulf Stream, off the Carolina coast. (*dilectus*, delightful.)

*Notosema dilecta*, GOODE & BEAN, Bull. Mus. Comp. Zool., x, No. 5, 193, 1883, Gulf Stream off the coast of South Carolina; GOODE & BEAN, Oceanic Ichthyology, 437, 1896.

*Paralichthys stigmatias*, GOODE, Nat. Hist. Aquat. Anim., 182, 1884; by inadvertence for *dilectus*.

*Ancylosetta dilecta*, JORDAN, Cat. Fish. N. A., 134, 1885; JORDAN & GOSS, Review Flounders and Soles, 250, 1889.

#### 1028. GASTROPSETTA, B. A. Bean.

*Gastropsetta*, BARTON A. BEAN, Proc. U. S. Nat. Mus. 1894, 633 (*frontalis*).

Body oblong-ovate, highly arched in front, covered with small, cycloid, embedded scales; lateral line arched in front, deflected downward on caudal peduncle. Teeth small, in a single series in each jaw. Dorsal fin beginning in advance of eye, its anterior rays produced, not connected by the irregular and broadly fringed membrane. Gill rakers very short, almost as broad as long, few in number. Ventral of eyed side produced, ending in a long filamentous ray in the young. This genus is closely allied to *Ancylosetta*, from which it differs in form of body, and especially in the entirely smooth scales, singularly branched and produced anterior dorsal rays, and very short and broad gill rakers. (*γαστήρ*, belly; *ψήτρα*, turbot or flounder.)

#### 3004. GASTROPSETTA FRONTALIS, B. A. Bean.

Head  $4\frac{1}{2}$ ; depth  $2\frac{1}{2}$ ; middle caudal rays  $2\frac{1}{2}$ ; eye large,  $3\frac{1}{2}$  in head. D. 60; A. 48; V. 6; P. I, 10. Mouth of moderate size, maxillary  $2\frac{1}{2}$  in head, the jaws curved; interorbital ridge prominent, very narrow. Dorsal beginning in front of eye on snout, its anterior rays singularly branched, the third and fourth longest, almost equaling length of head; anal fin beginning at vent, which is situated on blind side, its anterior rays scarcely produced; ventral of colored side much produced; middle caudal rays long. Color in spirits, light brown; 3 black spots on body, 2 along back, and 1 near anal base; fins with dusky blotches; several vertical stripes across eyes. A smaller specimen from Albatross Station 2317 has D. 62; A. 52; V. 6; P. I, 11. Gill rakers short, broad laminae, 2 + 7. Teeth weak, uniserial. Anterior rays of dorsal greatly produced, the third  $1\frac{1}{2}$  times as long as head. Ventral of eyed side very long, ending in a thread-like filament. Color as in the preceding. An example from Albatross Station 2373 near Apalachicola, is 224 mm. long; its depth 90 mm. D. 60; A. 49; P. I, 10; V. 6; C. 15. Vent situated in a deep notch, which forms the front margin of abdomen, and not on side, as in other specimens. Color darker than that of the Key West examples, being dark reddish brown; body spotted and fins blotched as in the preceding. Two specimens obtained by the Albatross, January 15, 1885, at Station 2317, Lat.  $24^{\circ} 25' 45''$  N., Long.  $81^{\circ} 46' 45''$  W., near Key West, Florida, in 45 fathoms of water, the type 8 inches long, the other one 6 inches. (B. A. Bean.) (*frontalis*, pertaining to the forehead.)

*Gastropsetta frontalis*, BARTON A. BEAN, Proc. U. S. Nat. Mus. 1894, 633, Key West. (Type, No. 37668, U. S. Nat. Mus. Coll. Albatross.)

1029. PLEURONICHTHYS, Girard.

*Pleuronichthys*. GIRARD, Proc. Ac. Nat. Sci. Phila. 1854, 139 (*cœnosus*).  
*Heteroproopon*, BLEEKER, Comptes Rendus Acad. Amsterdam, XIII, 1862, 8 (*cornutus*).  
*Parophrys*, GÜNTHER, Cat. Fishes, IV, 454, 1862; not of GIRARD.

Eyes and color on the right side. Body deep; head short, with very short, blunt snout; mouth small, with several series of slender, acute teeth, which are most developed on the blind side, and are often wanting in 1 or both jaws on the colored side; no teeth on vomer or palatines; lips thick, with several lengthwise folds within which is a series of short fringes. Lower pharyngeals narrow, each with a double row of very small teeth. Gill rakers wide set, very short and weak. Lateral line nearly straight, with a dorsal branch in our species. Scales small, cycloid, non-imbricate, embedded. Dorsal fin anteriorly twisted from the dorsal ridge toward the blind side; anal fin preceded by a spine; caudal fin convex behind. Intestinal canal elongate. Herbivorous species, feeding chiefly on algae. Pacific Ocean. This well-marked genus contains 3 American species, which are very closely related to each other. The Asiatic species, *Platessa cornuta*, Schlegel, of the coasts of China and Japan, is also a member of this group, having an accessory branch to the lateral line as in the American species. This species bears some resemblance to *Pl. verticalis*. The species of *Pleuronichthys* spawn in the spring, and live in comparatively deep water. (*πλαῦρον*, side; *ιχθῦς*, fish.)

- a. Dorsal fin beginning on the level of the lower lip, its first 9 rays on the blind side. DECURRENS, 3005.
- aa. Dorsal fin beginning on level of upper lip, its first 5 rays being on the blind side.
  - b. Interorbital ridge posteriorly with a strong spine directed backward, some tubercles on interorbital ridge. VERTICALIS, 3006.
  - bb. Interorbital ridge prominent, but without spines and conspicuous tubercles. CÆNOSUS, 3007.

3005. PLEURONICHTHYS DECURRENS, Jordan & Gilbert.

Head  $3\frac{1}{2}$ ; depth  $1\frac{1}{2}$ . D. 72; A. 40; scales 80; eye 3 in head; maxillary  $4\frac{1}{2}$ ; pectoral  $1\frac{1}{2}$ ; highest dorsal rays  $1\frac{1}{2}$ ; anal rays  $1\frac{1}{2}$ ; caudal 1. Body short and wide; mouth very small, the maxillary reaching nearly to pupil; teeth villiform, in moderate bands on blind side, a narrow band on eyed side of lower jaw; eyes very large, the upper edge of upper eye even with profile; snout extremely short; a blunt tubercle in front of upper eye, another at each end of the narrow interorbital ridge, the posterior largest, but usually not spine-like; 2 or 3 above the latter behind the upper eye; some prominences above the opercle; gill opening short, not extending above upper edge of pectoral. Dorsal beginning very low, on level of end of maxillary, its first 9 rays on the blind side; anal spine well developed, the origin of anal a little behind vertical from base of pectoral; pectoral of eyed side a little larger than its mate, both rounded behind; ventral of blind side shorter than that of eyed side, and placed slightly before it, caudal well rounded. Scales cycloid, embedded, a space between them anteriorly; lateral line without arch, slightly curved. Color brownish, usually much mottled with chocolate and grayish, often finely spotted with brownish on body and fins; all fins darker than body;

dorsal, anal, caudal, and ventrals narrowly edged with white; pectoral uniformly blackish. Pacific coast of United States, south to Monterey. This species is rather scarce along the California coast, being taken chiefly in deep water. It reaches a larger size than either *P. verticalis* or *P. cænosus*. Here described from a specimen from San Francisco market, 8 inches in length. (*decurrens*, running down.)

*Pleuronichthys cænosus*, LOCKINGTON, Proc. U. S. Nat. Mus. 1870, 97; not *Pleuronichthys cænosus*, GIRARD.

*Pleuronichthys quadrituberculatus*, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1880, 50, not of PALLAS; JOEDAN, Nat. Hist. Aquat. Anim., 189, 1884.

*Pleuronichthys decurrens*, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1880, 453, San Francisco; Monterey Bay (Coll. Jordan & Gilbert); JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1881, 69; JORDAN & GILBERT, Synopsis, 829, 1883; JORDAN & GOSS, Review Flounders and Soles, 282, 1889.

#### 3006. PLEURONICHTHYS VERTICALIS, Jordan & Gilbert.

Head 4 in body; depth 2. D. 65; A. 45; scales about 80; vertebrae 13 + 25 = 38. Form broad ovate, the outlines regular; head small, somewhat constricted behind the upper eye; eyes large, but smaller than in *P. decurrens*. Interorbital ridge narrow; a small tubercle or prominence in front of upper eye; a large one in front of upper edge of lower eye; another larger and sharper at interior edge of the interocular space; another at the posterior edge of interocular spine ridge; this latter developed into a long, sharp, triangular spine, which is nearly as long as the pupil, and is directed backward; a prominent tubercle at posterior lower angle of upper eye; upper edge of opercle somewhat uneven, but no other tubercles present. Mouth small, as in other species; the lips thick, with lengthwise plicae. Teeth in a broad band on the left (blind) side of each jaw; no teeth on the right side in either jaw. Gill rakers very small, weak, and flexible, about 10 in number. Scales essentially as in other species, small, cycloid, embedded, scarcely imbricated; lateral line nearly straight, with an accessory branch which extends to the middle of the dorsal fin. Dorsal fin beginning on blind side at level of premaxillary, there being but about 4 of its rays on left side of median line; vertical fins less elevated than in other species, the longest rays of dorsal about  $\frac{1}{2}$  length of head; anal fin preceded by a spine; caudal peduncle short and deep; caudal fin elongate, rounded behind; pectorals short, nearly equal; ventrals moderate, reaching anal spine. Color dark olive brown, with round grayish spots, the body and fins mottled with blackish. This species agrees in habits and general characters with *Pleuronichthys decurrens*. Coast of California, in rather deep water. The above description from the original type. (*verticalis*, pertaining to the vertex.)

*Pleuronichthys verticalis*, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1880, 49, San Francisco (Coll. Jordan & Gilbert); JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1881, 169; JORDAN & GILBERT, Synopsis, 829, 1883; JORDAN, Nat. Hist. Aquat. Anim., 189, 1884; JORDAN & GOSS, Review Flounders and Soles, 282, 1889.

#### 3007. PLEURONICHTHYS CÆNOSUS, Girard.

Head  $3\frac{1}{2}$ ; depth 2. D. 68; A. 49; scales 61; eye 3 in head; pectoral  $1\frac{1}{2}$ ; dorsal and anal rays  $1\frac{1}{2}$ ; caudal a little longer than head. Body ovate; snout scarcely produced; mouth small, maxillary reaching past front of

lower eye; 3 or 4 rows of teeth on blind side of jaws, 1 on eyed side of lower; eyes very large; interorbital a high, narrow ridge, somewhat angulated behind, but with no conspicuous spine or tubercle; snout very short, about  $\frac{1}{2}$  eye; gill opening not extending above upper edge of pectoral. Scales cycloid, embedded, some distance apart anteriorly, their edges not in contact; lateral line nearly straight, with a long dorsal branch which reaches past middle of body. Dorsal and anal high; origin of dorsal on blind side on a level with premaxillary, its first 5 rays on blind side; origin of anal under base of pectoral; pectoral of eyed side a little larger than its mate; caudal well rounded. Color dark brown, usually mottled, the colors variable; our specimens from Puget Sound, very dark, the fins colored like body, with light and dark spots; a conspicuous black spot on lateral line on middle of sides. Pacific coast, from Sitka to San Diego. This species is comparatively common in rather deep water and about rocks, being most abundant about Puget Sound. Its apparent abundance as compared with the other species of the genus is doubtless due to its inhabiting shallower waters than they. It is quite variable in form. The above description from a specimen, 6 inches long, from Seattle. (*canocus*, in: *ady.*)

*Pleuronichthys canosus*, GIRARD, Proc. Ac. Nat. Sci. Phila. 1854, 139, San Francisco; GIRARD, U. S. Pac. R. R. Surv., x, Fishes, 151, 1858; LOCKINGTON, Rep. Com. Fisheries California, 1878-79, 45; LOCKINGTON, Proc. U. S. Nat. Mus. 1879, 97; JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1880, 50; JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1886, 453; JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1881, 68; JORDAN & GILBERT, Synopsis, 830, 1883; JORDAN, Nat. Hist. Aquat. Anim., 189, 1884; JORDAN & GOSS, Review Flounders and Soles, 282, 1889; JORDAN, Proc. Cal. Ac. Sci. 1895, 852.

*Parophrys caenosa*, GÜNTHER, Cat., IV, 456, 1862.

### 1030. HYP SOPSETTA, Gill.

(DIAMOND FLOUNDERS.)

*Hypsopsetta*, GILL, Proc. Ac. Nat. Sci. Phila. 1864, 195 (*guttulatus*).

Eyes and color on the right side; body broad, ovate, rhomboid; mouth very small; teeth slender, equal, acute, in several series; lips thick, not plicate; lateral line nearly straight, with an accessory dorsal branch; scales small, smooth; dorsal fin beginning on the dorsal ridge, not turned to the blind side at its insertion; anal spine present; caudal fin convex; gill rakers little developed. This genus consists of a single species, abundant on the coast of California. It is very close to *Pleuronichthys*, from which it differs only in a few characters of comparatively minor importance. Its range is in shallower and warmer water than that of the species of *Pleuronichthys*, and, in accordance with this fact, its flesh is firmer and its number of vertebrae fewer than in the latter genus. (*ὄψι*, deep; *ψῆττα*, flounder.)

### 3008. HYSOPSETTA GUTTULATA (Girard).

(DIAMOND FLOUNDER.)

Head  $3\frac{1}{2}$ ; depth  $1\frac{1}{2}$ . D. 68; A. 50; scales 95. Body very deep, somewhat angulated near middle of back and belly; eyes moderate, separated by a flattish, raised area; head without spines or tubercles; scales of opercular

region little developed; those of blind side reduced; no teeth on right side of either jaw; accessory lateral line long,  $\frac{1}{2}$  length of body; anal spine small; pectorals about  $\frac{1}{2}$  length of head; ventrals rather short; caudal peduncle much deeper than long; caudal large, nearly as long as head. Brown, with numerous pale-bluish blotches in life, these disappearing in spirits; blind side white, with a strong tinge of yellow along profile of head; fins plain, sometimes with black specks. Coast of California and southward, Cape Mendocino to Magdalena Bay. This species is one of the most abundant in the shore waters of the California coast. It is a food-fish of fair quality. (*guttulatus*, with small spots.)

*Pleuronichthys guttulatus*, GIRARD, Proc. Ac. Nat. Sci. Phila. 1856, 137, Tomales Bay, California (Coll. E. Samuels); GIRARD, Journ. Boston Soc. Nat. Hist. 1857, pl. 25, figs. 1-4; GIRARD, U. S. Pac. R. R. Surv., x, Fishes, 152, 1858; LOCKINGTON, Rep. Com. Fisheries California, 1878-79, 44; LOCKINGTON, Proc. U. S. Nat. Mus. 1879, 94.

*Parophrys ayresi*, GÜNTHER, Cat., IV, 1862, 457, San Francisco. (Coll. Dr. W. O. Ayres.)

*Pleuroneetes guttulatus*, GÜNTHER, Cat., IV, 445, 1862.

*Hypsopsetta guttulata*, GILL, Proc. Ac. Nat. Sci. Phila. 1864, 195; JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1880, 453; JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1881, 68; JORDAN & GILBERT, Synopsis, 830, 1883; JORDAN, Nat. Hist. Aquat. Anim., 185, 1884; JORDAN & GOSS, Review of Flounders and Soles, 283, 1889.

### 1031. PAROPHRYS, Girard.

*Parophrys*, GIRARD, Proc. Ac. Nat. Sci. Phila. 1854, 139 (*vetulus*).

Body rather elongate, covered with small, cycloid scales; scales of the head roughish. Head rather pointed; mouth small, the teeth uniserial, all more or less blunt, compressed, incisor-like, close set. Lateral line with an accessory dorsal branch; upper eye on median line of top of head. A single species, on the Pacific coast of America. The narrow interorbital space and the vertical range of the upper eye give it a peculiar physiognomy, but in most regards it is not very different from some of the species of *Pleuroneetes*. (*παρά*, near; *ὄφρῦς*, eyebrow, from the narrow inter-bital.)

### 3009. PAROPHRYS VETULUS, Girard.

Head  $3\frac{1}{2}$ ; depth  $2\frac{1}{2}$ ; eye  $4\frac{1}{2}$  in head. D. 74 to 86; A. 54 to 68; scales 105 (tubes). Body elongate-elliptical; snout very prominent, much protruding, forming an abrupt angle with the descending profile; depth of head opposite middle of upper eye about equaling distance from middle of orbit to snout; eyes large, separated by a very narrow, high ridge, the upper with vertical range; mouth very small; maxillary not reaching pupil; teeth trenchant, small, and rather narrow, widened at tip, about 45 teeth on left side of lower jaw; few teeth on right side of lower jaw. Accessory lateral line long. Pectoral about  $\frac{1}{2}$  length of head; caudal truncate,  $1\frac{1}{2}$  in head; fin rays entirely scaleless; scales on body all cycloid, those on cheeks often slightly ciliated. Uniform light olive brown; the young somewhat spotted with blackish. Pacific coast of North America, Sitka to Santa Barbara. This small flounder lives in waters of moderate depth. It is, next to *Platichthys stellatus*, probably the most abundant of the flounders of the California coast. (*vetulus*, an old man.)

*Parophrys vetulus*, GIRARD, Proc. Ac. Nat. Sci. Phila. 1854, 140, California; GÜNTHER, Cat., IV, 455, 1862; LOCKINGTON, Rep. Com. Fish. Cal. 1878-79, 45; LOCKINGTON, Proc. U. S. Nat. Mus. 1879, 100; JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1880, 453; JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1881, 68; JORDAN, Nat. Hist. Aquat. Anim., 185, 1884; JORDAN & GOSS, Review Flounders and Soles, 284, 1889.

*Pleuronectes digrammus*, GÜNTHER, Cat., IV, 445, 1862, Victoria. (Coll. Earl Russell.)

*Parophrys hubbardi*, GILL, Proc. Ac. Nat. Sci. Phila. 1862, 281, San Francisco.

*Pleuronectes vetulus*, JORDAN & GILBERT, Synopsis, 831, 1883.

### 1032. INOPSETTA, Jordan & Goss.

*Inopsetta*, JORDAN & GOSS, in JORDAN, Cat. Fish. N. A., 136, 1885 (*ischyrus*).

This genus resembles *Parophrys*, differing chiefly in having the scales less imbricated, all strongly ctenoid, and having the eyes both lateral, the snout much less acute than in *Parophrys*. A single species, closely allied to *Platichthys stellatus*, but separated from it by the curious character common to many of our Pacific coast flounders, of having an accessory branch to the lateral line. ( $\zeta$ s, strength;  $\psi\eta\rho\tau\alpha$ , flounder.)

### 3010. INOPSETTA ISCHYRA (Jordan & Gilbert).

Head  $3\frac{1}{2}$ ; depth 2. D. 70 to 76; A. 52 to 57; V. 6; scales 85. Body oblong, robust; caudal peduncle rather long; snout projecting, forming an angle with the profile; mouth oblique, the chin projecting; teeth  $\frac{5+25}{10+22}$ , narrowly incisor-like, bluntish, in a single, rather close-set series; maxillary reaching past front of orbit,  $5\frac{1}{2}$  in head; eyes large; interorbital space rather broad, scaly, continuous with a ridge above opercle; head mostly covered with scales like those of the body, but smaller and rougher; gill rakers feeble; lower pharyngeals each with 2 rows of coarse, blunt teeth; scales thick and firm, adherent, not closely imbricated, those in front well apart; all the scales strongly ctenoid; blind side with similar scales, almost as strongly ctenoid; vertical fins mostly scaly; lateral line conspicuous, its scales less rough than the others; a distinct short accessory lateral line on both sides, extending to about the tenth dorsal ray, less than  $\frac{1}{2}$  head; a series of pores around lower eye behind; dorsal beginning over pupil, its anterior rays low, its highest rays nearly  $\frac{1}{2}$  length of head; caudal large, double truncate; pectoral of right side about  $\frac{1}{2}$  head. Light olive-brown, vaguely clouded with light and dark; fins reddish brown; a few roundish dusky blotches on dorsal and anal; pectoral and caudal tipped with dusky; blind side white, immaculate, or with small, round rusty spots; left side of head sometimes rusty tinged. Puget Sound. This species is known only from 4 specimens taken by Dr. Jordan at Seattle in 1880. It is a large, rough flounder, with firm, white flesh. (*ἰσχυρός*, robust.)

*Parophrys ischyrus*, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1880, 276 and 453, Puget Sound (Coll. Dr. Jordan); JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1881, 67; JORDAN, Nat. Hist. Aquat. Anim., 185, 1884.

*Pleuronectes ischyrus*, JORDAN & GILBERT, Synopsis, 832, 1883.

*Inopsetta ischyrus*, JORDAN, Cat. Fish. N. A., 136, 1885; JORDAN & GOSS, Review Flounders and Soles, 284, 1889.

## 1033. ISOPSETTA, Lockington.

*Isopsetta*, LOCKINGTON MS., in JORDAN & GILBERT, Synopsis, 832, 1883 (*isolepis*).

Body much compressed, elliptical in form; mouth rather large; the teeth chiefly uniserial, all more or less blunt, separated, not incisor-like; scales closely imbricated, all strongly ctenoid; lower pharyngeals each with a double row of bluntish teeth. A single species found on the coast of California. *Isopsetta* approaches in many respects very close to the large-mouthed flounders of the type of *Hippoglossoides*, and it may fairly be said to be intermediate between *Psettichthys* and *Lepidopsetta*. Its affinities on the whole are nearest the latter, but the close relation of the *Hippoglossine* and *Pleuronectine* is clearly shown. (ἴσος, equal; ψῆττα, flounder.)

## 3011. ISOPSETTA ISOLEPIS (Lockington).

Head 4; depth 2½. D. 88; A. 65; scales 88; vertebrae 10 + 32 = 42. Body elliptical, much compressed, moderately deep, the curvature very regular; head moderate, strongly compressed, the profile little depressed above the eye; eyes rather large; interorbital space broad, flattish, with several series of scales. Scales on cheeks similar to those on body, rather large, ctenoid, and closely imbricated. Mouth comparatively large, maxillary reaching pupil, 3½ in head; teeth not large, about  $\frac{11+14}{9+24}$ , conical, close set, in 1 somewhat irregular series, or partly in 2 series, those on colored side small; lower pharyngeals each with a double row of bluntish teeth. On the blind side the scales are more or less ctenoid, sometimes smooth; those on the cheeks weakly ctenoid; most of the opercle, the preopercle, interopercle, and subopercle on blind side naked; lateral line with a very slight arch in front, the depth of which is less than ½ the length; accessory branch nearly as long as head; fins rather low, mostly covered with ctenoid scales. Color brownish, mottled and blotched with darker. This small flounder is rather common off the coast of California, where it reaches a length of about 15 inches. It much resembles *Psettichthys melanostictus*, but its small mouth and blunt dentition indicate a real affinity with the small-mouthed flounders, among which it is here placed. Its nearest relative among our species is doubtless *Lepidopsetta bilineata*. Puget Sound to Point Concepcion, in rather deep water; not rare. (ἴσος, equal; λεπῖς, scale.)

*Lepidopsetta umbrosa*, LOCKINGTON, Proc. U. S. Nat. Mus. 1879, 106; not of GIRARD.

*Lepidopsetta isolepis*, LOCKINGTON, Proc. U. S. Nat. Mus. 1880, 325, San Francisco.

*Parophrys isolepis*, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1880, 453 and 1881, 67;

JORDAN & GILBERT, Synopsis, 832, 1883; JORDAN, Nat. Hist. Aquat. Anim., 186, 1884.

*Isopsetta isolepis*, JORDAN, Cat. Fish. N. A., 136, 1885; JORDAN & GOSS, Review Flounders and Soles, 285, 1889.

## 1034. LEPIDOPSETTA, Gill.

*Lepidopsetta*, GILL, Proc. Ac. Nat. Sci. Phila. 1864, 195 (*umbrosus*).

Body robust; mouth small. Teeth stout, conical, little compressed, bluntish, in 1 series, rather irregularly placed. Lateral line with a distinct arch in front and accessory dorsal branch; scales imbricated,



rough ctenoid, smooth in the very young. A single species, abundant on the Pacific coast of North America. It is close to *Inopsetta*, from which it is separated by the arch of the lateral line, and still closer to *Limanda*, from which the accessory branch of the lateral line alone separates it. (*λεπίς*, scale; *ψιττα*, flounder.)

## 3012. LEPIDOPSETTA BILINEATA (Ayres).

Head  $3\frac{1}{2}$ ; depth  $2\frac{1}{2}$ . D. 80; A. 60; teeth  $\frac{27+7}{25+10}$ ; scales 85. Vertebrae 11 + 29 = 40. Body broadly ovate, thickish; mouth moderate, turned toward the left side; teeth stout, conical, little compressed, bluntish, in 1 series, rather irregularly placed. Lower pharyngeals broad, with 2 rows of blunt teeth. Gill rakers few, very short, thick and weak, without teeth. Snout projecting; eyes large, separated by a prominent ridge, which, like the cheeks and upper portion of opercle, is covered with rough stellate scales; lower eye advanced; opercle, subopercle, and interopercle of left side scaly; preopercle naked. Scales rather small, mostly ctenoid, not closely imbricated, those on the blind side smooth; scales on cheeks and other parts of head very rough; scales of body smoother and less closely imbricated anteriorly, the degree of roughness variable, northern specimens (var. *umbrosus*) being roughest. Lateral line moderately arched anteriorly, with an accessory dorsal branch, which is less than  $\frac{1}{2}$  length of head; height of arch less than  $\frac{1}{2}$  its length. Dorsal beginning over eye, its anterior rays low; caudal convex; anal preceded by a spine; a concealed spine behind ventrals; rays of dorsal and anal all simple; dorsal and anal somewhat scaly; caudal  $\frac{3}{4}$  length of head; pectoral  $\frac{1}{2}$  head. Lower pharyngeals broad, each with 2 rows of blunt teeth. Yellowish brown, with numerous round pale blotches. Pacific coast of North America, Bering Strait to Monterey. This species is one of the commonest of the flounders of the Pacific coast, its abundance apparently increasing toward the northward. In Bering Sea it far outnumbered all other flounders. We have specimens from Bering Island, Medni Island, Unalaska, St. Paul, St. George, and Chignik Bay. It reaches a weight of 5 or 6 pounds and is an inhabitant of shallow waters. Specimens from Puget Sound and northward are rougher than southern specimens and constitute a slight geographical variety, for which the name *Lepidopsetta bilineata umbrosa* may be used. This is the same as *P. pararcuatus* of Cope. (*bilineatus*, two-lined.)

*Platessa bilineata*, AYRES, Proc. Ac. Nat. Sci. Cal. 1855, 40, San Francisco.

*Platichthys umbrosus*, GIRAUD, Proc. Ac. Nat. Sci. Phila. 1856, 136, Puget Sound.

*Pleuronectes pararcuatus*, COPE, Proc. Ac. Nat. Sci. Phila. 1873, 30, Unalaska.

*Pleuronectes umbrosus*, GÜNTHER, Cat., IV, 454, 1862.

*Pleuronectes bilineatus*, GÜNTHER, Cat., IV, 444, 1862; JORDAN & GILBERT, Synopsis, 833, 1883.

*Lepidopsetta bilineata*, GILL, Proc. A.S. Nat. Sci. Phila. 1864, 105; LOCKINGTON, Proc. U. S. Nat. Mus. 1879, 103; LOCKINGTON, Rep. Com. Fisheries California, 1878-79, 46; JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1880, 453; JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1881, 68; BEAN, Proc. U. S. Nat. Mus. 1881, 241; BEAN, Cat. Coll. Fish. U. S. Nat. Mus. 1883, 19; BEAN, Proc. U. S. Nat. Mus. 1883, 353; JORDAN, Nat. Hist. Aquat. Anim., pl. 50, 184, 1884; JORDAN & GOSS, Review Flounders and Soles, 286, 1889.

## 1035. LIMANDA, Gottsche.

(MUD DABS.)

*Limanda*, GOTTSCHÉ, Archiv für Naturgesch. 1835, 100 (*limanda*).*Myzopssetta*, GILL, Proc. Ac. Nat. Sci. Phila. 1864, 217 (*ferruginea*).

Teeth chiefly uniserial; lateral line with a distinct arch in front, and without accessory dorsal branch; scales imbricated, rough etenoid; vertebrae about 40. This genus is closely allied to *Pseudopleuronectes*, from which it differs only in the presence of an arch on the anterior part of the lateral line. (*Limanda*, an old name of the European Dab, *Limanda limanda*, from *limus*, mud.)

a. Head comparatively large,  $3\frac{1}{2}$  to  $4\frac{1}{2}$  in length.

b. Dorsal rays 85; anal rays 62.

c. Scales rather small, 90 to 100 in lateral line; scales of right side etenoid, closely imbricated, those of blind side mostly smooth; teeth conical, close set, forming a continuous series, about 11 + 30 in lower jaw; snout abruptly projecting, forming in front of upper eye a sharp angle with the descending profile. FERRUGINEA, 3013.

cc. Scales larger, wide apart, about 80 in lateral line; scales of blind side more or less rough. ASPERA, 3014.

bb. Dorsal rays 60 to 70; anal 47 to 53; scales small, 86 to 95; snout long, protruding; scales of blind side smooth. PROBOSCIDEA, 3015.

aa. Head very short,  $5\frac{1}{2}$  in length; snout very short; interorbital space very narrow. D. 64; A. 63; scales 88. DEANII, 3016.

## 3013. LIMANDA FERRUGINEA (Storer).

(RUSTY DAB.)

Head 4 in length; depth  $2\frac{1}{2}$ . D. 85; A. 62; scales 100. Body ovate-elliptical, strongly compressed; teeth small, conical, close set, in a single series on each side in each jaw, about 11 + 30 in the lower jaw; snout projecting, forming a strong angle above upper eye, with the descending profile; gill rakers of moderate length, very weak, not toothed; eyes moderate,  $4\frac{1}{2}$  in head, the lower slightly in advance of upper, separated by a high, very narrow ridge, which is scaled posteriorly, and is continued backward as an inconspicuous but rough ridge to the beginning of the lateral line; scales imbricate, nearly uniform, those on right side rough etenoid, those on left side nearly or quite smooth; scales on body rougher than on cheeks; caudal peduncle short, higher than long; dorsal inserted over middle of eye, its middle rays highest; pectoral less than  $\frac{2}{3}$  length of head; caudal fin rounded; anal spine present; lateral line simple, with a rather low arch in front, the depth of which is barely  $\frac{1}{2}$  the length; a concealed spine behind ventrals; ventral of colored side partly lateral, the other wholly so; anal spine strong. Brownish olive, with numerous, irregular, reddish spots; fins similarly marked; left side with caudal fin, caudal peduncle and margins of dorsal and anal fins lemon yellow. Atlantic coast of North America, Labrador to New York. This species is rather common northward on our Atlantic coast. It is allied to the European Dab, but has smaller scales and a more prominent snout. Our specimens are from the east coast of Massachusetts. (*ferrugineus*, rusty red.)

*Plateasa ferruginea*, D. H. STORER, Rept. Fish. Mass., 141, pl. 2, 1839, Cape Ann; DE KAY New York Fauna: Fishes, 297, pl. 48, fig. 155, 1842; STORER, Syn. Fish. N. A., 476, 1846.

*Plateasa rostrata*, H. R. STORER, Bost. Journ. Nat. Hist., VI, 1850, 268, Labrador.

*Pleuronectes ferrugineus*, GÜNTHER, Cat., IV, 447, 1862; JORDAN & GILBERT, Synopsis, 834, 1882.

*Myzopsetta ferruginea*, GILL, Proc. Ac. Nat. Sci. Phila. 1864, 217.

*Limanda ferruginea*, GOODE, Proc. U. S. Nat. Mus. 1880, 472; GOODE, Hist. Aquat. Anim., pl. 49, 1884; GOODE & BEAN, Oceanic Ichthyology, 427, 1890; JORDAN & GOSS, Review Flounders and Soles, 287, 1889.

*Limanda rostrata*, GILL, Proc. Ac. Nat. Sci. Phila. 1864, 217.

3014. LIMANDA ASPERA (Pallas).

(ALASKA DAB.)

Head  $3\frac{1}{2}$ ; depth 2. D. 69; A. 53; scales about 80. Form of *Lepidopssetta bilineata*. Teeth small, almost conical, on both sides of the mouth; interorbital space narrow, scaly; opercle and preopercle naked below; gill rakers very feeble; pharyngeals not very broad, their teeth bluntyish, not paved; scales small, wide apart, partly embedded, each one with 1 to 4 spinules, which are almost erect; anterior scales with 3 to 4 of these spinules; posterior mostly with 1; scales of blind side smoother; only middle rays of dorsal and anal scaly; no accessory lateral line; anal spine present; twentieth anal ray and thirty-seventh dorsal ray longest; caudal double truncate. Brown, nearly plain, the blind side with tinges of lemon yellow. Bering Sea, generally common, south to Vancouver Island and to the Okhotsk Sea. We have specimens from Petropaulski and Robben Reef, Bristol Bay, and Herendeen Bay. It is especially abundant in Bristol Bay, and, according to Dr. Gilbert, it is an excellent food-fish. Dr. Bean has also collected it in various localities in Alaska. Its scales are larger and rougher than in *L. ferruginea* which, in many respects, it resembles. A specimen from the island of Saghalien is in the museum at Cambridge. The above description is from examples taken by Dr. Bean. (*asper*, rough.)

*Pleuronectes asper*, PALLAS, Zoogr. Rosso-Asiat., III, 425, 1811, east coast of Siberia; GÜNTHER, Cat., IV, 454, 1862; STEINDACHNER, Plouronectiden, etc., aus Decanstris Bay, 1870-1875; JORDAN & GILBERT, Synopsis, 835, 1883.

*Limanda aspera*, BEAN, Proc. U. S. Nat. Mus. 1881, 242; BEAN, Cat. Coll. Fish, U. S. Nat. Mus. 1883, 20; BEAN, Proc. U. S. Nat. Mus. 1883, 354; BEAN, Hist. Aquat. Anim., 184, pl. 48, 1884; JORDAN & GOSS, Review Flounders and Soles, 288, 1889.

3015. LIMANDA PROBOSCIDEA, Gilbert.

Depth  $2\frac{1}{2}$  to  $2\frac{3}{4}$  in length; head large, 3 to  $3\frac{1}{2}$  in length in a specimen 7 inches long. D. 63 to 67; A. 47 to 49; scales 86 to 95. Resembling *L. ferruginea*, but having fewer rays in dorsal and anal, larger scales and longer snout. Profile sharply angulated above front of upper eye, the snout convexly protruding; form varying from very slender to broadly elliptical, the 2 outlines equally curved; caudal peduncle short, widening backward, its least depth twice its length; mouth oblique, maxillary reaching beyond front of lower eye, 4 in head; teeth narrow, little compressed, in a single series on both sides of the jaw, extending farther back on the blind side; eyes on right side; lower eye well in advance of upper, the diameter of

upper eye  $5\frac{1}{2}$  to 6 in head,  $1\frac{1}{2}$  in snout; vertical from front of upper eye, falling midway between front of orbit and front of pupil of lower eye; interorbital space a very narrow, sharp ridge, naked in females, with a single series of etenoid scales in males; gill rakers short, about equal to diameter of pupil, 13 or 14 in number, 9 or 10 on lower limb; scales loosely imbricated, etenoid in males on colored side, smooth in females; blind side of both sexes smooth; head scaled on eyed side in males; the opercle, subopercle, interopercle, and preopercle mostly naked in females; head on blind side naked; rays of vertical fins with a single series of etenoid scales; dorsal fin beginning slightly behind front of upper eye, the first 3 rays usually higher and with membranes more deeply incised than in those which follow; highest portions of both dorsal and anal fins behind the middle of the body; these fins about equal, their longest rays equal to the snout and eye; caudal  $\frac{3}{4}$  head; pectorals short,  $\frac{1}{4}$  in head; ventrals reaching beyond front of anal,  $3\frac{1}{2}$  in head; the usual small antrorse spine in front of anal fin. Color light grayish or brownish, thickly covered with small whitish spots; entire left side with margins of dorsal, caudal, and anal fins bright lemon yellow (as in *ferruginea*); vertical fins grayish, with an occasional dark-brown ray. Specimens described  $7\frac{1}{2}$  inches long. Bering Sea; several specimens from Albatross Stations 3239 and 3240, in Bristol Bay, in  $11\frac{1}{2}$  to  $14\frac{1}{2}$  fathoms; 1 young individual from Herendeen Bay. (Gilbert.) (*proboscideus*, having a long snout or proboscis.)

*Limanda proboscidea*, GILBERT, Report U. S. Fish Comm. 1893 (1896), 460, pl. 33, Bristol Bay and Herendeen Bay. (Coll. Albatross.)

#### 3016. LIMANDA BEANII, Goode.

Head  $5\frac{1}{2}$ ; depth  $2\frac{3}{4}$ . D. 64; A. 63; scales 88. Body elliptical, with angular outlines, strongly compressed; head very short; snout abbreviated; mouth small, subvertical; teeth small, apparently in two rows, chiefly on the blind side of lower jaw; eyes large, as long as mandible; interorbital space very narrow. Dorsal fin beginning about pupil, its rays long, wide apart, exerted; right ventral near the median line; caudal broad, fan-shaped. Lateral line with an abrupt curve, the length of which is twice its height and about equal to length of head, its scales highly specialized; lateral line on colored side less developed; scales small, strongly etenoid on the right side; larger and cycloid on the blind side. Grayish, mottled with darker; a conspicuous black blotch on the outer rays of caudal on each side. (Goode.) Deep water off the coasts of New England; not common. (Named for Dr. Tarleton Hoffman Bean.)

*Limanda beanii*, GOODE, Proc. U. S. Nat. Mus. 1880 (Feb. 16, 1881), 473, southern coast New England, Fish Hawk Stations, 875, 876; GOODE & BEAN, Oceanic Ichthyology, 428, pl. 102, figs. 355a and 355b, 1896.

*Pleuronectes beanii*, JORDAN & GILBERT, Synopsis, 835, 1883; JORDAN & GOSS, Review Flounders and Soles, 288, 1889.

#### 1036. PSEUDOPLEURONECTES, Bleeker.

(WINTER FLOUNDERS.)

*Pseudopleuronectes*, BLEEKER, Comptes Rendus Acad. Amst., Plenron., 7, 1862 (*planus*).

Body oblong, with firm flesh; the scales firm, regularly imbricated, strongly etenoid on eyed side in both sexes; fin rays scaly; mouth small;

teeth uniserial, incisor-like, close set, all more or less blunt; lower pharyngeals very narrow, each with 2 rows of separate, conical teeth. This genus is distinguished from *Pleuronectes* chiefly by the well-imbricated ctenoid scales, and from *Limanda*, which it more closely resembles, by the want of arch to the lateral line. Besides the typical species, we refer to this genus a second from the North Pacific. (*ψεδδος*, false; *Pleuronectes*.)

a. Dorsal rays 65; anal rays 48; scales 83; vertical fins nearly plain.

AMERICANUS, 3017.

aa. Dorsal rays 58; anal rays 38; scales 70; vertical fins with black bars.

PINNIFASCIATUS, 3018.

3017. PSEUDOPLEURONECTES AMERICANUS (Walbaum).

(COMMON FLATFISH; WINTER FLOUNDER.)

Head 4 in length; depth 2½. D. 65; A. 48; scales 83. Body elliptical; an angle above eye. Head covered above with imbricated, strongly ctenoid scales, similar to those on the body; blind side of head nearly naked; interorbital space rather broad, strongly convex, its width ½ eye, entirely scaled; teeth compressed, incisor-like, widened toward tips, close set, forming a continuous cutting edge; some of teeth often emarginate, sometimes movable; right side of each jaw toothless. Highest dorsal rays less than length of pectorals, and more than ¼ length of head; anal spines present. Dark rusty brown, spotted or nearly plain; young olive brown, more or less spotted and blotched with reddish. Atlantic coast of North America, from Labrador to Chesapeake Bay. This small flounder is one of the most abundant of the group on our Atlantic coast. It reaches a length of about 15 inches and a weight of less than 2 pounds. It is a very good food-fish and sells readily in the markets. Along the south coast of Massachusetts this species is more abundant than any other of the flatfishes. The specimens examined by us are from Labrador, Cape Breton, Anticosti, Grand Menan, Boston, Provincetown, Woods Hole, New Bedford, and Somers Point, New Jersey.

*Flounder*, SCHÖFF, Schrift. Gesellschaft Naturforscher Freunde, VIII, 1788, 148, New York.

*Pleuronectes americanus*, WALBAUM, Artedi Piscium, 113, 1792, based on the *Flounder* of

SCHÖFF; BLOCH & SCHNEIDER, Syst. Ichth., 150, 1801; GÜNTHER, Cat., IV, 443, 1862;

JORDAN & GILBERT, Synopsis, 637, 1883; STEARNS, Proc. U. S. Nat. Mus. 1883, 125.

*Pleuronectes planus*, MITCHELL, Trans. Lit. & Philos. Soc. N. Y., I, 1815, 387, New York.

*Platessa pusilla*, DE KAY, New York Fauna: Fishes, 296, pl. 47, fig. 153, 1842, New York;

STORER, Synopsis, 477, 1846.

*Platessa plana*, STOREY, Rept. Fishes Mass., 140, 1839; DE KAY, New York Fauna:

Fishes, 295, pl. 49, fig. 158, 1842; STORER, Synopsis, 476, 1846.

*Pseudopleuronectes planus*, BLEEKER, Comptes Rendus Amstord., XIII, 1862, 7.

*Pseudopleuronectes americanus*, GILL, Proc. Ac. Nat. Sci. Phila. 1864, 216; GOODE, Nat. Hist.

Aquat. Anim., 182, pl. 44, 1884; JORDAN & GOSS, Review Flounders and Soles, 289, 1889.

3018. PSEUDOPLEURONECTES PINNIFASCIATUS (Kner).

Head 3½ in body; depth 2½. D. 58; A. 38; scales 70; eye 5½ in head; snout 5; highest anal ray 2; pectoral 2; caudal 4½ in body. Body sub-elliptical, the snout rather pointed and not forming an angle above eye; mouth rather small, maxillary reaching scarcely to the middle of the lower eye; interorbital space rather broad, ¼ width of eye; a rather prominent

rugose ridge above opercle, with a smaller similar ridge behind it; both sides of jaws with teeth, those on blind side stronger; origin of dorsal over middle of upper eye. Color brown, with vague dusky spots; 6 or 7 blackish vertical bars on dorsal and anal; similar lengthwise blotches on caudal. Okhotsk Sea, east to Kamchatka. (Steindachner.) Not seen by us. From the excellent figure we conclude that it belongs to *Pseudopleuronectes*, although its pharyngeals have not been described. It seems to us nearer to *P. americanus* than to *Liopsetta glacialis*. (pinna, fin; *fasciatus*, banded.)

*Pleuronectes pinnifasciatus*, KNER, in STEINDACHNER, Ueber einige Pleuronectiden, etc., aus Decastris Bay, 2, pl. 1, fig. 1, 1870, Decastris Bay, mouth of Amur River; JORDAN & GOSS, Review Flounders and Soles, 290, 1889.

### 1037. PLEURONECTES (Artedi) Linnaeus.

(PLAICE.)

*Pleuronectes*, ARTEDI, Genera, etc., in part, 16, 1738.

*Pleuronectes*, LINNÆUS, Syst. Nat., Ed. X, 208, 1758 (*platessa*); included all known *Pleuronectidae*.

*Platessa*, CUVIER, Règne Animal, Ed. 1, II, 220, 1817 (*platessa*).

*Pleuronectes*, SWAINSON, Nat. Hist. Class'n Anim., II, 302, 1839 (*platessa*).

*Pleuronectes*, BLEEKER, Comptes Rendus Acad. Amsterd., XIII, 1862 (*platessa*); and of most recent authors.

Body oblong, with firm flesh. Mouth small, teeth uniserial, incisor-like, compressed, forming a continuous cutting edge. Lateral line straightish, without arch or accessory dorsal branch. Scales imperfectly imbricated, chiefly cycloid in both sexes; lower pharyngeals small and narrow, separate, each with 1 or 2 rows of small bluntish teeth. No stellate scales along bases of dorsal and anal. Species mostly European; valued as food. (*πλεῦρον*, side; *νέκτης*, swimmer.)

### 3019. PLEURONECTES QUADRITUBERCULATUS, Pallas.

Head  $3\frac{1}{2}$ ; depth 2. D. 68; A. 50; scales 78. Mouth very small, with small, incisor-like teeth, rounded at tip. Eyes separated by a narrow ridge; about 5 small, prominent, conical, obtuse, bony tubercles in a row above the opercle, continuous with the direction of the lateral line, which is straight, without accessory dorsal branch; tubercle above opercle largest. Scales small, cycloid in all specimens examined. Anal spine present. Grayish, mottled with paler and with round black spots; fins very dark. Bering Sea on both coasts, south to Kadiak; not common. Our specimens from Avatcha Bay, Bristol Bay, Herendeen Bay, Chernofsky Harbor, Grantley Harbor, Chignik Bay, and Robben Island. The above description from a small specimen (No. 28025, U. S. Nat. Mus.) collected by Mr. W. J. Fisher at Kadiak. The species proves, as suspected by Jordan & Goss, to be a true *Pleuronectes*, having the lower pharyngeals narrow, separate, with 2 rows of bluntish teeth. (*quadrītuberculatus*, having four tubercles.)

*Pleuronectes quadrītuberculatus*, PALLAS, Zoogr. Rosso-Asiat., III, 423, 1811, sea between Kamchatka and Alaska; BEAN, Proc. U. S. Nat. Mus. 1881, 241; JORDAN & GILBERT, Synopsis, 836, 1883.

*Pleuronectes pallasi*, STEINDACHNER, Ichth. Beitr., VIII, 45, 1879, Kamchatka.

*Parophrys quadrītuberculatus*, GÜNTHER, Cat., IV, 456, 1862.

*Platessa quadrītuberculata*, JORDAN & GOSS, Review Flounders and Soles, 292, 1889.

## 1038. LIOPSETTA, Gill.

(EEL-BACK FLOUNDERS.)

*Liopsetta*, GILL, Proc. Ac. Nat. Sci. Phila. 1864, 217 (*glaber*); females.*Euchalarodus*, GILL, Proc. Ac. Nat. Sci. Phila. 1864, 222 (*putnami*); males.

Teeth chiefly uniserial, incisor-like; scales imperfectly imbricated, rough ctenoid in the male, more or less cycloid in the female (fin rays scaly in the male, naked in the female); lower pharyngeals very large, more or less united in the adult, their surface somewhat concave, with teeth in 5 or 6 rows, large, blunt, close set; lateral line without arch or dorsal branch. This genus comprises several species of small flounders of the Arctic seas. The genus is distinguished by the large, half-united pharyngeals, as also by the peculiar squamation, the scales in the males being very rough, in the females smooth. This difference has given rise to the nominal genus *Euchalarodus*, based on the males, while *Liopsetta* was based on the smoother females, which were erroneously supposed to be scaleless. (*λειος*, smooth; *ψιττα*, flounder.)

a. Dorsal rays 55 or 56; anal 40 to 42.

b. Pectoral fin short,  $\frac{1}{2}$  length of head in males, shorter in females. GLACIALIS, 3020.bb. Pectoral fin long,  $1\frac{1}{2}$  in head in males, nearly 2 in females. PUTNAMI, 3021.aa. Dorsal rays 59 to 62; anal 45 or 46; pectoral  $1\frac{1}{2}$  in head in males. OBSCURA, 3022.

## 3020. LIOPSETTA GLACIALIS (Pallas).

(ARCTIC FLOUNDER.)

Head 4; depth  $2\frac{1}{2}$ . D. 56; A. 42. Form of *Liopsetta putnami*. A roughened ridge above the cheeks and opercles on the eyed side. Eyes separated by a narrow, smooth, bony ridge. Scales minute, embedded, nonimbricate, ctenoid in the males, smooth in the females; scales on blind side similar, less developed; scales of lateral line a little larger. Teeth colored, incisor-like, forming an even edge, mostly on blind side. An anal spine; pectorals short. Dark brown, the fins spotted. Arctic shores of Alaska and Siberia, south in Bering Sea to Petropaulski, St. Michaels, and Bristol Bay. Our specimens from Port Clarence, Petropaulski, Bristol Bay, mouth of Nushagak River, and Kotzebue Sound; the description from specimens from the last-named locality taken by Dr. Bean. It is said to be abundant in the Arctic Ocean and as far south as Bristol Bay. "Although small, its great abundance and fine flavor make it important as an article of food." The male is the rough fish described by Pallas as *P. cicatricosus*. The smoother female is Dr. Günther's *Pleuronectes franklinii*, the sexual differences being much as in *Liopsetta putnami*. *Liopsetta drinensis* of the northern coasts of Russia may be the same species. (*glacialis*, icy.)

*Pleuronectes glacialis*, PALLAS, Itin., III, App., 706, mouth of River Obi; BLOCH & SCHNEIDER, Syst. Ichth., 150, 1801; PALLAS, Zoogr. Rosso-Asiat., II, 424, 1811; RICHARDSON, Fauna Bor.-Amer., Fish., 258, 1836; DE KAY, N. Y. Fauna: Fishes, 302, 1842; STORER, Synopsis Fish. N. A., 479, 1846; BEAN, Proc. U. S. Nat. Mus. 1881, 241; JORDAN & GILBERT, Synopsis, 337, 1883; BEAN, Cat. Coll. Fish. U. S. Nat. Mus., 20, 1883; BEAN, Nat. Hist. Aquat. Anim., 184, pl. 47, 1884.

*Pleuronectes cicatricosus*, PALLAS, Zoogr. Rosso-Asiat., III, 424, 1811, male, sea between Kamchatka and Alaska.

- 1 *Platessa dvinensis*, LILLJEBORG, Vch. Ah. Handl. 1850, 360, tab. 20, mouth of River Dwina.  
*Pleuronectes franklinii*, GÜNTHER, Cat. Fish., IV, 442, 1862, Arctic seas of America, female;  
 BEAN, Proc. U. S. Nat. Mus. 1881, 241.  
*Liopsetta glacialis*, JORDAN & GOSS, Review Flounders and Soles, 295, pl. 17, 1889.

## 3021. LIOPSETTA PUTNAMI (GILL).

(EEL-BACK FROUNDER.)

Head  $3\frac{1}{2}$ ; depth 2. D. 55; A. 40; scales 70 (pores). Body oblong, ovate. Eyes rather small, separated by a naked elevated ridge. Jaws sometimes each with 2 distinct rows of teeth, the interrupted outer series of truncate, close set, thickish, incisor-like teeth, which are sometimes movable; the inner row of similar teeth more widely set and rather distant from the outer row (and often or generally wanting); about 20 teeth in outer row in lower jaw; right side of each jaw toothless; interorbital ridge continuous, with a broad, naked, smoothish, tuberculose ridge, which joins the lateral line. Scales small, distant, nonimbricate, smooth in the female, and more or less ctenoid in the male, those on blind side smaller. Fins moderate, somewhat scaly; anterior rays of dorsal low; pectoral a little more than  $\frac{1}{2}$  head; bases of vertical fins not tuberculate; anal spine present; lower pharyngeals separate, broad, with coarse teeth. Grayish brown, mottled with darker brown; fins with blackish spots. Length 10 inches. Atlantic coast of North America, from Cape Cod northward to Labrador and beyond; occasionally found in abundance. This species is rather common along the coast of northern Massachusetts and northward to Labrador. Specimens are frequently found in the markets, mixed with those of *Pseudopleuronectes americanus*. The numerous specimens in our possession were found in the markets of Indianapolis, having been sent thither from Boston. The remarkable sexual differences in the species have been fully discussed by Dr. Bean (Proc. U. S. Nat. Mus. 1878, 345), the form formerly called *Euchalarodus putnami* being the male, and that called *Pleuronectes glaber* being the female of the same species. These conclusions of Dr. Bean are fully corroborated by our series of specimens in which both sexes are fully represented.

Although *Liopsetta putnami* is abundant where found, its ascertained range is somewhat limited. The specimens in the United States National Museum represent localities from Salem, Massachusetts, to Belfast, Maine. In the Museum of Comparative Zoology the localities represented are Providence, Boston, Salem, Grand Manan, and Labrador. (Named for Prof. Frederic Ward Putnam.)

- Platessa glabra*, STOREY, Proc. Boston Soc. Nat. Hist. 1843, 130, female, Massachusetts; STOREY, Syn. Fish. N. A., 477, 1846; STOREY, Hist. Fish. Mass., 190, pl. 31, fig. 1, 1867; PUTNAM, Bull. Essex Inst., VI, 1874, 12; not *Platessa glabra* of RATHKE, 1837, a species of *Fleuss*.  
*Euchalarodus putnami*, GILL, Proc. Ac. Nat. Sci. Phila. 1864, 216-221, Salem, Massachusetts (Coll. F. W. Putnam), male; GILL, Report U. S. Fish Comm. 1873, 794; GOODE & BEAN, Amer. Journ. Sci. and Arts, XIV, 1877.  
*Liopsetta glabra*, GILL, Proc. Ac. Nat. Sci. Phila. 1864, 217.  
*Pleuronectes glaber*, GILL, Report U. S. Fish Comm. 1873, 794; GOODE & BEAN, Amer. Journ. Sci. and Arts, XIV, 1877, 476; XVII, 1879, 40; GOODE & BEAN, Proc. U. S. Nat. Mus. 1878, 347; JORDAN & GILBERT, Synopsis, 836, 1883; GOODE, Nat. Hist. Aquat. Anim., 183, pl. 45, 1884.  
*Liopsetta putnami*, JORDAN & GOSS, Review Flounders and Soles, 294, pl. 16, 1889.



## 3022. LIOPSETTA OBSCURA (Herzenstein).

To this species we refer 2 males from Shana Bay, Iturup Island. The scales on the colored side are everywhere strongly ctenoid and imbricated, while in Herzenstein's types (supposed to be females) they were cycloid. In our specimens the head is somewhat smaller,  $3\frac{1}{6}$  in length instead of  $3\frac{1}{5}$  to  $3\frac{2}{5}$ ; the depth is greater,  $2\frac{1}{2}$  in length instead of  $2\frac{3}{4}$  to  $2\frac{4}{5}$ ; the interorbital space is covered with very fine scales, not naked; the curve of the lateral line seems more marked, its cord contained 5 instead of 6 times in the straight portion. All of the fins are higher than in the female types, the pectoral of colored side being  $1\frac{3}{4}$  head, the caudal  $1\frac{1}{2}$ , the ventral  $\frac{1}{2}$  head, and the highest dorsal ray  $1\frac{1}{2}$ . Some of these differences may well be sexual. The lower pharyngeals are short and broad, the two closely appressed but united in our specimens, 27 and 29 cm. long. The teeth are large and very blunt, like cobble stones, and are arranged in 1 row along the outer edge; a row of larger teeth along the inner edge, and a short row along the posterior edge of the triangle. The arrangement is very similar is that found in *L. glacialis*, but here a few small teeth, without definite arrangement, are interposed in the middle of the bone, between the third series described. Dorsal 59 and 62; anal 45 and 46; tubes in the lateral line 79. Color on eyed side uniform dark brown on body and fins, the extreme tips of the fin rays white; on blind side yellowish white, with a few irregular scattered dark spots; dorsal and anal yellowish at base, becoming more or less mottled with dusky on distal half, the fins marked with broad dark bars parallel with the rays, about 7 on the anal fin, 10 or 11 on the dorsal; caudal light on basal half more or less blotched with darker, becoming black posteriorly. With this species we identify also a number of young individuals, 9 to 15 cm. long, from the same locality (Iturup Island). They are probably young females, but the viscera are in such condition as to prevent positive determination. The scales are perfectly smooth, but in other respects they agree perfectly with the adult males, except in their more varied coloration; head and body brownish, profusely spotted in coarser or finer pattern with light gray; also with a few scattered black spots edged with gray; markings on the fins as described for adults. In 7 specimens the dorsal contains 60, 62, 62, 62, 64, 65, and 66 rays; anal 45, 45, 45, 46, 47, 47, 48. Sea of Okhotsk. Our specimens from Shana Bay, Iturup Island, one of the Kurils; originally described from Manchuria. (*obscurus*, dark.)

*Pleuronectes obscurus*, HERZENSTEIN, Mélanges Biologiques, 127, 1890, Manchuria.  
*Liopsetta obscura*, JORDAN & GILBERT, Rep. Fur Seal Invest., 1898.

## 1039. PLATICHTHYS, Girard.

(STARRY FLOUNDERS.)

*Platichthys*, GIRARD, Proc. Ac. Nat. Sci. Phila. 1854, 136 (*rugosus*=*stellatus*).

Body very robust, broad, not greatly compressed. Mouth small; teeth chiefly uniserial, incisor-like. Scales all in both sexes and on both sides of body reduced to coarse scattered stellate tubercles, which are not

imbricated; similar tubercles between bases of dorsal and anal rays; lateral line without scales, with no anterior arch or accessory lateral line; lower pharyngeals broad, each with 3 rows of blunt coarse teeth. A single species, the largest of the small-mouthed flounders, and distinguished from related forms chiefly by the development of coarse stellate tubercles instead of scales. (*πλατύς*, flat; *λίθύς*, fish.)

## 3023. PLATICHTHYS STELLATUS (Pallas).

(GREAT FLOUNDER.)

Head 3 $\frac{1}{2}$ ; depth 2. D. 58; A. 42. Vertebrae 34. Body broad and short, the snout forming a slight angle with the profile; lower jaw projecting; interocular space rather broad, with very rough scales; large rough scales at base of dorsal and anal rays and on sides of head; similar but smaller scales scattered over the body; lateral line smooth; fins without scales; a cluster of bony prominences above opercle. Teeth incisor-like, truncate, rather broad,  $\frac{10+15}{13+16}$ . Lower pharyngeals broad, with coarse paved teeth. Dark brown or nearly black, with lighter markings; fins reddish brown; dorsal and anal with 4 or 5 vertical black bands; caudal with 3 or 4 black longitudinal bands. Pacific coast of America, from Point Conception to the Arctic Ocean and south to the Amur River. This is one of the largest of the American flounder, reaching a weight of 15 to 20 pounds. Of the small-mouthed flounders it is much the largest species known. It is an excellent food-fish, and from its size and abundance it is one of the most important of the group in the region where it is found, constituting half the total catch of flounders on our Pacific coast, and it is equally abundant in Bering Sea. It lives in shallow water and sometimes ascends the larger rivers. It is one of the most widely distributed of all the flounders, its range extending from San Luis Obispo, where it was obtained by Jordan & Gilbert, to the mouth of the Anderson and Colville rivers on the Arctic coast, where it was observed by Dr. Bean, and to Port Clarence, where Mr. Scoufield obtained specimens. We have also specimens from Petropaulski, Bering, Medni, and Robben islands and from Bristol Bay. A specimen from the island of Saghalien in Asia is in the museum at Cambridge. (*stellatus*, starry.)

*Pleuronectes stellatus*, PALLAS, Zoographia Rosso-Asiatica, III, 416, 1811, Kamchatka, Aleutian and Kuril Islands; GÜNTHER, Cat., IV, 443, 1862; STEINDACHNER, Pleur. von Decastria Bay, 1870, 1; JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1880, 453; JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1881, 68; BEAN, Proc. U. S. Nat. Mus. 1881, 420; JORDAN & GILBERT, Synopsis 835, 1883; BEAN, Proc. U. S. Nat. Mus. 1883, 353; BEAN, Cat. Coll. Fish. U. S. Nat. Mus. 1883, 20; JORDAN, Nat. Hist. Aquat. Anim., 184, pl. 46, 1884.

*Platichthys rugosus*, GIRARD, Proc. Ac. Nat. Sci. Phila. 1854, 139, 155, San Francisco; Presidio; Petaluma; GIBBARD, U. S. Pac. R. R. Surv., X, Fishes, 148, 1858.

*Platessa stellata*, DE KAY, N. Y. Fauna: Fishes, 301, 1842; STOREY, Synopsis, 478, 1846.

*Platichthys stellatus*, LOCKINGTON, Rep. Com. Fish. Cal. 1878-79, 43; LOCKINGTON, Proc. U. S. Nat. Mus. 1879, 91; JORDAN & COSS, Review Flounders and Soles, 296, 1889.

1040. MICROSTOMUS,\* Gottsche.

(SMEAR DABS.)

*Microstomus*, GOTTSCHÉ, Archiv für Naturgesch. 1835, 150 (*latidens*); not *Microstoma*, RISSO, 1826.

*Cynoglossus*, BONAPARTE, Fauna Italica, 1837, fasc., XIX (*cynoglossus*, NILSSON, not of L.).

*Cynoglossa*, BONAPARTE, Catalogo Metodico Pesci Europei, 48, 1848 (*microcephalus*); not *Cynoglossus*, HAMILTON, 1822.

*Brachyprosopeon*, BLEEKER, Comptes Rendus Acad. Sci. Amsterd., XIII, Pleuron., 7, 1862 (*microcephalus*).

*Cynoglossus*, JORDAN & GILBERT, Synopsis, 460, 1883 (*microcephalus*).

Body elongate, compressed; mouth very small; teeth broad, incisor-like, on blind side only; scales small, all cycloid; vertebrae numerous (48 to 52); dorsal rays 90 to 100; anal rays 70 to 85; anal spine obsolete; left side of skull normal, without mucous cavities; ventral fins with 5 rays each. Arctic seas. This genus is widely separated from *Pleuronectes*

\* We here retain the generic name *Microstomus*, although in accordance with recent usage of most ornithologists and ichthyologists, it should be suppressed, as identical with *Microstoma*. The two words are from the same root, and differ only in the termination. But is not this difference enough? The code of nomenclature of the American Ornithologists' Union very properly declares that "a name is only a name, and has no necessary meaning," and therefore no necessarily correct spelling, except the spelling selected by the writer from whom it dates its origin. As a result of this, the original spelling of each generic name is (undoubted misprints aside) the orthography to be adopted, regardless of all questions as to the correct etymology of the word. As a necessary sequence, it seems to us that all generic names, not actually preoccupied by names spelled in the same way, should be tenable. There is no other certain boundary line between names tenable and names untenable. We therefore regard all generic names as available unless used in zoology earlier and in exactly the same orthography. Among American genera of fishes we may therefore use the following, notwithstanding their earlier analogies:

<i>Microstomus</i> for	<i>Cynoglossus</i> notwithstanding the prior	<i>Microstoma</i> .
<i>Heterodontus</i>	<i>Cestracion</i>	<i>Heterodon</i> .
<i>Lucania</i>		<i>Lucanus</i> .
<i>Thymallus</i>	<i>Choregon</i>	<i>Thymallus</i> .
<i>Nebria</i>		<i>Nebria</i> .
<i>Xiphidion</i>	<i>Xiphister</i>	<i>Xiphidium</i> .
<i>Amitra</i>	<i>Moumitra</i>	<i>Amitrus</i> .
<i>Scytalina</i>	<i>Scytaliscus</i>	<i>Scytalinus</i> .
<i>Lagocheila</i>	<i>Quassilabia</i>	<i>Lagocheilus</i> .
<i>Auchenipterus</i>	<i>Cremnobates</i>	<i>Auchenipterus</i> .
<i>Liopsetta</i>		<i>Liopsetta</i> .
<i>Leucos</i>	<i>Myloleucus</i>	<i>Leucus</i> .
<i>Pterophrynus</i>	<i>Pterophrynoides</i>	<i>Pterophrynus</i> .
<i>Scaphirhynchus</i>	<i>Scaphirhynchops</i>	<i>Scaphirhynchus</i> .
<i>Lepidion</i>	<i>Haloporphyrus</i>	<i>Lipidia</i> .
<i>Gramma</i>		<i>Grammia</i> .
<i>Stenotomus</i>		<i>Stenotoma</i> .

If *Microstomus* be discarded, the next name in order of date is *Cynoglossus*.

The following is Bonaparte's definition of *Cynoglossus* as quoted by Gill (Proc. Ac. Nat. Sci. Phila. 1864, 222):

"Secundo è *Cynoglossus* nob. che come il *Pl. cynoglossus* L. ha in linea laterale retta, la bocca piccola, i denti come quello di sopra [*Platessa*] ma la mascelle uguale, con labbra turgide, e l'ano senza spina."

Later, in his Catalogo Metodico del Pesci Europei, Bonaparte changes this name from *Cynoglossus* to *Cynoglossa*, giving the sole species as *Cynoglossa microcephala*, and quoting as its synonym "*Pleuronectes cynoglossus*, N. Nilss.", showing that his identification of the Linnean species coincided with that of Nilsson, who at first used the name "*Pleuronectes cynoglossus*" for the present species instead of the species of *Glyptocephalus*. In Bonaparte's Catalogo, *Glyptocephalus*, Gottsche, is regarded by Bonaparte as synonymous with *Platessa*.

It is thus evident, as Dr. Gill has suggested, that Bonaparte meant to refer to the *Pleuronectes microcephalus* instead of *Pl. cynoglossus*, he "having followed Nilsson in his erroneous identification" of the latter with the former. In further evidence of this we have the fact that *Cynoglossus microcephalus* (kitt) has no anal spine, while such a spine is present in the species of *Glyptocephalus*. We would be, therefore, justified in the use of *Cynoglossus* instead of the later *Brachyprosopeon*, if *Microstomus* should be regarded as ineligibile on account of the prior name *Microstoma*. (Jordan & Goss.)

and its allies by its greatly increased number of vertebrae, a character accompanied by a similar increase in the number of fin rays. It is close to *Glyptocephalus*, but the lack of the cavernous structure of the bones of the head, a structure peculiar to the species of that genus, sufficiently distinguishes it. (*μικρός*, small; *στόμα*, mouth.)

a. Dorsal rays 85 to 93; anal rays 70 to 76; head very small, about 5 in length; eye 4 in head. KITT, 3024.

aa. Dorsal rays 102; anal ray 85; head  $4\frac{1}{2}$  in length; eye 3 in head. PACIFICUS, 3025.

3024. *MICROSTOMUS* KITT (Walbaum).

(SMEAR DAB.)

Head  $5\frac{1}{2}$  in length; depth  $2\frac{1}{2}$ . D. 85 to 93; A. 70 to 76; scales 130; caudal  $1\frac{1}{2}$  in head; pectoral  $1\frac{1}{2}$ . Body moderately elongate; mouth small, the maxillary not reaching to front of lower eye; teeth on blind side conical, rather compressed and blunted, 11 to 13 on either jaw; eyes close together, the lower slightly in advance; gill rakers short, not numerous. Origin of dorsal above middle of upper eye, its rays larger in the posterior half of body; pectorals about equal in size; no spine before anal; caudal rounded; head, except snout, entirely sealed; scales cycloid; lateral line with a small curve; vertebrae  $13 + 35 = 48$ . Color dull yellowish, blotched, and with dark spots, especially over the chest and along the base of anal fin; dark blotches and spots on anal, caudal, and ventral fins; dark base to pectoral, which has also some cloudy markings. (Day.) Seas of the north of Europe in rather deep water, south to Cornwall. Recorded by Steindachner (as *Pleuronectes gilli*), from the sea between Iceland and Greenland. This small flounder is rather common in the waters of northern Europe. It reaches the length of a foot or more, and is said to be excellent as food. Like its congener, *Microstomus pacificus*, this species is very slimy in life. *Pleuronectes gilli*, as described by Dr. Steindachner, seems to differ from *Microstomus kitt* only in the larger head, which is but  $4\frac{1}{2}$  in the length to base of caudal. It is probably not specifically distinct from the latter: Only a single specimen,  $10\frac{1}{2}$  inches long, is known. (Eu.) (The specific name "kitt," given by Walbaum on the authority of Jago's description, should be adopted for this species. According to Day, the species is still called "kitt" on the coast of Cornwall.)

*Rhombus laevis cornubiensis*, JAGO, in Ray, "Syn. Pisc., 162, tab. 1, fig. 1, 1713."

*The Smear Dab*, PENNANT, British Zoology, III, 230, pl. 41, 1776.

*Pleuronectes kitt*, WALBAUM, Artedii Piscium, III, 120, 1792, after RAY; the description in part confused with that of *Lepidorhombus*.

*Pleuronectes laevis*, SHAW, Gen'l Zool., IV, 299, 1803.

*Pleuronectes quensellii*, HÖLBÖLL, Bohusläns Fiske, IV, 59, 1821, Bohusläns, Sweden.

*Pleuronectes quadridens*, FABRICIUS, Kongl. Dansk. Vid. Selsk. Aftauidl., I, 39, 1824, Iceland.

*Microstomus latidens*, GOTTSCHKE, Archiv für Naturgesch. 1835, 150, Zealand.

*Pleuronectes gilli*, STEINDACHNER, Ichth. Notizen, VII, 40, 1868, Polar Sea north of Iceland.

*Pleuronectes microcephalus*, DONOVAN, British Fishes, II, pl. 42, 1802; GÜNTHER, Cat., IV, 447; STEINDACHNER, Ichth. Beitr., VIII, 47; DAY, Fishes Great Britain, II, 28, pl. 102;

COLLETT, Norges Fiske, 145, and of recent European writers generally.

*Pleuronectes microstomus*, FABER, Isis, 886, 1828.

*Platessa microcephala*, FLEMING, British Anim., 198, 1828, and of numerous writers.

*Cynoglossa microcephala*, BONAPARTE, Catalogo Metodico Pesci Ent., 48, 1845.

*Microstomus kitt*, JORDAN & GOSS, Review of Flounders and Soles, 1886, 298.

3025. *MICROSTOMUS PACIFICUS* (Lockington).

(SLIPPERY SOLE.)

Head  $4\frac{1}{2}$  in body; dept.  $2\frac{3}{8}$ . D. 102; A. 85; scales 140; eye  $3\frac{1}{2}$  in head; maxillary 5; pectoral  $1\frac{3}{8}$ ; greatest height of dorsal  $2\frac{1}{8}$ ; anal  $2\frac{1}{8}$ ; caudal  $1\frac{3}{8}$ ; vertebrae  $12 + 40 = 52$ . Body elongate, elliptical; mouth small, the maxillary reaching just past front of lower eye; teeth long and broad, forming a continuous cutting edge, on blind side only, about 10 teeth on lower jaw; eyes very large, nearly twice as long as snout, the upper even with profile above; interorbital a narrow scaly ridge; gill opening adnate to shoulder girdle above pectoral; gill rakers short, 8 below angle, 5 or 6 very small scarcely developed ones above; scales small, cycloid, not closely imbricated, lateral line nearly straight. Origin of dorsal slightly behind middle of upper eye, caudal truncate or slightly rounded. Color olive brown, blotched on body and fins with darker, all fins blackish toward the ends of the rays. Pacific coast of North America, Monterey to Unalaska, in rather deep water, 15 to 50 fathoms; common. Here described from a specimen, about 14 inches in length, from *Albatross* Station 2927, off the coast of California. This small flounder abounds in deep water about San Francisco, but comes near the shore farther north. It is exceedingly slimy when first taken. The large individuals are considered excellent as food; the smaller are thrown away. It rarely reaches the weight of a pound.

*Glyptocephalus pacificus*, LOCKINGTON, Rep. Cal. Com. Fisheries, 1878-79, 43, off Point Reyes, California; LOCKINGTON, Proc. U. S. Nat. Mus. 1879, 86; JORDAN, Nat. Hist. Aquat. Anim., 188, 1884.

*Cynicoglossus pacificus*, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1880, 453; JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1881, 68; JORDAN & GILBERT, Synopsis, 838, 1883.

*Microstomus pacificus*, JORDAN & GOSS, Review Flounders and Soles, 290, 1889.

1041. *EMBASSICHTHYS*, Jordan & Evermann.

*Embassichthys*, JORDAN & EVERMANN, Check-List Fishes, 506, 1896 (*bathybius*).

This genus is a deep sea representative of *Microstomus*, from which it differs in the increased number of vertebrae (63 instead of 48 to 52). Its fin rays are correspondingly increased, the body is deeper than in *Microstomus*, and it has teeth on both sides of the jaws, as in *Glyptocephalus*. (*ἐν*, in; *βάσσος*, for *βαθύς* deep; *ἰχθύς*, fish; a fish in the depths.)

3026. *EMBASSICHTHYS BATHYBIUS* (Gilbert).

Head 4 to  $4\frac{1}{2}$  in length; depth 2 to  $2\frac{1}{2}$ . D. 111 to 117; A. 96 to 98; vertebrae  $14 + 49 = 63$ . Body oval, very thin and deep, the greatest depth at anterior third of body; upper profile very abruptly angulated opposite hinder margin of upper pupil, the anterior half of head conspicuously protruding beyond general outline. Caudal nearly sessile, the peduncle very short. Mouth small, maxillary about  $\frac{1}{2}$  length of head in specimens 1 foot long. Teeth broad incisors, slightly notched at tip, nearly equally developed on blind and colored sides, 21 on blind side of lower jaw, 16 on

colored side. As in other members of this group, the lower jaw is the longer, the upper teeth included. Interorbital space wholly scaled, with a very high, rather sharp ~-shaped ridge. Eyes very large, the upper entering largely into the upper profile, the lower much in advance; front margin of upper orbit on vertical of front of lower pupil; diameter of upper eye  $2\frac{1}{4}$  to  $2\frac{3}{4}$  in head. Anterior nostrils of both sides in rather long tubes, the posterior margins produced to form short flaps. Preopercular margins adnate, as usual, concealed by scales. No conspicuous mucous excavations on blind side. Gill rakers weak and rather short, 10 or 11 on anterior of arch. Scales very small, cycloid, in about 165 cross rows, the tubes of lateral line much fewer, not regularly arranged; over 50 longitudinal rows above lateral line. Dorsal beginning over posterior edge of pupil; fins low, the highest dorsal rays behind middle of body,  $\frac{2}{3}$  length of head; caudal rounded,  $1\frac{1}{2}$  in head; pectorals 2 in head; ventrals small, each with 5 rays, as in *Microstomus pacificus*. (*Glyptocephalus cynoglossus* and *zachirus* have 6 rays in each ventral.) Color of eyed side warm brown, darker toward margins, becoming black on vertical fins; everywhere on body and fins coarsely blotched with light blue, the marks so arranged on upper and lower thirds of sides as to form 5 broad bars of bluish, alternating with those of the ground color, and corresponding above and below; lips and branchiostegal membranes black; blind side dusky brownish. This well-marked species differs from the species of *Microstomus* in its much greater depth and bright coloration, and in having teeth well developed on both sides of jaws, as in the species of *Glyptocephalus*. Two specimens from the Santa Barbara Channel, in deep water. (Gilbert.) (*βαθύς*, deep; *βίος*, life.)

*Cynoglossus bathybius*, GILBERT, Proc. U. S. Nat. Mus. 1890, 123, Santa Barbara Channel, at Albatross Station 2980, Lat.  $33^{\circ} 49' 45''$  N., Long.  $119^{\circ} 24' 30''$  W., in 603 fathoms. (Type in U. S. N. M. Coll. Gilbert.)

#### 1042. GLYPTOCEPHALUS, Gottsche.

(FLUKES.)

*Glyptocephalus*, GOTTSCHÉ, Archiv für Naturgesch. 1835, 156 (type *saxicola* = *cynoglossus*, L.).

Eyes and color on the right side. Body extremely elongate, more than twice as long as deep, much compressed. Head very small and short, its blind side with many excavations and mucous cavities in the skull, mandible, and preopercle. Mouth very small; teeth moderate, incisor-like, broad, equal, close set, in a single series; no teeth on vomer or palatines. Gill rakers short, weak. Lower pharyngeals narrow, with 1 or 2 rows of conical teeth. Lateral line nearly straight, simple; scales very small, smooth; dorsal and anal very long, there being more than 90 rays in the dorsal and more than 80 in the anal; caudal fin rounded; anal spine present; ventral rays 6. Vertebrae in increased number, 58 to 65. Northern seas, in deep water. This genus is one of the most strongly marked in the family, being distinguished from most of the genera by the greatly increased number of vertebrae, and from all of them by the remarkable cavernous

structure of the bones of the head. Two species known. (*γλυπτός*, sculptured; *κεφαλή*, head.)

a. Pectoral fins very short, not falcate, that of right side about  $\frac{1}{2}$  length of head; vertebrae 58. CYNOGLOSSUS, 3027.

aa. Pectoral fins of colored side falcate, longer than the head; vertebrae 65. ZACHIRUS, 3028.

3027. GLYPTOCEPHALUS CYNOGLOSSUS (Linnæus).

(CRAIG FLUKE; POLE FLOUNDER.)

Head 5 to 5 $\frac{1}{2}$  in body; depth 2 $\frac{1}{2}$  to 3. D. 101 to 112; A. 87 to 100; scales 125; V. 6; highest dorsal and anal rays 2 in head; pectoral a little more than 2; vertebrae 58. Body oblong, fusiform; head small, ovate; the profile slightly decurved; mouth very small, with the cleft oblique; teeth on blind side close set, with incisoral edges,  $\frac{1}{10}$ ; on the eyed side, distant, obtusely conic,  $\frac{2}{3}$ ; eyes moderate, the lower advanced, close together, 3 in head; scales regularly imbricated, lateral line straight; pectoral short, falcate; origin of dorsal above middle of upper eye; anal spine present; caudal convex or angulated behind; pectoral fins very short, not falcate, that of right side about  $\frac{1}{2}$  length of head; upper jaw with about 30 teeth; opercle adnate to the shoulder girdle for a short distance only. Color grayish brown; fins with dark spots; tip of pectoral dusky above. North Atlantic, on both coasts, chiefly in deep water, south to Cape Cod and France. This species is found in rather deep water on sandy bottoms. It reaches a length of 12 to 18 inches. This flounder has been taken in great numbers with the beam trawl in deep water off our New England coast. It is pronounced by the United States Fish Commission to be not inferior as a food-fish to the European sole. (Eu.) (*cynoglossus*, a sole; *κύων*, dog; *γλωσσο*, tongue.)

*Pleuronectes*, etc., *Corpore oblongo glabro*, GRONOW, Museum Ichthyol., i, iv, 39, etc., Belgium.

*Pleuronectes cynoglossus*, LINNÆUS, Syst. Nat., Ed. x, 269, 1758, after GRONOW; GÜNTHER, Cat., iv, 449, 1862; DAY, Fishes Great Britain, ii, 30, pl. 103; LILLJEBORG, Sveriges och Norges Fiske, ii, 386, 1891; and of European writers generally.

*Platessa pola*, CUVIER, Règne Animal, Ed. ii, Vol. 2, 339, 1829, after la Polo of DuRoiel.

*Pleuronectes saxicola*, FABER, Tidsskr. f. Naturv., 5 B., 244, 1828, Denmark.

*Pleuronectes nigromanus*, NILSSON, Prodr. Ichth. Scand., 55, 1832.

*Platessa elongata*, YARRELL, Hist. Brit. Fish., 619, 1859, young.

*Glyptocephalus acadianus*, GILL, Proc. Ac. Nat. Sci. Phila. 1873, 360, Nova Scotia. (Type, No. 12685.)

*Glyptocephalus cynoglossus*, GILL, Proc. Ac. Nat. Sci. Phila. 1873, 360; GOODE & BEAN, Proc. U. S. Nat. Mus. 1878, 21; GOODE, Proc. U. S. Nat. Mus. 1880, 337; GOODE, Proc. U. S. Nat. Mus. 1880, 475; COLLETT, Norske Nord-Havs Exped. 1880, 150; GOODE & BEAN, Bull. Mus. Comp. Zool., x, No. 5, 195, 1883; JORDAN & GILBERT, Synopsis, 838, 1883; GOODE, Nat. Hist. Aquat. Anim., 198, 1884; JORDAN & GOSS, Review Flounders and Soles, 300, pl. 19, 1889.

*Solea cynoglossa*, RAFINESQUE, Indice di Ittiologia Siciliana, 53, 1810; based on the Sole or *Cynoglossum* of RONDELET.

*Glyptocephalus saxicola*, GOTTSCHÉ, Archiv für Naturgesch. 1835, 156.

*Platessa saxicola*, KRÖYER, Danmark's Fiske, 338, 1843.

*Pleuronectes elongatus*, GÜNTHER, Cat., iv, 450, 1862.

*Glyptocephalus elongatus*, GILL, Proc. Ac. Nat. Sci. Phila. 1873, 362.

## 3028. GLYPTOCEPHALUS ZACHIRUS, Lockington.

(LONG-FINNED SOLE.)

Head  $5\frac{1}{2}$  to  $5\frac{1}{2}$ ; greatest width of body  $3\frac{1}{4}$  to  $3\frac{1}{2}$ ; eye  $3\frac{1}{2}$  in head; snout 8. D. 94 to 106; A. 79 to 89; P. 11 to 13; V. 6; vertebrae  $13+52=65$ . Body elongate-ovate, anterior portion of the oval shorter than posterior; snout declivous, almost vertical, its tip level with upper margin of lower eye, its curve uniting without sensible depression with that of nape; dorsal outline rising with a regular gentle curve from snout to about twenty-second dorsal ray, thence declining very gradually and regularly with but slight curvature to caudal peduncle; abdominal outline almost straight from knob of mandible to ventral; from thence to end of anal curved in same manner as dorsal outline; peduncle of tail expanded toward caudal, its least width about  $\frac{1}{4}$  of greatest depth of body; greatest distance from anal to lateral line less than length of head. Eyes large, elliptical, the lower in advance of the upper about  $\frac{1}{4}$  length of pupil, and scarcely reaching dorsal profile anteriorly. Interocular space very narrow, about  $\frac{1}{3}$  of longitudinal diameter of eye, smooth; not raised above the eye in a fresh fish; a slight ridge rising at its posterior part, forming lower posterior margin of upper eye, and dying out on cheek. Nostrils of right side level with upper margin of lower eye; anterior nostril with a short tube, the posterior with a raised margin, and vertical with the front margin of the lower orbit; posterior nostril of blind side in advance of eye; anterior nostril nearly as on colored side; nostrils small and inconspicuous. Gape of mouth very small on colored side, considerably larger on blind side; on the colored side the cleft is nearer vertical than horizontal; posterior end of maxillary reaching very little behind anterior margin of orbit of lower eye, and the symphysis of intermaxillaries about level with upper edge of orbit; mandible projecting in the closed mouth, short, not passing a vertical from front margin of pupil, with a prominent knob below the symphysis, and a smaller one at its posterior extremity. Teeth on both sides of jaws throughout the full length of the gape, in a single row, broad, but thick, forming a blunt, continuous edge, about 34 in lower jaw and rather fewer in the upper, in an individual  $11\frac{3}{8}$  inches long; in an example  $14\frac{1}{2}$  inches long there were 14 teeth on the colored side and 26 on the blind side of the mandible, the latter the larger; in the intermaxillaries, 13 on the colored side and 23 on the blind side; each lower pharyngeal with a double row of teeth, the inner larger than the outer; the 4 anterior teeth of outer row conspicuously larger than those following; about 12 teeth in each inner row; upper pharyngeals each with a close-set row of 6 or 7 blunt conical teeth. Branchiostegals 7; gill rakers few, flexible, very short. Dorsal commencing between front of orbit and pupil, considerably behind nostrils, long and low, forming a continuous arch of slightly greater curvature than dorsal outline, the longest rays in central portion, and ending opposite anal at about  $\frac{1}{3}$  of width of caudal peduncle from origin of caudal; anal with a horizontal spine, the first ray rather distant from the visible portion of the spine, and nearly length of ventral behind pectoral base, similar to the dorsal; almost all the rays of dorsal and anal directly backward; caudal convex on posterior margin, rather narrow, the rays once bifurcate,



sometimes bifurcate again near the tips; pectoral of colored side exceedingly long and lanceolate, about  $\frac{1}{2}$  of total length of fish; first 5 rays simple, the others once bifurcate; fourth ray longest, fifth nearly equal, sixth a little longer than third, thence diminishing rapidly. Usual proportion of the first 4 rays 3-8-10-12; pectoral of blind side lanceolate, rather more than  $\frac{1}{2}$  of length of that of colored side, and formed of the same number of rays, first 4 simple, the others once forked; fourth and fifth rays longest; ventrals inserted so that their hinder axil is vertical with, or a little posterior to, anterior axil of pectoral, their tips reaching to first anal ray; 4 posterior rays once bifurcate. Lateral line almost straight, rising very slightly anteriorly, formed of a double row of tubes, about 138 in number, excluding those upon caudal; a row of similar pores commencing at ridge under upper eye, and continuing around lower eye almost to its front margin; scales small, smooth, uniform over the body, and extending over the head to snout, on which they are smaller; intermaxillaries and mandible scaleless; scales on blind side similar; caudal scaly on both sides; no scales on the other fins. Color uniform brownish or cinereous; fins darker; the color formed by minute dark spots on the scales; membrane between the fin rays closely set with dark points; blind side whitish, the ground tint clouded with numerous black points. Deep waters of the Northern Pacific, from San Francisco northward; found throughout Bering Sea in 35 to 350 fathoms. This species is a thin, dry flounder, reaching a length of something over a foot. It is taken in the sweep nets in deep water about San Francisco. It is readily known by its long pectoral fin. ( $\zeta\alpha$ -, an intensive particle;  $\chi\epsilon\iota\rho$ , hand, from the long pectoral.)

*Glyptocephalus zachirus*, LOCKINGTON, Proc. U. S. Nat. Mus. 1879, 88, San Francisco; LOCKINGTON, Rep. Com. Fisheries California 1878-79, 42; JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1880, 453; JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1881, 68; JORDAN & GILBERT, Synopsis, 838, 1883; JORDAN, Nat. Hist. Aquat. Anim., 188, 1884; JORDAN & GOSS, Review Flounders and Soles, 301, 1889; GILBERT, Rept. U. S. Fish. Comm. 1893 (1896), 460.

#### 1043. LOPHOPSETTA, Gill.

(WINDOW PANES.)

*Lophopsetta*, GILL, Proc. Ac. Nat. Sci. Phila. 1862, 216 (*maculatus*).

Eyes and color on the left side. Body broadly ovate, strongly compressed, pellucid; mouth large, oblique, the maxillary reaching to beyond eye; teeth subequal, in narrow bands, or in single series; a small patch of teeth on the vomer. Scales small, cycloid, imbricate, the skin without bony tubercles. Lateral line strongly arched in front, without accessory branch. Dorsal fin beginning on the snout, its anterior rays exerted; anal fin not preceded by a spine; ventral of left side free from the anal, inserted nearly on the ridge of the abdomen, its base broad, the rays well separated; pectoral and ventral fins moderate. One species. Very close to the European genus *Bothus*, Rafinesque (= *Scophthalmus*, Rafinesque, = *Rhombus*, Cuvier, = *Passer*, Valenciennes), from which it differs in the more numerous gill rakers, pellucid body, and produced dorsal rays, all characters of minor importance. The European Turbot (*Psetta*, Swainson), is

also closely related, but the typical species, *Psetta maxima*, is a large robust fish, scaleless and beset with bony tubercles. (Λόφος, crest; ψήτρα, turbot.)

3029. *LOPHOPSETTA MACULATA* (Mitchill).

(WINDOW PANE.)

Head  $3\frac{1}{2}$ ; depth  $1\frac{1}{2}$ . D. 65; A, 52; scales 85; eye 4 in head; pectoral  $1\frac{1}{2}$ ; highest dorsal rays  $1\frac{1}{2}$ ; highest anal rays  $1\frac{1}{2}$ ; interorbital space  $\frac{1}{2}$  eye. Body broadly rhomboid, strongly compressed, translucent in life; mouth large, the maxillary reaching nearly to posterior margin of eye, maxillary of eyed side with a bony tubercle on its anterior end; jaws subequal, the lower with a sharp knob at symphysis; teeth in each jaw in 1 series laterally, in a very narrow band in front; interorbital space rather broad, slightly concave, its posterior third or fourth with scales; gill rakers short and slender, about 8 + 25; maxillary, mandibles, snout, and the greater part of interorbital naked; scales on head and body cycloid, loosely imbricated, those on the blind side a little smaller. Anterior rays of dorsal produced, their ends branched and free, the first on tip of snout, the rays at the beginning of posterior third of fin the highest; origin of anal directly under angle of preopercle; base of ventrals long, that of the eyed side extending along ridge of body from notch in isthmus to front of anal, base of ventral on blind side shorter; pectoral reaching past curve on eyed side, its mate much smaller; caudal rather long. Color light olive brown, almost translucent, everywhere marbled with paler, and with many small, irregular, sharply defined black spots; dorsal, anal, and caudal with larger, round, blended spots of dark brown; pectoral with brown, interrupted cross lines. This small flounder much resembles the European Brill (*Bothus rhombus*), but is smaller, thinner, and more translucent in body. Its weight rarely exceeds a pound or two, and its value as a food-fish is but slight; nevertheless, it is a near ally of the European Turbot (*Psetta maxima*), and in its technical characters it very closely agrees with the latter species. Atlantic coast of United States, from Casco Bay to South Carolina; common. (*maculatus*, spotted.)

*Pleuronectes maculatus*, MITCHILL, Rept. in part, Fish. N. Y., 9, 1814, New York; DE KAY, New York Fauna: Fishes, 301, pl. 47, fig. 151, 1842; STORER, Synopsis, 479, 1846; STORER, Hist. Fish. Mass., 204, 1867; JORDAN & GOSS, Review of Flounders and Soles, 258, 1889.

*Pleuronectes aquosus*, MITCHILL, Trans. Lit. and Phil. Soc. N. Y., 1, 1815, 389, pl. 2, fig. 3, New York.

*Lophopsetta maculata*, GILL, Proc. Ac. Nat. Sci. Phila. 1862, 216; *ibid.*, 1864, 220; JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1878, 371.

*Bothus maculatus*, JORDAN & GILBERT, Synopsis, 815, 1883.

*Rhombus aquosus*, GÜNTHER, Cat., IV, 411, 1862.

1044. *PLATOPHRYS*, Swainson.

*Solea*, RAFINESQUE, Indice di Ittiologia Siciliana, 52, 1810 (*rhomboide*); not of QUENSEL, 1806.

*Platophrys*, SWAINSON, Nat. Hist. Class'n Fishes, II, 302, 1839 (*ocellatus*).

*Peloria*, COCCO, Intorno ad Alcuni Pesci del mar di Messina, Giorn. del Gabin., 1844, 21-30, Lettere di Messina (*heckeli*, a larval form of *P. podas*); not *Pelorus* of MONTFORT, 1808.

! *Coccolus*,\* BONAPARTE, in COCCO, Alcuni Pesci Messina, 21, 1844 (*annectens*; larval form—probably of *P. podas*, with the right eye in transit to the left side).

*Bothus*, BONAPARTE, Catalogo Metodico, 49, 1846 (*podas*); not of RAFINESQUE.

*Rhomboidichthys*, BLEEKER, Act. Soc. Sci. Indo-Nederl. Manad. and Makassar, 67, 1357-58 (*myriaster*).

*Platophrys*, BLEEKER, Comptes Rendus Acad. Sci. Amsterd., XII, 1862, Pleuron., 5 (*ocellatus*).

Eyes and color on the left side. Body ovate, strongly compressed; mouth of the large type, but comparatively small; the maxillary  $\frac{1}{2}$  or less of the length of the head; teeth small, subequal, in 1 or 2 series; no teeth on vomer or palatines. Interorbital space broad and concave, broadest in adult males. Gill rakers moderate. Dorsal fin beginning in front of eye, all its rays simple; ventral of colored side on ridge of abdomen; caudal convex behind; pectoral of left side usually with 1 or more filamentous rays, longest in the male. Scales very small, ctenoid, adherent; lateral line with a strong arch in front. Coloration usually variegated.

This well-marked genus is widely diffused in the warm seas. The sexual differences are greater than usual among flounders, and the different sexes have often been taken for different species. As a rule, in the males, the pectoral fin on the left side is much prolonged, the interorbital area is much widened and very concave, and there are some tubercles about the snout and lower eye. The young fishes, as is usually the case, resemble the adult females. Lately, Dr. Emery has shown that the larval flounder, known as *Peloria heckeli*, is in all probability the young of *Pleuronectes podas*. The generic name, *Coccolus*, based on forms slightly more mature than those called *Peloria*, probably belongs here also. We have seen larval forms so young as those which have been described as *Peloria heckeli*. We have, however, examined small transparent flounders, one with the eyes quite symmetrical, taken in the Gulf Stream, and another with the eyes on the left side, taken at Key West. Both these may be larvae of *Platophrys ocellatus*. The figures published by Emery seem to make it almost certain that the corresponding European forms belong to *P. podas*, although some doubt as to this is expressed by Facciola. The species of *Platophrys* are widely distributed through the warm seas, no tropical waters being wholly without them. All the species of *Platophrys* are extremely closely related and can be distinguished with difficulty. On the other hand, the variations due to differences of age and sex are greater than in any other of our genera. The following analysis of the species of *Platophrys* is very unsatisfactory. There are certainly 3 species (*podas*, the European species, *maculifer*, and *lunatus*) which are known to be distinct in their adult state. The young forms of *maculifer* and *lunatus* are not well known, nor is it known how they differ from *ocellatus*, *spinosus*, and other species which presumably reach a smaller size. Only a thorough study of the species, in all stages of development in their native waters, can give us the characters by which the species can be really discriminated. (*πλατύς*, broad; *ὄφρὺς*, eyebrow.)

\* "Parvus mole et pleuronectiformis, medius inter Pleuronectidas et Bibroninos hic piscis videtur! Attamen dum illi oculos unilaterales habeant, iste vero bilaterales; in hoc novo genere oculi, alter a latere, alter in vertice vix ad appositum latus convenus positi sunt." (Bonaparte: quoted by Facciola, Su di Alcuni Rari Pleuronettidi.)

- a. Anal rays, at least anteriorly, each with a spinule at base (these formed by a slight widening of the tip of the interhemal spines, each being covered by a little rough scale); front of dorsal with similar projections.
- l. Color brown, with pale rounded spots; fins dotted with brown; a faint dark spot at first  $\frac{1}{4}$  of lateral line; snout with horny points; mouth small, the maxillary reaching front of eye. SPINOSUS, 3030
- aa. Anal rays without spinules at their base.
- c. Anterior profile of head convex before the interorbital area, the very short snout scarcely forming a reentrant angle at its base; form elliptic-ovate, the outlines more regular than in *P. lunatus*.
- d. Dorsal rays 85 to 95.
- e. Scales not very small, about 75 pores in lateral line; no blue markings, at least in the young.
- f. Mouth small, the maxillary 3 in head; no spines about the snout; eyes  $3\frac{1}{2}$  in length; interorbital width 3 in head (in the type); pectoral short; curve of lateral line 6 times in straight part. Color dark brown, with numerous stellate white spots, the most distinct of them with darker edgings; these generally scattered over the body; but some of them on sides of body gathered together in little rings; these spots blue rather than white in life. CONSTELLATUS, 3031
- ff. Mouth smaller, the maxillary  $3\frac{1}{2}$  in head. Color light grayish, tinged with reddish, with small round spots of darker gray, and with lighter rings inclosing spaces of ground color. OCELLATUS, 3032.
- ee. Scales smaller, 90 to 95 pores in lateral line. Color of adult, reddish gray, the body everywhere covered with rings formed of round, sky-blue spots, which are not confluent and not edged with black; besides these, very few detached spots or other blue markings. MACULIFER, 3033.
- dd. Dorsal rays 105; anal rays 80; pectoral short; interorbital space  $2\frac{1}{2}$  in head; depth  $1\frac{1}{2}$  in length; scales 91; body deep. Color (specimen  $4\frac{1}{2}$  inches long) grayish, much spotted and mottled with whitish, no blue in young example. ELLIPTICUS, 3034.
- cc. Anterior profile of head strongly concave before interorbital area, the projecting snout leaving a marked reentrant angle above it.
- j. Mouth not very small; maxillary 3 in head. Color dark olive, with many rings, curved spots, and small round dots of sky blue edged with darker on body, these largest near middle of sides, where some are as large as eye; 3 obscure dark blotches on straight part of lateral line. LUNATUS, 3035.
- gg. Mouth small; maxillary  $3\frac{1}{2}$  in head. Color highly variegated with different shades of gray, the pale blotches rounded, very irregular in size and position; no blue spots. LEOPARDINUS, 3036.

#### 3030. PLATOPHRYS SPINOSUS (Poey).

Depth  $1\frac{1}{2}$ . D. about 74; A. about 57; scales about 80. Anal rays, at least anteriorly, each with a spinule at base, these formed by a slight widening of the tips of the interhemal spines, each being covered by a little rough scale; front of dorsal with similar projections. Snout with horny points; mouth small, the maxillary reaching front of eye. Eyes very wide apart,  $2\frac{1}{2}$  in head, the interorbital space  $1\frac{1}{2}$  in head; pectoral fin short; curve of lateral line 5 in straight part. Color brown, covered with pale rounded spots; fins dotted with brown; a faint dark spot at first third of lateral line. Described from specimens from Cuba, probably the types,  $4\frac{1}{2}$  inches

long, which have been partly dried before being placed in alcohol. Cuba. The original description of this species is a very scanty one. In all respects, unless it be the color, it agrees with the European species, *Platophrys podas*. We have found 2 small specimens sent by Professor Poey to the Museum of Comparative Zoology, which may be the types of this species. They are  $4\frac{1}{2}$  inches long, and have been partly dried in the sun. A result of this has been to increase the prominence of the interorbital spines. Whether these be the original types or not, the species is an extremely doubtful one. The eyes are farther apart in these specimens than in any of *Platophrys ocellatus*, which we have examined. They agree in this respect with Agassiz's figure of *Rhombus ocellatus*. (*spinosus*, spinous.)

*Rhomboidichthys spinosus*, POEY, Synopsis, 409, 1868, Cuba; POEY, Enumeratio, 139, 1875. *Platophrys spinosus*, JORDAN & GOSS, Review Flounders and Soles, 266, 1889.

### 3031. PLATOPHRYS CONSTELLATUS, Jordan.

Head 4; depth  $1\frac{1}{2}$ ; eye  $3\frac{1}{2}$  in head; interorbital width 3. D. 89; A. 65; scales 75. Body elliptic-ovate, the outlines more regular than in *P. lunatus*; anterior profile of head convex before the interorbital area, the very short snout scarcely forming a reentrant angle at its base; anal rays without spinules at their base; mouth small, the maxillary 3 in head; no spines about the snout; pectoral short; curve of lateral line 6 times in straight part. Color dark brown, with numerous stellate white spots, the most distinct of them with darker edgings; these generally scattered over the body, but some of them on sides of body are gathered together in little rings (perhaps these spots are blue rather than white in life); fins mottled with dark brown, the pectoral finely barred. Specimens examined  $3\frac{1}{2}$  inches long. Galapagos Archipelago. Originally described from 3 specimens, the largest  $3\frac{1}{2}$  inches long, numbered 11146 on the register of the Museum of Comparative Zoology. They are from James Island, in the Galapagos. The species is closely related to *P. ocellatus* and others, but in color, at least, it is different, and its habitat is remote; locally common. (*constellatus*, with star-like spots.)

*Platophrys constellatus*, JORDAN, in JORDAN & GOSS, Review Flounders and Soles, 266, 1889, James Island, Galapagos Archipelago. (Types in M.C.Z.)

### 3032. PLATOPHRYS OCELLATUS (Agassiz).

Head 4 in length; depth  $1\frac{1}{2}$ ; eye (lower)  $3\frac{1}{2}$  in head; snout 5. D. 85; A. 64; scales 75 (pores); vertebrae 37. Body ovate, deep anteriorly, the profile descending steeply, rendered abruptly concave in front of interorbital space by the conspicuously projecting short snout. Mouth very small and oblique, the maxillary reaching vertical from front of lower eye,  $3\frac{1}{2}$  in head; tip of lower jaw entering the profile. Teeth fine, conical, in 2 series in the upper jaw, 1 in the lower, those of the outer row in upper jaw larger and more widely separated than those of the inner series. Snout very short, equaling interorbital width. Interorbital space narrow, deeply concave, closely scaled. Eyes large, the lower in advance of upper. Gill rakers obsolete, 7 rudiments on horizontal branch of anterior arch. Scales moderate, not extending on the fins, those on colored side

ctenoid, those on blind side smooth; arch of lateral line short and high, its base contained  $4\frac{1}{2}$  to 5 times in the straight portion. Dorsal fin beginning opposite anterior nostril, the rays nearly uniform in length, the longest about  $\frac{1}{2}$  head; pectoral of colored side  $4\frac{1}{2}$  in length; ventral of colored side beginning under middle of lower eye, with 6 rays; the right ventral with 5 rays. Color in life, light grayish with reddish tinge, covered with small round spots of darker gray and with lighter rings inclosing spaces of the ground color; vertical fins similarly colored, with a small black spot near base of each ninth or tenth ray; 2 black spots on median line of body divide the length into nearly equal thirds; some other small black spots scattered over colored side. Western Atlantic, from Long Island to Rio Janeiro, on sandy shores. Here described from Key West specimens, types of *P. nebularis*. This species is very common at Key West in clear, shallow water on sandy bottom. The largest of the numerous specimens taken is 3 inches in length. A specimen similar to those has been taken by Dr. Bean on the south coast of Long Island. This seems to be the same as the Cuban species called *Rhomboidichthys ocellatus* by Poey, and some of the specimens sent by Poey to the Museum of Comparative Zoology are apparently identical with the types of *P. nebularis*. In the Museum of Comparative Zoology we have compared specimens of the real *Platophrys ocellatus* (No. 11423, Rio Janeiro, Agassiz) with a representative specimen of *P. nebularis* (No. 26147, from the Tortugas, Florida), and are unable to find any differences. We adopt, therefore, the name *Platophrys ocellatus* for all, and regard it as one of the widely distributed flounders, like *Etropus crossotus* and *Citharichthys spilopterus*. (*ocellatus*, with eye-like spots.)

*Rhombus ocellatus*, AGASSIZ, Spix, Pisc. Brasil., 85, pl. 46, 1829, Brazil.

*Rhombus bahianus*, CASTELNAU, Anim. nouv. rares Amérique du Sud, 1855, Bahia.

*Platophrys nebularis*, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1884, 31, 143, Key West (Type, 34972. Coll. Dr. Jordan); GOODE & BEAN, Oceanic Ichthyology, 441, 1896.

*Platophrys ocellatus*, SWAINSON, Nat. Hist. Class'n Fishes, II, 302, 1839; JORDAN & GOSS, Review Flounders and Soles, 266, 1889.

*Rhomboidichthys ocellatus*, GÜNTHER, Cat., IV, 433, 1862; POEY, Synopsis, 408, 1868.

### 3033. PLATOPHRYS MACULIFER (Poey).

Head 4; depth  $1\frac{1}{2}$ . D. 90 to 95; A. 70; scales 90 to 95. Body elliptical, ovate. Mouth small, oblique, the maxillary  $3\frac{1}{2}$  in head; teeth in each jaw in 2 irregular series; filamentous rays of pectorals reaching very nearly to last rays of dorsal; arch of lateral line short and high, its length  $1\frac{1}{2}$  times its height and  $2\frac{1}{2}$  in head; snout very short, 4 in head; interorbital area  $3\frac{1}{2}$  in head. Color of adult reddish gray, the body everywhere covered with rings formed of round, sky-blue spots, which are not confluent and are not edged with black; besides these, very few detached spots or other blue markings; head with similar blue spots, but no rings; area inclosed in the blue rings not different from the ground color; caudal with blue spots, other fins with none; dorsal and anal mottled; a large, diffuse, dusky spot at front of straight part of lateral line; 1 better defined on middle of lateral line; a faint one farther back; pectorals grayish, with dark bars. Cuba. We identify specimens taken by Dr.

Jordan at Havana with this species. In the Museum of Comparative Zoology are other specimens similar to these, sent to Cambridge by Poey. In several respects these specimens agree fairly with Poey's *P. ellipticus*, but that species is said to have 104 dorsal rays. (*macula*, spot; *fero*, I bear.)

? *Pleuronectes maculiferus*, POEY, Memorias, II, 316, 1860, Cienfuegos. (Coll. Poey.)

? *Rhomboidichthys maculiferus*, POEY, Synopsis, 408, 1868; POEY, Enumeratio, 139, 1875.

*Platophrys maculifer*, JORDAN & GOSS, Review Flounders and Soles, 267, 1889.

*Platophrys ellipticus*, JORDAN, Proc. U. S. Nat. Mus. 1886, 51; not of POEY.

3084. PLATOPHRYS ELLIPTICUS (Poey).

Depth 1½. D. 105; A. 80; scales 91. Body elliptical, ovate; anterior profile of head convex before the interorbital area; pectoral short; interorbital space 2½ in head; body deep. Color (specimen 4½ inches long) grayish, much spotted and mottled with whitish; no blue (in young example). Cuba. Poey describes his *P. ellipticus* as having 104 dorsal rays. In none of our other species does the number of these rays reach 100. Among the specimens sent by Poey to the museum at Cambridge is 1, described above, 4½ inches long, which has 105 dorsal rays. We have therefore assumed that the species to which this specimen belongs is the real *P. ellipticus*, and that the one heretofore called *P. ellipticus* is Poey's *P. maculifer*. Both these assumptions are open to considerable doubt. (*ellipticus*, elliptical.)

? *Pleuronectes ellipticus*, POEY, Memorias, II, 315, 1860, Cuba. (Coll. Poey.)

? *Rhomboidichthys ellipticus*, GÜNTHER, Cat., IV, 434, 1862; POEY, Synopsis, 408, 1868; POEY, Enumeratio, 139, 1875.

*Platophrys ellipticus*, JORDAN & GOSS, Review Flounders and Soles, 267, 1889.

3085. PLATOPHRYS LUNATUS (Linnaeus).

(PEACOCK FLOUNDER.)

Head 3½ in length; depth 2. D. 93; A. 70; scales 90; lower eye 6 in head; maxillary 2½; interorbital 2½; highest dorsal rays 2½; highest anal rays 2½; caudal 1½; base of ventral of eyed side 3½. Vertebrae 9 + 30 = 39. Body elliptical, ovate, strongly compressed; anterior profile concave, the snout projecting, leaving a reentrant angle above it; mouth moderate, the maxillary reaching to middle of pupil of lower eye; jaws subequal, the lower with a well-developed knob at symphysis, teeth small, in an irregular double series in each jaw; anterior end of maxillary with a large blunt spine, pointing outward and forward, a smaller one behind it on upper edge of maxillary, pointing upward and backward; interorbital very wide and deeply concave; orbital rim, below on upper orbit, above on lower, broken up into blunt papillæ; gill rakers short and thick, 9 developed on lower part of arch, none on upper. Anterior part of interorbital, snout, maxillary, and mandible, naked; scales all cycloid; the rays of dorsal and anal with scales, a few on ventral of eyed side; arch of lateral line 5 in straight part. Pectoral of eyed side filamentous, reaching to base of caudal, its mate of opposite side shorter, about 1½ in head; origin of dorsal over snout; ventral of eyed side with a long base, extending from

angle at isthmus, along ridge of body, slightly past front of anal; base of ventral of blind side  $\frac{1}{4}$  the length of that of its mate; caudal with the middle rays produced, double convex. Color dark olive, with many rings, curved spots, and small round dots of sky blue edged with darker on body, these largest near middle of sides, where some are as large as the eye; 3 obscure dark blotches on straight part of lateral line; head and vertical fins with sharply defined blue spots, which are mostly round; spots on opercles larger and curved; pectorals with dark bars. West Indies, north to Florida; common. Here described from a specimen from Green Turtle Cay, Florida, 14 inches in length. This handsome and curiously colored species is not rare in the waters of the West Indies. The specimens examined by us are from Cuba, Sombrero, St. Thomas, and other localities in the West Indies. The original figure of this species published by Catesby is a very good one and leaves no room for doubt as to the species intended. The figure of Bloch, called *Pleuronectes argus*, is also fairly accurate, and can refer to no other species. This species reaches a length of some 18 inches, and is the largest in size of the American species of *Platophrys*. We have never seen any young examples which certainly belong to it, and till its development is traced some of the species known from small examples only must be doubtful. (*lunatus*, crescent-shaped, from the spots.)

*Solea lunata et punctata* (the Sole), CATESBY, Nat. Hist. Carolina, tab. 27, 1725, Bahamas. *Pleuronectes lunatus*, LINNÆUS, Syst. Nat., Ed. x, 269, 1758, Bahamas; based on CATESBY; and of the various copyists.

*Pleuronectes argus*, BLOCH, Ichthyol., tab. 48, 1783, Martinique; after Plumier.

?*Pleuronectes surinamensis*, BLOCH & SCHNEIDER, Syst. Ichth., 156, 1801, Surinam; "*satis parva et glabra*;" fins scaly; mouth small; lateral line arched in front; D. 96; A. 55.

*Rhomboidichthys lunatus*, GÜNTHER, Cat., IV, 433, 1862; POEY, Synopsis, 408, 1868.

*Rhomboidichthys lunulatus*, POEY, Enumeratio, 138, 1875.

*Platophrys lunatus*, JORDAN, Proc. U. S. Nat. Mus. 1886, 51; JORDAN & GOSS, Review Flounders and Soles, 267, 1889.

#### 3036. PLATOPHRYS LEOPARDINUS (Günther).

Head  $3\frac{1}{2}$  in length; depth  $1\frac{1}{2}$ ; eye (lower)  $3\frac{1}{2}$  in head. D. 86 to 88; A. 64 (62 to 66); scales about 80. Mouth very small, the maxillary  $3\frac{1}{2}$  in head; teeth very small, biserial above. Interorbital space concave, rather broad, its width  $3\frac{1}{2}$  in head. Eyes large, the lower considerably before the upper. Lateral line with a short sharp curve anteriorly. Gill rakers very small. Anterior rays of dorsal not elevated; left pectoral not produced, little longer than right,  $1\frac{1}{2}$  in head. Coloration highly variegated with different shades of gray, the pale blotches rounded, very irregular in size and position; no distinct black spots along the lateral line; a large whitish cloud between the eyes; blind side pale, scaled like the eyed side. Gulf of California. This species is known only from the original type from unknown locality, and from a single specimen,  $2\frac{1}{2}$  inches long, in the United States National Museum, taken by Mr. H. F. Emeric, at Guaymas, Sinaloa. From this the above description was taken. (*leopardinus*, leopard-like.)

*Rhomboidichthys leopardinus*, GÜNTHER, Cat. Fish., IV, 434, 1862, locality unknown.

*Platophrys leopardinus*, JORDAN, Proc. U. S. Nat. Mus. 1884, 260, specimen from Guaymas; JORDAN & GOSS, Review of Flounders and Soles, 268, 1889.



## 1045. PERISSIAS, Jordan &amp; Evermann.

*Perissias*, JORDAN & EVERMANN, new genus (*tænopterus*).

This genus differs from *Platophrys* in the larger scales, narrower interorbital, and especially in the greatly produced ribbon-like lobe at the front of the dorsal. From *Engyprosope* it differs in the short thick gill rakers and in the produced dorsal rays. The lateral line is wanting on the blind side. Deep sea. (*περισσόος*, strange.)

## 3037. PERISSIAS TÆNIOPTERUS (Gilbert).

Head  $3\frac{3}{4}$ ; depth  $2\frac{1}{4}$ . D. 86 to 88; A. 67 to 70; scales 60 to 65, the arch with 15 pores; 20 scales in a series running upward and backward from lateral line. Body elongate; caudal fin subsessile, the last anal and dorsal rays inserted near rudimentary caudal rays; height of caudal peduncle 1 in height of body; upper profile descending very obliquely anteriorly, a slight reentrant angle in front of lower eye; in males the profile slightly angulated in front of upper orbit, below which it ascends more steeply; lower eye much in advance of upper; in females eyes close together, the vertical from middle of lower eye passing through front of upper orbit; diameter of upper orbit  $3\frac{1}{4}$  in head; in males the lower eye may be entirely in advance of upper; in females 3 inches long, and in very young males the interorbital space is a narrow, concave, scaleless groove, less than diameter of pupil, running into a deep pit behind lower eye; in males 2 inches long the interorbital space has already widened, and in specimens  $3\frac{1}{4}$  inches long is as wide as longitudinal diameter of orbit; it is traversed by an oblique ridge running upward and backward from front of lower eye, separating the anterior scaleless portion from the deep scaleless pit behind; supraorbital ridge of lower eye serrated, forming a strong series of spines, less marked in females; anterior rim of upper orbit similarly but less strongly marked; a strong double spine on maxillary in front of nostrils; a spine near end of maxillary in males; mouth small, maxillary not reaching front of pupil, equaling diameter of orbit; teeth small, in a single close-set series in each jaw, equally developed on both sides, with enlarged canines; gill rakers very short and weak, 8 on horizontal limb; anterior nostrils with very short flaps; dorsal beginning above front of lower eye; in all specimens, females as well as males, the first 2 rays detached from the rest of fin, the second ray produced into a flat, ribbon-shaped filament about as long as head; dorsal and anal rays all unbranched; median caudal rays forked; no anal spine; ventral of colored side on ridge of abdomen, the 2 anterior rays in males connected by membrane at base only, produced into flat filaments as long as head, extending far beyond front of anal; pectoral of left side well developed, but small, slightly more than  $\frac{1}{2}$  head; that of blind side little developed, about  $\frac{1}{4}$  diameter of orbit; scales of left side strongly ctenoid, absent on interorbital space, snout, maxillary, and mandible; lateral line with strong curve anteriorly, the cord of which is contained 5 times in straight portion. Along lateral line are occasional broad cutaneous flaps, colored blue in life; scales of blind side cycloid, the tubes of lateral line obsolete, the course of lateral

line indistinctly indicated by pits at bases of scales and occasional pore-like markings; median rays of dorsal and anal on left side with series of ctenoid scales, otherwise scaleless; caudal rays with double series on both sides. Color on left side olive brown, with many small irregular spots of light gray, with darker border; 3 or 4 dark blotches along lateral line; along dorsal and ventral outlines about 5 pairs of light spots, broadly ocellated with blackish; males with a bright blue spot on anterior profile at base of each of first 10 or 12 dorsal rays and 1 on end of snout; blind side in males with a broad oblique bar covering about  $\frac{1}{2}$  of sides, bluish black in life, dark brown in spirits; from its upper anterior part a number of narrow parallel streaks run forward toward head, much as in *Engyophrys sancti-laurentii*; filamentous rays of dorsal and ventral white; fins all speckled; a small black spot at base of median caudal rays. Differing from all known species of *Platophrys* in the ribbon-shaped prolongations of second dorsal ray and first and second ventral rays of eyed side, and in the obsolete lateral line of blind side. Several specimens from the Gulf of California and the western coast of Lower California, in 40 fathoms. (Gilbert.) (*ταινία*, ribbon; *πτερόν*, fin.)

*Platophrys tenuiopterus*, GILBERT, Proc. U. S. Nat. Mus. 1890, 118, Gulf of California, north of La Paz, at Albatross Station 2998, Lat.  $24^{\circ} 51' N.$ , Long.  $110^{\circ} 39' W.$ , in 40 fathoms. (Type, No. 43095. Coll. Gilbert.)

#### 1046. ENGYOPHRYS, Jordan & Bollman.

*Engyophrys*, JORDAN & BOLLMAN, Proc. U. S. Nat. Mus. 1889, 176, (*sancti-laurentii*).

This genus is allied to *Platophrys*, Swainson, but differs from it in having the interorbital space very narrow and armed with a spine, and the scales of moderate size and ctenoid. Gill rakers obsolete. No anal spine. Gill membranes entirely separate. It is still nearer the genus *Engyprosopon*, Günther, but in that group the interorbital space is broader and the gill rakers are developed and slender. (*ἐγγύς*, near together; *ὄφρυς*, eyebrow.)

#### 3038. ENGYOPHRYS SANCTI-LAURENTII, Jordan & Bollman.

Head  $2\frac{1}{2}$  to  $2\frac{1}{2}$  (3 to  $3\frac{1}{2}$ ); depth  $1\frac{1}{2}$  to 2 (2 to  $2\frac{1}{2}$ ). D. 78 to 85; A. 68 to 72; scales 60 to 68, along lateral line. Body broadly ovate, much compressed; the greatest depth over pectorals; dorsal and ventral outlines equally curved; profile scarcely concave before eyes. Mouth very small, oblique, the maxillary reaching opposite pupil of lower eye, 4 to  $4\frac{1}{2}$  in head. Teeth present on blind side, well developed, close set and even, none on vomer. Snout short,  $4\frac{1}{2}$  to 5 in head. Interorbital space a very narrow, sharp, scaleless ridge, the ridge forking above pupil, leaving a very narrow concavity anteriorly; lower ridge armed with a strong spine, turned backward, inserted just above pupil of lower eye. Anterior orbital rim of upper eye rather high, entering profile. Eyes large, lower in advance of upper,  $3\frac{1}{2}$  to 4 in head. Gill rakers almost obsolete, represented by 5 or 6 small fleshy papillae. Scales moderately small, ctenoid, and not very firmly attached; small scales on rays of dorsal and

anal fins; arch of lateral line short and small, but abrupt, 4 to 5 times in straight part. Dorsal beginning on blind side just behind posterior nostril and in front of eye; pectoral of colored side 2 in head, that of blind side  $2\frac{1}{2}$  in head; ventrals of colored side slightly longest, 3 in head; that of colored side with 6 rays, of blind side with 5 or 6 rays. Color of left or eyed side, blackish brown, with scattered white and black spots, the latter most prominent along base of dorsal and anal fin; 3 large, black, noncellated blotches on straight part of lateral line, the first at beginning, second at middle, and third on peduncle; fins dusky; dorsal and anal with scattered white and black spots; caudal with 5 black spots arranged in a curved series; blind side with 5 or 6 curved parallel dusky bands as wide as eye, the first beginning on interopercle and curving across cheek to along base of dorsal; second beginning at throat and curving along posterior margin of preopercle, and extending on back, parallel with the first from vent; third curving around in front of pectorals, across posterior part of opercle, and extending to base of dorsal fin behind the middle; rest behind pectorals. All of these bands fade out behind middle of body, so that the posterior portion is immaculate. In young examples these bands are very faint or obsolete. Coast of Colombia, southwest of Panama. Numerous specimens, the largest about  $4\frac{1}{2}$  inches long, were dredged at Albatross Station 2795, at a depth of 33 fathoms, and at Albatross Station 2805 at a depth of  $51\frac{1}{2}$  fathoms. This peculiar species is distinguished from the species of *Platophrys* and *Engyprosopon* by its very narrow interorbital ridge, from the species of *Arnoglossus* by the form of the body, the short gill rakers, etc., and from all related species by the peculiar coloration of the blind side. (Named for St. Lawrence, in allusion to the gridiron-like markings of the blind side.)

*Engyphrys sancti-laurentii*, JORDAN & BOLLMAN, Proc. U. S. Nat. Mus. 1889, 176, Pacific Ocean, off coast of Colombia, at Albatross Station 2805, Lat.  $7^{\circ} 56' N.$ , Long.  $79^{\circ} 41' 30'' W.$ , and Station 2795, Lat.  $7^{\circ} 57' N.$ , Long.  $78^{\circ} 55' W.$  (Type, No. 41155.)

#### 1047. TRICHOPSETTA, Gill.

*Trichopsetta*, GILL, Proc. U. S. Nat. Mus. 1888, 603 (*ventralis*).

Body ovate, covered with rather large, ctenoid adherent scales; mouth moderate, the chin prominent; vomer toothless; teeth small, somewhat enlarged and hooked in front, uniserial; maxillaries obliquely truncated behind; interorbital area a narrow ridge, with a median groove in front; none of the dorsal rays produced; ventrals free from the anal; caudal fin subsessile; both pectoral fins present; right ventral much produced, the left on the ridge of the abdomen; lateral line with a strong arch in front. (*οπις*, hair; *ψηττα*, turbot, from the prolonged ventral.)

#### 3089. TRICHOPSETTA VENTRALIS (Goode & Bean).

Head 4 in body; depth  $2\frac{1}{2}$ . D. 93; A. 73; pectoral 11 (eyed side), 7 or 8 (blind side); scales 19-66-23; eye  $3\frac{1}{2}$  in head; maxillary scarcely 2; interorbital very narrow, scaleless, its width 8 in eye; scales strongly ctenoid; dorsal beginning upon snout upon the blind side, in advance of eyes, its

highest rays equaling length of mandible; origin of anal under base of pectoral, its longest ray equaling or slightly exceeding  $\frac{1}{2}$  the distance of its anterior ray from snout; caudal equal to length of head without snout; pectorals inserted considerably below origin of lateral line, close to gill opening, that of the eyed side 6 in length of body; that of the blind side almost as long as head. Color light brownish gray; a dark blotch as long as eye on the anterior rays of the anal, a few obscure on different parts, of lighter hue at the junction of the curved and straight portion of the lateral line. (Goode & Bean.) Deep waters of the Gulf of Mexico. (*ventralis*, pertaining to the ventrals.)

*Citharichthys ventralis*, GOODE & BEAN, Proc. U. S. Nat. Mus. 1885, 592, deep waters of Gulf of Mexico. (Coll. *Albatross*.)

*Arnioglossus ? ventralis*, JORDAN & GOSS, Review Flounders and Soles, 262, 1889.

*Trichopsetta ventralis*, GOODE & BEAN, Oceanic Ichthyology, 440, pl. 109, fig. 372, 1896.

#### 1048. SYACIUM, Ranzani.

*Syacium*, RANZANI, Novis Specibus Piscium, Diss. Sec., 20, 1840 (*micrurum*).

*Hemirhombus*, BLEEKER, Comptes Rendus Acad. Sci. Amsterd., XIII, Plenon., 4, 1862 (*ytineënsis*).

*Aramaca*, JORDAN & GOSS, in JORDAN, Cat. Fish. N. A., 133, 1885 (*pætulæ*).

Body elliptic-ovate, much compressed; interorbital space broad in the males and more or less concave, narrowed in the female; mouth moderate, the gape curved; teeth in the upper jaw biserial, in the lower uniserial; the front teeth of the upper jaw enlarged; vomer toothless; scales rather large, ciliate; lateral line without arch in front; pectoral fins on both sides present; septum of gill cavity below gill arches without foramen; a deep emargination near the isthmus; gill rakers short and thick; dorsal low, its anterior rays not elevated; pectorals both present; caudal sessile; no anal spine; pectorals produced in the males; ventral fins short, that of colored side on ridge of abdomen. This genus contains a considerable number of species, mostly American and African, which form a transition from *Platophrys* to *Citharichthys*. They fall readily into 2 groups distinguished by the width of the interorbital space. As this width is dependent on age, and as it is subject to various intergradations, the group *Aramaca* founded on it can not be admitted as a distinct genus. (*συάκιον*, diminutive of *σῦαξ*, a kind of pulse, the application unexplained.)

a. Snout and orbits without spines or spinous processes.

b. Scales rather large, 50 to 57 in the lateral line; interorbital space broad. Color nearly plain brown, with darker dots or mottlings, no ring-like spots or ocelli; fins mottled; left pectoral barred; blind side sometimes wholly or partly dusky, especially in northern specimens. PAPILLOSUM, 3040.

bb. Scales rather small, 58 to 70 in the lateral line.

c. Scales 65 to 70. Color dark brown, with many rings and spots of light gray and blackish, some of the dark rings with a black central spot; a diffuse dusky blotch on lateral line above pectoral, and 1 near base of caudal peduncle; fins with numerous inky spots and dark markings; blind side pale. MICRURUM, 3041.

cc. Scales 58 to 60.

d. Interorbital space in male broader than eye. Color light brown, with grayish and light bluish dots, some darker areas, and a few round brown spots ocellated with lighter; interorbital space with a vertical brown bar bordered by lighter; fins mottled and spotted.

LATIFRONS, 3042.

dd. Interorbital space not broader than pupil. Color light olive brown, nearly uniform, the vertical fins with elongate dark spots.

OVALE, 3043.

3040. SYACIUM PAPILLOSUM (Linnaeus).

Head  $3\frac{1}{2}$  in length; depth  $2\frac{1}{2}$ . D. 82; A. 63 to 70; scales 53; eye 5 in head; maxillary  $2\frac{3}{4}$ ; pectoral of eyed side  $1\frac{1}{2}$ ; caudal  $1\frac{1}{2}$ . Body elliptic-ovate, the anterior profile regularly decurved, forming an angle above the snout; mouth rather large, arched; maxillary extending to below middle of eye, its posterior end concave; teeth in upper jaw in 2 series, some of the outer forming small canines; lower teeth in 1 row; eye large, 4 in head; lower eye in advance of upper, especially in the adult; interorbital space broad, concave, greater than the long diameter of the eye in the males, about equal to the vertical diameter in the females; accessory scales very numerous; mandible, maxillary, and interorbital with scales; gill rakers short, scarcely as long as pupil, about  $2 + 8$ ; dorsal rather low, beginning slightly in front of lower eye, the first 3 or 4 rays on blind side, the anterior rays produced beyond the membrane; ventrals with moderate base, that of eyed side on ridge of body, that of blind side slightly in advance of its mate; anal beginning a little in advance of pectoral; pectoral of eyed side pointed behind, the upper rays filamentous (at least in the male); caudal double truncate. Vertebrae  $10 + 26 = 36$ . Color nearly plain brown, with darker dots or mottlings, no ring-like spots or ocelli; fins mottled; left pectoral barred; blind side sometimes wholly or partly dusky, especially in northern specimens. Charleston to Rio Janeiro, in rather deep water. Here described from an adult specimen from Charleston, a foot in length. Of the species found in the deep waters about Pensacola, and called by Dr. Bean *Hemirhombus patulus*, we have numerous specimens. Lately we have received from Mr. Charles C. Leslie, of Charleston, a specimen which shows its presence also in Carolina waters. It has not yet been recorded from Cuba, but in the Museum of Comparative Zoology is a specimen (26104) taken by Mr. Samuel Garman, at Kingston, Saint Vincent. But its range extends much farther to the southward, for among the collections made by Professor Agassiz, at Rio Janeiro, there are many specimens (11375, 4666), the largest about a foot long. These seem to be completely identical with Florida examples, differing only in having the blind side pale, it being usually partly blackish in northern samples. These Brazilian specimens agree very closely with the figure of *Rhombus soleaeformis*, except that Agassiz has represented that species as having a dusky blotch at the shoulder. No such marking is apparent in any of our specimens. The coloration and the breadth of the interorbital both render it unlikely that Agassiz's *soleaeformis* could have been *micrurum*. The *Aramaca* of Marcgrave, which is the sole basis of *Pleuronectes papillosus*, *Pleuronectes macrolepidotus*, and *Rhombus aramaca*, can not well be any known species other than the present one. According

to Maregrave's rude figure and his description, this species has the form of a sole, the eyes wide apart, the left pectoral produced, the mouth very large, the body oblong, and the coloration stone-like (sand color) on the left side and white on the eyed side. *Syacium micrurum* is not colored in that way, and its eyes are not noticeably far apart. We therefore adopt for this species the oldest name, *Syacium papillosum*. (*papillosum*, having papillae.)

*Aramaca*, MARCGRAVE, Hist. Brasil., 181, 1648, Brazil.

*Pleuronectes papillosum*, LINNÆUS, Syst. Nat., x, 271, 1758, Brazil; based on MARCGRAVE.

? *Pleuronectes macrolepidotus*, BLOCH, Ausländische Fische, vi, 25, tab. 190, 1787; apparently based on MARCGRAVE.

*Pleuronectes aramaca*, DONNDORF, Beiträge zur Ausgabe des Linnæischen Natursystems, xiii, 386, 1798; after MARCGRAVE.

*Rhombus aramaca*, CUVIER, Règne Animal, Ed. 2, ii, 341, 1829; after MARCGRAVE.

*Rhombus soleaformis*, AGASSIZ, Spix, Pisc. Brasil., 86, tab. 47, 1829, Atlantic Ocean.

*Hippoglossus intermedius*, RANZANI, Novis Speciebus Piscium Dissertatio Secundo, 1840, 14, pl. 4, Brazil.

*Hemirhombus soleaformis*, GÜNTHER, Cat. Fish., iv, 423, 1862.

*Hemirhombus pætilus*, BEAN MS., JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1882, 304, Pen-sacola (Coll. Silas Stearns); GOODE & BEAN, Proc. U. S. Nat. Mus. 1882, 414; BEAN, Cat. Coll. Fish. U. S. Nat. Mus. 1883, 45.

*Citharichthys aramaca*, JORDAN & GILBERT, Synopsis, 816, 1883.

*Citharichthys pætilus*, JORDAN & GILBERT, Synopsis, 964, 1883; JORDAN, Proc. U. S. Nat. Mus. 1884, 38; GOODE & BEAN, Oceanic Ichthyology, 448, pl. 109, fig. 373, 1896.

*Aramaca papillosa*, JORDAN, Proc. U. S. Nat. Mus. 1886, 602; synonymy confused with *S. micrurum*.

*Aramaca soleaformis*, JORDAN, Proc. U. S. Nat. Mus. 1886, 602.

*Syacium papillosum*, JORDAN & GOSS, Review Flounders and Soles, 268, 1889.

#### 3041. SYACIUM MICRURUM, Ranzani.

Head  $3\frac{1}{2}$  in length; depth  $2\frac{1}{2}$ . D. 87 to 92; A. 54 to 68; scales 65 to 70 (pores); eye 4 in head; maxillary  $2\frac{1}{2}$  to 3. Form regularly elliptical, the profile evenly convex to end of snout; eyes large, nearly even in front, the male with the interorbital space deeply concave, its width  $\frac{2}{3}$  the vertical depth of the eye (or more in Brazilian specimens); female with interorbital area much narrower, with a more or less perfect median groove, its width about equal to depth of pupil; mouth small, the maxillary reaching to below middle of eye; teeth small, slender, in 2 rows above, in 1 row below, the outer series in upper jaw somewhat enlarged, but hardly canine-like; gill rakers very short and thick, about 1 + 7 in number. Scales small, firm, moderately ctenoid; pectoral  $1\frac{1}{2}$  in head in the female, reaching nearly to base of caudal in the male; vertebrae 9 + 24 = 33. Color dark brown, with many rings and spots of light gray and blackish, some of the dark rings with a black central spot; a diffuse dusky blotch on lateral line above pectoral, and 1 near base of caudal peduncle; fins with numerous inky spots and dark markings; blind side pale. West Indian fauna, Key West to Rio Janeiro; rather common. We have found in the Museum of Comparative Zoology specimens purporting to be the types of *Hemirhombus ocellatus*, Poey (No. 11144; Poey's number, 88). These are female examples, and they differ from the types of *Hemirhombus athalion*, obtained in Cuba by Dr. Jordan, only in their greater size. Numerous

specimens (11373) from Rio Janeiro belong to the same species. Among these are males, which have the interorbital space much broader than in the types of *ocellatus* and *athalion*. Besides these specimens we have examined others from Hayti, Cuba, and Key West, and there can be no reasonable doubt of their identity, and that all are identical with Günther's *Hemirhombus aramaca*. This fish is described and fairly well figured by Ranzani under the name of *Syacium micrurum*. It is the type of his genus *Syacium*, a generic name which, strangely enough, has received no notice from subsequent authors until lately. (*μικρός*, small; *ὄψα*, tail.)

*Syacium micrurum*, RANZANI, Nov. Spec. Pisc. Diss. Sec., 20, pl. 5, 1840, Brazil; JORDAN & GOSS, Review Flounders and Soles, 269, 1889.

*Hypoglossus ocellatus*, POEY, Memorias, II, 314, 1860, Cuba.

*Hemirhombus aramaca*, GÜNTHER, Cat., IV, 42, 1862, Cuba; Jamaica; not *Rhombus aramaca*, CUVIER.

*Citharichthys athalion*, JORDAN, Proc. U. S. Nat. Mus. 1886, 52, Havana. (Type, No. 37748. Coll. D. S. Jordan.)

*Hemirhombus ocellatus*, POEY, Synopsis, 407, 1868; POEY, Enumeratio, 138, 1875.

*Citharichthys ocellatus*, JORDAN & GILBERT, Synopsis, 964, 1883; JORDAN, Proc. U. S. Nat. Mus. 1884, 143.

*Hemirhombus athalion*, JORDAN, Proc. U. S. Nat. Mus. 1886, 602.

#### 3042. SYACIUM LATIFRONS (Jordan & Gilbert).

Head 4; depth  $2\frac{1}{2}$ . D. 92; A. 72; scales 60. Body elliptical, the dorsal and ventral outlines equally arched; mouth placed low, below axis of body; snout with an abrupt constriction in front of upper orbit, the outline then again convex; eyes on left side, distant, the lower in advance of the upper; a vertical line from anterior margin of upper orbit passing through middle of lower; distance of upper eye from dorsal outline equaling  $\frac{2}{3}$  its vertical diameter; interorbital space concave, very wide, its width  $1\frac{1}{2}$  times diameter of orbit in a specimen 8 inches long, much narrower in the young; a ridge from upper angle in lower eye runs upward and backward to join a ridge from upper orbit. Nostrils on a level with upper margin of lower eye, the anterior with a flap, distant from the posterior, which is circular; length of snout to front of lower eye  $4\frac{1}{2}$  to 5 in head; mouth very oblique, the gape convex upward and backward; maxillary  $\frac{2}{3}$  length of head, reaching to middle of lower pupil, very narrow and covered with small scales; teeth small, the upper jaw with 2 series, the front teeth of the outer series somewhat enlarged; lower jaw with a single series; vomer and palatines toothless; gill rakers short and broad, the longest about  $\frac{1}{2}$  vertical diameter of pupil; about 7 on anterior limb of arch; pseudobranchiae present; preopercle with posterior margin nearly vertical, only the lower third free, the upper  $\frac{2}{3}$  grown fast to opercle and scaled over; the lower margin running very obliquely downward and forward, the angle thus an obtuse one; dorsal fin commencing on the snout in front of upper eye, the first 4 or 5 rays exerted and turned over to the blind side; the highest rays are behind the middle of the fin and are about  $\frac{2}{3}$  length of head; anal fin similar to dorsal, its origin under base of pectorals; caudal short, about  $\frac{2}{3}$  length of head, the middle rays the longest, the outer rays slightly prolonged; ventrals un-

symmetrical, that of colored side on the ridge of the abdomen, the other inserted in front of it; pectoral of colored side long, the rays very slender, the two upper prolonged and filamentous, the upper (in adults) more than  $\frac{1}{2}$  total length; pectoral of blind side more than  $\frac{2}{3}$  of length of head; scales ciliated, somewhat irregular, of moderate size, with small scales intermixed; snout naked, head and body otherwise scaly; scales on interorbital region very small; a series of small scales on basal half of each dorsal and anal ray; base of caudal thickly scaled, a series of small scales running nearly to tip of each ray, lateral line slightly rising anteriorly, but without distinct curve. Color light brown, with grayish and light bluish dots, some darker areas and a few round brown spots ocellated with lighter; interorbital space with a vertical brown bar bordered by lighter; fins mottled and spotted. This species is known only from the original types, taken by Professor Gilbert at Panama. The several variations in this species have not been studied. The species differs from *Syacium ovale* chiefly in the much broader interorbital space. We should regard this as unquestionably the adult male of *S. ovale* were it not that in making large collections of the latter species at Mazatlan we found not one referable to *S. latifrons*. (*latus*, broad; *frons*, forehead.)

*Oitharichthys latifrons*, JORDAN & GILBERT, Bull. U. S. Fish Comm. 1881, 334, Panama. (Coll. C. H. Gilbert.)

*Syacium latifrons*, JORDAN & GOSS, Review Flounders and Soles, 271, 1889.

#### 3043. SYACIUM OVALE (Günther).

Head  $3\frac{3}{4}$  in length; depth  $2\frac{1}{10}$ . D. 86; A. 69; scales 58; eye  $4\frac{1}{2}$  in head; maxillary  $2\frac{3}{5}$ ; pectoral  $1\frac{1}{2}$ ; caudal  $1\frac{1}{2}$ . Body elliptic-ovate, body outline from snout to caudal peduncle uniform, the snout not produced; mouth moderate; maxillary concave behind, reaching to middle of pupil of lower eye; lower jaw slightly included; teeth biserial in upper jaw, the inner series small and sharp, the outer much larger, irregular, uniserial in lower jaw; the lower eye slightly in advance of the upper; interorbital space narrow, as broad as pupil, concave; gill rakers as long as pupil, 2 + 8 in number. Scales strongly ctenoid; scales on mandible, maxillary, and a few in front of interorbital, the middle of which is naked; lateral line not curved. Dorsal beginning slightly in front of upper eye on blind side, the anterior rays produced a little beyond membrane; base of ventral of blind side wider than that of eyed side; caudal double lunate. Color light olive brown, nearly uniform, the vertical fins with elongate dark spots; caudal with large, irregular black spots. Pacific coast of tropical America; common at Mazatlan and Panama. Here described from specimens 6 or 7 inches in length, collected at Mazatlan, Mexico, by the Hopkins expedition to Sinaloa. None of these shows the broad interorbital area of *Syacium latifrons*. (*ovalis*, oval.)

*Hemirhombus ovalis*, GÜNTHER, Proc. Zool. Soc. Lond. 1864, 154, Panama; GÜNTHER, Fish. Centr. Amer., 472, pl. 80, fig. 1, 1869; JORDAN & GILBERT, Bull. U. S. Fish Comm. 1882, 108-111.

*Oitharichthys ovalis*, JORDAN, Proc. U. S. Nat. Mus. 1885, 301.

*Syacium ovale*, JORDAN & GOSS, Review Flounders and Soles, 271, 1889.



1049. CYCLOPSETTA, Gill.

*Cyclopsetta*, GILL, Proc. U. S. Nat. Mus., XI, 1888, 601 (*fimbriata*).

Mouth very large; jaws squarely truncated behind; teeth uniserial, those of the upper jaw moderate, of lower jaw enlarged and largest at sides; dorsal and anal almost symmetrical, dorsal commencing in front of eye on snout, scarcely deflected on blind side; caudal slightly pedunculate and convex; pectorals subequal and with a subtruncate free margin; ventrals nearly equal, the left on the preanal ridge, the right lateral, each with the inner ray connected by membrane to the body; interbranchial membrane imperforate; gill rakers tubercular and surmounted by blunt denticles. This genus differs from *Azevia* only in the smooth scales. (*κύκλος*, circle; *ψῆττα*, flounder, from the cycloid scales.)

a. Dorsal rays 91 to 95; anal 73 to 75; scales 90 to 95. Color nearly plain, the fins blotched. QUERNA, 3044.

aa. Dorsal rays 80 to 82; anal 62; dorsal and anal with dark ocelli.

b. Scales small, about 90; pectoral fin uncolored; anterior dorsal rays scarcely produced. CHITTENDENI, 3045.

bb. Scales larger, about 70; pectoral fin with black ocellus; anterior rays of dorsal somewhat produced. FIMBRIATA, 3046.

3044. CYCLOPSETTA QUERNA (Jordan & Bollman).

Head  $3\frac{1}{2}$  to  $3\frac{3}{4}$ ; depth  $2\frac{1}{2}$ . D. 91 to 95; A. 73 to 75; scales along lateral line 90 to 95. Body shaped as in *Azevia panamensis*. Mouth large, maxillary  $1\frac{1}{2}$  in head. Teeth as in *A. panamensis*, in single series, rather long and slender, the anterior somewhat more enlarged. Snout 5 in head, its tip hooked over the lower jaw so that the outer canines project. Interorbital space rather narrow, slightly concave, with a few small scales, its width a little less than pupil,  $\frac{1}{2}$  diameter of eye. Eyes moderate,  $5\frac{1}{2}$  in head, the upper somewhat in advance. Gill rakers short and broad, as in *A. panamensis*, each with 3 or 4 strong teeth. Scales small, cycloid on both sides, those below pectorals more reduced than in *A. panamensis*, about 65 in a cross series; anterior part of lateral line bent slightly upward, this portion about  $3\frac{1}{2}$  in straight part. Dorsal beginning above and between the nostrils, the anterior rays short, but with free tips; longest ray  $2\frac{1}{2}$  in head; pectoral of eye side  $1\frac{1}{2}$  to 2 in head, of blind side  $2\frac{1}{2}$  to  $2\frac{3}{4}$ ; ventrals subequal, each 6-rayed,  $2\frac{1}{2}$  in head, extending  $\frac{1}{2}$  their length beyond vent. Color plain brown, unspotted; fins dusky, thickly punctulate; young with 2 large oval indistinct dark spots on dorsal and anal; 3 on caudal, of which the middle is much larger. Distinguished from *A. panamensis* (Steindachner) by having much smaller cycloid scales on eyed side and by its plain coloration. Coast of Colombia. Numerous specimens, the largest about 8 inches in length, were dredged in 7 fathoms at Albatross Station 2800 and in 16 fathoms at Station 2802. (*quernus*, oaken, i. e., tanned.)

*Azevia querna*, JORDAN & BOLLMAN, Proc. U. S. Nat. Mus. 1889, 174, Pacific Ocean off coast of Colombia, at Albatross Station 2802, Lat.  $8^{\circ} 38' N.$ , Long.  $79^{\circ} 31' 30'' W.$  (Type, No. 41159.)

## 3045. CYCLOPSETTA CHITTENDENI, B. A. Bean.

Head  $3\frac{1}{2}$  in body; depth  $2\frac{1}{3}$ . D. 82; A. 62; scales 90; eye 5 in head. Mouth widely cleft, oblique, the jaws curved; cleft of mouth less than 2 in head. Teeth of each jaw in a single series, those of lower jaw strong and sharp, curved inward and backward, those of upper jaw not so large, and very irregular in size. Ventral fins well developed, that of eyed side being on abdominal ridge, and about  $\frac{1}{2}$  as long as pectoral; pectorals  $\frac{1}{2}$  as long as head, their length equaling a little more than  $\frac{1}{3}$  of body depth, posterior margin oblique; gill rakers very short, tubercular, almost as broad as long, 3 or 4 + 8 in number. Color brown; fins lighter, marked with blackish; 3 small faint blotches of black on first half of dorsal fin, and 3 rather distinct blotches on second half, last blotch extending to caudal peduncle; anal fin with 3 black blotches situated as and similar to those of dorsal fin; ventral of eyed side blackish, that of blind side pale; caudal fin with 3 black spots at its extremity; pectoral fin of colored side blackish; quite a large blotch of black on body under this fin. This species is distinguished from *Cyclopssetta fimbriata* by its shorter head, smaller and closely adhering scales, larger teeth, the little-produced anterior dorsal rays and by the oblique posterior margin of the pectorals. In *C. fimbriata* the scales are rather large and deciduous, the teeth small, the anterior rays of the dorsal considerably produced, and the posterior margin of the pectoral is subtruncate. A single specimen collected by Dr. John F. Chittenden, of the Victoria Institute, Port of Spain, Trinidad Island, and named in his honor. It is  $7\frac{1}{4}$  inches in length. (B. A. Bean.)

*Cyclopssetta chittendeni*, B. A. BEAN, Proc. U. S. Nat. Mus. 1894, 635, Trinidad. (Type, No. 44100. Coll. Dr. Chittenden.)

## 3046. CYCLOPSETTA FIMBRIATA (Goode &amp; Bean).

Head  $3\frac{1}{2}$  in length; depth nearly 2. D. 80; A. 60 or 61; pectoral 10; ventral 6; scales 25-70-31; maxillary 2 in head; caudal  $4\frac{1}{2}$  in total length; pectoral  $5\frac{1}{2}$ . Mouth very large, the upper jaw strongly curved, lower jaw included; teeth uniserial in each jaw, some of the anterior ones in the upper jaw being much larger than those following, while those in the lower jaw are still larger than these, some of the teeth in each jaw depressed; upper eye placed at a distance from profile equal to  $\frac{1}{2}$  its own diameter, which is a little less than 5 in head; eyes in the same vertical; interorbital ridge low, 4 in eye; gill rakers very short, tubercular, about 9 on lower part of angle. Scales cycloid; curve of lateral line slight, curve  $3\frac{1}{2}$  in straight part. Dorsal beginning on snout in advance of nostrils, first ray higher than second, highest rays behind middle of fin; origin of anal under base of pectoral, its highest rays behind middle of fin, higher than highest dorsal rays; ventral of eyed side on ridge of abdomen; middle caudal rays produced. Color grayish brown; dorsal and anal fins each with 2 round dark blotches upon their posterior halves, which are slightly larger than eye; a similar dark blotch upon middle of caudal, sometimes with smaller blotches, irregularly placed near its outer margin; pectoral with a very narrow dark band near its base, whole of outer half marked with a dark blotch, reticulated and mottled with

lighter; intervening portion pearly white with dark specks upon the rays; blind side cream colored. Deep waters of the Gulf of Mexico. (Goode & Bean.) (*fimbriatus*, fringed; from the produced dorsal rays.)

*Hemirhombus fimbriatus*, GOODE & BEAN, Proc. U. S. Nat. Mus. 1885, 591, deep waters of the Gulf of Mexico, between Mississippi Delta and Cedar Keys. (Type, No. 37330. Coll. Albatross.)

*Arnoglossus? fimbriatus*, JORDAN & GOSS, Review Flounders and Soles, 262, 1889.

*Cyclosetta fimbriata*, GOODE & BEAN, Oceanic Ichthyology, 451, fig. 368, 1896.

#### 1050. AZEVIA, Jordan.

*Azevia*, JORDAN, in JORDAN & GOSS, Review Flounders and Soles, 271, 1889 (*panamensis*).

Body elliptical, compressed, covered with small, firm, ctenoid scales; mouth large; teeth in both jaws uniserial; vomer without teeth; gill rakers very short and thick, tubercle-like; interorbital space very narrow in both sexes, the ridges coalescing between the eyes; lateral line without arch in front; ventrals free from the anal; septum of gill cavity below gill arches, without foramen; a deep emargination near isthmus. None of the fins especially modified or with elongate rays. This genus differs from *Citharichthys* in its tubercular gill rakers, as also in its small, firm scales, and other characters of minor importance. (*Azevia*, a Portuguese name for the sole, used at Lisbon, according to Brito-Capello. It probably corresponds to the Cuban name *Accidia*.)

#### 3047. AZEVIA PANAMENSIS (Steindachner).

Head  $3\frac{3}{8}$  in length; depth  $2\frac{1}{4}$ . D. 95; A. 73 to 78; scales 73 to 78; eye 5 in head, maxillary 2; pectoral  $1\frac{3}{8}$ ; caudal  $\frac{1}{4}$ . Body rather elongate; anterior profile evenly convex; mouth large, the maxillary reaching to posterior border of eye, the upper jaw somewhat hooked over the lower; about 3 teeth in upper jaw enlarged and hooked, canines in lower jaw long and sharp; eyes about even in head; interorbital space very narrow, less than diameter of pupil, a ridge along its middle; gill rakers divided into many sharp points around its edge, very short, as wide as long, about  $4 + 9$  in number. Scales on posterior part of interorbital, maxillary, and mandible; tip of snout, the greater part of interorbital, and tip of lower jaw naked; scales all strongly ctenoid; lateral line not curved anteriorly. Origin of dorsal at the vertical between tip of snout and front of eyes, scarcely on blind side, the anterior rays somewhat produced beyond membrane, the fin rather low; origin of anal below angle of opercle; pectorals short, that of eyed side pointed, its mate of the opposite side broadly rounded behind; caudal double lunate. Here described from a specimen collected by the Hopkins Expedition to Sinaloa, at Mazatlan, Mexico, about 11 inches in length. We have also examined specimens from Panama, in the museum at Cambridge, a part of the series of Dr. Steindachner's original types. Pacific coast of Central America; common at Mazatlan and Panama. (*panamensis*, from Panama.)

\**Citharichthys panamensis*, STEINDACHNER, Ichth. Beitr., III, 62, 1875, Panama; JORDAN & GILBERT, Bull. U. S. Fish Comm. 1882, 108 and 111; GILBERT, Bull. U. S. Fish Comm. 1882, 112.

*Azevia panamensis*, JORDAN & GOSS, Review Flounders and Soles, 272, 1889; JORDAN, Proc. Cal. Ac. Sci. 1895, 503.

## 1051. CITHARICHTHYS,\* Bleeker.

(WHIFFS.)

*Citharichthys*, BLEEKER, Comptes Rendus Acad. Sci. Amsterd., XIII, Pleuronectoidei, 6 1862 (*cayennensis* = *spilopterus*).

*Orthopsetta*, GILL, Proc. Ac. Nat. Sci. Phila. 1862, 330 (*sordidus*).

*Metoponops*, GILL, Proc. Ac. Nat. Sci. Phila. 1864, 198 (*cooperti* = *sordidus*).

Eyes and color on the left side. Body oblong; mouth of the large type, but comparatively small, with 1 series of small, sharp teeth in each jaw; no teeth on vomer or palatines. Gill rakers moderate, slender. Dorsal fin beginning just in front of eye; all the fin rays simple; ventrals of colored side on the ridge of the abdomen; no anal spine; caudal fin convex or double truncate behind; none of the fins produced. Scales thin, deciduous, slightly ctenoid. Lateral line nearly straight, simple. Lower pharyngeals separate, each with a single row of teeth. Vertebrae 30 to 40. This genus includes small flounders of weak organization, especially characteristic of the sandy shores of tropical America. The subgenus *Orthopsetta* includes species of more northern range and somewhat different form, and especially noteworthy as having an increased number of vertebrae. The two groups intergrade so perfectly that no sharp line of division can be drawn between them. (*Citharus*, an allied genus; *ιχθύς*, fish—a fish which lies on its *κίθαρος*, or ribs; that is, on its side.)

ORTHOSETTA (*ὀρθός*, straight; *ψῆτρα*, flounder):

a. Vertebrae 37 to 40; interorbital ridge sharply elevated; the head not closely compressed; eyes large; species of the North Pacific.

b. Interocular space concave, scaly, at least behind.

c. Gill rakers  $x + 16$  to 18.

d. Scales 65 to 70; dorsal rays 95; anal 77; depth  $2\frac{1}{2}$ . DUS, 3048.

dd. Scales 46 to 50; dorsal rays 83 to 87; anal 67 to 70; depth  $2\frac{3}{4}$  in length. FRAGILIS, 3049.

cc. Gill rakers  $x + 10$  or 11; dorsal rays about 84; anal 65; scales 50; depth  $2\frac{1}{4}$  in length. XANTHOSTIGMUS, 3050.

bb. Interocular space a sharp, naked ridge; dorsal rays 85 to 90; anal 68 to 72; scales 55 to 60; head  $3\frac{1}{4}$  in length; depth  $2\frac{1}{4}$ . STIGMÆUS, 3051.

CITHARICHTHYS:

aa. Vertebrae 33 to 36; interorbital ridge low and narrow, the head closely compressed. Species of the Atlantic or the Tropics.

c. Eyes large, 3 to  $4\frac{1}{4}$  in head.

f. Head large, 3 to  $3\frac{1}{4}$  in length.

g. Interorbital space very narrow, 5 in eye; snout with a spine; pectoral of eyed side elongate,  $\frac{1}{2}$  longer than head; maxillary  $2\frac{1}{4}$  in head. D. 91; A. 73; scales 48. DINOCEROS, 3052.

gg. Interorbital space very broad, 2 in eye; snout without spine; pectoral of eyed side shorter than head; maxillary  $2\frac{1}{4}$  in head. D. 78; A. 62; scales 43. PLATOPHRYS, 3053.

ff. Head smaller, about 4 in length.

h. Body comparatively elongate, the depth about  $2\frac{1}{2}$  in length; mouth very small; the maxillary  $3\frac{1}{4}$  in head; dorsal rays 83; anal 67; scales 40; eye 4 in head. ARCTIFRONS, 3054.

\* "As the name *Citharichthys* was introduced a short time before that of *Orthopsetta*, proposed for the *Psettichthys sordidus*, and was framed for a species related to that type, that name must be adopted if the *O. sordida* is not regarded as generically distinct." (Gill.)

hh. Body comparatively broad, the depth about  $\frac{1}{4}$  the length; mouth larger.

i. Snout with a strong, sharp spine on eyed side, above upper lip; eyes large, 3 in head; greatest depth of body over the pectorals; interorbital space with a wide ridge, about  $\frac{1}{4}$  diameter of eye. D. 74; A. 60; scales 40. UNICORNIS, 3055.

ii. Snout without distinct spine; eyes moderate,  $3\frac{1}{2}$  to  $4\frac{1}{4}$  in head; greatest depth of body under middle of dorsal; interorbital space a narrow, scaly ridge with a slight median groove; maxillary  $2\frac{1}{2}$  in head; teeth small, those in front slightly enlarged; body not very thin; gill rakers moderate, 6 + 13.

j. Dorsal rays 68; anal 52; scales smaller, the lateral line with about 53 pores; sides with whitish blotches.

UTLERI, 3056.

jj. Dorsal rays 80; anal 50; scales large, 41 in lateral line; sides and fins with dark blotches.

MACROPS, 3057.

ee. Eyes quite small, 5 to 6 in head; snout short, forming an angle with the profile; mouth moderate, oblique, the maxillary  $2\frac{1}{2}$  to  $2\frac{3}{4}$  in head; teeth small, the anterior somewhat enlarged; dorsal rays about 80; anal rays 60; body and fins speckled.

k. Scales not very large, 45 to 48 in lateral line; gill rakers long and slender, longer than pupil.

SPILOPTERUS, 3058.

kk. Scales large, 46 to 48 in lateral line; gill rakers short, not longer than pupil.

GILBERTI, 3059.

Subgenus ORTHOPSETTA, Gill.

3048. CITHARICHTHYS SORDIDUS (Girard).

(SOFT FLOUNDER.)

Head  $3\frac{3}{4}$ ; depth  $2\frac{1}{4}$ . D. 95; A. 77; scales 65 to 70. Form elliptical; interocular space concave, scaly, a conspicuous sharp ridge above the lower eye; mouth not large, the maxillary about 3 in length of head; teeth anteriorly subequal, growing much smaller behind. Gill rakers about 7 + 16. Lower pharyngeals narrow, each with 1 row of slender teeth. Scales rather large, thin, and membranaceous, readily deciduous, their edges slightly ciliate; accessory scales numerous. Eye large, much longer than snout,  $3\frac{1}{2}$  in head; depth of caudal peduncle less than  $\frac{1}{4}$  head; pectorals long, nearly  $\frac{2}{3}$  length of head. Vertebrae 11 + 29 = 40. Dull olive brownish of varying shade, the males with dull orange spots and blotches; each scale with a darker edge; dorsal and anal fins in the male blackish, with dull orange blotches, and edged anteriorly with yellowish; female paler, the fins nearly plain. Pacific coast of North America, in water of moderate depth; British Columbia to Lower California. This small flounder is one of the commonest species on the Pacific coast, being found in water of 10 fathoms or more depth in all localities from the Mexican boundary to British Columbia. Although much larger in size than any other species of the genus, it rarely exceeds 2 pounds in weight. In its deciduous scales and soft flesh it much resembles *Lyopsetta exilis* and *Atheresthes stomias*, 2 species which are often taken in company with it. Of all the species of *Citharichthys*, this one has the most extended range to the northward. (*sordidus*, sordid, from its dull coloration.)

- Psettichthys sordidus*, GIRARD, Proc. Ac. Nat. Sci. Phila., VII, 1854, 142, San Francisco; Tomales Bay; GIRARD, U. S. Pac. R. R. Surv., x, Fishes, 155, 1858.
- Metoponops cooperi*, GILL, Proc. Ac. Nat. Sci. Phila., 1864, 198, Santa Barbara; shrivelled specimen. (Type, No. 9407.)
- Orthopsetta sordida*, GILL, Proc. Ac. Nat. Sci. Phila., 1862, 330.
- Citharichthys sordidus*, LOCKINGTON, Rep. Com. Fisheries of California, 1878-79, 42; LOCKINGTON, Proc. U. S. Nat. Mus., 1879, 83; JORDAN & GILBERT, Proc. U. S. Nat. Mus., 1880, 453; JORDAN & GILBERT, Proc. U. S. Nat. Mus., 1881, 67; JORDAN & GILBERT, Synopsis, 817, 1883; BEAN, Proc. U. S. Nat. Mus., 1883, 353; JORDAN & GOSS, Review Flounders and Soles, 274, 1889.

## 3040. CITHARICHTHYS FRAGILIS, Gilbert.

Head  $3\frac{2}{3}$  to  $3\frac{3}{4}$  in length; depth  $2\frac{1}{4}$  to  $2\frac{3}{4}$  (in specimens 5 inches long.) D. 83 to 87; A. 67 to 70; scales 46 to 50. Vertebrae 10 + 27. Body elongate, posteriorly sharply wedge-shaped, tapering to base of caudal; anterior profile very conspicuously angulated above front of upper eye, the snout strongly projecting, its anterior profile nearly vertical; depth of caudal peduncle  $2\frac{1}{4}$  in head. Anterior nostril with a short tube and flap, the latter nearly obsolete on blind side. Mouth more oblique than in *C. sordidus*; maxillary reaching vertical from front of pupil,  $2\frac{2}{3}$  to  $2\frac{3}{4}$  in head. Teeth in a single series, close set, those anteriorly somewhat enlarged, but none of them canine-like. Eyes large, the vertical from front margin of upper eye falling through front of lower pupil; longest diameter of upper orbit  $2\frac{3}{4}$  in head; interorbital space narrow, concave, scaled, the lower ridge strongest, its width about  $\frac{1}{2}$  diameter of orbit. Symphyseal knob sharp. Gill rakers long, slender, close set (as in *C. sordidus*), 18 on anterior limb of arch, the longest  $\frac{1}{2}$  orbit. Scales large, deciduous, somewhat irregularly arranged, 12 or 13 series above lateral line; scales smooth on blind side, minutely spinous on eyed side; lateral line without anterior arch. Dorsal beginning slightly in advance of eye, the longest ray  $\frac{1}{2}$  head; pectorals long and narrow, with 11 rays on colored side. Color dusky olivaceous, with occasional slaty-blue spots. This species is closely related to *C. sordidus*, from which it differs in the fewer vertebrae and fin rays and in the larger scales. Many specimens from the Gulf of California in from 18 to 76 fathoms, at Albatross Stations 3011, 3016 to 3018, and 3033. (Gilbert.) (*fragilis*, fragile.)

*Citharichthys fragilis*, GILBERT, Proc. U. S. Nat. Mus., 1890, 120, Gulf of California, east coast of Lower California. (Type, No. 44409. Coll. Dr. Gilbert.)

## 3050. CITHARICHTHYS XANTHOSTIGMUS, Gilbert.

D. 81 to 86; A. 63 to 67; scales 50. Vertebrae 11 + 26. Body deep, varying from  $2\frac{1}{4}$  (in young, 3 inches long) to  $2\frac{1}{16}$  (7 inches long) in length of body. Profile angulated above front of upper eye, the snout convexly projecting. Depth of caudal peduncle  $\frac{1}{2}$  head. Lower eye in advance, the vertical from front of the upper passing through front of lower pupil. Mouth rather small, the outline somewhat curved, the maxillary reaching the vertical from front of lower pupil, 3 to  $3\frac{1}{4}$  in head; mandible with a sharp downward-directed point at symphysis. Teeth in a single close-set series in each jaw, growing slightly larger anteriorly, but without canines.

Anterior nostril with a short tube, and a narrow flap arising from its inner edge. Interorbital width  $3\frac{1}{2}$  to 5 in orbit, slightly concave, the lower ridge much stronger and higher than the upper, scaled posteriorly. Eye large, the orbit  $3\frac{1}{4}$  to  $3\frac{3}{8}$  in head. Gill rakers rather long and slender, coarsely dentate on inner margin, distant, 10 or 11 on anterior limb of arch. Scales large, in regular series, appearing cycloid, but the edges very minutely spinous; lateral line gently rising on anterior  $\frac{1}{3}$ , but without curve; fifty vertical series of scales, with as many pores in lateral line; 13 to 15 horizontal series above lateral line. Dorsal beginning immediately behind posterior nostril of blind side, ending so as to leave caudal peduncle free for a distance equaling  $\frac{1}{2}$  diameter of eye; ventrals long, reaching beyond origin of anal; pectoral very long and slender, normally with 9 rays, the longest ray on colored side longer than head, about  $\frac{1}{2}$  length of body. Color light olive brown, irregularly flecked with slaty, and with numerous bright yellow spots broadly ocellated with brownish black, a series of these usually on lateral line, and 2 others halfway between it and the dorsal and ventral outlines, respectively, those of the latter series forming pairs; fins not conspicuously marked, the pectorals sometimes with faint broad dusky cross bars. Both coasts of Lower California. In external appearance the species closely resembles *C. sordidus*, to which, however, it is not closely related, differing in number of scales, fin rays, and vertebrae, and in the size and number of gill rakers. (Gilbert.) Many specimens, from Albatross Stations 3039, 3043, and 3044, in 47 to 74 fathoms. (*ξανθός*, yellow; *στίγμα*, spot.)

*Citharichthys xanthostigma*, GILBERT, Proc. U. S. Nat. Mus. 1890, 120, Gulf of California, west coast of Lower California, and Magdalena Bay. (Type, No. 4408. Coll. Dr. Gilbert.)

3051. CITHARICHTHYS STIGMEUS, Jordan & Gilbert.

Head  $3\frac{1}{2}$  in length without caudal; depth 2 $\frac{1}{2}$ ; dorsal 87; anal 68; scales 54 (pores). Body moderately deep, the 2 profiles regularly and equally arched; snout short, gibbous, projecting a little beyond the outline; caudal peduncle very short, not high, its length (from end of last vertebra to vertical from last anal ray) about  $\frac{2}{3}$  its height, which is  $\frac{2}{3}$  length of head; caudal fin appearing sessile. Mouth moderate, very oblique, the maxillary reaching beyond front of pupil,  $2\frac{1}{2}$  in head; teeth in a single series, subequal in the two jaws, rather long, very slender and numerous, decreasing toward angle of mouth; about 40 teeth in the upper jaw, and 30 in the lower on blind side. Eyes large, close together, separated by a narrow, sharp, scaleless ridge; the upper eye largest, slightly behind the lower, with considerable vertical range; diameter of upper eye  $3\frac{1}{4}$  in head. Snout and lower jaw scaleless; end of maxillary and rest of head scaled. Gill rakers moderate, not strong, about 9 on anterior limb. Dorsal fin beginning on the vertical from front of upper eye, the first 3 rays being somewhat turned to blind side; the fin low, the highest at beginning of its posterior third, the longest ray nearly  $\frac{1}{2}$  length of head; anal spine present, very small; caudal rounded, about equaling length of head; pectoral of colored side  $1\frac{1}{2}$  in head, of blind side,  $2\frac{1}{2}$ . Scales

moderate, those forming the lateral line persistent, the others deciduous, those on colored side with ciliated margins, on blind side smooth; lateral line without anterior curve, the scales crowded and smaller anteriorly. Color in spirits uniform olivaceous, the scales dark edged; lips and some of the membrane bones of head margined with blackish; fins dusky, each seventh (to tenth) ray of vertical fins with a very small but conspicuous black spot on its middle. The above description is from the original type from Santa Barbara. Numerous specimens dredged by the *Albatross* in 9 to 41 fathoms off the coast of California show the following characters: Gill rakers  $x+9$ . Specimens 5 mm. long show white spots each with a black half ring on the outer side symmetrically arranged along bases of dorsal and anal; 4 distinct pairs of these, 2 unpaired ones more anteriorly along dorsal base, and a few fainter ones midway between these rows and the lateral line and alternating with them; there are some other scattered light spots. The abdomen is covered by a broad black streak; this, however, is wanting in specimens larger and smaller. Coast of California; rare; in rather deep water. The original type of this species is a young example, taken near Santa Barbara by Capt. Andrea Larco. In the Museum of Comparative Zoology are other specimens collected by Mr. Cary at San Francisco. These have 72 anal rays, while the original type had but 68. A few other specimens have been since obtained. Some of these are full of spawn at a length of 5 inches. (*στρυμαλιος*, speckled.)

*Citharichthys stigmæus*, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1882, 410, 411, Santa Barbara (Coll. A. Larco. Type, 31099 U. S. Nat. Mus.); JORDAN & GILBERT, Synopsis, 965, 1883; JORDAN & GOSS, Review Flounders and Soles, 274, 1889; GILBERT, Rept. U. S. Fish. Comm. 1893 (1896) 473.

#### Subgenus CITHARICHTHYS.

#### 3052. CITHARICHTHYS DINOCEROS, Goode & Bean.

Head  $3\frac{1}{2}$  in length; depth  $2\frac{1}{2}$ . D. 91; A. 73; scales 14-48-16; eye  $3\frac{1}{2}$  in head; maxillary a little less than 2; greatest height of dorsal 2; pectoral  $2\frac{1}{2}$  in body; caudal  $5\frac{1}{2}$ . Teeth uniserial in both jaws, those in the front much the largest; a strong spine upon the snout overhanging the upper lip, above this a second shorter spine; interorbital very narrow, its width less than 5 in eye, ridge rather prominent, narrow, sharp. Scales thin, deciduous, cycloid, large; lateral line slightly curved over the pectoral. Dorsal beginning on snout, in advance of eye, upon the blind side, its highest rays behind the middle; origin of anal under base of pectoral; third and fourth pectoral rays upon the eyed side elongate, the fin  $\frac{3}{8}$  longer than its mate of the opposite side; caudal subseesile, pointed. Color grayish brown above, white below. Vertebrae 33 to 36. West Indies, in deep water. The type specimen, 92 mm. long to base of caudal, was taken by the *Blake*, off Guadeloupe; others were taken off St. Lucie and Barbados, from 310 to 955 fathoms. (Goode & Bean.) (*δεινός*, terrible; *κέρας*, horn)

*Citharichthys dinoceros*, GOODE & BEAN, Bull. Mus. Comp. Zool., XII, No. 5, 157, 1886, off Martinique, St. Lucie, and Barbados; JORDAN & GOSS, Review Flounders and Soles, 275, 1889; GOODE & BEAN, Oceanic Ichthyology, 447, 1896.



3053. CITHARICHTHYS PLATOPHRYS, Gilbert.

Head 3; depth 2. D. 78; A. 62; scales 43. Body ovate; caudal fin subsessile, the free portion of caudal peduncle about  $\frac{1}{2}$  as long as diameter of pupil, its depth  $\frac{1}{2}$  length of head. Mouth very oblique; maxillary  $2\frac{1}{2}$  in head, reaching vertical from middle of lower eye. Teeth slender, close set, in a single series in each jaw, those in front of upper jaw largest, but not canine-like. Eyes large, the lower much in advance of the upper, their horizontal diameter  $3\frac{1}{2}$  in head. Interorbital space very wide for the genus, concave, divided by an oblique ridge running backward from middle of upper orbit; interorbital width  $8\frac{1}{2}$  in head, nearly  $\frac{1}{2}$  as wide as eye. Distance from tip of snout to front of lower eye  $\frac{3}{4}$  diameter of eye, from tip of snout to upper eye  $\frac{1}{2}$  head. Gill rakers short and very slender, less than diameter of pupil, 9 present on horizontal limb of outer arch. Scales large, those on blind side very weakly ctenoid. Dorsal beginning behind nostril on blind side of snout, its longest ray  $2\frac{1}{2}$  in head; pectoral of eyed side long and narrow, 4 in length, containing 11 rays, that of blind side but  $\frac{1}{2}$  its length; ventrals short; caudal rounded,  $1\frac{2}{3}$  in head. Color in spirits, uniform light brownish (olivaceous in life), without distinctive marks; fins somewhat dusky; ventral of eyed side jet-black, that of blind side blackish on distal portion of inner rays. (Gilbert.) One specimen known, from Albatross Station 2799, southwest of Panama. (*πλατύς*, broad; *ὄφρυς*, eyebrow.)

*Citharichthys platophris*, GILBERT, Proc. U. S. Nat. Mus. 1890, 454, Albatross Station 2799, southwest of Panama. (Coll. Albatross.)

3054. CITHARICHTHYS ARCTIFRONS, Goode.

Head 4 in body; depth  $2\frac{1}{2}$ . D. 82; A. 67; pectorals 9 or 10, 7; scales 8-40-8; eye 4 in head; maxillary  $3\frac{1}{2}$ ; caudal 1. Body comparatively elongate; mouth small; teeth small, the anterior scarcely enlarged; interorbital space narrow, sharp, scaleless; scales cycloid, deciduous; small scales on the rays of the ventral fins; lateral line sharply defined, straight. Dorsal beginning above front of upper eye, its highest ray about 3 times the distance from snout to first ray; origin of anal under base of pectoral; caudal subsessile, triangular; rays of vertical fins all exerted; pectoral inserted low, that of eyed side twice the length of the other. Color dirty light brown. Deep waters of the Gulf Stream. (*arctus*, contracted; *frons*, forehead.)

*Citharichthys arctifrons*, GOODE, Proc. U. S. Nat. Mus. 1890, 341, 472, Gulf Stream off southern coast of New England; GOODE & BEAN, Bull. Mus. Comp. Zool., Vol. x, No. 5, XIX, 194, 1883; JORDAN & GILBERT, Synopsis, 818, 1883; JORDAN & GOSS, Review Flounders and Soles, 275, 1889; GOODE & BEAN, Oceanic Ichthyology, 442, fig. 366, 1896.

3055. CITHARICHTHYS UNICORNIS, Goode.

Head 4 in length; depth a little less than length. D. 74; A. 60; P. 4 (right), 10 (left); scales 12-40-12; eye 3 in head; maxillary scarcely 2; highest dorsal ray 2. Body deep, its greatest height over the pectorals;

scales thin, deciduous; eye equal to snout or interorbital space; interorbital with a strong ridge; teeth minute, close set, in a single series, stronger on the blind side; a strong, sharp spine on the snout at the anterior termination of the ridge at lower margin of upper eye; caudal pointed, triangular, sessile; pectoral of left side twice as long as the eye, not  $\frac{1}{2}$  longer than right pectoral. Dorsal beginning at side of preorbital spine, its anterior rays being slightly upon the blind side; anal equal to dorsal in height. Ashy gray, with dark lateral line; eyes black. (Goode.) Deep waters of the Gulf Stream. (*unicornis*, having one horn.)

*Citharichthys unicornis*, GOODE, Proc. U. S. Nat. Mus. 1880, 342, Gulf Stream off southeast of New England; JORDAN & GILBERT, Synopsis, 818, 1883; JORDAN & GOSS, Review Flounders and Soles, 275, 1889; GOODE & BEAN, Oceanic Ichthyology, 444, fig. 369, A & B, 1896.

3056. CITHARICHTHYS UHLERI, Jordan.

D. 68; A. 52; scales 53 (pores). Body comparatively broad, regularly oval, without angle; greatest depth of body under middle of dorsal; eyes moderate,  $4\frac{1}{2}$  in head, close together, the orbital ridges coalescent, the lower larger. Teeth small, uniserial; maxillary  $2\frac{1}{2}$  in head; gill rakers short and very slender,  $x + 12$ . Color dark brown, with whitish blotches. The fins mottled. Hayti. A single specimen in the Museum of Comparative Zoology,  $4\frac{1}{2}$  inches in length. The species is close to *Citharichthys macrops*, but its fin rays and scales are considerably more numerous than in the latter. (Named for Mr. Philip Reese Uhler, the well-known entomologist, its discoverer.)

*Citharichthys uhleri*, JORDAN & GOSS, Review Flounders and Soles, 275, 1889, Hayti. (Coll. P. R. Uhler. Type in Mus. Comp. Zool.)

3057. CITHARICHTHYS MACROPS, Dresel.

Head 4 in body; depth scarcely 2. D. 80; A. 56; scales 14-41-16; lower eye 4 in head; maxillary  $2\frac{1}{2}$ ; highest dorsal rays a little over 2; pectoral of eyed side  $1\frac{3}{4}$ ; caudal 4 in body; vertebrae  $9 + 25 = 34$ . Body suboval; upper profile very convex, descending in a sharp curve from nape to front of upper eye, and forming an abrupt angle with the short, blunt snout; mouth moderate, very oblique and curved; maxillary reaching to below middle of eye; teeth minute, uniserial, slightly larger on blind side; interorbital narrow, with a scaleless ridge, which curves upward and backward to upper angle of gill opening; upper eye very close to profile, its anterior margin on the same vertical line with lower; snout shorter than eye; gill rakers about  $\frac{1}{2}$  the length of eye,  $6 + 13$  in number. Scales large, not ciliated, no accessory scales; origin of dorsal on blind side near tip of snout, anterior rays exerted, the first ray as long as eye, the fin highest at its middle portion; origin of anal under base of pectoral, its highest part a little higher than dorsal; caudal pointed; pectoral of blind side somewhat shorter than that of eyed side. Color in spirits, light-olive brown; body with some 20 dark-brown spots, the largest as large as eye, 4 of these arranged at equal intervals along the lateral line, the second near the middle the most prominent; dorsal and anal fins with a series of round, brown spots, 1 at the middle of every sixth or seventh ray, besides small irregular spots and mottlings;

caudal spotted and mottled with dark brown, and with 2 round, brown spots, 1 above the other at the base of the fin. (Dresel.) South Atlantic and Gulf coasts of the United States; rather common; a well-marked species. We have examined several specimens dredged in the harbor of Beaufort, N. C., by Prof. Oliver P. Jenkins. (*μακρός*, large; *ᾠψ*, eye.)

*Citharichthys macrops*, DRESEL, Proc. U. S. Nat. Mus. 1864, 539, Pensacola (Type No. 21500); JORDAN, Proc. U. S. Nat. Mus. 1884, 29; JORDAN & GOSS, Review Flounders and Soles, 275, 1889.

3058. *CITHARICHTHYS SPILOPTERUS*, Günther.

Head  $3\frac{1}{2}$  in body; depth  $2\frac{1}{6}$ . D. 75 to 80; A. 58 to 61; scales 45 to 48; eye 6 in head; maxillary  $2\frac{1}{2}$ ; pectoral  $2\frac{1}{2}$ ; highest dorsal and anal rays 2; caudal  $1\frac{1}{2}$ . Body moderately elongate, much compressed; snout short, forming an angle with the profile; jaws strongly curved, the upper somewhat hooked over the lower; lower jaw slightly included; maxillary reaching to posterior margin of lower orbit; teeth small, in a single row, the anterior a little enlarged; interorbital area a low, narrow ridge, which is divided only anteriorly; gill rakers short and rather slender, about 3 in eye, 4 + 12 in number; scales cycloid. Origin of dorsal above anterior edge of upper eye, very slightly on blind side, its highest rays in its posterior half; origin of anal slightly behind base of pectoral; pectoral of eyed side very slightly shorter than that of eyed side; vertebrae 34. Color olive brownish, somewhat translucent, with darker dots and blotches; a series of distant obscure blotches along bases of dorsal and anal. Atlantic coast of tropical America north to New Jersey; very common on sandy shores; not found in the Pacific, all west coast references belonging to *C. gilberti*. Here described from a specimen from Havana, 6 inches in length. This little flounder is almost everywhere abundant on the sandy shores of the warmer parts of the Western Atlantic, in shallow water. Careful comparison of specimens from South Carolina, Cuba, and Brazil shows no tangible difference, and we are compelled to regard all as forming a single species. It rarely exceeds 5 or 6 inches in length. It usually comes into the markets mixed with other shore fishes, and it nowhere receives any notice as a food-fish. This species is common in the markets of Havana, and it is evidently the original of Poey's *Hemirhombus fuscus*, although in Poey's description there seems to be some confusion, because the teeth are said to be biserial above, and 60 scales are counted in the lateral line. A specimen from Poey in the museum at Cambridge is labeled "*Hemirhombus fuscus*, type. Collector's number, 87." This belongs to *C. spilopterus*, and it has 48 scales in the lateral line. Bleeker's *C. guatemalensis* agrees in all respects with *C. spilopterus*. We are unable to find any description of *C. cayennensis*, if, indeed, the species has ever been described. Specimens of *C. spilopterus* are in the museum at Cambridge from Cuba, Pará, Sambaia, Pernambuco, Camaru, Rio das Velhas, Rio Janeiro, and San Matheo. (*σπίλος*, spot; *πτερόν*, fin.)

*Citharichthys spilopterus*, GÜNTHER, Cat., IV, 421, 1862, New Orleans, San Domingo, Jamaica; JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1882, 618; JORDAN & GILBERT, Synopsis, 817, 1883; JORDAN, Proc. U. S. Nat. Mus. 1886, 53; JORDAN & GOSS, Review Flounders and Soles, 276, 1889.

*Citharichthys cayennensis*, BLEEKER, Comptes Rendus Acad. Sci. Amsterd., XIII, 1862, 6, Cayenne; name only.

*Citharichthys guatemalensis*, BLEEKER, Neder. Tydschr. Dierk. 1864, 73, Guatemala; GÜNTHER, Fish. Centr. Amer., 472, 1869.

*Hemirhombus fuscus*, POEY, Synopsis, 406, 1868, Havana; POEY, Enumeratio, 138, 1875.

3059. CITHARICHTHYS GILBERTI, Jenkins & Evermann.

Head  $3\frac{1}{2}$  to  $3\frac{3}{4}$ ; depth of head 4; depth of body  $1\frac{5}{16}$  to  $2\frac{1}{4}$ . D. 77 to 82; A. 57 to 61; scales 18-40 to 46-19. Body comparatively broad, formed as in *C. spilopterus*, the two profiles about equally arched; snout slightly longer than longest diameter of eye, and without a distinct spine. Eyes on left side, equal in size, small, 5 to  $5\frac{1}{4}$  in head; interorbital space narrow,  $1\frac{1}{2}$  in eye, low, slightly grooved, and scaled on posterior portion only. Maxillary  $2\frac{3}{4}$  in head, reaching barely to posterior border of eye; upper jaw projecting. Teeth small, in a single series; gill rakers 4 + 13, short and slender, not longer than pupil, with a rather broad base, narrowing to a slender stalk. Dorsal fin beginning in front of upper eye, the first 3 rays growing from the blind side, the distance of origin from snout 7 in head; fin rays all simple,  $2\frac{3}{16}$  in head; pectorals nearly equal, the one on colored side being slightly longer,  $1\frac{5}{16}$  in head; rays on colored side 9; on blind side 8; ventrals  $2\frac{3}{16}$  in head; caudal rounded, caudal peduncle short, its depth 8 in the body, equaling height of anal; scales large, ciliated, pretty uniform, those toward head and margins of disk becoming smaller; lateral line gradually descending along the course of about 16 scales, from which point it is straight. Color light brown, with about 15 irregular dark blotches of various sizes, the largest being a pair on the latter third of the disk, 1 on each side of lateral line, as great in diameter as length of ventral fin. Specimens from fresh waters (*C. sumichrasti*) are much darker in color; gray, everywhere closely peppered with dark specks; pectoral and caudal mottled. Pacific coast of tropical America; very abundant in sandy bays from Guaymas to Panama, ascending all the streams. This species very closely resembles *C. spilopterus*, representing the latter on the Pacific coast, and it has been frequently recorded under the name *C. spilopterus*. *C. gilberti* differs mainly in the shorter gill rakers and in the slightly larger scales. Fresh-water specimens (as the type of *C. sumichrasti* from Rio Zanatenco, Chiapas, and numerous examples collected by us in Rio Presidio, near Mazatlan) differ considerably in color, being much darker, but there is no other difference. ("This species is dedicated to Prof. Charles H. Gilbert, whose collection and notes on fishes from Mazatlan, containing undescribed species, this among them, were destroyed by fire in 1883.")

*Citharichthys gilberti*, JENKINS & EVERMANN, Proc. U. S. Nat. Mus. 1888, 157, Guaymas, Mexico (Type, No. 39627. Coll. Jenkins & Evermann); JORDAN, Proc. Cal. Ac. Sci. 1895, 503.

*Citharichthys sumichrasti*, JORDAN & GOSS, Review Flounders and Soles, 276, 1889, Rio Zanatenco, Chiapas. (Coll. Prof. Francis E. Sumichrast. Type, 25299, M. C. Z.)

*Citharichthys spilopterus*, GÜNTHER, Fish. Centr. Amer., 471, pl. 80, fig. 2, 1869; JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1882, 382; JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1882, 630; JORDAN & GILBERT, Bull. U. S. Fish Comm. 1882, 108-111; not of GÜNTHER, 1862.

## 1052. ETROPUS, Jordan &amp; Gilbert.

*Etropus*, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1881, 364 (*crossotus*).

Eyes and color on left side. Body regularly oval, deep, and compressed. Head small; mouth very small, the teeth close set, slender, and pointed, somewhat incurved, mostly on the blind side; no teeth on vomer. Eyes small, separated by a narrow, scaleless ridge; margin of preopercle free. Ventrals free from anal, that of colored side inserted on ridge of abdomen, its base rather long. Dorsal fin beginning above eye; caudal double truncate; anal without spine. Scales thin, deciduous, ctenoid on left side, cycloid on blind side. Lateral line simple, nearly straight. Size small. This genus is very close to *Citharichthys*, from which it differs only in the very small size of the mouth, and in the correspondingly weak dentition. The 3 or 4 known species are similar in appearance to the species of *Citharichthys*, and they inhabit the same waters. Another genus extremely close to *Etropus* and *Citharichthys* is *Thysanopsetta*, a South American genus. The teeth in *Thysanopsetta* are, however, arranged in a band. The larval forms are translucent and symmetrical, as in *Platophrys*, *Monoleue*, *Arnoglossus*, etc. (*ἔτροπον*, abdomen; *πόυς*, foot; in allusion to the insertion of the ventrals, common to all the *Psettinae*, but not found in other small-mouthed species.)

a. Snout not acute; dorsal rays 75 to 85.

b. Body comparatively elongate, the depth rather less than  $\frac{1}{2}$  length.

c. Dorsal rays 81; anal 58; head  $4\frac{1}{2}$  in length; eye  $3\frac{3}{4}$  in head; maxillary 4.

MICROSTOMUS, 3060.

cc. Dorsal rays 77; anal 61; head 4 in length; eye  $3\frac{1}{2}$  in head; maxillary  $4\frac{1}{2}$ .

RIMOSUS, 3061.

bb. Body very deep, the depth more than  $\frac{1}{2}$  length; eye  $3\frac{3}{4}$  in head; maxillary 4;

head  $4\frac{1}{2}$ ; depth  $1\frac{1}{2}$  to 2; D. 76 to 85; A. 56 to 67; scales 42 to 48; cirri on subopercle of blind side very numerous, white; olive ground, with darker blotches; fins sanded.

CROSSOTUS, 3062.

## 3060. ETROPUS MICROSTOMUS (GILL).

"D. 81; A. 58; caudal 4, 6, 5, 3; pectoral 10; ventral 6. The height of the body enters about  $2\frac{3}{8}$  times (0.36-0.37) in the extreme length; that of the caudal peduncle about 11 times. The head forms a fifth of the length, is rather abbreviated, scarcely sinuous above the eyes, blunt at the snout, which scarcely exceeds  $\frac{1}{4}$  of the head's length and the rostral area is rhombic, and not higher than long. The eyes are even; the longitudinal diameter contained about  $3\frac{3}{8}$  times (0.05 $\frac{1}{2}$ ) in the head's length. The mouth is rather small, the length of the upper only equaling  $\frac{1}{2}$  of the length, and that of the lower  $\frac{2}{3}$  of the head's length. The teeth are very small, and close together; larger in front. The dorsal commences above the front of the orbit, and is highest and convergent near the fortieth ray, which equals about  $\frac{1}{10}$  of the total length; the anal is highest at about the twenty-fifth ray, and is as high or even higher than the dorsal. The caudal is rounded behind, and forms about  $\frac{1}{4}$  of the length. The pectoral fins are unequally developed, that of the dark side being prolonged, and contained only  $6\frac{3}{8}$  times in the total length, while that of the white side only equals  $\frac{1}{10}$  of the same; the rays are all simple.

The ventral fins are also unequally developed, the right being on the abdominal ridge at its origin, rather in advance of the opercular margin and with its longest rays contained about 14 times in the total length; stretched backward, it extends to the second anal ray; the fin on the white side is more advanced, wider, and its rays longer, contained less than 12 times in the length and extends backward to nearly the third anal ray. The scales are large, angular behind, covered with smaller ones, especially near the point of junction of contiguous ones, where alone they are developed on the blind side; the scales of the eyed side are mostly minutely ciliated behind, unarmed, however, near the lateral line, the scales of which last are quadrate and mostly covered; the scales of the blind side are less angular behind and unarmed. The lateral line runs through about 42 scales, while of longitudinal rows there are 10 above and 14 below the lateral line. The color is uniform reddish brown. A single specimen, little more than 3 inches long, was first obtained by Professor Baird, at Beesleys Point." (Gill.)

This species has not been certainly recognized by recent writers, unless, as supposed by Jordan & Goss, it is identical with *Etropus rimosus*. It is in any event certainly an *Etropus*. In the Museum of Comparative Zoology are numerous young specimens collected at Somers Point, New Jersey, by Dr. Stimpson. These seem to belong to the genus *Etropus*. The teeth are equal; the scales are 44, and the depth of the body is  $2\frac{1}{2}$  in its length. The eye is 4 in head, the dorsal rays 75 to 80, and the anal rays 56 or 57. The color is light brown, mottled and spotted with darker. These probably represent the *Citharichthys microstomus* of Gill, collected in the same neighborhood by the same naturalist. We are unable to distinguish them from *Etropus rimosus*. (*μικρός*, small; *στόμα*, mouth.)

*Citharichthys microstomus*, GILL, Proc. Ac. Nat. Sci. Phila. 1864, 223, Beesleys Point, New Jersey (Coll. Prof. S. F. Baird); GOODE & BEAN, Oceanic Ichthyology, 446, 1896; JORDAN, Proc. U. S. Nat. Mus. 1890, 332.

*Etropus microstomus*, JORDAN & GOSS, Review Flounders and Soles, 278, 1889.

#### 3061. ETROPUS RIMOSUS, Goode & Bean.

Head 4 in body; depth 2 to  $2\frac{1}{4}$ . D. 77; A. 61; scales 12-41-14; eye  $3\frac{1}{2}$  in head; maxillary  $4\frac{1}{2}$ ; snout 8; caudal 1. Body somewhat elongate, pear-shaped; mouth very small, its cleft less than  $\frac{1}{2}$  the orbit, its angle below anterior margin of lower eye; teeth well developed on blind side on each jaw, also on eyed side of lower jaw in front; eyes placed in the same vertical; upper eye close to dorsal profile, and separated from its mate by a space less than  $\frac{1}{2}$  its diameter; interorbital ridge low; nostrils in a line with the interorbital ridge, each in a short tube, the posterior the larger, the anterior midway between tip of snout and front of lower orbit; head entirely scaled; accessory scales numerous. Dorsal commencing on blind side at anterior margin of eye, the highest rays somewhat behind middle of fin, its length 7 times in total length; origin of anal under base of pectoral, its highest rays equal to those of dorsal; pectoral of eyed side longest, equal to head without snout; caudal fin rounded. Color gray, hoary above, with a few irregularly placed indistinct brownish blotches, none of them larger than eye; white below. West coast of Florida;

type, 100 mm. long, collected by the *Albatross* at Station 2408, depth 21 fathoms, between Pensacola and Cedar Keys, Florida. (Goode & Bean.) On reexamining our specimens of *Etropus*, we find that those obtained by Jordan & Evermann from Pensacola differ from the others in the greater elongation of the body and in the somewhat grayer coloration. These correspond fairly to the description and figure of *Etropus rimosus*. All other specimens from the United States coast collected by Dr. Jordan and his associates are, in our opinion, referable to *Etropus crossotus*. The original description of *Citharichthys microstomus*, Gill, fits this species better than any other known. The fish in question is much too elongate for *Etropus crossotus* (depth  $2\frac{3}{4}$  in total length), and the mouth is too small for any of the known species of *Citharichthys* (maxillary 4 in head; mandible  $2\frac{1}{2}$ ). We have little doubt of the identity of *Etropus rimosus* and *microstomus*, but leave the matter for further investigation. The separation of *E. rimosus* from *E. crossotus* is not beyond question. (*rimosus*, frosted.)

*Etropus rimosus*, GOODE & BEAN, Proc. U. S. Nat. Mus. 1885, 593, coast of Florida, between Pensacola and Cedar Keys, dredged at the depth of 21 fathoms; GOODE & BEAN, Oceanic Ichthyology, 455, pl. 104, figs. 360, 361, 1896. (The latter figure an excellent representation of the symmetrical, translucent larval form, before the right eye has crossed the forehead.)

*Etropus crossotus*, JORDAN & EVERMANN, Proc. U. S. Nat. Mus. 1886, 476; not of JORDAN & GILBERT.

3062. *ETROPUS CROSSOTUS*, Jordan & Gilbert.

Head  $4\frac{1}{2}$  in length; depth  $1\frac{1}{2}$  to 2. D. 76 to 85; A. 56 to 67; V. 6; scales 42 to 48; vertebrae  $9+25=34$ . Body oval, strongly compressed, with the dorsal and ventral curves nearly equal; both outlines strongly arched anteriorly, the body much deeper in adult specimens. Head very small; snout short; mouth very small, its cleft not so long as diameter of orbit. Teeth conical, pointed, close set, strongly incurved, in a single series, those in upper jaw on blind side only, those in lower jaw on both sides. Eyes large, the lower in advance of the upper, the two separated by a very narrow scaleless ridge, which extends backward above preopercle; edge of opercle on blind side, with a row of conspicuous white cilia. Upper nostril turned somewhat to blind side; anterior nostril on left side, with a very slender cirrus. Dorsal fin commencing over front of upper eye, its middle rays highest, the anterior not elevated; anal fin not preceded by a spine, its middle rays highest; caudal fin very sharply double-truncate, as long as head; pectorals short, that of left side the longer, about  $\frac{1}{2}$  length of head; ventral of colored side on ridge of abdomen, the membrane of its last rays nearly reaching base of first ray of anal; ventral of blind side longer than the other,  $\frac{1}{2}$  length of head, inserted farther forward than that of colored side. Vent lateral, with a well-developed anal papilla. Scales thin, large, ctenoid on colored side, smooth on blind side, those on the middle part of the body larger; head entirely scaly, except snout and interorbital ridge; rays of vertical fins with scales on the basal half, on colored side; lateral line developed equally on both sides, nearly straight. Color olive brown, with some darker blotches most distinct in the larger specimens; vertical fins finely mottled and streaked with black and gray; pectoral and ventral on left

side spotted. Tropical America on both coasts, north to Cerros Island and North Carolina, south to Panama and Rio Janeiro; the type a single specimen, about 5 inches long, taken with a seine in the Astillero at Mazatlan. This little fish seems to be abundant in all warm and sandy shores of tropical America. It is the smallest and feeblest of all our flounders, and has therefore been generally overlooked by collectors. In the Museum of Comparative Zoology are specimens of this species from Rio Janeiro, Santos, Victoria, Para, and Sambaia, in Brazil. The largest of these is 6 inches in length. Head 5 in length; depth  $1\frac{3}{5}$ ; scales 44; D. 85; A. 67. We have specimens from Charleston, Cedar Keys, New Orleans, Galveston, Beaufort, North Carolina, Mazatlan, Panama, and from several localities along both sides of the coast of Lower California. These vary in form, color, and squamation, but we are unable to point out specific distinctions among them. (*κροβδωρός*, fringed, from the cirri of the subopercle.)

*Etropus crossotus*, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1881, 364, Mazatlan; JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1882, 305; JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1882, 618; JORDAN & GILBERT, Bull. U. S. Fish Comm. 1882, 108-111; JORDAN & GILBERT, Synopsis, 839, 1883; BEAN, Cat. Fish. Intern. Exh. 1883, 44; JORDAN & SWAIN, Proc. U. S. Nat. Mus. 1884, 234; JORDAN & GOSS, Review Flounders and Soles, 278, 1889.

*Etropus microstomus*, JORDAN, Proc. U. S. Nat. Mus. 1886, 29; not *Citharichthys microstomus*, GILL.

#### 1053. MONOLENE, Goode.

*Monolene*, GOODE, Proc. U. S. Nat. Mus. 1880, 338 (*sessilicauda*).

*Thyris*, GOODE, Proc. U. S. Nat. Mus. 1880, 344 (*pellucidus*; larval form); name preoccupied.

*Delothyris*, GOODE, Proc. U. S. Nat. Mus. 1883, 110 (*pellucidus*); substitute for *Thyris*.

Body thin, elongate; eyes on the left side, very close together, near the profile; mouth moderate, the length of the maxillary less than  $\frac{1}{2}$  that of the head; teeth minute, in a single series, nearly equal on both sides; no teeth on vomer or palatines. Scales rather large, ctenoid on colored side, cycloid on blind side. Lateral line well marked, that of colored side strongly and angularly curved anteriorly, that of blind side nearly straight. Pectoral of blind side wholly absent; dorsal beginning on the snout, its rays all simple. Caudal fin sessile, almost confluent with dorsal and anal. Ventral fins normal, that of the left side on the ridge of the abdomen; gill rakers few, feeble. Vertebrae 43. Deep-sea fishes, closely allied to *Trichopsetta* and *Arnoglossus*, but with the right pectoral obsolete. The translucent larva of *Monolene* is similar to the larva of *Platophrys*. It was at first described as a distinct genus (*Thyris* = *Delothyris*) by Dr. Goode before its true character was recognized. The following are the characters ascribed to the larval genus *Delothyris*.\*

\* The following are the characters of the species, *Delothyris pellucidus*, Goode:

Colorless, translucent; 3 conspicuous, dusky, longitudinal lines on left side, the middle one faintest. Two streaks on right side; eyes black. Body thin, pellucid, divided into 3 longitudinal tracks by depressions at the bases of the rows of interspinous processes. Scales small, thin, caducous. Head very small; eyes small, protruding, their diameter equal to the interorbital space and  $\frac{1}{2}$  the length of the snout; mouth small, formed as in the soles, the upper jaw somewhat hook-shaped. Dorsal fin beginning in advance of the eye, of long, flexible, simple rays, the tips of which are much exerted. Pectorals inserted far below lateral line, that on blind side as long as orbit, the other as long as snout; ventrals reaching past front of anal. Head 5; depth 3. D. 100; A. 80; P. 12 (left), 4 (right). Length 3 inches. (Goode.) Gulf Stream, off the coast of Rhode Island.



Body elongate, soft, and translucent. Head very short; mouth small, toothless. Eyes sinistral, close together, the lower slightly advanced. Pectoral of blind side smallest; ventrals crowded together on median keel of body, their bases prolonged on this keel. Rays simple; dorsal beginning on the snout; caudal sessile, almost confluent with dorsal and anal. Scales very thin, easily detached, probably cycloid. Lateral line well marked, straight. (*μόνος*, one; *ώλένη*, arm.)

a. Dorsal rays 99 to 103; anal rays 79 to 81; scales 92.  
aa. Dorsal rays 124; anal rays 100; scales 105.

SESSILICAUDA, 3063.  
ATRIMANA, 3064.

3063. *MONOLENE SESSILICAUDA*, Goode.

Head 5 in body; depth 2½. D. 99 to 103; A. 79 to 84; scales 23-92-25; eye 4 in head; highest dorsal ray 2; highest anal rays slightly more than 3; pectoral 1½; caudal nearly 1. Body moderately elongate; maxillary extending slightly past front of lower eye, with uniserial, subequal teeth; lower eye in advance of upper; interorbital space very small, less than ½ the diameter of eye; head everywhere closely scaled; scales ctenoid; lateral line strongly curved over anterior ¾ of pectoral, the curve with 2 angles, 72 scales along straight portion; lateral line of blind side nearly straight. Origin of dorsal over anterior edge of lower eye, longest rays in the posterior fourth of the fin; origin of anal under base of pectoral; pectoral present only on eyed side. Color on left side ashy brown, with numerous more or less distinct darker brown spots; on blind side white, pectoral blackish with traces of lighter transverse bands. Specimens from shallow water near Key West (Coll. Prof. C. C. Nutting), according to Mr. Garman, are much more brightly colored. These are "grayish brown, with numerous spots of darker to blackish over head and body, the spots being ½ as large as the eye or smaller, arranged for the greater part in broad transverse bands as wide as the interspaces, of which bands the first and foremost passes from the nape to the opercle, the second lies immediately behind the pectoral, the third just in front, and the fourth just behind the middle of total length, and the fifth, more indistinct, crosses near the ends of the dorsal and anal; the caudal is crossed by 2 rather indefinite narrow streaks; the pectoral is white at its base and bears 3 or 4 narrow curved transverse bands of white, separating 3 or 4 similar bands of black, which with the white are more distinct in the lower half of the fin." Specimens from 150 fathoms or more have markings similar but less distinct. D. 104; A. 84; V. 6; P. 11; scales 22-93-24. Deep waters of the Gulf Stream, Cape Cod to Key West. (*sessilis*, sessile; *cauda*, tail.)

*Monolene sessilicauda*, GOODE, Proc. U. S. Nat. Mus. 1880, 337, 338, deep sea south of New England; GOODE, Proc. U. S. Nat. Mus. 1880, 472; JORDAN & GILBERT, Synopsis, 841, 1883; JORDAN & GOSS, Review Flounders and Soles, 280, 1889; GOODE & BEAN, Oceanic Ichthyology, 452, figs. 357 A & B, 1896; GARMAN, Bull. Iowa Lab. Nat. Hist. 1896, 91.

*Thyris pellucidus*, GOODE, Proc. U. S. Nat. Mus. 1880, 344, Gulf Stream off the coast of Rhode Island.

*Delothyris pellucidus*, GOODE, Proc. U. S. Nat. Mus. 1883, 109.

3064. *MONOLENE ATRIMANA*, Goode & Bean.

Head  $4\frac{1}{2}$  in length of body; depth about 3. D. 124; A. 100; scales 30-105-32; eye  $2\frac{1}{2}$  in head; maxillary 3; highest dorsal ray 2; left ventral  $3\frac{1}{2}$ ; pectoral  $4\frac{1}{2}$  in body; caudal 6. Body rather elongate; snout slightly produced. Mouth oblique, the maxillary extending to a little behind front of lower eye, teeth uniserial, well developed on both sides; lower eye in advance of upper; interorbital a very narrow ridge, about 9 in eye; nostrils in very short tubes in the same line with the interorbital ridge, the posterior one is slightly less distant from lower eye than the anterior one is from the snout; head everywhere scaly; lateral line strongly arched over anterior third of pectoral. Origin of dorsal on blind side above front of lower eye, longest rays in posterior fourth of fin; highest rays of anal a little higher than dorsal rays; pectoral on eyed side only; caudal sessile, rounded. Color light brownish gray, right ventral pale, other fins dusky; pectoral and eyelids black. West Indies. The type was taken by the *Blake* in 288 fathoms, off Barbados; its length is 114 mm. (Goode & Bean.) (*ater*, black; *manus*, hand.)

*Monolene atrimana*, GOODE & BEAN, Bull. Mus. Comp. Zool., XII, 155, 1896, deep waters off Barbados; JORDAN & GOSS, Review Flounders and Soles, 280, 1889; GOODE & BEAN, Oceanic Ichthyology, 455, fig. 358, 1896.

## Family CCXX. SOLEIDÆ.

## (THE SOLES.)

Body oblong or elongate, usually scaly; mouth very small, much twisted toward the eyed side; the teeth in villiform bands, very small or obsolete; eyes small, close together, with or without a bony ridge between them; edge of preopercle adnate, concealed by the skin and scales; gill openings narrow, the gill membranes adnate to the shoulder girdle above; pectoral fins small or wanting; ventral fins small, 1 or both sometimes wanting. Small fishes living on sandy bottoms, similar to the *Pleuronectidæ* in structure, but much degraded, the fins and teeth having lost many of their distinctive qualities. The vertebrae are usually in increased numbers.\*

\* The following are the numbers of vertebrae in several species of *Soleidæ*:

## I.—ACHIRINÆ.

<i>Achirus fasciatus</i> .....	8 + 20 = 28
<i>Achirus inscriptus</i> .....	9 + 19 = 28

## II.—SOLEINÆ.

<i>Synaptura zebra</i> .....	[8 + 41 = 49]
<i>Solea solea</i> .....	9 + 40 = 49
<i>Solea kleini</i> .....	10 + 37 = 47
<i>Solea aurantiaca</i> .....	[46]
<i>Quensella ocellata</i> .....	9 + 28 = 37
<i>Microchirus luteus</i> .....	8 + 29 = 37
<i>Monochirus hispidus</i> .....	9 + 25 = 34

## III.—CYNOGLOSSINÆ.

<i>Symphurus atricaudus</i> .....	10 + 42 = 52
<i>Symphurus nigrescens</i> .....	9 + 40 = 49
<i>Symphurus plagiata</i> .....	9 + 38 = 47

They are numerous in the warm seas, and those of sufficient size are valued as food. Genera about 12; species 150. The North American species belong to 2 subfamilies very different one from the other. The soles are naturally divisible into 3 subfamilies, each quite distinct from the others, and possibly independently descended or degraded from normal *Pleuronectida*. The *Achirinae*, or American soles, are apparently allied to the *Psettinae*, and as in the latter, the ventral fin of the eyed side extends along the ridge of the abdomen. The *Soleinae*, or European soles, show in the insertion of ventral and in other respects a strong resemblance to the *Pleuronectinae*. The more aberrant *Cynoglossinae*, or tongue fishes, are perhaps degraded *Soleinae*, but the eyes are sinistral, as in the *Psettinae*. In the *Soleinae* and *Achirinae* the eyes are dextral, as in the *Pleuronectinae*.

ACHIRINÆ:

I. Soles with the eyes on the right side and separated by a distinct bony ridge; the ventral with long base confluent with the anal. Body oblong or ovate, with the color on the right side; eyes moderate or small, the upper eye usually more or less in advance of the lower; mouth small, more or less twisted toward the blind side; teeth little developed, in villiform bands; edge of opercle adnate, usually concealed by the scales; gill openings more or less narrowed, the gill membranes adnate to the shoulder girdle above; blind side of head usually with fringes; pectoral fins small, sometimes wanting; ventral fins developed, one or both of them sometimes obsolete; scales usually ctenoid, rarely wanting; lateral line straight, usually single; right ventral with extended base, confluent with the anal fin.

a. Gill openings of moderate extent, confluent below; vertical fins well separated; body ovate in outline, the depth nearly  $\frac{1}{2}$  the length; pectoral fins rudimentary or wanting; lateral line straight; scales well developed, ctenoid, those on the head more or less enlarged, those of the blind side of the head with fringes; vertebrae about 28. ACHIRUS, 1054.

aa. Gill openings very small, separate, each reduced to a small slit below angle of opercle; right ventral beginning at the chin; pectoral fins minute or wanting; lateral line straight; snout dilated, the dorsal beginning upon it.

b. Scales present, ctenoid; caudal somewhat confluent with dorsal.

c. Left ventral rudimentary, with 2 rays. APIONICTHYS, 1055.

bb. Scales none; caudal free from dorsal and anal. GYMNACHIRUS, 1056.

CYNOGLOSSINÆ:

II. Soles with the eyes on the left side, not separated by a bony ridge. Body elongate, more or less lanceolate in outline, with the color on the left side; eyes small, very close together, with no distinct interorbital ridge between them; mouth small, twisted toward the blind side; teeth little developed, in villiform bands; gill openings narrow, the gill membranes adnate to the shoulder girdle above, joined together and free from the isthmus below; pectoral fins wanting (in the adult); ventral fins small, that of the blind side often wanting; vertical fins more or less confluent; scales ctenoid; lateral lines sometimes wanting, sometimes duplicated.

d. Ventral fin of eyed side only present, free from the anal; no pectoral fins; no lateral line; head without fringes. SYMPHURUS, 1057.

1054. ACHIRUS, Lacépède.

(AMERICAN SOLES.)

*Achirus*, LACÉPÈDE, Hist. Nat. Poiss., iv, 659, 1803 (*fasciatus*, etc.).

*Achirus*, CUVIER, Règne Animal, Ed. 2, II, 343, 1829 (restriction to *fasciatus*, etc.).

*Trinectes*, RAFINESQUE, Atlantic Journal and Friend of Knowledge, 1, 1832 (*scabra*).

*Grammichthys*, KAUP, Archiv für Naturgesch. 1858, 94 (*lineatus*; *fasciatus*); *Achirus* being restricted to *Tardachirus barbatus*, etc.

*Monochirus*, KAUP, Archiv für Naturgesch. 1858, 94 (*maculipinnis*); not of RAFINESQUE, 1814, a genus of *Soleina*.

? *Aseraggodes*, KAUP, Archiv für Naturgesch. 1858, 103 (*guttulata*).

*Baiostoma*, BEAN, Proc. U. S. Nat. Mus. 1882, 413 (*brachiale*).

*Beostoma*, JORDAN & GILBERT, Synopsis, 965, 1883; amended orthography.

Eyes and color on the right side. Body oblong, bluntly rounded anteriorly. Head small; eyes small, close together, the upper eye in advance of the lower, the two separated by a bony ridge; mouth small, somewhat turned toward the colored side; nasal flaps present, the nostril of the blind side fringed; lip of the colored side fringed; teeth very small, on blind side only; gill openings rather narrow, but confluent below, not reduced to a slit; the branchiostegal region scaled. Head closely scaled everywhere, the scales on the colored side similar to those on the body, those of the nape and chin much enlarged; scales on the blind side anteriorly with their pectinations more or less produced, forming cirri; scales of both sides extremely rough, extending on the fins. Lateral line straight, simple; edge of preopercle covered by the scales. Dorsal beginning on the snout, low in front and thickly scaled, its rays divided; anal fin similar, without spine; caudal fin free, convex; caudal peduncle very short and deep; pectoral fin of left side wanting, that of right side small or obsolete; ventral rays 3 or 4, the ventral fin of the colored side long, connected with the anal by a membrane. This strongly marked genus contains numerous species, all very closely related, and nearly all American. It has been united by Dr. Günther with *Solea*, but for no good reason, as the number of vertebrae is very much fewer than in the European soles, and the right ventral fin is decurrent along the abdomen and united with the anal in the American soles, while it is short and wholly free in all the European forms. The 2 groups belong in fact to distinct subfamilies. It is also worth noticing that the name *Achirus* is prior in date to that of *Solea*. The species with rudimentary pectoral fins have been set apart by Dr. Bean to form the genus *Baiostoma*, but the very slight development of these organs in some of the species and the evidently very close relationship of them all lead us to regard *Baiostoma* as a subgenus only. If we follow Kaup in restricting the name *Achirus* to the Asiatic group called *Pardachirus*, the present genus would receive the name of *Trinectes*. It seems to us, however, that both Lacépède and Cuvier regarded the species called by us *fasciatus* as the type of their genus *Achirus*. (*ἄχχιρ*, without hands; without pectoral fins.)

BAIOSTOMA (*βαίος*, small; *στόμα*, mouth):

a. Pectoral fins small, present at least on the right side.

b. Pectoral fin present on both sides, that of the left side rudimentary, of a single ray; that of the eyed side with about 3 rays.

c. Dorsal rays 60 to 67; anal rays about 48; scales 20; depth  $1\frac{1}{2}$  in length. Color brownish, irregularly spotted with darker, and with about 10 black vertical lines crossing the lateral line. ACHIRUS, 3065.

cc. Dorsal rays 53 to 57; scales 75 to 80; depth  $1\frac{1}{2}$  in length; scales not very rough, those of colored side with scattered, hair-like appendages, some black, others pale. Color olivaceous; head, body, dorsal, and anal fins covered with a network of dark lines; traces of about 8 dark cross streaks sometimes present. INSCRIPTUS, 3066.

- bb. Pectoral of right side only present.
- d. Dorsal rays 65 or 66; anal rays 48 to 51.
- e. Pectoral well developed, with about 6 rays; scales of eyed side without hair-like filaments; scales of lateral line 77 to 80; chin little prominent. KLUNZINGERI, 3067.
- dd. Dorsal rays 50 to 58; anal rays 38 to 48.
- f. Pectoral fin of 4 to 6 rays, considerably longer than eye; body with 8 to 10 narrow, vertical dark bars, these sometimes obsolete with age.
- g. Vertical fins all with round, dark spots, these usually especially distinct on the caudal fin; some of the scales of eyed side with black, hair-like appendages; pectoral fin with 5 or 6 rays, about 3 in head, its length equal to that from outer edge of 1 eye to outer edge of another. Head  $3\frac{1}{2}$ ; depth  $1\frac{1}{2}$ . Body with 8 narrow, vertical cross streaks. LINEATUS, 3068.
- gg. Vertical fins dark, without distinct markings. Body broad, ovate, the depth about  $1\frac{1}{2}$  in length; pectoral fin with 4 rays; scales of right side with numerous black, hair-like appendages. D. 56; A. 42; scales 70. MAZATLANUS, 3069.
- ff. Pectoral fins of 1 to 3 rays, about as long as eye.
- h. Body with 6 narrow, dark bands, these sometimes obsolete. Body rather narrowly ovate, its depth  $1\frac{1}{2}$  in length. D. 58; A. 44; scales 85. FONSECENSIS, 3070.
- hh. Body with about 10 black cross lines; depth  $1\frac{1}{2}$  in length. D. 61; A. 44; scales 60; pectoral of a single ray. FISCHERI, 3071.
- hhh. Body with very numerous (20 to 40) black cross bands, which are as broad as the interspaces.
- j. Blind side of snout with few fringes. Depth  $1\frac{1}{2}$  in length. D. 55; A. 48; scales 80. Body covered by many blackish, wavy bands; caudal with black spots. SCUTUM, 3072.

ACHIRUS:

- aa. Pectoral fins wholly wanting.
- k. Dorsal rays 46; anal rays 33; right lower lip with serrated fringes; nostril in a fringed tube; depth  $1\frac{1}{2}$  in length; head 3. Color brown, head and body with numerous large, rounded or kidney-shaped white spots, edged with dark brown; scales 70. FIMBRIATUS, 3073.
- kk. Dorsal rays 50 to 55; anal rays 37 to 46; right lower lip fringed; left nostril with some fringes; depth  $1\frac{1}{2}$  in length; head 4; none of the scales of eyed side with hair-like appendages. Color dusky olive, more or less mottled with about 8 dark vertical stripes, these varying very much in width and number; caudal spotted. FASCIATUS, 3074.
- kkk. Dorsal rays 59 or 60; anal rays 41 to 45; snout and chin without evident fringe or barbel; right lower lip fringed; head 4 in length; depth  $1\frac{1}{2}$ ; scales 64. Body with 12 black cross bands with narrower ones between; caudal spotted. PANAMENSIS, 3075.

Subgenus BAIOSTOMA, Bean.

3065. ACHIRUS ACHIRUS (Linnaeus).

D. 60 to 67; A. 48; P. right 3, left 1; scales 80. Pectorals rudimentary on both sides; right ventral fin composed of 5 rays, which are continuous with the anal. Scales on the nape and on chin twice as large as those on body; snout with a few fringes on blind side; right lower lip fringed. Height of body  $1\frac{1}{2}$  in total length (without caudal); width of interorbital space nearly equal to, or rather more than diameter of eye; upper eye

slightly in advance of lower; longest dorsal rays are in posterior fifth of the fin,  $\frac{3}{4}$  of length of head; caudal rounded, rather longer than head. Brownish, irregularly spotted with darker, and with about 10 black vertical lines crossing the lateral line. Coasts of Surinam. (Günther.) Not seen by us.

We know this species only from Dr. Günther's description. *Pleuronectes achirus*, Linnaeus, is based on a description by Gronow of some *Achirus* from Surinam. Gronow's fish agrees with the present species in having 60 dorsal rays and 48 anal rays, in being brown, with transverse black bands, with dark spots on the fins, as well as in coming from Surinam. But Gronow explicitly denies the presence of pectorals, and the present species has rudimentary pectoral fins on both sides. Probably these were overlooked by Gronow, and as no other species found in the same region has so large a number of rays, we feel justified in the use of the name *Achirus achirus* for this species. (*♂*-, without; *χείρ*, hand.)

*Pleuronectes oculis dextris, corpore glabro, pinnis pectoralibus nullis*, GRONOW, Museum, t. No. 42, Surinam.

*Pleuronectes achirus*, LINNAEUS, Syst. Nat., Ed. x, 268, 1758, Surinam; based on GRONOW.

*Solea gronovii*, GÜNTHER, Cat., IV, 472, 1862, Surinam.

*Achirus gronovii*, JORDAN, Proc. U. S. Nat. Mus. 1886, 602.

*Achirus achirus*, JORDAN & GOSS, Review Flounders and Soles, 311, 1889.

#### 3066. ACHIRUS INSCRIPTUS, Gosse.

Head  $3\frac{1}{2}$  in body; depth  $1\frac{1}{2}$ . D. 53 to 57; A. 40; scales 75 to 80; inter-orbital width less than eye; upper eye in advance of lower. Pectoral fin present on each side, that of the left side rudimentary, of a single ray; that of the eyed side with about 3; left ventral with 1 or 2 small rays, in some specimens entirely absent; right ventral joined to anal. Scales smaller and less rough than usual in this genus, those of nape scarcely enlarged on eyed side, those of blind side much fringed; scales of colored side with scattered hair-like appendages, some black, others pale. Color olivaceous; head, body, dorsal, and anal fins covered with a network of dark lines; traces of about 8 dark cross streaks sometimes present; caudal fin yellowish, nearly plain, or with a few dark dots or reticulations, its base dusky. Vertebrae  $8 + 20 = 28$ . West Indies north to Key West. Known to us from numerous specimens taken by Dr. Jordan at Key West, and from specimens from Hayti, in the museum at Cambridge. These specimens belong undoubtedly to the species called *reticulatus* by Poey, and this is apparently not different from the *inscriptus* of Gosse, as the agreement with the latter is even closer than with the former description. (*inscriptus*, written on.)

*Achirus inscriptus*, GOSSE, Nat. Sojourn Jamaica, 52, pl. 1, fig. 4, 1851, Jamaica; JORDAN, Proc. U. S. Nat. Mus. 1884, 143; JORDAN & GOSS, Review Flounders and Soles, 311, 1889.

*Monochir-reticulatus*, POEY, Memorias, II, 317, 1861, Cuba; POEY, Synopsis, 409; POEY, Enumeratio, 139.

*Solea reticulata*, GÜNTHER, Cat., IV, 472, 1862.

*Solea inscripta*, GÜNTHER, Cat., IV, 473, 1862.

*Bivotoma reticulatum*, BEAN & DRESEL, Proc. U. S. Nat. Mus. 1884, 152.

3067. ACHIRUS KLUNZINGERI (Steindachner).

Head  $3\frac{1}{2}$  in body; depth  $1\frac{1}{2}$ . D. 65; A. 51; scales 37-80-42; eye  $3\frac{1}{2}$  in head; height of dorsal and anal  $1\frac{1}{2}$ ; caudal 1. Body moderately broad; eyes in the same vertical line; interorbital as wide as length of eye; angle of mouth reaching a little past front of lower eye; right under lip fringed; scales near upper profile of head enlarged, all scales strongly ctenoid; scales of eyed side without hair-like filaments. Pectoral of right side only present, with about 6 rays; caudal round behind. Color brownish, with 9 or 10 narrow blackish cross lines; small rounded blackish spots on the membranes of each of the vertical fins, much as in *A. lineatus*. (Steindachner.) Pacific coast of tropical America; Panama to Guayaquil. (Named for Dr. C. B. Klunzinger, Professor of Zoology at Stuttgart, author of Memoirs on the Fishes of the Red Sea.)

*Solea klunzingeri*, STEINDACHNER, Zur Fische des Cauca und der Flüsse bei Guayaquil, 44, 1879, Guayaquil.

*Achirus klunzingeri*, JORDAN, Proc. U. S. Nat. Mus. 1885, 391; JORDAN & GOSS, Review Flounders and Soles, 312, 1889.

3068. ACHIRUS LINEATUS (Linnaeus).

Head  $3\frac{1}{2}$ ; depth about  $1\frac{1}{2}$ . D. 49 to 58; A. 38 to 44; scales 75 to 85. Pectoral fin of right side only developed, of 4 to 6 rays, considerably longer than eye. Body with 8 to 10 narrow vertical dark bars, these sometimes obsolete with age; vertical fins all with round dark spots, these usually especially distinct on the caudal fin; some of the scales of eyed side with black, hair-like appendages; pectoral fin with 5 or 6 rays, about 3 in head, its length equal to that from outer edge of one eye to outer edge of another. Color brown, the young spotted with whitish, the adult sometimes with darker; body with about 8 narrow vertical cross streaks of blackish. West Indies and Brazil, Florida Key to Guayaquil; common and variable.

We have placed the Florida species, *comifer* and *brachialis*, in the synonymy of *lineatus*. They differ from the latter only in the slightly smaller number of the scales and fin rays. The following table shows our count of a number of specimens from different localities:

Locality.	Species.	D.	A.	Scales.
Key West .....	<i>comifer</i> .....	50	35	55 to 67
Pensacola .....	<i>brachialis</i> .....	51	37	75 to 77
Cienfuegos .....	<i>lineatus</i> .....	54	43	85
Rio Janeiro .....	<i>maculipinnis</i> .....	57	42	85
Do.....	<i>maculipinnis</i> .....	54	44	72
Rio Grande de Sul.....	<i>maculipinnis</i> .....	49	38	70
Coary .....	<i>maculipinnis</i> .....	53	40	68
Mauacapuru .....	<i>maculipinnis</i> .....	55	42	75

It is evident from this table that neither the fin rays nor the scales form characters by which the subspecies can be absolutely distinguished. It is evident also, from the examination of large series of specimens, that the

coloration is subject to very great variations—as great as in *Achirus fasciatus*. In some of these the caudal is dark and immaculate, in others pale and usually profusely spotted. In some the ground color is nearly plain blackish, in others it is pale, usually with narrow dark cross bands, but sometimes closely spotted everywhere. The specimens examined by us are from Pensacola and Egmont Key (*brachialis*), Key West (*comifer*), Cienfuegos (Cuba, Poey), Coary, Teffy, Tapajos, Porto Alegre, Pernambuco, Cannarivieras, Manacapuru, Porto do Moz, Rio Grande do Sul, Rio Janeiro, San Matheo, Rosario, Itabapuna, Obidos, Xingu, Gurupa, Jutaby, Curaçao, Para, Bahia, Santarem, Iça, Fonteloa, San Paolo, Rio Trompetas, Sambaia, Manes, Javary, and Tabatinga. The species would appear to be one of the commonest in Brazil. (*lineatus*, striped.)

a. Var. *lineatus*.

*Passer lineis transversis notatus*, SLOANE, Jamaica, 2, 77, pl. 246, fig. 2, 1725, Jamaica.

*Pleuronectes fuscus subrotundus glaber*, BROWNE, Jamaica, 445, 1750, Jamaica.

*Pleuronectes lineatus*, LINNÆUS, Syst. Nat., Ed. X, 268, 1758, Jamaica; based on BROWNE and SLOANE; not of Ed. XII, which is *Achirus fasciatus*.

*Monochir maculipinnis*, AGASSIZ, Spix, Pisc. Brasil., 88, pl. 49, 1829, Brazil; POEY, Synopsis, 409, 1868.

*Monochir lineatus*, QUOY & GAIMARD, Voy. Uranic, Zool., 238, 1824.

*Achirus lineatus*, D'ORBIGNY, Voyage Amér. Mérid., Poiss., pl. 16, fig. 2, 1847; JORDAN & GOSS, Review Flounders and Soles, 312, 1889.

*Solea maculipinnis*, GÜNTHER, Cat., IV, 473, 1862; KNER, Novara Fische, III, 289, 1866.

*Achirus maculipinnis*, JORDAN, Proc. U. S. Nat. Mus. 1886, 602.

b. Var. *brachialis*.

*Baistoma brachialis*, BEAN, Proc. U. S. Nat. Mus. 1882, 413, Appalachicola Bay and South Florida. (Types, Nos. 26605 and 30463. Coll. Silas Stearns.)

*Baistoma brachiale*, JORDAN & GILBERT, Synopsis, 965, 1883.

*Achirus brachialis*, JORDAN, Proc. U. S. Nat. Mus. 1884, 149.

c. Var. *comifer*.

*Achirus comifer*, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1884, 31, Key West. (Coll. Dr. Jordan.)

3069. *ACHIRUS MAZATLANUS* (Steindachner).

(MEXICAN SOLE; LENGUADO DE RIO; TEIPALCATE.)

Head  $3\frac{1}{2}$  in body; depth  $1\frac{1}{2}$ . D. 56; A. 42; scales 70; eye  $7\frac{1}{2}$  in head; dorsal and anal rays  $3\frac{1}{2}$  in depth of body; caudal 3. Body broad, oval; eyes small, the upper in advance of the lower; interorbital about  $\frac{1}{2}$  the diameter of eye; nostril in a tube, placed just above middle of mouth; pectoral developed on eyed side only, with about 4 rays; origin of dorsal on tip of snout; greatest height of dorsal and anal behind their middle; scales of right side with numerous black hair-like appendages. Color brownish, with 8 or 9 narrow vertical black bars; fins dark, without distinct markings. West coast of Mexico, entering all streams; common and variable. Many specimens from Mazatlan and Rio Presidio examined by us, as also a specimen from Chiapas. (Name from Mazatlan, \* the river of the deer.)

\* "With eternal sun above thee,  
'T is not strange the tall deer loved thee,  
That he gave his name, Mazatl,  
To thy river, Mazatlan!"



*Solea mazatlanana*, STEINDACHNER, Ichth. Notizen, IX, 23, 1869, Mazatlan; JORDAN & GILBERT, Bull. U. S. Fish Comm. 1882, 108.

*Solea pilosa*, PETERS, Berliner Monatsber. 1869, 709, Mazatlan.

*Achirus mazatlanus*, JORDAN, Proc. U. S. Nat. Mus. 1885, 391; JORDAN & GOSS, Review Flounders and Soles, 313, 1889; JORDAN, Proc. Cal. Ac. Sci. 1895, 505.

3070. *ACHIRUS FONSECENSIS* (Günther).

Head  $3\frac{1}{2}$ ; depth  $1\frac{3}{4}$ . D. 58; A. 44; P. 2; scales about 85. No trace of a pectoral on the left side, that on the right not much longer than the eye; right ventral fin composed of 5 rays, which are continuous with the anal. Scales on the nape twice or thrice as large as those on the body. The upper part of the snout slightly overlaps the lower jaw. The left anterior part of the head with numerous tentacles; the right lower lip with very distinct slender fringes; nostril on the right side in a wide and short tube. The width of the interorbital space is less than the diameter of the eye; the upper eye is in advance of the lower. The rays of the vertical fins are branched; the longest dorsal rays are  $\frac{3}{4}$  of the length of the head. Caudal rounded, as long as the head. Brownish olive, with 6 pairs of deep brown vertical lines extending on the dorsal and anal fins. Pacific Coast of Tropical America. (Günther.) Described from 1 specimen  $4\frac{1}{2}$  inches long, from Gulf of Fonseca; 2 others since taken by us in Rio Presidio, near Mazatlan. (Name from Fonseca, the type locality.)

*Solea fonsecensis*, GÜNTHER, Cat., IV, 475, 1862, Gulf of Fonseca. (Coll. Sir John Richardson.)

*Achirus fonsecensis*, JORDAN & GOSS, Review Flounders and Soles, 314, 1889; JORDAN, Proc. Cal. Ac. Sci. 1895, 505.

3071. *ACHIRUS FISCHERI* (Steindachner).

(PEGE OJA.)

Head  $3\frac{1}{2}$  in body; depth  $1\frac{3}{4}$ ; caudal  $3\frac{1}{4}$ . D. 61; A. 44; P. right. 1; V. 5; scales 60 to 62. Pectoral wanting on left side, rudimentary on right, of a single ray scarcely longer than eye; right ventral connected with the anal. Scales on neck and lower portion of head  $1\frac{1}{2}$  to 2 times as large as those on body; right side of lower lip fringed; upper jaw not projecting forward over lower jaw; left side of head with dermal flaps only around corner of mouth and on lower jaw. Eyes small, the upper a little further forward than the other, and 2 in snout; breadth of forehead equaling diameter of eye. Dorsal rays increasing gradually in length to the forty-eighth, which is about  $\frac{5}{8}$  length of head. Scales strongly ctenoid, the teeth considerably largest at the middle; rays of all the fins, except of pectorals, scaled to their tips, the membranes less fully scaled; only the anterior third of the caudal membranes scaled, and between the last dorsal and anal rays the scales extend slightly upon the fins. A few black thread-like appendages on right side of body between scales. Color of right side dark gray; 2 or 3 blackish cross lines on head, about 10 on body, between them numberless spots of similar color; spots on fins, especially those on caudal, a little larger; a few large dark spots on body, irregular

and poorly defined; blind side reddish yellow. Total length about 10 cm. Rio Mamone, near Panama; known to us only from Steindachner's description and figure. (Steindachner.) Not seen by us. (Named for W. Fischer.)

*Solea fischeri*, STEINDACHNER, Beiträge Kenntniss Fluss-Fische Sudamer., 1, 13, 1879, Rio Mamone, near Panama.

*Achiris fischeri*, JORDAN, Proc. Ac. Nat. Sci. Phila. 1887, 391.

#### 3072. ACHIRUS SCUTUM (Günther).

Head  $3\frac{3}{8}$ ; depth  $1\frac{1}{4}$ . D. 55; A. 48; P. 3; scales 80. No trace of a pectoral on left side; right pectoral quite rudimentary, scarcely longer than the eye; right ventral composed of 5 rays, which are continuous with anal. Scales on nape nearly twice as large as those on the body; snout with scarcely any fringes on the blind side, right lower lip fringed. Width of interorbital space less than horizontal diameter of orbit; upper eye slightly in advance of lower. Longest dorsal rays in posterior third of fin,  $\frac{3}{8}$  length of head; caudal rounded, longer than head. Grayish; head, body, and fins with numerous blackish, irregular, waving, sometimes bifurcate, transverse bands, which are broader than the interspaces; caudal with rounded deep black spots; the left side uniform white. Pacific coast of Central America. (Günther.) Not seen by us. (*scutum*, a shield.)

*Solea scutum*, GÜNTHER, Cat., IV, 474, 1862, Gulf of Fonseca, Panama.

*Achirus scutum*, JORDAN & GOSS, Review Flounders and Soles, 314, 1889.

#### Subgenus ACHIRUS.

#### 3073. ACHIRUS FIMBRIATUS (Günther).

Head 3; depth  $1\frac{1}{4}$ . D. 46; A. 33; scales 70. Pectorals none; right ventral of 5 rays, which are continuous with the anal. Scales on nape 4 times, those on the chin twice, as large as on the body. Upper part of the snout slightly bent downward over the mandible and forming a short hook; right lower lip broadly fringed, each fringe being serrated; nostril in a short, wide, fringed tube. No tentacles on left side of head. Width of interorbital space equaling diameter of circular small orbit; upper eye slightly in advance of lower. Longest dorsal rays  $\frac{3}{8}$  of length of head; rays of vertical fins branched; caudal rounded, its length being  $\frac{1}{4}$  of the total. Brown; head and body with numerous large, rounded, or kidney-shaped white spots, edged with dark brown. Gulf of Fonseca, Central America. (Günther.) Known from 1 specimen,  $3\frac{1}{8}$  inches long. (*fimbriatus*, fringed.)

*Solea fimbriata*, GÜNTHER, Cat., IV, 477, 1862, Gulf of Fonseca. (Coll. Sir John Richardson.)

*Achirus fimbriatus*, JORDAN & GOSS, Review Flounders and Soles, 315, 1889.

#### 3074. ACHIRUS FASCIATUS, Lacépède.

(AMERICAN SOLE; HOG CHOKER.)

Head 4 in body; depth  $1\frac{1}{4}$ . D. 50 to 55; A. 37 to 46; scales 66 to 75; eye 7 in head; height of dorsal and anal nearly 2; caudal  $1\frac{1}{4}$ . Body broad, regularly elliptical; mouth moderate, reaching just past front of lower

eye; right lower lip fringed; eyes very small, the upper one in advance of the lower; nostril ending in a wide tube, nearer lower eye than tip of snout; interorbital space with scales, more than  $\frac{1}{2}$  eye; head and body scaled with strongly ctenoid scales, none of them with hair-like appendages; lateral line nearly straight; gill opening short, about twice as long as maxillary. Origin of dorsal on tip of snout; last few rays of dorsal and anal rapidly decreasing, giving the fins a truncate appearance posteriorly; pectorals wholly wanting; caudal rounded. Color dusky olive, more or less mottled, and with about 8 dark, vertical stripes, these varying very much in width and in number; vertical fins with the membrane of every second or third pair of rays blackish, besides dark cloudings at base of fin; caudal with numerous longitudinally oblong spots; blind side often with round, dark spots, especially in northern specimens, usually immaculate in southern ones (var. *browni*). Vertebrae 8 + 20 = 28. South Atlantic and Gulf coast, from Cape Ann to Brazos Santiago, ascending sandy streams in shallow water. The species is the best known of the American soles, and it is common along our coast, ascending the rivers for a considerable distance above tide water. It seldom exceeds 5 or 6 inches in length, and is of but little value as food on account of its small size. Here described from a specimen, 4 inches long, from Beaufort, North Carolina. This species has not yet been recorded from the West Indies. The form found along the Gulf coast has been described as a distinct species under the name *Solea browni*. The differences are not very evident. We have compared a number of specimens from Boston (*fasciatus*) with others from Pensacola, and find the following differences, none of which is constant: In the Gulf variety (*browni*) the blind side is always immaculate, while in almost all Atlantic examples (*fasciatus*) the blind side is profusely covered with round, dark spots. In 1 specimen, however (11360, Boston), the blind side is immaculate. The darker cross streaks on the eyed side are usually broader and more numerous in southern specimens, and the scales on the blind side of the head rougher. There are no constant differences either in the fin rays or in the scales. We have examined specimens of this species from Boston, Chestertown, Tarrytown, New York, Port Monmouth, Havre de Grace, Potomac River, Neuse River, Beaufort, Charleston, Pensacola, Mobile, and Galveston. In 1 large specimen from Pensacola (11482, M. C. Z.) there is a rudiment of a pectoral fin on the eyed side. It consists of a single ray  $\frac{2}{3}$  as long as the eye. (*fasciatus*, banded.)

*Pleuronectes lineatus*, LINNÆUS, Syst. Nat., Ed. XII, 458, 1766, on a specimen from Charleston, received from Dr. Garden; not *Pleuronectes lineatus* of Ed. X.

*Achirus fasciatus*, LACÉPÈDE, Hist. Nat., Poiss., IV, 659, 662, 1803, Charleston; excl. syn., description based entirely on the Linnaean account of the fish sent by Garden; JORDAN & GOSS, Review Flounders and Soles, 315, 1889.

*Pleuronectes molleis*, MITCHILL, Trans. Lit. and Phil. Soc. N. Y., I, 1815, 388, pl. 2, fig. 4, New York.

*Pleuronectes apoda*, MITCHILL, Amer. Monthly Mag. and Crit. Rev., Feb., 1818, 244, Straits of Bahama; perhaps *A. lineatus*.

*Trinectes scabra*, RAFINESQUE, Atlantic Journal and Friend of Knowledge, I, 1832, Pennsylvania, in fresh water.

*Solea browni*, GÜNTHER, Cat., IV, 477, 1862, New Orleans; Texas.

- Achirus lineatus*, CUVIER, Règne Animal, Ed. 2, II, 343, 1829; GILL, Cat. Fishes East Coast N. Am., in Rept. U. S. Fish Comm., 1871-72, 794; JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1878, 368; GOODE, l. c., 1879, 110; GOODE & BEAN, l. c., 1879, 123; BEAN, l. c., 1880, 77; JORDAN & GILBERT, l. c., 1882, 618; BEAN, l. c., 1883, 365.
- Grammichthys lineatus*, KAUP, Archiv für Naturgesch. 1858, 101.
- Achirus mollis*, STORER, Synopsis, 228, 1840; STORER, Hist. Fish. Mass., 206, pl. 32, 1867; DE KAY, New York Fauna: Fishes, 303, pl. 49, fig. 159, 1842.
- Achirus achirus mollis*, JORDAN, Cat. Fish. N. A., 137, 1885.
- Solea achirus*, GÜNTHER, Cat., IV, 470, 1862; not *Pleuronectes achirus* L.
- Achirus achirus*, JORDAN, Proc. U. S. Nat. Mus. 1885, 19; JORDAN, Cat. Fish. N. A., 137, 1885.
- Achirus lineatus*, var. *browni*, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1882, 305.

## 3075. ACHIRUS PANAMENSIS (Steindachner).

Head 4 in body; depth  $1\frac{1}{2}$ . D. 59; A. 45; scales 63 to 65; highest dorsal and anal spines 2 in head; caudal 1. Body broad, elliptical; angle of mouth below middle of lower eye; edge of lower lip, on the eyed side, fringed; eyes small, the upper in advance of the lower; interorbital scaled, scarcely as wide as diameter of eye; scales ctenoid; pectorals wholly wanting; origin of dorsal on end of snout; highest rays of anal and dorsal behind their middle; ventral rays short; middle rays of caudal the longest, fin sharply rounded behind. Color brown; about 12 dark cross bands on head and body; between these faint, paler cross bands, which form spots on dorsal and anal; caudal similarly spotted, the spots forming obscure cross bands. (Steindachner.) Pacific coast of tropical America, Panama.

*Solea panamensis*, STEINDACHNER, Ichthyol. Beiträge, v, 10, taf. II, 1870, Panama.

*Achirus panamensis*, JORDAN & GOSS, Review Flounders and Soles, 316, 1889.

## 1055. APIONICHTHYS, Kaup.

*Apionichthys*, KAUP, Archiv für Naturgesch. 1858, 104 (*dumertli*).

*Soleotalpa*, GÜNTHER, Cat., IV, 489, 1862 (*unicolor*).

Gill openings very small, separate, each reduced to a slight slit below angle of opercle; right ventral beginning at the chin, confluent with the anal; pectoral fins wanting or very small; lateral line present, straight; eyes small; snout dilated, the dorsal beginning upon it. Scales present, ctenoid; caudal fin somewhat confluent with dorsal. Left ventral rudimentary, with 2 rays. West Indies and Brazil. This genus is closely related to *Achiropsis*, Steindachner, of the rivers of Brazil, but in the latter genus the ventrals are both well developed. ( $\acute{\alpha}$ , not;  $\pi\acute{\iota}\omega\nu$ , fat;  $\lambda\chi\theta\upsilon\varsigma$ , fish.)

## 3076. APIONICHTHYS UNICOLOR (Günther).

D. 76; A. 57; V. right 5, left 2; scales 92. Body very flat and thin, its height being contained  $2\frac{1}{2}$  times in the total length (without caudal), the length of the head  $4\frac{1}{2}$  times. The upper part of the snout is dilated, bent downward like an aquiline nose, the end covering the symphysis of the mandibles; the cleft of the mouth is curved, the lower eye being immediately above its angle. The eyes are mere points, rather distant from

each other. The gill opening is reduced to a very small slit, the gill membrane being attached to the sides of the throat. The dorsal fin commences on the extremity of the snout and terminates at the root of the caudal, its rays are simple, and each is accompanied by a series of very small ctenoid scales; the longest rays are not quite  $\frac{1}{2}$  as long as the head, and occupy the middle and the third quarter of the fin. Caudal quite free, as long as the head, somewhat pointed. The right ventral appears as a mere continuation of the anal; the left is reduced to 2 minute rays near the vent. The scales on both sides are ctenoid, those on the neck and on the chin being twice the size of those on the body. Color uniform brownish gray. Coast of Surinam and Brazil. The above description from Günther, taken from the type of *Soleotalpa unicolor*. A specimen (No. 4677, M. C. Z.), from Obydos, Brazil, examined by us, differs in coloration, being pale brown, the body and fins profusely covered with round, dark spots of varying sizes, the largest as wide as from eye to eye. Head  $4\frac{1}{2}$ ; depth  $2\frac{3}{4}$ . D. 18; A. 56; scales 100; V. 2. Eyes reduced to points, the upper in advance of lower, near middle of length of head; gill openings small, subequal; right ventral beginning at the chin, continuous with anal; dorsal and anal slightly connected with caudal. Steindachner gives D. 72; A. 53; scales 95. Color brownish, mottled with darker spots. Probably Günther's specimen is faded. (*unicolor*, one-colored.)

*Apionichthys dumerili*, KAUP, Archiv für Naturgesch. 1858, 104, no locality; no description.

*Soleotalpa unicolor*, GÜNTHER, Cat., IV, 489, 1862, West Indies. (Coll. Scrivener.)

*Apionichthys nebulosus*, PETERS, Berliner Monatsberichte 1869, 709, Surinam.

? *Apionichthys blockeri*, HORST, Nederl. Tydschr. Dierk., IV, 30, 1878, locality unknown; specimen in Mus. Utrecht. (Description not seen by us.)

*Apionichthys unicolor*, JORDAN, Proc. U. S. Nat. Mus. 1886, 603.

*Apionichthys dumerili*, BLEEKER, Nederl. Tydschr. voor Dierkunde, II, 1805, 305; STEINDACHNER, Ichth. Beitr., VIII, 48, 1878.

#### 1056. GYMNACHIRUS, Kaup.

*Gymnachirus*, KAUP, Uebersicht der Soleine, Archiv für Naturgesch. 1858, 101 (*nudus*).

This genus differs from *Achirus* in the absence of scales; the dorsal and anal are free from the caudal. Brazil. (*γυμνός*, naked; *Achirus*.)

#### 3077. GYMNACHIRUS FASCIATUS, Günther.

Head  $4\frac{1}{2}$ ; depth  $1\frac{1}{2}$ . D. 68; A. 50; pectoral of right side present, very small, of 2 rays,  $\frac{1}{2}$  length of eye; jaws hidden in thick skin; lips and left side of head covered with fringes. Gill opening not extending upward as far as pectoral; vertical fins in thick skin. Olive, with 14 brown cross bands as broad as the interspaces, all extending on dorsal and anal, the first across snout, the second and third across eye; caudal with 3 brown bands. (Günther.) Locality unknown, probably Surinam or Brazil; a related species (*G. nudus*, Kaup; no pectoral fins. D. 51; A. 42), being described from Bahia. (*fasciatus*, banded.)

*Gymnachirus fasciatus*, GÜNTHER, Cat., IV, 488, 1862, locality unknown; JORDAN & GOSS, Review Flounders and Soles, 317, 1889.

## 1057. SYMPHURUS,\* Rafinesque.

(TONGUE-FISHES.)

*Symphurus*, RAFINESQUE, Indee d'Ittologia Siciliana, 52, 1810 (*nigrescens*).*Bibronia*, COCCO, Alcuni Pesci del mare di Messina, 15, 1844 (*ligulata*; larval form).*Plagusia*, CUVIER, Règne Animal, Ed. 2, 11, 344, 1829 (based on *Plagusia* of BROWN); name preoccupied in Crustaceans, Latreille, 1806.*Plagusia*, BONAPARTE, Catalogo Metodico, 51, 1846 (*lactea*); substitute for *Plagusia* preoccupied.*Aphoristia*, KAUP, Archiv für Naturgesch. 1858, 106 (*ornata*).*Ciliosichthys*, GILL, Cat. Fish. E. Coast N. A., 51, 1861 (*plagusia*).*Ammopleurops*, GÜNTHER, Cat., IV, 490, 1862 (*lacteus* = *nigrescens*).*Bascanius*, SCHMÖDTE, Naturhist. Tydsskr., v, 269, 1867 (*tædifer*; larval form).*Acedia*, JORDAN, in JORDAN & GOSS, Review Flounders and Soles, 321, 1889 (*nebulosus*).

Body elongate, more or less lanceolate in outline, with the eyes and color on the left side; eyes small, very close together, with no distinct interorbital ridge between them; mouth small, twisted toward the blind side; teeth little developed, in villiform bands; edge of preopercle covered by the scales; gill openings narrow, the gill membranes adnate to the shoulder girdle above, joined together and free from the isthmus below; pectoral fins wanting (in the adult); vertical fins more or less confluent; scales ctenoid; lateral line wanting. Ventral fin of eyed side only present, free from the anal; head without fringes. (σύν, together; φύω, to grow; οὐρά, tail; from the united vertical fins.)

## SYMPHURUS:

a. Scales not minute, ctenoid, 65 to 105 in number; dorsal rays 86 to 100; anal rays 70 to 87.

b. Scales rather large, about 6. Head  $4\frac{1}{2}$ ; depth  $4\frac{1}{2}$ ; color, clouded brown.

PIGER, 3078.

bb. Scales small, moderately ctenoid, 75 to 105 in a longitudinal series.

c. Dorsal and anal pale anteriorly, becoming more or less abruptly black posteriorly.

d. Caudal fin abruptly pale, at least at tip.

e. Body elongate, depth  $4\frac{1}{2}$  in length; head  $5\frac{1}{2}$ . D. 96 to 100; A. 86 or 87; scales 88 to 90. Color, grayish, speckled with brown; dorsal and anal fins black on last tenth, the caudal abruptly pale; tips of fin rays vermilion.

MARGINATUS, 3079.

ee. Body deeper, the depth  $3\frac{1}{2}$  to  $3\frac{3}{4}$  in length.

f. Color, light brown, irregularly barred and marbled with darker; dorsal and anal with 3 to 6 inky blotches posteriorly. D. 92 to 95; A. 75 to 78.

ATRAMENTATUS, 3080.

\* We follow Jordan & Goss in using the name *Symphurus* instead of *Aphoristia*, as the so-called *Ammopleurops lacteus* is a genuine member of the latter genus, and as it seems to be evident that the latter species is the original of *Symphurus nigrescens* of Rafinesque. The following is Rafinesque's description: "III. Gen. *Symphurus*. Ala caudale acuta, e rivinita all' ale dorsali, ed anali, ocej alla sinistra. Ossero. Si dovranno ragguagliare in questo genere due specie del genere *Achirus* di Lacepede, cioè gli *A. bilineatus*, e *A. ornatus*. Sp. no. 44. *Symphurus nigrescens*. Nerastro senza fuscio, allungato, una sola linea laterale da ogni lato."

This single lateral line assumed to distinguish *Ammopleurops* from *Aphoristia* is not a real lateral line, but a depression along the median line produced by the junction of the muscles. The species of *Symphurus* are somewhat numerous and very closely allied. With the exception of the European *Symphurus nigrescens*, all of them are American. The development of the species is imperfectly known. According to Giglioli, the larvae called *Bibronia*, may belong to this genus, and so possibly may *Charybdia*. The name *Plagusia* belongs properly to the present genus rather than to the type of *Plagusia bilineata*, to which it has been restricted by Kaup and Günther. It is, however, preoccupied in crustaceans, and in any case, both *Plagusia* and the substitute name *Plagusia* are antedated by the name *Symphurus*.

*ff.* Color, light olive, with numerous roundish brownish black spots much larger than eyes, dorsal and anal black, with narrow white margin. D. 94; A. 77; scales 95.

FASCIOLARIS, 3081.

*dd.* Caudal fin black, as is a large part of dorsal and anal, the black either continuous or in the form of large spots. Color brownish, often mottled, usually with more or less distinct darker cross bands and with longitudinal streaks along the rows of scales, sometimes nearly plain brown.

*g.* Scales quite small, 98 to 105.

*h.* Body decidedly elongate, the depth about  $4\frac{1}{2}$  in length. D. 97; A. 82; scales 98. ELONGATUS, 3082.

*hh.* Body less elongate, the depth  $3\frac{1}{2}$  in length; head  $5\frac{1}{2}$ ; longitudinal streaks very distinct. D. 100; A. 80; scales 105. ATRICAUDUS, 3083.

*gg.* Scales rather larger, 75 to 90.

*i.* Body rather elongate, the depth  $3\frac{1}{2}$  to  $4\frac{1}{2}$ ; dorsal rays 80 to 99; anal 80 to 85; opercular flap large; body with 3 or 4 dark cross bands. LEEI, 3084.

*ii.* Body less elongate, the depth  $3\frac{1}{2}$  to  $3\frac{3}{4}$  in length; the head  $5\frac{1}{2}$  to  $5\frac{3}{4}$ . D. 90 to 95; A. 75 to 80.

PLAUSIA, 3085.

*cc.* Dorsal and anal pale throughout, or more or less mottled or spotted with darker, the caudal similarly colored, not distinctly black. Body not very elongate, the depth 3 to  $3\frac{1}{2}$  in length. (Probably all varieties of *S. plagiosa*.)

*j.* Body with dark cross bands more or less distinct; the fins mottled or speckled; upper eye slightly in advance of lower.

*k.* Dorsal rays 86 to 95; anal rays 75 to 80; head 5 in length; depth  $3\frac{1}{2}$ ; scales 85 to 93; cross bands more distinct than in related species. PLAGUSA, 3086.

*kk.* Dorsal rays 78 to 85; anal rays 70 to 72; head 5 in length; depth  $3\frac{1}{2}$ ; scales 80 to 90. Color, light brown, with darker cross bars, which become obsolete with age. PUSILLUS, 3087.

*jj.* Body uniform grayish, without cross bands; last part of dorsal and anal with 3 or 4 oblong black blotches, each somewhat larger than the eye; upper eye directly above lower; head  $5\frac{3}{4}$  in length. Scales 85; D. 92; A. 75. DIOMEDEANUS, 3088.

*ddd.* Caudal and posterior part of dorsal and anal not black, scarcely darker than anterior part; scales 92; D. 93; A. 73. WILLIAMSII, 3089.

ACEDIA (Spanish name of *Symphurus plagusia* at Havana):

*aa.* Scales very small, ctenoid, each with a median dark streak, which simulates a keel, but is not a ridge; snout and jaws naked; fin rays in increased number.

*l.* Head  $5\frac{3}{4}$ ; depth  $4\frac{3}{4}$ . D. 110; A. 107; scales 120. Grayish, everywhere mottled with brown. NEBULOSUS, 3090.

#### Subgenus SYMPHURUS.

#### 3078. SYMPHURUS PIGER (Goode & Bean).

Head  $4\frac{1}{2}$  in total length; depth  $3\frac{1}{2}$ . D. 90; A. 69 to 75; ventral 4; scales 65-34 (transverse); eye 6 in head; snout  $4\frac{1}{2}$ ; mouth oblique, curved, its angle below middle of lower eye; teeth feeble, closely placed, a little stronger on colored side; nostril tubular, a little nearer eye than tip of snout; eyes moderate in size, very close together, the upper very slightly in advance, its distance from the dorsal outline equal to its diameter; scales large, ctenoid, deciduous. Dorsal beginning over middle of upper eye; longest dorsal and anal rays 3 in depth of body; pectorals obsolete.

Color grayish and brownish, with a submetallic luster upon the scales when examined separately; the denticulations of the scales dark and prominent, giving a clouded general aspect; some of the smaller specimens with large, irregular, brownish blotches above, and a dark subcircular blotch near the root of the tail, its diameter twice eye; colorless below. (Goode & Bean.) West Indies and Gulf of Mexico, in deep water; a well-defined species. (*piger*, sluggish.)

*Aphoristia pigra*, GOODE & BEAN, Bull. Mus. Comp. Zool., XII, 5, 154, 1886, St. Kitts, in about 250 fathoms (Coll. Blake); GOODE & BEAN, Oceanic Ichthyology, 460, 1896.  
*Symphurus piger*, JORDAN & GOSS, Review Flounders and Soles, 328, 1889.

#### 3079. SYMPHURUS MARGINATUS (Goode & Bean).

Head  $5\frac{1}{2}$  in total length; depth  $4\frac{1}{4}$ . D. 96 to 100; A. 86 or 87; ventral 4; scales 88 to 90; eye  $4\frac{1}{2}$  in head; snout  $4\frac{1}{2}$ . Body slender, lanceolate; mouth moderate, oblique, curved, its angle below front of pupil of upper eye; dentition feeble; eyes moderate, close together, the upper very slightly in advance; nostril in a long slender tube, midway between lower eye and tip of snout; scales moderate, strongly and sharply denticulate, not keeled; origin of dorsal above posterior margin of upper eye; anal scarcely so high as dorsal; median caudal rays short. Color in life, reddish gray, much speckled with brown; belly bluish gray; bases and membranes covering fin rays dark brown; caudal abruptly pale; tips of dorsal and anal rays and some of the membrane covering caudal rays vermilion. West Indies, in deep water. Described from a specimen, 102 mm. in length, collected by the Blake at Station CLXXXI, in 321 fathoms. (Goode & Bean.) (*marginatus*, edged.)

*Aphoristia marginata*, GOODE & BEAN, Bull. Mus. Comp. Zool., XII, No. 5, 153, 1886, off St. Vincent (Coll. Blake); GOODE & BEAN, Oceanic Ichthyology, 459, fig. 376, 1896.  
*Symphurus marginatus*, JORDAN & GOSS, Review Flounders and Soles, 323, 1889.

#### 3080. SYMPHURUS ATRAMENTATUS, Jordan & Bollman.

Head  $4\frac{2}{3}$  to 5; depth  $3\frac{1}{4}$  to  $3\frac{3}{4}$ . D. 92 to 95; A. 75 to 78; scales 95 to 100, 38 in a cross series. Body more elongate than in *S. atricaudus*. Eyes larger, the upper in advance of lower, vertical diameter of each  $3\frac{1}{2}$  to 4 in head. Cleft of mouth somewhat more curved than in *S. atricaudus*, otherwise similar. Scales larger than in *S. atricaudus*; spines on posterior margin not so strong. Ventral fins (measured from angle of gill opening)  $2\frac{2}{3}$  to 3 in head. Color light brown, irregularly barred and marbled with darker; several irregular grayish bars most distinct on posterior parts, a distinct narrow, dark bar behind gill opening; anterior part of dorsal and anal fin pale, posterior dark; anterior part with 4 to 7 dusky oblique areas, posterior part with 3 to 6 roundish inky-black spots; caudal black, narrowly tipped with white; each scale with a narrow dark edge. Length about  $4\frac{1}{2}$  inches. Pacific Ocean off Colombia, in water of moderate depth; common. Related to *Symphurus atricaudus* (Jordan & Gilbert), but distinguished by having 3 to 6 black oblong blotches on posterior part of dorsal



and anal; the general coloration darker; the scales and eyes larger. (*atramentatus*, inked.)

*Symphurus atramentatus*, JORDAN & BOLLMAN, Proc. U. S. Nat. Mus. 1889, 177, off coast of Colombia, at Albatross Station 2795, Lat. 7° 57' N., Long. 78° 55' W., in 33 fathoms. (Type, 41157, U. S. Nat. Mus. Coll. Albatross.)

3081. SYMPHURUS FASCIOLARIS, Gilbert.

Depth  $3\frac{3}{8}$  in length; head  $5\frac{1}{8}$ . D. 94; A. 77; scales 95. Eye small, 7 in head; cleft of mouth reaching to below middle of lower eye. Color light olive, with numerous roundish brownish-black spots much larger than eye, the largest arranged in 5 vertical dusky cross bars, the spots being connected by a darker ground color; a vertical dusky streak through eye; a wide dusky cross bar, bounded by darker lines on cheeks; dorsal and anal posteriorly black, with narrow white margin; caudal jet-black, with white edge; ventral white. Gulf of California, where several specimens were dredged by the *Albatross*, in shallow water. (Gilbert.) (*fasciolaris*, with narrow bands.)

*Symphurus fasciolaris*, GILBERT, Proc. U. S. Nat. Mus. 1891, 566, Gulf of California. (Coll. Dr. Gilbert.)

3082. SYMPHURUS ELONGATUS (Günther).

Head  $5\frac{1}{8}$  in body; depth  $4\frac{3}{8}$ . D. 97; A. 82; scales 98 to 105; eye 10 or 11 in head; gape of mouth  $3\frac{1}{4}$ ; caudal  $2\frac{1}{4}$ . Body extremely elongate; mouth strongly curved, reaching past lower eye; eyes in contact, the upper in advance; opercle vertical behind, divided into 2 convex flaps by a concave portion, its upper end hardly reaching axis of body; scales not keeled, ctenoid. Pectorals obsolete; dorsal beginning above eye; rays of dorsal and anal short, subequal, the fins confluent with the caudal, which ends in a sharp point; ventral of blind side obsolete, that of eyed side on the body ridge, separated from the anal. Color brownish, often mottled, usually with more or less distinct darker cross bands, and with longitudinal streaks along the rows of scales, sometimes nearly plain brown; caudal fin black, as is a large part of the dorsal and anal, the black either continuous or in the form of large spots. Pacific coast of Central America; not rare. Here described from a specimen, 6 inches long, from *Albatross* Station 2804, in Panama Bay, in 47 fathoms. (*elongatus*, elongate.)

*Aphoristia ornata*, var. *elongata*, GÜNTHER, Fishes Centr. Amer., 473, 1869, Panama.

*Aphoristia elongata*, JORDAN & GILBERT, Bull. U. S. Fish. Comm. 1882, 111.

*Symphurus elongatus*, JORDAN & GOSS, Review Flounders and Soles, 323, 1885.

3083. SYMPHURUS ATRICAUDUS (Jordan & Gilbert).

(SAN DIEGO SOLE.)

Head  $5\frac{1}{8}$ ; depth  $3\frac{1}{8}$ . D. 100; A. 80; scales 105. Body oblong-lanceolate, anteriorly somewhat blunt, regularly narrowed behind and ending in a point; the snout rather abruptly truncate; eyes and color on the left side. Eyes very small, nearly even behind, the upper eye the larger and extending farther forward. A single nostril in front of interorbital space, and

apparently a single smaller one below it. Mouth moderate, extending to opposite eye, somewhat turned toward eyed side; lips large, not fringed, the upper with a small black papilla in advance of lower eye, this apparently normal, but it may be a detached piece of skin, hardened by the alcohol; upper jaw scarcely produced, not forming a hook. Teeth small, on blind side only, the edge of the jaw on eyed side forming a smooth ridge. Gill openings narrow, not extending up to level of mouth. Scales very small, ctenoid, pretty regular over the body, much smaller on the head, the rows of scales rendered very distinct by black dots, the stripes converging toward the snout; scales on the 2 sides of the body similar; no lateral line on either side; about 105 scales (100 to 110) in a longitudinal series from the head to the tail, 45 to 50 in cross series. Dorsal fin beginning on head, continuous with anal around the tail; ventral fin of colored side only present, nearly on ridge of abdomen, and separated from the anal by an interval  $\frac{1}{2}$  longer than cleft of mouth; rays of middle parts of dorsal and anal fins with a fleshy border at base on blind side. Coloration brownish olive, with vertical dark half bars, irregular in size and position, some of them coming down from the back and others up from the belly, these posteriorly nearly meeting, but anteriorly alternating; streaks of dark points along the rows of scales, these forming very distinct longitudinal streaks; posterior part of dorsal and anal broadly edged with black; right side plain white. San Diego to Cape San Lucas, in sandy bays; common in the bay of San Diego, in which locality the numerous specimens before us were taken. A small specimen,  $\frac{1}{2}$  inches long, with light spots on the colored side and a pale ocellation on the black of the tail, taken by Mr. Lyman Belding near Cape San Lucas, probably belongs to the same species. (*ater*, black; *cauda*, tail.)

*Aphoristia atricauda*, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1880, 23, San Diego; JORDAN & GILBERT, Synopsis, 842, 1883; JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1882, 380; JORDAN, Proc. U. S. Nat. Mus. 1886, 54.

*Symphurus atricauda*, JORDAN & GOSS, Review Flounders and Soles, 324, 1889.

#### 3084. SYMPHURUS LEEI, Jordan & Bollman.

Head 4 to  $4\frac{1}{2}$  ( $4\frac{1}{3}$  to  $4\frac{2}{3}$ ); depth  $3\frac{1}{2}$  to 4 ( $4\frac{1}{3}$  to  $4\frac{2}{3}$ ). D. 95 to 100; A. 80 to 85; scales 80 to 90, 35 to 38 in a cross series; ventrals  $3\frac{1}{2}$  to  $3\frac{3}{4}$  in head. Body more elongate than in *S. atricaudus* or *S. atramentatus*, approaching that of *S. elongatus*; outline of under part of head more oblique than in the other Pacific coast species. Eyes larger than in the preceding species, the upper in advance of lower, their vertical diameter 5 to  $5\frac{1}{2}$  in head; cleft of mouth extending slightly farther back than in *S. atricaudus* or *atramentatus*, but not beyond eye as in *S. elongatus*; maxillary reaching posterior border of eye,  $3\frac{1}{2}$  to 4 in head; snout  $5\frac{1}{2}$  to  $5\frac{3}{4}$  in head. Scales comparatively large, not so firmly embedded as in *S. atricaudus* or *atramentatus*, those on opercles rather large. Opercular flap larger than in other Pacific species. Color light brown, speckled with darker, and with 3 or 4 broad black cross bands, width of median bands  $2\frac{1}{4}$  to 3 in head, the posterior band widest; caudal and the posterior  $\frac{2}{3}$  of the dorsal and anal

black; no black spots on dorsal; scales thickly punctulate, but with no distinct darker edgings. Related to *Symphurus atricaudus* (Jordan & Gilbert), but the body with 4 wide black cross bands, and the form more elongate. Bay of Panama. Many specimens of this species were obtained at Albatross Station 2804, at a depth of 47 fathoms. It is evidently very different from *S. atramentatus*, and needs comparison only with *S. elongatus*, from which it seems to be sufficiently distinct. Length of type 4½ inches. (Named for Prof. Leslie A. Lee and Mr. Thomas Lee, naturalists on board the Albatross when the species was discovered.)

*Symphurus leei*, JORDAN & BOLLMAN, Proc. U. S. Nat. Mus. 1889, 178, Lat. 8° 16' 30" N. Long. 79° 37' 45" W. (Type, No. 41134. Coll. Prof. L. A. Lee and Mr. Thomas Lee.)

## 3085. SYMPHURUS PLAGUSIA\* (Bloch &amp; Schneider).

(ACEDIA.)

Head 5½ to 5¾; depth 3¼ to 3½ in length. D. 90 to 95; A. 75 to 80; scales 75 to 85. Body rather elongate. Color brownish, often mottled, usually with more or less distinct darker cross bands, and with longitudinal streaks along the rows of scales, sometimes nearly plain brown; caudal black, including a large part of dorsal and anal, the black continuous as in the form of spots. West Indies to Brazil; Cuba to Rio Janeiro; common. The numerous specimens of this species examined by us are from Havana, Pernambuco, Santos, Rio Janeiro, Curuça, and Victoria. (*plagusia*, an old name, from *πλάγιος*, oblique.)

*Plagusia*, BROWNE, Jamaica, 445, No. 1, 1756, Jamaica.

*Pleuronectes plagusia*, BLOCH & SCHNEIDER, Syst. Ichth., 162, 1801, Jamaica; after BROWNE.  
*Achirus ornatus*, LACÉPÈDE, Hist. Nat. Poiss., IV, 659, 1803, on a specimen "presented by Holland to France."

*Plagusia tessellata*, QUOY & GAIMARD, Voyage Uranie, Zoologie, 240, 1824, Rio Janeiro.

*Plagusia brasiliensis*, AGASSIZ, Spix, Pisc. Brasil., 89, tab. 50, 1827, Brazil.

\* The synonymy of this species is somewhat doubtful. The original type of *Pleuronectes plagusia* was sent to Linnæus by Dr. Garden, of Charleston. It would therefore appear probable that this specimen represented the species of this genus which is found on the Carolina coast. But this typical specimen is still preserved in the rooms of the Linnean Society in London, where it has been examined by Goode and Bean. From their notes (Proc. U. S. Nat. Mus. 1885, 196) we quote:

"The type of this species may have come from Africa or India. There is considerable doubt as to its origin. (See Garden's Correspondence with Linné, p. 314.) D. ca 92, A. ca 80; scales 77. The species is more elongate than our specimens of *Aphoristia plagusia*, so called, the depth being contained in the total length without caudal 4½ times and the head 6 times."

As, however, no species of this genus are yet known from Africa or India, it is rather probable that Garden's fish actually came from Charleston. The greater slenderness of the original type is perhaps due to distortion, and the smaller number than usual of the scales does not afford a marked distinction. The name *Achirus ornatus* is also doubtful in its proper application. The only thing distinctive in the description of Lacépède is that the typical specimen was "given by Holland to France." Many of the species in this Dutch collection seem to have come from Surinam, and this is probably no exception. But Lacépède's description might apply as well to any other species of *Symphurus* as to this. The name *Pleuronectes plagusia*, given by Schneider to the species described by Browne, seems to admit of no doubt, as this is the only one of the group yet known from Jamaica. If, therefore, the name *Symphurus plagusia* be used for the northern species, or dropped altogether as not identified, the present species will stand as *Symphurus plagusia*. We have compared numerous specimens from Rio Janeiro (representing the nominal species *tessellata* or *brasiliensis*) with others (*plagusia*—*ornata*) from Havana. There is certainly no permanent difference. The Brazilian specimens are a little more slender on an average, but there are numerous exceptions, and all variations in color are found in both.

*Aphoristia ornata*, KAUP, Archiv für Naturgesch. 1858, 106; GÜNTHER, Cat., IV, 490, 1862; POEY, Synopses, 409, 1808; POEY, Enumeratio, 140, 1875; KNER, Novara Fische, III, 292; D. 90; A. 75; depth  $3\frac{1}{2}$  in length.

*Aphoristia plagiosa*, JORDAN, *Proc. U. S. Nat. Mus.* 1880, 59; not *S. plagiosa* of this paper.  
*Symphurus plagiosa*, JORDAN & GOSS, Review Flounders and Soles, 324, 1889.

3086. SYMPHURUS PLAGIOSA\* (*Aphoristia*)

(FORAGE FISH.)

Head 5; depth 3 to  $3\frac{1}{2}$ . D. 86 to 95; A. 75 to 80; scales 85 to 93. Body not very elongate. Body grayish, with dark cross bands more distinct than in related species; dorsal and anal more or less mottled or spotted with darker; caudal similarly colored, not distinctly black. South Atlantic and Gulf coasts of the United States, from Cape Hatteras to Pensacola and Key West, replacing *S. plagiosa* northward, the species as similar as the two names; very common on the sandy shores of our South Atlantic and Gulf States. Our numerous specimens are from Beaufort, Charleston, Pensacola, and Key West. Those from Key West nearly plain gray, as would be expected in fishes taken from the coral sands. (*πλάγιος*, oblique.)

*Pleuronectes plagiosa*, LINNÆUS, Syst. Nat., Ed. XII, 455, 1766, in a specimen from Dr. Garden, probably from Charleston, but the locality not quite certain; and of various copyists.

*Plagusia fasciata*, HOLBROOK MS., DE KAY, New York Fauna: Fishes, 304, 1842, Charleston.

*Glossichthys plagiosa*, GILL, Cat. Fish. East Coast N. Am., 51, 1861.

*Plagusia plagiosa*, GILL, Cat. Fish. East Coast N. Am., 794, 1873.

*Aphoristia plagiosa*, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1878, 368; JORDAN, l. c., 1880, 22; JORDAN & GILBERT, l. c., 1882, 305 and 618; JORDAN & GILBERT, Synopses, 842, 1883; JORDAN, Proc. U. S. Nat. Mus. 1884, 144.

*Aphoristia fasciata*, JORDAN, Proc. U. S. Nat. Mus. 1886, 53.

*Symphurus plagiosa*, JORDAN & GOSS, Review Flounders and Soles, 325, 1889.

## 3087. SYMPHURUS PUSILLUS (Goode &amp; Bean).

Head 5 in total length; depth  $3\frac{1}{2}$ . D. 78; A. 70; scales 85 to 90-95 (transverse); eye  $5\frac{1}{2}$  in head; snout  $5\frac{1}{2}$ ; length of gape of mouth  $1\frac{1}{2}$ . Body slender, lanceolate; mouth small, oblique, curved, its angle under anterior margin of pupil of lower eye; dentition feeble; eyes small, close together, in the same vertical line; tubular nostril midway between lower eye and tip of snout; scales small, strongly ctenoid; jaws and snout scaled. Dorsal beginning above middle of eye, its highest rays  $2\frac{1}{2}$  in depth of body; greatest height of anal 3; median caudal rays short; ventrals well separated from anal. Color light brown, with 6 or 7 cross bars of slightly

\* A specimen of *Symphurus*, nearly 6 inches long, collected at Beaufort, North Carolina, by Prof. O. P. Jenkins, seems referable to *Symphurus pusillus* rather than to the typical *plagiosa*. It is highly mottled in coloration, the body and fins being profusely speckled and blotched with blackish, besides 9 or 10 rather distinct cross bands. D. 82; A. 72; scales about 80. Depth  $3\frac{1}{2}$  in length. Another large specimen, 7 inches long, from the Florida Keys, is in the museum at Cambridge. This has: D. 82; A. 72; scales 76. Depth 3 in length. Color brown, almost plain, except that the fins are mottled, especially posteriorly; caudal fin not black. If these two specimens are really typical of *Symphurus pusillus*, it probably can not be separated as a species from *S. plagiosa*.

darker hue; fins pale, with dusky blotches; blind side white. (Goode & Bean.) Gulf Stream, in deep water. Very close to *Symphurus plagiusa*. (*pusillus*, weak.)

*Aphorista pusilla*, GOODE & BEAN, Proc. U. S. Nat. Mus. 1885, 590, Gulf Stream, Lat. 40° N., in deep water; GOODE & BEAN, Oceanic Ichthyology, 461, fig. 379, 1896.  
*Symphurus pusillus*, JORDAN, Proc. U. S. Nat. Mus. 1889, 651.

## 3088. SYMPHURUS DIOMEDEANUS (Goode &amp; Bean).

Head  $5\frac{2}{3}$  in body; depth  $3\frac{1}{2}$ . D. 96 (including  $\frac{1}{2}$  of caudal); A. 79; scales 85; eye 6 in head; snout 5; caudal 10 in total length. Mouth oblique, curved, its angle below front of eye, teeth very feeble; nostril tubular, nearer eye than tip of snout; eyes moderate, equal, very close together, upper eye directly over the lower; scales moderate, somewhat deciduous, ctenoid; jaws and snout with small thin scales. Origin of dorsal above middle of upper eye, highest rays  $3\frac{1}{2}$  times depth of body; ventrals well separated from the anal. Color uniform gray, lighter below, the scales above somewhat metallic in luster; the last fourth of dorsal with 3 oblong black blotches somewhat larger than eye, the anal with 4, similar in position; in the young there is a slight brownish marginal line upon each scale, and an appearance of indistinct cloudings of brown upon the colored side. Off Trinidad and Dominica and in the Gulf of Mexico. The specimen here described was collected by the *Albatross* at Station 2414, in the Gulf of Mexico, north of the Tortugas, at a depth of 26 fathoms; its length is 140 mm. Other specimens were dredged by the *Albatross* at Station 2362, in Lat. 22° 08' 30" N., Long. 86° 53' 30" W., in 25 fathoms, and at Stations 2191 and 2122, between Lat. 10° 37' 40" N., Long. 61° 42' 40" W., and Lat. 10° 37' N., Long. 61° 41' 22" W., in 31 to 34 fathoms. Specimens were also secured by the *Blake* at Stations XXIV and XXV, off Dominica. (Goode & Bean.) Evidently very close to *Symphurus plagiusa*. (*Diomedea*, the *Albatross*; from the name of the steamer by which most of the deep-sea explorations of the United States Fish Commission have been accomplished.)

*Aphorista diomedea*, GOODE & BEAN, Proc. U. S. Nat. Mus. 1885, 589, Gulf of Mexico, Lat. 25° 04' 30" N., Long. 82° 59' 15" W. (Type, No. 37347. Coll. *Albatross*); GOODE & BEAN, Oceanic Ichthyology, 460, fig. 378, 1896.

## 3089. SYMPHURUS WILLIAMSII, Jordan &amp; Culver.

Head  $4\frac{1}{2}$ ; depth  $3\frac{2}{3}$ . D. 93; A. 73; scales 92. Body more slender than in *S. plagiusa*, which it much resembles, but less slender than *S. elongatus*; upper eye slightly in advance of lower. Sand color in life; light gray, everywhere finely mottled with light and dark; traces of a few very narrow dark cross bands; fins all mottled; caudal and posterior part of dorsal and anal not black, scarcely darker than anterior part. Known only from Mazatlan, where 2 specimens, the larger about  $1\frac{1}{2}$  inches long, were obtained by Mr. T. M. Williams, in tide pools with sandy bottom in very shallow water near the estuary. (Named for Thomas Marion Williams, a student in biology in Stanford University, discoverer of the species.)

*Symphurus williamsii*, JORDAN, Proc. Cal. Ac. Sci. 1895, 506, pl. 55, Mazatlan. (Coll. Hopkins Exped. to Mazatlan.)

## Subgenus ACEDIA, Jordan.

## 3090. SYMPHURUS NEBULOSUS (Goode &amp; Bean).

Head  $5\frac{1}{2}$  in total length; depth  $4\frac{1}{2}$ . D. 119 (to middle of base of caudal); A. 107; V. 5; scales 120; eye  $7\frac{1}{2}$  in head; snout 5. Body slender; angle of mouth below front of lower pupil; teeth feeble, very slender, and rather closely placed, apparently equally developed on both sides; eyes small, close together, separated by a single row of scales, the upper one very slightly in advance; tubular nostril nearer eye than tip of snout; scales small, ctenoid, each with a median dark streak (but not keeled, as erroneously stated in the original description);\* jaws and snout naked. Origin of dorsal a little behind eyes, highest rays 3 in depth of body; longest anal rays twice length of snout; median caudal rays longest, twice length of snout; pectorals obsolete; ventrals well separated. Color grayish, everywhere mottled with brown; a dark median line on scales. (Goode & Bean.) Gulf stream. A well-marked species. The increased number of fin rays indicates a probability that the number of vertebrae will also be found similarly increased. (*nebulosus*, clouded.)

*Aphoristia nebulosa*, GOODE & BEAN, Bull. Mus. Comp. Zool., x, No. 5, 192, 1883, Gulf Stream, off the coast of Carolina; GOODE & BEAN, Oceanic Ichthyology, 458, fig. 375, 1896.

*Symphurus nebulosus*, JORDAN & GOSS, Review Flounders and Soles, 326, 1889.

## Order CC. PEDICULATI.

## (THE PEDICULATE FISHES.)

Carpal bones notably elongate, forming a kind of arm (pseudobrachium) which supports the broad pectoral. Gill opening reduced to a large or small foramen situated in or near the axil, more or less posterior to the pectorals. Ventral fins jugular if present; anterior dorsal reduced to a few tentacle-like, mostly isolated spines; soft dorsal and anal short; no scales. First vertebra united to cranium by a suture; epiotics united behind supraoccipital; elongate basal pectoral radii (actinosts) reduced in number; no interclavicles; post-temporal broad, flat, simple; upper pharyngeals 2, similar, spatulate, with anterior stem and transverse blade; basis of cranium simple; no air duct to the swim bladder. Marine fishes, chiefly of the tropics and the oceanic abysses. The group is an offshoot from the *Acanthopteri*, its chief modifications being in the elongation of the actinosts and in the position of the gill opening. Its nearest relatives among the spiny-rayed fishes are, perhaps, the *Batrachoidichthys*. (*pediculatus*, having a footstalk.)

## ANALYSIS OF FAMILIES OF PEDICULATI.

- a. Gill openings in or behind the lower axil of the pectoral; mouth large, terminal.
  - b. Pseudobranchic present; pseudobranchia with 2 actinosts; head broad, depressed, the enormous mouth with very strong teeth; ventrals present.

ЛОПННДЪ, ССХХІ.

\* The appearance of "keeled scales," described by Goode & Bean, is due to a black line on the skin under the center of each row of scales. There seems to be no real keel, and the species is congeneric with the other species of *Symphurus*.

bb. Pseudobranchie none; pseudobranchia with 3 aethiosts.

c. Ventrals present; arm angulate, the pseudobranchia elongate.

ANTENNARIIDÆ, CCXXII.

cc. Ventrals wanting, arm not angulate, the pseudobranchia moderate.

CERATHIDÆ, CCXXIII.

aa. Gill openings in or behind the upper axil of the pectoral; mouth small, usually inferior.

OCCOCEPHALIDÆ, CCXXIV.

Family CCXXI. LOPHIIDÆ.

(THE ANGLERS.)

Head wide, depressed, very large. Body contracted, conical, tapering rapidly backward from the shoulders. Mouth exceedingly large, terminal, opening into an enormous stomach; upper jaw protractile; maxillary without supplementary bone; lower jaw projecting; both jaws with very strong, unequal, cardiform teeth, some of the teeth canine-like, most of them depressible; vomer and palatines usually with strong teeth. Gill openings comparatively large, in the lower axil of the pectorals. Pseudobranchie present. Gill rakers none. Gills 3. Skin mostly smooth, naked, with many dermal flaps about the head. Spinous dorsal of 3 isolated, tentacle-like spines on the head, and 3 smaller ones behind, which form a continuous fin; second dorsal moderate, similar to the anal; pectoral members scarcely geniculated, each with 2 actinosts and with elongate pseudobranchia; ventrals jugular, I, 5, widely separated, large, much enlarged in the young. Young with the head spinous. Pyloric area present. Two genera, with 4 or 5 species, living on sea bottoms, at moderate or great depths; remarkable for their great voracity. (*Pediculati*, part, genus *Lophius*, Günther, Cat., III, 178-182, 1861.)

a. Vertebrae 27 to 31.

LOPHIUS, 1058.

aa. Vertebrae 18 or 19 only.

LOPHIOMUS, 1059.

1058. LOPHIUS (Artedi) Linnaeus.

(FISHING-FROGS.)

*Lophius* (ARTEDI) LINNÆUS, Syst. Nat., Ed. x, 1, 236, 1758 (*piscatorius*).

Characters of the genus included above. Vertebrae numerous, about 30 in number. (*Lophius*, the ancient name of *L. piscatorius*, from *λόφος*, a crest.)

3091. LOPHIUS PISCATORIUS, Linnaeus.

(COMMON ANGLER; FISHING-FROG; MONKFISH; GOSEFISH; ALL-MOUTH; BELLOWS-FISH.)

D. I-I-I, III-10; A. 9. Body depressed, tapering, scarcely longer than head. Humeral spine with 3 points, of which the posterior is the longest. Head surrounded with a fringe of barbels; top of head, in young, with many strong spines. Anterior dorsal spine elongate, fleshy at tip. Brownish, mottled, below white; mouth behind the hyoid bone immaculate; pectorals and caudal black at tip; peritoneum black. Length 3 feet. North

Atlantic, on both coasts; generally common, ranging southward along the shore to Cape Hatteras; found in deep water as far south as Barbados, in 209 fathoms, and to the Cape of Good Hope; northward to Norway and Nova Scotia. A well-known fish of singular ugliness of appearance, and of enormous voracity. (Eu.) (*piscatorius*, pertaining to an angler, in allusion to the baited dorsal spines which overhang the cavernous mouth.)

*Lophius piscatorius*,\* LINNÆUS, Syst. Nat., Ed. x, 1, 236, 1758, seas of Europe; after ARTEDI, *Lophius ore cirrhoso*, etc.; GÜNTHER, Cat., III, 179, 1861; GILL, Proc. U. S. Nat. Mus. 1878, 219; JORDAN & GILBERT, Synopsis, 844, 1883.

*Lophius americanus*, CUVIER & VALENCIENNES, Hist. Nat. Poiss., XII, 380, 1837, Philadelphia (Coll. Le Sueur); STORER, Hist. Fish. Mass., pl. 18, fig. 2, 101, 1867.

#### 1059. LOPHIOMUS, Gill.

*Lophiomus*, GILL, Proc. U. S. Nat. Mus. 1882, 552 (*setigerus*).

This genus is closely allied to *Lophius* in external characters, but it is strikingly distinguished by the reduced number of its vertebrae, which are only 18 or 19, a fact which is associated with its tropical habitat. One species from the Pacific. (*Lophius*; ἄμῶς, shoulder, in apparent allusion to the trifold humeral spine.)

#### 3092. LOPHIOMUS SETIGERUS (Vahl).

Dorsal III-III-9; A. 5. Head above orbits and laterally with numerous spines and prickles; humeral bone ending in 3 blunt points; numerous cirri scattered along sides of head and body. Vertebrae 18. Color dusky; floor of mouth black posteriorly, but without white spots; pectorals and ventrals pale on basal half, black distally; caudal and anal black, with some white spots; soft dorsal translucent, with black specks; first dorsal spine with its membranaceous tip white, the latter provided with 2 black eye-like spots. Pacific Ocean; not uncommon in rather deep water off coasts of China and Japan. Known on the American coast from 1 speci-

\* According to Professor Horace A. Hoffman this fish is called in Athens *Πεσκανδρίτζα* or *Πεσκαντρίτζα*. These names, "probably of Italian origin, meaning fisher; χλάσχα, at Chalcis, σκλημπού, and *Βατραχόψαρο* at Patras. The *βάτραχος ὁ ἄλιεύς* (the fisher frog) of Aristotle. (See Aristotle 305a 6b 4, 506b 16, 564b 18, 565b 29, 570b 30, 620b 11 ff, 695b 14, 696a 27, 749a 23, 754a 23 ff, 755a 9, 835b 13, 1527b 41-43, 540b 18.) Aristotle says with regard to the *βάτραχος*: 'Inasmuch as the flat front part is not fleshy, nature has compensated for this by adding to the rear and the tail as much fleshy substance as has been subtracted in front.' The *βάτραχος* is called the angler. He fishes with the hair-like filaments hung before his eyes. On the end of each filament is a little knob just as if it had been placed there for a bait. He makes a disturbance in sandy or muddy places, hides himself and raises these filaments. When the little fishes strike at them he leads them down with the filaments until he brings them to his mouth. The *βάτραχος* is one of the *σελάχη*. All the *σελάχη* are viviparous or ovoviviparous except the *βάτραχος*. The other flat *σελάχη* have their gills uncovered and underneath them, but the *βάτραχος* has its gills on the side and covered with skinny opercula, not with horny opercula like the fish which are not *σελαχῶδη*. Some fishes have the gall bladder upon the liver, others have it upon the intestine, more or less remote from the liver and attached to it by a duct. Such are *βάτραχος*, ἔλαψ, *συναγρίς*, *σμούραινα*, and *ξίφιας*. (This has been proved true of *Lophius piscatorius* by a dissection by Dr. C. H. Gilbert.) The *βάτραχος* is the only one of the *σελάχη* which is oviparous. This is on account of the nature of its body, for it has a head many times as large as the rest of its body, and spiny and very rough. For this same reason it does not afterwards admit its young into itself. The size and roughness of the head prevents them both from coming out (i. e., being born alive) and from going in (being taken into the mouth of the parent). The *βάτραχος* is most prolific of the *σελάχη*, but they are scarce because the eggs are easily destroyed, for it lays them in a bunch near the shore." (Hoffman & Jordan, Proc. Ac. Nat. Sci. Phila., 1892, 278.)



men, 2½ inches long, dredged at *Albatross* Station 2805, southwest of Panama. From this specimen, the above description is taken. Comparing this with a larger specimen taken at Tokio by Prof. K. Otaki, we find no differences likely to prove permanent. (*seta*, bristle; *gero*, I bear.)

*Lophius setigerus*, VAHL,\* Skrivt. Naturh., iv, 214, tab. 3, figs. 5 and 6, 1797, China Sea; CUVIER & VALENCIENNES, Hist. Nat. Poiss., xii, 383, 1837; GÜNTHER, Cat., iii, 180, 1861.

*Lophius viviparus*, BLOCH & SCHNEIDER, Syst. Ichth., 142, 1801, tab. 32, China Sea; after Vahl.

*Lophiomus setigerus*, GILBERT, Proc. U. S. Nat. Mus. 1890, 454.

Family CCXXII. ANTENNARIIDÆ.

(THE FROG-FISHES.)

Head and body more or less compressed. Mouth vertical or very oblique, opening upward; lower jaw projecting; jaws with eardiform teeth; premaxillaries protractile. Gill openings small, pore-like, in or behind the lower axils of the pectorals. No pseudobranchia. Gills 2½ or 3; skin naked, smooth, or prickly. Pectoral members forming an elbow-like angle. Pseudobranchia long, with 3 actinosts. Ventral fins present, jugular, near together. Spinous dorsal of 1 to 3 separated, tentacle-like spines; soft dorsal long, larger than anal. Pyloric caeca none. Genera about 5; species 50. Inhabitants of tropical seas, "living on floating seaweed, and enabled, by filling the capacious stomach with air, to sustain themselves on the surface of the water;" therefore widely dispersed by currents in the sea. (*Pediculati*, pt., Günther, Cat., III, 182 to 200, 1861.)

a. Head compressed; a rostral spine or tentacle, followed by 2 larger spines; palatine teeth developed; dorsal spines disconnected.

b. Skin naked and smooth; ventral fins elongate.

PTEROPHRYNE, 1060.

bb. Skin covered with prickles; ventral fins short.

ANTENNARIUS, 1061.

aa. Head enboid; a single rostral spine or tentacle, received in a groove; soft dorsal low.

CHAUNAX, 1062.

1060. PTEROPHRYNE, Gill.

(MOUSE-FISH.)

*Pterophryne*, GILL, Proc. Ac. Nat. Sci. Phila. 1863, 90 (*bougainvillei*).

*Pterophrynoides*, GILL, Proc. U. S. Nat. Mus., i, 1878, 216 (*histrion*); name a substitute for *Pterophryne*, if the latter be regarded as preoccupied by the earlier *Pterophrynus*.

Body smooth or scarcely granular, short, somewhat compressed, with tumid abdomen; mouth small, oblique; palate with teeth; wrist and pectoral fin slender; ventrals elongated; soft dorsal and anal vertically

\**Lophiomus setigerus*, is thus described by Dr. Günther:

"Dorsal III-III, 8 or 9; A. 6 or 7. Teeth arranged in 2 alternate series in the upper jaw, in 3 in the lower; 2 or 3 teeth on each side of the vomer; humeral spine terminating in 3 points; the mouth behind the hyoid bone purplish black, with white spots. Vertebrae 19, the anterior ones very short, the middle and posterior ones nearly equal in length. Coasts of China and Japan." (Günther.)

expanded. Small fishes of fantastic shape in the West Indies and Gulf Stream. (*πετρόν*, wing; *φρύνη*, toad.)

a. "Bait" on first dorsal spine bifurcate at tip.

HISTRIO, 3093.

aa. "Bait" on first dorsal spine bulbous, covered with fleshy filaments.

GIBBA, 3094.

### 3093. PTEROPHYNE HISTRIO (Linnaeus).

(MOUSE-FISH; SARGASSUM-FISH.)

Head  $2\frac{1}{2}$ ; depth  $1\frac{1}{2}$ . D. III-14; A. 7; V. 5. Skin of head and body, as well as dorsal fins, with fleshy tags, which are most numerous on the dorsal spines and abdomen. Wrist slender; ventrals large, nearly  $\frac{1}{2}$  as long as head. Dorsal and anal with the posterior rays not adnate to caudal peduncle; first dorsal spine bifurcate at tip. Yellowish, marbled with brown; 3 dark bands radiating from eye; vertical fins barred with brown; belly and sides with small white spots. Tropical parts of the Atlantic; abundant on our Gulf coast and occasional northward to Cape Hatteras or beyond, especially in floating masses of *Sargassum*. Once taken in Europe (Vadsö, Norway) in floating seaweed from the Gulf Stream. Recorded from the coast of Senegambia; its history and synonymy confused with that of the following species. A remarkable fish, excessively variable in coloration. (*histrío*, a harlequin.)

*Lophius tumidus*, OSBECK, Iter Chinensis, 400, 1757, Open Sea; pre-Linnaean.

*Lophius histrío*,\* LINNÆUS, Syst. Nat., Ed. x, 237, 1758, after various authors, especially

*Balistes quaperna* seu *chinensis*, LINNÆUS, Mus. Ad. Fr., 56.

*Pterophryne histrío*, GILL, Proc. U. S. Nat. Mus. 1878, 216; GOODE & BEAN, Oceanic Ichthyology, 486, 1896.

*Antennarius histrío*, JORDAN & GILBERT, Synopsis, 846, 1883; COLLETT, Campagnes Hiron-delle, 38, 1896.

\* Concerning the use of the name *histrío* for this species, Dr. Gill remarks:

"In 1794 (as appears from the dates on the plates), Shaw published a number of his 'Naturalists' Miscellany,' in which he described 3 fishes under the generic name of *Lophius*. These were described as (1) *Lophius striatus* (the Striated Lophius), pl. 175; (2) *Lophius pictus* (the Variegated Lophius), pl. 176, upper figure, and (3) *Lophius marmoratus* (the Marbled Lophius), pl. 176, lower figure. The originals of these are evidently the varieties (a, b, and c) of *Lophius histrío* admitted by Bloch & Schneider. It is quite clear that the first two were based on species of typical *Antennarius* (not *Pterophryne*), while the third is incomprehensible, and, if the figure is at all correct, must represent a factitious fish; it most certainly has nothing to do with *Pterophryne*. The other species, however, notwithstanding the bad figures, are readily identifiable. The *Lophius striatus* (as has recently been recognized by Günther) is the first name of an *Antennarius* peculiar to the Pacific, and quite distinct from the Caribbean *Antennarius scaber* (= *A. histrío* Günther), with which it was at first confounded by Günther. The *Lophius pictus* was evidently based on the species or variety of *Antennarius* which was afterwards named *Antennarius phymatodes* by Bleeker, and it agrees very closely, in the distribution of colors, with a specimen figured by that ichthyologist, and would probably be considered by Günther as a variety of his *Antennarius commersonii*. But whatever may be the value of the forms embraced under the name *Antennarius commersonii* by Günther—whether species or varieties—the name *Antennarius pictus* must be revived from Shaw, either especially for the *Antennarius phymatodes* of Bleeker or for the collection designated as *Antennarius commersonii*. It has thus been demonstrated (1) that the Linnaean name, *Lophius histrío*, was originally created for the common *Pterophryne*, and (2) that the names generally employed for the *Pterophryne* were originally applied to very different forms, and members of even a different genus. Hence if the laws of priority, as formulated by the British and American Associations for the Advancement of Science, are to guide us, there can be no question that the species of *Pterophryne* must hereafter be designated as *Pterophryne histrío*; if, however, it is allowable to go behind even the tenth edition of the *Systema Naturæ* and to take the oldest binomial name, without other considerations, the designation *tumidus* must be revived. It seems best, however, to follow general usage." (Gill, Proc. U. S. Nat. Mus., 1, 1878, 226.)

- Chironectes pictus*, CUVIER & VALENCIENNES, Hist. Nat. Poiss., XII, 393, 1837, Surinam.  
*Chironectes tumidus*, CUVIER & VALENCIENNES, Hist. Nat. Poiss., XII, 397, 1837, "Cabinet du Roi," Sargasso Sea. (Coll. Péron.)  
*Chironectes arcticus*, DÜBEN & KOREN, Kong. Vet. Akad. Abh. Stockholm 1844, 72, Vadsö, Norway, from a specimen carried northward in *Sargassum*; the only European record; fide COLLETT.  
*Antennarius marmoratus*, GÜNTHER, Cat., III, 185, 1861; in part; not of Cuvier.  
*Chironectes levigatus*, DE KAY, N. Y. Fauna: Fishes, 165, pl. 27, fig. 83, 1842; not of Cuvier.

3094. PTEROPHYRNE GIBBA (Mitchill).

Garman refers to this species certain specimens obtained in Gulf weed about Key West and the Tortugas. These resemble *P. histrio*, but "differ markedly in certain respects. The bait on the first dorsal spine, for instance, is bulbous and covered with slender fleshy filaments in our individuals, but in *P. histrio* it is bifurcate. *P. gibba* is fairly represented by Cuvier, 1817, in his *Chironectes levigatus*. The formula for the individuals in hand is D. III, 12; A. 7; V. 5; P. 10; C. 9." (Garman.) West Indies, north to Key West and the Tortugas; not examined by us; probably common, but hitherto confounded with *P. histrio*. (*gibbus*, gibbous.)

- Lophius gibbus*, MITCHILL, Trans. Lit. and Phil. Soc. N. Y. 1815, 1, pl. 4, f. 9, off St. Croix, Lat. 22° N., Long. 64° W. (Coll. Dr. John D. Jaques.)  
*Chironectes levigatus*, CUVIER, Mém. du Mus., III, 423, pl. 16, fig. 1, 1817, South Carolina (Coll. Bosc); CUVIER & VALENCIENNES, Hist. Nat. Poiss., XII, 399, 1837.  
*Pterophyrne levigata*, GILL, Proc. Ac. Nat. Sci. Phila. 1863, 90.  
*Chironectes sonntagii*,\* Baron J. W. VON MÜLLER, Reisen in den Vereinigten Staaten, Canada und Mexico, Band 1, 180, 1864, in floating seaweed; no exact locality stated.  
*Pterophrynoideus gibbus*, GARMAN, Bull. Iowa Lab. Nat. Hist. 1896, 81.

1061. ANTENNARIUS, Lacépède.

- Antennarius* (COMMERSON) LACÉPÈDE, Hist. Nat. Poiss., 1, 421, 1798 (*chironectes*).  
*Histrio*, FISCHER, Zoognosia, 78, 1813 (*histrio*, etc.). (No type; includes all known *Antennariidae*; description transposed with that of *Lophius* by error.)  
*Chironectes*, CUVIER, Règne Animal, Ed. 2, vol. II, 252, 1829 (*chironectes*); preoccupied in mammals, Illiger, 1811.

Body oblong, compressed, very deep through the occipital region, tapering behind; breast tumid; mouth rather large, more or less oblique, or even vertical; cardiform teeth on jaws, vomer, and palatines; eye small; skin with small granules or spinules, these usually forked, and numerous fleshy slips. First dorsal spine developed as a small rostral tentacle;

\* The following is the substance of the long account of *P. sonntagii* (Von Müller):

"D. II, 10 to 12; P. 10 or 11; C. 6 to 7 ('Strahlenpaare'); B. 6. Head and body slightly compressed; dorsal spines like little horns, covered over and over with spinous growths. Mouth wide, with numerous rows of small teeth; throat and belly with many fleshy slips. Pectorals produced on a long peduncle like the flippers of a tortoise; ventrals similar, but formed more like feet; anal fin like a rudder. Color clear yellowish green, with greenish brown stripes; a broad dark stripe across breast to root of pectoral; another on the back; another on the side, running backward in the form of a hammer, paler at last on lower part of back; several stripes and spots, more or less dusky, on the tail and other extremities; on the soft underside to the anal intense reddish golden yellow spots; between the dark streaks and the yellow ground color of the body are often white shades and markings; eye fiery orange. Atlantic Ocean or Gulf of Mexico; living in floating seaweed."

This species must be a *Pterophyrne*, and it is not evidently different from *Pterophyrne gibba*.

second and third dorsal spines strong, covered with skin, with numerous fleshy filaments; soft dorsal high and long; anal short and deep; caudal fin rounded, the peduncle free; pectoral fins wide, with a rather wide wrist, at the lower posterior angle of which are the very small gill openings; ventral fins short. Fantastic-looking fishes, often gaily colored; very numerous in warm seas. (*antenna*, *i.* feeler or tentacle.)

- a. Bulbous tip or "bait" of first dorsal spine simple, undivided at tip.
- b. Skin smoothish except about eyes; first dorsal spine short, second rough. Body brown, with whitish spots; no ocelli. INOPS, 3095.
- bb. Skin with prickles, velvety or shagreen-like.
- c. Prickles simple, none of them bifid.
- d. Color black; tips of pectorals and ventrals and one or two spots on side white (prickles undescribed). PRINCIPIS, 3096.
- dd. Color dusky; dorsal with 3 ocelli; caudal with many spots; first dorsal longer than second; no dermal flaps. TENERROSUS, 3097.
- ddd. Color reddish or grayish, reticulate with heavy black lines; first dorsal spine short. RETICULARIS, 3098.
- cc. Prickles or spinules on body mostly bifid.
- e. Body without ocelli; first dorsal spine filiform.
- f. Mouth immaculate within; body with numerous rosy and dusky tracts, the latter forming bars and concentric streaks below; fins barred. STRIGATUS, 3099.
- ff. Mouth largely black within; body blood red, with black spots on sides and below dorsal. SANGUINEUS, 3100.
- ee. Body with 3 large ocelli, 1 on dorsal, 1 on caudal, and 1 on middle of side, besides many black spots and streaks; tip of first dorsal spine fringed; mouth largely black within. OCELLATUS, 3101.
- aa. Bulbous tip or "bait" on first dorsal spine bifid at tip; skin shagreen-like.
- g. Color reddish, with brown spots, those about the eye radiating.
- h. Dermal flaps numerous on body; spinules on skin short and stiff, rendering the surface shagreen-like. SCABER, 3102.
- hh. Dermal flaps few; spinules on skin longer and slender, rendering the surface velvety. TIGRIS, 3103.
- gg. Color uniform black; surface of body rough, shagreen-like; inside of mouth white; first dorsal spine short, little longer than second. NUTTINGII, 3104.
- aaa. Bulbous tip or "bait" of first dorsal spine trifid.
- i. First dorsal ray twice as long as second and as long as caudal; sides with numerous black ocelli, besides other streaks and dark spots; skin smoothish. MULTIOCELLATUS, 3105.
- ii. First dorsal spine barely  $\frac{1}{2}$  longer than second; shorter than caudal; sides with dark streaks and reticulations; a large ocellus under middle of soft dorsal; body rough, with shagreen. RADIOSUS, 3106.

#### 3095. ANTENNARIUS INOPS, Poey.

Depth  $2\frac{1}{2}$  with caudal. Skin lustrous, smooth, except for some points behind and below eye; third of the first 3 dorsal rays largest, its membrane not reaching to vent; second ray also large, but shorter, placed between eyes; first spine developed as a fishing rod, filiform, ending in a small, membranaceous lobe, its base close to that of second, and, therefore, distant from end of snout, its spine short, the tip not reaching middle of second spine; short tentacles, like horns, on anterior part of third spine, over the nostrils, and under the mouth; caudal rounded; pectoral so joined that it can not be turned forward as usual in this group, but rising

obliquely backward and upward. Eye slightly longer than snout; mouth brown within. Color brown, with white spots on the body and median fins, 6 of the largest of these each with the center yellowish, the largest from once to twice diameter of eye; spots on dorsal fins small; eye golden. Porto Rico. (Poey.) Not seen by us. The type 70 mm. long. (*inops*, helpless. "I call this species '*inops*' on account of the miserable fishing rod which has fallen to its lot." Poey.)

*Antennarius inops*, POEY, Anal. Soc. Esp. Hist. Nat., x, 1881, 340, Porto Rico. (Coll. Don Juan Gündlach.)

3096. ANTENNARIUS PRINCIPIS (Cuvier & Valenciennes).

D. III-11; A. 7; P. 10. Anterior dorsal spine twice as long as second, ending in a small, slender lobe; membrane behind third spine extending to root of soft dorsal; last ray of dorsal not reaching caudal. Skin rough, covered with small spines; no cutaneous fringes. Black; tips of pectorals and ventrals white; a small white spot above pectoral. (Günther.) West Indies to Brazil; not seen by us. Günther's specimen, above described, from Para. (*principis*, of the prince. Named for its discoverer, Prince Maurice of Nassau.)

*Chironectes principis*,\* CUVIER & VALENCIENNES, Hist. Nat. Poiss., xii, 416, 1837, Brazil; on 2 drawings by Prince MAURICE, the second representing the present species to which Günther restricts the name *principis*. The first figure is more like *tigris*.

*Antennarius principis*, GÜNTHER, Cat., iii, 193, 1861.

3097. ANTENNARIUS TENEBROSUS (Poey).

D. III-12; A. 7; P. 11. Anterior dorsal spine longer than second, terminating in a simple and slender tentacle; soft dorsal fin terminating at some distance from caudal. Skin rough, covered with small spines, without cutaneous fringes. Blackish brown, marbled with darker and lighter; a series of 3 black, blue-edged ocelli on upper posterior part of dorsal fin; many similar ocelli on caudal fin, irregularly disposed. (Poey.) Cuba. Not seen by us. (*tenebrosus*, dusky.)

*Chironectes tenebrosus*, POEY, Memorias, i, 219, pl. 17, fig. 1, 1851, Cuba.

*Antennarius tenebrosus*, GÜNTHER, Cat., iii, 197, 1861.

3098. ANTENNARIUS RETICULARIS, Gilbert.


D. III-12; A. 7. First dorsal spine short, very slender and filiform, not reaching tip of second, terminating in a short, fleshy flap; second spine moderately robust, flexible, not curved backward, wholly free and with-



\* Concerning this nominal species and *Chironectes mentzelii*, both of which were based on drawings by Prince Maurice, Cuvier & Valenciennes remark:

"We here cite these figures, and we give them specific names only to fix the attention of travelers and to get them to find the species which have served as models for these figures."





The following is the substance of the original description of *Chironectes principis*:  
In the first figure, color very deep brown, speckled with black spots on body and fins; dorsal fin with only 1 series of spots. Filament of first spine twice as long as that of the second, and terminating in a little knob or bait, the second spine free from the first and similar in shape.

Second figure of the same form, the second dorsal longer, the first ray ending in a spiral, and the whole body white, with 2 white round spots on each side, one above the other.





**IMAGE EVALUATION  
TEST TARGET (MT-3)**



**Photographic  
Sciences  
Corporation**

23 WEST MAIN STREET  
WEBSTER, N.Y. 14580  
(716) 872-4503

15  
18  
22  
25  
28  
32  
36  
40  
45  
50  
56  
63  
72  
80  
90  
100

10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100



out membrane; third spine nearly erect, not free, depressible with difficulty; not curved as in *A. sanguineus*; spines on the body rather coarse and shagreen-like, with expanded, undivided tips. Color in spirits, top of head, including dorsal spines and front of soft dorsal, coral red, the body otherwise light gray, broadly reticulated on sides and below with heavy black lines, which inclose 5 or 6 large pale spots; pectorals, ventrals, and anal with narrow terminal and wide median black bars. Soft dorsal uniformly light. This species closely resembles *A. sanguineus*, but differs from it in the straight, erect spinules, the color, and the character of the plates on the body. Length  $1\frac{1}{2}$  inches. Gulf of California. (Gilbert.) Only the type known. (*reticularis*, netted.)

*Antennarius reticularis*, GILBERT, Proc. U. S. Nat. Mus. 1891, 566, Gulf of California, at Albatross Station 2825. (Coll. Gilbert.)

#### 3099. ANTENNARIUS STRIGATUS, GILL.

D. III-12; A. 7. First dorsal spine elongate, filiform, twice length of second, with very slender, dermal tip; third spine more robust than second, wholly concealed in the skin, its length equal to that of first spine. Lips, maxillary, and a large transverse area behind second dorsal spine naked, each side of this area with a few spinous tubercles; skin elsewhere covered with fine shagreen-like armature. Color in spirits, olivaceous everywhere on body and on inside of mouth, finely mottled with light olive brown; many irregular blackish areas on head and body, those on lower side of head showing a tendency to form concentric bars; some on sides forming irregular bars downward from back; posterior portion of body not darker than the anterior; terminal parts of all the fins largely blackish, but with distinct black bars; some scattered round blotches on sides, each consisting of a number of smaller black spots on an olive ground; head and body with numerous pinkish and rose-red spots and bars, the latter sinuous, irregular, with wavy margins; a pinkish bar behind maxillary; a broad, saddle-like pinkish blotch across interval between second and third dorsal spines; a third bar from in front of origin of second dorsal downward toward base of pectorals; a fourth across top of caudal peduncle; first dorsal spine narrowly barred with brown. Pacific coast of tropical America, from Cape San Lucas to Panama. Here described from an adult, 10 inches in length, from Panama. This differs considerably from the descriptions of the young (*strigatus, tenuifilis*) given by Gill and Günther.

Two young individuals, types of *A. strigatus*, are thus characterized by Dr. Gill:

"The anterior dorsal spine is very slender and filiform, without appendages; the second is straight and moderate; the third concealed and developed as a hump, obtuse behind. The spines which cover the body are small and mostly bifid. The back and front of the dorsal fin are reddish; the rest light brown, with black stripes which diverge downward above the pectorals, those in front being parallel with the profile and at right angles with those behind; around the pectoral fins and on the flanks, the streaks are generally blended to form a continuous black

area; a black dorsal saddle is in front of the dorsal fin, and a black band covers the posterior half of the caudal fin; the abdomen is broadly reticulated with black, and the brown intervals themselves are frequently striated with the same color; the interior of the mouth is immaculate." (*strigatus*, striped.)

*Antennarius strigatus*, GILL, Proc. Ac. Nat. Sci. Phila. 1863, 92, Cape San Lucas (Coll. J. Xantus), young; JORDAN & GILBERT, Proc. U. S. Nat. Mus., 1882, 650, adult; JORDAN, Cat. Fishes, 138, 1885.

*Antennarius tenuifilis*, GÜNTHER, Fishes Centr. Amer., 440, 1869, Panama; young.

3100. ANTENNARIUS SANGUINEUS, Gill.

Anterior dorsal spine very slender,  $2\frac{1}{2}$  in length of caudal fin, terminating in a flap extended on each side, lacinated outward; second spine rough, robust, and curved strongly backward at its end; third not free, but apparent as a hump pointed backward, and extending  $\frac{1}{2}$  of the distance from its insertion to that of dorsal fin; skin covered with small bifid spines, whose prongs diverge considerably and are acute. Color blood red, except on abdomen, but with several more or less distinct black spots under origin of dorsal fin and on sides; abdomen light or yellowish brown, spotted with black; intervals between caudal and anal rays also marked with black; floor of mouth behind tongue with 2 lateral black bands converging toward the front, while the posterior margin of the tongue itself is also sometimes lined with black. (Gill.) Pacific coast of tropical America, Cape San Lucas to Panama; scarce. (*sanguineus*, bloody.)

*Antennarius sanguineus*, GILL, Proc. Ac. Nat. Sci. Phila. 1863, 91, Cape San Lucas (Coll. Xantus); JORDAN, Cat. Fishes, 138, 1885.

*Antennarius leopardinus*, GÜNTHER, Proc. Zool. Soc. London, 1864, 151, Panama.

3101. ANTENNARIUS OCELLATUS (Bloch & Schneider).

Depth  $1\frac{1}{2}$  in length. D. II-I-14; A. 8; P. 11; orbit equaling snout, eye much smaller; maxillary  $3\frac{1}{2}$  in body; pectoral rays  $5\frac{1}{2}$ ; caudal 4. Body short, oblong, compressed, very deep through occipital region; mouth large, subvertical; teeth small, sharp, cardiform, in wide bands on jaws, vomer, and palatines; maxillary extending downward to below axis of body; a very large knob at symphysis; lower part of head with many large, thick tentacles. First 2 dorsal spines on interorbital space, the first slender, terminating in a fringed lobe, the second shorter and much thicker, behind it a smooth depression; the third spine exceedingly rough and thick, blunt at tip and adnate to body; soft dorsal long and low, its origin in front of middle of body, tips of last rays reaching base of caudal; anal posterior, tips of its rays coterminous with dorsal rays, its height equal to its length; pectorals near middle of body, placed far below axis; ventrals short, the rays thickened, their position under posterior edge of eye. Skin covered with minute bifurcate spines, running upon dorsal, anal, and caudal rays; gill opening in front and below pectoral, its length about equal to snout. Color brown, marbled with

lighter, and with scattered black dots, especially on belly and outer portions of dorsal and caudal; each side with 3 large black spots ocellated with brownish, 1 on dorsal near its base, a second immediately below it on the sides, and a third in the middle of the caudal fin; mouth behind tongue black, with yellow lines. West Indies, north to Florida. Common in the West Indies; the most abundant of the American species. Here described from a specimen from off Pensacola, Florida, about 15 inches in length.

Mr. Garman gives the following note on *Antennarius ocellatus*:

"The species was tolerably figured by Parra, but has not been recognized by some of the subsequent writers. On 5 specimens before me the amount of variations in markings is comparatively small. The 3 large ocelli, on dorsal, caudal, and middle of side, are present on each, as is also the case with the numerous small spots of black on the ventral portions of the body and on the outer portions of dorsal and caudal. The dorsal ocellus lies between the sixth and seventh rays, on the middle of the fin; that on the flank is situated on the vent, and that on the caudal between the fourth and fifth rays, from the top, near the middle of the fin. The black portion of either of these spots is larger than the orbit, which latter is rather small when contrasted with that of other species. The white circle around the black, again, is surrounded by a narrow one of brown. On the caudal, at each side of the ocellus, there are transverse streaks. The first ray of the dorsal is as long as the second, and is covered by scales. The bulb apparently is simple, and bears numerous laciniæ. The second dorsal spine is shorter than the third; both are club shaped. The space behind the second dorsal spine is covered by scales." (*ocellatus*, with eye-like spots.)

*Pescador*, PARRA, Dif. Piezas, Hist. Nat., 1, tab. 1, 1780, Cuba.

*Lophius histrio*, var. *ocellatus*, BLOCH & SCHNEIDER, Syst. Ichth., 142, 1801; after PARRA.

*Antennarius pleurophthalmus*, GILL, Proc. Ac. Nat. Sci. Phila. 1863, 92, Key West; JORDAN & GILBERT, Synopsis, 846, 1883; JORDAN, Cat. Fishes, 138, 1885; GOODE & BEAN, Oceanic Ichthyology, 487, 1893.

*Antennarius ocellatus*, POEY, Synopsis, 105, 1868; GARMAN, Bull. Iowa Lab. Nat. Hist. 1896, 82.

### 3102. ANTENNARIUS SCABER (Cuvier).

D. III-12; A. 7; P. 9 or 10. Anterior dorsal spine as long as second, and provided with 2 long and thick cutaneous flaps at its tip; third dorsal spine not continuous with the soft dorsal; soft dorsal fin terminating at some distance from the caudal, its last ray not extending to root of caudal, if laid backward; dorsal spines, head, back, and sides of the body with more or less numerous cutaneous fringes, those of dorsal spines sometimes forming a dense cluster; skin very rough, covered with small spines. Ground color yellowish or reddish, with numerous brown spots, those around the eye forming radiating streaks; dorsal and anal fins with 3 series of round brown spots, the middle of which is formed by the largest and most constant spots; sometimes uniform brown. Caribbean Sea. (Günther.) A small specimen from Port Castries, St. Lucia, has the body

light brown, clouded with darker, fins all with round black spots, those of the base of the dorsal somewhat larger than others; ventrals tipped with black. (*scaber*, rough.)

*Chironectes scaber*, CUVIER, Mém. Mus., III, 425, pl. 16, fig. 2, 1817, Martinique (Coll. Plée); CUVIER & VALENCIENNES, Hist. Nat. Poiss., XII 412, 1837.

*Lophius spectrum*, GRONOW, Cat. Fish., Ed. Gray, 49, 1854, Antilles; after *Lophius acute scabra*, GRONOW, Zoophyl., 210, 1781.

*Antennarius scaber*, JORDAN, Proc. U. S. Nat. Mus. 1889, 652, specimen from Port Castries, St. Lucia.

*Antennarius histrio*, GÜNTHER, Cat., IV, 188, 1861; not *Lophius histrio*, LINNÆUS.

### 3103. ANTENNARIUS TIGRIS, Poey.

D. III-12; A. 7; P. 11. Anterior dorsal spine longer than second, terminating in 2 long cutaneous flaps; third dorsal spine connected with soft dorsal by a broad membrane, the latter terminating at some distance from the caudal, and its last ray not extending to root of caudal if laid backward. Skin rough, covered with small spines, without cutaneous fringes. Ground color yellow, with numerous brown spots and streaks, the latter radiating from the eye; dorsal fin irregularly spotted, with a series of large round brown spots. (Poey.) Cuba. Not seen by us. According to Mr. Garman, *Antennarius scaber* and *A. tigris* "are closely allied, but if placed side by side the squamation and filaments suffice to distinguish them, great similarity in color notwithstanding. *A. scaber* has coarser scales, with shorter, rougher spines, the scales are farther apart, and the cutaneous flaps appear on the body much as figured by Cuvier. On *A. tigris* there are few of the cutaneous appendages, the scales are closer together, the spines are longer and more slender, giving rise to an appearance more like velvet, and the head and body are more compressed." (*tigris*, tiger.)

*Chironectes tigris*, POEY, Memorias, I, 217, pl. 17, fig. 2, 1851, Cuba.

*Antennarius tigris*, GARMAN, Bull. Iowa Lab. Nat. Hist. 1896, 83.

### 3104. ANTENNARIUS NUTTINGII, Garman.

D. 2+12; A. 7; V. 5; P. 11; C. 9. In form this species is shorter, more massive anteriorly, and less compressed than either *A. ocellatus* or *A. radiosus*. A transverse section across the middle of the body is a nearly equilateral triangle. Caudal region short. Head nearly as wide as high; cheeks swollen; forehead rather broad, converging forward on the edges. Occipital concavity wide and deep, free from scales in a wide space below the ends of the first and second dorsal rays, this bare space being apparently for the reception of the fleshy bait bulb, which latter has 2 elongate lobes. Snout as long as the orbit, broad, truncate; chin vertical; symphyseal knob prominent. Mouth wide, subvertical. Eye small; orbit twice as long, hardly more than  $\frac{1}{2}$  the interorbital space. First and second dorsal rays equal in length, not inclusive of the 2 elongate fleshy fringed lobes surmounting the first. The base of the first ray stands forward prominently over the mouth, being free for some distance. The

greater portion of the second ray is free, while the third is connected with the dorsum, by the skin, from base nearly to tip. This last ray is larger than either of its fellows. Soft dorsal large; middle rays longest, as long as the distance from the maxillary to the hind edge of the operculum, or as long as the rays of the caudal fin; fin not reaching back to the bases of the caudal rays, fringed. Hind margin of caudal convex, fringed. Anal moderate, rays prominent in the margin, fin with a blunt angle on the outer edge, subtending, when laid up against the tail,  $\frac{1}{2}$  or more of the length of the caudal rays. The rays on the pectoral fins extend out beyond the margins more noticeably than those of the other fins. Ventrals small, in most instances with 6 points on the outer margin, in one case having but 5. Greatest length of the caudal nearly  $\frac{1}{2}$  of the total length. Length of each maxillary  $\frac{2}{3}$  of the caudal. Scales short, small, close set, harsh to the touch, having none of the velvety appearance. Uniform black; inside of the mouth black; "bait" white. Great Bahama Banks. Besides the specimens in Nutting's collection there are several others in that of the Mus. Com. Zool. "This species is readily separated from *A. principis* of authors by the short first dorsal spine." ("The specific name is given in honor of Prof. C. C. Nutting, to whom science is so much indebted for the origination and successful accomplishment of the expedition.")

*Antennarius nuttingii*, GARMAN, Bull. Iowa Lab. Nat. Hist. 1896, 83, pl. II, Great Bahama Banks. (Coll. C. C. Nutting.)

??*Chironectes mentzelii*,\* CUVIER & VALENCIENNES, Hist. Nat. Poiss., XII, 417, Brazil; on a drawing by Prince MAURICE.

### 3105. ANTENNARIUS MULTIOCELLATUS (Cuvier & Valenciennes).

(MARTIN PESCADOR.)

Mouth large, vertical. First dorsal spine slender and straight, nearly equal to length of caudal, terminating in 3 simple tentacles; second dorsal spine curved at the middle and extending to the base of the third; third dorsal spine partly embedded in the skin, reaching halfway to dorsal; wrists and pectorals widened; ventrals short. Skin covered with bifid spines. Fawn color, lighter below; many black spots ocellated with white, both on the body and fins; body with several pink areas, 1 of which forms a triangular saddle in front of the dorsal and another a broad ring around the base of the caudal fin; angles of mouth with a pink spot. West Indies, north to Florida Keys; common. This description (after Gill) from the type of *Antennarius annulatus* from Garden Key. According to Mr. Garman, "this species is distinguished by the trifid bulb and the long first dorsal ray, near twice as long as the second and quite as long as the caudal, by the high nape, by the large third dorsal ray, much larger and more swollen than the second, and by the coloration. The eye is very small. The black centers of the largest of the ocelli are smaller than the eye. Besides the ocellus on the soft dorsal that on the anal and the 3

\* The following is the substance of the very brief description of this nominal species: "First dorsal filament not longer than the second, and ending in a small bait or knob. Body black, with large marblings."

forming a triangle on the caudal, there are others scattered over the caudal and other fins, and over the sides of the body. Below the eye on the cheek and under the chin and the chest the spots are little more than black dots. Over the sides, a specimen in hand, the type of *A. corallinus*, Poey, is freckled with lighter rounded spots. Behind the pectoral, on the side, there is a small ocellus with a black center. On each side in the same position, a short distance above the pectoral, there is a brown ocellus larger than the orbit, in the center of which there is a white dot. A brown streak passes back from the upper part of the orbit and curves down toward the anal ocellus, another passes back from the middle of the eye and curves down toward the pectoral, and a third below the third dorsal spine runs down and then forward toward the lower end of the maxillary. The forehead is comparatively narrow; behind the second dorsal ray the bare space is hardly large enough to receive the bait."

Poey thus describes *Antennarius corallinus*, which according to Garman, is the same as *A. multiocellatus*:

"D. II-1, 12; A. 7; V. I, 5; C. 17. Two dorsal spines in front of eye, formed like horns, another higher on the nape; gill opening spiral, at the lower base of the pectoral; general form of the fish almost globular; the mouth vertical; tongue marbled with black and white; caudal rounded; eyes very small; pectoral low, reaching middle of body without caudal; ventrals short; vent near anal; first dorsal spine ending in a single short filament. Color reddish with black spots; 2 of these spots eye-like, with a larger black center and the iris of the color of the body, surrounded with a black circle; 1 spot at the base of the soft dorsal at the second third of its length, a very weak one at base of anal; between first spot and pectoral fin 3 small inconspicuous ocellate spots, of which the pupil is a small point; middle of the body with dusky spots; dorsal, anal, and caudal with black points; tubercles about eye and on the cheek, but not spinous. Type 95 mm. long. Cuba." (Poey.) (*multus*, many; *ocellus*, an eye-like spot.)

*Chironectes multiocellatus*, CUVIER & VALENCIENNES, Hist. Nat. Poiss., XI, 422, 1837, Martinique. (Coll. M. Garnot.)

*Antennarius annulatus*, GILL, Proc. Ac. Nat. Sci. Phila. 1863, 91, Garden Key, Florida (Coll. Llent. Wright); JORDAN & GILBERT, Synopsis, 846, 1883.

*Antennarius corallinus*, POEY, Repertorio, 1, 188, 1865, Cuba. (Coll. Poey.)

*Antennarius multiocellatus*, GÜNTHER, Cat., III, 194, 1861; GARMAN, Bull. Iowa Lab. Nat. Hist. 1896, 82.

#### 3106. ANTENNARIUS RADIOSUS, Garman.

D. 3+13; A. 8; V. 5; P. 11; C. 9. Resembling *A. tigris*, Poey, in shape, squamation, etc., but differing in coloration and in possession of a much longer dorsal ray. The staff in this ray is very slender, much longer than the second ray, and bears a small, trifid "bait." Second and third dorsal rays shorter than the first, the third well tied down by the skin. Scales uniform, sharp. No cutaneous fringes on large specimens. Grayish or brownish white, darker on nape and dorsal fin, with numerous spots of light color, as large as the orbit, surrounded by more or less complete

edgings of brown, producing a semblance to reticulation, or to spottings by drops of liquid; 7 streaks of brown radiate from the eye; as in *A. tigris*, they are continued upon the head and down toward the ventrals; a large spot of black, white-edged, a little larger than the orbit,  $\frac{1}{2}$  on the fin and  $\frac{1}{2}$  on the muscles of the body, occupies the space between the eighth and the tenth rays of the soft dorsal fin; the light areas vary in intensity, and lie close together over nearly the whole of body and fins; belly lighter, with faint indications of lines of brownish, radiating from the head; caudal with oblique transverse cloudings of brownish; hindmost  $\frac{1}{2}$  light. The color in life was probably reddish or yellowish. Secured off Key West, in about 50 fathoms. A young individual, of less than an inch, taken opposite Havana, is of lighter gray, and has a large ocellus, of light color in the center, between the black one at the base of the dorsal and the upper end of the humerus. There are small cutaneous fringes on the flanks. (Garman.) (*radiosus*, rayed.)

*Antennarius radiosus*, GARMAN, Bull. Lab. Nat. Hist. Iowa Univ. 1896, 85, pl. 1, off Key West, in 50 fathoms. (Coll. C. C. Nutting.)

#### 1062. CHAUNAX, Lowe.

*Chaunax*, LOWE, Trans. Zool. Soc. Lond., III, 1846, 339 (*pictus*).

Head very large, depressed, cuboid. Mouth large, subvertical; jaws and palate with bands of small teeth. Skin with small, sharp spines. Spinous dorsal reduced to a small tentacle above the snout, retractile into a groove; soft dorsal moderate, low; anal short; ventrals small. Gills  $2\frac{1}{2}$ ; no pseudobranchiae. Muciferous channels very conspicuous, the lateral line prominent, undulate; another series of mucous tubes extending from lower jaws to axil; still another extends backward from snout and maxillary to a point behind eye, when it ceases, uniting with a vertical line which extends from the lateral line to the lower line; these lines thus inclose a quadrate area on the cheek. Gill opening small, well behind pectoral under front of soft dorsal. Deep seas. (*χαύραξ*, one who gapes.)

a. Dorsal rays 11; anal 5; depth  $2\frac{1}{2}$  in length.

PICTUS, 3107.

aa. Dorsal rays 13; anal 7; depth  $2\frac{3}{4}$  in length.

NUTTINGII, 3108.

#### 3107. CHAUNAX PICTUS, Lowe.

Head  $1\frac{1}{2}$ ; depth  $2\frac{1}{2}$ . D. I, 11; A. 5; P. 11; V. 4; C. 7. Rostral tentacle short, pedicellate; muciferous channels appearing as chain-like rows of pits. Bright orange above; sides rosy; fins vermilion. Deep waters of the Atlantic; recorded from Madeiro, Soudan, Cape Verdes, Barbados, off Rhode Island, and elsewhere in the Gulf Stream, in 130 to 428 fathoms. A similar species (*Chaunax fimbriatus*), regarded by Günther as the same, occurs in the Japan Seas, Bay of Bengal and the Fiji Islands. (*pictus*, painted.)

*Chaunax pictus*, LOWE, Trans. Zool. Soc. Lond. 1846, 339, Camera de Lobos, Madeira; GÜNTHER, Cat., III, 200, 1861; GOODE, Proc. U. S. Nat. Mus. 1880, 470; JORDAN & GILBERT, Synopsis, 847, 1883; GOODE & BEAN, Oceanic Ichthyology, 487, fig. 398, 1896.

? *Chaunax fimbriatus*, HILGENDORF, Sitzber. Ges. Naturf. Freunde 1879, 80, Sea of Japan.

? *Chaunax nuttingii*, GARMAN, Bull. Lab. Nat. Hist. Iowa Univ. 1896, 85, pl. III, fig. 2, near Sand Key Light, Florida, in 120 fathoms. (Coll. C. C. Nutting.)

## 3108. CHAUNAX NUTTINGII, Garman.

B. 6. D. II, 13; A. 7; V. 4; P. 14; C. 9. Form resembles that of *Chaunax pictus*, but is shorter, broader, and possessed of more fin rays. Anteriorly it is broad and depressed, posteriorly compressed. From head to soft dorsal on the nape it is arched very little. Head broader than high, flattened or slightly concave on the occiput, nearly vertical on the chin. Snout short, broad, truncate. Eye medium, the length of the scaleless area covering it equals the width of that between the canals on the inter-orbital space, or about  $\frac{2}{3}$  of the space itself; the distance from the maxillary is about the ocular width. The niche in which the first dorsal spine is received is subelliptical and about  $\frac{2}{3}$  as long as the eye; the tentacle is little more than  $\frac{1}{2}$  as long as the niche, is broad near the base, tapers rapidly and bears a 2-lobed "bait" with slender fringes. Mouth wide, oblique, maxillary about 3 times as long as the eye, widened and rounded at the outer end; intermaxillaries alone forming upper border of mouth. Teeth small, slender, sharp, in villiform bands. Origin of soft dorsal in the middle of the distance from the rostral tentacle to the base of the caudal fin, fourth ray above the gill opening, anterior rays shorter. Vent below the seventh ray of the second dorsal. Pectorals short, broad, rounded. The canals of the lateral system are in the main like those of *C. pictus*, but have stronger curves; they begin to curve outward immediately behind the niche, not remaining parallel or converging as in Lowe's species. Scales very fine, sharp and close together. In life this fish was probably red or yellowish with transverse cloudings or blotches of brownish, it is now dingy brownish white; one of the blotches lies just behind the eye, another lies below the orbit, and apparently 3 transverse bands cross the back through the soft dorsal; orbit blackish; tentacular niche black. The coloration of the individual described indicates a habitat within reach of the effects of sunlight. Florida Keys; the type dredged nearly 8 miles south of Sand Key Light, Florida, in about 120 fathoms. (Garman.) This species is evidently not very different from *Chaunax pictus* and may be the same. (Named for C. C. Nutting, professor of zoology in the University of Iowa, director of the Bahama Expedition of 1893.)

*Chaunax nuttingii*, GARMAN, Bull. Lab. Nat. Hist. Iowa Univ. 1896. 86, pl. III, f. 2, Sand Key Light, Florida. (Coll. C. C. Nutting.)

## Family CCXXIII. CERATIIDÆ.

## (THE SEA DEVILS.)

Head and body compressed. Mouth terminal, more or less oblique. Gill openings small, in the lower part of the axils. No pseudobranchiæ. Spinous dorsal represented by 1 or more tentacles. Pectoral members not geniculated, with short pseudobranchia and 3 actinosts. No ventral fins. Fishes of the open seas, usually inhabiting considerable depths; 13 genera and 15 species known. All are uniform blackish in color.

"The bathybial sea devils," writes Günther, "are degraded forms of *Lophius*; they descend to the greatest depths of the ocean. Their bones



are of an extremely light and thin texture, and frequently other parts of their organization, their integuments, muscles, and intestines are equally loose in texture when the specimens are brought to the surface. In their habits they probably do not differ in any degree from their surface representative, *Lophius*. The number of the dorsal spines is always reduced, and at the end of the series of these species only 1 spine remains, with a simple, very small lamella at the extremity (*Melanocetus johnsonii*, *Melanocetus murrayi*). In other forms sometimes a second cephalic spine, sometimes a spine on the back of the trunk, is preserved. The first cephalic spine always retains the original function of a lure for other marine creatures, but to render it more effective a special luminous organ is sometimes developed in connection with the filaments with which its extremity is provided (*Ceratias bispinosus*, *Oncirodes eschrichtii*). So far as it is known at present these complicated tentacles attain to the highest degree of development in *Himantolophus* and *Egwonichthys*. In other species very peculiar dermal appendages are developed, either accompanying the spine on the back or replacing it. They may be paired or form a group of 3, are pear-shaped, covered with common skin, and perforated at the top, a delicate tentacle sometimes issuing from the foramen." (*Pediculati*, genus *Ceratias*, Günther, Cat., III, 205, 1861; *Ceratiidae*, Gill, Proc. U. S. Nat. Mus. 1878, 216.)

a. Mouth moderate.

b. Gills in  $2\frac{1}{2}$  pairs.

CERATIINÆ:

c. Cleft of mouth nearly vertical; skin prickly.

d. Cephalic spine single.

e. Dorsal spine present; lateral caruncles present; no teeth on vomer.

CERATIAS, 1063.

ee. Dorsal spine wanting; caruncles present.

f. Caruncles remote from soft dorsal.

MANCALIAS, 1064.

ff. Caruncles close to soft dorsal.

CRYPTOPSARAS, 1065.

ONEIRODINÆ:

cc. Cleft of mouth nearly horizontal; skin smooth; 1 cephalic spine and 1 postcephalic spine.

ONEIRODES, 1066.

HIMANTOLOPHINÆ:

bb. Gills in  $\frac{1}{2}$   $2\frac{1}{2}$  pairs; body with scattered tubercular scutella; no second dorsal spine.

g. Body and head compressed; mouth oblique; joint of mandible below or behind eye; eye rudimentary.

h. Body oblong, oval; dorsal rays 9; pectoral 12.

HIMANTOLOPHUS, 1067.

hh. Body short and deep; dorsal rays 4; pectoral about 17.

CORYNOLOPHUS, 1068.

aa. Mouth with enormous gape.

MELANOCETINÆ:

i. Cleft of mouth nearly vertical; pectoral small, in advance of dorsal and of gill opening; second dorsal spine wanting; gills in  $2\frac{1}{2}$  pairs.

j. Gular tentacle wanting; no teeth on vomer.

LIOCETUS, 1069.

jj. Gular tentacle present; 1 tooth on the vomer.

LINOPHRYNE, 1070.

CAULOPHRYNINÆ:

ii. Cleft of mouth nearly horizontal; pectorals below dorsal and behind gill opening; gills in  $\frac{1}{2}$   $2\frac{1}{2}$  pairs.

k. Dorsal and anal greatly produced; skin naked; head and body with many luminous filaments.

CAULOPHRYNE, 1071.

1063. CERATIAS, Kröyer.

*Ceratias*, KRÖYER, Naturhist. Tidsskrift. 2 Række, 1, 639, 1844 (*holbølli*).

Head and body much compressed and elevated, oblong, covered with prickly skin. Mouth wide, its cleft nearly vertical; teeth in jaws conic, movable, of moderate size; no teeth on vomer or palatines. Gills 24; gill arches unarmed. Spinous dorsal reduced to 2 spines, 1 on the head, the other on the back, the basal element of the second spine exerted; the cephalic spine much elongate; soft dorsal and anal short; pectorals very short, broad, of about 20 rays. Caudal fin much produced, fan-shaped, with exerted rays. Pyloric caeca 2, small. Skeleton soft, fibrous. Greenland. (*κερατίας*, one that has horns.)

3109. CERATIAS HOLBØLLI, Kröyer.

D. I-I, 4; A. 4; P. 19; C. 10. Head 2½; depth nearly 2; head deeper than long; eyes small, not more than ¼ the length of the head; free rays of the head a little shorter than to the base of the caudal fin; the forked part of the caudal fin shorter than the length of the fish; the length of the pectoral fin equals almost ⅔ the entire length of the fish, the membranes from the dorsal and anal fins posteriorly extend almost to the base of the caudal fin. Color entirely black. (Kröyer.) North Atlantic; 4 specimens known; 3 from Greenland, 1 from Nova Scotia. (Named for C. Holbøll, a Danish naturalist.)

*Ceratias holbølli*, KRÖYER, Naturh. Tidsskr. 1844, 639, Greenland; GÜNTHER, Cat., III, 205, 1861; JORDAN & GILBERT, Synopsis, 847, 1883; GOODE & BEAN, Oceanic Ichthyology, 489, pl. 117, fig. 394, 1896.

1064. MANCALIAS, Gill.

*Mancalias*, GILL, Proc. U. S. Nat. Mus., 1, 1878, 227 (*uranoscopus*).

*Typhlopsaras*, GILL, Forest and Stream, New York, 1883, Nov. 8, 284 (*shufeldti*).

General characters of *Ceratias*, but with the spinous dorsal reduced to a single rostral spine, and 2 fleshy claviform tubercles or caruncles behind it. Pectoral fins narrow, with 10 to 15 slender rays. (*mancus*, defective, "with a quasi-diminutive termination to correspond with *Ceratias*.")

a. Dorsal caruncles placed before dorsal fin a distance 6 times in length of trunk from gill opening to base of caudal. URANOSCOPUS, 3110.

aa. Dorsal caruncles placed nearer dorsal, the distance from dorsal 4½ times in trunk, as above. SHUFELDTI, 3111.

3110. MANCALIAS URANOSCOPUS (Murray).

D. I, 3 or 4; A. 4; C. 8; P. 10. Anterior spine of first dorsal produced in a long filament, ending in a pear-shaped bulb, terminating in a semitransparent whitish spot, this spine originating on posterior part of head, and reaching, when depressed, nearly to the tip of tail; far behind this are 2 short, fleshy tubercles, lying in a depression in front of second dorsal. Teeth moderate, depressible. Skin everywhere with minute embedded conical spines. Eye very small, placed high on the middle of the head.

Color uniform black. (Murray.) To this description Goode & Bean add from the same specimen: "Anal opposite second dorsal, the 4 median caudal rays being much larger than the others and bifid; pectorals small, above the gill opening; the upper jaw is formed by the intermaxillaries, and is armed, together with the lower jaw, with a series of teeth of moderate size, which can be depressed as in *Lophius*. The skin is thickly covered with minute embedded conical spines; the eyes are very small and are placed high upon the middle of the head." Mid-Atlantic, in very deep water; 2 specimens known, the type in 2,400 fathoms, taken between the Canary and Cape Verde islands, the second (26159) in 372 fathoms off the coast of Rhode Island in the Gulf Stream. (*ὄρανοσόκος*, star-gazing, from the upturned eyes.)

*Ceratas uranoscopus*, MURRAY, in Wyville Thompson, *The Atlantic*, II, 67, fig. 20, 1878, mid-Atlantic, between Canary and Cape Verde Islands in 2,400 fathoms; GÜNTHER, *Challenger Report*, XXII, 54, pl. 11, fig. C, 1887.

*Mancalias uranoscopus*, GILL, *Proc. U. S. Nat. Mus.* 1878, 228; GOODE, *Proc. U. S. Nat. Mus.* 1880, 469; JORDAN & GILBERT, *Synopsis*, 848, 1883; GOODE & BEAN, *Oceanic Ichthyology*, 490, 1896.

#### 3111. MANCALIAS SHUFELDTI (Gill).

Maxillary  $\frac{1}{2}$  the length from gill opening to caudal base; intermaxillary  $3\frac{1}{2}$  times in this length. Form more slender than that figured by Günther, with 4 rays in the dorsal, and apparently 15 in pectoral. There are no vomerines; intermaxillary and mandible armed with a narrow band of depressible teeth of various lengths. The skin with a fine granular appearance and everywhere covered with minute prickles. The caruncles only 2 in number and situated as in *Mancalias uranoscopus*, as figured in the *Challenger* fishes. Length of dorsal spine, without the joint bearing the pear-shaped appendage, equaling distance from gill opening to root of tail; the joint bearing the appendage is  $\frac{2}{3}$  of this distance; in *Mancalias uranoscopus* (No. 26159) the first dorsal, without the joint bearing the appendage, contains the distance from the gill opening to the root of the tail  $1\frac{1}{2}$  times. The joint containing the appendage is  $\frac{1}{2}$  as long as the distance from the gill opening to root of tail. Dermal caruncles distant from the dorsal a space equal to  $\frac{1}{2}$  of distance from the gill opening to root of tail. In *M. shufeldti* the caruncles are placed at a distance from the dorsal a space contained  $4\frac{1}{2}$  times in the distance from the gill opening to the root of the tail. In the specimens described by Goode & Bean as *Mancalias uranoscopus* (No. 26159), the length  $3\frac{1}{2}$  inches, the length of the maxillary is  $\frac{1}{2}$  of length from gill opening to root of tail, and the intermaxillary  $3\frac{1}{2}$  times in same distance. Teeth in jaws depressible, in narrow bands, and of unequal size; vomer toothless. Two small caruncles not far from front of dorsal fin, and instead of being placed opposite each other, according to the usual arrangement, one is placed behind the other. Skin covered with minute granules or papillae, each one surmounted by a slender prickle, as in *Typhlopsaras*. The pectoral of the individual described contains 15 rays. The pectorals of *T. shufeldti* are imperfect. (Goode & Bean.) Gulf Stream, off the coast of southern New England;

1 specimen known. (Named for Dr. Robert W. Shufeldt, United States Army, the well-known ornithotomist.)

*Typhlopsaras shufeldti*,\* GILL, Forest and Stream, Nov. 8, 1883, Western Atlantic (Type, No. 33552); JORDAN, Cat. Fishes, 138, 1885.

*Ceratius shufeldti*, GÜNTHER, Challenger Report, XXII, Deep-Sea Fishes, 54, 1887.

*Mancalia shufeldti*, GOODE & BEAN, Oceanic Ichthyology, 490, fig. 401, 1896.

### 1065. CRYPTOPSARAS, Gill.

*Cryptopsaras*, GILL, Forest and Stream, Nov. 8, 1883, 284 (*coesii*).

Body shortened; back longitudinally convex, eyes small but conspicuous; anterior spine with concealed basal joint and elongated terminal joint; a large intermediate globular and a pair of subpedunculated lateral dorsal appendages or caruncles close to the front of the dorsal fin; pectorals well developed, of about 15 rays. Deep-seas. (*κρυπτός*, concealed; modern Greek *ψαράς*, fisherman, in reference to the concealed rod bearing the dorsal spine or fishing apparatus.)

### 3112. CRYPTOPSARAS COESII, GILL.

The basal joint of the rod-like spine is almost entirely concealed and procumbent, and the distal joint alone free, reaching backward to the dorsal tubercle; the bulb is pyriform, and surmounted by a long whitish filament; dorsal and anal each with 4 spines, the caudal 8 (the 4 middle dichotomous) and the pectorals each about 15 rays. (Gill.) A specimen of *Cryptopsaras* (No. 33558, U. S. Nat. Mus.) was obtained, by the *Albatross*, from Station 2101, in Lat. 38° 18' 30" N., Long. 68° 24' W., at a depth of 1,686 fathoms. The type of *Cryptopsaras coesii* is only 35 mm. long. The caudal is imperfect. The length without caudal is 30 mm. and contains the greatest height 2½ times. The bulb on the dorsal spine when laid backward can be made to reach to the dermal caruncles on the back. The length of the upper jaw is about ¼ of the length without caudal; gill opening nearly midway between front of head and root of tail; mouth placed vertically; intermaxillary teeth occupying about entire length of bone; mandibular teeth unequal in size; at symphysis of mandible a pair of minute spines closely connected at base and slightly separated at the extremity. Specimen No. 39183 is 58 mm. long; 47 mm. to base of caudal. Greatest height 2½ in length without caudal. Gill opening a little nearer end of caudal than to front of head; distal portion of dorsal spine about ½ length without caudal; median dermal caruncle very much

\* The following is the original account of *Typhlopsaras*:

"*Typhlopsaras*.—Ceratiines with an elongated trunk, rectilinear back, obsolete or no eyes, far exerted basal joint of the anterior spine and shortened terminal joint, a small intermediate and a pair of pedunculated dorsal appendages some distance in advance of the dorsal fin, and reduced pectoral fin with about 5 or 6 rays.

"*Typhlopsaras shufeldti*.—The first joint of the rod-like spine reaches to the axil of the dorsal fin, and the bulb to the base of the caudal fin, when the spine is bent backward; the bulb is pear-shaped and without any appendages; the dorsal has 4 rays, the anal 4, the caudal 8 (the median 4 of which are forked), and there are 4 or 5 pectoral rays. A single specimen was found. I have dedicated the species to my esteemed friend, Dr. R. W. Shufeldt, U. S. A., the well-known ornithotomist.

"The name, *Typhlopsaras*, is a compound from the Greek τυφλος (blind) and ψαράς (angler), meaning 'blind angler.'"

larger than the two lateral ones; skin covered with minute granules of uniform size; pectoral with 16 rays, its length about  $\frac{1}{2}$  that of head; length of upper jaw about  $\frac{1}{2}$  of total without caudal; pair of spines at symphysis of mandible replaced by a very small knob; teeth in intermaxillary very small, diminishing in number toward the symphysis, apparently uniserial. On each side of head of vomer 2 or 3 depressible teeth; palatines apparently wanting. We have seen something like traces of similar teeth on the vomer of *Mancalias uranoscopus*, but owing to the condition of the specimen can not be certain about this character. (Goode & Bean.) Gulf Stream, off the coast of New England. (Named for the eminent ornithologist, Dr. Elliott Coues.)

*Cryptoparas couesii*, GILL, Forest and Stream, Nov. 8, 1883, 284, Gulf Stream off New England (Coll. Albatross); JORDAN, Cat Fishes, 138, 1885; GOODE & BEAN, Oceanic Ichthyology, 491, fig. 402, 1896.

*Cerattias couesii*, GÜNTHER, Challenger Report, xxii, 55, 1887.

*Cerattias carunculatus*, GÜNTHER, Challenger Report, xxii, 55, pl. 11, fig. d, 1887, south of Yezo, Japan, in 345 fathoms;  $1\frac{1}{2}$  inches long. (Coll. Challenger.)

#### 1066. ONEIRODES, Lütken.

*Oneirodes*, LÜTKEN, Overs. Dansk. Vidensk. Selsk. Forhandl. 1871, 56 (*eschrichtii*).

Body compressed, oval, short, covered with smooth skin. Head compressed, very large. Mouth moderate, almost horizontal, the joint of mandible behind eyes. Teeth unequal, depressible; vomer with teeth. Gill arches unarmed; gills in  $2\frac{1}{2}$  pairs. Spinous dorsal represented by a cephalic spine, the basal element of which is procumbent and subcutaneous, the tip bulbous, and a second spine about midway between the rostral spine and the soft dorsal; soft dorsal and anal short; no ventrals; no pyloric caeca. Greenland. (*ὄνειρώδης*, dream-like, in illusion to the small, almost covered, eyes.)

#### 3118. ONEIRODES ESCHRICHTII, Lütken.

D. I-I, 6; A. 8; C. 8. Terminal half of the bulb of the cephalic spine whitish. Cephalic spine with a bulbous termination, surmounted by slender filaments in several transverse rows. Caudal fin shorter than trunk without head. Color black. Deep sea, off Greenland. Known from a single specimen 8 inches long. (Gill.) (Named for D. F. Eschricht, a Danish naturalist, a student of the Cetacea.)

*Oneirodes eschrichtii*, LÜTKEN, Overs. Dansk. Vidensk. Selsk. Forhandl. 1871, 56, 9-18, pl. 2, deep sea off Greenland; GILL, Proc. U. S. Nat. Mus. 1878, 218; JORDAN & GILBERT, Synopsis, 848, 1883; GOODE & BEAN, Oceanic Ichthyology, 492, 1896.

#### 1067. HIMANTOLOPHUS, Reinhardt.

*Himantolophus*, REINHARDT, Dansk. Vidensk. Selsk. Nat. 1837, 74 (*greenlandicus*).

Head large, compressed. Skin thick, with scattered, round, prickly scales. Body oval, compressed. Mouth moderate, the cleft oblique, the joint of the mandible below or behind the eyes. Gills in  $\frac{1}{2}$   $2\frac{1}{2}$  pairs; gill arches armed with dentigerous tubercles. Spinous dorsal represented

only by a single long rostral spine, the basal element of which is procumbent and subcutaneous; the extremity with numerous long filaments. Soft dorsal short, with 9 rays; anal short; pectoral rather broad, with 12 rays. Greenland. (*ιμάς*, a thong; *λόφος*, crest.)

3114. HIMANTOLOPHUS GRÆNLANDICUS, Reinhardt.

Depth  $2\frac{1}{2}$  in total length. D. I-9; P. 12. Body oblong oval. Cephalic ray provided with about 11 tentacles. (Gill.) Greenland. "This species has never been fully described, the only existing example being an imperfect one, 23 inches long, obtained off the coast of Greenland, about 1837."

*Himantolophus grænlændicus*, REINHARDT, Dansk. Vid. Selsk. Nat. Math. Afh. 1837, 74, Greenland; GILL, Proc. U. S. Nat. Mus. 1878, 218; JORDAN & GILBERT, Synopsis, 849, 1883.

1068. CORYNOLOPHUS, Gill.

*Corynolophus*, GILL, Proc. U. S. Nat. Mus. 1878, 219 (*reinhardtii*).

This genus is scarcely distinct\* from *Himantolophus*, differing in the short oval form, the short dorsal of about 5 rays, and the broader pectoral with about 17 rays each. (*κορύνη*, club; *λόφος*, crest.)

3115. CORYNOLOPHUS REINHARDTI (Lütken).

Depth  $1\frac{1}{2}$ . D. I-5; P. 17. Body short, oval; cephalic ray  $\frac{2}{3}$  length of head, with about 8 tentacles, which branch out forming a brush at tip; skin sparsely covered with thorn-like prickles. Greenland. One specimen known, 14 inches long. (Named for Prof. Johann Reinhardt, naturalist at the University of Copenhagen.)

*Himantolophus reinhardtii*, LÜTKEN, Kong. Dansk. Vidensk. Selsk. 1878, 321, Greenland. *Corynolophus reinhardtii*, GILL, Proc. U. S. Nat. Mus. 1878, 219.

1059. LIOCETUS, Günther.

*Liocetus*, GÜNTHER, Challenger Report, xx., 57, 1887 (*murrayi*).

Mouth enormous, the cleft nearly vertical; pectorals small, in advance of dorsal and of gill opening; second dorsal spine wanting; gills in  $2\frac{1}{2}$  pairs; no gular tentacle. This genus is similar to *Melanocetus*, differing in having no teeth on the vomer, a greater projection of the mandible, and a smaller mouth. Deep sea. (*λεῖος*, smooth; *κῆτος*, whale.)

3116. LIOCETUS MURRAYI (Günther).

D. I-13; A. 4; C. 9; P. 14. (Günther.) Extremely similar to *Melanocetus johnsonii*, but without trace of vomerine teeth, while there is no distinction between the two species as regards dentition of jaws; posterior angle

\*Dr. Gill, replying to certain strictures as to the validity of this genus, made by Lütken (who calls it a "wanton" subdivision), states that the "differences alleged to exist between *Himantolophus* and *Corynolophus* are very marked. If they do exist as stated there can be no doubt that the two should be kept apart. I know of no reason except the singularity and greatness of the difference specified for doubting the correctness of Reinhardt's observations."

of mandible projecting more and forming a salient point; mouth comparatively less wide, and the maxillary considerably shorter, being about  $\frac{2}{3}$  of total length, without caudal, while it is rather more than  $\frac{1}{2}$  in the Madeira species. Eye rudimentary. One cephalic spine, shorter than maxillary; last dorsal ray connected by a short and delicate membrane with caudal fin; most of the caudal rays bifid, the longest shorter than maxillary; pectoral fin as much developed as in *Melanocetus johnsonii*. Entirely black. Total length 44 lines; length of mandible 14 lines; length of maxillary 12 lines; length of caudal fin  $10\frac{1}{2}$  lines. A young individual, 44 lines in length, was taken by H. M. S. *Challenger* in the mid-Atlantic, at a depth of 1,850 fathoms (Station 106); another of 13 lines at the depth of 2,450 fathoms (Station 348). (Goode & Bean.) (Named for Dr. John Murray, second director of the civilian staff on board H. M. S. *Challenger*.)

*Melanocetus bispinosus*, GÜNTHER, Study of Fishes, 473, 1880; name only.

*Melanocetus (Liooetus) murrayi*, GÜNTHER, Challenger Report, XXII, 57, pl. 11, fig. A, 1887, mid-Atlantic.

*Liooetus murrayi*, GOODE & BEAN, Oceanic Ichthyology, 495, fig. 407, 1896.

#### 1070. LINOPHRYNE, Collett.

*Linophryne*, COLLETT, Proc. Zool. Soc. London 1886, 138 (*lucifer*).

Head enormous; the body slender, compressed, mouth oblique. Spinous dorsal reduced to a single cephalic tentacle, the basal part of which is erect, not procumbent. Teeth in the jaws on the vomer and the upper pharyngeals. Gill openings exceedingly narrow, situated a little below the root of the pectoral. Soft dorsal and anal very short; ventrals none. Abdominal cavity forming a sac, suspended from the trunk. Skin smooth; a long tentacle on the throat. This genus differs from *Melanocetus* in the presence of the gular tentacle. (*λίπος*, linsn, net; *φρύνη*, a toad.)

#### 3117. LINOPHRYNE LUCIFER, Collett.

D. 1-3; A. 2; C. 9; P. 14 or 15. A spinous projection or horn above each orbit. Cephalic tentacle black, with a large ovate bulb, the upper half of which is white; gular tentacle much larger, terminating in 2 tongue-like appendages, which are furnished on the upper edge with a row of round, white papillæ. (Goode & Bean.) Mid-Atlantic, northwest of Madeira, Lat. 36° N., Long. 20° W. One specimen known. (*Lucifer*, an evil spirit; *lux*, light; *fero*, I bear.)

*Linophryne lucifer*, COLLETT, Proc. Zool. Soc. London 1886, 138, pl. 15, mid-Atlantic, between Madeira and the West Indies (Coll. Capt. P. Andresen. Mus. Univ. Christiania); GÜNTHER, Challenger Report, XXII, 57, 1887; GOODE & BEAN, Oceanic Ichthyology, 496, fig. 408, 1896.

#### 1071. CAULOPHRYNE, Goode & Bean.

*Caulophryne*, GOODE & BEAN, Oceanic Ichthyology, 496, 1896 (*Jordani*).

Head large, compressed; mouth with the cleft nearly horizontal; body short, much compressed. Spinous dorsal reduced to a single cephalic tentacle, which is supported on a short procumbent base. Teeth of unequal size in the intermaxillary and the mandible; vomer, palatines, and upper

pharyngeals toothed. Gill openings narrow, horizontal slits placed below and in front of root of pectorals. Branchiæ in  $\frac{1}{2}$   $2\frac{1}{2}$  pairs. Branchial arches armed with denticigerous tubercles. Skin naked. Numerous luminous filaments on head and body. Soft dorsal and anal many-rayed, the rays greatly produced; caudal long, tapering; ventrals none; pectorals very broad, sessile, postmedian, under dorsal fin, with numerous rays. Pyloric appendage reduced to 1 small rudiment. Air bladder absent. (*καυλός*, stem; *φρόνη*, toad, from the many stems or fin rays.)

3118. CAULOPHYNE JORDANI, Goode & Bean.

Depth about 2 in length without caudal, the greatest height occurring behind the head. Cephalic appendage with a pale tuft at its tip, the length of the distal portion 3 in length of body. The tuft somewhat mutilated, but showing no evidence of a laminated structure; basal portion of cephalic appendage about twice as long as the very small eye; maxillary very slender, narrow, extending about as far backward as intermaxillary; intermaxillary slightly protractile and with about 10 teeth on each side, several of which are nearly twice as large as the rest, its length  $2\frac{1}{2}$  in body; mandible as long as head without snout, with 8 teeth on each side, the anterior pair and several other pairs along shaft of bone being greatly enlarged; a pair of enlarged teeth on head of vomer; several similar teeth on palatines; upper pharyngeals armed with several strong teeth; eye very small, inconspicuous, its distance from tip of snout equaling nearly  $\frac{1}{2}$  its distance from soft dorsal origin. Intestine shorter than length without caudal. Soft dorsal with 16 rays, all of which, except the last 4, are greatly produced; the second, third, and fourth rays longest, nearly twice as long as body. Anal with 14 rays, all of which, except last 3, are much produced, the fin not quite perfect, yet its anterior rays are longer than body; caudal with 8 rays, the 4 inner ones divided, the rest simple; middle rays of caudal as long as distance from tip of lower jaw to base of pectoral; pectoral comparatively short, with 16 simple articulated rays, the longest about  $\frac{1}{2}$  as long as head. About 9 luminous filaments on each side of head, 7 more between nape and dorsal, and about 12 on sides; the filaments nearly twice as long as eye. Head and body black; caudal, cephalic tuft, and most of the rays pale. Gulf Stream. The type of the species (No. 39265) was taken by the steamer *Albatross*, September 19, 1887, in Lat.  $39^{\circ} 27' N.$ , Long.  $71^{\circ} 15' W.$ , 1,276 fathoms. (Named for David Starr Jordan.)

*Caulophryne jordani*, GOODE & BEAN, *Oceanic Ichthyology*, 496, pl. 21, fig. 409, 1896, Gulf Stream, off Carolina, in 1,276 fathoms. (Coll. *Albatross*; the plate named *Caulophryne setosus*, by slip in proof reading.)

Family CCXXIV. OGCOEPHALIDÆ.

(THE BAT-FISHES.)

Head very broad and depressed, the snout more or less elevated, the trunk short and slender. Mouth not large, subterminal or inferior, the lower jaw included; teeth villiform or cardiform. Gill openings very



small, above and behind the axils of the pectoral fins. Body and head covered with bony tubercles or spines. Spinous dorsal reduced to a small rostral tentacle, which is retractile into a cavity under a prominent process on the forehead; in 1 genus the rostral tentacle is obsolete; soft dorsal and anal fins small and short; ventrals well developed; pectoral fin well developed, its base strongly angled, with long pseudobranchia and 3 actinosts. Branchiostegals 5; no pseudobranchia. Genera 8; species about 30, chiefly American, some of them in the deep sea. (*Pediculati*, part; genera *Malthe* and *Halieutæa*, Günther, III, 200-205, 1861.)

## OGCOCEPHALINÆ:

- a. Disk with the frontal region elevated and the snout more or less produced forward, the tail stout; orbits lateral; teeth on vomer and palatines; rostral tentacle present.  
 b. Gills 2½; disk longer than broad. OGCOCOPHALUS, 1072.  
 bb. Gills 2¼; disk broader than long. ZALIEUTES, 1073.

## HALIEUTINÆ:

- aa. Disk with the frontal region depressed, not elevated above the rest; eyes partly superior; snout rounded, obtuse in front; tail slender.  
 c. Dorsal fin present.  
 d. Vomer and palatines with teeth. HALIEUTICETHYS, 1074.  
 dd. Vomer and palatines toothless.  
 e. Disk subcircular; gills 2¼.  
 f. Mouth rather large, subvertical; prickles rather strong. HALIEUTÆA, 1075.  
 ff. Mouth rather small, terminal; prickles feeble. HALIEUTELLA, 1076.  
 ee. Disk subtriangular; gills 2; prickles very strong. DIBRANCHUS, 1077.

## 1072. OGCOCEPHALUS, Fischer.

## (SEA-BATS.)

*Ogcocephalus*, FISCHER, Zoognosia, 78, 1813 (*vespertilio*).

*Oncocephalus*, GILL, modified spelling.

*Malthe*, CUVIER, Règne Animal, Ed. I, II, 311, 1817 (*vespertilio*).

*Malthæa*, corrected spelling.

Body stoutish, tapering backward; head very broad and depressed, triangular in form, the forehead elevated and produced. Eyes large, lateral. Mouth rather small, subinferior under the snout; villiform teeth in bands, on jaws, vomer, and palatines. Skin covered with rough, bony tubercles. Dorsal and anal fins very small; rostral tentacle present, retractile into a cavity under a bony prominence on the forehead; ventrals present, I, 5, well separated; pectorals large, placed horizontally. Gills 2½. No air bladder; no pyloric caeca. Tropical America, in shallow water; small fishes of singular form, often regarded by the ignorant as venomous. (*ὄγκος*, hook; *κεφαλή*, head; properly written *Oncocephalus*, but Fischer chose the above monstrous spelling.)

- a. Snout produced, the rostral process pointed, 6 to 10 in length of body. VESPERTILIO, 3119.  
 aa. Snout short, the rostral process 12 to 15 times in length of body. NASUTUS, 3120.  
 aaa. Snout short, the rostral tubercle reduced to a button-like tubercle, which is about 25 times in length of body. RADIATUS, 3121.

## 3119. OGCOCEPHALUS VESPERTILIO (Linnaeus).

(BAT FISH; DIABLO.)

Head, from tip of upper jaw to gill opening, nearly  $\frac{1}{2}$  the length; depth 5 in length from upper jaw to base of caudal; width  $1\frac{1}{2}$ . D. 4; A. 4; rostral process from 6 to 10 (9 in our specimens from Havana); P.  $4\frac{1}{2}$ ; V. 6; C.  $4\frac{1}{2}$ . Body stoutish, much depressed, rostral process longer than in the other species, variable in length; mouth small, the maxillary reaching nearly to posterior margin of eye; villiform teeth in bands, on jaws, vomer, and palatines; interorbital flattish, its width less between anterior edge of eyes than posterior edge; rostral groove longer than broad; body covered with bony protuberances, variable in size, and not very definite in position, lower parts with a shagreen-like covering; posterior edge of pectorals much behind middle of body; ventrals long, reaching outward to edge of the disk-like, anterior part of body; origin of dorsal over posterior edge of pectoral; anal under the vertical of tips of dorsal rays, anal reaching nearly to base of caudal. Pale grayish brown above, reddish below; back with round black spots, conspicuous in life, but growing fainter and sometimes disappearing in spirits; belly in life a coppery red; pectorals nearly plain dusky. Length 12 inches. West Indies, north to the Florida Keys; common in shallow water. Here described from a specimen from Havana, Cuba, about 10 inches in length. The length of the snout is subject to great variation, but it is never short and button-like, as in *O. radiatus*. (*vespertilio*, a bat.)

*Lophius vespertilio*, LINNÆUS, Syst. Nat., Ed. x, 1, 236, 1758, American Seas; after *Lophius fronti unicorni* of ARTEDI.

*Malthæa vespertilio*, CUVIER & VALENCIENNES, Hist. Nat. Poiss., XII, 440, 1837; prominence on snout 10 in length; DE KAY, N. Y. Fauna: Fishes, 167, 1842; GÜNTHER, Cat., III, 200, 1861; JORDAN & GILBERT, Synopsis, 850, 1883; JORDAN & SWAIN, Proc. U. S. Nat. Mus. 1884, 234.

*Lophius rostratus*, SHAW, Zool., IV, 383, pl. 163, 1803.

*Malthæa longirostris*, CUVIER & VALENCIENNES, Hist. Nat. Poiss., XII, 452, 1837, Bahia (Coll. Blanchet); snout 6 in length.

## 3120. OGCOCEPHALUS NASUTUS (Cuvier &amp; Valenciennes).

Head 2. D. 4; A. 4. Rostral process short, about 12 to 15 times in length of body. Cavity of nostril tentacle higher than broad; width of body 2 in length; vent behind middle of body. Dusky above, with round black spots, edged with whitish. West Indies. (Lütken); not seen by us; perhaps a variation of *O. vespertilio*. (*nasutus*, long-nosed.)

*Malthæa nasuta*, CUVIER & VALENCIENNES, Hist. Nat. Poiss., XII, 452, 1837, Martinique.

*Malthæa nasuta*, LÜTKEN, Nat. For. Vid. Medd. 1865, 4; JORDAN & GILBERT, Synopsis, 850, 1883.

*Malthæa notata*, CUVIER & VALENCIENNES, Hist. Nat. Poiss., XII, 453, 1837, Surinam; snout 15 in length; body spotted.

## 3121. OGCOEPHALUS RADIATUS (Mitchill).

(SHORT-NOSED BAT-FISH.)

Head 2. Dorsal 4; A. 4. Rostral cavity somewhat broader than high, or width equal to height; distance between anterior angles of orbits about equal to that between the posterior angles; eye a little wider than interorbital width; snout, exclusive of rostral tubercle, not produced beyond the rostral cavity, but with a cylindrical button-like tubercle, slightly contracted at base, pointing obliquely upward and forward, its length 25 times in body; posterior edge of pectoral slightly nearer base of caudal than upper jaw; caudal peduncle very thick and heavy; vent about midway between tip of jaw and base of caudal fin. Color brownish, with dark round spots sometimes edged with white; pectorals with a network of white lines dividing the dark color into dark brown spots; tip of caudal blackish, belly coppery red. Length 8 to 12 inches. Coast of Florida and neighboring waters; very common in shallow bays among weeds, especially about the Florida Keys. Here described from a specimen from Cedar Key, Florida, 7 to 8 inches in length. (*radiatus*, rayed.)

*Lophius radiatus*, MITCHILL, Amer. Monthly Mag., March, 1818, 326, Straits of the Bahamas.

*Malthæ cubifrons*, RICHARDSON, Fauna Bor.-Amer., III, 103, 1836, said to be from Labrador (Coll. J. J. Audubon), but this is certainly an error; Audubon collected also in Carolina and Florida; GÜNTHER, Cat., III, 203, 1861; JORDAN & GILBERT, Synopsis, 850, 1893; JORDAN, Cat. Fishes, 139, 1885.

? *Malthæa truncata*, CUVIER & VALENCIENNES, Hist. Nat. Poiss., XII, 454, 1837, America; snout wholly obsolete; perhaps a species of *Zalieutes*.

*Malthæa angusta*, CUVIER & VALENCIENNES, Hist. Nat. Poiss., XII, 454, 1837, Dutch Guiana; snout more than 20 in length.

## 1073. ZALIEUTES, Jordan &amp; Evermann.

*Zalieutes*, JORDAN & EVERMANN, Check-List Fish. N. and M. A., 511, 1896 (*elater*).

Disk wider than long, about as long as rest of body (including caudal fin); middle line of head elevated, but the forehead not projecting beyond mouth; rostral tentacle present, the cavity about as wide as high; mouth small; minute teeth on vomer and palatines. Gills 2½. Eastern Pacific. The genus is very close to *Malthopsis*, Alcock,\* but the latter, like *Ogcocephalus*, has the disk longer than broad, but the gills are reduced to 2. (ζάλη, surge of the sea; ἀλιευτής, fisher.)

## 3122. ZALIEUTES ELATER (Jordan &amp; Gilbert).

Body very broad and depressed, the disk considerably broader than long, its width 1½ times in length of body; back and snout considerably raised above rest of body; greatest depth of body scarcely more than width of mouth. Mouth small, its width ½ greater than diameter of orbit; snout very short, scarcely projecting beyond mouth, its length

\* *Malthopsis*, ALCOCK, Ann. and Mag. Nat. Hist. 1891, 26; *Malthopsis luteus*, from the Andaman Sea; *Malthopsis mitrifer*, GILBERT & CRAMER, Proc. U. S. Nat. Mus. 1896, 434, with plate, off Hawaiian Islands.

about equal to interorbital width, shorter than its own width in front. Eye rather large, much longer than snout, wider than interorbital area. Process representing first dorsal spine present, small. Skin covered with spines, which are comparatively slender and sharp, their stellate bases inconspicuous, those on snout and middle of back and tail largest, much slenderer and sharper than in *Ogcocephalus vespertilio*; no spines on ocelli of back; belly rough; under side of tail with tubercular plates; tail depressed toward base of fin. Pectorals  $\frac{1}{2}$  longer than ventrals, their length  $1\frac{1}{2}$  width of mouth; caudal a little longer than pectoral,  $4\frac{1}{2}$  in body. Color light olive, above everywhere thickly and uniformly covered with small round spots of dark brown, these about as large as the pupil and about as wide as the lighter interspaces; a conspicuous ocellus, larger than eye, on each side of back, this ocellus with a bright yellow spot in the center, surrounded by a black ring, around which is a pale ring, and finally a fainter dark one; under parts plain white; pectorals spotted; caudal yellowish at base, with a terminal blackish band. Length 4 inches. Pacific coast of Mexico, south to Panama, in water of moderate depth; very rare near the shore, but obtained in abundance by the Albatross at Stations 2794 and 2795, near Panama. (*Elater*, the spring beetle, from the resemblance of the ocelli to the eye-like spots on the back of *Elater*.)

*Milthe elater*, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1881, 365, Mazatlan. (Type, No. 28127. Coll. Dr. J. U. Bastow.)

*Ogcocephalus elater*, JORDAN, Proc. Cal. Ac. Sci. 1895, 536.

#### 1074. HALIEUTICHTHYS, Poey.

*Halieutichthys*, POEY, Proc. Ac. Nat. Sci. Phila. 1863, 83 (*reticulatus*).

Disk subcircular, anteriorly cordiform, the head merging into the body, very large and much depressed; cranial portion not elevated; interorbital space low and narrow; eyes partly superior; mouth terminal, horizontal, the jaws subequal, the lower jaw nearly semicircular; teeth fine, on jaws and palate. Gills  $2\frac{1}{2}$ ; no gill rakers; gill openings anterior to pectoral; rostral tentacle very small, retractile; dorsal and anal few-rayed; pectorals large, the carpus slender; caudal rounded; skin above sparsely armed with stellate tubercles; lower surface smooth. (*ἁλιευτήρ*, fisher; *ἰχθύς*, fish.)

a. Surface of body covered with brownish reticulations; fins not barred with black.

ACULEATUS, 3123.

aa. Surface of body blackish, not reticulate; pectorals with a broad black bar mesially, the tip pale; caudal blackish toward the tip.

CARIBBEUS, 3124.

#### 8123. HALIEUTICHTHYS ACULEATUS (Mitchill).

D. I, 4 or 5; A. 4; V. I, 5; P. 16 to 18; C. 9; gills  $2\frac{1}{2}$ . Disk cordiform, about as wide as long, its length more than  $\frac{1}{2}$  that of body. Body covered above with stout conical spines with stellular bases, largest upon the trunk, upon which they are arranged in about 2 irregular longitudinal

rows on each side of the dorsal; upon the disk they are placed above the principal bones of the skeleton, most abundant upon its cranial portion; a single row of stout spines, usually 3-pointed, on the outer margin of disk, a particularly large one at each outer angle; body entirely smooth below; snout very short, obtuse; bridge over the rostral cavity covered in front with a 3-pointed spine, having on each side a simple spine; short, stout, simple spines upon each supraorbital margin, the front of which is immediately above and behind the cavity containing the nostrils; vertex with several similar spines; many spines closely placed upon the humeral area; numerous short tentacles upon margin of disk and on sides of trunk; supraoral cavity elliptical, small (horizontal diameter  $\frac{7}{8}$  diameter of orbit), containing a well-developed, club-shaped, very perceptible tentacle; width of opening of anterior nostril, which is in a short tube,  $\frac{1}{2}$  that of posterior nostril, which is not tubular; width of mouth much less than distance between pupils and equaling diameter of orbit. Diameter of orbit  $8\frac{1}{2}$  times in distance from snout to base of caudal, 6 times in distance from snout to origin of soft dorsal,  $6\frac{1}{2}$  times in distance to origin of anal, 3 times in distance to base of ventrals, and 6 times in distance to angle between pectorals and trunk,  $4\frac{1}{2}$  times in distance from snout to gill opening, 6 in greatest width of disk, and nearly 2 in that of trunk; width of interorbital area  $\frac{2}{3}$  diameter of orbit. First dorsal ray longest, equal to diameter of orbit; anal fin inserted under third ray of dorsal, with 4 rays, the third or longest very slightly longer than the longest dorsal ray; ventral fins inserted nearly under the middle of the disk, with 1 rudimentary and 5 dorsal rays, increasing in length posteriorly, the last and longest contained 5 times in total length; distances between origins of ventrals  $6\frac{1}{2}$  in total length; pectorals with peduncles entirely included in common membrane, with blades far back, horizontal, lying close to trunk, composed of 16 rays, the middle or longest  $3\frac{1}{2}$  in total length; caudal fin rounded, composed of 9 rays, the external rays, 1 above and 2 below, simple, the others bifid; length of middle ray equal to that of trunk (measured from junction of pectorals to base of caudal rays) and slightly exceeding the longest pectoral ray. Length of intestine contained  $1\frac{1}{2}$  times in total length. Color, body covered above with reticulations of brown, the general hue varying from light yellowish gray to grayish brown, the markings being darker upon darker specimens; pectoral and caudal fins with about 3 dark bars; the terminal bars in young very black; body beneath milky white (Goode & Bean.) West Indies, Gulf of Mexico, and Gulf Stream, in water of moderate depth; taken by the *Blake* and the *Albatross* at numerous stations in depths ranging from 10 to 95 fathoms. "As in *Halicutaea*, *Dibranchus*, and allies, a rostral tentacle is present in this genus. Among specimens belonging to the Museum of Comparative Zoology there is evidence of the existence of a couple of distinct forms in the West Indian waters. The true *H. aculeatus* is much the lighter in the ground colors and has brownish reticulations across the back, 2 or 3 narrowish transverse bands of the same color across the pectorals, and 2 or 3 similar bands appear on the caudal, the posterior being darkest. The margins of the fins are light in color. The rostrum is acute; it ends in a spine which

turns upward, and, seen from above, it is hardly long enough to cover the tentacular niche. Evidently this type belongs to the shallower waters. The localities noted carry its distribution from the Bahamas to the Yucatan Banks, in depths of 40 fathoms and less." (Garman.) (*aculeatus*, with needle-like spines.)

*Lophius aculeatus*, MITCHILL, Amer. Mon. Mag., II, 1818, 325, Straits of Bahama.  
*Halieutichthys reticulatus* (POEY MS.) GILL, Proc. Ac. Nat. Sci. Phila. 1863, 91, Cuba (Coll. Prof. Felipe Poey); JORDAN & GILBERT, Synopsis, 851, 1883.  
*Halieutichthys aculeatus*, GOODE, Proc. U. S. Nat. Mus. 1879, 109; GOODE & BEAN, Oceanic Ichthyology, 594, pl. 122, fig. 414a and b, 1896; GARMAN, Bull. Lab. Nat. Hist. Iowa Univ. 1896, 87, pl. 4, fig. 1.

3124. HALIEUTICHTHYS CARIBBÆUS, Garman.

D. I-5; A. 4; V. 5; P. 17; C. 9. Color darker than *H. aculeatus*; the reticulations are not present; the outer half of the pectoral, except at the margin, is black; and, excepting the narrow posterior margin, the hinder fifth of the caudal fin is black; the upper surface is clouded brownish without traces of the network pattern common to *H. aculeatus*. On the specimens described, the rostrum is acute, and the spine extends forward to cover the cavity receiving the tentacle so that it is not visible when viewed from above. West Indies. As now known, this species ranges from Jamaica to Barbados in depths of 70 to 150 fathoms or more. (Garman.) (*caribbæus*, from the Caribbean Sea.)

*Halieutichthys caribbæus*, GARMAN, Bull. Lab. Nat. Hist. Iowa Univ. 1896, 87, pl. 4, fig. 2, Jamaica to Barbados.

1075. HALIEUTÆA, Cuvier & Valenciennes.

*Halieutæa*, CUVIER & VALENCIENNES, Hist. Nat. Poiss., XII, 455, 1837 (*stellatus*).

Head very large, broad, depressed, its outline nearly circular; cleft of the mouth wide, horizontal; jaws with small cardiform teeth; no teeth on vomer or palatines. Skin everywhere covered with small, stellate spines. Forehead with a transverse bony ridge, beneath which is a tentacle, retractile into a cavity, the only rudiment of the spinous dorsal fin; soft dorsal and anal very short, far back. Gills  $2\frac{1}{2}$ , the anterior gill arch without laminae. Branchiostegals 5; vertebrae 17. Pacific Ocean. (*άλιευτής*, one who fishes.)

3125. HALIEUTEA SPONGIOSA, Gilbert.

D. 6; A. 4; C. 9; V. 4; P. 12 or 13. This species is remarkable for the soft, spongy texture of the body, and the membranaceous or cartilaginous character of its bones. Width of head  $1\frac{1}{2}$  in its length; tail long and slender, the vent midway between base of caudal and articulation of mandible; width of base of tail  $4\frac{1}{2}$  in its length; mouth little or not at all overpassed by the snout, its width  $2\frac{1}{2}$  to 3 in that of head, lower jaw usually not included; gape of mouth oblique, almost wholly anterior. Teeth in wide cardiform bands in the jaws, none of them enlarged; palate toothless. Interorbital width slightly greater than length of snout, 5 in width of

head; eye  $1\frac{1}{2}$  in interorbital width. Rostral tentacle short, with an expanded 3-lobed tip; front of dorsal midway between base of caudal and occiput; caudal long, rounded, the lower rays more shortened than the upper, the longest nearly  $\frac{1}{2}$  width of head; anal rays high, closely bound together, the fin slender, shaped like the intromittent organ of *Gambusia*, the length of its base equaling  $\frac{2}{3}$  diameter of orbit, its longest ray reaching base of caudal; pectorals long, the posterior rays rapidly shortened, the longest  $\frac{1}{2}$  width of head; head and body everywhere with broadly conical, tubercular plates, varying in size, marked with strong lines, radiating from the center; the apex sometimes blunt, more often provided with a slender spine, sometimes bifid or trifid; on the tail these spines become longer and are directed backward; plates along edge of disk not compressed nor specially modified. A deep groove-like channel just behind mandible and following curve of latter, becoming continuous with another deeper channel running just below edge of disk to near base of pectorals; a third groove runs backward from nostrils, uniting with the others, these grooves spanned at intervals by pairs of fleshy tentacles with fringed tips, which spring from the edges of the grooves and meet across them; at the bottom of the grooves under each pair of tentacles is a small fleshy tubercle. Fin rays, at least at base, with series of small curved prickles. Color uniform dusky, the tail sometimes lighter; fins blackish, more or less edged with white. One specimen with the body and tail uniformly light. Pacific coast of Mexico in deep water. Numerous specimens, the largest  $4\frac{1}{2}$  inches long, from Albatross Station 2992, in 460 fathoms. (Gilbert.) (*spongiosus*, spongy.)

*Halieutæa spongiosa*, GILBERT, Proc. U. S. Nat. Mus. 1890, 124, west of Revillagigedo Islands, at Albatross Station 2992, Lat.  $18^{\circ} 17' 30''$  N., Long.  $114^{\circ} 43' 15''$  W., in 460 fathoms. (Coll. Gilbert.)

#### 1076. HALIEUTELLA, Goode & Bean.

*Halieutella*, GOODE & BEAN, Proc. Biol. Soc. Washington, 1882, 88 (*lappa*).

Body subcircular, depressed, its width equal to its length, covered with flaccid, inflatable skin. Spines feebler and less numerous than in *Halieutæa*. Head merged in body; forehead with a transverse bony ridge; no perceptible supraoral cavity; no tentacle. Mouth small, terminal; lower jaw slightly curved forward. Teeth in the jaws minute, cardiform, not discernible on palate, though possibly present. Carpus broad, slightly exerted; pectoral fins remote from tail, obliquely placed, with membranes subvertical. Branchial aperture posterior to carpus, upon the disk, and not remote from its margin. Gills  $2\frac{1}{2}$ . Dorsal fin 5-rayed, inserted at junction of disk with caudal peduncle; anal fin 4-rayed, originating at root of caudal peduncle. (*ἀλιευτήρ*, a fisherman.)

#### 3126. HALIEUTELLA LAPPA, Goode & Bean.

D. 5; A. 4; C. 9; P. 15; V. 5. Disk subcircular, more than  $\frac{2}{3}$  as long as the body. Body covered with a loose, flaccid, inflatable skin, which so obscures its proportion, that it is impossible to determine its exact height,

but it is not nearly so much depressed as in the related genera. When the body is inflated the height and length of the disk is nearly equal. Spines rather feeble; about 10 between snout and dorsal fin; about 6 strong spines, with conical bases and stellular tips, on outer margin of disk on each side, the anterior of them being opposite the eye; in front of these spines on the discal margin, and between them and the snout, are several small, simple spines, pointing backward; belly armed with spines similar to those on the back, but weaker; a stellate spine upon tip of snout, with 2 weaker, simple spines on each side; nasal openings midway between eye and tip of snout; mouth small, upon the margin of the disk; upper jaw shorter than diameter of eye. Teeth as described in the generic diagnosis. Dorsal fin inserted at posterior limit of disk, with 5 simple, articulated rays, its longest ray  $\frac{1}{2}$  as long as disk; anal fin with 4 simple, articulated rays, inserted directly beneath fourth ray of dorsal, its second and longest ray  $\frac{1}{2}$  as long as disk; caudal twice as long as anal, and slightly longer than caudal peduncle, with 9 simple, articulated rays. Carpus inserted at a distance from snout equal to twice length of longest pectoral ray, which is slightly greater than distance of posterior margin of carpus, at its junction with disk, from vent; number of pectoral rays 15; ventral inserted at a point equidistant from the snout and origin of anal, its longest ray (the fourth) equal to  $\frac{1}{2}$  distance of anal fin from snout. Color yellowish white. Gulf Stream. A single specimen,  $1\frac{1}{2}$  inches long, known. (*Iappa*, the burdock, from its prickles.)

*Haliutella Iappa*, GOODE & BEAN, Proc. Biol. Soc. Washington, II, 1882, 88, Gulf Stream, at Fish Hawk Station 1151, Lat.  $39^{\circ} 58' 30''$  N., Long.  $70^{\circ} 37'$  W., in 125 fathoms; GOODE & BEAN, Oceanic Ichthyology, 500, pl. 122, figs. 512a and 512b, 1896.

### 1077. DIBRANCHUS, Peters.

*Dibranchus*, PETERS, Monatsber. Kon. Akad. Wiss. Berlin 1876, 736 (*atlanticus*).

Head merged in body, very large, much depressed, forming a broadly ovate disk, with margin laterally prolonged; cranial portion not elevated; the interorbital area low, narrow, with orbits partly superior; supraoal cavity large, protected above by a transverse bony ridge. Mouth terminal, horizontal, wide; lower jaw convex; teeth in cardiform bands, none on vomer or palatines. Gills 2; no gill rakers; gill openings small, anterior to pectorals. Rostral tentacle retractile, trilobate at tip. Skin with numerous strong stellate spines above and below, those at margins of disk especially strong, 3-pointed. Atlantic; distinguished from related genera by the reduction of the gills to 2 pairs. (*δῖς*, two; *βράγχος*, gill.)

### 3127. DIBRANCHUS ATLANTICUS, Peters.

D. 6 or 7; A. 4; C. 9; P. 13 to 15; V. I, 5; Br. 6; gills 2. Disk orbicular, nearly as wide as long, its length about  $\frac{1}{2}$  that of body, its lateral outline prolonged on each side, and terminating in a strong spine armed at the tip with a group of irregularly arranged acicular spinelets. Body covered above with numerous stout conical spines with stellular bases, these



largest upon the trunk, where they are arranged approximately in about 4 irregular longitudinal rows upon each side of the dorsal fin; closely set rows of these stout spines mark the outer margin of the disk, and there is also a cluster of 5 to 7 upon each carpal peduncle; outside of these marginal spines, upon each side, is an irregular marginal row of 5 depressed knife-like spines, each tipped with a crown of 3 acicular spinelets; on the anterior margin of the disk the 2 rows coalesce to a greater or less extent and form a bristling array of closely set spines, some pointing dorsally, some laterally, some ventrally; 2 kinds of spines upon the dorsal surface, in addition to the large ones already described, some large, somewhat remote from each other, conical, stellular; others, much more numerous and filling the interspaces, pickle-like, stellular; belly armed with numerous closely-set spines of a similar kind; snout somewhat projecting, armed with 3 many-tipped spines; a spine-armed ridge in front of the eyes, over the top of the snout; in this 4 spines are conspicuous, 1 in front of each eye, and between these a larger pair in front of the supraorbital ridges; from these last mentioned spines extend spine-armed ridges along the upper margins of each orbit; under the snout is a cavity (horizontal diameter  $\frac{1}{2}$  that of the orbit) containing a barbel, pedicled, with thick, club-shaped, trilobate tip; on each side of this cavity are the nasal openings, which are as in *Halieutichthys*. Width of mouth equal to distance between centers of pupils of eyes. Diameter of orbit contained as follows in other dimensions of the body: In total length  $9\frac{1}{2}$ ; in distance from snout to dorsal 6; same to anal 7; the base of ventrals 3; to angle between pectorals and trunk  $5\frac{1}{2}$ ; to gill openings 5; in greatest width of disk  $5\frac{1}{2}$ ; of trunk 4. Width of interorbital area in diameter of orbit  $\frac{2}{3}$ . Dorsal fin with 6 or 7 rays, the longest (third)  $1\frac{1}{2}$  times diameter of orbit and 6 times in total length; anal fin inserted entirely behind dorsal, with 4 rays, the longest (third) about as long as longest in dorsal fin; ventral fins inserted nearly under middle of disk, a little nearer vent than to mandibular symphysis, with 1 rudimentary and 5 well-developed rays, increasing in length posteriorly, the last and longest  $6\frac{1}{2}$  times in total; distance between ventral organs  $7\frac{1}{2}$  in total length. Pectorals with peduncles slightly exerted, bases included in common membrane, composed of 13 to 15 rays, the longest third or fourth  $4\frac{2}{3}$  in total. Caudal fin rounded, consisting of 9 rays, all bifid except the 2 external ones; length of middle ray about  $\frac{1}{2}$  that of trunk and exceeding that of pectoral, being contained  $4\frac{1}{2}$  times in total length. Stomach egg-shaped, intestine somewhat longer than body; liver very wide and large. Color uniform reddish, gray above, slightly lighter below. Deep waters of the Atlantic; very abundant, in about 300 fathoms. Known from the west coast of Africa, off the Cape Verdes, off Barbados, and north in the Gulf Stream to Newport. (*atlanticus*, of the Atlantic.)

*Dibranchius atlanticus*, PETERS, Monatsber. Kon. Akad. Wiss. Berlin 1876, 736, with plate, West Africa, Lat.  $10^{\circ}$  N., Long.  $17^{\circ}$  W., in 360 fathoms (Coll. H. M. S. *Gazelle*); GÜNTHER, Challenger Report, XXII, 59, 1887; VAILLANT, Travalleur, etc., 343, 1883; GOODE & BEAN, Oceanic Ichthyology, 501, pl. 122, fig. 413, 1896.

## ADDENDA.

Page 12. After *Entosphenus tridentatus* add:

### 11(a). ENTOSPHEMUS CAMTSCHATICUS (Tilesius).

A lamprey taken by Steller in the Bolschaya River, Kamchatka, has not been recorded by subsequent writers. It is reported by Steller as 13½ inches in length; the head ½ of an inch; mouth long, with 2 teeth above, 6 below; dorsals 2. Color shining brassy, dark above; sides with dusky serpentine lines. A figure published by Tilesius shows the upper teeth as bifid, and 9 teeth below.

Pallas describes specimens from the sea at Petropaulski as 7 inches long, not marbled nor variegated. The species of Steller is probably an *Entosphenus*. That of Pallas may be the same, or it may be a *Lampetra*, allied to or identical with *L. aurea*.

*Petromyzon marinus camtschaticus*, TILESIIUS, Mém. Acad. St. Petersburg 1809, 240, with plate, Kamchatka.

*Petromyzon camtschaticus*, TILESIIUS, l. c., 241.

*Lampetra variegata* (STELLER MS.) TILESIIUS, l. c., 247, Bolschaya River, Kamchatka.

! *Petromyzon marinus camtschaticus*, PALLAS, Zoogr. Ross. Asiat., III, 1810, 67, Petropaulski

Page 14. From the synonymy of *Lampetra wilderi* omit "*Petromyzon branchialis*, Günther, Cat., VIII, 504, 1870," and after the last synonym add: Not *P. branchialis*, Linnaeus, which is the larva of some European species, perhaps of *P. marinus*.

Page 25. In the description of *Catulus uter* the teeth should read 88 instead of 83.

Page 27. In the key, under *dd*, read: Root of tail with a conspicuous notch above.

Page 28. The following key to West Coast species of *Galeus* and *Mustelus* will prove helpful.

- a. Eye large, spiracle small, the latter not more than ¼ major diameter of orbit.
  - b. Mouth broad, snout broadly rounded, mandibular angle of 90° or more. Fins less deeply incised; the lower caudal lobe rounded; pectoral and ventral margins nearly straight. GALEUS CALIFORNICUS, 33.
  - bb. Mouth narrow, snout long, acute, mandibular angle 60° to 65°. Fins deeply incised; lower caudal lobe acute; pectoral and ventral margins concave. MUSTELUS LUNULATUS, 30.
- aa. Eye small, spiracle large, the latter ½ to ¾ the major diameter of orbit. Snout sharp, mouth narrow, the mandibular angle about 70°. Terminal lobe of caudal broad, obliquely truncate posteriorly. Nostrils very large, their width nearly equalling width of interspace. Fins less incised. GALEUS DORSALIS, 32.

Page 37. After *Carcharhinus heulei* add:

45(a). *CARCHARHINUS CERDALE*, Gilbert, new species.

Body moderately compressed, not elevated, the depth at front of dorsal not more than  $\frac{1}{2}$  greater than the oblique anterior margin of dorsal fin, less than distance from the nostril to first gill slit. Head depressed, the snout flattened, long and narrow, acute; length of snout beyond mouth  $\frac{1}{2}$  to  $\frac{1}{10}$  greater than distance between angles of mouth in all but one (the largest) of our specimens, in which it is slightly less than width of mouth;  $\frac{2}{3}$  to  $\frac{1}{2}$  greater than distance from tip of lower jaw to a line connecting angles of mouth;  $\frac{1}{2}$  to  $\frac{1}{4}$  greater than width of snout opposite outer angle of nostrils. Interorbital width equaling distance from tip of snout to front of eye in young, to middle or posterior border of eye in older individuals; less than  $\frac{1}{2}$  distance to first gill opening. Middle of eye nearer nostril than angle of mouth by  $\frac{1}{2}$  to  $\frac{1}{4}$  its diameter; distance from eye to nostril  $\frac{1}{2}$  or slightly more than  $\frac{1}{2}$  distance from nostril to tip of snout; middle of nostrils much nearer front of mouth than tip of snout; nasal flap with a very narrow, short, acute lobe, placed at end of inner third of flap; outer angle of nostrils nearly at margin of snout, the inner angles separated by a distance equaling or slightly exceeding that between inner angle of nostril and back of eye. Lips very little developed, the lower entirely concealed in closed mouth, the upper visible as a very short fold. Teeth in lower jaw narrow, erect, serrulate on both margins, more coarsely so toward base; the serration more conspicuous in our smallest specimens (450 mm.), and is obsolescent on some of the teeth in adults; teeth in upper jaw broadly triangular, in front of jaw narrower and erect, those in sides of jaw growing at once broader and more and more oblique; the lateral teeth with a strong notch on the outer side; both margins strongly serrate, the serrations increasing toward base, one or more of those below notch sometimes enlarged and cusp-like in adults; teeth about  $\frac{1}{2}$ . Conspicuous areas of large and of small pores on underside of head. Gill openings of moderate width, the longest equaling distance between eye and nostril, the fifth much shortened, about  $\frac{1}{2}$  length of first. Eye small, equaling length of nasal opening,  $1\frac{1}{2}$  to 2 in middle gill slit. Pectoral short and broad, the posterior margin not strongly incurved; tip of fin extending to a vertical intersecting dorsal base at origin of its posterior third or fourth; anterior margin of pectoral 3 times length of inner or posterior margin, the latter less than width of base; first dorsal beginning behind a vertical from axil of pectoral a distance about equaling that which separates eye from nostril; free margin of fin gently concave, the anterior angle extending to a point midway between base and tip of posterior lobe, when the fin is depressed; base of first dorsal  $2\frac{1}{2}$  to  $2\frac{3}{4}$  in interspace between dorsals; base of second dorsal 7 in interspace between dorsals,  $2\frac{1}{2}$  in its distance from anterior margin of pit; origin of second dorsal falling over or behind middle of anal base, the fin but slightly concave, with rounded anterior angle, its posterior angle much produced, the posterior margin exceeding base of fin, which about equals length of anterior margin; anal inserted more ante-

riorly than second dorsal, its base longer, its margin much more deeply concave, the length of base contained about  $1\frac{1}{2}$  times in its distance from lower caudal lobe; lower caudal pit in advance of the upper; caudal broad throughout, the lower lobe not falcate, slightly less ( $\frac{1}{10}$  to  $\frac{1}{4}$ ) than  $\frac{1}{2}$  length of upper lobe, which is about  $\frac{1}{4}$  in total length. Shagreen coarse. Color varying from light to dark gray above, the belly and lower part of sides whitish; fins all dusky or grayish, the caudal often with a blackish border; pectoral with or without a black tip, the latter when present not as conspicuous as in *C. ethalorus*, usually not extended into inner face of fin. A specimen 730 mm. long has the claspers undeveloped, extending slightly beyond margin of ventrals. Another specimen, 850 mm. long, has the claspers fully developed, extending beyond the margin of the ventrals for a distance of 50 mm. Strongly resembling *C. ethalorus*, with which it is associated in the Bay of Panama. It is distinguishable at sight by the narrower gill slits, broader and less falcate fins, and by the much less conspicuous black tips to the pectorals. The dentition is very dissimilar in the two, and makes it necessary to arrange them in different subgenera. Abundant at Panama, where numerous specimens were secured. (Gilbert.) (*κερδάλη*, wary, fox-like.)

*Carcharhinus*, sp. Indeser., JORDAN, Proc. U. S. Nat. Mus. 1885, 363.

*Carcharhinus cordale*, GILBERT, Fishes of Panama, MS. 1898, Panama. (Type, No. 11884, L. S. Jr. Univ. Mus. Coll. Dr. Gilbert.)

**Page 39.** *Carcharhinus nicaraguensis* or its marine original form was found in abundance in the Bay of Panama by Dr. Gilbert.

**Page 41.** After *Carcharhinus oxyrhynchus* add:

54(b). *CARCHARHINUS VELOX*, Gilbert, new species.

Preoral portion of snout slightly more than  $1\frac{1}{2}$  times width of mouth, 5 times distance between nostrils,  $1\frac{1}{2}$  times width of snout opposite outer angles of nostrils,  $1\frac{1}{2}$  times interorbital width,  $2\frac{1}{2}$  times distance from chin to line joining angles of mouth. Nostrils transverse in position, the inner angle nearer mouth than tip of snout by a distance slightly less than length of nostril. Front of eye equidistant from nostril and front of mouth, the middle of eye nearer angle of mouth than nostril; diameter of eye less than nostril, slightly more than  $\frac{1}{2}$  longest gill slit. Snout very porous. Folds at angle of mouth slightly longer than usual. Gill slits rather wide, the middle one  $1\frac{1}{2}$  times diameter of orbit. Teeth of lower jaw very narrow, erect, very minutely serrulate, appearing entire except under a lens. Teeth in upper jaw very oblique, wide at base, with a deep notch on outer margin, the terminal cusp rather narrowly triangular. Pectoral broadly falcate, the anterior margin convex, the distal edge concave, both angles rounded; tip of pectoral reaching a short distance beyond base of first dorsal; anterior margin of pectoral  $2\frac{3}{4}$  times the posterior (inner) margin, about  $1\frac{1}{2}$  times the distal edge; first dorsal inserted about the diameter of orbit behind a vertical from axil of pectoral, nearer pectoral, therefore, than ventral; anterior margin concave basally, convex on distal half, the anterior angle rounded; free margin concave, largely owing to the much pro-

duced acute posterior lobe; vertical height of fin exceeding length of base, the anterior lobe very high, extending beyond tip of posterior when the fin is declined, equaling  $\frac{3}{4}$  length of anterior margin of pectoral; posterior margin of dorsal  $3\frac{1}{2}$  in the anterior margin; base of first dorsal contained  $2\frac{1}{2}$  times in interspace between dorsals; base of second dorsal 6 times; margin of second dorsal gently concave; front margin low, the angle broadly rounded, barely reaching posterior end of base when fin is declined; posterior lobe much produced and acute, slightly longer than base of fin, the latter  $1\frac{1}{2}$  in the distance from its base to front of caudal pit; upper lobe of caudal  $3\frac{3}{8}$  in total length, the lower lobe  $2\frac{1}{2}$  in the upper; terminal lobe of caudal  $3\frac{3}{8}$  in the upper lobe; anal larger than second dorsal, higher, with deeply incurved margin, its base a little longer, its origin slightly in advance of that of second dorsal, the posterior insertions of the two fins nearly opposite; length of anal base  $1\frac{3}{8}$  in its distance from anterior edge of caudal pit. Color bluish above, whitish or grayish below; free margin of pectoral narrowly white, the anterior edge narrowly bordered with black, most evident when seen from the outer surface, the inner surface being dusky; first dorsal unmarked, the second dorsal with dusky anterior lobe; upper edge of caudal black, the lower margin faintly dusky; fins otherwise unmarked. A single specimen, a female 4 feet long, was procured from the Panama market. As preserved, it is partially skinned. The following measurements were taken when the specimen was intact, before preservation. Where not agreeing with dimensions given above, the latter will be found more reliable:

	Inches.
Tip of snout to insertion of dorsal.....	10 $\frac{1}{2}$
Base of first dorsal.....	4 $\frac{3}{8}$
Distance between dorsals.....	11
Base of second dorsal.....	1 $\frac{3}{8}$
From second dorsal to front of caudal pit.....	2 $\frac{3}{8}$
Front of caudal pit to tip of caudal.....	13 $\frac{3}{8}$
Tip of snout to axil of pectoral.....	15
Axil of pectoral to front of base of ventrals.....	11 $\frac{1}{8}$
Front of ventrals to front of anal.....	6 $\frac{1}{2}$
Front of anal to front of caudal pit.....	4 $\frac{1}{8}$
Girth at front of first dorsal.....	17 $\frac{3}{8}$

Distiguished from other known sharks of the Pacific coast of America by the excessively long, slender, acute snout, the slender body, and the very long caudal fin. Panama; only the type known. (*velox*, swift.)

*Carcharhinus velox*, GILBERT, Fishes of Panama, MS. 1898, Panama. (Type, No. 11893, L. S. Jr. Univ. Mus. Coll. Dr. Gilbert.)

**Page 42.** *Scoliodon longurio* has the teeth serrulate at base only. The base of the first dorsal is  $2\frac{1}{2}$  in the interspace between the dorsals.

**Page 44.** *Sphyrna tiburo* occur also in the Pacific. We have recently secured specimens at Mazatlan, where *S. tudes* and *S. zygaena* are also found.

**Page 47.** *Carcharias littoralis* reaches a length of 8 feet 7 inches. (Specimens from Beaufort, North Carolina. Coll. H. H. Brimley.)

**Page 49.** *Lamna cornubica*, the salmon shark, is abundant and destructive to salmon on the coasts of southern Alaska, especially about Kadiak, where it was seen by us.

**Page 53.** Under *Squalidae*, read ovoviviparous for "oviparous."

**Page 54.** *Squalus sucklii* has been but once recorded from Bering Sea. (Bering Island. Coll. Dr. L. Stejneger.) It is very abundant at Kadiak.

**Page 60.** *Pristis perrotteti* is not authentically known except from the rivers of Africa. Our west coast species is doubtless distinct and should stand as *Pristis zephyreus*. *Pristis pectinatus* occurs northward at least to Beaufort, North Carolina. (Specimen 12½ feet long. Coll. H. H. Brimley.)

**Page 61.** After the synonymy of *Pristis pectinatus* insert:

80(a). PRISTIS ZEPHYREUS, Jordan & Starks.

(PEZ DE ESPADA.)

Snout to nostrils, 3 in length to base of caudal; breadth of saw at anterior end between first 2 pairs of teeth  $\frac{1}{2}$  breadth of its base behind the last pairs; teeth on saw trenchant behind, arranged in 22 pairs; hinder teeth wide apart, the interspaces 5 times their base; posterior teeth turned slightly backward, a groove on their posterior edge; front teeth not quite  $\frac{1}{2}$  as long as the saw is broad at their base; distance between first and second tooth 3 times base of first. (Other specimens examined for us by Dr. G. W. Rogers show 18 to 21 pairs of teeth.) Eye equal to spiracle, contained 3 times in base of saw just behind last pair of teeth; width of mouth a little greater than base of saw; mouth with about 65 series of blunt teeth; slant height of pectoral in front a little more than half distance from tip of snout to mouth. Dorsals subequal; first dorsal inserted in advance of ventrals, about  $\frac{1}{2}$  its base over ventrals; caudal with a lower lobe, which is equal to slant height of pectoral; tail with a keel on side. Color plain olive gray above, light below. Measurements: Length 59 inches; caudal 7 inches; pectoral 7 inches; dorsal front 5½ inches; snout without nostril 11 inches. Type: A skin in L. S. Jr. Univ. Museum. Common in brackish waters at the mouth of the Rio Presidio, where 1 fine specimen was obtained. The species is also recorded (as *Pristis perrotteti*) by Dr. Gilbert from Mazatlan, and by Dr. Günther from Chiapas. Dr. Günther identifies this species with *Pristis perrotteti* described by Müller & Henle, from the Senegal River. In view of the great difference in the fauna of the Gulf of California from that of equatorial Africa this identification may be questioned, especially as there are several details in which the description of *P. perrotteti* differs from our specimen.

*Pristis zephyreus*, JORDAN & STARKS, Fishes of Sinaloa, 383, 1895, Mazatlan, Mexico. (Coll. Hopkins Exped. to Sinaloa.)

h of base,  
when the  
oral; pos-  
rst dorsal  
d dorsal 6;  
low, the  
hen fin is  
nger than  
of caudal  
2½ in the  
rger than  
a little  
the pos-  
l base 1½  
h above,  
white, the  
hen seen  
orsal un-  
of caudal  
arked. A  
Panama  
measure-  
ervation.  
be found

Inches.  
..... 16½  
..... 4½  
..... 11  
..... 13  
..... 2½  
..... 13½  
..... 15  
..... 11½  
..... 6½  
..... 4¾  
..... 17½

America  
and the  
ift.)

No. 11893,

ly. The

recently  
are also

(Spec-

Page 62. After *Rhinobatus lentiginosus*, add:

81(a). RHINOBATUS STELLIO, Jordan & Rutter.

Disk triangular, its greatest width a little less than  $\frac{1}{2}$  the distance from snout to dorsal, and equal to distance from snout to a line connecting points of greatest width. Sides of disk straight, tip of snout rounded, posterior point of pectoral more broadly rounded than snout. Length of snout equal to, or a little less than,  $\frac{1}{2}$  greatest width of disk, equal to distance between outer points of anterior gill openings; interorbital width 4 to 4 $\frac{1}{2}$  in snout, a little less than length of eye and spiracle, but about equal to length of nostril; internasal width equal to orbit; spiracle  $\frac{1}{2}$  length of eye, a prominent curved papilla and a slight ridge in its posterior side. Anterior nasal valve with a long slender flap extending across the nostril; 3 broad flaps on posterior side. Rostral ridges separate for their entire length, width between them at base equal to width of spiracle. Mouth nearly straight, its width 2 $\frac{1}{2}$  in its distance from snout and equal to distance between inner folds on posterior side of spiracle. Eye 4 $\frac{1}{2}$  to 5 $\frac{1}{2}$  in snout. Width of body at axil of pectorals 1 $\frac{1}{2}$  in snout. Dorsal fins about equal in size and shape, the distance between them 2 $\frac{1}{2}$  times base of first, the distance between the origins of the two fins equal to snout and about equal to distance from axil of pectoral to origin of first dorsal. Sides of tail with a conspicuous fold. Skin above with a fine uniform shagreen, nearly smooth below except near margins of the disk. A series of very small spines above eye and spiracle, 1 or 2 minute spines on shoulder girdle; the largest spines of body situated along median line of back, extending beyond first dorsal; no spine on snout, but in 2 of the 3 specimens there is a pair of minute spineless plates near its tip. Color dusky brown above, about 7 faint dusky bars on the side of the tail behind first dorsal; uniform pale below; large translucent areas on each side of the snout; back with numerous small light spots, much smaller than pupil, arranged symmetrically but not in the same pattern on the 3 type specimens; 2 or 3 pairs between eyes, a few pairs behind eyes near median line, some below eye, where they approach nearest the margin of disk, usually 1 or 2 on median line, sometimes 2 are confluent, about 40 or 45 pairs in all; axil of pectoral in 1 specimen with a dusky blotch on upper side. This species is most nearly related to *Rhinobatus glaucostigma* of the Pacific coast, differing in having a narrow interorbital, narrower body behind disk, and in the very different color. The description is based on 3 specimens, each about 20 inches. Jamaica. (*stellio*, the starry one.)

*Rhinobatus stellio*, JORDAN & RUTTER, Proc. Ac. Nat. Sci. 1897, 91, Kingston, Jamaica.  
(Type, No. 11851, L. S. Jr. Univ. Mus. Coll. Joseph Sead Roberts.)

Page 66. To the synonymy of *Raja* add: *Cephaloetherus*, Rafinesque, Indice, 61, 1810 (*maculatus*).

The genus *Cephaloetherus*, Rafinesque, was, as Dr. Gill has shown (in lit.), probably based on a monstrous example of the genus *Raja*, in which the pectoral fins were not developed on the snout. It should be transferred to the synonymy of *Raja*, leaving *Myliobatis* as the generic name of the Eagle Rays.

Page 74. After *Raja equatorialis* add:

104(a). *RAJA ROSISPINIS*, Gill & Townsend.

Snout moderately produced, with a soft, moderately narrow, rostral cartilage and a bluntish tip; interorbital space nearly plane; snout with a number of plates having stellate bases about middle, and many smaller asperities, leaving only the borders of the pectorals and ventrals naked; larger spines with stellate bases are interspersed between the disk and the pectoral rays; back with sparse, coarse prickles; a row of about 26 thorn-like spines, with radiating ridges, extends from the interhumeral area to the dorsal fins; 2 spines on each shoulder, 1 spine above antocular region, another above postocular region, and another behind it about  $\frac{1}{2}$  the distance; skeleton soft. Bering Sea; only the type known. (*roseus*, rosy; *spinus*, spine.)

*Raja rosispinis*, GILL & TOWNSEND, Proc. Biol. Soc. Wash., xi, 1897 (Sept. 17, 1897), 231, Bering Sea. (Type, No. 48762, U. S. Nat. Mus. Coll. Albatross.)

*Raja obtusa*,\* GILL & TOWNSEND, Proc. Biol. Soc. Wash., xi, 1897 (Sept. 17, 1897), 231, Bering Sea. (Type, No. 48763, U. S. Nat. Mus. Coll. Albatross.)

104(b). *RAJA INTERRUPTA*, Gill & Townsend.

Snout moderately produced, with a very soft attenuated rostral cartilage and a blunt tip; interorbital space concave; mouth small; the width equal to  $\frac{1}{2}$  preoral area; entire back covered with very small embedded spines, extending nearly uniformly over the disk and snout, leaving only the tip of the latter naked; a row of compressed, acutely curved, smooth spines along middle of back, extending from the interhumeral region to dorsal, but interrupted along the posterior half of disk, where the spines are absent or obsolete; about 4 spines are in the anterior portion and the series recommences on a line with the emargination of the disk; a single spine on each shoulder and occasionally a rudimentary second; no specialized supraorbital spines. Bering Sea; only the type known. (*interruptus*, interrupted.)

*Raja interrupta*, GILL & TOWNSEND, Proc. Biol. Soc. Wash., xi, 1897 (Sept. 17, 1897), 232, Bering Sea. (Type, No. 48760, U. S. Nat. Mus. Coll. Albatross.)

Page 75. *Raja aleutica* and *Raja abyssicola* were described by Gilbert (not Gilbert & Thoburn) in Rept. U. S. Fish. Comm. 1893 (Dec. 9, 1896), 396 and 397, pls. 20 and 21.

\* The following is the original description of this nominal species:

Snout not at all produced, but very bluntly rounded; interorbital space narrow; mouth small, rectilinear; minute distant prickles on the snout, the anterior portion of disk and interorbital area, as well as in a broad median band extending on tail to dorsal and commencing at the interhumeral area; a row of scarcely enlarged acute spines above the eye; an uninterrupted row of unguiform spines with smooth bases extending from the interhumeral area to dorsal fin; 2 similar spines on each shoulder. Bering Sea; only the type known. (Gill and Townsend.) To which we add: Spines in longitudinal series 23 to 25; width of mouth  $\frac{1}{2}$  in preoral area; width of disk  $1\frac{1}{2}$  times its length; tail a little longer than disk; interorbital width 3 in snout; snout from eye  $3\frac{1}{2}$  in disk to end of base of ventrals. Color plain brown, rather pale. One specimen 11 inches long, a very young male in very bad condition. Evidently the young of *R. rosispinis*.



Page 78. After *Narcine brasiliensis* insert:

112(a). NARCINE ENTEMEDOR, Jordan & Starks.

(ENTEMEDOR.)

Snout  $3\frac{1}{2}$  in length of disk; preocular part of snout equals preoral; interocular space in snout  $1\frac{1}{2}$ ; width of mouth  $2\frac{1}{2}$ . Eye much smaller than spiracle; spiracles edged with small tubercles. Length of disk equal to its width; disk equal to length of tail, without caudal fin; tail with a loose fold of skin on each side. First and second dorsals equal, rounded behind; ventrals large, ending midway between posterior edge of disk and caudal fin. Color pale olive brown, a little clouded with darker; second dorsal edged with pale; dots on head dusky. Two specimens taken in the estuary at Mazatlan, and a third procured by Mr. James A. Richardson in the harbor of La Paz. Specimens had also been obtained by Dr. Gilbert, at Panama, in 1883, but having been destroyed by fire, the species has remained undescribed until recently. Length of largest specimen 20 inches. (The Spanish name *Entemedor* seems to be equivalent to *Intimidator*.)

*Narcine entemedor*, JORDAN & STARKS, Fishes of Sinaloa, 386, 1895, Mazatlan, Mexico. (Type, No. 1699, L. S. Jr. Univ. Mus. Coll. Hopkins Exped. to Sinaloa.)

Page 81. After *Urolophus nebulosus* add:

115(a). UROLOPHUS UMBRIFER, Jordan & Starks.

Disk round, not wider than long, its length greater than tail; snout pointed, not exerted. Snout from eye  $4\frac{1}{2}$  in disk; eyes equal to spiracles; mouth 2 in distance to tip of snout; caudal spine inserted in front of middle of tail; skin perfectly smooth. Color brown above, with blackish cross shades or bars, radiating from the shoulder; a dark band behind eyes, and 1 from eyes; caudal fin dark. Mazatlan. One adult female specimen, the uterus containing 4 young. Occasionally taken with *Urolophus mundus*, but much less common. This is probably not identical with Garman's *Urolophus nebulosus*, being perfectly smooth and different in color. (*umbra*, shade; *fero*, I bear.)

*Urolophus umbrifer*, JORDAN & STARKS, Fishes of Sinaloa, 389, 1895, Mazatlan, Mexico. (Coll. Hopkins Exped. to Sinaloa.)

Page 82. *Urolophus asterias*, Jordan & Gilbert, is identical with *Urolophus mundus* (Gill), as is shown by specimens recently collected by D. Gilbert at Panama.

120(a). UROLOPHUS ROGERSI, Jordan & Starks.

Disk broader than long by a distance  $2\frac{1}{2}$  times the interorbital width; anterior margins of disk nearly straight, the tip of snout projecting; snout from eye  $3\frac{1}{2}$  in length of disk; eye little smaller than spiracles; width of mouth  $2\frac{1}{2}$  times in preoral part of snout; caudal spine inserted in front of middle of tail; skin with minute prickles on margin of pectorals and on middle of back, leaving smooth areas near middle of pectorals and

over branchial arches; 16 to 20 large spinules along median line of back and tail. Color plain brown; caudal fin darker, edged with white. This species differs from *Urolophus asterias* in having a wider disk, more acute snout, much smaller prickles, and fewer spinules on back and tail. Mazatlan. Three specimens obtained in the Astillero, the longest 18 inches in entire length. (This species is named for Dr. George Warren Rogers, a scholarly physician, native of Vermont, but long resident in Mazatlan.)

*Urolophus rogersi*, JORDAN & STARKS, Fishes of Sinaloa, 388, 1895, Mazatlan, Mexico. (Type, No. 1700, L. S. Jr. Univ. Mus. Coll. Hopkins Exped. to Sinaloa.)

Page 88. In the key at top of page for *Rhinopterinae* read *Myliobatinae*.

132. AETOBATUS LATICEPS (Gill).

This species is probably not different from *A. narinari* and may be omitted. We find no differences between specimens from Mazatlan and the West Indies. The following description is based upon Mazatlan specimens:

Length of disc  $1\frac{1}{2}$  in width; proximal  $\frac{1}{2}$  of anterior margin of pectoral fins straight, distal  $\frac{1}{2}$  convex; posterior margin concave, the end of each ray forming a small scallop; lateral angle sharp. Snout forming an angle, from its tip to division of nasal lobes,  $1\frac{1}{2}$  times breadth of head; width of snout  $1\frac{1}{2}$  times distance from its tip to the division of nasal lobes; nasal lobes projecting back over the mouth; width of mouth  $1\frac{1}{2}$  its distance to tip of snout; numerous blunt buccal papillae around upper dental plate and on ridge between nostrils; interorbital  $4\frac{1}{2}$  in disk; eyes smaller than spiracles, which are as long as base of dorsal. Ventrals well rounded,  $3\frac{3}{4}$  in length of disk; tail  $3\frac{1}{2}$  times disk. First caudal spine equals base of dorsal, which is  $\frac{1}{2}$  second spine. Color bluish black with many round yellowish spots scattered equally over the back and ventral fins; spots about as large as eye on back, smaller on head, sometimes two spots run together forming an elliptical spot, about 16 spots from eye along anterior margin of pectoral to lateral angle; posterior margin of pectoral very narrowly margined with white; ventral side pearly white. From the description of *Actobatus laticeps* this species differs in the following respects: Disk not so broad; tail not so long; width of head and snout less; ventrals not truncated behind; pectorals not margined with blackish; spots on ventrals not assuming the form of ocelli. (Jordan.)

Page 87. For the description of *Picroplatea crebrispunctata* in text substitute the following:

Width of disk twice length to posterior end of anal slit; snout forming a regular curve from a little in front of middle of pectorals, a very small blunt projection at tip; anterior margin of disk convex near snout and lateral angles, pectorals concave medially; posterior margin weakly convex; posterior angle broadly rounded; lateral angle sharply rounded; distance from snout to a line drawn through lateral angles,  $2\frac{1}{2}$  times in distance to tip of tail. Interorbital a little wider than its distance to tip of snout; eyes twice spiracles; mouth equals snout,  $6\frac{1}{2}$  in disk. Tail rat-

like, with a scarcely perceptible fold of skin on its dorsal side. Ground color olive brown, everywhere with small dark points, not so close set as in *Pteroplatea rara*, indistinct grayish spots,  $\frac{1}{2}$  as large as iris, scattered over the body among the dark points, these spots more distinct on anterior edge of disk; tail mottled with darker; lower parts light. Markings nowhere so distinct as in *P. rara*. Very common on sandy shores everywhere about Mazatlan, from which locality it was originally described; also taken by Dr. Gilbert.

**Page 87.** After *Pteroplatea marmorata* add:

**130(a).** PTEROPLATEA RAYA, Jordan & Starks.

(MANTARIA COLORADA.)

Length of disk  $1\frac{1}{2}$  width; snout forming an angle which is almost a right angle; pectorals slightly concave medially; posterior margin of disk weakly convex; posterior angle not broadly rounded, but curved in somewhat suddenly; lateral angles acute. A line drawn through lateral angles would bisect a line from snout to tip of tail. Interorbital  $1\frac{1}{2}$  in snout; eye  $1\frac{1}{2}$  in spiracles; mouth 7 in disk,  $1\frac{1}{2}$  in snout; tail straight and slender, with a very slight fold on dorsal side. Ground color light olive brown, thickly set with sharp-cut black points; conspicuous gray or white spots,  $\frac{1}{2}$  as large as iris, scattered over the body, around which the black spots form rings; brighter yellowish spots and half-spots around anterior edge of disk; tail mottled above with darker; lower parts chiefly light orange red or rust-colored in life. All the markings are very distinct and clear cut, the reddish of the belly conspicuous. Mazatlan. One specimen 12 inches long. (*ravus*, reddish.)

*Pteroplatea raya*, JORDAN & STARKS, Fishes of Sinaloa, 390, 1895, Mazatlan, Mexico. (Type, No. 1587, L. S. Jr. Univ. Mus. Coll. Hopkins Exped. to Sinaloa.)

**Page 90.** After *Myliobatis californicus* add:

**134(a).** MYLIOBATIS ASPERRIMUS, Gilbert, new species.

Upper surface of head and body, excepting the snout, an area on outer side of spiracle, the pectoral margin and its posterior angle, and the ventral fins thickly covered with minute, usually stellate prickles, of uniform size, most numerous on median portions of head and back; those on basal  $\frac{1}{2}$  or  $\frac{2}{3}$  of pectoral least crowded and arranged in definite longitudinal series, corresponding with the muscle bands; tail very rough throughout, covered with similar stellate prickles and also crossed by numerous narrow grooves, or indented lines, mostly convex forward, somewhat irregular in position and direction, and not corresponding on the two sides. In the type they follow at an average interval of about 10 mm. Lower side of disk mostly smooth, with some prickles on the basal part of pectorals anteriorly, arranged in lengthwise series, and other patches on lower side of head, belly, and base of ventrals. Color dusky brown above, the anterior portion of pectorals with 8 to 10 narrow transverse bars of bluish

white, most of which break up into series of spots toward outer margin of disk, the posterior ones also breaking up toward middle line; the bars and spots fainter anteriorly, becoming whiter and more intense posteriorly; toward outer angles of disk the bars sometimes separated by intermediate series of ligate round spots, the bars usually failing to meet across the back; posterior portion of disk including base of tail and upper surface of ventrals covered with round white spots not much larger than pupil, some of those immediately succeeding the bars showing a transverse serial arrangement; top of head with one or more pairs of indistinct light spots; margin of snout and of pectorals blackish; spiracular border black; dorsal with a black blotch posteriorly; underside of head and disk bright white; proximal portion of tail blackish above, lighter below, the entire tail becoming black more posteriorly.

Dimensions of type specimen.

	Millimeters.
Length of disk to front of anus.....	272
Length of disk to posterior edge of pectorals.....	338
Width of disk.....	345
Length of tail (not perfect).....	1,215
Greater width of head at origin of pectorals.....	79
Width of cranium between orbits.....	45
Width of snout opposite front of eye.....	55
Tip of snout to middle of nasal flap.....	60
Length of nasal flap.....	26
Greatest width of nasal flap.....	35
Diameter of iris.....	10½
Width of mouth.....	33
Distance between anterior gill openings.....	75
Distance between posterior gill openings.....	45
Distance from anterior to posterior gill openings.....	45
Length of spiracle.....	26
Length of fontanelle.....	60
Greatest width of fontanelle (at anterior end).....	23

Rostrofrontal fontanel scarcely constricted anteriorly, the bounding ridges diverging abruptly at their anterior ends. Nasal flap with a shallow median notch, covering the mouth except the median portion of lower dental plate; posterior margin coarsely fringed. Teeth in each jaw in 1 broad median row, and 3 lateral rows, those of median row about 5 times as broad as long anteroposteriorly.

One specimen, a male, with undeveloped claspers which do not nearly reach edge of ventrals, from Panama. (Gilbert.) (*asperrimus*, very rough.)

*Myliobatis asperrimus*, GILBERT, Fishes of Panama, MS. 1898. (Type, No. 11805, L. S. Jr. Univ. Mus. Coll. Dr. Gilbert.)

134(b). MYLIOBATIS GOODEI, Garman.

Disk about  $\frac{3}{4}$  as long as broad; lateral angles acute, bluntly rounded at the apices; posterior angles of pectorals nearly right; snout very broad, short, with a slight prominence in front; the fin, or flange, beneath the eye at the side of the head is very wide, much wider than in either *M. freminvillei* or *M. californicus*; eye very small, without a prominence above in either

male or female (immature specimens); tail less than 2 and more than  $1\frac{1}{2}$  times length of the disk; dorsal fin smaller than that of *freminvillei*; teeth in 7 series, much shorter and narrower than those of *freminvillei*, third row about 2 and middle row about 4 times as wide as long. Body smooth. Entire length 29 inches; snout to end of ventrals 11.5, vent to end of tail 18.5, and width of disk 17.5 inches. Olivaceous, darker on the center; white below. The Museum of Comparative Zoölogy has a large specimen which agrees well with this description. Compared with *M. freminvillei*, this species has very small eyes, the pectoral below the orbit is wider than the eyeball, and the fin in front of the skull is but little wider than at its sides. In *freminvillei* the eyeball is twice as wide as the fin beneath it, and the fin in front of the skull is much wider than below the eye. Comparing specimens of about the same size, of the same sex, of *freminvillei*, *californicus*, and *goodei*, the latter is readily distinguished from the former two by the broad flange at the side of the head, the small eyes, the small teeth, and the broader lateral angles of the pectorals. Central America. (Garman); probably on the Atlantic Coast. (Named for George Brown Goode.)

*Myliobatis goodei*, GARMAN, Proc. U. S. Nat. Mus. 1885, 39. Central America. (Types, Nos. 9524 male, and 9529 female.)

**Page 91.** Family XXVIII should stand as *Aodontidae*, the name *Mantidae* being used for a family of *Orthoptera*.

**Page 92.** After *Aodon hypostomus* insert:

58 (a). CERATOBATIS, Boulenger.

*Ceratobatis*, BOULENGER, Ann. Mag. Nat. Hist., ser. 6, vol. xx, August, 1897, 227 (*robertsii*).

Characters of *Dicerobatis*, Blainville, but the teeth restricted to the upper jaw. (*Κεράς*, horn; *Βαρίς*, ray.)

138 (a). CERATOBATIS ROBERTSII, Boulenger.

Band of teeth occupying only  $\frac{1}{3}$  width of mouth, its width 10 times in its length; teeth tessellated, hexagonal, 2 to 3 times as broad as long, rugose with numerous obtuse ridges; mouth inferior, wide. Pupil vertically elliptic. Body smooth; pectoral fins with nearly straight, slightly convex anterior and slightly concave posterior border; cephalic fins measuring a little less than width of mouth; spiracles behind the eyes; space between last branchial clefts  $\frac{1}{2}$  that between first; dorsal fin between the ventrals; tail slender, without spine, nearly twice length of body.

	Millimeters.
Length of disk, without cephalic appendages.....	350
Width of disk.....	780
Cephalic fin.....	90
Width of mouth.....	105
Diameter of eye.....	12
Ventral fin.....	70
Tail.....	620

Black above, white beneath. Jamaica. One specimen known. This ray grows to a very great size, but specimens are almost impossible to obtain,

owing to the superstitious fear of the fishermen. (Named for Rev. Joseph Seed Roberts.)

*Ceratobatis robertsi*, BOULENGER, Ann. Mag. Nat. Hist., ser. 6, vol. xx, August, 1897, 227, Jamaica. (Type in British Mus. Coll. J. S. Roberts.)

**Page 105.** There is no truth in the statement that *Acipenser medirostris* is poisonous. It is a good food-fish, and on the coast of Washington it is somewhat abundant.

**Page 116.**

#### ADDITIONAL NOTES ON TACHYSURINÆ.

In the text of this work, pages 116 to 133, the descriptions of the species of *Tachysurinæ* are for the most part too brief to render certain the discrimination of species. The following additional descriptions of these species will be found useful as supplementary to those given in the text. A slight change in the arrangement of the genera has been found desirable, and 3 new species are added.

#### 70. SCIADEICHTHYS, Bleeker.

Dorsal shield much enlarged, formed like an armorial shield; teeth on palate villiform; posterior nasal openings not connected by membrane; band of palatine teeth extended backward.

#### 165. SCIADEICHTHYS TROSCHELI (Gill).

Head  $3\frac{1}{2}$  ( $3\frac{1}{2}$  in total with caudal); width of head  $4\frac{1}{2}$  ( $4\frac{1}{2}$  in total); depth 5 ( $6\frac{1}{2}$ ). D. I, 7; P. I, 12; A. 18. Body comparatively robust, broad anteriorly; head not much depressed, broader than high; eye moderate, 7 to 8 times in length of head; width of interorbital space  $1\frac{1}{2}$ ; breadth of mouth  $1\frac{1}{2}$ ; length of snout  $3\frac{1}{2}$ . Teeth all villiform; band of vomerine teeth simple, trapezoidal, quadrangular, longer than broad, without division on median line; band of palatine teeth very large, each separated in young specimens from the vomerine band by a narrow toothless line; in old specimens the vomerine and palatine bands are wholly confluent; each palatine band with a narrow backward prolongation on the median line; band of premaxillary teeth broad, about six times as long as wide; lower jaw included. Maxillary barbel nearly or quite reaching gill opening; outer mental barbels about  $\frac{2}{3}$  head, the inner nearly  $\frac{1}{2}$ . Dorsal shield much larger than in most species, shaped like an armorial shield, its posterior margin concave; its anterior end acute, wedged into a deep emargination of the occipital process, the two becoming coossified with age; length of antedorsal plate on the median line 5 to 6 in head, a little more than its width; occipital process short and broad, much broader than long, its median line with a broad keel, its edges nearly straight. Shields all coarsely granular, the granulations anteriorly forming radiating striæ. Fontanel large, claviform, broadest posteriorly, its posterior end about midway between tip of snout and front of dorsal, its greatest

breadth about equal to the diameter of the eye, and  $\frac{1}{2}$  its length, a short groove extending backward from its obtuse tip; sides of fontanel bony and granulated for its whole length, the granules extending forward to opposite nostrils. Dorsal spine strong,  $1\frac{1}{2}$  in head, moderately compressed; pectoral spine  $1\frac{1}{2}$  in head. Axillary pore obsolete. Humeral process coarse, granular, broad, nearly  $\frac{1}{2}$  length of pectoral spine; base of adipose fin scarcely  $\frac{1}{2}$  length of anal, its posterior margin little free; caudal deeply lunate, small, its upper lobe slightly the longer and narrower,  $1\frac{1}{2}$  in head; ventrals not quite reaching anal; vent much nearer base of ventrals than anal. Dark brown, with strong bronze luster above, white below; dorsal dusky, especially above; pectorals blackish; anal dark; caudal rather pale; ventrals usually dark toward the tip, their inner side pale; maxillary barbel dusky; mental barbels pale. This species is not rare along the Pacific coast of tropical America, specimens having been observed at Mazatlan, Punta Arenas, and Panama.

*Sciades troscheli*, GILL, Proc. Ac. Nat. Sci. Phila. 1863, 171, Panama. (Coll. Capt. Dow.)  
*Arius brandtii*, JORDAN & GILBERT, Bull. U. S. Fish Comm., 11, 1882, 39; description from 28230, U. S. Nat. Mus., 24 inches in length.

The following is the original description of *Sciades troscheli*, Gill: "Dorsal I, 7; anal 16; caudal 11, I, 6; 7, I, 11. The greatest height is contained about  $4\frac{1}{2}$  times in the length to the base of the caudal fin, and  $5\frac{1}{2}$  times in the total. The caudal peduncle, behind the anal, equals the interval between the snout and the eye, and its least height that between the center of the anterior nostril and the eye. The head in front and on the sides is smooth, and a smooth, oblong, triangular area extends nearly to the vertical from the upper angle of the preoperculum; a triangular area on each side is incurved externally to the narrow anterior extremity, and covered with white pisiform granulations. The dorsal buckler is a pentagon, with a semicircular excavation behind and with its surface rugose. The head enters 3 times in the length before the end of the anal fin and more than 4 times in the total; its width equals the interval between the snout and upper angle of preoperculum, and the interocular area equals  $\frac{1}{2}$  the head's length. The eye is elliptical, and its diameter is contained  $6\frac{1}{2}$  times in the head's length. The distance of the posterior nostril from it equals a diameter. The maxillary barbels extend to about the middle of the pectoral; the outer mental to its base, and the inner mental are  $\frac{2}{3}$  as long as the outer. There are 3 villiform patches on the palate which are almost contiguous, and together describe an arch in front; the median patch is small, rather transverse, and widest toward the front; the outer are oblong, subtriangular. The band of the upper jaw is nearly uniform and quite wide; the lower, interrupted at the symphysis, is nearly  $\frac{1}{2}$  as wide as the upper, and is narrowed toward its ends. The dorsal spine enters  $1\frac{1}{2}$  times in the head's length; has in front, first, minute teeth pointed downward, and then a row of small pisiform tubercles; teeth pointed downward on its hinder border. The first ray is little higher than the spine. The anal commences at a distance from the snout  $3\frac{1}{2}$  times as great as that from the base of the caudal

fin; its length enters  $6\frac{1}{2}$  in the length, exclusive of the caudal, and when bent back it reaches to the supernumerary caudal rays; the greatest height nearly equals the length. The pectoral fins extend rather beyond the base of the dorsal and exceed  $\frac{1}{2}$  of the length, exclusive of the caudal; the spine equals that of the dorsal. The ventrals are inserted midway between the base of the pectoral spines and the axil of the anal, and extend to the origin of the anal. The fins are almost blackish." A single specimen is in the collection of Captain Dow from Panama. The type of *Sciades troscheli* is now lost. At our request, Dr. Gill has again considered this description, in connection with the species now known from the coast. He is positive that his type of *troscheli* had the large dorsal shield characteristic of *brandtii*. Apparently Dr. Eigenmann is right in regarding *troscheli* and *brandtii* as identical.

160. SCIADEICHTHYS EMPHYSETUS (Müller & Troschel).

Head  $3\frac{1}{2}$ ; depth 6. D. I, 7; A. 18. Closely related to *Sciadeichthys troscheli*. Depth little greater than the width. Profile straight, less steep than in *S. troscheli*. Depth of the head  $1\frac{1}{2}$  in its length, its width  $1\frac{1}{2}$ . Top of the head sparsely and coarsely granular, the granulation extending forward only to middle of cheek; fontanel bordered anteriorly by smooth ridges; occipital process coarsely and closely granular, without a prominent keel, its margins convex, its tips emarginate, not coossified with the dorsal plate; dorsal plate shield-shaped, not keeled, its surface irregularly pitted, its margin more finely graven, its length about  $3\frac{1}{2}$  in the head. In the specimen examined the dorsal plate seems to have been at some time slightly broken in front, a small, narrow, sharp process of the occipital process fitting into the split. Eye small, 3 in the snout, 11 in the head, 5 in the interocular. Maxillary barbels flattened, reaching to below middle or end of dorsal fin, postmentals not quite to base of pectorals. Upper jaw slightly projecting; all the teeth minute, villiform, the vomerine patch emarginate in front and behind, joined to the subtriangular palatine patches; pterygoid patches long-elliptical. Gill membrane with a narrower free margin than in *troscheli*. Distance of dorsal fin from tip of snout  $2\frac{1}{2}$  in the length, the dorsal spine  $1\frac{1}{2}$  in the head, granular in front, recurved teeth on its inner margin. Distance between the dorsal and adipose fins  $3\frac{1}{2}$  in the length; adipose fin about as long as the dorsal fin. Caudal deeply forked, the upper lobe longer,  $3\frac{1}{2}$  in the length; anal little longer than high; ventrals reaching nearly to the anal, about 2 in the head; pectoral spine  $1\frac{1}{2}$  in the head, its outer margin granular, the inner rather finely toothed. The skin on the dorsal surface of the head and humeral region finely reticulate with mucous canals. Yellowish brown, lighter below, the fins yellowish, finely punctulate. One specimen 0.51 m. Surinam. (Eigenmann & Eigenmann, Nematog-nathi, 53.)



167. *SCIADEICHTHYS TEMMINCKIANUS* (Cuvier & Valenciennes), text, p. 122.

168. *SCIADEICHTHYS FLAVESCENS* (Cuvier & Valenciennes), text, p. 123.

169. *SCIADICHTHYS MESOPS* (Cuvier & Valenciennes).

170. *SCIADEICHTHYS PROOPS* (Cuvier & Valenciennes).

Head 4 to 4½; depth 7. D. I, 7; A. 18; eye 1¼ to 1½ in snout, 5½ to 8 in head, 1¾ to 2¾ in the interorbital, 2¼ to 3¾ in the interocular. Slender and elongate, broader than deep. Head depressed, its width 1½ in its length, its depth 2, width at mouth 2; anterior portion of the head flat above; top of the head, humeral process, and dorsal plate coarsely granular, the granules arranged in series along the fontanel. Occipital process mucronate, broader than long; dorsal plate large, butterfly-shaped. Opercle striate; fontanel 1½ times as long as the eye, its center in front of the middle of the eye, continued as a shallow groove. Jaws subequal; teeth all villiform, the intermaxillary band very wide and shallow; teeth on the roof of the mouth in 6 contiguous patches. Gill membranes meeting in an angle, forming a broad fold across the isthmus; gill rakers 5 + 10. Pectoral pores large; vertical series of pores. Distance of dorsal spine from the snout 2½ in the length; the dorsal spine granular in front, striate on the sides, weakly serrate behind, its length 1¼ to 1½ in head; space between dorsal and adipose fins 2¾ to 3 in length, the adipose fin little shorter than the dorsal, the posterior margin free. Caudal deeply forked, its upper lobe longer, 4 to 4½ in the length; anal emarginate, as high as long, 2 to 2½ in head; ventrals 2 in head; pectoral spine roughened or granular in front, serrated behind, 1½ to 1¾ in head. Plumbeous above, with blue luster, white below; maxillary barbels dark, the mental barbels white; fins all more or less dotted with brown. Five specimens 0.25 to 0.46 m. long. Pernambuco. (Hartt & Fletcher.) Northern coast of South America to Pernambuco. (Eigenmann & Eigenmann, *Nematognathi*, 57.)

171. *SCIADEICHTHYS FASSANY* (Cuvier & Valenciennes), text, p. 124.

172. *SCIADEICHTHYS ALBICANS* (Cuvier & Valenciennes), text, p. 124.

#### 71. *SELENASPIS*, Bleeker.

Dorsal shield much enlarged, truncate before, in the adult; palatine teeth villiform, the patch extended backward in the adult; posterior nasal openings connected by membrane.

173. *SELENASPIS HERZBERGII* (Bleeker).

Head 3¾ to 3½; depth 5 to 6. D. I, 7; A. 18; eye 1¼ to 2½ in snout, 5½ to 8 in head, 2¾ to 4 in interocular. Elongate, the width as great or greater than the depth. Width of the head 1¼ to 1½ in its length, at the angle of the mouth about 2; depth 1¾ to 1½ in its length. Humeral process, dorsal plate, top of head to between the eyes, granular. Occipital

process wider than long, scarcely keeled. Fontanel not continued behind the eyes, and without backward projecting groove; posterior nostrils connected by a membrane. Barbels flattish, those of the maxillary reaching to near the ventrals, to middle of pectorals in older individuals; postmental to or beyond base of pectoral, mental to gill opening. Teeth villiform; vomerine and palatine patches of about equal size and shape in the young; a separate patch behind the palatines is developed later. Gill membranes meeting in an angle, forming a fold across the isthmus; gill rakers 6 + 10. Distance of dorsal spine from snout  $2\frac{1}{2}$  to  $2\frac{3}{4}$  in the length; dorsal and pectoral spines subterete, the outer margins roughened, the sides striate; the dorsal spine slightly serrate behind, a little shorter than the pectoral spine,  $1\frac{1}{2}$  to  $1\frac{3}{4}$  in the head; pectoral spine strongly serrate behind; space between dorsal and adipose fins  $3\frac{1}{2}$  to 4 in the length; adipose fin as long as the dorsal; upper caudal lobe longer, about 4 in the length; anal as high as long, 2 in head; ventrals  $1\frac{1}{2}$  to 2 in head; pectoral pore minute; sides with vertical series of pores. Color plumbeous above, silvery on sides; fins dusky. The specimens examined measure from 0.14 to 0.32 meters. Para; Curuca; Bahia. (Eigenmann & Eigenmann, Nematognathi, 59.)

## 174. SELENASPIS DOWII (Gill).

Head  $3\frac{1}{2}$ ; depth 6. D. I, 7; A. 19; eye small, elliptical, 3 in snout, 12 in head, 6 in interocular. A narrow flap of skin across the snout connecting the posterior nasal openings. Width below the dorsal spine a little greater than the depth, less than the width at the humeral process, which equals the greatest width of the head measured at the opercles. Head depressed, its depth at base of occipital process  $1\frac{1}{2}$  in the greatest width, becoming gradually more depressed forward; width at angle of mouth  $1\frac{1}{2}$  in length of head, its greatest width about  $1\frac{1}{2}$  in its length; snout short, 4 in head. Top of head coarsely granular, the granules forming striae in front, vermiculations posteriorly or, in places, more or less regular striae. Occipital process truncate, its width at tip greater than its length, the dorsal plate large, saddle-shaped, its bony tubercles forming striae which are parallel with the strongly convex margin of the "saddle"; opercular bones granular striate, the humeral process with bony tubercles. Fontanel nearly obsolete, the granular bony surface being separated in front by thick skin, which covers an elongate area about 7 times longer than wide. Maxillary barbels reaching beyond humeral process; postmental barbels beyond gill opening, the mentals shorter. Upper jaw produced, equal to the short diameter of the eye. Teeth of the intermaxillaries in a villiform band which is narrowed in front, not produced backward to the angle of the mouth; vomer with a rather broad band confluent with the much wider subquadrate palatine patches which are produced backward in an angle; ovate patches on the pterygoids separate from the palatine teeth; teeth of the lower jaw in a comparatively shallow band, tapering very gradually to the angle of the mouth; the teeth of the jaws minute villiform, those of the palate and

vomer bluntly conical. Gill membranes broadly united, meeting in an angle, joined to the isthmus, but with a free margin; gill rakers 9 + 15. Distance of dorsal from end of snout  $2\frac{1}{2}$  in the length. Dorsal spine granular on sides and in front, about  $\frac{1}{2}$  the length of the head in height; distance of the adipose fin from the dorsal  $3\frac{1}{2}$  in the length, the height of the adipose fin about  $2\frac{1}{2}$  in its length, which is contained  $2\frac{3}{4}$  in the length of the head. Caudal deeply forked,  $5\frac{1}{2}$  in the length; anterior  $\frac{2}{3}$  of the anal strongly convex, the posterior  $\frac{1}{3}$  slightly emarginate, the highest ray about  $2\frac{1}{2}$  in head. Ventrals reaching to anal, about 2 in head, their distance behind the dorsal equal to the length of dorsal and  $\frac{1}{2}$  the dorsal plate; pectoral spine granulose on sides, the outer margin with a series of larger granules which become recurved notches toward the tip, the inner edge with recurved hooks, its height  $1\frac{3}{4}$  in the length of the head; a small pectoral pore; no evident series of vertical pores. Bluish gray above, becoming white below; the fins brownish with dots. Description from the type of *Arius alatus*, .68 m. long, from Panama; collected by Dr. Steindachner. (Eigenmann & Eigenmann, Nematognathi, 61.)

*Selencapsis dowii* is thus characterized by Jordan & Gilbert:\* Head 4 ( $4\frac{1}{2}$  with caudal); depth  $6\frac{1}{2}$  ( $7\frac{1}{2}$ ); width of head  $5\frac{1}{2}$ . D. I, 8; A. 4, 12. Length (29529, U. S. Nat. Mus.) 10 inches. Body elongate, narrow, and slender; the caudal peduncle  $1\frac{1}{2}$  in head. Head low and narrow, tapering anteriorly, the snout subtruncate. Eye small, 7 in head, placed rather high; interorbital space little arched, with ridges and depressions,  $2\frac{1}{2}$  in head; snout  $3\frac{1}{2}$  in head; breadth of mouth  $2\frac{1}{2}$  in head. Mouth moderate, with thinnish lips; teeth villiform, blunty; vomerine teeth forming 2 smallish, rounded patches, separated by a moderate interspace; each patch confluent with the neighboring palatine patch, which is rounded and rather large; the suture marked by a constriction; palatine bands without backward prolongation; premaxillary band of teeth broad. Barbels very long; maxillary barbel extending well beyond tip of pectoral fin; outer mental barbel reaching well past front of pectoral; inner  $2\frac{1}{2}$  in head. Dorsal shield comparatively large, not distinctly crescent-shaped, its divisions produced backward, their length about twice the length of the shield on the median line; anterior margin with 2 emarginations, the point fitting into an emargination of the occipital process; dorsal shield without keel. Occipital process very broad and short, its edges nearly straight, its breadth at base considerably greater than its length; its median line with a rather low keel. Fontanel broad and very short, ending obtusely at a point not far behind eye, the distance from this point to tip of snout  $1\frac{1}{2}$  in its distance from base of dorsal; each side of fontanel with a conspicuous smooth ridge, the 2 ridges converging anteriorly; shields of head rather finely granulated, few of the granulations forming lines, none of them extending farther forward than posterior margin of eye. Opercle striate. Gill membranes meeting below in a sharp angle, forming a rather broad fold across isthmus. Dorsal spine very short, its length a trifle less than pectoral spine,  $2\frac{1}{2}$  in head. Axillary pore obsolete. Humeral process granulated, rather narrowly triangular, a

\* Bull. U. S. Fish Comm., II, 1882, 50.

little less than  $\frac{1}{2}$  length of pectoral spine, which extends barely  $\frac{1}{2}$  the distance to the ventral fins; adipose fin long and low, very nearly or quite coterminous with the anal; caudal narrow, rather short, the upper lobe the longer,  $1\frac{1}{2}$  in head; anal rather low and short; ventrals short, the vent not far behind them. Color dusky above, pale below, the fins all more or less dusky; maxillary barbels dusky, others pale. A single young male was obtained at Panama.

71(a). **ASPITOR**, Jordan & Evermann, new genus.

*Aspistor*, JORDAN & EVERMANN, new genus (*luniscutis*).

This genus differs from *Selenaspis* in the presence of granular teeth on the palate and in the absence of a membranaceous flap connecting the posterior nostrils. (*ἀσπίστηρ*, a shielded warrior.)

175. **ASPITOR LUNISCUTIS** (Cuvier & Valenciennes).

Head  $3\frac{1}{2}$ ; depth  $5\frac{1}{2}$  to 6. D. I, 7; A. 16 to 19; eye 2 to 3 in snout, 6 to 9 in head, 3 to  $4\frac{1}{2}$  in the interocular. Body comparatively stout, the greatest width equaling the greatest depth. Head large, flattish above; profile descending; width of head  $1\frac{1}{2}$  in its length, width at the mouth 2 to  $2\frac{1}{2}$ , its depth at the base of the occipital process scarcely less than its greatest width; top of head coarsely granular in young, the granules becoming finer and more regularly arranged in the adult; opercles smooth; humeral process with radiating lines of granules. Occipital process variable in shape, broader than long, the posterior margin convex; dorsal plate variable in outline, rounded anteriorly, saddle-shaped, either broader than long or longer than broad; middle of the fontanel above the posterior margin of the eye; the fontanel divided into 3 by 2 bony ridges, the middle portion being more than  $\frac{1}{2}$  of its whole length. Sides of head with reticulating mucous canals. No skinny flap connecting the posterior nostrils. Maxillary barbels extending little beyond the base of the pectoral, or shorter; mental barbels short. Upper jaw little produced; teeth in the jaws rather large, conical; teeth of vomer and palatines finely granular, the vomerine patches separated from each other and from the palatine patches in the young, united and covering almost the entire roof of the mouth in the adult; the inner margins of the palatine patches approximated, sometimes a small elliptical patch of teeth between. Gill membranes forming a broad marginal flap across the isthmus. Gill rakers 3 to 4 + 7 to 9. Axillary pore minute or wanting; vertical series of pores present. Distance of dorsal from tip of snout  $2\frac{1}{2}$  in the length; the spine  $1\frac{1}{2}$  to  $1\frac{3}{4}$  in the head, granular in front, scarcely serrate behind; distance of adipose fin from the dorsal  $3\frac{1}{2}$  to 4 in the length, the adipose fin twice as long as high, adnate, as long as the dorsal fin; caudal forked, the upper lobe longer,  $4\frac{1}{2}$  to  $4\frac{1}{4}$  in the length; anal fin about as long as high,  $2\frac{1}{2}$  to  $2\frac{1}{4}$  in head; ventrals  $1\frac{1}{2}$  to 2 in head; pectoral spine stout,  $1\frac{1}{2}$  to  $1\frac{1}{4}$  in head, granular in front (serrate in the very young), striate on sides, serrate along inner margin. Color purplish brown above, sprinkled with brown dots below; fins about the color of the back. Numerous specimens

examined 0.11 to 0.44 m. long. Porto Alegre; Bahia; Nazareth, near Bahia; Rio Janeiro; Pará; Porto Seguro; São Muthcos; Cannavieras. (Eigenmann & Eigenmann, Nematognathi, 63.)

176. *SELENASPIS PARKERI* (Traill), text, p. 125.72. *NETUMA*, Bleeker.

Dorsal shield small, lunate; teeth on palate villiform, the patch on each side with a backward-extending process or angle.

Subgenus *NOTARIUS*, Gill.

Occipital process constricted at base.

177. *NETUMA GRANDICASSIS* (Cuvier & Valenciennes).

Head  $3\frac{3}{4}$  to  $3\frac{1}{2}$ ; depth  $5\frac{3}{4}$  to 6. D. I, 7; A. 18; eye 3 to  $3\frac{1}{2}$  in snout,  $8\frac{1}{2}$  to 10 in head, 4 to  $4\frac{1}{2}$  in interocular. Body cylindrical in front, tapering to a slender caudal peduncle. Head greatly depressed, profile almost straight, descending, the width of the head  $1\frac{1}{4}$  to  $1\frac{1}{2}$  in its length, its depth  $1\frac{1}{4}$  to 2 in its length. Occipital process with a deep constriction where it joins the occiput, shaped like a clover leaflet, much as *Felichthys panamensis*, sometimes broader than long, sometimes much longer than broad, sometimes keeled. Center of the fontanel over the middle of eye, the fontanel not continued backward as a groove; occipital process, top of head, and humeral process granular; interorbital region with 4 ridges, the inner ones bounding the fontanel, the outer ones running obliquely backward from near the posterior nasal opening. Maxillary barbels reaching to the base of the pectoral, mentals to gill opening, postmentals a little longer. Upper jaw projecting a diameter of the eye or more, the lip very wide, especially in front, making the nose pointed; teeth of both jaws rather large, those on the palate somewhat smaller; the depth of the intermaxillary band 7 to 9 in its width; the mandibular band very shallow; vomerine teeth none in 3 of the examples, a small patch on one side in another specimen, and a small patch on each side in another; palatine patches triangular, produced backward. Gill membranes meeting in an angle, forming a fold across the isthmus; gill rakers 6 + 10. Distance of dorsal spine from snout  $2\frac{1}{4}$  to  $2\frac{1}{2}$  in the length, the spine broken in the specimen studied. Distance of adipose fin from the dorsal  $3\frac{3}{4}$  to 4 in the length; adipose fin at least as long as the dorsal fin, adnate. Caudal fin forked, the upper lobe longer, about 5 in the length, the tips broken; anal fin apparently longer than high, but the rays are somewhat worn off; ventrals small; pectoral pore large, slit-like. Color light brown above, somewhat smutty below from the occurrence of minute scattered dots. We have examined 4 specimens from 0.23 m. to 0.33 m. long, collected by Agassiz & Bourget, Thayer Expedition, at Maranhao, and a fifth, 0.21 m. long, collected by Professor Agassiz at Bahia. (Eigenmann & Eigenmann, Nematognathi, 65.)

178. NETUMA STRICTICASSIS (Cuvier & Valenciennes), text, p. 126.

Subgenus NETUMA.

Occipital process not constricted.

179. NETUMA DUBIA (Bleeker), text, p. 126.

180. NETUMA KESSLERI (Steindachner).

Head  $3\frac{1}{2}$  (4 in total with caudal); depth 6 (7 in total); width of head 4. D. I, 7; A. IV, 13. Length (29252, U. S. Nat. Mus.) 14 inches. Body rather long and low; the head long, broad and much depressed, much broader than deep. Eye very small, about 10 in head, placed well above the mouth. Interorbital space 2 in head; snout  $3\frac{1}{2}$ ; breadth of mouth 2. Mouth large, with thickish lips, the upper jaw considerably projecting. Teeth all villiform, rather pointed. Vomerine patches rather large, roundish, usually fully confluent into a trapezoidal band, without division on the median line, and separated by a very narrow groove from the palatine bands. Palatine bands very large, broadly triangular, with a backward prolongation from the inner margin. (Teeth on vomer and palatines all forming one continuous band in old specimens, according to Steindachner.) Bands of teeth in jaws broad, the jaws strong. Barbels rather short and very slender, the maxillary barbels reaching little past base of pectoral; outer mental barbels about reaching gill opening; inner about as long as snout. Antedorsal shield short, crescent-shaped, rough, but without median keel. Occipital process long, narrowly triangular, its edges straight, its length  $\frac{1}{2}$  to  $\frac{1}{3}$  more than its width at base, its median line sharply keeled. Fontanel broad and shallow, its posterior end obtuse or almost truncate, its tip not prolonged in a groove, its edge bounded by a bony ridge, which is not granulated in front of middle of eye; end of fontanel about midway between tip of snout and front of dorsal, its greatest width about equal to length of eye. Shields of head all very coarsely granular, the roughnesses extending forward about to the eye. Gill membranes forming a broad free fold across isthmus. Dorsal spine moderate, a little more than  $\frac{1}{2}$  head, about equal to pectoral spine; humeral process triangular, granular, not quite  $\frac{2}{3}$  length of pectoral spine; axillary pore obsolete; adipose fin long and low, its posterior margin little free; caudal short and broad, the upper lobe the longer,  $1\frac{1}{2}$  in head; anal and ventrals rather small, the vent close behind the latter. Color dark brown, with bronze reflections; belly white; fins all dusky in 1 specimen, in the other mostly pale; maxillary barbels dusky, others pale. Two large specimens obtained at Panama by Gilbert. (Jordan & Gilbert. Bull. U. S. Fish Comm., II, 1882, 40.)

181. NETUMA INSCULPTA (Jordan & Gilbert).

Head 4 ( $4\frac{1}{2}$  in total); depth  $5\frac{1}{2}$  ( $6\frac{1}{2}$ ); width of head  $4\frac{1}{2}$ . D. I, 6; A. IV, 14. Length (29415)  $13\frac{1}{2}$  inches. Body moderately elongate, little compressed, the caudal peduncle slender and short. Head shortish, low and

broad, anteriorly depressed. Eye rather large,  $6\frac{1}{2}$  in head, placed rather high. Interorbital space flat and nearly smooth, 2 in head; snout 3; breadth of mouth  $1\frac{5}{6}$ ; snout very bluntly rounded, almost truncate in front. Mouth large; teeth all villiform; vomerine bands of teeth large (fully confluent with each other in the type, partly separated in smaller examples), and with the large, club-shaped band on the palatines, from which they are separated by a slight furrow and constriction; palatine band of teeth with a backward prolongation; premaxillary band of teeth large; maxillary barbel long, somewhat compressed, extending to middle of pectoral spine; outer mental barbel reaching base of pectoral spine, inner 2 in head. Dorsal shield short, crescent-shaped, without median keel, its tips produced, its length on the median line about  $\frac{2}{3}$  the length of 1 of its halves. Occipital process about as broad at base as long, with a moderate median keel, its lateral margins somewhat concave; fontanel becoming gradually contracted at a point a little nearer base of dorsal than tip of snout, thence forming a narrow groove, which extends to within a diameter of the pupil of the base of the occipital process; this groove sometimes nearly obsolete; greatest width of fontanel about  $\frac{2}{3}$  diameter of eye. Granulated striæ extending along the sides of the fontanel to a point opposite or in front of middle of eye. Shield of head finely and evenly granulated, the roughnesses more uniform than usual, and many of them arranged in lines, especially anteriorly; opercle not striate, the skin marked with fine vermiculations; gill membranes forming a broad fold across the isthmus. Dorsal and pectoral spines long, about equal,  $1\frac{1}{2}$  in head. No axillary pores; humeral process very large, triangular, finely granular, about  $\frac{1}{2}$  as long as pectoral spine; adipose fin large, without free tip; upper lobe of caudal the longer,  $1\frac{1}{2}$  in head; anal and ventrals moderate, the vent close behind the latter. Color rather pale; belly pale; fins and barbels all pale, or but slightly tinged with dusky. A single adult male was obtained by Dr. Gilbert at Panama. Two smaller ones are in the Museum collection, also from Panama. (Jordan & Gilbert, Bull. U. S. Fish Comm., II, 1882, 41.)

182. NETUMA PLANICEPS (Steindachner).

Head 4 ( $4\frac{1}{2}$  in total); depth  $5\frac{1}{2}$  ( $5\frac{3}{4}$ ); width of head 5. D. I, 7; A. IV, 13. Length (29417) 11 inches. Body comparatively elongate; the head small, rather narrow, depressed anteriorly; the snout rather narrow and moderately rounded. Eye moderate, placed well above mouth, its length  $5\frac{1}{2}$  in head. Interorbital space flat and smooth,  $2\frac{1}{2}$  in head; snout  $3\frac{1}{2}$ ; breadth of mouth 2. Mouth rather large, with thickish lips; teeth villiform; vomerine bands moderate, confluent with each other and with the much larger ovate palatine bands, a slight constriction or furrow making the divisions; palatine bands each with a backward prolongation; premaxillary band moderate; barbels very short; maxillary barbel scarcely or not reaching to base of pectoral; outer mental barbel scarcely past gill opening below; inner shorter than snout. Dorsal shield short, anteriorly truncate, not keeled, the length on the median line about  $\frac{1}{4}$  of 1 of its halves. Occipital process subtriangular, rather narrow, truncate

behind, its margins straight, becoming concave forward, its width at base about equal to its length; fontanel an almost obsolete groove, its posterior end not reaching base of occipital process by about the diameter of the eye, the groove extending forward to a point about midway between tip of snout and base of dorsal spine; anterior to this point is an equilateral triangle, flat, covered with smooth skin, the base of the triangle formed by the smooth, flattish interorbital area. Shields of head rather coarsely granular-striate, the granulations beginning anteriorly about opposite posterior margin of eye; opercle scarcely striate; gill membranes forming a moderate fold across the isthmus. Dorsal spine high, about equal to pectoral spine, and but little shorter than head; no axillary pore; humeral process triangular, granulated, a little more than  $\frac{1}{2}$  length of pectoral spine; adipose fin rather long; upper lobe of caudal the longer, a little shorter than head; ventrals and anal moderate. Color brownish, not very dark; belly pale, thickly speckled with brown; fins more or less dusky; maxillary barbels black; mental barbels pale. Two specimens were obtained by Dr. Gilbert at Panama. They disagree in several details from Steindachner's description, and it is barely possible that they belong to a different species. The head in Steindachner's types is  $3\frac{1}{4}$  to  $3\frac{1}{2}$  in length, and the occipital process is narrower and less widened anteriorly. (Jordan & Gilbert, Bull. U. S. Fish Comm., II, 1882, 42.)

183. NETUMA PLATYOGON (Günther).

Head  $3\frac{3}{4}$  ( $4\frac{1}{2}$ ); width of head  $4\frac{1}{2}$ ; depth  $5\frac{1}{2}$  ( $6\frac{3}{8}$ ). D. I, 7; A. IV, 14. Length (28286)  $15\frac{1}{2}$  inches. Body rather elongate, the head not very broad nor much depressed, a little broader than deep. The rather large, 5 to 6 in head. Interorbital space slightly more than  $\frac{1}{2}$  head, a trifle less than width of mouth; length of snout  $3\frac{1}{2}$  in head. Teeth all pointed; bands of vomerine teeth small, roundish, their boundaries traceable by a slight depression in the young, in the adult fully confluent with each other and with the palatine bands; palatine bands broad, ovate, several times as large as the patches on vomer, continued backward over the pterygoid region; premaxillary band rather broad, 5 to 6 times as broad as long; maxillary barbel reaching past base of pectoral in the young, not to gill opening in the adult, its base a little broader and more compressed than usual; outer mental barbels 2 in head; inner  $2\frac{1}{2}$ . Dorsal shield very short, lunate, subtruncate in front, its breadth more than 3 times its length on the median line; occipital process long, triangular, with straight margins, its length about  $1\frac{1}{2}$  times its width in front, its broad median line rather sharply keeled. In the young it is proportionally shorter, little longer than broad. At the beginning of this keel is the end of the long, narrow, groove-like fontanel, which extends forward to a point just behind the eye, where it merges into the flattish and smooth anterior part of the head. Shields of the head all finely granular, the granules rarely forming distinct lines. Dorsal spine long,  $1\frac{1}{2}$  to  $1\frac{1}{2}$  in head, the soft rays projecting beyond the spine; pectoral spine about as long as dorsal, sharply serrate behind, the anterior



serræ not very sharp; axillary pore small or absent; humeral process nearly smooth, rather narrow and short,  $\frac{1}{2}$  length of pectoral spine; adipose fin short and rather high, its base barely  $\frac{1}{2}$  length of base of anal; caudal deeply forked, its upper lobe the longer and slightly falcate, about as long as head; ventrals very short, reaching anal in females, shorter in the males; vent nearer base of ventrals than anal. Color in life very pale olive brown, with bronze and blue reflections, white below; fins all pale, the tip of anal and edges of caudal somewhat dusky; female with fins rather darker, the upper edge of the pectorals and ventrals largely black; in the males these fins are pale, or somewhat brown above; maxillary barbels blackish; lower pale. Generally abundant along the Pacific coast of tropical America. Specimens were observed by Dr. Gilbert at Mazatlan, Libertad, Punta Arenas, and Panama. It reaches a length of about 18 inches, and is seldom eaten. It resembles *Galeichthys gilberti*, but is readily distinguished by the small, pale ventrals, as also by the generic character of the dentition. The males of this species, according to Dr. Steindachner, carry the eggs in their mouths until after hatching. (Jordan & Gilbert, Bull. U. S. Fish Comm., 11, 1882, 44.)

184. NETUMA OSCULA (Jordan & Gilbert).

Head  $3\frac{1}{4}$  ( $4\frac{1}{4}$  in total); depth  $6\frac{1}{4}$  ( $7\frac{1}{4}$ ); width of head  $4\frac{1}{4}$ . D. I, 7; A. IV, 14. Body moderately elongate, the head short, rather narrow, tapering forward, considerably broader than deep. Eye small,  $7\frac{1}{4}$  in head, placed well above the mouth. Interorbital space  $1\frac{1}{10}$  in head; snout 3; breadth of mouth  $2\frac{1}{4}$ . Mouth very small for the genus, with thick lips. Teeth on vomer and palatines villiform, but rather coarse and bluntish. Vomerine patches small, rather longer than broad, separated on the median line, and each also separated by a narrow groove from the large and roundish palatine bands, which have a distinct backward prolongation. Premaxillary band of teeth very broad, barely 3 times as long as wide. Barbels short, the maxillary barbels reaching slightly beyond base of pectorals, the outer mental barbels scarcely past gill opening below; inner mental barbels about as long as snout. Dorsal shield short, crescent-shaped, granulated, but without median keel, its length about  $\frac{1}{2}$  its breadth. Occipital process narrow, its edges almost parallel until abruptly widened at base; the narrow part considerably longer than broad, with curved edges; a well-developed median keel. Fontanel broad and shallow, abruptly contracted at a point midway between tip of snout and end of occipital process, thence continued backward as a narrow groove to a point less than an eye's diameter in front of the base of the occipital process. Greatest width of fontanel about  $\frac{1}{2}$  eye. Shields of top of head all coarsely and rather sparsely granular, and anteriorly striate. Interorbital space nearly plane, with a few low, smooth ridges. Opercles scarcely rugose. Gill membranes forming a narrow fold across isthmus posteriorly. Dorsal spine very high,  $1\frac{1}{2}$  in head, a little longer than pectoral spine; humeral process granular, not quite  $\frac{1}{2}$  length of pectoral spine; no axillary pore; adipose fin adnate posteriorly; caudal long,

its upper lobe the longer, somewhat falcate,  $1\frac{1}{6}$  in head; anal rather high. Color brown, with bluish reflections; lower parts dusky, with dark punctulations; fins all blackish; maxillary and outer mental barbels dusky. A single male example 11 inches long was obtained at Panama by Dr. Gilbert. (Jordan & Gilbert, Bull. U. S. Fish Comm., II, 1882, 46.)

## 185. NETUMA FLATTURA (Jordan &amp; Gilbert).

Head  $3\frac{3}{4}$  ( $4\frac{1}{4}$  in total); depth  $5\frac{1}{2}$  ( $6\frac{1}{2}$ ); width of head  $4\frac{3}{4}$ . D. I, 6; A. IV, 14. Length (29408, U. S. Nat. Mus.)  $12\frac{1}{2}$  inches. Body low, not very elongate, the head rather short and very broad, much broader than deep, the snout depressed and very broadly rounded, almost truncate. Eye moderate, placed rather high, its diameter 7 in head. Interorbital space  $2\frac{1}{2}$  in head; snout  $3\frac{3}{4}$ ; breadth of mouth  $1\frac{3}{4}$ . Mouth large, with thickish lips, the upper jaw considerably projecting. Teeth on vomer and palatines villiform, but bluntly conical, less acute than in most of the species. Vomerine patches oblong, small, separated by a narrow interspace from each other and from the palatine bands, which are roundish and comparatively small, with a backward prolongation. Teeth in jaws in broad bands. Barbels rather short, the maxillary barbels reaching a little past base of pectorals, the outer mental barbels a little past gill opening, the inner a little more than  $\frac{1}{2}$  head. Dorsal shield not very short, crescent-shaped, with a distinct median keel, its length on the median line about  $\frac{1}{2}$  its breadth. Occipital process short, broadly triangular, with concave sides which spread out abruptly near the base, forming a sort of shoulder, its length scarcely equal to its width at base. Median keel well developed. Fontanel broad and shallow, abruptly narrowed posteriorly at a point a little nearer base of dorsal than tip of snout, but extending as a groove to a point distant less than a diameter of the eye from the base of the occipital process, this groove indistinct in the smaller specimen. Greatest width of fontanel scarcely more than  $\frac{1}{2}$  the eye. Shields of head granular-striate, the roughness less coarse than in *A. kessleri*. Interorbital space with 2 prominent ridges and numerous striae, none of them granular, the granulations chiefly confined to the region behind widest part of fontanel. Opercle striate. Gill membranes forming a moderate fold across isthmus. Dorsal spine low, shorter than pectoral spine, which is  $1\frac{1}{2}$  in head, the anterior edges of both bluntly serrate; humeral process broadly triangular, granulated, not  $\frac{2}{3}$  length of pectoral spine, much smaller than in *A. insculptus*; no axillary pore; adipose fin long and low, without free posterior margin; lower fins of moderate length; vent much nearer ventrals than anal. Caudal short, the upper lobe longest,  $1\frac{3}{4}$  in head (a little more than  $\frac{1}{2}$  head in the smaller specimens). Color dusky above, the lower parts soiled with dark points; fins all more or less dusky with dark points; maxillary barbels dusky, others pale. One male individual (29408) was obtained at Panama by Dr. Gilbert; another (30995) at Panama by Mr. Rowell. (Jordan & Gilbert, Bull. U. S. Fish Comm., II, 1882, 45.)

## 185 (a). NETUMA INSULARUM, Flora Hartley Greene.

Head  $3\frac{3}{4}$  in length; width of head  $4\frac{3}{4}$  in length; interorbital space in length 7; interorbital space in head scarcely 2; snout in head 3; breadth of mouth in head 2; eye in head  $6\frac{3}{4}$ . D. 1, 7; A. 17. Head much broader than deep; snout depressed and broadly rounded; eye above the level of the mouth. Upper jaw projecting. Teeth on vomer and palatines villiform and bluntly conical, the 2 vomerine patches forming together a band almost as long and slightly broader than the premaxillary band, the 2 sides separated by a narrow interspace; palatine teeth well separated from the vomerine teeth and in 2 large triangular patches which extend backward over the pterygoid region; each triangle has a sharp notch in its anterior side; its antero-posterior length is twice its lateral width; teeth of lower jaw in a narrower band than the upper jaw. Maxillary barbel extending to end of first third of the length of the pectoral spine; outer mental barbel to base of pectoral; inner mental barbel past gill opening,  $2\frac{1}{2}$  in head. Dorsal shield crescent-shaped, without median keel, length on median line  $2\frac{3}{4}$  in distance between the horns of the crescent; 2 notches on its anterior side to meet the corresponding points from the occipital process. Occipital process broadly triangular, with the outer sides concave and 2 small projections at its posterior end. Median keel evident, rather short. Occipital process much broader at base than long; its length  $3\frac{1}{2}$  in head; posterior breadth 2 in length of process. Fontanel broad and shallow, narrowed gradually posteriorly to a point halfway between snout and base of dorsal spine. A narrow line runs back from it the distance of a long diameter of the eye. Greatest width of the fontanel equals the short diameter of the eye. Shields of the head granular-striate, the striae evident and extending to the middle of the interorbital space, and on the side to meet the humeral process at the top of the gill opening. Opercles nearly smooth. Gill membranes forming a fold across the isthmus. Dorsal and pectoral spines crenulate in front and sharply decurved serrate behind. Dorsal shorter than pectoral, which is  $1\frac{1}{4}$  in head. No axillary pore evident. Adipose fin long and low with posterior margin attached. Vent much nearer ventrals than anal. Color in alcohol, dark blue above, light blue on side, and white below; maxillary barbel dusky; fins all dusky. The type of this species (No. 47577, U. S. Nat. Mus.) was collected by the *Albatross* in the Galapagos Archipelago, being part of the collection studied by Jordan & Bollman in 1889. It was recorded by them (Proc. U. S. Nat. Mus. 1889, 179) as "*Tachysurus elatturus* (var?)." Its relations to *Netuma elattura* are close, but its fins are better developed and there are several differences in details of structure.

*Netuma insularum*, FLORA HARTLEY GREENE in GILBERT, Proc. U. S. Nat. Mus. 1896 (Feb. 5, 1897), 439, Galapagos Archipelago.

## 69. GALEICHTHYS, Cuvier &amp; Valenciennes.

Dorsal shield small, lunate; teeth on palate villiform, the patches on each side not extending backward over the pterygoid region.

Subgenus GALEICHTHYS.

Shields of head mostly covered by soft skin, hiding the granulations.

163. GALEICHTHYS LENTIGINOSUS (Eigenmann & Eigenmann).

Head 4 to  $4\frac{1}{2}$ ; depth 5 to 6. D. I, 6; A. 22. Eye  $2\frac{1}{2}$  in snout,  $8\frac{1}{2}$  in head,  $4\frac{1}{2}$  in interocular,  $2\frac{1}{2}$  in interorbital. Body nearly terete anteriorly, becoming compressed backward; the width, above the pectorals, a little greater than the depth. Head flat, depressed, its depth at base of occipital process  $1\frac{1}{2}$  in its greatest width, which is about  $1\frac{1}{2}$  in its length. Occipital process somewhat roughened, about twice as long as its greatest width, its margin straight and oblique; the middle of the fontanel above the posterior part of the eye. Head everywhere covered with skin; sides of the head and opercle with vermiculating canals. Snout somewhat pointed; upper jaw very little projecting; lips thick; teeth all villiform; the intermaxillary band strongly curved; vomerine teeth in 2 oval patches joined to the larger patches of the palatines; mandibular band of teeth separated in front, the outer margins, if continued forward, forming an angle at the symphysis. Maxillary barbels reaching beyond base of pectorals; mental barbels reaching about  $\frac{2}{3}$  toward the gill opening; the postmentals to the gill opening in 1 specimen, a little before in the other. Gill membranes forming a broad, free margin across the isthmus. Gill rakers 3 + 4. Pectoral pore minute; humeral process pointed behind. Distance of dorsal from snout  $2\frac{1}{2}$  in the length, the dorsal spine covered with a membrane, its outer margin granular, its height  $1\frac{1}{2}$  in the head, the first soft ray  $1\frac{3}{4}$  the length of the fin; distance of adipose fin from the dorsal  $3\frac{3}{4}$  to 4 in the length, the fin adnate, longer than the dorsal; caudal lunate, the upper lobe longer, somewhat falcate,  $4\frac{1}{2}$  to 5 in the length; anal fin twice as long as high, the highest ray 2 to  $2\frac{1}{2}$  in the head; ventrals short and broad,  $1\frac{1}{2}$  in the head; pectoral spine covered with a membrane,  $1\frac{1}{2}$  to  $1\frac{3}{4}$  in head. Light brown, becoming nearly white below, the sides freckled; fins reddish. Panama. (Eigenmann & Eigenmann, Nematognathi, 50.)

164. GALEICHTHYS PERUVIANUS, Lütken.

Head  $3\frac{1}{2}$ ; depth  $4\frac{1}{2}$  to  $5\frac{1}{2}$ . D. I, 7; A. 14 to 16. Eye 2 in snout, 7 in head, 4 in the interocular, 2 in the interorbital. Subterete, tapering to a long, slender caudal peduncle; the greatest width about equal to the greatest depth. Head not much depressed; interorbital area flattened, the greatest depth of the head  $1\frac{1}{2}$  in its length, its greatest width  $1\frac{1}{2}$  to  $1\frac{3}{4}$ ; the width at angle of mouth 2 in its length; the surface of the cranial bones longitudinally furrowed, covered with muscle and skin. Occipital process more than 3 times as long as wide; anterior fontanel elongate, its center over the middle of the eye, continued as a deep groove to the base of occipital process; a small opening a pupil's distance behind the anterior fontanel, and a larger one in the occipital bone at the end of the groove. Snout, upper part of the neck, and the opercle sometimes with conspicuous reticulating mucous canals. Snout blunt, decurved. Maxillary barbels extending beyond base of pectoral, mentals about to gill

openings, the postmentals about 1 diameter of the eye farther. Jaws subequal, the upper longer; teeth all fine, villiform; intermaxillary band of teeth very wide, its depth about 8 in its width; 2 small patches on the vomer; palatine patches very wide and shallow, tapering to a point. Gill membranes meeting at an acute angle, forming a fold across the isthmus; gill rakers 3+10. Humeral process very thin, covered with skin, more than  $\frac{1}{4}$  as long as the pectoral spine, broadly expanded and rounded behind; pectoral pore present. Distance of dorsal spine from snout  $2\frac{1}{2}$  to  $2\frac{3}{4}$  in the length; the dorsal spine broken off in all the specimens; distance of adipose from the dorsal  $3\frac{1}{2}$  to  $3\frac{3}{4}$  in the length; adipose fin as long as the dorsal fin, adnate; caudal fin broadly lunate, the upper lobe longer, falcate,  $3\frac{1}{4}$  in the length; anal fin higher than long, the highest ray 2 to  $2\frac{1}{2}$  in head; ventrals reaching to the anal,  $1\frac{1}{2}$  to 2 in head; pectoral spines broken in all the specimens. Back, top of head, and a band from humeral process to the lower caudal lobe, blue black; a broad conspicuous, bluish-silvery band along the lateral line; lower parts white; fins blackish; ventrals and anal sometimes with light areas. Eleven specimens, 0.25 to 0.35 m. long. Callao, Peru; Haslar Expedition. (Eigenmann & Eigenmann, *Nematognathi*, 5.)

Subgenus *HEXANEMATICHTHYS*, Bleeker.

Shields not entirely covered by soft skin, the granulations evident, especially in the male.

187. *GALEICHTHYS SEEMANNI* (Günther).

D. I, 7; A. 19; P. I, 10. The height of the body is contained  $4\frac{1}{2}$  times in the total length (without caudal); the length of the head  $2\frac{1}{2}$ ; head much broader than high, its greatest width being equal to its length without snout. Eyes of moderate size, much nearer to the end of the snout than to the operculum; the length of the snout is  $\frac{2}{3}$  of the width of the interorbital space. The median longitudinal fonticulus on the upper side of the head extends to the base of the occipital process. Teeth on the vomer separated in the middle by a short interspace, forming a pair of small subquadrangular patches which are confluent with those of the palatines. The latter are much longer than broad, elliptical. The band of intermaxillary teeth is  $5\frac{1}{2}$  times as broad as long. The maxillary barbels extend nearly to the end of the head, and are about twice as long as the outer ones of the mandible. Crown of the head, and nape finely granular; occipital process broader than long, with a prominent ridge along its middle. The basal bone of the dorsal spine is small, with a few fine granules. Dorsal spine of moderate strength, more than  $\frac{1}{4}$  as long as the head, serrated along both edges; the first soft ray is as high as the body. Adipose fin rather shorter than dorsal. The upper caudal lobe is the longer,  $\frac{2}{3}$  of total length. Porus axillaris present. Ventral fin shorter than pectoral. Sides of the body silvery; basal half of the inner side of the paired fins black. Central America. A fine specimen 12 inches long, from the Haslar collection, collected by Dr. Seemann. (Günther.)

*Arius seemanni*, GÜNTHER, *Cat.*, v, 147, 1864, Central America.

Jordan (Proc. Ac. Nat. Sci. Phila. 1883, 282) adds the following note on the type of this species: "Fontanel extending backward in a deep and narrow groove which reaches the occipital process. Middle of top of head smooth, much as in *A. platypogon*."

The following account is given by Eigenmann & Eigenmann (Nematognathi, 78):

Head  $3\frac{1}{2}$ ; depth 5. D. I, 7; A. 18. Body about as deep as wide, tapering to a slender peduncle. Head flat, depressed in front, top of the head coarsely granular; opercles smooth or with faint striations; humeral process slightly granular, covered with skin; the greatest depth of the head  $1\frac{1}{2}$  in its length, greatest width  $1\frac{1}{2}$  to  $1\frac{2}{3}$ ; the width at angles of the mouth  $2\frac{1}{2}$ . Occipital process wider than long; fontanel open to above the posterior margin of the eye, with a deep backward-extending groove. Interorbital area smooth, without ridges. Eye 2 in snout, 7 in head,  $3\frac{1}{2}$  to 4 in the interocular,  $2\frac{1}{2}$  in the interorbital. Maxillary barbels reaching slightly beyond base of pectorals, mental barbels  $\frac{2}{3}$  toward the gill opening, the postmentals  $\frac{1}{2}$  a diameter of the eye behind the gill opening or farther. Upper jaw longer; teeth all villiform; vomerine teeth in 2 small ovate patches, which are separated from each other but joined to the much larger palatine patches. Gill membranes forming a moderate fold across the isthmus. Gill rakers 5 + 12. Pectoral pore large; vertical series of pores present. Distance of dorsal fin from tip of snout  $2\frac{1}{2}$  to  $2\frac{3}{4}$  in the length, the spine rather stout,  $1\frac{1}{2}$  in head, its outer edge granular toothed, its inner edge with short, recurved teeth; distance of the adipose fin from the dorsal  $3\frac{1}{2}$  to  $3\frac{3}{4}$  in the length; adipose fin slightly longer than high, shorter than the dorsal fin. Caudal 4 in the length; anal emarginate, little longer than high; ventrals  $1\frac{1}{2}$  to 2 in the length of the head; pectoral spine  $1\frac{1}{2}$  in head, its anterior margin granular toothed, its inner edge with long, straight teeth. Plumbeous, silvery below; fins dusky, inner surface of ventrals and pectorals dark. One female 0.28 m., Panama. One male 0.21 m., Panama.

187(a). GALEICHTHYS GILBERTI, Jordan & Williams.

Head  $3\frac{1}{2}$  to 4; width of head  $5\frac{1}{2}$ ; depth 5. D. I, 7; A. IV, 14. Body comparatively elongate, the head depressed but not very broad, somewhat broader than high; eye rather large, 5 to 6 in length of head; width of interorbital space  $2\frac{1}{2}$  in head; breadth of mouth  $1\frac{1}{2}$ ; length of snout  $3\frac{1}{2}$ . Teeth all villiform; bands of vomerine teeth separated by a rather wide interval, each small, roundish, confluent with the neighboring palatine band, the junction marked by a slight constriction; palatine band ovate, broad behind, varying considerably in size and somewhat in form, the width ranging from  $\frac{1}{2}$  diameter of eye to  $\frac{2}{3}$ , being generally largest in adults; band of palatine teeth without backward prolongation; band of maxillary teeth rather broad and short, its length about 5 times its breadth. Maxillary barbel flattened at base, reaching a little past base of pectoral in young, scarcely to gill opening in adult; outer mental barbels 2 in head, inner 3. Gill rakers 4 + 12. Dorsal shield very short, narrowly crescent-shaped, its length on median line not more than  $\frac{1}{2}$  that of one

of its sides. Occipital process subtriangular, not quite as long as broad at base, with a strong median keel, its sides slightly curved. A short distance in front of the beginning of the keel is the end of the very narrow groove-like fontanel, which is somewhat widened anteriorly, finally merging into the broad, flat, smooth interorbital area, the boundaries of which are not well defined; shields of head unusually smooth, all finely and very sparsely granular, the granules not forming distinct lines. Some specimens (probably females) about as smooth as in the subgenus *Galeichthys*. Gill membranes forming a rather broad fold across isthmus. Dorsal spine long, usually, but not always, shorter than the pectoral spine, about  $1\frac{1}{2}$  in head; axillary pore absent; humeral process rather broadly triangular, not much produced backward, less than  $\frac{1}{2}$  length of pectoral spine, its surface not granular, covered with skin; adipose fin  $\frac{1}{2}$  length of anal, its posterior margin little free; upper lobe of caudal the longer and somewhat falcate, about as long as head; ventrals long, about reaching anal in females, rather shorter in males; vent much nearer base of ventrals than anal. Color olive green, with bluish luster, white below; upper fins dusky olivaceous; caudal yellowish dusky at tip; anal yellowish with a median dusky shade; ventrals yellowish, the basal half of upper side abruptly black; pectorals similarly colored, the black area rather smaller; maxillary barbel blackish; other barbels pale. Length 12 to 18 inches. Coast of Sinaloa; very common; by far the most abundant species at Mazatlan; not recorded from localities farther south.

*Arius assimilis*, JORDAN & GILBERT, Bull. U. S. Fish Comm., II, 1882, 47; not of GÜNTHER. *Galeichthys gilberti*, JORDAN & WILLIAMS, Rept. Fishes Sinaloa, in Proc. Cal. Ac. Sci. 1895, 395, pl. 26, Mazatlan. (Type, No. 29213. Coll. Chas. H. Gilbert.)

#### 188. GALEICHTHYS JORDANI (Eigenmann & Eigenmann).

Head  $3\frac{3}{8}$  to  $3\frac{3}{4}$ ; depth  $5\frac{1}{4}$  to  $5\frac{3}{8}$ . D. I, 7; A. 18; eye large,  $1\frac{1}{4}$  in snout,  $5\frac{1}{4}$  in head, 2 in the interorbital,  $2\frac{1}{4}$  to 3 in the interocular. The specimens agree very closely with the description of *assimilis* by Jordan & Gilbert (*gilberti* of the present paper). They differ in the width of the mouth and in having a pectoral pore. Rather robust, the width little less than the depth; caudal peduncle compressed. Head heavy, little broader than high, its height  $1\frac{1}{2}$  in its length, its width  $1\frac{3}{8}$  to  $1\frac{1}{2}$ , width at the angle of the mouth 2 to  $2\frac{1}{4}$ ; interorbital area flat and smooth; posterior portion of the head finely and sparsely granular; opercle and humeral process smooth; occipital process about as long as broad, unusually sharply keeled; fontanel extending to above the posterior part of the orbit, continuing as a deep groove to the base of the occipital process; maxillary barbels extending to the pectoral pore, postmentals at least to the gill opening, mental about  $\frac{2}{3}$  as long as the postmental barbels; snout blunt, decurved; upper jaw a little produced; teeth all villiform, those on the vomer forming 2 small, separate, ovate patches, which are contiguous to the twice or thrice as large palatine patches; gill membranes forming a fold across the isthmus; gillrakers 6+9; pectoral pore large; vertical series of pores present; distance of dorsal spine from tip of snout  $2\frac{1}{2}$  to  $2\frac{3}{4}$  in the length; the spine of the dorsal and pectoral fins granular on the

basal half of their outer margin; almost the entire inner margins serrate, the spines of equal length,  $1\frac{1}{2}$  in head; distance of adipose fin from the dorsal  $3\frac{3}{8}$  in the length; the adipose more than  $\frac{1}{2}$  as long as the dorsal fin, its posterior margin free; caudal deeply forked, the upper lobe longer, somewhat falcate,  $3\frac{3}{4}$  to 4 in length; anal fin about as long as high, deeply emarginate, its highest ray  $2\frac{3}{8}$  in head; ventral fins not reaching to the anal, 2 in head. Dorsal surface dark blue, with metallic luster, becoming silvery below; lower caudal lobe dusky; basal half of the inner surface of the paired fins black. (Eigenmann & Eigenmann, Nematognathi, 79.) Panama; known only from specimens in the Museum of Comparative Zoology. The specimens from Mazatlan referred to *jordani*, *seemanni*, and *assimilis* by authors belong to *Gleichthys gilberti*.

## 188(a). GALEICHTHYS AZUREUS, Jordan &amp; Williams.

Head  $3\frac{1}{2}$ ; width of head  $4\frac{1}{2}$ ; depth 9. D. I, 7; P. I, 10; A. IV, 14. Gill rakers 6 + 13. Body robust, its width anteriorly greater than its depth; caudal peduncle short, stout; distance from end of anal fin to base of median caudal rays about  $\frac{1}{2}$  length of head. Head flat, very broad, its depth at posterior angle of jaw about  $\frac{1}{2}$  its width; interorbital region flat, smooth anteriorly and granulated posteriorly; fontanel almost obsolete, wide anteriorly and ending in a short groove posteriorly at a point  $\frac{1}{2}$  distance from tip of snout to posterior end of occipital process; top of head, occipital process, and antedorsal shield finely granular, granulations mostly arranged in radiating striae and extending forward to a line with the pupils; nostrils very large and close together, posterior one with a broad valve; occipital process pentagonal, its length  $4\frac{1}{2}$  in head, about as long as wide, with a very low ridge; dorsal shield crescent-shaped, with points extending back on each side of fin, its median length about  $\frac{1}{2}$  the length of the side; eye small, about 9 in head; interorbital width almost 2 in head; snout almost 4 in head; breadth of mouth  $2\frac{3}{10}$  in head; maxillary barbel slender, thick at base,  $1\frac{1}{2}$  in head; outer mental barbel reaching to posterior angle of jaw, about  $2\frac{3}{8}$  in head; inner mental barbel about 4 in head; teeth all villiform; premaxillary band narrow, about  $\frac{1}{4}$  as wide as long; vomerine and palatine bands of teeth fully confluent on each side, forming together a crescent-shaped patch, narrowly divided on the median line of the vomer; form of vomerine bands similar to that of the palatine bands but smaller; palatine band of teeth without backward prolongations; opercle with radiating ridges; humeral process granular, triangular, lower posterior corner prominent; axillary pore very small; gill membranes forming a broad fold across isthmus; dorsal fin short, base not including spine equal to base of adipose dorsal; dorsal spine robust, but little shorter than pectoral spine, about 2 in head, its anterior serrae small and tubercle-like, its posterior edge, as well as that of pectoral, retrorsely serrate; soft rays of dorsal extending but little beyond spine, the longest about  $\frac{2}{3}$  length of head; adipose dorsal about  $\frac{1}{2}$  as high as long; caudal lobes unequal, the upper lobe about  $\frac{1}{2}$  longer than lower lobe; anal short, of medium height; distance from vent to base of ventrals  $\frac{1}{2}$  distance from



origin of anal; pectoral spine very strong, its anterior margin with serrations toward the tip, becoming small tubercles toward base; soft rays but little longer than spine, which reaches slightly beyond  $\frac{1}{2}$  distance from origin to base of ventrals. Color dark blue, with silvery reflections on sides; belly pale; mental barbels dusky; maxillary barbels light below and black above; paired fins darkest on inner side; other fins almost uniformly dusky. One specimen, 19 $\frac{1}{4}$  inches long, taken by the Hopkins Expedition at Mazatlan; probably not distinct from *G. guatemalensis*.

*Galeichthys azureus*, JORDAN & WILLIAMS, Rept. Fishes Sinaloa, in Proc. Cal. Ac. Sci. 1895, 398, pl. 27, Mazatlan. (Type, No. 1575, L. S. Jr. Univ. Mus. Coll. Hopkins Exped. to Sinaloa.)

189. GALEICHTHYS CERULESCENS (Günther).

D. I, 7; A. 17; P. I, 10. The height of the body is contained about 5 times in the total length (without caudal), the length of the head  $3\frac{1}{2}$  or  $3\frac{3}{4}$  times; head much broader than high, its greatest width being  $\frac{2}{3}$  of its length. Eyes rather small, their diameter being  $\frac{1}{2}$  of the extent of the snout,  $\frac{2}{3}$  of their distance from the gill opening, and  $\frac{1}{4}$  of the width of the interorbital space. The teeth on the palate form a slightly curved band, composed of 2 vomerine patches which are much broader than long, and of a pair of palatine patches which are subcontinuous with, scarcely broader and longer than, those of the vomer. The barbels of the maxillaries extend to the middle, the outer ones of the mandible to the base of the pectoral. Crown of the head granular; occipital process broader than long, subtriangular, subtruncated behind, and slightly raised along the median line; the basal bone of the dorsal spine is subtriangular, small. Dorsal spine of moderate strength, more than  $\frac{1}{2}$  as long as the head, granulated in front and slightly serrated behind; the first soft ray is as high as the body; adipose fin shorter than the dorsal; caudal deeply forked, with the upper lobe the longer, its length being nearly equal to that of the head; pectoral spine serrated along its inner edge and on the extremity of its outer edge; it is as long as the head without snout. Ventral fin shorter than pectoral. Sides steel-blue iridescent, blackish toward the back, and silvery below; vertical fins black; inner side of the paired fins blackish. Guatemala. a-b. Five specimens, 12 inches long. Hnamuchal. From the collection of Messrs. Godman and Salvin. (Günther, Cat., v, 149.)

The following note on the types of this species is given by Jordan (Proc. Ac. Nat. Sci. Phila. 1883, 282): "Head more depressed than in *A. assimilis*. Fontanel very short, ending abruptly behind and not produced in a groove behind the smooth area of the top of the head, the boundary of the smooth area being rather abruptly convex. Occipital process broader than long, its edges nearly straight. Bands of palatine teeth small, not produced backward on the inner margin. Paired fins black at base above." No recent collector has found this species.

## 189(a). GALEICHTHYS XENAUCHEN, Gilbert, new species.

Head  $3\frac{3}{4}$  in length; depth at front of dorsal  $5\frac{1}{2}$ ; anal 23. Width of head at opercle  $1\frac{1}{2}$  in its length; width at front of eyes 2 in head; width of mouth at inner angles  $2\frac{3}{4}$  in head; interorbital width  $2\frac{1}{10}$ , eye very small, 9 in head,  $3\frac{1}{4}$  in its distance from tip of snout,  $4\frac{5}{8}$  in postocular part of head,  $4\frac{1}{4}$  in interorbital width. Teeth all villiform; mandibular bands well separated on middle line, very broad mesially, rapidly tapering to a point laterally, the band produced beyond angle of mouth, its greatest width  $2\frac{1}{2}$  times in its length; premaxillary band very convexly curved, following the outline of the snout, its width  $5\frac{3}{8}$  in its length; vomerine patches roundish, separated by an evident medial groove, marked off from the palatine patches by a narrower groove and a constriction; the palatine patches are equal in width to the vomerine patches, and less than twice as long, of nearly equal width throughout. Maxillary barbels very slender, reaching slightly beyond the base of the pectoral spine; the mental barbels do not reach edge of gill membrane, the outer pair equaling length of snout and  $\frac{1}{2}$  of eye. Nostrils very large, the anterior broadly oval, with widely reflexed rim; the posterior widely elliptical, not concealed by the valve; distance from anterior nostril to tip of snout equaling that from posterior nostril to front of eye. Fontanel wide, with nearly parallel edges on frontal region, abruptly narrowing at front of occiput, where it is continuous with a narrow and shallow groove; the latter fails to reach base of occipital process by a distance equaling  $\frac{1}{2}$  diameter of eye. The raised margins of the fontanel continuous with a pair of sharp ridges bounding the groove, these accompanied by a pair of lower ridges on their outer sides and parallel with them; posteriorly these ridges roughened with granules and merging into the granulated area on posterior part of occiput; occipital process granulated, the granules arranged in more or less definite lines radiating backward and downward on each side from median point of base; lateral portions of occiput with an area of radiating striae separated from the central ridges by a smooth groove-like depression; a narrow granulated area extending forward on each side of fontanel to above back of orbits; occipital process very long and narrow, its width opposite its middle being but  $\frac{2}{3}$  of its length, abruptly expanding near base, the basal width being  $\frac{1}{2}$  its length plus that of dorsal plate on median line; opercles and humeral plate weakly striate. Gill membranes with a wide, free fold posteriorly; gill rakers weak and short, 1+1 movable ones; no evident axial pore. Dorsal spine slender, with a series of sharp granulations on anterior edge, minutely roughened, not serrate, behind, broken in the type, but its length was about  $\frac{2}{3}$  that of head; pectoral spines rather slender, rough granular on outer margins, with short, fine serrae within, both mutilated in the type, but their length was about equal to that of dorsal spine; pectoral extending nearly  $\frac{2}{3}$  distance to ventrals, the ventrals nearly to origin of anal; distance from anus to base of ventrals  $\frac{2}{3}$  its distance from front of anal; anal fin very long, its base  $1\frac{1}{4}$  in head, its longest ray  $\frac{1}{2}$  head; distance between dorsals  $3\frac{1}{4}$  in length; adipose fin long, highest about opposite the middle, a short, almost verti-

cal, free posterior margin, its vertical height  $3\frac{5}{8}$  in its length; the latter over twice its distance from rudimentary caudal rays, greater than base of first dorsal, equal to  $\frac{1}{2}$  length of head; caudal with broad lobes, the lower rounded; the upper mutilated in the type, but evidently acute and longer than the lower. Color purplish above, more bluish anteriorly, the lower parts silvery. coarsely punctate with brown; fins all blackish except the lower surface of the paired fins. In appearance most closely allied to species of *Netuma*, having the low, depressed head with the lateral outlines converging forward to the narrow pointed snout, and a long, largely adherent adipose dorsal. The palatine patches are, however, narrow and without backwardly projecting lobes. The species is distinguished from all those known from the Pacific coast of America by the long and extraordinarily narrow occipital process. Type, a female 380 mm. long, from Panama. (*ξέρως*, strange; *αυχίην*, nape.)

*Galeichthys xenauchen*, GILBERT, Fishes of Panama, MS. 1898, Panama.

190. GALEICHTHYS GUATEMALENSIS (Günther).

Head  $3\frac{5}{8}$  ( $4\frac{1}{2}$  in total); width of head 5 ( $6\frac{1}{2}$ ); depth  $6\frac{1}{2}$  (7). D. I, 6; P. I, 10; A. III, 15. Length (28140, U. S. Nat. Mus.)  $12\frac{1}{2}$  inches. Body slender, its width anteriorly greater than depth; caudal peduncle compressed, short; distance from end of anal to base of median caudal rays about  $\frac{1}{2}$  length of head. Head depressed, not very broad, its depth at posterior margin of branchiostegal membranes less than  $\frac{3}{8}$  its width; interorbital region flat, smooth, the smooth area forming a broad equilateral triangle, its base at the interorbital space, the apex at a point  $\frac{1}{4}$  the distance from snout to dorsal, the triangle forming the termination of the almost obsolete fontanel; top of head, occipital process, and antedorsal shield finely granular, some of the anterior granulations only arranged in lines, none of them in radiating striae. Occipital process broadly trapezoidal, its width slightly greater than the length of its side, with a slight or obsolete median carina; its posterior margin truncated; its sides slightly convex posteriorly, concave toward the front. Dorsal shield small, narrow, crescent-shaped, its median length about  $\frac{1}{2}$  the length of its side. Eye small, 6 in head; interorbital width  $2\frac{7}{8}$  in head; snout 4 in head; breadth of mouth 2. Maxillary barbel very slender, reaching base of pectoral spine; outer mental barbel to well beyond margin of branchiostegal membranes, its length about  $\frac{1}{2}$  head; inner mental barbel 3 in head. Teeth all villiform; width of premaxillary band about  $\frac{1}{2}$  its length; vomerine and palatine bands of teeth fully confluent on each side, forming together a crescent-shaped patch, narrowly divided on the median line of the vomer; form of vomerine band similar to that of the palatine band; palatine band of teeth without backward prolongation; opercle with radiating ridges; humeral process granular, narrow, produced backward, not quite  $\frac{1}{2}$  length of pectoral spines; no axillary pore. Gill membranes forming a narrow fold across isthmus. Dorsal short, its base about equal to that of the adipose dorsal; dorsal spine robust, but little shorter than the pectoral spine, about  $\frac{3}{8}$  length of head, its anterior serrae small and tubercle-like; its posterior edge, as well as that of the pectoral, retrorsely serrate; soft rays of dorsal extending much beyond the spine,

the longest about  $\frac{2}{3}$  length of head; adipose dorsal about  $\frac{1}{2}$  as high as long, its posterior margin largely free; caudal very widely forked, the upper lobe falcate, nearly  $\frac{1}{2}$  longer than the lower, as long as head; anal short and low; distance from vent to base of ventrals slightly more than  $\frac{1}{2}$  its distance from origin of anal; pectoral spine very strong, much stronger than dorsal spine, its anterior margin with serræ toward the tip, becoming small tubercles toward base; inner edge with strong retrorse serræ, the soft rays longer than spines, reaching  $\frac{2}{3}$  distance to base of ventrals. Color very dark bluish or greenish above; sides with bronze luster; belly silvery; mental barbels white, with black edge; maxillary barbels blackish; fins all blackish, the caudal nearly uniform; paired fins darkest on inner side; sides with vertical series of mucous pores, conspicuous in life. This species is not uncommon at Mazatlan, where several specimens were obtained by Dr. Gilbert. Four specimens from Colima are also in the National Museum. It has not been observed at Panama. The original description of this species is brief and not entirely correct. That it was intended to refer to the species here described we have ascertained by the examination of Dr. Günther's original types in the British Museum. (Jordan & Gilbert, Bull. U. S. Fish Comm., II, 1882, 48.)

191. GALEICHTHYS ASSIMILIS (Günther).

D. I, 7; A. 19; P. I, 10. The height of the body is contained  $4\frac{2}{3}$  times in the total length (without caudal), the length of head  $3\frac{2}{3}$ ; head much broader than high, its greatest width being  $\frac{2}{3}$  of its length. Eyes rather small, situated nearer to the end of snout than to that of operculum; the length of snout is  $\frac{2}{3}$  of the width of interorbital space. The median longitudinal fenticulus on the upper side of the head does not extend to the base of occipital process. Teeth on vomer but slightly separated in the middle, forming a pair of oblong transverse patches which are confluent with those on the palatine bones; the latter are short, club-shaped. The band of intermaxillary teeth is 5 times as broad as long. All the teeth villiform. The maxillary barbels extend nearly to the end of head; the length of the outer ones of the mandible is  $\frac{1}{2}$  or  $\frac{2}{3}$  that of the head. Crown of the head granular, the granulations being arranged in radiating streaks. Occipital process broader than long, triangular, with its hinder end concave. The basal bone of the dorsal spine of moderate size, crescent-shaped. Dorsal spine of moderate strength, more than  $\frac{1}{2}$  as long as head, granulated in front and slightly serrated behind; the first soft ray longer than spine and as high as body; adipose fin shorter than dorsal; caudal deeply forked, with the upper lobe the longer, its length being contained  $5\frac{1}{2}$  times in the total; pectoral spine serrated along its inner edge and on the extremity of the outer edge; ventral fin shorter than pectoral. Sides of the body silvery; vertical fins grayish; basal half of the inner side of the paired fins black. Guatemala. A fine specimen, 13 inches long. Lake of Yzabal. From the collection of Messrs. Godman and Salvin. (Günther, Cat., v, 146.)

Jordan (Proc. Ac. Nat. Sci. Phila. 1883, 281) has the following note on the type of this species: "Area between the eyes smooth, extending backward in the form of a rather narrow triangle, which is moderately

obtuse behind. Fontanel narrow and short, ending far in front of occipital process, not extending backward as a groove behind the smooth area of the top of the head; posterior end of fontanel midway between tip of snout and middle of dorsal shield. Occipital process broad, its edges not straight. Band of palatine teeth large, but not produced backward on the inner margin. \* \* \* There is no evidence of the occurrence of the true *A. assimilis* in Pacific waters."

192. *GALEICHTHYS SURINAMENSIS* (Bleeker), text, p. 129.

193. *GALEICHTHYS DASYCEPHALUS* (Günther).

Head  $4\frac{1}{2}$  ( $5\frac{2}{3}$  in total); depth 6 ( $7\frac{1}{2}$  in total); width of head  $5\frac{1}{2}$ . D. I, 7; A. IV, 17. Length (29400) 11 inches. Body elongate, compressed behind, the head small, narrow, and moderately depressed anteriorly, the snout not very blunt. Eye rather large, placed somewhat above level of angle of mouth, its length 5 in head; width of interorbital space  $2\frac{1}{2}$  in head; breadth of mouth  $2\frac{1}{2}$ ; length of snout  $3\frac{1}{4}$ . Teeth villiform, those of vomer and palatines rather coarse, bluntly conic; bands of vomerine teeth separated by a rather broad area, each confluent with the neighboring palatine band, the two forming a small oblong patch much smaller than the eye, the division between the palatine and vomer scarcely appreciable. Palatine bands without backward prolongation. Bands of teeth in jaws short and broad. Maxillary barbel reaching about to middle of pectoral spine; outer mental barbel to base of pectoral; inner slightly more than  $\frac{1}{2}$  head. Dorsal shield short, crescent-shaped, a little more than 3 times as broad as long on the median line. Occipital process subtriangular, its sides straight, slightly longer than broad, its median line rather sharply keeled. Close in front of its base begins the deep fontanel, which is narrow and groove-like posteriorly, becoming rather abruptly broader above the opercle, then gradually narrowed anteriorly. Ridges bounding fontanel prominent anteriorly to a point just behind vertical from nostrils, coarsely granular for their whole length, the granules mostly arranged in 1 series. Between these ridges and the eye on each side is another ridge extending obliquely backward and inward from above front of eye, likewise very coarsely granular, the granules mostly in 2 series. Shields of head all rough granular, the granules forming irregular lines. Gill membranes forming a narrow fold across isthmus. Dorsal spine moderate, about equal to pectoral spine,  $1\frac{1}{2}$  in head; axillary pore present, small; humeral process broad, scarcely granular, about  $\frac{2}{3}$  pectoral spine; adipose fin rather long and low; caudal long, the upper lobe the longer, somewhat longer than head; anal long and high, its outline emarginate, its longest rays a little more than  $\frac{1}{2}$  head; ventrals long, the vent nearer their base than that of anal. Color dusky, the entire ventral surface soiled with dark points; fins all largely blackish; barbels black. Two specimens were obtained at Panama by Dr. Gilbert. This species may be known at once by the 4 granulated ridges, which extend the length of the interorbital space. In the female, later taken, the granulations on the head are largely covered by soft skin.

## 194. GALEICHTHYS LONGICEPHALUS (Eigenmann &amp; Eigenmann).

Head  $3\frac{3}{4}$ ; depth  $6\frac{1}{4}$ . D. I, 7; A. 20. Elongate, slender, greatest width little greater than the depth. Head long and depressed, its greatest width  $1\frac{1}{2}$  in its length, its greatest depth little more than  $\frac{1}{2}$  its length. Top of head with faint granules almost entirely concealed by the skin; interorbital area flat and with 4 ridges which are obscurely granular, the inner two bordering the fontanel, the outer ones curved in front extending obliquely backward from near the posterior nasal opening; occipital process as long as broad, its margins concave; fontanel produced as a deep groove to the base of the occipital process; opercle faintly striate; humeral process entirely covered with thick skin, not granular. Eye lateral, well above the angle of the mouth, its diameter  $1\frac{1}{2}$  in snout, 6 in head, 3 in interocular; snout depressed and rounded in front. Maxillary barbels extending scarcely beyond base of pectoral, mentals not to gill opening. Upper jaw little projecting; width of the mouth  $2\frac{1}{2}$  in the head; intermaxillary teeth long and slender, the depth of the band  $4\frac{1}{2}$  in its width; vomerine and palatine teeth obtusely conical, the vomerine patches separate, contiguous to, but not confluent with, the palatine patches. Gill membranes not forming an angle where they meet, with a rather broad, free margin. Gill rakers short and thick, 4 + 5. Pectoral pore small; vertical series of pores present; distance of dorsal fin from tip of snout  $2\frac{3}{4}$  in the length, the spine  $1\frac{1}{2}$  in the head, its outer margin granular-toothed near its base, its inner margin with short teeth; distance of adipose fin from the dorsal  $3\frac{1}{2}$  in the length; adipose fin much longer than high, as long as the dorsal fin; caudal forked, the upper lobe  $\frac{1}{2}$  longer than the lower, very nearly as long as head,  $3\frac{1}{2}$  in the length; anal fin emarginate, scarcely longer than high, its height  $2\frac{1}{2}$  in the head; ventrals reaching almost to the anal, about 2 in head; pectoral spine a little longer than the dorsal spine  $1\frac{3}{4}$  in the head; its outer edge roughened, inner edge with rather sharp teeth. Brown above, the sides silvery, entire ventral surface sprinkled with brown dots; a black median line on the back; fins dusky; barbels blackish. One specimen, a male, .29 m. long (No. 4972, M. C. Z.). Panama. Steindachner. (Eigenmann & Eigenmann, Nematognathi, 82.)

## 195. GALEICHTHYS FLUGISPINIS (Cuvier &amp; Valenciennes)

Head  $3\frac{1}{2}$  to 4; depth  $5\frac{1}{2}$  to 6. D. I, 7; A. 19 to 21. Slender, compressed on the tail. Head broad and depressed, tapering forward; width of the head  $1\frac{1}{2}$  to  $1\frac{3}{4}$  in its length, at the angle of the mouth  $2\frac{3}{4}$  to  $2\frac{1}{2}$ ; depth of head  $1\frac{3}{4}$  to 2; profile rather steep. Top of head, humeral process, front and sides of spines, and dorsal plate granular, the granulation not extending forward to above middle of cheeks. Occipital process triangular, about as long as broad, the median ridge not very prominent. Middle of fontanel behind the eye, the posterior portion separated by a bridge, not continued backward as a groove; interorbital region with 4 ridges. Eye small, 3 in snout, 10 in head,  $3\frac{1}{2}$  in the interocular. Barbels villiform. Maxillary barbel reaching to or beyond base of pectoral; postmental to gill opening, mental barbels much shorter. Mouth inferior, lower jaw

included, lips thick; teeth villiform, the anterior ones in the jaws longer; depth of the intermaxillary band 4 in its width; palatine patches 1 diameter of eye apart, the width of the patches less than 1 diameter of eye. Gill membranes meeting in an angle, forming a fold across the isthmus. Gill rakers 6 + 11. Pectoral pore none; vertical series of pores present. Distance of dorsal spine from the snout  $2\frac{1}{2}$  to  $2\frac{3}{4}$  in the length, the spine broken in the specimens examined. Space between dorsal and adipose fins 4 to  $4\frac{1}{2}$  in the length. Adipose fin adnate, as long as the anal fin; ventrals  $2\frac{1}{2}$  in the head; pectoral spine serrated behind (broken). Two specimens 0.22 m. and 0.26 m. long. Para. Agassiz and Bourget. (Eigenmann & Eigenmann, Nematognathi, 83.)

196. *GALEICHTHYS PHRYGIATUS* (Cuvier & Valenciennes), text, p. 130.

74. *TACHYSURUS*, Lacépède.

Teeth on palate granular; dorsal shield small; palatine bands of teeth without backward projecting angle.

197. *TACHYSURUS NUCHALIS* (Günther).

D. I, 7; A. 21; P. I, 10. The height of the body  $\frac{1}{2}$  of the total length (without caudal), the length of head  $\frac{1}{2}$ . Head as broad as high, its greatest width being  $\frac{2}{3}$  its length; its upper surface granulated; occipital process triangular, as long as broad, with the lateral margins slightly concave; it is elevated into an obtuse ridge running along the middle; the longitudinal groove in the middle of the forehead is rather wide, narrow behind, and does not extend to the base of occipital process. Teeth on palate are coarsely granular, and form 2 subtriangular patches of moderate extent, which, sometimes, are subcontinuous with their anterior angles. The maxillary barbels extend nearly to end of pectoral. Dorsal spine of moderate strength, slightly serrated along both edges,  $\frac{5}{8}$  length of head; adipose fin small, shorter than dorsal; pectoral spine as long as, but stronger than, that of dorsal; pectoral fin shorter than head. British Guiana. a-c. Six inches long. Purchased of Mr. Scrivener. d-f. Young. Presented by Sir R. Schomburgk. (Günther, Cat., v, 171.)

198. *TACHYSURUS FISSUS* (Cuvier & Valenciennes).

D. I, 7; A. 20 or 21. Length of head  $\frac{1}{2}$  of the total (without caudal). The distance between the end of snout and that of occipital process  $\frac{1}{2}$  of the total length (with caudal); basal bone of dorsal spine small. The teeth on the palate form 2 separate subovate patches. The maxillary barbel extends to, or nearly to, the middle of pectoral fin. Adipose fin small. Cayenne. a-b. Presented by Prof. R. Owen. These specimens having had the cavity of the mouth and of the gills extended in an extraordinary manner, I was induced to examine the cause of it, when, to my great surprise, I found them filled with about 20 eggs, rather larger than an ordinary pea, perfectly uninjured, and with the embryos in a forward state of

development. The specimens are males, from 6 to 7 inches long, and in each the stomach was almost empty. Although the eggs might have been put into the mouth of the fish by their captor, this does not appear probable. On the other hand, it is a well-known fact that the American Siluroids take care of their progeny in various ways; and I have no doubt that in this species and in its allies the males carry the eggs in their mouth, depositing them in places of safety, and removing them when they fear the approach of danger or disturbance. (Günther, Cat., v, 172.)

## 199. TACHYSURUS SPIXII (Agassiz).

Head  $3\frac{3}{4}$  to 4; depth 5 to  $5\frac{1}{2}$ . D. I, 7; A. 21. Body compressed, especially toward the caudal fin, the depth greater than the width. Head narrowed forward, its greatest width  $1\frac{1}{2}$  in its length, its greatest depth  $1\frac{1}{2}$ ; width at the mouth  $2\frac{1}{2}$  in the length of the head. Top of the head granular in the young, the granules becoming more or less united in the adult, forming fine reticulating ridges, especially on the occipital process, longer than broad, with a blunt median ridge, the margins concave. Fontanel narrow, without interruptions, continued as a deep tapering groove to near the base of the occipital process; interorbital area with 4 ridges; opercles and humeral process roughened, covered with skin; sides of the head, and snout with reticulating mucous canals. Eye  $1\frac{1}{2}$  to 2 in the snout, 5 to  $6\frac{1}{2}$  in the head,  $2\frac{3}{4}$  to 3 in the interocular. Maxillary barbels varying in extent, from about the middle of the pectoral to the base of the ventrals; postmental barbels extending to the base of pectoral or to near its tip; mentals to edge of gill membrane or to beyond base of pectoral. Upper jaw projecting; lips more or less papillose; teeth on the intermaxillary and the outer ones of the mandible, villiform; the inner series of the mandible and the palate with granular teeth; the palatine patches of teeth small, subovate, sometimes contiguous in front. Gill membranes united, joined to the isthmus, not forming a free margin across it; gill rakers 6 + 11 to 13. Pectoral pore moderate; distance of dorsal spine from snout  $2\frac{1}{4}$  to  $2\frac{1}{2}$  in the length; the spine  $1\frac{1}{2}$  to  $1\frac{1}{2}$  in head, serrated on its inner margin, granular or almost smooth on its outer margin. Distance of adipose from the dorsal fin  $3\frac{1}{2}$  to  $3\frac{3}{4}$  in the length, the adipose fin shorter than the dorsal fin, free posteriorly; caudal forked, the upper lobe slightly the longer, 4 to 5 in the length; anal fin scarcely longer than high, its highest ray about 2 in head; ventral fin  $1\frac{3}{4}$  to 2 in head; pectoral spine strong, about as long as the dorsal spine, serrated on its inner margin, granular or scarcely roughened on the outer margin. Color brownish above, sides and ventral surface silvery, sometimes with brown dots. We have examined over 70 specimens measuring from 0.07 to 0.24 m. from Maranhao, Bahia, Rio Janeiro, Para, Santos in São Paulo, Abrolhos, Brazil. The specimens from Para are much darker in color, the lips more papillose, the barbels longer than those of other specimens. The Santos specimens are ashy above, white below the lateral line, with rather large brown dots on sides, becoming fewer below. (Eigenmann & Eigenmann, Nematogathi, 89.)



## 200. TACHYSURUS MELANOPUS (Günther).

D. I, 7; A. 21; P. I, 10. The height of the body is contained 5 times in the total length (without caudal), the length of the head  $4\frac{1}{2}$  times; head somewhat broader than high, its greatest width being  $\frac{3}{4}$  of its length; the occiput and nape are finely granulated; occipital process subtriangular, as long as broad, with the lateral margins somewhat concave, and with the median ridge a little elevated. The longitudinal groove in the middle of the crown of the head is indistinct, narrow, linear behind, scarcely extending to the base of the occipital process. The teeth on the palatines are obtusely conical, and form 2 rather small subovate patches, apart from each other, and situated on the front part of the palate. The maxillary barbels do not quite extend to the middle of the pectoral fin. Dorsal spine of moderate strength, scarcely serrated anteriorly, equal in length to the distance of the gill opening from the anterior margin of the orbit, or even somewhat shorter; adipose fin small, the length of its base being less than that of the dorsal; pectoral spine nearly as long and strong as that of the dorsal fin, very strongly serrated anteriorly. Porous axillaries nearly as wide as a nasal opening. The upper (inner) surface of the ventral fins deep black, the lower (outer) white; the inner surface of the pectorals blackish. Rio Motagua (east slope). a-b. From 8 to 9 inches long. From Mr. Salvin's collection. (Günther, Cat., v, 172.) The specimens from the Pacific Coast mentioned in the text (page 132) belong to the following species.

## 200(a). TACHYSURUS LIROPUS, Susan B. Bristol.

Head  $3\frac{3}{4}$  to  $3\frac{1}{2}$ ; depth  $4\frac{3}{4}$  to  $5\frac{1}{10}$ . D. I, 6; A. II, 19; P. I, 9 or 10. Body elongate, its width anteriorly a little less than depth, the posterior portion much compressed; back elevated at front of dorsal; anterior profile from front of dorsal to tip of snout oblique; head flat, very broad, its width  $1\frac{1}{2}$  in its length; snout broad, rounded,  $1\frac{1}{10}$  to  $1\frac{1}{8}$  in interorbital width; eye rather large, laterally placed, its width about  $1\frac{1}{8}$  in its length,  $4\frac{1}{10}$  to  $4\frac{9}{10}$  in head; mouth small, upper jaw considerably projecting, its breadth  $2\frac{3}{4}$  to 3 in head; jaws thin; wide bands of minute pointed teeth present on both jaws; vomerine bands widely separated, and indistinguishable from the palatine band, which is small, oblong-ovate, and scarcely prolonged backward; interval separating vomerine bands about  $2\frac{1}{4}$  or 3 in eye; the teeth on these bands larger than those on jaws, and very bluntly conical. Interorbital space broad,  $2\frac{1}{4}$  to 3 in head. Barbels long and slender, the maxillary barbel extending to, nearly to, or, in some cases, past base of pectoral,  $1\frac{1}{4}$  to  $1\frac{1}{2}$  in head; outer mental barbel  $1\frac{1}{2}$  to  $1\frac{1}{4}$  in head; inner mental barbel  $2\frac{1}{2}$  to  $2\frac{5}{8}$  in head. Antedorsal shield very short, narrowly crescent-shaped, its length on the median line about 2 or 3 in its width; occipital process subtriangular, a little longer than broad at base, its edge slightly concave, its median keel strong. The long, narrow groove of the fontanel beginning abruptly a short distance in front of occipital keel, the distance from its end to base of dorsal  $1\frac{1}{2}$  to 2 in the distance to tip of snout. Shields of head rather smooth, finely granular, the granules forming distinct lines anteriorly. The flat

area between eyes triangular, with a median groove extending from fontanel forward to tip of snout, its posterior end a little behind eye, the granulations on each side of it extending forward as far as posterior border of pupil; opercles with no radiating striae. Gill membranes forming a very narrow fold across the isthmus. Gill rakers 5 + 12. Nostrils 2 on either side, large, placed close together and near tip of snout, the posterior with a large flap; axillary pore well developed; humeral process smooth, very short, 4 to  $5\frac{1}{2}$  in pectoral spine. Base of dorsal  $2\frac{3}{8}$  to  $2\frac{3}{10}$  in head; dorsal spine long and very strong,  $1\frac{1}{2}$  to  $1\frac{3}{4}$  in head, its upper anterior serræ small and tubercle-like, its upper posterior and its lower edges retroseely serrate; the soft rays extending considerably beyond the spine,  $1\frac{1}{2}$  to  $1\frac{1}{4}$  in head; adipose fin small, its base  $3\frac{1}{2}$  to  $4\frac{1}{2}$  in head, its height  $1\frac{3}{8}$  to 2 in its base; caudal widely forked, the upper lobe, measured from base of caudal to its tip, the longer, about  $1\frac{1}{2}$  in head; base of anal  $1\frac{3}{8}$  to  $1\frac{9}{10}$  in head, its longest ray  $2\frac{1}{2}$  to  $2\frac{3}{8}$  in head; ventrals reaching  $\frac{2}{3}$  to  $\frac{7}{8}$  the distance to origin of anal; vent about midway between origin of ventrals and origin of anal; pectoral spine  $1\frac{1}{2}$  in head, serrate, the serræ on inner edge larger and sharper than those on outer, the upper anterior serræ tubercle-like, the rays a little longer than spine. Bluish silvery, light yellowish below; top of head and back brown; fins dusky olive, lighter at base, all margined with darker; ventrals pale; adipose fin covered with minute black dots; maxillary barbels dark brown, with bluish silvery luster; other barbels lighter; eye yellowish. Here described from 6 specimens from San Juan Lagoon, mouth of Rio Ahome, Sonora, Mexico (No. 47584, U. S. Nat. Museum). Length  $7\frac{1}{2}$  to 9 inches.

*Tachysurus liropus*, BRISTOL, in GILBERT, Proc. U. S. Nat. Mus. 1896 (Feb. 5, 1897), 438, San Juan Lagoon, near mouth of Rio Ahome, Sonora, Mexico.

200(b). TACHYSURUS EMMELANE, Gilbert, new species.

Head  $3\frac{3}{8}$  in length ( $4\frac{1}{10}$  in total); depth 5 (6 in total). A. 27 (3 + 24). Eye 7 in head,  $2\frac{1}{2}$  in its distance from tip of snout, 4 in postorbital part of head,  $3\frac{3}{8}$  in interorbital width,  $2\frac{1}{2}$  in frontal width opposite middle of eyes. Mouth of moderate width, gently convex, the distance between its angles (measured internally)  $2\frac{3}{8}$  in head. Teeth in premaxillary and front of mandible finely villiform; posterior mandibular teeth stronger than those in front, bluntly conic, not, however, granular or flat and pavement-like, as are the posterior mandibular teeth in *T. furthii*, *T. melanopus*, and *T. liropus*. Mandibular bands with a wide interspace mesially, each widest near symphysis, rapidly tapering laterally, and extending beyond angle of mouth. The width of the bands is less than in related species,  $\frac{1}{4}$  eye at their widest point. The length of 1 of the mandibular bands is slightly greater ( $1\frac{1}{10}$ ) than length of eye. Premaxillary band very short, its length but  $\frac{1}{2}$  greater than that of 1 of the mandibular bands, extending on each side less than  $\frac{1}{2}$  distance from median line to angle of mouth; width of band  $\frac{1}{2}$  its length. Palatine teeth granular, in small oblong patches, which taper to a point laterally, and are widely separated on medial line, the patches agreeing in size and shape with those in *T. liropus*. Head depressed, tapering and at the same time narrowing

anteriorly, as in other species of *Tachysurus*; profile rising in a uniform, gently convex curve to occiput, where it becomes concave, owing to the more rapidly ascending outline of the occipital process. Eye low, but little above angle of mouth, the interorbital space decidedly convex. Barbels slender, the maxillary barbels reaching edge of gill membrane in front of pectoral spine, the outer mental barbels extending beyond gill membrane,  $1\frac{1}{2}$  in head, the inner not to edge of membrane. Gill membrane widely attached to isthmus, without free edge. Occipital region with very fine granulations, those on middle of occiput forming parallel series along the fontanel groove, those on median portion of occipital process in series which diverge backward from the median line. The sculptured area extends forward to a vertical which traverses the cheek at a distance of its own diameter behind the eye; anterior edge of granulated area equidistant between tip of snout and front of dorsal plate; fontanel produced backward as a deep, narrow groove, which fails to reach base of occipital process by a distance equaling  $\frac{1}{2}$  the length of the process on the median line; the groove widening but little anteriorly; an area behind and on each side of the groove with parallel series of granulations, and marked off from the rest of the head by a shallow trench; base of occipital process similarly indicated by a transverse indented line; occipital process not keeled, very wide at base, becoming abruptly very narrow behind, its posterior  $\frac{1}{2}$  having parallel margins and being as wide as long, the lateral margins therefore deeply concave; width of process at base equaling its length on median line, plus that of dorsal plate, its hinder edge deeply incised to receive the anterior rounded wedge process of the dorsal plate, the latter finely granulated anteriorly, the lateral wings concealed under the smooth skin; a narrow groove as long as eye occupies the anterior end of the fontanel; no similar groove found in *T. furthii*, a short roundish one present in the type of *T. liropus*, and a continuous one the entire length of fontanel in the specimen which we identify with *T. melanopus*; opercle without radiating ridges; a short, slit-like axillary pore present; humeral process short, the exposed portion not broadly triangular, the surface smooth, or indistinctly rough. Gill rakers 6+13, of moderate length and thickness, the longest below the angle,  $\frac{2}{3}$  diameter of eye. Dorsal spine with a series of obtuse granulations in front and very weak retrorse serræ behind, its length to tip of calcified portion  $1\frac{2}{3}$  in head; longest soft ray  $1\frac{2}{3}$  in head; adipose dorsal not adnate, its anterior insertion about over middle of anal; distance between dorsals equal to length of head; base of adipose dorsal much greater than its height, less than base of first dorsal; pectoral spine strong, ridged and granulated in front, the hinder edge with very strong serræ; length of spine  $1\frac{1}{2}$  in head, the fin projecting beyond tip of spine and reaching  $\frac{2}{3}$  distance from axil to base of ventrals; ventrals reaching to or nearly to origin of anal; vent midway between base of ventrals and front of anal; base of anal equaling length of pectoral spine; margin of anal gently concave, the longest ray  $2\frac{1}{3}$  in head; caudal with pointed lobes, the lower longest in the type,  $1\frac{1}{2}$  in head. Color dark steel blue or brownish above, becoming bright silvery below; posterior  $\frac{2}{3}$  of anal white.

the anterior portion black with a narrow white edge; pectorals and ventrals with anterior (outer) face white or slightly dusky; pectorals with inner face of upper rays black; a black blotch covers all of inner face of ventrals, except terminal half of inner rays; barbels blackish. Closely related to *T. melanopus* and *T. multiradiatus*, differing from the former in the longer anal fin, from the latter in the black markings on lower fins. The description of the type of *T. multiradiatus* (*Bagrus? arioides*) Kner & Steindachner, Abhandl. der K. Bayer. Akad. der Wissen., X, I, 1864, indicates a species with much rougher sculpturing of the head, a longer fontanel groove, narrower occipital process, and more anteriorly inserted adipose dorsal. The type is a single specimen, 280 mm. long, from Panama. (Gilbert.) (*ἔν*, in *μελανν*, ink.)

*Tachysurus emmelane*, GILBERT, Fishes Panama, MS. 1898, Panama.

201. TACHYSURUS FURTHII (Steindachner).

Head  $3\frac{1}{2}$  to  $3\frac{3}{4}$ ; depth 5 to  $5\frac{1}{2}$ . D. I, 7; A. 20. Body compressed posteriorly; profile slightly convex. Head broad, tapering forward, its greatest width  $1\frac{1}{2}$  to  $1\frac{1}{4}$  in its length; width, at the angle of the mouth,  $2\frac{3}{8}$  to  $2\frac{1}{2}$  in the head. Top of head densely covered with fine granules. Occipital process about as long as broad, with a median ridge, emarginate on its sides and at tip; interorbital region with 4 smooth ridges, the inner bordering the fontanel, the other extending obliquely backward from near the posterior nasal opening; sides of head and snout with vermiform mucous pores. Middle of the fontanel over the pupil. Eye strictly lateral, not entirely above the angle of the mouth, its center in front of the posterior end of the mandible, 2 in snout, 6 to 7 in head,  $3\frac{1}{2}$  to 4 in the distance between the eyes. Maxillary barbels thin, reaching to the middle of the pectoral or shorter, postmentals beyond base of pectorals, or sometimes not beyond edge of gill membrane; mentals to edge of gill membrane or shorter. Jaws about equal, the upper rather thin; teeth on the intermaxillaries villiform; the mandible with villiform teeth except the inner 2 or 3 series, which are granular; like the palatine patches irregular, suboval, sometimes the anterior end, sometimes the posterior, and sometimes both ends pointed. Gill membranes united, joined to the isthmus without a free margin. Gill rakers long and slender, 4 + 11. Axillary pore small; vertical series of pores present. Distance of dorsal from snout  $2\frac{1}{2}$  to  $2\frac{3}{8}$  in the length; the dorsal spine  $1\frac{1}{2}$  to  $1\frac{3}{4}$  in the head, on sides and front granular, with small, sharp teeth on its inner margin; the first soft ray little, if any, higher than the spine. Distance of adipose fin from the dorsal 3 to  $3\frac{1}{4}$  in the length, the fin longer than high, shorter than the dorsal fin. Caudal fin forked, the lobes rounded,  $4\frac{1}{2}$  in the length. Anal little longer than high, the highest ray 2 in the length of the head. Ventrals short,  $1\frac{1}{2}$  to  $2\frac{1}{2}$  in head. Pectoral spine long and slender,  $1\frac{1}{2}$  to  $1\frac{3}{4}$  in the head, outer margin granular, inner margin with short teeth. Ashy above, white below. We have examined 15 specimens, the largest measuring 0.29 m. The sexes do not differ externally. Panama. (Eigenmann & Eigenmann, Nematognathi, 90.)

202. *TACHYSURUS VARIOLOSUS* (Cuvier & Valenciennes), text, p. 132.

203. *TACHYSURUS MULTIRADIATUS* (Günther), text, p. 132.

75. *CATHOROPS*, Jordan & Gilbert, text, p. 133.

204. *CATHOROPS HYPOPHthalmus* (Steindachner), text, p. 133.

205. *CATHOROPS GULOSUS* (Eigenmann & Eigenmann), text, p. 133.

**Page 134.** After *Ictalurus furcatus* add:

206(a). *ICTALURUS ANGUILLA*, Evermann & Kendall.

(EEL CAT; WILLOW CAT.)

Head 4; depth  $4\frac{1}{2}$ ; eye 7 in head; snout  $2\frac{1}{2}$ ; interorbital  $1\frac{1}{2}$ ; maxillary (without barbel) 3; free portion of maxillary barbel longer than head; dorsal spine 2 in head; pectoral spine 2; width of mouth 2. D. I, 6; A. 24; vertebrae 42. Head large, broad, and heavy; the mouth unusually broad; cheeks and periocular portion of top of head very prominent; interorbital space flat, a broad, deep groove extending backward to origin of dorsal fin; body stout, compressed posteriorly; back scarcely elevated. Eye small; maxillary barbel long, reaching considerably past gill opening; other barbels short. Origin of dorsal fin equidistant between snout and origin of adipose fin, its distance from snout  $2\frac{3}{4}$  in length of body; base of dorsal fin  $3\frac{1}{2}$  in head; longest dorsal ray  $1\frac{1}{4}$  in head; dorsal spine strong, entire both before and behind; pectoral spine strong, entire in front, a series of strong, retrorse serrae behind; humeral process  $2\frac{1}{2}$  in pectoral spine; ventrals barely reaching origin of anal, their length 2 in head; anal fin long and low, the longest rays about  $2\frac{1}{2}$  in head; base of fin greater than head,  $3\frac{1}{2}$  in body; caudal moderately forked, the middle rays about  $2\frac{1}{2}$  in outer rays, which are about  $1\frac{3}{4}$  in head. Color uniform pale yellowish or olivaceous; no spots anywhere.

An examination of the 6 cotypes shows that there is not much variation, all the important characters remaining quite constant. The maxillary barbel varies somewhat in length, in some individuals scarcely reaching gill opening, and the number of anal rays varies from 24 to 26.

A comparison of the skull of this species with that of *I. furcatus* and of *I. punctatus* of the same size shows a number of very marked differences. Nearly all the bones in *I. anguilla* are heavier than in the other species; the supraoccipital is broadly triangular, and its upper surface finely grooved, while in each of the other species it is much longer and narrower and the upper surface nearly smooth.

From the blue cat (*Ictalurus furcatus*) this species differs chiefly in the fewer rays in the anal fin, the wider mouth, the shorter, heavier head, the much longer maxillary barbel, and in the cranial characters already given. From the spotted cat (*I. punctatus*) it may be distinguished by its wider mouth, more blunt snout, heavier head, the color, and the cranial characters already mentioned.

The eel cat rarely attains a greater weight than 5 pounds, and usually does not exceed 3 pounds. Its flesh is firm and of excellent flavor. The spawning season appears to be during the spring, as several of the individuals examined were in mature spawning condition.\* Lower Mississippi Valley; thus far known only from the Atchafalaya River, Louisiana and the Ohio River at Louisville. (*anguilla*, the generic name of the eel.)

<sup>1</sup> *alurus anguilla*, EVERMANN & KENDALL, Bull. U. S. Fish Comm. 1897 (Feb. 9, 1898), 125, pl. 6, fig. 1, Atchafalaya River, Louisiana. (Type, No. 48788. Coll. Evermann & Chamberlain.)

Recent studies of the catfishes of the Lower Mississippi Valley by Dr. Evermann have shown that the most abundant and most important species of catfish in that region is *Ictalurus furcatus* (Le Sueur), and not *Ameiurus lacustris* (Walbaum), as has hitherto been supposed. The large specimen described by Dr. Bean as *Ameiurus ponderosus* is an *Ictalurus* (as shown by the skeleton now in the United States National Museum) and apparently *I. furcatus*. The common names "Great Fork-tailed Cat," "Mississippi Cat," and "Blue Cat" all belong to *I. furcatus*.

**Page 138.** The species called *Ameiurus dugesii* belongs to the genus *Villarius*, Rutter.

**Page 142.** After *Ameiurus nigrilabris* add:

77(a). VILLARIUS, Rutter.

*Villarius*, RUTTER, Proc. Cal. Ac. Sci., ser. 2, vol. VI, 1896, 256 (pricei).

Allied to *Ameiurus*, differing in the presence of scattered cilia on the sides. Backward process from occipital short, broad, emarginate, connected by ligament with the first interspinal buckler; in adults the distance between this process and the buckler is equal to the length of the former; in young examples the process overlaps the keel on the underside of the buckler. Head narrow, width of intermaxillary band of teeth  $\frac{1}{2}$  of head; caudal deeply forked, the upper lobe the longer; barbels long, those of the maxillary extending past the gill opening. Sides with scattered hair-like cirri; these are very noticeable under a lens, but not readily distinguished by the naked eye. This genus differs from all others of the family in having hair-like cirri on the sides. It differs from *Ictalurus* in having the occipital process and the interspinal buckler widely separated and connected by ligament; from *Ameiurus* in having a narrow head and a deeply forked caudal. Two species known, the following and *Villarius dugesii* (Bean). (*villus*, a hair.)

\* This species is well known to the fishermen of the Atchafalaya River, by whom it is usually called the "eel cat," though the name "willow cat" is sometimes applied to it. It was explained by the fishermen that the name "eel cat" was given on account of the long feelers (i. e., barbels) and the name "willow cat" because it is most frequently found about the roots of willow trees. The eel cat is not an abundant species in the Atchafalaya River. During six days (April 19-24) spent at Morgan City several hundred catfish were examined at the three fish houses, and the total number of eel cats seen was fewer than twenty-five. The fishermen report that this proportion is about as great as at any time of the year. Of the four commercial species of catfishes handled on this river the most abundant one is the blue cat (*Ictalurus furcatus*), and the next is the yellow cat or goujon (*Leptops oivaris*); the eel cat comes next and the spotted cat (*Ictalurus punctatus*) last. The blue cat and the yellow cat probably constitute 98 per cent of the entire catch.

## 220(a). VILLARIUS PRICEI, Rutter.

B. 8; D. I, 6; A. 22 or 23; C. 17; P. I, 9; V. 8. Head  $3\frac{1}{4}$  to  $3\frac{3}{4}$  in body; eye 5 to 7 in head; snout  $2\frac{3}{8}$ ; maxillary  $5\frac{1}{2}$  to 6. Maxillary barbel very long, reaching beyond the pectoral spine, in the adult about to its tip when depressed, 3 to 4 times as long as the barbel at nostril. Origin of dorsal midway between snout and middle of base of adipose fin; pectorals inserted halfway between snout and ventrals; longest dorsal ray 6 to 7 times in length of body; spine of dorsal longer than its base, equal to base of adipose fin; longest pectoral ray about half of head, pectoral spine  $2\frac{1}{2}$  to 3 in head, with about 12 distinct hooked serræ behind, these fewer and somewhat smaller in the young; base of anal 3 times in its distance from snout, its longest ray equal to length of ventral; caudal deeply forked. Lateral line faint. This species differs from *V. dugesii* (Bean) in having very prominent serrations on the pectoral spines, the types of *dugesii* having the pectoral spines without serræ. We have examined a specimen of *dugesii*, 4 inches long, from Salamanca, Mexico, which is in the type basin; it has the cirri minute and light in color, a row of papillæ along the lateral line, and the pectoral spines with 4 or 5 degenerate serræ. (Named for William Wightman Price, who collected the type specimen.)

*Villarius pricei*, RUTTER, Proc. Cal. Ac. Sci., ser. 2, vol. VI, 1896, 257, San Bernardino Creek, a tributary of the Yaqui River, southern Arizona. (Type, No. 4826, L. S. Jr. Univ. Mus.)

**Page 143.** *Leptops olivaris* is known as the *Goujon* in Louisiana, where it is an important food-fish.

**Page 146.** In *Schilbeodes gyrynus* the anal rays are 14 to 16; not 13.

**Page 152.** Under *Rhamdia salvinii* read "Osbert Salvin" for "Oscar Salvin."

**Page 170.** *Pantosteus arizonæ*, Gilbert, is described and figured in Proc. U. S. Nat. Mus. 1898, 488, pl. 36.

**Page 174.** *Catostomus discobolus* is distinct from the true *C. latipinnis*. The two species are confused in the description of *C. latipinnis* given by us. They may each be described as follows:

## 279. CATOSTOMUS LATIPINNIS, Baird &amp; Girard.

Head 4; depth about  $5\frac{1}{2}$ ; eye high up and small, 5 to 7 in head, 3 to  $3\frac{1}{2}$  in snout,  $2\frac{1}{8}$  to  $2\frac{3}{8}$  in interorbital space; interorbital width  $2\frac{3}{8}$  in head. D. 14 or 15; A. 7; scales 19 or 20-89 to 102-16 to 18, 46 to 50 transverse rows in front of dorsal fin. Head depressed and flat above, its greatest depth  $1\frac{1}{2}$  in its length, the depth below lower edge of orbit 3 in its length. Least depth of caudal peduncle  $4\frac{1}{2}$  in head, or  $3\frac{1}{2}$  in its own length. Fins very large, the dorsal with its upper margin concave; ventrals and pectorals rounded; dorsal as long as its longest ray,  $1\frac{1}{10}$  in head, its last ray a little less than  $\frac{1}{2}$  the length of the first ray; origin of dorsal fin nearer tip of snout than base of caudal; ventrals not reaching quite to vent,  $1\frac{1}{8}$  in head. Muzzle not projecting; about 6 rows of short, thick papillæ on upper lip, the smallest above; lower lip large, incised to its base, with

about 12 rows of short, thick papillae, posteriorly quite small; distance from front of upper lip to back of lower  $1\frac{1}{2}$  in snout; jaws with a slight cartilaginous sheath; width of preorbital a little less than  $\frac{1}{2}$  its length. Reaches a length of about 2 feet. Lower Colorado River basin. This description by Gilbert & Scofield, based upon specimens from the Gila River at Tempe, Arizona.

279(a). CATOSTOMUS DISCOBOLUS, Cope.

Head  $3\frac{3}{4}$  to  $4\frac{1}{4}$ ; depth about  $5\frac{1}{4}$ ; eye small, high up,  $5\frac{1}{2}$  to 6 in head,  $2\frac{3}{4}$  in snout,  $2\frac{3}{4}$  in interorbital width, which is  $8\frac{1}{2}$  in head; width of preorbital less than  $\frac{1}{2}$  its length; least depth of caudal peduncle  $2\frac{1}{2}$  in its length, or 2 in head; greatest depth of head  $1\frac{3}{8}$  in its length; depth from lower edge of orbit  $3\frac{1}{4}$  in head. D. 12 or 13; A. 7; scales 19 to 21—101 to 109—17 to 21, 52 to 63 in front of dorsal. Muzzle projecting slightly beyond upper lip. Upper margin of dorsal very slightly concave, the length of its base  $1\frac{1}{2}$  in its longest ray, or  $1\frac{1}{2}$  in head; last dorsal ray  $\frac{1}{2}$  length of first; origin of dorsal midway between tip of snout and base of caudal; ventral rounded,  $1\frac{3}{8}$  in head, not quite reaching vent. Mouth as in *C. latipinnis* except that the posterior tubercles on lower lip are long and not nearly so closely set, there being 9 or 10 rows; jaws with a slight cartilaginous pellicle. Upper portion of the Colorado River basin. Attains the length of a foot or more. The above description by Gilbert & Scofield from specimens from Green River at Green River Station, Wyoming.

*Catostomus discobolus*, COPE, Hayden's Geol. Surv. Wyo., 435, 1870, Green River, Wyoming; GILBERT & SCOFIELD, Proc. U. S. Nat. Mus. 1898, 490.

Page 175. After *Catostomus griseus* add:

280(a). CATOSTOMUS RETROPINNIS, Jordan.

A doubtful species which is, however, not yet shown to be invalid. Head  $4\frac{1}{2}$ ; depth  $5\frac{1}{2}$ ; eye  $6\frac{1}{2}$  in head; snout 2; interorbital  $2\frac{1}{2}$ . D. 11; A. 7; scales 17—108—14. Body slender, head slender, snout very long, caudal peduncle long, its least depth less than snout,  $2\frac{1}{2}$  in head; dorsal profile very little elevated; mouth large, wholly inferior, overhung by the pig-like projecting snout; lips thin but very broad, lower lip incised nearly to base, with about 6 rows of moderate papillae; lobes of lower lip very long, about  $\frac{1}{2}$  of snout; gill rakers short and weak. Origin of dorsal a little nearer base of caudal than tip of snout; base of dorsal equal to snout; longest dorsal ray a little greater than base of fin; anal fin long and pointed, the fourth ray longest,  $1\frac{1}{10}$  in head; caudal lunate, the middle ray  $1\frac{1}{2}$  in outer rays; pectoral somewhat falcate, the longest  $1\frac{1}{2}$  in head; ventrals rather short, not reaching vent by more than an eye's diameter. Length 14 inches.

*Catostomus retropinnis*, JORDAN, Bull. U. S. Nat. Mus., XII, 178, 1878, Milk River, Montana. (Type, No. 21197. Coll. Dr. Elliott Coles.)



Page 176. After *Catostomus catostomus* add:

282(a). *CATOSTOMUS RIMICULUS*, Gilbert & Snyder.

Head  $4\frac{1}{2}$  in body; depth 5; depth of caudal peduncle  $2\frac{1}{2}$  in head; eye  $7\frac{1}{2}$ ; dorsal rays 11; scales 18-91-13, before dorsal 42. D. 11; A. 7; pectoral 17. Head as deep as wide. Both lips full, the lobe of lower lip broadly rounded behind, the cleft not nearly reaching base of lip, the portion between mandible and apex of cleft with 4 series of tubercles; tubercles coarse and blunt, becoming reduced in size toward margins of lips, but less so than in related species; upper lip with 5 rows of tubercles. Eyes very small, the front of the eye nearly midway of head; interorbital space convex,  $2\frac{1}{2}$  in head. Scales comparatively smooth, gradually growing smaller posteriorly. Dorsal fin inserted midway between end of snout and base of caudal, first ray preceded by 2 short, simple ones; last ray divided to base; length of base of fin equal to the height, which is contained  $6\frac{1}{2}$  times in the body; height of anal twice the length of the base, 5 in body; length of pectorals  $4\frac{1}{2}$  in body; ventrals  $6\frac{1}{2}$  in body; caudal  $4\frac{1}{2}$ . Color above dusky, the central parts of scales lighter; under parts white; dorsal and caudal fins dusky, others white. This species belongs to the *C. catostomus* type, with very small scales, and is most nearly related to *C. tahoensis*. From the latter it differs in the smaller eye, less deeply cleft lower lip, blunter labial tubercles, larger scales, and the much smaller fontanel, which is reduced in adults to a very narrow linear slit, or more commonly entirely obsolete. Lower portion of the Klamath River basin, northern California. (Diminutive of *rimus*, crevice, from the small fontanel.)

*Catostomus rimiculus*, GILBERT & SNYDER, Bull. U. S. Fish Comm. 1897 (Jan. 6, 1898), 3, Trinity River, Humboldt County, California. (Type, No. 5634, L. S. Jr. Univ. Mus. Coll. Capt. W. E. Dougherty.)

Page 177. *Catostomus rex* is identical with *Deltistes luxatus* and should be added to the synonymy of that species, p. 183.

The type of *Catostomus labiatus* did not come from Klamath Lake, but from the Sacramento River, at Stockton, California. It is identical with *C. occidentalis*. The species from Klamath Lake has been recently described as

285. *CATOSTOMUS SNYDERI*, Gilbert.

Head  $4\frac{1}{2}$  in length; snout  $2\frac{3}{4}$  in head, equaling interorbital width; eye  $5\frac{1}{2}$ . D. 11; A. 7; scales 13 or 14-69 to 77-10 or 11. Mouth very small, the width between angles but  $\frac{1}{2}$  length of snout in our largest specimen; greatest width of lobe of lower lip  $\frac{2}{3}$  diameter of eye; lower lip deeply incised, with 1 or 2 papillae between symphysis and base of cleft; upper lip narrow, with 5 or 6 papillae in a cross series, the uppermost becoming very small; basal portion of the lower lip with coarse tubercles, those toward posterior margin becoming very fine and arranged in evident series separated by grooves. Mucous canals on head forming conspicuous raised ridges with prominent pores, the system much more conspicuously developed than in any related species. Origin of dorsal fin constantly

nearer snout than base of caudal; the dorsal fin short, its base not exceeding the height of the longest ray, usually less. In our specimens the pectorals reach scarcely  $\frac{3}{8}$  distance to ventrals and the ventrals scarcely  $\frac{3}{8}$  distance to vent; the anal may extend beyond base of rudimentary caudal rays. Scales strongly ridged, their margins crenate; the anterior scales are smaller, but do not appear greatly crowded; the average number of tubes in the lateral line is about 73, the number varying from 69 to 77. Dusky, the lower part of sides with coarse black specks, the under parts white; fins all dusky. (Gilbert.)

A larger specimen has been described as follows: Head  $4\frac{1}{5}$ ; depth 4; eye  $6\frac{1}{2}$  in head; snout  $2\frac{1}{2}$ ; maxillary  $3\frac{1}{2}$ ; mandible  $2\frac{1}{2}$ ; interorbital  $2\frac{1}{2}$ ; width of mouth  $3\frac{1}{2}$  in head, more than  $\frac{1}{2}$  length of snout; greatest width of lower lip  $\frac{1}{2}$  diameter of eye. D. II, 11; A. 7; scales 13-70-11. Body rather slender; head long, mouth moderate, horizontal; lips thick papillose, the upper with about 4 or 5 rows of papillæ, lower with about 7; lower lip divided nearly to base, leaving only 1 row of papillæ crossing the symphysis; premaxillary not much projecting and not forming a prominent hump; maxillary rather short, not reaching vertical at front of anterior nostril; eye equally distant between snout and posterior edge of opercle; mucous canals on head forming raised ridges, the pores conspicuous. Fins moderate; origin of dorsal a little nearer snout than base of caudal, sixth spine over insertion of ventrals; pectoral  $1\frac{1}{2}$  in head, reaching slightly more than  $\frac{3}{8}$  distance to ventrals; ventrals not quite reaching vent, the seventh ray longest,  $1\frac{3}{4}$  in head; anal long, pointed, reaching to base of caudal,  $1\frac{1}{5}$  in head. Scales crowded anteriorly, about 32 transverse rows in front of dorsal, strongly ridged, the margins crenate. (Evermann & Meek.) Length 1 to 2 feet. Klamath Lakes, Oregon; specimens examined from Upper Klamath Lake, Lost River, and Williamson River. (Named for Mr. John O. Snyder, instructor in Zoology in Stanford University.)

*Catostomus snyderi*, GILBERT, Bull. U. S. Fish Comm. 1897 (Jan. 6, 1898) 3. Upper Klamath Lake, Oregon (Type, No. 48222. Coll. Gilbert, Cramer & Otaki); EVERMANN & MEEK, Bull. U. S. Fish Comm. 1887, 69.

Page 178. After *Catostomus occidentalis* add:

286(a). *CATOSTOMUS TSILTCOOSENSIS*, Evermann & Meek.

Head  $4\frac{1}{2}$ ; depth 5; eye  $6\frac{1}{2}$  in head; snout 2. D. 13; A. 7; scales 13-65-8, 34 before the dorsal. Pectoral  $1\frac{1}{2}$  in head; longest dorsal ray  $1\frac{1}{2}$ ; base of dorsal  $1\frac{1}{2}$ ; longest anal ray  $1\frac{1}{2}$ ; ventral  $1\frac{1}{2}$ . Body rather slender, subterete; head small, snout long and pointed; mouth inferior, overhung by the projecting snout; lips rather thin, 1 row of large papillæ on upper lip, and about 2 irregular rows of smaller ones behind or inside of it; lower lip incised nearly to base, 1 or 2 rows of small papillæ across the isthmus; lobes of lower lip moderately long and thin, the bases with papillæ merging into plications toward the tips. Eye quite small, the anterior edge of orbit at middle of head. Top of head flat or very slightly convex between the eyes. Fins small; pectorals short and rounded; ventrals short, rounded,

the middle rays but little longer than the others; anal small, somewhat pointed; margin of dorsal somewhat concave; caudal lunate, not deeply forked. Muciferous canals on head not strongly developed. Scales moderately large; lateral line nearly straight, not running upward toward nape. This species differs from *C. occidentalis*, to which it is related, in the smaller head, longer, more pointed snout, smaller eye, larger scales, and its much smaller fins. In *C. occidentalis* the pectoral fins are falcate, while in this species they are more rounded; the ventrals also are less pointed. Length a foot or less. Coastal streams of middle western Oregon; known from Tsiltcoos Lake and the Siuslaw River. (*tsiltcoosensis*, from the type locality.)

*Catostomus tsiltcoosensis*, EVERMANN & MEEK, Bull. U. S. Fish Comm. 1897 (Jan. 6, 1898), 68, fig. 1, Tsiltcoos Lake, Lane County, Oregon. (Type, No. 48479. Coll. Dr. Seth E. Meek.)

**Page 180.** The species called *Catostomus fecundus* in the text belongs in the genus *Chasmistes*, to which it should be transferred as *Chasmistes fecundus* (Cope & Yarrow).

**Page 182.** The species of *Chasmistes* are not confined to the Great Basin. One species (*C. brevirostris*) occurs in the Klamath Lakes basin.

**Page 183.** The species called *Chasmistes luxatus* in the text belongs to a genus distinct from *Chasmistes*, which may be characterized as follows:

93(a). DELTISTES, Seale.

*Deltistes*, SEALE, Proc. Cal. Ac. Sci., ser. 2, vol. VI, 1896, 269 (*luxatus*).

This genus is close to *Chasmistes*, agreeing with it in every respect except in the peculiar structure of the gill rakers. In *Chasmistes* they are as in *Catostomus*, while in *Deltistes* they are broad, shaped like the Greek letter Δ (delta), and their edges are unarmed and entire. Lower pharyngeals weak, with numerous small teeth. *Deltistes luxatus* (Cope) is the single known species. (δέλτα, the Greek letter corresponding to D.)

After *Chasmistes cujus* add:

297(a). CHASMISTES STOMIAS, Gilbert.

Head  $4\frac{1}{2}$ ; depth  $4\frac{1}{2}$ ; eye 7; snout  $2\frac{3}{4}$ ; maxillary (measured from free end to tip of snout)  $3\frac{1}{2}$ ; mandible  $2\frac{1}{4}$ . D. II, 11; A. I, 7; scales 13-85-10; inter-orbital width  $2\frac{1}{4}$ ; vertical depth of head at mandibular articulation  $2\frac{1}{2}$ . Head small, body heavy forward, the back strongly and regularly arched from snout to origin of dorsal fin, thence declined in a nearly straight line to base of caudal; ventral surface nearly straight. Premaxillary spines strongly protruding, forming a prominently projecting snout; mouth rather small, inclined upward at an angle of about  $40^\circ$ , maxillary scarcely reaching vertical from front of anterior nostril; width of mouth  $1\frac{3}{4}$  in snout or  $4\frac{3}{4}$  in head; upper lip thin, without papillae; lower lip thin, interrupted at symphysis, forming narrow lateral lobes, the width of which is about  $2\frac{3}{4}$  times in their length; faint indications of a few papillae; mucous canals forming ridges, the pores conspicuous; gill rakers long,

narrowly triangular at the tip when viewed from behind, densely tufted on the anterior edge; fontanel narrow, its length  $2\frac{3}{4}$  in the snout, its width about  $\frac{1}{2}$  its length. Fins all large; the origin of the dorsal a little nearer tip of snout than base of caudal, the sixth ray over base of ventral, its base  $1\frac{3}{4}$  in head, the free edge nearly straight, the last ray  $1\frac{3}{4}$  in the first, which is  $1\frac{1}{2}$  in head; pectorals scarcely falcate, reaching a little more than  $\frac{3}{8}$  distance to base of ventrals, their length  $1\frac{1}{4}$  in head; ventrals long, reaching vent, the rays gradually increasing in length from the outer to the seventh and eighth, which are longest, the ninth and tenth being but slightly shorter, the length of the longest ray  $1\frac{3}{4}$  in head or about  $\frac{1}{2}$  longer than the first; anal long and pointed, the fourth ray longest, reaching base of caudal,  $1\frac{1}{2}$  in head; each ray of anal fin with 8 to 12 strong tubercles; caudal lobes about equal, their length  $1\frac{1}{4}$  times the middle ray. Length a foot or more. Upper Klamath Lake, Oregon, where it is abundant and of some importance as a food-fish. The Klamath Indian name is K-ahp-tu. (*στρούαξ*, large-mouthed.)

*Chasmistes stomias*, GILBERT, Bull. U. S. Fish Comm. 1897 (Jan. 6, 1898), 5, with plate, Upper Klamath Lake, Oregon (Type, No. 48223. Coll. Gilbert, Cramer & Otaki); EVERMANN & MECK, Bull. U. S. Fish Comm. 1897, 70.

297(b). CHASMISTES COPEI, Evermann & Meek.

Head  $3\frac{3}{4}$ ; depth 4; eye  $6\frac{1}{2}$ ; snout  $2\frac{1}{2}$ ; maxillary (measured from free end to tip of snout) 3; mandible  $2\frac{3}{4}$ . D. II, 10; A. I, 7; scales 13-80-12; inter-orbital width  $2\frac{1}{4}$ ; vertical depth of head at mandibular articulation  $2\frac{1}{2}$ . Head large, cheek very deep, the depth equal to distance from tip of snout to nostril; body stout, back scarcely elevated, caudal peduncle rather short and stout; ventral surface somewhat convex. Premaxillary spines less protruding than in *C. stomias*, not forming a prominent hump; mouth large, inclined upward at an angle of  $45^\circ$ , maxillary not nearly reaching vertical at front of anterior nostril; width of mouth  $1\frac{3}{4}$  in snout, or 4 in head; upper lip thin, without papillae; lower lip thin, entirely without papillae, interrupted at symphysis, forming rather broad lateral lobes; pores on head very conspicuous; gill rakers larger than in *C. stomias*, broadly triangular at tip when viewed from behind, densely tufted on anterior edge, each appendage more or less bifid and club-shaped, closely resembling those of *C. liorus*; fontanel narrow, its length  $2\frac{1}{2}$  in snout, width  $\frac{1}{2}$  its length. Fins all small; origin of dorsal a little nearer snout than base of caudal, its sixth ray over base of ventrals, free edge straight, base  $2\frac{1}{2}$  in head, last ray a little less than 2 in first, which is 2 in head; pectorals somewhat falcate, reaching slightly more than  $\frac{1}{2}$  distance to ventrals, their length  $1\frac{3}{4}$  in head; ventrals very short, reaching only  $\frac{3}{8}$  distance to vent, free end nearly straight; outer ray longest,  $2\frac{7}{8}$  in head; inner shortest,  $3\frac{1}{4}$  in head; anal fin short, bluntly pointed, not reaching base of caudal, third and fourth rays longest,  $1\frac{3}{4}$  in head; no tubercles on anal rays; caudal lobes equal, length about  $1\frac{1}{4}$  times the middle ray. Scales small and crowded anteriorly, about 14 rows downward and backward from front of dorsal to lateral line, 11 vertically upward from base of ventral to lateral line, about 38 oblique series before

dorsal; lateral line nearly straight, with about 80 scales. Entire upper parts of head and body, and sides nearly to level of base of pectorals, dark olivaceous; under parts abruptly whitish or yellowish in alcohol; a dark spot in upper part of axil; dorsal and caudal dark; pectorals dark on inner surface; ventrals and anal plain. From *Chasmistes stomias* this species is readily distinguished by its larger head, larger, more oblique mouth, less prominent snout, and very small fins. The differences in the fins are very great, particularly in the ventrals. It differs from *C. brevirostris*, as characterized by Dr. Gilbert, in its much larger, more oblique mouth, the absence of papillæ on the lips, and shorter fins. Length 2 feet. Upper Klamath Lake, Oregon. Klamath Indian name "Tswani." (Named for the late Prof. Edward Drinker Cope, who wrote the first paper on the fishes of Upper Klamath Lake.)

*Chasmistes copei*, EVERMANN & MEEK, Bull. U. S. Fish Comm. 1897 (Jan. 6, 1898), 70, fig. 3, Pelican Bay, Upper Klamath Lake, Oregon. (Type, No. 48224. Coll. Meek & Alexander.)

**Page 205.** *Campostoma pricei* can not be distinguished by us from *C. ornatum*. See Rutter, Proc. Cal. Ac. Sci., ser. 2, vol. VI, 1896, 259.

**Page 211.** After *Algansea tinella* add:

337(a). ALGANSEA TARASCORUM, Steindachner.

Head  $3\frac{3}{8}$ ; depth  $4\frac{2}{3}$ ; eye less than 5; snout about 4; interorbital 3. D. III, 7; A. III, 6; P. 17; V. 9; scales 84 or 85, 18 or 19. Body stouter than in *A. lacustris*, head shorter, lateral line more decurved and nearer ventral line at middle of body, and scales more numerous. Mouth very oblique, lower jaw not projecting, maxillary not quite reaching vertical at anterior edge of eye. Teeth 4-4, hooked, and with narrow grinding surface. Origin of dorsal in advance of ventrals, equally distant between base of caudal and middle of eye; height of dorsal twice its base. Ventrals not reaching anal fin by an eye's diameter; caudal deeply notched. A dark gray longitudinal band with metallic luster extending from opercle to caudal fin, lying chiefly above lateral line; color otherwise plain. Length  $5\frac{1}{2}$  inches. Lake Pátzcuaro, Mexico. (Steindachner.)

*Algansea tarascorum*, STEINDACHNER, Einige Fischarten Mex., II, pl. 3, figs. 2-2c, 1895, Lake Pátzcuaro, Mexico. (Coll. Princess Theresa von Bayern.)

**Page 218.** To the synonymy of *Pimephales notatus* add:

*Spinicephalus fibulatus*, LE SUEUR, in VAILLANT, Bull. Soc. Philom., VIII, 1896, 29, pl. 26.

**Page 225.** After *Ptychocheilus oregonensis* add:

358. PTYCHOCHEILUS GRANDIS (Ayres).

(SACRAMENTO PIKE.)

This species differs from *P. oregonensis* principally in the larger size of the scales above the lateral line, the smaller number of rays in the dorsal fin, and the lighter and slenderer pharyngeal bones. Head  $3\frac{3}{8}$  to  $3\frac{1}{2}$  in length; depth 5 to  $5\frac{1}{10}$ ; eye  $3\frac{3}{8}$  to 4 in head; scales 13 to 16 above lateral

line, 70 to 80 transverse rows along lateral line (16 to 18 above lateral line, 69 to 72 transverse rows in *P. oregonensis*). D. 8; A. 8. In other respects similar to *P. oregonensis*. *Ptychocheilus harfordi* is apparently not distinct from *P. grandis*, being based on a specimen with very small scales. *P. grandis* is confined to waters of California, *P. oregonensis* to Washington, Idaho, and Oregon.

*Gila grandis*, AVRES, Proc. Cal. Ac. Nat. Sci. 1854, 18, San Francisco.

*Ptychocheilus major*, AGASSIZ, Am. Jour. Sci. Arts 1855, 229, San Francisco.

*Ptychocheilus harfordi*, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1881, 72, Sacramento River (Type, No. 27246. Coll. Jordan & Gilbert); JORDAN & GILBERT, Synopsis, 226 1883.

Page 239. After *Leuciscus balteatus* add:

376(a). LEUCISCUS SIUSLAWI, Evermann & Meek.

Head  $4\frac{1}{2}$ ; depth  $4\frac{1}{2}$ ; eye 4; snout  $3\frac{1}{2}$ ; maxillary  $3\frac{2}{5}$ . D. II, 9; A. II, 12 or 13; scales 11-58-8; teeth 2, 4-5, 2, somewhat hooked. Body rather slender, slightly elevated and somewhat compressed; head small and pointed, cheek not deep; snout pointed, somewhat longer than eye; mouth moderate, somewhat oblique, maxillary just reaching vertical at front of orbit; jaws subequal, the lower sometimes slightly projecting; eye large, not as great as snout. Origin of dorsal fin behind base of ventrals and much nearer base of caudal than tip of snout, the longest ray  $1\frac{1}{2}$  in head, greater than base of fin; origin of anal fin under last dorsal ray but 2, its height equal to that of dorsal, its base equal to its longest ray; free edges of dorsal and anal nearly straight; pectoral  $1\frac{1}{2}$  to  $1\frac{1}{4}$  in head, not reaching insertion of ventrals; ventrals short,  $1\frac{1}{4}$  in pectoral, reaching anus; caudal deeply forked; lateral line complete, decurved. Color in spirits, brownish or olivaceous above, middle of side with a broad dark band involving the lateral line anteriorly and posteriorly, but lying chiefly above it mesially; middle of side from gill opening to beneath dorsal fin with a broad rosy band, following closely beneath the lateral line; lower part of sides and under parts silvery, dusted over with fine dark specks; a light yellowish band extending backward from upper posterior border of eye nearly halfway to origin of dorsal fin; cheek with a silvery or golden crescent; top of head dark; opercles dusky silvery; snout dusky; fins plain, dorsal and caudal somewhat dusky. This species is close to *L. balteatus*, but has smaller anal and dorsal fins, a more slender body, smaller and more slender head, and longer, more pointed snout. The extent of variation in proportional measurements and in the number of anal fin rays appears to be much less than in *L. balteatus*. It also resembles *L. cooperi*, but has a much shorter lower jaw and a more pointed snout. Known only from the Siuslaw River and Tsiltcoos Lake, western Oregon, where it is common. (*siuslawi*, of the Siuslaw River.)

*Leuciscus siuslawi*, EVERMANN & MEEK, Bull. U. S. Fish Comm. 1897 (Jan. 6, 1898), 72, fig. 4, Siuslaw River, Mapleton, Oregon. (Type, No. 48480. Coll. Dr. Meek.)

Page 240. After *Leuciscus elongatus* add:

378(a). *LEUCISCUS NACHTRIEBI*, Cox.

Head  $4\frac{1}{2}$  to  $4\frac{3}{4}$ ; depth 5 ( $4\frac{1}{2}$  to  $5\frac{1}{2}$ ); eye 4; snout  $4\frac{3}{4}$ . D. 8; A. 8. Body rather heavy, not greatly compressed; back slightly elevated, its curve a little greater than that of the belly; caudal peduncle rather stout, its depth  $\frac{1}{2}$  the length of the head. Head rather short, not any more compressed than the body, upper surface slightly flattened; snout quite blunt in mature individuals, its length  $1\frac{1}{2}$  times width of eye; mouth not very large, but little oblique, lower jaw included; maxillary scarcely reaching to front of orbit; pharyngeal teeth 2, 4-5, 2. Dorsal fin inserted nearer base of caudal than tip of snout, also slightly back of ventrals; caudal fin forked; anal slightly smaller than dorsal; ventrals small, not reaching vent by  $\frac{1}{2}$  their length; pectorals inserted rather high, not reaching the ventrals by  $\frac{1}{4}$  their length; scales small, 12-72-9, lateral line complete on mature individuals, decurved, the pores extending on head in several lines, 1 passing back of eye, another down to nostril. General color dusky, darkest on back; sides above lateral line dull silvery, below lateral line light silvery; a faint dark dorsal band in some specimens, in others absent; no black lateral band, but some specimens have a very faint dusky shade along lateral line; no light stripe above lateral line; upper portion of opercles with a dusky shade, lower part bright silvery; upper part of head dark-colored; all the above colors typical in the young as well as adults. Length 4 inches. *L. nachtriebi* differs from *L. neogvus* in having a well-developed lateral line, a smaller eye, fewer scales, less oblique mouth, a shorter maxillary, and in being a larger fish and differently colored. It differs from *L. elongatus* in having a smaller mouth, the lower jaw never projecting, head less pointed, a shorter maxillary, finer scales, and the absence of the black lateral band. Lakes of northern Minnesota; at present known from Mille Lacs, Man Trap, Mud and Elbow lakes. (Named for Prof. Henry F. Nachtrieb, State zoologist of Minnesota.)

*Leuciscus nachtriebi*, Cox, Rept. U. S. Fish Comm. 1894 (Dec. 14, 1896), 615, Mille Lacs Lake, Aikin County, Minnesota. (Type, No. 47688. Coll. Minn. Nat. Hist. Surv.)

**Page 241.** To the synonymy of *Leuciscus neogvus* add:

*Oyprinus burtonianus*, LE SUEUR in VAILLANT, Bull. Soc. Philom., VIII, 1896, 28, with plate, Burton Mine, Missouri.

**Page 244.** *Leucos* and *Myloleucus* can not be maintained as subgenera, the characters of the teeth not being constant.

The following notes on *Rutilus olivaceus* as seen at Emerald Bay, Lake Tahoe, may prove useful.

385. *RUTILUS OLIVACEUS* (Cope).

(TAHOE CHUB.)

This species is very different from *Rutilus symmetricus*, looking like *Leuciscus lineatus*. Very common; reaches 2 to 3 pounds weight; devours eggs of trout. No doubt the records of *Leuciscus lineatus (atarius)* from Lake Tahoe belong to this species. Head 4; depth  $4\frac{1}{2}$  to  $4\frac{3}{4}$ . D. 8; A. 8; scales 11-56-6; teeth always 5-5, with broad grinding surface. Body oblong,

moderately compressed, the back somewhat elevated anteriorly in old examples. Head conical, rounded above; eye moderate,  $1\frac{1}{2}$  in snout (6 inches long), 5 in head; about as long as maxillary. Mouth terminal, very oblique, the lower jaw included; the snout not prominent; the short maxillary not reaching eye. Dorsal high and pointed; anal short, rather high; pectoral long, reaching  $\frac{2}{3}$  distance to ventrals, which reach vent; ventrals below front of dorsal, which is behind middle of body. Scales with edges largely exposed; lateral line running low, complete. Dusky olive above and on sides to level of ventrals, with brassy luster everywhere; middle of belly only white, a pale yellowish area between pectorals and ventrals; head brassy, dusky above, closely dotted above and on sides; body everywhere closely dotted with black, except on middle line below; fins all dusky, with dark points. This species is well separated from all the *R. symmetricus* tribe.

**Page 247.** *Luxilinus occidentalis* is the young of *Larinia exilicauda*, Baird & Girard (p. 209), and must be placed in the synonymy of that species. *Luxilinus* is a pure synonym of *Larinia*.

**Page 249.** Under *Opsopæodus bollmani*, for "Buckland Creek" read "Buckhead Creek."

**Page 254.** For *Azteca*, line 22, substitute *Aztecula*, Jordan & Evermann, new subgenus. The former name is preoccupied by *Azteca*, Forel, 1878, a genus of ants. The same substitution to be made in the key on page 255 and on page 258.

**Page 260.** Before *Notropis cayuga* insert:

404(a). NOTROPIS WELAKA, Evermann & Kendall.

Head  $4\frac{1}{2}$ ; depth 5; eye 3 in head; snout  $3\frac{1}{2}$ . D. 8; A. 8 or 9; scales 6-35-3; teeth 4-4, hooked. Body rather slender, moderately compressed; head short, snout bluntly pointed; mouth moderate, somewhat oblique, lower jaw slightly included, maxillary scarcely reaching front of eye; premaxillaries protractile. Eye large; posterior edge of pupil at middle of longitudinal length of head; interorbital width greater than eye; caudal peduncle long and slender. Dorsal fin inserted well behind base of ventrals, a little nearer base of caudal than tip of snout, its longest rays shorter than head, but slightly longer than longest anal rays; anterior dorsal and anal rays longest; pectoral  $1\frac{1}{2}$  in head; ventrals reaching origin of anal; caudal deeply notched, the lobes long and pointed. Scales large, lateral line incomplete, developed only on 6 to 10 scales. Back olivaceous; side with a broad black band extending from snout through eye, and ending in a rather distinct black spot on base of caudal, the black spot in some specimens (probably mature males) surrounded by orange; the black line bordered above by a narrow orange or reddish line, less distinct, or even whitish, in females and immature individuals; under parts plain; fins all plain; dorsal and caudal somewhat dusky; dusky specks on body along base of anal and under side of caudal peduncle; lower jaw tipped with dusky. This species resembles *Notropis anogenus*, but differs in having the mouth somewhat larger and less oblique,



the lower jaw more included, the body more slender, the lateral line less developed, the dorsal fin more posterior, and the anal rays more numerous. It was found in considerable abundance in the St. Johns River, near Welaka, Florida. (*welaka*, from the type locality.)

*Notropis welaka*, EVERMANN & KENDALL, Bull. U. S. Fish Comm. 1897 (Feb. 9, 1898), 126, pl. 6, fig. 2, St. Johns River, near Welaka, Florida. (Type, No. 48786. Coll. Dr. W. C. Kendall.)

**Page 262.** After *Notropis bleunius* add:

408(a). **NOTROPIS BUCHANANI**, Meek.

Head 4; depth 4. D. 8; A. 8; scales 6-31-2; teeth 4-4. Body rather robust, back considerably elevated, snout blunt, mouth small and nearly horizontal. Snout short, about  $\frac{3}{4}$  diameter of eye. Preorbital bone slightly longer than broad. Eye moderate, 3 in head. Lateral line complete, or nearly so; about 12 scales in a series before dorsal fin. Dorsal fin slightly nearer tip of snout than base of caudal; pectorals reaching ventrals; ventrals reaching anal. Color light olivaceous, a faint silvery lateral band; no dark lateral band or black caudal spot. This species belongs to the *N. bleunius* type. It is a smaller species, lighter in color, and has fewer scales in the lateral line. Poteau River, Arkansas. (Named for Dr. John L. Buchanan, president of the Arkansas Industrial University.)

*Notropis buchanani*, MEEK, Bull. U. S. Fish Comm. 1895 (April 13, 1896), 342, small creek near Poteau, Indian Territory. (Type, No. 47532. Coll. Dr. Meek.)

**Page 267.** Under *Notropis nux*; *nuxce*, not *neche*, is nut in Spanish.

**Page 274, line 11,** for *luxoides*, read *luriloides*.

**Page 287.** After *Notropis lutipinnis* insert:

466(a). **NOTROPIS CHAMBERLAINI**, Evermann, new species.

Head  $4\frac{1}{2}$ ; depth  $4\frac{1}{2}$ ; eye 4; snout 4. D. 7; A. 9; scales 7-39-3, about 15 before the dorsal. General form much like that of *Hybognathus*; body only moderately compressed, dorsal and ventral outlines slightly arched; head rather small, pointed; mouth small, a little oblique, the maxillary scarcely reaching anterior border of orbit, lower jaw slightly included; snout equal to eye; eye in axis of body. Fins all rather small; origin of dorsal slightly behind vertical at insertion of ventrals; free edge of dorsal fin somewhat concave, the anterior rays about equal to length of head; pectoral short, slightly falcate, the longest rays about  $1\frac{1}{2}$  in head; ventrals shorter than pectoral, barely reaching vent; anal similar to dorsal, the rays shorter; caudal widely forked, the middle rays  $2\frac{1}{2}$  in the outer, the lobes as long as head, the lower lobe slightly longer than the upper. Scales moderately imbricated, the exposed portions not deeper than long; lateral line complete, somewhat decurved. Teeth 2, 4-4, 2 or 1, rather weak, hooked, and with small grinding surface. Intestine short; peritoneum silvery. General color light straw; middle of side with a broad, well-defined silvery band from upper end of gill opening to middle of

base of caudal fin, the anterior half lying wholly above the lateral line, the posterior portion lying partly below it; this silvery band bounded above by a narrow dark border; cheeks and opercles silvery; a darkish band along median line of back; fins all plain straw color or pale lemon. Fourteen examples of this species, 2 to 3 inches in length, were obtained from the Atchafalaya River at Melville, Louisiana, by Mr. Fred M. Chamberlain, for whom the species is named.

*Notropis chamberlaini*, EVERMANN MS., Atchafalaya River, Melville, Louisiana. (Type, No. 48901.)

**Page 291.** *Notropis scopifer*, Eigenmann & Eigenmann, is identical with *Notropis hudsonius selene* (Jordan), (p. 269), and should be omitted.

**Page 294.** After *Notropis dilectus* insert:

487(a). **NOTROPIS LOUISIANE**, Evermann, new species.

Head  $4\frac{1}{2}$ ; depth  $5\frac{1}{2}$ ; eye 3; snout 3. D. 7; A. 11; scales 7-37-3, 19 or 20 before the dorsal. Teeth 1, 4-4, 2, little hooked. Body long and slender, back not arched; head short, but pointed; mouth rather large, oblique, maxillary scarcely reaching orbit, lower jaw somewhat included; eye large, equal to or greater than snout. Fins rather small; origin of dorsal far behind insertion of ventrals, its longest rays  $1\frac{1}{2}$  in head; pectorals short, their length equal to height of anal; ventrals very short, 2 in head; caudal deeply forked. Scales firm, moderately imbricated; lateral line complete, gently decurved. Color pale; side with a faint plumbeous band; back and upper part of sides with numerous dark specks chiefly on the margins of the scales, thus forming cross-hatchings; a narrow dark vertebral band on caudal peduncle; peritoneum silvery, with numerous minute round black specks. Length  $2\frac{1}{2}$  inches. This species resembles *Notropis dilectus*, but has a much smaller mouth, blunter snout, and in being less silvery along the side. Known only from the Atchafalaya River, Louisiana.

*Notropis louisiana*, EVERMANN MS., Atchafalaya River, Melville, Louisiana. (Type, No. 48902. Coll. Fred M. Chamberlain.)

**Page 348.** *Anguilla chrysypa* is abundant in the Gulf of St. Lawrence, according to Dr. Wm. Wakeham.

**Page 355.** The original type of *Congermuræna* is *C. habenata*, Kaup, a species with blunt or granular teeth. The American species all belong to a distinct genus, *Congrellus*, Ogilby (type *balearica*), distinguished by the villiform teeth. These genera are characterized by Mr. Ogilby in a paper as yet unpublished.

**Pages 356 and 357.** In *Congermuræna flava* the upper jaw projects far beyond the lower. By a slip in the original description the reverse is said to be the case.

**Page 359.** *Murænoax coniceps* is called Culevra Blanca at Mazatlan, and reaches a length of 7 feet.

**Page 368.** *Avocettina gilli*, Bean, should probably stand as a species distinct from *Avocettina infans*. The description in the footnote on page

368 is sufficiently full. See Jordan, Proc. Cal. Ac. Sci., ser. 2, vol. vi, 1896, 206, pl. 21.

**Page 369.** No. 604, *Labichthys elongatus*, is a true *Arocettina*, having the vent far behind the head. It should stand as—

602(a). *AVOCETTINA ELONGATA* (Gill & Ryder).

**Page 376.** After *Myrichthys tigrinus*, Girard, add:

615(a). *MYRICHTHYS XYSTURUS*, Jordan & Gilbert.

Teeth all more or less blunt and granular; a band of 3 or 4 series on each side of lower jaw; a band of 2 rows on each side of upper jaw; vomer with a long series divided into 2 for about  $\frac{1}{2}$  its length. Anterior nasal tubes conspicuous, turned downward. Eye  $2\frac{1}{2}$  in snout; front of eye above middle of gape, the length of which is a little more than  $\frac{1}{3}$  of head; the angle of mouth well behind eye. Interorbital width about  $\frac{2}{3}$  length of the rather long and slender snout, which projects much beyond lower jaw, the tip of the latter about reaching middle of snout. Length of head contained  $4\frac{1}{2}$  times in that of trunk; head and trunk together shorter than tail, and contained  $2\frac{1}{2}$  to  $2\frac{3}{4}$  times in total length. Pectoral very small, its length about equal to depth of gill opening. Dorsal beginning close behind nape, much in front of gill opening; fins low; tail pointed, the tip sharp. Color light olive; sides each with 3 series of large round brown spots, those of the 2 upper series of equal size, those of lower scarcely  $\frac{1}{2}$  as large, faint, and often obsolete anteriorly; the spots irregular in their arrangement, those of the upper series usually twice as numerous as those of the next; those of the upper series along base of dorsal fin extending partly on the base of the fin; lower series of spots along base of anal, some of them extending on the fin or even entirely upon it; on the belly are sometimes small dark spots, scarcely arranged in series; dorsal fin with a terminal series of dark spots, which are partly confluent, the fin narrowly margined with white; anal reddish, with a lighter margin; pectoral with a blackish blot; head covered with round black spots, which become smaller and more numerous toward the snout; lower jaw with dark spots; iris light yellow. Pacific coast of Mexico; common among the rocks about Mazatlan. (*Ζύστουρ*, a spike; *ὄψρις*, tail.)

A species distinct from *M. tigrinus*, which is known only from the original type figured by Jordan & Davis, and described in the text of Part I of this work, page 376. This specimen, said to be from "Adair Bay, Oregon," may not be American, as there is no such bay in Oregon, and no second specimen of the true *Myrichthys tigrinus* has been found anywhere.

*Ophichthys xysturus*, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1881, 346, Mazatlan, Mexico. (Type No. 28142. Coll. Dr. Gilbert.)

615(b). *MYRICHTHYS PANTOSTIGMIUS*, Jordan & McGregor.

Head  $3\frac{1}{2}$  in trunk; head and trunk  $1\frac{1}{2}$  in tail; cleft of mouth 3 in head; eye  $2\frac{3}{4}$  in snout, which is 5 in head; pectorals 2 in snout; anterior nasal tube equal to the eye. Color olivaceous, with distinct rows of roundish

blackish spots, some oblong, smaller on head and covering the whole belly; 39 spots in the dorsal row, these spots usually alternating each with its fellow on the other side of dorsal, sometimes opposite; spots of second row usually opposite; spots of third row smaller and more numerous, extending from the cheeks to opposite the vent, thence running along base of anal, not running on fin, most of the spots of this row little more than  $\frac{1}{2}$  length of snout; 2 rows of smaller spots along belly from gill opening to front of anal; spots on nape rather large, on head larger and more numerous than in *M. xysturus*; pale color of head reduced to reticulations; chin and throat spotted as much as head; no pale centers to any of the spots; dorsal without spots or with only a few, which come up from back; from beginning to end the dorsal has a broad black margin about  $\frac{1}{4}$  height of fin; anal mostly pale, but toward tip having some black markings; pectoral with upper half jet-black, a white margin posteriorly, a small black spot in lower corner. This species is distinguishable from all others by the great number of spots of small size and without pale centers; the black edge of dorsal; the black spot on the rather large pectoral, and especially by having the belly spotted as much as the other parts. Clarion Island. One specimen, about a foot long, known. ( $\pi\acute{\alpha}\varsigma$ , whole, entire, all;  $\sigma\tau\acute{\iota}\gamma\mu\alpha$ , spot.)

*Myrichthys pantostigmus*. JORDAN & MCGREGOR, Rept. U. S. Fish Comm. 1898, pl. 4, Clarion Island. (Type, No. 5710, U. S. Jr. Univ. Mus. Coll. R. C. McGregor.)

Page 377. After *Pisoodonophis eruentifer* add:

618(a). *PISOODONOPHIS DASPILLOTUS*, Gilbert, new species.

Brownish above, gray below, the head and body usually thickly covered with black spots smaller than the eye; these are smaller and more numerous on the head, fewer and fainter on the lighter interior surface, and become indistinct or entirely disappear on the terminal portion of tail. In 1 specimen the head and trunk are spotted and the entire tail unicolor. In another no spots are present, the upper parts being a uniform dark brown, the under parts lighter brown, a few dark freckles only being present on sides of head. In all specimens the snout and lower jaw are blackish. The anus is near the middle of the total length, sometimes nearer the tip of snout, sometimes nearer tip of tail. The cheeks are not greatly swollen. The gape extends behind the eye, its length, measured from tip of lower jaw to angle of mouth, being contained  $4\frac{1}{2}$  to  $4\frac{3}{4}$  in head. The snout projects beyond the lower jaw for a distance about equaling diameter of orbit. Eye 2 to  $2\frac{1}{2}$  in snout,  $1\frac{1}{2}$  to  $2\frac{1}{2}$  in interorbital width. Tubes of anterior nostrils about  $\frac{1}{2}$  diameter of eye, directed downward near tip of snout. Posterior nostrils under front of eye, concealed in the upper lip as usual. Teeth all bluntly conic, in rather wide bands on jaws and vomer; they are usually not disposed in regular series within the bands, but each band has about the width of 4 series, and these are sometimes distinguishable. The mandibular teeth become larger on approaching the symphysis, those at point of mandible and those on head of vomer being much the largest teeth present. The patch on shaft of vomer tapers

backward to a point considerably behind angle of month. Origin of dorsal entirely behind tip of pectorals, its distance from snout  $\frac{1}{2}$  to  $\frac{1}{3}$  greater than length of the head. The tip of the tail is compressed, acute, horny, used for defense. Pectoral very short, from a wide base which slightly exceeds length of gill slit. The fin rapidly narrows downward, the longest portion contained 10 to 14 times in length of head. The width of gill slit is about  $\frac{1}{2}$  head.

The following table gives measurements of 4 specimens in millimeters.

Total length.	Head and trunk.	Tail.	Head.	Gape.	Eye.	Intraorbital width.	Width at cheeks.	Length of snout.	Projection of snout beyond mandible.	Length of pectoral.	Basal width of pectoral.	Distance from snout to dorsal.	Vertical height of dorsal.	Depth of body.
362	177	185	38	8	3	5	9 $\frac{1}{2}$	6 $\frac{1}{2}$	2 $\frac{1}{2}$	3 $\frac{1}{2}$	5	4 $\frac{1}{2}$	3	12
401	203	198	48	10 $\frac{1}{2}$	3	7	11	7 $\frac{1}{2}$	3 $\frac{1}{2}$	4	6	5 $\frac{1}{2}$	4	14 $\frac{1}{2}$
492	248	244	52	11	3 $\frac{1}{2}$	7 $\frac{1}{2}$	16 $\frac{1}{2}$	8 $\frac{1}{2}$	3 $\frac{1}{2}$	3 $\frac{1}{2}$	6 $\frac{1}{2}$	6 $\frac{1}{2}$	5 $\frac{1}{2}$	18
494	255	239	56	12	4 $\frac{1}{2}$	7	16	8 $\frac{1}{2}$	4 $\frac{1}{2}$	4	6	6 $\frac{1}{2}$	5 $\frac{1}{2}$	16

Four specimens were secured, 3 obtained in brackish water at the mouth of a small stream which empties into Panama Bay, the fourth in a fresh-water pond at Miraflores. There is some reason to suppose that they burrow in the mud.

*Pisodonophis daspilotus*, GILBERT, Fishes of Panama, MS. 1898, Panama.

**Page 382.** *Muraena ophis*, Linnaeus, is without much doubt the original *Ophichthus harannensis*. The species would therefore stand as—

626. OPHICHTHES OPHIS (Linnaeus).

**Page 396.** *Sidera castanea*, Jordan & Gilbert, should be removed from the synonymy of *Lycodontis fuchris*. It is apparently a valid species and should be inserted as—

650(a). LYCODONTIS CASTANEUS (Jordan & Gilbert).

(MORENA PRIETA.)

Tail about as long as rest of body, or slightly longer. Head  $2\frac{1}{2}$  in length of trunk; cleft of mouth wide,  $2\frac{1}{3}$  to  $2\frac{1}{2}$  in head. Teeth everywhere uniserial or nearly so, those on sides of mandible small, compressed, close set, subtriangular, directed backward, about 18 in number on each side; mandible with about 4 large canines anteriorly; upper jaw with the teeth partly in 2 series, some of the teeth being movable, the other mostly stronger, caninelike, especially anteriorly; front of vomer with 2 very long, slender canines, behind them a single series of small teeth; teeth all entire. Eye large, slightly nearer tip of snout than angle of mouth, its diameter 2 to  $2\frac{1}{2}$  in snout; gill opening  $\frac{1}{2}$  wider than orbit; tube of anterior nostril short, less than  $\frac{1}{2}$  diameter of orbit; posterior nostril with-

out tube; occiput not especially elevated, the anterior profile scarcely concave (perfectly straight in young 2 feet long). Dorsal fin commencing much in advance of gill opening, becoming unusually high posteriorly, where its vertical height is more than  $\frac{1}{2}$  greatest depth of body; the length of the longest ray more than greatest depth of body. Color light brownish chestnut, slightly paler on abdomen; no spots or bands anywhere; fins without dark margins; no dark spot on gill opening or at angle of mouth; no black about eye; head without conspicuous pores. The specimen here described is 44 inches in length; others about 2 feet in length agree very closely. This enormous eel is very common among the rocks about Mazatlan, where it reaches a length of 6 feet. It is close to the West Indian species, *L. faucrii*, but the colors are not the same, *faucrii* being a greenish black, while *castaneus* is a purplish chestnut, without shades of olive or green. (*castaneus*, chestnut.)

*Sidera castaneus*, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1882, 647, Mazatlan. (Type, Nos. 28246, 29535 and 29591. Coll. C. H. Gilbert.)

After *Lycodontis mordax* add:

649(a). LYCODONTIS PICTUS (Ahl).

Head  $\frac{1}{2}$  in trunk; tail about as long as body; eye  $2\frac{1}{2}$  in snout, situated midway between snout and angle of mouth; cleft of mouth  $2\frac{3}{4}$  in head; snout  $5\frac{1}{2}$  in head; anterior nasal tube 5 in snout; gill opening 11 in head. Teeth in each jaw in a single series; palatine series either parallel with these or divergent; no distinct canines; teeth comparatively small; anterior vomerine 1 or 2 in number, bluntnish and conical; posterior vomerine teeth rather blunt. Anterior nasal tubes moderate. Dorsal low anteriorly and beginning in front of gill opening. Color brownish gray or purplish, everywhere covered with small purplish black spots, which are not confluent; in the adult the spots are arranged in roundish or ring-like blotches on the sides; fins colored like body, without dark edges; young pale with black ring-shaped markings; variation in color and form of markings numerous. East Indies; everywhere common. East to offshore islands of Mexico. Two specimens, about 3 feet in length, taken at Clarion Island by Mr. R. C. McGregor. (*pietus*, painted.)

*Muræna picta*, AHL, Le Muræna et ophichtho, vi, 6, tab. 2, f. 2; GÜNTHER, Cat. Fish, VIII, 116.

*Uyjinnothorax pictus*, BLEEKER, At. Ichth., Muræna, 87, tab. 26, 28, 29, 45.

*Murænophis pantherina*, LACÉPÈDE, Hist. Nat. Poiss., v, 628, 1803.

*Muræna variegata*, QUOY & GAIMARD, Voy. Uranie, Zool., 246, pl. 52, f. 1.

*Muræna lita*, RICHARDSON, Voy. Erebus and Terror, 84, Moluccas.

*Muræna siderra*, RICHARDSON, Voy. Erebus and Terror, 85, pl. 48, f. 1 5, Australia.

*Muræna pfeifferi*, BLEEKER, Nat. Tyds. Ned. Ind., v, 173, Celebes.

*Sidera pfeifferi*, KAUP, Apodes, 70.

Page 401. After *Muræna argus* insert:

660(a). MURÆNA CLEPSYDRA, Gilbert, new species.

Closely related to *M. insularum* and *M. argus*, from the tropical Pacific, differing from both in color. Nostrils tubular, of almost equal length.

Mouth closing completely, the teeth entirely concealed by the lips. Gape straight, horizontal, extending to well behind the eyes,  $2\frac{1}{2}$  to  $2\frac{3}{4}$  in head. Teeth in jaws large, compressed, and wide at base, tapering uniformly to an acute point, directed backward, close set, everywhere uniserial; those in sides of mandible noticeably smaller than those of upper jaw, the teeth in both jaws increasing in size anteriorly; as many as 18 or 20 teeth may be present in the half of either jaw, but many of them are usually wanting, leaving gaps in the series; a single row of small teeth on shaft of vomer, beginning opposite front of eye; head of vomer with 2 long canines, larger than any of the other teeth, one or both of these usually wanting in larger specimens, having apparently fallen out. Head  $2(1\frac{1}{2}$  to  $2\frac{1}{2})$  in trunk; head and trunk  $1\frac{1}{2}$  to  $1\frac{1}{4}$  in tail; depth at anus approximately  $\frac{1}{2}$  length of head; eye small, its diameter contained 12 to 16 times in head; snout 5 to  $5\frac{1}{2}$ . Dorsal beginning on the head, its distance from snout  $1\frac{1}{2}$  to  $1\frac{1}{4}$  in head. Color dark brown, lighter on belly, dull whitish on under side of head; head, body, and fins closely covered with white spots, those on posterior parts larger, with some smaller ones intermingled, the larger spots with a more or less evident central constriction which makes them hourglass-shaped; toward the head the spots become very small and crowded, not more than  $\frac{1}{2}$  as large as pupil; fins indistinctly light margined; a large elliptical jet-black blotch surrounds the gill slit, distinctly margined by a series of confluent white spots; the longitudinal diameter of the blotch is contained 5 to  $5\frac{1}{2}$  times in the length of the head; angle of mouth with a small black blotch, often obscure, preceded by a pale spot on mandible; the throat is marked with a number of parallel lengthwise folds, the bottom of each fold with a dark line.

The following table gives measurements in millimeters of 5 specimens:

Total length.	Head and trunk.	Tail.	Head.	Gape.	Snout.	Eye.	Depth at anus.	Distance from snout to origin of dorsal.
675	311	364	106	45	$20\frac{1}{2}$	7	59	72
630	289	341	96	38	$18\frac{1}{2}$	$6\frac{1}{2}$	$52\frac{1}{2}$	70
612	287	325	98	39	19	6 $\frac{1}{2}$	47	64
473	203	250	66	28	13	$5\frac{1}{2}$	40	50
397	177	220	58	21	11	$4\frac{1}{2}$	27	39

This species is abundant at Panama, where it is frequently brought to market. About 25 specimens were seen during our visit, all essentially alike in coloration. The type is 397 millimeters long (see table of measurements), and has the spots on body less numerous than in larger specimens. (*clepsydra*, κλεψύδρα, an hourglass, from its markings.)

*Murena clepsydra*, GILBERT, Fishes of Panama. MS. 1899, Panama.

**Page 410.** It is probable that several species are confounded under the name *Elops saurus*. According to Ogilby the Australian species has only 63 vertebrae.

Page 411. 199. Genus ALBULA, Bloch & Schneider.

The proper binomial authority for this generic name, as well as for the names *Synodus*, *Umbra*, and *Anableps*, is Scopoli, as Dr. Gill informs us. These pre-Linnaean names, with others, were first used in binomial nomenclature as names of genera by Scopoli, *Introd. His. Nat.* 1777, pp. 449 (*Synodus*) and 450 (*Albula*, *Umbra*, *Anableps*). The genera should then stand as follows:

Page 411. 199. ALBULA (Gronow) Scopoli.

Page 533. 248. SYNODUS (Gronow) Scopoli.

Page 623. 298. UMBRA (Krümer) Scopoli.

Page 684. 312. ANABLEPS (Artedi) Scopoli.

Page 414. To the description of *Chanos chanos* the following may be added:

The skeletal peculiarities of *Chanos* are numerous and remarkable, many archaic characters persisting. The following account of the skeleton has been prepared by Mr. Starks:

SKELETON OF CHANOS CHANOS.

a. Cranium:

The frontals are very large, covering nearly the whole top of the head, and extending over the dorsoanterior part of the parietals, supraoccipital and the parotic process. On the side of the skull there is an area bounded by the supraoccipital, the opisthotic and the sphenotic, which is not ossified, but is composed of cartilage. Between the frontals, at about their middle, there is a place in which the bone is fibrous and largely cartilaginous; it is easily broken through. The basal cavity under the brain cavity is large. On the upper part of the operculum is a large scale-like bone. The suborbitals are well developed and plate-like, extending back nearly to the posterior edge of the preopercle.

b. Vertebral column:

There are 42 vertebrae in the spinal column. The first vertebra is co-ossified to the skull, and apparently bears no ribs; the second vertebra supports a pair of very small, slender ribs, which articulate directly with the sides of the vertebra; the third vertebra supports the first pair of large ribs; they are articulated with the transverse processes. The first 14 or 15 neural spines and pairs of transverse processes are articulated with the vertebrae by sutures; they are easily separated from the vertebrae by boiling or maceration. The vertebrae gradually increase in size and reach their largest size about  $\frac{3}{4}$  of the distance from the anterior to the posterior end of the spinal column, where they are 3 or 4 times the size of the anterior ones. This character is more marked in the adult than in the young.

s. Gape  
in head.  
ormly to  
al; those  
jaw, the  
20 teeth  
usually  
on shaft  
2 long  
usually  
Head 2  
at anus  
ed 12 to  
its dis-  
n belly,  
covered  
er ones  
onstric-  
e spots  
pil; fins  
rrounds  
ots; the  
length  
bscure,  
a num-  
rk line.  
imens:

ance  
om  
ut to  
in of  
sal.

72  
70  
64  
50  
39

ght to  
ntially  
easure-  
imens.

or the  
s only



c. Shoulder girdle:

The shoulder girdle is exceedingly well braced, the post-temporal is widely forked, and strongly articulated to the epiotic processes of the skull. The supraclavicle is long and slender, its posterior face is hollowed out and attached some distance from the upper end of the clavicle, which projects upward. This projecting upper end of the clavicle is braced to the skull by two long bones. The first bone is very slender, at its anterior end it is connected to the exoccipital; near its middle it is connected with the posterior end of the post-temporal, at which point it turns at a sharp angle and runs to the clavicle. The second bone is much larger; it is articulated to the basioccipital. Its posterior edge is nearly straight for its whole length, but its anterior edge is produced and much swollen near its middle, and joins the post-temporal over the first bone, then runs to the upper end of the clavicle. The inner part of the clavicle and the coracoid are thin and pierced by many holes, so that the bone in places is little more than network. The hypercoracoid has a very large foramen; at its posterior edge is a projection which supports a thin bone, probably a dermal bone. The mesocoracoid is well developed. There are 4 actinosts; the first is long, but they rapidly decrease in size to the fourth, which is short and triangular. The first ray of the pectoral is large at the basal end, and hollowed out; it works directly on the hypercoracoid.

d. Branchial apparatus:

The branchial apparatus is peculiar in the adult, in having gill rakers somewhat resembling the filaments of a feather, on both sides of each arch and on the basibranchial. They meet in a middle line between the arches and unite, forming a continuous lattice-work screen, through which nothing but the very smallest bodies can pass. The pharyngeals have no teeth, but have gill rakers similar to those on the arches; they are inclosed in sac-like projections on each side. This description is taken from the skeleton of a large specimen 4 feet long. The gill rakers are not united in young individuals.

e. Other parts:

The septa between the myotomes are ossified about  $\frac{1}{4}$  an inch under the skin, forming long, slender rays of bone. There is an upper series running from the middle of the sides up on the back, and a lower series from the sides down on the belly, and form a sort of a basket around the the body. Those below have a single branch near the middle of each, the ones above have 2 branches each; these branches are lost toward the posterior end. These bones are not present in the young. The large caudal fin is attached very firmly to the hypural, the long rays of each lobe join the hypural at about the same oblique angle, the base of each ray is deeply divided and articulated immovably with the hypural. The middle short rays are all nearly horizontal and are much less firmly fastened. The first interspinal ray of the anal is hollow and cone-shaped, the posterior end of the air bladder runs into it as in the genera *Eucinostomus* and *Calamus*. The scales are very thick and closely imbricated; the skin anteriorly is  $\frac{1}{4}$  inch thick. (Jordan, Fishes of Sinaloa, 404-409.)

Page 417. After *Dorosoma petenense* add:

202(a). SIGNALOSA, Evermann & Kendall.

*Signalosa*, EVERMANN & KENDALL, Bull. U. S. Fish Comm. 1897 (Feb. 9, 1898), 127 (*Atchafalaya*).

Body short, deep, and compressed, the form somewhat elliptical; ventral outline more strongly curved than the dorsal; head rather large, snout sharp and pointed, not tumid; mouth small, oblique, the lower jaw scarcely included; maxillary of 3 pieces, broad and curved, but without notch in the outer margin as in *Dorosoma*; caudal peduncle short and deep. Branchiostegals 5; pseudobranchia large; gill rakers short and very numerous, about 340 in number. No teeth; adipose eyelid present; stomach gizzard-like; scutes about 17 + 10. Last ray of dorsal very long and filamentous. This genus is allied to *Dorosoma*, from which it is plainly distinguished by the absence of the notch in the maxillary, the more pointed snout, the less-included lower jaw, the shorter anal fin, larger scales, and the fewer scutes. It differs from *Alosa* in the very numerous gill rakers, the character of the dorsal fin, and in other respects. (*signum*, a flagstaff or pole; *Alosa*, the shad; a reference to the long dorsal ray.)

679(a). SIGNALOSA ATCHAFALAYE, Evermann & Kendall.

Head  $3\frac{3}{4}$ ; depth  $3\frac{1}{4}$ ; eye  $3\frac{1}{4}$  in head; snout  $5\frac{1}{4}$ ; maxillary  $3\frac{1}{8}$ . D. I, 12; A. I, 24; scales 42-15; scutes 17 + 10. Body oblong-elliptical, compressed, the back in front of dorsal narrow; ventral edge sharp, serrate; head small, mouth terminal, oblique, lower jaw slightly included; snout rather pointed, not blunt, as in *Dorosoma cepedianum*; maxillary in 3 pieces, long and curved, reaching vertical at front of pupil, the outer edge not notched; no teeth. Caudal peduncle short, compressed, and deep. Origin of dorsal fin over base of ventrals, much nearer tip of snout than base of caudal, the last ray filamentous, about  $\frac{1}{4}$  longer than head and nearly reaching base of caudal; the first dorsal ray about 2 in the last one; pectoral  $1\frac{1}{4}$  in head, reaching base of ventrals; ventrals short, reaching only halfway to vent, their length  $1\frac{1}{2}$  in pectorals; anal rays short, base of fin  $1\frac{1}{8}$  in head; scutes moderate; caudal widely forked, the lower lobe the longer; scales large, thin, deciduous, somewhat crowded anteriorly; accessory scales at bases of pectorals and ventrals; base of caudal with small scales. Color bluish black or dark olivaceous on back and sides to level of the jet-black humeral spot; rest of sides and under parts bright silvery; dorsal and caudal dusky; other fins plain. The cotypes from Grand Etang Bayou are 2 females with ripe roe. They are  $4\frac{1}{4}$  and  $5\frac{1}{2}$  inches long, respectively, and differ from the types only in the deeper body and the much darker coloration of the upper parts.

The amount of variation in this species, shown by the material at hand, is exhibited in the following table:

No.	Head.	Depth.	Eye.	Snout.	Max.	Dorsal.	Anal.	Scutes.	Scales.	Locality.
1	4½	3	3¾	5½	3½	I, 12	I, 24	16 + 11	40-15	} Grand Plains Bayou, Miss.
2	4	2½	3¼	5½	3½	I, 12	I, 24	16 + 11	42-15	
3	4	3	3¼	5½	3¾	I, 12	I, 24	16 + 10	42-14	} Molville, La.
4	3¾	3½	3½	5½	3½	I, 12	I, 24	17 + 10	42-15	
5	3¾	3	4	5	3½	I, 12	I, 24	16 + 11	43-15	} Grand Plains Bayou, Miss.
6	3¾	3	3¾	5½	3	I, 12	I, 24	16 + 10	41-15	
7	3¾	3	3½	5	3½	I, 12	I, 24	17 + 9	41-15	} Black Bayou, Miss.
8	3¾	3	3½	5	3	I, 12	I, 24	17 + 9	41-15	
9	3¾	2½	3½	5½	3	I, 12	I, 24	17 + 9	40-15	

This species appears to be rather common in the larger lowland streams and bayous of Louisiana and Mississippi. It probably does not reach a large size, adult examples being less than 6 inches long. It is not used as food, but is of considerable value as bait in the catfish fishery of the Atchafalaya River and its connecting lakes and bayous. Length 4 to 6 inches. (*atchafalaya*, from the type locality.)

*Signalosa atchafalaya*, EVERMANN & KENDALL, Bull. U. S. Fish Comm. 1897 (Feb. 9, 1898), 127, pl. 7, fig. 4, Atchafalaya River, Melville, Louisiana. (Type, No. 48790. Coll. Fred M. Chamberlain.)

**Page 425.** The statement that *Pomolobus mediocris* does not ascend rivers to spawn is not correct. This species is known to ascend the St. Johns River, Florida, at least as far as Lake Monroe, during the winter. They usually run somewhat earlier than the shad.

**Page 427.** After *Alosa*, Cuvier, add:

- a. Gill rakers numerous, 93 to 120; upper jaw with sharp, deep notch at tip; lower jaw not projecting. SAPIDISSIMA, 693.  
 aa. Gill rakers fewer than 76; notch in upper jaw smaller; lower jaw more strongly projecting. ALABAME, 693(a).

**Page 428.** After synonymy of *Alosa sapidissima* add:

693(a). ALOSA ALABAME, Jordan & Evermann.

(ALABAMA SHAD; GULF SHAD.)

Head 4¾; depth 3; snout 4¾; eye 4¾; maxillary 2¼. D. 15; A. 20; scales 55, —16 in a crosswise series; scutes 21 + 15; vertebrae 54; gill rakers 56 to 68. Body deep; back gently and evenly arched from tip of snout to origin of dorsal fin, thence descending in a regular curve to base of caudal fin; ventral outline nearly straight from tip of mandible to ventrals, and also from there to base of caudal. Head small, snout pointed; upper lip with a small notch, into which fits the tip of the slightly projecting lower jaw; maxillary narrow; cheek much deeper than long; teeth on tongue and maxillary scarcely perceptible. Origin of dorsal nearer snout than base of caudal, the fin low, the longest ray shorter than the base, or about equal to snout and eye; base of anal somewhat greater

than that of dorsal, or equal to length of pectoral. Gill rakers 68, the longest about equal to length of snout. Peritoneum pale. Color as in *Alosa sapidissima*; the caudal, dorsal, and pectoral fins rather darker tipped. The male differs from the female only in being somewhat more slender. This species differs from *Alosa sapidissima* chiefly in the fewer gill rakers, its sharper, more pointed snout, smaller notch in upper jaw, more projecting mandible, and more slender maxillary. It seems to reach maturity at a much smaller size than the common shad. Streams tributary to the Gulf of Mexico; known from Tuscaloosa, Alabama, and Pensacola, Florida.

*Alosa alabama*, JORDAN & EVERMANN, in EVERMANN, Rept. U. S. Fish Comm. 1895 (Dec. 28, 1896), 203, Black Warrior River, Tuscaloosa, Alabama (Type, Nos. 47689 and 47690. Coll. J. H. Fitts); EVERMANN & KENDALL, Bull. U. S. Fish Comm. 1897, 127, pl. 7, figs. 5 and 6.

According to Ogilby, *Kowala* is a genus distinct from *Sardinella*.

**Page 436.** *Ilisha panamensis* is not separable from *I. furthi*. The latter name has priority.

**Page 437.** *Opisthopterus lutipinnis* is very abundant on the outer sand beaches about Mazatlan.

**Page 445.** Species 728, *Stolephorus poeyi*, is a species of *Lycengraulis*, and should stand as—

743(a). LYCENGRAULIS POEYI (Kner & Steindachner).

Numerous specimens lately taken by Dr. Gilbert at Panama. A large species used as food. The teeth are unequal in *Lycengraulis*, but none of them can be properly described as canine-like.

**Page 447.** After *Stolephorus lucidus* add:

732(a). STOLEPHORUS BASTRALIS, Gilbert & Pierson, new species.

Head 3.16 (3.1 to 3.3); depth 3.8 (3.5 to 4.2); eye 3.4 in head (3.33 to 4). D. 14 (12 to 15); A. 26 to 32. Body much compressed and deep; belly sharply keeled in front of ventrals; dorsal outline much less curved than ventral, the lower profile rising very rapidly from a point opposite middle of pectorals to tip of snout, in shape of head thus closely resembling the species of *Cetengraulis*. Maxillary reaching almost but not quite to gill opening; snout high, compressed, its length  $\frac{1}{2}$  to  $\frac{3}{4}$  diameter of eye. Gill rakers averaging in larger examples 51 + 64, in smaller specimens 44 + 50; the largest about as long as eye. Insertion of dorsal fin variable, but never posterior to a point midway between base of caudal and middle of eye; pectoral fins reaching to or nearly to insertion of ventrals, the latter not to vent. Color olivaceous, the lower part of side with violet reflections; sides of head silvery; a conspicuous silvery lateral band varying in width from about  $1\frac{1}{2}$  times length of orbit in the largest examples to less than  $\frac{1}{2}$  orbit in the smaller specimens; the band is widest before dorsal, tapering to  $\frac{1}{2}$  or less than  $\frac{1}{2}$  its greatest width on caudal peduncle, where it frequently disappears in the young. In larger specimens the ventral edge of this band is frequently ill-defined anteriorly; top of head with

widely spaced black specks; a dark vertebral streak, more or less of it often consisting of 2 narrow lines; tips of caudal lobes often blackish; fins otherwise unmarked. This species differs from closely allied species in the following characters: From *Stolephorus lucidus*, in the much longer head, more compressed body, well-defined lateral stripe, and smaller eye; from *S. compressus*, in the longer head and wider lateral band; from *S. panamensis* and *S. mundeolus*, in the much more numerous gill rakers, and the more anterior position of the dorsal relative to the anal, the origin of the anal being under the middle of the dorsal, while in *S. panamensis* the origins of the two fins lie in the same vertical. Length 2 to 3 inches. Panama. Many specimens. (Gilbert & Pierson.) (*rastrum*, a rake, from the long gill rakers.)

*Stolephorus rastralis*, GILBERT & PIERSON, Fishes of Panama, MS. 1898, Panama.

732(b). *STOLEPHORUS MUNDEOLUS*, Gilbert & Pierson, new species.

Head 4.15 (4 to 4.25); depth 3.77 (3.40 to 4.25); eye 3.41 in head (3.12 to 3.70). D. 13 or 14; A. 33 (33 to 35); scales 36 (35 to 39). Dorsal and ventral contours about equally and gradually rounded from the middle region of body to the tip of snout and base of caudal fin. Snout short, high, compressed, blunt at tip, its length 1.8 in eye. Eye very large. Maxillary broad, tapering to a sharp point, which reaches margin of gill opening. Gill rakers 17 to 22 + 21 to 24; the longest 1.5 to 2 in eye. Anterior insertion of dorsal fin varying from a point midway between base of caudal and middle of eye, to a point midway between the caudal and tip of snout. In 10 examples its insertion is before that of the anal. Anal fin long, averaging 33 rays, its origin beneath the anterior third of the dorsal; length of base shorter than in *S. panamensis*, being 3.04 in length, while in the latter its length is contained 2.5 in length. Pectoral long, reaching well beyond the insertion of the ventrals, equaling length of head behind front of pupil; a large axillary scale; ventrals scarcely reaching vent. Uniform light olive, with silvery reflections; a faint, narrow, silvery-gray lateral stripe, sometimes scarcely distinguishable; sides of head plain silvery; upper margin of orbital rim black; dorsal region blackish; a faint, narrow dark line on each side of the light mid-dorsal streak; caudal slightly dusky; fins otherwise unmarked. This species is closely allied to *Stolephorus panamensis* and *S. compressus*, but may be distinguished from the former by its longer head, larger eye, greater depth, fewer scales along the lateral line, and its much shorter anal base; also by the much fainter lateral silvery stripe. The eye is contained 14 to 16 times in length, excluding the caudal, while in *S. panamensis* the length contains the eye 16 to 20 times. From *S. compressus* it differs in the relative length of the head and maxillary. In *S. mundeolus* the maxillary is contained in the head 1.27 times (1.19 to 1.37); in *compressus* 1.48 times (1.30 to 1.81). In *mundeolus* the head is contained 4.15 times in the length; in *compressus* 4.44 times. Length 4 to 6½ inches. Panama; many specimens. (Gilbert.) (*mundeolus*, somewhat shining, from *mundus*, neat or clean.)

*Stolephorus mundeolus*, GILBERT & PIERSON, Fishes of Panama, MS. 1898, Panama.

## 732(e). STOLEPHORUS NASO, Gilbert &amp; Pierson, new species.



Head 3.3 to 3.5; depth 4.7 to 5.8; eye  $4\frac{1}{2}$  to 5 in head. D. 14 or 15; A. 22 to 24; lateral line about 35 (?). Dorsal and ventral outlines weakly arched; body slender, its greatest depth 1.5 in head, compressed; belly carinated in front of ventrals, and sometimes behind them in larger specimens. Head long and slender, its greatest width 1.5 to 1.7 in its length, the lower profile much more oblique than the upper. Snout long, compressed, bluntly rounded, its length exceeding the small eye. Cheek with a very acute posterior angle. Opercle narrow, oblique. Maxillary rather bluntly pointed, failing to reach gill opening by about  $\frac{1}{2}$  diameter of pupil. Teeth on the maxillary quite prominent and directed forward. Gill rakers short, 17 + 20 in number, the longest 1.5 in eye. Scales large, thin, deciduous, only a few scattering ones remaining in our specimens. Dorsal fin inserted midway between front or middle of orbit and base of median caudal rays. Origin of anal under or slightly behind middle of dorsal; length of anal base about equal to the distance from front of orbit to base of ventral fin; pectorals not reaching ventrals, their length about  $\frac{1}{2}$  length of head. Length of ventrals equaling or slightly exceeding distance from tip of snout to middle of pupil. Color light olive, with the usual bright reflections; a large dark patch of brown dots on occiput; a double series of dots along median line posterior to dorsal, this absent in some specimens; large specimens with a bright, well-defined silvery streak, slightly narrowing anteriorly and on caudal peduncle, its greatest width about equaling diameter of eye; in the young, this band is fainter and narrower; a conspicuous series of black dots at base of anal. Characterized by the slender form, well-defined silvery streak, sharply carinated breast, the small eye, and the very long, compressed, deep, and rather bluntly rounded snout. Most closely resembling *S. starksi*, from which it differs in the smaller eye, longer snout, and slightly longer anal. Length 2 to 2 $\frac{1}{4}$  inches. Panama; common. (Gilbert & Pierson.) (*naso*, long-nosed.)

*Stolephorus naso*, GILBERT & PIERSON, Fishes of Panama, MS. 1898, Panama.





## 732(d). STOLEPHORUS STARKSI, Gilbert &amp; Pierson, new species.

Head 3.3 to 3.6; depth 4.8 to 5.5 in length,  $1\frac{1}{2}$  in head; eye 3 to 3.5 in head. D. 15 or 16; A. 17 to 22; scales about 41. Body long and slender, slightly deeper and more compressed than in *S. ischannus*, which much resembles this species. Dorsal outline very little arched; ventral outline nearly straight from gill opening to insertion of anal fin, the lower profile of head oblique, nearly straight. Belly compressed, keeled for anterior  $\frac{2}{3}$  of distance anterior to base of ventrals. Head long and pointed, its width  $1\frac{1}{2}$  times in its length; maxillary abruptly widened at the mandibular joint, tapering posteriorly to a blunt point, which reaches almost to the gill opening, its length equal to length of base of anal; snout long, sharp, and projecting, abruptly compressed in its terminal portion as seen from above, its length  $\frac{2}{3}$  diameter of orbit, or slightly more. Branchiostegal membranes united at base for a very short distance. In 4 exam





**IMAGE EVALUATION  
TEST TARGET (MT-3)**



**Photographic  
Sciences  
Corporation**

23 WEST MAIN STREET  
WEBSTER, N.Y. 14590  
(716) 872-4503



15 128  
16 132  
17 125  
18 122  
19 120

01

ples examined as to this point, the gill rakers are as follows: 20 + 25, 23 + 24, 21 + 23, 19 + 30, the longest contained  $1\frac{1}{2}$  to  $1\frac{3}{4}$  in eye. Scales large, thin, and deciduous, a few only remaining on the specimens at hand. Origin of the dorsal fin equally distant from the base of the caudal fin and tip of snout or front of eye. Anal inserted under beginning of posterior third of base of dorsal; pectorals not reaching ventrals, the latter  $\frac{3}{4}$  distance to front of anal. Color light olive, with broad, well-defined lateral silvery streak of nearly uniform width, usually narrowing anteriorly and on middle of caudal peduncle, its width in our largest specimen  $\frac{3}{4}$  diameter of eye; the silvery streak has a slight golden tinge; a narrow dark vertebral line, which widens on the nape; occiput blackish. Vertebrae 40, counted in 1 example only. This species differs from *Stolephorus cultratus* in its slenderer body, shorter snout, wider opercle and smaller teeth; the belly is also not sharply carinate, the dorsal is more anteriorly placed, the ventrals are farther back, and the silvery streak is wider anteriorly. It differs from *S. delicatissimus* in its longer, slenderer head and body, smaller eye, longer, sharper snout, and much wider, better-defined silvery streak. Length  $1\frac{1}{2}$  to  $2\frac{1}{2}$  inches. Panama; common. (Gilbert & Pierson.) (Named for Edwin Chapin Starks.)

*Stolephorus starksii*, GILBERT & PIERSON, Fishes of Panama, MS. 1898, Panama.

Page 448. After *Stolephorus spinifer* add:

737(a). *STOLEPHORUS SCOFIELDI*, Jordan & Culver.

Head  $3\frac{3}{4}$  to  $3\frac{9}{10}$  in length to base of caudal; depth  $4\frac{1}{2}$  to 5; eye  $3\frac{1}{4}$  to 4 in head. D. 12; A. 25 or 26; scales 41 or 42. Body somewhat compressed and elevated, the belly not carinated nor serrated. Teeth in both jaws, and on palatines, a few on vomer; maxillary covered with teeth its entire length and reaching beyond base of mandible, but not to opercular margin. Gill rakers 10 + 12, the longest a little more than  $\frac{1}{2}$  the eye. Origin of dorsal midway between base of median caudal rays and center of eye; anal not quite as long as head, its origin below the middle of dorsal; lower caudal lobe longer than upper; longest ray equaling length of the head; shortest caudal ray  $2\frac{1}{2}$  in longest. Pectorals not reaching ventrals,  $1\frac{1}{4}$  in head. Both anal and dorsal fins preceded by a rudimentary spine, not  $\frac{1}{2}$  length of first true ray. Color translucent, with a distinct broad silvery stripe as wide as the eye, growing more diffuse at lower anterior edge, narrowing on caudal peduncle, and becoming fan-shaped on the base of caudal; tip of snout black; a distinct median band of black specks extending from tip of snout to base of caudal; no distinct black markings on fins. Close to *Stolephorus delicatissimus*, but with larger head, wider lateral band, and greater number of dorsal and anal rays. Length 3 inches. Found in the Astillero at Mazatlan; not very abundant. (Named for Mr. Norman Bishop Scofield, a member of the Hopkins expedition to Sinaloa.)

*Stolephorus scofieldi*, JORDAN & CULVER, Fishes of Sinaloa, 410, 1895 Mazatlan, Mexico. (Type, No. 2941, I. S. Jr. Univ. Mus. Coll. Hopkins Exped. to Sinaloa.)

737(b). *STOLEPHORUS ASTILBE*, Jordan & Rutter.

Head  $4\frac{1}{2}$  in length; depth  $4\frac{1}{2}$  to 5. D. 12; A. 19 to 22; eye  $3\frac{1}{2}$  in head; pectoral  $1\frac{3}{4}$ ; base of anal  $1\frac{1}{2}$ . Body rather elongate, not greatly compressed; edge of belly moderately sharp; head sharp; snout projecting beyond lower jaw, shorter than diameter of eye; tip of lower jaw reaching a little past anterior edge of orbit; maxillary reaching gill opening, its end tapering to a sharp point; eye longer than snout, nearly 2 in post-orbital part of head; gill rakers  $\frac{3}{4}$  eye; a slight keel on top of head. Origin of dorsal midway between base of caudal and eye; scales caducous. Translucent, head silvery; sides without lateral band; a dark spot on top of head; back with black points. This species is similar to *Stolephorus brownii*, but more slender, head shorter, and lateral silvery stripe wanting. Length 3 inches. Jamaica. Numerous specimens obtained. ( $\alpha$ , not;  $\sigma\tau\iota\lambda\beta\eta$ , shining.)

*Stolephorus astilbe*, JORDAN & RUTTER, Proc. Ac. Nat. Sci. Phila. 1897, 95, Jamaica. (Type, No. 4854, L. S. Jr. Univ. Mus. Coll. Joseph Seed Roberts.)

737(c). *STOLEPHORUS ROBERTSI*, Jordan & Rutter.

Head 3 in length; depth 4. D. 14; A. 23; scales about 35; eye 4 in head; pectoral  $2\frac{3}{4}$ ; base of anal  $1\frac{3}{4}$ ; caudal  $1\frac{1}{2}$ . Body deep, strongly compressed, abdomen compressed to an edge, head large, compressed, the snout rather sharp, projecting beyond lower jaw, a little shorter than eye; cheek triangular, opercle large; distance from lower angle of cheek to edge of opercle equal to distance from same point to posterior edge of eye; maxillary short, not reaching root of mandible, its end rounded; lower jaw not reaching beyond anterior edge of orbit; gill rakers longer than eye, as long as orbit. Origin of dorsal midway between base of caudal and front of eye; scales caducous. Color translucent; head silvery, punctulate above; a silvery lateral band nearly as broad as eye; caudal with dark points, other fins colorless. This species seems to be related to *Stolephorus opercularis*, but the lateral band is distinct and the opercle is shorter. Jamaica; only the type, 2 inches long, known. (Named for Rev. Joseph Seed Roberts, of Kingston, Jamaica, who collected the type specimen.)

*Stolephorus robertsi*, JORDAN & RUTTER, Proc. Ac. Nat. Sci. Phila. 1897, 95, Jamaica. (Type, No. 4853, L. S. Jr. Univ. Mus.)

**Page 449.** *Anchoria* can not be maintained as a distinct genus. The name must be placed as a synonym of *Stolephorus*.

**Page 450.** Add:

741(a). *CETENGRAULIS ENGYMEN* Gilbert & Plerson, new species.

Head 3 to 3.3; depth 4 to 4.9; eye 4 in head. D. 14 or 15; A. 20 to 23; B. 7 (9); vertebrae 41. Body compressed, fusiform, not so deep as in *C. mysticetus* or *C. edentulus*. The dorsal and ventral outlines being about equally and regularly curved in the larger specimens; in the smaller specimens the ventral contour is more nearly straight. Belly trenchant, but

not carinate nor serrate; caudal peduncle moderate, its depth being contained 1.5 times in its length. Head similar to *C. mysticetus*; the snout longer, contained 5.5 to 7 times in head,  $1\frac{1}{2}$  times in eye (the snout contained 8 to 9 times in head in *C. mysticetus*). Both jaws bear minute teeth, those on the maxillary largest. Branchiostegal membranes united for only  $\frac{2}{3}$  to  $\frac{3}{4}$  of the distance between tip of mandible and mandibular articulation, wholly free from the isthmus. Tip of mandible directly beneath the anterior border of orbit. Gill rakers long,  $\frac{1}{10}$  diameter of eye, 20 to 30 on the upper limb, 25 to 30 on the lower limb; in 5 examples as follows: 25 + 30, 27 + 25, 30 + 26, (23 + 29 to 20 + 25), 25 + 30. The origin of the dorsal is midway between base of median caudal rays and a point varying between front and middle of the eye. Insertion of anal below the posterior fourth or third of the dorsal, its length equaling the distance from the posterior border of the eye to insertion of pectoral. The pectoral is short,  $2\frac{1}{4}$  to  $2\frac{1}{2}$  in head, failing to reach the insertion of the ventrals by  $\frac{1}{2}$  or nearly  $\frac{1}{2}$  its length. Caudal deeply forked, its median rays  $2\frac{1}{2}$  to 3 times in head. Color uniformly silvery with a distinct, well-defined lateral silvery band extending from upper angle of gill opening to base of caudal, its greatest width equaling the diameter of orbit, becoming narrower on caudal peduncle. This species differs from *C. mysticetus* in the much narrower union of the gill membranes, the less numerous gill rakers, and in the longer snout. Length  $1\frac{1}{2}$  to  $2\frac{1}{2}$  inches. Panama Bay. Not rare. (Gilbert & Pierson.) (*εγγύς*, near; *δμήρ*, membrane.)

**Page 451.** *Lycengraulis* has the teeth large and somewhat unequal, but none of them is properly described as "canine-like."

**Page 459.** Add:

229(a). **ERICARA**, Gill & Townsend.

*Ericara*, GILL and TOWNSEND, Proc. Biol. Soc. Wash., XI, 1897 (Sept. 17, 1897), 232 (*salmonea*).

Alepocephalids with small, perfectly smooth, imbricated cycloid scales, wide cranium, projecting snout, deeply cleft mouth, uniserial and acrodont teeth on vomer and anterior portion of palatines, and dorsal and anal of normal extent and opposite each other. Bering Sea. (*έρή*, an intensive particle; *καρά*, head.)

758(a). **ERICARA SALMONEA**, Gill & Townsend.

D. 17; A. 24. Maxillary extending to vertical of posterior border of orbit; head large; length  $8\frac{1}{2}$ ; depth 5; width  $4\frac{1}{2}$ . Bering Sea; only the type known, a large example in good condition.

*Ericara salmonsea*, GILL & TOWNSEND, Proc. Biol. Soc. Wash., XI, 1897 (Sept. 17, 1897), 232, Bering Sea, southwest of Pribilof Islands, at Albatross Station 3603, in 1,771 fathoms. (Type, No. 48769, U. S. Nat. Mus. Coll. *Albatross*.)

**Page 465.** Dr. G. A. Boulenger has kindly sent us the following note regarding the types of *Coregonus richardsoni* which are in the British Museum:

I have examined the types (dry) of *Coregonus richardsoni*. There are about 20 gill rakers on the lower part of the anterior arch, the longest  $\frac{1}{4}$  the diameter of the eye. The maxillary extends to below anterior border of eye, and its length is 4 times in length of

head as stated by Günther, therefore a little shorter than in *C. clupeiformis*. Tongue with 4 series of teeth, as in *C. labradoricus*. It seems to agree best with *C. nelsoni* (description), but has fewer scales in lateral line. In short, I can not identify *C. richardsonii* with any of the forms known to me.

Page 471. After *Argyrosomus lauretta* add:

768(a). ARGYROSOMUS ALASCANUS, Scofield, new species.

Head  $4\frac{1}{2}$ ; depth about 4. D. 12; A. 12; scales 10-85-9. Eye a little shorter than snout, 5 in head,  $1\frac{1}{2}$  in interorbital space. Head wedge-shaped, the upper and lower profiles straight and meeting with a sharp angle at the snout. Viewed from above the snout is blunt, almost square, with the narrow, pale rounded tip of the lower jaw slightly projecting. Mouth oblique; the distance from tip of snout to tip of maxillary is equal to the distance from tip of snout to center of pupil; the maxillary from its anterior articulation is contained  $3\frac{1}{2}$  in the head, its width 3 in its length, its upper anterior edge closing under the preorbital; mandible  $2\frac{1}{2}$  in head, its articulation with the quadrate bone beneath the posterior edge of the eye; width of supplemental bone a little more than  $\frac{1}{2}$  width of maxillary. Preorbital broad, its greatest width equaling  $\frac{2}{3}$  its length or diameter of pupil; width of supraorbital equals  $\frac{2}{3}$  its length. Gill rakers 12 to 14 + 21 to 23, long and slender, the longest  $\frac{2}{3}$  diameter of the eye. The tongue, vomer, and palatines without teeth. Distance from tip of snout to nape equaling  $\frac{1}{2}$  the distance from nape to the front of the dorsal or  $\frac{2}{3}$  length of head. Adipose fin large; ventral scale  $\frac{1}{2}$  length of fin; longest dorsal ray  $1\frac{1}{2}$  in head; longest anal ray 2 in head; the pectorals reach more than  $\frac{1}{2}$  to the ventrals; the ventrals reach  $\frac{2}{3}$  to vent; the caudal is forked for a little more than  $\frac{1}{2}$  its length. Color dusky above, silvery beneath; the dorsal, adipose fin, tips of caudal rays and upper side of anterior pectoral rays dusky; the rest pale. This species appears nearest related to *Argyrosomus arctici*, from which it differs chiefly in the number of gill rakers.\* Length about a foot. Northern Alaska near Bering Straits; 3 specimens known, 1 from salt water at Point Hope, the others from fresh water at Grantley Harbor.

*Argyrosomus alascanus*, SCOFIELD, in JORDAN, Rept. U. S. Fur Seal Investigations, 1898, Point Hope and Grantley Harbor, Alaska. (Coll. Scofield & Seale.)

Page 482. Beginning with line 10 from the bottom, the statement that the small form of the redfish has been traced from the mouth of the Columbia to Wallowa Lake is not true. The remark was meant to apply to the large form. The question as to whether the small form descends to the sea is still unsettled.

\* The fin formulae, etc., of these 3 specimens are as follows:

Locality.	Length.	Gill rakers.		Dorsal.	Anal.	Scales.
Grantley Harbor.....	8 $\frac{1}{2}$ in.	14 + 23	14 + 22	12	14	88
Do .....	9 in.	14 + 22	12 + 22	12	12	87
Point Hope, Alaska.....	11 in.	13 + 21	12 + 21	12	12	85

## Page 492:

## 779. SALMO MYKISS, Walbaum.

(MYKISS; SOMKA; KAMCHATKA SALMON TROUT.)

By an unfortunate error the writers have heretofore used the name *Salmo mykiss* for the Cutthroat Trout of the Northwest. It was known that the Cutthroat is the only true or black-spotted trout in Alaska, and it was assumed that its range extended along the coast to all streams in Bering Sea. But our recent explorations have shown that it probably does not occur in Bering Sea, nor is there any undoubted record to the north of Wrangell. If it reaches Kadiak or Sitka or Prince William Sound, it is only rarely, and the streams of the Aleutian Islands and the east coast of Bering Sea contain no species of *Salmo*. The name *Salmo mykiss* must, therefore, be restricted to the Kamchatkan species, while the species of the American rivers heretofore called *Salmo mykiss* must stand as *Salmo clarkii*. We have, therefore, studied with great interest a specimen of the genuine *Salmo mykiss*, the first on record since the times of Pallas, Krasheninnikof, and Steller. The specimen, an adult male, 960 mm. long, was taken by Dr. Leonhard Stejneger, September 15, 1897, in the Kalakhtyrka River, near Petropaulski, Kamchatka. It was called "Sonka" or "Somka" by the natives. It is said to occur rarely and to be found in but few rivers, the Kalakhtyrka among them. It is considered to be superior as food to other *Salmonidae*, except the King Salmon (*O. tshawytscha*). Head 4 in length; depth 4½. D. 11. A. 10 (developed rays); scales 125-24. Mouth large, the maxillary 1½ in head, being somewhat produced at tip; vomerine teeth few, evidently deciduous, only 3 being present. Eye 8½ in head; snout 2¾. Pectoral 2 in head, longest anal ray 2¾. Anal fin high and somewhat falcate; ventrals inserted under anterior third of dorsal, reaching about halfway to vent; adipose fin over posterior end of anal; caudal lunate. Color dark grayish above, sides silvery; a few small, faint, round black spots on back and on top of head, these sparse and obscure; a few faint spots on base of dorsal, and some on adipose dorsal; spots on caudal small, but distinct, especially in middle of fin; no trace of red at throat, in example preserved in formalin, and doubtless none in life. The specimen is now a half skin, in good condition.

The following measurements were taken from the fresh specimen by Dr. Stejneger:

	Millimeters.
Total length .....	960
Total length without caudal .....	853
Head .....	215
Tip of nose to anterior end of dorsal .....	400
Length of base of dorsal .....	190
Posterior end of dorsal to anterior end of adipose fin .....	167
Length of base of adipose fin .....	17
Posterior end of adipose to caudal .....	81
Posterior end of anal to caudal .....	81
Length of base of anal .....	71
Anterior end of anal to posterior end of ventral .....	165
Height of body in front of dorsal .....	195
Height of body at posterior end of adipose and anal fins .....	105

Millimeters.

Height of body at beginning of caudal ..... 77  
 Ventrals under anterior third of dorsal; adipose fin over posterior end of anal; ventrals reach about  $\frac{1}{2}$  distance to vent; 24 scales in transverse series from origin of dorsal to lateral line; 125 scales in lateral line. Color silvery gray on back, black spots obsolete.

This species is evidently a close ally of the Atlantic salmon, belonging to the restricted subgenus *Salmo*. From *Salmo salar* it differs in the slightly larger mouth and rather different coloration and in very little else.

*Mykiss*, PENNANT, Arctic Zool., Intro., 126, 1792, Kamchatka; after KRASHENINNIKOF, etc. *Salmo mykiss*, WALBAUM, Artedi Piscium, 59, 1792, Kamchatka; based on *Mykiss* of PENNANT.

*Salmo penshinensis*, PALLAS, Zool. Rosso-Asiat., III, 1811, Gulf of Peshin.

*Salmo purpuratus*, PALLAS, Zool. Rosso-Asiat., III, 374, 1811, Bering Sea.

The correct names of the American Cutthroat Trout and its numerous known varieties are the following:

- 780. SALMO CLARKII (Richardson).
- 780(a). SALMO CLARKII LEWISI (Girard).
- 780(b). SALMO CLARKII GIBBSII (Suckley).
- 780(c). SALMO CLARKII HENSHAWI (Gil & Jordan).
- 780(d). SALMO CLARKII VIRGINALIS (Girard).
- 780(e). SALMO CLARKII SPILURUS (Cope).
- 780(f). SALMO CLARKII PLEURITICUS (Cope).
- 780(g). SALMO CLARKII BOUVIERI (Bendire).
- 780(h). SALMO CLARKII STOMIAS (Cope).
- 780(i). SALMO CLARKII MACDONALDI, Jordan & Evermann.

Page 500. Before *Salmo irideus* insert the following:

781(b). SALMO GAIRDNERI BEARDSLEEI, Jordan & Seale.

(BLUEBACK TROUT OF LAKE CRESCENT.)

Head  $3\frac{1}{2}$  in length to base of caudal; depth about 4; eye  $4\frac{1}{2}$  in head,  $1\frac{1}{2}$  in snout; scales 24-130-20, 130 cross series, those in front of dorsal numerous, about 70 if counted along median line, 60 if the rows along upper side are counted; dorsal with 10 branched rays; anal with 11 branched rays; branchiostegals 11; gill rakers 8 + 13, rather long and slender, the longest nearly  $\frac{1}{2}$  in length, 7 to 9 in maxillary. Head pointed; mouth rather large; maxillary extending to hinder margin of eye,  $1\frac{1}{2}$  in head, with about 20 teeth; snout  $3\frac{1}{2}$  in head; preorbital very narrow, the maxillary almost touching the orbit; posterior suborbitals shorter than eye, about 6 in head; opercle not very broad, equal to eye, its free part  $4\frac{1}{2}$  in head; interorbital width  $3\frac{1}{2}$  in head, equal to snout; several large teeth along margin of tongue; no hyoid teeth; teeth on vomer in zigzag series. Origin of dorsal in middle of the length, margin slightly concave, the first ray  $1\frac{1}{2}$  times the last, the last ray being pointed, slightly greater than base,  $2\frac{1}{10}$  in head. Origin of anal midway between origin of dorsal and base of caudal, margin straight, the tip of the last ray slightly exerted; anterior rays  $3\frac{1}{2}$  times posterior, and equal to base of fin,  $2\frac{1}{2}$  in head. Adi-

Millimeters.  
 ..... 960  
 ..... 853  
 ..... 215  
 ..... 400  
 ..... 100  
 ..... 167  
 ..... 17  
 ..... 81  
 ..... 81  
 ..... 71  
 ..... 165  
 ..... 165  
 ..... 105

pose fin high and slender, situated above or anterior to end of anal. Pectorals  $1\frac{1}{2}$  in head; ventrals under middle of dorsal,  $2\frac{1}{2}$  in head. Caudal broad, nearly truncate, the middle portions abruptly lunate when spread open, with pointed angles, each lobe being somewhat convex on its edge; longest rays  $1\frac{1}{2}$  in head. Least depth of caudal peduncle  $2\frac{1}{2}$  in head. Pyloric caeca 50 to 60, short and thick, the longest about 3 in head. Color in spirits very dark blue above, sides abruptly brighter, with many scales abruptly silvery; below white, lower jaw white, its margin dusky; cheeks below suborbitals very dark; sides, top of head, dorsal, and caudal fins spotted, the spots all very small; pectorals and ventrals nearly colorless, without spots, and slightly dusky; adipose fin with 2 spots; tips of lower fins faintly tinged with yellowish. Two specimens, each 16 inches long, Nos. 1861 and 1862, L. S. Jr. Univ. Mus. They were taken on March 12 and 16, 1896, in Lake Crescent, by Mrs. George E. Mitchell, of Fairholme, and sent to us by Mr. M. J. Carrigan, of Port Angeles.

A third specimen shows the following characters: Head  $3\frac{3}{8}$ ; depth  $3\frac{3}{8}$ . D. 12; A. 12 branched rays; branchiostegals 11 or 12; scales 23-123-26, 61 before dorsal; snout  $2\frac{3}{8}$ ; eye  $7\frac{3}{8}$ ; maxillary  $1\frac{1}{2}$  in head, its depth 8 in its length. Body robust, little compressed; head large; maxillary moderate, extending beyond eye; opercle moderate, its width  $5\frac{3}{8}$  in head. Last ray of dorsal pointed. Caudal subtruncate, lunate mesially, each lobe somewhat convex, pointed at tip. Caudal peduncle short and thick. Series of vomerine teeth long, in double row. Color above dark green, with black spots, which are small and sparse on body, extending to below lateral line; many small spots on head, dorsal, and caudal; spots not more numerous behind than before; sides and belly bright silvery; no red on lower jaw; a faint pink shade along lateral line; pectorals colorless except the upper ray; ventrals and anal colorless; flesh pale; gill rakers removed. This specimen, male, was taken in Lake Crescent. Length  $26\frac{1}{2}$  inches; weight in life 14 pounds. This specimen differs from a large *gairdneri* most in the large scales. In addition the head is much larger, and the body deeper.

A fourth, still larger, specimen (No. 1865, L. S. Jr. Univ.), an old spent male, 27 inches long, has been still later received. It shows the following characters: D. 11; A. 12. Head  $3\frac{3}{8}$  in length; gill rakers 8 + 12, of medium size, rather broad but sharp pointed; opercle  $3\frac{1}{2}$  in head; eye 7 in head; branchiostegals 11; maxillary long, reaching beyond eye,  $1\frac{1}{2}$  in head, its width  $9\frac{1}{2}$  in length. A double row of sharp teeth extending to within a short distance of end, where they are replaced by a single row of slightly larger teeth; teeth on tongue rather large; no teeth on hyoid; teeth on vomer in zigzag series. Scales 137-26. This specimen, a spent male, has the flabby muscles and slimy, half-concealed scales of the spent male salmon. The dark dots are very numerous and small and show very distinctly on back and sides, as also on head and fins; there is a dull red lateral band on head and body—this is about an inch broad, its outlines diffuse; a black blotch on cheek; maxillary dusky with a red blotch toward its tip; lower jaw and branchiostegals dusky; pectoral, ventral, and anal dark; back dark green, belly dusky.



The following account of the life coloration of *Salmo beardaleei* is given by Mr. George E. Mitchell:

The Blueback Trout caught in Lake Crescent are on the back a deep dark-blue ultramarine color of a peculiar transparency, dotted with small round black spots from the size of a pin's head to a little larger. The 2 fins on the top of the back are a dark smoky color, also dotted as on back end, and are transparent. The tail is the color and transparency with dots also—same as the top fins. The side fins and the bottom fins are dead white and sometimes faintly tinged with a pinkish hue at the edges; the belly is white. Looking at the fish sideways the sides of the fish show the scales to be iridescent, the red flash predominating. The head has very much the polish of mother-of-pearl around the lower jaws and jowls, red and pale-blue colors predominating; under the eyes a few black spots; on top of head the blue much darker than on top of back—so dark, in fact, that the black spots on it look blacker than the rest. The nearer the shore these fish are caught the lighter the blue on back, the fish often having an impression of the surroundings distinctly marked on them.

The following notes are added by Admiral Beardslee:

HABITS.

The Blueback is a deep-water dweller; those taken by me in late October were caught at depths varying from 30 to 50 feet, on large spoons. They fought hard until brought near the surface, then gave up, and when landed were found puffed up with air. Specimens taken in spring and put in pools in mountain streams with other trout died very soon, while the others lived. The trout caught by Mr. Mitchell, in March, was taken near bottom, by a large spoon, and it is not on record that at so early a date one has previously been caught.

FLESH.

Light lemon color before cooking; devoid of the oily salmon flavor, and very excellent; whitening by cooking.

OVA.

October 28. The eggs in the large fish were in individual size, and in size of cluster much smaller than those of a salmon of the same size.

The following extracts from a letter from Mr. Carrigan, dated Port Angeles, April 30, are of much interest:

\* \* \* Answering your direct inquiries: The Beardslees and Crescents are readily distinguishable, and can always be told apart. There are no red spots at the points indicated on the Crescent trout—no markings to suggest the Cutthroat trout.

(Named for Admiral L. A. Beardslee, U. S. N., in recognition of his active and intelligent interest in American game fishes.)

*Salmo gairdneri beardaleei*, JORDAN & SEALE, Proc. Cal. Ac. Sci., ser. 2, vol. vi, 1896, 209, pl. 23, Crescent Lake, Clallam County, Washington. (Coll. Mrs. George E. Mitchell. Type, No. 1864, L. S. Jr. Univ. Mus.)

780(c). SALMO GAIRDNERI CRESCENTIS, Jordan & Beardslee.

(SPECKLED TROUT OR LAKE CRESCENT.)

Head  $3\frac{1}{2}$  in length to base of caudal; depth 5; exposed portion of eye 6 in head,  $1\frac{1}{2}$  in snout; scales 32-151-34, 151 cross series, 83 in front of dorsal; dorsal with 10 branched rays, anal with 11; branchiostegals 10; gill rakers 6 + 11, counting rudiments, these very short and thick, the longest but  $\frac{3}{8}$  inch in length,  $18\frac{1}{2}$  in maxillary; mouth large, maxillary extending much beyond eye,  $1\frac{1}{2}$  in head, with about 20 teeth; tongue with

the usual teeth; teeth on vomer in zigzag series; hyoid region of tongue without teeth. Snout  $3\frac{1}{2}$  in head; preorbital very narrow, not so wide as maxillary adjacent to it; the posterior suborbitals longer than eye,  $5\frac{1}{2}$  in head; opercle and subopercle very narrow, scarcely as wide as eye, the free part of opercle  $6\frac{1}{8}$  in head; interorbital width  $4\frac{1}{2}$  in head. Origin of dorsal in middle of length of body, its margin straight, anterior  $2\frac{1}{2}$  times posterior, and slightly longer than base,  $2\frac{1}{2}$  in head; last ray of dorsal pointed. Origin of anal midway between origin of dorsal and base of caudal, margin irregular, anterior rays 3 times length of posterior and equal to base of fin,  $2\frac{3}{8}$  in head. Adipose fin high and slender, situated immediately behind anal; pectoral  $1\frac{1}{2}$  in head; ventrals under middle of dorsal,  $2\frac{3}{8}$  in head; caudal broad, slightly emarginate, nearly truncate when spread, its corners not rounded, its longest rays  $1\frac{1}{2}$  in head; least depth of caudal peduncle  $3\frac{3}{8}$  in head. Pyloric caeca about 51, the longest about  $1\frac{1}{2}$  in head, and very slender. Color in alcohol, very dark steel blue above, becoming paler below, nearly white anteriorly on belly, where only the margins of the scales are punctate; no silvery anywhere; lower jaw dusky, a large black blotch on cheek between suborbital and premaxillary; sides, back, top of head, dorsal and caudal fins with few small dark spots; pectorals dusky, slightly spotted at base; anal slightly dusky, without spots; ventrals dusky with a few spots in middle; adipose fin with a few spots; lower fins all tipped with pale, probably yellowish red in life; spots all very small and faint, not confined to posterior part of body. The specimen before us, No. 1863, L. S. Jr. Univ., is a male, 18 $\frac{1}{2}$  inches long. It was taken at Fairholme on Lake Crescent, Clallam County, Washington, March 12, 1896, by Mrs. G. E. Mitchell, of Fairholme. (Named for Crescent Lake, Washington, the type locality.)

*Salmo gairdneri crescentia*, JORDAN & BEARDSLEE, Proc. Cal. Ac. Sci., ser. 2, vol. VI, 1896, 207, pl. 22, Crescent Lake, Clallam County, Washington. (Coll. Mrs. George E. Mitchell. Type, No. 1863, L. S. Jr. Univ. Mus.)

**Page 504.** Under *Cristiromer* for "Eastern North America" read "Northern North America." The genus occurs also in the lakes of Alaska and British Columbia.

**Page 508.** Before *Salvelinus alpinus* insert:

784(a). *SALVELINUS KUNDSCHA*, Pallas.

This seems to be a species very distinct from *S. malma*. A specimen in the United States National Museum (No. 33814) from Petropaulski has been described by Bean & Bean as follows:

Similar in form to *S. malma*, but the body stouter and less elongate. Head  $4\frac{1}{2}$  to  $4\frac{3}{8}$  ( $4\frac{1}{2}$  in the Tareinsky Bay specimen); depth  $4\frac{1}{2}$  to  $4\frac{3}{8}$ ; eye  $5\frac{1}{2}$  in head, 2 in interorbital, or  $1\frac{1}{2}$  in snout; maxillary reaching to or beyond vertical through posterior edge of orbit; upper jaw nearly  $\frac{1}{2}$  length of head; lower jaw slightly shorter than upper. Hyoid teeth feebly developed. Scales small, 36-195, 122 pores. Fins all short; origin of dorsal about midway between tip of snout and base of upper caudal lobe, the base of the fin nearly as long as the longest ray, or  $\frac{1}{2}$  as long as head, its

upper margin very slightly concave, the last ray 2 in the longest; adipose fin over end of anal, its width about  $\frac{1}{2}$  its length, which is about equal to eye; pectoral 7 to  $7\frac{1}{2}$  in body length; ventral under middle of dorsal, not nearly reaching vent, its length 2 in head; caudal emarginate, its middle rays  $\frac{1}{2}$  the outer; anal scarcely concave when expanded. Pyloric caeca 22; branchiostegals 12; gill rakers 6 + 10, the longest less than  $\frac{1}{2}$  eye. Color bluish gray above, whitish below; the sides with numerous large white spots, some of which are  $\frac{3}{4}$  as large as eye. (Bean & Bean.)

This species is said to be common from Kamchatka northward, but only 6 specimens are actually extant, 4 obtained at Petropaulski by Dr. Leonhard Stejneger and 1 by Col. N. Grobnitski, and now in the United States National Museum, and 1 obtained from Tarcinsky Bay by Mr. Gerald E. H. Barrett-Hamilton and now in the museum of Stanford University.

*Salmo kundtsha*, PALLAS, Zoogr. Rosso-Asiat., III, 250, 1811, Kamchatka.

*Salmo leucomenis*, PALLAS, Zoogr. Rosso-Asiat., III, 250, 1811, Kamchatka.

*Salmo curitus*, PALLAS, Zoogr. Rosso-Asiat., III, 251, 1811, Kuril Islands.

The true *Salvelinus malma* is very common at Unalaska, Kadiak, Komandorski Islands, and Petropaulski. Specimens from these various places are all alike. Head  $4\frac{1}{2}$  to  $4\frac{1}{2}$ ; depth  $4\frac{1}{2}$  to  $4\frac{3}{4}$ . Spots grayish, tinged with red, much smaller than eye. Caudal well forked; lower fins short; pectoral reaching halfway to vent. Hyoid teeth present. The head seems much shorter than in examples from the United States. The dwarf form from the little brook (Pyramid Creek) at the head of Captains Harbor agrees fully in form with large examples taken in the sea about Unalaska. The small ones are brighter in color and mature at 4 to 6 inches. The form occurring throughout the northwestern United States, and described on page 508 as *Salvelinus malma*, should apparently be regarded as a species distinct from *S. malma*, and would stand as—

784(a). SALVELINUS PARKEI (Suckley).

Page 515. Add this footnote to *Salvelinus oquassa marstoni*:

A specimen of *Salmo marstoni* sent me some days ago indicates a more distinct species than was at first supposed. This is the most slender of our charrs, apparently the swiftest. The male is gorgeous; brilliant red extends upon the back and onto the dorsal and caudal fins as well as upon the other fins. Though quite distinct, the species is nearer to *S. oquassa* than any other. (Garman, in lit., March 24, 1895.)

Page 524. After *Osmerus dentex* add:

704(a). OSMERUS ALBATROSSIS, Jordan & Gilbert, new species.

(KADIAK SMEIT.)

Head  $4\frac{3}{4}$ ; depth  $5\frac{1}{2}$ . D. 2, 10; A. 1, 20; scales 75; maxillary  $2\frac{1}{2}$ ; eye  $5\frac{1}{2}$ ; snout  $3\frac{3}{4}$ ; mandible 2; pectorals  $1\frac{1}{2}$ ; ventrals  $1\frac{1}{2}$ ; dorsal  $1\frac{3}{4}$ . Body elongate, moderately compressed; back elevated at nape so that anterior profile is somewhat depressed between and behind eyes; interorbital space  $3\frac{3}{4}$  in head. Mouth large, lower jaw heavy, strongly projecting; opercle with concentric striae; pectorals moderate; ventrals long; dorsal high; anal fin low, very long, its longest ray  $2\frac{3}{4}$  in head; caudal moderate, well forked; ventrals inserted before dorsal.

Scales small, deciduous, those on back still smaller; lateral line distinct. Gill rakers long and slender, about 12 below angle of arch, longest about as long as eye. Tongue with moderate teeth, the anterior 2 to 4 small hooked canines; upper jaw with small sharp teeth similar to those in lower jaw, none of them canine-like; small teeth on palatines and pterygoids; vomer with 2 very small canines scarcely fang-like. Color bluish above with bright reflections; scales margined with dark points; sides silvery with golden and coppery luster; inside of gill openings dusky; fins white, somewhat dotted. About Kadiak Island, Alaska. Two specimens caught in the upward haul of a dredge in Shelikof Straits, north of Karluk, Kadiak Island, Alaska, at *Albatross Station No. 3675*. The depth of the dredge haul was 109 fathoms, but these fishes were no doubt taken from near the surface. One specimen is 8, the other about 7 inches in length. The species is allied to *Osmerus dentex*, the Rainbow Smelt, but differs in the extremely long anal and in the very weak vomerine and lingual canines. The flesh is firm, as in *O. dentex*. (Named for the U. S. Fish Commission steamer *Albatross*.)

*Osmerus albatrossis*, JORDAN & GILBERT, Rept. Fur. Seal Invest., MS. 1898, Shelikof Straits, north of Karluk, Alaska.

*Mesopus* should replace *Hypomesus*. It is originally characterized on page 14 (not 168) Proc. Ac. Nat. Sci. Phila. 1862, *Hypomesus* on page 15. The ventrals are inserted below front of dorsal in *Mesopus* as in allied genera, and there are 8 branchiostegals as in allied groups. The feeble teeth distinguish *Mesopus* from *Osmerus*. The statement that the stomach is caecal in *Argentinidae* is true of a few genera only, and the character has no high systematic value. In *Mesopus pretiosus* and *Osmerus dentex*, the stomach is siphonal, as in *Salmonidae*. In *Thaleichthys pacificus*, however, the stomach forms a blind sac. The small number of pyloric caeca and the peculiar structure of the ovaries remain to define *Argentinidae* as a family distinct from *Salmonidae*.

**Page 525.** To the synonymy of *Hypomesus olidus* add:

*Osmerus oligodon*, KNER.

The species ranges south to Amur River.

**Page 530.** After *Bathylagus pacificus* add:

804(a). *BATHYLAGUS BOREALIS*, Gilbert.

Head  $4\frac{1}{2}$  to base of caudal; depth  $5\frac{3}{5}$ ; eye  $2\frac{1}{2}$  in head; snout  $2\frac{1}{2}$  in eye. D. 8; A. 19; ventral 8; pectoral 8. Scales in about 40 rows, judging from the scars; head scaleless. Interorbital width grooved, the groove widening posteriorly, opening onto the flat occipital region, which is not swollen. Width of cartilaginous portion of interorbital space  $\frac{1}{2}$  orbit; including the thin membranaceous plates which overarch the orbits, the interorbital width is  $\frac{3}{4}$  orbit. The anterior profile of snout declines gently, bringing the mesial portion of premaxillaries on a level with lower margin of pupil. Distance from tip of snout to end of maxillary slightly exceeding length of snout,  $2\frac{1}{2}$  in orbit. Opercle with 2 strong ridges diverging downward and backward from behind the eye. Front of dorsal midway

between front of snout and adipose fin; base of dorsal contained  $3\frac{1}{2}$  times in length of head. Ventrals inserted under posterior portion of dorsal. Free portion of adipose fin very long and narrow, rising above the base of the second and third anal rays before the last, its tip reaching rudimentary caudal rays when depressed; anal fin rather long, the base  $1\frac{1}{2}$  in head, the vent immediately before it. Length of tail much exceeding head,  $3\frac{1}{2}$  in total length without caudal. Uniform blackish brown on sides, the head and ventral region blue black. Differing from *B. pacificus* in its much greater depth, longer tail, longer anal fin, and flat occiput. Length 132 mm. Bering Sea, in deep water north of Unalaska; 2 specimens known. (*borealis*, northern.)

*Bathylagus borealis*, GILBERT, Rept. U. S. Fish Comm. 1893 (Dec. 9, 1896), 402, Bering Sea at Albatross Station 327, north of Unalaska, in 322 fathoms.

804(b). BATHYLAGUS MILLERI, Jordan & Gilbert, new species.

Distinguished by the posterior insertion of the dorsal fin and the greatly swollen occipital region provided with a median keel. The type is in very poor condition, the skin being largely denuded from head and body. No traces remain of the scales, the pectoral and ventral fins are lost and the others greatly mutilated. Enough remains, however, to demonstrate that it is distinct from all known species and to furnish characters by which the species may be recognized. The interorbital space is converted into a very deep channel by 2 vertical thin lamellæ which arise on either side, and mark off the narrow interorbital space from the contiguous supraocular areas. From the base of these vertical lamellæ arise externally the thin supraocular plates, which extend outward and upward and roof over the orbit. A deep narrow channel is included between the lamellæ and the plates. The floor of the interorbital groove is raised mesially into a sharp ridge, which is continuous anteriorly with the ethmoidal ridge and posteriorly with a ridge running along middle of occiput. On anterior half of occiput this ridge is a high strong keel; posteriorly, it becomes lower and rounded. The occipital region is swollen and prominent, much higher than the interorbital space. It is bounded laterally by 2 strong rounded ridges which originate at the upper posterior margin of the orbit and converge rapidly backward. The occipital cartilage is heavy and strong, not yielding readily to pressure. The width of interorbital space is  $\frac{1}{2}$  orbit; the distance between outer margins of orbital plates above middle of eyes is  $\frac{3}{4}$  diameter of eye. The opercle is marked with delicate striæ diverging downward and backward, but is without strong ridges. The front of dorsal is midway between adipose fin and gill opening, slightly nearer base of caudal than tip of snout. The fin contains 8 rays. Anal badly mutilated, containing at least 24 rays. The mutilated condition of the type will not permit further description. Length 155 mm. Cortez Banks off San Diego, California, in deep water; known only from the type taken by the *Albatross* at Station 3627, in 776 fathoms. (Named for Walter Miller, professor of classical philology in Leland Stanford Jr. University, in recognition of his intelligent interest in zoological nomenclature.)

Page 531. In key under *h*, for "incomplete" read "complete."

Page 537. We can not separate *Synodus jenkinsi* from *Synodus scituliceps*, and the former name should probably be abandoned.

Page 555. *Macrostoma angustidens* and related species need further study. The synonymy and application of the names *angustidens*, *elongatus*, and *resplendens* are uncertain. *Macrostoma brachycheir* is probably a good species.

In *M. caudispinosum* the dorsal has 20, not 36, rays.

Page 580. The generic name *Bonapartia*, Goode & Bean, is preoccupied in ornithology. For its use in fishes the name *Zaphotias* is proposed, taking the same species (*pedaliota*) as type. The genus and its species would then stand as follows:

274. ZAPHOTIAS, Goode & Bean, new generic name.

(*Zaphotias*, having organs which emit light; ζα, intensive particle; φῶς, light.)

872. ZAPHOTIAS PEDALIOTUS (Goode & Bean).

Page 522. *Cyclothone microdon* occurs also in Bering Sea in very deep water.

Page 586. *Astroaesthes* is from ἄστρον, star; ἐσθής, vestment.

Page 594. *Plagyodus* (Steller) should probably supersede *Alepisaurus*, in which case the family becomes *Plagyodontidae*.

Page 603. *Sternoptyx diaphma* is common off both the Japanese and Hawaiian islands.

Page 608. For *Aldrorandia*, Goode & Bean, substitute the earlier name *Halosauropsis*, Collett.

*Halosauropsis*, COLLETT, Camp. Sci. Hirondelle, June, 1896, 143 (*macrocheir*).

Page 618. Add:

916(a). MACDONALDIA ALTA, Gill & Townsend.

D. 32; A. 31 to end of dorsal, 52 spines, 125 rays. Body comparatively high, greatest height equal to  $3\frac{1}{2}$  the distance between vent and tip of snout; pectoral fin with its root twice as far from upper cleft of branchial aperture as from the lateral line, and much nearer to the posterior end of operculum than to lateral line. Bering Sea; only the type known. (*altus*, deep.)

*Macdonaldia alta*, GILL & TOWNSEND, Proc. Biol. Soc. Wash., xi, 1897 (Sept. 17, 1897), 232, Bering Sea, Lat. N. 54° 54', Long. W. 168° 59', Albatross Station 36c, Aug. 13, 1895 in 1,401 fathoms. (Type, No. 4877a U. S. Nat. Mus. Coll. Albatross.)

916(b). MACDONALDIA LONGA, Gill & Townsend.

D. 33; A. 26 to opposite end of dorsal, 55 spines, 111 rays. Body comparatively slender, with the greatest height about  $\frac{1}{2}$  distance between vent and tip of snout; pectoral fin with its root 3 times as far from upper cleft of branchial aperture as from lateral line, and very much nearer

lateral line than end of operculum. Bering Sea; only the type known. (*longus*, long.)

*Macedonalia longa*, GILL & TOWNSEND, Proc. Biol. Soc. Wash., XI, 1897 (Sept. 17, 1897), 232, Bering Sea, Albatross Station 3607, 1895, in 900 fathoms. (Type, No. 48775, U. S. Nat. Mus. Coll. Albatross.)

Page 627. *Lucius vermiculatus* occurs also in Texas, specimens having been obtained in both the Trinity and Neches rivers near Palestine, by Evermann & Seovell.

Page 632. *Aplocheilus* = *Apocheilichthys* = *Haplocheilus* = *Panchar*, is a genus distinct from *Fundulus*, and should be erased from the synonymy of the latter. The genus is defined by the flat, much produced snout, and the long anal fin.

To the synonymy of *Fundulus* add:

*Planterus*, GARMAN, Monogr. Cyprinodonts, in Mem. M. C. Z., XIX, No. 1, 96, 1895 (*kansee* = *zebrinus*.)

Page 635. In the key, under *aa*, the phrase "inhabiting mountain springs and brooks" applies only to Nos. 943 and 944. It should be transferred and made a part of *s*.

Page 637. *Fundulus punctatus* and *F. vinetus* are wrongly placed by Garman in the synonymy of *F. parvipinnis*.

Page 638. *Fundulus pallidus* is placed by Garman in the synonymy of *F. grandis*, to which it bears but little resemblance.

Page 639. To the synonymy of *Fundulus majalis* add:

*Hydrargyra formosa*, STORER, Proc. Bost. Soc. Nat. Hist. 1837, 76.

Page 641. To the synonymy of *Fundulus heteroclitus macrolepidotus* add:

*Hydrargyra ornata*, LE SUEUR, Journ. Ac. Nat. Sci. Phila., I, 1817, 131, Delaware River, near Philadelphia. (Coll. G. Ord.)

Garman regards *Fundulus grandis* as a good species. We have recently compared specimens from Cape Cod with others from Tampa, and reach the same conclusion.

Page 642. Before *Fundulus ocellaris* insert:

932(c). *FUNDULUS HETEROCLITUS BADIUS*, Garman.

(This is the form found about Grand Manan, named but not characterized by Garman.)

Garman refers *Fundulus ocellaris* to the synonymy of *Fundulus grandis*, which is very doubtful.

Page 643. Garman refers *Fundulus fonticola* also to the synonymy of *F. grandis*, which is not correct. He also wrongly regards *Fundulus bernuda* as a variety of *heteroclitus*.

Page 644. *Fundulus robustus* is referred, probably by error, by Garman to the synonymy of *F. labialis*, which is certainly incorrectly made a variety of *F. parvipinnis*.

Page 645. Garman refers *Fundulus zebra*, *zebrinus*, and *extensus* to the synonymy of *Fundulus adinia*, all of which is certainly wrong. Such ref-

erences defy all our knowledge of the geographic distribution of these fishes. For example, *F. extensus* is a brackish-water fish of Cape San Lucas; *F. zebra*, which is the basis of *F. zebrinus*, is a fish of the mountain streams of New Mexico, Colorado, and northeastward, while *F. adinia* is found near the mouth of the Rio Grande. There is no doubt that the original *Fundulus zebra* is the species called *zebrinus* by us and *kansae* by Garman. It came from some point between "Fort Union and Fort Defiance." In other words, it came from the head waters of the Canadian River or the Rio Grande. No species of this type has been recorded from the upper Rio Grande, but the species called *zebrinus* and *kansae* is in all the upper waters of the Arkansas basin, to which the Canadian River belongs, and doubtless in the streams above Fort Union.

**Page 646.** To the synonymy of *Fundulus zebrinus* add:

*Fundulus kansae*, GARMAN, Monogr. Cyprinodonts, 103, pl. 2, fig. 10, 1895, Kansas.

This species (*F. zebrinus*) is rightly made the type of a new subgenus, or possibly genus, *Plancterus*, by Garman. It has long, convoluted intestines and very small pharyngeals. *Fundulus seminolis* (subgenus *Fontinus*) has short intestines and coarse pharyngeals.

**Page 648.** *Fundulus stellifer* is wrongly referred by Garman to the synonymy of *F. catenatus*.

**Page 649.** *Fundulus lineatus* is referred by Garman to the synonymy of *F. sciadicus*, which reference seems to be correct.

*Fundulus albofasciatus*, which Garman also refers to *F. sciadicus*, seems to be a perfectly good species. It is certainly not *F. sciadicus*.

Garman refers *Fundulus confluentus* to the synonymy of *F. grandis*, which is probably not correct.

**Page 650.** Garman's reference of *Fundulus funduloides* to the synonymy of *F. grandis* may be correct.

The species called *Fundulus dovii* in the text is an *Aplocheilus* and should stand as:

968(a). *APLOCHEILUS DOVII* (Günther)

Garman recognizes *Zygonectes* as a distinct genus, but its boundaries are not easily defined.

The description of *Fundulus confluentus* should be modified to include the following, taken from the type: Head  $3\frac{3}{4}$ ; depth  $4\frac{1}{2}$ . D. 11; scales 44 or fewer. A black spot on middle of membrane of last 3 dorsal rays. This species resembles *F. diaphanus* rather than *F. majalis*. *Fundulus ocellaris* seems to be identical with *F. confluentus*.

**Page 651.** Garman refers *Fundulus macdonaldi* to the synonymy of *F. sciadicus*, which is probably correct, but the reference of *F. floripinnis* to the same synonymy is certainly wrong.

**Page 652.** Garman refers *F. pulvereus* to the synonymy of *F. grandis*, which is without warrant.

**Page 655.** To the synonymy of *Fundulus chrysotus* add:

*Gambusia arlingtonia*, GOODE & BEAN, Proc. U. S. Nat. Mus. 1879, 118, Arlington River, Florida. (Type, No. 21308. Coll. Dr. Goode.)



*Zygonectes henshalli*, JORDAN, Proc. U. S. Nat. Mus. 1879, 237, San Sebastian River, Florida. (Type, No. 23449. Coll. Dr. James A. Henshall.)

To the synonymy of *Fundulus cingulatus* add:

*Zygonectes rubrifrons*, JORDAN, Proc. U. S. Nat. Mus. 1879, 237, San Sebastian River, Florida. (Type, No. 23450. Coll. Dr. James A. Henshall.)

*Zygonectes auroguttatus*, HAY, Proc. U. S. Nat. Mus. 1885, 556, Westville, Florida. (Type, No. 37362. Coll. Mann & Davison.)

An examination of a large amount of material recently collected in Florida by Drs. Evermann and Kendall shows that the synonymy of these species should stand as indicated above.

Examination of the type of *Gambusia arlingtonia* shows it to be the young of the form hitherto known as *Z. henshalli*, which, from an examination of the type and other specimens, proves to be the female of *Fundulus chrysotus*. The dorsal in *Gambusia arlingtonia* is not inserted so far back as the sixth anal ray, but is rather over the third or fourth. Both *G. arlingtonia* and *Z. henshalli*, agree with descriptions of *F. chrysotus* except in coloration. Both are females, as shown by form of anal fin. All specimens examined of the *henshalli* form are females, as shown in part by dissection and by the form of the anal fin. All specimens examined of the form agreeing with descriptions of *F. chrysotus* prove to be males, as shown partly by dissection and by the form of the anal fin. Front series of teeth much enlarged in all; anal fin usually with 11 rays.

The type of *Z. rubrifrons* differs from that of *Z. henshalli* in having a heavier head, really longer snout, mandible more oblique, giving the muzzle a truncated appearance, and the slope of the back to the snout beginning farther forward.

The type of *Z. rubrifrons* agrees with the description of *F. cingulatus*, except in the number of anal rays, there being 10 instead of 8, as given in the description, which is a redescription of the type of *F. cingulatus*. Cuvier & Valenciennes, however, give 10 anal rays in the original description. Specimens in the United States National Museum labeled *Zygonectes cingulatus*, from Pensacola, Florida, agree with the type of *Z. rubrifrons*, with the exception of 1 specimen, which has 11 anal rays.

National Museum specimens collected by Dr. Shufeldt at New Orleans, labelled *Zygonectes chrysotus*, contain both the *Z. henshalli* and *Z. chrysotus* forms, i. e., those with pearly spots and no cross bars (females) and those with dark cross bars (males)—that is, male and female of *Fundulus chrysotus*. Comparison of specimens collected at Tampa and Welaka, Florida, reveal 2 color forms. Most of those from Tampa have the heavier head, truncated muzzle, and outlines of *F. cingulatus*. The 2 color forms are those with dark cross bars, all males as shown by dissection and form of anal fin, and those with no cross bars and no pearly spots, which are all females. The majority of individuals have 10 anal rays each.

Most of the Welaka specimens have more slender and pointed head, pre-orbital less deep, really shorter snout, and the curve of the body toward the snout beginning farther back than in the preceding, and the majority have each 11 anal rays. The 2 color forms represent the 2 sexes—females with pearly spots and no cross bars, and males with dark cross bars and many with small brown spots.

While a few of the *chrysotus* form are found in the Tampa collection, and a few of the *cingulatus* form with the Welaka lot, they can be easily distinguished. A very few of the *cingulatus* form have 11 anal rays and a very few of *chrysotus* 10, but they can be otherwise distinguished. Whereas the females of *F. cingulatus* have no trace of pearly spots the females of *F. chrysotus* almost invariably have them. As a rule, the cross bars in the male, *F. cingulatus*, are narrower and more numerous than in the male of *F. chrysotus*, though young individuals of the latter do not differ in this respect. In *Fundulus cingulatus* there are often faint spots on the scales of the back forming longitudinal lines which seem to be absent in *F. chrysotus*. The teeth in the front row of *F. chrysotus* are larger than in *F. cingulatus*.

**Page 658.** Garman refers *Fundulus guttatus* to *F. nottii*, which is very doubtful, but he is right in so referring *F. hieroglyphicus*. He also refers *F. dispar* to *F. nottii*, which is probably wrong.

**Page 658.** *Fundulus guttatus* (Agassiz) can not be separated from *Fundulus nottii* (Agassiz).

**Page 659.** *Fundulus melapleurus* is, as Garman observes, a *Gambusia*, and should stand as *Gambusia melapleura*.

*Adinia guatemalensis* and *A. pachycephala* are recklessly referred by Garman to the synonymy of *Fundulus parvipinnis*. They might just as well have been placed at random under any other species of a totally different fauna.

**Page 660.** Before *Adinia* insert:

300(a). APLOCHEILUS, McClelland.

Snout flat, both jaws much depressed. Bones of mandible firmly united; upper jaw protractile; each jaw with a narrow band of villiform teeth. Body oblong, depressed anteriorly, compressed posteriorly. Dorsal fin short, commencing behind the origin of the anal, which is more or less elongate. Intestinal tract but slightly convoluted; air bladder present. (Günther.)

*Aplocheilus*, McCLELLAND, Ind. Cypr. As. Res., XIX, 301, 1839 (*chrysoatigmus* = *panchax*).  
*Panchax*, CUVIER & VALENCIENNES, Hist. Nat. Poiss., XVIII, 380, 1846 (*panchax*).  
*Haplochilus*, GÜNTHER, Cat., VI, 310, 1866, corrected spelling.

968(a). APLOCHEILUS DOVII (Günther).

For description and synonymy see p. 650.

**Page 662.** According to Garman the air bladder is present in *Rivulus*. He refers *R. marmoratus* to the synonymy of *R. cylindraceus*, which is probably correct.

Add the following species:

973(a). RIVULUS ISTHMENSIS, Garman.

Head  $3\frac{1}{2}$  in body; eye 3 in head; snout 6. D. 9; A. 11; V. 6; P. 15; scales 32-8. Elongate, compressed posteriorly, depressed forward; head broad, much depressed, flattened on the crown; snout medium, blunt;

interorbital width greater than eye. Origin of dorsal fin over middle of base of anal,  $\frac{1}{2}$  distance from snout to base of caudal; origin of anal fin midway between head and caudal, the last ray nearly as far back as that of dorsal; caudal elongate, pointed, as long as head. Light olivaceous, with a dark blotch at base of dorsal and another on back above or in front of first anal ray; apparently a light, transverse streak at base of caudal. Rio San Jose, Costa Rica.

*Rivulus isthmensis*, GARMAN, The Cyprinodonts, Mem. Mus. Comp. Zool., XIX, No. 1, July, 1895, 140, Rio San Jose, Costa Rica. (Type in M. C. Z.)

**Page 663.** *Lucania ommata* is wrongly referred to the synonymy of *Heterandria formosa* by Garman.

**Page 664.** The species called *Lucania goodei* in the text has 2 rows of teeth and is a true *Fundulus*, or rather *Zygonectes*, as Garman has shown. It may stand as *Fundulus goodei*.

**Page 665.** *Lucania venusta* is wrongly referred by Garman to *L. parva*, to which, however, it is closely related.

**Page 668.** Garman wrongly refers *Characodon bilineatus* and *C. variatus* to the synonymy of *C. lateralis*.

**Page 669.** Add:

883(a). CHARACODON EISENI, Rutter.

Head  $3\frac{1}{2}$ ; depth  $3\frac{1}{2}$ ; eye 3. D. 11 to 13; A. 13; scales 30 to 32-12. Snout shorter than eye, lower jaw projecting. About 9 teeth in upper jaw and about 14 in lower; teeth strongly bicuspid, the villiform teeth not developed. Mouth almost vertical when closed, mandible about  $\frac{1}{2}$  length of eye; interorbital space flat, the anterior part equal to orbit, wider posteriorly. Insertion of dorsal in middle of total length; anal inserted under fourth ray of dorsal; pectoral reaching past insertion of ventral; tips of depressed dorsal and anal in vertical through middle of caudal peduncle; caudal broad, truncate, length of middle rays equal to length of top of caudal peduncle. Head about  $\frac{1}{2}$  of total; greatest depth of body above ventrals; depth of caudal peduncle  $\frac{1}{2}$  its length. Color in alcohol, male with a broad indefinite lateral band; female with dark blotches on sides which in 1 of 3 specimens form distinct cross bands. This species is most closely related to *Characodon variatus*, Bean. It differs from that species in having fewer rays and scales, much fewer teeth, larger eye, much more posterior position of dorsal, and in color. Length  $1\frac{1}{2}$  inches. Rio Grande de Santiago, Tepic, Mexico.

*Characodon eiseni*, RUTTER, Proc. Cal. Ac. Sci. 1896, 266, Rio Grande de Santiago, Tepic, Mexico. (Type, No. 4999, L. S. Jr. Univ. Mus. Coll. Dr. Gustav Eisen.)

**Page 670.** Add the following:

The specimens from Parras, Mexico, referred by Garman to *C. lateralis*, appear to be new. They may be described as follows:

984(a). CHARACODON GARMANI, Jordan & Evermann, new species.

B. 4; D. 12; A. 12; V. 6; P. 17; scales 32-11 or 12; vertebrae 15 + 18. Body compressed, moderately stout, caudal pedicel deep, back gently

arched. Head about  $\frac{1}{3}$  of length to base of caudal; very little arched transversely. Snout short, not as long as the eye; chin steep. Mouth medium; upper jaw protractile. Teeth in outer series bicuspid. Eye large, nearly equal to interorbital space,  $\frac{1}{3}$  longer than snout,  $\frac{2}{7}$  of head. The specimen examined had 4 branchiostegal rays on each side; whether this is normal must be decided from others. Fins small; dorsal origin about  $\frac{2}{3}$  of the distance from snout to caudal; anal opposed to dorsal; posterior margin of caudal subtruncate. Olive to reddish brown, with scattered small spots of darker on the back, a darker band with or without spots of dark along the flank, more distinct posteriorly. Fins with fine dots of dark color. Parras, Coahuila, Mexico. (Named for Prof. Samuel Garman of the Museum of Comparative Zoology, in recognition of his valuable studies of the *Cyprinodonts*.)

*Caracodon lateralis*, GARMAN, The Cyprinodonts, Mem. Mus. Comp. Zool., XIX, No. 1, pl. 1, fig. 9, 1895, Parras, Coahuila, Mexico; not of GÜNTHER.

984(b). CHARACODON LUITPOLDII, Steindachner.

Head  $4\frac{1}{3}$  to  $4\frac{2}{3}$ ; depth 3 to  $3\frac{1}{2}$ ; eye 4 to  $4\frac{1}{2}$  in head; snout 3 to  $3\frac{1}{2}$ ; interorbital  $1\frac{1}{2}$  to 2. D. 14; A. 15 or 16; P. 15 or 16; V. 6; scales 40-17. Body moderately slender; caudal peduncle strongly compressed; head short; upper profile slightly arched, somewhat depressed at occiput; ventral outline more convex; bases of anal and dorsal quite oblique. Dorsal rounded, longest ray  $1\frac{1}{2}$  in head; anal somewhat smaller. Outer teeth slender, movable, broadened toward front of jaw which is notched; behind these a band of minute teeth, scarcely distinguishable. Two rows of scales below eye; preorbital, jaws, and narrow border of preopercle scaleless. Pectoral shorter than head, not reaching ventrals, which are nearer snout than base of caudal; origin of dorsal nearer base of caudal than gill opening; anal slightly behind dorsal. Color in alcohol, upper half of body light brown or brownish gray, lighter gray or silvery gray below, fading to yellowish white toward ventral line; a silvery gray band along middle of side, not well defined, its width that of 1 or 2 scales. Lake Pátzcuaro, Mexico. (Steindachner.)

*Characodon luitpoldii*, STEINDACHNER, Einige Fischarten Mexico, 12, pl. 2, figs. 3-3b, 1895, Lake Pátzcuaro, Mexico. (Coll. Princess Theresa von Bayern.)

**Page 675.** Garman refers *Cyprinodon elegans* to the synonymy of *C. eximius* and *C. felicianus* to that of *C. riverendi*, both of which seem to be correct.

**Page 680.** *Gambusia infans* is probably identical with *G. gracilis*, as indicated by Garman.

**Page 681.** Garman calls our *Gambusia affinis G. patruelis* and makes *G. holbrooki*, the northern form, a distinct species, neither of which views seems to be justifiable.

**Page 682.** *Gambusia nobilis* and *G. nicaraguensis* are referred by Garman to the synonymy of *G. gracilis*, which is questionable; but his reference to *G. punctulata* of *G. picturata* is probably correct.

Page 682. After *Gambusia affinis* add:

1000(a). GAMBUSIA TRIDENTIGER, Garman.

Head  $4\frac{1}{2}$ ; depth at anal  $4\frac{1}{2}$ ; snout short, not as long as eye, narrow, rounded forward, and blunt. D. 7 or 8; A. 10; V. 6; P. 12; scales 28 to 30-8; vertebrae 14 + 17. Mouth medium, directed obliquely upward; lower jaw longer than the upper, which is short, narrow, and protractile. Teeth in the outer series larger, strongly hooked, pointed, broadened somewhat toward the apex; inner series very small, in bands, tricuspid as in *Pacilia*; pharyngeal with a shoulder. Eye large, longer than snout, 3 in head. Fins small, excepting the caudal; dorsal smaller than anal and farther back, its origin about midway from occiput to end of caudal, nearly above the hindmost anal ray, 17 or 18 scales from the head; anal origin midway between snout and end of caudal; farther forward on the male, between the ventrals, and the fin is modified to form an intromittent organ about  $\frac{1}{2}$  length of entire fish; caudal deep, as long as head, rounded on hinder margin. Scales large, median series on flank as wide as eye. Intestine short. Light olivaceous, yellowish or brownish, with 7 or 8 vertical bars of brownish, separated by light or silvery spaces of equal width, on the sides of the caudal portion, edges of scales darker, the centers or median series more or less silvery; belly and lower surface of head silvery or golden; peritoneum black, showing through abdominal wall; occiput dark; top of snout light; a dark line between anal and caudal; dorsal with a faint spot or group of punctuations behind the middle near the base; other fins plain to dark tipped. (Garman.) Isthmus of Panama, in fresh water (*tridentiger*, bearing trifid teeth).

*Gambusia tridentiger*, GARMAN, Cyprinodonts, Mem. Mus. Comp. Zool., XIX, No. 1, 89, pl. 4, fig. 10, 1895, Isthmus of Panama.

Pages 688 and 689. *Heterandria versicolor* and *H. occidentalis* are correctly referred by Garman to the genus *Pacilia*. It is not improbable that *H. versicolor* is the same as *Pacilia vivipara*, Bloch & Schneider.

*Lebistes* is doubtless identical with *Pacilia*, as is also *Acropacilia*. *Acropacilia tridens* is probably identical with *Pacilia dominicensis*, as stated by Garman.

Page 691. Garman wrongly refers *Pacilia butleri* to the synonymy of *P. sphenops*.

Most of the Mexican and Central American species are imperfectly known and imperfectly described. Of these Garman refers the following to the synonymy of *P. sphenops*, whether correctly or not only a study of adequate material can determine: *Pacilia mexicana*, *P. thermalis*, *P. petenensis*, *P. dovii*, *P. conchiana*, *P. plumbeus*, *P. fasciatus*, and *P. spilurus*.

*Pacilia paronina* is referred, perhaps correctly, to the synonymy of *P. rittata*.

Page 696. Garman thinks that *Pacilia vandepolli* is identical with *P. reticulata*, Peters, which may be described as follows:

1032. PACILIA RETICULATA, Peters.

D. 7 or 8; A. 8 or 9; V. 5; scales 26 to 28-8. Depth of body  $\frac{2}{3}$  and length of head nearly  $\frac{1}{2}$  of the length to the base of the caudal. Males rather more

slender. Eye longer than snout, not quite  $\frac{1}{2}$  of head,  $\frac{3}{4}$  of interorbital space. Forehead flat. Dorsal origin somewhat nearer to end of snout than to end of caudal, opposite first ray of anal on females. Anal of male advanced, between the ventrals, which are elongate; anal process as long as the head, without hooks. Caudal large, rather longer than head, obtusely rounded; free portion of tail somewhat elongate, base of anal being  $\frac{1}{2}$  of its distance from the caudal; ventrals reaching anal; pectorals as long as the head, not reaching ventrals. Female yellowish olive, scales with a narrow blackish edge, belly silvery, trunk above the belly blackish. Male with 2 brown streaks along the trunk, sometimes confluent into a band, a brown streak along the middle of the side of the tail, a round black spot behind the shoulder, another at the commencement of the caudal streak, and a third at the root of the caudal; 1 or 2 of these spots may be absent. Trinidad; Venezuela (*reticulatus*, netted).

The male from Venezuela differs in color from those from Trinidad. It has large silvery patches between the brown streaks, and a large ovate black spot in the middle of the side of the tail. (Günther.)

NOTE.—The following is the original description: "Grüngelblich mit einem schwarzen Netzwerk, dessen Maschen den Rändern der Schuppen parallel liegen, am Bauche eiförmig. Schuppen in 7 Längs- und in 27 Querreihen; obwohl einige derselben durchbohrt erscheinen, ist doch keine deutliche Seitenlinie zu sehen. Ganze Länge 39, Höhe 9, Länge des Kopfes 7 Millimeter. D. 8; A. 10. Caracas; in dem Guayre-Flusse von Gollner gesammelt."

*Pocilia reticulata*, PETERS, Monatsb. Berl. Ak. 1859, 412, Caracas; GARMAN, Cyprinodonte, 63, 1895.

*Girardinus guppii*, GÜNTHER, Cat., VI, 353, 1866, Trinidad; Venezuela; EIGENMANN, Proc. U. S. Nat. Mus. 1891, 65.

*Girardinus vandepolli*, VAN LINDTH DE JEUDE, Notes from Leyden Museum, IX, 137, 1887, Curaçao, one of the Leeward Islands.

*Pocilia vandepolli arubensis*, VAN LINDTH DE JEUDE, Notes from Leyden Museum, IX, 137, 1887, Aruba, one of the Leeward Islands.

*Pocilia branneri*, EIGENMANN, Ann. N. Y. Ac. Sci. 1894, 629.

Page 697. Garman refers *Pocilia elongata*, one of the best marked species of large size, and marine in its habitat, to the synonymy of *P. gillii*. This is certainly wrong, as is also the reference to *P. gillii* of *P. chisoyensis* and *P. boucardi*.

*P. melanogaster* is probably correctly referred to *P. dominicensis*.

Page 698. Add:

1087 (a). *POCILIA CUNEATA*, Garman.

B. 5; D. 8 to 10; A. 10 or 9; V. 6; P. 15 or 16; scales 28 or 29-9. Short and deep; caudal pedicel deep. Head depressed, broad, flat on the crown, equaling depth between dorsal and anal, or  $\frac{1}{2}$  of the length to the base of the caudal; snout as long as the eye, broad, truncate; chin short, steep; mouth wide, directed upward; jaws weak, loosely joined, lower short, upper shorter, protractile; outer series of teeth slender, oar-shaped, hooked, movable; inner in bands, small, pointed; eye large, as long as snout,  $\frac{1}{2}$  of interorbital space,  $\frac{3}{4}$  of head. Dorsal larger than anal, origin midway from head to base of caudal, over third ray of anal, 13 scales behind the occiput. Anal small, acute angled, third ray longest; on the

male the base of the anal is forward of that of the dorsal, the fin is modified to form a sharp-pointed organ in which the rays are less changed than in most species; its length is less than that of the head. Ventrals small, not reaching the anal. Pectorals reaching back over 7 scales. Caudal deep, as long as the head, hind margin rounded. Scales large. Intestine long. Brownish, olive tinted, bases of scales dark, back darker, and top of head darkest; more or less of the hind margin, or  $\frac{1}{2}$  of the scale, is whitish to silvery on the scales of the flank; lighter to silvery under head and abdomen; dorsal with 1 to several transverse series of small spots of black; fin sometimes black tipped; a brownish streak extending back and upward on the opercle behind the eye; caudal with small spots of black on the basal half, or with a couple of clouded transverse bands; other fins uniform or punctulate; very small ones are lighter with a faint silvery band along the middle of the flank, but without vertical bars; a large one has numerous small white spots, somewhat like *Fundulus heteroclitus*. Females  $2\frac{1}{2}$  and males  $1\frac{1}{2}$  inches. Turbo, Gulf of Darien.

*Pœcilia cuneata*, GARMAN, Cyprinodonts, 62, pl. 5, fig. 3, 1895, Turbo, Gulf of Darien.

**Page 704.** After *Typhlichthys*, Girard, add:

a. No scleral cartilages; no pigment in or about the eye; retinal elements readily separable into ganglionic, inner reticular, and nuclear layers, the nuclear and outer reticular layers rarely distinguishable; diameter of eye about .150 mm.

SUBTERRANEUS, 1047.

aa. Scleral cartilages large, forming a hood over front of eye; a mass of pigment in front of eye; pigment layer of retina with more or less pigment; eye a mere vestige, about .040 mm. in diameter.

ROSE, 1047(a).

**Page 706.** After *Typhlichthys subterraneus* add:

1047(a). *TYPHLICHTHYS ROSE*, Eigenmann.

Extremely close to *T. subterraneus*, from which it seems to differ only in the less development of the eye. Scleral cartilages large, forming a hood over the front of the eye; a mass of pigment in front of eye; pigment layer of retina with more or less pigment; eye a mere vestige,  $\frac{1}{2}$  the size of that of *T. subterraneus*, about .040 mm. in diameter. The types of this species are 2 small, thoroughly dissected specimens, in the Museum of Indiana University, collected from a cave in Jasper County, Missouri, by Miss Ruth Hoppin. (Named for Mrs. Rosa Smith Eigenmann.)

*Typhlichthys rosæ*, EIGENMANN, Science, N. S., vol. VII, No. 164, 227, February 18, 1898, cave near Sarcoxie, Jasper County, Missouri.

**Page 723.** *Hemiramphus balao* is a valid species as defined.

**Page 729:**

*Exocetus volitans*, Linnæus, as Lönnberg has shown, is identical with *E. evolans* L. As the genus *Exocetus*, Syst. Nat., Ed. x, 316, is based solely on *Exocetus volitans*, the name *Exocetus* must go with this species, taking the place of *Halocypselus*. The ordinary flying fishes must therefore be called *Cypsilurus*. The species with long anal fin may, however, be held as generically distinct from the type of *Cypsilurus*, and for them (*ersiliens*, *rondeletii*, etc.) the name *Exonantes* has been proposed by Jordan & Evermann, Check List, 322. (Type, *ersiliens*.) (ἔξω, out of; ταυρῆς, swimmer.)

Our species of *Exonautes* are the following:

1080. *EXONAUTES EXSILIENS* (Müller).  
 1081. *EXONAUTES RONDELETHI* (Cuvier & Valenciennes).  
 1082. *EXONAUTES VINCIGUERRE* (Jordan & Meek).  
 1083. *EXONAUTES SPECULIGER* (Cuvier & Valenciennes).  
 1084. *EXONAUTES RUFIPINNIS* (Cuvier & Valenciennes).

To these should be added the following:

1084(a). *EXONAUTES AFFINIS* (Günther).

Head 4; depth 6; eye  $3\frac{1}{2}$ ; snout  $3\frac{1}{4}$ . D. 11 to 13; A. 11 to 13; scales 6-50 to 52, 35 before dorsal. Interorbital space flat, slightly greater than eye. Pectoral fin extending scarcely beyond dorsal and anal; base of ventral midway between eye and base of caudal, its rays reaching beyond middle of base of anal; dorsal opposite anal, its anterior rays  $2\frac{1}{2}$  in head. Pectoral with an oblique white blotch across its lower half, and with a narrow whitish margin; ventrals grayish. Cuba? Atlantic; West Africa. (Günther.) Probably distinct from *E. speculiger*.

*Exocoetus affinis*, GÜNTHER, Cat., VI, 288, 1860, Cuba?

The species of *Cypsilurus* are the following:

1085. *CYPSILURUS HETERURUS* (Rafinesque).  
 1086. *CYPSILURUS LUTKENI* (Jordan & Evermann).  
 1087. *CYPSILURUS FURCATUS* (Mitchill).  
 1088. *CYPSILURUS NIGRICANS* (Bennett).  
 1089. *CYPSILURUS XENOPTERUS* (Gilbert).  
 1090. *CYPSILURUS LINEATUS* (Cuvier & Valenciennes).

Under this species (p. 739) for Corea (in 3 places) read Gorea.

1091. *CYPSILURUS CYANOPTERUS* (Cuvier & Valenciennes).

This is a good species. The specimens recorded from James Island belong to *C. bahiensis*.

1092. *CYPSILURUS BAHIENSIS* (Ranzani).  
 1093. *CYPSILURUS CALIFORNICUS* (Cooper).  
 1094. *CYPSILURUS CALLOPTERUS* (Günther).  
 1095. *CYPSILURUS GIBBIFRONS* (Cuvier & Valenciennes).

Page 732. In the key, for "jj" read "hh," for "jjj" read "hhh," for "kk" read "ii," for "ii" read "gg," and for "hh" read "f."

Page 746. According to the studies of Mr. Rutter and Dr. Gilbert all the forms of *Gasterosteus* should probably be reduced to a single species (*Gasterosteus aculeatus*), having 3 or 4 geographic varieties, each running into a number of forms which differ in the degree of armature of the body.

Page 749. After *Gasterosteus bispinosus curieri*, insert:

1100(a). *GASTEROSTEUS GLADIUNCULUS*, Kendall.

Head  $3\frac{1}{2}$ ; depth  $3\frac{1}{2}$ ; D. II-I, 10; A. I, 8. Head rather long; eye about 3 times in head; opercle not striate; body deep, compressed, with 5 lateral



dermal plates anteriorly counting from pectoral fin, none posteriorly; caudal peduncle short, naked, not keeled; innominate bone lanceolate, its width about 3 times in length; ventral spines rather long, about  $1\frac{1}{2}$  times in head, serrated above and below, a strong cusp at base on both upper and lower edge. Color in life, grass green, mottled and finely punctated with black on top of head and back; sides of head and body golden, with dark blotches; breast silvery, ventrals scarlet. In alcohol the back becomes smoky black, the mottling and black dots more distinct, the golden hue of the sides fades, becoming more or less silvery, the dark blotches more pronounced. Coast of Maine and Woods Hole, Massachusetts. (*gladiunculus*, little sword; sticklebacks being called by the boys about Portland, Maine, "Little swordfish.")

*Gasterosteus gladiunculus*, KENDALL, Proc. U. S. Nat. Mus. 1395, 623, off Seguin Island, Maine. (Type, No. 47589. Coll. *Grampus*.)

Page 754. *Aulostomus maculatus* is pinkish-red in life.

Page 757. *Fistularia tabacaria* has been recorded by Storer from Holmes Hole, Massachusetts, and H. M. Smith records it from Buzzards Bay, near Quisset, and from about Woods Hole.

Page 762. In the key to species of *Siphostoma* read:

eee. Dorsal covering 4 or 5 caudal (not body) rings.

o. Rings 16 to 18 + 29 to 33.

q. Rings 16 + 30 to 33; dorsal 30 to 34, on 3 + 5 rings.

Page 767. It is doubtful if *Siphostoma pelagicum* occurs in America. *S. roussseai* has probably been sometimes mistaken for it.

Page 768. After *Siphostoma jonesi* add:

1124(a). *SIPHOSTOMA ROBERTSI*, Jordan & Rutter.

Head  $7\frac{1}{2}$  in length; depth  $2\frac{1}{2}$  in head; eye  $5\frac{1}{2}$  in head. Dorsal 20, on 4 rings; segments 17 + 32. Snout  $2\frac{1}{2}$  in head, with a slight keel; a slight keel on top of head, another above opercle, and 1 on anterior side of opercle, but not reaching posterior edge; shields without spines; lateral keel ending on last body segment; ventral keel on next to last; upper body keel extending nearly to end of dorsal fin, upper caudal beginning below it on first caudal segment; all ridges of body very prominent, the tail with 4 plain ridges; caudal pouch 3 in total length. Color mottled brown, paler below, the membrane connecting the segments pale bluish, forming cross stripes which are especially marked on the egg pouch; prominent pale cross bars on lower side of head; dorsal colorless, except that the base is finely dusted with brown; caudal thickly dusted with brown, except near base. This species is most closely related to *Siphostoma jonesi*, differing in having a shorter dorsal with more rays, and in the lateral keel ending distinct from lower caudal keel. Jamaica; 1 specimen,  $4\frac{1}{2}$  inches long, known. (Named for Rev. Joseph Seed Roberts, who collected the type.)

*Siphostoma robertsi*, JORDAN & RUTTER, Proc. Ac. Nat. Sci. Phila. 1897, 97, Kingston, Jamaica. (Type, No. 4988, L. S. Jr. Univ. Mus.)

## 1124(b). SIPHOSTOMA STARKSII, Jordan &amp; Culver.

Head  $10\frac{1}{2}$ ; depth 21. Dorsal 38, on 0 + 10 or 11 rings. Rings 13 or 14 + 37 or 38. Head and body in tail 2. Snout  $2\frac{3}{4}$  in head. Dorsal  $\frac{1}{2}$  longer than head. Body rather stout. Head scarcely carinate above. Snout with a slight smooth carina. Two lateral keels, confluent into 1 behind. Belly slightly keeled; no keel on opercle. Color dark olive, much mottled with darker but without distinct markings; yellow below. Male and female common in the fresh waters of Rio Presidio at Mazatlan, among algae; not seen in salt or brackish water. The pouch of the male teeming with eggs in January. Length 4 to 6 inches. Mazatlan, Mexico. Common in the Rio Presidio in sluggish water, on the bottom, about a mile below the village of Presidio. The species is probably found in brackish and fresh waters rather than in the sea.

*Siphostoma starksii*, JORDAN & CULVER, Fishes Sinaloa, in Proc. Cal. Ac. Sci. 1895, 416, pl. 30, Rio Presidio, Mazatlan. (Type, No. 2686, L. S. Jr. Univ. Mus. Coll. Hopkins Exped. to Mazatlan.)

## 1124(c). SIPHOSTOMA SINALOAE, Jordan &amp; Starks, new species.

Allied to *Siphostoma arctum* Jenkins & Evermann.

Head  $8\frac{1}{2}$  in length to base of caudal; depth  $3\frac{1}{2}$  in head. Dorsal 26, on  $1\frac{1}{2}$  + 5 rings, 14 + 35. Snout  $1\frac{1}{2}$  in head, a strong median ridge above running to between middle of eyes, a ridge on each side from angle of mouth to below eye, occipital and nuchal plates keeled, a slight keel on anterior part of opercle; dorsal keels ceasing in front of the last 4 or 5 rays of dorsal, the lateral ridge running up and continuing as dorsal ridges; belly with a keel on each side. Preanal part of belly  $1\frac{1}{2}$  in postanal part; pectoral shorter than eye, caudal 3 in head. Color olive brown above, abruptly lighter below lateral ridges anteriorly, the edges of the plates dark, forming reticulations on lower parts of body; between every 4 rings is a narrow white cross bar; from each eye is a narrow light bar running upward and backward to occiput; caudal dark. The 2 type specimens, 1 of which was sent to the British Museum, collected by the Hopkins Expedition at Mazatlan. They were erroneously referred to *Siphostoma arctum* in our paper on the Fishes of Sinaloa. Type, No. 2945, L. S. Jr. Univ. Museum.

Page 772. *Corythoichthys*, Kaup, should apparently be recognized as a genus distinct from *Siphostoma*. The species belonging in it are the following:

## 1134. CORYTHOICHTHYS ALBIROSTRIS, Heckel.

## 1135. CORYTHOICHTHYS CAYANNENSIS (Sauvage).

## 1135(a). CORYTHOICHTHYS CAYORUM, Evermann &amp; Kendall.

Head  $8\frac{3}{4}$ ; depth  $12\frac{3}{4}$ ; snout  $3\frac{1}{2}$  in head; eye  $4\frac{1}{2}$ . D. 21 rays, on  $1\frac{1}{2}$  +  $3\frac{1}{2}$  rings; A. 3, on first caudal ring; C. 10; P. 10. Rings  $17 + 26 = 43$ . Body short and stout; head short, snout very short; tail but little longer than head and trunk. Cranial ridges strong; a high, sharp keel on snout, the occipital keel very high, its edge convex, notched near the middle, not

continuous with keel on snout; a strong supraocular ridge, beginning opposite posterior end of nasal keel and continuing backward with 1 hiatus upon upper edge of opercle; just below this on the opercle another longer but scarcely stronger ridge; another short ridge on anterior part of opercle at level of lower part of eye; opercles very convex, as if swollen outward; keels on body and tail all strong; the 2 lateral keels on body terminating on third caudal ring; the 2 lateral keels on tail beginning on the last body ring, thus overlapping the body keels; median keel on side well developed, terminating on sixteenth body ring; ventral keels strong; abdominal keel very strong. Egg sac on first 18 caudal rings. Color yellowish brown, with darker punctulations; tip of snout white; cheek, throat, and under parts of snout white, crossed by about 7 or 8 irregular brownish bars extending downward and backward; opercles brown; fins pale. This species is related to *C. albirostris* of Heckel, differing from it chiefly in the shorter snout, smaller dorsal, and fewer rings. Key West, Florida. (*cayorum*, of the Keys; from Cayo Hueso, Bone Key, the original Spanish name of the island of Key West.)

*Corythoichthys cayorum*, EVERMANN & KENDALL, Bull. U. S. Fish Comm. 1897 (Feb. 9, 1898), 128, pl. 7, fig. 7, near Crawfish Bar, Key West, Florida. (Type, a male 3½ inches long, No. 48784. Coll. Drs. Evermann & Kendall.)

Page 774. *Syngnathus aquoreus* is doubtfully American. Until a comparison of specimens can be made our species may stand as—

1188. SYNGNATHUS HECKELI (Kaup).

Page 792. *Lethostole*, Jordan & Evermann, is identical with *Chirostoma*, and the definition assigned is that of *Chirostoma*.

To the synonymy of *Chirostoma estor* add:

*Atherinichthys albus*, STEINDACHNER, Anzeiger der Kais. Akad. Wiss. Wien, 1894, 148, Lake Pátzcuaro, Mexico. (Coll. Princess Therese von Bayern.)

Page 793. In *Chirostoma humboldtianum* the scales are serrulate. After this species insert the following:

1155(a). CHIROSTOMA GRANDOCULE (Steindachner).

Head 4; depth 5½; eye 3½ in head; interorbital width 4½; pectoral fin 1½; ventral 2½; caudal 1½; anal base 1½, its greatest height 1½. D. V-I, 10; A. I, 20; P. 15 or 16; scales 60 to 62-15 or 16. Upper profile of head merging gradually into that of back, rising slightly toward beginning of second dorsal. Lower jaw slightly projecting; posterior end of upper jaw reaching eye. Teeth on maxillary sharp, brush-like, in 3 or 4 rows, the inner teeth of the maxillary and the outer teeth of lower jaw somewhat enlarged and close set. Cheek narrower than in *C. humboldtianum* and *C. estor*, and with 4 rows of scales. Origin of first dorsal midway between anterior border of eye and base of caudal, the second dorsal ¼ diameter of eye nearer base of caudal than hinder border of eye; greatest height of second dorsal scarcely greater than base of fin. Longest anal ray about 1½ in base of fin; dorsal and anal concave on free border; origin of anal nearly an eye's diameter in front of that of second dorsal; caudal deeply incised, the mid-

dle rays about 2 in the longest; caudal peduncle more than  $4\frac{1}{2}$  in body, its least depth somewhat more than 2 in greatest depth of body. Scales slightly ctenoid. Side with a broad, sharply defined silvery-gray band. Body much more slender, snout shorter, and eye larger than in *C. humboldtianum* or *C. estor*. Length 5 inches. Lake Patzcuaro, Mexico.

*Atherinichthys grandoculis*, STEINDACHNER, Anzeiger der Kais. Akad. d. Wissensch. Wien. 1894, 149, Lake Patzcuaro, Mexico. (Coll. Princess Thereso von Bayern.)

### 354(a). ESLOPSARUM, Jordan & Evermann.

*Eslopsarum*, JORDAN & EVERMANN, Check-List Fishes, 330, 1896 (*Jordani*).

This genus is close to *Chirostoma*, from which it differs in the large entire scales. To it belong the 2 following species:

1156. ESLOPSARUM BARTONI (Jordan & Evermann).

1157. ESLOPSARUM JORDANI (Woolman).

To the synonymy of this species should be added

*Atherinichthys brevis*, STEINDACHNER, Anzeiger der Kais. Akad. d. Wissensch. Wien. 1894, 149, Lake Cuitzeo, Mexico. (Coll. Prinzessin Therese von Bayern.)

Page 793. In *Eslopsarum jordani* the anal is I, 16, not I, 6.

Page 795. *Kirtlandia laciniata* has been found to intergrade with *K. vagrans* and should stand as—

1158a. KIRTLANDIA VAGRANS LACINIATA (Swain).

Page 796. Under *d* in the key read:

*d*. Snout about equal to eye, which is 3 to  $3\frac{1}{2}$  in head.

Page 800. An examination of numerous specimens of *Menidia* from various places between Florida and Halifax shows that *M. notata* and *M. menidia* intergrade perfectly. The first will therefore stand as—

1167a. MENIDIA MENIDIA NOTATA (Mitchill).

Page 801. *Menidia guatemalensis* and *Menidia pachylepis* belong in the genus *Thyrina*, Jordan & Culver.

Page 819. *Agonostomus nasutus* has the anal usually II, 10, sometimes II, 9.

Page 821. Add the following:

In the Transactions of the Jamaica Society of Arts for 1855, Mr. Richard Hill gives a paper on "Fishes of the Jamaica Shores and Rivers" which has been overlooked by subsequent writers. The list is chiefly a nominal one, but it contains a number of vernacular names not elsewhere given. The only new species are given under the head of *Labrax* (page 142) and *Mugil* (page 143), and these are named rather than described. They are the following:

There is another *Labrax*, common enough in the Kingston market when the rains send strong freshets from the river into the harbor. The fishermen call it the river chub, and confound it with the *mucronatus*. It is a different species; it is marked with

bands like the *Perca fluviatilis* of Europe, and the *Perca granulata* of America. We will call it the *Labrax pluvialis*, rainy weather chub.

*Mugil petrosus*—rock mullet;  
*lineatus*—short mullet, 1;  
*albula*—short mullet, 2;  
*eurema*—long mullet;  
*equinoculus*—horse-eye mullet;  
*capitulinus*—drab mullet, long ears;  
*plumieri*—pond mullet;  
*liza*—callipeva;  
*Dajaus monticola*—mountain mullet;  
*choirorynchus*—hog-nose mullet.

The *Dajaus monticola* inhabits only the mountain streams; the *choirorynchus* or hog-nose mullet is a fish of double the size of the *monticola*, and found in the same waters. The *mugil liza* is the largest of the mullets, from 20 inches to 3 feet long; the callipeva is the name by which it is exclusively known. This is, no doubt, its Indian name. The *equinoculus* and *capitulinus*, known in the market as long mullets, are readily distinguishable from each other by the size of the head, and especially by the size of the eye: the horse-eye mullet has the large eye, the *capitulinus* unusually small. The *plumieri*, Plumier's mullet of Cuvier & Valenciennes, is a long mullet; and the *lineatus* and *albula* what the market people distinguish as short mullets. The callipeva is a river mullet seldom extending further than the embouchure of streams, or into the ponds and marshes. The *eurema* is a large mullet found on the sea banks; it is the most highly colored of all the mullets, the back is a golden green and it has scales on the second dorsal fin.

*Dajaus choirorynchus* is identical with *Agonostomus nasutus*, but the scanty description hardly justifies the substitution of this name for the later one. The other new species we fail to identify. *Labrax pluvialis* we do not recognize.

Page 823. The great *Barracuda* should stand as—

1199. SPHYRENA BARRACUDA (Walbaum).

To its synonymy add:

*Esox barracuda*, WALBAUM, *Artedl Piscium*, III, 94, 1792; after CATESBY.

Page 827. The ventrals in the *Polynemidae* are truly thoracic, the long pubic bone being attached to the shoulder girdle. This family is probably nearest allied to the *Scientidae*.

Page 833. After *Anmodytes personatus* add:

372(a). RHYNCHIAS, Gill, new genus.

*Rhynchias*, GILL, MS., new genus (*septipinnis*).

This generic name is provisionally given to a species known only from a description of Pallas, and supposed to differ from *Anmodytes* in the presence of ventral fins. It may prove to belong to some different family. (ρύγχος, snout.)

1214(a). RHYNCHIAS SEPTIPINNIS (Pallas).

This species has not been recognized by any recent collector, and it is not certain to what family it belongs. The following is the substance of Pallas's description:

D. 43; A. 24; V. 8; P. 16; C. 24. Form of *Anmodytes tobianus*. Head 3030—101

compressed; snout long, slender, depressed. Maxillary with fine teeth; rictus long. Branchiostegals 4. Body compressed, slender, with transverse streaks. Scales inconspicuous; 1 lateral line. Pectoral large, unarmed. Dorsal short, well backward, lower posteriorly; caudal sub-bifurcate. Color white, the dorsal edged with darker. Kamchatka. (Pallas.)

If we can trust the description, this fish would seem to represent a distinct genus of *Ammodytidae*, characterized by the presence of ventral fins, but it may be that the account is erroneous in this regard and that Pallas had in mind *Ammodytes personatus*. (*septem*, seven; *pinna*, fin.)

*Ammodytes septipinnis*, PALLAS, Rosso-Asiat., III, 1811, Kamchatka.

Page 833. *Ammodytes alascanus* is not separable from *A. personatus*.

Page 839. *Caulolepis longidens* occurs also in the Pacific, specimens having been collected by the *Albatross* at Cortez Banks, off San Diego, California, in 1896.

Page 847. Add:

1230(a). MYRIPRISTIS CLARIONENSIS, Gilbert.

Head  $3\frac{1}{2}$  in length; depth  $2\frac{5}{8}$ . D. X-I, 14; A. IV, 12; scales  $3\frac{1}{2}$ -41-7. Least depth of caudal peduncle  $\frac{1}{2}$  length of snout and eye. Greatest (oblique) diameter of eye  $2\frac{1}{4}$  in head. Least interorbital width equaling length of snout,  $4\frac{1}{4}$  in head. Mouth less oblique than in related species, the line of upper jaw with a more pronounced double curve. Lower jaw the longer, with well-developed symphyseal knob. Teeth finely villiform, very slightly enlarged toward middle of both jaws; wide patches of similar teeth on head of vomer and on palatine bones. Length of maxillary (measured from front of upper jaw) very slightly (about  $\frac{1}{10}$ ) less than length of snout and eye. Color before immersion in spirits, reddish, the upper parts dusky, especially on top of head and on the margins of the scales; evident horizontal dusky streaks between the rows of scales; opercular membrane blackish; fins all light, without dark markings. Differing from all known American species of *Myripristis* in having  $3\frac{1}{2}$  series of scales between the lateral line and the base of the spinous dorsal, instead of  $2\frac{1}{2}$ . Length  $6\frac{1}{2}$  inches. Revillagigedo Islands; only the type known.

*Myripristis clarionensis*, GILBERT, Proc. U. S. Nat. Mus. 1898, 441, pl. 69, Clarion Island, Revillagigedo Archipelago. (Type, No. 47746. Coll. Dr. Gilbert.)

Page 852. Insert the following description by Jordan & Rutter of *Holocentrus marianus*, based upon specimen 6 inches long from Jamaica:

Head  $2\frac{3}{8}$ ; depth 2 in length; eye  $2\frac{1}{4}$  in head. D. XI, 13; A. IV, 9; scales 4-45-7. Dorsal outline much more curved than ventral; mouth low, but little oblique, the lower jaw projecting and entering upper profile; maxillary to below middle of eye; eye large, lower margin of orbit cut by a line connecting tip of snout and upper base of pectoral; angle of opercle high, higher than top of pupil, with 3 sharp teeth, small teeth along the margin next the subopercle; subopercle long and narrow, dentate near upper end; preopercle very finely serrate, with a strong spine at angle; a single row of scales on opercle along margin of preopercle; suborbital

bones very narrow, finely serrate; premaxillary groove on top of head as long as eye; length of pectoral equals head behind middle of eye; spinous dorsal depressible into a groove, highest (anterior) rays of soft dorsal equal to ventrals, longer than soft rays of anal; third anal spine very long and heavy, as long as pectorals; caudal forked almost to base, the lobes equal, as long as pectorals. Each row of scales with a red band, yellow lines between the rows; fins all yellowish. This is a strongly marked species, very different from *Holocentrus ascensionis*, perhaps the type of a distinct genus, characterized by the large mouth and projecting chin.

**Page 855.** Dr. Bean reports the Red Mullet or Goat Fish (*Mullus auratus*) as being plentiful at Sandy Hook in September and October.

**Page 857.** The nominal genus *Mulloides* can not be separated from *Upeneus*.

**Page 866.** In *Scomber colias* read: Head about 3; depth  $4\frac{1}{2}$ ; first dorsal longer than high.

**Page 873.** To the synonymy of *Scomberomorus* add:

*Polipterus*, RAFINESQUE, Anal. de la Nature 1815, 84; substitute for *Scomberomorus*.

**Page 874.** In line 12 of description of *Scomberomorus maculatus*, for "side" read "part."

**Page 878.** *Bipinnula*, Jordan & Evermann, is a synonym of *Escolar*, Jordan and Evermann, in Goode and Bean, Oceanic Ichthyology, 519, 1896. The error resulted from Goode & Bean taking our original MS. name *Escolar*, for which we afterwards substituted *Bipinnula*.

This genus and its species will therefore stand as follows:

396. ESCOLAR, Jordan & Evermann.

*Escolar*, JORDAN & EVERMANN, in GOODE & BEAN, Oceanic Ichthyology, 519, Aug. 23, 1896 (*violaceus*).

*Bipinnula*, JORDAN & EVERMANN, Fishes North and Middle Amer., 878, Oct. 3, 1896 (*violaceus*).

1267. ESCOLAR VIOLACEUS (Bean).

**Page 886.** Instead of *Lepidopus caudatus*, which is not yet known to occur in American waters, insert:

1276. LEPIDOPUS XANTUSI, Goode & Bean.

Head  $4\frac{1}{2}$  in body; depth 3 in head; eye  $5\frac{1}{2}$ ; interorbital space  $8\frac{1}{2}$ ; snout 3; maxillary  $3\frac{1}{2}$ . D. 82; A. 11, 45. Jaws with long, sharp teeth in front, followed by single rows of weaker ones, arranged in groups of twos and threes. Height of dorsal, near middle of body, 3 in head. Anal preceded by 2 scutes, the first minute, the second wide, strongly keeled, its length  $\frac{1}{2}$  the diameter of eye. Pectorals of 12 rays, length 2 in head. Each ventral consists of a flat keeled spine followed by a minute ray. This species is known from 2 small mutilated specimens, both found on the beach near San Jose del Cabo, Cape San Lucas. The type was taken by John Xantus, about 1860, and recorded by Jordan & Gilbert as *Lepidopus caudatus*. The second, of about the same size ( $5\frac{1}{2}$  inches), was taken by

Richard C. McGregor, in 1897. From the latter the above account was taken. The species differs from *Lepidopus caudatus* in the much shorter dorsal and longer anal. D. 103; A. 24. (Named for John Xantus de Vesey.)

*Lepidopus caudatus*, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1882, 358; not of EUPHRASEN *Lepidopus xantusi*, GOODE & BEAN, Ocean. Ichth., 519, 1896; same type; no description.

**Page 889.** *Trichiurus lepturus* is recorded by Storer from Buzzards Bay (1840) and Wellfleet, Massachusetts (1845), and H. M. Smith records it from Woods Hole (1897).

**Page 892.** The synonymy at top of page under *Tetrapturus imperator* belongs to the footnote on same page.

**Page 899.** Add:

1286(a). *OLIGOPLITES MUNDUS*, Jordan & Starks, new species.

Head 4; depth  $2\frac{3}{4}$ ; eye  $4\frac{1}{2}$ . D. V-I, 19; A. II-I, 20. Body deep and compressed. Length of head about  $\frac{1}{6}$  greater than its depth at nape; eye equal to snout and to interorbital; maxillary extending considerably beyond vertical from hinder margin of eye, its length  $1\frac{1}{2}$  in head; second suborbital not over  $\frac{1}{2}$  as wide as lowest, and much shorter, thus forming a prominent notch in posterior margin of suborbital bones; a slight emargination in opercle in front of pectoral. Teeth small, sharp, in a band in each jaw, narrow in upper. Origin of soft dorsal midway between snout and base of caudal, the anal opposite; the anterior rays of both somewhat produced; second soft ray of each equal to head behind pupil, and equal to pectoral; ventrals equal to  $\frac{2}{3}$  of pectorals, their inner margins fastened to body; caudal deeply forked, the middle rays  $3\frac{1}{2}$  in longest, which are longer than head. Lateral line nearly straight, but forming a broad angle above pectoral. Color silvery on sides, becoming darker above; fins colorless. This species differs from *Oligoplites altus* in the much larger mouth and in having the suborbital bones notched posteriorly. *Oligoplites saliens* of the West Indies seems to be more elongate in body and with the suborbitals even behind as in *O. altus*. Pacific coast of tropical America.

This description is based on a specimen 11 inches long from San Juan Lagoon, Mexico, at the mouth of Ahome River, collected by the *Albatross*. Three other specimens from Algodones Lagoon, Mexico (*Albatross* Coll.), agree in every respect, except that 1 of them has but 4 free spines in front of dorsal.

Numerous other specimens have been since brought by Dr. Gilbert from Panama.

*Oligoplites mundus*, JORDAN & STARKS, in JORDAN & EVERMANN, Check-List Fishes, 344, 1896, Mazatlan, Mexico; name only.

**Page 909.** The Californian species *Trachurus symmetricus* is probably a species distinct from *T. picturatus*, described from Madeira. The two forms have never been properly compared.

**Page 912.** The identity of *Hemicaranx amblyrhynchus* with *Caranx falcatulus*, Holbrook, needs proof. The latter species, if distinct, may be described as follows:



## 1305(a). HEMICARANX FALCATUS (Holbrook).

Head 6 in total length; depth about 3. D. VII-I, 28; A. II-I, 25; C. 19; V. 5; P. 16; lateral line with 50 plates. Body oval, compressed; the head short, the facial outline descending in a gentle curve to snout, which is rounded though narrow. Eye large, in the middle third of the head, the posterior margin rather nearer snout than posterior margin of opercle; nostrils close together, nearly midway between eye and snout, and on a line within the orbit, the posterior larger, subround, the anterior ovoidal. Mouth small; each jaw with a single row of slender, conical teeth; a small patch of minute teeth on the vomer, and a small, narrow group of similar teeth on the palatines; tongue small, narrow, a few minute teeth near its base; pharyngeal bones armed with numerous cari-like teeth, longer than those of the jaws. Soft dorsal long and low, the first 3 or 4 rays moderately elevated, the fin sealed at base; pectoral falcate, very long, extending to anterior third of soft dorsal; ventral small, very short, reaching beyond vent; anal shaped like the soft dorsal; caudal very long and widely forked, the upper lobe more than  $\frac{1}{2}$  longer than the lower. Lateral line at first almost semicircular; at origin of soft dorsal descending to median plane, then straight; plates beginning with the soft dorsal increasing in size to the thirty-fifth, whence they decrease rapidly; scales minute, those of lateral line elongated quadrilateral, with 1 angle prolonged and rounded. Color, upper part of head and body above lateral line pale brown with slight bluish tint; lower jaw, opercle, and side yellowish; belly silvery, with a slight golden tint; anterior dorsal transparent; posterior transparent but with a yellowish tint; caudal yellowish. Known certainly only from Charleston, South Carolina.

*Caranx falcatus*, HOLBROOK, Ichth. South Carolina, 92, pl. 13, fig. 2, Charleston, South Carolina.

Page 914. Add:

## 1306(a). HEMICARANX ZELOTES, Gilbert, new species.

Head 4 to 4 $\frac{1}{2}$ ; depth 2 $\frac{2}{3}$  to 2 $\frac{3}{4}$ . D. VII-I, 26 to 29; A. II-I, 23 to 25; P. 20 to 22; scutes about 52. Body regularly elliptical, its greatest depth about in middle of its length, exclusive of caudal peduncle. Head small; anterior profile more decurved, and hence the snout is blunter than in *H. atrimanus*; depth of head just behind eye about  $\frac{2}{3}$  its length. Jaws subequal, tip of lower slightly projecting; maxillary narrow, not quite reaching anterior margin of pupil, about  $3\frac{1}{2}$  in head ( $3\frac{1}{2}$  to  $3\frac{3}{4}$  in *atrimanus*). A single series of small, close-set, subequal teeth in each jaw; no teeth on vomer, palatines, or tongue. Orbit considerably greater than snout,  $3\frac{1}{2}$  to  $3\frac{3}{4}$  in head. Interorbital width (taken at anterior margin of orbit) slightly less than orbit. Occiput with an evident carina. Distance from snout to first dorsal spine greater than length of pectoral. Spinous dorsal very low, the highest spine considerably less than orbit (greater than orbit in *atrimanus*); a well-developed antrorse spine before the dorsal; soft dorsal and anal similar, not falcate, the rays decreasing in size from the first; highest ray of soft dorsal 2 to 2 $\frac{1}{2}$  in head; highest ray of anal about 2 $\frac{1}{2}$  in head; dorsal and anal depressible into a high sheath of scales, the last 3 or 4 rays

uncovered; caudal fin wide, well forked, the upper lobe the longer, the longest ray not quite  $\frac{1}{2}$  total length of body; pectoral fin long,  $3\frac{1}{4}$  to  $3\frac{1}{2}$  in body ( $2\frac{3}{4}$  to  $2\frac{1}{2}$  in *atrimanus*); ventrals  $2\frac{3}{4}$  to  $2\frac{1}{2}$  in head. Scales as in *atrimanus*; lateral line with a very strong curve anteriorly, the height of the curve  $2\frac{3}{4}$  to  $3\frac{1}{2}$  in its length; its length  $2\frac{1}{4}$  to  $2\frac{1}{2}$  in the straight portion; entire length of straight portion with scutes, which are very small in front and behind; scutes considerably wider and lower than in *atrimanus*, the widest about  $\frac{1}{2}$  diameter of orbit (about  $\frac{1}{4}$  diameter of orbit in *atrimanus*). Coloration much as in *H. atrimanus*, but darker, and the fins without yellow; blackish olive above, dusky silvery below; top of head and snout black; spinous dorsal and the broad margins of soft dorsal and anal black; caudal dark, margined with black; pectorals very dark, black inside, the extreme lower rays light; a large jet-black blotch at base, on each side of pectorals, extending for about  $\frac{1}{2}$  the whole length of the fin; axil black. Closely related to *Hemicarax atrimanus*. Like it, it has a large jet-black area at axil and base of pectoral, and differs from it in the following characters: In having a shorter pectoral, shorter ventrals, profile of snout more rounded, a lower spinous dorsal, a shorter maxillary, a higher, shorter curve in lateral line, wider scutes, which are less sharply carinated, and darker fins. Panama. (*ζελορός*, an imitator.)

**Page 921.** *Caranx crysos* and *Caranx pisquetus* are probably distinct species, the former ranging from New York to Florida, the latter from the West Indies to Brazil.

In *Caranx pisquetus* the pectoral fins are very long, as in the Pacific species *Caranx caballus*, from which we can not separate it. The species need further study.

**Page 934.** After *Vomer setipinnis* add:

1329(a). **VOMER SPIXII** (Swainson).

Head  $2\frac{1}{2}$ ; depth  $1\frac{3}{4}$ ; D. VI-I, 22; A. I, 18; eye  $3\frac{1}{2}$  in head; maxillary  $2\frac{1}{2}$ ; snout  $1\frac{1}{2}$ ; caudal 1. Body very deep, in form much like *Selene arstedii*; profile very steep, almost vertical; snout slightly protruding. Mouth oblique, maxillary reaching to the vertical from front of eye; gill rakers 7 to 27, the longest a little more than  $\frac{1}{2}$  eye. Lateral line strongly arched in front, the arch  $1\frac{1}{2}$  the straight part; plates of lateral line little differentiated; pectoral falcate, as long or slightly longer than head; ventrals small, under base of pectorals. Color bluish above, sides silvery, fins except ventrals and anal dusky. Here described from specimens from Jamaica about 10 inches in length. These specimens are evidently different from the Northern *Vomer setipinnis* (= *Vomer browni*), the body in specimens of the same length being much deeper. It corresponds to the figure given by Agassiz of *Vomer browni*, this figure being the basis of *Vomer spixii* of Swainson. Probably all West Indian records of *Vomer setipinnis* belong to *Vomer spixii*. (Named for Jean Baptiste Spix, of Munich, naturalist and explorer.)

*Platyomus spixii*, SWAINSON, Class. Fishes, III, 250 and 406, 1839, Brazil; after AGASSIZ & SPIX.

*Vomer gabonensis*, GUICHENOT, Ann. Soc. Maine et Loire, 1865, 42, Gaboon.

**Page 938.** To the synonymy of *Chloroscombrus chrysurus* add:

*Seriola cosmopolita*, CUVIER, Règne Animal, Ed. 2, vol. II, 1829, Gorea; after *Scomber chloris*, BLOCH.

Add the following species:

1334(a). *CHLOROSCOMBRUS ECTENURUS*, Jordan & Osgood.

Head  $3\frac{3}{8}$ ; depth  $2\frac{3}{8}$ . D. VIII-I, 27; A. II-I, 26. Snout slightly shorter than eye, which is  $3\frac{1}{2}$  in head. Chord of curved part of lateral line  $1\frac{1}{2}$  in straight part. Depth of caudal peduncle 2 in its length, measuring from the base of the last dorsal ray to the base of the first caudal ray. Pectorals long and falcate, 3 in length; ventrals short,  $2\frac{3}{8}$  in head, extending beyond the vent, which is situated in a groove in which these fins fit. Depth of head equal to or slightly less than its length; maxillary reaching anterior edge of eye,  $2\frac{3}{8}$  in head. Lateral line unarmed; curve of ventral outline very slightly more pronounced than that of the dorsal; dorsal and anal fin sheaths well developed. Tips of upper spines and rays dusky; a black blotch at base of upper rays of caudal, and a black axillary and opercular spot. The species is closely related to *Chloroscombrus chrysurus*, the common species of the South Atlantic and Gulf States, which it evidently represents in the West Indies. The species *chrysurus* is deeper in every way, having a deeper body, a deeper head, and a deeper caudal peduncle. In *chrysurus* also the eye is larger, the mouth more nearly vertical, and the arch of the lateral line higher. When specimens of equal size from Florida and Havana are compared the characters are very evident. In 2 specimens, each  $7\frac{1}{2}$  inches in length, from Havana and Florida, respectively, the depth of the body of the one is contained  $1\frac{1}{2}$  times in that of the other, the depth of the head  $1\frac{1}{2}$ , the depth of the caudal peduncle  $1\frac{1}{2}$ , and the length of the eye  $1\frac{1}{2}$ . The names *chrysurus* (South Carolina), *latus* (Carolina), and *caribeus* (Texas) evidently all belong to the species of the United States coast. The type of *chloris* came from Acará, in Guinea, and *cosmopolita* of Cuvier was originally as a mere substitute for *chloris*. Until the African species can be examined, it is better not to use the name for either of the American forms. Probably *Chloroscombrus chloris*, when studied, will be found distinct from either. If not, that name would take the place of *ectenurus*. West Indies; known from Jamaica and Cuba. (*ἔκτενής*, extended; *οὐρά*, tail.)

*Chloroscombrus ectenurus*, JORDAN & OSGOOD, Proc. Ac. Nat. Sci. Phila. 1897, 101, Jamaica. (Coll. J. S. Roberts.)

The validity of *Chloroscombrus ectenurus* is still doubtful.

**Page 942.** After *Trachinotus falcatus* add:

1337(e). *TRACHINOTUS RHOMBOIDES* (Bloch).

Head  $3\frac{3}{8}$ ; depth  $1\frac{3}{8}$  in length; eye  $3\frac{1}{2}$  in head. D. VI-I, 20; A. II-I 18. Back much elevated, but not angulated at origin of soft dorsal; end of snout not vertical, curved; head slightly concave at occiput. Maxillary to below anterior margin of pupil; eye on level of lower edge of premaxillary and axil of pectoral. Origin of soft dorsal behind tip of pectoral,

its lobe much elongated, extending to middle of caudal; lobe of anal reaching to below base of caudal; caudal lobes equal,  $2\frac{1}{2}$  in body; pectoral rounded,  $1\frac{1}{2}$  in head; ventrals  $2\frac{1}{2}$  in head. Scales minute, large posteriorly near lateral line. Pale olive above, becoming silvery on belly; lobes of vertical fins dusky. This West Indian species is apparently different from the northern *Trachinotus falcatus* with which it has been confounded. *Trachinotus falcatus* seems to be confined to the coasts of the United States. In specimens of the same size the vertical fins are much higher in the West Indian species. ( $\rho\acute{o}\mu\beta\acute{o}\varsigma$ , rhomb;  $\epsilon\acute{\iota}\delta\omega\varsigma$ , resemblance.)

*Chaetodon rhomboides*, Bloch, Ichth., 1787, pl. 209, Martinique.

Page 945. After *Trachinotus paloma* insert:

428(a). ZALOCYS, Jordan & McGregor.

*Zalocys*, JORDAN & MCGREGOR, Rept. U. S. Fish Comm. 1898 (*stilbe*).

This genus is closely allied to *Hypodis*, Rafinesque (= *Lichia*, Cuvier), differing in the absence of a procurent spine before the dorsal, and in the cultrate thoracic region. From *Trachinotus* it is distinguished by the same characters and also by the lower forehead and nonfalcate dorsal and anal fins. *Hypodis* is scarcely different from *Trachinotus*, the only tangible characters being the larger teeth, the low dorsal, and the less elevated forehead. *Porthmeus*, Cuvier (= *Lichia amia* and *L. radijo*) is a well-defined genus, distinguished by the large mouth and projecting lower jaw. ( $\zeta\acute{\alpha}\lambda\eta$ , surge of the sea;  $\acute{\alpha}\nu\acute{\upsilon}\varsigma$ , swift.)

1844(a). ZALOCYS STILBE, Jordan & McGregor.

Head  $4\frac{1}{2}$ ; depth  $2\frac{1}{2}$ . D. VI-I, 26; A. II-I, 23. Body elliptical, deeper than in *Hypodis glaucus*; belly sharply compressed; ventral outline similar to that of dorsal; anterior profile of the head elevated and sharp, the eye being rather below than above its middle; eye 5 in head, with conspicuous adipose eyelid before and behind; posterior nostril much larger than anterior; vertically oblong maxillary broad, without supplemental bone, extending to pupil,  $2\frac{3}{8}$  in head. Mouth moderate, oblique; each jaw with bands of villiform teeth; similar teeth on vomer, palatines, and tongue. Preopercle very broad; cheek moderate; suborbital narrow; preorbital very narrow, 4 in eye. No pseudobranchiae. Gill rakers very long and slender, numerous. No procumbent spine before dorsal; spines low and separate, progressively higher; soft dorsal and anal each with a sheath of scales; first rays of dorsal very slightly elevated,  $2\frac{1}{2}$  in head; anal without distinct anterior lobe, longest ray  $2\frac{1}{2}$  in head; caudal peduncle long and slender; depth  $3\frac{1}{2}$  in head; length below  $2\frac{1}{2}$  in head; caudal fin widely forked; lobes long and slender, upper a little the longer, more than  $\frac{1}{2}$  longer than the head and  $2\frac{3}{8}$  in body; pectoral moderate,  $1\frac{1}{2}$  in head; ventrals very small,  $6\frac{1}{2}$  in head; snout  $3\frac{1}{2}$  in head; premaxillary protractile. Color dark steel blue or blackish above; lower parts soiled white; axil and base of pectoral within jet-black; dorsal and anal each with a narrow whitish edging; caudal black, each lobe with a narrow whitish edging within. Body covered with small smooth scales, much as in *Trachinotus*;

lateral line undulate, very slightly arched anteriorly. Clarion Island; 1 specimen, 16 inches in length, known. (*στῆλη*, shining.)

*Zalocys stilbe*, JORDAN & MCGREGOR, Rept. U. S. Fish Comm. 1898, pl. 5, Clarion Island, Revillagigedo Archipelago. (Type, No. 11996, L. S. Jr. U. M. Coll. R. C. McGregor.)

**Page 965.** *Rhombus*, *Palometa*, and *Poronotus* should probably stand as distinct genera. The species placed in *Rhombus* in the text would then stand as follows:

- 1363. RHOMBUS PARU (Linnæus).
- 1364. RHOMBUS XANTHURUS (Quoy & Gaimard).
- 1365. PALOMETA PALOMETA (Jordan & Bollman).
- 1366. PALOMETA MEDIA (Peters).
- 1367. PALOMETA SIMILLIMA (Ayres).
- 1368. PORONOTUS TRIACANTHUS (Peck).

The identity of the South Atlantic Coast *Rhombus alepilotus* with the West Indian *Rhombus paru* is very doubtful.

**Page 973.** The genus *Acrotus*, Bean, represents a family distinct from *Icosteida*.

Family CXXXVI(a). ACROTIDÆ.

Two additional specimens of *Acrotus willoughbyi* have lately come to light—the one from Port Townsend, the other from Monterey.

After *Acrotus willoughbyi* insert:

Family CXXXVI(b). ZAPRORIDÆ.

Body robust, moderately compressed, the back not elevated, the belly not carinate. Body covered with small adherent cycloid scales, which cover the membranes of all the fins except the distal third, as also the gill membranes, lower jaw, cheeks, opercles, and nuchal region. No lateral line; no spinules. Head short, the nape not elevated, the forehead broad and abruptly convex in profile; eye moderate, placed high; preopercle, parietal region, and region about eye with very large open mucous pores. No spines on head; edges of membrane bones of head covered with thick scaly skin. Mouth moderate, terminal, oblique, its cleft mainly anterior; upper jaw protractile, but not movable; maxillary rather narrow, simple; lower jaw very heavy, its thick lip projecting beyond upper jaw. Teeth alike in both jaws, rather strong, blunt, even, close set, forming a uniform cutting edge; no teeth on vomer, palatines, or tongue, the tongue very thick. Lower pharyngeals narrow, with bluntish teeth, those on the edge larger; upper pharyngeals rather large, with small, blunt, velvety teeth; no distinct tooth-like processes in the œsophagus; pseudobranchie present; gill rakers very slender and flexible, rather short; gills 4, a large slit behind the fourth; gill membranes separate, free from the isthmus; opercle adnate to shoulder girdle above its angle; coracoids not largely developed. Pectoral fin long, rounded, attached a little nearer

ventral than dorsal outline; ventrals wholly wanting. Dorsal fin beginning above gill opening, composed entirely of simple inarticulate rays or spines, these moderately flexible, attached to the membrane to their tips, and all except the first and last of about equal length. Caudal peduncle short and stout, not contracted, the large caudal subtruncate or rounded at tip, and without procurent rays; vent nearly median. Anal much shorter than dorsal, somewhat higher, and composed of soft rays, subequal in length. Skeleton rather limp and flexible, but much less so than in *Icosteus*.

## 445(a). ZAPRORA, Jordan.

*Zaprora*, JORDAN, Proc. Cal. Ac. Sci. 1890, 202 (*silenus*).

Characters of the genus included above.

This genus bears some resemblance to *Icichthys*, but differs in the stout caudal peduncle, absence of ventrals and lateral line, and in the form and structure of the head. Among the genera known to us it seems to come nearest to *Icichthys*, and it might be placed among the *Icosteidae* were it not for the presence of pharyngeal teeth. (ζά, an intensive particle; πρῶρα, prow.)

## 1372(a). ZAPRORA SILENUS, Jordan.

Head  $5\frac{1}{2}$  in length to base of caudal; depth  $4\frac{1}{2}$ . D. LVI; A. 27; P. 20 to 22; C. 22; scales about 200-85. Greatest thickness of body about  $\frac{2}{3}$  its depth; length of caudal peduncle  $1\frac{2}{3}$  in its least depth, which is  $1\frac{1}{10}$  in head. Eye  $5\frac{1}{2}$  in head; snout  $5\frac{1}{2}$ ; interorbital space 3; maxillary  $2\frac{1}{2}$ , ending under front of pupil; mandible  $2\frac{1}{2}$ , its depth  $4\frac{1}{2}$ ; teeth about  $\frac{1}{8}$  on each side; lips, snout, and bones about eye naked; rest of head covered with small scales. Lower jaw with a thick lip, slightly fringed on its edge, and with a mesial frenum; the rounded tip entering the profile when the mouth is closed. Three large pores on each ramus of mandible; behind these 3 others in a line on horizontal limb of preopercle; 3 on vertical limb; 2 close together in front of eye; 1 near the nostrils, so similar to them that there seem to be 3 nasal openings; 7 on suborbitals; 4 in 2 rows behind eye; 1 above eye, and before upper edge of preopercle; a horizontal row of 5 along temporal region, the last and largest of all in opercular flap above gill opening; 1 at vertex; 1 between vertex and eye, and 2 on each side of nape. Gill rakers 8+20, the longest  $\frac{1}{2}$  eye. No trace of lateral line. Scales small, resembling those of a salmon, covering the membranes of all the fins on the basal two-thirds. Pectoral as long as head, its base  $2\frac{1}{2}$  in head; longest dorsal spine  $1\frac{1}{2}$ ; caudal  $1\frac{1}{10}$ ; longest anal ray  $1\frac{1}{2}$ . Color in spirits uniform dusky, without markings on the body, the belly pale, and the side of the head irregularly blotched with lemon yellow, apparently bright in life, and brightest about the pores of the head. Coast of British Columbia; only the type, 29 inches long, known. (σειλένος, a drunken demigod, covered with slime, in allusion to the open mucous pores.)

*Zaprora silenus*, JORDAN, Proc. Cal. Ac. Sci. 1896, 203, pl. 20, Nanaimo, Vancouver Island. (Type in Provincial Museum at Victoria. Coll. H. T. Stainton.)

**Page 982.** Prof. Harrison Garman records *Elassoma zonatum* from Waccamaw River, Whitesville, North Carolina, and Little Pedee River, South Carolina. Vertebrae 29; scales 34 to 36; D. IV, 9; A. III, 5.

**Page 1019.** Under *kk* read: "gill membranes narrowly or broadly connected."

**Page 1047.** Before *Ulocentra* insert:

1436(a). COTTOGASTER CHENEYI, Evermann & Kendall.

Head 4; depth 6; eye 4 in head; snout 4; maxillary  $3\frac{1}{2}$ ; interorbital width  $5\frac{1}{2}$ . D. XI-12; A. II, 8; scales 7-56-6. Body rather stout, heavy forward, compressed behind; head heavy; mouth moderate, slightly oblique, lower jaw included, maxillary reaching front of pupil; premaxillaries protractile. Cheeks, opercles, breast, and nape entirely naked; scales of body large and strongly stenoid; lateral line complete, straight; median line of belly naked anteriorly, with ordinary scales posteriorly. Fins large; dorsals separated by a space equal to  $\frac{1}{2}$  diameter of eye; origin of spinous dorsal a little nearer origin of soft dorsal than tip of snout, its base about equal to length of head; longest dorsal spine  $2\frac{1}{2}$  in head, the outline of the fin gently and regularly rounded; soft dorsal higher than spinous portion, the second to tenth rays about equal in length, scarcely 2 in head, the first, eleventh, and twelfth rays but slightly shorter than the others; anal moderate, its origin under base of third dorsal ray, the spines slender, the second a little longer than the first, whose length is  $3\frac{1}{2}$  in head; longest anal rays about  $2\frac{1}{2}$  in head; caudal lunate, the lobes more produced and pointed than usual among darters; pectorals long and pointed, the middle rays longest, about  $1\frac{1}{2}$  in head, reaching tips of ventrals; ventrals well separated, not nearly reaching vent, the longest rays  $1\frac{1}{2}$  in head. Color in alcohol, back dark brownish, covered with irregular spots and blotches of darker; side with about 8 or 9 large dark spots lying on the lateral line; belly pale; top of head dark; snout black; lower jaw and throat dark; a broad black line downward from eye to throat; cheek and opercles rusty; spinous dorsal crossed by a median dark line; ventrals blue black; other fins pale, but dusted with rusty specks. An examination of the 14 cotypes shows some variation in the species. In 2 examples there is a well-developed frenum, rendering the premaxillaries nonprotractile, and in a third specimen the frenum is partially developed; in some individuals the origin of the spinous dorsal is exactly midway between the tip of snout and origin of soft dorsal. The females and immature males are less highly colored than the adult male described above. Length  $1\frac{1}{2}$  to  $2\frac{1}{2}$  inches. This species is most closely related to *Cottogaster shumardi*, from which it may be readily distinguished by the shorter snout, the naked cheeks and opercles, the smaller soft dorsal, the smaller anal, and the different coloration. Fifteen examples of this interesting darter were obtained July 18, 1894, by Messrs. Evermann & Bean in the Racket River near Norfolk, St. Lawrence County, New York. It did not seem to be very common, as only 15 examples resulted from numerous hauls of the collecting seine. (Named for Mr. A. Nelson Cheney, State fish-culturist of New York, in recognition

of his valuable contributions to our knowledge of the food and game fishes of that State.)

*Cottogaster cheneys*, EVERMANN & KENDALL, Bull. U. S. Fish Comm. 1897 (Feb. 9, 1898), 129, pl. 8, fig. 8, Racket River near Norfolk, New York. (Type, No. 48781. Coll. Evermann & Bean.)

Page 1049. After *Ulocentra gilberti* add:

1488(a). *ULOCENTRA MEADIE*, Jordan & Evermann, new species.

Head  $3\frac{1}{2}$ ; depth  $4\frac{1}{2}$ ; eye  $3\frac{1}{2}$  in head; snout  $3\frac{1}{2}$ ; interorbital 5. D. XII-12; A. II, 7; scales 7-48-6. Body rather heavy, somewhat fusiform; head large; snout blunt, decurved, profile rising abruptly to interorbital, thence nearly horizontal to origin of dorsal, from which it descends gently in a straight line to caudal peduncle; opercular spine small but sharp; mouth low, horizontal, rather large, the maxillary reaching vertical at front of orbit; premaxillaries protractile; branchiostegal membranes not connected, free from the isthmus; ventral fins close together, the space separating their bases about  $\frac{1}{2}$  diameter of orbit; fins all moderate; distance from tip of snout to origin of spinous dorsal 3 in body; spinous and soft dorsals close together, the space separating them about 2 in orbit; longest soft dorsal rays  $1\frac{1}{2}$  in head, about equaling those of anal; the two anal spines of about equal length, the first the stouter; pectorals long, longer than head, their tips passing those of ventrals but not reaching vent; ventrals short,  $1\frac{1}{2}$  in head; caudal slightly lunate when expanded. Scales rather large, strongly ctenoid; cheeks and breast naked; opercle scaled above, naked below; nape scaled; lateral line complete, straight; ventral line of body covered with ordinary adherent scales. Color in alcohol, yellowish or olivaceous above and on sides, the back with 6 dark saddle-like blotches, the first just anterior to origin of spinous dorsal, the second under the fifth and sixth spines, the third under the last two spines, the fourth under the sixth and seventh soft rays, the fifth just posterior to the last dorsal ray, and the sixth, which is quite small, upon the caudal peduncle at the base of the caudal fin; sides blotched with dark, 6 to 8 larger dark blotches along side just below lateral line, sometimes more or less continuous with the dark dorsal blotches; a dark blotch at base of middle caudal rays; belly pale; top of head dark; a dark spot at lower posterior angle of eye and a smaller one back of it on upper edge of opercle; a dark band downward from eye; opercle dark; upper lip dark, interrupted by a light line at the symphysis; spinous dorsal pale, with a broad dark band through its lower third; soft dorsal crossed by 3 or 4 irregular lines of dark specks; caudal with about 4 broad dark cross bars; other fins pale. Length 2 inches. This species somewhat resembles *U. gilberti*, but differs from it in the larger head, stouter body, larger scales, naked cheeks, larger mouth, and in other respects. Known only from Indian Creek, basin of Powell River, east Tennessee, where 3 examples were collected October 17, 1893. (Named for Mrs. Meadie Hawkins Evermann.)

*Ulocentra meadia*, JORDAN & EVERMANN, new species, Indian Creek, Cumberland Gap, Tennessee. (Type, No. 48903. Coll. Dr. R. R. Gurley.)



**Page 1051.** To the synonymy of *Ulocentra simotera* add:

*Etheostoma duryi*, HENSHELL, Journ. Clin't Soc. Nat. Hist., April, 1889, 32, small tributary of Tennessee River at Whiteside, Tennessee. (Type in Mus. Clin't Soc. Nat. Hist. Coll. Charles Dury.)

**Page 1089.** To the synonymy of *Etheostoma caruleum* add:

*Etheostoma formosa*, HENSHELL, Journ. Clin't Soc. Nat. Hist., April, 1889, 32, small tributary of Tennessee River at Whiteside, Tennessee. (Type in Mus. Clin't Soc. Nat. Hist. Coll. Charles Dury.)

**Page 1109.** Add:

1501(a). APOGON ATRICAUDUS, Jordan & McGregor.

Head  $2\frac{1}{2}$ ; depth 3. D. VI-I, 9; A. II, 8; scales largely ctenoid; eye  $3\frac{1}{2}$  in head; second dorsal spine stoutest, about 2 in head; gill rakers 17, moderate. Body similar in shape to *A. retrosellus*. Jaws reaching to posterior border of eye,  $1\frac{1}{2}$  in head. Pectoral reaching to opposite front of anal,  $1\frac{1}{2}$  in head. Color rosy, darkened with dusky points; more or less olivaceous above; head and throat verging on orange; first dorsal black; second dorsal rosy; caudal dusky, more or less flushed with rosy, other fins paler; no black spot on head or on base of caudal, there being no definite markings anywhere except the dusky red of the tail. West coast of Mexico. Numerous specimens collected at San Benedicto, Socorro, and Clarion islands. Usual length 3 to 4 inches. (ater, black; cauda, tail.)

*Apogon atricaudus*, JORDAN & MCGREGOR, Rept. U. S. Fish Comm. 1898, Socorro, Clarion and San Benedicto islands. (Coll. R. C. McGregor.)

**Page 1125.** *Centropomus affinis* can not be separated from *C. ensiferus*.

**Page 1148.** To the synonymy of *Epinephelus* add:

*Phrynotitan*, GILL, Stand. Nat. Hist., III, 255, 1885 (*Batrachus gigas*).

**Page 1150.** In the key under *dd*, read: Lower jaw strongly projecting.

**Page 1156.** Add:

1551(a). EPINEPHELUS NIPHOBLES, Gilbert & Starks.

Head  $2\frac{1}{2}$  in body; depth  $2\frac{1}{2}$ . D. XI, 14; A. III, 9; scales 16-116-40; eye 5 in head; maxillary 2; third dorsal spine  $2\frac{1}{2}$ ; middle dorsal rays  $2\frac{1}{2}$ ; highest anal rays 2; third anal spine  $3\frac{1}{2}$ ; pectoral  $1\frac{1}{2}$ ; ventrals  $1\frac{1}{2}$ ; caudal  $1\frac{1}{2}$ . Form rather robust, moderately compressed; dorsal outline uniformly curved from tip of snout to caudal peduncle; mouth large, the maxillary reaching to below posterior orbital rim; lower jaw strongly projecting; teeth conical and sharp, in 1 or 2 bands at sides of jaws, 3 or 4 in front; upper jaw with a rather strong canine on each side of front; snout longer than eye; nostrils close together, the posterior one the larger, a little in front of the vertical from front of eye, the anterior in a short, wide tube with a flap behind; vertical and horizontal limbs of preopercle meeting at right angles, its edge with blunt serrae, those at angle enlarged; opercle with 3 flat spines before the flap; gill rakers moderate, nearly  $\frac{1}{2}$  eye, 8+16 in number. Top of head, orbitals, maxillary, and mandible, naked; fine scales on cheeks and opercles; scales on body ctenoid;

fins without scales. Dorsal beginning a little in front of the vertical from pectoral base, the third spine a little the highest, but the ones behind it not much shortened; soft dorsal higher than spinous, its outline rounded; pectoral rounded behind, reaching to below the base of eighth dorsal spine; third anal spine the longest, not nearly so long as the soft rays, the anal fin similar in shape to the soft dorsal; ventrals reaching past vent, scarcely to front of anal, their ends rounded, as are all the fins; caudal broadly rounded. Color in spirits brownish red, sides with clear-cut, distinct, white spots about as large as pupil, about 6 at base of dorsal, 6 or 7 along lateral line, following its arch, a horizontal series of 4 extending back from opercular flap, about 3 from base of pectoral following curve of ventral outline, 2 at base of anal, 1 behind lower edge of caudal peduncle and 1 above anus; a well-marked streak above maxillary following its outline; lips colored like rest of head; dorsal dusky, with vague white spots; ventrals and anal nearly black, with a reddish tinge; anal with a narrow white border below; pectoral and caudal uniform yellowish. Magdalena Bay, Lower California; only the type, 6 inches long known. (*νιφοβής*, snowed over, from the white spots.)

*Epinephelus niphobles*, GILBERT & STARKS, Proc. U. S. Nat. Mus. 1896, 442, Magdalena Bay, Lower California. (Type, No. 47532. Coll. *Albatross*.)

**Page 1164.** Species 1558 should probably be called *Alphetses chloropterus* (Cuvier & Valenciennes). The name *afer*, given to a specimen from Guinea, may belong to some other species.

**Page 1168.** Add:

1560(a). *DERMATOLEPIS ZANCLUS*, Evermann & Kendall.

Head  $2\frac{3}{4}$ ; depth  $2\frac{1}{6}$ ; eye 8 in head; snout  $3\frac{1}{4}$ ; maxillary 3; mandible 2. D. XI, 19; A. III, 10; scales difficult to count, but about 30-130-35, those above lateral line counted obliquely backward and downward from origin of dorsal, those below from origin of anal upward and forward to lateral line. Branchiostegals 8; gill rakers 8 + 12, short and stout, the longest  $1\frac{1}{2}$  in orbit. Body stout, compressed, oblong-elliptical, the dorsal and ventral outlines about equally curved; head moderate, the profile rising from tip of snout to origin of dorsal fin, thence descending in a regular, gentle curve to caudal peduncle; a depression above nostrils and a slight one on nape; interorbital very narrow, equal to orbit; mouth moderate, somewhat oblique; premaxillaries protractile; maxillary broad at tip, reaching vertical at posterior edge of the pupil; supplemental bone well developed; lower anterior edge of maxillary covered by the broad dermal flap of the premaxillary; eye small, high up; nostrils close together and close to eye, the anterior small and round, the posterior oblong-oval, much larger than the other. Small cardiform teeth on each jaw, those in front movable, scarcely canine-like; similar teeth on vomer and a long, narrow band on each palatine. Preopercle coarsely serrate, the serrae short and blunt, more or less obscured by the skin; opercle with a broad dermal border, somewhat produced at lower angle. Fins all large; origin of dorsal slightly in advance of base of pectoral, its distance from tip of snout equal

to length of head; third dorsal spine longest, its length about  $2\frac{1}{2}$  in head or  $2\frac{1}{2}$  times length of first ray; interspinal membranes of the spinous dorsal deeply incised, the anterior portion of each somewhat produced beyond its spine; soft dorsal high, the middle rays longest,  $1\frac{1}{2}$  in head, the anterior portion of the fin gently convex, the posterior slightly concave; pectoral short, broad, and rounded, barely reaching origin of anal, the length  $1\frac{1}{2}$  in head; ventral pointed, the second and third rays longest,  $1\frac{1}{2}$  in pectoral, the fin somewhat falcate; anal fin strongly falcate, the fourth and fifth rays longest, longer than pectoral,  $1\frac{1}{2}$  in head,  $2\frac{1}{2}$  times length of last anal ray; second anal spine short,  $5\frac{1}{2}$  in head; caudal shallowly lunate, the lobes  $1\frac{1}{2}$  in head. Scales small, smooth, and thin, closely but irregularly imbricated; nape, opercles, and cheeks scaled, snout and lower jaw naked; bases of all the fins except the ventrals densely scaled; lateral line beginning at upper angle of opercle, gently arched above pectoral fin, following approximately the curvature of the back and on median line of caudal peduncle. General color of body in life brown, with large, irregular blotches of dirty white on back and upper part of sides, these blotches with small rusty spots; lower part of sides, belly, and caudal peduncle with irregular whitish spots; belly brassy brown; snout and nape with numerous small, round dark spots; cheek with large blotches of whitish overlaid with black and brassy spots; lips whitish, with dark spots; spinous dorsal blotched with white, olivaceous and black; soft dorsal brown, with numerous white spots and a few black ones, the posterior rays tipped with white and orange; anal olivaceous, with irregular white spots, greenish at edge, the produced rays black toward distal ends; pectoral dark olivaceous, with greenish white splotches, the edge yellowish; ventral rays greenish white, the membranes black; inside of mouth white; eye brown. Related to *D. inermis* (Cuvier & Valenciennes), but differing notably from that species in the shorter, stouter gill rakers, the emarginate caudal, the shorter anal spines, and the strongly falcate anal fin. Length 20 inches. Key West; only the type known. (*ζάγκλον*, a scythe or sickle, from the falcate anal fin.)

*Dermatolepis zancus*, EVERMANN & KENDALL, Bull. U. S. Fish Comm. 1897 (Feb. 9, 1898), 129, pl. 8, fig. 9, Key West, Florida. (Type, No. 48843. Coll. Drs. Evermann & Kendall.)

Page 1186. Add:

1576(a). MYCTEROPERCA HOPKINSI, Jordan & Rutter.

Head  $2\frac{1}{2}$ ; depth  $4\frac{1}{2}$ . D. XI, 15; A. III, 11; scales about 125; eye 6 in head,  $1\frac{1}{2}$  in snout. Body long, not much compressed; angle of preopercle sharply serrate; gill rakers 6 + 9, counting rudiments; nostrils close together, the posterior larger, with a horizontal septum across base; profile concave above nostrils; maxillary nearly to posterior margin of eye,  $2\frac{1}{2}$  in head; lower jaw projecting; 2 anterior canines of upper jaw very strong; third and fourth dorsal spines longest; posterior portion of anal truncate; caudal concave. Pectorals 2, ventrals  $2\frac{1}{2}$ , and caudal  $1\frac{1}{2}$  in head. Color of alcoholic specimen nearly uniform brownish, side of jaws paler; soft dorsal, anal, ventrals, and caudal with a narrow pale edging,

these fins otherwise brownish olive, with a subterminal band of black; pectorals pale, darker in middle. Allied to *Mycteroperca calliura*, differing in having fewer gill rakers, more slender body, smaller scales, and a less lunate caudal. Jamaica; only 1 specimen, 6 inches long, known. (Named for Timothy Hopkins.)

*Mycteroperca hopkinsi*, JORDAN & RUTTER, Proc. Ac. Nat. Sci. Phila. 1897, 105, Jamaica. (Type, No. 5073, L. S. Jr. Univ. Mus. Coll. J. S. Roberts.)

Page 1187. Insert:

1576(a). MYCTEROPERCA BOULENGERI, Jordan & Starks.

Head  $2\frac{1}{2}$  in length; depth  $2\frac{3}{8}$ . D. XI, 14 or 15; A. III, 9 or 10; scales about 90, 20 above and 42 below; snout  $3\frac{1}{2}$  in head; maxillary  $2\frac{1}{2}$ ; eye  $5\frac{1}{2}$ ; pectoral  $1\frac{1}{4}$ ; ventral  $1\frac{1}{6}$ ; longest anal ray  $1\frac{3}{8}$ ; caudal  $1\frac{3}{8}$ ; longest dorsal spine  $2\frac{1}{2}$ ; gill rakers short, about 6 + 17, the longest about  $\frac{3}{8}$  eye; longest dorsal ray 2 in head. Body short and deep, compressed; head moderate, compressed, its profile not steep, nearly straight, a depression before eye. The supraoccipital and temporal crests are high, the supraoccipital crest extending to the posterior margin of orbit; the temporal crests are parallel to each other, and extending to pupil; interorbital space concave. Upper canines moderate, the lower quite small. Nostrils small, well separated, the anterior slightly larger. Lower jaw very strongly projecting; maxillary reaching opposite posterior edge of pupil. Preopercle slightly notched, the angle slightly salient, with enlarged teeth. Dorsal not deeply notched, the fourth spine not much elevated; second dorsal high, not long, its angle not rounded; caudal scarcely lunate, the upper lobe long, the lower truncate; anal very high, strongly elevated, its posterior border incised, the anterior rounded; pectoral and ventral moderate. Scales smoothish, not very small. Color olive gray, covered everywhere with oblong irregular markings of black, between which the ground color forms rivulations; gray lines radiating from the eye; a black blotch below maxillary; pectoral olive yellow; other fins blackish, clouded with pale; first dorsal with faint small black spots. Mazatlan, Mexico. (Named for George Albert Boulenger, ichthyologist of the British Museum, in recognition of his epoch marking work on the Percoid fishes.)

*Mycteroperca boulengeri*, JORDAN & STARKS, Fishes Sinaloa, 445, pl. 38, 1895. Mexico. Sinaloa, Mexico. (Type, No. 1621, L. S. Jr. Univ. Mus. Coll. Hopkins Exped. Sinaloa.)

Page 1235. The original type of *Lobotes* is *surinamensis*, not *erate*.

*Lobotes erate* is a species distinct from *L. surinamensis*, inhabiting the coasts of India and China. *Lobotes farkhari* and *L. incurvus* are probably identical with *L. erate*, and all 3 should be erased from the synonymy of *L. surinamensis*.

Page 1236. After *Lobotes surinamensis* add:

The *Lobotes* of the Pacific coast of Central America is distinguished from the other known species, *L. surinamensis* and *L. erate*, by the small

size of the preopercular serrations, those at the angle not elongated and spine-like, even in the young. The following description is furnished by Dr. Gilbert:

## 1023(a). LOBOTES PACIFICUS, Gilbert, new species.

(BERRUGATE.)

Head  $2\frac{1}{2}$  in length; depth  $2\frac{1}{2}$  to  $2\frac{1}{10}$  (to base of caudal rays); depth of caudal peduncle  $2\frac{1}{2}$  in head. D. XII, 15; A. III, 11; pectoral 15. Scales 11-46 (+6 on base of caudal)-18; vertebrae 12+12; Br. 6. Body more elongated than *L. surinamensis*, agreeing in this respect with *L. erate*, the depth less than  $\frac{1}{2}$  the length. Upper profile deeply concave at occiput, thence strongly convex to front of dorsal; head shorter and narrower than in *L. surinamensis*, the interorbital width but slightly longer than snout,  $3\frac{1}{10}$  to 4 in head ( $3\frac{1}{2}$  to  $3\frac{3}{4}$  in head in *L. surinamensis*). Eye small,  $6\frac{3}{4}$  to  $7\frac{1}{4}$  in. head, 2 or  $2\frac{1}{10}$  in interorbital width. Mandible strongly protruding, but without symphyseal knob; maxillary narrow, not concealed in closed mouth, its tip reaching vertical from middle of pupil,  $2\frac{1}{4}$  to  $2\frac{3}{10}$  in head. Upper jaw with a moderate villiform band of teeth, in front of which is a single series of conical, close-set canines; lower jaw with a single series, similar to outer series of upper jaw, and behind them a very narrow band of villiform teeth, which grow slightly larger toward symphysis; palate toothless. Posterior margin of preopercle vertical, the angle protruding but little in the young. In 5 young examples, 7 to 11 inches long, the preopercular teeth are fine, acute, short, and inconspicuous, about as in species of *Pomadasis*. They increase but little in size toward the angle, where they are never spine-like; on lower limb they are perceptible only in the immediate vicinity of the angle, the remainder of the horizontal limb being entire. In the adult the vertical limb is finely and evenly toothed, the angle and lower limb slightly roughened or entire; opercle with 2 short spinous points, behind the lower of which a narrow tongue-shaped process of the subopercle extends to near the edge of opercular membrane; humeral process very weakly toothed, contrasting with the strong serrate condition in *L. surinamensis*. Gill rakers short,  $2\frac{1}{2}$  in eye in young, comparatively shorter in adults, 6 on vertical limb, all but one of which are broad, firmly fixed tubercles, 14 on horizontal limb, the anterior 2 or 3 tubercular. Spinous dorsal low, with gently rounded outline; notch between dorsals shallow, the eleventh spine  $\frac{2}{3}$  the length of the longest, which is contained 2 to  $2\frac{1}{2}$  times in head in the young, 3 times in adults; when declined the spines are partially received within a scaly groove; soft dorsal, anal, and caudal with dorsal portions densely scaled and with series of scales running up on membrane to beyond middle of fin; soft dorsal and anal of equal height, forming bluntly rounded lobes, the longest rays of which are about  $\frac{1}{2}$  head in adults,  $1\frac{1}{2}$  to  $1\frac{3}{4}$  in young; third anal spine about  $\frac{1}{2}$  length of longest ray; pectorals shorter than ventrals, 2 to  $2\frac{1}{2}$  in head; ventrals  $1\frac{1}{2}$  in head in young, shorter in adults. Scales less strongly ctenoid than in *L. surinamensis*; tubes of lateral line mostly simple, occasionally with 1 to 3 branches. Color grayish or brownish, with plumbeous or silvery reflections. The youngest examples show faintly the dark streaks so con-

spicuous in young of *L. surinamensis*, viz, a pair running backward from interorbital space; a pair from upper posterior border of eye converging toward front of dorsal, and a broader band from eye downward and backward across cheek; soft dorsal, anal, and caudal uniform blackish, or the caudal with an ill-defined lighter edge; pectorals translucent; ventrals blackish. Abundant at Panama, where it is known as *Berrugate*.

*Lobotes auctororum*, STEINDACHNER, Ichth. Beltr., IV, 6, 1875; not of GÜNTHER.

*Lobotes surinamensis*, JORDAN & GILBERT, Bull. U. S. Fish Comm., 11, 1882, 110; GILBERT, l. c., 112; JORDAN, Proc. U. S. Nat. Mus. 1885, 378; not *L. surinamensis* of BLOCH.

*Lobotes pacificus*, GILBERT, Fishes of Panama, 1898 MS., Panama. (Type, No. 5883, L. S. Jr. Univ. Mus. Coll. Gilbert.)

**Page 1238.** After *Priacanthus cruentatus* add the following:

1625(a). **PRIACANTHUS CAROLINUS**, Lesson.

This species is very close to *Priacanthus cruentatus*, distinguished by the larger spine on preopercle, which reaches the edges of the opercle and is  $2\frac{1}{2}$  in eye; that of *P. cruentatus* not reaching opercle and measuring 4 in eye, its edge less rough. Body a little deeper than that of *P. cruentatus*; depth of the latter 3 in the length. In *P. carolinus* the depth is  $2\frac{3}{4}$  in the length; caudal truncate. In color and general appearance the 2 species are similar. The distinctness of this species from *P. cruentatus* is very doubtful. Abundant at Clarion Island, where it was taken by Mr. R. C. McGregor. (*carolinus*, from Caroline Islands.)

*Priacanthus carolinus*, LESSON, Voyage Coquille, Poiss., 204, 1826, Caroline Islands.

*Priacanthus schlegelii*, HILGENDORF, Sitzgber. Ges. Naturf. 1879, 79, Japan.

**Page 1262.** In the first line of the description of *Neomænis vivanus* read: Head  $2\frac{3}{4}$  to  $2\frac{1}{2}$ ; depth  $2\frac{1}{2}$  to 3. D. X, 14; A. III, 8 or 9; eye  $4\frac{1}{2}$  in head; scales (7) 8-72-17, 50 pores.

**Page 1264.** After *Neomænis vivanus* add:

1639(a). **NEOMÆNIS HASTINGSI**, Bean.

(Bermuda Silk Snapper.)

Head 3; depth 3; least depth of caudal peduncle 9 in length of type to caudal base. D. X, 14; A. III, 8; V. I, 5; P. 16; scales 8 or 9-65-17. Maxillary reaching scarcely past front of eye, 3 in head. Vomerine teeth in an arrow-shaped patch, with a backward extension which is fully  $\frac{1}{2}$  as long as the eye; canines in upper jaw very feeble; 2 or 3 posterior teeth of mandible are weak canines; 7 rows of scales on cheeks, 9 rows on gill cover. Least interorbital width equal to eye, which is  $1\frac{1}{2}$  in snout and 4 in head. Gill rakers 7 + 9, the one in the angle conspicuously longest, about 2 in eye. First dorsal spine 7 in head; fifth and longest spine about 3 in head; last dorsal spine equal to eye in length; longest ray of soft dorsal equal to maxillary, or 3 in head; first anal spine 8, the second and third about 4 in head, the second slightly longer than third; anal base nearly  $2\frac{1}{2}$  in head; third and longest anal ray about equal to anal base; pectoral extending to vent; ventral not reaching vent by a space  $\frac{1}{2}$  as long as the eye. Colors in life, ground color vermilion, the upper parts over-

laid with coppery brown, lower parts vermilion; 4 or 5 narrow golden stripes below lateral line; caudal dark brown with a narrow black margin; anal dusky, the spines and the membranes of last 2 rays pale; a narrow black blotch at pectoral base; ventral pale, somewhat mingled with dusky; membranes of spinous and soft dorsal uniformly dark; snout copper color; eye lemon yellow; pupil blue black; many scales, especially on front of body, with a minute brown dot at base; brownish spots on scales forming many oblique streaks above lateral line. Some living examples show a faint dark lateral blotch much like that of *N. synagris*, and similarly placed. In spirits the body is pink with the upper parts brownish; the dusky color remains on the anal and the black blotch at base of pectoral; black margin of caudal becoming merged with the general dark color of the fin. (Bean.) Most closely related to *N. vivanus*. Length of type 11½ inches. Bermuda, where numerous specimens were obtained in 1897. (Named for General Russell Hastings of Soney, Bermuda.)

*Neomænis hastingsi*, BEAN, Bull. Amer. Mus. Nat. Hist., x, Article III, 45, 1898, Bermuda.

Page 1290. Under *g* read: Anal fin short, its rays III, 7 to III, 13.

Page 1413. In first line of description of *Cynoscion phorocephalus* for "A. III, 10" read "A. II, 10."

Page 1416. In last line of description of *Sagenichthys ancyodon* for "companion" read "comparison."

Page 1605. Instead of *Chlorichthys* read:

#### 640. THALASSOMA, Swainson.

*Thalassoma*, SWAINSON, Nat. Hist. Class. Fishes, II, 224, 1839 (*purpureus*).  
*Julis*, GÜNTHER, Cat., IV, 179, 1862; not of CUVIER & VALENCIENNES.

The species of *Thalassoma* (*pavo*, *unimaculatus*, *bifasciatus*) examined have 3 anal spines, as is the case with the American species referred to *Chlorichthys*. The first spine, small and hidden in the skin, is easily overlooked. There is therefore no distinction between *Thalassoma* and *Chlorichthys*, and all the American species must be referred to the former genus.

The species will stand as follows:

- 2014. THALASSOMA LUCASANUM (Gill).
- 2015. THALASSOMA SOCORROENSE, Gilbert.
- 2016. THALASSOMA NITIDUM (Günther).
- 2017. THALASSOMA NITIDISSIMUM (Goode).
- 2018. THALASSOMA STEINDACHNERI (Jordan).
- 2019. THALASSOMA BIFASCIATUM (Bloch).
- 2020. THALASSOMA GRAMMATICUM, Gilbert.
- 2021. THALASSOMA VIRENS, Gilbert.

Page 1670. In *Pomacanthus* (*P. paru*, species examined by Mr. E. C. Starks) and *Chatodon* the air bladder is wholly contained in the body cavity, while in *Holacanthus* and *Angelichthys* (*A. ciliaris* species examined) it is posteriorly separated from the body cavity. The 2 latter genera con-

stitute the subfamily *Holacanthinae*, distinct alike from *Chatodontinae* and *Pomacanthinae*.

**Page 1717.** *Ceratacanthus*, including *Osbeckia*, should stand as a valid genus distinguished from *Alutera* by the convex or lanceolate caudal. The species will then stand as follows:

2135. *CERATACANTHUS SCHÖEPLII* (Walbaum).

2136. *CERATACANTHUS PUNCTATUS* (Agassiz).

2137. *CERATACANTHUS SCRIPTUS* (Osbeck).

2138. *ALUTERA MONOCEROS* (Osbeck).

**Page 1741**, line 17, read: Swainson takes "*κάνθα*" to mean spine, not "*ἀκάνθα*," which is the correct word for spine. There is no classical warrant for *Cantherines* and *Cunthigaster*, unless derived from *κάνθος*, the uss.

**Page 1776.** In sixth line from bottom read "increased" for "self."

**Page 1786.** Note on *Sebastodes rufus*:

This species is ovate in form, like *S. ovalis*, from which it differs in color and form of mouth and head. Its depth is  $2\frac{3}{4}$  in length, not  $3\frac{1}{2}$ , as stated (through misprint) by Dr. Eigenmann. A fine specimen before us was taken by Dr. Gilbert off San Diego.

**Page 1790.** The type number of *Sebastodes hopkinsi* is 2282, not 2286.

**Page 1795.** The subgenus *Zalopyr* does not include *Sebastodes atrovirens* nor *S. atrovirens*. These 2 species belong in the subgenus *Rosicola*.

**Page 1799.** In first line under *Sebastodes crameri* for "P. 19<sub>10</sub>" read "P. 19."

**Page 1805.** After line 4 add: (*intro*, within; *niger*, black.)

**Page 1815.** The type of *Sebastodes zacentrus* came from *Albatross Station* 2946, not 2996.

**Page 1829.** After line 2 insert Subgenus *Sebastosomus*.

**Page 1831.** Specimens of *Sebastodes taczanowskii* were obtained in 1896 by the *Albatross* at the Kuril Islands, and this species should therefore go in the regular text.

**Page 1832.** After line 15 insert Subgenus *Sebastomus*.

**Page 1833.** Before *Sebastodes matsubaræ* (Hilgendorf) insert Subgenus *Zalopyr*.

**Page 1833.** In fifth line from bottom insert Subgenus *Pteropodus* after "*nebulosus*."

**Page 1836.** In key at bottom of page, to *a* add: mouth plumbeous within. To *aa* add: mouth black within.

**Page 1837**, line 2, for "*Cardonniera*" read "*Cardouniera*." For "*Scorfanudi Funal*" read "*Scorfana di Funal*."

**Page 1840.** Above "*a*. Breast scaly," insert *Parascorpaena* (*παρά*, near; to *Scorpana*). Before 2236. *Scorpana agassizii*, Goode & Bean, insert Subgenus *Parascorpana*, Bleeker.

**Page 1850.** In last line of synonymy of *Scorpana mystes* for 1501 read 1601.



Page 1854. After *Scorpana inermis* add:

2247(a). SCORPENA NEMATOPHTHALMUS (Günther).

Head 3 in total length; depth 3½; eye 4 in head; snout rather less than 4. D. XII, 10; A. III, 5; scales 40 or 41. Dorsal outline much arched at greatest depth of body. Eye placed high, entering upper outline of head. Intermaxillaries styliform, armed, like the dentaries, with a rather narrow band of villiform teeth; band of vomerine teeth angularly bent, produced forward at the angle; maxillaries styliform at superior extremity, moderately dilated at the lower. Head scaled to posterior angle of orbit above and to the preorbital and angle of mouth laterally. Spines on head very prominent and acute in the young, more obtuse in older examples; 2 turbinal spines; on each side of the occiput a series of 5 spines between orbit and nape; 2 between eye and scapula; preorbital armed with 2 strong, recurved spines at the inferior margin; 3 spines on interorbital ridge; preopercular margin rounded, with 4 spines, the uppermost and strongest opposite end of interorbital ridge; opercle with 2 flat spines; a pair of spines at throat. The only skinny appendage is a long, slender, tapering filament above posterior angle of orbit. Origin of dorsal immediately behind vertical from suprascapula, its distance from occiput equaling length of first spine, which is about ½ length of second; third and fourth spines longest, 2¾ in head; the following spines gradually decreasing to the eleventh, which equals the first; twelfth spine much longer, apparently belonging to the soft portion, which is supported by it; margin of soft portion rounded, very little higher than the spinous, posteriorly fixed to the back of the tail by a membrane; caudal subtruncated; origin of anal somewhat behind that of soft dorsal, its second spine strong, rather longer than the third dorsal, and with a longitudinal groove; pectoral reaching anal; ventral reaching vent. Scales of moderate size, rather irregularly arranged. Color probably uniform red. Supposed to be from the West Indies. (Günther.) Only the type known. (*νημα*, thread. *ὄφθαλμός*, eye.)

*Sebastes nematophthalmus*, GÜNTHER, Cat., II, 99, 1860, West Indies; the exact locality unknown. (Coll. Sir R. Schomburgk.)

Page 1862. *Anoplopoma fimbria* is occasionally taken off Santa Catalina in deep water. A specimen was seen by us at Redondo Beach.

Page 1866, line 7 from bottom, read "always," not "usually."

Page 1867. In the key, *b* should read as follows:

- b*. Fourth line of pores forking in advance of base of ventrals, the lower branch running to base of ventral fin, the upper to middle of ventral. OCTOGRAMMUS, 2259.

In the footnote for "Keinosuke Otaki" read "Keinoske Otaki."

Page 1879, line 27, for "jointed" read "joined." Line 37, after "hypercoracoid" add "and hypocoracoid."

Page 1880. In the key read:

- m*. Lateral line armed with a series of bony plates; preopercular antler-like processes usually numerous.

**Page 1881.** The key should be modified to read:

q. Interorbital space deeply concave or grooved. Head with cirri (in *lateralis*), or without (in *asperulus*). ARTEDIUS, 712.

A better distinction between *Artedius* and *Axyrias* is found in the presence of patches of etenoid scales on the head in the latter.

**Page 1884.** In line 19, for *Hexagrammidæ* read *Zantolepidinæ*.

**Page 1898.** The type number of *Icelinus strabo* is 5045, not 5451.

**Page 1902.** *Artedius asperulus* is better separated from *A. lateralis* by the coalescence of the bands of scales behind the dorsal and their continuance upon the caudal peduncle. In the description of the genus *Artedius* the bands of scales are said not to meet behind the dorsal. This applies to *A. lateralis* only.

**Page 1902.** To the description of *Artedius* add: No patches of etenoid scales on the head.

**Page 1903.** To the description of *Artedius asperulus* add: Head without cirri.

**Page 1906.** The type of *Artediellus atlanticus* is No. 448 L. S. Jr. Univ. Mus.

**Page 1940.** In the key, under *a*, add: Preopercular spine with 3 hooks above. Under *aa*, add: Preopercular spine with 6 or 7 hooks above.

**Page 1958.** *Cottus aleuticus* extends southward in the Coast Range to Monterey.

**Page 1964.** In key under *h* for "anal" read "axil."

**Page 2000.** After *Porocottus tentaculatus* add:

2371(a). POROCOTTUS BRADFORDI, Rutter, new species.

Head 3; depth 3¼ to 4¼; eye 4. D. IX, 15\* or 16; A. 11 to 13; P. 13 or 14; B. 6. Head broad, somewhat depressed; bones of head cavernous; lower jaw included, maxillary to below middle or hinder edge of pupil, 2½ in head; teeth in jaws and in a narrow crescent on vomer; eye equal to snout; nasal spines blunt, covered by the skin; no ocular, opercular, nor suprascapular spines; preopercular spines 3, upper slender, curved inward, lower straight, pointing downward, middle 1 short and blunt, a mere tubercle; a very slight tubercle represents the fourth spine belonging to the genus; no slit behind last gill; 3 pairs of cirri on top of head, 1 above eye, multifid, another at occiput, single or bifid, the other between them, trifid to multifid; a minute barbel on tip of maxillary; whole top and side of head, lower jaw, and edge of preopercle thickly covered with pores; a double series of pores, 34 to 36 each, along lateral line with many accessory pores, these arranged in groups of 1 to 5 between the pairs of the

	D.				A.		
	IX, 15	IX, 16	IX, 17	VIII, 17	11	12	13
Fin rays.....	11	12	1	1	1	22	2
Number specimens.....							

lateral line; nostrils with short tubes; dorsals united at base, the spines with short filaments, middle spines 3 in head, middle rays of soft dorsal  $2\frac{1}{2}$  in head; caudal and ventrals  $1\frac{1}{2}$  in head, ventrals usually reaching vent or anal, but sometimes falling short of each; pectoral  $1\frac{1}{2}$  in head, reaching to or beyond anal. Color dusky, below colorless, a pale bar across occiput (often absent), another between dorsals, 2 across body under soft dorsal and another behind soft dorsal; sometimes the pale color predominates and the dusky portion is left as 4 bars; sometimes plain dusky without cross bars; spinous dorsal dusky with 3 or 4 colorless spaces on the web; other fins barred with series of dusky blotches, ventrals sometimes colorless; 5 to 8 oval white spots behind pectoral, sometimes obscure; males with inner ray or rays of ventral tuberculate or serrate. This species differs from *Porocottus sellaris* in the presence of cirri on top of head; it has more numerous fin rays and more cirri on head than *Porocottus quadrifilis*. This species is the most common fish in the rock pools at Karluk, where many specimens were taken. These are in the U. S. National Museum, in the collection of the U. S. Fish Commission, and in that of Leland Stanford Junior University. (Named for Mr. William B. Bradford, secretary of the Alaska Packers' Association, from whom the collector received many favors.)

Page 2015. Before *Oxycottus* insert the following:

745(a). SIGMISTES, Rutter, new genus.

*Sigmistes*, RUTTER, MS., new genus (*caulias*).

This genus differs from *Oxycottus*, to which it is most closely related, in the deep compressed body, strongly arched lateral line, long dorsal fin, and large mouth. Body deep and compressed; skin smooth; lateral line complete, strongly arched anteriorly; gill membranes united, free from isthmus; no slit behind last gill; preopercular spine simple, short, strongly curved upward, anal papilla large; vent immediately behind ventral fins, about  $\frac{2}{3}$  of distance from gill membrane to anal; ventral rays I, 3. (*σιγμα*, the letter S, from the form of the lateral line.)

2382(a). SIGMISTES CAULIAS, Rutter, new species.

Head  $3\frac{1}{2}$ ; depth  $3\frac{1}{2}$ . D. IX, 20 (IX, 21 in 1 specimen); A. 15 (14 in each of 2 specimens); P. 13. Back elevated, body compressed; eyes lateral,  $4\frac{1}{2}$  in head; snout  $3\frac{1}{2}$ ; cleft of mouth lateral; maxillary 2 in head, reaching to below pupil (only a little past front of eye in 1 specimen). Teeth coarse, cardiform, the inner row of upper jaw enlarged, almost canine-like; a similar pair in inner series of mandible, near symphysis; a small patch on vomer, and 1 on front of palatines; preopercular spine small, sharp, appressed, strongly curved upward, the preopercular margin without spines or tubercles below it; nostrils in short tubes, 1 pair directly behind nasal spines, 1 pair lateral, directly in front of eyes; nasal spines strong, sharp; a pair of tufted cirri above eyes, a pair simple or branched at occiput, and a pair of simple ones halfway between these; a filament on nasal spine, a series of 3 or 4 short ones on margin of preopercle and 1

A.	
12	13
22	2

nt opercular angle; a series of pores around under side of jaw and along edge of preopercle, 2 concentric series under eye and across cheek, and others scattered on head behind eyes; skin smooth, lateral line strongly arched. Dorsal fins connected at base, third spine longest,  $2\frac{1}{2}$  in head, margin of fin even from third to sixth spines, origin of spinous dorsal over upper edge of gill opening; soft dorsal higher, longest rays 2 in head, base of soft dorsal  $\frac{1}{2}$  length; tips of anal rays all free, longest  $2\frac{1}{2}$  in head; origin of anal under third ray of soft dorsal; longest pectoral ray a little longer than head; caudal truncate,  $1\frac{1}{2}$  in head; ventral about reaching anal, about same length as anal papilla; tail slender, least depth slightly less than eye, length from anal  $1\frac{1}{2}$  in head, its length from dorsal about equal to its depth. Color in life, pale pinkish; spinous dorsal dusky, nearly black along margin; soft dorsal plain or with dusky cross bars; anal with about 7 dusky cross bars, extending downward and forward almost at right angles to the rays; 3 or 4 pale blotches surrounded by a black ring along base of dorsal, 1 between dorsals, 1 at end of soft dorsal, and others at base of soft dorsal (some or all sometimes absent); a curved dark line from snout through eye to preopercular spine. Six specimens, 1 each 2,  $2\frac{1}{2}$ , and 3 inches long, and 3  $1\frac{1}{2}$  inches long. From the rock pools at Karluk, on the Island of Kadiak. Coll. Cloudsley Rutter. The type is in Leland Stanford Junior University Museum. Cotypes are in the U. S. Fish Commission and U. S. National Museum. ( $\kappa\alpha\upsilon\lambda\omicron\varsigma$ , stem, from the many dorsal rays.)

**Page 2015.** In third line under *Oxycottus acuticeps* instead of "region" read "reaching."

*Oxycottus* is much nearer *Blennicottus* than *Oligocottus*, and perhaps is best placed as a subgenus of *Blennicottus*. There is no slit behind the last gill in any of the species. In the subgenus *Oxycottus* should be placed:

2883. *BLENNICOTTUS ACUTICEPS* (Gilbert).

2884. *BLENNICOTTUS EMBRYUM* (Jordan & Gilbert).

**Page 2042.** In line 9 for *Phalangistes* substitute *Brachyopsis*.

**Page 2051.** In first line of footnote, for "Dr. Gilbert" read "Seofield & Seale."

**Page 2071.** In line 14 of the description of *Aceruncus sterletus* read "upward," not "downward."

**Page 2108.** The synonymy on this page and the last synonym on page 2107 all belong with the footnote.

**Page 2113.** The type number of *Neoliparis greeni* is 3019, not 3010.

**Page 2128.** Before *Bathypasma* insert the following:

785(a). *CRYSTALLICHTHYS*, Jordan & Gilbert, new genus.

*Crystallichthys*, JORDAN & GILBERT, new genus (*mirabilis*).

Closely allied to *Liparis*, but with nostril single. A single dorsal fin; a well-developed sucking disk; wide bands of teeth, many of which are trilobate near tip; an inferior mouth, much overhung by the produced

conical snout; a single nostril, corresponding to the anterior nostril of other Liparids, the posterior opening being wholly wanting. The typical species, *C. mirabilis*, differs from all known species of *Liparis* except *L. cyclostigma* in its large size, compressed form, and translucent gelatinous texture. (*κρυστάλλος*, crystal; *ἰχθύς*, fish.)

2458(a). CRYSTALLICHTHYS MIRABILIS, Jordan & Gilbert, new species.

Head 4 in length; depth  $2\frac{1}{2}$ ; snout  $2\frac{1}{2}$  in head; eye  $3\frac{1}{2}$  in snout; width of mouth  $\frac{1}{2}$  length of head; length of gill slit  $\frac{1}{2}$  snout, equaling distance from front of eye to front of nostril tube; P. 33. Head and body compressed, especially along upper profile, which descends in a gentle, nearly even curve to tip of snout; lower profile less curved, nearly straight and horizontal on anterior third of body; snout conical, tapering to a sharp tip, its lower profile nearly horizontal, protruding beyond the mouth for a distance (measured axially) equaling  $\frac{2}{3}$  its length; mandibular symphysis vertically below nostril tube; upper jaw strongly arched anteriorly, the mandible much shorter, nearly transverse in position. When the mouth is closed, there is exposed the entire width of the thick upper lip and the anterior portion of the band of fringes which precedes the premaxillary teeth. Teeth slender, shorter than in *Liparis cyclostigma*, arranged in about 25 oblique series in the  $\frac{1}{2}$  of each jaw; the posterior longer teeth more or less distinctly 3-lobed in both jaws, the anterior teeth shorter, simple. A deep cleft on lower side of snout running from its tip to front of premaxillaries, deepening backward, opening into the deep groove above premaxillaries; from base of cleft arises a high free fold, the sharp edge of which nearly reaches the margins of the cleft; a series of 3 large pores along each side of this cleft, with 3 more equally spaced on each side and parallel with front of mouth; belonging to this series but distant from them and much smaller, is another on middle of cheek below eye, and 1 halfway between eye and middle of gill slit; a pore behind eye, and a series of 4 on each side of nape complete the pores of the head; no pore in the position of the posterior nasal opening; a second series of 6 on each side of mandible and preopercle; no other pores on head. Nostril single, in a distinct wide tube, as long as the diameter of pupil; distance from eye to angle of mouth  $3\frac{1}{2}$  in head; vertical from angle of mouth, passing through front of orbit. Gill cleft narrow, reaching base of first pectoral ray, its length  $4\frac{1}{2}$  in head. Lateral line rising in an abrupt curve from upper end of gill opening, decurved again behind pectorals, to reach middle of sides, on the posterior half of which it becomes obsolete; anteriorly the lateral line is accompanied above by a second series of pores which is not curved, but runs straight forward from just above the summit of the curve. Dorsal and anal fins enveloped anteriorly in thick gelatinous tissue, so that their points of origin and number of fin rays can not be determined, the fins high, the longest anal ray equaling length of snout and eye; 32 dorsal and 33 anal rays can be distinguished in the posterior transparent portions of the fins, the total number of rays being greater; last anal ray joining outer caudal ray at middle of length of the latter; dorsal joined narrowly to

base of caudal at end of basal seventh of outer caudal ray; longest caudal ray  $2\frac{1}{2}$  in head; lower 7 pectoral rays thickened, forming a lobe, the distal third of each ray free from the membrane; longest pectoral ray  $1\frac{1}{2}$  in head; disk of moderate size, anteriorly placed, its posterior margin under the gill slit, its length  $\frac{1}{2}$  that of head. Color translucent, apparently light grayish or purplish in life, the dorsal region, including dorsal fin, marked with many large round spots, probably reddish in life, each spot surrounded with a faint darker ring. A large species, soft and gelatinous in texture, the color translucent grayish or purplish, marked on back with many large light circles which were probably reddish in life. Type, a specimen 330 mm. long, from *Albatross Station 3643*, off Provostmaya, Kamchatka, at a depth of 100 fathoms. Bering Sea; 1 specimen from Kamchatka, a smaller one dredged off St. Paul Island, Pribilof group. (*mirabilis*, wonderful.)

*Crystallichthys mirabilis*, JORDAN & GILBERT, Rept. Fur Seal Invest., MS., 1898, Provostmaya, Kamchatka. (Coll. *Albatross*.)

Page 2129. *Allurus* is preoccupied by *Allurus*, Forster, 1862, a genus of *Hymenoptera*; also by *Allurus*, Eisen, 1874, a genus of worms. We substitute for our use of it the name *Allinectes*.

*Allinectes*, JORDAN & EVERMANN, new subgenus (*ectenes*).

Page 2131, line 7, for "*Allurus*" read "*Allurus*." In lines 4 and 7, for "*ἀλλός*" read "*ἄλλος*."

Page 2137. After *Careproctus ectenes* add:

787(a). **PROGNURUS**, Jordan & Everman w genus.

*Prognurus*, JORDAN & EVERMANN, new genus (*cypselurus*).

This genus is distinguished from *Careproctus* by the very elongate caudal which is forked at the tip. (*πρόγυρη*, a swallow or martin; *οὐρά*, tail.)

2469(a). **PROGNURUS CYPSELURUS**, Jordan & Gilbert, new species.

Head  $4\frac{3}{8}$  in length; depth  $4\frac{1}{8}$ ; cleft of mouth  $1\frac{1}{2}$  in head,  $\frac{3}{4}$  distance from symphysis of lower jaw to angle of mouth; total interorbital width  $2\frac{1}{2}$  in head; eye large, equaling length of snout,  $3\frac{1}{2}$  in head; gill opening entirely above base of pectoral, not reaching base of upper ray, its length 3 in head; opercular lobe broadly rounded. Snout blunt, broadly rounded, the mouth horizontal along its lower margin, scarcely overlapped by it; upper lip wide. Teeth acute, without cusps, in about 27 oblique rows in 1 side of each jaw; maxillary reaching the vertical from posterior edge of the pupil; nostril opening in a wide, low tube. Front margin of ventral disk very slightly behind angle of mouth, its diameter  $\frac{2}{3}$  that of eye, about  $\frac{1}{2}$  length of head. Pectorals broadly rounded, regularly shortened below, not deeply notched, the lower 7 rays thickened and exerted; the longest free ray about  $\frac{1}{2}$  length of head; upper portion of fin with 26 rays, the tips only protruding, the longest equaling length of head; dorsal beginning shortly behind vertical from gill slit, its distance from tip of snout  $3\frac{3}{4}$  in length; dorsal with about 58 rays; caudal very long and narrow, only its basal third connate with last

rays of dorsal and anal. Unlike all other Liparids, the caudal is forked at tip, the terminal notch involving about  $\frac{1}{4}$  of fin. Translucent dusky, darker around snout, gill openings, and on the fins, the vertical fins largely jet-black; mouth and gill cavity dusky, not black. This species is most nearly related to *Careproctus melanurus*, from which it differs in darker coloration and shorter gill slit. From all known species of *Careproctus* it differs in the very elongate caudal fin which is forked at the tip. Bering Sea and North Pacific. The type, a single specimen, 21 cm. long, dredged at Albatross Station 3614, off Bogoslof Island, at a depth of 664 fathoms. A second specimen was obtained by the Albatross in 1889, at Station 3074, off the coast of Washington, in 877 fathoms, but it was too seriously mutilated to admit of description. (*κυψελός*, a swift; *ὀψά*, tail.)

**Page 2175.** The genus *Chelidonichthys* should be compared with *Trigla* rather than with *Prionotus*.

*Chelidonichthys pictipinnis* is probably not American, and should be omitted or, at most, admitted only in a footnote.

The genus *Chelidonichthys* differs from *Trigla* in the absence of lateral plates.

**Page 2183.** To the synonymy of *Cephalacanthus* add:

*Cephalacandia*, RAFINESQUE, Anal. de la Nature 1815, 85; substitute for *Cephalacanthus*.

**Page 2196**, line 5, for "Pañeca" read "Puñeca."

**Page 2207.** *Sicya* being preoccupied in *Lepidoptera* we substitute for our use of it the name *Sicyonus*.

*Sicyonus*, JORDAN & EVERMANN, new subgenus (*gymnogaster*).

**Page 2207.** Add the following species:

2531(a). *SICYDIUM PUNCTATUM*, Perugia.

D. VI-I, 11; A. I, 10; scales 56. Head  $5\frac{1}{2}$  in total length without caudal, its width equaling its height or  $\frac{3}{4}$  that of body under first dorsal. Scales of body larger than those of head or nape; maxillary reaching posterior border of eye. Eye 4 in head, or  $1\frac{1}{2}$  in interorbital space. Snout 4; pectoral equaling head in length; spines of first dorsal somewhat elongated, the longest (third) twice height of body; second dorsal as high as body and like the anal. Teeth of upper jaw fine, very slender, and ending behind in an obtuse angle; lower jaw with conic robust teeth and minute horizontal ones. Color grayish, the ventral gill color (giallognolo); under part of head with numerous small black spots; scales strongly ciliated and each with a brown spot in the center; dorsals brown; anal transparent, with a narrow black line; ventral disk yellowish. Length 8 cm. This species is not *S. plumieri* of Cuvier & Valenciennes, nor is it *S. antillarum*, Ogilvie-Grant, because of the difference in the number of scales, the different proportions and a different coloration. The type was collected by Captain Guiseppe Capurro at St. Pierre, Martinique. (Perugia.)

*Sicydium punctatum*, PERUGIA, Annali del Museo Civico di Storia Naturale di Genova, ser. 2, vol. xvi, 1896, 18, Martinique. (Coll. Capt. Guiseppe Capurro.)

**Page 2226.** *Gobius zebra* has 26 scales. Many fine specimens of this species, 3 to 4 inches long, from Clarion Island, are in the museum of Stanford University.

**Page 2227.** The type locality of *Gobius bosci* is Martinique.

**Page 2230,** second line, read "ÉMÉRAUDE" for "EMERANDE."

**Page 2241.** Insert the following synonymy after No. 2572:

*Gobius lucretiae*, EIGENMANN & EIGENMANN, Proc. Cal. Ac. Sci., ser. 2, vol. I, Jan. 25, 1888, 57, Pearl Island, Gulf of Panama.

**Page 2263.** To the synonymy of *Gobioides* add:

*Plecopodus*, RAFINESQUE, Anal. de la Nature 1815, 87; substitute for *Gobioides*.

**Page 2269.** Omit the last reference but one.

**Page 2314.** To the synonymy of *Batrachoides* add:

*Batrictius*, RAFINESQUE, Anal. de la Nature 1815, 82; substitute for *Batrachoides*.

**Page 2350.** After *Enneanectes carminalis* insert:

#### 868(a). DIALOMMUS, Gilbert.

*Dialommus*, GILBERT, Proc. U. S. Nat. Mus. 1890, 452 (*fuscus*).

Teeth conic, strong, in a narrow band in the front of both jaws, thin narrowing to a single series laterally; outer teeth enlarged in both jaws. Teeth on vomer in a single series; palatines smooth. A single slender tentacle above orbits, and 1 on each side of nape. Body with moderate cycloid scales; lateral line high in front, declining behind pectoral fins, not strongly developed, evident on a few scales near head, the remainder of its course traceable by occasional pores on bases of scales or by their notched margins. Dorsal beginning on the nape, its anterior  $\frac{1}{2}$  composed of slender flexible spines, the remainder of soft rays, unbranched; anal without spines; caudal distinct, rounded; ventrals well developed, I, 3. Eyes as in *Anableps*, the cornea divided by an oblique pigmented band into an anterior lower and a posterior upper half. One species known. (*δία γύω*, to loose one from another, to part asunder; *ὄμμα*, eye.)

#### 2687(a). DIALOMMUS FUSCUS, Gilbert.

Head 5 in length; depth 6 to 7. D. XXV, 13 or 14; A. I, 28; lateral line, 52. Elongate, slender, scarcely tapering. Head short, transversely evenly rounded, with very short, blunt, decurved snout; width of head greater than its depth, and more than  $\frac{1}{2}$  its length. Mouth horizontal at lower outline of snout, the maxillary nearly reaching vertical from posterior margin of orbit,  $2\frac{1}{2}$  in head. Teeth strong, conical, the outer series enlarged in both jaws, a narrow band of villiform teeth behind the outer series; vomer with a single series; palatines toothless. Eyes large, round, closely approximated, their diameter greater than length of snout, twice the width of interorbital space,  $3\frac{1}{2}$  in head. Gill membranes very widely joined, free from isthmus. No hook on inner edge of shoulder girdle. Dorsal fin beginning on the nape, over front of opercle, its spines slender and flexible, much lower than soft rays; height of anterior and middle spines about equal,  $\frac{1}{2}$  length of head; the posterior spines shortened, about  $\frac{1}{2}$  that length; height of soft rays  $\frac{1}{2}$  head; first anal ray short and spinous, the succeeding rays articulated, but not branched (like those of dorsal). Interradial membranes of anal fin very deeply incised; caudal



fin wholly free, rounded, its length nearly equaling that of head; pectorals slightly shorter than head, posteriorly pointed, the longest rays below the middle of fin; ventrals comparatively broad, inserted but little in front of pectorals, their bases separated by a space equal to  $\frac{1}{4}$  diameter of orbit. Color in spirits, brownish above and on sides, becoming blackish on head; under side of head, belly, and a line along each side of anal fin light; back with traces of about 10 black cross bars, which invade base of dorsal fin and extend onto middle of sides; in 1 specimen the scales of the interspaces are marked each with a light spot (probably blue in life); fins all dusky, the caudal variegated with lighter in fine pattern; ventrals light at base. Two specimens from the Galapagos Islands—1 from Duncan Island, 72 mm. long, the other from Albemarle Island, 75 mm. long.

*Dialommus fuscus*, GILBERT, Proc. U. S. Nat. Mus. 1890, 452, Galapagos Islands. (Coll. Albatross)

**Page 2352.** *Gibbonsia crides* intergrades with *Gibbonsia elegans* and must apparently be regarded as a subspecies of the latter.

**Page 2356.** Under *Malacoctenus ocellatus* for "A. II, 8" read "A. II, 18."

**Page 2413.** Genus 904 should read *Ulvicola*, Gilbert, not "Gilbert & Starks."

**Page 2421.** *Anoplarchus alectrolophus* is the type of a new genus:

908(a). ALECTRIAS, Jordan & Evermann.

*Alectrias*, JORDAN & EVERMANN, new genus (*alectrolophus*).

This genus is distinguished from *Anoplarchus* by having the gill membranes free from the isthmus, as in *Cebedichthys*. (*ἀλέκτωρ*, cock; from the crest.)

**Page 2422**, under *Anoplarchus alectrolophus*: The 2 specimens obtained in Monterey Bay by Arthur W. Greeley were erroneously referred to this species. They are the young of *Cebedichthys*.

**Page 2470.** To synonymy of *Lycenchelys* add:

*Lycodopsis*, VAILLANT, Exp. Sci. Trav. et Talisman, 311, 1888 (*albus*).

**Page 3472.** *Furcella* is preoccupied by *Furcella*, Lamarck, 1801, a genus of Mollusca. We substitute for our use of it the name *Furcimanus*.

*Furcimanus*, JORDAN & EVERMANN, new genus (*diapterus*).

**Page 2473.** In lines 33, 36, and 37 read "in 282 to 376 fathoms" instead of the depths given.

**Page 2475.** In lines 26 and 28, for "coast of Alaska" read "coast of California."

**Page 2480.** The locality for *Melanostigma pammelas* is coast of California, not Alaska.

**Page 2567.** In first line of footnote read "fin," not "spine."

**Page 2601**, line 20, for "*Trachyterus*" read "*Trachypterus*."

**Page 494.** The large "Silver Trout" of Lake Tahoe, a specimen of which is described on page 494, should probably be separated subspecific-

ally from its parent form, *Salmo clarkii henshawi*. It may be described as follows:

780(d). SALMO CLARKII TAHOENSIS, Jordan & Evermann, new subspecies.

(SILVER TROUT OF LAKE TAHOE.)

Head  $4\frac{1}{5}$ ; depth  $3\frac{1}{2}$ ; eye  $7\frac{2}{3}$  in head. D. 9; A. 12; Br. 10; scales 33-205-40; 140 pores. Pectoral  $1\frac{1}{2}$  in head; maxillary  $1\frac{1}{2}$ . Body very robust, compressed, unusually deep for a trout, the outline elliptical. Head large, rather more compressed than in typical *Salmo clarkii henshawi*; eye small, silvery. Vomerine teeth in 2 long series, those of the 2 series alternating in position; hyoid teeth distinct, in a rather long series; gill rakers short, thickish, 5 + 13. Mouth large, the maxillary extending well beyond the eye. Preopercle moderate, its lower posterior edge not evenly rounded, but with a slightly projecting, rounded lobe and a slight concavity above and below, this character not strongly marked; opercle evenly, but not strongly, rounded. Scales small, reduced above and below, those in or near lateral line largest. Fins moderate, the anal rather high, with 1 more ray than usual; caudal slightly lunato, almost truncate when spread open. Color dark green above, belly silvery; sides with a broad coppery shade covering cheeks and opercles; sides of lower jaw yellowish; fins olivaceous, a little reddish below; orange dashes between rami of lower jaw moderately conspicuous; back, from tip of snout to tail, closely covered with large, unequal black spots; spots on top of head and nape round; posteriorly the spots run together, forming variously shaped markings, usually vertically oblong; these may be regarded as formed of 3 or 4 spots placed in a series, or with 1 or 2 at the side of the other; the longest of the oblong markings not quite as long as eye; along side of head and body the spots are very sparse, those on head round, those behind vertically oblong; belly profusely covered with small black spots which are nearly round; still smaller round spots numerous on lower jaw; all the spots on caudal peduncle vertically oblong or curved; dorsal and caudal densely covered with oblong spots, smaller than those on the body; anal with rather numerous round spots; pectorals and ventrals with a few small spots, the first ray in each case with a series of faint small spots; adipose fin spotted. The above description from a specimen 2 feet 4 inches long and weighing  $7\frac{1}{2}$  pounds. This form attains a weight of 10 to 30 pounds and spawns only in the depths of the lake. *Salmo clarkii henshawi* reaches a much smaller size, is much darker in color, and spawns in the streams. Thus far known only from the deep waters of Lake Tahoe. (*tahoensis*, from Lake Tahoe.)

*Salmo clarkii tahoensis*, JORDAN & EVERMANN, new subspecies, Lake Tahoe. (Coll. A. J. Bayley.)

**Page 518.** It is wholly uncertain where Valenciennes got the specimen which he called *Thymallus ontariensis*. It is probably the ordinary Grayling (*Thymallus thymallus*) of Europe, erroneously attributed to Milbert's New York collection. In any case, its identity with the Michigan Grayling is more than doubtful, as the rivers in which the latter occurs were then unexplored. The American Graylings would therefore stand as follows:

787. THYMALLUS SIGNIFER (Richardson).

(ARCTIC GRAYLING; POISSON BLEU.)

788. THYMALLUS TRICOLOR, Cope.

(MICHIGAN GRAYLING.)

788(a). THYMALLUS TRICOLOR MONTANUS (Milner).

(MONTANA GRAYLING.)

To the synonymy of the Montana Grayling add the following wholly unnecessary synonym:

*Thymallus lewisii*, HENSHALL, Forest and Stream, July 23, 1898, 70, headwaters Jefferson River, Red Rock Lake, Montana; after notes of Lewis & Clark.

Dr. J. C. Merrill, U. S. A., informs us that this Grayling is found also in Sun River at Fort Shaw, Montana.

Page 852. The species called *Holocentrus marianus*, Cuvier & Valenciennes, is the type of a distinct genus:

382 (a). FLAMMEO, Jordan & Evermann, new genus.

*Flammeo*, JORDAN & EVERMANN, new genus (*marianus*).

This genus is distinguished by the very large mouth and projecting chin. The lower jaw is considerably more than  $\frac{1}{2}$  the length of the head, and the chin projects beyond the upper jaw. In the species properly referable to *Holocentrus*, the lower jaw is slightly included and its length is less than  $\frac{1}{2}$  the head. The single known species of this genus is

2238. FLAMMEO MARIANUS (Cuvier & Valenciennes).

Page 858. In line 13 the interorbital space in *Upeneus* is said to be "concave;" this should, of course, read "convex."

Page 2013. *Oligocottus maculosus*, Girard, was evidently based on a specimen from the Farallones of the species called by us *Oligocottus borealis*, Jordan & Snyder. The name *maculosus* must therefore be transferred to the latter species. The species called by us *Oligocottus maculosus* must therefore be renamed and may stand as

2381. OLIGOCOTTUS SNYDERI, Greeley, new species.

*Oligocottus snyderi*, GREELEY, MS. 1898.

Page 2126, lines 13 and 14. The specimens from St. Paul Island and Petropaulski, referred by us to *Liparis cyclostigma*, Gilbert, belong to *Crystallichthys mirabilis*, Jordan & Gilbert, described on page 2865.

Page 2626. Before *Paralichthys estuarius*, Gilbert & Seefield, insert:

2991 (a). PARALICHTHYS MAGDALENE, Abbott, new species.

Head  $3\frac{1}{2}$ ; depth 2 $\frac{1}{2}$ . D. 80; A. 64; scales 120. Body oval-elliptical, the dorsal outline evenly bowed, the greatest depth in the middle of the body.

Ventral outline straighter anteriorly. Mouth large; mandible somewhat projecting, about  $1\frac{1}{2}$  in head; maxillary large, extending considerably beyond orbit; snout (measuring from upper orbit) about 4 in head,  $\frac{3}{4}$  length of eye; interorbital about  $9\frac{1}{2}$  in head; eye 6 in head. Teeth moderately strong, the anterior ones in the lower jaw somewhat larger than the others; gill rakers 7 + 17, slender, weakly serrate, the longest a trifle less than eye. Scales cycloid, on cheeks, opercles and maxillary; snout, interorbital, and mandible naked; accessory scales present, especially prominent among the small crowded scales in the region below the pectoral; arch of lateral line  $3\frac{1}{2}$  in straight part; pores about 38 + 82. Ventral  $3\frac{1}{2}$  in head; pectoral 2; dorsal beginning above anterior rim of orbit; middle rays of anal and dorsal longest,  $2\frac{1}{2}$  in head, equaling width of caudal peduncle; caudal double lunate, the middle rays the longest; pectoral of blind side rounded,  $2\frac{1}{2}$  in head. Color dark reddish brown, closely peppered with darker dots; a series of indistinct white spots, 4 or 5 in number, following margins of the body, as in *P. californicus*; traces of darker mottling along sides of body. Length 17 inches.

This species, represented by a specimen from Magdalena Bay, Lower California, is closely related to *P. californicus*, resembling it in the large number and close arrangement of the gill rakers, but differing from it in having cycloid scales, a greater number of fin rays, somewhat narrower interorbital, and greater depth in proportion to the length.

*Paralichthys magdalenae*, ABBOTT MS., new species, Magdalena Bay, Lower California. (Type, No. 10196, L. S. Jr. Univ. Mus. Coll. Charles H. Gilbert.)

Page 2627. Instead of *Paralichthys adspersus* (Steindachner), read :

2994. PARALICHTHYS SINALOE, Jordan & Abbott, new species.

The specimens described in the text (from Mazatlan and La Paz), under the name *Paralichthys adspersus*, belong to a distinct species, thus far known only from the west coast of Mexico and Central America and which may be called *Paralichthys sinaloe*, one of the many specimens taken by the Hopkins Expedition at Mazatlan and La Paz being taken as type, and the following as cotypes: Nos. 11726, 11727, 11728, 11729, 11730, 11731, L. S. Jr. Univ. Mus., all from Mazatlan. *Paralichthys adspersus* is known only from the coast of Peru, the specimens before us being from Callao. (Coll. Admiral L. A. Beardslee.)

*Paralichthys sinaloe* is distinguished from *P. adspersus* by its cycloid scales and broader interorbital space. The gill rakers in *P. adspersus* are close-set, rather long and slender, about  $\frac{1}{4}$  to  $\frac{1}{3}$  of eye, and with rather slender spinules on the inner margin. In *P. sinaloe* they are set farther apart on the limb of the gill arch, are shorter and thicker—about 2 in eye—and have the inner edge armed with coarser teeth. All specimens of *P. sinaloe* have each 14 or 13 gill rakers on the lower limb of the arch, while in 4 examples of *P. adspersus*, from Callao, there are 16 or 17 gill rakers on the lower limb (19 in 1 specimen), and 7 or 6 above (5 in 1 specimen), showing that while there may be variation in the number yet it is confined within limits which do not intergrade and the average number

in the 2 species is quite different. In the single known specimen of *P. woolmani* the number of gill rakers given is 5 + 11, which makes it probable that the present species is not the same. The more striking difference between *P. adspersus* and *P. sinaloa* lies in the scales, which in the latter are cycloid, while in the true *P. adspersus* they are strongly toothed as stated by Steindachner. The specimens from Callao referred by us in the text to *adspersus* belong to that species, but they are not original types of *adspersus*, belonging to the later collections of Agassiz and Steindachner.

*Paratichthys sinaloa*, JORDAN & ABBOTT MS., new species, Mazatlan, Sinaloa, Mexico. (Type, No. 2930, L. S. Jr. Univ. Mus. Coll. Hopkins Exped. to Mazatlan.)

Page 686. After *Platypecilus maculatus* add:

1009(a). *PLATYPECILUS QUITZEENSIS*, B. A. Bean.

Head  $3\frac{1}{2}$ ; depth  $2\frac{1}{2}$ ; eye  $3\frac{1}{2}$  in head; snout 4; interorbital width  $2\frac{3}{4}$ . D. 13; A. 13; scales 30, 10. Body compressed, back elevated, head small and depressed, flat on top; snout short. Mouth small, cleft oblique, the lower jaw heavy, projecting; teeth conic, those in upper jaw in an irregular series, those below very small, apparently irregularly arranged and close-set. Origin of dorsal fin in advance of that of anal, midway between tip of upper jaw and end of caudal rays, the first ray of anal being under sixth dorsal ray. Color in alcohol light brown, with traces of darker on back; interorbital space and edge of scales dark brown; 3 dark bars on posterior part of body, the first extending from median line to origin of anal, the second from median line to end of anal base, the third midway between end of anal and origin of caudal; 2 dark spots on end of caudal peduncle; fins all pale. Lake Quitzeo, Michoacan, Mexico. Only the type known.

*Platypecilus quitzeensis*, B. A. BEAN, Proc. U. S. Nat. Mus. 1898, 540, with text figure, Lake Quitzeo, Michoacan, Mexico. (Type, No. 48209. Coll. E. W. Nelson.)



**ARTIFICIAL KEY TO THE FAMILIES OF THE TRUE  
FISHES OR TELEOSTEI.**

The following key is intended simply to facilitate the identification of species of the true fishes. No attempt is made to indicate the natural characters or relations of the families, and only those species of any group which are included in the present work are taken into consideration. Most of the ordinary fishes can be readily placed by its means, but it should not be trusted in the study of ichthyological rarities, or of fishes from the deep seas.

**I.—VENTRAL FINS PRESENT, ABDOMINAL.**

- A. Back with an adipose fin behind the single rayed dorsal fin.
- B. Adipose fin composed of a single spine with a thin membrane; body mailed. . . . .XXXV, LORICARIIDÆ, 155.
- BB. Adipose fin without spine.
- C. Head with 4 to 8 long barbels about the mouth and nostrils; body scaleless; a single spine in each pectoral and in dorsal fin. . . . .XXXIV, SILURIDÆ, 115.
- CC. Head without barbels as described above.
- D. Sides of body without photophores or luminous glands; no barbel at throat.
- E. Body scaleless; teeth very strong, some of them fang-like.
- F. Dorsal fin very long and high, occupying nearly whole length of back. . . . .LXXXI, ALEPISAUROIDÆ, 593.
- FF. Dorsal fin short, median or posterior. . . . .LXXXII, ODONTOSTOMATIDÆ, 597.
- EE. Body scaly.
- G. Pseudobranchiæ present.
- H. Dorsal, anal, and ventrals each with a small but distinct spine; scales ctenoid. . . . .CIV, PERCOPSIDÆ, 783.
- HH. Dorsal, anal and ventral without distinct spine.
- I. Head naked.
- J. Branchiostegals 6 to 20.
- K. Dorsal fin long and high, of about 24 rays. . . . .LXV, THYMALLIDÆ, 517.
- KK. Dorsal fin moderate, of fewer than 20 rays.
- L. Stomach with many pyloric cæca. . . . .LXIV, SALMONIDÆ, 460.
- LL. Stomach with few pyloric cæca; size small. . . . .LXVI, ARGENTINIDÆ, 519.
- JJ. Branchiostegals 3 or 4; mouth very small. . . . .LXVII, MICROSTOMATIDÆ, 527.
- II. Head scaly on sides.
- M. Maxillary very narrow, rudimentary, or obsolete; hypocoracoids not divergent. . . . .LXVIII, SYNODONTIDÆ, 532.
- MM. Maxillary well developed, dilated behind; pectorals normal; hypocoracoids mostly divergent. . . . .LXIX, AULOPODIDÆ, 541.
- GG. Pseudobranchiæ absent.
- N. Pectorals normally formed, teeth incisor-like or else rudimentary; pseudobranchiæ absent. . . . .XXIX, CHARACINIDÆ, 331.

- NN. Pectorals not normally formed.  
   O. Pectorals undivided, subhumeral; pseudobranchiæ absent.  
     LXX, BENTHOSAURIDÆ, 543.  
 OO. Pectoral rays elongate, arranged in two groups.  
     LXXI, BATHYPTEROIDIDÆ, 544.  
 DD. Sides of body with photophores more or less developed.  
   P. Barbel at throat present, very long; body naked.  
     LXXVIII, ASTRONESTHIDÆ, 586.  
 PP. Barbels none.  
   Q. Vertebral spines projecting through skin of back before dorsal  
     fin; body short and deep, greatly compressed.  
     LXXXIV, STERNOPTYCHIDÆ, 603.  
 QQ. Vertebral spines not exerted in front of dorsal.  
   R. Pseudobranchiæ present.  
   S. Premaxillaries forming entire margin of upper jaw; body scaly;  
     opercles complete.  
   T. Form elongate, the snout pointed, barracuda-like; photophores  
     very small. . . . . LXXXIII, PARALEPIDIDÆ, 599.  
 TT. Form oblong, the snout not much produced; photophores con-  
     spicuous. . . . . LXXV, MYCTOPHIDÆ, 550.  
 SS. Premaxillaries not forming the whole margin of upper jaw, the  
     maxillary entering into it; body naked; opercular appa-  
     ratus incomplete. . . . . LXXXVI, MAUROLOCIDÆ, 576.  
 RR. Pseudobranchiæ absent; mouth large, with canine teeth; scales  
     deciduous or wanting. . . . . LXXVII, CHAULIODONTIDÆ, 578.  
 AA. Back without adipose fin.  
   B. Back with a single dorsal fin made up of rays and not preceded  
     by a series of free spines or followed by finlets.  
   C. Tail evidently strongly heterocercal.  
   D. Body naked; snout with a spatulate blade; mouth wide, without  
     barbels. . . . . XXX, POLYODONTIDÆ, 101.  
 DD. Body with 5 series of body shields; mouth, inferior, toothless,  
     preceded by 4 barbels. . . . . XXXI, ACIPENSERIDÆ, 102.  
 DDD. Body scaly.  
   E. Scales cycloid; a broad bony gular plate; dorsal fin many rayed.  
     XXXIII, AMIIDÆ, 112.  
   EE. Scales ganoid; no gular plate; dorsal fin short.  
     XXXII, LEPISOSTEIDÆ, 108.  
 CC. Tail not evidently heterocercal.  
   F. Tail tapering to a point, without caudal fin; anal fin very long,  
     of about 200 rays; body scaly.  
     LXXXVI, HALOSAURIDÆ, 606.  
 FF. Tail not tapering to a point; caudal fin developed.  
   G. Body naked.  
   H. Throat with a long barbel; no caudal filament; mouth large.  
   I. Barbel free at tip. . . . . LXXIX, STOMIDÆ, 587.  
   II. Barbel connecting throat with symphysis of lower jaw.  
     LXXX, MALACOSTEIDÆ, 592.  
 HH. Throat without barbel.  
   J. Caudal fin with a long filament; body elongate; mouth very  
     small. . . . . CI, FISTULARIIDÆ, 755.  
 JJ. Caudal fin without filament.  
   K. Pectorals present.  
   L. Gill membranes joined to the isthmus; opercles complete.  
     XXXVII, CYPRINIDÆ, 199.  
 LL. Gill membranes free from isthmus; opercles incomplete.  
     LXXXIII, RONDELETIDÆ, 547.  
 KK. Pectorals wanting; body snake-like; dorsal long and low.  
     LXXXV, IDIACANTHIDÆ, 604.  
 GG. Body scaly.



- M. Head with a large divided luminous plate in place of eyes. LXXII, IPNOPIDÆ, 546.
- MM. Head with eyes concealed beneath the skin; vent at the throat. XCIII, AMBLYOPSIDÆ, 702.
- MMM. Head with normally developed eyes.
- N. Body with a coat of mail; maxillary with barbels. XXXV, LORICARIIDÆ, 155.
- NN. Body with ordinary scales.
- O. Anal fin with many spines; mouth toothless, sucker-like. LXXXVIII, LIPOGENYIDÆ, 619.
- OO. Anal fin without distinct spines.
- P. Pectoral fins inserted high, near axis of body; lower pharyngeals united; lateral line along sides of belly.
- Q. Jaws each with long sharp teeth mixed with smaller ones. XCIV, ESOCIDÆ, 708.
- QQ. Jaws with small equal teeth, conic or tricuspid.
- R. Lower jaw more or less produced; teeth tricuspid. XCV, HEMIRAMPHIDÆ, 718.
- RR. Lower jaw a little produced; teeth conic; pectorals elongate, forming an organ of flight..... XCVII, EXOCETIDÆ, 726.
- PP. Pectoral fins inserted below axis of body; lower pharyngeals separate.
- S. Gill membranes broadly joined to the isthmus; head naked; no teeth in jaws.
- T. Lower pharyngeal teeth very numerous, in 1 row like the teeth of a comb. (Suckers.)..... XXXVI, CATOSTOMIDÆ, 161.
- TT. Lower pharyngeal teeth few, fewer than 8, in 1 to 3 rows. (Carp; Chubs; Minnows.)..... XXXVII, CYPRINIDÆ, 199.
- SS. Gill membranes free from the isthmus.
- U. Throat with a long barbel; sides with phosphorescent spots. LXXXIX, STOMIATIDÆ, 587.
- UU. Throat without barbels.
- V. Phosphorescent spots present; teeth unequal. LXXXVII, CHAULIODONTIDÆ, 578.
- VV. Phosphorescent spots none.
- W. Head scaly, more or less.
- X. Maxillaries connate with premaxillaries; jaws long. LXVII, SYNODONTIDÆ, 532.
- XX. Maxillaries distinct.
- Y. Upper jaw not protractile, its lateral margins formed by the maxillaries; lateral line more or less developed.
- Z. Teeth cardiform; jaws depressed, prolonged. XCI, LUCIDÆ, 624.
- ZZ. Teeth villiform; jaws short; no lateral line.
- a. Pectoral very broad, of about 35 rays. LXXXIX, DALLIIDÆ, 620.
- aa. Pectoral narrow, of about 13 rays..... XC, UMBRIDÆ, 622.
- YY. Upper jaw protractile, its margin formed by premaxillaries alone; no lateral line..... XCII, PÆCILLIDÆ, 630.
- WW. Head naked.
- a. Anterior vertebrae coalesced and modified; no pseudobranchiae; jaws with strong canines..... XXXVIII, ERYTHRINIDÆ, 330.
- aa. Anterior vertebrae normal, not modified.
- b. Dorsal fin inserted more or less before anal (rarely slightly behind it); shore fishes or river fishes, usually si very in coloration and with skeleton firm; air bladder all developed.
- c. Gular plate present, between branches of lower jaw; mouth large; teeth present, all pointed; axillary scales and sheaths large..... LVI, ELOPIDÆ, 408.
- cc. Gular plate none.
- d. Lateral line well developed.
- e. Teeth present, no accessory branchial organ.

- f.* Mouth small, horizontal; posterior part of tongue and roof of mouth covered with coarse-paved teeth.  
LVII, ALBULIDÆ, 410.
- ff.* Mouth large, the teeth all pointed, some of them canine, none paved or molar .....LVIII, HIODONTIDÆ, 412.
- ee.* Teeth none; an accessory branchial organ behind gill cavity.  
LIX, CHANIDÆ, 414.
- dd.* Lateral line wanting; no gular plate.
- g.* Mouth small, inferior, toothless, the maxillary simple or nearly so; stomach gizzard-like.....LX, DOROSOMATIDÆ, 415.
- gg.* Mouth moderate, terminal, the maxillary of about 3 pieces; stomach not gizzard-like.....LXI, CLUPEIDÆ, 417.
- ggg.* Mouth subinferior, very large, below a tapering, pig-like snout; maxillary very long.....LXII, ENGRAULIDIDÆ, 439.
- bb.* Dorsal fin posterior, opposite anal; deep-sea fishes, of loose organization; mostly blackish in color; mouth small, with small pointed teeth; air bladder wanting.  
LXIII, ALEPOCEPHALIDÆ, 451.
- BB.* Dorsal fin single, preceded by free spines.
- h.* Body scaleless, naked or with bony plates.
- i.* Ventral fins I, 1, the spine strong; snout moderate.  
XCVIII, GASTEROSTEIDÆ, 742.
- ii.* Ventral fins, I, 5, the spine slender; snout prolonged.  
XCIX, AULORHYNCHIDÆ, 752.
- hh.* Body scaly; snout tubular .....C, AULOSTOMIDÆ, 771.
- BBB.* Dorsal fin composed of free spines; ventrals with 1 or 2 spines each; body elongate.....LXXXVII, NOTACANTHIDÆ, 613.
- BBBB.* Dorsal fins 2, the anterior of spines only, the posterior chiefly of soft rays.
- j.* Pectoral fin with 5 to 8 lowermost rays detached and filamentous .....CIX, POLYNEMIDÆ, 827.
- jj.* Pectoral fin entire.
- k.* Snout tubular, bearing the short jaws at the end; body compressed .....CII, MACRORHAMPHOSIDÆ, 758.
- kk.* Snout not tubular.
- l.* Teeth strong, unequal; lateral line present.  
CVIII, SPIRYRÆNIDÆ, 822.
- ll.* Teeth small or wanting; lateral line obsolete.
- m.* Dorsal spines 4, stout; anal spines 3.....CVII, MUGILIDÆ, 808.
- mm.* Dorsal spines 4 to 8, slender; anal spine single.  
CVI, ATHERINIDÆ, 788.
- BBBBB.* Dorsal fin soft-rayed, followed by a series of detached finlets.  
XCVI, SCOMBROSOCIDÆ, 724.

**II.—VENTRAL FINS PRESENT, THORACIC OR SUBJUGULAR, THE NUMBER OF RAYS DEFINITELY 1, 5.**

- A. Gill openings in front of the pectoral fins.
- B. Body more or less scaly or armed with bony plates.
- C. Ventral fins completely united; gill membranes joined to the isthmus; no lateral line .....CLXXXVIII, GOMIDÆ, 2188.
- CC. Ventral fins separate.
- D. Suborbital with a bony stay, which extends across the cheek to or toward the preopercle; cheeks sometimes entirely mailed.
- E. Pectoral fin with 3 lower rays detached and free; head bony.  
CLXXXIV, TRIGLIDÆ, 2147.
- EE. Pectoral fin with 2 lower rays detached and free; body mailed.  
CLXXXV, PERISTEIIDÆ, 2177.
- EEE. Pectoral fin entire.
- F. Sit behind fourth gill small or wanting.

- G. Dorsal spines 4; lips fringed; eyes superior.  
CXC VII, URANOSCOPIIDÆ, 2305.
- GG. Dorsal spines 8 to 17.
- H. Anal spines 3; body scaly.....CLXXVI, SCORPÆNIDÆ, 1758.
- HH. Anal spines obsolete; body partly or wholly naked.  
CLXXIX, COTTIDÆ, 1879.
- FF. Slit behind fourth gill large; body scaled.
- I. Nostril single on each side, a small pore above it; dorsal fin continuous.....CLXXVIII, HEXAGRAMMIDÆ, 1863.
- II. Nostrils 2 on each side; dorsal fins 2, separate, except in the genus *Erelepis*.....CLXXVII, ANOPILOMATIDÆ, 1861.
- DD. Suborbital stay wanting; cheeks not mailed.
- J. Spinous dorsal transformed into a sucking disk on top of head, composed of 8 to 30 transverse plates.  
CLXXXIX, ECHENEIDIDÆ, 2265.
- JJ. Spinous dorsal (if present) not transformed into a sucking disk.
- K. Dorsal spines all or nearly all disconnected from each other.
- L. Body elongate, spindle-shaped..CXXVII, RACHYCENTRIDÆ, 947.
- LL. Body oblong or ovate, compressed.
- M. Caudal peduncle very slender, the fin widely forked; preopercle entire.....CXXV, CARANGIDÆ, 895.
- MM. Caudal peduncle stoutish, the fin little forked.
- N. Gill membranes free from the isthmus; preopercle serrulate.  
CXXXIV, CENTROLOPHIDÆ, 962.
- NN. Gill membranes broadly united to the isthmus; preopercle entire.  
CLXIV, EPHIPPIDÆ, 1666.
- KK. Dorsal spines (if present) all, or most of them, connected by membrane.
- O. Pectoral fin with 4 to 9 lowermost rays detached and filiform.  
CIX, POLYNEMIDÆ, 827.
- OO. Pectoral fin entire.
- P. Dorsal and anal each with 1 or more detached finlets.
- Q. Anal preceded by 2 free spines.....CXXV, CARANGIDÆ, 895.
- QQ. Anal not preceded by 2 free spines.
- R. Caudal peduncle keeled.....CXVIII, SCOMBRIDÆ, 863.
- RR. Caudal peduncle not keeled.....CXIX, GEMPYLIDÆ, 877.
- PP. Dorsal and anal without finlets.
- S. Lateral line armed posteriorly with a series of keeled plates; 2 free anal spines; gill membranes free from isthmus.  
CXXV, CARANGIDÆ, 895.
- SS. Lateral line armed posteriorly with a sharp, movable, lancet-like spine, or with a few bony tubercles; scales small, rough; gill membranes adherent to isthmus.  
CLXVII, TEUTHIDIDÆ, 1688.
- SSS. Lateral line unarmed.
- T. Throat with 2 long barbels (placed just behind chin); dorsal fins 2.....CXVII, MULLIDÆ, 855.
- TT. Throat without long barbels.
- U. Head with a short bony horn before each eye; gill membranes united to isthmus; scales very small, rough.  
CLXVI, ZANCLIDÆ, 1687.
- UU. Head without bony prominence or horns.
- V. Anal fin preceded by 2 free spines (these obsolete in the very old, joined by membrane in the very young).
- W. Preopercle entire; teeth moderate if present.  
CXXV, CARANGIDÆ, 895.
- WW. Preopercle serrate; teeth unequal, some of them very strong.  
CXXXVI, POMATOMIDÆ, 945.
- VV. Anal fin not preceded by free spines.
- X. Nostril single on each side; lateral line interrupted; lower pharyngeals united.

- Y. Anal spines 2.....CLIX, POMACENTRIDÆ, 1543.  
 YY. Anal spines 3 to 11. Fresh-water fishes. CLVIII, CICHLIDÆ, 1512.  
 XX. Nostril double on each side.  
 Z. Lateral line extending to tip of middle rays of caudal.  
 a. Anal spines 3, the second strong.  
 b. Dorsal fins 2, separate; body elongate.  
     CXLV, CENTROPOMIDÆ, 1116.  
 bb. Dorsal fin continuous.....CL, HÆMULIDÆ, 1289  
 aa. Anal spines 1 or 2, the second large or small.  
     CLV, SCLÆNIDÆ, 1392.  
 ZZ. Lateral line not extending beyond base of caudal fin.  
 c. Gills 3½, the slit behind the last very small or wanting.  
 d. Mouth not vertical, the lips not fringed; dorsal fin continuous, the spines 8 to 18; scales cycloid; lower pharyngeals united.  
 e. Teeth in each side of each jaw united, forming a sort of beak.  
     CLXI, SCARIDÆ, 1620.  
 ee. Teeth distinct or nearly so, the anterior usually more or less canine.....CLX, LAHRIDÆ, 1571.  
 dd. Mouth nearly vertical, the lips with fleshy fringes; dorsal divided, the spinous part short, of about 4 spines; lower pharyngeals separate....CXCVII, URANOSCOUIDÆ, 2305.  
 cc. Gills 4, a long slit behind the fourth.  
 f. Teeth setiform, like the teeth of a brush; body elevated, longer than deep, the soft fins completely scaled; gill membranes attached to the isthmus.  
 g. Dorsal fin continuous.....CLXV, CHÆTODONTIDÆ, 1669.  
 gg. Dorsal fin divided.....CLXIV, EPHIPPIDÆ, 1666.  
 ff. Teeth not setiform.  
 h. Body deeper than long, covered with rough scales; dorsal spines 8; anal spines 3; soft fins very long.  
     CLXIII, CAPROIDÆ, 1663.  
 hh. Body longer than deep.  
 i. Gill membranes broadly joined to isthmus; body long and low; no lateral line.....CLXXXVII, GOBIDÆ, 2188.  
 ii. Gill membranes free from isthmus or very nearly so.  
 j. Premaxillaries excessively protractile, their basal process very long, in a groove at top of cranium.  
 k. Teeth small; scales large, silvery; spines strong.  
     CLIII, GERRIDÆ, 1366.  
 kk. Teeth none; spines slender.....CLII, MENIDÆ, 1364.  
 jj. Premaxillaries moderately protractile or not protractile.  
 l. Lower pharyngeals united; scales large; anal fin with 3 spines and more than 15 soft rays; preopercle entire. (Viviparous fishes of the Californian fauna.)  
     CLVII, EMBIOTOCIDÆ, 1493.  
 ll. Lower pharyngeals separate.  
 m. Body elongate, not compressed, covered with hard grooved scales; jaws box-like..CXXXVIII, TETRAGONURIDÆ, 975.  
 mm. Body not as above.  
 u. Lateral line incomplete or interrupted, running close to dorsal fin; dorsal spines very slender, continuous with the soft rays; body low, covered with small scales; anal fin very long.  
 o. Anal rays fewer than 30; maxillary produced behind.  
     CXCI, OPISTHOGNATHIDÆ, 2279.  
 oo. Anal rays more than 30; maxillary not produced behind.  
     CXCH, BATHYMASTERIDÆ, 2287.  
 nn. Lateral line, if present, not as above.  
 p. Scales circular, cycloid, nonimbricate, each with 1 or 2 erect spines; dorsal spines obsolete.  
     CXII, STEPHANOBERYCIDÆ, 835.

1543.  
1512.  
  
1116.  
1289  
  
1392.  
  
tinu-  
geals  
  
eak.  
1620.  
r less  
1571.  
dorsal  
lower  
2305.  
  
l, lon-  
mem-  
  
1669.  
1666.  
  
dorsal  
  
1663.  
  
low;  
2188.  
  
very  
  
1366.  
1364.  
  
spines  
vivip-  
  
1493.  
  
ooved  
e, 975.  
  
dorsal  
e soft  
a very  
  
2279.  
  
2287.  
  
erect  
  
835.

- pp. Scales not as above.  
 q. Anal fin much longer than dorsal; body much compressed, the belly prominent.  
 r. Dorsal spines none; scales cycloid. .CXI, BATHYCLUPEIDÆ, 834.  
 rr. Dorsal spines few, graduated; anal spines 3. CXXXIX, PEMPHERIDIDÆ, 977.  
 qq. Anal fin not much, if any, longer than dorsal.  
   a. Pseudobranchiæ wanting or covered by skin.  
   t. Dorsal fin of soft rays, only beginning as a crest on the head; caudal widely forked. Pelagic fishes. CXXXIX, CORYPLENIDÆ, 951.  
 tt. Dorsal fin with spines anteriorly, not beginning on the head. Fresh-water fishes.  
   u. Anal spines 3 to 10.  
   v. Dorsal spines 6 to 12; lateral line well developed. CXLI, CENTRARCHIDÆ, 984.  
 vr. Dorsal spines about 4; no lateral line; length less than 2 inches. CXL, ELASSOMATIDÆ, 981.  
 uu. Anal spines 1 or 2; body oblong or elongate; length less than 8 inches. . . . . CXLIII, PERCIDÆ, 1015.  
 ss. Pseudobranchiæ developed.  
 w. Spinous dorsal of 2 or 3 short spines only; anal without spines; scales small, smooth. . . . . CXLVI, SERRANIDÆ, 1126.  
 ww. Spinous dorsal, if present, not as above.  
   x. Opercle ending in a long scaly flap; snout depressed, spatulate; mouth very large, the lower jaw projecting. CXCV, CHELONICHTHYIDÆ, 2293.  
   xx. Opercle not ending in a long scaly flap; snout not greatly depressed.  
   y. Pectoral fin broad, its lower rays thickened and not branched. CLVI, CIRRIITIDÆ, 1490.  
 yy. Pectoral rather narrow at base, its lower rays branched, like the others.  
   z. Dorsal fin continuous, the spines few, slender; maxillary usually with an enlarged tooth behind; nape sometimes with an adipose appendage; anal fin long, even. CXC, MALACANTHIDÆ, 2274.  
 zz. Dorsal fin continuous or divided, not as above.  
   a. Perch-like fishes, the caudal peduncle not very slender, the scales well developed, etenoid or cycloid; the dorsal with distinct spines; the anal with at least 1 spine, its soft rays usually few.  
   b. Maxillary not sheathed by the preorbital, or only partially covered by the edge of the latter; ventral with its accessory scale very small or wanting; pectoral without accessory scale; sheath at base of spinous dorsal little developed; vomer usually with teeth; opercle usually ending in a spine.  
   c. Precaudal vertebrae with transverse processes from the third or fourth to the last; ribs all but the last 1 to 4 sessile, inserted on the centra behind the transverse processes; anal spines 3; species silvery in color, the dorsal deeply notched, with 10 spines; vertebrae 10 + 15 = 25. CXLII, KUHLIDÆ, 1013.  
 cc. Precaudal vertebrae normal, anteriorly without transverse processes; all or most of the ribs inserted on the transverse processes when these are developed.  
   d. Anal spines 2 or 1; pseudobranchiæ small; preopercle with a hook-like spine below; vertebrae increased in number (30 to 46). Fresh-water fishes. . . . . CXLIII, PERCIDÆ, 1015.  
 dd. Anal spines 2, rarely 3; vertebrae 24 or 25; dorsal fin divided. Marine fishes . . . . . CXLIV, CHEILODIPTERIDÆ, 1105.

- ddd. Anal spines 3, never 2 nor 1; dorsal fin continuous or divided; vertebrae 24 to 35.
- e. Vomer, and usually palatines also, with teeth.
- f. Anal fin shorter than dorsal; head not everywhere covered with rough scales; postocular part of head not shortened.  
CXLVI, SERRANIDÆ, 1126.
- ff. Anal fin scarcely shorter than dorsal and similar to it; head and body everywhere covered with rough scales; body deep, compressed, the posterior part of head shortened.  
CXLVIII, PRIACANTHIDÆ, 1236.
- ee. Vomer without teeth; dorsal fin continuous; body deep, compressed..... CXLVII, LOBOTIDÆ, 1235.
- bb. Maxillary slipping for most of its length under the edge of the preorbital, which forms a more or less distinct sheath; ventrals with an accessory scale; opercle without spines; maxillary without supplemental bone; anal spines 3, rarely 2.
- g. Fishes carnivorous; intestines of moderate length; teeth in jaws not all incisor-like; vertebrae usually 24 or 25.
- h. Vomer with teeth, these sometimes very small; maxillary long.  
CXLIX, LUTIANIDÆ, 1241.
- hh. Vomer without teeth; palatines and tongue toothless.
- i. Teeth on sides of jaws not molar; maxillaries formed essentially as in the *Serranidæ*; preopercle mostly serrate.  
CL, HÆMULIDÆ, 1289.
- ii. Teeth on sides of jaws molar; maxillaries peculiar in form and in articulation; anterior teeth conical or else more or less incisor-like; preopercle entire... CL, SPARIDÆ, 1343.
- gg. Fishes herbivorous; intestinal canal elongate; anterior teeth in jaws incisor-like; no molars or canines; premaxillaries moderately protractile..... CLIV, KYPHOSIDÆ, 1380.
- aa. Mackerel-like fishes, with the caudal peduncle usually very slender, the fin widely forked, the scales various, usually not ctenoid; the dorsal spines various, anal fin long.
- j. Scales firm, linear, parchment-like; body compressed; bones of head rough; dorsal spines few; mouth small.  
CXXXVII, GRAMMICOLEPIDIDÆ, 973.
- jj. Scales not linear, mostly cycloid.
- k. Dorsal spines numerous, most of them produced in long filaments; pectorals very long... CXXXIV, NEMATISTIDÆ, 891.
- kk. Dorsal spines mostly low, not more than 2 of them filamentous.
- l. Dorsal fin very long, all the rays soft; skeleton soft.  
CXXXVI, ICOSTEIDÆ, 968.
- ll. Dorsal fin with 3 or more spines.
- m. Dorsal fin divided, the spines 6 to 12 in number.
- n. Scales weak, cycloid; jaws without canines.  
CXXXVIII, NOMEIDÆ, 948.
- nn. Scales ciliate; jaws with canines... CXXXVI, POMATOMIDÆ, 945.
- nnn. Scales firm, each with a median ridge; no canines.  
CXXXIII, STEINEGERIIDÆ, 960.
- mm. Dorsal spines 3 or 4, the fin not divided.
- o. Scales minute, body oblong, the shoulder girdle moderate.  
CXXXIV, CENTROLOPHIDÆ, 962.
- oo. Scales rather large, firm; body broad, ovate, the shoulder girdle very strong..... CXXXII, BRAMIDÆ, 956.
- BB. Body scaleless, smooth or armed with tubercles, prickles, or scattered bony plates.
- C. Breast with a sucking disk.
- D. Gill membrane free from the isthmus; no spinous dorsal; a large sucking disk between the ventral fins.  
CXCIX, GORIESCIDÆ, 2326.

- DD. Gill membranes joined to the isthmus; a sucking disk formed of the ventral fins.  
 E. Skin perfectly smooth; spinous dorsal not distinct. CLXXXIII, LIPARIDIDÆ, 2105.  
 EE. Skin with tubercles or spines, or else with a distinct spinous dorsal. CLXXXII, CYCLOPTERIDÆ, 2094.  
 CC. Breast without sucking disk.  
 F. Gill membranes broadly attached to the isthmus.  
 G. Ventrals completely united. CLXXXVIII, GOBIDÆ, 2188.  
 GG. Ventrals widely separated; body depressed; preopercle with a strong spine. CLXXXVII, CALLIONYMIDÆ, 2184.  
 FF. Gill membranes nearly or quite free from the isthmus.  
 H. Anal preceded by 2 free spines (these lost with age; connected by membranes in the very young). CXXV, CARANGIDÆ, 895.  
 HH. Anal without free spines.  
 I. Dorsal and anal fins followed by finlets. CXVIII, SCOMBRIDÆ, 863.  
 II. Dorsal and anal without finlets.  
 J. Suborbital with a bony stay; no free anal spines. CLXXIX, COTTIDÆ, 1879.  
 JJ. Suborbital without bony stay.  
 K. Mouth very large, nearly horizontal, the teeth sharp; no pseudo-branchiæ. CXCIII, CHIASMODONTIDÆ, 2291.  
 KK. Mouth large, nearly vertical; body compressed; preopercle armed with spines. CXCIV, TRICHODONTIDÆ, 2295.  
 AA. Gill openings small, behind, above, or below the pectoral fins, which are more or less pediculate.  
 L. Gill openings in or behind upper axil of pectorals; mouth small. CCXXIV, OGCOEPHALIDÆ, 2735.  
 LL. Gill openings in or behind lower axil of pectoral; mouth large.  
 M. Head compressed; no pseudo-branchiæ. CCXXII, ANTENNARIDÆ, 2715.  
 MM. Head depressed; pseudo-branchiæ present. CCXXI, LOMPIDÆ, 2713.

III.—VENTRAL FINS PRESENT, THORACIC OR JUGULAR, THE NUMBER OF RAYS NOT DEFINITELY 1, 5.

- A. Eyes unsymmetrical, both on the same side of head.  
 B. Eyes large, well separated; edge of preopercle usually evident. CCXIX, PLEURONECTIDÆ, 2602.  
 BB. Eyes small, very close together; edge of preopercle hidden by skin; mouth very small. CCXX, SOLEIDÆ, 262.  
 AA. Eyes symmetrical, one on each side of the head.  
 C. Ventral rays with or without spine, the number of soft rays more than 5.  
 D. Caudal fin wanting; scales spinous. CCXV, MACROURIDÆ, 2561.  
 DD. Caudal fin well developed.  
 E. Tail isocercal, the vertebrae progressively smaller to base of caudal; ventrals jugular; no spines in any of the fins.  
 F. Jaws and vomer with strong canines; second dorsal and anal deeply notched; no barbel. CCXIII, MERLUCCIDÆ, 2529.  
 FF. Jaws and vomer without distinct canines; chin usually with a barbel. CCXIV, GADIDÆ, 2531.  
 EE. Tail not isocercal, the last vertebra not reduced in size.  
 G. Ventral rays about 15; dorsal fin single, elevated. CXXX, LAMPRIDIDÆ, 953.  
 GG. Ventral rays I, 3 or I, 5; dorsal very high. CXXXI, PTERACLIDIDÆ, 955.  
 GGG. Ventral rays I, 6 to 1, 10; dorsal with spines.

- H. Vent anterior; dorsal spines 3 or 4; scales ctenoid.  
CV, APHREDODERIDÆ, 785.
- III. Vent normal.  
I. Chin with two long barbels, behind symphysis; dorsal continuous, with 5 spines .....CXVI, POLYMIXIDÆ, 851.
- II. Chin without barbels.  
J. Dorsal fin divided, the anterior part of a single slender spine; ventrals elongate.....CCXII, BREGMACEROTIDÆ, 2525.
- JJ. Dorsal fin divided, the anterior part of many spines.  
K. Body covered with firm serrated scales; anal spines 4; dorsal spines not elevated.....CXV, HOLOCENTRIDÆ, 845.
- KK. Body naked or covered with small scales, besides bony plates or warts.....CLXII, ZEIDÆ, 1659.
- KKK. Body uniformly covered with cycloid scales; dorsal spines mostly very high and filamentous.  
CXXIV, NEMATISTIDÆ, 894.
- JJJ. Dorsal fin continuous, its spines 2 to 8.  
L. Suborbitals narrow, not covering the cheeks.  
CXIV, BERYCIDÆ, 837.
- LL. Suborbitals very broad, covering the cheeks.  
CXIII, TRACHICHTHYIDÆ, 836.
- CC. Ventral fins with or without spine, the number of soft rays fewer than 5.  
M. Gill opening before the pectoral fin.  
N. Anal fin present; caudal fin not directed upward.  
O. Upper jaw not prolonged into a sword.  
P. Dorsal fin with some spines or simple rays.  
Q. Dorsal fin without soft rays, composed of spines only.  
CC, BLENNIIDÆ, 2344.
- QQ. Dorsal fin with soft rays anteriorly, with spines posteriorly; gill membranes joined to isthmus..CCVI, ZOARCIDÆ, 2455.
- QQQ. Dorsal fin of spines anteriorly, with soft rays posteriorly.  
R. Dorsal spines all separate and unconnected; body scaleless, naked, or with bony plates; ventral with a sharp spine.  
XCVIII, GASTEROSTEIDÆ, 742.
- RR. Dorsal spines connected by membrane.  
S. Suborbital with a bony stay, extending across the cheek, to or toward the preopercle, the cheek sometimes entirely covered with a coat of mail.  
T. Pectoral fin divided into 2 parts, 1 of them very long; head bony.....CLXXXVI, CEPHALACANTHIDÆ, 2182.
- TT. Pectoral fin not divided.  
U. Body entirely covered with an armor of bony plates; head bony.  
CLXXXI, AGONIDÆ, 2031.
- UU. Body naked, or more or less rough or scaly, not entirely covered by bony plates.  
V. Gill opening very small, not extending below upper edge of pectoral; skin everywhere prickly; head very large, bony above.....CLXXX, RHAMPHOCOTTIDÆ, 2029.
- VV. Gill opening large, extending downward nearly or quite to lowest pectoral ray.....CLXXIX, COTTIDÆ, 1879.
- SS. Suborbital without bony stay.  
W. Dorsal spines 2 to 4 only; head very broad, depressed; gills 3; gill membranes broadly united to the isthmus.  
x. Ventrals each a strong spine; teeth incisor-like; scales shagreen-like.....CLXVIII, TRIACANTHIDÆ, 1697.
- xx. Ventrals not reduced each to a single spine.  
CXCVIII, BATRACHOIDIDÆ, 2313.
- WW. Dorsal spines numerous; gills 4.  
X. Gill membranes separate, free from the isthmus.  
Y. Body greatly elongate; lower jaw with a slit at base to permit free motion; lips not fringed.



- Z. Soft dorsal and anal with a distinct lobe anteriorly, distinct from spinous part. ....CXIX, GEMPYLIDÆ, 877.
- ZZ. Soft dorsal and anal without anterior lobe, continuous with spinous part. ....CXX, LEPIDOPODIDÆ, 884.
- YY. Body moderately elongate; opercles and lips fringed; eyes superior. ....CXCVI, DACTYLOSCOPIDÆ, 2297.
- XX. Gill membranes broadly united, attached to the isthmus or not.
- x. Gill opening moderate or large. ....CC, BLENNIDÆ, 2344.
- xx. Gill openings very small, reduced to oblique slits before the pectoral fins. ....CCIII, CERDALIDÆ, 2448.
- PP. Dorsal fins of soft rays only.
- a. Breast with a large sucking dish between ventral fins. ....CXCIX, GOMIESOCIDÆ, 2326.
- aa. Breast without sucking disk.
- b. Body covered with a coat of mail; dorsal very short. ....CLXXXI, AGONIDÆ, 2031.
- bb. Body not mailed; dorsal many-rayed.
- c. Lateral line and base of dorsal beset with prickles; skeleton very soft; body compressed ...CXXXVI, ICOSTEIDÆ, 968.
- cc. Lateral line unarmed.
- d. Tail isocercal, the vertebral column pointed behind, the last vertebre very small; hypercoracoid not perforate; no pseudobranchiæ.
- e. Caudal fin present. ....CCXIV, GADIDÆ, 2531.
- ee. Caudal fin wanting. ....CCXV, MACROURIDÆ, 2561.
- dd. Tail not isocercal, truncate at base of caudal; hypercoracoid perforate.
- f. Gill membranes joined to the isthmus; pseudobranchiæ present.
- g. Ventral fins under shoulder girdle. ....CCVI, ZOARCIDÆ, 2455.
- gg. Ventral fins inserted below the eyes. ....CCVII, DEREPODICHTHYIDÆ, 2480.
- ff. Gill membranes free from the isthmus.
- h. Ventral fins inserted below or before the eyes; pseudobranchiæ generally well developed. ....CCVIII, OPHIDIIDÆ, 2481.
- hh. Ventral fins inserted below shoulder girdle; no pseudobranchiæ. ....CCXI, BROTLIDÆ, 2498.
- OO. Upper jaw prolonged into a bony sword; dorsal fin long and high; size large. ....CXXII, ISTIOPHORIDÆ, 890.
- NN. Anal fin wanting; caudal fin distorted or directed upward; body ribbon-like.
- i. Ventral fins each of a few slender rays. ....CCXVII, TRACHYPTERIDÆ, 2597.
- ii. Ventral fins each reduced to a long slender filament. ....CCXVI, KEGALECIDÆ, 2595.
- MM. Gill openings behind the pectoral fins.
- j. Gill openings above and behind pectorals; mouth small, low. ....CCXXIV, OGCOCYPHALIDÆ, 2735.
- jj. Gill openings below and behind pectorals; mouth large, nearly vertical. ....CCXXII, ANTENNARIIDÆ, 2715.

IV.—VENTRAL FINS WHOLLY WANTING.

- A. Promaxillary and maxillary wanting or grown fast to the palatines; body greatly elongate, eel-shaped; gill openings restricted to the sides; scales minute or wanting; scapular arch not attached to the skull. Eels.
- B. Gill openings not very far behind cranium; gape not inordinately distensible; gill arches 4 pairs.
- C. Gill openings well developed, leading to large interbranchial slits; tongue present; opercles and branchial bones well developed; scapular arch present.

- D. Skin covered with rudimentary embedded scales, usually linear in form, arranged in small groups, and placed obliquely at right angles to those of the neighboring groups; pectorals and vertical fins well developed, the latter confluent about the tail; lateral line present; posterior nostril in front of eyes; tongue with its margins free.
- E. Gill openings well separated; branchiostegals long, bent upward behind.
- F. Gill openings lateral and vertical; snout conic, the jaws not very heavy; gape longitudinal; lips thick; lower jaw projecting; teeth in cardiform bands on jaws and vomer; eggs minute.....XLIII, ANGUILLIDÆ, 346.
- FF. Gill openings horizontal, inferior.
- G. Snout very blunt, with very strong jaws; gape transverse; lips obsolete; teeth blunt, in 1 series, on jaws only.  
XLIV, SIMENCHELYIDÆ, 348.
- GG. Snout conical and slender, the jaws of moderate strength; gape lateral; lips obsolete; tongue but little developed; teeth acute, in bands on jaws and vomer.  
XLV, ILYOPHIDIDÆ, 349.
- EE. Gill openings inferior, very close together, apparently confluent; branchiostegal rays abbreviated behind; head conical; tongue small; posterior nostrils in front of eye.  
XLVI, SYNAPHOBRANCHIDÆ, 350.
- DD. Scales wholly wanting; eggs (so far as known) of moderate size, much as in ordinary fishes.
- H. Tip of tail with a more or less distinct fin, the dorsal and anal fins confluent around it; the tail sometimes ending in a long filament. Coloration almost always plain, brownish, blackish, or silvery, the fins often black-margined.
- I. Posterior nostril without tube, situated entirely above the upper lip.
- J. Tongue broad, largely free anteriorly and on sides; vomerine teeth moderate.
- K. Pectoral fins well developed; body not excessively elongate; lower jaw not projecting; anterior nostril remote from eye.....XLVII, LEPTOCEPHALIDÆ, 352.
- JJ. Tongue narrow, adnate to the floor of the mouth or only the tip slightly free; vomerine teeth well developed, sometimes enlarged.
- L. Jaws not attenuate and recurved at tip; gill openings well separated; anterior nostril remote from eye.
- M. Pectoral fins well developed; skin thick; skeleton firm; snout moderate; tail not ending in a filiform tip.  
XLVIII, MURENESOCIDÆ, 358.
- MM. Pectoral fins wholly wanting; snout and jaws much produced, the upper longer; jaws straight; skin thin and skeleton weak; tail ending in a filiform tip; gill openings small, subinferior; teeth sharp, subequal, recurved, a long series on the vomer. Deep-sea eels, soft in body, black in color.....XLIX, NETTASTOMATIDÆ, 361.
- LL. Jaws long and slender, tapering to a point, recurved at tip; nostrils large, both pairs close in front of eye; gill openings convergent forward, separate or confluent; pectorals and vertical fins well developed; membranes of fins thin, not enveloping the rays; skeleton well developed. Deep-sea eels.....L, NEMICHTHYIDÆ, 366.
- II. Posterior nostril close to the edge of the upper lip; tongue more or less fully adnate to the floor of the mouth; teeth subequal.....LI, MYRIDÆ, 370.
- III. Tip of tail without rays, projecting beyond the dorsal and anal fins (not filiform); posterior nostril on the edge of the

- upper lip; anterior nostril near tip of snout, usually in a small tube; tongue usually adnate to the floor of the mouth. Coloration frequently variegated.
- LII, OPHICHTHYIDÆ, 372.
- CC. Gill openings small, roundish, leading to restricted interbranchial slits; tongue wanting; pectoral fins (typically) wanting; opercles feebly developed; fourth gill arch modified, strengthened, and supporting pharyngeal jaws.
- N. Scapular arch obsolete or represented by cartilage; heart not far back; pectorals wanting; (skin thick; coloration often variegated) .....LIII, MURÆNIDÆ, 388.
- BB. Gill openings far behind cranium; gape of mouth inordinately distensible; gill arches 5 or 6 pairs; tail excessively long, tapering to a point.
- O. Distance from gill opening to vent much greater than that from tip of snout to gill opening.
- LIV, SACCOPHARYNGIDÆ, 405.
- OO. Distance from gill opening to vent much less than from tip of snout to gill opening..... LV, EURYPHARYNGIDÆ, 406.
- AA. Premaxillary and maxillary present, often immovably united to rest of cranium.
- P. Gill openings united in a single slit below throat; no pectoral fins; body eel-shaped.....XLI, SYMBRANCHIDÆ, 342.
- PP. Gill openings not united in a longitudinal slit.
- Q. Dorsal fin wanting; anal fin very long; vent near the head; caudal obsolete; body band-like.....XL, GYMNOTIDÆ, 340.
- QQ. Dorsal fin present.
- R. Body eel-shaped, contracted at the neck; the vertical fins confluent around the tail; premaxillary and maxillary immovably united to the skull...XLII, DERICHTHYIDÆ, 343.
- RR. Body eel-shaped, ending in a long filament, longer than rest of body.
- X. No anal or caudal fin.....CCXVIII, STYLEPHORIDÆ, 2601.
- XX. No caudal fin; anal present.....CCIV, PTILICHTHYIDÆ, 2451.
- RRR. Body not truly eel-shaped.
- S. Gill openings far behind pectoral fins; mouth oblique, very large; spinous dorsal represented by fleshy tentacles.
- CCXXIII, CERATHIDÆ, 2727.
- SS. Gill openings before pectoral fins.
- T. Gill membranes broadly united to the isthmus, restricting the gill openings to the sides.
- U. Snout tubular, bearing the short, toothless mouth at the end; body mailed.....CIII, SYNGNATHIDÆ, 760.
- UU. Snout not tubular.
- V. Breast without sucking disk.
- W. Dorsal fin single, of spines or undivided rays only.
- X. Jaws and vomer with coarse molar teeth.
- CCII, ANARHICHADIDÆ, 2445.
- XX. Jaws and vomer without molars.
- Y. Mouth nearly vertical; dorsal spines slender, rather high.
- CCI, CRYPTACANTHODIDÆ, 2442.
- YY. Mouth not nearly vertical; dorsal spines moderate or low, some or all of them usually pungent.....CC, BLENNIIDÆ, 2344.
- WW. Dorsal fins 2, the anterior of spines, the posterior of soft rays; body short and deep.
- Z. Spinous dorsal of 2 or 3 spines; scales rather large, rough or bony .....CLXIX, BALISTIDÆ, 1698.
- ZZ. Spinous dorsal of 1 or 2 spines; scales minute, rough, forming a velvety covering.....CLXX, MONACANTHIDÆ, 1712.
- WWW. Dorsal fin continuous, of soft rays only.
- a. Body oblong or elongate, the back not elevated; dorsal and anal joined to caudal.

- b. Pectoral rather narrow, the lower rays similar to the others.  
CCVI, ZOARCID.E, 2453.
- bb. Pectorals very broad, the lower rays procurrent and produced  
at tip ..... CLXXXIII, LIPARIDID.E, 2105.
- aa. Body short, not elongate; dorsal and anal free from caudal.
- c. Teeth in each jaw confluent into 1.
- d. Body compressed, rough ..... CLXXV, MOLID.E, 1752.
- dd. Body not compressed, spinous ..... CLXXIV, DIODONTID.E, 1712.
- cc. Teeth in each jaw confluent into 2.
- e. Back broadly rounded ..... CLXXXII, TETRAODONTID.E, 1726.
- ee. Back with a sharp median ridge.  
CLXXXIII, CANTHIGASTERID.E, 1740.
- ccc. Teeth separate; body enveloped in a bony box.  
CLXXI, OSTRACID.E, 1721.
- VV. Breast with a sucking disk.
- e. Skin perfectly smooth; dorsal continuous or slightly notched.  
CLXXXIII, LIPARIDID.E, 2105.
- ee. Skin more or less tubercular; dorsal usually divided.  
CLXXXII, CYCLOPTERID.E, 2091.
- TT. Gill membranes free from the isthmus.
- f. Vent at the throat.
- g. Vertical fins confluent; body elongate, almost eel-shaped.  
CCX, FIERASFERID.E, 2491.
- gg. Vertical fins separate; body oblong, scaly.  
XCIII, AMBLYOPSID.E, 702.
- ff. Vent posterior, not at the throat.
- h. Caudal fin wanting; body naked, greatly elongate.  
CXXI, TRICHURID.E, 888.
- hh. Caudal fin present.
- i. Upper jaw prolonged into a sword; size very large.  
CXXIII, XIPHIID.E, 893.
- ii. Upper jaw not prolonged into a sword.
- j. Belly with a series of bony scutes along its edge; body much  
compressed. .... LXI, CLUPEID.E, 417.
- jj. Belly not armed with scutes.
- k. Mouth inordinately large, formed like the mouth of a whale,  
with sharp teeth; no scales. .... LXXIV, CETOMIMID.E, 548.
- kk. Mouth not inordinately large, not peculiar in form.
- l. Body ovate, much compressed.
- m. Scales small, cycloid, silvery ..... CXXXV, STROMATEID.E, 964.
- mm. Scales wanting; caudal peduncle very slender.  
CXXXVI, ICOSTEID.E, 968.
- l. Body oblong or elongate, much longer than deep.
- n. Gill membranes broadly united; teeth present.
- o. Dorsal fin of spines only ..... CC, BLENNIID.E, 2344.
- oo. Dorsal fin of soft rays only; body eel-shaped.  
CCV, SCYTALINID.E, 2453.
- ooo. Dorsal fin single, the posterior half of soft rays, the anterior of  
spines; body elongate, covered with small scales.  
CC, BLENNIID.E, 2344.
- oooo. Dorsal fins 2, the anterior of slender spines, posterior soft, body  
naked ..... CLXXIX, COTTID.E, 1879.
- nn. Gill membranes separate.
- p. Jaws toothless, the lower jaw projecting; body scaly, with cross  
folds of skin ..... CX, AMMODYTID.E, 831.
- pp. Jaws with teeth.
- q. Body naked, without folds of skin; no pseudobranchiae.  
CCIX, LYCODAPODID.E, 2491.
- qq. Body with small scales; pseudobranchiae present; head with  
very large mucous pores; lower jaw very strong.  
CXXXVIIb, ZAPRORID.E, 2849.

GLOSSARY OF TECHNICAL TERMS.\*

- Abdomen.* Belly.
- Abdominal.* Pertaining to the belly; said of the ventral fins of fishes when inserted considerably behind the pectorals, the pelvic bones to which the ventral fins are attached having no connection with the shoulder girdle.
- Abortive.* Remaining or becoming imperfect.
- Actinosts.* A series of bones at the base of the pectoral rays.
- Acuminate.* Tapering gradually to a point.
- Acute.* Sharp-pointed.
- Adipose fin.* A peculiar, fleshy, fin-like projection behind the dorsal fin, on the backs of salmons, catfishes, etc.
- Adult.* A mature animal.
- Airbladder.* A sac filled with air, lying beneath the backbone of fishes, corresponding to the lungs of higher vertebrates.
- Alisphenoid.* A small bone on the anterior lateral wall of the brain case.
- Amphicoelian.* Double concave; said of vertebrae.
- Anadromous.* Running up; said of marine fishes which run up rivers to spawn.
- Anal.* Pertaining to the anus or vent.
- Anal fin.* The fin on the median line behind the vent, in fishes.
- Anchyloused.* Grown firmly together.
- Angular.* A small bone on the posterior end of the mandible.
- Antorse.* Turned forward.
- Anus.* The external opening of the intestine; the vent.
- Arterial bulb.* The muscular swelling, at the base of the great artery, in fishes.
- Articular.* The bone of the mandible supporting the dentary.
- Articulate.* Jointed.
- Atlas.* The first vertebra.
- Atrophy.* Nondevelopment.
- Attenuate.* Long and slender, as if drawn out.
- Auditory capsule.* The ventrolateral swelling of the skull.
- Barbel.* An elongated fleshy projection, usually about the head, in fishes.
- Basal.* Pertaining to the base; at or near the base.
- Basibranchials.* A lower median series of bones of the branchial arches.
- Basioccipital.* A median posterior ventral bone of the skull to which the atlas is attached.
- Basis cranii.* Formed by shelves of bone developed from the inner sides of the prootics which meet and form a roof to the myelome and a floor to the brain cavity.
- Bicolor.* Two-colored.
- Biocupid.* Having 2 points.
- Brachial ossicles.* Synonymous with actinosts, q. v.
- Branchiae.* Gills; respiratory organs of fishes.
- Branchial.* Pertaining to the gills.
- Branchiials.* Small bones at base of gill arches.
- Branchiostegals.* The bony rays supporting the branchiostegal membranes, under the head of a fish, below the opercular bones, and behind the lower jaw.

\* In the preparation of this Glossary the authors are indebted to Mr. Edwin Chapin Starks for valuable assistance.

- Bristle.** A stiff hair, or hair-like feather.
- Buccal.** Pertaining to the mouth.
- Caducous.** Falling off early.
- Cæcal.** Of the form of a blind sac.
- Cæcum.** An appendage of the form of a blind sac, connected with the alimentary canal at the posterior end of the stomach, or pylorus.
- Canines.** The teeth behind the incisors—the “eye-teeth;” in fishes, any conical teeth in the front part of the jaws, longer than the others.
- Cardiform (teeth).** Teeth coarse and sharp, like wool cards.
- Carinate.** Keeled; having a ridge along the middle line.
- Carotid.** The great artery running to the head.
- Carpus.** The wrist.
- Catadromous.** Running down; said of fresh-water species which run down to the sea to spawn.
- Caudal.** Pertaining to the tail.
- Caudal fin.** The fin on the tail of fishes and whales.
- Caudal peduncle.** The region between the anal and caudal fins in fishes.
- Cavernous.** Containing cavities, either empty or filled with a mucous secretion.
- Centrum.** The body of a vertebra.
- Cephalic fins.** Fins on the head of certain rays; a detached portion of the pectoral.
- Ceratobranchials.** Bones of the branchial arches just below their angle.
- Ceratohyal.** One of the hyoid bones.
- Chiasma.** Crossing of the fibers of the optic nerve.
- Chin.** The space between the rami of the lower jaw.
- Ciliated.** Fringed with eyelash-like projections.
- Cirri.** Fringes.
- Claspers.** Organs attached to the ventral fins in the male of sharks, skates, etc.
- Clavicle.** The collar bone, or lower anterior part of shoulder girdle, not entering into socket of arm.
- Compressed.** Flattened laterally.
- Condyle.** Articulating surface of a bone.
- Coracoid.** The principal bone of the shoulder girdle in fishes; otherwise a bone or cartilage on the ventral side, helping to form the arm socket. Synonymous with hypercoracoid, q. v.
- Cranial.** Pertaining to the cranium or skull.
- Ctenoid.** Rough-edged; said of scales when the posterior margin is minutely spinous or pectinated.
- Cycloid.** Smooth-edged; said of scales not ctenoid, but concentrically striate.
- Deciduous.** Temporary; falling off.
- Deurved.** Curved downward.
- Dentary.** The principal or anterior bone of the lower jaw or mandible, usually bearing the teeth.
- Dentate.** With tooth-like notches.
- Denticle.** A little tooth.
- Depressed.** Flattened vertically.
- Depth.** Vertical diameter (usually of the body of fishes).
- Dermal.** Pertaining to the skin.
- Diaphanous.** Translucent.
- Distal.** Remote from point of attachment.
- Dorsal.** Pertaining to the back.
- Dorsal fin.** The fin on the back of fishes.
- Emarginate.** Slightly forked or notched at the tip.
- Endoskeleton.** The skeleton proper; the inner bony framework of the body.
- Enteron.** The alimentary canal.
- Epibranchials.** The bones directly above the angle of the branchial arches.
- Epithyal.** One of the hyoid bones.
- Epipteurals.** Rays of bone attached to the ribs and anterior vertebrae usually touching the skin in the vicinity of the lateral line.

- Erectile.** Susceptible of being raised or erected.
- Ethmoid.** A median anterior bone of the skull.
- Exoccipitals.** Two bones of the skull, 1 on each side of the foramen magnum.
- Ezooskeleton.** Hard parts (scales, scutes) on the surface of the body.
- Erserted.** Projecting beyond the general level.
- Extralimital.** Beyond the limits (of this book).
- Facial.** Pertaining to the face.
- Falcate.** Scythe-shaped; long, narrow, and curved.
- Falciform.** Curved, like a scythe.
- Fauna.** The animals inhabiting any region, taken collectively.
- Femoral.** Pertaining to the femur, or proximal bone of the hinder leg.
- Filament.** Any slender or thread-like structure.
- Filiform.** Thread form.
- Fontanel.** An unossified space on top of head covered with membrane.
- Foramen.** A hole or opening.
- Foramen magnum.** The aperture in the posterior part of the skull for the passage of the spinal cord.
- Forehead.** Frontal curve of head.
- Forficat.** Deeply forked; scissors-like.
- Fossae (nasal).** Groves in which the nostrils open.
- Frontal bone.** Anterior bone of top of head, usually paired.
- Fulera.** Rudimentary spine-like projections extending on the anterior rays of the fins of ganoid fishes.
- Furcate.** Forked.
- Fusiform.** Spindle-shaped; tapering toward both ends, but rather more abruptly forward.
- Ganglion.** A nerve center.
- Ganoid.** Scales or plates of bone covered by enamel.
- Gape.** Opening of the mouth.
- Gill arches.** The bony arches to which the gills are attached.
- Gill openings.** Openings leading to or from the branchiae.
- Gill rakers.** A series of bony appendages, variously formed, along the inner edge of the anterior gill arch.
- Gills.** Organs for breathing the air contained in water.
- Glabrous.** Smooth.
- Glossohyal.** The tongue bone.
- Graduated (spines).** Progressively longer backward, the third being as much longer than the second as second is longer than first.
- Granulate.** Rough with small prominences.
- Gular.** Pertaining to the *gula*, or upper foreneck.
- Hæmal arch.** An arch under a hæmal spine for the passage of a blood vessel.
- Hæmal canal.** The series of hæmal arches as a whole.
- Hæmal spine.** The lowermost spine of a caudal vertebra, in fishes.
- Hæmopophyses.** Appendages on the lower side of abdominal vertebrae, in fishes.
- Height.** Vertical diameter.
- Heterocercal.** Said of the tail of a fish when unequal; the backbone evidently running into the upper lobe.
- Homocercal.** Said of the tail of a fish when not evidently unequal; the backbone apparently stopping at the middle of the base of the caudal fin.
- Humerus.** Bone of the upper arm.
- Hyoid.** Pertaining to the tongue.
- Hyoid apparatus.** Formed by a series of bones extending along the inner side of the mandible and supporting the tongue.
- Hyomandibular.** A bone by which the posterior end of the suspensorium is articulated with the skull; the supporting element of the suspensorium, the mandible, the hyoid apparatus, and the opercular apparatus.
- Hypercoracoid.** The upper of the 2 bones attached to the clavicle, indirectly bearing the pectoral fin.

- Hyppleural.** The modified last vertebra supporting the caudal fin.
- Hypobranchials.** Bones of the branchial arches below the ceratobranchials.
- Hypocoracoid.** The lower of the 2 bones attached to the clavicle behind.
- Hypohyals.** Small bones, usually 4, by which the respective sides of the hyoid apparatus are joined.
- Imbricate.** Overlapping, like shingles on a roof.
- Imperforate.** Not pierced through.
- Inarticulate.** Not jointed.
- Incisors.** The front or cutting teeth.
- Inferior pharyngeals.** Synonymous with pharyngeals, q. v.
- Infraoral.** Below the mouth.
- Interhamal spines.** Elements supporting the anal fin.
- Interhamals.** Bones to which anal rays are attached, in fishes.
- Interhyal.** Upper hyoid bone attached to hyomandibular.
- Intermusculars.** Synonym of epipleurals, q. v.
- Interneural spines.** Elements supporting the dorsal fins.
- Interspinous bones.** The interneurals and the interhamals.
- Intermaxillaries.** The premaxillaries; the bones forming the middle of the front part of the upper jaw, in fishes.
- Interneurals.** Bones to which dorsal rays are attached, in fishes.
- Interopercle.** Membrane bone between the preopercle and the branchiostegals.
- Interorbital.** Space between the eyes.
- Interspinals.** Bones to which fin rays are attached (in fishes); inserted between neural spines above and hamal spines below.
- Isocercal (tail).** Last vertebrae progressively smaller and ending in median line of caudal fin, as in the codfish.
- Jugular.** Pertaining to the lower throat; said of the ventral fins, when placed in advance of the attachment of the pectorals.
- Keeled.** Having a keel along the middle line.
- Lacustrine.** Living in lakes.
- Lamella.** Plate-like processes like those inside the bill of a duck.
- Larva.** An immature form, which must undergo change of appearance before becoming adult.
- Lateral.** To or toward the side.
- Lateral line.** A series of muciferous tubes forming a raised line along the sides of a fish.
- Lateral processes.** Synonym of parapophyses, q. v.
- Laterally.** Sidewise.
- Lunate.** Form of the new moon; having a broad and rather shallow fork.
- Mandible.** Under jaw.
- Maxilla, or maxillary.** Upper jaw.
- Maxillaries.** Outermost or hindmost bones of the upper jaw, in fishes; they are joined to the premaxillaries in front, and usually extend farther back than the latter.
- Mesethmoid.** Synonym of ethmoid, q. v.
- Mesopterygoid.** A bone of the suspensorium.
- Metapterygoid.** A bone of the suspensorium, or chain supporting the lower jaw.
- Molars.** The grinding teeth; posterior teeth in the jaw.
- Muciferous.** Producing or containing mucus.
- Myocomma.** A muscular band.
- Myodome.** Cavity under the brain cavity for the reception of the rectus muscles of the eye.
- Nape.** Upper part of neck, next to the occiput.
- Nares.** Nostrils, anterior and posterior.
- Nasal.** Pertaining to the nostrils.
- Nasal plate.** Plate in which the nostrils are inserted.
- Neural arch.** An opening through the base of the neural spine for the passage of the spinal cord.



- Neural canal.* The neural arches as a whole.  
*Neural processes.* Two plates rising vertically, 1 on each side of the centrum of the vertebra, which unite toward their ends and form a spine.  
*Neural spine.* The uppermost spine of a vertebra.  
*Nictitating membrane.* The third or inner eyelid of birds, sharks, etc.  
*Notochord.* A cellular chord which in the embryo precedes the vertebral column.  
*Nuchal.* Pertaining to the nape or *nucha*.  
*Obsolete.* Faintly marked; scarcely evident.  
*Obtuse.* Blunt.  
*Occipital.* Pertaining to the occiput.  
*Occipital condyle.* That part of the occipital bone modified to articulate with the atlas.  
*Occiput.* Back of the head.  
*Ocellate.* With eye-like spots, generally roundish and with a lighter border.  
*Oid* (suffix). Like; as *Percoid*, perch-like.  
*Opercle, or operculum.* Gill cover; the posterior membrane bone of the side of the head, in fishes.  
*Opercular bones.* Membrane bones of the side of the head, in fishes.  
*Opercular flap.* Prolongation of the upper posterior angle of the opercle, in sunfishes.  
*Opiethocelcian.* Concave behind only; said of vertebrae which connect by ball-and-socket joints.  
*Opiethotic.* A bone of the skull to which the lower limb of the post-temporal usually articulates.  
*Orbicular.* Nearly circular.  
*Orbit.* Eye socket.  
*Osseous.* Bony.  
*Ossicula auditus.* Bones of the ear, in fishes.  
*Osteology.* Study of bones.  
*Oviparous.* Producing eggs which are developed after exclusion from the body, as in all birds and most fishes.  
*Ovoviviparous.* Producing eggs which are hatched before exclusion, as in the dogfish and garter snake.  
*Ovum.* Egg.  
*Palate.* The roof of the mouth.  
*Palatines.* Membrane bones of the roof of the mouth, 1 on each side extending outward and backward from the vomer.  
*Palustrine.* Living in swamps.  
*Papilla.* A small fleshy projection.  
*Papillose.* Covered with papillae.  
*Parapophyses.* The lateral projections on some of the abdominal vertebrae to support ribs.  
*Parasphenoid.* Bone of roof of mouth behind the vomer. Synonym of prefrontal.  
*Parietal.* Bone of the side of head above.  
*Parotic process.* A posterior lateral process of the skull formed by the pterotic and opisthotic.  
*Pectinate.* Having teeth like a comb.  
*Pectoral.* Pertaining to the breast.  
*Pectoral fins.* The anterior or uppermost of the paired fins, in fishes, corresponding to the anterior limbs of the higher vertebrates.  
*Pelagic.* Living on or in the high seas.  
*Pelvic girdle.* The bones supporting the ventral fins or pelvises.  
*Pelvis.* The bones to which the hinder limbs (ventral fins in fishes) are attached.  
*Perforate.* Pierced through.  
*Peritoneum.* The membrane lining the abdominal cavity.

- Pharyngeal bones.** Bones behind the gills and at the beginning of the esophagus of fishes, of various forms, almost always provided with teeth; usually 1 pair below and 2 pairs above. They represent a fifth gill arch.
- Pharyngobranchials.** Upper elements of the branchial arches, usually bearing teeth.
- Pharyngognathous.** Having the lower pharyngeal bones united.
- Physoclistous.** Having the air bladder closed.
- Physostomous.** Having the air bladder connected by a tube with the alimentary canal.
- Pigment.** Coloring matter.
- Pineal body.** A small ganglion in the brain; a rudiment of an optic lobe, which in certain lizards (and in extinct forms) is connected with a third or median eye.
- Pituitary body.** A small ganglion in the brain.
- Plicate.** Folded; showing transverse folds or wrinkles.
- Plumbeous.** Lead colored; dull bluish gray.
- Polygamous.** Mating with more than 1 female.
- Postclavicle.** A ray composed of 1 or 2 bones attached to the inner upper surface of the clavicle and extending downward.
- Postorbital.** Behind the eye.
- Post-temporal.** The bone, in fishes, by which the shoulder girdle is suspended to the cranium.
- Præcoracoid.** A portion of coracoid more or less separated from the rest.
- Præcoracoid arch.** An arch in front of the coracoid in most soft-rayed fishes.
- Prefrontals.** Bones forming lateral projections at the anterior end of the skull.
- Premaxillaries.** The bones, 1 on either side, forming the front of the upper jaw in fishes. They are usually larger than the maxillaries and commonly bear most of the upper teeth.
- Premolars.** The small grinders; the teeth between the canines and the true molars.
- Preocular.** Before the eye.
- Preopercle.** The membrane bone lying in front of the opercle and more or less nearly parallel with it.
- Preorbital.** The large membrane bone before the eye, in fishes.
- Procalian.** Concave in front only.
- Procurrent (fin).** With the lower rays inserted progressively farther forward.
- Projectile.** Capable of being thrust forward.
- Prootic.** A bone forming an anterolateral ossification of the brain case.
- Protractile.** Capable of being drawn forward.
- Proximal.** Nearest.
- Pseudobranchia.** Small gills developed on the inner side of the opercle, near its junction with the preopercle.
- Pterotic.** A bone at the posterior lateral process of the skull.
- Pterygoids.** Bones of roof of mouth in fishes, behind the palatines.
- Pubic bones.** Same as pelvic bones, q. v.
- Pubis.** Anterior lower part of pelvis.
- Pulmonary.** Pertaining to the lungs.
- Punctate.** Dotted with points.
- Pyloric caeca.** Glandular appendages in the form of blind sacs opening into the alimentary canal of most fishes at the pylorus, or passage from the stomach to the intestine.
- Quadrate.** A bone of the suspensorium on which the mandible is hinged.
- Quincunx.** Set of 5 arranged alternately, thus
- \*           \*
- \*           \*
- Radius.** Outer bone of forearm.
- \*           \*

- Ray.** One of the cartilaginous rods which support the membrane of the fin of a fish.
- Recurved.** Curved upward.
- Reticulate.** Marked with a network of lines.
- Retrorse.** Turned backward.
- Rudimentary.** Undeveloped.
- Rugose.** Rough with wrinkles.
- Sacral.** Pertaining to the *sacrum*, or vertebrae of the pelvic region.
- Scapula.** Shoulder blade; in fishes, the bone of the shoulder girdle below the post-temporal.
- Scapular arch.** Shoulder girdle.
- Scute.** Any external bony or horny plate.
- Second dorsal.** The posterior or soft part of the dorsal fin, when the two parts are separated.
- Septum.** A thin partition.
- Serrate.** Notched, like a saw.
- Stemless.** Without a stem or peduncle.
- Setaceous.** Bristly.
- Setiform.** Bristle-like.
- Shaft.** Stiff axis of a quill.
- Shoulder girdle.** The bony girdle posterior to the head, to which the anterior limbs are attached (post-temporal, scapula, and coracoid or clavicle).
- Soft dorsal.** The posterior part of the dorsal fin in fishes, when composed of soft rays.
- Soft rays.** Fin rays which are articulate and usually branched.
- Spatulate.** Shaped like a spatula.
- Sphenoid.** Basal bone of skull.
- Sphenotic.** A lateral bone of the skull.
- Spine.** Any sharp projecting point; in fishes those fin rays which are unbranched, inarticulate, and usually, but not always, more or less stiffened.
- Spinous.** Stiff or composed of spines.
- Spinous dorsal.** The anterior part of the dorsal fin when composed of spinous rays.
- Spiracles.** Openings in the head and neck of some fishes and batrachians.
- Sicellate.** Star-like; with radiating ridges.
- Striate.** Striped or streaked.
- Sub** (in composition). Less than; somewhat; not quite; under, etc.
- Subcaudal.** Under the tail.
- Subopercle.** The bone immediately below the opercle (the suture connecting the two often hidden by scales).
- Suborbital.** Below the eye.
- Suborbital stay.** A bone extending from one of the suborbital bones in certain fishes, across the cheek, to or toward the preopercle.
- Subulate.** Awl-shaped.
- Superciliary.** Pertaining to the region of the eyebrow.
- Superior pharyngeals.** Synonym of pharyngobranchials, q. v.
- Supplemental maxillary.** A small bone lying along upper edge of the maxillary in some fishes.
- Supraclavicle.** A bone interposed between the clavicle and the post-temporal.
- Supraoccipital.** The bone at posterior part of skull in fishes, usually with a raised crest above.
- Supraoral.** Above the mouth.
- Supraorbital.** Above the eye.
- Suprascapular.** The post-temporal or bone by which the shoulder girdle in fishes is joined to the skull.
- Suspensorium.** The chain of bones from the hyomandibular to the palatine.
- Suspensory bones.** Bones by which the lower jaw, in fishes, is fastened to the skull.

*Suture.* The line of union of 2 bones, as in the skull.

*Symphysis.* Point of junction of the 2 parts of lower jaw; tip of chin.

*Symplectic.* The bone in fishes that keys together the hyomandibular and quadrate posteriorly.

*Synonym.* A different word having the same or a similar meaning.

*Synonymy.* A collection of different names for the same group, species, or thing; "A burden and a disgrace to science." (Coes.)

*Tail.* In fishes (usually), the part of the body posterior to the anal fin. (Often used more or less vaguely.)

*Temporal.* Pertaining to the region of the temples.

*Terete.* Cylindrical and tapering.

*Terminal.* At the end.

*Tessellated.* Marked with little checks or squares, like mosaic work.

*Thoracic.* Pertaining to the chest; ventral fins are thoracic when attached immediately below the pectorals, as in the perch, the pelvic bones being fastened to the shoulder girdle.

*Transverse.* Crosswise.

*Trenchant.* Compressed to a sharp edge.

*Truncate.* Abrupt, as if cut squarely off.

*Tubercle.* A small excrescence, like a pimple.

*Type (of a genus).* The species upon which was based the genus to which it belongs.

*Type (of a species).* The particular specimen upon which the original specific description was based.

*Type locality.* The particular place or locality at which the type specimen was collected.

*Typical.* Of a structure the most usual in a given group.

*Ultimate.* Last or farthest.

*Unicolor.* Of a single color.

*Vent.* The external opening of the alimentary canal.

*Ventral.* Pertaining to the abdomen.

*Ventral fins.* The paired fins behind or below the pectoral fins in fishes, corresponding to the posterior limbs in the higher vertebrates.

*Ventral plates.* In serpents or fishes, the row of plates along the belly between throat and vent.

*Ventricle.* One of the thick-walled chambers of the heart.

*Versatile.* Capable of being turned either way.

*Vertebra.* One of the bones of the spinal column.

*Vertical.* Up and down.

*Vertical fins.* The fins on the median line of the body; the dorsal, anal, and caudal fins.

*Villiform.* Said of the teeth of fishes when slender and crowded into velvety bands.

*Viscous.* Slimy.

*Viviparous.* Bringing forth living young.

*Vomer.* In fishes, the front part of the roof of the mouth; a bone lying immediately behind the premaxillaries.

*Zygapophysis.* Points of bone affording to the vertebrae more or less definite articulation with each other.

# INDEX.

	Page.		Page.
abacurus, Eleotris .....	2200	acanthias, Squalus .....	54
Abadejo .....	1184	Acanthidium .....	55
abboti, Syngnathus .....	764	pusillum .....	55
abboti, Osmerus mordax .....	524	Acanthinion .....	939
abbreviata, Chimera .....	95	rhomboides .....	942
abbreviatus, Nauclerus .....	900	acanthistius, Bodianus .....	1147
Abeona .....	1496	Acanthocheilodon .....	1682
aurora .....	1497	Acanthoclinus .....	2374
minima .....	1497	chaperi .....	2375
trowbridgii .....	1497	Acanthocottus .....	1970, 1971
aberrans, Hypoplectrus unicolor .....	1193	labradoricus .....	2001
Lilopropoma .....	1136	laticeps .....	1989
Perca .....	1136	mucosus .....	1975
abildgaardii, Scarus .....	1635	ocellatus .....	1976
Sparisoma .....	1635	patris .....	2009
Aboua .....	2240	profundorum .....	1991
chiquita .....	2241	sellaris .....	1998
de Mar .....	2195	variabilis .....	1975
de Rio .....	2236	virginianus .....	1976
etheostoma .....	2240	Acanthocybinae .....	865
lucretiae .....	2241	Acanthocybium .....	876
Abramidopsis .....	249	petus .....	877
Abramis .....	249	solandri .....	876
americanus .....	250	Acanthoderma .....	879, 1713
balteatus .....	239	temminkii .....	880
crysolenca .....	250	acanthoderma, Thyrsites .....	880
bosci .....	251	Acantholabrus exoletus .....	1576
gardoneus .....	251	Acantholebias .....	1866
lateralis .....	239	nubulosus .....	1872
leptosomus .....	250	Acantholepis .....	525
occidentalis .....	247	Acanthonotus .....	614
versicolor .....	250	nasus .....	615
Abudofduf .....	1500	acanthophorus, Serranus .....	1196
analogus .....	1503	Acanthopteri .....	779, 1241
declivifrons .....	1502	Acanthorhinus .....	59
rudis .....	1503	Acanthosoma .....	1753
saxatilis .....	1501	carinatum .....	1754
taurus .....	1503	Acanthostracion .....	1721, 1722, 1724
abyssiola, Raja .....	76, 2751	polygonus .....	1725
acadian, Cottus .....	2023	Acanthurus .....	1689
acadianus, Glyptocephalus .....	2657	aliaha .....	1694
Hemipteris .....	2023	bahianus .....	1693
Acantharchus .....	989	brooksii .....	1691
pomotis .....	989	broussoneti .....	1691
Acanthias .....	53	ceruleus .....	1691
americanus .....	54	chirurgus .....	1692
blainvilliei .....	57	glaucopterus .....	1694
sucklii .....	54	hepatus .....	1692
vulgaris .....	54	hirundo .....	1691

	Page.		Page.
<i>Acanthurus matoides</i> .....	1693	<i>Acipenser aleutensis</i> .....	104
<i>nigricans</i> .....	1692	<i>alexandri</i> .....	105
<i>phlebotomus</i> .....	1692	<i>anasimos</i> .....	106
<i>subarcticus</i> .....	1691	<i>anthracinus</i> .....	106
<i>tractus</i> .....	1693	<i>telaspis</i> .....	106
<i>tristegus</i> .....	1691	<i>atillus</i> .....	105
<i>zobra</i> .....	1691	<i>ayresi</i> .....	106
<i>Acanthopoma</i> .....	579	<i>burdi</i> .....	105
<i>asteropeleus</i> .....	579	<i>californicus</i> .....	104
<i>gonostomus</i> .....	579	<i>brevirostris</i> .....	106
<i>Acara aya</i> .....	1264	<i>buffalo</i> .....	106
<i>bartoni</i> .....	1510	<i>carbonarius</i> .....	106
<i>caeruleopunctatus</i> .....	1514	<i>caryi</i> .....	104
<i>fusco-maculata</i> .....	1540	<i>cataphractus</i> .....	107
<i>pinima</i> .....	1323	<i>cincinnati</i> .....	106
<i>pitamba</i> .....	1276	<i>copei</i> .....	106
<i>rectangularis</i> .....	1515	<i>dekayi</i> .....	106
<i>tetracanthus</i> .....	1540	<i>girardi</i> .....	105
<i>acara</i> , <i>Pristipoma pinima</i> .....	1323	<i>holbrookii</i> .....	105
<i>accensum</i> , <i>Plectropoma</i> .....	1193	<i>honeymani</i> .....	106
<i>accensum</i> , <i>Hypoplectrus</i> .....	1193	<i>hopctis</i> .....	106
<i>unicolor</i> .....	1193	<i>hospitus</i> .....	105
<i>accipiter</i> , <i>Podotheus</i> .....	2055	<i>kennicottii</i> .....	105
<i>acclivis</i> , <i>Larimus</i> .....	1422	<i>kirtlandi</i> .....	106
<i>Acedia</i> .....	2704, 2705, 2709, 2712	<i>levis</i> .....	106
<i>Acentrogobius</i> .....	2210	<i>lucymerus</i> .....	102
<i>Acentrolophus</i> .....	962	<i>lunari</i> .....	106
<i>maculosus</i> .....	963	<i>latirostris</i> .....	105
<i>acervum</i> , <i>Cybbium</i> .....	875	<i>lecontei</i> .....	105
<i>achigan</i> , <i>Bodianus</i> .....	1011	<i>lesueuri</i> .....	106
<i>Achirine</i> .....	2693	<i>lichtensteini</i> .....	105
<i>Achirophichthys</i> .....	387	<i>macrorhinus</i> .....	105
<i>typus</i> .....	388	<i>macrostomus</i> .....	106
<i>Achirus</i> .....	2693, 2695, 2700	<i>maculosus</i> .....	106
<i>achirus</i> .....	2695	<i>medrostris</i> .....	104, 2757
<i>brachialis</i> .....	2698	<i>megalaspis</i> .....	105
<i>comifer</i> .....	2698	<i>microrhynchus</i> .....	106
<i>fasciatus</i> .....	2700	<i>milberti</i> .....	105
<i>fimbriatus</i> .....	2700	<i>mitchilli</i> .....	105
<i>fischeri</i> .....	2699, 2700	<i>nerthinianus</i> .....	106
<i>fonsecensis</i> .....	2699	<i>obtusirostris</i> .....	106
<i>gronovii</i> .....	2696	<i>ohiensis</i> .....	100
<i>inscriptus</i> .....	2690	<i>oligopeltis</i> .....	105
<i>klonzingeri</i> .....	2697	<i>oxyrhynchus</i> .....	105
<i>lineatus</i> .....	2697, 2698, 2702	<i>paranusimos</i> .....	105
<i>maculipinnis</i> .....	2698	<i>platyrhynchus</i> .....	107
<i>mazatlanus</i> .....	2698	<i>platyrhinus</i> .....	106
<i>ornatus</i> .....	2709	<i>putnami</i> .....	104
<i>panamensis</i> .....	2702	<i>rainesqui</i> .....	106
<i>scutum</i> .....	2700	<i>raubi</i> .....	106
<i>achirus</i> , <i>Achirus</i> .....	2695	<i>rhynchaus</i> .....	106
<i>mollis</i> .....	2702	<i>richardsoni</i> .....	106
<i>Plenronectes</i> .....	2696	<i>rosarium</i> .....	106
<i>Solea</i> .....	2702	<i>rostellum</i> .....	106
<i>Acipenser</i> .....	103	<i>rubicundus</i> .....	106
<i>acatirostris</i> .....	104, 105	<i>rupertianus</i> .....	106
<i>agassizii</i> .....	105		

Page.		Page.		Page.
104	<i>Acipenser serotinus</i> .....	106	<i>acuminatus, Eques</i> .....	1487
105	<i>storeri</i> .....	105	<i>umbrosus</i> .....	1487
106	<i>sturio</i> .....	105	<i>Etrumeus</i> .....	419
106	<i>thompsoni</i> .....	105	<i>Myrichthys</i> .....	376
106	<i>transmontanus</i> .....	104	<i>Ophichthys</i> .....	377
105	<i>yarrelli</i> .....	105	<i>Ophisurus</i> .....	377
106	<i>Acipenseridae</i> .....	102	<i>Pareques</i> .....	1487
105	<i>acipenserinus, Agonus</i> .....	2062	<i>acuta, Loricaria</i> .....	158
104	<i>Aspidophorus</i> .....	2062	<i>Myliobatis</i> .....	89
106	<i>Paragonus</i> .....	2062	<i>Ophisoma</i> .....	356
106	<i>Phalangistes</i> .....	2062	<i>Perca</i> .....	1024
108	<i>Podotheicus</i> .....	2061, 2062	<i>acuticeps, Blennicottus</i> .....	2864
104	<i>ackleyi, Raja</i> .....	70	<i>Oligocottus</i> .....	2015, 2016
107	<i>Acomus</i> .....	173	<i>acutirostris, Acipenser</i> .....	104, 105
106	<i>griseus</i> .....	175	<i>Cerna</i> .....	1181
106	<i>Acoupa</i> .....	1403	<i>Corvina</i> .....	1437
106	<i>acoupa, Cestreus</i> .....	1404	<i>Lutjanus</i> .....	1259
105	<i>Cheilodipterus</i> .....	1404	<i>Pristis</i> .....	61
105	<i>Cynoscion</i> .....	1403	<i>Serranus</i> .....	1181
106	<i>Cynoscium</i> .....	1404	<i>Acutomentum</i> .....	1765, 1774, 1785
106	<i>Aerocheilus</i> .....	207	<i>macdonaldi</i> .....	1787
105	<i>aleutaceus</i> .....	208	<i>acutum, Hæmu'on</i> .....	1299
105	<i>acrolepis, Macrourus</i> .....	2585	<i>acutus, Exocoetus</i> .....	728
106	<i>Macrurus</i> .....	2585	<i>Fodiator</i> .....	728
106	<i>actinotus, Carcharhinus</i> .....	36	<i>Pseudoscarus</i> .....	1652
102	<i>Squalus</i> .....	36	<i>Scarus</i> .....	1652
106	<i>Actinopus carneus</i> .....	1692	<i>Acus</i> .....	774
105	<i>peruleatus</i> .....	1691	<i>acus, Sphyrana</i> .....	717
105	<i>fuscus</i> .....	1692	<i>Tylosurus</i> .....	716
106	<i>nigriculus</i> .....	1693	<i>Adinia</i> .....	660; 2830
105	<i>Acropocilia</i> .....	690	<i>dugesi</i> .....	661
105	<i>tridens</i> .....	690, 2833	<i>guatemalensis</i> .....	660, 2830
106	<i>Acrotidae</i> .....	2849	<i>multifasciata</i> .....	661
106	<i>Acrotina</i> .....	969	<i>pachycephala</i> .....	660, 2830
104, 2757	<i>Acrotus</i> .....	973, 2849	<i>adinia, Fundulus</i> .....	645
105	<i>willoughbyi</i> .....	973, 2849	<i>adirondacus, Salmo</i> .....	505
106	<i>Actinochir</i> .....	2114, 2116, 2127	<i>adiobe, Agosia</i> .....	310
105	<i>major</i> .....	2128	<i>Adonis</i> .....	2377
105	<i>aculeata, Mola</i> .....	1754	<i>cristatus</i> .....	2383
106	<i>aculeatus, Chelmon</i> .....	1671	<i>adscenionis, Epinephelus</i> .....	1152, 1154
106	<i>Chrysops</i> .....	1347	<i>Scomber</i> .....	927
100	<i>Olinus</i> .....	2433	<i>Trachinus</i> .....	1153
105	<i>Doryichthys</i> .....	773	<i>adspersus, Ctenolabrus</i> .....	1577
105	<i>Gasterosteus</i> .....	747, 2836	<i>Labrus</i> .....	1577
106	<i>cataphra-</i>		<i>Paralichthys</i> .....	2627
107	<i>tas</i> .....	750	<i>Pseudorhombus</i> .....	2627
106	<i>Haliutichthys</i> .....	2730	<i>Tautogolabrus</i> .....	1577
104	<i>Lophius</i> .....	2741	<i>adusta, Corvina</i> .....	1448
106	<i>Lumpenus</i> .....	2433	<i>Eupomacentrus</i> .....	1551
106	<i>Prognathodes</i> .....	1671	<i>Gobiesox</i> .....	1494
106	<i>Stenotomus</i> .....	1346	<i>Julidion</i> .....	1497
106	<i>Stichans</i> .....	2433	<i>Ophioscion</i> .....	1447
106	<i>acuminata, Gobius longissimus</i> .....	2230	<i>Pomacentrus</i> .....	1552
106	<i>Jenkinsia</i> .....	419	<i>Pseudojulis</i> .....	1603
106	<i>Maffona</i> .....	377	<i>adustus, Conesius</i> .....	325
106	<i>Selena</i> .....	1488	<i>Æglefinus</i> .....	2542

	Page.		Page.
<i>Æglefinus linnei</i> .....	2543	<i>affinis, Clinostomus</i> .....	239
<i>æglefinus, Gadus</i> .....	2543	<i>Cremnobates</i> .....	2372
<i>Melanogrammus</i> .....	2542, 2543	<i>Esox</i> .....	628
<i>Morrhua</i> .....	2543	<i>Exocoetus</i> .....	735, 2836
<i>Ælurichthys</i> .....	116	<i>Exonates</i> .....	2836
<i>ælurus, Amiurus</i> .....	140	<i>Fierasfer</i> .....	2495
<i>ævea, Cichla</i> .....	990	<i>Gambusia</i> .....	680
<i>æneolus, Notropis</i> .....	206	<i>Gila</i> .....	228
<i>æneus, Contrachus</i> .....	900	<i>Heros</i> .....	1529
<i>Cottus</i> .....	1973	<i>Heterandria</i> .....	681
<i>Myoxocephalus</i> .....	1972	<i>Hypoplectrus</i> .....	1193
<i>Pimelodus</i> .....	143	<i>unicolor</i> .....	1193
<i>Tetragonopterus</i> .....	333	<i>Isopisthus</i> .....	1399
<i>ænigmaticus, Teosteus</i> .....	972	<i>Lenciscus</i> .....	240
<i>æpyterus, Ammocetes</i> .....	11	<i>Lucania</i> .....	665
<i>æquatoris, Talismania</i> .....	456	<i>Pimelodus</i> .....	134
<i>Æquidens</i> .....	1513	<i>Scopelus</i> .....	571
<i>cærulopunctatus</i> .....	1514	<i>Stomias</i> .....	588
<i>æquidens, Cilius</i> .....	2202	<i>Synphobranchus</i> .....	351
<i>Eleotris</i> .....	2202	<i>Syngnathus</i> .....	769
<i>Prionodes</i> .....	1210	<i>Thynnus</i> .....	869
<i>Serranus</i> .....	1211	<i>afra, Murena</i> .....	396
<i>æquorens, Syngnathus</i> .....	774	<i>Perca</i> .....	1833
<i>æreus, Sebastodes</i> .....	1807	<i>africana, Scorpena</i> .....	1833
<i>æsculapius, Alepisaurus</i> .....	595	<i>afrum, Plectropoma</i> .....	1166
<i>æsopus, Boleosoma</i> .....	1057	<i>agassizii, Acipenser</i> .....	105
<i>æstivalis, Clupea</i> .....	427	<i>Alepocephalus</i> .....	453
<i>Gobio</i> .....	316	<i>Amphysticus</i> .....	1502
<i>Hybopsis</i> .....	316	<i>Anopus</i> .....	541
<i>marconis</i> .....	316	<i>Bathysaurus</i> .....	540
<i>Pomolobus</i> .....	426	<i>Brama</i> .....	959
<i>æstuarinus, Paralichthys</i> .....	2626	<i>Chlorophthalmus</i> .....	541
<i>æthalion, Citharichthys</i> .....	2673	<i>Chologaster</i> .....	704
<i>Hemirhombus</i> .....	2673	<i>Cratinus</i> .....	1188
<i>æthalarus, Carcharhinus</i> .....	40	<i>Cylindrostens</i> .....	111
<i>Æthopora</i> .....	565	<i>Dieromita</i> .....	2506
<i>ellulgens</i> .....	566	<i>Holconotus</i> .....	1506
<i>Inelda</i> .....	565	<i>Hyperprosopon</i> .....	1502
<i>Actobatina</i> .....	88	<i>Liparis</i> .....	2121
<i>Actobatis</i> .....	88	<i>Pimephales</i> .....	217
<i>Actobatus</i> .....	88	<i>Salmo</i> .....	507
<i>laticeps</i> .....	88, 2753	<i>Salvelinus fontinalis</i> .....	507
<i>narinari</i> .....	88	<i>Scorpena</i> .....	1840
<i>afer, Alphestes</i> .....	1164	<i>Serranus</i> .....	1189
<i>Epinephelus</i> .....	1165	<i>Xenichthys</i> .....	1287
<i>Gymnothorax</i> .....	305	<i>aggregatum, Ditrema</i> .....	1499
<i>aflno, Myctophum</i> .....	570	<i>aggregatus, Cymatogaster</i> .....	1498
<i>Plectropoma</i> .....	1193	<i>Micrometrus</i> .....	1499
<i>Siphostoma</i> .....	769, 770	<i>angilis, Gadus</i> .....	2534
<i>aflnis, Atherinops</i> .....	807	<i>agnus</i> .....	2306
<i>Achenopterus</i> .....	2371	<i>anoplus</i> .....	2308
<i>Carapus</i> .....	2497	<i>Agonide</i> .....	2031
<i>Caulolatilus</i> .....	2277	<i>Agonina</i> .....	2033
<i>Centropomus</i> .....	1124	<i>Agonomalus</i> .....	2036
<i>Cheilodipterus</i> .....	1113	<i>proboschialis</i> .....	2037
<i>Chimera</i> .....	95	<i>Agonopsis</i> .....	2068



Page.		Page.		Page.
239	Agonopsis chiloensis .....	2060	algula, Lachnolaimus .....	1580
2372	Agonostoma globiceps .....	821	Ailurichthys .....	116
628	nasutum .....	820	bagre .....	117
735, 2836	Agonostominae .....	809	oudouxii .....	118
2836	Agonostomus .....	818	filamentosus .....	118
2495	microps .....	820	gronovii .....	117
680	monticola .....	819	longispinis .....	119
228	nasutus .....	819, 2840, 2841	marinus .....	118
1529	percoides .....	819	nuchalis .....	117
681	Agonus .....	2064	panamensis .....	117
1193	acipenserinus .....	2062	pinnimaculatus .....	117
1193	anna .....	2043	ailurus, Pimelodus .....	140
1309	barkani .....	2044	Aimaras .....	330
240	entaphraetus .....	2065, 2067	aix, Pallasina .....	2050
665	chiloensis .....	2069	Aka nevo .....	1833
134	curlicus .....	2036	Aka soi .....	1830
571	decagonus .....	2053	Alabama Shad .....	2810
588	dodecaedron .....	2046	alabamae, Alosa .....	2810
351	gilberti .....	2060	Etheostoma .....	1095
769	japonicus .....	2036	whipplei .....	1095
869	laevigatus .....	2048	Notropis .....	298
396	niger .....	2069	Alalunga, Albacora .....	871
1833	quadricornis .....	2041	alalunga, Oreynus .....	871
1833	rostratus .....	2048	Alalunga .....	871
1166	spinosissimus .....	2054	alalunga, Germa .....	871
105	stegophthalmus .....	2036	alascanus, Ammodytes .....	832
453	vulsus .....	2068	Argyrosomus .....	2817
1502	Agosia .....	308, 309, 313	Sobastolobus .....	1761
541	adobe .....	310	Xenochirus .....	2081
540	chrysogaster .....	313	Alaska Blackfishes .....	620, 621
959	conesii .....	310	Codfish .....	2541
541	falcata .....	313	Dab .....	2645
704	shuswap .....	313	Dog salmon .....	478
1188	metallica .....	314	Greenfish .....	1869
111	nevadensis .....	310	Stickleback .....	749
2506	novemradiata .....	312	Alaskan Pollacks .....	2535
1506	nubila .....	311, 312	alatinga, Scomber .....	871
1502	carringtonii .....	311	alatus, Arius .....	125
2121	oscula .....	309	Lampamycetus .....	559
217	shuswap .....	313	Mugil .....	733
507	umatilla .....	313	Prionotus .....	2159
507	volifera .....	212	Alausa .....	427
1840	yarrowi .....	309	californica .....	423
1189	agua-bonita, Salmo iridens .....	503	striata .....	431
1287	mykiss .....	504	Alausella .....	424
1499	Aguaji .....	1174, 1177	alba, Rogenia .....	422
1498	aguaji, Trisotropis .....	1175	Albacora .....	869
1499	Aguja Blanca .....	892	albacaga .....	871
2534	de Casta .....	715, 892	thynnus .....	870
2306	de Paladar .....	892	albacora, Thynnus .....	871
2308	Prleta .....	891	Albacore, Great .....	870
2031	Voladora .....	891	Long-finned .....	871
2033	Agujon .....	714, 715, 716	Albacores .....	870
2036	Aguilha, Peixe .....	711	albacore Scomber .....	870
2037	A dia .....	370	Albatrossia .....	2573
2068	egmontis .....	370	pectoralis .....	2573

	Page.		Page.
albatrossis, Osmerus .....	2823	Alburnollus porcobromus .....	295
albeolus, Notropis megalops....	259, 283, 284	rubrifrons .....	295
albescens, Echeueis .....	2272	sinus .....	267
Remora .....	2272	umbratilis .....	299
albicans, Bagrus .....	124	Alburnops .....	254, 256, 261
Schadeichthys .....	124, 2760	blonnius .....	262
Tachisurus .....	124	heterodon .....	261
albicauda, Echeueis .....	2269	illecebrosus .....	269
albidactylus, Exocoetus .....	739	longirostris .....	267
albidum, Hamulon .....	1299	nubilus .....	215
Moxostoma .....	192	plumbeolus .....	283
albidus, Amirus .....	138	saludanus .....	270
Gadus .....	2531	shumardi .....	268
Ictalurus .....	138	taurocephalus .....	253
Labrax .....	1132	Alburnus amabilis .....	291
Osmerus .....	538	americanus .....	1475
Pimelodus .....	132, 138	dilectus .....	294
Ptychostomus .....	192	formosus .....	280
Tetrapturus .....	892	lepidulus .....	294
albigutta, Cathetostoma .....	2313	lineolatus .....	263
Kathetostoma .....	2312	megalops .....	291
albiguttus, Paralichthys .....	2631	nitidus .....	293
albirostre, Siphostoma .....	772	oligaepis .....	294
albirostris, Corythoichthys .....	772, 2838	rubellus .....	293
Prionotus .....	2163	rubrifrons .....	295
Syngnathus .....	772	socius .....	292
albolineatus, Fundulus .....	649	umbratilis .....	299
albomaculatus, Paralabrax .....	1197	zonatus .....	285
Serranus .....	1197	alburnus, Centropomus .....	1475
Albramis oligaepris .....	294	Monticirrhus .....	1475
Albula .....	411, 2807	Perca .....	1475
conorynchus .....	411	Umbrina .....	1475
erythrocheilos .....	412	albus, Atherinichthys .....	2839
fosteri .....	412	Centropomus .....	1135
goreensis .....	412	Cestrens .....	1411
neoguinaica .....	412	Coregonus .....	466
parva .....	411	Cynoscion .....	1411
rostrata .....	412	Gymnotus .....	340
seminuda .....	412	Lepisosteus .....	110
vulps .....	411	Otolithus .....	1411
albula, Mugil .....	812, 2841	Ptychostomus .....	191
Albulidae .....	410	Aleidea .....	1886
albulus, Bryttus .....	1067	thoburni .....	1887
Lepomis .....	1007	Aldrovandia .....	608, 2826
album, Hamulon .....	1295, 1296	goodei .....	610
Moxostoma .....	191	gracilis .....	610
Alburnellus .....	254	macrochir .....	609
altipinnis .....	287	pallida .....	611
amabilis .....	291	rostrata .....	609
amoenus .....	206	Alecrin .....	32
argo .....	294	Alectis .....	931
jaculus .....	293	ciliaris .....	931
jemezianus .....	294	crinitus .....	932
matutinus .....	301	Alectrias .....	2869
megalops .....	291	alectrolophus, Anoplarchus ..	2421, 2422, 2869
micropteryx .....	297	Blennius .....	2422
		Centronotus .....	2422

Page.		Page.		Page.
295	<i>Alectrolophus</i> , Gunnellus.....	2422	<i>Algansea sallæi</i> .....	212
295	<i>Aledon</i> .....	1753	<i>tarascorum</i> .....	2796
267	<i>capensis</i> .....	1754	<i>tincella</i> .....	211, 2796
299	<i>storeri</i> .....	1754	<i>alger.ensis</i> , <i>Gasterosteus</i> .....	748
4, 256, 261	<i>Alepos</i> .....	915	<i>Algoma</i> .....	212
262	<i>Alepidosaurus</i> .....	594	<i>amara</i> .....	215
261	( <i>Caulopus borealis</i> ).....	597	<i>fluvialis</i> .....	215
269	( <i>Caulopus poeyi</i> ).....	596	<i>alia</i> , <i>Labrus fauogæ</i> .....	1570
267	( <i>Caulopus serra</i> ).....	597	<i>allala</i> , <i>Acanthurus</i> .....	1694
215	<i>alepidotum</i> , <i>Gobiosoma</i> .....	2259	<i>Teuthis</i> .....	1693
283	<i>alepidotus</i> , <i>Chætodon</i> .....	966	<i>alicæ</i> , <i>Leuciscus</i> .....	236
270	<i>Derepodichthys</i> .....	2480	<i>Squalius</i> .....	236
268	<i>Gobius</i> .....	2259	<i>altiotulus</i> , <i>Trachurus</i> .....	904
253	<i>Lucioblennius</i> .....	2404	<i>Aliflonghi</i> .....	871
291	<i>Rhombus</i> .....	966	<i>alipes</i> , <i>Salmo</i> .....	509
1475	<i>Stromateus</i> .....	966	<i>Salvelinus alpinus</i> .....	509
294	<i>Alepisauridæ</i> .....	593	<i>alleganiensis</i> , <i>Salmo</i> .....	507
280	<i>Alepisaurus</i> .....	594, 2826	<i>alleterata</i> , <i>Gymnosarda</i> .....	869
294	<i>æsculapinus</i> .....	595	<i>alleteratus</i> , <i>Scomber</i> .....	869
263	<i>ativelis</i> .....	596	<i>allidus</i> , <i>Merluccius</i> .....	2531
291	<i>azureus</i> .....	595	<i>Alligator Gar</i> .....	111
293	<i>borealis</i> .....	596	<i>Allinectes</i> .....	2866
294	<i>fexox</i> .....	595	<i>alliteratus</i> , <i>Ethynnus</i> .....	869
293	<i>serra</i> .....	597	<i>Orcynus</i> .....	869
295	<i>Alepocephalidæ</i> .....	451	<i>All-Mouth</i> .....	2713
292	<i>Alepocephalus</i> .....	452	<i>Allochir</i> .....	2129, 2131, 2135
299	<i>agassizii</i> .....	453	<i>Allosomus</i> .....	467, 473
285	<i>bairdii</i> .....	454	<i>Alurus</i> .....	2129, 2131, 2136, 2866
1475	<i>macropterus</i> .....	458	<i>alm:ida</i> , <i>Belone</i> .....	715
1475	<i>productus</i> .....	452	<i>Tylosurus</i> .....	715
1475	<i>tenobrosus</i> .....	453	<i>Almejero</i> , <i>Mojarra</i> .....	1294
1475	<i>Aleposomus</i> .....	459	<i>Alopecias vulpes</i> .....	45, 46
2839	<i>copel</i> .....	459	<i>alopecias</i> , <i>Squalus</i> .....	46
1135	<i>aleutensis</i> , <i>Acipenser</i> .....	104	<i>Alopias</i> .....	45
1411	<i>Lyconectes</i> .....	2444	<i>macrourus</i> .....	46
466	<i>Aleutera</i> .....	2860	<i>vulpes</i> .....	45
1411	<i>alentianus</i> , <i>Sebastes</i> .....	1795	<i>Alopiidæ</i> .....	45
340	<i>alantica</i> , <i>Raja</i> .....	75; 2751	<i>Alosa</i> .....	427, 2810
110	<i>alenticus</i> , <i>Cottus</i> .....	1957	<i>alabamæ</i> .....	2810
1411	<i>Alewife</i> .....	426	<i>apicalis</i> .....	429
191	<i>Alewives</i> .....	424	<i>bishopi</i> .....	430
1886	<i>alexandri</i> , <i>Acipenser</i> .....	105	<i>cyanonoton</i> .....	427
1887	<i>alexandrini</i> , <i>Orthragoriscus</i> .....	1754	<i>lineata</i> .....	426
608, 2826	<i>Alexurus</i> .....	2202	<i>menbaden</i> .....	432
610	<i>armigor</i> .....	2203	<i>præstabilis</i> .....	428
610	<i>Alfione</i> .....	1507	<i>sapidissima</i> .....	427, 428, 2810
609	<i>Alfoncino</i> .....	1107	<i>teres</i> .....	420
611	<i>Alfonsin a Casta Cumpurida</i> .....	844	<i>alosoites</i> , <i>Amphiodon</i> .....	413
609	<i>Larga</i> .....	844	<i>Ilodon</i> .....	413
32	<i>Alfonsines</i> .....	844	<i>Alphestes</i> .....	1164
931	<i>Algansea</i> .....	211	<i>afer</i> .....	1164
931	<i>antica</i> .....	245	<i>chloropterus</i> .....	2854
932	<i>bicolor</i> .....	245	<i>multiguttatus</i> .....	1165
2869	<i>dugesi</i> .....	211	<i>alpinus</i> , <i>Salmo</i> .....	509, 514
2422, 2869	<i>formosa</i> .....	246	<i>Salmo, Æivalis</i> .....	509
2422	<i>obesa</i> .....	246	<i>Salvelinus</i> .....	508

	Page.		Page.
<i>alpinus</i> , <i>Salvelinus alpinus</i> .....	509	<i>Alvarius</i> .....	1099
<i>arcturus</i> .....	510	<i>fonticola</i> .....	1105
<i>aureolus</i> .....	511	<i>lateralis</i> .....	1099
<i>stagnalis</i> .....	510	<i>alveata</i> , <i>Trygonorhina</i> .....	65
<i>Alpismaris</i> .....	533	<i>alvordii</i> , <i>Cottus</i> .....	1952
<i>risso</i> .....	537	<i>Alvordius</i> .....	1028, 1029, 1030
<i>alta</i> , <i>Chola</i> .....	322	<i>aspro</i> .....	1033
<i>Macdonaldia</i> .....	2826	<i>crassus</i> .....	1034
<i>alter</i> , <i>Atinga minor orbicularis</i> .....	1749	<i>evides</i> .....	1037
<i>alternans</i> , <i>Scarus</i> .....	1651	<i>macrocephalus</i> .....	1031
<i>alternata</i> , <i>Perca mitchilli</i> .....	1133	<i>maculatus</i> .....	1032, 1034
<i>alticolus</i> , <i>Catostomus</i> .....	179	<i>nevisensis</i> .....	1034
<i>Alticus</i> .....	2390	<i>phoxocephalus</i> .....	1031
<i>altifrons</i> , <i>Heros</i> .....	1538	<i>spillmani</i> .....	1039
<i>altipinna</i> , <i>Bolone</i> .....	717	<i>variatus</i> .....	1034
<i>altipinnis</i> , <i>Alburnellus</i> .....	287	<i>Alysia</i> .....	568
<i>Micropogon</i> .....	1464	<i>loricata</i> .....	569
<i>Minnilus</i> .....	287	<i>amabilis</i> , <i>Alburnellus</i> .....	291
<i>Notropis</i> .....	287	<i>Alburnus</i> .....	291
<i>altivelis</i> , <i>Alepisaurus</i> .....	596	<i>Minnilus</i> .....	291
<i>Anchenopterus</i> .....	2370	<i>Notropis</i> .....	291
<i>Cremnolates</i> .....	2371	<i>amara</i> , <i>Algoma</i> .....	215
<i>Sebastolobus</i> .....	1763	<i>Hybognathus</i> .....	215
<i>altus</i> , <i>Bubalichthys</i> .....	165	<i>Auarilla</i> , <i>Chopa</i> .....	1386
<i>Chorinomus</i> .....	899	<i>Gnativore</i> .....	1144, 1145
<i>Hudsonius</i> .....	322	<i>Salmonete</i> .....	859
<i>Hybopsis</i> .....	321	<i>Amarillas</i> , <i>Mojarra de las Aletas</i> .....	1376
<i>Oligoplites</i> .....	899	<i>Amarillo</i> , <i>Cibi</i> .....	919
<i>Priacanthus</i> .....	1240	<i>Purgo</i> .....	1200
<i>Pseudopriacanthus</i> .....	1239	<i>Ronco</i> .....	1303
<i>alusis</i> , <i>Muraena</i> .....	403	<i>amarus</i> , <i>Hudsonius</i> .....	270
<i>aluta</i> , <i>Bairdiella</i> .....	1437	<i>Hybognathus</i> .....	215
<i>Sciæna</i> .....	1438	<i>Notropis hudsonius</i> .....	270
<i>alutaceus</i> , <i>Aerochellus</i> .....	208	<i>amazonica</i> , <i>Sciæna</i> .....	1419
<i>Alutarius amphacanthus</i> .....	1720	<i>amazonicus</i> , <i>Johnius</i> .....	1419
<i>maeracanthus</i> .....	1720	<i>ambassis</i> , <i>Sargus</i> .....	1346
<i>obliteratus</i> .....	1720	<i>Amber-fish</i> , <i>Great</i> .....	903
<i>Alutera</i> .....	1717, 1718, 1720	<i>Amber-fishes</i> .....	901
<i>cinerea</i> .....	1720	<i>Amber Jack</i> .....	903
<i>cuspicanda</i> .....	1718	<i>ambiguus</i> , <i>Lutjanus</i> .....	1272
<i>guntheriana</i> .....	1720	<i>Merluccius</i> .....	2530
<i>monoceros</i> .....	1720, 2860	<i>Mesopriou</i> .....	1272
<i>pieurata</i> .....	1719	<i>Neomanis</i> .....	1271
<i>punctata</i> .....	1718, 1719	<i>Amblodon</i> .....	1483
<i>schoepfi</i> .....	1718	<i>bubalus</i> .....	165
<i>scripta</i> .....	1719	<i>concinuus</i> .....	1484
<i>Aluteres</i> .....	1717	<i>grunniens</i> .....	1484
<i>berardi</i> .....	1720	<i>lineatus</i> .....	1484
<i>pareva</i> .....	1719	<i>neglectus</i> .....	1484
<i>Aluterus anginosus</i> .....	1720	<i>niger</i> .....	169
<i>eutifrons</i> .....	1718	<i>saturnus</i> .....	1456
<i>holbrookii</i> .....	1718	<i>Ambloplites</i> .....	989
<i>venosus</i> .....	1719	<i>interruptus</i> .....	991
<i>alutus</i> , <i>Apogon</i> .....	1110	<i>pomotis</i> .....	989
<i>Apogonichthys</i> .....	1110	<i>rupestris</i> .....	990
<i>Sebastodes</i> .....	1700	<i>cavifrons</i> .....	990



Page.		Page.		Page.
1099	amblops, <i>Ceratiichthys</i> .....	321	americana, <i>Stilbo</i> .....	250
1105	<i>Hybopsis</i> .....	320	<i>Tantoga</i> .....	1579
1099	<i>Nocomis</i> .....	321	americanus, <i>Abramis</i> .....	250
65	<i>Rutilus</i> .....	321	<i>Acanthias</i> .....	54
1952	<i>Amblygobius</i> .....	2210	<i>Alburnus</i> .....	1475
8, 1029, 1030	<i>Amblyopsidae</i> .....	702	<i>Ammolytes</i> .....	833
1033	<i>Amblyopsis</i> .....	706	<i>Amphiprion</i> .....	1139
1034	<i>spelaus</i> .....	706	<i>Apogonichthys</i> .....	1107
1037	<i>amblyopsis</i> , <i>Culius</i> .....	2200	<i>Balistes</i> .....	1707
1031	<i>Electris</i> .....	2199, 2200	<i>Blennius</i> .....	2457
1032, 1034	<i>Amblyopus brasiliensis</i> .....	2264	<i>Carcharias</i> .....	47
1034	<i>brevis</i> .....	2263	<i>Cyprinus</i> .....	250, 251, 1475
1031	<i>mexicanus</i> .....	2264	<i>Enehelyopus</i> .....	2457, 2555
1039	<i>pernunnus</i> .....	2265	<i>Eques</i> .....	1490
1034	<i>sagitta</i> .....	2263	<i>Esox</i> .....	626
568	<i>Amblypomacentrus</i> .....	1549	<i>lucius</i> .....	626
569	<i>amblyrhynchus</i> , <i>Caranx</i> .....	913	<i>Hemitripterus</i> .....	2023
291	<i>Hemleuranx</i> .....	912	<i>Hippoglossus</i> .....	2612
291	<i>Amblyscelon</i> .....	1420, 1421	<i>Histiophorus</i> .....	891
291	<i>argenteus</i> .....	1421	<i>Labrax</i> .....	1135
201	<i>amboynensis</i> , <i>Balistes</i> .....	1704	<i>Labrus</i> .....	1579
215	<i>Ameiurus</i> .....	135, 136, 139	<i>Leucosomus</i> .....	250
215	<i>bolli</i> .....	140	<i>Lophius</i> .....	2714
1386	<i>catus</i> .....	138	<i>Lucius</i> .....	626
1144, 1145	<i>dugesi</i> .....	138	<i>Menticirrhus</i> .....	1474
859	<i>dugei</i> .....	2789	<i>Notemigonus</i> .....	251
1376	<i>erobennus</i> .....	139	<i>Odontaspis</i> .....	47
919	<i>lacustris</i> .....	137	<i>Petromyzon</i> .....	10
1260	<i>lupus</i> .....	137	<i>Phycis</i> .....	2555
1303	<i>marmoratus</i> .....	141	<i>Platycephalus</i> .....	2029
270	<i>melas</i> .....	141	<i>Pleuronectes</i> .....	2647
215	<i>mispilliensis</i> .....	141	<i>Polynemus</i> .....	830
270	<i>natalis</i> .....	139	<i>Polyprion</i> .....	1139
1419	<i>nebulosus</i> .....	140	<i>Pseudopleuronectes</i> .....	2647
1419	<i>catulus</i> .....	141	<i>Squalus</i> .....	47
1346	<i>mauratus</i> .....	141	<i>amethystinus</i> , <i>Salmo</i> .....	505
903	<i>nigrilabris</i> .....	142, 2789	<i>amethystinus-punctatus</i> , <i>Mauro-</i>	
901	<i>okcechobensis</i> .....	138	<i>licus</i> .....	577
903	<i>platycephalus</i> .....	142	<i>Amia</i> .....	112, 1106
1272	<i>vulgaris</i> .....	140	<i>calva</i> .....	113
2530	<i>xanthocephalus</i> .....	141	<i>canina</i> .....	113
1272	American Eel .....	348	<i>cinerea</i> .....	113
1271	Perch .....	1023	<i>immaculata</i> .....	411
1483	Pike Perches .....	1020	<i>lentiginosa</i> .....	113
165	Shad .....	427	<i>marmorata</i> .....	113
1484	Smelt .....	523	<i>occellicauda</i> .....	113
1484	Sole .....	2700	<i>occidentalis</i> .....	113
1484	Soles .....	2693	<i>ornata</i> .....	113
1484	americana, <i>Cherna</i> .....	1160	<i>pliquottii</i> .....	113
169	<i>Lucioperca</i> .....	1022	<i>reticulata</i> .....	113
1456	<i>Manta</i> .....	93	<i>retrosella</i> .....	1109
989	<i>Morone</i> .....	1134, 1135	<i>subcaerulea</i> .....	113
991	<i>Morrhua</i> .....	2540, 2541	<i>thompsoni</i> .....	113
989	<i>Perca</i> .....	1024, 1135	<i>Amia viridis</i> .....	113
990	<i>Raia</i> .....	69	<i>Amiatus</i> .....	113
990	<i>Scorpana</i> .....	2023	<i>Amlichthys</i> .....	1113
			<i>dapterus</i> .....	1113

	Page.		Page.
<i>Amiidae</i> .....	112	<i>Ammodytes vittatus</i> .....	833
<i>Amitra</i> .....	2138	<i>Ammodytidae</i> .....	831
<i>liparina</i> .....	2138, 2139	<i>Ammodytoidei</i> .....	781, 831
<i>Amitrichthyis</i> .....	2139, 2141	<i>Ammopleurops</i> .....	2701
<i>Amitrinae</i> .....	2106	<i>amoenus</i> , <i>Albumellus</i> .....	296
<i>Amiurus</i> .....	135	<i>Notropis</i> .....	296
<i>alurus</i> .....	140	<i>Amore guaco</i> .....	2259
<i>albidus</i> .....	138	<i>pixuma</i> .....	2261
<i>bolli</i> .....	140	<i>amorea</i> , <i>Gobius</i> .....	2261
<i>borealis</i> .....	137	<i>Amorphocephalus</i> .....	1617
<i>brachyacanthus</i> .....	141	<i>granulatus</i> .....	1619
<i>brunneus</i> .....	142	<i>amphacanthus</i> , <i>alutarius</i> .....	1720
<i>catus</i> .....	141	<i>Amphiodon</i> .....	412, 413
<i>caulifurcatus</i> .....	135	<i>alosoidea</i> .....	413
<i>cragini</i> .....	141	<i>amphiodon</i> , <i>Hyodon</i> .....	413
<i>furcatus</i> .....	134	<i>Amphioxii</i> .....	2
<i>lophius</i> .....	138	<i>Amphioxys</i> .....	3
<i>meridionalis</i> .....	135	<i>lanceolatus</i> .....	3
<i>natalis analis</i> .....	140	<i>amphioxys</i> , <i>Monacanthus</i> .....	1717
<i>nigrilabris</i> .....	142	<i>Pseudomonacanthus</i> .....	1717
<i>niveiventris</i> .....	138	<i>Amphiprion americanus</i> .....	1139
<i>obesus</i> .....	141	<i>matejuelo</i> .....	849
<i>ponderosus</i> .....	137	<i>Amphistichus</i> .....	1503
<i>prosthistius</i> .....	139	<i>agassizi</i> .....	1502
<i>pullus</i> .....	141	<i>argenteus</i> .....	1503, 1504
<i>vulgaris</i> .....	140	<i>heermanni</i> .....	1504
<i>Ammocætes</i> .....	9	<i>similis</i> .....	1501
<i>æpyterus</i> .....	11	<i>amplexicaollis</i> , <i>Sarothrodus</i> .....	1674
<i>aureus</i> .....	13	<i>amplus</i> , <i>Scarus</i> .....	1635
<i>bicolor</i> .....	10	<i>Tetrapterus</i> .....	892
<i>borealis</i> .....	11	<i>ampullaceus</i> , <i>Ophiognathus</i> .....	406
<i>branchialis</i> .....	14	<i>Saccopharynx</i> .....	406
<i>cibarlus</i> .....	13	<i>Anablepinae</i> .....	632
<i>concolor</i> .....	11	<i>Anableps</i> .....	684, 2807
<i>niger</i> .....	14	<i>dovii</i> .....	685
<i>tridentatus</i> .....	12	<i>Anacanthini</i> .....	782, 2528
<i>unicolor</i> .....	10	<i>Anacanthus</i> .....	82
<i>Ammocætus</i> .....	9	<i>Anacyrtus guatemalensis</i> .....	338
<i>Ammocrypta</i> .....	1061	<i>anagallinus</i> , <i>Lepomis</i> .....	1004
<i>asprella</i> .....	1061	<i>anale</i> , <i>Ditrema</i> .....	1501
<i>beanii</i> .....	1064	<i>analogutta</i> , <i>Pomacentrus</i> .....	1554
<i>gelida</i> .....	1064	<i>analis</i> , <i>Amiurus natalis</i> .....	140
<i>pellucida</i> .....	1062	<i>Caranx</i> .....	927
<i>pellucida clara</i> .....	1063	<i>Centridermichthys</i> .....	2013
<i>vivax</i> .....	1063	<i>Clinocottus</i> .....	2012
<i>vitrea</i> .....	1065	<i>Conger</i> .....	356
<i>ammocryptus</i> , <i>Tetrodon</i> .....	1735	<i>Eupomacentrus</i> .....	1554
<i>Ammodytes</i> .....	832	<i>Holenotus</i> .....	1501
<i>alascanus</i> .....	832, 2842	<i>Hyperprosopon</i> .....	1501
<i>americanus</i> .....	833	<i>Hypocritichthys</i> .....	1500, 1501
<i>aureus</i> .....	13	<i>Lutjanus</i> .....	1267
<i>dubius</i> .....	832	<i>Mesoprion</i> .....	1266
<i>personatus</i> .....	833, 2841, 2842	<i>Neomansis</i> .....	1265
<i>septipinnis</i> .....	2842	<i>Notacanthus</i> .....	615
<i>tobianus</i> .....	2841	<i>Oligocottus</i> .....	2013



Page.		Page.		Page.
833	analis, Ophisoma .....	356	Anelyopsetta quadrocellata .....	2635
831	Orthragoriscus .....	1754	Aneylodon .....	1416
781, 831	Pomacentrus .....	1555	aneylodon .....	1416
2701	Scyris .....	932	jaculidens .....	1416
296	Umbrina .....	1468	parvipinnis .....	1399
296	analogus, Abudedefduf .....	1563	aneylodon, Leuchurus .....	1416
2236	Epinephelus .....	1152	Sagenichthys .....	1416
2201	Eusichistodus .....	1563	andree, Rhinoscopelus .....	569
2201	Kyphosus .....	1385	Scopelus .....	569
1617	Pinnolepterus .....	1386	Stenobranchius .....	569
1619	analostana, Cliola .....	279	andrei, Gobius .....	2218
1720	Cyprinella .....	279	Pomadasis .....	1332
412, 413	analostanus, Leuciscus .....	279	Pristipoma .....	1332
413	Notropis .....	279	Anged .....	414
413	Anarhichadidae .....	2445	Angel, Black .....	1679
2	Anarhichas .....	2445	Angel fish .....	58, 1668, 1684, 1685
3	latifrons .....	2446	Angel Sharks .....	58
3	lupus .....	2447	Angelichthys .....	1684, 2859
1717	Anarmostus .....	1291	ciliaris .....	1684, 1685
1717	Anarrhichas dentifolatus .....	2446	iodocus .....	1686
1139	karrak .....	2446	isabelita .....	1685
849	leopardus .....	2446	angelus, Squatina .....	59
1503	maculatus .....	2446	anginosus, Aluternus .....	1720
1502	minor .....	2446	Angler, Common .....	2713
1503, 1504	orientalis .....	2447	Anglers .....	2713
1504	pantherinus .....	2446	anglorum, Lumpus .....	2097
1501	strigosus .....	2447	anguiformis, Ophichthys (Sphage-	
1674	vomerrinus .....	2447	branchus) .....	374
1635	Anarrhichthyinae .....	2445	anguiformis, Sphagebranchus .....	374
892	Anarrhichthys .....	2447	Anguilla .....	347
406	felis .....	2448	aterrima .....	348
406	ocellatus .....	2448	blephura .....	348
632	anasimos, Acipenser .....	106	chrysypa .....	348, 2801
684, 2807	anceps, Cottus (Acanthocottus) .....	1973	cubana .....	348
685	Plesiopeca .....	1039	laticauda .....	348
782, 2528	Anchisomus .....	1729	lutea .....	348
82	augusticeps .....	1731	novae-terrae .....	348
338	caudicinctus .....	1742	nova-orleanensis .....	348
1094	geometricus .....	1736	oceanica .....	355
1501	reticularis .....	1735	punctatissima .....	348
1554	Anchoa pelada .....	436	rostrata .....	348
140	Anchovia .....	449, 2815	tenuirostris .....	348
927	macrolepidota .....	449	texana .....	348
2013	Sardinella .....	429	tyrannus .....	348
2012	anchovia, Clupea .....	429	wabashensis .....	348
356	Anchovies .....	439, 448	xanthomelas .....	348
1554	Silvery .....	439	anguilla, Anguilla rostrata .....	348
1501	Anchovy, California .....	448	latalurus .....	2788
1501	Striped .....	443	anguillaris, Blennius .....	2436, 2457
1500, 1501	Anchybopsis .....	243	Gunnellus .....	2436
1267	Ancistrus .....	160	Lumpenus .....	2436
1266	chagresi .....	160	Stichaeus .....	2436
1265	Anelyopsetta .....	2634	Zoarces .....	2457
615	dendritica .....	2633	Anguillidae .....	346
2013	Anelyopsetta dilecta .....	2636	anguilliformis, Pholidichthys .....	2405







**IMAGE EVALUATION  
TEST TARGET (MT-3)**



**Photographic  
Sciences  
Corporation**

23 WEST MAIN STREET  
WEBSTER, N.Y. 14580  
(716) 872-4503

128  
125  
132  
122  
120  
118

110  
108  
106  
104  
102  
100

	Page.		Page.
anguina, Murrena .....	390	anomalus, Caulolatilus .....	2277
anguineus, Chlamydoselachus .....	16	Rutilus .....	206
Nerophis .....	774	Anoplagonus .....	2088, 2089, 2093
angulifer, Heros .....	1517	incermis .....	2094
anguliterum, Cichlasoma .....	1517	Anoplarchus .....	2421
angusta, Malthua .....	2738	alectrolophus . 2421, 2422, 2869	
angusticeps, Belone .....	712	atropurpureus .....	2422, 2423
Coregonus .....	466	cristagalli .....	2423
Spheroides .....	1731	purpurescens .....	2423
Tetrodon .....	1731	Anoplogaster .....	839
Tylosurus .....	712	cornutus .....	840
angustidens, Macrostoma .....	555, 2826	Anoplogastrina .....	838
angustifrons, Dermatolepis .....	1159	Anoplopoma .....	1861
Serranus .....	1159	fimbria .....	2801, 1862
angustus, Platycephalus .....	2029	merlangus .....	1862
Ahl. ....	1193	Anoplopomatidæ .....	1861
Anisarchus .....	2435	Anoplopomatinae .....	1861
medius .....	2436	anoplos, Astroscopus .....	2308
Anisochartodon .....	1672	Uranoscopus .....	2308
Anisotremus .....	1314, 1315, 1318	anoplus, Agnus .....	2308
bicolor .....	1319	Astroscopus .....	2308
bilineatus .....	1319	Anopsus .....	7
caesius .....	1316	Anosmius .....	1741
davidsonii .....	1321	Antacens .....	103
dovii .....	1317	antecessor, Gasterosteus .....	900
interruptus .....	1319	Antennariide .....	2715
pacifici .....	1316	Antennarius .....	2717
scapularis .....	1320	annulatus .....	2725
serrula .....	1323	corallinus .....	2725
spleniatus .....	1321	histrio .....	2716, 2723
surinamensis .....	1318, 1319	inops .....	2718
interruptus .....	1319	leopardinus .....	2721
tæniatus .....	1322	marmoratus .....	2717
trilineatus .....	1320	multicellatus .....	2724
virginicus .....	1322, 1323	nuttingii .....	2723
anisurum, Moxostoma .....	190, 196	ocellatus .....	2721
anisurus, Catostomus .....	190	pleurophthalmus .....	2722
anna-carolina, Mugilomorus .....	410	principis .....	2719
anna, Agonus (Brachyopsis) .....	2042	radiosus .....	2725
Cottus .....	1960	reticularis .....	2719
annularis, Centropristes .....	1214	sauguiueus .....	2721
Nauclerus .....	900	scaber .....	2722
Pomoxis .....	987	strigatus .....	2720
Serranus .....	1214	tenebrosus .....	2719
annulata, Melanura .....	624	tennifilis .....	2721
annulatum, Exoglossum .....	327	tigris .....	2723
annulatus, Antennarius .....	2725	antennatus, Chilomycterus .....	1750
Spheroides .....	1735	Diodon .....	1750
Spheroides politus .....	1736	Anthias .....	1226
testudineus .....	1736	aquilonaris .....	1283
Tetrodon .....	1736	asperilingois .....	1227
anogenus, Notropis .....	250, 260	caballerote .....	1257
anolls, Saurus .....	535	chema .....	1157
anomali, Dekaya .....	2277	formosus .....	1304
anomalum, Campostoma .....	205	fureifer .....	1222
		jocu .....	1258

Page.		Page.		Page.
2277	<i>Anthias multifasciatus</i> .....	1226	<i>Aphredoderus</i> .....	786
206	<i>oculatus</i> .....	1283	<i>gibbosus</i> .....	787
2088, 2089, 2095	<i>pernanus</i> .....	1223	<i>saxanus</i> .....	786
2093	( <i>Hermianthias</i> ) <i>pernanus</i> ..	1223	<i>Aphyoninae</i> .....	2499
2421	<i>quartus rondeleti</i> .....	1266	<i>Aphyonus</i> .....	2525
2421, 2422, 2869	<i>rabirabia</i> .....	1276	<i>mollis</i> .....	2525
2422, 2423	<i>saponaceus</i> .....	1232	<i>apia</i> , <i>Pirati</i> .....	1174
242..	<i>striatus</i> .....	1157	<i>apiarius</i> , <i>Petrometopon</i> ..	1142
2423	<i>trifurcus</i> .....	1202	<i>Saranus</i> .....	1142
839	<i>vivans</i> .....	1224	<i>apiatus</i> , <i>Lepomis</i> .....	998
840	<i>Anthilinae</i> .....	1131	<i>apicalis</i> , <i>Alosa</i> .....	429
838	<i>anthracinus</i> , <i>Acipenser</i> .....	106	<i>Clupea</i> .....	429
1861	<i>antica</i> , <i>Algarsea</i> .....	245	<i>Echeneis</i> .....	2268
2861, 1862	<i>antiana</i> , <i>Leucus</i> .....	245	<i>Sardinella</i> .....	429
1862	<i>Antignia</i> .....	1664	<i>Apionichthys</i> .....	2702
1861	<i>capros</i> .....	1665	<i>bleekeri</i> .....	2703
2308	<i>mulleri</i> .....	1665	<i>dumerli</i> .....	2703
2308	<i>Antigoninae</i> .....	1663	<i>nebalosus</i> .....	2703
2308	<i>antillanus</i> , <i>Conodon</i> .....	1324	<i>unicolor</i> .....	2702
2368	<i>antillarum</i> , <i>Caranx</i> .....	921	<i>Aplecion</i> .....	1010
7	<i>Chilomycterus</i> .....	1749	<i>pottsi</i> .....	1083
1741	<i>Sicydium</i> .....	2206	<i>Aplites</i> .....	1010
103	<i>Talismania</i> .....	455	<i>Aplocheilus</i> .....	632; 2827; 2828; 2830
900	<i>Antimora</i> .....	2544	<i>dovii</i> .....	2828, 2830
2715	<i>microlepis</i> .....	2545	<i>Aplodinotinae</i> .....	1337
2717	<i>viola</i> .....	2544	<i>Aplodinotus</i> .....	1483
2725	<i>antiqorum</i> , <i>Hippocampus</i> .....	776	<i>grunniens</i> .....	1484
2725	<i>antiquus</i> , <i>Hippocampus</i> .....	776	<i>Aplurus</i> .....	879
2725	<i>antistius</i> , <i>Chenobryttus</i> .....	992	<i>simplex</i> .....	880
2716, 2723	<i>antoniensis</i> , <i>Pimelodus</i> .....	140	<i>Apocheilichthys</i> .....	633, 2827
2718	<i>Antonino</i> .....	909	<i>Apocope</i> .....	308, 309
2721	<i>antrostomus</i> , <i>Idiacanthus</i> .....	605	<i>carringtonii</i> .....	312
2717	<i>Aodon</i> .....	91	<i>conesii</i> .....	310
2724	<i>hypostomus</i> .....	92, 2756	<i>henshavi</i> .....	312
2723	<i>Aodontidae</i> .....	2756	<i>ubila</i> .....	311
2721	<i>Apeltes</i> .....	752	<i>oscula</i> .....	309
2722	<i>quadraeus</i> .....	752	<i>ventricosa</i> .....	309
2719	<i>apeltes</i> , <i>Gasterosteus</i> .....	752	<i>vulnerata</i> .....	312
2725	<i>Apeltnae</i> .....	743	<i>apoda</i> , <i>Perca</i> .....	1259
2719	<i>aper</i> , <i>Labrus</i> .....	1586	<i>Pleuronectes</i> .....	2701
2721	<i>Aphanopinae</i> .....	885	<i>Apodes</i> .....	344
2722	<i>Aphanopus</i> .....	885	<i>Apodichthys</i> .....	2411
2720	<i>minor</i> .....	885	<i>flavidus</i> .....	2411
2719	<i>Aphododerus cookianus</i> .....	787	<i>fuconum</i> .....	2413
2721	<i>Aphoristia</i> .....	2704	<i>inornatus</i> .....	2412
2723	<i>atricauda</i> .....	2708	<i>sanguineus</i> .....	2412
1750	<i>diomedea</i> .....	2711	<i>univittatus</i> .....	2412
1750	<i>elongata</i> .....	2707	<i>violaceus</i> .....	2427
1236	<i>fasciata</i> .....	2710	<i>virescens</i> .....	2412
1283	<i>marginata</i> .....	2706	<i>Apodontis</i> .....	873
1227	<i>nebulosa</i> .....	2712	<i>apodus</i> , <i>Neomensis</i> .....	1258
1257	<i>ornata</i> .....	2707, 2710	<i>Apogon</i> .....	1106
1157	<i>pigra</i> .....	2706	<i>alutus</i> .....	1110
1304	<i>plagiata</i> .....	2710	<i>atricaudus</i> .....	2853
1222	<i>pusilla</i> .....	2711	<i>hinotatus</i> .....	1109
1258	<i>Aphredoderidae</i> .....	785	<i>dovfi</i> .....	1108
			<i>imberbis</i> .....	1107

	Page.		Page
Apogon maculatus .....	1109	arabatsch, Salmo .....	482
pigmentarius .....	1109	arabicus, Chanos .....	419
retrosella .....	1108	aracanga, Pseudoscarus .....	164
rex-mullorum .....	1107	Scarus .....	1642, 1647
ruber .....	1107	Sparisoma .....	1647
Apogonichthys .....	1110	araca, Atherina .....	799
alutus .....	1110	areopus, Catostomus .....	172
americanus .....	1107	Pantosteus .....	172
puncticulatus .....	1111	Aramaca .....	2679
stellatus .....	1110	papillora .....	2672
Apomotis .....	995	soleceformis .....	2672
chaetodon .....	995	aramaca, Citharichthys .....	2672
cyanellus .....	996	Hemirhamphus .....	2672
ischyrus .....	997	Pleuronectes .....	2672
murius .....	996	Rhombus .....	2626, 2672
obesus .....	993	arangoi, Chaerophilus .....	1597
phenax .....	997	Arara, Bonaci .....	1174
punctatus .....	997	arara, Haemulon .....	1306
symmetricus .....	998	Rouco .....	1304
apos, Gunnellus .....	2430	Serranus .....	1159, 1175
appendiculatus, Centropomus .....	1119	aratus, Lutjanus .....	1274
Exocoetus .....	736	Mesoprion .....	1274
appendix, Lepomis .....	1005	Neomysis .....	1273
Petromyzon .....	10	Arbaciosa .....	2340
approximans, Polydactylus .....	829	eos .....	2343
Polynemus .....	829	humeralis .....	2341
Pomadasys .....	1333	rhossodon .....	2340
Trichidion .....	829	rupestris .....	2341
Aprion .....	42, 1279	zebra .....	2341
ariommus .....	1278	arcansanum, Etheostoma zonale .....	1675
macrophthalmus .....	1280, 1281	arcansana, Notropis telescopus .....	292
aprion, Gerres .....	1373	archidium, Bairdiella .....	1432
Aprionodon .....	42	Eltattarchus .....	1431
isodon .....	42	Odontoscium .....	1422
Aprodon .....	2460	Archistes .....	1900
corfezianus .....	2461	plumarius .....	1900, 1901
Apsicephalus .....	1729	Archocentrus .....	1514, 1515, 1525
Apsilus .....	1278	Archoperca .....	1169, 1171
dentatus .....	1278	Archoplites .....	990
Aptericthys .....	373	interruptus .....	991
selachops .....	374	Archosargus .....	1358, 1359, 1361
Apterurus .....	91	aries .....	1361
apua, Bodianus .....	1174	pourtalesii .....	1360
Epinephelus .....	1159	probatocephalus .....	1361, 1362
Mycteroperca venenosa .....	1173, 1174	tridens .....	1360
Serranus .....	1158	unimaculatus .....	1359, 1360
apus, Centronotus .....	2430	Archosion parvipinnis .....	1399
Platyroctes .....	458	remifer .....	1399
aqua-auleis, Gymnothorax .....	391	Arctic Flounder .....	2649
Muræna .....	391	Grayling .....	517
Rabula .....	390	Sculpin .....	1973
Aquavira .....	1204	arctica, Liparis .....	2121
aquilonaris, Anthias .....	1283	arcticum, Benthosema .....	574
Etelis .....	1283	arcticus Chironectes .....	2717
aquosus, Pleuronectes .....	2660	Salmo .....	521
Rhombus .....	2660	arcticus Scopelus .....	574

Page.		Page.		Page.
483	arctifrons, Calamus .....	1355	argentem, Hyperprosopon .....	1502
419	Citharichthys .....	2683	punctatum ..	1502
1643	Arctoscopus .....	2237	argenteus, Amblyscion .....	1421
1642, 1647	japonicus .....	2297	Amphistichus .....	1503, 1504
1642	Arciozenus .....	601	Centronotus .....	899
790	borealis .....	601	Diplodus .....	1363
172	cornucanus .....	601	Eucinostomus .....	1371
2670	arctum, Siphostoma .....	771	Gerres .....	1371
2672	arcturus, Salmo .....	510	Hyperprosopon .....	1501, 1502
2672	Salvehnus alpinus .....	510	Ichthyomyzon .....	11
2672	arcuata, Harengula .....	431	Larimus .....	1421
2672	arcuatum, Ditrema .....	1502	Leuciscus .....	221
2673	Hamulon .....	1305	Micropogon .....	1403
2672	Hyperprosopon .....	1502	Pagrus .....	1357
2626, 2672	arcuatus, Bathygadus .....	2564	Petromyzon .....	11
1597	Chaetodon .....	1680	Pimelodus .....	125
1174	Hyperprosopon .....	1502	Sarchirus .....	110
1306	Pomacanthus .....	1679, 1681	Sargus .....	1383
1304	ardens, Catoctomus .....	179	Sparus .....	1357
1159, 1175	Hypsilepis .....	301	Synodus .....	411
1274	Leuciscus .....	301	Trachinotus .....	944
1274	Mitsunilus .....	301	Trachynotus .....	944
1273	Notropis umbratilis .....	301	Trichirus .....	889
2340	ardeola, Belone .....	713	Argentina .....	525
2343	Tylosurus .....	713	carolina .....	410
2341	ardesiaca, Gila .....	237	glossodonta .....	411
2340	ardesiacus, Lepomis .....	1006	maculata .....	410
2341	Squalus .....	237	menidia .....	443
2341	arenata, Umbrina .....	1474	pennanti .....	577
ale..... 1675	arenatus, Arius .....	132	pretiosa .....	525
ous..... 292	Priacanthus .....	1237, 1238	sialis .....	526
1432	Rhinichthys .....	308	silus .....	526
1432	Rypticus .....	1232	striata .....	526
1432	arenicola, Fierasfer .....	2496	syrtesium .....	526
1900	Gilléllus .....	2299	Argentines .....	525
1900, 1901	arenosus, Gadus .....	2541	Argentiniæ .....	519
1514, 1515, 1525	argalus, Belone .....	713	argentinus, Pimelodus .....	135
1169, 1171	arge, Alburnellus .....	294	argentipinnis, Rhombus .....	966
990	Kuhlia .....	1014	argentissimus, Gasterosteus .....	747
991	Notropis .....	294	Plageopterus .....	329
1358, 1359, 1361	argentata, Ciliata .....	2559	argentiventris, Lutjanus .....	1261
1361	Couchia .....	2559	Lutjanus .....	1261
1360	Motella .....	2559	Mesoprion .....	1261
1361, 1362	argentatus, Astyanax .....	336	Neomysis .....	1260
1360	Galdropsarus .....	2559	argenti-vittatus, Thynnus .....	871
1350, 1360	Merluccius .....	2530	argentosa, Dionda .....	215
1399	Plargyrus .....	283	Argo .....	957
1399	Tetragonopterus .....	336	argus, Muræna .....	401
2649	argentea, Bathylupea .....	835	Pleuronectes .....	2066
517	Chimæra .....	95	Squalus .....	26
1973	Muræna .....	348	Argyrea .....	796
2121	Selene .....	936	Argyreiosus .....	935
574	Sphyræna .....	826	gabonensis .....	935
2717	Steindachneria .....	2568	pacificus .....	936
521	argenteum, Ditrema .....	1504		
574				

	Page.		Page.
<i>Argyreosus setipinnis</i> .....	934	<i>argyrosomus, Damalichthys</i> .....	1509
<i>unimaculatus</i> .....	934	<i>Argyrotaenia</i> .....	832
<i>vomer</i> .....	936	<i>vittata</i> .....	833
<i>argyreosus, Leucosomus</i> .....	224	<i>argyrurus, Coryphaena</i> .....	955
<i>Pogonichthys</i> .....	224	<i>argyrus, Pimelodus</i> .....	135
<i>Symmetrurus</i> .....	224	<i>aries, Archosargus</i> .....	1361
<i>Argyreus</i> .....	305	<i>Sargus</i> .....	1362
<i>dulcis</i> .....	307	<i>arioides, Bagrus</i> .....	133
<i>nasutus</i> .....	306	<i>ariomnus, Aprion</i> .....	127-
<i>notabilis</i> .....	309	<i>Minnilus</i> .....	290
<i>nubilus</i> .....	311	<i>Notropis</i> .....	290
<i>osculus</i> .....	309	<i>Photogenis</i> .....	290
<i>rubripinnis</i> .....	282	<i>Ariopsis</i> .....	119
<i>argyreus, Fario</i> .....	480	<i>Ariosoma</i> .....	353
<i>Lepidopus</i> .....	887	<i>Arius</i> .....	119
<i>Salmo</i> .....	480	<i>alatus</i> .....	125
<i>Argyrosius capillaris</i> .....	936	<i>arenatus</i> .....	132
<i>spixii</i> .....	936	<i>assimilis</i> .....	120, 2774
<i>triacanthus</i> .....	936	<i>brandtii</i> .....	122, 2758
<i>Argyrosius brevoorti</i> .....	936	<i>caeruleus</i> .....	129
<i>filamentosus</i> .....	936	<i>dasycephalus</i> .....	130
<i>mauricei</i> .....	936	<i>dowi</i> .....	125
<i>mitchilli</i> .....	936	<i>dubius</i> .....	127
<i>setifer</i> .....	936	<i>elatturus</i> .....	128
<i>argyritis, Hybomachus</i> .....	214	<i>emphysetus</i> .....	122
<i>Argylepes</i> .....	915	<i>equestris</i> .....	128
<i>Argyrocottus</i> .....	1995	<i>felis</i> .....	128
<i>zandori</i> .....	1995	<i>fissus</i> .....	131
<i>argyrolenca, Bairdiella</i> .....	1434	<i>flavesceus</i> .....	123
<i>Corvina</i> .....	1434	<i>furthii</i> .....	132
<i>argyrolencus, Bodianus</i> .....	1433	<i>grandicassis</i> .....	126
<i>argyromelas, Scriola</i> .....	650	<i>guatemalensis</i> .....	129
<i>Argyropelecus</i> .....	603	<i>herzbergii</i> .....	125
<i>durvillii</i> .....	604	<i>hypophthalmus</i> .....	133
<i>hemigymnus</i> .....	604	<i>insculptus</i> .....	127
<i>olfersi</i> .....	604	<i>kessleri</i> .....	127
<i>argyrophanus, Engraulis</i> .....	445	<i>laticeps</i> .....	132
<i>Stolephorus</i> .....	444	<i>luniscutis</i> .....	125
<i>argyropomus, Gastrosteus</i> .....	748	<i>melanopus</i> .....	132
<i>Argyrops caprinus</i> .....	1345	<i>mesops</i> .....	123
<i>argyrops, Sparus</i> .....	1346	<i>milberti</i> .....	128
<i>argyrosoma, Damalichthys</i> .....	1510	<i>multiradiatus</i> .....	133
<i>Eniblotoca</i> .....	1510	<i>nuchalis</i> .....	131
<i>Argyrosomus</i> .....	407	<i>oscula</i> .....	127
<i>alascanus</i> .....	2817	<i>parkeri</i> .....	126
<i>artedi</i> .....	468	<i>passany</i> .....	124
<i>sisco</i> .....	469	<i>phrygiatus</i> .....	131
<i>hoii</i> .....	469, 472	<i>planiceps</i> .....	127
<i>lanrettae</i> .....	471, 2817	<i>platypogon</i> .....	127
<i>laevidus</i> .....	470	<i>puncticulatus</i> .....	131
<i>nigripinnis</i> .....	472	<i>quadriscutis</i> .....	126
<i>osneriformis</i> .....	468	<i>rugispinis</i> .....	130
<i>prognathus</i> .....	471	<i>seemani</i> .....	128
<i>pusillus</i> .....	470	<i>seemanni</i> .....	2772
<i>tullibee</i> .....	473	<i>stricticassis</i> .....	126
<i>bisselli</i> .....	473	<i>surinamensis</i> .....	130

Page.		Page.		Page.
1509	<i>Arius tenaninckii</i> .....	123	<i>ascensionis</i> , Scomber .....	925
832	<i>valenciennesi</i> .....	124	<i>ascita</i> , <i>Myxus</i> .....	155
833	<i>variolosus</i> .....	132	<i>asellus</i> , <i>Cheilichthys</i> .....	1740
953	<i>arizonicus</i> , <i>Pantosteus</i> .....	170	<i>Aseraggodes</i> .....	2694
135	<i>Arlina</i> .....	1054	<i>asper</i> , <i>Centridermichthys</i> .....	1944
1361	<i>atripinus</i> .....	1051	<i>Cottus</i> .....	1944
1362	<i>effulgens</i> .....	1058	<i>Diodon</i> .....	1744, 1752
137	<i>arlingtonia</i> , <i>Gambusia</i> .....	652, 2828, 2829	<i>Exerpes</i> .....	2367
127-	<i>arlingtonius</i> , <i>Fundulus</i> .....	652	<i>Hexagrammos</i> .....	1872
290	<i>armata</i> , <i>Bairdiella</i> .....	1436	<i>Macrurus</i> .....	2572
290	<i>Corvina</i> .....	1437	<i>Plouropectes</i> .....	2645
290	<i>armatus</i> , <i>Aspidophornis</i> .....	2067	<i>aspera</i> , <i>Limanda</i> .....	2645
119	<i>Contridermichthys</i> .....	2012	<i>Uranidea</i> .....	1944
353	<i>Centropomus</i> .....	1123	<i>asperilinguis</i> , <i>Anthias</i> .....	1227
119	<i>Serranus</i> .....	1165	<i>Odontanthias</i> .....	1227
125	<i>armiger</i> , <i>Alexurus</i> .....	2203	<i>asperrimus</i> , <i>Balistes</i> .....	1706
132	<i>Arnillo</i> .....	1278	<i>Myllobatis</i> .....	2754
129, 2774	<i>arnillo</i> , <i>Mesoprion</i> .....	1279	<i>aspersus</i> , <i>Epinephelus</i> .....	1154
122, 2758	<i>Tropidinius</i> .....	1279	<i>Serranus</i> .....	1153
129	<i>Arnillos</i> .....	1278	<i>asperulus</i> , <i>Ariodius</i> .....	1903
130	<i>arnillus</i> , <i>Lutjanus</i> .....	1279	<i>Aspidorus quadricornis</i> .....	2041
125	<i>Arnoglossus fimbriatus</i> .....	2077	<i>Aspicottus</i> .....	1937, 1938
127	<i>ventralis</i> .....	2670	<i>bison</i> .....	1938
128	<i>Aroidei</i> .....	119	<i>aspidolepis</i> , <i>Chaetostomus</i> .....	159
122	<i>Arothron</i> .....	1738	<i>Hemiancistrus</i> .....	159
128	<i>orethizon</i> .....	1739	<i>Aspidophoroides</i> .....	2088
128	<i>Arrow-toothed Halibut</i> .....	2609	<i>bartoni</i> .....	2092
131	<i>artedi</i> , <i>Argyrosomus</i> .....	468	<i>grœnlandicus</i> .....	2092
123	<i>siseo</i> .....	469	<i>guntheri</i> .....	2090
132	<i>Polynemus</i> .....	828	<i>incermis</i> .....	2093
126	<i>Artediellus</i> .....	1905	<i>monopterygius</i> .....	2091, 2092
129	<i>atlanticus</i> .....	1906, 2862	<i>olriki</i> .....	2089
125	<i>paefiens</i> .....	1906	<i>tranquebar</i> .....	2092
133	<i>uncinatus</i> .....	1905, 1906	<i>Aspidophoroidinae</i> .....	2033
127	<i>Artedius</i> .....	1902, 2862	<i>Aspidophornis</i> .....	2064
127	<i>asperulus</i> .....	1903, 2862	<i>acipenserinus</i> .....	2002
132	<i>fenestralis</i> .....	1900	<i>armatus</i> .....	2067
125	<i>lateralis</i> .....	1902	<i>cataphractus</i> .....	2067
132	<i>pugetensis</i> .....	1590	<i>chiloensis</i> .....	2009
123	<i>quadriseriatus</i> .....	1897	<i>decagonus</i> .....	2054
128	<i>arteste</i> , <i>Etheostoma</i> .....	1094	<i>doliceædrus</i> .....	2046
133	<i>Pœcilichthys</i> .....	1094	<i>europæus</i> .....	2067
131	<i>arabensis</i> , <i>Pœclia vanderpolti</i> .....	696, 2834	<i>lisiza</i> .....	2036
127	<i>arundinaceus</i> , <i>Syngnathus</i> .....	765	<i>malmoides</i> .....	2054
126	<i>ascani</i> , <i>Salmo</i> .....	509	<i>niger</i> .....	2009
124	<i>Silus</i> .....	526	<i>proboscivalis</i> .....	2038
131	<i>Ascelichthyinæ</i> .....	1883	<i>rostratus</i> .....	2048
127	<i>Ascelichthys</i> .....	2024	<i>spinosissimus</i> .....	2054
127	<i>rhodorus</i> .....	2025	<i>supercilliosus</i> .....	2036
131	<i>ascendens</i> , <i>Siphostoma</i> .....	768	<i>aspidurus</i> , <i>Urolophus</i> .....	81
126	<i>Syngnathus</i> .....	768	<i>Aspistor</i> .....	2763
130	<i>ascensionis</i> , <i>Caranx</i> .....	925	<i>luniventris</i> .....	2763
128	<i>Epinephelus</i> .....	1154	<i>Aspisurus</i> .....	1689
2772	<i>Holocentrus</i> .....	848	<i>asprella</i> , <i>Ammocrypta</i> .....	1001
126	<i>rufus</i> .....	849	<i>Crystallaria</i> .....	1001
130	<i>Perca</i> .....	840	<i>asprellus</i> , <i>Etheostoma</i> .....	1001



	Page.		Page.
asprellus, <i>Plourolepis</i> .....	1061	<i>Atherina</i> .....	789
<i>Radulinus</i> .....	1920	<i>arva</i> .....	790
asprigenis, <i>Paecllichthys</i> .....	1085	<i>bosci</i> .....	801
aspro, <i>Alvordius</i> .....	1033	<i>brownii</i> .....	443
<i>Hadropterus</i> .....	1032	<i>carolina</i> .....	791
<i>Percina</i> .....	1833	<i>harringtonensis</i> .....	791
<i>Asproperca</i> .....	1024	<i>humboldtiana</i> .....	793
<i>zebra</i> .....	1027	<i>insularum</i> .....	807
assimilis, <i>Arius</i> .....	129, 2774	<i>laticeps</i> .....	790
<i>Galeichthys</i> .....	2779	<i>martinea</i> .....	795
<i>Hexanematichthys</i> .....	129	<i>menidia</i> .....	801
<i>Astichthys</i> .....	1006	<i>microps</i> .....	791
<i>caeruleus</i> .....	1089	<i>mordax</i> .....	523
<i>zonalis</i> .....	1075	<i>notata</i> .....	800
asterias, <i>Blennius</i> .....	2383	<i>stipes</i> .....	790
<i>Mustelus</i> .....	29	<i>storeri</i> .....	807
<i>Urolophus</i> .....	82, 2752	<i>veliana</i> .....	790
<i>Asternopteryx</i> .....	2420	<i>viridescens</i> .....	800
<i>gunelliformis</i> .....	2420	<i>vomerina</i> .....	793
<i>Asternotremia</i> .....	786	<i>Atherizella</i> .....	805
<i>mesotrema</i> .....	787	<i>crystallina</i> .....	805
<i>Astrospondyli</i> .....	19	<i>oriarcha</i> .....	803
<i>Asthenurus</i> .....	2526	<i>evermanni</i> .....	804
<i>atripinnis</i> .....	2527	<i>panmensis</i> .....	805
<i>astilbe, Stolephorus</i> .....	2815	<i>Atherinichthys</i> .....	792
<i>astori, Ichthyomyzon</i> .....	12	<i>albus</i> .....	2839
<i>Lampetra</i> .....	12	<i>brevis</i> .....	2840
<i>Petromyzon</i> .....	12	<i>californiensis</i> .....	807
<i>Astracion tricornis</i> .....	1725	<i>gracilis</i> .....	797
<i>Astrolytes</i> .....	1898	<i>grandoculis</i> .....	2840
<i>notospilotus</i> .....	1899	<i>guatemalensis</i> .....	801
<i>Astronesthes</i> .....	586	<i>humboldtii</i> .....	793
<i>barbatus</i> .....	586	<i>menidia</i> .....	800
<i>gemmifer</i> .....	586	<i>notata</i> .....	800
<i>niger</i> .....	580	<i>pachylepis</i> .....	801
<i>richardsoni</i> .....	587	<i>Atherinidae</i> .....	788
<i>Astronesthidae</i> .....	586	<i>Atherinoides</i> .....	792
<i>Astroscopus</i> .....	2306	<i>atherinoides, Chlorodorus</i> .....	719
<i>anoplos</i> .....	2308	<i>Clupea</i> .....	451
<i>anoplus</i> .....	2309	<i>Engraulis</i> .....	451
<i>guttatus</i> .....	2310	<i>Notropis</i> .....	254, 293
<i>y-graecum</i> .....	2307	<i>Pterengraulis</i> .....	450
<i>zephyreus</i> .....	2309	<i>Atherinops</i> .....	807
<i>Astyanax</i> .....	333	<i>affinis</i> .....	807
<i>argentatus</i> .....	336	<i>insularum</i> .....	807
<i>Asymmetron</i> .....	4	<i>regis</i> .....	808
<i>lucayanum</i> .....	4	<i>Atherinopsis</i> .....	806
<i>ataniatus, Chaetodon</i> .....	1676	<i>californiensis</i> .....	806
<i>Sarothrodus</i> .....	1676	<i>tenulis</i> .....	802
<i>atehafalaya, Signalosa</i> .....	2800	<i>Athlennes</i> .....	717
<i>atelaspia, Aclipenser</i> .....	100	<i>hians</i> .....	718
<i>aterrima, Anguilla</i> .....	348	<i>Atimostoma</i> .....	950
<i>Muraena</i> .....	396	<i>Atinga</i> .....	1750
<i>Thyrsoidea</i> .....	396	<i>Atinga alter minor orbicularis</i> .....	1749
<i>Atheresthes</i> .....	2600	<i>Chilomycterus</i> .....	1750
<i>stomias</i> .....	2609	<i>Diodon</i> .....	1746
		<i>atinga, Guamaiaacu</i> .....	1749

Page.		Page.		Page.
789	Atka-fish .....	1864	atripinnis, Bregmaceros .....	2527
790	atkinsi, Gasterosteus bispinosus ..	748	Goodea .....	685
801	atlantica, Elacato .....	948	atrocandalla, Notropis cayuga .....	260
443	Emblemaria .....	2402	atrocyaneus, Pomacentrus .....	1552
791	atlanticum, Oreosoma .....	1663	atromaculata, Esox .....	629
791	atlanticus, Artediellus .....	1906	Etheostoma .....	1057
793	Benthodesmus .....	887	atromaculatus, Cyprinus .....	222
807	Bregmaceros .....	2527	Semotilus .....	222
790	Callorhynchus .....	95	thoreaul-	
795	Dibranchius .....	2743	anus ..	223
801	Epinephelus .....	1154	atronasus, Cyprinus .....	307
791	Megalops .....	409	Rhinichthys .....	307
523	Neoliparis .....	2107	croceus ..	308
800	Prometheus .....	883	lunatus ..	308
790	Promethichthys .....	883	meleagris ..	308
807	Rupiscartes .....	2397	atropurpureum, Ophidium .....	2423
790	Salaris .....	2397	atropurpureus, Anoplarchus .....	2422, 2423
800	Sparus .....	1153	Atropus .....	920
793	Tarpon .....	409	atrorubens, Sebastodes .....	1706
805	Tetragonurus .....	970	atrovirens, Sebastichthys .....	1798
805	Thynnus .....	871	Sebastodes .....	1797
803	atomarium, Sparisoma .....	1631	att. auata, Vincigurria .....	577
804	atomarius, Scarus .....	1631	attenuatus, Maurolicus .....	577
805	Atopoclinus .....	2376	Merluccius .....	2546
792	ringens .....	2376	Osmerus .....	523
2839	Atractoperca .....	1194	attilus, Acipenser .....	105
2840	clathrata .....	1198	atwoodi, Carcharias .....	50
807	Atractosion .....	1402, 1413	aubrieti, Lutjanus .....	1271
797	nobilis .....	1413	Auchenionchus .....	2360
2840	Atractosteus .....	109, 111	Auchenopterus .....	2369, 2371
801	lucius .....	111	atlanticus .....	2371
795	tropicus .....	111	altivelis .....	2370
800	atramentatus, Sympniurus .....	2706	asper .....	2368
801	atraria, Perca .....	1200	fasciatus .....	2373
788	Sibona .....	233	integripinnis .....	2372
792	atrarius, Centropristis .....	1200	marmoratus .....	2371
719	Pimelodus .....	140	monophthalmus .....	2372
451	Serranus .....	1200	nigripinnis .....	2369
451	Squalius .....	233	nox .....	2373
254, 293	Xenomystax .....	361	auctorum, Lobotes .....	1236, 2858
450	atricauda, Aphoristia .....	2708	Auctospina .....	1765, 1776, 1817
807	Hydrargira .....	624	audens, Menidia .....	798
807	atricaudus, Apogon .....	2853	augusticeps, Anechisomus .....	1731
807	Symphurus .....	2707	Aulastome maregravii .....	757
808	atrilatus, Zygonectes .....	682	Auliscops .....	753
806	atrilobatus, Chromis .....	1546	spinescens .....	754
806	atrimana, Monolene .....	2692	auliscus, Siphostoma .....	767
802	atrimanus, Caranx .....	914	Aulopida .....	541
717	Hemlecaranx .....	913	Aulepus agassizii .....	541
718	atripes, Ditrema .....	1507	Aulorhynchidae .....	752
950	Lythrurus .....	300	Aulorhynchus .....	753
1750	Mimulus .....	300	flavidus .....	754
1749	Notropis umbratilis .....	300	Aulostoma .....	754
1750	Phanerodon .....	1507	cinereum .....	755
1746	atripinnis, Arlinia .....	1051	coloratum .....	755
1749	Atheuurus .....	2527	Aulostomidae .....	754

	Page.		Page.
Aulostomus .....	754	aurolineatum, Hamulon .....	1310
cinereus .....	755	aurolineatus, Diabasis .....	1309
maculatus .....	754, 2837	auropunctatus, Callyodon .....	1624
aurantiacum, Etheostoma .....	1041	Cryptotomus .....	1624
aurantiacus, Balistes .....	1718	aurora, Abeona .....	1497
Ceratacanthus .....	1718	Capropomus .....	1665
Cottogaster .....	1041	Catoatomus .....	176
Hadropterus .....	1041	Fario .....	499
Hypohomus .....	1040	Salmo .....	493
aurata, Cliola .....	272	Sebastichthys .....	1803
Coryphaena .....	953	Sebastodes .....	1802
Moniana .....	272	aurorubens, Centropriaris .....	1278
Saleima .....	1380	Lutjanus .....	1278
auratum, Pristipoma .....	1324, 1343	Mesoprion .....	1278
auratus, Carassius .....	201	Rhomboplites .....	1277, 1278
Centropomus .....	1107	aurovittatus, Mesoprion .....	1276
Cyprinus .....	201	Ocyrops .....	1276
Gadus .....	2542	australe, Etheostoma .....	1081
Holocentrus .....	1145	Zophendum .....	212
Mullus .....	856	australis, Echenis .....	2269, 2271
barbatus .....	856	Esox .....	628
Notemigonus .....	250	Icelus .....	1918
Psenes .....	951	Tenthis .....	1691
Serranus .....	1145	Remilegia .....	2270
aurea, Brevoortia tyrannus .....	434	austrina, Myxostoma .....	192
Clupea .....	434	austrinum, Minytrema .....	192
Lampetra .....	13	Moxostoma .....	192
aureolum, Moxostoma .....	192	Auxis .....	867
aureolus, Catoatomus .....	192, 196	rochei .....	868
Gerres .....	1375	tapeinosoma .....	868
Salvelinus alpinus .....	511	thazard .....	867
Xenotis .....	1003	thyroides .....	868
auroruber, Serranus .....	1635	vulgaris .....	868
auroriviridis, Sphyræna .....	1119	Averruncus .....	2069, 2864
aureus, Aumocetes .....	13	emmelane .....	2060
Caranx .....	923	sterleus .....	2071
Chaetodon .....	1680	averruncus, Kathetostoma .....	2311
Chupanodon .....	434	avitus, Chologaster .....	704
Enpomotis .....	1010	avocetta, Nemichthys .....	369
Fundulus .....	659	Avocettina .....	367
Haplochilus .....	659	elongata .....	2802
Heros .....	1533	gilli .....	2801
Pomacanthus .....	1680	infans .....	367, 2801
Sparus .....	1010	Awa .....	414
auriculatus, Sebastodes .....	1817, 1818	Awaous .....	2234
dalli .....	1818	flavus .....	2235
auriga, Dules .....	1220	mexicanus .....	2237
Monacanthus .....	1716	nelsoni .....	2235
Serranus .....	1221	talasica .....	2236
auritus, Labrus .....	1001	axillare, Pristipoma .....	1328
Lepomis .....	1001, 1009	axillaris, Boreoottus .....	1981
solis .....	1001	Brachydeuterus .....	1328
aurofrenatum, Sparisoma .....	1694	Cottus .....	1981
aurofrenatus, Serranus .....	1634	Gerres .....	1378
auroguttatus, Zygonectes .....	654, 2829	Myoxocephalus .....	1980, 1981
aurolineatum, Bathystoma .....	1310	Pomadasis .....	1328

Page.		Page.		Page.
1310	axinophrys, Xystes .....	2076	Bagrus macronemus .....	117
1309	Azyrius .....	1903, 2662	mesops .....	123
1624	harringtoni .....	1904	passany .....	124
1624	aya, Acanth .....	1264	pomecus .....	125
1497	Bodlanus .....	1265	proops .....	124
1665	Chaetodon .....	1075	temminckianus .....	123
176	Lutjanus .....	1265	valenciennesi .....	124
499	Mesoprion .....	1204	Bahama Lanelet .....	4
493	Neomans .....	1264	bahamensis, Pisces viridis .....	1638
1801	Aylopon .....	1226	Vulpes .....	411
1802	martinicensis .....	1228	bahianus, Acanthurus .....	1693
1278	ayresi, Aelpenser .....	104	Teuthis .....	1693
1278	Centropristes .....	1205	bahiensis, Cypselurus .....	2836
1278	Parophrys .....	2640	Exocoetus .....	730
1277, 1278	Petromyzon .....	13	Felichthys .....	118
1276	Ayresia .....	1545, 1548	Galeichthys .....	119
1276	punctipinnis .....	1548	Rhombus .....	2064
1081	ayresii, Sebastodes .....	1808	baileyl, Cyprinodon .....	675
212	azalea, Runula .....	2377	bailloni, Gasterosteus .....	747
2269, 2271	Azevia .....	2677	Bulone .....	506
628	panamensis .....	2677	Baiostoma .....	2694, 2095
1918	querna .....	2675	brachialis .....	2608
1691	azorica, Coryphæna .....	953	bairdi, Aelpenser .....	105
2270	Azteca .....	254, 255, 258, 2799	Callionymus .....	2185
192	Aztecula .....	2799	Cottus punctulatus .....	1950
192	aztecus, Notropis .....	258	bairdianum, Siphostoma .....	765, 770
192	Azul, Pescado .....	1553	bairdianus, Sphyrænops .....	1114
867	Azules, Pescados .....	1549	Syngnathus .....	770
868	azurea, Hermosilla .....	1383, 1384	Bairdiella .....	1432, 1433
868	Galeichthys .....	595	aluta .....	1437
868	Azurina .....	2775	archidinn .....	1432
868	hirundo .....	1544	argyroleuca .....	1434
2069, 2804	azurissimus, Microspathodon .....	1544	armata .....	1436
2069	dorsa .....	1570	chrysoleuca .....	1438
2071	lis .....	1570	chrysur .....	1433
2311	Bacalao .....	1184, 1185	ensifera .....	1434
704	bacalaus, Gobius .....	2230	icistia .....	1435
309	Bacalhao sabura .....	2230	punctata .....	1434
367	Bachelor .....	987	ronchus .....	1436
2802	badius, Euctenogobius .....	2227	bairdii, Alepocephalus .....	454
2801	Fundulus heteroclitus .....	2827	Bathymyzon .....	9
307, 2801	Gobius .....	2227	Cottus .....	1951
414	Rhynchichthys .....	308	Gastrostomus .....	406
2234	bagre, Alurichthys .....	117	Macrourus .....	2583
2235	Felichthys .....	117	Microspathodon .....	1560, 1567
2235	Silurus .....	117	Mitchillina .....	454
2236	Bagre Colorado .....	122	Petromyzon .....	9
1328	Sapo .....	2319	Pomacentrus .....	1507
1981	Bagres de Rio .....	149	Salmo .....	508
1328	Bagrus albicans .....	124	Salvelinus .....	508
1981	arioides .....	133	Bajonado .....	1352
1378	coelestinus .....	125	bajonado, Calamus .....	1352, 1353
1980, 1981	emphysetus .....	122	Pagellus .....	1352
1328	flavescens .....	123	Sparus .....	1352
			balantiophthalmus, Scomber .....	911
			balao, Hemiramphus .....	723, 2835

	Page.		Page.
Balaos .....	718, 722, 723	Balistes punctatus .....	1702
balearica, Conger-muræna .....	356	ringens .....	1709, 1711
Conger-muræna .....	356	rufus .....	1707
Muræna .....	356	schmittii .....	1705
balearicum, Ophlosoma .....	356	schepfli .....	1718
balias, Chirus .....	1873	scelopax .....	759
Baliste, Le Bridé .....	1704	scriptus .....	1719
Balistes .....	1609, 1700, 1703	serraticornis .....	1720
amboinensis .....	1704	sobaco .....	1706
americanus .....	1707	spilotopterygius .....	1702
asperrimus .....	1706	sullamen .....	1706
aurantiacus .....	1718	taeniopterus .....	1702
barbatus .....	1720	unicornus .....	1720
bellus .....	1703	urantiacus .....	1718
broccus .....	1716	vetula .....	1703
buniva .....	1701, 1711	Balistidæ .....	1698
capistratus .....	1704	Ballerus .....	249
caprinus .....	1702	balteatum, Chichlasoma .....	1521
caprisens .....	1701	balteatus, Abramis .....	239
carolinensis .....	1701	Cyprinus (Abramis) .....	239
cicatricosus .....	1709	Eques .....	1490
ciliaris .....	1702	Heros .....	1522
ellipticus .....	1715	Leuciscus .....	238
curassavicus .....	1709	Pomacanthus .....	1689
equestris .....	1702	Richardsonius .....	239
forcipatus .....	1702	Thynnus .....	871
frenatus .....	1705	Upeneus .....	860
fuliginosus .....	1702	banana, Butyrinus .....	411
guttatus .....	1702	Gobius .....	2236
heckeli .....	1709	Banana-fish .....	411
hippe .....	1705	bancrofti, Torpedo .....	78
hispidus .....	1716	Banded Pickerei .....	626
kleinei .....	1720	Suntishes .....	994
laevis .....	1719	Bang .....	423
liberensis .....	1702	banksi, Citula .....	927
lineo-punctatus .....	1709	Barathrolemus .....	2517
linguata .....	1720	manatinus .....	2517
longus .....	1707	Barathronus .....	2524
macrops .....	1706	bicolor .....	2524
macropterus .....	1707	barbara, Siphostoma .....	765
maculatus .....	1707, 1708	barbata, Brotula .....	2500
melanopterus .....	1707	Loricaria .....	158
mento .....	1710	Pallasina .....	2049
mitis .....	1705	barbatula, Lemonema .....	2557
monoceros .....	1719, 1720	barbatulum, Lemonema .....	2556
moribundus .....	1702	barbatum, Echistoma .....	587
naufragium .....	1730, 1701	Lycenema .....	2474
niger .....	1711	barbatus, Astronesthes .....	586
nigra .....	1711	Balistes .....	1720
nitidus .....	1709	Enchelyopus .....	2500
notatus .....	1709	Gadus .....	2541
oblonginseculus .....	1720	Liparis .....	2118
oculatus .....	1707	Lonchurus .....	1482
ornatus .....	1719	Mullus auratus .....	856
piceus .....	1711	Siphagonus .....	2050
polylepis .....	1700	Barber .....	1601
powelli .....	1702	Barbeiro .....	1093, 1226

Page.		Page.	
1702	Barbero .....	1691	Bass, Oswego .....
1709, 1711	Negro .....	1692	Rock .....
1707	Barboso, Congre.....	155	Rock Sea.....
1705	Barbu .....	829	Round.....
1718	Barbuda, Lija .....	1720	Sea .....
759	Barbudo.....	150, 829	Stone .....
1719	Barbudos.....	828, 854	Strawberry.....
1720	Barbulifer.....	2260	Striped.....
1706	cauthucous.....	2260	White.....
1702	papillosus .....	2261	White Lake.....
1706	barbulifer, Rhinopolaris.....	2145	"White Sea" of California .....
1702	bardus, Minomus .....	171	Yellow .....
1720	Barfish .....	987	Bassogigas .....
1718	barikani, Agonus (Brachyopsis) .....	2044	gillii.....
1703	Barndoor Skate .....	71	stelliferoides .....
1698	baronis-mulleri, Pimelodus .....	151	Bassozetus .....
249	Rhamdia .....	151	catena .....
1521	Barracouta .....	826	compressus .....
230	Barracuda.....	2841	normalis .....
239	California .....	826	tenia .....
1490	European .....	826	Bastard Halibut .....
1522	Great.....	823	Margaret.....
238	Northern.....	825	Weakfish .....
1689	barracuda, Esox .....	823	batabana, Corvula.....
239	Sphyraena .....	2841	batabanus, Johnius .....
871	Barracudas .....	822	Larimus.....
860	barrattii, Boleosoma.....	1102	Batfish .....
411	Hololepis .....	1102	Short-nosed .....
2236	Paciliolepis .....	1102	Bat-Fishes .....
411	barreto, Gobioides.....	2264	Bathyaegonus.....
626	Barretos.....	2263	nigripinnis .....
994	bartholomae, Caranx .....	919	bathybius, Cynoglossus .....
423	bartoni, Acara .....	1516	Eubassichthys .....
927	Aspidophoroides .....	2092	Histiobranchus .....
2517	Chirostoma .....	793	Synaphobranchus.....
2517	Cichlasoma .....	1515	Bathyclupea .....
2524	Cylindrosteus .....	111	argentea .....
2524	Eslopaarum .....	2840	Bathyclupeidae .....
765	Bascanichthys .....	378	Bathygadina.....
2500	bascanium .....	379	Bathygadus .....
158	peninsulae .....	379	arcuatus .....
2049	scuticaris .....	378	cavernosus .....
2557	bascantium, Bascanichthys .....	379	favosus .....
2556	Cœcula .....	380	longifilis .....
587	Bascanius .....	2704	macrops .....
2474	Bashaw .....	143	Bathylaco .....
586	basilaris, Heros .....	1532	nigricans .....
1720	Basking Sharks .....	50	Bathylagina .....
2500	Bass, Bayou .....	1012	Bathylagus .....
2541	Black .....	1010	benedicti .....
2118	Sea.....	1108	borealis .....
1482	Calico .....	987	euryops .....
856	Channel .....	1453	milleri .....
2050	Common Rock .....	990	pacificus .....
1601	Grass .....	987	Bathymaster .....
1693, 1926	Green .....	1012	hypoplectus .....

	Page.		Page.
Batiymasterjordanii .....	2239	Baya .....	1176
signatus .....	2238	bayanus, Pomadasis .....	1331
bathymaster, Bregmaceros .....	2527	Bayou Bass .....	1012
Bathymastoridae .....	2287	bdellium, Petromyzon .....	11
Bathymyzon .....	9	Rdellostoma dombey .....	6
bairdii .....	9	polytrema .....	6
Bathynectes .....	2507	stouti .....	6
compressus .....	2508	beardiei, Synechoglanis .....	135
laticeps .....	2523	beani, Caranx .....	920
Bathyonus .....	2507	Heros .....	1538
compressus .....	2509	Ophidion .....	2487
taenia .....	2510	Pleuronectes .....	2646
Bathyopsis .....	605	Pæcilichthys .....	1057
ferox .....	605	Triglops .....	1924
Bathyphasma .....	2128, 2864	beanii, Ammocrypta .....	1064
ovigerum .....	2128	Limanda .....	2646
bathyphila, Cyclothone .....	582	Melamphaes .....	843
bathyphilum, Neostoma .....	583	Plectromus .....	842
Bathypteroidae .....	544	Prionotus .....	2170
Bathypterois .....	544	Sorrivomer .....	367
longipes .....	546	beardaleei, Salmo gairdneri .....	2819
quadrifilis .....	545	Bear Lake Bullhead .....	1954
Bathysaurus .....	539	Beau Gregory .....	1555
agassizii .....	540	Beauty, Rock .....	1684
ferox .....	539	beekwithi, Cyprinella .....	273
Bathysebastes .....	1860	Becuna .....	823
Bathyatoma .....	1308	becuna, Sphyraena .....	823
aurolineatum .....	1310	beldingii, Cottus .....	1958
rimator .....	1308	belengeri, Caranx .....	923
striatum .....	1310	belisanus, Beloneox .....	684
Bathytroctes .....	454	belizianus, Eleotris (Cullis) .....	2201
stomias .....	454	bella, Hypoclydonia .....	1113
Batis .....	66	Bellator .....	2173
Batoides .....	59	egretta .....	2174
Batrachoides .....	2314, 2868	militaris .....	2173
pacifici .....	2314	bellicus, Nocomia .....	3213
surinamensis .....	2314	Bellows-fish .....	759, 2713
tau .....	2314	bellus, Ballistes .....	1703
vernulus .....	2316	Minnilus .....	297
variegatus .....	2316	Notropis .....	297
Batrachoididae .....	2313	Belly, Yellow .....	1001
Batrachops .....	1740	Belone almeida .....	715
Batrachus .....	2314, 2315	altipinna .....	717
celatus .....	2316	angusticeps .....	712
guavina .....	2195	ardeola .....	714
magaritatus .....	2323	argalus .....	713
pacifici .....	2315	caribbea .....	717
porosissimus .....	2321	oignonella .....	713
surinamensis .....	2314	crassa .....	716
tau .....	2316	depressa .....	711, 713
beta .....	2316	diplotania .....	712
pardus .....	2317	exilis .....	714
variegatus .....	2316	galeata .....	716
Batrachius .....	2868	gerania .....	716
bat'tarae, Orthragoriscus .....	1754	guyanensis .....	715
Bay Shark .....	37	hians .....	718

Page.		Page.		Page.
1176	<i>Belone jonesi</i> .....	717	<i>beryllina</i> , <i>Menidia</i> .....	797
1331	<i>latimana</i> .....	717	<i>gracilis</i> .....	797
1012	<i>maculata</i> .....	718	<i>beryllinum</i> , <i>Chirostoma</i> .....	798
11	<i>melanochira</i> .....	716	<i>beryllinus</i> , <i>Cryptotomus</i> .....	1624
6	<i>microps</i> .....	712	<i>Beryx</i> .....	834
6	<i>notata</i> .....	711	<i>decadactylus</i> .....	844
6	<i>pacifica</i> .....	716	<i>splendens</i> .....	844
135	<i>raphidoma</i> .....	716	<i>Besau</i> .....	1687
920	<i>scrutator</i> .....	714	<i>Beshow</i> .....	1862
1538	<i>stolzmanni</i> .....	713	<i>Besugo</i> .....	1356
2487	<i>subtruncata</i> .....	711	<i>beta</i> , <i>Batrachus tau</i> .....	2316
2646	<i>timucu</i> .....	715	<i>betaurus</i> , <i>Cirrhitus</i> .....	1492
1057	<i>truncata</i> .....	714, 715	<i>biaeuleatus</i> , <i>Gasterosteus</i> .....	748
1924	<i>belone</i> <i>Esox</i> .....	714	<i>Biajaiba</i> .....	1270
1064	<i>Histiophorus</i> .....	892	<i>de lo Alto</i> .....	1140
2646	<i>Tetrapturus</i> .....	892	<i>Bibronia</i> .....	2704
843	<i>Belonesox</i> .....	884	<i>biaudalis</i> , <i>Lactophrys</i> .....	1723
842	<i>belizanus</i> .....	884	<i>Ostracion</i> .....	1723
2170	<i>Belonichthys</i> .....	773	<i>bicolor</i> , <i>Algansea</i> .....	245
367	<i>Belonopsis</i> .....	369	<i>Ammocetes</i> .....	10
2819	<i>leuchtenbergii</i> .....	369	<i>Anisotremus</i> .....	1319
1954	<i>bendirei</i> , <i>Potamocottus</i> .....	1965	<i>Barathronus</i> .....	2524
1555	<i>Uranidea</i> .....	1964	<i>Exocoetus</i> .....	738
1684	<i>benedicti</i> , <i>Bathylagus</i> .....	529	<i>Grammiconotus</i> .....	726
273	<i>benoiti</i> , <i>Myctophum</i> .....	573	<i>Leuciscus</i> .....	232, 245
823	<i>Scopelus</i> .....	573	<i>Leucos</i> .....	245
823	<i>Benthocometes</i> .....	2514	<i>Pristipoma</i> .....	1320
1958	<i>robustus</i> .....	2514	<i>Rhypticus</i> .....	1232
923	<i>Benthodesmus</i> .....	887	<i>Rondeletia</i> .....	548
684	<i>atlanticus</i> .....	887	<i>Rutilus</i> .....	244
2201	<i>elongatus</i> .....	988	<i>Rypticus</i> .....	1231
1115	<i>Benthosaurida</i> .....	543	<i>Smecticus</i> .....	1232
2173	<i>Benthosaurus</i> .....	543	<i>Squalius</i> .....	232
2174	<i>grallator</i> .....	543	<i>Tigoma</i> .....	232
2173	<i>Benthozema</i> .....	573	<i>bicornis</i> , <i>Centridermichthys</i> .....	1913
3213	<i>arcticum</i> .....	574	<i>Cottus</i> .....	1913
759, 2713	<i>mulleri</i> .....	574	<i>Icelus</i> .....	1911
1703	<i>berardi</i> , <i>Ainteros</i> .....	1720	<i>Bielaya Ryba</i> .....	480
297	<i>Bergall</i> .....	1577	<i>bifasciatum</i> , <i>Cichlasoma</i> .....	1521
297	<i>Berg-gylt</i> .....	1577	<i>Thalassoma</i> .....	1610, 2550
1001	<i>berglax</i> , <i>Macrourus</i> .....	2581	<i>bifasciatus</i> , <i>Chlorichthys</i> .....	1600
715	<i>berlandieri</i> , <i>Lepidosteus</i> .....	111	<i>Heros</i> .....	1521
717	<i>Mugil</i> .....	812	<i>Julis</i> .....	1610
712	<i>Bermuda Catfish</i> .....	882	<i>Labrus</i> .....	1609
713	<i>Chub</i> .....	1387	<i>Thalassoma</i> .....	1610
713	<i>bermuda</i> , <i>Fundulus</i> .....	643	<i>bifrenata</i> , <i>Hemiltemia</i> .....	259
717	<i>bermudensis</i> , <i>Hierasfer</i> .....	2497	<i>bifrenatus</i> , <i>Hypopsis</i> .....	259
713	<i>Lefroyia</i> .....	2497	<i>Notropis</i> .....	253
716	<i>bernardini</i> , <i>Catostomus</i> .....	178	<i>bifurca</i> , <i>Chetodon cauda</i> .....	1562
711, 713	<i>Berrigate</i> .....	2857	<i>Big Eye</i> .....	1238
712	<i>bertheloti</i> , <i>Crius</i> .....	971	<i>Big-eyed Herring</i> .....	410, 426
714	<i>Berycida</i> .....	837	<i>Scad</i> .....	911
716	<i>Berycina</i> .....	838	<i>Big-headed Gurnard</i> .....	2171
716	<i>Berycoid Fishes</i> .....	833	<i>Big Skate</i> .....	68
715	<i>Berycoidel</i> .....	781, 833, 834	<i>of California</i> .....	72
718	<i>Berycoids</i> .....	837	<i>biguttata</i> , <i>Cochlognathus</i> .....	252



	Page.		Page.
biguttatus, <i>Ceratichthys</i> .....	323	bisua, <i>Scomber</i> .....	867
<i>Labrisomus</i> .....	2360	bivittata, <i>Elucate</i> .....	948
<i>Malacoctenus</i> .....	2360	<i>Haiparca</i> .....	1205
<i>Semotilus</i> .....	322	bivittatus, <i>Centropristis</i> .....	1205
bilinearis, <i>Merluccius</i> .....	2530	<i>Chaurojulis</i> .....	1597
<i>Merluccius</i> .....	2531	<i>Hallecheres</i> .....	1597
<i>Stomodon</i> .....	2531	<i>Hybopsis</i> .....	233
bilineata, <i>Lepidopsetta</i> .....	2643	<i>Iridio</i> .....	1595
<i>Plateasa</i> .....	2643	<i>Labrus</i> .....	1596
bilineatum, <i>Pristipoma</i> .....	1319	<i>Minnilus</i> .....	233
bilineatus, <i>Anisotremus</i> .....	1319	<i>Platyglossus</i> .....	1597
<i>Characodon</i> .....	668	<i>Serranus</i> .....	1205
<i>Pleuronectes</i> .....	2643	bixanthopterus, <i>Caranx</i> .....	926
Billfish .....	109, 714, 725, 892	Black and yellow Rockfish .....	1825
<i>billingsiana</i> , <i>Cliola</i> .....	272	<i>Angel</i> .....	1679
<i>Cyprinella</i> .....	272	Black-banded Rockfish .....	1827
<i>biloba</i> , <i>Corvina</i> .....	1460	<i>Sunfish</i> .....	995
<i>Pachypops</i> .....	1460	<i>Bass</i> .....	1010
<i>bilobus</i> , <i>Blepsias</i> .....	2018	<i>Large-mouthed</i> .....	1012
<i>Histiocottus</i> .....	2018	<i>Small-mouthed</i> .....	1011
<i>Peropus</i> .....	2018	Black-belly .....	426
bimaculata, <i>Percina</i> .....	1027	<i>Bullhead</i> .....	141
<i>Pileoma</i> .....	1027	<i>Croaker</i> .....	1456
bimaculatus, <i>Chetodon</i> .....	1674	<i>Drums</i> .....	1454
<i>Clinus</i> .....	2358	Blackfin .....	472
<i>Malacoctenus</i> .....	2358	<i>Snapper</i> .....	1261
<i>Pæcilioides</i> .....	678	Black-fish .....	207, 963, 1199, 1200, 1578
<i>Pseudoxiphophorus</i> .....	678	blackfish, <i>Labrus</i> .....	1578
<i>Sayris</i> .....	725	Blackfishes, <i>Alaska</i> .....	620, 261
<i>Xiphophorus</i> .....	678	<i>Groupers</i> .....	1161, 1174
binoculata, <i>Raja</i> .....	72	<i>Grunt</i> .....	1297
<i>Uraptera</i> .....	73	<i>Guativeres</i> .....	1146
binotatus, <i>Apogon</i> .....	1109	<i>Harry</i> .....	1189
Bipinnula .....	878, 2843	Black-head Minnow .....	217
<i>violacea</i> .....	878	Blackhorse .....	168
bipinnulata, <i>Seriola</i> .....	907	Black Jewfish .....	1161
bipinnulatus <i>Elagatis</i> .....	906	<i>Muray</i> .....	396
birostratus, <i>Prionotus</i> .....	2152, 2156	<i>nosed Dace</i> .....	305, 307
<i>Manta</i> .....	92	<i>Oldwife</i> .....	1711
bishopi, <i>Aloss</i> .....	430	<i>Perch</i> .....	1504
<i>Sardinella</i> .....	430	<i>Pilot</i> .....	1555
bison, <i>Aspicottus</i> .....	1938	<i>Rockfish</i> .....	1784
<i>Carpiodes</i> .....	106	<i>Rudder Fishes</i> .....	963
<i>Enophrys</i> .....	1938	<i>Ruffe</i> .....	963
<i>Lepisosteus</i> .....	110	<i>Ruffs</i> .....	962
bispinosus, <i>Gasterosteus</i> .....	748	<i>Sculpin</i> .....	1985
<i>atkinsi</i> .....	748	<i>Sea Bass</i> .....	1198
<i>cuvieri</i> .....	749	Black-sided Darter .....	1028, 1032
<i>Melanocetus</i> .....	2734	Black-spotted Trout .....	487
<i>Melichthys</i> .....	1711	<i>Swallowers</i> .....	2291
<i>Myliobatis</i> .....	89	<i>Will</i> .....	1109
bisselli, <i>Argyrosomus tullibee</i> .....	473	blackfordi, <i>Lutjanus</i> .....	1265
<i>Coregonus tullibee</i> .....	473	<i>Yarella</i> .....	584
bistriplinus, <i>Bodianus</i> .....	1234	blainvillei, <i>Acanthias</i> .....	51
<i>Ryptilus</i> .....	1233	Blakea .....	2351
		<i>elegans</i> .....	2353

Page.		Page.		Page.
867	Blanca, Aguja .....	892	Blennius flicicornis .....	2381
948	Chopa .....	1388	fimbriatus .....	2457
1205	Lisa .....	813	fucorum .....	2379
1205	Mojarra .....	1372	geminatus .....	2385
1597	Pesca .....	321	gentilis .....	2388
1597	Sardina .....	332	gracilis .....	2438
233	Blancas, Mojarras .....	1372	gunnellus .....	2419
1595	blanchardi, Gasterosteus .....	740	hentzi .....	2399
1596	Neoolinus .....	2354	herminier .....	2362
233	Blanco, Burro .....	1328	labrosus .....	2457
1597	Matajuelo .....	2275, 2276	lampetraformis .....	2438
1205	Pescado de Chapala .....	792	lumpeus .....	2438
926	Ronco .....	1297	marmoreus .....	2381
1825	Blancos, Pescados .....	792	microstomus .....	2385
1679	Blanquillo .....	2276, 2278	multifilis .....	2385
1827	Blanquillos .....	2274, 2276	murænoides .....	2419
995	bleekeri, Apionichthys .....	2703	nuchifilis .....	2383
1010	bleekeriana, Ilisha .....	436	oceanicus .....	2379
1012	Peliona .....	436	pilicornis .....	2380
1011	Blenitrachus .....	2391	polaris .....	2469
426	Blenniocottus .....	2016, 2864	polyactocephalus .....	2409
141	acuticeps .....	2864	punctatus .....	2390, 2440
1456	embryum .....	2016, 2864	regius .....	2553
1454	globiceps .....	2017	rosous .....	2420
472	bryosus .....	2017	serpentinus .....	2439
1261	Blennies .....	2344, 2377	stearnsi .....	2379
0, 1200, 1578	Snake .....	2435	striatus .....	2388
1578	Blenniidae .....	2314	tenia .....	2418
620, 261	Blenninæ .....	2346	torak .....	2561
1161, 1174	Blennioclinus .....	2360	truncatus .....	2381
1297	Blennoid Fishes .....	2343	vinctus .....	2382
1146	Blennioidea .....	2343	(Zoarces?) polaris .....	2460
1109	Blennioidei .....	782	blennius, Alburnops .....	262
217	blennioidea, Diplesion .....	1053	Etheostom .....	1072, 1073
168	Etheostoma .....	1033	Minnilus .....	262
1161	Blenniulus .....	2386, 2390	Notropis .....	261
396	blennioperca, Hyostoma .....	1053	Blennophis .....	2400
305, 307	Blenniophidium .....	2428	webbii .....	2401
1711	petropauli .....	2430	Blenny, Snake .....	2438
1504	Blennius .....	2377, 2378, 2553	Blepharichthys .....	931
1555	alectrolophus .....	2422	crinitus .....	932
1784	americanus .....	2457	Blepharis .....	931
963	anguillaris .....	2436, 2457	crinitus .....	932
963	asterias .....	2383	major .....	932
962	bosquianus .....	2394	antor .....	932
1985	brevipinnis .....	2391	blepharis, Carangoides .....	932
1198	capiti laevi .....	2438	blephura, Anguilla .....	348
1028, 1032	carolinus .....	2378	Blepsias .....	2018
487	chuss .....	2555	bilobus .....	2018
2201	ciliatus .....	2457	cirrhosus .....	2018
1199	(Clinus) lumpeus .....	2438	oculofasciatus .....	2021
1265	crinitus .....	2383	trifolobus .....	2019
584	cristatus .....	2382, 2383	ventricosus .....	1930
51	dolichogaster .....	2417	Blepsina .....	1883
2351	europæus .....	2410	Bleu, Poisson .....	517
2353	favosus .....	2380	Blindfish, Cuban .....	2501

	Page.		Page.
Blindfish, of the Mammoth Cave...	706	<i>Bodianus blistripinus</i> .....	1234
Small.....	704	<i>blochii</i> .....	1583
Blind Fishes.....	702	<i>bodianus</i> .....	1583
Gobies.....	2261	<i>costatus</i> .....	1402
Goby of Point Loma.....	2262	<i>cruentatus</i> .....	1142
Bleater.....	471	<i>diplotænia</i> .....	1582
Blot.....	1950	<i>dubius</i> .....	1146
<i>blochii</i> , <i>Bodianus</i> .....	1583	<i>exiguus</i> .....	1433
<i>Caranx</i> .....	919	<i>flavescens</i> .....	1024
<i>Galeichtbys</i> .....	118	<i>fulvus</i> .....	1144
<i>Orthogoriscus</i> .....	1754	<i>punctatus</i> .....	1146
<i>Pimelodus</i> .....	155	<i>ruber</i> .....	1145, 1146
<i>Piramutana</i> .....	155	<i>guativero</i> .....	1145
Blower.....	1733	<i>jaguar</i> .....	849
Blow-fish, Spring-back.....	1734	<i>margnatus</i> .....	1174
Blue-back.....	420	<i>pallidus</i> .....	1433
Mullet.....	813	<i>pauaensis</i> .....	1141
Salmon.....	481	<i>pectoralis</i> .....	1582
Trout.....	514	<i>pentacanthus</i> .....	849
Trout of Lake Crescent..	2819	<i>pulchellus</i> .....	1584
Blue Bream.....	1005	<i>punctatus</i> .....	1146
Cut.....	134	<i>punctiferus</i> .....	1147
Cod.....	1875	<i>ruber</i> .....	1265
Darter.....	1088	<i>rufus</i> .....	1135, 1583
Herring.....	425	<i>rupestris</i> .....	990
Mullet.....	191	<i>stellifer</i> .....	1443
Parrot-fish.....	1636, 1652	<i>striatus</i> .....	1259
Perch.....	1505, 1577	<i>tænlops</i> .....	1144
Pike.....	1021	<i>trivrus</i> .....	1236
Sharks.....	33	<i>vivanet</i> .....	1257
Sunfish.....	1005	<i>bodianus</i> , <i>Bodianus</i> .....	1583
Surgeon.....	1691	<i>Cosyphus</i> .....	1583
Tang.....	1691	<i>Bodicon</i> .....	1867
Blue-breasted Darter.....	1076	<i>Bocostoma brachiale</i> .....	2698
Bluefin.....	472	<i>reticulatum</i> .....	2696
Bluefish, California.....	1410	<i>Boga</i> .....	1365
Bluefishes.....	945, 946	<i>Bogoslovius</i> .....	2574
Blue-gill.....	1005	<i>clarki</i> .....	2575
Blueheaded Sucker.....	171	<i>firmisquamis</i> .....	2575
Blue-spotted Sunfish.....	996	<i>Bola</i> .....	1455
Blunt-nosed Minnow.....	218	<i>Boleichthys</i> .....	1101
Shiner.....	984	<i>eos</i> .....	1102
Boar-fishes.....	1063	<i>exilis</i> .....	1103
Boar Grunt.....	1303	<i>fusiformis</i> .....	1101
Bobo.....	821	<i>warreni</i> .....	1103
Boca Dulce.....	29	<i>whippili</i> .....	1096
<i>Negra</i> .....	1837	<i>boleoides</i> , <i>Cottus</i> .....	1968
Bocaccio.....	1780	<i>Radulinus</i> .....	1919
Bocou.....	442, 450	<i>Uranidea</i> .....	1968
bocous, <i>Sardinia</i> .....	449	<i>Boleosoma</i> .....	1054
<i>Bodianus</i> .....	1143, 1581	<i>æopos</i> .....	1057
<i>acanthistius</i> .....	1147	<i>barrattii</i> .....	1102
<i>achigan</i> .....	1011	<i>camurum</i> .....	1060
<i>apua</i> .....	1174	<i>chlorosoma</i> .....	1060
<i>argyroleucus</i> .....	1433	<i>copelandi</i> .....	1040
<i>aya</i> .....	1265	<i>fusiformis</i> .....	1102

Page.		Page.		Page.
1234	<i>Boleosoma gracile</i> .....	1102	Bonito California.....	872
1583	<i>lepida</i> .....	1089	<i>Oceanic</i> .....	868
1583	<i>lepidum</i> .....	1089	Bonitos.....	871
1462	<i>longimanns</i> .....	1054	Bonnet-head.....	44
1142	<i>maculatum</i> .....	1057, 1077	Bony-fish.....	410, 433
1582	<i>mutatum</i> .....	1057	Bony Fishes.....	113
1146	<i>nigrum</i> .....	1056	Ganoïds.....	107
1433	<i>effulgens</i> .....	1058	Bony-tail.....	236
1024	<i>maculaticeps</i> ....	1058	Boohoo.....	891
1144	<i>mesæum</i> .....	1059	boops, <i>Caranx</i> .....	922
1146	<i>olmstedii</i> .....	1057	<i>Centaurus</i> .....	1755
1145, 1146	<i>vexillare</i> .....	1058	<i>Myctophum</i> .....	572
1145	<i>olmstedii brevipinnis</i> ....	1057	<i>Notropis</i> .....	268
849	<i>phlox</i> .....	1052	<i>Ostracion</i> .....	1755
1174	<i>podostemone</i> .....	1055	<i>Scopelus</i> .....	572
1433	<i>pottsi</i> .....	1083	<i>Trachurus</i> .....	922
1141	<i>punctulatum</i> .....	1091	Borborys.....	633
1582	<i>shumardi</i> .....	1047	borea, <i>Lucioperca</i> .....	1022
849	<i>etigmaeum</i> .....	1048	boreale, <i>Etheostoma</i> .....	1082
1584	<i>susane</i> .....	1059	borealis, <i>Alepidosaurus (Canlopis)</i> ..	597
1146	<i>tessellatum</i> .....	1046, 1057	<i>Alepisaurus</i> .....	596
1147	<i>variatum</i> .....	1070	<i>Amiurus</i> .....	137
1265	<i>whipplei</i> .....	1006	<i>Ammocætes</i> .....	11
1135, 1583	<i>boleosoma Gobius</i> .....	1102	<i>Arctozenns</i> .....	601
990	<i>bolli, Amelurus</i> .....	140	<i>Bathylagus</i> .....	2824, 2825
1443	<i>bollmani, Hippoglossina</i> .....	2621	<i>Chimæra</i> .....	95
1259	<i>Opsopceodus</i> .....	249	<i>Fierasfer</i> .....	2443
1144	<i>Scarus</i> .....	1646	<i>Icelinus</i> .....	1896
1236	<i>Bollmannia</i> .....	2237	<i>Lamargus</i> .....	57
1257	<i>chlamydes</i> .....	2238	<i>Manrolicus</i> .....	577
1583	<i>macropoma</i> .....	2239	<i>Notorhynchus</i> .....	18
1583	<i>ocellata</i> .....	2238	<i>Oligocottus</i> .....	2014
1867	<i>stigmatura</i> .....	2259	<i>Paralepis</i> .....	601
2698	<i>bombifrons, Lepomis</i> .....	1003	<i>Petromyzon</i> .....	13
2696	<i>Pomotis</i> .....	1003	<i>Pimelodus</i> .....	137
1365	<i>Bonaci Arará</i> .....	1174	<i>Pœcilichthys</i> .....	1082
2574	<i>Cardenal</i> .....	1173, 1174	<i>Scopelus</i> .....	577
2575	<i>de Piedra</i> .....	1172	<i>Sphyræna</i> .....	825
2575	<i>Gato</i> .....	1187	<i>Squirlus</i> .....	57
1455	<i>bonaci, Epinephelus</i> .....	1175	<i>Sudis</i> .....	601
1101	<i>Mycteroperca</i> .....	1174	Boregat.....	1867
1102	<i>xanthosticta</i> .....	1176	Boreocottus.....	1970
1103	<i>Serranus</i> .....	1175	<i>axillaris</i> .....	1981
1101	<i>Trisotropis</i> .....	1175	Boreogadus.....	2533
1103	<i>Bonapartia</i> .....	580, 2826	<i>polaris</i> .....	2534
1096	<i>pedaliota</i> .....	580, 2826	<i>salda</i> .....	2593
1968	<i>bonapartii, Scopelus</i> .....	557	Boreogaleus.....	32
1910	<i>bonariense, Hæmulon</i> .....	1297	Borer.....	7
1068	<i>bonariensis, Halatractus</i> .....	905	Borors.....	5
1054	<i>Seriola</i> .....	905	boreum, <i>Stizostedion canadense</i> ....	1022
1057	<i>bonasus, Bubalichthys</i> .....	164	boreus, <i>Esox</i> .....	628
1102	<i>Raja</i> .....	90	Borlae.....	963
1060	<i>Rhinoptera</i> .....	90	bosci, <i>Abramis crysoleucas</i> .....	251
1060	<i>Bone-fish</i> .....	411	<i>Atherina</i> .....	801
1046	<i>Bone Shark</i> .....	51	<i>Gobiosoma</i> .....	2259
1102	<i>Bonito</i> .....	869, 948	<i>Gobius</i> .....	2227, 2259

	Page.		Page.
bosci, Halatractus .....	905	brachycephalum, Siphostoma.....	769
Leuciscus .....	251	brachycephalus, Exocætus.....	733
Pimelepterus .....	1388	Syngnathus.....	769
Zonichthys .....	905	Uranichthys.....	382
boscianus, Chasmodon.....	2394	brachychir, Macrostoma .....	2826
boscii, Seriola .....	905	Myctophum .....	556
bosqui, Cyphosus .....	1388	brachychirus, Trachurops .....	911
bosquianus, Blennius .....	2394	Brachyconger .....	359
Chasmodon.....	2394	savanna .....	360
bosqui, Pimelepterus .....	1388	Brachydeuterus.....	1325
bostonensis, Catostomus .....	179	axillaris.....	1328
bostoniensis, Muræna .....	348	corvinaformis .....	1326
Botete.....	1731	leuciscus.....	1327
Bothragonus .....	2086	nitidus .....	1326
swanii.....	2086, 2088	Brachygenys.....	1307
Bothrocara .....	2475	chrysgyreus.....	1307
mollis .....	2476	tæniata.....	1308
pusilla .....	2476	Brachyistius.....	1490
Bothrolæmus.....	939	frenatus.....	1490
Bothus.....	2661	Brachymullus.....	858
maculatus.....	2660	Brachyopsis.....	2046, 2864
Boucane .....	1261	annæ .....	2043
boucardi .....	247	barkani.....	2044
.....	247	decayonis .....	2054
Pæcilia.....	695	dodecaedrus.....	2046
Pristipoma.....	1334	rostratus.....	2046, 2048
Rutilus .....	247	segallensis.....	2048
boulengeri, Mycteroperca .....	1171, 2856	verrucosus.....	2044
Bout de Tabac.....	1215	xyosternus .....	2043
bouvieri, Salmo clarkii .....	2819	Brachyospina.....	2032
mykiss .....	496	brachypoda, Gasterosteus pungitius	746
purpuratus .....	496	Pygosteus pungitius..	746
bovinum, Plectropoma .....	1193	Brachypomacentrus.....	1549
bovinus, Cyprinodon .....	673	Brachyprosope.....	2653
Hypoplectrus.....	1193	brachyptera, Echeneis .....	2272
unicolor.....	1193	Remora.....	2272
Bowfin .....	113	Rhamdia .....	151
Bowfin .....	111, 112	brachypterus, Holocentrus.....	852
bowmani, Plargyrus .....	283	Lutjanus.....	1268
Box, Tobacco.....	68	Neomænis.....	1268
Boxaodon .....	1365	Pimelodus .....	152
brachiale, Baostoma.....	2698	Remoropsis .....	2272
Sparisoma.....	1641	Thynnus.....	870
brachialis, Achirus .....	2698	Zygonectes.....	682
Ammocætes .....	14	brachyrhinchus, Acipenser .....	104
Balostoma.....	2698	brachyrhinus .....	1221
Petromyzon .....	14, 2745	creolus .....	1222
Scarus.....	1641	furcifer.....	1222
brachiatus, Diodon.....	1746	Brachysomphis.....	387
Brachioptilon .....	92	crocodilinus .....	388
hamiltoni.....	93	horridus .....	388
brachirus, Gymnotus .....	346	brachysomus, Calamus.....	1353
brachiusculus, Grammicolepis .....	974	Epinephelus .....	1154
brachyacanthus, Amiurus.....	141	Sparus.....	1353
brachycentrus, Gasterosteus.....	748	brachyurus, Oxydontichthys.....	385
Nauclerus .....	900	bradfordi, Porocottus .....	2862

Page.		Page.		Page.
769	Brama	958	brasilensis, Mugil	810, 814, 816
733	agassizii	959	Murena seu conger	403
769	brevoortii	959	Narcine	78, 2752
382	chilensis	960	Paralichthys	2626
2826	duseumieri	960	Plagusia	2709
556	orcini	960	Pseudorhombus	2626
911	parra	1586	Scorpena	1842
359	rafi	958, 959	Serranus	1221
360	raji	960	Thynnus	869
1325	sausurii	958	Torpedo	78
1328	brama, Cynædus	1360	Vomer	934
1326	Bramble Sharks	57	braytoni, N-tropis	204
1327	Bramidae	956	Bream	250, 1009, 1358, 1300
1326	Bramocharax	338	Bream, Blue	1005
1307	bransfordi	339	Copper-nosed	1005
1307	Bramopsis	1501	Redbreast	1001
1308	mento	1502	Bregmaceros	2528
1499	Branch Herring	426	atlanticus	2527
1490	branchialis, Ammocetes	14	atripinnis	2527
858	Petromyzon	14	bathymaster	2527
2046, 2864	Branchiostegus	2578	maclellandii	2526
2043	Branchiostoma	3, 4	Bregmacerotidae	2525
2044	californiense	4	Bresson	125
2054	californiensis	4	brevibarbe, Lepophidium	2485
2046	caribaëum	3	Ophidium	2485
2046, 2048	lanceolatum	3	brevicauda, Pomoxys	987
2048	lubricum	3	Salmo	493
2044	Branchiostomatidae	2	brevicaudata, Brevoortia tyrannus	434
2043	Branchiostomidae	2	Breviceps	116
2032	Branderius	373	breviceps, Evorthodus	2208
746	brandti, Arius	122	Gasterosteus	746
746	Cottus	1984	Larimus	1423
1549	Myoxocephalus	1984	Moxostoma	196
2653	Tachisurus	122	Pomotis	1003
2272	brandtii, Arius	2753	Ptychostomus	196
2272	branicki, Pomadasis	1333	brevidens, Gonostoma	579
151	Pristipoma	1334	brevimanus, Gorres	1377
852	branneri, Pœcilia	2834	Tetragonopterus	335
1268	bransfordi, Bramocharax	339	brevipes, Lycodes	2467
1268	Loricaria	158	brevipinna, Scymnus	57
152	Rhamdia	151	Somnioaus	57
2272	brasilianus, Gerres	1378	brevipinne, Ditrema	1499
870	brasiliense, Pristipoma	1320	Pristipoma	1341
682	brasiliensibus, Capenna	1311	brevipinnis, Biennius	2391
104	Guabi coara	1305	Boleosoma olmstedii	1057
1221	brasiliensis, Amblyopus	2264	Hypsoblennius	2390
1222	Centropristia	1221	Isaciella	1341
1222	Chirostoma	794	Orthopristia	1341
387	Chlorichthys	1591	Thynnichthys	869
388	Clupea	411	Thynnus	869
388	Conger	360	brevirostris, Chasmistes	183, 199
1353	Esox	723	Cololabis	726
1154	Hemiramphus	722	Gerres	1376
1353	Hippoglossus	2626	Hippocampus	776
385	Labrus	1591	Histiophorus	892
2862	Menidia	801	Hypoprion	41

	Page.		Page.
brevirostris, <i>Macrognathus</i> .....	723	<i>Brosomphycis ventralis</i> .....	2503
<i>Saurus</i> .....	533	<i>Brotula</i> .....	2500
<i>Scombresox</i> .....	726	<i>barbata</i> .....	2500
<i>Syngnathus</i> .....	705	<i>Brotullidæ</i> .....	2498
brevirostrum, <i>Acipenser</i> .....	106	<i>Brotulinae</i> .....	2498
<i>Hæmulon</i> .....	1300	<i>Brotuloid Fishes</i> .....	2498
brevis, <i>Acanthurus</i> .....	1091	<i>broussoneti</i> , <i>Gobioides</i> .....	2264
<i>Amblyopus</i> .....	2263	<i>broussonetii</i> , <i>Acanthurus</i> .....	1691
<i>Atherlichthys</i> .....	2840	<i>Umbrina</i> .....	1466
<i>Centropomus</i> .....	1125	<i>broussonnetii</i> , <i>Gobiodes</i> .....	2263
<i>Cephalus</i> .....	1754	<i>Brown Cat</i> .....	142
<i>Cetengraulis</i> .....	450	<i>Hind</i> .....	1142
<i>Engraulis</i> .....	450	<i>Rockfish</i> .....	1817
<i>Tyntlastes</i> .....	2262	<i>browni</i> , <i>Atherina</i> .....	443
brevispinis, <i>Sebasticthys</i> .....	1788	<i>Engraulis</i> .....	443
<i>Sebastes</i> .....	1787	<i>Hemirhamphus</i> .....	723
brevoorti, <i>Argyrosus</i> .....	936	<i>Solea</i> .....	2701
<i>Euleptorhamphus</i> .....	724	<i>Stolephorus</i> .....	443
Brevoortia.....	433	<i>Vomer</i> .....	934
<i>tyrannus</i> .....	433	<i>Brown-winged Sea-robin</i> .....	2167
<i>aurea</i> .....	434	<i>brucus</i> , <i>Squalus</i> .....	58
<i>brevicaudata</i> .....	434	<i>brunnea</i> , <i>Maynea</i> .....	2476
<i>patrenus</i> .....	434	<i>brunneus</i> , <i>Amiurus</i> .....	142
brevoortii, <i>Brama</i> .....	959	<i>Catulus</i> .....	24
bricei, <i>Chaetodon</i> .....	1678	<i>Gobius</i> .....	2218
bristolæ, <i>Emmion</i> .....	2375	<i>Ilyophis</i> .....	350
Broad Shad.....	1372	<i>Serranus</i> .....	1175
<i>Whitefish</i> .....	464	<i>Trisotropis</i> .....	1175
Broad-head, <i>Grubber</i> .....	447	<i>Brycon</i> .....	337
broccus, <i>Belistes</i> .....	1716	<i>dentex</i> .....	337
<i>Monacanthus</i> .....	1716	<i>striatulus</i> .....	337
Brochet de Mer.....	1118	<i>bryoporus</i> , <i>Spratelloides</i> .....	422
brodanius, <i>Cottus</i> .....	2066	<i>Bryostemma</i> .....	2408
Brook Lampreys.....	12	<i>nugator</i> .....	2410
<i>Silverside</i> .....	805	<i>polyactocephalum</i> ..	2408, 2409
<i>Stickleback</i> .....	744	<i>bryosus</i> , <i>Blennicottus globiceps</i> ...	2017
<i>Sucker</i> .....	178	<i>Bryaseteres</i> .....	2328
<i>Trout</i> .....	506	<i>pluniger</i> .....	2328
<i>Trout of western Oregon</i> ...	501	<i>Bryasophilus</i> .....	2329, 2330
Brosme.....	2561	<i>Bryttus</i> .....	995
<i>brosme</i> .....	2561	<i>albulus</i> .....	1007
<i>brosme</i> , <i>Brosme</i> .....	2561	<i>fasciatus</i> .....	993
<i>Brosmius</i> .....	2561	<i>gloriosus</i> .....	994
<i>Enchelyopus</i> .....	2561	<i>humilis</i> .....	1004
<i>Gadus</i> .....	2561	<i>mineopas</i> .....	996
<i>brosmiana</i> , <i>Lota</i> .....	2551	<i>oculatus</i> .....	1004
<i>Brosminæ</i> .....	2532	<i>punctatus</i> .....	998
<i>Brosmius</i> .....	2561	<i>reticulatus</i> .....	998
<i>brosme</i> .....	2561	<i>signifer</i> .....	996
<i>flavesceus</i> .....	2561	<i>unicolor</i> .....	1001
<i>flavesny</i> .....	2561	<i>Bubalichthys</i> .....	163
<i>marginatus</i> .....	2502	<i>altus</i> .....	165
<i>vulgaris</i> .....	2561	<i>bonasus</i> .....	164
<i>Brosomphycinæ</i> .....	2498	<i>bubalinus</i> .....	165
<i>Brosomphycis</i> .....	2502	<i>bubalus</i> .....	516
<i>marginatus</i> .....	3502	<i>niger</i> .....	164

Page.		Page.		Page.	
2503	Bubalichthys urus.....	164	Burnstickle.....	747	
2500	bubalina, Cliola.....	273	Burr-fish, Common.....	1748	
2500	Cyprinella.....	273	Burr-fishes.....	1747	
2498	bubalinus, Bubalichthys.....	165	Burrito.....	1333, 1327	
2498	Lenciscus.....	273	Burritos.....	1325	
2498	Notropis.....	273	Burro.....	1332	
2264	bubalis, Cottus.....	1972	Blanco.....	1328	
1691	Myoxocephalus.....	1971	Burros.....	1329	
1466	bubalus, Amblyodon.....	165	burtonianus, Cyprinus.....	2798	
2263	Bubalichthys.....	165	busculus, Prionodes.....	1211	
142	Catostomus.....	165	Butirinus maderaspatensis.....	415	
1142	Ichthyobus.....	164	butleri, Pœcilia.....	691	
1817	Ictiobus.....	164	butleriannus, Pœciliichthys.....	1102	
443	Bubbler.....	1484	Butter-fish.....	967, 1144, 2416, 2419	
443	buccanella, Lutjanus.....	1262	utter-fishes.....	965	
723	Mesoprion.....	1262	Butterfly.....	1677	
2701	Neomænis.....	1261	Butterfly-fishes.....	1660, 1672	
443	buccata, Eriocymba.....	302	Butterfly Ray.....	86	
934	bucciferus, Labrisomus.....	2363	Butyrinus.....	411	
2167	bucco, Moxostoma.....	190	banana.....	411	
58	Ptychostomus.....	191	Bythites.....	2504	
2476	bucculentus, Chonophorus.....	2236	fuscus.....	2504	
142	buchanani, Notropis.....	2800	Bythitinae.....	2498	
24	Buñalo Cod.....	1875			
2218	Common.....	163	caballa, Cybium.....	876	
350	fishes.....	163	Scomberomorus.....	876	
1175	Mongrel.....	164	Caballerote.....	1255, 1257	
1175	Razor-backed.....	104	caballerote, Anthias.....	1257	
337	Red mouth.....	163	Lutjanus.....	1257	
337	Small-mouthed.....	164	Mesoprion.....	1257	
337	Sucker-mouthed.....	164	caballus, Caraux.....	921	
422	buffalo, Acipenser.....	106	Cabezon.....	1423, 1889, 2321	
2408	bufo, Lophius.....	2316	Smooth.....	2012	
2410	Scorpana.....	1849	Cabezones.....	1889	
2408, 2409	Bugara.....	1508	Cabezote.....	790	
2017	Bugfish.....	433	Cabezuda, Lisa.....	811	
2328	Bull Red-fish.....	1453	caboverdianus, Ginglymostoma.....	26	
2328	Bull Trout.....	507	Cabra Mora.....	1152	
2329, 2330	bullaris, Cyprinus.....	221	Cabrilla.....	1158, 1197, 1832	
995	Semotilus.....	222	Calamaria.....	1184	
1007	bulleri, Prionodes.....	1213	de Raizero.....	1171	
993	Serranus.....	1214	Pirlita.....	1181	
994	Bullhead.....	1050	Cabrilla, Spotted.....	1196	
1004	Bullhead, Bear Lake.....	1054	Cabrillas Verdes.....	1194	
996	Black.....	141	Cabrillo de Astillero.....	1176	
1004	Common.....	140	Cachucho.....	1282	
998	Prickly.....	1944	Caçonetta.....	40	
998	Rocky Mountain.....	1949	Cæcilia.....	373	
996	Bullhead Shark.....	20	cælolepis, Centroscymnus.....	55	
1001	Sharks.....	19	cænicola, Pœcilia.....	641	
163	Bullon.....	1650	cænosus, Pimelodus.....	140	
165	Bumper.....	938	cærulea, Cliola.....	277	
164	buniva, Ballistes.....	1701, 1711	Clupea.....	421	
165	Burbots.....	2550	Codoma.....	277	
510	Burgall.....	1577	Coryphæna.....	1653	
164	burgall, Ctenolabrus.....	1577	Maletta.....	423	



	Page.		Page.
<i>cærulea</i> , <i>Novacula</i> .....	1653	<i>California Barracuda</i> .....	826
<i>Tautoga</i> .....	1577	<i>Big Skate</i> .....	72
<i>cæruleatus</i> , <i>Acronurus</i> .....	1691	"Bluefish" .....	1410
<i>cæruleo-aureus</i> , <i>Harpe</i> .....	1585	<i>Bonito</i> .....	872
<i>cærneopunctatus</i> , <i>Acara</i> .....	1514	<i>Conger Eel</i> .....	395
<i>cærulescens</i> , <i>Arins</i> .....	129	<i>Dogfish</i> .....	54
<i>Galeichthys</i> .....	2776	<i>Hagfish</i> .....	6
<i>Hexanematchichthys</i> ...	129	<i>Herring</i> .....	422
<i>Pimelodus</i> .....	135	<i>Jewfish</i> .....	1137
<i>cæruleus</i> , <i>Acanthurus</i> .....	1691	<i>Lancelet</i> .....	4
<i>Carcharhinus</i> .....	37	<i>Pompauo</i> .....	967
<i>Carcharias</i> .....	37	<i>Redfish</i> .....	1585
<i>Cheonida</i> .....	232	<i>Sardine</i> .....	423
<i>Clupanodon</i> .....	423	<i>Smelt</i> .....	866
<i>Ctenolabrus</i> .....	1577	<i>Sole</i> .....	2613
<i>Cyclopterus</i> .....	2097	<i>Stickleback</i> .....	751
<i>Notropis</i> .....	277	<i>Stingray</i> .....	89
<i>Photogenis</i> .....	277	<i>Tomcod</i> .....	2539
<i>Pseudoscarius</i> .....	1654	<i>Torpedo</i> .....	77
<i>Scarus</i> .....	1652, 1654	<i>Whiting</i> .....	1476
<i>Squalus</i> .....	232	<i>californica</i> , <i>Alausa</i> .....	423
<i>Squalus</i> .....	33	<i>Morrhua</i> .....	2539
<i>Teuthis</i> .....	1601	<i>Squatina</i> .....	59
<i>cærneopunctatus</i> , <i>Equidens</i> .....	1514	<i>Tetronarce</i> .....	77
<i>Cæsar</i> .....	1308	<i>Torpedo</i> .....	78
<i>Cæsiomorus</i> .....	939	<i>Uropsetta</i> .....	2626
<i>Cæsiosoma californiense</i> .....	1391	<i>californiens</i> , <i>Cypsilurus</i> .....	2836
<i>cæsius</i> , <i>Anisotremus</i> .....	1316	<i>Exocoetus</i> .....	730, 740
<i>Pomadasy</i> .....	1317	<i>Gadus</i> .....	2539
<i>Cagon de lo Alto</i> .....	1277	<i>Galeus</i> .....	30
<i>Callieu</i> .....	429	<i>Halichares</i> .....	1601
<i>Callieu-tassart</i> .....	432	<i>Hippoglossus</i> .....	2626
<i>Calman</i> .....	2216	<i>Mustelus</i> .....	30
<i>Caji</i> .....	1258	<i>Myllobatis</i> .....	89
<i>Calafate</i> .....	1711	<i>Oxyjulis</i> .....	1601
<i>Calamaria</i> , <i>Cabrilla</i> .....	1184	<i>Paralichthys</i> .....	2625, 2626
<i>Calamus</i> .....	1347, 1349	<i>Pseudojulis</i> .....	1601
<i>arctifrons</i> .....	1355	<i>Pseudorhombus</i> .....	2626
<i>hajonado</i> .....	1352, 1353	<i>Stereolepis</i> .....	1138
<i>brachysomus</i> .....	1353	<i>californiense</i> , <i>Branchiostoma</i> .....	4
<i>calamus</i> .....	1349	<i>Cæsiosoma</i> .....	1391
<i>leucostens</i> .....	1353	<i>Myctophum</i> .....	572
<i>macrops</i> .....	1350, 1354	<i>Siphostoma</i> .....	764
<i>medius</i> .....	1356	<i>californiensis</i> , <i>Atherinichthys</i> .....	807
<i>megacephalus</i> .....	1350, 1351	<i>Atherinops</i> .....	806
<i>penna</i> .....	1354, 1355	<i>Branchiostoma</i> .....	4
<i>pennatula</i> .....	1351	<i>Chilomycterus</i> .....	1751
<i>plummatula</i> .....	1352	<i>Cyprinodon</i> .....	674
<i>proridens</i> .....	1350	<i>Diapterus</i> .....	1370
<i>taurinus</i> .....	1354	<i>Doryichthys</i> .....	774
<i>calamus</i> , <i>Calamus</i> .....	1349	<i>Doryrhamphus</i> .....	773
<i>Chrysophrys</i> .....	1350	<i>Eucinostomus</i> .....	1369
<i>Pagellus</i> .....	1350	<i>Gerres</i> .....	1370
<i>calcarata</i> , <i>Scorpena</i> .....	1854	<i>Medialuna</i> .....	1391
<i>Calico Bass</i> .....	987	<i>Ophæurus</i> .....	384
<i>California Anchovy</i> .....	448	<i>Otolithus</i> .....	1413

Page.		Page.		Page.
826	californiensis, Polynemus.....	829	Calloptilum.....	2526
72	Scorpiæ.....	1301	mirum.....	2527
1410	Syngnathus.....	764	Callorhynchus atlanticus.....	95
872	Typhlogobius.....	2262	centrinus.....	95
395	Xenichthys.....	1286	Callyodon.....	1621, 1642, 1651
54	Xenistius.....	1286	aeropunctatus.....	1624
6		2541	flavescens.....	1640
422	callarias, Gadus.....	508	psittacus.....	1638
1137	Callaus.....	1455	nustus.....	1624
4		378	callyodon, Cyclopterus.....	2110
967	Callechelys.....	378	Liparis.....	2111
1585	murena.....	379	Neoliparis.....	2110
423	peninsula.....	2360	calopteryx, Serranus.....	1213
806	Calliclinus.....	1621, 1642, 1644, 1650	Calotomus.....	1626
2613	dentiens.....	1623	xenodon.....	1626
751	gibbosus.....	1296	calva, Amla.....	113
89	lineatus.....	1651	Calycilepidotus.....	1936
2539	retractus.....	1623	lateralis.....	1899
77	Calliodon, Liparis.....	2120	spinosus.....	1937
1476	Callionymidae.....	2184	Camarina.....	1381
423	Callionymus.....	2185	nigricans.....	1382
259	bairdi.....	2185	camopardalis, Mycteroperca tigris	1187
59	calliurus.....	2187	Serranus.....	1187
77	himantophorus.....	2186	campbelli, Moxostoma.....	186
78	pauciradiatus.....	2188	Salmo.....	508
2626	pelagicus.....	2184	Campbellite.....	987
2836	Callipteryx, Campostoma.....	206	campechianus, Lutjanus.....	1285
730, 740	Callisema, Cliola.....	273	Mesoprion.....	1295
2539	Codoma.....	273	camperi, Scombreox.....	725
30	Eplisma.....	273	Campostoma.....	204
1601	Notropis.....	272	anomalum.....	205
2626	Callisoma, Herpetoichthys.....	384	callipteryx.....	206
30	Callistia, Cliola.....	276	dubium.....	206
89	Codoma.....	276	formosulum.....	206
1601	Callistina, Notropis.....	276	gobionum.....	206
2625, 2626	Photogenis.....	276	hippops.....	206
1601	Callinre, Cliola.....	275	mormyrus.....	206
2626	Cyprinella.....	275	nasutum.....	206
1138	Etheostoma.....	1011	ornatum.....	205
4	Mycteroperca.....	1186	pricei.....	205, 2796
1391	Calliurus.....	995, 1010	prolixum.....	206
572	diaphanus.....	996	Campostominae.....	202
764	floridensis.....	992	Campylodon.....	614
807	formosus.....	996	fabricii.....	615
806	longulus.....	996	camtschatica, Lampetra.....	13
4	melanops.....	992	camtschaticus, Entosphenus.....	2745
1751	microps.....	996	Pteromyzon.....	2745
674	murius.....	996	marinus.....	2745
1370	punctulatus.....	992, 1011	camura, Cliola.....	280
774	Calliurus, Callionymus.....	2187	Vaillantia.....	1060
773	Ioglossus.....	2193	camurum, Boleosoma.....	1060
1369	Trisotropis.....	1186	Etheostoma.....	1076
1370	Callogobius.....	2210	camurus, Nanostoma.....	1076
1391	callolepis, Harengula.....	430	Notropis.....	279
384	callopterus, Cypsilurus.....	2836	camurus, Peccilichthys.....	1076
1413	Exocoetus.....	740	Cafia-bota.....	19

	Page.		Page.
canada, Elacato.....	948	capenna, Hæmulon .....	1311
canadonæ, Stizostedion .....	1022	Hæmylum .....	1311
borium.....	1022	Serranus .....	1311
griseum .....	1022	capillaris, Argyrosia .....	936
canadensis, Luciopecca .....	1022	Zona.....	936
Salmo.....	507	capiliatus, Clinus.....	2362
canadus, Gasterosteus .....	948	Labrisomus .....	2362
Rachycentron .....	948	capistratus, Baliates .....	1704
canaliculatus, Icelus.....	1917	Chaetodon .....	1677
canariensis, Clinus.....	2302	Paclymnathus .....	1704
canidissima, Leptocephalus.....	354	Sarothrudus .....	1678
Candil .....	846	Tetrodon .....	1742
Candlefish .....	521	Capitaino.....	1579
Cane di Mare .....	48	capite, Eleotris plagioplateo.....	2201
canescens, Chaetodon.....	1688	Labrus obtuso .....	1609
Zanclus.....	1688	capiti lævi, Biennius.....	2438
canina, Amla.....	113	capito, Poronitra.....	840
caninianus, Scopelus.....	570	capitulinus, Mugil .....	2811
caninus, Caranx.....	921	caprinus, Argyrops .....	1345
Lachnolaimus.....	1580	Ballistes .....	1702
Pagellus.....	1352	Catoptomus .....	168
Canis carcharias .....	38	Otrynter .....	1345
canis, Mustelus.....	29	Stenotomus .....	1345
Salmo.....	479	Capriscus .....	1699, 1700
Squalus .....	29	capriscus, Bullata .....	1701
cauna, Hæmulon.....	1297, 1299	mulum dentibus minutis .....	1720
Cannorhynchus .....	756	caprodes, Etheostoma .....	1027
cantharinum, Pristipoma .....	1340	Percina .....	1026
cantharinus, Orthopristis.....	1339, 1340	manitou.....	1028
Cantharus nigromaculatus.....	987	zebra .....	1027
Cantherines.....	1713	Sciæna .....	1027
carole.....	1713	Caproidæ.....	1663
pullus.....	1713	Caproidea .....	1663
Canthidormis .....	1705	Caprophonus.....	1664
maculatus .....	1706, 1707	aurora .....	1665
sobaco.....	1705	capros, Antigonis .....	1665
sufflamen .....	1706	caramura, Murenophis .....	395
willughbelli.....	1707	Carangichthys .....	916, 917, 922
Canthigaster.....	1741	Carangidæ.....	895
caudicinctus.....	1742	Carangoides.....	928
lobatus .....	1732	blepharis .....	932
punctatissimus .....	1741	cibi.....	920
rostratus .....	1741	dorsalis.....	930
Canthigasteridæ.....	1740	gallichthys .....	932
Canthirhynchus.....	2088	iridinus .....	919
monopterygius.....	2092	orthogrammus.....	928
Canthorhinus .....	1713	carangoides, Seriolophus .....	895
Cantileña, Mojarra .....	1360	Carangops.....	912
Capelin .....	520	heteropygus.....	913
capensis, Aledon.....	1754	secundus .....	914
Carcharodon .....	50	carangua, Caranx .....	920
Elops.....	410	Carangus .....	915, 916
Scorpiæna.....	1833	chrysos .....	921
Sebastes.....	1833	esculentus .....	921
Sebastodes .....	1833	fallax .....	923
Capenna brasiliensis .....	1311	hippos .....	921

Page.		Page.		Page.
.....	1311	.....	929	
.....	1311	.....	930	
.....	1311	.....	928	
.....	936	.....	923	
.....	936	.....	923	
.....	2362	.....	910	
.....	2362	.....	921, 2846	
.....	1704	.....	927	
.....	1677	.....	912	
.....	1704	.....	928	
.....	1678	.....	908	
.....	1742	.....	923	
.....	1579	.....	916	
.....	2201	.....	908	
.....	1609	.....	908	
.....	2438	.....	914	
.....	840	.....	923	
.....	2841	.....	911	
.....	1345	.....	934	
.....	1702	.....	934	
.....	168	.....	927	
.....	1345	.....	928	
.....	1345	.....	926	
.....	1609, 1700	.....	910	
.....	1701	.....	910	
.....	1720	.....	910	
.....	1027	.....	910	
.....	1026	.....	910	
.....	1028	.....	918	
.....	1027	.....	921	
.....	1027	.....	952	
.....	1663	.....	911	
.....	1663	.....	2579	
.....	1664	.....	340	
.....	1665	.....	341	
.....	395	.....	340	
.....	916, 917, 922	.....	2497	
.....	895	.....	341	
.....	928	.....	341	
.....	932	.....	201	
.....	920	.....	201	
.....	930	.....	1145	
.....	932	.....	1027	
.....	919	.....	1300	
.....	928	.....	106	
.....	895	.....	2534	
.....	912	.....	2535	
.....	913	.....	509	
.....	914	.....	919	
.....	920	.....	1300	
.....	915, 916	.....	1283	
.....	921	.....	28	
.....	921	.....	33, 35, 37, 2747	
.....	923	.....	36	
.....	921	.....	40	
.....	923	.....	37	
.....	921	.....		

	Page.		Page
<b>Carcharhinus</b>		<b>Cardenal</b>	1108
<i>cerdale</i>	2746, 2747	<i>Bonaci</i>	1173, 1174
<i>commersoni</i>	38	<i>Mojarra</i>	850
<i>falciformis</i>	36	<b>Cardinal fishes</b>	1165
<i>fronto</i>	39	<i>cardinalis</i> , <i>Serranus</i>	1174
<i>glancus</i>	33	<i>Trisotropis</i>	1174
<i>henlei</i>	37, 2746	<b>Cardonniera</b>	1837
<i>lamia</i>	38	<b>Caroliparis</b>	2114, 2115
<i>lamiella</i>	37	<b>Caremitra</b>	2129, 2130, 2131
<i>leucos</i>	38	<b>Carenchelyi</b>	343
<i>limbatus</i>	40	<b>Careproctus</b>	2129, 2130, 2131
<i>milberti</i>	37	<i>colletti</i>	2131
<i>nicaraguensis</i>	39, 2747	<i>ectenes</i>	2136, 2866
<i>obscurus</i>	35	<i>gelatinosus</i>	2134, 2135
<i>oxyrhynchus</i>	40, 2747	<i>melanurus</i>	2135, 2867
<i>perezi</i>	36	<i>ostentum</i>	2134
<i>platyodon</i>	39	<i>phasma</i>	2132
<i>platyrhynchius</i>	36	<i>ranula</i>	2134
<i>remotus</i>	37	<i>reinhardi</i>	2133, 2134
<i>velox</i>	2747, 2748	<i>simus</i>	2131
<b>Carcharias</b>	33, 46	<i>spectrum</i>	2133
<i>americanus</i>	47	<i>caribbeum</i> , <i>Branchiostoma</i>	3
<i>atwoodi</i>	50	<i>caribæus</i> , <i>Diplodus</i>	1360
<i>cæruleus</i>	37	<i>Sargus</i>	1360
<i>falcijunius</i>	35	<i>caribbæa</i> , <i>Belone</i>	717
<i>fronto</i>	39	<i>caribbæus</i> , <i>Chloroscombrus</i>	938
<i>glaucus</i>	33	<i>Halicichthys</i>	2741
<i>griseus</i>	47	<i>Tylosurus</i>	717
<i>henlei</i>	37	<i>carinatum</i> , <i>Acanthosoma</i>	1754
<i>isodon</i>	42	<i>Siphostoma</i>	763
<i>lalandi</i>	43	<i>carinatus</i> , <i>Diodon</i>	1754
<i>lamia</i>	38	<i>Labichthys</i>	368
<i>leucos</i>	38	<i>Placopharynx</i>	198
<i>limbatus</i>	40	<i>Salmo</i>	493
<i>littofalls</i>	40, 2748	<i>carminale</i> , <i>Tripterygium</i>	2350
<i>longurio</i>	42	<i>carminalis</i> , <i>Eunoanectes</i>	2350
<i>microps</i>	40	<i>carminatus</i> , <i>Cælorhynchus</i>	2588
<i>milberti</i>	37	<i>Macrurus</i>	2589
<i>mulleri</i>	40	<i>Macrurus (Cælorhynchus)</i>	2589
<i>obscurus</i>	35	<i>carinatus</i> , <i>Sebastichthys</i>	1825
<i>oxyrhynchus</i>	41	<i>Sobnostodes</i>	1824
<i>porosus</i>	37	<i>carneus</i> , <i>Acronurus</i>	1692
<i>punctatus</i>	42	<i>Gobiesox</i>	2337
<i>terra-novæ</i>	43	<i>Sicyases</i>	2337
<i>trigris</i>	49	<i>carole</i> , <i>Cantherines</i>	1713
<i>verus</i>	50	<i>carolina</i> , <i>Argentina</i>	410
<b>carcharias</b> , <i>Canis</i>	38	<i>Atherina</i>	791
<i>Carcharodon</i>	50	<i>Trigla</i>	2156, 2172
<i>Squalus</i>	38, 50	<b>Carolina Whiting</b>	1474
<b>Carchariidae</b>	46	<i>caroline</i> , <i>Potamocottus</i>	1952
<b>Carcharodon</b>	50	<i>carolinensis</i> , <i>Ballistes</i>	1701
<i>capensis</i>	50	<i>Cestrens</i>	1409
<i>carcharias</i>	50	<i>Clupea</i>	434
<i>rondéleti</i>	50	<i>Cynoscion</i>	1409
<i>smithi</i>	50		
<b>Carcharodontinae</b>	47		

Page.		Page.		Page.
1108		2218	carolinensis Gobina.....	1505
1173, 1174		988	Hyperistius.....	356
850		117	Mystus.....	844
1195		1409	Otolithus.....	844
1174		902	Seriola zonata.....	959
1174		2378	carolinus, Blennius.....	396
1837		944	Gasterosteus.....	960
2114, 2115		1578	Labrus.....	556
20, 2130, 2131		2379	Pholis.....	11
343		2858	Priacanthus.....	2804
20, 2130, 2131		2156, 2157	Prionotus.....	556
2131		956	Pteraclis.....	11
2136, 2866		944	Trachinotus.....	436
2134, 2135		944	Trachinotus.....	111
2135, 2867		167	Carp, Lake.....	1856
2131		166	Sucker.....	1856
2132		165	Suckers.....	31
2134		166	carpio, Carpiodes.....	22
2133, 2134		166, 190	Catostomus.....	134
2131		675	Cyprinodon.....	142
2133		201	Cyprinus.....	134
3		190	Moxostoma.....	138
1360		165	Carpoides.....	134
1360		166	bison.....	101
717		166	carpio.....	2788
908		107	cutisanserinus.....	137
2741		167, 168	cyprinus.....	137
717		167	damalis.....	137
1754		166	difformis.....	137
763		167	grayi.....	142, 143
1754		166	nummifer.....	143
368		167	selene.....	140
198		165	taurus.....	140
493		167	thompsoni.....	101
2350		167	tumidus.....	144
2350		164	urus.....	134, 138
2588		168	vacca.....	2788
2589		167	vellifer.....	130, 143
yn-		165	vitulus.....	554
2589		160	Catablemeia.....	2504
1825		161, 199	Catetyx.....	2505
1824		231	rubrirostris.....	1322
1692		2589	Catalina.....	1684
2337		2590	Catalineta.....	1682
2337		1152	Catalinetas.....	1237, 1238
1713		311	Catalufa.....	978
410		312	de lo Alto.....	1238
791		100	cataufa, Priacanthus.....	1236
2156, 2172		2732	Catalufas.....	977
1474		104	Deep-water.....	781
1952		1509	Cataphracti.....	2067
1701		1509	Cataphractus schoneveldii.....	107
1409		1508, 1509	cataphractus, Acipenser.....	2065, 2067
434		938	Agonus.....	2067
1409		937	Aspilophorus.....	2053, 2066
			Coitus.....	

	Page.		Page.
cataphractus, <i>Gasterosteus</i> .....	749	<i>Catostomus commersoni</i> .....	178
<i>aculeatus</i> .....	750	<i>communis</i> .....	179
<i>Scaphirhynchus</i> .....	107	<i>congestus</i> .....	192
cataractæ, <i>Ceratichthys</i> .....	306	<i>cypho</i> .....	184
<i>Goblo</i> .....	306	<i>discobolus</i> .....	172, 175, 2791
<i>Rhinichthys</i> .....	306	<i>duquesni</i> .....	193
<i>dulcis</i> .....	306	<i>duquesnii</i> .....	198
cataractus, <i>Leucosomus</i> .....	221	<i>elongatus</i> .....	169
catena, <i>Bussozetes</i> .....	2509	<i>erythrurus</i> .....	193
catenata, <i>Echidna</i> .....	403	<i>fasciatus</i> .....	187
<i>Muræna</i> .....	403	<i>fasciolaris</i> .....	186
<i>Pœcilia</i> .....	648	<i>fecundus</i> .....	180, 2794
<i>Xenisma</i> .....	648	<i>flexuosus</i> .....	179
catenatus, <i>Fundulus</i> .....	648	<i>forsterianus</i> .....	176
<i>Gymnothorax</i> .....	403	<i>generosus</i> .....	170
catenula, <i>Murænophis</i> .....	403	<i>gibbosus</i> .....	186
catesbæi, <i>Scarus</i> .....	1638	<i>gila</i> .....	180
<i>Sparisoma</i> .....	1638	<i>gracilis</i> .....	179
catesbæi, <i>Pomotis</i> .....	1010	<i>griseus</i> .....	175, 2791
catesby, <i>Scarus</i> .....	1638	<i>guzmaniensis</i> .....	171
catesbyi, <i>Sparisoma</i> .....	1638	<i>hudsonius</i> .....	176
Catfish of the Lakes.....	137	<i>insignis</i> .....	180
Catfish, Bermuda.....	882	<i>labiatus</i> .....	177, 2792
Sea.....	118, 119, 128	<i>lactarius</i> .....	175
Small .....	140, 141	<i>latipinnis</i> .....	174, 2790
Catfishes .....	114, 115	<i>lesueurii</i> .....	195
<i>Ga<sup>o</sup>-topsall</i> .....	116	<i>longirostris</i> .....	176
<i>catharina</i> , <i>Fristipoma</i> .....	1323	<i>longirostrum</i> .....	176
<i>catheplateo</i> , <i>Ostracion oblongus</i> ..	1728	<i>macrocheilus</i> .....	178
<i>Cathetostoma albigutta</i> .....	2313	<i>macrolopidotus</i> .....	194
<i>Cathorops</i> .....	133, 2788	<i>mæulosus</i> .....	181
<i>gulosus</i> .....	133, 2788	<i>megastomus</i> .....	181
<i>hypophthalmus</i> .....	133, 2788	<i>melanops</i> .....	187
<i>Catochaenum</i> .....	1373	<i>melanotus</i> .....	206, 218, 322
<i>Catonotus</i> .....	1066	<i>nanomyzon</i> .....	177
<i>fasciatus</i> .....	1098	<i>nebulifer</i> .....	171
<i>fiacillatus</i> .....	1098	<i>nebuliferus</i> .....	171
<i>kennicotti</i> .....	1098	<i>nigricans</i> .....	181
<i>lineatus</i> .....	1099	<i>etowaanus</i> .....	181
<i>Catostomida</i> .....	161	<i>occidentalis</i> .....	178, 2793
<i>Catostomiæ</i> .....	162	<i>onelda</i> .....	193
<i>Catostomus</i> .....	173, 174	<i>pallidus</i> .....	179
<i>alticolus</i> .....	179	<i>planiceps</i> .....	181
<i>anisurus</i> .....	190	<i>plebeius</i> .....	171
<i>areopus</i> .....	172	<i>pocatello</i> .....	175
<i>ardens</i> .....	179	<i>reticulatus</i> .....	179
<i>aureolus</i> .....	193, 196	<i>retropluvius</i> .....	175, 2791
<i>aurora</i> .....	176	<i>rex</i> .....	177, 2792
<i>bernardini</i> .....	178	<i>rhothæcus</i> .....	181
<i>bostonensis</i> .....	179	<i>rimiculus</i> .....	2792
<i>bubalus</i> .....	165	<i>snyderi</i> .....	2792
<i>caprinus</i> .....	168	<i>sucklii</i> .....	179
<i>carpio</i> .....	166, 190	<i>tahoensis</i> .....	177
<i>catostomus</i> .....	176, 2792	<i>teres</i> .....	179
<i>chloropteron</i> .....	179	<i>texanus</i> .....	192
<i>clarki</i> .....	172	<i>tsiltoosensis</i> .....	2793

Page.		Page.		Page.
178	<i>Catostomus tuberculatus</i> .....	186	<i>caudispinosus</i> , <i>Scopelus</i> .....	55
179	<i>ntawana</i> .....	179	<i>Caulurehus</i> .....	2327
192	<i>velifer</i> .....	167	• <i>maandricus</i> .....	2328
184	<i>xanthopus</i> .....	181	<i>caulias</i> , <i>Sigmistes</i> .....	2863
172, 175, 2791	<i>catostomus</i> , <i>Catostomus</i> .....	176	<i>Caulolatilina</i> .....	2275
193	<i>Cyprinus</i> .....	176	<i>Caulolatilus</i> .....	2276
198	<i>Phenacobius</i> .....	304	<i>affinis</i> .....	2277
169	<i>Cats</i> , <i>Channel</i> .....	133	<i>anomalus</i> .....	2277
193	<i>Mud</i> .....	142	<i>chrysops</i> .....	2277
187	<i>Stone</i> .....	143	<i>cyanops</i> .....	2278
186	<i>Catulus</i> .....	23, 24	<i>microps</i> .....	2277
180, 2794	<i>brunneus</i> .....	24	<i>princeps</i> .....	2276, 2277
179	<i>cephalus</i> .....	25	<i>Caulepis</i> .....	838
176	<i>retifer</i> .....	25	<i>longidens</i> .....	839, 2842
170	<i>uter</i> .....	25, 2745	<i>Caulophryne</i> .....	2734
186	<i>xanirus</i> .....	24	<i>jordani</i> .....	2735
180	<i>catulus</i> , <i>Ameiurus nebulosus</i> .....	141	<i>Caulophrynine</i> .....	2728
179	<i>Evorthodus</i> .....	2218	<i>Caulopus</i> .....	594, 596
175, 2791	<i>Gobius</i> .....	2218	<i>caulopus</i> , <i>Homesthes</i> .....	2394
171	<i>Pimelodus</i> .....	141	<i>caurinus</i> , <i>Cyprinus (Leuciscus)</i> .....	220
176	<i>catus</i> , <i>Ameiurus</i> .....	138	<i>Leucosomus</i> .....	220
180	<i>Amiurus</i> .....	141	<i>Mylocheilus</i> .....	219, 220
177, 2792	<i>Epinephelus</i> .....	1159	<i>Sebastes</i> .....	1821
175	<i>Pimelodus</i> .....	140	<i>Sebastodes</i> .....	1821
174, 2790	<i>Serranus</i> .....	1159	<i>Cavalla</i> .....	875
195	<i>Silurus</i> .....	138	<i>cavalla</i> , <i>Cybius</i> .....	876
176	<i>cauda bifurca</i> , <i>Chetodon</i> .....	1562	• <i>Scomberomorus</i> .....	875
176	<i>Gobius longissima acuminata</i> .....	2230	<i>Cavally</i> .....	920
178	<i>Percu nigra</i> .....	1303	<i>cavernosus</i> , <i>Bathygadus</i> .....	2581
194	<i>Turdus convexa</i> .....	1145	<i>Hymenocephalus</i> .....	2580
181	<i>caudacuta</i> , <i>Motella</i> .....	2560	<i>cavifrons</i> , <i>Ambloplites rupestris</i> ..	990
181	<i>Rhinonemus</i> .....	2560	<i>Diagramma</i> .....	1343
187	<i>candafurcatus</i> , <i>Amiurus</i> .....	135	<i>Hemitripterus</i> .....	2023
206, 218, 322	<i>Pimelodus</i> .....	135	<i>Icelinus</i> .....	1892
177	<i>caudalis</i> , <i>Halichares</i> .....	1600	<i>Tarandichthys</i> .....	1891
171	<i>Iridio</i> .....	1599	<i>Caxis</i> .....	1259
171	<i>Jullis</i> .....	1599	<i>caxis</i> , <i>Lutjanus</i> .....	1260
181	<i>Platyglossus</i> .....	1599, 1600	<i>Mesoprion</i> .....	1260
181	<i>Pomacentrus</i> .....	1556	<i>Sparus</i> .....	1259
178, 2793	<i>caudanotatus</i> , <i>Mesoprion</i> .....	1262	<i>cayanus</i> , <i>Pristigaster</i> .....	438
193	<i>caudata</i> , <i>Lamna</i> .....	37	<i>cayennense</i> , <i>Siphostoma</i> .....	772
179	<i>caudatus</i> , <i>Lepidopus</i> .....	887, 2844	<i>cayennensis</i> , <i>Citharichthys</i> .....	2686
181	<i>Triobichrus</i> .....	887	<i>Corythoichthys</i> .....	2938
171	<i>caudicinctus</i> , <i>Prilonotus (Anchiso-</i>	1742	<i>Lutjanus</i> .....	1404
175	<i>mus)</i> .....	1742	<i>Otolithus</i> .....	1404, 1411
179	<i>Tetrodon</i> .....	1742	<i>Syngnathus</i> .....	773
175, 2791	<i>caudlenla</i> , <i>Conger</i> .....	355	<i>Trachinotus</i> .....	945
177, 2792	<i>Leptocephalus</i> .....	355	<i>Cayennia</i> .....	2265
181	<i>caudilimbatus</i> , <i>Conger</i> .....	355	<i>gulchenoti</i> .....	2265
2792	<i>Echelms</i> .....	355	<i>cayennsis</i> , <i>Vomer</i> .....	934
179	<i>Leptocephalus</i> .....	355	<i>cayorum</i> , <i>Corythoichthys</i> .....	2838
177	<i>caudimacla</i> , <i>Diplodus</i> .....	1363	<i>Ogilbia</i> .....	2503
170	<i>Hæmulon</i> .....	1290, 1302, 1309	<i>cayuga</i> , <i>Eucalia inconstans</i> .....	744
192	<i>Sargus</i> .....	1363	<i>Notropis</i> .....	260
2793	<i>caudispinosum</i> , <i>Macrostoma</i> .....	556, 2820	<i>utrocaudalis</i> .....	260
	<i>caudispinosus</i> , <i>Notoscoelus</i> .....	556	<i>Cazon de Playa</i> .....	36



	Page.		Page.
Cebedichthyina .....	2349	Centronotus gunelliformis .....	2421
Cebedichthys .....	2426, 2869	gunnellus .....	2419
cristagalli .....	2427	islandicus .....	2438
violaceus .....	2427	letus .....	2420
Céfaio .....	811	nebulosus .....	2414
celatus, Batrachus .....	2316	(O p i s t h o c e n t r u s )	
Cenisophius .....	243	quinquemaclatus .....	2430
Centaurus .....	1753	pictus .....	2416
boops .....	1755	roscoi .....	2420
Centrarchidae .....	984	spinosus .....	948
Centrarchinae .....	985	taczanowskii .....	2416
Centrarchus .....	988	Centrophorus caeleolepis .....	55
aeneus .....	900	centropleura, Cottus semiscabra ...	1945
fasciatus .....	1012	Centropomidae .....	1116
hexacanthus .....	987	Centropomus .....	1117
interruptus .....	991	affinis .....	1124, 2853
macropterus .....	988	alburnus .....	1475
maculosus .....	991	albus .....	1135
pentacanthus .....	990	appendiculatus .....	1119
pomotis .....	989	armatus .....	1123
tetracanthus .....	1540	auratus .....	1107
viridis .....	992	brevis .....	1125
centrarchus, Cichlasoma .....	1526	cuvieri .....	1121
Heros .....	1526	ensiferus .....	1125, 2853
Centridermichthys analis .....	2013	grandoculatus .....	1120
asper .....	1944	luteus .....	1024
bicornis .....	1913	medius .....	1120
globiceps .....	2017	mexicanus .....	1121
gulosus .....	1945	nigrescens .....	1119
maculosus .....	2014	parallelus .....	1122
uncinatus .....	1906	pectinatus .....	1122
centrina, Callorhynchus .....	95	pedimacula .....	1119
Centriscus .....	759	robalito .....	1123
scolopax .....	759	rubens .....	1107
Centroblennius .....	2435	scaber .....	1124
nubilus .....	2438	undecimalis .....	1118
centrognathus, Zanclus .....	1098	undecimradiatus .....	1119
Centrolabrus .....	1575	unionensis .....	1122
exoletus .....	1576	viridis .....	1118
Centrolophidae .....	962	Centropriestes .....	1198
Centrolophinae .....	962	annularis .....	1214
Centrolophus .....	962	atrorius .....	1200
liparis .....	963	aurorubens .....	1278
morio .....	963	ayresi .....	1205
niger .....	963	bivittatus .....	1205
pompius .....	963	brailliensis .....	1221
Centronotus .....	900, 2414	dispilurus .....	1220
alectrolophus .....	2422	fascicularis .....	1208
apus .....	2430	fuscus .....	1211
argenteus .....	899	luciopercannus .....	1216
conductor .....	900	macrophthalmus .....	1281
cristagalli .....	2423	m. tropoma .....	1206
dolichogaster .....	2417	merus .....	1162
dybowski .....	2431	nigricans .....	1209
fasciatus .....	2418	oculatus .....	1283
gardenii .....	948	ocyurus .....	1200

Page.		Page.		Page.
.....	2421	1201	cephalus, Semotilus .....	222
.....	2419	1212	Cepolophs .....	2477
.....	2438	1214	Cepphus .....	2540
.....	2420	1213	cerapalus, Opeanus .....	2316
.....	2414	1205	cerasinus, Gobiesox .....	2336
.....	2430	1208	Hypsilepis cornutus .....	283
.....	2416	1199	Notropis .....	283
.....	2420	1199	Ceratacanthus .....	1717, 1718, 2860
.....	948	1219	aurantiacus .....	1718
.....	2416	1215	punctatus .....	2860
.....	55	1202	schœpfi .....	2860
.....	1945	1202	scriptus .....	2860
.....	1116	1682	Ceratias .....	2729
.....	1117	56	carunculatus .....	2732
.....	1124, 2853	56	conesii .....	2732
.....	1475	54	holbholi .....	2729
.....	1135	55	uraoscopus .....	2730
.....	110	83	Ceratichthys .....	252, 314
.....	1123	83	amblops .....	321
.....	1107	83	biguttatus .....	323
.....	1125	83	cataractæ .....	306
.....	1121	381	cumingii .....	318
.....	1121	2012	cyclotis .....	323
.....	1125, 2853	416	dissimilis .....	319
.....	1120	416	gelidus .....	317
.....	1024	416	hyalinus .....	321
.....	1120	416	hypsinosus .....	320
.....	1121	1238	labrosus .....	319
.....	1119	2867	leptocephalus .....	323
.....	1122	2182	lucens .....	321
.....	1122	2183, 2867	micropogon .....	323
.....	1119	2183	monacus .....	318
.....	1123	89, 2750	nubilus .....	312
.....	1107	119	physignathus .....	326
.....	1124	2210	plumbens .....	324
.....	1118	2008	prothemius .....	324
.....	1119	1143	rubrifrons .....	320
.....	1122	92	sallæi .....	212
.....	1118	93	aquamilentus .....	323
.....	1198	93	sterletus .....	316
.....	1214	93	stigmaticus .....	323
.....	1200	92	symmetricus .....	246
.....	1278	93	ventricosus .....	309
.....	1205	92	vigilax .....	253
.....	1205	23, 25	zanemus .....	319
.....	1221	228, 1753	Ceratiidæ .....	2727
.....	1220	1754	Ceratiinæ .....	2728
.....	1208	1756	Ceratius shufeldti .....	2731
.....	1211	1756	Ceratobatis .....	2756
.....	1216	1754	robertsii .....	2756
.....	1281	1754	Ceratocottus .....	1939
.....	1206	1756	diceraus .....	1940, 1941
.....	1162	25	lucasi .....	1940
.....	1200	2332	Ceratoptera .....	92
.....	1283	811	vampyrus .....	93
.....	1200	2141	Ceratoscopelus .....	557
.....	1200	1201	cephalus, Semotilus .....	222
.....	2419	1212	Cepolophs .....	2477
.....	2438	1214	Cepphus .....	2540
.....	2420	1213	cerapalus, Opeanus .....	2316
.....	2414	1205	cerasinus, Gobiesox .....	2336
.....	2430	1208	Hypsilepis cornutus .....	283
.....	2416	1199	Notropis .....	283
.....	2420	1199	Ceratacanthus .....	1717, 1718, 2860
.....	948	1219	aurantiacus .....	1718
.....	2416	1215	punctatus .....	2860
.....	55	1202	schœpfi .....	2860
.....	1945	1202	scriptus .....	2860
.....	1116	1682	Ceratias .....	2729
.....	1117	56	carunculatus .....	2732
.....	1124, 2853	56	conesii .....	2732
.....	1475	54	holbholi .....	2729
.....	1135	55	uraoscopus .....	2730
.....	110	83	Ceratichthys .....	252, 314
.....	1123	83	amblops .....	321
.....	1107	83	biguttatus .....	323
.....	1125	83	cataractæ .....	306
.....	1121	381	cumingii .....	318
.....	1121	2012	cyclotis .....	323
.....	1125, 2853	416	dissimilis .....	319
.....	1120	416	gelidus .....	317
.....	1024	416	hyalinus .....	321
.....	1120	416	hypsinosus .....	320
.....	1121	1238	labrosus .....	319
.....	1119	2867	leptocephalus .....	323
.....	1122	2182	lucens .....	321
.....	1122	2183, 2867	micropogon .....	323
.....	1119	2183	monacus .....	318
.....	1123	89, 2750	nubilus .....	312
.....	1107	119	physignathus .....	326
.....	1124	2210	plumbens .....	324
.....	1118	2008	prothemius .....	324
.....	1119	1143	rubrifrons .....	320
.....	1122	92	sallæi .....	212
.....	1118	93	aquamilentus .....	323
.....	1198	93	sterletus .....	316
.....	1214	93	stigmaticus .....	323
.....	1200	92	symmetricus .....	246
.....	1278	93	ventricosus .....	309
.....	1205	92	vigilax .....	253
.....	1205	23, 25	zanemus .....	319
.....	1221	228, 1753	Ceratiidæ .....	2727
.....	1220	1754	Ceratiinæ .....	2728
.....	1208	1756	Ceratius shufeldti .....	2731
.....	1211	1756	Ceratobatis .....	2756
.....	1216	1754	robertsii .....	2756
.....	1281	1754	Ceratocottus .....	1939
.....	1206	1756	diceraus .....	1940, 1941
.....	1162	25	lucasi .....	1940
.....	1200	2332	Ceratoptera .....	92
.....	1283	811	vampyrus .....	93
.....	1200	2141	Ceratoscopelus .....	557

	Page.		Page.
<i>Ceratoscopelus madeirensis</i> .....	557	<i>Cetomimus storeri</i> .....	550
<i>cercostrigma</i> , <i>Cyprineilla</i> .....	275	<i>Cetorhinidae</i> .....	50
<i>Minnilius</i> .....	275	<i>Cetorhinus</i> .....	51
<i>Notropis</i> .....	274	<i>maximus</i> .....	51
<i>Cerdale</i> .....	2448	<i>shavianus</i> .....	51
<i>ionthas</i> .....	2449	<i>ceuthœcum</i> , <i>Gobisoma</i> .....	2261
<i>cerdale</i> , <i>Carcharhinus</i> .....	2746, 2747	<i>ceuthœcus</i> , <i>Barbulifer</i> .....	2260
<i>Scytalina</i> .....	2454	<i>Chænichthyidae</i> .....	2293
<i>Cerdalidae</i> .....	2448	<i>Chænobryttus</i> .....	991
<i>Cerna</i> .....	1148	<i>antistius</i> .....	992
<i>acutirostris</i> .....	1181	<i>gulosus</i> .....	992
<i>gigas</i> .....	1154	<i>Chænomugil</i> .....	816
<i>macrogenis</i> .....	1181	<i>proboacidens</i> .....	816
<i>nebulosa</i> .....	1181	<i>Chænopsinae</i> .....	2347
<i>sicana</i> .....	1162	<i>Chænopsis</i> .....	2403
<i>Cernier</i> .....	1139	<i>ocellatus</i> .....	2403
<i>cernium</i> , <i>Polyprion</i> .....	1139	<i>Chærojulis</i> .....	1587
<i>Cero</i> .....	875	<i>arangoi</i> .....	1597
<i>cervinum</i> , <i>Lepophidium</i> .....	2484, 2485	<i>bivittatus</i> .....	1597
<i>Moxostoma</i> .....	197	<i>cinctus</i> .....	1593
<i>cervinus</i> , <i>Ptychostomus</i> .....	197	<i>crotophus</i> .....	1598
<i>Teretulus</i> .....	197	<i>cyanosigma</i> .....	1591
<i>cervus</i> , <i>Synauecia</i> .....	1941	<i>grandisquamis</i> .....	1597
<i>Cestracion</i> .....	43	<i>humeralis</i> .....	1597
<i>francisci</i> .....	21	<i>internasalis</i> .....	1594
<i>pantherinus</i> .....	21	<i>maculiplana</i> .....	1595
<i>quoyi</i> .....	21	<i>radiatus</i> .....	1591
<i>Cestraciont Sharks</i> .....	19	<i>ruptus</i> .....	1583
<i>Cestreus acoupa</i> .....	1404	<i>Chætodipterus</i> .....	1667
<i>albus</i> .....	1411	<i>faber</i> .....	1668
<i>carolinensis</i> .....	1409	<i>zonatus</i> .....	1668
<i>lefarchus</i> .....	1415	<i>Chætodon</i> .....	1672, 1673, 1677, 1679, 2859
<i>microlepidotus</i> .....	1415	<i>alepidotus</i> .....	966
<i>nebulosus</i> .....	1409	<i>arcuatus</i> .....	1680
<i>nobilis</i> .....	1413	<i>atæniatus</i> .....	1676
<i>nothus</i> .....	1407	<i>aureus</i> .....	1680
<i>obliquatus</i> .....	1405	<i>ays</i> .....	1675
<i>othonopterus</i> .....	1405	<i>bimaculatus</i> .....	1674
<i>parvipinnis</i> .....	1410	<i>bricei</i> .....	1678
<i>phoxocephalus</i> .....	1414	<i>canescens</i> .....	1688
<i>regalis</i> .....	1407	<i>capistratus</i> .....	1677
<i>thalassinus</i> .....	1408	<i>cauda bifurca</i> .....	1562
<i>reticulatus</i> .....	1409	<i>chirurgus</i> .....	1692
<i>squamipinnis</i> .....	1404	<i>ciliaris</i> .....	1685
<i>stolzmanni</i> .....	1412	<i>cornutus</i> .....	1688
<i>xanthinum</i> .....	1411	<i>conaga</i> .....	1691
<i>Cestrorhinus</i> .....	43	<i>cyprinaceus</i> .....	1388
<i>cetaceus</i> , <i>Squaleus</i> .....	51	<i>faber</i> .....	1668
<i>Cetengranulis</i> .....	450	<i>glaucus</i> .....	941
<i>brevis</i> .....	450	<i>gracilis</i> .....	1675
<i>edentulus</i> .....	450	<i>humeralis</i> .....	1674
<i>engymen</i> .....	2815	<i>lanceolatus</i> .....	1490
<i>mysticetus</i> .....	450	<i>littoricola</i> .....	1680
<i>Cetomimidae</i> .....	549	<i>lutescens</i> .....	1680
<i>Cetomimus</i> .....	549	<i>macrolepidotus</i> .....	1677
<i>gillii</i> .....	549	<i>maculocinctus</i> .....	1874

Page.		Page.		Page.
550	<i>Chaetodon marginatus</i>	1562	<i>chalybeius, Hyphalonedrus</i>	542
50	<i>mauriti</i>	1562	<i>chamaeleonticeps, Lopholatilus</i>	2278
51	<i>nigrirostris</i>	1673, 1674	<i>chamberlaini, Notropis</i>	2800
51	<i>ocellatus</i>	1674	<i>Chani</i>	414
51	<i>oviformis</i>	1668	<i>Chanida</i>	414
2291	<i>parrae</i>	1685	<i>Channel Bass</i>	1453
2260	<i>paru</i>	1681	<i>Cat</i>	134
2293	<i>plumieri</i>	1668	<i>Cat of the Potomac</i>	138
991	<i>rhomboides</i>	942, 2848	<i>Cats</i>	133
992	<i>sargoides</i>	1562	<i>Channomuraena</i>	404
992	<i>saxatilis</i>	1562	<i>cubensis</i>	404
816	<i>sedentarius</i>	1675	<i>vittata</i>	404
816	<i>squamulosus</i>	1685	<i>Chanos</i>	414
2347	<i>striatus</i>	1677	<i>arabicus</i>	415
2403	<i>tricolor</i>	1684	<i>chanos</i>	414, 2807
2403	<i>tristegus</i>	1691	<i>chloropterus</i>	415
1587	<i>unicolor</i>	1676	<i>cyprinella</i>	415
1597	<i>zebra</i>	1691	<i>indicus</i>	415
1597	<i>chaetodon, Apomotis</i>	995	<i>mento</i>	415
1593	<i>Cottus</i>	2316	<i>nuchalis</i>	415
1598	<i>Mesogonistius</i>	995	<i>orientalis</i>	415
1591	<i>Pomotis</i>	995	<i>salmonens</i>	415
1597	<i>Chaetodontidae</i>	1669, 1670	<i>chanos, Chanos</i>	414
1597	<i>Chaetodontinae</i>	2860	<i>Mugil</i>	415
1594	<i>Chaetodontops</i>	1672, 1673	<i>chantenay, Raja</i>	71
1595	<i>Chaetostomus</i>	160	<i>chaperi, Acanthoclinus</i>	2375
1591	<i>aspidolepis</i>	159	<i>Paraclinus</i>	2374
1593	<i>fischeri</i>	160	<i>Chapin</i>	1722, 1723
1667	<i>gaucharote</i>	150	<i>Chapinus</i>	1721, 1722, 1723
1668	<i>chagresensis, Chaicinopsis</i>	337	<i>Chappaul</i>	224
1668	<i>chagresi, Ancistrus</i>	160	<i>Characinidae</i>	331
1677, 1670, 2859	<i>Pimelodella</i>	154	<i>Characininae</i>	332
966	<i>Pimelodus</i>	154	<i>Characius</i>	331
1680	<i>ohalceum, Pristopoma</i>	1338	<i>Characodon</i>	667
1676	<i>chalcens, Orthopristis</i>	1337	<i>bilineatus</i>	668, 2831
1680	<i>Chalcinopsis</i>	337	<i>eiseni</i>	2831
1675	<i>chagresensis</i>	337	<i>ferrugineus</i>	669
1674	<i>dentex</i>	337	<i>furcoidens</i>	669
1678	<i>striatulus</i>	337	<i>garmani</i>	2832
1688	<i>chalcogramma, Theragra</i>	2535	<i>lateralis</i>	668, 2831
1677	<i>chalcogrammus, Gadus</i>	2536	<i>luitoldii</i>	2832
1562	<i>Pollachius</i>	2536, 2537	<i>variatus</i>	669, 2831
1692	<i>chalinus, Epinephelus</i>	1181	<i>Charr, European</i>	508
1685	<i>Chalinura</i>	2576	<i>Greenland</i>	508, 510
1688	<i>filifera</i>	2577	<i>Long-finned</i>	509
1691	<i>serrula</i>	2576	<i>Oregon</i>	507
1388	<i>simula</i>	2578	<i>Charrs</i>	508
1668	<i>Chalinurus</i>	2576	<i>charybdis, Lepomis</i>	992
941	<i>Chalisona</i>	1699	<i>Chasmistes</i>	182
1675	<i>velata</i>	1703	<i>brevirostris</i>	183, 199
1674	<i>challengeri, Macdonaldia</i>	617	<i>copel</i>	2795
1490	<i>Notacanthus</i>	618	<i>cujus</i>	183, 2794
1680	<i>chalybeus, Hybopsis</i>	288	<i>fecundus</i>	2794
1680	<i>Minnis</i>	288	<i>liorus</i>	183
1677	<i>Notropis</i>	288	<i>luxatus</i>	183, 2794
1674	<i>chalybeius, Chlorophthalmus</i>	542	<i>stomias</i>	2794

	Page.		Page.
Chasmodes .....	2301	cherna, Anthias .....	1157
boschianus .....	2394	Cherna Criolla .....	1157
bosquianus .....	2394	de Vivero .....	1160
jenkinsi .....	2391, 2392	Cherno de lo Alto .....	1151
novemlineatus .....	2393	cheateri, Phycis .....	2556
quadrifasciatus .....	2392	Urophycis .....	2556
sahurrae .....	2392	Chevalier, Ombre .....	508
Chatoeuss .....	415	chevola, Gallichthys .....	932
cepedianus .....	416	Chi .....	209
ellipticus .....	416	Chiasmodon .....	2291
eumorphus .....	433	niger .....	2291, 2292
mexicanus .....	416	Chiasmodontidae .....	2291
petenensis .....	417	Chiasmodus .....	2291
signifer .....	433	Chicarro .....	911
Chauffe-soleil .....	1548	chickasavensis, Luxilus .....	275
Chauffe-soleils .....	1545	Chicolar .....	879
Chauliodontidae .....	578	Chigh .....	209
Chauliodontinae .....	578	chinahua, Notropis .....	265
Chauliodus .....	584	Chilara .....	2488
macouni .....	585	taylori .....	2489
richardsoni .....	587	chilensis, Brama .....	960
schneideri .....	585	Caranx .....	927
setinotus .....	585	Exocoetus .....	730
sloanei .....	585	Pelamys .....	873
Chaunax .....	2726	Sarda .....	872
fimbriatus .....	2726	chiliticus, Hybopsis .....	287
nuttingii .....	2726, 2727	Nctropis .....	287
pictus .....	2726	Chilodipterus .....	1112
Cheilichthys .....	1729; 1730; 1734	chiloensis, Agonopsis .....	2069
asellus .....	1740	Agonus .....	2069
psittacus .....	1740	Aspidorporus .....	2069
pachygaster .....	1738	Chilomycterus .....	1747, 1748, 1750
Cheilodipteridae .....	1105	antennatus .....	1750
Cheilodipterinae .....	1105	antillarum .....	1749
Cheilodipterus .....	946, 1112	atinga .....	1750
acoupa .....	1404	californiensis .....	1751
affinis .....	1113	fuliginosus .....	1749
chrysepterus .....	1324	geometricus .....	1749
heptacanthus .....	947	puncticulatus .....	1750
Cheilonemus .....	220	reticulatus .....	1751
pulchellus .....	222	spinosus .....	1749
Cheilotrema .....	1455, 1456	Chilorhinus .....	372
Cheiragonus .....	2038	suensonii .....	372
gradiens .....	2041	Chimara .....	94
Chelidonichthys .....	2175, 2867	abbreviata .....	95
pictipinnis .....	2175, 2867	affinis .....	95
Chelmo pelta .....	1671	argentea .....	95
Chelmon aculeatus .....	1671	borealis .....	95
chemnitzii, Notacanthus .....	614	cristata .....	95
cheuwei, Cottogaster .....	2851	mediterranea .....	95
Cheonda .....	228, 230, 236	monstrosa .....	94
caeruleus .....	232	plumbea .....	95
cooperi .....	236	Chimæras .....	93
modesta .....	234	Chimæridæ .....	93
Cherna .....	1157	Chimærinae .....	94
americana .....	1180	Chimæroides .....	93

Page.		Page.		Page.
1157	Chimæroids	93	Chirus pictus	1873
1157	Chino, Escolar	1114, 1284	trigammus	1872
1160	Mojarra	1377	chirus, Xiphister	2424
1151	Chinook Salmon	479	Xiphistes	2424
2556	chiosictetus, Entomacrodus	2398	Chisel-mouths	207, 208
2556	Salarias	2398	chisoensis, Pœcilia	693
508	chiquita, Aboma	2241	Chltonotus	1889
932	Gobius	2241	Chitonotus megacephalus	1891
209	Chirivita	1679	pugetensis	1890, 1891
2291	Chirivitas	1679	chittendeni, Cyclosetta	2676
2291, 2292	Chiro	410	Chivo	800
2291	Chirocentrodon	435	chlamydes, Bollmannia	2238
2291	tænolatus	435	Chlamydoselachidæ	16
911	Chirolophinae	2347	Chlamydoselachus	16
275	Chirolophis polyactocephalus	2409	angineus	16
879	Chirolophis japonicus	2409	chleverages, Gymnothorax	399
209	Chironectes	2717	Lycodontia	398
265	arcticus	2717	Sidera	399
2488	lævigatus	2717	Chlopsis	364
2489	mentzelii	2724	equatorialis	364
960	multicellatus	2725	chlora, Cliola	263
927	pictus	2717	Chlorichthys	1605; 2859
730	principis	2719	bifasciatus	1609, 1610
873	scaber	2723	brasiliensis	1591
872	sonntagii	2717	grammaticus	1610
287	tenebrosus	2719	lucasanus	1607
287	tigris	2723	nitidissimus	1608
1112	tumidus	2717	nitidus	1608
2069	Chiropsis	1866	socorroensis	1607
2069	constellatus	1868	steindachneri	1609
2069	guttatus	1869	virens	1610
747, 1748, 1750	nebulosus	1872	chlora, Pseudoscaris	1648
1750	Chirostoma	792, 2839, 2840	Pseudoscarus	1654
1749	bartoni	793	Scarus	1637, 1640
1750	brasiliensis	794	Scomber	938
1751	beryllinum	798	chloristia, Cliola	278
1749	estor	792, 2839	Codoma	278
1749	grandocule	2839	chloristius, Notropis niveus	278
1750	humboldtianum	793, 2839	chlorocephalus, Hybopsis	286
1751	jordani	793	Minnilus	286
1749	peninsulæ	797	Notropis	286
372	sicculum	806	chloropteron, Catostomus	179
372	vagrans	795	chloropteron, Plectropoma	1165
94	chirurgus, Acanthurus	1692	chloropterus, Chanos	415
95	Chaetodon	1692	Prospinus	1165
95	Chirus	1866	Chlorophthalmus	541
95	balius	1873	agassizii	541
95	constellatus	1869	chalybeius	542
95	decagrammus	1869	Chloroscombrinae	897
95	guttatus	1868	Chloroscombrus	937
94	hexagrammus	1872	caribbeus	938
95	maculoseriatus	1868	chlora	2847
93	monopterygius	1866	chrysurus	938, 2847
93	nebulosus	1872	ectenurus	2847
94	octogrammus	1870	orqueta	937
93	ordinatus	1870	stirurus	938

	Page.		Page.
chlorosoma, <i>Boleosoma</i> .....	1060	<i>Chornia</i> Ryba .....	621
<i>Vaillantia</i> .....	1060	<i>Chorophthalmus</i> <i>agassizii</i> .....	542
<i>chlorostictus</i> , <i>Sebastichthys</i> .....	1812	<i>truculentus</i> .....	542
<i>Sebastodes</i> .....	1811	<i>chouicha</i> , <i>Oncorhynchus</i> .....	480
<i>chlorostomus</i> , <i>Triostropis</i> .....	1170	<i>Chriodorus</i> .....	719
<i>chlorurum</i> , <i>Plectropoma</i> .....	1193	<i>atherinoideus</i> .....	719
<i>Chlorurus</i> .....	1642	<i>Chriolax</i> .....	2148
<i>chlorurus</i> , <i>Hypoplectrus</i> .....	1193	<i>Chriolepis</i> .....	2205
<i>unicolor</i> .....	1198	<i>minutillus</i> .....	2205
<i>Serranus</i> .....	1193	<i>Chriomitra</i> .....	873
<i>Chenopsetta</i> .....	2624	<i>concolor</i> .....	874
<i>dentata</i> .....	2630	<i>Chriope</i> .....	254, 255, 258
<i>oblonga</i> .....	2633	<i>Chromides</i> .....	781, 1511
<i>ocellaris</i> .....	2630	<i>Chromis</i> .....	946, 1545
<i>Cheroichthys</i> .....	773	<i>atrilobatus</i> .....	1546
<i>cherostomus</i> , <i>Engraulis</i> .....	444	<i>cyaneus</i> .....	1547
<i>Stolporus</i> .....	444	<i>enchrysurus</i> .....	1548
<i>Chetopterus</i> .....	1279	<i>epicurorum</i> .....	947
<i>Chogset</i> .....	1577	<i>fenestrata</i> .....	1518
<i>chogset</i> , <i>Ctenolabrus</i> .....	1577	<i>fusco-maculatus</i> .....	1540
<i>Labrus</i> .....	1577	<i>insolatus</i> .....	1548
<i>fulvus</i> .....	1577	<i>marginatus</i> .....	1546
<i>Choice</i> , Sailor's .....	1297, 1338	<i>multilineatus</i> .....	1547
<i>choirorhynchus</i> , <i>Dajaus</i> .....	2841	<i>nebulifer</i> .....	1524
<i>Choker</i> , Hog .....	2700	<i>punctipinnis</i> .....	1548
<i>Chologaster</i> .....	703	<i>chromis</i> , <i>Dibasis</i> .....	1299
<i>agassizii</i> .....	704	<i>Hæmulon</i> .....	1299
<i>avitus</i> .....	704	<i>Labrus</i> .....	1483
<i>cornutus</i> .....	703	<i>Chronophorus</i> <i>mexicanus</i> .....	2237
<i>papilliferus</i> .....	704	<i>Chrosomus</i> .....	209
<i>Chondroganoidea</i> .....	98, 100	<i>dakotensis</i> .....	210
<i>Chondrostei</i> .....	102	<i>eos</i> .....	210
<i>Chondrostoma</i> <i>gardoneum</i> .....	251	<i>erythrogaster</i> .....	209
<i>pullum</i> .....	206	<i>eos</i> .....	210
<i>Chondrostominae</i> .....	202	<i>oreas</i> .....	211
<i>Chonophorus</i> .....	2234	<i>pyrrhogaster</i> .....	210
<i>bucculentus</i> .....	2236	<i>chrosomus</i> , <i>Hybopsis</i> .....	288
<i>Chonophorus</i> <i>flavus</i> .....	2235	<i>Hydrophlox</i> .....	288
<i>taiasica</i> .....	2237	<i>Mianilus</i> .....	288
<i>Chopa</i> .....	1387	<i>Notropis</i> .....	288
<i>Amarilla</i> .....	1386	<i>chrysaargyreus</i> , <i>Hæmulon</i> .....	1308
<i>Blanca</i> .....	1387	<i>chrysa</i> <i>gyreus</i> , <i>Brachygenys</i> .....	1307
<i>Chopa Spina</i> .....	1357, 1358	<i>chryseus</i> , <i>Rhinoberyx</i> .....	847
<i>Chopas</i> .....	1384	<i>chrysitis</i> , <i>Dionda</i> .....	214
<i>chorlatus</i> , <i>Saccopharynx</i> .....	406	<i>chrysocephalus</i> , <i>Luxilus</i> .....	282
<i>Stylephorus</i> .....	2601	<i>chrysochloris</i> , <i>Pomolobus</i> .....	425
<i>Choregon</i> .....	517	<i>Clupea</i> .....	425
<i>Chorinemus</i> <i>altus</i> .....	899	<i>chrysoaster</i> , <i>Agosia</i> .....	313
<i>inornatus</i> .....	899	<i>chrysoleuca</i> , <i>Bairdiella</i> .....	1438
<i>occidentalis</i> .....	898	<i>Corvina</i> .....	1439
<i>palometa</i> .....	899	<i>Solana</i> .....	1439
<i>quiebra</i> .....	899	<i>chrysoleucus</i> , <i>Notemigonus</i> .....	250
<i>saliens</i> .....	899	<i>chrysomeiaunrus</i> , <i>Sparus</i> .....	1157
<i>saltans</i> .....	899	<i>chrysomelas</i> , <i>Sebastichthys</i> .....	1826
<i>Chorististlum</i> .....	1136	<i>Sebastodes</i> .....	1825, 1826
<i>rubrum</i> .....	1136	<i>Chrysophrys</i> <i>calamus</i> .....	1350

Page.		Page.		Page.
621	Chrysophrys cyanoptera	1354	Chub of Utah Lake	232
542	<i>taurina</i>	1394	Chuckle-headed Cat	134
542	Chrysops aculatus	1347	Chuss	2555
480	chrysops, Caulolatilus	2277	chuss, Phycis	2555
719	Latilus	2278	Urophycis	2555
719	Ophichthys	385	Chylomycterus schœpfi	1748
2148	Ophlurus	385	cibaria, Lampetra	13
2205	Perca	1132	cibarius, Ammocetes	13
2205	Roccus	1132	Cibi Amarillo	910
873	Sparus	1346	Mancho	919
874	Stenotomus	1346	cibi, Carangoides	920
254, 255, 258	chrysopsis, Hyodon	413	Caranx	920
781, 1511	chrysoptera, Perca	1339	oicatricosus, Baliates	1709
946, 1545	chrysopteron, Hæmulon	1309	Pleuronectes	2640
1540	chrysopteron, Hæmulon	1309	Xanthichthys	1709
1547	Sparisoma	1036, 1637	Cichla anea	990
1548	chrysopterus, Cheilodipterus	1324	<i>fasciata</i>	1012
947	Diabasis	1309	<i>floridana</i>	1012
1518	Leuciscus	221	<i>minima</i>	1012
1540	Orthopristsis	1338	<i>ohioensis</i>	1012
1548	Scarus	1637	<i>storeria</i>	987
1540	chrysos, Carangus	921	Cichlasoma	1514, 1515
1547	chrysoctenus	954	<i>anguliferum</i>	1517
1524	chrysoctus, Fundulus	655	<i>balteatum</i>	1521
1548	Haplochilus	656	<i>bartoni</i>	1515
1299	Zygocetes	656	<i>bifasciatum</i>	1521
1299	chrysaure, Bairdiella	1433	<i>centrarchus</i>	1526
1483	Schena	1434	<i>deppii</i>	1524
2237	chrysurus, Chlorocombrus	938	<i>fenestratum</i>	1518
209	Coryphæna	952	<i>goodmanni</i>	1516
210	Dipteron	1433	<i>helleri</i>	1521
209	Glyphidodon	1567	<i>intermedium</i>	1517
210	Lutjanus	1276	<i>lentiginosum</i>	1524
210	Mesoprion	1276	<i>longimanus</i>	1520
211	Micropteryx	938	<i>macracanthum</i>	1518
210	Microspathodon	1567	<i>margaritifera</i>	1519
288	Ocyurus	1275	<i>melanopogon</i>	1523
288	Scomber	938	<i>melanurum</i>	1523
288	Sparus	1276	<i>montezuma</i>	1518
1308	chrysus, Caranx	921	<i>multispinosum</i>	1525
1307	chryssya, Anguilla	348	<i>nebuliferum</i>	1524
847	Chub	1387	<i>nigrofasciatum</i>	1525
214	Bermuda	1387	<i>parva</i>	1519
282	Columbia	219	<i>rectangulare</i>	1515
425	Flat-headed	326	<i>rostratum</i>	1522
425	Great	232	<i>sieboldii</i>	1516
313	Indian	322	<i>spilurum</i>	1520
1438	Nigger	327	Cichlidae	1512
1439	River	322	Cichlids	1512
1439	Sacramento	231	Ciego, Pez	2501
1157	Silver	221, 320	Cigar-fish	907
1826	Steelbacked	205	cigonella, Belone	713
1825, 1826	Tahoe	2798	ciliaris, Alectis	931
1350	Chub Mackerel	866	<i>Angelichthys</i>	1684, 1685
	Chub of the Rio Grande	233	<i>Balistes</i>	1702
	Suckers	184	<i>Chetodon</i>	1685



	Page.		Page.
<i>ciliaris</i> , <i>Holacanthus</i> .....	1685	<i>Cirrimens</i> .....	1460
<i>Pomacanthus</i> .....	1685, 1686	<i>cirris</i> , <i>Cottus plurimis</i> .....	2066
<i>Zeus</i> .....	932	<i>Cirrostromes</i> .....	2
<i>Ciliata argenteata</i> .....	2559	<i>Cisco</i> .....	468
<i>ciliatus</i> , <i>Balistes</i> .....	1715	<i>Moon-eye</i> .....	469
<i>Blennius</i> .....	2457	<i>Cisco of Lake Michigan</i> .....	469
<i>Epinephelus</i> .....	1784	<i>Tippecanoe</i> .....	469
<i>Monacanthus</i> .....	1714, 1715	<i>Ciscoes</i> .....	467
<i>Petromyzon</i> .....	12	<i>cismontanus</i> , <i>Coregonus williamsoni</i>	463
<i>Sebastodes</i> .....	1783	<i>Citharedus</i> .....	1672
<i>cimbria</i> , <i>Motella</i> .....	2560	<i>Citharichthys</i> .....	2678, 2682
<i>cimbricus</i> , <i>Enchelyopus</i> .....	2561	<i>aethalion</i> .....	2673
<i>cimbricus</i> , <i>Enchelyopus</i> .....	2560	<i>aramaca</i> .....	2672
<i>Gadus</i> .....	2560	<i>arctifrons</i> .....	2683
<i>Onos</i> .....	2561	<i>cayennensis</i> .....	2686
<i>Rhinomenus</i> .....	2561	<i>dinoceros</i> .....	2682
<i>cincinnati</i> , <i>Acipenser</i> .....	106	<i>fragilis</i> .....	2680
<i>cinctus</i> , <i>Chærojulia</i> .....	1593	<i>gilberti</i> .....	2686
<i>Julia</i> .....	1593	<i>guatemalensis</i> .....	2686
<i>cinerea</i> , <i>Alutera</i> .....	1720	<i>latifrons</i> .....	2674
<i>Amia</i> .....	113	<i>macrops</i> .....	2684
<i>Etheostoma</i> .....	1078	<i>microstomus</i> .....	2688
<i>cinereum</i> , <i>Aulostoma</i> .....	755	<i>ocellatus</i> .....	2673
<i>Etheostoma</i> .....	1078	<i>ovalis</i> .....	2674
<i>Xystema</i> .....	1372	<i>panamensis</i> .....	2677
<i>cinereus</i> , <i>Aulostomus</i> .....	755	<i>platophrys</i> .....	2683
<i>Gerres</i> .....	1370	<i>pœtulus</i> .....	2672
<i>Marorourus</i> .....	2585	<i>sordidus</i> .....	2679
<i>Microspathodon</i> .....	1570	<i>spilopterus</i> .....	2685, 2686
<i>dorsalis</i> ..	1570	<i>stigmæus</i> .....	2681
<i>Mugil</i> .....	1373	<i>sumichrasti</i> .....	2686
<i>Nothonotus</i> .....	1078	<i>uhleri</i> .....	2681
<i>Turdus peltatus</i> .....	1773	<i>unicornis</i> .....	2683
<i>cingulatus</i> , <i>Fundulus</i> .....	656	<i>ventralis</i> .....	2670
<i>Pomacanthus</i> .....	1680	<i>xanthostigmus</i> .....	2680
<i>Zygonectes</i> .....	655, 656	<i>Citharus</i> .....	2614
<i>circumnotatus</i> , <i>Scarus</i> .....	1641	<i>platessoides</i> .....	2615
<i>cirratum</i> , <i>Ginglymostoma</i> .....	26	<i>citrinellus</i> , <i>Heros</i> .....	1534
<i>cirratus</i> , <i>Milvus</i> .....	2183	<i>Citula</i> .....	929
<i>Phycis</i> .....	2554	<i>banksi</i> .....	927
<i>Squalus</i> .....	26	<i>dorsalis</i> .....	930
<i>Urophycis</i> .....	2553	<i>clucara</i> , <i>Echelus</i> .....	356
<i>Cirrhiosomus</i> .....	1729	<i>civillis</i> , <i>Hybognathus</i> .....	215
<i>Cirrhites</i> .....	1491	<i>Clam Cracker</i> .....	83
<i>betaurus</i> .....	1492	<i>Clamagore</i> .....	1652
<i>rivulatus</i> .....	1491	<i>clara</i> , <i>Ammocrypta pellucida</i> .....	1063
<i>Cirrhitichthys</i> .....	1491	<i>Menidia</i> .....	801
<i>rivulatus</i> .....	1492	<i>clarías</i> , <i>Pimelodus</i> .....	155
<i>Cirrhitiidæ</i> .....	1490	<i>Silurus</i> .....	155
<i>Cirrhitoid Fishes</i> .....	1490	<i>Claricola</i> .....	1066, 1069, 1093
<i>Cirrhitoide</i> .....	781	<i>clarionensis</i> , <i>Holacanthus</i> .....	1683
<i>Cirrhitoidei</i> .....	1490	<i>Myripristis</i> .....	2842
<i>Cirrhitoids</i> .....	1490	<i>clarionis</i> , <i>Xesurus</i> .....	1695
<i>cirrhosum</i> , <i>Lepisma</i> .....	2362	<i>clarki</i> , <i>Bogoslouius</i> .....	2575
<i>cirrhosus</i> , <i>Blepsias</i> .....	2018	<i>Catostomus</i> .....	173
<i>Trachinus</i> .....	2019	<i>Fario</i> .....	501

Page.		Page.		Page.
1460	<i>clarki</i> , <i>Pantostens</i> .....	172	<i>Clinus macrocephalus</i> .....	2364
2006	<i>clarkii</i> , <i>Salmo</i> .....	402, 2819	<i>maculatus</i> .....	2433
2	<i>bouvieri</i> .....	2819	<i>medius</i> .....	2435
468	<i>gibbsii</i> .....	2819	<i>mohri</i> .....	2438
469	<i>henshawi</i> .....	2819	<i>nebulosus</i> .....	2438
469	<i>lewisii</i> .....	2819	<i>nigripinnis</i> .....	2370
467	<i>macdonaldi</i> .....	2819	<i>nuchipinnis</i> .....	2362
Amsoni	<i>pleniticus</i> .....	2819	<i>ocellatus</i> .....	2357
463	<i>spilurus</i> .....	2819	<i>ocellifer</i> .....	2353
1672	<i>stomias</i> .....	2819	<i>pectinifer</i> .....	2362
2678, 2692	<i>tahoensis</i> .....	2870	<i>phillipi</i> .....	2359
2073	<i>virginialis</i> .....	2819	<i>præcius</i> .....	2441
2672	<i>clarum</i> , <i>Etheostoma pellucidum</i> ..	1063	<i>punctatus</i> .....	2440
2683	<i>clathrata</i> , <i>Atraptera</i> .....	1198	<i>unimaculatus</i> .....	2441
2686	<i>clathratus</i> , <i>Labrax</i> .....	1198	<i>zonifer</i> .....	2359
2682	<i>Paralabrax</i> .....	1197, 1198	<i>Cliola</i> .....	252
2680	<i>Serranus</i> .....	1198	<i>alta</i> .....	322
2686	<i>claudalus</i> , <i>Hydon</i> .....	413	<i>analostana</i> .....	279
2686	<i>claviformis</i> , <i>Moxostoma</i> .....	186	<i>aurata</i> .....	272
2674	<i>claviger</i> , <i>Cottus</i> .....	1939	<i>billingsiana</i> .....	272
2684	<i>Enophrys</i> .....	1038	<i>bubalina</i> .....	273
2688	<i>clepydra</i> , <i>Murena</i> .....	2805	<i>caerulea</i> .....	277
2673	<i>Clepticus</i> .....	1574	<i>callisema</i> .....	273
2674	<i>Clepticus</i> .....	1586	<i>callistia</i> .....	276
2677	<i>genzara</i> .....	1587	<i>calliura</i> .....	275
2683	<i>parre</i> .....	1586	<i>camura</i> .....	280
2672	<i>clevelandii</i> , <i>Phoxinus</i> .....	237	<i>chlora</i> .....	263
2679	<i>Clevelandia</i> .....	2254	<i>chloristia</i> .....	278
2685, 2686	<i>ios</i> .....	2254	<i>coblis</i> .....	305
2681	<i>longipinnis</i> .....	2255	<i>deliciosa</i> .....	272
2686	<i>rose</i> .....	2255	<i>euryopa</i> .....	270
2684	<i>Cling-fishes</i> .....	2326, 2329	<i>eurystoma</i> .....	277
2683	<i>Clininae</i> .....	2344	<i>forbesi</i> .....	272
2670	<i>Clinocottus</i> .....	2012	<i>formosa</i> .....	271
2680	<i>analis</i> .....	2612	<i>fretensis</i> .....	281
2614	<i>Clinostomus</i> .....	228, 230, 239	<i>galactura</i> .....	279
2615	<i>affinis</i> .....	239	<i>gibbosa</i> .....	272
1534	<i>elongatus</i> .....	240	<i>gunnisoni</i> .....	273
929	<i>funduloides</i> .....	239	<i>hæmatura</i> .....	218
927	<i>hydrophlox</i> .....	238	<i>hudsonia</i> .....	269
930	<i>margarita</i> .....	241	<i>hypsoptera</i> .....	280
356	<i>montanus</i> .....	238	<i>iris</i> .....	272
215	<i>pandora</i> .....	234	<i>jugalis</i> .....	272
83	<i>phlegethontis</i> .....	243	<i>leonina</i> .....	271
1652	<i>proriger</i> .....	240	<i>lepida</i> .....	273
1063	<i>tænia</i> .....	238	<i>lineolata</i> .....	263
801	<i>Clinus aculeatus</i> .....	2433	<i>longirostris</i> .....	267
155	<i>bimaculatus</i> .....	2358	<i>ludibunda</i> .....	273
155	<i>canariensis</i> .....	2362	<i>lutrensis</i> .....	272
66, 1069, 1093	<i>capillatus</i> .....	2362	<i>microstoma</i> .....	264
1683	<i>delalandii</i> .....	2359	<i>missuriensis</i> .....	262
2842	<i>evides</i> .....	2353	<i>montiregis</i> .....	272
1695	<i>gilli</i> .....	2358	<i>nigrotæniata</i> .....	264
2575	<i>gobio</i> .....	2365	<i>nives</i> .....	278
173	<i>hermineri</i> .....	2362	<i>notata</i> .....	274
501	<i>lumpenus</i> .....	2438	<i>nubila</i> .....	215

	Page.		Page.
<i>Cliola ornata</i> .....	271	<i>Clupea macrophthalma</i> .....	430
<i>procne</i> .....	264	<i>matto vacca</i> .....	426
<i>pyrrhomelas</i> .....	281	<i>mediocri</i> .....	426
<i>rubripinna</i> .....	281	<i>megalops</i> .....	426
<i>sallei</i> .....	212	<i>membras</i> .....	421
<i>saludana</i> .....	270	<i>menhaden</i> .....	434
<i>sima</i> .....	267	<i>minima</i> .....	422
<i>siathii</i> .....	253	<i>mirabilis</i> .....	422
<i>spectruncula</i> .....	265	<i>neglecta</i> .....	434
<i>stigmatura</i> .....	275	<i>pallasii</i> .....	422
<i>storeriana</i> .....	270	<i>parvula</i> .....	426
<i>straminea</i> .....	262	<i>pseudoharengus</i> .....	426
<i>suavis</i> .....	272	<i>pseudohispanica</i> .....	424
<i>taurocephala</i> .....	253	<i>pusilla</i> .....	426
<i>trichroistia</i> .....	276	<i>sadina</i> .....	420
<i>tuditana</i> .....	253	<i>sagax</i> .....	423
<i>umbrosa</i> .....	273	<i>stolifera</i> .....	432
<i>urostigma</i> .....	275	<i>thriasa</i> .....	432
<i>velox</i> .....	253	<i>thriassina</i> .....	431
<i>venusta</i> .....	274	<i>tyrannus</i> .....	434
<i>vigilax</i> .....	253	<i>vernalis</i> .....	426
<i>vittata</i> .....	258	<i>villosa</i> .....	521
<i>vivax</i> .....	253	<i>virescens</i> .....	426
<i>whipplei</i> .....	279	<i>vittata</i> .....	421
<i>xænura</i> .....	280	<i>Clupeidae</i> .....	417
<i>Clodalus</i> .....	412	<i>clupeiformis</i> , <i>Coregonus</i> .....	465, 469
<i>clodalus</i> , <i>Hiodon</i> .....	413	<i>Salmo</i> .....	466
<i>Clupanodon</i> .....	422	<i>Clupeinae</i> .....	418
<i>aureus</i> .....	434	<i>Clupeoidea</i> .....	407
<i>cæruleus</i> .....	423	<i>clupeoides</i> , <i>Engraulis</i> .....	447
<i>pseudohispanicus</i> .....	423	<i>Stolephorus</i> .....	447
<i>Clupea</i> .....	421	<i>clupeola</i> , <i>Harengula</i> .....	429, 430
<i>æstivalis</i> .....	427	<i>Sardinella</i> .....	429
<i>anchovia</i> .....	429	<i>Clupeonia</i> .....	428
<i>apicalis</i> .....	429	<i>Clypeocottus</i> .....	1937
<i>atherinoides</i> .....	451	<i>robustus</i> .....	1938
<i>aurea</i> .....	434	<i>Coal-fish</i> .....	1862, 2534
<i>brasiliensis</i> .....	411	<i>coara</i> , <i>Guabi brasiliensibus</i> .....	1305
<i>cærulea</i> .....	421	<i>Coast Range Trout</i> .....	500
<i>carolinensis</i> .....	434	<i>Coballito del Mar</i> .....	776, 777
<i>chrysochloris</i> .....	425	<i>Cobbler</i> .....	640, 641, 931
<i>elongata</i> .....	421	<i>Cobessicentic Smelt</i> .....	524
<i>esca</i> .....	421	<i>Cobia</i> .....	948
<i>fasciata</i> .....	426	<i>Cobitis heteroclitia</i> .....	641
<i>halac</i> .....	421	<i>killifish</i> .....	641
<i>harengus</i> .....	421, 422	<i>majalis</i> .....	639
<i>heterura</i> .....	416	<i>cobitis</i> , <i>Cliola</i> .....	305
<i>hudsonia</i> .....	289	<i>Leuciscus</i> .....	305
<i>humeralis</i> .....	431	<i>Tiaroga</i> .....	305
<i>indigena</i> .....	428	<i>coccineus</i> , <i>Lycodes</i> .....	2469
<i>lamprotecula</i> .....	419	<i>Scarus</i> .....	1635
<i>latulus</i> .....	422	<i>coccogenis</i> , <i>Hypsellepis</i> .....	285
<i>leachi</i> .....	422	<i>Leuciscus</i> .....	285
<i>lbertatis</i> .....	423	<i>Minnilus</i> .....	285
<i>lineolata</i> .....	422	<i>Notropis</i> .....	284
<i>macrocephala</i> .....	411	<i>coccol</i> , <i>Rhinoscopelus</i> .....	568

Page.		Page.		Page.
430	cocci, <i>Scopelus</i> .....	560	cæruleum, <i>Etheostoma spectabile</i> ..	1089
426	<i>Stenobranchius</i> .....	569	cæruleus, <i>Astatichthys</i> .....	1089
426	cocherani, <i>Cephalus</i> .....	1756	<i>Pœcilichthys</i> .....	1089
426	Cochinito.....	1694	cognata, <i>Uranidea</i> .....	1955
421	Cochino.....	1703	cognatus, <i>Cottus</i> .....	1954
434	Cochlognathus.....	251	Coho Salmon.....	480
422	<i>biguttata</i> .....	252	colias, <i>Scomber</i> .....	866
422	<i>ornata</i> .....	252	colii, <i>Salmo</i> .....	579
434	<i>ornatus</i> .....	252	colinus, <i>Gadus</i> .....	2535
422	Cocinera.....	918	Coliscus.....	217
426	Cocinero Dorado.....	921	<i>parietalis</i> .....	217
426	Cock and hen Paddo.....	2096	collapsum, <i>Moxostoma</i> .....	190
424	Cockeyo Pilot.....	1555, 1561	collapsus, <i>Ptychostomus</i> .....	190
426	Ccuyo.....	1709	colletti, <i>Careproctus</i> .....	2131
420	Cod, Blue.....	1875	Colletia.....	567
423	Buffalo.....	1875	<i>nocturna</i> .....	567
432	Cultus.....	1875	<i>rafinesquei</i> .....	567
432	Green.....	2534	colliet, <i>Hydrolagus</i> .....	95
431	Wachna.....	2537	Colocephali.....	346, 388
434	Codfish, Alaska.....	2541	Cololabis.....	726
426	<i>Commodus</i> .....	2541	<i>brevirostris</i> .....	726
521	Greenland.....	2542	Colomesina.....	1727
426	Codfishes.....	2531, 2540	Colomesus.....	1740
421	Codling.....	2552, 2555	<i>psittacus</i> .....	1740
417	Codoma.....	254, 256, 270	colonus, <i>Serranus</i> .....	1222
465, 469	<i>cærulea</i> .....	277	Colorada, <i>Maurararia</i> .....	2754
466	<i>callisema</i> .....	273	colorada, <i>Liza</i> .....	1713
418	<i>callistia</i> .....	276	<i>Vieja</i> .....	1639
407	<i>chloristia</i> .....	278	Colorado, Bagre.....	122
447	<i>corystoma</i> .....	285	Pargo.....	1264, 1267, 1356
447	<i>ornata</i> .....	271	Perro.....	1583
429, 430	<i>pyrrhomelas</i> .....	281	Pescado.....	1453
429	<i>stigmatura</i> .....	275	colorado, <i>Lutjanus</i> .....	1268
428	<i>trichroistia</i> .....	276	<i>Lutjanus</i> .....	1268
1937	<i>vittata</i> .....	258	<i>Neomenis</i> .....	1267
1938	<i>xanura</i> .....	280	Colorado River Trout.....	496
1862, 2534	Codorniz.....	1467	coloratum, <i>Aulostoma</i> .....	755
1305	Cods, Cultus.....	1875	coluber, <i>Gempylus</i> .....	884
500	Cœclophis.....	381	Columbia.....	784
776, 777	Cœcula <i>bascanium</i> .....	380	Chub.....	219
0, 641, 631	<i>scuticaris</i> .....	379	River Sucker.....	178
524	<i>teres</i> .....	370	Trout.....	492
948	cœcus, <i>Gastrobranchus</i> .....	8	Salmon.....	479
641	cælestinus, <i>Bagrus</i> .....	125	Columbia <i>transmontana</i> .....	784
641	<i>Pseudoscarus</i> .....	1655, 1656	columbiana, <i>Pantosteus</i> .....	172
639	<i>Scarus</i> .....	1656	<i>Vomer</i> .....	934
305	Coelho.....	882	comatus, <i>Cypselurus</i> .....	736
305	cœlolepis, <i>Centrophorus</i> .....	55	<i>Exocoetus</i> .....	736
305	Cœlorhynchus.....	2587	comes, <i>Roccus</i> .....	1407
2469	<i>carrinatus</i> .....	2588, 2589	commersoni, <i>Carcharhinus</i> .....	38
1635	<i>carribæus</i> .....	2589	commersonii, <i>Catostomus</i> .....	178
285	<i>occa</i> .....	2587	<i>Cyprinus</i> .....	179
285	<i>scaphopsis</i> .....	2590, 2591	<i>Fistularia</i> .....	758
285	cœnosa, <i>Parophrys</i> .....	2639	Common Alligator Fish.....	2061
284	cœnosus, <i>Pleuronichthys</i> .....	2638, 2639	American Sea-Horse.....	777
568	cæruleum, <i>Etheostoma</i> .....	1088	Angler.....	2713

	Page.		Page.
Common Atlantic Salmon .....	486	concatenatus, Ostracion .....	1723
Buffalo Fish .....	163	Conchognathus .....	349
Bullhead .....	140	grimaldii .....	349
Burr-fish .....	1748	concinus, Amblodon .....	1484
Cobbler .....	641	Gasterosteus .....	745
Codfish .....	2541	concolor, Ammocetes .....	11
Dolphin .....	952	Chromitra .....	874
Eastern Pickerel .....	627	Euschistodus .....	1559
Stickleback .....	748	Ichthyomyzon .....	11
Flatfish .....	2647	Lycodes .....	273
Gar Pike .....	109	Nexilarius .....	1559
Grunt .....	1304	Petromyzon .....	11
Gurnard .....	2156	Scomberomorus .....	873
Half-beak .....	721	Thyrsoidea .....	396
Herring .....	421	Condorado, Ronco .....	1306
Killifish .....	640	conductor, Centronotus .....	900
Mackerel .....	865	Conejo .....	596, 882
Mullet .....	811	Coney .....	1141
Pámpano .....	944	confertus, Hyborhynchus .....	217, 218
Pike .....	628	Pimephales promelas .....	217
Pipefish .....	770	confinia, Pimelodus .....	141
Rat-tail .....	2583	Salmo .....	505
Red Horse .....	192	confuentus, Fuululus .....	650
Rock Bass .....	990	Salmo .....	480
Sawfish .....	60	conformis, Lavinia .....	231
Scup .....	1346	Lenciscus .....	231
Shad .....	427	Squalius .....	231
Skate .....	68	Tigoma .....	231
of California .....	73	congener, Parn brasiliense .....	966
Spotted Moray .....	395	Conger .....	353
Sting Ray .....	83	analis .....	356
Sturgeon .....	105	brasiliensis .....	360
Sucker .....	178	caudicaula .....	355
Sunfish .....	1009	caudilimbatus .....	355
Surf-fish .....	1504	esculentus .....	355
Surgeon .....	1691	impressus .....	356
Swordfish .....	894	limbatus .....	360
Trunk-fish .....	1723	macrops .....	355
Weakfish .....	1407	microstomus .....	356
Whitefish .....	465	mordax .....	387
communis, Catostomus .....	179	niger .....	355
Leucosomus .....	326	occidentalis .....	355
Liparis .....	2118	opisthophthalmus .....	356
Platygobio .....	326	orbignyianus .....	355
Pogonichthys .....	326	rubescens .....	355
complanata, Cyprinella .....	272	verreauxi .....	355
Montana .....	272	verus .....	355
compressa, Lota .....	2551	vulgaris .....	355
compressus, Bassozetus .....	2508	Conger Eel of California .....	395
Bathynoctes .....	2500	Eels .....	352, 354
Bathyonus .....	2509	conger, Leptocephalus .....	354
Engraulis .....	447	Muræna .....	354
Gadus .....	2551	Conger muræna .....	355
Nanclerus .....	900	balearica .....	356
Rutilus .....	282, 290	flava .....	357, 2501
Stolephorus .....	447	macrura .....	359

Page.		Page.		Page.
1723	Congermuræna mellissii	356	copei, Aleposomus	459
349	nitens	357	Chasmistes	2795
349	prorigera	357	Cottus	1968
1484	congestum, Moxostoma	192	Paraliparis	2143
745	congestus, Catostomus	192	Squalius	236
11	Congresox	359	Copelandellus	1100
874	Congro Barbosa	155	qulesceus	1100
1559	Congros Barbosus	154	copelandi, Beleosoma	1046
11	Congrus	353, 381	Cottogaster	1045
2173	curvidens	360	Rheocrypta	1046
1559	leucophæus	355	Copelandia	992
11	coniceps, Murænesox	359	eriarcha	994
873	conico, Ostracion oblongus	1745	copii, Leuciscus	293
396	coniferum, Oreosoma	1663	Copper-nosed Bream	1005
1306	Conocara	456	Corallicola	2360
900	macdonaldi	457	corallina, Narcine brasiliensis	78
598, 882	macroptera	457, 458	corallinum, Cryptotrema	2366
1141	conocephala, Gila	219	corallinus, Antennarius	2725
217, 218	conocephalus, Mylopharodon	210	Corbineta	1435
217	Conodon	1324	Cordylus	865
141	antillanus	1324	Coregoni	461
505	nobilis	1324	Coregoninae	461
650	pacifæi	1316	coregonoides, Paralopis	602
480	plumieri	1324	Coregonus	461, 462, 465
231	sorrifer	1324	allus	466
231	Conorhynchus	411	angusticeps	466
231	conorynchus, Albula	411	clupeiformis	465, 469
231	conspersa, Muræna	397	couesii	463
966	Tigoma	234	coulterii	462
353	conspersus, Gymnothorax	397	harengus	469
356	Lycodontis	397	hoylei	468, 470
360	Serranus	1156	kennicotti	464
355	Squalius	234	labradoricus	466
355	Constantino de las Aletas Prictas	1119	latior	466
355	constellatus, Chiropsis	1868	lucidus	471
356	Piatophrys	2663	merckii	470
360	Sebastichthys	1807	nelsonii	466
355	Sebastodes	1806	neohantoniensis	466
356	consuetus, Salmo	479	nigripinnis	472
387	continuum, Hamulon	1297	nova-angliae	465
355	contractus, Rhinogobius	2236	osmeriformis	468
355	corrainii, Tylosurus	717	otsego	466
355	conus, Moxostoma	196	prognathus	472
355	Ptychostomus	196	quadrilateralis	465
355	convexa Turdus cauda	1145	richardsonii	465, 2816
355	convexifrons, Pomotis	1003	ruher	538
355	Cony, Horny	1715	sapidissimus	466
355	Cook, Rock	1375, 1576	signifer	518
385	cookianus, Aphododerus	787	thymalloides	518
352, 354	cooperi, Cheonda	236	tullibee	473
354	Leuciscus	236	bisselli	473
354	Metoponopus	2680	williamsoni	463
355	Raia	73	cismontanus	463
356	Salmo	483	coregonus, Moxostoma	191
357, 2601	Squalius	236	Ptychostomus	191
279	copei, Acipenser	196	coretta, Thynnus	870

	Page.		Page.
coriaceus, Eleutheractis .....	1233	Corvina armata .....	1437
Rypticus .....	1233	biloba .....	1460
corinus, Hexanchus .....	18	chrysoleuca .....	1439
cornura, Thyrsoides .....	394	deliciosa .....	1456
Cornet-fishes .....	755	dentex .....	1426
Corneta .....	757	fulgens .....	1435
cornifer, Achirus .....	2698	furerea .....	1460
cornubica, Lamna .....	40, 2749	furthi .....	1441
cornubiens, Squalus .....	49	macrops .....	1428
cornubiensis, Lepidogaster .....	2108	microps .....	1445
Pimelepterus .....	964	monacantha .....	1419
Rhombus kevia .....	2654	neglecta .....	1484
cornutus, Anoplogaster .....	840	ocellata .....	1454
Chaetodon .....	1688	ophioscion .....	1448
Chologaster .....	703	oscula .....	1484
Cyclichthys .....	1749	oxyptera .....	1222
Cyprinus .....	282	richardsoni .....	1484
Holecanthus .....	1685	ronchus .....	1436
Hypsilepis .....	283	saturna .....	1457
cerasinus .....	283	stearnsi .....	1458
cyaneus .....	283	stellifera .....	1445
gibbus .....	283	subequalis .....	1429
Leuciscus .....	283	trispinosa .....	1443
Minnilus .....	283	vermicularis .....	1453
Notropis .....	281	Corvina de las Aletas Amarillas .....	1410
cyaneus .....	283	corvinaeforme, Brachydeuterus .....	1326
frontalis .....	283	Haemulon .....	1327
Sillurus .....	759	Pomadasis .....	1327
Zanclus .....	1687, 1688	Corvinus (Johnius) jacobi .....	1457
coro, Pristipoma .....	1324	Corvula .....	1427
Sclera .....	1324	batabana .....	1430
coroides, Umbrina .....	1466	macrops .....	1427, 1428
Coronado .....	903	sanctae-luce .....	1429
coronata, Seriola .....	905	sialis .....	1428
coronatus, Cyclopterus .....	2097	subequalis .....	1429
Enneacetrus guttatus .....	1142	Corynolophus .....	2733
Halatractus .....	905	reiuhardtii .....	2733
Petrometopon cruentatus .....	1142	Coryphæna .....	952
Serranus .....	1142	argyrurus .....	953
Zonibthys .....	905	aurata .....	953
Corporal .....	221	azorica .....	953
corporalis, Cyprinus .....	221, 222	caerulea .....	1653
Leucosomus .....	222	chrysurus .....	952
Semotilus .....	221, 222	dolfyn .....	953
Corpore oblongo glabro .....	2657	dorado .....	953
Corsair .....	1808	equisetis .....	953
cortezianus, Aprodon .....	2461	fasciolatus .....	952
coruscans, Aretozonus .....	601	hippurns .....	952
Paralepis .....	602	immaculata .....	953
Sudis .....	602	imperialis .....	952
coruscus, Holocentrus .....	851	leasonii .....	953
Corvalos .....	1477	lineata .....	1619
Corvina .....	1408, 1425, 1455, 1461	maregravii .....	953
acutirostris .....	1437	novacula .....	1619
adusta .....	1448	percliformis .....	964
argyroleuca .....	1434	plumieri .....	2276

Page.		Page.		Page.
1437	<i>Coryphaena punctulata</i> .....	953	<i>Cottus aleuticus</i> .....	1957, 2862
1460	<i>scomberoides</i> .....	953	<i>alvordii</i> .....	1952
1439	<i>sueuri</i> .....	953	<i>anceps</i> .....	1973
1456	<i>virgata</i> .....	953	<i>anna</i> .....	1960
1426	<i>vlamingii</i> .....	953	<i>asper</i> .....	1944
1435	<i>Coryphaenidae</i> .....	951	<i>axillaris</i> .....	1981
1460	<i>Coryphanoides</i> .....	2578	<i>bairdi</i> .....	1950
1441	<i>carapinus</i> .....	2579	<i>beldingii</i> .....	1958
1428	<i>norvegicus</i> .....	2579	<i>bicornis</i> .....	1913
1445	<i>rupestris</i> .....	2578	<i>boleoides</i> .....	1968
1419	<i>Coryphaena lineolata</i> .....	1619	<i>brandti</i> .....	1984
1484	<i>nigrescens</i> .....	1200	<i>brodamus</i> .....	2006
1454	<i>psittacus</i> .....	1619	<i>hubalis</i> .....	1972
1448	<i>Coryphopterus</i> .....	2210	<i>cataphractus</i> .....	2053, 2066
1484	<i>glaucofrœnum</i> .....	2220	<i>cephaloides</i> .....	2008
1222	<i>Corythoichthys</i> .....	761, 763, 772, 2838	<i>cha-tolon</i> .....	2316
1484	<i>albirostris</i> .....	772, 2838	<i>cirris plurimis</i> .....	2066
1436	<i>cayannensis</i> .....	2838	<i>claviger</i> .....	1939
1457	<i>cayorum</i> .....	2838	<i>cognatus</i> .....	1954, 1955
1458	<i>cosmopolita</i> , <i>Micropteryx</i> .....	938	<i>copei</i> .....	1968
1445	<i>Seriola</i> .....	938, 2847	<i>eriniger</i> .....	2013
1429	<i>Cossyphus</i> .....	1581	<i>decastrensis</i> .....	1983
1443	<i>bodianus</i> .....	1583	<i>dicercaus</i> .....	1941
1453	<i>darwinii</i> .....	1586	<i>elegans</i> .....	1939
1410	<i>diplothenia</i> .....	1582	<i>evermanni</i> .....	1945
1326	<i>eclancheri</i> .....	1583	<i>fabricii</i> .....	2009
1327	<i>pectoralis</i> .....	1582	<i>formosus</i> .....	1969
1327	<i>puellaris</i> .....	1584	<i>franklini</i> .....	1967
1457	<i>pulchellus</i> .....	1584	<i>glaber</i> .....	2316
1427	<i>rufus</i> .....	1583	<i>glacialis</i> .....	1976
1430	<i>terres</i> .....	1583	<i>gobio</i> .....	1941, 1968, 2009
1427, 1428	<i>costatesi</i> , <i>Smaragdus</i> .....	2225	<i>gobioides</i> .....	1968
1429	<i>costatus</i> , <i>Bodianus</i> .....	1462	<i>gracilis</i> .....	1968
1428	<i>Micropogon</i> .....	1462	<i>grœnlandicus</i> .....	1975
1429	<i>costellatus</i> , <i>Chirus</i> .....	1869	<i>gulosus</i> .....	1944
2733	<i>Cottidae</i> .....	1879	<i>honilepidotus</i> .....	1936
2733	<i>Cottinae</i> .....	1882	<i>hexacornis</i> .....	2003
952	<i>Cottogaster</i> .....	1044	<i>hirundo</i> .....	2011
953	<i>aurantiacus</i> .....	1041	<i>hispidus</i> .....	2023
953	<i>chenevi</i> .....	2851	<i>humilis</i> .....	1979
953	<i>copelandi</i> .....	1045	<i>ietalops</i> .....	1950
1653	<i>patnami</i> .....	1046	<i>indicus</i> .....	2092
952	<i>shumardi</i> .....	1046, 2851	<i>jaok</i> .....	1978
953	<i>uraniden</i> .....	1044, 1045	<i>japonicus</i> .....	2036
953	<i>Cottopsis</i> .....	1942	<i>klamathensis</i> .....	1955
953	<i>asper</i> .....	1944	<i>labradoricus</i> .....	2004
952	<i>gulosus</i> .....	1945	<i>leipomus</i> .....	1962
952	<i>parvus</i> .....	1945	<i>maculatus</i> .....	1972
953	<i>semiscaber</i> .....	1950	<i>marginatus</i> .....	1966
952	<i>Cottunculus</i> .....	1992	<i>marmoratus</i> .....	1983
953	<i>microps</i> .....	1992	<i>meridionalis</i> .....	1951
1619	<i>thomsonii</i> .....	1993	<i>uertensii</i> .....	1986
953	<i>torvus</i> .....	1994	<i>miutus</i> .....	1958
1619	<i>Cottus</i> .....	1493, 1941, 1953, 1970	<i>mitchilli</i> .....	1973
964	<i>acadian</i> .....	2023	<i>monopterygius</i> .....	2092
2276	<i>aeneus</i> .....	1973	<i>niger</i> .....	1983, 1986



	Page.		Page.
<i>Cottus nigricans</i> .....	1973	<i>Couchii</i> , Serranus .....	1129
<i>nivosus</i> .....	1985	Couchu .....	160
<i>octodecimspinosus</i> .....	1976	<i>couesii</i> , Agosia .....	310
<i>onychus</i> .....	1953	<i>Apocope</i> .....	310
<i>pachypus</i> .....	1973	<i>Ceratias</i> .....	2732
<i>perplexus</i> .....	1955	<i>Coregonus</i> .....	463
<i>philonips</i> .....	1959	<i>Cryptopsaras</i> .....	2731
<i>pistilliger</i> .....	2008	<i>Prosopium</i> .....	463
<i>platycephalus</i> .....	1983, 1988	<i>Conesius</i> .....	323
<i>polaris</i> .....	1999	<i>adustus</i> .....	325
<i>pollicaris</i> .....	1941, 1953	<i>dissimilis</i> .....	324
<i>polyacanthocephalus</i> .....	1977	<i>greeni</i> .....	324
<i>porosus</i> .....	1975	<i>physignathus</i> .....	326
<i>princeps</i> .....	1962	<i>plumbeus</i> .....	323
<i>punctulatus</i> .....	1948, 1951	<i>prothemius</i> .....	324
<i>quadricornis</i> .....	2001	<i>squamilentus</i> .....	323
<i>quadrifilis</i> .....	1998, 2000	<i>Couia</i> .....	183
<i>rhotheus</i> .....	1946	<i>coulterii</i> , <i>Coregonus</i> .....	462
<i>ricel</i> .....	1952	<i>courbina</i> , <i>Pogonathus</i> .....	1483
<i>richardsoni</i> .....	1951	<i>Pogonias</i> .....	1483
<i>scorpio</i> .....	1973	<i>cromis</i> .....	1483
<i>scorpioides</i> .....	1973	<i>Courpata</i> .....	976
<i>scorpius</i> .....	1974	<i>courtadei</i> , Serranus .....	1152
<i>groenlandicus</i> .....	1975	<i>courvina</i> , Johnius .....	1419
<i>semiscaber</i> .....	1949	<i>Sciæna</i> .....	1419
<i>semiscabra centroleura</i> .....	1945	<i>Cow-fish</i> .....	1724
<i>shasta</i> .....	1947	<i>Cow-nose Ray</i> .....	90
<i>spilotus</i> .....	1961	<i>Cow-pilot</i> .....	1561
<i>stelleri</i> .....	1941	<i>Cow Shark</i> .....	19
<i>taeniopterus</i> .....	1979, 1988	<i>Sharks</i> .....	17
<i>tentaculatus</i> .....	2000	<i>Crab Eater</i> .....	948
<i>thomsonii</i> .....	1994	<i>Crabra</i> .....	1837
<i>trachurus</i> .....	1936	<i>Cracker, Clam</i> .....	83
<i>tricuspis</i> .....	2009	<i>ragini</i> , <i>Amiurus</i> .....	141
<i>tripterygius</i> .....	2023	<i>Etheostoma</i> .....	1091
<i>uncinatus</i> .....	1906	<i>Craig Fluke</i> .....	2656
<i>ventralis</i> .....	2008, 2009	<i>crameri</i> , <i>Leuresthes</i> .....	802
<i>verrucosus</i> .....	1980	<i>Sebastes</i> .....	1799
<i>villosus</i> .....	2022	<i>Crampfish</i> .....	77
<i>virginianus</i> .....	1976	<i>Cranlomi</i> .....	781, 2146
<i>viscosus</i> .....	1968	<i>Crappet</i> .....	987
<i>wilsoni</i> .....	1952	<i>Crappies</i> .....	986, 987
<i>Cotylis nigripinnis</i> .....	2332	<i>crassa</i> , <i>Bcione</i> .....	716
<i>nuda</i> .....	2331	<i>Tigoma</i> .....	231
<i>stannii</i> .....	2332	<i>crassicauda</i> , <i>Lavinia</i> .....	231
<i>stelleri</i> .....	2104	<i>Lenciscus</i> .....	231
<i>ventricosus</i> .....	2104	<i>Sibona</i> .....	231
<i>Cotylopus</i> .....	2207	<i>crassiceps</i> , <i>Melamphaes</i> .....	843
<i>gymnogaster</i> .....	2207	<i>Plectronus</i> .....	843
<i>salvini</i> .....	2208	<i>Scopelus</i> .....	843
<i>couchi</i> , <i>Dionda</i> .....	216	<i>crassilabre</i> , <i>Moxostoma</i> .....	194, 196
<i>Montana</i> .....	272	<i>crassilabris</i> , <i>Embryx</i> .....	2458
<i>Couchia argentata</i> .....	2550	<i>Geophagus</i> .....	1543
<i>couchiana</i> , <i>Limla</i> .....	695	<i>Lycodopsis</i> .....	2458
<i>Pœclia</i> .....	695	<i>Lycolia</i> .....	2869
<i>couchii</i> , <i>Pœclia</i> .....	695	<i>Ptychostomus</i> .....	194

Page.		Page.		Page.
1129	<i>crassilabris</i> , <i>Satanoperca</i> .....	1542	<i>cristagalli</i> , <i>Gobius</i> .....	2209
160	<i>crasna</i> , <i>Alvordius</i> .....	1034	<i>cristata</i> , <i>Chimera</i> .....	95
310	<i>Esox</i> .....	627	<i>cristatus</i> , <i>Adonis</i> .....	2383
310	<i>Lepidosteus</i> .....	110	<i>Blennius</i> .....	2382, 2383
2732	<i>Squalius</i> .....	231	<i>cristiceps</i> , <i>Melamphaes</i> .....	844
463	<i>Tylosurus</i> .....	716	<i>Plectromus</i> .....	843
2731	<i>craticula</i> , <i>Zygonectes</i> .....	657	<i>Cristivomer</i> .....	504
463	<i>Cratinus</i> .....	1188	<i>namaycush</i> .....	504
323	<i>agassizi</i> .....	1188	<i>siscowet</i> .....	505
325	<i>Cravo</i> .....	954	<i>cristulata</i> , <i>Scorpena</i> .....	1841
324	<i>Crawl-a-bottom</i> .....	181, 1038	<i>Crius</i> .....	970
324	<i>Crayracion</i> .....	1746	<i>bertheloti</i> .....	971
326	<i>crebripunctata</i> , <i>Pteroplatea</i> .....	87	<i>Croaker</i> .....	1460, 1484
323	<i>Creek Chub</i> .....	222	<i>Black</i> .....	1456
324	<i>Creekfsh</i> .....	185	<i>White</i> .....	1397
323	<i>Cremnobates</i> .....	2369	<i>Yellow-tailed</i> .....	1467
183	<i>affinis</i> .....	2372	<i>croaker</i> , <i>Sciema</i> .....	1462
462	<i>altivelis</i> .....	2371	<i>Croakers</i> .....	1392, 1461
1483	<i>fasciatus</i> .....	2373	<i>croceus</i> , <i>Leuciscus</i> .....	308
1483	<i>integripennis</i> .....	2373	<i>Rhinichthys atronaso</i> .....	308
1483	<i>marmoratus</i> .....	2371	<i>crocodilinus</i> , <i>Braichysomopsis</i> .....	388
976	<i>monophthalmus</i> .....	2372	<i>Ophichthys</i> .....	388
1152	<i>nox</i> .....	2374	<i>Ophisurus</i> .....	388
1419	<i>cremnobates</i> , <i>Labrosomus</i> .....	2366	<i>crocodilus</i> , <i>Gasteropolecus</i> .....	558
1419	<i>Starksia</i> .....	2365, 2366	<i>Lampanyctus</i> .....	558
1724	<i>Crenilabrus</i> .....	1581	<i>Scopelus</i> .....	558
90	<i>microstoma</i> .....	1576	<i>crocota</i> , <i>Plectropoma</i> .....	1192
1561	<i>crenulare</i> , <i>Myctophum</i> .....	575	<i>crocotus</i> , <i>Hypoplectrus</i> .....	1192
19	<i>crenularis</i> , <i>Tarletonbeania</i> .....	575	<i>unicolor</i> .....	1192
17	<i>crenulatus</i> , <i>Rhombus</i> .....	966	<i>crocro</i> , <i>Pomadasis</i> .....	1333
948	<i>Creole</i> .....	1586	<i>Pristipoma</i> .....	1333
1837	<i>Fish</i> .....	1221	<i>croicensis</i> , <i>ichthys</i> .....	1651
83	<i>croolus</i> , <i>Brachyrhinus</i> .....	1222	<i>Scarus</i> .....	1650
141	<i>Paranthias</i> .....	1222	<i>Cromileptes</i> .....	1148
1091	<i>Serranus</i> .....	1222	<i>cromis</i> , <i>Labrus</i> .....	1483
2656	<i>crestantale</i> , <i>Gobiosoma</i> .....	2259	<i>Pogonias</i> .....	1482
802	<i>crestantalis</i> , <i>Gebiosoma</i> .....	2260	<i>courbina</i> .....	1483
1799	<i>Pomacanthus</i> .....	1682	<i>crossotus</i> , <i>Etropus</i> .....	2689
77	<i>cresecentia</i> , <i>Salmo gairdneri</i> .....	2821	<i>erotalina</i> , <i>Lycolia</i> .....	2869
781, 2146	<i>Crested Gobies</i> .....	2209	<i>crotalinus</i> , <i>Embryx</i> .....	2458
987	<i>crestonis</i> , <i>Teuthis</i> .....	1692	<i>Lucodopsis</i> .....	2459
986, 987	<i>Crevallé</i> , <i>Horse</i> .....	920	<i>Crotalopsis</i> .....	386
716	<i>Crevallés</i> .....	915, 920, 921	<i>mordax</i> .....	387
231	<i>crinigerum</i> , <i>Siphostoma</i> .....	771	<i>punctifer</i> .....	387
231	<i>crinitus</i> , <i>Alectis</i> .....	932	<i>crotophus</i> , <i>Cherojulius</i> .....	1598
231	<i>Blennius</i> .....	2383	<i>Julis</i> .....	1531, 1598
231	<i>Blepharichthys</i> .....	932	<i>PlatyGLOSSUS</i> .....	1598
843	<i>Blepharis</i> .....	932	<i>Crucian Carps</i> .....	201
843	<i>Caranx</i> .....	932	<i>cruentatus</i> , <i>Bodianus</i> .....	1142
843	<i>Gallichthys</i> .....	932	<i>Labrus</i> .....	1238
194, 196	<i>Zeus</i> .....	932	<i>Petrometopon</i> .....	1141
2458	<i>Criolla</i> , <i>Cherna</i> .....	1157	<i>coronatus</i> .....	1142
1543	<i>Criollo</i> , <i>Pargo</i> .....	1265	<i>Priacanthus</i> .....	1238, 2858
2458	<i>cristagalli</i> , <i>Anoplarchus</i> .....	2423	<i>Sparus</i> .....	1142
2869	<i>Cebedichthys</i> .....	2427	<i>cruentifer</i> , <i>Pisoodonopsis</i> .....	377
194	<i>Centronotus</i> .....	2423	<i>crumenophthalmus</i> , <i>Caranx</i> .....	911

	Page.		Page.
crumenophthalmus, Scouber .....	911	Cubera .....	1254
Trachurops .....	911	cubera, Lutjanus .....	1255
crureum Xiphidion .....	2425	Cubiceps .....	950
crureus, Squalius .....	233	indicus .....	951
Cryptacanthodes .....	2443	multiradiatus .....	951
inornatus .....	2443	pauciradiatus .....	951
maculatus .....	2443	cubifrons, Malthe .....	2738
Cryptacanthodide .....	2442	Cub-shark .....	38
Cryptops .....	341	Cuckold .....	1724
Cryptopsarus .....	2731	cuculus, Trigla .....	2177
cousii .....	2731	cucuri, Prionodon .....	40
Cryptopterus .....	381, 382	Cucuyo .....	1701, 1709
puncticeps .....	382	Cugupgnacu .....	1158, 1163
cryptosus, Stromatosus .....	968	cujus, Chasmistes .....	183
Cryptotomus .....	1621	Culius .....	2199
aeropunctatus .....	1624	aequidens .....	2202
beryllinus .....	1624	amblyopsis .....	2200
dentiens .....	1623	belizianus .....	2201
retractus .....	1623	perniger .....	2201
rosceus .....	1626	cultrata, Novacula .....	1619
rustus .....	1624	cultratus, Stolephorus .....	443
Cryptotrema .....	2366	Xyrichtus .....	1619
corallinum .....	2366	cultriferum, Pristipoma .....	1333
crysoleucas, Abramis .....	250	cultrifrons, Alutera .....	1718
bosci .....	251	Cultus Cod .....	1875
Cyprinus .....	250	culveri, Trachinotus .....	942
crynos, Caranx .....	921	cumberlandicum, Etheostoma fla-	
Scomber .....	921	bellare .....	1098
Crystallaria .....	1060	cuningil, Ceratichthys .....	318
asprella .....	1061	Hybopsis .....	318
Crystallichthys .....	2864	cuneata, Pécilia .....	2834
mirabilis .....	2865	Cunner .....	1576, 1577
crystallina, Atherinella .....	805	cupreoides, Pimelodus .....	140
Thyrius .....	804	cupreus, Pimelodus .....	140
Crystalligobiinae .....	2192	Silurus .....	140
Ctenodax .....	975	Trachinotus .....	944
Ctenodon .....	2432	Trachynotus .....	944
maculatus .....	2433	curassavicus, Ballstes .....	1709
Utenogobius .....	2210, 2211, 2218	curema, Mugil .....	813, 2841
fasciatus .....	2223	curlicus, Agonus .....	2036
Ctenolabrus adpersus .....	1577	curilus, Salmo .....	508, 2853
burgull .....	1577	Curimata .....	332
caeruleus .....	1577	magdalene .....	332
chogaet .....	1577	Curimatella .....	332
uninotatus .....	1577	Curimatine .....	331
cuba, Vomer .....	934	Curimatus .....	332
Cuban Blindfish .....	2501	magdalene .....	332
cubana, Anguilla .....	348	curtus, Stolephorus .....	445
Murena .....	348	Vomer .....	934
cubanus, Engraulis .....	442	curvidens, Congrus .....	360
Epinephelus .....	1159	curvillneata, Murenophis .....	395
Stolephorus .....	442	curvus, Tetrodon .....	1728
cubensis, Channomurena .....	404	Cusk Eels .....	2481, 2487
Hynnis .....	932	Cusks .....	2561
Lutia .....	692	cuspicauda, Alutera .....	1718
Pécilia .....	692	cutisauerinus, Carpiodes .....	167

Page.		Page.		Page.
1254	Cutlass Fishes.....	888	Cylichthys cornutus.....	1740
1255	Cut-lips.....	199, 327	Cycloganoidea.....	111
950	Cut-throat Trout.....	487, 492, 493	Cyclogaster.....	2114
951	ouvieri, Caranx.....	910	pulchellus.....	2127
951	Centropomus.....	1121	cyclogaster, Liparis.....	2118
951	Gasterosteus bispinosus.....	749	Cyclogobius.....	2240
2738	Tetragonurus.....	976	cyclolepis, Microgobius.....	2247
38	Trachurus.....	910	Moseleya.....	2570
1724	cuzamile, Scarus.....	1648	Nematonurus.....	2571
2177	cyanea, Furcaria.....	1547	Zalypnus.....	2246
40	cyanelus, Apomotis.....	996	Cyclonarce.....	78
1701, 1709	Ichthyobus.....	164	cyclopomatus, Serranus.....	1175
1158, 1163	Lepomis.....	996	Cyclopaetia.....	2675
183	cyaneus, Chromis.....	1547	chittendeni.....	2676
2199	Heliastes.....	1547	fimbriata.....	2676
2202	Hypsilepis cornutus.....	283	querna.....	2675
2200	Notropis cornutus.....	283	Cyclopteichthys.....	2103
2201	Cyanichthys.....	1747	glaber.....	2104
2201	cyanocephalus, Iridio.....	1594	stelleri.....	2104
1619	Labrus.....	1594	ventricosus.....	2104
443	Lythrus.....	300	Cyclopteridae.....	2094
1619	Minnilus.....	300	Cyclopterinae.....	2095
1333	Notropis umbratilis.....	300	Cyclopteroidea.....	2102
1718	cyanoguttatus, Herichthys.....	1538	gyrinops.....	2102
1875	Heroa.....	1537	Cyclopterus.....	2096
942	cyanolene, Sparisoma.....	1633	caeruleus.....	2097
cyanonoton, Alosa.....	427	callyodon.....	2110	
Cyanoperca.....	1022	coronatus.....	2097	
cyanophrys, Naucrates.....	900	gelatinosus.....	2135	
Paenes.....	950	lineatus.....	2118	
cyanops, Caulolatilus.....	2278	liparis.....	2118, 2123	
1570, 1577	cyanopectera, Chrysophrys.....	1354	major.....	2118
140	cyanopterus, Cypsilurus.....	2836	minor.....	2121
140	Exocoetus.....	739	liparoides.....	2108
140	Lutjanus.....	1255	lumpus.....	2096, 2097
944	Mesoprion.....	1255	minutus.....	2097
944	Neomænis.....	1254	montacuti.....	2108
1709	cyanostigma, Charojulis.....	1591	musculus.....	2118
813, 2841	Julis.....	1591	nudus.....	2336
2036	PlatyGLOSSUS.....	1591	orbis.....	2100
508, 2823	Cybium.....	873	pavoninus.....	2097
332	acervum.....	875	pyramidatus.....	2097
332	caballa.....	876	spinosus.....	2099, 2100
332	cavalla.....	876	stelleri.....	2104
331	immaculatum.....	876	ventricosus.....	2104
332	maculatum.....	874	cyclops, Liparis.....	2112, 2118
332	petus.....	877	Cyclospindylus.....	52, 53
445	regale.....	875	Cyclospindylous Sharks.....	52
934	sara.....	877	cyclostigma, Liparis.....	2125
360	secundri.....	877	Cyclothone.....	581, 582
395	verany.....	877	bathypbila.....	582
1728	Cycleptinae.....	102	elongata.....	583
2481, 2487	Cycleptus.....	108	lusca.....	582
2561	elongatus.....	168	microdon.....	582, 2826
1718	nigrescens.....	169	cyclotis, Ceratichthys.....	323
167	Cylichthys.....	1747, 1748	cylindraceus, Rivulus.....	062

	Page.		Page.
Cylindrosteus.....	109, 110	Cynoscion thalassinus.....	1407
agassizii.....	111	vireacens.....	1415
bartoni.....	111	xanthulus.....	1410
castelnaui.....	111	Cynoscion acoupa.....	1404
productus.....	111	cypho, Catostomus.....	184
raffinesquiei.....	111	Esox.....	627
zadocki.....	111	Cyphosus bosqui.....	1388
Cymatogaster.....	1498, 1502	cyprinaceus, Chaetodon.....	1388
aggregatus.....	1498	Cyprinella.....	254, 256, 273
ellipticus.....	1503	analostana.....	279
larkinsii.....	1503	beckwithii.....	273
minimus.....	1497	billingsiana.....	272
pulchellus.....	1503	bubalina.....	273
rosacens.....	1500	callura.....	275
1053		cercostigma.....	275
cymatogramma, Plecoma.....	1053	complanata.....	272
cyuatotænia, Etheostoma (Hadrop- terns).....	1042	forbesi.....	272
Hypohomus.....	1041	gunnisoni.....	273
Cynædus brama.....	1360	lepidæ.....	273
Cynichthys.....	1148	ludibunda.....	273
Cynicoglossus.....	2653	lugubris.....	274
bathybius.....	2656	luxoides.....	274
pacificus.....	2655	macrostoma.....	274
Cynocephalus.....	33	notata.....	274
cynodon, Lutjanus.....	1255	rubripinna.....	281
Mesoprion.....	1255, 1266	suavis.....	272
Cynoglossa.....	2653	texana.....	274
microcephala.....	2655	umbrosa.....	273
cynoglossa, Solea.....	2657	venusta.....	274
Cynoglossina.....	2693	whippilii.....	279
cynoglossus, Glyptocephalus.....	2656	cyprinella, Chanos.....	415
Pleuronectes.....	2611	Ictiobus.....	163
Cynoperca.....	1020, 1021	Sclerognathus.....	164
Cynoponticus.....	359	Cyprinidae.....	199, 200
ferox.....	360	Cyprinina.....	201
Cynoscion.....	1401, 1403	Cyprinodon.....	670
acoupa.....	1403	baileyi.....	675
albus.....	1411	bovinus.....	673
carolinensis.....	1409	californiensis.....	674
jamaicensis.....	1406	carpio.....	675
leiarcbus.....	1414	elegans.....	675, 2832
macdonaldi.....	1411	eximius.....	673, 2832
maculatum.....	1400	felicianus.....	676, 2832
microlepidotus.....	1415	gibbosus.....	672
nebulosus.....	1409	latifasciatus.....	676
nobilis.....	1413	macularius.....	674
nothus.....	1406	baileyi.....	675
obliquatus.....	1405	martæ.....	675
othonopterus.....	1404	mydrus.....	676
parvipinnis.....	1410	nevadensis.....	674
phoxocephalus.....	1413, 2859	parvus.....	666
regale.....	1407	riverendi.....	673, 2832
regalis.....	1407	variegatus.....	671
reticulatus.....	1408	Cyprinodontina.....	631
squampinnis.....	1404, 1405	cyprinoides, Gobius.....	2209
stolzmanni.....	1412	Lophogobius.....	2209

Page.		Page.		Page.
1407	Cyprinus	201	Dacentrus	1495
1415	americanus	250, 251, 1475	luconis	1496
1410	atromaculatus	222	Daetylagmus	2304
1404	atrouasus	307	mundus	2304
184	auratus	291	daetyloptera, Scorpaena	1837
627	balteatus	239	Daetylopterus	2183
1388	bullaris	221	communis	2184
1388	burtonianus	2798	pirapela	2183
54, 256, 273	carpio	291	volitans	2183
279	catostomus	176	daetylopterus, Helicolenus	1837
273	caurinus	220	Sebastes	1837, 1838
272	commersonii	179	Sebastopus	1837
273	cornutus	282	Dactyloscopidae	2297
275	corporalis	221, 222	Dactyloscopus	2300, 2301
275	crysoloucas	250	lunaticus	2302
272	gracilis	326	pectoralis	2301
272	hemiplus	250	poeyi	2302
273	maxillingna	327	tridigitatus	2301
273	megalops	282	zelotes	2303
273	melanurus	282	daetylosus, Paraliparis	2144
274	oblongus	186	Daddy Sculpin	1974
274	oregonensis	225	Daector	2325
274	pala	415	dowi	2325
274	smithii	413	Dajaus	818, 819
281	sneceta	186	choirorynchus	2841
272	sueuril	195	microps	820
274	teres	179	monticola	2841
273	tolo	415	dakotensis, Chrosomus	210
274	vittatus	307	Dalatidae	56, 620
279	cyprinus, Carpiodes	107, 168	dalli, Gobius	2230
415	Cypselurus	730, 731, 735, 2835, 2836	Dallia	621
163	bahiensis	2836	delicatissima	621
164	californicus	2836	pectoralis	621
199, 200	callopterus	2836	dallii, Pteropodus	1819
201	comatus	736	Sebastodes auriculatus	1818
670	cyanopterus	2836	Damalichthys	1509
675	furcatus	737, 2836	argyrosoma	1510
673	gibbifrons	2836	argyrosomus	1509
674	heterurus	2836	vacca	1510
675	lineatus	2836	damalis, Carpiodes	107
675, 2832	nigricans	2836	d'Amplora, Maire	555
673, 2832	xenopterus	2836	Dark-green Parrot-fish	1638
676, 2832	ypselurus, Prognurus	2866	Darter, Black-sided	1028, 1032
672	Cyttinae	1660	Blue	1088
676	Cyttus hololepis	1662	Blue-breasted	1076
674			Fan-tailed	1079
675			Green-sided	1053
675	Dab, Alaska	2645	Johnny	1056
676	Rusty	2644	Least	1104
674	Smear	2654	Manitou	1027
666	Dabbler, Mud	640	Rainbow	1086
673, 2832	Dabs, Mud	2644	Tessellated	1057
671	Smear	2653	Sand	1061, 1062
631	Dace	228, 281	darwini, Cossyphus	1586
2209	Black-nosed	305, 307	Pimelometopon	1586
2209	Long-nosed	306	Sebastes	1832
2209	Dace, Red-bellied	209		

	Page.		Page.
darwini, Sebastodes .....	1832	declivifrons, Abudedefduf.....	1562
Trochocopus .....	1586	Euschistodus .....	1562
Dasibatis .....	82	Glyphidodon.....	1562
centrura.....	83	declivis, Seriola.....	905
diptera.....	85	Zonichthys.....	905
hastata .....	84	Decodon .....	1584
longa .....	85	puellaris.....	1584
sabina .....	85	decoratus, Entomacrodus .....	2390
sayi .....	86	Promicropterus.....	1234
tuberculata .....	84	Rhypticus .....	1234
dasypilotus, Pseudonopha.....	2803	decurrens, Pleuronichthys.....	2637, 2638
Dasyatida .....	79	de Casta, Aguja.....	892
Dasyatinae .....	79	Mojarra.....	1372
Dasyatis .....	82, 83	de Chapala, Pescado Blanco.....	792
centrura .....	83	de dos Coloros, Pescado Azul .....	1557
dipterna.....	85	Vaqueta .....	1684
gymnura .....	84	de España, Sardina.....	423
hastata .....	83	de Gallo, Pez.....	895
longa .....	85	de la Alto, Isabelito.....	1074
sabina .....	84	de la Piedras, Mojarra.....	1681
say .....	86	de las Piedras, Pez.....	1700
Dasybatus .....	60, 82	de Ley, Mojarra.....	1370
dipterus.....	85	de lo Alto, Cagon.....	1277
dasycephalus, Arius.....	130	Pargo.....	1262
Galichthys .....	2780	Seal.....	1261
Hexanemateichthys ..	130	de Mar, Aboma .....	2195
Dasyacopelus .....	574	Esmeralda .....	2204
spinosa.....	575	Esmeraldas .....	2203
Dasycottus .....	1991	de Marais, Poisson .....	113
setiger .....	1991	de Paladar, Aguja.....	892
daubentonii, Caranx .....	920	de Perdriz, Liza Ojo.....	814
daivisonii, Anisotremus .....	1321	de Playa, Cazon.....	36
Mouacanthus .....	1715	de Pluma, Pez.....	1347
Pomadasya.....	1321	de Raizero, Pargo.....	1273
Pristipoma.....	1321	de Río, Aboma .....	2236
daivisoni, Etheostoma .....	1049	Bagres .....	149
Ulocentra.....	1049	Lenguado .....	2698
Decactylus.....	173, 174, 177	de Vivero, Cherna.....	1160
decadactylus, Beryx .....	844	Deep-water Catalufas .....	977
decagonus, Agonus.....	2053	Gurrnards .....	2177
Aspidophorus .....	2054	Porgies .....	1344
Brachyopsis .....	2054	defensor, Caranx .....	921
Leptagonus .....	2052	de Gato Pai .....	1837
decagrammus, Chirus .....	1869	Dekaya .....	2276
Hexagrammos .....	1867	anomala.....	2277
Hexagrammus .....	1875	dekayi, Aclipenser.....	106
Labrax .....	1868	Gasterosteus .....	746
Decapterus .....	907	Isuropsis.....	48
hypodus .....	908	Isurus .....	48
macarellus .....	909	Phycis .....	2555
punctatus.....	907	Pimelodus.....	140
sanctae-helenae .....	908	Scomber.....	867
scombrinus .....	908	Syngnathus .....	771
Decaptus .....	906	delalandi, Labrisomus.....	2359
decastrensis, Cottus.....	1983	Malacotenus.....	2358
decimalis, Serranus .....	1175	delalandii, Citinus .....	2350

Page.		Page.		Page.
1562	de Ley, Sardina .....	430	denticulatus, Anarrhichas .....	2446
1562	delicatissima, Dallia .....	621	denticus, Calliodon .....	1623
1562	Umbra .....	621	Cryptotonus .....	1623
905	delicatissimus, Engraulis .....	444	denudatum, Gonostoma .....	570
905	Stolephorus .....	444	deppii, Cichlasoma .....	1524
1584	deliciosa, Clloa .....	272	Heros .....	1524
1584	Corvina .....	1456	deprandus, Esox .....	628
2399	Moniana .....	262	depressa, Belone .....	211, 713
1234	Selenis .....	1455	Fistularia .....	757
1234	Delolepis .....	2442	depressus, Lonchurus .....	1482
2637, 2638	virgatus .....	2442	de Raizero, Cabrilla .....	1171
892	Delothyris .....	2690	Derepodichthyidae .....	2480
1372	pellucidus .....	2691	Derepodichthys .....	2480
792	delphinus, Minomus .....	171	alepidotus .....	2480
1557	Pantosteus .....	171	Derichthyidae .....	343
1684	del Rey, Pescadillos .....	807	Derichthys .....	343
423	Pescado .....	806	serpentinus .....	343
895	Pez .....	799	dermatina, Lycopodus .....	2492
1674	Deltentosteus .....	2210	Salmo .....	479
1681	Deltistes .....	2794	Dermatolepis .....	1166, 1168
1700	luxatus .....	2792	angustifrons .....	1169
1370	de Mer, Brochet .....	1118	incermis .....	1167, 2855
1277	Demoiselle .....	1543, 1561	punctatus .....	1163
1262	dendritica Ancylosetta .....	2634	zancus .....	2854
1261	Ramularia .....	2633	dermatolepis, Epinephelus .....	1169
2195	denegatus, Pomacentrus .....	1567	Dermatostethus .....	761, 763
2204	dennyi, Liparis .....	2124	punctipinnis .....	763
2203	dentata, Chemonopsetta .....	2630	desmarestia, Raja .....	71
113	Platessa .....	2615, 2630	detersor, Julis .....	1610
892	Pomatopsetta .....	2615	detrusus, Gillichthys .....	2251
814	Stygicola .....	2500	Devilfish .....	92
36	dentatus Apsilus .....	1278	Devil, Sea .....	91, 92, 2727
1347	Grammatostomias .....	590	Dibasis .....	1291
1273	Hippoglossoides .....	2615	aurolineatus .....	1309
2236	Lucifuga .....	2500	chromis .....	1299
149	Lutjanus .....	1255	chrysopterus .....	1300
2698	Mesoprius .....	1279	elegans .....	1304
1160	Paralichthys .....	2629, 2630	flavolineatus .....	1306
977	Pleuronectes .....	2630	fremebundus .....	1297
2177	Pseudorhombus .....	2630, 2632	maculicauda .....	1314
1344	Stygicola .....	2500	obliquatus .....	1304
921	Tropidinius .....	1279	parra .....	1299
1837	Upeneus .....	859	plumieri .....	1306
2276	Dentex .....	1288	scudleri .....	1300
2277	filamentosus .....	1289	steindachneri .....	1302
106	dentex, Brycon .....	337	trivittatus .....	1311
746	Caranx .....	927	Diablo .....	2737
48	Chalcinopsis .....	337	Diabolichthys .....	92
48	Corvina .....	1426	alliotti .....	93
2555	Engraulis .....	451	diabolus, Raja marinus .....	93
140	Larimus .....	1426	Diacope .....	1247
867	Menidia .....	801	viridis .....	1246
771	Odontoscion .....	1425	diadema, Pseudosearus .....	1646
2359	Osmerus .....	524	Scarus .....	1646
2358	Seomber .....	927	Diagramma cavifrons .....	1343
2359	Dentleine .....	1244	melanospilum .....	1321



	Page.		Page.
Dialommus.....	2868	dimidiatus, Leucns.....	244
fuscus.....	2868	PlatyGLOSSUS.....	1594
diaphana, Hydrargyra.....	645	Serranus.....	1179
Raia.....	71	Syngnathus.....	765
Sternoptyx.....	603, 2826	Trisotropis.....	1179
Diaphanichthys.....	353	Dinectus.....	103
diaphanus, Calliurus.....	996	truncatus.....	106
Fundulus.....	645	Dinematichthys marginatus.....	2502
menona.....	645	ventralis.....	2503
Diaphasia.....	2495	Dinemus.....	854
Diaphus.....	564	venustus.....	854
theta.....	564, 565	dinemus, Minnilus.....	293
diaptera, Furcella.....	2472	dinoceros, Citharichthys.....	2682
Diapterus.....	1373, 1375	Diodon.....	1744, 1747
californiensis.....	1370	antennatus.....	1750
dowi.....	1368	asper.....	1744, 1752
gracilis.....	1370	atinga.....	1746, 1750
homonymus.....	1371	brachiatus.....	1746
lefroyi.....	1372	carinatus.....	1754
diapterus, Amlichthys.....	1113	echinus.....	1746
Lycodes.....	2473	fuliginosus.....	1749
Dibranchus.....	2743	geometricus.....	1748, 1749
atlanticus.....	2743	holocanthus.....	1746
diceraus, Ceratocottus.....	1940, 1941	hystrix.....	1744, 1746
Cottus.....	1941	liturosus.....	1746
Enophrys.....	1941	maculatus.....	1746
Dicerobatis.....	2756	maculifer.....	1747
Dicerobatus.....	92	maculostriatus.....	1748
Dick, Nigger.....	327	melanopsis.....	1746
Slippery.....	1595	meulini.....	1748
Dicrolene.....	2522	multimaenulatus.....	1746
intronigra.....	2522	nigrolineatus.....	1749
Dicromita.....	2506	novemmaculatus.....	1746
agassizii.....	2506	pilosus.....	1744, 1752
Dicrotus.....	882	punctatus.....	1746
parvipinnis.....	883	quadrifaculatus.....	1746
Dictyosomatina.....	2349	reticulatus.....	1751
diego, Scombor.....	867	rivulatus.....	1748
dienceus, Eupomacentrus.....	1552	schaeppi.....	1748
difformis, Carpiodes.....	175	seemaculatus.....	1746
digitatus, Lycodes.....	2466	spinosissimus.....	1746
digitis, Trigla vicensis.....	2183	spinous.....	1749
digrammus, Pleuronectes.....	2041	verrucosus.....	1749
dilecta, Anclpsetta.....	2336	Diodontidae.....	1742
Notosema.....	2636	diomedea, Aphoristia.....	2711
dilectum, Notosema.....	2635	diomedeanus, Hoplunnis.....	361
dilectus, Alburnus.....	294	Symphurus.....	2711
Notropis.....	294	Dionda.....	212, 213, 214
Dimalocentrus.....	1613	argentosa.....	215
di Mare, Cane.....	48	chrysitis.....	214
dimidiata, Mycteroperca.....	1179	couchl.....	216
dimidiatus, Epinephelus.....	1179	episcopa.....	215
Gasterosteus.....	749	grisca.....	216
Halichares.....	1594	melanops.....	216
Icthycallus.....	1594	papalis.....	214
Julis.....	1594	plumben.....	216

Page.		Page.		Page.
244	<i>Dionda punctifer</i> .....	215	<i>diacobolus, Catostomus</i> .....	172, 175, 2791
1504	<i>serena</i> .....	214	<i>Discocephali</i> .....	781, 2265
1179	<i>spadicea</i> .....	216	<i>Discopyge</i> .....	78
765	<i>texensis</i> .....	215	<i>omnata</i> .....	78
1179	<i>Dioplites</i> .....	1010	<i>diapar, Fundulus</i> .....	658
103	<i>nucensis treculii</i> .....	1012	<i>Zygonectes</i> .....	659
106	<i>variabilis</i> .....	1012	<i>dispilurus, Centropriasis</i> .....	1220
2502	<i>dipnemia, Hypailepis</i> .....	300	<i>Dules</i> .....	1219
2503	<i>diplemius, Minullus</i> .....	300	<i>dispilus, Halichæres</i> .....	1598
854	<i>Diplanchias</i> .....	1753	<i>Iridio</i> .....	1597
854	<i>nasus</i> .....	1754	<i>PlatyGLOSSUS</i> .....	1598
293	<i>Diplectrum</i> .....	1203, 1204, 1207	<i>dissimilis, Ceratichthys</i> .....	319
2682	<i>euryplectrum</i> .....	1206	<i>Conesius</i> .....	324
1744, 1747	<i>faeculare</i> .....	1208	<i>Hybopsis</i> .....	318
1750	<i>formosum</i> .....	1207	<i>Leucosomus</i> .....	324
1744, 1752	<i>macropoma</i> .....	1205	<i>Luxilus</i> .....	319
1746, 1750	<i>radiale</i> .....	1204	<i>distichus, Salmo</i> .....	509
1746	<i>radialis</i> .....	1205	<i>distinctum, Sparisoma</i> .....	1635, 1636
1754	<i>radians</i> .....	1208	<i>distinctus, Scarus</i> .....	1636
1746	<i>sciurus</i> .....	1204	<i>Ditrema</i> .....	1510
1749	<i>diplemius, Semotilus</i> .....	222	<i>aggregatum</i> .....	1499
1749, 1749	<i>Diplesion</i> .....	1052	<i>anale</i> .....	1501
1746	<i>blennioides</i> .....	1053	<i>arenatum</i> .....	1502
1744, 1746	<i>fasciatus</i> .....	1081	<i>argenteum</i> .....	1504
1746	<i>Diplesium blennioides</i> .....	1053	<i>atripes</i> .....	1507
1747	<i>Diptodus</i> .....	1362	<i>brevipinne</i> .....	1499
1748	<i>argentus</i> .....	1363	<i>caryi</i> .....	1509
1746	<i>caribæus</i> .....	1360	<i>furcatum</i> .....	1506
1748	<i>caudimacula</i> .....	1363	<i>jacksoni</i> .....	1505
1746	<i>flavolineatus</i> .....	1360	<i>lave</i> .....	1511
1748	<i>holbrookii</i> .....	1362	<i>laterale</i> .....	1506
1746	<i>probatocephalus</i> .....	1361	<i>megalops</i> .....	1502
1749	<i>rhomboidea</i> .....	1358	<i>orthonotus</i> .....	1507
1746	<i>sargus</i> .....	1363	<i>rhodoterum</i> .....	1503
1744, 1752	<i>Diplolepis</i> .....	1418	<i>temminckii</i> .....	1510, 1511
1746	<i>squamosissimus</i> .....	1419	<i>toxotes</i> .....	1508
1751	<i>diploproa, Sebastichthys</i> .....	1802	<i>vacca</i> .....	1510
1748	<i>Sebastodea</i> .....	1801	<i>Diver, Sand</i> .....	535
1748	<i>Diplospondyli</i> .....	16	<i>Dobula</i> .....	228
1746	<i>diploptenia, Belone</i> .....	712	<i>Doctor-fish</i> .....	1689, 1691
1746	<i>Bodianus</i> .....	1582	<i>dodecaedron, Agonus</i> .....	2046
1749	<i>Cossyphus</i> .....	1582	<i>Occa</i> .....	2044
1749	<i>Harpe</i> .....	1582	<i>dodecaedrus, Aspidophorus</i> .....	2046
1742	<i>diploptenia, Tylosurus</i> .....	712	<i>Brachyopsis</i> .....	2046
2711	<i>Dipterodon</i> .....	1106	<i>Dodecagrammos</i> .....	1866
361	<i>hexacanthus</i> .....	1107	<i>Dogfish</i> .....	113, 623
2711	<i>ruber</i> .....	1107	<i>California</i> .....	53, 54
212, 213, 214	<i>Dipteron chrysurus</i> .....	1433	<i>Dog Salmon</i> .....	478
215	<i>dipterna, Dasibatis</i> .....	85	<i>of Alaska</i> .....	478
214	<i>Dasyatis</i> .....	85	<i>Shark</i> .....	28, 29
216	<i>dipternus, Dasybatis</i> .....	85	<i>Snapper</i> .....	1257
215	<i>Dipterygonotus</i> .....	1365	<i>dolfyn, Coryphæna</i> .....	953
216	<i>Dipturus</i> .....	66	<i>dolichocephalus, Gobius</i> .....	2237
216	<i>dipus, Microdesmus</i> .....	2450	<i>dolichogaster, Blennius</i> .....	2417
216	<i>Discoboli</i> .....	1758	<i>Centronotus</i> .....	2417
214	<i>Discoboli liparidina</i> .....	2105	<i>Gunnellus</i> .....	2417

	Page.		Page.
dolichogaster, <i>Muraenoides</i> .....	2417	dorsalis <i>Citula</i> .....	930
<i>Pholis</i> .....	2416	<i>Galeus</i> .....	30
<i>Doliodon</i> .....	939	<i>Halatractus</i> .....	902
<i>Dollardee</i> .....	1005	<i>Hybopsis</i> .....	262
<i>Dollar-fish</i> .....	967	<i>Hypsypops</i> .....	1570
<i>Dollfish</i> .....	1374	<i>Macrurus</i> .....	2585
<i>Dolly Varden Trout</i> .....	507	<i>Microspathodon</i> .....	1568, 1570
<i>dolomieu</i> , <i>Micropterus</i> .....	1011	<i>Mustelus</i> .....	30
<i>Dolphin</i> , Common .....	952	<i>Pomatopron</i> .....	1570
Small .....	953	<i>Semotilus</i> .....	222
<i>Dolphins</i> .....	951, 952	<i>Seriola</i> .....	902
<i>dombey</i> , <i>Bdeliostoma</i> .....	6	<i>Umbrina</i> .....	1469
<i>Gastrobranchus</i> .....	6	<i>Vomer</i> .....	934
<i>Le Gastrobranco</i> .....	6	<i>dorsatus</i> , <i>Petromyzon marinus</i> .....	10
<i>Polistotrema</i> .....	6	<i>dorso</i> , <i>Perca monapterygia</i> .....	1433
<i>Dómine</i> .....	880	<i>dorsomacula</i> , <i>Girella</i> .....	1382
<i>dominicensis</i> , <i>Poecilia</i> .....	696	<i>dorsopunicans</i> , <i>Pomacentrus</i> .....	1557
<i>Vomer</i> .....	934	<i>Dorsuarius</i> .....	1384
<i>dominus</i> , <i>Protoporus</i> .....	233	<i>Dory</i> .....	1021
<i>Doncella</i> .....	1590, 1595	<i>Doryichthys</i> .....	773
<i>Doncella</i> .....	1587	<i>aculeatus</i> .....	773
<i>Dorada</i> , <i>Mojarra</i> .....	928	<i>californiensis</i> .....	774
<i>Dorado</i> .....	952	<i>lineatus</i> .....	773
<i>Cocinero</i> .....	921	<i>Doryrhamphus</i> .....	773
<i>dorado</i> , <i>Coryphæna</i> .....	353	<i>californiensis</i> .....	773
<i>Doratonotus</i> .....	1611	<i>lineatus</i> .....	773
<i>megalepis</i> .....	1611	<i>Dough-belly</i> .....	205
<i>thalassinus</i> .....	1612	<i>Dourado</i> .....	952
<i>Dories</i> , <i>John</i> .....	1059	<i>Dovetail Fish</i> .....	1563
<i>Dormeur</i> .....	1235	<i>dovii</i> , <i>Anableps</i> .....	683
<i>Dormitator</i> .....	2195	<i>Anisotremus</i> .....	1318
<i>gundlachi</i> .....	2198	<i>Aplocheilus</i> .....	2828, 2830
<i>latifrons</i> .....	2197	<i>Apogon</i> .....	1108
<i>lineatus</i> .....	2198	<i>Fundulus</i> .....	650
<i>maculatus</i> .....	2196, 2198	<i>Gymnothorax</i> .....	397
<i>microphthalmus</i> .....	2198	<i>Haplochilus</i> .....	650
<i>dormitator</i> , <i>Philypnus</i> .....	2195	<i>Heros</i> .....	1535
<i>Platycephalus</i> .....	2195	<i>Lycodontis</i> .....	397
<i>dormitatrix</i> , <i>Electris</i> .....	2195	<i>Muraena</i> .....	397
<i>dormitor</i> , <i>Gobiomorus</i> .....	2198	<i>Opisthopterus</i> .....	437
<i>Philypnus</i> .....	2194	<i>Poecilia</i> .....	695
<i>Dorosoma</i> .....	415	<i>Pomadasis</i> .....	1318
<i>cepedianum</i> .....	416	<i>Pristigaster</i> .....	437
<i>exile</i> .....	416	<i>Pristipoma</i> .....	1318
<i>insociabilis</i> .....	416	<i>Sidera</i> .....	397
<i>mexicanum</i> .....	416	<i>dowi</i> , <i>Arius</i> .....	125
<i>notata</i> .....	416	<i>Dæctor</i> .....	2325
<i>petenense</i> .....	417, 2809	<i>Diapterus</i> .....	1368
<i>Dorosomida</i> .....	415	<i>Eucinostomus</i> .....	1367
<i>dorsale</i> , <i>Hemulon</i> .....	1303	<i>Exocoetus</i> .....	735
<i>dorsalis</i> <i>azurissimus</i> , <i>Microspatho-</i>		<i>Gerres</i> .....	1368
<i>don</i> .....	1570	<i>Leptarius</i> .....	125
<i>Carangoides</i> .....	930	<i>Selenaspis</i> .....	125, 2761
<i>Caranx</i> .....	930	<i>Tachisurus</i> .....	125
<i>setipinnis</i> .....	934	<i>Thalassophryne</i> .....	2326
<i>cinereus</i> , <i>Microspathodon</i> .....	1570	<i>Daydixodon</i> .....	1382

Page.		Page.		Page.
930	Doydixodon fasciatum	1383, 1384	duquesni, Catostomus	193
30	freminvillei	1382, 1384	Placopharynx	198
902	Drachinus trichodon	2297	durvillii, Argyropelecus	604
262	Dragonets	2184	duryi, Etheostoma	2853
1570	Drepanopsetta	2614	Dusky Shark	35
2585	plutessoides	2615	dussumieri, Brama	060
1568, 1570	Drum	1482	Seriola	900
30	Fresh-water	1484	Dussumieria stoffera	419
1570	Drummer, Ground	1436	Dussumieriinae	417
222	Jewsharp	1473	dux, Lachnolaimus	1580
902	Mongolar	1406	dvinensis, Platassa	2650
1469	White-mouth	1462	dybowski, Centronotus	2431
934	drummond-hayi, Epinephelus	1159	Pholidapus	2430
10	drummondii, Otolithus	1409	Eagle Rays	87, 89
1833	Drums, Black	1454	earlli, Phycis	2555
1382	Red	1453	Urophycis	2554
1557	River	1483	Easter Mackerel	866
1384	Sea	1482	Eastern Carp Sucker	168
1021	Drunken-fish	1722	Mud Minnow	624
773	dubia, Netuma	126, 2765	Eater, Crab	948
773	Seriola	905	Écaille, Grande	400
774	dubium, Campostoma	206	Echelus	353
773	Exoglossum	206	candilimbatus	355
773	dubius, Anuodytes	832	ciuciara	356
773	Arius	127	Echeneididm	2265
773	Bodianus	1146	Echeneis	2268, 2271
205	Elerasfer	2496	albescens	2272
952	Menephorus	1147	albicauda	2269
1563	Serranus	1147	apicalis	2268
685	Tachisurus	127	australis	2269, 2271
1318	Dublin Pound Trout	507	brachyptera	2272
2828, 2830	Duck-bill Cat	101	fusca	2270
1108	ductor, Gasterosteus	900	gnaiaean	2270
650	Naucrates	900	holbrooki	2270
397	dugesii, Adinia	661	jacobaea	2272
650	Algansea	211	lineata	2268, 2270
1535	Ameiurus	138	lunata	2269
397	Fundulus	661	metallica	2270
397	Dulee, Boca	29	naucrateoides	2270
437	duleis, Argyreus	307	naucrates	2269
695	Rhynchichthys	307	naucrates	2269
1318	cataracte	306	newhoffi	2272
437	Dules	1217	osteochlr	2273
1318	auriga	1220	pallida	2272
397	dispilurus	1219	postica	2272
125	flaviventris	1221	quatuordecimlamellatus	2272
2325	subllgarius	1218	romora	2272
1368	dumerilii, Aplonichthys	2703	romoroides	2272
1367	Caranx	904	scutata	2271
735	Paralichthrus	1478	sexdecimlamellata	2272
1368	Polycirrus	1470	sphyranarum	2268
125	Seriola	903, 904	squallipeta	2272
125, 2761	Squatina	59	tetraputorum	2273
125	duodecim, Engraulis	446	trepla	2268
2326	duplex, Orthopristis	1339	verticalis	2270
1382	duquesnel, Ptychostomus	103		

	Page.		Page.
<i>Echenels vittata</i> .....	2269	<i>egmontis</i> , <i>Abhia</i> .....	370
<i>Echidna</i> .....	402	<i>Myrophis</i> .....	371
<i>catenata</i> .....	403	<i>egregia</i> , <i>Tigoma</i> .....	237
<i>flavoscripta</i> .....	403	<i>egregius</i> , <i>Leuciscus</i> .....	237
<i>fuscumaculata</i> .....	403	<i>Squalius</i> .....	237
<i>nocturna</i> .....	402	<i>egretta</i> , <i>Bellator</i> .....	2174
<i>echinatum</i> , <i>Lelodon</i> .....	57	<i>Prionotus</i> .....	2175
<i>echinatus</i> , <i>Orbis</i> .....	1745	<i>eigenmanni</i> , <i>Evara</i> .....	304
<i>Echinorhinda</i> .....	57	<i>Gobiesox</i> .....	2339
<i>Echinorhinus</i> .....	57	<i>Gobius</i> .....	2218
<i>obeus</i> .....	58	<i>Rimicola</i> .....	2339
<i>spinus</i> .....	58	<i>Sebastodes</i> .....	1789
<i>echinus</i> , <i>Diodon</i> .....	1746	<i>Eigenmannia</i> .....	341
<i>Echiodon</i> .....	2405	<i>humboldti</i> .....	341
<i>Echiopsis</i> .....	386	<i>eiseni</i> , <i>Characodon</i> .....	2831
<i>Echiostoma</i> .....	589	<i>ekala</i> , <i>Caranx</i> .....	921
<i>barbatum</i> .....	589	<i>ekstromi</i> , <i>Liparis</i> .....	2108
<i>margarita</i> .....	589	<i>elaborata</i> , <i>Murana</i> .....	389
<i>eclancheri</i> , <i>Cossyphus</i> .....	1583	<i>elaboratus</i> , <i>Gymnothorax</i> .....	389
<i>Harpe</i> .....	1583	<i>Lycodontis</i> .....	389
<i>ectenes</i> , <i>Careproctus</i> .....	2136	<i>Elacate</i> .....	948
<i>Micropogon</i> .....	1463	<i>atlantica</i> .....	948
<i>ectenurus</i> , <i>Chloroscombrus</i> .....	2847	<i>blvittata</i> .....	948
<i>edentula</i> , <i>Platirostra</i> .....	102	<i>canada</i> .....	948
<i>edentulus</i> , <i>Cetengraulis</i> .....	450	<i>falcipinnis</i> .....	948
<i>Engraulis</i> .....	450	<i>malabarica</i> .....	948
<i>edwardi</i> , <i>Sciæna</i> .....	1490	<i>motta</i> .....	948
<i>Stilbiscus</i> .....	363	<i>nigra</i> .....	948
<i>Eel</i> , <i>American</i> .....	348	<i>pondiceriana</i> .....	948
<i>Conger</i> of <i>California</i> .....	395	<i>Elagatis</i> .....	906
<i>Fresh-water</i> .....	348	<i>bipinnulatus</i> .....	906
<i>Lamprey</i> .....	10	<i>pinnulatus</i> .....	907
<i>Sand</i> .....	833	<i>Elanura</i> .....	1930
<i>Snipe</i> .....	369	<i>forcata</i> .....	1930
<i>Eel Cat</i> .....	2788	<i>Elaphocottus</i> .....	2006
<i>Eel-back Flounder</i> .....	2649, 2650	<i>pistilliger</i> .....	2008
<i>Eel-pout</i> .....	2453, 2455, 2456, 2457	<i>Elapsopsis</i> .....	381
<i>Eels</i> .....	344, 346, 347	<i>elasoohir</i> , <i>Noturus</i> .....	147
<i>Conger</i> .....	352, 354	<i>elassodon</i> , <i>Hippoglossoides</i> .....	2615
<i>Cusk</i> .....	2481, 2487	<i>Elassoma</i> .....	982
<i>Long-necked</i> .....	343	<i>evergladei</i> .....	982
<i>Ooze</i> .....	349	<i>zonatum</i> .....	982, 2851
<i>Snake</i> .....	372	<i>Elassomidae</i> .....	981
<i>Snipe</i> .....	366	<i>Elastoma</i> .....	1281
<i>Snub-nosed</i> .....	348	<i>macrophthalmus</i> .....	1281
<i>Spiny</i> .....	612	<i>elater</i> , <i>Malthe</i> .....	2739
<i>Symbranchoid</i> .....	342	<i>Ogcocephalus</i> .....	2739
<i>True</i> .....	340	<i>Zallcutes</i> .....	2738
<i>Worm</i> .....	370	<i>Elattarchus</i> .....	1431
<i>eeltenke</i> , <i>Myliobatis</i> .....	88	<i>archidium</i> .....	1431
<i>effulgens</i> , <i>Æthoprora</i> .....	566	<i>Elattonistius</i> .....	412
<i>Arlina</i> .....	1058	<i>elattura</i> , <i>Netuna</i> .....	128, 2769
<i>Boleosoma nigrum</i> .....	1058	<i>elatturus</i> , <i>Arius</i> .....	128
<i>Larimus</i> .....	1421	<i>Electric Rays</i> .....	76
<i>eglanteria</i> , <i>Raja</i> .....	68, 71	<i>Star-gazers</i> .....	2306
<i>Raja</i> .....	71	<i>electricus</i> , <i>Rhinobatus</i> .....	63

Page.		Page.		Page.
370	<i>Electris dormitatrix</i> .....	2195	<i>ellipsoidea, Lebias</i> .....	672
371	<i>elegans, Blakea</i> .....	2353	<i>ellipticus, Chatoessus</i> .....	416
237	<i>Cottus</i> .....	1939	<i>Cymatogaster</i> .....	1503
237	<i>Cyprinodon</i> .....	675	<i>Platophrys</i> .....	2065
237	<i>Diabasis</i> .....	1304	<i>Pleuronectes</i> .....	2665
2174	<i>Etheostoma</i> .....	1074	<i>Rhomboidichthys</i> .....	2065
2175	<i>Gasterosteus</i> .....	748	<i>Ellwife</i> .....	426
304	<i>Gibbonsia</i> .....	2353, 2860	<i>elongata, Aporistia</i> .....	2707
2339	<i>Gila</i> .....	226	<i>Clupea</i> .....	421
2218	<i>Hemulon</i> .....	1304	<i>Cyclothone</i> .....	583
2339	<i>Kyphosus</i> .....	1387	<i>Platessa</i> .....	2657
1789	<i>Labeo</i> .....	186	<i>Poecilia</i> .....	697
341	<i>Leuciscus</i> .....	227	<i>Umbrina</i> .....	1476
341	<i>Mesoprion</i> .....	1278	<i>elongatum, Gonostoma</i> .....	583
2831	<i>Myxodes</i> .....	2353	<i>elongatus, Avocettina</i> .....	2802
921	<i>Nanostoma</i> .....	1075	<i>Benthodesmus</i> .....	888
2108	<i>Orthrorogiscus</i> .....	1754	<i>Catostomus</i> .....	169
389	<i>Pimalepterus</i> .....	1387	<i>Cephalus</i> .....	1756
389	<i>Rhomboplites</i> .....	1278	<i>Clinostomus</i> .....	240
38	<i>Sebastes</i> .....	1830	<i>Cycleptus</i> .....	168
948	<i>Sebastodes</i> .....	1830	<i>Labichthys</i> .....	369
918	<i>Eleginus</i> .....	2537	<i>Leuciscus</i> .....	240
948	<i>navaga</i> .....	2537	<i>Luxilus</i> .....	240
948	<i>Electridinae</i> .....	2188	<i>Megalops</i> .....	409
948	<i>Electris</i> .....	2190	<i>Menticirrhus</i> .....	1476
948	<i>abacurus</i> .....	2200	<i>Ophiodon</i> .....	1875
948	<i>aquidens</i> .....	2202	<i>Osmerus</i> .....	525
948	<i>amblyopsis</i> .....	2199, 2200	<i>Pleuronectes</i> .....	2657
948	<i>belizianus</i> .....	2201	<i>Pomadasis</i> .....	1328
906	<i>capito plagioplateo</i> .....	2201	<i>Pomotis</i> .....	1001
906	<i>grandisquama</i> .....	2198	<i>Sclerognathus</i> .....	169
907	<i>guavina</i> .....	2199	<i>Scopelus</i> .....	555
1930	<i>gyrinus</i> .....	2201	<i>Sebastes</i> .....	1816
1930	<i>lateralis</i> .....	2195	<i>Sebastodes</i> .....	1815
2006	<i>latifrons</i> .....	2198	<i>Squalius</i> .....	240
2008	<i>longiceps</i> .....	2195	<i>Symphurus</i> .....	2707
381	<i>mauricii</i> .....	950	<i>Zoarces</i> .....	2457
147	<i>mugiloides</i> .....	2198	<i>Eloplide</i> .....	408
2615	<i>omocyaneus</i> .....	2198	<i>Elopine</i> .....	408
982	<i>perniger</i> .....	2201	<i>Elops</i> .....	409
982	<i>pictus</i> .....	2201	<i>capensis</i> .....	410
982, 2851	<i>pisonis</i> .....	2200, 2201	<i>inermis</i> .....	410
981	<i>semiluda</i> .....	2204	<i>purpurascens</i> .....	410
1281	<i>sima</i> .....	2198	<i>saurus</i> .....	410, 2806
1281	<i>smaragdus</i> .....	2204	<i>elucens, Siphostoma</i> .....	768
2739	<i>somnolentus</i> .....	2198	<i>Syngnathus</i> .....	768
2739	<i>Elephant Fish</i> .....	95	<i>El Verde</i> .....	817
2738	<i>Fishes</i> .....	94	<i>emarginatum, Scarus</i> .....	1641
1431	<i>Shark</i> .....	51	<i>Sparisoma</i> .....	1641
1431	<i>elephas, Squalus</i> .....	51	<i>emarginatus, Lobotes</i> .....	1257
412	<i>Eleutheractis</i> .....	1220	<i>Serranus</i> .....	1181
128, 2769	<i>coriaceus</i> .....	1233	<i>Embassichthys</i> .....	2655
128	<i>eletherus, Noturus</i> .....	148, 149	<i>bathyblus</i> .....	2655
76	<i>Schilbeodes</i> .....	148	<i>Embiotoca</i> .....	1504
2306	<i>Ellips</i> .....	133	<i>argyrosona</i> .....	1510
63	<i>elliotti, Diabolichthys</i> .....	93	<i>caryi</i> .....	1509

	Page.		Page.
<i>Embiotoca cassidyl</i> .....	1505	<i>Enchelycore</i> .....	389
<i>jacksoni</i> .....	1504, 1505	<i>curyrhina</i> .....	390
<i>lateralis</i> .....	1506	<i>nigricans</i> .....	389
<i>lineata</i> .....	1506	<i>Enchelyopus</i> .....	989, 2456, 2540, 2560
<i>ornata</i> .....	1506	<i>americanus</i> .....	2457, 2555
<i>perspicabilis</i> .....	1506	<i>barbatus</i> .....	2560
<i>webbi</i> .....	1505	<i>brosmie</i> .....	2561
<i>Embiotocidæ</i> .....	1493	<i>cimbricus</i> .....	2561
<i>Embiotocine</i> .....	1494	<i>cimbricus</i> .....	2560
<i>Emblemaria</i> .....	2401	<i>regalis</i> .....	2553
<i>atlantica</i> .....	2402	<i>Enchrasicholus</i> .....	448
<i>nivipes</i> .....	2402	<i>enchrysurus</i> , <i>Chromis</i> .....	1548
<i>oculocirris</i> .....	2403	<i>Endormi Emerande</i> .....	2230
<i>Emblemariinae</i> .....	2347	<i>Enodrias</i> .....	2411
<i>emblematicus</i> , <i>Gobius</i> .....	2247	<i>nebulosus</i> .....	2414
<i>Lepidogobius</i> .....	2247	<i>Engraulididæ</i> .....	439
<i>Scarus</i> .....	1654	<i>engraulinus</i> , <i>Photogenis leucops</i> ...	296
<i>Zalypnus</i> .....	2247	<i>Engraulis</i> .....	448
<i>embryum</i> , <i>Blennicot-</i>		<i>argyrophanus</i> .....	445
<i>tus</i> ... 2016, 2864		<i>atherinoides</i> .....	451
<i>Oligocot-</i>		<i>brevis</i> .....	450
<i>tus</i> .....	2017	<i>brownii</i> .....	443
<i>Embryx</i> .....	2458	<i>chærostomus</i> .....	444
<i>crassilabris</i> .....	2458	<i>clupeoides</i> .....	447
<i>crotalinus</i> .....	2458	<i>compressus</i> .....	447
<i>embryx</i> , <i>Gerres</i> .....	1379	<i>cubanus</i> .....	442
<i>Emerald Fish</i> .....	2229	<i>delicatissimus</i> .....	444
<i>Emerande</i> , <i>Endormi</i> .....	2230	<i>dentex</i> .....	451
<i>Emichthys megalops</i> .....	1502	<i>duodecim</i> .....	446
<i>emilia</i> , <i>Opsopneodus</i> .....	248	<i>edentulus</i> .....	450
<i>Emmeekia</i> .....	1601	<i>grossidens</i> .....	451
<i>venusta</i> .....	1602	<i>janeiro</i> .....	451
<i>emmelane</i> , <i>A verruncus</i> .....	2069	<i>lemniscatus</i> .....	443
<i>Tachysurus</i> .....	2785	<i>louisiana</i> .....	446
<i>Emmelas</i> .....	1765, 1773, 1777	<i>macrolepidotus</i> .....	449
<i>Lepophidium</i> .....	2483	<i>mitchilli</i> .....	446
<i>Emmellethyinæ</i> .....	1364	<i>mordax</i> .....	448
<i>Emmelichthys</i> .....	1365	<i>mysticetus</i> .....	450
<i>vittatus</i> .....	1365, 1366	<i>nanus</i> .....	449
<i>Emmion</i> .....	2375	<i>panameusis</i> .....	448
<i>bristolæ</i> .....	2375	<i>per fasciatus</i> .....	442
<i>Emmine</i> .....	2345	<i>pluquinga</i> .....	443
<i>emoryi</i> , <i>Gila</i> .....	227	<i>poeyi</i> .....	445
<i>Leuciscus</i> .....	226	<i>productus</i> .....	447
<i>Emperador</i> .....	894	<i>spinifer</i> .....	448
<i>Empetrichthys</i> .....	666	<i>surinamensis</i> .....	447
<i>merriami</i> .....	667	<i>tricolor</i> .....	443
<i>Emplycus</i> .....	2552, 2554	<i>engymen</i> , <i>Cetengraulis</i> .....	2815
<i>omphysetus</i> , <i>Arius</i> .....	122	<i>Engyophrys</i> .....	2668
<i>Bagrus</i> .....	122	<i>sancti-laurentii</i> .....	2668
<i>Sciadeichthys</i> .....	122, 2759	<i>enigmaticus</i> , <i>Schedophilus</i> .....	372
<i>Tachysurus</i> .....	122	<i>Enjambre</i> .....	1141
<i>Enantioliparis</i> .....	2114	<i>Enjambres</i> .....	1140
<i>encæromus</i> , <i>Gobius</i> .....	2223	<i>Enneacanthus</i> .....	992
<i>Encheilopus</i> .....	2457	<i>eriarchus</i> .....	994
<i>Enchelycephali</i> .....	345, 346	<i>gloriosus</i> .....	993

Page.	Page.	Page.
389	Enneacanthus margarotis .....	994
390	obesus .....	993
389	pinniger .....	994
2540, 2560	simulans .....	994
2457, 2555	Enneacentrus .....	1143
2500	fulvus .....	1145
2561	ortalibi .....	1146
2561	guttatus .....	1142
2560	coronatus .....	1142
2553	panamensis .....	1141
448	punctatus .....	1146
1548	taeniops .....	1144
2230	enneagrammus, Ernogrammus .....	2441
2411	Stichæus .....	2441
2414	Enneanectes .....	2349
439	cardinalis .....	2350, 2868
296	Enneistius .....	1143, 1147
448	Enneichthys .....	1501
415	Enophrys .....	1937, 1938
451	bison .....	1938
450	claviger .....	1938
443	diceraus .....	1941
444	Enseigne, Porto .....	1687
447	ensenade, Rhinoptera .....	91
447	ensifera, Bairdiella .....	1434
442	Schena .....	1435
444	ensiferus, Centropomus .....	1125
451	ensiformis, Trichiurus .....	887
446	ensis, Gaidropsarus .....	2558
450	Motella .....	2559
451	Onos .....	2559
451	Sphyræna .....	824
443	Entemedor .....	2752
446	entemedor, Narcine .....	2752
449	Entomacrodus .....	2397
446	ebiosictus .....	2398
448	decoratus .....	2399
450	margaritaceus .....	2398
449	nigricans .....	2399
448	entomelas, Sebastichthys .....	1786
412	Sebastodes .....	1785
413	Entosphenus .....	11
445	cristaticus .....	2745
447	epihexodon .....	12
448	tridentatus .....	12
447	Entoxychirus .....	53
443	Enxaréu .....	926
2815	Enypnius .....	2231, 2233
2668	Eopsetta .....	2613
2668	jordani .....	2613
372	Arbaclosa .....	2343
1141	Boleichthys .....	1102
1140	Chrosomus .....	210
992	erythrogaster .....	210
994	Goblesox .....	2343
993	Orthonops .....	2262
	eos, Pcecilichthys .....	1102
	Pronotogrammus .....	1225
	Sebastodes .....	1810
	Eosebastes .....	1765, 1775, 1793
	Eporlannus .....	522
	Ephippide .....	1666
	Ephippinæ .....	1667
	ephippium, Plectropoma .....	1192
	Ephippus faber .....	1668
	gigas .....	1668
	zonatus .....	1669
	Epicopus .....	2529
	gayi .....	2530
	epicurorum, Chromis .....	947
	Epigonichthys .....	4
	Epigonus .....	1111
	occidentalis .....	1112
	epihexodon, Entosphenus .....	12
	Lampetra .....	12
	Epinephelina .....	1128
	Epinephelus .....	1148, 1152, 2853
	adscensionis .....	1152, 1154
	ater .....	1165
	analogus .....	1152
	apus .....	1159
	ascensionis .....	1154
	asperus .....	1154
	atlanticus .....	1154
	bonaci .....	1175
	brachysomus .....	1154
	callurus .....	1186
	catus .....	1159
	chalinus .....	1181
	ciliatus .....	1784
	cubanus .....	1158
	dermatolepis .....	1169
	dimidiatus .....	1179
	drummond-hayi .....	1159
	falcatus .....	1185
	flavolimbatus .....	1155
	galeus .....	1164
	gigas .....	1154
	guaza .....	1154
	guttatus .....	1142, 1159
	inermis .....	1168
	interstitialis .....	1179
	jordani .....	1177
	labrifomis .....	1155
	lunulatus .....	1159
	maculosus .....	1158
	merus .....	1162
	microlepis .....	1178
	morio .....	1160
	multiguttatus .....	1166
	mystacinus .....	1151
	nigritus .....	1162



	Page.		Page.
<i>Epinochelus niphobles</i> .....	2853	<i>Eriocaula</i> .....	2816
<i>niveatus</i> .....	1156	<i>salmonca</i> .....	2816
<i>olfax</i> .....	1183	<i>Erioceta</i> .....	999
<i>ordinatus</i> .....	1155	<i>Eriocosma</i> .....	1028, 1030, 1036
<i>pammensis</i> .....	1141	<i>Erieymba</i> .....	302
<i>pardalis</i> .....	1183	<i>buccata</i> .....	302
<i>punctatus</i> .....	1146, 1154	<i>erleymba, Schla-na</i> .....	1445
<i>quingefasciatus</i> .....	1164	<i>Eriepilinae</i> .....	1831
<i>rosaceus</i> .....	1184	<i>Erileps</i> .....	1862
<i>ruber</i> .....	1181	<i>zonifer</i> .....	1863
<i>sellicanda</i> .....	1155	<i>Erimystax</i> .....	314, 315
<i>striatus</i> .....	1157, 1208	<i>Erimyzon</i> .....	184
<i>taenlops</i> .....	1144	<i>goodii</i> .....	186
<i>tigris</i> .....	1187	<i>sucotta</i> .....	185, 186
<i>venucosus</i> .....	1172	<i>oblongus</i> .....	186
<i>xenarchus</i> .....	1180	<i>erinacea, Rain</i> .....	68
<i>Epimula</i> .....	880	<i>erinacens, Trichocyclus</i> .....	1741
<i>magistralis</i> .....	880	<i>Eritrema</i> .....	308
<i>episcopa, Dianda</i> .....	215	<i>Erizo</i> .....	1745
<i>Hybognathus</i> .....	215	<i>Guambana</i> .....	1746
<i>episcopi, Gumbusia</i> .....	685	<i>Ernogrammus euneogrammus</i> .....	2411
<i>episcopus, Hybognathus</i> .....	215	<i>erochrons, Hololepis</i> .....	1102
<i>Episoma</i> .....	254	<i>Pecilichthys</i> .....	1102
<i>callisema</i> .....	273	<i>Erogala</i> .....	251
<i>jejuna</i> .....	290	<i>Erotelis</i> .....	2203
<i>Epitrichys</i> .....	1023	<i>smaragdus</i> .....	2204
<i>epaetus, Esox</i> .....	443	<i>valenciennesi</i> .....	2204
<i>epatorialis, Cllopsis</i> .....	364	<i>Erychthys</i> .....	1642
<i>Raja</i> .....	74	<i>croicensis</i> .....	1651
<i>Eques</i> .....	1485, 1489	<i>erythreus</i> .....	1531
<i>acuminatus</i> .....	1487	<i>Heros</i> .....	1531
<i>umbrosus</i> .....	1487	<i>Erythrichthys</i> .....	1365
<i>americanus</i> .....	1490	<i>vittatus</i> .....	1366
<i>baltentus</i> .....	1490	<i>Erythrinidae</i> .....	330
<i>lanceolatus</i> .....	1489, 1490	<i>erythrinoides, Scarus</i> .....	1635
<i>lineatus</i> .....	1487	<i>erythrocheilus Albulu</i> .....	412
<i>pulcher</i> .....	1489	<i>erythrogaster, Chrosomus</i> .....	209
<i>punctatus</i> .....	1488, 1489	<i>cos</i> .....	210
<i>viola</i> .....	1486	<i>Leuciscus</i> .....	210
<i>equestris, Arius</i> .....	128	<i>Serranus</i> .....	1160
<i>Balistes</i> .....	1703	<i>erythrogastrum, Pencilosoma</i> .....	1089
<i>Equietus</i> .....	1485	<i>erythrops, Gobiesox</i> .....	2336
<i>equinoculus, Migil</i> .....	2841	<i>Ichthels</i> .....	990
<i>equirostrum, Scombrosox</i> .....	726	<i>erythroptera, Pimelodus</i> .....	135
<i>equisetis, Coryphaena</i> .....	953	<i>erythrorhynchus, Salmo</i> .....	508
<i>Equitina</i> .....	1397	<i>erythrinus, Caranx</i> .....	920
<i>erute, Lobotes</i> .....	1236	<i>Catostomus</i> .....	193
<i>erebennus, Ameiurus</i> .....	139	<i>Ptychostomus</i> .....	193
<i>erebus, Murena</i> .....	396	<i>esca, Clupea</i> .....	421
<i>erethizon, Arothron</i> .....	1739	<i>escambia, Zygonectes</i> .....	658
<i>Ovoides</i> .....	1739	<i>escamuda, Sardina</i> .....	431
<i>Tetradon</i> .....	1739	<i>eschrichtii, Oneirodes</i> .....	2732
<i>eriartha, Atherinella</i> .....	803	<i>Escolar</i> .....	879, 2843
<i>Copelandia</i> .....	904	<i>Chino</i> .....	1114, 1283
<i>Eurystole</i> .....	803	<i>de Natura</i> .....	976
<i>eriarthus, Euneacanthus</i> .....	904		

Page.		Page.		Page.
2816	Escolor violaceus.....	4843	Esox saurus.....	725
2816	Escolares.....	879	scomberius.....	626
999	Escolars.....	877	spel.....	826
030, 1036	Escribano.....	720, 722	aphyrana.....	826
302	esulentus, Curangus.....	921	stomias.....	585
302	Couger.....	355	synodus.....	536
1445	Merluccius.....	2530	timucus.....	711
1841	Esox sarum.....	2840	tridecemlineatus.....	628
1862	bartoni.....	2840	tristechus.....	111
1863	jordani.....	2840	umbrosus.....	627
314, 315	Esiocopus.....	2300, 2303	vern. leulatus.....	627
184	esmarkii, Lycodes.....	2463	viridis.....	110
186	Esmeralda.....	2227, 2230	vittatus.....	628
185, 186	de Mar.....	2203, 2204	vulpes.....	411
186	Negra.....	2204	zonatus.....	639
68	Esoctide.....	708	Espuda.....	894
1741	esopus, Labco.....	186	Pez de.....	2749
311	Esox.....	625	Espadon.....	894
308	affinis.....	628	Espagnol, Quatilibi.....	1140
1745	americanus.....	620	Espino, Puerco.....	1745
1746	atromaculata.....	629	estor, Chirostoma.....	702
241	australis.....	628	Esox.....	628
1162	barracuth.....	823, 2841	Gila.....	240
1162	belone.....	714	Lethostole.....	792
251	boreus.....	628	Leuciscus.....	240
2203	brasiliensis.....	723	Squalius.....	240
2204	crassus.....	627	Estrella.....	1054
2204	cypho.....	627	Eteline.....	1243
1642	deprandus.....	628	Etelis.....	1281
1651	epsetus.....	443	aquilonaris.....	1283
1531	estor.....	628	carbonculus.....	1283
1531	fasciatus.....	626	oculatus.....	1282
1365	flavulus.....	639	Etheostoma.....	1028, 1066, 1069, 1097
1366	immaculatus.....	630	alabama.....	1095
330	imperialis.....	717	artesian.....	1094
1635	lineatus.....	627	asprellus.....	1001
412	longirostris.....	714	atromaculata.....	1057
209	lucioides.....	628	aurantiacum.....	1041
210	lucius.....	628	australe.....	1081
210	lugubrosus.....	628	biennioides.....	1033, 1053
1160	marinus.....	714	biennius.....	1072, 1073
1089	masquinongy.....	629	boreale.....	1082
2336	immaculatus.....	630	ceruleum.....	1080, 2853
990	niger.....	626	spectabile.....	1089
135	nobilior.....	629	callura.....	1011
508	obtensis.....	630	camurum.....	1076
920	ornatus.....	626	caprodes.....	1027
193	osseus.....	110	cinerea.....	1078
193	ovinus.....	672	cinereum.....	1078
421	phaleratus.....	628	cragini.....	1091
658	pisciculus.....	641	cymatolania.....	1042
431	pisculentus.....	641	davisoni.....	1049
2732	porosus.....	627	duryi.....	2853
879, 2843	raveneli.....	626	elegans.....	1074
1114, 1281	reticulatus.....	628	evides.....	1037
976	salmonus.....	538, 627, 620	exile.....	1103

	Page.		Page.
<i>Etheostoma flabellare</i> .....	1007	<i>Etheostoma schumardi</i> .....	1047
<i>cumberland-</i>		<i>sclerum</i> .....	1038
<i>ieum</i> .....	1098	<i>scovellii</i> .....	1082
<i>lineolatum</i> .....	1098	<i>squamatus</i> .....	1040
<i>flabellaria</i> .....	1097	<i>squamiceps</i> .....	1096
<i>flabellata</i> .....	1007	<i>stigmaeum</i> .....	1048
<i>fonticola</i> .....	1105	<i>swannanoa</i> .....	1070
<i>fontinulis</i> .....	1097	<i>tessellatum</i> .....	1078
<i>formosa</i> .....	2853	<i>thalassinum</i> .....	1071
<i>fusiforme</i> .....	1103	<i>tippecanoe</i> .....	1090
<i>guntheri</i> .....	1034	<i>tucumbia</i> .....	1100
<i>histris</i> .....	1051	<i>urumidea</i> .....	1045
<i>inscriptum</i> .....	1072	<i>variatum</i> .....	1069
<i>ioe</i> .....	1084	<i>verecundum</i> .....	1050
<i>lowe</i> .....	1083	<i>vexillare</i> .....	1058
<i>jessie</i> .....	1084	<i>virgatum</i> .....	1093
<i>jordani</i> .....	1079, 1080	<i>vulneratum</i> .....	1077
<i>julie</i> .....	1093	<i>whipplei alabamæ</i> .....	1095
<i>laterale</i> .....	1099	<i>whippili</i> .....	1095
<i>lepidogenys</i> .....	1087	<i>wrighti</i> .....	1047
<i>lepidum</i> .....	1089	<i>zonale</i> .....	1075
<i>linsleyi</i> .....	1097	<i>arcansanum</i> .....	1075
<i>longimana</i> .....	1054	<i>etheostoma, Aboma</i> .....	2240
<i>luteovinctum</i> .....	1086	<i>Etheostominae</i> .....	1018
<i>lyuceum</i> .....	1075	<i>elton, Syngnathus</i> .....	767
<i>macrocephalum</i> .....	1031	<i>Etmopterus</i> .....	55
<i>maculatum</i> .....	1077	<i>pusillus</i> .....	55
<i>microperca</i> .....	1104	<i>etowana, Catostomus nigricans</i> ..	181
<i>micropterus</i> .....	1083	<i>Etropus</i> .....	2687
<i>nevisense</i> .....	1034	<i>crossotus</i> .....	2689
<i>nianguæ</i> .....	1043	<i>microstomus</i> .....	2687, 2690
<i>splotum</i> .....	1044	<i>rimosus</i> .....	2688
<i>nigrofasciatum</i> .....	1039	<i>Etrumens</i> .....	419
<i>nigrum</i> .....	1057	<i>acuminatus</i> .....	419
<i>notatum</i> .....	1070	<i>sadira</i> .....	420
<i>obeyense</i> .....	1092	<i>teres</i> .....	420
<i>ohmstedti</i> .....	1057	<i>Eucalia</i> .....	743
<i>ouachito</i> .....	1035	<i>inconstans</i> .....	744
<i>pagei</i> .....	1092	<i>cayuga</i> .....	744
<i>parvipinna</i> .....	1096	<i>pygmaea</i> .....	744
<i>pellucidum clarum</i> .....	1063	<i>Eucentrarchus</i> .....	988
<i>peltatum</i> .....	1034	<i>Euchalarodus</i> .....	2649
<i>phoxocephalum</i> .....	1031	<i>putnami</i> .....	2650
<i>podostemone</i> .....	1055	<i>Encinostomus</i> .....	1367
<i>pottsii</i> .....	1082	<i>argenteus</i> .....	1371
<i>preharae</i> .....	1104	<i>californiensis</i> .....	1369
<i>preliaris</i> .....	1104	<i>dowi</i> .....	1367
<i>punctulatum</i> .....	1090	<i>gula</i> .....	1370
<i>quappella</i> .....	1084	<i>gulula</i> .....	1371
<i>quiescens</i> .....	1101	<i>harengulus</i> .....	1368
<i>rex</i> .....	1026	<i>lefroyi</i> .....	1372
<i>roanoka</i> .....	1036	<i>productus</i> .....	1372
<i>rufilineatum</i> .....	1079	<i>pseudogula</i> .....	1368
<i>rufolineatum</i> .....	1079	<i>Euctenogobius</i> .....	2210, 2215, 2226
<i>rupestre</i> .....	1073	<i>badius</i> .....	2227
<i>sagitta</i> .....	1080	<i>latus</i> .....	2237

Page.		Page.		Page.
1047	<i>Euotengobius lyricus</i> .....	2225	European Hake .....	2530
1038	<i>sagittula</i> .....	2229	Lancelets .....	3
1082	<i>Eucyclogobius</i> .....	2248	Porgies .....	1350
1040	<i>newberryi</i> .....	2248	Sculpin .....	1074
1096	<i>endouxii</i> , <i>Allurichthys</i> .....	118	Stickleback .....	747
1048	<i>Fellethys</i> .....	118	<i>Eurymyetera</i> .....	392
1070	<i>Galeichthys</i> .....	118	<i>euryopa</i> , <i>Chlola</i> .....	270
1078	<i>Eugaleus</i> .....	31	<i>Hudsonius</i> .....	270
1071	<i>Eugomphodus</i> .....	46	<i>euryops</i> , <i>Bathylagus</i> .....	529
1090	<i>Ittoralls</i> .....	47	<i>Icelus</i> .....	1915, 1916
1100	<i>Eulachon</i> .....	521	<i>Myxostoma</i> .....	193
1045	<i>Eulamia</i> .....	33	<i>Tylosurus</i> .....	711
1069	<i>lamia</i> .....	38	<i>euryorus</i> , <i>Eupomotis</i> .....	1008
1050	<i>longimana</i> .....	38	<i>Lepomis</i> .....	1000
1058	<i>millberti</i> .....	37	<i>Eurypharyngida</i> .....	406
1093	<i>nicaraguensis</i> .....	30	<i>euryplectrum</i> , <i>Diplectrum</i> .....	1206
1077	<i>platyrhynchus</i> .....	36	<i>euryrhina</i> , <i>Enchelycore</i> .....	390
1095	<i>eulepis</i> , <i>Microgobius</i> .....	2244	<i>Eurystole</i> .....	802
1095	<i>Euleptorhamphus</i> .....	723	<i>eriarcha</i> .....	803
1047	<i>brevoorti</i> .....	724	<i>eurystole</i> , <i>Stolephorus</i> .....	445
1075	<i>longirostris</i> .....	724	<i>eurystoma</i> , <i>Chlola</i> .....	277
1075	<i>velox</i> .....	724	<i>Codoma</i> .....	285
2240	<i>Eumesogrammus</i> .....	2441	<i>Eurystomus</i> .....	173
1018	<i>praecius</i> .....	2441	<i>eurystomus</i> , <i>Notropis</i> .....	277
767	<i>subfluvireatus</i> .....	2440	<i>Photogenis</i> .....	277
55	<i>Eumicrotremus</i> .....	2097	<i>Enacarus</i> .....	1627, 1620, 1639
55	<i>orbis</i> .....	2099, 2100	<i>Eusehistodus</i> .....	1560, 1562
181	<i>spliosus</i> .....	2098, 2099	<i>analogus</i> .....	1503
2687	<i>eumorphus</i> , <i>Chatoessus</i> .....	433	<i>concolor</i> .....	1550
2689	<i>Eupomacentrus</i> .....	1549, 1550, 1551	<i>deellivifrons</i> .....	1562
2687, 2690	<i>adustus</i> .....	1551	<i>Eusebastes</i> .....	1760
2688	<i>annalis</i> .....	1554	<i>Eusphyra</i> .....	43
419	<i>dieneus</i> .....	1552	<i>Eustomatodus</i> .....	907
419	<i>flavilatus</i> .....	1557	<i>Euthynnus</i> .....	808
420	<i>flaviventer</i> .....	1557	<i>aliteratus</i> .....	869
420	<i>fuscus</i> .....	1552	<i>pelamys</i> .....	869
743	<i>leucornis</i> .....	1551	<i>Eutychelithus</i> .....	1483
744	<i>leucostictus</i> .....	1555	<i>evansi</i> , <i>Hybognathus</i> .....	213
744	<i>otophorus</i> .....	1555	<i>Evapristis</i> .....	1334, 1336, 1340
744	<i>partitus</i> .....	1558	<i>Evarra</i> .....	304
988	<i>planifrons</i> .....	1559	<i>eigenmanni</i> .....	304
2649	<i>rectifrenum</i> .....	1553	<i>Eventognathi</i> .....	161
2650	<i>Eupomotis</i> .....	1006, 1007	<i>Evepigymnus</i> .....	907
1367	<i>aureus</i> .....	1010	<i>evergladei</i> , <i>Elassoma</i> .....	982
1371	<i>euryorus</i> .....	1008	<i>evermanni</i> , <i>Atherinella</i> .....	804
1369	<i>gibbosus</i> .....	1009	<i>Cottus</i> .....	1945
1367	<i>heros</i> .....	1007	<i>Searus</i> .....	1651
1370	<i>holbrookii</i> .....	1008	<i>Synodus</i> .....	535
1371	<i>humilis</i> .....	1004	<i>Thyrina</i> .....	804
1368	<i>macrochirus</i> .....	1005	<i>Evermannia</i> .....	2256
1372	<i>pallidus</i> .....	1006	<i>longipinnis</i> .....	2256
1372	<i>europeus</i> , <i>Aspidophorus</i> .....	2067	<i>zosterura</i> .....	2256
1368	<i>Blennius</i> .....	2419	<i>Evertzens</i> , <i>Jacob</i> .....	1143
15, 2226	<i>Trachurus</i> .....	911	<i>evides</i> , <i>Alvordius</i> .....	1037
2227	European <i>Barracuda</i> .....	826	<i>Clinus</i> .....	2353
2237	<i>Charr</i> .....	508	<i>Etheostoma</i> .....	1037

	Page.		Page.
evides, Gibbonsia .....	2352, 2869	Exocoetus fasciatus .....	733
Hudropterus .....	1036	fureatus .....	737
Plectobranchus .....	2432	georgianus .....	730
evionthas, Ophichthus .....	381	gibbifrons .....	741
Quassiremus .....	380	gryllus .....	729
evolans, Exocoetus .....	730	heterurus .....	735
Halocypræus .....	729	hillianus .....	729
Prionotus .....	2167, 2168, 2169	hucellifer .....	733
Trigla .....	2160	lutkeni .....	736
Evoplites .....	1245	maculipinnis .....	737
viridis .....	1246	melanurus .....	735, 736
Evoorthodus .....	2208	mesogaster .....	729
breviceps .....	2208	monocellus .....	730
catulus .....	2218	nigricans .....	737
Evoxymetopon .....	885	oveboracensis .....	735, 736
teniatus .....	886	nuttalli .....	737
exasperata, Platyrrhina .....	65	obtusirostris .....	730
Syrhina .....	65	orbignanus .....	729
exasperatus, Rhuobatus .....	65	parræ .....	740
Zapteryx .....	64	proene .....	737
Exerpes .....	2367	quadrifemis .....	735
asper .....	2367	roberti .....	735
exiguus, Bodiannus .....	1433	robustus .....	736
Stolephorus .....	442	rondeletii .....	733, 734
exilo, Dorosoma cepedianum .....	416	rubescens .....	734
Etheostoma .....	1103	rufipinnis .....	735
exilicauda, Lavinia .....	208	scylla .....	735
Lenciscus .....	209	speculiger .....	734
exsiliens, Exocoetus .....	732, 734	spilonopterus .....	740
exilis, Belone .....	714	spilopus .....	738
Boleichthys .....	1103	splendens .....	730
Hippoglossoides .....	2613	vermiculatus .....	740
Lyopsetta .....	2613	vinciguerra .....	734
Noturus .....	147	volador .....	733
Pæcillethys .....	1103	volitans .....	734, 736, 2835
Schilbeodes .....	147	zenopterus .....	738
Tylosurus .....	714	Exoglossine .....	204
eximius, Cyprinodon .....	673	Exoglossum .....	327
Exocetidae .....	726	annulatum .....	327
Exocoetus .....	730, 731, 732, 734	dubium .....	206
acutus .....	728	lesueurianum .....	327
affinis .....	735, 2836	maxilligua .....	327
albinaetylus .....	739	mirabile .....	303
appendiculatus .....	736	nigrescens .....	327
bahiensis .....	739	spinicephalum .....	206
bicolor .....	738	vittatum .....	327
brachycephalus .....	733	exoletus, Acantholabrus .....	1576
californicus .....	730, 740	Centrolabrus .....	1576
callopterus .....	740	Labrus .....	1576
chilensis .....	730	Exonantes .....	2835
comatus .....	736	affinis .....	2836
cyanopterus .....	739	exsiliens .....	2836
dowi .....	735	rondeletii .....	2836
evolans .....	730, 2835	rufipinnis .....	2836
exsiliens .....	732, 734		

Page.		Page.		Page.
733	<i>Exonantes speculiger</i> .....	2836	<i>Fario stellatus</i> .....	492
737	<i>vinciguerræ</i> .....	2836	<i>tsupptich</i> .....	493
730	<i>expansum, Ostracion</i> .....	1724	<i>farkharil, Lobotes</i> .....	1230
741	<i>exsiliens, Exomantes</i> .....	2836	<i>fasciata, Aphoristia</i> .....	2710
729	<i>extensus, Fundulus</i> .....	646	<i>Cichla</i> .....	1012
735	<i>Lycodapus</i> .....	2494	<i>Clupea</i> .....	426
729	<i>faber, Chætodipterus</i> .....	1668	<i>Molnesia</i> .....	695
733	<i>Chætodon</i> .....	1668	<i>Plagusia</i> .....	2710
739	<i>Ephippus</i> .....	1668	<i>Pocilia</i> .....	641
736	<i>Faber marinus</i> .....	1668	<i>Seriola</i> .....	904
737	<i>fabricii, Campylodon</i> .....	615	<i>Trigla</i> .....	2183
735, 736	<i>Centrosecyllium</i> .....	56	<i>fasciatum, Doydixodon</i> .....	1383, 1384
729	<i>Cottus</i> .....	2009	<i>Pristipoma</i> .....	1330
730	<i>Gadus</i> .....	2534	<i>fasciatus, Aehlrus</i> .....	2700
737	<i>Gunnellus</i> .....	2438	<i>Anchenopterus</i> .....	2373
735, 736	<i>Liparis</i> .....	2121, 2128	<i>Bryttus</i> .....	993
737	<i>Lumpenus</i> .....	2437	<i>Caranx</i> .....	914
730	<i>Macrorus</i> .....	2582	<i>Carapus</i> .....	341
729	<i>Spinax</i> .....	56	<i>Catonotus</i> .....	1098
740	<i>falcata, Agosia</i> .....	313	<i>Catostomus</i> .....	187
737	<i>shuswap</i> .....	313	<i>Centrarchus</i> .....	1012
735	<i>Mycteroperca</i> .....	1184	<i>Centronotus</i> .....	2418
735	<i>phenax</i> .....	1185	<i>Cromnobates</i> .....	2373
736	<i>Seriola</i> .....	905	<i>Ctenogobius</i> .....	2223
733, 734	<i>falcatus, Caranx</i> .....	913, 2845	<i>Diplesion</i> .....	1081
734	<i>Epinephelus</i> .....	1185	<i>Esox</i> .....	626
735	<i>Hemicaranx</i> .....	2845	<i>Exocoetus</i> .....	733
735	<i>Labrus</i> .....	942	<i>Genyonemus</i> .....	1479
734	<i>Lachnolaimus</i> .....	1580	<i>Giton</i> .....	340
740	<i>Serranus</i> .....	1185	<i>Gobiesox</i> .....	2338
738	<i>Sparus</i> .....	1583	<i>Gobius</i> .....	2322
730	<i>Trachinotus</i> .....	941	<i>Gunnellus</i> .....	2418
740	<i>Trisotropis</i> .....	1185	<i>Gymnachirus</i> .....	2703
734	<i>falciformis, Carebarhinus</i> .....	36	<i>Gymnotus</i> .....	340
733	<i>Careharius</i> .....	35	<i>Halatractus</i> .....	904
736, 2835	<i>Platypodon</i> .....	36	<i>Harpurus</i> .....	1691
738	<i>faleipinnis, Elacate</i> .....	948	<i>Hemirhamphus</i> .....	720
204	<i>fallax, Carangus</i> .....	923	<i>Larimus</i> .....	1424
327	<i>Caranx</i> .....	923	<i>Murenoides</i> .....	2418
327	<i>Pontotis</i> .....	1003	<i>Mytilophagus</i> .....	1504
206	<i>Trachurus</i> .....	910	<i>Orthagoriscus</i> .....	1754
327	<i>Fallfish, Red</i> .....	286	<i>Pholis</i> .....	2417
327	<i>Fall-fishes</i> .....	220, 221	<i>Pimephales</i> .....	217
303	<i>Fall Herring</i> .....	425	<i>Pogonias</i> .....	1483
327	<i>Fanejal</i> .....	1837	<i>Prionoles</i> .....	1212
206	<i>fanfanas, Naucrates</i> .....	900	<i>Scomber</i> .....	904
327	<i>Fanguito</i> .....	692	<i>Selastes</i> .....	1761, 1827
1576	<i>Fan-tailed Darter</i> .....	1097	<i>Sicyasus</i> .....	2338
1576	<i>Mullet</i> .....	816	<i>Syngnathus</i> .....	771
1576	<i>Fario</i> .....	483	<i>Synodus</i> .....	536
2835	<i>argyreus</i> .....	480	<i>Tetragonopterus</i> .....	334
2836	<i>aurora</i> .....	499	<i>Trachurus</i> .....	904
2836	<i>clarkii</i> .....	501	<i>Trachynotus</i> .....	941
2836	<i>gairdneri</i> .....	499	<i>fasciulare, Diplectrum</i> .....	1208
2836	<i>Fario newberryi</i> .....	499	<i>fascicularis, Centropristis</i> .....	1208

	Page.		Page.
fascicularis, Hippocampus .....	778	Fierasfer borealis .....	2443
Serranus .....	1208	dubius .....	2496
fasciolaris, Catostomus .....	186	Ærasfer, Lycodapus .....	2493
Notropis umbratilis .....	301	Fierasferidae .....	2494
Sebastichthys .....	1827	filamentosus, Ailurichthys .....	118
Symphurus .....	2707	Argyriosus .....	336
fasciolatus, Coryphaena .....	952	Dentex .....	1289
Fatback .....	433, 946	Felichthys .....	118
Fat-head .....	217, 1585	Hemirhamphus .....	723
Father-lasher .....	1971	Icelinus .....	1893
favosus, Bathygadus .....	2565	Monacanthus .....	1716
Blennius .....	2380	Scomber .....	932
fecundus, Catostomus .....	180	Tarandichthys .....	1862
Felichthys .....	116	Filefish .....	1712, 1715, 1717, 1718
bagre .....	117	Orange .....	1718
bahiensis .....	118	filicornis, Blennius .....	2381
endouxii .....	118	Alifera, Chalinura .....	2577
filamentosus .....	118	fimbria, Anoplopoma .....	1862
marinus .....	118	Gadus .....	1862
panamensis .....	117	fimbriata, Cyclosetta .....	2676
pinnimaculatus .....	117	Kaia .....	93
felicianus, Cyprinodon .....	676	Solea .....	2700
Trifarctus .....	676	Squatina .....	59
felinus, Pimelodus .....	140	fimbriatus, Achirus .....	2700
Serranus .....	1187	Arnoglossus .....	2677
felis, Anarrhichthys .....	2448	Blennius .....	2457
Arius .....	128	Chanmax .....	2726
Hexanematichthys .....	128	Hemirhombus .....	2677
Mustelis .....	31	Icelinus .....	1894
Pmelodus .....	141	Serranus .....	1151
Silurus .....	128	Zoarcus .....	2457
fenestralis, Artedius .....	1900	Fimbriatorpedo .....	77
fenestrata, Chronis .....	1518	Fine-sealed Sucker .....	173, 178
fenestratum, Cichlasoma .....	1518	firmsquamis, Bogoslovius .....	2575
fenestratus, Heros .....	1518	Macrurus .....	2576
ferox, Alepisaurus .....	595	fischeri, Achirus .....	2700
Bathyophis .....	605	Achirus .....	2699
Bathysaurus .....	539	Chaetostomus .....	160
Cynoponticus .....	360	Solea .....	2700
Idiacanthus .....	605	Tetragonopterus .....	334
Lepisosteus .....	111	Fish, Angel .....	58
Stomias .....	588	Bat .....	2737
ferruginea, Limanda .....	2644	Butter .....	2419
Myzopselta .....	2645	Cobbler .....	931
Platessa .....	2645	Common Alligator .....	2061
ferrugineus, Characodon .....	609	Common Buffalo .....	163
Pleuronectes .....	2645	Creek .....	185
fenille, Polyodon .....	102	Creole .....	1221
Fiatolus .....	964	Devil .....	92
fibulatus, Spinicephalus .....	2796	Dismal Swamp .....	703
Fiddler Fish .....	63	Doll .....	1674
fieldii, Stomias .....	586	Dovetail .....	1563
Fierasfer .....	2495	Elephant .....	95
affinis .....	2495	Emerald .....	2229
arenicola .....	2496	Flo .....	1715, 1718
bermudensis .....	2497	Fiddler .....	63

Page.		Page.		Page.
2443	Fish, Fool.....	1715, 1718	Fishes, Mackerel-like.....	860
2496	Glance.....	954	Mail-cheeked.....	1756
2493	Globe.....	1734	Milk.....	414
2494	Good.....	487	Parrott.....	1620
118	Guitar.....	93	Pediculate.....	2712
336	Hand-saw.....	596	Perch-like.....	979
1289	Harvest.....	965	Pike-like.....	622
118	Indian.....	1680	Plectognathous.....	1696
723	Leather.....	1714, 1715	Poreupine.....	1742, 1744
1893	Lion.....	1859	Rag.....	968
1716	Lizard.....	538	Rudder.....	1380
932	Log.....	964	Scorpion.....	1839
1892	Mutton.....	1376	Sergeant.....	947
1717, 1718	Oil.....	879	Spiny-rayed.....	779
1718	Portuguese Man-of-War.....	949	Syngnathous.....	707
2381	Prick.....	555	Trachinoid.....	2273
2577	Priest.....	1784	Truo.....	97
1862	Rabbit.....	882	Trunk.....	1720
1862	Rainwater.....	665	Fishing-Frogs.....	2713
2676	Red-mouth Buffalo.....	161	fissuratus, Neoliparis.....	2113
93	Red Parrot.....	1635	fissus, Arius.....	131
2700	Ribbon.....	1489, 1490	Tachisurus.....	131
59	San Pedro.....	954	Tachysurus.....	131, 2782
2700	Scabbard.....	887, 889	Fistularia.....	756
2677	Scour.....	879	commersonii.....	758
2457	Sergeant.....	948	depressa.....	757
2726	Singing.....	2321	immaculata.....	758
2677	Soldier.....	1083	neoboracensis.....	757
1894	Tongue.....	2710	petimba.....	758
1151	Tyrant.....	886	serrata.....	758
2457	Ugly.....	137	tabacaria.....	757, 758, 2837
77	Unicorn.....	1719	Fistulariidae.....	755
173, 178	Yellow.....	1144	fistularis, Flagellaria.....	757
2575	Fishes.....	14	fistulatum, Siphostoma.....	765
2576	Angel.....	58	fistulatus, Syngnathus.....	765
2700	Atka.....	1864	flabellare, Etheostoma.....	1097
2699	Black Rudder.....	963	cumberlandi- eum.....	1098
160	Blennioid.....	2343	hueolatum.....	1098
2700	Blind.....	702	flabellaris, Etheostoma.....	1097
334	Bony.....	113	flabellata, Etheostoma.....	1097
58	Brotuloid.....	2498	flabellatus, Catenofus.....	1098
2737	Buffalo.....	163	Flag, Spanish.....	1817
2419	Cardinal.....	1105	Flagellaria.....	756
931	Carp-like.....	160	flstularis.....	757
2061	Cirrhitoid.....	1490	flagellum, Ruda.....	88
163	Cutlass.....	888	Succopharynx.....	406
185	Elephant.....	94	Flags, Spanish.....	1130
1221	Fllo.....	1712, 1717	Flumenco.....	1209
92	Four-eyed.....	684	Flammaeo.....	2871
1674	Ganoid.....	100	marianus.....	2871
1563	Guitar.....	61	flammans, Leuciscus.....	242
95	Isospondylous.....	407	Phoxinus.....	242
2229	Jugular.....	2528	Flannel-mouth Cut.....	137
15, 1718	Lanceet.....	593, 594, 595	Sucker.....	174
63	Lantern.....	530, 550	Flusher.....	1235
	Lizard.....	531		



	Page.		Page.
Flatfish .....	1680, 2602	flavus, Turulus .....	1583
Common .....	2647	Flesh-colored Rockfish .....	1824
Flat-headed Chub .....	326	fleurieu, Osterlinchus .....	1107
flava, Congermurena .....	357	flexularis, Lepomis .....	1011
flavescens, Arius .....	123	flexuosus, Cntostomus .....	179
Bagrus .....	123	Flier .....	988
Bodianus .....	1024	Fliona .....	1793
Brosimius .....	2501	florae, Neoliparis .....	2111
Callyodon .....	1640	florealis, Platyglossus .....	1597
Mesoprion .....	1260	Florida Cat .....	137
Morone .....	1024	floridae, Jordanella .....	677
Perea .....	1023	Siphostoma .....	766
fluviatilis .....	1024	floridana, Cichla .....	1012
Prionodes .....	1215	floridanus, Phycis .....	2554
Scarus .....	1640	Urophycis .....	2554
Setadeichtys .....	123, 2760	floridensis, Calliurus .....	992
Sparisoma .....	1639, 1640	Fundulus .....	642
Tachisurus .....	123	floripinnis, Fundulus .....	651
flavesny, Brosimius .....	2501	Haplochilus .....	651
flavicauda, Hyporthodus .....	1156	Zygonectes .....	651
flavidus, Apodichthys .....	2411	Flounder, Arctic .....	2649
Aulorhynchus .....	754	Eel-back .....	2650
Sebastichthys .....	1782	Four-spotted .....	2632
Sebastodes .....	1781	Great .....	2652
flaviguttatum, Haemulon .....	1312	Gulf .....	2631
Lythron .....	1312	Peacock .....	2665
flaviguttatus, Haemulon .....	1312	Pole .....	2657
flavilatus, Eupomacentrus .....	1557	Soft .....	2679
Pomacentrus .....	1558	Southern .....	2630
flavipinnis, Hybognathus .....	215	Starry .....	2651
Hisha .....	435	Summer .....	2629
Pellona .....	436	Winter .....	2646
Pristigaster .....	436	Fluke, Craig .....	2056, 2657
flavirostre, Siphostoma .....	768	fluviatilis, Algoma .....	215
flavirostris, Syngnathus .....	768	Hudsonius .....	269
flavissimus, Forcipiger .....	1671	Hybognathus .....	215
flaviventer, Eupomacentrus .....	1557	Perea flavescens .....	1024
flaviventris, Dules .....	1221	Sargosomus .....	1496
Serranus .....	1221	Fly-fish .....	1809
flavolimbatus, Epinephelus .....	1155	Flying-fish, Great .....	740
flavolmeatum, Haemulon .....	1306	Sharp-nosed .....	728
flavolineatus, Diabasis .....	1306	Flying-fishes .....	726, 730
Diplodus .....	1360	Flying Gurnard .....	2182, 2183
Pimblepterus .....	1386	Robin .....	2183
Sargus .....	1360	Foditor .....	727
flavomarginatus, Pseudoscarnus .....	1652	acutus .....	728
Scarus .....	1652	fodiator, Tylosurus .....	715
flavoscripta, Eclidna .....	403	fotens, Salmo .....	538
flavoscriptus, Gymnothorax .....	395	Saurus .....	538
flavovittatus, Mulloides .....	860	Synodus .....	538
Upeneus .....	860	folium, Polyodon .....	102
flavulus, Esox .....	619	fonsecensis, Achirus .....	2699
flavus, Awaous .....	2235	Solea .....	2699
Chonophorus .....	2235	fonticola, Alvarius .....	1105
Gobius .....	2235	Etheostoma .....	1105
Noturus .....	144	Fundulus .....	643
		Microperca .....	1104

## Index.

2979

Page.		Page.		Page.
1583	fontinalis, Etheostoma .....	1097	fraterculus, Mylocheilus .....	220
1824	Salmo .....	507	fremebundum, Hamulon .....	1297
1107	Salvelinus .....	506	fremebundus, Diabasis .....	1297
1011	agassizii .....	507	freminvillei, Doydixodon .....	1382, 1384
179	Foutinus .....	633, 634, 645	Myliobatis .....	89
988	Foolfish .....	1715, 1718	frenatus, Balistes .....	1705
1793	forbesi, Cliola .....	272	Brachyistius .....	1499
2111	Cyprinella .....	272	Micrometrus .....	1499
1597	Orthopristis .....	1336	Odontopyxis .....	2675
137	foroipatus, Balistes .....	1702	Sarritor .....	2073
677	Forcipiger .....	1671	Zanlolepis .....	1877
766	flavissimus .....	1671	French Grunt .....	1306
1012	forficata, Elanura .....	1930	Mullet .....	813
2554	Gnaperva lata .....	1702	Frère Jacques .....	846
2554	formosa, Algnsea .....	246	Fresh-water Drum .....	1484
902	Cliola .....	271	Eel .....	348
642	Etheostoma .....	2853	fretensis, Cliola .....	261
651	Heterandria .....	687	Hybopsis .....	261
651	Hydrargyra .....	2827	Notropis .....	261
651	Leucos .....	246	Friars .....	789
2649	Mollienisia .....	699	friedrichsthalii, Heros .....	1528
2650	Moniana .....	271	Frigate Mackerels .....	867
2632	Perca .....	1208	frigida, Moniana .....	271
2652	Uranidea .....	1969	trigidus, Notropis .....	271
2631	formosulum, Campostoma .....	206	Lycodes .....	2465
2665	formosum, Diplectrum .....	1207	Friiled Sharks .....	16
2657	Hamulon .....	1305	Frog Fishes .....	2715
2679	formosus, Alburnus .....	280	frondosum, Sparisoma .....	1641, 1642
2630	Anthias .....	1304	frondosus, Searus .....	1636, 1642
2651	Calliurus .....	996	frontalis, Caranx .....	925
2629	Cottus .....	1969	Gastropsetta .....	2636
2646	Girardinus .....	688	Leuciscus .....	283
2656, 2657	Holacanthus .....	1685	pis cornutus .....	283
215	Leuciscus .....	246	carecharhinus .....	39
269	Leucus .....	246	Carebarias .....	39
215	Notropis .....	271	Frostfish .....	2540
1024	Spheroides .....	1736	Frost Fishes .....	886
1496	Tetrodon .....	1737	Fry, Hog-month .....	444
1809	forakali, Glossodus .....	411	fucensis, Liparis .....	2119
740	forsteri, Sphyræna .....	824	Theragra .....	2536
728	forsterianus, Catostomus .....	176	fucorum, Apodichthys .....	2413
726, 730	fosteri, Albula .....	412	Blennius .....	2379
182, 2183	Caranx .....	923	Nererpes .....	2413
2183	Sombrosax .....	726	fulgens, Corvina .....	1435
727	Four-Bearded Rocklings .....	2560	Myriopristis .....	840
728	Four-eyed Fishes .....	684	Priacanthus .....	1238
715	Four-spotted Flounder .....	2632	fulgida, Meda .....	329
538	Fox Shark .....	45	fuliginosus, Balistes .....	1702
538	fragilis, Chtharichthys .....	2680	Chilomycterus .....	1749
102	Francesa, Lisa .....	410	Diodon .....	1749
2699	francisci, Crestacion .....	21	Holeconotus .....	1505
2699	Gyroleurodus .....	20	Symbranchus .....	342
1105	franklini, Cottus .....	1967	fulva, Labrus chogset .....	1577
1105	Plenrometes .....	2650	fulvomaculatum, Pristipoma .....	1339
643	Uranidea .....	1967	fulvomaculatus, Labrus .....	1339
1104	Fraser River Saluon .....	481	fulvum, Ginglymostoma .....	26

	Page.		Page.
<i>Fulvus</i> , <i>Bodianus</i> .....	1144	<i>Fundulus</i> <i>multifasciatus</i> .....	645
<i>punctatus</i> .....	1146	<i>nigrofasciatus</i> .....	641
<i>ruber</i> .....	1145, 1146	<i>notatus</i> .....	659
<i>Enneacentrus</i> .....	1145	<i>nottii</i> .....	656, 2830
<i>outalibi</i> .....	1146	<i>ocellaris</i> .....	642, 2827, 2828
<i>Labrus</i> .....	1145	<i>pachycephalus</i> .....	661
<i>Physiculus</i> .....	2547	<i>pallidus</i> .....	638, 2827
<i>fumeus</i> , <i>Notropis</i> .....	294	<i>parvipinnis</i> .....	640, 2827, 2830
<i>Fonal</i> , <i>Scorfanudi</i> .....	1837	<i>pisicentus</i> .....	641
<i>Funcinita</i> .....	1107	<i>pulvereus</i> .....	652, 2828
<i>Fundulina</i> .....	631	<i>punctatus</i> .....	637, 2827
<i>funduloides</i> , <i>Clinostomus</i> .....	239	<i>rathbuni</i> .....	649
<i>Fundulus</i> .....	650	<i>rhizophore</i> .....	644
<i>Leucisens</i> .....	240	<i>robustus</i> .....	644, 2827
<i>Squalius</i> .....	240	<i>rubrifrons</i> .....	653
<i>Zygonectes</i> .....	650	<i>scartes</i> .....	654
<i>Fundulus</i> .....	632, 633, 637, 2827	<i>scialicens</i> .....	654, 2828
<i>adina</i> .....	645, 2827	<i>seuinolis</i> .....	647, 2828
<i>albolineatus</i> .....	649, 2828	<i>similis</i> .....	638
<i>arlingtonius</i> .....	652	<i>stellifer</i> .....	648, 2828
<i>aureus</i> .....	659	<i>swampina</i> .....	645
<i>bermude</i> .....	643, 2827	<i>tenellus</i> .....	659
<i>catenatus</i> .....	648, 2828	<i>vinetus</i> .....	637, 2827
<i>chrysolus</i> .....	655, 2828	<i>viridescens</i> .....	641
<i>cingulatus</i> .....	656, 2829	<i>xenicus</i> .....	662
<i>confluentus</i> .....	650, 2828	<i>zebra</i> .....	641, 647, 2827
<i>diaphnus</i> .....	645, 2828	<i>zebrinus</i> .....	646, 2827, 2828
<i>menona</i> .....	645	<i>zonatus</i> .....	657
<i>dispar</i> .....	658, 2830	<i>funebria</i> , <i>Gobiesox</i> .....	2334
<i>dovii</i> .....	650, 2828	<i>Gymnothorax</i> .....	396
<i>dingesii</i> .....	661	<i>Lycolontis</i> .....	396
<i>extensus</i> .....	646, 2827	<i>Noturus</i> .....	147
<i>floridensis</i> .....	642, 651, 2828	<i>Schilbeodes</i> .....	147
<i>fonticola</i> .....	643, 2827	<i>Sidera</i> .....	396
<i>funduloides</i> .....	650, 2828	<i>furca</i> , <i>Perca</i> .....	1200
<i>fuscus</i> .....	624	<i>Furcaria</i> .....	1545, 1546
<i>goodii</i> .....	2831	<i>cyanea</i> .....	1547
<i>grandis</i> .....	2827, 2828	<i>puncta</i> .....	1547
<i>guatemalensis</i> .....	660	<i>furcatum</i> , <i>Ditrema</i> .....	1506
<i>guttatus</i> .....	658, 2830	<i>furcatus</i> , <i>Amiurus</i> .....	134
<i>henshalli</i> .....	653	<i>Cypselurus</i> .....	737, 2836
<i>heteroclitus</i> .....	640	<i>Exocoetus</i> .....	737
<i>badus</i> .....	2827	<i>Ictalurus</i> .....	134
<i>grandis</i> .....	641	<i>Phaenodon</i> .....	1500
<i>macrolepidotus</i> .....	641, 2827	<i>Pimelodus</i> .....	134
<i>hieroglyphicus</i> .....	658, 2830	<i>Furcella</i> .....	2472, 2869
<i>jenkinsi</i> .....	651	<i>diaptera</i> .....	2472
<i>kansu</i> .....	2828	<i>furcoides</i> , <i>Characodon</i> .....	669
<i>labialis</i> .....	644, 2727	<i>furcifer</i> , <i>Anthias</i> .....	1222
<i>limbatus</i> .....	643, 649, 2828	<i>Brachyrhinus</i> .....	1222
<i>luce</i> .....	654	<i>Paranthias</i> .....	1227
<i>macdonaldi</i> .....	650, 2828	<i>Pimelodus</i> .....	135
<i>majalis</i> .....	636, 2827, 2828	<i>Sorranus</i> .....	1222
<i>melapleurus</i> .....	659, 2830	<i>furciger</i> , <i>Icelus</i> .....	1913
<i>mudfish</i> .....	641	<i>Furcimanus</i> .....	2869
		<i>furcata</i> , <i>Corvina</i> .....	1460

Page.		Page.		Page.
645	<i>furcraea</i> , <i>Perca</i> .....	1460	<i>gabonensis</i> , <i>Caranx setipinnis</i> .....	935
641	<i>furcraeus</i> , <i>Pachypops</i> .....	1459	<i>Vomer</i> .....	934, 2846
650	<i>Pachyurus</i> .....	1460	<i>Gadella</i> .....	2545
956, 2830	<i>furiosus</i> , <i>Noturus</i> .....	149	<i>Gadidae</i> .....	2531
927, 2828	<i>Schilbeoides</i> .....	149	<i>Gadinae</i> .....	2531
661	<i>furnieri</i> , <i>Micropogon</i> .....	1462	<i>Gadus</i> .....	2540
938, 2827	<i>Umbrina</i> .....	1463	<i>aglefinus</i> .....	2543
927, 2830	<i>furthi</i> , <i>Arius</i> .....	132	<i>agilis</i> .....	2534
641	<i>Caranx</i> .....	914	<i>albidus</i> .....	2531
952, 2828	<i>Corvina</i> ( <i>Homoprion</i> ) .....	1441	<i>arenosus</i> .....	2541
937, 2827	<i>Hemicaranx</i> .....	914	<i>auratus</i> .....	2542
649	<i>Ilisha</i> .....	436	<i>barbatus</i> .....	2541
644	<i>Pellona</i> .....	436	<i>brosme</i> .....	2561
944, 2827	<i>Pristipoma</i> .....	1319	<i>californicus</i> .....	2539
653	<i>Sciæna</i> .....	1441	<i>callarias</i> .....	2541
654	<i>Sphaeroides</i> .....	1737	<i>carbonarius</i> .....	2534
954, 2828	<i>Spheroides</i> .....	1737	<i>chalcogrammus</i> .....	2536
947, 2828	<i>Stellifer</i> .....	1441	<i>cimbrius</i> .....	2560
638	<i>Tachisurus</i> .....	132	<i>colinus</i> .....	2535
948, 2828	<i>Tachysurus</i> .....	132, 2787	<i>compressus</i> .....	2551
645	<i>furvus</i> , <i>Serranus</i> .....	1200	<i>fabricii</i> .....	2534
659	<i>fusca</i> , <i>Echeneis</i> .....	2270	<i>timbria</i> .....	1862
937, 2827	<i>Hydrargyra</i> .....	624	<i>glacialis</i> .....	2534
641	<i>Labrus tautoga</i> .....	1579	<i>gracilis</i> .....	2538
662	<i>Sciæna</i> .....	1483	<i>heteroglossus</i> .....	2541
947, 2827	<i>fuscatus</i> , <i>Silurus</i> .....	140	<i>lacustris</i> .....	137, 2551
927, 2828	<i>fuscoauratus</i> , <i>Tetragonopterus</i> .....	334	<i>longipes</i> .....	2555
657	<i>fusco-maculata</i> , <i>Acanth</i> .....	1540	<i>lubb</i> .....	2561
2334	<i>Echidna</i> .....	403	<i>macrocephalus</i> .....	2541
396	<i>fusco-maculatus</i> , <i>Chromis</i> .....	1540	<i>maculosus</i> .....	2551
396	<i>fuscula</i> , <i>Halioperca</i> .....	1211	<i>maius</i> .....	2541
147	<i>fusculus</i> , <i>Centropristes</i> .....	1211	<i>nivaldi</i> .....	2546
147	<i>fuscum</i> , <i>Siphostoma</i> .....	770	<i>merluccius</i> .....	2530
396	<i>fuscus</i> , <i>Aconurus</i> .....	1692	<i>merlus</i> .....	2530
1200	<i>Bythites</i> .....	2504	<i>molva</i> .....	2552
45, 1546	<i>Dialomus</i> .....	2868	<i>morhua</i> .....	2541
1547	<i>Eupomacentrus</i> .....	1552	<i>ogac</i> .....	2542
1547	<i>Fundulus</i> .....	624	<i>ogat</i> .....	2542
1506	<i>Gadus tomcodus</i> .....	2540	<i>periscopus</i> .....	2536
134	<i>Hemirhombus</i> .....	2686	<i>polymorphus</i> .....	2540
97, 2836	<i>Pomacentrus</i> .....	1552	<i>productus</i> .....	2531
737	<i>Psenes</i> .....	951	<i>proximus</i> .....	2539
134	<i>Serranus</i> .....	1181	<i>prinosus</i> .....	2540
1500	<i>Syngnathus</i> .....	770	<i>punctatus</i> .....	2553
134	<i>Trachinotus</i> .....	942	<i>pygmaeus</i> .....	2542
2, 2869	<i>fusiforme</i> , <i>Etheostoma</i> .....	1103	<i>raptor</i> .....	2552
2472	<i>fusiformis</i> , <i>Boleichthys</i> .....	1101	<i>ruber</i> .....	2530
669	<i>Boleosoma</i> .....	1102	<i>rupestris</i> .....	2541
1222	<i>Holoiepis</i> .....	1102	<i>saida</i> .....	2534
1222	<i>Phaianglites</i> .....	2048	<i>tau</i> .....	2316
1227	<i>Pœcilichthys</i> .....	1102	<i>tenuis</i> .....	2555
135	<i>fyllæ</i> , <i>Raja</i> .....	69	<i>tomcod</i> .....	2540
1222			<i>tomcodus</i> .....	2540
1913			<i>fuscus</i> .....	2540
2869	<i>Gabilan</i> .....	91	<i>luteus</i> .....	2540
1460	<i>gabonensis</i> , <i>Argyreolus</i> .....	935	<i>mixtus</i> .....	2540

	Page.		Page.
<i>Gadus torsk</i> .....	2561	<i>Galeocerdo tigrinus</i> .....	32
<i>vertagus</i> .....	2541	<i>galeoides</i> , <i>Otophidium</i> .....	2491
<i>virens</i> .....	2534	<i>Galeorhininae</i> .....	27
<i>Gaff-top-sail</i> .....	118	<i>Galeorhinus</i> .....	31
Catfishes .....	116	<i>zyopterus</i> .....	32
Pampano .....	940	<i>Galeus</i> .....	29, 31, 2745
<i>Gag</i> .....	1177	<i>californicus</i> .....	30
<i>Galdropsarime</i> .....	2532	<i>dorsalis</i> .....	32
<i>Gaidropsarus</i> .....	2557	<i>maculatus</i> .....	30
<i>argentatus</i> .....	2559	<i>galens</i> , <i>Epinephelus</i> .....	1164
<i>ensis</i> .....	2558	<i>Serranus</i> .....	1164
<i>septentrionalis</i> .....	2559	<i>Gallichthys</i> .....	931
<i>gaimardianus</i> , <i>Mugil</i> .....	814	<i>gallinula</i> , <i>Monacanthus</i> .....	1716
<i>gairdneri</i> , <i>Fario</i> .....	499	<i>Galliwasp</i> .....	538
<i>Salmo</i> .....	497	<i>Gallus</i> .....	931
<i>beardsleei</i> .....	2819	<i>virescens</i> .....	932
<i>creescens</i> .....	2821	<i>gallus</i> , <i>Zen</i> .....	936
<i>kamloops</i> .....	499	<i>galtia</i> , <i>Squalius</i> .....	237
<i>shasta</i> .....	502	<i>Gambusia</i> .....	678
<i>stonei</i> .....	503	<i>affinis</i> .....	680, 2832, 2833
<i>galaetura</i> , <i>Chloa</i> .....	279	<i>arlingtonia</i> .....	652, 2828, 2829
<i>galaeturus</i> , <i>Hysilepis</i> .....	279	<i>episcopi</i> .....	683
<i>Notropis</i> .....	279	<i>gracilis</i> .....	682, 683, 2832
<i>Galafate</i> .....	1711	<i>holbrooki</i> .....	681, 2832
<i>galapagorum</i> , <i>Umbrina</i> .....	1468	<i>humilis</i> .....	682
<i>galata</i> , <i>Belone</i> .....	716	<i>infans</i> .....	680, 2832
<i>galatus</i> , <i>Gymnocanthus</i> .....	2010	<i>melapleura</i> .....	2830
<i>Tylosurus</i> .....	716	<i>modesta</i> .....	693
<i>Galei</i> .....	21	<i>nicaranguensis</i> .....	682, 2832
<i>Galeichthys</i> .....	119, 122, 2770, 2771	<i>nobilis</i> .....	682, 2832
<i>assimilis</i> .....	2779	<i>patrnelis</i> .....	682, 2832
<i>azureus</i> .....	2775	<i>picturata</i> .....	683, 2832
<i>bahiensis</i> .....	119	<i>plumbea</i> .....	695
<i>blochii</i> .....	118	<i>punctata</i> .....	679
<i>caerulescens</i> .....	2776	<i>puncticulata</i> .....	680, 2832
<i>chevola</i> .....	932	<i>senilis</i> .....	682
<i>crinitus</i> .....	932	<i>speciosa</i> .....	681
<i>dasycephalus</i> .....	2780	<i>tridentiger</i> .....	2833
<i>eydoxii</i> .....	118	<i>Gambusiinae</i> .....	632
<i>gilberti</i> .....	2773	<i>Gambusia</i> .....	633, 635, 648
<i>gronovi</i> .....	117	<i>gambodon</i> , <i>Oxyrhina</i> .....	49
<i>gnatemalensis</i> .....	2778	<i>Ganoid Fishes</i> .....	100
<i>jordani</i> .....	2771	<i>Ganoidei</i> .....	100
<i>lentiginosus</i> .....	122, 2771	<i>Ganoids</i> , <i>Bony</i> .....	107
<i>longicephalus</i> .....	2781	<i>Cartilaginous</i> .....	100
<i>peruvianus</i> .....	122, 2771	<i>Gar</i> , <i>Alligator</i> .....	111
<i>phrygiatus</i> .....	2782	<i>Great</i> .....	111
<i>rugispinis</i> .....	2781	<i>Long-nosed</i> .....	109
<i>seemanni</i> .....	2772	<i>Short-nosed</i> .....	110
<i>surinamensis</i> .....	2780	<i>Gar Pikes</i> .....	108
<i>xenauchen</i> .....	2777	<i>garabata</i> , <i>Mojarra</i> .....	1353
<i>galeichthys</i> , <i>Carangoides</i> .....	932	<i>gardeniana</i> , <i>Hiatula</i> .....	1578
<i>Galeidae</i> .....	27	<i>gardenii</i> , <i>Centronotus</i> .....	948
<i>Galeina</i> .....	27	<i>Sternopyx</i> .....	966
<i>Galeocerdo</i> .....	32	<i>Stromateus</i> .....	966
<i>maculatus</i> .....	32	<i>gardoneum</i> , <i>Choudrostoma</i> .....	251

Page.	Page.	Page.
32	gardonensis, Abramis.....	251
2491	Leuciscus.....	251
27	Notemigonus.....	251
31	Gardonus.....	243
32	Garfish.....	714
29, 31, 2745	Garibaldi.....	1564
30	Garibaldis.....	1564
32	Garlopa.....	1186
30	garmani, Characodon.....	2832
1164	Gobius.....	2225
1164	Lepomis.....	1002
931	Notropis.....	281
1716	Garmanuia.....	2231, 2232
538	hemigymna.....	2233
931	paradoxa.....	2232
932	seminuda.....	2233
936	garnoti, Halicheres.....	1593
237	Iridio.....	1593
678	Julis.....	1593
0, 2832, 2833	Platygllossus.....	1593
2, 2828, 2829	Garrapa.....	1161, 1797
683	nigrita.....	1161
32, 683, 2832	Gascon.....	910
681, 2832	Gaspereau.....	426
682	Gaspergon.....	1484
680, 2832	Gasteracanthus.....	746
2830	Gasteropelecus.....	337
693	acanthurus.....	579
682, 2832	crocodilus.....	558
682, 2832	humboldti.....	572
682, 2832	maculatus.....	338
683, 2832	Trachinus.....	2297
695	Gasterosteia.....	745
679	blanchardi.....	746
680, 2832	Gasterosteida.....	742
682	Gasterosteinae.....	743
681	Gasterosteus.....	746
2833	aculeatus.....	747, 2836
632	cataphractus.....	750
33, 635, 648	algeriensis.....	748
49	antecessor.....	900
100	apeltes.....	752
100	argentissimus.....	747
107	argyropomus.....	748
100	bailloni.....	747
111	biaculeatus.....	748
111	bispinosus.....	748
109	atkinsi.....	748
110	cuvieri.....	749, 2836
108	brachycentrus.....	748
1353	breviceps.....	746
1578	canadus.....	948
948	carolinus.....	944
966	cataphractus.....	749
966	concinuus.....	745
251	dekeyi.....	746
	Gasterosteus, dimidiatus.....	740
	ductor.....	900
	elegans.....	748
	gladiocentrus.....	2836
	globiceps.....	744
	gymnurus.....	748
	inconstans.....	744
	inscriptus.....	750
	intermedius.....	750
	islandicus.....	748
	hevis.....	745
	lelurns.....	747
	loricatus.....	747
	locharingus.....	746
	mainensis.....	745
	micropus.....	744
	millipunctatus.....	752
	nebulosus.....	746
	neustriannus.....	747
	obolaris.....	750
	occidentalis.....	745
	plebeius.....	751
	ponticus.....	747
	pugetti.....	751
	pungitius.....	745
	brachypoda.....	746
	pygmaeus.....	744
	quadraeus.....	752
	saltatrix.....	947
	semilarmatus.....	747
	semiloricatus.....	747
	serratus.....	750
	spinulosus.....	748
	tetracanthus.....	748
	trachurus.....	747
	wheatlandi.....	749
	williamsoni.....	750
	microcephalus.....	750
	Gastrobranchus.....	7
	ciccus.....	8
	dombey.....	6
	Gastrophysus.....	1727
	Gastropsetta.....	2636
	frontalis.....	2636
	Gastrostomus.....	406
	bairdii.....	406
	Gata.....	26
	Gato.....	28
	Bonaei.....	1187
	gaucharate, Chaetostomus.....	159
	Hemiancistrus.....	159
	Hypostomus.....	159
	gavails, Lepisosteus.....	110
	gayi, Epicopus.....	2530
	gelatinosum, Melanostigma.....	2479

	Page.		Page.
gelatinosus, Careproctus .....	2134, 2135	Gerres .....	1373, 1377
Cyclopterus .....	2135	apriou .....	1373
Liparis .....	2134, 2135	argenteus .....	1371
gelida, Ammoerypta .....	1064	aureolus .....	1375
gefidus, Ceratichthys .....	317	axillaris .....	1378
Gobio .....	317	brasiliannus .....	1378
Hypopsis .....	316, 317	brevimanus .....	1377
geminatus, Bleennius .....	2385	brevirostris .....	1376
Hypenrochilus .....	2385	californiensis .....	1370
gemua, Hypoplectrus .....	1193	cinereus .....	1370
gemmifer, Astronesthes .....	586	dowi .....	1368
Lampanyctes .....	559	embryx .....	1379
Gempylidae .....	877	gracilis .....	1370
Gempylinae .....	878	gula .....	1371
Gempylus .....	883	harengulus .....	1369
coluber .....	884	jonesi .....	1368
ophidiannus .....	884	lineatus .....	1377
prometheus .....	883	mexicanus .....	1380
serpens .....	884	olisthostoma .....	1377
solandri .....	883	olisthostomus .....	1376
generosus, Catostomus .....	170	patuo .....	1378
Pantosteus .....	170	peruvianus .....	1376
Genicanthus .....	1682	plumieri .....	1379
tricolor .....	1684	pseudogula .....	1368
Genizara .....	1586	rhombus .....	1374
genizara, Clepticus .....	1587	squampinnis .....	1373
Rabirubia .....	1586	zebra .....	1373
gentilis, Bleennius .....	2388	Gerridae .....	1366
Hypsoblennius .....	2387	ghini, Orthragoriscus .....	1754
Isosthes .....	2388	Ghost-fish .....	2443
Genyatremus .....	1342	gibba, Pterophryne .....	2717
luteus .....	1342	gibber, Salmo .....	478
Genyonemus .....	1460	gibbiceps, Heros .....	1536
fasciatus .....	1479	gibbifrons, Cypselurus .....	2836
lineatus .....	1460	Exocoetus .....	741
Genyoroce .....	1247	Gibbonsia .....	2351
viridis .....	1246	elogans .....	2353, 2869
Genypterus omostigma .....	2490	evides .....	2352, 2869
Genytremus .....	1314	gibbonsii, Holocentrus .....	1509
interruptus .....	1319	gibbosa, Ciliola .....	272
geometricus, Anchisonus .....	1736	Gila .....	235
Chilomycterus .....	1749	Moniana .....	272
Diodon .....	1748, 1749	Perca .....	1009, 1236
Tetrodon .....	1735, 1736	marina .....	1295
Zeus .....	936	Tigoma .....	235
Geophagus .....	1542	gibbosum, Haemulon .....	1296
crassilabris .....	1543	gibbosus, Aphredoderus .....	787
georgianus, Caranx .....	927	Calliodon .....	1296
Exocoetus .....	730	Catostomus .....	186
georgii, Tetrapturus .....	892	Cyprinodon .....	672
gerania, Belone .....	716	Eupomotis .....	1009
germannus, Notropis .....	261	Holocentrus .....	1319
germo .....	870	Leuciscus .....	231
alalunga .....	871	Pomotis .....	1005
germo, Scomber .....	871	Squalius .....	231
Germon .....	871	gibbsii, Hemilipidotus .....	1936





	Page.		Page.
glaber, Cycloptericthys .....	2104	Glossodus forskali .....	411
Ostracodon oblongus .....	1735	Glossoplites .....	991
Pleuronectes .....	2650	gloveri, Salmo .....	487
glabra, Liopsetta .....	2650	Glut Herring .....	426
Platessa .....	2650	glutnosa, Myxine .....	7
glabro, Corpore oblongo .....	2657	Glyphidodon .....	1560
Ostracodon subrotundus ven- tro .....	1749	chrysurus .....	1567
glaciale, Myctophum .....	574	declivifrons .....	1562
glacialis, Cottus .....	1976	rudis .....	1563
Gadus .....	2534	saxatilis .....	1562
Liopsetta .....	2649, 2650	taurus .....	1563
Pleuronectes .....	2649	trosebelli .....	1562
Scopelus .....	574	Glyphisodon .....	1560, 1561
Squalus .....	57	moucharra .....	1562
gladiunculus, Gasterosteus .....	2836	rubicundus .....	1565
gladius, Trichurus .....	887	Glyptocephalus .....	2656
Tylosurus .....	710	acanthinus .....	2617
Xiphias .....	894	cynoglossus .....	2656
Glance Fish .....	954	pacificus .....	2655
Glasseye .....	1021	saxicola .....	2657
glauca, Prionace .....	33	zachirus .....	2658
glaucofrænum, Coryphopterus .....	2220	Gnathanodon .....	927
Gobius .....	2219	Gnathobolus .....	437
glaucoides, Trachinotus .....	941	mucronatus .....	438
glancopareius, Acanthurus .....	1694	Gnathocentrum .....	1687
glancos, Sebastodes .....	1777	Gnathodor speciosus .....	928
glaucoctictus, Rhinobatus .....	63	gnathodus, Pseudoscarius .....	1650
glancostigma, Rhinobatus .....	62	Scarus .....	1650
glancus, Carcharhinus .....	33	Gnatholepis .....	2210
Carcharias .....	33	Gnathophis .....	355
Chaetodon .....	941	Gnathypops .....	2283
Isuropsis .....	48	macrops .....	2284
Squalus .....	33	maxillosa .....	2284
Trachinotus .....	940	maxillosus .....	2284
Trachynotus .....	941	mystacina .....	2286
Glaustegus .....	61	mystacinus .....	2286
Globefish .....	1734	rhomalea .....	2285
globiceps, Agonostoma .....	821	scops .....	2283
Blennicottus .....	2017	snyderi .....	2285
bryosus .....	2017	Goatfish, Red .....	858
Centridermichthys .....	2017	Yellow .....	859
Gasterosteus .....	744	Goatfishes .....	857
Oligocottus .....	2017	Gobies .....	2184, 2188, 2210
globosa, Lyosphaera .....	1752	Blind .....	2261
gloriosus, Bryttus .....	994	Crested .....	2209
Enneacanthus .....	993	Half-naked .....	2231
Glossamia .....	1111	Naked .....	2257
pandionis .....	1111	Gobiesocidae .....	2326
Glossichthys .....	2704	Gobiesocinae .....	2327
plagiusa .....	2710	Gobiesox .....	2320, 2330, 2331
Glossodon .....	412	adustus .....	2334
harengoides .....	413	carneus .....	2337
heterurus .....	413	cephalus .....	2332
glossodonta, Argentina .....	411	cerasimus .....	2336
Glossodus .....	411	eigenmanni .....	2339
		eoss .....	2343

Page.		Page.		Page.
411	<i>Gobiesox erythropterus</i>	2336	<i>Gobiosoma alepidotum</i>	2259
901	<i>fasciatus</i>	2338	<i>boscii</i>	2259
487	<i>funebria</i>	2334	<i>centrocentrum</i>	2261
426	<i>gyrinus</i>	2331	<i>crescentale</i>	2259
7	<i>haeres</i>	2337	<i>crescentalis</i>	2260
1560	<i>humeralis</i>	2341	<i>histrio</i>	2258
1567	<i>macrophthalmus</i>	2335	<i>ios</i>	2255
1562	<i>nigripinnis</i>	2332	<i>longipinna</i>	2256
1563	<i>nudus</i>	2331	<i>molestum</i>	2258
1562	<i>papillifer</i>	2330	<i>multifasciatum</i>	2260
1563	<i>pinniger</i>	2329	<i>zosterurum</i>	2257
1562	<i>poecilophthalmus</i>	2335	<i>Gobius oblongus</i>	2264
1560, 1561	<i>punctulatus</i>	2338	<i>Gobius</i>	2209, 2211, 2216
1562	<i>reticulatus</i>	2328	<i>alepidotus</i>	2259
1565	<i>rhesodon</i>	2340	<i>amorea</i>	2201
2656	<i>rhodospilus</i>	2335	<i>andrei</i>	2218
2677	<i>rubiginosus</i>	2337	<i>bucalus</i>	2230
2656	<i>rupestris</i>	2341	<i>badius</i>	2227
2655	<i>strumosus</i>	2333	<i>banana</i>	2236
2657	<i>tudes</i>	2333	<i>boleosoma</i>	2221
2658	<i>virgatulus</i>	2333	<i>boscii</i>	2227, 2259, 2268
927	<i>zebra</i>	2342	<i>brunneus</i>	2218
437	<i>Gobiichthys</i>	2210	<i>cauda longissima acuminata</i>	2230
438	<i>Gobiidae</i>	2188	<i>carolinensis</i>	2218
1687	<i>Gobiinae</i>	2190	<i>catulus</i>	2218
928	<i>Gobio aestivalis</i>	316	<i>chignita</i>	2241
1650	<i>cataractae</i>	306	<i>cristagalli</i>	2209
1650	<i>gelidus</i>	317	<i>cyprinoides</i>	2209
2210	<i>plumbeus</i>	324	<i>dalli</i>	2230
355	<i>vernalis</i>	321	<i>dolichocephalus</i>	2237
2283	<i>gobio, Clinus</i>	2365	<i>eigemannii</i>	2218
2284	<i>Cottus</i>	1941, 1968, 2000	<i>emblematicus</i>	2247
2284	<i>Gobioclinus</i>	2365	<i>encanemus</i>	2223
2284	<i>Gobiodellus</i>	2364	<i>fasciatus</i>	2222
2286	<i>gobio</i>	2365	<i>flavus</i>	2235
2286	<i>Gobioidea</i>	2184	<i>garmani</i>	2225
2285	<i>Gobioidei</i>	781	<i>glaucofrantum</i>	2219
2283	<i>Gobioides</i>	2263, 2268	<i>gracilis</i>	2240
2285	<i>barreto</i>	2264	<i>gronovii</i>	949
858	<i>broussoneti</i>	2264	<i>gulosus</i>	2244
859	<i>broussonetii</i>	2263	<i>hastatus</i>	2229
857	<i>peruanus</i>	2264	<i>hemigyunus</i>	2233
188, 2210	<i>gobioidea, Cottus</i>	1968	<i>kraussi</i>	2228
2261	<i>Uranidea</i>	1968	<i>lacertus</i>	2218
2209	<i>Gobioiidae</i>	2192	<i>lancoletus</i>	2220, 2230
2231	<i>Gobiomorus</i>	2194	<i>lepidus</i>	2249
2257	<i>dornitator</i>	2195	<i>lineatus</i>	2218, 2260
2326	<i>dormitor</i>	2195	<i>longicauda</i>	2229
2327	<i>gronovianns</i>	950	<i>lucretiae</i>	2268
330, 2331	<i>Gobionellus</i>	2210, 2215, 2227	<i>lyricus</i>	2224
2334	<i>hastatus</i>	2229	<i>manglicola</i>	2220
2337	<i>oceanicus</i>	2230	<i>mapo</i>	2218
2332	<i>amaragdus</i>	2228	<i>murtiniensis</i>	2236
2336	<i>stigmaticus</i>	2224	<i>mexicanus</i>	2237
2339	<i>gobionium, Campostoma</i>	200	<i>microdon</i>	2227
2343	<i>Gobiosoma</i>	2257	<i>minutus</i>	2097

	Page.		Page.
<i>Gobius nelsoni</i> .....	2235	<i>Gonocephalus macrocephalus</i> .....	2184
<i>newberryi</i> .....	2248	<i>Gonocheilodon</i> .....	1672
<i>nicholsi</i> .....	2218	<i>Gonopterus</i> .....	1687
<i>oceanicus</i> .....	2230	<i>merens</i> .....	1688
<i>paradoxus</i> .....	2232	<i>Gonostoma</i> .....	578
<i>pisonsi</i> .....	2201	<i>brevidens</i> .....	570
<i>plumieri</i> .....	2206	<i>denddatum</i> .....	579
<i>poeyi</i> .....	2226	<i>elongatum</i> .....	583
<i>quadriflorus</i> .....	2221	<i>microndon</i> .....	582
<i>sagittula</i> .....	2228	<i>Gonostominae</i> .....	578
<i>seminudus</i> .....	2234	<i>Gonostomus acanthurus</i> .....	579
<i>shufeldti</i> .....	2221	<i>Goodea</i> .....	685
<i>amaragdus</i> .....	2227	<i>atriphnis</i> .....	685
<i>smyrnensis</i> .....	2118	<i>goodei</i> , <i>Aldrovandia</i> .....	610
<i>soporator</i> .....	2216	<i>Erlmyzon</i> .....	186
<i>stigmaticus</i> .....	2224	<i>Halosaurus</i> .....	610
<i>stigmaturus</i> .....	2220	<i>Hymenocephalus</i> .....	2572
<i>strigatus</i> .....	2228	<i>Lucania</i> .....	664
<i>talasica</i> .....	2236	<i>Macrurus</i> .....	2572
<i>thalassius</i> .....	2245	<i>Myliobatis</i> .....	2755
<i>townsendi</i> .....	2250	<i>Nematonurus</i> .....	2571
<i>viridipallidus</i> .....	2259	<i>Paralichthys</i> .....	1480
<i>wurdemanni</i> .....	2225	<i>Ptilichthys</i> .....	2452
<i>zebra</i> .....	2226	<i>Sebastichthys</i> .....	1780
<i>gobius</i> , <i>Liparis</i> .....	2108	<i>Sebastodes</i> .....	1779
<i>goboides</i> , <i>Hypsicometes</i> .....	2294	<i>Splivomer</i> .....	367
<i>Goby</i> , Long-jawed .....	2250	<i>Trachinotus</i> .....	943
Naked .....	2259	<i>Urolophus</i> .....	81
Sharp-tailed .....	2220	<i>Goodeinae</i> .....	632
<i>goddefroyi</i> , <i>Percichthys</i> .....	1197	<i>Goodfish</i> .....	487
<i>godmanni</i> , <i>Cichlasoma</i> .....	1516	<i>Goodios</i> .....	1458
<i>Heros</i> .....	1516	<i>Goody</i> .....	1458
<i>Pimelodus</i> .....	152	<i>Goosefish</i> .....	2713
<i>Rhamdia</i> .....	152	<i>Gorbusecha</i> .....	478
<i>Goggle-eye</i> .....	990, 992	<i>gorbuscha</i> , <i>Oncorhynchus</i> .....	478
<i>Jack</i> .....	911	<i>Salmo</i> .....	478
<i>Goggler</i> .....	911	<i>Gordichthys</i> .....	363
<i>Golden Shiner</i> .....	250	<i>irretitus</i> .....	363
Trout of Mount Whitney ..	503	<i>gorensis</i> , <i>Albula</i> .....	412
<i>Goldfish</i> .....	201	<i>Trachynotus</i> .....	943
<i>Golet</i> .....	507	<i>Vomer</i> .....	934
<i>Goltra</i> .....	508	<i>gonani</i> , <i>Lepidopus</i> .....	887
<i>Goma soi</i> .....	1833	<i>Goujon</i> .....	2790
<i>gomesii</i> , <i>Ophichthys</i> .....	384	<i>Gourd-seed Sucker</i> .....	168
<i>Ophichthys</i> .....	385	<i>gracile</i> , <i>Boleosoma</i> .....	1102
<i>Ophisurus</i> .....	385	<i>Myctophum</i> .....	572
<i>Gonenion</i> .....	946	<i>Peristedion</i> .....	2179
<i>serra</i> .....	947	<i>gracilis</i> , <i>Aldrovandia</i> .....	610
<i>Goniobatis</i> .....	88	<i>Atherinichthys</i> .....	797
<i>macroptera</i> .....	88	<i>Blennius</i> .....	2438
<i>Goniodus</i> .....	57	<i>Catostomus</i> .....	179
<i>Gonionarce</i> .....	78	<i>Chaetodon</i> .....	1675
<i>Gonioperca</i> .....	1194	<i>Cottus</i> .....	1968
<i>Gonioplectrus</i> .....	1139	<i>Cyprinus</i> .....	326
<i>hispanus</i> .....	1140	<i>Dapterus</i> .....	1370
<i>Gonocephalus</i> .....	2183	<i>Galus</i> .....	2538

Page.		Page.		Page.
2184	<i>gracilis</i> , <i>Gambusia</i> .....	682, 683	<i>Grammistes acuminatus</i> .....	1487
1072	<i>Gerres</i> .....	1370	<i>hepatus</i> .....	1343
1087	<i>Gila</i> .....	227	<i>mauribii</i> .....	1323
1088	<i>Gobius</i> .....	2249	<i>trivittatus</i> .....	1311
578	<i>Hippocampus</i> .....	777	<i>ulmaeulatus</i> .....	1360
579	<i>Hybopsis</i> .....	321	<i>Grand Oranche</i> .....	1057
579	<i>Lepidogobius</i> .....	2249	<i>Grande Écaille</i> .....	409
583	<i>Lepisosteus</i> .....	110	<i>grandicussis</i> , <i>Arius</i> .....	126
582	<i>Leptocephalus</i> .....	354	<i>Nefima</i> .....	126, 2764
578	<i>Leuciscus</i> .....	281, 326	<i>Tachisurus</i> .....	126
579	<i>Lycodes</i> .....	2465	<i>stricticus-</i>	
685	<i>Menidia</i> .....	797	<i>sis</i> .....	126
685	<i>beryllina</i> .....	797	<i>grandicornis</i> , <i>Scorpaena</i> .....	1850
610	<i>Moniana</i> .....	272	<i>grandipinnis</i> , <i>Photogenis</i> .....	280
186	<i>Porca</i> .....	1024	<i>grandis</i> , <i>Fundulus</i> .....	2827
610	<i>Photoneutes</i> .....	591	<i>heteroclitus</i> .....	641
2572	<i>Pinelodus</i> .....	135	<i>Gila</i> .....	225, 2707
664	<i>Platygobio</i> .....	326	<i>Leuciscus</i> .....	225
2572	<i>Pleurogadus</i> .....	2538	<i>Ptychocheilus</i> .....	225, 2796
2755	<i>Poecilichthys</i> .....	1103	<i>grandisquamis</i> , <i>Eleotris</i> .....	2198
2571	<i>Ptychocheilus</i> .....	225	<i>grandisquamis</i> , <i>Chacrojulis</i> .....	1597
1480	<i>Scomber</i> .....	867	<i>PlatyGLOSSUS</i> .....	1597
2452	<i>Scopelus</i> .....	572	<i>Upeneus</i> .....	860
1780	<i>Septogunnellus</i> .....	2436	<i>grandoculatus</i> , <i>Centropomus</i> .....	1120
1770	<i>Tigona</i> .....	236	<i>grandocule</i> , <i>Chirostoma</i> .....	2839
367	<i>Tilesia</i> .....	2538	<i>grandoculis</i> , <i>Atherinichthys</i> .....	2840
943	<i>Umbrina</i> .....	1474	<i>granulata</i> , <i>Percu</i> .....	1024
81	<i>Uranidea</i> .....	1968	<i>Rala</i> .....	72
632	<i>Xiphophorus</i> .....	683	<i>granulatus</i> , <i>Amorphocephalus</i> .....	1619
487	<i>graculosus</i> , <i>Pinelodus</i> .....	135	<i>granulosa</i> , <i>Pristis</i> .....	61
1458	<i>gradlens</i> , <i>Hypsagonus</i> ( <i>Cheira-</i>		<i>Graodus</i> .....	254
1458	<i>gonus</i> ) .....	2041	<i>nigrotentatus</i> .....	264
2713	<i>graeli</i> , <i>Ophidion</i> .....	2488	<i>Grass Bass</i> .....	987
478	<i>grahami</i> , <i>Gila</i> .....	227	<i>Porgy</i> .....	1355
478	<i>Leuciscus</i> .....	228	<i>Rockfish</i> .....	1819
478	<i>Oligocephalus</i> .....	1089	<i>Gray Grunt</i> .....	1296
363	<i>grallator</i> <i>benthosaurus</i> .....	543	<i>Pike</i> .....	1022
363	<i>Gramma</i> .....	1228	<i>Snapper</i> .....	1255
412	<i>loreto</i> .....	1229	<i>grayi</i> , <i>Carploides</i> .....	167
943	<i>Grammatens</i> .....	1347, 1348, 1353	<i>Lepidosteus</i> .....	111
934	<i>humilis</i> .....	1355	<i>Salmo</i> .....	509
887	<i>medius</i> .....	1356	<i>Grayling</i> , <i>Aretic</i> .....	517, 2871
2700	<i>grammaticum</i> , <i>Thalassoma</i> .....	1610, 2859	<i>European</i> .....	2870
168	<i>grammaticus</i> , <i>Chlorichthys</i> .....	1610	<i>Michigan</i> .....	518, 2871
1102	<i>Grammatopleurus</i> .....	1866	<i>Montana</i> .....	519, 2871
572	<i>lagocephalus</i> .....	1875	<i>Graylings</i> .....	517
2179	<i>Grammatostomias</i> .....	590	<i>Great Albacore</i> .....	870
610	<i>dentatus</i> .....	590	<i>Amber-fish</i> .....	903
797	<i>Grammichthys</i> .....	2693	<i>Barracuda</i> .....	823
2438	<i>lineatus</i> .....	2702	<i>Bear Lake Herring</i> .....	470
179	<i>Grammicolepididæ</i> .....	973	<i>Blue Shark</i> .....	33
1675	<i>Grammicolepis</i> .....	974	<i>Chub</i> .....	232
1968	<i>brachiusculus</i> .....	974	<i>Flounder</i> .....	2652
326	<i>Grammiconotus</i> .....	725	<i>Flying-fish</i> .....	740
1370	<i>bicolor</i> .....	726	<i>Fork-tailed Cat</i> .....	137
2538	<i>Gramminæ</i> .....	1131	<i>Gar</i> .....	111

	Page.		Page.
Great Lake Trout .....	504	grenlandicus, <i>Cottus scorpius</i> .....	1975
Northern Pike .....	630	Gunnellus .....	2418
Pámpano .....	943	Himantolophus .....	2733
Pike .....	629	Hippoglossus .....	2611
Pipefish .....	764	Microstomus .....	528
Sculpin .....	1976	Myoxocephalus .....	1974
Sculpins .....	1970	Salmo .....	521
Sea Lamprey .....	10	Gronias .....	135, 136, 142
Tunnies .....	869	nigrilabris .....	142
White Shark .....	50	gronovianus, <i>Gobiomorus</i> .....	950
grebnitskii, <i>Pholidapus</i> .....	2431	gronovii, <i>Achirus</i> .....	2696
Green Bass .....	1012	Allurichthys .....	117
Cod .....	2534	Galeichthys .....	117
Parrot-fish .....	1638, 1657	Gobius .....	949
Pike .....	627	Nomets .....	949
Sturgeon .....	104	Ostracion .....	1725
Sunfish .....	996	Solea .....	2606
Green-back Trout .....	497	Zoarcetes .....	2457
Green-fish .....	1382	grossidens, <i>Engraulis</i> .....	451
Alaska .....	1869	Lycengraulis .....	451
greeni, <i>Uranidea</i> .....	1965	Ground Drummer .....	1436
greeni, <i>Conesius</i> .....	324	Spearing .....	533
Neoliparis .....	2112	Groupers, Black .....	1161, 1174
Greenland Charr .....	508, 510	Mangrove .....	1171
Codfish .....	2542	Nassau .....	1157
Halibut .....	2611	Red .....	1160
Greenling .....	1871	Yellow .....	1183
Cecalings .....	1863, 1866	Yellow-finned .....	1155, 1172
Green-sided Darter .....	1053	Groupers .....	1148
Gregory, Bean .....	1555	Grubber Brood-head .....	447
Grenadiers .....	2561	Grubby .....	1973
grex, <i>Scomber</i> .....	867	grunniens, <i>Ambloplites</i> .....	1484
grimaldii, <i>Conchognathus</i> .....	349	Aplodinotus .....	1484
Grindle .....	113	Haploidonotus .....	1484
grisea, <i>Dionda</i> .....	216	Labrus .....	1483
Lucioperca .....	1022	Mugil .....	1483
Sciama .....	1484	Grunt, Black .....	1297
Umbracabertura .....	342	Boar .....	1303
griseolineatum, <i>Siphostoma</i> .....	764	Compton .....	1304
griseolineatus, <i>Syngnathus</i> .....	764	French .....	1306
griseum, <i>Stizostedion canadense</i> .....	1022	Gray .....	1296
griseus, <i>Acomus</i> .....	175	Margaret .....	1295
Carcharias .....	47	Open-mouthed .....	1306
Catostomus .....	175	Red-mouthed .....	1308
Hexanchus .....	19	Striped .....	1296
Labrus .....	1257	White .....	1310
Lutjanus .....	1257	Yellow .....	1303
Mesoprion .....	1257	Grunters .....	1289
Neomans .....	1255	Grunts .....	1291
Notidanus .....	19	Striped .....	1313
Saurus .....	537	gryllus, <i>Exocoetus</i> .....	729
Squalus .....	19	Grystes .....	1010
grollsi, <i>Ophiodon</i> .....	2487	lineatus .....	1868
gronlandien, <i>Nausenia</i> .....	528	megastoma .....	1012
gronlandicus, <i>Aspidophoroides</i> .....	2092	nobilis .....	1012
Cottus .....	1975	nuceensis .....	1012

Page.	Page.	Page.
1975	Guabi coara brasiliensis..... 1305	guavina, Guavina..... 2198
2418	Guacamaia..... 1657, 1658	Guavinas..... 2194
2733	guacamaia, Hemistoma..... 1659	guaymasia, Gillichthys..... 2252
2611	Pseudoscarus..... 1656, 1659	guaza, Epinephelus..... 1154
528	Scarus..... 1656, 1658	Labrus..... 1154
1974	Guacamaias..... 1655	Gudlax..... 954
521	guachancho, Sphyræna..... 824	Guebucu..... 891
5, 136, 142	Guachinango, Pargo..... 1264	guebucu, Skeponopodus..... 891
142	guaco, Amore..... 2236	guentheri, Percina..... 1034
950	Guaguanche..... 824	Guerubaco..... 2198
2696	Pelon..... 824	Gueule, Petite..... 1370
117	guaiacan, Echeinis..... 2270	guianensis, Belone..... 715
117	Guajacon..... 679	guichenoti, Cayennia..... 2265
949	Guajacones..... 678	guineensis, Ostracion..... 1725
949	Guajica..... 652	Guiritinga..... 119
1725	Guamajacu atinga..... 1749	Guitarfish..... 63
2696	Guamajacu guara..... 1745	Guitarfishes..... 61
2457	guanabana, Erizo..... 1746	Guitarro..... 62
451	Guapena..... 1489	gula, Eucinostomus..... 1370
451	Guaperva..... 1703	Gerres..... 1371
1436	lataforcipata..... 1702	Gila..... 234
533	guara, Caranx..... 926	Squalius..... 234
1161, 1174	Guamajacu..... 1745	Gulf Flounder..... 2631
1171	Scomber..... 927	Menhaden..... 434
1157	Guarapucu..... 876	Shad..... 2810
1160	Guardfish..... 715	gulo, Holocentrus..... 1139
1183	Guasa..... 1154, 1162	gulonellus, Lencosomus..... 326
1155, 1172	guasa, Promicrops..... 1164	Pogonichthys (Platy-
1148	Serranus..... 1164	gobio)..... 236
447	Guaseta..... 1164	gulosa, Uranidea..... 1945
1973	guatemalensis, Arius..... 129	gulosus, Cathorops..... 1948
1484	Adinia..... 660	Centridermichthys..... 1943
1484	Anacyrtus..... 338	Chenobryttus..... 992
1484	Atherinichthys..... 801	Cottopsis..... 1945
1483	Ctenarichthys..... 2686	Cottus..... 1944
1483	Fundulus..... 660	Gobius..... 2244
1297	Galeichthys..... 2778	Lepidogobius..... 2244
1303	Hexanemafichthys..... 129	Pomotis..... 992
1304	Menidia..... 801	Tachisurus..... 133
1306	Pimelodus..... 152	Gulpers..... 404
1296	Rhamdia..... 152	gulula, Eucinostomus..... 1371
1295	Ro-boides..... 338	gummigutta, Hypoplectrus..... 1192
1306	Guativere..... 1144, 1145	unicolor..... 1192
1308	Amarilla..... 1144, 1145	Plectropoma..... 1192
1296	Black..... 1146	gundlachi, Dormitator..... 2198
1310	Red..... 1145	guelliformis, Asternopteryx..... 2420
1303	guativere, Bodianus..... 1145	Centronotus..... 2421
1289	Serranus..... 1145	Muraenoides..... 2421
1291	Guatucupa juba..... 1323	Gumellus ingens..... 2419
1313	Guavina..... 2194, 2198, 2201	macrocephalus..... 2419
729	guavina..... 2198	Gunnell..... 2419
1010	Hoyem..... 2236	Gunnellops..... 2420
1868	Mupo..... 2196	rosens..... 2420
1012	Tétard..... 2200	Gunnellus alectrolophus..... 2422
1012	guavina, Batrachus..... 2195	anguillaris..... 2436
1012	Eleotris..... 2199	apos..... 2430

	Page.		Page.
<i>Gunnellus dolichogaster</i> .....	2417	<i>guttatus</i> , Johnius.....	1174
<i>labrici</i> .....	2438	<i>Lampris</i> .....	955
<i>fasciatus</i> .....	2418	<i>Lutjanus</i> .....	1269
<i>grendlandicus</i> .....	2418	<i>Lutjanus</i> .....	1269
<i>blunthius</i> .....	2430	<i>Mesoprion</i> .....	1269
<i>marinoides</i> .....	2418	<i>Muraenoides</i> .....	2419
<i>polyodon</i> .....	2414	<i>Nemipops</i> .....	1269
<i>perforatus</i> .....	2420	<i>Ophisurus</i> .....	382
<i>punctatus</i> .....	2440	<i>Percopsis</i> .....	784
<i>ruberrimus</i> .....	2417	<i>Potrometopon</i> .....	1117
<i>vulgaris</i> .....	2419	<i>Pomotis</i> .....	994
<i>gunnellus</i> , Blennius.....	2419	<i>Promicrops</i> .....	1162
<i>Centronotus</i> .....	2419	<i>Sebastapistes</i> .....	1848
<i>Muraenoides</i> .....	2419	<i>Upsilonphorus</i> .....	231
<i>Pholis</i> .....	2419	<i>Zeus</i> .....	95
<i>Gunnels</i> .....	2414	<i>Zygonectes</i> .....	658
<i>gunneri</i> , Scorpaenidae.....	955	<i>guttavarium</i> , <i>Plectropoma</i> .....	119
<i>gunnerianus</i> , <i>Squalus</i> .....	51	<i>guttavarius</i> , <i>Hypoplectrus</i> .....	119
<i>gunnisoni</i> , <i>Clupea</i> .....	273	<i>unicolor</i> .....	119
<i>Cyprinella</i> .....	273	<i>guttifer</i> , <i>Ophichthus</i> .....	384
<i>guntheri</i> , <i>Aspidophoroides</i> .....	2090	<i>Ophichthus</i> .....	384
<i>Etheostoma</i> .....	1034	<i>guttulatus</i> , <i>Hippocampus</i> .....	778
<i>Hadropterus</i> .....	1033	<i>Pisodonophis</i> .....	377
<i>Halosaurus</i> .....	608	<i>Pleuronectes</i> .....	2649
<i>Hoplopagrus</i> .....	1244	<i>Pleuronichthys</i> .....	2649
<i>Lampayetius</i> .....	559	<i>guzmanicensis</i> , <i>Catostomus</i> .....	171
<i>Mugil</i> .....	812	<i>Pantosteus</i> .....	171
<i>Sireno</i> .....	2523	<i>Gymnachirus</i> .....	249
<i>Sphyriena</i> .....	824	<i>fasciatus</i> .....	2703
<i>Xiphophorus</i> .....	702	<i>Gymnelcotris</i> .....	2291
<i>guntheriana</i> , <i>Alutera</i> .....	1720	<i>seminuda</i> .....	2291
<i>guppil</i> , <i>Girardinus</i> .....	2834	<i>seminudus</i> .....	2294
<i>Gurnard</i> , Big-headed.....	2171	<i>Gymnelina</i> .....	2456
Common.....	2156	<i>Gymnelis</i> .....	2477
Flying.....	2183	<i>pietus</i> .....	2477
Northern Striped.....	2167, 2168	<i>stigma</i> .....	2477
Red.....	2177	<i>viridis</i> .....	2477
<i>Gurnards</i> .....	2147, 2148, 2152	<i>Gymnepignathus</i> .....	907
Deep-water.....	2177	<i>Gymnocanthus</i> .....	2000
Flying.....	2182	<i>galeatus</i> .....	2010
Mottled.....	2176	<i>pistilliger</i> .....	2006, 2008, 2009
Small-scaled.....	2175	<i>tricuspis</i> .....	2008
<i>Gurnardus</i> .....	2148	<i>Gymnocephalus</i> .....	962
<i>Gusas</i> .....	1162	<i>ruber</i> .....	1146
<i>guttata</i> , <i>Mycteroperca venenosa</i> .....	1174	<i>Gymnodontes</i> .....	781, 1729
<i>Perca</i> .....	1142, 1161	<i>Gymnogaster</i> .....	888
<i>Scorpena</i> .....	1847	<i>gymnogaster</i> , <i>Cotylopus</i> .....	2207
<i>guttatus</i> , <i>Astroscopus</i> .....	2310	<i>Sicydium</i> .....	2208
<i>Ballistes</i> .....	1702	<i>Silyopterus</i> .....	2208
<i>Chiropsis</i> .....	1869	<i>Gymnomuraena</i> .....	402, 406
<i>Chirus</i> .....	1868	<i>nectura</i> .....	404
<i>Enneacentrus</i> .....	1142	<i>vittata</i> .....	404
<i>coronatus</i> .....	1142	<i>Gymnonoti</i> .....	339
<i>Epinephelus</i> .....	1142, 1159	<i>Gymnopsis</i> .....	402
<i>Fundulus</i> .....	658	<i>Gymnosarda</i> .....	848
<i>Hippocampus</i> .....	776	<i>alleterata</i> .....	869

Page.		Page.		Page.
1174	<i>Gymnosarda pelamis</i> .....	868	Haddo .....	478
955	<i>Gymnostethus, Prionotus</i> .....	2153	Haddock, Jerusalem .....	954
1269	<i>Gymnothorax</i> .....	392, 400, 401	Haddocks .....	2542
1269	<i>afer</i> .....	395	Hadropterus .....	1028, 1030, 1038
1269	<i>aga e-dulcis</i> .....	391	<i>aspro</i> .....	1032
2419	<i>catenatus</i> .....	403	<i>aurantiacus</i> .....	1041
1269	<i>chlevastes</i> .....	399	<i>cymatotania</i> .....	1042
382	<i>consersus</i> .....	397	<i>evides</i> .....	1036
781	<i>dovii</i> .....	397	<i>guntheri</i> .....	1033
1117	<i>elaboratus</i> .....	389	<i>macrocephalus</i> .....	1031
991	<i>flavoscriptus</i> .....	395	<i>maedatus</i> .....	1031, 1034
1162	<i>funcbris</i> .....	396	<i>nianguae</i> .....	1043
1818	<i>longicauda</i> .....	392	<i>nigrofasciatus</i> .....	1038
2311	<i>marmoratus</i> .....	391	<i>onachita</i> .....	1035
95	<i>miliaris</i> .....	398	<i>peltatus</i> .....	1034
65	<i>mordax</i> .....	396	<i>phoxocephalus</i> .....	1030
119	<i>moringua</i> .....	395	<i>ronoka</i> .....	1036
119	<i>nigrocastaneus</i> .....	390	<i>sciurus</i> .....	1037
119	<i>obscuratus</i> .....	389	<i>serrula</i> .....	1038
383	<i>ocellatus</i> .....	399	<i>shumardi</i> .....	1047
383	<i>nigromar-</i>		<i>squamatus</i> .....	1040
778	<i>ginatus</i> .....	400	<i>tessellatus</i> .....	1070
377	<i>saxicola</i> .....	399	<i>variatus</i> .....	1070
2649	<i>panamopsis</i> .....	391	<i>haematura, Cliola</i> .....	218
2649	<i>pecturatus</i> .....	395	<i>haematurus, Hybopsis</i> .....	218
171	<i>pectus</i> .....	2805	<i>Leuciscus</i> .....	218
171	<i>polygonus</i> .....	394	<i>Haemulide</i> .....	1289
279	<i>rostratus</i> .....	395	<i>Haemulon</i> .....	1291
2793	<i>santa-helene</i> .....	397	<i>acutum</i> .....	1299
2294	<i>scriptus</i> .....	398	<i>albidum</i> .....	1299
2294	<i>umbrosus</i> .....	390	<i>album</i> .....	1295, 1296
2294	<i>verrilli</i> .....	394	<i>arara</i> .....	1306
2456	<i>versipunctatus</i> .....	394	<i>arcuatum</i> .....	1305
2477	<i>vicinus</i> .....	394	<i>aurolineatum</i> .....	1310
2477	<i>virescens</i> .....	391	<i>bonariense</i> .....	1297
2477	<i>Gymnotida</i> .....	340	<i>brevirostrum</i> .....	1300
2477	<i>Gymnotorpedo</i> .....	77	<i>cana</i> .....	1299
997	<i>Gymnotus albus</i> .....	340	<i>canna</i> .....	1297
200	<i>hachirus</i> .....	340	<i>capeuna</i> .....	1311
2010	<i>carapo</i> .....	341	<i>carbonarium</i> .....	1300
2008, 2009	<i>fasciatus</i> .....	340	<i>caudimacula</i> .....	1299, 1302, 1309
2008	<i>putaol</i> .....	41	<i>chromis</i> .....	1299
962	<i>gymnura, Dasyatis</i> .....	84	<i>chrysargyreum</i> .....	1308
1146	<i>Trygon</i> .....	84	<i>chrysopteron</i> .....	1309
781, 1726	<i>gyrans, Querimana</i> .....	818	<i>continuum</i> .....	1297
88	<i>Gyrinichthys</i> .....	2137	<i>curvinaeforme</i> .....	1327
2207	<i>minytrems</i> .....	2137	<i>dorsale</i> .....	1308
2208	<i>gyrinops, Cyclopteroides</i> .....	2102	<i>elegans</i> .....	1304
2208	<i>gyrinus, Eleotris</i> .....	2201	<i>flaviguttatum</i> .....	1312
402, 403	<i>Gobiesox</i> .....	2331	<i>flaviguttatus</i> .....	1312
401	<i>Noturus</i> .....	146	<i>flavolineatum</i> .....	1306
404	<i>Schilbeodes</i> .....	146	<i>formesum</i> .....	1305
339	<i>Silurus</i> .....	146	<i>fremebundum</i> .....	1297
402	<i>Gyroleurodus</i> .....	20	<i>gibbosum</i> .....	1296
818	<i>Gyroleurodus francisci</i> .....	20	<i>heterodon</i> .....	1306
869	<i>quoyi</i> .....	21	<i>bians</i> .....	1394



	Page.		Page.
<i>Hæmulon jeniguano</i> .....	1310	Half-beak, Common.....	721
<i>labridum</i> .....	1319	Halfbeaks.....	719
<i>lateum</i> .....	1304	Half-moon.....	1301
<i>macrostoma</i> .....	1297	Half-naked Gobies.....	2231
<i>macrostomum</i> .....	1296	<i>Halias</i> .....	2502
<i>maculicauda</i> .....	1314	<i>marginatus</i> .....	2502
<i>maculosum</i> .....	1295	Halibut.....	2611
<i>margaritifera</i> .....	1312	Arrow-toothed.....	2609
<i>mazatlanum</i> .....	1314	Bastard.....	2625
<i>melanurum</i> .....	1302, 1303	Greenland.....	2611
<i>microphthalmum</i> .....	1296	Monterey.....	2625
<i>modestum</i> .....	1340	Halibuts, Bastard.....	2624
<i>multilineatum</i> .....	1304	<i>Halicampus</i> .....	761
<i>notatum</i> .....	1297	<i>Halibaeres</i> .....	1587
<i>obtusum</i> .....	1319	<i>bivittatus</i> .....	1597
<i>parra</i> .....	1287	<i>californicus</i> .....	1601
<i>parrie</i> .....	1297, 1309	<i>candialis</i> .....	1600
<i>plumieri</i> .....	1304	<i>dimidiatus</i> .....	1594
<i>quadrilineatum</i> .....	1309, 1311	<i>dispilus</i> .....	1598
<i>quinquelineatum</i> .....	1311	<i>gurnoti</i> .....	1593
<i>retrocurvus</i> .....	1297	<i>maculipinna</i> .....	1595
<i>rimator</i> .....	1309	<i>nicholsi</i> .....	1592
<i>schrauki</i> .....	1302, 1303	<i>poeyi</i> .....	1598
<i>selurus</i> .....	1303	<i>radiatus</i> .....	1591
<i>scudderi</i> .....	1299, 1300	<i>sellifer</i> .....	1592
<i>serratum</i> .....	1299	<i>semicinctus</i> .....	1593
<i>sexfasciatum</i> .....	1294	<i>Halieutea</i> .....	2741
<i>sexfasciatus</i> .....	1295	<i>spongiosa</i> .....	2742
<i>similis</i> .....	1304	<i>Halieutella</i> .....	2742
<i>steindachneri</i> .....	1301	<i>lappa</i> .....	2742
<i>striatum</i> .....	1311	<i>Halientichtlys</i> .....	2739
<i>subarcuatum</i> .....	1306	<i>aculeatus</i> .....	2739
<i>tenuatum</i> .....	1308	<i>caribbeus</i> .....	2741
<i>undecimale</i> .....	1309	<i>reticulatus</i> .....	2741
<i>xanthopteron</i> .....	1307	<i>Halientine</i> .....	2736
<i>xanthopteron</i> .....	1307	<i>Haliperca</i> .....	1203, 1204
<i>Hæmulopsis</i> .....	1325	<i>bivittata</i> .....	1205
<i>Hæmylum capenna</i> .....	1311	<i>fuscula</i> .....	1211
<i>haeres, Gobiesox</i> .....	2337	<i>juvone</i> .....	1215
<i>Hagfish</i> .....	7	<i>phoebe</i> .....	1212
California.....	6	<i>prestigiator</i> .....	1214
<i>Hagfishes</i> .....	5, 7	<i>tabacaria</i> .....	1215
<i>Hairtails</i> .....	889	<i>halleri, Urolophus</i> .....	80
<i>Hake, European</i> .....	2530	<i>Halocypselus</i> .....	729, 2835
New England.....	2530	<i>evolans</i> .....	729
Silver.....	2530	<i>obtusirostris</i> .....	730
White.....	2555	<i>Haloporphyrus viola</i> .....	2544
<i>Hakes</i> .....	2529	<i>Halosauridae</i> .....	606
<i>Halaehurus</i> .....	23	<i>Halosauropsis</i> .....	2826
<i>Halatractus</i> .....	901	<i>Halosaurus</i> .....	607
<i>bonariensis</i> .....	905	<i>goodei</i> .....	610
<i>bosci</i> .....	905	<i>guntheri</i> .....	608
<i>coronatus</i> .....	905	<i>macrochlr</i> .....	610
<i>dorsalis</i> .....	902, 906	<i>oweni</i> .....	607
<i>fasciatus</i> .....	904	<i>rostratus</i> .....	699
<i>halec, Clupea</i> .....	421	<i>hamatus, Icelus</i> .....	1913

Page.	Page.	Page.
721	hamiltoni, Brachioptilon .....	93
719	Hippoglossoides .....	2616
1391	Hamlet .....	395, 1157
2231	hamlini, Podotheicus .....	2056
2592	Hammer-head .....	181
2592	Shark .....	45
2611	hammondi, Percopsis .....	784
2609	Pimelodus .....	135
2625	Semotilus .....	222
2611	Hand-saw Fish .....	596
2625	Hannahill .....	1199
2624	Haplocheilus .....	633, 2827, 2830
761	Haplochilus aureus .....	659
1587	chrysotus .....	656
1597	dovii .....	650
1601	floripinnis .....	651
1600	luciae .....	655
1594	melanopleurus .....	660
1598	melanops .....	682
1594	pulchellus .....	659
1595	sciadicus .....	654
1592	Haplodoci .....	782, 2313
1598	haplognathus, Lepomis .....	1004
1591	Haploidonotus .....	1483
1592	grammens .....	1484
1593	Haplomi .....	622
2741	Harder .....	949
2742	Hardhead .....	497
2742	Hardheads .....	719
2742	Hardmouth .....	208
2739	Hardtail .....	921
2749	Hare-lip Snucker .....	199
2741	harengoides, Glossodon .....	413
2711	Harengula .....	428, 430
2736	arcuata .....	431
1203, 1204	callolepis .....	430
1205	elupeola .....	429, 430
1211	hammeralis .....	431
1215	jaguana .....	430
1212	macrophthalma .....	430
1214	maculosa .....	430
1215	pensacolo .....	431
80	sardina .....	430
729, 2855	harengulus, Encinostomus .....	1368
729	Gerres .....	1369
730	harengus, Clupea .....	421, 422
2544	Coregonus .....	469
606	Lavinia .....	209
2826	Myxus .....	818
607	Querimann .....	817
610	Salmo .....	469
608	harfordi, Ptychocheilus .....	225, 2797
610	Harpe .....	1581
607	caeruleo-aureus .....	1583
499	dilatata .....	1582
1913	oelancheri .....	1583
	Harpe pectoralis .....	1582
	pulchella .....	1584
	pulchra .....	1585
	rufa .....	1583
	Harpinae .....	1572, 1574
	Harpurus .....	1689
	fasciatus .....	1691
	harringtonensis, Atherina .....	791
	harringtoni, Axyrius .....	1904
	Harriotta .....	96
	raleighana .....	96
	Harriotina .....	94
	Harry, Black .....	1199
	Harvest Fish .....	965, 967
	hasselti, Paraserranus .....	1205
	hastata, Dasibatis .....	84
	Dasyatis .....	83
	Trygon .....	84
	hastatus, Gobionellus .....	2229
	Gobius .....	2229
	hastingsi, Neomenis .....	2858
	Haustor .....	135, 136, 137
	havannensis, Murena .....	382
	Ophichthus .....	382
	Uranichthys .....	382
	haydeni, Ptychostomus .....	187
	hayi, Hybognathus .....	214
	Hay-ko .....	478
	Head-fish .....	1753
	Head-fishes .....	1752, 1753
	hearni, Salmo .....	519
	heberi, Scomber .....	923
	Hechdo .....	447
	heckeli, Balistes .....	1705
	Nerophis .....	774
	Syngnathus .....	2839
	Hectoria .....	1138
	heermanni, Amphistichus .....	1504
	Heliasas .....	1545, 1546, 1548
	insolatus .....	1548
	multilineatus .....	1547
	Heliasias .....	1545
	cyanus .....	1547
	Helicolenus .....	1836
	ductylopterus .....	1837
	maderensis .....	1837
	Helioperca .....	999, 1004
	helleri, Cichlasoma .....	1521
	Heros .....	1521
	Xiphophorus .....	701, 702
	Helmichthys .....	353
	Helmictis .....	353
	helolepis, Trachyrincus .....	2568
	Helops .....	193
	helvemaclatus, Sebastodes .....	1828
	Hemidurgan .....	1760

	Page.		Page.
Hemiancistrus .....	159	Hemirhamphus unifasciatus .....	720, 721
<i>aspidolepis</i> .....	159	Hemirhombus .....	2670
<i>gaucharoto</i> .....	159	<i>athalon</i> .....	2673
Hemianthias, <i>peruanus</i> .....	1222	<i>aramaca</i> .....	2673
<i>vivans</i> .....	1223	<i>ambriatus</i> .....	2677
Hemiaris .....	119	<i>fuscus</i> .....	2686
Hemibranchii .....	741	<i>ocellatus</i> .....	2673
Hemibranchs .....	741	<i>ovalis</i> .....	2674
Hemibrycon .....	333	<i>petulus</i> .....	2672
Hemicarax .....	912	<i>soleiformis</i> .....	2672
<i>amblyrhynchus</i> .....	912, 2844	Hemirhamphus .....	722
<i>atrimanus</i> .....	913, 2846	Hemistoma .....	1612
<i>falcatus</i> .....	2845	<i>gnacamaia</i> .....	1659
<i>furthii</i> .....	914	Hemitremia .....	228, 230, 242
<i>leucurus</i> .....	914	<i>bifrenata</i> .....	259
<i>secundus</i> .....	914	<i>heterodon</i> .....	261
<i>zelotes</i> .....	2845	<i>maculata</i> .....	259
Hemihætodon .....	1672	<i>vittata</i> .....	242
Hemigobius .....	2210	Hemitriptera .....	1883
Hemigrammus .....	333	Hemitripterus .....	2022
hemigramma, <i>Garmannia</i> .....	2233	<i>acadianus</i> .....	2023
hemigrammus, <i>Argyroleucus</i> .....	604	<i>americanus</i> .....	2023
<i>Gobius</i> .....	2233	<i>cavifrons</i> .....	2023
Hemilepidotina .....	1880	<i>marmoratus</i> .....	1889, 2022
Hemilepidotus .....	1934	Hemitygon .....	82, 83
<i>gibbsii</i> .....	1936	hemphillii, <i>Stathmonotus</i> .....	2408
<i>hemilepidotus</i> .....	1935	henlei, <i>Carcharhinus</i> .....	37, 2746
<i>jordani</i> .....	1934	<i>Carcharias</i> .....	37
<i>spinosus</i> .....	1937	<i>Rhinotriacis</i> .....	31
<i>tileii</i> .....	1936	<i>Triacis</i> .....	31
hemilepidotus, <i>Cottus</i> .....	1936	henshalli, <i>Fundulus</i> .....	653
<i>Hemilepidotus</i> .....	1935	<i>Zygonectes</i> .....	653, 2829
Hemiodon .....	156, 157	henshavi, <i>Apocope</i> .....	312
Hemiorlicaria .....	156	<i>Rhinichthys</i> .....	312
Hemioplites <i>simulans</i> .....	994	henshawi, <i>Salmo clarkii</i> .....	2819
Hemioplites .....	992	<i>mykiss</i> .....	493
Hemiplus .....	249	hentz, <i>Blennius</i> .....	2390
<i>lacustris</i> .....	250	<i>Hypsoblennius</i> .....	2390
hemiplus, <i>Cyprinus</i> .....	250	hentzi, <i>Hypsoblennius</i> .....	2390
Hemiramphida .....	718	<i>Isestes</i> .....	2390
Hemiramphus .....	722	Hepatus .....	1689
<i>balao</i> .....	723, 2835	hepatus, <i>Acanthurus</i> .....	1692
<i>brasiliensis</i> .....	722	<i>Grammistes</i> .....	1343
<i>browni</i> .....	723	<i>Teuthis</i> .....	1693
Hemirhamphus <i>fasciatus</i> .....	720	heptacanthus, <i>Cheilodipterus</i> .....	947
<i>filamentosus</i> .....	723	heptagonus, <i>Hippocampus</i> .....	775, 777
<i>longirostris</i> .....	724	heptatremide .....	5
<i>macrochirus</i> .....	723	Heptranchias <i>mæculatus</i> .....	18
<i>maerorhynchus</i> .....	724	heraldi, <i>Tetrodon</i> .....	1736
<i>marginatus</i> .....	723	herichthys .....	1526
<i>picarti</i> .....	720	<i>cyanoguttatus</i> .....	1538
<i>pleii</i> .....	723	hermanni, <i>Sternopyx</i> .....	603
<i>poeyi</i> .....	720	hermineri, <i>Clina</i> .....	2362
<i>richardi</i> .....	720	herminder, <i>Blennius</i> .....	2362
<i>roberti</i> .....	721	herminiger, <i>Labrisomus</i> .....	2361
<i>rosei</i> .....	722	Hermosilla .....	1383, 1384

Page.		Page.		Page.
720, 721	<i>Hermosilla azurea</i> .....	1383, 1384	<i>Herpetoleichthys, callisoma</i> .....	384
2670	<i>Heros</i> .....	1526	<i>ocellatus</i> .....	384
2673	<i>affinis</i> .....	1529	<i>sulcatus</i> .....	382
2673	<i>altifrons</i> .....	1538	<i>Herring, Big-eyed</i> .....	410, 426
2677	<i>angulifer</i> .....	1517	<i>Blue</i> .....	425
2686	<i>aureus</i> .....	1533	<i>Branch</i> .....	426
2673	<i>halteatus</i> .....	1522	<i>California</i> .....	422
2674	<i>basilaris</i> .....	1532	<i>Common</i> .....	421
2672	<i>beani</i> .....	1538	<i>Fall</i> .....	425
2672	<i>bifasciatus</i> .....	1521	<i>Glut</i> .....	426
722	<i>centrarchus</i> .....	1526	<i>Great Bear Lake</i> .....	470
1612	<i>citrinellus</i> .....	1534	<i>Lake</i> .....	468
1659	<i>cyanoguttatus</i> .....	1537	<i>Michigan</i> .....	468
228, 230, 242	<i>deppli</i> .....	1524	<i>Mountain</i> .....	463
259	<i>dovii</i> .....	1535	<i>Rainbow</i> .....	524
261	<i>erythreus</i> .....	1531	<i>Round</i> .....	420
259	<i>erythreus</i> .....	1531	<i>Summer</i> .....	426
242	<i>fenestratus</i> .....	1518	<i>Tailor</i> .....	425
1883	<i>friedrichshali</i> .....	1528	<i>Thread</i> .....	432
2022	<i>gibbiceps</i> .....	1536	<i>Toothed</i> .....	413
2023	<i>godmanni</i> .....	1516	<i>Wall-eyed</i> .....	426
2023	<i>holleri</i> .....	1521	<i>Herrings</i> .....	417, 421
2023	<i>intermedius</i> .....	1517	<i>herchelii, Histiothorus</i> .....	892
1889, 2022	<i>irregularis</i> .....	1541	<i>Tetrapturus</i> .....	892
82, 83	<i>labiatus</i> .....	1530	<i>herchelinius, Liparis</i> .....	2123
2408	<i>lentiginosus</i> .....	1524	<i>herzbergii, Arlus</i> .....	125
37, 2716	<i>lobochilus</i> .....	1531	<i>Selenaspis</i> .....	124, 2760
37	<i>longimanus</i> .....	1521	<i>Silurus</i> .....	125
31	<i>macracanthus</i> .....	1519	<i>Tachysurus</i> .....	125
31	<i>maculipinnis</i> .....	1529	<i>Hesperanthias</i> .....	1281
653	<i>managuensis</i> .....	1533	<i>oculatus</i> .....	1283
653, 2829	<i>margaritifer</i> .....	1520	<i>Heterandria</i> .....	686
312	<i>melanopogon</i> .....	1523	<i>affinis</i> .....	681
312	<i>melanurus</i> .....	1524	<i>formosa</i> .....	687, 2831
2819	<i>microphthalmus</i> .....	1536	<i>holbrookii</i> .....	681
493	<i>montezuma</i> .....	1518	<i>metallica</i> .....	687
2390	<i>motaguensis</i> .....	1531	<i>nobilis</i> .....	682
2390	<i>multispinosus</i> .....	1526	<i>occidentalis</i> .....	689, 2833
2390	<i>nicaraguaensis</i> .....	1532	<i>ommata</i> .....	664
1689	<i> nigrofasciatus</i> .....	1525	<i>patruelis</i> .....	681
1692	<i>oblongus</i> .....	1535	<i>pleurospilus</i> .....	688
1343	<i>parma</i> .....	1519	<i>uninotata</i> .....	687
1693	<i>pavonaceus</i> .....	1538	<i>versicolor</i> .....	688, 2833
947	<i>rostratus</i> .....	1523	<i>heteroclitus, Cobitis</i> .....	641
775, 777	<i>salvini</i> .....	1528	<i>heteroclitus, Fundulus</i> .....	640, 2827
5	<i>sieboldii</i> .....	1517	<i>badus</i> .....	2827
18	<i>spilurus</i> .....	1520	<i>grandis</i> .....	641
1736	<i>tetraacanthus</i> .....	1539	<i>macroleptode-</i>	
1526	<i>triagramma</i> .....	1529	<i>tus</i> .....	641
1538	<i>trimaculatus</i> .....	1529	<i>heterodon, Alburnops</i> .....	261
603	<i>troscheli</i> .....	1537	<i>Hemulon</i> .....	1306
2362	<i>urophthalmus</i> .....	1537	<i>Hemitrema</i> .....	261
2362	<i>heros, Eupomotis</i> .....	1007	<i>Hypsopsis</i> .....	261
2361	<i>heros, Lepomis</i> .....	1008	<i>heterodon, Lencisus</i> .....	261
1383, 1381	<i>Pomotis</i> .....	1007	<i>Notropis</i> .....	261
	<i>Herpetoleichthys</i> .....	381	<i>Heterodontidae</i> .....	19

	Page.		Page.
heteroglossus, Gadus .....	2541	Hexanematichthys phrygiatus.....	150
Heterognathi .....	329	rugispinis .....	130
Heterognathus .....	792	seemani .....	128
heterolepis, Jolinus .....	1419	surinamensis ..	129
Notropis .....	260	hians, Athlennes.....	748
Plagiocion .....	1419	Belone.....	718
Seiema .....	1419	Hemulon .....	1304
Heteromi.....	612	Myctophum .....	572
Heteroprosopon .....	2637	Sayris.....	725
heteropygus, Carangops.....	913	Tylosurus .....	718
Caranx.....	913	Hiatula .....	1577
Heterosomata .....	782, 2602	gardeniana.....	1578
Heterostichus .....	2350	hiatula .....	1579
rostratus .....	2351	oultis.....	1579
heterura, Clupea .....	416	hiatula, Hiatula .....	1579
heterurus, Cypselurus .....	2836	Labrus .....	1578
Exocoetus.....	735	Hickory Shad .....	416, 425
Glossodon .....	413	hieroglyphicus, Fundulus.....	658
hexacanthus, Centrarchus .....	987	Zygocetes .....	658
Dipterodon .....	1107	Hilgendorfia .....	2139, 2140, 2141
hexacornis, Cottus .....	2003	hillianus, Exocoetus.....	729
Onecottus .....	2002	Spinax .....	55
Hexagrammidae .....	1863, 2862	Himantolophinae.....	2728
Hexagramminae.....	1864	Himantolophus .....	2732
Hexagrammos .....	1866	greenlandicus.....	2733
asper.....	1872	reinhardti .....	2733
decagrammus .....	1867	himantophorus, Callionymus.....	2186
hexagrammus .....	1872	Himantura .....	82
lagocephalus.....	1873	Hind, Brown.....	1142
stelleri.....	1871	Red.....	1141, 1158
superciliatus.....	1872	Rock .....	1152
Hexagrammus, decagrammus.....	1875	Speckled .....	1159
lagocephalus .....	1875	hinnullus, Squalus.....	29
monopterygius .....	1866	Hiodon .....	412, 413
octogrammus .....	1869	alosoides .....	413
ordinatus.....	1870	clodulus.....	413
otakii.....	1867	selenops.....	414
scaber.....	1873	tergisus.....	413
hexagrammus, Chirus.....	1872	Hiodontidae .....	412
Hexagrammus .....	1872	Hipohemus spilotus.....	1043
Labrax .....	1872	hippe, Balistes.....	1705
Ozorthus.....	2441	Hippocampinae.....	761
Sticheus.....	2441	Hippocampus .....	775
Hexanchidae .....	17	antiquorum .....	776
Hexanchus .....	18	antiquus .....	776
corinus .....	18	brevirostris.....	776
griseus .....	19	fascicularis.....	778
Hexanematichthys.....	119, 121, 128, 2772	gracilis.....	777
assimilis.....	129	guttatus.....	776
caeruleus .....	129	guttulatus .....	778
dasycephalus ..	130	heptagonus.....	775, 777
fells .....	128	hippocampus .....	775
guatemalensis..	129	hudsonius.....	777
hymenorrhinus..	125	ingens.....	776
jordani.....	129	knda .....	778
longicephalus ..	130	levicaudatus.....	777



Page.		Page.		Page.
130	<i>Hippocampus longirostris</i> .....	778	<i>hirundo</i> , <i>Leleocottus</i> .....	2011
130	<i>marginalis</i> .....	778	<i>Hispania</i> , <i>Serrana</i> .....	1488
128	<i>punctulatus</i> .....	777	<i>hispanis</i> , <i>Serrana</i> .....	1489
129	<i>stylifer</i> .....	778	<i>Hispaniscus</i> .....	1765, 1776, 1813
718	<i>zosteræ</i> .....	778	<i>hispanum</i> , <i>Plectropoma</i> .....	1140
718	<i>hippocampus</i> , <i>Hippocampus</i> .....	775	<i>hispanus</i> , <i>Gonioplectrus</i> .....	1140
1304	<i>Syngnathus</i> .....	775	<i>hispidus</i> , <i>Balistes</i> .....	1716
572	<i>Hippoccephalus</i> .....	2033	<i>Cottus</i> .....	2023
725	<i>japonicus</i> .....	2036	<i>Monacanthus</i> .....	1715
718	<i>supercilius</i> .....	2036	<i>Tetrodon</i> .....	1733
1577	<i>Hippoglossina</i> .....	2620	<i>Histiobranchus</i> .....	351
1578	<i>boliviani</i> .....	2621	<i>bathybius</i> .....	352
1579	<i>macrops</i> .....	2621	<i>infernalis</i> .....	352
1579	<i>stomata</i> .....	2620	<i>Histiocottus</i> .....	2018
1579	<i>Hippoglossinae</i> .....	2605	<i>bilobus</i> .....	2018
1578	<i>Hippoglossoides</i> .....	2614	<i>Histiophorus</i> .....	890
416, 425	<i>dentatus</i> .....	2615	<i>americanus</i> .....	891
658	<i>elassodon</i> .....	2615	<i>belone</i> .....	892
658	<i>exilis</i> .....	2613	<i>brevirostris</i> .....	892
2140, 2144	<i>hamiltoni</i> .....	2616	<i>herchelii</i> .....	892
729	<i>jordani</i> .....	2614	<i>pulchellus</i> .....	891
55	<i>linanda</i> .....	2615	<i>Histrio</i> .....	2717
2728	<i>linandoides</i> .....	2615	<i>histrio</i> , <i>Antennarius</i> .....	2716, 2723
2732	<i>melanostictus</i> .....	2618	<i>Eltheostoma</i> ( <i>Ulocentra</i> )....	1051
2733	<i>platessoides</i> .....	2614	<i>Gobiosoma</i> .....	2258
2733	<i>robustus</i> .....	2616	<i>Lophius</i> .....	2716, 2722
2186	<i>hippoglossoides</i> , <i>Platysomatichthys</i>	2611	<i>Petrophryne</i> .....	2716
82	<i>Pleuronectes</i> .....	2611	<i>Scorpena</i> .....	1843, 1846
1142	<i>Reinhardtius</i> .....	2611	<i>Ulocentra</i> .....	1050, 1051
1141, 1158	<i>Hippoglossus</i> .....	2611	<i>Hutch</i> .....	209
1152	<i>americanus</i> .....	2612	<i>hiuleus</i> , <i>Stelephorus</i> .....	443
1159	<i>brasiliensis</i> .....	2626	<i>Hog Choker</i> .....	2700
29	<i>californicus</i> .....	2626	<i>Molly</i> .....	181, 1026
412, 413	<i>gigas</i> .....	2612	<i>Sucker</i> .....	181
413	<i>greenlandicus</i> .....	2611	<i>Hogfish</i> .....	1026, 1338, 1579
413	<i>hippoglossus</i> .....	2611	<i>Spanish</i> .....	1583
414	<i>intermedius</i> .....	2672	<i>Hog-mouth Fry</i> .....	444
413	<i>maximus</i> .....	2612	<i>Holacanthinae</i> .....	2860
412	<i>ocellatus</i> .....	2673	<i>Holacanthus</i> .....	1682, 1729, 2850
1043	<i>pinguis</i> .....	2611	<i>cellaris</i> .....	1685
1705	<i>ponticus</i> .....	2612	<i>clarionensis</i> .....	1683
761	<i>vulgaris</i> .....	2612	<i>cornutus</i> .....	1685
775	<i>hippoglossus</i> , <i>Hippoglossus</i> .....	2611	<i>formosus</i> .....	1685
776	<i>Pleuronectes</i> .....	2612	<i>iodocus</i> .....	1687
776	<i>hippops</i> , <i>Campostoma</i> .....	206	<i>leleothos</i> .....	1735
776	<i>hippos</i> , <i>Carangus</i> .....	921	<i>melanoflus</i> .....	1728
778	<i>Caranx</i> .....	920, 923	<i>passer</i> .....	1682
777	<i>Scomber</i> .....	908, 920	<i>strigatus</i> .....	1683
776	<i>hippuroides</i> , <i>Lepimphid</i> .....	952	<i>tricolor</i> .....	1684
778	<i>hippurus</i> , <i>Coryphaena</i> .....	952	<i>holacanthus</i> , <i>Diodon</i> .....	1746
775, 777	<i>hirudo</i> , <i>Ichthyomyzon</i> .....	11	<i>Ostracion oblongus</i> .....	1746
775	<i>hirundinaceus</i> , <i>Squidus</i> .....	33	<i>Holanthias martinicensis</i> .....	1228
777	<i>Hirundo</i> .....	2183	<i>holholli</i> , <i>Ceratias</i> .....	2729
776	<i>Acanthurus</i> .....	1691	<i>holbrookii</i> , <i>Aelpenser</i> .....	105
778	<i>Azurina</i> .....	1544	<i>Aluterus</i> .....	1718
777	<i>Cottus</i> .....	2011	<i>Echeneis</i> .....	2270

	Page.		Page.
holbrookii, Eupomotis .....	1008	Holocentrus tigrinus .....	1211
Gambusia .....	681	unicolor .....	1192
Heterandria .....	681	vexillarius .....	852
Ophidium .....	2487	Holocephali .....	93
Ophidium .....	2488	holocyanneos, Searus .....	1654
Pomotis .....	1008	Hole'epis barratti .....	1102
holbrookii, Diplodus .....	1362	erocrous .....	1102
Lepomis .....	1008	fusiformis .....	1102
Sargus .....	1363	hololepis, Cyttus .....	1662
Holconoti .....	781, 1493	Zenion .....	1661
Holconotus .....	1502, 1505	holomelas, Paraliparis .....	2140
agassizii .....	1506	Holoporphyrus .....	2543
analis .....	1501	Holorhinus .....	89
fuliginosus .....	1505	vespertilio .....	90
gibbonsii .....	1509	Holostei .....	98, 107
megalops .....	1502	holotrachys, Macrourus .....	2582
rhodoterus .....	1502	Homalogrystes .....	1148
trowbridgii .....	1497	Homalopomus .....	2529
Holia .....	478	trowbridgii .....	2531
hollardi, Hollardia .....	1698	Homesthes .....	2394
Hollardia .....	1697	Homesthes caulopus .....	2394
hollardi .....	1698	homianus, Squalus .....	51
holoacanthus, Diodon .....	1746	homonymus, Diapterus .....	1371
Holocenthrus .....	847	Homoprion .....	1439
Holocentride .....	845	acutirostris .....	1437
Holocentrum .....	847	furthi .....	1441
longipinne .....	849	lanceolatus .....	1444
perlatum .....	853	subtruncatus .....	1434
productum .....	852	xanthurus .....	1434, 1459
prospinosum .....	853	honeymani, Acipenser .....	106
retrospinis .....	853	hoodii, Salmo .....	505
riparium .....	852	hoodii, Salmo .....	507, 510
rostratum .....	852	Hoopid Salmon .....	489
sicciferum .....	850	hopkinsi, Hyunis .....	953
vexillarium .....	852	Myeteroperca .....	2855
Holocentrus .....	847	Plagiogrammus .....	2428
ascensionis .....	848, 2843	Sebastodes .....	1789
rufus .....	849	Hopladelphus .....	142
auratus .....	1145	olivaris .....	143
brachypterus .....	852	Hoplarchus .....	1526
coruscus .....	851	hopliticus, Paricelinus .....	1886
gibbosus .....	1319	hoplomystax, Sparisoma .....	1632
gulo .....	1139	Hoplopagrine .....	1242
marianus .....	852, 2842	Hopiopagrus .....	1244
merou .....	1154	guntheri .....	1244
osculus .....	853	Hoplostethus .....	837
pentacanthus .....	849	japonicus .....	837
punctatus .....	1153	mediterraneus .....	837
rostratus .....	849	Hoplunnis .....	361
sancti-pauli .....	853	diomedianus .....	361
sanguineus .....	1761	schmidtii .....	361
siccifer .....	849	Horned Dace .....	222
sogo .....	849	Pout .....	135, 140
striatus .....	849	Horny Cony .....	1715
suborbitalis .....	850	Hornyhead .....	322
surinameensis .....	1236	Hornyheads .....	314



Page.	Page.	Page.
1211	2172	humeralis, Platyglossus . . . . . 1507
1192	388	Sardinella . . . . . 431
852	920	Searus . . . . . 1641
94	870, 909	Serranus . . . . . 1197
1654	923	humeri-maculatus, Sargus . . . . . 1360
1102	934	humile, Pristipoma . . . . . 1331
1102	936	humilis, Bryttus . . . . . 1004
1102	814	Cottus . . . . . 1979
1692	105	Eupomotis . . . . . 1004
1661	135	Gambusia . . . . . 682
2110	29	Grammateus . . . . . 1355
2513	715, 716	Lepomis . . . . . 1004
89	2236	Pagellus . . . . . 1355
90	469, 472	Pomadasis . . . . . 1331
98, 107	468, 470	Tetragonopterus . . . . . 335
2582	141	Humpback Salmon . . . . . 478
1118	1969	Sucker . . . . . 184
2529	2641	Whitefish . . . . . 466
2531	269	hunting, Molva . . . . . 2551
2394	269	Huro . . . . . 1010
2394	507	nigricans . . . . . 1012
51	254, 256, 266	huronensis, Lepisosteus . . . . . 110
1371	322	Huso . . . . . 103
1439	270	hyalinus, Ceratichthys . . . . . 321
1437	270	hyalope, Squalius . . . . . 222
1411	269	Hybognathus . . . . . 212, 213
1411	212	amara . . . . . 215
1331	176	aeneus . . . . . 215
1434, 1159	777	argyritis . . . . . 214
106	269	civilis . . . . . 215
505	269	episcopa . . . . . 214
507, 510	269	episcopus . . . . . 215
480	270	evansi . . . . . 213
953	270	flavipinnis . . . . . 215
2855	269	fluvialtilis . . . . . 215
2428	793	hayi . . . . . 214
1789	341	melanops . . . . . 216
142	572	nigrotentata . . . . . 214
143	236	nubila . . . . . 215
1526	571	nuchalis . . . . . 213
1886	572, 577	osmerinus . . . . . 213
1632	237	perspicuus . . . . . 218
1242	341	placitus . . . . . 213
1244	237	plumbea . . . . . 216
1214	793	proene . . . . . 204
837	793, 2839	punctifer . . . . . 215
837	2341	regius . . . . . 213
837	1597	serena . . . . . 214
361	1074	stramineus . . . . . 262
361	431	volucellus . . . . . 263
361	2341	Hybopsis . . . . . 314, 315, 319
222	431	festivialis . . . . . 316
135, 140	1596	marconis . . . . . 316
1715	1459	altus . . . . . 321
322	1097	amblops . . . . . 320
314	1196, 1197	bifrenatus . . . . . 259







**IMAGE EVALUATION  
TEST TARGET (MT-3)**



**Photographic  
Sciences  
Corporation**

23 WEST MAIN STREET  
WEBSTER, N.Y. 14580  
(716) 872-4503

15 128  
16 132  
17 122  
18 120

19 110  
20 100

	Page.		Page.
Hybopsis bivittatus .....	233	Hydrargyra multifasciata .....	645
chalybeus .....	288	nigrofasciata .....	641
chiliticus .....	287	ornata .....	2827
chlorocephalus .....	286	similis .....	639
chrosomus .....	288	swampina .....	641, 645
cumingii .....	318	trifasciata .....	639
dissimilis .....	318	vernalis .....	639
dorsalis .....	262	zebra .....	647
fretensis .....	261	Hydrolagus .....	95
gelidus .....	316, 317	collei .....	95
gracilis .....	321	Hydrophlox .....	254, 257, 284
hamaturus .....	218	chrosomus .....	288
heterodon .....	261	Intipinnis .....	287
hudsonius .....	269	rubricroceus .....	286
hyostomus .....	316	Hydrophlox, Clinostomus .....	238
hypsinotus .....	320	Leuciscus .....	238
kentuckiensis .....	322	Squalus .....	238
labrosus .....	319	hygomii, Myctophium .....	573
lacertosus .....	284	Scopelus .....	573
longiceps .....	264	Hylomyzon .....	173
meeki .....	317	nigricans .....	181
missuriensis .....	262	Hymenocephalus .....	2580
monacus .....	318	cavernosus .....	2580
montanus .....	317	goodii .....	2572
niveus .....	278	longifilis .....	2567
phaenna .....	270	Hymenoptera .....	2866
procne .....	264	hymenorrhinus, Hexanematicithys ..	125
rubricroceus .....	286	Hynnus .....	932
rubrifrons .....	320	cubensis .....	932
scylla .....	263	hopkinsi .....	933
spectrunculus .....	265	Hyodon amphiodon .....	413
storerianus .....	270, 321	chrysoptis .....	413
stramineus .....	262	claudalus .....	413
tetraneus .....	315	vernalis .....	413
timpanogensis .....	223	Hyoganoidea .....	98
tuditanus .....	253	Hyostoma .....	1052
volucellus .....	263	blemmlopera .....	1053
watauga .....	319	newmani .....	1053
winchelli .....	321	sluoterum .....	1051
xenocephalus .....	289	hyostomus, Hybopsis .....	316
Hyborhynchus .....	217	Nocomis .....	316
confertus .....	217, 218	hypacanthus, Psednobleinius .....	240.6
nigellus .....	217	Hypargyrus .....	252
notatus .....	218	tuditanus .....	253
pauiceus .....	218	Hypenens .....	858
siderius .....	314	Hyperistius .....	986
supercilliosus .....	218	carolinensis .....	988
tenellus .....	218	Hyperoartil .....	8
Hydrargira .....	632	Hyperotreti .....	5
atricauda .....	624	Hyperprosope .....	1501
Hydrargyra diaphana .....	645	agassizi .....	1502
formosa .....	2827	annalis .....	1501
fusca .....	624	arcuatum .....	1502
limi .....	624	arcuatus .....	1502
lucie .....	655	argenteum .....	1502
majalis .....	630	puncta- tum .....	1502

Page.		Page.		Page.
645	Hyperprosopon argentens	1501, 1502	Hypoplectrus unicolor primivarinus	1192
641	Hypilepis cornutus cerasinus	283	puella	1192
2827	Hypentelium	173, 174, 181	vitulinus	1192
639	macropteron	181	hypoplectrus, Bathymaster	2290
641, 645	Hyperchæristus	589	Hypoplites	1247
639	tanneri	589	Hypoprion	41
647	Hyphalonedrus	541	brevirostris	41
95	chalybeius	542	longirostris	41
95	Hyplenrochilus	2385	signatus	41
254, 257, 284	geminatus	2385	Hypoprionodon	41
288	multifilis	2385	Hyporhamphus	719
287	punctatus	2390	roberti	721
286	Hypocaranx	927	rosæ	721
238	Hypoclydonia	1115	trienspidatus	720
238	bella	1115	unifasciatus	720
238	Hypocritichthys	1500	Hyporthodus	1148
573	analis	1500, 1501	flavicanda	1156
573	Hypodis	915, 2848	Hyposerranus	1148
173	glancus	2848	Hypostominae	173
181	hypodus, Decapterus	908	Hypostomus gaucharoto	159
2580	Hypogymnæobius	2210	hypostomus, Aodon	92
2580	Hypohortulus	1039, 1040	Cephalopterus	92
2572	aurantiacus	1040	Hypsagonus	2038
2567	cymatotenias	1041	gradiens	2041
2866	niangne	1042	quadriceornis	2038, 2041
125	sqnamatus	1040	swanii	2038
932	Hypomesus	524	hypsoleptera, Chiola	280
932	olidus	525, 2824	hypsolepterus, Lenciscus	280
933	pretiosus	525	Notropis	280
413	hypophthalmus, Arius	133	hypselurus, Pimelodus	152
413	Cathorops	133, 2798	Rhamdia	152
413	hypoplecta, Rathibunelia	2290	Hypsicometes	2293
413	Hypoplectrus	1187	goboides	2294
98	accensus	1193	Hypsitaro	474, 477, 481
1052	affinis	1193	kennerlyi	481
1053	bovinus	1193	Hypsilepis ardens	301
1053	chlorurus	1193	coccogenis	285
1051	crocotus	1192	cornutus	283
316	gemma	1193	cyanus	283
316	gummigutta	1192	gibbus	283
2416	guttavarius	1192	diplamla	300
252	indigo	1193	galacturus	279
253	lamprurus	1190	iris	272
858	maculiferus	1192	kentuckiensis	279
986	puella	1192	Hypsinotus	1664
988	unicolor	1190, 1192	rubescens	1965
5	aberrans	1193	hypsinotus, Ceratichthys	320
5	accensus	1193	Hybopsis	320
1501	affinis	1193	Hypsoblemnus	2386
1502	bovinus	1193	brovipinnis	2390
1501	chlorurus	1192	gentilis	2387
1502	crocotus	1192	gilberti	2386
1502	gummigutta	1192	hantz	2390
1502	guttavarius	1192	hentzi	2390
1502	indigo	1193	lonthas	2388
1502	nlgricans	1193	striatus	2388, 2392

	Page.		Page.
Hypsoiepis .....	254	Icichthys lockingtonii .....	969
Hypsopsetta .....	2639	icistia, Bairdiella .....	1435
Hypsurus .....	1508	Sciama .....	1436
caryi .....	1508, 1509	Icosteidae .....	968, 2849
Hypsypops .....	1564	Icosteinae .....	969
dorsalis .....	1570	Icosteus .....	972
rubicundus .....	1564, 1565	enigmaticus .....	972
Hysteroecarpinae .....	1494	Ictalops, Cottus .....	1950
Hysteroecarpus .....	1495	Pegedictis .....	1951
traski .....	1496	Ictalurinae .....	115
hystrix, Diodon .....	1744, 1746	Ictalurus .....	131
Icelinus .....	1894	albidus .....	138
borealis .....	1896	angulla .....	2788
cavifrons .....	1892	furcatus .....	134, 2788
filamentosus .....	1893	kevincii .....	138
fimbriatus .....	1894	lacustris .....	137
oculatus .....	1895	lophius .....	138
quadriseriatus .....	1897	lupus .....	137
strabo .....	1897, 2862	macaskeyi .....	138
tenuis .....	1894	meridionalis .....	135
Icelus .....	1911	nigricans .....	137
australis .....	1918	niveiventris .....	138
bicornis .....	1911	okeechobeensis .....	139
canaliculatus .....	1917	ponderosus .....	137
euryops .....	1915, 1916	punctatus .....	134
furciger .....	1913	robustus .....	135
hamatus .....	1913	simpsoni .....	135
megacephalus .....	1891	Icthyophis .....	403
pugetensis .....	1891	vittatus .....	404
quadriseriatus .....	1897	Ictiobinae .....	162
sentiger .....	1910	Ictiobus .....	163, 164
spiniger .....	1914	bubalus .....	164
uncinatus .....	1906	cyprinella .....	163
vicinalis .....	1916	meridionalis .....	164
Icthalurus punctatus .....	135	urus .....	164
Ictholis .....	999	velifer .....	167
erythrops .....	990	Idiacanthidae .....	604
megalotis .....	1003	Idiacanthus .....	605
melanops .....	996	antrostomus .....	605
Ictheloides, Lepomis .....	990	ferox .....	605
Icthyapus .....	374	Idol, Moorish .....	1687
selachops .....	374	Ilietis .....	142
Icthyacallus .....	1587	Ilisha .....	435
dimidiatus .....	1594	bleekeriana .....	436
Icthyobus .....	163	flavipinnis .....	435
bubalus .....	164	furthi .....	436
cyanellus .....	164	panamensis .....	436, 2811
Icthyomyzon .....	10	Illecebrosus, Alburnops .....	269
argenteus .....	11	Notropis .....	268
astori .....	12	Stellifer .....	1442
castaneus .....	11	Ilyophidae .....	349
Icthyomyzon concolor .....	11	Ilyophis .....	349
hirudo .....	11	brunneus .....	350
tridentatus .....	12	Ilypnus .....	2253
Icichthys .....	969	gilberti .....	2253
		Imberbe, Peristedion .....	2182

Page.		Page.		Page.
969	imberbi, Ophidium .....	2443	Inermia vittata .....	1366
1435	imberbis, Apogon .....	1107	inermis, Anoplagonus .....	2094
1436	Mullus .....	1107	Aspidophoroides .....	2093
968, 2849	Sciæna .....	1454	Dermatolepis .....	1167
969	Vulsiculus .....	2181	Elops .....	410
972	imiceps, Ophiocion .....	1451	Epinephelus .....	1168
972	Sciæna .....	1451	Lioperca .....	1168
1950	immaculata, Amia .....	411	Lutjanus .....	1275
1951	Coryphæna .....	953	Mesoprion .....	1275
115	Fistularia .....	758	Ostracion .....	1723
133	Percæ .....	1135	Rabirubia .....	1274
138	Unibranchapertura .....	342	Raia inornata .....	73
2788	immaculatum, Cybium .....	876	Scorpena .....	1853
134, 2788	immaculatus, Esox .....	630	Sebastes .....	1820
138	masquinongy .....	630	Sebastodes .....	1829
137	Lincius masquinongy .....	530	Serranus .....	1168
138	Salmo .....	507	infans, Avocettina .....	367
137	Symbranchus .....	342	Gambusia .....	680
138	Imostoma .....	1044, 1046	Nemichthys .....	368
135	shumardi .....	1047	infernalis, Histiobranchus .....	352
137	Imperator, Tetrapterus .....	892	Muraena .....	396
138	Xiphias .....	892	Synaphobranchus .....	352
139	Imperial, Serran .....	1837	infimus, Novaculichthys .....	1616
137	imperialis, Coryphæna .....	952	Xyrichtys .....	1616
134	Esox .....	717	ingens, Gunnellus .....	2410
135	Sebastes .....	1837, 1838	Hippocampus .....	776
135	Trachurus .....	927	Intistius .....	1619
403	Zeus .....	955	mundicarpus .....	1620
404	impetiginosus, Serranus .....	1153	Iniomi .....	530
162	impressus, Conger .....	356	Innominado .....	382
163, 164	inequilabius, Carapus .....	341	innominatus, Girardinichthys .....	666
164	inequilobus, Leucosomus .....	224	inopa, Antennarius .....	2718
163	Pogonichthys .....	224	Inosetta .....	2641
164	incilia, Mugil .....	813	ischyra .....	2641
164	incisor, Pimelepterus .....	1386	inornata, Lota .....	2551
167	Pomotis .....	1005	Raja .....	73
604	Inconnu .....	473, 474	Raja .....	73
605	inconstans, Eucalla .....	744	inornatus, Apodichthys .....	2412
605	cayuga .....	744	Chorinemus .....	899
605	pygmaea .....	744	Cryptacanthodes .....	2443
1687	Gasterosteus .....	744	Microlepidotus .....	1341
142	incrassatus, Leucosomus .....	222	Oligoplites .....	899
435	incurus, Lobotes .....	1236	Orthopristsis .....	1342
436	indefatigabile, Otophidium .....	2490	Pseudojullia .....	1604
435	Indian Chub .....	322	inscripta, Solea .....	2696
436	Fish .....	1080	inscriptum, Etheostoma .....	1072
436, 2811	indicus, Chanos .....	415	Nanostoma .....	1072
269	Cottus .....	2092	inscriptus, Achirus .....	2696
268	Cubiceps .....	951	Notbonotus .....	1072
1442	Nauarates .....	900	Pomotis .....	1003
349	Tetrapterus .....	892	inseulpta, Netuma .....	127, 2705
349	indigena, Clupea .....	428	inseulptus, Arins .....	127
350	indigo, Hypoplectrus .....	1193	inseulptus, Gasterosteus .....	750
2253	unicolor .....	1193	Luelocharax .....	339
2253	Plectropoma .....	1193	insigne, Pimelodus .....	147
2182	Inermia .....	1365	insignis, Catostomus .....	180

	Page.		Page.
insignis, <i>Noturus</i> .....	147	<i>Ioglossus</i> .....	2192
<i>Schilbeodes</i> .....	147	<i>calliurus</i> .....	2193
insociabilis, <i>Dorosoma</i> .....	416	ionthas, <i>Cerdale</i> .....	2449
insolatus, <i>Chromis</i> .....	1548	<i>Ilypsoblennius</i> .....	2388
<i>Ileliases</i> .....	1548	<i>Isestes</i> .....	2389
insula-sanctae-erueis, <i>Searus</i> .....	1651	ios, <i>Clevelandia</i> .....	2254
insularum, <i>Atherina</i> .....	807	<i>Gobiosoma</i> .....	2255
<i>Atherinops</i> .....	807	<i>Iotichthys</i> .....	228, 231, 243
<i>Muraena</i> .....	400	<i>iowa</i> , <i>Etheostoma</i> .....	1083
<i>Netuma</i> .....	2770	<i>Ipnopidae</i> .....	546
integripinnis, <i>Auchenopterus</i> .....	2372	<i>Ipnops</i> .....	546
<i>Cremonobates</i> .....	2372	<i>murrayi</i> .....	547
intermedia, <i>Tigoma</i> .....	235	<i>irideus</i> , <i>Labrus</i> .....	988
intermedium, <i>Cichlasoma</i> .....	1517	<i>Salmo</i> .....	500
intermedius, <i>Gasterosteus</i> .....	750	<i>agna-bonita</i> .....	503
<i>Heros</i> .....	1517	<i>gilberti</i> .....	502
<i>Hippoglossus</i> .....	2672	<i>nasoni</i> .....	501
<i>Leuciscus</i> .....	235	<i>shasta</i> .....	502
<i>Paralepis</i> .....	600	<i>stonei</i> .....	503
<i>Pomoxys</i> .....	987	<i>iridinus</i> , <i>Carangoides</i> .....	919
<i>Saurus</i> .....	535	<i>Caranx</i> .....	919
<i>Squalius</i> .....	235	<i>Iridio</i> .....	1587
<i>Sudis</i> .....	600	<i>bivattatus</i> .....	1595
<i>Synodus</i> .....	535, 536	<i>caudalis</i> .....	1599
internasalis, <i>CherogJulis</i> .....	1594	<i>cyanocephalus</i> .....	1594
<i>Julis</i> .....	1594	<i>dispilus</i> .....	1597
<i>Platyglossus</i> .....	1594	<i>garnoti</i> .....	1593
interrupta, <i>Morone</i> .....	1134	<i>kirschii</i> .....	1598
<i>Perea mitchilli</i> .....	1133	<i>maculipinna</i> .....	1591
<i>Raja</i> .....	2751	<i>nicholsi</i> .....	1591
interruptus, <i>Ambloplites</i> .....	991	<i>pictus</i> .....	1599
<i>Anisotremus</i> .....	1319	<i>pocoyi</i> .....	1599
<i>surinam-</i> <i>ensis</i> .....	1319	<i>radiatus</i> .....	1596
<i>Archoplites</i> .....	991	<i>sellifer</i> .....	1592
<i>Centrarchus</i> .....	991	<i>semicinctus</i> .....	1592
<i>Gentremus</i> .....	1819	<i>iris</i> , <i>Chiola</i> .....	272
<i>Luxilus</i> .....	282	<i>Ilypsilepis</i> .....	272
interstitialis, <i>Epinephelus</i> .....	1179	<i>Leuciscus</i> .....	222
<i>Mycteroperca</i> .....	1178	<i>Irish Lord</i> .....	1934
<i>Serranus</i> .....	1179	<i>Pompano</i> .....	1376
<i>Trisotropis</i> .....	1179	<i>irradians</i> , <i>Serranus</i> .....	1268
intertinctus, <i>Mystriophis</i> .....	386	<i>irregularis</i> , <i>Heros</i> .....	1541
<i>Ophichthys</i> .....	387	<i>Theraps</i> .....	1540
<i>Ophisura</i> .....	387	<i>irretitus</i> , <i>Gordichthys</i> .....	363
introniger, <i>Sebastichthys</i> .....	1805	<i>Mugil</i> .....	819
<i>Sebastodes</i> .....	1805	<i>irroratus</i> , <i>Monacanthus</i> .....	1713
intronigra, <i>Dicrolene</i> .....	2522	<i>isabelita</i> .....	1684
inurus, <i>Zygonectes</i> .....	682	<i>isabelita</i> , <i>Angellchthys</i> .....	1685
ioa .....	1064	<i>isabelito de la Alto</i> .....	1674
<i>vigil</i> .....	1065	<i>Isaciella</i> .....	1340
<i>vitrea</i> .....	1064	<i>brevipinnis</i> .....	1341
ioa, <i>Etheostoma</i> .....	1084	<i>ischannus</i> , <i>Notemigonus</i> .....	251
iolocus, <i>Angellchthys</i> .....	1686	<i>Stolephorus</i> .....	442
<i>Holacanthus</i> .....	1687	<i>ischinagi</i> , <i>Megaperca</i> .....	1138
		<i>ischyra</i> , <i>Inopsetta</i> .....	2641



Page.		Page.		Page.
2192	ischyrus, Apomotis .....	997	Jack Goggle-eye .....	911
2193	Lepiopomus .....	997	Horse-eye .....	923
2449	Lepocottus .....	997	Yellow .....	919
2388	Parophrys .....	2641	Jack Salmon .....	1021
2389	Pleuronectes .....	2641	Jacket, Leather .....	1701
2254	Isesthes .....	2386	Jackets, Leather .....	898
2255	gentilis .....	2388	Jacksoni, Ditrema .....	1505
8, 231, 243	gilberti .....	2387	Embiotoca .....	1504, 1505
1083	hontzi .....	2390	Jacob Evertzens .....	1143
546	fontinas .....	2389	Jacobae, Echeueis .....	2272
546	punctatus .....	2390	Remora .....	2272
547	scrutator .....	2389	Jacobi, Corvius (Johnius) .....	1457
988	striatus .....	2388	Sciæna .....	1457
500	islandicus, Centronotus .....	2438	Jacobus, Myrripristis .....	846
503	Gasterosteus .....	748	Jacome .....	1215
502	Gunnellus .....	2439	Jacome, Haliperca .....	1215
501	Stichæus .....	2439	Serranus .....	1215
502	isodon, Aprionodon .....	42	jaculidens, Ancyodon .....	1416
503	Carcharias .....	42	jaculus, Alburnellus .....	293
919	Mesoprion .....	1267	Jaguacaguaro .....	1502
919	isotus, Squalus .....	51	jaguana, Harengula .....	430
1587	Isogomphodon .....	33, 35, 40	jaguar, Bodianus .....	849
1595	limbatus .....	40	Jullao .....	1295
1599	maculipinnis .....	40	jamaicensis, Cynoscion .....	1406
1594	isolepis, Isopsetta .....	2642	Otolithus .....	1406
1597	Lepidopsetta .....	2642	Raja .....	81
1593	Parophrys .....	2642	Urolophus .....	81
1598	Sternotremula .....	787	Janeiro, Engraulis .....	451
1591	isopisthus .....	1399	Janissary .....	1586
1591	affinis .....	1399	Januaria, Umbrina .....	1474
1599	parvipinnis .....	1399	Jaok, Cottus .....	1978
1599	renifer .....	1399	Myoxocephalus .....	1977
1596	Isopsetta .....	2642	japonensis, Salmo .....	479
1592	isolepis .....	2642	japonica, Squatina .....	59
1592	Isospondyli .....	407	japonicus, Agonus .....	2036
272	Isospondylous Fishes .....	407	Aretoscopus .....	2297
272	isthmensis, Rivulus .....	2830	Chirolophus .....	2409
222	Istiphorida .....	890	Cottus .....	2036
1934	Istiophorus .....	890	Hippocephalus .....	2036
1376	nigricans .....	891	Hoplotelus .....	837
1208	Isuropsis .....	47, 48	Pereis .....	2034
1541	dekayi .....	48	Phalangistes .....	2036
1540	glauca .....	48	Physiculus .....	2549
363	Isurus .....	47, 48	Trichodon .....	2297
819	dekayi .....	48	Jaqueta .....	1561
1713	oxyrinchus .....	48, 49	Jaquette, Petitio .....	1559
1684	spallanzani .....	49	Jarrovi, Lepidomeda .....	328
1685	Italara .....	1142, 1162	Minomus .....	170
1674	itaiara, Promlerops .....	1164	Pantosteus .....	171
1340	Serranus .....	1164	Javanicus, Psenes .....	951
1341	Jabon .....	1232	Jaw-fishes .....	2279
251	Jahoncello .....	1232	Jejuna, Episema .....	290
442	Jack .....	627, 920, 1780	Jejunus, Minnilus .....	290
1138	Amber .....	903	Notropis .....	290
2641			jemezans, Alburnellus .....	294

	Page.		Page.
jemezianus, Minnilus.....	294	Jordanella.....	677
Jeniguana.....	1302	<i>floridae</i> .....	677
Jenignano.....	1310	Jordanii, Bathymaster.....	2289
Jenignauo, Hiemulon.....	1310	<i>Canlophryno</i> .....	2735
Jenkinsi, Chasmodes.....	2391, 2392	<i>Chirostoma</i> .....	793
<i>Fundulus</i> .....	651	<i>Eopsetta</i> .....	2613
<i>Synodus</i> .....	537, 2826	<i>Epinephelus</i> .....	1177
<i>Zygonectes</i> .....	652	<i>Esloparum</i> .....	2840
Jenkinsia.....	418	<i>Etheostoma</i> .....	1079, 1080
<i>acuminata</i> .....	419	<i>Guleichthys</i> .....	2774
<i>lamprotenia</i> .....	419	<i>Hemilepidotus</i> .....	1934
<i>stolifera</i> .....	419	<i>Hexanematichthys</i> .....	129
Jenny, Silver.....	1370	<i>Hippoglossoides</i> .....	2614
Jerker.....	222	<i>Mycteroperca</i> .....	1176
Jerusalem Haddock.....	954	<i>Neomenis</i> .....	1251
Jessie, Etheostoma.....	1094	<i>Notropis</i> .....	259
<i>Pencilichthys</i> .....	1085	<i>Pantosteus</i> .....	171
<i>Xenocys</i> .....	1285	<i>Raia</i> .....	73
<i>Nyriichthys</i> .....	1613	<i>Ronquilus</i> .....	2289
<i>Xyrula</i> .....	1612, 1613	<i>Sebastes</i> .....	1778
Jewfish, Black.....	1161	<i>Tachisurus</i> .....	129
<i>California</i> .....	1137	Jordania.....	1884
Jewfishes.....	1137	<i>zonope</i> .....	1884
Jewsharp Drummer.....	1473	Jordaniine.....	1880
Jignagua.....	920	Jorobado.....	934, 936
Jocu.....	1257	Josephi, Ophidion.....	2488
Jocu, Anthias.....	1258	Joturo.....	821
<i>Lutjanus</i> .....	1258	Joturus.....	820
<i>Mesoprion</i> .....	1258	<i>plchardi</i> .....	821
<i>Neomenus</i> .....	1257	<i>stipes</i> .....	821
John A. Grindle.....	113	Joyneri, Sebastes.....	1829
<i>Dorles</i> .....	1659	Juba, Guatucupa.....	1323
<i>Mariggle</i> .....	410	<i>Perca</i> .....	1323
<i>Paw</i> .....	1159	Jug-fish.....	1728
Johni, Cephaloptera.....	93	Jugalis, Cliola.....	272
Johnius.....	1455	<i>Moriana</i> .....	272
<i>amazonicus</i> .....	1419	Juglar Fishes.....	2528
<i>latabanus</i> .....	1431	Julie, Etheostoma.....	1093
<i>crouvina</i> .....	1419	Julidine.....	1572, 1574
<i>guttatus</i> .....	1174	Julidio.....	1602
<i>heterolepis</i> .....	1419	<i>adustus</i> .....	1602
<i>jacobi</i> .....	1457	<i>notospilus</i> .....	1603
<i>nobilis</i> .....	1413	Julis.....	2859
<i>ocellatus</i> .....	1454	<i>bifasciata</i> .....	1610
<i>regalis</i> .....	1407	<i>bifasciatus</i> .....	1610
<i>saxatilis</i> .....	1475	<i>caudalis</i> .....	1599
Johnny.....	2013	<i>cinctus</i> .....	1593
<i>Darter</i> .....	1056	<i>crotaphus</i> .....	1591, 1598
<i>Verlo</i> .....	1195	<i>cyanostigma</i> .....	1591
Jolt-head Porgy.....	1352	<i>detersor</i> .....	1610
Jonesi, Belone.....	717	<i>dimidiatus</i> .....	1594
<i>Gerres</i> .....	1368	<i>gamotl.</i> .....	1593
<i>Mollienisia</i> .....	699	<i>gillians</i> .....	1610
<i>Siphostoma</i> .....	768	<i>humeralis</i> .....	1596
<i>Syngnathus</i> .....	768	<i>internasalis</i> .....	1594
Jopaton.....	1341	<i>lucasanus</i> .....	1607, 1608

Page.		Page.		Page.
677	<i>Julis maculipinna</i> .....	1595	Killer, Salmon .....	749
677	<i>melanochr</i> .....	1609	Killifish.....	639, 641
2289	<i>pedunculatus</i> .....	1601	Common.....	640
2735	<i>nitida</i> .....	1608	killifish, <i>Cobitis</i> .....	641
793	<i>nitidissima</i> .....	1608	Killifishes.....	630, 632
2613	<i>opalina</i> .....	1591	killinensis, <i>Salmo</i> .....	509
1177	<i>patatus</i> .....	1591	King of the Mackerels.....	1755
2840	<i>pletus</i> .....	1000	Mulletts.....	1106
1079, 1080	<i>principis</i> .....	1591	Salmon.....	470
2774	<i>psittaculus</i> .....	1597	Kingfish.....	875, 1460, 1469, 1475
1934	<i>semicinctus</i> .....	1503	<i>kirschii</i> , <i>Iridio</i> .....	1598
129	Jumping Mullet.....	197	<i>kirtlandi</i> , <i>Acipenser</i> .....	106
2614	Jump-rocks.....	197	<i>Kirilandlu</i> .....	794
1176	June Sucker of Utah Lake.....	183	<i>laciniata</i> .....	795, 2840
1251	Jurel.....	899, 921, 923	<i>martinea</i> .....	795
259	<i>Jurvucapeba</i> .....	1142	<i>vagrans</i> .....	794, 2840
171			<i>laciniata</i> .....	2840
73	Kalog.....	1976	<i>Kisutch</i> .....	480
2289	Kamchatka Salmon Trout.....	2818	<i>kisutch</i> , <i>Oncorhynchus</i> .....	480
1778	kamloops, <i>Salmo gairdneri</i> .....	499	<i>Salmo</i> .....	481
129	Kamloops Trout.....	499	kitt, <i>Microstomus</i> .....	2654
1884	kanawa, <i>Notropis</i> .....	264	<i>Pleuronectes</i> .....	2654
1884	kansae, <i>Fundulus</i> .....	2828	kiamathensis, <i>Cottus</i> .....	1955
1880	karrak, <i>Anarrhichas</i> .....	2446	kleinii, <i>Balistes</i> .....	1720
934, 936	Kathetostoma.....	2311	klauzingeri, <i>Achirus</i> .....	2697
2488	<i>albigutta</i> .....	2312	<i>Solea</i> .....	2697
821	<i>averruncus</i> .....	2311	kneri, <i>Pristopoma</i> .....	1338
820	Kathetostomatinae.....	2306	Kodiak Smeit.....	2823
821	kaupi, <i>Physiculus</i> .....	2548	kœlreuteri, <i>Scomber</i> .....	900
821	kaupii, <i>Synaphobranchus</i> .....	351	Koguneso.....	1833
1829	Kolpfish.....	1592, 2351, 2352	Kowala.....	428, 2811
1323	<i>Spofted</i> .....	2353	Krasnaya Ryba.....	481
1323	kendalli, <i>Spiagebranchus</i> .....	375	kraussi, <i>Gobius</i> .....	2228
1728	<i>Verma</i> .....	375	Krohnus.....	2587
272	kennedyi, <i>Trachinotus</i> .....	942	kroyeri, <i>Scopelus</i> .....	556
272	keenerlyi, <i>Hypsifario</i> .....	483	kuda, <i>Hippocampus</i> .....	778
2528	<i>Moxostoma</i> .....	186	Kuhliu.....	1013
1083	<i>Oncorhynchus nerka</i> .....	483	<i>arge</i> .....	1014
1572, 1574	<i>Salmo</i> .....	483	<i>xenura</i> .....	1015
1602	kenicotti, <i>Acipenser</i> .....	105	Kuhlidae.....	1013
1602	<i>Catonotus</i> .....	1098	kumlien, <i>Uranidea</i> .....	1967
1603	<i>Coregonus</i> .....	464	kundscha, <i>Salmo</i> .....	2823
2859	Kenoza.....	625, 626	<i>Salvelinus</i> .....	2822
1610	kentuckiensis, <i>Hybopsis</i> .....	322	Kuro Sol.....	1834
1610	<i>Hypsilepis</i> .....	279	Kyach.....	426
1599	<i>Leuciscus</i> .....	279	Kyphoside.....	1380
1593	<i>Luxilus</i> .....	279, 322	Kyphosine.....	1381
1591, 1598	Kern River Trout.....	502	Kyphosus.....	1384
1591	Keshimingo.....	1833	<i>analogus</i> .....	1385
1610	kessleri, <i>Arins</i> .....	127	<i>elegans</i> .....	1387
1594	<i>Netuma</i> .....	127, 2765	<i>incisor</i> .....	1386
1593	<i>Tachisurus</i> .....	127	<i>lutescens</i> .....	1388
1610	keta, <i>Oncorhynchus</i> .....	478	<i>ocyrus</i> .....	1390
1596	keta vel kayko, <i>Salmo</i> .....	479	<i>sectatrix</i> .....	1387
1594	kevinskii, <i>Ictalurus</i> .....	138		
607, 1608	Kleye of Lake Michigan.....	469	Labeo elegans.....	186

	Page.		Page.
<i>Labeo esopus</i> .....	186	<i>Labrosomus microlepidotus</i> .....	2361
<i>longatus</i> .....	186	<i>pectinifer</i> .....	2362
<i>labialis</i> , <i>Fundulus</i> .....	644	<i>xanti</i> .....	2363
<i>labiatus</i> , <i>Catostomus</i> .....	177	<i>labrosus</i> , <i>Bleennius</i> .....	2457
<i>Heros</i> .....	1530	<i>Ceratichthys</i> .....	319
<i>Labichthys</i> .....	368	<i>Hybopsis</i> .....	319
<i>carinatus</i> .....	368	<i>Zoarces</i> .....	2457
<i>elongatus</i> .....	360, 2802	<i>Labrus adpersus</i> .....	1577
<i>gilli</i> .....	368	<i>americanus</i> .....	1579
<i>Labidesthes</i> .....	805	<i>aper</i> .....	1586
<i>siccus</i> .....	805	<i>auritus</i> .....	1601
<i>Labracopsis</i> .....	1135	<i>bifasciatus</i> .....	1609
<i>labradoricus</i> , <i>Acanthocottus</i> .....	2001	<i>bivittatus</i> .....	1596
<i>Coregonus</i> .....	466	<i>blackfish</i> .....	1578
<i>Cottus</i> .....	2004	<i>brasiliensis</i> .....	1591
<i>Oncocottus</i> .....	2004	<i>capite obtuso</i> .....	1609
<i>Labrax</i> .....	1866, 2840	<i>carolinus</i> .....	1578
<i>albidus</i> .....	1132	<i>chogset</i> .....	1577
<i>americanus</i> .....	1135	<i>fulva</i> .....	1577
<i>clathratus</i> .....	1198	<i>chronis</i> .....	1483
<i>decagrammus</i> .....	1868	<i>chromis</i> .....	1483
<i>hexagrammus</i> .....	1872	<i>eruentatus</i> .....	1238
<i>lagocephalus</i> .....	1875	<i>cyancephalus</i> .....	1591
<i>lineatus</i> .....	1113	<i>exoletus</i> .....	1576
<i>monopterygius</i> .....	1866	<i>falcatus</i> .....	942
<i>mucronatus</i> .....	1135	<i>fulvomaculatus</i> .....	1339
<i>multilineatus</i> .....	1132	<i>fulvus</i> .....	1145
<i>nebulifer</i> .....	1195	<i>griseus</i> .....	1257
<i>nigricans</i> .....	1135	<i>grunneus</i> .....	1483
<i>notatus</i> .....	1132	<i>guaza</i> .....	1154
<i>octagrammus</i> .....	1870	<i>hiatula</i> .....	1578
<i>osculatii</i> .....	1132	<i>iridens</i> .....	988
<i>pallidus</i> .....	1135	<i>macropterus</i> .....	988
<i>pluvialis</i> .....	2841	<i>maximus</i> .....	1580
<i>rufus</i> .....	1135	<i>onitis</i> .....	1578
<i>stelleri</i> .....	1872	<i>ornatus</i> .....	1610
<i>superciliatus</i> .....	1873	<i>pallidus</i> .....	1005
<i>Labridae</i> .....	1571	<i>pentacanthus</i> .....	1576
<i>labridum</i> , <i>Hemulon</i> .....	1319	<i>plumieri</i> .....	1305
<i>labriformis</i> , <i>Epinephelus</i> .....	1155	<i>psittaculus</i> .....	1596
<i>Serranus</i> .....	1155	<i>pulcher</i> .....	1585
<i>Labrinae</i> .....	1572-1573	<i>radians</i> .....	1633
<i>Labrisomus</i> .....	2360	<i>radiatus</i> .....	1591
<i>biguttatus</i> .....	2360	<i>rostrum reflexo</i> .....	1677
<i>bucciferus</i> .....	2363	<i>rufus</i> .....	1583
<i>capillatus</i> .....	2362	<i>salmoides</i> .....	1012
<i>delalandi</i> .....	2350	<i>semiruber</i> .....	1583
<i>herminiger</i> .....	2361	<i>sparoides</i> .....	987
<i>microlepidotus</i> .....	2363	<i>squeteague</i> .....	1407, 1109
<i>nuchipinnis</i> .....	2362	<i>striatus</i> .....	1200
<i>xanti</i> .....	2362	<i>subfuscus</i> .....	1578
<i>Labroid Fishes</i> .....	1571	<i>tautoga</i> .....	1579
<i>Labroperca</i> .....	1148	<i>alla</i> .....	1579
<i>Labrosomus</i> .....	2360	<i>fusca</i> .....	1579
<i>cremnobates</i> .....	2366	<i>rubens</i> .....	1579
<i>macrocephalus</i> .....	2364	<i>tessellatus</i> .....	1578

Page.		Page.		Page.
2361	<i>Labrus torquatus</i> .....	1609	<i>laevigata</i> , <i>Pterophryno</i> .....	2717
2362	<i>versicolor</i> .....	1340	<i>laevigatus</i> , <i>Agonus</i> .....	2048
2363	<i>Lac de Marbre</i> Trout .....	515	<i>Chironectes</i> .....	2717
2457	<i>lacera</i> , <i>Lagochila</i> .....	199	<i>Lagocephalus</i> .....	1728
319	<i>Quassilabia</i> .....	199	<i>Phalangistes</i> .....	2048
319	<i>lacerta</i> , <i>Lampanyetus</i> .....	500	<i>Salmo</i> .....	508
2457	<i>Myctophum</i> .....	560	<i>Tetodon</i> .....	1728
1577	<i>Synodus</i> .....	537	<i>Laeviraja</i> .....	66
1579	<i>lacertinus</i> , <i>Synodus</i> .....	536	<i>lovis</i> , <i>Acipenser</i> .....	100
1586	<i>Lacerto</i> .....	537, 867	<i>Balistes</i> .....	1719
1901	<i>lacertosus</i> , <i>Hypobysis</i> .....	284	<i>Gasterosteus</i> .....	745
1609	<i>Minnilus</i> .....	284	<i>Orbis variegatus</i> .....	1735
1596	<i>Notropis</i> .....	284	<i>Pleuronectes</i> .....	2654
1578	<i>lacertus</i> , <i>Gobius</i> .....	2218	<i>Raja</i> .....	71
1591	<i>Seomber</i> .....	867	<i>Rhombus cornubiensis</i> .....	2654
1609	<i>Lachnolemus</i> .....	1579	<i>Squatina</i> .....	59
1578	<i>maximus</i> .....	1580	<i>La Fayette</i> .....	967, 1458
1577	<i>Lachnolepinus</i> .....	1579	<i>Lagarto</i> .....	533, 538
1577	<i>agula</i> .....	1580	<i>lagenarius</i> , <i>Acipenser</i> .....	102
1483	<i>caninus</i> .....	1580	<i>Lagocephalus</i> .....	1727
1483	<i>dux</i> .....	1580	<i>laevigatus</i> .....	1728
1298	<i>faleatus</i> .....	1580	<i>pachycephalus</i> .....	1729
1591	<i>maximus</i> .....	1579	<i>lagocephalus</i> , <i>Grauniatopleurus</i> .....	1875
1576	<i>psittacus</i> .....	1580	<i>Hexagrammos</i> .....	1873
942	<i>sullus</i> .....	1580	<i>Hexagrammus</i> .....	1875
1539	<i>lachrymalis</i> , <i>Ptychostomus</i> .....	194	<i>Labrax</i> .....	1875
1145	<i>laciniata</i> , <i>Kirtlandia</i> .....	795, 2840	<i>Oncorhynchus</i> .....	479
1257	<i>vagrans</i> .....	2840	<i>Salmo</i> .....	470
1483	<i>Menidia vagrans</i> .....	795	<i>Lagochila</i> .....	198
1154	<i>lacrimosum</i> , <i>Sparisoma</i> .....	1632	<i>lacera</i> .....	199
1578	<i>lacrimosus</i> , <i>Searns</i> .....	1632	<i>Lagodon</i> .....	1357
988	<i>lactarius</i> , <i>Catostomus</i> .....	175	<i>rhomboides</i> .....	1358
988	<i>Lactophrys</i> .....	1721, 1722, 1723	<i>Lake Carp</i> .....	167
1580	<i>bicaudalis</i> .....	1723	<i>Crescent Speckled Trout</i> .....	2821
1578	<i>oviceps</i> .....	1724	<i>Herring</i> .....	408
1610	<i>tricornis</i> .....	1724	<i>Lawyer</i> .....	2550
1005	<i>trigonus</i> .....	1724, 1724	<i>Sheepshead</i> .....	1484
1576	<i>triqueter</i> .....	1722	<i>Sturgeon</i> .....	106
1305	<i>lacustris</i> , <i>Ameiurus</i> .....	137	<i>Tahoe Trout</i> .....	403, 2870
1596	<i>Gadus</i> .....	137, 2551	<i>lalandi</i> , <i>Carcharias</i> .....	43
1585	<i>Hemiplus</i> .....	250	<i>Seriola</i> .....	902, 903
1633	<i>Ietalurus</i> .....	137	<i>lamarrii</i> , <i>Acipenser</i> .....	100
1591	<i>Pomolobus pseudoharen-</i> <i>gus</i> .....	426	<i>lamellifer</i> , <i>Exocoetus</i> .....	733
1677	<i>Lady-fish</i> .....	411, 1583	<i>Lamia</i> .....	38, 49, 50
1583	<i>Spanish</i> .....	1583	<i>lamia</i> , <i>Carcharhinus</i> .....	38
1012	<i>Lady-fishes</i> .....	410, 1581	<i>Carcharias</i> .....	38
1583	<i>Læmargus</i> .....	50	<i>Eulamia</i> .....	38
987	<i>borealis</i> .....	57	<i>lamiella</i> , <i>Carcharhinus</i> .....	37
1407, 1409	<i>Læmonema</i> .....	2556	<i>Lamiopsis</i> .....	33
1578	<i>barbatula</i> .....	2557	<i>Lamna</i> .....	49
1579	<i>barbatulum</i> .....	2556	<i>caudata</i> .....	37
1579	<i>melanurum</i> .....	2557	<i>cornubica</i> .....	49, 2740
1579	<i>laetabilis</i> , <i>Moniana</i> .....	272	<i>punctata</i> .....	48
1579	<i>letus</i> , <i>Centronotus</i> .....	2420	<i>spallanzani</i> .....	49
1578	<i>levicaudatus</i> , <i>Hippocampus</i> .....	777	<i>Lamnidae</i> .....	47
			<i>Lamninae</i> .....	47
			<i>lanotteni</i> , <i>Petromyzon</i> .....	10

	Page.		Page.
Lampadena .....	560	lanceolatus, Chastodon .....	1490
speculigera .....	561	Eques .....	1489, 1490
Lampanyctus .....	557	Gobius .....	2229, 2230
a'atus .....	559	Homoprion .....	1411
crocodillus .....	558	Limux .....	3
gemmifer .....	559	Lonchurus .....	1482
guntheri .....	559	Lonchurus .....	1482
lacerta .....	560	Stellifer .....	1413
resplendens .....	555	Lancet-fish .....	1691
townsendi .....	558	Lancet Fishes .....	503, 504, 505
Lamperina .....	6	Landlocked Salmon .....	487
Lampetra .....	12	Lane Snapper .....	1270
astori .....	12	Langburn .....	2433
aurea .....	13	Lant .....	833
camtschatica .....	13	Lantern Fishes .....	530, 550
cibaria .....	13	Lapon .....	1819
epihexodon .....	12	lappa, Halieutella .....	2742
plumbea .....	13	La Quescho .....	413
spadicea .....	13	Large-mouthed Black Bass .....	1012
tridentata .....	12	Large-sealed Sucker .....	192
variegata .....	2745	Larimus .....	1420, 1421
wilderi .....	13, 2745	acelivis .....	1422
lampetrisformis, Blennius .....	2438	argenteus .....	1421
Lampenus .....	2438	bafabannus .....	1431
Lamprey Eel .....	10	breviceps .....	1423
Lamprey, Great Sea .....	10	dentex .....	1426
Silvery .....	11	effulgens .....	1421
Small Black .....	13	fasciatus .....	1424
Lampreys .....	4, 8, 9	pacificus .....	1424
Brook .....	12	stahl. .....	1423
River .....	10	larkinsii, Cymnogaster .....	1503
Lamprida .....	953	lata, Gnaptera forcipata .....	1702
Lampris .....	954	Latebrus .....	1114
guttatus .....	955	oculatus .....	1115
lauta .....	955	latepletus, Serranus .....	1175
luna .....	954	laterale, Ditrema .....	1506
regius .....	955	Etheostoma .....	1099
lamprotentia, Clupea .....	419	lateralis, Abramis .....	239
Jenkinsia .....	419	Alvarius .....	1099
Spatelloides .....	419	Arctedius .....	1902
lamprurus, Hypoplectrus .....	1190	Calycilepidotus .....	1900
Serranus .....	1190	Caracodon .....	2832
Lampugus .....	952	Characodon .....	668
neapolitanus .....	953	Eleotris .....	2195
punctulatus .....	953	Embiotoca .....	1506
siculus .....	953	Leuciscus balteatus .....	239
lanatus, Merluccius .....	2530	Mylocheilus .....	220
Lancelet, Bahama .....	4	Notropis .....	263
California .....	4	Phanerodon .....	1506
West Indian .....	3	Philypnus .....	2195
Lancelets .....	2, 3	Pimelodus .....	135
European .....	3	Pencilthys .....	1319
lanceolata, Perca .....	1482	Richardsonius .....	239
Sciama .....	1444	Searus .....	1657
lanceolatum, Branchiostoma .....	3	Scorpenichthys .....	1902
lanceolatus, Amphioxus .....	3	Tanlotoca .....	1505, 1506

Page.		Page.		Page.
1490	lateralis, Zygonectes.....	650	Leather-sided Minnow.....	230
1489, 1490	laticauda, Anguilla.....	348	Le Baliste Bridé.....	1704
2220, 2220	Rhamdia.....	1512	Leblas ellipsoides.....	672
1444	laticaudus, Pimelodus.....	1512	ovinus.....	672
3	laticeps, Acanthocottus.....	1980	rhomboidalis.....	672
1482	Aetobatus.....	88, 2753	Lebistes.....	689, 2833
1482	Arius.....	132	pencilhodes.....	680
1443	Atherina.....	790	Lebius.....	1866
1691	Bathynectes.....	2523	Lebrancho.....	810
93, 504, 505	Megalocottus.....	1988	lebranchus, Mugil.....	811
487	Misonus.....	2523	lecontei, Acipenser.....	105
1270	latielavivus, Prienurus.....	1696	Lo Diodon.....	1746
2433	Xesurus.....	1695	Orbe.....	1740
833	latidens, Microstomus.....	2654	Tacheté.....	1740
530, 530	latifasciatus, Cyprinodon.....	676	leel, Symphurus.....	2708
1819	latifrons, Anarhichas.....	2446	lefroyi, Diapterus.....	1372
2742	Citharichthys.....	2674	Encinostomus.....	1372
413	Dormitor.....	2197	Ulma.....	1371
1012	Eleotris.....	2198	Lefroyia.....	2495
192	Syacium.....	2673	hermudensis.....	2497
1420, 1421	Xenochirus.....	2082	Le Gastrobrancho Dombey.....	6
1422	latilimo.....	2275	leiarclus, Cestrens.....	1415
1421	latilus chrysops.....	2278	Cynoseion.....	1414
1131	princeps.....	2277	Otolithus.....	1415
1423	latiuaculatus, Ophisurus.....	376	Leiolabrus.....	61, 79
1426	latimama, Belono.....	717	sloani.....	81
1121	lator, Coregonus.....	466	Leiocottus.....	2010
1421	latipinna, Mollienisia.....	700	hirundo.....	2011
1421	latipinnis, Catostomus.....	174, 2790	Leiodon.....	56
1123	Zaniolepis.....	1876	echinatum.....	57
1506	latirostris, Acipenser.....	105	Leioglossus.....	916
1702	Lepidosteus.....	111	leionothis, Holacanthus.....	1735
1114	latulus, Clupea.....	422	leipomus, Cottus.....	1962
1115	latus, Caranx.....	922	Lelostomus.....	1558
1175	Euctenogobius.....	2237	humeralis.....	1459
1506	Scomber.....	938	lineatus.....	1460
1099	launces, Sand.....	831, 832, 833	obliquus.....	1459
239	laurette, Argyrosomus.....	471	xanthurus.....	1458
1099	Laurida.....	533	Leirus.....	746
1902	mediterranea.....	537	leirus, Gasterosteus.....	747
1900	laurito, Sparisoma.....	1637	Le Kai Salmon.....	478
2832	lauta, Lampris.....	955	Lembus.....	2194
668	lavaretns, Salmo.....	464	lemonii, Squalus.....	225
2195	Lavinia.....	208	lemniscatus, Engraulis.....	443
1506	conformis.....	231	Osmerus.....	533
239	crassicauda.....	231	Pimelodus.....	147
220	exilicauda.....	208, 2799	Lemnisoma.....	883
263	harengus.....	209	thyrsitoides.....	884
1506	Lawyer.....	113, 1255	Lenguado de Rio.....	2698
2195	Lawyer, Lako.....	2550	lenibus, Ostracion triangulatus.....	1724
135	leachi, Clupea.....	422	lentiginosa, Amia.....	113
1319	leachianus, Thynnus.....	869	Murena.....	402
239	Least Darter.....	1104	lentiginosum, Cichlasoma.....	1524
1637	Leath Fish.....	1714, 1715	lentiginosus, Galeichthys.....	122, 2771
1902	Leather Jacket.....	1701	Heres.....	1524
1505, 1506	Jackets.....	898	Rhinobatus.....	62, 2750

	Page.		Page.
lentiginosus, Tachysurus .....	122	Lepidosteus grayi .....	111
leonensis, Oligocephalus .....	1089	latrostris .....	111
leonina, Cliola .....	271	leptorhynchus .....	110
Meniana .....	272	manjuari .....	111
Leopard Shark .....	31	oculatus .....	111
leopardinus, Antennarius .....	2721	otarius .....	110
Platophrys .....	2666	viridis .....	111
Rhomboidichthys .....	2666	lepidulus, Alburnus .....	291
leopardus, Acanthichas .....	2446	lepidum, Boleosoma .....	1085
Lepadogaster cornubiensis .....	2108	Etheostoma .....	1089
nudus .....	2331	lepidus, Gobius .....	2249
reticulatus .....	2328	Lepidogobius .....	2249
testar .....	2332	Pocilloichthys .....	1089
Lepibema .....	1131	Lepinophilis .....	952
lineatum .....	1133	hippuroides .....	952
mitchilli .....	1133	Lepiopomus .....	399
lepada, Boleosoma .....	1089	ischyrus .....	997
Cliola .....	273	Lepisoma .....	2360
Cyprinella .....	273	cirrhosum .....	2362
Lepidamla .....	1106	Lepiosteidae .....	108
Lepidion .....	2543	Lepiosteus .....	109
verecundum .....	2543	albus .....	110
Lepidochatodon .....	1672	bison .....	110
Lepidocybium .....	873	ferox .....	111
Lepidogaster macandricus .....	2328	gavialis .....	110
lepidogenys, Etheostoma .....	1087	gracilis .....	110
Lepidogobius .....	2249	huronensis .....	110
omblematicus .....	2247	lineatus .....	110
gilberti .....	2254	longirostris .....	110
gracilis .....	2249	osseus .....	109
galosus .....	2244	oxyurus .....	110
lepidus .....	2249	platostomus .....	110
newberryi .....	2248	platyrhincus .....	111
thalassinus .....	2245	platystomus .....	110
Lepidolepis .....	2568	semiradintus .....	110
norvegicus .....	2579	spatula .....	111
Lepidomeda .....	328	tristechus .....	111
jarrovi .....	328	tropicus .....	111
vittata .....	328	Lepodus .....	958
Lepidomegas .....	901	saragus .....	960
Lepidopidae .....	884	Lepominae .....	985
Lepidopline .....	885	Lepomis .....	999, 1010
Lepidopsetta .....	2642	albus .....	1007
bilineata .....	2643	annagallinus .....	1004
isolepis .....	2642	aplatus .....	998
umbrosa .....	2642	appendix .....	1005
Lepidopus .....	886	ardesiatus .....	1006
argyreus .....	887	auritus .....	1001, 1009
caudatus .....	887, 2844	sollis .....	1001
gouani .....	887	bombifrons .....	1003
petonii .....	887	charybdis .....	992
xantasi .....	2843, 2844	cyanellus .....	996
Lepidosoma .....	2568	euryurus .....	1009
Lepidosteus .....	109	flexularis .....	1011
berlandieri .....	111	garmani .....	1002
crassus .....	110	gillii .....	992



Page.		Page.		Page.
111	<i>Lepomis haplognathus</i> .....	1004	<i>Leptocephalus conger</i> .....	354
111	<i>heros</i> .....	1008	<i>gracilis</i> .....	354
116	<i>holbrookii</i> .....	1008	<i>morrisi</i> .....	354
111	<i>humilis</i> .....	1004	<i>spallanzanii</i> .....	354
111	<i>ichtheloides</i> .....	990	<i>leptocephalus, Ceratichthys</i> .....	323
116	<i>ischyrus</i> .....	997	<i>Merlangus</i> .....	2535
111	<i>llrus</i> .....	1007	<i>Leptoclinus</i> .....	2432
294	<i>longispinis</i> .....	1006	<i>maculatus</i> .....	2433
1085	<i>macrochirus</i> .....	1005	<i>Leptoconger</i> .....	362
1089	<i>marginatus</i> .....	1003	<i>prolongus</i> .....	363
2249	<i>megaleotis</i> .....	1002	<i>Leptocottus</i> .....	2111
2249	<i>miniatus</i> .....	1002	<i>armatus</i> .....	2012
1085	<i>mystacalis</i> .....	1001	<i>Leptodes</i> .....	584
952	<i>notata</i> .....	1011	<i>Leptogunnellus</i> .....	2435
952	<i>notatus</i> .....	1008	<i>Leptophidium</i> .....	2482
39	<i>ophthalmicus</i> .....	1001	<i>marmoratum</i> .....	2483
997	<i>pallida</i> .....	1012	<i>microlepis</i> .....	2486
2360	<i>pallidus</i> .....	1005	<i>prorates</i> .....	2485
2362	<i>peltastes</i> .....	1003	<i>Leptops</i> .....	142
108	<i>phenax</i> .....	997	<i>olivaris</i> .....	143, 2700
109	<i>punctatus</i> .....	998	<i>Leptorhinophis</i> .....	381
110	<i>purpurescens</i> .....	1006	<i>leptorhynchum, Siphostoma</i> .....	704
110	<i>salmonea</i> .....	1011	<i>Leptorhynchus</i> .....	369
111	<i>symmetricus</i> .....	999	<i>lenchtenbergii</i> .....	369
110	<i>trifasciata</i> .....	1011	<i>leptorhynchus, Lepidosteus</i> .....	110
110	<i>Lepomotis nephielus</i> .....	1005	<i>Odontopyxis</i> .....	2076
110	<i>Lepophidium</i> .....	2482	<i>Sarritor</i> .....	2075
110	<i>brevibarbe</i> .....	2485	<i>Syngnathus</i> .....	765
110	<i>cervinum</i> .....	2484, 2485	<i>leptoscenus, Abramis</i> .....	250
109	<i>emmelas</i> .....	2483	<i>Luxilus</i> .....	250
110	<i>marmoratum</i> .....	2482	<i>Notemigonus</i> .....	250
110	<i>microlepis</i> .....	2486	<i>Lepturus</i> .....	889
111	<i>pardale</i> .....	2486	<i>lepturus</i> .....	889
110	<i>profundorum</i> .....	2484	<i>lepturus, Caranx</i> .....	923
110	<i>stigmatistium</i> .....	2483	<i>Lepturus</i> .....	889
111	<i>leptacanthus, Noturus</i> .....	146	<i>Macrourus</i> .....	2584
111	<i>Schilbeoides</i> .....	146	<i>Trichinrus</i> .....	869
111	<i>Lontagonus</i> .....	2052	<i>Le Sphéroïde Tuberculé</i> .....	1733
958	<i>decagonus</i> .....	2052	<i>Tetrodon Plumier</i> .....	1733
960	<i>spinosissimus</i> .....	2054	<i>Les Alutères</i> .....	1717
985	<i>Lep'tarius</i> .....	119	<i>Batrachopes</i> .....	1740
999, 1010	<i>dowl</i> .....	122	<i>Brosme</i> .....	2561
1007	<i>Leptaspis</i> .....	310	<i>Curimates</i> .....	332
1004	<i>Leptecheneis</i> .....	2268	<i>Dichotomycetères</i> .....	1738
998	<i>naucrateoides</i> .....	2270	<i>Dilobomycetères</i> .....	1738
1005	<i>naucrates</i> .....	2269	<i>Elacates</i> .....	948
1006	<i>Leptoblenius</i> .....	2435	<i>Lottes</i> .....	2550
1009	<i>nubilus</i> .....	2438	<i>Mustèles</i> .....	2557
1001, 1009	<i>serpentinus</i> .....	2439	<i>Ovoïdes</i> .....	1738
1001	<i>Leptocardi</i> .....	2	<i>Pristipomes</i> .....	1329
1003	<i>Leptocephalichthys</i> .....	353	<i>Pronecocephales</i> .....	1727
992	<i>Leptocephalide</i> .....	352	<i>Sphéroïdes</i> .....	1729
996	<i>Leptocephalus</i> .....	353	<i>Stellifères</i> .....	1439
1009	<i>candlissimus</i> .....	354	<i>Stenomotopes</i> .....	1729
1011	<i>caudicula</i> .....	355	<i>lessoni, Caranx</i> .....	923
1002	<i>caudilimbatus</i> .....	355	<i>Tetrapturus</i> .....	892
992				

	Page.		Page.
lessonii, <i>Coryphæna</i> .....	953	<i>Leuciscus gracilis</i> .....	283, 326
lesueurii, <i>Aclpenser</i> .....	106	<i>grabami</i> .....	228
<i>Moxostoma</i> .....	194	<i>grandis</i> .....	225
lesueurianum, <i>Exoglossum</i> .....	327	<i>luenaturnus</i> .....	218
lesueurii, <i>Catostomus</i> .....	195	<i>heterodon</i> .....	261
Letharcus.....	375	<i>hudsonius</i> .....	269
<i>velifer</i> .....	375	<i>humboldtii</i> .....	236
lethopristis, <i>Orthopristis</i> .....	1340	<i>hydrophlox</i> .....	238
lethostigma, <i>Paralichthys</i> .....	2630	<i>hypselepterus</i> .....	280
lethostigmus, <i>Paralichthys</i> .....	2630	<i>intermedius</i> .....	235
Lethostole.....	792, 2839	<i>iris</i> .....	222
<i>estor</i> .....	792	<i>kentuckiensis</i> .....	279
Lethotremus.....	2100	<i>luceatus</i> .....	232
<i>muticus</i> .....	2101	<i>lutrensis</i> .....	272
<i>vinolentus</i> .....	2101	<i>macrolepidotus</i> .....	224
leuchtenbergii, <i>Belonopsis</i> .....	369	<i>margarita</i> .....	241
<i>Leptorhynchus</i> .....	369	<i>milnerianus</i> .....	242
leuciodus, <i>Minnilus</i> .....	291	<i>montanus</i> .....	238
<i>Notropis</i> .....	291	<i>nachtriebi</i> .....	2798
<i>Photogenis</i> .....	291	<i>nasutus</i> .....	306
Leuciscinae.....	202	<i>neogeus</i> .....	240, 2798
Leuciscus.....	238, 252	<i>niger</i> .....	235
<i>affinis</i> .....	240	<i>nigrescens</i> .....	233
<i>aliciae</i> .....	236	<i>nitidus</i> .....	221
<i>analostanus</i> .....	279	<i>obosus</i> .....	246, 282
<i>ardens</i> .....	301	<i>oreutti</i> .....	241
<i>argenteus</i> .....	221	<i>oregonensis</i> .....	225
<i>balteatus</i> .....	238, 2797	<i>phlegethontis</i> .....	243
<i>lateralis</i> .....	239	<i>photogenis</i> .....	296
<i>bicolor</i> .....	232, 245	<i>puceae</i> .....	264
<i>boscl</i> .....	251	<i>productus</i> .....	240
<i>boucardi</i> .....	247	<i>prolixus</i> .....	206
<i>bubalinus</i> .....	273	<i>proxiger</i> .....	240
<i>chrysopterus</i> .....	221	<i>pulchelloides</i> .....	222
<i>coblitis</i> .....	305	<i>pulchellus</i> .....	221
<i>coccogenis</i> .....	285	<i>purpureus</i> .....	234
<i>conformis</i> .....	231	<i>pygmaeus</i> .....	624
<i>cooperi</i> .....	236	<i>robustus</i> .....	228
<i>copli</i> .....	293	<i>rotengulus</i> .....	221
<i>cornutus</i> .....	283	<i>rubellus</i> .....	293
<i>crassicauda</i> .....	231	<i>rubrifrons</i> .....	295
<i>croceus</i> .....	308	<i>siuslawi</i> .....	2797
<i>egregius</i> .....	237	<i>spilopterus</i> .....	279
<i>elegans</i> .....	227	<i>spirlingulus</i> .....	282
<i>elongatus</i> .....	240, 2797	<i>storeri</i> .....	222
<i>emorii</i> .....	227	<i>storerianus</i> .....	270
<i>erythrogaster</i> .....	210	<i>telescopus</i> .....	292
<i>estor</i> .....	240	<i>tincella</i> .....	217
<i>oxilicauda</i> .....	209	<i>tuditanus</i> .....	253
<i>flammeus</i> .....	242	<i>vandoisulus</i> .....	239
<i>formosus</i> .....	246	<i>vittatus</i> .....	282
<i>frontalis</i> .....	283	<i>volucellus</i> .....	263
<i>funduloides</i> .....	240	<i>zeylonicus</i> .....	415
<i>gardoneus</i> .....	251	<i>zunnensis</i> .....	227
<i>gibbosus</i> .....	231	<i>leuciscus</i> , <i>Brachyleuterus</i> .....	1327
<i>gilli</i> .....	239	<i>Pomadasis</i> .....	1328

Page.		Page.		Page.
283, 326	leuciscus, Pomadasys.....	1328	lewis, Squatina.....	59
228	Pristipoma.....	1328	lewisi, Salar.....	493
225	leucomenis, Salmo.....	2823	Salmo clarkii.....	2819
218	leucopneus, Congrus.....	355	mykiss.....	493
261	leucopsarum, Myctophum (Steno-		Zygæna.....	45
269	brachius).....	562	iberiensis, Balistes.....	1702
236	Nannobranchium.....	562	libertate, Opisthonema.....	433
238	leucops, Photogenis.....	296	libertatis, Clupea.....	433
280	engraulinus.....	290	Meletta.....	433
235	leucopus, Photogenis.....	277	Opisthonema.....	433
222	Rhamphoberyx.....	847	Lichia quiebra.....	890
279	leucorhynchus, Rhinobatus.....	62	lichtensteini, Acipenser.....	105
292	leucorus, Eupomacentrus.....	1551	ligulata, Seriola.....	905
272	Leucos.....	243, 244, 2798	Lija.....	1714, 1715, 1718
224	bicolor.....	245	Barbuda.....	1720
241	formosa.....	246	Colorado.....	1713
242	obesus.....	246	Trompa.....	1710
238	leucos, Carcharhinus.....	38	Lilo.....	428, 429, 431
2768	Carcharias.....	38	lima, Loricaria.....	158
306	Leucosomus.....	220, 221, 250	Limamurena.....	400
240, 2798	americanus.....	250	melanotis.....	402
235	argyreus.....	224	Limanda.....	2644
236	catarnctus.....	221	aspera.....	2645
221	caurinus.....	220	beanii.....	2646
246, 282	communis.....	326	ferruginea.....	2644
241	corporalis.....	222	proboscidea.....	2645
225	dissimilis.....	324	rostrata.....	2645
243	gulonellus.....	326	limanda, Hippoglossoides.....	2615
296	inaequilobus.....	224	limandoites, Hippoglossoides.....	2615
264	incrassatus.....	223	Pleuronectes.....	2615
240	occidentalis.....	247	Limax lanceolatus.....	3
206	pallidus.....	222	limbatus, Carcharhinus.....	40
240	pulchellus.....	222	Carcharias.....	40
222	rhotheus.....	222	Conger.....	360
221	symmetricus.....	246	Fundulus.....	643
234	leucosteus, Calamus.....	1353	isogomphodon.....	40
624	leucostictus, Eupomacentrus.....	1555	Oxydenticthys.....	385
228	Pomacentrus.....	1556	Saurus.....	533
221	leucurus, Caranx.....	915	limi, Hydrargyra.....	624
293	Heulcaranx.....	014	Umbra.....	623
295	Naucleus.....	900	pygmaea.....	624
2797	Leucus anteus.....	245	Limia.....	690
279	boncardi.....	247	couchiana.....	695
282	dimidiatus.....	244	cubensis.....	692
222	formosus.....	246	matamorensis.....	700
270	olivaceus.....	244	pavonia.....	692
292	tincella.....	211	Pecilloides.....	700
211	Leuresthes.....	801	venusta.....	665
253	cramerii.....	802	Limnurgus.....	666
239	tenis.....	802	variegatus.....	666
282	Lenroglossus.....	527	limosa, Myxino.....	8
293	stilbius.....	527	limosus, Pylodictis.....	142
415	Leurynnis.....	2460	Pylodictis.....	143
227	paucidens.....	2460	Silurus.....	143
1327	levis, Sebastichthys.....	1816	limca, Mesoprion.....	1260
1328	Sebastodes.....	1816	Siphonoma.....	768

	Page.		Page.
linea, Syngnathus.....	768	lineolatus, Metrogaster.....	1499
lineata, Alosa.....	426	Pseudoscarus.....	1651
Coryphæna.....	1619	Tetrodon.....	1728
Eoheneis.....	2268, 2270	lineopinnis, Murena.....	396
Embiotoca.....	1506	lineo-punctatus, Balistes.....	1709
Morone.....	1133	Ling.....	2550
Novacula.....	1619	Lings.....	2551
Sciæna.....	1133, 1460	linguata, Balistes.....	1720
Tigoma.....	233	Pleuronectes.....	2615
Trigla.....	2167	linnaei, Eglofinus.....	2543
Unibranchapertura.....	342	Merluccius.....	2530
lineatum, Lepibema.....	1133	Molva.....	2552
lineatus, Achirus.....	2697, 2698, 2702	Trachurus.....	911
Amblodon.....	1484	Linophora.....	1672
Calliodon.....	1651	Linophryne.....	2734
Cyclopterus.....	2118	lucifer.....	2734
Cypselurus.....	2836	linsleyi, Etheostoma.....	1097
Dormitator.....	2108	Liocetus.....	2733
Doryichthys.....	773	murrayi.....	2733
Doryrhamphus.....	773	Lioglossina.....	2622
Eques.....	1487	tetrophthalmæ.....	2622
Esox.....	627	liolepis, Paralichthys.....	2624
Exocoetus.....	739	Xystreurus.....	2623
Funnulus.....	649	Liomonacanthus.....	1713
Genyonemus.....	1460	Lionfish.....	1850
Gerres.....	1377	Lioniscus.....	103
Gobius.....	2218, 2260	liopeltis, Acipenser.....	106
Grammichthys.....	2702	Lioperca.....	1166
Grystea.....	1868	inermis.....	1168
Labrax.....	1113	Liopropoma.....	1135
Leiostomus.....	1460	aberrans.....	1136
Leplaoateus.....	110	rubra.....	1137
Leuciscus.....	232	Liopropomina.....	1127
Liparis.....	2118	Liopsetta.....	2649
multistriatus.....	2118	glabra.....	2650
Micropogon.....	1461	glacialis.....	2640, 2650
Monoclinr.....	2698	obscura.....	2651
Mugil.....	812, 2841	putnami.....	2650
Phtheirichthys.....	2268	lorus, Chasmistes.....	183
Pleuronectes.....	2698, 2701	loaternus, Phenacobius teretulus..	303
Prionotus.....	2167	Liparididæ.....	2105
Rocens.....	1113, 1132	Liparidina, Discoboli.....	2105
Smaris.....	1378	Liparidinae.....	2105
Squallus.....	233	Liparlua, Amitra.....	2138, 2139
Trichodon.....	2297	Monomitra.....	2139
Xyrichthys.....	1619	Liparis.....	2114, 2115, 2116, 2118
Zygonectes.....	649, 657	agassizii.....	2121
lluolata, Cliola.....	263	arctica.....	2121
Clupea.....	422	barbatus.....	2118
Coryphæna.....	1619	calliodon.....	2120
Mollenisla.....	700	callyodon.....	2111
Pelamys.....	873	(Careproctus) reinhardi.....	2134
Pæcilla.....	700	communis.....	2118
lineolatum, Etheostoma flabellare..	1098	cyclogaster.....	2118
lineolatus, Alburnus.....	263	cyclopus.....	2112, 2118
Catonotus.....	1099	cyclostigma.....	2125, 2865

Page.		Page.		Page.
1499	<i>Liparis dennyi</i> .....	2124	Little Tunny .....	869
1651	<i>ekstromi</i> .....	2108	Little-head Porgy .....	1350
172-	<i>fabricii</i> .....	2121, 2128	Little-mouth Porgy .....	1354
396	<i>fucensis</i> .....	2119	<i>littoralis</i> , <i>Carcharias</i> .....	46, 2748
1709	<i>gelatinosus</i> .....	2134, 2135	<i>Engomphodus</i> .....	47
2550	<i>gibbus</i> .....	2123	<i>Menticirrhus</i> .....	1477
2551	<i>gobius</i> .....	2108	<i>Squalus</i> .....	47
1720	<i>herschelii</i> .....	2123	<i>Umbrina</i> .....	1477
2615	<i>lineatus</i> .....	2118	<i>littoricola</i> , <i>Chaetodon</i> .....	1680
2543	<i>multistriatus</i> .....	2118	<i>litura</i> , <i>Mesoprion</i> .....	1258
2530	<i>liparis</i> .....	2116, 2118	<i>liturosus</i> , <i>Diodon</i> .....	1746
2552	<i>maculatus</i> .....	2108	<i>lividus</i> , <i>Petromyzon</i> .....	12
911	<i>major</i> .....	2127	<i>Silurus</i> .....	140
1672	<i>montagui</i> .....	2107, 2108	<i>Liza</i> .....	810
2734	<i>mucosus</i> .....	2111	<i>liza</i> , <i>Mugil</i> .....	811, 2841
2734	<i>nostras</i> .....	2118	<i>Liza Ojo de Perdiz</i> .....	814
1097	<i>ophioides</i> .....	2118	Lizardfish .....	538
2733	<i>pulchellus</i> .....	2126	Lizard Fishes .....	532, 533
2733	<i>ranula</i> .....	2134	<i>lobatus</i> , <i>Canthogaster</i> .....	1732
2622	<i>reticulata</i> .....	2108	<i>Spheroides</i> .....	1731, 1732
2622	<i>stellatus</i> .....	2118	<i>lobochilus</i> , <i>Heros</i> .....	1531
2624	<i>tunicata</i> .....	2121, 2128	<i>Lobotes</i> .....	1235
2623	<i>tunicatus</i> .....	2120	<i>auctorum</i> .....	1236, 2858
1713	<i>vulgaris</i> .....	2118	<i>emarginatus</i> .....	1257
1850	<i>liparis</i> , <i>Centrolophus</i> .....	963	<i>erate</i> .....	1236, 2856
103	<i>Cyclopterus</i> .....	2123	<i>farkharii</i> .....	1236, 2856
106	<i>major</i> .....	2128	<i>incurvus</i> .....	1236, 2856
1166	<i>minor</i> .....	2121	<i>pacificus</i> .....	2857, 2858
1168	<i>Liparis</i> .....	2116, 2118	<i>somnolentus</i> .....	1236
1135	<i>liparoides</i> , <i>Cyclopterus</i> .....	2108	<i>surinamensis</i> .....	1235, 2856, 2858
1136	<i>Liparops</i> .....	2104	<i>Lobotidae</i> .....	1235
1137	<i>stelleri</i> .....	2104	<i>lockingtonii</i> , <i>Teichthys</i> .....	969
1127	<i>Liparopsina</i> .....	2095	<i>Lodde</i> .....	520
2649	<i>Lipogenis</i> .....	619	<i>lueve</i> , <i>Ditrema</i> .....	1511
2650	<i>Lipogenyidae</i> .....	610	<i>Log Fish</i> .....	964
2649, 2650	<i>Lipogenys gillii</i> .....	619	<i>Perches</i> .....	1024, 1028
2651	<i>Lipophrys</i> .....	2377, 2378	<i>Lonchichurus</i> .....	1481
2650	<i>lipopus</i> , <i>Tachysaurus</i> .....	2784	<i>lanceolatus</i> .....	1482
183	<i>Lirus perciformis</i> .....	964	<i>Lonchopisthus</i> .....	2286
303	<i>lirus</i> , <i>Lepomis</i> .....	1007	<i>micrognathus</i> .....	2287
2105	<i>Minulus</i> .....	298	<i>lonchurus</i> , <i>Opisthognathus</i> .....	2281
2105	<i>Notropis</i> .....	297	<i>Lonchurus ancyrodon</i> .....	1416
2105	<i>Lisa Blanca</i> .....	813	<i>barbatus</i> .....	1482
2138, 2139	<i>Cahezuda</i> .....	811	<i>depressus</i> .....	1482
2139	<i>Francesa</i> .....	410	<i>lanceolatus</i> .....	1482
5, 2116, 2118	<i>lisiga</i> , <i>Aspidophorus</i> .....	2036	<i>lonchurus</i> , <i>Opisthognathus</i> .....	2281
2121	<i>Lisitu</i> .....	814	<i>Longjaw</i> .....	471
2121	<i>listeri</i> , <i>Ostracion</i> .....	1725	<i>Long-jawed Goby</i> .....	2250
2118	<i>lita</i> , <i>Murena</i> .....	2805	<i>Long-jaws</i> .....	710, 711
2120	<i>Litholepis tristæchus</i> .....	111	<i>Long Mingo</i> .....	1718
2111	<i>Little Pickerel</i> .....	627	<i>longa</i> , <i>Dasibatis</i> .....	85
2134	<i>Red-eye</i> .....	996	<i>Dasyatis</i> .....	85
2118	<i>Roncador</i> .....	1460	<i>Macdonaldia</i> .....	2826
2118	<i>Skate</i> .....	68	<i>longatus</i> , <i>Labeo</i> .....	186
2112, 2118	<i>Smelt</i> .....	807	<i>Longe</i> .....	504
2125, 2865	<i>Tunnies</i> .....	868	<i>Long-eared Sunfish</i> .....	1002

	Page.		Page.
Long-finned Albacore.....	871	longulus, Calliurus.....	996
Charr.....	509	Pomotis.....	996
Sole.....	2658	longurio, Carcharias.....	42
longicauda, Gobius.....	2220	Scoliodon.....	42, 2748
Gymnothorax.....	392	longus, Ballstes.....	1707
Murana.....	392	Ophisurus.....	377
Rabula.....	391	Pisodonophis.....	377
longicephalus, Galeichthys.....	781	Look-Down.....	936
Hexanematischthys.....	130	lophar, Perca.....	947
Tachisurus.....	130	Lopharis.....	946
longiceps, Eleotris.....	2195	mediterraneus.....	947
Hybopsis.....	264	Lophiidae.....	2713
Sibona.....	233	Lophlomus.....	2714
longicollis, Myrophis.....	371	setigerus.....	2714
longidens, Caulolepis.....	839	Lophius.....	2713
longitilis, Bathygadus.....	2566	aculeatus.....	2741
Hymenocephalus.....	2567	americanus.....	2714
longimana, Etheostoma.....	1054	bufo.....	2316
Eulamia.....	38	gibbus.....	2717
longimanus, Boleosoma.....	1054	histrion.....	2716, 2722
Cichlasoma.....	1520	ocellatus.....	2722
Heros.....	1521	piscatorius.....	2713
Squalus.....	38	radiatus.....	2738
Xystroplites.....	1008	rostratus.....	2737
longipes, Bathypterois.....	546	setigerus.....	2715
Gadus.....	2555	spectrum.....	2723
longipinna, Gobiosoma.....	2256	tinnulus.....	2716
Holocentrum.....	849	viviparus.....	2715
longipinnis, Clevelandia.....	2255	lophius, Amiurus.....	138
Evermannia.....	2256	Ictalurus.....	138
Rhombus.....	966	Lophobranchii.....	759
Stromateus.....	966	Lophobranchs.....	759
longirostris, Alburnops.....	267	Lophogobius.....	2209
Catostomus.....	176	cyprinoides.....	2209
Ctiola.....	267	Lopholatilus.....	2278
Esox.....	714	chamaeleonticeps.....	2278
Euleptorhamphus.....	724	Loplosetta.....	2659
Hemirhamphus.....	724	maculata.....	2666
Hippocampus.....	778	Lord, Irish.....	1934
Hypoprion.....	41	lordii, Salmo.....	508
Lepisosteus.....	110	loreto, Gramma.....	1229
Malthua.....	2737	Loricaria.....	156, 157, 159
Notropis.....	267	acuta.....	158
Saurus.....	538	barbata.....	158
Tylosurus.....	714	bransfordi.....	158
longirostrum, Catostomus.....	176	lima.....	158
longispathum, Peristedion.....	2178	panamensis.....	157
longispinis, Ailurichthys.....	119	rostrata.....	157
Lepomis.....	1006	strigilata.....	158
Pontinus.....	1858	uracantha.....	158
Long-necked Eels.....	343	variegata.....	159
Long-nosed Dace.....	306	Loricariichthys.....	156
Gar.....	109	Loricariidae.....	155
Sucker.....	176	Loricariinae.....	156
Long-spined Sculpin.....	1076	loricata, Alysia.....	569
Long-tail Shark.....	45	Loricati.....	1756

Page.		Page.		Page.
906	loricatus, <i>Gasterosteus</i> .....	747	Lucifuginae .....	2408
956	<i>Macrognaathus</i> .....	110	Luciida .....	624
42	<i>Phalangistes</i> .....	2040	Lucioblennius .....	2404
42, 2748	Loro .....	1652, 1653, 1655, 1657	<i>alepidotus</i> .....	2404
1707	loro, <i>Scarus</i> .....	1654	Lucioceps, <i>Saurus</i> .....	530
377	Loros .....	1642	<i>Synodus</i> .....	530
377	Lota .....	2550	Luciocharax .....	339
936	<i>brosimiana</i> .....	2551	<i>insculptus</i> .....	339
947	<i>compressa</i> .....	2551	Lucioides, <i>Esox</i> .....	628
946	<i>inornata</i> .....	2551	Lucioperca americana .....	1022
947	<i>maculosa</i> .....	2550	<i>borea</i> .....	1022
2713	Lotella .....	2546	<i>canadensis</i> .....	1022
2714	<i>maxillaris</i> .....	2546	<i>grisea</i> .....	1022
2714	lotharingus, <i>Gasterosteus</i> .....	746	<i>pepinus</i> .....	1022
2713	Lotinae .....	2532	<i>vitrea</i> .....	1022
2714	loubina, <i>Perca</i> .....	1119	Luciopercana, <i>Mentiperca</i> .....	1216
2316	louisiana, <i>Ingraulis</i> .....	446	Luciopercanus, <i>Centropristis</i> .....	1216
2717	louisiana, <i>Notropis</i> .....	2801	<i>Prionodes</i> .....	1216
0716, 2722	<i>Siphonotoma</i> .....	770	<i>Serranus</i> .....	1216
2722	<i>Syngnathus</i> .....	770	Luciopercinae .....	1018
2713	lowei, <i>Polymixia</i> .....	854	Luciotrutta .....	473
2738	lowii, <i>Omosudis</i> .....	598	<i>nackenzii</i> .....	474
2737	loxias, <i>Prionotus</i> .....	2156	Lucius .....	625, 626, 628
2715	lubb, <i>Gadus</i> .....	2561	<i>americanus</i> .....	626
2723	lubricum, <i>Branchiostoma</i> .....	3	<i>lucius</i> .....	628
2716	Lucania .....	663, 666	<i>maquinongy</i> .....	629
2715	<i>affinis</i> .....	665	<i>immaculatus</i> .....	630
138	<i>goodell</i> .....	664, 2831	<i>ohiensis</i> .....	629
138	<i>ommata</i> .....	663, 2831	<i>reticulatus</i> .....	627
759	<i>parva</i> .....	665, 2831	<i>vermiculatus</i> .....	627, 2827
759	<i>venusta</i> .....	665, 2831	Lucius, <i>Atractosteus</i> .....	111
2209	lucasana, <i>Sphyrna</i> .....	826	<i>Esox</i> .....	628
2209	lucasanus, <i>Thalassoma</i> .....	1607, 2859	<i>americanus</i> .....	626
2278	lucasanus, <i>Chlorichthys</i> .....	1607	Lucius .....	628
2278	<i>Julis</i> .....	1607, 1608	<i>Ptychocheilus</i> .....	225
2659	lucasi, <i>Ceratocottus</i> .....	1940	Lucky Proach .....	1971
2660	lucayanum, <i>Asymmetron</i> .....	4	Lucretia, <i>Aboma</i> .....	2241
1931	Lucius vorax .....	628	<i>Gobius</i> .....	2868
508	lucens, <i>Ceratichthys</i> .....	321	ludibunda, <i>Cliola</i> .....	273
1229	<i>Dacentrus</i> .....	1496	<i>Cyprinella</i> .....	373
0, 157, 159	luciae, <i>Fundulus</i> .....	654	ludibundus, <i>Notropis</i> .....	273
158	<i>Haplochilus</i> .....	655	Lugger, Stone .....	181
158	<i>Hydrargyra</i> .....	655	lugubris, <i>Curanx</i> .....	924
158	<i>Zygonectes</i> .....	655	<i>Cyprinella</i> .....	274
157	lucida, <i>Ethioprora</i> .....	565	<i>Malacoctenus</i> .....	2357
157	lucidus, <i>Argyrosomus</i> .....	470	<i>Melanphaes</i> .....	842
157	<i>Coregonus</i> .....	471	<i>Myxodes</i> .....	2357
158	<i>Luxilus</i> .....	299	<i>Plectronus</i> .....	842
158	<i>Notemigonus</i> .....	299	lugubrosus, <i>Esox</i> .....	628
159	<i>Salmo</i> ( <i>Coregonus</i> ) .....	471	luitpoldii, <i>Characodon</i> .....	2832
156	<i>Stolephorus</i> .....	447	lumbicus, <i>Morone</i> .....	342
155	Lucifer .....	591	<i>Myrophis</i> .....	371
156	lucifer, <i>Linophryne</i> .....	2734	Lumpetinae .....	2349
569	Lucifuga .....	2501	Lumpenus .....	2435, 2436
1756	<i>dentatus</i> .....	2500	<i>aculeatus</i> .....	2433
	<i>subterraneus</i> .....	2501	<i>anguillaris</i> .....	2436

	Page.		Page.
<i>Lumpenus fabricii</i> .....	2437	<i>lutens, Genyatremus</i> .....	1342
<i>Lumpetæformis</i> .....	2438	<i>Lutianus</i> .....	1343
<i>mackayi</i> .....	2436	<i>Notrus</i> .....	141
<i>maculatus</i> .....	2433	<i>Rhinichthys</i> .....	307
<i>medius</i> .....	2435	<i>Lutianidæ</i> .....	1241
<i>nubilus</i> .....	2438	<i>Lutianinæ</i> .....	1242
<i>oculeatus</i> .....	2433	<i>Lutianus</i> .....	1247
<i>lumpenus, Blennius</i> .....	3438	<i>argentiiventris</i> .....	1261
<i>Clinus</i> .....	2438	<i>colorado</i> .....	1268
<i>Sticheus</i> .....	2438	<i>guttatus</i> .....	1269
<i>Lumpfish</i> .....	2096	<i>lutens</i> .....	1343
<i>Lump Sucker</i> .....	2094, 2096	<i>novemfasciatus</i> .....	1253
<i>Lumpus</i> .....	2096	<i>stearnsi</i> .....	1256
<i>anglorum</i> .....	2097	<i>Lutipinnis, Hydrophlox</i> .....	287
<i>spinosus</i> .....	2099	<i>Minuilus</i> .....	287
<i>vulgaris</i> .....	2097	<i>Notropis</i> .....	286
<i>lumpus, Cyclopterus</i> .....	2096, 2097	<i>Opisthopterus</i> .....	437
<i>Luna, piscis</i> .....	1754	<i>Pristigaster</i> .....	437
<i>Poz</i> .....	1753	<i>lutjanoides, Lutjanus</i> .....	1261
<i>luna, Caranx</i> .....	927	<i>Neomenis</i> .....	1261
<i>Lampris</i> .....	954	<i>Ocyurus</i> .....	1261
<i>Pomotis</i> .....	1006	<i>Lutjanus acutirostris</i> .....	1259
<i>Zeus</i> .....	955	<i>ambiguus</i> .....	1272
<i>lunaris, Orthragoriscus</i> .....	1754	<i>analis</i> .....	1267
<i>lunata, Echeneis</i> .....	2269	<i>aratus</i> .....	1274
<i>lunaticus, Dactyloscopus</i> .....	2302	<i>argentiiventris</i> .....	1261
<i>lunatus, Platophrys</i> .....	2665	<i>arnillus</i> .....	1279
<i>Pleuronectes</i> .....	2666	<i>anbrieti</i> .....	1271
<i>Rhinichthys</i> .....	308	<i>aurorbens</i> .....	1278
<i>atronsasus</i> .....	308	<i>aya</i> .....	1265
<i>Rhomboidichthys</i> .....	2666	<i>blackfordi</i> .....	1265
<i>Lune, Poisson</i> .....	954	<i>brachypterus</i> .....	1268
<i>Tetrodon</i> .....	1754	<i>buccanella</i> .....	1262
<i>luniscutis, Arius</i> .....	125	<i>caballeroto</i> .....	1257
<i>Aspistor</i> .....	2763	<i>campechianus</i> .....	1265
<i>Selenaspis</i> .....	125	<i>caxis</i> .....	1260
<i>Tachisurus</i> .....	125	<i>cayennensis</i> .....	1404
<i>lunulatus, Epinephelus</i> .....	1150	<i>chrysurus</i> .....	1276
<i>Lutjanus</i> .....	1158	<i>colorado</i> .....	1268
<i>Mustelus</i> .....	28	<i>cyanopterus</i> .....	1255
<i>Rhomboidichthys</i> .....	2666	<i>cynodon</i> .....	1255
<i>Serranus</i> .....	1150	<i>enbera</i> .....	1255
<i>lupus, Ancyurus</i> .....	137	<i>dentatus</i> .....	1255
<i>Anarhichas</i> .....	2447	<i>griseus</i> .....	1257
<i>Ictalurus</i> .....	137	<i>guttatus</i> .....	1269
<i>Pimelodus</i> .....	137	<i>inermis</i> .....	1275
<i>lusca, Cyclothone</i> .....	582	<i>jocu</i> .....	1258
<i>lusitaniensis, Vandellius</i> .....	887	<i>lunulatus</i> .....	1158
<i>lusca, Anguilla</i> .....	348	<i>lutjanoides</i> .....	1261
<i>luteovinctum, Ethicostoma</i> .....	1080	<i>mahogoni</i> .....	1273
<i>lutescens, Chætodon</i> .....	1680	<i>melanurus</i> .....	1276
<i>Kyphosus</i> .....	1388	<i>novemfasciatus</i> .....	1253
<i>Pimelepterus</i> .....	1389	<i>oiano</i> .....	1273
<i>luteum, Hæmulon</i> .....	1304	<i>pacificus</i> .....	1253
<i>lutens, Centropomus</i> .....	1024	<i>prieto</i> .....	1254
<i>Gadus tomcodus</i> .....	2540	<i>profundus</i> .....	1264



Page.		Page.		Page.
1342	<i>Lutjanus purpureus</i> .....	1264	<i>Lycodapus dermatinus</i> .....	2492
1343	<i>rosaceus</i> .....	1267	<i>extensus</i> .....	2479
144	<i>stearnsi</i> .....	1257	<i>ferasfer</i> .....	2493
307	<i>surinamensis</i> .....	1319	<i>parviceps</i> .....	2493
1241	<i>synagris</i> .....	1271	<i>Lycodes</i> .....	2461, 2462
1242	<i>torridus</i> .....	1264	<i>brevipis</i> .....	2467
1247	<i>triangulum</i> .....	1454	<i>coccineus</i> .....	2469
1261	<i>tridens</i> .....	1202	<i>concolor</i> .....	2463
1268	<i>trilobus</i> .....	1200	<i>diapterus</i> .....	2473
1269	<i>uninotatus</i> .....	1271	<i>digitatus</i> .....	2466
1343	<i>verres</i> .....	1583	<i>esmarkii</i> .....	2463
1253	<i>viridis</i> .....	1246	<i>frigidus</i> .....	2465
1256	<i>vivanus</i> .....	1264, 1265	<i>gracilis</i> .....	2465
287	<i>lutkeni</i> , <i>Exocoetus</i> .....	736	<i>mucosus</i> .....	2470
287	<i>Lutodeira</i> .....	414	<i>nebulosus</i> .....	2468
286	<i>lutrensis</i> , <i>Cliola</i> .....	272	<i>pacificus</i> .....	2460
437	<i>Leuciscus</i> .....	272	<i>paris</i> .....	2466
437	<i>Notropis</i> .....	271	<i>paxilloides</i> .....	2471
1261	<i>luxatus</i> , <i>Chasmodon</i> .....	183	<i>paxillus</i> .....	2471
1261	<i>Luxilius</i> .....	247	<i>perapicillum</i> .....	2465
1261	<i>occidentalis</i> .....	247, 2799	<i>polaris</i> .....	2460
1259	<i>Luxilus</i> .....	250, 254, 257, 281	<i>porifer</i> .....	2472
1272	<i>chickasawensis</i> .....	275	<i>reticulatus</i> .....	2465
1267	<i>chrysocephalus</i> .....	282	<i>rossi</i> .....	2465
1274	<i>dissimilis</i> .....	319	<i>seminudus</i> .....	2468
1261	<i>elongatus</i> .....	240	<i>terra-novæ</i> .....	2466
1279	<i>erythrogaster</i> .....	210	<i>turneri</i> .....	2469
1271	<i>interruptus</i> .....	282	<i>vablii</i> .....	2463
1278	<i>kentuckiensis</i> .....	279, 322	<i>verrillii</i> .....	2471
1265	<i>leptosomus</i> .....	250	<i>zoarchus</i> .....	2464
1265	<i>lucidus</i> .....	299	<i>Lycodidae</i> .....	2455
1268	<i>occidentalis</i> .....	247	<i>Lycodiura</i> .....	2456
1262	<i>roseus</i> .....	288	<i>Lycodontis</i> .....	392, 393
1257	<i>seco</i> .....	250	<i>castaneus</i> .....	2804
1265	<i>selene</i> .....	269	<i>chlevestes</i> .....	398
1260	<i>zonistius</i> .....	285	<i>conspersus</i> .....	397
1404	<i>luxoides</i> , <i>Cyprinuella</i> .....	274	<i>dovii</i> .....	397
1276	<i>lyadon</i> , <i>Oncorhynchus</i> .....	481, 483	<i>elaboratus</i> .....	398
1268	<i>Salmo</i> .....	483	<i>funebri</i> .....	396, 2804
1255	<i>Lycenchelys</i> .....	2469, 2470	<i>miliaris</i> .....	397
1255	<i>paxillus</i> .....	2471	<i>mordax</i> .....	395, 2805
1255	<i>porifer</i> .....	2471	<i>moringa</i> .....	395
1255	<i>verrillii</i> .....	2470, 2471	<i>obscuratus</i> .....	389
1257	<i>Lycengraulis</i> .....	451, 2811, 2816	<i>ocellatus</i> .....	399
1269	<i>grossidens</i> .....	451	<i>nigromargina-</i>	
1275	<i>poeyi</i> .....	2811	<i>tua</i> .....	399
1258	<i>lychnus</i> , <i>Myripristis</i> .....	847	<i>saxicola</i> .....	399
1158	<i>Lycia</i> .....	2461, 2463, 2468	<i>pictus</i> .....	2805
1261	<i>Lycocara</i> .....	2478	<i>polygonius</i> .....	394
1273	<i>parii</i> .....	2478	<i>sanctæ-helenæ</i> .....	397
1276	<i>Lycodalepis</i> .....	2468	<i>verrilli</i> .....	393
1253	<i>mucosus</i> .....	2470	<i>vicinus</i> .....	394
1273	<i>polaris</i> .....	2468	<i>vlrescens</i> .....	394
1253	<i>turneri</i> .....	2469	<i>Lycodonus</i> .....	2473
1253	<i>Lycodapodidae</i> .....	2491	<i>mirabilis</i> .....	2474
1264	<i>Lycodapus</i> .....	2492	<i>Lycodopsis</i> .....	2800

	Page.		Page.
Lycodopsis.....	2460	Macondaldia rostrata .....	617
crassilabris .....	2458	macellus, Prionistius .....	192-
crotalinus .....	2450	macer, Polyprosopus .....	51
pacifcus .....	2460	Machera .....	890
paucidens .....	2460	machete, Sardina .....	433
Lycolla .....	2869	machuta, Argentina .....	410
crassilabris .....	2869	Machyo .....	811
crotalina .....	2869	Machuelo .....	432
Lyconectes .....	2444	Machuto .....	811
aleutensis .....	2444	mackayi, Lumpenus .....	2436
Lyconoma .....	2474	Siphostoma .....	766
barbatum .....	2474	mackenzii, Luciostrutta .....	474
lynceum, Etheostoma .....	1075	Salmo .....	474
lynx, Pinelodus .....	138	Stenodus .....	474
Lyoliparis .....	2114, 2116, 2126	Mackerel, Chub .....	866
Lyomeri .....	404	Common .....	865
Lyopomi .....	600	Easter .....	866
Lyopsetta .....	2612	Horse .....	870
exilis .....	2612	Monterey Spanish .....	873
Lysophaera .....	1751	Snap .....	946
globosa .....	1751	Spanish, of England .....	866
lyricus, Ectenogobius .....	2225	Spanish .....	874
Gobius .....	2224	Thimble-eyed .....	866
lythrochloris, Xenotis .....	1003	Tinker .....	866
Lythirulon .....	1311	Yellow .....	921
flaviguttatum .....	1312	Mackerel-like Fishes .....	860
opalescens .....	1312	Mackerel Scads .....	907
Lythrurus .....	254, 258, 297	Shad .....	909
atrilpes .....	300	Shark .....	48, 49
cyanocephalus .....	300	Sharks .....	47
lythrurus, Notropis .....	300	Mackerels .....	863, 865
umbratilis .....	300	Frigate .....	867
Lythrypnus .....	2210, 2216, 2230	King of .....	1755
Macabi .....	411	Snake .....	883
Macana .....	341	Mackinaw Trout .....	504
macarellus, Caranx .....	909	maclura, Pteroplatea .....	86
Decapterus .....	909	Raia .....	87
Macarla .....	890	Mucolor .....	1247
macaskeyi, Ictalurus .....	318	macouni, Chauliodus .....	585
maclollandii, Bregmaceros .....	2526	macracanthum, Clethrasoma .....	1518
macondaldi, Acutomentum .....	1787	Pristipoma .....	1332
Conocara .....	457	macracanthus, Alutarius .....	1720
Cynoscion .....	1411	Heros .....	1519
Fundulus .....	651	Pomadasis .....	1332
Nannobranchium .....	563	macrocephala, Clupea .....	411
Notropis .....	284	Murena .....	348
Penopus .....	2521	macrocephalum, Etheostoma .....	1031
Salmo clarkii .....	2819	Macrocephalus .....	1117
mykiss .....	497	macrocephalus, Alvordius .....	1031
Sebastodes .....	1786	Clinus .....	2364
Zygonectes .....	651	Gadus .....	2541
Macondaldia .....	616	Gonocephalus .....	2184
alta .....	2826	Gunellus .....	2410
challengeri .....	617	Hadropterus .....	1031
longa .....	2826	Labrosomus .....	2364
		Mniertes .....	2364

Page.		Page.		Page.
617	macrocephalus, Percina .....	1031	macrops, Ballistes .....	1706
102-	Semotilus .....	222	Bathygadus .....	2560
51	macrocerus, Monacanthus .....	1713	Calamus .....	1350, 1534
890	macrocheilus, Catostomus .....	178	Citharichthys .....	2084
433	macrochir, Atheroandia .....	609	Conger .....	355
410	Halosaurus .....	610	Corvina .....	1428
811	Sebastolobus .....	1763	Corvula .....	1427, 1428
432	macrochirus, Eupomotis .....	1005	Gnathypops .....	2284
811	Hemirhamphus .....	723	Hippoglossina .....	2621
2436	Lepomis .....	1005	Opisthognathus .....	2284
786	Macrodon .....	330	Opisthopterus .....	437
474	malabaricus .....	330	Pristigaster .....	437
474	microlepis .....	330	Sclera .....	1428
866	Macrodonophis .....	380	macroptera, Conocara .....	457, 458
865	mordax .....	387	Gonlobatis .....	88
866	macroodus, Squalus .....	47	macropteron, Hypentellum .....	181
870	macrogenis, Cerna .....	1181	macropterus, Alepocephalus .....	458
873	macrognathum, Opisthognathus .....	2281	Ballistes .....	1707
946	Macrognathus .....	759	Centrarchus .....	988
866	brevirostris .....	723	Labrus .....	988
874	loricatus .....	110	Thynnus .....	871
866	macrognathus, Opisthognathus .....	2282	macroptus, Mulacoetemus .....	2357
866	macrolepidota, Anchovia .....	449	Myxodes .....	2357
921	Pecilia .....	641	Macroruhmphoside .....	758
860	macrolepidotum, Moxostoma .....	193	Macroshamphosus .....	759
907	macrolepidotus, Catostomus .....	194	scolopax .....	759
909	Chetodon .....	1677	macrorhinus, Acipenser .....	105
48, 49	Engraulis .....	449	macrorhynchus, Hemirhamphus .....	724
47	Fundulus hetero-		macrospila, Piramutana .....	155
863, 865	clitus .....	641	Macrostoma .....	554
867	Lenciscus .....	224	angustidens .....	555, 2826
1755	Notropis .....	299	brachychir .....	2826
883	Pleuronectes .....	2672	castaneum .....	556
504	Pogonichthys .....	223	caudispinosum .....	559, 2826
86	Stolephorus .....	449	margaritifera .....	555
87	macrolepis, Pontinus .....	1855	quercinum .....	554
1247	macrolema, Pimelodus .....	155	macrostoma, Cyprinella .....	274
585	macrolema, Bagrus .....	117	Hemulon .....	1297
1518	Nemipterus .....	1289	Sahn .....	481
1332	Polynemus .....	828	macrostomum, Hemulon .....	1296
1720	Synagris .....	1289	macrostomus, Acipenser .....	106
1519	macrophthalmus, Clupea .....	430	Notropis .....	274
1332	Harengula .....	430	Macrouridae .....	2561
411	macrophthalmus, Aprion .....	1280, 1281	Macrourinae .....	2562
348	Caranx .....	911	Macrounus .....	2581
1031	Centropristis .....	1281	acrolepis .....	2585
1117	Elastomus .....	1281	baiensis .....	2583
1031	Gobiosox .....	2335	bergii .....	2581
2364	Priacanthus .....	1238	clereus .....	2586
2541	Sardinella .....	430	fabrii .....	2582
2184	Scomber .....	867	holotrachys .....	2582
2410	macropona, Bollmannia .....	2239	lepturus .....	2584
1031	Centropristis .....	1206	rupestris .....	2582
2364	Diplectrum .....	1205	stelgidolepis .....	2585
2364	Macrops .....	1281	stromii .....	2579
	oculatus .....	1283	macrourus, Alopias .....	46

	Page.		Page.
Macrozoarces .....	2456	maculatus, Leptocephalus .....	2433
macrura, Congermurena .....	356	Liparis .....	2108
Macruropus .....	2581	Lumpenus .....	2433
macrurus, Ophisoma .....	357	Monoprius .....	1109
Macrurus .....	2581	Mullhypeneus .....	859
acrolepis .....	2585	Mullus .....	859
asper .....	2572	Nomus .....	950
carminatus .....	2580	Nothonotus .....	1077
carrilbeneus .....	2590	Notchynchus .....	17
dorsalis .....	2585	Notropis .....	29
firmisquamis .....	2576	Ostracion .....	1725
goodii .....	2572	Pimelodus .....	135, 155
(Nematonurra) mugnus ..	2574	Platypræcius .....	686
occa .....	2588	Plouronectes .....	2660
pectoralis .....	2574	Procerus .....	102
rupestris .....	2579	Psenes .....	951
scaphopsis .....	2590	Rhypticus .....	1231
slunius .....	2578	Scomber .....	867, 874
suborbitalis .....	2573	Scomberomorus .....	874, 875
macrurus, Oxydenticithys .....	385	Serranus .....	1153
macularius, Cyprinodon .....	674	Spherides .....	1733
baileyi .....	675	Spheroides .....	1733
maculata, Apogon .....	1109	Stichæus .....	2433
Belone .....	718	Upeneus .....	858
Hemitremia .....	259	maculicauda, Diabasis .....	1314
Lophopsetta .....	2660	Hæmulon .....	1314
Morone .....	1010	Orthostechus .....	1313
Muraena nigra .....	395	maculiter, Diodon .....	1747
Nerophis .....	774	Platophrys .....	2661
Percia .....	1153	Pleuronectes .....	2665
Sciæna .....	2198	maculiferus, Hypoplectrus .....	1192
maculaticeps, Boleosoma nigrum ..	1058	Rhomboidichthys .....	2665
maculatofasciatus, Paralabrax .....	1196	maculipinna, Chærojulis .....	1595
Serranus .....	1196	Halicheres .....	1595
maculatum, Boleosoma .....	1057, 1077	Iridio .....	1594
Cybium .....	874	Julis .....	1595
Cynoscion .....	1409	Platygllossa .....	1595
Etheostoma .....	1077	maculipinnis, Achirus .....	2698
maculatus, Aivordius .....	1032, 1034	Exocoetus .....	737
Anarrhichas .....	2446	Heros .....	1529
Apogon .....	1109	Isogomphodon .....	40
Anlostomus .....	754	Monochir .....	2698
Balistes .....	1707, 1708	Muraena .....	394
Bothus .....	2660	Solca .....	2698
Canthidermis .....	1706, 1707	Thyrsoidea .....	394
Clinus .....	2433	maculocinctus, Chætodon .....	1674
Cottus .....	1972	Sarothrodus .....	1674
Cryptacanthodes .....	2443	maculosa, Harengula .....	436
Ctenodon .....	2433	Lota .....	2550
Diodon .....	1746	Molva .....	2551
Dormitator .....	2196, 219	Muraena .....	382
Galeocerdo .....	32	Thalassop'ryne .....	2324
Galeus .....	32	maculoseriatus, Chirus .....	1808
Gasteropelecus .....	338	nuculoseriatus, Diodon .....	1748
Hadropterus .....	1031, 1034	maculosum, Hæmulon .....	1295
Heptanchias .....	18	maculosus, Acentrolophus .....	963

Page.		Page.		Page.
2433	<i>maenulosus</i> , <i>Acipenser</i> .....	106	<i>makua</i> , <i>Ranzania</i> .....	1755
2168	<i>Catostomus</i> .....	181	<i>malabarica</i> , <i>Elacate</i> .....	948
2432	<i>Centrarchus</i> .....	991	<i>malabarius</i> , <i>Macrodon</i> .....	330
1109	<i>Contridermichthys</i> .....	2014	<i>Malacanthidae</i> .....	2274
859	<i>Epinephelus</i> .....	1158	<i>Malacanthinae</i> .....	2275
859	<i>Gadus</i> .....	2551	<i>Malacanthus</i> .....	2275
950	<i>Nomeus</i> .....	950	<i>plumieri</i> .....	2275
1077	<i>Oligocottus</i> .....	2013	<i>trachinus</i> .....	2276
17	<i>Paralichthys</i> .....	2026	<i>Malachorhinus</i> .....	66
259	<i>Pimphales</i> .....	217	<i>Malucocephalus</i> .....	2560
1725	<i>promelas</i> .....	217	<i>occidentalis</i> .....	2570
135, 155	<i>Pleuronectes</i> .....	2026	<i>Malucoctenus</i> .....	1994
686	<i>Serpens marinus</i> .....	382	<i>zonurus</i> .....	1994
2660	<i>Serranus</i> .....	1159	<i>Malucoctenus</i> .....	3356
102	<i>madeirensis</i> , <i>Ceratoscopelus</i> .....	557	<i>biguttatus</i> .....	2360
951	<i>Scopelus</i> .....	557	<i>bimaculatus</i> .....	2358
1234	<i>Mademoiselle</i> .....	1433	<i>delalandi</i> .....	2358
867, 874	<i>maderaspuntensis</i> , <i>Butirius</i> .....	415	<i>gillii</i> .....	2358
874, 875	<i>maderensis</i> , <i>Helicolenus</i> .....	1837	<i>lugubris</i> .....	2357
1153	<i>Madregal</i> .....	905	<i>macropus</i> .....	2357
1732	<i>Mad Tom</i> .....	147	<i>ocellatus</i> .....	2356, 2869
1733	<i>Toms</i> .....	144	<i>varius</i> .....	2357
2433	<i>meandricus</i> , <i>Caularchus</i> .....	2328	<i>versicolor</i> .....	2359
858	<i>Lepidogaster</i> .....	2328	<i>Malucostolde</i> .....	592
1314	<i>Menidia</i> .....	1364	<i>Malucosteus</i> .....	592
1314	<i>Mantinae</i> .....	1364	<i>niger</i> .....	593
1312	<i>magdalenae</i> , <i>Curimata</i> .....	332	<i>Malapterinae</i> .....	1572
1747	<i>Otolithus</i> .....	1410	<i>malariaoides</i> , <i>Aspidophorus</i> .....	2054
2664	<i>Paralichthys</i> .....	2872	<i>Muleta cariden</i> .....	423
2665	<i>Selena</i> .....	1420	<i>maliger</i> , <i>Sebasticthys</i> .....	1823
1192	<i>magistralis</i> , <i>Epinnula</i> .....	880	<i>Sebastes</i> .....	1822
2665	<i>magnioculis</i> , <i>Ophichthys</i> .....	385	<i>malleus</i> , <i>Squalus</i> .....	45
1595	<i>Ophichthys</i> .....	385	<i>Zygana</i> .....	45
1595	<i>Scytalophis</i> .....	385	<i>Mallotus</i> .....	520
1594	<i>magnus</i> , <i>Macrurus</i> ( <i>Nematonurus</i> ).....	2574	<i>villosus</i> .....	520
1595	<i>Mahogany Snapper</i> .....	1272	<i>Malmu</i> .....	507, 508
1595	<i>mahogoni</i> , <i>Lutjanus</i> .....	1273	<i>malma</i> , <i>Sahno</i> .....	508
2698	<i>Mesoprion</i> .....	1273	<i>Salvelinus</i> .....	507, 2823
737	<i>Neomenis</i> .....	1272	<i>Malthua</i> .....	2736
1529	<i>Mail-cheeked Fishes</i> .....	1750	<i>angusta</i> .....	2738
40	<i>Mailed Gurnards</i> .....	2176	<i>longirostris</i> .....	2737
2698	<i>mainensis</i> , <i>Gasterosteus</i> .....	745	<i>nasuta</i> .....	2737
394	<i>Maire d'Amplora</i> .....	555	<i>notata</i> .....	2737
2698	<i>majalis</i> , <i>Cobitis</i> .....	639	<i>truncata</i> .....	2738
394	<i>Fundulus</i> .....	639, 2827	<i>vespertilio</i> .....	2737
1674	<i>Hydrargyra</i> .....	639	<i>Malthe</i> .....	2736
1674	<i>Majarra</i> , <i>Ralada</i> .....	1561	<i>cubifrons</i> .....	2738
436	<i>major</i> , <i>Actinoclin</i> .....	2128	<i>oliter</i> .....	2739
2550	<i>Blepharis</i> .....	932	<i>Mammoth Cave Blindfish</i> .....	706
2551	<i>Cyclopterus liparis</i> .....	2128	<i>Mammy</i> .....	205
382	<i>Liparis</i> .....	2127	<i>managuensis</i> , <i>Heros</i> .....	1533
2324	<i>Ptychocheilus</i> .....	225, 2797	<i>Pimelodus</i> .....	153
1868	<i>Major, Sergeant</i> .....	1561	<i>Rhamdia</i> .....	153
1748	<i>Makaira</i> .....	890	<i>manatia</i> , <i>Raia</i> .....	93
1295	<i>nigricans</i> .....	891	<i>manatinus</i> , <i>Barathrodemus</i> .....	2517
963	<i>makaira</i> , <i>Xiphias</i> .....	891	<i>Maucalias</i> .....	2729

	Page.		Page.
Mancalis shufeldti .....	2730	marginatum, Ophidium .....	2489
uranoscopus .....	2729	marginatus, Bodianus .....	1174
Mancho, Cibi .....	919	Brosmius .....	2502
Man-eater Sharks .....	50	Brosomphycis .....	2502
maniglicola, Gobit s. ....	2220	Carenx .....	922
mango, Polynemus .....	830	Chaetodon .....	1562
Mangrove Grouper .....	1171	Chronis .....	1546
Minnow .....	643	Cottus .....	1966
Snapper .....	1255	Dinematichthys .....	2502
Manitou Darter .....	1027	Halias .....	2502
manitou, Percina caprodes .....	1928	Hemirhamphus .....	723
Manjua .....	443	Lepomis .....	1003
Manjuari .....	111	Neobythites .....	2513
manjuari, Lepidosteus .....	111	Noturus .....	147
mannii, Zygonectes .....	664	Phycis .....	2555
Manta .....	92	Pimelodus .....	135
americana .....	93	Pomotis .....	1003
birostris .....	92	Serranus .....	1154
manta, Cephaloptera .....	93	Symphurus .....	2706
Mantararia Colorado .....	2754	Maria Molle .....	1552
Mantida .....	91, 2756	Marian .....	852
manus, Gadus .....	2154	marianus, Flammeo .....	2871
Mapo .....	2216	Holocentrus .....	852, 2871
Guavina .....	2196	Maria-prieta .....	1319
mapo Gobius .....	2218	Marina, Perca .....	1761
maraldi, Gadus .....	2546	cauda nigra .....	1305
Uraleptus .....	2545	gibbosa .....	1295
maregravii, Aulastome .....	757	pinnis .....	1259
Coryphæna .....	953	sectatrix .....	1388
Rhinobatus .....	63	venenosa .....	1172
marconis, Hybopsis testivalis .....	516	Umbla minor .....	823
Mareño, Pargo .....	1252	marinus, Ailurichthys .....	118
Margaret, Bastard .....	1297	Esox .....	714
Margaret Grunt .....	1295	Faber .....	1668
margarita, Clinostomus .....	241	Felichthys .....	118
Echiostoma .....	589	Petromyzon .....	10
Leuciscus .....	241	camtschaticus .....	2745
margaritaceus, Entomacrodus .....	2398	dorsatus .....	10
Sairias .....	2399	unicolor .....	10
margaritatus, Batrachus .....	2323	Raja diabolus .....	93
Porichthys .....	2322	Sobastos .....	1760, 1761
margaritifera, Heros .....	1520	viviparus .....	1761
Notoscopelus .....	555	Serpens maculosus .....	382
Serranus .....	1156	Silurus .....	118
margaritifera, Cichlasoma .....	1519	Tylosurus .....	714
Hamulon .....	1312	Mariposas .....	953, 954
Macrostoma .....	555	marmorata, Amia .....	113
margaritus, Phoxinus .....	241	Pteroplatea .....	87
Squalius .....	241	Unibranchapertura .....	342
margarotis, Enneacanthus .....	994	marmoratum, Lepophidium .....	2482, 2483
Margate-fish .....	1295	marmoratus, Ameiurus .....	141
marginalis, Hippocampus .....	778	nebulosus .....	141
marginata, Aphoristia .....	2706	Antennarius .....	2717
Rissola .....	2489	Auchenopterus .....	2371
Thyrsoidea .....	394	Cottus .....	1983
Uranidea .....	1965		

	Page.		Page.
marmoratus, Cromnobates .....	2371	matejuelo, mphiprion .....	849
2489 Hemitripterus .....	1889, 2022	mathematicus, Tetrodon .....	1728
1174 Pimelodus .....	141	Mathemeg .....	137
2502 Rhinichthys .....	306	matoides, Acanthurus .....	1693
2502 Rivulus .....	663, 2830	matubaræ, Sebastodes .....	1833
922 Scorpænichthys .....	1889	Mattowacca .....	425
1562 Spheroides .....	1734	matowacca, Clupea .....	426
1546 Symbranchus .....	342	matutinus, Alburnellus .....	301
1966 Tirus .....	537	Minniles .....	301
2502 marmorca, Rabula .....	391	Notropis .....	301
2502 marmoreum, Siphostoma .....	768	umbratilis .....	301
723 marmoreus, Blennius .....	2381	matubaræ, Sebastodes .....	1796
1003 Gymnothorax .....	391	mauricei, Argyrosus .....	996
2513 Murenophis .....	391	mauritii, Chaetodon .....	1562
147 Syngnathus .....	768	Eleotris .....	950
2555 Marsipobranchii .....	4	Grammistes .....	1323
135 marstoni, Salmo .....	516	manrolici, Scopelus .....	577
1003 Salvelinus oquassa .....	515	Mauroleicida .....	576
1154 martii, Cyprinodon .....	675	Maurolicus .....	576
2706 martii, Pristigaster .....	438	amethystino-punctatus .....	577
1552 Martin Pescador .....	2724	attenuatus .....	577
852 martinica, Atherina .....	795	borealis .....	577
2871 Kirtlandia .....	795	mulleri .....	577
2, 2871 Menidia .....	795	pennanti .....	577
1319 Spicara .....	1364, 1365	tripunctulatus .....	578
1761 martinicensis, Ayoalon .....	1228	maxillaris, Lotella .....	2546
1307 Holanthias .....	1228	Murenoides .....	2418
1295 Menticirrhus .....	1473	Maxillingua .....	327
1259 Nerephis .....	774	maxillingua Cyprinus .....	327
1388 Novacula .....	1617	Exoglossum .....	327
1172 Novaenlichthys .....	1616	maxillosa, Gnathypops .....	2284
823 Ocyanthias .....	1228	maxillosus, Gnathypops .....	2284
118 Odontanthias .....	1228	Opisthognathus .....	2284
714 Umbrina .....	1474	Rhinichthys .....	307
1668 Vomer .....	934	maxima, Selache .....	51
118 Xyrichthys .....	1617	maximus, Cetorhinus .....	51
10 marthicus, Gobius .....	2236	Hippoglossus .....	2612
2745 Smaris .....	1365	Labrus .....	1580
10 Upeneus .....	859	Lachnolemus .....	1580
10 Masamacush .....	504	Lachnolaimus .....	1579
93 Mascalongus .....	625, 626, 629	Selachus .....	51
60, 1761 maschalespilos, Searus .....	1642	Squalus .....	51
1761 Sparisoma .....	1641	maycus, Salmo omisco .....	487
382 Mackinongy .....	629	Mayfish .....	639
118 masoni, Salmo .....	501	Maynea brunnea .....	2476
714 masquihongy, Esox .....	629	pusilla .....	2476
953, 954 immaculatus .....	630	May Sucker .....	199
113 Lucius .....	629	mazatlanæ, Seriola .....	904
87 immaculatus .....	630	Solea .....	2699
342 ohlensis .....	629	mazatlanum, Hamulon .....	1314
82, 2483 mussachusettensis, Monacanthus .....	1716	mazatlanus, Achirus .....	2698
141 mulliensis, Scorpæna .....	1139	McCloud River Rainbow Trout .....	502
141 Masticura .....	59, 79	meadæ, Ulocentra .....	2852
2717 matejuelo .....	848	meanyi, Ruscarius .....	1908
2371 Blanco .....	2275, 2276	Meda .....	328
1983 Real .....	410	fulgida .....	329

	Page.		Page.
media, Palometa .....	2849	megalops, Opsopoeodus .....	248
Medialuna .....	1390, 1391	Pinnelodus .....	135
californiensis .....	1391	Trycherodon .....	249
Medialunas .....	1390	megalotis, Ichthelis .....	1069
melocri, Clupea .....	426	Lepomis .....	1002
mediocris, Pomolebus .....	425	Megaperca .....	1187
medirostris, Acipenser .....	104	ischinagi .....	1188
mediterranea, Chimera .....	95	Megaphalus .....	2329
Laurida .....	537	megastoma, Grystes .....	1012
Sarda .....	872	Opisthognathus .....	2282
mediterraneus, Hoplostethus .....	837	megastomus, Catostomus .....	181
Lopharis .....	947	Melanphaea beanii .....	843
Scomber .....	872	crassiceps .....	843
Sternoptyx .....	604	cristiceps .....	844
Thynnus .....	870	lugubris .....	842
medius, Anisarchus .....	2436	Melanphaima .....	838
Calamus .....	1355	melampterus, Salmo .....	489
Centropomus .....	1120	melampygus, Caranx .....	925
Clinus .....	2436	Melanichthys .....	1381, 1711
Grammateus .....	1356	melanocephalus, Plargyrus .....	217
Lampemus .....	2435	Melanocetina .....	2728
Myxocephalus .....	1983	Melanocetus bispinosus .....	2734
Rhombus .....	967	(Liocetus) murrayi .....	2734
Stichans .....	2436	melanochir, Julis .....	1609
Medregal .....	904	melanochira, Belone .....	716
medusicola, Caranx .....	924	melanogaster, Pleuronectes .....	2620
medusophagus, Schedophilus .....	970	Paecilia .....	696
meeki, Hybopsis .....	317	Melanogrammus .....	2542
megacephalus, Calamus .....	1350, 1351	aglefinus .....	2542, 2543
Chitonotus .....	1891	melanopsis, Diodon .....	1746
Iccius .....	1891	melanopleurus, Haplochilus .....	660
Megaderus .....	402	melanopogon, Cichlasoma .....	1523
megalaspis, Acipenser .....	105	Heros .....	1523
megalepis, Doratonotus .....	1912	melanopoma, Polynemus .....	831
Megalobrycon .....	337	melanops, Calliurus .....	992
Megalocottus .....	1987	Catostomus .....	187
laticeps .....	1988	Dionda .....	216
platycephalus .....	1987	Haplochilus .....	682
megulodon, Pristis .....	61	Hyboganthus .....	216
Megalopinae .....	408	Ichthelis .....	996
Megalops atlanticus .....	409	Minytrema .....	187
cepedina .....	416	Sebaste .....	1783
elongatus .....	409	Sebastodes .....	1782, 1783
notata .....	432	Zygonecetes .....	682
oglina .....	432	melanopterum, Pristipoma .....	1319
thrissoides .....	409	melanopterus, Balistes .....	1707
megalops, Alburnellus .....	291	melanopus, Arius .....	132
Alburnus .....	291	Tachysurus .....	132, 274
Clupea .....	426	melanorhina, Plectropoma .....	1192
Cyprinus .....	282	melanospilum, Diagramma .....	1321
Ditrema .....	1502	melanostictus, Hippoglossoides .....	2618
Eunichthys .....	1502	Paettichthys .....	2618
Holeconotus .....	1502	Melanostigma .....	2478
Micropogon .....	1463	gelatinosum .....	2479
Minnilus .....	291	pammelas .....	2479, 2869
Notropis albeolus .....	284	Melanostigmatina .....	2456



Page.		Page.		Page.
248	melanostomus, Sebastodes .....	1803	Menidia guatemalensis .....	801, 2840
135	melanotilus, Holacanthus .....	1728	martinica .....	795
249	melanotis, Limnamuriens .....	402	menida .....	800
1069	Murana .....	401	notata .....	2840
1002	Pseudojulis .....	1605	notata .....	800, 2840
1137	Scarus .....	1638	pachylepis .....	801, 2840
1138	melanotus, Catostomus .....	206, 218, 322	peninsule .....	797
2329	Melanura .....	623	sardina .....	799
1012	annulata .....	624	vagrans .....	795
2282	melanurum, Cichlasoma .....	1523	laciniata .....	795
181	Heanlon .....	1302, 1303	menidia, Argentina .....	443
843	Lemonema .....	2557	Atherina .....	801
843	Perca .....	1303	Atherinichthys .....	800
844	melanurus, Careproctus .....	2135	Menidia .....	800
842	Cyprinus .....	282	notata .....	2840
838	Exocoetus .....	735, 736	Menominee Whitefish .....	465
483	Heros .....	1524	menona, Fundulus diaphanus .....	645
925	Lutjanus .....	1276	mentalis, Platypocilus .....	686
381, 1711	Rutilus .....	193	Menticirrus .....	1469, 1470
217	melapleura, Pectilia .....	660	alburnus .....	1475
2728	melapleurus, Fundulus .....	659	americanus .....	1474
2734	melas, Ameiurus .....	141	elongatus .....	1476
2734	Silurus .....	141	littoralis .....	1477
1609	meleagris, Murana .....	399	martinicensis .....	1473
716	Priodonophis .....	399	nasus .....	1472, 1473
2630	Rhinichthys .....	308	nebulosus .....	1475
696	atronasus .....	308	panmensis .....	1473
2542	Meletta .....	424	saxatilis .....	1475
2542, 2543	luberfatis .....	433	simus .....	1472
1746	suerii .....	425	undulatus .....	1476
660	venosa .....	426	Mentiperca .....	1208, 1209, 1214
1523	Melichthys .....	1711	lucioperca .....	1216
1523	bispinosus .....	1711	mento, Ballistes .....	1710
831	picus .....	1711	Bramopsis .....	1502
992	ringens .....	1711	Chanos .....	415
187	Melletes .....	1932	Paraliparis .....	2142
216	papilio .....	1932	Xanthichthys .....	1710
682	mellissil, Congromurana .....	356	meitzelii, Chironectes .....	7224
216	Membras .....	789	Serranus .....	1154
996	membras, Clupea .....	421	merkil, Coregonus .....	470
187	Menephorus .....	1143, 1146	meridionalis, Amlurus .....	135
1783	dubius .....	1147	Cottus .....	1951
1782, 1783	punctiferus .....	1147	Ictalurus .....	135
682	menhaden, Alosa .....	434	Ictolobus .....	164
1319	Clupea .....	434	Sclerognathus .....	164
1707	Gulf .....	434	Merlangus .....	2529
132	Menhadens .....	433	leptocephalus .....	2535
132, 2784	Menidia .....	443, 796, 2840	polaris .....	2534
1192	audens .....	798	productus .....	2531
1321	beryllina .....	797	purpureus .....	2535
2618	brasiliensis .....	801	merlangus, Anoplopoma .....	1862
2618	clara .....	801	Merluccidae .....	2529
2478	dentex .....	801	Merluccius .....	2520
2479	gilberti .....	798	ambiguus .....	2530
2479, 2869	gracilis .....	797	argentatus .....	2530
2456	beryllina .....	797	bilinearis .....	2530, 2531

	Page.		Page.
<i>Merluccius esculentus</i> .....	2530	<i>Mesoprion linea</i> .....	1260
<i>linnaei</i> .....	2530	<i>litura</i> .....	1258
<i>merluccius</i> .....	2530	<i>mahogoni</i> .....	1273
<i>productus</i> .....	2531	<i>oianco</i> .....	1273
<i>sinuatus</i> .....	2530	<i>pacificus</i> .....	1253
<i>smiridus</i> .....	2530	<i>pargus</i> .....	1255
<i>vulgaris</i> .....	2530	<i>profundus</i> .....	1263
<i>Merluccius, Gadus</i> .....	2530	<i>ricardi</i> .....	1273
<i>Merluccius</i> .....	2530	<i>rosaceus</i> .....	1267
<i>Merluccius albidus</i> .....	2531	<i>sobra</i> .....	1266
<i>attenuatus</i> .....	2546	<i>uninotatus</i> .....	1271
<i>lanatus</i> .....	2530	<i>vivanus</i> .....	1263
<i>Merlus</i> .....	2529	<i>vorax</i> .....	1281
<i>merlus, Gadus</i> .....	2530	<i>mesops, Arins</i> .....	123
<i>Mero</i> .....	1154, 1162	<i>Bagrus</i> .....	123
<i>de lo Alto</i> .....	1161	<i>Sciadeichthys</i> .....	123, 2760
<i>Merou</i> .....	1154, 1780	<i>Tachisurus</i> .....	123
<i>merriami, Empetrichthys</i> .....	667	<i>Mesopus</i> .....	524
<i>mertensii, Cottus</i> .....	1986	<i>olidus</i> .....	525
<i>meru, Holocentrus</i> .....	1154	<i>mesotrema, Asternotremia</i> .....	787
<i>Merulius</i> .....	2148, 2149, 2156	<i>metallica, Agosiu</i> .....	314
<i>Merus</i> .....	1148	<i>Echeneis</i> .....	2270
<i>merus, Centropristis</i> .....	1162	<i>Heterandria</i> .....	687
<i>Epinephelus</i> .....	1162	<i>metallicus, Girardinus</i> .....	687
<i>mesæum, Boleosoma nigrum</i> .....	1059	<i>Notrops</i> .....	297
<i>mesæus, Pœclichthys</i> .....	1059	<i>metamorensis, Limia</i> .....	700
<i>mesogaster, Exocoetus</i> .....	729	<i>Metoponops</i> .....	2678
<i>Parexocetus</i> .....	728	<i>cooperi</i> .....	2680
<i>Mesogobius</i> .....	2210	<i>Metrogaster</i> .....	1498
<i>Mesogonistius</i> .....	994	<i>lincolatus</i> .....	1499
<i>chatodon</i> .....	995	<i>menlini, Diodon</i> .....	1748
<i>Mesoprion</i> .....	1247	<i>Me Waru</i> .....	1829
<i>ambiguus</i> .....	1272	<i>Tokenoko</i> .....	1829
<i> analis</i> .....	1266	<i>Mexican Sole</i> .....	2698
<i>aratus</i> .....	1274	<i>mexicana, Pœcilia</i> .....	692
<i>argentiventris</i> .....	1261	<i>mexicanum, Dorosoma</i> .....	416
<i>arnillo</i> .....	1279	<i>Myctophum</i> .....	563
<i>aurorubens</i> .....	1278	<i>Nannobranchium</i> .....	563
<i>aurrovittatus</i> .....	1276	<i>mexicanus, Amblyopus</i> .....	2264
<i>aya</i> .....	1264	<i>Awaous</i> .....	2237
<i>buccanella</i> .....	1262	<i>Centropomus</i> .....	1121
<i>caballerote</i> .....	1257	<i>Chatoessus</i> .....	416
<i>campechanus</i> .....	1265	<i>Chronophorus</i> .....	2237
<i>caudanotatus</i> .....	1262	<i>Gerres</i> .....	1380
<i>caxis</i> .....	1260	<i>Gobius</i> .....	2237
<i>chrysurus</i> .....	1276	<i>Mugil</i> .....	813
<i>cyanopterus</i> .....	1255	<i>Pempheris</i> .....	978
<i>cynodon</i> .....	1255, 1260	<i>Saurus</i> .....	538
<i>dentatus</i> .....	1279	<i>Tetragonopterus</i> .....	355
<i>elegans</i> .....	1278	<i>miarchus, Stolephorus</i> .....	441
<i>flavescens</i> .....	1260	<i>Michigan Clisco</i> .....	469
<i>griseus</i> .....	1257	<i>Gruyling</i> .....	518
<i>guttatus</i> .....	1269	<i>Herrling</i> .....	469
<i>inermis</i> .....	1275	<i>Kieyo</i> .....	469
<i>isodon</i> .....	1267	<i>Micristius</i> .....	633
<i>jocu</i> .....	1258	<i>Micristodus</i> .....	52

Page.		Page.		Page.
1260	<i>Micristodus punctatus</i> .....	52	<i>microperca</i> , <i>Etheostoma</i> .....	1104
1258	<i>microcephala</i> , <i>Cynoglossa</i> .....	2655	<i>Microphis</i> .....	773
1273	<i>Platessa</i> .....	2654	<i>mic. ophthalmum</i> , <i>Hæmulon</i> .....	1296
1273	<i>microcephalus</i> , <i>Gasterosteus wil-</i>		<i>microphthalmus</i> , <i>Dormitator</i> .....	2198
1253	<i>liamsoni</i> .....	751	<i>Heros</i> .....	1536
1255	<i>Pleuronectes</i> .....	2654	<i>Tetragonopterus</i> .....	334
1263	<i>Somniosus</i> .....	57	<i>Micropogon</i> .....	1461
1273	<i>Squalus</i> .....	57	<i>altiplinnis</i> .....	1464
1267	<i>Microdesmus</i> .....	2450	<i>argenteus</i> .....	1463
1266	<i>dipus</i> .....	2450	<i>costatus</i> .....	1462
1271	<i>retropinnis</i> .....	2450	<i>ectenes</i> .....	1463
1263	<i>microdon</i> , <i>Cyclothone</i> .....	582, 2826	<i>furnieri</i> .....	1462
1281	<i>Gobius</i> .....	2227	<i>lineatus</i> .....	1461
123	<i>Gonostoma</i> .....	582	<i>megalops</i> .....	1463
124	<i>Osmerus</i> .....	521	<i>opercularis</i> .....	1461
123, 2760	<i>Pseudo-riakis</i> .....	27	<i>undulatus</i> .....	1461
123	<i>Microdonophis</i> .....	381	<i>micropogon</i> , <i>Ceratichthys</i> .....	323
524	<i>Microgadus</i> .....	2538	<i>microps</i> , <i>Agonostomus</i> .....	820
525	<i>proximus</i> .....	2530	<i>Atherina</i> .....	791
787	<i>toncod</i> .....	2540	<i>Belone</i> .....	712
314	<i>microgathus</i> , <i>Lonchomisthus</i> .....	2287	<i>Callinurus</i> .....	996
2270	<i>Opisthognathus</i> .....	2287	<i>Caroharias</i> .....	40
687	<i>Microgobius</i> .....	2242	<i>Caulolatilus</i> .....	2277
687	<i>cyclolepis</i> .....	2247	<i>Corvina</i> .....	1445
297	<i>eulepis</i> .....	2244	<i>Cottunculus</i> .....	1992
700	<i>gulosus</i> .....	2243	<i>Dajaus</i> .....	820
2678	<i>signatus</i> .....	2246	<i>Nebris</i> .....	1417
2680	<i>thalassinus</i> .....	2245	<i>Otolithus</i> .....	1415
1498	<i>microlepidota</i> , <i>Gila</i> .....	207	<i>Pagellus</i> .....	1355
1499	<i>Microlepidotus</i> .....	1341	<i>Rhypticus</i> .....	1232
1748	<i>inornatus</i> .....	1341	<i>Stellifer</i> .....	1445
1829	<i>microlepidotus</i> , <i>Cestrens</i> .....	1415	<i>Tylosurus</i> .....	712
1829	<i>Cynoscion</i> .....	1415	<i>microptera</i> , <i>Rhandia</i> .....	153
2698	<i>Labrisomus</i> .....	2363	<i>Micropterinae</i> .....	986
692	<i>Labrosomus</i> .....	2364	<i>Micropterus</i> .....	1010
416	<i>Orthodon</i> .....	207	<i>dolomieu</i> .....	1011
563	<i>Otolithus</i> .....	1415	<i>salmoides</i> .....	1012
562	<i>Microlepis</i> .....	228	<i>micropterus</i> , <i>Etheostoma</i> .....	1083
2264	<i>microlepis</i> , <i>Antimora</i> .....	2545	<i>Pimeiodus</i> .....	153
2237	<i>Epinephelus</i> .....	1178	<i>Micropteryx</i> .....	901, 937
1121	<i>Lepophidium</i> .....	2486	<i>chrysurus</i> .....	938
416	<i>Macrondon</i> .....	330	<i>cosmopolita</i> .....	938
2237	<i>Mycteroperca</i> .....	1177	<i>micropteryx</i> , <i>Alburnellus</i> .....	297
1380	<i>Trisotropis</i> .....	1178	<i>Minulus</i> .....	297
2237	<i>microlophus</i> , <i>Pomotis</i> .....	1008	<i>Notropis</i> .....	206
813	<i>Micromesus</i> .....	90, 91	<i>Platysomus</i> .....	934
978	<i>Micrometrus</i> .....	1496, 1498	<i>micropus</i> , <i>Gasterosteus</i> .....	744
538	<i>aggregatus</i> .....	1499	<i>microrhynchus</i> , <i>Acipenser</i> .....	106
325	<i>frenatus</i> .....	1499	<i>microrrhinos</i> , <i>Pseudoscarus</i> .....	1655
441	<i>rosaceus</i> .....	1500	<i>Microspathodon</i> .....	1565
469	<i>miconema</i> , <i>Peristethus</i> .....	2182	<i>azurissimus</i> .....	1570
518	<i>miconemus</i> , <i>Peristedion</i> .....	2182	<i>bairdi</i> .....	1566, 1567
469	<i>Microperca</i> .....	1103	<i>chrysurus</i> .....	1567
469	<i>fonticola</i> .....	1104	<i>cinereus</i> .....	1570
633	<i>praellaris</i> .....	1103	<i>dorsalis</i> .....	1568, 1570
52	<i>punctulata</i> .....	1104	<i>azurissimus</i> .....	1570

	Page.		Page.
Microspathodon dorsalis cinereus ..	1570	Minellus .....	254
niveatus .....	1567	minima, Abeona .....	1497
Microspatbodontina .....	1544	Cichla .....	1012
microstignius, Myrophis .....	371	Clupea .....	422
microstoma, Chola .....	264	Perca .....	1057
Crenilabrus .....	1576	minimus, Cymatogaster .....	1497
Scartella .....	2384	miniofrenatus, Searus .....	1634
Tetragonopterus .....	334	Mink, Sea .....	1475
Uranileca .....	1958	Minnilus, altipinnis .....	287
Microstomido .....	527	anahalalis .....	291
Microstominae .....	527	ardens .....	301
Microstomus .....	2653	ariomunus .....	290
grœnlandicus .....	528	atripes .....	300
kitt .....	2654	bellus .....	297
latidens .....	2654	bivittatus .....	233
pacificus .....	2655	blennius .....	262
microstomus, Blennius .....	2385	cercoostigma .....	275
Citharichthys .....	2688	chalybœus .....	288
Conger .....	356	chlorocephalus .....	286
Etropus .....	2687, 2690	chrosomus .....	288
Minnilus .....	262	coccogenis .....	285
Pleuronectes .....	2654	cornutus .....	283
micurum, Syacium .....	2672	cyanoccephalus .....	300
Midshipman .....	2317, 2321	dinenus .....	293
Miklas .....	2819	dipleminis .....	300
milberti, Acpenser .....	105	jejunus .....	290
Arius .....	128	jemezanus .....	294
Carcharhinus .....	37	lacertosus .....	284
Carcharias .....	37	leucodus .....	291
Eulamia .....	37	lirus .....	298
milbertianus, Syngnathus .....	771	lutipinnis .....	287
miles, Porogadus .....	2520	matutinus .....	301
Prionotus .....	2160	megalops .....	291
milesi, Pimephales .....	217	micropteryx .....	297
miliaris, Bellator .....	2173	microstomus .....	262
Gymnothorax .....	398	nigripinnis .....	299
Lycodontis .....	397	notatus .....	218
Muraena .....	398	oligaspis .....	294
Thrysoldea .....	398	percobromus .....	295
Milk Fishes .....	414	plumbeolus .....	283
milkschitch, Salmo .....	481	punctulatus .....	302
milleri, Bathylagus .....	25	roseus .....	288
Miller's Thumb .....	1041, 1050	rubellus .....	293
mllipunctatus, Gasterosteus .....	752	rubricoccus .....	286
milneri, Nocomis .....	324	rubrifrons .....	295
Pagellus .....	1355	rubripinnis .....	298
Sparus .....	1355	scabriceps .....	268, 290
milnerianus, Leuclaus .....	242	septicus .....	296
Phoxinus .....	242	selene .....	269
Milvus cirratus .....	2183	shumardi .....	268, 269
mineopas, Bryttus .....	996	stilbus .....	293
Mingo, Jong .....	1718	telescopus .....	292
minlatu n, Peristeddon .....	2178	timpanogenesis .....	233
minlatus, Lepomis .....	1002	umbratilis .....	299
Sebastichthys .....	1795	xænocephalus .....	289
Sebastodes .....	1794	xænurus .....	280

Page.		Page.		Page.
254	Minnilus, zonistius .....	285	mitchilli, Perca .....	1133
1497	Minnow, Black-head .....	217	alternata .....	1133
1012	Blunt-nosed .....	218	interrupta .....	1133
422	Eastern Mud .....	624	Stolephorus .....	446
1057	Leather-sided .....	236	Mitchillina .....	453
1497	Mangrove .....	643	bairdii .....	454
1634	Shoalshead .....	671	mitis, Balistes .....	1705
1475	Silver-sided .....	238	mitzukurii, Sebastodes .....	1831
287	Silvery .....	213	miurus, Mystriophis .....	387
201	Spot-tailed .....	269	Noturus .....	148
301	Spotted-tail .....	275	Ophichthys .....	387
290	Star-headed .....	656	Schilbeodes .....	148
300	Straw-colored .....	261	Scytalichthys .....	387
297	Top .....	659, 680	Mixonus .....	2523
233	Minnows, Mud .....	623	laticeps .....	2523
262	Pursy .....	670, 671	mixtus, Gadus tomcodus .....	2540
275	Minomus .....	173	Mnierpes .....	2364
288	bardus .....	171	macrocephalus .....	2364
286	delphinus .....	171	Mobula .....	91
288	jarrovi .....	170	molesta, Cheonda .....	234
285	platyrhynchus .....	170	Gambusia .....	693
283	minor, Anarrhichas .....	2446	Pimelodella .....	154
300	Anarrhichas .....	2446	modestum, Hemulon .....	1340
293	Aphanopus .....	885	modestus, Julis .....	1601
300	Atinga alter orbicularis .....	1749	Oxyjulis .....	1601
290	Cyclopterus liparis .....	2121	Pimelodus .....	154
294	Stellifer .....	1442	Pomadasys .....	1321
284	Umbla marina .....	823	Pseudojulis .....	1601
291	minutillus, Chriolepis .....	2205	Squalius .....	234
298	minutus, Cottus .....	1958	Xyrichtys .....	1619
287	Cyclopterus .....	2097	Mobia .....	2510
301	Gobius .....	2097	promelas .....	2511
291	Minytrema .....	186	mcerens, Gonopterus .....	1688
297	austrinum .....	192	Moharra .....	1373, 1374
262	melanops .....	187	mohri, Clinus .....	2438
299	minytremus, Gyrinichthys .....	2137	Mojarra .....	1379
218	Mionurus .....	1106	Almejero .....	1294
294	mirabile, Exoglossum .....	303	Blanca .....	1372
295	mirabilis, Clupea .....	422	Cantilleña .....	1369
283	Crystallichthys .....	2865	Cardenal .....	850
302	Gillichthys .....	2250	China .....	1377
288	Lycodonus .....	2474	de Casta .....	1372
293	Phenacobius .....	303	de la Piedras .....	1681
286	mirum, Calloptilum .....	2527	de las Aletas Amarillas .....	1376
295	mispilliensis, Ameiurus .....	141	du Ley .....	1370
298	Mississippi Cat .....	137	Dorada .....	928
268, 290	mississippiensis, Morone .....	1134	Garabata .....	1353
296	Pristis .....	61	Prieta .....	1299
269	Missouri Suoker .....	168	Verde .....	1538
263, 269	missuriensis, Cliola .....	262	Mojarras .....	1366, 1373
293	Hybopsis .....	262	Mojarritas .....	1367
292	mitchilli, Argyrosus .....	936	Mojarron .....	1319
233	Acipenser .....	105	Mola .....	1753
290	Cottus .....	1973	aculeata .....	1754
289	Engraulis .....	446	mola .....	1753
280	Lepibema .....	1133	nasus .....	1754

	Page.		Page.
<i>Mola planci</i> .....	1756.	<i>Monacanthus setifer</i> .....	1716
<i>retzii</i> .....	1754	<i>signifer</i> .....	1716
<i>rotunda</i> .....	1754	<i>spilonotus</i> .....	1716
<i>mola, Mola</i> .....	1753	<i>stratus</i> .....	1713
<i>Orthogoriscus</i> .....	1754	<i>ruppellii</i> .....	1713
<i>Tetrodon</i> .....	1754	<i>varius</i> .....	1716
<i>Molacanthus</i> .....	1753	<i>monacanthus, Plectropoma</i> .....	1165
<i>Molarii</i> .....	402	<i>monacans, Ceratichthys</i> .....	318
<i>molestum, Gobiosoma</i> .....	2258	<i>Hybopsis</i> .....	318
<i>Mollidae</i> .....	1752	<i>mona, Stephanoberyx</i> .....	836
<i>Molinae</i> .....	1752	<i>monapterygia, Pesca dorso</i> .....	1833
<i>molle, Maria</i> .....	1552	<i>Monda</i> .....	899
<i>Mollenisia</i> .....	698	<i>monensis, Squalus</i> .....	49
<i>fasciata</i> .....	695	<i>monasticus, Salmo</i> .....	509
<i>formosa</i> .....	699	<i>Mongolar Drummer</i> .....	1406
<i>jonesi</i> .....	699	<i>Mongrel Buffalo</i> .....	164
<i>latipinna</i> .....	700	<i>Whitefish</i> .....	473
<i>lineolata</i> .....	700	<i>Moniana</i> .....	254, 256, 271
<i>petenensis</i> .....	700	<i>aurata</i> .....	272
<i>mollis, Achirus mollis</i> .....	2702	<i>complanata</i> .....	272
<i>Aphyonus</i> .....	2525	<i>couchi</i> .....	272
<i>Bothrocara</i> .....	2476	<i>deliciosa</i> .....	262
<i>Pleuronectes</i> .....	2701	<i>formosa</i> .....	271
<i>Molly, Hog</i> .....	181, 1026	<i>frigida</i> .....	271
<i>Molva</i> .....	2551	<i>gibbosa</i> .....	272
<i>huntingia</i> .....	2551	<i>gracilis</i> .....	272
<i>Innrei</i> .....	2552	<i>jugalis</i> .....	272
<i>maculosa</i> .....	2551	<i>laetabilis</i> .....	272
<i>molva</i> .....	2551	<i>leonina</i> .....	272
<i>vulgaris</i> .....	2552	<i>nitida</i> .....	265
<i>molva, Gadus</i> .....	2552	<i>proserpina</i> .....	272
<i>Molva</i> .....	2551	<i>pulchella</i> .....	272
<i>monacantha, Corrius</i> .....	1419	<i>rutila</i> .....	272
<i>Monacanthidae</i> .....	1712	<i>tristis</i> .....	272
<i>Monacanthus</i> .....	1714	<i>Monkfish</i> .....	58, 2713
<i>amphioxys</i> .....	1717	<i>monoceros, Alutera</i> .....	1720, 2860
<i>auriga</i> .....	1716	<i>Balistes</i> .....	1719, 1720
<i>broccus</i> .....	1716	<i>Monacanthus</i> .....	1720
<i>ciliatus</i> .....	1714, 1715	<i>Monochir lineatus</i> .....	2698
<i>davidsoni</i> .....	1715	<i>maculipinnis</i> .....	2698
<i>filamentosus</i> .....	1716	<i>reticulatus</i> .....	2696
<i>gallinula</i> .....	1716	<i>Monochirus</i> .....	2694
<i>hispidus</i> .....	1715	<i>monocirrus, Exocetus</i> .....	730
<i>irroratus</i> .....	1713	<i>Monodactylinae</i> .....	1667
<i>macrocerus</i> .....	1713	<i>Monolene</i> .....	2690
<i>massachusettsensis</i> .....	1716	<i>atrimana</i> .....	2692
<i>monoceros</i> .....	1720	<i>sessilicauda</i> .....	2691
<i>occidentalis</i> .....	1715	<i>Monomitra</i> .....	2138
<i>oppositus</i> .....	1716	<i>lparina</i> .....	2139
<i>pardalis</i> .....	1713	<i>monophthalmus, Auchenopterus</i> .....	2372
<i>parraianus</i> .....	1713	<i>Crennobates</i> .....	2372
<i>piraca</i> .....	1715	<i>Monoprion</i> .....	1106
<i>proboscideus</i> .....	1719	<i>maculatus</i> .....	1109
<i>pullus</i> .....	1713	<i>pigmentarius</i> .....	1109
<i>punctatus</i> .....	1713, 1719	<i>Monopterhinus</i> .....	18
<i>scriptus</i> .....	1719	<i>monopterygius, Aspidophoroidea</i> .....	2091, 2092

Page.		Page.		Page.
1716	monopterygius, <i>Canthirhynchus</i> . . .	2092	<i>moringa</i> , <i>Lycodontis</i> . . . . .	395
1716	<i>Chirus</i> . . . . .	1869	<i>Muraena</i> . . . . .	395
1716	<i>Cottus</i> . . . . .	2092	<i>Sidera</i> . . . . .	395
1713	<i>Hexagrammus</i> . . . . .	1866	<i>moringua</i> , <i>Gymnothorax</i> . . . . .	395
1713	<i>Labrax</i> . . . . .	1866	<i>morio</i> , <i>Centrolophus</i> . . . . .	963
1716	<i>Pneurogrammus</i> . . . . .	1864	<i>Epinephelus</i> . . . . .	1160
1165	<i>Monosira stahli</i> . . . . .	1423	<i>Serranus</i> . . . . .	1160
318	<i>morstrosa</i> , <i>Chimera</i> . . . . .	91	<i>mormyrus</i> , <i>Campostoma</i> . . . . .	206
318	<i>montacuti</i> , <i>Cylopterus</i> . . . . .	2108	<i>Morone</i> . . . . .	1133
836	<i>montagui</i> , <i>Liparis</i> . . . . .	2107, 2108	<i>americana</i> . . . . .	1134, 1135
1833	<i>Montana Grayling</i> . . . . .	519, 2871	<i>flavescens</i> . . . . .	1024
899	<i>montana</i> , <i>Gila</i> . . . . .	238	<i>interrupta</i> . . . . .	1134
49	<i>montanus</i> , <i>Clinostomus</i> . . . . .	238	<i>lineata</i> . . . . .	1133
509	<i>Hybopsis</i> . . . . .	317	<i>maculata</i> . . . . .	1010
1406	<i>Leuciscus</i> . . . . .	238	<i>mississippiensis</i> . . . . .	1134
164	<i>Squalius</i> . . . . .	238	<i>multilineata</i> . . . . .	1132
473	<i>Thymallus ontariensis</i> . . . . .	519	<i>pallida</i> . . . . .	1135
254, 256, 271	<i>signifer</i> . . . . .	519	<i>rufa</i> . . . . .	1135
272	<i>tricolor</i> . . . . .	2871	<i>Moronina</i> . . . . .	1127
272	<i>Monterey Halibut</i> . . . . .	2025	<i>Moronopsis</i> . . . . .	1013
272	Spanish Mackerel . . . . .	837	<i>Morrhua</i> . . . . .	2540
262	<i>montezuma</i> , <i>Cichlasoma</i> . . . . .	1518	<i>seglefinus</i> . . . . .	2543
271	<i>Heros</i> . . . . .	1518	<i>americana</i> . . . . .	2540, 2541
272	<i>monticola</i> , <i>Agonostomus</i> . . . . .	819	<i>californica</i> . . . . .	2539
272	<i>Dajaus</i> . . . . .	2841	<i>punctatus</i> . . . . .	2543
272	<i>Mugil</i> . . . . .	819	<i>morrhua</i> , <i>Gadus</i> . . . . .	2541
272	<i>montiregia</i> , <i>Chiola</i> . . . . .	272	<i>morrissi</i> , <i>Leptocephalus</i> . . . . .	354
272	<i>Moon-eye</i> . . . . .	413	<i>Moseleya</i> . . . . .	2570
272	<i>Cisco</i> . . . . .	469	<i>cyclolepis</i> . . . . .	2570
265	<i>Moon-eyes</i> . . . . .	412	<i>moseri</i> , <i>Verasper</i> . . . . .	2619
272	<i>Moonfish</i> . . . . .	934, 954	<i>Mosabunker</i> . . . . .	433
272	<i>Moonfishes</i> . . . . .	935	<i>motaguensis</i> , <i>Heros</i> . . . . .	1534
272	<i>Moorish Idol</i> . . . . .	1687	<i>Pinelodus</i> . . . . .	151
272	<i>Mora</i> , <i>Cabr.</i> . . . .	1152	<i>Rhambia</i> . . . . .	151
58, 2713	<i>Moray, Black</i> . . . . .	396	<i>Motella</i> . . . . .	2558
1720, 2860	Common Spotted . . . . .	395	<i>argentata</i> . . . . .	2559
1719, 1720	Spotted . . . . .	399	<i>caudacuta</i> . . . . .	2560
1720	<i>Morays</i> . . . . .	388, 400	<i>cimbria</i> . . . . .	2560
2698	<i>mordax</i> , <i>Atherina</i> . . . . .	523	<i>ensis</i> . . . . .	2559
2698	Conger . . . . .	387	<i>reinhardtii</i> . . . . .	2559
2696	<i>Crotalopsis</i> . . . . .	387	<i>septentrionalis</i> . . . . .	2560
2694	<i>Engraulis</i> . . . . .	448	<i>Mother of Eels</i> . . . . .	2457
730	<i>Gymnothorax</i> . . . . .	396	<i>motta</i> , <i>Elacate</i> . . . . .	948
1667	<i>Lycodontis</i> . . . . .	395	<i>moucharra</i> , <i>Glyphisodon</i> . . . . .	1562
2600	<i>Macrodonophis</i> . . . . .	387	<i>Mountain Herring</i> . . . . .	463
2692	<i>Muraena</i> . . . . .	396	<i>Suckers</i> . . . . .	169, 170
2691	<i>Osmerus</i> . . . . .	523	<i>Mouse-Fish</i> . . . . .	2715, 2716
2138	<i>abbotti</i> . . . . .	524	<i>Moxostoma</i> . . . . .	185, 187
2139	<i>spectrum</i> . . . . .	523	<i>albidum</i> . . . . .	192
2372	<i>Sidera</i> . . . . .	396	<i>album</i> . . . . .	191
2372	<i>Morena Pinta</i> . . . . .	402	<i>anisurum</i> . . . . .	190, 196
1106	<i>Pintita</i> . . . . .	397	<i>aureolum</i> . . . . .	192
1109	<i>Prieta</i> . . . . .	2804	<i>austrinum</i> . . . . .	192
1109	<i>Verde</i> . . . . .	306	<i>breviceps</i> . . . . .	196
18	<i>moribundus</i> , <i>Balistes</i> . . . . .	1702	<i>bucco</i> . . . . .	190
2091, 2092	<i>Morinae</i> . . . . .	2532	<i>campbelli</i> . . . . .	186

	Page.		Page.
<i>Moxostoma carpio</i> .....	190	<i>Mugilcephalus</i> .....	811
<i>cervinum</i> .....	197	<i>chanos</i> .....	415
<i>claviformis</i> .....	186	<i>cincrous</i> .....	1373
<i>collapsum</i> .....	190	<i>ourema</i> .....	813, 2841
<i>congestum</i> .....	192	<i>equinoculus</i> .....	2841
<i>conus</i> .....	196	<i>galmardianus</i> .....	814
<i>coregonus</i> .....	191	<i>gigas</i> .....	1483
<i>crassilabro</i> .....	194, 196	<i>grunulens</i> .....	1483
<i>kuinnerlyi</i> .....	186	<i>guntheri</i> .....	812
<i>lesneuri</i> .....	194	<i>hospes</i> .....	814
<i>macrolepidotum</i> .....	193	<i>inellii</i> .....	813
<i>oblongum</i> .....	186	<i>irretitus</i> .....	819
<i>oneida</i> .....	193	<i>lebranchus</i> .....	811
<i>papillosum</i> .....	189	<i>lineatus</i> .....	812, 2841
<i>philense</i> .....	191	<i>liza</i> .....	811, 2841
<i>poecilurum</i> .....	196	<i>mexicanus</i> .....	813
<i>robustum</i> .....	193	<i>monticola</i> .....	819
<i>rupiscartes</i> .....	196	<i>nigro-strigatus</i> .....	817
<i>tenne</i> .....	186	<i>obliquus</i> .....	1459
<i>thalassinum</i> .....	191	<i>petrosus</i> .....	814, 2841
<i>tristignatum</i> .....	179	<i>plumieri</i> .....	812, 2841
<i>valenciennesi</i> .....	190	<i>proboscideus</i> .....	816
<i>velatum</i> .....	190	<i>rammelsbergii</i> .....	812
<i>victoriae</i> .....	187	<i>salmoneus</i> .....	415
<i>mucosum</i> , <i>Xiphidion</i> .....	2425	<i>setosus</i> .....	815
<i>Xiphister</i> .....	2425	<i>tang</i> .....	812
<i>mucosus</i> , <i>Acanthocottus</i> .....	1975	<i>thoburni</i> .....	813
<i>Liparis</i> .....	2111	<i>trichodon</i> .....	816
<i>Lycodalepis</i> .....	2470	<i>Mugilidae</i> .....	808
<i>Lycodes</i> .....	2401	<i>Mugilinae</i> .....	809
<i>Neoliparis</i> .....	2111	<i>mugiloides</i> , <i>Eleotris</i> .....	2198
<i>mucronata</i> , <i>Odontognathus</i> .....	438	<i>Mugilomorus</i> .....	409
<i>Perca</i> .....	1135	<i>anna-carolina</i> .....	410
<i>mucronatum</i> , <i>Ophidium</i> .....	2419	<i>mulkias</i> , <i>Salmo</i> .....	492
<i>mucronatus</i> , <i>Gnathobolus</i> .....	438	<i>Muksun of the Russians</i> .....	464
<i>Labrax</i> .....	1135	<i>muksun</i> , <i>Salmo</i> .....	464
<i>Neoconger</i> .....	362	<i>mulleri</i> , <i>Antigonia</i> .....	1665
<i>Odontognathus</i> .....	438	<i>Benthoema</i> .....	574
<i>Pristigaster</i> .....	438	<i>Carcharias</i> .....	40
<i>Mud Cat</i> .....	142, 143	<i>Maurolleus</i> .....	577
<i>Dabblers</i> .....	640	<i>Pempheris</i> .....	978
<i>Dabs</i> .....	2644	<i>Salmo</i> .....	574
<i>Minnows</i> .....	622	<i>Scopelus</i> .....	579, 574
<i>Parrot</i> .....	1639	<i>Mullet</i> .....	192
<i>Sunfish</i> .....	989	<i>Blue</i> .....	191
<i>Mudfish</i> .....	113, 640, 1649	<i>Blue-back</i> .....	813
<i>mudfish</i> , <i>Fundulus</i> .....	641	<i>Common</i> .....	811
<i>Mudfishes</i> .....	623	<i>Fan-tail</i> .....	816
<i>Muffle-jaw</i> .....	1950	<i>French</i> .....	813
<i>Muger</i> , <i>Vieja</i> .....	1639	<i>Jumping</i> .....	197
<i>Mugil</i> .....	809	<i>Red-eye</i> .....	814
<i>alatus</i> .....	733	<i>Snip-nose</i> .....	964
<i>albula</i> .....	812, 2841	<i>Striped</i> .....	811
<i>berlandieri</i> .....	812	<i>Whirligig</i> .....	818
<i>brasiliensis</i> .....	810, 814, 816	<i>White</i> .....	189, 813
<i>capitulinus</i> .....	2841	<i>Mullet of Utah Lake</i> .....	179



Page.		Page.		Page.
811	Mullets .....	808	<i>Murena aquae-dulcis</i> .....	391
415	<i>Mullhypenus</i> .....	858	<i>urgentes</i> .....	348
1373	<i>maculatus</i> .....	850	<i>argus</i> .....	401, 2805
813, 2841	Mullidae .....	855	<i>aterrima</i> .....	396
2841	<i>Mulloides</i> .....	857, 2843	<i>balearica</i> .....	350
814	<i>flavovittatus</i> .....	800	<i>bostoniensis</i> .....	348
1483	<i>rathbuni</i> .....	857	<i>cassini</i> .....	356
1483	Mullus .....	856	<i>cutenata</i> .....	403
812	<i>auratus</i> .....	856, 2843	<i>clepsydra</i> .....	2805
814	<i>barbatus auratus</i> .....	856	<i>conger</i> .....	354
813	<i>imberbis</i> .....	1107	<i>coniceps</i> .....	359
819	<i>maculatus</i> .....	859	<i>conspersa</i> .....	397
811	<i>multifasciata</i> , <i>Adinia</i> .....	661	<i>cubana</i> .....	348
812, 2841	<i>Hydrargyra</i> .....	645	<i>dovii</i> .....	397
811, 2841	<i>Scelena</i> .....	1459	<i>elaborata</i> .....	380
813	<i>multifasciatus</i> , <i>Anthias</i> .....	1226	<i>erebus</i> .....	396
810	<i>Fundulus</i> .....	645	<i>havennensis</i> .....	382
817	<i>Pronotogrammus</i> .....	1226	<i>infernalis</i> .....	396
1459	<i>multifilis</i> , <i>Blennius</i> .....	2385	<i>insularum</i> .....	400
814, 2841	<i>Hypleurochilus</i> .....	2385	<i>lenticinosa</i> .....	402
812, 2841	<i>multiguttatus</i> , <i>Alphestes</i> .....	1165	<i>lineopinnis</i> .....	396
816	<i>Epinephelus</i> .....	1166	<i>lita</i> .....	2805
812	<i>Plectropoma</i> .....	1166	<i>longicauda</i> .....	392
415	<i>multiineata</i> , <i>Morone</i> .....	1132	<i>lumbicus</i> .....	342
815	<i>Pecilia</i> .....	700	<i>macrocephala</i> .....	348
812	<i>multilineatum</i> , <i>Hemulon</i> .....	1304	<i>maculata nigra</i> .....	395
813	<i>multimaculatus</i> , <i>Chromis</i> .....	1547	<i>maculipinnis</i> .....	394
816	<i>Diodon</i> .....	1746	<i>maculosa</i> .....	382
808	<i>Heliases</i> .....	1547	<i>melanatis</i> .....	402
809	<i>Labrax</i> .....	1132	<i>melanotis</i> .....	401, 402
2198	<i>multicellata</i> , <i>Murena</i> .....	398	<i>meleagria</i> .....	399
409	<i>multicellatus</i> , <i>Antennarius</i> .....	2724	<i>miliaris</i> .....	398
410	<i>Chironectes</i> .....	2725	<i>mordax</i> .....	396
492	<i>multiradiatus</i> , <i>Arius</i> .....	133	<i>moringa</i> .....	395
464	<i>Cubiceps</i> .....	951	<i>multicellata</i> .....	398
464	<i>Tachysurus</i> .....	132, 2788	<i>nigra</i> .....	355
1065	<i>multispinosum</i> , <i>Cichlasoma</i> .....	1525	<i>nigricans</i> .....	390
574	<i>multispinosus</i> , <i>Heros</i> .....	1526	<i>ocellata</i> .....	399
40	<i>multistriatus</i> , <i>Liparis lineatus</i> .....	2118	<i>ophis</i> .....	382, 2804
577	<i>Mummichog</i> .....	640	<i>panamensis</i> .....	391
978	<i>mundecolus</i> , <i>Stolephorus</i> .....	2812	<i>pfeifferi</i> .....	2805
574	<i>mundiceps</i> , <i>Novacula</i> .....	1618	<i>picta</i> .....	2805
570, 574	<i>Xyrichtys</i> .....	1618	<i>pinnata</i> .....	351
192	<i>mundicorpus</i> , <i>Iniiatius</i> .....	1620	<i>pinta</i> .....	402
191	<i>Novacula</i> .....	1620	<i>phititi</i> .....	397
813	<i>mundus</i> , <i>Dactyagnus</i> .....	2304	<i>punctata</i> .....	395
811	<i>Oligoplites</i> .....	2844	<i>retifera</i> .....	401
816	<i>Urolophus</i> .....	81	<i>rostrata</i> .....	348
813	<i>Urotrygon</i> .....	81	<i>saetae-helenae</i> .....	397
197	<i>Muñeca</i> .....	1674	<i>sanguinea</i> .....	390
814	<i>Mupine</i> .....	962	<i>savanna</i> .....	360
964	<i>Murena</i> .....	347, 400	<i>serpentina</i> .....	348
811	<i>acuminata</i> .....	377	<i>sen conger brasiliensis</i> .....	403
818	<i>afra</i> .....	306	<i>sidera</i> .....	2805
189, 813	<i>aluisi</i> .....	403	<i>sordida</i> .....	403
170	<i>anguina</i> .....	390	<i>variegata</i> .....	2805

	Page.		Page.
Muraena vicina.....	394	mutatum, Boleosoma.....	1057
muraena, Callochelys.....	378	muticus, Lethotremus.....	2101
Muraenesocidae.....	358	Mutton-fish.....	1376, 1265, 2457
Muraenesocinae.....	358	Mycteroperca.....	1169, 1171, 1183
Muraenox.....	359	bonaci.....	1174
coniceps.....	2801	xanthostieta.....	1170
savanna.....	360	boulengeri.....	1171, 2856
Muraenidae.....	345, 388	calliura.....	1786, 2856
Muraenoblenna.....	7, 403	dimidiata.....	1179
nectura.....	404	falcata.....	1184
Muraenoides.....	2414	phenax.....	1185
dolichogaster.....	2417	hopkinsi.....	2855
fasciatus.....	2418	interstitialis.....	1178
gunelliformis.....	2421	jordani.....	1176
gunnellus.....	2419	microlepis.....	1177
guttatus.....	2419	olfax.....	1183
maxillaris.....	2418	ruberrima.....	1183
ornatus.....	2420	pardalis.....	1181
stujef.....	2419	reticulata.....	1187
tonia.....	2418	rosacea.....	1184
muraenoides, Blennius.....	2419	rubor.....	1180
Gunnellus.....	2418	selenga.....	1181
Muraenophis.....	381, 383, 400	tigris.....	1187
caramura.....	395	camelopardalis.....	1187
catenula.....	403	venadorum.....	1180
curvilineata.....	395	venosa.....	1172
marmoreus.....	391	apua.....	1173, 1174
ocelletus.....	384	guttata.....	1174
pantherina.....	2805	xenarcha.....	1180
punctata.....	397	Myctophidae.....	550
triserialis.....	384	Myctophum.....	560
undulata.....	403	attine.....	570
vicina.....	394	benoiti.....	573
Murotelago.....	2183	boops.....	572
muricatus, Orbis.....	1749, 1750	brachychr.....	556
murinus, Apomotis.....	996	californiense.....	572
Calliurus.....	996	crenulare.....	575
muroadsi, Caranx.....	908	glaciale.....	574
murrayi, Ipnops.....	547	gracile.....	572
Liocetus.....	2733	hians.....	572
Melanocetus.....	2734	humboldti.....	571
Muscalonge.....	629	hygomi.....	573
muscarum, Rimicola.....	2338	laeerta.....	560
musculus, Cyclopterus.....	2118	(Stenobrachius) leu-	
Muskallunge.....	629	copsarum.....	562
Musquaw River Whitefish.....	466	mexicanum.....	563
Mustela.....	2558	nannochir.....	562
Mustellus stellatus.....	29	nocturnum.....	568
Mustelus.....	28, 2745	opalinum.....	571
asterias.....	29	procellarum.....	575
californicus.....	30	protoculus.....	565
canis.....	29	punctatum.....	570
dorsalis.....	30	regale.....	563
felis.....	31	remiger.....	573
lunulatus.....	28	townsendi.....	558
plebejus.....	29	Myctophus rafinesquei.....	567
vulgaris.....	29		

	Page.		Page.
mydrus, Cyprinodon .....	676	Myoxocephalus stelleri .....	1981
Mykiss .....	492, 2818	verrucosus .....	1079
mykiss, Salmo .....	487, 492, 2818	Myrichthys .....	375
agua-bonita .....	504	acuminatus .....	376
bouvieri .....	496	oculatus .....	376
gibbsii .....	493	pantostigmus .....	2802
henshawi .....	493	tigrinus .....	376, 2802
lewisii .....	493	xysturus .....	2802
macdonaldi .....	407	Myridae .....	370
pleuritiens .....	490	Myriolepis .....	1862
spilurus .....	495	zoulier .....	1863
stomias .....	497	Myriosteon .....	60
virginalis .....	493	Myripristis .....	846
Mylobatidae .....	87	clarionensis .....	2842
Mylobatinae .....	2753	fulgens .....	846
Mylobatis .....	89, 2750	jacobus .....	846
acuta .....	89	lychnus .....	847
asperrimus .....	2754	occidentalis .....	847
bispinosus .....	80	paecilopus .....	847
californicus .....	80, 2754	trachypoma .....	846
eastenkee .....	88	Myrophis .....	371
fremenvillei .....	89	egmontis .....	371
goodii .....	2755	longicollis .....	371
sayi .....	86	lumbricus .....	371
Mylocheilus .....	219	microstigmus .....	371
canrinus .....	219, 220	punctatus .....	371
fraterculus .....	220	vafer .....	372
lateralis .....	220	mystacalis, Lepomis .....	1001
Myloleucus .....	243, 244, 2798	mystacina, Gnathypops .....	2286
parovanus .....	246	mystacinus, Epinephelus .....	1151
pulverulentus .....	246	Gnathypops .....	2286
thalassinus .....	245	Schistorus .....	1151
Mylopharodon .....	218	Serranus .....	1151
conocephalus .....	219	Mystaconurus .....	2580
robustus .....	219	mystes, Scorpaena .....	1849
Mylopharodontinae .....	202	mysticetus, Cetengraulis .....	450
Mylorhina .....	90	Engraulis .....	450
myops, Salmo .....	533	mystinus, Sebastichthys .....	1785
Saurus .....	533	Sebastodes .....	1784, 1785
Synodus .....	533	Mystriophis .....	386
Trachinocephalus .....	533	intertinctus .....	386
Myoxocephalus .....	1970, 1971, 1976	minrus .....	387
aeneus .....	1972	Mystus .....	116
axillaris .....	1980, 1981	ascita .....	155
brandti .....	1984	carolinensis .....	117
bubalis .....	1971	Mytilophagus .....	1503
grœnlandicus .....	1974	fasciatus .....	1504
jaok .....	1977	Myxine .....	7
medinus .....	1983	glutinosa .....	7
niger .....	1985	lmosa .....	8
nivosus .....	1984	Myxinidae .....	7
octodecimspinosus .....	1976	Myxodagnus .....	2305
polyacanthocephalus .....	1976	opercularis .....	2305
scorpioides .....	1973	Myxodes elegans .....	2353
scorpius .....	1974	lugubris .....	2357
		macropus .....	2357

	Page.		Page.
<i>Myxodes varius</i> .....	2357	<i>nasus</i> , <i>Acanthonotus</i> .....	615
<i>versicolor</i> .....	2359	<i>Diplanchias</i> .....	1754
<i>Myxostoma austrinum</i> .....	192	<i>Menticirrhus</i> .....	1472, 1473
<i>eurypus</i> .....	193	<i>Mola</i> .....	1754
<i>pæcilura</i> .....	196	<i>Notacanthus</i> .....	615
<i>Myxus harengus</i> .....	818	<i>Squalus</i> .....	49
<i>Myzopsetta</i> .....	2644	<i>Umbrina</i> .....	1473
<i>ferruginea</i> .....	2645	<i>nasuta</i> , <i>Malthæa</i> .....	2737
<i>Naccaysh</i> .....	413	<i>nasutum</i> , <i>Agonostoma</i> .....	820
<i>nachtriebi</i> , <i>Leuciscus</i> .....	2798	<i>Campostoma</i> .....	206
<i>nacrea</i> , <i>Gila</i> .....	228	<i>nasutus</i> , <i>Agonostomus</i> .....	819, 2841
<i>Naked Gobies</i> .....	2257	<i>Argyreus</i> .....	306
<i>Goby</i> .....	2259	<i>Leuciscus</i> .....	306
<i>nalnal</i> , <i>Sparactodon</i> .....	947	<i>Ogcocephalus</i> .....	2737
<i>Namayush</i> .....	504	<i>Rhluichthys</i> .....	306
<i>Salmon</i> .....	505	<i>Trachynotus</i> .....	941
<i>nanayush</i> , <i>Cristivomer</i> .....	504	<i>natalis</i> , <i>Amciurus</i> .....	139
<i>siscowet</i> ..	505	<i>analis</i> .....	140
<i>Salvelinus</i> .....	505	<i>Pimelodus</i> .....	140
<i>siscowet</i> ....	505	<i>Naucleus abbreviatus</i> .....	900
<i>Naunobranchium</i> .....	561	<i>annularis</i> .....	900
<i>leucopsarum</i> .....	562	<i>brachycentrus</i> .....	900
<i>macdonaldi</i> .....	563	<i>compressus</i> .....	900
<i>mexicanum</i> .....	563	<i>leucurus</i> .....	900
<i>nannochoir</i> .....	562	<i>triacanthus</i> .....	900
<i>regale</i> .....	563	<i>naucruteoides</i> , <i>Echeneis</i> .....	2270
<i>nannochoir</i> , <i>Myctophum</i> .....	562	<i>Leptecheneis</i> .....	2270
<i>Nannobranchium</i> .....	562	<i>Naucrates</i> .....	900
<i>nanomyzon</i> , <i>Catostomus</i> .....	177	<i>cyanophrys</i> .....	900
<i>Nanostoma</i> .....	1066, 1067, 1070	<i>ductor</i> .....	900
<i>camurus</i> .....	1076	<i>fanfarus</i> .....	900
<i>elegans</i> .....	1075	<i>indicus</i> .....	900
<i>inscriptum</i> .....	1072	<i>novboracensis</i> .....	900
<i>vinctipes</i> .....	1075	<i>serratus</i> .....	900
<i>zonale</i> .....	1075	<i>naucrates</i> , <i>Echeneis</i> .....	2269
<i>Nansenia</i> .....	528	<i>Leptecheneis</i> .....	2269
<i>grœnlandica</i> .....	528	<i>naufragium</i> , <i>Balistes</i> .....	1700, 1701
<i>nanus</i> , <i>Engraulis</i> .....	449	<i>Nautichthys</i> .....	2020
<i>Naracacion</i> .....	77	<i>oculofasciatus</i> .....	2020, 2021
<i>Narcine</i> .....	78	<i>pribilovius</i> .....	2020
<i>brasiliensis</i> .....	78, 2752	<i>Nautiscus</i> .....	2019
<i>corallina</i> .....	78	<i>pribilovius</i> .....	2019
<i>entemedor</i> .....	2752	<i>nautopædium</i> , <i>Porichthys</i> .....	2323
<i>nigra</i> .....	78	<i>navagu</i> , <i>Eloginus</i> .....	2537
<i>umbrosa</i> .....	78	<i>Navarchus</i> .....	956
<i>Narcobatidæ</i> .....	76	<i>Neolotus</i> .....	881
<i>Narcobatus</i> .....	77	<i>tripes</i> .....	881
<i>naresi</i> , <i>Salmo</i> .....	515	<i>neapolitanus</i> , <i>Lampugus</i> .....	953
<i>Salvelinus</i> .....	515	<i>nebrascensis</i> , <i>Nocomis</i> .....	323
<i>oquassa</i> .....	515	<i>Nebris</i> .....	1416
<i>narinari</i> , <i>Aetobatus</i> .....	88	<i>microps</i> .....	1417
<i>Raja</i> .....	88	<i>zestus</i> .....	1417
<i>Etosadon</i> .....	88	<i>nebularis</i> , <i>Platophrys</i> .....	2664
<i>naso</i> , <i>Stolephorus</i> .....	2813	<i>nebulifer</i> , <i>Catostomus</i> .....	171
<i>Nassau Grouper</i> .....	1157	<i>Chromis</i> .....	1524
		<i>Labrax</i> .....	1195

Page.	Page.	Page.
615	nebulifer, <i>Peralabrax</i> .....	1195, 1196
1754	<i>Serranus</i> .....	1196
1473	nebuliferum, <i>Cichlasoma</i> .....	1524
1754	nebuliferus, <i>Catostomus</i> .....	171
615	nebulosa, <i>Aphoristia</i> .....	2712
49	<i>Cerna</i> .....	1181
1473	<i>Percina</i> .....	1027
2737	<i>Pileoma</i> .....	1027
820	<i>Umbrina</i> .....	1475
206	nebulosum, <i>Gunnellus</i> .....	2414
9, 2841	nebulosus, <i>Acantholebius</i> .....	1872
306	<i>Ameiurus</i> .....	140
306	<i>catulus</i> .....	141
2737	<i>marmoratus</i> .....	141
306	<i>Apionichthys</i> .....	2703
941	<i>Centronotus</i> .....	2414
139	<i>Cestrens</i> .....	1409
140	<i>Chiropsis</i> .....	1872
140	<i>Chirus</i> .....	1872
900	<i>Clinus</i> .....	2438
900	<i>Cynoscion</i> .....	1409
900	<i>Enedrias</i> .....	2414
900	<i>Gasterosteus</i> .....	746
900	<i>Lyeodes</i> .....	2468
900	<i>Menticirrhus</i> .....	1475
2270	<i>Otolithus</i> .....	1409
2270	<i>Pimelodus</i> .....	140
900	<i>Sebastes</i> .....	1826
900	<i>Silurus</i> .....	143
900	<i>Symphurus</i> .....	2712
900	<i>Urolophus</i> .....	80, 2752
900	<i>Nector</i> .....	1436
900	<i>nectura</i> , <i>Gymnomuraena</i> .....	404
900	<i>Murenoblema</i> .....	404
2269	<i>necturus</i> , <i>Uropterygius</i> .....	404
2269	<i>Needlefishes</i> .....	714
700, 1701	<i>Needlefishes</i> .....	708
2020	<i>Neetroplus</i> .....	1541
020, 2021	<i>nematopus</i> .....	1541
2020	<i>nicaraguensis</i> .....	1542
2019	<i>nefastus</i> , <i>Pomotis</i> .....	1003
2019	<i>neglecta</i> , <i>Clupea</i> .....	434
2323	<i>Corvina</i> ( <i>Amblyodon</i> ) .....	1484
2537	<i>neglectus</i> , <i>Amblyodon</i> .....	1484
956	<i>Negra</i> , <i>Boca</i> .....	1837
881	<i>Esmeralda</i> .....	2204
881	<i>Negro</i> .....	1100
953	<i>Petite</i> .....	1142
323	<i>Negro</i> , <i>Barbero</i> .....	1692
1416	<i>Pargo</i> .....	1252
1417	<i>Negro-fish</i> .....	1146
1417	<i>negromaculatus</i> , <i>Rhypticus</i> .....	1233
2664	<i>nelsoni</i> , <i>Awaous</i> .....	2235
171	<i>Gobius</i> .....	2235
1524	<i>nelsonii</i> , <i>Coregonus</i> .....	466
1195	<i>Nematistidae</i> .....	894
	<i>Nematistius</i> .....	895
	<i>pectoralis</i> .....	895
	<i>Nematognathi</i> .....	114
	<i>Nematonurus</i> .....	2571
	<i>cyclolepis</i> .....	2571
	<i>goodei</i> .....	2571
	<i>Macrurus magnus</i> .....	2574
	<i>suborbit-</i>	
	<i>alis</i> .....	2572, 2573
	<i>Nematonus</i> .....	2518
	<i>pectoralis</i> .....	2518
	<i>nematophthalmus</i> , <i>Scorpena</i> .....	2861
	<i>Sebastes</i> .....	2861
	<i>nematopus</i> , <i>Neetroplus</i> .....	1541
	<i>Physiculus</i> .....	2548
	<i>Nematostoma</i> .....	774
	<i>Nemianthyidae</i> .....	366
	<i>Nemichthys</i> .....	369
	<i>avocetta</i> .....	369
	<i>infans</i> .....	368
	<i>scelopacea</i> .....	369
	<i>scelopaceus</i> .....	369
	<i>Nemipterus</i> .....	1288
	<i>macroneurus</i> .....	1289
	<i>Nemobrama</i> .....	854
	<i>Nemocampsis</i> .....	1010
	<i>neoboracensis</i> , <i>Pistularia</i> .....	757
	<i>Neobythites</i> .....	2512
	<i>gillii</i> .....	2513
	<i>maginatus</i> .....	2513
	<i>ocellatus</i> .....	2513
	<i>robustus</i> .....	2515
	<i>stelliferoides</i> .....	2516
	<i>Neoclinus</i> .....	2354
	<i>blanchardi</i> .....	2354
	<i>satiricus</i> .....	2355
	<i>Neoconger</i> .....	362
	<i>mucronatus</i> .....	362
	<i>perlongus</i> .....	363
	<i>verruiformis</i> .....	362
	<i>Ncoditrema</i> .....	1511
	<i>ransonnetii</i> .....	1511
	<i>neogaeus</i> , <i>Leuclenus</i> .....	240
	<i>Phoxinus</i> .....	241
	<i>neoguinaea</i> , <i>Albula</i> .....	412
	<i>neohantoniensis</i> , <i>Coregonus</i> .....	466
	<i>Neoliparis</i> .....	2106
	<i>atlanticus</i> .....	2107
	<i>callyodon</i> .....	2110
	<i>fissuratus</i> .....	2113
	<i>flora</i> .....	2111
	<i>greeni</i> .....	2112
	<i>mucosus</i> .....	2111
	<i>rutteri</i> .....	2108
	<i>Neomenis</i> .....	1247, 1248, 1251
	<i>ambiguus</i> .....	1271

	Page.		Page.
Neomænis analls	1265	nevadensis, Agonia	310
apodus	1258	Cyprinodon	674
aratus	1273	Rhinichthys (Apocope)	311
argentiventris	1260	nevisense, Etheostoma	1034
ayu	1264	nevisensis, Alvordius	1034
brachypterus	1268	newberryi, Encyclogobius	2248
buccanella	1261	Fario	499
colorado	1267	Gobius	2248
cyanopterus	1254	Leplogobius	2248
griseus	1255	New England Hake	2530
guttatus	1269	New Light	987
hastingsi	2858	newmani, Hyostoma	1053
jocu	1257	New York Smelt	468
jordani	1251	Nexilarius	1559
lutjanoides	1261	concolor	1559
mahogoni	1272	nianguæ, Etheostoma (Hadrop- terns)	1043
novemfasciatus	1252	Etheostoma spilotum	1044
synagris	1270	Hypohomus	1042
vivanus	1262, 2858	nicaraguensis, Carcharhinus	39, 2747
Neomuraena	392	Eulamia	39
nigromarginata	400	Gambusia	682
Neosebastes	1839	Heros	1532
Neostoma	581	Neotroplus	1542
hathyphilum	583	Pimelodus	152
Neozærces	2426	Rhandia	152
pulcher	2426	nicholsi, Halicheres	1592
nepheus, Lepenotus	1005	Iridio	1591
Tetradon	1733	Platyglossus	1592
turgidus	1733	nicholsii, Gobius	2218
nerka, Oncorhynchus	481	nienhofii, Echenels	2272
kemnerlyi	483	nigellus, Hyberhynchius	217
Salmo	483	niger, Agonus	2069
Nerophis	774	Amblodon	169
anguinens	774	Ammocetes	14
heckeli	774	Aspidorporus	2069
maculata	774	Astronesthes	586
martinicensis	774	Balistes	1711
nerinianns, Aelpenser	106	Bubalichthys	164
Nestia	818	Centrolophus	963
Nettastoma procerum	366	Chiasmodon	2291
Nettastomidie	364	Chiasmodus	2292
Netuma	119, 120, 126, 2704, 2765	Conger	355
dubia	126, 2765	Cottus	1983, 1986
elattura	128, 2769	Esox	626
graniticassis	126, 2764	Leuciscus	235
insculpta	127, 2765	Malacosteus	503
insularum	2770	Myoxocephalus	1985
keesleri	127, 2765	Perca	963
osculi	127, 2768	Scomber	948
planiceps	127, 2766	Sparus	960
platypogon	127, 2767	Squalius	235
proops	124	Tantoga	1577
quadriscutis	126	Thyrstites	879
strictiennisis	126, 2765	Zona	936
neucræta, Echenels	2269	Nigger Chub	327
neustrianns, Gasterosteus	747		

	Page.		Page.
<b>Nigger Dick</b> .....	327	<b>nigrirostris</b> , <i>Chaetodon</i> .....	1673, 1674
<b>Nigger-fish</b> .....	1144, 1146	<i>Sarothroilus</i> .....	1674
<b>nigra</b> , <i>Balistes</i> .....	1711	<b>nigrita</b> , <i>Garrupa</i> .....	1161
<i>Elacate</i> .....	948	<b>nigritus</b> , <i>Epinephelus</i> .....	1162
<i>Gila</i> .....	235	<i>Serranus</i> .....	1161
<i>Muraena</i> .....	355	<b>nigrocastaneus</b> , <i>Gymnothorax</i> .....	390
<i>maculata</i> .....	395	<b>nigrocinctus</b> , <i>Sebastes</i> .....	1828
<i>Narcine</i> .....	78	<i>Sebastichthys</i> .....	1828
<b>nigrescens</b> , <i>Centropomus</i> .....	1119	<i>Sebastodes</i> .....	1827
<i>Coryphæna</i> .....	1200	<b>nigrofasciata</b> , <i>Hydrargyra</i> .....	641
<i>Cycleptus</i> .....	169	<b>nigrofasciatum</b> , <i>Cichlasoma</i> .....	1525
<i>Exoglossum</i> .....	327	<i>Etheostoma</i> .....	1030
<i>Leuciscus</i> .....	233	<b>nigrofasciatus</b> , <i>Fundulus</i> .....	641
<i>Pimelodus</i> .....	137	<i>Hadropterus</i> .....	1038
<i>Salmo</i> .....	507	<i>Heros</i> .....	1525
<i>Serranus</i> .....	1200	<b>nigrolineatus</b> , <i>Diodon</i> .....	1749
<i>Squalius</i> .....	234	<b>nigromaculatus</b> , <i>Cantharus</i> .....	987
<i>Tigoma</i> .....	234	<b>nigromannus</b> , <i>Pleuronectes</i> .....	2657
<b>nigricans</b> , <i>Acanthurus</i> .....	1692	<b>nigromarginata</b> , <i>Neomuraena</i> .....	400
<i>Bathylaco</i> .....	540	<i>Sidera</i> .....	400
<i>Camarina</i> .....	1382	<b>nigromarginatus</b> , <i>Gymnothorax</i>	
<i>Catostomus</i> .....	181	<i>ocellatus</i> .....	400
<i>etowannus</i> .....	181	<i>Lycodontis ocellatus</i> .....	399
<i>Centrepristes</i> .....	1200	<b>nigropunctata</b> , <i>Perca (Pomacampus)</i>	1021
<i>Cottus</i> .....	1973	<b>nigrostrigatus</b> , <i>Mugil</i> .....	817
<i>Cypselurus</i> .....	2836	<b>nigrotaniata</b> , <i>Chloa</i> .....	264
<i>Enchelycore</i> .....	389	<i>Hybognathus</i> .....	214
<i>Entomacrolus</i> .....	2399	<b>nigrotaniatus</b> , <i>Graodius</i> .....	264
<i>Exocoetus</i> .....	737	<i>Notropis</i> .....	264
<i>Girella</i> .....	1382	<b>nigrum</b> , <i>Boleosoma</i> .....	1056
<i>Huro</i> .....	1012	<i>effulgens</i> .....	1058
<i>Hylomyzon</i> .....	161	<i>maculaticeps</i> .....	1058
<i>Hypoplectrus unicolor</i> ..	1193	<i>mesæum</i> .....	1059
<i>Ictalurus</i> .....	137	<i>olmstedii</i> .....	1057
<i>Istiophorus</i> .....	891	<i>Etheostoma</i> .....	1057
<i>Labrax</i> .....	1135	<i>vexillare</i> .....	1058
<i>Makaira</i> .....	891	<i>Petromyzon</i> .....	14
<i>Muraena</i> .....	390	<b>Nine-spined Stickleback</b> .....	745
<i>Petromyzon</i> .....	10	<b>niphobles</b> , <i>Epinephelus</i> .....	2853
<i>Pimelodus</i> .....	137	<i>Sparisoma</i> .....	1633
<i>Plectropoma</i> .....	1193	<b>Nissnee Trout</b> .....	503
<b>nigriculus</b> , <i>Serranus</i> .....	1153	<b>nitens</b> , <i>Congermuraena</i> .....	357
<i>Acronurus</i> .....	1693	<i>Ophisoma</i> .....	357
<b>nigriclavis</b> , <i>Amelurus</i> .....	142	<b>nitida</b> , <i>Julis</i> .....	1608
<i>Amiurus</i> .....	142	<i>Moniana</i> .....	265
<i>Gronias</i> .....	142	<i>Pomotis</i> .....	1003
<b>nigriplunis</b> , <i>Argyrosomus</i> .....	472	<b>nitidissima</b> , <i>Julis</i> .....	1608
<i>Auchenopterus</i> .....	2369	<b>nitidissimum</b> , <i>Thalassoma</i> .....	2859
<i>Bathygonus</i> .....	2078	<b>nitidissimus</b> , <i>Chlorichthys</i> .....	1608
<i>Cfinus</i> .....	2370	<b>nitidum</b> , <i>Pristipoma (Hamulopsis)</i> ..	1326
<i>Coregonus</i> .....	472	<i>Thalassoma</i> .....	1608, 2859
<i>Cotylin</i> .....	2332	<b>nitidus</b> , <i>Alburnus</i> .....	293
<i>Gobiesox</i> .....	2332	<i>Balistes</i> .....	1709
<i>Minnilus</i> .....	299	<i>Brachydeuterns</i> .....	1326
<i>Rhypticus</i> .....	1234	<i>Chlorichthys</i> .....	1608
<i>Rypticus</i> .....	1234		

	Page.		Page.
nitidus, Leuciscus .....	221	normalis, Bassozetus .....	2507
Pomadasis .....	1326	Northern Barracuda .....	825
Promoxis .....	987	Striped Gurnard .....	2167
Salmo .....	509	Sucker .....	176
Salvelinus .....	509	Whiting .....	1475
nivalis, Salmo alpinus .....	509	North-River Shad .....	427
nivea, Cliola .....	278	norvegica, Perca .....	1761
niveatus, Epinephelus .....	1156	norvegicus, Coryphænoides .....	2579
Microspathodon .....	1567	Lepidoleprus .....	2579
Pomacentrus .....	1568	Sebastes .....	1761
Serranus .....	1150	norwegianus, Squalus .....	57
niveiventris, Amlurus .....	138	norwegica, Perca .....	1761
Ictalurus .....	138	No-shee Trout .....	503
niveus, Hybopsis .....	278	nostras, Liparis .....	2118
Notropis .....	277	notabilis, Argyreus .....	309
chloristinus .....	278	Notacanthinae .....	613
Photogenis .....	278	Notacanthus .....	614
Nivica .....	1066, 1082	nalis .....	615
nivipes, Emblemaria .....	2402	challengeri .....	618
nivosus, Cottus .....	1985	chemnitzii .....	614
Myoxocephalus .....	1984	nasus .....	615
Sebastes .....	1834	phasganornus .....	616
Sebastes .....	1833	rissoanus .....	618
nobilior, Esox .....	629	rostratus .....	617
nobilis, Atractoscion .....	1413	Notarins .....	119, 2764
Costreus .....	1413	notata, Atherina .....	800
Conodon .....	1324	Atherinichthys .....	800
Cynoscion .....	1413	Belone .....	711
Esox .....	629	Cliola .....	274
Gambusia .....	682	Cyprinella .....	274
Grystes .....	1012	Dorosoma .....	416
Heterandria .....	682	Lepomis .....	1011
Johinus .....	4413	Malthua .....	2737
Perca .....	1324	Megalops .....	432
Nocomis .....	314, 315, 322	Meuidia .....	800
amblops .....	321	menidia .....	2840
bellicus .....	323	Perca .....	1024
hyostomus .....	316	notatum, Etheostoma .....	1070
milneri .....	324	Hæmulon .....	1297
nobrascensis .....	323	Pristipoma .....	1321
rubrifrons .....	320	notatus, Balistes .....	1709
nocomis, Notropis .....	268	Fundulus .....	659
nocturna, Collettia .....	567	Hyborhynchus .....	218
Echidna .....	402	Labrax .....	1132
noctarnus, Noturnus .....	146	Lepomis .....	1008
Pæcilophis .....	403	Minnilus .....	218
Schilbeodes .....	146	Notropis .....	274
Noire, Oreille .....	1261	Pimeledus .....	135
no Mai, Yanagl .....	1830	Pimephales .....	218
Nomeidæ .....	948	Pomotis .....	1008
Nomeinæ .....	949	Porichthys .....	2321
Nomeus .....	949	Semotilus .....	659
gronovii .....	949	Tylosurus .....	710, 711
maculatus .....	950	Zygonectes .....	659
maculosus .....	950	notemigonoides, Notropis .....	292
oxyurus .....	950	Notemigonus .....	249, 250



Page.		Page.		Page.
2507	<i>Notemigonus americanus</i> .....	251	<i>Notropis atherinoides</i> .....	254, 293
825	<i>auratus</i> .....	250	<i>aztecus</i> .....	258
2167	<i>chrysoleucus</i> .....	250	<i>bellus</i> .....	297
176	<i>gardoneus</i> .....	251	<i>bifrenatus</i> .....	258
1475	<i>ischanus</i> .....	251	<i>blennius</i> .....	251, 2800
427	<i>leptosomus</i> .....	250	<i>boops</i> .....	268
1761	<i>lucidus</i> .....	299	<i>braytoni</i> .....	264
2579	<i>occidentalis</i> .....	247	<i>bubalinus</i> .....	273
2579	<i>nothochir</i> , <i>Ophichthys</i> .....	380	<i>buchanani</i> .....	2800
1761	<i>Quassiremus</i> .....	380	<i>cæruleus</i> .....	277
57	<i>Nothonotus</i> .....	1066, 1067, 1076	<i>callisema</i> .....	272
1761	<i>cinereus</i> .....	1078	<i>callistius</i> .....	276
503	<i>inscriptus</i> .....	1072	<i>camurus</i> .....	279
2118	<i>jordani</i> .....	1080	<i>cayuga</i> .....	260, 2799
309	<i>maculatus</i> .....	1077	<i>atrocaudalis</i> .....	260
613	<i>rufilineatus</i> .....	1079	<i>cerasinus</i> .....	283
614	<i>sanguifluus</i> .....	1077	<i>cercostigma</i> .....	274
615	<i>tessellatus</i> .....	1078	<i>chalybeus</i> .....	288
618	<i>thalassinum</i> .....	1072	<i>chamberlaini</i> .....	2800
614	<i>vulneratus</i> .....	1077	<i>chihuahua</i> .....	265
615	<i>nothus</i> , <i>Cestrens</i> .....	1407	<i>chiliticus</i> .....	287
616	<i>Cynoscion</i> .....	1406	<i>chlorocephalus</i> .....	286
618	<i>Otolithus</i> .....	1407	<i>chromomus</i> .....	288
617	<i>Notidanoid Sharks</i> .....	16	<i>coccogenis</i> .....	284
119, 2764	<i>Notidanus</i> .....	18	<i>cornutus</i> .....	281
800	<i>griseus</i> .....	19	<i>cyanens</i> .....	283
800	<i>Notistium</i> .....	890	<i>frontalis</i> .....	283
711	<i>Notocanthida</i> .....	613	<i>dilectus</i> .....	294, 2801
274	<i>Notoglanis</i> .....	149	<i>eurytomus</i> .....	277
274	<i>Notogrammus</i> .....	2430	<i>formosus</i> .....	271
416	<i>rothrocki</i> .....	2440	<i>frotensis</i> .....	261
1011	<i>Notorhynchus</i> .....	17	<i>frigidus</i> .....	271
2737	<i>borealis</i> .....	18	<i>fumeus</i> .....	204
432	<i>maculatus</i> .....	17	<i>galacturus</i> .....	279
800	<i>Notoscopelus</i> .....	554	<i>garmani</i> .....	281
2840	<i>castanens</i> .....	556	<i>germanus</i> .....	261
1024	<i>caudispinosus</i> .....	556	<i>gilberti</i> .....	266
1070	<i>margaritifera</i> .....	555	<i>heterodon</i> .....	261
1297	<i>quereinus</i> .....	555	<i>heterolepis</i> .....	260
1321	<i>Notosema</i> .....	2635	<i>hudsonius</i> .....	269
1709	<i>dilecta</i> .....	2636	<i>amarus</i> .....	270
653	<i>dilectum</i> .....	2635	<i>salदानus</i> .....	270
218	<i>notospilotus</i> , <i>Astrolytes</i> .....	1899	<i>scelene</i> .....	269, 2801
1132	<i>Julldio</i> .....	1603	<i>hypaelopterus</i> .....	280
1008	<i>Pseudojulis</i> .....	1603	<i>illecebrosus</i> .....	268
218	<i>Notropis</i> .....	254, 257-258, 290	<i>jejunus</i> .....	290
274	<i>tenecus</i> .....	266	<i>jordani</i> .....	250
135	<i>albamae</i> .....	298	<i>kanawha</i> .....	264
218	<i>albeolus</i> .....	259, 283	<i>lacertosus</i> .....	284
1008	<i>altipinnis</i> .....	287	<i>lateralis</i> .....	263
2321	<i>amabilis</i> .....	291	<i>leuciodus</i> .....	291
659	<i>amœnus</i> .....	296	<i>lirus</i> .....	297
710, 711	<i>analostanus</i> .....	279	<i>longirostris</i> .....	267
659	<i>anogonus</i> .....	259, 260	<i>louisiana</i> .....	2801
292	<i>argo</i> .....	294	<i>ludibundus</i> .....	273
249, 250	<i>ariommus</i> .....	290	<i>lutipinnis</i> .....	280, 2800

	Page.		Page.
<i>Notropis lutrensis</i> .....	271	<i>Notropis umbratilis umbratilis</i> .....	209
<i>lythrus</i> .....	300	<i>umbrifer</i> .....	274
<i>macdonaldi</i> .....	284	<i>venustus</i> .....	274, 275
<i>macrolepidotus</i> .....	299	<i>volucellus</i> .....	263
<i>macrostomus</i> .....	274	<i>welaka</i> .....	2799
<i>maculatus</i> .....	259	<i>whiplii</i> .....	278
<i>matutinus</i> .....	301	<i>xenoccephalus</i> .....	289
<i>megalops albeolus</i> .....	284	<i>xenurus</i> .....	280
<i>metallicus</i> .....	297	<i>zonatus</i> .....	285
<i>micropteryx</i> .....	296	<i>zonistius</i> .....	277, 285
<i>nigroteniatus</i> .....	264	<i>nottii</i> , <i>Fundulus</i> .....	656
<i>niveus</i> .....	277	<i>Zygonectes</i> .....	657
<i>chloristius</i> .....	278	<i>Noturus</i> .....	143
<i>nocomis</i> .....	268	<i>elassochir</i> .....	147
<i>notatus</i> .....	274	<i>eleutherus</i> .....	148, 149
<i>notemigonoides</i> .....	292	<i>exilis</i> .....	147
<i>nux</i> .....	267	<i>flavus</i> .....	144
<i>orca</i> .....	289	<i>funebri</i> .....	147
<i>ornatus</i> .....	270	<i>furius</i> .....	149
<i>ozarcanus</i> .....	265	<i>gilberti</i> .....	148
<i>phenacobius</i> .....	263	<i>gyrinus</i> .....	146
<i>photogenis</i> .....	295, 296	<i>insignis</i> .....	147
<i>plptolepis</i> .....	260	<i>leptacanthus</i> .....	146
<i>procne</i> .....	264	<i>luteus</i> .....	144
<i>proserpina</i> .....	272	<i>marginatus</i> .....	147
<i>pyrrhomelas</i> .....	280	<i>miurus</i> .....	148
<i>reticulatus</i> .....	262	<i>nocturnus</i> .....	146
<i>roseipinnis</i> .....	298	<i>occidentalis</i> .....	144
<i>roseus</i> .....	237	<i>platycephalus</i> .....	144
<i>rubricroceus</i> .....	286	<i>sialis</i> .....	146
<i>rubrifrons</i> .....	295	<i>Novacula</i> .....	1617
<i>sabinæ</i> .....	262	<i>cærulea</i> .....	1653
<i>scabriceps</i> .....	290	<i>cultrata</i> .....	1619
<i>scepticus</i> .....	296	<i>lineata</i> .....	1619
<i>scopifer</i> .....	291	<i>martincensis</i> .....	1617
<i>scylla</i> .....	263	<i>mundiceps</i> .....	1618
<i>shumardi</i> .....	268	<i>mundicorpus</i> .....	1620
<i>simus</i> .....	267	<i>novacula</i> , <i>Coryphæna</i> .....	1619
<i>socius</i> .....	292	<i>Xyrichtys</i> .....	1619
<i>spectrunculus</i> .....	265	<i>Novaculichthys</i> .....	1613
<i>stigmaturus</i> .....	275	<i>infirmus</i> .....	1616
<i>stilbius</i> .....	293	<i>murtincensis</i> .....	1616
<i>swalmi</i> .....	290	<i>rosipes</i> .....	1614
<i>telescopus</i> .....	292	<i>ventralis</i> .....	1615
<i>arcansanus</i> .....	292	<i>novæ-angliæ</i> , <i>Coregonus</i> .....	465
<i>toxanus</i> .....	274	<i>novæ-orleanensis</i> , <i>Anguilla</i> .....	348
<i>topeka</i> .....	266	<i>novæ-terræ</i> , <i>Anguilla</i> .....	348
<i>trichroistius</i> .....	275	<i>novæboracensis</i> , <i>Exocætus</i> .....	735, 736
<i>umbratilis</i> .....	298	<i>Naucrates</i> .....	900
<i>ardens</i> .....	301	<i>Vomer</i> .....	934
<i>atripes</i> .....	300	<i>novemfasciatus</i> , <i>Lutjanus</i> .....	1253
<i>cyanoccephalus</i> .....	300	<i>Neomænis</i> .....	1252
<i>fasciolaris</i> .....	301	<i>novemlineatus</i> , <i>Chasmodes</i> .....	2393
<i>lythrus</i> .....	300	<i>rhollis</i> .....	2393
<i>matutinus</i> .....	301	<i>novemmaculatus</i> , <i>Diodon</i> .....	1746
<i>punctulatus</i> .....	301	<i>novemradiata</i> , <i>Agosia</i> .....	312

Page.		Page.		Page.
299	nox, Anchenopterus .....	2373	obliquatus, Cestrens .....	1405
274	Cremnobates .....	2374	Cynoscion .....	1405
74, 275	nubila, Agosia .....	212, 311	Diabasis .....	1304
263	carringtonii .....	311	Otolithus .....	1405
2799	Apocope .....	311	obliquus, Mugil .....	1459
278	Chola .....	215	oblonga, Chenopsetta .....	2633
289	Hybognathus .....	215	Platessa .....	2630
280	nubilus, Alburnops .....	215	oblongior, Plmelepterus .....	1388
285	Argyreus .....	311	oblongiuaculus, Balistes .....	1720
277, 285	Centrobrennius .....	2438	oblongus, Corpore glabro .....	2657
656	Ceraticthys .....	312	oblongum, Moxostoma .....	186
657	Leptobrennius .....	2438	oblongus, Cyprinus .....	186
143	Lumpenus .....	2438	Erinymzon sucetta .....	186
147	Stichæus .....	2438	Gobius .....	2264
148, 149	nuchalis, Ailurichthys .....	117	Heros .....	1535
147	Arius .....	131	Ortbagoriscus .....	1756
144	Chanos .....	415	Ostracion cathelopteleo..	1728
147	Hybognathus .....	213	conico .....	1745
149	Pseudoscarus .....	1654	glaber .....	1735
148	Searus .....	1654	holacanthus .....	1746
146	Tachisurus .....	131	Paralichthys .....	2632
147	Tachysurus .....	131, 2782	Pleuronectes .....	2633
146	nuchifilis, Blennius .....	2383	Pseudorhombus .....	2630
144	nuobipinnis, Clinus .....	2362	Sebastes .....	1830
147	Labrisomus .....	2362	Sebastes .....	1830
148	nuda, Cotylis .....	2331	Sparus .....	2276
146	nudus, Cyclopterus .....	2336	obscura, Liopsetta .....	2651
144	Gobiesox .....	2331	obscuratus, Gymnothorax .....	389
144	Lepadogaster .....	2331	Lycodontis .....	389
146	nuecensis, Dioplites treenlii .....	1012	Pomacentrus .....	1552, 1555
1617	Grystes .....	1012	obscurus, Carcharhinus .....	35
1653	nugator, Bryostomma .....	2410	Carcharias .....	35
1619	Numbfish .....	77	Centrarchus .....	1012
1619	nummifer, Carpiodes .....	166	Pleuronectes .....	2651
1617	Salmo .....	508	Pomotis .....	1006
1618	Nurse .....	57	Squalus .....	35
1620	Nurse Shark .....	25, 26	obtusa, Labrus capite .....	1609
1619	nuttalli, Exocoetus .....	737	Ophisoma .....	355
1619	nuttingii, Antennarius .....	2723	Raia .....	2751
1613	Chaunax .....	1726, 2727	obtusirostris, Acipenser .....	106
1616	nux, Notropis .....	267	Exocoetus .....	730
1616	Nyctophus .....	569	Halocyprsolus .....	730
1614	obesa, Aigalsea .....	246	obtusum, Hemulon .....	1319
1615	Tigoma .....	233	obtusus, Pseudoscarus .....	1654
465	obesus, Amirus .....	141	Rhinichthys .....	308
348	Aponotis .....	993	Searus .....	1654
348	Echinorhinus .....	85	Squalus .....	39
735, 736	Ennacanthus .....	993	obvelatus, Prionodon .....	35
900	Leuciscus .....	246, 282	Occa .....	2043
934	Leucos .....	246	dodecaedron .....	2044
1253	Pomotis .....	993	verru cosa .....	2043
1252	Squalius .....	233	occa, Cælorhynchus .....	2587
2393	obeyense, Ethcostoma .....	1092	Macrurus .....	2588
2393	oblarius, Gasterosteus .....	750	Pristis .....	61
1746	obliqua, Sciaena .....	1459	occidentalis, Abramis .....	247
312	obliteratus, Alutarius .....	1720	Amia .....	113

	Page.		Page.
occidentalis, <i>Catostomus</i> .....	178	ocellatus, <i>Johnius</i> .....	1454
Chorinemis.....	898	Lophius.....	2722
Conger.....	355	Lycodontis.....	399
Epigonus.....	1112	nigromargina-	
Gasterosteus.....	745	tus.....	399
Girardinus.....	689	saxicola.....	399
Heterandria.....	689	Malacoctenus.....	2356, 2809
Leucosomus.....	247	Murrenophis.....	384
Luxilinus.....	247	Neobythites.....	2513
Luxilus.....	247	Ophichthus.....	383
Malacocephalus.....	2570	Ophichthys.....	384
Monacanthus.....	1715	Opisthocentrus.....	2429
Myripristis.....	847	Platophrys.....	2663
Notemigonus.....	247	Priodonophis.....	399
Noturus.....	144	Rhomboidichthys.....	2664
Oligoplites.....	898	Rhombus.....	2664
Scombroides.....	899	Sciaenops.....	1453
Tetronarce.....	77	Zenopsis.....	1660
Torpedo.....	77	Zeus.....	1661
occipitalis, <i>Scorpena</i> .....	1854	ocellicauda, <i>Amia</i> .....	113
Ocean Pipefish.....	774	ocellifer, <i>Clinus</i> .....	2353
Tang.....	1693	octodecimspinosus, <i>Cottus</i> .....	1976
Turbot.....	1706	Myoxocephalus.....	1976
Oceanic Bonito.....	868	octofilis, <i>Polynemus</i> .....	830
oceanica, <i>Anguilla</i> .....	355	Trichidion.....	830
oceanicus, <i>Blennius</i> .....	2379	Octogrammus.....	1866
Gobionellus.....	2230	pallasi.....	1870
Gobius.....	2230	octogrammus, <i>Chirus</i> .....	1870
ocella, <i>Rhinichthys</i> .....	307	Hexagrammos.....	1869
ocellaris, <i>Chaenopsetta</i> .....	2630	Labrax.....	1870
Fundulus.....	642, 2827	octonemus, <i>Polydactylus</i> .....	830
Platessa.....	2630	Polynemus.....	830
Pseudorhombus.....	2630	Trichidion.....	830
ocellata, <i>Bollmannia</i> .....	2238	oculata, <i>Sebastes</i> .....	1832
Corvina.....	1454	Squatina.....	59
Murena.....	399	oculatus, <i>Anthias</i> .....	1283
Perca.....	1454	Balistes.....	1707
Raja.....	69	Bryttus.....	1094
Raja.....	68	Centropristis.....	1283
Sciaena.....	1454	Etelis.....	1282
Sidera.....	399	Hesperanthias.....	1283
ocellatum, <i>Ophidion</i> .....	2430	Icelinus.....	1895
ocellatus, <i>Acanthoectus</i> .....	1976	Latebrus.....	1115
Anarrhichthys.....	2448	Lepidosteus.....	111
Antennarius.....	2721	Lumpenus.....	2433
Chaenopsis.....	2403	Macrops.....	1283
Clavodon.....	1674	Myrichthys.....	376
Citharichthys.....	2673	Pisodonophis.....	376
Clinus.....	2357	Orthogoriscus.....	1754
Gymnothorax.....	399	Scombrops.....	1114
nigromar-		Sebastes.....	1832
ginatus.....	400	Serranus.....	1283
saxicola.....	399	oculocirris, <i>Emblemaria</i> .....	2403
Hemirhombus.....	2673	oculofasciatus, <i>Blepsias</i> .....	2021
He. petoichthys.....	384	Nautichthys.....	2020, 2021
Hippoglossus.....	2673	oculo-radiato, <i>Turdus</i> .....	1591

Page.		Page.		Page.
1454	oculo, <i>Turdus radlato</i> .....	1703	Ojanco .....	1272
2722	<i>Ocyanthias</i> .....	1227	ojanco, <i>Lutjanus</i> .....	1273
309	<i>martineensis</i> .....	1228	<i>Mesoprion</i> .....	1273
399	<i>Ocyurus</i> .....	1275	okeechobeensis, <i>Ameiurus</i> .....	139
399	<i>aurovittatus</i> .....	1270	<i>Ictalurus</i> .....	139
2869	<i>chrysurus</i> .....	1275	Old Wench .....	1703
384	<i>lutjanoides</i> .....	1261	Wife .....	940, 1458, 1649, 1703
2513	<i>rijgersmoei</i> .....	1276	Black .....	1711
383	<i>ocyurus</i> , <i>Centropriates</i> .....	1200	<i>olfax</i> , <i>Epinephelus</i> .....	1183
384	<i>Kyphosus</i> .....	1300	<i>Mycteroperca</i> .....	1183
2429	<i>Pimelepterus</i> .....	1300	<i>ruberrima</i> .....	1183
2663	<i>Sectator</i> .....	1389	<i>Serranus</i> .....	1183
399	<i>Serranus</i> .....	1201	<i>olfersi</i> , <i>Argyropoecilus</i> .....	604
2664	<i>Odontanthias asperilingula</i> .....	1227	<i>Pleurothyrus</i> .....	604
2664	<i>martineensis</i> .....	1228	<i>Sternoptyx</i> .....	604
1453	<i>Odontaspis</i> .....	46	<i>olidus</i> , <i>Hypomesus</i> .....	525
1660	<i>americanus</i> .....	47	<i>Mesopus</i> .....	525
1661	<i>Odontognathus</i> .....	437	<i>Salmo</i> ( <i>Osmerus</i> ) .....	525
113	<i>mucronata</i> .....	438	<i>oligopsis</i> , <i>Abramis</i> .....	294
2353	<i>mucronatus</i> .....	438	<i>Alburnus</i> .....	294
1976	<i>panamensis</i> .....	438	<i>Minnilus</i> .....	294
1976	<i>Odontogobius</i> .....	2210	<i>Oligocephalus</i> .....	1066, 1068, 1083
830	<i>Odontopyxis</i> .....	2085	<i>grahami</i> .....	1089
830	<i>frenatus</i> .....	2075	<i>humeralis</i> .....	1097
1866	<i>leptorhynchus</i> .....	2076	<i>leonensis</i> .....	1089
1870	<i>trispinosus</i> .....	2085	<i>pulchellus</i> .....	1089
1870	<i>trispinosus</i> .....	2086	<i>Oligocottus</i> .....	2013, 2864
1869	<i>Odontoscium</i> .....	1425	<i>aenticeps</i> .....	2016
1870	<i>dentex</i> .....	1425	<i>analis</i> .....	2013
830	<i>xanthops</i> .....	1427	<i>borealis</i> .....	2014
830	<i>Odontoscium archidium</i> .....	1432	<i>embryum</i> .....	2017
830	<i>Odontostomidae</i> .....	597	<i>globiceps</i> .....	2017
1832	<i>arstedii</i> , <i>Selene</i> .....	935	<i>maculosus</i> .....	2013
59	<i>Tetragonopterus</i> .....	334	<i>snyderi</i> .....	2871
1283	<i>ogac</i> , <i>Gadus</i> .....	2542	<i>oligodon</i> , <i>Osmerus</i> .....	2824
1707	<i>ogat</i> , <i>Gadus</i> .....	2542	<i>Polynemus</i> .....	830
1004	<i>Ogcocephalidae</i> .....	2735, 2739	<i>Oligolepis</i> .....	2210
1283	<i>Ogcocephalus</i> .....	2736	<i>oligopeltis</i> , <i>Acipenser</i> .....	105
1282	<i>elater</i> .....	2739	<i>Oligoplites</i> .....	898
1283	<i>nasutus</i> .....	2737	<i>altus</i> .....	899, 2844
1895	<i>radiatus</i> .....	2738	<i>inornatus</i> .....	899
1115	<i>veapertilio</i> .....	2737	<i>mundus</i> .....	2844
111	<i>Ogilbia</i> .....	2502	<i>occidentalis</i> .....	898
2433	<i>cayorum</i> .....	2503	<i>saliens</i> .....	899, 2844
1283	<i>ventralis</i> .....	2503	<i>pulometa</i> .....	899
376	<i>oglina</i> , <i>Megalops</i> .....	432	<i>saurus</i> .....	898
376	<i>oglinum</i> , <i>Opisthonema</i> .....	432	<i>Oligopodus</i> .....	955
1754	<i>Ognichodes</i> .....	2263	<i>olisthostoma</i> , <i>Gerres</i> .....	1377
1114	<i>ohiensis</i> , <i>Acipenser</i> .....	100	<i>olisthostomus</i> , <i>Gerres</i> .....	1376
1832	<i>Esox</i> .....	630	<i>olivacea</i> , <i>Pecilia</i> .....	659
1283	<i>Lucius masquinongy</i> .....	629	<i>olivaceus</i> , <i>Leucus</i> .....	244
2403	<i>Ohio Sturgeon</i> .....	106	<i>Rutilus</i> .....	244
2021	<i>oidoensis</i> , <i>Cichla</i> .....	1012	<i>olivaris</i> , <i>Hopladelus</i> .....	143
20, 2021	<i>Oil Fish</i> .....	879	<i>Leptops</i> .....	143
1501	<i>Shark</i> .....	32	<i>Pelodichthys</i> .....	143
	<i>Oja</i> , <i>Pego</i> .....	2699	<i>Pilodictis</i> .....	143

	Page.		Page.
olivaris, Silurus .....	143	ontariensis, Thymallus .....	518
ohustedii, Bolcosoma brevipinnis ...	1057	montanus ..	519
nigrum .....	1057	signifer ....	519
Etheostoma .....	1057	Onus .....	2529, 2558
olriki, Aspidophoroides .....	2089	riall .....	2530
Ombre Chevalier .....	508	onychus, Cottus .....	1953
omnata, Discopyge .....	78	Oolachan .....	521
Heterandria .....	664	Oonidus .....	1738
Lucania .....	663	Ooze Eels .....	349
Opisthognathus .....	2283	Opah .....	954
ommatum, Opisthognathus .....	2282	opah, Zeus .....	955
omnatus, Paralichthys .....	2635	opalescens, Lythrulon .....	1312
omnisco, Salmo mayens .....	487	opalina, Julia .....	1591
omocyaneus, Eleotris .....	2198	opalinum, Myctophum .....	571
omostigma, Genypterus .....	2400	opallnus, Platyglossus .....	1591
Otophidium .....	2490	Open-mouthed Grunt .....	1306
omosudis .....	598	opercularis, Micropogon .....	1461
lowii .....	598	Myxodagnus .....	2305
Oncoccephalus .....	2736	Polydactylus .....	831
Oncocottus .....	2000	Polynemus .....	831
hexacornis .....	2002	Sciæna .....	1461
labradoricus .....	2004	Stolephorus .....	445
quadricornis .....	2001	Trichidion .....	831
Oncorhynchus .....	474, 477, 478	Ophichthys .....	381
ehouicha .....	480	ovionthas .....	381
gorbuscha .....	478	gomesii .....	384
keta .....	478	guttifer .....	383
kisutch .....	480	havannensis .....	382, 2804
lagocephalus .....	479	magniloculis .....	385
lycaolon .....	481, 483	ocellatus .....	383
nerka .....	481	ophis .....	2804
kennerlyi .....	483	parilis .....	386
orientalis .....	480	puncticeps .....	382
paucidens .....	483	retropinnis .....	383
protens .....	478	rugifer .....	384
quinnat .....	480	triserialis .....	384
sanguinolentus .....	481	zophochir .....	385
sconleri .....	478	Ophichthyidæ .....	372
tschawytscha .....	479	Ophichthys .....	381, 382
tsuppitch .....	481	acuminatus .....	377
oncida, Catostomus .....	193	(Sphagobranchnus) an-	
Moxostoma .....	193	guiformis .....	374
Ptychostomus .....	193	chrysops .....	385
Oncirodes .....	2732	crocodilinus .....	388
oschrichtii .....	2732	gomesii .....	385
Oncirodinae .....	2728	guttifer .....	383
ongus, Serranus .....	1154	intertinctus .....	387
onitis, Hiatula .....	1579	magniloculis .....	385
Labrus .....	1578	mirus .....	387
Tautoga .....	1578, 1579	nothochir .....	380
Onos .....	2558	ocellatus .....	384
cimbrinus .....	2561	pardalis .....	376
ensis .....	2559	parilis .....	386
reinhardti .....	2559	pauciporis .....	386
rufus .....	2559	pisavarius .....	377
septentrionalis .....	2560	puncticeps .....	382

Page.	Page.	Page.
518	<i>Ophlethys punctifer</i> .....	387
519	<i>retropinnis</i> .....	383
519	<i>schneideri</i> .....	387
2558	<i>triserialis</i> .....	384
2530	<i>xysturus</i> .....	2802
1953	<i>zophochir</i> .....	385
521	<i>ophidianus, Gempylus</i> .....	894
1738	<i>Oplidido</i> .....	2481
349	<i>Ophidoidea</i> .....	2453
954	<i>Ophidoidei</i> .....	782
955	<i>Ophidion</i> .....	2487
1312	<i>beani</i> .....	2487
1591	<i>grællsi</i> .....	2488
571	<i>holbrookii</i> .....	2487
1591	<i>josephi</i> .....	2488
1306	<i>Ophidium</i> .....	2487
1461	<i>atropurpureum</i> .....	2423
2305	<i>brevibarbe</i> .....	2485
831	<i>grællsi</i> .....	2487
831	<i>holbrookii</i> .....	2488
1461	<i>imberbi</i> .....	2443
445	<i>josephi</i> .....	2489
831	<i>marginatum</i> .....	2489
381	<i>mucronatum</i> .....	2419
381	<i>ocellatum</i> .....	2430
384	<i>parii</i> .....	2478
383	<i>pellucidum</i> .....	354
2, 2804	<i>profundorum</i> .....	2484
385	<i>taylori</i> .....	2489
383	<i>unernak</i> .....	2477
2804	<i>viride</i> .....	2477
386	<i>ophidoidea, Liparis</i> .....	2118
382	<i>Ophioblennina</i> .....	2347
383	<i>Ophioblennius</i> .....	2400
384	<i>steindachneri</i> .....	2401
384	<i>webbii</i> .....	2401
385	<i>Ophiodon</i> .....	1875
372	<i>elongatus</i> .....	1875
81, 382	<i>pantherinus</i> .....	1876
377	<i>Ophiodontina</i> .....	1884
	<i>Ophionathus</i> .....	405
	<i>ampullaceus</i> .....	408
	<i>Ophiocion</i> .....	1446, 1447
	<i>adustus</i> .....	1447
	<i>imiceps</i> .....	1451
	<i>sciurus</i> .....	1452
	<i>simulus</i> .....	1449
	<i>strabo</i> .....	1448
	<i>typicus</i> .....	1448
	<i>vermicularis</i> .....	1452
	<i>ophioscion, Corvina</i> .....	1448
	<i>Sciæna</i> .....	1448
	<i>ophis, Muræna</i> .....	382
	<i>Ophichthus</i> .....	2804
	<i>Ophisoma</i> .....	353, 355
	<i>acuta</i> .....	356
	<i>Ophisoma analis</i> .....	356
	<i>balearicum</i> .....	356
	<i>macrurum</i> .....	357
	<i>nitens</i> .....	357
	<i>obtusa</i> .....	355
	<i>prorigerum</i> .....	357
	<i>Ophisomus</i> .....	2414
	<i>Ophisternon</i> .....	342
	<i>Ophisura intertinctus</i> .....	387
	<i>sugillatus</i> .....	387
	<i>Ophisuraphis</i> .....	374
	<i>Ophisurus</i> .....	375, 381
	<i>acuminatus</i> .....	377
	<i>californiensis</i> .....	384
	<i>chrysops</i> .....	385
	<i>crocodilius</i> .....	388
	<i>gomesii</i> .....	385
	<i>guttatus</i> .....	382
	<i>latimaculatus</i> .....	376
	<i>longus</i> .....	377
	<i>parilis</i> .....	386
	<i>remiger</i> .....	384
	<i>xysturus</i> .....	376
	<i>ophryas, Paralichthys</i> .....	2630
	<i>Prionotus</i> .....	2104
	<i>ophthalmicus, Lepomis</i> .....	1001
	<i>Opisthistilus</i> .....	1384
	<i>Opisthocentrina</i> .....	2349
	<i>Opisthocentrus</i> .....	2428
	<i>ocellatus</i> .....	2428
	<i>quinquemaculatus</i> .....	2430
	<i>tennis</i> .....	2430
	<i>Opisthognathida</i> .....	2279
	<i>Opisthognathus</i> .....	2280
	<i>lonchurum</i> .....	2281
	<i>lonchurus</i> .....	2281
	<i>macrognathum</i> .....	2281
	<i>macrognathus</i> .....	2282
	<i>macrops</i> .....	2284
	<i>maxillosus</i> .....	2284
	<i>megastoma</i> .....	2282
	<i>micrognathus</i> .....	2287
	<i>omnata</i> .....	2283
	<i>ommatum</i> .....	2282
	<i>punctatum</i> .....	2281
	<i>punctatus</i> .....	2281
	<i>rhomaleus</i> .....	2285
	<i>scaphiurus</i> .....	2282
	<i>Opisthonema</i> .....	432
	<i>libertate</i> .....	433
	<i>libertatis</i> .....	433
	<i>oglinum</i> .....	432
	<i>Opisthopterus</i> .....	436
	<i>dovii</i> .....	437
	<i>lutipinnis</i> .....	437, 2811
	<i>macrops</i> .....	437

	Page.		Page.
opisthophthalmus, Conger .....	350	oreas, Chrosomus .....	211
Opladelus .....	142	Oregon Brook Trout .....	501
Oplopoma .....	1875	Charr .....	507
pantherina .....	1876	Sturgeon .....	104
oppositus, Monocanthus .....	1716	oregonensis, Cyprinus (Leuciscus) ..	225
Opsanus .....	2315	Leuciscus .....	225
cerapellus .....	2316	Ptychocheilus .....	224, 2796
pardus .....	2316	Oreille Noire .....	1261
tau .....	2315	Oreosoma .....	1662
Opsopoea .....	247, 248, 249	atlanticum .....	1662
Opsopoeodus .....	247, 248	confertum .....	1663
bollmani .....	249	Orestinae .....	631
emilie .....	248	orientalis, Anarrhichas .....	2447
megalops .....	248	Chunos .....	415
oseni .....	248	Oncorhynchus .....	480
Ophthalmolophus .....	2360	Pelamys .....	873
Oquassa Trout .....	514	Salmo .....	480
oquassa, Salmo .....	515	ornata, Amia .....	113
Salvelinus .....	514, 515	Aphoristia .....	2707, 2710
marstoni .....	515	Chiola .....	271
naresi .....	515	Cochlognathus .....	252
Oranheco, Grand .....	1057	Codoma .....	271
Orango Filefish .....	1718	Embiotoca .....	1506
Rockfish .....	1793	Hydrargyra .....	2827
Orbe, Le Diodon .....	1749	Raja .....	70
orbicularis, Atinga alter minor .....	1749	ornatum, Campostoma .....	205
Rhombas .....	966	ornatus, Achirus .....	270 <sup>c</sup>
Orbidus .....	1729	Balistes .....	1 <sup>m</sup>
orbignanus, Exocoetus .....	729	Cochlognathus .....	1 <sup>m</sup>
o'bignyana, Pellona .....	436	Esox .....	626
Platessa .....	2626	Gillellus .....	2299
orbignyana, Conger .....	355	Gnnellus .....	2420
Orbis echinatus .....	1745	Labrus .....	1610
levis variegatus .....	1735	Murenoides .....	2420
muricatus .....	1749	Notropis .....	270
reticulatus .....	1750	Pholis .....	2419
orbis, Cyclopterus .....	2100	Tetradon .....	1742
Eumicrotremus .....	2009, 2100	Ornichthys .....	2148
orbitalis, Pagellus .....	1350	Orqueta .....	937
Sparus .....	1350	orqueta, Chloroscombrus .....	937
orca, Notropis .....	289	orsini, Ozodura .....	1754
Orcella .....	254, 257, 289	orthagoricus, Cephalus .....	1754
orcini, Brama .....	960	Orthagoricus .....	1754
orentti, Leuciscus .....	241	Orthichthys .....	759
Phoxinus .....	242	Orthodon .....	206
Oreynus .....	869, 870	microlepidotus .....	207
alalonga .....	871	orthogrammus, Carangoides .....	928
alliteratus .....	869	Caranx .....	929
pelamys .....	869	Orthonops eos .....	2262
schlegelii .....	870	orthonotus, Ditrema .....	1507
subulatus .....	871	Orthopristis .....	1334, 1335, 1336
thunna .....	869	breviplinnis .....	1341
thynnus .....	870	cantharinus .....	1339, 1340
ordinatus, Chirus .....	1870	chalcus .....	1337
Epinephelus .....	1155	chrysopterus .....	1338
Hexagrammus .....	1870	duplex .....	1339





	Page.		Page.
<i>Otolithus drummondii</i> .....	1409	<i>Oxyottus acuticeps</i> .....	2015, 2864
<i>jamaicensis</i> .....	1406	<i>Orygenseum</i> .....	207
<i>loiarchus</i> .....	1415	<i>pulvernulentum</i> .....	207
<i>magdalenae</i> .....	1410	<i>oxygenius, Polyprion</i> .....	1139
<i>microlepidotus</i> .....	1415	<i>Oxyjulis</i> .....	1601
<i>microps</i> .....	1415	<i>californicus</i> .....	1601
<i>nebulosus</i> .....	1409	<i>modestus</i> .....	1601
<i>nothus</i> .....	1407	<i>Oxylabrax</i> .....	1117
<i>obliquatus</i> .....	1405	<i>Oxylebiinae</i> .....	1864
<i>regalis</i> .....	1407	<i>Oxylebius</i> .....	1878
<i>reticulatus</i> .....	1409	<i>pictus</i> .....	1878
<i>rhomboidalis</i> .....	1404	<i>Oxyloricaria</i> .....	156
<i>squamipinnis</i> .....	1404	<i>Oxymacurus</i> .....	2587
<i>stolzmanni</i> .....	1412	<i>Oxymetopontinae</i> .....	2188
<i>thalassinus</i> .....	1498	<i>Oxyodontichthys</i> .....	381
<i>toeroe</i> .....	1404	<i>brachyurus</i> .....	385
<i>virescens</i> .....	1415	<i>limbatus</i> .....	385
<i>Otophidium</i> .....	2490	<i>macurus</i> .....	385
<i>galeoides</i> .....	2491	<i>oxyptera, Corvina</i> .....	1222
<i>indefatigabile</i> .....	2490	<i>Oxyrhina</i> .....	47
<i>omostigma</i> .....	2490	<i>gamphodon</i> .....	49
<i>otophorus, Eupomacentrus</i> .....	1555	<i>spallanzani</i> .....	49
<i>Pomacentrus</i> .....	1555	<i>oxyrhynchus, Acipenser</i> .....	105
<i>Otrynter</i> .....	1344	<i>Carcharhinus</i> .....	40
<i>caprinus</i> .....	1345	<i>Carcharias</i> .....	41
<i>otrynter, Caranx</i> .....	930	<i>Isurus</i> .....	48, 49
<i>otsego, Coregonus</i> .....	466	<i>Tetrodon</i> .....	1741
<i>ouachita, Haloproterus</i> .....	1035	<i>Oxyurichthys</i> .....	2210
<i>ouananiche, Salmo salar</i> .....	487	<i>Oxyurus</i> .....	353
<i>Ouatilibi</i> .....	1145	<i>oxyurus, Lepisosteus</i> .....	110
<i>Español</i> .....	1140	<i>Nomenus</i> .....	950
<i>outalibi, Eumeacentrus fulvus</i> .....	1146	<i>Oyster-fish</i> .....	1578, 2315
<i>Sorranus</i> .....	1146	<i>ozarcenus, Notropis</i> .....	265
<i>ovale, Syacium</i> .....	2074	<i>Ozodura</i> .....	1753
<i>ovalis, Citharichthys</i> .....	2074	<i>orsini</i> .....	1754
<i>Hemirhombus</i> .....	2074	<i>ozodura, Orthragoriscus</i> .....	1755
<i>Sebastichthys</i> .....	1789	<i>Ozorthus hexagrammus</i> .....	2441
<i>Sebastodes</i> .....	1788	<i>pachycephala, Adinia</i> .....	660
<i>ovatus, Trachynotus</i> .....	942	<i>pachycephalus, Fundulus</i> .....	661
<i>ovicephalus, Sparus</i> .....	1361	<i>Lagocephalus</i> .....	1729
<i>oviceps, Lactophrys</i> .....	1724	<i>Tetrodon</i> .....	1729
<i>oviformis, Chaetodon</i> .....	1668	<i>pachygaster, Spheroides</i> .....	1738
<i>ovigerum, Bathypasma</i> .....	2128	<i>Tetrodon</i> .....	1738
<i>ovinus, Esocx</i> .....	672	<i>Pachylabrus</i> .....	1507
<i>Lebias</i> .....	672	<i>variegatus</i> .....	1508
<i>ovis, Sargus</i> .....	1361	<i>pachylepis, Atherinichthys</i> .....	801
<i>Ovoides</i> .....	1738	<i>Menidia</i> .....	801
<i>orethizon</i> .....	1739	<i>Pachynathus</i> .....	1703
<i>setosus</i> .....	1739	<i>capistratus</i> .....	1704
<i>Ovum</i> .....	1738	<i>triangularis</i> .....	1705
<i>oweni, Halosaurus</i> .....	607	<i>Pachypops</i> .....	1459
<i>Oxybeles</i> .....	2495	<i>biloba</i> .....	1460
<i>oxybrachium, Sparisoma</i> .....	1034	<i>furcatus</i> .....	1459
<i>oxybrachius, Scarus</i> .....	1635	<i>pachypops, Coftus</i> .....	1973
<i>Oxycephas</i> .....	2568	<i>Pachyurus furcatus</i> .....	1460
<i>Oxycottus</i> .....	2015, 2863, 2864		

Page.		Page.		Page.
15, 2864	<i>Pachyrurus squamosissimus</i> .....	1419	<i>pallasii</i> , <i>Pleuronectes</i> .....	2648
207	<i>pacifica</i> , <i>Belone</i> .....	716	<i>Pallasina</i> .....	2648
207	<i>pacifici</i> , <i>Anisotremus</i> .....	1316	<i>aix</i> .....	2050
1139	<i>Batrachoides</i> .....	2314	<i>barbata</i> .....	2049
1601	<i>Batrachus</i> .....	2315	<i>pallida</i> , <i>Aldrovandia</i> .....	611
1601	<i>Conodon</i> .....	1316	<i>Echeneis</i> .....	2272
1601	<i>Pomadasis</i> .....	1316	<i>Lepomis</i> .....	1012
1117	<i>pacificus</i> , <i>Argyreus</i> .....	936	<i>Morone</i> .....	1135
1864	<i>Arteidiellus</i> .....	1906	<i>pallidus</i> , <i>Bodianus</i> .....	1433
1878	<i>Bathylagus</i> .....	530	<i>Catostomus</i> .....	179
1878	<i>Cynicoglossus</i> .....	2655	<i>Eupomotis</i> .....	1006
156	<i>Glyptocephalus</i> .....	2655	<i>Fundulus</i> .....	638, 2827
2587	<i>Larimus</i> .....	1424	<i>Labrax</i> .....	1135
2188	<i>Lobotes</i> .....	2857, 2858	<i>Labrus</i> .....	1005
381	<i>Lutjanus</i> .....	1253	<i>Lepomis</i> .....	1005
385	<i>Lycodes</i> .....	2460	<i>Leucosomus</i> .....	222
385	<i>Lycodopsis</i> .....	2460	<i>Pimelodus</i> .....	135
385	<i>Mesoprion</i> .....	1253	<i>Platygobio</i> .....	326
1222	<i>Microstomus</i> .....	2655	<i>Pomotis</i> .....	1007
47	( <i>Mallotus</i> ) <i>Salmo</i> .....	521	<i>Salmo</i> .....	505
49	<i>Thaleichthys</i> .....	521	<i>palmipes</i> , <i>Prionotus</i> .....	2157
49	<i>Thynnus</i> .....	871	<i>Trigla</i> .....	2150
105	<i>Tylosurus</i> .....	716	<i>paloma</i> , <i>Trachinotus</i> .....	945
40	<i>Paddle, Cock and Hen</i> .....	2096	<i>Palometa</i> .. 940, 941, 942, 943, 965, 966, 967, 2849	
41	<i>Paddle-fish</i> .....	101	<i>media</i> .....	2849
48, 49	<i>Paddle-fishes</i> .....	100	<i>palometa</i> .....	2849
1741	<i>patulus</i> , <i>Hemirhombus</i> .....	2672	<i>simillima</i> .....	2849
2210	<i>pagei</i> , <i>Etheostoma</i> .....	1092	<i>palometa</i> , <i>Chorinemus</i> .....	899
353	<i>Pagellus bajonado</i> .....	1352	<i>Oligoplites saliens</i> .....	899
110	<i>calamus</i> .....	1350	<i>Palometa</i> .....	2849
950	<i>caninus</i> .....	1352	<i>Rhombus</i> .....	966
1578, 2315	<i>humilis</i> .....	1355	<i>Stomateus</i> .....	967
265	<i>microps</i> .....	1355	<i>Palu Brasiliense congener</i> .....	966
1753	<i>milneri</i> .....	1355	<i>palustris</i> , <i>Pecellichthys</i> .....	1102
1754	<i>orbitalis</i> .....	1350	<i>Pammelas</i> .....	963
1755	<i>penna</i> .....	1355	<i>perciformis</i> .....	964
2441	<i>Pagrus</i> .....	1356	<i>pammelas</i> , <i>Melanostigma</i> .....	2479, 2869
660	<i>argenteus</i> .....	1357	<i>Pampanito</i> .....	941
661	<i>pagrus</i> .....	1356	<i>Pámpano</i> .....	930, 933
1729	<i>vulgaris</i> .....	1357	Common .....	944
1729	<i>pagrus</i> , <i>Pagrus</i> .....	1356	<i>Gulf-topail</i> .....	940
1738	<i>Sparus</i> .....	1357	Great .....	943
1507	<i>Pai de Gato</i> .....	1837	Round .....	941
1508	<i>Pajarito</i> .....	721	<i>Pámpanos</i> .....	895, 939
801	<i>pala</i> , <i>Cyprinus</i> .....	415	<i>pampanus</i> , <i>Trachynotus</i> .....	944
801	<i>palearis</i> , <i>Lycodes</i> .....	2466	<i>panamense</i> , <i>Pristipoma</i> .....	1331
1703	<i>Palinurichthys</i> .....	663	<i>panamensis</i> , <i>Achirus</i> .....	2702
1704	<i>perciformis</i> .....	964	<i>Ailurichthys</i> .....	117
1705	<i>Painurnus</i> .....	963	<i>Atherinella</i> .....	805
1450	<i>perciformis</i> .....	964	<i>Azevia</i> .....	2677
1460	<i>pallasii</i> , <i>Octogrammus</i> .....	1870	<i>Bodianus</i> .....	1141
1459	<i>Pallasia</i> .....	1754	<i>Caranx</i> .....	928
1973	<i>Pallasia</i> .....	1753	<i>Citharichthys</i> .....	2677
1460	<i>pallasii</i> .....	1754	<i>Engraulis</i> .....	448
1973	<i>pallasianus</i> , <i>Cophalus</i> .....	1754	<i>Enneacentrus</i> .....	1141
1460	<i>pallasii</i> , <i>Clupea</i> .....	422	<i>Epinephelus</i> .....	1141

	Page.		Page.
panamensis, Felichthys .....	117	papillosum, Syacium .....	2671
Gymnothorax .....	391	papillosus, Barbalifer .....	2661
Ilisha .....	436	Pleuronectes .....	2672
Loricaria .....	157	Ptychostomus .....	189
Menticirrhus .....	1473	Paraclinus .....	2374
Muraena .....	391	chaperi .....	2374
Odontognathus .....	438	Paraconodon .....	1314, 1315, 1316
Parapsettus .....	1069	Paradiodon .....	1744
Pellona .....	436	quadrifasciatus .....	1746
Petrometopon .....	1141	paradoxa, Garmannia .....	2232
Piabncina .....	332	paradoxus, Gobius .....	2232
Pomadasis .....	1331	Psychrolutes .....	2026
Pristigaster (Odontog- nathus) .....	438	Paradules .....	1013
Rabula .....	391	Paragonus .....	2054
Serranus .....	1141	neipenserinus .....	2062
Sidera .....	391	sturioides .....	2063
Solea .....	2702	Parahemiodon .....	156, 157, 158
Stolephorus .....	448	Parahubra .....	1194
Tetragonopterus .....	334	albomaculatus .....	1197
Umbrina .....	1473	clathratus .....	1197, 1198
Panchax .....	633, 2827, 2830	humeralis .....	1196, 1197
pauciradiatus, Cubiceps .....	957	maculatofasciatus .....	1196
pandionis, Glossamia .....	1111	nebulifer .....	1195, 1196
pandora, Clinostomus .....	234	Paralepidide .....	599
Squalius .....	234	Paralepine .....	599
Pañeca .....	2196	Paralepis .....	602
pauvosa, Scorpaena .....	1845	borealis .....	601
pantherina, Muraenophis .....	2805	coregonoides .....	602
Oplopoma .....	1876	coruscans .....	602
pantherinus, Anarchichas .....	2446	internediis .....	600
Costracion .....	21	Paralichthys .....	2624
Ophiodon .....	1876	adpersus .....	2627, 2872
Pseudariodes .....	155	aetnarius .....	2626, 2872
Pantostens .....	169	albiguttus .....	2631
arcopus .....	172	brasiliensis .....	2626
arizona .....	170, 2790	californicus .....	2625, 2626
clarki .....	172	dentatus .....	2629, 2630
columbianus .....	172	lethostigma .....	2630
delphinus .....	171	lethostigmus .....	2630
generosus .....	170	llolepis .....	2624
guzmanensis .....	171	maculosus .....	2626
jarrovi .....	171	magdalena .....	2872
jordanii .....	171	oblongus .....	2632
platyrhynchus .....	170	ommatius .....	2635
plebeius .....	171	ophryas .....	2630
virescens .....	171, 172	sinaloae .....	2872
pantostigmus, Myrichthys .....	2802	squamilentus .....	2631
Papagallo .....	895	stigmatias .....	2636
Papagallos .....	894	woolmanii .....	2628
papalis, Blonda .....	214	Paraliparis .....	2139, 2140
papilio, Molletes .....	1932	cephalus .....	2141
papillifer, Gobiesox .....	2330	copei .....	2143
papilliferus, Chologaster .....	704	ductylosus .....	2144
papillosa, Aramaca .....	2672	holmcelsus .....	2140
papillosum, Moxostoma .....	189	liparinus .....	2139
		mento .....	2142

Page.		Page.		Page.
2671	<i>Paraliparis rosaceus</i> .....	2142	<i>parilis, Ophichthys</i> .....	386
2261	<i>uloclair</i> .....	2144	<i>Ophichthys</i> .....	386
2672	<i>parallelus, Centropomus</i> .....	1122	<i>Ophisurus</i> .....	386
189	<i>Paralonchurus</i> .....	1477, 1478	<i>parkei, Salmo</i> .....	508
2374	<i>dumerili</i> .....	1478	<i>Salvelinus</i> .....	2823
2374	<i>goodei</i> .....	1480	<i>parkeri, Arius</i> .....	126
1315, 1316	<i>petorsi</i> .....	1481	<i>Selenaspis</i> .....	125, 2764
1744	<i>rathbuni</i> .....	1479	<i>Silurus</i> .....	126
1746	<i>Paramacrurus</i> .....	2587	<i>Trachisurus</i> .....	126
2232	<i>Paramia</i> .....	1112	<i>Parua rubicunda</i> .....	1565
2232	<i>paranashmos, Acipenser</i> .....	106	<i>parua, Cichlasoma</i> .....	1519
2026	<i>Paranthias</i> .....	1221	<i>Heros</i> .....	1519
1013	<i>ereolus</i> .....	1222	<i>parmifera, Raja</i> .....	75
2054	<i>farcifer</i> .....	1221	<i>Raja</i> .....	74
2062	<i>Parapomacentrus</i> .....	1549, 1550	<i>parnatus, Setarches</i> .....	1860
2063	<i>Parapsettus</i> .....	1669	<i>Parophrys</i> .....	2637, 2640
156, 157, 158	<i>panamensis</i> .....	1669	<i>ayresi</i> .....	2640
1194	<i>Parques</i> .....	1485	<i>cornosa</i> .....	2639
1197, 1198	<i>Parascorpena</i> .....	1839, 2860	<i>hubbardi</i> .....	2641
1196, 1197	<i>Paraserranus hasselti</i> .....	1205	<i>ischyrus</i> .....	2641
1196	<i>parasiticus, Simenichelys</i> .....	349	<i>isolepis</i> .....	2642
1195, 1196	<i>paraspistes, Caranx</i> .....	923	<i>quadrituberculatus</i> .....	2648
599	<i>Paratractus</i> .....	916, 917, 921	<i>vetulus</i> .....	2640
599	<i>pisquetus</i> .....	921	<i>parovanus, Cyprinodon</i> .....	666
602	<i>Parché</i> .....	1674, 1677	<i>Myloleucus</i> .....	246
601	<i>pardale, Leptopidium</i> .....	2486	<i>Upeneus</i> .....	859
602	<i>pardalis, Epinephelus</i> .....	1183	<i>parra, Diabasis</i> .....	1299
602	<i>Monacanthus</i> .....	1713	<i>Hamulon</i> .....	1297
600	<i>Myeteropereca</i> .....	1181	<i>parrae, Albula</i> .....	411
2624	<i>Ophichthys</i> .....	376	<i>Brama</i> .....	1586
2627, 2872	<i>pardus, Batrachus tan</i> .....	2317	<i>Chetodon</i> .....	1685
2626, 2872	<i>Opsanus</i> .....	2316	<i>Clepticus</i> .....	1586
2631	<i>Parepinephelus</i> .....	1169, 1170, 1180	<i>Exocoetus</i> .....	740
2626, 2626	<i>Parques</i> .....	1485, 1486	<i>Hamulon</i> .....	1297, 1309
2630	<i>aemulatus</i> .....	1487	<i>Parraseranus</i> .....	1203
2630	<i>parva, Aluterus</i> .....	1719	<i>parrianus, Monacanthus</i> .....	1713
2629, 2630	<i>Paraxocetus</i> .....	728	<i>Parrot-fish, Blue</i> .....	1636, 1652
2630	<i>mesogaster</i> .....	728	<i>Dark-green</i> .....	1638
2624	<i>Pargo</i> .....	1244, 1265	<i>Green</i> .....	1657
2626	<i>Amarillo</i> .....	1260	<i>Parrot Fishes</i> .....	1620, 1642
2872	<i>Colorado</i> .....	1264, 1267, 1356	<i>Parrot, Mud</i> .....	1639
2632	<i>Criollo</i> .....	1265	<i>Rose-back</i> .....	1635
2635	<i>de lo Alto</i> .....	1262	<i>parryi, Rhamdella</i> .....	153
2630	<i>de Raizero</i> .....	1273	<i>Rhamdia</i> .....	153
2872	<i>Guachinango</i> .....	1264	<i>partitus, Epomacentrus</i> .....	1558
2631	<i>Mareño</i> .....	1252	<i>Pomacentrus</i> .....	1558
2636	<i>Negro</i> .....	1252	<i>Paru</i> .....	1680
2628	<i>Prieto</i> .....	1252	<i>paru, Chetodon</i> .....	1680, 1681
2139, 2140	<i>parvus, Mesoprion</i> .....	1255	<i>Pomacanthus</i> .....	1680
2141	<i>Paricellinus</i> .....	1885	<i>Rhombus</i> .....	965, 2849
2143	<i>hopliticus</i> .....	1886	<i>Stromateus</i> .....	966
2144	<i>thoburni</i> .....	1888	<i>Parupeneus</i> .....	858
2140	<i>parietalis, Collisus</i> .....	217	<i>parva, Lucania</i> .....	665
2139	<i>parif, Lycovora</i> .....	2478	<i>purvleaps, Lycodapus</i> .....	2493
2142	<i>Ophidium</i> .....	2478	<i>parvipinne, Etheostoma</i> .....	1096
	<i>Uronectes</i> .....	2478	<i>parvipinnis, Archosion</i> .....	1399

	Page.		Page.
parvipinnis, <i>Cestrens</i> .....	1410	pectoralis, <i>Dalla</i> .....	621
<i>Cynoscion</i> .....	1410	<i>Harpe</i> .....	1582
<i>Dierotus</i> .....	883	<i>Maerurus</i> ( <i>Malacocephalus</i> ) .....	2574
<i>Fundulus</i> .....	640, 2827	<i>Nematistius</i> .....	895
<i>Isopisthus</i> .....	1399	<i>Nematonus</i> .....	2518
<i>Promethichthys</i> .....	883	<i>Pedalion</i> .....	1753
<i>parvula</i> , <i>Clupea</i> .....	426	<i>pedaliota</i> , <i>Bonapartia</i> .....	580
<i>parvus</i> , <i>Cotopsis</i> .....	1945	<i>pedaliotus</i> , <i>Zaphotias</i> .....	2826
<i>passany</i> , <i>Arius</i> .....	124	<i>pediculate</i> Fishes .....	2712
<i>Bagrus</i> .....	124	<i>Pediculati</i> .....	2712
<i>Sciaenichthys</i> .....	124, 2760	<i>pelimacra</i> , <i>Centropomus</i> .....	1119
<i>Tachysurus</i> .....	124	<i>Pega</i> .....	2269
<i>passer</i> , <i>Holacanthus</i> .....	1682	<i>Pegador</i> .....	2269
<i>Pomacanthus</i> .....	1683	<i>Pego Oja</i> .....	2699
<i>Pastinaca</i> .....	82	<i>Pegedictis</i> .....	1941, 1942, 1944
<i>Pastor</i> .....	949	<i>ictalops</i> .....	1951
<i>Patao</i> .....	1378	<i>Peixe Agulha</i> .....	711
<i>putao</i> , <i>Gerres</i> .....	1378	<i>Peixe-fonda</i> .....	1312
<i>patatus</i> , <i>Julis</i> .....	1591	<i>Peixe Rey</i> .....	806
<i>patris</i> , <i>Acanthocottus</i> .....	2009	<i>pelada</i> , <i>Anchoa</i> .....	436
<i>patronus</i> , <i>Brevoortia tyrannus</i> .....	434	<i>pelagicum</i> , <i>Siphostoma</i> .....	767
<i>patruelis</i> , <i>Gambusia</i> .....	682	<i>pelagicus</i> , <i>Callionymus</i> .....	2184
<i>Heterandria</i> .....	681	<i>Scomber</i> .....	952
<i>paucidens</i> , <i>Leurynnis</i> .....	2460	<i>Syngnathus</i> .....	770
<i>Lycodopsis</i> .....	2460	<i>pelamides</i> , <i>Scomber</i> .....	869
<i>Oncorhynchus</i> .....	483	<i>pelamis</i> , <i>Gymnosarda</i> .....	868
<i>Salmo</i> .....	483	<i>Scomber</i> .....	869
<i>pauciperis</i> , <i>Ophichthys</i> .....	386	<i>pelamitus</i> , <i>Scomber</i> .....	872
<i>pauciradiatus</i> , <i>Callionymus</i> .....	2168	<i>Pelamys</i> .....	871
<i>paucispinis</i> , <i>Aneyledon</i> .....	1399	<i>chilensis</i> .....	873
<i>Sebastes</i> .....	1781	<i>hneolata</i> .....	873
<i>Sebastes</i> .....	1780	<i>orientalis</i> .....	873
<i>pavonacens</i> , <i>Heros</i> .....	1538	<i>sarda</i> .....	872
<i>pavonia</i> , <i>Limia</i> .....	632	<i>pelamys</i> , <i>Ethynnus</i> .....	869
<i>Pucilia</i> .....	692	<i>Oreynus</i> .....	869
<i>pavoninus</i> , <i>Cyclopterus</i> .....	2097	<i>Scomber</i> .....	872
<i>Paw</i> , <i>John</i> .....	1159	<i>Thynnus</i> .....	869
<i>paxilloides</i> , <i>Lycodes</i> .....	2471	<i>pelegrinus</i> , <i>Squalus</i> .....	51
<i>paxillus</i> , <i>Lycenchelys</i> .....	2471	<i>Pel6rin</i> .....	51
<i>Lycodes</i> .....	2471	<i>Pellona</i> .....	435
<i>Peacock Flounder</i> .....	2665	<i>bleekeriana</i> .....	436
<i>Pea-lip Sucker</i> .....	199	<i>castelneana</i> .....	436
<i>Pearl-fish</i> .....	2495	<i>flavipinnis</i> .....	436
<i>Pearl-fishes</i> .....	2494	<i>furthi</i> .....	436
<i>Pêche-pêche</i> .....	338	<i>orbignyana</i> .....	436
<i>Pêche-Prêtre</i> .....	1784	<i>panamensis</i> .....	436
<i>peckianus</i> , <i>Syngnathus</i> .....	771	<i>pellucida</i> , <i>Ammocrypta</i> .....	1062
<i>peckil</i> , <i>Syngnathus</i> .....	770	<i>clara</i> .....	1063
<i>pectinatae</i> , <i>Centropomus</i> .....	1122	<i>vivax</i> .....	1063
<i>Pristis</i> .....	60, 61, 2749	<i>Salmoperca</i> .....	784
<i>pectinifer</i> , <i>Clinus</i> .....	2362	<i>Tbyris</i> .....	2691
<i>Labrosomus</i> .....	2362	<i>pellucidum</i> , <i>Etheostoma clarum</i> .....	1063
<i>pectoralis</i> , <i>Albūtrossia</i> .....	2573	<i>Ophidium</i> .....	354
<i>Bodianus</i> .....	1582	<i>pellucidus</i> , <i>Delothyris</i> .....	2691
<i>Cossyphus</i> .....	1582	<i>Osphyolax</i> .....	775
<i>Dactyloscopus</i> .....	2301		

Page.		Page.		Page.
621		1063	<i>pellucidus</i> , <i>Pleurolepis</i> .....	2841
1582		350	<i>Paenes</i> .....	1024
	<i>pha-</i>	142	<i>Pelodichthys</i> .....	1208
2574		143	<i>olivaris</i> .....	1200
895		824	<i>Pelon</i> , <i>Guaguancho</i> .....	1460
2518		2660	<i>Peloria</i> .....	1009, 1296
1753		1671	<i>pelta</i> , <i>Chelmo</i> .....	1154
580		1003	<i>peltastes</i> , <i>Lepomis</i> .....	1024
2826		1034	<i>pellata</i> , <i>Percina</i> .....	1024, 2841
2712		1034	<i>peltatum</i> , <i>Etheostoma</i> .....	1142, 1164
2712		1034	<i>peltatus</i> , <i>Hadropterus</i> .....	1135
1119		1373	<i>Turdus cinereus</i> .....	1923
2269		125	<i>pemecus</i> , <i>Bagrus</i> .....	1482
2269		976	<i>Pempheridae</i> .....	947
2699		977	<i>Pempheris</i> .....	1119
1941, 1942, 1944		978	<i>mexicanus</i> .....	1153
1951		978	<i>mulleri</i> .....	1761
711		979	<i>poeyi</i> .....	1303
1312		978	<i>schomburgki</i> .....	1295
806		379	<i>peninsula</i> , <i>Bascanichthys</i> .....	1259
436		379	<i>Callechelys</i> .....	1146
767		797	<i>Chirostoma</i> .....	1388
2184		797	<i>Menidia</i> .....	1172
952		1354, 1355	<i>penna</i> , <i>Calamus</i> .....	1303
770		1355	<i>Pagellus</i> .....	1057
869		577	<i>pennanti</i> , <i>Argentina</i> .....	1133
868		577	<i>Maurolicus</i> .....	1133
869		49	<i>Squalus</i> .....	1133
872		1351	<i>pennatula</i> , <i>Calamus</i> .....	1135
871		2520	<i>Penopus</i> .....	963
873		2521	<i>macdonaldi</i> .....	1021
873		431	<i>pensacola</i> , <i>Harengula</i> .....	1324
872		508, 2819	<i>penshinensis</i> , <i>Salmo</i> .....	1761
869		8 '9	<i>pentacanthus</i> , <i>Bodianus</i> .....	1024
869		990	<i>Centrarchus</i> .....	1454
872		849	<i>Holocentrus</i> .....	1202
869		1576	<i>Labrus</i> .....	1433
51		2081	<i>Xenochirus</i> .....	1107
51		828	<i>Pentanemus</i> .....	1154
435		828	<i>quinquarius</i> .....	1133
436		1022	<i>pepinus</i> , <i>Lucioperca</i> .....	849
436		965	<i>Peprilus</i> .....	1021
436		2643	<i>perarcuatus</i> , <i>Pleuronectes</i> .....	947, 1388
436		1024	<i>Perca</i> .....	1133
436		1130	<i>aberrans</i> .....	1388
436		1024	<i>acuta</i> .....	1133
436		1833	<i>afra</i> .....	1024
1062		1475	<i>albunus</i> .....	1153
1063		1024, 1145	<i>americana</i> .....	1311
1063		1259	<i>apoda</i> .....	1153
784		849	<i>ascensionis</i> .....	1202
2691		1200	<i>atralia</i> .....	1462
1063		1132	<i>chrysops</i> .....	1192
354		1339	<i>chrysoptera</i> .....	1360
2691		1833	<i>dorso nonapterygia</i> .....	1200
775		1023	<i>flavescens</i> .....	1784, 1796
			<i>Perca fluviatilis</i> .....	
			<i>flavescens</i> .....	
			<i>formosa</i> .....	
			<i>furca</i> .....	
			<i>furcraea</i> .....	
			<i>gibbosa</i> .....	
			<i>gigas</i> .....	
			<i>gracilis</i> .....	
			<i>granulata</i> .....	
			<i>guttata</i> .....	
			<i>immaculata</i> .....	
			<i>juba</i> .....	
			<i>lanceolata</i> .....	
			<i>lophar</i> .....	
			<i>longina</i> .....	
			<i>maculata</i> .....	
			<i>marina</i> .....	
			<i>cauda nigra</i> .....	
			<i>gibbosa</i> .....	
			<i>pinnis</i> .....	
			<i>puncticulata</i> .....	
			<i>sectatrix</i> .....	
			<i>venenosa</i> .....	
			<i>melanurum</i> .....	
			<i>minima</i> .....	
			<i>mitchilli</i> .....	
			<i>alternata</i> .....	
			<i>interrupta</i> .....	
			<i>mucronata</i> .....	
			<i>niger</i> .....	
			( <i>Pomacauopsis</i> ) <i>nigropunctata</i> .....	
			<i>nobilis</i> .....	
			<i>norwegica</i> .....	
			<i>notata</i> .....	
			<i>ocellata</i> .....	
			<i>philadelphia</i> .....	
			<i>punctata</i> .....	
			<i>pusilla</i> .....	
			<i>robusta</i> .....	
			<i>rock-fish</i> .....	
			<i>rufa</i> .....	
			<i>salmonea</i> .....	
			<i>saltatrix</i> .....	
			<i>saxatilis</i> .....	
			<i>sectatrix</i> .....	
			<i>septentrionalis</i> .....	
			<i>serratoannulata</i> .....	
			<i>stello</i> .....	
			<i>striata</i> .....	
			<i>tota maculis</i> .....	
			<i>trifurca</i> .....	
			<i>undulata</i> .....	
			<i>unicolor</i> .....	
			<i>unimaculata</i> .....	
			<i>varia</i> .....	
			<i>variabilis</i> .....	

	Page.		Page.
<i>Percia vitrea</i> .....	1021	<i>perfasciatus, Engraulis</i> .....	442
<i>percellens, Raja</i> .....	63	<i>Stolephorus</i> .....	441, 445
<i>Rhinobatus</i> .....	63	<i>Perichthys godeffroyi</i> .....	1197
<i>Percesoces</i> .....	781, 787	<i>periscopus, Gadus</i> .....	2539
<i>Perch, American</i> .....	1023	<i>perisii, Salmo</i> .....	509
Black .....	1504	<i>Perissias</i> .....	2667
Blue .....	1505, 1577	<i>taeniopterus</i> .....	2667
Pike .....	1021	<i>Peristediidae</i> .....	2177
Pirate .....	785, 786	<i>Peristedion</i> .....	2178
Raccoon .....	1023	<i>gracile</i> .....	2179
Ringed .....	1023	<i>Imberbe</i> .....	2182
River .....	1023	<i>longispathum</i> .....	2178
Sacramento .....	991	<i>miconemus</i> .....	2182
Trout .....	782, 784	<i>miniatum</i> .....	2178
Viviparous .....	1498	<i>platycephalum</i> .....	2189
White .....	1133, 1134, 1484, 1501, 1509	<i>Peristethus</i> .....	2178
Yellow .....	1023	<i>miconema</i> .....	2182
Perch-like fishes .....	979	<i>peristethus, Podothecus</i> .....	2062
<i>Perches</i> .....	1015	<i>Perkinsia</i> .....	420
<i>Perches, American Pike</i> .....	1020	<i>othonops</i> .....	420
Log .....	1024	<i>perlatum, Holoцентrum</i> .....	853
<i>Percidae</i> .....	1015	<i>perlongus, Neoconger</i> .....	363
<i>Percidinae</i> .....	2032	<i>Permit</i> .....	943
<i>perciformis, Coryphæna</i> .....	964	<i>perniger, Cullius</i> .....	2201
<i>Lirus</i> .....	964	<i>Electris</i> .....	2201
<i>Palinurichthys</i> .....	964	<i>peroni, Caranx</i> .....	923
<i>Palinurus</i> .....	964	<i>peronii, Lepidopus</i> .....	867
<i>Pammelas</i> .....	964	<i>Peropus</i> .....	2018
<i>Percina</i> .....	1024, 1026	<i>bilobus</i> .....	2018
<i>aspro</i> .....	1033	<i>perplexus, Cottus</i> .....	1955
<i>bimaculata</i> .....	1027	<i>Perrio</i> .....	1659
<i>caprodes</i> .....	1026	<i>perrio, Pseudoscarnus</i> .....	1659
<i>manitou</i> .....	1028	<i>Scarus</i> .....	1659
<i>zebra</i> .....	1027	<i>Perro Colorado</i> .....	1583
<i>guentheri</i> .....	1034	<i>Perro</i> .....	1579
<i>macrocephalus</i> .....	1031	<i>perroteti, Pristis</i> .....	60, 2749
<i>nebulosa</i> .....	1027	<i>personatus, Anmodytes</i> .....	833
<i>peltata</i> .....	1034	<i>perspicabilis, Embiotoca</i> .....	1506
<i>phoxocephala</i> .....	1031	<i>perspicillum, Lycodes</i> .....	2465
<i>rex</i> .....	1025	<i>perspicuus, Hybognathus</i> .....	218
<i>roanoka</i> .....	1036	<i>pertheatus, Stolephorus</i> .....	442
<i>Percine</i> .....	1018	<i>peruanus, Amblyopus</i> .....	2265
<i>Percis</i> .....	2033	<i>Anthias</i> .....	1223
<i>japonicus</i> .....	2034	<i>Gobioides</i> .....	2264
<i>percobromus, Alburnellus</i> .....	295	<i>Hemianthias</i> .....	1222
<i>Minnilus</i> .....	295	<i>Promotogrammus</i> .....	1223
<i>Percoidea</i> .....	979, 1241	<i>peruvianus, Galeichthys</i> .....	122, 2771
<i>Percoidei</i> .....	781	<i>Gerres</i> .....	1376
<i>percoides, Agonostomus</i> .....	819	<i>Tachysurus</i> .....	122
<i>Percoids, Spariform</i> .....	1241	<i>Pesca Blanca</i> .....	321
<i>Percopside</i> .....	783	<i>Verniglia</i> .....	1811
<i>Percopsis</i> .....	783	<i>Pescadillo del Red</i> .....	1416
<i>guttatus</i> .....	784	<i>Pescadillos del Rey</i> .....	807
<i>lanumondi</i> .....	784	<i>Pescadito</i> .....	231
<i>perezi, Carecharlinus</i> .....	36	<i>Pescado Azul</i> .....	1553
<i>Platypodon</i> .....	36	<i>Azul de dos Colores</i> .....	1557



Page.  
 442  
 441, 445  
 1197  
 2536  
 509  
 2667  
 2667  
 2177  
 2178  
 2179  
 2182  
 2178  
 2182  
 2178  
 2180  
 2178  
 2182  
 2062  
 420  
 420  
 853  
 363  
 943  
 2201  
 2201  
 923  
 887  
 2018  
 2018  
 1955  
 1659  
 1659  
 1659  
 1583  
 1579  
 60, 2749  
 833  
 1506  
 2465  
 218  
 442  
 2265  
 1223  
 2264  
 1222  
 1223  
 122, 2771  
 1376  
 122  
 321  
 1811  
 1416  
 807  
 233  
 1553  
 1557

	Page.
Pescalo Blanco de Chapala .....	792
Colorado .....	1453
del Rey .....	806
Pescador .....	2722
Martin .....	2724
Pescados Azules .....	1549
Blancos .....	792
Pesce Re .....	806
Tondo .....	48
petenense, Dorosoma .....	417
petenensis, Chatoessus .....	417
Mollienista .....	700
Pimelodus .....	153
Pocilia .....	694
Rhamdia .....	153
Tetragonopterus .....	335
Petenla .....	1513
splendida .....	1513
petersi, Paralenchurus .....	1481
petimba, Fistularia .....	758
Petimbubaba .....	757
Petite Gueule .....	1370
Jaquette .....	1559
Nègre .....	1142
Scie .....	1323
Petos .....	876
Petrometopon .....	1140
apiarius .....	1142
cruentatus .....	1141
coronatus .....	1142
guttatus .....	1142
panamensis .....	1141
Petromyzon .....	9
americanus .....	10
appendix .....	10
argenteus .....	11
astori .....	12
ayresi .....	13
hairlii .....	9
bdellum .....	11
borealis .....	13
branchialis .....	14, 2745
cautschatiensis .....	2745
castaneus .....	11
ciliatus .....	12
concolor .....	11
lamotteni .....	10
Hvidus .....	12
marius .....	10
chauscha tensis .....	2745
dorsatus .....	10
unicolor .....	10
nigricans .....	10
nigrum .....	14
plumbens .....	13

	Page.
Petromyzon tridentatus .....	12
Petromyzonidae .....	8
Petronason .....	1642
petropauli, Blenniophidium .....	2430
petrosus, Mugil .....	284, 814
Trisotropis .....	1172
petus, Acanthocybium .....	877
Cybium .....	877
Pez Ciego .....	2501
Pez de Espada .....	2749
Gallo .....	895
Pluma .....	1347, 1349, 1350
del Rey .....	769, 808
Luna .....	1753
Puerco .....	1700, 1704
Sierra .....	60
pfeifferi, Murana .....	2805
Sidera .....	2805
phaenna, Hybopsis .....	270
Phaenodon .....	586
ringens .....	586
phaeton, Pristigaster .....	438
phalena, Umbrina .....	1475
Phalangistes .....	2064, 2864
acipenserinus .....	2062
fusiformis .....	2048
japonicus .....	2036
levigatus .....	2048
loricatus .....	2046
phaleratus, Esox .....	628
Phaneron .....	1506
atripes .....	1507
fureatus .....	1506
lateralis .....	1506
Pharyngognathi .....	781, 1571
phasganorus, Notacanthus .....	616
phasma, Careproctus .....	2132
Phenacobius .....	302
cutostomus .....	304
mirabilis .....	303
scopifer .....	303
scopiferus .....	303
teretulus .....	303
hosternus .....	303
aranops .....	304
phenacobius, Notropis .....	263
phenax, Apomotis .....	997
Lepomis .....	997
Myeteroperca falcata .....	1185
philadelphia, Perca .....	1202
philadelphicus, Centropistes .....	1201
Serranus .....	1202
phillipi, Clinus .....	2359
philonips, Cottus .....	1960
Philosopho .....	1693
Philypnus .....	2194

	Page.		Page.
Philypnus dormitator .....	2195	Photonectinae .....	587
dormitor .....	2194	Phoxinus .....	228, 230, 240
lateralis .....	2195	clevelandi .....	237
phlebotomus, Acanthurus .....	1692	flammeus .....	242
phlegethontis, Clinostomus .....	243	margaritus .....	241
Gila .....	243	milnerianus .....	242
Leuciscus .....	243	neogaus .....	241
Phoxinus .....	243	oreutti .....	242
phlox, Boleosoma .....	1052	phlegethontis .....	243
Ulocentra .....	1052	phoxocephala, Percina .....	1031
Phobetor .....	2000	phoxocephalum, Etheostoma .....	1031
tricuspis .....	2000	phoxocephalus, Alvordius .....	1031
Phoebe .....	1211	Cestreus .....	1414
phaebe, Centropristis .....	1212	Cynoscion .....	1413
Prionodes .....	1211	Hadropterus .....	1030
Serranus .....	1212	phrygiatus, Arius .....	131
Pholidapus .....	2430	Hexanematchthys .....	130
dybowskii .....	2430	Tachisurus rugispinis .....	131
grobniiskii .....	2431	Phrynotitan .....	2853
Pholidichthys .....	2347	Phtheichthys .....	2268
anguilliformis .....	2405	lineatus .....	2268
Pholidinae .....	2348	Phycinae .....	2532
Pholis .....	2377, 2414, 2415, 2417	Phycis .....	2552
carolinus .....	2379	americanus .....	2555
dolicogaster .....	2416	chesteri .....	2556
fasciatus .....	2417	chuss .....	2555
gunnellus .....	2419	cirratus .....	2554
novemlineatus .....	2393	dekayi .....	2555
ornatus .....	2410	earlli .....	2555
pictus .....	2415, 2416	floridanus .....	2554
quadrifasciatus .....	2392, 2394	marginatus .....	2555
ruberrimus .....	2417	punctatus .....	2553
subbifurcata .....	2440	rogalis .....	2553
taczanowskii .....	2416	regius .....	2553
Photogenis .....	254	rostratus .....	2555
arionmus .....	290	tenuis .....	2555
caerulons .....	277	Physiculus .....	2547
callistius .....	276	fulvus .....	2547
engraullus .....	296	japonicus .....	2549
eurystomus .....	277	kaupi .....	2547
grandipinnis .....	280	nematopus .....	2548
leuciodus .....	291	rastrolliger .....	2549
leucops .....	296	physignathus, Ceratichthys .....	326
loucopus .....	277	Conesius .....	326
niveus .....	278	Platygobio .....	324
pyrrhomelas .....	281	Physogaster .....	1727
scabriceps .....	290	Piabucina .....	332
spilopterus .....	279	panamensis .....	332
stigmaturus .....	275	Plearels .....	1364
telescopus .....	202	picarti, Hemirhamphus .....	720
photogenis, Leuciscus .....	296	Piceonou .....	194
Notropis .....	295, 296	piceus, Balistes .....	1711
Squalus .....	296	Melichthys .....	1711
Photonectes .....	591	pichardi, Joturus .....	821
gracilis .....	591	Pickereel .....	628
		Banded .....	626

Page.		Page.		Page.
587	Pickereel Common Eastern.....	627	Pike Peroh.....	1021
230, 240	Little.....	627	Pikea.....	1135
237	Pickering.....	1022	Pikes.....	624
242	Picorellus.....	625	Gar.....	108
241	pieta, Muraena.....	2805	pilatus, Prionotus.....	2156
242	pietipinnis, Chelidonichthys.....	2175	Pileoma.....	1024
241	Trigla.....	2176	bimaculata.....	1027
242	picturata, Alutera.....	1719	carbonari.....	1027
243	Gambusia.....	683	cymatogramma.....	1053
1031	Seriola.....	910	nebulosa.....	1027
1031	picturatum, Siphostoma.....	768	semifasciatum.....	1027
1031	picturatus, Curanx.....	910	zebra.....	1028
1414	Gymnothorax.....	395	pilicornis, Blennius.....	2380
1413	Syngnathus.....	768	Pilodletis limosus.....	142
1030	Trachurus.....	909	olivaris.....	143
131	pictus, Centrotonus.....	2416	pilosa, Solea.....	2699
130	Chaunax.....	2726	pilosus, Diodon.....	1744, 1752
131	Chirocentos.....	2717	Trichodiodon.....	1743, 1744
2853	Chirus.....	1873	Pilot, Black.....	1555
2268	Eleotris.....	2201	Cockeye.....	1555, 1561
2268	Gymnelis.....	2477	Pilot-fish.....	465
2532	Gymnothorax.....	2805	Pilot-fishes.....	900
2552	Iridis.....	1599	Pilot, Shark's.....	902
2555	Julls.....	1600	Pimclepterus.....	1384
2556	Lycodontis.....	2805	analogus.....	1386
2555	Oxylebius.....	1878	bosci.....	1388
2554	Pholis.....	2415, 2416	bosquii.....	1388
2555	Platygllossus.....	1600	cornubiensis.....	964
2555	Torpedo.....	78	elegans.....	1387
2554	Urocentrus.....	2416	flavolineatus.....	1386
2555	Picuda.....	823	incisor.....	1386
2553	picuda, Sphyræna.....	823	lutescens.....	1389
2553	Picudilla.....	824	oblongior.....	1388
2553	picudilla, Sphyræna.....	824	ocyrus.....	1390
2555	pidlense, Moxostoma.....	191	Pimelodella.....	158
2555	pidlensis, Ptychostomus.....	191	chagresi.....	154
2547	piger, Symphurus.....	2705	modesta.....	154
2547	Pigfish.....	1338	Pimelodina.....	116
2549	Pigfishes.....	1334	Pimelodus.....	116, 154
2547	pigmentarius, Apogon.....	1109	æneus.....	143
2548	Monoprion.....	1109	affinis.....	134
2549	Pigny Sunfishes.....	981	ailurus.....	140
326	pigra, Aphoristia.....	2706	albidus.....	132, 138
326	Pigus.....	243	antoniensis.....	140
324	Pike, Blue.....	1021	argenteus.....	125
1727	Common.....	628	argentinus.....	135
332	Gar.....	109	argyrus.....	135
332	Gray.....	1022	atrarius.....	140
1364	Great.....	629	baronis-mulleri.....	151
720	Northeru.....	630	blochii.....	155
191	Green.....	627	borealis.....	137
1711	Sacramento.....	224, 2790	brachypterus.....	152
1711	Sand.....	1022	cænosus.....	140
821	Wall-eyed.....	1021	cærulescens.....	135
628	Yellow.....	1021	catulus.....	141
626	Pike-like Fishes.....	622	catus.....	140

	Page.		Page.
<i>Pimelodus caudafurcatus</i> .....	135	<i>Pimelometopon pulcher</i> .....	1585
<i>chagresi</i> .....	154	<i>Pimelonofus</i> .....	149
<i>clarias</i> .....	155	<i>Pimephales</i> .....	216
<i>confinis</i> .....	141	<i>agassizii</i> .....	217
<i>cupreoides</i> .....	140	<i>fasciatus</i> .....	217
<i>cupreus</i> .....	140	<i>maculosus</i> .....	217
<i>dekayi</i> .....	140	<i>nilesi</i> .....	217
<i>erythroptera</i> .....	135	<i>notatus</i> .....	218, 2796
<i>felinus</i> .....	140	<i>promelas</i> .....	217
<i>folis</i> .....	141	<i>confertus</i> .....	217
<i>furcatus</i> .....	134	<i>maculosus</i> .....	217
<i>furcifer</i> .....	135	<i>Pincers</i> .....	431
<i>godmani</i> .....	152	<i>Pinfish</i> .....	1358
<i>gracilis</i> .....	135	<i>pingelii</i> , <i>Triglops</i> .....	1923, 1925
<i>graciosus</i> .....	135	<i>pinguis</i> , <i>Hippoglossus</i> .....	2611
<i>guatemalensis</i> .....	152	<i>Platysomatichthys</i> .....	2611
<i>hammondi</i> .....	135	<i>Pleuronectes</i> .....	2611
<i>houghii</i> .....	135	<i>pini</i> , <i>Trigla</i> .....	2177
<i>hoeyi</i> .....	141	<i>pinima</i> , <i>Acara</i> .....	1323
<i>hypselerus</i> .....	152	<i>Pristipoma acara</i> .....	1323
<i>insigne</i> .....	147	<i>Pink-fish</i> .....	2262
<i>lateralis</i> .....	135	<i>pinnata</i> , <i>Muraena</i> .....	351
<i>latcaudus</i> .....	152	<i>pinnatus</i> , <i>Synphobranchius</i> .....	351
<i>lemniscatus</i> .....	147	<i>pinnifasciatus</i> , <i>Pseudopleuronectes</i> .....	2647
<i>lupus</i> .....	137	<i>pinniger</i> , <i>Bryssetaeres</i> .....	2328
<i>lynx</i> .....	138	<i>Emneacanthus</i> .....	994
<i>macronema</i> .....	155	<i>Gobiosox</i> .....	2329
<i>maculatus</i> .....	135, 155	<i>Sebastichthys</i> .....	1794
<i>managuensis</i> .....	153	<i>Sebastodes</i> .....	1733, 1794
<i>marginatus</i> .....	135	<i>Sebastosomus</i> .....	1794
<i>marmoratus</i> .....	141	<i>pinnimaculatus</i> , <i>Allurichthys</i> .....	117
<i>megalops</i> .....	135	<i>Felichthys</i> .....	117
<i>micropterus</i> .....	151	<i>pinnis</i> , <i>Ireia marina</i> .....	1259
<i>modestus</i> .....	154	<i>Furdus branchialibus</i> .....	1257
<i>motaguensis</i> .....	151	<i>pinnivarius</i> , <i>Hypoplectrus unicolor</i> .....	1192
<i>natalis</i> .....	140	<i>pinnulata</i> , <i>Seriola</i> .....	907
<i>nebulosus</i> .....	140	<i>pinnulatus</i> , <i>Elegatis</i> .....	907
<i>nicaraguensis</i> .....	152	<i>Pinta</i> , <i>Carilla</i> .....	1152
<i>nigrescens</i> .....	137	<i>Morena</i> .....	402
<i>nigricans</i> .....	137	<i>Pintado</i> .....	875
<i>notatus</i> .....	135	<i>Pintano</i> .....	1561
<i>pallidus</i> .....	135	<i>Pintanos</i> .....	1560
<i>petenensis</i> .....	153	<i>pintita</i> , <i>Morena</i> .....	397
<i>platycephalus</i> .....	142	<i>pintiti</i> , <i>Muraena</i> .....	397
<i>polycaulus</i> .....	153	<i>Pipe</i> .....	758
<i>pullus</i> .....	141	<i>Pipefish</i> , <i>Common</i> .....	770
<i>punctulatus</i> .....	143	<i>Great</i> .....	764
<i>rigidus</i> .....	155	<i>Ocean</i> .....	774
<i>salvini</i> .....	152	<i>Pipefishes</i> .....	760
<i>spixii</i> .....	132	<i>Pipor</i> .....	723
<i>vulgaris</i> .....	140	<i>piptolepis</i> , <i>Notropis</i> .....	266
<i>vulpeculus</i> .....	141	<i>Piquier</i> .....	1687
<i>vulpes</i> .....	135	<i>Piquitinga</i> .....	443
<i>wagneri</i> .....	151	<i>piquitinga</i> , <i>Engraulis</i> .....	443
<i>Pimelometopon</i> .....	1585	<i>piquottii</i> , <i>Amia</i> .....	113
<i>darwinii</i> .....	1586	<i>piraaca</i> , <i>Mouacanthus</i> .....	1715

Page.		Page.		Page.
1585	Pirabebé .....	2183	Plagopterus argentissimus .....	329
149	Piracoaba .....	830	Plagusia .....	2704, 2709
216	Piramutana blochii .....	155	brasiliensis .....	2709
217	macrospila .....	155	fasciata .....	2710
217	pirapeda, Dactylopterus .....	2183	plagusa .....	2710
217	Pira-pixanga or Gat-visch .....	1153	tescellata .....	2709
217	Pirate Perches .....	785, 786	plagusia, Pleuronectes .....	2709
118, 2796	Piratiapia .....	1174	Symphurus .....	2709
217	piritita, Cabrilla .....	1181	Plagyodontidae .....	2826
217	pisavarius, Ophichthys .....	377	Plagyodus .....	594, 599, 2826
217	piscatorius, Lophius .....	2713	Pluice .....	2048
431	piscatrix, Pseudorhamdia .....	155	Plain-tail .....	879
1358	Places .....	14, 1241	plana, Platessa .....	2647
923, 1925	pisces, Unicornu bahamensis .....	1710	planci, Mola .....	1750
2611	pisciculus, Esox .....	641	Typanonidium .....	1754
2611	piscis, Luna .....	1754	Plancterus .....	2827, 2828
2611	Piscis viridis bahamensis .....	1638	planiceps, Arius .....	127
2177	pisculentus, Esox .....	641	Catostomus .....	181
1323	Fundulus .....	641	Netuma .....	127, 2766
1323	pisonis, Eleotris .....	2201, 2200	Rhynchobatus .....	64
2262	Gobius .....	2201	planifrons, Eupomacentrus .....	1559
351	Pisodonophis .....	375, 377	Pomacentrus .....	1559
351	eruentifer .....	377, 2803	Planirostra .....	101
2647	daspilotus .....	2803	spatula .....	102
2328	guttnlatus .....	377	planus, Pleuronectes .....	2047
994	longus .....	377	Pseudopleuronectes .....	2647
2329	oculatus .....	376	Plargyrus .....	250, 254
1791	xysturus .....	376	argentatus .....	283
1793, 1794	pisquetus, Caranx .....	921	bowmani .....	283
1791	Paratractus .....	921	melanocephalus .....	217
117	typicus .....	283	plargyrus, Rutilus .....	282
117	Elaphocottus .....	2008	Plate-fish .....	1722
1259	Gymnoctanthus .....	2006, 2008, 2009	Platessa .....	2648
1257	Pitamba, Acara .....	1276	bilineata .....	2643
1192	pituitosus, Rhypticus .....	1234	dentata .....	2615, 2630
907	pixanga, Serranus .....	1153	divinensis .....	2650
907	pixuma, Amoro .....	2201	elongata .....	2657
1152	placitus, Hybognathus .....	213	ferruginea .....	2645
402	Placopharynx .....	197	glabra .....	2650
875	carinatus .....	198	microcephala .....	2654
1501	duquesnii .....	198	oblonga .....	2630
1560	Plagiogrammus .....	2427	ocellaris .....	2630
397	hopkinsi .....	2428	orbignyana .....	2626
397	plagioplateo, Eleotris capito .....	2201	plana .....	2647
758	Plagioscion .....	1418	pola .....	2657
770	heterolepis .....	1419	pusilla .....	2647
764	squamosissimus .....	1418	quadrituberculata .....	2648
774	surinamensis .....	1419	quadrocellata .....	2633
760	Plagusia .....	2704	rostrata .....	2645
723	plagusia, Aphoristia .....	2710	stellata .....	2652
266	Glossichthys .....	2710	platessa, Caranx .....	927
1687	Plagusia .....	2710	platessoides, Citharus .....	2615
443	Pleuronectes .....	2710	Drepanopssetta .....	2615
443	Symphurus .....	2710	Hippoglossoides .....	2614
113	Plagopterinæ .....	204	Pleuronectes .....	2615
1715	Plagopterus .....	320		

	Page.		Page.
Platichthys .....	2551	Platygobio physignathus .....	325
rugosus .....	2652	Platylinus .....	1270, 1280
stellatus .....	2652	vorax .....	1281
umbrosus .....	2643	platyodon, Carcharhinus .....	39
Platirostra .....	101	Squalus .....	30
edentula .....	102	Platypodon .....	33, 34, 35
Platiphrys .....	2660, 2661	falciformis .....	36
constellatus .....	2663	perezii .....	36
ellipticus .....	2665	Platypocellus .....	685
leopardinus .....	2666	maculatus .....	686
lunatus .....	2665	mentalis .....	686
maculifer .....	2664	quitzoensis .....	2873
nubularis .....	2664	platyptogon, Arius .....	127
ocellatus .....	2663	Netuma .....	127, 2767
spinosus .....	2662	Tachisurus .....	127
tenuopterus .....	2668	Platyrhinia exasperata .....	65
platiphrys, Charichthys .....	2683	triseriata .....	66
Platopterus .....	66	platyrrhineus, Lepisosteus .....	111
platorynchus, Aelpenser .....	107	Platyrrhinoideis .....	65
Scaphirhynchus .....	107	triseriatus .....	65, 66
platostomus, Lepisosteus .....	110	platyrrhinus, Aelpenser .....	106
platycephalum, Peristelion .....	2180	platyrrhynchus, Carcharhinus .....	36
Platycephalus .....	2028	Eulamia .....	36
americanus .....	2029	Minomus .....	170
angustus .....	2029	Pantosteus .....	170
dormitator .....	2195	Scaphirhynchus .....	107
platycephalus, Amelurus .....	142	platyrrhynchus, Scaphirhynchops .....	107
Cottus .....	1983, 1988	Platysomatichthys .....	2610
Megalocottus .....	1987	hippoglossoides .....	2611
Noturus .....	144	pinguis .....	2611
Pimelodus .....	142	stomias .....	2610
Platygaster .....	435	Platysomus .....	933
PlatyGLOSSUS bivittatus .....	1597	micropteryx .....	934
candalis .....	1509, 1600	spixii .....	934, 2846
erotaphus .....	1598	Platysqualus .....	43, 44
cyanostigma .....	1591	platystomus, Lepisosteus .....	110
dimidiatus .....	1594	Platytrictes .....	458
dispilus .....	1598	apus .....	458
florealis .....	1597	plebeius, Catostomus .....	171
garnoti .....	1593	Gasterosteus .....	751
grandisquamis .....	1597	Pantosteus .....	171
humeralis .....	1597	plebeius, Mustelus .....	29
internasalis .....	1594	Plecopodus .....	2263, 2868
maculipinna .....	1595	Plectobrachina .....	2349
nicholsi .....	1592	Plectobrachus .....	2431
opalinus .....	1591	evides .....	2432
pictus .....	1600	Plectognathi .....	1696
poeyi .....	1599	Plectognathous Fishes .....	1696
principis .....	1591	Plectospondyli .....	160
radiatus .....	1591, 1597	plectrodon, Perichthys .....	2321
reptus .....	1593	Plectromus .....	840
semicinctus .....	1593	beanii .....	842
Platygobio .....	325	crassiceps .....	843
communis .....	326	cristiceps .....	843
gracilis .....	326	lugubris .....	842
pallidus .....	326	suborbitalis .....	841

Page.	Page.	Page.
325	Plectropoma accensum .....	1193
9, 1280	aflne .....	1193
1281	afrum .....	1166
39	bovinum .....	1193
39	chloroptera .....	1165
34, 35	chlororum .....	1193
36	erocota .....	1192
36	ephippium .....	1192
685	gummigutta .....	1192
686	guttuvarium .....	1192
686	hispanum .....	1140
2873	indigo .....	1193
127	melanorhina .....	1192
27, 2767	monacanthus .....	1165
127	multiguttatus .....	1166
65	nigricans .....	1193
66	puella .....	1192
111	vitulinum .....	1192
65	Plectryps .....	853
65, 66	retrosipilis .....	853
106	pleianus, Pseudoscarus .....	1656
36	Scarus .....	1656
36	pleii, Hemirhamphus .....	723
170	Plesiopercu .....	1028
170	anceps .....	1039
107	Pleurocromylen .....	29
107	pleuriticus, Salmo clarkii .....	2819
2610	mykiss .....	496
2611	Pleurogadus .....	2537
2611	gracilis .....	2538
2610	Pleurogrammus .....	1864
933	monopterygius .....	1864, 1866
934	Pleurolepis .....	1061
34, 2846	asprellus .....	1061
43, 44	pellucidus .....	1063
110	Pleuronectes .....	2648
458	achirus .....	2696
458	americanus .....	2647
171	apoda .....	2701
751	aquosus .....	2600
171	aracaca .....	2672
29	argus .....	2666
263, 2868	asper .....	2645
2349	beani .....	2646
2431	bilineatus .....	2643
2432	cicatricosus .....	2649
1696	cynoglossus .....	2611, 2657
1696	dentatus .....	2630
169	digrammus .....	2641
2321	ellipticus .....	2665
840	elongatus .....	2657
842	ferrugineus .....	2645
843	franklinii .....	2650
843	gilli .....	2654
842	glaber .....	2650
841	glacialis .....	2649
	Pleuronectes guttulatus .....	2640
	hippoglossoides .....	2611
	bippoglossus .....	2612
	ischyrus .....	2641
	kitt .....	2654
	levis .....	2654
	luandoides .....	2615
	lineatus .....	2698, 2701
	linguacula .....	2615
	lanatus .....	2666
	macrolepidotus .....	2672
	maculatus .....	2660
	maculifer .....	2665
	maculosus .....	2626
	melanogaster .....	2630
	microcephalus .....	2654
	ulcerostomus .....	2654
	mollis .....	2701
	nigromanus .....	2657
	oblongus .....	2633
	obscurus .....	2651
	pallasi .....	2648
	papillosum .....	2672
	perareuatus .....	2643
	pinguis .....	2611
	plagiata .....	2710
	plagiata .....	2709
	plaus .....	2647
	platessoides .....	2615
	quadridens .....	2654
	quadrifuberculatus .....	2648
	quenseli .....	2654
	saxicola .....	2657
	stellatus .....	2652
	surinamensis .....	2666
	umbrosus .....	2643
	vetulus .....	2641
	Pleuronectidae .....	2602
	Pleuronectinae .....	2607
	Pleuronichtys .....	2637
	cuenosus .....	2638, 2639
	decurrans .....	2637, 2638
	guttulatus .....	2640
	quadrifuberculatus .....	2638
	verticalis .....	2638
	pleurophthalmus, Antennarius .....	2722
	pleurospilus, Girardinus .....	688
	Heterandria .....	688
	pleurostictus, Triglops .....	1923
	Pleurothyris .....	603
	olfersi .....	604
	plumaris, Archistes .....	1900, 1901
	plumatula, Calamus .....	1352
	plumbea, Chimera .....	95
	Dionda .....	216
	Gambusia .....	695

	Page.		Page.
<i>plumbea</i> , <i>Hybognathus</i> .....	216	<i>Pocilia chisoyensis</i> .....	693, 2834
<i>Lampetra</i> .....	13	<i>couchiana</i> .....	695, 2833
<i>plumbeolus</i> , <i>Alburnops</i> .....	283	<i>conchii</i> .....	695
<i>Minnulus</i> .....	283	<i>cubensis</i> .....	692
<i>plumbeum</i> , <i>Zophendum</i> .....	216	<i>euneata</i> .....	2834
<i>plumbens</i> , <i>Ceratichthys</i> .....	324	<i>dominicensis</i> .....	696, 2833, 2834
<i>Conesius</i> .....	323	<i>dovii</i> .....	695, 2833
<i>Gobio</i> .....	324	<i>elongata</i> .....	697, 2834
<i>Petromyzon</i> .....	13	<i>fasciata</i> .....	641
<i>plumier</i> , <i>Le Tetrodon</i> .....	1733	<i>fasciatus</i> .....	2833
<i>plumieri</i> , <i>Caranx</i> .....	912	<i>gillii</i> .....	692, 2834
<i>Chaetodon</i> .....	1668	<i>lineolata</i> .....	700
<i>Couodon</i> .....	1324	<i>macrolepidota</i> .....	641
<i>Coryphæna</i> .....	2276	<i>melanogaster</i> .....	696, 2834
<i>Diabasis</i> .....	1336	<i>melapleura</i> .....	666
<i>Gerres</i> .....	1379	<i>mexicana</i> .....	692, 2833
<i>Gobius</i> .....	2206	<i>multilineata</i> .....	700
<i>Hæmulon</i> .....	1304	<i>olivacea</i> .....	659
<i>Labrus</i> .....	1305	<i>pavonia</i> .....	692, 2833
<i>Malacanthus</i> .....	2275	<i>petenensis</i> .....	694, 2833
<i>Mugil</i> .....	812, 2841	<i>plumbeus</i> .....	2833
<i>Sciæna</i> .....	1324	<i>presilionis</i> .....	697
<i>Scomber</i> .....	911	<i>reticulata</i> .....	2833
<i>Scoræberonornis</i> .....	875	<i>schneideri</i> .....	691
<i>Scorpena</i> .....	1848	<i>sphenops</i> .....	694, 2833
<i>Sicydium</i> .....	2206	<i>spilurus</i> .....	697, 2833
<i>Tetrodon</i> .....	1733	<i>surinamensis</i> .....	691
<i>Trachurops</i> .....	912	<i>thermalis</i> .....	693, 2833
<i>Trichidion</i> .....	230	<i>(Acropocilia) tridens</i> .....	690
<i>plumierianus</i> , <i>Caranx emornis</i> .....	911	<i>vandepolli</i> .....	696, 2833
<i>plumierii</i> , <i>Polyductylus</i> .....	830	<i>arubensis</i> .....	696, 2834
<i>Polygenus</i> .....	830	<i>vittata</i> .....	692, 2833
<i>plurimis</i> , <i>Cottus cirris</i> .....	2066	<i>vivipara</i> .....	691, 2833
<i>plutonina</i> , <i>Raja</i> .....	69, 70	<i>Pocillichthys</i> .....	1066, 1067, 1069
<i>pluvialis</i> , <i>Labrax</i> .....	2841	<i>artesia</i> .....	1094
<i>Poacher</i> , <i>Sea</i> .....	2091	<i>asprigens</i> .....	1085
<i>Poachers</i> , <i>Sea</i> .....	2031	<i>barratti</i> .....	1102
<i>pocafello</i> , <i>Catostomus</i> .....	175	<i>beani</i> .....	1057
<i>podostemone</i> , <i>Boleosoma</i> .....	1055	<i>borealis</i> .....	1082
<i>Etheostoma</i> .....	1055	<i>budlerianus</i> .....	1102
<i>Podotheus</i> .....	2054	<i>camurus</i> .....	1076
<i>maculifer</i> .....	2055	<i>cæruleus</i> .....	1089
<i>acipenserinus</i> .....	2061, 2062	<i>ees</i> .....	1102
<i>gilberti</i> .....	2058	<i>erochrons</i> .....	1102
<i>hualini</i> .....	2056	<i>exilis</i> .....	1103
<i>peristethus</i> .....	2062	<i>fusiformis</i> .....	1102
<i>sturioides</i> .....	2063	<i>gracilis</i> .....	1103
<i>thompsoni</i> .....	2030	<i>jessho</i> .....	1085
<i>veterans</i> .....	2063, 2064	<i>lateralis</i> .....	1099
<i>vulsus</i> .....	2068	<i>lepidus</i> .....	1089
<i>Pocilia</i> .....	690, 2833	<i>mesæus</i> .....	1059
<i>boncardi</i> .....	695, 2834	<i>palustris</i> .....	1102
<i>braueri</i> .....	2834	<i>punctulatus</i> .....	1091
<i>butleri</i> .....	691, 2833	<i>quifacæus</i> .....	1101
<i>caerulea</i> .....	641	<i>rutilineatus</i> .....	1079
<i>catenata</i> .....	648	<i>sagitta</i> .....	1081



Page.		Page.		Page.
693, 2834	<i>Pocilichthys sanguifluus</i> .....	1077	<i>Pogonichthys inaequilobus</i> .....	224
695, 2833	<i>saxatilis</i> .....	1048	<i>naerolepidotus</i> .....	223
695	<i>spectabilis</i> .....	1089	<i>symmetricus</i> .....	246
692	<i>swaini</i> .....	1086	Pogy .....	433
2834	<i>versicolor</i> .....	1089	Po-he-wa .....	238
893, 2834	<i>virgatus</i> .....	1093	Point Loma, Blind Goby of.....	2262
695, 2833	<i>vitreus</i> .....	1065	Poison Toad-fishes .....	2323, 2325
697, 2834	<i>vulneratus</i> .....	1077	Poisson Bleu .....	517
641	<i>warreni</i> .....	1103	de Marais .....	113
2833	<i>zonalis</i> .....	1075	Lune .....	954
692, 2834	Pociliidae.....	630	pola, Platessa .....	2657
700	Pocilina .....	632	polaris, Blennius .....	2469
641	Pocilioides .....	678	Boreogadus .....	2534
696, 2834	<i>bimaculatus</i> .....	678	Cottus .....	1999
660	Pocilocephalus .....	381	Lycodalepis .....	2468
692, 2833	pocilooides, Lebistes .....	689	Lycodes .....	2469
700	<i>Limia</i> .....	700	Merlangus .....	2534
659	Pocilophis .....	462	Pollachius .....	2534
692, 2833	<i>nocturnus</i> .....	403	Porocottus .....	1998
694, 2833	<i>preilophthalmus</i> , Gobiesox.....	2335	Pole Flounder .....	2657
2833	pocilopus, Myrripristis .....	847	Polistotrema .....	6
697	<i>Rhamphoberyx</i> .....	847	<i>dombey</i> .....	6
2833	Pocilosoma .....	1066	<i>stontii</i> .....	6
691	<i>erythrogastrum</i> .....	1089	<i>politus</i> , Scirphus .....	1397
694, 2833	<i>transversum</i> .....	1089	<i>Splaceroides</i> .....	1736
697, 2833	pocilurn, Myxostoma .....	196	<i>annulatus</i> .....	1736
691	Pocilurichthys .....	333	Tetrodon .....	1736
693, 2833	pocilurum, Moxostoma .....	196	Pollachius .....	2534
690	poculus, Citharichthys .....	2672	<i>carbonarius</i> .....	2535
696, 2833	poyei, Alepidosaurus (Canlopus)...	596	<i>chalcogrammus</i> .....	2536, 2537
696, 2834	Dactyloscopus .....	2302	<i>polaris</i> .....	2534
692, 2833	<i>Engraulis</i> .....	445	<i>virens</i> .....	2534
691, 2833	Gobius .....	2226	Pollaek .....	2534
1067, 1069	<i>Halichares</i> .....	1598	Puget Sound .....	2536
1094	<i>Hemirhamphus</i> .....	720	Wall-eyed .....	2536
1085	Iridio .....	1599	Pollacks .....	2534
1102	<i>Lycengraulis</i> .....	2811	Alaskan .....	2535
1057	Orthopristis .....	1339	<i>pollicaris</i> , Cottus .....	1941, 1953
1082	Peupheris .....	979	Uraniden .....	1954
1102	PlatyGLOSSUS .....	1599	<i>pollux</i> , Pontinus .....	1857
1076	Siphostoma .....	766	<i>poloosoo</i> , Caranx .....	928
1089	Stolephorus .....	445	<i>polyacanthocephalus</i> , Cottus .....	1977
1102	Synodus .....	536	Myoxocephalus .....	1976
1102	Pogge .....	2065	<i>Polyacanthonotina</i> .....	613
1103	Pogonathus .....	1482	<i>polyacetocephalum</i> , Bryostemma..	2408, 2409
1102	<i>corbina</i> .....	1483	<i>polyacetocephalus</i> , Blennius .....	2400
1103	Pogonius .....	1482	<i>Chirolophis</i> .....	2409
1085	<i>corbina</i> .....	1483	<i>polycanulus</i> , Pimelodus .....	153
1099	<i>crocoides</i> .....	1482	<i>Rhamdia</i> .....	153
1089	<i>corbina</i> .....	1483	Polycirrhus .....	1477
1059	<i>fasciatus</i> .....	1483	<i>dumerili</i> .....	1479
1102	Pogonichthys .....	223	<i>rathbuni</i> .....	1479
1091	<i>argyrosus</i> .....	224	Polyclemus .....	1477, 1478
1101	<i>communis</i> .....	326	Polyductylus .....	828
1079	( <i>Platygobio</i> ) <i>gulonellus</i> .....	326	<i>approximans</i> .....	829
1081				

	Page.		Page.
Polydactylus octonemus .....	830	Pomacanthus paru .....	1680, 1681, 2859
opercularis .....	830	passer .....	1683
plumieri .....	830	quinquecinctus .....	1680
virginicus .....	829	tricolor .....	1684
polygonius, Acanthostracion .....	1725	zonipectus .....	1681
Gymnothorax .....	394	Pomacentridæ .....	1543
Lycodontis .....	394	Pomacentrinæ .....	1544
polylepis, Balistes .....	1700	Pomacentrus adustus .....	1552
Polymixia .....	854	analgutta .....	1554
lowei .....	854	nalis .....	1555
Polymixiidae .....	854	atrocyaneus .....	1552
polymorphus, Gadus .....	2540	bairdii .....	1567
Polynemidae .....	827, 2841	caudalis .....	1556, 1557
Polyneumus .....	828	denegatus .....	1567
americanus .....	830	dorsopunicans .....	1557
approximans .....	829	flavilatus .....	1558
artedi .....	828	fuscus .....	1552
californiensis .....	829	leucostictus .....	1556
macronemus .....	828	niveatus .....	1568
mango .....	830	obscuratus .....	1552, 1555
melanopoma .....	831	otophorus .....	1555
octofiliis .....	830	paritus .....	1558
octonemus .....	830	planifrons .....	1559
oligodon .....	830	quadrigutta .....	1570
opercularis .....	831	rectifranum .....	1554
plumieri .....	830	rubicundus .....	1565
quinqarius .....	828	variabilis .....	1552
sexradiatus .....	2183	xanthurus .....	1557
tridigitatus .....	2177	Pomadasis .....	1329
virginicus .....	830	andrei .....	1332
Polyodon .....	101	axillaris .....	1328
feuille .....	102	bayanus .....	1331
folium .....	102	branioki .....	1333
spathula .....	101, 102	corvineformis .....	1327
Polyodontidae .....	101	crocro .....	1333
Polyprion .....	1138	dovii .....	1318
americanus .....	1139	elongatus .....	1328
cernium .....	1139	humilis .....	1331
oxygenius .....	1139	leuciscus .....	1328
Polyprionine .....	1128	macracanthus .....	1332
Polyprosopus .....	51	nitidus .....	1326
macer .....	51	pacifici .....	1316
Polypterichthys .....	754	panamensis .....	1331
polytrema, Bdellostoma .....	6	productus .....	1332
Polyuranodon .....	392	ramosus .....	1334
Pomacampsis .....	1020	Pomadasyx .....	1333
Pomacanthinae .....	1670, 2860	cusius .....	1317
Pomacanthodes .....	1679, 1681	davidsoni .....	1321
zonipectus .....	1682	leuciscus .....	1328
Pomacanthus .....	1679, 2859	modestus .....	1321
arcuatus .....	1679, 1680, 1681	virginicus .....	1323
aureus .....	1680	Pomatapriion .....	1565
balteatus .....	1680	dorsalis .....	1570
ciliaris .....	1685, 1686	Pomatomichthys .....	1111
cingulatus .....	1680	Pomatomidae .....	945
crescentalis .....	1682	Pomatomus .....	946, 1111

Page.		Page.
31, 2859	<i>Pomatomus saltator</i> .....	947
1683	<i>saltatrix</i> .....	946
1680	<i>skib</i> .....	947
1684	<i>Pomatopsetta</i> .....	2614
1681	<i>dentata</i> .....	2615
1543	<i>Pomatoschistus</i> .....	2210
1544	<i>Pomfrets</i> .....	956, 958, 959
1552	<i>Pomolobus</i> .....	424
1554	<i>estivalis</i> .....	426
1555	<i>chrysochloris</i> .....	425
1552	<i>medicrils</i> .....	425, 2810
1567	<i>pseudoharongus</i> .....	426
556, 1557	<i>lacus-</i>	
1567	<i>trisa</i> .....	426
1557	<i>vernalis</i> .....	426
1558	<i>Pomotis</i> .....	999, 1006
1552	<i>bombifrons</i> .....	1003
1556	<i>brevicaps</i> .....	1003
1568	<i>catesbei</i> .....	1010
1552, 1555	<i>chaetodon</i> .....	995
1555	<i>convexifrons</i> .....	1003
1558	<i>elongatus</i> .....	1001
1559	<i>fallax</i> .....	1003
1570	<i>gibbosus</i> .....	1005
1554	<i>gulosus</i> .....	992
1565	<i>guttatus</i> .....	993
1552	<i>heros</i> .....	1007
1557	<i>holbrookii</i> .....	1008
1329	<i>incisor</i> .....	1005
1332	<i>inscriptus</i> .....	1003
1328	<i>longulus</i> .....	996
1331	<i>luna</i> .....	1006
1333	<i>marginatus</i> .....	1003
1327	<i>microlophus</i> .....	1008
1333	<i>nefastus</i> .....	1003
1318	<i>nitida</i> .....	1003
1328	<i>notatus</i> .....	1008
1331	<i>obesus</i> .....	993
1328	<i>obscurus</i> .....	1006
1332	<i>pallidus</i> .....	1007
1326	<i>popeii</i> .....	1003
1316	<i>ravenelii</i> .....	1010
1331	<i>rubicauda</i> .....	1001
1332	<i>sanguinolentus</i> .....	1003
1334	<i>solis</i> .....	1001
1333	<i>speciosus</i> .....	1000, 1008
1317	<i>vulgaris</i> .....	1010
1321	<i>pomotis</i> , <i>Acantharechus</i> .....	989
1328	<i>Ambloplites</i> .....	989
1321	<i>Centrarechus</i> .....	989
1323	<i>Pomoxis</i> .....	986
1565	<i>annularis</i> .....	987
1570	<i>sparoides</i> .....	987
1111	<i>Pomoxys</i> .....	986
945	<i>brevicauda</i> .....	987
946, 1111	<i>intermedius</i> .....	987

	Page.
<i>Pomoxys protacanthus</i> .....	987
<i>sparoides</i> .....	988
<i>Pompano, California</i> .....	967
Common.....	944
Irish.....	1376
<i>Pomphilus</i> .....	900
<i>Pompilus</i> .....	962
<i>pompilus, Centrolophus</i> .....	963
<i>Thynnus</i> .....	900
<i>Pompon</i> .....	1318
<i>Ponco Prieto</i> .....	1297
<i>Pond Smelt</i> .....	525
<i>ponderosus, Amiurus</i> .....	137
<i>Ictalurus</i> .....	137
<i>pondiceriana, Elacate</i> .....	948
<i>ponticus, Gasterosteus</i> .....	747
<i>Hippoglossus</i> .....	2612
<i>Poatinus</i> .....	1854
<i>castor</i> .....	1856
<i>longispinis</i> .....	1858
<i>macrolepis</i> .....	1855
<i>pollux</i> .....	1857
<i>rathbuni</i> .....	1857
<i>sierra</i> .....	1859
<i>popeli, Pomotis</i> .....	1003
<i>Pop-eye</i> .....	2586
<i>Porbeagles</i> .....	49
<i>porca, Scorpena</i> .....	1839
<i>Porcupine-fish</i> .....	1742, 1744
<i>Porgeo</i> .....	1509
<i>Porgies</i> .....	1343
<i>F water</i> .....	1344
.....	1356
<i>Porgy</i> .....	1346
<i>Grass</i> .....	1355
<i>Jolt-head</i> .....	1352
<i>Little-head</i> .....	1350
<i>Little-mouth</i> .....	1354
<i>Red</i> .....	1356
<i>Saucer-eye</i> .....	1349
<i>Shad</i> .....	1355
<i>Sheepshead</i> .....	1354
<i>Southern</i> .....	1349
<i>White-bone</i> .....	1353
<i>Porichthys</i> .....	2317
<i>margaritatus</i> .....	2322
<i>nautopædium</i> .....	2323
<i>notatus</i> .....	2321
<i>plectrodon</i> .....	2321
<i>porosissimus</i> .....	2310, 2321
<i>porifer, Lycopohelys</i> .....	2471
<i>Lycodes</i> .....	2472
<i>Porkfish</i> .....	1322
<i>Porobronchus</i> .....	2495
<i>Poroclinus</i> .....	2433
<i>rothrocki</i> .....	2434

	Page.		Page.
Porocottus .....	1996	Priacanthus .....	1237
bradfordi .....	2862	altus .....	1240
polaris .....	1998	arenatus .....	1237, 1238
quadratus .....	1998	carolinus .....	2858
quadrifilis .....	1999, 2000, 2863	catalufa .....	1238
sellaris .....	1996, 2863	cepodianus .....	1238
tentaculatus .....	2000, 2862	cruentatus .....	1238, 2858
Poroderma .....	23	fulgens .....	1238
Porogadus .....	2519	macrophthalmus .....	1238
miles .....	2520	schlegeli .....	2858
promelas .....	2512	serrula .....	1239
Porogobius .....	2210	pribilovius, Nautichthys .....	2020
Poromitra .....	840	Nautiscus .....	2019
capito .....	840	pricei, Campostoma .....	205
Poronotus .....	965, 967, 2849	Villarius .....	2790
simillimus .....	967	Prick Fish .....	555
triacanthus .....	2849	Prickly Bullhead .....	1944
porosissimus, Batrachus .....	2321	Priest Fish .....	1781
Porichthys .....	2319, 2321	Prieta, Aguja .....	891
porosus, Carcharias .....	37	Mojarra .....	1269
Cottus .....	1975	Morena .....	2801
Esox .....	627	Prieto Pargo .....	1252
Porto Enseigne .....	1687	Robalo .....	1119
Portuicus .....	2848	Ronco .....	1297
Portugais .....	1679	prieto, Lutjanus .....	1253
Portuguese Man-of-War Fish .....	949	Prilonotus .....	1741, 1742
Post Croaker .....	1458	(Auchisomus) caudicinc-	
postica, Echoneis .....	2272	tus .....	1742
Potamocottus .....	1942	Prinospinus .....	1765, 1774, 1783
bendirei .....	1965	princeps, Caulolatilus .....	2276, 2277
carolinus .....	1952	Cottus .....	1962
punctatus .....	1949	Lutilus .....	2277
zopherus .....	1952	principis, Antennarius .....	2719
Potomac Shad .....	427	Chironectes .....	2719
pottsii, Apleston .....	1083	Julis .....	1591
Boleosoma .....	1083	PlatyGLOSSUS .....	1591
Etheostoma .....	1082	Prinodon .....	670
pourtalesii, Archosargus .....	1360	Priodonophis .....	392, 393, 399
Sargus .....	1360	meleagris .....	399
Pout, Horned .....	135, 140	ocellatus .....	399
powelli, Balistes .....	1702	Prionace .....	33
praecisus, Clinus .....	2441	glauca .....	33
Eunnesogrammus .....	2441	Prioulstius .....	1927
priestabilis, Alosa .....	428	macellus .....	1928
prestigiator, Centropristis .....	1214	Prionodes .....	1208, 1209, 1210
Serranus .....	1214	aequidens .....	1210
presidionis, Pecilia .....	697	bulleri .....	1213
pretiosa, Argentina .....	525	fasciatus .....	1212
pretiosus, Hypomesus .....	525	flavescens .....	1215
Osmerus .....	525	fuscus .....	1211
Ruvettus .....	879	lucepercanus .....	1216
Thyrstites .....	880	pnebe .....	1211
Trachichthys .....	837	stillbostigma .....	1216
Prétre, Pêche .....	1784	tabacarius .....	1215
Priacanthichthys .....	1148	tigrinus .....	1211
Princanthidae .....	1236	Prionodon cucurli .....	40

Page.		Page.
1237		35
1240		2148, 2150, 2160, 2867
1237, 1238	<i>Prionotus</i>	2150
2858	<i>alatus</i>	2163
1238	<i>albirostris</i>	2170
1238	<i>beanii</i>	2152, 2156
1238, 2858	<i>birostratus</i>	2156, 2157
1238	<i>carolinus</i>	2175
1238	<i>egretta</i>	2107, 2168, 2169
2858	<i>evolans</i>	2153
1239	<i>gymnostethus</i>	2172
2020	<i>borreus</i>	2167
2019	<i>lineatus</i>	2155, 2156
205	<i>loxias</i>	2160
2790	<i>miles</i>	2164
555	<i>ophryas</i>	2157
1944	<i>palmipes</i>	2156
1784	<i>pilatus</i>	2158, 2164, 2169
891	<i>punctatus</i>	2161
1299	<i>quiescens</i>	2158
2804	<i>roseus</i>	2164
1252	<i>rubio</i>	2169
1119	<i>sarritor</i>	2157
1297	<i>scitulus</i>	2166
1253	<i>stearnsi</i>	2161
1741, 1742	<i>stephanophrys</i>	2167
inc-	<i>strigatus</i>	2171, 2172
1742	<i>tribulus</i>	2154
15, 1774, 1783	<i>xenisma</i>	1696
2276, 2277	<i>Prionurus</i>	1695
1962	<i>latielvius</i>	60
2277	<i>punctatus</i>	438
2719	<i>Pristida</i>	438
2719	<i>cayanus</i>	437
1591	<i>dovii</i>	436
1591	<i>flavipinnis</i>	437
670	<i>lutipinnis</i>	437
392, 393, 399	<i>macrops</i>	438
399	<i>martii</i>	438
399	<i>mucronatus</i>	438
33	( <i>Odontognathus</i> )	438
33	<i>panamensis</i>	438
1927	<i>phaeton</i>	418
1928	<i>Pristigasterina</i>	1329, 1331
08, 1209, 1210	<i>Pristipoma</i>	1323
1210	<i>acira pinima</i>	1332
1213	<i>andrei</i>	1321, 1343
1212	<i>auratum</i>	1328
1215	<i>axillare</i>	1320
1211	<i>bicolor</i>	1319
1216	<i>bilineatum</i>	1334
1211	<i>boucardi</i>	1334
1216	<i>brunicki</i>	1320
1215	<i>brasiliense</i>	1341
1211	<i>brevipinne</i>	1340
40	<i>catharinum</i>	1323
	<i>catharine</i>	

	Page.
<i>Pristipoma chalcenum</i>	1338
<i>coro</i>	1324
<i>crocro</i>	1333
<i>cultiferum</i>	1333
<i>davidsonii</i>	1321
<i>dovii</i>	1318
<i>fasciatum</i>	1339
<i>fulvomaculatum</i>	1339
<i>furthi</i>	1319
<i>humile</i>	1331
<i>kneri</i>	1338
<i>leuciscus</i>	1328
<i>maeracanthum</i>	1332
<i>melanopterum</i>	1319
( <i>Hemulopsis</i> )	1326
<i>notatum</i>	1321
<i>panamense</i>	1331
<i>productum</i>	1332
<i>ramosum</i>	1334
<i>rodo</i>	1323
<i>scapulare</i>	1321
<i>serrula</i>	1324, 1343
<i>splendatum</i>	1322
<i>surinamense</i>	1319
<i>trilineatum</i>	1320
<i>virginicum</i>	1323
<i>Pristipomoides</i>	1279
<i>Pristis</i>	60
<i>aenirostris</i>	61
<i>granulosa</i>	61
<i>megulodon</i>	61
<i>mississippiensis</i>	61
<i>oeca</i>	61
<i>pectinatus</i>	60, 61, 2749
<i>perroteti</i>	60, 2740
<i>zephyreus</i>	2740
<i>Pristobatus</i>	60
<i>Pristocantharus</i>	1334
<i>Proach, Lucky</i>	1971
<i>Proamblys</i>	1247
<i>Proarthri</i>	10
<i>probatocephalus, Archosargus</i>	1361, 1362
<i>Diplodus</i>	1361
<i>Sparus</i>	1361
<i>proboscoidalis, Agonomalus</i>	2637
<i>Aspidophorus</i>	2638
<i>proboscidea, Limanda</i>	2645
<i>proboscideus, Chaenomugil</i>	816
<i>Monacanthus</i>	1719
<i>Mugil</i>	816
<i>procellarum, Myctophum</i>	575
<i>procerus, Venefica</i>	365
<i>Procerus</i>	101
<i>vittatus</i>	102
<i>procerum, Nettastoma</i>	366
<i>Procerus maculatus</i>	102

	Page.		Page.
Prochilus .....	2105	Promicropterus .....	1229, 1231, 1233
proene, Cliola .....	264	decoratus .....	1234
Exocoetus (Cypselurus)....	737	Promoxis nitidus .....	987
Hybognathus .....	264	Prouotogrammus .....	1224
Hybopsis .....	264	eos .....	1224
Leuciscus .....	264	multifasciatus ..	1226
Notropis .....	264	peruanus .....	1223
productum, Holocentrum .....	852	vivanus .....	1224
Pristipoma .....	1332	proops, Bagrus .....	124
productus, Alepocephalus .....	452	Notuma .....	124
Cylindrosteus .....	111	Sciadeichthys .....	123, 2760
Engraulis .....	447	Tachisurus .....	124
Eucinostomus .....	1372	Propterygia .....	66
Gadus .....	2531	prorates, Leptophidium .....	2485
Leuciscus .....	240	proridens, Calamus .....	1350
Merlangus .....	2531	proriger, Clinostomus .....	240
Merluccius .....	2531	Leuciscus .....	240
Pomadasis .....	1332	Sebasteichthys .....	1788, 1793
Rhinobatus .....	63	Sebastes .....	1787, 1792
Stolephorus .....	447	Squalus .....	240
proliare, Etheostoma .....	1104	prorigera, Congermurana .....	357
proliaris, Etheostoma .....	1104	prorigerum, Ophisoma .....	357
Microperca .....	1103	proserpina, Moniana .....	272
profundorum, Acanthocottus .....	1991	Notropis .....	272
Leptophidium .....	2484	Prosopium .....	461, 462
Ophidium .....	2484	conesii .....	463
Scyllorhinus .....	22	prospinosum, Holocentrum .....	853
Zesticelus .....	1990	Prospinus .....	1164
profundus, Lutjanus .....	1264	chloropterus .....	1165
Mesoprius .....	1263	prosthemius, Ceratichthys .....	324
Prognathodes .....	1071	Cousius .....	324
aculeatus .....	1071	prosthistsius, Amiurus .....	139
prognathus, Argyrosomus .....	471	protacanthus, Pomoxys .....	987
Coregonus .....	472	protens, Oncorhynchus .....	478
Prognurus .....	2866	Salmo .....	478
cypselurus .....	2866	protoclus, Myctophum .....	565
prolixum, Campostoma .....	206	Protoporus .....	228
prolixus, Leuciscus .....	206	dominus .....	233
prolongus, Leptoconger .....	363	proxima, Seriola .....	904
promelas, Mœbia .....	2511	proximus, Gadus .....	2539
Pimephales .....	217	Microgadus .....	2539
confortus .....	217	pruinosis, Gadus .....	2540
maculosus .....	217	Pseudoblennius .....	2406
Porogadus .....	2512	hypacanthus .....	2406
Prometheus .....	882	Psaenes .....	950
Prometheus atlanticus .....	883	auratus .....	951
prometheus, Gempylus .....	883	cyanophrys .....	950
Promethichthys .....	882	fuscus .....	951
Promethichthys .....	882	javanicus .....	951
atlanticus .....	883	maculatus .....	951
parvipinnis .....	883	pellucidus .....	950
prometheus .....	882	regulus .....	951
Promicrops .....	1162	Psetticthys .....	2617
guasa .....	1164	melanostictus .....	2618
guttatus .....	1162	sordidus .....	2680
italara .....	1164	Psettiue .....	2608

Page.		Page.		Page.
231, 1233	<i>Pseudariodes</i> .....	154	<i>Pseudoscarus</i> <i>lincolatus</i> .....	1651
1234	<i>pamtherinus</i> .....	155	<i>microrrhinus</i> .....	1655
987	<i>Pseudarius</i> .....	119	<i>nuchalis</i> .....	1654
1224	<i>Pseudobastes</i> .....	1839	<i>obtusus</i> .....	1654
1224	<i>Pseudocanthicus</i> .....	159	<i>perrieo</i> .....	1659
1226	<i>pseudocrocodillus</i> , <i>Scopelus</i> .....	556	<i>pleianus</i> .....	1656
1223	<i>pseudogula</i> , <i>Eucinostomus</i> .....	1363	<i>punctulatus</i> .....	1646
1224	<i>Gerres</i> .....	1368	<i>psittacus</i> .....	1647
124	<i>pseudoharengus</i> , <i>Clupea</i> .....	426	<i>quadrispinosus</i> .....	1648
124	<i>Pomolobus</i> .....	426	<i>rostratus</i> .....	1658
123, 2760	<i>lacustris</i> .....	426	<i>sanctae-crucis</i> .....	1651
124	<i>Pseudohemiodon</i> .....	156	<i>simplex</i> .....	1656
66	<i>pseudohispanica</i> , <i>Clupea</i> .....	424	<i>superbus</i> .....	1650
2485	<i>Sardinia</i> .....	424	<i>taniopterus</i> .....	1646, 1647
1350	<i>pseudohispanicus</i> , <i>Clupanodon</i> .....	423	<i>trispinosus</i> .....	1648
240	<i>Pseudojulis</i> .....	1604	<i>turchesius</i> .....	1659
240	<i>adustus</i> .....	1603	<i>vetula</i> .....	1650
1788, 1793	<i>californicus</i> .....	1601	<i>Pseudosciena</i> <i>surinamensis</i> .....	1420
1787, 1792	<i>inornatus</i> .....	1604	<i>Pseudoscopelus</i> .....	2292
240	<i>melanotis</i> .....	1605	<i>scriptus</i> .....	2292
357	<i>modestus</i> .....	1601	<i>Pseudotriakide</i> .....	26
357	<i>notospilus</i> .....	1603	<i>Pseudotriakis</i> .....	27
272	<i>venustus</i> .....	1602	<i>microdon</i> .....	27
272	<i>Pseudoloricaria</i> .....	156	<i>Pseudoxiphophorus</i> .....	678
461, 462	<i>Pseudomonacanthus</i> .....	1717	<i>bimaculatus</i> .....	678
463	<i>amphioxys</i> .....	1717	<i>reticulatus</i> .....	678
853	<i>Pseudomuraena</i> .....	392	<i>Pseudupeneus</i> .....	858
1164	<i>Pseudophoxinus</i> .....	243	<i>Pylonotus</i> .....	1741
1165	<i>Pseudopleuronectes</i> .....	2646	<i>punctatissimus</i> .....	1741
324	<i>americanus</i> .....	2647	<i>psittacinus</i> , <i>Centropristis</i> .....	1213
324	<i>pinnafasciatus</i> .....	2647	<i>Serranus</i> .....	1213
139	<i>planus</i> .....	2647	<i>psittaculus</i> , <i>Julis</i> .....	1597
987	<i>Pseudopriacanthus</i> .....	1239	<i>Labrus</i> .....	1596
478	<i>altus</i> .....	1239	<i>psittacus</i> , <i>Callyodon</i> .....	1638
478	<i>serrula</i> .....	1239	<i>Cheilichthys</i> .....	1740
565	<i>Pseudorhamdia</i> .....	153, 154	<i>Colomesus</i> .....	1740
228	<i>piscatrix</i> .....	155	<i>Coryphæna</i> .....	1619
233	<i>Pseudorhombus</i> .....	2624	<i>Lachnolaimus</i> .....	1580
904	<i>adpersus</i> .....	2627	<i>Pseudoscarus</i> .....	1647
2539	<i>brasilienis</i> .....	2626	<i>Scarus</i> .....	1647
2539	<i>californicus</i> .....	2626	<i>Tetrodon</i> .....	1740
2540	<i>dentatus</i> .....	2630, 2632	<i>Xyrichtys</i> .....	1618, 1619
2406	<i>oblongus</i> .....	2630	<i>Psychrolutes</i> .....	2025
2406	<i>ocellaris</i> .....	2630	<i>paradoxus</i> .....	2026
950	<i>quadrocellatus</i> .....	2635	<i>zebra</i> .....	2027
951	<i>vorax</i> .....	2626	<i>Psychrolutina</i> .....	1883
950	<i>Pseudoscarus</i> <i>acutus</i> .....	1652	<i>Psychromaster</i> .....	1099
951	<i>aracanga</i> .....	1648	<i>tuscumbia</i> .....	1109
951	<i>cæruleus</i> .....	1654	<i>Pteraclide</i> .....	955
950	<i>chloris</i> .....	1648, 1654	<i>Pteraclis</i> .....	955
951	<i>cælestius</i> .....	1655, 1656	<i>carolinus</i> .....	956
2617	<i>diadema</i> .....	1646	<i>trichopterus</i> .....	956
2018	<i>fluvomarginatus</i> .....	1652	<i>Pterengraulis</i> .....	450
2680	<i>gnathodus</i> .....	1650	<i>atherinoides</i> .....	450
2608	<i>guacamaja</i> .....	1656, 1657, 1659	<i>Pterocephala</i> .....	92
			<i>Pterognathus</i> .....	2354, 2355

	Page.		Page.
Pteronotus.....	149	puellaris, Decodon.....	1584
Pterophryne.....	2715	Puerco Espino.....	1745
glbba.....	2717	Pez.....	1700, 1704
histrion.....	2716	Puffer.....	1733
kevigata.....	2717	Smooth.....	1728
Pterophrynoides.....	2715	Southern.....	1732
Pteroplatea.....	86	Puffers.....	1726
erebripunctata.....	87, 2753	Sharp-nosed.....	1740
umclura.....	80	pugetensis, Arctedius.....	1890
marmorata.....	87, 2754	Chitonotus.....	1890, 1891
rava.....	2754	Icelus.....	1891
Pteropodus.....	1765, 1776, 1819, 2860	Puget Sound Pollack.....	2536
dallii.....	1819	pugetti, Gasterosteus.....	751
Ptilichthyidae.....	2451	pulchella, Gila.....	234
Ptilichthys.....	2452	Harpe.....	1584
goodiei.....	2452	Montana.....	272
Ptychocheilus.....	224	pulchelloides, Leuciscus.....	222
gracilis.....	225	pulchellus, Bodianus.....	1584
grandis.....	225, 2796	Cheilonemus.....	222
harfordi.....	225, 2797	Cossyphus.....	1584
lucius.....	225	Cyclogaster.....	2127
major.....	225, 2707	Cymatogaster.....	1503
oregonensis.....	224, 2796	Haplocheilus.....	659
rapax.....	225	Histiophorus.....	891
vorax.....	227	Leuciscus.....	221
Ptycholepis.....	414	Leucosomus.....	222
Ptychostomus.....	187	Liparis.....	2126
albidus.....	192	Oligocephalus.....	1089
albus.....	191	Squalius.....	231
hreviceps.....	196	Zygonectes.....	659
bucco.....	191	putcher, Eques.....	1489
cervinus.....	197	Labrus.....	1585
collapsus.....	190	Neozoarches.....	2426
conus.....	196	Pimelometopon.....	1585
coregonus.....	191	Semicossyphus.....	1585
crassilabris.....	194	Squalius.....	234
duquesnei.....	193	Trochocopus.....	1585
erythrus.....	193	pulchra, Harpe.....	1585
haydeni.....	187	Tigoma.....	234
lachrymalis.....	194	pullum, Chondrostoma.....	206
oneida.....	193	pullus, Amiurus.....	141
papillosus.....	189	Cantherines.....	1713
pidiensis.....	191	Monacanthus.....	1713
robustus.....	193	Pimelodus.....	141
thalassius.....	192	pulverus, Fundulus.....	652
velatus.....	190	Zygonectes.....	652
Ptyonotus.....	2005	pulvorulentum, Oxygeneum.....	207
thompsonii.....	2005	pulvorulentus, Mylolenus.....	246
Pudding-wife.....	1590	Pumpkin Seed.....	1009
Pudiano.....	1583	Puuaru.....	2397
Verde.....	1590, 1591	puneta, Furcaria.....	1547
Vermelho.....	1583	punctata, Alutera.....	1718, 1719
puella, Hypoplectrus.....	1192	Bairdiella.....	1434
unicolor.....	1192	Gambusia.....	679
Plectropoma.....	1192	Lamna.....	48
puellaris, Cossyphus.....	1584	Murana.....	395



Page.
1584
1745
1700, 1704
1733
1728
1732
1726
1740
1890
1890, 1891
1891
2536
751
234
1584
272
222
1584
222
1584
2127
1503
659
891
221
222
2126
1089
231
659
1489
1585
2426
1585
1585
234
1585
1585
234
206
141
1713
1713
141
652
652
207
246
1009
2397
1547
1718, 1719
1434
679
48
395

	Page.
punctata, <i>Muraenopsis</i> .....	397
<i>Perca</i> .....	1145, 1146, 1433
<i>Schena</i> .....	1434
<i>Trigla</i> .....	2170
punctatissima, <i>Anguilla</i> .....	348
punctatissimus, <i>Canthigaster</i> .....	1741
<i>Tetrodon</i> .....	1741
punctatum, <i>Hyperprosopon argenteum</i> .....	1502
<i>Myctophum</i> .....	570
<i>Opisthognathus</i> .....	2281
<i>Sicydium</i> .....	2867
punctatus, <i>Apomotis</i> .....	997
<i>Balistes</i> .....	1702
<i>Blennius</i> .....	2390, 2440
<i>Bodianus</i> .....	1146
<i>fulvus</i> .....	1146
<i>Bryttus</i> .....	998
<i>Caranx</i> .....	908
<i>Carcharias</i> .....	41
<i>Ceratacanthus</i> .....	2860
<i>Chinus</i> .....	2440
<i>Decapterus</i> .....	907
<i>Dermatolepis</i> .....	1168
<i>Diodon</i> .....	1746
<i>Enneacentrus</i> .....	1146
<i>Epinophelus</i> .....	1154, 1146
<i>Eques</i> .....	1488, 1489
<i>Fundulus</i> .....	637, 2827
<i>Gadus</i> .....	2553
<i>Gunnellus</i> .....	2440
<i>Holocentrus</i> .....	1153
<i>Hyplocrochilus</i> .....	2390
<i>Ichthaelurus</i> .....	135
<i>Ictalurus</i> .....	134
<i>Isesthes</i> .....	2390
<i>Lepomis</i> .....	998
<i>Micristodus</i> .....	52
<i>Monacanthus</i> .....	1713, 1719
<i>Morrhua</i> .....	2543
<i>Myrophis</i> .....	371
<i>Oriisthognathus</i> .....	2281
<i>Priouotus</i> .....	2158, 2164, 2169
<i>Prionurus</i> .....	1695
<i>Silurus</i> .....	135
<i>Squalus</i> .....	26, 43
<i>Stichæus</i> .....	2439
<i>Tetrodon</i> .....	1735
<i>Trachinus</i> .....	1153
<i>Upeneus</i> .....	859
<i>Xesurus</i> .....	1694, 1695
puncticeps, <i>Cryptopterus</i> .....	382
<i>Ophichthys</i> .....	382
<i>Ophichthys</i> .....	382
puncticulata, <i>Gambusia</i> .....	680
<i>Perca marina</i> .....	1146

	Page.
puncticulatus, <i>Apogonichthys</i> .....	1111
<i>Arius</i> .....	131
<i>Chilomycterus</i> .....	1750
punctifer, <i>Crotalopsis</i> .....	387
<i>Dionda</i> .....	215
<i>Ophichthys</i> .....	387
<i>Hybognathus (Dionda)</i> ..	215
punctiferus, <i>Bodianus</i> .....	1147
<i>Menephorus</i> .....	1147
punctipinne, <i>Siphostoma</i> .....	763
punctipinnis, <i>Ayresia</i> .....	1548
<i>Chromis</i> .....	1548
<i>Dermatostethus</i> .....	763
punctulata, <i>Coryphaena</i> .....	953
<i>Micropora</i> .....	1104
<i>Uranidea</i> .....	1949
punctulatum, <i>Boleosoma</i> .....	1091
<i>Etheostoma</i> .....	1090
punctulatus, <i>Calliurus</i> .....	992, 1011
<i>Cottus</i> .....	1948
<i>hairdi</i> .....	1950
<i>Gobiosox</i> .....	2338
<i>Hippocampus</i> .....	777
<i>Laupagus</i> .....	953
<i>Mimulus</i> .....	302
<i>Notropis umbratilis</i> ..	301
<i>Pimelodus</i> .....	143
<i>Pacificichthys</i> .....	1091
<i>Potamocottus</i> .....	1949
<i>Pseudoscirus</i> .....	1646
<i>Scarus</i> .....	1645
<i>Sicyases</i> .....	2338
<i>Squalus</i> .....	26
Puñceas .....	2195
pungitius, <i>Gasterosteus</i> .....	745
<i>brachypoda</i> .....	746
<i>Pygosteus</i> .....	745
<i>brachypoda</i> ..	746
puncens, <i>Hyborhynchus</i> .....	218
Puraque .....	63
purpurascens, <i>Elops</i> .....	410
purpuratus, <i>Salmo</i> .....	492, 499, 2819
<i>houvieri</i> .....	496
purpurea, <i>Tigoma</i> .....	234
purpurescens, <i>Anoplarchus</i> .....	2423
<i>Lepomis</i> .....	1006
<i>Salpa variegata</i> .....	1271
purpurens, <i>Leuciscus</i> .....	234
<i>Lutjanus</i> .....	1264
<i>Merlangus</i> .....	2535
<i>Sebastichthys</i> .....	1826
<i>Squalus</i> .....	234
Pursy Minnows .....	670, 671
pusilla, <i>Aphorista</i> .....	2711
<i>Bothrocara</i> .....	2476
<i>Clupea</i> .....	426

	Page.		Page.
<i>pusilla</i> , <i>Maynea</i> .....	2476	<i>quadriporus</i> , <i>Gobius</i> .....	2221
<i>Perca</i> .....	1107	<i>quadripunctatus</i> , <i>Scomber</i> .....	809
<i>Platessa</i> .....	2647	<i>quadriremis</i> , <i>Exocoetus</i> .....	735
<i>pusillum</i> , <i>Acanthidium</i> .....	55	<i>quadriscutis</i> , <i>Arius</i> .....	126
<i>pusillus</i> , <i>Argyrosomus</i> .....	470	<i>Netuma</i> .....	126
<i>Etmopterus</i> .....	55	<i>quadriseriatus</i> , <i>Artedius</i> .....	1897
<i>Spinax</i> .....	55	<i>Icelinus</i> .....	1897
<i>Symphurus</i> .....	2710	<i>Icelus</i> .....	1897
<i>putaol</i> , <i>Gymnotus</i> .....	341	<i>quadrispinosus</i> , <i>Pseudosearus</i> .....	1648
<i>putnami</i> , <i>Aelpenser</i> .....	104	<i>Searus</i> .....	1648
<i>Cottogaster</i> .....	1046	<i>quadrituberculata</i> , <i>Platessa</i> .....	2648
<i>Euchalirodus</i> .....	2650	<i>quadrituberculatus</i> , <i>Parophrys</i> .....	2648
<i>Liopsetta</i> .....	2650	<i>Pleuronectes</i> .....	2648
<i>pygmaea</i> , <i>Eucalia inconstans</i> .....	744	<i>Pleuronichthys</i> .....	2638
<i>Umbra</i> .....	624	<i>quadrocellata</i> , <i>Anclyopsetta</i> .....	2635
<i>lini</i> .....	624	<i>Platessa</i> .....	2633
<i>pygmaeus</i> , <i>Cladus</i> .....	2542	<i>quadrocellatus</i> , <i>Pseudorhombus</i> .....	2635
<i>Gasterosteus</i> .....	744	<i>quappella</i> , <i>Etheostoma</i> .....	1804
<i>Leuciscus</i> .....	624	<i>quartus</i> , <i>Anthias rondeleti</i> .....	1266
<i>Pygosteus</i> .....	745	<i>Quasky</i> .....	514
<i>pungitius</i> .....	745	<i>Quassilabla</i> .....	198
<i>brachypoda</i> .....	745	<i>lacera</i> .....	199
<i>Pylodictis ilmosus</i> .....	143	<i>Quassiremus</i> .....	380
<i>pyramidatus</i> , <i>Cyclopterus</i> .....	2097	<i>oviventris</i> .....	380
<i>pyrrhogaster</i> , <i>Chrosomus</i> .....	210	<i>nothochir</i> .....	380
<i>pyrrhouelae</i> , <i>Cliola</i> .....	281	<i>quatuordecimlaminatus</i> , <i>Echeneis</i> .....	2272
<i>Codoma</i> .....	281	<i>Queentish</i> .....	1397
<i>Notropis</i> .....	280	<i>quonseii</i> , <i>Pleuronectes</i> .....	2654
<i>Photogenis</i> .....	281	<i>querclunus</i> , <i>Macrostoma</i> .....	554
<i>Pythonichthys</i> .....	390	<i>querinus</i> , <i>Notoscopelus</i> .....	555
<i>sanguineus</i> .....	390	<i>Queriman</i> .....	810
<i>quadracus</i> , <i>Apeltes</i> .....	752	<i>Querimma</i> .....	817
<i>Gasterosteus</i> .....	752	<i>gyrans</i> .....	818
<i>quadrangularis</i> , <i>Selene</i> .....	1668	<i>harengus</i> .....	817
<i>quadratus</i> , <i>Porocottus</i> .....	1998	<i>querna</i> , <i>Azevia</i> .....	2675
<i>Zens</i> .....	1668	<i>Cyclopacta</i> .....	2675
<i>quadricornis</i> , <i>Ostracium</i> .....	1725	<i>Quia-quia</i> .....	907
<i>quadricornis</i> , <i>Agonus</i> .....	2041	<i>Quiebru</i> .....	898
<i>Aspidophorus</i> .....	2041	<i>quiebru</i> , <i>Chorinemus</i> .....	899
<i>Cottus</i> .....	2001	<i>Lelbia</i> .....	899
<i>Hypsogonus</i> .....	2038, 2041	<i>quiescens</i> , <i>Copelandellus</i> .....	1100
<i>Ostracion</i> .....	1725	<i>Etheostoma</i> .....	1101
<i>quadridens</i> , <i>Pleuronectes</i> .....	2654	<i>Pectlichthys</i> .....	1101
<i>quadrifasciatus</i> , <i>Chasmodes</i> .....	2392	<i>Prionotus</i> .....	2161
<i>Pholis</i> .....	2392, 2394	<i>Uranidea</i> .....	1968
<i>quadrifilis</i> , <i>Bathypterois</i> .....	545	<i>Quietula</i> .....	2251
<i>Cottus</i> .....	1998, 2000	<i>y-cauda</i> .....	2251, 2252
<i>Porocottus</i> .....	1999, 2000	<i>Quillback</i> .....	167
<i>quadrigutta</i> , <i>Pomacentrus</i> .....	1570	<i>Quill-fishes</i> .....	2451
<i>quadrilateralis</i> , <i>Coregonus</i> .....	465	<i>Quimmat Salmon</i> .....	474, 479
<i>quadrilineatus</i> , <i>Hamulon</i> .....	1309, 1311	<i>quinnat</i> , <i>Oncorhynchus</i> .....	480
<i>quadriloba</i> , <i>Rala</i> .....	90	<i>Salmo</i> .....	480
<i>Rhinoptera</i> .....	90	<i>quinquarius</i> , <i>Pentanemus</i> .....	828
<i>quadrimaculatus</i> , <i>Diodon</i> .....	1746	<i>Polynemus</i> .....	828
<i>Paraliodon</i> .....	1746	<i>quinqueaculeata</i> , <i>Raja</i> .....	88

Page.		Page.
2221	quinquecinctus, Pomacanthus.....	1080
860	quinquefasciatus, Epinephelus.....	1164
735	Serranus .....	1164
126	quinquefasciatus, Hemulon.....	1311
126	quinquemaculatus, Centronotus.....	2430
1897	Oplithocentrus.....	2430
1897	Qulsutsch .....	480
1897	quoyi, Cestracion.....	21
1648	Gyroleurodus .....	21
1648		
2648	Rabbit-fish .....	882, 1748
2648	Rabbit-mouth Sucker .....	108, 109
2648	Rabdophorus .....	1072
2648	Rabbla .....	144, 145, 146
2638	Rabirubin .....	1274, 1275
2635	de lo Alto .....	1221
2633	gnudzara .....	1586
2635	inermis .....	1274
1804	rabirubra, Anthias .....	1276
1266	Rabirubins .....	1275
514	Rabula .....	390
198	aque-dulcis .....	390
199	longicauda .....	391
380	marmorata .....	391
380	panamensis .....	391
380	Raccoon Perch .....	1023
2272	Rachycentridae .....	946
1397	Rachycentron .....	948
2654	canadus .....	948
554	radiale, Diplectrum .....	1204
555	radialis, Centropristis .....	1205
810	Diplectrum .....	1205
817	Serranus .....	1205
818	radians, Centropristis .....	1208
817	Diplectrum .....	1208
2675	Lubrus .....	1633
2675	Serranus .....	1632, 1633
907	Serranus .....	1208
898	Sparisoma .....	1632
899	radiata, Raja .....	69
899	Raja .....	69
1100	radiato, Turdus oculo .....	1703
1101	radiatus, Chaerophilus .....	1591
1101	Halicichares .....	1591
2161	Iridio .....	1590
1968	Lubrus .....	1591
2251	Lophius .....	2738
2251, 2252	radiatus, Ogocephalus .....	2738
167	Platygllossus .....	1591, 1597
2451	Sparus .....	1596
474, 479	radiosus, Antennarius .....	2725
480	Radulinus .....	1919
480	asprellus .....	1920
828	boleoides .....	1919
828	rafinesquei, Acipenser .....	106
88	Collettia .....	567
	Cylindrosteus .....	111

	Page.
rafinesquei, Myctophus .....	567
Scaphirhynchus .....	107
Scopelus .....	567
Rafinesquellus.....	1066, 1068, 1082
Rag Fishes.....	968
Rala .....	66
americana .....	69
bivostrius .....	93
chantenay .....	71
cooperi .....	73
desmarestia .....	71
diaphana .....	71
eglanteria .....	68, 71
erinacea .....	68
fimbriata .....	93
flagellum .....	88
inornata .....	73
inermis .....	73
jordani .....	73
maclura .....	87
manatla .....	93
narinari .....	88
obtusata .....	2751
ocellata .....	69
parvifera .....	75
quadriflora .....	90
radiata .....	69
rhina .....	72
trachura .....	76
tuberculata .....	84
Raiada, Majarra.....	1561
Raiado, Roncador .....	1301, 1313
Sargo .....	1361
Raic tuberculée .....	84
raii, Brama.....	958, 959
Sparus .....	960
Rainbow Darter.....	1088
Herring .....	524
Trout .....	500
Rainwater Fish .....	665
Raizero .....	1247, 1251, 1273
Raja .....	66, 2750
abyssicola .....	76, 2751
ackleyi .....	70
alantica .....	75, 2751
binoculata .....	72
bonasus .....	90
centrura .....	83
diabolus marinus .....	93
eglanteria .....	71
equatorialis .....	74, 2751
erinacea .....	69
fylke .....	68
granulata .....	72
inornata .....	73
interrupta .....	2751

	Page.		Page.
<i>Raja jamaicensis</i> .....	81	<i>rathbuni</i> , Pontinus .....	1857
<i>hevils</i> .....	71	<i>Upeneus</i> .....	857
<i>ocellata</i> .....	68	<i>Raton</i> .....	829
<i>ornata</i> .....	71	<i>Rat-tail</i> , Common .....	2583
<i>parmifera</i> .....	74	<i>rauehi</i> , Aelpenser .....	100
<i>percellens</i> .....	63	<i>raucus</i> , Sargus .....	1364
<i>plutonla</i> .....	69, 70	<i>rava</i> , Pteroplatea .....	2751
<i>quinqueoculenta</i> .....	88	<i>Raven</i> , Sea .....	970, 2023
<i>rallata</i> .....	69	<i>raveneli</i> , Esox .....	626
<i>rhina</i> .....	72	<i>raveneli</i> , Pomotis .....	1010
<i>rosispinis</i> .....	2751	<i>Ravens</i> , Sea .....	2022
<i>suy</i> .....	86	<i>Ray</i> , Butterfly .....	80
<i>sentia</i> .....	71	<i>California Sting</i> .....	89
<i>sloani</i> .....	81	<i>Common Sting</i> .....	83
<i>stellulata</i> .....	75	<i>Cow-nose</i> .....	90
<i>trachura</i> .....	75	<i>Southern Sting</i> .....	86
<i>raji</i> , Brama .....	960	<i>Spotted Sting</i> .....	88
<i>Rajidae</i> .....	66	<i>Rays</i> .....	59
<i>raufelhana</i> , Harriotta .....	96	<i>Eagle</i> .....	87, 89
<i>raunfelsbergii</i> , Mugil .....	812	<i>Electric</i> .....	76
<i>ramosum</i> , <i>Pristipoma</i> .....	1334	<i>Round Sting</i> .....	79
<i>ramosus</i> , <i>Pomadasis</i> .....	1334	<i>Sting</i> .....	79, 82
<i>Ranularia</i> .....	2633	<i>Thick-tailed</i> .....	60
<i>dendritica</i> .....	2633	<i>Whip-tailed</i> .....	79
<i>ransennetii</i> , <i>Neoditrema</i> .....	1511	<i>Razor-back Buffalo</i> .....	164
<i>ranula</i> , <i>Careproctus</i> .....	2134	<i>Sucker</i> .....	184
<i>Liparis</i> .....	2134	<i>Razor-fish</i> .....	1617, 1618
<i>Ranzania</i> .....	1755	<i>Real</i> , <i>Matajnelo</i> .....	410
<i>makua</i> .....	1755	<i>rectangulare</i> , <i>Cichlasoma</i> .....	1515
<i>truncata</i> .....	1755, 1756	<i>rectangularis</i> , <i>Acara</i> .....	1515
<i>Ranzaniae</i> .....	1752	<i>rectifrenum</i> , <i>Enpomacentrus</i> .....	1553
<i>rapax</i> , <i>Ptychocheilus</i> .....	225	<i>Pomacentrus</i> .....	1554
<i>raphidoma</i> , <i>Belono</i> .....	716	<i>recurvirostris</i> , <i>Sayris</i> .....	725
<i>Tylosurus</i> .....	715	<i>Red-bellied Dace</i> .....	209
<i>raptor</i> , <i>Gadus</i> .....	2552	<i>Redbreast Bream</i> .....	1001
<i>Raro</i> .....	404	<i>reddingi</i> , <i>Orthopristis</i> .....	1336
<i>rarus</i> , <i>Rhinoscopelus</i> .....	569	<i>Red Drums</i> .....	1453
<i>Scopelus</i> .....	569	<i>Red-eye</i> .....	990
<i>Rascacio</i> .....	1848	<i>Little</i> .....	996
<i>rascacio</i> , <i>Scorpena</i> .....	1849	<i>Mullet</i> .....	814
<i>Rasciera</i> .....	1794	<i>Red Fallfish</i> .....	286
<i>Rasher</i> .....	1794	<i>Red-fin</i> .....	281, 298
<i>rashleighanus</i> , <i>Squalus</i> .....	51	<i>Redfish</i> .....	481, 1453, 1760
<i>rastralis</i> , <i>Stolephorus</i> .....	2811	<i>Bull</i> .....	1453
<i>rastrelliger</i> , <i>Physiculus</i> .....	2549	<i>California</i> .....	1585
<i>Sebastichthys</i> .....	1820	<i>Little</i> .....	482
<i>Sebastodes</i> .....	1819, 1821	<i>Red Goatfish</i> .....	858
<i>Rastrinus</i> .....	1909	<i>Grouper</i> .....	1160
<i>scutigera</i> .....	1909	<i>Guativera</i> .....	1145
<i>Ratfish</i> .....	95	<i>Gurnard</i> .....	2177
<i>Rathbunella</i> .....	2289	<i>Hind</i> .....	1141, 1158
<i>hypoplecta</i> .....	2290	<i>Redhorse</i> .....	187
<i>rathbuni</i> , <i>Fundulus</i> .....	649	<i>Common</i> .....	192
<i>Mulloides</i> .....	857	<i>Texas</i> .....	192
<i>Paralanchurus</i> .....	1479	<i>redi</i> , <i>Orthrorogiscus</i> .....	1754
<i>Polycirrus</i> .....	1479	<i>Red-mouth Buffalo Fish</i> .....	163

Page.		Page.		Page.
1857	Red-mouth Gunt	1308	Remora remora	2271
857	Red Parrot Fish	1035	remora, Echeueis	2272
820	Porgy	1356	Remora	2271
2583	Rockfish	1805	Remoras	2205, 2271
106	Rock Trout	1872	Remorina	2271, 2272
1364	Roncador	1450	remoroides, Echeueis	2272
2754	Sculpin	1935	Remoropsis	2271, 2272
976, 2023	Snapper	1204	brachypterus	2272
623	Sturgeon	100	remotus, Carcharhinus	37
1010	Sucker	176	Serranus	1160
2022	Red-sided Shiner	240	Renicops fiburo	44
86	Red-spotted Sunfish	1004	repandus, Serranus	1187
89	Trout	507	Requiem	38
83	Red-tail Snapper	1270	Sharks	27
90	Red-winged Sea-robin	2156	Requin	38
86	reflexo, Labrus rostro	1077	resplendens, Lampanyctus	555
88	regale, Cybllum	875	reticularis, Anchomus	1735
50	Cynoscion	1407	Antennarius	2719
87, 89	Myctophum	563	reticulata, Amia	113
76	Nannobranchium	563	Liparis	2108
79	regalis, Cestreus	1407	Mycteroperca	1187
79, 82	thalassinus	1408	Paeilia	2833
60	Cynoscion	1407	Solea	2696
70	Euchelyopus	2553	Sputularia	102
164	Johnius	1407	Thalassophryne	2325
184	Otolithus	1407	reticulatum, Bæostoma	2696
1617, 1618	Phycis	2553	reticulatus, Bryttus	998
410	Scomber	875	Catostomus	179
1515	Scomberomorus	875	Cestreus	1409
1515	regis, Atherinops	808	Chilomycterus	1751
1553	regius, Bleinius	2553	Cynoscion	1408
1554	Hybognathus	213	Diodon	1751
725	Laupris	955	Esox	628
299	Phycis	2553	Gobiesox	2328
1001	Urophycis	2553	Halientichthys	2741
1336	Zeus	955	Lepadogaster	2328
1453	regulus, Paenes	951	Lucius	627
990	Sebastes	1761	Lycodes	2465
996	Reina	1815	Monoclin	2696
814	reinhardi, Careproctus	2133, 2134	Notropis	262
286	reinhardtii, Corynolophus	2733	Orbis	1750
281, 298	Himantolophus	2733	Otolithus	1409
481, 1453, 1760	Motella	2559	Pseudoxiphophorus	678
1453	Onos	2559	Trisotropis	1187
1585	Reinhardtius	2610	retifer, Catulus	25
482	hippoglossoides	2611	Seylliorhinus	25
858	remifer, Archosion	1399	retifera, Muraena	401
1160	Isopisthus	1399	retiferum, Seyllium	25
1145	remiger, Myctophum	573	retractus, Calliodon	1623
2177	Ophisurus	384	Cryptotomus	1623
1141, 1158	Remilegia	2270	retrocurrens, Haemulon	1297
187	australis	2270	retropinnis, Catostomus	175, 2791
192	Remora	2271	Microdesmus	2450
192	albescens	2272	Ophichthys	383
1754	brachyptera	2272	Ophichthys	383
163	jacobea	2272	retrosella, Amia	1109

	Page.		Page.
retrosella, Apogon .....	1108	Rhinichthys badius .....	308
retrospinis, Holocentrum .....	853	cataractae .....	306
Plectrypops .....	853	dulcis .....	306
retzii, Mola .....	1754	dulcis .....	307
Orthragoriscus .....	1754	heushavii .....	312
rex, Catoctonus .....	177	lunatus .....	308
Etheostoma .....	1026	luteus .....	307
Percina .....	1025	marmoratus .....	306
rex-mullorum, Apogon .....	1107	maxillosus .....	307
Rey, Peixe .....	806	meleagris .....	308
Rhacochilus .....	1507	nasutus .....	306
toxotes .....	1507	(Apocope) nevadensis .....	311
Rhamdella .....	149, 150, 151	obtusus .....	308
parryi .....	153	ocella .....	307
Rhamdia .....	149, 150	simus .....	307
baronis-mulleri .....	151	transmontanus .....	307
brachyptera .....	151	(Apocope) velifer .....	312
brausefordi .....	151	Rhinobatidae .....	61
godmani .....	152	Rhinobatus .....	61
guatemalensis .....	152	electricus .....	63
hypselurus .....	152	exasperatus .....	65
latucauca .....	152	glaucoctictus .....	63
managuensis .....	153	glaucoctigma .....	62
microptera .....	153	lentiginosus .....	62, 2750
motaguensis .....	151	leucorhynchus .....	62
nicaraguensis .....	152	maregravii .....	63
parryi .....	153	percellens .....	63
petenensis .....	153	planiceps .....	64
polycaulus .....	153	productus .....	63
salvini .....	152, 2790	spinosus .....	63
wagneri .....	150, 151	stellio .....	2750
Rhamphoberyx .....	846	triseriatus .....	66
leucopus .....	847	undulatus .....	63
pocilopus .....	847	Rhinoberyx .....	818
Rhamphocottidae .....	2029	chryseus .....	847
Rhamphocottus .....	2030	Rhinodontina .....	52
richardsoni .....	2030	Rhinogobius contractus .....	2236
Rhegnopteri .....	781, 827	Rhinoliparis .....	2145
Rhencus .....	1329, 1331	barbulifer .....	2145
Rheocrypta .....	1044	Rhinotemus .....	2560
copelandi .....	1046	caudacuta .....	2560
Rhessodon, Arbacia .....	2340	chubrius .....	2561
Gobiesox .....	2340	Rhinoptera .....	90
Rhina .....	58	bonasus .....	90
squatina .....	59	ensenada .....	91
Rhina, Raja .....	72	quadriloba .....	90
Raja .....	72	steindachneri .....	91
Rhinoleicaria .....	156, 157, 158	vesperfilio .....	90
Rhinonanus .....	1721, 1722	Rhinopterina .....	88, 2753
Rhinichthyoides, Tigoma .....	312	Rhinoscion .....	1455
Rhinichthys .....	305	saturus .....	1457
aronatus .....	308	Rhinoscopelus .....	568
atronasus .....	307	andree .....	569
croceus .....	308	coccol .....	568
lunatus .....	308	rurus .....	569
meleagris .....	308	Rhinoseymnus .....	56

Page.		Page.		Page.
308	Rhinotriacis	30	Rhombus simillimus	967
306	henlei	31	soleiformis	2672
306	rhizophora, Fundulus	644	triacanthus	967
307	rhodochloris, Sebastichthys	1810	xanthurus	966, 2849
312	Sebastodes	1809	Rhonciscus	1329, 1330, 1333
308	rhodopus, Trachinotus	941, 943	Rhotheuca	1096
307	rhodorus, Ascelichthys	2025	blennius	1073
306	rhodespilus, Gobiosox	2335	rhothea, Uranidea	1947
307	rhodoterna, Ditrema	1503	rhotheus, Cottus	1946
308	rhodoterns, Holconotus	1502	rhotheus, Catostomus	181
306	Rhodymeichthys	2114, 2415, 2416	rhyncheus, Acipenser	106
311	ruberrimus	2417	Rhynchias	2841
308	rhomalea, Gnathypops	2285	septipinnis	2841
307	rhomuleus, Opisthognathus	2285	Rhynchichthys	847
307	Squalius	233	Rhynchotus	1741
307	rhombus, Gorros	1374	Rhypticus	1229
312	Rhombochirus	2273	bicolor	1232
61	osteocheir	2273	decoratus	1234
61	Rhombogamoidea	108	maculatus	1234
63	rhomboidalis, Lebias	672	microps	1232
65	Otolithus	1404	nigripinnis	1234
63	Turdus	1691	nigromaculatus	1233
62	rhomboides, Acanthion	942	pifuitosus	1234
62, 2750	Chaetodon	942, 2848	saponaceus	1232
62	Diplodus	1358	subfrenatus	1233
63	Lagodon	1358	xanti	1231
63	Sargus	1358	Rhytidostomus	368
64	Sparus	1358	Riado, Sargo	1321
63	Trachinotus	2247	riahi, Onus	2530
63	Trachinotus	942	Ribbon Fish	1490
2750	Rhomboidichthys	2661	Ribbon Fish	1459
66	ellipticus	2665	Ribbon Fishes	1485
63	leopardinus	2666	richardi, Mesoprion	1273
818	lunatus	2666	ricei, Cottus	1952
847	lunulatus	2666	Uranidea	1953
52	maeniferus	2665	richardi, Caranx	923
2236	ocellatus	2664	Hemirhamphus	720
2145	spinosus	2663	Salmo	483
2145	Rhomboplites	1276	richardsoni, Acipenser	106
2560	aurorubens	1277, 1278	Astronesthes	587
2560	elegans	1278	Chaetodus	587
2561	Rhombotides	1689	Corogonus	465
90	Rhombus	965, 2849	Corvina	1484
90	alpidotus	966, 2849	Cottus	1951
91	aquosus	2650	Rhanaphorotus	2030
90	nramaca	2626, 2672	Trachidermis	1944
91	argentipinnis	966	Uranidea	1952
90	bahiensis	2664	Richardsonius	228, 230, 238
88, 2753	crenulatus	966	balticus	239
1455	levis cornubiensis	2654	lateralis	239
1457	longipinnis	966	rigidus, Pimelodus	155
568	medius	967	rjggersmed, Ocyrus	1276
569	ocellatus	2664	rim, Scomber	928
568	orbicularis	966	rimulor, Bathystoma	1368
569	palmata	966	Hamulon	1300
56	paru	965, 2849	Rhincola	2378

	Page.		Page.
<i>Rimicola eigenmanni</i> .....	2499	Robin, Flying .....	2183
<i>Rimicola</i> .....	2438	Round .....	907
<i>rimiculus</i> , <i>Catostomus</i> .....	2792	<i>rolineta</i> , <i>Gila</i> .....	227
<i>rimosus</i> , <i>Etheostomus</i> .....	2688	<i>Puffin</i> .....	1154
Ringed Perch .....	1023	<i>robustus</i> , <i>Moxostoma</i> .....	193
<i>ringens</i> , <i>Atopoclinus</i> .....	2376	<i>robustus</i> , <i>Benthoicyetes</i> .....	2514
<i>Balistes</i> .....	1709	<i>Glyptocephalus</i> .....	1938
<i>Melicthys</i> .....	1711	<i>Excavatus</i> .....	746
<i>Phenodon</i> .....	586	<i>Fundulus</i> .....	644
<i>Stolephorus</i> .....	449	<i>Hippoglossoides</i> .....	2616
<i>Sudis</i> .....	601	<i>Ictalurus</i> .....	135
<i>Xanthichthys</i> .....	1709	<i>Leuciscus</i> .....	228
<i>Rio Grande Chub</i> .....	233	<i>Mylopharodon</i> .....	219
Trout .....	495	<i>Neobythites</i> .....	2515
<i>riparium</i> , <i>Holocentrum</i> .....	852	<i>Psychostomus</i> .....	193
<i>risso</i> , <i>Alpismaris</i> .....	537	<i>Rocens</i> .....	1131, 1132
<i>rissoanus</i> , <i>Notacanthus</i> .....	618	<i>chrysopterus</i> .....	1132
<i>rissol</i> , <i>Trachurus</i> .....	910	<i>comes</i> .....	1407
<i>Rissola</i> .....	2489	<i>lineatus</i> .....	1113, 1132
<i>marginata</i> .....	2489	<i>saxatilis</i> .....	1133
<i>rivalis</i> , <i>Salmo</i> .....	509, 510	<i>atriatus</i> .....	1133
<i>River Chub</i> .....	322	<i>rocheanus</i> , <i>Thynnus</i> .....	868
Drum .....	1483	<i>rochel</i> , <i>Anxela</i> .....	868
Lampreys .....	10	<i>Scomber</i> .....	867
Perch .....	1023	<i>Rock</i> .....	1132
Perch of New York .....	1135	Bass .....	989, 1197
<i>riverendi</i> , <i>Cyprinodon</i> .....	673	Common .....	990
<i>variegatus</i> .....	673	Beauty .....	1684
<i>Trifareus</i> .....	673	Cook .....	1575, 1576
<i>riroliana</i> , <i>Seriola</i> .....	904	Hind .....	1152
<i>rivularis</i> , <i>Salmo</i> .....	500	Salmon .....	905
<i>rivulatus</i> , <i>Cirrhitus</i> .....	1491	Sea Bass .....	1201
<i>Cirrhitichthys</i> .....	1492	Shellfish .....	1722
<i>Diodon</i> .....	1748	Sturgeon .....	106
<i>Serranus</i> .....	1187	Trout .....	1866, 1867, 1872
<i>Rivulus</i> .....	662, 2830	<i>Rockfish</i> .....	639, 1026, 1132, 1172
<i>cylindraceus</i> .....	662, 2830	Black .....	1784
<i>isthmensis</i> .....	2830	Black and yellow .....	1825
<i>marmoratus</i> .....	663, 2830	Black-banded .....	1827
<i>Roach</i> .....	250	Brown .....	1817
<i>Roaches</i> .....	243	Flesh-colored .....	1824
<i>roanoka</i> , <i>Etheostomus</i> .....	1036	Grass .....	1819
<i>Hadropterus</i> .....	1036	Orange .....	17, 3
<i>Percina</i> .....	1036	Percy .....	1133
<i>robalito</i> , <i>Centropomus</i> .....	1123	Red .....	1805
<i>Robalito de las Aletas Amarillas</i> .....	1123	Spotted .....	1806
<i>Prietas</i> .....	1119	Yellow backed .....	1822
<i>Robalo</i> .....	1118	Yellow-spotted .....	1826
<i>Prieto</i> .....	1119	Yellow-tail .....	1781
<i>Robalos</i> .....	1117	<i>Rock-fishes</i> .....	1758, 1765
<i>roberti</i> , <i>Excavatus</i> .....	735	<i>Rocklings</i> , Four-Bearded .....	2560
<i>Hemirhamphus</i> .....	721	Three-Bearded .....	2377
<i>Hyporhamphus</i> .....	721	<i>Rocky Mountain Bullhead</i> .....	749
<i>robertsii</i> , <i>Ceratobatis</i> .....	2756	Trout .....	487
<i>Siphostoma</i> .....	2837	Whitish .....	463
<i>Stolephorus</i> .....	2815	<i>roda</i> , <i>Pristigaster</i> .....	1323



Page.		Page.		Page.
2183	<i>Reboides</i> .....	338	<i>rosa</i> , <i>Hemirhamphus</i> .....	722
907	<i>guatemalensis</i> .....	338	<i>Hyporhamphus</i> .....	721
227	<i>Rogenia</i> .....	421	<i>Typhlichthys</i> .....	2835
1154	<i>alba</i> .....	422	<i>rosarium</i> , <i>Acipenser</i> .....	106
193	<i>rogersi</i> , <i>Urolophus</i> .....	2752, 2753	<i>Rose-back Parrot</i> .....	1635
2514	<i>Roller</i> , <i>Stone</i> .....	181, 204	<i>Rose-fish</i> .....	1760
1938	<i>Rollers</i> , <i>Sund</i> .....	783, 784	<i>Rose-fishes</i> .....	1760
1/3b	<i>Romero</i> .....	900	<i>roscephalus</i> , <i>Notropis</i> .....	298
641	<i>Roncadina</i> .....	1461	<i>rosens</i> , <i>Blennius</i> .....	2420
2616	<i>Roncador</i> .....	1457	<i>Centronotus</i> .....	2420
135	<i>Little</i> .....	1460	<i>Cryptotomus</i> .....	1626
228	<i>Raiado</i> .....	1301, 1313	<i>Gunnellops</i> .....	2420
219	<i>Red</i> .....	1456	<i>Luxilus</i> .....	288
2515	<i>Yellow-finned</i> .....	1467	<i>Minnilus</i> .....	288
193	<i>Roncador stearnsi</i> .....	1457	<i>Notropis</i> .....	287
1131, 1132	<i>roncador</i> , <i>Umbrina</i> .....	1467	<i>Prionotus</i> .....	2159
1132	<i>ronchus</i> , <i>Bairdiella</i> .....	1436	<i>Rosicola</i> .....	1765, 1775, 1793
1407	<i>Corvina</i> .....	1436	<i>rosipes</i> , <i>Novaeenclithys</i> .....	1614
1113, 1132	<i>Sclerata</i> .....	1436	<i>Xyrichtys</i> .....	1615
1133	<i>Ronco</i> .....	1436	<i>rosispinus</i> , <i>Raja</i> .....	2751
1133	<i>Amarillo</i> .....	1303	<i>rossi</i> , <i>Lycodes</i> .....	2465
868	<i>Arará</i> .....	1304	<i>rossi</i> , <i>Salvelinus</i> .....	510
868	<i>Blanco</i> .....	1297	<i>rossii</i> , <i>Salmo</i> .....	510
867	<i>Carbonero</i> .....	1300	<i>rostellum</i> , <i>Acipenser</i> .....	106
1132	<i>Condado</i> .....	1306	<i>rostrata</i> , <i>Albula</i> .....	412
989, 1197	<i>Prieto</i> .....	1297	<i>Aldrovandia</i> .....	609
990	<i>Ronco</i> .....	1304	<i>Anguilla</i> .....	348
1684	<i>Roncos</i> .....	1291	<i>anguilla</i> .....	348
1575, 1576	<i>Rondamin</i> .....	959	<i>Limanda</i> .....	2645
1152	<i>rondelii</i> , <i>Anthias</i> <i>quartus</i> .....	1266	<i>Loricaria</i> .....	157
905	<i>Carcharodon</i> .....	50	<i>Macdonaldia</i> .....	617
1201	<i>Scombrosox</i> .....	726	<i>Muraena</i> .....	348
1722	<i>Niphias</i> .....	894	<i>Platessa</i> .....	2645
106	<i>Rondeletia</i> .....	548	<i>rostratum</i> , <i>Ciclasoma</i> .....	1522
1866, 1867, 1872	<i>bicolor</i> .....	548	<i>Holocentrum</i> .....	852
1026, 1132, 1172	<i>rondelii</i> , <i>Exocoetus</i> .....	733, 734	<i>rostratus</i> , <i>Agonus</i> .....	2048
1781	<i>Exonotus</i> .....	2836	<i>Aspidophorus</i> .....	2048
1625	<i>Orthogomus</i> .....	1754	<i>Brachyopsis</i> .....	2046, 2048
1827	<i>Sargus</i> .....	1364	<i>Canthigaster</i> .....	1741
1817	<i>Rondeletidae</i> .....	547	<i>Gymnothorax</i> .....	395
1824	<i>Ronquil</i> .....	2289	<i>Halosaurus</i> .....	609
1819	<i>Ronquils</i> .....	2287	<i>Heros</i> .....	1523
173	<i>Ronquilus</i> .....	2289	<i>Heterostichus</i> .....	2351
1133	<i>jordani</i> .....	2289	<i>Holocentrus</i> .....	849
1805	<i>rosacea</i> , <i>Myceroperca</i> .....	1184	<i>Lophius</i> .....	2737
1806	<i>rosaceus</i> , <i>Cymatogaster</i> .....	1500	<i>Notacanthus</i> .....	617
1822	<i>Epinephelus</i> .....	1184	<i>Phycis</i> .....	2555
1826	<i>Lutjanus</i> .....	1267	<i>Pseudoscopus</i> .....	1659
1781	<i>Mesoprion</i> .....	1267	<i>Searus</i> .....	1658
1758, 1765	<i>Micrometrus</i> .....	1500	<i>Sphagebranchius</i> .....	373
5660	<i>Paraliparis</i> .....	2142	<i>Squalus</i> .....	49
257	<i>Sebastes</i> .....	1794	<i>Tetradon</i> .....	1742
449	<i>Sebastodes</i> .....	1608	<i>Zelus</i> .....	936
487	<i>Trisetropis</i> .....	1184	<i>rostru</i> , <i>Labrus</i> <i>reflexo</i> .....	1677
463	<i>Zalemblus</i> .....	1500	<i>rotengulus</i> , <i>Leuciscus</i> .....	221
1323	<i>rosae</i> , <i>Clevelandia</i> .....	2255	<i>rotheus</i> , <i>Leucosomus</i> .....	222

	Page.		Page.
rothrocki, <i>Notogrammus</i> .....	2440	<i>Rubio Volador</i> .....	2164
<i>Poroclinus</i> .....	2434	<i>rubra, Liopropoma</i> .....	1137
rotunda, <i>Mola</i> .....	1754	<i>Sclena</i> .....	849
Round Bass.....	988	<i>rubricrocea, Hybopsis</i> .....	286
Herring.....	420	Hydrophlox.....	286
Pámpano.....	941	Minnilus.....	286
Robin.....	907	Notropis.....	286
Sting Rays.....	79	<i>rubrifrons, Alburnellus</i> .....	295
Sunfish.....	988	Alburnus.....	295
Whitefish.....	465	Ceraticthys.....	320
Round-tail.....	227	Fundulus.....	653
rousseau, <i>Siphostoma</i> .....	767	Hybopsis.....	320
<i>Syngnathus</i> .....	767	Leuciscus.....	295
Roussettes.....	22	Minnilus.....	295
Rovetto.....	879	Nocomis.....	320
Rovetus <i>temminkii</i> .....	880	Notropis.....	295
rubella, <i>Sciæna</i> .....	1418	Zygonectes.....	654, 2829
rubellus, <i>Alburnus</i> .....	293	<i>rubripinna, Ciliola</i> .....	281
Leuciscus.....	293	Cyprinella.....	281
Minnilus.....	293	<i>rubripinna, Sparisoma</i> .....	1640
rubens, <i>Centropomus</i> .....	1107	<i>rubripinnis, Argyreus</i> .....	282
<i>Labrus fauotoga</i> .....	1579	Minnilus.....	298
<i>Trigla tota</i> .....	2177	Scarus.....	1640
ruber, <i>Apogon</i> .....	1107	<i>rubrirostris, Catactyx</i> .....	2505
<i>Bodianus</i> .....	1265	<i>rubrivinctus, Sebastichthys</i> .....	1817
<i>finvus</i> .....	1146	Sebastodes.....	1817
<i>Caranx</i> .....	919	<i>rubropunctatus, Salarias</i> .....	2396
<i>Coregonus</i> .....	538	<i>Seartichthys</i> .....	2396
<i>Dipterodon</i> .....	1107	<i>rubrum, Chorististium</i> .....	1136
<i>Epinephelus</i> .....	1181	<i>Rudder-fish</i> .....	902, 1387
<i>Gadus</i> .....	2530	<i>Rudder-fishes</i> .....	962, 964, 1380
<i>Gymnocephalus</i> .....	1146	<i>radis, Abudedefduf</i> .....	1563
<i>Myeteroperca</i> .....	1180	<i>Glyphidodon</i> .....	1563
<i>Rutilus</i> .....	300	<i>rufa, Harpe</i> .....	1583
<i>Scomber</i> .....	919	<i>Morone</i> .....	1135
<i>Sebastes</i> .....	1818	<i>Percu</i> .....	849
<i>Sebastodes</i> .....	1806	<i>Ruffa, Black</i> .....	962, 963
<i>Serranus</i> .....	1181	<i>rufilineatum, Ethicostoma</i> .....	1079
<i>ruberrima, Myeteroperca olfax</i> .....	1183	<i>rufilineatus, Nothonotus</i> .....	1079
<i>ruberrimus, Gunnellus</i> .....	2417	<i>Pocillichthys</i> .....	1079
<i>Pholis</i> .....	2417	<i>rufipinnis, Exocoetus</i> .....	735
<i>Rhodymenichthys</i> .....	2417	<i>Exonantes</i> .....	2836
<i>Sebastodes</i> .....	1805, 1806	<i>rufolineatum, Ethicostoma</i> .....	1079
<i>rubescens, Conger</i> .....	355	<i>rufus, Balistes</i> .....	1707
<i>Exocoetus</i> .....	734	<i>Bodianus</i> .....	1135, 1583
<i>Hypsiniotus</i> .....	1665	<i>Centropistes</i> .....	1199
<i>Steinogeria</i> .....	961	<i>Cossyphus</i> .....	1583
<i>rubicauda, Pomotus</i> .....	1001	<i>Holocentrus ascensionis</i> .....	849
<i>rubicunda, Parma</i> .....	1565	<i>Labrax</i> .....	1135
<i>rubicundus, Aepenser</i> .....	196	<i>Labrus</i> .....	1583
<i>Glyphisodon</i> .....	1565	<i>Onos</i> .....	2559
<i>Hypsypops</i> .....	1564, 1565	<i>Sebastodes</i> .....	1786
<i>Pomacentrus</i> .....	1565	<i>rugifer, Ophichthys</i> .....	384
<i>rubiginosus, Gobiesox</i> .....	2337	<i>rugispinis, Arius</i> .....	130
<i>Sicyases</i> .....	2337	<i>Galeichthys</i> .....	2781
<i>rubio Prionotus</i> .....	2164	<i>Hexanemullichthys</i> .....	130

Page.		Page.		Page.
2164	<i>rugispinis</i> , <i>Tachisurus</i> .....	130	<i>Ryba</i> , <i>Bielaya</i> .....	480
1137	.....	131	<i>Chornia</i> .....	621
849	..... <i>phrygiatus</i> .....	2652	<i>Krasnaya</i> .....	481
286	<i>rugosus</i> , <i>Platichthys</i> .....	898, 906, 921	<i>Rypticine</i> .....	1131
286	<i>Runner</i> .....	2377	<i>Rypticus</i> .....	1229, 1230
286	..... <i>azalea</i> .....	2377	..... <i>arenatus</i> .....	1232
286	<i>Runulina</i> .....	2346	..... <i>bicolor</i> .....	1231
295	.....	106	..... <i>blatrispinus</i> .....	1233
295	<i>rupertianus</i> , <i>Aelpenser</i> .....	1073	..... <i>coriaceus</i> .....	1233
320	<i>rupestre</i> , <i>Etheostoma</i> .....	2426	..... <i>nigripinis</i> .....	1234
653	..... <i>Xiphidion</i> .....	990	..... <i>saponaceus</i> .....	1232
320	<i>rupestris</i> , <i>Ambloplites</i> .....	990	..... <i>xanti</i> .....	1231
295	..... <i>cavifrons</i> .....	2341	<i>Sabalo</i> .....	409, 414
295	..... <i>Arbaciosa</i> .....	990	<i>Sabara</i> , <i>Bacalhao</i> .....	2230
320	..... <i>Bodianus</i> .....	2579	<i>sabina</i> , <i>Dasibatis</i> .....	85
295	..... <i>Coryphaenoides</i> .....	2541	..... <i>Dasyatis</i> .....	84
054, 2829	..... <i>Gadus</i> .....	2341	..... <i>Trygon</i> .....	85
281	..... <i>Gobiesox</i> .....	2582	<i>sablur</i> , <i>Notropis</i> .....	262
281	..... <i>Macrourus</i> .....	2579	<i>Sable</i> .....	889
1640	..... <i>Macrurus</i> .....	1813	<i>saburra</i> , <i>Chasmodes</i> .....	2302
282	..... <i>Sebastichthys</i> .....	1812	<i>Sac-à-Lait</i> .....	638, 987
298	..... <i>Sebastodes</i> .....	1174	<i>Saccopharyngidus</i> .....	405
1640	..... <i>Serranus</i> .....	2341	<i>Saccopharynx</i> .....	405, 406
2505	..... <i>Sicyaees</i> .....	2426	..... <i>ampullaceus</i> .....	406
1817	..... <i>Xiphidion</i> .....	2396	..... <i>chordatus</i> .....	406
1817	<i>Rupiscartus</i> .....	2397	..... <i>flagellum</i> .....	406
2306	..... <i>atlanticus</i> .....	196	<i>Saccostoma</i> .....	2249
2396	<i>rupiscartus</i> , <i>Moxostoma</i> .....	1713	<i>Sacramento Cat</i> .....	140
1136	<i>ruppelii</i> , <i>Monacanthus</i> .....	1593	..... <i>Chub</i> .....	231
002, 1387	<i>ruptus</i> , <i>Cherophilus</i> .....	1593	..... <i>Perch</i> .....	991
064, 1180	..... <i>PlatyGLOSSUS</i> .....	1908	..... <i>Piko</i> .....	224, 2796
1563	<i>Ruscarius</i> .....	143	..... <i>Salmon</i> .....	479
1563	..... <i>meanyi</i> .....	464	..... <i>Sturgeon</i> .....	104
1583	<i>Russian Cat</i> .....	1851	..... <i>Sucker</i> .....	178
1135	<i>Russians</i> , <i>Muksun of the</i> .....	2644	<i>sadina</i> , <i>Clupea</i> .....	420
849	<i>russula</i> , <i>Scorpaena</i> .....	272	..... <i>Etrumeus</i> .....	420
062, 963	<i>Rusty Dab</i> .....	243	<i>sagax</i> , <i>Clupea</i> .....	423
1079	<i>rutila</i> , <i>Moniana</i> .....	321	<i>Sagenichthys</i> .....	1416
1079	<i>Rutilus</i> .....	200	..... <i>ancy lodon</i> .....	1416, 2859
1079	..... <i>amblops</i> .....	244	<i>sagitta</i> , <i>Amblyopus</i> .....	2263
735	..... <i>anomalous</i> .....	247	..... <i>Ellicostoma</i> .....	1080
2836	..... <i>bicolor</i> .....	282, 206	..... <i>Poecilichthys</i> .....	1081
1079	..... <i>boucardi</i> .....	193	..... <i>Tylosurus</i> .....	711
1707	..... <i>compressus</i> .....	244, 2798	..... <i>Tynlastes</i> .....	2263
15, 1583	..... <i>melanurus</i> .....	282	<i>sagittula</i> , <i>Euctenogobius</i> .....	2229
1199	..... <i>olivaceus</i> .....	300	..... <i>Gobius</i> .....	2228
1583	..... <i>plargyrus</i> .....	321	<i>Saibling</i> .....	598
849	..... <i>ruber</i> .....	245	<i>snida</i> , <i>Boreogadus</i> .....	2533
1135	..... <i>storerianus</i> .....	245	..... <i>Gadus</i> .....	2534
1543	..... <i>symmetricus</i> .....	509	<i>Saignour</i> .....	1091
2559	..... <i>thalassinus</i> .....	334	<i>Sailfish</i> .....	167, 890
1786	<i>rutilus</i> , <i>Salmo</i> .....	2108	<i>Sailor's Choice</i> .....	1297, 1338, 1358
384	..... <i>Tetragonopterus</i> .....	879	<i>Salar</i> .....	483
130	<i>rutteri</i> , <i>Neoliparis</i> .....	879	..... <i>lewisii</i> .....	493
2781	<i>Ruvetto</i> .....	879	..... <i>virginalis</i> .....	495
130	<i>Ruvottus</i> .....	879		
	..... <i>pretiosus</i> .....			

	Page.		Page.
salar, Salmo.....	486	Salmo clarkii pleuriticus.....	2819
onananicus.....	487	spilurus.....	2819
sebago.....	487	stomias.....	2819
Salarlae.....	2377	tahoensis.....	2870
Salarlas.....	2397	virginalis.....	2810
atlanticus.....	2397	elnpeiiformis.....	466
chiostictus.....	2398	colii.....	509
margaritaceus.....	2399	conflus.....	505
rabropunctatus.....	2396	confluentus.....	480
textilis.....	2400	consuetus.....	479
Salaricthys.....	2400	cooperi.....	483
textilis.....	2400	(Coregonus) harengus.....	469
Salarifina.....	2346	lucidus.....	471
Salarinus vomerinus.....	2400	tullibee.....	473
Sälbling.....	508	curilus.....	508, 2223
Saleima.....	1384	dermatinus.....	479
aurata.....	1386	distichus.....	509
Salema.....	1358, 1359, 1385	erythrorhynchus.....	508
saliens, Chorinemus.....	899	fontens.....	538
Oligoplites.....	899	fontinalis.....	507
palometa.....	899	gairdneri.....	497
Scomber.....	899	beardsleei.....	2819
salin, Sparus.....	1360	crescentis.....	2821
sallavi, Algansea.....	212	kamloops.....	499
Ceraticthys.....	212	shasta.....	502
Chiola.....	212	stonei.....	503
Hudsonius.....	212	gibber.....	478
salmarinus, Salmo.....	509	gibbsii.....	493
Salmo.....	483, 486	gloveri.....	487
adirondacus.....	505	gorbuscha.....	478
agassizii.....	507	grayi.....	509
alipes.....	509	grenlandicus.....	521
alleghanicus.....	507	hearnei.....	510
alpinus.....	509, 514	hoodi.....	505
nivalis.....	509	hoodii.....	507, 510
amethystinus.....	505	hudsoniens.....	507
arabatsch.....	483	immaculatus.....	507
areticus.....	521	iridens.....	500, 2819
areturns.....	510	agna-bonita.....	503
argyreus.....	480	gilberti.....	502
ascanti.....	509	masoni.....	501
aurora.....	493	shasta.....	502
bairdii.....	508	stonei.....	503
breviceauda.....	493	japonensis.....	479
callaris.....	508	kennerlyi.....	484
campbelli.....	508	keta vel kayko.....	479
canadensis.....	507	killinensis.....	509
canis.....	479	kisutch.....	481
carbonarius.....	509	kundscha.....	2823
carinatus.....	493	lavignus.....	508
clarkii.....	492, 2819	lagocephalus.....	479
bouvieri.....	2819	lavaretus mucksun.....	464
gibbsii.....	2819	leucomenis.....	2823
henshawi.....	2819	lordii.....	508
lowisi.....	2819	lycaodon.....	483
macdonaldi.....	2819	mackenzii.....	474

Page.		Page.		Page.
2819	<i>Salmo macrostoma</i> .....	481	<i>Salmo siscowet</i> .....	505
2819	( <i>Mallotus pacificus</i> ) .....	521	siskawitz .....	505
2819	malmu .....	508	socialis .....	521
2870	marstoni .....	516	spectabilis .....	508
2819	mas .....	487	stagnalis .....	510
466	masoni .....	501	stellatus .....	493
509	melampterus .....	485	striatus .....	481
505	milkschitch .....	481	symmetricus .....	505
480	monestichus .....	509	tapdisma .....	483
479	mutkisi .....	492	( <i>Thymallus</i> ) signifer .....	518
483	muksun .....	464	toma .....	505
469	mulleri .....	574	trachinus .....	533
471	mykiss .....	487, 492, 2818	truncatus .....	499
473	agua-bonita .....	504	trutta .....	487
508, 2223	bouvieri .....	496	tschawytscha .....	480
479	clarkii .....	492	tschawytschiformis .....	478
509	gibbsii .....	493	tsuppitch .....	481, 495
508	henshawi .....	493	tudes .....	508
538	lewisi .....	493	umbra .....	509
507	macdonaldi .....	497	ursinus .....	505
497	pleniticus .....	496	utah .....	495
2819	spilurus .....	495	ventricosus .....	509
2821	stomias .....	497	warreni .....	483
499	virginalis .....	495	willoughbi .....	509
502	myops .....	533	salmoides, Labrus .....	1012
503	naresi .....	515	Micropterus .....	1012
478	nerka .....	483	Salmon .....	483
493	nigrescens .....	507	Atlantic .....	486
487	nitidus .....	509	Blue-back .....	481
478	nummifer .....	508	Chinook .....	479
509	omisco maycus .....	487	Coho .....	480
521	oquassa .....	515	Columbia .....	479
510	orientalis .....	480	Common Atlantic .....	486
505	( <i>Osmerus</i> ) olidus .....	525	Dog .....	478
507, 510	pallidus .....	505	Family .....	460
507	parkei .....	508	Fraser River .....	481
507	paucidens .....	483	Hoopid .....	480
500, 2819	penshinensis .....	508, 2819	Humpback .....	478
503	perisii .....	509	Jack .....	1021
502	proteus .....	478	Killer .....	749
501	purpuratus .....	492, 499, 2819	King .....	479
502	bouvieri .....	496	Landlocked .....	487
503	quinnat .....	480	Le kai .....	478
479	richardi .....	483	Nanaycush .....	505
483	rivalis .....	509, 510	Quinnat .....	474, 479
479	rivularis .....	500	Rock .....	905
509	rossii .....	510	Sacramento .....	479
481	rutilus .....	509	Saw-qui .....	481
2823	salar .....	486	Silver .....	480
508	oumaniche .....	487	Trout .....	497, 2818
479	sehago .....	487	Tyee .....	479
464	salmarinus .....	509	White, of the Colorado .....	225
2823	salvelinus .....	509	salmonea, Ericaria .....	2816
508	sanguinolentus .....	481	Lepomis .....	1011
483	saurus .....	537	Perca .....	1021
474	seouleri .....	478, 481	Salmonete .....	858


	Page.		Page.
Salmonete Amarilla .....	859	sancta-helene, Caraux .....	908
salmonens, Clinanos .....	415	Decapterus .....	908
Esox .....	538, 627, 629	Gyunothorax .....	397
Mugil .....	415	Lycodontis .....	397
Scombrocottus .....	1862	Murana .....	397
Salmonidae .....	460	sancta-lucie, Corvula .....	1429
Salmonidea .....	408	sancta-marthae, Vomer .....	934
Salmonina .....	461	sancta-petri, Vomer .....	934
Salmo .....	783	sancta-rosae, Ulvicola .....	2413
pellucida .....	784	sancti-laurentii, Euryphrys .....	2668
Salmo .....	780, 782	sancti-pauli, Holocentrus .....	853
Salpa purpurascens variegata .....	1271	Sand Dab .....	2614
saltans, Chorinemus .....	899	Darters .....	1061, 1062
saltator, Pomatomus .....	947	Diver .....	535
Scomberoides .....	899	Eel .....	833
Temnodon .....	947	Lannces .....	831, 832, 833
saltatrix, Gasterosteus .....	947	Pike .....	1022
Perca .....	947, 1388	Rollers .....	783, 784
Pomatomus .....	946	Shark .....	40
saludana, Cliola .....	270	Star-gazers .....	2297
saludaus, Alburnops .....	270	Sucker .....	1476
Notropis hudsonius .....	270	Whiting .....	1474
Salvelini .....	506	Sand-fish .....	1207, 2295
Salvelinus .....	506	Sanducha .....	411
alpinus .....	508, 2822	sanguifluus, Nothonotus .....	1077
alipes .....	509	Pacilichthys .....	1077
arcturus .....	510	sanguinea, Murana .....	390
aureolus .....	511	sanguineus, Antemarius .....	2721
stagnalis .....	510	Apodichthys .....	2412
bairdii .....	508	Holocentrus .....	1761
fontinalis .....	506	Pythonichthys .....	390
agassizii .....	507	sanguinolentus, Oncochelychus .....	481
kundscha .....	2822	Pomotis .....	1003
malma .....	507, 2823	Salmo .....	481
nanayensh .....	505	Sa-pen-que .....	487
siscowet .....	505	sapidissima, Alosa .....	427, 428
naresi .....	515	sapidissimus, Coregonus .....	466
nitidus .....	509	Sapo .....	2314, 2315, 2316, 2321
oquassa .....	514, 515	Bagre .....	2319
marstoni .....	515, 2823	saponaceus, Anthias .....	1232
naresi .....	515	Rhypticus .....	1232
parkei .....	2823	Rypticus .....	1232
rossi .....	510	sara, Cybium .....	877
spectabilis .....	508	sargus, Lepodus .....	960
stagnalis .....	509	Sararus apiarius .....	1142
salvelinus, Salmo .....	509	Sarchirus .....	109
salvini, Cotylopus .....	2208	argenteus .....	110
Heros .....	1528	vittatus .....	110
Pimelodus .....	152	Sarcidium .....	302
Rhamdia .....	152	scopiferum .....	303
Sicydium .....	2208	Sarchua .....	887
Sieyopterus .....	2208	Sarcura .....	59, 60
San Diego Solo .....	2707	Sarda .....	871
San Pedro Fish .....	954	chiliensis .....	872
sancte crucis, Pseudoscarnus .....	1051	mediterranea .....	872
Scarus .....	1051	sarda .....	872



Page.		Page.		Page.
908	sarda, Pelamys.....	872	Sarothrodus capistratus.....	1678
908	Scomber.....	872	maenlocinctus.....	1674
397	Sardina Blanca.....	332	nigrirostris.....	1674
397	Bocona.....	440	sedentarius.....	1675
397	de España.....	423	siriatum.....	1677
1420	de Ley.....	430	Sarritor.....	2072
934	Escamuda.....	431	frenatus.....	2073
934	Machete.....	433	leptorhynchus.....	2075
2413	pseudohispanica.....	424	sarritor, Prionotus.....	2160
2668	sardina, Hurengula.....	430	Satanoperca.....	1542
853	Menidla.....	799	crassilabris.....	1542
2614	Sardinella.....	430	satiriens, Neoclinus.....	2355
1061, 1062	Sardine.....	864	saturna, Sciaen.....	1456
535	Sardino, California.....	423	saturnus, Amblodon.....	1456
833	Sardinella.....	428, 429, 2811	Rhinoscion.....	1457
1, 832, 833	unchovia.....	429	Sancer-eyo Porgy.....	1349
1022	apicalis.....	429	Sanger.....	1022
783, 784	bishopi.....	430	Sault Whitefish.....	466
46	clupeola.....	429	Saurels.....	909, 910
2297	humeralis.....	431	Saurenehelys.....	364
1476	macrophthalmus.....	430	Sauries.....	724, 725
1474	sardina.....	430	Saurus.....	533
1207, 2295	stolifera.....	431	anolis.....	535
411	thirssina.....	430	brovirostris.....	533
1077	Sardines, Sealed.....	428	fotens.....	538
1077	True.....	422	grisons.....	537
390	Sardinia.....	422	intormedius.....	535
2721	Sargassum Fish.....	2716	limbatus.....	533
2412	Sargo.....	1363	longirostris.....	538
1761	Raiado.....	1321, 1361	lucioceps.....	539
390	sargoides, Chatodon.....	1562	mexicanus.....	538
481	Sargosomus.....	1495	myops.....	533
1003	fluviatilis.....	1496	spixianus.....	538
481	Sargus.....	1362	synodus.....	536
487	ambassis.....	1346	truncatus.....	533
427, 428	argenteus.....	1363	varius.....	536
466	aries.....	1362	saurus, Elops.....	410
2316, 2321	caribaëus.....	1360	Esox.....	725
2319	caudimacula.....	1363	Oligoplites.....	898
1232	flavolineatus.....	1360	Salmo.....	537
1232	holbrookii.....	1363	Scomber.....	898
1232	humeri-maculatus.....	1360	Scombrosox.....	725
877	ovis.....	1361	Synodus.....	537
960	poutalesii.....	1360	Trachurus.....	911
1142	raucus.....	1364	Saury.....	725
109	rhomboides.....	1358	saussurii, Brama.....	958
110	rondoletii.....	1364	Taractes.....	957
110	tridens.....	1364	Sauteur.....	898, 899
302	unifaculatus.....	1360	Savalle.....	409
303	variegatus.....	1364	Savanilla.....	409
887	vitula.....	1364	savanna, Brachyconger.....	360
59, 60	sargus, Diplodus.....	1363	Murena.....	360
871	Sparus.....	1364	Muranesox.....	360
872	Sarothrodus.....	1672	Savola.....	889
872	unplexicollis.....	1674	Saw-belly.....	426
872	ateniatus.....	1670	Sawfish, Common.....	60

	Page.		Page.
Saw-kwey.....	479	scapularis, Anisotremus.....	1320
Saw-qui Salmon.....	481	Tylosurus.....	711
saxatilis, A. hudeledni.....	1561	Scaridae.....	1572, 1620
Chaetodon.....	1562	Searnus.....	1621
Glyphidodon.....	1562	Scartella.....	2384
Gymnothorax ocellatus..	309	microstoma.....	2384
Johnius.....	1475	Scartes.....	2395
Lycodontis ocellatus.....	390	scartes, Fundul. s.....	654
Menticirrhus.....	1475	Scartichthys.....	2395
Perca.....	1133	rubropunctatus.....	2396
Platessa.....	2657	Searnus.....	1627, 1642, 1643, 1645
Percilichthys.....	1048	abildgaardi.....	1635
Roccus.....	1133	acutus.....	1632
saxicola, Pleuronectes.....	2657	alternans.....	1651
Sebastichthys.....	1790	amplus.....	1635
Sebastes.....	1798	aracanga.....	1642, 1647, 1648
say, Dusyatis.....	86	atomarius.....	1631
Raja.....	86	auroorbis.....	1635
sayanus, Aphredodorus.....	786	aurofrenatus.....	1634
Sceloporus.....	787	bollmani.....	1646
sayi, Dasibatis.....	86	brachialis.....	1641
Myliobatis.....	86	caerulens.....	1652, 1654
Trygon.....	86	catesbei.....	1638
Sayris.....	725	catesby.....	1638
bimaculatus.....	725	chloris.....	1637, 1640
hians.....	725	chrysopterus.....	1637
recurvirostris.....	725	circumnotatus.....	1641
serratus.....	726	coecineus.....	1635
Scabbard Fish.....	887, 889	collesinus.....	1656
scaber, Antennarius.....	2722	croicensis.....	1650
Centroponus.....	1125	euzanile.....	1648
Chironectes.....	2723	diadema.....	1646
Hexagrammus.....	1873	distinctus.....	1636
scabra, Trinectes.....	2701	emarginatum.....	1641
scabriceps, Mimus.....	268, 290	emblematicus.....	1654
Notropis.....	290	erythrinoides.....	1635
Photogenis.....	290	evermanni.....	1651
scabripinnis, Tetragonopterus.....	335	flavescens.....	1640
Sead.....	907	flavomarginatus.....	1652
Big-eyed.....	911	frondosus.....	1636, 1642
Mackerel.....	907	gnathodus.....	1650
Sealed Sardines.....	428	guacamaia.....	1656, 1658
Sealy-fins.....	1665	holocyanæos.....	1654
Seamp.....	1184, 1185	hoplomystax.....	1633
Scaphirhynchops platyrhynchus..	107	humeralis.....	1641
Scaphirhynchus.....	107	insula-sanctæ-crucis.....	1651
cataphractus.....	107	lacrimosus.....	1632
platyrhynchus.....	107	lateralis.....	1637
platyrhynchus.....	107	loro.....	1654
rafinesquei.....	107	maschalespilos.....	1642
scaphiurus, Opisthognathus.....	2282	melanotis.....	1638
scaphopsis, Caelorhynchus.....	2590	miniofrematus.....	1634
Maerurus (Caelorhynchus)	2591	nuchalis.....	1654
Scaphyrhynchops.....	107	obtusus.....	1654
scapulare, Pristipoma.....	1321	oxybrachius.....	1635
		perrico.....	1659





Page.		Page.		Page.
1320	<i>Scarus pleianus</i> .....	1658	<i>scheepii</i> , <i>Diodon</i> .....	1748
711	<i>psittacus</i> .....	1647	<i>scholaris</i> , <i>Thyrstites</i> .....	880
1572, 1620	<i>punctulatus</i> .....	1645	<i>schomburgkii</i> , <i>Pempheris</i> .....	978
1621	<i>quadrispinosus</i> .....	1648	<i>schoneveldii</i> , <i>Cataphractus</i> .....	2067
2384	<i>radians</i> .....	1632, 1633	Schoolmaster .....	1258
2384	<i>rostratus</i> .....	1658	<i>schranki</i> , <i>Hemulon</i> .....	1302, 1303
2395	<i>rubripinnis</i> .....	1640	<i>schumardi</i> , <i>Etheostoma</i> .....	1047
654	<i>sancti-crucis</i> .....	1651	Schnylkill Cat. ....	140
2305	<i>simplex</i> .....	1656	<i>Sciadeichthys</i> .....	119, 120, 122, 2757
2396	<i>spinidens</i> .....	1637	<i>albicans</i> .....	124, 2760
1643, 1645	<i>squalidus</i> .....	1640	<i>eniphysetus</i> .....	122, 2750
1635	<i>strigatus</i> .....	1639	<i>davescens</i> .....	123, 2760
1652	<i>superbus</i> .....	1650	<i>mesops</i> .....	123, 2760
1651	<i>taeniopterus</i> .....	1646	<i>passany</i> .....	124, 2760
1635	<i>triolabatus</i> .....	1654	<i>proops</i> .....	123, 2760
1647, 1648	<i>trispinosus</i> .....	1648	<i>temminckianus</i> .....	122, 2760
1631	<i>truncatus</i> .....	1641	<i>troschell</i> .....	122, 2757
1635	<i>turcesius</i> .....	1658	<i>Sciades troschell</i> .....	122, 2758
1634	<i>vetula</i> .....	1647, 1649	<i>sciadicus</i> , <i>Fundulus</i> .....	654
1646	<i>virens</i> .....	1640	<i>Haplochilus</i> .....	654
1641	<i>virginalis</i> .....	1647	<i>Zygonectes</i> .....	654
1652, 1654	<i>viridis</i> .....	1638	<i>Sciæna</i> .....	1454, 1465
1638	<i>scepticus</i> , <i>Mimulus</i> .....	296	<i>acuminata</i> .....	1488
1638	<i>Notropis</i> .....	296	<i>aluta</i> .....	1438
1637, 1640	<i>Triglops</i> .....	1925	<i>amazonica</i> .....	1419
1637	<i>Schedophilinae</i> .....	969	<i>caprodes</i> .....	1927
1641	<i>Schedophilopsis</i> .....	972	<i>chrysolenca</i> .....	1439
1635	<i>spinosus</i> .....	972	<i>chrysuræ</i> .....	1434
1656	<i>Schedophilus</i> .....	970	<i>coro</i> .....	1324
1650	<i>eugnatiens</i> .....	972	<i>croker</i> .....	1462
1648	<i>medusophagus</i> .....	970	<i>(Corvina) adusta</i> .....	1448
1646	<i>Schilbeodes</i> .....	144, 145, 146	<i>crouvina</i> .....	1419
1636	<i>eleutherus</i> .....	148	<i>deliciosa</i> .....	1455
1641	<i>exilis</i> .....	147	<i>edwardi</i> .....	1490
1654	<i>funebri</i> .....	147	<i>ericymba</i> .....	1445
1635	<i>furiosus</i> .....	149	<i>ensifera</i> .....	1435
1651	<i>gilberti</i> .....	148	<i>furthi</i> .....	1441
1640	<i>gyrius</i> .....	146, 2790	<i>fusca</i> .....	1483
1652	<i>insignis</i> .....	147	<i>gigas</i> .....	1483
636, 1642	<i>leptacanthus</i> .....	146	<i>grisea</i> .....	1484
1650	<i>minus</i> .....	148	<i>heterolepis</i> .....	1419
656, 1658	<i>nocturnus</i> .....	146	<i>icistia</i> .....	1436
1654	<i>Schistorus</i> .....	1148, 1151	<i>imberbis</i> .....	1451
1633	<i>mystacinus</i> .....	1151	<i>imiceps</i> .....	1451
1641	<i>schlegeli</i> , <i>Oreynus</i> .....	870	<i>jacobi</i> .....	1457
1651	<i>Præacanthus</i> .....	2858	<i>lanceolata</i> .....	1444
1632	<i>Sebastodes</i> .....	1834	<i>lineata</i> .....	1133, 1460
1637	<i>schmidti</i> , <i>Hoplunnis</i> .....	361	<i>macrops</i> .....	1428
1654	<i>schmittii</i> , <i>Balistes</i> .....	1705	<i>maculata</i> .....	2198
1642	<i>schneideri</i> , <i>Chauliodus</i> .....	585	<i>magdalene</i> .....	1420
1638	<i>Ophichthys</i> .....	387	<i>multifasciata</i> .....	1459
1634	<i>Pecilia</i> .....	691	<i>obliqua</i> .....	1459
1654	<i>scheepii</i> , <i>Alutera</i> .....	1718	<i>ocellata</i> .....	1454
1654	<i>Balistes</i> .....	1718	<i>opercularis</i> .....	1461
1635	<i>Ceratacanthus</i> .....	2860	<i>ophioscion</i> .....	1448
1659	<i>Chylomycterus</i> .....	1748	<i>oscitans</i> .....	1441





**IMAGE EVALUATION  
TEST TARGET (MT-3)**



**Photographic  
Sciences  
Corporation**

23 WEST MAIN STREET  
WEBSTER, N.Y. 14580  
(716) 872-4503

18  
20  
22  
25  
28  
32  
36

CT

	Page.		Page.
<i>Sciæna oscula</i> .....	1484	<i>Scomber balantiophthalmus</i> .....	911
<i>plumiferi</i> .....	1324	<i>bisus</i> .....	867
<i>punctata</i> .....	1434	<i>carangus</i> .....	920
<i>ronchus</i> .....	1436	<i>chloris</i> .....	938
<i>rubella</i> .....	1418	<i>chrysurus</i> .....	938
<i>rubra</i> .....	849	<i>collas</i> .....	866, 2843
<i>saturna</i> .....	1456	<i>crumenophthalmus</i> .....	911
<i>sciæra</i> .....	1452	<i>crysos</i> .....	921
<i>squamosissimus</i> .....	1418	<i>dekayi</i> .....	867
<i>stellifer</i> .....	1444	<i>dentex</i> .....	927
( <i>Stelliferus</i> ) <i>stellifer</i> .....	1443	<i>diego</i> .....	867
<i>surinamensis</i> .....	1420	<i>fasciatus</i> .....	904
<i>typica</i> .....	1448	<i>filamentosus</i> .....	932
<i>undecimalis</i> .....	1119	<i>germo</i> .....	871
<i>vermicularis</i> .....	1452, 1453	<i>gracilis</i> .....	867
<i>xanthurus</i> .....	1450	<i>grex</i> .....	867
<i>Sciænida</i> .....	1392	<i>guara</i> .....	927
<i>Sciænops</i> .....	1453	<i>gunneri</i> .....	955
<i>ocellatus</i> .....	1453	<i>heberi</i> .....	923
<i>sciæra</i> , <i>Sciæna</i> .....	1452	<i>hippos</i> .....	908, 920
<i>scierum</i> , <i>Etheostoma</i> .....	1038	<i>kreütereri</i> .....	900
<i>scierus</i> , <i>Hadropterus</i> .....	1037	<i>lacertus</i> .....	867
<i>serrula</i> .....	1038	<i>latus</i> .....	938
<i>Ophioscion</i> .....	1452	<i>macrophthalmus</i> .....	867
<i>Sciænga</i> .....	1180	<i>maculatus</i> .....	867, 874
<i>sciænga</i> , <i>Mycteroperca</i> .....	1181	<i>mediterraneus</i> .....	872
<i>scitnliceps</i> , <i>Synodus</i> .....	537, 2826	<i>niger</i> .....	848
<i>scitulus</i> , <i>Prionotus</i> .....	2157	<i>pelagicus</i> .....	952
<i>sciurus</i> , <i>Diplectrum</i> .....	1204	<i>pelamides</i> .....	869
<i>Hæmulon</i> .....	1303	<i>pelamis</i> .....	869
<i>Serranus</i> .....	1204	<i>pelamitus</i> .....	872
<i>Sparus</i> .....	1304	<i>pelamys</i> .....	872
<i>Sclæniæna</i> .....	1394	<i>plumieri</i> .....	911
<i>Sclerodermi</i> .....	781, 1697	<i>pneumatophorus</i> .....	867
<i>Sclerognathus</i> .....	163	<i>quadripunctatus</i> .....	869
<i>cyprinella</i> .....	164	<i>regalis</i> .....	875
<i>elongatus</i> .....	160	<i>rim</i> .....	928
<i>meridionalis</i> .....	164	<i>rochei</i> .....	867
<i>urus</i> .....	165	<i>ruber</i> .....	919
<i>scofieldi</i> , <i>Stolephorus</i> .....	2814	<i>saliens</i> .....	899
<i>Sculeosoma</i> .....	10	<i>sarda</i> .....	872
<i>Scoliodon</i> .....	42	<i>saurus</i> .....	898
<i>longurio</i> .....	42, 2748	<i>scombrus</i> .....	865
<i>terræ-novæ</i> .....	43	<i>sloanci</i> .....	870
<i>scolopacea</i> , <i>Nemichthys</i> .....	369	<i>speciosus</i> .....	928
<i>scolopaceus</i> , <i>Nemichthys</i> .....	369	<i>thazard</i> .....	867
<i>scolopax</i> , <i>Ballates</i> .....	759	<i>thynnus</i> .....	870
<i>Centriscaus</i> .....	759	<i>trachurus</i> .....	910
<i>Macrorhamphosus</i> .....	759	<i>undulatus</i> .....	867
<i>Scolopsis saianus</i> .....	787	<i>vernalis</i> .....	866
<i>Scomber</i> .....	865	<i>zonatus</i> .....	902
<i>adacensionis</i> .....	927	<i>scomberius</i> , <i>Esox</i> .....	926
<i>alatunga</i> .....	871	<i>Scomberodon</i> .....	873
<i>albacoës</i> .....	870	<i>Scomberoides</i> , <i>saltator</i> .....	899
<i>alleteratus</i> .....	809	<i>Coryphæna</i> .....	953
<i>ascensionis</i> .....	925	<i>Scomberomorus</i> .....	873, 4843

Page.		Page.		Page.
911	<i>Scomberomorus caballa</i> .....	875, 876	<i>scopifer</i> , <i>Phenacobius</i> .....	303
867	<i>concolor</i> .....	873	<i>scopiferum</i> , <i>Sarcidium</i> .....	303
920	<i>maculatus</i> .....	874, 875, 4843	<i>scopiferus</i> , <i>Phenacobius</i> .....	303
938	<i>plumieri</i> .....	875	<i>scops</i> , <i>Gnathypops</i> .....	2283
938	<i>regalis</i> .....	875	<i>Scorfanudi Funal</i> .....	1837
66, 2843	<i>sierra</i> .....	874	<i>Scorpaena</i> .....	1839
911	<i>Scombresocidae</i> .....	724	<i>africana</i> .....	1833
921	<i>Scombresox</i> .....	725	<i>agassizii</i> .....	1840, 2860
867	<i>brevirostris</i> .....	726	<i>americana</i> .....	2023
927	<i>camperi</i> .....	725	<i>brasiliensis</i> .....	1842
867	<i>equirostrum</i> .....	726	<i>bufo</i> .....	1849
904	<i>fosteri</i> .....	726	<i>calcarata</i> .....	1854
932	<i>rondeleti</i> .....	726	<i>capensis</i> .....	1833
871	<i>saurus</i> .....	725	<i>castor</i> .....	1856
867	<i>scutellatum</i> .....	726	<i>crisolata</i> .....	1841
867	<i>storeri</i> .....	726	<i>dactyloptera</i> .....	1837
927	<i>Scombrida</i> .....	863	<i>granlicornis</i> .....	1850
955	<i>Scombrina</i> .....	864	<i>guttata</i> .....	1847
923	<i>scombrinus</i> , <i>Caranx</i> .....	908	<i>histro</i> .....	1843, 1846
908, 920	<i>Decapterus</i> .....	908	<i>inermis</i> .....	1853, 2861
900	<i>Scorbrocotus</i> .....	1861	<i>massiliensis</i> .....	1139
867	<i>salmoneus</i> .....	1862	<i>mystes</i> .....	1849
938	<i>Scombroidel</i> .....	781, 860	<i>nematophthalmus</i> .....	2861
867	<i>Scombroides occidentalis</i> .....	890	<i>occipitalis</i> .....	1854
867, 874	<i>Scombroidina</i> .....	896	<i>pannosa</i> .....	1845
872	<i>Scombroptina</i> .....	1106	<i>plumieri</i> .....	1848
848	<i>Scombrops</i> .....	1114	<i>porca</i> .....	1839
952	<i>oculatus</i> .....	1114	<i>rascacio</i> .....	1849
869	<i>scombrus</i> , <i>Scomber</i> .....	865	<i>russula</i> .....	1851
869	<i>Scopelus</i> .....	569	<i>sierra</i> .....	1860
872	<i>affinis</i> .....	571	<i>sonora</i> .....	1852
872	<i>andreae</i> .....	569	<i>stearnsi</i> .....	1843
911	<i>aretiens</i> .....	574	<i>Scorpenichthys</i> .....	1880
867	<i>benoiti</i> .....	573	<i>Scorpenichthys</i> .....	1880
869	<i>bonaparti</i> .....	557	<i>lateralis</i> .....	1902
875	<i>boops</i> .....	572	<i>marmoratus</i> .....	1889
928	<i>borealis</i> .....	577	<i>Scorpenidae</i> .....	1758
867	<i>caninianus</i> .....	570	<i>Scorpeninae</i> .....	1759
919	<i>caudispinosus</i> .....	556	<i>Scorpeno</i> .....	1847
899	<i>coccoi</i> .....	569	<i>scorpio</i> , <i>Cottus</i> .....	1973
872	<i>crassiceps</i> .....	843	<i>scorpioides</i> , <i>Cottus</i> .....	1973
898	<i>crocodilus</i> .....	558	<i>Myoxocephalus</i> .....	1973
865	<i>elongatus</i> .....	555	<i>Scorpion</i> .....	1847
870	<i>gracilis</i> .....	572, 574	<i>Fishes</i> .....	1839
928	<i>huraboldti</i> .....	572, 577	<i>Scorpius californensis</i> .....	1391
867	<i>hygomi</i> .....	573	<i>Scorpius virginianus</i> .....	1976
870	<i>kroyeri</i> .....	556	<i>scorpius</i> , <i>Cottus</i> .....	1974
910	<i>madelrensis</i> .....	557	<i>Cottus grenlandicus</i> .....	1975
867	<i>maurolei</i> .....	577	<i>scouteri</i> , <i>Oncorhynchus</i> .....	478
866	<i>mulleri</i> .....	570, 574	<i>Salmo</i> .....	478, 481
902	<i>pseudocrocodilus</i> .....	556	<i>Scour Fish</i> .....	879
626	<i>rafinesquel</i> .....	567	<i>scovelli</i> , <i>Etheostoma</i> .....	1082
873	<i>rarus</i> .....	569	<i>Siphostoma</i> .....	769
899	<i>spinosus</i> .....	575	<i>scripta</i> , <i>Alutera</i> .....	1719
953	<i>tenorei</i> .....	577	<i>scriptus</i> , <i>Balistes</i> .....	1719
73, 4843	<i>scopifer</i> , <i>Notropis</i> .....	291	<i>Ceratacanthus</i> .....	2860

	Page.		Page.
scriptus, Gymnotherax	398	Sea Bass	1126
Monacanthus	1719	Bats	2736
Pseudoscopolus	2292	Catfish	118, 119, 128
scutator, Belone	714	Devil	91, 92, 2727
Iscæthes	2389	Drums	1482
scudderi, Diabasis	1300	Mink	1475
Hæmulon	1299, 1300	Poacher	2031, 2065, 2091
Sculpin	1847	Raven	976, 2622, 2023
Arctic	1973	Serpent	384
Black	1985	Snail	2105, 2114, 2116
Daddy	1974	Snipe	714
European	1974	Trout	1407
Great	1976	Spotted	1409
Long-spined	1976	Sea-horse	775
Red	1935	Common American	777
Yellow	1934	Sea Robin, Brown-winged	2167
Sculpins	1879	Red-winged	2156
Great	1970	sebago, Salmo salar	487
Spineless	2025	Sebastapistes	1839
Stone	1937	guttatus	1848
Scup, Common	1346	Sebastes	1760
Scuppaug	1346	auriculatus	1818
scutata, Echeueis	2371	capensis	1833
scutellatum, Scombrosox	728	caurinus	1821
Scutica	403, 404	dactylopterus	1837, 1838
scuticaris, Bascanichthys	378	darwinii	1832
Cœcula	379	elegans	1830
Sphagebranchus	379	elongatus	1816
scutiger, Icelus	1910	fasciatus	1761, 1827
Rastrinus	1909	helvomaculatus	1808
scutum, Achirus	2700	imperialis	1837, 1838
Solea	2700	inermis	1829
scylla, Exocætes	735	marinus	1760, 1761
Hybopsis	263	viviparus	1761
Notropis	263	melanops	1783
Scylliorhinide	22	neumatophthalmus	2861
Scylliorhinine	22	nigrocinctus	1828
Scylliorhinus	22	nivosus	1834
profundorum	22	norwegicus	1761
retifer	25	oblongus	1890
Scyllium	22	oculata	1832
retiferum	25	paucispinis	1781
ventriosum	25	regulus	1761
Scoymnoid Sharks	56	rosaceus	1794
Scymnus brevipinna	57	ruber	1818
Seyphius	774	septentrionalis	1761
Seyris	931	ateludachneri	1830
Seyris analis	932	taczanowskii	1832
Seytalichthys	387	variabilis	1784
minor	387	ventricosus	1820
Scytalina	2454	viviparus	1761
cerdale	2454	Sebastichthys	1765, 1777, 1827
Scytalinide	2453	atrovirens	1798
Scytaliscus	2454	aurora	1803
Scytalopsis	381, 384	brevispinis	1788
magniloculis	385	carnatus	1825

Page.		Page.		Page.
	Sebastichthys chlorostictus	1812	Sebastodes glaucus	1777
1126	chrysomelas	1826	goodiei	1779
2736	constellatus	1807	hopkinsi	1780, 2860
3, 119, 128	diploproa	1802	incermis	1829
1, 92, 2727	entomelas	1786	introniger	1805
1482	fasciolaris	1827	jordani	1778
1475	flavidus	1782	joyneri	1829
2065, 2091	goodiei	1780	levis	1816
2022, 2023	introniger	1805	macdonaldi	1786
384	levis	1816	maliger	1822
2114, 2116	maliger	1823	matsubarae	1796, 1833, 2860
714	miniatus	1795	melanops	1782, 1783
1407	mystinus	1785	melanostomus	1803
1409	nigrocinctus	1828	miniatus	1794
775	ovalis	1789	mitzukurii	1831
777	pinniger	1794	mystinus	1784, 1785
2167	proriger	1788, 1793	nebulosus	1826
2156	purpureus	1826	nigrocinctus	1827
487	rastrelliger	1820	nivosus	1833
1839	rhodochloris	1810	oblongus	1830
1848	rubrivinctus	1817	oculatus	1832
1760	rupestris	1813	ovalis	1788
1818	saxicola	1799	paucispinis	1780
1833	serriceps	1827	pinniger	1798, 1794
1821	sinensis	1814	proriger	1787, 1792
1837, 1838	umbrosus	1807	rastrelliger	1810, 1820
1832	vexillaris	1822	rhodochloris	1809
1830	Sebastina	1759, 1771	rosaceus	1808
1816	Sebastodes	1765, 1773, 1778	ruher	1806
1761, 1827	ureus	1807	ruberrimus	1805, 1806
1808	aleutianus	1795	rubrivinctus	1817
1837, 1838	alutus	1790	rufus	1786, 2860
1829	atrorubens	1790, 2860	rupestris	1812
1760, 1761	atrovirens	1797, 2860	saxicola	1798
1761	auriculatus	1817, 1818	sehlegelii	1834
1783	dallii	1818	semicinctus	1800
2861	aurora	1802	serranoides	1782
1828	ayresii	1808	serriceps	1827
1834	brevispinis	1787	sinensis	1813
1761	capensis	1833	steindachneri	1830
1830	carnatus	1824	taczanowskii	1831, 2860
1832	caurinus	1821	trivittatus	1834
1781	chlorostictus	1811	umbrosus	1807
1761	chrysomelas	1825, 1826	ventricosus	1829
1794	ciliatus	1783, 1784	vexillaris	1821
1818	constellatus	1806	vulpes	1835
1761	cramerii	1790, 2860	zacentrus	1814, 2860
1830	darwini	1832	Sebastolobus	1761
1832	diploproa	1801	ulascanus	1761
1784	eigenmanni	1789	ultivelis	1763
1829	elegans	1830	macrochir	1763
1761	elongatus	1815	Sebastomus	1765, 1775, 1805
35, 1777, 1827	entomelas	1785	Sebastoplus	1854
1798	eos	1810	dactylopterus	1837
1803	flavidus	1781	Sebastopais	1835
1788	gilberti	1823	xyris	1835
1825	gilli	1811		



	Page.		Page.
Sebastosomus .....	1765, 1774, 1781	sellicauda, Epinephelus .....	1155
pinniger .....	1794	selfer, Halicheres .....	1592
simulans .....	1783	Iridio .....	1592
seco, Luxilus .....	250	sem, Caranx .....	923
Sectator .....	1389	Sema .....	1498
ocyurus .....	1389	signifer .....	1490
sectatrix, Kyphosus .....	1387	semimarmatus, Gasterosteus .....	747
Perca .....	1388	semicinctus, Gilletus .....	2298
marina .....	1388	Halicheres .....	1593
secundo-dorsalis, Thynnus .....	876	Iridio .....	1592
secundus, Carangops .....	914	Jalis .....	1593
Caranx .....	914	PlatyGLOSSUS .....	1593
Hemicaranx .....	914	Sebastodes .....	1806
sedentarius, Chaetodon .....	1675	semicoronata, Seriola .....	904
Sarothrodus .....	1675	Semicossyphus pulcher .....	1585
seemanni, Arius .....	128, 2772	semifasciatum, Pileoma .....	1027
Galeichthys .....	2772	Triakis .....	31
Hexanematichthys .....	128	semifasciatus, Serranus .....	1197
Tachlaurus .....	129	semiloricatus, Gasterosteus .....	747
segalienis, Brachyopsis .....	2048	semiluna, Sparus .....	1276
Siphagonus .....	2048	seminolis, Fundulus .....	647
Syngnathus .....	2048	seminuda, Albulia .....	412
Segundo .....	914	Eleotris .....	2204
Selache .....	51	Garmannia .....	2233
maxima .....	51	Gila .....	228
Selachii .....	15	Gymneleotris .....	2204
selachops, Apterichthys .....	374	seminudus, Gobius .....	2234
Ichthyapus .....	374	Gymneleotris .....	2204
Sphagobranchus .....	374	Lycodes .....	2468
Selachostomi .....	100	semiradiatus, Lepisosteus .....	110
Selachus maximus .....	51	semiruber, Labrus .....	1583
Selanonius .....	49	semiscaber, Cottopsis .....	1950
walkeri .....	49	Cottus .....	1949
selauonus, Squalus .....	49	semiscabra, Cottus centroleura .....	1945
Selar .....	916, 918	Uranidea .....	1950
Selaroides .....	916	semispinosus, Caranx .....	911
Selenaspis .....	119	Semitapicis .....	332
Selenaspis .....	120, 124, 2760	Semotilus .....	220, 221, 222
dowii .....	125	atromaculatus .....	222
dowii .....	2761	thoreauianus .....	223
herzbergii .....	124, 2760	biguttatus .....	322
luniscutis .....	125	bullaris .....	222
parkeri .....	125, 2764	cephalus .....	222
Selene .....	935	corporalis .....	221, 222
argentea .....	936	diplemius .....	222
teratodii .....	935	dorsalis .....	222
quadrangularis .....	1068	haumondi .....	222
setiplinnis .....	934	macrocephalus .....	222
vomer .....	936	notatus .....	659
selene, Carpiodes .....	107	spectosus .....	222
Luxifas .....	260	thoreauianus .....	223
Minnilus .....	260	senegalensis, Vomer .....	934
Notropis hudsonius .....	260	senilis, Gambusia .....	682
selenops, Hiodon .....	414	Sennet .....	826
sellaria, Acanthoecottus .....	1998	Señorita .....	1592, 1601, 2352
Porocottus .....	1096		

Page.		Page.		Page.
1155	septa, Raja.....	71	serra, Alepisaurus.....	597
1592	septentrionalis, Gaidropsarus.....	2559	Gonenion.....	947
1592	Motella.....	2560	Serran Imperial.....	1837
923	Onos.....	2560	Serrana.....	1489, 1490
1498	Pera.....	1133	Hispana.....	1488
1499	Sebastes.....	1761	hispanis.....	1489
747	septipinnis, Ammodytes.....	2842	Serranidn.....	1126
2298	Rhynchias.....	2841	Serraninae.....	1129
1593	Septogunnellus gracilis.....	2436	Serrano.....	1207
1592	serena, Dionda.....	214	serranoides, Sebastodes.....	1782
1593	Hybognathus.....	214	Serranos.....	2208
1593	Sergeant Fish.....	947, 948	Serranus neanthophorus.....	1196
1806	Major.....	1561	acutirostris.....	1181
904	Seriola.....	901	aequidens.....	1211
1585	argyromelas.....	950	agassizii.....	1189
1027	bipinnulata.....	907	albomaculatus.....	1197
31	bonariensis.....	905	angustifrons.....	1159
1197	bosci.....	905	annularis.....	1214
747	coronata.....	905	apua.....	1158
1276	cosmopolita.....	938, 2847	arara.....	1159, 1175
647	declivis.....	905	armatus.....	1165
412	dorsalis.....	902	aspersus.....	1153
2204	dubia.....	905	atrinus.....	1200
2233	dumerili.....	903, 904	auratus.....	1145
228	dussumieri.....	900	auriga.....	1221
2204	fulcata.....	905	bivittatus.....	1205
2234	fasciata.....	904	bonaci.....	1175
2204	gigas.....	903	brasiliensis.....	1221
2468	halandi.....	902, 903	brunneus.....	1175
110	ligulata.....	905	bulleri.....	1214
1583	mazzatlanana.....	904	calopteryx.....	1213
1950	picturata.....	910	capenna.....	1311
1949	pinnulata.....	907	carana.....	1146
1945	proxima.....	904	cardinalis.....	1174
1950	rivoliiana.....	904	camelopardulis.....	1187
911	semicomata.....	904	catus.....	1159
332	stearnsii.....	903	chlorurus.....	1193
220, 221, 222	succincta.....	900	clathratus.....	1198
222	zonata.....	902	celonus.....	1222
223	carolinensis.....	902	conspersus.....	1154
223	Seriolichthys.....	906	coronatus.....	1142
322	Seriolina.....	896	conchii.....	1139
222	Seriolophus.....	895	courtaiei.....	1152
222	carangoides.....	895	creolus.....	1222
221, 222	Seriphus.....	1397	cycloponatus.....	1175
222	politus.....	1397	decimalis.....	1175
222	serotinus, Acipenser.....	106	dimidiatus.....	1179
222	serpens, Genopylus.....	884	dubius.....	1146
222	Serpens marinus maculosus.....	382	emarginatus.....	1181
659	Serpent, Sea.....	384	erythrogaster.....	1100
222	serpentina, Muræna.....	348	falentus.....	1185
223	serpentinus, Blennius.....	2439	fascicularis.....	1208
934	Derichthys.....	343	felinus.....	1187
682	Leptoblennius.....	2439	fimbriatus.....	1154
826	Serra.....	597	flaviventris.....	1221
2, 1601, 2352	serra, Alepidosaurus (Caulopus)...	597	furcifer.....	1222

	Page.		Page.
<i>Serranus furvus</i> .....	1200	<i>Serranus subligarius</i> .....	1219
<i>fuscus</i> .....	1211	<i>tabacarius</i> .....	1215
<i>fuscus</i> .....	1181	<i>teniops</i> .....	1144
<i>galeus</i> .....	1164	<i>tigrinus</i> .....	1214
<i>glgas</i> .....	1154	<i>tigris</i> .....	1187
<i>guasa</i> .....	1164	<i>tinea</i> .....	1181
<i>gnativero</i> .....	1145	<i>trifurcus</i> .....	1201, 1202
<i>humeralis</i> .....	1197	<i>undulosus</i> .....	1181
<i>impetiginosus</i> .....	1153	<i>unicolor</i> .....	1192
<i>inermis</i> .....	1168	<i>varius</i> .....	1153
<i>interstitialis</i> .....	1170	<i>Serraria</i> .....	1028, 1030, 1037
<i>irradians</i> .....	1208	<i>serrata, Fistularia</i> .....	758
<i>itaiara</i> .....	1164	<i>serraticornis, Balistes</i> .....	1720
<i>jacome</i> .....	1215	<i>serratogramulata, Perca</i> .....	1024
<i>labriformis</i> .....	1155	<i>serratum, Hæmulon</i> .....	1299
<i>lampyrus</i> .....	1190	<i>serratus, Gasterosteus</i> .....	750
<i>latepietus</i> .....	1175	<i>Nauerates</i> .....	900
<i>lucopecanus</i> .....	1216	<i>Sayris</i> .....	726
<i>lumulatus</i> .....	1150	<i>serriceps, Sebastichthys</i> .....	1827
<i>maculatofasciatus</i> .....	1196	<i>Sebastodes</i> .....	1827
<i>maculatus</i> .....	1153	<i>serrifer, Conodon</i> .....	1324
<i>maculosus</i> .....	1159	<i>Serrivomer</i> .....	367
<i>margaritifer</i> .....	1156	<i>beanii</i> .....	367
<i>marginatus</i> .....	1154	<i>serrula, Anisotremus</i> .....	1323
<i>mentzeli</i> .....	1154	<i>Chalinura</i> .....	2576
<i>morio</i> .....	1160	<i>Hadropterus sclerus</i> .....	1638
<i>mystacinus</i> .....	1151	<i>Priacanthus</i> .....	1239
<i>nebulifer</i> .....	1196	<i>Pristipoma</i> .....	1324, 1343
<i>nigrescens</i> .....	1200	<i>Pseudopriacanthus</i> .....	1239
<i>nigriculus</i> .....	1153	<i>Seserinus xanthurus</i> .....	966
<i>nigritus</i> .....	1161	<i>Seşi de lo Alto</i> .....	1261
<i>niventus</i> .....	1156	<i>sessilicauda, Monolene</i> .....	2691
<i>oculatus</i> .....	1283	<i>Setarches</i> .....	1860
<i>oeyurus</i> .....	1201	<i>parmatus</i> .....	1860
<i>olfax</i> .....	1183	<i>setifer, Argyriosus</i> .....	936
<i>ongus</i> .....	1154	<i>Monacanthus</i> .....	1716
<i>ouantalibi</i> .....	1146	<i>Stephanolepis</i> .....	1716
<i>panamensis</i> .....	1141	<i>setiger, Dasycottus</i> .....	1991
<i>philadelphicus</i> .....	1202	<i>setigerus, Lophionus</i> .....	2714
<i>phæbe</i> .....	1212	<i>Lophius</i> .....	2715
<i>pixanga</i> .....	1153	<i>setinotus, Chauliodus</i> .....	585
<i>prestigiator</i> .....	1214	<i>setipinnis, Argyreiosus</i> .....	934
<i>psittacinus</i> .....	1213	<i>Caranx dorsalis</i> .....	934
<i>quinquefasciatus</i> .....	1164	<i>gabonensis</i> .....	935
<i>radialis</i> .....	1205	<i>Selene</i> .....	934
<i>radians</i> .....	1208	<i>Vomer</i> .....	934
<i>remotus</i> .....	1160	<i>Zeus</i> .....	934
<i>repandus</i> .....	1187	<i>setosus, Mugil</i> .....	815
<i>rivulatus</i> .....	1187	<i>Ovoides</i> .....	1739
<i>ruber</i> .....	1181	<i>Tetraodon</i> .....	1740
<i>rupestris</i> .....	1174	<i>seu conger, Muræna brasiliensis</i> ...	403
<i>scirtus</i> .....	1204	<i>sexcornutus, Ostracion</i> .....	1725
<i>semifasciatus</i> .....	1107	<i>sexdecemlamellata, Echenis</i> .....	2272
<i>stadthouderi</i> .....	1159	<i>sexfasciatum, Hæmulon</i> .....	1294
<i>stillbostigma</i> .....	1217	<i>sexfasciatus, Hæmulon</i> .....	1295
<i>striatus</i> .....	1157	<i>sexmaculatus, Diodon</i> .....	1746

Page.		Page.		Page.
1210	<i>sexradiatus</i> , <i>Polynemus</i> .....	2183	Sharks, Requiem .....	27
1215	Shad .....	427	Sand .....	46
1144	Alabama .....	2810	Scymnoid .....	56
1214	American .....	427	Thresher .....	45
1187	Broad .....	1372	True .....	21
1181	Common .....	427	Typical .....	19
1201, 1202	Gulf .....	2810	W hale .....	52
1181	Hickory .....	416, 425	Sharp-nosed Flying-fish.....	728
1192	Mackerel .....	909	Puffers .....	1740
1153	Potomac .....	427	Shark .....	43
1030, 1037	Shad Porgy .....	1355	Sharp-tailed Goby .....	2329
758	Shad-waiter .....	405	shasta, <i>Cottus</i> .....	1047
1720	Shade, Gizzard .....	415	<i>Salmo gairdneri</i> .....	502
1024	Shark, Bay .....	37	<i>iridens</i> .....	502
1299	Bone .....	51	shavianus, <i>Cetorhinus</i> .....	51
750	Bullhead.....	20	Sheepshead .....	1358, 1361
900	Cat .....	31	Minnow .....	671
726	Cow .....	19	Porgy .....	1354
1827	Dusky .....	35	Sheepshead, Lake .....	1484
1827	Elephant.....	51	Shellfish .....	1723
1324	Great Blue .....	33	Shellfish, Rock .....	1722
367	White.....	50	Shi Shidai.....	1665
367	Hammer-headed .....	45	Shidai, Shi .....	1665
1323	Leopard .....	31	Shima Soi .....	1834
2576	Long-tail .....	45	Sh'ner .....	269, 281
1638	Mackerel .....	48	Blunt-nosed .....	934
1239	North liver .....	427	Golden .....	250
1324, 1343	Nurse .....	26	Red-sided .....	240
1239	Oil .....	32	Spotted .....	318
966	Sand .....	47	Shiners .....	254
1261	Sharp-nosed .....	43	Short-nosed Bat-Fish .....	2738
2691	Shovel-head .....	44	Gar .....	110
1860	Shovel-nosed .....	18	Sturgeon .....	106, 107
1860	Sleeper .....	57	shufeldti, <i>Ceratius</i> .....	2731
936	Soup-fin .....	32	<i>Gobius</i> .....	2221
1716	Swell .....	25	<i>Mancalias</i> .....	2730
1716	Tiger .....	32	<i>Typhlopsaras</i> .....	2731
1991	Shark Pilot .....	902	shumardi, <i>Alburnops</i> .....	268
2714	Sucker .....	2260	<i>Boleosoma</i> .....	1047
2715	Sharks .....	15	<i>Cottogaster</i> .....	1046
585	Angel .....	58	<i>Hadropterus</i> .....	1047
934	Basking .....	50	<i>Imostoma</i> .....	1047
934	Blue .....	33	<i>Minnilus</i> .....	268, 269
935	Bramble .....	57	<i>Notropis</i> .....	268
934	Bullhead .....	19	shuswap, <i>Agosia</i> .....	313
934	Cat .....	22	<i>falcata</i> .....	313
934	Cestraciant .....	19	sialis, <i>Argentina</i> .....	526
815	Cow .....	17	sialis, <i>Corvula</i> .....	1428
1739	Cyclospondylous .....	52	<i>Noturus</i> .....	146
1740	Dog .....	28	sibbaldi, <i>Syngnathus</i> .....	774
403	Fridged .....	16	Siboma .....	228, 231
1725	Hammer-headed .....	43	<i>atraria</i> .....	233
2272	Mackerel .....	47	<i>crassicauda</i> .....	231
1294	Man-eater .....	50	<i>longiceps</i> .....	233
1205	Notidanoid .....	16	sicaua, <i>Cerna</i> .....	1162
1746	Nurse .....	25	siceifer, <i>Holocentrus</i> .....	849

	Page.		Page.
sicciferum, <i>Holocentrum</i> .....	850	signatus, <i>Bathymaster</i> .....	2288
sicculum, <i>Chirostoma</i> .....	806	<i>Hypoprion</i> .....	41
sicculus, <i>Labidesthes</i> .....	805	<i>Microgobius</i> .....	2246
siculus, <i>Lampugnus</i> .....	953	signifer, <i>Bryttus</i> .....	996
Sicya .....	2207, 2867	<i>Chatoossus</i> .....	433
Sicyases .....	2329, 2330, 2336	<i>Coregonus</i> .....	518
<i>carneus</i> .....	2337	<i>Monacanthus</i> .....	1716
<i>fasciatus</i> .....	2338	<i>Saimo (Thymallus)</i> .....	518
<i>punctulatus</i> .....	2338	<i>Sema</i> .....	1499
<i>rubiginosus</i> .....	2337	<i>Stypodon</i> .....	220
<i>rupestris</i> .....	2341	<i>Thymallus</i> .....	517, 2871
Sicydiinae .....	2190	<i>lewisi</i> .....	2871
Sicydium .....	2205	<i>montanus</i> .....	519
<i>antillarum</i> .....	2206, 2867	<i>ontariensis</i> .....	519
<i>gymnogaster</i> .....	2208	<i>tricolor</i> .....	519
<i>plumieri</i> .....	2206, 2867	sigolutes, <i>Gilbertina</i> .....	2028
<i>punctatum</i> .....	2867	<i>silenus, Zaprora</i> .....	2850
<i>salviui</i> .....	2208	Silk Snapper .....	1262, 2858
<i>viragus</i> .....	2206	<i>Siluridu</i> .....	115
<i>vincente</i> .....	2207	<i>Silurus bagre</i> .....	117
Sicyogaster .....	2229	<i>catus</i> .....	138
Sicyopterus <i>gymnogaster</i> .....	2208	<i>clarus</i> .....	155
<i>salvini</i> .....	2208	<i>cornutus</i> .....	750
Sicyosus .....	2837	<i>cupreus</i> .....	140
Sidera .....	392	<i>felis</i> .....	128
<i>castaneu</i> .....	306, 2804	<i>furcatus</i> .....	140
<i>chlevastes</i> .....	399	<i>gyrinus</i> .....	146
<i>dovij</i> .....	397	<i>herzbergii</i> .....	125
<i>funebria</i> .....	396	<i>limosus</i> .....	143
<i>merdax</i> .....	396	<i>lividus</i> .....	140
<i>moringa</i> .....	395	<i>marinus</i> .....	118
<i>nigromarginata</i> .....	400	<i>melas</i> .....	141
<i>ocellata</i> .....	399	<i>nebulosus</i> .....	143
<i>panamensis</i> .....	391	<i>olivaris</i> .....	143
<i>pfeifferi</i> .....	2805	<i>parkeri</i> .....	126
<i>verrilli</i> .....	394	<i>punctatus</i> .....	135
<i>vicina</i> .....	394	<i>viscosus</i> .....	143
<i>siderea, Muræna</i> .....	2805	<i>xanthocephalus</i> .....	141
<i>siderium, Zophendum</i> .....	314	<i>Silus</i> .....	525
<i>siderius, Hyborhynchus</i> .....	314	<i>ascanii</i> .....	526
<i>sieboldii, Cichlasoma</i> .....	1510	<i>silus, Argentina</i> .....	526
<i>Heros</i> .....	1517	<i>Silver Chub</i> .....	221, 320
Sierra .....	874, 875	<i>Hake</i> .....	2530
Sierra, <i>Pez</i> .....	60	<i>Jenny</i> .....	1370
sierra, <i>Pontinus</i> .....	1859	<i>Salmon</i> .....	480
<i>Scomberomorus</i> .....	874	<i>Trout</i> .....	493
<i>Scorpena</i> .....	1860	<i>Whiting</i> .....	1477
Sierrita .....	713	<i>Silver-fn</i> .....	278
sierrita, <i>Tylosurus</i> .....	713	<i>Silver-fish</i> .....	409, 795, 889
Signistes .....	2863	<i>Silverside</i> .....	800
<i>caulias</i> .....	2863	<i>Brook</i> .....	805
Sigmops .....	581, 582	<i>Silver-sided Minnow</i> .....	238
<i>stigmaticus</i> .....	583	<i>Silversides</i> .....	788, 796
Sigmurus .....	1446, 1447, 1452	<i>Silvery Anchovies</i> .....	439
Signalosa .....	2809	<i>Lamprey</i> .....	11
<i>atchafalaye</i> .....	2809	<i>Minnow</i> .....	213

Page.		Page.		Page.
2288	<i>sima</i> , <i>Cliola</i> .....	287	<i>Siphostoma</i> <i>elucens</i> .....	768
41	<i>Eleotris</i> .....	2198	<i>fistulatum</i> .....	765
2246	<i>Simencheilyda</i> .....	348	<i>flavirostre</i> .....	768
996	<i>Simencheyls</i> .....	340	<i>florida</i> .....	766
433	<i>parasiticus</i> .....	349	<i>fuscum</i> .....	770
518	<i>similis</i> , <i>Amphistichus</i> .....	1504	<i>griseolineatum</i> .....	764
1716	<i>Fundulus</i> .....	638	<i>jonesi</i> .....	768, 2837
518	<i>Hemulon</i> .....	1304	<i>leptorhynchum</i> .....	764
1409	<i>Hydrargyra</i> .....	639	<i>lina</i> .....	768
220	<i>simillima</i> , <i>Palometa</i> .....	2849	<i>louisiana</i> .....	770
17, 2871	<i>simillimus</i> , <i>Poronotus</i> .....	967	<i>mackayi</i> .....	766
2871	<i>Rhombus</i> .....	967	<i>marmoratum</i> .....	768
519	<i>Stromateus</i> .....	967	<i>pelagicum</i> .....	767, 2837
519	<i>simotera</i> , <i>Ulocentra</i> .....	1051	<i>plecturatum</i> .....	768
519	<i>simotrum</i> , <i>Ilyostomus</i> .....	1051	<i>poeyi</i> .....	766
2028	<i>simplicis</i> , <i>Aplurus</i> .....	880	<i>punctipinne</i> .....	763
2850	<i>Pseudoscarus</i> .....	1656	<i>robertsi</i> .....	2837
262, 2858	<i>Scarus</i> .....	1656	<i>rousseau</i> .....	767, 2837
115	<i>Tetragonurus</i> .....	880	<i>scovelli</i> .....	769
117	<i>simpsoni</i> , <i>Ictalurus</i> .....	135	<i>sinaloe</i> .....	2838
138	<i>simula</i> , <i>Chalinura</i> .....	2578	<i>starksi</i> .....	771, 2838
155	<i>simulans</i> , <i>Enneacanthus</i> .....	994	<i>zatropis</i> .....	772
759	<i>Hemiplites</i> .....	994	<i>siragus</i> , <i>Sicydium</i> .....	2206
140	<i>Sebastosomus</i> .....	1783	<i>Sirajo</i> .....	2206
128	<i>simulus</i> , <i>Macrurus</i> .....	2578	<i>Siremboguntheri</i> .....	2523
140	<i>Ophioscion</i> .....	1449	<i>sisco</i> , <i>Argyrosomus</i> <i>artedi</i> .....	469
146	<i>simus</i> , <i>Alburnellus</i> .....	627	<i>Sisco</i> of Lake Tippecanoe.....	469
125	<i>Careproctus</i> .....	2131	<i>siscowet</i> , <i>Cristivomer</i> <i>namayensh</i> ..	505
143	<i>Menticirrhus</i> .....	1472	<i>Salmo</i> .....	505
140	<i>Notropis</i> .....	267	<i>Salvelinus</i> <i>namayensh</i> ...	505
118	<i>Rhinichthys</i> .....	307	<i>siskawitz</i> , <i>Salmo</i> .....	505
141	<i>sinaloe</i> , <i>Paralichthys</i> .....	2872	<i>siuslawi</i> , <i>Leuciscus</i> .....	2797
143	<i>Siphostoma</i> .....	2838	<i>Skate</i> , <i>Barndoor</i> .....	71
143	<i>Umbrina</i> .....	1438	<i>Big</i> .....	68, 72
126	<i>sinesis</i> , <i>Sebastichthys</i> .....	1814	<i>Common</i> .....	68
135	<i>Sebastodes</i> .....	1813	<i>Little</i> .....	68
143	<i>Singing Fish</i> .....	2321	<i>Skates</i> .....	15, 66
141	<i>sinuatus</i> , <i>Merluccius</i> .....	2530	<i>Skeponopodus</i> .....	891
525	<i>Siphagonus</i> .....	2046	<i>guebucu</i> .....	891
526	<i>barbatus</i> .....	2050	<i>typus</i> .....	892
520	<i>segaliensis</i> .....	2048	<i>skib</i> , <i>Pomatomus</i> .....	947
221, 320	<i>Siphateles</i> .....	243	<i>Skil</i> .....	1862
2530	<i>vittatus</i> .....	244	<i>Skil-fishes</i> .....	1861
1370	<i>Siphostoma</i> .....	761, 763, 2837	<i>Skimback</i> .....	167
480	<i>affine</i> .....	769, 770	<i>Skipjack</i> .....	425, 805, 872, 946
493	<i>albirostre</i> .....	772	<i>Skipper</i> .....	725
1477	<i>arctum</i> .....	771, 2838	<i>Skowitz</i> .....	480
278	<i>ascendens</i> .....	768	<i>Sleeper</i> .....	2194, 2200, 2216
795, 889	<i>auliscus</i> .....	767	<i>Shark</i> .....	57
800	<i>bairdianum</i> .....	765, 770	<i>Slimer</i> .....	2315
805	<i>barbara</i> .....	765	<i>Slippery Dick</i> .....	1595
238	<i>brachycephalum</i> .....	769	<i>Sole</i> .....	2655
788, 796	<i>californiense</i> .....	764	<i>sloanei</i> , <i>Chauliodus</i> .....	585
439	<i>carinatum</i> .....	763	<i>Scomber</i> .....	870
11	<i>cayennense</i> .....	772	<i>sloani</i> , <i>Leiobatus</i> .....	81
213	<i>erinigerum</i> .....	771	<i>Raja</i> .....	81

	Page.		Page.
Small Black Lamprey.....	13	Snipefishes.....	758
Blindfish.....	704	Snip-nose Mullet.....	964
Catfish.....	140, 141	Snook.....	1118
Dolphin.....	953	Snub-nosed Eels.....	848
Small-mouthed Black Bass.....	1011	nyderi, <i>Catostomus</i> .....	2702
Buffalo.....	164	<i>Gnathypops</i> .....	2285
Small-scaled Gurnards.....	2175	<i>Oligocottus</i> .....	2871
Smaragdus.....	2210	Soapfish.....	538, 1229, 1232
<i>costalesi</i> .....	2225	Sobaco.....	1705, 1706
<i>stigmaticus</i> .....	2224	sobaco, <i>Balistes</i> .....	1706
<i>valenciennel</i> .....	2228	<i>Canthidornis</i> .....	1705
smaragdus, <i>Eleotris</i> .....	2204	Sobacos.....	1705
<i>Gobionellus</i> .....	2228	sobra, <i>Mosoprion</i> .....	1266
<i>Gobius</i> .....	2227	sociata, <i>Salmo</i> .....	521
Smaris.....	1364	socius, <i>Alburnus</i> .....	292
<i>lineatus</i> .....	1378	<i>Notropis</i> .....	292
<i>martiniens</i> .....	1365	socorroense, <i>Thalassoma</i> .....	1608, 2850
Snear Dab.....	2653, 2654	socorroensis, <i>Chlorichthys</i> .....	1607
Smecticus.....	1229	Soft Flounders.....	2670
<i>bicolor</i> .....	1232	sogo, <i>Holocentrus</i> .....	849
Smelt, American.....	523	Soi, Aka.....	1830
California.....	806	Gona.....	1833
Cobessicentic.....	524	Kuro.....	1834
Kodiak.....	2823	Shima.....	1834
Little.....	807	solandri, <i>Acanthocybium</i> .....	876
Pond.....	525	<i>Cybium</i> .....	877
Wilton.....	523	<i>Gempylus</i> .....	883
Smelt of the New York Lakes.....	468	solaris, <i>Orthogoriscus</i> .....	1754
Smelts.....	519, 522	Soldado.....	848
Surf.....	524	Soldier Fish.....	1088
smiridus, <i>Merluccius</i> .....	2530	Sole, American.....	2700
smithi, <i>Carcharodon</i> .....	50	California.....	2613
smithii, <i>Chloa</i> .....	253	Long-finned.....	2658
<i>Cyprinus (Abramis) ?</i> .....	413	Mexican.....	2698
Smooth Cabezon.....	2012	San Diego.....	2707
Hound.....	29	Slippery.....	2655
Puffer.....	1728	Solea.....	2660
smyrnenensis, <i>Gobius</i> .....	2118	<i>achirus</i> .....	2702
Snail, Sea.....	2105, 2114, 2116	<i>browni</i> .....	2701
Snake Blennies.....	2435, 2438	<i>cynoglossa</i> .....	2657
Eels.....	372	<i>flmbriata</i> .....	2700
Mackerel.....	883	<i>fischeri</i> .....	2790
Snap Mackerel.....	946	<i>fousecensis</i> .....	2699
Snapper.....	1760	<i>gronovii</i> .....	2696
Black-fin.....	1261	<i>inscripta</i> .....	2696
Dog.....	1257	<i>klunzingeri</i> .....	2697
Gray.....	1255	<i>maculipinnis</i> .....	2698
Lane.....	1270	<i>mazatlana</i> .....	2699
Mahogany.....	1272	<i>panamensis</i> .....	2702
Mangrove.....	1255	<i>pilosa</i> .....	2699
Red.....	1264	<i>reticulata</i> .....	2696
Red-tail.....	1270	<i>scutum</i> .....	2700
Silk.....	1262, 2858	solea, <i>Caranx</i> .....	927
Snappers.....	1241, 1247	Soleidae.....	2692
Snipe Eel.....	366, 369	Solenostomus.....	754, 756
Snipe, Sea.....	714	<i>soleiformis</i> , <i>Aramaca</i> .....	2672

Page.	Page.	Page.
758	2672	Sparisoma catesbyi..... 1638
904	2672	chrysopterum..... 1636, 1637
1118	2702	cyanolene..... 1633
848	2703	distinctum..... 1635, 1636
2702	2692	emarginatum..... 1641
2285	2693	flavescens..... 1639, 1640
0, 1232	1001	frondosum..... 1641, 1642
5, 1706	1001	hoplomystax..... 1632
1706	2818	lucinosum..... 1632
1705	50	lurido..... 1637
1705	56	maschalespilos..... 1641
1266	57	niphobles..... 1633
521	57	oxybrachium..... 1634
292	2198	radians..... 1631
292	1236	rubripinne..... 1640
08, 2859	2717	strigatum..... 1639
1607	1852	vride..... 1638
2679	689	xystrodon..... 1630
849	2216	Sparisometime..... 1621
1830	364	sparoides, Labrus..... 987
1833	493	Pomoxis..... 987
1834	2680	Sparopsis..... 1279
1834	2679	Sparus..... 1356, 1361
876	2680	argenteus..... 1357
877	1284	argyrops..... 1346
883	32	atlanticus..... 1153
1754	2630	aureus..... 1010
848	1346	bajonado..... 1352
1088	1732	brachysomus..... 1353
2700	86	castaneola..... 960
2613	2168	caxis..... 1250
2658	101, 1666, 1668	chrysomelanurus..... 1157
2698	216	chrysops..... 1346
2707	13	chrysurus..... 1276
2655	49	erectatus..... 1142
2660	49	falcatus..... 1585
2702	49	milneri..... 1355
2701	1139, 1140, 1817	niger..... 960
2657	1583	oblongus..... 2276
2700	1583	orbitalis..... 1350
2700	874	ovicephalus..... 1361
2699	866	pagrus..... 1357
2696	946	probratocephalus..... 1361
2696	947	radiatus..... 1506
2697	1498	raii..... 960
2698	1343	rhomboides..... 1358
2690	1241	salin..... 1360
2702	1343	sargus..... 1364
2699	1625, 1627, 1630	sclurus..... 1364
2700	1635	semiluna..... 1276
927	1642	synagris..... 1271
2692	1631	tetracanthus..... 1257
754, 756	1634	vermicularis..... 1271
2672	1641	virginicus..... 1323
	1638	vittatus..... 1323
		xanthurus..... 1346



	Page.		Page.
spathula, Polyodon .....	101	Spheroides furthi .....	1737
Squalus .....	102	lobatus .....	1731, 1732
spatula, Lepisosteus .....	111	maculatus .....	1733
Planirostra .....	102	marmoratus .....	1734
Spatularia .....	101	pachygaster .....	1738
reticulata .....	102	spengleri .....	1732, 1733
Spawn-eater .....	269	testudineus .....	1734
Spearfish .....	167, 891, 892	annulatus .....	1736
Spearing, Ground .....	533	trichocephalus .....	1737
speciosa, Gambusia .....	681	Sphorodius .....	1729
speciosus, Caranx .....	923	Sphyræna .....	822
Gnathedon .....	923	acus .....	717
Pomotis .....	1006, 1008	argentea .....	826
Scomber .....	928	aureoviridis .....	1119
Semotilus .....	222	barracuda .....	2841
Speck .....	1047	becuna .....	823
Speckled Hind .....	1159	bercalis .....	825
Trout .....	508	ensis .....	824
of Lake Crescent .....	2821	forsteri .....	824
spectabile, Etheostoma coeruleum ..	1089	guachancho .....	824
spectabilis, Percichthys .....	1089	guntheri .....	824
Salmo .....	508	lucasana .....	826
Salvelinus .....	508	picuda .....	823
spectrum, Careproctus .....	2133	picudilla .....	824
Lophius .....	2723	spet .....	820
Osmerus mordax .....	523	sphyraena .....	823, 826
spectruncula, Clio'la .....	265	viridescens .....	826
spectrunculus, Hybopsis .....	265	vulgaris .....	826
Notropis .....	265	sphyraena, Esox .....	826
speculiger, Exocoetus .....	734	Sphyræna .....	823, 826
Exonantes .....	2836	sphyraenarum, Echeneis .....	2268
speculigera, Lampadena .....	561	Sphyrænidæ .....	822
spelens, Amblyopsis .....	700	Sphyrænops .....	1114
spengleri, Spheroides .....	1732, 1733	bairdiannus .....	1111
Tetrodon .....	1733	Sphyrna .....	43, 44, 45
Spet .....	826	tiburo .....	44, 2748
spet, Esox .....	826	tudes .....	44
Sphyræna .....	826	zygæna .....	45
Sphærina .....	822	Sphyrnida .....	43
Spheroides furthi .....	1737	Spicara .....	1364
politus .....	1736	martinica .....	1361, 1365
trichocephalus .....	1738	Spikefish .....	891
tuberculatus .....	1733	spillmani, Alvordius .....	1039
Sphagebranchus .....	373	spilonotus, Exocoetus .....	740
anguliformis .....	374	spilonotus, Monacanthus .....	1716
kendalli .....	375	spilopterus, Citharichthys .....	2685, 2686
rostratus .....	373	Leuciscus .....	279
scuticaris .....	379	Photogenis .....	279
selachops .....	374	spilopus, Exocoetus .....	738
teres .....	379	spilota, Uranidea .....	1953, 1962
sphenops, Percilia .....	604	spilotopterygins, Balistes .....	1702
Spheroides .....	1729, 1731	spilotum, Etheostoma niangua .....	1044
angusticeps .....	1731	spilotus, Cottus .....	1961
annulatus .....	1735	Hypchomus .....	1043
politus .....	1736	spilurum, Cichlasoma .....	1520
formosus .....	1736	spilurus, Heros .....	1520

	Page.		Page.
1737		<i>splendens</i> , <i>Exocoetus</i> .....	720
1732		<i>splendida</i> , <i>Petonia</i> .....	1513
1733		<i>spleniatus</i> , <i>Pristipomus</i> .....	1322
1734		<i>spleniatus</i> , <i>Anisotremus</i> .....	1321
1738		<i>Split-mouth Sucker</i> .....	199
1733		<i>Split tail</i> .....	223
1734		<i>spongiosa</i> , <i>Haliutaea</i> .....	2742
1736		<i>Spoon-bill Cat</i> .....	101
1737		<i>Spot</i> .....	1458
1729		<i>Spot-tailed Minnow</i> .....	269
822		<i>Spotted Cabrilla</i> .....	1196
717		<i>Jewfish</i> .....	1162
826		<i>Kolptfish</i> .....	2353
1119		<i>Moray</i> .....	399
2841		<i>Rockfish</i> .....	1806
823		<i>Sea Trout</i> .....	1409
825		<i>Shiner</i> .....	318
824		<i>Sting Ray</i> .....	88
824		<i>Suckers</i> .....	186, 187
824		<i>Trunk-fish</i> .....	1723
824		<i>Weakfish</i> .....	1429
826		<i>Spotted-tail Minnow</i> .....	275
823		<i>Sprat</i> .....	432, 450
824		<i>Spratella</i> .....	424
826		<i>Spratelloides bryoporus</i> .....	422
823, 826		<i>lamprotenia</i> .....	419
826		<i>Springfish</i> .....	1950
826		<i>Squalidae</i> .....	53, 2749
826		<i>squalidus</i> , <i>Scarus</i> .....	1640
823, 826		<i>squalipeta</i> , <i>Echenels</i> .....	2272
2268		<i>Squalus</i> .....	228
822		<i>alicia</i> .....	236
1114		<i>ardesiacus</i> .....	237
1114		<i>atrarius</i> .....	233
3, 44, 45		<i>bicolor</i> .....	232
44, 2748		<i>caeruleus</i> .....	232
44		<i>canis</i> .....	29
45		<i>conformis</i> .....	231
43		<i>consersus</i> .....	234
1364		<i>cooperi</i> .....	236
364, 1365		<i>copel</i> .....	236
891		<i>crassus</i> .....	231
1039		<i>crureus</i> .....	233
740		<i>egregius</i> .....	237
1716		<i>elongatus</i> .....	240
2085, 2686		<i>estor</i> .....	240
279		<i>funduloides</i> .....	240
279		<i>galthe</i> .....	237
738		<i>gibbosus</i> .....	231
1953, 1962		<i>gula</i> .....	234
1702		<i>humboldti</i> .....	237
1044		<i>hyalope</i> .....	222
1961		<i>hydrophox</i> .....	238
1043		<i>intermedius</i> .....	235
1520		<i>lemmoni</i> .....	235
1520		<i>lineatus</i> .....	233
		<i>spilurus</i> , <i>Pecilia</i> .....	697
		<i>Salmo clarkii</i> .....	2819
		<i>mykiss</i> .....	495
		<i>Spina</i> , <i>Chopa</i> .....	1357, 1358
		<i>Spinax</i> .....	55
		<i>fabricii</i> .....	56
		<i>hillianus</i> .....	55
		<i>pusillus</i> .....	55
		<i>sucklii</i> .....	54
		<i>Spineless Sculpin</i> .....	2025
		<i>spinescens</i> , <i>Auliscops</i> .....	754
		<i>spiuicephalum</i> , <i>Exoglossum</i> .....	206
		<i>Spiuicephalus flabellatus</i> .....	2796
		<i>spinidens</i> , <i>Scarus</i> .....	1637
		<i>spinifer</i> , <i>Engraulis</i> .....	448
		<i>Stolephorus</i> .....	448
		<i>spiniger</i> , <i>Icelus</i> .....	1914
		<i>Spinivomer</i> .....	367
		<i>goodoi</i> .....	307
		<i>spinosissimus</i> , <i>Agonns</i> .....	2054
		<i>Aspidophorus</i> .....	2054
		<i>Diodon</i> .....	1746
		<i>Leptagonus</i> .....	2054
		<i>spinosus</i> , <i>Calycepidotus</i> .....	1937
		<i>Centronotus</i> .....	948
		<i>Chilomycterus</i> .....	1749
		<i>Cyclopterus</i> .....	2099, 2100
		<i>Dasyscopelus</i> .....	575
		<i>Diodon</i> .....	1749
		<i>Ehinorhinus</i> .....	58
		<i>Emmicerotremus</i> .....	2098, 2099
		<i>Hemilepidotus</i> .....	1937
		<i>Lampus</i> .....	2099
		<i>Orthogoriscus</i> .....	1754
		<i>Platophrys</i> .....	2662
		<i>Rhinobatus</i> .....	63
		<i>Rhomboidichthys</i> .....	2663
		<i>Schedophilopsis</i> .....	972
		<i>Scopelus</i> .....	575
		<i>Squalus</i> .....	58
		<i>Trachinotus</i> .....	942
		<i>spinulosus</i> , <i>Gasterosteus</i> .....	748
		<i>Sphyn-back Blowfish</i> .....	1734
		<i>Splyn Eels</i> .....	612
		<i>Spiny-rayed Fishes</i> .....	779
		<i>Spirinchus</i> .....	522
		<i>spiringulus</i> , <i>Leuciscus</i> .....	282
		<i>spixianus</i> , <i>Saurus</i> .....	538
		<i>Synodus</i> .....	533
		<i>spixli</i> , <i>Argyrosus</i> .....	930
		<i>Pimelodus</i> .....	132
		<i>Platysonus</i> .....	934, 2846
		<i>Tachisurus</i> .....	132
		<i>Tachysurus</i> .....	131, 2783
		<i>Vomer</i> .....	2846
		<i>splendens</i> , <i>Beryx</i> .....	844

	Page.		Page.
<i>Squalus margaritus</i> .....	241	<i>Squalus rostratus</i> .....	49
<i>modestus</i> .....	234	<i>selanonus</i> .....	49
<i>montanus</i> .....	238	<i>spathula</i> .....	102
<i>uiger</i> .....	235	<i>spinosis</i> .....	58
<i>ni-grescens</i> .....	234	<i>squatina</i> .....	59
<i>obesus</i> .....	233	<i>sucklii</i> .....	54, 2749
<i>pendora</i> .....	254	<i>tiburo</i> .....	36, 44
<i>photogenis</i> .....	296	<i>vulpes</i> .....	46
<i>proriger</i> .....	240	<i>vulpinus</i> .....	46
<i>pulchellus</i> .....	234	<i>zygæna</i> .....	45
<i>pulcher</i> .....	234	<i>squamata</i> , Tigoma .....	233
<i>purpureus</i> .....	234	<i>squamatus</i> , <i>Etheostoma</i> ( <i>Hadrop-</i>	
<i>rhomaleus</i> .....	233	<i>terus</i> ) .....	1040
<i>squamatus</i> .....	233	<i>Hypohomus</i> .....	1040
<i>tænia</i> .....	238	<i>Squalius</i> .....	233
<i>vandoisulus</i> .....	240	<i>squamiceps</i> , <i>Etheostoma</i> .....	1096
<i>Squalus</i> .....	53	<i>squamilentus</i> , <i>Ceraticthys</i> .....	323
<i>acanthias</i> .....	54	<i>Conesius</i> .....	323
<i>acronotus</i> .....	36	<i>Paralichthys</i> .....	2631
<i>alopecias</i> .....	46	<i>Squamipinnes</i> .....	781, 1665
<i>americanus</i> .....	47	<i>squamipinnis</i> , <i>Cestreus</i> .....	1404
<i>argus</i> .....	26	<i>Cynoscion</i> .....	1404, 1405
<i>borealis</i> .....	57	<i>Gerres</i> .....	1373
<i>bruceus</i> .....	58	<i>Otolithus</i> .....	1404
<i>cæruleus</i> .....	33	<i>squamosissimus</i> , <i>Diplolepis</i> .....	1410
<i>carcharias</i> .....	38, 50	<i>Pachyrurus</i> .....	1410
( <i>Carcharias</i> ) <i>terre-novæ</i> .....	43	<i>Plagioscion</i> .....	1418
<i>cetaceus</i> .....	47	<i>Schiæna</i> .....	1418
<i>citratus</i> .....	26	<i>squamosus</i> , <i>Trachurus</i> .....	921
<i>cornubicus</i> .....	49	<i>squamulosus</i> , <i>Chetodon</i> .....	1685
<i>elephas</i> .....	51	<i>Square-mouth</i> .....	208
<i>glacialis</i> .....	57	<i>Square-tails</i> .....	975, 976
<i>glaucus</i> .....	33	<i>Squatina</i> .....	58
<i>griseus</i> .....	19	<i>angelus</i> .....	59
<i>gunnerianus</i> .....	51	<i>californica</i> .....	59
<i>hinnullus</i> .....	29	<i>dumerill</i> .....	59
<i>hirundinæus</i> .....	33	<i>fimbriata</i> .....	59
<i>homiæus</i> .....	51	<i>japonica</i> .....	59
<i>isodus</i> .....	51	<i>lævis</i> .....	59
<i>littoralis</i> .....	47	<i>lewis</i> .....	59
<i>longimanus</i> .....	38	<i>oculata</i> .....	59
<i>macrodon</i> .....	47	<i>squatina</i> .....	58
<i>malleus</i> .....	45	<i>vulgaris</i> .....	59
<i>maximus</i> .....	51	<i>squatina</i> , <i>Rhina</i> .....	59
<i>microcephalus</i> .....	57	<i>Squalus</i> .....	50
<i>monensis</i> .....	49	<i>Squatina</i> .....	58
<i>nasus</i> .....	49	<i>Squatiniide</i> .....	58
<i>norwegianus</i> .....	57	<i>Squato</i> .....	58
<i>obscurus</i> .....	35	<i>Squaw-fish</i> .....	224
<i>obtusus</i> .....	39	<i>Squeteague</i> .....	1407
<i>pelegrinus</i> .....	51	<i>squeteague</i> , <i>Labrus</i> .....	1407, 1409
<i>pennanti</i> .....	49	<i>Squirrel-fish</i> .....	845, 847, 1203, 1207
<i>platyodon</i> .....	39	<i>Squirrel Hako</i> .....	2555
<i>punctatus</i> .....	26, 43	<i>stadthouderi</i> , <i>Serranus</i> .....	1150
<i>punctulatus</i> .....	26	<i>stagnalis</i> , <i>Salmo</i> .....	510
<i>ræhleghanus</i> .....	57	<i>Salvelinus</i> .....	509

Page.		Page.		Page.
40	<i>stagnalis</i> , <i>Salvelinus alpinus</i> .....	510	<i>stellatus</i> , <i>Mustellus</i> .....	29
49	<i>stahli</i> , <i>Larimus</i> .....	1423	<i>Plectichthys</i> .....	2652
102	<i>Monosira</i> .....	1423	<i>Pleuronectes</i> .....	2652
58	<i>stannii</i> , <i>Cotyliis</i> .....	2332	<i>stellatus</i> , <i>Salmo</i> .....	493
59	<i>Star-gazers</i> .....	2305	<i>stelleri</i> , <i>Cottus</i> .....	1941
54, 2749	Electric .....	2306	<i>Cotyliis</i> .....	2104
36, 44	Sand .....	2297	<i>Cyclopterichthys</i> .....	2104
46	<i>Star-headed Minnow</i> .....	656	<i>Cyclopterus</i> .....	2104
46	<i>starksi</i> , <i>Siphostoma</i> .....	771	<i>Hexagrammos</i> .....	1871
45	<i>Stolephorus</i> .....	2814	<i>Labrax</i> .....	1872
233	<i>Starksia</i> .....	2365	<i>Liparops</i> .....	2104
	<i>cremnobates</i> .....	2365, 2336	<i>Myoxocephalus</i> .....	1981
	<i>starksi</i> , <i>Siphostoma</i> .....	2838	<i>Trichodon</i> .....	2297
1040	<i>Starry Flounders</i> .....	2651	<i>Stellerina</i> .....	2041
1040	<i>Stathmonotinae</i> .....	2347	<i>xyosterna</i> .....	2042
233	<i>Stathmonotus</i> .....	2408	<i>Stellicarens</i> .....	1439, 1440, 1445
1096	<i>hemphillii</i> .....	2408	<i>Stellifer</i> .....	1439, 1443
323	<i>stearnsi</i> , <i>Blennius</i> .....	2379	<i>ericymba</i> .....	1444
323	<i>Corvina</i> .....	1458	<i>furthi</i> .....	1441
2631	<i>Lutjanus</i> .....	1256	<i>ilicebrosus</i> .....	1442
781, 1665	<i>Lutjanus</i> .....	1257	<i>lanceolatus</i> .....	1443
1404	<i>Prionotus</i> .....	2166	<i>microps</i> .....	1445
104, 1405	<i>Roncador</i> .....	1457	<i>minor</i> .....	1442
1373	<i>Scorpaena</i> .....	1843	<i>oscitans</i> .....	1440
1404	<i>stearnsii</i> , <i>Seriola</i> .....	903	<i>stellifer</i> .....	1443
1419	<i>Steel-backed Chub</i> .....	205	<i>zestocarus</i> .....	1445
1410	<i>Steelhead</i> .....	497	<i>stellifer</i> , <i>Bodianus</i> .....	1443
1418	<i>stegophthalmus</i> , <i>Agonus</i> .....	2036	<i>Fundulus</i> .....	048
1418	<i>steindachneri</i> , <i>Chlorichthys</i> .....	1609	<i>Sciæna</i> .....	1444
921	<i>Diabasis</i> .....	1302	<i>Stellifer</i> .....	1443
1685	<i>Hæmulon</i> .....	1301	<i>stelliferoides</i> , <i>Bassogigas</i> .....	2516
208	<i>Ophioblennius</i> .....	2401	<i>Neobythites</i> .....	2516
975, 976	<i>Rhinoptera</i> .....	91	<i>stellio</i> , <i>Perca</i> .....	1153
58	<i>Sebastes</i> .....	1830	<i>Rhinobatus</i> .....	2750
59	<i>Sebastodes</i> .....	1830	<i>stellulata</i> , <i>Raja</i> .....	75
59	<i>Thalassoma</i> .....	1609, 2859	<i>Stenobranchius</i> .....	561
59	<i>Steindachnerella</i> .....	2567	<i>andreae</i> .....	560
59	<i>Steindachneria</i> .....	2567	<i>coccol</i> .....	569
59	<i>argentea</i> .....	2568	<i>Stenodus</i> .....	473
59	<i>Stejnegeria</i> .....	960	<i>mackenzii</i> .....	474
59	<i>rubescens</i> .....	961	<i>Stenogobius</i> .....	2210
58	<i>Stejnegeriidae</i> .....	960	<i>Stenotomus</i> .....	1345
59	<i>stejnegeri</i> , <i>Stelgistrum</i> .....	1921	<i>aculeatus</i> .....	1346
59	<i>stelgidolepis</i> , <i>Macrourus</i> .....	2585	<i>caprinus</i> .....	1345
59	<i>Stelgis</i> .....	2067	<i>chrysops</i> .....	1346
59	<i>vulsus</i> .....	2067	<i>Stephanoberycidae</i> .....	835
58	<i>Stelgistrum</i> .....	1921	<i>Stephanoberyx</i> .....	836
58	<i>stejnegeri</i> .....	1921	<i>gillii</i> .....	836
58	<i>stellifera</i> , <i>Corvina</i> .....	1445	<i>mona</i> .....	836
224	<i>Sciæna</i> ( <i>Stelliferus</i> ) .....	1443	<i>Stephanolepis</i> .....	1714
1407	<i>Xenisma</i> .....	048	<i>setifer</i> .....	1716
407, 1409	<i>stellata</i> , <i>Platessa</i> .....	2652	<i>stephanophrys</i> , <i>Prionotus</i> .....	2161
203, 1207	<i>stellatus</i> , <i>Apogonichthys</i> .....	1110	<i>Stereolepis</i> .....	1137
2555	<i>Caranx</i> .....	926	<i>californicus</i> .....	1138
1150	<i>Fario</i> .....	492	<i>gigas</i> .....	1137
510	<i>Liparis</i> .....	2118	<i>Sterletus</i> .....	103
509				

	Page.		Page.
storletus, <i>Averruncus</i> .....	2071	<i>Stilbe americana</i> .....	250
<i>Ceratichthys</i> .....	316	<i>stilbe</i> , <i>Zalocys</i> .....	2848
<i>Sternias</i> .....	1926	<i>Stilbiscine</i> .....	359
<i>xenostethus</i> .....	1927	<i>Stilbiscus</i> .....	363
<i>Sternoptychida</i> .....	603	<i>edwardsi</i> .....	363
<i>Sternoptyx</i> .....	603	<i>Stilbins</i> .....	249, 250
<i>diaphana</i> .....	603, 2826	<i>stilbins</i> , <i>Leuroglossus</i> .....	527
<i>gardenii</i> .....	966	<i>Minnilus</i> .....	293
<i>hermannii</i> .....	603	<i>Notropis</i> .....	293
<i>mediterraneus</i> .....	604	<i>stilbostigma</i> , <i>Prionodes</i> .....	1216
<i>olforsii</i> .....	604	<i>Serranus</i> .....	1217
<i>Sternopygus humboldti</i> .....	341	<i>stimpsoni</i> , <i>Triglopsis</i> .....	2005
<i>Sternotremia</i> .....	786	<i>Sting Rays</i> .....	79, 82
<i>isolepis</i> .....	787	<i>Stingareo</i> .....	83
<i>stevensi</i> , <i>Thaleichthys</i> .....	521	<i>stipes</i> , <i>Atherina</i> .....	790
<i>Stichæina</i> .....	2349	<i>Joturua</i> .....	821
<i>Stichæus</i> .....	2439	<i>Stipivisch</i> .....	1702
<i>aculeatus</i> .....	2433	<i>stirurus</i> , <i>Chloroscoubrus</i> .....	938
<i>anguillaris</i> .....	2436	<i>Stit-tao</i> .....	499
<i>encecagrammus</i> .....	2441	<i>Stizostedion</i> .....	1020, 1021
<i>hexagrammus</i> .....	2441	<i>canadense</i> .....	1022
<i>islandicus</i> .....	2439	<i>boreum</i> .....	1022
<i>lumpenus</i> .....	2438	<i>griseum</i> .....	1022
<i>maculatus</i> .....	2438	<i>vitreum</i> .....	1021
<i>medius</i> .....	2436	<i>Stizostichium</i> .....	1020
<i>nubilus</i> .....	2438	<i>Stomolou</i> .....	88
<i>punctatus</i> .....	2439	<i>narinari</i> .....	88
<i>unimaculatus</i> .....	2441	<i>Stolephorus</i> .....	439
<i>Stickleback</i> , Alaska .....	749	<i>argyrophanus</i> .....	444
<i>Brook</i> .....	744	<i>astilbe</i> .....	2815
<i>California</i> .....	751	<i>brownii</i> .....	443
<i>Common Eastern</i> .....	748	<i>cherostomus</i> .....	444
<i>European</i> .....	747	<i>clupeoides</i> .....	447
<i>Nine-spined</i> .....	745	<i>compressus</i> .....	447
<i>Two-spined</i> .....	748	<i>cubanus</i> .....	442
<i>Sticklebacks</i> .....	742, 746	<i>cultratus</i> .....	443
<i>stigma</i> , <i>Gymnelis</i> .....	2477	<i>curtus</i> .....	445
<i>stigmaea</i> , <i>Ulocentra</i> .....	1047	<i>delicatissimus</i> .....	444
<i>stigmæum</i> , <i>Boleosoma</i> .....	1048	<i>engymen</i> .....	2815
<i>Etheostoma</i> .....	1048	<i>enrystole</i> .....	445
<i>stigmæus</i> , <i>Citharichthys</i> .....	2681	<i>exiguus</i> .....	442
<i>stigmatias</i> , <i>Paralichthys</i> .....	2636	<i>hiuleus</i> .....	443
<i>stigmaticus</i> , <i>Ceratichthys</i> .....	323	<i>ischaus</i> .....	442
<i>Gobionellus</i> .....	2224	<i>lucidus</i> .....	446, 2811
<i>Gobins</i> .....	2224	<i>macrolepidotus</i> .....	449
<i>Sigmops</i> .....	583	<i>miarchus</i> .....	441
<i>Smaragdus</i> .....	2224	<i>mitschilli</i> .....	446
<i>stigmatiscum</i> , <i>Lepophidium</i> .....	2483, 2484	<i>mundeolus</i> .....	2812
<i>Stigmatogobins</i> .....	2210	<i>naso</i> .....	2813
<i>stigmatura</i> , <i>Bolmannia</i> .....	2239	<i>opercularis</i> .....	445
<i>Cliola</i> .....	275	<i>panamensis</i> .....	448
<i>Codoma</i> .....	275	<i>per fasciatus</i> .....	441, 445
<i>stigmaturus</i> , <i>Gobins</i> .....	2220	<i>pertheatus</i> .....	442
<i>Notropis</i> .....	275	<i>poeyi</i> .....	445, 2811
<i>Photogenis</i> .....	275	<i>productus</i> .....	447
<i>Stilbe</i> .....	249, 250	<i>rastralis</i> .....	2811



	Page.		Page.
strumosus, Gobiesox .....	2333	subterraneus, Typhlichthys .....	704
Studfish .....	648	subtruncata, Belono .....	711
Sturgeon, Common .....	105	subtruncatus, Homopriion .....	1434
Green .....	104	Tylosurus .....	711
Lake .....	106	subulatus, Oreynus .....	871
Ohio .....	106	succincta, Seriola .....	900
Oregon .....	104	sucetta, Cyprinus .....	186
Red .....	106	Erimyzon .....	185, 186
Rock .....	106	oblongus .....	186
Sacramento .....	104	Sucker, Blue-headed .....	171
Short-nosed .....	106	Brook .....	178
Stone .....	106	Carp .....	166
White .....	104, 107	Columbia River .....	178
Sturgeons .....	102	Common .....	178
Shovel-nose .....	107	Eastern Carp .....	168
Sturio .....	103	Fine-scaled .....	178
vulgaris .....	105	Flannel-mouthed .....	174
sturio, Acipenser .....	105	Gourd-seed .....	168
sturioides, Paragonus .....	2063	Hare-lip .....	199
Podothecus .....	2063	Hog .....	181
Sturisoma .....	156, 157	Hump-backed .....	184
Stygicola .....	2500	June, of Utah Lake .....	183
dentata .....	2500	Large-scaled .....	192
dentatus .....	2500	Long-nosed .....	176
Stylephoridae .....	2601	Lump .....	2096
Stylephorus .....	2601	May .....	199
chordatus .....	2601	Missouri .....	168
stylifer, Hippocampus .....	778	Northern .....	176
Stypodon .....	220	Pea-lip .....	190
signifer .....	220	Rabbit-month .....	198, 199
suareus, Caranx .....	008	Razor-back .....	184
suavis, Chiola .....	272	Red .....	176
Cyprinella .....	272	Sacramento .....	178
subaequalis, Corvina .....	1429	Sand .....	1476
Corvina .....	1429	Split-mouth .....	199
subarcuata, Zygaena .....	45	Tahoe .....	177
subarcuatum, Haemulon .....	1306	Webug .....	180
subarmatus, Acanthurus .....	1691	White .....	178, 192
Subatka .....	595	Winter .....	187
subbifrenatus, Rhypticus .....	1233	Suckerel .....	168
subbifurcata, Pholis .....	2440	Sucker-mouthed Buffalo .....	164
Uivaria .....	2440	Suckers .....	161
subbifurcatus, Eumesogrammus .....	2440	Carp .....	165
subcaernea, Amia .....	113	Chub .....	185
subfuscus, Labrus .....	1578	Fine-scaled .....	173
subligarius, Centropristis .....	1219	Lump .....	2094
Dules .....	1218	Mountain .....	169, 170
Serranus .....	1219	Suckers Spotted .....	186, 187
suborbitalis, Holocentrus .....	850	White-nosed .....	190
Macurus (Nematoneurus) .....	2573	Suck-fish .....	2328
Nematoneurus .....	2572	Sucking-fish .....	2269
Plectromus .....	841	sucklii, Acanthias .....	54
subrotundus, Ostracion ventre glabro .....	1749	Catostomus .....	179
subterraneus, Lucifuga .....	2501	Spinax .....	54
		Squalus .....	54, 2749
		Sudis .....	599

Page.		Page.		Page.
704	<i>Sudis borealis</i> .....	601	<i>surinamensis</i> , <i>Batrachoides</i> .....	2314
711	<i>cornucans</i> .....	602	<i>Batrachus</i> .....	2314
1434	<i>intermedius</i> .....	600	<i>Engraulis</i> .....	447
711	<i>ringens</i> .....	601	<i>Galeichthys</i> .....	2780
871	<i>sucersonii</i> , <i>C. florhinus</i> .....	372	<i>Hexanemataichthys</i> ..	129
900	<i>suenri</i> , <i>Coryphæna</i> .....	953	<i>Holocentrus</i> .....	1236
186	<i>Cyprinus (Catostomus)</i> .....	195	<i>Lobotes</i> .....	1235, 2858
85, 186	<i>sufflumen</i> , <i>Ballates</i> .....	1706	<i>Lutjanus</i> .....	1319
186	<i>Canthidermis</i> .....	1706	<i>Plagioscion</i> .....	1419
171	<i>sugillatus</i> , <i>Ophisura</i> .....	387	<i>Pleuronectes</i> .....	2666
178	<i>Suillus</i> .....	1580	<i>Pœcilia</i> .....	691
166	<i>sullus</i> , <i>Lachnolaimus</i> .....	1580	<i>Pseudosciaena</i> .....	1420
178	<i>sujef</i> , <i>Muraenoides</i> .....	2419	<i>Sciæna</i> .....	1420
178	<i>sulcatus</i> , <i>Herpetoichthys</i> .....	382	<i>Stolephorus</i> .....	447
168	<i>Trachonurus</i> .....	2591	<i>Tachisurus</i> .....	130
178	<i>sumichrasti</i> , <i>Citharichthys</i> .....	2686	<i>Surmulletts</i> .....	855, 856
174	<i>Summer Flounder</i> .....	2629	<i>susanne</i> , <i>Boleosoma</i> .....	1059
168	<i>Herring</i> .....	426	<i>sutor</i> , <i>Blepharis</i> .....	932
199	<i>Sunapee Trout</i> .....	511	<i>Caranx</i> .....	932
181	<i>Sunfish</i> .....	931, 1753	<i>swaini</i> , <i>Notropis</i> .....	290
184	<i>Black-banded</i> .....	995	<i>Pœciliichthys</i> .....	1086
183	<i>Blue</i> .....	1005	<i>Swainia</i> .....	1039, 1040
192	<i>Blue-spotted</i> .....	996	<i>Swallowers</i> , <i>Black</i> .....	2291
176	<i>Common</i> .....	1009	<i>swampina</i> , <i>Hydrargira</i> .....	641
2096	<i>Green</i> .....	996	<i>Hydrargyra</i> .....	645
199	<i>Long-eared</i> .....	1002	<i>Fundulus</i> .....	645
168	<i>Mud</i> .....	989	<i>swanii</i> , <i>Bothragonus</i> .....	2086, 2088
176	<i>Red-spotted</i> .....	1004	<i>Hypsagonus</i> .....	2088
199	<i>Round</i> .....	988	<i>swannanoa</i> , <i>Etheostoma</i> .....	1070
198, 199	<i>Sunfishes</i> .....	984, 999	<i>Swellfish</i> .....	1729, 1748
184	<i>Banded</i> .....	994	<i>Swell Shark</i> .....	25
176	<i>Pigmy</i> .....	981	<i>Toad</i> .....	1732, 1733, 1748
178	<i>Sunny</i> .....	1009	<i>Swingle Tail</i> .....	45
1476	<i>suceri</i> , <i>Meletta</i> .....	425	<i>Swordfish</i> , <i>Common</i> .....	894
199	<i>superbus</i> , <i>Pseudoscarius</i> .....	1650	<i>Swordfishes</i> .....	893
177	<i>Scarus</i> .....	1650	<i>Syacinum</i> .....	2670
180	<i>supercilius</i> , <i>Aspidophorus</i> .....	2036	<i>latifrons</i> .....	2673
178, 102	<i>Hexagrammos</i> .....	1872	<i>micurum</i> .....	2672
187	<i>Hippocephalus</i> .....	2036	<i>ovale</i> .....	2674
168	<i>Hyborhynchus</i> .....	218	<i>papillosum</i> .....	2671
164	<i>Labrax</i> .....	1873	<i>Symbranchia</i> .....	341
161	<i>Surf Smelts</i> .....	524	<i>Symbranchide</i> .....	342
165	<i>Whiting</i> .....	1477	<i>Symbranchoid Eels</i> .....	342
185	<i>Surf-fish</i> .....	1503	<i>Symbranchus</i> .....	342
173	<i>Common</i> .....	1504	<i>innaculatus</i> .....	342
2094	<i>Striped</i> .....	1505	<i>marmoratus</i> .....	342
69, 170	<i>Wall-eyed</i> .....	1493, 1501	<i>vittatus</i> .....	342
86, 187	<i>White</i> .....	1506	<i>symmetricus</i> , <i>Apomotis</i> .....	998
190	<i>Surgeon</i> , <i>Blue</i> .....	1691	<i>Carunx</i> .....	910
2328	<i>Common</i> .....	1691	<i>Ceraticthys</i> .....	246
2269	<i>Surgeon-fishes</i> .....	1688	<i>Lepomis</i> .....	999
54	<i>surinamense</i> , <i>Pristipoma</i> .....	1319	<i>Leucosomus</i> .....	246
179	<i>surinamensis</i> , <i>Anisotremus</i> .....	1318, 1319	<i>Pogonichthys</i> .....	246
54	<i>inter-</i>		<i>Rutilus</i> .....	245
2749	<i>ruptus</i> .....	1319	<i>Salmo</i> .....	505
599	<i>Arius</i> .....	130	<i>Trachurus</i> .....	910



	Page.		Page.
Symmotrus	223	Syngnathus elucens	768
argyreus	234	ethon	767
Sympharus	2704, 2705	fasciatus	771
atramentatus	2706	fistulatus	765
atricaudus	2707	flavirostris	768
dionedeanus	2711	fuscus	770
elongatus	2707	griscollineatus	764
fasciolaris	2707	heckeli	2839
leei	2708	hippocampus	775
marginatus	2706	jonesi	768
nebulosus	2712	leptorhynchus	765
piger	2705	linea	768
plagiata	2710	louisiane	770
plagiata	2709	marmoreus	768
pusillus	2710	milbertianus	771
williamsi	2711	pockianus	771
Synagris	1288	peckii	770
macronemus	1289	pelagicus	770
synagris, Lutjanus	1271	picturatus	768
Neomans	1270	rousseau	767
Sparus	1271	segaliensis	2048
Synaphobranchidae	350	sibbaldi	774
Synaphobranchus	351	tenuis	766
affinis	351	viridescens	771
bathybius	352	Synodontidae	532
infernalis	352	Synodus	533, 2807
kanpii	351	argenteus	411
plunatus	351	evermanni	535
Synapteron	545	fasciatus	536
Synanceia ceronus	1941	foetens	538
Sybranchius fuliginosus	342	intermedius	535, 536
transversalis	342	jenkinsi	537, 2826
Synchirinae	1883	lacerta	537
Synchirus	2023	lacertinus	536
gilli	2024	lucioceps	539
Synecoglanis	133	myops	533
beadlei	135	poeyi	536
Syentognathi	707	saurus	537
Syentognathous Fishes	707	scituliceps	537, 2826
Syngnathi	760	spixianus	538
Syngnathidae	760	synodus	536
Syngnathinae	760	synodus, Esox	536
Syngnathus	701, 774	Saurus	536
abboti	764	Synodus	536
aequoreus	774, 2839	Sypterus	946
affinis	769	Syrrhina	61
albirostris	772	exasperata	65
arundinaceus	765	syrtensium, Argentina	526
ascendens	768	tabacaria, Fistularia	757, 758
bairdianus	770	Haliperca	1215
brachycephalus	769	tabacarius, Centropristes	1215
brevirostris	765	Prionodes	1215
californiensis	764	Serranus	1215
cayennensis	773	tachete, Le Diodon	1746
dekayi	771	Tachisurus albicaeus	124
dimidiatus	765		

ge.		Page.		Page.
768	<i>Tachysurus brandti</i> .....	122	<i>tænlotus</i> , <i>Tetragonopterus</i> .....	334
767	<i>dowi</i> .....	125	<i>Tæniphis</i> .....	392
771	<i>dubius</i> .....	127	<i>westphali</i> .....	300
765	<i>fissus</i> .....	131	<i>tænlops</i> <i>Bodianus</i> .....	1144
768	<i>flavescens</i> .....	123	<i>Enneacentrus</i> .....	1144
770	<i>furthii</i> .....	132	<i>Epinephelus</i> .....	1144
764	<i>grandicassis</i> .....	126	<i>Serranus</i> .....	1144
2839	<i>atricticassis</i> .....	126	<i>tæniopterus</i> , <i>Balistes</i> .....	1702
775	<i>gulosus</i> .....	133	<i>Cottus</i> .....	1979, 1988
768	<i>jordani</i> .....	120	<i>Perissias</i> .....	2607
765	<i>kesleri</i> .....	127	<i>Platophrys</i> .....	2068
768	<i>longicephalus</i> .....	130	<i>Pseudoscærus</i> .....	1640
770	<i>luniscutis</i> .....	125	<i>Scarus</i> .....	1646
768	<i>mesops</i> .....	123	<i>Tæniosomi</i> .....	782
771	<i>nuchalis</i> .....	131	<i>Tæniotoca</i> .....	1505
770	<i>oscula</i> .....	127	<i>lateralis</i> .....	1505, 1506
770	<i>platypogon</i> .....	127	<i>Tæhee</i> <i>Chub</i> .....	2798
768	<i>proops</i> .....	124	<i>Lake Trout</i> .....	493, 2870
767	<i>rugispinis</i> .....	130	<i>Sucker</i> .....	177
2048	<i>phrygiatus</i> .....	131	<i>tahoensis</i> , <i>Catostomus</i> .....	177
774	<i>seemani</i> .....	120	<i>Salmo clarkii</i> .....	2870
760	<i>spixii</i> .....	132	<i>talasica</i> , <i>Awaous</i> .....	2236
771	<i>surinamensis</i> .....	130	<i>Chonophorus</i> .....	2237
532	<i>temminckianus</i> .....	123	<i>Gobius</i> .....	2236
2807	<i>variolosus</i> .....	132	<i>Tall, Hard</i> .....	921
411	<i>Tachysurinae</i> .....	115, 2757	<i>Tailor Herring</i> .....	425
535	<i>Tachysurus</i> .....	119, 121, 131, 2782	<i>Tails, Square</i> .....	975, 976
536	<i>emmelane</i> .....	2785	<i>Talismania</i> .....	455
538	<i>emphysetus</i> .....	122	<i>æquatoris</i> .....	456
5, 536	<i>fissus</i> .....	131, 2782	<i>antillarum</i> .....	455
2826	<i>furthii</i> .....	132, 2787	<i>Tally-wag</i> .....	1199
537	<i>herzbergii</i> .....	125	<i>Tambor</i> .....	1732, 1734, 1805
536	<i>lentiginosus</i> .....	122	<i>Tang</i> .....	1691
539	<i>liropus</i> .....	2784	<i>Blue</i> .....	1691
533	<i>melanopus</i> .....	132, 2784	<i>Ocean</i> .....	1693
536	<i>multiradiatus</i> .....	132, 2788	<i>tang</i> , <i>Mugil</i> .....	812
537	<i>nuchalis</i> .....	2782	<i>Tangbrosme</i> .....	2438
2826	<i>nuchalus</i> .....	131	<i>tanneri</i> , <i>Hyperchoristus</i> .....	589
538	<i>passany</i> .....	124	<i>tardisma</i> , <i>Salmo</i> .....	483
536	<i>peruvianus</i> .....	122	<i>tapeinosoma</i> , <i>Auxis</i> .....	868
536	<i>spixii</i> .....	131, 2783	<i>Taractes</i> .....	957
536	<i>variolosus</i> .....	132, 2788	<i>sansurii</i> .....	957
536	<i>taczanowskii</i> , <i>Centronotus</i> .....	2416	<i>Tarandichthys</i> .....	1891
946	<i>Pholis</i> .....	2416	<i>cavifrons</i> .....	1891
61	<i>Sebastodes</i> .....	1831	<i>filamentosus</i> .....	1892
65	<i>tænia</i> , <i>Bassozetus</i> .....	2510	<i>tarascorum</i> , <i>Algansea</i> .....	2796
526	<i>Bathynna</i> .....	2510	<i>Tarentola</i> .....	537
758	<i>Blennius</i> .....	2418	<i>Tarletonbeania</i> .....	575
1215	<i>Clinostomus</i> .....	238	<i>crenularis</i> .....	575
1215	<i>Murenoides</i> .....	2418	<i>tenua</i> .....	575
1215	<i>Squalus</i> .....	238	<i>Tarpon</i> .....	409
1215	<i>tæniatum</i> , <i>Hæmulon</i> .....	1308	<i>atlanticus</i> .....	409
1215	<i>tæniatus</i> , <i>Anisotremus</i> .....	1322	<i>Tarpons</i> .....	408
1746	<i>Chirocentron</i> .....	435	<i>Tarpum</i> .....	409
124	<i>Evoxymetopon</i> .....	886	<i>Tates, Tom</i> .....	1308
			<i>tau</i> , <i>Batrachus</i> .....	2310

	Page.		Page.
Tau, Batrachoides .....	2314	tenorei, Scopelus .....	577
Batrachus beta .....	2316	Ten-pounder .....	410
pardus .....	2317	tentacunda, Trigla .....	2183
Gadus .....	2316	tentaculatus, Cottus .....	2000
Opsanus .....	2315	Porocottus .....	2000
Tauridon .....	1942, 1943, 1952	tenua, Tarletonbeania .....	575
taurina, Chrysophrys .....	1354	tonne, Moxostoma .....	186
taurinus, Calamus .....	1354	tonulilis, Antennarius .....	2721
taurocephala, Chloa .....	253	tenuirostris, Anguilla .....	348
taurocephalus, Alburnops .....	253	tenuis, Atherinopsids .....	802
taurus, Abulefiduf .....	1563	Gadus .....	2555
Carpoides .....	165	Icelinus .....	1894
Glyphidodon .....	1563	Leuresthes .....	802
Tautoga .....	1577	Opisthocentrus .....	2430
americana .....	1579	Phycis .....	2555
caerulea .....	1577	Syngnathus .....	706
niger .....	1577	Uranidea .....	1966
onitis .....	1578, 1579	Urophycis .....	2555
tessellata .....	1579	teres, Alosa .....	420
tautoga, Labrus .....	1579	Catostomus .....	179
fusca .....	1579	Cœnula .....	379
rubens .....	1579	Cyprinus .....	179
Tantogolabrus .....	1576	Etrumeus .....	420
adpersus .....	1577	Sphagebranchus .....	379
Tantogs .....	1577, 1578	Teretulus .....	187
taylori, Chilara .....	2489	cervinus .....	197
Ophidium .....	2489	teretulus, Phenacobius .....	803
Tchaviche .....	479	Hosternus .....	303
Tectospondyli .....	53, 58	tergisus, Hiodon .....	413
Telpalcate .....	2698	terra-novæ, Carcharias .....	43
Teleostei .....	113	Lycodes .....	2466
Teleostomi .....	97, 1241	Scollodon .....	43
Teloscops .....	1111	Squalus (Carcharias) .....	43
telescopus, Leuciscus .....	292	tessellata, Plagnia .....	2709
Miunilus .....	292	Tautoga .....	1579
Notropis .....	292	Tessellated Darter .....	1057
arcansanus .....	292	tessellatum, Holoosoma .....	1046, 1057
Photogenis .....	292	Etheostoma .....	1078
Telestes .....	228	tessellatus, Hadropterus .....	1070
Telipomis .....	995	Labrus .....	1578
temmnickianus, Bagrus .....	123	Nothonotus .....	1078
Sciadeichthys .....	122, 2760	Testar .....	2332
Tachisurus .....	123	testar, Lepadogaster .....	2332
temminckii, Arius .....	123	testudinens, Spheroides annulatus .....	1736
Ditrema .....	1510, 1511	Spheroides .....	1734
temminckii, Acanthoderma .....	880	Tetraodon .....	1735
Rovctus .....	880	Tetrodon .....	1735
Temniatia .....	1934	Tétard .....	2332
ventricosa .....	1936	tetard, Gnavina .....	2200
Temnodon .....	946	Tête-de-roche .....	1323
saltator .....	947	Tetrabanchus .....	342
tenebrosus, Alepoccephalus .....	453	tetracanthus, A cara .....	1540
Antennarius .....	2710	Centrarchus .....	1540
Chironectes .....	2719	Gasterosteus .....	748
tenellus, Fundulus .....	659	Heros .....	1539
Hyborhynchus .....	218	Sparus .....	1257

Page.		Page.
577	tetradens, <i>Zipotheca</i> .....	887
410	<i>Tetraodon erolithizon</i> .....	1739
2183	<i>ornatus</i> .....	1742
2000	<i>Tetragonopterine</i> .....	331
2000	<i>Tetragonopterus</i> .....	333
575	<i>areneus</i> .....	333
186	<i>argentatus</i> .....	330
2721	<i>brevimanus</i> .....	335
348	<i>fasciatus</i> .....	334
802	<i>flaeheri</i> .....	334
2555	<i>fuscoauratus</i> .....	334
1894	<i>humilis</i> .....	335
802	<i>mexicanus</i> .....	335
2430	<i>microphthalmus</i> .....	334
2555	<i>microstoma</i> .....	334
706	<i>cerstedli</i> .....	334
1906	<i>panamensis</i> .....	334
2555	<i>petenensis</i> .....	335
420	<i>rutilus</i> .....	334
179	<i>scabripinnis</i> .....	335
379	<i>taenlatus</i> .....	334
179	<i>Tetragonoptus</i> .....	1072
420	<i>Tetragonuridae</i> .....	975
379	<i>Tetragonurus</i> .....	975
187	<i>atlanticus</i> .....	976
197	<i>cuvieri</i> .....	976
303	<i>simplex</i> .....	880
303	<i>tetranemus</i> , <i>Hybopsis</i> .....	315
303	<i>Tetraodon setosus</i> .....	1740
413	<i>testudineus</i> .....	1735
43	<i>tetraodon</i> , <i>Ostracion</i> .....	1740
2466	<i>Tetraodontidae</i> .....	1726, 1727
43	<i>Tetraodontinae</i> .....	1727
43	<i>Tetrapterus amplus</i> .....	892
2709	<i>imperator</i> .....	892, 2844
1579	<i>indicus</i> .....	892
1057	<i>tetrapturorum</i> , <i>Echeneis</i> .....	2273
1057	<i>Tetrapturus</i> .....	891
1078	<i>albidus</i> .....	892
1070	<i>belono</i> .....	892
1578	<i>georgii</i> .....	892
1078	<i>herschelli</i> .....	892
2332	<i>lessoni</i> .....	892
2332	<i>tetraspilus</i> , <i>Upeneus</i> .....	860
1736	<i>Tetrodon</i> .....	1727
1734	<i>ammocryptus</i> .....	1735
1735	<i>angusticeps</i> .....	1731
1735	<i>annulatus</i> .....	1730
2332	<i>capistratus</i> .....	1742
2200	<i>caudicinctus</i> .....	1742
1323	( <i>Cheilichthys</i> ) <i>pachygaster</i> .....	1738
342	<i>curvus</i> .....	1728
1540	<i>formosus</i> .....	1737
1540	<i>furthi</i> .....	1737
748	<i>geometricus</i> .....	1735, 1730
1539	<i>heraldi</i> .....	1730
1257		
	<i>Tetrodon hispidus</i> .....	1733
	<i>hevigatus</i> .....	1728
	<i>lineolatus</i> .....	1728
	<i>lune</i> .....	1754
	<i>mathematicus</i> .....	1728
	<i>mola</i> .....	1754
	<i>nephelus</i> .....	1733
	<i>oxyrhynchus</i> .....	1741
	<i>pachycephalus</i> .....	1729
	<i>pachygaster</i> .....	1738
	<i>plumieri</i> .....	1733
	<i>pollus</i> .....	1736
	<i>psittacus</i> .....	1740
	<i>punctatissimus</i> .....	1741
	<i>punctatus</i> .....	1735
	<i>rostratus</i> .....	1742
	<i>spengleri</i> .....	1733
	<i>testudineus</i> .....	1735
	<i>trichocephalus</i> .....	1738
	<i>truncatus</i> .....	1756
	<i>turgidus</i> .....	1733
	<i>nephelus</i> .....	1733
	<i>Tetronarce</i> .....	77
	<i>californica</i> .....	77
	<i>occidentalis</i> .....	77
	<i>trophthalma</i> , <i>Lioglossina</i> .....	2622
	<i>Tetroras</i> .....	51
	<i>Tenthididae</i> .....	1688
	<i>Tenthis</i> .....	1685
	<i>allala</i> .....	1693
	<i>australis</i> .....	1691
	<i>bahianus</i> .....	1693
	<i>caeruleus</i> .....	1691
	<i>crestonis</i> .....	1692
	<i>hepatus</i> .....	1692
	<i>tractus</i> .....	1693
	<i>trioctegus</i> .....	1690
	<i>Teuthys</i> .....	1689
	<i>texana</i> , <i>Anguilla</i> .....	348
	<i>Cyprinella</i> .....	274
	<i>texanus</i> , <i>Catostomus</i> .....	192
	<i>Notropis</i> .....	274
	<i>Texas Redhorse</i> .....	192
	<i>texensis</i> , <i>Dionda</i> .....	215
	<i>textilis</i> , <i>Salaria</i> .....	2400
	<i>Salariaichthys</i> .....	2400
	<i>Thaerondotis</i> .....	392
	<i>thalassinum</i> , <i>Ethoastoma</i> .....	1071
	<i>Moxostoma</i> .....	191
	<i>Nothouotus</i> .....	1072
	<i>thalassinus</i> , <i>Cestreus regalis</i> .....	1408
	<i>Cynoscion</i> .....	1407
	<i>Doratonotus</i> .....	1612
	<i>Gobius</i> .....	2245
	<i>Lepidogobius</i> .....	2245
	<i>Microgobius</i> .....	2245

	Page.		Page.
<i>thalassinus</i> , <i>Myloleucus</i> .....	245	<i>Thrissa</i> .....	422
<i>Otolithus</i> .....	1408	<i>thrissa</i> <i>Chupea</i> .....	432
<i>Ptychostomus</i> .....	192	<i>thrissina</i> , <i>Clupea</i> .....	431
<i>Rutilus</i> .....	245	<i>Sardinella</i> .....	430
<i>Thalassoma</i> .....	2850	<i>thrissoides</i> , <i>Megalops</i> .....	409
<i>bifasciatum</i> .....	1610, 2850	<i>Thumb</i> , <i>Miller's</i> .....	1941, 1950
<i>bifasciatus</i> .....	1610	<i>Thunder-pumper</i> .....	1484
<i>grammaticum</i> .....	1610, 2850	<i>thunnia</i> , <i>Orcynus</i> .....	809
<i>lucasanum</i> .....	1007, 2850	<i>Thynnichthys</i> .....	809
<i>nitidissimum</i> .....	2850	<i>Thynnus</i> .....	860
<i>nitidum</i> .....	1608, 2850	<i>Thunnus</i> .....	809
<i>socorroense</i> .....	1608, 2850	<i>thynnus</i> .....	870
<i>steindachneri</i> .....	1609, 2850	<i>Thymallidae</i> .....	517
<i>virens</i> .....	1611, 2850	<i>thymulloides</i> , <i>Coregonus</i> .....	518
<i>Thalassophryne</i> .....	2323	<i>Thymellus</i> .....	517
<i>dowi</i> .....	2326	<i>lewisii</i> .....	2871
<i>maculosa</i> .....	2324	<i>ontariensis</i> .....	518
<i>reticulata</i> .....	2325	<i>montanus</i> ... ..	519
<i>Thaleichthys</i> .....	521	<i>signifer</i> .....	517, 2871
<i>pacificus</i> .....	521	<i>montanus</i> .... ..	519
<i>stevensi</i> .....	521	<i>ontariensis</i> ... ..	519
<i>thaleichthys</i> , <i>Osmernus</i> .....	522	<i>tricolor</i> .....	519, 2871
<i>thazard</i> , <i>Auxis</i> .....	867	<i>monti-</i>	
<i>Scomber</i> .....	867	<i>nus</i> .. ..	2871
<i>Theragra</i> .....	2535	<i>tricolor</i> .....	519
<i>chalcogramma</i> .....	2535	<i>Thynnichthys</i> .....	868
<i>fucensis</i> .....	2536	<i>breviplinnis</i> .....	869
<i>Theraps</i> .....	1540	<i>thunnia</i> .....	869
<i>irregularis</i> .....	1540	<i>thynnoides</i> , <i>Auxis</i> .....	868
<i>thermalis</i> , <i>Pæcilla</i> .....	693	<i>Thynnus</i> .....	868, 869
<i>theta</i> , <i>Diaphnus</i> .....	565	<i>affinis</i> .....	869
<i>Diaphus</i> .....	564	<i>argenti-vittatus</i> .....	871
<i>Theuthis</i> .....	1689	<i>atlanticus</i> .....	871
<i>Thick-tailed Rays</i> .....	60	<i>balteatus</i> .....	871
<i>Thimble-eyed Mackerel</i> .....	866	<i>brachypterus</i> .....	870
<i>thoburni</i> , <i>Alcidaea</i> .....	1887	<i>brasiliensis</i> .....	869
<i>Mugil</i> .....	813	<i>breviplinnis</i> .....	869
<i>Tholichthys</i> .....	1672	<i>coretta</i> .....	870
<i>thomponi</i> , <i>Acipenser</i> .....	105	<i>leachianus</i> .... ..	869
<i>Amla</i> .....	113	<i>macropterus</i> .....	871
<i>Carpilodes</i> .....	167	<i>mediterraneus</i> .....	870
<i>Podothecus</i> .....	2060	<i>pacificus</i> .....	871
<i>Ptyonotus</i> .....	2005	<i>pelamys</i> .....	869
<i>Triglopis</i> .....	2005	<i>pompilus</i> .....	900
<i>thomsonii</i> , <i>Cottunculus</i> .....	1993	<i>rocheanus</i> .....	868
<i>Cottus</i> .....	1994	<i>secundo-dorsalis</i> .....	870
<i>thoreautanus</i> , <i>Semotilus</i> .....	223	<i>thunnia</i> .....	869
<i>atromaculatus</i> .....	223	<i>vulgaris</i> .....	870
<i>Threadfins</i> .....	827	<i>thynnus</i> , <i>Albacora</i> .....	870
<i>Threadfishes</i> .....	931	<i>Orcynus</i> .....	870
<i>Thread Herring</i> .....	432	<i>Scomber</i> .....	870
<i>Three-angled Trunk-fishes</i> .....	1721	<i>Thunnus</i> .....	870
<i>Three-bearded Rocklings</i> .....	2557	<i>Thyrina</i> .....	803, 2840
<i>Thresher</i> .....	45	<i>crystallina</i> .....	804
<i>Sharks</i> .....	45	<i>evermanni</i> .....	804
		<i>Thyris</i> .....	2690

Page.		Page.		Page.
422	<i>Thyriscus pellucidus</i> .....	2691	<i>Tilesia gracilis</i> .....	2538
432	<i>Thyriscus acanthoderma</i> .....	880	<i>tilesii</i> , <i>Hemilepidotus</i> .....	1936
431	<i>niger</i> .....	879	<i>timpanogensis</i> , <i>Hybopsis</i> .....	233
430	<i>protiosus</i> .....	880	<i>Minnilus</i> .....	233
409	<i>acholaris</i> .....	880	<i>Timucu</i> .....	711, 715
41, 1050	<i>Thyriscinus</i> .....	877	<i>tinucu</i> , <i>Belone</i> .....	715
1484	<i>thyrsitoides</i> , <i>Lemulasma</i> .....	884	<i>Esox</i> .....	711
869	<i>Thyriscus violaceus</i> .....	879	<i>tinea</i> , <i>Serranus</i> .....	1181
869	<i>Thyrsoidea atorrhina</i> .....	396	<i>tinocella</i> , <i>Algaussea</i> .....	211, 2796
869	<i>concolor</i> .....	396	<i>Leuciscus</i> .....	211
869	<i>cormura</i> .....	394	<i>Leucus</i> .....	211
870	<i>maculiphinis</i> .....	394	<i>Tinker Mackerel</i> .....	866
517	<i>marginata</i> .....	394	<i>Tinosa</i> .....	924
518	<i>millaris</i> .....	398	<i>tippecanoe</i> , <i>Etheostoma</i> .....	1090
517	<i>Tiaroga</i> .....	305	<i>Tippecanoe Sisco</i> .....	469
2871	<i>cobitis</i> .....	305	<i>Tirantes</i> .....	885
518	<i>tiburo</i> , <i>Renciceps</i> .....	44	<i>Tiru</i> .....	537
519	<i>Sphyrna</i> .....	44, 2748	<i>Tirus</i> .....	538
17, 2871	<i>Squalus</i> .....	36, 44	<i>marmoratus</i> .....	537
519	<i>Zygona</i> .....	44	<i>Toad</i> , <i>Swell</i> .....	1732, 1733
519	<i>Tiburon</i> .....	39	<i>Toadfish</i> .....	1733, 1748, 2313, 2315
519, 2871	<i>Ticky-ticky</i> .....	659	<i>Toad-fishes</i> , <i>Poison</i> .....	2323
	<i>Tiger Shark</i> .....	32	<i>Tobacco Box</i> .....	68
	<i>Tigoma</i> .....	228, 230, 231	<i>Toeroc</i> .....	1403
2871	<i>bicolor</i> .....	232	<i>toeroc</i> , <i>Otolithus</i> .....	1404
519	<i>conformis</i> .....	231	<i>Togue</i> .....	504
868	<i>conspersa</i> .....	234	<i>Tokenoko mo waru</i> .....	1829
869	<i>crassa</i> .....	231	<i>Tom</i> , <i>Mad</i> .....	144, 147
868	<i>egregia</i> .....	237	<i>tom</i> , <i>Salmo</i> .....	505
808, 869	<i>gibbosa</i> .....	235	<i>Tomcod</i> , <i>California</i> .....	2539
869	<i>gracilis</i> .....	236	<i>tomcod</i> , <i>Gadus</i> .....	2540
871	<i>humboldti</i> .....	237	<i>Microgadus</i> .....	2540
871	<i>intermedia</i> .....	235	<i>Tomcods</i> .....	2538, 2540
871	<i>lineata</i> .....	233	<i>tomcodus</i> , <i>Gadus</i> .....	2540
870	<i>nigrescens</i> .....	234	<i>fuscus</i> .....	2540
869	<i>obesa</i> .....	233	<i>luteus</i> .....	2540
869	<i>pulchra</i> .....	234	<i>mixtus</i> .....	2540
870	<i>purpurea</i> .....	234	<i>Tomicodon</i> .....	2329
869	<i>rhinichthyoides</i> .....	312	<i>Tomato</i> .....	1308
871	<i>aquamata</i> .....	233	<i>Tondo</i> , <i>Pesce</i> .....	48
870	<i>tigrinus</i> , <i>Galeocerdo</i> .....	32	<i>Tongue Fish</i> .....	2704, 2710
871	<i>Holocentrus</i> .....	1214	<i>Toothed Herring</i> .....	413
869	<i>Myrichthys</i> .....	376	<i>Top Minnow</i> .....	659, 680
900	<i>Prionodes</i> .....	1214	<i>topeka</i> , <i>Notropis</i> .....	266
868	<i>Serranus</i> .....	1214	<i>Topea</i> .....	31
870	<i>tigris</i> , <i>Antennarius</i> .....	2723	<i>Toro</i> .....	920, 1724
869	<i>Carcharias</i> .....	49	<i>torpedinus</i> , <i>Trygonobatus</i> .....	81
870	<i>Chironectes</i> .....	2723	<i>Urolophus</i> .....	81
870	<i>Epinephelus</i> .....	1187	<i>Torpedo</i> .....	77
870	<i>Mycteroperca</i> .....	1187	<i>bancrofti</i> .....	78
870	<i>camelopardalis</i> .....	1187	<i>brasiliensis</i> .....	78
870	<i>Serranus</i> .....	1187	<i>californica</i> .....	78
803, 2840	<i>Triastropis</i> .....	1187	<i>occidentalis</i> .....	77
804	<i>Tigrone</i> .....	32, 39	<i>pictus</i> .....	78
804	<i>Tilefish</i> .....	2278	<i>torpedo</i> , <i>California</i> .....	77
2690	<i>Tilesia</i> .....	2537	<i>torquatus</i> , <i>Labrus</i> .....	1609

	Page.		Page.
Torrentaria .....	1066, 1068, 1080	Trachurus .....	909
torridus, Lutjanus .....	1264	alicolus .....	904
torsk, Blennius .....	2561	boops .....	922
Gadus .....	2561	ouvieri .....	910
torvus, Cottunculus .....	1994	europaeus .....	911
tota, Perca maculis .....	1153	fallax .....	910
Toter .....	181	fasciatus .....	904
toto, Cyprinus .....	415	imperialis .....	927
Totuaa .....	1411	limnei .....	911
townaendi, Gobius .....	2250	picturatus .....	909, 2844
Lamparyctus .....	558	risoi .....	910
Myctophum .....	558	saurus .....	911
toxotes, Ditrema .....	1508	squamosus .....	921
Rhacochilus .....	1507	symmetricus .....	910, 2844
Trachelocirrus .....	950	trachurus .....	910
Trachichthyidae .....	836	trachurus, Caranx .....	910
Trachichthys pretiosus .....	837	Cottus .....	1936
Trachidermis richardsoni .....	1944	Gasterosteus .....	747
Trachinocephalus .....	533	Hemlepidotus .....	1936
myops .....	533	Scomber .....	910
Trachinoid Fishes .....	2273	Trachurus .....	910
Trachinoidea .....	2273	Trachynotus argenteus .....	944
Trachinoidei .....	781	carolinus .....	944
Trachinotina .....	897	cupreus .....	944
Trachinotus .....	939	fasciatus .....	941
argentens .....	944	glaucoides .....	941
carolinus .....	944	glaucus .....	941
cayennensis .....	945	goreensis .....	843
culveri .....	942	nasutus .....	941
cupreus .....	944	ovatus .....	942
falcatus .....	941, 2847	pampanus .....	944
fuscus .....	942	rhomboides .....	942
glaucus .....	940	trachypoma, Myripristis .....	846
goodei .....	943	Trachypterus .....	2870
kennedyi .....	942	trachyurus .....	2601
paloma .....	945, 2848	Trachyrhamphus .....	761, 2568
rhodopus .....	911, 943	Trachyrhynchus .....	2562
rhomboides .....	2847	Trachyrhynchus .....	2568
spinosus .....	942	Helolepis .....	2568
Trachinus adsenclonius .....	1153	trachyurus, Trachypterus .....	2601
cirrhosus .....	2019	tractus, Acanthurus .....	1693
gasteropeleus .....	2297	Tenthis .....	1693
osbeck .....	1153	Trahiras .....	330
punctatus .....	1153	tranquebar, Aspidophoroides .....	2092
trichodon .....	2296	transmontana, Columbia .....	784
trachinus, Malacanthus .....	2270	transmontanus, Actpenser .....	104
Salmo .....	523	Rhinichthys .....	307
Trachisurus parkeri .....	126	transversalis, Synbranchus .....	342
Trachourus .....	2591	transversum, Pectilosoma .....	1089
sulcatus .....	2591	traski, Hysteroecarpus .....	1496
trachura, Raja .....	70	treulli, Dioplites nuceensis .....	1012
Raja .....	75	Trefish .....	1827
Trachurops .....	911	Trematopsis .....	1753
brachycheirus .....	911	willugbei .....	1754
crumenophthalmus .....	911	Triacanthidae .....	1697
plumieri .....	912	Triacanthodidae .....	1697

Page.		Page.		Page.
909	trilacanthus, <i>Argyriosus</i> .....	936	tricolor, <i>Chavodon</i> .....	1684
904	<i>Nauclerus</i> .....	900	<i>Engraulis</i> .....	443
922	<i>Poronotus</i> .....	2849	<i>Genicanthus</i> .....	1684
910	<i>Rhombus</i> .....	967	<i>Holacanthus</i> .....	1684
911	<i>Stromateus</i> .....	968	<i>Pomacanthus</i> .....	1684
910	<i>Xenochirus</i> .....	2084	<i>Thymallus</i> .....	519
904	<i>Triakis henlei</i> .....	31	<i>signifer</i> .....	519
927	<i>triagramma</i> , <i>Heros</i> .....	1529	<i>Tricopterus</i> .....	915, 917, 920
911	<i>Triakis</i> .....	31	<i>tricornis</i> , <i>Lactophrys</i> .....	1724
009, 2844	<i>semifasciatum</i> .....	31	<i>Ostracion</i> .....	1725
910	<i>triangularis</i> , <i>Ostracion tuberculus</i> .....	1723	<i>tricuspidatus</i> , <i>Hyporhamphus</i> .....	720
911	<i>Pachynathus</i> .....	1705	<i>tricuspis</i> , <i>Cottus</i> .....	2009
921	<i>triangulum</i> , <i>Lutjanus</i> .....	1454	<i>Gymnocanthus</i> .....	2008
010, 2844	<i>Tribe</i> , <i>Flounder</i> .....	2607	<i>Phobos</i> .....	2009
910	<i>Halibut</i> .....	2605	<i>tridecemlineatus</i> , <i>Esox</i> .....	628
910	<i>Turbot</i> .....	2608	<i>tridens</i> , <i>Acropocilia</i> .....	690
1936	<i>tribunus</i> , <i>Prionotus</i> .....	2171, 2172	<i>Archosargus</i> .....	1360
747	<i>Trigla</i> .....	2172	<i>Centropristis</i> .....	1292
1936	<i>Trichidion</i> .....	828	<i>Lutjanus</i> .....	1292
910	<i>approximans</i> .....	829	<i>Poecilia (Acropocilia)</i> .....	690
910	<i>octofilis</i> .....	830	<i>Sargus</i> .....	1361
944	<i>oconemus</i> .....	830	<i>tridentata</i> , <i>Lampetra</i> .....	12
944	<i>opercularis</i> .....	831	<i>tridentatus</i> , <i>Ammocetus</i> .....	12
944	<i>plumieri</i> .....	830	<i>Entosphenus</i> .....	12
941	<i>Trichiuridae</i> .....	838	<i>Ichthyomyzon</i> .....	12
941	<i>Trichurus</i> .....	889	<i>Petromyzon</i> .....	12
941	<i>argenteus</i> .....	889	<i>tridentiger</i> , <i>Gambusia</i> .....	2833
243	<i>caudatus</i> .....	887	<i>tridigitatus</i> , <i>Dactyloscopus</i> .....	2301
941	<i>ensiformis</i> .....	887	<i>Polynemus</i> .....	2177
942	<i>gladius</i> .....	887	<i>Trifarcus</i> .....	670
944	<i>lepturus</i> .....	889, 2844	<i>felicianus</i> .....	676
942	<i>trichocephalus</i> , <i>Sphaeroides</i> .....	1738	<i>riverendi</i> .....	673
840	<i>Tetrodon</i> .....	1738	<i>trifasciata</i> , <i>Hydrargyra</i> .....	639
2870	<i>Trichocycilus</i> .....	1743, 1744	<i>Lepomis</i> .....	1011
2601	<i>erinaceus</i> .....	1744	<i>trifurca</i> , <i>Centropristes</i> .....	1202
761, 2568	<i>Trichoderma</i> .....	1714	<i>Perca</i> .....	1202
2562	<i>Trichodiodon</i> .....	1743, 1744	<i>trifurcus</i> , <i>Anthias</i> .....	1202
2568	<i>pilosus</i> .....	1743, 1744	<i>Centropristis</i> .....	1202
2568	<i>Trichodon</i> .....	2295	<i>Serranus</i> .....	1201, 1202
2601	<i>japonicus</i> .....	2297	<i>trigammus</i> , <i>Chirus</i> .....	1872
1093	<i>lineatus</i> .....	2297	<i>Trigger Fishes</i> .....	1698, 1699
1093	<i>stelleri</i> .....	2297	<i>Trigla</i> .....	2176, 2867
330	<i>trichodon</i> .....	2295	<i>carolina</i> .....	2156, 2172
2092	<i>trichodon</i> , <i>Drachinus</i> .....	2297	<i>euculus</i> .....	2177
784	<i>Mugil</i> .....	816	<i>digitis vicensis paluatis</i> .....	2183
104	<i>Trachinus</i> .....	2296	<i>evolans</i> .....	2169
307	<i>Trichodon</i> .....	2295	<i>fasciata</i> .....	2183
342	<i>Trichodontidae</i> .....	2295	<i>lineata</i> .....	2197
1089	<i>Trichonotus</i> .....	409	<i>palmipes</i> .....	2156
1496	<i>Trichopsetta</i> .....	2669	<i>pletipinnis</i> .....	2176
1012	<i>ventralis</i> .....	2669	<i>pini</i> .....	2177
1827	<i>trichopterus</i> , <i>Pteraclis</i> .....	956	<i>punctata</i> .....	2170
1753	<i>trichreletta</i> , <i>Cliola</i> .....	276	<i>strigata</i> .....	2167
1754	<i>Codoma</i> .....	276	<i>tentabunda</i> .....	2183
1697	<i>trichroistilus</i> , <i>Notropis</i> .....	275	<i>tota rubens</i> .....	2177
1697	<i>tricocephalus</i> , <i>Spheroides</i> .....	1737	<i>tribulus</i> .....	2172



	Page.		Page.
<i>Trigla voltans</i> .....	2183	<i>Trisotropis petrosus</i> .....	1172
<i>Triglida</i> .....	2147	<i>reticulatus</i> .....	1187
<i>Triglochis</i> .....	46	<i>rosacens</i> .....	1184
<i>Triglops</i> .....	1923	<i>atomias</i> .....	1178
<i>beani</i> .....	1924	<i>tigris</i> .....	1187
<i>pingelli</i> .....	1923	<i>trispinosa</i> , <i>Corvina</i> .....	1443
<i>plourostictus</i> .....	1923	<i>trispinosus</i> , <i>Odontopyxis</i> .....	2085
<i>scepticus</i> .....	1925	<i>Pseudoscarus</i> .....	1648
<i>xonostethus</i> .....	1927	<i>Scarus</i> .....	1648
<i>Triglopsis</i> .....	2005	<i>trispinosus</i> , <i>Odontopyxis</i> .....	2086
<i>stimpsoni</i> .....	2005	<i>tristis</i> , <i>Montana</i> .....	272
<i>thompsoni</i> .....	2005	<i>tristaeclus</i> , <i>Esox</i> .....	111
<i>Trigonobatus</i> .....	82	<i>Lepisosteus</i> .....	111
<i>trigonum</i> , <i>Ostracium</i> .....	1724	<i>Litholepis</i> .....	111
<i>trigonus</i> , <i>Lactophrys</i> .....	1723, 1724	<i>trirus</i> , <i>Bodianus</i> .....	1236
<i>Ostracium</i> .....	1724	<i>trivittatus</i> , <i>Diabasis</i> .....	1311
<i>trilineatum</i> , <i>Pristipoma</i> .....	1320	<i>Grammistes</i> .....	1311
<i>trilineatus</i> , <i>Anisotremus</i> .....	1320	<i>Sebastodes</i> .....	1834
<i>trilobatus</i> , <i>Scarus</i> .....	1654	<i>Trochocopus darwini</i> .....	1586
<i>Triloburus</i> .....	1198, 1199, 1201	<i>pulcher</i> .....	1585
<i>trilobus</i> , <i>Blepsias</i> .....	2019	<i>Trompa</i> .....	1653
<i>Lutjanus</i> .....	1200	<i>Lija</i> .....	171
<i>trimaculatus</i> , <i>Heros</i> .....	1529	<i>Trompetero</i> .....	754, 757
<i>Trinectes</i> .....	2693	<i>tropica</i> , <i>Echeneis</i> .....	2268
<i>scabra</i> .....	2701	<i>tropicus</i> , <i>Atractostens</i> .....	111
<i>trioctegus</i> , <i>Acanthurus</i> .....	1691	<i>Lepisostens</i> .....	111
<i>Chaetodon</i> .....	1691	<i>Tropidichthys</i> .....	1741
<i>Teuthis</i> .....	1690	<i>Tropidinius</i> .....	1278
<i>tripes</i> , <i>Nealotus</i> .....	881	<i>arnillo</i> .....	1279
<i>Triple-tail</i> .....	1235	<i>dentatus</i> .....	1279
<i>Tripteronotus</i> .....	461	<i>Tropidodus</i> .....	20
<i>Tripterygium carminale</i> .....	2350	<i>troschelli</i> , <i>Heros</i> .....	1537
<i>tripterygius</i> , <i>Cottus</i> .....	2023	<i>Sciadichthys</i> .....	122, 2757
<i>tripunctulatus</i> , <i>Maurollicus</i> .....	578	<i>Sciades</i> .....	122, 2758
<i>Valenciennellus</i> .....	578	<i>troschelli</i> , <i>Glyphidodon</i> .....	1562
<i>triqueter</i> , <i>Lactophrys</i> .....	1722	<i>Trout</i> .....	483
<i>Ostracium</i> .....	1723	<i>Black-spotted</i> .....	487
<i>triserialis</i> , <i>Muraenopsis</i> .....	384	<i>Blue-back</i> .....	514, 2819
<i>Ophichthus</i> .....	384	<i>Brook</i> .....	506
<i>Ophichthys</i> .....	384	<i>Bull</i> .....	507
<i>triseriata</i> , <i>Platyrrhina</i> .....	60	<i>Coast Range</i> .....	500
<i>triseriatus</i> , <i>Platyrrhinoids</i> .....	65, 66	<i>Colorado River</i> .....	496
<i>Rhinobatus</i> .....	66	<i>Columbia River</i> .....	492
<i>tresignatum</i> , <i>Mexostoma</i> .....	179	<i>Cut-throat</i> .....	487, 492, 493
<i>Trisotropis</i> .....	1160, 1172	<i>Dolly Varden</i> .....	507
<i>aguajl</i> .....	1175	<i>Dublin Pound</i> .....	507
<i>bonaci</i> .....	1175	<i>Gila</i> .....	226
<i>brunneus</i> .....	1175	<i>Golden of Mount Whitney</i> .....	503
<i>calliurus</i> .....	1186	<i>Great Lake</i> .....	504
<i>camelopardalis</i> .....	1137	<i>Green-back</i> .....	497
<i>cardinulis</i> .....	1174	<i>Kamchatka Salmon</i> .....	2818
<i>chlorostomus</i> .....	1170	<i>Kamloops</i> .....	499
<i>dimidiatus</i> .....	1179	<i>Kern River</i> .....	502
<i>falcatus</i> .....	1185	<i>Lac de Marbre</i> .....	515
<i>interstitialis</i> .....	1179	<i>Lake Tahoe</i> .....	493, 2870
<i>microlepis</i> .....	1178	<i>Mackinaw</i> .....	504

Page.		Page.		Page.
1172	Trout McCloud River Rainbow	502	Trygon sayi	86
1187	Nissae	503	tuberculata	84
1184	No-she	503	Trygonobatus torpedinus	81
1178	of Utah Lake	495	Trygonorhina alveata	65
1187	Oquassa	514	Tschawytscha	479
1443	Rainbow	500	tschawytscha, Oncorhynchus	479
2085	Red Rock	1872	Salmo	480
1648	Red-spotted	507	tschawytschniformis, Salmo	478
1648	Rio Grande	495	tsiltcoosensis, Catostomus	2793
2086	Rock	1866, 1867	tsuppitch, Fario	493
272	Rocky Mountain	487	Oncorhynchus	481
111	Salmon	497	Salmo	481, 495
111	Sea	1407	tuberculata, Dasibatis	84
111	Silver	493	Enia	84
1236	Speckled	506	Trygon	84
1311	of Lake Crescent	2821	tuberculatus, Catostomus	186
1311	Sunapee	511	Spheroides	1733
1834	Truckee	493	tuberculé, Le Sphéroïde	1733
1586	Waha Lake	496	tuberculé, Raie	84
1585	Yellow-fin	496	tudes, Goblesox	2333
1653	Yellowstone	493	Salmo	508
171	Tront Perch	782, 784	Sphyrna	44
754, 757	troubridgii, Abeona	1497	Zygena	44
2268	Holconotus	1497	tuditana, Clhola	253
111	Homalopomus	2531	tuditanis, Hybopsis	253
111	Trucha	819	tuditanus, Hypargurus	253
1741	Truckee Trout	493	Leuciscus	253
1278	truculentus, Cherophthalmus	542	Tullibee	473
1279	True Eels	346	tullibee, Argyrosomus	473
1279	Fishes	97	bisselli	473
20	Sardines	422	Coregonus	473
1537	Sharks	21	bisselli	473
122, 2757	Trumpet-fish	754, 756, 759	Salmo (Coregonus)	473
122, 2758	truncata, Belone	714, 715	tumidus, Carpiodes	167
1552	Malthæa	2738	Chironectes	2717
483	Ranzania	1755, 1756	Lophius	2716
487	truncatus, Blennius	2381	Tuna	870
514, 2819	Dinectus	106	tunicata, Liparis	2121, 2128
500	Orthogoriscus	1756	tunicatus, Liparis	2120
507	Salmo	499	Tunnies	869
500	Saurus	533	Little	868
496	Scarus	1641	Tunny	870
492	Tetrodon	1756	Little	869
7, 492, 493	Trunk-fish	1720, 1721, 1722, 1723	Turbot	1701
507	Spotted	1723	Ocean	1706
507	Three-angled	1721	Tribo	2608
226	Trutta	483, 486, 487	turcesius, Pseudoscarus	1659
503	trutta, Salmo	487	Scarus	1658
504	Trutte	483	Turdus cauda convexa	1145
497	Trycherodon	247	cinereus peltatus	1373
2818	megalops	240	flavus	1583
499	Trygon	82	oculo radiato	1591, 1703
502	gymnura	84	pinis branchialibus	1257
515	hastata	84	rheuboidalis	1691
493, 2870	osteosticta	84	turgidus, Tetrodon	1733
504	sabina	85	nepelus	1733

	Page.		Page.
turneri, <i>Lycodalepis</i> .....	2469	tyrannus, <i>Clupea</i> .....	434
<i>Lycodes</i> .....	2469	Tyrant Fish .....	886
tnscumbia, <i>Etheostoma</i> .....	1100		
<i>Psychromaster</i> .....	1100	Ugly Fish .....	137
Two-spined Stickelback .....	748	uhleri, <i>Citharichthys</i> .....	2684
Tyee Saluon .....	479	Uhema .....	1371
Tylosurus .....	708	<i>lefroyi</i> .....	1371
<i>acus</i> .....	716	Ulca .....	2021
<i>almeida</i> .....	715	<i>marmorata</i> .....	2021
<i>angusticeps</i> .....	712	Ulcina .....	2088
<i>ardeola</i> .....	713	Ulka .....	1974
<i>caribbeus</i> .....	717	Ulke .....	1974
<i>contrainii</i> .....	717	Ulocentra .....	1047, 2851
<i>crassus</i> .....	716	<i>davisonii</i> .....	1049
<i>diploænia</i> .....	712	<i>gilberti</i> .....	1049, 2852
<i>enryops</i> .....	711	<i>histrio</i> .....	1050, 1051
<i>exilis</i> .....	714	<i>meadlie</i> .....	2852
<i>fodiator</i> .....	715	<i>phlox</i> .....	1052
<i>galeatus</i> .....	716	<i>simotera</i> .....	1051, 2853
<i>gladius</i> .....	716	<i>stigmata</i> .....	1047
<i>hians</i> .....	718	<i>verecunda</i> .....	1047
<i>longirostris</i> .....	714	ulochir, <i>Paraliparis</i> .....	21..
<i>marinus</i> .....	714	ulvæ, <i>Xiphidion</i> .....	2424
<i>microps</i> .....	712	<i>Xiphistes</i> .....	2423
<i>notatus</i> .....	710, 711	Ulvaria .....	2440
<i>pacificus</i> .....	716	<i>subbifurcata</i> .....	2440
<i>raphidoma</i> .....	715	Ulvicola .....	2413, 2860
<i>sagitta</i> .....	711	<i>sanctæ-rosæ</i> .....	2413
<i>scapularis</i> .....	711	umatilla, <i>Agosia</i> .....	313
<i>sierrita</i> .....	713	Umbla .....	506
<i>stolzmanni</i> .....	713	<i>minor marina</i> .....	823
<i>subtruncatus</i> .....	711	umbra, <i>Salmo</i> .....	509
<i>timucu</i> .....	711	Umbræ .....	623, 2807
Tympanomium .....	1753	<i>delicatissima</i> .....	621
<i>planci</i> .....	1754	<i>limi</i> .....	623
Tyntlastes .....	2262	<i>pygmea</i> .....	624
<i>brevis</i> .....	2262	<i>pygmaea</i> .....	624
<i>sagitta</i> .....	2263	umbratilis, <i>Alburnellus</i> .....	299
Typhlichthys .....	704, 2835	<i>Alburnus</i> .....	299
<i>rosæ</i> .....	2835	<i>Minnilus</i> .....	299
<i>subterraneus</i> .....	704, 2835	<i>Notropis</i> .....	298
Typhlogobius .....	2261	<i>ardens</i> .....	301
<i>californiensis</i> .....	2262	<i>atripes</i> .....	300
Typhlopaaras .....	2729	<i>cyanocephalus</i> .....	300
<i>shufeldti</i> .....	2731	<i>fasciolaris</i> .....	301
typica, <i>Sciæna</i> .....	1448	<i>lythrinus</i> .....	300
Typical Sharks .....	19	<i>matutinus</i> .....	301
typicus, <i>Ophioseion</i> .....	1448	<i>punctulatus</i> ..	301
<i>Plargyrus</i> .....	283	<i>umbratilis</i> .....	299
typus, <i>Achirophichthys</i> .....	388	Umbride .....	622
<i>Skeponopodus</i> .....	892	umbrifer, <i>Notropis</i> .....	274
tyrannus, <i>Anguilla</i> .....	348	<i>Urolophus</i> .....	2752
<i>Brevoortia</i> .....	433	Umbrina .....	1405
<i>aurea</i> .....	434	<i>alburnus</i> .....	1475
<i>brevicandata</i> ..	434	<i>analis</i> .....	1468
<i>patronus</i> .....	434	<i>arenata</i> .....	1474

Page.		Page.		Page.
434	<i>Umbrina broussonetii</i> .....	1466	<i>unicolor</i> , <i>Bryttus</i> .....	1001
886	<i>coroides</i> .....	1466	<i>Chaetodon</i> .....	1676
	<i>dorsalis</i> .....	1469	<i>Holocentrus</i> .....	1192
137	<i>elongata</i> .....	1476	<i>Hypoplectrus</i> .....	1190, 1192
2684	<i>furnleri</i> .....	1463	<i>aberrans</i> .....	1193
1371	<i>galapagorum</i> .....	1468	<i>occensus</i> .....	1193
1371	<i>gracilis</i> .....	1474	<i>atkinsi</i> .....	1193
2021	<i>januaria</i> .....	1474	<i>bovinus</i> .....	1193
2021	<i>littoralis</i> .....	1477	<i>chlorurus</i> .....	1193
2088	<i>martinicensis</i> .....	1474	<i>crocotus</i> .....	1192
1974	<i>nasus</i> .....	1473	<i>gummigutta</i> .....	1192
1974	<i>nebulosa</i> .....	1475	<i>guttavarius</i> .....	1192
47, 2851	<i>panamensis</i> .....	1473	<i>indigo</i> .....	1193
1049	<i>phalena</i> .....	1475	<i>negricans</i> .....	1192
49, 2852	<i>roncador</i> .....	1407	<i>pinnivarius</i> .....	1192
50, 1051	<i>sinsloe</i> .....	1468	<i>puella</i> .....	1192
2852	<i>undulata</i> .....	1467, 1476	<i>vitulinus</i> .....	1192
1052	<i>xanti</i> .....	1467, 1468	<i>Perca</i> .....	1192
51, 2853	<i>umbrosa</i> , <i>Cliola</i> .....	273	<i>Petromyzon marinus</i> .....	10
1047	<i>Cyprinella</i> .....	273	<i>Serranus</i> .....	1192
10	<i>Lepidopsetta</i> .....	2642	<i>Solca</i> .....	2702
21	<i>Narcine</i> .....	78	<i>Soleotalpa</i> .....	2703
2424	<i>umbrosus</i> , <i>Eques acuminatus</i> .....	1487	<i>Unicorn Fish</i> .....	1719
2423	<i>Esox</i> .....	627	<i>unicornis</i> , <i>Citharichthys</i> .....	2683
2440	<i>Gymnothorax</i> .....	390	<i>Unicornu pisces bahamensis</i> .....	1719
2440	<i>Platichthys</i> .....	2643	<i>unicornus</i> , <i>Balistes</i> .....	1720
13, 2869	<i>Pleuronectes</i> .....	2643	<i>unifasciatus</i> , <i>Hemirhamphus</i> .....	720, 721
2413	<i>Sebastichthys</i> .....	1807	<i>Hyporhamphus</i> .....	720
313	<i>Sebastodes</i> .....	1807	<i>unimaculata</i> , <i>Perca</i> .....	1360
506	<i>Umbrula</i> .....	1469, 1471, 1476	<i>unimaculatus</i> , <i>Archosargus</i> .....	1359, 1360
823	<i>Unbarana</i> .....	411	<i>Argyreosus</i> .....	934
509	<i>uncinatus</i> , <i>Artediellus</i> .....	1905, 1906	<i>Clinus</i> .....	2441
23, 2807	<i>Centridermichthys</i> .....	1906	<i>Grammistes</i> .....	1360
621	<i>Cottus</i> .....	1906	<i>Sargus</i> .....	1360
623	<i>Icolus</i> .....	1906	<i>Stichæus</i> .....	2441
624	<i>uncompagre</i> , <i>Xyranchen</i> .....	184	<i>uninotata</i> , <i>Heterandria</i> .....	687
624	<i>undecimale</i> , <i>Hemulon</i> .....	1300	<i>uninotatus</i> , <i>Ctenlabrus</i> .....	1577
299	<i>undecimilis</i> , <i>Centropomus</i> .....	1118	<i>Girardinus</i> .....	687
299	<i>Sclera</i> .....	1119	<i>Lutjanus</i> .....	1271
299	<i>undecimradiatus</i> , <i>Centropomus</i> .....	1119	<i>Mesoprion</i> .....	1271
298	<i>undulata</i> , <i>Muraenopsis</i> .....	403	<i>unionensis</i> , <i>Centropomus</i> .....	1122
301	<i>Perca</i> .....	1462	<i>univittatus</i> , <i>Apodichthys</i> .....	2412
300	<i>Umbrina</i> .....	1467, 1476	<i>Upeneus</i> .....	857, 2843
300	<i>undulatus</i> , <i>Monticirrhus</i> .....	1476	<i>baiteatus</i> .....	860
301	<i>Micropogon</i> .....	1461	<i>dentatus</i> .....	859
300	<i>Ostracion</i> .....	1724	<i>flavovittatus</i> .....	860
301	<i>Rhinobatus</i> .....	63	<i>grandisquamis</i> .....	860
301	<i>Scomber</i> .....	867	<i>maculatus</i> .....	858
299	<i>undulosus</i> , <i>Serranus</i> .....	1181	<i>martinicus</i> .....	859
622	<i>unerarak</i> , <i>Ophidium</i> .....	2477	<i>parvus</i> .....	859
274	<i>Unibranchapertura</i> .....	342	<i>punctatus</i> .....	858
2752	<i>grisea</i> .....	342	<i>rathbuni</i> .....	857
1465	<i>humaculata</i> .....	342	<i>tetraspilus</i> .....	860
1475	<i>lineata</i> .....	342	<i>xanthogrammus</i> .....	860
1468	<i>marmorata</i> .....	342	<i>Upselonphorus</i> .....	2306
1474	<i>unicolor</i> , <i>Ammocetes</i> .....	10	<i>guttatus</i> .....	2310

	Page.		Page.
Upselonophorus y-graecum .....	2308	Urolophus jamaicensis .....	81
uracantha, Loricaria .....	158	mundus .....	81
Uraleptus .....	2545	nebulosus .....	80, 2752
maraldi .....	2545	rogersi .....	2752, 2753
Uranichthys .....	381	torpedinus .....	81
brachycephalus .....	382	umbifer .....	2752
havannensis .....	382	Uronectes parii .....	2478
Uranidea .....	1963	urophthalmus, Heros .....	1537
aspera .....	1944	Urophycis .....	2552
bendirei .....	1964	chesteri .....	2556
boleoides .....	1968	chuss .....	2555
cognata .....	1955	cirratus .....	2553
formosa .....	1960	earli .....	2554
franklini .....	1967	floridanus .....	2554
gobioides .....	1968	regius .....	2553
gracilis .....	1968	tenulis .....	2555
greenel .....	1965	Uropsetta .....	2624
gulosa .....	1945	californica .....	2626
hoylei .....	1969	Uropterygius .....	403
kumlienii .....	1967	necturus .....	404
marginata .....	1905	urostigma, Cliola .....	275
microstoma .....	1958	Urotrygon .....	80
pollicaris .....	1954	mundus .....	81
punctulata .....	1949	Uroxis .....	82
quiescens .....	1968	ursinus, Salmo .....	505
rhothea .....	1947	urus, Bubalichthys .....	164
ricel .....	1953	Carploides .....	164
richardsoni .....	1952	Ictiobus .....	164
semiscabra .....	1950	Sclerognathus .....	165
splota .....	1953, 1962	ustus, Callyodon .....	1624
tenuis .....	1966	Cryptotomus .....	1624
vheeleri .....	1950	Utah Lake Chub .....	232
viscosa .....	1968	Mullet .....	179
uranidea, Cottogaster .....	1044	Trout .....	495
Etheostoma .....	1045	utah, Salmo .....	495
uranops, Phenacobius .....	304	utawana, Catostomus .....	179
Uranoscopidae .....	2305	uter, Catulus .....	25
Uranoscopinae .....	2306	Uwo Aka .....	1833
Uranoscopus anoplos .....	2308	Vaca .....	1190
y-graecum .....	2308	Vacas .....	1189
uranoscopus, Ceratias .....	2730	vacca, Capriodes .....	168
Mancalias .....	2729	Damalichthys .....	1519
Uraptera .....	66	Ditrema .....	1510
binoculata .....	73	Vacuocna .....	1427
Uraspis .....	916, 918, 926	vafer, Myrophis .....	372
Uriphaton .....	1143	vagrans, Chirostoma .....	795
Urocentrus .....	2414, 2415	Kirtlandia .....	794
pictus .....	2416	lacinlata .....	2840
Uroconger .....	358	Menidia .....	795
vicinus .....	358	lacinlata .....	795
Urolophinae .....	70	vahlhi, Lycodes .....	2463
Urolophus .....	70	Vallantia .....	1054, 1060
aspidurus .....	81	camura .....	1060
asterias .....	82, 2752	chlorosoma .....	1060
goulei .....	81	valenciennel, Smnragdus .....	2228
halleri .....	80	Valenciennellus .....	577

	Page.		Page.
81		<i>velifera</i> , <i>Agosia</i> .....	212
81		<i>velox</i> , <i>Carcharhinus</i> .....	2747, 2748
2752		<i>Chiola</i> .....	253
2753		<i>Euleptorhamphus</i> .....	724
81		<i>venadorum</i> , <i>Mycteropereca</i> .....	1186
2752		<i>Venetica</i> .....	365
2478		<i>procera</i> .....	365
1537		<i>venenosus</i> , <i>Mycteropereca</i> .....	1172
2552		<i>apna</i> .....	1173, 1174
2556		<i>guttata</i> .....	1174
2555		<i>marina</i> .....	1172
2553		<i>venenosus</i> , <i>Epinophelus</i> .....	1172
2554		<i>venosa</i> , <i>Meletta</i> .....	426
2554		<i>venosus</i> , <i>Aluterus</i> .....	1719
2553		<i>ventralis</i> , <i>Arnoglossus</i> .....	2670
2555		<i>Brosuophycis</i> .....	2503
2624		<i>Citharichthys</i> .....	2670
2626		<i>Cottus</i> .....	2008, 2009
403		<i>Dinenaticthys</i> .....	2503
404		<i>Novaculichthys</i> .....	1615
275		<i>Ogilbia</i> .....	2503
80		<i>Trichopsetta</i> .....	2669
81		<i>Xyrichtys</i> .....	1616
82		<i>ventre</i> , <i>Ostracion subrotundus</i>	
505		<i>glabro</i> .....	1749
164		<i>ventricosa</i> , <i>Apocope</i> .....	300
164		<i>Temnistia</i> .....	1936
164		<i>ventricosus</i> , <i>Blepsias</i> .....	1936
165		<i>Ceratichthys</i> .....	309
1624		<i>Cotylin</i> .....	2104
1624		<i>Cyclopterichthys</i> .....	2104
232		<i>Cyclopterus</i> .....	2104
179		<i>Salmo</i> .....	509
495		<i>Sebastes</i> .....	1829
495		<i>Sebastodes</i> .....	1829
179		<i>ventriosum</i> , <i>Scyllium</i> .....	25
25		<i>venusta</i> , <i>Chiola</i> .....	274
1833		<i>Cyprinella</i> .....	274
1190		<i>Emmcekia</i> .....	1602
1189		<i>Limia</i> .....	665
168		<i>Lucania</i> .....	665
1519		<i>venustus</i> , <i>Dinemus</i> .....	854
1510		<i>Notropis</i> .....	274, 275
1427		<i>Pseudojullis</i> .....	1602
372		<i>Xyrichtys</i> .....	1610
795		<i>verany</i> , <i>Cybbium</i> .....	877
794		<i>Verasper</i> .....	2618
2840		<i>moseri</i> .....	2619
795		<i>Verde</i> , <i>Johnny</i> .....	1195
795		<i>Mojarra</i> .....	1538
2463		<i>Morena</i> .....	396
1060		<i>Pudiano</i> .....	1590, 1591
1060		<i>Verdes</i> , <i>Cabrillas</i> .....	1194
2228		<i>verecunda</i> , <i>Ulocentra</i> .....	1049
577		<i>verecundum</i> , <i>Etheostoma</i> .....	1050
		<i>Lepidon</i> .....	2543

	Page.		Page.
Verilus.....	1283	vetula, Pseudoscarus.....	1650
sordidus.....	1284	Scarus.....	1647, 1649
Verma.....	374	vetulus, Parophrys.....	2640
kendalli.....	375	Pleuronectes.....	2641
Vermelho, Pudianio.....	1583	vexillare, Boleosoma nigrum.....	1058
vernicularis, Corvina.....	1453	Etheostoma nigrum.....	1058
Ophioacton.....	1452	vexillaris, Sebastelchthys.....	1822
Sciæna.....	1452, 1453	Sebastodes.....	1821
Sparus.....	1271	vexillarium, Holocentrum.....	852
vernivolatus, Esox.....	627	vexillarius, Holocentrus.....	852
Exocoetus.....	740	Vexillifer.....	2495
Lucius.....	627, 2827	vheeleri, Uranidea.....	1950
Xyrichthys.....	1619	Viajaca.....	1539
vermiformis, Neoconger.....	302	vicensis, Trigla digitis.....	2183
Vernigilla, Pesca.....	1811	vicina, Murena.....	394
vernalis, Clupea.....	426	Murenophis.....	394
Gobio.....	321	Sidera.....	394
Hydrargyra.....	639	vicinalis, Icelus.....	1916
Hyodon.....	413	vicinus, Gymnothorax.....	304
Pomolobus.....	426	Lycodontis.....	394
Scomber.....	806	Uroconger.....	358
vernullas, Batrachoides.....	2316	victoriae, Moxostoma.....	187
verreauxi, Conger.....	355	Vicuda.....	824 *
verros, Coosyphus.....	1583	Vieja.....	1635, 1636, 1649
Lutjanus.....	1583	Colorada.....	1639
verrilli, Gymnothorax.....	394	Muger.....	1639
Lycenchelys.....	2470, 2471	Viejas.....	1627
Lycodes.....	2471	vigil, Ioa.....	1065
Lycodontis.....	393	vigilax, Ceratichthys.....	253
Sidera.....	394	Ctiola.....	253
verrucosa, Occa.....	2043	Villarius.....	2780
verrucosus, Brachyopsis.....	2044	dugesii.....	2789
Cottus.....	1980	pricei.....	2790
Diodon.....	1749	villosa, Clupea.....	521
Myoxocephalus.....	1970	villosus Cottus.....	2022
Verrugata.....	1476	Mailotus.....	520
Verrugato.....	1462, 1463	vincente, Sicydium.....	2207
versicolor, Abramis.....	250	vinoiguerra, Exocoetus.....	734
Girardinus.....	689	Exonantes.....	2836
Heterandria.....	688	Vinciguerra.....	577
Labrus.....	1346	attenuata.....	577
Malacocterus.....	2359	vinctipes, Nanostoma.....	1075
Myxodes.....	2359	vinctus, Blennius.....	2382
Pecilichthys.....	1089	Caranx.....	918
versipunctatus, Gymnothorax.....	394	Fundulus.....	637, 2827
vertagus, Gadus.....	2541	vinolentus, Lethotremus.....	2101
verticalis, Echenois.....	2270	viola, Antimora.....	2544
verticalis, Pleuronichthys.....	2638	Eques.....	1486
verus, Carcharias.....	50	Haloporphyrus.....	2544
Conger.....	355	violacea, Biptnula.....	878
vespertilio, Holoridnus.....	90	violaceus, Apodichthys.....	2427
Malthuca.....	2737	Cebedichthys.....	2427
Ogocephalus.....	2737	Escolar.....	4843
Rhinoptera.....	90	Thyrstops.....	879
veternus, Podothecus.....	2063, 2064	Viper-fishes.....	578, 584
vetula, Ballistes.....	1703	virens, Chlorichthys.....	1610

Page.		Page.		Page.
1650	<i>virens</i> , <i>Gadus</i> .....	2534	<i>vitrea</i> , <i>Perca</i> .....	1021
1649	<i>Pollachius</i> .....	2534	<i>vitreum</i> , <i>Stizostedion</i> .....	1021
2640	<i>Scarus</i> .....	1640	<i>vitreus</i> , <i>Poecilichthys</i> .....	1065
2641	<i>Thalassoma</i> .....	1611, 2859	<i>vitta</i> , <i>Xyrichtys</i> .....	1617
1058	<i>virescens</i> , <i>Apodichthys</i> .....	2412	<i>vittata</i> , <i>Argyrotenia</i> .....	833
1058	<i>Clupea</i> .....	426	<i>Channomuræna</i> .....	404
1822	<i>Cynoscion</i> .....	1415	<i>Cluola</i> .....	258
1821	<i>Gallus</i> .....	932	<i>Clupea</i> .....	421
852	<i>Gymnothorax</i> .....	394	<i>Codoma</i> .....	258
852	<i>Lycodontis</i> .....	394	<i>Echeneis</i> .....	2269
2495	<i>Otolithus</i> .....	1415	<i>Gymnomuræna</i> .....	404
1950	<i>Pantosteus</i> .....	171, 172	<i>Hemitremia</i> .....	242
1539	<i>virgata</i> , <i>Coryphæna</i> .....	953	<i>Inermia</i> .....	1366
2183	<i>virgatulus</i> , <i>Goblesox</i> .....	2333	<i>Lepidomeda</i> .....	328
394	<i>virgatum</i> , <i>Etheostoma</i> .....	1093	<i>Poecilia</i> .....	692
394	<i>virgatus</i> , <i>Delolepis</i> .....	2442	<i>vittatum</i> , <i>Exoglossum</i> .....	327
394	<i>Poecilichthys</i> .....	1093	<i>vittatus</i> , <i>Amniodytes</i> .....	833
1916	<i>virginalis</i> , <i>Salar</i> .....	495	<i>Cyprinus</i> .....	307
304	<i>Salmo clarkii</i> .....	2819	<i>Emmlichthys</i> .....	1305, 1366
394	<i>mykiss</i> .....	495	<i>Erythrichthys</i> .....	1366
358	<i>Scarus</i> .....	1647	<i>Esox</i> .....	628
187	<i>virginianus</i> , <i>Acanthocottus</i> .....	1976	<i>Ichthyophis</i> .....	404
824	<i>Cottus</i> .....	1976	<i>Leuciscus</i> .....	282
1, 1649	<i>Scorpius</i> .....	1076	<i>Proceros</i> .....	102
1639	<i>virginicum</i> , <i>Pristipoma</i> .....	1323	<i>Sarchirus</i> .....	110
1639	<i>virginicus</i> , <i>Anisotremus</i> .....	1322, 1323	<i>Siphoteles</i> .....	244
1627	<i>Polydactylus</i> .....	829	<i>Sparus</i> .....	1323
1065	<i>Polynemus</i> .....	830	<i>Symbranchus</i> .....	342
253	<i>Pomadays</i> .....	1323	<i>vitula</i> , <i>Sargus</i> .....	1364
253	<i>Sparus</i> .....	1323	<i>vitulinum</i> , <i>Plectropoma</i> .....	1192
2780	<i>viride</i> , <i>Ophidium</i> .....	2477	<i>vitulinus</i> , <i>Hypoplectrus unicolor</i> .....	1192
2789	<i>Sparisoma</i> .....	1678	<i>vitulus</i> , <i>Carpiodes</i> .....	165
2790	<i>viridescens</i> , <i>Atherina</i> .....	800	<i>Viuva</i> .....	1788
521	<i>Fundulus</i> .....	641	<i>vivanet</i> , <i>Bodianus</i> .....	1257
2022	<i>Osmerus</i> .....	523	<i>vivanus</i> , <i>Anthias</i> .....	1224
520	<i>Sphyræna</i> .....	826	<i>Hemianthias</i> .....	1229
2207	<i>Syngnathus</i> .....	771	<i>Lutjanus</i> .....	1264, 1265
734	<i>viridipallidus</i> , <i>Gobius</i> .....	2259	<i>Mesoprion</i> .....	1263
2836	<i>viridis</i> , <i>Amia</i> .....	113	<i>Neomenis</i> .....	1262
577	<i>Centrarchus</i> .....	992	<i>Pronotogrammus</i> .....	1224
577	<i>Centropomus</i> .....	1118	<i>vivax</i> , <i>Ammocrypta pellucida</i> .....	1063
1075	<i>Diacope</i> .....	1246	<i>Cluola</i> .....	253
2382	<i>Esox</i> .....	110	<i>vivipara</i> , <i>Poecilia</i> .....	691
918	<i>Evoplites</i> .....	1246	<i>Viviparus Perch</i> .....	1498
7, 2827	<i>Genyerogeto</i> .....	1246	<i>viviparus</i> , <i>Lophius</i> .....	2715
2101	<i>Gymnelis</i> .....	2477	<i>Sebastes</i> .....	1761
2544	<i>Lepidosteus</i> .....	111	<i>vlamingii</i> , <i>Coryphæna</i> .....	953
1486	<i>Lutjanus</i> .....	1246	<i>Voilier</i> .....	891
2544	<i>Piscis bahamensis</i> .....	1638	<i>Volador</i> .....	740, 2183
878	<i>Scarus</i> .....	1638	<i>volador</i> , <i>Exocoetus</i> .....	733
2427	<i>viscosa</i> , <i>Uranidea</i> .....	1968	<i>Rubio</i> .....	2164
2427	<i>viscosus</i> , <i>Cottus</i> .....	1968	<i>Voladora</i> , <i>Aguja</i> .....	891
4843	<i>Silurus</i> .....	143	<i>Volantin</i> .....	914
879	<i>vitrea</i> , <i>Ammocrypta</i> .....	1065	<i>volitans</i> , <i>Cephalacanthus</i> .....	2183
78, 584	<i>Ioa</i> .....	1064	<i>Dactylopterus</i> .....	2183
1610	<i>Lucloperca</i> .....	1022	<i>Exocoetus</i> .....	734, 736



	Page.		Page.
voltans, Trigla .....	2183	vulneratum, Etheostoma .....	1077
volucellus, Hybognathus .....	263	vulneratus, Nothonotus .....	1077
Hybopsis .....	263	Pælicthys .....	1077
Leuciscus .....	263	vulpeculus, Pimelodus .....	141
Notropis .....	263	Vulpes bahamensis .....	411
Vomer .....	933	vulpes, Albula .....	411
brasiliensis .....	934	Alopias .....	46
browni .....	934, 2846	Alopias .....	45
cayennensis .....	934	Esox .....	411
columbianus .....	934	Pimelodus .....	135
cubæ .....	934	Sebastodes .....	1835
curtus .....	934	Squalus .....	46
dominicensis .....	934	vulpinus, Squalus .....	46
dorsalis .....	934	Vulsiculus .....	2181
gabonensis .....	934, 2846	imberbis .....	2181
goreensis .....	934	vulsus, Agonns .....	2068
martinicensis .....	934	Podothecus .....	2068
noveboracensis .....	934	Stelgis .....	2067
sanctæ murtine .....	934	wabaahensis, Anguilla .....	348
sanctæ petri .....	934	Wachna Cod .....	2537
senegalensis .....	934	wagneri, Pimelodus .....	151
setipinnis .....	934, 2846	Rhandin .....	150, 151
spixii .....	2846	Waha Lake Trout .....	496
vomer, Argyreosus .....	936	Wahoo .....	876
Seleno .....	936	walkerii, Selanotius .....	49
Zeus .....	936	Wall-eyed Herring .....	426
vomerina, Athorina .....	793	Piko .....	1021
vomerinus, Anarrhichias .....	2447	Pollack .....	2536
Salarius .....	2400	Surf-fishes .....	1501
vorax, Lucius .....	628	Warmouths .....	991, 992
Mosoprion .....	1281	warreni, Boleichthys .....	1103
Platyulius .....	1281	Pælicthys .....	1103
Pseudorhombus .....	2026	Salmo .....	483
Ptychocheilus .....	227	Warn, Me .....	1829
Voraz .....	1280	watauga, Hybopsis .....	319
vulgaris, Acanthias .....	54	Weakfish, Bastard .....	1406
Ameiurus .....	140	Common .....	1407
Amiurus .....	140	Spotted .....	1409
Auxis .....	868	webbi, Embiotoca .....	1505
Brosminus .....	2561	webbii, Blennophis .....	2401
Conger .....	355	Ophiobleunius .....	2401
Gunnellus .....	2419	Webbug Sucker .....	180
Hippoglossus .....	2612	welaka, Notropis .....	2799
Liparis .....	2118	Welshman .....	848
Lumpus .....	2097	Wench, Old .....	1703
Merluccius .....	2530	West Indian Lancelet .....	3
Molva .....	2552	westphali, Teniophis .....	396
Mustelus .....	29	Whale Sharks .....	52
Pagrus .....	1357	wheatlandi, Casterosteus .....	749
Pimelodus .....	140	Whiffs .....	2678
Pomotis .....	1010	whipplei, Boleosoma .....	1096
Sphyrna .....	826	Cluola .....	279
Squatina .....	59	Etheostoma alabamæ .....	1005
Sturio .....	105	whippilii, Boleichthys .....	1006
Thynnus .....	870	Cyprinuella .....	279
vulnerata, Apocope .....	312		

Page.	Page.	Page.
1077	whiplil, <i>Etheostoma</i> .....	1095
1077	<i>Notropis</i> .....	278
1077	Whip-tailed Rays.....	70
141	Whirligig Mullet.....	818
411	White Bass.....	1132
411	Cat.....	134, 138
46	Croaker.....	1307
45	Grunt.....	1310
411	Hake.....	2555
135	Lake Bass.....	1132
1835	Mullet.....	189, 813
40	Perch.....	1133, 1134, 1484, 1501, 1509
46	Salmon of the Colorado.....	225
2181	Sea Bass of California.....	1413
2181	Sturgeon.....	104, 107
2068	Sucker.....	178, 192
2068	Surf-fish.....	1506
2067	White-bill.....	431
348	White-bone Porgy.....	1353
2537	White-eye.....	1021
151	Whitefish.....	433, 461, 2276
0, 151	Broad.....	404
496	Common.....	465
876	Humpback.....	466
49	Menominee.....	465
426	Mongrel.....	473
1021	Musquaw River.....	466
2536	Rocky Mountain.....	463
1501	Round.....	465
1, 992	Sault.....	466
1103	White-mouthed Drummer.....	1462
1103	White-nosed Suckers.....	190
483	Whiting.....	2530
1829	California.....	1477
319	Carolina.....	1474
1406	Northern.....	1475
1407	Sand.....	1474
1409	Silver.....	1477
1505	Surf.....	1477
2401	Whiting of Lake Winnipiscogee.....	460
2401	Widow-fish.....	1788
2401	Wife, Old.....	940, 1703
180	wilderi, <i>Lampetra</i> .....	13, 2745
2799	Will, Black.....	1199
848	williamsi, <i>Symphurus</i> .....	2711
1703	williamsoni, <i>Coregonus</i> .....	463
3	<i>clermontanus</i> .....	463
396	<i>Gasterosteus</i> .....	750
52	<i>Gasterosteus micro-</i>	
749	<i>cephalus</i> .....	751
2678	willoughbyi, <i>Acrotus</i> .....	973
1096	Willow Cat.....	2788
279	willoughbeii, <i>Canthidermis</i> .....	1707
1095	<i>Salmo</i> .....	509
1096	<i>Trematopsis</i> .....	1754
279	wilsoni, <i>Cottus</i> .....	1952
	Wilton Smelt.....	523
	winchell, <i>Hybopsis</i> .....	321
	Wind-fish.....	221
	Window Pines.....	2650, 2660
	Winniniah.....	487
	Winnipiscogee, Whiting.....	406
	Winter Flounders.....	2646
	Sucker.....	187
	Wolf Eel.....	2448
	Wolf-fish.....	595, 2445, 2447
	woolmani, <i>Paralichthys</i> .....	2628
	Worm Eels.....	370
	Wrasse-fishes.....	1571
	Wreckfish.....	1138, 1139
	wrighti, <i>Etheostoma</i> .....	1047
	Wry-mouth.....	2442, 2443
	wurdemanni, <i>Gobius</i> .....	2225
	xænocephalus, <i>Hybopsis</i> .....	289
	<i>Minnilus</i> .....	289
	<i>Notropis</i> .....	289
	xænura, <i>Chloa</i> .....	280
	<i>Codoma</i> .....	280
	xænurus, <i>Minnilus</i> .....	280
	<i>Notropis</i> .....	280
	xaniurus, <i>Catulus</i> .....	24
	Xanthichthys.....	1708
	<i>cicatricosus</i> .....	1709
	<i>mento</i> .....	1760
	<i>ringens</i> .....	1709
	xanthocephalus, <i>Ameiurus</i> .....	141
	<i>Silurus</i> .....	141
	xanthogrammus, <i>Upeneus</i> .....	860
	xanthomelas, <i>Anguilla</i> .....	348
	xanthops, <i>Odontoscion</i> .....	1427
	xanthopteron, <i>Hemulon</i> .....	1307
	xanthopteron, <i>Hemulon</i> .....	1307
	xanthopus, <i>Catostomus</i> .....	181
	xanthopygus, <i>Caranx</i> .....	921
	xanthosticta, <i>Mycteroperca bonaci</i> .....	1176
	xanthostignus, <i>Citharichthys</i> .....	2680
	xanthulum, <i>Cestrens</i> .....	1411
	xanthulus, <i>Cynoscion</i> .....	1410
	xanthurus, <i>Homoprion</i> .....	1434, 1459
	<i>Leiostomus</i> .....	1458
	<i>Pomacentrus</i> .....	1557
	<i>Rhombus</i> .....	966, 2849
	<i>Sciæna</i> .....	1459
	<i>Seserinus</i> .....	966
	<i>Sparus</i> .....	1346
	xanti, <i>Labrisomus</i> .....	2362, 2363
	<i>Rhyptions</i> .....	1231
	<i>Rypticus</i> .....	1231
	<i>Umbrina</i> .....	1467, 1468
	<i>Xenichthys</i> .....	1288
	xantusi, <i>Lepidopus</i> .....	2844, 4843

	Page.		Page.
<i>xenarcha</i> , Mycteroperca.....	1180	Xiphidae.....	893
<i>Xenarehi</i> .....	780, 785	<i>Xiphister</i> .....	2424
<i>xenachus</i> , Epinophelus.....	1180	<i>chirus</i> .....	2424
<i>xenanchon</i> , Galeichthys.....	2777	<i>mucosum</i> .....	2425
<i>Xenichthyma</i> .....	1244	<i>rupestris</i> .....	2426
<i>Xenichthys</i> .....	1287	<i>Xiphistes</i> .....	2423
<i>agassizii</i> .....	1287	<i>chirus</i> .....	2424
<i>californiensis</i> .....	1286	<i>ulvæ</i> .....	2423
<i>xunti</i> .....	1288	<i>Xiphophorus</i> .....	701
<i>xenops</i> .....	1288	<i>binaculatus</i> .....	678
<i>xenurus</i> .....	1015	<i>gillii</i> .....	692
<i>xenicus</i> , Fundulus.....	662	<i>gracilis</i> .....	683
<i>Xenisma</i> .....	633, 635, 648	<i>guntheri</i> .....	702
<i>catenata</i> .....	648	<i>helleri</i> .....	701, 702
<i>stellifer</i> .....	648	<i>Xurel</i> .....	909, 923
<i>xenisma</i> , Prionotus.....	2154	<i>del Castilla</i> .....	937
<i>Xenistius</i> .....	1286	<i>xyosterna</i> , Stellerina.....	2042
<i>californiensis</i> .....	1286	<i>xyosternus</i> , Brachyopsis.....	2043
<i>Xenochirus</i> .....	2079	<i>Xyrauchen</i> .....	184
<i>alascanus</i> .....	2081	<i>cypho</i> .....	184
<i>latifrons</i> .....	2082	<i>uncompa gre</i> .....	184
<i>pentacanthus</i> .....	2080	<i>Xyrichthyinae</i> .....	1575
<i>triacanthus</i> .....	2084	<i>Xyrichthys</i> .....	1017
<i>Xenocys</i> .....	1285	<i>cultratus</i> .....	1619
<i>jessiae</i> .....	1285	<i>infirmus</i> .....	1616
<i>xenodon</i> , Calotomus.....	1626	<i>jessiae</i> .....	1613
<i>Xenomi</i> .....	620	<i>lineatus</i> .....	1619
<i>Xenomystax</i> .....	360	<i>martincensis</i> .....	1617
<i>atrarius</i> .....	361	<i>modestus</i> .....	1610
<i>xenops</i> , Xenichthys.....	1288	<i>mundiceps</i> .....	1618
<i>xenopterus</i> , Cypsilurus.....	2830	<i>novacula</i> .....	1619
<i>Xenopterygii</i> .....	782, 2326	<i>psittacus</i> .....	1618, 1619
<i>xenostethus</i> , Sternias.....	1927	<i>rosipes</i> .....	1615
<i>Triglops</i> .....	1927	<i>ventralis</i> .....	1616
<i>Xenotis</i> .....	999, 1000, 1002	<i>venustus</i> .....	1610
<i>aureolus</i> .....	1003	<i>vermiculatus</i> .....	1610
<i>lythrochloris</i> .....	1003	<i>vitta</i> .....	1617
<i>xenura</i> , Kuhlla.....	1015	<i>xyris</i> , Sebastopsis.....	1835
<i>xenurus</i> , Xenichthys.....	1015	<i>Xyrula</i> .....	1612
<i>Xererpes</i> .....	2413	<i>jessiae</i> .....	1612, 1013
<i>fucorum</i> .....	2413	<i>Xystæma</i> .....	1372
<i>Xesurus</i> .....	1694	<i>cinereum</i> .....	1372
<i>clarionis</i> .....	1695	<i>Xyster</i> .....	1384
<i>latioclavius</i> .....	1695	<i>xyster</i> , Zapteryx.....	65
<i>punctatus</i> .....	1694, 1695	<i>Xystes</i> .....	2076
<i>Xiphias</i> .....	893	<i>axinophrys</i> .....	2076
<i>gladius</i> .....	894	<i>Xystophorus</i> .....	900
<i>imperator</i> .....	892	<i>Xystreurus</i> .....	2023
<i>makaira</i> .....	891	<i>hoplepis</i> .....	2023
<i>rondeleti</i> .....	894	<i>xystrodon</i> , Sparisoma.....	1630
<i>Xiphidinae</i> .....	2348	<i>Xystroperca</i> .....	1169, 1170, 1181
<i>Xiphidion</i> .....	2424	<i>Xystroplites</i> .....	1006
<i>cruciforme</i> .....	2425	<i>gillii</i> .....	1007
<i>mucosum</i> .....	2425	<i>longimanus</i> .....	1008
<i>rupestre</i> .....	2426	<i>xysturus</i> , Myrichthys.....	2802
<i>ulvæ</i> .....	2424	<i>Ophichthys</i> .....	2802

Page.		Page.		Page.
893	<i>xyeturus</i> , <i>Ophisurus</i> .....	376	<i>zancus</i> , <i>Dermatolepis</i> .....	2854
2424	<i>Pisodonophis</i> .....	376	<i>zanderi</i> , <i>Argyrocottus</i> .....	1995
2424			<i>zanemus</i> , <i>Ceraticichthys</i> .....	310
2425	<i>yalei</i> , <i>Ostracion</i> .....	1724	<i>Zanolepidae</i> .....	2862
2426	<i>yanagi</i> , <i>Nomal</i> .....	1830	<i>Zanolepis</i> .....	1876
2423	<i>Yarella</i> .....	583	<i>frenatus</i> .....	1877
2424	<i>blackfordi</i> .....	584	<i>Latipianis</i> .....	1876
2423	<i>yarrelli</i> , <i>Acipenser</i> .....	105	<i>Zapater</i> .....	898
701	<i>yarrowi</i> , <i>Agosia</i> .....	309	<i>Zaphotias</i> .....	2826
678	<i>y-canda</i> , <i>Gillichthys</i> .....	2252	<i>pedaliotus</i> .....	2826
692	<i>Quiletula</i> .....	2251, 2252	<i>Zaprora</i> .....	2850
693	<i>Yellow-backed Rockfish</i> .....	1822	<i>silenus</i> .....	2850
702	<i>Yellow Bass</i> .....	1134	<i>Zaproridae</i> .....	2849
701, 702	<i>Belly</i> .....	1001	<i>Zapteryx</i> .....	64
909, 923	<i>Cat</i> .....	139, 143	<i>exasperatus</i> .....	64
937	<i>Yellow-finned Grouper</i> .....	1155, 1172	<i>xyster</i> .....	65
2042	<i>Roncador</i> .....	1467	<i>zatropis</i> , <i>Siphostoma</i> .....	772
2043	<i>Trout</i> .....	497	<i>zebra</i> , <i>Acanthurus</i> .....	1691
184	<i>Yellow Fish</i> .....	1144	<i>Arbuciosa</i> .....	2341
184	<i>Gontfish</i> .....	859	<i>Asproperca</i> .....	1027
184	<i>Grouper</i> .....	1183	<i>Chetodon</i> .....	1691
1575	<i>Grunt</i> .....	1303	<i>Fundulus</i> .....	641, 647
1617	<i>Jack</i> .....	919	<i>Gerres</i> .....	1373
1619	<i>Mackerel</i> .....	921	<i>Goblesox</i> .....	2342
1616	<i>Perch</i> .....	1023	<i>Gobius</i> .....	2226
1613	<i>Pike</i> .....	1021	<i>Hydrargyra</i> .....	647
1619	<i>Sculpin</i> .....	1934	<i>Percina caprodes</i> .....	1627
1617	<i>Yellow-spotted Rockfish</i> .....	1826	<i>Pileoma</i> .....	1028
1619	<i>Yellowstone Trout</i> .....	493	<i>Psychrolutes</i> .....	2027
1618	<i>Yellow-tail</i> .....	902, 906, 1275, 1433	<i>zebrinus</i> , <i>Fundulus</i> .....	646
1619	<i>Creaker</i> .....	1407	<i>Zeide</i> .....	1659
18, 1619	<i>Rockfish</i> .....	1781	<i>Zeinae</i> .....	1660
1615	<i>y-grucum</i> , <i>Astroscopus</i> .....	2307	<i>zelotes</i> , <i>Dactyloscopus</i> .....	2303
1616	<i>Upsilonphorus</i> .....	2308	<i>Hernicranx</i> .....	2845
1619	<i>Uranoscopus</i> .....	2308	<i>Zenion</i> .....	1661
1619	<i>Yuriria</i> .....	314, 315, 321	<i>hololepis</i> .....	1661
1617			<i>Zenopsis</i> .....	1660
1835	<i>zacentrus</i> , <i>Sebastesichthys</i> .....	1815	<i>ocellatus</i> .....	1660
1612	<i>zachirus</i> , <i>Glyptocephalus</i> .....	2658	<i>zenopterus</i> , <i>Exocoetus</i> .....	738
12, 1613	<i>Zaclemus</i> .....	1477, 1478, 1480	<i>Zeoidea</i> .....	1950
1372	<i>zadocki</i> , <i>Cylindrosteus</i> .....	111	<i>zephyreus</i> , <i>Astroscopus</i> .....	2300
1372	<i>Zalemblus</i> .....	1499	<i>Pristis</i> .....	2749
1384	<i>rosaceus</i> .....	1500	<i>Zeisticelus</i> .....	1990
65	<i>Zalientes</i> .....	2738	<i>profundorum</i> .....	1990
2076	<i>elater</i> .....	2738	<i>Zestidium</i> .....	1439, 1442
2076	<i>Zalocys</i> .....	2848	<i>Zestis</i> .....	1439, 1440
900	<i>stilho</i> .....	2848	<i>zestocarus</i> , <i>Stellifer</i> .....	1445
2623	<i>Zalopyr</i> .....	1795, 2680	<i>zestus</i> , <i>Nebris</i> .....	1417
2623	<i>Zalypnus</i> .....	2246	<i>Zeus capillaris</i> .....	936
1630	<i>cyclolepis</i> .....	2246	<i>ciliaris</i> .....	932
70, 1181	<i>emblematicus</i> .....	2247	<i>crinitus</i> .....	932
1006	<i>Zancurus</i> .....	890	<i>gallus</i> .....	936
1007	<i>Zanclus</i> .....	1687	<i>geometricus</i> .....	936
1008	<i>canescens</i> .....	1688	<i>guttatus</i> .....	955
2802	<i>centrognathus</i> .....	1688	<i>imperialis</i> .....	955
2802	<i>ornatus</i> .....	1687, 1688	<i>luna</i> .....	955

	Page.		Page.
Zoos niger .....	936	zonurus, Malacocottus .....	1994
ocellatus .....	1061	Zophendum .....	308
opah .....	955	australe .....	212
quadratus .....	1668	plumbeum .....	216
regius .....	955	siderium .....	314
rostratus .....	936	zopherus, Potamocottus .....	1952
setipinnis .....	934	zophochir, Ophichthus .....	385
stromii .....	955	Ophichthus .....	385
vomere .....	936	zostera, Hippocampus .....	778
zeylonicus, Leuciscus .....	415	zosterura, Evermannia .....	2256
Ziphotheca .....	887	zosterurum, Gobiosoma .....	2257
tetradena .....	887	zunnensis, Leuciscus .....	227
Zoarcas .....	2456	Zygæna .....	43
anguillaris .....	2457	lewini .....	45
elongatus .....	2457	malleus .....	45
fimbriatus .....	2457	subarcuata .....	45
gronovii .....	2457	tiburo .....	44
labrosus .....	2457	tudes .....	44
polaris .....	2469	zygæna, Sphyrna .....	45
Zoarchidæ .....	2455	Squalus .....	45
Zoarchus .....	2456	Zygobatis .....	90
zoarchus, Lycodes .....	2464	Zygonectes .....	650, 2828
Zoarcidæ .....	2455	atrilatus .....	682
Zoarcina .....	2455	auroguttatus .....	654, 2829
zonale, Etheostoma .....	1075	brachypterus .....	632
arcansanum .....	1075	chrysotus .....	656
Nanostoma .....	1075	cingulatus .....	655, 656
zonalis, Astatichthys .....	1075	craticula .....	657
Pælicthys .....	1075	dispar .....	650
zonata, Seriola carolinensis .....	902	escambia .....	658
zonatum, Elasmoma .....	982	floripinnis .....	651
zonatus, Alburnus .....	285	funduloides .....	650
Chaetodipterus .....	1668	guttatus .....	658
Ephippus .....	1669	henshalli .....	653, 2829
Esox .....	639	hieroglyphicus .....	658
Fundulus .....	657	inurus .....	682
Notropis .....	285	jenkinsi .....	652
Scumber .....	902	lateralis .....	659
Zygonectes .....	659	lineatus .....	649
Zonichthys .....	901, 904	lineolatus .....	657
bosci .....	905	lucia .....	655
coronatus .....	905	macdonaldi .....	651
declivis .....	905	mannii .....	664
gigas .....	903	melanops .....	682
zonifer, Clivus .....	2359	notatus .....	659
Erilepis .....	1863	notii .....	657
zonifer, Myriolepis .....	1863	pulchellus .....	659
Zygonectes .....	657	pulvereus .....	652
zonipectus, Pomacanthodes .....	1682	rubrifrons .....	654, 2829
Pomacanthus .....	1681	scadicus .....	654
zonistius, Luxilus .....	285	zonatus .....	659
Minnilus .....	285	Zygonectes zonifer .....	657
Notropis .....	277, 285	Zygonectes .....	633, 635
Zonogobius .....	2210	zyopterus, Galeorhinus .....	32
zonopæ, Jordania .....	1884	Zyphothyca .....	883
Zonoscion .....	1477, 1478, 1479		

Page.

1904  
308  
212  
216  
314  
1952  
385  
385  
778  
2256  
2257  
227  
43  
45  
45  
45  
44  
44  
45  
45  
90  
2828  
682  
2829  
682  
656  
656  
657  
650  
658  
651  
650  
658  
2829  
658  
682  
652  
659  
649  
657  
655  
651  
604  
682  
659  
657  
659  
652  
2829  
654  
659  
657  
635  
32  
883

