




IMAGE EVALUATION TEST TARGET (MT-3)



6"



Photographic
Sciences
Corporation

23 WEST MAIN STREET
WEBSTER, N.Y. 14580
(716) 872-4503

18
20
22
24
26
28
30
32
34
36
CIHM/ICMH
Microfiche
Series.

CIHM/ICMH
Collection de
microfiches.



Canadian Institute for Historical Microreproductions / Institut canadien de microreproductions historiques

© 1982

Technical and Bibliographic Notes/Notes techniques et bibliographiques

The Institute has attempted to obtain the best original copy available for filming. Features of this copy which may be bibliographically unique, which may alter any of the images in the reproduction, or which may significantly change the usual method of filming, are checked below.

- Coloured covers/
Couverture de couleur
- Covers damaged/
Couverture endommagée
- Covers restored and/or laminated/
Couverture restaurée et/ou pelliculée
- Cover title missing/
Le titre de couverture manque
- Coloured maps/
Cartes géographiques en couleur
- Coloured ink (i.e. other than blue or black)/
Encre de couleur (i.e. autre que bleue ou noire)
- Coloured plates and/or illustrations/
Planches et/ou illustrations en couleur
- Bound with other material/
Relié avec d'autres documents
- Tight binding may cause shadows or distortion
along interior margin/
La reliure serrée peut causer de l'ombre ou de la
distortion le long de la marge intérieure
- Blank leaves added during restoration may
appear within the text. Whenever possible, these
have been omitted from filming/
Il se peut que certaines pages blanches ajoutées
lors d'une restauration apparaissent dans le texte,
mais, lorsque cela était possible, ces pages n'ont
pas été filmées.
- Additional comments:/
Commentaires supplémentaires:

L'Institut a microfilmé le meilleur exemplaire qu'il lui a été possible de se procurer. Les détails de cet exemplaire qui sont peut-être uniques du point de vue bibliographique, qui peuvent modifier une image reproduite, ou qui peuvent exiger une modification dans la méthode normale de filmage sont indiqués ci-dessous.

- Coloured pages/
Pages de couleur
- Pages damaged/
Pages endommagées
- Pages restored and/or laminated/
Pages restaurées et/ou pelliculées
- Pages discoloured, stained or foxed/
Pages décolorées, tachetées ou piquées
- Pages detached/
Pages détachées
- Showthrough/
Transparence
- Quality of print varies/
Qualité inégale de l'impression
- Includes supplementary material/
Comprend du matériel supplémentaire
- Only edition available/
Seule édition disponible
- Pages wholly or partially obscured by errata
slips, tissues, etc., have been refilmed to
ensure the best possible image/
Les pages totalement ou partiellement
obscures par un feuillet d'errata, une pelure,
etc., ont été filmées à nouveau de façon à
obtenir la meilleure image possible.

This item is filmed at the reduction ratio checked below/
Ce document est filmé au taux de réduction indiqué ci-dessous.

10X	14X	18X	22X	26X	30X
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

12X 16X 20X 24X 28X 32X

The copy filmed here has been reproduced thanks to the generosity of:

The Nova Scotia Legislative Library

The images appearing here are the best quality possible considering the condition and legibility of the original copy and in keeping with the filming contract specifications.

Original copies in printed paper covers are filmed beginning with the front cover and ending on the last page with a printed or illustrated impression, or the back cover when appropriate. All other original copies are filmed beginning on the first page with a printed or illustrated impression, and ending on the last page with a printed or illustrated impression.

The last recorded frame on each microfiche shall contain the symbol → (meaning "CONTINUED"), or the symbol ▽ (meaning "END"), whichever applies.

Maps, plates, charts, etc., may be filmed at different reduction ratios. Those too large to be entirely included in one exposure are filmed beginning in the upper left hand corner, left to right and top to bottom, as many frames as required. The following diagrams illustrate the method:



L'exemplaire filmé fut reproduit grâce à la générosité de:

The Nova Scotia Legislative Library

Les images suivantes ont été reproduites avec le plus grand soin, compte tenu de la condition et de la netteté de l'exemplaire filmé, et en conformité avec les conditions du contrat de filmage.

Les exemplaires originaux dont la couverture en papier est imprimée sont filmés en commençant par le premier plat et en terminant soit par la dernière page qui comporte une empreinte d'impression ou d'illustration, soit par le second plat, selon le cas. Tous les autres exemplaires originaux sont filmés en commençant par la première page qui comporte une empreinte d'impression ou d'illustration et en terminant par la dernière page qui comporte une telle empreinte.

Un des symboles suivants apparaîtra sur la dernière image de chaque microfiche, selon le cas: le symbole → signifie "A SUIVRE", le symbole ▽ signifie "FIN".

Les cartes, planches, tableaux, etc., peuvent être filmés à des taux de réduction différents. Lorsque le document est trop grand pour être reproduit en un seul cliché, il est filmé à partir de l'angle supérieur gauche, de gauche à droite, et de haut en bas, en prenant le nombre d'images nécessaire. Les diagrammes suivants illustrent la méthode.

NOVA SCOTIA
LEGISLATIVE LIBRARY



PROVINCE HOUSE

359

345

LEGIS

HALIFAX,
THE
1871

CANADIAN NATURALIST

AND

Quarterly Journal of Science.

ON PORTIONS OF THE SKELETON OF A WHALE
FROM GRAVEL ON THE LINE OF THE CANADA
PACIFIC RAILWAY, NEAR SMITH'S FALLS, ON-
TARIO.

By J. W. DAWSON, LL.D., F.R.S.

Bones of large whales are of not infrequent occurrence on the less elevated terraces of the Pleistocene period on the Lower St. Lawrence. I have seen them at several places in the neighborhood of Metis, on the lowest sea terrace, now elevated only a few feet above the level of the sea, and they are reported to have been found on the second terrace at an elevation of 69 to 70 feet. Mr. Richardson, late of the Geological Survey, informs me that he has seen them in several other places on the lower terraces. It has also been reported that bones of a whale were found on Mt. Camille in rear of Metis at a considerable elevation; but Mr. Richardson, who visited the locality, failed to verify the statement. The bones found on the lower, and therefore modern terraces are usually in a good state of preservation and have a very recent appearance. The above statements relate to remains of the larger whalebone whales.

Remains of the *Beluga* or small white whale were found by the late Dr. Zadok Thompson, author of the "Natural History of Vermont," in the marine clay in the township of Charlotte, Vermont, at an elevation of 150 feet above the sea. They were associated with shells of *Saxicava* and *Leda*. The species was supposed to be distinct from the *B. Cetodon*, Gray, and was named by Thompson *B. Vermontana*. I have found detached bones of *Beluga* in the Post-pliocene clays of Rivière du Loup, and considerable portions of a skeleton were found in the

excavations for the Intercolonial Railway, on the south side of the Baie des Chaleurs, and were described by Gilpin in the Transactions of the Nova Scotia Institute of Natural Science.* Bones have also been found in the brick-clays near Montreal, and a specimen was discovered several years ago in sand holding *Saxicola*, near Cornwall, Ontario. The last-named specimen was studied by Mr. Billings, and its bones compared with those of the modern species in the McGill College Museum. On this evidence Mr. Billings concluded that it belonged to the modern species, and I believe extended this conclusion to Dr. Thompson's specimen, the distinctive characters of which, as stated by that naturalist, seem not to exceed the individual differences in modern specimens.

But though the *Beluga*, which now extends its excursions far up the St. Lawrence, and has even been captured in the vicinity of Montreal, occurs as far west as Cornwall, no remains of the larger whales have, so far as I am aware, been found so far inland until the discovery of the specimens referred to in the present note. These were found, as I am informed by Archer Baker, Esq., General Superintendent of the Canada Pacific Railway, "in a ballast pit, at Welshe's, on the line of the C. P. Railway, three miles north of Smith's Falls, and thirty-one miles north of the St. Lawrence River, in the Township of Montague, County of Lanark. They occurred in gravel at a depth of 30 feet from the surface, and about 50 feet back from the original face of the pit."

Mr. Peterson, C.E., has been kind enough to obtain for me the elevation of the place where the remains were found, as indicated by the railway levels. It is 420 feet above the level of the St. Lawrence at Hochelaga, or as nearly as possible 440 feet above sea level. It is interesting to observe that this corresponds exactly with the height of one of the sea terraces on the Montreal mountain, and is only 30 feet lower than the well-marked beach with sea shells above Côte des Neiges, on the west side of the Mountain. The highest level at which Post-pliocene marine shells are known to occur on Montreal Mountain, is near the park-keeper's house, at an elevation of about 520 feet. These marine deposits of Montreal are of the same geological period with the Cetacean remains in question, so that the animal to which these belonged may have sailed past the rocky islet which

then represented Montreal Mountain at an elevation of 400 feet above the lower levels of the city, and in a wide sea which then covered all the plain of the lower St. Lawrence.

The deposit in which the remains occurred is no doubt the equivalent of the Saxicava sand and gravel, and was probably a beach or bank near the base of the Laurentian hills, forming the west side of a bay which then occupied the Silurian country between the Laurentian hills north of the Ottawa, and those extending southward toward the Thousand Islands, and which opened into a wide extension of the Gulf of St. Lawrence, reaching to the hills of Eastern Canada and New England, and westward, perhaps, to the Niagara escarpment at the head of Lake Ontario. Such a sea might well be frequented by whales in the summer season, and individuals might occasionally be stranded on shallows or driven ashore by gales or by the pressure of floating ice.

The bones secured consist of two vertebrae and a fragment of another with a portion of a rib, and others are stated to have been found. They are in good preservation but have become white and brittle through the loss of their animal matter. On comparison with such remains of whales as exist in the Peter Redpath Museum, and with the figures and descriptions of other species, I have little doubt that they belong to the Humpback whale, *Megaptera longimana* of Gray, *Balaena hoopis* of Fabricius, a species still common in the Gulf of St. Lawrence, and which extends its range some distance up the River, and is more disposed than most others of the large whales to haunt inland waters, and to approach the shores. I have seen it as far up the river as the mouth of the Saguenay, and there is reason to believe that occasionally it runs up much further. It is a species well known to the Gaspé whalers and often captured by them. Of course with so little material it is not possible to be absolutely certain as to the species, but I think it may safely be referred to that above named. The larger of the two vertebrae, a lumbar one, has the centrum eleven inches in transverse diameter and is seven inches in length. The smaller, a dorsal, is ten inches in its greater diameter and four in length. Through the kindness of Mr. Baker, the specimens have been deposited in the Peter Redpath Museum of McGill University.

POLYZOA OF THE QUEEN CHARLOTTE ISLANDS. PRELIMINARY NOTICE OF NEW SPECIES.

By the Rev. THOMAS HINCKS, B.A., F.R.S.

In this paper I propose to give a diagnosis of a number of Polyzoa from the Queen Charlotte Islands, entrusted to me by Dr. G. M. Dawson, on behalf of the Geological Survey of Canada.

These forms will be more fully described and figured in a special report on the Polyzoa of these islands, which I hope to publish hereafter. As the preparation of the plates may occupy some time, it seems better to record the new species at once, and so avoid the risk of being anticipated after much labor has been expended on the work.

All critical notes on the species will be reserved for the Report.

Family *Membraniporidae*.

MEMBRANIPORA, De Blainville.

MEMBRANIPORA NIGRANS, n. sp.

Zoecia ovate (variable, sometimes arched above and narrowing downwards, sometimes broad-ovate, sometimes oval), irregularly disposed, margins much elevated, crenate, the whole front of the cell covered by a rather coarse stout membrane of a black color; oral valve large; on each side at the top a pointed *avicularium*, placed on the margin, depressed at the base, the beak sloping upwards, mandible directed obliquely downwards; very large *orientaria*, slightly raised in front, with a broad triangular mandible, which is bent abruptly in the middle, scattered amongst the zoecia. *Orium* very shallow, just covering the extremity of the cell, smooth, with a raised rib across it a little above the oral margin.

Zoarium of a deep black color, forming a large irregularly spreading crust.

Loc. Houston-Stewart Channel, Queen Charlotte Islands (Dr. G. M. Dawson).

MEMBRANIPORA EXILIS, n. sp.

Zoecia oblong, quincuncial, subtruncate above and below, set closely together, of considerable size and delicate material, margin thin, a good deal raised, the front wall wholly membranous; at the top of the cell 2 spines, and 3 or 4 on each side (or some-

times a smaller number), situated on the upper half of the cell, pointed, slender, suberect, jointed to a tubular base; a sessile *avicularium* on the margin at one side (often absent), just below the top, beak sloping upwards, scarcely bent at the extremity, mandible blunt, directed obliquely outwards. *Oocia* (?)

Loc. Houston-Stewart Channel, Queen Charlotte Islands, enveloping *Celoria horalis*, Busk, with a very thin crust (*Dr. G. M. Dawson*).

MEMBRANIPORA CONFERTA, N. SP.

Zoecia oval, quincuncial, set closely together, front wall wholly membranous, margin thin, smooth; on each side about 4 sharply pointed spines, and (often) a central one below, which bend rather abruptly over the area and meet in the middle; an *avicularium* at each side on the margin, just below the upper end, slightly raised, pointed, the mandible directed upwards, a small erect spine below the *avicularia*; at the bottom of the cell a single pointed *avicularium* with triangular mandible variously turned, *Ocium* rounded, smooth, with a variously shaped depressed area (or *tessa*) in front, composed of thinner material than the rest of the surface, and appearing dark-colored as compared with the surrounding dense white crust.

Loc. Houston-Stewart Channel, Queen Charlotte Islands (*Dr. G. M. Dawson*).

MEMBRANIPORA LEVATA, N. SP.

Zoecia small, oval, distinct, quincuncial, margin very slightly raised, thin, smooth, the whole front closed in by a smooth, light-colored, and rather glossy membrane, which lies very much on a level with the edge of the cell; above each zoecium, on a somewhat quadrate area, a small nodule, with a pointed *avicularium* on one side of it, the mandible directed transversely upwards. *Ocium* rounded, smooth, umboonate.

Loc. Houston-Stewart Channel, Queen Charlotte Islands (*Dr. G. M. Dawson*).

MEMBRANIPORA ECHINUS, N. SP.

Zoecia quincuncial, oval, distinct, separated by rather deep interspaces, 2 spines at the top and 7-8 slender, pointed, and rather tall spines down each side, which slant inwards but do not meet in the centre; on each side, springing from below the second spine from the top, a pedicellate *avicularium*, the upper part large and much swollen (closely resembling a "bird's

head,"), very slightly hooked at the extremity, apparently jointed to an extremely thin pedicle, mandible slender, pointed, *Orcium* (?).

Loc. Houston-Stewart Channel, Queen Charlotte Islands (*Dr. G. M. Dawson*).

Family *Cribribinidae*.

CRIBRILINA, Gray.

CRIBRILINA FURCATA, n. sp.

Zoecia ovate, quincuncial, very regularly disposed, moderately convex; surface smooth and lustrous, often of a reddish-brown color, on each side four to six shallow grooves, radiating to a median line, and a central one below, which are occupied by a row of roundish pores set very closely together, the ridges between them slightly raised, usually bearing several elliptical pores; orifice arched above, straight below, much broader than high, on each side a stout bifid spine (occasionally simple); peristome much thickened in front and rising into a central muero, *Avicularia* none. *Orcium* large (covering about half the cell above it), rounded, taller than broad, depressed in front, with a shallow oral arch; surface smooth, rather thickly punctured.

Loc. Off Cumshewa Harbor, Queen Charlotte Islands (*Dr. G. M. Dawson*).

CRIBRILINA HIPPOCREPIS, n. sp.

Zoecia ovate, quincuncial; surface lustrous, flattish (sutures very shallow), traversed by radiating ridges (three to five on each side), which pass from the sides to the centre (no median keel), the grooves between them occupied by a line of rather large oblong pores; at the margin of each ridge an elliptical foramen, covered in by a delicate membrane; orifice large, well arched above, constricted a little above the lower margin, which is straight; operculum of a rich reddish brown; peristome not elevated, lower margin much thickened, usually terminating on each side in a knob; large, elongate, depressed spatulate *avicularia* scattered amongst the cells. *Orcium* (?).

Surface of zoarium flat; color brown, with a tinge of red, in old states white and highly calcified.

Loc. Cumshewa Harbor and Houston-Stewart Channel, Queen Charlotte Islands, on shell (*Dr. G. M. Dawson*).

, apparently
lder, pointed.

Islands (Dr.

osed, mode-
f a reddish-
s, radiating
eepened by
ridges be-
ell elliptical
roader than
ple); peri-
tral muero,
half the cell
ont, with a
ctured.
ds (Dr. G.

h (sutures
ve on each
ian keel),
large ob-
foramen,
ell arched
which is
e not ele-
g on each
vicularia

of red, in
l, Queen

Family *Myriozoidae* (part.), Smitt.

SCHIZOPORELLA, Hincks.

SCHIZOPORELLA CRASSILABRIS, N. SP.

Zoaria large, elongate, ovate, quincuncial, very distinct, convex, sutures not very deep; surface dense, punctured (the punctures often obliterated by the calcification); orifice subacute suborbicular, with a broad rounded sinus occupying nearly the whole of the lower margin; peristome raised and thickened, forming a wall round the orifice, often massive in front, where it is carried out into a broad projection which is notched or sinuated in the centre. *Arcularia* none. *Ocicium* large, rounded, broader than high, with rather large punctures.

Loc. Houston-Stewart Channel, Queen Charlotte Islands 15-20 fms. (Dr. G. M. Dawson).

SCHIZOPORELLA LONGIROSTRATA, N. SP.

Zoaria ovate, disposed in lines, moderately convex, (sutures shallow); surface roughened or minutely granulated, covered with an epitheca; orifice arched above, lower margin extended into a wide rounded sinus; peristome thin, elevated at each side; on one side, generally a little below the orifice, an elongate, slender, pointed, dependent *arcularium*, the mandible (which is broad at the base and tapering above) directed obliquely downwards, usually turned slightly outwards. *Ocicium* rounded flattened in front, thickly punctured, with a shallow oral arch.

Loc. Off Cumshewa Harbor, on shell (Dr. G. M. Dawson).

SCHIZOPORELLA INSCULPTA, N. SP.

Zoarium foliaceous and bilaminar or inerusting. *Zoaria* large, ovate, quincuncial, depressed separated by raised lines, sutures shallow; surface vitreous, glossy, thickly covered over its whole extent with punctures; orifice arched above, the lower margin almost entirely occupied by a wide very shallow sinus; peristome thin, moderately raised, extended in front (beyond the sinus) so as to form a small chamber, in which is a rounded orifice (? *arcularium*). *Ocicium* profusely developed, very large (covering about two thirds of the cell above), elongate, rounded above, with a tall oral arch, thickly covered with slightly granulated ridges, which radiate from the opening to the base.

Loc. Queen Charlotte Islands under 30 fms., attached to a stem, and on shell (Dr. G. M. Dawson).

SCHIZOPORELLA MACULOSA, N. SP.

Zonaria quinquecival, rather small, moderately convex, sutures shallow; surface shining, covered with small punctures, which are closed in by a brownish membrane, and give a spotted appearance to the front wall; orifice arched above, with a shallow bluntly-pointed sinus below, not contracted at the opening; peristome slightly thickened, on one side just below the orifice (or occasionally on both sides) a small rounded *articularium* on a prominent boss. *Ocicium* (?).

Loc. Queen Charlotte Islands, on shell (*Dr. G. M. Dawson*).

SCHIZOPORELLA TEMULOSA, N. SP.

Zonaria quinquecival, very regularly arranged, very convex, ovate, much elevated centrally below the mouth, the wall sloping steeply down to the margin of the cell; surface dense, smooth, rather glossy, areolated round the edge, ridges radiating towards the centre; orifice orbicular, with a small central sinus, not contracted at the opening, peristome not elevated; immediately below the orifice, at one side of the sinus a rostrum, bearing on one side a pointed *articularium*, the beak very slightly bent at the extremity, mandible directed upwards, the rostrum rising into a short incurved point behind the *articularium*; very commonly on the front of the cell near the bottom a much raised *articularium* (mounted on a prominent elevation), with a pointed mandible directed straight outwards. *Ocicium* rounded, smooth, much broader than high, with a tall oral arch, filled in by a calcareous plate.

Loc. Off Cumshewa Harbor, Queen Charlotte Islands, in 20 fms., forming a brownish spreading crust (*Dr. G. M. Dawson*).

SCHIZOPORELLA DAWSONI, N. SP.

* *Zonaria* ovate, quinquecival, very moderately convex, separated by raised lines, lightly calcified, vitreous; surface reticulato-punctate (punctures appearing as deep shafts in the vitreous crust); orifice arched above, much broader than high (narrow between the upper and inferior margins); a shallow rounded sinus in the centre of the lower margin, not contracted at the opening; peristome perfectly simple, not raised. *Articularia* none. *Ocicium* rounded, closely united to the cell above, somewhat depressed in front, glossy, covered with rather large punctures; a prominent, thickened border round the opening.

* *Loc.* Virago Sound, Queen Charlotte Islands (*Dr. G. M. Dawson*).



