



**IMAGE EVALUATION  
TEST TARGET (MT-3)**



**Photographic  
Sciences  
Corporation**

23 WEST MAIN STREET  
WEBSTER, N.Y. 14580  
(716) 872-4503

18  
20  
22  
25  
28

**CIHM/ICMH  
Microfiche  
Series.**

**CIHM/ICMH  
Collection de  
microfiches.**



**Canadian Institute for Historical Microreproductions / Institut canadien de microreproductions historiques**

10  
11  
12

**© 1984**

Technical and Bibliographic Notes/Notes techniques et bibliographiques

The Institute has attempted to obtain the best original copy available for filming. Features of this copy which may be bibliographically unique, which may alter any of the images in the reproduction, or which may significantly change the usual method of filming, are checked below.

L'Institut a microfilmé le meilleur exemplaire qu'il lui a été possible de se procurer. Les détails de cet exemplaire qui sont peut-être uniques du point de vue bibliographique, qui peuvent modifier une image reproduite, ou qui peuvent exiger une modification dans la méthode normale de filmage sont indiqués ci-dessous.

- Coloured covers/  
Couverture de couleur
- Covers damaged/  
Couverture endommagée
- Covers restored and/or laminated/  
Couverture restaurée et/ou pelliculée
- Cover title missing/  
Le titre de couverture manque
- Coloured maps/  
Cartes géographiques en couleur
- Coloured ink (i.e. other than blue or black)/  
Encre de couleur (i.e. autre que bleue ou noire)
- Coloured plates and/or illustrations/  
Planches et/ou illustrations en couleur
- Bound with other material/  
Relié avec d'autres documents
- Tight binding may cause shadows or distortion  
along interior margin/  
La reliure serrée peut causer de l'ombre ou de la  
distortion le long de la marge intérieure
- Blank leaves added during restoration may  
appear within the text. Whenever possible, these  
have been omitted from filming/  
Il se peut que certaines pages blanches ajoutées  
lors d'une restauration apparaissent dans le texte,  
mais, lorsque cela était possible, ces pages n'ont  
pas été filmées.
- Additional comments:  
Commentaires supplémentaires:

- Coloured pages/  
Pages de couleur
- Pages damaged/  
Pages endommagées
- Pages restored and/or laminated/  
Pages restaurées et/ou pelliculées
- Pages discoloured, stained or foxed/  
Pages décolorées, tachetées ou piquées
- Pages detached/  
Pages détachées
- Showthrough/  
Transparence
- Quality of print varies/  
Qualité inégale de l'impression
- Includes supplementary material/  
Comprend du matériel supplémentaire
- Only edition available/  
Seule édition disponible
- Pages wholly or partially obscured by errata  
slips, tissues, etc., have been refilmed to  
ensure the best possible image/  
Les pages totalement ou partiellement  
obscurcies par un feuillet d'errata, une pelure,  
etc., ont été filmées à nouveau de façon à  
obtenir la meilleure image possible.

This item is filmed at the reduction ratio checked below/  
Ce document est filmé au taux de réduction indiqué ci-dessous.

10X	14X	18X	22X	26X	30X
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
12X	16X	20X	24X	28X	32X

The copy filmed here has been reproduced thanks to the generosity of:




Douglas Library  
Queen's University

The images appearing here are the best quality possible considering the condition and legibility of the original copy and in keeping with the filming contract specifications.

Original copies in printed paper covers are filmed beginning with the front cover and ending on the last page with a printed or illustrated impression, or the back cover when appropriate. All other original copies are filmed beginning on the first page with a printed or illustrated impression, and ending on the last page with a printed or illustrated impression.

The last recorded frame on each microfiche shall contain the symbol  $\rightarrow$  (meaning "CONTINUED"), or the symbol  $\nabla$  (meaning "END"), whichever applies.

Maps, plates, charts, etc., may be filmed at different reduction ratios. Those too large to be entirely included in one exposure are filmed beginning in the upper left hand corner, left to right and top to bottom, as many frames as required. The following diagrams illustrate the method:



L'exemplaire filmé fut reproduit grâce à la générosité de:

Douglas Library  
Queen's University

Les images suivantes ont été reproduites avec le plus grand soin, compte tenu de la condition et de la netteté de l'exemplaire filmé, et en conformité avec les conditions du contrat de filmage.

Les exemplaires originaux dont la couverture en papier est imprimée sont filmés en commençant par le premier plat et en terminant soit par la dernière page qui comporte une empreinte d'impression ou d'illustration, soit par le second plat, selon le cas. Tous les autres exemplaires originaux sont filmés en commençant par la première page qui comporte une empreinte d'impression ou d'illustration et en terminant par la dernière page qui comporte une telle empreinte.

Un des symboles suivants apparaîtra sur la dernière image de chaque microfiche, selon le cas: le symbole  $\rightarrow$  signifie "A SUIVRE", le symbole  $\nabla$  signifie "FIN".

Les cartes, planches, tableaux, etc., peuvent être filmés à des taux de réduction différents. Lorsque le document est trop grand pour être reproduit en un seul cliché, il est filmé à partir de l'angle supérieur gauche, de gauche à droite, et de haut en bas, en prenant le nombre d'images nécessaire. Les diagrammes suivants illustrent la méthode.

errata  
to

pelure,  
en à



32X

Queen's University  
Library

KINGSTON, ONTARIO

SO

HI

PRINCIPLES  
OF THE  
SOLUTION OF EQUATIONS  
OF THE  
HIGHER DEGREES, WITH APPLICATIONS.

BY GEORGE PAXTON YOUNG,  
TORONTO, CANADA.

*[Read before the Canadian Institute, March 3rd, 1883].*

---

TORONTO :  
PRINTED BY ROWSELL & HUTCHISON.

1884.

1828  
1834 Y7

## SOLUTION

1. Cor  
expression

2. The  
under wh  
rational i  
cognate fo  
cognate fo

3. Det  
primitive  
must have  
tive  $n$  re  
may leave  
subsist am

4. If  
irreducible  
rank ( $\S 3$ )  
the index  
simplified  
of the sur  
multiple o

5. Tw

6. The  
of the  $m^{\text{th}}$   
tions, is of

PRINCIPLES  
OF THE  
SOLUTION OF EQUATIONS OF THE HIGHER DEGREES,  
WITH APPLICATIONS.

---

BY GEORGE PAXTON YOUNG,  
*Toronto, Canada.*

---

CONTENTS.

1. Conception of a simple state to which every algebraical expression can be reduced. §6.

2. The unequal particular cognate forms of the generic expression under which a given simplified expression falls are the roots of a rational irreducible equation; and each of the unequal particular cognate forms occurs the same number of times in the series of the cognate forms. §9, 17.

3. Determination of the form which a rational function of the primitive  $n^{\text{th}}$  root of unity  $\omega_1$  and of other primitive roots of unity must have, in order that the substitution of any one of certain primitive  $n$  roots of unity,  $\omega_1, \omega_2, \omega_3$ , etc., for  $\omega_1$  in the given function may leave the value of the function unaltered. Relation that must subsist among the roots  $\omega_1, \omega_2$ , etc., that satisfy such a condition. §20.

4. If a simplified expression which is the root of a rational irreducible equation of the  $N^{\text{th}}$  degree involve a surd of the highest rank (§3) not a root of unity, whose index is  $\frac{1}{m}$ , the denominator of the index being a prime number,  $N$  is a multiple of  $m$ . But if the simplified root involve no surds that are not roots of unity, and if one of the surds involved in it be the primitive  $n^{\text{th}}$  root of unity,  $N$  is a multiple of a measure of  $n - 1$ . §28.

5. Two classes of solvable equations. §30.

6. The simplified root  $r_1$  of a rational irreducible equation  $F(x) = 0$  of the  $m^{\text{th}}$  degree,  $m$  prime, which can be solved in algebraical functions, is of the form

120169



$$r_1 = \frac{1}{m} \left( g + \Delta_1 \frac{1}{m} + a_1 \Delta_1 \frac{2}{m} + b_1 \Delta_1 \frac{3}{m} + \dots + e_1 \Delta_1 \frac{m-2}{m} + h_1 \Delta_1 \frac{m-1}{m} \right);$$

where  $g$  is rational, and  $a_1, b_1, \dots$ , involve only surds subordinate to  $\Delta_1 \frac{1}{m}$ . §38, 47.

7. The equation  $F(x) = 0$  has an auxiliary equation of the  $(m-1)$ <sup>th</sup> degree. §35, 52.

8. If the roots of the auxiliary be  $\Delta_1, \delta_2, \delta_3, \dots, \delta_{m-1}$ , the  $m-1$  expressions in each of the groups

$$\begin{array}{l} \Delta_1 \frac{1}{m} \frac{1}{\delta_{m-1}^m}, \quad \delta_2 \frac{1}{m} \frac{1}{\delta_{m-2}^m}, \dots, \quad \delta_{m-1} \frac{1}{m} \frac{1}{\Delta_1^m}, \\ \Delta_1 \frac{2}{m} \frac{1}{\delta_{m-2}^m}, \quad \delta_2 \frac{2}{m} \frac{1}{\delta_{m-4}^m}, \dots, \quad \delta_{m-1} \frac{2}{m} \frac{1}{\delta_2^m}, \\ \Delta_1 \frac{3}{m} \frac{1}{\delta_{m-3}^m}, \quad \delta_2 \frac{3}{m} \frac{1}{\delta_{m-6}^m}, \dots, \quad \delta_{m-1} \frac{3}{m} \frac{1}{\delta_3^m}, \end{array}$$

and so on, are the roots of a rational equation of the  $(m-1)$ <sup>th</sup> degree.

The  $\frac{m-1}{2}$  terms

$$\Delta_1 \frac{1}{m} \frac{1}{\delta_{m-1}^m}, \quad \delta_2 \frac{1}{m} \frac{1}{\delta_{m-2}^m}, \dots, \quad \delta_{\frac{m-1}{2}} \frac{1}{m} \frac{1}{\delta_{\frac{m+1}{2}}^m},$$

are the roots of a rational equation of the  $\left(\frac{m-1}{2}\right)$ <sup>th</sup> degree. §39, 44, 55.

9. Wider generalization. §45, 57.

10. When the equation  $F(x) = 0$  is of the first class, the auxiliary equation of the  $(m-1)$ <sup>th</sup> degree is irreducible. §35. Also the roots of the auxiliary are rational functions of the primitive  $m$ <sup>th</sup> root of unity. §36. And, in the particular case when the equation  $F(x) = 0$  is the reducing Gaussian equation of the  $m$ <sup>th</sup> degree to the equation  $x^n - 1 = 0$ , each of the  $\frac{m-1}{2}$  expressions,

$$\Delta_1 \frac{1}{m} \frac{1}{\delta_{m-1}^m}, \quad \delta_2 \frac{1}{m} \frac{1}{\delta_{m-2}^m}, \quad \&c.,$$

has the rational value  $n$ . §41. Numerical verification. §42.

11. Solv

12. Ana  
The auxilia  
sub-auxilia  
three case  
for the roo

§1. It w  
paper have

unless wh

regarded as  
understood

For instanc

When a sur  
it may be p  
present in t

§2. In su

nate to the  
cipal surd i

§3. A su

be of the fi  
( $a-1$ )<sup>th</sup> r  
 $a$ <sup>th</sup> rank.  
indices of  
numbers.

§4. An a

surd may b

$m$ <sup>th</sup>, thus,

$$\frac{1}{m} (g_1 + k$$

where  $g_1, k$

## 11. Solution of the Gaussian. §43.

## 12. Analysis of solvable irreducible equations of the fifth degree.

The auxiliary biquadratic either is irreducible, or has an irreducible sub-auxiliary of the second degree, or has all its roots rational. The three cases considered separately. Deduction of Abel's expression for the roots of a solvable quintic. §58-74.

## PRINCIPLES.

§1. It will be understood that the surds appearing in the present paper have *prime numbers* for the denominators of their indices, unless where the contrary is expressly stated. Thus,  $2^{\frac{1}{15}}$  may be regarded as  $h^{\frac{1}{5}}$ , a surd with the index  $\frac{1}{5}$ ,  $h$  being  $2^{\frac{1}{3}}$ . It will be understood also that no surd appears in the denominator of a fraction.

For instance, instead of  $\frac{2}{1 + \sqrt{-3}}$  we should write  $\frac{1 - \sqrt{-3}}{2}$ .

When a surd is spoken of as occurring in an algebraical expression, it may be present in more than one of its powers, and need not be present in the first.

§2. In such an expression as  $\sqrt{2} + (1 + \sqrt{2})^{\frac{1}{2}}$ ,  $\sqrt{2}$  is *subordinate* to the *principal* surd  $(1 + \sqrt{2})^{\frac{1}{2}}$ , the latter being the only principal surd in the expression.

§3. A surd that has no other surd subordinate to it may be said to be of the *first rank*; and the surd  $h^{\frac{1}{a}}$ , where  $h$  involves a surd of the  $(a - 1)^{\text{th}}$  rank, but none of a higher rank, may be said to be of the  $a^{\text{th}}$  rank. In estimating the rank of a surd, the denominators of the indices of the surds concerned are always supposed to be prime numbers. Thus,  $3^{\frac{1}{2}}$  is a surd of the second rank.

§4. An algebraical expression in which  $\Delta_1^{\frac{1}{m}}$  is a principal (see §2) surd may be arranged according to the powers of  $\Delta_1^{\frac{1}{m}}$  lower than the  $m^{\text{th}}$ , thus,

$$\frac{1}{m} \left( g_1 + k_1 \Delta_1^{\frac{1}{m}} + a_1 \Delta_1^{\frac{2}{m}} + b_1 \Delta_1^{\frac{3}{m}} + \dots + e_1 \Delta_1^{\frac{m-2}{m}} + h_1 \Delta_1^{\frac{m-1}{m}} \right) (1)$$

where  $g_1, k_1, a_1$ , etc., are clear of  $\Delta_1^{\frac{1}{m}}$ .

§5. If an algebraical expression  $r_1$ , arranged as in (1), be zero, while the coefficients  $g_1, k_1$ , etc., are not all zero, an equation

$$\omega \Delta_1^{\frac{1}{m}} = l_1 \tag{2}$$

must subsist; where  $\omega$  is an  $m^{\text{th}}$  root of unity; and  $l_1$  is an expression involving only such surds exclusive of  $\Delta_1^{\frac{1}{m}}$  as occur in  $r_1$ . For, let the first of the coefficients  $h_1, e_1$ , etc., proceeding in the order of the descending powers of  $\Delta_1^{\frac{1}{m}}$ , that is not zero, be  $n_1$ , the coefficient of  $\Delta_1^{\frac{s}{m}}$ . Then we may put

$$mr_1 = n_1 \left\{ f \left( \Delta_1^{\frac{1}{m}} \right) \right\} = n_1 \Delta_1^{\frac{s}{m}} + \text{etc.} = 0.$$

Because  $\Delta_1^{\frac{1}{m}}$  is a root of each of the equations  $f(x) = 0$  and  $x^m - \Delta_1 = 0$ ,  $f(x)$  and  $x^m - \Delta_1$  have a common measure. Let their H. C. M., involving only such surds as occur in  $f(x)$  and  $x^m - \Delta_1$ , be  $\varphi(x)$ . Then, because  $\varphi(x)$  is a measure of  $x^m - \Delta_1$ , the roots of the equation

$$\varphi(x) = x^c + p_1 x^{c-1} + p_2 x^{c-2} + \text{etc.} = 0$$

are  $\Delta_1^{\frac{1}{m}}, \omega_1 \Delta_1^{\frac{1}{m}}, \omega_2 \Delta_1^{\frac{1}{m}}, \dots, \omega_{c-1} \Delta_1^{\frac{1}{m}}$ ; where  $\omega_1, \omega_2$ , etc., are distinct primitive  $m^{\text{th}}$  roots of unity. Therefore,

$$\Delta_1^{\frac{c}{m}} (\omega_1 \omega_2 \dots) (-1)^c = p_c.$$

Now  $c$  is a whole number less than  $m$  but not zero; and, by §1,  $m$  is prime. Therefore there are whole numbers  $n$  and  $h$  such that

$$\Delta_1^{\frac{cn}{m}} (\omega_1 \omega_2 \dots)^n (-1)^{cn} = \Delta_1^{\frac{1}{m}} \Delta_1^{\frac{h}{m}} (\omega_1 \omega_2 \dots)^n (-1)^{cn} = p_c^n.$$

Therefore, if  $(\omega_1 \omega_2 \dots)^n = \omega$ , and  $l_1 \Delta_1^{\frac{h}{m}} (-1)^{cn} = p_c^n$ ,  $\omega \Delta_1^{\frac{1}{m}} = l_1$ .

§6. Let  $r_1$  be an algebraical expression in which no root of unity having a rational value occurs in the surd form  $\Delta_1^{\frac{1}{m}}$ . Also let there be in  $r_1$  no surd  $\Delta_1^{\frac{1}{m}}$  not a root of unity, such that

where  $e_1$   
 $\Delta_1^{\frac{1}{m}}$  excc

the same  
said to ha

§7. Som  
root 8<sup>1</sup> ca  
2 $\omega$ ,  $\omega$  bein  
the inadmi  
simplified  
 $\sqrt{5} = 2$   
more, a ro  
(2 +  $\sqrt{5}$ )  
(2 -  $\sqrt{3}$ )

§8. Let

where  $\Delta_1^{\frac{1}{m}}$   
 $p_2$ , etc., inv  
are roots  
from  $\Delta_1^{\frac{1}{m}}$

For, by §5,  
 $m^{\text{th}}$  root o  
 $\Delta_1^{\frac{1}{m}}$ ; an

§9. The  
generic sy  
 $r_1, r_2, r_3$ , e  
surd invo  
surd is uni  
account.  
may be ca  
 $r_1 = 1 \pm \sqrt{5}$

$$\Delta_1^{\frac{1}{m}} = e_1, \quad (3)$$

where  $e_1$  is an expression involving no surds of so high a rank as  $\Delta_1^{\frac{1}{m}}$  except such as either are roots of unity, or occur in  $r_1$  being at the same time distinct from  $\Delta_1^{\frac{1}{m}}$ . The expression  $r_1$  may then be said to have been *simplified* or to be *in a simple state*.

§7. Some illustrations of the definition in §6 may be given. The root  $8^{\frac{1}{3}}$  cannot occur in a simplified expression  $r_1$ ; for its value is  $2\omega$ ,  $\omega$  being a third root of unity; but the equation  $8^{\frac{1}{3}} = 2\omega$  is of the inadmissible type (3). Again, the root  $\sqrt[5]{5}$  cannot occur in a simplified expression; for,  $\omega_1$  being a primitive fifth root of unity,  $\sqrt[5]{5} = 2(\omega_1 + \omega_1^4) + 1$ ; an equation of the type (3). Once more, a root of the cubic equation  $x^3 - 3x - 4 = 0$ , in the form  $(2 + \sqrt{3})^{\frac{1}{3}} + (2 - \sqrt{3})^{\frac{1}{3}}$ , is not in a simple state, because  $(2 - \sqrt{3})^{\frac{1}{3}} = (2 - \sqrt{3})(2 + \sqrt{3})^{\frac{1}{3}}$ .

$$\text{§8. Let } p_1 \Delta_1^{\frac{m-1}{m}} + p_2 \Delta_1^{\frac{m-1}{m}} + \dots + p_m = 0; \quad (4)$$

where  $\Delta_1^{\frac{1}{m}}$  is a surd occurring in a simplified expression  $r_1$ ; and  $p_1, p_2, \dots$ , involve no surds of so high a rank as  $\Delta_1^{\frac{1}{m}}$ , except such as either are roots of unity, or occur in  $r_1$  being at the same time distinct from  $\Delta_1^{\frac{1}{m}}$ . The coefficients  $p_1, p_2, \dots$ , must be zero separately.

For, by §5, if they were not, we should have  $\omega \Delta_1^{\frac{1}{m}} = l_1$ ,  $\omega$  being an  $m^{\text{th}}$  root of unity, and  $l_1$  involving only surds in (4) distinct from  $\Delta_1^{\frac{1}{m}}$ ; an equation of the inadmissible type (3).

§9. The expression  $r_1$  being in a simple state, we may use  $R$  as a generic symbol to include the various particular expressions, say  $r_1, r_2, r_3, \dots$ , obtained by assigning all their possible values to the surds involved in  $r_1$ , with the restriction that, where the base of a surd is unity, the rational value of the surd is not to be taken into account. These particular expressions, not necessarily all unequal, may be called the *particular cognate forms* of  $R$ . For instance, if  $r_1 = 1^{\frac{1}{m}}$ ,  $R$  has two particular cognate forms, the rational value of the

third root of unity not being counted. If  $r_1 = (1 + \sqrt{2})^{\frac{1}{2}}$ ,  $R$  has six particular cognate forms all unequal. Should  $r_1 = (2 + \sqrt{3})^{\frac{1}{2}} + (2 - \sqrt{3})^{\frac{1}{2}}$ ,  $R$  has six particular cognate forms, but only three unequal, each of the unequal forms occurring twice.

§10. PROPOSITION I. An algebraical expression  $r_1$  can always be brought to a simple state.

For  $r_1$  may be cleared of all surds such as  $1^{\frac{1}{m}}$  having a rational value. Suppose that  $r_1$  then involves a surd  $\Delta_1^{\frac{1}{m}}$ , not a root of unity, by means of which an equation such as (3) can be formed. Substitute for  $\Delta_1^{\frac{1}{m}}$  in  $r_1$  its value  $e_1$  as thus given. The result will be to eliminate  $\Delta_1^{\frac{1}{m}}$  from  $r_1$  without introducing into the expression any new surd as high in rank as  $\Delta_1^{\frac{1}{m}}$ , and at the same time not a root of unity. By continuing to make all the eliminations of this kind that are possible, we at last reach a point where no equation of the type (3) can any longer be formed. Then because, by the course that has been pursued, no roots of the form  $1^{\frac{1}{m}}$  having a rational value have been left in  $r_1$ ,  $r_1$  is in a simple state.

§11. It is known that, if  $N$  be any whole number, the equation whose roots are the primitive  $N^{\text{th}}$  roots of unity is rational and irreducible.

§12. Let  $N$  be the continued product of the distinct prime numbers  $n, a, b$ , etc. Let  $\omega_1$  be a primitive  $n^{\text{th}}$  root of unity,  $\theta_1$  a primitive  $a^{\text{th}}$  root of unity, and so on. Let  $\omega$  represent any one indifferently of the primitive  $n^{\text{th}}$  roots of unity,  $\theta$  any one indifferently of the primitive  $a^{\text{th}}$  roots of unity, and so on. Let  $f(\omega_1, \theta_1, \text{etc.})$  be a rational function of  $\omega_1, \theta_1, \text{etc.}$  Then a corollary from §11 is, that if  $f(\omega_1, \theta_1, \text{etc.}) = 0, f(\omega, \theta, \text{etc.}) = 0$ . For  $t_1$  being a primitive  $N^{\text{th}}$  root of unity, and  $t$  representing any one indifferently of the primitive  $N^{\text{th}}$  roots of unity, we may put

$$f(\omega_1, \theta_1, \text{etc.}) = a_1 t_1^{N-1} + a_2 t_1^{N-2} + \text{etc.} = 0,$$

$$\text{and } f(\omega, \theta, \text{etc.}) = a_1 t^{N-1} + a_2 t^{N-2} + \text{etc.};$$

where the coefficients  $a_1, a_2, \text{etc.}$ , are rational. Should these coefficients be all zero,  $f(\omega, \theta, \text{etc.}) = 0$ . Should they not be all zero, let  $a_r$  be the first that is not zero. Then we may put

$$f(\omega_1, \theta_1, \text{etc.}) = a_r \{ \varphi(t_1) \} = a_r t_1^{N-r} + \text{etc.} = 0.$$

Therefore  
the same  
whose ro  
 $\varphi(x)$  ha  
Therefore  
 $\psi(x) =$

§13. A

$f(x)$   
where  $h_1$   
equal to  
 $f(\omega, \theta_1,$   
 $\omega \{ f(\omega,$   
Then, in  
 $nh_1 = h,$   
In like m

§14. P  
particular  
 $F(x) =$   
equation.

For, le  
be establ  
of unity.

of unity  
number.  
that do  
of unity  
 $n$  distinc

still to l

not a r  
Then  $F'$   
the exp  
surds in

$r_1$  to  $r_2$

Therefore,  $t_1$  is a root of the rational equation  $\varphi(x) = 0$ , being at the same time a root of the rational equation  $\psi(x) = 0$ , whose roots are the primitive  $N^{\text{th}}$  roots of unity. Hence  $\psi(x)$  and  $\varphi(x)$  have a common measure. But by §11,  $\psi(x)$  is irreducible. Therefore it is a measure of  $\varphi(x)$ ; and the roots of the equation  $\psi(x) = 0$  are roots of the equation  $\varphi(x) = 0$ . Therefore,

$$f(\omega, \theta, \text{etc.}) = a_r \{ \varphi(t) \} = 0.$$

§13. Another corollary is, that if

$$f(\omega_1, \theta_1, \text{etc.}) = h_1 \omega_1^{n-1} + h_2 \omega_1^{n-2} + \dots + h_n = 0,$$

where  $h_1, h_2, \text{etc.}$ , are clear of  $\omega_1$ , the coefficients  $h_1, h_2, \text{etc.}$ , are all equal to one another. For, by §12, because  $f(\omega_1, \theta_1, \text{etc.}) = 0$ ,  $f(\omega, \theta_1, \text{etc.}) = 0$ . Therefore  $\omega \{ f(\omega, \theta_1, \text{etc.}) \} = 0$ . In  $\omega \{ f(\omega, \theta_1, \text{etc.}) \}$  give  $\omega$  successively its  $n - 1$  different values.

Then, in addition,

$$nh_1 = h_1 + h_2 + \dots + h_n. \text{ Similarly, } nh_2 = h_1 + h_2 + \dots + h_n. \therefore h_1 = h_2.$$

In like manner all the terms  $h_1, h_2, \text{etc.}$ , are equal to one another.

§14. PROPOSITION II. If the simplified expression  $r_1$ , one of the particular cognate forms of  $R$ , be a root of the rational equation  $F(x) = 0$ , all the particular cognate forms of  $R$  are roots of that equation.

For, let  $r_2$  be a particular cognate form of  $R$ . By §12, the law to be established holds when there are no surds in  $r_1$  that are not roots of unity. It will be kept in view that, according to §1, when roots

of unity are spoken of, such roots are meant as  $1^{\frac{1}{m}}$ ,  $m$  being a prime number. Assume the law to have been found good for all expressions that do not involve more than  $n - 1$  distinct surds that are not roots of unity; then, making the hypothesis that  $r_1$  involves not more than  $n$  distinct surds that are not roots of unity, the law can be shown

still to hold; in which case it must hold universally. For, let  $\Delta_1^{\frac{1}{m}}$ ,

not a root of unity, be a surd of the highest rank (see §3) in  $r_1$ . Then  $F(r_1)$  may be taken to be the expression (1), and  $F(r_2)$  to be the expression formed from (1) by selecting particular values of the surds involved under the restriction specified in §9. In passing from

$r_1$  to  $r_2$ , let  $\Delta_1^{\frac{1}{m}}$ ,  $a_1, \text{etc.}$ , become respectively  $\Delta_2^{\frac{1}{m}}$ ,  $a_2, \text{etc.}$  Then

$$m \{ F(r_1) \} = h_1 \Delta_1^{\frac{m-1}{m}} + e_1 \Delta_1^{\frac{m-2}{m}} + \text{etc.} = 0,$$

$$\text{and } m \{ F(r_2) \} = h_2 \Delta_2^{\frac{m-1}{m}} + e_2 \Delta_2^{\frac{m-2}{m}} + \text{etc.}$$

By §8, because  $r_1$  is in a simple state, and  $F(r_1) = 0$ , the coefficients  $h_1, e_1$ , etc., are zero separately. But  $h_1$  is clear of the surd  $A_1^{\frac{1}{m}}$ . It therefore does not involve more than  $n - 1$  distinct surds that are not roots of unity. Therefore, on the assumption on which we are proceeding, because  $h_1 = 0, h_2 = 0$ . In like manner,  $e_2 = 0$ , and so on. Therefore  $F(r_2) = 0$ .

§15. *Cor.* Let the simplified expression  $r_1$  be the root of an equation  $F(x) = 0$  whose coefficients involve certain surds

$z_1^{\frac{1}{n}}, u_1^{\frac{1}{s}}$ , etc., that have the same determinate values in  $r_1$  as in  $F(x)$ . Then, if  $r_2$  be a particular cognate form of  $R$  in which the

surds  $z_1^{\frac{1}{n}}, u_1^{\frac{1}{s}}$ , etc., retain the determinate values belonging to them in  $r_1, r_2$  is a root of the equation  $F(x) = 0$ . For,  $F(r_1) = 0$ . Therefore, by the Proposition,  $F(R) = 0$ . Let  $R$ , restricted by the

condition that the surds  $z_1^{\frac{1}{n}}, u_1^{\frac{1}{s}}$ , etc., retain the determinate values belonging to them in  $r_1$ , be  $R'$ . Then  $F(R') = 0$ . A particular case of this is  $F(r_2) = 0$ . The corollary established simply means that

the surds  $z_1^{\frac{1}{n}}, u_1^{\frac{1}{s}}$ , etc., may be taken to be rational for the purpose in hand.

§16. The simplified expression  $r_1$  being one of the particular cognate forms of  $R$ , let  $r_1, r_2$ , etc. (5)

be the entire series of the particular cognate forms of  $R$ , not necessarily unequal to one another. Then, if the equation whose roots are the terms in (5) be  $X = 0, X$  is rational. In like manner, if those particular cognate forms of  $R$ , not necessarily unequal, that

are obtained when certain surds  $z_1^{\frac{1}{n}}, u_1^{\frac{1}{s}}$ , etc., retain the determinate values belonging to them in  $r_1$ , be

$$r_1, r_2, \text{ etc.} \tag{6}$$

and if the equation whose roots are the terms in (6) be  $X' = 0, X'$

involves only surds found in the series  $z_1^{\frac{1}{n}}, u_1^{\frac{1}{s}}$ , etc. This is substantially proved by Legendre in his *Théorie des Nombres*, §487, third edition.

§17. For the general case, the roots are the roots of an equation whose roots are unequal to one another. The series

As in §15, let the terms of the series be the terms of the series, all the terms of the series being  $X = 0$ . It has a rational root, all the roots of the equation being irrational, while at the same time the roots of the equation are unequal to one another.

Because  $X$  is a rational number,  $F(x)$  is a rational number, its roots are the roots of the equation  $F(x) = 0$ . Since  $F(x)$  is a rational number,  $F(x)$  is a rational number, and  $X$  is a rational number. The forms of  $R$  are cognate forms of  $R$ .

§18. *Co*

forms of  $R$  are the terms of the series, not necessarily unequal to one another. (6) occurs in (6) are

involve only surds found in the series  $z_1^{\frac{1}{n}}, u_1^{\frac{1}{s}}$ , etc. This is substantially proved by Legendre in his *Théorie des Nombres*, §487, third edition.

§17. PROPOSITION III. The unequal particular cognate forms of  $R$ , the generic expression under which the simplified expression  $r_1$  falls, are the roots of a rational irreducible equation; and each of the unequal particular cognate forms occurs the same number of times in the series of the cognate forms.

As in §16, let the entire series of the particular cognate forms of  $R$  be the terms in (5), the equation that has these terms for its roots being  $X = 0$ . By §16,  $X$  is rational. Should  $X$  not be irreducible, it has a rational irreducible factor, say  $F(x)$ , such that  $r_1$  is a root of the equation  $F(x) = 0$ . By Prop. II., because  $r_1$  is in a simple state, all the terms in (5) are roots of the equation  $F(x) = 0$ , while at the same time, because  $F(x)$  is a factor of  $X$ , all the roots of the equation are terms in (5). And the equation  $F(x) = 0$ , being irreducible, has no equal roots. Therefore its roots are the unequal terms in (5). Should  $F(x)$  not be identical with  $X$ , put

$$X = \{F(x)\} \{\varphi(x)\}.$$

Because  $X$  and  $F(x)$  are rational,  $\varphi(x)$  is rational. Then, since  $\varphi(x)$  is a measure of  $X$ , and the equation  $F(x) = 0$  has for its roots the unequal roots of the equation  $X = 0$ , the equations  $F(x) = 0$  and  $\varphi(x) = 0$  have a root in common. Consequently, since  $F(x)$  is irreducible, it is a measure of  $\varphi(x)$ . Therefore  $\{F(x)\}^2$  is a measure of  $X$ . Going on in this way we ultimately get  $X = \{F(x)\}^N$ ; which means that each of the particular cognate forms of  $R$  has its value repeated  $N$  times in the series of the particular cognate forms.

§18. Cor. 1. The series (6) consisting of those particular cognate forms of  $R$  in which certain surds  $z_1^{\frac{1}{n}}, u_1^{\frac{1}{s}}$ , etc., retain the determinate values belonging to them in  $r_1$ , each of the unequal terms in (6) occurs the same number of times in (6); and the unequal terms in (6) are the roots of an irreducible equation whose coefficients

involve only surds found in the series  $z_1^{\frac{1}{n}}, u_1^{\frac{1}{s}}$ , etc. Should  $X'$  not be irreducible, by which in such a case is meant incapable of being broken into lower factors involving only surds occurring in  $X'$ , let it have the irreducible factor  $X''$ . That is to say,  $X''$  involves only surds occurring in  $X'$ , and has itself no lower factor involving only surds that occur in  $X'$ . We may take  $r_1$  to be a root of the equation  $X'' = 0$ . Then, by Cor. Prop. II., all the terms in (6) are roots of that equation, all the roots of the equation being at the same time terms in (6). And the equation  $X'' = 0$  being irreducible, has no equal roots. Therefore its roots are the unequal terms in (6). Put



$X' = (X'')(X''')$ . Then, by the line of reasoning followed in the Proposition,  $X''$  has a measure identical with  $X''$ . And so on. Ultimately  $X' = (X'')^N$ .

§19. Cor. 2. If  $r_2$ , one of the particular cognate forms of  $R$ , be zero, all the particular cognate forms of  $R$  are zero. For, by the proposition, the particular cognate forms of  $R$  are the roots of a rational irreducible equation  $F(x) = 0$ . And  $r_2$ , one of the roots of that equation, is zero, but the only rational irreducible equation that has zero for a root is  $x = 0$ . Therefore  $F(x) = x = 0$ . In fact, in the case supposed, the simplified expression  $r_1$  is zero, and  $R$  has no particular cognate forms distinct from  $r_1$ .

§20. PROPOSITION IV. Let  $N$  be the continued product of the distinct prime numbers  $n, a$ , etc. Let  $\omega_1$  be a primitive  $n^{\text{th}}$  root of unity,  $\theta_1$  a primitive  $a^{\text{th}}$  root of unity, and so on. Then if the equation

$$F(x) = x^d + b_1x^{d-1} + b_2x^{d-2} + \text{etc.} = 0$$

be one in which the coefficients  $b_1, b_2$ , etc., are rational functions of  $\omega_1, \theta_1$ , etc., and if all the primitive  $n^{\text{th}}$  roots of unity, which, when substituted for  $\omega_1$  in  $F(x)$ , leave  $F(x)$  unaltered, be

$$\omega_1, \omega_2, \dots, \omega_s, \tag{7}$$

the series (7) either consists of a single term or it is made up of a cycle of primitive  $n^{\text{th}}$  roots of unity,

$$\omega_1, \omega_1^\lambda, \omega_1^{\lambda^2}, \dots, \omega_1^{\lambda^{s-1}}; \tag{18}$$

that is to say, no term in (8) after the first is equal to the first, but  $\omega_1^{\lambda^s} = \omega_1$ . Also, if (let it be kept in view that  $n$  is prime) the cycle that contains all the primitive  $n^{\text{th}}$  roots of unity be

$$\omega_1, \omega_1^\beta, \omega_1^{\beta^2}, \dots, \omega_1^{\beta^{n-2}}, \tag{9}$$

and if  $C_1$  be the sum of the terms in the cycle (8), the form of  $F(x)$  is

$$F(x) = x^d - (p_1C_1 + p_2C_2 + \dots + p_mC_m) x^{d-1} + (q_1C_1 + q_2C_2 + \text{etc.}) x^{d-2} + \text{etc.} \tag{10}$$

where each of the expressions in the series  $C_1, C_2, C_3$ , etc., is what the immediately preceding term becomes by changing  $\omega_1$  into  $\omega_1^\beta, C_m$  through this change becoming  $C_1$ ; and  $p_1, p_2, q_1$ , etc., are clear of  $\omega_1$ .

For, assuming that there is a term  $\omega_2$  in (7) additional to  $\omega_1$ , we may take  $\omega_2$  to be the first term in (9) after  $\omega_1$  that occurs in (7).

and it may be considered to be  $\omega_1^{\beta^m}$ , which may be otherwise written  $\omega_1^\lambda$ . Then, if  $F(x)$  be written  $\varphi(\omega_1)$ , we have by hypothesis

$\varphi(\omega_1) =$   
 $\varphi(\omega_1^\lambda)$   
 $\varphi(\omega_1^{\lambda^2})$ ,  
 negative.  
 these term  
 in (7), say  
 $\varphi(\omega_1) =$   
 because s  
 Because a  
 it is not l  
 which, w  
 Therefore  
 less than  
 $z = - (h$   
 which, be  
 in (9) aft  
 unaltere  
 (8), whil  
 in (7).  
 the cycle  
 expressio  
 (9). Th  
 Because  
 $F(x) =$   
 where p  
 $\omega_1$  is ch  
 Theref  
 Here,  
 the sa

$\varphi(\omega_1) = \varphi(\omega_1^\lambda)$ . Therefore, by §12, changing  $\omega_1$  into  $\omega_1^\lambda$ ,  $\varphi(\omega_1) = \varphi(\omega_1^{\lambda^2})$ . Therefore  $\varphi(\omega_1) = \varphi(\omega_1^{\lambda^3})$ . And thus ultimately  $\varphi(\omega_1) = \varphi(\omega_1^{\lambda^z})$ , or  $\varphi(\omega_1) = \varphi(\omega_1^{\beta^{mz}})$ ,  $z$  being any whole number positive or

negative. But  $\omega_1^{\lambda^z}$  includes all the terms in (8). Therefore each of these terms is a term in (7). Suppose if possible that there is a term

in (7), say  $\omega_1^{\beta^h}$ , which does not occur in (8). Then, just as we deduced  $\varphi(\omega_1) = \varphi(\omega_1^{\beta^{mz}})$  from the equation  $\varphi(\omega_1) = \varphi(\omega_1^{\beta^m})$ , we can, because still farther  $\varphi(\omega_1) = \varphi(\omega_1^{\beta^h})$ , deduce  $\varphi(\omega_1) = \varphi(\omega_1^{\beta^{mz+hu}})$ .

Because  $\omega_1^{\beta^h}$  lies outside the cycle (8),  $h$  is not a multiple of  $m$ . And it is not less than  $m$ , because  $\omega_1^{\beta^m}$  is the first term in (9) after  $\omega_1$ , which, when substituted for  $\omega_1$  in  $\varphi(\omega_1)$ , leaves  $\varphi(\omega_1)$  unaltered. Therefore  $h = qm + v$ , where  $q$  and  $v$  are whole numbers, and  $v$  is less than  $m$  but not zero. Put

$$(7) \quad z = -(h+q), \text{ and } u = m+1 \therefore mz+hu = v \therefore \varphi(\omega_1) = \varphi(\omega_1^{\beta^v});$$

which, because  $v$  is less than  $m$  but not zero, and  $\omega_1^{\beta^m}$  is the first term in (9) after  $\omega_1$  which, when substituted for  $\omega_1$  in  $\varphi(\omega_1)$ , leaves  $\varphi(\omega_1)$  unaltered, is impossible. Hence, no term in (7) lies outside the cycle (8), while it has also been shown that all the terms in (8) are terms in (7). Therefore the terms in (7) are identical with those constituting the cycle (3). We have now to determine the form of  $F(x)$ . The expressions,  $C_1, C_2$ , etc., taken together, are the sum of the terms in (9). Therefore  $C_1 + C_2 + \dots + C_m = -1$ . (11)

Because (9) contains all the primitive  $n^{\text{th}}$  roots of unity, we may put

$$F(x) = x^d - \{p + (p+p_1)\omega_1 + (p+p_2)\omega_1^\beta + \text{etc.}\} x^{d-1} + \text{etc.}; (12)$$

where  $p, p_1$ , etc., are clear of  $\omega_1$ . But  $F(x)$  remains unaltered when  $\omega_1$  is changed into  $\omega_1^{\beta^m}$ . Therefore

$$F(x) = x^d - \{p + (p+p_1)\omega_1^{\beta^m} + \text{etc.}\} x^{d-1} + \text{etc.} (13)$$

Therefore, equating the coefficients of  $x^{d-1}$  in (12) and (13),

$$(p-p_1) + \dots + (p_{m+1}-p_1)\omega_1^{\beta^m} + \text{etc.} = 0.$$

Here, by §13, the coefficients of the different powers of  $\omega_1$  have all the same value. And one of them,  $p-p_1$ , is zero. Therefore

$p_{m+1} = p_1$ . That is to say, the coefficient of  $\omega_1^{\beta^m}$  or  $\omega_1^\lambda$  is the same as that of  $\omega_1$ . In like manner the coefficients of all the terms in (8) are the same. Therefore one group of the terms that together make up the coefficient of  $x^d - 1$  in (12) is properly represented by  $-(p + p_1)C_1$ . In the same way another group is properly represented by  $-(p + p_2)C_2$ , and so on. Hence

$$F(x) = x^d - \{p + (p + p_1)C_1 + (p + p_2)C_2 + \text{etc.}\}x^{d-1} + \text{etc.}$$

And by (11) this is equivalent to (10). The form of  $F(x)$  has been deduced on the assumption that the series (7) contains more than one term; but, should the series (7) consist of a single term, the result obtained would still hold good, only in that case each of the expressions  $C_1, C_2$ , etc., would be a primitive  $n^{\text{th}}$  root of unity.

§21. A simplified expression will not cease to be in a simple state, if we suppose that any surd that can be eliminated from it, without the introduction of any new surd, has been eliminated.

§22. PROPOSITION V. In the simplified expression  $r_1$ , one of the particular cognate forms of  $R$ , modified according to §21, let the surd  $\Delta_1^{\frac{1}{m}}$  of the highest rank be not a root (see §1) of unity. Then,

if the particular cognate forms of  $R$  obtained by changing  $\Delta_1^{\frac{1}{m}}$  in  $r_1$  successively into the different  $m^{\text{th}}$  roots of the determinate base  $\Delta_1$ , be

$$r_1, r_2, \dots, r_m, \quad (14)$$

these terms are all unequal.

For the terms in (14) are all the particular cognate forms of  $R$  obtained when we allow all the surds in  $r_1$  except  $\Delta_1^{\frac{1}{m}}$  to retain the determinate values belonging to them in  $r_1$ . Therefore, by Cor. 1, Prop. III., each of the unequal terms in (14) has its value repeated the same number of times in that series. Let  $u$  be the number of the unequal terms in (14), and let each occur  $c$  times. Then  $uc = m$ . Suppose if possible that  $u = 1$ . This means that all the terms in (14) are equal. Therefore,  $r_1$  being the expression (1),

$$mr_1 = r_1 + r_2 + \dots + \text{etc.} = gr_1.$$

Therefore the surd  $\Delta_1^{\frac{1}{m}}$  can be eliminated from  $r_1$  without the introduction of any new surd; which, by §21, is impossible. Therefore  $u$  is not unity. But, by §1,  $m$  is a prime number. And  $m = uc$ . Therefore  $c = 1$  and  $u = m$ . This means that all the terms in (14) are unequal.

§23. Co

$R$ ; and become in

forms of  $R$  different  $m$

These term

$r_1$ , and  $r_2$

value is  $m$

be taken

except the

no surds d

in the sim

Let  $r_a + 2$

Then  $r_a +$

expression

Therefore

would, by

impossible

manner al

§24. Co

terms in (

clear of t

roots of  $u$

unity in  $u$

$c^{\text{th}}$  roots  $c$

be respect

selected,

§23. *Cor. 1.* Let  $r_{a+1}$  be any one of the particular cognate forms of  $R$ ; and let  $J_{a+1}^{\frac{1}{m}}, h_{a+1}$ , etc., be respectively what  $J_1^{\frac{1}{m}}, h_1$ , etc., become in passing from  $r_1$  to  $r_{a+1}$ . Also let the  $m$  particular cognate forms of  $R$ , obtained by changing  $J_{a+1}^{\frac{1}{m}}$  in  $r_{a+1}$  successively into the different  $m^{\text{th}}$  roots of  $J_{a+1}$ , be

$$r_{a+1}, r_{a+2}, \dots, r_{a+m}. \quad (15)$$

These terms are all unequal. For, because  $J_1^{\frac{1}{m}}$  is a principal surd in  $r_1$ , and  $r_2$  is what  $r_1$  becomes when  $J_1^{\frac{1}{m}}$  is changed into a surd whose

value is  $\omega_1 J_1^{\frac{1}{m}}$ ,  $\omega_1$  being a primitive  $m^{\text{th}}$  root of unity, the view may be taken that  $r_2$  involves no surds additional to those found in  $r_1$ , except the primitive  $m^{\text{th}}$  root of unity  $\omega_1$ . Therefore  $r_1 - r_2$  involves no surds distinct from primitive  $m^{\text{th}}$  roots of unity that are not found in the simplified expression  $r_1$ . Therefore  $r_1 - r_2$  is in a simple state.

Let  $r_{a+2}$  be what  $r_{a+1}$  becomes by changing  $J_{a+1}^{\frac{1}{m}}$  into  $\omega_1 J_{a+1}^{\frac{1}{m}}$ . Then  $r_{a+1} - r_{a+2}$  is a particular cognate form of the generic expression under which the simplified expression  $r_1 - r_2$  falls. Therefore  $r_{a+1} - r_{a+2}$  cannot be zero; for, if it were,  $r_1 - r_2$  would, by *Cor. 2, Prop. III.*, be zero; which, by the proposition, is impossible. Hence, the first two terms in (15) are unequal. In like manner all the terms in (15) are unequal.

§24. *Cor. 2.* Let  $X_1 = 0$  be the equation whose roots are the terms in (14). When  $X_1$  is modified according to §21, it is, by §16, clear of the surd  $J_1^{\frac{1}{m}}$ . Should it involve any surds that are not roots of unity, take  $z_1^{\frac{1}{c}}$  a surd of the highest rank not a root of unity in  $X_1$ ; and, when  $z_1^{\frac{1}{c}}$  is changed successively into the different  $c^{\text{th}}$  roots of the determinate base  $z_1$ , let

$$X_1, X_1', X_1'', \dots, X_1^{(c-1)}, \quad (16)$$

be respectively what  $X_1$  becomes. Any term in (16), as  $X_1$ , being selected, the  $m$  roots of the equation  $X_1 = 0$  are unequal particular

cognate forms of  $R$ . For,  $z_2^{\frac{1}{c}}$  being a  $c^{\text{th}}$  root of  $z_1$  distinct from  $\frac{1}{z_1^{\frac{1}{c}}}$ , let  $r_{a+1}$  be what  $r_1$  becomes when  $\frac{1}{z_1^{\frac{1}{c}}}$  becomes  $z_2^{\frac{1}{c}}$ ; the expressions  $\Delta_1^{\frac{1}{m}}$ ,  $h_1$ , etc., at the same time becoming  $\Delta_{a+1}^{\frac{1}{m}}$ ,  $h_{a+1}$ , etc. Then we may put

$$X_1 = x^m + (bz_1^{\frac{c-1}{c}} + dz_1^{\frac{c-2}{c}} + \text{etc.}) x^{m-1} + \text{etc.}; \quad (17)$$

where  $b, d$ , etc., are clear of  $z_1^{\frac{1}{c}}$ . Therefore, because  $r_1$  is a root of the equation  $X_1 = 0$ ,

$$\left\{ \frac{1}{m} (h_1 \Delta_1^{\frac{m-1}{m}} + \text{etc.}) \right\}^m + (bz_1^{\frac{c-1}{c}} + dz_1^{\frac{c-2}{c}} + \text{etc.}) \left\{ \frac{1}{m} (h_1 \Delta_1^{\frac{m-1}{m}} + \text{etc.}) \right\}^{m-1} + \text{etc.} = 0.$$

All the surds in this equation occur in the simplified expression  $r_1$ . Therefore, by Prop. II.,

$$\left\{ \frac{1}{m} (h_{a+1} \Delta_{a+1}^{\frac{m-1}{m}} + \text{etc.}) \right\}^m + (bz_2^{\frac{c-1}{c}} + dz_2^{\frac{c-2}{c}} + \text{etc.}) \left\{ \frac{1}{m} (h_{a+1} \Delta_{a+1}^{\frac{m-1}{m}} + \text{etc.}) \right\}^{m-1} + \text{etc.} = 0.$$

Therefore  $\frac{1}{m} (h_{a+1} \Delta_{a+1}^{\frac{m-1}{m}} + \text{etc.})$  or  $r_{a+1}$  is a root of the equation

$$X_1 = x^m + (bz_2^{\frac{c-1}{c}} + \text{etc.}) x^{m-1} + \text{etc.} = 0. \quad (18)$$

Therefore also, by Cor. Prop. II., all the terms in (15) are roots of that equation. And, by Cor. 1, the terms in (15) are all unequal.

Therefore the equation  $X_1 = 0$  has  $m$  unequal particular cognate forms of  $R$  for its roots.

§25. Cor. 3. No two of the expressions in (16), as  $x_1$  and  $X_1$ , are identical with one another. For, in order that  $X_1$  and  $X_1$  might be identical, the coefficients of the several powers of  $x$  in  $X_1$  would need to be equal to those of the corresponding powers of  $x$  in  $X_1$ ; but, if

one of the coefficients in  $X_1$  in all unequal  
 §26. C  
 selected, t  
 For, supp  
 Taking th  
 the equati  
 All the su  
 that  $z_2^{\frac{1}{c}}$   
 $\theta_1$  being a  
 $\frac{1}{c}$   
 and  $z_2^{\frac{1}{c}}$   
 Since  $z_2^{\frac{1}{c}}$   
 We may,  
 which  $\Delta_1^{\frac{1}{m}}$   
 $\varphi$   
 +  
 where  $p_1$ ,  
 $m^{\text{th}}$  root of  
 whose valu

distinct from  
 $\frac{1}{z_2^c}$ ; the ex-  
 $\frac{1}{m}$   
 $a+1, h_{a+1}$ , etc.

+ etc.; (17)

$r_1$  is a root of

$m-1$  + etc. = 0.

expression  $r_1$ .

$m-1$  + etc. = 0.

of the equation

= 0. (18)

(15) are roots of  
 are all unequal.

lar cognate forms

as  $x_1$  and  $X_1$ , are

nd  $X_1$  might be

in  $X_1$  would need

$x$  in  $X_1$ ; but, if

one of the coefficients of  $X_1$  be selected in which  $z_1^{\frac{1}{c}}$  is present, this coefficient can be shown to be unequal to the corresponding coefficient in  $X_1$  in the same way in which the terms in (15) were proved to be all unequal.

§26. Cor. 4. Any two of the terms in (16), as  $X_1$  and  $X_1'$ , being selected, the equations  $X_1 = 0$  and  $X_1' = 0$  have no root in common. For, suppose, if possible, that these equations have a root in common. Taking the forms of  $X_1$  and  $X_1'$  in (17) and (18), since  $r_1$  is a root of the equation  $X_1' = 0$ ,

$$r_1^m + (bz_2^{\frac{c-1}{c}} + \text{etc.}) r_1^{m-1} + \text{etc.} = 0. \quad (19)$$

All the surds in this equation except  $z_2^{\frac{1}{c}}$  occur in  $r_1$ . It is impossible that  $z_2^{\frac{1}{c}}$  can occur in  $r_1$ ; for,  $z_1^{\frac{1}{c}}$  occurs in  $r_1$ ; and  $z_2^{\frac{1}{c}} = \theta_1 z_1^{\frac{1}{c}}$ ,  $\theta_1$  being a primitive  $c^{\text{th}}$  root of unity; but this equation, if both  $z_1^{\frac{1}{c}}$  and  $z_2^{\frac{1}{c}}$  occurred in  $r_1$ , would be of the inadmissible type (3).

Since  $z_2^{\frac{1}{c}}$  does not occur in  $r_1$ , it is a principal (see §2) surd in (19). We may, therefore, keeping in view that  $r_1$  is the expression (1) in

which  $J_1^{\frac{1}{m}}$  is a principal surd, arrange (19) thus,

$$\begin{aligned} \varphi(J_1^{\frac{1}{m}}) &= J_1^{\frac{m-1}{m}} (p_1 z_2^{\frac{c-1}{c}} + p_2 z_2^{\frac{c-2}{c}} + \text{etc.}) \\ &+ J_1^{\frac{m-2}{m}} (q_1 z_2^{\frac{c-1}{c}} + q_2 z_2^{\frac{c-2}{c}} + \text{etc.}) + \text{etc.} = 0; \quad (20) \end{aligned}$$

where  $p_1, q_1, \text{etc.}$ , are clear of  $z_2^{\frac{1}{c}}$ . Then,  $\omega_1$  being a primitive  $m^{\text{th}}$  root of unity such that, by changing  $J_1^{\frac{1}{m}}$  into the  $m^{\text{th}}$  root of  $J_1$  whose value is  $\omega_1 J_1^{\frac{1}{m}}$ ,  $r_1$  becomes  $r_2$ ,

$$\begin{aligned} \varphi(\omega_1 \Delta_1^{\frac{1}{m}}) &= \omega_1^{m-1} \Delta_1^{\frac{m-1}{m}} (p_1 z_2^{\frac{c-1}{c}} + \text{etc.}) \\ &+ \omega_1^{m-2} \Delta_1^{\frac{m-1}{m}} (q_1 z_2^{\frac{c-1}{c}} + \text{etc.}) + \text{etc.} \end{aligned} \quad (21)$$

The coefficients of the several powers of  $\Delta_1^{\frac{1}{m}}$  in  $\varphi(\Delta_1^{\frac{1}{m}})$  cannot be all zero; for, if they were, we should have, from (21),  $\varphi(\omega_1 \Delta_1^{\frac{1}{m}}) = 0$ . This means that  $r_2$  is a root of the equation  $X_1 = 0$ . But in like manner all the terms in (14) would be roots of that equation, and  $X_1$  would be identical with  $X$ ; which, by Cor. 3, is impossible. Since the coefficients of the different powers of  $\Delta_1^{\frac{1}{m}}$  in  $\varphi(\Delta_1^{\frac{1}{m}})$  are not all zero, the equation (20) gives us, by §5,  $\omega \Delta_1^{\frac{1}{m}} = l_1$ ,  $\omega$  being an  $m^{\text{th}}$  root of unity, and  $l_1$  involving only surds in  $\varphi(\Delta_1^{\frac{1}{m}})$  exclusive of  $\Delta_1^{\frac{1}{m}}$ . In  $l_1$  we may conceive  $z_2^{\frac{1}{c}}$  changed into  $\theta_1 z_1^{\frac{1}{c}}$ . Then  $l_1$  involves only surds distinct from  $\Delta_1^{\frac{1}{m}}$ , all of them except the primitive  $c^{\text{th}}$  root of unity  $\theta_1$  being surds that occur in  $r_1$ . This makes the equation  $\omega \Delta_1^{\frac{1}{m}} = l_1$  of the inadmissible type (3). Hence the equations  $X_1 = 0$  and  $X_1 = 0$  have no root in common.

§27. Cor. 5. Let  $X_2$  be the continued product of the terms in (16).

Then  $X_2$ , modified according to §21, is clear of  $z_1^{\frac{1}{c}}$ , in the same way in which  $X_1$  is clear of  $\Delta_1^{\frac{1}{m}}$ . Also since, by Cor. 2, each of the equations  $X_1 = 0$ ,  $X_1 = 0$ , etc., has  $m$  unequal particular cognate forms of  $R$  for its roots, and since, by Cor. 4, no two of these equations have a root in common, the  $mc$  roots of the equation  $X_2 = 0$  are unequal particular cognate forms of  $R$ .

§28. Pr  
according

$m^{\text{th}}$  degree  
of the high  
ly surds  
 $m^{\text{th}}$  root of

First. le  
in  $r_1$ . T  
§24, and  
surd  $\Delta_1^{\frac{1}{m}}$ .

$X_2$  be form

at  $X_1$ , an  
equation  $X_1$   
second step

surd  $z_1^{\frac{1}{c}}$ ,  
unequal pa  
the series  
enter that  
 $X_2 = 0$  be  
modified a  
putting  $d$

$X_2 = x_d - ($

where, one  
of unity a  
the sum o

or  $\frac{n -$   
 $m$

$n^{\text{th}}$  roots o  
and  $C_2$  to  
at the clos  
which is t  
the roots o  
subject to  
 $C_1$  in  $X_e$

§28. PROPOSITION VI. Let the simplified expression  $r_1$ , modified according to §21, be a root of the rational irreducible equation of the  $n^{\text{th}}$  degree,  $F(x) = 0$ . Then if  $\Delta_1^{\frac{1}{m}}$ , not a root of unity, be a surd of the highest rank in  $r_1$ ,  $N$  is a multiple of  $m$ . But if  $r_1$  involve only surds that are roots of unity, one of them being the primitive  $n^{\text{th}}$  root of unity,  $N$  is a multiple of a measure of  $n - 1$ .

(21)

First, let  $\Delta_1^{\frac{1}{m}}$ , not a root of unity, be a surd of the highest rank in  $r_1$ . Taking the expression (1) to be  $r_1$ , let  $X_1$  be formed as in §24, and let it be modified according to §21. It is clear of the

surd  $\Delta_1^{\frac{1}{m}}$ . Should it involve a surd that is not a root of unity, let  $X_2$  be formed as in §27. Setting out from  $r_1$  we arrived by one step at  $X_1$ , an expression clear of  $\Delta_1^{\frac{1}{m}}$ , and such that the roots of the equation  $X_1 = 0$  are unequal particular cognate forms of  $R$ . A second step brought us to  $X_2$ , an expression clear of the additional surd  $z_1^{\frac{1}{o}}$ , and such that the  $mc$  roots of the equation  $X_2 = 0$  are unequal particular cognate forms of  $R$ . Thus we can go on till, in the series  $X_1, X_2$ , etc., we reach a term  $X_e$  into which no surds enter that are not roots of unity, the  $mc \dots l$  roots of the equation  $X_e = 0$  being unequal particular cognate forms of  $R$ . Should  $X_e$  be modified according to §21, not be rational, its form, by Prop. IV., putting  $d$  for  $mc \dots l$ , is

$$X_e = x^d - (p_1 C_1 + \dots + p_m C_m) x^{d-1} + (q_1 C_1 + \dots + q_m C_m) x^{d-2} + \text{etc.};$$

where, one of the roots occurring in  $X_e$  being the primitive  $n^{\text{th}}$  root of unity  $\omega_1$ , the coefficients  $p_1, q_1$ , etc., are clear of  $\omega_1$ ; and  $C_1$  is the sum of the cycle of primitive  $n^{\text{th}}$  roots of unity (8) containing

$\frac{n-1}{m}$  terms; and, the cycle (9) containing all the primitive

$n^{\text{th}}$  roots of unity, the change of  $\omega_1$  into  $\omega_1^{\beta}$  causes  $C_1$  to become  $C_2$ , and  $C_2$  to become  $C_3$ , and so on,  $C_m$  becoming  $C_1$ . As was explained at the close of §20, the cycle (8) may be reduced to a single term, which is then identical with  $C_1$ . It will also not be forgotten that the roots of unity such as the  $n^{\text{th}}$  here spoken of are, according to §1, subject to the condition that the numbers such as  $n$  are prime. When  $C_1$  in  $X_e$  is changed successively into  $C_1, C_2$ , etc., let  $X_e$  become

$$X_e, X_e', X_e'', \dots, X_e^{(m-1)} \tag{22}$$



If  $X_{e+1}$  be the continued product of the terms in (22), the  $dm$  roots of the equation  $X_{e+1} = 0$  can be shown to be unequal particular cognate forms of  $R$ . For, no two terms in (22) as  $X_e$  and  $X_e$  are identical; because, if they were,  $X_e$  would remain unaltered by the change of  $\omega_1$  into  $\omega_1^\beta$ ; which, by Prop. IV., because  $\omega_1^\beta$  is not a term in the cycle (8), is impossible. It follows that no two of the equations  $X_e = 0$ ,  $X_e = 0$ , etc., have a root in common. For, if the equations  $X_e = 0$ , and  $X_e = 0$  had a root in common, since  $X_e$  and  $X_e$  are not identical,  $X_e$  would have a lower measure involving only surds found in  $X_e$ , because the surds in  $X_e$  are the same with those in  $X_e$ . Let  $\varphi(x)$  be this lower measure of  $X_e$ , and let  $r_1$  be a root of the equation  $\varphi(x) = 0$ . Then, by Cor. Prop. II., all the  $d$  roots of the equation  $X_e = 0$  are roots of the equation  $\varphi(x) = 0$ ; which is impossible. In the same way it can be proved that no equation in the series  $X_e = 0$ ,  $X_e = 0$ , etc., has equal roots. Since no one of these equations has equal roots, and no two of them have a root in common, the  $dm$  roots of the equation  $X_{e+1} = 0$  are unequal particular cognate forms of  $R$ . Also  $X_{e+1}$ , modified according to §21, is clear of the primitive  $n^{\text{th}}$  roots of unity. Should  $X_{e+1}$  not be rational, we can deal with it as we did with  $X_e$ . Going on in this way, we ultimately reach a rational expression  $X_z$  such that the  $dm \dots g$  roots of the equation  $X_z = 0$  are unequal particular cognate forms of  $R$ . This equation must be identical with the equation  $F(x) = 0$  of which  $r_1$  is a root. For, by Prop. III., the equation  $F(x) = 0$  has for its roots the unequal particular cognate forms of  $R$ . Therefore, because the roots of the equation  $X_z = 0$  are all unequal and are at the same time particular cognate forms of  $R$ ,  $X_z$  must be either a lower measure of  $F(x)$  or identical with  $F(x)$ . But  $F(x)$ , being irreducible, has no lower measure. Therefore  $X_z$  is identical with  $F(x)$ . Therefore, the equation  $F(x) = 0$  being the  $N^{\text{th}}$  degree,  $N = mc \dots lm \dots g$ . Hence  $N$  is a multiple of  $m$ . This is the

result arrived at when  $r_1$  involves a surd of the highest rank  $\Delta_1^{\frac{1}{m}}$  not a root of unity. Should  $r_1$  involve no surds except roots (see §1) of unity, we should then have set out from  $X_e$  regarded as identical with  $x - r_1$ . The result would have been  $N = m \dots g$ . Therefore  $N$  is a multiple of  $m$ ; and, because  $m$  is here the number of cycles of  $s$  terms each, that make up the series of the primitive  $n^{\text{th}}$  roots of unity,  $ms = n - 1$ . Therefore  $N$  is a multiple of a measure of  $n - 1$ .

§29. Cor. Let  $N$  be a prime number. Then, if  $r_1$  involve a surd of the highest rank  $\Delta_1^{\frac{1}{m}}$  not a root (see §1) of unity,  $N = m$ ; for,

the series  
reduced  
unity,  $n$   
because  $N$   
 $n - 1 =$

THE SOLV

§30. Th  
by an exa  
 $m^{\text{th}}$  degree  
though it  
under a co  
root  $r_1$  is,  
roots of u  
denominat  
may be sai  
of the seco

§31. In  
§21, if one  
unity  $\omega_1$ ,  
 $X_e$  in Pro

The  $m$  roo  
have

For, by I  
the expre  
equation.  
For, tak  
( $p_m - 1$ )  
each of  
 $p_1, p_2,$   
that the  
§21, is ir  
 $F(x) =$

the series of integers  $m, c$ , etc., of which  $N$  is the continued product, reduced to its first term. If  $r_1$  involve only surds that are roots of unity,  $n - 1$  is a multiple of  $N$ ; for  $N = m \dots g$ ; therefore, because  $N$  is prime, it is equal to  $m$ ; but  $ms = n - 1$ ; therefore  $s - 1 = sN$ .

**THE SOLVABLE IRREDUCIBLE EQUATION OF THE  $m^{\text{th}}$  DEGREE,  $m$  PRIME.**

§30. The principles that have been established may be illustrated by an examination of the solvable irreducible rational equation of the  $m^{\text{th}}$  degree  $F(x) = 0$ ,  $m$  being prime. Two cases may be distinguished, though it will be found that the roots can in the two cases be brought under a common form; the one case being that in which the simplified root  $r_1$  is, and the other that in which it is not, a rational function of roots of unity, that is, according to §1, of roots of unity having the denominators of their indices prime numbers. The equation  $F(x) = 0$  may be said to be in the former case of the *first class*, and in the latter of the *second class*.

**THE EQUATION  $F(x) = 0$  OF THE FIRST CLASS.**

§31. In this case, by Cor. Prop. VI.,  $r_1$  being modified according to §21, if one of the roots involved in  $r_1$  be the primitive  $n^{\text{th}}$  root of unity  $\omega_1$ ,  $n - 1$  is a multiple of  $m$ . Also the expression written  $X_s$  in Prop. VI. is reduced to  $x - r_1$ , so that

$$r_1 = p_1C_1 + p_2C_2 + \dots + p_mC_m.$$

The  $m$  roots of the equation  $F(x) = 0$  being  $r_1, r_2$ , etc, we must have

$$\left. \begin{aligned} r_1 &= p_1 C_1 + p_2 C_2 + \dots + p_m C_m, \\ r_2 &= p_m C_1 + p_1 C_2 + \dots + p_{m-1} C_m, \\ &\dots\dots\dots \\ r_m &= p_2 C_1 + p_3 C_2 + \dots + p_1 C_m. \end{aligned} \right\} \quad (23)$$

For, by Prop. II., because  $r_1$  is a root of the equation  $F(x) = 0$ , all the expressions on the right of the equations (23) are roots of that equation. And no two of these expressions are equal to one another. For, take the first two. If these were equal, we should have  $(p_m - p_1) C_1 + (p_1 - p_2) C_2 + \text{etc.} = 0$ . Therefore, by §13, each of the terms  $p_m - p_1, p_1 - p_2$ , etc., is zero. This makes  $p_1, p_2$ , etc., all equal to one another. Therefore  $r_1 = - p_1$ ; so that the primitive  $n^{\text{th}}$  root of unity is eliminated from  $r_1$ ; which, by §21, is impossible. Hence the values of the  $m$  roots of the equation  $F(x) = 0$  are those given in (23).

§32. Let  $r_1$  be one of the particular cognate forms of the generic expression  $R$  under which the simplified expression  $r_1$  falls. Then, because, by Prop. II., all the particular cognate forms of  $R$  are roots of the equation  $F(x) = 0$ ,  $r_1$  is equal to one of the  $m$  terms  $r_1, r_2, \dots$ , etc., say to  $r_s$ . I will now show that the changes of the surds involved that cause  $r_1$  to become  $r_1$ , whose value is  $r_s$ , cause  $r_2$  to receive the value  $r_{s+1}$ , and  $r_3$  to receive the value  $r_{s+2}$ , and so on. This may appear obvious on the face of the equations (23); but, to prevent misunderstanding, the steps of the deduction are given. Any changes made in  $r_1$  must transform  $C_1$  into  $C_s$ , one of the  $m$  terms  $C_1, C_2, \dots$ . In passing from  $r_1$  to  $r_1$ , while  $C_1$  becomes  $C_s$ , let  $r_2$  become  $r_2$ , and  $p_1$  become  $p_1$ , and  $p_2$  become  $p_2$ , and so on. The change that causes  $C_1$  to become  $C_s$  transforms  $C_2$  into  $C_{s+1}$ , and  $C_3$  into  $C_{s+2}$ , and so on. Therefore, it being understood that  $p_{m+1}, C_{m+1}, \dots$ , are the same as  $p_1, C_1, \dots$ , respectively,

$$r_1 = p_1 C_s + p_2 C_{s+1} + \text{etc.},$$

$$\text{and } r_2 = p_m C_s + p_1 C_{s+1} + \text{etc.};$$

which may be otherwise written

$$\left. \begin{aligned} r_1 &= p_{m+2-s} C_1 + p_{m+3-s} C_2 + \text{etc.}, \\ r_2 &= p_{m+1-s} C_1 + p_{m+2-s} C_2 + \text{etc.} \end{aligned} \right\} \quad (24)$$

Therefore, from (24) and (23),

$$C_1(p_{m+2-s} - p_{m+2-s}) + C_2(p_{m+3-s} - p_{m+3-s}) + \text{etc.} = 0.$$

Therefore, by §13,  $p_{m+2-s} = p_{m+2-s}, p_{m+3-s} = p_{m+3-s}, \dots$ , etc.

Hence the second of the equations (24) becomes

$$r_2 = p_{m+1-s} C_1 + p_{m+2-s} C_2 + \text{etc.} = r_{s+1}.$$

Thus  $r_2$  is transformed into  $r_{s+1}$ . In like manner  $r_3$  receives the value  $r_{s+2}$ , and so on.

§33. By Cor. Prop. VI., the primitive  $n^{\text{th}}$  root of unity being one of those involved in  $r_1$ ,  $n - 1$  is a multiple of  $m$ . In like manner, if the primitive  $a^{\text{th}}$  root of unity be involved in  $r_1$ ,  $a - 1$  is a multiple of  $m$ , and so on. Therefore, if  $t_1$  be the primitive  $m^{\text{th}}$  root of unity,  $t_1$  is distinct from all the roots involved in  $r_1$ .

§34. For  $r_m$ , be a etc., and if,  $t_1^m$

$r_s + r_c + r_s = r_s$ . is reached and so on  $r_1, r_2, \dots$  by §12,  $m^{\text{th}}$  roots  $mr_c - (r_1$

§35. P

the terms are the  $r \varphi(x) = F(x) =$  For, cognate particular  $m^{\text{th}}$  root in  $r_1, r_{s+1}, \dots$

contain number indiffer

The co can pre



from the different values of  $t$  that are taken in  $\mathcal{J}'$ , while the expressions  $r_1, r_2, \text{etc.}$ , remain unaltered. And  $t$  has not more than  $m - 1$  values. Hence there are not more than  $m - 1$  unequal particular cognate forms of  $\mathcal{J}$ . But the  $m - 1$  forms obtained by taking the different values of  $t$  in  $\mathcal{J}'$  are all unequal. For, selecting  $t_1$  and  $t_1^a$ , two distinct values of  $t$ , suppose if possible that

$$(r_1 + t_1 r_2 + \text{etc.})^m = (r_1 + t_1^a r_2 + \text{etc.})^m$$

$$\therefore t_1^s (r_1 + t_1 r_2 + \text{etc.}) = r_1 + t_1^a r_2 + \text{etc.},$$

$s$  being a whole number. This may be written

$$r_{m+1-s} + r_{m+2-s} t_1 + \text{etc.} = r_1 + t_1^a r_2 + \text{etc.} \quad (28)$$

Therefore, by §34,  $r_{m+1-s} = r_1$ . This means, since all the  $m$  terms  $r_1, r_2, \text{etc.}$ , are unequal, that  $s = 0$ . Hence (28) becomes

$$r_1 + r_2 t_1 + \text{etc.} = r_1 + r_2 t_1^a + \text{etc.}$$

Therefore

$$\begin{aligned} r_2 + r_3 t_1^a + \text{etc.} &= r_2 t_1^{1-a} + r_3 t_1^{2-a} + \text{etc.} \\ &= r_{a+1} + r_{a+2} t_1 + \text{etc.} \end{aligned}$$

Therefore, by §35,  $r_2 = r_{a+1}$ . Therefore, because all the  $m$  terms  $r_1, r_2, \text{etc.}$ , are unequal,  $a = 1$ ; which, because  $t_1$  and  $t_1^a$  were supposed to be distinct primitive  $m^{\text{th}}$  roots of unity, is impossible. Therefore no two of the terms in (27) are equal to one another. And it has been proved that there is no particular cognate form of  $\mathcal{J}$  which is not equal to a term in (27). Therefore the terms in (27) are the unequal particular cognate forms of  $\mathcal{J}$ . Therefore, by Prop. III, they are the roots of a rational irreducible equation.

§36. PROPOSITION VIII. The roots of the equation  $\varphi(x) = 0$  auxiliary (see §35) to  $F(x) = 0$  are rational functions of the primitive  $m^{\text{th}}$  root of unity.

For, let the value of  $\mathcal{J}_1$ , obtained from (26), and modified according to §21, be

$$\mathcal{J}_1 = k_1 + k_2 t_1 + k_3 t_1^2 + \dots + k_m t_1^{m-1},$$

where  $k_1, k_2, \text{etc.}$ , are clear of  $t_1$ . Suppose if possible that  $k_1, k_2, \text{etc.}$ , are not rational. We may take the primitive  $n^{\text{th}}$  root of unity  $\omega_1$  to be present in these coefficients. But  $\omega_1$  occurs in  $r_1, r_2, \text{etc.}$ , and therefore also in  $\mathcal{J}_1$ , only in the expressions  $C_1, C_2, \text{etc.}$ . Therefore  $\mathcal{J}_1 = d_1 C_1 + \dots + d_m C_m$ ; where  $d_1, \text{etc.}$ , are clear of  $\omega_1$ . The coefficients  $d_1, d_2, \text{etc.}$ , cannot all be equal; for this would make  $\mathcal{J}_1 = -d_1$ ; which, by §21, is impossible. Hence  $m$  unequal

values of  
successively

To show t  
If these w

Therefore  
because d  
Since the  
by Prop.  
not lower  
Therefore  
in (27) is

§37. C  
which is s  
generic ex  
for  $t_1$  succ  
 $k_1, k_2, \text{e}$   
the Propo  
function  
 $\mathcal{J}_1$  satisfie

§38. F  
 $F(x) =$

For, s

$p_2 = ($

Multipl

$p_2 = ($

Hence  
 $r_2, \text{ and}$

values of the generic expression  $J$  are obtained by changing  $C_1$  successively into  $C_1, C_2, \text{ etc.}$ , namely,

$$\begin{aligned} & d_1 C_1 + d_2 C_2 + \dots + d_m C_m, \\ & d_m C_1 + d_1 C_2 + \dots + d_{m-1} C_m, \\ & \dots\dots\dots \\ & d_2 C_1 + d_3 C_2 + \dots + d_1 C_m. \end{aligned}$$

To show that these expressions are all unequal, take the first two. If these were equal, we should have

$$(d_m - d_1) C_1 + (d_1 - d_2) C_2 + \text{etc.} = 0.$$

Therefore, by §13,  $d_m - d_1 = 0, d_1 - d_2 = 0$ , and so on; which, because  $d_1, d_2, \text{ etc.}$ , are not all equal to one another, is impossible. Since then  $J$  has at least  $m$  unequal particular cognate forms,  $J_1$  is, by Prop. III., the root of a rational irreducible equation of a degree not lower than the  $m^{\text{th}}$ ; which, by Prop. VII., is impossible. Therefore  $k_1, k_2, \text{ etc.}$ , are rational. Hence each of the expressions in (27) is a rational function of  $t_1$ .

§37. Cor. Any expression of the type  $k_1 + k_2 t_1 + k_3 t_1^2 + \text{etc.}$ , which is such that all the unequal particular cognate forms of the generic expression under which it falls are obtained by substituting for  $t_1$  successively the different primitive  $m^{\text{th}}$  roots of unity, while  $k_1, k_2, \text{ etc.}$ , remain unaltered, is a rational function of  $t_1$ . For, in the Proposition,  $J_1$  or  $k_1 + k_2 t_1 + \text{etc.}$  was shown to be a rational function of  $t_1$ , the conclusion being based on the circumstance that  $J_1$  satisfies the condition specified.

§38. PROPOSITION IX. If  $g$  be the sum of the roots of the equation  $F(x) = 0$ ,

$$\begin{aligned} r_2 = \frac{1}{m} (g + J_1^{\frac{1}{m}} + a_1 J_1^{\frac{2}{m}} + b_1 J_1^{\frac{3}{m}} + \dots \\ + c_1 J_1^{\frac{m-2}{m}} + h_1 J_1^{\frac{m-1}{m}}); \end{aligned} \tag{29}$$

For,  $\varepsilon$  being one of the whole numbers,  $1, 2, \dots, m - 1$ , put

$$p_\varepsilon = (r_1 + t_1^\varepsilon r_2 + t_1^{2\varepsilon} r_3 + \text{etc.}) (r_1 + t_1 r_2 + t_1^2 r_3 + \text{etc.})^{-\varepsilon}. \tag{30}$$

Multiply the first of its factors by  $t_1^{-\varepsilon}$  and the second by  $t_1^\varepsilon$ . Then

$$p_\varepsilon = (r_2 + t_1^\varepsilon r_3 + t_1^{2\varepsilon} r_4 + \text{etc.}) (r_2 + t_1 r_3 + t_1^2 r_4 + \text{etc.})^{-\varepsilon}. \tag{31}$$

Hence  $p_\varepsilon$  does not alter its value when we change  $r_1$  into  $r_2, r_2$  into  $r_3$ , and so on. In like manner it does not alter its value when we

change  $r_1$  into  $r_a$ ,  $r_2$  into  $r_{a+1}$ , and so on. Therefore, by §33,  $p_z$  is not changed by any alterations that may be made in  $r_1, r_2$ , etc., while  $t_1$  remains unaltered. Consequently, if  $p_z$  be a particular cognate form of  $P$ , all the unequal particular cognate forms of  $P$  are obtained by substituting for  $t_1$  successively in  $p_z$  the different primitive  $m^{\text{th}}$  roots of unity, while  $r_1, r_2$ , etc., remain unaltered. Therefore, by Cor., Prop. VIII.,  $p_z$  is a rational function of  $t_1$ . When  $z = 2$ , let  $p_z = a_1$ ; when  $z = 3$ , let  $p_z = b_1$ , and so on. Then, from

$$(26) \text{ and } (30), J_2^{\frac{1}{m}} = a_1 J_1^{\frac{1}{m}}, J_3^{\frac{1}{m}} = b_1 J_1^{\frac{1}{m}} \text{ and so on. But, from}$$

(27), since  $g$  is the sum of the roots of the equation  $F(x) = 0$ ,

$$r_1 = \frac{1}{m} (g + J_1^{\frac{1}{m}} + J_2^{\frac{1}{m}} + \dots + J_{m-1}^{\frac{1}{m}}).$$

By putting  $a_1 J_1^{\frac{1}{m}}$  for  $J_2^{\frac{1}{m}}$ ,  $b_1 J_1^{\frac{1}{m}}$  for  $J_3^{\frac{1}{m}}$  and so on, this becomes

(29). Because  $a_1, b_1$ , etc., are rational functions of  $t_1$ , while  $J_1$ , the root of a rational irreducible equation of the  $(m-1)^{\text{th}}$  degree, is also a rational function of  $t_1$ , the coefficients  $a_1, b_1$ , etc., involve no surd that is not subordinate to  $J_1^{\frac{1}{m}}$ .

§39. PROPOSITION X. If the prime number  $m$  be odd, the expressions

$$J_1^{\frac{1}{m}} J_{m-1}^{\frac{1}{m}}, J_2^{\frac{1}{m}} J_{m-2}^{\frac{1}{m}}, \dots, J_{\frac{m-1}{2}}^{\frac{1}{m}} J_{\frac{m+2}{m}}^{\frac{1}{m}}, \tag{32}$$

are the roots of a rational equation of the  $\left(\frac{m-1}{2}\right)^{\text{th}}$  degree.

By §32, when  $r_1$  is changed into  $r_2$ ,  $r_2$  becomes  $r_{2+1}$ ,  $r_3$  becomes  $r_{2+2}$ , and so on. Hence the terms  $r_1 r_2, r_2 r_3, \dots, r_m r_1$ , form a cycle, the sum of the terms in which may be denoted by the symbol  $\Sigma_1^1$ . In like manner the sum of the terms in the cycle  $r_1 r_3, r_2 r_4, \dots, r_m r_2$ , may be written  $\Sigma_3^1$ . And so on. In harmony with this notation, the sum of the  $m$  terms  $r_1^2, r_2^2$ , etc., may be written  $\Sigma_1^2$ . Now  $r_1$  can only be changed into one of the terms  $r_1, r_2$ , etc.; and we have seen that, when it becomes  $r_2$ ,  $r_2$  becomes  $r_{2+1}$ , and so on. Such changes leave the cycle  $r_1 r_2, r_2 r_3$ , etc., as a whole unaltered.

Therefo  
rational  
has a ra  
But  $\Sigma_1^1$   
Therefo  
 $J_1^{\frac{1}{m}}$   
Hence, s  
(32) are  
sively in  
roots of  
§40. F  
number s  
obtain th  
one of it  
nth roots  
reducing  
§41. F  
ducing  
 $x^n - 1$   
Let th  
(8), wh  
symbol  
since s i  
us reaso  
pairs of  
cycle co

e, by §33,  $p_x$  is  
 n  $r_1, r_2, \dots$ ,  
 e a particular  
 forms of  $P$  are  
 different primi-  
 altered. There  
 of  $t_1$ . When  
 n. Then, from

Therefore, by Prop. III.,  $\Sigma_2^1$  is the root of a simple equation, or has a rational value. In like manner each of the expressions

$$\Sigma_1^1, \Sigma_2^1, \Sigma_3^1, \dots, \Sigma_m^1, \tag{33}$$

has a rational value. From (26), by actual multiplication,

$$J_1^{\frac{1}{m}} J_{m-1}^{\frac{1}{m}} = \Sigma_1^1 + (\Sigma_2^1) t_1 + (\Sigma_3^1) t_1^2 + \text{etc.}$$

on. But, from

But  $\Sigma_2^1, \Sigma_3^1, \dots$ , etc., are respectively identical with  $\Sigma_m^1, \Sigma_{m-1}^1, \dots$ . Therefore

$$J_1^{\frac{1}{m}} J_{m-1}^{\frac{1}{m}} = \Sigma_1^1 + (\Sigma_2^1) (t_1 + t_1^{-1}) + (\Sigma_3^1) (t_1^2 + t_1^{-2}) + \text{etc.} \tag{34}$$

- 1).

on, this becomes

Hence, since the terms in (33) are all rational, and since the terms in (32) are respectively what  $J_1^{\frac{1}{m}} J_{m-1}^{\frac{1}{m}}$  becomes by changing  $t_1$  successively into the  $\frac{m-1}{2}$  terms  $t_1, t_1^2, \dots$ , the terms in (32) are the roots of a rational equation of the  $\left(\frac{m-1}{2}\right)^{\text{th}}$  degree.

$t_1$ , while  $J_1$ , the  
 )th degree, is also  
 , involve no surd

m be odd, the

§40. For the solution of the equation  $x^n - 1 = 0$ ,  $n$  being a prime number such that  $m$  is a prime measure of  $n - 1$ , it is necessary to obtain the solution of the equation of the  $m^{\text{th}}$  degree which has for one of its roots the sum of the  $\frac{n-1}{m}$  terms in a cycle of primitive  $n^{\text{th}}$  roots of unity. This latter equation will be referred to as the *reducing Gaussian equation* of the  $m^{\text{th}}$  degree to the equation

$$J_{\frac{m+2}{n}}^{\frac{1}{m}}, \tag{32}$$

$$x^n - 1 = 0.$$

)th degree.

§41. PROPOSITION XI. When the equation  $F(x) = 0$  is the reducing Gaussian (see §40) of the  $m^{\text{th}}$  degree to the equation  $x^n - 1 = 0$ , each of the  $\frac{m-1}{2}$  expressions in (32) is equal to  $n$ .

$r_{x+1}, r_3$  become  
 ...  $r_m r_1$ , form  
 ted by the symbo  
 cycle  $r_1 r_3, r_2 r_4$

In harmony with  
 may be written  $\Sigma_1^1$   
 $r_1, r_2, \dots$ ; and  
 $r_{x+1}$ , and so on  
 a whole unaltered

Let the sum of the primitive  $n^{\text{th}}$  roots of unity forming the cycle (8), which sum has in preceding sections been indicated by the symbol  $C_1$ , be the root  $r_1$  of the equation  $F(x) = 0$ . This implies, since  $s$  is the number of the terms in (8), that  $ms = n - 1$ . Let us reason first on the assumption that the cycle (8) is made up of pairs of reciprocal roots  $\omega_1$  and  $\omega_1^{-1}$ , and so on. Then, because the cycle consists of  $\frac{s}{2}$  pairs of reciprocal roots,  $C_1^2$  or  $r_1^2$  is the sum of



$s^2$  terms, each an  $n^{\text{th}}$  root of unity. Among these unity occurs  $s$  times. Let  $\omega_1$  occur  $h_1$  times; and let  $\omega_1^\wedge$  the second term in (8) occur  $h'$  times. Since  $\omega_1^\wedge$  may be made the first term in the cycle (8), it must, under the new arrangement, present itself in the value of  $r_1^2$ , precisely where  $\omega_1$  previously appeared. That is to say,  $h' = h_1$ . In like manner each of the terms in (8) occurs exactly  $h_i$  times in the expression for  $r_1^2$ . The cycle (9) being that which contains all the primitive  $n^{\text{th}}$  roots of unity, let us, adhering to the notation of previous sections, suppose that, when  $\omega_1$  is changed into  $\omega_1^\beta$ ,  $C_1$  or  $r_1$  becomes  $C_2$  or  $r_2$ ,  $C_2$  or  $r_2$  becomes  $C_3$  or  $r_3$ , and so on. On the same grounds on which every term in (8) occurs the same number of times in the value of  $r_1^2$ , each term in the cycle of terms whose sum is  $C_2$  occurs the same number of times; and so on. Therefore

$$\begin{aligned} r_1^2 &= s + h_1 C_1 + h_2 C_2 + \dots + h_m C_m. \\ r_2^2 &= s + h_m C_1 + h_1 C_2 + \dots + h_{m-1} C_m, \\ &\dots\dots\dots \\ r_m^2 &= s + h_2 C_1 + h_3 C_2 + \dots + h_1 C_m. \end{aligned}$$

Therefore, keeping in view (11),  $\Sigma_1^1 = ms - (h_1 + h_2 + \dots + h_m)$ . But  $s^2 - s$  is the number of the terms in the value of  $r_1^2$  which are primitive  $n^{\text{th}}$  roots of unity. And this must be equal to

$$s(h_1 + \dots + h_m).$$

Therefore

$$h_1 + h_2 + \dots + h_m = s - 1 \dots \Sigma_1^1 = ms + 1 - s = n - s.$$

Again, because  $r_1$  is made up of pairs of reciprocal roots, and because therefore unity does not occur among the  $s^2$  terms of which  $r_1 r_2$  is the sum,

$$\begin{aligned} r_1 r_2 &= k_1 C_1 + k_2 C_2 + \dots + k_m C_m, \\ r_2 r_3 &= k_m C_1 + k_1 C_2 + \dots + k_{m-1} C_m, \\ &\dots\dots\dots \\ r_m r_1 &= k_2 C_1 + k_3 C_2 + \dots + k_1 C_m; \end{aligned}$$

where  $k_1, k_2$ , etc., are whole numbers whose sum is  $s$ . Therefore  $\Sigma_2^1 = -s$ . In like manner each of the terms in (33) except the first is equal to  $-s$ . Therefore (34) becomes

$$\frac{1}{\Delta_1^m} \frac{1}{\Delta_{m-1}^m} = (n - s) - s(t_1 + t_1^2 + \text{etc.}) = n.$$

Let us re  
of pair  
roots. B  
ards the  
ny  $C_2$ , r  
sum are r  
 $C_1$  to  $C_2$ ,  
This bein  
 $\Sigma_1^1 = n -$   
 $C_2$  be sele  
of those o  
from (34)

In like ma  
have the v

§42. Tw  
preceding  
the third d  
which give

The next  
Equations  
equation  
which give

Let us reason now on the assumption that the cycle (8) is not made up of pairs of reciprocal roots. It contains in that case no reciprocal roots. By the same reasoning as above we get  $\Sigma_1^1 = -s$ . As regards the terms in (33) after the first, one of the terms  $C_1, C_2, \text{etc.}$ , say  $C_s$ , must be such that the  $n^{\text{th}}$  roots of unity of which it is the sum are reciprocals of those of which  $C_1$  is the sum. In passing from  $C_1$  to  $C_s$ , we change  $r_1$  into  $r_s$ . In fact,  $C_1$  being  $r_1$ ,  $C_s$  is  $r_s$ . This being kept in view, we get, by the same reasoning as above,  $\Sigma_s^1 = n - s$ . But, if any of the expressions  $C_1, C_2, \text{etc.}$ , except  $C_a$ , be selected, say  $C_a$ , none of the roots in (8) are reciprocals of any of those of which  $C_a$  is the sum. Therefore  $\Sigma_a^1 = -s$ . Therefore, from (34)

$$\begin{aligned} J_1^{\frac{1}{m}} J_{m-1}^{\frac{1}{m}} &= -s + (n-s) t_1^{z-1} \\ -s \left\{ (t_1 + t_1^2 + \dots + t_1^{m-1}) - t_1^{z-1} \right\} &= n. \end{aligned}$$

In like manner every one of the expressions in (34) can be shown to have the value  $n$ .

§42. Two numerical illustrations of the law established in the preceding section may be given. The reducing Gaussian equation of the third degree to the equation  $x^{10} - 1 = 0$  is  $x^9 - x^2 - 6x - 7 = 0$ ; which gives

$$\begin{aligned} r_1 &= \frac{1}{3} (-1 + J_1^{\frac{1}{3}} + J_2^{\frac{1}{3}}), \\ 2J_1 &= 19 (7 + 3 \sqrt{3}), \\ 2J_2 &= 19 (7 - 3 \sqrt{3}), \\ J_1^{\frac{1}{3}} J_2^{\frac{1}{3}} &= 19. \end{aligned}$$

The next example is taken from Lagrange's Theory of Algebraical Equations, Note XIV., §30. The Gaussian of the fifth degree to the equation  $x^{11} - 1 = 0$  is  $x^5 + x^4 - 4x^3 - 3x^2 + 3x + 1 = 0$ ; which gives

$$\begin{aligned} r_1 &= \frac{1}{5} (-1 + J_1^{\frac{1}{5}} + J_2^{\frac{1}{5}} + J_3^{\frac{1}{5}} + J_4^{\frac{1}{5}}); \\ 4 J_1 &= 11 (-89 - 25 \sqrt{5} + 5p - 45q), \\ 4 J_2 &= 11 (-89 + 25 \sqrt{5} - 45p - 5q), \\ 4 J_3 &= 11 (-89 - 25 \sqrt{5} - 5p + 45q), \\ 4 J_4 &= 11 (-89 + 25 \sqrt{5} + 45p + 5q), \\ p &= \sqrt{(-5 - 2 \sqrt{5})}, \\ q &= \sqrt{(-5 + 2 \sqrt{5})}, \\ pq &= -\sqrt{5} \therefore J_1 J_4 = 11^5. \end{aligned}$$



The  $m - 1$  terms in the first of the groups (37) are the  $\frac{m-1}{2}$  terms in (32) each taken twice. Therefore, by Prop. X., the law enunciated in the present Proposition is established so far as this groupe is concerned. The general proof is as follows. By (30) in

§38, taken in connection with (26),  $p_{m-z} J_1^{\frac{m-z}{m}} = J_{m-z}^{\frac{1}{m}}$ . There-

fore  $J_1^{\frac{z}{m}} J_{m-z}^{\frac{1}{m}} = p_{m-z} J_1$ . But, by §38,  $p_{m-z}$  is a rational function of  $t_1$ ; and, by Prop. VIII.,  $J_1$  is a rational function of  $t_1$ .

Therefore  $J_1^{\frac{z}{m}} J_{m-z}^{\frac{1}{m}}$  is a rational function of  $t_1$ . Also from the manner in which  $p_{m-z}$  is formed, when  $t_1$  in  $p_{m-2} J_1$  is changed successively into  $t_1 t_1^2, \dots, t_1^{m-1}$ , the expression  $J_1^{\frac{z}{m}} J_{m-z}^{\frac{1}{m}}$  is changed successively into the  $m - 1$  terms of that one of the groups (37) whose first term is  $J_1^{\frac{z}{m}} J_{m-z}^{\frac{1}{m}}$ . Therefore the terms in that group are the roots of a rational equation.

§45. Cor. The law established in the Proposition may be brought under a yet wider generalization. The expression

$$J_1^{\frac{a}{m}} J_2^{\frac{b}{m}} J_3^{\frac{c}{m}} \dots J_{m-1}^{\frac{s}{m}} \tag{38}$$

is the root of a rational equation of the  $(m - 1)^{\text{th}}$  degree, if

$$a + 2b + 3c + \dots + (m - 1)s = Wm,$$

$W$  being a whole number. For, by (30) in connection with (26),

$$J_2^{\frac{1}{m}} = p_2 J_1^{\frac{2}{m}}, J_3^{\frac{1}{m}} = p_3 J_3^{\frac{1}{m}}, \text{ and so on. Therefore (38) has}$$

the value

$$(p_2 p_3 \dots) J_1^{\frac{a + 2b + 3c + \dots + (m-1)s}{m}}, \text{ or } (p_2 p_3 \dots) J_1^{\frac{W}{m}}.$$

This is a rational function of  $t_1$ , and therefore the root of a rational equation of the  $(m - 1)^{\text{th}}$  degree.

aining the pr  
of the reduc  
 $x^n - 1 = 0$   
etc. Therefor  
-1). (35  
 $t_1$ . Therefor  
 $m-1$   
1  
 $2(m-1)$   
.....  
 $k_m t_1$ ; } (36  
f equations (26

minate functio  
quantities. Fo  
of  $C_1, C_2, \text{ etc}$   
 $y_{m-1}$ ;  
But, by §13, th  
to one anothe  
etc., are know  
Therefore, fro

in Prop. X fall  
1 expressions i

$J_1^{\frac{1}{m}}$  )  
 $J_2^{\frac{1}{m}}$  )  
 $J_3^{\frac{1}{m}}$  )  
( $m - 1$ )<sup>th</sup> degr-

THE EQUATION  $F(x) = 0$  OF THE SECOND CLASS.

§46. We now suppose that the simplified root  $r_1$  of the rational irreducible equation  $F(x) = 0$  of the  $m^{\text{th}}$  degree,  $m$  prime, involves when modified according to §21, a principal surd not a root of unity. It must not be forgotten that, when we thus speak of roots of unity, we mean, according to §1, roots which have prime numbers for the denominators of their indices. In this case conclusions can be established similar to those reached in the case that has been considered. The root  $r_1$  is still of the form (29). The equation  $F(x) = 0$  has still an auxiliary of the  $(m - 1)^{\text{th}}$  degree, whose roots are the  $m^{\text{th}}$  powers of the expressions

$$\frac{1}{\Delta_1^m}, a_1 \Delta_1^{\frac{2}{m}}, b_1 \Delta_1^{\frac{3}{m}}, \dots, e_1 \Delta_1^{\frac{m-2}{m}}, h_1 \Delta_1^{\frac{m-1}{m}}, \quad (39)$$

though the auxiliary here is not necessarily irreducible. Also, substituting the expressions in (39) for  $\Delta_1^{\frac{1}{m}}, \Delta_2^{\frac{1}{m}}$ , etc., in (37), the law of Proposition XIII. still holds, together with corollary in §45.

§47. By Cor. Prop. VI., the denominator of the index of a surd of the highest rank in  $r_1$  is  $m$ . Let  $\Delta_1^{\frac{1}{m}}$  be such a surd. By §21, the coefficients of the different powers of  $\Delta_1^{\frac{1}{m}}$  in  $r_1$  cannot be all zero. We may take the coefficient of the first power to be distinct from zero and to be  $\frac{1}{m}$  for, if it were  $\frac{k_1}{m}$ , we might substitute  $s^{\frac{1}{m}}$  for  $k_1 \Delta_1^{\frac{1}{m}}$  and so eliminate  $\Delta_1^{\frac{1}{m}}$  from  $r_1$ , introducing in its room the new surd  $s^{\frac{1}{m}}$  with  $\frac{1}{m}$  for the coefficient of its first power. We may then put

$$r_1 = \frac{1}{m} (g + \Delta_1^{\frac{1}{m}} + a_1 \Delta_1^{\frac{2}{m}} + \dots + e_1 \Delta_1^{\frac{m-2}{m}} + h_1 \Delta_1^{\frac{m-1}{m}}); \quad (40)$$

where  $g, a_1$ , etc., are clear of  $\Delta_1^{\frac{1}{m}}$ . When  $\Delta_1^{\frac{1}{m}}$  is changed successively into  $\Delta_1^{\frac{1}{m}}, t_1^{-1} \Delta_1^{\frac{1}{m}}, t_1^{-2} \Delta_1^{\frac{1}{m}}$ , etc., let

$$r_1, r_2, \dots, r_m, \quad (41)$$

be respecti  
By Prop.  
 $F(x) = 0$   
 $R$ , let  $\Delta_1^{\frac{1}{m}}$   
ing from  
different m  
By Prop. I  
and, by §2  
some order  
(41) is  $g$ .  
§48. PR  
only princi  
Suppose  
from  $\Delta_1^{\frac{1}{m}}$   
in view th  
denominat  
numbers.)  
of  $e_1$ , let  
By Prop.  
putting  $t_n$   
Therefore,  
 $\Delta_1^{\frac{1}{m}}$

CLASS.  
of the rational  
prime, involves  
a root of unity  
roots of unity  
numbers for the  
ons can be estab  
been considered  
F(x) = 0 has  
roots are the m<sup>th</sup>

be respectively what r<sub>1</sub> becomes, t<sub>1</sub> being a primitive m<sup>th</sup> root of unity.  
By Prop. VI., the terms in (41) are the roots of the equation  
F(x) = 0. Taking r<sub>n</sub>, any one of the particular cognate forms of  
R, let J<sub>n</sub><sup>1/m</sup>, a<sub>n</sub>, etc., be respectively what J<sub>1</sub><sup>1/m</sup>, a<sub>1</sub>, etc., become in pass-  
ing from r<sub>1</sub> to r<sub>n</sub>; and when J<sub>n</sub><sup>1/m</sup> is changed successively into the  
different m<sup>th</sup> roots of the determinate base J<sub>n</sub>, let r<sub>n</sub> become

$$r_n, r_n', r_n'', \dots, r_n^{(m-1)}. \tag{42}$$

By Prop. II., the terms in (42) are roots of the equation F(x) = 0 ;  
and, by §23, they are all unequal. Therefore they are identical, in  
some order, with the terms in (41). Also, the sum of the terms in  
(41) is g. Therefore g is rational.

$\frac{m-1}{m}$   
D<sub>1</sub><sup>1/m</sup>, (39)  
ible. Also, sub

in (37), the law  
ary in §45.

§48. PROPOSITION XIV. In r<sub>1</sub>, as expressed in (40), J<sub>1</sub><sup>1/m</sup> is the  
only principal (see §2) surd.

index of a surd of  
rd. By §21, the

Suppose, if possible, that there is in r<sub>1</sub> a principal surd z<sub>1</sub><sup>1/c</sup> distinct  
from J<sub>1</sub><sup>1/m</sup>. And first, let z<sub>1</sub><sup>1/c</sup> be not a root of unity. (It will be kept  
in view that when, in such a case, we speak of roots of unity, the  
denominators of their indices are understood, according §1, to be prime  
numbers.) When z<sub>1</sub><sup>1/c</sup> is changed into z<sub>2</sub><sup>1/c</sup>, one of the other c<sup>th</sup> roots  
of z<sub>1</sub>, let r<sub>1</sub>, a<sub>1</sub>, etc., become respectively r<sub>1</sub>', a<sub>1</sub>', etc. Then

$$mr_1' = g + J_1^{1/m} + a_1 J_1^{2/m} + \text{etc} \tag{43}$$

om the new surd  
We may then put

By Prop. II., r<sub>1</sub> is equal to a term in (41), say to r<sub>n</sub>. And, by §48,  
putting t<sub>n-1</sub> for t<sub>1</sub><sup>1-n</sup>,

$$mr_n = g + t_{n-1} J_1^{1/m} + t_{n-1}^2 a_1 J_1^{2/m} + \text{etc}. \tag{44}$$

- k<sub>1</sub> D<sub>1</sub><sup>1/m</sup>); (40)

changed succes-

Therefore,

$$(41) \quad J_1^{1/m} (1 - t_{n-1}) + J_1^{2/m} (a_1 - a_1 t_{n-1}^2) + \text{etc.} = 0. \tag{45}$$

This equation involves no surds except those found in the simplified expression  $r_1$ , together with the primitive  $m^{\text{th}}$  root of unity. Therefore the expression on the left of (45) is in a simple state. Therefore

by §8, the coefficients of the different powers of  $J_1^{\frac{1}{m}}$  are separately zero. Therefore  $t_{n-1} = 1$ ,  $a_1' = a_1$ ,  $b_1' = b_1$ , and so on. But, as was shown in Prop. V.,  $z_1^{\frac{1}{c}}$  being a principal surd not a root of unity

in the simplified expression  $a_1$ ,  $a_1$  cannot be equal to  $a_1'$  unless  $z_1^{\frac{1}{c}}$  can be eliminated from  $a_1$  without the introduction of any new surd

In like manner  $b_1$  cannot be equal to  $b_1'$  unless  $z_1^{\frac{1}{c}}$  can be eliminated from  $b_1$ . And so on. Therefore, because  $a_1 = a_1'$ , and  $b_1 = b_1'$ ,

and so on,  $z_1^{\frac{1}{c}}$  admits of being eliminated from  $r_2$  without the introduction of any new surd, which, by §21, is impossible. Next, let

$z_1^{\frac{1}{c}}$  be a root (see §1) of unity, which may be otherwise written  $\theta_1$ . Let the different primitive  $c^{\text{th}}$  roots of unity be  $\theta_1, \theta_2$ , etc.; and when  $\theta_1$  is changed successively into  $\theta_1, \theta_2$ , etc., let  $r_1$  become successively  $r_1, r_1'$ , etc. Suppose it possible that the  $c - 1$  terms  $r_1, r_1',$  etc., are all equal. Since  $z_1^{\frac{1}{c}}$  is a principal surd in  $r_1$ , we

may put  $r_1 = h\theta_1^{c-1} + k\theta_1^{c-2} + \dots + l$ ; where  $h, k$ , etc., are clear of  $\theta_1$ . Therefore  $(c - 1) r_1 = cl - (h + k + \text{etc.})$  Thus

$z_1^{\frac{1}{c}}$  may be eliminated from  $r_1$  without the introduction of any new surd; which by §21 is impossible. Since then the terms  $r_1, r_1',$  etc., are not all equal, let  $r_1$  and  $r_1'$  be unequal. Then  $r_1'$  is equal to a term in (41) distinct from  $r_1$ , say to  $r_n$ . Expressing  $nr_1$  and  $nr_n$  as in (43) and (44), we deduce (45); which, as above, is impossible.

§49. PROPOSITION XV. Taking  $r_1, r_n, J_n^{\frac{1}{m}}$ , etc., as in §47, an

equation 
$$t J_n^{\frac{1}{m}} = p J_1^{\frac{c}{m}} \tag{46}$$

can be for  
number le  
note (see §  
By §47,  
it is imma  
(h  
The coeffic  
for the coe  
equation t  
volving onl  
XIV.,  $J_1^{\frac{1}{m}}$   
except  $J_n^{\frac{1}{m}}$   
 $l_1 =$   
where  $d, d$   
where  $d,$   
§8, since  $J$   
 $d' - J_n, c$   
5

can be formed; where  $t$  is an  $m^{\text{th}}$  root of unity, and  $c$  is a whole number less than  $m$  but not zero, and  $p$  involves only surds subordinate (see §3) to  $J_n^{\frac{1}{m}}$  or  $J_n^{\frac{1}{m}}$

By §47, one of the terms in (42) is equal to  $r_1$ . For our argument it is immaterial which be selected. Let  $r_n = r_1$ . Therefore

$$(h_n J_n^{\frac{m-1}{m}} + e_n J_n^{\frac{m-2}{m}} + \dots + J_n^{\frac{1}{m}}) - (h_1 J_1^{\frac{m-1}{m}} + e_1 J_1^{\frac{m-2}{m}} + \dots + J_1^{\frac{1}{m}}) = 0. \quad (47)$$

The coefficients of the different powers of  $J_n^{\frac{1}{m}}$  here are not all zero, for the coefficient of the first power is unity. Therefore by §5, an

equation  $t J_n^{\frac{1}{m}} = l_1$  subsists,  $t$  being an  $m^{\text{th}}$  root of unity, and  $l_1$  involving only surds exclusive of  $J_n^{\frac{1}{m}}$  that occur in (47). By Prop.

XIV.,  $J_1^{\frac{1}{m}}$  is a surd of a higher rank (see §3) than any surd in (47) except  $J_n^{\frac{1}{m}}$ . Therefore we may put

$$l_1 = d + d_1 J_1^{\frac{1}{m}} + d_2 J_1^{\frac{2}{m}} + \dots + d_{m-1} J_1^{\frac{m-1}{m}};$$

where  $d, d_1, \text{etc.}$ , involve only surds lower in rank than  $J_1^{\frac{1}{m}}$ . Then

$$J_n = l_1^m = (d + d_1 J_1^{\frac{1}{m}} + \text{etc.})^m \\ = d' + d'_1 J_1^{\frac{1}{m}} + d'_2 J_1^{\frac{2}{m}} + \text{etc.};$$

where  $d', d'_1, \text{etc.}$ , involve only surds lower in rank than  $J_1^{\frac{1}{m}}$ . By

§8, since  $J_1^{\frac{1}{m}}$  is a surd in the simplified expressions  $r_1$ , the coefficients

(46)  $d', d'_1, \text{etc.}$ , in the equation



$$(d' - J_n) + d'_1 \Delta_1^{\frac{1}{m}} + d'_2 \Delta_1^{\frac{2}{m}} + \text{etc.} = 0 \tag{48}$$

are separately zero. Therefore  $(d + d_1 \Delta_1^{\frac{1}{m}} + \text{etc.})^m = d'$ . And  $t_1$  being a primitive  $m^{\text{th}}$  root of unity,

$$(d + d_1 t_1 \Delta_1^{\frac{1}{m}} + \text{etc.})^m = d' + d' t_1 \Delta_1^{\frac{1}{m}} + \text{etc.} = d'$$

Therefore,

$$(d + d_1 t_1 \Delta_1^{\frac{1}{m}} + \text{etc.}) = t_1^a (d + d_1 J_1^{\frac{1}{m}} + d_2 J_1^{\frac{2}{m}} + \text{etc.}),$$

$t_1^a$  being one of the  $m^{\text{th}}$  roots of unity. In the same way in which

the coefficients of the different powers of  $\Delta_1^{\frac{1}{m}}$  in (48) are separated

zero, each of the expressions  $d(1 - t_1^a)$ ,  $d_1(t_1 - t_1^a)$ , etc., must be

zero. But not more than one of the  $m - 1$  factors,  $t_1 - t_1^a$ ,

$t_1^2 - t_1^a$ , etc., can be zero. Therefore not more than one of the

$m - 1$  terms  $d_1, d_2, \text{etc.}$ , is distinct from zero. Suppose if possible

that all these terms are zero. Then  $t \Delta_n^{\frac{1}{m}} = d$ . Therefore the

different powers of  $\Delta_n^{\frac{1}{m}}$  can be expressed in terms of the surds in-

volved in  $d$  and of the  $m^{\text{th}}$  root of unity. Substitute for  $\Delta_n^{\frac{1}{m}}, \Delta_n^{\frac{2}{m}}$ ,

etc., in (47), their values thus obtained. Then (47) becomes

$$Q - (h_1 \Delta_1^{\frac{m-1}{m}} + \dots + \Delta_1^{\frac{1}{m}}) = 0; \tag{49}$$

where  $Q$  involves no surds, distinct from the primitive  $m^{\text{th}}$  root

of unity, that are not lower in rank than  $\Delta_1^{\frac{1}{m}}$ ; which, because

the coefficient of the first power of  $\Delta_1^{\frac{1}{m}}$  in (49) is not zero, is, by

impossible. Hence there must be one, while at the same time there can

only one of the  $m - 1$  terms,  $d_1, d_2, \text{etc.}$ , distinct from zero. L

$d_1$  be the  
 $1 - t_1^a$  is  
 $t \Delta_n^{\frac{1}{m}} =$   
 §50. C  
 $\Delta_n^{\frac{1}{m}}$  can b  
 $t \Delta_n^{\frac{1}{m}} =$   
 where  $p, q,$   
 etc., are w  
 the numb  
 with mul  
 series 1, 2  
 $m$ . Ther  
 in a certai  
 §51. P  
 forms of  $d$   
 $t \Delta_n^{\frac{1}{m}}$   
 are severa  
 $m^{\text{th}}$  roots  
 By §47  
 it is imm  
 XV., the  
 different  
 By Cor.  
 order, wi

$= 0$  (48)  $d_e$  be the term that is not zero. Then  $t_1^e - t_1^e = 0$ . Therefore  $1 - t_1^e$  is not zero. Therefore  $d = 0$ . Therefore, putting  $p$  for  $d_e$ ,

$t \Delta_n^{\frac{1}{m}} = p \Delta_1^{\frac{e}{m}}$ . And

50. Cor. By the proposition, values of the different powers of  $\Delta_n^{\frac{1}{m}}$  can be obtained as follows :

$$t \Delta_n^{\frac{1}{m}} = p \Delta_1^{\frac{c}{m}}, t^2 \Delta_n^{\frac{2}{m}} = q \Delta_1^{\frac{s}{m}}, t^3 \Delta_n^{\frac{3}{m}} = k \Delta_1^{\frac{z}{m}}, \text{ etc.;} \quad (50)$$

where  $p, q, \text{ etc.}$  involve only surds that occur in  $\Delta_1$  or  $\Delta_n$ ; and  $c, s, z, \text{ etc.}$  are whole numbers in the series  $1, 2, \dots, m - 1$ . No two of the numbers  $c, s, \text{ etc.}$ , can be the same; for they are the products, with multiples of the prime number  $m$  left out, of the terms in the series  $1, 2, \dots, m - 1$ , by the whole number  $c$  which is less than  $m$ . Therefore the series  $c, s, z, \text{ etc.}$ , is the series  $1, 2, \dots, m - 1$ , in a certain order.

51. PROPOSITION XVI. If  $r_n$  be one of the particular cognate forms of  $R$ , the expressions

$$t \Delta_n^{\frac{1}{m}}, t^2 a_n \Delta_n^{\frac{2}{m}}, \dots, t^{m-2} e_n \Delta_n^{\frac{m-2}{m}}, t^{m-1} h_n \Delta_n^{\frac{m-1}{m}}, \quad (51)$$

are severally equal, in some order, to those in (39),  $t$  being one of the  $m^{\text{th}}$  roots of unity.

By §47, one of the terms in (42) is equal to  $r_1$ . For our argument it is immaterial which be chosen. Let  $r_n = r_1$ . By Cor. Prop. XV., the equations (50) subsist. Substitute in (47) the values of the different powers of  $\Delta_n^{\frac{1}{m}}$  so obtained. Then

$$(t^{-1} p \Delta_1^{\frac{c}{m}} + t^{-2} q a_n \Delta_1^{\frac{s}{m}} + \text{etc.}) - (\Delta_1^{\frac{1}{m}} + a_1 \Delta_1^{\frac{2}{m}} + \text{etc}) = 0. \quad (52)$$

By Cor. Prop. XV., the series  $\Delta_1^{\frac{c}{m}}, \Delta_1^{\frac{s}{m}}, \text{ etc.}$ , is identical, in some order, with the series  $\Delta_1^{\frac{1}{m}}, \Delta_1^{\frac{2}{m}}, \text{ etc.}$  Also, by §8, since  $\Delta_1^{\frac{1}{m}}$  is a

surd occurring in the simplified expression  $r_1$ , and since besides  $\Delta_1^{\frac{1}{m}}$  there are in (52) no surds, distinct from the primitive  $m^{\text{th}}$  root of unity, that are not lower in rank than  $\Delta_1^{\frac{1}{m}}$ , if the equation (52) were arranged according to the powers of  $\Delta_1^{\frac{1}{m}}$  lower than the  $m^{\text{th}}$  the coefficients of the different powers of  $\Delta_1^{\frac{1}{m}}$  would be separately zero. Hence  $\Delta_1^{\frac{1}{m}}$  is equal to that one of the expressions,

$$t^{-1} p \Delta_1^{\frac{c}{m}}, t^{-2} q a_n \Delta_1^{\frac{s}{m}}, \text{ etc.} \tag{53}$$

in which  $\Delta_1^{\frac{1}{m}}$  is a factor. In like manner  $a_1 \Delta_1^{\frac{2}{m}}$  is equal to that one of the expressions (53) in which  $\Delta_1^{\frac{2}{m}}$  is a factor. And so on. Therefore the terms  $\Delta_1^{\frac{1}{m}}, a_1 \Delta_1^{\frac{2}{m}}, \text{ etc.}$ , forming the series (39), are severally equal, in some order, to the terms in (53), which are those forming the series (51.)

§52. PROPOSITION XVII. The equation  $F(x) = 0$  has a rational auxiliary (Compare Prop. VII.) equation  $\varphi(x) = 0$ , whose roots are the  $m^{\text{th}}$  powers of the terms in (39).

Let the unequal particular cognate forms of the generic expression  $\Delta$  under which the simplified expression  $\Delta_1$  falls be

$$\Delta_1, \Delta_2, \dots, \Delta_c. \tag{54}$$

By Prop. XVI., there is a value  $t$  of the  $m^{\text{th}}$  root of unity for which the expressions

$$t \Delta_2^{\frac{1}{m}}, t^2 a_2 \Delta_2^{\frac{2}{m}}, \dots, t^{m-2} e_2 \Delta_2^{\frac{m-2}{m}}, t^{m-1} h_2 \Delta_2^{\frac{m-1}{m}} \tag{55}$$

are severally equal, in some order, to those in (39). Therefore  $\Delta_2$  is equal to one of the terms

$$\Delta_1, a_1^m \Delta_1^2, \dots, e_1^m \Delta_1^{m-2}, h_1^m \Delta_1^{m-1}. \tag{56}$$

In like ma  
And, becau  
to differen  
roots of a  
from the se  
the remain  
irreducible  
be the con  
terms in (3  
§53. The  
of the exp  
to the eq  
auxiliaries  
§54. Pr  
becomes  $\Delta$   
only surds  
 $a_n, b_n, e$   
 $\Delta$ , there c  
forms of  $\Delta$   
case of  $b_1$   
For, jus  
there are p  
 $\tau \Delta$   
are equal  
assigning  
equal to  
 $\Delta$   
Suppose  
 $\frac{1}{m}$   
of  $\Delta_1^{\frac{1}{m}}$   
a 2  $m^{\text{th}}$   
than  $\Delta_1^{\frac{1}{m}}$   
rational

In like manner each of the terms in (54) is equal to a term in (56). And, because the terms in (54) are unequal, they are severally equal to different terms in (56). By Prop. III., the terms in (54) are the roots of a rational irreducible equation, say  $\psi_1(x) = 0$ . Rejecting from the series (56) the roots of the equation  $\psi_1(x) = 0$ , certain of the remaining terms must in the same way be the roots of a rational irreducible equation  $\psi_2(x) = 0$ . And so on. Ultimately, if  $\varphi(x)$  be the continued product of the expressions  $\psi_1(x), \psi_2(x), \text{etc.}$ , the terms in (56) are the roots of the rational equation  $\varphi(x) = 0$ .

§53. The equations  $\psi_1(x) = 0, \psi_2(x) = 0, \text{etc.}$ , formed by means of the expressions  $\psi_1(x), \psi_2(x), \text{etc.}$ , may be said to be *sub-auxiliary* to the equation  $F(x) = 0$ . It will be observed that the sub-auxiliaries are all irreducible.

§54. PROPOSITION XVIII. In passing from  $r_1$  to  $r_n$ , while  $\Delta_1$  becomes  $\Delta_n$ , the expressions  $a_1, b_1$ , which, by Prop. XIV., involve only surds occurring in  $\Delta_1$ , must severally receive determinate values,  $a_n, b_n, \text{etc.}$  In other words,  $a_1$  being a particular cognate form of  $A$ , there cannot, for the same value of  $\Delta_n$ , be two particular cognate forms of  $A$ , as  $a_n$  and  $a_N$ , unequal to one another. And so in the case of  $b_1, e_1, \text{etc.}$

For, just as each of the terms in (42) is equal to a term in (41), there are primitive  $m^{\text{th}}$  roots of unity  $\tau$  and  $T$  such that the expressions

$$\tau \Delta_n^{\frac{1}{m}} + \tau^2 a_n \Delta_n^{\frac{2}{m}} + \text{etc.}, T \Delta_N^{\frac{1}{m}} + T^2 a_N \Delta_N^{\frac{1}{m}} + \text{etc.},$$

are equal to one another. Therefore, if  $\Delta_N = \Delta_n$ , in which case, by assigning suitable values to  $\tau$  and  $T$ ,  $\Delta_N^{\frac{1}{m}}$  may be taken to be

equal to  $\Delta_n^{\frac{1}{m}}$ ,

$$\Delta_n^{\frac{1}{m}} (\tau - T) + \Delta_n^{\frac{2}{m}} (a_n \tau^2 - a_N T^2) + \text{etc.} = 0. \quad (57)$$

Suppose if possible that the coefficients of the different powers

of  $\Delta_n^{\frac{1}{m}}$  in (57) are not all zero. Then, by §5,  $t \Delta_n^{\frac{1}{m}} = l_1$ ;  $t$  being an  $m^{\text{th}}$  root of unity; and  $l_1$  involving only surds of lower ranks

than  $\Delta_n^{\frac{1}{m}}$ . Hence, by Prop. XV. and Cor. Prop. XV.,  $\Delta_n^{\frac{1}{m}}$  is a

rational function of surds of lower ranks than  $\Delta_n^{\frac{1}{m}}$  and of the

primitive  $m^{\text{th}}$  root of unity; which, by the definition in §6, is impossible. Since then the coefficients of the different powers of  $\Delta_n^{\frac{1}{m}}$  in (57) are separately zero,  $\tau = T, a_n \tau^2 = a_N T^2$ , therefore  $a_n = a_N$ .

§55. PROPOSITION XIX. Let the terms in (39) be written respectively

$$\Delta_1^{\frac{1}{m}}, \delta_2^{\frac{1}{m}}, \delta_3^{\frac{1}{m}}, \dots, \delta_{m-1}^{\frac{1}{m}}. \tag{58}$$

The symbols  $\Delta_1, \delta_2, \delta_3$ , etc., are employed instead of  $\Delta_1, \Delta_2, \Delta_3$ , etc. because this latter notation might suggest, what is not necessarily true, that the terms in (56) are all of them particular cognate forms of the generic expression under which  $\Delta_1$  falls. Then (compare Prop. XIII.) the  $m - 1$  expressions in each of the groups

$$\left. \begin{aligned} & \left( \Delta_1^{\frac{1}{m}} \delta_{m-1}^{\frac{1}{m}}, \delta_2^{\frac{1}{m}} \delta_{m-2}^{\frac{1}{m}}, \delta_3^{\frac{1}{m}} \delta_{m-3}^{\frac{1}{m}}, \dots, \delta_{m-1}^{\frac{1}{m}} \Delta_1^{\frac{1}{m}} \right), \\ & \left( \Delta_1^{\frac{2}{m}} \delta_{m-2}^{\frac{1}{m}}, \delta_2^{\frac{2}{m}} \delta_{m-4}^{\frac{1}{m}}, \delta_3^{\frac{2}{m}} \delta_{m-6}^{\frac{1}{m}}, \dots, \delta_{m-1}^{\frac{2}{m}} \delta_2^{\frac{1}{m}} \right), \\ & \left( \Delta_1^{\frac{3}{m}} \delta_{m-3}^{\frac{1}{m}}, \delta_2^{\frac{3}{m}} \delta_{m-6}^{\frac{1}{m}}, \delta_3^{\frac{3}{m}} \delta_{m-9}^{\frac{1}{m}}, \dots, \delta_{m-1}^{\frac{3}{m}} \delta_3^{\frac{1}{m}} \right) \end{aligned} \right\} \tag{59}$$

and so on, are the roots of a rational equation of the  $(m - 1)^{\text{th}}$  degree. Also (compare Prop. X.) the first  $\frac{m-1}{2}$  terms in the first of the groups (59) are the roots of a rational equation of the  $\left(\frac{m-1}{2}\right)^{\text{th}}$  degree.

In the enunciation of the proposition the remark is made that the series (54) is not necessarily identical with the series

$$\Delta_1, \delta_2, \delta_3, \dots, \delta_{m-1}.$$

The former consists of the unequal particular cognate forms of  $\Delta$ ; the latter consists of the roots of the auxiliary equation  $\varphi(x) = 0$ . These two series are identical only when the auxiliary is irreducible. To prove the first part of the proposition, take the terms forming the second of the groups (59). Because  $\delta_{m-2}^{\frac{1}{m}}$  represents  $e_1 \Delta_1^{\frac{m-2}{m}}$ ,

Let  $E$  be  $e_1$  falls. In the term  $e_1, e_2, \dots$  determinat

There is t equal to a root of uni some order

is equal be as in P $^m - 2 e_2 \Delta_1^m$  and  $M - q_1^2 k_1 \Delta_1^m$  respective

In other second of equal to unequal t

Then, by irreducib distinct t be shown rational mately,  $f_2(x)$ , et roots of applies second p groups ( with the

$$e_1 J_1 = J_1^{\frac{2}{m}} \delta_{m-1}^{\frac{1}{m}}$$

Let  $E$  be the generic symbol under which the simplified expression  $e_1$  falls. By Prop. XVIII., when  $J_1$  is changed successively into the  $c$  terms in (54),  $e_1$  receives successively the determinate values  $e_1, e_2, \dots, e_c$ ; and therefore  $e_1 J_1$  receives successively the determinate values

$$e_1 J_1, e_2 J_2, \dots, e_c J_c. \tag{60}$$

There is therefore no particular cognate form of  $EJ$  that is not equal to a term in (60). By Prop. XVI. there is a value of the  $m^{\text{th}}$  root of unity  $t$  for which the terms in (55) are severally equal, in some order, to those in (39). Let the term in (39) to which  $t J_2^{\frac{1}{m}}$

is equal be  $q_1 J_1^{\frac{n}{m}}$ . Then, applying the principle of Cor. Prop. XV., as in Prop. XVI., it follows that the term in (39) to which  $e^{m-2} e_2 J_2^{\frac{m}{m-2}}$  in (55) is equal is  $k_1 J_1^{\frac{M-2n}{m}}$ ,  $M$  being a multiple of  $m$ , and  $M - 2n$  being less than  $m$ . Therefore  $e_2 J_2$  is equal to  $q_1^2 k_1 J_1^{\frac{M}{m}}$ , which is the product of two of the terms in (39) occurring respectively at equal distances from opposite extremities of the series.

In other words,  $e_2 J_2$  is equal to an expression  $\delta_{\frac{2}{m}}^{\frac{1}{m}} \delta_{m-2n}^{\frac{1}{m}}$  in the second of the groups (59). In like manner every term in (60) is equal to an expression in the second of the groups (59). Let the unequal terms in (60) be

$$e_1 J_1, \text{ etc.} \tag{61}$$

Then, by Prop. III., the terms in (61) are the roots of a rational irreducible equation, say  $f_1(x) = 0$ . Rejecting these, which are distinct terms in the second of the groups (59), it can in like manner be shown that certain other terms in that group are the roots of a rational irreducible equation, say  $f_2(x) = 0$ . And so on. Ultimately, if  $f(x)$  be the continued product of the expressions  $f_1(x), f_2(x)$ , etc., the terms forming the second of the groups (59) are the roots of a rational equation of the  $(m-1)^{\text{th}}$  degree. The proof applies substantially to each of the other groups. To prove the second part, it is only necessary to observe that, in the first of the groups (59), the last term is identical with the first, the last but one with the second, and so on.

§56. *Cor. 1.* The reasoning in the proposition proceeds on the assumption that the prime number  $m$  is odd. Should  $m$  be even, the series  $\Delta_1, \delta_1, \text{etc.}$ , is reduced to its first term. The law may be considered even then to hold in the following form. The product  $\frac{1}{\Delta_1^m} \frac{1}{\Delta_1}$  is the root of a rational equation of the  $(m - 1)^{\text{th}}$  degree or is rational. For this product is  $\Delta_1$ , which, by Prop. XVII., is the root of an equation of the  $(m - 1)^{\text{th}}$  degree.

§56. *Cor. 2.* I merely notice, without farther proof, that the generalization in §45 in the case when the equation  $F(x) = 0$  is of the first (see §30) class holds in the present case likewise.

ANALYSIS OF SOLVABLE EQUATIONS OF THE FIFTH DEGREE.

§58. Let the solvable irreducible equation of the  $m^{\text{th}}$  degree, which we have been considering, be of the fifth degree. Then, by Prop. IX. and §47, whether the equation belongs to the first or to the second of the two classes that have been distinguished, assuming the sum of the roots  $g$  to be zero,

$$r_1 = \frac{1}{5} (\Delta_1^{\frac{1}{5}} + a_1 \Delta_1^{\frac{2}{5}} + e_1 \Delta_1^{\frac{3}{5}} + h_1 \Delta_1^{\frac{4}{5}}), \tag{6}$$

though, when the equation is of the first class, the root, as then presented, is not in a simple state.

§59. PROPOSITION XX. If the auxiliary biquadratic has a rational root  $\Delta_1$  not zero, all the roots of the auxiliary biquadratic are rational.

Because  $\Delta_1$  is rational, the auxiliary biquadratic  $\varphi(x) = 0$  is irreducible. Therefore, by Prop. VII., the equation  $F(x) = 0$  is of the second (see §30) class. Therefore, by Prop. XIV.,  $\Delta_1^{\frac{1}{5}}$  is the principal surd in  $r_1$ . Consequently, because  $\Delta_1$  is rational,  $a_1, e_1$  and  $h_1$  are rational. Therefore  $\Delta_1, a_1^5 \Delta_1^2, e_1^5 \Delta_1^3, h_1^5 \Delta_1^4$  which are the roots of the auxiliary biquadratic, are rational.

§60. PROPOSITION XXI. If the auxiliary biquadratic has a quadratic sub-auxiliary  $\psi_1(x) = 0$  with the roots  $\Delta_1$  and  $\Delta_2$ , the  $\Delta_2 = h_1^5 \Delta_1^4$ , and  $\Delta_1 = h_2^5 \Delta_2^4$ ; and  $h_1 \Delta_1$  is rational.

As in §52,  $t$  being a certain fifth root of unity, each term in (55) is equal to a term in (39). The first term in (55) cannot be equal to the first in (39), for this would make  $\Delta_2 = \Delta_1$ . Suppose if possible that the first in (55) is equal to the second in (39). Then, equations (50), applied as in Prop. XVI.,

Now  $a_1^5$  .  
And  $a_1^5 \Delta_1^2$  .  
state. T  
 $\psi_1(x) =$   
roots of t  
are roots  
distinct fr  
zero, and t  
From this  
another ;  
have  $e_1 t$   
§8, is imp  
roots ; wh  
Therefore  
In the sam  
Therefore  
(39) is eqv  
 $\Delta_1^{\frac{1}{5}} = t^i$   
that, the  
 $\Delta_1, \Delta_2,$  c  
term in t  
 $h_1$  be a pa  
form of //  
Hence, si  
different i  
rational.

$$\left. \begin{aligned}
 t J_2^{\frac{1}{5}} &= a_1 J_1^{\frac{2}{5}}, & t^2 a_2 J_2^{\frac{2}{5}} &= h_1 J_1^{\frac{4}{5}}, \\
 t^3 e_2 J_2^{\frac{3}{5}} &= J_1^{\frac{1}{5}}, & t^4 h_2 J_2^{\frac{4}{5}} &= e_1 J_1^{\frac{3}{5}}, \\
 \text{therefore } J_2 &= a_1^5 J_1^2, & a_2^5 J_2^2 &= h_1^5 J_1^4, \\
 e_2^5 J_2^3 &= J_1, & h_2^5 J_2^4 &= e_1^5 J_1^3.
 \end{aligned} \right\} (63)$$

Now  $a_1^5 J_1^2$ , being equal to  $J_2$ , is a root of the equation  $\psi_1(x) = 0$ .

And  $a_1^5 J_1^2$ , involving only surds that occur in  $r_1$ , is in a simple

state. Therefore, by Prop. III.,  $a_2^5 J_2^2$  is a root of the equation

$\psi_1(x) = 0$ . Therefore  $h_1^5 J_1^4$ , and therefore also  $h_2^5 J_2^4$  or  $e_1^5 J_1^3$ , are roots of that equation. Hence all the terms

$$J_1, a_1^5 J_1^2, e_1^5 J_1^3, h_1^5 J_1^4, \tag{64}$$

are roots of the equation  $\psi_1(x) = 0$ . But  $a_1, e_1, h_1$ , are all distinct from zero; for, by (63), if one of them was zero, all would be

zero, and therefore  $J_1^{\frac{1}{5}}$  would be zero; which by §6, is impossible.

From this it follows that no two terms in (64) are equal to one another; for taking  $a_1^5 J_1^2$  and  $e_1^5 J_1^3$ , if these were equal, we should

have  $e_1 t J_1^{\frac{1}{5}} = a_1$ ,  $t$  being a fifth root of unity; which; which by §8, is impossible. This gives the equation  $\psi_1(x) = 0$  four unequal

roots; which, because it is of the second degree, is impossible.

Therefore the first term in (55) is not equal to the second in (39). In the same way it can be shown that it is not equal to the third. Therefore it must be equal to the fourth. In like manner the first in

(39) is equal to the fourth in (55). Because then  $t J_2^{\frac{1}{5}} = h_1 J_1^{\frac{4}{5}}$ , and

$J_1^{\frac{1}{5}} = t^4 h_2 J_2^{\frac{4}{5}}$ ,  $h_2 J_2 = h_1 J_1$ . But, just as it was proved in §56

that, the roots of the sub-auxiliary  $\psi_1(x) = 0$  being the  $c$  terms  $A_1, A_2$ , etc., there is no particular cognate form of  $EJ$  that is not a

term in the series  $e_1 J_1, e_2 J_2, \dots, e_c J_c$ , it follows that, if  $h_1$  be a particular cognate form of  $H$ , there is no particular cognate

form of  $HJ$  that is not equal to one of the terms  $h_1 J_1$  and  $h_2 J_2$ . Hence, since  $h_1 J_1 = h_2 J_2$ ,  $HJ$  has no particular cognate form

different in value from  $h_1 J_1$ . Therefore, by Prop. III.,  $h_1 J_1$  is rational.



§61. PROPOSITION XXII. The auxiliary biquadratic  $\varphi(x) = 0$  either has all its roots rational, or has a sub-auxiliary (see §53) of the second degree, or is irreducible.

It will be kept in view that the sub-auxiliaries are, by the manner of their formation, irreducible. First, let the series (54), containing the roots of the sub-auxiliary  $\psi_1(x) = 0$  consist of a single term  $J_1$ . Then, by Prop. III.,  $J_1$  is rational. Therefore, by Prop. XX., all the roots of the auxiliary are rational. Next, let the series (54) consist of the two terms  $J_1$  and  $J_2$ . By this very hypothesis, the auxiliary biquadratic has a quadratic sub-auxiliary. Lastly, let the series (54) contain more than two terms. Then it has the three terms  $J_1, J_2, J_3$ . We have shown that these must be severally equal to terms in (64). Neither  $J_2$  nor  $J_3$  is equal to  $J_1$ . They cannot both be equal to  $h_1^5 J_1^1$ . Therefore one of them is equal to one of the terms  $a_1^5 J_1^2, e_1^5 J_1^3$ . But in §60 it appeared that, if  $J_2$  be equal either to  $a_1^5 J_1^2$  or to  $e_1^5 J_1^3$ , all the terms in (64) are roots of the irreducible equation of which  $J_1$  is a root. The same thing holds regarding  $J_3$ . Therefore, when the series (54) contains more than two terms, the irreducible equation which has  $J_1$  for one of its roots has the four unequal terms in (64) for roots; that is to say, the auxiliary biquadratic is irreducible.

§62. Let  $5u_1 = J_1^{\frac{1}{5}}, 5u_2 = a_1 J_1^{\frac{2}{5}}, 5u_3 = e_1 J_1^{\frac{3}{5}}, 5u_4 = h_1 J_1^{\frac{4}{5}}$ ; and,  $n$  being any whole number, let  $S_n$  denote the sum of the  $n^{\text{th}}$  powers of the roots of the equation  $F(x) = 0$ . Then

$$S_1 = 0; S_2 = 10(u_1 u_4 + u_2 u_3); S_3 = 15 \left\{ \sum (u_1 u_2^2) \right\};$$

$$S_4 = 20 \left\{ \sum (u_1^3 u_2) \right\} + 30(u_1^2 u_4^2 + u_2^2 u_3^2) - 120 u_1 u_2 u_3 u_4;$$

$$S_5 = 5 \left\{ \sum (u_1^5) \right\} + 100 \left\{ \sum (u_1^3 u_3 u_4) \right\} + 150 \left\{ \sum (u_1 u_3^2 u_4^2) \right\};$$

where such an expression as  $\sum (u_1 u_2^2)$  means the sum of all such terms as  $u_1 u_2^2$ ; it being understood that, as any one term in the circle  $u_1, u_2, u_4, u_3$ , passes into the next, that next passes into its next,  $u_3$  passing into  $u_1$ .

THE ROOTS OF THE AUXILIARY BIQUADRATIC ALL RATIONAL.

§63. Any rational values that may be assigned to  $J_1, a_1, e_1$ , and  $h_1$  in  $r_1$ , taken as in (62), make  $r_1$  the root of a rational equation of the fifth degree, for  $r_1$  by render the values of  $S_1, S_2$ , etc., in §62, rational. In fact,  $S_1 = 0, 25 S_2 = 10 J_1 (h_1 + a_1 e_1)$ , and so on.

THE A

§64. may be degree, must b

where  $\psi_1(x) = a_2 = b$

roots J

By E class, th we are sub-aux p and k  $\psi_1(x) = q$  for h Therefo enuncia

a fifth u unity, v

$J_2^{\frac{1}{5}}, J$  what a

of  $e_1 J$

the sam

be mor princip  $a_1$  inv  $\sqrt{(p^2}$   $a_1 = l$   $a_1$  becc in the

§65. in  $r_1,$

## THE AUXILIARY BIQUADRATIC WITH A QUADRATIC SUB-AUXILIARY.

§64. PROPOSITION XXIII. In order that  $r_1$ , taken as in (62), may be the root of an irreducible equation  $F(x) = 0$  of the fifth degree, whose auxiliary biquadratic has a quadratic sub-auxiliary, it must be of the form

$$r_1 = \frac{1}{5} \{ (\mathcal{J}_1^{\frac{1}{5}} + \mathcal{J}_2^{\frac{1}{5}}) + (a_1 \mathcal{J}_1^{\frac{2}{5}} + a_2 \mathcal{J}_2^{\frac{2}{5}}) \}; \quad (65)$$

where  $\mathcal{J}_1$  and  $\mathcal{J}_2$  are the roots of the irreducible equation  $\zeta_1(x) = x^2 - 2px + q^5 = 0$ ; and  $a_1 = b + d\sqrt{(p^2 - q^5)}$ ,  $a_2 = b - d\sqrt{(p^2 - q^5)}$ ;  $p$ ,  $b$  and  $d$  being rational; and the roots  $\mathcal{J}_1^{\frac{1}{5}}$  and  $\mathcal{J}_2^{\frac{1}{5}}$  being so related that  $\mathcal{J}_1^{\frac{1}{5}} \mathcal{J}_2^{\frac{1}{5}} = q$ .

By Prop. VII., when a quintic equation is of the first (see §30) class, the auxiliary biquadratic is irreducible. Hence, in the case we are considering, the quintic is of the second class. The quadratic sub-auxiliary may be assumed to be  $\zeta_1(x) = x^2 - 2px + k = 0$ ,  $p$  and  $k$  being rational. By Prop. XXI., the roots of the equation  $\zeta_1(x) = 0$  are  $\mathcal{J}_1$  and  $h_1^5 \mathcal{J}_1^4$ . Therefore  $k = (h_1 \mathcal{J}_1)^5$ ; or, putting  $q$  for  $h_1 \mathcal{J}_1$ ,  $k = q^5$ . By the same proposition,  $h_1 \mathcal{J}_1$  is rational. Therefore  $q$  is rational. Hence  $\zeta_1(x)$  has the form specified in the enunciation of the proposition. Next, by Proposition XVI., there is

a fifth root of unity  $t$  such that  $t \mathcal{J}_2^{\frac{1}{5}} = h_1 \mathcal{J}_1^{\frac{4}{5}}$ . If we take  $t$  to be unity, which we may do by a suitable interpretation of the symbol

$\mathcal{J}_2^{\frac{1}{5}}$ ,  $\mathcal{J}_2^{\frac{1}{5}} = h_1 \mathcal{J}_1^{\frac{4}{5}}$ . This implies that  $e_1 \mathcal{J}_1^{\frac{3}{5}} = a_2 \mathcal{J}_2^{\frac{2}{5}}$ ,  $a_2$  being what  $a_1$  becomes in passing from  $\mathcal{J}_1$  to  $\mathcal{J}_2$ . Substituting these values

of  $e_1 \mathcal{J}_1^{\frac{3}{5}}$  and  $h_1 \mathcal{J}_1^{\frac{4}{5}}$  in (62), we obtain the form of  $r_1$  in (65), while at the same time  $\mathcal{J}_1^{\frac{1}{5}} \mathcal{J}_2^{\frac{1}{5}} = h_1 \mathcal{J}_1 = q$ . The forms of  $a_1$  and  $a_2$  have to

be more accurately determined. By Prop. XIV.,  $\mathcal{J}_1^{\frac{1}{5}}$  is the only principal surd that  $r_1$ , as presented in (62), contains. Therefore  $a_1$  involves no surd that does not occur in  $\mathcal{J}_1$ ; that is to say,  $\sqrt{(p^2 - q^5)}$  is the only surd in  $a_1$ . Hence we may put  $a_1 = b + d\sqrt{(p^2 - q^5)}$ ;  $b$  and  $d$  being rational. But  $a_2$  is what  $a_1$  becomes in passing from  $\mathcal{J}_1$  to  $\mathcal{J}_2$ . And  $\mathcal{J}_2$  differs from  $\mathcal{J}_1$  only in the sign of the root  $\sqrt{(p^2 - q^5)}$ . Therefore

$$a_2 = b - d\sqrt{(p^2 - q^5)}.$$

§65. Any rational values that may be assigned to  $b$ ,  $d$ ,  $p$  and  $q$  in  $r_1$ , taken as in (65), make  $r_1$  the root of a rational equation of the

fifth degree; for they render the values of  $S_1, S_2$ , etc., in §62, rational. In fact,  $S_1 = 0, 25 S_2 = 10 \{q + q^2 b^2 - q^2 d^2 (p^2 - q^5)\}$ , and so on.

THE AUXILIARY BIQUADRATIC IRREDUCIBLE.

§66. When the auxiliary biquadratic is irreducible, the unequal particular cognate forms of  $J$  are, by Prop. III., four in number,  $J_1, J_2, J_3, J_4$ . As explained in §55, because the equation  $\varphi(x) = 0$  is irreducible, these terms are severally identical with  $J_1, \delta_2, \delta_3, \delta_4$ . Hence, putting  $m = \delta$ , the first two terms in the first of the groups (59) may be written in the notation of (37),

$$J_1^{\frac{1}{5}} J_4^{\frac{1}{5}}, J_2^{\frac{1}{5}} J_3^{\frac{1}{5}}; \tag{66}$$

and the second and third groups may be written

$$\left. \begin{aligned} (J_1^{\frac{2}{5}} J_3^{\frac{1}{5}}, J_2^{\frac{2}{5}} J_1^{\frac{1}{5}}, J_3^{\frac{2}{5}} J_4^{\frac{1}{5}}, J_4^{\frac{2}{5}} J_2^{\frac{1}{5}}) \\ (J_1^{\frac{3}{5}} J_2^{\frac{1}{5}}, J_2^{\frac{3}{5}} J_4^{\frac{1}{5}}, J_3^{\frac{3}{5}} J_1^{\frac{1}{5}}, J_4^{\frac{3}{5}} J_3^{\frac{1}{5}}) \end{aligned} \right\} \tag{67}$$

§67. PROPOSITION XXIV. The roots of the auxiliary biquadratic equation  $\varphi(x) = 0$  are of the forms

$$\left. \begin{aligned} J_1 = m + n \sqrt{z} + \sqrt{s}, J_2 = m - n \sqrt{z} + \sqrt{s_1}, \\ J_4 = m + n \sqrt{z} - \sqrt{s}, J_3 = m - n \sqrt{z} - \sqrt{s_1}; \end{aligned} \right\} \tag{68}$$

where  $s = p + q \sqrt{z}$ , and  $s_1 = p - q \sqrt{z}$ ;  $m, n, z, p$  and  $q$  being rational; and the surd  $\sqrt{s}$  being irreducible.

By Propositions XIII. and XIX., the terms in (66) are the roots of a quadratic. Therefore  $J_1 J_4$  and  $J_2 J_3$  are the roots of a quadratic. Suppose if possible that  $J_1 J_3$  is the root of a quadratic. By

Propositions IX. and XIX.,  $J_3^{\frac{1}{5}} = e_1 J_1^{\frac{3}{5}}$ . Therefore  $e_1^5 J_1^4$  is the root of a quadratic. From this it follows (Prop. III.) that there are not more than two unequal terms in the series,

$$e_1^5 J_1^4, e_2^5 J_2^4, e_3^5 J_3^4, e_4^5 J_4^4. \tag{69}$$

But suppose if possible that  $e_1^5 J_1^4 = e_2^5 J_2^4$ . Then,  $t$  being one of the fifth roots of unity,  $t e_1 J_1^{\frac{4}{5}} = e_2 J_2^{\frac{4}{5}}$ . But, by Propositions IX. and XIX.,  $J_2^{\frac{1}{5}} = h_1 J_1^{\frac{4}{5}}$ . Therefore,  $t e_1 J_1^{\frac{4}{5}} = e_2 h_1^4 J_1^3 J_1^{\frac{1}{5}}$ . There-

fore, the biquadratic equation is irreducible. The roots are those

§68. pressed By P (67) are powers

are the  $J_4$  in (6) follows:

$$\begin{aligned} J_1^2 J_3 &= \\ + \\ J_2^2 J_1 &= \\ - \\ J_4^2 J_2 &= \\ + \\ J_3^2 J_4 &= \\ - \end{aligned}$$

where  $h$  four exp roots of rational. in terms = 0.

$$(J_1^2 J_2$$

fore, by §8,  $e_1 = 0$ . Therefore one of the roots of the auxiliary biquadratic is zero; which because the auxiliary biquadratic is assumed to be irreducible, is impossible. Therefore  $e_1^5 J_1^4$  and  $e_2^5 J_2^4$  are unequal. In the same way all the terms in (69) can be shown to be unequal; which, because it has been proved that there are not more than two unequal terms in (69), is impossible. Therefore  $J_1 J_3$  is not the root of a quadratic equation. Therefore the product of two of the roots,  $J_1$  and  $J_4$ , of the auxiliary biquadratic is the root of a quadratic equation, while the product of a different pair,  $J_1$  and  $J_3$ , is not the root of a quadratic. But the only forms which the roots of an irreducible biquadratic can assume consistently with these conditions are those given in (68).

§68. PROPOSITION XXV. The surd  $\sqrt{s_1}$  can have its value expressed in terms of  $\sqrt{s}$  and  $\sqrt{z}$ .

By Propositions XIII. and XIX, the terms of the first of the groups (67) are the roots of a biquadratic equation. Therefore their fifth powers

$$J_1^2 J_3, J_2^2 J_1, J_3^2 J_4, J_4^2 J_2, \tag{70}$$

are the roots of a biquadratic. From the values of  $J_1, J_2, J_3$  and  $J_4$  in (68), the values of the terms in (70) may be expressed as follows:

$$\left. \begin{aligned} J_1^2 J_3 &= F + F_1 \sqrt{z} + (F_2 + F_3 \sqrt{z}) \sqrt{s} \\ &\quad + (F_4 + F_5 \sqrt{z}) \sqrt{s_1} + (F_6 + F_7 \sqrt{z}) \sqrt{s} \sqrt{s_1}, \\ J_2^2 J_1 &= F - F_1 \sqrt{z} + (F_2 - F_3 \sqrt{z}) \sqrt{s} \\ &\quad - (F_4 - F_5 \sqrt{z}) \sqrt{s} - (F_6 - F_7 \sqrt{z}) \sqrt{s} \sqrt{s_1}, \\ J_3^2 J_4 &= F - F_1 \sqrt{z} - (F_2 - F_3 \sqrt{z}) \sqrt{s} \\ &\quad + (F_4 - F_5 \sqrt{z}) \sqrt{s} - (F_6 - F_7 \sqrt{z}) \sqrt{s} \sqrt{s_1}, \\ J_4^2 J_2 &= F + F_1 \sqrt{z} - (F_2 + F_3 \sqrt{z}) \sqrt{s} \\ &\quad - (F_4 + F_5 \sqrt{z}) \sqrt{s_1} + (F_6 + F_7 \sqrt{z}) \sqrt{s} \sqrt{s_1}, \end{aligned} \right\} \tag{71}$$

where  $F, F_1$ , etc., are rational. Let  $\Sigma(J_1^2 J_3)$  be the sum of the four expressions in (70). Then, because these expressions are the roots of a biquadratic,  $\Sigma(J_1^2 J_3)$  or  $4F + 4F_1 \sqrt{z}$  must be rational. Suppose if possible that  $\sqrt{s_1}$  cannot have its value expressed in terms of  $\sqrt{s}$  and  $\sqrt{z}$ . Then, because  $\sqrt{s} \sqrt{s_1}$  is not rational,  $\Sigma(J_1^2 J_3) = 0$ . By (68), this implies that  $z = 0$ . Let

$$\begin{aligned} (J_1^2 J_3)^2 &= L + L_1 \sqrt{z} + (L_2 + L_3 \sqrt{z}) \sqrt{s} \\ &\quad + (L_4 + L_5 \sqrt{z}) \sqrt{s_1} + (L_6 + L_7 \sqrt{z}) \sqrt{s} \sqrt{s_1}, \end{aligned}$$

where  $L, L_1, \dots$ , are rational. Then, as above,  $L_7 = 0$ . Keeping in view that  $n = 0$ , this means that  $m^2 q = 0$ . But  $q$  is not zero, for this would make  $\sqrt{s} = \sqrt{s_1}$ ; which, because we are reasoning on the hypothesis that  $\sqrt{s_1}$  cannot have its value expressed in terms of  $\sqrt{s}$  and  $\sqrt{z}$ , is impossible. Therefore  $m$  is zero. And it was shown that  $n$  is zero. Therefore  $J_1 = \sqrt{s}$ , and  $J_3 = -\sqrt{s}$ . Therefore  $J_1 J_3 = -\sqrt{(\rho^2 - q^2 z)}$ ; which, because it has been proved that  $J_1 J_3$  is not the root of a quadratic equation, is impossible. Hence  $\sqrt{s_1}$  cannot be a rational function of  $\sqrt{s}$  and  $\sqrt{z}$ .

§69. PROPOSITION XXVI. The form of  $s$  is

$$h(1 + e^2) + h\sqrt{(1 + e^2)}, \tag{72}$$

$h$  and  $e$  being rational, and  $1 + e^2$  being the value of  $z$ .

By Prop. XXV.,  $\sqrt{s_1} = v + c\sqrt{s}$ ,  $v$  and  $c$  being rational functions of  $\sqrt{z}$ . Therefore  $s_1 = v^2 + c^2 s + 2vc\sqrt{s}$ . By Prop. XXIV.,  $\sqrt{s}$  is irreducible. Therefore  $vc = 0$ . But  $c$  is not zero, for this would make  $\sqrt{s_1} = v$ , and thus  $\sqrt{s_1}$  would be the root of a quadratic equation. Therefore  $v = 0$ , and  $\sqrt{s_1} = c\sqrt{s} = (c_1 + c_2\sqrt{z})\sqrt{s}$ ,  $c_1$  and  $c_2$  being rational. Therefore

$$\begin{aligned} \sqrt{(ss_1)} &= \sqrt{(\rho^2 - q^2 z)} = (c_1 + c_2\sqrt{z})(\rho + q\sqrt{z}) \\ &= (c_1\rho + c_2qz) + \sqrt{z}(c_1q + c_2\rho) = P + Q\sqrt{z} \end{aligned}$$

Here, since  $\rho^2 - q^2 z$  is rational, either  $P = 0$  or  $Q = 0$ . As the latter of these alternatives would make  $\sqrt{(\rho^2 - q^2 z)}$  rational, and therefore would make  $\sqrt{(\rho + q\sqrt{z})}$  or  $\sqrt{s}$  reducible, it is inadmissible. Therefore  $c_1\rho + c_2qz = 0$ , and

$$\sqrt{(\rho^2 - q^2 z)} = (c_1q + c_2\rho)\sqrt{z}.$$

Now  $qz$  is not zero, for this would make  $\sqrt{(ss_1)} = \pm p$ ; which, because  $\sqrt{s}$  is irreducible, is impossible. Therefore  $c_2 = 0$ . But, by hypothesis,  $c_1 \neq 0$ ; therefore  $\sqrt{s_1}$ , which is equal to  $(c_1 + c_2\sqrt{z})\sqrt{s}$ , is zero; which is impossible. Hence  $c_1$  cannot be zero. We may therefore put  $ce = 1$ , and  $h(1 + e^2) = \rho$ . Then  $s = \rho + q\sqrt{z} = h(1 + e^2) + h\sqrt{(1 + e^2)}$ . Having obtained this form, we may consider  $z$  to be identical with  $1 + e^2$ ,  $q$  with  $h$ , and  $p$  with  $h(1 + e^2)$ .

§70. The reasoning in the preceding section holds good whether the equation  $F(x) = 0$  be of the first (see §30) or of the second class. If we had had to deal simply with equations of the first class, the proof given would have been unnecessary, so far as the form of  $z$  is concerned; because, in that case, by Prop. VIII.,  $J_1$  is a rational function of the primitive fifth root of unity.

§71  
establis  
Crelle  
For,  
the exp

are the  
to, it is  
of a bi  
corollar  
roots o  
respecti

Then J

Substit

This, v  
only w  
terms  
Theref  
appear  
That i  
it was  
functi

§71. PROPOSITION XXVII. Under the conditions that have been established, the root  $r_1$  takes the form given without deduction in *Crelle* (Vol. V., p. 336) from the papers of Abel.

For, by *Cor. Prop. XIII.* (compare also *Cor. 2, Prop. XIX.*) the expressions

$$\begin{aligned} \sqrt[5]{J_1^5 J_3^2 J_4^3 J_2^3}, \sqrt[5]{J_2^5 J_1^2 J_3^3 J_4^3}, \\ \sqrt[5]{J_3^5 J_4^2 J_2^3 J_1^3}, \sqrt[5]{J_4^5 J_2^2 J_1^3 J_3^3}, \end{aligned} \quad (73)$$

are the roots of a biquadratic equation. In the corollaries referred to, it is merely stated that each of the expressions in (73) is the root of a biquadratic; but the principles of the propositions to which the corollaries are attached show that the four expressions must be the roots of the same biquadratic. Let the terms in (73) be denoted respectively by

$$5A_1^{-1}, \quad 5A_2^{-1}, \quad 5A_3^{-1}, \quad 5A_4^{-1}.$$

Then  $\sqrt[5]{J_1^5 J_3^2 J_4^3 J_2^3} = \sqrt[5]{J_4^5} (\sqrt[5]{J_1^5} \sqrt[5]{J_3^2} \sqrt[5]{J_4^3} \sqrt[5]{J_2^3})$  is an identity. Therefore

$$\frac{1}{5} \sqrt[5]{J_4^5} = A_1 (\sqrt[5]{J_1^5} \sqrt[5]{J_3^2} \sqrt[5]{J_4^3} \sqrt[5]{J_2^3}). \quad \text{Similarly,}$$

$$\frac{1}{5} \sqrt[5]{J_3^5} = A_3 (\sqrt[5]{J_3^5} \sqrt[5]{J_4^2} \sqrt[5]{J_2^3} \sqrt[5]{J_1^3})$$

$$\frac{1}{5} \sqrt[5]{J_2^5} = A_2 (\sqrt[5]{J_2^5} \sqrt[5]{J_1^2} \sqrt[5]{J_3^3} \sqrt[5]{J_4^3}), \text{ and}$$

$$\frac{1}{5} \sqrt[5]{J_1^5} = A_4 (\sqrt[5]{J_4^5} \sqrt[5]{J_2^2} \sqrt[5]{J_1^3} \sqrt[5]{J_3^3}).$$

Substituting these values in (62), we get

$$\begin{aligned} r_1 = A_1 (\sqrt[5]{J_1^5} \sqrt[5]{J_3^2} \sqrt[5]{J_4^3} \sqrt[5]{J_2^3}) + A_2 (\sqrt[5]{J_2^5} \sqrt[5]{J_1^2} \sqrt[5]{J_3^3} \sqrt[5]{J_4^3}) \\ + A_3 (\sqrt[5]{J_3^5} \sqrt[5]{J_4^2} \sqrt[5]{J_2^3} \sqrt[5]{J_1^3}) + A_4 (\sqrt[5]{J_4^5} \sqrt[5]{J_2^2} \sqrt[5]{J_1^3} \sqrt[5]{J_3^3}). \end{aligned} \quad (74)$$

This, with immaterial differences in the subscripts, is Abel's expression; only we need to determine  $A_1, A_2, A_3$  and  $A_4$  more exactly. These terms are the reciprocals of the terms in (73) severally divided by 5. Therefore they are the roots of a biquadratic. Also, no surds can appear in  $A_1$  except those that are present in  $J_1, J_2, J_3$  and  $J_4$ . That is to say,  $A_1$  is a rational function of  $\sqrt{s}, \sqrt{s_1}$  and  $\sqrt{z}$ . But it was shown that  $\sqrt{s_1} \sqrt{s} = h e \sqrt{z}$ . Therefore  $A_1$  is a rational function of  $\sqrt{s}$  and  $\sqrt{z}$ . We may therefore put

$$A_1 = K + K' J_1 + K'' J_4 + K''' J_1 J_4,$$

$K, K', K''$  and  $K'''$  being rational. But the terms  $A_1, A_2, A_3, A_4$  circulate with  $J_1, J_2, J_3, J_4$ . Therefore

$$\begin{aligned} A_2 &= K + K' J_2 + K'' J_3 + K''' J_4, \\ A_4 &= K + K' J_4 + K'' J_1 + K''' J_2, \\ A_3 &= K + K' J_3 + K'' J_2 + K''' J_1, \end{aligned}$$

These are Abel's values.

§72. Keeping in view the values of  $J_1, J_2$ , etc., in (67), and also that  $z = 1 + \epsilon^2$ , and  $s = hz + h\sqrt{z}$ , any rational values that may be assigned to  $m, n, e, h, K, K', K''$  and  $K'''$  make  $r_1$ , as presented in (74), the root of an equation of the fifth degree. For, any rational values of  $m, n$ , etc., make the values of  $S_1, S_2$ , etc., in §62, rational.

§73. It may be noted that, not only is the expression for  $r_1$  in (74) the root of a quintic equation whose auxiliary biquadratic is irreducible, but on the understanding that the surds  $\sqrt{s}$  and  $\sqrt{z}$  in  $J_1$  may be reducible, the expression for  $r_1$  in (74) contains the roots both of all equations of the fifth degree whose auxiliary biquadratics have their roots rational, and of all that have quadratic sub-auxiliaries. It is unnecessary to offer proof of this.

§74. The equation  $x^5 - 10x^3 + 5x^2 + 10x + 1 = 0$  is an example of a solvable quintic with its auxiliary biquadratic irreducible. One of its roots is

$$\omega^{\frac{1}{5}} + \omega\omega^{\frac{2}{5}} + \omega^3\omega^{\frac{3}{5}} + \omega^4\omega^{\frac{4}{5}},$$

$\omega$  being a primitive fifth root of unity. It is obvious that this root satisfies all the conditions that have been pointed out in the preceding analysis as necessary. A root of an equation of the seventh degree of the same character is

$$\omega^{\frac{1}{7}} + \omega^4\omega^{\frac{2}{7}} + \omega^6\omega^{\frac{3}{7}} + \omega^2\omega^{\frac{4}{7}} + \omega^5\omega^{\frac{5}{7}} + \omega^3\omega^{\frac{6}{7}},$$

$\omega$  being a primitive seventh root of unity. The general form under which these instances fall can readily be found. Take the cycle that contains all the primitive  $(m^2)^{\text{th}}$  roots of unity,

$$\theta, \theta^\beta, \theta^{\beta^2}, \text{ etc.} \quad (75)$$

$m$  being prime. The number of terms in the cycle is  $(m-1)^2$ . Let  $\theta_1$  be the  $(m+1)^{\text{th}}$  term in the cycle (75),  $\theta_2$  the  $(2m+1)^{\text{th}}$  term, and so on. Then the root of an equation of the  $m^{\text{th}}$  degree, including the instances above given, is

$$r_1 = (\theta + \theta^{-1}) + (\theta_1 + \theta_1^{-1}) + \dots + (\theta_{m-3} + \theta_{m-3}^{-1}).$$

$A_2, A_1,$

, and also  
that may  
presented  
y rational  
2, rational.

$r_1$  in (74)  
ic is irre-  
nd  $\sqrt{z}$  in  
the roots  
quadratics  
ratic sub-

$= 0$  is an  
ratic irre-

at this root  
e preceding  
nth degree

form under  
e cycle that

(75)

$(m - 1)^2,$   
 $(2m + 1)^{\text{th}}$   
 $m^{\text{th}}$  degree,

$\left(\frac{-1}{\frac{m-3}{2}}\right).$



