

**CIHM
Microfiche
Series
(Monographs)**

**ICMH
Collection de
microfiches
(monographies)**



Canadian Institute for Historical Microreproductions / Institut canadien de microreproductions historiques

© 1997

Technical and Bibliographic Notes / Notes techniques et bibliographiques

The Institute has attempted to obtain the best original copy available for filming. Features of this copy which may be bibliographically unique, which may alter any of the images in the reproduction, or which may significantly change the usual method of filming are checked below.

- Coloured covers / Couverture de couleur
- Covers damaged / Couverture endommagée
- Covers restored and/or laminated / Couverture restaurée et/ou pelliculée
- Cover title missing / Le titre de couverture manque
- Coloured maps / Cartes géographiques en couleur
- Coloured ink (i.e. other than blue or black) / Encre de couleur (i.e. autre que bleue ou noire)
- Coloured plates and/or illustrations / Planches et/ou illustrations en couleur
- Bound with other material / Relié avec d'autres documents
- Only edition available / Seule édition disponible
- Tight binding may cause shadows or distortion along interior margin / La reliure serrée peut causer de l'ombre ou de la distorsion le long de la marge intérieure.
- Blank leaves added during restorations may appear within the text. Whenever possible, these have been omitted from filming / Il se peut que certaines pages blanches ajoutées lors d'une restauration apparaissent dans le texte, mais, lorsque cela était possible, ces pages n'ont pas été filmées.
- Additional comments / Commentaires supplémentaires:

L'Institut a microfilmé le meilleur exemplaire qu'il lui a été possible de se procurer. Les détails de cet exemplaire qui sont peut-être uniques du point de vue bibliographique, qui peuvent modifier une image reproduite, ou qui peuvent exiger une modification dans la méthode normale de filmage sont indiqués ci-dessous.

- Coloured pages / Pages de couleur
- Pages damaged / Pages endommagées
- Pages restored and/or laminated / Pages restaurées et/ou pelliculées
- Pages discoloured, stained or foxed / Pages décolorées, tachetées ou piquées
- Pages detached / Pages détachées
- Showthrough / Transparence
- Quality of print varies / Qualité inégale de l'impression
- Includes supplementary material / Comprend du matériel supplémentaire
- Pages wholly or partially obscured by errata slips, tissues, etc., have been refilmed to ensure the best possible image / Les pages totalement ou partiellement obscurcies par un feuillet d'errata, une pelure, etc., ont été filmées à nouveau de façon à obtenir la meilleure image possible.
- Opposing pages with varying colouration or discolourations are filmed twice to ensure the best possible image / Les pages s'opposant ayant des colorations variables ou des décolorations sont filmées deux fois afin d'obtenir la meilleure image possible.

This item is filmed at the reduction ratio checked below / Ce document est filmé au taux de réduction indiqué ci-dessous.

10x	14x	18x	22x	26x	30x
12x	16x	20x	24x	28x	32x

The copy filmed here has been reproduced thanks to the generosity of:



Legislative Library
Victoria

The images appearing here are the best quality possible considering the condition and legibility of the original copy and in keeping with the filming contract specifications.

Original copies in printed paper covers are filmed beginning with the front cover and ending on the last page with a printed or illustrated impression, or the back cover when appropriate. All other original copies are filmed beginning on the first page with a printed or illustrated impression, and ending on the last page with a printed or illustrated impression.

The last recorded frame on each microfiche shall contain the symbol → (meaning "CONTINUED"), or the symbol ▽ (meaning "END"), whichever applies.

Maps, plates, charts, etc., may be filmed at different reduction ratios. Those too large to be entirely included in one exposure are filmed beginning in the upper left hand corner, left to right and top to bottom, as many frames as required. The following diagrams illustrate the method:



L'exemplaire filmé fut reproduit grâce à la générosité de:

Legislative Library
Victoria

Les images suivantes ont été reproduites avec le plus grand soin, compte tenu de la condition et de la netteté de l'exemplaire filmé, et en conformité avec les conditions du contrat de filmage.

Les exemplaires originaux dont la couverture en papier est imprimée sont filmés en commençant par le premier plat et en terminant soit par la dernière page qui comporte une empreinte d'impression ou d'illustration, soit par le second plat, selon le cas. Tous les autres exemplaires originaux sont filmés en commençant par la première page qui comporte une empreinte d'impression ou d'illustration et en terminant par la dernière page qui comporte une telle empreinte.

Un des symboles suivants apparaîtra sur la dernière image de chaque microfiche, selon le cas: le symbole → signifie "A SUIVRE", le symbole ▽ signifie "FIN".

Les cartes, planches, tableaux, etc., peuvent être filmés à des taux de réduction différents. Lorsque le document est trop grand pour être reproduit en un seul cliché, il est filmé à partir de l'angle supérieur gauche, de gauche à droite, et de haut en bas, en prenant le nombre d'images nécessaire. Les diagrammes suivants illustrent la méthode.

MICROCOPY RESOLUTION TEST CHART

(ANSI and ISO TEST CHART No. 2)



APPLIED IMAGE Inc

100 East Main Street
Rochester, New York 14609 USA
(716) 482-0300 Phone
(716) 288-5989 Fax

GEOLOGICAL SURVEY OF CANADA
ROBERT BELL, M.D., D.Sc., LL.D., F.R.S.

ON TRIONYX FOVEATUS, LEHDY, AND TRIONYX
VAGANS, COPE, FROM THE CRETACEOUS
ROCKS OF ALBERTA

BY

LAWRENCE M. LAMBE, F.G.S., F.R.S.C. (S)

Assistant Palaeontologist

Reprinted from the Summary Report for the year 1901.



OTTAWA
PRINTED BY S. E. DAWSON, PRINTER TO THE KING'S MOST
EXCELLENT MAJESTY
1902





3 3298 00024 2330

ON TRIONYX FOVEATUS, LEIDY, AND TRIONYX
VAGANS, COPE, FROM THE CRETACEOUS
ROCKS OF ALBERTA.

BY LAWRENCE M. LAMBE

(With four plates.)

During the past summer whilst engaged, on behalf of the Geological Survey, in making a collection of vertebrate remains from the Cretaceous of the Red Deer river, Alberta, to supplement the material obtained in two former years, the writer was fortunate enough to secure two almost complete carapaces of turtles, one referable to *Trionyx foveatus*, Leidy, the other to *T. vagans*, Cope. In the Red Deer river district, referred to, the remains of these two species are abundant and are found associated with a number of other species of Chelonia, of which one is *Adovus variolosus*,* Cope (sp.), remarkable alike for its size and the strikingly rugose character of the sculpture of its shell. The rocks holding these remains belong to the Belly river series of the Cretaceous brackish water deposits underlying the Pierre Fox-Hills formation, by which they are separated from the still higher Laramie series.

Trionyx foveatus[†] was originally described in 1860 by Leidy from small fragments of costal and sternal bones from the Judith river beds of Nebraska.

*Proceedings of the Academy of Natural Sciences of Philadelphia, vol. XXVIII, p. 257, 1876 (Cope), and Ottawa Naturalist, vol. XV, p. 63, plates iii, iv, v and vi (Lambe).

†Transactions of the American Philosophical Society, vol. XI, p. 148, pl. xi, figs. 1 and 2. Extinct Vertebrata from the Judith river and Great Lignite formations of Nebraska.

The Red Deer river specimen of *T. forcatus* consists of the carapace, in which the nuchal plate is missing, without any part of the plastron. Sternal plates, probably referable to this species, were found not far distant, but further reference will not be made to them here.

The carapace (Plate II, fig. 2) is a little less than one fourth broader than long, and is only slightly convex. The eight costals of the left side are practically entire, but the first right costal is wanting. Five neural plates are preserved with part of another that lay principally between the first costals, its posterior end being between the front inner borders of the second costals. If this neural were as long in proportion to its breadth as the plate immediately following it, its front margin probably effected a sutural union with the nuchal plate, and its outline was presumably as is shown by the dotted lines in the figure.* Neural two, three and four are six sided, neural five is oblong, and neural six, lying for the most part between the sixth costals, is shield shaped, coming to a point behind. The seventh costals are sutureally united at their inner ends, where they develop a breadth sufficient to separate the eighth costals from each other. These latter are subtriangular in shape, with three convex sides. The nuchal plate was evidently small, as the lateral termination of the suture (*a* in the figure) between it and the left first costal indicates a side extension scarcely past a point in advance of the mid length of the first costal. Small, shallow, rounded depressions mark the surface of the neutrals and the inner ends of the costals. In the latter, as the distance from the neutrals increases, the depressions gradually grow larger and more decided, becoming often reniform or oval, and frequently coalescing, until in the distal ends of the costals a few more or less continuous furrows are formed parallel with the outer margins of the plates. These furrows are a conspicuous feature in the sculpture; they are not so well marked on the posterior margin of the carapace, but they are well developed near the front edges of the first costals. In the neutrals and inner halves of the costals there is a narrow, smooth strip devoid of sculpture bordering the sutures. The rib-heads are well developed.

*Since the above was written some misplaced fragments have been added to the anterior end of this carapace; they reveal the fact that the first costals are in reality separated by a divided first neural. The writer wishes here to express his indebtedness to Professor Osborn and Dr. Hay, of the American Museum of Natural History, for the loan of the type of *Trionyx raganus* and the opportunity thus afforded him of making a direct comparison of the type with the Red Deer specimen.

Measurements

	M.
Estimated length of carapace along median line, in inches.	170
Distance from front margin of first costal to back margin of seventh costal (slightly curved) in inches.	107
Maximum breadth of carapace (8 th inchest).	211
Length of second neural.	023
Maximum breadth of second neural.	017
Thickness of fourth costal at centre in millimetres.	005
Thickness of fourth costal at centre on anterior end.	005
Thickness of eighth costal at centre.	005

Trionyx planus,* Owen and Bell, a British Lower Eocene species, described from the posterior half of the carapace, bears a strong general resemblance to *T. foreatus* as regards the sculpture and also in the absence in both species of the seventh and eighth neurals accompanied by a similar curtailment of the sixth neural. The strong development of the seventh costals found in the Red Deer river carapace resulting in a reduction of the eighth pair of costals may be an irregularity of growth of no specific importance.

The second species of *Trionyx*, represented by an almost complete carapace is referred to *T. rayanus*, Cope, first described by that authority in 1874† from 'a number of fragments of costal bones and perhaps of sternals also,' from the 'Lignite cretaceous of Colorado; near the mouth of the Big Horn river, Montana; Long lake, Nebraska, found at the last two localities by Dr. Hayden.' Later, in 1875, in 'The Vertebrata of the Cretaceous Formations of the West' the same description appears: this time with figures of two fragments of costal bones. Dr. O. P. Hay, of the American Museum of Natural History, New York, has kindly compared part of a costal plate from the Red Deer river with fragments labelled *T. rayanus* by Professor Cope in the New York collections. Owing to the insufficiency of the published description and figures of this species, the writer could scarcely judge whether the Canadian specimens were or were not referable to *T. rayanus*, but Dr. Hay writes that he thinks they probably are.

The carapace (Plate IV, fig. 4) is broader than long, the breadth exceeding the length by more than one-sixth and it is only moderately convex. In outline, as seen from above, it is flat behind with the sides

*Monograph of the British Reptilia of the London Clay, part I., p. 58, tab. XIXC, 849. Palaeontographical Society.

†Bulletin of the United States Geological Survey of the Territories, 1874, No. 2, p. 27.

curving broadly to the front margin, at the centre of which there is a shallow concavity. The shell protrudes where the ribs pass outward from beneath, causing the lateral margin to be sinuous, the sinuosity being most marked toward the front in the first, second and third pairs of costal plates. Of the eight pairs of costals, the first costals are the broadest at the inner ends, whilst the fifth are the broadest distally. The seventh costals are extremely narrow throughout their length and the eighth pair is well developed. The first costals increase in breadth rather suddenly at their outer ends and are separated by a divided first neural plate. The neutrals gradually decrease in breadth to the fifth, their sides being not so nearly parallel to each other as those of the corresponding plates in *T. arcatus*. The sixth and last neural is very much reduced in size and is irregularly oval in outline. Of the protruding rib ends, all the six of the left side were secured, in a fair state of preservation, except the one belonging to the first costal and it was only lined in part. The rib heads are well developed. In the figure, the nuchal plate is represented as entire. Of the carapace under consideration, the central part only of the nuchal plate, extending from the margin in front to the suture behind, was found, but fortunately the left end of a nuchal, of another individual of similar size, showing the left front margin and the suture between the plate and the first costal with part of the latter adherent, supplied the deficiency. The sculpture consists of a network of narrow ridges ramifying and inclosing so as to enclose small, sunken areas of irregular shape and size, the areas being generally wider than the ridges are broad. The frequent confluence of a varying number of areas results in a more open pattern, the ridges at times shewing a tendency to run in parallel lines. The sculpture is not so distinctly defined near the sides of the carapace as it is toward and at the centre and anteriorly, but in the hinder part it is more decidedly rugose, the ridges being here higher and the enclosed areas larger. Near the intercostal sutures, more particularly in the inner halves of the costal bones, the sculpture is partially effaced and consists of low, poorly defined parallel ridges at right angles to the sutures, forming a distinct border, with a maximum breadth of about 5 centimeters. A smooth border, broadest at the sides of the carapace and narrowest in front, extends along the whole of the peripheral edge.

As regards a divided first neural in species of this genus, it is interesting to note that Lydekker in describing *T. melitensis*, from the Miocene of Malta, in 1891, (Quarterly Journal of the Geological Society, vol. XLVII, p. 37, fig. 1) mentions the occurrence in the Miocene species of a divided first neural and remarks (p. 37) that 'all the fossil species

hitherto described, of which the entire carapace is known, agree with the normal type in having but a single long neural between the first pair of costals.¹

Measurements

	M.
Length of carapace along median line (8½ inches)	470
Maximum breadth of carapace (23½ inches)	500
Breadth of first costal at inner end	075
Thickness of same near inner end	009
Thickness of same at outer end	012
Breadth of fifth costal at outer end	119
Thickness of same at outer end	013
Breadth at mid length of seventh costal	036
Thickness at centre of eighth costal	010
Maximum breadth of neural I _a	056
Maximum breadth of neural I _a	039
Length of same	013
Thickness of median plate at left end	018
Length of vertebral centrum	045

EXPLANATION OF PLATES.

PLATE I.

FIGURE 1. Upper surface of the carapace of *Trionyx forsteri*, Leidy; from the Cretaceous of Alberta. One-half natural size. The sinuous lines indicate the sutures between the bones; the dotted ones parts restored; N1, nucha bone; N 1, 2, &c., neural bones; C 1, 2, &c., costal bones.

FIGURE 1a. Outline of the transverse curve of the upper surface of the carapace.

PLATE II.

FIGURE 2. Upper surface of the carapace shewn in plate I; from a photograph. Considerably over one-half the natural size.

PLATE III.

FIGURE 3. Upper surface of the carapace of *Trionyx vagans*, Cope; from the Cretaceous of Alberta. One eighth the natural size. Lines and letters as in figure 1 of plate I.

FIGURE 3a. Outline of the transverse curve of the upper surface of the carapace.


FIGURE 3b. The sculpture of part of the upper surface of the same carapace; from a photograph. Natural size.


PLATE IV.

FIGURE 4. Upper surface of the carapace shewn in plate III; from a photograph. Slightly less than one-fourth natural size.

GEOLOGICAL SURVEY OF CANADA


PLATE I.





GEOLOGICAL SURVEY OF CANADA.

PLATE II.





2.




GEOLOGICAL SURVEY OF CANADA

PLATE III.



3b.



GEOLOGICAL SURVEY OF CANADA.

PLATE IV.



