





**IMAGE EVALUATION
TEST TARGET (MT-3)**



**Photographic
Sciences
Corporation**

23 WEST MAIN STREET
WEBSTER, N.Y. 14580
(716) 872-4503

15 28 25
18 22 20
18

**CIHM/ICMH
Microfiche
Series.**

**CIHM/ICMH
Collection de
microfiches.**



Canadian Institute for Historical Microreproduction / Institut canadien de microreproductions historiques

11
10
11

© 1981

Technical and Bibliographic Notes/Notes techniques et bibliographiques

The Institute has attempted to obtain the best original copy available for filming. Features of this copy which may be bibliographically unique, which may alter any of the images in the reproduction, or which may significantly change the usual method of filming, are checked below.

L'Institut a microfilmé le meilleur exemplaire qu'il lui a été possible de se procurer. Les détails de cet exemplaire qui sont peut-être uniques du point de vue bibliographique, qui peuvent modifier une image reproduite, ou qui peuvent exiger une modification dans la méthode normale de filmage sont indiqués ci-dessous.

- Coloured covers/
Couverture de couleur
- Covers damaged/
Couverture endommagée
- Covers restored and/or laminated/
Couverture restaurée et/ou pelliculée
- Cover title missing/
Le titre de couverture manque
- Coloured maps/
Cartes géographiques en couleur
- Coloured ink (i.e. other than blue or black)/
Encre de couleur (i.e. autre que bleue ou noire)
- Coloured plates and/or illustrations/
Planches et/ou illustrations en couleur
- Bound with other material/
Relié avec d'autres documents
- Tight binding may cause shadows or distortion along interior margin/
La reliure serrée peut causer de l'ombre ou de la distorsion le long de la marge intérieure
- Blank leaves added during restoration may appear within the text. Whenever possible, these have been omitted from filming/
Il se peut que certaines pages blanches ajoutées lors d'une restauration apparaissent dans le texte, mais, lorsque cela était possible, ces pages n'ont pas été filmées.
- Additional comments:/
Commentaires supplémentaires:

- Coloured pages/
Pages de couleur
- Pages damaged/
Pages endommagées
- Pages restored and/or laminated/
Pages restaurées et/ou pelliculées
- Pages discoloured, stained or foxed/
Pages décolorées, tachetées ou piquées
- Pages detached/
Pages détachées
- Showthrough/
Transparence
- Quality of print varies/
Qualité inégale de l'impression
- Includes supplementary material/
Comprend du matériel supplémentaire
- Only edition available/
Seule édition disponible
- Pages wholly or partially obscured by errata slips, tissues, etc., have been refilmed to ensure the best possible image/
Les pages totalement ou partiellement obscurcies par un feuillet d'errata, une pelure, etc., ont été filmées à nouveau de façon à obtenir la meilleure image possible.

This item is filmed at the reduction ratio checked below/
Ce document est filmé au taux de réduction indiqué ci-dessous.

10X	14X	18X	22X	26X	30X
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>
12X	16X	20X	24X	28X	32X

The copy filmed here has been reproduced thanks to the generosity of:




Legislative Library of
British Columbia

The images appearing here are the best quality possible considering the condition and legibility of the original copy and in keeping with the filming contract specifications.

Original copies in printed paper covers are filmed beginning with the front cover and ending on the last page with a printed or illustrated impression, or the back cover when appropriate. All other original copies are filmed beginning on the first page with a printed or illustrated impression, and ending on the last page with a printed or illustrated impression.

The last recorded frame on each microfiche shall contain the symbol \rightarrow (meaning "CONTINUED"), or the symbol ∇ (meaning "END"), whichever applies.

Maps, plates, charts, etc., may be filmed at different reduction ratios. Those too large to be entirely included in one exposure are filmed beginning in the upper left hand corner, left to right and top to bottom, as many frames as required. The following diagrams illustrate the method:



L'exemplaire filmé fut reproduit grâce à la générosité de:

Legislative Library of
British Columbia

Les images suivantes ont été reproduites avec le plus grand soin, compte tenu de la condition et de la netteté de l'exemplaire filmé, et en conformité avec les conditions du contrat de filmage.

Les exemplaires originaux dont la couverture en papier est imprimée sont filmés en commençant par le premier plat et en terminant soit par la dernière page qui comporte une empreinte d'impression ou d'illustration, soit par le second plat, selon le cas. Tous les autres exemplaires originaux sont filmés en commençant par la première page qui comporte une empreinte d'impression ou d'illustration et en terminant par la dernière page qui comporte une telle empreinte.

Un des symboles suivants apparaîtra sur la dernière image de chaque microfiche, selon le cas: le symbole \rightarrow signifie "A SUIVRE", le symbole ∇ signifie "FIN".

Les cartes, planches, tableaux, etc., peuvent être filmés à des taux de réduction différents. Lorsque le document est trop grand pour être reproduit en un seul cliché, il est filmé à partir de l'angle supérieur gauche, de gauche à droite, et de haut en bas, en prenant le nombre d'images nécessaire. Les diagrammes suivants illustrent la méthode.

tails
du
odifier
une
mage

rrata
o

elure,
n à

32X



GEOLOGICAL SURVEY OF CANADA.

ALFRED R. C. SELWYN, C.M.G. F.R.S., Etc., Director.

CONTRIBUTIONS

TO

CANADIAN PALÆONTOLOGY.

VOLUME III. (Quarto).

ON VERTEBRATA FROM THE TERTIARY AND CRETACEOUS ROCKS
OF THE NORTH WEST TERRITORY.

BY

E. D. COPE.

1.—The Species from the Oligocene or Lower Miocene beds of the Cypress Hills.



PRINTED FOR THE GOVERNMENT OF CANADA.

WILLIAM FOSTER BROWN & CO., PUBLISHERS.

MONTREAL.

1891.

1629

211

60

V. 2. 11


The present Report is the first of a series of descriptive and illustrated quarto memoirs on the Vertebrata of the Tertiary and Cretaceous rocks of the Canadian North West Territory, kindly prepared for the Survey by Professor E. D. Cope of Philadelphia. It is exclusively devoted to a consideration of the species from the Lower Miocene deposits of the Cypress Hills in the district of Alberta, and consists of twenty-seven pages of letter press, illustrated by fourteen full page lithographic plates.

Part 2, which will contain illustrated descriptions of the vertebrates of the Laramie formation of the North West Territory, by the same author, is now in course of preparation.

ALFRED R. C. SELWYN.

GEOLOGICAL SURVEY DEPARTMENT,

Ottawa, 28th February, 1891.



GEOLOGICAL SURVEY OF CANADA.

THE VERTEBRATA OF THE TERTIARY AND CRETACEOUS ROCKS OF THE NORTH WEST TERRITORY.

BY E. D. COPE.

1. *The Species from the Oligocene or Lower Miocene beds of the Cypress Hills.*

The collection on which the present report is based was made by Messrs. R. G. McConnell and T. C. Weston, in 1883-84, in the Cypress Hills, North-West Territory, about long. 109°, lat. 49° 40'. The Cypress Hills, in the District of Assiniboia, were examined geologically by Mr. R. G. McConnell, of the Dominion Geological Survey in 1885. He found them to constitute a plateau of considerable extent, consisting largely of beds of conglomerate (see p. 31 C. Report Geol. Survey of Canada, 1885), chiefly quartzitic, and evidently derived from the harder and older rocks of the Rocky Mountains. The conglomeritic character of the beds accounts for the generally broken condition of the fossils.

Dr. A. R. C. Selwyn, Director of the Survey, having sent the fossils to me for identification, I gave a preliminary list of the species in the *American Naturalist* for February, 1885. It was then pointed out that the genera and species obtained by Messrs. McConnell and Weston proved the beds in question to belong to the White River Oligocene series. The presence of a genus of well-marked *Creodonts* (*Hemipsalodon*, Cope) was regarded as an indication that the Cypress Hills, Swift Current Creek beds are probably somewhat older than those of the typical locality on the White River of Dakota and Nebraska. But the presence of this genus may be yet ascertained in the latter locality. Explorations set on foot during the year 1888 resulted in the obtaining by Mr. T. C. Weston, of the Canadian Survey of a number of additional species, some of which are of considerable interest. These were described in the *American Naturalist* of 1889. Most of these specimens were also in a fragmentary condition, owing to the conglomeritic nature of the deposit. The total number of species is twenty-five.

PISCES.

HALECOMORPHI.

AMIA L.

The *Amiidae* appear in the Laramie formation, and are represented in the Puerco and in the Wasatch Eocene. They occur abundantly in the Bridger Eocene, but were not known from any later formation, until discovered by Mr. Weston in the Cypress Hills Oligocene beds, as here recorded. As *Amia* is a well known genus of the present period, it is to be looked for in all the beds between the Bridger Eocene and the Pleistocene, inclusive.

AMIA WHITEAVESIANA. Sp. nov.

Founded on an anterior vertebra of large size, which differs in various respects from that of the Eocene species. There are no diapophyses, and there are tuberosities external to and adjoining the neuropophysial facets. The latter are so much worn that it is impossible to make out their character. There is a minute, round notochordal foramen above the middle of the centrum. In outline the centrum is a transverse oval, more transverse than in any other species of the genus at present known. External walls smooth. Two fossae, separated by a narrow median rib, on the inferior surface.

<i>Measurements.</i>		<i>Mm.</i>
Diameters of centrum {	Anteroposterior	9
	Vertical.....	26
	Transverse.....	40

This species is dedicated to Mr. J. F. Whiteaves, of Ottawa, the distinguished palaeontologist of the Geological Survey of Canada.

AMIA MACROSPONDYLIA. Sp. nov.

This species is also indicated by an anterior vertebra. It agrees with the last in lacking diapophyses, and in having a minute *foramen corbo-dorsalis*, which is, however, nearer the middle of the centrum than in the *A. whiteavesiana*. The *A. macrospondylia* differs much from the latter in its proportions. The vertebra, while absolutely smaller, has a larger anteroposterior diameter, as shown by the measurements below. The neuropophysial facets are too much worn for description. The wearing has left the appearance of a pair of parallelogrammic fossae at opposite positions of the external wall of the centrum. Where unworn, the surface displays delicate wrinkles in the direction of the long axis of the vertebral column.

<i>Measurements.</i>		<i>Mm.</i>
Diameters of centrum {	Anteroposterior.....	12
	Vertical.....	22
	Transverse.....	26

NEMATOGNATHI.

? RHINEASTES, Cope.

A species of Siluroid is referred with doubt to this genus, since the parts essential for exact determination are wanting. The disposition of the tissue of the vertebral centrum is in vertical laminae, parallel to those of the articular faces, as is the case with such vertebrae of Rhineastes as are known. Diapophyses are present, and the neuropophysies are coossified with the centrum. The lateral walls of the centrum below the diapophysis are not excavated by fossae.

This genus has been known hitherto from the Eocene only.

RHINEASTES RILEYI. Sp. nov.

Represented by a median abdominal vertebra, with which is associated a second, as belonging to the same genus and perhaps species.

The centrum of the former is nearly entire. Its articular faces are a little deeper than wide, and are marked with smooth concentric striae. No notochordal foramen. Centrum not flattened below, and furnished with a single median fossa. Neural canal with a longitudinal fossa on each side. Two large fossae on each side between the bases of the neurapophysis and diapophysis. No fossae on the side of the centrum except a small one below and at the base of the diapophysis. The exposed edges of the vertical laminae are close together at the middle of the side of the centrum, and they turn forwards below, converging to a point on each side of the inferior fossa. They are less numerous just below the fossa below the diapophysis, and are connected by longitudinal bars. The bases of the diapophyses (all that remains of them) are hollow.

Measurements.

	Min.	
Diameters of centrum {	Anteroposterior	12
	Vertical	31
	Transverse	24

The second centrum is fragmentary and may represent another species. It differs in the presence of a very large fossa immediately below the diapophysis, and in the presence of one very large one above the diapophysis. The concentric lines of the articular faces are more prominent. Size similar.

AMIURUS, Raf.

Two species have left vertebrae in the formation of the Cypress Hills, which resemble those of this genus of Siluroids, which still inhabits North America. That they belong to it cannot be positively asserted, since important diagnostic parts of the skeleton are unknown. Neurapophyses coossified; lamination longitudinal; base of diapophyses hollow.

AMIURUS CANCELATUS. Sp. nov.

Two vertebral centra represent this species—one from the median dorsal region, and the other probably from the caudal.

The first-mentioned and best-preserved centrum has sub-round articular faces, a little flattened above and below, and is without *foramen coraco-dorsalis*. The floor of the neural canal and of the space between the neurapophyses and the diapophyses are excavated, and the floor of both spaces is composed of longitudinal laminae, which separate rather coarse and deep fossae, the whole resembling cancellous tissue. Inferior median line with a principal median fossa. Dense tissue of the articular faces, reverted on the sides of the centrum. The space between them with minute longitudinal pores. These become coarser and more cancellous as we approach the base of the diapophysis, where a larger cavity, more or less divided by cancellae, is placed. Concentric layers of articular faces with distinct edges, producing a slight rugosity.

Measurements.

		Mm.
Diameters of centrum	Anteroposterior	11
	Vertical	24
	Transverse	25

The second centrum is perhaps caudal, as it resembles those of *Amiurus* in having a lateral fossa below as well as above the median lateral portion. The latter is coarsely reticulate cancellous, and a deep fossa separates the two ridges which represent the bases of either the neural or the haemal spines. The anteroposterior diameter is relatively greater than in the vertebra last described.

Measurements.

		Mm.
Diameters of centrum	Anteroposterior	12
	Vertical (restored)	28
	Transverse	23

AMURUS MACONNELLII. Sp. nov.

This species appear to have been larger than the last. It differs from it in the much finer cancellation of the lateral surfaces, over which the articular dense layer is much less reverted. Instead of a cancellated tract between the diapophysis and the neural spine, there are two deep fossae, and there are two large shallow fossae on the inferior face instead of one deep one. Near the articular borders a few cancellous lines are vertical in direction. The species presents some of the characters of the *Rhineastus chicus*. Vertebrae, only, of this species were found. They indicate large size, the measurements being:

		Mm.
Diameters of centrum	Vertical	40
	Transverse	38
	Anteroposterior	11.5

This centrum has a subquadrate outline, so far as can be determined in the absence of the superior border. It has a minute transverse notochordal foramen. The diapophysis is rudimental and superior in position; it presents no articular facet. Both above and below it, is a shallow fossa. Except a pair of shallow fossae on the inferior face, the lateral walls are without fossae, but are finely impressed with minute longitudinal pits.

A second centrum, probably caudal, belongs to a still smaller individual. Its outline is subpentagonal, the apex inferior and without a single median fossa. No diapophyses. Three superior fossae—one large median and a smaller one on each side. Minute pits of external walls sub-round.

		Mm.
Diameters	Vertical	21
	Transverse	21
	Anteroposterior	7.5

This catfish equalled in dimensions the large *A. nigricans* of the Mississippi. It is dedicated to the able geologist Mr. R. G. McConnell, of the Dominion Survey.

REPTILIA.

TESTUDINATA.

TRIONYX, Geoffr.

TRIONYX LEUCOPOTAMICUS. Sp. nov.

This species is represented by a part of a single costal bone, with the sculpture well preserved. This specimen would scarcely serve as a basis for a specific description, but I have a number of well-preserved fragments of apparently the same turtle from the White Buttes of Dakota, which render important aid. The latter furnished me with the first indication of the presence of this genus in beds of White River age, as they have not been yet found in the best known regions of this formation in Southern Dakota and Eastern Colorado.

The costal bones are flat and rather thin, but thickest medially, as usual. The sculpture consists of large, sub-round and sub-oval fossae, which are separated by ridges much narrower than themselves. In the Cypress Hills specimen the fossae tend to form longitudinal series towards the distal extremity of the bone. The sutural borders are not preserved, but in the Dakota specimens the sculpture continues to show, and does not give place to a band of nearly smooth and transversely-lined surface, as is the case in the *T. punctiger*,¹ Cope, also from the White River bed of Central Dakota.

This species resembles in its sculpture the *Trionyx niutawensis* of Leidy from the Bridger Eocene. The sole difference which the latter presents is the presence of bands of grooves parallel to the long axis of the carapace, along the intercostal sutures. It is probable that the costals are thicker in the Bridger species, a character still more strongly marked in those of the Wasatch series.

STYLEMYS, Leidy.

(?) STYLEMYS NEBRASCENSIS, Leidy.

A few fragments, including a marginal bone, not distinguishable from this species, were obtained by Mr. Weston.

MAMMALIA.

GLIRES.

PALÆOLAGUS, Leidy.

PALÆOLAGUS TURGIDUS, Cope.

Report U. S. Geol. Survey Terrs., III, Bk. I., p. 882, pl. Ixvi., f. 28; Ixvii., 13-27.

Mandibular rami identical in character with those from the White River beds of Dakota and Colorado.

¹ Besides the character above described, the sculpture of the costal bones of the *T. punctiger* is much less distinct than in the *T. leucopotamicus*. It is both punctate and groove-like, and the ridges are thickened and irregular, and towards the distal ends of the bones obscure. While not wider than those of the latter species, the costals are thicker and more curved, the thickness continuing to the free margin, where they are bevelled off, with a prominent rib-end in the middle. Measurements of No. 1:—Width of costal, 18 mm.; thickness at lateral suture, 5 mm.; at middle, 9 mm. No. 2:—Width of costal, 52 mm.; thickness at middle, 11 mm. This is *Trionyx* sp. 2, Cope Proceeds. Am. Philos. Soc., 1883, p. 217.

BUNOTHERIA.

CREODONTA.

HEMIPSALODON, Cope.

This genus belongs to the Oxyænidæ, and is the only one of that family that has been found in beds higher than the Bridger Eocene. The only known species is the largest of the Creodonts, and the jaw from which it is known is more robust than that of any existing carnivore. Its dimensions are about those of the *Achænodon insolens* of the Bridger beds. The genus *Hemipsalodon* differs from the others of the family in the presence in the lower jaw of the full dental series of four premolars and three true molars without diastema behind the canine. Incisors three. The only crown perfectly preserved is the last true molar. It is of the type of *Oxyæna*, but has probably no internal tubercle or metacone; (specimen worn at that point). It has a heel more or less cutting.

This remarkable genus resembles, so far as the characters preserved permit us to judge, the *Stypolophus* of the Eocenes, which embraces numerous species, none of which exceeded a red fox in size. It is, however, most nearly related to *Pterodon*, with which Schlosser believes it to be identical. That possesses but three inferior premolars, but the value of this character in this case is not yet certainly known. If no other character distinguish *Hemipsalodon* but the four inferior premolars, the *Pterodon quercyi* of Filhol, from the French phosphorites, must be placed in it. The blade of the last inferior molar is more oblique to the long axis of the jaw than is represented to be the case in the *Pterodon dasyroides*, making an angle of 30° with it, while the edge of the heel is slightly oblique to it in the opposite direction. This may indicate some peculiarity in the superior molar, or it may be but a specific character.

HEMIPSALODON GRANDIS, Cope.

American Naturalist, 1885, p. 163; Annual Report of the Geol. and Nat. History Survey of Canada, 1885, Appendix C, p. 2.

The species is characterized by the deep compressed form of the ramus, and the long symphysis. The incisor teeth are crowded, the first and third being external and close together, and the second internal in position. The canine tooth is of enormous size, and is directed upwards. The section of the summit of the root is a wide oval without angles. The premolars are all two-rooted, except the fourth. The first is longer than the first true molar. The true molars increase in size posteriorly. The third is very robust, and has elevated cusps, with a sub-triangular section, the median exceeding the anterior. The sectorial edges are very steep, forming together a V. The heel is quite short, and has a cutting keel, which is the summit of the external face, and is nearly median. The coronoid process rises at a very short distance posterior to it. The masseteric fossa does not extend downwards to the inferior edge of the ramus. The latter is indented on the inner side as far posterior as below the middle of the coronoid process, where it is broken off.

Length of the dental series, M. .212; of premolars, .108; of true molars, .108; diameters of last true molar: anteroposterior .034, transverse .021; do. of canine at base: anteroposterior .040, transverse .029. Depth of ramus at M. 3, .036; length of symphysis, .131.

This species was the largest flesh-eater of the epoch of the White River beds, and the size of its canine teeth proves it to have been a dangerous animal. Its molars are interesting on account of their illustrating a primitive form of a sectorial tooth.

Two femora in the collection probably belong to this species, as they are creodont in the presence of the third trochanter, and their dimensions are appropriate to the lower jaw above described. One of them is larger than the other, and I give its dimensions in comparison with those of two of the largest Carnivora, the grizzly bear and the lion, in millimeters:—

	Hemipusodon.	Bear.	Lion.
Length	413	145	379
Width at great trochanters.....	129	112	95
“ “ little “	71	57	45
“ “ third “ or same position	63	40	35
“ “ condyles	90	90	81
Depth at rotular ridges	98	73	77

The third trochanter is low, much as in *Protopsalis tigrinus*, and has a thickened, rough edge. It gives the shaft of the femora an external convexity, which is greater than the gentle concavity of the internal border. The great trochanter projects to the horizontal line of the head and not beyond. It is obliquely truncate externally, and narrowly at the extremity. It encloses a deep trochanteric fossa posteriorly, through the strong recurvature of the posterior border. This border continues as a transverse convexity to the little trochanter. The latter is a rounded prominent tuberosity, and has a superior position, as in Carnivora generally, and is not placed low down on the shaft as in *Protopsalis*. The shaft is flattened from before backwards, with the external edge angulated both above and below the third trochanter. The rotular groove is remarkably elevated and rather narrowed, giving the distal extremity of the femora the massive character of that of an ungulate mammal. Borders of rotular groove subequally prominent, its surface continuous with that of the condyles. Internal face of condyle with a pronounced fossa. Internal condyle a little more prominent than the external. Intercondylar fossa broadly rounded anteriorly.

The deeper trochanteric fossa and more elevated position of the little trochanter distinguish these femora from that of *Protopsalis tigrinus*. It also considerably exceeds that of the latter animal in dimensions.

ANCYLOPODA.

CHALICOTHERIUM, Kaup.

Macrotherium, Lartet, teste Forsyth-Major and Filhol.

The remarkable character of this genus, as discovered by Filhol, has been mentioned in the American Naturalist.¹ It has little relation to the family of Perissodactyla to which it has given the name, and which it so resembles in molar dentition. It must form a family by itself, and the genera with which it has been associated must form a family to which the name Lambdotheriidae has been applied. The anterior ungual phalanges of Chalicotherium are of prehensile character and not ungulate, but rather unguiculate. The phalanges resemble those of the Edentata, but the carpus and tarsus are according to Filhol, diarthrous in structure, while the Edentata are taxepodous. We have in the Chalicotheriidae the antithesis of the Condylarthra. While the latter is

¹ Cope, American Naturalist, 1889, p. 153; American Journal of Morphology, 1889, p. 112.

² Osborn on Chalicotherium, American Naturalist, 1888, p. 728.

ungulate with an unguiculate carpus and tarsus, the former is unguiculate with an-ungulate (diplarthrous) carpus and tarsus. Thus the Chalicotheriidae must be referred to a distinct order of unguiculate Mammalia, which I have called the Ancylopoda, with the above definition. Two genera belong to the single family, the Chalicotheriidae, viz., Chalicotherium Kaup, and Ancylotherium Gaudry. Marsh has not yet shown how his genus *Moropus* differs from Ancylotherium. The species described by Marsh under this name are from the Loup Fork bed of Kansas.

Although the species of Chalicotherium from the Cypress Hills is the first one described from North American beds, it is not the first discovery of the genus. Professor Scott showed me a series of superior molars from the Loup Fork formation of Kansas, from the Agassiz Museum, which he identified as belonging to this genus. The present species is of larger size than the Kansas form, and is apparently equal to the *C. goldfussii* of the Upper Miocene of Europe. The occurrence of this form in the Oligocene or Lower Miocene (White River), as well as the Upper Miocene (Loup Fork), of this country, is a noteworthy fact, but is parallel to its history in Europe. Described from the Upper Miocene by Kaup, it was afterwards found in the Middle Miocene (*C. grande*) by Dartet, and in the Upper Eocene (*C. modicum*) by Gaudry.

CHALICOTHERIUM BILOBATUM, Cope.

American Naturalist, 1889, p. 151.

Founded on a mandibular symphysis and part of the left ramus of an adult animal, which contains the alveoli of the anterior four molars and part of that of the fifth. All the premolars are two-rooted, showing that they are but three in number. Canines and incisors wanting, the anterior alveolar margin thin and prominent and bilobed, with a median emargination. Symphysis coossified, with an angulate inferior margin, posteriorly with a fossa on each side of the median line, sloping regularly upwards to the alveolar margin, and concave above behind the margin. Minute traces of alveoli of a canine and two incisors on each side, which were probably present in the fetus. Length of symphysis above, 120 mm.; depth posteriorly, 48 mm. Length of symphysis in front of p. m. iii. Length of premolar series, 75 mm. Length of m. i., 40 mm.

DIPLARTHRA.

PERISSODACTYLA.

MENODUS, Pomel.

This genus has received numerous accessions within the last few years through the labors of Professors Scott, Osborn¹ and Marsh. To these may be added the species obtained from the Cypress Hills area of the White River series. As remarked by Professors Scott and Osborn, the species of *Menodus* present parallel relations to those of *Symborodon*, which latter may be regarded as descendants of the former, of more special-

¹ Preliminary Account of the Fossil Mammals from the White River Formation in the Mus. Comp. Zoology, Bull. of the Mus., Vol. xlii, No. 5, 1887; Amer. Jour. Sci. and Arts, 1887, p. 323; Cope, Amer. Naturalist, 1887, p. 926.

ized character and probably later age. The nine species of *Menodus* may be compared with five of those of *Symborodon*, as follows:—

	Nasals long; horns short.	Nasals and horns intermediate.	Nasals short; horns long.	
<i>Menodus</i>	Nasals transversely angulated.....	<i>M. americanus</i> ,	<i>M. syceras</i> ,	
		<i>M. coloradensis</i> .	<i>M. proutii</i> ,	
	Nasals not angulated....	<i>M. angustigenis</i> .	<i>M. tichoeras</i> .	<i>M. platyceras</i> ,
		<i>M. schwynianus</i> .		<i>M. dolichoeras</i> .
<i>Symborodon</i>	Nasals angulated.....	<i>S. trigonoceus</i> ,		
	Nasals not angulated....	<i>S. bucco</i> ,	<i>S. altirostris</i> ,	
			<i>S. accr.</i>	

Apart from the generic characters, the *Menodus americanus*, Leidy, appears to be an enlarged *S. trigonoceus*, Cope; and the *M. tichoeras*, S. and O., an enlarged *S. altirostris*, Cope. Apart from these, the correspondences are not so close. A definite character which divides *Menodus* into two groups is the presence of an internal cingulum of the premolars in some of the species. The same character divides the genus *Symborodon*. According to this character, the species may be grouped as follows, so far as they are known in this respect:—

	<i>Menodus</i> .	<i>Symborodon</i> .
Without cingulum.....	{ <i>M. proutii</i>	<i>S. bucco</i>
	{ <i>M. tichoeras</i>	<i>S. altirostris</i>
	{ <i>M. americanus</i>	<i>S. trigonoceus</i>
With cingulum.....	{ <i>M. coloradensis</i>	<i>S. heloeras</i>
	{ <i>M. angustigenis</i>	

The collections of the Geological Survey include fragments of skeletons of five species of *Menodontida*, which I shall refer, some of them provisionally, to the genus *Menodus*. The *M. americanus* is the only one known to have possessed horns of triangular section. The section is oval in the *M. proutii* and *M. angustigenis*. In *M. schwynianus* the horns are unknown.

In the Annual Report of the United States Geological Survey of the Territories for 1874, p. 489, I gave a general account of the osteology of the genus *Symborodon*, which applies equally well to the nearly allied genus *Menodus*. I will now add a few points not referred to in that report, derived from the specimens here described.

The condyles of the humerus have no trace of trochlear crest. The olecranon is expanded laterally and vertically at the extremity. The head of the radius is only convex below. Its carpal extremity is narrowed inwards, and it is bounded below its middle by a fossa. The scaphoid and lunar parts of the surface are not distinguished.

The third trochanter of the femur is not a process but an angle, projecting but little beyond the external face of the femur above it, but bounding a contraction of the diameter below it. The little trochanter is insignificant. The rotular surface is grooved medially, and the lateral ridges are prominent, especially above and proximally, where they rise abruptly from the shaft, which has at this point, and between them, a fossa.

smaller than those of the species before mentioned. There is a considerable popliteal callosity, and, what is especially characteristic, the crest is divided by a vertical median groove, as in the genera *Aphelops* and *Rhinoceros*. This type of tibia corresponds to that of some of the species of *Symborodon*, e. g., *S. bucco*, and perhaps others. There is no tibia nor any other bone, except perhaps part of a humerus, which corresponds with the smallest astragalus.

The bones of the two largest species are probably those of the *M. americanus* and *M. proutii*, but which belongs to which I cannot certainly determine. As the *M. americanus* is, in the characters of its muzzle and horns, the nearest to the *M. angustigenis*, it will be probably safe to refer to it the tibia which approaches the corresponding bone of that species. That will be the second of the above description. The third corresponds in size to the *M. angustigenis*, and agrees with it in approaching the species of *Symborodon* in some respects. I give corresponding measurements of the tibia, astragalus and calcanea of the two species:—

	<i>M. proutii.</i>	<i>M. americanus.</i>
I. TIBIA.		
Total length.....	387	355
Diameters of head {		
Anteroposterior.....	135	150
Transverse.....	144	135
Diameters of shaft {		
Anteroposterior.....	60	55
Transverse.....	60	60
Distal diameters.. {		
Anteroposterior.....	72	65
Transverse.....	100	100 ¹
II. CALCANEUM.		
Length.....	155	135
Length of tuber.....	90	80
Width distally.....	75	57
Diameters of cuboid facet {		
Vertical.....	35	42
Transverse.....	55	30
III. ASTRAGALUS.		
Diameters of trochlea {		
Transverse.....	80	70
Anteroposterior {		
External roller.....	70	57
Internal roller.....	57	
Diameters of navicular facet {		
Vertical.....	54	
Transverse.....	49	
Oblique length of cuboid facet.....	47	

From the above it is evident that the hind foot of the *M. americanus* is relatively and absolutely larger than in the *M. proutii*. In the former the external cotylus has a greater transverse, and the internal a smaller anteroposterior diameter than in that of the *M. proutii*.

¹ Partly restored.

I describe here a humerus which I suppose to belong to one or the other of these large species. The great tuberosity is of huge dimensions, exceeding in anteroposterior diameter the head and lesser tuberosity together. Its external face forms a triangular area which terminates at its inferior apex in a large compressed process which is directed outwards. Below this the smooth surface of the shaft winds spirally from a posterior position to the front, where it is bounded below by a narrow, deep, transverse coronoid fossa. It is bounded externally below by a ridge which is produced upwards from the internal epicondyle in a short tuberosity. The olecranon fossa is very deep and has abrupt sides. The external epicondylar region is flat.

Measurement of Humerus.

Length		Mm.	
		460	
Anteroposterior diameter	{	of entire head..... 210	
		of condyle only..... 110	
Transverse diameter	{	of head..... 158	
		of condyle..... 93	
Diameter at narrow part of shaft	{	Anteroposterior..... 80	
		Transverse..... 72	
Diameter of distal condyles	{	Transverse..... 130	
		{	External..... 98
			Internal..... 72
Distal width, with epicondyle.....		150	

I compare the only entire femora in the Cypress Hills collection with one nearly entire from Colorado in my collection. The latter was found near other bones which belong probably to the *M. americanus*, and the Cypress Hills specimen not improbably belongs to the *M. proutii*. The head and condyles are smaller than those of the *M. americanus*. The great trochanter extends as far proximad as the line of the head, while it falls considerably short of it in the *M. americanus*. The third trochanter is not prominent, but forms the inferior extremity of the wide proximal two-fifths of the bone, a slight concavity of the external border existing between it and the great trochanter. Below it the shaft is abruptly contracted on the external side. It then expands equally on each side to the condyles, giving the posterior face a wide, slightly concave face proximad to the latter. Their articular surfaces are continuous with each other, and with that of the rotular surface, at the point of junction of which their external borders are notched. The rotular groove and its lateral ridges are injured. Enough remains to show that there is a fossa at its proximal border on the shaft. The inferior part of the femora of the *M. americanus* does not differ except in its superior dimensions. The following are the measurements:—

Measurements of Femora.

	<i>M. proutii.</i>	<i>M. americanus.</i>
	Mm.	Mm.
Length	600	
Anteroposterior diameter of head.....	90	119
Width at great trochanter.....	187	182
“ “ third “	115	
“ (least) below third trochanter.....	76	96
“ above condyles	150	180
“ of condyles.....	125	150
Anteroposterior diameter (least) below third trochanter.....	55	60

Three scaphoid bones of graduated sizes are contained in the Cypress Hills collection, and there are two magnums which correspond in size to the middle-sized scaphoid and to one bone that size and the largest. These may be referred, comparing them with the other bones already described, as follows:—Scaphoid No. 1, *M. americanus*; magnum, No. 1, *M. proutii*; scaphoid and magnum No. 2, *M. angustigenis*; scaphoid No. 3, *M. (?) scerces*. Comparisons will be made under the head of *M. angustigenis*.

Teeth of the large species of Menodus are rare in the collection, a last inferior molar and premolar representing them. Most of the teeth preserved belong to the size appropriate to the *M. angustigenis*.

MENODUS ANGUSTIGENIS, Cope.

Annual Report Geol. and Nat. Hist. Survey of Canada, 1885, C, p. 81; *Haplocodon angustigenis*, Cope, American Naturalist, 1889, p. 153.

This large Mammal is represented by numerous specimens. I select for present description two maxillary bones from the same skull, each of which contains the first premolar and the true molars; and two lower jaws from second and third individuals. One of these consists of little more than the symphysis. The other includes part of the symphysis and part of the left ramus, which contains all the molar teeth except the first and last.

I refer the species to Menodus, because both lower jaws have, like the *Menodus proutii*, Leidy, two incisor teeth on each side. The specimen in which the ramus is present has a small alveolus for the first premolar on each side; the side of the other specimen, where this part is preserved, has no such alveolus. These specimens show the identity of the supposed genus Brontotherium with Menodus. In the contracted shape of its mandibular symphysis this species resembles the species of Symborodon rather than the *Menodus proutii*, and it resembles the smaller species of Symborodon in its inferior dimensions. It resembles the species of Menodus in the wide internal cingulum of the superior premolars. The species of Symborodon which present this character are the *S. trigonoceras* and the *S. heloceras*, Cope. Its measurements are inferior to those of the *S. trigonoceras*, and the superior molars are of different form. In the species just named their outline is oblong, the anteroposterior diameter exceeding the transverse in all three of them. In the *M. angustigenis* the molars are nearly square in outline.

The superior molars of the *S. trigonoceras* are characterized by the flatness of the middle portion of the external face of the external Vs. This surface is neither excavated, nor is it keeled, excepting a slight convexity on the middle of the anterior V of the first molar. The middle lines of the external faces of the Vs of the fourth premolar are slightly convex. There is a prominent vertical angle descending from the apex of each external V, and no lateral ones, so that there are no lateral pits at the internal base of the V on each side of the apex, as is seen in the *Symborodon trigonoceras*. The internal cones of the first superior premolar are not well distinguished. The only traces of cingula on the true molars are just in front of the median external vertical rib.

Probably belonging to the same species, and perhaps to the same individual, are the greater part of the frontal and nasal bones, with horns, and the right posterior base of the skull, with part of the zygoma. The first-mentioned fragment shows that the species differs from the *Symborodontes trigonoceras* and *acer*, Cope, and the *Menodus americanus* Leidy, in the absence of angulation above, between the free and other parts of the nasal bones. It also clearly differs from the *S. trigonoceras* in the semi-erect horns, with little pronounced triangular section. From the *S. bucco* the lack of expansion of the zygomatic bones distinguishes it. As compared with the *S. altirostris*, Cope, it has much longer and wider nasal bones, and the horns are more widely separated. The compression makes their apices anteroposterior, while they are transverse in the *S. altirostris*.

The ascription of long bones in the collection to this species is more or less uncertain, although a considerable number of pieces which correspond in size with the parts already described probably belong to it. A nearly complete tibia, and the proximal part of a second, may be with great probability assigned here. As already remarked, it differs from those previously described in having the crest divided vertically with a deep groove. The external side is rather the most prominent. A considerable callus occupies the popliteal region posterior to the spine and the external cotylus, from the latter of which it is marked off by a groove. The anteroposterior diameter of the external cotylus is greater than the transverse. Below the crest the section of the shaft is triangular, with obtuse angles anterior and exterior. This tibia is much like one which I obtained in Colorado, near to the skulls of several species of Symborodonts. A femur which accompanies the Cypress Hill tibia is of smaller size than that described under *M. proutii*, and is appropriate to the former. A similar femur accompanies the Colorado tibia. It resembles the one described under the *M. proutii* in having the great trochanter considerably less prominent than the head.

Measurements of Femur, in Mm.

	Cypress Hills.	Colorado.	
Anteroposterior diameter of head	×	95	
" " of great trochanter	×	80	
Width of head and great trochanter	×	115	
Distance from head to little trochanter, inclusive	×	2 × 5	
Width at condyles	110	×	
Diameters of shaft, 175 mm. above condyles, inclusive {	Anteroposterior	43	×
	Transverse	65	×

Measurements of Tibia.

Total length (with spine)	325	415	
Proximal diameters. {	Anteroposterior (with crest)	125	115
	Transverse, about	115	110
Least diameters of shaft {	Anteroposterior	47	60
	Transverse	55	55
Distal diameters. {	Anteroposterior (at middle)	53	67
	Transverse (at middle)	78	80

Four astragali and three calcanea are appropriate in size to the *M. angustigenis*. The former have the ridge dividing the cuboid and navicular facets more prominent than in either of the two large species first described, so that it forms a sharply-angulate protuberance. The cuboid facet forms a wide and less oblique triangle. The navicular facet

is quite concave. The sustentacular facet is continuous with it, and forms an oval in the longitudinal axis of the astragalus. The cruseular facet is transverse and concave. The tuber is not very robust and is moderately oblique.

Measurements.

I. CALCANEUM.

	<i>Min.</i>
Total length.....	123
" " of tuber.....	80
Width of tuber at base.....	43
" at sustentaculum.....	84
Diameters of cuboid facet {	
Vertical.....	26
Transverse.....	46

II. ASTRAGALUS.

Width {	of trochlea.....	67
	below trochlea.....	75
Length {	inner side.....	60
	outer ".....	55
Width of distal facets.....		70
" " navicular facet.....		42
" " cuboid ".....		29
Vertical diameter of navicular facet, at middle.....		40
" " of cuboid " ".....		25

The scaphoid appropriate to this species has, like the others, the facet for the magnum at the extremity of a step-like projection. The trapezium facet is but little smaller than that for the trapezoides. Both are subtriangular, while that for the magnum is antero-posteriorly oval, with straight internal and posterior borders, which unite at an angle. Area about equal to that of the trapezoid facet. The bone differs from the large scaphoid before mentioned, in having the lower facet abbreviated posteriorly by a fossa, which is small and upward-looking in the latter.

The magnum referred to the *M. angustigenis* has the anterior face wider than deep. The corresponding bone of the large species differs from it in having the metacarpal facet relatively a little wider, and in having at the lateral posterior edge of that facet a narrow anteroposterior oblique facet, apparently for an internal facet of the proximal end of the fourth metacarpal.

Measurements of Scaphoides and Magnum, in *Mm.*

I. SCAPHOIDES.

	<i>M. (?) americanus.</i>	<i>M. angustigenis.</i>		
Anteroposterior.....	90	75		
Diameters {	Vertical {	at trapezoides.....	24	23
		at magnum.....	50	45
	Transverse at middle.....	61	50	
Width of facet for magnum (transverse).....	39	27		
" " " trapezoides (anteroposterior).....	19	27		
" " " trapezium (").....	22	27		

H. MENSUM.

	<i>M. (?) proutii.</i>	<i>M. angustigenis.</i>
Diameters of face.....	Vertical	29 31
	Transverse.....	54 50
Diameters scaphoid facet ..	Anteroposterior	75 60
	Transverse (in front)	36 28
Diameters metacarpal facet ¹	Anteroposterior	50 45
	Transverse (at front).....	48 42

This species appears to be nearest the *M. coloradoensis*, Leidy, which is only known from a nasal portion of the skull, with the attached horns. The form of the coossified nasals is different in the two specimens, the *M. angustigenis* having at the extremity an oblique truncation on each side of a median notch, approximately as in *Symborodon trigonoceras* (*Menops varians*, Marsh), while those of the *M. coloradoensis* have a gradually acuminate outline. The value of this character is uncertain, but I incline to think it important. What is more important is that the nasals in *M. angustigenis* have a deep longitudinal concavity below, bounded on each side by a strong marginal rib, both which appear to be wanting in the *M. coloradoensis*, as Leidy says¹ the nasals "are thinner" at these borders "than elsewhere." In one point they differ from those of any of the species of this genus or of *Symborodon* which I have seen. The lateral antrum is divided anteriorly into two fossae by a vertical column, which materially strengthens the parts at the anterior base of the horn-core.

MENODUS SELWYNIANUS, Cope.

American Naturalist, 1889, p. 628.

This species is founded on the free parts of the coossified nasal bones of a single individual of smaller size than any of those already described in this report. These parts differ from those of any species known to me.

The nasals, which are completely coossified, are prominent, narrowed and vaulted. The lateral borders are nearly parallel, and the extremity is rounded. Owing to their great thickness, their profile descends abruptly at the extremity, and the surface is roughened as though for the attachment of thickened integument, a small dermal horn, or enlarged muscles. From this tuberosity the surface descends steeply to a thin border. A short distance posterior to the extremity the lateral margins are decurved, forming lateral walls to a deep, longitudinal, median, gutter-like concavity. This is deeper than in any other species. The horns are broken off, and the upper surface of the nasal bones is so little recurved that it is evident that they are not only small, but more posteriorly placed than in the other species.

Measurements of Nasal Bones.

	<i>M.</i>
Length of fragment above.....	130
" of nasal border.....	70
Width at nareal notch.....	80
" near extremity.....	65
Depth at apical tuberosity.....	25

¹ Report U. S. Geol. Surv. Terrs., I, p. 240, plate i, figs 2, 3; ii, fig. 2.

A humerus, scaphoid and astragalus may be referable to this species. They do not differ from those of *M. angustigenis*, except in their smaller dimensions.

It is dedicated to Dr. A. R. C. Selwyn, Director of the Geological Survey of the Dominion of Canada, and formerly of the Geological Survey of Victoria, Australia.

MENODUS SYCERAS, Cope.

American Naturalist, 1889, p. 628.

This species is represented by three pairs of coossified nasal bones, one of which supports a horn, in the collection, which I cannot refer to any known species. One of them is smaller and less robust than the other, and as a trace of the median suture still remains, probably indicates a younger animal.

These nasal bones are flat and not decurved anteroposteriorly, nor vaulted transversely. Their free portion is shorter than wide. Accordingly the lateral margins soon begin to thicken downwards. The inferior surface is divided into two grooves by a median, longitudinal, obtuse ridge, which is not found in the *M. selwynianus*. The free border is regularly rounded, with a median notch, and its edge is subacute. The nasal bones are considerably decurved, making an angle with the frontal surface, from which their plane is separated by an obtuse angle which connects the bases of the horns. The bases of the horns are near together, while the horns themselves are erect and moderately divergent. They are of moderate length, and are like those of *M. coloradensis* and *M. angustigenis*, with a longitudinally oval section, flattened at the base on the external side. The apex is sub-round in section.

This species differs from the *M. angustigenis* in the shorter decurved nasal bones, separated by a transverse angle from the frontal plane. The horns, though similar, are much closer together. It differs in the same characters from the *M. coloradensis*.

Measurements.

No. 1.

	<i>M.</i>
Width of nasals at base of horns.....	116
Length " from " "	70
Diameters of bases of horns {	94
	67
Width between external bases of horns	160
Elevation of horn from nasal border	160

No. 2.

Length of fragment above.....	120
Width at 30 mm. from extremity.....	80
Thickness " " "	25

No. 3.

Width at 53 mm. from end.....	80
Probable depth of nasal notch.....	60

This species belongs to the intermediate group of *Menodus*, as defined by Messrs. Scott and Osborn, in which both the nasal bones and the horns are of moderate length. They place here *M. proutii*, Leidy, and *M. tithoceras*, S. & O. From both of these *M.*

syceus differs in the angulation of the profile between the horns, and in the very close position of the horns. The entire width of the muzzle at the horns is not greater than the elevation of the horns themselves.

I may add that the bones of the skeleton described under the head of the *M. angustigenis* may well belong to the present species, or, *vice versa*, the horns and muzzle here described may belong to the species represented by the lower jaws, to which the name *M. angustigenis* has been given. In that case the name *M. syceus* becomes a synonym, and the front and nasal bones described under the *M. angustigenis* must be referred to a different species; perhaps the *M. coloradoensis*.

CÆNOPUS, Cope.

American Naturalist, 1887, pp. 925, 1060.

CÆNOPUS OCCIDENTALIS, Leidy.

Cope l. c., *Rhinoceros occidentalis*, Leidy, Proceeds. Acad. Philadel., 1850, p. 119; 1851, p. 276; Ancient Fauna of Nebraska, 1853, p. 81, plates xii, xiii; *Aceratherium occidentale*, Leidy, Proceeds. Acad. Philadel., 1854, p. 157; Extinct Mammalia Dakota and Nebraska, p. 220, pl. xxi, fig. 31; xx, xxiii, 1-3.

The only specimen from the Cypress Hills which represents this species is a fragment of a right mandibular ramus supporting a damaged last molar tooth. The dimensions of the parts are as follows:—

		<i>Mm.</i>
Diameters of M. III. {	Anteroposterior	38
	Transverse	26
Width of ramus at base of coronoid		33

CÆNOPUS MITIS, Cope.

American Naturalist, 1887, p. 925; *Aceratherium mite*, Cope, Annual Report U. S. Geol. Survey Terrs., 1873 (1875), p. 493.

Fragments of mandibular rami of two individuals represent this species. The measurements are as follow:—

Measurements.

		<i>Mm.</i>
No. 1.		
Diameters of base of M. III. {	Anteroposterior	25
	Transverse	16.5
Depth of ramus M. III.		42
Width of ramus at base of coronoid		20
No. 2.		
Diameters of M. I. {	Anteroposterior	21
	Transverse	19
Depth of ramus at M. I.		43

CÆNOPUS PUMILUS, Cope.

Aceratherium pumilus, Cope, American Naturalist, 1885, p. 103 (name only); Annual Report G. & N. H. Survey, Canada, 1885, App. C, p. 83.

Portions of mandibles of two individuals represent this, the smallest of the Rhinocerotidae. One of the rami possesses the alveoli of the large recumbent canine teeth, indicating

that the species is not a Hyracodon. The molar teeth are unfortunately broken away. The other ramus supports the second premolar, the last deciduous molar, with the first two permanent true molars.

The anterior (? fourth) premolar has a single large root, with a deep groove on the external side. In the true molars the V-shaped crests are fully developed, and there is a low cross-crest at the anterior border of the crown. There is no complete cingulum, but short sections opposite the valleys on both the internal and external bases of the crown, on the external side near the front, and at the posterior base. The measurements show how much smaller this species is than the *C. mitis*, and that it does not exceed the *Hyracodon nebrascensis*.

Measurements.

No. 1.		M.
Width between bases of P. m. i.033
Length of base of anterior three premolars042
Depth of ramus at diastema036
" " " third premolar042
No. 2.		
Length of molars i and ii039
Diameter M. ii.	{ Anteroposterior	.020
	{ Transverse	.012
Depth of ramus at front of M. ii033

ANCHITHERIUM, Kaup.

ANCHITHERIUM WESTONII, Cope.

American Naturalist, 1889, p. 153.

This species is represented by a single superior molar and two inferior molars, the latter in place in a part of the mandible. The teeth are smaller than those of the *A. lairdii*, from which they also differ in their greater transverse as compared with their anteroposterior diameters. The intermediate tubercle of the posterior cross-crest is more distinct than that of the anterior, and the posterior intermediate cingular cusp, so prominent in the *A. lairdii*, is here wanting. The posterior cingulum continues round the internal base of the posterior internal cusp. Diameters of superior molar: Transverse, 13.5 mm.; anteroposterior, 10 mm. Diameters of inferior molar: Transverse, 8 mm.; anteroposterior, 10.5 mm. This species, interesting for its primitive character in the absence of the posterior cingular cusp, is dedicated to Mr. T. C. Weston, the most successful collector in the region from which these fossils were obtained.

ARTIODACTYLA.

ELOTHERIUM, Pomel.

Eutelodon, Aymard.

ELOTHERIUM ARCTATUM, Cope.

American Naturalist, 1889, p. 629. *Elothorium mertonii*, Leidy, Cope, Report G. & N. H. Surv., Canada, 1885, p. 154.

Established on a left mandibular ramus which supports all the molar teeth, but

lacks the canine and incisors, and its inferior border anterior to the first premolar. From the last circumstance it results that the presence or character of the mandibular tuberosities cannot be ascertained. Some separate molar teeth of the same species I have recorded as above as the *E. mortoni*, to which the *E. arctatum* is nearly allied.

The present species is a little inferior in dimensions to the *E. mortoni* and *E. magnum*, and it differs from the former in the absence of diastemata between the premolar teeth, except a very short one between the third and fourth. The latter is separated by a very short space from the canine. The second premolar is larger than the first, and the third and fourth are abruptly smaller than either. The latter has but one, a compressed, laterally-grooved root. The crowns of all the premolars are compressed, and have fore and aft cutting edges. The true molars are peculiar in the elevation of the anterior part of the crown of two tubercles, above the posterior part. The latter consists of three tubercles, the third or posterior median tubercle being better developed than in the *E. mortoni*, especially on the last molar. As the jaw is adherent to a block of conglomerate rock, only the internal sides of the teeth are visible. These are without cingula, and the enamel of the molars is smooth, while that of the premolars is wrinkled.

A wide space separates the last molar from the base of the coronoid process, which is low. The condyle is in line with the crowns of the molars. The ramus is compressed and not very deep.

Measurements.

	<i>Min.</i>
Length from condyle to edge of canine alveolus.....	295
" " " " " last molar.....	125
" of true molar series.....	67
Diameters of m. iii.	23
{ Longitudinal.....	23
{ Transverse.....	13
Diameters of M. i.	22
{ Longitudinal.....	22
{ Transverse.....	13
Diameters of P. m. i.	20
{ Vertical.....	20
{ Longitudinal at base.....	28
Diameters of P. m. ii.	21
{ Vertical.....	21
{ Longitudinal at base.....	28
Diameters of P. m. iv.	13
{ Vertical.....	13
{ Longitudinal at base.....	13
Elevation of coronoid above border.....	57
Depth of ramus at m. i.	55

This species is interesting as presenting a more primitive character than the American species so far known, in the closely-placed premolar teeth. The form of the true molars is also more primitive in the elevation of the anterior pair of cusps, almost resembling some species of *Miocænus* in this respect. The large fifth tubercle has a similar significance. The type specimen was found by Mr. T. C. Weston near the head of the Swift Current Creek in August, 1889.

? OREODON, Leidy.

The scarcity of Oreodontidae in the collections from the Cypress Hills is a matter of surprise. An inferior fourth (first) premolar is the only indication of their presence. The genus is not determinable.

HYPERTRAGULUS, Cope.

Bulletin U. S. Geol. Survey Terrs., 1, 1874, p. 26; Annual Report do., 1873 (1874), p. 502; American Naturalist, 1889, p. 111, pl. vi.

HYPERTRAGULUS TRANSVERSUS, Cope.

American Naturalist, 1889, p. 154.

Indicated by two superior molar teeth of old individuals. They are of nearly twice the linear dimensions of the only known species, *H. calcaratus*, Cope. The external cusps are subconical, and the external rib which separates them in the *Leptomeryx* is wanting here. Anterior cingular cusp small. The anterior horn of the posterior internal crescent enters the notch between the external cusps, but does not fuse with either of them. Slight cingula on the anterior and posterior sides of the internal lobes, which do not pass round their internal sides. No external cingulum. Diameters: anteroposterior, 12 mm.; transverse (at base), 15 mm. Crown very brachyodont.

LEPTOMERYX, Leidy.

Proceeds. Academy Philada., 1853, p. 394; 1857, p. 89; Extinct Mammalia Dakota and Nebraska, 1869, p. 165; Cope, Proceeds. Amer. Philos. Soc., 1884, p. 23; *Ibid.*, 1887, p. 389.

LEPTOMERYX ESULCATUS, Cope.

American Naturalist, 1889, p. 154.

A single superior molar indicates this species, which is of about the dimensions of the *L. evansii*. It differs distinctly from this *Tragulid* in the greater convexity of the external face of the external cusps, and the absence of the sulci which define an external rib of that surface in the *L. evansii*. The rib which defines the external faces of the cusps from each other is present. Anterior external cingular cusp small, continuous with anterior cingulum. No internal nor external cingulum. Diameters of crown: Anteroposterior, 6.5 mm.; transverse, 7.5 mm.

LEPTOMERYX MAMMIFER, Cope.

Report G. & N. H. Survey, Canada, 1885, App. C. p. 84; American Naturalist, 1889, p. 154.

This species is represented by a fragment of the mandible which supports the last two molars. Four superior molars add to the characters already derived from mandibular teeth, but their reference to the same species is provisional, as they were found separately. A tarsal cannon bone in the collection may belong to the same species. It is not certain that it belongs to a *Leptomeryx*, but I leave it there until further information enables me to make a final determination.

The crowns of the inferior molars are not prismatic, nor are they brachyodont. The crown is well distinguished, and expands but little. The sections of the internal columns are lenticular, while the external are crescentic. There are no basal columns or cingula between the latter. In the second true molar, the horns of the anterior internal crescent join the anterior external crescent early on wearing, while the junction comes later in the case of the two posterior columns. In the third true molar the anterior horn of the posterior crescent does not reach the posterior external column, but only touches the anterior internal column. In the same way, the posterior horn does not reach the external

column, but is separated from it by a distinct mammary tubercle or short column, which has an anteroposteriorly short oval section. The heel of this tooth is broken off, but it was small, judging by the fragments of its base.

The peculiar column intercalated between the heel and the posterior internal column distinguishes this species from all the selodont Artiodactyla known to me.

The enamel is slightly wrinkled. The half-worn condition of the crowns show that the animal was adult.

<i>Measurements.</i>		<i>Min.</i>
Diameters of M. ii.	Anteroposterior0098
	Transverse0070
	Vertical (of enamel).....	.0015

In the superior molars the median and anterior external cingular cusps are large and obtusely subconical. The anterior external cusp has a very strong median external rib, while the posterior has a very weak one. The anterior horns of the internal crescents are much produced; the posterior but little. The cingula are slight, and are not continued round the internal base. Diameter of superior molar: anteroposterior, 11 mm.; transverse, 11.5 mm.

LEPTOMERYX SEMICINCTUS, Cope.

American Naturalist, 1889, p. 154.

A large species possessing twice the linear dimensions of the *L. evansii* in the superior molar teeth, is represented by three of the teeth designated. In these the external crescents are more compressed and less conical than the two species above described, resembling more nearly those of the *L. evansii*. The posterior has a weak vertical rib; the anterior a strong one. The external cingular cusps are thoroughly fused with the external crescents, forming their anterior horns. The anterior horns of the internal crescents are a little more produced than the posterior. No external or posterior cingulum; a much interrupted anterior cingulum, which is continued round the internal base of the anterior crescent, and which is further continued on the anterior side of the internal base of the posterior crescent. Enamel finely wrinkled. Diameters: anteroposterior, 11 mm.; transverse, at base, 15 mm.

EXPLANATION OF PLATES.

PLATE I.

Bones of fishes and tortoises, natural size.

- Fig. 1. *Amia whitearsiana*, Cope, anterior vertebra, articular face; *a* from above; *b* from below.
 Fig. 2. *Amia macrospandyla*, Cope, anterior vertebra, articular face; *a* from above; *b* from below.
 Fig. 3. *Rhincastrus rheus*, Cope, dorsal vertebra, articular face; *a* from above; *b* the side.
 Fig. 4. *Amiurus cancellatus*, Cope, dorsal vertebra, articular face; *a* from above; *b* from below.
 Fig. 5. *Amiurus cancellatus*? caudal vertebra, articular face; *a* from above; *b* from side.
 Fig. 6. *Amiurus macconnelli*, Cope, dorsal vertebra, articular face; *a* from below; *b* from side.
 Fig. 7. *Amiurus macconnelli*? vertebra, articular face; *a* from side.
 Fig. 8. *Triomys lucopotamicus*, Cope, costal bone from above; from White Buttes, Dakota, U. S.
 Fig. 9. *Triomys lucopotamicus*, Cope, part of costal bone from above; typical specimen.

PLATE II.

Right mandibular ramus (lacking posterior part) of the *Hemipsetodon grandis*, Cope, natural size; *a* superior view.

PLATE III.

Hemipsetodon grandis, Cope, femora, two-thirds natural size.

- Fig. 1. Anterior view; trochanter restored from smaller specimen.
 Fig. 2. Posterior view; " " " " "
 Fig. 3. Proximal extremity, end view; taken from the second and smaller specimen.
 Fig. 4. Distal end left side; from specimen No. 1.

PLATE IV.

- Fig. 1. *Chadivobrium babatuna*, Cope, symphyseal part of mandible, the right alveolar ridge wanting, left side; two-thirds natural size; *a* symphysis, posterior view; *b* from above.
 Fig. 2. *Canoqus mitis*, Cope, part of mandibular ramus, with crowns of third and fourth premolars, two-thirds natural size, from above.
 Fig. 3. *Canoqus mitis*, Cope, symphysis and part of left ramus mandibuli, from above; *a* from left side.
 Fig. 4. *Canoqus pumilus*, Cope, part of right mandibular ramus from above, natural size.

PLATE V.

- Fig. 1. *Menodus angustigenis*, Cope, posterior four superior molars, three-eighths natural size, from below.
 Fig. 2. *Menodus angustigenis*, Cope, mandibular ramus of another individual (also figured on plate viii), three-eighths natural size.
 Fig. 3. *Menodus scleroticus*, Cope, nasal bones from above, three-eighths natural size; *a* the left side; *b* from below.

PLATE VI.

- Fig. 1. *Menodus? americanus*, Leidy? female; nasal and part of frontal bone, from above; one-half natural size.
 Fig. 2. *Menodus angustigenis*, Cope, left ramus mandibuli, external view; three-eighths natural size; *a* internal side.

PLATE VII.

- Fig. 1. *Menodus angustigenis*, Cope, symphyseal part of right mandibular ramus, from the right side, two-thirds natural size; *a* from below; *b* from front. Type of the species.
 Fig. 2. *Menodus angustigenis?* Cope, nasal bones and right horn, from the front; one-half natural size. Type of *M. syceras*, Cope.

PLATE VIII.

- Fig. 1. *Menodus? americanus*, Leidy? female, nasal bones, from the front; one-half natural size.
 Fig. 2. *Menodus? americanus*, specimen represented in fig. 1, left side; one-half natural size.
 Fig. 3. Same as fig. 2, outline of section of base of left horn.
 Fig. 4. *Menodus angustigenis*, Cope, (*M. syceras*) nasal bones and horn, right side; one-half natural size; external view.
 Fig. 5. Outline of section of base of horn of specimen fig. 4.

PLATE IX.

- Fig. 1. *Menodus ? proutii*, O. N. & E. horn of right side; three-eighths natural size; external view; a section of horn at base.
 Fig. 2. *Menodus americanus*, Leidy, horn of left side; external view; three-eighths natural size; a base of same with adjacent extensions, from below.
 Fig. 3. *Menodus ? americanus*, head of tibia, proximal view, figured at No. 1, plate xii.

PLATE X.

- Fig. 1. *Menodus*, sp. humerus; two-fifths natural size; posterior view.
 Fig. 2. *M. nodus ? americanus*, Leidy, scaphoid bone; two-fifths natural size, from front; a proximal face; b distal face.
 Fig. 3. *Menodus ? proutii*, O. N. & E. tibia; proximal view of head; two-fifths natural size. From tibia figured at No. 2, plate xii.

PLATE XI.

Bones of *Menodus*, three-eighths natural size.

- Fig. 1. *Menodus ? angustigenis*, Cope, femur, from front; rotular face injured; a posterior view; b proximal view of head.
 Fig. 2. *Menodus*, sp. humerus, figured on plate x.; head, proximal view; a condyles, distal view.

PLATE XII.

Tibiae of *Menodus*, three-eighths natural size.

- Fig. 1. *Menodus ? americanus*, Leidy, front view; a posterior view; c distal end.
 Fig. 2. *Menodus ? proutii*, O. N. & E. front view; a posterior view.
 Fig. 3c. *Menodus ? proutii*, second individual; distal extremity.
 Fig. 4. *Menodus ? angustigenis*, Cope, tibia, from front; b proximal view of the same; c distal view of the same, another specimen.

PLATE XIII.

Bones of the feet of *Menodus*, three-eighths natural size.


- Fig. 1. *Menodus ? angustigenis*, Cope, scaphoid, from front; a proximal side; b inferior side.
 Fig. 2. *Menodus*, least species; scaphoid bone, from front; a proximal view; b distal view.
 Fig. 3. *Menodus ? proutii*, os magnum from above; a from front.
 Fig. 4. *Menodus ? angustigenis*, os magnum from above; a from front.
 Fig. 5. *Menodus americanus*, Leidy, calcaneum from above; sustentaculum wanting; a distal extremity.
 Fig. 6. *Menodus ? angustigenis*, calcaneum from above; a distal view.
 Fig. 7. *Menodus ? americanus*, astragalus from above; a distal view.
 Fig. 8. *Menodus angustigenis*, astragalus from above; a distal view.
 Fig. 9. *Menodus*, least species; astragalus from above; a distal view.

PLATE XIV.

Chiefly teeth of *Diphthera*, natural size, except figure 3, which is two-thirds natural size.

- Fig. 1. *Anechtherium westoni*, Cope, superior molar from below; external wall restored.
 Fig. 2. *Anechtherium westoni*, fragment of lower jaw, with two molar teeth, from above; a do, external side.
 Fig. 3. *Blatherium arctatum*, Cope, left ramus of mandible, lacking the anterior extremity, and part of inferior border, internal view; a dentition, viewed from above.
 Fig. 4. *Hypertrogulus transiens*, Cope, worn superior molar tooth, from below; a external view.
 Fig. 5. *Leptomeryx colubus*, Cope, superior molar, † natural size; a natural size; external wall of anterior crescent restored.
 Fig. 6. *Leptomeryx mammifer*, Cope, superior molar; a external view.
 Fig. 7. *Leptomeryx mammifer*, Cope, portion of mandible, with second and parts of the first and third true molars, from above; a from external side. Typical specimen.
 Fig. 8. *Leptomeryx semicinctus*, Cope, superior molar, from above; a from external side.
 Fig. 9. *Petrologus targulus*, Cope, left mandibular ramus, external view; a superior view.






PISCES and TESTUDINATA.




III MIPSA LODON GRANDIS, COOP






HEMIPSALODON GRANDIS, Cope






ANCYLOPODA and PERISSODACTYLA.











PLATE VI.


GEOLOGICAL SURVEY OF CANADA






MENODIUS






1 MENODUS ANGUSTIGENIS. C. P. M. SYCLERAS. Cope.





1-4 MENODUS AMERICANUS, Ledy 5 M. SYCLERAS, Cope






MENODUS.






MENODUS.






MENODUS.








MENODUS.





PERISSODACTYLA ARTIODACTYLA, and GLIRES

