

**CIHM
Microfiche
Series
(Monographs)**

**ICMH
Collection de
microfiches
(monographies)**



Canadian Institute for Historical Microreproductions / Institut canadien de microreproductions historiques

© 1998

The copy filmed here has been reproduced thanks to the generosity of:



McGill University
Health Sciences Library
Montreal

The images appearing here are the best quality possible considering the condition and legibility of the original copy and in keeping with the filming contract specifications.

Original copies in printed paper covers are filmed beginning with the front cover and ending on the last page with a printed or illustrated impression, or the back cover when appropriate. All other original copies are filmed beginning on the first page with a printed or illustrated impression, and ending on the last page with a printed or illustrated impression.

The last recorded frame on each microfiche shall contain the symbol \rightarrow (meaning "CONTINUED"), or the symbol ∇ (meaning "END"), whichever applies.

Maps, plates, charts, etc., may be filmed at different reduction ratios. Those too large to be entirely included in one exposure are filmed beginning in the upper left hand corner, left to right and top to bottom, as many frames as required. The following diagrams illustrate the method:



L'exemplaire filmé fut reproduit grâce à la générosité de:

McGill University
Health Sciences Library
Montreal

Les images suivantes ont été reproduites avec le plus grand soin, compte tenu de la condition et de la netteté de l'exemplaire filmé, et en conformité avec les conditions du contrat de filmage.


Les exemplaires originaux dont la couverture en papier est imprimée sont filmés en commençant par le premier plat et en terminant soit par la dernière page qui comporte une empreinte d'impression ou d'illustration, soit par le second plat, selon le cas. Tous les autres exemplaires originaux sont filmés en commençant par la première page qui comporte une empreinte d'impression ou d'illustration et en terminant par la dernière page qui comporte une telle empreinte.

Un des symboles suivants apparaîtra sur la dernière image de chaque microfiche, selon le cas: le symbole \rightarrow signifie "A SUIVRE", le symbole ∇ signifie "FIN".

Les cartes, planches, tableaux, etc., peuvent être filmés à des taux de réduction différents. Lorsque le document est trop grand pour être reproduit en un seul cliché, il est filmé à partir de l'angle supérieur gauche, de gauche à droite, et de haut en bas, en prenant le nombre d'images nécessaire. Les diagrammes suivants illustrent la méthode.


MICROCOPY RESOLUTION TEST CHART

(ANSI and ISO TEST CHART No. 2)



APPLIED IMAGE Inc

1653 East Main Street
Rochester, New York 14609 USA
(716) 482 - 0300 - Phone
(716) 288 - 5989 - Fax



**THE ORGANS
OF INTERNAL SECRETION**

27

.

.

100

THE ORGANS OF INTERNAL SECRETION

THEIR DISEASES AND
THERAPEUTIC APPLICATION

A BOOK FOR GENERAL PRACTITIONERS

BY

IVO GEIKIE COBB, M.D., M.R.C.S.

LATE ASSISTANT TO OUT-PATIENT PHYSICIAN, THE MIDDLESEX
HOSPITAL



TORONTO

J. F. HARTZ CO., LTD.

1917

Printed in England

PREFACE

THIS book is founded upon articles which appeared in the *Medical Press and Circular* during the summer and autumn of 1916. The author's object in writing these articles was to lay before the busy practitioner the important points in the study of the Endocrine Glands. At the same time he was anxious to make them as complete as possible, and yet keep them within the limits of such publications.

After the appearance of these articles, he received requests for reprints, and as they had not been reprinted, and the interest in them appeared to warrant it, he decided to publish them in book form. The present volume comprises these articles, with slight alterations.

Hormone-therapy is already established as a recognized therapeutic agent, yet as the books which deal with the ductless glands and their secretions are exhaustive studies including the results of laboratory research in detail, and require far more time to read than the general practitioner has at his disposal, it is difficult for him to glean the salient facts from these lengthy works. It seemed, therefore, to the author that a small book which contained an account of the

diseases and therapeutic application of extracts of these glands might prove useful.

In publishing this small volume, however, the author wishes to state that it makes no claim to be considered an exhaustive and complete account of the endocrine glands, neither can it be considered as a comprehensive therapeutic guide to the administration of the organic extracts. Rather does it aim at being a guide to other practitioners as to the rôle which the ductless glands play in promoting bodily health; while it endeavours to point out those morbid states of health in which organic extracts may be utilized with success.

Among the glands possessing an internal secretion, the thyroid is perhaps the one which has attracted most attention, partly on account of the well-recognized disorders which arise in connection with it, and partly because, of all the endocrine glands, the thyroid has been most utilized therapeutically. In order to emphasize the signs and symptoms associated with morbid conditions of this gland, the author has devoted three chapters to the consideration of Exophthalmic Goitre and Thyroid Deficiency. The last of these three chapters contains references to the administration of thyroid extract; while this subject is again referred to in Chapter X.

A chapter has been devoted to the Pituitary Gland, and the subject of pituitary-therapy has been briefly reviewed. In Chapter VI. the Adrenal Glands are described, and the clinical conditions of hyperadrenia

and hypoadrenia are discussed. Chapter VII. deals with the Pancreas, and describes its structure, physiological functions, and relation to glycosuria, and concludes by referring to the therapeutic possibilities of preparations of this gland.

The subject of the Internal Secretions of the Sexual Organs, dealt with in Chapter VIII., has been compressed into as small a space as was possible; a full account would have filled a large volume. Nevertheless, this chapter reviews the outstanding features, suggests ways of administering extracts of the genital glands, and describes morbid conditions in which they may be helpful.

The Internal Secretions of Digestion form the subject of Chapter IX.; and here the author has endeavoured to lay emphasis upon the therapeutic aspect, as he believes that in the near future this branch of organo-therapy will find a very wide field of utility, and will succeed in alleviating many morbid conditions which have hitherto proved resistant to treatment.

The Therapeutic Application of Hormones is summarized in Chapter X. This chapter has been enlarged since its appearance in the *Medical Press and Circular*, and it endeavours to epitomize the subject for those readers who are mainly concerned with the therapeutic aspect of this subject. It necessarily summarizes the conclusion of each chapter, but in addition references will be found to preparations and doses not included in the previous chapters.

Chapter XI. is devoted to a survey of the whole subject, and deals with the present position of hormone-therapy.

As the therapeutic aspect of this subject was dealt with at the end of each article, it has been decided to leave this arrangement unaltered. Likewise, the references have been left in their original positions; but a bibliography has been included at the end of Chapter XI., so that those desirous of a fuller account of the ductless glands and organo-therapy will be able to refer to the works enumerated there.

In conclusion, the author desires to thank the editor and proprietors of the *Medical Press and Circular* for permission to reproduce these articles in book form.

IVO GEIKIE COBB.

SEYMOUR STREET, W.,
November, 1916.

CONTENTS

CHAPTER	PAGE
PREFACE	v
INTRODUCTION	1
I. THE THYROID AND PARATHYROID GLANDS	16
<i>dissect</i> II. EXOPHTHALMIC GOITRE (HYPERTHYROIDISM)	31
III. THYROID DEFICIENCY	54
IV. THYROID DEFICIENCY (<i>continued</i>)	70
V. THE PITUITARY BODY	94
VI. THE ADRENAL GLANDS	114
VII. THE PANCREAS	131
VIII. THE SEXUAL ORGANS AND THEIR INTERNAL SECRETIONS	147
IX. THE INTERNAL SECRETIONS OF DIGESTION	163
X. THE THERAPEUTIC APPLICATION OF HORMONES	183
XI. CONCLUSIONS AND BIBLIOGRAPHY	213
INDEX	223



THE ORGANS OF INTERNAL SECRETION

INTRODUCTION

THE place which those glands possessing internal secretions now occupy in the practice of medicine cannot be over-estimated. Although it is only in recent years that their importance has been understood, every day brings to our knowledge fresh evidence of their vital influences upon the general bodily and mental health. When it was discovered that exophthalmic goitre owed its existence to an over-action, associated with hypersecretion, of the thyroid gland, the first milestone had been passed in the path which led us to the discovery of the important part which the hormones play in our lives.

What we may call the grosser lesions due to disturbances in the normal ratio between the various internal secretory glands are nowadays matters of almost popular knowledge. What we desire to emphasize in this book are the smaller signs and symptoms which, to the eye trained to observe, show minor disturbances, either of one or more glands or of the

2 THE ORGANS OF INTERNAL SECRETION

balance between these glands. Many of these details, trifling in themselves, have been proved to be sufficiently characteristic to justify medical science in including them among the constant features which owe their origin to the ductless glands. When the pioneer work was being done on this subject, the evidences of the slighter disturbances of the endocrine glands were regarded by many observers as too fanciful to merit serious consideration. Thus, when Levi and Rothschild first pointed to the "eyebrow" sign as indicative of deficient thyroidism, it was hard for many students of these subjects to convince themselves that this sign was of any value whatsoever.

Nevertheless, this and other manifestations of equally slight nature are now admitted to be of the greatest value in diagnosing deficient thyroid secretion. Many of the signs of kindred nature are so slender as to require the most minute study and the most careful observation before we can say, with any degree of probability, which particular gland is at fault. The importance of this lies in the fact that many cases of "functional disturbances," which we have been content previously to cosset with various preparations, and to class as neurotic or neurasthenic according to the depths of our ignorance, are now recognized as originating in abnormal functioning of the endocrine glands. This much is now generally known and universally admitted. Only relatively few observers recognize that this is not the *fons et origo mali*, but only one stage on the journey.

To take a concrete instance, let us suppose that a patient exhibits signs that the thyroid is not functioning adequately. The patient exhibits many of the well-known signs pointing to this deficiency. Most of us are well content to leave it at that, and even to spend time (which should be occupied in delving still farther and asking ourselves why such a deficiency is present) on self-adulation at our extraordinary deductive powers. It is certainly true that we can benefit our patients by the exhibition of one or other of the organo-therapeutic products, but we shall do so to a much larger extent if we realize one or two simple facts.

When advising a patient to take, let us say, thyroid extract for a time, we are constantly asked by the patient the *rationale* of such a prescription. Having explained that we have reason to believe that this gland is not supplying an adequate amount of secretion, on more than one occasion we have been asked by the logical patient, "Have I got to continue this medicine for the rest of my life?" or, "Will the need for its administration be overcome in due time?" This is an important matter, because it makes us think for ourselves, and not blindly prescribe drugs for indefinite times without reasoning as to their method and length of administration.

I think that we are justified in considering for a few moments what underlies the deficiency in secretion. It is highly improbable that one or other of the endocrine glands would suddenly and without any stimulus

4 THE ORGANS OF INTERNAL SECRETION

refrain from supplying, or determine to over-supply, its valuable contents to the blood-stream. It is much more probable that some cause, be it mental or bodily, has determined this upset, and it behoves us to realize that this is the case, and not to consider our diagnosis completed when we have made up our mind that such and such a gland is defective or over-active.

In his book on "Intestinal Stasis," Lane states that one of the effects of intestinal intoxication is atrophy of the thyroid. Here we have a definite attempt to go to the root of the trouble, although, unfortunately, there is little evidence at present for this statement. For the present, at any rate, let us assume that this is one of the causes which underlie submyxœdema. What others may there possibly be? Among the causes of diabetes which find their place in most current textbooks on medicine is worry and anxiety. Now, this is at once assuming that mental causes may upset a bodily function or functions, and thereby disorganize metabolism. And the writer would be the last to wish to deny this. If, therefore, we may assent to this detail of etiology with regard to diabetes, why should we not include such a cause when we are discussing disorganization of the endocrinic glands? Many cases which we can call to mind at the moment afford us the strongest possible support for such a theory. Prolonged anxiety, business worry, a sudden shock, are said to be capable of producing diabetes, just as puncture of the floor of the fourth ventricle is, experimentally, capable

of doing the same. It would seem at least equally probable that the same causes can, and do, upset the mechanism which governs the balance between the ductless glands.

In the severe cases of this nature (*e.g.*, Graves' disease), this fact is sufficiently recognized. But is it, has it been, when we come to consider slighter derangements of these important glands? It is obvious that the entire study of this subject is of too recent a date to have made the minute clinical study of the ultimate cause a feasible proposition. Nevertheless, we now come to the time when our patients ask us the why and the wherefore of such prescribing.

We have suggested two possible causes which may underlie the disorganization of the endocrinic system—namely, intestinal toxæmia (as suggested by Lane); and mental causes, such as worry, anxiety, and mental strain of any kind. But are these the sole causes which may be held responsible? We are all familiar with the damaged health which may result from a long illness, or, alternatively, from a sudden short attack, such as influenza. The patient recovers but slowly, the strength returns not, the mind is clouded, and in countless other ways the individual shows the effects of the illness. Hitherto we have referred to such cases as post-influenzal debility, when this disease has been at the root of the trouble; or as "neurasthenia," when we could not find a cause even as tangible as influenza. Our remedies have been confined, certainly in many instances, to a change

6 THE ORGANS OF INTERNAL SECRETION

to the sea or spa, and a generous addition to the diet. But we have rarely asked ourselves what factor underlies these "delayed recoveries."

Why should not the toxins of influenza, in like manner to the toxins generated by the inhabitants of the bowel, or the adverse mental influences which exert their harmful action where mental strain is present, be capable of producing an endocrinic disorganization? Hypothetically, at any rate, such an occurrence is at least probable, and it would give us a reason for the sudden or gradual withdrawal of the internal secretion which happens to be deficient in the particular case.

From the practical standpoint, moreover, we must advance some such hypothesis as this in order to account for the train of symptoms, which, certainly in many cases, owes its origin to a disturbance of the normal ratio which exists in health between the various endocrinic glands. Again, most of us are familiar with the cases of delayed convalescence following an operation. The patient invariably presents a similar picture to that designated "post-influenzal debility." In theory, at any rate, he ought to respond to the administration of one or other of the preparations of the hormones, and in many cases he does.

On more than one occasion I have had the opportunity of putting this theory into practice. One lady consulted me some years ago for neurasthenia following a severe abdominal operation. This condition had resisted a wealth of treatment; many and

diverse remedies had been tried without relief. I hoped that I might be enabled to afford relief by the administration of an organo-therapeutic preparation. On considering her syndrome, I came to the conclusion that she might derive benefit from the exhibition of pituitary extract. In spite of the laboratory evidence, which should convince us that it is useless to give an extract of this gland by the mouth, I took my courage in both hands and prescribed it. The result surpassed even my optimistic expectations. The lady recovered her strength and health; her digestion righted itself; her functions became normal, and she regained perfect health. This result is striking, for it followed many other remedies, and the patient herself always refers to this medicine as the "magic mixture."

I mention this case as it exemplifies our hypothesis that many and diverse causes may produce a change in the normal functioning of these glands. Whether, in this particular case, it was the shock of the operation, the anæsthetic, or the changes in diet necessitated by these procedures, it is impossible to say. Another example of a similar attack of "thyroid deficiency" is the case of a lady who, having nursed her husband through a long and trying illness, which resulted fatally, consulted me for symptoms which, upon investigation, were shown to be due to deficient thyroid secretion. Upon the administration of thyroid extract, she made a capital recovery and was restored to health.

These cases both help us to answer the question so

8 THE ORGANS OF INTERNAL SECRETION

often put to us when we recommend an extract of one or other of the ductless glands. It would seem that, certainly in many instances, the administration of the extract either by the mouth or hypodermically serves to act as a stimulus to the normal secretion, so that it is unnecessary to continue artificially its administration for lengthy periods. And, again, the prescribing of the requisite extract at the right time is the "shortest cut" to health which exists.

It is even possible that many of the benefits which we all recognize to accrue from a change of climate, from a course of spa treatment, of from a sea-voyage, are really largely efficacious because they stimulate into activity the gland (or glands) which has been temporarily inhibited by the illness, operation, or other cause. In this connection we may refer to another instance where "post-influenzal debility" has yielded to organo-therapy. A lady consulted me for this condition, which had been in existence for five years, and which had resisted all treatment. It had commenced after a bad attack of influenza five years before. Her condition was much improved from the first by the administration of thyroid extract, and she made a most satisfactory recovery.

The point which seems to need emphasizing is this: x that prolonged illness, shock, mental anxiety, and many other causes, produce effects which owe their origin to a disturbance of the relation between the hormones. The indications are rarely broad. Sometimes they require the eye of a medical "detective"

before their significance is realized. But the signs and symptoms are rarely wanting if they are looked for.

Neurasthenia has been likened to influenza (inasmuch as any intangible condition has received this label), and has been dubbed the "dustbin of the neurologist." Nevertheless it is a real and concrete disease—concrete in the sense that it is not a hotch-potch of other diseases. Doubtless some of the diagnoses which have been made under this name would in reality have received another title did we but realize what was the underlying pathology. Some patients who have been called neurasthenics are in reality neurasthenics, but they are neurasthenics because they are suffering from a deficiency of hormones.

I may perhaps be allowed to mention one other case as illustrating the relation between neurasthenia and the endocrinic glands. During the course of last year I was consulted by a doctor who informed me that he was a neurasthenic, and brought me a typewritten account of his symptoms in support of this statement. I need not describe the case in detail, as the few points I mention will serve our purpose.

The patient complained that he was slow mentally, became extremely tired after comparatively small exertion, was exhausted after sexual connection, unable to concentrate for any length of time, and so on. On examination, I discovered a very slow pulse (barely fifty to the minute), a dry and rough skin,

10 THE ORGANS OF INTERNAL SECRETION

prematurely grey hair, especially over the temples (the patient was in the thirties), trophic changes in the skin appendages, and many other minor signs which I will not waste time by enumerating. Suffice it to say that the patient presented a typical picture of submyxœdema, and I advised small doses of thyroid. Some months later he advised me that he had suffered from some "extraordinary sinking feelings" when he had taken the thyroid, and in consequence had been forced to abandon it. I explained that these were most certainly due to the stimulating effect which the thyroid would produce upon the circulation, and I encouraged him to persevere with it. The interest of this case lies in the fact that all the symptoms had gradually supervened after an attack of influenza, and that their real nature had never been diagnosed.

Such a case as this will serve to show how a certain proportion of patients who have been treated, and only too frequently dismissed as incurable under some such name as neurasthenia, may be helped by the judicious administration of these extracts.

I need not offer an account of the symptoms which make up these diseases, nor is there need for me to describe in detail what is so well known about individual symptoms of deficiency in thyroid, in pituitary, or in adrenals. But one or two points have come to my notice about these conditions that I should like to mention in passing.

There can be little doubt that deficiency in thyroid secretion comes on more or less suddenly in some

cases. I recall the case of a young male subject who developed this complaint after a hazardous season on the Stock Exchange. His condition, when he came under my observation, was typical, and he made speedy progress under thyroid medication. Again, I have seen a typical attack of submyxœdema develop after one of the exanthemata; likewise excessive thyroid secretion ensue after such a disease as rheumatism.

It may be of interest to note at this place the extraordinary intolerance to tobacco which develops when the thyroid secretion is deficient. I have on several occasions observed that the patient, a heavy smoker previously, has had to abandon the fragrant weed at or about the time when his illness commenced. I have been told that even one cigarette is followed by unpleasant sensations, and I have ascertained that blood-pressure has been lowered as much as ten points after one cigarette.

These occurrences all point to the fact that, given suitable conditions, it is not a difficult matter to upset the normal ratio between the various hormones. It is necessary in these cases to study the antecedent conditions with as much care as we should when taking the previous history of, let us say, a case of tuberculosis. For in these patients we can often discover some occurrence which may well have some bearing upon the etiology, and may give us valuable information both as to the actual cause and as to the particular gland at fault.

12 THE ORGANS OF INTERNAL SECRETION

Unfortunately, the therapeutics of the other glands are scarcely in such a satisfactory state as that of the thyroid gland, and we do not, in practice, obtain results as striking as those which so often ensue from the administration of thyroid. But, before leaving the subject, we may say a few words as to the prescribing of the adrenals.

The extract of the adrenals may be given in the form of dry extract, and it is often of great benefit to those patients who are weakly, debilitated, with a low blood-pressure and constant fatigue. These are the cases of neurasthenia in which this extract should be tried. Again, the preparations of one or other part of the pituitary gland are in some of these cases more beneficial; while it is sometimes of service to use what Leonard Williams calls a "mitrailleuse"—*i.e.*, a preparation containing the extracts of many glands. Such a one is Hormotone, and it is claimed that its exhibition is followed by marked benefit in many indefinite conditions.

The method of treating disease by means of extracts of the endocrine glands is, relatively, still in its infancy, so we must not be hypercritical at that part of this medication which works without the support of the laboratory. But, as has been admitted elsewhere by a physiological chemist, clinical experience often is at variance with laboratory results, and clinical results are not always in the wrong. However much we may condemn the indiscriminate and speculative use of these extracts without adequate reasoning, if

we abide by laboratory results, and never test these by practical endeavour, we are liable to remain with little added knowledge on these subjects as the years roll on.

The practitioner must perforce use his eyes before he prescribes thyroid extract; he must be familiar with the small signs which go to make up the picture of deficient or excessive action of this gland; and he must not hesitate to prescribe this substance, although the signs are slight. He must be familiar with the diagnostic features, and he must be equally *au fait* with what is now known about treatment. He must choose a carefully standardized preparation; be careful that his preparation is new, and not several months older than when the local chemist purchased it from the wholesale house; and, finally, he must understand that the dose is a matter deserving the closest attention. Thyroid, to quote one example, is not a drug to use from 3 to 10 grains, but in fractions of a grain—certainly to commence with. Had it not been for the fact that it has been utilized to reduce weight, the probability is that it would not have obtained that popularity which it now possesses in the lay mind. This has made its administration a matter of danger, especially when we consider that the dose, or rather the initial dose, is often far too large.

Convalescence is frequently accompanied by deficiency in the thyroid gland, and its administration in minimal doses is of very real help. During the months which follow a serious illness, such as pneumonia,

14 THE ORGANS OF INTERNAL SECRETION

the administration of a "mitrailleuse" is frequently indicated. The tendency to obesity, or alternatively to undue loss of flesh, which is usual after severe illnesses, would point to an upset in the hormone balance; and this may be remedied by a careful study of the symptom-complex and the prescription of a suitable organo-therapeutic extract.

To all those who treat neurasthenia, who have to lighten the lot of those unfortunate sufferers from what are known as "the functional neuroses," the importance of organo-therapy cannot well be over-estimated. When administered with intelligence, with the patient under observation, there need be no risk in such prescribing. Rather is there a risk in prolonging a morbid condition owing to neglect in the faculty of observation or an oversight as to the underlying causation.

Again, what a difficult matter it is to "fatten up" some patients! Rest cures, hyperalimentation, malt extracts, digestive ferments—all seem to be of no avail. But the extracts of the ductless glands will often be found of service, when taken with regularity. It should, however, be remembered that thyroid must be prescribed with great care in such cases as these. Extracts of the brain and spinal cord, of the pancreas and liver, of the testes and seminal vesicles, are of much help. I have recently had under my care a man suffering from advanced neurasthenia, whose weight had dropped from eleven stone to between eight and nine. Under the administration of a mixed

extract his weight is now (one month after the commencement of treatment) nearly ten and a half stone, although nothing had previously been able to stop the loss of weight.

In the *Practitioner* for January and February, 1915, will be found many able articles from the pens of experts on the subject of the endocrinic glands. These articles discuss fully the available material both from the standpoint of the laboratory and the bedside. A perusal of these numbers will well repay the time occupied.

With these introductory remarks, we will leave the general survey of the possibilities of opotherapy, and will pass on to a consideration of the thyroid and parathyroid glands. In the next chapter we shall deal briefly with the histology and physiology of these glands, and offer a short account of the functions which have been attributed by different observers to these bodies.

The grosser disorders of these glands are discussed in subsequent chapters.

CHAPTER I

THE THYROID AND PARATHYROID GLANDS

IN the Introduction we briefly reviewed the present place of the endocrinic glands in medicine, and suggested some practical points in connection with hormone-therapy. We now propose to discuss in a little more detail the more important of the ductless glands, laying emphasis upon the known results of administration of the extract of these glands, and the signs pointing to deficiency or excess of secretion.

Naturally, the thyroid gland first deserves our attention, and for several reasons. More is known of its characteristics, secretion, functions, and disorders; it takes a particularly important place in medicine on account of the frequency with which disorders of its functions are encountered; and, finally, of all the ductless glands whose secretions have been utilized in the form of extracts for oral or hypodermic injections, the thyroid gland has given the best and most astonishing results. From the moment when G. R. Murray first published the results achieved by administering thyroid extract to a patient suffering from myxœdema (October 10, 1891), great interest has been aroused in the practice of organo-therapy,

THYROID AND PARATHYROID GLANDS 17

an interest which Brown-Séquard's previous statements on the results of feeding with testicular extract had failed to arouse.

In 1890, Vassale in Italy and Gley in France experimented with injections of thyroid extract to animals who had been deprived of their thyroids, and demonstrated that these animals could be kept alive by such injections. This was followed shortly afterwards by the application of this discovery to therapeutics, when Murray treated his case of myxœdema by thyroid extracts and established that this condition could be cured by these means. Previous to this date, Gull and Ord in England had investigated myxœdema, and Kocher had described a condition following the removal of the thyroid, which, from the similarity of its symptom-complex, was shown to be identical with the spontaneous condition already named myxœdema.

As far back as 1859, Schiff discovered the fatal result which accrued from the total removal of this gland, and later, when Kocher described "cachexia strumipriva," or the condition of post-operative myxœdema which followed so many of his early operations for goitre, this fact attracted wide attention. Schiff and subsequent observers also discovered that transplanting the gland beneath the skin relieved the symptoms. Unfortunately, it has now been shown that an implanted gland is very liable to absorption, and thus to lose its utility.

These researches had established the fact that

18 THE ORGANS OF INTERNAL SECRETION

removal of the thyroid gland was followed by a train of symptoms which constituted the condition named "myxœdema."* A fresh factor then arose, on account of the discovery of two pairs of small glands, situated on either side of, and deep to, the lateral lobes of the thyroid, which received the name of "parathyroids." It was maintained by some authorities that the symptoms which followed the removal of the thyroid gland were due to the removal of these small glands, and in support of this theory it was adduced that, while removal of the thyroid alone was not usually fatal, removal of both thyroid and parathyroids proved rapidly fatal. On the other hand, some observers maintained that it was impossible to remove the parathyroids without also interfering with the integrity of the thyroid. Forsyth concludes, from the result of his investigations, that the parathyroids have no connection with tetany, and in this he is in opposition to most observers. We shall have more to say about this when we are dealing with the functions of the parathyroids.

Histology and Physiology of the Thyroid Gland.

The thyroid gland belongs to the group of ductless glands—that is to say, its secretion is poured into the blood-stream direct, and not by means of a duct. It is situated in the neck, and consists of two lobes,

* This name came into existence on account of the believed existence of excess of mucin in the subcutaneous tissues in these cases.

THYROID AND PARATHYROID GLANDS 19

one on each side of the trachea, extending upwards to the thyroid cartilage, covering its inferior cornu and part of its body. Joining these two lobes is the isthmus. The gland is surrounded by a capsule, and is composed of vesicles lined with cylindrical or cubical cells. These vesicles contain the typical colloid material, iodothyron, which is composed of iodine in combination with an active principle which has the characters of a globulin. Its recognition we owe to Baumann of Freiburg, who discovered its existence in 1896. Herbivora possess it in abundance, as most vegetables contain iodine. The weight of the gland varies between 32 and 60 grammes; it is larger in females than in males, and undergoes changes in size at the menstrual periods, and at such times as puberty and the climacteric; it is liable to swell under marked excitement and during emotional strain.

The colloid with which the vesicles are filled plays an important part in general bodily metabolism. In man there is nearly always sufficient for it to be discovered analytically (in normal health), but Wells states that the actual amount varies with the locality. It is scanty in young children and in parenchymatous goitre; unusually high in exophthalmic goitre. It has been suggested that an important function of the thyroid gland is to control the iodine metabolism of the body.

Now, what are the results of removal of the gland in man? When we come to review the knowledge which experimental physiology has given us, we are

20 THE ORGANS OF INTERNAL SECRETION

at first rather bewildered by the apparent contradiction which faces us. Early observations upon the effects which followed removal of the gland in animals showed that in carnivora the effects were very severe, and often fatal, tetany supervening and the animal dying from cachexia. In herbivora, however, the results of such an operation were often exceedingly slight, and were confined to a slight cachexia, but sometimes a rapidly fatal result ensued. The first inference to be drawn from this was that a meat diet was the important factor which accounted for the difference between the effects in carnivorous and herbivorous animals. Gley, however, showed that the explanation was quite different, and he proved that the parathyroids (one pair, now usually known as the "external parathyroids") were responsible for the wide difference in the results. He demonstrated that these small glands were so situated in carnivora that, when the thyroid was removed, the parathyroids were also removed; whereas in herbivora they were as habitually left behind. When the operations were made identical the results were approximated.

This originated the view now widely held, which regards the nervous symptoms following removal as attributable to the loss of the parathyroids—the effects of thyroid removal alone being, in the young a condition analogous to cretinism, in the adult a cachectic condition. Indeed, in the young animal the effects of thyroid extirpation are much more marked than in the adult animal. A few of the changes

THYROID AND PARATHYROID GLANDS 21

produced are striking. As we have already said, a cretinoid condition results, with delayed ossification of epiphyses, diminished development of the bones of the skull, protuberant abdomen, relaxation of the ligaments of the spinal column, and sexual infantilism. Alterations in the growth of the hair are observed in some species, and, most interesting of all, an extensive atheromatous degeneration of the aorta is found, which bears out the hypothesis that the function of this gland is concerned with calcium metabolism. Horsley had surmised previously to this discovery that many of the features of thyroid inadequacy resembled those which characterize senile decay.

Where thyroid secretion is excessive, we know that there is increased calcium leaving the body; and, in those patients exhibiting deficiency of thyroid, we may assume that increase of the calcium salts at the disposal of the metabolism results in an atheroma—a laying down of this substance in the walls of the arteries.

After thyroid extirpation there are several changes which deserve special mention. There is a marked retardation of metabolism in general, nitrogenous metabolism is greatly reduced, fat metabolism is likewise diminished, as the tendency to the deposition of fat in myxœdema shows. The animal without a thyroid exhibits excessive tolerance to carbohydrates; it can take abnormally large amounts without suffering from glycosuria. The reduction in the metabolic exchanges naturally diminishes heat production, and this point is of interest from the practical standpoint,

22 THE ORGANS OF INTERNAL SECRETION

human beings with deficient thyroid secretion being more susceptible to cold, and suffering increased discomfort in the winter months. As opposed to this, patients with Graves' disease rarely suffer any inconvenience from cold, but feel the heat of the summer months to be almost insupportable. As is well known, they are nervous and restless, and the increased tissue wastage produces a rapid loss of flesh, and in some cases actual emaciation.

The Parathyroid Glands.

Turning now to the parathyroids, whose existence, it will be remembered, is of more recent discovery, we find that there are two pairs of these glands, situated, as we have already said, in close proximity to the lateral lobes of the thyroid gland itself. In structure they bear some resemblance to this latter gland, resembling it, however, more in its embryonic characteristics. They are formed of columns of granular epithelium cells, and show a very vascular connective tissue between these. It has been stated that, if these glands are left when the thyroid has been removed, they undergo a marked hypertrophy.

To Sandström belongs the credit of first accurately describing these little glands. His view was that they were in reality embryonic rests of thyroid tissue proper. At the present time, although our knowledge of their functions has advanced somewhat, much of the knowledge we possess may be described as more

THYROID AND PARATHYROID GLANDS 28

speculative than proven. MacCullum came to the conclusion that they controlled in some way the calcium metabolism of the body. Calcium moderates the activity of nerve cells; therefore, when the secretion of the parathyroids is deficient, and calcium is lost from the nerve cells, the patient will suffer from an exaggeration of nervous excitability. This corresponds to the surmises as to the causation of the nervous symptoms which are manifest when the thyroid has been extirpated. In support of this view, the administration of extracts of the parathyroid glands in cases of tetany has been proved to be efficacious in controlling the convulsions. It will be remembered that tetany is a prominent symptom after extirpation operations in animals. Likewise the administration of calcium salts will control the symptoms of tetany.

Opposed to the views just quoted, some observers believe that the parathyroids are portions of the true thyroid gland; that they have become separated from the gland itself, or that they secrete the same colloid, although they have not as yet formed vesicles; or that they develop into normal thyroid tissue, intermediate types having been noticed.

It is stated that if all four parathyroids are removed the animal succumbs rapidly, no matter whether the thyroid is left intact or not. If one parathyroid be left, death does not usually ensue, although tetany may be brought on. It has also been said that changes in these glands are very common in cases of tetany in children, in pregnant women, and in

24 THE ORGANS OF INTERNAL SECRETION

osteomalacia (a disease associated with defective calcium metabolism). This would seem to be supported by the beneficial effects of calcium administered by the mouth, as it would tend to replace the excessive loss of calcium due to the defective parathyroid secretion.

With the few facts at our disposal, and the many speculative theories based thereon, it is natural that many diseases should have been attributed to an abnormality of these glands. To take one example, it has been suggested that paralysis agitans may be due to disease of the parathyroid glands, but, unfortunately, it does not appear to benefit by the exhibition of extracts of these glands. This is, of course, not tantamount to a refutation of this theory, as it may mean that the extract undergoes changes during its passage through the body which rob it of its natural properties, so that it is unable to replace the normal secretion which should be delivered into the blood-stream. The only instance which, so far as I am aware, has been published in which parathyroid extract appears to have been absolutely successful, and to have been, if one may use the expression, specific, is described by Hertz in the *Practitioner* for January, 1915, in an article on the parathyroids. Briefly, this case was a man of middle age, who had suffered from an enlargement of the thyroid gland, and had undergone an operation in which the greater part of it was removed. He remained well for two years, and then suddenly became

THYROID AND PARATHYROID GLANDS 25

very restless, tremulous, and developed fibrillary tremors and other signs which closely resembled Graves' disease. His hair ceased to grow, he became impotent, his bowels were loose, and his weight decreased very rapidly.

No treatment was of much avail, and the administration of dry thyroid gland aggravated the symptoms. He was given dry ox parathyroid by the mouth, and from that day he began to improve, became stronger, increased very markedly in weight, and returned to work. Not until he had been taking the parathyroid gland for six months did his sexual powers return. Hertz thus concludes the account of this very interesting case: "His weight in February, 1914, had risen to 189 pounds; he felt perfectly well and strong, and no trace of nervousness remained, although he was working very hard."

Here, apparently, was a case in which the secretion of the parathyroids was deficient, and where its administration by the mouth was able to replace the normal secretion.

If this latter surmise be correct, it would seem that it is possible for an analogous treatment to that of myxœdema to be initiated, if only we could arrive at an accurate diagnosis. It would support the school which believes in the separate functioning of the thyroid and parathyroid, and might eventually show us where we are lacking in the treatment of exophthalmic goitre. It is certain that the removal of part of the thyroid does not cure this disease.

26 THE ORGANS OF INTERNAL SECRETION

although it may ameliorate the symptoms for a variable time. If the nervous symptoms are due to excess of thyroid secretion, then removal of part of the gland should counteract the excess. But if this group of the many symptoms of Graves' disease owes its existence to an abnormality of the parathyroids, we must alter our surgical treatment of this disease.

On the other hand, we have yet to study the views of that school which has returned to Gley's original belief: that the parathyroids are part of the thyroid gland; that they are an embryonic and partly developed thyroid tissue. In support of this view, it is stated that post-operative tetany is greatly benefited by administering thyroid gland by the mouth, and that pure parathyroid, even in larger amounts, has not given the same results. It would appear from this that tetany owes its origin to deficient thyroid, and not necessarily deficient parathyroid, although the ordinary sheep's thyroid gland contains parathyroid substance.

This school, therefore, regards the parathyroids as part of the thyroid, and not separate structures; their function would be similar, and it follows from this hypothesis that the diseases which have been tentatively attributed to the parathyroids (mainly on account of the similarity which their symptom-complexes bear to those produced experimentally) must in reality owe their origin to the thyroid—assuming that they are in any way connected with this gland. This is comforting, as it narrows the field

of inquiry: it leaves us one set of riddles to solve instead of two. But we must admit that this view is not widely accepted, mainly on account of the many small facts which have been collecting, albeit slowly, to show us that the parathyroids are undeserving of neglect.

Functions of the Thyroid Gland.

We must now leave the subject of the parathyroid glands, and, before concluding this chapter, briefly review the theories which endeavour to account for the work of the normal thyroid gland. What is the function of the thyroid? Does it govern metabolism? Is it a "vital antiseptic"? Or is it concerned with growth alone? The only way in which we can attempt an answer to these questions is to give some few facts and more theories dealing with the function of this gland.

Firstly, then, the thyroid gland possesses the peculiar property—peculiar in the sense that it is not shared, so far as we know, by the other endocrinic glands—of being able to store its secretion. This is proven by the fact that, in cases where the gland has atrophied or been removed, its secretion can be replaced by artificial ingestion. We quite naturally turn to the colloid as being the stored-up secretion, and, indeed, we are justified in so doing, as there is evidence to show that this substance arises as droplets in the epithelial cells lining the vesicles (Dale). Again, this secretion contains a relatively large percentage

28 THE ORGANS OF INTERNAL SECRETION

of iodine (0.2 per cent.), and on this fact, or partly on this fact, has arisen the theory that the thyroid has a phagocytic or antitoxic action. In young animals, however, the iodine content is very small, and in adult animals it appears to bear a direct ratio to the iodine-content of the food. On the other hand, there is no reason to suppose that the power of the body to resist intoxication is raised when iodine or the iodides are given by the mouth; although Hunt found that young thyroids had some power to raise the resistance to a particular substance (acetonitrile), and that this power ran more or less roughly parallel to the iodine-content.

In this connection, the action of iodides upon gummata is of interest; and, as pointed out by Rendle Short, the beneficial action of these drugs is in reality due to the increased action of the thyroid which is engendered by the administration. He says: "I have found thyroid extract quite as effectual as iodide of potassium in healing tertiary syphilitic ulcers." In speaking of the action of iodides on gummata and atheroma, he says: "In cases of myxœdema arteriosclerosis is early and intense. The same is true in animals after removal of the thyroid."¹ Eiselsberg gives a number of very convincing photographs of intense atheroma of the aorta in his cretin lambs in which the thyroid had been removed in early life. In the second place, thyroid extract has a wonderful power over young connective tissue, as is seen by the way in which it absorbs the subcutaneous thickening of myxœdema and cretinism. It is not surprising,

THYROID AND PARATHYROID GLANDS 29

therefore, that it should be able to deal also with gummata and atheroma.

The theory of toxin-neutralization states that the secretion of the thyroid has the power to neutralize toxins which find their way into the blood-stream. It goes on to maintain that these toxins (of albuminoid nature) are absorbed from the alimentary canal, and undergo iodization in the thyroid by means of its secretion. It will be remembered that Lane maintains that atrophy of the thyroid gland is one of the features of chronic intestinal stasis. Is this the result of overwork? Hardly, for overwork produces hypertrophy, not atrophy; and yet, if the symptoms and signs attributed to intestinal stasis owe their origin to toxæmia, and the thyroid is the neutralizer, then (to reconcile the two theories) the thyroid should be enlarged, *not* atrophied. In other words, it should be overworked, not idle.

We know, however, several facts of importance about the thyroid to balance this tangle of theories. Firstly, we know that a train of symptoms follows its deficiency or absence, whether produced experimentally or arising spontaneously, and that these symptoms will yield to thyroid feeding. The results of an extirpation operation in young animals differ only slightly from the condition which we know as cretinism, while the adult analogy of this we encounter as myxœdema.

Thyroid is therefore concerned with the growth of bone, with the development of the body, and with a normal circulation. It has been suggested that the

30 THE ORGANS OF INTERNAL SECRETION

reason why the colloid is scanty in the young is that it is used as rapidly as it can be manufactured by the gland. As we shall see when discussing thyroid deficiency, the clinical pictures of the slighter forms of inadequacy require discrimination to discover, and are as yet not sufficiently definite for their wide acceptance. But the above facts lend themselves to further investigation.

Again, in the adult, there can now be little doubt that absence or diminution of the secretion produces, or helps to produce, a condition of secondary anæmia. Whether this argues any direct connection with the hæmatopoietic system, we are unable at present to say. That the thyroid is a direct circulatory stimulant there can be little doubt; for the slow pulse, cold extremities, sluggish circulation, and deficient action of the sweat glands, in submyxœdema, are very well recognized.

The interaction of the thyroid with the other ductless glands is at present unknown, but the future will probably show that the relation between the thyroid and some of the other endocrinic glands, notably the spleen, is a close one.

We must here leave the discussion of the thyroid and its small neighbours, realizing only too well that many blanks still remain to be filled up, which further light on this most difficult subject can alone do.

REFERENCE.

- ¹ The Newer Physiology in Surgical and General Practice, by A. Rendle Short, third edition, pp. 82 83

CHAPTER II

EXOPHTHALMIC GOITRE

Synonyms : Graves' Disease; Basedow's Disease:
Hyperthyroidism.

WITH increasing knowledge, the conception of this disease has become a much more difficult matter than when it was regarded as being solely due to an over-action of the thyroid gland. Modern views as to the pathology of Graves' disease would have us believe that, although derangement of this gland is present in this malady, the thyroid is not the sole organ at fault, neither is a hypersecretion of this gland alone responsible for the symptoms.

Definition.

Let us for a moment refer to some definitions of this condition which have been current, and then compare these with modern views as to the etiology and pathology. In one textbook of medicine we meet the following definition: "A disease characterized by enlargement of the thyroid, exophthalmos, increased action of the heart, tremor, and nervous instability."¹ Again: "The four classical symptoms

32 THE ORGANS OF INTERNAL SECRETION

of Graves' disease are—A staring appearance of the eyes, generally spoken of as exophthalmos, though there need be no actual protrusion of the eyeballs; moderate and almost symmetrical enlargement of the thyroid gland; a pulse-rate between 120 and 180 per minute—usually about 140 when the attack is moderately severe; and extreme nervousness, with fine tremor of the outstretched fingers. When all these symptoms are present at the same time, there can be little doubt as to the diagnosis, but very often some of them are absent, and it is possible for tachycardia to be the only symptom of the disease.”² “There are three prominent symptoms: Protrusion of the eyeballs, enlargement of the thyroid gland, and frequent action of the heart.”³ “We are accustomed to recognize three cardinal symptoms in this disease—namely, (1) tachycardia, (2) goitre, and (3) exophthalmos; but we must remember that these are not the only symptoms.”⁴

These definitions represent the general views which have been held on the nature of the disease and on the most constant symptoms.

The condition was first recognized by the celebrated Dublin physician whose name it now bears about the year 1835, although Von Basedow in 1840 published a paper on the subject. Consequently the disease is in Germany and some other parts of the Continent still referred to as “Basedowsche Krankheit,” or “Von Basedow's disease.” However, as early as 1825 Caleb Parry, of Bath, drew attention to the condi-

tion, and, according to Osler, to him belongs the credit of first describing the disease. Many of the early conceptions of the malady have, in the fulness of time, given place to views which have been promulgated from the results of the extensive practical physiological research which has been undertaken in order to establish the causation of the disease. But it was recognized then, and it is believed now, that emotional strain can precipitate the disease. Thus, Trousseau refers to a lady who was suffering great grief at the death of her father, and had been crying for a long time; she "suddenly felt her eyes swell and lift up her eyelids." This was accompanied by copious epistaxis, violent palpitation, and throbbing and enlargement of the thyroid. A few days later the nature of the disease was recognized. Again, Stokes describes the case of a man who developed the disease from long-continued bleeding from piles; and many other records show that the etiology of the disease was universally regarded as a wide one.

From the early thirties of the last century the nature of the condition was recognized, but it is only comparatively recently that the diagnosis has been narrowed. Thus, it will be seen, from the definitions quoted above, that it is not necessary, as used to be thought, for all the classical symptoms to be present in any given case. Nevertheless, there is one symptom without which, as Mackenzie rightly insists, the condition cannot be diagnosed, and that is persistent tachycardia.

84 THE ORGANS OF INTERNAL SECRETION

So we come down to this: that exophthalmic goitre may, and in the opinion of many observers does, exist without the exophthalmic symptom. It is, therefore, somewhat unfortunate that the name of the disease should be inseparably connected with a symptom which is by no means constant. But we cannot, at the moment at any rate, suggest any other name which is free from objection. As we shall see later, the modern nomenclature which is sometimes used—namely, “hyperthyroidism”—is open to an equally serious objection, as it implies that it is always an over-action of the thyroid alone which is responsible for the features of the disease, of which we must reckon proptosis, when present, as one. Again, it has been pointed out by one observer that this latter symptom is significant of over-action, not of the thyroid, but of the adrenals.⁵ Therefore, hyperthyroidism is a no more suitable label for the disease than is exophthalmic goitre. Although there seems to be an objection to utilizing the name of the discoverer of a disease to designate that disease,* in the present

* It may be affirmed that this mode of naming a disease after its discoverer is to be deprecated, partly on account of the difference of opinion as to whom the credit of the discovery really belongs, and partly because it is, perhaps, an unfortunate reward for the happy pioneer in the particular disease. Again, if a disease is discovered more or less synchronously by different observers in different countries, a nomenclature is adopted which varies with the different countries, and this all leads to confusion. If a disease is called by its most distinctive feature, this will probably be similar in different languages, so where such a course is possible it is certainly to be preferred.

instance it would seem, for the moment at any rate, to be the most satisfactory way to designate this malady. "Graves' disease" has the merit of being non-committal as to symptoms, and less unwieldy than many of the other names by which the disease has been known.

In the majority of textbooks, the malady under discussion is referred to as "exophthalmic goitre," in contradistinction to ordinary goitre. When the swelling in the neck, if present, is accompanied by the other well-recognized signs of Graves' disease, such as persistent tachycardia, exophthalmos, tremor, and other symptoms and signs to be described anon, then we diagnose the presence of this disease as opposed to simple goitre.

Etiology and Symptoms.

Graves' disease (as we shall call the malady in this book) is seen more commonly in the female sex than in the male, and more commonly in young people than in elderly. It has been seen, however, in an infant only two and a half years of age, and several cases are on record of the disease occurring in children of both sexes. Von Graefe stated that the proportion of females to males was 6 to 1, while Eulenburg said that the ratio was at least 2 to 1. Trousseau's cases show a ratio of 50 to 8, Henoch's 23 to 4, and Præel's 28 to 1. Whatever figures we study, there is plainly a vast preponderance of females over males. The

36 THE ORGANS OF INTERNAL SECRETION

commonest time for the disease to develop is in the decades twenty to forty, and its features are often made manifest at critical times in the history of the patient, such as puberty, the catamenia, and the menopause. Indeed, it is said that in normal persons of the female sex the thyroid is liable to swell and to exhibit increased vascularity at these times and during sexual excitement.

As we have already indicated, there can be little doubt that, certainly in individuals prone to the disease (by this I mean persons who exhibit signs indicative of thyroid instability), a mental strain or a sudden anxiety seems to be capable of precipitating the disease. Again, there is a sufficiency of evidence to show that a parenchymatous goitre can develop into an exophthalmic goitre, given suitable opportunities. The symptom-complex which is produced by thyroid feeding on a large scale differs in many details from the symptom-complex of this disease, so, as Biedl says, we must regard the similarity of the two pictures, not as conclusive evidence that the thyroid is the organ responsible for the disease, but as very strong presumptive evidence.

When we come to study the symptoms of Graves' disease, we see three or four definite and fairly constant features, and a multitude of smaller and somewhat vaguer signs. There can be little doubt that tachycardia deserves the first mention, as it is very constant, and may be looked upon as a fundamental sign of this disease. Indeed, it is one of the symptoms which

can always be produced by the ingestion of thyroid extract; and there must be very few cases of undoubted exophthalmic goitre which do not show this sign. The pulse is usually rapid, ranging from 100 to 140 or more per minute. The pulse-wave is not always thin; indeed, in many cases a full and bounding pulse is observed. Again, it is usually regular, but the rate increases upon very slight exertion.

In this connection it is interesting to refer to the theory which Eppinger and Hess brought forward. They consider that the symptoms seen in Graves' disease enable us to divide these cases into two groups. In the first the symptoms of sympathetic excitement predominate, and these they describe as sympathetico-tonic; while the second group is described as vago-tonic, from the fact that the symptoms seem to proceed from disorganization of the autonomous system. "Falta, Eppinger, and Rudinger, assume a polyvalency of the thyroid secretion, and they regard the hyperthyroidism of Graves' disease as the outcome of a simultaneous though probably independent stimulation of both the sympathetic and autonomous nervous systems."⁶

There seems to be little doubt that tachycardia is only one of the many symptoms seen in this disorder which can be justly attributed to the sympathetic nervous system. We shall refer to these later.

The symptom which merits consideration next is the local enlargement of the thyroid gland. Although not so constant as tachycardia, it is nevertheless

88 THE ORGANS OF INTERNAL SECRETION

present in the majority of cases of this disease. The thyroid is generally moderately enlarged, the right side being perhaps more so than the left. In some cases, however, the enlargement is scarcely perceptible, while, on the other hand, it may be very great. In the early stages the gland is soft and elastic from vascular engorgement, but later it becomes harder from fibrous hyperplasia.

Histologically the gland presents a picture of diffuse enlargement, with a great increase in the vascular supply. There is definite new formation of tissue, which runs hand in hand with certain retrogressive processes, notably cell desquamation. Young follicles are seen associated with older cells, and in the latter may be seen breaking-down processes. As a rule a thrill can be detected, certainly in the larger tumours, and as the disease progresses the thyroid may alter in size, diminishing in favourable cases. It is said that the histological appearance in Graves' disease is typical, and Erdheim maintains that the young cell formations with fat granules are characteristic of Graves' disease.

For a long time the proptosis was looked upon as an essential feature of this disease, and it is only recently that we have come to realize that it is by no means always present. Its origin or, rather, the changes which underlie the exophthalmos are still unknown. It has been suggested that it is due to a deposit of fat behind the eyeballs; to a venous congestion in the posterior part of the orbit; to dilatation

of the retrobulbar arteries; or to contraction of Müller's muscle. It is pointed out that the sympathetic system is largely connected with the symptoms of this disease, and that a disorganization of this system will account for many, indeed for most, of the features, always excepting the changes in the gland itself. On this theory it is much more probable that the proptosis is due to circulatory changes than to either a contraction of Müller's muscle or a deposit of fat in the orbit.

There are certain classical signs which one is accustomed to look for in association with the exophthalmos, and we must briefly refer to these. The widening of the palpebral fissure gives an appearance of great protrusion to the eyeball, and this is partly due to the retraction of the upper lid. This is known as Stellwag's sign. Although the eyeball in many cases appears to be prominent, it is in reality less so than it appears, on account of the uncovering of the eyeball due to the retraction of the upper lid. There is also a diminution of the reflex excitability of the eye, so that there is less irritation of the globe than would otherwise be the case. Another sign in connection with the lid is known as Von Graefe's sign, and consists in a lagging of the upper lid behind the globe during the downward descent of the eye.

Sometimes there is an insufficiency of the internal recti muscles of the eye, as a result of which convergence of the eyes in near vision is imperfect (Möbius). Occasionally blepharo-clonus is present.

Tremor is another constant sign of Graves' disease,

40 THE ORGANS OF INTERNAL SECRETION

and is quite characteristic. It is usually very fine, and confined to the hands, although occasionally it is seen in the muscles of the trunk, so that by laying a hand on the shoulder or trunk the observer can feel a quiver of the whole body (Oppenheim). The tremor of this disease is usually excited by movement or by nervousness, but is also present during rest. The rate of movement is somewhere about nine per second, and can best be seen when the patient extends the hands with the palms downwards.

Before turning to the discussion of the mental changes present in Graves' disease, there remain a few less constant symptoms and signs which we have to mention. These may be classed under the headings vasomotor, secretory, and trophic. The subject of Graves' disease is very liable to suffer from profuse perspiration; indeed, in some cases this amounts to a definite hyperidrosis. As we stated in the previous chapter, these patients feel the heat greatly, and naturally this symptom is worse in hot weather. The sweating may be more or less local, or it may be general. One point of interest in this connection is with regard to the electrical resistance of the skin in cases of Graves' disease. Vigoroux first discovered that in patients suffering from exophthalmic goitre the electric resistance of the skin was diminished. This is now generally believed to be due to the undue moisture of the skin owing to the increased sweating.

Flushing of the skin, quite irregular in distribution, erythemata of a patchy nature, pigmentation—

particularly of the margins of the lids of the eyes, and sometimes as marked as that seen in Addison's disease—are all phenomena of this condition. As is well known, there is a tendency to loose evacuations of the bowels, sometimes of actual diarrhoea; while the secretion of urine is also increased. The digestive system is often upset, and attacks of sickness, with bulimia or, alternatively, loss of appetite, are about equally common.

The reflexes are often altered—more usually increased, although they may be diminished, or even, in rare cases, absent.

Cases are occasionally encountered in which the symptoms present a strange combination of those seen in exophthalmic goitre and those characteristic of thyroid deficiency. Thus, patches of lipomatosis are sometimes met with in a typical case of Graves' disease; while occasionally the condition of the skin approximates much more nearly to that typical of submyxœdema. Leonard Williams believes that there is often, if not always, a combination of excess and deficiency in this disease. Again, some of the symptoms which we are accustomed to regard as indicative of disease in the chromaffin system, such as pigmentation, would suggest the presence of a combination of thyroid and adrenal disturbance.

But as important, or nearly as important, as the physical symptoms of this complaint is the mental change which characterizes exophthalmic goitre. The main features of the sufferer from this disease are well

42 THE ORGANS OF INTERNAL SECRETION

known, but perhaps a brief sketch of the mental make-up of the patient may be useful. Whatever is the agent at work in these cases, whether the disturbance is originally in the sympathetic system, or to be attributed to an excess of thyroid secretion due to a cause or causes unknown, it is one fraught with evil for the peace of mind both of the patient and her relatives. From being gentle and docile, it may be, she changes to an intractable, selfish, restless, and inconsiderate being. The medical attendant as a rule receives the full benefit of this, and little he can do is right. He is either old-fashioned when he explains that the reason for rest in bed, for example, is to avoid straining an already weakened heart; or an ignoramus if he insists that rest combined with hygienic principles offers the best hope for alleviating the disease. If he suggests trying a new remedy, he is experimenting with her; if he persists with the old, he is a "stick-in-the-mud."

Perhaps the saddest of the changes wrought by this disease are the changes for the worse in the psyche, and it behoves the medical man to be very tactful with such patients, and to remember that the mind is the victim of a disordered bodily functioning, and is no more to blame for its vagaries than the victim of, let us say, puerperal mania.

The mental symptoms are, like the physical, capable of entirely clearing up; but even so, Leonard Williams doubts whether the individual is ever quite the same again. There can be little doubt that this disease

plays havoc with the patient's character. It changes the quiet to the restless, the unselfish to the self-centred, and the amiable to the perverse. That these patients are notoriously difficult to treat is well known; that they make the treatment of their condition doubly difficult by their attitude to what is being done for them is almost equally well realized by everyone connected with the case except the patient herself.

Apart from the changes in character which we have outlined, more serious symptoms are seen in connection with Graves' disease. Apart from actual insanity, such as melancholia, mania, hallucinatory confusion, and obsessions, there are the minor changes which are apt to lead to these more serious psychoses. Thus, the patient is excitable, wildly restless, sleepless, and confused. She is incapable of organized thought, of successful memory, or of attentive control; while her general mental habits have been well spoken of by Reynolds as "mental chorea." It is quite obvious that it is but a short step from this stage to that of insanity.

Needless to say, fortunately only a relatively small percentage passes from the mentality characteristic of Graves' disease to that typical of insanity. Nevertheless, the picture we have drawn of the mental condition of these sufferers is not exaggerated. It is true that in many diseases which the physician is called upon to treat to-day the patient's worst enemy is himself; and in no disease is this more true than in exophthalmic goitre. Consequently, it is no rare

44 THE ORGANS OF INTERNAL SECRETION

occurrence for such a patient to pass from one doctor to another, giving no one a fair chance to improve the patient's condition. This is the analogue in the mental apparatus of the restlessness in the physical system which prohibits the patient from resting, which is the one thing most to be desired. Thus is her temperament at the time her worst enemy, and it is this which needs treatment quite as much as the syndrome of symptoms which we have been discussing.

Such, then, is the bare outline of what constitutes Graves' disease. What is the prognosis, and what can we do to benefit that part of suffering humanity afflicted with Graves' disease?

Prognosis.

First as to prognosis. The course of the disease is very variable; it may in rare cases be rapidly fatal, or it may linger for many years, sometimes showing improvement, at other times retrogressing. Mackenzie gives the mortality at approximately 25 per cent. In patients who eventually recover, there is often a period when no improvement seems to take place and the disease seems to be stationary. Even in these cases there is a prolonged convalescence, and sometimes years afterwards some slight trace of the disturbance can be found.

In the early stages the malady is very liable to be diagnosed wrongly, some such diagnosis as neurasthenia being made. This probably partly accounts for the

advances which this disease makes in the early stages, owing to a misconception of the nature of the trouble. Also the prognosis must be based upon the length of duration of the disease, the prominence of individual symptoms, the means of the patient, and, last but by no means least, the degree of mental abnormality which exists.

But, speaking generally, patients who have had the disease for a long time, especially where treatment has been of little avail, are not the most hopeful of subjects, and there seems little reason upon which to base hopes of complete recovery. Indeed, in such cases as these it is doubtful whether they are ever entirely restored to the *status quo ante*. Although it is usual to see amelioration of the symptoms, even a vanishing of the proptosis, or at least a diminution of the amount of protrusion, under successful treatment, such patients are extremely liable to a relapse. Their nervous symptoms recur from time to time; indeed, it would seem that it is very difficult for them to return to the successful functioning of this system or Graves' disease has developed.

There have, however, been many reports of recovery of long duration. Cheadle reports recovery lasting twenty years; while Oppenheim says that, in one case of his, recovery lasted "for twenty-seven years, in another for eighteen, in four for six to eight years."⁷ The chief difficulty lies in keeping up the faith of the patients, in exacting obedience, and in counteracting individual symptoms.

46 THE ORGANS OF INTERNAL SECRETION

Sudden death has taken place during the course of this disease, and when we remember the strain upon the circulation, the dilatation of the heart, and the disorganization of the neuro-musculature of this organ, it is not to be wondered at. Occasionally, as a sequel to this malady, atrophy of the thyroid may take place, with resulting myxœdema.

Treatment.

We shall have to refer at some length to the treatment of this malady, for it necessitates a discussion of methods which have been utilized for many years, as well as those which have found favour recently. Modern physiological research has enlightened us upon many points in this connection, and some of these have pointed to different methods of treatment, a few of which, at any rate, have shown signs of success.

But whatever treatments we may initiate—whether we incline to the old practice of counteracting the symptoms by controlling the exuberance of the heart's action, by producing adequate sleep, by anointing the thyroid gland, or by countless other small attentions—one factor remains constant, and that is the insistence upon sufficient rest, even complete rest in bed in bad cases. The application of the principles of general hygiene cannot be insisted upon too firmly, for this should be the bed-rock upon which all subsequent treatment is based. The difficulty is that the restlessness which these patients so often exhibit makes this

stipulation a difficult matter to enforce. Nevertheless, it is not so much a matter of the practice of medicine as of the practice of common-sense which suggests this; for it must be obvious that exertion, particularly undue exertion, which is always followed, as it is in this disease, by an increase in the symptoms, must be wrong, and therefore in no circumstances to be considered. When tachycardia is present, as it invariably is, our first aim must be to reduce the work of the heart to a minimum, and in doing this to reduce the tachycardia. Even if the rate of the heart's action is not excessive, the patient must be made to take periodic rests in the course of the day. She should rise late and retire to bed early, resting after meals, and avoiding hurry and perturbation. Where tachycardia is a marked feature, and more especially where the signs of general debility with marked loss of flesh are evident, complete rest in bed is essential.

There can be little doubt, moreover, that these patients require every hygienic advantage which can be given them. Thus, where possible, they should reside in the country or by the sea, but not in too bracing a place. They should rest in the open air, lie in the sun for the greater part of the day, and attempt no exercise of a strenuous nature. I have on several occasions had occasion to emphasize the harmful effects which absence of rest has produced; but it is often extremely difficult to make the patient understand this. Again, the diet must be regulated to the individual case, and with some thought to the

48 THE ORGANS OF INTERNAL SECRETION

nature of the disease. Thus, we must bear in mind what underlies this malady, especially what has been discovered in recent years. Firstly, the condition of the gastro-intestinal tract must be considered: if the digestion is good, and there is little tendency to diarrhoea, vegetables may be permitted, but this item of diet must remain under the control of the medical attendant.

Again, in this connection we should bear in mind another important fact—namely, that iodine is excessive in the colloid in Graves' disease, and is also contained in most vegetables. Therefore we do not want to add to the amount already present in the body. In this connection Rendle Short says: "We see also that exophthalmic goitre is due to hypersecretion of the iodothylin, as is proved by the artificial imitation of the disease by excessive thyrcid feeding, by the excess of iodine present in the colloid in Graves' disease, and by the character of the histological changes. Thus, we have reason to expect good from partial removal, which has been very successful in the hands of Kocher, the Mayos, and others. It would be reasonable also to try the effect of iodine starvation by eliminating vegetables and ordinary tap-water from the dietary, and substituting for the latter the water of a goitre well. It is well known that exophthalmic goitre and parenchymatous goitre show a sort of geographical antagonism, and the effect of the water in reducing the amount of iodine for conversion into iodothylin would be valuable."⁸

It must be borne in mind that parenchymatous goitre is supposed to occur owing to a deficiency of the iodothyron, so that the thyroid hypertrophies in an endeavour to supply that deficiency. If, therefore, there is little of this element in the thyroid of sufferers from parenchymatous goitres, they may be helped by the ingestion of vegetables. The water of the well-known goitre wells is supposed to produce parenchymatous goitres by containing a substance which deprives the body of the iodine by forming a combination with it. In an effort to counterbalance the diminished output of iodine the gland hypertrophies. On this assumption, then, it is recommended to try the water on patients suffering from exophthalmic goitre, in the hope that the unknown substance in the water will utilize some of the excessive secretion of the thyroid.

Turning now from the consideration of the general treatment of this disease, we find a host of other remedies which have from time to time been recommended. Apart from symptomatic remedies, over the consideration of which we shall spend but little time, we have before us a choice of many drugs, both for internal and external application. Of these, preference seems to lie with belladonna, aspirin, the salicylates, arsenic, and iron salts (where indicated). As we have already said, Graves' disease is often associated with anæmia, and therefore some benefit may be expected by the administration of these latter salts.

50 THE ORGANS OF INTERNAL SECRETION

Of cardiac tonics, some authorities prefer strophanthus, while others recommend digitalis and nuxvomica. But the help which the physician may expect from these drugs is strictly limited to their local action upon tachycardia.

Leonard Williams has reported good results in several cases from the hypodermic or intramuscular injection of bile salts. The *rationale* of this treatment lies in the fact that it is well known that these salts circulating in the blood produce a sedative effect on the brain and a slowing of the pulse.

Of recent years more and more stress has been laid on the chance of finding an organo-therapeutic compound which will help to neutralize the excessive action of the thyroid. Extracts of the adrenals have been tried, as also the extracts of pituitary, the ovaries, the parathyroids, and the spleen. Unfortunately, these have not given good results, only the parathyroid holding out any hope in this direction.

But there remain several external remedies to mention, some of which have been stated to give really good results in some cases. Of these, the X rays is very well spoken of when applied to the exterior of the gland, and it has been said on many occasions to have limited the exuberant activity of the thyroid. Again, galvanism and faradism, applied either to the gland itself or to the sympathetic in the neck, has been tried, sometimes with success. Or a compress of adrenalin applied to the goitre is a help in some cases. I have had good results in some patients from the

application of a mercury ointment to the thyroid. A small portion rubbed in every night, associated with internal treatment, has on more than one occasion resulted in an amelioration of the symptoms.

Again, since 1884 preparations have been manufactured from thyroidectomized animals. Of these, the more important are "antithyroidin" (Möbius), "hæmato-éthyroidine," and "thyroidectin." Beebe has evolved a method of treating Graves' disease by an antiserum produced by the inoculation of a thyroid preparation in animals.⁹

It is too soon to speak of the results of this latter treatment, but its value is said to have been proved in a large number of cases.

Some observers, notably H. Campbell,¹⁰ advocate the administration of calcium salts to patients suffering from exophthalmic goitre, and claim better results from treatment in which this medication is included than from any other treatment.

Again, preparations of thymus gland have been recommended for this disease, but so far no very encouraging reports have been published.

Many authorities speak very highly of the beneficial effects of X rays, and claim that treatment by this method successfully counteracts the over-action of the thyroid; while some observers maintain that the X rays should always be administered before an operation is finally decided upon.

With regard to the question of operative interference, there can be little doubt that it should be reserved

52 THE ORGANS OF INTERNAL SECRETION

for the grave cases only, and that it should only be performed after all other remedies have failed.

The medical man must insist upon hygienic principles as a *sine qua non*, and he, as well as the patient, must regard the disease as one requiring the tacit obedience which is given to the orders of the medical man in a disease such as pneumonia or typhoid. If this is done, the patient is prevented from straining an already overworked organ, and much possible mischief may be prevented.

In conclusion, we must emphasize the fact that, of all diseases which the medical man is called upon to treat, probably no one makes such calls on his perseverance and patience. For he has to treat an irresponsible patient, and one who is not the best judge of her condition. He requires infinite tact and an everlasting patience if he is to see the fruits of his labours. Furthermore, he must perforce try any remedies which hold out a chance of help, and he must discriminate nicely between those which are logically futile and those which are based upon reason.

There is some ground for hoping that in the near future, as our knowledge of the disease and its causation widens, we shall evolve a more satisfactory mode of treatment. For the present we must make up by our resourcefulness what we lack in our nicety of knowledge.

REFERENCES.

- ¹ A Textbook of Medical Practice, edited by Bain, p. 315.
- ² Index of Differential Diagnosis of Main Symptoms, edited by French. p. 772.
- ³ The Practice of Medicine, by Taylor, p. 860.
- ⁴ A Textbook of Nervous Diseases, Oppenheim. p. 1343.
- ⁵ Maurice, *Lyon Médical*, quoted by Leonard Williams in *Practitioner*, January, 1915, p. 96.
- ⁶ The Internal Secretary Organs, by Biedl, p. 98.
- ⁷ Oppenheim, Textbook of Nervous Diseases, p. 1340.
- ⁸ The Newer Physiology in Surgical and General Practice, by A. Rendle Short, pp. 85, 86.
- ⁹ The Diseases of the Ductless Glands, for Practitioners of Medicine and Surgery, by S. P. Beebe.
- ¹⁰ H. Campbell, Exophthalmic Goitre, *Clinical Journal*, 1915, xliv. 329.

CHAPTER III

THYROID DEFICIENCY

IN the previous chapter we dealt with the disease variously known as "Graves' disease," "exophthalmic goitre," and "hyperthyroidism," and showed that there is strong evidence to support the view that the symptoms are largely caused by excess of the thyroid secretion. In this chapter we propose to discuss thyroid deficiency, laying emphasis on the lesser degrees.

The history of thyroid-therapy has already been given, and we have described the earlier operations of Kocher upon the thyroid, and referred to the condition of "cachexia strumipriva" which was liable to ensue. Likewise we have mentioned the earlier work upon the functions of this gland, and the discovery by Gull and Ord of the disease myxœdema. Reference was also made to the important fact elicited by G. H. Murray, that this condition yielded to the administration of extract of thyroid gland.

The discovery was of the greatest importance, and for two reasons: First, because it proved beyond question that the peculiar disease to which the name "myxœdema" had been assigned was largely due to

a deficiency or absence of the thyroid secretion; second, and even more important, it enabled complete relief to be afforded to the unfortunate sufferers.

Since that time a great deal of work has been done on the thyroid, both by laboratory workers and clinicians, with the result that myxœdema has emerged from the nebulæ of ignorance, and has taken its place as a curable disease.

But quite recently workers in all fields of medicine have been observing cases which, while they were by no means comparable to myxœdema, were sufficiently similar to appear to resemble this condition in certain points. As a countenance reminds one of a more familiar face by reason of similarity in one feature, so do these cases, sometimes because of an item we can barely specify, make one think of the more marked disorder.

In this chapter, then, we wish to include those cases whose vagueness is a defiance to conclusive diagnosis, as well as the more definite disease which has received the name of "myxœdema." As we have already in a previous chapter discussed "cachexia strumipriva," we shall here confine our attention to that variety which arises apparently idiopathically, and not ensuing after operative procedures.

Myxœdema.

We will briefly review the features of this condition, but will not attempt to describe them in any detail, as they can be found discussed in any textbook of medicine.

As we have already said, Gull and Ord, G. R. Murray, Kocher, Mayo, and others, are the workers responsible for unearthing the cause of the symptom-complex which is known as myxœdema, and to Murray belongs the credit of discovering that the administration of thyroid extract relieved the symptoms.

Myxœdema is one of those interesting clinical conditions which are diagnosed rather by the summary of the effects produced by the individual symptoms than by one predominant feature. Thus, we find that a typical instance shows to the observer a slowly moving, mentally sluggish, and prematurely aged individual, with localized deposits of fat, more marked in certain areas, a dry, rough skin, characteristic facies, and uncertain gait. The patient is altered in most ways; indeed, the disease, in common with many, if not most, of the disturbances of the endocrinous glands, changes the personality of the patient almost beyond recognition; but this alteration is, in the majority of cases, gradual.

The onset of the disease is insidious, and may occupy several years, although in a few instances its onset is more acute. The initial symptoms vary in different cases, but most usually commence with

lassitude, debility, disinclination for exertion, and marked susceptibility to cold and chills.

The patient appears a changed man, and, to anyone who has not seen him for a protracted period, almost a different being. He becomes much stouter, and the actual body-weight is increased, but the fat is deposited in a characteristic manner. Pads of adipose tissue are situated in the clavicular regions, over the cervical spinal area, and in the neighbourhood of the lower ribs and flanks. But perhaps the facies is even more characteristic. The eyelids are thickened, with resilient bags of fluid under the lower lids; the lips, ears, and tongue, are enlarged; and the whole appearance of the face is one of extreme coarseness. At the same time there is an absolute lack of any intelligent expression in the face; the features are immobile, with, in some cases, a slight expression of surprise owing to the eyebrows being somewhat raised.

The skin loses its normal moisture, and becomes harsh, dry, and rough. The teeth are frequently in a state of decay, the nails are ridged, and the hair loses its gloss and shows evidence of trophic change—in fact, the epidermal appendages undergo atrophic degeneration.

The individual becomes comparatively lifeless. His extremities change; his hands are carried in a peculiar manner, which has received the name of "spade-hands," while his feet enlarge (not, however, in a marked manner, as is seen in gigantism), and are

58 THE ORGANS OF INTERNAL SECRETION

carried in a clumsy manner. Even the fingers show a change, for they are shaped somewhat like sausages, and are not narrower between the joints as are the fingers of a normal person.

Small warty growths are liable to appear on the surface of the skin; while patches of dry eczema, similar to psoriasis, make their appearance.

The changes in the habits are equally characteristic. The speech is slow, and a long time elapses before an answer is given to a question. Co-ordination is interfered with, and the gait is "waddling." The mental attitude is almost exactly antagonistic to that which is typical of exophthalmic goitre. Thus, the patient is slow in every act, whether this act involves mental or bodily exertion. The finer co-ordinations of the muscles are deficient, a greater length of time is required even for simple acts, while the mental hebetude is shown by the set and unintelligent expression of the face.

This disease depends upon an atrophy of the thyroid gland, which can be demonstrated by an examination of the gland in the neck. It will be much harder to localize than in health, and to the examining finger its presence at all can only be detected by requesting the patient to swallow.

That the disease owes its origin to an absence or diminution of the thyroid secretion there can be no doubt; for the administration of an extract of this gland removes, or markedly lessens, the symptoms.

Lesser Degrees of Hypothyroidism.

Now let us turn for a moment to the study of the lesser degrees of deficiency in thyroid output, and we will then study the physiological effects which the thyroid produces upon the body.

Needless to say, it is a more difficult matter to deal with these lesser degrees than it is to diagnose those marked instances known as myxœdema. There can hardly be any mistaking a case of myxœdema, unless a particular instance is very atypical. Once a typical instance is seen, the peculiar characteristics cannot well be forgotten. When, however, we come to describe the symptoms and signs which would lead us to arrive at a diagnosis of hypothyroidism, the difficulty arises that the various signs are widely divergent (according to modern views), and that their presence at all may be transitory, or, again, the physical signs may be very few. We shall endeavour to describe those signs which are usually present in a case of sub-myxœdema, laying especial emphasis upon those which are characteristic. In this way it is hoped that those cases requiring thyroid medication will be more readily recognized.

As is to be expected, on the whole the clinical picture is a modified version of myxœdema, with this difference, that the disease—*i.e.*, the thyroid deficiency—is still probably in its youth, and the more advanced changes, which are dependent upon structural alterations of the grosser kind, are absent.

60 THE ORGANS OF INTERNAL SECRETION

It is not possible with our present knowledge of this subject to do more than hazard a guess as to the cause of thyroid insufficiency, and this we have already done.¹ Whether prolonged illness on the physical side, and anxiety on the mental, are capable of initiating a thyroid disturbance we do not know for certain. With the evidence at our disposal, it would seem probable; for, in the experience of most medical men, hypothyroidism is more liable to ensue after illness, worry, and similar harmful processes, than it is to arise spontaneously.* Whatever its genesis, it behoves the medical man to be on the lookout for its signs, more especially after acute illness. Let us briefly review its features, and outline the signs of thyroid insufficiency. Before doing this, however, it is better to state that there is a certain difference of

* I described a case of submyxœdema in a young male subject in the *British Medical Journal* for June 20, 1914. This followed a prolonged period of financial stress on the Stock Exchange, and was quite typical in its features. The patient was a young man, who had been out of sorts for some weeks, and had gone about his usual occupations, although he had noticed that he had been putting on weight rapidly. Briefly, his symptoms consisted in dyspnoea on exertion, bradycardia, eczematous rashes, pads of fat in the clavicular region, subnormal temperature, slow pulse, and trophic changes in the hair. Even the "eyebrow" sign was present. This man was first treated with cardiac stimulants, rest, and general hygienic measures, but made no progress. The day after he commenced thyroid medication the temperature approximated to the normal, the pulse-rate increased, he became more comfortable in himself, and after three weeks' treatment his good health was restored, with a reduction of weight. As may be imagined, his temperature chart was a most instructive one.

opinion as to what constitutes the typical picture of thyroid insufficiency. We shall, therefore, at this place confine our remarks to those signs about which there can be little dispute, and in conclusion we shall briefly review the many points in this connection about which opinion is unsettled.

Signs of Thyroid Insufficiency.

When studying the main features of Graves' disease, we saw that the rate of bodily metabolism was vastly increased, that the nutritive exchanges were accelerated, and that there was a loss of weight in consequence. Again, we noted that the vasomotor system was irritable, that perspiration was easily induced, erythematata were frequent, and that the tendency to the production of glycosuria was increased. The general picture of thyroid insufficiency is exactly opposite (I am speaking of the main features) to that seen in exophthalmic goitre. Let us take the salient points one by one:

Temperature.—In this condition the bodily temperature is usually subnormal, in extreme cases as low as 96° F., more generally about 97° to 97.5° or 98° F. In any case, if the temperature be taken consistently, it will rarely be found to be normal. In this connection we must remember that these patients feel the cold in a marked manner, and are in a more or less constant state of chilliness. A patient who constantly complains of his "bad circulation" should be suspected of thyroid inadequacy.

62 THE ORGANS OF INTERNAL SECRETION

Pulse.—The pulse-rate in submyxœdema is consistently slow, although I am aware that many authorities differ from this. I would therefore modify this statement by saying that, where other signs of thyroid insufficiency are present, and the pulse-rate is not slower than normal, or even faster, this is a sign that the condition present is not one of simple thyroid inadequacy, but is a condition of complicated endocrinous disturbance—a condition in which there may be more or less concomitant disturbance of the thyroid and some other internal secretion, or in which there may be a concurrent hypo- and hyper-thyroidism.

General Nutrition.—In submyxœdema, just as in the larger disease, there is a storage of products of digestion, as is seen by the larger amounts of sugar which can be consumed before glycosuria is produced. The bodily weight is therefore increased, but the deposit of fat is more or less local, as in myxœdema, and certain areas are more affected than others. Thus, the neck and shoulders are thickened, the clavicular regions contain pads of fat, while the feet and hands, ankles and thighs, are often found to be unaffected. Again, the hypochondrium is another situation which increases in size, while the abdominal wall frequently contains masses of fat.* A sudden increase of weight,

* The condition known as "chronic subcutaneous fibrosis," about which Stockman of Glasgow has written, although not attributed to deficient thyroid secretion, nevertheless yields to the ingestion of thyroid extract. This condition is characterized by the increase of fat, but in the subcutaneous tissue are also found masses of fibrous tissue, and this latter tissue in this complaint is very tender to the touch.

without obvious cause, should make us at once suspect some thyroid disturbance.

Skin.—The skin is dry, rough, patchy in places, and may be in an eczematous condition. The internal surfaces of the tibiæ, the sternum, the forearm, and the back, are the main situations where ichthyotic rashes are seen. There is usually present in thyroid insufficiency an itching, which is sometimes so intolerable as to make the patient wish to scratch every part of his body at the same time. There may, on the other hand, be little or nothing to see, but the proof that the skin irritation is due to the diminution of thyroid secretion is that it gradually yields when thyroid feeding is instituted, although not until some time after the beginning of the treatment. If the hair be examined, it will be found to be dull and without its usual lustre, sparse in places, prematurely grey (in cases where the condition has persisted for some time), with patches of alopecia. Léopold-Levi and H. de Rothschild draw attention to the "eyebrow sign"—*i.e.*, the scarcity of hair in the eyebrows, with a marked diminution of the outer third of each eyebrow. This sign the present writer believes to be fairly constant, as he has observed it in many patients showing other signs of thyroid deficiency.

Constipation.—The subjects of this condition are nearly always constipated. As in thyroid excess the reverse is the case and the motions are on the loose side and frequent, so in submyxœdema are the bowels costive.

64 THE ORGANS OF INTERNAL SECRETION

Other Characteristic Features.—If we can describe a type of "thyroid deficient"—that is to say, subjects of long-standing submyxœdema—we must lay emphasis on one or two points. The individuals are usually small, with deepset eyes, scanty hair, dry skin, and with the appearance of premature senility. They require abundant sleep, and are particularly liable to somnolence after meals.

They are inclined to obesity, with the deposits of fat which characterize this condition, thickset, with an expressionless face, possibly carious teeth, and gingivitis or actual pyorrhœa.

Again, the subjects of thyroid deficiency are very prone to fatigue, which develops without an undue expenditure of energy. In this way they resemble the neurasthenic, just as in their somnolence after food they remind the observer of the lithæmic subject. As we pointed out in a previous chapter, these patients frequently exhibit signs of premature senility, the arteries may be thickened, the skin wrinkled, the joints have lost their suppleness, as they are wont to do in old age. We may refer in passing to the supposed relationship between the parathyroid glands and calcium metabolism, as it is assumed by many observers that derangements of these small glands permit the retention in the body of the lime salts, which are deposited in various parts of the body. It may be that thyroid deficiency is related in many of these patients with a parathyroid deficiency, and that it is the diminution in the latter secretion and not

in the former which causes the symptoms of premature senility.

It may be helpful at this stage of our study to refer to the work of Léopold-Levi and H. de Rothschild on this subject. These observers combine physiological research with clinical utility, so that the two aspects of thyroid deficiency are, so to speak, brought into focus.² These authors refer to "thyroid instability"—*i.e.*, to the liability to disturbance in the normal working of this gland, whether in the direction of excess or that of deficiency. They assert that a simple excess (slight hyperthyroidism) is very liable to be succeeded by a deficiency, and that the two conditions,—hyperthyroidism and subthyroidism—are frequently associated in the same individual. Thus, when these two conditions are present in the same patient, the prominent features of both might equally well be noticed. Those pointing to excess of the secretion of the gland, such as palpitation, nervousness, exophthalmos, and tremors, are combined with those indicating deficiency, such as constipation, chilliness, sclerodermia, scanty hair, etc. In another place³ Léopold-Levi refers to the "hyperthyroidism associated with thyroid deficiency," which he says may be classified as follows:

(a) Paroxysmal derangements of a hyperthyroidic nature, which appear as simple reactions and may be classed as symptoms.

(b) Reactions of a more complex nature, the mani-

66 THE ORGANS OF INTERNAL SECRETION

festation of which includes other symptoms, and which may be classed as syndromes.

(c) Reactions affecting the thyroid gland itself (endogenous goitre).

To emphasize the characteristics which we have already discussed as being present in thyroid deficiency, we will quote from this same article what this author gives as being the symptoms of a "slight degree of hyperthyroidism," as by this means we shall serve to impress the two different clinical pictures.

1. Hypertrichosis, more particularly of the eyebrow.
2. Hyperthermia, flushings, febricula of thyroid origin.
3. Acro-erythrosis, with cutaneous humidity.
4. Tendency to diarrhœa, the stools being soft and frequent.
5. Great physical restlessness, with a sensation of being hurried.
6. Insomnia.
7. Emaciation.
8. Excessive height.
9. Developmental precocity.
10. The syndrome which I have named "syndrome of persistent juvenility."
11. Extreme nervous irritability, with cardiac and cutaneous excitability, etc.
12. Large brilliant protruding eyes, with nystagmic movements.
13. The reactions are generally excessive, and there is exaggeration of the nutritional exchanges.

14. Hyperplasia of the entire thyroid gland.⁴

It is, however, quite obvious that many of these symptoms can only be present in those patients where the condition has been congenital, has commenced during adolescence, or has been in existence for a very long time; for, to take only one example, the stature could scarcely be altered in an adult patient who develops the condition after, let us say, a severe illness. But, on the other hand, there can be little doubt that many of the symptoms are quite constant, and can be observed in a large proportion of cases. At the present time the writer of this book has under his observation a lady who is undoubtedly suffering from deficiency of the thyroid, but has also the syndrome which points to excessive action of this gland. Thus, while she has typical deposits of fat, ichthyotic skin, characteristic hair, somnolence, and mental turpitude, she also has slight tachycardia, rather prominent eyes, an almost unnoticeable tremor, some degree of nervousness, and transitory restlessness. In fact, in this case the picture changes from time to time—sometimes the signs of excess in the thyroid secretion being uppermost, at other times the signs of deficiency being more noticeable. There can be little doubt that in many cases both conditions are present, but not necessarily at the same time. It is quite possible that a hyperthyroidism succeeds a transitory deficiency, that while this condition is present the signs of its presence are marked, but that the reaction brings with it exactly opposite symptoms.

68 THE ORGANS OF INTERNAL SECRETION

The view expressed by Léopold-Levi—namely, that the thyroid can be in an unstable condition, so that both these two opposite syndromes can be combined—is one which is supported by many observers. He believes, however, that the two conditions are rarely combined at the same time, but that the fundamental state is one of "thyroid instability." This, however, may constitute a condition in which the thyroid wobbles, so to speak, between outbursts of hard work and spasms of idleness. About such a state he says:

"Associated with thyroid insufficiency is a large number of syndromes, the characteristic features of which are that they are precipitated suddenly in the form of thyroid crises, and that they yield to treatment with thyroid extract.

"Although associated with thyroid inadequacy, these syndromes are the clinical expression of a hyperthyroidism which is, in itself, the reactionary manifestation of the fundamental subthyroidic condition.

"They are—migraine, ophthalmic migraine, asthma, nasal asthma, attacks of chronic rheumatism and gout, mucous enteritis, urticaria and other skin affections (acne, eczema), mental symptoms."⁵

From this quotation the reader will see that the alteration in the thyroid is believed, by this authority at least, to present many features which are sufficiently capable of recognition to enable the diagnostician to arrive at an accurate judgment. Before closing this chapter, however, it may be well to remember

that not all these symptoms can be expected to be present in the same patient. Rather must the physician be on the lookout for any of them, remembering that, so far as the endocrinous glands are concerned, diagnosis must be aided by ingenious detection.

REFERENCES.

- ¹ See Introduction, pp. 3-6.
- ² See *La Petite Insuffisance Thyroïdienne et son Traitement*, by Léopold-Levi and H. de Rothschild, published by O. Doin et Fils, Paris, 1913.
- ³ *Practitioner*, vol. xciv., No. 2, p. 211.
- ⁴ *Ibid.*, vol. xciv., No. 2, p. 212.
- ⁵ *Ibid.*, p. 213.

CHAPTER IV

THYROID DEFICIENCY (*Continued*)

WE have now discussed the broad outlines of thyroid excess and thyroid deficiency. But this latter subject is of such importance, if we are to use to the full the powers which a knowledge of organo-therapy gives us, that we propose to devote this chapter to a more detailed study of the signs and symptoms of hypothyroidism.

Owing to the importance of this subject, which, indeed, increases from day to day in vital importance as our knowledge of the lesions which may be attributed to disturbances in the hormonal system becomes wider, it is necessary for us to discuss at some length this aspect of the endocrinous glands. For thyroid deficiency is of the most vital interest to all students of medicine, if only that, when recognized early, many aberrations from normal health may be remedied, or even averted, by the administration of extract of the thyroid gland.

For this reason alone it behoves us to make ourselves *au fait* with the more marked symptoms of this disorder, for many of these will yield to rational therapy. Again, as in every other treatment, there

is a right and a wrong way of proceeding with thyroid administration, but, unfortunately, this fact is not widely appreciated. Probably from the erroneous impression as to the dosage which is given in textbooks of medicine and in posological tables, the dose of thyroid extract is usually far too high. In any case it is necessary to proceed cautiously when prescribing this drug, for the personal factor is never more important than in thyroid therapy. We shall discuss this in more detail at the end of this chapter.

In the last chapter we reviewed the opinions which are held as to the symptomatology of thyroid deficiency, and quoted the views of Léopold-Levi and H. de Rothschild. We referred to "thyroid instability," to the syndromes which these authors believe to be characteristic of thyroid excess and thyroid deficiency, and to their belief that the two conditions may be present successively or even concurrently in the same patient. The opinions expressed by these observers are important, for they have made a careful study of the thyroid gland and its disorders, and their conclusions are of the greatest help to anyone wishing to familiarize himself with thyroid dosage.

Harrower quotes in his book on "Practical Hormone-Therapy" from Léopold-Levi and H. de Rothschild's book on "La Petite Insuffisance Thyroïdienne."¹ These authors say: "Considerable importance attaches to minor thyroid insufficiency, for, unlike myxœdema, it is very frequent. It should be also especially interesting to the physician because of its usually

72 THE ORGANS OF INTERNAL SECRETION

rapid response to treatment. Clinical experience is the basis of these deductions, just as it was the original basis of the present knowledge of the stigmata of hypothyroidism."

Now, let us spend some time in studying those points which are generally considered to be typical of thyroidic insufficiency. We have already studied the main symptoms, and the manner in which normal physiological processes are altered in the absence or perversion of thyroid secretion. Our object now is to study the type which makes one look for symptoms rather than the symptoms themselves.

The following may be said to be points in bodily construction which suggest faulty thyroid secretion. As we have already said, these patients are small. Their stature is frequently diminutive, while the development of their soft structures is excessive in proportion. There is also present, according to Hertoghe, a relaxed condition of the articular ligaments, which permits of over-extension of the joints. Thus, we shall look for what is known in popular language as "double-jointed" people. The patient to whom we referred in the last chapter (in whom symptoms suggestive both of excess and insufficiency were present) showed this symptom remarkably well. She was able to bend her fingers back until the nails almost touched the back of the hand. In this connection, Hertoghe refers to a man who had been under his care for a number of years suffering from thyroid insufficiency, in whom there was spontaneous painless

dislocation of the patella. So impressed is Hertoghe with the changes in this respect that he says "it is possible, with very little practice, to judge of the degree of thyroid inadequacy merely by squeezing the patient's hand."² He says that these hands give the impression, when squeezed, of a "glove filled with clay."

The present writer is certainly of the opinion that the relaxation of the ligamentous structures in submyxœdema is something more than an artificial symptom. He has noticed it in a number of patients affected with this disease, in which the diagnosis was confirmed by treatment.

In children certain signs of thyroid deficiency have to be added to those which one must look for in an adult. Thus, while growth is impaired and stature infantile, we have learned to expect a protuberant abdomen, a lumbar lordosis, due to relaxation of the spinal ligaments, sometimes abnormalities of development, such as umbilical hernia, undescended testes, and so on.

Again, as such children grow up, they continue to show these and similar signs, which are more noticeable owing to adolescence. The sexual development is delayed, immature sexual organs only being present in many cases, the stature remains stunted, and there is faulty mental development. The exhibition of thyroid extract serves to increase stature, remedy faulty development in other directions, and even to abolish symptoms such as umbilical hernia. If omitted too soon, or if given in insufficient doses, the improvement is apt to cease and the growth to stop.

74 THE ORGANS OF INTERNAL SECRETION

This, then, is what to expect in subthyroidic children; if exaggerated, we meet cretinism. Such children are deficient in hair, and may even show more or less generalized alopecia. This, when combined with the infantile expression of the face, should at once make one suspect thyroid deficiency. The child may be small for his age; he may present one or more of the stigmata referred to above; and, finally, the diagnosis is clinched by the exhibition of thyroid extract and the resulting improvement.

So much, then, for thyroid deficiency in a child. What produces this is a more difficult question. There can be little doubt that in many cases the tendency is an inherited one. These children often come from stock which can be shown, on careful examination and questioning, to be subthyroidic. Hertoghe quotes a case of infantilism of the Lorain type, the mother of the child being a myxœdematous subject of a benign type.

Like many other diseases, prenatal conditions must be blamed for many of these cases, and it will have already been noticed how some of the defects, due to imperfect development, are more or less directly traceable to the thyroid gland, as they are remedied after birth by the administration of the extract of this gland. In support of this, witness the umbilical hernia in one of Hertoghe's cases, and note how it cleared up when the child had been taking thyroid for some time. Again, it will be remembered that such stigmata as imperfect development of the bones are among the

signs of inherited syphilis; while it has been shown³ that the administration of thyroid extract is as efficacious as iodide of potassium in healing tertiary syphilitic lesions.

Two other functions of the thyroid are pertinent to our study. First, we have to remember the theory which deals with the "antitoxic" power of this gland, and to bear in mind that, if this is true, the subjects of any kind of thyroid deficiency must needs be more open to bodily infection, either from so-called auto-genetic or heterogenetic sources. Second, as pointed out by Hertoghe, the life of each cell of the body, considered as an individual, is relatively a short one. It serves its purpose, degenerates, and is excreted from the body by one of the normal channels. For this to be performed, however, in a normal manner, the presence of thyroid secretion is necessary; for in its absence such cell elimination does not follow its usual course. Hence the vast increase in the subcutaneous tissue which is present in myxœdema, and to a less degree in submyxœdema. As is so well known, this gives rise to œdema, or, to be accurate, to increased subcutaneous tissue, which pits on pressure. Extract of thyroid gland diminishes this abnormal subcutaneous tissue, and in so doing lessens the bodily weight. The excretion of nitrogen is increased by this extract, so that the elimination of broken-down cells is restored to normal—another reason for the reduction in weight under thyroid feeding.

As we have said already, the submyxœdematous

76 THE ORGANS OF INTERNAL SECRETION

patient is habitually constipated, and this may have an effect to which we drew attention in a former chapter. These patients show many of the signs upon which Lane lays emphasis as indicative of intestinal stasis; in fact, one of these signs is atrophy of the thyroid. The question to be answered is this: Is the thyroid primarily responsible for the inertia of the intestine, or does this latter factor produce atrophy of the thyroid?

I have seen cases which would argue that Lane's view is the correct one; on the other hand, many patients impress one more in favour of the theory that the thyroid atrophy is responsible for the signs to which both the supporters of the alimentary theory and those who uphold the "primary thyroid deficiency" theory lay claim. A short time ago I saw a lady who was complaining of vague pains in the extremities, dry and itching skin, languor and lack of mental alertness, and constipation. On examination I found a swelling of the thyroid gland, a slow pulse, and an ichthyotic condition of the epidermis. Furthermore, there was a good deal of evidence of intestinal stasis, such as meteorism, offensive flatus, insufficient motions even after a purge, and digestive disturbances. To many of those students who have read Sir Arbuthnot Lane's latest work upon the subject of intestinal stasis, the signs indicative of intestinal stasis were all (I refer to the main signs, and not to the smaller indications) present. At the same time the student of opotherapy would find most of the signs which he

had learnt to regard as pathognomonic of thyroid deficiency present in this case. The stature was small, the hair atrophic, the skin dry, the pulse slow, the bowels costive. I need hardly enumerate the other well-known signs.

There I must leave the discussion of the primary cause, as there is insufficient evidence to warrant us in dogmatizing further on the why and the wherefore of thyroid atrophy.

In reading the mass of literature which has accumulated of recent years on this subject, the reader is struck with the many different signs which are believed by the majority of observers of organo-therapy to be caused by thyroid deficiency. Even if at first glance one is inclined to believe that the clinical picture of submyxœdema has been exaggerated, yet if one realizes the signs which a severe degree of thyroid deficiency produces, one can scarcely doubt that a minor degree might very well alter the individual, although in a slighter degree; indeed, in the very mild cases to so small a degree as to make the change all but imperceptible.

The main object of this chapter is to bring the practical points in connection with hypothyroidism before the notice of those whose time does not permit of the perusal of the latest physiological and clinical research, so that a greater percentage of patients who might be helped by organo-therapy (and who will in all probability be helped by no other means)

78 THE ORGANS OF INTERNAL SECRETION

shall benefit by our knowledge, which has increased every year since Murray's original discovery. The study of the changes wrought in the various parts of the body by an absence or diminution in the secretion of the thyroid are so numerous that it will be best to deal with them separately, classing them under the different bodily systems. We will therefore study them in this way.

The Influence of the Thyroid Secretion upon the Cell.

Throughout life the living cell requires certain hormones if it is to follow the normal cycle of life. Thus, if starved of thyroid its growth is impaired; administer to the individual organism the extract of this gland, and growth is resumed. This is shown by the behaviour of cretins to thyroid feeding; they improve in every way under such treatment, but relapse if it is stopped or diminished. We have already pointed out that broken-down cells which should be eliminated are retained and stored (in a form that originally was thought to be mucinous) if the thyroid secretion is diminished. Thus, the cycle of growth and decay are both interfered with if the endocrinous system is disturbed. In association with this, we must explain one or two important facts. The subject of thyroid deficiency is never "fit"; he is generally tired, he finds work a burden, and, whether his employment is physical or mental, he requires a "push" before he can make himself take up his tools. Again,

his mental apparatus fails him, his memory is unreliable, he cannot recall either recent or past events; his attention wanders; his concentration is never at its brightest when he wishes to solve a problem; and, in a word, he is sluggish mentally as he is languid physically. He is neurasthenoid.

The muscular system, in association with the subcutaneous, ligamentous, and fascial systems, undergoes infiltration; while the nerve cell "suffers derangement of nutrition." It becomes "infiltrated, and it also undergoes compression, as the result of the infiltration of the connective tissue surrounding it. Hence, the transmission of motor and sensory impulses, both voluntary and involuntary, though delayed, is in no sense abolished. The reflexes are tardy, but they are present."⁴

Therefore the cells of the nervous system are embarrassed in a double manner—from within and without. The infiltration of fascia, of muscle, and of sheath, must of necessity exaggerate the difficulty of exertion. This would apply both to physical and mental exertion.

Symptoms Referable to Nervous Tissue.

Apart from the mental sluggishness, which we have already described, these patients suffer from a variety of ailments more or less directly referable to the nervous system. Thus, they are very liable to headache, which is usually frontal, but occasionally occipital. The two forms, however, are rarely merged. The



MICROCOPY RESOLUTION TEST CHART

(ANSI and ISO TEST CHART No. 2)



4.5

5.0

5.6

6.3

7.1

8.0

9.0

10

11.2

12.5

14

16

18

20

22.5

25

28

31.5

36

40

45

50

56

63



APPLIED IMAGE Inc

1653 East Main Street
Rochester, New York 14609 USA
(716) 482 - 0300 - Phone
(716) 288 - 5989 - Fax

80 THE ORGANS OF INTERNAL SECRETION

headache of thyroid insufficiency is, in nearly every case, worse in the morning, improving as the day wears on.

Giddiness is another constant symptom, with which is associated, in some cases, a "swimming" sensation. Tinnitus aurium, either in the form of buzzing, roaring, whistling, or shrieking noises, is common, while these patients sometimes experience even more serious sensory disturbances, such as hearing voices, the ringing of bells, etc.⁶ In practice, many such cases have been benefited by the administration of thyroid extract. So far as I am aware, there is no evidence that true vertigo is seen, although I have prescribed this extract for one or two patients in whom the giddiness and dizziness were very similar to true vertigo.

We have already referred to the functional nervous changes which are seen in thyroid deficiency, of which somnolency is, perhaps, one of the most marked. These patients seem to be able to sleep very soundly at night, after every meal, and at any other time when the opportunity offers. Most subthyroidics complain of their marked tendency to slumber at all times, and the more intelligent find that it is sufficiently marked to be abnormal. In spite of this, these patients do not wake up refreshed, and they feel worse in the morning than they do at night. Hertoghe suggests that in all probability the thyroid neutralizes the toxins generated during the hours of activity during the night—that is, under normal physiological

conditions—but that this does not happen in the subject of thyroid deficiency.⁶

We have already dealt with the changes in the mental apparatus which must be looked for in submyxœdema; to the confusion of mind and the lack of mental initiative we must attribute the slowness of thought, as well as the lack of confidence, which these patients show. To these facts also we must attribute the undue exertion which any work demands from these patients.

The Subcutaneous and Muscular Tissue.

The thickening seen in thyroid inadequacy is characteristic. It involves all tissue, according to most observers, but in certain areas it is laid down more or less in "heaps." Thus, the nape of the neck down to the last cervical vertebra shows a thickening, while in the last-named area is almost always seen a mass of fat which resembles a lipoma. Particularly is this so in adult patients, more especially in the female sex. The head is held somewhat forward ("cassowary neck"), so that the cervical deposit appears more prominent than is really the case. The shoulders of such patients are broad, while the clavicular regions contain pads of fat.

Such deposits of fat as these are more or less dependent on sluggish circulation, and therefore regions such as the abdominal wall are apt to suffer. The flanks and back are, in adult patients, enlarged by rolls of fat, while the bones of the extremities are

82 THE ORGANS OF INTERNAL SECRETION

sunken in an envelope of adipose tissue. At the same time this deposit is not solely fatty, nevertheless it adds to the body-weight considerably and hampers bodily activity. The entire subcutaneous integument appears to be thickened, and this thickening can be made to pit on pressure, although the pitting disappears as soon as the pressure is relaxed.

As regards the changes in the muscular system, we can by reason of these changes explain several symptoms from which these patients suffer. The muscles are burdened with fat and mucinous infiltration, so that their size is increased and their drainage insufficient. Muscular movement is hampered, so that the metabolic exchanges are even more restricted. Hence the dislike of active exercise which these patients invariably show. Again, the increase in the volume of the musculature leads to discomfort, if not actual pain, which latter may be vastly relieved by the exhibition of thyroid extract.

It is well known that children who are the subjects of thyroid inadequacy are very prone to the development of adenoids, and that this is one of the characteristic stigmata of this condition. Again, in such subjects the enlargement of lymphatic glands is common, and some thickening of these structures is frequently observed.

Sweating is practically unknown in submyxœdema, *per se*, and in consequence there is some increase in the secretion of the kidneys, with a tendency to nocturnal enuresis. The solids of the urine are, of course,

diminished, but increase enormously when the patient is taking thyroid. The nocturnal enuresis from which these patients suffer is in part, at any rate, due to the extreme heaviness with which they sleep; and for some years now thyroid extract has been prescribed empirically for this condition, although the underlying causation was not understood in the same way as it now is. This medication has given very good results in many cases, but it is almost infallible where the underlying condition is one of thyroid inadequacy.

The absence of adequate secretion from the skin—and these subjects rarely sweat—results in a stagnant condition of the liver; and, according to Hertoghe, this results in itching of the skin, the icterus of myxœdema. It may also assist in the intense drowsiness, if not also in the slow pulse; but even if so, it would only be a contributory factor. It is, however, quite conceivable that the constipation is due to diminution of the normal biliary secretion in the intestine, the rational natural laxative.

The Osseous and Cartilaginous Systems.

The delayed growth resulting from a deficient thyroid secretion is too well known to demand lengthy description. So long as the epiphyses are not ossified, there remains the possibility of renewal of growth under thyroid feeding. This can be ascertained by the Röntgen rays.

Again, myxœdema produces its characteristic in-

84 THE ORGANS OF INTERNAL SECRETION

filtration in the joints and their ligamentous structures. This causes crackling in the joints, and gives the impression of a rheumatoid condition. When this sign is present, therefore, a careful search should be made for other symptoms pointing to thyroid deficiency. It has been said that delayed union after fracture is very common in subthyroidic subjects, and that the bone unites on the exhibition of thyroid extract.

Epidermal Symptoms.

The dry, rough, and ichthyotic skin which characterizes these subjects has already been described. From their point of view, this is probably one of the most troublesome features, as the inconvenience of a generalized itching cannot well be exaggerated. There is little to be seen on examining the skin of these patients beyond what we have already described. The hair suffers in common with other epidermal appendages. The nails are brittle and grooved, or they may be actually cracked. As we have already said, the "eyebrow" sign, first described by Léopold-Levi and H. de Rothschild, is of considerable diagnostic significance. There is a sparseness of the outer third of each eyebrow, with a scarceness and partial falling out of the whole eyebrow. This feature can be seen in many subthyroidics, and should always be remembered when looking for confirmatory evidence of a suspected inadequacy of this secretion.

Pigmentation of the skin, perhaps more commonly

seen on the forehead, and raising of the eyebrows, are both characteristic; while the lips are usually thickened, and the mouth tends to open. The changes in the facial characteristics give an unusually stupid appearance to the patient, but this changes under thyroid feeding. The more marked cases of thyroid insufficiency develop countenances which resemble nothing so much as a porker; and the change which is brought about in the expression as treatment proceeds is most gratifying to the patient himself, as well as his relatives and friends.

In adult subthyroidics the hair is prematurely grey, and this change in colour is usually more marked over the temples. In the case which I published in the *British Medical Journal* of June 20, 1914, this feature was well shown, although the patient was a young man about twenty-eight years of age. If the hair is a normal colour, it is frequently without gloss, and is brittle and "lifeless." Patches of alopecia are common, and the scalp as a whole is dry and scaly.

It must be remembered that pigmentation of the skin is seen in thyroid excess as well as in thyroid deficiency, and that it is a characteristic feature of Addison's disease.* As a diagnostic symptom, therefore, it is not of great value.

* See a case of marked facial pigmentation which is described in Chapter X., pp. 199-201.

Sexual Symptoms.

During childhood the diminution or absence of the thyroid secretion prevents the normal development of the genital organs at puberty. In fact, such patients grow up to adult years without any change from the infantile character of these organs. Indeed, in some cases more serious defects are seen; for cryptorchidism, either complete or unilateral, may be present. In cretinism, of course, there is no development of any sexual characteristics, and the child remains in this respect unchanged.

Graves' disease produces a diminution of the periods, either with scanty menstruation or amenorrhœa. The reverse is true in submyxœdema, for here there is a tendency towards menorrhagia, although the periods do not as a rule develop until the patient is well past the usual time. The uterus remains undeveloped, or at all events partially so, while the ovaries are likewise immature. There is sometimes a marked retroversion.

The changes brought about in thyroid inadequacy by pregnancy are for the best; while the patient is pregnant the thyroid secretion is increased. But, unfortunately, this improvement is not permanent, and she retrogresses, so far as this is concerned, after delivery. Immediately after delivery there is too much secretion, but later this ceases, and the former condition of subthyroidism is reverted to. According to Hertoghe, some of the increased thyroid secretion

is directed towards establishing uterine involution, while another function of this important secretion at such times is to establish lactation.

Gastro-Intestinal Symptoms.

In considering the gastro-intestinal symptoms of thyroid inadequacy, we must refer again to the syndrome, which Lane attributes to intestinal stasis, but which observers in organo-therapy claim to be caused by the thyroid. "The symptoms are hypothermia, uncontrollable headache, rheumatoid pain and neuralgia, mental depression, dyspnoea, asthmatic attacks, premature greyness and baldness, dental caries, cholelithiasis, and brownish pigmentation of the skin." Thus Hertoghe refers to this syndrome, and he attributes the gastro-intestinal symptoms of thyroid insufficiency to infiltration of the muscular tissue of the intestines, causing partial paresis. This in turn leads to ptosis, or partial ptosis, and then we have kink formation and the other features so carefully described in Lane's book.⁷

We must emphasize, in passing, what we believe to be a very important aspect of this condition—namely, weakening of the abdominal wall, which fails to give that support upon which the intestines have come to rely. This may be due to degenerative processes of an infiltrative nature, or to some other cause, but in any case the weakening helps the intestinal derangements. This in turn affects the digestive functions;

88 THE ORGANS OF INTERNAL SECRETION

appetite, which is notoriously capricious in subthyroidics, is poor; digestion is imperfect, and assimilation lessened. These patients are always weak and incapable of sustained effort. This is probably due in part to the causes already enumerated, and partly to deficient assimilation of food.

The association between subthyroidic conditions and septic affections of the alimentary tract, such as appendicitis, has often been commented upon, while it is said that the relation between tonsillitis and appendicitis is a close one.

Rectal hæmorrhage is said to be common in this condition, and, according to Hertoghe, it is due to premature thickening of the rectal veins and to the deficient coagulation of the blood. Just as the monthly period is apt to be excessive, so is the hæmorrhage due to piles, etc., liable to be copious.

Treatment.

We have now outlined the main changes which may be observed in subthyroidic patients, and the chief signs by which the condition may be recognized. All that now remains to be done is to summarize our conclusions in a few words, and to outline the treatment.

Whether the condition is in evidence during childhood or not until after adolescence, the main features are characteristic. The change in the bodily temperature is constant (a subnormal temperature is nearly

always present), likewise the pulse is slow. These two points should always be investigated. Having determined the condition of the temperature and pulse, the skin and hair should be examined, and a careful investigation undertaken to determine the bodily weight and the characteristics of the subcutaneous tissue. The deposits of fat, if present, should be localized. On the other hand, the patient may not necessarily be obese if the degree of thyroid deficiency is slight. In such cases careful observation must be made to ascertain whether other signs are in evidence, and, if so, whether the sum of the signs and symptoms points to submyxœdema. The condition of the digestion and the bowels should be ascertained, the nature of the monthly period in women, the presence or absence of sensations of chilliness, of flushings, of pains in the limbs, and of rachialgia, determined.

Having reached this stage in the investigation, the further points which suggest themselves for examination deal with the minor changes which are characteristic of this disease. Any change in the voice, in the expression, or in the mental outlook, which either the patient or her relatives have noticed, is of importance. The functional efficiency of the patient should be studied, bodily weakness and fatigability examined, and the patient questioned as to the length of time these changes have been noticed.

All the minor points are important, as the question of treatment depends upon these. We cannot emphasize too strongly the fact that, so far as submyxœdema

90 THE ORGANS OF INTERNAL SECRETION

is concerned, the initial dose should be small. Even if this should be so small as to appear to the inexperienced prescriber to be useless, it is infinitely preferable to order a small dose, and to continue this over a lengthy period, than to commence with a big dose and have to suspend treatment on account of untoward reaction.

In severe cases—*i.e.*, where definite myxœdema is present—the initial dose may be larger; but such cases do not nowadays form the majority of patients, as they are usually far outnumbered by the patients who show the signs of minor degrees of hypothyroidism. Where myxœdema is present, the patient may be given one, two, or three 5-grain tablets of thyroid substance in a day; or, if it is preferred, a liquid preparation may be prescribed, such as Elixir Colloid (Squire) or the Liquor Thyroidei (B.P.). The dose of Elixir Colloid is from 1 to 2 drachms, but even smaller quantities may be given (1 drachm represents $1\frac{1}{2}$ drachms of the dry extract); while the Liquor Thyroidei is given in doses ranging from 5 to 15 drops. Needless to say, these doses must be given entirely according to the individual case, as under no circumstances should thyroid medication be undertaken by rule of thumb. Each case must be treated on its individual characteristics, the length of time during which the disease has been manifest, the degree of thyroid deficiency, and the alterations in the patient which this has produced.

When we come to the treatment of the lesser degrees

we cannot speak so definitely. The initial dose should always, as we have already indicated, be small. The drug should be increased after some days or weeks, and it should be regarded as a failure of technique if the prescriber ever has to reduce the dose owing to commencing with too large a quantity. He should, so to speak, feel his way, looking carefully for the changes which will show that the drug is doing its work. The temperature will approximate to normal almost from the first, certainly as the circulation feels the result of the additional thyroid extract. The pulse will increase in rate, and, if charted, will show a steady approximation to the normal. It must be remembered that these patients are often small, and that small people have, under normal conditions, a more rapid pulse than large individuals. When we find, therefore, that a small patient has a slow pulse, we should seek for the reason. Again, the subjective symptoms will improve under thyroid medication; the feelings of languor, the extreme chilliness, the depression, and the general discomfort, will diminish in severity.

Provided that such changes are looked for and noted as they appear, it will be safe to increase the dose should they be delayed or incomplete. The administration of thyroid is both an art and a science: an art, because of the skill required to apply correctly the knowledge gained of thyroid-therapy; and a science, because of the exactitude of such application, which is essential if we wish to achieve the best results.

92 THE ORGANS OF INTERNAL SECRETION

In cases of submyxœdema, it is far better to err on the side of caution—*i.e.*, of too small doses—than of rashness. The former can be remedied as the reaction of the patient indicates; while the latter may not only lose the faith of the patient in this remedy (which is, if the diagnosis be correct, the only remedy), but may even make the prescriber doubt his own judgment. One-tenth to one-quarter of a grain should form an initial dose, certainly in the majority of cases, which, when all is said and done, are in the main minor degrees of deficiency. This may be given two or three times a day, and increased to $\frac{1}{2}$ grain, 1 grain, or even 2 grains, three times a day. Generally speaking, this is quite a sufficiently large dose. There are cases of submyxœdema which require larger doses, but they should only attain to these after a long administration.

One other word of warning, and we may conclude. The prescriber must be able to rely upon his preparation. He must know that the percentage composition is constant, that it contains a due proportion of iodine, and that it is fresh and has not reposed on a shelf in a local chemist's shop for many moons. Thyroid extract degenerates, and when this happens it becomes inert; and it does not take long for this to happen. If the patient fails to react, we must remember that the quality of the drug may be at fault, and not the diagnosis. So long as the drug is fresh, it matters little whether it be given in the form of dry extract, liquid extract, or one of the many prettier preparations on the market. The prescriber

must know what he is prescribing, or he will be unable properly to check progress.

Finally, as the treatment proceeds, a careful watch must be kept upon the temperature, pulse, and weight, for mainly on these points can the supply of thyroid extract be regulated.

REFERENCES.

- ¹ Chapter XIII., p. 192.
- ² *Practitioner*, vol. xciv., No. 1, p. 30.
- ³ *The Newer Physiology in Surgical and General Practice*, by Rendle Short, p. 85.
- ⁴ *Practitioner*, vol. xciv., No. 1, pp. 40, 41.
- ⁵ Hertoghe quotes Murray in saying that these patients show a marked disinclination to discuss these symptoms, and asks whether this is due to the fact that such hallucinations show a marked resemblance to those associated with alcoholism. They are due, Hertoghe believes, to infiltration of the nervous centres, or to more local infiltration, such as thickening of the aural mucosa, Eustachian tubes, and the naso-pharynx.
- ⁶ In myxedema Brun and Mott have shown the presence of subacute general chromatolysis of nerve cells.
- ⁷ Chronic Intestinal Stasis.

CHAPTER V

THE PITUITARY BODY

THE existence of this body has been known for centuries, but until recently it has been regarded as of no functional importance—as a vestigial relic. Now, however, the very reverse has been shown to be the case, and since 1895 it has been known that this small gland exerts an active influence upon the general bodily metabolism. Its intimate functions are still far from clear, and experiments have not infrequently led to contrary results. However, the light is beginning to dawn, and clinicians are now reaping the benefits which laboratory workers have made possible by their investigations of the structure and functions of this gland.

These investigations have been made largely with a view to ascertain the results of extirpation experiments, and the conclusion arrived at is this, that, far from being a vestigial remnant and unnecessary to life, its influence upon the health is of prime importance, and that its normal functioning is essential to health.

Its full relation to bodily efficiency has still to be explained, as there are aspects and features about

which we should like more enlightenment. The French were wont to refer to the pituitary as "l'organ énigmatique," and its mysteries more than justify such a name. The ancients, while cognizant of its existence, regarded it from a different standpoint. Thus, Galen and Vesalius believed that it was concerned in the formation of the nasal secretion; hence the name "pituitary." Vieussens and Sylvius considered that it was concerned with the manufacture of the cerebro-spinal fluid. The first hint as to its real functions—namely, the manufacture of an internal secretion—is conveyed in a paper to which Cushing calls attention. This was written by Lower, and was called "Dissertatio de Origine Catarrhi," 1672, and contained the statement: "For whatever serum is separated into the ventricles of the brain and tissues out of them through the infundibulum to the glandula pituitaria distills not upon the palate, but is poured again into the blood and mixed with it." Here, then, we have the precursor of our knowledge of hormone-therapy.

As the space at our disposal is limited, we must confine our remarks to the more important and substantiated facts about the pituitary body; and these we will study from the standpoints of structure, physiological functions, effects of excess or deficiency of the secretion of the pituitary, and therapeutic considerations.

Structure of the Pituitary Body.

The pituitary body consists of three parts: an anterior part (glandular), an intermediate portion, and a posterior or nervous part. The anterior part is composed of a network of epithelial cells, between which run many wide bloodvessels with thin walls. This part of the gland is derived from the buccal invagination (that portion known as "Rathke's pouch"). As Swale Vincent points out, the structure is similar to the adrenal cortex, the islets of Langerhans, the thyroid, the thymus (in its epithelial stage), and the interstitial tissue of the ovary and testis.¹ Next to this anterior part comes the pars intermedia, which, while it is derived from Rathke's pouch, and resembles the pars glandularis in structure, differs from it in that "its cells are less glandular and its bloodvessels much less numerous."² This portion of the gland is separated from the pars anterior by a well-defined cleft; but it is not always markedly separate from the pars posterior. This latter portion of the gland is derived from the infundibulum, or stalk of the pituitary. This is attached by its terminal portion to the floor of the third ventricle; while its other extremity expands and forms the posterior part of the pituitary body. This is known as the pars nervosa, and consists of neuroglial fibres and cells.

Histologically the anterior part of the gland consists, as already stated, of "trabecular masses of epithelium-like cells between which are very numerous sinus-like

blood-capillaries lying in intimate relation to the cells, which are, indeed, sometimes set closely round the blood-spaces."³ The pars intermedia resembles the pars anterior in general formation, but its cells are less granular and it is not so vascular. The pars posterior is formed almost entirely of neuroglial fibres, with neuroglial cells scattered between the fibres. Between these fibres is found a hyaline matter, which is more abundant in the neighbourhood of the stalk. Schäfer maintains that there can be little doubt that the activity of extracts of this part of the gland is due to this substance. Herring denies that any nerve cells are present in the pars posterior, and believes that there are very few nerve fibres.

Physiological Functions.

In studying the physiological action of extracts of the pituitary, we are faced with certain apparent contradictions. If an extract of the whole gland be injected, there is an immediate rise of blood-pressure, associated with a slowing of the heart's action. In this connection the effect of pituitary is similar to that produced by adrenin. The arterial system is affected by this extract, as is all the involuntary tissue of the body—in other words, vaso-constriction is increased; and it is this property which makes pituitary extract such a valuable remedy in shock. We shall refer to this factor when discussing the therapeutic aspect of extracts of the pituitary body.

98 THE ORGANS OF INTERNAL SECRETION

Certain observers believe that the substance which exerts this tonic action exists in the posterior part of the gland only⁴ (which includes in this connection the pars intermedia as well as the pars nervosa), and, in point of fact, it has been shown that this effect is produced by either the pars intermedia or the pars posterior, but more by the latter than by the former.

As well as this tonic effect upon the heart and the bloodvessels (which, while it resembles that produced by adrenin, differs from it in that this latter extract causes increase in the rapidity of the heart's beat, whereas pituitary extract causes slowing of the pulse), pituitary extract acts upon other plain muscle, such as the stomach, the intestines, the uterus, and the bladder.

As opposed to this general stimulation, there are one or two notable exceptions. The kidney, when stimulated by the administration of extract of the pituitary, reacts in a different manner. Thus, the renal arteries dilate, there is diuresis, and this is partly due "to a selective influence of one of the contained hormones upon the cells of the renal tubules. . . . This action has been supposed to be due to a special diuretic hormone from the pars intermedia, and it is not as yet certain that this is not the case; but it seems more probable that all the effects that it causes are due to the active principles of the posterior lobe."⁵

Again, the effects produced upon lactation by

injections of this gland are interesting. Ott and Scott, in 1910-11, pointed out that it possessed a marked galactagogue action, and since that time many observers have supported this statement. There have, however, been a few dissentients, who maintain that extracts of pituitary have no such effect; these are, however, in the minority.*

One other feature, which at first sight appears rather contradictory, is the effect of pituitary extract upon amenorrhœa. It has been on several occasions utilized to counteract cessation of menstruation, and with marked success. This is, of course, contrary to what we should expect, but can only be explained, as suggested by Harrower, on the assumption that the gland is acting vicariously in carrying on this function. It is known that the pituitary undergoes hypoplasia during pregnancy, and Fromme tried injecting an extract of the gland in amenorrhœa, with some degree of success.

Now as regards the active principles of the pituitary. First of all in importance, certainly from the therapeutic standpoint, is the fact that the majority of observers are agreed that the posterior part of the gland contains the active principle. In other words, better and more constant results have been obtained by the use of some preparation of this part of the

* Schäfer maintains that the effect of pituitary extract upon lactation is one of the most striking experiments in physiology; but Heaney does not believe in the galactogogic effect of pituitary extract (quoted in Harrower's "Practical Hormone-Therapy").

gland than by the exhibition of extracts of the whole gland or by the attempted isolation of extracts of the other portions. Harrower states that a definite salt has been isolated from the pituitary—*hypophysin sulphate*—which is available for use, and standardized in 1 in 1,000 solutions. This has been utilized by many observers, who report favourably on its use (Hertzberg, Fuhner, and Harrower).

The Effects Produced by Disease of the Pituitary Body.

In discussing the thyroid gland, we saw that morbid conditions due to its faulty functioning might be classified into those due to hypersecretion (hyperthyroidism), and those due to deficient secretion (hypothyroidism). In like manner, when discussing diseases due to abnormal working of the hypophysis cerebri, we may adopt this division—namely, diseases due to excessive action of the pituitary and those due to deficiency.

The best recognized disease associated with this gland is that to which Marie gave the name "acromegaly" (*ἄκρον* = point; *μέγας* = large). It is now believed that this condition owes its origin to a hypersecretion of the pituitary, and that part of the symptoms are directly due to an excess of the hormone elaborated by the anterior part of this gland.* It is

* Schäfer says: "That the acromegalic skeletal growth is produced by hypertrophy and over-secretion of the anterior lobe is highly probable, both as the result of partial extirpations in animals and from the effect of operative removal of pituitary tumours in man" ("The Endocrinic Glands," p. 113).

also now generally believed that a common sequence in the production of this disease is a primary hyperpituitarism, the cause of which may or may not be clear, followed in some cases by a hypopituitarism.

Before proceeding to outline the features of these two clinical conditions, we will briefly describe the symptoms of acromegaly. This disease usually commences with a marked narrowing of the field of vision, so that ocular symptoms are a prominent part of the complaint in its early stages. This feature may be due to the presence of a tumour in the gland pressing upon the region of the optic chiasma. Sooner or later are developed the characteristic features of the complaint which caused Marie to describe it under the name "acromegaly." These consist in a marked overgrowth of the head and extremities. The face enlarges and coarsens in its expression; while the shape of the head tends to alter, becoming flatter and squarer. The fingers, if viewed by the X rays, show an enlargement of the bones, with a "mushrooming" of the terminal phalanges. The entire skeleton increases in size, so that the individual becomes more or less generally enlarged.

Certain other changes are common, and these develop concurrently with the skeletal enlargement. There is a tendency to hypertrichosis, with a general increase of the hair all over the body. Sexual activity diminishes, as a rule, early in the course of the disease; although, in those cases where hyperpituitarism is succeeded by hypopituitarism, sexual activity may

102 THE ORGANS OF INTERNAL SECRETION

be primarily increased, and subsequently diminished. A tendency to glycosuria is common, which, again, may be followed by a diminished glycogenesis. The integument is also thickened, in a somewhat similar manner to that seen in myxœdema; while if the disease commences before ossification of the epiphyses has taken place, the stature becomes enormous and the individual turns into a giant.

It is now generally believed that acromegaly is due to an overgrowth of the pituitary; indeed, in many cases actual tumours are present, and have been demonstrated post mortem. Sometimes, in these cases, the sella turcica can be shown by a skiagram to be definitely enlarged. On the other hand, after the glandular enlargement, a secondary atrophy may ensue, in which case the symptoms become those of hypopituitarism. It is at this stage that symptoms contradicting the earlier clinical picture may make their appearance, so that, whereas at first there may have been a tendency to undue glycogenesis, there is now increased sugar storage, and whereas at first there had been hypertrichosis, there is now a change in the hair, the distribution often tending towards the feminine type.

Again, it does not follow that every case of what Cushing calls dyspituitarism may commence as a hyperpituitarism, and subsequently change to a hypopituitarism; for some cases evidence symptoms pointing to the latter condition from the first. The state described by Frohlich, and named by Bartels "Dystrophia

adiposogenitalis," is essentially a typical example of hypopituitarism. The symptoms pointing to this state are, as is to be expected, the opposite of those typical of hyperpituitarism. Thus, the stature is small (if the disease commence before adolescence), adiposity is excessive, sexual development is delayed or deficient, the development of hair is wanting, and there is a tendency to the formation of feministic characteristics. Thus, the hair on the pubes is more localized than is the case in man, but the development of the hair on the head is frequently abundant. The sweat is usually deficient, although the skin exhibits a smooth and even texture.

Cushing deals very fully with the production of this state in his book, "The Pituitary Body and its Disorders." He believes that the tendency to the deposition of fat, which subjects of hypopituitarism show, owes its origin to deficiency of the secretion from the posterior lobe. Likewise the unusually high tolerance for sugar and the increased assimilation of carbohydrates are due to the same cause.

Again, in hypopituitarism there are other changes which may be expected in the symptoms. The temperature is usually subnormal, the pulse slow, the blood pressure low, and the mental attitude sluggish, with a tendency to somnolence. It will be seen that the clinical picture is not unlike that seen in myxœdema. The parallel is, in fact, a close one, certainly in many features.

It is important to remember that the symptoms of hypopituitarism will differ according to whether the

104 THE ORGANS OF INTERNAL SECRETION

deficiency first appears before adolescence, or is only in action after adult life has commenced. Once the ossification of the epiphyses has taken place, there is no possibility of further increase of stature. But the increase in the integuments, the alteration in the circulation, the changes in the mental outlook—these all point to the underlying cause. Schäfer says that the symptoms differ widely, according to whether the anterior or posterior lobe of the gland is mainly affected; if the lesion is concerned with the anterior part of the pituitary, the chief effects will be upon the stature, while affections of the posterior lobe produce fat formation and deficient sexual development. Needless to say, the two sometimes occur together; the important point to emphasize is that the anterior lobe in excess produces enlarged stature, in deficiency the reverse; while the posterior lobe concerns itself with the regulation of carbohydrate metabolism, and, if deficient in secretion, undue fat formation results *inter alia*. A total deficiency of the pituitary secretion, appearing during childhood, would produce a dwarfed stature,* with the other symptoms which are believed to be due to deficient posterior secretion. Moreover, it is possible that with one part of the gland functioning adequately, and the other deficient, the clinical picture might correspond to what is expected when the particular part is abnormal. Supposing the anterior part of the gland to be normal and the posterior deficient, we

* Cases have been recorded in which the pituitary gland has been found atrophied in dwarfs, although this is not invariable.

should expect a fully-grown individual, but with excessive deposits of fat, slow circulation, sluggish mentality, and the other features of hypopituitarism, with the exception of stunted skeletal development.

Although much of this particular study leads us into realms of speculation, nevertheless there is strong probability that these suppositions are near the truth. Although the pituitary is by no means always atrophied in dwarfs, this is not tantamount to saying that its secretion is normal qualitatively as well as quantitatively; for it might be enlarged by tumour formation, or by hypertrophy, without secreting the normal amount.

Here we must leave the discussion of abnormalities in the pituitary secretion, and discuss briefly the treatment of such disorders.

Treatment of Pituitary Disorders.

In a brief review of the treatment of pituitary disorders, or, rather, of such disorders as come within the scope of this chapter, we must perforce spend some time on the study of both surgical and medical treatment. For, as regards that class of disorder which is caused by a radical change in the structure of the gland or by a tumour interfering with the adequate functioning of the gland either by direct pressure, or by raising the intracranial tension, surgery is not always even a matter of choice, but a matter of necessity.

As we have already said, some cases of acromegaly

106 THE ORGANS OF INTERNAL SECRETION

have shown the existence of tumours of the pituitary gland which have upset its normal secretion. Thus, a cystic growth of the gland, when removed, may produce hypopituitarism, which, again, can be remedied in some cases by pituitary feeding. But previous to the operative measures the symptoms have naturally been those of excessive secretion, sometimes accompanied by localizing symptoms due to an increase of intracranial tension.

Cushing has reviewed the treatment of pituitary disorders very fully in his book,⁶ and he deals with the question of surgical interference and with the choice of treatment in a thorough manner. Apart from those cases which show the presence of a tumour, where surgical treatment is generally the only hope at the present state of our knowledge—*i.e.*, with the uncertainty of the minutiae of the diagnosis—we can scarcely recommend surgical interference for hyperpituitarism, on the same grounds on which some authorities recommend partial removal of the thyroid gland for hyperthyroidism. In this connection Cushing says:

“In view of the fact, therefore, that hyperpituitarism, so far as glandular secretion is concerned, is a condition which tends to right itself, it must remain for the time being a matter of uncertainty as to whether or not, in the absence of a degree of hyperplasia sufficient to cause neighbourhood symptoms, operative measures can hold out any promise of permanently controlling the disorder.

When, however, neighbourhood symptoms have

arisen owing to the extreme enlargement of the gland, due to the formation of an adenomatous struma, whether or not there have been antecedent symptoms of acromegaly, the surgical aspects of the matter stand on firmer ground.'⁷

Owing to the situation of the pituitary gland, and the difficulty of operation, we see, therefore, that its partial removal, in the absence of urgent symptoms, is not to be considered, even as a possibility, in the hope of benefiting symptoms of hyperpituitarism. Fortunately, there seems to be a good deal of evidence to point to the assumption that states of hypersecretion are often transitory, so that we must treat the early stages of such a condition with care and watchfulness. It is opposed to our present knowledge to administer extract of the pituitary gland to patients suffering from acromegaly; for as thyroid extract makes patients suffering from Graves' disease worse, so does pituitary extract affect acromegalics.⁸ The subsequent hypopituitarism may well be met by organo-therapy; sometimes, indeed, one feels tempted to employ the extracts of other of the endocrine glands in the treatment of hyperpituitarism, in the hope that the hormone balance will be restored. In the present state of our knowledge, however, this latter form of therapy for this disorder must needs be empirical; nevertheless, much has in the past been achieved by empiricism, and doubtless in the future much more will have to be placed to its credit.

Turning now to the study of the treatment of hypopituitarism, we find that medicine naturally takes a

more prominent place than surgery. The administration of pituitary substance may be carried out in three ways—namely, by ingestion, by injection, or by grafting. For simplicity of administration, the first is undoubtedly the method of choice, and consequently is the one mainly relied upon in actual practice. From the experimental standpoint, Goetsch found that the subcutaneous injection of the extract in dogs was more effective in the ratio of 4 to 1 than the oral administration, while the intravenous injection was even more effective than the subcutaneous in the ratio of 2 to 1. These tests were made with the standard devised by Cushing, from whose book they are quoted. This standard is based upon the increased tolerance for sugar which patients suffering from hypopituitarism exhibit; for in hypopituitarism "the rational dosage of glandular extract to be administered by mouth can possibly be determined by giving the individual daily an amount of glucose or lævulose sufficient to produce a temporary mellituria in a normal individual of equal bodily weight; meanwhile an increasing amount of the extract is administered daily, until, under the conditions of increased carbohydrate tolerance which the patient exhibits, hyperglycæmia occurs, with a trace of sugar in the urine."⁹

Having arrived, then, at the diagnosis of hypopituitarism, we next have to decide upon the treatment, both as to the method to adopt and as to the particular preparation to employ. We are concerned here more with the routine administration of pituitary extract (in

one of its many forms) than we are with its employment in urgent conditions, such as shock, obstetrical emergencies, etc.; so that this account must be understood to deal chiefly with the treatment of definite hypopituitarism.

If it is decided that pituitary extract is required over a lengthy period, it is far preferable to administer it by the mouth, as there are many disadvantages attached to both the other methods. Where it has to be given repeatedly by injection, the intramuscular method is the one for choice, as subcutaneous injection is undesirable on account of the local anæmia produced. For a very rapid result, it may be administered by intravenous injection well diluted with normal saline solution.

As to the various preparations which are on the market, a wide selection is before the prescriber. *Pituitrin* may be administered by intramuscular injection or by the mouth. The dose in the latter method is from 10 to 30 minims.

This preparation is said to be especially cardio-tonic in action, "and in a less degree hypertensive."¹⁰ Again, Hypophysin (Fuhner) is stated to possess, from the physiological standpoint, "the essential properties of the extract of the macerated posterior lobe."¹¹ These and other liquid preparations are usually put up in ampoules, should it be desired to administer them hypodermically or intramuscularly. The dose of the liquid preparations is usually from 8 to 16 minims. A preparation which will be found useful in practice is

110 THE ORGANS OF INTERNAL SECRETION

the Elixir Hypophysis (Squire), which is manufactured from the whole gland, 1 teaspoonful representing $\frac{1}{2}$ grain of the dried and powdered extract. The dose is from 1 to 2 drachms two or three times daily.

Again, pituitary extract is manufactured in solid form, either as a separate preparation or in combination with other of the endocrine gland extracts. These products are very numerous, and it would be tedious to mention them in detail; suffice it to say that most of the wholesale chemists manufacture such extracts in tablet or powder form.

It is, of course, as important in dealing with this extract as it is in dealing with thyroid extract that there should be a known standard of strength, and that the prescriber should be sure that his preparation is fresh. As a rule this is easily ascertained, and the dosage regulated with some degree of certainty.

Many manufacturers place on the market preparations specified as having been made from the anterior or posterior part solely, and thus another variety of this treatment is introduced. The majority of cases appear to indicate treatment with posterior extract; indeed, the majority of the preparations are expressly stated to have been manufactured from the pars posterior.

Now as to small points in the treatment of pituitary disorders, and also in the utilization of this extract in counteracting symptoms. First in importance is undoubtedly the contra-indication. As pituitary extract raises the arterial tension, it (or any preparation con-

taining it) should be carefully avoided in all states indicating hypertension. It should be administered with caution over long periods, and frequent sphygmomanometric readings taken to control any undue rise in blood-pressure. The present writer has known it work wonders in some neurasthenic conditions, likewise in states of weakness following long-continued strain, in impotence and such disorders. Again, in delayed convalescence, in persistent low blood-pressure, in depressive states, and the like, a course of pituitary extract, wisely controlled, will frequently be found to be invaluable.

The requisite dose varies enormously, and so wide is the range that it appears probable that some at least of the failures are due to the dose being unsuitable. In speaking of the dosage required in some of these cases, Cushing emphasizes the vast range of dose which is effectual in different cases. Thus, one boy exhibiting the signs of hypopituitarism in a marked manner underwent a "complete mental, moral, and physical awakening" by taking 18 grains of whole gland preparation daily. But another case quoted by the same author required as much as 300 grains daily (a prohibitive dose, as Cushing remarks) to give the same subjective benefits which other patients experienced with far smaller doses.

In undertaking the treatment of a patient by means of pituitary extract, it is wise to progress slowly, and to endeavour to ascertain the dosage required by the individual case, as well as the reaction to the

112 THE ORGANS OF INTERNAL SECRETION

extract. In practice, it is often necessary to rest content with methods which are, perhaps, not so precise as those which are open to laboratory workers; and it is therefore often only possible to watch carefully the progress made under the treatment, at the same time looking for any symptoms the amelioration of which may justifiably be claimed to be due to the drug.

Cases are sometimes encountered, however, especially in neurasthenic subjects, in which irritability is produced by the administration of pituitary extract. Often this appears to be due to the rise in blood-pressure which follows such treatment; but it is well to be careful to reduce the dose should this symptom appear, or even to cease its administration altogether, and after a few days' cessation to recommence the treatment with small doses. This will generally be successful in combating the irritability, and will enable the treatment to be persevered with after the rest.

In Chapter X. the use of pituitary extract in combination with other organic extracts is fully discussed, also the joint administration of this substance and thyroid extract is considered.

REFERENCES.

- ¹ Swale Vincent, *Practitioner*, vol. xciv., No. 1, p. 149.
- ² Schäfer, *The Endocrine Glands*, p. 77.
- ³ *Ibid.*
- ⁴ Howell, *Jour. Exper. Med.*, 1898, iii. 245-258.
- ⁵ Harrower, *Practical Hormone-Therapy*, p. 305.
- ⁶ H. Cushing, p. 294 *et seq.*
- ⁷ *Ibid.*
- ⁸ Experiments of Renon and Delile, quoted by Cushing, p. 315
- ⁹ Cushing, pp. 317, 318.
- ¹⁰ Harrower, p. 313.
- ¹¹ *Ibid.*

CHAPTER VI

THE ADRENAL GLANDS

THE study of the chromaffin system involves a careful investigation of its most important member—namely, the adrenal or suprarenal glands. These bodies are now known to possess such vital properties in the body physiology, and to perform such an indispensable part in the maintenance of the “tonic” state of all vascular and plain muscular structures, that they deserve a foremost place in any study of the endocrine glands.

Although as long ago as the sixteenth century the great anatomist Eustachius discovered the existence of the adrenal bodies, their importance remained unknown until recent years. In 1849 Addison discovered that, in some cases of illness characterized by definite pigmentation of the skin, the adrenal glands were found to be diseased, often with a tuberculous infection. From that day to almost fifty years after, our knowledge of these structures may be said to have increased but little; but in 1894 the famous discovery by Oliver and Schäfer, that extracts of the glands possessed a marked blood-pressure-raising property, inaugurated the interest which has led to the really remarkable discoveries of the properties which the

chromaffin system in general, and the adrenals in particular, possess.

Before commencing the study of the therapeutic value of extracts of the adrenals, we must briefly outline our knowledge of these glands, both from the anatomical and physiological standpoint.

Anatomy and Physiology of the Adrenal Glands.

The suprarenal glands are two flattened bodies more or less globular in shape, of a yellowish colour, and are situated behind the peritoneum in front of the upper part of each kidney. Their size in normal health varies in different individuals, but as a rule the left is slightly larger than the right, and situated somewhat higher up. The former is semilunar in shape, while the latter is more triangular, and somewhat resembles a "cocked hat."

In structure these glands are made up of a central portion, or medulla, and a peripheral part, or cortex. The former is composed of highly vascular cells, embedded in a venous plexus, which secrete adrenalin; owing to this property they stain more or less deeply with chromate salts, which has earned for them the name of "chromaffin cells." According to the depth of stain which the cells of the medulla take, the degree of activity of their secretion can be roughly estimated.

The medulla of the suprarenal gland has very close nervous relations, which, from a morphological standpoint, would be expected. For early in embryonic

life neuroblastic cells emigrate, and while some settle in front of the spinal cord, forming the ganglia of the sympathetic chain, others eventually become the visceral ganglia; while another group, not converted into nerve ganglia, remain in close connection with the kidney, forming the medullary cells of the adrenals, which secrete its active principle. As well as the medulla, which is formed in this manner, certain clumps of cells stray farther afield, forming accessory bodies external to the main gland.

The importance of these facts is shown by two characteristics of the gland; the first is that it is these cells, originally formed from the same embryological tissue as the great ganglia, which secrete adrenalin; the second is that this secretion stimulates only that plain muscle in the body which is supplied by the sympathetic system. That this is so can be verified by electrical stimulation of the sympathetic nerves, which corresponds exactly with the result of adrenalin stimulation.

The exact composition of adrenalin has been discovered by Takamine, to whom we owe the isolation of this substance; and he shows that it is ortho-dioxy-phenyl-ethanol-methylamine. Its secretion into the blood-stream is controlled by the splanchnic sympathetic nerves. Moreover, the amount of secretion has been shown to depend largely on other factors, both mental and physical. Thus, mental agitation—fright, emotion, hurried exertion, etc.—all exert an influence upon the secretion of this substance. It is

to the cells of the medulla, therefore, that we owe this important hormone—in this case a true hormone, as it is an excito-tonic chemical. These cells, it must be remembered, have an extremely intimate connection with the sympathetic nervous system; this is important, as we shall see when we discuss the effects of hypo-adrenia.

The other part of the gland is of different structure, and is composed of fatty material; a doubly refractive lipid is found in the cortex, with which this part of the gland is loaded. The cortex is very much larger than the medulla; in fact, it forms about 90 per cent, of the whole gland.

Included in the chromaffin system are the following bodies: The medulla of the suprarenal gland, the carotid gland, and the intercarotid body, the accessory adrenals, and some of the cells of the anterior lobe of the pituitary body, which give the same reaction to chromate salts.

While the medulla is in reality a part of the nervous system, at all events morphologically, the cortex is not in any way connected or controlled by the sympathetic nerves. From an embryological standpoint, its cells are derived from the same neighbourhood as the sexual glands, and as the testes or ovaries descend they carry with them processes from the same area. Elliott maintains that there is reason to believe that some buds of this nature are embedded in the sex gland itself, giving rise to the lutein cells of the ovary or the interstitial cells of the testis.¹

118 THE ORGANS OF INTERNAL SECRETION

The cortical cells, however, are glandular structures, and appear to supply some secretion which influences the growth and reproductive powers of the individual. This has been demonstrated clinically, for cases are on record where tumours of the adrenal cortex have been found in cases where the characteristics of the opposite sex have developed; in one girl, quoted by Elliott, the menses ceased, a beard commenced to grow, and the body took on masculine characters.

Adrenalin.

The active principle which is secreted by these glands has been named "adrenalin." It possesses certain very definite characteristics, and is a necessary secretion for the continuance of life, as loss of both adrenals is fatal, although one adrenal can be removed without any apparent ill effects.

With regard to adrenalin itself, its most striking characteristic is its power of raising blood-pressure, even when injected in small doses. Experimentally, about one-twentieth of a milligramme when injected into an animal is sufficient to cause a considerable rise in the sphygmomanometric reading; which, again, is even more noticeable if the vagus is put out of action, as this substance also causes a slowing of the pulse when injected. If this is done, the blood-pressure may rise in quite a remarkable manner. This effect is produced by a vaso-constricting action upon the walls of the peripheral bloodvessels. Oliver

and Schäfer showed, by means of oncometric readings, that the volume of the viscera was very markedly reduced by adrenalin, while the volume in the extremities was increased owing to the excess of blood reaching them from the splanchnic area.

A full account of the chemistry and physiology of adrenalin will be found in Biedl's book on the internal secretory organs, in Swale Vincent's book, and in many other studies of this subject. Sufficient has now been said to show that this secretion is very important in bodily metabolism, and exerts a considerable influence upon the cardio-vascular system. The cortex is supposed to have an internal secretory function, which is concerned in neutralizing the poisonous products of muscular activity. In other words, the medullary secretion exerts an angio-tonic influence, while the cortical is concerned with neutralizing toxins. There seems to be a good deal of evidence that the adrenal secretion is in some way concerned with muscular energy; it is a well-known fact that in Addison's disease, where hypo-adrenia exists, muscular asthenia is a marked symptom; also that muscular power is raised "after adrenal secretion is invoked, or after epinephrin is injected."² Where the adrenal glands are injured (which sometimes happens during operative procedures in the neighbourhood of the kidney) or diseased, there can be little doubt that muscular efficiency is diminished. Langlois has shown that the adrenals are concerned in the destruction of muscle poisons.³ So we are justified in assuming that one

120 THE ORGANS OF INTERNAL SECRETION

function of the secretion of these glands is to keep up cardio-vascular tone, while another is to neutralize poisons elaborated during muscular energy.

It will have been noticed that we have laid stress here upon the intimate relation between the adrenals and the sympathetic system. The present writer has on several occasions laid stress upon the sympathetic symptoms so commonly present in neurasthenia, and has hazarded a theory that in many instances of this condition the underlying cause is sympathetic involvement.⁴ The low blood-pressure, the generalized asthenia, the vasomotor symptoms, are, certainly in the majority of cases of true neurasthenia, the most striking features.

Following up this line of argument, we should expect to see much improvement in these patients after the administration of a vaso-tonic hormone, such as adrenalin. And in many cases, in the experience of the writer, this has happened. Doubtless some patients suffer from a combined endocrinous derangement, and these would not of necessity benefit from adrenal administration alone. But, nevertheless, the similarity between hypo-adrenia and neurasthenia is sufficiently striking to be noted.

The features of a disorganization of the adrenal system are daily receiving more attention, and Sargent "has distinguished three different types of adrenal insufficiency—the chronic, the subacute, and the acute."⁵ He maintains that many of the manifestations of acute illnesses are due to adrenal insufficiency. It is

obvious that such features might conceivably be due to this cause, and when further light has been shed upon the endocrine glands, it is probable that we shall have to revise many of our existing views on the manifestations which accompany febrile disturbances. We shall certainly have a better opportunity of explaining factors hitherto regarded as necessary but inexplicable features of many diseases.

Adrenal Insufficiency.

Now for a little fuller description of the signs and symptoms of adrenal insufficiency. Of prime importance is a low blood-pressure; an asthenic condition, of mind and body; a subnormal temperature; a headache of a "vacuum" nature; and the dermographic sign described by Sargent.* This same observer refers to certain complications of convalescence, which he regards as due to a subacute inflammation of the adrenals; while other observers also lay stress upon the exhaustion which may supervene during the course of a fever. This may presumably be due to an exhaustion of the adrenals, produced by their efforts to safeguard the organism from toxæmia.⁶ It is quite possible that these small glands play an important part in all such febrile states, in neutralizing toxins, and in helping to maintain blood-pressure. In any case, it is

* This consists in a white line which appears on the skin of the abdomen when lightly stroked with the finger.

122 THE ORGANS OF INTERNAL SECRETION

stated that their administration in such conditions can be relied upon to counteract these effects.

The clinical state of the patient with Addison's disease needs no detailed description here. The flabby atony and muscular weakness, the vasomotor debility, as shown in the coldness of the skin surface, as well as the characteristic pigmentation, are too well known to require minute detailing. As Elliott points out, all the symptoms, except the pigmentation, can be explained by the vasomotor derangement and the paralysis of the sympathetic nerves.⁷

There is also in this disease gastro-intestinal disturbance, as shown by vomiting and diarrhoea; but the danger undoubtedly lies in the risk of heart-failure. Other symptoms are anorexia with deficient appetite, sometimes constipation, attacks of vertigo, syncopal attacks, subnormal temperature, with coldness of the extremities. There are frequently found a persistent thymus, an enlarged spleen, more or less generalized hypertrophy of the lymphoid tissue of the body, particularly of the stomach and intestines. There is usually some degree of wasting; indeed, in the more severe cases a considerable degree of emaciation may be present. These constitute the graver form of the disease. As a rule, although muscular strength is very markedly diminished, general nutrition is not affected. Death occurs from asthenia, delirium or convulsions, sudden syncope, or, in some cases, generalized miliary tubercle. The main features of Addison's disease are the markedly feeble pulse, the extreme asthenia, the pigmentation, and the gastro-intestinal disturbance.

The milder degrees of hypo-adrenia are those to which we have already referred, and they are frequently labelled "neurasthenia." That is to say that the main features are asthenia, vasomotor disturbances, and a low blood-pressure. In speaking of these forms, Harrower says: "The disorders of the adrenals, accompanied by anatomic changes, are fortunately not very common, while functional hypo-adrenia (sometimes termed 'hypo-epinephrinia'), varying very greatly in degree and the consequent manifestations, is of frequent occurrence. In general, the evidence of the presence of such conditions is lack of vascular tone, hypotension, myasthenia, and instability of the sympathetic nervous system. This class of disorders is still frequently overlooked, and, because the intimate relations of the endocrinous organs, almost invariably associated with, and complicated by, disturbances in other internal secretory organs."⁸

It might be expected that the extreme weakness and asthenia which are associated with this condition would be accompanied by great bodily wasting. But this is by no means always so; in fact, the bodily weight usually is not lost in any marked degree. It may be that this is due to an increased storage of fat or carbohydrate in the body, due to the diminished glycogenesis; for we know that an injection of adrenalin increases the conversion of glycogen into glucose, and may produce glycosuria; therefore a diminished adrenalin content in the blood-stream might prevent a utilization of the stored-up glycogen, and produce a diminished

combustion. This would be probable, seeing that the bodily heat is lowered, with an extreme degree of subjective feelings of chilliness.

Treatment.

Having now very briefly reviewed the symptoms of hypo-adrenia, we will turn to the study of the therapeutics of the adrenals. Opinion is divided as to the beneficial effects which result from the administration of adrenalin to patients suffering from gross deficiency of this hormone—*i.e.*, Addison's disease. Thus, Elliott says: "Neither with animal experiments nor in man has treatment with adrenalin been proved to be of value in prolonging life. I have used hypodermic and intravenous injections of adrenalin without obvious benefit. Still, the substance is worth a trial." On the other hand, Harrower says: "Addison's disease was naturally first among the disorders in which this form of treatment was tried; and while the results have been by no means uniform, numerous cases are recorded by such men as Oliver, Langlois, Osler, Robin, and others, in which the general condition was improved, pigmentation was diminished, and nutrition was benefited. In most cases the results were temporary, though Beclere obtained a complete and permanent cure in one case. He credits the treatment with causing a compensatory hypertrophy of the unaffected areas of adrenal tissue."⁹

Again, it has been suggested that in some cases the

causation is solely referable to the adrenal glands, while in others the sympathetic system is at fault, with or without adrenal disturbance; and that in the first class of case benefit is to be expected from adrenal therapy, while in the second we cannot and do not see improvement from the administration of the hormone. As it is generally believed that 80 per cent. of the patients who suffer from this disease are the subjects of tubercle, the general antitubercular treatment should be insisted upon, whatever hormonal treatment they receive. In any case, the dose of adrenal substance would have to be large, and Harrower says that as much as 2 or 3 grains of desiccated gland substance should be given three times a day.

We must believe, therefore, from the summation of evidence, that, so far as is at present known, the administration of adrenalin has not been the success in Addison's disease that it was expected to be. But the explanation probably is that, if the larger part of the gland is destroyed by tubercular or other mischief, the substitution of its secretion is not sufficient, and the disease progresses in intensity.

With regard to the utilization of adrenal extracts for other conditions, it has been tried and recommended for a variety of diseases. Thus, it has been administered as a vaso-tonic in cholera, with, according to the particular author, who believes that this disease is an "acute hypo-adrenia due to poisoning of the adrenals by toxins of the cholera spirillum," immediate and gratifying success.¹⁰

It has likewise been utilized in the vomiting of pregnancy, in the belief that the adrenals stand between the body and the toxæmias of pregnancy which arise in the chorionic villi. Again, in the condition known as "cyclic vomiting" it has been tried with apparent success.

It will be seen from these quotations that, so far, the diseases mentioned have been acute, and often present no known etiology. It is, therefore, more than possible that the chromaffin system is at fault. But there are encountered other conditions of a more chronic nature which appear to yield to adrenalin, and these are more particularly seen in children. Thus, a boy or girl who develops asthenia, in some cases after an acute specific illness, with languor, hypopiasis, and "backwardness" at school, is in all probability suffering from hypoadrenia. And it has been stated that administration of adrenal extract counteracts this condition, renders the patient more active, and banishes the indolence.

Again, after acute illnesses in adults, where asthenia and delayed convalescence is a marked feature, the exhibition of adrenalin should deserve consideration. It is not improbable that a disturbance of the adrenal hormone (which, it must be remembered, is a hormone in the real sense of the word) contributes to the slow recuperation and hinders recovery. Therefore, an extract of the adrenals seems to be indicated. Or, alternatively, a pluriglandular preparation should be selected if it is considered, on all the evidence, to be desirable.

Adrenalin, or an adrenal extract, has been tried in exophthalmic goitre, but, so far as the present writer's experience is concerned, the results have not been encouraging.

Blair Bell believes that osteomalacia should be treated by means of suprarenal extract—not necessarily because this extract is antagonistic to the ovarian hormone, but because he believes there is often suprarenal inadequacy in this disease.¹¹ He thinks that the secretions of the adrenals are of value during pregnancy in “assisting the absorption and retention of lime.” In this connection it is interesting to note that certain observers have experimented upon the action of adrenalin upon the growth of bone, and have found that animals from whose bones rings of tissue have been removed, if subjected to the ingestion of this extract, heal with bony union much more rapidly than the control animals who are not subjected to the action of this drug. It would seem, therefore, that adrenalin exerts a helpful influence upon the growth of osseous tissue.

Now as to the methods of administering the adrenal preparations, and the different varieties which may be utilized. The extract of the gland may be given either separately or in combination with other hormones. Thus, Houssay recommends the combination of adrenalin and hypophysin, and maintains that given together a smaller dose of the former extract is required, which, as it is the more toxic of the two, is a decided advantage. “The combination may be used success-

fully in fulminating intoxications associated with hypotension, in tachycardia, and in myocarditis toxica, either subcutaneously or by the mouth. The combination produces a greater and more persistent local ischæmia than adrenalin alone. The adrenalin neutralizes the powerful enterokinetic action of hypophysin, while the active constituent of the posterior lobe of the hypophysis counteracts the mydriatic effect of the adrenal extract."¹² Swann has found that moderate doses of a 1 in 1,000 solution of adrenalin administered hypodermically completely cured, in two doses, urticaria.

The extract of this gland may be given either as a liquid preparation or as a solid extract. Adrenalin or adrenalin chloride may be administered by the mouth, hypodermically, or intravenously (the latter in emergencies). The first of these methods is the routine for the majority of cases—*i.e.*, where the treatment is required in small doses, probably over a lengthy period. Again, where it is to be utilized, not because there is definite evidence that the adrenal substance is deficient, but either as a counteraction to other hormones which may be presumed to be in excess or simply for its therapeutic action, this preparation may be relied on. Orally it may be given in doses ranging from 2 to 3 minims up to $\frac{1}{2}$ drachm. The dose hypodermically should be about half—that is to say, up to 15 minims.

This extract has also been used in the treatment of asthma, with a considerable degree of success. It may

be given hypodermically or locally as a spray or douche to the nasal mucous membrane.

Preparations of the gland are also made in the form of dry extracts, but the dose appears to vary with different authors. Thus, it is sometimes recommended in doses ranging from $\frac{1}{4}$ to 3 grains (for mild hypoadrenia), and it is possible that this variance is to be explained by the fact that different authorities are referring to different preparations. Again, some manufacturers supply a tablet containing the extract of several of the ductless glands, such as thyroid, pituitary, and adrenals. If it is desired to give a pluriglandular preparation, such as this, it is important to know the exact dose of each extract which the tablet contains.

We may mention here a possible method of counteracting hyperadrenia. The patient with over-action of the adrenals is by no means uncommonly met with, but the accurate treatment of such a case is difficult. He is usually thin, nervous, with digestive instability and disturbed sleep. It is always worth while to treat such a patient with pancreatic extracts, as the antagonism between the adrenals and the pancreas is well proven. It may be assumed that such extracts will help to antagonize the excessive secretion from the suprarenal glands, and thus assist in the maintenance of a normal balance.

Finally as to the contra-indications of adrenal therapy. It will be quite obvious, from what has already been said, that hypertension is an absolute

contra-indication; also that it should never be administered in diabetes, as it is well known that the output of sugar is increased by the administration of adrenalin. Again, if it is given at all in tuberculosis, especially in phthisis, it must be stopped as soon as any pulmonary hæmorrhage manifests itself.

Provided that it is given under the careful observation of a medical man, who can observe the changes which accrue as the treatment progresses, more particularly with regard to the vascular tension and pulse, it is not only a safe remedy, but a particularly successful one. We shall refer in a subsequent chapter to the important points in pluriglandular therapy, and to the factors which decide our choice between the prescription of a single hormone or a preparation containing the extracts of several of the ductless glands.

REFERENCES.

- ¹ Elliott, *Practitioner*, vol. xciv., No. 1, p. 126.
- ² Edit. *Jour. Med. Amer. Assoc.*, 1913, lxi. 123.
- ³ Harrower, *Practical Hormone-Therapy*, p. 263.
- ⁴ Geikie Cobb, *Neurasthenia: Its Causes and Treatment*, *Practitioner*, August, 1915; see also article on the Diagnosis of Neurasthenia in *Practitioner*, April, 1913.
- ⁵ Sargent, quoted in *Practical Hormone-Therapy*, p. 266.
- ⁶ Sajous, *Hypo-adrenia as a Cause of Death in Infections, and its Treatment*, *Mo. Cyc. and Med. Bull.*, 1911, iv. 725.
- ⁷ Elliott, *loc. cit.*, pp. 127, 128.
- ⁸ Harrower, *loc. cit.*, p. 265.
- ⁹ Harrower, *loc. cit.*, p. 264.
- ¹⁰ Harrower, *loc. cit.*, quoted from Naame, *Jour. de Pract.*, 1912, xxiv. 87.
- ¹¹ Blair Bell, *The Internal Secretions and Female Characteristics*, *Proc. Roy. Soc. Med.*, 1913, vi. 74.
- ¹² Quoted in Harrower's book, pp. 275, 276.

CHAPTER VII

THE PANCREAS

THE next gland which we must study in connection with the endocrine glands is the pancreas. And from its importance in the bodily metabolism, an importance which is receiving fresh proofs every day, it deserves a high place in any series which deals with the internal secretions. Apart from the vital nature of its external secretion—that is to say, its digestive ferments—there is an ever-accumulating mass of evidence to show that the pancreas is concerned with the regulation of carbohydrate metabolism in an even more important relation than are the thyroid and pituitary glands.

For this reason alone, therefore, it behoves us to have a practical knowledge of the pancreas and its functions, and to familiarize ourselves with the results of recent physiological research. We shall therefore devote this chapter to a consideration of the structure, functions, and pathology of disorders, of this gland, and conclude by outlining the therapeutic possibilities.

Anatomy and Physiology of the Pancreas.

The pancreas is a tubulo-racemose gland, situated transversely across the posterior wall of the abdomen, at the back part of the epigastric and left hypochondriac regions. In structure it resembles the salivary glands, although it is of softer consistency and less compact than these organs. It is composed of a head, a neck, a body, and a tail; while its duct has received the name of "the canal of Wirsung." This duct opens into the duodenum, either in common with the ductus communis choledochus or else by an independent orifice.

The gland is composed of alveoli, loosely held together by connective tissue, containing in its meshes cells known as the "islets of Langerhans." These cells are of great importance, for, as we shall see later, they are concerned, as physiological research has shown, in the production of a hormone which governs carbohydrate metabolism. They are plentifully supplied with bloodvessels, arranged in the form of a compact network of capillary vessels.

For many years these cells have been known to exist, but until recently their function has been hidden. Now, however, it is becoming increasingly clear that they, in all probability, secrete a hormone which is absolutely necessary for the continuance of life. Previous to the recent work on internal secretion, the rôle played by the pancreas was supposed to be solely one of digestion; but Bayliss and Starling showed as long ago as 1902 that the digestive juices of the pancreas

were capable of being stimulated by a substance known as secretin, which is derived from pro-secretin by treatment with an acid. In normal digestion, pro-secretin, when it comes into contact with the acid of the gastric juices, is converted into secretin, which sets up pancreatic activity.

The external secretion of the pancreas is composed (according to Halliburton) of water, organic solids, and inorganic salts. The organic solids consist of the ferments trypsin, amyllopsin, steapsin, and a milk-curdling ferment, together with other less important constituents, such as traces of leucine, tyrosine, and a small amount of proteid matter.¹

These ferments are most important in carrying on the digestive processes inaugurated in the mouth and stomach; but as their chemical actions are physiological facts not directly connected with our study, we will not deal at any length with them here. What does concern us in this connection is the internal secretion of the pancreas, and the influence it exerts upon metabolism in general.

As we have already said, the islets of Langerhans are of great interest in studying the internal secretion of this gland, as recent work has confirmed the theory that it is this part of the pancreas which is so essential to life.

The Internal Secretion of the Pancreas.

In 1889, V. Mering and Minkowski made the important discovery that removal of the whole or of a large part of the pancreas was followed by hyperglycæmia, and eventually fatal diabetes. That this is not due to the abolition of the external secretion is proved by the discovery that, if pancreatic substance is grafted into an animal, glycosuria is prevented, although the secretion of ferments into the alimentary canal is absent. Again, if the pancreatic duct be tied, the pancreatic tissue atrophies, but it is believed that the islets of Langerhans remain in a sufficiently normal condition to furnish the body with the internal secretion. For although diabetic symptoms do not ensue after ligation of the duct (in spite of the atrophy of the pancreas), if the organ be subsequently removed glycosuria results.

Again, in support of the belief that the islets are responsible for an internal secretion, it has been noted that these cells are degenerated in some cases of diabetes, which is strong presumptive evidence that pancreatic diabetes owes its origin to an interference with an internal secretion provided normally by the cells in the islets of Langerhans. Schäfer says that these cells have a common origin with the cells of the duct, although they are not connected with the alveoli or ducts.²

Without going into elaborate details of pancreatic physiology, we must give the bare outline of the struc-

ture and behaviour of this gland, in order that its bearing on the hormone balance and its relation to carbohydrate metabolism may be clear.

To summarize the modern view as to the relationship which the pancreas bears to diabetes, we may say that it may fairly be assumed that the islets of Langerhans secrete a substance which, circulating in the bloodstream, prevents an undue amount of glucose from accumulating in this fluid. Hedon has shown, by passing the blood of normal animals through a de-pancreatized dog, that glycosuria may be prevented. This serves to show that the blood of normal animals contains some substance or substances which regulate carbohydrate metabolism.

The Relation of the Pancreas to Diabetes.

This evidence, when added to the facts about the islets to which we have already referred, is sufficiently convincing to warrant us in assuming that the pancreas possesses an internal secretion which, by its presence in the blood, regulates sugar storage, although its influence upon the glycogenetic properties of the liver has still to be determined. In this connection, Schäfer suggests that "the substance produced by the pancreas islets may be a chalonic autacoid which tends to inhibit the formation of glucose from glycogen, and incidently to promote the storage of glycogen, so that in its absence the glycogen which is present in the liver is rapidly converted into glucose, and the sugar

absorbed from the alimentary canal formed in the body is not stored by the liver. The result again will be hyperglycæmia and glycosuria."³ Schäfer proceeds to say that the "hypothesis that there exists a chalonic or inhibitory agent in the internal secretion of the pancreas which affects carbohydrate metabolism is probably the correct one."⁴

In any case, the evidence as to the part played by the islets of Langerhans in the production of diabetes is very complete; the discovery by Opie that a man who had died of diabetes possessed atrophy of these cells was a strong confirmation of the theory that these islets possessed some such function. This has been confirmed by numerous autopsies, so that one of the greatest authorities on this subject, Professor Van Noorden, now maintains that no diabetes can exist without disease of Langerhans' islands.⁵ This author in the same place divides diseases of the islands into two groups: those which result from diseases of their environment, and those resulting from specific disease of the islands independent of their surrounding tissues. The first of these groups would include such affections as inflammatory conditions of metastatic origin, or inflammation proceeding up Wirsung's duct, and pancreatic arteritis; while the second group includes "congenital constitutional atony of the island system" and "acquired atony of the same system." In association with this classification—that is to say, in support of the idea that some cases of diabetes owe their origin to disturbances from without the pancreas—Morris

believes that a bacterial infection from the duodenum may cause pressure upon the islets, and thus set up a pancreatitis. When the pancreatitis is relieved, the pressure disappears and likewise the glycosuria.⁶

The present writer has quoted these views for the reason that they offer a feasible theory as to the production of pancreatic diabetes in man, and give some idea as to the manner in which interference with Langerhans' islets may be brought about.

Apart from the interest of the recent discoveries which physiologists have made about the functions of the pancreas, these discoveries throw an added light upon the part which this gland displays in the general bodily metabolism, as well as its importance in supporting the hormonal balance. Again, our newly found knowledge explains many facts which were hitherto known as facts, but lacked an explanation. As an example of this assertion, we may mention one factor which characterizes glycosuria—namely, the increased liability to infection. It has been known for many years that the diabetic possesses little phagocytic power, and that he is not only a bad subject for operation, but also extremely prone to develop concurrent diseases, such as tubercle. Many observers have drawn attention to this fact, and have emphasized the lowered resistance to the tubercle bacillus which diabetics possess.

H. R. Harrower published an article in the *Practitioner*, in which he dealt very fully with the influence of the pancreas and its extracts upon infection; and he

maintained that this gland is concerned with defending the body against bacterial infection. This is reprinted with additions in his recent book,⁷ and the reader is referred to this chapter for an able and comprehensive account of this action of the pancreas.

In discussing the action of the pancreatic hormone, we must bear in mind that it, like all other internal secretions, has a relation to its fellow-hormones, and that some facts bearing upon this have been recently brought to light. Thus, it has been shown that the thyroid as well as the parathyroids influence pancreatic activity, but in opposite relations. The thyroid appears to be antagonistic to the pancreatic secretion, for its removal tends to prevent pancreatic diabetes; while the removal of the parathyroids assists hyperglycæmia. We have already seen that there is an increased tendency to glycosuria in Graves' disease, while there is always an increased sugar tolerance in the opposite condition, submyxœdema.

Another factor which must be mentioned when discussing sugar metabolism is the permeability of the renal filter—that is to say, the amount which the kidney is capable of passing through its meshes. The normal sugar content of the blood is about 0.1 per cent., and to this amount the renal apparatus is impermeable. But if the amount exceeds this figure the kidney is unable to resist the hyperglycæmia, and sugar passes into the urine. A marked degree of hyperglycæmia is always present as the result of extirpation of the pancreas, and this is the immediate cause of the glyco-

suria. The immediate production of sugar in the blood in excessive quantities following upon pancreas removal is doubtless concerned with several facts which bear upon the hyperglycæmia. The glycogen in the liver is one of the first substances to be altered by the operation, and this is very markedly reduced shortly after the removal of the gland. After generous feeding with carbohydrates, moreover, only traces of glycogen are able to be discovered. In contrast to this, the leucocytes exhibit a very high glycogen content. It is possible that this freeing of the glycogen reserves is one source of the increased amount of sugar in the blood, but it can hardly be the only one, for it would soon be exhausted. The additional sugar must come from albuminoids and fats, and this would in some measure account for the extreme wasting which these cases show.

So much for a few facts bearing upon the relation between the pancreas and the symptoms of diabetes. Now we will discuss a little more fully the internal secretion of this gland before turning to the therapeutic consideration of the pancreas.

Origin of the Pancreatic Internal Secretion.

It was at first supposed that, if an internal secretion existed at all in this gland, it was formed by the same cells which manufactured the ferments—in fact, that such cells performed a double action. In 1869 Langerhans first drew attention to those groups of cells

140 THE ORGANS OF INTERNAL SECRETION

which have since borne his name, and from their characteristics it is now assumed that they are responsible for the internal secretion. They are composed of badly defined polygonal cells, with large nuclei, and a very free blood-supply which comes from capillary bloodvessels in intimate relation to the cells. It is generally believed that these cells have no communication with the acini of the gland, but the question is not definitely settled. In some of the lower animals proof is forthcoming that there exists some connection between the excretory ducts and the cells of the islands.* Again, it is believed by some workers that the islets play a part in the production of the external pancreatic ferments, or, again, that the islets and the acini are not functionally differentiated.

The action of adrenalin upon pancreatic activity is of interest, for it is well known that adrenalin when injected into the blood-stream sets up a temporary glycosuria; but it is not equally well known that some observers have found that the administration of adrenalin to animals has been followed by the appearance of necrosed areas in the islets of Langerhans. Although the action of adrenalin in producing glycosuria is apparently mainly through the pancreas, that this is not solely so is evidenced by the fact that there is an increase of sugar even in depancreatized animals after the administration of adrenalin. Again, hyper-

* Some observers have been able to inject the islands through the ducts, while others have observed connections between lumina in the islands and the acini.

glycæmia can be produced by means other than disorder of the pancreatic functions and the administration of adrenalin, such as puncture of the fourth ventricle and stimulation of the splanchnics.

A fact which is interesting from the promise of therapeutic possibilities in the future is that elicited by Zuelzer—namely, that adrenalin glycosuria can be overcome by the administration of pancreatic extracts and juices; for, whether due to an antagonistic action on the adrenal hormone or to its effects on the permeability of the kidney, it shows promise of further therapeutic developments.

It has been stated that adrenalin instilled into the eye of a depancreatized animal causes dilatation of the pupil, which does not occur in the normal animal, and this test has been made use of for the diagnosis of pancreatic diabetes, as it has been shown that it is likewise present in man in some cases of diabetes. In any case, it seems that it will not be long before further light will be thrown both upon the relation of the pancreas to the production of diabetes, and also upon the site of production and the more exact nature of the internal secretion.

Therapeutic Considerations.

We now come to a brief consideration of the therapeutic possibilities of extracts of pancreas. It has been necessary in this chapter to deal somewhat fully with diabetes, as the study of this condition is directly

142 THE ORGANS OF INTERNAL SECRETION

related to any consideration of the internal secretion of the pancreas; and our knowledge at the present time has been mainly gained by experiments in the production of hyperglycæmia and glycosuria and their relation to the pancreas.

In dealing with the therapeutic side of the pancreas, we must realize that a good deal of difference of opinion exists as to the efficacy or otherwise of this particular form of hormone-therapy. First, with regard to the administration of pancreatic extracts in cases of diabetes, Schäfer, writing doubtless from the experimental standpoint, says: "It is not found that pancreas extracts have the effect of antagonizing the results of pancreas extirpation; in this respect they offer a parallel to the negative results of suprarenal extracts in antagonizing the effects of adrenal deprivation."⁸ It must be remembered that it is not maintained that all cases of diabetes which are encountered clinically are of necessity due to pancreatic disorders, in spite of the opinion of Van Noorden, which we have already quoted. It is therefore only reasonable to assume that some cases of diabetes which we treat with pancreatic extracts may be due to causes other than pancreatic. If this is so, we must not expect to improve every patient suffering from glycosuria who comes under our care by the administration of pancreatic preparations.

The reasons why we may expect that this treatment will not be universally successful are summarized by Harrower as follows:

"First, not in every case of diabetes is the pan-

creatic factor present or prominent. Second, very few preparations of pancreas are made with the object of specifically supplying as much as possible of the islet substance. Third, it is no easy matter to supply the lacking hormone, for it is normally supplied continuously to the blood, and we have no means of knowing how much is elaborated, or, in disease, how much is lacking, in each case. Fourth, the antagonism of the chromaffin hormone must be considered; for the diabetes may be due not simply to a deficiency of the pancreatic internal secretion, but to an exaggeration of that of the adrenals or a combination of both. Fifth, evidently several other factors are concerned, and hence the hormone treatment of diabetes *alone* cannot be instituted with hopes of success."⁹

Having given the reasons why this method of treating diabetes is at present not entirely successful, we may perhaps be allowed to quote a few opinions which are favourable to pancreas-therapy. Thus, the same author a little farther on in the same chapter says: "It is evident that the pancreas ferment preparations may have an excellent influence upon glycosuria."¹⁰ Harrower explains this thus: "This is probably due to the restorative action on the pancreas caused by the reinforcement of digestion, and the removal, in part or altogether, of the pancreatic disability, which is not alone reflected upon its external, but also on its internal secretion."

Again, Harrower refers to trypsogen (a combination of the ferments trypsin and amylopsin with the internal

144 THE ORGANS OF INTERNAL SECRETION

secretion) as having benefited several cases of undoubted diabetes mellitus; occasionally the sugar has quite disappeared, more often the symptoms characteristic of this complaint have been much benefited, weight has been increased, polyuria quite controlled, and sugar reduced."¹¹

There are also on the market other preparations of pancreas, some of which are composed of one or more of the pancreatic ferments, while others are extracts of the whole gland. Before proceeding farther with the discussion of this form of therapy, we must emphasize that it is absolutely essential it should be regarded, in so far as diabetes is concerned, as an adjunct rather than as a sole form of treatment. For it is quite agreed that where there is excessive glycaemia there is every reason to relieve materially the strain upon the pancreas by limiting the ingestion of carbohydrates. Therefore, so far as this disease is concerned, diet should always be the primary consideration, and an endeavour to replace a lost or diminished internal secretion a justifiable secondary one.

Another preparation which contains the extracts of liver and pancreas is Kinazyme, and it is recommended by its manufacturers to increase weight; it is also said to be a valuable adjunct to the treatment of tuberculosis. It may be presumed that this not only is believed to encourage normal digestive processes, but also to raise the bodily resistance to infection.

It has also been stated that extracts of pancreas are useful in hypertension, presumably for their counter-

acting power on the chromaffin system. In cases where there is some evidence of hyperadrenia, such treatment should be given a trial.

Finally, while insisting with as much stress as formerly upon dietetic and hygienic principles in the treatment of diabetic and kindred disorders, the practitioner should bear in mind preparations of this gland; for it is quite possible that such preparations will turn the scale in favour of the patient, while if left solely to the conservative method of withholding carbohydrate food it might turn in favour of the disease. Such therapy as is offered by extracts of the pancreas is rational and harmless, and it must be confessed that there is an opening for any form of treatment which will make the lot of the diabetic an easier one. For this reason alone, even if no other is forthcoming, it is advisable to try this form of opotherapy in those diseases where it may be assumed that the pancreas is at fault; not only in definite diabetes, but in deficient digestion (the so-called carbohydrate dyspepsia), either alone or combined with preparations of the intestinal secretions, to which the pancreatic ferments have so close a relation.

These diseases, whose name is legion, offer a legitimate scope for the rational practice of organo-therapy, and so far as assimilation is concerned the pancreas may be regarded as the foundation, which, if upset, even by temporary or slight disturbances, is capable of upsetting the balance without which normal metabolism is impossible.

REFERENCES.

- ¹ Halliburton, *Handbook of Physiology*, p. 491.
- ² Schäfer, *The Endocrinic Glands*, p. 125.
- ³ *Ibid.*, p. 129.
- ⁴ *Ibid.*, p. 130.
- ⁵ Van Noorden, *Practitioner*, vol. xciv., No. 2, p. 237.
- ⁶ Morris, *Jour. Am. Med. Assoc.*, lx., 1913.
- ⁷ H. R. Harrower, *Practical Hormone-Therapy*, ch. viii.
- ⁸ Schäfer, *loc. cit.*, p. 128.
- ⁹ Harrower, *loc. cit.*, p. 104.
- ¹⁰ Harrower, *loc. cit.*, p. 106.
- ¹¹ Harrower, *loc. cit.*, p. 107.

CHAPTER VIII

THE SEXUAL ORGANS AND THEIR INTERNAL SECRETIONS

It will only be possible to review briefly the main features of the sexual organs and their hormones. For the subject is so vast, and the difference of opinion which exists about many of these organs and their secretions is so great, that it would necessitate a lengthy study, which is not possible with the space at our disposal. We shall therefore enumerate the main facts with regard to the hormones elaborated by these glands, and deal briefly with the relations between these secretions and those of other of the endocrine glands.

For this reason, if some of the views may appear dogmatic, it must be remembered that we are unable, owing to limitation of space, to give the results obtained by different observers, but must content ourselves with an expression of the views which have obtained more or less general assent.

We have mentioned in previous chapters that an intimate relation exists between many of the sexual secretions and the hormones elaborated by some of the other ductless glands. As an example, the adrenals

148 THE ORGANS OF INTERNAL SECRETION

have an intimate relation with the ovaries and testes, and more than a few cases are recorded which bear this out. Elliott mentions the case of a young woman, perfectly normal sexually, who at the age of twenty lost her menses; this was followed by the appearance of secondary sexual characteristics of the opposite sex, such as the growth of a beard and the development of a masculine type. Four years later she was operated upon and a tumour of the right adrenal gland removed.¹

Again, a case described by Alberti shows a similar relationship, only the perversion is differently connected. The patient was a young girl who was perfectly normal sexually, but who gradually reverted to the masculine type. There was a loss of adipose tissue, the hair increased, more especially upon the body, the breasts became soft, and the voice deepened. A few years later she was operated upon for ovarian tumour, and eventually died of peritonitis. These two examples are selected, for they refer to relations between two different systems, and show that an intimate relationship does exist between the sexual glands and other of the internal secretory organs.

Internal Secretion of the Testis.

Recent experiments have shown that the internal secretion of the testis has a vast influence in animals belonging to the species which has a periodical rutting season, for changes in various parts of the body as well as in the reproductive organs have been noticed at

such times. If there is an internal secretion provided by the testis and the ovary, where is this formed? It is now believed that the "interstitial cells of Leydig" in the testicle, and the corresponding cells in the ovarian stroma, are the sites where this secretion is manufactured.

The changes which appear at puberty owe their origin to an internal secretion which "sensitizes" the nervous reflexes concerned with the performance of the sexual act. That the various nervous reflexes are developed to a high degree of sensitiveness has been shown by experiments on frogs during the spawning season, which proved conclusively that these animals develop additional nervous mechanisms during the period of sexual activity.

Again, it has been abundantly established that a large number of internal secretions are concerned in the development of the sexual glands; for sexual precocity can be associated with tumours of the adrenals and with disturbances in the hormonal balance; while the influences upon sexual development which are exerted by the hypophysis have already been discussed in a previous chapter.² It would seem, however, that the adrenal gland is especially concerned with the regulation of the sexual secretions, for, in some post-mortem examinations of cases of sexual precocity, out of seventeen such cases, and four surgical cases, suprarenal tumours were found in eleven, nine being girls and two boys.³ There are many more such cases which have been published which certainly

justify us in assuming that the relation between the genital glands and the chromaffin system is a close one, and morphologically this is to be expected.⁴

The Uterus and Mammary Gland.

Now for a few words about the relation between the genital gland of the female and the mammary gland. There can be little doubt that the development of this latter organ depends upon a secretion from the genital apparatus. It was at one time thought that the connection between the uterus and the breast was a nervous one; but this has long since ceased to be believed, as experiments have shown that destruction of the nervous connection has no influence in inhibiting the flow of milk in a pregnant animal. The connection is therefore by means of a chemical messenger; and it is now believed that this messenger emanates from the ovaries, and not from the uterus, as was formerly supposed. Experiments have shown that the mammary gland does not develop after removal of the ovaries in young animals, but that transplantation of ovarian substance produces normal development.

Much work has been done recently upon the relation between the mammary gland and the pregnant uterus. The question to be solved is, From what source comes the hormone which activates lactation? Is it from the maternal or the foetal organs? Does it come, as maintained by Halban, from the placenta, or does the foetus provide it, or is it a product of the uterus or

ovary? We have not space to enter into the details of the physiological work which has been done on this subject; we must therefore content ourselves with outlining the views of authorities.

Basch, from the results of experiments in which he injected placental extracts, came to the conclusion that mammary activity during pregnancy depended upon a substance found in the placenta. Starling and Lane-Claypon experimented with injections of foetal extract, extract of uterine mucous membrane, and ovarian extract. They produced mammary hyperplasia by injecting foetal extract, but the other extracts were inert in this respect. This has been confirmed by experiments carried out by Foa, and also by Biedl and Königstein.

Without entering into any further details of these experiments, we may quote in full the conclusion at which Biedl arrives on this subject, for it gives a good hypothesis upon which to base our views. "The secretion of the hyperplastic mammæ is not difficult of explanation. . . . The growth of the gland is the expression of an assimilatory increase of material with simultaneous inhibition of disassimilatory decomposition. The growing gland produces no secretion, or very little (colostrum). With the suppression of the assimilatory hormone supplied by the foetus—that is to say, at birth or by termination of the pregnancy during the second half—disassimilatory decomposition, as expressed by the secretion, is enabled to proceed unchecked."⁵

We need not spend much time in studying the effects of castration upon development, for these are well known. There are, however, one or two facts which are important.

Effects of Castration.

The removal of the ovaries in young animals produces certain features of arrested development, among these being a rudimentary condition of the Fallopian tubes and uterus. It has been shown, however, that the transplantation of ovarian tissue under the skin remedies all these defects and the animal develops normally. This shows definitely that the controlling influence is chemical (hormonic), and not reflex. Again, the effects of castration upon young male animals, and upon youths, as is practised in the East, shows that there is a definite controlling agent in the secretion of the genital glands which, directly or indirectly, governs growth. Thus, changes take place in the bones, particularly in the skull and the pelvis, which produce an increase of growth, with retarded ossification of the epiphyses.

Pregnancy appears to be accompanied by reduced ovarian secretion, and it has been noted that young females grow during pregnancy, which, accompanied as it is by diminished ovarian secretion, would appear to show that the restraining influence of this secretion is temporarily removed. Or, to put it in another way, the sexual glands elaborate a hormone which is concerned in the ossification of epiphyses; if this be with-

drawn before ossification is complete, increased skeletal growth ensues.

Again, castration affects the hormonal balance, as is shown by the changes which are produced in the thyroid, thymus, pituitary, etc., by removal of the testes: and Fichera showed, in 1906, that the enlargement of the pituitary which followed castration in animals could be reduced by the injection of extract of the testicle. Again, it has been shown that the thyroid under such conditions is atrophied, the thymus persistent, and the hypophysis enlarged. Only a few experiments have been performed on this aspect of the relationship of the genital glands with other of the internal secretion organs; but in one instance of a eunuch dissected by Tandler and Gross these changes in the thyroid, thymus, and pituitary, were noted.

Again, the general metabolism is profoundly affected by changes in the reproductive organs; and the obesity, and in some cases mental sluggishness or other psychic changes which occur in senility, are too well known to require more than mention. Löwy and Richter investigated the metabolism in castrated animals, and showed that there was a reduction in proportion to body-weight, as well as a reduction in the total metabolism of the body.

The relation between ovarian secretion and osteomalacia has been the subject of much speculation. Biedl thinks that, while it cannot be maintained that we have definite proof of the causative rôle played by the ovarian secretion in the production of this disease

the probability is that the ovary, rather than any of the other endocrine glands, such as the thyroid, thymus, etc., is the main gland concerned in the production of the disease; although there may be several glands concerned, the ovary is certainly the chief.

The pioneer work which Brown-Séquard undertook in administering to himself extracts of testicles, and the report which he made as to the results, may be said to have founded the theory and practice of organotherapy, however empirical this treatment was and, to a certain extent, still is. His results are very well known, and need not be stated here at any length; suffice it to say that he found his strength much increased by this administration, he was less liable to fatigue, his excretory apparatus was improved, and the effects were in general satisfactory. Testicular extract is said to contain nucleoprotein and toxic bases, and it has been stated by some observers that the results of administering such an extract to animals are widespread and important; also it has recently been said to play a part in the production of the genito-vesicular reflex (Serralach and Pares, quoted by Biedl). On the other hand, it has been stated that *ovarian* extract, when injected, lowers blood-pressure, causes in the thyroid gland a preliminary reduction followed by an enormous increase in volume, and reduces the volume of the kidney.

The Internal Secretions of the Sexual Organs.

As we have already said, the cells of Leydig are supposed to be the seat of formation of the internal secretion of the testis; and an interesting fact in connection with this is that puberty produces an enormous development of these cells, while old age sees a marked degree of atrophy of the same cells. It is stated that, in those cases where cryptorchidism in man is associated with the development of secondary feminine characteristics, there is complete sclerosis of the testicle, and that the cells of Leydig are reduced to a few scanty groups (Biedl).

Without going farther into the physiology of the testicle, it is sufficient to say that there can be very little doubt that these organs play a vital part in the production of secretions which, although we are ignorant of their full importance, are necessary for the development of a normal man: not only in so far as sexual characteristics are concerned, but even with regard to the adequate working of his nervous system.

The secretions of the prostate are likewise not fully known, certainly so far as its internal secretion is concerned. Its external secretion exerts an influence upon the spermatozoa, and dilutes the testicular fluid; but little is known of the properties of its hormone, except that injections of prostatic extract made from animals are extremely toxic. Therapeutically, extracts of prostate gland are not prescribed much in this country,

although they are supposed to be beneficial in post-operative melancholic conditions following removal of this gland.

Therapeutics of the Sexual Glands.

We must now turn our attention to the therapeutic possibilities of extracts of the genital glands, and we shall endeavour to give a résumé of the possibilities of such therapy.

A very wide range of treatment is open for the administration of preparations made from the gonads, and it is now established that, in conjunction with other organo-therapeutic preparations, should these be indicated, a great deal of benefit may accrue to patients suffering from a variety of diseases which it would be difficult to ameliorate by other means.

Let us commence our study by considering the help which may be given to that very widely encountered disease, neurasthenia. It has been conclusively proved that the exhibition of some preparation of the testes (such as Spermin) has the power of raising the amount of muscular work available, of increasing the strength of the individual, and of banishing ready fatigue. In other words, the conclusions arrived at by Brown-Séquard have been substantiated, and it is now believed that the administration of testicular preparations is followed by improvement in these cases. Some manufacturers combine this substance with other secretions in the form which Leonard Williams aptly calls a "mitrailleuse."

Unfortunately, there appears to exist some degree of opposition to this remedy, and in consequence it has never been widely used. There is, however, a good deal to be said for its powers, and Harrower summarizes its utility in the following words:

“ A course of treatment with testicular extract frequently accomplishes such a number of changes of such remote organs, and influences such widely disconnected ones, that to report actual results might even be sufficient to discount the position of this method. The extent of these results is undoubtedly due to the stimulating action of the testicular hormone upon the endocrinous system as a whole, and as a means of revitalizing an individual with deficient oxidation, poor nutrition, nervous inactivity, and general weakness, the system is of undoubted value. There are few single remedies which can compare with active and properly prepared testicular extracts.”⁶

The therapeutic possibilities of the ovaries are numerous. Thus, it is well known that an intimate relation exists between the pituitary gland and the ovaries, and consequently, in undertaking the treatment of disorders of the pituitary, this fact should be remembered. The administration of ovarian extract is frequently indicated in such cases, more specially where the symptoms point to excessive bony growth (it has already been said that the ovary is presumed to exert an influence on the ossification of the epiphyses). In this connection Blair Bell, speaking of imperfect development, says:

"Nearly all the patients are adipose, whether the deficiency arise in the thyroid, pituitary, or ovaries. If there be pituitary insufficiency, the skin will be found to be soft and smooth; if there be thyroid insufficiency, it is harsh and rough. With pituitary insufficiency the blood-pressure tends to be below 110 mm. Hg; but with thyroid insufficiency it is usually normal (125 mm. Hg), or slightly above normal."

Further, if it be true that both thyroid and pituitary insufficiency induce ovarian insufficiency, then we can, as Blair Bell points out, treat the ovarian deficiency by means of either thyroid or pituitary extract. This simplifies treatment in so far as this class of disorder is concerned; but, nevertheless, for other instances of ovarian deficiency, such as are exemplified in derangements of the catamenia, ovarian extract will be found to be very useful. Again, Blair Bell himself drew attention to the influence of the ovaries upon calcium metabolism, and showed that the calcium in the urine was diminished after oöphorectomy by one-half;⁸ which again bears out what we know with reference to the beneficial effects which follow (in cases of osteomalacia) when double oöphorectomy is performed.

Harrower says that better results are obtained by the exhibition of extracts of the corpus luteum than are recorded by the ingestion of extracts of the whole gland; and other observers are of the opinion that the superiority of this extract over the total extract of the gland is incontrovertible. Extract of corpus luteum will be found useful in neurasthenia in women, in

difficulties arising at the menopause, after removal of the ovaries for cystic or other disease, and in conjunction with thyroid or pituitary for incomplete development, etc.

Several writers, notably Leduc and Blair Bell, lay great stress upon the calcium metabolism of the body, and its relation to disorders of the nervous system (neurasthenia, hystero-epilepsy). Thus, we know that there is a storage of calcium salts in hypothyroidism, and an increase in the calcium metabolism in Graves' disease; also that at times like the menopause a disturbance of more than one gland may take place. The signs in any individual case may point to which of the endocrinous glands is chiefly at fault; but the best results are probably obtained, as Blair Bell points out, by the administration of ovarian extracts, in company with whichever of the other glands appears to be at fault. Again, this writer suggests that the mental depression from which some women suffer at the climacteric can be much relieved by the administration of thyroid extract. This in itself would point to the thyroid as being one of the glands deficient in secretion at the menopause.

One more fact in connection with the ovaries, and we can pass on to the consideration of other secretions. It is assumed that menorrhagia may be due to excessive action of the ovaries—*i.e.*, hyperoöphorism, with which may be associated hyperthyroidism. If this may be presumed to be so in any given case, it is obviously wrong treatment, and may even do harm,

160 THE ORGANS OF INTERNAL SECRETION

to administer ovarian extracts; for the same reason extract of thyroid gland must be avoided. On the other hand, where amenorrhœa is present, especially where the appearance of the menses at puberty is delayed, we may with advantage prescribe thyroid extract with or without ovarian extract, remembering the stimulating effect which this extract exerts upon this function. Therefore, where obesity is associated with amenorrhœa the possibility of hypo-oöphorism should be borne in mind.

We must refer in passing to the mammary gland, and state briefly the modern conception as to its utility as a therapeutic agent. We have already pointed out that Starling and Lane-Clayton experimented with foetal extracts, and came to the conclusion that such extracts were capable of producing hyperplasia of the mammæ. Therefore it would seem from this that the foetus is the seat of formation of a hormone which stimulates the secretion of the breasts. That there is a close relation between the uterus and the mammary gland has been known for a long time; what that relation is exactly we are better able to say now than formerly. The obvious connection between the functioning of the breast and the uterus is exemplified by such occurrences as the enlargement of the mammæ at puberty and pregnancy, the intimacy of its nerve connections, and the pain in the breasts during menstruation. The following facts also bear upon this study: The placental hormone, to which we have already referred, when injected into another animal,

produces a flow of milk; and this will likewise take place in transplanted portions of the gland.

The relation of the ovary to the mammary gland has something in its bearing of an antagonistic nature. Thus, although complete ovariectomy may result in disappearance of the breasts, which latter glands will reappear upon grafting ovarian tissue, some observers believe that ovarian secretion is antagonistic to mammary. In this connection it is interesting to note that some investigators have found that the rupture of the ripe follicle in the ovary results in the growth of the mammary glands.

From the therapeutic standpoint, moreover, the balance of evidence is in favour of administering mammary extract to promote lactation. It is also stated to be useful in combating dysmenorrhœa and menorrhagia; it is not improbable that it is able to render aid to the uterine functions, either on account of its supposed correlation with the thyroid gland, or on account of the known sympathy which exists between the mammæ and the uterus.

We have now outlined the facts with reference to the reproductive glands and their internal secretions, and we have indicated the lines along which treatment by opotherapy of these disorders may be conducted.

162 THE ORGANS OF INTERNAL SECRETION

REFERENCES.

- ¹ Elliott, *Practitioner*, vol. xciv., No. 1, p. 132.
- ² Chapter V., pp. 101, 102.
- ³ Biedl, *The Internal Secretary Organs*, p. 371.
- ⁴ Chapter VI., p. 117.
- ⁵ Biedl, *loc. cit.*, p. 376.
- ⁶ Harrower, *Practical Hormone-Therapy*, p. 341.
- ⁷ Blair Bell, *Practitioner*, vol. xciv., No. 2, p. 270.
- ⁸ Blair Bell, *The Genital Functions of the Ductless Glands in the Female*, *Lancet*, 1913, i, 809.

CHAPTER IX

THE INTERNAL SECRETIONS OF DIGESTION

No book dealing with the ductless glands and their internal secretions would be complete without due reference to the part played by chemical messengers in the processes of digestion. For while our knowledge in this field is very incomplete, we know now that many of the factors which contribute to normal digestion are chemical, and not, as was previously thought, nervous in origin. It was previously assumed that the functions of digestion were regulated by either the sympathetic or central nervous system, and the influence of hormones was unrecognized. But recent work on digestion has shown that even in this field the work performed by autacoids is by no means unimportant.

For this reason, therefore, we propose to discuss briefly the internal secretions concerned in digestion, and to outline the main features of their work. We shall also, wherever possible, draw attention to any relationships which may be presumed to exist between the organs of digestion and other glandular structures of the body.

The salivary glands have been supposed to secrete

a hormone which plays a part in digestion either by stimulating the later stages of digestion, or by carrying on salivary digestion in the stomach. There is now no doubt that this stage of digestion continues in the stomach, certainly in that part of the bolus which remains alkaline—that is to say, until it is soaked through with the gastric juices; and therefore the main work performed by the ferment may be said to be carried on in the stomach.

Again, certain authorities lay stress upon the parotid juice, and the possibility that it may be influenced by chemical messengers; while some clinicians have worked with preparations of the salivary glands, and have reported that they have influenced some of the abnormal conditions arising in other and remote organs. Thus, they have been used for ovarian disorders (the close connection existing between the parotid gland and the gonads is shown by the metastatic inflammations arising in the course of parotitis in the testicles and ovaries), and have been said to do good. Apart from this, which we have mentioned in passing, there is no evidence, not even presumptive, that the salivary glands are useful from the therapeutic standpoint.

When we turn to the question of gastric and duodenal digestion, we find that we are at once in quite a different position. At the risk of wearying our readers, we will briefly recapitulate the work which led to the important discovery that pancreatic digestion is dependent upon chemical stimulation.

INTERNAL SECRETIONS OF DIGESTION 165

Following upon the researches of certain observers, it was discovered that the introduction of dilute hydrochloric acid into the alimentary canal was followed by a marked increase in pancreatic secretion; that this was due to the acid coming into contact with the walls of the duodenum; and that this produced a hormone which possessed the property of stimulating pancreatic secretion. It was the discovery of this last factor, a discovery made by Bayliss and Starling in 1902, which may be said to have laid the foundation of our present knowledge of the importance of the chemical messenger in keeping that balance in organic metabolism which alone can produce perfect health.

This substance has been named "*secretin*," and is released from its precursor *pro-secretin* by the entrance into the alkaline duodenum of the acid chyme. This substance, which has been called the principal alimentary hormone, is present in all animals, and is not only a human property; in fact, it has been stated that the human intestine is relatively and actually lower in its secretin-content than is that of the sheep.¹ An analogous secretion is furnished by the pyloric walls, and has been called gastric secretin or *gastrin*; and it acts upon the oxyntic and peptic glands of the stomach. Edkins was the first to demonstrate the presence of this substance in the stomach, and to show that its object is to stimulate into activity the secreting cells of the stomach. This hormone has a precursor, known as *pro-gastrin*, from which the hormone is formed, either by the products of digestion or by other stimulus.

166 THE ORGANS OF INTERNAL SECRETION

In this way, then, do we find that a chemical messenger exists to activate from latent chemicals an active substance; and that in both cases—the gastric and the duodenal—the key which unlocks this chemical is a normal and usual concomitant of digestion. In the case of the duodenal “secretin,” the acid unlocks the secretion either by neutralizing a substance which inhibits the flow of secretin, or else by directly producing the substance “secretin” from its precursor “pro-secretin.” Whichever is the case is not a matter which need detain us here, and we will pass on to discuss other properties of this important hormone.

It appears that secretin acts not only in the upper intestinal tract in producing pancreatic secretion, but also on the lower intestinal walls—*i.e.*, the ileum—and assists in the production of *succus entericus*. It likewise stimulates bile production, and it is said that the typical stools of jaundice have been rendered normal in colour by a few doses of secretin. In this way, therefore, this substance may be regarded as an excitant of the motor functions of the bowel; for it has a laxative power partly dependent upon its action upon the intestinal juices and partly owing to its action in increasing the normal laxative, the bile. But it is necessary that a certain element should be present in the blood, as Hustin has shown that secretin alone is incapable of stimulating the pancreatic juice in the absence of blood. “The pro-secretin of the duodenal wall is first changed into

INTERNAL SECRETIONS OF DIGESTION 167

secretin by contact with hydrochloric acid; it is then carried in the blood-stream to the pancreas, where, as a result of chemical change, its characteristic features are lost, secretin combining with amylopsinogen and steapsinogen, to form the ferments amylopsin and steapsin, and with the precursor protrypsinogen, to form trypsinogen. This last becomes the ferment trypsin only after it has come into contact with the entero-kinase in the intestinal canal."*2

It will be seen from what we have already said that the acid content of the stomach is of great importance in the production of the secretion of the pancreas, and to a lesser degree of the other intestinal and hepatic secretions. It will also be seen that it is not unlikely that much of the good which has long been recognized to accrue from the prescription of acid mixtures in dyspeptic and constipated subjects may in reality, as pointed out by Harrower, be due to the release by such acids of the internal secretions of the bowel and stomach. The hydrochloric acid of the gastric mucosa has several important functions to perform, among these being its work as an antiseptic, its work in opening the pyloric sphincter, its power to activate pepsin, and lastly its action as a stimulant to the pancreatic secretion. In its absence pancreatic digestion cannot be carried on, for secretin cannot be formed from pro-secretin.

Although the facts which we have related with regard

* Although this is believed to be the sequence of events, there is still some divergence of opinion on the subject.

168 THE ORGANS OF INTERNAL SECRETION

to secretin are well known and generally admitted as proven, little is known as to the actual constitution of this substance. It is not a ferment, but exactly what it is we do not know. It is, like the other hormones, a substance of low molecular weight, and, as we have already seen, it stimulates the formation of all the ferments of the pancreas. Its work in the complex scheme of digestion is balanced to a nicety. Thus, it is not secreted until there is work for it to do—in other words, until the food has entered the duodenum in its acid state; thus, there is no chance that it will irritate the walls of the intestine by producing pancreatic ferment-action in the absence of food. Even so, trypsin is not formed until it meets entero-kinase, which again is not formed until food is present.

There is no positive evidence that secretin stimulates the internal as well as the external secretion of the pancreas, for it has yet to be proved that there is an absence or diminution of secretin in diabetes. In fact, it has been shown in one case of diabetic coma that there was abundant pro-secretin in the body.³

Turning now to another intestinal secretion, the succus entericus, there is still some question as to its formation. Pawlow thinks that its formation depends upon two factors, the distension of the bowel and the presence of pancreatic juice. Succus entericus contains several ferments: *entero-kinase*, which acts upon the trypsinogen of the pancreatic juice; *erepsin*, which completes the digestion of proteids; and several sugar-splitting ferments. There is a good deal of

INTERNAL SECRETIONS OF DIGESTION 169

dissension as to the exact nature of its formation, some observers believing that it acts as its own hormone, others that it is stimulated by the pancreatic juice, while others believe that additional factors other than those mentioned are required to bring about the secretion of this juice. As regards the proteolytic action of the intestinal juices, these latter require the presence of pancreatic juice to exert this action.

Turning now to other secretions of the alimentary canal, we must refer in passing to the supposed secretion of the appendix, also to that which it has been suggested emanates from the lymphoid tissue of the intestine, such as Peyer's patches. Some observers believe that the appendix secretes a hormone which stimulates peristalsis, and in support of this they quote the constipation which in some cases follows the removal of the appendix as proof that a normal intestinal excitant has been removed or inhibited. There is no definite proof of this, although the future may show that in ruthless appendicectomies we are removing a necessary stimulant, in spite of the fact that so many regard this organ as a useless relic.

The liver, whose chief function is the production of bile, nevertheless performs many other and important functions which contribute to the normal working of the body. For many years observers have been struck by the enormous size of this gland, which, they argued, could hardly be justified by its bile-producing properties alone. This was before its storage and filtering functions were realized; and, even after

170 THE ORGANS OF INTERNAL SECRETION

the facts collected by the experiments in which Eck's fistula was utilized, much remained to be explained as to the rôle which the liver played in bodily metabolism. We have now reached the stage where we have to consider whether the liver possesses an internal secretion, and whether extracts of this organ can be utilized therapeutically.

Let us first summarize the functions of the liver, and with the facts thus gleaned at our disposal we shall be better able to form an opinion as to the secretions produced by this organ. The hepatic cells are responsible for the production of the bile, with the bile-salts, pigments, and ferments; they are also concerned in the regulation of the sugar content of the blood, and in the regulation of nitrogenous metabolism; they are concerned, moreover, in regulating the power of coagulation of the blood, and they appear to have a definite rôle in the destruction of toxins, and in the prevention of such material from reaching the cells of the body. In this latter relation they act as filters between the gastro-intestinal tract and the general blood-stream.

We see, therefore, that modern physiology teaches us that the liver has other functions to explain its vast size beyond the mere production of bile. What evidence, then, have we that it possesses an internal secretion? It would seem that the secreting powers of the liver are analogous to those of the pancreas, inasmuch as they depend upon stimulation by chemical means for the adequate working of their secretions.

INTERNAL SECRETIONS OF DIGESTION 171

This has been shown by experiments which have proved that all the nerves to the liver can be cut without interfering with the biliary secretions, and that excitation of the cut nerves does not influence the secretions either by accelerating or retarding such secretions. We saw when discussing the pancreatic secretion that it was stimulated by the production of secretin from pro-secretin in the presence of the acid chyme—i.e., that it was dependent upon a humoral messenger—and in like manner may we assume that the secretion of the liver is affected. Indeed, we know that the production of bile is increased by secretin, and that other of the digestive juices are stimulated by this same hormone. If this secretion of the liver is stimulated by a hormone, it is more than probable that others emanating from the same gland are also so activated.

Again, the extreme value of cod-liver oil has been known for many years, but only recently has it been recognized that it must owe its power to something more than its fat-content. In fact, recent work has shown that some of its value can be removed by processes of refining. Leonard Williams, speaking of the properties of cod-liver oil, lays stress upon this aspect of its value, and says: ". . . it is very probable that this biliary secretion of the cod, when introduced into the human economy, acts as a stimulant to one of the normal internal secretory glands, and that the secretion of the one so stimulated is inimical to the development of the tubercle bacillus. . . ." This

172 THE ORGANS OF INTERNAL SECRETION

shows us another reason for its great value in tuberculosis, and explains its power of building up the body, while preventing toxins from reaching remote organs.

Other observers have experimented with cod-liver oil, and have sought to show that it is not the animal fat which it contains, so much as other less recognizable constituents, to which it owes its value. The internal secretory theory of the liver extracts certainly opens the way to many new applications of hormone-therapy, and also explains many hitherto unexplained facts about the value of such substances as cod-liver oil.

Here we must leave this subject and pass on to a brief study of the value of extracts of liver. For some years such extracts have been used on the Continent, and many observers speak highly of their beneficial influence in countless morbid conditions. Thus, Gilbert and Carnot, of Paris, have studied this subject intimately, and have arrived at certain conclusions about this organ and its secretions. Others, following on these lines, have arrived at similar conclusions, and we will here refer briefly to one of the theories about the liver which has been recently expressed. Some of the workers in this subject believe that the liver possesses an internal secretion which filters the blood coming from the digestive tract, and neutralizes, or renders harmless, toxins emanating therefrom. This is supposed to be due to some chemical present in the liver cells itself, and not only in the bloodstream. In this way the liver may be regarded as a vast filter standing between the chemical exchanges

INTERNAL SECRETIONS OF DIGESTION 173

occurring in the gastro-intestinal tract—which may possibly be harmful if admitted into the systemic circulation without due filtering—and the body. It is only to be expected that the organ which has such important work to perform should be large, and these and other recent views (even if not at present established beyond criticism) nevertheless help to explain the work which this gland is called upon to perform.

Before discussing the value of liver extracts, let us consider for a moment the therapeutic properties of secretin. It will at once be obvious that, provided this chemical can be prescribed for oral administration, and that when taken in this way it still exerts its characteristic influence, we have to hand a remedy of the very greatest possibilities.

Now, it appears that secretin-containing preparations can be so prescribed, and that they are active when taken by the mouth. As we have already seen, secretin requires an acid medium for its production, and that in the absence of an adequate degree of gastric acidity the normal secretion of this substance cannot take place. It would seem, therefore, that in cases where hypochlorhydria is suspected, especially where secondary digestive disturbances are present, the administration of an acid mixture with meals might be sufficient to insure the production of an adequate amount of secretin.

This is one aspect of the subject, and, there can be little doubt, an important one; for the benefit which so often accrues from an acid mixture taken about



MICROCOPY RESOLUTION TEST CHART

(ANSI and ISO TEST CHART No. 2)



1.45

1.50

1.56

1.63

1.71

1.80

1.88

1.96

2.05

2.14

2.24

2.34

2.44

2.54

2.64

2.74

2.84

2.94

3.04

3.14

3.24

3.34



APPLIED IMAGE Inc

1653 East Main Street
Rochester, New York 14609 USA
(716) 482 - 0300 - Phone
(716) 288 - 5989 - Fax

meal-times is frequently out of all proportion to what might be expected, if we only regard it as an adjuvant to gastric acidity.

Another question which arises is, Can we give secretin by the mouth? Undoubtedly the preparations which are on the market achieve excellent results in many cases, and the dry secretin-containing tablets and powders appear to exert a beneficial action in a wide variety of cases, which range from intestinal meteorism to glycosuria. But the dose of the substances requires regulating, as many observers emphasize the fact that small doses often succeed where large doses fail.

Secretogen, which the makers describe as "the master-key which unlocks the alimentary glands," is such a preparation, and many excellent results have been reported after its use. Again, secretin is sometimes combined with extracts of other alimentary glands, such as the liver, pancreas, spleen, etc.

It seems to the present writer that these preparations find a particular field of usefulness in those cases where intestinal symptoms are combined with signs indicative of pancreatic or hepatic insufficiency. Thus, intestinal stasis in combination with glycosuria would be a condition where a secretin-containing preparation might improve the symptoms in a quite remarkable manner, for it might in very truth "unlock" the duodenal hormone (or replace it so far as its pancreatic stimulus is concerned), and thus start the alimentary secretions, stimulating in its turn the

bile, succus entericus, etc., and supplying the missing hormone, the absence of which may have produced the entire syndrome.

From this, let us turn to the consideration of the value of prescribing liver extracts, and here we have to accept the testimony of those who have worked on these lines, and they are not numerous, as certainly in this country such administration has not been widely supported.

First of all we must enumerate the actions which liver extracts are supposed to possess, and the disorders in which they have been employed. Hepatic extract has been employed in functional disturbances of the liver, in those cases not always recognized in textbooks of medicine, but which are very well known in practice, and to which we may refer as "congestion of the liver," "hepatic inactivity," or "hepatic insufficiency." We may assume that in such cases, and it must be emphasized that we are referring only to functional disturbances of this organ, for some reason the secretion of the hepatic cells (not only the bile) is deficient, and that possibly not only is the conversion of the products of digestion incomplete, but that the filter-function of this organ is deficient. It used to be supposed that the symptoms of a "congested liver" owed their presence to bile in the blood-stream; but in the light of recent work it is much more probable that other and more complicated processes are responsible.

Extracts of liver have been prescribed in such cases,

much along the lines that other glandular extracts have been prescribed in diseases which we have already discussed—that is to say, with a view to stimulating the production of the necessary secretions, and to prevent the appearance of the symptoms which we know must appear if the absence of the secretion concerned is allowed to persist. But this is only one example of the uses to which these extracts have been put. Hepatic cirrhosis has been treated by the administration of liver extract, and, more especially in the atrophic variety, has met with some degree of success.

It has been stated that such administration has a marked action upon nutrition, and assists the urinary excretion, and that favourable results have followed its use in cirrhosis of the liver, as well as in functional congestion, and even in cancer. It is only rational to assume that, provided it can reach its goal without being destroyed on the journey, it will have the power either to replace normal secretion, or to stimulate that part of the secreting mechanism of the organ which has not been destroyed. That it must have some action in replacing the deficiency has been shown by those observers who have demonstrated that the changes in excretion alone have been marked after the use of these extracts. Thus, they have shown that the urea index is markedly increased, as well as the total quantity of urine; that anasarca has diminished, and that the tendency to hæmorrhages, which is common in cirrhosis, has been lessened.

Again, its use has been recommended in cases of hepatic diabetes, and where the diagnosis has been accurately made its employment should do good. Harrower suggests that a good plan to adopt when in doubt as to the nature of glycosuria is to commence treatment with pancreatic preparations, and to administer these for one month, and then if no favourable result be obtained to try liver preparations. He goes on to emphasize that the form of diabetes associated with liver disturbances has peculiar characteristics, notably that the glycosuria occurs at definite times, as after a meal or in the evenings; that the major symptoms of the complaint, such as thirst and polyuria, are not so marked, but that the tendency to infection and gangrene is well seen. From this he infers that the liver is failing in its antibacterial powers. It is in these cases, according to this author, that hepatic extracts are useful.

It will be quite obvious that in so far as the treatment of diabetes by liver extracts is concerned, diagnosis is the essential point; and that whatever reliance be placed upon opotherapy, general medicinal treatment should be insisted upon. For, as we have said elsewhere, the more we learn about this important subject, the more are we impressed with the fact that a regulated diet (in order to spare, so far as is possible, strain upon those organs concerned with sugar metabolism) is the first and most important point.

The dose of liver extracts is variable, but most authorities are agreed that it should be administered

in generous and regular doses. It is said that extract of this organ, administered orally or per rectum, is practically non-toxic, and that it may be given in doses ranging from 15 or 20 grains of the dried extract, given every hour, and increased to as much as 60 grains three or four times daily.

A few words about bile may not be out of place here. Bile has been administered to patients suffering from "sluggish liver," and with excellent results. It has been shown that the administration of even small doses of bile is followed by a marked increase in the amount of bile secreted; and it is along these lines that treatment by biliary extracts has been followed. It certainly seems more rational to administer the normal stimulus than to irritate the papilla of Vater with calomel. For relatively small doses of bile extract are cathartic—*i.e.*, a dose of 10 grains—while 5 grains in repeated doses is frequently laxative. The administration of the biliary extracts is followed in many cases by the disappearance of the signs of intestinal putrefaction, such as meteorism and the passage of offensive flatus, and by the re-establishment of regular and formed motions. Bile, although not in itself an antiseptic, acts in the intestine as such, for it prevents stagnation by its power of stimulating the secretions of the bowel.

In conclusion we must refer to the hormone elaborated by Zuelzer, and placed upon the market under the name of "Hormonal." This observer examined gastric mucous membrane, and after many experiments

succeeded in discovering a substance which proved to possess the power of increasing peristalsis. Further, he found that this substance was elaborated in the stomach and small bowel during digestion, carried in the blood, and for the most part stored in the spleen. As a result of experiments it was found that this substance, when injected into the blood, produced a marked increase in peristalsis. In another chapter we deal more fully with this hormone, so that at this place we will content ourselves with saying that the administration of this substance was followed by certain symptoms of an undesirable character, and that its use was discontinued in favour of "*neo-hormonal*," which is stated to be free from the objectionable features. *Neo-hormonal* is usually given by intramuscular injection, but it has recently been given by the mouth, and the results have been promising.

It is probable that the stomach and duodenum produce a substance which has the power of stimulating intestinal movements, and that this hormone is largely responsible for the regular and daily evacuations which are characteristic of health. If further experience shows that *neo-hormonal* can rectify an intestinal musculature which has become sluggish, it will prove of the greatest value to therapeutics.

Among the abdominal organs which have been said to influence digestion is the spleen. The exact function of this organ is still shrouded in mystery, although certain attributes are now recognized. That it is concerned in the formation of certain of the

white corpuscles, notably the lymphocytes, is now admitted, while it is believed that it also manufactures large mononuclear leucocytes. It is said to be the filter which removes broken-down corpuscles, both red and white, and it also drains the blood-stream, but its hæmopoietic function has long been established.

The question which concerns us here is, How is the spleen related to digestion? It would appear that it exerts a definite if somewhat remote effect upon digestion, for experiments upon animals have shown that splenectomized animals do not obtain nourishment from their food in the same way that normal animals do. In other words, this organ enables the main digestive glands to perform their work more adequately than is possible in the absence of the spleen.

It has been assumed from the results of these and other experiments that, while the spleen is not an organ that is essential to life, its purpose is partly to regulate the blood-stream and its contents, and partly to influence digestion, although our knowledge of its exact action in this latter respect is nil.

Its function as a filter is clearer, for does not the spleen enlarge in many acute infections, notably enteric? It is possible that it is concerned in preventing the products of intestinal fermentation from remaining in the blood-stream, and that it secretes a chemical which is destructive to micro-organisms. The enlargement which is so commonly seen in disease would then be accounted for by a hypertrophy

INTERNAL SECRETIONS OF DIGESTION 181

in response to a sudden demand for additional defences against infection.

This, of course, is purely speculative, and we have, at present, no evidence in support of this. But it is hardly conceivable that an organ of this size can be for no purpose, beyond the contributory hæmopoietic function which it shares with other organs. In spite of the fact that splenectomy is not followed by fatal results, and that after this operation other glands take on the known functions of the spleen, it seems highly probable that the future will disclose some fresh facts about the splenic function, and will prove beyond doubt that it elaborates a hormone which may be included among the internal secretions of digestion.

Splenic extract has been prescribed in a wide variety of cases. Thus, it has been given in typhoid with good results; in splenomegaly, in malaria, also in anæmia, and in morbid states associated with a low blood-count. Some observers speak enthusiastically of its influence in reducing the size of an enlarged spleen, more especially in what we may speak of as "idiopathic" splenomegaly.

The dose of splenic substance varies widely with different authorities, but large doses of the fresh substance are said to be badly tolerated. The dry extracts may be given in doses of a few grains, and as much as 15 grains daily are usually well tolerated. It undoubtedly finds a useful field of activity in exhaustive maladies, and in patients showing signs of a diminished resistance to infection. In the diseases just referred

182 THE ORGANS OF INTERNAL SECRETION

to it may confidently be administered, with the object of raising the resistance to the invading organism, assisting metabolism, and thus maintaining the strength of the patient.

This, then, is a brief survey of the internal secretions concerned in digestion, and of their relations to each other. We have attempted also to indicate the lines along which treatment by administration of the extracts of these organs can be carried out, and we have tried to show where such therapy may succeed. As such treatment is still in its infancy, and as there are a good many reasons why it cannot be expected to be successful in every case, this should make us all the more careful to diagnose, so far as lies in our power, the exact causation of every case which we treat with organo-therapy; and we should remember that, while certain of these preparations are toxic, others are not so. As an adjuvant to ordinary methods, the extracts referred to in this chapter should be found very useful.

REFERENCES.

- ¹ Sweet and Pemberton, Exp. Observ. on "Secreti.," quoted in *Practitioner*; H. R. Harrower, Practical Hormone-Therapy.
- ² Harrower, *ibid.*, pp. 64-65.
- ³ Langdon Brown, *Practitioner*, vol. xciv., No. 2, p. 252.
- ⁴ L. Williams, The Therapeutic Promise of the Internal Secretions, *Practitioner*, 1911, lxxxvii. 605.

CHAPTER X

THE THERAPEUTIC APPLICATION OF HORMONES

So far we have dealt more with the physiological and pathological aspects of the ductless glands and their secretions than we have with the therapeutic application, while some of the chapters have concerned themselves exclusively with diseases associated with one or other of the ductless glands. In this chapter we propose, so far as space will admit, to deal with the uses to which preparations of these glands can be put in practice; and we shall hope to outline the more commonly used extracts, and attempt to give some indications of their utility.

From this latter standpoint there can be little doubt that the thyroid must first claim our attention; for this, of all the endocrine glands, gives the best results to administration of its extracts—that is to say, administration over a duration of time. Starling divides hormones into two groups, the acute and the chronic—*i.e.*, whether their action is immediate or constant.

Now, what is the rationale of hormone-therapy, and with what object in view do we administer these extracts? Harrower epitomizes these questions in

his book, and he divides "the availability of hormone action" into four groups. The first is "substitution therapy"—the supplying of a missing hormone whose loss is due to destructive lesions or other cause of absence of the organ; the second is "supplementary therapy"—"the augmentation, directly or reflexly, of a presumed deficiency"; the third is "specific physiological therapy"—the production of a definite physiological influence which is indicated, but which is not due to any change in the normal hormone production; and the fourth is "empirical therapy."¹

Now, with this distinction emphasized we can commence to study the facts which have to be considered when we propose to commence the administration of extracts of the ductless glands.

First of all, we know that, with certain exceptions, these extracts are not toxic, although some of them have to be given in small doses and with great care. Thus, thyroid extract (as we have frequently pointed out) must on no account be given in the doses which are still found in books—that is to say, from 3 to 10 grains as one dose. Its administration should take the form of fractions of a grain, certainly to commence with. We shall go into the details of thyroid administration later. Again, extract of prostate is highly toxic; and if prescribed it is essential that the initial dose should be small, and its reactions carefully watched. Likewise, the extracts of adrenals and pituitary glands require judicious dosage, and cannot be prescribed in a haphazard manner.

On the other hand, the preparations of such organs as the liver, spleen, pancreas, testicle, ovary, etc., are in no way poisonous, and large doses can be given with comparative safety. In many instances, however, the administration of these latter extracts partakes of empirical therapy, for, frequently, the decision to endeavour to remedy certain symptoms rests more on a supposed deficiency than upon definite signs of inadequacy of a particular gland. Thus, in cases where intestinal stasis appears to be present, it may be assumed that this depends upon a deficiency of one of the alimentary secretions, to remedy which we prescribe a preparation of the gland in question. This, although empirical, has much to recommend it, and is surely preferable to constant dosing with purgatives, accompanied, as such treatment always is, by unpleasant sequelæ.

Another aspect of organo-therapeutic dosage has been defined by Hallion, and concerns itself with the stimulant action of an extract, administered to man or animal, on the organ secreting that extract. "Extracts of an organ exert on the same organ an exciting influence, which lasts for a longer or shorter time. When the organ is insufficient, it is conceivable that this influence augments its action, and, when it is injured, that it favours its restoration." It is here assumed that, so long as a part of a gland remains unaffected, it is possible to regenerate it from the secreting point of view, so that administration of its extract will have more than a temporary influence.

Again, even supposing that the gland is absent or totally destroyed, the administration of its extract will, in some instances—*e.g.*, thyroid—replace the natural secretion.

Still another aspect of treatment by extracts of the endocrine glands is what may be termed "antagonistic action." By this we mean the power of neutralizing an excessive secretion of one gland by the exhibition of an extract of another, assumed to be antagonistic. This adjustment of secretions in order to obtain the hormone balance is one of the most important features of organo-therapeutics, and in all probability the future will see its utility widely extended. In the past, as is well known, when Graves' disease was supposed to depend primarily upon an excess of thyroid secretion, treatment by means of adrenal administration was much vaunted, especially by the late Dr. Gibson, who spoke very highly of the results accruing therefrom.

There is also, in addition to the uses already mentioned, the physiological administration of organic extracts. As an example of this method we may cite the obstetrical use of hypophyseal extract to promote uterine contraction, and the prescribing of the same substance to counteract shock. These uses belong to the class referred to by Starling as "acute," only in the therapeutic sense. Starling, by his division of the hormones into acute and chronic, referred to those hormones which under physiological stimulus secrete suddenly an increased amount of a particular hormone

—*e.g.*, the pouring into the blood of adrenalin under the stimulus of fright, and the constant supply of thyroid for metabolic purposes, respectively.

It will be seen from this brief survey of the rationale of hormone-therapy that its uses are many, and that its application is logical. It is obvious that at present we are only on the fringe of the possibilities of such treatment, and that the near future will see its development to an immense degree. We do not mean that the empirical and sometimes illogical dosing by organic extracts, or the exhibition of every extract of the human body in the vain hope of "hitting the target somewhere," will be, or, for the matter of that, ought to be, widely developed. What we infer is that, with increased knowledge of the functions of the endocrine organs and their relations to each other, fresh fields of scientific prescribing will be open for use, and that many of the cases which we now treat with inorganic medicines, in the blind hope that Nature will step in and cure while we are gaining time, will be positively treated by organic extracts, with the almost certain knowledge that definite symptoms will disappear, and that normal features will reappear.

We will now consider the uses to which these extracts may be put in practice, and we will commence with the extract of thyroid.

In a previous chapter we dealt with the methods of prescribing thyroid extract for myxœdema and sub-myxœdema, and described the dosage. We will deal here with other uses to which this extract can be put,

but must say at once that many of these are founded on empirical reasoning.

For some years the nocturnal enuresis from which so many children suffer has been treated with thyroid extract, and, in many instances, with striking success. Not all these cases, it is needless to say, suffer from thyroid inadequacy, but many of them doubtless do so suffer, and the increased thyroid content of the blood effectually helps this enuresis. As Hertoghe has pointed out, a condition of thyroid deficiency nearly always produces frequency of micturition, owing to the poor nutrition of the walls of the bladder, which, in consequence, desquamate rapidly, rendering the bladder unduly sensitive to stimuli. In such cases thyroid extract is urgently needed; and unless it is supplied in increasing doses from within, or the internal supply augmented from without, it is difficult to see how the symptoms arising therefrom can be remedied.

This, then, is one of the uses to which this extract can be put, and another use is connected with this, inasmuch as the condition is frequently found associated with enuresis, obviously because the causation is identical. We refer to adenoids and enlarged tonsils. Hertoghe has rendered an inestimable service to medicine by explaining the underlying pathology of these cases. He emphasizes the induration and thickening which is always present where the thyroid secretion is deficient; not only in the subcutaneous tissue, but in places where formerly its existence had scarcely

been suspected. Thus, the tinnitus aurium and deafness of subthyroidics may be assumed to be due to infiltration of the aural passages and apparatus; the growth of adenoids to the same cause; the headaches and nervous symptoms to neural infiltration; and the enlargements of the tonsils to fibrous hyperplasia.

These, which are only a few examples, will serve to indicate the lines along which thyroid-therapy can be conducted. If, therefore, a child exhibits one or more of these symptoms, confirmatory evidence should be looked for. The subject of nocturnal enuresis may show the presence of adenoids, may have enlarged tonsils, or exhibit other signs of thyroid deficiency. The "adenoid facies"—*i.e.*, the mental condition plus the physical conformation of countenance—would be much more correctly described as the "subthyroidic facies." This being so, it will be obvious that such a child owes urinary symptoms of this nature to deficiency of thyroid, and not to local conditions *per se*; but we must emphasize in passing the importance of excluding certain local factors, such as calculus and phimosis. Fascinating as this form of treatment is, and almost limitless as the vista of healing is which is opened by its conception, diagnosis must always precede treatment.

For these patients a small dose of thyroid, increased gradually as is required, is indicated. The initial dose should be a fraction of a grain, according to the age of the patient. Thus, one-sixth to one-quarter of the dry extract or a minim or two of the liquor thyroidei

190 THE ORGANS OF INTERNAL SECRETION

is enough to commence with, and a careful watch is essential to ascertain the reaction of the system to the extract.

Again, obese patients should be examined for signs of thyroid inadequacy, as where such increase in weight is of recent occurrence the treatment should be at once instituted—the sooner the better—once the diagnosis has been made. Even if the signs of thyroid deficiency are of long duration, the prognosis is equally, or almost equally, good; for although the treatment must be spread over a much longer time, nevertheless the benefit accruing is equally great.

Some of these cases are not solely due to thyroid inadequacy, but are a mixture of thyroid and pituitary deficiency. It is a little difficult to describe the diagnostic features in a few words, but the indications of thyroid inadequacy are a slow pulse, dry and rough skin, subnormal temperature, and obesity; while the signs of hypophyseal deficiency are an increase in weight, a smooth skin—as opposed to the rough skin of submyxœdema—a low blood-pressure, and a disturbance of function, as exemplified by constipation, with often a marked degree of meteorism and faulty digestion, which latter is often seen in subthyroidics.

For such cases thyroid should be prescribed, and when the dose has reached what may be regarded as sufficient for the individual case, but still symptoms remain which do not improve, then pituitary extract may be added to the thyroid. Such a procedure will often help where thyroid by itself fails; and, provided

the dose of both substances is regulated to the individual case and not by rule of thumb, no ill results should accrue.

The preparations of thyroid which may be utilized have already been mentioned, but in order that this chapter may be helpful in actual practice, we will enumerate them once more.

The total extract of thyroid may be given, or the dry extract chosen, if preferred; one part of the dry extract is approximately equivalent to five parts of the other extract. The dose of extracts should be minimal to commence with, and opinions vary as to the margin of safety. So far as the experience of the present writer is concerned, one-half to one grain of the total extract three times a day is a good dose to commence with, and can be increased with due observance of the alteration in weight, pulse-rate, and blood-pressure. Again, if a liquid preparation be preferred, liquor thyroidei, in doses of from a few minims to about seven or eight, is a satisfactory method of administering this extract, provided that the preparation be fresh. Elixir Colloid (Squire) presents this gland in a palatable form, and it can be combined with a pituitary extract if desired. The same firm manufactures a liquid preparation of the pituitary gland, under the name of Elixir Hypophysis Cerebri, with a dose which may range from a few minims to two drachms. These preparations are a convenient way to administer the thyroid and pituitary glands to the same patient at the same time, and the relative doses may be regulated with ease.

The hypophysis may be given in tablets, and several firms manufacture this extract either by itself or in conjunction with other extracts. There can be little doubt that many patients whose symptom-complex resists almost all kinds of treatment will receive very marked benefits if treated by these means.

From considerations of space, we must now consider other organic extracts and their methods of administration. We are not so much concerned here with the urgent use of these extracts, as exemplified by their utilization in shock, obstetrical emergencies, and surgical cases, and we shall therefore pass over the employment of adrenalin, pituitarin, and other preparations of these glands for local administration, and deal with their use from the routine standpoint only.

The extract of the adrenal gland has definite and powerful properties, and, if used, requires great care in its administration. It is not suitable for prolonged administration, and there are certain cases in which its use is quite contra-indicated. As it raises the blood-pressure and causes a rapid rise in the pulse-rate, it should never be given in cases of arterio-sclerosis, or in hyperpiesis where there has been hæmoptysis, or in glycosuria. In fact, its prolonged administration is not desirable, and, considering that extract of the pituitary is much less toxic, and that the results are almost as favourable, the administration of adrenal extract has not become popular. Its local action, either alone or in combination with a local anæsthetic,

is of great use; but so far as we are concerned at this place, we need not spend further time on its discussion.

Extracts of the gonads, which are sometimes manufactured in combination with extracts of the central nervous system, will be found to have a wide applicability. Thus, in neurasthenic cases, in patients who are run down, and in delayed convalescence, they are frequently of the greatest use. As Brown-Séguard was the first to show with reference to testicular extract, they increase the capacity for muscular work, revive the tired nervous system, and have a general "tonic" influence upon the functions. The present writer has utilized them for several different types of disease (or perhaps the word "disorder" is more suitable), such cases ranging from marasmic wasting to impotence. It is a little difficult to give definite indications for their use, and perhaps it will be better to cite one or more cases in which their use at the hands of the present writer has been attended with success. One was a man aged thirty, with rapid wasting and low blood-pressure, who had been in a very depressed mental condition for over two years. His treatment consisted, *inter alia*, in rest in bed for six weeks, with hyperalimentation and this form of organo-therapeutic treatment. At the end of a month his weight had increased by over one stone, and his improvement in every way was most striking. By the end of six weeks there was a continuance of this improvement, and when he left the nursing home his weight had increased by a total of two and a half stone, his mentality was, for all

intents and purposes, normal, and his physical health quite restored.

Another instance of the same treatment was in a man of middle age, prematurely old, with impotence, neurasthenia, hypopiesis, and depression. One month's treatment by these methods restored him to normal, raised his blood-pressure, and restored his nervous functioning. In this case, moreover, there was a flabby condition of the genitals, a loss of sexual power, and testicular sensation was decreased.

A preparation called "spermin," isolated by Von Poehl, and believed to benefit a number of conditions, most of which emanate from auto-intoxication, has been tried with some success. It has also been suggested that extracts of the gonads are useful in raising the resistance of the body to infection; there can be little doubt that they are valuable in "revivifying" the overworked organism.

In dealing with the hormones which may be supplied from animals, we must refer to those emanating from the alimentary canal and the abdominal organs. Thus, we encounter "secretin," which was isolated by Starling from the duodenal mucous membrane. In normal health this substance is set free from this mucous membrane as soon as the acid chyme enters the bowel. It is difficult to prescribe, for it appears to be destroyed when given by the mouth. Nevertheless, there have been some good results published, notably by Harrower, and this should encourage its further trial. As this hormone is a direct stimulant

to the pancreatic juice, its absence or diminution must be attended with some degree of deficient metabolism. A manufacturer's preparation known as "secretogen" is on the market, and the present writer has employed this with some benefit.

Again, Zuelzer, experimenting to ascertain some substance which would have a decided action in increasing peristalsis, discovered a hormone, prepared from the gastric and duodenal mucous membrane, which showed this property. He considers that this hormone is produced in the stomach and duodenum, carried off in the blood, and stored in the spleen. A preparation known as "hormonal" was then manufactured, mainly from the spleen of animals, and administered with or without pituitarin. Apart from cases of spastic constipation and the like, this is, as Harrower points out, a useful remedy for deficient peristalsis.

As there was some danger from the use of hormonal, which was apparently due to the presence of an albumose, this remedy has now been discontinued, and a fresh substance, with the harmful constituent removed, has been placed on the market, and the results quoted in Harrower's book speak very highly of its action, more especially in cases of intestinal paresis.*

Another combination of an internal secretion with the external secretion is to be found in "trypsogen," which contains the pancreatic secretion and ferments. This is recommended for use in cases of diabetes, or

* This preparation has received the name "neo-hormonal"

where it may be presumed that the pancreas is at fault. Another use for the same preparation is in hypertension; but there are other methods of treating this condition, which must only receive organo-therapeutic treatment of any kind when its origin has been determined.

We will pass over the administration of bile-salts as a rational laxative, as the subject is too large to admit of adequate treatment in a chapter of this length. Suffice it to say here that bile-salts are clearly indicated in some cases of constipation, more particularly where there is reason to suspect a disordered liver, with deranged bile secretion into the bowel. A few grains of bile-salts, about three to five, form an excellent laxative, while in larger doses such an extract is cathartic.

Before concluding this rapid survey of hormonal preparations, we must refer to the less definite internal secretions and their administration. Of these, ovarian extract, either whole extract or else extract of the corpus luteum, has been given in a variety of conditions, either alone or with one of the other endocrine preparations. It has been found useful in disorders of the catamenia, at the climacteric, for post-operative disorders, such as those which follow double oöphorectomy, and as an antagonist to certain of the other internal secretions. In similar manner, the extract of the mammary gland finds a field of utility, and has been prescribed for uterine conditions (it will be remembered that there is a close connection between

the uterus and the mammæ) and as a galactagogue. Some observers speak highly of its efficacy in this connection.

We have already mentioned the use of prostatic extract, and stated that it is a toxic substance, and therefore of limited utility. The present writer has had insufficient experience of this extract to speak definitely; but it is stated to be useful in controlling the peculiar melancholic condition which sometimes supervenes after removal of the prostate.

General Survey of Organo-therapy.

We will now briefly recapitulate what we have said with reference to the uses of organic extracts in medicine, and amplify our remarks by quoting some typical cases in which its administration has proved beneficial. As we have arranged this book so that a brief summary of treatment will be found at the end of each chapter, it may be convenient to summarize our conclusions in one chapter, so that, if desired, the therapeutics dealt with in this volume may be gleaned from a perusal of one chapter.

Organo-therapy holds out the hope that it will ameliorate the unhappy lot of many patients suffering from diseases which have hitherto been intractable to most forms of treatment. It has been welcomed with open arms; its component extracts have been prescribed in many maladies, and the reports as to the results of this medication have been various and frequently con-

tradietory. So far as the lesser-known organic preparations are concerned, this book has not had the requisite space to deal with them; neither do they seem, if we may say so, to come within the scope of a work such as this.

The author has endeavoured to describe the grosser lesions in which organic preparations may be employed with a prospect of success, and he has attempted to outline morbid conditions in which such extracts can be prescribed. It must be remembered that an organic preparation may be used in several ways; thus, it may be given to replace the same secretion, which is assumed to be deficient (substitution therapy); it may be administered to increase a secretion which is presumably not supplied by the gland in sufficient quantity; it may be prescribed in the hope of antagonizing the excessive secretion of one or more of the other internal secretory glands; or it may be given purely for its physiological action.

A vast range of utility is thus opened for preparations of the organic extracts; and now that the signs and symptoms of deranged secretion are more widely known, the application of the theoretical knowledge of the endocrine glands to actual therapeutical practice is becoming easier than when such prescribing was almost solely empirical. As an example, the many different abnormalities which thyroid extract will correct, in one or other of its applications, are daily receiving additions, as we discover fresh symptoms of thyroid insufficiency.

Originally its therapeutic use was confined to the treatment of myxœdema. When this disease was seen to be remedied by thyroid feeding, a mild type of the same disease was recognized as yielding to the persuasive powers of this extract. Submyxœdema as a clinical entity has been supplemented by a variety of symptoms, any one of which, when present, may sometimes be ascribed to hypothyroidism. Thus, we may say that a consistently slow pulse points to a deficiency in the thyroid output, although other symptoms must necessarily be present before we can diagnose "submyxœdema." But it does not follow, of course, that every person with a slow pulse has hypothyroidism, although, by the physiological use of this extract, it might be justifiable to accelerate this slow pulse-rate, were it desirable, by the judicious administration of thyroid.

Again, any of the symptoms which are so well recognized when grouped together in the same patient, as pointing to the thyroid as the organ at fault, may be treated by means of this extract, provided that a careful watch is kept of the reaction of the patient to this remedy.

A few examples will explain how this principle may be applied to practice.

CASE 1.—M. W., *æt.* 33, married, with one child. Shortly after she became pregnant (three years before the date she first consulted me), she noticed she was feeling better in her general health than she had done for some years, as she had previously suffered from

200 THE ORGANS OF INTERNAL SECRETION

indigestion and constipation, and her "nerves had been bad." She informed me that shortly after her child was born she relapsed, and that about this time brown patches appeared upon her face. These had continued ever since, and her health had never been restored to normal.

On examination, very marked patches of a light brown colour were seen on the forehead and around the mouth. These were so marked as to cover nearly the whole forehead, although there were clear areas of normal skin between. Her pulse was slow, although not markedly so, and the thyroid gland was prominent, the neck measuring $13\frac{1}{2}$ inches at the level of this gland. She was quite abnormally sleepy, and became unduly tired.

I prescribed one half-grain tablet of thyroid gland daily, to be increased in a week to one grain in the day. This was gradually increased until she was taking three one-grain tablets daily.

The result was most gratifying. Three months from the commencement of treatment the patches had almost disappeared, and small areas only of slightly darker colour than the surrounding skin could be seen in a good light. She stated that her general health was much better, and she remarked that "she felt fitter than she had since her baby was born." The thyroid had decreased in size, and the neck measurement was barely 13 inches.

Note.—This case is instructive for the following reasons:

1. The thyroid secretion had evidently increased during pregnancy, when she experienced better health, but had diminished after parturition, which is often the case.

2. The patches of pigmentation yielded at once to the ingestion of thyroid.

3. The general health improved *pari passu*, both the digestion and the bowels functioning more normally.

4. The decrease in the thyroid gland under this treatment pointed to the swelling as a hypertrophy of the gland, an effort to increase the deficient secretion.

CASE 2.—J. W., *æt.* 47, complained of an increasing weight, accompanied by dyspnoea on exertion, broken sleep, digestive disturbances, and constipation. This condition had been of gradual onset, but had been rapidly increasing of late.

On examination, the increase of weight was largely accounted for by deposits of fat in the regions of the abdomen and shoulders; the pulse was about 70, the temperature subnormal, the blood-pressure was 145 mm. Hg, and the urine contained large quantities of phosphates. The skin was dry and rather rough, and the patient most markedly bald; this latter feature he had treated by the adoption of a wig.

I restricted his carbohydrate food, ordered an occasional saline, and commenced with small doses of thyroid. In a month's time he had lost 9 pounds, and felt and looked better. His digestion improved, his sleep became more normal, and his energy returned.

202 THE ORGANS OF INTERNAL SECRETION

This is a year ago, and he still remains in good health, although a relapse in weight occasioned a fresh course of thyroid medication.

CASE 3.—An analogous instance is the case of Dr. A. This gentleman consulted me for similar symptoms, but certain points in his case were different. Thus, his inertia was greater, fatigue on exertion was most marked, his blood-pressure was lower, and his general syndrome seemed to me to point to a pluriglandular deficiency.

He took, at my suggestion, small doses of thyroid combined with dry extract of the adrenals. He improved markedly, but had to continue the medication at intervals in order to avoid a recurrence of the symptoms.

Note.—Both the above instances presented neurasthenoid symptoms, but the improvement in each case will afford evidence that many so-called “neurasthenics” are in reality sufferers from a disturbed hormone balance, and in consequence may be helped by this form of treatment.

CASE 4.—M. P., *æ*t. 34, complained of undue fatigue, of feeling very unwell each morning on waking, and of a “quite extraordinary tendency to fatigue.” He informed me that he had “always had a very slow pulse, as his father had before him.” This was certainly true, as his pulse-rate at the first consultation was about 50 per minute. His blood-pressure was low, his temperature subnormal, but his weight was about normal for his height.

There is little further need to discuss this case, for the diagnosis was obvious. His symptoms all yielded to the administration of thyroid extract, which increased appetite, regulated sleep, and approximated the pulse-rate to normal, so that his circulation improved and he was able to resume his normal life, which entailed a considerable amount of active exercise.

CASE 5.—An interesting case came under my notice some time ago. A young girl was brought to me whose mental development was deficient, although physically she was quite normal, with the exception of a markedly protuberant abdomen. In this case, as growth proceeded, she had exhibited signs of a lack of mental development, and an eminent surgeon had decided to operate and implant a thyroid graft, in the hope that the thyroid secretion would replace the hypothyroidism which was assumed to exist.

When I saw this patient it was five years since the operation, and the main features of the case were as follows:

Her stature was decidedly above the average, and her weight over 12 stone. She was fully developed physically, but her circulation was so poor that she suffered from chilblains on the hands, feet, and legs. Her pulse was slow, about 48 per minute, her temperature markedly subnormal, her blood-pressure low, but there was nothing abnormal in the heart. In the epigastrium and umbilical regions was a large swelling, tympanitic on percussion, which displaced the

liver, and cardiac dulness upwards. An X-ray examination in conjunction with a bismuth meal had failed to locate this. The skin was harsh, dry, and scaly. The bowels were regulated by means of liquid paraffin.

This case appeared to be due to thyroid deficiency, but it will readily appear that this was not the only secretion at fault. The patient's father came of a family of giants, and it seemed probable that there was an element of *dyspituitarism* in this case; that is to say, that the patient may have suffered during childhood from a condition of unbalanced pituitary action, which had been followed during adolescence by hypopituitarism.

Acting upon this assumption, I prescribed a mixture containing a small dose of thyroid extract and a large dose of pituitary. The improvement has been marked, particularly as regards the circulation, but enough time has not elapsed to enable a definite result to be recorded. The abdominal condition is slowly improving, and the tympanitic mass is responding to the stimulation of the alimentary canal by the hypophyseal extract. This swelling I assumed to be a dilated transverse colon which was consequent upon the deficient innervation of the bowel.

CASE 6.—W. P., a case of advanced neurasthenia with marked loss of weight. This patient suffered from mental depression and digestive atony, and accompanying neurasthenic symptoms.

There was also evidence of a lowered blood-pressure and a sluggish cardiac action. Although this patient

had been under treatment for some time by many different methods, no improvement had taken place. However, a short course of treatment by a pluri-glandular preparation was followed by a disappearance of nearly all the abnormalities, and the patient returned to his home and resumed his normal life, which he had been unable to do for three years previously.

The above cases will show the lines upon which treatment by the more important organic extracts may be regulated. The legitimate uses to which thyroid extracts may be put are not confined to myxœdema, or even definite submyxœdema; there are a number of symptoms, sometimes isolated, which may be remedied by this substance. Case 1, quoted above, will show that one symptom will often point the way to the underlying pathology.

But, as was seen in Cases 3 and 5, the full benefit was derived by a combination of extracts—in the former by the association of thyroid and suprarenal extracts, in the latter by that of thyroid and pituitary. It is not always easy to decide on the particular combination to employ, neither can definite reasons always be given for the choice. But if the signs which accompany one abnormality appear to be contradictory, either the preliminary diagnosis may be at fault, or else more than the one gland may be at the root of the trouble. As an example, thyroid deficiency produces a harsh, dry skin, hypopituitarism a smooth one. Again, the condition of the hair would lend assistance,

as it is scanty and "lifeless" in submyxœdema, but, so far as the head is concerned, is usually the reverse in deficiency of the pituitary secretion.

Again, where thyroid deficiency is associated with hypopiesis and loss of flesh, with languor and undue fatigue, adrenal extract may wisely be combined with thyroid. Such treatment will frequently afford relief in this type of patient, who is very commonly encountered in practice.

There are on the market many pluriglandular preparations, and to these we have already referred. It is scarcely necessary to say that they should only be given where the prescriber knows the exact dose in each tablet or solution; for, if he is in ignorance of the composition and proportion of the constituents, he may well be unable to account for any untoward symptoms which arise. It is probable that the improvement which follows the employment of a pluriglandular preparation is sometimes due to the stimulating effect it has upon the endocrine glands as a whole. Thus, cases of acromegaly have been reported as cured, or much relieved, by preparations containing the extracts of the thyroid, sexual glands, spinal cord, etc., presumably by the restoration of the hormonal balance produced by the antagonistic extracts.

And now a few words to summarize the utility of the more important organic extracts. We will turn our attention first to the thymus gland.

Extracts of this gland have been used in a variety of diseases, ranging from exophthalmic goitre to de-

ficient development. Some observers speak well of its employment in simple as well as exophthalmic goitre, in chlorosis, in rickets, in delayed union of fractures, in rheumatoid arthritis, and in deficient growth. One fact stands out pre-eminently from the mass of supposition in relation to this gland and its functions, and that is that it undoubtedly plays an important part in the calcium metabolism of the body.

It is generally admitted that this gland atrophies about puberty, except in exceptional cases—in fact, there appears to be an antagonism existing between the thymus and the sexual organs. Ovariectomy is followed by thymus hypertrophy, and the removal of the thymus in some animals produces rapid development of the sexual organs.*

Several authorities also report the changes in the skeleton which have followed experimental thymectomy, and these consist in deprivation of the calcium of the bones, resulting in spontaneous fractures, osteomalacia, and other abnormalities.

It would seem, therefore, that the thymus is concerned with the regulation of the lime-salts in the body, and it is mainly upon this assumption that its extract has been administered to patients suffering from exophthalmic goitre. In this disease the calcium metabolism is defective, and many authorities maintain that the administration of calcium salts should form part of the treatment in every instance. Likewise, the

* See article by Noël Paton, "The Relation of the Thymus to the Sexual Organs," *Journ. Physiol.*, 1905, xxxii. 26.

prescribing of extract of thymus might help in this direction, for it evidently exerts a favourable influence in rickets, where the lime-salts are deficient in quantity. It is usually prescribed in doses ranging from a few grains to half a drachm daily.

We have already discussed the uses of parathyroid extract, but perhaps we may here recapitulate the subject. It has been assumed that these small glands regulate in part the supply of calcium to the cells of the body, and that it is the withdrawal of the calcium which produces the tetany following experimental parathyroidectomy. On this assumption, the extract has been prescribed in a variety of diseases, notably Graves' disease, paralysis agitans, and epilepsy. The dose is small—a fraction of a grain of the dry extract—and should be given once or twice daily. We have already referred* to a most interesting case, reported by Hertz, in which definite parathyroid insufficiency was cured by the administration of parathyroid extract.

In conclusion, let us briefly enumerate the organic extracts which are at the disposal of the practitioner, and their approximate doses. It must be understood that the doses are only approximate, and that no attempt is made to give an exhaustive list, only the more important of the extracts being mentioned.

Thyroid.—The preparations are the dry and the liquid extracts, the dose of the former being from $\frac{1}{4}$ to 1 grain for an initial dose; of the latter 2 or 3 minims. Many products are upon the market in the form of

* Chapter I., p. 24.

elixirs, capsules, and tablets, either alone or in combination with other organic extracts.

Pituitary.—Two classes of pituitary preparations are in use, one manufactured from a part of the hypophysis (usually the pars posterior), the other from the entire gland. The preparations of the posterior part of the gland are more commonly prescribed, and the dose ranges from 1 to 5 grains of the entire extract. The dose of the liquid hypophyseal extracts ranges from 8 to 15 minims, which may be repeated at frequent intervals.

Adrenals.—As we have already said, the extract of these glands is more toxic than that of the pituitary. The liquid preparation may be prescribed in the form of adrenalin, the dose being from 5 minims to $\frac{1}{2}$ drachm of 1 in 1,000 solution. This preparation may be administered hypodermically. The dry extract may be given in doses of 1 to 3 grains.

Pancreas.—The average dose of preparations of the pancreas is 3 grains. The preparation "trypsin," referred to in the text, is put up in 5-grain tablets.

Testicles.—Orchitic extract is given in doses ranging from 2 to 4 grains, according to the individual preparation. There are also many liquid preparations on the market, likewise products combining this extract with that of the central nervous system. (See Pluriglandular Extracts.)

Ovary.—Extracts of the corpus luteum are usually administered. The average dose is 3 grains.

Mammæ.—The mammary gland preparations have been mainly used for menorrhagia, as an antagonist

210 THE ORGANS OF INTERNAL SECRETION

to the ovarian secretion (see note on pluriglandular preparations), and in the treatment of uterine fibroids. The dose for the first of these purposes should begin at 5 grains and be increased to about 10 grains; to antagonize hypo-oöphorism, the dose must necessarily be elastic; while for the treatment of uterine fibroids the dose is about the same as that for menorrhagia, but varies according to different authorities.

Prostate.—The preparations of the prostate gland are given in doses ranging from $\frac{1}{2}$ to 3 grains.

Thymus.—The dose varies from 3 to 10 grains. It is stated that care should be exercised in prescribing this extract to gouty patients, on account of its richness in nucleins.

Intestine.—Preparations of the duodenum containing pro-secretin are given in doses of 1 to 3 grains. "Secretogen" tablets contain duodenal pro-secretin and succus entericus.

Liver.—The dose varies enormously with different preparations. Large doses may be given, however, even up to 1 drachm, of the vacuum-dried extract. An average dose, however, is 15 grains three or four times a day.

The dose of bile is from 3 to 5 grains; 10 grains is a purgative dose.

Spleen.—Of the solid extract, 5 grains is an average dose; up to 15 minims of the liquid preparations.

Parathyroid.—In exophthalmic goitre, paralysis agitans, chorea, and epilepsy, the dose is from $\frac{1}{30}$ — $\frac{1}{10}$ grain.

Pluriglandular Preparations.— These, of course, present many different combinations. Perhaps the most usual combination is that of the extracts of nervous tissue—the brain and spinal cord—with extracts of the gonads. To such a preparation some manufacturers add extracts of the thyroid, pituitary, or thymus. In the author's opinion, this is to be deprecated, as the reaction of the patient is difficult to judge in the presence of such complicating compounds. The correlation of extracts of the ovary and breasts is likewise unwise, as their actions are by no means similar; in fact, in some cases—*e.g.*, the use of mammary extract in menorrhagia—their effects are opposed. On the other hand, there is much to be said in favour of pluriglandular therapy, as many "neuras-thenic" conditions are undoubtedly examples of hypo-endocrinism, and as such benefit from this form of therapy.

Care is needed, however, in commencing "tonic" treatment by these means, as a condition of irritability—the reaction from the torpor of hypo-endocrinism—is apt to result in a certain proportion of cases when these preparations are first administered. Nevertheless, the help they can give is very great, especially in functional neuroses; but the preparation must be chosen with care, and the reaction of the patient to this form of medication noted, so that any changes may not escape notice, as it may be necessary to alter the component parts of the product, and re-arrange, so to speak, the proportions of its constituents.

212 THE ORGANS OF INTERNAL SECRETION

The present writer is in the habit of picking out the extracts which he considers are required in an individual case, and combining them in the proportions which seem most suitable. In this way a more accurate dosage is possible, and the constituents may be readjusted should this be necessary.

This, then, is a brief summary of the more important organic extracts which are available for administration, although the list is by no means exhaustive. It is hoped that the preparations here enumerated will assist in the cure of many cases hitherto regarded as "chronic."

Every medical man encounters from time to time cases which are not relieved or even benefited by ordinary methods of treatment. In many of these patients relief would be afforded, if not actual cure, by organo-therapy. It is hoped that this small work will succeed in interesting those who have not hitherto adopted this method of treating disease, in order that a large number of so-called incurable diseases, and a larger proportion of functional disorders, may be remedied by these means.

REFERENCE.

- ¹ Harrower, Practical Hormone-Therapy, pp. 22, 23.

CHAPTER XI

CONCLUSIONS

HAVING dealt with the pathology and treatment of the organs of internal secretion, and having briefly outlined the organic extracts which may be utilized in the treatment of disease, it may not be out of place at this point to discuss the present place of hormone-therapy in medicine.

Not many years ago an eminent physician, in speaking of the use of medicines, said that in prescribing a drug a medical man was putting a substance of whose properties he knew little into a body about which he knew less. This may be only too true with reference to the administration of inorganic chemicals, or, rather, with reference to certain inorganic chemicals; but now that organo-therapy has become an established method of treatment, it can scarcely be applied indiscriminately to all forms of prescribing. For it cannot be denied that much drug-therapy is empirical, and is resorted to in the hope of counteracting morbid conditions of the body, and that much of such treatment would be quite useless were it not accompanied by the hope and faith of the patient.

In prescribing organic extracts, however, we are

214 THE ORGANS OF INTERNAL SECRETION

endeavouring to counteract morbid states by administering the substance the deficiency of which we believe to have caused the condition. We gather together the salient features of an individual case, and compare the syndrome with that which is known to be present in derangements definitely due to disturbances of the hormone balance. If this points decidedly to a deficiency of one hormone the path is clear, and the result will be steady improvement to cure. But if a multiple disturbance is present, and it must be remembered that this is very often the case, the course of the treatment will not be so smooth, neither can the first combination be expected to prove successful.

Nevertheless, even with this class of case, the adoption of organo-therapy is less speculative than is the prescription of many inorganic remedies, which, in ameliorating one symptom, are prone to initiate another. With obvious hypothyroidism or hypopituitarism, the only remedy which will give a speedy and satisfactory cure is the administration of the organic extract whose deficiency has produced the symptoms. But, apart from such straightforward cases as these, there are a host of physical and mental deviations from normal health which may be rectified by this method of treatment.

We have already outlined these, but we desire to emphasize in this chapter that the use of organic preparations is now an established method of treating disease, and one which will undoubtedly form a large part of treatment in the future.

In an article dealing with the question of the treatment of disease which was published recently in a prominent journal, the opinion was expressed that the majority of chronic diseases of middle age could only be palliated—that their cure was impossible. The writer proceeded to emphasize the fact that these diseases arise during early youth, and that our ignorance of the origin of chronic disease is profound. This latter statement is, unfortunately, correct; but the first clause which we have quoted was probably truer twenty years ago than it is now. For every year that passes increases our knowledge of the efficacy of modern methods of treatment, and not least among these we must class organo-therapy. Before the discovery that myxœdema owed its origin to atrophy of the thyroid gland, and that the symptoms caused thereby yielded to the administration of extracts of this gland, the features of this malady were hopelessly intangible, and their treatment, either as individual symptoms or a syndrome, impossible. This disease may be included among those which the writer of this article had in mind; but although the actual cause of the atrophy is unknown, it is quite untrue to say that its palliation only is possible. For both in the grosser disease myxœdema and in the lesser malady submyxœdema, the administration of thyroid gland is curative—in so far as the disappearance of symptoms constitutes a cure. It is quite true that the treatment must be continued, even throughout life, but so long as it prevents the disease from persevering, we have achieved what may be called a cure.

This is only one disease, and one among thousands; but it is nowadays curable, and it is probable that, with the advance of our knowledge on these lines, many other lesions included in the past among the incurable or chronic disorders will yield to modern therapy. Not only the vague neuroses—the neurasthenic, the neurotic, or the hypersensitive patients, whose existence is due to the civilization we boast of—but diseases which, in the past, have left their victims more or less crippled for life, will in the future probably be treated so that they no longer maim permanently.

Many of these diseases are accountable for the derangements seen in middle age, and by the time advice is sought the machine can only be repaired, not restored. But the repairs will be less needed when our knowledge of the treatment of the diseases incidental to childhood is improved, and it may be that the endocrine glands will figure largely in such improvement. If, for example, we can increase the defences of the body against infection by the administration of an organic extract (as is now done in the case of tuberculosis), we shall have developed a weapon against which the onslaught of micro-organisms will be weakened, if not defeated. In which case the germs responsible for the exanthemata, with their damaging sequelæ, will find their task less easy than formerly.

Even the organic changes which we are accustomed to regard as inseparable from middle and old age may be remedied or averted by the employment of organic extracts. An example will serve to make our meaning

clearer. The complex metabolism of the body appears to be regulated by the organs of internal secretion; a deficiency or excess of one of these important correlators is able so to upset the metabolism that substances, often of a noxious nature, are stored in the body instead of being eliminated by the usual channels. In course of time this gradual retention may produce a metabolic disease such as gout or diabetes, and it may assist in the well-known accompanying changes such as are present in chronic interstitial nephritis, in hepatic cirrhosis, in fibrositis, and numberless other deviations from normal health. If deficient elimination is treated in the future from the commencement—*i.e.*, when the first signs of metabolic disturbance are present—by the exhibition of one of the organic extracts which regulate metabolism, instead of by occasional purgation, we may well hope that the “diseases of middle age, which arise during infancy or adolescence,” may no longer be regarded as a necessary concomitant of advancing years.

This, then, is what we mean by the statement that our knowledge of organo-therapy may conceivably be extended to the avoidance of organic lesions in the near future. We have already pointed out that the thyroid, pituitary, and pancreas, to mention three out of many, are intimately concerned in the control of metabolism, and that a deficiency in secretion of either of the first two glands results in a storage of waste products, in thickenings of the subcutaneous tissues, and in the retention of chemical substances which

should leave the body. Such conditions disappear upon the administration of the extracts of these organs.

By comparing these syndromes with that which accompanies, let us say, "idiopathic" obesity, or the gouty diathesis, we must at once be struck by the similarity of the two pictures. Therefore it does not seem to be stretching a point to say that, were these latter diseases suitably treated in the early stages, it would be possible so to regulate metabolism and to effect adequate elimination that the system would be restored to the *status quo ante*. The advanced stages of faulty elimination which are now encountered in the out-patient and consulting-room could thereby be avoided.

This is, doubtless, looking far ahead; but if we are justified in assuming that the majority of the diseases of middle age date from childhood, and are in consequence incurable, it behoves us to look around for a means of avoiding such a lamentable state of affairs.

Slight degrees of hypo-endocrinism usually recover without aid from opo-therapy. If, however, they do not, it is conceivable that the perversion of metabolism commences from such a condition, initiated, let us say, by an infectious disease. The faulty elimination is at first slight, but increases; the individual takes less exercise as his age advances, and when he reaches his fourth decade he is probably prematurely old, physically unfit, with yet no definite signs of organic disease. Nevertheless, another ten years may have altered his condition for the worse: the blood-pressure

may be high, the urine show a trace of albumin and a decidedly acid reaction, the joints be stiff, the hirsutes scanty, and the skin dry and its appendages lifeless.

Every practitioner is familiar with such a type—a man aged at fifty! Even now he will feel benefit in many cases by an organic remedy. But twenty years before what a difference the adjustment of his hormone balance would have made to his future health and comfort!

The present writer has seen many ailments, slight in themselves, but of some considerable tenacity, disappear upon the administration of thyroid extract. He has also seen patients, who admit that "there is nothing wrong with them," attain a degree of health which they had not known for years from the same treatment. He is convinced, therefore, that a great future lies in front of organo-therapy, but it must be given every chance: to wait until the patient's system has become altered by years of faulty elimination is to deny the rational remedy a fair chance of success.

This, then, is a plea for the early use of organic extracts in every case where their administration appears to be indicated. By their prompt use there can be small doubt that many derangements will be righted before they can become diseases, and it will not be necessary for our successors in medical practice to make the lamentable confession recorded by the writer of the article referred to above.

* * * * *

This little book has aimed at presenting to the reader a short survey of the organs of internal secretion, their diseases, and the uses to which preparations of the hormone-producing glands, and of organic tissue generally, may be put. At the close of this chapter will be found a short bibliography, and the reader anxious for a fuller account of this subject is referred to the works there mentioned. Many of these deal with one gland only, while some are general surveys of organo-therapy.

Among the large treatises which have been published in the last few years is Cushing's book on the pituitary gland; Léopold-Levi and H. de Rothschild's study of thyroid deficiency; and a monumental work by Allen on glycosuria and diabetes. The study of lengthy and exhaustive treatises such as these is often impossible for the busy practitioner, and this small work has attempted to summarize the subject in as brief a space as is possible.

For those desirous of obtaining the latest information which is available on the practical side of the subject, Harrower's book well repays a careful perusal. Not only has this author collected a large amount of literature dealing with the internal secretions, but he has produced a book which is of really practical help to those desirous of prescribing organic extracts.

For a histological survey, Schäfer's recent work entitled "The Endocrine Glands" should be consulted; while Biedl's book on the "Internal Secretary Organs" is a careful and thorough account of the

subject from all standpoints, although it does not deal with hormone-therapy.

A list of notable contributions to this subject is appended.

BIBLIOGRAPHY.

- The Diseases of the Ductless Glands; for Practitioners of Medicine and Surgery. By S. P. Beebe, Ph.D., M.D. Phil.
- The Nervous and Chemical Regulators of Metabolism. By D. Noel Paton, M.D., B.Sc. Macmillan, London and New York.
- Organo-therapy; or, Treatment by Means of Preparations of Various Organs. By H. Batty Shaw, M.D. Lond., F.R.C.P. Cassell and Co., London.
- Glandes Surrénales et Organes Chromaffines. By Lucien et Parisot. Gittler, Paris.
- The Pituitary Body and its Disorders. By Harvey Cushing, M.D. Lippincott, Philadelphia.
- The Practitioner*, London. Special numbers for January and February, 1915, vol. xciv., Nos. 1 and 2.
- Glycosuria and Allied Conditions. By P. J. Cammidge, M.D. Lond. Arnold, London.
- Diseases of the Thyroid Gland. By George R. Murray, M.A., M.D. Lewis, London.
- New Aspects of Diabetes. By Professor Carl Von Noorden, Vienna. Wright, Bristol.
- The Endocrine Organs. By E. A. Schäfer. Longmans, Green and Co., London.
- Practical Hormone-Therapy. By H. R. Harrower, M.D., Bailliére, Tindall and Cox.

222 THE ORGANS OF INTERNAL SECRETION

Internal Secretion and the Ductless Glands. By Swale Vincent, M.D., D.Sc. Arnold, London.

The Internal Secretions and the Principles of Medicine. By Charles de M. Sajons, M.D., LL.D. Davis, Philadelphia; Phillips, London.

The Internal Secretary Organs: Their Physiology and Pathology. By Professor Dr. Arthur Biedl. With an Introductory Preface by Leonard Williams, M.D., M.R.C.P. Translated by Linda Forster.

INDEX

A				PAGES
ADDISON'S disease				122
Adrenal glands				114
" anatomy and physiology of				115
" cortex, composition of				117
" extract of				209
" insufficiency of				121
" medulla of				115-117
" therapeutics of				124, 127-128
Adrenalin				118
" characteristics of				118
" composition of				116
" discovery of				116
" effects of administration of				120
B				
Basch, experiments with placental extract				151
Basedow's disease (<i>see</i> Graves' Disease)				31
Baumann, and thyroid colloid				19
Bayliss and Starling, and secretin				165
Bibliography				221
Bile, therapeutics of				178, 210
Blair Bell, and calcium metabolism				158
" and osteomalacia				127
" and treatment by ovarian extract				159
Brown-Séquard, and testicular extract				193
C				
Cachexia strumipriva				17, 54
Campbell, H., and Graves' disease				51
Cassowary neck				81
Cheadle, and Graves' disease				45
Chromaffin system				117
Cod-liver oil, value of				171

224 THE ORGANS OF INTERNAL SECRETION

	PAGES
Cushing and hyperpituitarism	106
" and hypopituitarism	111
D	
Diabetes, islets of Langerhans in	134
" relativa to pancreas	135
" treatment of, by opo-therapy	142
" Von Noorden and	136
Digestion, internal secretions of	163 <i>et seq.</i>
" relation of spleen to	179
Dyspituitarism	102
" case of	203, 204
E	
Edkins, and gastrin	165
Endocrine glands, antagonistic action of	186
Entero-kinase	168
Eppinger and Hess, and Graves' disease	37
Erdheim, quoted	38
Erepsin	168
Exophthalmic goitre (<i>see Graves' Disease</i>)	31
F	
Falta, Eppinger, and Rudinger, and hyperthyroidism	37
G	
Gibson, and treatment of Graves' disease	186
Gley, experiments with thyroid extract	17
Glycosuria	134 <i>et seq.</i>
" and adrenalin	123, 130
" and removal of pancreas	134, 139
Goetsch, and pituitary extract	108
Gonads, extracts of	193
Graves' disease	31 <i>et seq.</i>
" definitions of	31, 32
" electrical resistance of skin in	40
" etiology	35
" histological changes in thyroid in	38
" history of	32
" insanity in	43
" mental changes in	41, 42
" prognosis in	44
" treatment in	46 <i>et seq.</i>
" tremor in	39-40
Gull and Ord, discovery of myxœdema by	56

INDEX

225

H	PAGES
Halliburton, W. D., quoted	133
Harrower, H. R., quoted .. 99, 109, 119, 123, 124, 125, 137, 143, 157, 167, 184, 195	135
Hedon, and glycosuria	72, 73, 74, 80, 83, 87, 88
Hertoghe, quoted	24, 25, 208
Hertz, quoted	178, 195
Hormonal	183-187
Hormone-therapy, rationale of	127
Houssay, and treatment by adrenal extract	129
Hyperadrenia, symptoms of	100
Hyperpituitarism	106
" treatment of	196
Hypertension, treatment of	31 <i>et seq.</i> , 65, 66
Hyperthyroidism (<i>see Graves' Disease</i>)	123
Hypo-adrenia	121
" symptoms of	124
" treatment of	109
Hypophysin	103
Hypopituitarism	103, 104
" symptoms	105 <i>et seq.</i>
" treatment of	54
Hypothyroidism (<i>see Thyroid Deficiency</i>)	65
" association with hyperthyroidism	59
" lesser degrees of	88-93, 198, 199, 208
" treatment of	
I	
Intestine, extracts of	165, 168, 210
" paresis of, treatment	195
K	
Kinazyme	144
Kocher, and cachexia strumipriva	17, 54
L	
Lane, Sir W. Arbuthnot, quoted	4, 76
Langerhans, islets of	132
Léopold-Levi and H. de Rothschild, quoted	65 <i>et seq.</i> , 71
Liver, extracts of	173, 175, 176, 210
" functions of	169-171
" production of bile by	171
" treatment of diabetes by extracts of	177
	15

226 THE ORGANS OF INTERNAL SECRETION

M				PAGE
Mackenzie, and Graves' disease	33
Mammary gland	150
" extracts of	200, 210
" relation to ovary	161
" relation to uterus	150
Mental chorea	43
Minkowski and V. Mering, and diabetes	134
Morris, and etiology of glycosuria	130-137
Murray, G. R., quoted	16, 54, 56
Muscular work, effects of orchitic extracts on	193
Myxœdema	56
" Murray's treatment of	54
" origin of name	18
" pathology of	17, 18, 20, 55
" symptoms of	57, 58
N				
Neo-hormonal	179, 195
O				
Opie, and diabetes	136
Osteomalacia, treatment by suprarenal extract	127
Ovaries, effects of removal of	152
" relation to mammary gland	161, 196
" relation to menorrhagia	159
" relation to osteomalacia	153
" therapeutics of	157, 158, 196, 209
P				
Pancreas, the	131 <i>et seq.</i>
" anatomy and physiology of	132
" origin of internal secretion	139
" relation to diabetes of	135
" secretion, external	133
" " internal	134
" therapeutic use of preparations of	141, 209
Paralysis agitans and parathyroid glands	24
Parathyroid glands	22
" " and calcium metabolism	23
" " and tetany	23
" " deficiency	25
" " extracts of	208, 210
Pawlow, and succus entericus	162

	PAGES
Pituitary body, the	94 <i>et seq.</i>
.. .. . effects produced by disease of	100
.. .. . physiology of	97
.. .. . structure of	96
.. .. . test for deficient secretion of	108
.. .. . treatment of disorders of	105
.. .. . by extracts of	180, 209
Pituitrin	100
Pluriglandular therapy	206-211
Pro-gastrin	165
Proptosis, causation of, in Graves' disease	38
Prostate, external secretion of	155
.. .. . internal secretion of	155
.. .. . preparation of	197, 210
R	
Reuon and Delile, experiment of	107
Reynolds, and "mental chorea"	43
S	
Sargent, and dermatographic sign	121
.. .. . and hypo-adrenia	120
Schäfer, Sir E., quoted	135, 142
Schiff, and removal of thyroid gland	17
Secretin	165-168
Secretogen	195
Sexual organs, internal secretions of	147
Short, A. Rendle, quoted	28, 48
Sleep in submyxœdema	80
Spermin	194
Spleen, extracts of	181, 182, 210
.. .. . functions as filter of	180
.. .. . relation to digestion	179-180
Stellwag's sign in Graves' disease	39
Stokes, and Graves' disease	33
Succus entericus	168
T	
Takamine and adrenaline	116
Tandler and Gross, a case described by	153
Testis, changes in disease of	155
.. .. . effects of removal	152
.. .. . external secretion of	155
.. .. . extracts of	157, 209
.. .. . internal secretion of	148

228 THE ORGANS OF INTERNAL SECRETION

	PAGES
Thymus, extracts of	206-208, 210
Thyroid and calcium metabolism	21
" and cell excretion	75
" antitoxic function of	75
" deficiency	54 <i>et seq.</i> , 71 <i>et seq.</i>
" " in children	73, 74
" " treatment of	88 <i>et seq.</i>
" diseases of	31 <i>et seq.</i> , 54 <i>et seq.</i>
" excess of	31 <i>et seq.</i>
" extract of	90-93, 188-191, 208
" " combined with adrenal extract	206
" functions of	27 <i>et seq.</i>
" instability	68
" results of removal of	20, 21
" weight of	19
Trousseau, and Graves' disease	33
Trypsogen	143, 195

V

Vassale, experiments with thyroid extract	17
Vigorous, and electrical resistance of skin in Graves' disease	40
Von Graefe's sign in Graves' disease	30
Von Mering and Minkowski, and diabetes	134
Von Noorden, and diabetes	136
Von Poehl's spermin	194

W

Williams, Leonard, quoted	41, 42, 50, 171
Wirsung, canal of	132

Z

Zuelzer's hormone	178, 195
---------------------------	----------

3
4
5
6
7
8
9
0
1
2
3
4
5

7
0
C
4
6
4

1
2

5

