

**CIHM
Microfiche
Series
(Monographs)**

**ICMH
Collection de
microfiches
(monographies)**



Canadian Institute for Historical Microreproductions / Institut canadien de microreproductions historiques

© 1997

Technical and Bibliographic Notes / Notes techniques et bibliographiques

The Institute has attempted to obtain the best original copy available for filming. Features of this copy which may be bibliographically unique, which may alter any of the images in the reproduction, or which may significantly change the usual method of filming are checked below.

- Coloured covers / Couverture de couleur
- Covers damaged / Couverture endommagée
- Covers restored and/or laminated / Couverture restaurée et/ou pelliculée
- Cover title missing / Le titre de couverture manque
- Coloured maps / Cartes géographiques en couleur
- Coloured ink (i.e. other than blue or black) / Encre de couleur (i.e. autre que bleue ou noire)
- Coloured plates and/or illustrations / Planches et/ou illustrations en couleur
- Bound with other material / Relié avec d'autres documents
- Only edition available / Seule édition disponible
- Tight binding may cause shadows or distortion along interior margin / La reliure serrée peut causer de l'ombre ou de la distorsion le long de la marge intérieure.
- Blank leaves added during restorations may appear within the text. Whenever possible, these have been omitted from filming / Il se peut que certaines pages blanches ajoutées lors d'une restauration apparaissent dans le texte, mais, lorsque cela était possible, ces pages n'ont pas été filmées.
- Additional comments / Commentaires supplémentaires:

L'Institut a microfilmé le meilleur exemplaire qu'il lui a été possible de se procurer. Les détails de cet exemplaire qui sont peut-être uniques du point de vue bibliographique, qui peuvent modifier une image reproduite, ou qui peuvent exiger une modification dans la méthode normale de filmage sont indiqués ci-dessous.

- Coloured pages / Pages de couleur
- Pages damaged / Pages endommagées
- Pages restored and/or laminated / Pages restaurées et/ou pelliculées
- Pages discoloured, stained or foxed / Pages décolorées, tachetées ou piquées
- Pages detached / Pages détachées
- Showthrough / Transparence
- Quality of print varies / Qualité inégale de l'impression
- Includes supplementary material / Comprend du matériel supplémentaire
- Pages wholly or partially obscured by errata slips, tissues, etc., have been refilmed to ensure the best possible image / Les pages totalement ou partiellement obscurcies par un feuillet d'errata, une pelure, etc., ont été filmées à nouveau de façon à obtenir la meilleure image possible.
- Opposing pages with varying colouration or discolourations are filmed twice to ensure the best possible image / Les pages s'opposant ayant des colorations variables ou des décolorations sont filmées deux fois afin d'obtenir la meilleure image possible.

This item is filmed at the reduction ratio checked below / Ce document est filmé au taux de réduction indiqué ci-dessous.

10x	14x	18x	22x	26x	30x
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
12x	16x	20x	24x	28x	32x

The copy filmed here has been reproduced thanks to the generosity of:



Library
Agriculture Canada

The images appearing here are the best quality possible considering the condition and legibility of the original copy and in keeping with the filming contract specifications.

Original copies in printed paper covers are filmed beginning with the front cover and ending on the last page with a printed or illustrated impression, or the back cover when appropriate. All other original copies are filmed beginning on the first page with a printed or illustrated impression, and ending on the last page with a printed or illustrated impression.

The last recorded frame on each microfiche shall contain the symbol \rightarrow (meaning "CONTINUED"), or the symbol ∇ (meaning "END"), whichever applies.

Maps, plates, charts, etc., may be filmed at different reduction ratios. Those too large to be entirely included in one exposure are filmed beginning in the upper left hand corner, left to right and top to bottom, as many frames as required. The following diagrams illustrate the method:



L'exemplaire filmé fut reproduit grâce à la générosité de:



Bibliothèque
Agriculture Canada

Les images suivantes ont été reproduites avec le plus grand soin, compte tenu de la condition et de la netteté de l'exemplaire filmé, et en conformité avec les conditions du contrat de filmage.

Les exemplaires originaux dont la couverture en papier est imprimée sont filmés en commençant par le premier plat et en terminant soit par la dernière page qui comporte une empreinte d'impression ou d'illustration, soit par le second plat, selon la cas. Tous les autres exemplaires originaux sont filmés en commençant par la première page qui comporte une empreinte d'impression ou d'illustration et en terminant par la dernière page qui comporte une telle empreinte.


Un des symboles suivants apparaîtra sur la dernière image de chaque microfiche, selon le cas: le symbole \rightarrow signifie "A SUIVRE", le symbole ∇ signifie "FIN".

Les cartes, planches, tableaux, etc., peuvent être filmés à des taux de réduction différents. Lorsque le document est trop grand pour être reproduit en un seul cliché, il est filmé à partir de l'angle supérieur gauche, de gauche à droite, et de haut en bas, en prenant le nombre d'images nécessaire. Les diagrammes suivants illustrent la méthode.




MICROCOPY RESOLUTION TEST CHART

(ANSI and ISO TEST CHART No. 2)



APPLIED IMAGE Inc

1653 East Main Street
Rochester, New York 14609 USA
(716) 482 - 0300 - Phone
(716) 288 - 5989 - Fax



JAN 5 1915

PLEASE RETURN.

CIRCULAR No. 41
(New Horticultural Series).



233

PROVINCE OF BRITISH COLUMBIA.

DEPARTMENT OF AGRICULTURE
(HORTICULTURAL BRANCH).

THE OYSTER-SHELL SCALE

(*Lepidosaphes ulmi* Linn.).

THE oyster-shell scale is a familiar object to fruit-growers. It occurs commonly in the Lower Kootenays, in the Lower Fraser Valley, and in the developed sections of Vancouver Island. It is



Twig infested by oyster-shell scale. (Author's illustration.)

present also in the Dry Belt fruit areas of the Okanagan, but it finds circumstances more suited to its taste in districts where the atmospheric conditions are moist and humid.

It derives its name, first, from the fact that it belongs to that class of insects which develops a waxy shield over its body, so-called a "scale," under which it lives, protected from any adverse conditions; secondly, from the fact that the waxy scale resembles an oyster-shell in shape. In colour this insect may be one of several shades of brown, depending very largely upon the colour of the bark upon which it is living. Fruit-growers, therefore, will be able to identify this scale by its colour, by its shape, by its size, which is approximately $\frac{1}{8}$ inch in length, and from the fact that it often occurs in great numbers on a piece of bark, often overlapping and forming deep encrustations. At times the fruit itself may possess an occasional specimen.

Food-plants.

With this insect, possibly more so than with any other insect which attacks cultivated fruit-trees in this Province, it is correct to agree with the common expression that the "woods are full of it." It is known to attack a wide variety of native plants, among which may be mentioned, Nuttall's dogwood, broom, willow, poplar, vine-maple, hawthorn, amelanchier, flowering currant, mountain-ash, and the cascara-tree. It is essentially, however, a fruit-infesting insect, attacking varieties of apple, crab-apple, pear, peach, apricot, cherry, and currant. Its presence is noted more often in apple-orchards than in plantations of the other fruits, and in certain sections of the Province its numbers have increased to such an extent that apple-trees, naturally weakened by several contributing causes, have finally succumbed to its attacks.

Life-history.

In the Province of British Columbia the oyster-shell scale has only one generation in the year. Its life-history is as follows: If during winter some of the familiar scales are overturned on the point of a penknife, a number of shining, white, oval objects will be seen, varying in number from 20 to 100. These are eggs, and it is in this stage that the winter is passed. As the spring arrives these eggs commence to hatch, and usually at the time the apple-trees are in blossom, small white or cream-coloured objects may be seen issuing from beneath the scale. These are the larvæ, or young scale-insects. Owing to differences in climate, which the various fruit districts of the Province present, and owing very often to a marked dissimilarity in the spring seasons at any one locality, one year with another, the period of hatching varies considerably. However, at some time in the four weeks surrounding the period of full bloom, hatching takes

place and the young larvæ become discernible to the naked eye as small cream-coloured dots, moving, in proportion to their size, comparatively rapidly over the bark. They are wingless and very delicate at this stage, and their sole object is to find a suitable spot upon which to settle and feed. After a few hours' activity they establish themselves, insert minute sucking mouth-organs, with which they are provided, into the bark and commence to suck the sap. Once the female has settled on a particular spot she never moves again to take up any other position on the tree. The young larvæ gradually excrete a waxy substance over their bodies which forms and hardens into the familiar scale or protective shield. In the autumn or during the winter the female, which remains for ever wingless, dies beneath the scale she has constructed, leaving behind her complement of eggs. The male insect undergoes a similar metamorphosis in the early part of the summer, but later in the year, about July, emerges with wings which apparently are only strong enough to enable it to reach and fertilize the stationary female.

Means of Control.

A study of the life-history of this insect reveals the facts (1) that it is single-brooded, (2) that it has a reproductive capacity of an approximate average of 60 eggs, and (3) that it is not capable of much movement. Thus a commercial orchardist has no right to claim this insect as an important fruit-tree pest, on the understanding that practical horticulture to-day demands proper attention to timely applications of spray. A thorough application of lime-sulphur, 1-30 or 35 (specific gravity 1.009), immediately after the petals fall, under ordinary circumstances suffices to hold this insect in check. Moderately severe cases of infestation may be treated also with a winter strength lime-sulphur solution at the time the buds are bursting, to act as an additional means of control.

In some of the older orchards of the Province, where trees are encrusted with superfluous bark and growth of lichens (locally referred to as "moss"), an application of Gillett's lye at the rate of 1 lb. to 4 gallons of water will be found advantageous. This may be performed in the autumn and applied direct to the trees by the most convenient method.

Victoria, B.C., issued March, 1918.

This circular has been prepared by R. C. Treherne, Field Officer for British Columbia, Entomological Branch, Dominion Department of Agriculture, at the request of the Horticultural Branch.

Copies of this circular may be obtained free of charge on application to the Horticultural Branch, Department of Agriculture, Victoria, B.C., or from local branch offices of the Department.

VICTORIA, B.C. :
Printed by **WILLIAM H. CULLIN**, Printer to the King's Most Excellent Majesty.
1918.



