




**IMAGE EVALUATION
TEST TARGET (MT-3)**



**Photographic
Sciences
Corporation**

23 WEST MAIN STREET
WEBSTER, N.Y. 14580
(716) 872-4503

18
20
22
25
28

**CIHM/ICMH
Microfiche
Series.**

**CIHM/ICMH
Collection de
microfiches.**



Canadian Institute for Historical Microreproductions / Institut canadien de microreproductions historiques

01

© 1981

Technical and Bibliographic Notes/Notes techniques et bibliographiques

The Institute has attempted to obtain the best original copy available for filming. Features of this copy which may be bibliographically unique, which may alter any of the images in the reproduction, or which may significantly change the usual method of filming, are checked below.

L'Institut a microfilmé le meilleur exemplaire qu'il lui a été possible de se procurer. Les détails de cet exemplaire qui sont peut-être uniques du point de vue bibliographique, qui peuvent modifier une image reproduite, ou qui peuvent exiger une modification dans la méthode normale de filmage sont indiqués ci-dessous.

- Coloured covers/
Couverture de couleur
- Covers damaged/
Couverture endommagée
- Covers restored and/or laminated/
Couverture restaurée et/ou pelliculée
- Cover title missing/
Le titre de couverture manque
- Coloured maps/
Cartes géographiques en couleur
- Coloured ink (i.e. other than blue or black)/
Encre de couleur (i.e. autre que bleue ou noire)
- Coloured plates and/or illustrations/
Planches et/ou illustrations en couleur
- Bound with other material/
Relié avec d'autres documents
- Tight binding may cause shadows or distortion along interior margin/
La reliure serrée peut causer de l'ombre ou de la distortion le long de la marge intérieure
- Blank leaves added during restoration may appear within the text. Whenever possible, these have been omitted from filming/
Il se peut que certaines pages blanches ajoutées lors d'une restauration apparaissent dans le texte, mais, lorsque cela était possible, ces pages n'ont pas été filmées.
- Additional comments:/
Commentaires supplémentaires:

- Coloured pages/
Pages de couleur
- Pages damaged/
Pages endommagées
- Pages restored and/or laminated/
Pages restaurées et/ou pelliculées
- Pages discoloured, stained or foxed/
Pages décolorées, tachetées ou piquées
- Pages detached/
Pages détachées
- Showthrough/
Transparence
- Quality of print varies/
Qualité inégale de l'impression
- Includes supplementary material/
Comprend du matériel supplémentaire
- Only edition available/
Seule édition disponible
- Pages wholly or partially obscured by errata slips, tissues, etc., have been refilmed to ensure the best possible image/
Les pages totalement ou partiellement obscurcies par un feuillet d'errata, une pelure, etc., ont été filmées à nouveau de façon à obtenir la meilleure image possible.

This item is filmed at the reduction ratio checked below/
Ce document est filmé au taux de réduction indiqué ci-dessous.

10X	14X	18X	22X	26X	30X
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
12X	16X	20X	24X	28X	32X

The c
to the

The li
possib
of the
filming

Original
beginning
the la
sion,
other
first p
sion,
or illu

The li
shall
TINU
which

Maps
differ
entire
begin
right
requir
metho

The copy filmed here has been reproduced thanks to the generosity of:


Library of the Public
Archives of Canada

The images appearing here are the best quality possible considering the condition and legibility of the original copy and in keeping with the filming contract specifications.

Original copies in printed paper covers are filmed beginning with the front cover and ending on the last page with a printed or illustrated impression, or the back cover when appropriate. All other original copies are filmed beginning on the first page with a printed or illustrated impression, and ending on the last page with a printed or illustrated impression.

The last recorded frame on each microfiche shall contain the symbol \rightarrow (meaning "CONTINUED"), or the symbol ∇ (meaning "END"), whichever applies.

Maps, plates, charts, etc., may be filmed at different reduction ratios. Those too large to be entirely included in one exposure are filmed beginning in the upper left hand corner, left to right and top to bottom, as many frames as required. The following diagrams illustrate the method:



L'exemplaire filmé fut reproduit grâce à la générosité de:


La bibliothèque des Archives
publiques du Canada

Les images suivantes ont été reproduites avec le plus grand soin, compte tenu de la condition et de la netteté de l'exemplaire filmé, et en conformité avec les conditions du contrat de filmage.

Les exemplaires originaux dont la couverture en papier est imprimée sont filmés en commençant par le premier plat et en terminant soit par la dernière page qui comporte une empreinte d'impression ou d'illustration, soit par le second plat, selon le cas. Tous les autres exemplaires originaux sont filmés en commençant par la première page qui comporte une empreinte d'impression ou d'illustration et en terminant par la dernière page qui comporte une telle empreinte.

Un des symboles suivants apparaîtra sur la dernière image de chaque microfiche, selon le cas: le symbole \rightarrow signifie "A SUIVRE", le symbole ∇ signifie "FIN".

Les cartes, planches, tableaux, etc., peuvent être filmés à des taux de réduction différents. Lorsque le document est trop grand pour être reproduit en un seul cliché, il est filmé à partir de l'angle supérieur gauche, de gauche à droite, et de haut en bas, en prenant le nombre d'images nécessaire. Les diagrammes suivants illustrent la méthode.



208
Extracted from THE OTTAWA NATURALIST, Vol. XI. No. 12, pp. 209-220,
Ottawa, March, 1898.

**NOTES ON THE PLEISTOCENE GEOLOGY OF A
FEW PLACES IN THE OTTAWA VALLEY.**

BY

W. J. WILSON, Ph. B.,
of the Geological Survey of Canada.

1898(6)

1898(6)

...

...

75357

NOTES ON THE PLEISTOCENE GEOLOGY OF A
FEW PLACES IN THE OTTAWA VALLEY.

By W. J. WILSON, Ph. B.,
Of the Geological Survey of Canada.

The Ottawa River rises near the height of land in Latitude 48° N. and Longitude 76° W., whence it flows westwardly for a distance of 250 miles to Lake Temiscaming. From this lake its course is south-east till it reaches the St. Lawrence River. Its whole length is about 650 miles, but from its source to the St. Lawrence River in a direct line is less than 200 miles. Its branches on the north or Quebec side are the Dumoine, Black, Coulonge, Gatineau, Lièvre, and Rouge, all of which flow nearly south. From the west or Ontario side it receives the waters of the Montreal, Mattawa, Petewawa, Bonnechere, Madawaska, Mississippi, Rideau, and South Nation, all of which flow eastward. The whole area drained by the Ottawa is approximately fifty-six thousand square miles.

It is the purpose of this paper to record the result of observations made at a few points extending from the city of Ottawa to Pembroke, a distance of eighty miles, during the summers of 1895-96, more especially the district included in the townships of Ros3, Westmeath, Stafford, Wilberforce, Bromley, S. Algona, Sebastopol and Grattan, in Renfrew county.

Everywhere in this district there is abundant evidence of ice action, both in the polished and striated rocks and in the general distribution of boulder-clay and boulders. The question of the direction of the ice movement is comparatively simple. Near the Ottawa River the striæ follow closely the course of the

valley as seen north of Allumette Island in two places, at Vinton, Portage du Fort, and near Bristol, on the Quebec side, where the course is from S. 10° E. to S. 30° E.* On the south side the same courses were seen north of Pembroke, at several places in Ross and Bromley townships; at Shamrock, south of Renfrew, and at Galetta. In Ottawa city, "Barrack Hill,"† the course is S. 45° E., and Dr. Ami reports striæ on Park Avenue and Nicholas Street almost due east and west.‡ Near Hintonburg on the Ottawa, Arnprior and Parry Sound Railway, the course is S. 87° E. While the south-east course is constant in the valley close to the river, at a distance of ten to twenty miles back the ice moved west of south. On the south side of the river a large number of observations show the direction to be from S. 2° W. to S. 35° W., the most common being from S. 15° W. to S. 25° W. It is probable that the south-west course is the older of the two, and that the south-east course was produced when the ice had become so thin that it was deflected by the minor irregularities of the surface, and so followed the course of the river. At an earlier stage the ice had evidently been thick enough to over-ride irregularities of surface of considerable size, the course being south-west, as above stated, where the present drainage is to the east. The south-east and the south-west courses were not observed on the same surface in the area under consideration, but in the vicinity of Lake Temiscaming these two courses are frequently seen crossing each other, and Mr. Barlow states that the oldest course is about S. 20° W., while the more recent courses follow the river valleys.

Speaking generally of the glaciation of the district particularly referred to, it may be said that good exposures of rock are common where the results of ice movement can be studied.

* The bearings are referred to the true meridian.

† Geology of Canada, p. 892.

‡ Ottawa Naturalist, 1887, p. 69.

Splendid examples of stossing are numerous, and even where the rock is so weathered that no striæ are visible, the rounded and smooth appearance of the north side of exposures, and the abrupt and sharp edges on the south, so characteristic of glacier action, may be seen and often enable us to determine with considerable accuracy the direction in which the ice moved. Boulder-clay, boulders, Leda-clay, sand and gravel are also abundant, while the less common phenomena of surface geology, viz.: kames, *asar* or eskers and moraines are occasionally seen.

BOULDER-CLAY.

Till, or boulder-clay has been described as a "firm, tough tenaceous clay which gives evidence of having been subjected to great pressure. Often the accumulation becomes coarser and sandier. Again it may be described as a coarse agglomeration of subangular and angular stones set in a scanty matrix of coarse earthy grit and sand. Sometimes the stones in the till are so numerous that hardly any matrix of clay is visible." It will be seen that the term boulder-clay embraces deposits whose appearance differs widely, but however it may vary in appearance and composition it can usually be recognized by the peculiar shape and striation of the stones contained in it. Typical boulder-clay may be seen in many places near Ottawa, as in the cutting on the Ottawa, Arnprior and Parry Sound Railway near Hintonburg already referred to, at Hog's Back, and in very many places in the area under consideration. In connection with the boulder-clay a word about the distribution of boulders will be in place. This district is no exception to the general rule that most of the boulders in the boulder-clay and those scattered over a country are from rocks near at hand, and only a few of the harder kinds are carried to a great distance. In this district the boulders commonly seen are limestone, usually flat and angular, and gneiss, granite, etc., more rounded and worn. Dr. Ells has already referred to the great blocks of Black

River limestone which occur along the Opeongo Road on a ridge south of Clear Lake, at a height of nearly fourteen hundred feet above sea-level.* Some of these blocks are five to eight feet through and very angular. As far as known the Black River formation nowhere occurs in the vicinity of Clear Lake at a greater height than eight hundred feet, so that if the relative levels of the country have not changed since the glacial period, these boulders have been carried upward a distance of five to six hundred feet. Travelled boulders in similar positions have been frequently noted. Sir J. W. Dawson records large Laurentian boulders on Montreal Mountain which, he says, must have been carried probably a hundred miles from the Laurentian region to the north-east.† Dana states that Mount Katahdin in Maine has many boulders on its northern face derived from the Devonian rock of the low country to the north, three thousand feet below it in level.‡ In Nova Scotia sandstone boulders are common on the Cobequid Mountains at a considerable height above the present level of the Carboniferous beds, from which they were derived. The position of these boulders at such heights forms a most interesting subject for study, and many theories have been advanced in regard to it. Among the theories put forward the following may be mentioned. Some writers claim that these erratics were placed in their present position by floating ice. They claim that the land was submerged to a depth sufficient to allow icebergs or ice jams to pass over or become stranded on the higher ground, where they deposited whatever material was embedded in them, or carried on their surface. Another explanation is that the land was covered by a glacier to a depth equal to or more than the greatest height at which these boulders are found, and that this mass moved over the country, carrying boulders, etc., along with

* The Ottawa Naturalist, December, 1896, p. 171.

† Canadian Ice Age, p. 201.

‡ Manual, p. 690.

it and deposited them wherever it chanced to be when the ice ceased to move forward. In connection with this it is urged that blocks of stone, entering the bottom of the glacier under certain conditions, gradually rise to the surface and in this way often reach a height far above their original position. It may be remarked that more or less serious objections have been raised against all these explanations.

LEDA CLAY.


The next formation in ascending order is the Leda clay, which is common in all the lower levels of the Ottawa Valley, where it forms broad stretches of level country. It is a bluish-gray clay, somewhat unctuous, and when dry becomes very hard and cracks into square or oblong shaped blocks. In Ross and Bromley townships there are large areas covered with this clay. In one place in Ross is a plain five miles across which extends lengthwise for a much greater distance. The approximate height of these plains is from four hundred to four hundred and fifty feet. Though the roads traversed passed over many miles of this formation the only fossil found was a fragment of a shell which was too small to be identified.

For a description of this deposit below the City of Ottawa, I quote the following. "Along the south bank of the Ottawa River from the city of Ottawa to Hawkesbury, and again from Point Fortune to its junction with the St. Lawrence the lower clay is seen in banks of from twenty to forty feet in height. . . . The greatest breadth of the level clay surface which has been observed here is in the township of L'Orignal, where it extends about fifteen miles back from the river. . . . On the north side of the Ottawa, from Hull to Isle Jesus, the clay often covers a considerable breadth between the river and the Laurentian Hills and extends among these for several miles up the larger tributaries."* Mr. A. E. Barlow states that this clay is abundant at

* Geology of Canada, p. 916.

Lake Temiscaming, the level of which at high water is five hundred and ninety-six feet above the sea. Perhaps the most remarkable thing about this clay is the scarcity of marine shells even where it is known to be far below the level reached by the sea during the Champlain subsidence. Marine fossils are recorded at Montreal at a height of five hundred and sixty feet, at Smith's Falls four hundred and twenty feet, near Galetta four hundred and seventy-five and Chelsea four hundred and twenty-five, so that the land in this valley during the time the Leda clay was laid down must have been six hundred feet lower than at present. We can therefore, I think, fairly assume that the stratified clays which are not more than five hundred or six hundred feet above sea level are marine. The marked resemblance of the clays on the higher levels to those on the lower, where fossils are found, is strong corroborative evidence of a similar origin. Even at the lower levels fossils are by no means common in this clay. In the city of Ottawa where excavations are frequently made and large quantities of clay are thrown out, I have seen fossils in two places only. At Mohr's Corners, about a mile from the village of Galetta, there is a sand terrace abounding in marine shells. Underlying the sand there is a bed of this clay, well stratified, twenty to thirty feet deep, and although there was a section ten feet deep on the roadside for a quarter of a mile, a careful examination revealed no fossils. Sir J. W. Dawson says: "Where the Leda clay is thick and well developed it admits of sub-division into a lower Leda clay, unfossiliferous or with only shells of *Leda glacialis* and *Macoma Greenlandica*, and an upper Leda clay, usually more sandy and holding a rich boreal fauna identical with that of the northern part of the Gulf of St. Lawrence at present."* Mr. F. B. Taylor in a recent article says "Near the city of Ottawa the upper limit

* Canadian Ice Age, p. 60.



of the Champlain submergence was not less than four hundred and seventy-five feet above the sea and was probably higher. Beaches undoubtedly belonging to this submergence have been found near Renfrew, and also at about four hundred and fifty feet at Pembroke, and also at about five hundred and thirty feet at Mackey's Station on the Ottawa River. Between Hudson Bay and Lake Superior shells of the same age have been found up to four hundred and fifty feet above the sea, and within one hundred and fifty miles of the lake. The upper limit of the submergence was probably still higher. It seems almost certain that during the Champlain submergence the sea extended far up the Ottawa valley, probably reaching the head of Lake Temiscaming." He says, however, that within the area no continuous tracing of the marine beaches has been made.

SAXICAVA SAND.

The Saxicava sand is also well represented in the Ottawa valley. It is a shallow water or shore deposit, and where a section is complete rests on the Leda clay. It is yellow or brownish and varies from fine sand to coarse gravel. Sometimes there is a distinct line between this formation and the Leda clay, while in other places the one runs into the other. It is at the junction of the two that the fossils are most plentiful. Exposures of this sand are abundant everywhere in this valley, and in many places nothing else is seen for a long distance. Many such areas were noted in Renfrew county, but no fossils were seen. In a cutting a short distance north of Chelsea Station on the Gatineau Valley Railway there is a narrow seam of coarse yellow sand which in places is full of shells of *Saxicava rugosa* and *Macoma fragilis*. Near this seam, if not in it, a small perfect shell of *Leda arctica* was found, and in a higher bank of Leda clay a fragment of a *Balanus*, probably *Hameri*. These deposits attain a height of four hundred and twenty-five feet at this point. About half a mile west of Carp station this sand forms

a terrace known as Johnston's Grove, in which *Saxicava rugosa* and *Macoma fragilis* are abundant at a height of about three hundred and fifty feet. These shells are also found in a cutting at Carp Station, as stated by Dr. Ells. At Mohr's Corners, already referred to, there is a sand terrace which rises to a height of four hundred and seventy-five feet and contains the above shells in great abundance. In a cutting through a gravel bed on the Electric Railway west of Hintonburg, *Leda arctica*, *Saxicava rugosa* and a fragment of a *Balanus* were found. This is so different from the other localities that it deserves special mention. It is composed of a beach-like gravel, distinctly stratified and well water-worn. The pebbles are generally one to two inches in diameter, with a small number of larger ones, the interstices being filled with sand. *Leda arctica* is the most abundant here, while at the other places named, only one specimen was found. The shells are small but well preserved, and in some cases the two valves are joined. The few specimens of *Saxicava rugosa* which were found were also well preserved. This cutting is on the edge of a terrace which extends back to the Ottawa, Arnprior and Parry Sound Railway and is about two hundred and thirty feet above sea level.

TERRACES.

Terraces and old shore lines or beaches have been described as occurring in many places along the Ottawa Valley. A good example of a cut terrace may be seen on the Montreal road near Green's Creek, at a height of about two hundred feet, but this may be of fluvial origin. At Chelsea there is a terrace at a height of three hundred and fifty feet, and one round the base of King's Mountain seven hundred and five feet high, North of the west end of Muskrat Lake on the road leading to Beachburg, a fairly well defined shore-line occurs at a height of from four hundred to four hundred and fifty feet. In many places where the level tracts of Leda clay occur there are sloping

ridges rising on each side fifty to one hundred feet, and just where one would expect to find a cut terrace, but none were seen that could safely be put down as such. On the north-west side of Lake Doré there is a well defined gravel terrace at a height of four hundred and fifty feet above sea level and forty feet above the level of the lake.

A few sand and gravel ridges occur in this district, though none of any great length were seen. Two small ridges occur in Westneath township. They are composed largely of sand and gravel, and are ten to fifteen feet high. They lie about S. 30° E. Another low ridge was seen near Caldwell Station on the Ottawa, Arnprior and Parry Sound Railway, though the material in this one is more of the character of boulder clay. On the road between Concessions XXI and XXII, Lot 12, Wilberforce township, just east of the stream from Green Lake there are four ridges running parallel to each other, course S. 23° E. The first is only a few rods from the stream which cuts through it south of the road. In this part it is very regular in shape and rises about twenty feet above the surrounding ground. Where the road cuts through it, it is about two hundred and twenty-five feet wide, flanked on each side by a low ridge four or five feet high. The sides are steep and meet in a sharp ridge at the top. Between the first and second ridges there is a peat bog two hundred feet wide. The second ridge is about the same size as the first, then another narrow bog, and the third crosses the road. It is similar to the other two only not quite as high. A short distance from this is a fourth ridge, but it is low and rounded at the top. A good section is shown in each where the road crosses. They are composed of well rounded water-worn pebbles, sand and gravel and a few angular fragments of rock, but the second one mentioned differs considerably from the others. A section in descending order is—rounded pebbles, sand and gravel, five feet; fine sand well stratified

twelve feet, covered near the base with boulder clay. Striae on the same road a short distance to the east run S. 2° W. These ridges do not extend for any great distance as far as could be seen from the road, but on the next Concession road to the north-west, I saw short ridges composed partly of boulder-clay and partly of water-worn material, which were in the same line with those just described but not continuous. Gravel ridges and mounds occur near Beachburg and eastward and enclose small lakes. Mounds partaking of the character of drumlins occur on the road between Concessions VIII and IX, Wilberforce township.

MORAINES.

The Geology of Canada mentions ridges of glacial drift or boulders running north and south and resembling moraines as occurring at the Hudson Bay Company's post eighteen miles from the head of Lake Temiscaming, at Long Sault just below the Lake, at the mouth of the Maganissippi, about twenty miles above the Mattawa, at Roche Capitaine, near Green's Creek, and at L'Original where six ridges occur in less than a mile. Along the south side of Mink Lake in Wilberforce township there is an irregular mass which extends continuously more than a mile and rises in places to thirty feet in height. It has an average breadth of three hundred and fifty feet. It does not rise to a sharp ridge like those at Green Lake just referred to, but is flat on the top and in some parts there are two or more ridges parallel, or diverging at an angle of from 10° to 20°. At the west end it is broken up into hillocks or mounds and at the east end it is lower and bends toward the north following the shore of the lake, but it does not continue across the marsh at the north-east end of the lake. A low ridge which may be a continuation of it was seen to the east. Many large angular blocks of limestone are scattered over the surface, and sections both transverse and lateral showed that it was composed of clay and

angular or sub-angular stones such as are seen in ordinary till. Water-worn or stratified material was seen in it. Its approximate height is five hundred feet above sea level, course nearly east and west. Striæ on road along side of ridge. Course S 7° E.

In Hull there are several moraine-like ridges. One crosses Bridge street about half a mile from the Ottawa River, and is a mass of large limestone blocks mixed with a small quantity of sand and gravel. It reaches a height of ten to twenty feet and is three hundred feet wide. It lies in an east and west direction. It extends east of Bridge street a short distance and west to the next street where there is a break, but about two hundred feet to the north another ridge of similar character begins and continues westward to Brewery Creek. Two short ridges strike off from this at right angles to the south but extend only for a short distance. A section is exposed along the south side of the long ridges and clearly shows their composition. They are made up of large, flat, angular blocks of limestone (some are ten feet long and nearly as wide), arranged in layers overlapping each other and generally dipping south at different angles. This feature is very noticeable in walking over the ridges and is also seen in the lateral sections. On the surface no fine material was observed, but down in the mass there is a small quantity of water-worn gravel and stones, an occasional rounded boulder from the Laurentian rocks occurs. As far as observed these are well rounded and smooth but not striated.

The underlying limestone is polished and bears distinct striæ, course S. 60° E. Not more than one hundred and fifty feet from the western and longest part of the ridge an almost parallel ridge lies to the north. This is composed almost entirely of well rounded Laurentian boulders, gneiss, granite, etc., with an occasional flat limestone block. There are no openings in it, so that it is impossible to say what it is like below the surface, but judging from what can be seen there is more clay mixed with the boulders than in the first mentioned, It is crescent shaped

the concave side, facing the north, and is highest midway tapering and spreading out towards both ends. Further observation has shown that these ridges with slight breaks extend eastward through the city of Hull toward Lake Flora. Northward as far as the Gatineau River there are many low ridges of rounded boulders of much the same material as in the most northerly ridge referred to above.

The diversity both in the character and condition of the material composing the first ridges described, taken in connection with their proximity, makes it somewhat difficult to account for their origin. Those lying to the north and extending to the Gatineau River are probably moraines, and were left in their present position by ice moving down the Gatineau Valley.

The angular condition of the blocks in the most southerly ridge shows that they have not been transported any very great distance. The following section is from this ridge near Chaudiere street and is in descending order,

- | | |
|---|---------|
| 1. Large angular blocks of limestone mixed with sand and gravel, and an occasional rounded boulder of granite, etc..... | 8 feet |
| 2. Fine sand and gravel..... | 2 " |
| 3. Fine tough bluish stratified clay (Leda clay)..... | 1 1/4 " |
| 4. Boulder clay..... | 3 " |
| 5. Limestone rock in place, glaciated, striæ, course S. 60° E. | |

If No. 1 of this section is of morainic origin, then after the ice which glaciated the underlying rock had receded, leaving the boulder-clay, the land remained submerged long enough to admit of the deposition of the clay and sand. (Nos 3 and 2 of the above section) after which the ice again advanced, and without displacing the underlying material deposited the limestone blocks composing the ridge. Another explanation is that it was formed by ice jams in the Ottawa River when it flowed through the channel where Brewery Creek now runs.

[Faint, illegible text, possibly bleed-through from the reverse side of the page. The text is too light to transcribe accurately.]

